
0037854

ROYAL BOTANIC GARDENS,
KEW.

DUPLICATE 02

Digitized by the Internet Archive
in 2013

http://archive.org/details/indexkewensispla11hook

INDEX KEWENSIS

INDEX KEWENSIS

AN ENUMERATION OF THE GENERA AND SPECIES

OF

FLOWERING PLANTS

FROM THE TIME OF LINNAEUS TO THE YEAR 1885 INCLUSIVE
TOGETHER WITH THEIR AUTHORS' NAMES, THE WORKS IN
WHICH THEY WERE FIRST PUBLISHED, THEIR NATIVE COUNTRIES
AND THEIR SYNONYMS

COMPILED AT THE EXPENSE OF
THE LATE CHARLES ROBERT DARWIN
UNDER THE DIRECTION OF JOSEPH D. HOOKER

BY

B. DAYDON JACKSON

VOLUME I

OXFORD
AT THE CLARENDON PRESS
M DCCC XCV

INDEX KEWENSIS

PLANTARUM PHANEROGAMARUM

NOMINA ET SYNONYMA OMNIUM GENERUM ET SPECIERUM

A LINNAEO USQUE AD ANNUM MDCCCLXXXV

COMPLECTENS

NOMINE RECEPTO AUCTORE PATRIA

UNICUIQUE PLANTAE SUBJECTIS

SUMPTIBUS BEATI CAROLI ROBERTI DARWIN

DUCTU ET CONSILIO JOSEPHI D. HOOKER

CONFECIT B. DAYDON JACKSON

TOMUS I

OXONII

E PRELO CLARENDONIANO

M DCCCXCV

PREFACE

SHORTLY before his death, Mr. DARWIN informed me of his intention to devote a considerable sum in aid or furtherance of some work of utility to biological science; and to provide for its completion, should this not be accomplished during his lifetime. He further informed me that the difficulties he had experienced in accurately designating the many plants which he had studied, and ascertaining their native countries, had suggested to him the compilation of an Index to the Names and Authorities of all known Flowering Plants and their Countries, as a work of supreme importance to Students of Systematic and Geographical Botany, and to Horticulturists, and as a fitting mode of fulfilling his intentions.

I have only to add that, at his request, I undertook to direct and supervise such a work; and that it is being carried out at the HERBARIUM OF THE ROYAL GARDENS, KEW, with the aid of the staff of that establishment.

JOS. D. HOOKER

Royal Gardens, Kew: July, 1893.

MONITUM

PAULLO ante quam mortuus est, ostendit mihi vir clarissimus CAROLUS ROBERTUS DARWIN sibi esse in animo pecuniam satis magnam in opus aliquod conferre quod usui foret scientiae biologicae studiosis, eamque pecuniam ita collocare ut quodcunque opus designavisset ad finem perduceretur etiamsi se vivo non potuisset absolvi. Illud praeterea addidit, expertum se, cum tot plantarum naturas exploraret, quam difficile esset nomen proprium unicuique tribuere et patriam indagare, de indice conficiendo cogitasse, qui plantarum omnium phanerogamarum, quotquot cognitae essent, nomina auctores patrias complecteretur: nihil certe maioris pretii fore aut Botanicae systematicae et geographicae studiosis aut hortorum cultoribus, nihil voto suo et consilio convenientius.

Restat ut dicam suscepisse me eius rogatu operis propositi regimen, et id agere hodie in HERBARIO HORTI REGII KEWENSIS, adiuvantibus eiusdem curatoribus, ut numeris omnibus absolvatur.

JOS. D. HOOKER

Scribebam in Horto Regio Kewensi, m. Julio, a. 1893.

LONDINI

APUD HENRICUM FROWDE

NOVI EBORACI

APUD A. MACMILLAN ET SOCIOS

OXONII: EXCUDEBAT HORATIUS HART, TYPOGRAPHICUS ACADEMICUS

INDEX KEWENSIS

INDEX KEWENSIS

AN ENUMERATION OF THE GENERA AND SPECIES

OF

FLOWERING PLANTS

FROM THE TIME OF LINNAEUS TO THE YEAR 1885 INCLUSIVE
TOGETHER WITH THEIR AUTHORS' NAMES, THE WORKS IN
WHICH THEY WERE FIRST PUBLISHED, THEIR NATIVE COUNTRIES
AND THEIR SYNONYMS

COMPILED AT THE EXPENSE OF
THE LATE CHARLES ROBERT DARWIN
UNDER THE DIRECTION OF JOSEPH D. HOOKER

BY

B. DAYDON JACKSON

PART 1

OXFORD
AT THE CLARENDON PRESS
M DCCC XCIII

INDEX KEWENSIS

PLANTARUM PHANEROGAMARUM

NOMINA ET SYNONYMA OMNIUM GENERUM ET SPECIERUM

A LINNAEO USQUE AD ANNUM MDCCCLXXXV

COMPLECTENS

NOMINE RECEPTO AUCTORE PATRIA

UNICUIQUE PLANTAE SUBJECTIS

SUMPTIBUS BEATI CAROLI ROBERTI DARWIN

DUCTU ET CONSILIO JOSEPHI D. HOOKER

CONFECIT B. DAYDON JACKSON

FASCICULUS I

OXONII

E PRELO CLARENDONIANO

MDCCCXCIII

PREFACE

SHORTLY before his death, Mr. DARWIN informed me of his intention to devote a considerable sum in aid or furtherance of some work of utility to biological science; and to provide for its completion, should this not be accomplished during his lifetime. He further informed me that the difficulties he had experienced in accurately designating the many plants which he had studied, and ascertaining their native countries, had suggested to him the compilation of an Index to the Names and Authorities of all known Flowering Plants and their Countries, as a work of supreme importance to Students of Systematic and Geographical Botany, and to Horticulturists, and as a fitting mode of fulfilling his intentions.

I have only to add that, at his request, I undertook to direct and supervise such a work; and that it is being carried out at the HERBARIUM OF THE ROYAL GARDENS, KEW, with the aid of the staff of that establishment.

JOS. D. HOOKER

MONITUM

PAULLO ante quam mortuus est, ostendit mihi vir clarissimus CAROLUS ROBERTUS DARWIN sibi esse in animo pecuniam satis magnam in opus aliquod conferre quod usui foret scientiae biologicae studiosis, eamque pecuniam ita collocare ut quodcunque opus designavisset ad finem perduceretur etiamsi se vivo non potuisset absolvi. Illud praeterea addidit, expertum se, cum tot plantarum naturas exploraret, quam difficile esset nomen proprium unicuique tribuere et patriam indagare, de indice conficiendo cogitasse, qui plantarum omnium phanerogamarum, quotquot cognitae essent, nomina auctores patrias complecteretur: nihil certe maioris pretii fore aut Botanicae systematicae et geographicae studiosis aut hortorum cultoribus, nihil voto suo et consilio convenientius.

Restat ut dicam suscepisse me eius rogatu operis propositi regimen, et id agere hodie in HERBARIO HORTI REGII KEWENSIS, adiuvantibus eiusdem curatoribus, ut numeris omnibus absolvatur.

JOS. D. HOOKER

OXONII
Excudebat Horatius Hart
Typographus academicus

INDEX KEWENSIS

PLANTARUM PHANEROGAMARUM

NOMINA ET SYNONYMA OMNIUM GENERUM ET SPECIERUM

A LINNAEO USQUE AD ANNUM M.DCCC.LXXXV

COMPLECTENS

AA—ABELMOSCHUS.

AA, Rcichb. f. Xen. Orchid. i. 18 (1858) = **Altensteinia**, H. B. & K. (Orchid.).
argyrolepis, Reichb. f. l. c. = Alt. argyrolepis.
paleacea, Reichb. f. l. c. = Alt. paleacea.

AAMA, Hassk. Cat. Hort. Bog. Alt. 171 (1844), sphalm. = ADAMIA, Wall. = **Dichroa**, Lour. (Saxifr.).

ABACOSA, Alef. in Bonplandia, ix. (1861) 102 = **Vicia**, Linn. (Legumin.).
americana, Alef. l. c. 104 = V. americana.
canescens, Alef. l. c. 103 = V. canescens.
dumetorum, Alef. l. c. 104 = V. dumetorum.
Kingii, Alef. l. c. 103 = V. Kingii.
micrantha, Alef. l. c. 104 = V. micrantha.
onobrychioides, Alef. l. c. 103 = V. onobrychioides.
pallida, Alef. l. c. = V. pallida.
patagonica, Alef. l. c. = V. patagonica.

ABALON, Adans. Fam. ii. 511 (err. typ. 512) (1763) = **Helonias**, Linn. (Liliac.).
albiflorum, Rafin. New Fl. Am. i. 33.—Am. bor.

ABALUM, Adans. Fam. 47 (1763) = praec.

ABAMA, Adans. Fam. ii. 47 (1763) = **Narthecium**, Moehr. (Liliac.).
anthericoides, Steud. Nom. ed. I. 550, in syn. (1821) = N. ossifragum.
glutinosa, Rafin. New Fl. Am. i. 36 = Tofieldia glutinosa.
ossifraga, DC. Fl. Fr. iii. 171 = N. ossifragnm.
pubens, Rafin. New Fl. Am. i. 35 = Tofieldia pubens.
pusilla, Rafin. l. c. 36.—Am. bor.
scabra, Rafin. l. c. 35; Fl. Tellur. 28 = Tofieldia pnbens.

ABANDION, Adans. Fam. ii. 54 (1763) = **Bulbocodium**, Linn. (Liliac.).

ABAPHIUS, Rafin. Fl. Tellur. iv. 19 (1836) = seq.

ABAPUS, Adans. Fam. ii. 57 (1763) = **Gethyllis**, Linn. (Amaryll.).
spiralis, Rafin. Fl. Tellur. iv. 19 = G. spiralis.

ABASICARPON, Andrz. ex Reichb. in Moessl. Handb. ed. II. 1150 (1858) = **Arabis**, Linn. (Crucif.).
patulum, Andrz. ex Reichb. l. c. = Arab. auriculata.

ABASOLOA, La Llave, in Nov. Veg. Desc. fasc. i. 11 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 362. Taboada, *La Llave, l. c.*—Mexic.

ABATIA, Ruiz & Pav. Prod. 78. t. 14 (1794). *SAMYDACEAE*, Benth. & Hook. f. i. 799.
GRANIERA, Mand. & Wedd. ex Benth. & Hook. f. l. c.
MYRIOTRICHE, Turcz. in Bull. Soc. Nat. Mosc. (1863), I. 554.
RALEIGHIA, Gardn. in Hook. Lond. Journ. Bot. iv. (1845). 97.

†

ABATIA:—
americana, *Eichl. in Mart. Fl. Bras.* xiii. I. 510.—Bras.
brasiliensis, *Baill. Adansonia*, x. (1872) 245.—Bras.
parviflora, *Ruiz & Pav. Syst.* i. 136; *Prod.* t. 14.—Peruv.; N. Granat.
rugosa, *Ruiz & Pav. ll. cc.*—Peruv.
tomentosa, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. I. 509.—Bras.
verbascifolia, H. B. & K. Nov. Gen. et Sp. v. 358 = parviflora.

ABAURIA, Becc. Malesia, i. (1877) 169. *LEGUMINOSAE*.
excelsa, *Becc. l. c.*—Malaya.

ABAZICARPUS, Andrz. ex DC. Syst. ii. 213 (1821) = **Arabis**, Linn. (Crucif.).

ABBEVILLEA, Berg, in Linnaea, xxvii. 425 (1856) = **Campomanesia**, Ruiz & Pav. (Myrtac.).
cerasoides, Berg, in Mart. Fl. Bras. xiv. I. 436.—Bras.
chrysophylla, Berg, l. c. 437 = Psidium chrysophyllum.
eugenioides, Berg, l. c. 436.—Bras.
Fenzliana, Berg, l. c. 433.—Bras.
Gardneriana, Berg, l. c. 436.—Bras.
Guaviroba, Berg, l. c. 432 = C. Guaviroba.
Klotzschiana, Berg, l. c. 434.—Bras.
Langsdorffiana, Berg, l. c. 608.—Bras.
Martiana, Berg, l. c. 435.—Bras.
maschalantha, Berg, l. c. 432.—Bras.
neriiflora, Berg, l. c. 431.—Bras.
phaea, Berg, l. c. 435.—Bras.
plana, Berg, in Linnaea, xxvii. (1854) 427 (sphalm. = *phaea*).
punctulata, Berg, in Mart. Fl. Bras. xiv. I. 432.—Bras.
recurvata, Berg, l. c. 438.—Bras.
Regeliana, Berg, l. c. 608.—Bras.
rufa, Berg, l. c. 437 = Psidium rufum.
rugosa, Berg, l. c.—Bras.
Schlechtendahliana, Berg, l. c. 435.—Bras.
Sellowiana, Berg, l. c. 431.—Bras.
Widgreniana, Berg, in Linnaea, xxxi. (1861–62) 260.—Bras.

ABBOTIA, Rafin. New Fl. Am. i. 36 (1836) = **Triglochin**, Linn. (Naiad.).
filiformis, Rafin. l. c. = T. triandrum.
palustris, Rafin. l. c. 37 = T. palustre.
pumila, Rafin. l. c. = T. palustre.

ABBOTTIA, F. Muell. Fragm. ix. 181 (1875) *RUBIACEAE*.
singularis, *F. Muell. l. c.*—Anstral.

ABELIA, R. Br. in Abel, Chin. App. 376 (1818) ii. 4. *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 4.
adenotricha, *Hance, in Journ. Bot.* ix. (1871) 132.—China.
biflora, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) VII. 152.—China.
chinensis, *R. Br. in Abel, Chin. App.* 376.—China.

ABELIA:—
coriacea, *Hemsl. Diagn. Pl. Nov.* 53.—Mexic.
corymbosa, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 608.—As. temp.
Davidii, Hance, in Seem. Journ. Bot. vi. (1868) 329 = biflora.
floribunda, *Decne. in Fl. des Serres*, Sér. I. ii. (Janv. 1846).—Mexic.
Hanceana, Mart. ex Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 216 = chinensis.
hirsuta, Walp. Rep. vi. 3 = speciosa.
rupestris, Lindl. in Jonrn. Hort. Soc. i. (1846) 63 = chinensis.
serrata, Sieb. & Zucc. Fl. Japon. i. 76. t. 34 = uniflora.
spathulata, *Sieb. & Zucc. l. c.* 77. t. 34.—Japon.
speciosa, *Decne. in Fl. des Serres*, ii. (1846) snb t. 5.—Mexic.
splendens, Hort. ex C. Koch, Dendrol. ii. I. 20 = Lonicera caprifolioides.
triflora, *R. Br. in Wall. Pl. As. Rar.* i. 14. t. 15.—Reg. Himal.
uniflora, *R. Br. in Wall. l. c.*—China.

ABELICEA, Reichb. Consp. 84 (1828); Baill. Hist. Pl. vi. 185 (1875) = **Zelkova**, Spach (Urtic.).
aquatica, Rafin. New Fl. Am. iii. 41 = Planera aqnatica.
crenata, Baill. Hist. Pl. vi. 181 = Z. crenata.
cretica, Rafin. New Fl. Am. iii. 41 = Z. cretica.
Davidii, Buek, Ind. Gen. DC. Prod. iv. 1 = Hemiptelea Davidii.
sibirica, Rafin. New Fl. Am. iii. 41 = Z. crenata.

ABELMOSCHUS, Medic. Malv. 45 (1787) = **Hibiscus**, Linn. (Malvac.).
achanioides, Trncz. in Bnll. Soc. Nat. Mosc. xxxi. (1858) I. 196 = H. achanioides.
aculeatus, Walp. Rep. i. 308 = H. surattensis, cannabinus.
acuminatus, C. Muell. in Walp. Ann. vi. 407.—Java.
albo-rubens, Benth. Fl. Anstral. i. 209 = seq.
albo-ruber, F. Mnell. Fragm. i. 67 = H. ficulneus.
angulosus, Wight & Arn. Prod. 53 = H. angulosus.
angustifolius, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 152 = H. angustifolius.
aquaticus, Walp. Rep. i. 311 = H. roseus.
ascendens, Walp. l. c. 310 = H. physaloides.
Bammia, Webb, Fragm. Fl. Aethiop. 48 = H. esculentus.
betulifolia, Wall. Cat. p. 87 et n. 1918 = H. Abelmoschus.
cancellatus, Wall. l. c. p. 87 et n. 1920 = H. cancellatus.
chinensis, Wall. l. c. p. 87 et n. 1916 = H. Abelmoschus.
ciliaris, Walp. Rep. ii. 308 = H. ciliaris.
cisplatinus, Walp. l. c. 309 = H. cisplatinus.
congener, Walp. l. c. 308 = H. cannabinus.
crinitus, Wall. Pl. As. Rar. i. 39 = H. cancellatus.
cruentus, Walp. Rep. i. 310 = H. cruentus.
cryptocarpus, Walp. l. c. = H. Abelmoschus.
cubensis, Walp. l. c. 309 = H. Abelmoschus.
cucurbitaceus, Walp. l. c. 310 = H. Abelmoschus.
decandrus, Medic. Malv. 46 = Pavonia praemorsa.
divaricatus, Walp. Rep. i. 309 = H. divaricatus.
esculentus, Moench, Meth. 617 = H. esculentus.

ABELMOSCHUS :—
ficulneoides, Walp. Rep. i. 309 = H. ficulneoides.
ficulneus, Wight. & Arn. Prod. i. 53 = H. ficulneus.
fusiformis, Wall. Cat. p. 87 et n. 1921 = H. cancellatus.
Genevii, Walp. Rep. i. 310 = H. liliiflorus.
guineensis, Walp. l. c. = H. tiliaceus.
Haenkeanus, Presl, Rel. Haenk. ii. 134.—Ins. Luzon.
laxiflorus, Walp. Rep. i. 310 = H. laxiflorus.
longifolius, Kostel. Allg. Med.-Pharm. Flora, v. 1859 = H. Abelmoschus.
maculatus, Bartl. in Linnaea, xii. (1838) Litt. 81 = H. Bartlingianus.
magnificus, Wall. Cat. p. 87 et n. 1919.—Ind. or.
Manihot, Medic. Malv. 46 = H. Manihot.
marianus, Presl, Rel. Haenk. ii. 134.—Ins. Marian.
moschatus, Medic. Malv. 46 = H. Abelmoschus.
multiformis, Wall. Cat. p. 87 et n. 1917.—Ind. or.
officinalis, Endl. ex Heynh. Nom. ii. 1 = H. esculentus.
palustris, Walp. Rep. i. 311 = H. Moscheutos.
panduraeformis, Hassk. in Tijdschr. Nederl. Ind. i. (1843) 134 = H. panduraeformis.
pentaphyllus, Voigt, Hort. Suburb. Calc. 119 = H. pentaphyllus.
persicifolius, Walp. Rep. i. 308 = H. persicifolius.
Pseudo-abelmoschus, Blume, Bijdr. 70 = H. Abelmoschus.
pungens, Wall. Cat. p. 87 et n. 1924 = H. pungens.
quinquelobus, Walp. Rep. i. 310 = H. quinquelobus.
racemosus, Wall. Cat. p. 87 et n. 1929 = H. cancellatus.
rhodopetalus, F. Muell. Fragm. ii. 113 = H. rhodopetalus.
roseus, Walp. Rep. i. 311 = H. Moscheutos.
rostellatus, Walp. l. c. 308 = H. rostellatus.
rugosus, Wight & Arn. Prod. i. 43 = H. rugosus.
splendens, Walp. Rep. i. 309 = H. splendens.
strictus, Voigt, Hort. Suburb. Calc. 119 = H. ficulneus.
sublobatus, Presl, Rel. Haenk. ii. 134.—Ins. Pacif.
tetraphyllus, Wall. Cat. p. 87 et n. 1925 = H. tetraphyllus.
theiodes, Voigt, Hort. Suburb. Calc. 120 = H. sulphureus, *Voigt*.
tulipaeflorus, Walp. Rep. v. 92 = H. tulipaeflorus.
venustus, Walp. l. c. i. 309 = H. mutabilis.
verrucosus, Walp. l. c. 308 = H. cannabinus.
Vriesianus, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 134; xii. (1845) 88.—Malaya.
warreensis, Dalz. in Hook. Kew Journ. iii. (1851) 123 = H. tetraphyllus.
Zollingeri, C. Muell. in Walp. Ann. vi. 407.—Java.

ABENA, Neck. Elem. i. 296 (1790) = **Stachytarpheta**, Vahl (Verbenac.).

ABEREMOA, Aubl. Pl. Gui. i. 610, t. 245 (1775) = **Duguetia**, St. Hil. (Anonac.).
furfuracea, Baill. Hist. Pl. i. 204.—Bras.
guianensis, Aubl. Pl. Gui. i. 610.—Guiana.
lanceolata, Warm. in Kjoeb. Vidensk. Meddel. (1855) 146; (1873) 146.—Bras.
longifolia, Baill. Hist. Pl. i. 205 = D. longifolia.

ABERIA, Hochst. in Flora, xxvii. (1844) Bes. Beil. 2.
BIXINEAE, Benth. & Hook. f. i. 128.
ATELESTE, Sond. in Linnaea, xxiii. (1850) 9, 10.
abyssinica, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 236.—Afr. trop.
caffra, *Harv. & Sond. Fl. Cap.* ii. *Add.* 584.—Afr. austr.
edulis, T. Anders. in Journ. Linn. Soc. vii. (1862) 67 = caffra.
Gardnerii, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 235.—Zeylan.
longispina, *Harv. in Harv. & Sond. Fl. Cap.* ii. 585.—Afr. austr.
macrocalyx, *Oliver, Fl. Trop. Afr.* i. 122.—Afr. trop.
mollis, *Oliver, l. c.*—Afr. trop.
tristis, *Sond. in Linnaea*, xxiii. (1850) 9.—Afr. austr.
verrucosa, *Hochst. in Flora*, xxvii. (1844) *Bes. Beil.* 2.—Abyss.
Zeyheri, *Sond. in Linnaea*, xxiii. (1850) 10.—Afr. austr.

ABESINA, Neck. Elem. i. 33 (1790) = **Verbesina**, Linn. (Compos.).

ABIES, D. Don, in Loud. Arb. Brit. iv. 2329 (1838) = **Picea**, Link (Conif.).

ABIES, [Tourn.] Linn. Fl. Lapp. 277 (1737); Link, in Linnaea, xv. (1841) 525. *CONIFERAE*, Benth. & Hook. f. iii. 441.
PICEA, D. Don, in Loud. Arb. Brit. iv. 2293 (1838).
ajanensis, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 212 = Picea ajanensis.

ABIES :—
alba, Michx. Fl. Bor. Am. ii. 207 = Picea alba.
alba, Mill. Gard. Dict. ed. VIII. n. 1 (1768) = pectinata (1805).
Albertiana, A. Murr. in Proc. Hort. Soc. iii. (1863) 149 = Tsuga Mertensiana.
Alcockiana, Lindl. ex C. Koch, Dendrol. ii. II. 245 = Picea Alcockiana.
Alcoqueana, Gord. Pinet. Supp. 8 ('desc. falsa') = praec.
amabilis, *Forb. Pin. Wob.* 125. t. 44.—Am. bor. occ.
amabilis, Parl. in DC. Prod. xvi. II. 426, in parte = subalpina.
americana, Mill. Gard. Dict. ed. VIII. n. 6 = Tsuga canadensis.
Apollinis, Link, in Linnaea, xv. (1841) 528 = cephalonica.
Araragi, Siebold, in Verh. Batav. Gen. xii. (1830) 12 = Tsuga Sieboldii.
araucana, Poir. Encyc. Suppl. vi. 524 = Araucaria excelsa.
arctica, A. Cunn. ex Gord. Pinet. 11 = Picea rubra.
arctica, A. Murr. in Journ. Bot. v. (1867) 253 = Picea alba.
argentea, De Chambr. Traité Prat. Arb. Résin. 17 = pectinata.
aromatica, Rafin. Atl. Journ. 119 = balsamea ?
atlantica, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 214 = Cedrus atlantica.
baborensis, Coss. ex Parl. in DC. Prod. xvi. II. 423 = numidica.
baldensis, Zuccagn. ex Nym. Consp. 674 = pectinata.
balsamea, *Mill. Gard. Dict.* ed. VIII. n. 3.—Am. bor.
balsamifera, Michx. Fl. Bor. Am. ii. 207 = balsamea.
batavorum, Siebold.—Cf. Carr. Conif. ed. II. 229 = Cunninghamia sinensis.
bicolor, Maxim. in Bull. Acad. Pétersb. x. (1866) 488 = Picea Alcockiana.
bifida, Sieb. & Zucc. Fl. Jap. ii. 18 = firma.
bifolia, A. Murr. Syn. Var. Conif. 29 = lasiocarpa.
brachyphylla, *Maxim. in Bull. Acad. Pétersb.* x. (1866) 488.—Japon.
bracteata, *Hook. & Arn. Bot. Beech. Voy.* 394.—Calif.
Bridgesii, Kellogg, in Proc. Calif. Acad. ii. (1863) 8 = Tsuga Mertensiana.
Brunoniana, Lindl. in Penny Cyclop. i. 30 = T. Brunoniana.
californica, Hort. ex Steud. Nom. ed. II. i. 1 = Pseudotsuga Douglasii.
campylocarpa, A. Murr. in Trans. Bot. Soc. Edinb. vi. (1860) 370 = magnifica.
canadensis, Michx. Fl. Bor. Am. ii. 206 = Tsuga canadensis.
canadensis, Mill. Gard. Dict. ed. VIII. n. 4 = Picea alba.
candicans, Fisch. ex Endl. Syn. Conif. 96 = pectinata.
cedroides, Griff. ex Carr. Conif. ed. I. 188 = Tsuga Brunoniana.
Cedrus, Poir. Encyc. vi. 510 = Cedrus Libani.
cephalonica, *Link, in Linnaea*, xv. (1841) 529.—Graecia.
chiloënsis, Hort. ex Carr. Conif. ed. I. 222 = A. Pindrow.
chilrowensis, Hort. ex Parl. in DC. Prod. xvi. II. 425 = Webbiana.
chlorocarpa, Purk. ex Nym. Consp. 674 = Picea excelsa.
cilicica, *Ant. & Kotschy, in Oestr. Wochenschr.* (1853) 409.—As. Min.
cilicica, *Carr. Conif.* ed. I. 229.—As. Min.
cinerea, Borkh. Forstb. i. 398 = Picea excelsa.
Clanbrassiliana, Hort. ex Loud. Encyc. Pl. 804 = P. excelsa.
coerulea, Lodd. ex Forbes, Pin. Wob. 99 = P. alba.
coerulescens, Hort. ex C. Koch, Dendrol. ii. II. 242 = P. rubra.
Columbaria, Desf. Tabl. ed. II. 247 = Araucaria imbricata.
commutata, Gord. Pinet. ed. II. 5 = Picea Engelmanni.
concolor, *Lindl. & Gord. in Journ. Hort. Soc.* v. (1850) 210.—Am. bor. occ.
cupressoides, Poir. Encyc. vi. 524 = Libocedrus tetragona.
curvifolia, Salisb. in Trans. Linn. Soc. viii. (1807) 315 = Picea alba.
Dammara, Poir. Encyc. Suppl. v. 35 = Agathis loranthifolia.
Davidiana, *Franch. in Nouv. Mém. Mus. Par.* vi. (1884) 96.—As. or.
decidua, Wall.—Cf. C. Koch, Dendrol. ii. II. 253 = Tsuga Brunoniana.
densa, Griff. Notul. iv. 19.—Cf. Webbiana.
denticulata, Poir. Encyc. vi. 520 = Picea nigra.
Deodara, Lindl. in Penny Cyclop. i. 34 = Cedrus Deodara.

ABIES :—
diversifolia, Maxim. in Bull. Acad. Pétersb. xii. (1868) 229 = Tsuga diversifolia.
Douglasii, Lindl. in Penny Cyclop. i. 32 = Pseudotsuga Douglasii.
Drummondi, Hort. ex Gord. Pinet. ed. I. 16 = praec.
dumosa, Loud. Arb. Brit. iv. 2325 = Tsuga Brunoniana.
Eichleri, Lauche, in Berl. Gartenz. i. (1882) 63 = Veitchii.
elegans, Sm. ex Knight, Syn. Conif. 36 = Picea excelsa.
Engelmanni, Parry, in Trans. Acad. St. Louis, ii. (1868) 212 = Picea Engelmanni.
eremita, Hort. Gall. ex C. Koch, Dendrol. ii. II. 237 = Picea excelsa.
erythrocarpa, Purk. ex Nym. Consp. 674 = P. excelsa.
excelsa, Link, in Abh. Acad. Berl. (1827) 182 = pectinata.
excelsa, Poir. Encyc. vi. 518 = P. excelsa.
extrema, T. Fries, in Bot. Notiser (1867), 56 = P. excelsa.
falcata, *Rafin. Atl. Journ.* 120.—Am. bor.
finedonensis, Hort. ex Gord. Pinet. Suppl. 4 = P. excelsa.
Finhonnoskiana, R. Neum. ex Parl. in DC. Prod. xvi. II. 431 = homolepis.
firma, *Sieb. & Zucc. Fl. Jap.* ii. 15. t. 107.—Japon.
Fischeri, *Lindl. ex Ledeb. Fl. Ross.* ii. 172.—Rossia.
Fortunei, A. Murr. Pin. Japan. 49 = Keteleeria Fortunei.
Fraseri, *Lindl. in Penny Cyclop.* i. 30.—Am. bor.
gigantea, Sm. ex Carr. Conif. ed. II. 327 = Picea excelsa.
glauca, Roezl, ex Gord. Pinet. 149 = religiosa.
glaucescens, Roezl, Cat. Gr. Conif. Mex. (1858–59) 5 = religiosa.
Glehni, F. Schmidt, Reisen Amurl. 176 = Picea Glehnii.
Gmelini, Rupr. in Beitr. Pfl. Russ. Reich. ii. 56 = Larix dahurica.
Gordoniana, Carr. Conif. ed. II. 298 = grandis.
grandis, *Lindl. in Penny Cyclop.* i. 30.—Am. bor. occ.
Gregoryana, Low, ex Gord. Pinet. Suppl. 4 = P. excelsa.
Griffithiana, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 214 = Larix Griffithii.
Hanburyana, Hort. ex Gord. Pinet. ed. II. 32, 421 = Tsuga Sieboldii ?
Harryana, McNab, in Proc. Roy. Irish Acad. (1876) 689 = homolepis.
heterophylla, C. Koch, in Linnaea, xxii. (1849) 295 = sibirica.
heterophylla, Rafin. Atl. Journ. 119 = Tsuga Mertensiana.
hirtella, Lindl. in Penny Cyclop. i. 31 = religiosa.
hispanica, De Chamb. Trait. Prat. Arb. Résin. 339 = A. Pinsapo.
holophylla, *Maxim. in Bull. Acad. Pétersb.* x. (1866) 487.—As. temp.
homolepis, *Sieb. & Zucc. Fl. Jap.* ii. 17.—Japon.
Hookeriana, A. Murr. in Edinb. N. Phil. Journ. N. S. i. (Jan.-Apr. 1855). 289 = Tsuga Hookeriana.
Hudsonia, Bosc, ex Carr. Conif. ed. I. 200 = Fraseri.
humilis, La Pyl. in Mém. Soc. Linn. Par. iv. (1826) 437 = Fraseri.
inverta, R. Sm. ex Gord. Pinet. Suppl. 4 = P. excelsa.
jezoënsis, Lindl. & Paxt. Flow. Gard. i. (1850–51) 42. f. 26 = Keteleeria Fortunei.
jezoënsis, Sieb. & Zucc. Fl. Jap. ii. 19. t. 110 = Picea Alcockiana.
Kaempferi, Lindl. in Penny Cyclop. i. 34 = Pseudolarix Kaempferi.
kamtschatica, Rupr. in Beitr. Pfl. Russ. Reich. ii. 57 = Larix dahurica.
Khutrow, Loud. Encyc. Trees, 1032 = Picea Morinda.
Kotschyana, *Fenzl, ex Tchihat. Asie Min. Bot.* ii. 495.—As. Min.
lanceolata, Poir. Encyc. vi. 523 = Cunninghamia sinensis.
Larix, Poir. Encyc. Suppl. vi. 511 = Larix europaea.
lasiocarpa, *Hook. Fl. Bor. Am.* ii. 163.—Am. bor. occ.
lasiocarpa, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 210 = grandis, Lowiana.
laxa, C. Koch, Dendrol. ii. II. 243 = Picea alba.
Ledebourii, Rupr. in Beitr. Pfl. Russ. Reich. ii. 56 = Larix dahurica.
leioclada, Stev. ex Gord. Pinet. 152 = Nordmanniana.
Lemoniana, Booth, ex Gord. l. c. ed. I. 5 = Picea excelsa.
leptolepis, Sieb. & Zucc. Fl. Japon. 12. t. 103 = Larix japonica.
Lindleyana, Roezl, ex Carr. Conif. ed. II. 275 = religiosa.
Lowiana, *A. Murr. Syn. Var. Conif.* 27.—Am. bor. occ.

ABIES:—
Luscombeana, Loud. Arb. Brit. iv. 2325 = cephalonica.
magnifica, *A. Murr. in Proc. Roy. Hort. Soc.* iii. (1863) 318.—Calif.
Mariana, Mill. Gard. Dict. ed. VIII. n. 5 = Picea nigra.
Mariesii, *Mast. in Gard. Chron.* xii. (1879) 789.—As. or.
Maximowiczii, R. Neum. Cat. (1865); ex Parl. in DC. Prod. xvi. II. 431 = Picea Maximowiczii.
medioxima, Lawson, Pin. Brit. 159 = P. excelsa.
Menziesii, Lindl. in Penny Cyclop. i. 32 = Picea Menziesii.
Merkiana, Fisch. ex Parl. in DC. Prod. xvi. II. 418 = Picea Menziesii.
Mertensiana, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 211 = Tsuga Mertensiana.
metensis, Hort. Par. ex Gord. Pinet. ed. II. 211 = pectinata.
microcarpa, Lindl. in Penny Cyclop. i. 33 = Larix microcarpa.
microphylla, *Rafin. Atl. Journ.* (1820) 119.—Am. bor.
microsperma, Lindl. in Gard. Chron. (1861) 22 = Picea ajanensis.
miniata, Knight, ex Gord. Pinet. ed. II. 8 = P. excelsa.
minor, Duham. ex Gord. I. c. ed. I. 143 = balsamea.
minor, Gilib. Exerc. Phytol. ii. 412 = pectinata.
minuta, Poir. Encyc. vi. 523 = P. rubra.
Momi, Siebold, in Verh. Batav. Gen. xii. (1830) 12 = firma.
montana, Nym. Syll. Suppl. 602 = P. excelsa.
Morinda, Loud. Encyc. Pl. 1294 = P. Morinda.
mucronata, Rafin. Atl. Journ. 120 = Pseudotsuga Douglasii.
mucronata, Rauch, ex Carr. Conif. ed. II. 329 = Picea excelsa.
nephrolepis, Maxim. in Bull. Acad. Pétersb. x. (1866) 486 = Veitchii.
nigra, Desf. Hist. Arbr. et Arb. ii. 580 = Picea nigra.
nigra, Duroi, Harb. Baumz. ed. Pott, ii. 182 = P. nigra.
nigra, Engelm. in Am. Journ. Sc. xxxiii. (1838) 330 = Picea Engelmanni.
nobilis, *Lindl. in Penny Cyclop.* i. 30.—Am. bor. occ.
Nordmanniana, *Spach, Hist. Vég. Phan.* xi. 418.—Reg. Caucas.
Novae-Angliae, C. Koch, Dendrol. ii. II. 240 = Picea nigra.
numidica, *De Lannoy, ex Carr. in Rev. Hortic.* (1866) 106.—Afr. bor.
obliqua, Bong. ex Gord. Pinet. Supp. 10 = Pseudotsuga Douglasii.
obliquata, Rafin. ex Gord. l. c. = praec.
obovata, Loud. Arb. Brit. iv. 2329 = Picea obovata.
Omorika, Nym. Consp. 673 = Picea Omorika.
orientalis, Poir. Encyc. vi. 518 = Picea orientalis.
panachaica, Heldr. ex Parl. in DC. Prod. xvi. II. 422 = cephalonica.
Parsonsiana, Hort. ex Mast. in Gard. Chron. (1880) I. 648 = Lowiana.
parvula, Knight, Syn. Conif. 36 = Picea excelsa.
Pattoni, Jeffrey, ex Gord. Pinet. Suppl. 6 = seq.
Pattoniana, Jeffrey, ex A. Murr. in Bot. Exped. Oregon. Circ. = Tsuga Hookeriana.
pectinata, *DC. Fl. Fr.* iii. 276.—Europ.
pectinata, Gilib. Exerc. Phytol. ii. 411 = P. excelsa.
pectinata, Poir. Encyc. vi. 523 = Picea rubra.
peloponnesiaca, Haage, ex Parl. in DC. Prod. xvi. II. 422 = cephalonica.
pendula, Griff. ex Gord. Pinet. Suppl. 7 = Picea Morinda.
pendula, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 213 = Larix microcarpa.
Picea, Lindl. in Penny Cyclop. i. 29 = pectinata.
Picea, Mill. Gard. Dict. ed. VIII. n. 2 = Picea excelsa.
Pichta, Forbes, Pin. Wob. 113. t. 39 = sibirica.
Pindrow, *Spach, Hist. Vég. Phan.* xi. 423.—Reg. Himal.
Pinsapo, *Boiss. in Bibl. Univ. Genève* (1838); *ex ej. Voy. Bot. Espagne*, ii. 584.—Hispan.
polita, Sieb. & Zucc. Fl. Jap. ii. 20. t. 111 = Picea polita.
Reginae-Amaliae, Heldr. in Regel, Gartenfl. ix. (1860) 313 = cephalonica.
religiosa, *Lindl. in Penny Cyclop.* i. 31.—Mexic.
Rinzi, Hort. Par. ex Gord. Pinet. Suppl. 50 = cilicica.
Romania, Loud. ex Steud. Nom. ed. II. i. 1 = Pinus Laricio.
rubra, Poir. Encyc. vi. 520 = P. rubra.
sachalinensis, *Mast. in Gard. Chron.* (1879) II. 588.—As. or.
sacra, *Franch. in Nouv. Arch. Mus. Par.* vii. (1882) 290.—As. or.
Schrenkiana, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 212 = Picea obovata.

ABIES:—
selinusia, Carr. in Fl. des Serres (1859), 69 = cilicica.
sibirica, *Ledeb. Fl. Alt.* iv. 202.—Sibir.
sitchensis, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 212 = Picea Menziesii.
Smithiana, Lindl. in Penny Cyclop. i. 31 = P. Morinda.
spectabilis, Spach, Hist. Vég. Phan. xi. 422 = Webbiana.
spinulosa, Griff. Itin. 259, 265, 275 = Picea Morinda.
Standishiana, C. Koch, Dendrol. ii. II. 257 = Tsuga Douglasii.
subalpina, Engelm. ex L. Ward, in Am. Natural. x. (1876) 555 = lasiocarpa.
subarctica, Nym. Consp. 674 = Picea excelsa.
sumatrana, Desf. Hort. Par. ed. III. 356 = Agathis loranthifolia.
taxifolia, Desf. l. c. = pectinata.
taxifolia, Drum. ex Gord. Pinet. 16 = Pseudotsuga Douglasii.
taxifolia, Jeffr. ex Gord. l. c. 18 = Tsuga Mertensiana.
taxifolia, Presl, Epim. Bot. 237 = Pseudotsuga Douglasii.
taxifolia, Rafin. New Fl. Am. i. 38.—Am. bor.
Tchugatskoi, Lawson, ex Gord. Pinet. Suppl. 50 = cilicica.
Thunbergii, Lindl. in Penny Cyclop. i. 34 = Picea polita.
Tlapalcatuda, Roezl, Cat. Gr. Conif. Mex. (1858–59) 5 = religiosa.
Torano, Siebold, in Verh. Bat. Gen. xii. (1830) 12 = Picea polita.
trigona, Rafin. Atl. Journ. (1820) 119 = Sequoia sempervirens?
Tschonoskiana, Regel, in Ind. Sem. Hort. Petrop. (1865) 32 = homolepis.
Tsuga, Sieb. & Zucc. Fl. Jap. ii. 14. t. 106 = Tsuga Sieboldii.
Veitchii, *Lindl. in Gard. Chron.* (1861) 23.—Japon.
venusta, C. Koch, Dendrol. ii. II. 210 = bracteata.
viminalis, Wahlenb. Fl. Suec. 630 = Picea excelsa.
virescens, Hinterh. ex Nym. Consp. 673 = P. alba.
vulgaris, Poir. Encyc. vi. 514 = pectinata.
Webbiana, *Lindl. in Penny Cyclop.* i. 30.—Reg. Himal.
Williamsonii, Bridges, ex Gord. Pinet. Suppl. 12 = Tsuga Mertensiana.
Williamsonii, Newberry, in Williamson, in Pacif. Rail. Rep. vi. 53. t. 7 = Tsuga Hookeriana.

ABIGA, St. Lag. in Ann. Soc. Bot. Lyon, vii. 85 (1880) = **Ajuga**, Linn. (Labiat.).
cistifolia, St. Lag. l. c. 85, 119 = Ajuga Iva.

ABILDGAARDIA, Vahl, Enum. ii. 296 (1806) = **Fimbristylis**, Vahl (Cyperac.).
aphylla, Kunth, Enum. ii. 248 = F. bahiensis.
Baeothryon, St. Hil. Voy. ii. 389 = F. bahiensis.
barbata, Beauv. Fl. Oware, ii. 47. t. 86. f. 1.—Afr. trop.
brevifolia, Steud. Syn. Pl. Cyp. 72 = F. setacea.
cinnamometorum, Thw. Enum. Pl. Zeyl. 347 = F. cyperoides.
compressa, J. & C. Presl, Rel. Haenk. i. 179 = F. monostachya.
cyperoides, Nees & Meyen, in Wight, Contrib. 95 = F. cyperoides.
Eragrostis, Nees & Meyen, l. c. = F. Eragrostis.
fimbristyloides, F. Muell. Fragm. viii. 273 = F. disticha.
fulvescens, Thw. Enum. Pl. Zeyl. 347 = F. fulvescens.
fusca, F. Muell. Fragm. viii. 273 = F. cyperoides.
fusca, Nees, in Wight, Contrib. 95.—Reg. Himal.
geminata, Schrad. ex Nees, in Mart. Fl. Bras. ii. I. 71 = F. bahiensis.
indica, Nees, in Linnaea, ix. (1835) 289 = F. monostachya.
javana, Nees, l. c. = F. monostachya.
javanica, Steud. Syn. Pl. Cyp. 72 = F. juncea.
laevigata, Willd. ex Steud. Nom. ed. II. i. 1 = F. monostachya.
lanceolata, Schum. Guin. Pl. 33 = F. tristachya.
monostachya, Vahl, Enum. ii. 296 = F. monostachya.
nervosa, J. & C. Presl, Rel. Haenk. i. 180 = F. schoenoides.
pauciflora, Kunth, Enum. Pl. ii. 249.—Ins. Molucc.
pilosa, Nees, in Linnaea, ix. (1835) 289.—Afr. trop.
polycephala, Brongn. in Duperr. Voy. Bot. 176. t. 32 = Androtrichum polycephalum.
polygama, Spreng. Nov. Prov. 45.—Bras.
polystachya, Spreng. Fl. Hal. Mant. 29 = Pharus angustifolius.
pubescens, J. & C. Presl, Reliq. Haenk. 180 = F. exilis.
Rottboelliana, Nees, ex Hook. & Arn. Bot. Beech. Voy. 272 = F. monostachya.
schoenoides, R. Br. Prod. 229 = F. squarrulosa.

ABILDGAARDIA:—
scirpoides, Nees, in Linnaea, ix. (1835) 289 = F. bahiensis.
setacea, Griseb. Cat. Pl. Cub. 238.—Cuba.
ternata, Schrad. ex Nees, in Mart. Fl. Bras. ii. I. 71 = F. bahiensis.
tristachya, Vahl, Enum. ii. 297 = F. tristachya.
vaginata, R. Br. Prod. i. 229 = F. Brownii.

ABILDGARDIA, Reichb. Consp. 56 (1828) = praec.

ABILGAARDIA, Poir. in Dict. Sc. Nat. i. Suppl. 3 (1816) = praec.

ABLANIA, Aubl. Pl. Gui. i. 585. t. 234 (1775) = **Sloanea**, Linn. (Tiliac.).
digitata, Spreng. Syst. iv. Cur. Post. 210.—Bras.
guianensis, Aubl. Pl. Gui. i. 585. t. 234 = S. guianensis.
laurifolia, Pers. Syn. ii. 81 = S. guianensis.

ABOBRA, Naud. in Rev. Hortic. (1862) 111. *CUCURBITACEAE*, Benth. & Hook. f. i. 834.
tenuifolia, *Naud. l. c.*—Bras.
viridiflora, *Naud. l. c.*—Am. austr.

ABOLA, Adans. Fam. ii. 31 (1763) = **Cinna**, Linn. (Gramin.).

ABOLA, Lindl. Fol. Orch. (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 566.
radiata, *Lindl. l. c.*—N. Granat.

ABOLARIA, Neck. Elem. i. 105 (1790) = **Globularia**, Linn. (Selagin.).

ABOLBODA, Humb. & Bonpl. Pl. Aequin. ii. 109. t. 114 (1809). *XYRIDEAE*, Benth. & Hook. f. iii. 842.
CHLOERUM, Willd. ex Link, in Spreng. Jahrb. i. III. 74 (1833).
POARCHON, Mart. ex Seub. in Mart. Fl. Bras. iii. I. 223 (1855).
Aubletii, *Kunth, Enum. Pl.* iv. 28.—Guian.
brasiliensis, *Kunth, l. c.* 26.—Bras.
grandis, *Griseb. in Linnaea*, xxi. (1848) 281.—Bras.
imberbis, *H. B. & K. Nov. Gen. et Sp.* i. 256.—Guian.
inermis, Link, ex Steud. Nom. ed. II. i. 2 = imberbis.
Poarchon, *Seub. in Mart. Fl. Bras.* iii. I. 223.—Bras.
Poeppigii, *Kunth, Enum. Pl.* iv. 27.—Bras.
pulchella, *Humb. & Bonpl. Pl. Aequin.* ii. 110. t. 114.—Venezuela.

ABROCHIS, Neck. Elem. iii. 130 (1790) = **Orchis**, Tourn.

ABROMA, Jacq. Hort. Vind. iii. t. 1 (1776). *STERCULIACEAE*, Benth. & Hook. f. i. 224.
AMBROMA, Linn. f. Suppl. 341 (1781).
alata, *Blanco, Fl. Filip.* ed. I. 605.—Ins. Philipp.
angulata, Lam. Encyc. i. 126 = augusta.
angulosa, Poir. Encyc. Suppl. i. 322 = augusta.
augusta, *Linn. f. Suppl.* 341 (Ambroma).—As. trop.
communis, Blanco, Fl. Filip. ed. I. 606 = augusta.
denticulata, *Miq. Pl. Jungh.* 288.—Java.
elongata, Lam. Encyc. i. 127 = augusta.
fastuosa, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 409.—As. trop.; Austral.
fastuosa, Gaertn. Fruct. i. 307. t. 64 = augusta.
fastuosa, Jacq. Hort. Vind. iii. 3. t. 1 = augusta.
javanica, *Miq. Fl. Ind. Bat.* i. II. 183.—Java.
Mariae, *Mart. in Denkschr. Bayer. Ges.* iii. (1837–43) 297.—Java.
mollis, *DC. Prod.* i. 485.—Ins. Molucc.
nitida, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 73.—Chili.
obliqua, *Presl, Rel. Haenk.* ii. 143.—Ins. Philipp.
tomentosa, Steud. Nom. ed. II. i. 2, sphalm.—Cf. Ambora tomentosa.
Wheleri, Retz. Obs. v. 27 = augusta.

ABRONIA, Juss. Gen. 448 (1789). *NYCTAGINEAE*. Benth. & Hook. f. iii. 7.
CYCLOPTERA, Nutt. ex A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 319.
TRICRATUS, L'Hérit. ex Willd. Sp. Pl. i. 807 (1799).
arenaria, *Menzies, ex Hook. Exot. Fl.* t. 193.—Calif.
californica, Raeusch. Nom. ed. III. 191 = umbellata
Crux-Maltae, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 71. f. 16.—Calif.
Cycloptera, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 319.—Texas.
fragrans, *Nutt. ex Hook. Kew Journ.* v. (1853) 261.—Calif.
glauca, *Menzies, ex Hook. Exot. Fl.* t. 194.—Calif.
gracilis, *Benth. Bot. Voy. Sulph.* 44.—Calif.
latifolia, *Eschsch. in Mem. Acad. Petersb.* v. (1826) 281.—Calif.

ABRONIA:—
maritima, *Nutt. ex S. Wats. Bot. Calif.* ii. 4.—Calif.
mellifera, *Dougl. in Hook. Bot. Mag.* t. 2879.—Calif.
Menziesii, *Dougl. ex Hook. Fl. Bor. Am.* ii. 125.—Calif.
micrantha, Choisy, in DC. Prod. xiii. II. 436 = A. Cycloptera.
nana, *S. Wats. in Proc. Am. Acad.* xiv. (1870) 294.—Am. bor. occ.
parviflora, *H. B. & K. Nov. Gen. et Sp.* ii. 216. t. 128.—N. Granat.
rosea, Hartweg, ex Loud. Hort. Brit. Suppl. iii. 479 = umbellata?
rotundifolia, Gaertn. f. Fruct. iii. t. 214 = umbellata.
speciosa, *Buckl. in Proc. Acad. Sc. Philad.* (1863) 7.—Am. bor. occ.
turbinata, *Torr. ex S. Wats. in Bot. King's Exped.* 285. t. 31.—Am. bor. occ.
umbellata, *Lam. Illustr.* i. 469. t. 105.—Calif.
vespertina, Dougl. ex Hook. Fl. Bor. Am. ii. 125 = mellifera.
villosa, *S. Wats. in Am. Naturalist*, vii. (1873) 302.—Am. bor.

ABROPHAËS, Rafin. Sylva Tellur. 99 (1838) = **Fothergilla**, Linn. f. (Hamamel.).
mirabilis, Rafin. l. c. = F. alnifolia.

ABROPHYLLUM, Hook. f. in Benth. Fl. Austral. ii. 437 (1864). *SAXIFRAGEAE*, Benth. & Hook. f. i. 647.
BRACHYNEMA, F. Muell. Fragm. iii. 90. (1862).
ornans, *Hook. f. l. c.*—Austral.

ABROTANELLA, Cass. in Dict. Sc. Nat. xxxvi. 27 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 431.
CERATELLA, Hook. f. Fl. Antarct. i. 23. t. 18 (1844).
SCLEROLEIMA, Hook. f. in Lond. Journ. Bot. v. (1846) 444. t. 14.
TRINEURON, Hook. f. Fl. Antarct. i. 23. t. 17 (1844).
emarginata, *Cass. in Dict. Sc. Nat.* xxxvi. 27 (1825).—Ins. Falkl.
forsterioides, *Hook. f. Handb. N. Zeal. Fl.* 139.—N. Zel.; Austral.
inconspicua, *Hook. f. l. c.* 140.—N. Zel.
linearifolia, *A. Gray, in Proc. Am. Acad.* v. (1861) 137.—Fuegia.
linearis, *Bergg. in Minneskr. Fisiog. Sällsk. Lund* (1877), VIII. p. 14.—N. Zel.
nivigena, *F. Muell. in Benth. Fl. Austral.* iii. 554.—Austral.
pusilla, *Hook. f. Handb. N. Zeal. Fl.* 139.—N. Zel.
rhynchocarpa, *Balf. f. in Phil. Trans.* clxviii. (1879) 352.—Ins. Rodr.
rosulata, *Hook. f. Handb. N. Zeal. Fl.* 139.—N. Zel.
scapigera, *F. Muell. ex Benth. Fl. Austral.* iii. 554.—Austral.
spathulata, *Hook. f. Handb. N. Zeal. Fl.* 139.—N. Zel.
submarginata, *A. Gray, in Proc. Am. Acad.* v. (1861) 137.—Reg. Magell.

ABROTANUM, [Tourn.] Linn. Syst. ed. I (1735) = **Artemisia**, Tourn. (Compos.).
alpestre, ambiguum, brachylobium, congestum, incanescens, pauciflorum, pedunculare, platylobum, pulverulentum, rhodanicum, suave, virgatum, viridulum, xerophilum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 69–72 = Art. Abrotanum.
frutescens, Gilib. Fl. Lituan. i. 173 = Art. campestris.

ABRUS, Linn. Hort. Cliff. 488 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 527.
ZAGA, Rafin. Fl. Tellur. ii. 33 (1836).
acutifolius, Blume, ex Miq. Fl. Ind. Bat. i. 160 = pulchellus.
arboreus, Vell. Fl. Flum. 303; vii. t. 99 = Ormosia nitida.
canescens, *Welw. in Oliver, Fl. Trop. Afr.* ii. 175.—Afr. trop.
cantoniensis, *Hance, in Journ. Bot.* vi. (1868) 112.—China.
fruticulosus, *Wall. Cat.* n. 5820.—Ind. or.
laevigatus, E. Mey. Comm. i. 126 = pulchellus.
lusorius, *Vell. Fl. Flum.* 302; vii. t. 97ª.—Bras.
maculatus, Noronha, in Verh. Batav. Gen. v. (1790), Art. IV. 7 = precatorius.
melanospermus, Hassk. Cat. Pl. Hort. Bog. Alt. 282 = pulchellus.
minor, Desv. in Ann. Sc. Nat. Sér. I. xi. (1826) 418 = precatorius.
mollis, *Hance, in Journ. Bot.* ix. (1871) 430.—China.
pauciflorus, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 418 = precatorius.

ABRUS:—
precatorius, *Linn. Syst.* ed. XII. 472.—Reg. trop.
pulchellus, *Wall. Cat.* n. 5819.—Ind. or.; Afr. austr.
squamulosus, E. Mey. Comm. i. 126 = precatorius.
tenuiflorus, *Benth. in Mart. Fl. Bras.* xv. I. 216.—Bras.

ABRYANTHEMUM, Neck. Elem. ii. 82 (1790) = **Mesembryanthemum**, Linn. (Ficoid.).

ABSINTHIUM, Tourn. ex Linn. Syst. ed. I (1735) = **Artemisia**, Tourn. (Compos.).
alpinum, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 240 = Art. lanata.
arborescens, Moench, Meth. 579 = Art. arborescens.
argenteum, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 272 = Art. argentea.
bipedale, Gilib. Fl. Lituan. i. 174 = Art. Absinthium.
boreale, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 265 = Art. borealis.
campestre, Dulac, Fl. Hautes-Pyr. 502 = Art. campestris.
camphoratum, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 230 = Art. camphorata.
canariense, Bess. l. c. 229 = Art. canariensis.
candidum, Bieb. ex DC. Prod. vi. 125 = Art. frigida.
cernuum, Moench, Meth. 579 = Art. crithmifolia.
chinense, DC. Prod. vi. 118 sub n. 139 = Art. chinensis.
congestum, Lam. Fl. Fr. ii. 46 = Art. glacialis.
corymbosum, DC. Prod. vi. 122. sub n. 161 = Art. camphorata.
dentatum, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 265 = Art. rupestris.
divaricatum, Fisch. ex Bess. l. c. 263 = Art. anethifolia.
frigidum, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 251 = Art. frigida.
glaciale, Lam. Illustr. iii. 273. t. 695 = Art. glacialis.
grandiflorum, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 232 = Art. grandiflora.
grandiflorum, Roehl. ex Steud. Nom. ed. II. i. 2 = Art. rupestris.
lagocephalum, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 233 = Art. lagocephala.
lanatum, Bess. l. c. 265 = Art. lanata.
laxum, Lam. Fl. Fr. ii. 46 = Art. Mutellina.
mollissimum, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 265 = Art. mollissima.
Moxa, Bess. l. c. 228 = Art. Moxa.
Mutellina, Roehl. ex Steud. Nom. ed. II. i. 2 = Art. Mutellina.
nitens, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 235 = Art. sericea.
officinale, Brot. Fl. Lusit. i. 357 = Art. Absinthium.
pedunculare, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 63 = Art. splendens.
petrosum, Baumg. Enum. Stirp. Transs. iii. 90 = Art. Baumgarteni.
ponticum, Bess. Enum. Pl. Volh. 76 = Art. pontica.
rupestre, Schrank, Prim. Fl. Salisb. 203 = Art. rupestris.
santonicum, Gueldenst. ex Ledeb. Fl. Ross. ii. 573 = Art. maritima.
saxatile, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 231 = Art. camphorata.
sericeum, Bess. l. c. 237 = Art. sericea.
Sieversianum, Bess. l. c. 259 = Art. Sieversiana.
spathulatum, Noronha, in Verh. Batav. Gen. v. (1790) Art. IV. 6 = Dichrocephala latifolia.
spicatum, Baumg. Enum. Stirp. Transs. iii. 90 = Art. spicata.
splendens, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 238 = Art. splendens.
tanacetifolium, Gaertn. Fruct. ii. 393 = Art. tanacetifolia.
trifidum, Schrank, Baier. Fl. ii. 362 = Art. austriaca.
virescens, Juss. ex Bess. in Bull. Soc. Nat. Mosc. viii. (1835) 29 = Art. Jussieana.
virgatum, Turcz. ex Ledeb. Fl. Ross. ii. 597 = Art. rupestris.
viride, Bess. in Bull. Soc. Nat. Mosc. i. (1829) 249 = Art. rupestris.
viridiflorum, Bess. l. c. 246 = Art. rupestris.
vulgare, Dulac, Fl. Hautes-Pyr. 502 = Art. vulgaris.
vulgare, Lam. Fl. Fr. ii. 45 = Art. Absinthium.

ABSYNTHIUM, Gaertn. Meyer & Scherb. Fl. Wett. iii. 196 (1801) = praec.

ABULFALI, Adans. Fam. ii. 190 (1763) = **Thymbra**, Linn. (Labiat.).

ABUMON, Adans. Fam. ii. 54 (1763) = **Agapanthus**, L'Hérit. (Liliac.).

ABUTA, [Barr.] Aubl. Pl. Gui. i. 618. t. 250 (1775). *MENISPERMACEAE*, Benth. & Hook. f. i. 35.
ANELASMA, Miers, in Ann. Nat. Hist. Ser. II. vii. (1851) 42.
BATSCHIA, Mutis, ex Thunb. in Nov. Act. Soc. Sc. Upsal. v. (1792) 120. t. 2.
TRICHOA, Pers. Syn. ii. 634 (1807).
acutifolia, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 259.—Bras.
amara, Aubl. Pl. Gui. i. 620 = Aristolochia glaucescens.
barbata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 258.—Guiana.
candicans, *Rich. ex DC. Syst.* i. 543.—Guiana.
Candollei, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 47.—Guiana.
concolor, Benth. in Journ. Linn. Soc. v. (1861) App. 49 = guianensis.
concolor, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 64. t. 188.—Am. austr.
Grisebachii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 47.—Bras.
guianensis, *Eichl. in Flora*, xlvii. (1864) 390.—Guiana.
heterophylla, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 258.—Am. austr.
Imene, *Eichl. in Flora*, xlvii. (1864) 389.—Bras.
macrophylla, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 258.—Am. austr.
oblonga, *Miers, l. c.*—Am. austr.
oblongifolia, *Miers, l. c.*—Am. austr.
panurensis, *Eichl. in Flora*, xlvii. (1864) 390.—Bras.
platyphylla, Mart. ex Eichl. in Mart. Fl. Bras. xiii. I. 200 (= *Botryopsis platyphylla*).—Bras.
racemosa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 48.—N. Granat.
rufescens, *Aubl. Pl. Gui.* i. 618. t. 250.—Guiana.
rufescens, DC. Syst. i. 542 = Candollei.
rufescens, Griseb. in Journ. Linn. Soc. iii. (1859) 109 = Grisebachii.
scandens, DC. Syst. i. 542 in syn. = rufescens.
Seemanni, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 50.—Darien.
Selloana, *Eichl. in Flora*, xlvii. (1864) 389.—Bras.
spicata, *Triana & Planch. in Ann. Sc. Nat.* Sér. II. xvii. (1682) 119.—N. Granat.
tomentosa, Sagot, ex Benth. in Journ. Linn. Soc. v. (1861) Suppl. ii. 51 (Coccolobae sp.).—Guian.
umbellata, Sagot, ex Benth. l. c. = Sciadotaenia cayennensis.

ABUTILAEA, F. Muell. in Linnaea, xxv. (1852) 379 = **Abutilon**, Tourn. (Malvac.).
cryptantha, F. Muell. l. c. = Abutilon Mitchelli.

ABUTILON, Tourn. ex Adans. Fam. ii. 398 (1763). *MALVACEAE*, Benth. & Hook. f. i. 204.
ABUTILAEA, F. Muell. in Linnaea, xxv. (1852) 379.
BELOËRE, Shuttlew. ex Benth. & Hook. f. Gen. i. (1862) 204.
BOGENHARDEA, Reichb. Nom. 200 (1841).
HERISSANTIA, Medic. Phil. Bot. i. 90 (1789).
LASS, Adans. Fam. ii. 400 (1763).
PRESTONIA, Scop. Introd. 281 (1777).
acerifolium, G. Don, Gen. Syst. i. 504 = Sida acerifolia.
affine, *G. Don, l. c.* 503.—Reg. Argent.
albescens, Miq. Pl. Jungh. 285 = crispum.
albidum, Sweet, Hort. Brit. ed. I. 54.—Ins. Canar.
albidum, *Walp. Rep.* i. 325.—Mexic.
albidum, Webb & Berth. Phyt. Canar. i. 39. t. 2 = fruticosum.
ambiguum, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 205 = Sida Lindeniana.
americanum, Sweet, Hort. Brit. ed. I. 53 = indicum.
amplexifolium, *G. Don, Gen. Syst.* i. 502.—Mexic.
amplum, *Benth. Fl. Austral.* i. 200.—Austral.
Anderssonianum, *Garcke, ex Anderss. Galap. Oar. Vegetat.* 230.—Ins. Galap.
Andrieuxii, *Hemsl. Diagn. Pl. Nov.* 24.—Mexic.
angulatum, Mast. in Oliver, Fl. Trop. Afr. i. 183 = intermedium.
anodoides, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 49.—Bras.
arborescens, Medic. Malv. 29 = indicum.
arboreum, *Sweet, Hort. Brit.* ed. I. 53.—Peruv.
Arnottianum, Walp. Rep. i. 324 = Sida Arnottiana.
asiaticum, *G. Don, Gen. Syst.* i. 503.—Reg. trop.
asiaticum, Guill. & Perr. Fl. Seneg. Tent. i. 67 = glaucum.
asiaticum, Sweet, Hort. Brit. ed. I. 53 = indicum.
atropurpureum, *G. Don, Gen. Syst.* i. 502.—Java.

ABUTILON:—
aurantiacum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 357.—Calif.
aureum, G. Don, in Sweet, Hort. Brit. ed. III. 80 = Sida aurea.
auritum, *Sweet, l. c.* ed. I. 53.—Geront. trop.
australe, *Presl, Bot. Bemerk.* 20; *Garcke, Ind. Sem. Hort. Berol.* (1861) 10.—Austral.
Avicennae, *Gaertn. Fruct.* ii. 251. t. 135.—Geront. trop.
Bedfordianum, *A. St. Hil. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 48.—Bras.
Behrianum, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 13 = A. Avicennae.
benedictum, *Bunb. in Proc. Linn. Soc.* i. (1841) 109.—Bras.
bidentatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 68.—Abyss.; Ind. or.
blandum, *Fenzl, Delect. Sem. Hort. Bot. Vindob.* (1858).—Mexic.
californicum, Benth. Bot. Voy. Sulph. 8 = A. Avicennae.
calycinum, *Presl, Rel. Haenk.* ii. 115.—Peruv.
carneum, *A. St. Hil. Fl. Bras. Mer.* i. 205.—Bras.
ceratocarpum, *Hook. Bot. Misc.* iii. (1833) 154.—Chili.
Chapelieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 508.—Madag.
cinereum, *Griseb. in Goett. Abh.* xxiv. (1879) 45.—Reg. Argent.
circinnatum, *G. Don, Gen. Syst.* i. 502.—Bras.
cistiflorum, *G. Don, l. c.* 503.—Java.
coccineum, *Hort. ex Regel, Cat. Pl. Hort. Aksakov.* (1860) 1, nomen.—Hab.?
confertiflorum, *A. Rich. Ess. Fl. Cub.* i. 153.—Cuba.
contractum, Sweet, Hort. Brit. ed. II. 64 = Sida contracta.
cornutum, Sweet, l. c. 65 = Sida cornuta.
crassifolium, *G. Don, Gen. Syst.* i. 504.—Ind. occ.
crispum, *Sweet, Hort. Brit.* ed. I. 53.—Reg. trop.
cryptopetalum, *F. Muell. ex Benth. Fl. Austral.* i. 201.—Austral.
Cunninghami, *Benth. l. c.* 205.—Austral.
cymosum, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 185.—Am. trop.
cysticarpum, Hance, in Walp. Ann. ii. 157 = indicum.
Darwini, *Hook. f. Bot. Mag.* t. 5917.—Bras.
decarpum, *Rafin. New Fl. Am.* i. 40.—Am. bor.
deflexum, *Moench, Meth.* 620.—Hab.?
densiflorum, *Walp. Rep.* i. 324.—Chili.
denticulatum, Webb, Fragm. Fl. Aethiop. 51 = fruticosum.
depauperatum, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 232.—Ins. Galap.
dianthum, *Presl, Reliq. Haenk.* ii. 114.—Peruv.
diplotrichum, F. Muell. in Linnaea, xxv. (1852) 380 = Fraseri.
discissum, *Schlecht. in Linnaea*, 218.—Reg. Argent.
divaricatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 204.—Mexic.
domingense, *Turcz. l. c.* 205.—Ins. S. Doming.
elaeocarpoides, Webb, Fragm. Fl. Aethiop. 53 = ramosum.
elatum, *Griseb. Fl. Brit. W. Ind.* 79.—Ind. occ.; Mexic.
elegans, *A. St. Hil. Fl. Bras. Mer.* i. 207.—Bras.
elegans, Walp. Rep. i. 325 = Sida elegans.
ellipticum, *Schlecht. in Linnaea*, xi. (1837) 368.—Mexic.
elongatum, Moench, Meth. Suppl. 205 = indicum.
crosum, *Schlecht. in Linnaea*, xi. (1837) 367.—Mexic.
esculentum, *A. St. Hil. Fl. Bras. Mer.* i. 240.—Bras.
excelsior, G. Don, Gen. Syst. i. 500 = Wissadula excelsior.
exonemum, *F. Muell. Fragm.* xi. 63.—Austral.
exstipulare, *G. Don, Gen. Syst.* i. 503.—Ins. Borbon.
falcatum, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 48.—Bras.
ferrugineum, H. B. & K. Nov. Gen. et Sp. v. 271 = Sida ferruginea.
Figarianum, Webb, Fragm. Fl. Aethiop. 52 = graveolens.
floribundum, *Schlecht. in Linnaea*, xi. (1837) 366.—Mexic.
foetidum, *Moench, Meth. Suppl.* 206.—Hab.?
Fraseri, *Walp. Ann.* ii. 158.—Austral.
frutescens, Medic. Malv. 29 = indicum.
fruticosum, *Guill. & Perr. Fl. Seneg. Tent.* i. 70.—As. et Afr. trop.
furfurellum, *Miq. Fl. Ind. Bat.* i. II. 144.—Java.
geminiflorum, H. B. & K. Nov. Gen. et Sp. v. 274. t. 474 = ibarrense.
geranioides, *Benth. Fl. Austral.* i. 202.—Austral.

ABUTILON:—
giganteum, *Presl, Rel. Haenk.* ii. 116.—Am. trop.
glaucum, *G. Don, Gen. Syst.* i. 504.—Afr. trop.
glaucum, Sweet, Hort. Brit. ed. I. 54 = muticum.
glechomaefolium, *A. St. Hil. Fl. Bras. Mer.* i. 198. t. 411.—Bras.
globiflorum, *G. Don, Gen. Syst.* i. 502.—Maurit.
Goudotianum, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 184.—N. Granat.
grandiflorum, G. Don, Gen. Syst. i. 504 = indicum.
grandifolium, Sweet, Hort. Brit. ed. I. 53 = molle.
graveolens, Seem. Bot. Voy. Herald, 365 = indicum.
graveolens, *Wight & Arn. Prod.* 56.—Geront. trop.
Grevilleanum, *Walp. Rep.* i. 324.—Chili.
Guichenotianum, *Decne. in Herb. Timor. Desc.* 106 (1835).—Ins. Timor.
gymnanthemum, *Griseb. in Goett. Abh.* xxiv. (1879) 47.—Reg. Argent.
Haenkeanum, *Presl, Rel. Haenk.* ii. 115.—Mexic.
halophilum, F. Muell. in Linnaea, xxv. (1852) 381 = Fraseri.
hernandioides, Sweet, Hort. Brit. ed. I. 53 = Sida hernandioides.
heterotrichum, Hochst. ex Mast. in Oliver, Fl. Trop. Afr. i. 187 = hirtum.
hirsutissimum, Moench, Meth. Suppl. 205 = indicum.
hirtum, *G. Don, Gen. Syst.* i. 503.—Reg. trop.
hirtum, Sweet, Hort. Brit. ed. I. 53 = graveolens.
holosericeum, *Scheele, in Linnaea*, xxi. (1847) 471.—Texas.
Hulseanum, *Torr. ex A. Gray, Pl. Fendl.* 23.—N. Mexic.
hypoleucum, *A. Gray, Pl. Wright.* i. 20.—Texas.
ibarrense, *H. B. & K. Nov. Gen. et Sp.* v. 272.—Ecuador.
imberbe, G. Don, Gen. Syst. i. 502 = crispum.
inaequilaterum, *A. St. Hil. Fl. Bras. Mer.* i. 198. t. 40.—Bras.
incanum, *Sweet, Hort. Brit.* ed. I. 53.—Ins. Sandvic.; Mexic.
indicum, *Sweet, l. c.* 54.—Geront. trop.
insigne, *Planch. in Van Houtte, Fl. des Serres*, Sér. I. vi. (1850–51) 41.—N. Granat.
integerrimum, *Hook. Bot. Mag.* t. 4360.—Mexic.
intermedium, *Hochst. ex Schweinf. Beitr. Fl. Aethiop.* 49.—Geront. trop.
Jacquini, *G. Don, Gen. Syst.* i. 503.—Am. bor.
Julianae, *Endl. Prod. Fl. Norf.* 135.—Ins. Norfolc.
Kotschyi, Hochst. in Webb, Fragm. Fl. Aethiop. 52 = hirtum.
lanatum, *Miq. in Linnaea*, xxii. (1849) 553.—Bras.
laxiflorum, Guill. & Perr. Fl. Seneg. Tent. i. 66 = Wissadula rostrata.
leiospermum, *Griseb. Fl. Brit. W. Ind.* 79.—Ind. occ.
Lemmoni, *S. Wats. in Proc. Am. Acad.* xx. (1885) 357.—Calif.
lepidum, F. Muell. Fragm. vi. 168 = Sida lepida.
Leschenaultianum, Sweet, Hort. Brit. ed. I. 53 = Wissadula Leschenaultiana.
Leucanthemum, *A. St. Hil. Fl. Bras. Mer.* i. 200.—Bras.
leucopetalum, *F. Muell. ex Benth. Fl. Austral.* i. 200.—Austral.
lignosum, G. Don, Gen. Syst. i. 501 = Jacquini.
longicuspe, *Hochst. ex A. Rich. Fl. Abyss.* i. 69.—Abyssin.
longilobum, *F. Muell. Fragm.* ix. 130.—Austral.
lucianum, *Sweet, Hort. Brit.* ed. I. 53.—Ind. occ.
macranthum, *Peyr. in Linnaea*, xxx. (1859–60) 59.—Mexic.
macranthum, *A. St. Hil. Fl. Bras. Mer.* i. 208.—Bras.
macrocarpum, *Guill. & Perr. Fl. Seneg. Tent.* i. 69.—Afr. trop.
macrocarpum, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 47.—Bras.
macrophyllum, *St. Hil. & Naud. l. e.*—Bras.
macropodum, *Guill. & Perr. Fl. Seneg. Tent.* i. 69. t. 14.—Afr. trop.
macrum, *F. Muell. Fragm.* ix. 59.—Austral.
malachroides, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 49.—Bras.
marmoratum, Hort. ex Lem. in Illustr. Hortic. iii. (1856) t. 82 = Hibiscus marmoratus.
mauritanicum, Medic. Malv. 28 = scq.
mauritianum, Sweet, Hort. Brit. ed. I. 53 = indicum.
megapotamicum, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 49.—Am. trop.
melanocarpum, *St. Hil. & Naud. l. c.* 48.—Bras.
mendocinum, *Phil. Sert. Mend. Alt.* 6.—Chili.
Menziesii, *Seem. Fl. Vit.* 15.—Ins. Sandvic.
mexicanum, *Presl, Rel. Haenk.* ii. 115.—Mexic.

ABUTILON:—
micranthum, G. Don, Gen. Syst. i. 502, sphalm. = macranthum.
micropetalum, *Benth. Fl. Austral.* i. 201.—Austral.
microphyllum, A. Rich. Fl. Abyss. i. 70 = fruticosum.
microspermum, *G. Don, Gen. Syst.* i. 501.—Hab.?
Mitchelli, *Benth. Fl. Austral.* i. 201.—Austral.
molle, *Sweet, Hort. Brit.* ed. II. 65.—Peruv.
mollicomum, *Sweet, l. c.* ed. I. 54.—Hab.?
mollissimum, *Sweet, l. c.* 53.—Peruv.
montanum, *A. St. Hil. Fl. Bras. Mer.* i. 207.—Bras.
mucronulatum, A. Gray, in Proc. Am. Acad. v. (1862) 175 = Wissadula mucronulata.
muticum, *Sweet, Hort. Brit.* ed. II. 65.—Geront. trop.
muticum, Webb, Fragm. Fl. Aethiop. 51 = glaucum.
neilgherrense, *Munro, in Wight, Ill.* t. 66.—Ind. or.
neurocarpum, *Miq. Pl. Jungh.* 285.—Java.
Newberryi, *S. Wats. in Proc. Am. Acad.* xi. (1876) 125.—Am. bor.
niveum, *Griseb. in Goett. Abh.* xix. (1874) 92.—Reg. Argent.
notolophium, *A. Gray, in Proc. Am. Acad.* v. (1875) 175.—Mexic.
nudiflorum, Sweet, Hort. Brit. ed. I. 53 = Wissadula nudiflora.
Nuttallii, *Torr. & Gray, Fl. N. Am.* i. 231.—Am. bor.
obtusum, *G. Don, Gen. Syst.* i. 501.—Hab.?
occidentale, Medic. Malv. 29 = Gaya subtriloba.
Ochsenii, *Phil. in Linnaea*, xxviii. (1856) 613.—Chili.
orbiculatum, *G. Don, Gen. Syst.* i. 504.—China.
otocarpum, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 13.—Austral.
oxycarpum, *F. Muell. Fragm.* ii. 12.—Austral.
oxypetalum, *Planch. & Linden, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 184.—N. Granat.
oxyphyllum, Presl, Bot. Bemerk. 20 = polyandrum.
paeoniaeflorum, *Walp. Rep.* v. 95.—Bras.
Palmeri, *A. Gray, in Proc. Am. Acad.* viii. (1870) 289.—Am. bor.
pannosum, Webb, Fragm. Fl. Aethiop. 52 = glaucum.
paranthemoides, *Griseb. in Goett. Abh.* xxiv. (1879) 46.—Reg. Argent.
paranthemum, *Griseb. l. c.*—Reg. Argent.
Parishii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 357.—Arizona.
parviflorum, *A. St. Hil. Fl. Bras. Mer.* i. 201.—Bras.
parvulum, *A. Gray, Pl. Wright.* i. 21.—Texas.
patens, *A. St. Hil. Fl. Bras. Mer.* i. 200.—Bras.
pauciflorum, *A. St. Hil. l. c.* 206.—Bras.
pedunculare, *H. B. & K. Nov. Gen. et Sp.* v. 273.—Bras.
peraffine, Shuttl. ex Chapm. Fl. S. U. St. 56 = Jacquini.
periplocifolium, Sweet, Hort. Brit. ed. I. 53 = Wissadula rostrata.
permolle, *Sweet, l. c.*—Florida; Cuba.
petiolare, *H. B. & K. Nov. Gen. et Sp.* v. 272.—N. Granat.
pictum, *Walp. Rep.* i. 324.—Reg. Argent.
planiflorum, *C. Koch & Bouché, in C. Koch, Allg. Gartenz.* (1857) 98.—Mexic.
polyandrum, *G. Don, Gen. Syst.* i. 500.—Ind. or.
polyanthon, Sweet, Hort. Brit. ed. I. 53 = Sida polyantha.
populifolium, Sweet, l. c. = indicum.
pubescens, Moench, Meth. 620 = A. Avicennae.
pulchellum, Sweet, Hort. Brit. ed. I. 53 = scq.
pulchrum, G. Don, Gen. Syst. i. 503 = Plagianthus pulchellus.
pyramidale, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 203.—N. Granat.
racemosum, *Schlecht. in Linnaea*, xi. (1837) 367.—Mexic.
ramiflorum, *A. St. Hil. Fl. Bras. Mer.* i. 199.—Bras.
ramosissimum, *Presl, Rel. Haenk.* ii. 116.—Mexic.
ramosum, G. Don, Gen. Syst. i. 501 = Sida cordifolia.
ramosum, *Guill. & Perr. Fl. Seneg. Tent.* i. 68.—Geront. trop.
reflexum, *Sweet, Hort. Brit.* ed. I. 53; *Presl, Rel. Haenk.* ii. 114.—Mexic.
Regnellii, *Miq. in Linnaea*, xxii. (1849) 554.—Bras.
rigidum, *G. Don, Gen. Syst.* i. 504.—Afr. trop.
rivulare, *A. St. Hil. Fl. Bras. Mer.* i. 202.—Bras.
rufescens, G. Don, Gen. Syst. i. 502 = rufinerve.
rufescens, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) 1. 202.—Mexic.
rufinerve, *A. St. Hil. Fl. Bras. Mer.* i. 205. t. 42.—Bras.
rufinerve, Seem. Bot. Voy. Herald, 83 = cymosum.
Sellovianum, *Regel, in Ann. Sc. Nat.* Ser. IV. xii. (1859) 379.—Bras.
septemlobum, *Miq. in Linnaea*, xxii. (1849) 554.—Bras.
sessiliflorum, *Presl, Rel. Haenk.* ii. 113.—Mexic.

ABUTILON:—
sidoides, Dalz. & Gibs. Bomb. Fl. 18 = ramosum.
sidoides, *Hemsl. Diagn. Pl. Nov.* 24.—Mexic.
Sonneratianum, *Sweet, Hort. Brit.* ed. I. 54.—Afr. austr.
Sonorae, *A. Gray, Pl. Wright.* ii. 23.—Mexic.
sparmannioides, Guill. & Perr. Fl. Seneg. Tent. i. 70 = ramosum.
speciosum, G. Don, Gen. Syst. i. 502 = Sida speciosa.
spicatum, H. B. & K. Nov. Gen. et Sp. v. 271 = Wissadula spicata.
stenopetalum, *Garcke, in Bot. Zeit.* viii. (1850) 683.—N. Granat.
stipulare, *Presl, Rel. Haenk.* ii. 114.—Luzon.
striatum, *Dicks. in Bot. Reg.* (1839) *Misc.* 39.—Bras.
subviscosum, *Benth. Fl. Austral.* i. 202.—Austral.
sundaicum, *G. Don, Gen. Syst.* i. 500 (1831).—Java.
terminale, *A. St. Hil. Fl. Bras. Mer.* i. 203.—Bras.
texense, Torr. & Gray, Fl. N. Am. i. 231 = incanum.
Theophrasti, Medic. Malv. 28 = A. Avicennae.
Thurberi, *A. Gray, Pl. Nov. Thurb.* 307.—Texas.; N. Mexic.
thyrsodendron, *Griseb. in Goett. Abh.* xxiv. (1879) 48.—Reg. Argent.
tiliaefolium, Sweet, Hort. Brit. ed. I. 53 = A. Avicennae.
timoriense, *G. Don, Gen. Syst.* i. 500 (1831).—Ins. Timor.
tomentosum, Wight & Arn. Prod. 56 = muticum.
tortuosum, Guill. & Perr. Fl. Seneg. Tent. i. 66 = graveolens.
trichodum, A. Rich. Ess. Fl. Cub. i. 155 = crispum.
trilobatum, *Hemsl. Diagn. Pl. Nov. Mex.* 24.—Mexic.
triquetrum, *Presl, Reliq. Haenk.* ii. 115.—Mexic.
triquetrum, *Sweet, Hort. Brit.* ed. I. 53.—Ind. occ.
truncatum, *G. Don, Gen. Syst.* i. 503.—Ins. S. Doming.
tubulosum, *Walp. Ann.* ii. 158.—Austral.
umbellatum, *Sweet, Hort. Brit.* ed. I. 53.—Ind. occ.
umbelliferum, G. Don, Gen. Syst. i. 501 = seq.
umbelliflorum, *A. St. Hil. Fl. Bras. Mer.* i. 204.—Bras.
uratocarpum, Walp. Rep. i. 324 = ceratocarpum.
velutinum, *G. Don, Gen. Syst.* i. 504.—Afr. trop.
velutinum, A. Gray, Illustr. Gen. Pl. Am. Bor. ii. 67. t. 125 = holosericeum.
venosum, *Lem. Hortic. Univ. Misc.* 1844; *ex Fl. des Serres*, ii. (Mars 1846) t. 5.—Mexic.
verbascoides, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 202.—Venezuela.
vesicarium, Sweet, Hort. Brit. ed. I. 54 = A. Avicennae.
vexillarium, E. Morr. Belg. Hortic. (1864) 289 = megapotamicum.
virens, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 48.—Bras.
virgatum, *Sweet, Hort. Brit.* ed. I. 53.—Hab.?
vitifolium, *Presl, Rel. Haenk.* ii. 116.—Chili.
Wagenerianum, *Regel, Cat. Pl. Hort. Aksakov.* (1860) 1, nomen.—Hab.?
wissadifolium, *Griseb. in Goett. Abh.* xxiv. (1879) 47.—Reg. Argent.
Wrightii, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 162.—Texas.
zanzibaricum, *Boj. ex Mast. in Oliver, Fl. Trop. Afr.* i. 186.—Afr. trop.

ABUTUA, Lour. Fl. Cochinch. 630 (1790) = **Gnetum**, Linn.
africana, Lour. l. c. 631 (Quid?).—Afr. or.
indica, Lour. l. c. 630 = G. funiculare.

ACACALLIS, Lindl. Fol. Orchid. 1853. *ORCHIDEAE*, Benth. & Hook. f. iii. 544.
cyanea, *Lindl. l. c.*—Bras.

ACACIA, [Tourn.] Linn. Fl. Zeyl. 217 (1737); Willd. Sp. Pl. iv. 1049 (1805). *LEGUMINOSAE*, Benth. & Hook. f. i. 594.
ARTHROSPRION, Hassk. Retzia, i. 212 (1855).
CHITONANTHUS, Lehm. Pl. Preiss. ii. 368 (1847).
CUPARILLA, DREPAPHYLA, EBURNAX, ESCLERONA, GUMIFERA, HECATANDRA, POPONAX, SENEGALIA, ZIGMALOBA, Rafin. Sylva Tellur. 118-120 (1838).
FARNESIA, Gasp. Desc. Nuov. Gen. (1838).
HOOPESIA, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 433, partim.
PHYLLODOCE, Link, Handb. ii. 132 (1831).
TETRACHEILOS, Lehm. Pl. Preiss. ii. 868 (1847).
VACHELLIA, Wight & Arn. Prod. 272 (1834).
abietina, Willd. Sp. Pl. iv. 1051 = linifolia.
abstergens, Steud. Nom. ed. II. i. 3 = concinna.
abyssinica, *Hochst. ex Benth. in Hook. Lond. Journ. Bot.* v. (1846) 97.—Abyssin.

ACACIA:—
acanthocarpa, Willd. Enum. Hort. Berol. 1057 = Mimosa acanthocarpa.
acanthoclada, *F. Muell. Fragm.* iii. 127.—Austral.
acantholoba, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1089 = Mimosa acantholoba.
acanthophora, Steud. Nom. ed. II. i. 3 = Mimosa acanthocarpa.
acapulcensis, Kunth, Mim. 78. t. 24 = Lysiloma acapulcensis.
acatlensis, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 513.—Mexic.
acicularis, R. Br. in Ait. Hort. Kew. ed. II. v. 450 = juniperina.
acicularis, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 1056 = Farnesiana.
acinacea, *Lindl. in Mitch. Three Exped.* ii. 267.—Austral.
aciphylla, *Benth. in Linnaea*, xxvi. (1853-55) 627.—Austral.
Acradenia, F. Muell. in Journ. Linn. Soc. iii. (1859) 142 = umbellata.
acuifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 496.—Ins. Baham.
aculeaticarpa, Lag. Gen. et Sp. Nov. 16 = Mimosa acanthocarpa.
acuminata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 373.—Austral.
acutangula, Buch.-Ham. ex Wall. Cat. sub n. 5271 = Pithecolobium angulatum.
acutistipula, Mart. in Flora, xx. (1837) Beibl. ii. 107 = Mimosa acutistipula.
Adansonii, Guill. & Perr. Fl. Seneg. Tent. i. 249 = arabica.
adenantheroides, Mart. & Gal. in Bull. Acad. Brux. xii. (1846) 311 = Mimosa adenanthcroides.
adenocarpa, Walp. Rep. i. 924, sphalm. = seq.
adenopa, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 206 = A. Farnesiana.
adenophora, Spreng. Syst. iii. 140 = decurrens.
adenostylis, Fenzl, in Flora, xxvii. (1844) 312 = Dichrostachys nutans.
adhaerens, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 517.—Bras.
adiantoides, Spreng. Syst. iii. 146 = Piptadenia laxa.
adnata, *F. Muell. in Melb. Chemist* (July 1882).—Austral.
adstringens, A. Cunn. in G. Don, Gen. Syst. ii. 405 = falciformis.
adstringens, Mart. Reise Bras. i. 548—Stryphnodendron Barbatimam.
aduncа, A. Cunn. ex G. Don, Gen. Syst. ii. 406 = crassiuscula.
aegyptiaca, *Baill. Adansonia*, iv. (1863-64) 82.—Aegypt.
affinis, Sweet, Hort. Brit. ed. I. 102 = dealbata.
alata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 464.—Austral.
alba, Colla, Hort. Rip. App. ii. 339 = Calliandra portoricensis.
alba, *De Vriese, in Miq. Pl. Jungh.* 270.—Malaya.
alba, Willd. Enum. Hort. Berol. Suppl. 68 = leucophloea.
albescens, Steud. Nom. ed. II. i. 3 = tortuosa.
albicans, Kunth, Mim. 87. t. 27 = Pithecolobium albicans.
albida, *Delile, Fl. Egypt.* 286. t. 52. f. 3.—Afr. bor.
albida, Lindl. Bot. Reg. t. 1317 = tortuosa.
alliacea, Buch.-Ham. in Wall. Cat. n. 5258 = caesia.
alpina, *F. Muell. Fragm.* iii. 129.—Austral.
amara, Willd. Sp. Pl. iv. 1074 = Albizzia amara.
amazonica, *Benth. in Trans. Linn. Soc.* xxx. (1875) 523.—Am. trop.
ambigua, *Hoffmgg. Verz. Pfl. Nachtr.* iii. 15.—Afr. trop.
ambigua, *Vogel, in Linnaea*, x. (1835-36) 600.—Am. trop.
amblycarpa, R. Grah. in Wall. Cat. n. 5260 = pennata.
amblygona, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 332.—Austral.
amblyphylla, *F. Muell. in Wing, South Sc. Record* (July 1882).—Austral.
amentacea, *DC. Prod.* ii. 455.—Am. bor.
amentifera, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 141.—Austral.
amoena, Sieber, ex Walp. Rep. i. 897 = rubida.
amoena, *Wendl. Comm. Acac.* 16. t. 4.—Austral.
amythethophylla, *Steud. ex A. Rich. Fl. Abyss.* i. 245.—Abyssin.
anceps, *DC. Mém. Lég.* xii. 446.—Austral.
anceps, Hook. Ic. Pl. t. 167 = complanata.
aneura, *F. Muell. in Linnaea*, xxvi. (1853-55) 627.—Austral.
Angico, Mart. Mat. Med. Bras. 53 = Piptadenia rigida.
angulata, Desv. Journ. Bot. iii. (1814) 68 = discolor.

ACACIA:—
angulosa, Bertol. Fl. Guatim. 42 = filicina.
angustifolia, Lodd. Bot. Cab. t. 768 = longifolia.
angustifolia, Meissn. in Lehm. Pl. Preiss. i. 13 = triquetra.
angustifolia, Wendl. Comm. Acac. 34 = suaveolens.
angustiloba, DC. Prod. ii. 470 = Piptadenia peregrina.
angustisiliqua, Desf. Cat. Hort. Par. ed. III. 300 = Desmanthus virgatus.
anthelmintica, Baill. Hist. Pl. ii. 56 = Albizzia anthelmintica.
apiculata, Meissn. in Lehm. Pl. Preiss. i. 17 = subcaerulea.
arabica, *Willd. Sp. Pl.* iv. 1085.—Afr.; As. trop.
Arar, Buch.-Ham. in Wall. Cat. n. 5249 = A. Intsia.
arborea, Benth. Pl. Hartw. 13 = Lysiloma microphylla.
arborea, Willd. Sp. Pl. iv. 1064 = Pithecolobium filicifolium.
Arceuthos, Spreng. Syst. iii. 134 = juniperina.
arcuata, *Decne. Herb. Timor. Desc.* 133.—Ins. Timor.
arcuata, Sieber, ex Spreng. Syst. iii. 135 = melanoxylon.
arenosa, Willd. Sp. Pl. iv. 1054 (err. typ. 1060) = Mimosa arenosa.
argyrophylla, Hook. ex F. Muell. Pl. Vict. ii. 18 = brachybotrya.
arida, *A. Cunn. ex G. Don, Gen. Syst.* ii. 404.—Austral.
armata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 463.—Austral.
armata, Heyne, ex Wall. Cat. n. 2564 = A. Farnesiana.
armatoides, Hort. ex Walp. Rep. i. 888 = armata.
Aroma, Gill. in Hook. Bot. Misc. iii. (1833) 206 = macracantha.
aromatica, Poepp. ex Benth. in Trans. Linn. Soc. xxx. (1875) 502 = A. Farnesiana.
Arrophula, D. Don? in Wall. Cat. n. 5257 = Albizzia stipulata?
Arrophula, D. Don. Prod. Fl. Nep. 247 = pennata.
Asak, Willd. Sp. Pl. iv. 1077 = hamulosa?
asparagoides, *A. Cunn. in Field, N. S. Wales*, 343.—Austral.
aspera, *Lindl. in Mitch. Three Exped.* ii. 139.—Austral.
asperulacea, F. Muell. in Journ. Linn. Soc. iii. (1859) 123 = lycopodifolia.
aspidioides, G. F. W. Mey. Prim. Fl. Esseq. 165 = Pentaclethra filamentosa.
asplenioides, Nees, in Flora, iv. (1821) 303 = Calliandra mertensioides.
astringens, A. Cunn. ex G. Don, Gen. Syst. ii. 405 = penninervis.
ataxacantha, *DC. Prod.* ii. 459.—Afr. trop.
ataxiphylla, *Benth. in Linnaea*, xxvi. (1853-55) 605.—Austral.
atomiphylla, Burch. Trav. i. 341 = haematoxylon.
atramentaria, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 392.—Chili.
Aucheri, Benth. l. c. 499 = nubica.
aulacocarpa, *A. Cunn. ex Benth. l. c.* 379.—Austral.
aurea, *Noronha, in Verh. Batav. Gen.* v. (1790) Art. IV. 5.—Java.
auriculaeformis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 377.—Austral.
auriculata, Buch.-Ham. ex Wall. Cat. sub n. 5237 = Albizzia stipulata.
auriculata, Charp. ex Fourn. in Bull. Soc. Bot. Fr. xii. (1865) 400 = Albizzia auriculata.
aurita, Schlecht. in Linnaea, xii. (1838) 565 = Lysiloma aurita.
auronitens, *Lindl. Swan River App.* 15.—Austral.
Ausfeldi, Regel, Ind. Sem. Hort. Petrop. (1866) 106; Gartenfl. xvi. (1867) 225 = aspera.
axillaris, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 341.—Tasmania.
Azak, G. Don, Gen. Syst. ii. 412 = hamulosa?
Bagsteri, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 327.—Austral.
bahamensis, Griseb. Fl. Brit. W. Ind. 221 = Lysiloma latisiliqua.
bahiensis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 525.—Am. trop.
Balbisii, Spreng. Syst. iii. 140 = Pithecolobium fragrans.
Bancroftiana, Bert. ex Colla, Hort. Rip. 1; DC. Prod. ii. 473 = Caesalpinia bijuga.
barbinervis, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 326.—Austral.
basaltica, F. Muell. & Benth. in Journ. Linn. Soc. iii. (1859) 146 = Albizzia basaltica.
Baueri, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 344.—Austral.
Baxteri, *Benth. l. c.* 327.—Austral.

ACACIA :—

Baxteri, Meissn. in Lehm. Pl. Preiss. i. 7 = congesta.
bella, Mart. ex Spreng. Syst. iii. 141 = Calliandra bella.
Benthamii, Meissn. in Lehm. Pl. Preiss. 11 = cochlearis.
Berlandieri, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 522.—Mexic.
Bernieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 363. —Madag.
Berteriana, Balb. ex DC. Prod. ii. 470 = Pithecolobium fragrans.
Berteriana, *Spreng. Syst.* iii. 138.—Jamaica.
biceps, Willd. Sp. Pl. iv. 1075 = Leucaena glauca.
bidentata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 333.—Austral.
Bidwilli, *Benth. in Linnaea*, xxvi. (1853–55) 629.— Austral.
biflora, *R. Br. in Ait. Hort. Kew.* ed. II. v. 463.— Austral.
biflora, Paxt. Mag. ix. (1842) 221 = decipiens.
biglandulosa, Meissn. in Lehm. Pl. Preiss. ii. 205 = pentadenia.
Bilaura, Buch.-Ham. ex Wall. Cat. sub n. 5234 = odoratissima.
bimucronata, DC. Prod. ii. 469 = Mimosa sepiaria.
binervata, *DC. l. c.* 452.—Austral.
binervosa, DC. Mém. Lég. xii. 448 = bivenosa.
bivenosa, *DC. Prod.* ii. 452.—Austral.
Blomei, Ohlend. in Neue Allgem. Gartenz. (1845) 369 = saligna.
Boivini, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 363.— Madag.
bombycina, Benth. ex Paxt. Flow. Gard. ii. (1851–52). 101 = brachybotrya.
bonariensis, *Gill. in Hook. Bot. Misc.* iii. (1833) 207. —Reg. Argent.
Bonplandii, Gill. ex Hook. l. c. 205 = Mimosa Bonplandii.
bossiaeoides, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 323.—Austral.
bossiaeoides, Seem. Eingef. Acac. t. 1 = glaucoptera.
botrycephala, Desf. Cat. Hort. Par. ed. III. 300 = discolor.
brachyacantha, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 1055 = Mimosa acanthocarpa.
brachybotrya, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 347.—Austral.
brachyloba, Willd. Sp. Pl. iv. 1071 = Desmanthus brachylobus.
brachyphylla, *Benth. ex Schlecht. in Linnaea*, xxvi. (1853-55) 615.—Austral.
brachyptera, Benth. in Hook. Lond. Journ. Bot. i. (1842) 325 = incurva.
brachystachya, *Benth. Fl. Austral.* ii. 403.—Austral.
brasiliensis, *Spreng. Syst.* iii. 142.—Bras.
brevifolia, *Benth. Fl. Austral.* ii. 395.—Austral.
brevifolia, Lodd. Bot. Cab. t. 1235 = lunata.
brevipes, A. Cunn. in Bot. Mag. t. 358 = melanoxylon.
Browniana, Wendl. in Flora, ii. (1819) 139 = strigosa.
Brownii, Steud. Nom. ed. I. 2 = juniperina.
brunioides, *A. Cunn. ex G. Don, Gen. Syst.* ii. 404.— Austral.
Buchananiana, Voigt, Hort. Suburb. Calc. 293 = pinnata.
Burkei, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 98.—Afr. austr.
Burkittii, *F. Muell. ex Benth. Fl. Austral.* ii. 400.— Austral.
Burmanniana, *DC. Prod.* ii. 461.—Ind. or.
buxifolia, *A. Cunn. in Field, N. S. Walcs*, 344.—Austral.
Bynocana, *Benth. in Linnaea*, xxvi. (1853-55) 614.— Austral.
caduca, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1089 = Minosa caduca.
caesia, Roxb. ex Wall. Cat. n. 5253 A = pennata.
caesia, *Wight & Arn. Prod.* 278.—Ind. or.
caesia, Willd. Sp. Pl. iv. 1090, ex parte = A. Intsia.
caffra, *Willd. l. c.* 1078.—Afr. austr.
calamifolia, Sieber, ex Steud. Nom. ed. II. i. 4 = quadrilateralis.
calamifolia, *Sweet, ex Bot. Reg.* t. 839.—Austral.
Calcyi, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 347 = podalyriaefolia.
callicoma, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 104.—Afr. austr.
calligera, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 141 = Wickhami.
Callistemon, Schlecht. in Linnaea, xii. (1838) 568 = Calliandra grandiflora.
callosa, *Bertol. Pl. Nov. Hort. Bon.* i. 9.—Hab.?

ACACIA :—

callosa, Spreng. Syst. iii. 138 = Piptadenia communis.
calyculata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 379.—Austral.
Campbellii, Arn in Nov. Act. Nat. Cur. xviii. 1. (1836) 333 = planiformis.
campylacantha, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 242 = A. Suma.
campylophylla, *Benth. ex Schlecht. in Linnaea*, xxvi. (1853–55) 605.—Austral.
Candolleana, Meissn. in Lehm. Pl. Preiss. ii. 206 = Drummondii.
canescens, R. Grah. in Wall. Cat. n. 5256 = pennata.
canescens, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 314 = Mimosa depauperata.
capensis, Burch. Trav. i. 195 = horrida.
capensis, *Colla, in Mem. Acc. Torin.* xxxv. (1831) 175. —Afr. austr.
caracasana, Willd. Sp. Pl. iv. 1068 = Calliandra caracasana.
carbonaria, Schlecht. in Linnaea, xii. (1838) 571 = villosa.
cardiophylla, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 385.—Austral.
Caringa, Buch.-Ham. in Wall. Cat. n. 5239 = Leucaena glauca.
cassioides, *Willd. Enum. Hort. Berol.* 1051.—Hab.?
Catechu, Brandis, For. Fl. 186 = A. Sundra.
Catechu, Wight & Arn. Prod. 272 = A. Suma.
Catechu, *Willd. Sp. Pl.* iv. 1079.—Ind. or.
catechuoides, Benth. in Hook. Lond. Journ. Bot. i. (1842) 510 = A. Catechu.
caudata, DC. Prod. ii. 456 = Entada polystachya.
Cavenia, Bert. ex Bull. Férussac, xx. (1830) 108 = A. Farnesiana.
Cebil, *Griseb. in Goett. Abh.* xix. (1874) 136.—Reg. Argent.
cedroides, *Benth. ex Schlecht. in Linnaea*, xxvi. (1853–55) 615.—Austral.
celastrifolia, Benth. in Hook. Lond. Journ. Bot. i. (1842) 349 = myrtifolia.
centrophylla, *DC. Prod.* ii. 465.—Jamaica.
centrota, Spreng. ex DC. l. c. = praec.
Ceratonia, Willd. Sp. Pl. iv. 1091 = Mimosa Ceratonia.
chlorantha, Zucc. in Abh. Akad. Muench. ii. (1832) 347 = filicina.
chordophylla, F. Muell. ex Benth. in Linnaea, xxvi. (1853-55) 612 = rigens.
choriophylla, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 495.—Ins. Baham.
chrysantha, *DC. Prod.* ii. 471.—Guiana.
chrysobotrys, Meissn. Ind. Sem. Hort. Basil. (1842) 1 = spectabilis.
chrysocoma, Miq. in Fl. Ind. Bat. i. 1. 6 = tomentosa.
Chundra, Willd. Sp. Pl. iv. 1078 = A. Catechu.
cibaria, *F. Muell. in Melb. Chemist* (July 1882).— Austral.
ciliata, R. Br. in Ait. Hort. Kew. ed. II. v. 465 = strigosa.
ciliata, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 1055.—Am. austr.
cincinnata, *F. Muell. Fragm.* xi. 35.—Austral.
Cineraria, Willd. Sp. Pl. iv. 1057 = Prosopis spicigera.
cinerascens, Sieber, ex DC. Prod. ii. 454 = glaucescens.
cinerea, Spreng. Syst. iii. 143 = Dichrostachys cinerea.
Clausseni, Benth. in Hook. Lond. Journ. Bot. i. (1842) 518 = paniculata.
clavata, Schlecht. in Linnaea, xx. (1847) 662 = montana.
clavigera, *E. Mey. Comm. Pl. Afr. Austr.* 168.—Afr. austr.
Coa, Walp. Ann. iv. 626 = A. Koa.
cochlearis, *Wendl. Comm. Acac.* 15.—Austral.
cochliacantha, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1081.—Am. austr.
cochliocarpa, *G. Don, Gen. Syst.* ii. 422.—Bras.
cochlocarpa, *Meissn. in Bot. Zeit.* xiii. (1855) 10.— Austral.
Colleana, Presl, Rep. Bot. Syst. 183 = Calliandra portoricensis.
colletioides, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 336.—Austral.
colletioides, F. Mucll. Pl. Vict. ii. 5 = continua.
colubrina, Mart. Herb. Fl. Bras. 107 = Piptadenia colubrina.
comorensis, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 363.—Afr. trop.
complanata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 369.—Austral.
compressa, Schult. ex Colla, Hort. Ripul. App. ii. 393 = visncoides.

ACACIA :—

concinna, *DC. Prod.* ii. 464.—As. trop.
concinna, *Phil. in Anal. Univ. Chil.* (1870) II. 170.— Chili.
concinna, Wall. Cat. n. 5250 D = pennata?
concordiana, Loud. Hort. Brit. 409 = Pithecolobium umbellatum.
conferta, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 345.—Austral.
congesta, *Benth. l. c.* 327.—Austral.
conjunctifolia, *F. Muell. Fragm.* xi. 68.—Austral.
conspersa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 140.—Austral.
conspicua, A. Cunn. ex G. Don, Gen. Syst. ii. 405 = vestita.
constricta, *Benth. in A. Gray, Pl. Wright.* i. 66.— N. Mexic.
continua, *Benth. Fl. Austral.* ii. 322.—Austral.
contorta, DC. Prod. ii. 470 = Piptadenia contorta.
Cooleyi, Eaton, Man. ed. VI. (1833) 1 = Desmanthus Jamesii.
cordata, *Steud. Nom.* ed. II. i. 4.—Austral.
coriacea, *DC. Mém. Lég.* xii. 446.—Austral.
coriaria, Buch.-Ham. ex Wall. Cat. sub n. 5264 B = A. Farnesiana.
cornigera, Willd. Sp. Pl. iv. 1080 = spadicigera, sphaerocephala.
coronillaefolia, *Desf. Tabl.* ed. II. 307.—Afr. bor.
costata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 339.—Austral.
Coulteri, *Benth. ex A. Gray, Pl. Wright.* i. 66.—N. Mexic.
Courrantiana, DC. Prod. ii. 465 = Mimosa acantholoba.
crassicarpa, *A. Cunn. ex Benth in Hook. Lond. Journ. Bot.* i. (1842) 379.—Austral.
crassifolia, *A. Gray, in Mem. Am. Acad.* N. S. v. (1855) 317.—Mexic.
crassistipula, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 326.—Austral.
crassiuscula, Meissn. in Lehm. Pl. Preiss. i. 16 = pycnophylla.
crassiuscula, *Wendl. Comm. Acac.* 31. t. 8.—Austral.
crispula, *Benth. ex Schlecht. in Linnaea*, xxvi. (1853–55) 606.—Austral.
cultrata, *Ait. ex Steud. Nom.* ed. II. i. 4.—Austral.
cultriformis, *A. Cunn. ex G. Don, Gen. Syst.* ii. 406.— Austral.
cumanensis, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1058 = Prosopis juliflora.
Cumingii, Benth. in Hook. Lond. Journ. Bot. i. (1842) 525 = villosa.
cuneata, *Benth. in Enum. Pl. Hueg.* 42.—Austral.
Cunninghamii, *Hook. Ic. Pl.* t. 165.—Austral.
Cunninghamii, Sweet, Hort. Brit. ed. II. 164 = trinervata.
curvifolia, *Bonpl. Desc. Pl. Malm.* 99. t. 40.—Hab.?
cuspidata, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 337 = diffusa.
cuspidata, Schlecht. in Linnaea, xii. (1838) 573 = filicina.
cyanophylla, *Lindl. Bot. Reg.* (1839) *Misc.* 45.— Austral.
cyclophylla, Schlecht. in Linnaea, xx. (1847) 663 = obliqua.
Cyclops, *A. Cunn. ex G. Don, Gen. Syst.* ii. 404.— Austral.
cyclosperma, DC. Prod. ii. 456 = Albizzia fulgens.
cycnorum, Benth. in Hook. Lond. Journ. Bot. i. (1842) 388 = pulchella.
cycnorum, Hook. Bot. Mag. t. 4653 = obscura.
cygnorum, Meissn. in Lehm. Pl. Preiss. i. 22 = pulchella.
cylindriflora, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 311.—Mexic.
cymbacantha, Zucc. ex Benth. in Trans. Linn. Soc. xxx. (1875) 501 = cochliacantha?
cyperophylla, *F. Muell. ex Benth. Fl. Austral.* ii. 400. —Austral.
Daintreana, F. Muell. Fragm. iv. 6 = excelsa.
Dalea, Desv. Journ. Bot. iii. (1814) 69 = Dichrostachys cinerea.
Dallachiana, *F. Muell. Fragm.* i. 7.—Austral.
daphnifolia, Meissn. in Bot. Zeit. xiii. (1855) 11 = microbotrya.
dasyphylla, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 359 = lineata.
daviesiaefolia, *A. Cunn. ex Loud. Hort. Brit.* 406.— Austral.

ACACIA :—

daviesioides, A. Cunn. ex G. Don, Gen. Syst. ii. 402 = diffusa.
dealbata, A. Cunn. in Field, N. S. Wales, 345 = lnnata.
dealbata, *Link*, *Enum. Hort. Berol.* 445.—Austral.
Decaisnei, Steud. Nom. ed. II. i. 4 = arcnata.
decipiens, *R. Br. in Ait. Hort. Kew.* ed. II. v. 463.—Ind. or.; Austral.
decora, *Reichb. Ic. Exot.* t. 199.—Austral.
decurrens, *Willd. Sp. Pl.* iv. 1072.—Austral.
decussata, Tenore, Cat. Ort. Nap. (1845) 77 = longifolia.
delibrata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 374.—Anstral.
deltoidea, *A. Cunn. ex G. Don, Gen. Syst.* ii. 401.—Anstral.
demostachya, Benth. Pl. Hartw. 12 = acapulcensis.
Dempsteri, *F. Muell. Fragm.* xi. 65.—Austral.
densa, Wall. Cat. n. 5262 = leucophloea.
densifolia, Benth. in Hook. Lond. Journ. Bot. i. (1842) 360 = aspera.
denticulosa, *F. Muell. Fragm.* x. 32.—Austral.
dentifera, *Benth. in Maund, Botanist*, iv. t. 179.—Anstral.
denudata, Lehm. Del. Sem. Hort. Hamb. (1842) = pulchella.
depauperata, Mart. ex Steud. Nom. ed. II. i. 4 = Desmanthus virgatns.
dependens, A. Cunn. ex Benth. in Hook. Lond. Jonrn. Bot. i. (1842) 372 = longifolia.
dermatophylla, *Benth. Fl. Austral.* ii. 346.—Austral.
desmostachya, Benth. Pl. Hartw. 13 = Lysiloma acapulcensis.
detinens, *Burch. Trav.* i. 310.—Afr. anstr.
dictyocarpa, Benth. in Linnaea, xxvi. (1853–55) 616 = brachybotrya.
dictyophleba, *F. Muell. Fragm.* iii. 128.—Austral.
Dietrichiana, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 149.—Austral.
diffusa, *Lindl. Bot. Reg.* t. 634.—Anstral.
dilatata, *Benth. in Linnaea*, xxvi. (1853–55) 608.—Austral.
dilatata, *Steud. Nom.* ed. II. i. 4.—Austral.
dimidiata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 381.—Austral.
dineura, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 130 = binervata.
diptera, *Lindl. Swan River App.* 15.—Austral.
diptera, Willd. Enum. Hort. Berol. 1051 = Prosopis juliflora.
discolor, Whitley, ex Steud. Nom. ed. II. i. 4 = prismatica.
discolor, *Willd. Sp. Pl.* iv. 1068.—Austral.
dissitiflora, Benth. in Hook. Lond. Journ. Bot. i. (1842) 371 = longifolia.
dissoneura, *F. Muell. in Wing, South Sc. Record* (July 1882).—Austral.
distachya, Moç. & Sesse, ex DC. Prod. ii. 456 = Mimosa distachya.
divaricata, A. Cunn. ex Steud. Nom. ed. II. i. 4 = dilatata.
divaricata, Willd. Sp. Pl. iv. 1070 = Lysiloma Schiedeana.
divergens, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 331.—Austral.
diversifolia, Schlecht. in Linnaea, xii. (1838) 571 = Leucaena diversifolia.
dodonaeifolia, *Willd. Enum. Hort. Berol. Suppl.* 68.—Austral.
dolabriformis, Colla, Hort. Ripul. 1 = decipiens.
dolabriformis, A. Cunn. in Hook. Ic. Pl. t. 169 = dimidiata.
dolabriformis, Wendl. Acac. 56 = Daviesia incrassata.
domingensis, Bert. ex DC. Prod. ii. 464 = Mimosa domingensis.
doratoxylon, *A. Cunn. in Field, N. S. Wales*, 345.—Austral.
doratoxylon, Meissn. in Lehm. Pl. Preiss. i. 19 = acuminata.
drepanocarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 137.—Austral.
Drummondii, *Benth. in Lindl. Swan River App.* 15.—Austral.
dnmetorum, *DC. Prod.* ii. 458.—Bras.
dumosa, Wight & Arn. Prod. 274 = latronum.
dura, *Benth. in Linnaea*, xxvi. (1853–55) 622.—Anstral.
Durandiana, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 453 = Greggii.
dysophylla, Benth. in Hook. Lond. Journ. Bot. i. (1842) 346 = undulaefolia.
eburnea, *Willd. Sp. Pl.* iv. 1081.—Oriens; Ind. or.

ACACIA :—

echinula, DC. Prod. ii. 449 = juniperina.
Edgeworthii, *T. Anders. in Journ. Linn. Soc.* v. (1860) *Suppl.* 18.—Ind. or.
edulis, Humb. & Bonpl. ex Willd. Ennm. Hort. Berol. 1056 = A. Farnesiana.
eglandulosa, DC. Mém. Lég. xii. 445 = cochlearis.
Ehrenbergiana, *Hayne, Arzneigew.* x. t. 29.—Afr. trop.
Ehrenbergii, T. Nees, Pl. Offic. t. 334 = praec.
Ehrenbergii, *Steud. Nom.* ed. II. i. 5.—Mexic.
elata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 383.—Austral.
elata, R. Grah. in Wall. Cat. n. 5233 = Albizzia procera.
elegans, Schlecht. in Linnaea, xii. (1838) 569 = filicina.
elephantina, Burch. Trav. ii. 236 = Elephantorrhiza Burchelli.
Elephantorrhiza, Burch. ex DC. Prod. ii. 457 = praec.
elliptica, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 347 = bivenosa.
elongata, *Sieber, ex DC. Prod.* ii. 451.—Anstral.
emarginata, Wendl. Comm. Acac. 27 = stricta.
Emoryana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 522.—Am. bor.
Endlicheri, Meissn. ex Lehm. Pl. Preiss. i. 21 = strigosa.
ephedroides, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 370.—Austral.
eriadenia, *Benth. l. c.* v. (1846) 98.—Afr. austr.
eriantha, *Desv. Journ. Bot.* iii. (1814) 68.—Ind. or.?
ericaefolia, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 345.—Austral.
erinacea, *Benth. l. c.* 360.—Anstral.
erioclada, *Benth. in Linnaea*, xxvi. (1853–55) 606.—Austral.
erioloba, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1215 = Edgeworthii.
erioloba, E. Mey. Comm. Pl. Afr. Austr. i. 171 = giraffae.
eriophylla, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1532 = viscidula.
erubescens, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 343.—Afr. trop.
erythrantha, Steud. ex A. Rich. Tent. Fl. Abyss. i. 243 = A. Suma.
erythrocephala, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 362 = aspera.
erythropus, Tenore, Ind. Sem. Hort. Nap. (1842) 12 = Riceana.
esculenta, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 314 = Leucaena pulverulenta.
esculenta, Moç. & Sesse, ex DC. Prod. ii. 470 = Lencaena esculenta.
Esterhazia, *Mackay, ex Regel, Cat. Pl. Hort. Aksakov.* 2.—Austral.
estrophiolata, *F. Muell. in Wing, South Sc. Record* (July 1882).—Austral.
etbaica, *Schweinf. in Linnaea*, xxxv. (1867–68) 330. t. 7, 8.—Afr. trop.
excelsa, *Benth. in Mitch. Trop. Austral.* 225.—Austral.
explanata, Jungh. Reise, 260; ex Miq. Fl. Ind. Bat. I. 9 = leucophloea.
exsudans, Lindl. in Mitch. Three Exped. ii. 214 = verniciflua.
extensa, *Lindl. Swan River App.* 15.—Austral.
fagonioides, Benth. in Hook. Lond. Journ. Bot. i. (1842) 387 = pulchella.
falcata, Desf. Tabl. ed. II. 207 = Prosopis juliflora.
falcata, Sieber, ex Steud. Nom. ed. II. i. 5 = plagiophylla.
falcata, *Willd. Sp. Pl.* iv. 1053.—Austral.
falcifolia, Hoffmgg. Verz. Pfl. Nachtr. 125, 201 = procumbens.
falciformis, DC. Prod. ii. 452 = penninervis.
falcinella, Meissn. in Bot. Zeit. xiii. (1855) 11 = pycnantha.
falcinella, *Tausch, in Flora*, xix. (1836) 419.—Austral.
fallax, E. Mey. Comm. Pl. Afr. Austr. i. 169 = caffra.
farinosa, *Lindl. in Mitch. Three Exped.* ii. 146.—Austral.
Farnesiana, Wall. Cat. n. 5264 = planifrons.
Farnesiana, *Willd. Sp. Pl.* iv. 1083.—Reg. trop.
fasciculata, R. Br. in Salt, Abyss. App. 65 = A. Lahal.
fasciculata, Guill. & Perr. Fl. Seneg. Tent. i. 252 = tortilis.
fasciculata, Kunth, Mim. 75. t. 23 = Mimosa fasciculata.
fasciculifera, *F. Muell. ex Benth. Fl. Austral.* ii. 361.—Austral.
ferox, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 97.—Afr. anstr.
ferox, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 312.—Mexic.
ferruginea, *DC. Prod.* ii. 458.—Ind. or.

ACACIA :—

filicina, *Willd. Sp. Pl.* iv. 1072.—Mexic.
filicoma, Mart. in Flora, xx. (1837) Beibl. ii. 109 = Piptadenia nitida.
filifolia, Benth. in Hook. Lond. Journ. Bot. i. (1842) 369 = ephedroides.
fimbriata, A. Cunn. ex G. Don, Gen. Syst. ii. 406 = prominens.
Fistula, Schweinf. in Linnaea, xxxv. (1867–68) 344 = A. Seyal.
flaccida, *Spreng. Syst.* iii. 137.—Bras.
flava, Spreng. ex DC. Prod. ii. 469 = Piptadenia flava.
flavescens, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 381.—Austral.
flexicaulis, *Benth. l. c.* 505.—Texas.
flexifolia, *A. Cunn. ex Benth. l. c.* i. 359.—Austral.
flexuosa, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1082 = macracantha.
flexuosa, Lag. Gen. et Sp. Nov. 16 = Prosopis juliflora.
floribunda, Willd. Sp. Pl. iv. 1051 = longifolia.
fluviatilis, Spreng. Syst. v. Ind. 3 = polyphylla.
foetida, H. B. & K. Nov. Gen. et Sp. vi. 265 = Piptadenia foetida.
foliolosa, R. Grah. in Wall. Cat. n. 5241 = Albizzia myriophylla.
formosa, Kunth, Mim. 102. t. 32 = Calliandra formosa.
fragrans, Tenore, Cat. Ort. Nap. 1 = Calliandra caracasana.
Fraseri, Hook. Ic. Pl. t. 171 = podalyriaefolia.
frondosa, Willd. Sp. Pl. iv. 1076 = Lencaena glauca.
frutescens, *Steud. Nom.* ed. II. i. 5.—Ind. or.
fruticosa, Loud. Hort. Brit. 409 = frutescens.
fruticosa, Mart. Herb. Fl. Bras. 107 = Piptadenia latifolia.
fruticosa, Voigt, Hort. Suburb. Calc. 261 = Mimosa fruticosa, *Roxb.*
fulgens, Charp. ex Fonrn. in Bull. Soc. Bot. Fr. xix. (1865) 400 = Albizzia Charpentieri.
fulgens, Labill. Sert. Anstr. Caled. 68. t. 67 = Albizzia fulgens.
furcata, Desv. Journ. Bot. iii. (1814) 67 = Prosopis juliflora.
furcata, *Gill. in Hook. Bot. Misc.* iii. (1833) 206.—Am. austr.
furcifera, Lindl. in Mitch. Three Exped. ii. 267 = armata.
furfuracea, G. Don, Gen. Syst. ii. 405 = lunata.
galioides, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 344.—Austral.
gemella, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 40; iii. 16.—Hab.?
genistifolia, Link, Enum. Hort. Berol. ii. 442 = diffusa.
genistoides, *A. Cunn. ex Benth. Fl. Austral.* ii. 330.—Austral.
Gerrardi, *Benth. in Trans. Linn. Soc.* xxx. (1875) 508.—Afr. austr.
gigantea, Noronha, in Verh. Batav. Gen. v. (1790) Art. IV. 5 = Parkia speciosa.
Gilberti, *Meissn. in Lehm. Pl. Preiss.* ii. 204.—Austral.
Gilesiana, *F. Muell. in Melb. Chemist* (July 1882).—Austral.
Gilliesii, Steud. Nom. ed. II. i. 5 = fnrcata.
Giraffa, Sieber, ex Steud. l. c. = A. Seyal.
giraffae, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 238 = stenocarpa.
giraffae, *Willd. Enum. Hort. Berol.* 1054.—Afr. austr.
glabrata, Schlecht. in Linnaea, xii. (1838) 569 = filicina.
gladiiformis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 354.—Austral.
glandulosa, Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 360 = Leucaena Forsteri.
glandulosa, Willd. Sp. Pl. iv. 1071 = Desmanthus brachylobus.
glauca, Moench, Meth. 466 = Lencaena glauca.
glaucescens, *Willd. Sp. Pl.* iv. 1052.—Austral.
glaucifolia, Banmg. ex Lehm. Pl. Preiss. i. 14 in nota = cnltriformis.
glaucophylla, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 243.—Afr. trop.
glaucoptera, *Benth. in Linnaea*, xxvi. (1853–55) 604.—Austral.
glomerosa, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 521.—Am. austr.
glutinosa, F. Muell. Fragm. iv. 6 = ixiophylla.
Gnidinm, *Benth. Fl. Austral.* ii. 359.—Austral.
gonoacantha, Mart. in Flora, xx. (1837) Beibl. ii. 109 = Piptadenia communis.

ACACIA:—
gonocarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 136.—Austral.
gonoclada, *F. Muell. l. c.* 140.—Austral.
gonophylla, *Benth. in Linnaea*, xxvi. (1853-55) 613.—Austral.
gracilis, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 311 = Calliandra formosa.
graminea, Lehm. Del. Sem. Hort. Hamb. (1842) = extensa.
grandiflora, Willd. Sp. Pl. iv. 1074 = Calliandra grandiflora.
grandis, Henfr. in Gard. Mag. Bot. iii. (1851) 177 = pulchella.
grandisiliqua, Benth. in Hook. Lond. Journ. Bot. i. (1842) 518 = paniculata.
grandistipula, *Benth. l. c.* 511.—Bras.
granulosa, Labill. Fl. N. Caled. 67. t. 66 = Albizzia granulosa.
grata, Willd. Enum. Hort. Berol. 1056 = Piptadenia macrocarpa.
graveolens, A. Cunn. ex G. Don, Gen. Syst. ii. 404 = verniciflua.
Greggii, *A. Gray, Pl. Wright.* i. 65.—N. Mexico.
Gregorii, *F. Muell. Fragm.* iii. 47.—Austral.
Greveana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 364.—Madag.
Guachapele, *H. B. & K. Nov. Gen. et Sp.* vi. 281 = Lysiloma Guachapele.
guadalupensis, DC. Prod. ii. 464 = riparia.
gualparensis, Steud. Nom. ed. II. i. 5 = Albizzia stipulata.
guatemalensis, Hook. & Arn. Bot. Beech. Voy. 419 = Mimosa guatemalensis.
guayaquilensis, Desf. Tabl. Hort. Par. ed. II. 208 = Mimosa guayaquilensis.
guianensis, Willd. Sp. Pl. iv. 1061 = Stryphnodendron guianense.
Guilandinae, DC. Prod. ii. 465 = Mimosa obovata.
gummifera, Delile, Fl. Aeg. Illustr. 79 = spirocarpa.
gummifera, *Willd. Sp. Pl.* iv. 1056.—Afr. bor.
Gunnii, Benth. in Hook. Lond. Journ. Bot. i. (1842) 332 = vomeriformis.
guttata, *Hoffmgg. Verz. Pfl. Nachtr.* i. 201, 125.—Hab.?
guttulifera, *Hoffmgg. l. c.* iii. 16 = Martinicus.
gyrocarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 241 = albida.
Habbas, Link, Enum. Hort. Berol. ii. 444 = Mimosa aspera?
habbasioides, Boj. Hort. Maurit. 116 = concinna.
hadiensis, *DC. Prod.* ii. 472.—Arab.
haematomma, Bert. ex DC. Mém. Lég. xii. 448. t. 68 = Calliandra hacmatomma.
Haematoxylon, *Willd. Enum. Hort. Berol.* 1056.—Afr. austr.
hakeoides, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 354.—Austral.
Hamiltoni, Desv. in Ham. Prod. Fl. Ind. Occ. 59 = Calliandra portoricensis.
hamulosa, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 509.—Arab.
harpophylla, *F. Muell. ex Benth. Fl. Austral.* ii. 389.—Austral.
Hartwegi, Benth. Pl. Hartw. 13 = filicina.
Harveyi, *Benth. Fl. Austral.* ii. 368.—Austral.
hastulata, *Sm. in Rees, Cyclop.* xxxix. *Suppl.*—Austral.
Hayesii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 524.—Am. trop.
hebecephala, A. Cunn. ex G. Don, Gen. Syst. ii. 403 = elongata.
hebeclada, *DC. Cat. Hort. Monsp.* 73; *Prod.* ii. 461.—Afr. austr.
hecatophylla, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 242.—Afr. trop.
hemignosta, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 134.—Am. trop.
hemiteles, Benth. in Linnaea, xxvi. (1853-55) 619 = subcaerulea.
Hemu, Walp. Rep. v. 598 (sphalm. pro *A. Nemu*).
heteracantha, *Burch. Trav.* i. 389.—Afr. austr.
heterocarpa, Delile, Fl. Aeg. Illustr. 79 = Prosopis Stephaniana.
heteroclita, *Meissn. in Lehm. Pl. Preiss.* i. 18.—Austral.
heteroneura, *Benth. in Linnaea*, xxvi. (1853-55) 624.—Austral.

ACACIA:—
heterophylla, Hook. & Arn. Bot. Beech. Voy. 81 = A. Koa.
heterophylla, *Willd. Sp. Pl.* iv. 1054.—Ins. Mascar.
Hindsii, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 504.—Mexico.
hippuroides, *Heward, ex Benth. l. c.* 344.—Austral.
hirsuta, Schlcht. in Linnaea, xii. (1838) 572 = filicina.
hirta, *Boj. Hort. Maurit.* 115.—Ins. Maurit.
hirta, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 318 = Mimosa Galcottii.
hirta, Nutt. in Torr. & Gray, Fl. N. Am. i. 404 = filicina.
hirtella, *E. Mey. Comm. Pl. Afr. Austr.* 167.—Afr. austr.
hispidissima, DC. Prod. ii. 455 = pulchella.
hispidula, *Willd. Sp. Pl.* iv. 1054.—Austral.
Hoffmannseggii, *DC. Prod.* ii. 471.—Bras.
holocarpa, *Benth. Fl. Austral.* ii. 408.—Austral.
holosericea, *A. Cunn. ex G. Don, Gen. Syst.* ii. 407.—Austral.
homaloclada, *F. Muell. Fragm.* xi. 34.—Austral.
homalophylla, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 365.—Austral.
homomalla, Wendl. Acac. 49. t. 13 = glaucescens.
Hookeri, Meissn. in Lehm. Pl. Preiss. i. 11 = ericifolia.
Hooperiana, Zipp. ex Miq. Fl. Ind. Bat. i. 10 = concinna.
horrida, Span. in Linnaea, xv. (1841) 199 = pennata.
horrida, *Willd. Sp. Pl.* iv. 1082.—Afr. austr.
horridula, *Meissn. in Lehm. Pl. Preiss.* i. 9.—Austral.
hostilis, Mart. Reise, i. 555 = Mimosa hostilis.
Houstoni, Willd. Sp. Pl. iv. 1062 = Calliandra Houstoni.
Huegelii, *Benth. in Enum. Pl. Hueg.* 42.—Austral.
Humboldtii, Desv. in Ham. Prod. Fl. Ind. Occ. 60 = macracantha.
humifusa, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 382.—Austral.
humilis, Schlecht. in Linnaea, xii. (1838) 567 = Calliandra humilis.
Hunteri, *Oliver, in Hook. Ic. Pl.* t. 1350.—Arab.
hybrida, Lodd. Bot. Cab. t. 1342 = armata.
hypacantha, *Hoffmgg. ex Heynh. Nomencl.* i. 3, nomen.—Afr. austr.
idiomorpha, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 329.—Austral.
ignorata, *C. Koch, in Allg. Gartz.* (1858) 195.—Hab.?
imbricata, F. Muell. Fragm. i. 65 = lineata.
implexa, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 368.—Austral.
impressa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 133.—Austral.
impressa, Lindl. in Bot. Reg. t. 1115 = penninervis.
incana, Spreng. Syst. iii. 137 = Mimosa incana.
incrassata, Hook. Ic. Pl. t. 370 = decipiens.
incurva, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 325.—Austral.
indica, Desv. Journ. Bot. i. (1814) 69 = Farnesiana.
ingrata, *Benth. Fl. Austral.* ii. 331.—Austral.
insignis, Hoffmgg. Verz. Pfl. 159 = Albizzia lophantha.
insignis, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 313 = filicina.
insularis, *A. Rich. Ess. Fl. Cub.* 464.—Ind. occ.
insularum, *Guill. in Ann. Sc. Nat.* Sér. II. vii. (1837) 360 = Leucaena Forsteri.
intermedia, A. Cunn. ex Hook. Bot. Mag. t. 3203 = longifolia.
intermedia, *Pasq. in Atti R. Acad. Nap.* ix. (1882) 7.—Hab.?
intertexta, Sieber, ex DC. Prod. ii. 454 = longifolia.
Intsia, Willd. Sp. Pl. iv. 1091 = caesia.
intsioides, DC. Prod. ii. 464 = praec.
inundata, Mart. Reise, i. 555 = Pithecolobium multiflorum.
irrorata, Sieber, ex Spreng. Syst. iii. 141 = dealbata.
Isenbergiana, Schimp. ex A. Rich. Tent. Fl. Abyss. i. 236 = Albizzia Julibrissin.
iteaphylla, F. Mucll. ex Benth. in Linnaea, xxvi. (1853-55) 617 = neriifolia.
ixiophylla, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 364.—Austral.
Jacquemontii, *Benth. l. c.* 499.—Ind. or.
Jaubertiana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 363.—Madag.
javanica, *DC. Prod.* ii. 471.—Java.
Julibrissin, Willd. Sp. Pl. iv. 1065 = Albizzia Julibrissin.

ACACIA:—
Julibrissin, Sieber, ex Benth. in Trans. Linn. Soc. xxx. (1875) 529 = paniculata.
julifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 374.—Austral.
julifera, F. Muell. in Journ. Linn. Soc. iii. (1859) 139 = torulosa.
juliflora, Willd. Sp. Pl. iv. 1076 = Prosopis juliflora.
juncifolia, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 341.—Austral.
juniperina, *Willd. Sp. Pl.* iv. 1049.—Austral.
Jupunba, Willd. l. c. 1067 = Pithecolobium trapezifolium.
Jurema, *Mart. Reise*, ii. 788; *Syst. Mat. Med. Fl. Bras.* 53.—Bras.
Kalkora, G. Don, in Loud. Hort. Brit. 409 = Albizzia Julibrissin.
Kantuffa, Bruce, ex Steud. Nom. ed. I. 2 = Mimosa Kantuffa.
Karroo, Hayne, Arzneigew. x. t. 33 = horrida.
Kempeana, *F. Muell. in Melb. Chemist* (July, 1882).—Austral.
Kirkii, *Oliver, Fl. Trop. Afr.* ii. 350.—Afr. trop.
Koa, *A. Gray, Bot. U. S. Expl. Exped.* i. 480.—Ins. Sandvic.
Kraussiana, *Meissn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 515.—Afr. austr.
Kunthii, *Dehnh. Rivist. Nap.* i. 169; *ex Walp. Rep.* i. 924.—N. Granat.
laccrans, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 511.—Bras.
lachnophylla, *F. Muell. in Wing, South. Sc. Record* (July, 1882).—Austral.
lacustris, Desf. Cat. Hort. Par. ed. III. 301 = Neptunia oleracea.
laeta, *R. Br. in Salt, Abyss. App.* 65.—Afr. bor.; Arab.
laevigata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1059 = Prosopis juliflora.
laevis, Hort. ex Steud. Nom. ed. II. i. 6 = Gleditschia triacanthos.
Lahai, *Steud. & Hochst. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 506.—Abyss.
Lambertiana, D. Don, in Bot. Reg. t. 721 = Calliandra Lambertiana.
lanata, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 315.—Mexico.
Langsdorfii, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 521.—Bras.
lanigera, *A. Cunn. in Field, N. S. Wales*, 345.—Austral.
lanuginosa, Hort. ex Regel, in Gartenfl. (1854) 155 = pulchella?
laricina, *Meissn. in Lehm. Pl. Preiss.* i. 6.—Austral.
lasiocarpa, Benth. in Enum. Pl. Hueg. 43 = pulchella.
lasiopetala, *Oliver, Fl. Trop. Afr.* ii. 346.—Afr. trop.
lasiophylla, *Benth. in Trans. Linn. Soc.* xxx. (1875) 527.—Am. trop.
latescens, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 380.—Austral.
latifolia, *Benth. l. c.* 382.—Austral.
latifolia, Desf. Tabl. Hort. Par. ed. II. 207 = A. melanoxylon.
latifrons, Hort. ex Scem. & Schmidt, in Flora, xxvii. (1844) 495 = alata.
latipes, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 334.—Austral.
latisiliqua, Willd. Sp. Pl. iv. 1067 = Lysiloma latisiliqua.
latispinosa, Desf. Cat. Hort. Par. ed. III. 299 = Mimosa latispinosa.
Latrobei, Meissn. in Lehm. Pl. Preiss. i. 10 = acinacea.
latronum, *Willd. Sp. Pl.* iv. 1077.—Ind. or.
laurifolia, *Willd. l. c.* 1053.—Ins. Pacif.
Lawsoni, *Ait. ex Steud. Nom.* ed. II. i. 6.—Austral.
laxa, *Willd. Sp. Pl.* iv. 1069 = Calliandra laxa.
laxiflora, DC. Prod. ii. 455 = Pithecolobium laxiflorum.
laxophylla, vide *loxophylla*.
Lebbeck, Willd. Sp. Pl. iv. 1066 = Albizzia Lebbek.
lebbekoides, DC. Prod. ii. 467 = Albizzia lebbekoides.
Leichardtii, *Benth. Fl. Austral.* ii. 372.—Austral.
leiophylla, Benth. in Hook. Lond. Journ. Bot. i. (1842) 351 = Albizzia saligna.
lenticellata, F. Muell. in Journ. Linn. Soc. iii. (1859) 147 = Farnesiana.
lenticularis, *Buch.-Ham. in Wall. Cat.* n. 5244.—Ind. or.

ACACIA:—
lentiscifolia, *Desf. Tabl. Hort. Par.* ed. II. 208.—Mexico.
lentiscifolia, *A. Rich. Ess. Fl. Cub.* 469.—Cuba.
lepidota, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 205 = Mimosa Bonplandii.
leprosa, *Sieber, ex DC. Prod.* ii. 450.—Austral.
leptocarpa, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 376.—Austral.
leptoclada, *A. Cunn. ex Benth. l. c.* 385.—Austral.
leptoneura, *Benth. l. c.* 341.—Austral.
leptopetala, *Benth. in Linnaea*, xxvi. (1853–55) 619.—Austral.
leptophleba, F. Muell. in Journ. Linn. Soc. iii. (1859) 143 = aulacocarpa.
leptophylla, DC. Cat. Hort. Monsp. 74 = Farnesiana.
leptophylla, F. Muell. Fragm. xix. 9 = A. Bynoeana.
leptosperma, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 265.—Ind. occ.
leptospermoides, *Benth. in Linnaea*, xxvi. (1853–55) 626.—Austral.
leptostachya, *Benth. Fl. Austral.* ii. 406.—Austral.
leucacantha, Bert. ex Spreng. Syst. iii. 144 = tortuosa.
leucacantha, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 276.—Afr. trop.
leucadendron, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 374 = glaucescens.
leucocephala, Link, Enum. Hort. Berol. ii. 444 = Leucaena glauca.
leucolobia, *Sweet, Hort. Brit.* ed. II. 165.—Austral.
leucophloea, *Willd. Sp. Pl.* iv. 1083.—As. trop.
leucophylla, Colvill, ex Sweet, Hort. Brit. ed. I. 101 = holosericea.
leucophylla, Lindl. in Mitch. Three Exped. ii. 13 = pendula.
ligulata, *Ait. ex Steud. Nom.* ed. II. i. 6.—Austral.—Cf. seq.
ligulata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 362 = salicina.
ligustrina, *Meissn. in Lehm. Pl. Preiss.* ii. 203.—Austral.
limbata, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 145.—Austral.
linarioides, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 371.—Austral.
Lindleyi, *Meissn. in Lehm. Pl. Preiss.* i. 14.—Austral.
linearis, Desv. in Ham. Prod. Fl. Ind. Occ. 591 = Calliandra portoricensis.
linearis, *Sims, Bot. Mag.* t. 2156.—Austral.
lineata, *A. Cunn. ex G. Don, Gen. Syst.* ii. 403.—Austral.
lineolata, *Benth. ex Schlecht. in Linnaea*, xxvi. (1853–55) 626.—Austral.
lingvatouana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 362.—Madag.
linifolia, *Willd. Sp. Pl.* iv. 1051.—Austral.
litakunensis, *Burch. Trav.* ii. 452.—Afr. austr.
littoralis, *A. Rich. Ess. Fl. Cub.* 467.—Cuba.
lomatocarpa, DC. Prod. ii. 467 = Albizzia stipulata.
longifolia, Paxt. Mag. Bot. xii. (1846) 269 = dentifera.
longifolia, *Willd. Sp. Pl.* iv. 1052.—Austral.
longispicata, Benth. in Mitch. Journ. Trop. Austr. 298 = Cunninghamii.
longissima, Wendl. Comm. Acac. 45. t. 11 = linearis.
lophantha, Willd. Sp. Pl. iv. 1070 = Albizzia lophantha.
lophanthoides, DC. Prod. ii. 457 = villosa.
loxocarpa, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 377.—Austral.
loxophylla, *Benth. in Linnaea*, xxvi. (1853–55) 622.—Austral.
lucens, *Boj. Hort. Maurit.* 116.—Madag.
lucida, Baill. Hist. Pl. ii. 61 = Albizzia lucida.
Ludwigii, Ohlend, Verz. (1844) 74; Walp. Ann. ii. 453 = myrtifolia.
Luehmanni, *F. Muell. Fragm.* xi. 116.—Austral.
lunata, Hort. Prag. ex Steud. Nom. ed. II. i. 6 = falcinella.
lunata, Lodd. Bot. Cab. t. 324 = myrtifolia.
lunata, *Sieber, ex DC. Prod.* ii. 452.—Austral.
lutea, Leavenw. in Am. Journ. Sc. vii. (1824) 61 = Neptunia lutea.
lycopodifolia, *A. Cunn. in Hook. Ic. Pl.* t. 172.—Austral.
lycopodioides, Desv. Journ. Bot. iii. (1814) 69 = Neptunia plena.
lysiploia, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 137.—Austral.

ACACIA:—
macocentra, Walp. Rep. i. 876. in syn.—sphalm.—Cf. malacocentra.
macracantha, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1080.—Am. austr.
macracanthoides, Bert. in DC. Prod. ii. 463 = macracantha.
macradenia, *Benth. in Mitch. Trop. Austral.* 360.—Austral.
macrocentra, Walp. Rep. iv. 200.—Cf. malacocentra.
macroloba, Willd. Sp. Pl. iv. 1054 = Pentaclethra filamentosa.
macrophylla, Bunge, Enum. Pl. Chin. Bor. 20 = Albizzia Lebbek.
macrostachya, *Reichb. ex G. Don, Gen. Syst.* ii. 411.—Senegal.
Magdalenae, Bert. ex DC. Prod. ii. 455 = Calliandra Magdalenae.
Maitlandi, F. Muell. Fragm. iii. 46 = patens.
malacocentra, Mart. Herb. Fl. Bras. 106 = Mimosa malacocentra.
malacophylla, *Benth. in A. Gray, Pl. Wright.* i. 64.—Mexico.
malacophylla, Steud. ex A. Rich. Tent. Fl. Abyss. i. 235 = Albizzia pallida.
mammifera, *Schlecht. in Linnaea*, xii. (1838) 563.—Mexico.
Mangium, *Willd. Sp. Pl.* iv. 1053.—Ins. Molucc.
marginata, Buch.-Ham. in Wall. Cat. n. 5243 = Albizzia stipulata.
marginata, Steud. Nom. ed. I. 3 = Albizzia odoratissima.
marginata, R. Br. in Ait. Hort. Kew. ed. II. iii. 462 = myrtifolia.
maritima, Benth. in Hook. Lond. Journ. Bot. i. (1842) 384 = discolor.
Martii, *Benth. l. c.* 519.—Bras.
martinicensis, Presl, Bot. Bemerk. 65 = paniculata.
maschalocephala, *Griseb. Cat. Pl. Cub.* 82.—Cuba.
mauroceana, *DC. Cat. Hort. Monsp.* 74.—Marocco.
media, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 316 = Calliandra angelina.
megaladena, Desv. Journ. Bot. iii. (1814) 69 = pennata.
megalantha, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 143.—Austral.
megaphylla, F. Muell. l. c. 117 = anceps.
Meissneri, *Lehm. Del. Sem. Hort. Hamb.* (1842).—Austral.
melanoceras, *Beurling, in Vet. Akad. Handl. Stockh.* (1854) 53.—Panama.
melanochaetes, Zoll. in Nat. en Geneesk. Arch. Neel. Indie, iii. (1846) 80 = leucophloea.
melanoxylon, *R. Br. in Ait. Hort. Kew.* ed. II. v. 462.—Austral.
mellifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 507.—Abyss.
metrosideriflora, Schlecht. in Linnaea, xii. (1838) 567. = Calliandra Houstoni.
metrosiderosifolia, Hort. ex Steud. Nom. ed. II. i. 6 = dodonaeifolia.
micracantha, A. Dietr. in Otto & Dietr. Allg. Gartenz. i. (1833) 83 = armata.
micracantha, Desv. Journ. Bot. iii. (1814) 69 = Pithecolobium parvifolium.
micradena, Benth. in Hook. Lond. Journ. Bot. i. (1842) 518 = adhaerens.
micrantha, *Benth. in Trans. Linn. Soc.* xxx. (1875) 526.—Mexico.
micrantha, Desv. in Ham. Prod. Fl. Ind. Occ. 59 = seq.
microcantha, G. Don, Gen. Syst. ii. 476 = Pithecolobium parvifolium.
microbotrya, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 353.—Austral.
microcarpa, *F. Muell. 2nd Gen. Report*, 7.—Austral.
microcephala, A. Rich. Ess. Fl. Cub. 469 = paniculata.
microcephala, R. Grah. in Wall. Cat. n. 5263 = leucophloea.
microcephala, Macfad. Fl. Jamaic. i. 316 = macracantha.
microneura, *Meissn. in Lehm. Pl. Preiss.* i. 19.—Austral.
microphylla, Roxb. ex Steud. Nom. ed. II. i. 6 = Albizzia myriophylla.
microphylla, Willd. Sp. Pl. iv. 1073 = Piptadenia peregrina.
Miersii, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 522.—Bras.

ACACIA:—
Mikanii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 526.—Am. trop.
minutifolia, *F. Muell. Fragm.* viii. 243.—Austral.
Mirbeli, *Dehnh. Rivist. Nap.* i. 168; *ex Walp. Rep.* i. 922.—Austral.
misera, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 275.—Afr. trop.
Mitchelli, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 387.—Austral.
modesta, *Wall. Pl. As. Rar.* ii. 27. t. 130.—Ind. or.
moesta, Lindl. Bot. Reg. (1846) t. 67 = verticillata.
mollicula, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 313.—Mexico.
mollis, Wall. Pl. As. Rar. ii. 76. t. 177 = Albizzia Julibrissin.
mollissima, Willd. Enum. Hort. Berol. 1053 = decurrens.
monacantha, *Willd. l. c.* 1056.—Bras.
moniliformis, *Griseb. in Goett. Abh.* xix. (1874) 136.—Reg. Argent.
montana, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 360.—Austral.
montana, Jungh. in Nat. en Geneesk. Arch. Neerl. Indie, ii. (1845) 35 = Albizzia montana.
mossambecensis, Bolle, in Peters, Reise Mossamb. Bot. i. 15 = albida.
mucronata, Willd. Enum. Hort. Berol. Suppl. 68 = longifolia.
Muelleri, Benth. in Linnaea, xxvi. (1853–55) 603 = anceps.
multiflora, H. B. & K. Nov. Gen. et Sp. vi. 277 = Pithecolobium multiflorum.
multiflora, Spreng. Syst. iii. 142 = polyantha.
multifoliolata, *Bertol. Misc. Bot.* xvi. 26.—Ins. S. Doming.
multijuga, *Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 105.—Afr. austr.
multinervia, DC. Mém. Lég. xii. 445; Prod. ii. 250 = lanigera.
multispicata, *Benth. Fl. Austral.* ii. 400.—Austral.
muricata, *Willd. Sp. Pl.* iv. 1058.—Ind. occ.
Murrayana, *F. Muell. ex Benth. Fl. Austral.* ii. 370.—Austral.
myriadenia, Bert. ex Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 359 = Serianthes myriadena.
myriobotrya, Meissn. in Lehm. Pl. Preiss. i. 15 = microbotrya.
myriophylla, R. Grah. in Wall. Cat. n. 5242 = Albizzia myriophylla.
myrtifolia, *Willd. Sp. Pl.* iv. 1054.—Austral.
natalitia, *E. Mey. Comm. Pl. Afr. Austr.* 167.—Afr. austr.
Nebroeb, *Baill. Adansonia*, iv. (1863–64) 117.—Afr. trop.
Nellyrenza, R. Grah. in Wall. Cat. n. 5240 = amara.
nematophylla, F. Muell. ex Benth. in Linnaea, xxvi. (1853–55) 612 = calamifolia.
Nemu, Willd. Sp. Pl. iv. 1065 = Albizzia Julibrissin.
neriifolia, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 357.—Austral.
Nernstii, F. Muell. Fragm. iv. 3 = amblygona.
nervosa, *DC. Mém. Lég.* xii. 444.—Austral.
Neuciana, Buckl. ex Benth. in Trans. Linn. Soc. xxx. (1875) 592 sphalm. = Pithecolobium brevifolium.
neurocarpa, A. Cunn. in Hook. Ic. Pl. t. 168 = holosericea.
nigra, Clos, in C. Gay, Fl. Chil. ii. 253 = Calliandra chilensis.
nigrescens, *Oliver, Fl. Trop. Afr.* ii. 340.—Afr. trop.
nigricans, *R. Br. in Ait. Hort. Kew.* ed. II. v. 465.—Austral.
nilotica, Delile, Fl. Aegypt. Illustr. 79; Desf. Tabl. Hort. Par. ed. II. 208 = arabica.
Niopo, H. B. & K. Nov. Gen. et Sp. vi. 282 = Piptadenia peregrina.
nitida, Willd. Sp. Pl. iv. 1086 = Pithecolobium umbellatum.
nitidula, *Benth. Fl. Austral.* ii. 381.—Austral.
nodiflora, *Benth. in Linnaea*, xxvi. (1853–55) 621.—Austral.
nodosa, Willd. Sp. Pl. iv. 1016 = Cassia bacillaris.
notabilis, *F. Muell. Fragm.* i. 6.—Austral.
nubica, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 498.—Nubia.
nudiflora, *Willd. Sp. Pl.* iv. 1058.—Ind. occ.
Nuceiana, Buckl. in Proc. Ac. Sc. Philad. '1861' (1862) 453 = Pithecolobium brevifolium.
nutans, Spreng. ex DC. Prod. ii. 429 = Mimosa leiocarpa?
nyssophylla, F. Muell. Fragm. iv. 4 = colletioides.

ACACIA:—
obliqua, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 334.—Austral.
obliqua, Desv. Journ. Bot. iii. (1814) 67 = suavcolens.
obliquifolia, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 317 = Pithecolobium dulce.
obovata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 329.—Austral.
obscura, *A. DC. Not.* vi. *Pl. Rar. Jard. Gen.* 23. t. 3.—Austral.
obtusa, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1087 = macraeantha.
obtusata. *Sieber, ex DC. Prod.* ii. 453.—Austral.
obtusifolia, A. Cunn. in Field, New South Wales, 345 = longifolia.
odorata, Desv. Journ. Bot. iii. (1814) 67 = suaveolens.
odoratissima, Buch.-Ham. ex Wall. Cat. n. 5233 = elata.
odoratissima, Willd. Sp. Pl. iv. 1063 = Albizzia odoratissima.
offiicnalis, *Dehnh. Rivist. Nap.* i. 170; *ex Walp. Rep.* i. 923.—Tasman.
Oldfieldii, F. Muell. Fragm. iv. 17 = acuminata.
oleaefolia, A. Cunn. ex G. Don, Gen. Syst. ii. 405 = lunata.
oligoneura. *F. Muell. in Journ. Linn. Soc.* iii. (1859) 139.—Austral.
oligophylla, *Hoffmgg. Verz. Pfl. Nachtr.* i. 201.—Hab.?
Oliveri, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 274.—Afr. trop.
omalophylla = homalophylla.
oncinocarpa, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 378.—Austral.
oncinophylla, *Benth. in Lindl. Swan Riv. App.* 15.—Austral.
oraria, *F. Muell. Fragm.* xi. 66.—Austral.
ornithophora, Sweet, Fl. Austral. t. 24 = armata.
orthocarpa, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 136 = xylocarpa.
Oswaldi, *F. Muell. Pl. Vict.* ii. 27.—Austral.
Ouyrarema, *DC. Prod.* ii. 469.—Guiana.
ovoidea, Benth. in Hook. Lond. Journ. Bot. i. (1842) 339 = verticillata.
Oxycedrus, *Sieber, ex DC. Prod.* ii. 453.—Austral.
oxyclada, *F. Muell. ex Benth. Fl. Austral.* ii. 341.—Austral.
oxyphylla, R. Grah. in Wall. Cat. n. 5252 = caesia.
pachycarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 139.—Austral.
pachyloma, Mart. Herb. Fl. Bras. n. 1108 = Calliandra parviflora.
pallida, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 147.—Austral.
pallida, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1059 = Prosopis juliflora.
Palmeri, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 350.—Am. bor. occ.
paludosa, Blume, ex Miq. Fl. Ind. Bat. i. 14 = pennata.
paniculaeflora, Steud. in Flora, xxvi. II. (1843) 760 = Mimosa myriadenia.
paniculata, *Willd. Sp. Pl.* iv. 1074.—Bras.
papulaeformis, Walp. Rep. i. 898 = buxifolia.
papyrocarpa, *Benth. Fl. Austral.* ii. 338.—Austral.
paradoxa, DC. Cat. Hort. Monsp. 74 = armata.
parvifolia, Hort. ex Steud. Nom. ed. II. i. 7 = guttata.
parvifolia, Willd. Sp. Pl. iv. 1086 = Pithecolobium parvifolium.
patens, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 120.—Austral.
patula, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 1055.—Am. austr.
pauciflora, *A. Rich. Ess. Fl. Cub.* 461.—Cuba.
Pawlikowskyana, Ohlend. in Neue Allg. Gartenz. (1845) 369 = myrtifolia.
pectinata, Hoffmgg. Verz. Pfl. Nachtr. i. 201 = Hoffmannseggiana.
pectinata, H. B. & K. Nov. Gen. et Sp. vi. 282 = Parkia pectinata.
pedicellata, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 522.—Bras.
pedunculata, Roxb. ex Steud. Nom. ed. II. i. 7 = Parkia biglandulosa?
pedunculata, Willd. Sp. Pl. iv. 1084 = Farnesiana.
pellacantha, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 19 = macracantha.
pendula, *A. Cunn. ex G. Don, Gen. Syst.* ii. 404.—Austral.
penicillifera, Lag. Gen. et Sp. Nov. 16 = filicina.

ACACIA:—
pennata, *Willd. Sp. Pl.* iv. 1090.—As. et Afr. trop.
pennatula, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 390.—Mexico.
penninervis, *Sieber, ex DC. Prod.* ii. 452.—Austral.
pennivenia, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 404.—Ins. Socotra.
pentadenia, *Lindl. in Bot. Reg.* t. 1521.—Austral.
pentaedra, Regel, Gartenfl. (1852) 228, t. 23 = extensa.
pentagona, Hook. f. in Hook. Nig. Fl. 331 = pennata.
pentaptera, Welw. Apont. 584 = pennata.
peregrina, Kunth, Mimos. Humb. 96. t. 30 = Piptadenia colubrina.
peregrina, Willd. Sp. Pl. iv. 1073 = Piptadenia peregrina.
Perrottetii, Steud. Nom. ed. II. i. 7 = tortilis.
persica, Sterl. Hort. Nymph. ex Steud. Nom. ed. II. i. 7 = Prosopis Stephaniana.
personioides, *Ait. ex Steud. Nom.* ed. II. i. 7.—Austral.
Pervillei, *Benth. in Trans. Linn. Soc.* xxx. (1875) 521.—Madag.
peruviana, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 1056.—Peruv.
Petersiana, *Bolle, in Peters, Reise Mossamb. Bot.* i. 4.—Mossamb.
petiolaris, Lehm. Novit. Pl. Hort. Hamb. (1851) 7; Linnaea, xxv. (1852) 306 = pycnantha.
Peuce, *F. Muell. Fragm.* iii. 151.—Austral.
Pharnesiana, Risso, Fl. Nicc, 166 = Farnesiana.
philippinarum, Benth. in Hook. Lond. Journ. Bot. i. (1842) 514 = concinna.
phlebocarpa, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 119.—Austral.
phlebophylla, F. Muell. 1st Gen. Rep. 12 = longifolia.
piauhiensis, *Benth. in Trans. Linn. Soc.* xxx. (1875) 523.—Bras.
piligera, A. Cunn. in Hook. Bot. Mag. sub t. 3394 = undulaefolia.
pilosa, *Benth. in Linnaea*, xxvi. (1853–55) 607.—Austral.
pilosa, Bert. ex DC. Prod. ii. 455 = Calliandra haematomma?
pinifolia, Benth. in Mitch. Trop. Austral. 342 = juncifolia.
pinnata, Dalz. & Gibs. Bomb. Fl. 87 = pennata.
pinnata, Link, Enum. Hort. Berol. ii. 446 = tamarindifolia.
pityoides, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 135.—Austral.
plagiophylla, F. Muell. l. c. 131 = undulaefolia.
plagiophylla, Spreng. Syst. iii. 135 = falcata.
planifrons, *Wight & Arn. Prod.* 276.—Ind. or.
platyacantha, *Schlecht. in Linnaea*, xii. (1838) 565.—Mexico.
platycarpa, F. Muell. in Journ. Linn. Soc. iii. (1859) 145 = sericata.
platyloba, *Spreng. Syst.* iii. 144.—N. Granat.
platyloba, Bert. ex DC. Prod. ii. 472 = Pithecolobium sericiflorum.
platyphylla, *Sweet, Hort. Brit.* ed. I. 102, nomen.—Austral.
platyptera, Lindl. in Bot. Reg. (1841) Misc. 3 = alata.
plectocarpa, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 375.—Austral.
plumosa, *Lowe, in Bot. Mag.* t. 3366.—Bras.
pluricapitata, *Steud. Nom.* ed. II. i. 7.—As. trop.
podalyriaefolia, *A. Cunn. in G. Don, Gen. Syst.* ii. 405.—Austral.
polyacantha, Willd. Sp. Pl. iv. 1079 = A. Catechu.
polyantha, Spreng. ex Benth. in Trans. Linn. Soc. xxx. (1875) 591 = Pithecolobium multiflorum.
polyantha, Zipp. ex Span. in Linnaea, xv. (1841) 199 = juncifolia.
polybotrya, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 384.—Austral.
polycephala, DC. Prod. ii. 473 = concinna.
polycephala, R. Grah. in Wall. Cat. n. 5255 = pluricapitata.
polymorpha, *Pasq. Cat. Ort. Bot. Nap.* 1.—Austral.
polyphylla, Clos, in C. Gay, Fl. Chil. ii. 254 = Lysiloma polyphylla.
polyphylla, *DC. Cat. Hort. Monsp.* 74; *Prod.* ii. 469.—N. Granat.
polystachya, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 376.—Austral.
portoricensis, Willd. Sp. Pl. iv. 1069 = Calliandra portoricensis.
praecox, *Griseb. in Goett. Abh.* xix. 136.—Reg. Argent.

ACACIA:—
praelongata, *F. Muell. in Melb. Chemist*, iii. (Aug. 1883).—Austral.
pravifolia, F. Muell. Fragm. i. 4 = sublanata.
pravissima, *F. Muell. 1st Gen. Report*, 12.—Austral.
Preissiana, Lehm. Del. Sem. Hort. Hamb. (1842) = obscura.
prensans, Lowe, in Bot. Mag. t. 3408 = pennata.
prismatica, *Hoffmgg. Verz. Pfl.* 159.—Hab.?
procera, Willd. Sp. Pl. iv. 1063 = Albizzia procera.
prominens, *A. Cunn. ex G. Don, Gen. Syst.* ii. 406.—Austral.
prona, *Noronha, in Verh. Batav. Gen.* v. (1790) Art. iv. 5, nomen.—Java.
propinqua, A. Rich. Ess. Fl. Cub. i. 466 = Albizzia Lebbek.
prosopoides, Moç. & Sesse, ex DC. Prod. ii. 460 = Piptadenia patens.
prosopoma, *Schnyder, in Anal. Soc. Cient. Argent.* iii. (1877) 152.—Reg. Argent.
prostrata, Lodd. Bot. Cab. t. 631 = diffusa.
pruinescens, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 296, 298.—Burma.
pruinosa, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 383.—Austral.
prunifolia, C. Mucll. in Walp. Ann. vii. 937, sphalm.—Cf. pravifolia.
pseudoarabica, *Blume, ex Miq. in Fl. Ind. Bat.* i. 8.—N. Guinea.
pseudointsia, Miq. l. c. 12 = cacsia.
pseudotrichodes, *DC. Prod.* ii. 466.—Ins. S. Doming.
psilostachya, DC. l. c. 457 = Piptadenia psilostachya.
Psoralea, DC. l. c. 464 = Mimosa Psoralea.
pteridifolia, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 521.—Am. trop.
pterigoidea, Secm. in Verh. Gartenb. (1846) 11 = microbotrya.
pterocarpa, F. Muell. in Journ. Linn. Soc. iii. (1859) 134 = excelsa.
pterocarpa, Steud. Nom. ed. I. 3 = Gagnebina tamariscina.
pteroclada, F. Muell. Fragm. iv. 3 = trigonophylla.
pterophylla, Hoffmgg. Verz. Pfl. (1824) 202 = pennata.
pterygocarpa, Hochst. ex Benth. in Hook. Lond. Journ. Bot. v. (1846) 96 = nubica.
ptychophylla, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 142.—Austral.
pubescens, *R. Br. in Ait. Hort. Kew.* ed. II. v. 467.—Austral.
pubescens, *Schlecht. in Linnaea*, xii. (1838) 563.—Mexico.
pugioniformis, *Wendl. Comm. Acac.* 38. t. 9.—Austral.
pugioniformis, Wendl. in Flora, ii. (1819) 139 = juniperina.
pulchella, *R. Br. in Ait. Hort. Kew.* ed. II. v. 464.—Austral.
pulcherrima, Willd. Sp. Pl. iv. 1061 = Stryphnodendron floribundum.
pulverulenta, A. Cunn. ex Benth. in Hook. Lond. Journ. Bot. i. (1842) 342 = calamifolia.
pulverulenta, Schlecht. in Linnaea, xii. (1838) 571 = Leucaena pulverulenta.
punctata, Desf. Cat. Hort. Par. ed. III. 300 = Neptunia plena.
punctata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1084 = macracantha.
pungens, Spreng. Syst. iii. 134 = juniperina.
purpurascens, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 277.—Afr. trop.
purpurea, *Bolle, in Peters, Reise Mossamb. Bot.* 6.—Afr. trop.
purpurea, Steud. Nom. ed. II. i. 7 = Inga tergemina.
pycnantha, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 351.—Austral.
pycnophylla, *Benth. Fl. Austral.* ii. 368.—Austral.
pycnostachya, *F. Muell. Pl. Vict.* ii. 33.—Austral
pyrifolia, *DC. Mém. Lég.* xii. 447.—Austral.
quadrangularis, Link, Enum. Hort. Berol. ii. 445 = Calliandra tetragona.
quadriglandulosa, Mart. in Flora, xx. (1838); Beibl. ii. 110 = riparia.
quadrilateralis, DC. Prod. ii. 451 = pugioniformis.
quadrilateralis, Decne. Herb. Tim. 177 = juncifolia?
quadrimarginea, *F. Muell. Fragm.* x. 31.—Austral.
quadrisulcata, *F. Muell. l. c.* iii. 127.—Austral.
Raddiana, Savi, Sopra Alc. Acac. Egiz. 1 = tortilis.
ramosissima, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 356 ex parte.—Austral.

ACACIA:—
ramosissima, Benth. in Hook. Lond. Journ. Bot. i. (1842) 356 ex parte = Harveyi.
reclinata, F. Muell. 1st Gen. Rep. 12 = leprosa.
recurva, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 519.—Bras.
reficiens, *Wawra & Peyr. Sert. Benguel.* 15.—Afr. trop.
reniformis, *Benth. in Hook. Ic. Pl.* t. 1165.—Mexico.
restiacea, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 323.—Austral.
reticulata, *Willd. Sp. Pl.* iv. 1056.—Afr. austr.
retinervis, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 379.—Austral.
retinodes, Schlecht. in Linnaea, xx. (1847) 664 = neriifolia.
retivenea, *F. Muell. Fragm.* iii. 128.—Austral.
retrorsa, Meissn. in Bot. Zeit. (1855) 10 = sphacelata.
revoluta, Kunth, Mimos. Humb. 84. t. 26 = Mimosa revoluta.
rhigiophylla, *F. Muell. ex Schlecht. Benth. in Linnaea*, xxvi. (1853–55) 611.—Austral.
rhodacantha, *Desf. Tabl. Hort. Par.* ed. II. 208.—Hab.?
rhombifolia, *Desf. ex Benth. in Trans. Linn. Soc.* xxx. (1875) 637.—Hab.?
rhombifolia, *Poir. ex Steud. Nom.* ed. I. 3.—Ins. Trinit.
Riceana, *Hensl. in Maund, Botanist*, t. 135.—Tasman.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 482. t. 53.—Ins. Fiji.
rigens, *A. Cunn. ex G. Don, Gen. Syst.* ii. 403.—Austral.
rigidula, Benth. in Hook. Lond. Journ. Bot. i. (1842) 504 = amentacea.
riparia, Bert. ex Spreng. Syst. iii. 142 = polyphylla.
riparia, *H. B. & K. Nov. Gen. et Sp.* vi. 276.—N. Granat.
robusta, *Burch. Trav.* ii. 442.—Afr. austr.
Roemeriana, *Scheele, in Linnaea*, xxi. (1848) 456.—N. Mexico.
Rohriana, DC. Prod. ii. 457 = nudiflora.
rostellifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 356.—Austral.
rostellifera, Seem. Eingef. Acac. t. 2 = microbotrya.
rostrata, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 1054 (err. typ. 1060) = Lysiloma rostrata.
rotundata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 521.—Am. trop.
rotundifolia, Hook. Bot. Mag. t. 4041 = obliqua.
Rovumae, *Oliver, Fl. Trop. Afr.* ii. 353.—Afr. trop.
Roxburghii, Kostel. Allg. Med. pharm. Flora, iv. 1367 = Albizzia myriophylla.
Roxburghii, Wight & Arn. Prod. i. 276 = planifrons et eburnea?
rubescens, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 315 = Calliandra laxa.
rubida, *A. Cunn. in Field, New South Wales*, 344.—Austral.
rufa, Hassk. Cat. Pl. Hort. Bog. 290 = Albizzia rufa.
rugata, Buch.-Ham. in Wall. Cat. n. 5251 = concinna.
runciformis, A. Cunn. ex G. Don, Gen. Syst. ii. 404 = lineata.
rupestris, Stocks, ex Boiss. Fl. Orient. ii. 638 = A. Senegal.
rupicola, *F. Muell. ex Benth. in Linnaea*, xxvi. (1853–55) 610.—Austral.
ruscifolia, A. Cunn. ex G. Don, Gen. Syst. ii. 407 = verticillata.
rutaefolia, *Ait. ex Steud. Nom.* ed. II. i. 8.—Austral.
rutaefolia, Link, Enum. Hort. Berol. ii. 444 = nigricans.
Sabeana, Buckl. in Proc. Ac. Sc. Philad. '1861' (1862) 453 = Leucaena retusa.
saccharata, Benth. in Hook. Lond. Journ. Bot. i. (1842) 505 = albida.
salicina, *Lindl. in Mitch. Three Exped.* ii. 20.—Austral.
saligna, *Wendl. Comm. Acac.* 26.—Austral.
salinarum, DC. Prod. ii. 456 = Prosopis juliflora.
Saltiana, Steud. Nom. ed. II. i. 8 = A. Lahai.
saltuum, Jungh. in Bot. Zeit. iii. (1845) 703 = Albizzia montana.
Sandra, Bedd. Fl. Sylv. t. 50 = A. Sundra.
sanguinea, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 245 = venosa.
sapindoides, A. Cunn. ex Sweet, Hort. Brit. ed. III. 198 = Pithecolobium pruinosum.
saponaria, Heyne, ex. Benth. in Trans. Linn. Soc. xxx. (1875) 531 = concinna.
sarmentosa, Desv. Journ. Bot. iii. (1814) 70 = caesia.
sarmentosa, Griseb. Fl. Brit. W. Ind. 221 = riparia.
Sassa, *Baill. Hist. Pl.* ii. 47.—Madag.

ACACIA:—
scabra, *Benth. in Linnaea*, xxvi. (1853–55) 605.—Austral.
scalpelliformis, *Meissn. in Lehm. Pl. Preiss.* ii. 200.—Austral.
scandens, Benth. in Hook. Lond. Journ. Bot. i. (1842) 517 = plumosa.
scandens, Willd. Sp. Pl. iv. 1057 = Entada scandens.
scandens, Willd. Enum. Hort. Berol. 1057 = polyphylla.
scapuliformis, A. Cunn. in G. Don, Gen. Syst. ii. 405 = cultriformis.
schinoides, Benth. in Hook. Lond. Journ. Bot. i. (1842) 382 = pruinosa.
Schottii, *Torr. in Bot. Mex. Bound.* 62.—Am. bor.
scirpifolia, *Meissn. in Bot. Zeit.* (1855) 10.—Austral.
sclerophylla, *Lindl. in Mitch. Three Exped.* ii. 139.—Austral.
sclerosperma, *F. Muell. in Wing, South. Sc. Record*, (July, 1882).—Austral.
scleroxyla, *Tuss. Fl. Antill.* i. t. 21.—Ind. occ.
secundiflora, Juss. ex DC. Prod. ii. 425 = Entada polystachya.
Seeressa, Roxb. ex Steud. Nom. ed. II. i. 8 = Albizzia Lebbek.
selenicarpa, Desv. in Ham. Prod. Fl. Ind. Occ. 60 = albida.
Selloi, Spreng. Syst. iii. 137 = Calliandra brevipes.
semicordata, *Roxb. ex Steud. Nom.* ed. II. i. 8.—Ind. or.
semispinosa, Willd. ex Steud. l. c. = Mimosa semispinosa.
semitriptera, *Mart. ex Steud. l. c.*—Bras.
semiverticillata, Knowl. & Westc. Fl. Cab. ii. (1838) 27 = verticillata.
Senegal, *Willd. Sp. Pl.* iv. 1077.—Geront. trop.
Sentis, *F. Muell. 2nd Gen. Report*, 7 (1854).—Austral.
sericata, Ait. ex Steud. Nom. ed. II. i. 8 = seq.?
sericata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 380.—Austral.
sericea, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 311 = acatlensis.
sericocephala, Fenzl, in Flora, xxvii. (1844) 312 = Albizzia amara.
sericophylla, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 122 = stenophylla.
Serra, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 519.—Bras.
sertiformis, A. Cunn. in Bot. Mag. sub t. 3394 = undulaefolia.
sessiliceps, *F. Muell. in Melb. Chemist* (July, 1882).—Austral.
sessilis, Benth. in Hook. Lond. Journ. Bot. i. (1842) 336 = sphacelata.
setigera, Hook. Ic. Pl. t. 361 = undulaefolia.
setosa, *Dehnh. Rivist. Nap.* i. 169; *ex Walp. Rep.* i. 924.—Hab.?
setosa, *Spreng. Syst.* iii. 137.—Bras.
setulifera, *Benth. in Linnaea*, xxvi. (1853–55) 625.—Austral.
Seyal, *Delile, Fl. Aegypte*, 286. t. 52. f. 2.—Aegypt.
Shuttleworthii, *Meissn. in Lehm. Pl. Preiss.* i. 7.—Austral.
siculiformis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 337.—Austral.
Sieberi, Tausch, ex Steud. Nom. ed. II. i. 8 = crassiuscula.
Sieberiana, *DC. Prod.* ii. 453.—Senegal.
Sieberiana, Scheele, in Linnaea, xxvii. (1843) 337 = discolor.
Sieberiana, Tausch, in Flora, xix. (1836) 420 = crassiuscula.
signata, *F. Muell. Fragm.* iv. 7.—Austral.
Siliquastrum, Lag. Gen. et Sp. Nov. 16 = Prosopis juliflora.
similis, *Zoll. & Morr. in Nat. en Geneesk. Arch. N. Ind.* iii. (1846) 81.—Malaya.
Simsii, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 368.—Austral.
Sing, Guill. & Perr. Tent. Fl. Seneg. i. 251 = Sieberiana.
Sirissa, Buch.-Ham. in Wall. Cat. n. 5265 = Albizzia Lebbek.
smilacifolia, Field. & Gardn. Sert. Pl. t. 3 = urophylla.
Smithiana, Roxb. ex Wall. Cat. n. 5237 = Albizzia stipulata.
Sobeana, *Buckl. Proc. Acad. Nat. Sc. Philad.* (1861) 453.—Am. bor.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 511.—Ins. Socotra.

ACACIA:—
Solandri, *Benth. Fl. Austral.* ii. 406.—Austral.
somalensis, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 274.—Afr. trop.
somnians, Sterler, ex Steud. Nom. ed. II. i. 8 = Mimosa somnians.
Sophorae, R. Br. in Ait. Hort. Kew. ed. II. v. 462 = longifolia.
spadicigera, *Cham. & Schlecht. in Linnaea*, v. (1830) 594.—Jamaica.
spathulata, *F. Muell. ex Benth. Fl. Austral.* ii. 356.—Austral.
spathulata, Tausch, in Flora, xix. (1836) 420 = longifolia.
speciosa, Hort. ex Steud. Nom. ed. II. i. 8 = Albizzia lophantha.
speciosa, Willd. Sp. Pl. iv. 1066 = Albizzia Lebbek.
spectabilis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 383.—Austral.
sphacelata, *Benth. l. c.* 338.—Austral.
sphaerocephala, *Cham. & Schlecht. in Linnaea*, v. (1830) 594.—Mexico.
spinescens, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 323.—Austral.
Spini, *Balb. in Spin. Cat. Suppl.* (1823) 8.—Ins. Guadaloup.
spinosa, E. Mey. Comm. Pl. Afr. Austr. i. 170 = Dichrostachys nutans.
spinosissima, *Benth. ex Schlecht. in Linnaea*, xxvi. (1853–55) 621.—Austral.
spinulosa, *Sweet, ex Steud. Nom.* ed. II. i. 8.—Austral.
spirocarpa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 239.—Afr. trop.
spirorbis, *Labill. Sert. Austr. Caled.* 69. t. 69.—N. Caled.
spondylophylla, *F. Muell. Fragm.* viii. 243.—Austral.
Sprengelii, G. Don, Gen. Syst. ii. 410 = Berteriana.
Sprengelii, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 205 = Mimosa incana.
squamata, *Lindl. Swan Riv. App.* 15.—Austral.
squamata, Morr. in Ann. Soc. Gand, iii. (1847) 209. t. 134 = restiacea.
squarrosa, Mart. ex Benth. in Hook. Lond. Journ. Bot. iii. (1844) 104 = Calliandra squarrosa.
stellata, *Willd. Sp. Pl.* iv. 1078.—Arab.
stenocarpa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 238.—Abyss.
stenophylla, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 386.—Austral.
stenoptera, *Benth. l. c.* 325.—Austral.
stenostachya, Desv. ex Ham. Prod. Fl. Ind. Occ. 59 = Piptadenia psilostachya.
Stephaniana, Willd. Sp. Pl. iv. 1088 = Prosopis Stephaniana.
stereophylla, *Meissn. in Lehm. Pl. Preiss.* ii. 203.—Austral.
stigmatophylla, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 377.—Austral.
stipellata, Schlecht. in Linnaea, xii. (1838) 574 = filicina.
stipulata, DC. Prod. ii. 469 = Albizzia stipulata.
stipuligera, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 144.—Austral.
stipulosa, *F. Muell. l. c.* 119.—Austral.
stolonifera, Burch. Trav. ii. 241 = hebeclada.
striata, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 1089.—Am. austr.
striatula, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 336.—Austral.
stricta, *Willd. Sp. Pl.* iv. 1052.—Austral.
strigosa, Lindl. in Mitch. Three Exped. ii. 185 = aspera.
strigosa, *Link, Enum. Hort. Berol.* ii. 444.—Austral.
strigosa, Spreng. Syst. iii. 137 = strigulosa.
strigulosa, *Steud. Nom.* ed. II. i. 8.—Peruv.
strombulifera, Willd. Sp. Pl. iv. 1055 = Prosopis strombulifera.
strongylophylla, *F. Muell. Fragm.* viii. 226.—Austral.
Stuartiana, F. Muell. ex Benth. in Linnaea, xxvi. (1853–55) 609 = siculiformis.
suaresensis, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 363.—Madag.
suaveolens, *Willd. Sp. Pl.* iv. 1050.—Austral.
subalata, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 276.—Afr. trop.
subbinervia, Meissn. ex Lehm. Plant. Preiss. i. 16 = rostellifera.
subcaerulea, *Lindl. in Bot. Reg.* t. 1075.—Austral.

ACACIA:—
subdimidiata, Splitgerb. in Hoev. & De. Vr. Tijdschr. ix. (1842) 112 = Pithecolobium multiflorum.
suberosa, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 499.—Austral.
subfalcata, Meissn. in Bot. Zeit. (1855) 11 = microbotrya.
subinermis, Berr. ex DC. Prod. ii. 463 = macracantha.
sublanata, *Benth. in Enum. Pl. Hueg.* 42.—Austral.
subporosa, *F. Muell. Pl. Vict.* ii. 24.—Austral.
subrhombea, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 364.—Madag.
subternata, *F. Muell. & Benth. in Journ. Linn. Soc.* iii. (1859) 124.—Austral.
subtilifolia, H. B. & K. Nov. Gen. et Sp. vi. 268 = Piptadenia subtilifolia.
subtilinervis, *F. Muell. Pl. Vict.* ii. 32.—Austral.
subtilis, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 43.—Austral.
subulata, *Bonpl. Jard. Malm.* 110. t. 45.—Austral.
sulcata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 460.—Austral.
sulcipes, Sieber, ex G. Don, Gen. Syst. ii. 419 = decurrens.
Suma, Buch.-Ham. ex Wall. Cat. sub n. 5227 = A. Sundra.
Suma, *Kurz, ex Brandis. For. Fl.* 187.—Ind. or.
Sundra, *DC. Prod.* ii. 458.—Ind. or.
Sundra, Roxb. in Wall. Cat. n. 5227 = A. Suma.
taitensis, *Vatke, in Oest. Bot. Zeit.* xxx. (1880) 278.—Afr. trop.
tamarindifolia, Griseb. Cat. Pl. Cub. 82 = Mimosa polyancistra.
tamarindifolia, *Willd. Sp. Pl.* iv. 1092.—Am. austr.
tamariscina, Willd. l. c. 1062 = Gagnebina tamariscina.
taxifolia, A. Cunn. in Field, New South Wales, 344 = trinervata.
taxifolia, Lodd. Bot. Cab. t. 1225 = Riceana.
taxifolia, *Willd. Sp. Pl.* iv. 1050.—Cochinch.
Tayloriana, *F. Muell. in Wing, South. Sc. Record* (July, 1882).—Austral.
tenerrima, Miq. Fl. Ind. Bat. i. 14. t. 1 = pennata.
tenuiflora, Willd. Sp. Pl. iv. 1088 = Mimosa Cabrera.
tenuifolia, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 37.—Austral.
tenuifolia, Willd. Sp. Pl. iv. 1091 = paniculata.
tenuissima, F. Muell. ex Benth. in Journ. Linn. Soc. iii. (1859) 135 = xylocarpa.
tephroloba, A. Gray, Pl. Wright. i. 65 = Berlandieri.
teretifolia, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 326.—Austral.
tetragona, Willd. Sp. Pl. iv. 1069 = Calliandra tetragona.
tetragonocarpa, *Meissn. in Lehm. Pl. Preiss.* i. 4.—Austral.
tetragonophylla, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 121.—Austral.
texensis, Torr. & Gray, Fl. N. Am. i. 404 = filicina.
thegonocarpa, A. Cunn. ex Sweet, Hort. Brit. ed. III. 198, sphalm. = trigonocarpa.
Thibaudiana, DC. Prod. ii. 456 = Piptadenia moniliformis.
Thozetiana, F. Muell. Fragm. iv. 3 = Albizzia Thozetiana.
Tomentilla, Zipp. ex Span. in Linnaea, xv. (1841) 199 = pennata.
tomentosa, *Willd. Sp. Pl.* iv. 1087.—Ind. or.
torquata, Lag. Gen. et Sp. Nov. 16, n. 206 = Prosopis torquata.
tortilis, *Hayne, Arzneigew.* ix. t. 31.—Afr. trop.; Arab.
tortuosa, *Willd. Sp. Pl.* iv. 1083.—Am. austr.
torulosa, *Benth. in Journ. Linn. Soc.* iii. (1859) 139.—Austral.
translucens, *A. Cunn. in Hook. Ic. Pl.* t. 160.—Austral.
trapezoidea, DC. ex Steud. Nom. ed. II. i. 8 = decipiens.
triacantha, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 244 = glaucophylla.
triangularis, Benth. Enum. Pl. Hueg. 44 = biflora.
trichandra, Zucc. Pl. Nov. Fasc. ii. 41 = Leucaena diversifolia.
trichodes, Willd. Sp. Pl. iv. 1063 = Leucaena trichodes.
trichophylloides, Macfad. Fl. Jamaic. 320 = Piptadenia peregrina.
trigona, A. DC. Not. viii. Pl. Rar. Jard. Gen. 20 = myrtifolia.
trigonocarpa, A. Cunn. ex Lond. Hort. Brit. 407 = longifolia.

ACACIA:—
trigonophylla, *Meissn. in Lehm. Pl. Preiss.* ii. 199.—Austral.
trinervata, *Sieber, ex DC. Prod.* ii. 451.—Austral.
trinervis, *Desv. Journ. Bot.* iii. (1814) 70.—Hab.?
trincura, *F. Muell. Pl. Vict.* ii. 25.—Austral.
triptera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 325.—Austral.
triptycha, *F. Muell. ex Benth. Fl. Austral.* ii. 337.—Austral.
triquetra, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 358.—Austral.
trisperma, Mart. Herb. Fl. Bras. 108 = Piptadenia trisperma.
trispinosa, Stokes, Bot. Mat. Med. iii. 168 = A. Senegal.
trissoneura, F. Muell. Fragm. iv. 6 = heteroclita.
tristis, R. Grah. in Bot. Mag. t. 3420 = armata.
tristis, *Welw. ex Oliver, in Fl. Trop. Afr.* ii. 349.—Afr. trop.
truncata, Hort. ex Hoffmgg. Pfl. Verz. 34 = decipiens?
tuberosa, *Sterler, Hort. Nymph. ex Steud. Nom.* ed. II. i. 8.—Hab.?
tubulifera, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 520.—Peruv.
tucumanensis, *Griseb. in Goett. Abh.* xix. (1874) 135.—Am. austr.
tumida, *F. Muell. in Journ. Linn. Soc.* iii. (1859) 144.—Austral.
Tuundra, Steud. Nom. ed. II. i. 8, sphalm. = Sundra.
ulicina, *Meissn. in Lehm. Pl. Preiss.* ii. 202.—Austral.
umbellata, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 378.—Austral.
umbellifera, Kunth, Mimos. Humb. 100. t. 13 = filicina.
umbraculata, Wight, in Wall. Cat. n. 5245 = latronum.
umbrosa, A. Cunn. ex G. Don, Gen. Syst. ii. 405 = binervata.
uncifera, *Benth. in Mitch. Trop. Austr.* 341.—Austral.
uncinata, Lodd. Bot. Cab. t. 909 = undulaefolia.
uncinella, *Benth. in Linnaea*, xxvi. (1853–55) 613.—Austral.
uncinella, DC. ex Steud. Nom. ed. II. i. 8 = acanthocarpa.
uncinella, Desf. Cat. Hort. Par. ed. III. 299 = Mimosa uncinella.
undulata, Willd. Enum. Hort. Berol. Suppl. 68 = armata.
undulaefolia, *A. Cunn. ex G. Don, Gen. Syst.* ii. 404.—Austral.
ungulata, Desv. Journ. Bot. iii. (1814) 68 = Calliandra portoricensis.
uniglandulosa, Seem. & Schmidt, in Flora, xxvii. (1844) 495 = alata.
urophylla, *Benth. ex Lindl. in Bot. Reg.* (1841), *Misc.* 24.—Austral.
vaga, *Willd. Sp. Pl.* iv. 1066.—Bras.
Valenzuelana, A. Rich. Ess. Fl. Cub. 462 = villosa.
varians, Benth. in Mitch. Journ. Trop. Austral. 132 = salicina.
velutina, Benth. in Hook. Lond. Journ. Bot. i. (1842) 512 = monacantha.
velutina, *Bertol. Pl. Nov. Hort. Bonon.* ii. 13. t. 4.—Ind. occ.
velutina, *DC. Prod.* ii. 459.—Bras.
venosa, *Hochst. ex Benth. in Hook. Lond. Journ. Bot.* v. (1846) 98.—Abyss.
venulosa, *Benth. l. c.* 366.—Austral.
venusta, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1858) 53.—Hab.?
venusta, Willd. Enum. Hort. Berol. 1052 = Calliandra portoricensis.
vera, Willd. Sp. Pl. iv. 1085 = arabica.
Verek, Guill. & Perr. Tent. Fl. Senegamb. i. 245. t. 56 = A. Senegal.
verniciflua, *A. Cunn. in Field, New South Wales*, 344.—Austral.
verticillata, Sieber, ex Benth. Fl. Austral. ii. 332 = juniperina.
verticillata, *Willd. Sp. Pl.* iv. 1049.—Austral.
verugera, *Schweinf. in Linnaea*, xxxv. (1867–68) 340. t. 9, 10.—Afr. trop.
vespertina, Macfad. Fl. Jamaic. i. 318 = Calliandra portoricensis.
vestita, *Ker-Gawl. in Bot. Reg.* t. 698.—Austral.
viarum, *Allem. ex Tenore, Cat. Ort. Napol.* 77.—Bras.
vicioides, Ferr. & Galin. Voy. Abyss. iii. 119 = mellifera.

ACACIA:—
Victoriae, Benth. in Mitch. Journ. Trop. Austral. 333 = sentis.
villosa, *Willd. Sp. Pl.* iv. 1067.—Jamaica.
viminalis, *Ait. ex Steud. Nom.* ed. II. i. 8.—Austral.
viminea, *Dietr. ex Steud. l. c.* 9.—Austral.
Vincentis, Griseb. Fl. Brit. W. Ind. 222 = Pithecolobium tortum.
Virchowiana, *Vatke & Hildeb. in Oest. Bot. Zeit.* xxx. (1880) 275.—Afr. trop.
virescens, *DC. Cat. Hort. Monsp.* 74.—Am. austr.
virgata, Gaert. Fruct. ii. 317. t. 148 = Desmanthus virgatus.
virgata, Lodd. Bot. Cab. t. 1246 = verniciflua.
virgata, *Rafin. Fl. Ludov.* 136.—Am. bor.
virgultosa, Vahl, ex Walp. Rep. i. 929 = Inga virgultosa.
viridiflora, Kunth, Mimos. Humb. 81. t. 25 = Piptadenia subtilifolia.
viridiflora, Tweed. ex Benth. in Hook. Lond. Journ. Bot. iii. (1844) 98 = Calliandra viridiflora.
viridiramis, Burch. Trav. i. 300 = Xerocladia Zeyheri.
viridis, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 363.—Madag.
viscidula, Benth. in Maund, Botanist, iv. t. 179 = elongata.
viscidula, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 363.—Austral.
visciflua, F. Muell. Pl. Vict. ii. 24 = dodonaeifolia.
Visco, *Lorentz, ex Griseb. in Goett. Abh.* xxiv. (1879) 122.—Reg. Argent.
viscosa, Schrad. ex Wendl. Comm. Acac. 30. t. 7 = dodonaeifolia.
Visite, *Griseb. in Goett. Abh.* xix. (1874) 135.—Reg. Argent.
visneoides, *Colla, in Linnaea*, iv. (1829) *Lit.* 56.—Austral.
Vogeliana, Steud. Nom. ed. II. i. 9 = ambigua.
volubilis, *F. Muell. Fragm.* x. 98.—Austral.
vomeriformis, *A. Cunn. ex Benth. in Hook. Lond. Journ. Bot.* i. (1842) 332.—Austral.
vulcanica, Korth. Hort. Bogor. ex Flora, xxx. (1847) 705 = Albizzia montana.
Wallichiana, DC. Prod. ii. 458 = A. Catechu.
Wattsiana, *F. Muell. ex Benth. Fl. Austral.* ii. 374.—Austral.
Welwitschii, *Oliver, Fl. Trop. Afr.* ii. 341.—Afr. trop.
Westiana, DC. Prod. ii. 464 = riparia.
Whanii, *F. Muell. ex Benth. Fl. Austral.* ii. 386.—Austral.
Wickhami, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 379.—Austral.
Wightiana, R. Grah. in Wall. Cat. n. 5259 = Albizzia amara.
Wightii, *Baker, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 506.—Ind. or.
Wightii, Wight & Arn. Prod. 274 = Albizzia amara.
Wilhelmiana, F. Muell. in Trans. Phil Soc. Victoria, i. (1855) 37 = calamifolia.
Willdenowana, Wendl. Verz. K. Berggart. (1845) 5 = diptera.
Wrightii, *Benth. ex A. Gray, in Pl. Wright.* i. 64.—Texas.
xanthina, Benth. in Hook. Lond. Journ. Bot. i. (1842) 355 = bivenosa.
xanthophloea, *Benth. in Trans. Linn. Soc.* xxx. (1875) 511.—Afr. trop.
xiphocarpa, Hochst. ex Benth. in Hook. Lond. Journ. Bot. v. (1846) 96 = abyssinica.
xylocarpa, *A. Cunn. ex Benth. l. c.* i. (1842) 370.—Austral.
xylocarpa, Willd. Sp. Pl. iv. 1055 = Xylia dolabriformis.
Zygia, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 364.—Madag.
zygioides, *Baill. l. c.* 363.—Madag.

ACAENA, Linn. Mant. ii. 145 (1771).—*ROSACEAE*, Benth. & Hook. f. i. 623.
ANCISTRUM, Forst. Char. Gen. 3. t. 2. (1776).
LASIOCARPUS, Banks & Soland. ex Hook. f. Fl. Antaret. 264 (1847).
SPHAERULA, Anderss. ex Hook. f. l. c. 268 (1847).
aculeata, *Meyen, Reise*, i. 348.—Chili.
adscendens, *Vahl, Enum.* i. 297.—Am. austr.; Austral.
affinis, Hook. f. Fl. Antaret. 268. t. 96 = adscendens.

ACAENA:—
agrimonioides, *H. B. & K. Nov. Gen. et Sp.* vi. 231. —Mexico.
alpina, *Poepp. in Nov. Act. Nat. Cur.* xix. Suppl. i. (1843) 326.—Chili.
andicola, Gill. ex Walp. Rep. ii. 43 = pinnatifida.
andina, *Phil. in Linnaea*, xxviii. (1856) 636.—Chili.
antarctica, *Hook. f. Fl. Antarct.* 269.—Reg. Magell.
argentea, Bert. ex Steud. Nom. ed. II. i. 9 = Berteriana.
argentea, *Ruiz & Pav. Fl. Per.* i. 67. t. 103. f. b.—Chili.
Behriana, Schlecht. in Linnaea, xx. (1847) 660 = ovina.
Berteriana, *Bernh. ex Steud. Nom.* ed. II. i. 9.—Chili.
Buchanani, *Hook. f. Handb. New Zeal. Fl.* 57.—N. Zel.
Cadilla, *Hook. f. Fl. Antarct.* 269.—Patagonia.
caespitosa, *Gill. in Hook. Bot. Misc.* iii. 307 (1833).—Chili.
Calcitrapa, *Phil. in Linnaea*, xxxiii. (1864–65) 67.—Chili.
canescens, *Phil. Fl. Atac.* 18.—Chili.
Closiana, *C. Gay, Fl. Chil.* ii 298.—Chili.
Coxi, *Phil. in Linnaea*, xxxiii. (1864-65) 63.—Chili.
cuneata, *Hook. & Arn. in Hook. Bot. Misc.* iii. 307 (1833) = Reg. Magell.
cylindristachya, *Ruiz & Pav. Fl. Per.* i. 68 t. 104. f. 2.—Peruv.
decumbens, Menzies, in Hook. f. Fl. Antarct. 9 = Sanguisorbae.
depressa, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 548. —N. Zeland.
digitata, *Phil. in Anales Univ. Chil.* (1862) ii. 445.—Chili.
echinata, Lehm. Pl. Preiss. i. 95 = ovina.
elegans, *C. Gay, Fl. Chil.* ii. 297.—Chili.
elongata, *Linn. Mant.* ii. 200.—Mexico.
Eupatoria, *Schlecht. in Linnaea*, ii. (1827) 30.—Uruguay.
exigua, *A. Gray, Bot. U. S. Expl. Exped.* i. 498.—Ins. Sandvic.
glaberrima, Phil. in Linnaea, xxviii. (1856) 685 = pumila.
glabra, *J. Buch. in Trans. N. Z. Inst.* iv. (1872) 226. t. 14.—N. Zel.
Grahamiana, Gill. ex Walp. Rep. ii. 43 = adscendens.
hirsuta, *Phil. in Anales Univ. Chil.* (1872) 712.—Chili.
Huttoni, *R. Br.* [*ter.*] *in Trans. N. Z. Inst.* xvi. (1884) 382.—N. Zeland.
incisa, Lindl. Bot. Reg. t. 1271 = pinnatifida.
inermis, *Hook. f. Fl. N. Zel.* i. 54.—N. Zel.
integerrima, *Gill. in Hook. Bot. Misc.* iii. (1833) 306. —Chili.
Krausei, Phil. in Anales Univ. Chil. (1862) 711 = laevigata.
laevigata, *Ait. Hort. Kew.* ed. II. i. 68.—Reg. Magell.
lappacea, *Ruiz & Pav. Fl. Per.* i. 66. t. 103. f. a.—Peruv.
latebrosa, *Ait. Hort. Kew.* ed. II. i. 67.—Afr. austr.
laxiflora, *Willd. ex Steud. Nom.* ed. II. i. 9.—Am. austr.
leptacantha, *Phil. in Linnaea*, xxxiii. (1864–65) 66.—Chili.
lucida, *Vahl, Enum.* i. 296.—Ins. Falkl.
lucida, Willd. ex Steud. Nom. ed. II. i. 9 = pinnatifida.
macrocephala, *Poepp. Fragm. Syn.* 25.—Chili.
macrorhiza, *Hook. f. Fl. Antarct.* 265.—N. Granat.
macrostemon, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1846) 476, 269.—Chili.
magellanica, *Vahl, Enum.* i. 297.—Reg. Magell.
microcephala, *Schlecht. in Linnaea*, xxviii. (1856) 463. —Chili.
microphylla, *Hook. f. Fl. N. Zel.* i. 55.—N. Zel.
montana, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1846) 276.—Tasman.
montana, *Phil. in Linnaea*, xxviii. (1856) 686.—Chili.
montevidensis, Hook. f. Fl. Antarct. 265 = A. Eupatoria.
multifida, A. Dietr. Sp. Pl. i. 619 = trifida.
multifida, *Hook. f. Fl. Antarct.* 265.—Patagonia.
myriophylla, *Lindl. Bot. Reg. t.* 1271.—Reg. Argent.
Novae-Zelandiae, *T. Kirk, in Trans. N. Z. Inst.* iii. (1871) 177.—N. Zel.
ochreata, *Wedd. Chlor. And.* ii. 240.—Peruv.
oligacantha, *Phil. in Linnaea*, xxxiii. (1864–65) 66.—Chili.
ovalifolia, *Ruiz & Pav. Fl. Par.* i. 67. t. 103.—Am. austr.

ACAENA:—
ovalis, Pers. Syn. i. 142 = ovalifolia.
ovina, *A. Cunn. in Field, New South Wales*, 358.—Austral.
pectinata, *C. Presl, Epim. Bot.* 201.—Chili.
pinnatifida, Bert. ex Steud. Nom. ed. II. i. 9 = cuneata, myriophylla.
pinnatifida, Hort. ex Steud. l. c. = A. Berteriana.
pinnatifida, Lindl. Bot. Reg. t. 1271 = trifida.
pinnatifida, *Ruiz & Pav. Fl. Per.* i 68. t. 104.—Chili.
Poeppigiana, *C. Gay, Fl. Chil.* ii. 284.—Chili.
polycarpa, *Griseb. in Goett. Abh.* xix. (1874) 138.—Reg. Argent.
poterioides, *Willd. ex Steud. Nom.* ed. II. i. 9.—Am. austr.
pumila, *Phil. in Linnaea*, xxxiii. (1864–65) 67.—Chili.
pumila, *Vahl, Enum.* i. 298.—Reg. Magell.
quinquefida, *Phil. in Linnaea*, xxviii. (1856) 686.—Chili.
Sanguisorbae, *Vahl, Enum.* i. 294.—Austral.
sarmentosa, Carmich. in Trans. Linn. Soc. xii. (1818) 502 = A. Sanguisorbae.
sericea, *Jacq. f. Ecl.* i. 81. t. 55.—Mexico.
splendens, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 306.—Chili.
stellaris, *Meyen, Reise*, i. 348.—Chili.
stricta, *Griseb. in Goett. Abh.* xix. (1874) 138.—Reg. Argent.
suaveolens, A. Gray, Bot. U. S. Expl. Exped. i. 748, sphalm. = Acacia suaveolens.
subincisa, *Wedd. Chlor. And.* ii. 242.—Ecuador.
tridactyla, *Presl, Epim. Bot.* 201.—Mexico.
trifida, *Ruiz & Pav. Fl. Per.* i. 67. t. 104. f. c.—Calif.; Peru; Chili.
trifida, Steud. & Hochst. ex Steud. Nom. ed. II. i. 9 = Berteriana.
valdiviana, *Phil. in Linnaea*, xxxiii. (1864–65) 65.—Chili.
venulosa, *Griseb. in Goett. Abh.* vi. (1854) 118.—Reg. Magell.

ACAENOPS, Schrad. ex Steud. Nom. ed. II. 9 (1840) = **Dipsacus**, Linn.
pilosus, Fourr. in Ann. Soc. Linn. Lyon. N.S. xvi. (1868) 400 = D. pilosus.
vulgaris, Schrad. ex Steud. Nom. ed. II. i. 9 = D. pilosus.

ACAJOU, Tourn. ex Adans. Fam. ii. 344 (1763) = seq.

ACAJUBA, Gaertn. Fruct. i. 192. t. 40. (1788) = **Anacardium**, Linn.
occidentalis, Gaert. l. c. = Anac. occidentale.

ACALYPHA, Linn. Coroll. Gen. 19 (1737). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 311.
CALYPTROSPATHA, Klotzsch, ex Baill. Etudes Euph. 440 (1858).
CATURUS, Linn. ex Schreb. Gen. Pl. 677 (1791).
CUPAMENI, Adans. Fam. ii. 356 (1763).
GALURUS, Spreng. Anleit. ii. 364 (1817).
GYMNALYPHA, Griseb. in Bonplandia, vi. (1858) 2.
LINOSTACHYS, Klotzsch, ex Schlecht. in Linnaea, xix. (1847) 235.
ODONTEILEMA, Turcz. in Bull. Soc. Nat. Mosc. xxi. 1. (1848) 587.
USTERIA, Dennst. Schluess. Hort. Malab. 31 (1818).
abortiva, Hochst. ex Baill. Etud. Gén. Euphorb. 443 = indica.
accedens, *Muell. Arg. in Linnaea*, xxiv. (1865–66) 35.—Bras.
acuminata, *Benth. in Hook. Kew Journ.* vi. (1854) 329. —Bras.
acuminata, Vahl, ex Baill. Adansonia, i. (1860) 267 = spiciflora.
acuta, Thunb. Fl. Cap. ed. Schult. 546 = Adenocline acuta.
adenophora, Griseb. in Goett. Nachr. (1865) 175 = membranacea.
adenostachya, *Muell. Arg. in Linnaea*, xxxiv. (1865) 21.—Mexico.
adenotricha, *A. Rich. Tent. Fl. Abyss.* ii. 248.—Abyss.
adscendens, Hornem. Hort. Hafn. Suppl. 108 = chamaedrifolia.
affinis, Klotzsch, in Nov. Act. Acad. Nat. Cur. xix. Suppl. i. (1843) 416 = stipulacea.
agrimonioides, D. Dietr. Syn. v. 377 err. typ. = anemioides.
alnifolia, Poir. Encyc. vi. 203 = Poiretii.

ACALYPHA:—
alnifolia, *Klein, ex Willd. Sp. Pl.* iv. 525.—Ind. or.
alnifolia, Wall. Cat. n. 7782 A = capitata.
alopecuroidea, *Jacq. Ic. Rar.* iii. 19. t. 620; *Coll.* iii. 196. —Venezuela.
amblyodonta, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 365.—Bras.
amboynensis, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 233 = grandis.
amentacea, Roxb. Fl. Ind. iii. 676 = fruticosa.
andina, *Muell. Arg. in Flora*, lv. (1872) 26.—Reg. Andin.
anemioides, *H. B. & K. Nov. Gen. et. Sp.* ii. 94.—Mexico.
angatensis, *Blanco, Fl. Filip.* ed. I. 750.—Ins. Philipp.
angolensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 335.—Afr. trop.
angustata, Sond. in Linnaea, xxiii. (1850) 115 = pedunculares.
angustifolia, *Sw. Prod. Veg. Ind. Occ.* 99.—Ins. S. Doming.
anisodonta, *Muell. Arg. in DC. Prod.* xv. II. 819.—Ins. Pacif.
arborea, *Comm. ex Poir. Encyc.* vi. 205.—Ins. Masc.; St. Helen.
Arciana, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 362.—Bras.
aristata, *H. B. & K. Nov. Gen. et Sp.* ii. 93.—N. Granat.
arvensis, *Poepp. & Endl. Nov. Gen. & Sp.* iii. 21.—Peruv.
australis, *Linn. Sp. Pl.* 1004.—Reg. trop.
Bailloniana, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 44.—Afr. trop.
Baroni, *Baker, in Journ. Linn. Soc.* xx. (1884) 254.—Madag.
benguelensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 335.—Afr. trop.
Berteroana, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 33.—Porto-Rico.
betulaefolia, Sw. Prod. Veg. Ind. Occ. 99 = scabrosa.
betulina, Retz. Obs. Fasc. v. 30 = fruticosa.
betulina, Schweinf. Pl. quaedam Nilot. 13 = Cephalocroton cordofanus.
betulina, Sond. in Linnaea, xxxiii. (1850) 116 = glabrata.
betuloides, Klotzsch, ex Baill. in Adansonia, v. (1864–65) 228 = variabilis.
betuloides, Pav. ex Klotzsch, in Seem. Bot. Herald, 101 = diversifolia.
Bilbergiana, Klotzsch, l. c. = diversifolia.
bipartita, *Muell. Arg. in Flora*, xlvii. (1864) 538.—Afr. trop.
bisetosa, *Spreng. Syst.* iii. 879.—Venezuel.
boehmerioides, *Miq. Fl. Ind. Bat. Suppl.* i. 459.—Ins. Pacif.
Boiviniana, *Baill. Adansonia*, i. (1860–61) 272.—Afr. trop.
boliviensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 162.—Bolivia.
Botteriana, *Muell. Arg. l. c.* 46.—Mexico.
brachiata, Krauss, in Flora, xxvii. (1845) 83, nomen = tenuis.
brachyandra, *Baill. in Adansonia*, v. (1864–65) 232. —Bras.
brachyclada, *Muell. Arg. in DC. Prod.* xv. II. 862.—Peruv.
brachystachya, *Hornem. Hort. Hafn.* ii. 909.—China.
bracteata, *Miq. Fl. Ind. Bat.* ii. 406.—Malaya.
brasiliensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 37.—Bras.
brevibracteata, *Muell. Arg. in DC. Prod.* xv. II. 855.—Ins. Pacif.
brevicaulis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 13.—Mexico.
brevipes, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 348.—Bras.
brevipes, *Rafin. New Fl. Am.* i. 44.—Am. bor.
Buchenavii, *Muell. Arg. in Bremen, Abh.* vii. (1880) 27.—Afr. trop.
bullata, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 17. —Peruv.
californica, *Benth. Bot. Voy. Sulph.* 51.—Calif.
callosa, *Benth. Pl. Hartw.* 252.—N. Granat.
calyciformis, Wight, ex Wall. Cat. n. 7786 = brachystachya.
cancana, *Muell. Arg. in Flora*, xlvii. (1864) 438. —N. Granat.
canescens, Wall. Cat. n. 7785 = indica.

ACALYPHA:—
capensis, D. Dietr. Syn. v. 378, sphalm. = corensis.
caperonioides, Baill. Adansonia, iii. (1862–63) 157 = peduncularis.
capillipes, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 40.—Austral.
capitata, Wall. Cat. n. 7783. A, B = fruticosa.
capitata, *Willd. Sp. Pl.* iv. 525.—Ind. or.
caroliniana, Blanco, Fl. Filip. ed. I. 748 = indica.
caroliniana, *Ell. Sketch.* ii. 645.—Am. bor.
caroliniana, Walt. Fl. Carol. 238 = virginica.
carpinifolia, Poepp. ex Baill. Etud. Gén. Euphorb. 442 = leptostachya.
carpinifolia, *Poir. in Lam. Encyc.* vi. 203. (1804).—Ins. S. Doming.
carthagenensis, *Jacq. Enum. Pl. Carib.* 32.—N. Granat.
Caturus, *Blume, Bijdr.* 629.—Java.
caudata, H. B. & K. Nov. Gen. et Sp. ii. 95 = macrostachya.
chamaedrifolia, Muell. Arg. in DC. Prod. xv. II. 879 = reptans.
Chibomboa, *Baill. Adansonia*, i. (1860–61) 269.—Ins. Comor.
chinensis, Roxb. Fl. Ind. iii. 677 = australis.
chorisandra, *Baill. Adansonia*, v. (1864–65) 235.—Bras.
ciliata, *Forsk. Fl. Aegypt. Arab.* 162.—Geront. trop.
ciliata, Wall. Cat. n. 7779 J = indica.
cincta, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 20.—Mexico.
cireinnata, A. Gray, ex Seem. List Vit. Pl. 11 = Wilkesiana.
Clausseni, *Baill. Adansonia*, v. (1864–65) 226.—Bras.
codonocalyx, *Baill. l. c.* i. (1860–61) 271.—Ins. Comor.
colorata, Spreng. Syst. iii. 879 = integrifolia.
Commersoniana, Baill. Adansonia, i. (1860–61) 267 = inregrifolia.
communis, Muell. Arg. in Linnaea, xxxiv. (1865–66) 23 = variabilis.
conferta, Roxb. Fl. Ind. iii. 677 = brachystachys.
consimilis, *Muell. Arg. in DC. Prod.* xv. II. 807.—Ins. Pacif.
conspicua, *Muell. Arg. l. e.* 832.—Mexico.
contermina, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 46.—Peruv.
corchorifolia, *A. Rich. in Sagra, Hist. Cuba*, iii. 203.—Ind. occ.
corchorifolia, Vahl, ex Baill. Etud. Gén. Euphorb. 443 = fallax.
corchorifolia, Willd. Sp. Pl. iv. 524 = reptans.
cordata, Drège, ex Baill. Adansonia, iii. (1863) 158 = Eckloni.
cordata, Thunb. Prod. Pl. Cap. 117 = decumbens.
cordifolia, Hook. f. in Trans. Linn. Soc. xx. (1847) 186 = parvula.
cordifolia, *Griseb. in Goett. Abh.* xix. (1874) 97.—Reg. Argent.
cordifolia, Anderss. Galap. Veg. 103 = parvula.
cordobensis, Griseb. in Goett. Abh. xix. (1874) 97 = seq.
cordoviensis, *Muell. Arg. in Journ. Bot.* xii. (1874) 228.—Reg. Argent.
corensis, *Jacq. Enum. Pl. Carib.* 32.—Ind. occ.
erassa, Meissn. ex Krauss, in Flora, xxviii. (1845) 83 = peduncularis.
crenata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 245.—Afr. trop.
crenulata, Rafin. New Fl. Am. i. 44 = virginica.
cucullata, Poir. Encyc. Suppl. iv. 683 = macrostachya.
cuneata, *Poepp. & Endl. Nov. Gen.* iii. 22.—Am. austr.
Cunninghami, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 35.—Austral.
cuspidata, *Jacq. Hort. Schoenb.* ii. 63. t. 243.—Am. austr.
cylindrica, Roxb. Fl. Ind. iii. 678 = Poiretii.
decidua, *Forsk. Fl. Aegypt. Arab.* 161.—Arab.
decumbens, *Thunb. Prod. Pl. Cap.* 117.—Afr. austr.
densiflora, Blume, Bijdr. 628 = hispida.
dentata, Schum. & Thonn. Beskr. Guin. Pl. 410 = Mallotus oppositifolius.
denudata, *Muell. Arg. in DC. Prod.* xv. II. 819.—Ins. Pacif.
depauperata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 160.—Mexico.
Deppeana, Schlecht. in Linnaea, vii. (1832) 385 = Schiedeana.

ACALYPHA:—
dictyoneura, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 12.—Peruv.
diffusa, Anderss. Galap. Veget. 104 = parvula.
digyneia, *Rafin. Fl. Ludov.* 112.—Louisian.
digynostachya, *Baill. Adansonia*, v. (1864–65) 233.—Bras.
dimorpha, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 354.—Bras.
discolor, Boj, Hort. Maurit. 286 = integrifolia.
discolor, E. Mey. Zwei Pfl. Docum. 129 (n. 20); in Flora, xxviii. (1845) 84 = decumbens.
distans, *Muell. Arg. in DC. Prod.* xv. II. 820.—Cuba.
divaricata, Baill. in Adansonia, v. (1864–65) 234 = gracilis.
divaricata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 34.—Peruv.
divaricata, *Rafin. New. Fl. Am.* i. 44.—Am. bor.
diversifolia, *Jacq. Hort. Schoenb.* ii. 63. t. 244.—Am. austr.
domingensis, Spreng. Syst. Veg. iii. 880 = carpinifolia.
dumetorum, *Muell. Arg. in Journ. Bot.* ii. (1864) 334—Afr. trop.
Dupraeana, Baill. Adansonia, v. (1864–65) 229 = accedens.
echinata, *Rafin. New. Fl. Am.* i. 45.—Am. bor.
Eckloni, *Baill. Adansonia*, iii. (1862–63) 158.—Afr. austr.
elegantula, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 246 = brachystachya.
elliptica, *Sw. Prod. Veg. Ind. Occ.* 99.—Jamaica.
emirnensis, *Baill. Adansonia*, i. (1860–61) 270.—Madag.
eremorum, *Muell. Arg. in Flora*, xlvii. (1864) 440.—Austral.
erythrostachya, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 51.—N. Granat.
estrellana, Baill. Adansonia, v. (1863–64) 237 = brasiliensis.
exaltata, *Baill. l. e.* ii. (1861–62) 225.—N. Caled.
fallax, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 43.—As. trop.
fasciculata, *Muell. Arg. l. c.* 31.—Ins. Mascar.
filiformis, Poir. Encyc. vi. 205 = arborea.
filiformis, Klotzsch, ex Schlecht. in Linnaea, xix. (1847) 235 = Schlechtendaliana.
fimbriata, Baill. Adansonia, i. (1860–61) 272 = Bailloniana et crenata.
fimbriata, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 245.—Abyss.
fimbriata, Schum. & Thonn. Beskr. Guin. Pl. 409 = ciliata.
firmula, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 21.—Am. centr.
fissa, Herb. Madr. ex. Wall. Cat. n. 7786. B. = brachystachya.
flaccida, Hook. f. in Trans. Linn. Soc. xx. (1847) 186 = parvula.
flexuosa, Wight, ex Steud. Nom. ed. II. i. 9 = lanceolata.
Forsteriana, *Muell. Arg. in DC. Prod.* xv. II. 807.—Ins. Pacif.
Fournieri, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 162.—Mexico.
Frederici, *Muell. Arg. in DC. Prod.* xv. II. 828.—Mexico.
fruticosa, *Forsk. Fl. Aegypt. Arab.* 161.—Geront. trop.
fruticulosa, *Rafin. Fl. Ludov.* 112.—Louisiana.
fuscescens, *Muell. Arg. in DC. Prod.* xv. II. 821.—Afr. trop.
gemina, Spreng. Syst. iii. 880 = australis.
glabrata, *Thunb. Prod. Pl. Cap.* 117.—Afr. austr.
glabrata, Vahl, ex A. Juss. Tent. Euph. 105 = Thecacoris madagascariensis.
glandulifolia, Meissn. ex Krauss, in Flora, xxviii. (1845) 83 = peduncularis.
glandulosa, *Blanco, Fl. Filip.* ed. I. 749.—Ins. Philipp.
glandulosa, *Cav. in Anal. Hist. Nat. Madr.* ii. (1800) 141.—Mexico.
glechomaefolia, A. Rich. Fl. Cub. Fanerog. ii. 205 = reptans.
Goudotiana, Baill. Adansonia, i. (1860–61) 268 = arborea.
gracilens, A. Gray, Man. Bot. N. Unit. States, 390 = virginica.
gracilipes, Baill. Adansonia, i. (1860–61) 273 = integrifolia.

ACALYPHA:—
gracilis, *Spreng. Syst.* iv; *Cur. Post.* 315.—Bras.
grandidentata, *Muell. Arg. in DC. Prod.* xv. II. 823.—Afr. austr.
grandifolia, Poir. Encyc. vi. 204 = Claoxylon crassifolium.
grandis, *Benh. in Hook. Lond. Journ. Bot.* ii. (1843) 232.—Malaya.
Hartwegiana, Benth. ex Baill. Etud. Gén. Euphorb. 442 = diversifolia.
havanensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 49.—Cuba.
hederacea, *Torr. in Bot. Mex. Bound.* 200.—N. Mexico.
hernandifolia, *Sw. Prod. Veg Ind. Occ.* 99.—Jamaica.
heterodonta, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 12.—Am. trop.
hirsuta, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 248 = villicaulis.
hirsutissima, Willd. Sp. Pl. iv. 528 = macrostachya.
hirta, Cav. in Anal. Hist. Nat. ii. (1800) 141 = phleoides.
hirta, Spreng. Syst. iv; Cur. Post. 315 = variabilis.
hispida, Blume, Bijdr. 628 = boehmerioides.
hispida, *Burm. f. Fl. Ind.* 303. t. 61.—Ind. or.
hispida, Wall. Cat. n. 7780. pro parte = fallax.
hispida, Willd. Sp. Pl. iv. 523 = Poiretii.
Hochstetteriana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 39.—Afr. trop.
hologyna, *Baker, in Journ. Linn. Soc.* xxi. (1885) 441.—Madag.
Hystrix, Balb. ex Spreng. Syst. iii. 883 = aristata.
indica, *Linn. Sp. Pl.* 1003.—As. & Afr. trop.
indica, Vell. Fl. Flum. x. t. 64 = Poiretii.
infesta, *Poepp. & Endl. Nov. Gen.* iii. 22.—Peruv.
infestans, Muell. Arg. in Linnaea, xxxiv. (1865–66) 23 = praec.
insulana, *Muell. Arg. in Flora* (1864) 439.—Ins. Pacif.
integrifolia, *Willd. Sp. Pl.* iv. 530.—Ins. Mascar.
interrupta, Schlecht. in Linnaea, vii. (1832) 386 = Bernardia interrupta.
jamaicensis, *Rafin. New Fl. Am.* 46.—Jamaica.
japonica, Houtt. ex Steud. Nom. ed. I. 4 = Boehmeria spicata.
Jardini, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 36.—Ins. Pacif.
Karwinskii, *Muell. Arg. in Flora*, lv. (1872) 41.—Am. centr.
Klotzschiana, Muell. Arg. in DC. Prod. xv. II. 857 = prunifolia.
Klotzschii, Baill. Adansonia, v. (1864–65) 231 = longifolia.
Kraussiana, Buching. ex Krauss, in Flora, xxvii. (1845) 84 = decumbens.
laevifolia, *Muell. Arg. in DC. Prod.* xv. II. 853.—Ins. Pacif.
laevigata, *Sw. Prod.* 99.—Jamaica.
laevigata, Willd. Sp. Pl. iv. 527 = carpinifolia.
Lagascana, *Muell. Arg. in Flora*, lv. (1872) 27.—Am. centr.
lagoensis, *Muell Arg. in Mart. Fl. Bras.* xi. II. 367.—Bras.
lamiifolia, *Scheele, in Linnaea*, xxv. (1852) 587.—Afr. austr.
lanceolata, Wall. Cat. n. 7789 = australis.
lanceolata, *Willd. Sp. Pl.* iv. 524.—Ind. or.
Langiana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 159.— Mexico.
languida, E. Mey. & Sond. in Linnaea, xxiii. (1850) 116 = petiolaris.
languida, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 29.—Afr. austr.
lantanaefolia, Boj. Hort. Maurit. 286 = reticulata.
latifolia, *Muell. Arg. in DC. Prod.* xv. II. 817.—Ins. Pacif.
laxiflora, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 19.—Ind. occ.
leonensis, *Benth. in Hook. Nig. Fl.* 504 = Mareya micrantha.
Lepinei, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 14.—Ins. Pacif.
leptoclada, *Benth. in Bot. Voy. Sulph.* 164.—Mexico.
leptopoda, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 39.—Mexico.
leptorhachis, *Muell. Arg. l. c.* 7.—Cuba.
leptostachya, A. Rich. in Sagra, Hist. Cuba, iii. 205 = laxiflora.
lepostachya, H. B. & K. Nov. Gen et Sp. ii. 96 = diversifolia.
Liebmanni, Muell. Arg. in DC. Prod. xv. II. 820 = seq.

ACALYPHA:—

Liebmanniana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 161.—Mexico.
Lindeniana, *Muell. Arg. in DC. Prod.* xv. II. 827.—Mexico.
Lindheimeri, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 47.—Mexico.
Linostachys, Baill. Adansonia, v. (1864–65) 235 = villosa.
Livingstoniana, *Muell. Arg. in Flora*, xlvii. (1864) 440.—Afr. trop.
longestipularis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 51.—Mexico.
longifolia, Baill. Etud. Gen. Euph. 443 = cuneata.
longifolia, *Klotzsch, ex Baill. Adansonia*, v. (1864–65) 231.—Bras.
longispicata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 163.—Mexico.
Lyallii, *Baker, in Journ. Linn. Soc.* xx. (1884) 255.—Madag.
macrodonta, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 51.—Peruv.
macrophylla, H. B. & K. Nov. Gen. et Sp. ii. 96 = macrostachya.
macrosperma, Muell. Arg. in Linnaea, xxxiv. (1865–66) 18 = membranacea.
macrostachya, *Jacq. Hort. Schoenb.* ii. 63. t. 245.—Am. austr.
macrostachyos, Poir. Encyc. vi. 208 = Poiretii.
macrostachyoides, *Muell. Arg. in DC. Prod.* xv. II. 809.—Mexico.
major, Baill. Adansonia, v. (1864–65) 236 = Weddelliana.
malabarica, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 42.—Ind. or.
Mandoni, *Muell. Arg. l. c.* 162.—Bolivia.
Manniana, *Muell. Arg. in Flora*, xlvii. (1864) 441.—Afr. occ.
Mappa, Willd. Sp. Pl. iv. 526 = Macaranga Mappa.
marginata, *Spreng. Syst.* iii. 879.—Ind. or.
Martiana, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 359.—Bras.
melochiaefolia, *Muell. Arg. in DC. Prod.* xv. II. 821.—Mexico.
membranacea, *A. Rich. Fl. Cub. Fanerog.* ii. 204.—Cuba.
mexicana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 41.—Mexico.
micrantha, Benth. in Hook. Nig. Fl. 505 = Mareya micrantha.
microcephala, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 160.—Mexico.
microgyne, Poepp. & Endl. Nov. Gen. iii. 21 = diversifolia.
microphylla, Klotzsch, in Seem. Bot. Voy. Herald, 278 = reptans.
microstachya, Benth. Pl. Hartw. 71 = mollis.
mollis, *H. B. & K. Nov. Gen. et. Sp.* ii. 94.—Mexico.
monococca, Engelm. ex A. Gray, Man. Bot. N. U. States, 390 = virginica.
monostachya, Benth. Pl. Hartw. 15 = anemioides.
monostachya, *Cav. in Anal. Hist. Nat. Madr.* ii. (1800) 138.—Mexico.
multicaulis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 53.—Bras.
multipartita, Moench, Meth. Suppl. 122 = tomentosa.
muralis, *Zipp. ex Span. in Linnaea*, xv. (1841) 350, nomen.—Timor.
nana, Griseb. in Goett. Nachr. (1865) 176 = reptans.
nemorum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 38.—Austral.
neocaledonica, *Muell. Arg. in DC. Prod.* xv. II. 812.—N. Caled.
neogranatensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 15.—N. Granat.
neomexicana, *Muell. Arg. l. c.* 19.—N. Mexico.
neptunica, *Muell. Arg. in Bremen Abh.* vii. (1880) 26.—Afr. trop.
nigritiana, *Muell. Arg. in Flora*, xlvii. (1864) 440.—Afr. trop.
obovata, Benth. in Bot. Voy. Sulph. 163. t. 53 = cuneata.
obscura, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 163.—Mexico.
obtusa, Thunb. Fl. Cap. ed. Schult. 546 = Leidesia obtusa.
obtusata, Spreng. ex Steud. Nom. ed. II. i. 10. [Adenocline, sp.].—Afr. austr.
ocymoides, *H. B. & K. Nov. Gen. et. Sp.* ii. 93.—Mexico.
odorata, Steud. Nom. ed. II. ii. 696 sphalm. = integrifolia.

ACALYPHA:—

oligantha, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 159.—Mexico.
oligodonta, *Muell. Arg. in DC. Prod.* xv. II. 831.—Mexico.
ornata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 247.—Abyss.
ostryaefolia, Riddell, Syn. Fl. W. St. 33 = caroliniana.
ovalifolia, Baill. Adansonia, i. (1860–61) 269 = arborea.
oxyodonta, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 367.—Mexico.
padifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 97.—Am. austr.
panamensis, Klotzsch, in Seem. Bot. Voy. Herald, 101 = diversifolia.
Pancheriana, *Baill. Adansonia*, ii. (1861–62) 225.—N. Caled.
paniculata, *Miq. Fl. Ind. Bat.* i. II. 406. (1859).—Geront. trop.
parvifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 161.—Mexico.
parvula, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 185.—Ins. Galap.
Pastoris, DC. ex Willd. Enum. Hort. Berol. 993 = phleoides.
Pastoris, Schrank, in Denkschr. Bayer. Ges. ii. (1822) 39 = phleoides.
patens, *Muell. Arg. in DC. Prod.* xv. II. 847.—Afr. austr.
pauciflora, Hornem. Hort. Hafn. ii. 909 = australis.
Pavoniana, Muell. Arg. in Linnaea, xxxiv. (1865–66) 50 = arvensis.
Peckoltii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 365.—Bras.
pedunculares, E. Mey. Zwei Pfl. Docum. 136 n. 26 = Zeyheri.
pedunculares, *Meissn. ex Krauss, in Flora*, xxviii. (1845) 82.—Afr. austr.
pendula, Wright, ex Griseb. in Goett. Nachr. (1865) 176 = reptans.
persimilis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 25.—Ind. occ.
peruviana, *Muell. Arg. l. c.* 17.—Peruv.
Pervilleana, Baill. Adansonia, i. (1860–61) 273 = arborea.
petiolaris, *Krauss, in Flora*, xxviii. (1845) 83.—Afr. austr.
petiolaris, Sond. in Linnaea, xxiii. (1850) 117 = Sonderiana.
phleoides, *Cav. in Anal. Hist. Nat. Madr.* ii. (1800) 139.—Mexico.
phleoides, Torr. Bot. Mex. Bound. Surv. 199 = Lindheimeri.
pilifera, Klotzsch, ex Baill. Adansonia, v. (1860–61) 227 = variabilis.
pilosa, *Cav. in Anal. Nat. Hist. Madr.* ii. (1800) 136.—Am. trop.; Ins. Philipp.
pinnata, Poir. Encyc. vi. 205 = Tragia pinnata.
platyphylla, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 6.—Peruv.
plicata, *Muell. Arg. in DC. Prod.* xv. II. 855.—Bolivia.
Pohliana, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 360.—Bras.
Poiretii, *Spreng. Syst.* iii. 879.—Am. trop.
polymorpha, *Muell. Arg. in Journ. Bot.* ii. (1864) 335.—Afr. trop.
polystachya, *Jacq. Hort. Schoenb.* ii. 64. t. 246.—Peruv.; Mexico.
popayanensis, H. B. & K. Nov. Gen. et Sp. vii. 173 = diversifolia.
portoricensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 22.—Ind. occ.
Pringlei, *S. Wats. in Proc. Am. Acad.* xx. (1885) 373.—Am. bor.
prunifolia, H. B. & K. Nov. Gen. et Sp. ii. 92 = phleoides.
prunifolia, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 37.—Bras.
pruriens, *Nees & Mart. l. c.* 36.—Bras.
psilostachya, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 246.—Abyss.
pubiflora, *Baill. Adansonia*, i. (1860–61) 268.—Afr. trop.; Madag.
punctata, Meissn. ex Krauss, in Flora, xxvii. (1845) 83 = peduncularis.
purpurascens, *H. B. & K. Nov. Gen. et. Sp.* ii. 97.—Mexico.
pygmaea, A. Rich. Fl. Cub. Fanerog. ii. 205 = reptans.
racemosa, Heyne, in Wall. Cat. n. 7784 = paniculata.

ACALYPHA:—

radians, *Torr. in Bot. Mex. Bound.* 200.—N. Mexico.
radicans, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 39.—Bras.
Radula, *Baker, in Journ. Linn. Soc.* (1884) 254.—Madag.
reflexa, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 33.—Peruv.
reniformis, Hook. f. in Trans. Linn. Soc xx. (1847) 187 = parvula.
repanda, *Muell. Arg. in Flora*, xlvii. (1864) 439.—Ins. Pacif.
reptans, *Sw. Prod. Veg. Ind. Occ.* 99.—Ind. occ.
reticulata, Muell. Arg. in DC. Prod. xv. II. 851 = arborea.
rhombifolia, Baill. Adansonia, v. (1864–65) 230 = Poiretii.
rhombifolia, *Schlecht. in Linnaea*, vii. (1832) 382.—Mexico.
rhomboidea, *Rafin. New. Fl. Am.* i. 45.—Am. bor.
Richardiana, *Baill. in Adansonia*, i. (1860–61) 268.—Madag.
Riedeliana, *Baill. l. c.* v. (1864–65) 231.—Bras.
rivularis, *Seem. List Vit. Pl.* 11.—Ins. Fiji.
rottleroides, *Baill. Adansonia*, i. (1860–61) 270.—Madag.
rotundifolia, Vahl, ex Baill. Etud. Gén. Euphorb. 442 = infesta.
rubra, Roxb. in Beatson, Tracts St. Helena Append. 295 = arborea.
rubra, Wight, ex Wall. Cat. n. 7781 = ciliata.
rubra, Willd. Enum. Hort. Berol. 992 = phleoides.
Ruiziana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 16.—Peruv.
salicifolia, *Muell. Arg. in Flora*, xlvii. (1864) 438.—Peruv.
salviaefolia, *Baill. Etud. Gén. Euphorb.* 443, nomen.—Ins. Mascar.?
samydaefolia, *Poepp. & Endl. Nov. Gen.* iii. 21. t. 224.—Peruv.
scabra, Vahl, ex Baill. Adansonia, i. (1860–61) 280 = Claoxylon parviflorum.
scabrosa, *Sw. Prod. Veg. Ind. Occ.* 100.—Jamaica.
scandens, *Benth. in Hook. Kew Journ.* vi. (1854) 329.—Am. trop.
Schiedeana, *Schlecht. in Linnaea*, vii. (1832) 384.—Mexico.
Schlechtendahliana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 6.—Mexico.
Schlumbergeri, *Muell. Arg. in DC. Prod.* xv. II. 861.—Mexico.
Seemanni, Klotzsch, in Seem. Bot. Voy. Herald, 102 = macrostachya.
segetalis, *Muell. Arg. in Journ. Bot.* ii. (1864) 336.—Afr. occ.
seminuda, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 360.—Bras.
senensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 96.—Afr. occ.
senilis, *Baill. Adansonia*, v. (1864–65) 228.—Am. austr.
septemloba, *Muell. Arg. in Flora*, lv. (1872) 27.—Amer. centr.
sericea, Anderss. Galapag. Veget. 103 = parvula.
sessilis, Poir. Encyc. vi. 204 = australis.
setosa, *A. Rich. Fl. Cub. Fanerog.* ii. 204.—Cuba.
sidaefolia, *A. Rich. Tent. Fl. Abyss.* ii. 249.—Abyss.
sidaefolia, H. B. & K. Nov. Gen. et Sp. ii. 95 = macrostachya.
somalium, Muell. Arg. in Bremen Abh. vii. (1880) 27 = segetalis.
Sonderiana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 9.—Afr. austr.
Spachiana, *Baill. Adansonia*, i. (1860–61) 272.—Madag.
spicata, Anderss. Galap. Veget. 104 = parvula.
spicata, Forsk. Fl. Aegypt. Arab. 161 = indica.
spiciflora, *Burm. f. Fl. Ind.* 303 (i. e. 203). t. 61.—Madag.
spiciflora, Thou. ex Baill. Adansonia, i. (1860–61) 281 = Claoxylon Linostachys.
spiciflora, Poir. Encyc. vi. 206 = Claoxylon parviflorum.
spicigera, Klotzsch, in Seem. Bot. Voy. Herald, 101 = diversifolia.
spinescens, *Benth. in Hook. Ic. Pl.* t. 1291.—Malaya.
stenoloba, *Muell. Arg. in Flora*, lv. (1872) 41.—Peruv.
stipulacea, *Klotzsch, in Nov. Act. Acad. Nat. Cur.* xix. (1843) Suppl. I. 416.—Ins. Molucc.

ACALYPHA:—
strieta, *Poepp. & Endl. Nov. Gen.* iii. 21. t. 225.—Peruv.
strobilifera, Hook. f. in Trans. Linn. Soc. xx. (1847) 187 = parvula.
subtomentosa, *Lag. Elench. Hort. Reg. Matr.* (1805); *Gen. et Sp. Nov.* 21.—Mexico.
subvillosa, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 341.—Bras.
supera, Forsk. Fl. Aegypt. Arab. 162 = brachystachya.
tarapotensis, *Muell. Arg. in DC. Prod.* xv. II. 808.—Peruv.
tenuicaulis, Baill. Adansonia, v. (1864-65) 234 = multicaulis.
tenuifolia, *Muell. Arg. in DC. Prod.* xv. II. 863.—Venezuela.
tenuiramea, *Muell. Arg. l. c.* 858.—Ins. Pacif.
tenuis, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 30.—Afr. austr.
tiliaefolia, Poir. Encyc. vi. 203. (An Euphorbiacea ?).—Ins. S. Doming.
tomentosa, *Blanco, Fl. Filip.* ed. I. 750.—Ins. Philipp.
tomentosa, Boj. Hort. Maurit. 285 = marginata.
tomentosa, *Sw. Prod. Veg. Ind. Occ.* 99.—Ins. S. Domingo.
trachyloba, *Muell. Arg. in Flora*, lv. (1872) 25.—Mexico.
tricholoba, *Muell. Arg. in Linnaea*, xxxiv. (1865-66). —Mexico.
tricolor, *Seem. Fl. Vit.* 225.—Ins. Pacif.
triloba, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 23. —Mexico.
tristis, Poepp. & Endl. Nov. Gen. et Sp. iii. 22 = macrostachya.
ulmifolia, Benth. Pl. Hartw. 252 = diversifolia.
unibracteata, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 160.—Mexico.
urophylla, *Boiv. ex Baill. Adansonia*, i. (1860-61) 273.—Afr. trop.
urostachya, *Baill. l. c.* v. (1864-65) 229.—Bras.
urticaefolia, *Poir. Encyc.* vi. 208.—Am. austr.
urticoides, Klotzsch, ex Baill. Adansonia, v. (1864-65) 227 = communis.
vagans, *Cav. in Anal. Hist. Nat. Madr.* ii. (1800) 139.—Mexico.
Vahliana, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 43.—Afr. trop.
variabilis, *Klotzsch, ex Baill. Adansonia*, v. (1864-65) 226.—Bras.
Vedeliana, Baill. Adansonia, ii. (1861-62) 224 = Macaranga Vedeliana.
velamea, *Baill. l. c.* v. (1864-65) 228 = Bras.
velutina, Hook. f. in Trans. Linn. Soc. xx. (1847) 186 = parvula.
venosa, Poir. Encyc. vi. 204 = Leptonema venosum.
vestita, Benth. in Bot. Voy. Sulph. 164 = cuspidata.
villicaulis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 248.—Abyss.
villosa, *Jacq. Enum. Pl. Carib.* 32.—Ind. occ.
villosa, Vahl, ex Baill. Etud. Gén. Euphorb. 442 = Poirctii.
virgata, Forst. f. Prod. 67 = Forsteriana.
virgata, *Linn. Pugill. Jamaic. Pl.* 28 (*Am. Acad.* v. 410).—Jamaica.
virgata, Vell. Fl. Flum. x. t. 63 = variabilis.
virgata, Thunb. Fl. Jap. 268 = australis.
virginica, *Linn. Sp. Pl.* 1003.—Am. bor.
virginica, Wall. Cat. n. 7779 G = fallax.
Wallichiana, Muell. Arg. in DC. Prod. xv. II. 802 = seq.
Wallichii, Thw. Enum. Pl. Zeyl. 271 = paniculata.
Weddelliana, Baill. Adansonia, v. (1864-65) 232 = brasiliensis.
Welwitschiana, *Muell. Arg. in Journ. Bot.* ii. (1864) 334.—Afr. trop.
Wightiana, Muell. Arg. in Linnaea, xxxiv. (1865-66) 43 = lanceolata.
Wilkesiana, Muell. Arg. in DC. Prod. xv. II. 817 = tricolor.
zambesica, *Mucll. Arg. in Flora*, xlvii. (1864) 440.—Afr. trop.
Zeyheri, *Baill. Adansonia*, iii. (1862-63) 156.—Afr. austr.
zeylanica, *Rafin. New Fl. Amer.* i. 46, nomen.—Zeylan.
Zollingeri, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 40.—Sumbawa.

ACAMPE, Lindl. Fol. Orch. (1853).
ORCHIDEAE, Benth. & Hook. f. iii. 579.
Cephalotes, *Lindl. l. c.* 3.—Ind. or.
congesta, *Lindl. l. c.* 2.—Ind. or.
dentata, *Lindl. l. c.* 3.—Ind. or.
excavata, *Lindl. l. c.* 3.—Ind. or.
Griffithii, *Reichb. f. in Flora*, lv. (1872) 277.—Ind. or.
intermedia, *Reichb. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 217.—Hab. ?
longifolia, *Lindl. Fol. Orchid. Acampe*, 1.—Ind. or.
multiflora, *Lindl. l. c.*—China.
pachyglossa, *Reichb. f. Otia Bot. Hamb.* 76; *et in Bot. Zeit.* xxxix. (1881) 449.—Afr. trop.
papillosa, *Lindl. Fol. Orch. Acampe*, 2.—Ind. or.
Renschiana, *Reichb. f. Otia Bot. Hamb.* 77; *et in Bot. Zeit.* xxxix. (1881) 449.—Madag.
Wightiana, *Lindl. Fol. Orch. Acampe*, 2.—Ind. or.

ACAMPTOPAPPUS, A. Gray, in Proc. Am. Acad. viii. (1873) 634 = **Aplopappus**, Cass.—(Compos.).
Shockleyi, A. Gray, l. c. xvii. (1881-82) 208.—Am. bor.
sphaerocephalus, A. Gray, l. c. viii. (1873) 634.—Am. bor.

ACANOS, Adans. Fam. ii. 116 (1763) = **Onopordon**, Linn. (Compos.).
Spina, Scop. Fl. Carn. ed. II. ii. 132 = O. Acanthium.

ACANTACARYX, Arruda, ex Koster, Trav. Bras. 491 (1816) = **Caryocar**, Linn. (Ternstroem.).
pinguis, Arruda, ex Koster, l. c.—Bras.

ACANTHELLA, Hook. f. in Benth. & Hook. f. Gen. i. 748 (1867).—*MELASTOMACEAE*, Benth. & Hook. f. i. 748.
Sprucci, *Hook. f. l. c.*—Bras.

ACANTHINOPHYLLUM, Fr. Allem. in Rev. Brazil. i. (1858) 368 = **Sahagunia**, Liebm. (Urtic.).
strepitans, Fr. Allem. l. c. 369.—Bras.

ACANTHIUM, Hall. Enum. Stirp. Helv. ii. 674 (1742) = **Onopordon**, Linn. (Compos.).
Onopordon, Gueldenst. Reise Russl. ii. 69 = O. Acanthium.

ACANTHOBOTRYA, Eckl. & Zeyh. Enum. 192 (1836) = **Lebeckia**, Thunb. (Legumin.).
angustifolia, Eckl. & Zeyh. l. c. 194 = L. multiflora.
armata, Eckl. & Zeyh. l. c. 193 = L. microphylla.
cinerea, Eckl. & Zeyh. l. c. 193 = L. microphylla.
decumbens, Eckl. & Zeyh. l. c. 193 = L. humilis.
disticha, Eckl. & Zeyh. l. c. 193 = L. disticha.
linearifolia, Benth. in Ann. Wien. Mus. ii. (1838) 142 = L. linearifolia.
microphylla, Benth. l. c. = L. microphylla.
pungens, Eckl. &. Zeyh. Enum. 193 = L. pungens.
sericea, Eckl. & Zeyh. l. c. 193 = L. microphylla.
sessilifolia, Eckl. & Zeyh. l. c. 193 = L. marginata.

ACANTHOCARPEA, Klotzsch, in Peters, Reise Mossamb. Bot. 137. t. 24 (1861) = **Limeum**, Linn. (Ficoid.)
scabrida, Klotzsch, l. c. 139.—Mossamb.
sulcata, Klotzsch, l. c. 138. t. 24 = L. linifolium.

ACANTHOCARPUS, Lehm. Pl. Preiss. ii. 274 (1847).
JUNCACEAE, Benth. & Hook. f. iii. 864.
fimbriatus, *F. Muell. Census*, 119.—Austral.
Preissii, *Lehm. Pl. Preiss.* ii. 274.—Austral.
Serra, *F. Muell. Census*, 119.—Austral.

ACANTHOCARYA, Arruda, ex Endl. Gen. 1076 (1840) = **Caryocar**, Linn. (Ternstroem.).

ACANTHOCAULON, Klotzsch, in Endl. Gen. Suppl. iv. III. 88 (1850) = **Platygyne**, Mercier (Euphorb.).
pruriens, Klotzsch (sphalm. pruricus), ex Endl. l. c. = P. urens.

ACANTHOCEPHALUS, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 127. *COMPOSITAE*, Benth. & Hook. f. ii. 510.
HARPACHAENA, Bunge, Del. Sem. Hort. Dorp. (1845).
HARPOCARPUS, Endl. Gen. Suppl. iii. 10 (1843).
amplexifolius, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 128.—Sibir. altaic.
Benthamianus, *Regel, Pl. Nov. Fedsch.* 47.—Turkestan.

ACANTHOCHITON, Torr. in Sitgr. Rep. 170. t. 13 (1853). *AMARANTACEAE*, Benth. & Hook. f. iii. 29.
Wrightii, *Torr. l. c.*—N. Mexico.

ACANTHOCLADIUM, F. Muell. Fragm. ii. 155 (1861) = **Helichrysum**, Vaill. (Compos.).
Dockerii, F. Muell. l. c. 156 = H. Dockerii.

ACANTHOCLADUS, Klotzsch, ex Hassk. in Miq. Ann. Mus. Lugd. Bat. i. 184 (1864). *POLYGALEAE*, Benth. & Hook. f. i. 974.
albicans, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 46. —Bras.
brasiliensis, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Lugd. Bat.* i. 184.—Bras.
microphylla, *Griseb. in Goett. Abh.* xxiv. (1879) 24.—Reg. Argent.

ACANTHODIUM, Delile, Fl. Egypte, 241. t. 33 (1812) = **Blepharis**, Juss. (Acanth.).
angustum, Nees, in DC. Prod. xi. 273.—Afr. austr.
capense, Nees, in Linnaea, xv. (1841) 361.—Afr. austr.
carduifolium, Nees, in DC. Prod. xi. 278 = B. carduifolia.
Delilii, Buck, Ind. DC. Prod. iii. 1 = B. edulis.
dispermum, E. Mey. ex Nees, in DC. Prod. xi. 278 = Acanthopsis disperma.
diversispinum, Nees, l. c. 275.—Afr. austr.
furcatum, Nees, l. c. 276.—Afr. austr.
glabrum, Nees, l. c. 278.—Afr. austr.
glaucum, Nees, l. c. 277.—Afr. austr.
grossum, Nees, l. c. 274.—Afr. austr.
grossum, Wight, Ic. t. 1535, 1536 = B. scindica.
hirtinervium, Nees, in DC. Prod. xi. 277.—Afr. austr.
hirtum, Hochst. ex Nees, l. c. 274 = B. scindica.
Hoffmanseggianum, Nees, l. c. xi. 277.—Afr. austr.
macrum, Nees, l. c. 276.—Afr. austr.
marginatum, Nees, l. c. 275.—Afr. austr.
plumulosum, E. Mey. ex Nees, l. c. 278 = Acanthopsis horrida.
procumbens, Nees, in Linnaea, xv. (1841) 362.—Afr. austr.
serrulatum, Nees, in DC. Prod. xi. 275.—Afr. austr.
sinuatum, Nees, in Linnaea, xv. (1841) 362.—Afr. austr.
spathulare, Nees, in DC. Prod. xi. 277.—Afr. austr.
spicatum, Acerbi, ex Nees, l. c. 67 = Nelsonia canescens.
spicatum, Delile, Fl. Egypte, 241. t. 33. f. 3 = B. edulis.
squarrosum, Nees, in DC. Prod. xi. 275.—Afr. austr.

ACANTHODUS, Rafin. Princ. Somiol. 26 (1814) = **Acanthus**, Tourn. (Acanthac.).

ACANTHOGLOSSUM, Blume, Bijdr. 381 (1825) = **Pholidota**, Lindl. (Orchid.).
nervosum, Blume, l. c. 382 = P. nervosa.

ACANTHOGONUM, Torr. Pacif. Rail. Rep. iv. 132 (1856) = **Chorizanthe**, R. Br. (Polygon.).
corrugatum, Torr. & Gray, l. c. v. 364 = C. corrugata.
rigidum, Torr. l. c. iv. 133 = C. rigida.

ACANTHOLEPIS, Less. in Linnaea, vi. (1831) 88.
COMPOSITAE, Benth. & Hook. f. ii. 463.
orientalis, *Less. l. c.*—As. Min.; Persia.

ACANTHOLINUM, C. Koch, in Linnaea, xxi. (1848) 715, sphalm. = seq.

ACANTHOLIMON, Boiss. Diagn. Ser. I. vii. 69 (1846). *PLUMBAGINEAE*, Benth. & Hook. f. ii. 625.
accrosum, *Boiss. l. c.* Ser. I. vii. 80.—As. Min.
acmostegium, *Boiss. & Buhse, in Nouv. Mém. Mosc.* xii. (1860) 182.—Persia.
alatavicum, *Bunge, Gatt. Acanthol.* 40.—Sibir.
Alberti, *Regel, in Act. Hort. Petrop.* vi. (1879) 389.—Turkestan.
androsaceum, *Boiss. Diagn.* Ser. I. vii. 73.—Orient.
araxanum, *Bunge, Gatt. Acanthol.* 33.—Persia.
aristulatum, *Bunge, l. c.* 60.—Persia.
armenum, *Boiss. & Huet, Diagn.* Ser. II. iv. 64.—Armenia.
aspadanum, *Bunge, Gatt. Acanthol.* 50.—Persia.
assyriacum, Boiss. Diagn. Ser. I. vii. 81 = venustum.

ACANTHOLIMON:—
atropatanum, *Bunge, Gatt. Acanthol.* 29.—Persia.
auganum, *Bunge, l. c.*—Afghan.
avenaceum, *Bunge, l. c.* 25.—Persia.
Balansae, Boiss. ex Bunge, l. c. 36 = armenum.
baltanense, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 838.—As. Min.
Bodeanum, *Bunge, Gatt. Acanthol.* 42.—Persia.
brachyphyllum, *Boiss. in DC. Prod.* xii. 628.—Afghan.
brachystachyum, *Boiss. ex Bunge, Gatt. Acanthol.* 51.—Kurdistan.
bracteatum, *Boiss. Diagn.* Ser. I. vii. 70.—Armen.; Persia.
breviscapum, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 840 = Kotschyi.
bromifolium, *Boiss. ex Bunge, Gatt. Acanthol.* 23.—Kurdistan.
cabulicum, *Boiss. in DC. Prod.* xii. 623.—Afghan.
caesareum, *Boiss. & Bal. Diagn.* Ser. II. iv. 66.—Cappadoc.
calocephalum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1881) 77.—Afghan.
Calverti, *Boiss. Diagn.* Ser. II. iv. 65.—Armenia.
caryophyllaceum, *Boiss. Diagn.* Ser. I. vii. 78.—Kurdistan.
caryophyllaceum, Haussk. ex Bunge, Gatt. Acanthol. 37 = armenum.
cataonicum, Bunge, l. c. 43 = lycaonicum.
cephalotum, St. Lag. in Ann. Soc. Bot. Lyon. vii. (1880) 118 = A. Cephalotes.
Cephalotes, *Boiss. in DC. Prod.* xii. 622.—Afghan.
curviflorum, *Bunge, Gatt. Acanthol.* 57.—Persia.
cymosum, *Bunge, l. c.* 15.—Persia.
desertorum, *Regel, in Act. Hort. Petrop.* vi. (1879) 391.—Turkestan.
diapensioides, *Boiss. in DC. Prod.* xii. 624.—Afghan.
distachyum, *Boiss. l. c.* 623.—Afghan.
Echinus, Bunge, Gatt. Acanthol. 46 = androsaceum.
erythraeum, *Bunge, in Act. Hort. Petrop.* iii. (1875) 99.—Turkestan.
eschkerense, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 844.—Persia.
fasciculare, *Boiss. in DC. Prod.* xii. 626.—Afghan.
Fausti, *Trautv. in Act. Hort. Petrop.* ii. (1873) 482.—Persia.
ferox, Boiss. Diagn. Ser. I. vii. 81 = A. Scorpius.
Fetisowi, *Regel, Dec. Pl. Nov.* 8.—Turkest.
festucaceum, *Boiss. Diagn.* Ser. I. vii. 76.—Persia.
flexuosum, *Boiss. ex Bunge, Gatt. Acanthol.* 51.—Persia.
genistoides, *Boiss. Diagn.* Ser. I. vii. 76.—Persia.
glumaceum, *Boiss, l. c.* 75.—Armenia.
Griffithianum, *Boiss. in DC. Prod.* xii. 623.—Afghan.
gulistanum, *Bunge, Gatt. Acanthol.* 67.—Persia.
Hausknechti, Bunge, l. c. 37 = armenum.
heratense, *Bunge, l. c.* 55.—Afghan.
Hohenackeri, *Boiss. Diagn.* Ser. I. vii. 75.—Reg. Caucas.
Hohenackeri, Ledeb. Fl. Ross. iii. 470 = glumaceum.
horridum, *Bunge, Gatt. Acanthol.* 64.—Persia.
Huetii, *Boiss. Diagn.* Ser. II. iv. 65.—Armenia.
Hystrix, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 71.—Persia.
iconicum, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iv. 839.—As. Min.
incomptum, Boiss. & Buhse, in Nouv. Mém. Mosc. xii. (1860) 183 = A. Scorpius.
Karelinii, *Bunge, Gatt. Acanthol.* 58.—Armenia.
kokandense, *Bunge, in Act. Hort. Petrop.* iii. (1875) 99.—Turkest.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 74.—Cilicia.
kurdicum, Bunge, Gatt. Acanthol. 52 = brachystachyum.
latifolium, *Boiss. Diagn.* Ser. II. iv. 61.—Kurdistan.
latifolium, Rupr. Sert. Thiansch. 69 = Ruprechti.
laxiflorum, Boiss. Fl. Orient. iv. 832 = venustum.
laxiflorum, *Boiss. ex Bunge, Gatt. Acanthol.* 28.—Syria.
leptostachyum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1881) 76.—Afghan.
lepturoides, *Boiss. Diagn.* Ser. I. vii. 77.—Reg. Caucas.
leucacanthum, Boiss. l. c. 81 = A. Scorpius.
libanoticum, *Boiss. in DC. Prod.* xii. 630.—Syria.
Listoniae, Boiss. l. c. 631 = acerosum.
longiflorum, *Boiss. Diagn.* Ser. II. iv. 62.—Afghan.
lycaonicum, *Boiss. & Heldr. ex Boiss. in DC. Prod.* xii. 627.—As. Min.

ACANTHOLIMON:—
lycopodioides, *Boiss. in DC. Prod.* xii. 632.—As. centr.; Kashmir.
Maewskianum, *Regel, in Act. Hort. Petrop.* vi. (1879) 391.—Samarkand.
melananthum, *Boiss. Diagn.* Ser. I. vii. 74.—Persia.
Munroanum, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1881) 76.—Afghan.
oliganthum, *Boiss. Diagn.* Ser. I. vii. 77.—Persia.
Olivieri, *Boiss. l. c.* 80.—Persia.
parviflorum, *Regel, Pl. Nov. Fedsch.* 73.—As. temp.
Peronini, *Boiss. Fl. Orient.* iv. 842.—Cilicia.
petraeum, *Boiss. ex Bunge, Gatt. Acanthol.* 28.—Kurdistan.
phrygium, Boiss. Diagn. Ser. I. vii. 79 = acerosum.
Pinardi, Boiss. l. c. = acerosum.
polystachyum, *Boiss. in DC. Prod.* xii. 624.—Afghan.
pterostegium, *Bunge, Gatt. Acanthol.* 15.—Persia.
puberulum, Boiss. & Bal. Diagn. Ser. II. iv. 62 = androsaceum.
quinquelobum, *Bunge, Gatt. Acanthol.* 56.—Persia.
restiaceum, *Bunge, l. c.* 22.—Persia.
roseum, *Boiss. Diagn.* Ser. I. vii. 71.—Persia; Afghan.
rubellum, Boiss. ex Bunge, Gatt. Acanthol. 66 = roseum.
Ruprechti, *Bunge, Gatt. Acanthol.* 20.—Turkestan.
sahendicum, *Boiss. & Buhse, in Nouv. Mém. Mosc.* xii. (1860) 183.—Persia.
sarawschanicum, *Regel, Pl. Nov. Fedsch.* 73.—As. temp.
scabrellum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 845.—Persia.
schahrudicum, *Bunge, Gatt. Acanthol.* 65.—Persia.
schirazianum, *Boiss. Diagn.* Ser. I. vii. 72.—Persia.
scirpinum, *Bunge, Gatt. Acanthol.* 22.—Persia.
scorpioideum, St. Lag. in Ann. Soc. Bot. Lyon. vii. (1880) 118 = A. Scorpius.
Scorpius, *Boiss. Diagn.* Ser. I. vii. 81.—Persia.
senganense, *Bunge, Gatt. Acanthol.* 29.—Persia.
setiferum, *Bunge, l. c.* 68.—Turkestan.
splendidum, Bunge, l. c. 17 = bracteatum.
Stocksii, *Boiss. Diagn.* Ser. II. iv. p. 63.—Beluchistan.
subulatum, *Boiss. in DC. Prod.* xii. 629.—Afghan.
Szovitzii, Boiss. & Buhse, in Nouv. Mém. Mosc. xii. (1860) 184 = Karelinii.
talagonicum, *Boiss. in DC. Prod.* xii. 625.—Persia.
tataricum, *Boiss. l. c.* 623.—Turkestan.
Tchihatcheffi, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. (1854) i. 30 = acerosum.
tenuiflorum, *Boiss. Diagn.* Ser. I. vii. 78.—Trans Caucas.
tibeticum, Hook. f. & Thoms. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 479 = lycopodioides.
tomentellum, *Boiss. Diagn.* Ser. I. vii. 72.—Persia.
Tournefortii, *Boiss. l. c.* 74.—Armenia.
tragacanthinum, *Boiss. l. c.* 81.—Persia.
tragacanthinum, Griff. ex Bunge, Gatt. Acanthol. 64 = fasciculare.
truncatum, *Bunge, l. c.* 56.—Persia.
ulicinum, *Boiss. in DC. Prod.* xii. 627.—Syria.
venustum, *Boiss. Diagn.* Ser. I. vii. 80.—As. Min.
viscidulum, *Boiss. l. c.* 71.—Persia.
Wiedemanni, *Bunge, Gatt. Acanthol*, 45.—Anatolia.

ACANTHOLIPPIA, Griseb. in Goett. Abh. xix. (1874) 244 = **Lippia**, Linn. (Verben.).
hastulata, Griseb. l. c. xxiv. (1879) 279 = L. hastulata.
salsoloides, Griseb. l. c. xix. (1874) 244.—Reg. Argent.

ACANTHOLOMA, Gaudich. ex Baill. Adansonia, vi. (1866) 231. t. i = **Pachystroma**, Klotzsch (Euphorb.).
spinosum, Baill. l. c. 237 = P. ilicifolium.

ACANTHOMINTHA, A. Gray, in Syn. Fl. N. Am. ii. 365 (1878).—*LABIATAE*, Benth. & Hook. f. ii. 1192.
ilicifolia, *A. Gray, l. c.*—Calif.
lanceolata, *Curran, in Bull. Calif. Acad.* i. (1884) 13.—Calif.

ACANTHONEMA, Hook. f. Bot. Mag. t. 5339 (1862). *GESNERACEAE*, Benth. & Hook. f. ii. 1021.
strigosum, *Hook. f. l. c.*—Afr. trop.

ACANTHONOTUS, Benth. in Hook. Nig. Fl. 293 (1849) = **Indigofera**, Linn. (Legum.).
echinatus, Benth. l. c. = I. echinata.

ACANTHONYCHIA, Rohrb. in Mart. Fl. Bras. xiv. ii. 249. t. 56 (1872) = **Pentacaena**, Bartl. (Illecebr.).
polycnemoides, Rohrb. l. c. 250 = P. polycnemoides.
ramosissima, Rohrb. l. c. 249. t. 56 = P. polycnemoides.

ACANTHOPANAX, Miq. Ann. Mus. Lugd. Bat. i. 10 (1863). *ARALIACEAE*, Benth. & Hook. f. i. 939.
KALOPANAX, Miq. l. c. (1863).
PLECTRONIA, Lour. Fl. Cochinch. 162 (1790).
aculeatum, *Seem. in Journ. Bot.* v. (1867) 238.—As. trop.
asperatum, Franch. & Sav. Enum. Pl. Jap. ii. 378 = divaricatum.
divaricatum, *Seem. in Journ. Bot.* v. (1867) 239.—Japon.
innovans, *Franch. & Sav. Enum. Pl. Jap.* i. 193.—Japon.
japonicum, *Franch. & Sav. l. c.* ii. 377.—Japon.
ricinifolium, *Seem. in Journ. Bot.* vi. (1868) 140.—Japon.
sciadophylloides, *Franch. & Sav. Enum. Pl. Jap.* ii. 378.—Japon.
sepium, Seem. in Journ. Bot. v. (1867) 239 = aculeatum.
sessiliflorum, *Seem. l. c.*—China.
spinosum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 10.—Japon.
trichodon, *Franch. & Sav. Enum. Pl. Jap.* ii. 377.—Japon.

ACANTHOPHIPPIUM, Blume, Bijdr. 353. t. 47 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 512.
bicolor, *Lindl. Bot. Reg.* xx. (1835) 1730.—Zeylan.
Curtisii, *Reichb. f. in Gard. Chron. N. S.* xv. (1881) 1. 169.—Malaya.
javanicum, *Blume, Bijdr.* 354.—Java.
parviflorum, *Hassk. Cat. Bog. Alt.* 43.—Malaya.
ringiflorum, Griff. Notul. iii. 347 = sylhetense.
striatum, *Lindl. Bot. Reg.* (1838) *Misc.* 41.—Reg. Himal.
sylhetense, *Lindl. Gen. et Sp. Orch.* 177.—Reg. Himal.

ACANTHOPHOENIX, H. Wendl. in Fl. des Serres. Sér. II. vi. (1865–67) 181. t. 1706. *PALMEAE*, Benth. & Hook. f. iii. 898.
DECKENIA, H. Wendl. ex Seem. in Gard. Chron. (1870) 561.
crinita, *H. Wendl. l. c.*—Ins. Mascarens.
rubra, *H. Wendl. l. c.* sub t. 1706.—Ins. Mascarens.

ACANTHOPHYLLUM, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 37 = **Nassauvia**, Juss. (Compos.).
axillare, Hook. & Arn. l. c. (= Strongyloma glomerulosa).
rosulatum, Hook. & Arn. l. c. 43 = N. rosulata.

ACANTHOPHYLLUM, C. A. Mey. Verz. Pfl. Cauc. 210 (1831). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 145.
ALLOCHRUSA, Bunge, in Boiss. Fl. Orient. i. 559 (1867).
JORDANIA, Boiss. Diagn. Ser. I. viii. 93 (1849).
Alexandrae, *Smirn. ex Regel, in Act. Hort. Petrop.* v. (1877) 250.—Reg. Amur.
bracteatum, *Boiss. Diagn.* Ser. I. i. 43.—Persia.
Bungei, *Trautv. in Act. Hort. Petrop.* ii. (1873) 511.—Persia.
caespitosum, *Boiss. Diagn.* Ser. I. i. 42.—Persia.
crassifolium, *Boiss. l. c.*—Persia.
elatius, *Bunge, in Boiss. Fl. Orient.* i. 561.—Persia; Turkest.
Fontanesii, Boiss. & Buhse, in Nouv. Mém. Mosc. xii. (1860) 39 = glandulosum.
Fontanesii, *Boiss. Diagn.* Ser. I. i. 41.—Persia.
glandulosum, *Bunge, ex Boiss. Fl. Orient.* i. 565.—Persia.
glaucum, Bunge, ex Boiss. l. c. 566 = sordidum.
gracile, *Bunge, ex Boiss. l. c.* 562.—Persia.
grandiflorum, *Stocks, in Hook. Kew Journ.* iv. (1852) 142.—Baluchist.
gypsophiloides, *Regel, Pl. Nov. Fedsch.* 15.—Turkest.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 249.—Turkest.
laxiflorum, Boiss. Diagn. Ser. II. i. 182 = Gypsophila erinacea.
macrodon, *Edgew. in Hook. Fl. Brit. Ind.* i. 216.—Ind. or.; Afghan.
microcephalum, *Boiss. Diagn.* Ser. I. i. 43.—Persia.
mite, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. (1854). i. 35. t. 1.—As. Min.

ACANTHOPHYLLUM:—
mucronatum, *C. A. Mey. Verz. Pfl. Cauc.* 210.—Reg. Caucas.
mucronatum, Fenzl, in Ann. Wien. Mus. i. (1836) 37 = Tournefortii.
paniculatum, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) i. 539.—Turkest.
pungens, Boiss. Fl. Orient. i. 561 = spinosum.
sordidum, *Bunge, ex Boiss. Fl. Orient.* i. 565.—Persia.
spinosum, Bunge, in Mém. Sav. étr. Pétersb. vii. (1851) 227 = elatius.
spinosum, *C. A. Mey. Verz. Pfl. Cauc.* 210.—Reg. Caucas.; Sibir.
squarrosum, *Boiss. Diagn.* Ser. II. i. 81.—Persia.
Stocksianum, Boiss. l. c. = squarrosum.
Tournefortii, *Fenzl, in Ann. Wien. Mus.* ii. (1838) 310.—Armenia.
versicolor, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 31.—Persia.
verticillatum, *C. A. Mey. Verz. Pfl. Cauc.* 210.—Armenia.

ACANTHOPHYTON, Less. Syn. Comp. 128 (1832) = **Cichorium**, Linn. (Compos.).
spinosum, Less. l. c. = C. spinosum.

ACANTHOPLEURA, C. Koch, in Bot. Zeit. vii. (1849) 408 = **Polylophium**, Boiss. (Umbell.).
involucrata, C. Koch, l. c. = P. orientale.

ACANTHOPRASIUM, Spenn. in T. Nees, Gen. Fl. Germ. Gamop. ii. (1843) = **Ballota**, Linn. (Labiat.).
frutescens, Spenn. l. c. = B. spinosa?

ACANTHOPSIS, Harv. in Hook. Lond. Journ. Bot. i. (1842) 28. *ACANTHACEAE*, Benth. & Hook. f. ii. 1089.
disperma, *Harv. l. c.*—Afr. austr.
horrida, *Nees, in DC. Prod.* xi. 278.—Afr. austr.

ACANTHORHIZA, H. Wendl. in Kerch. Palmiers, 230 (1878); Bot. Zeit. (1879) 147. *PALMAE*, Benth. & Hook. f. iii. 925.
aculeata, *H. Wendl. in Kerch. Palmiers*, 230; *Illust. Hortic.* xxvi. (1879) t. 367.—Mexico.
Chuco, *Drude, in Mart. Fl. Bras.* iii. ii. 554.—Bras.
Mocinni, *Benth. & Hook. f. Gen.* iii. 925.—Mexico.
stauracantha, H. Wendl. ex Linden, Cat. n. 87 (1871); Illust. Hortic. xxviii. (1881) 15 = aculeata.
Wallisi, *H. Wendl. ex Regel, in Gartenfl.* xxviii. (1879) 163. t. 977. f. 2.—Hab.?
Warscewiczii, *H. Wendl. in Kerch. Palmiers*, 230.—Panama.

ACANTHOSICYOS, Welw. in Trans. Linn. Soc. xxvii. (1869) 30. *CUCURBITACEAE*, Benth. & Hook. f. i. 824.
horrida, *Welw. l. c.* 31.—Afr. trop.

ACANTHOSPERMA, Vell. Fl. Flum. viii. t. 152 (1827) = **Acicarpha**, Juss. (Calycer.).
littorale, Vell. l. c. = Acicarpha spathulata.

ACANTHOSPERMUM, Schrank, Pl. Rar. Hort. Monac. t. 53 (1819). *COMPOSITAE*, Benth. & Hook. f. ii. 349.
CENTROSPERMUM, H. B. & K. Nov. Gen. et Sp. iv. 270. t. 397 (1820).
ECHINODIUM, Poiteau, ex Cass. in Dict. Sc. Nat. lix. 235 (1829).
ORCYA, Vell. Fl. Flum. 344 (1825).
brasilum, *Schrank, Pl. Rar. Hort. Monac.* ii. 53.—Bras.
hirsutum, DC. Prod. v. 522 = brasilum.
hispidum, *DC. l. c.* 522.—Bras.
humile, *DC. l. c.* 522.—Ind. occ.
xanthioides, DC. l. c. 521 = brasilum.

ACANTHOSPORA, Spreng. Anleit. ii. i. 255 (1817); Syst. ii. 25 (1830) = **Tillandsia**, Linn. (Bromeliac.).
conantha, Spreng. Syst. ii. 25 = T. strobilantha.
juncea, Spreng. l. c. 25 = T. setacea.
vittata, D. Dietr. Syn. Pl. ii. 1056 = Cryptanthus bivittatus?

ACANTHOSTACHYS, Link, Klotzsch & Otto, Ic. Pl. Rar. i. 21. t. 9 (1841). *BROMELIACEAE*, Benth. & Hook. f. iii. 663.
strobilacea, *Link, Klotzsch, & Otto, l. c.* 22. t. 9.—Bras.

ACANTHOSTEMMA, Blume, Rumphia, iv. 29 (1848) = **Hoya**, R. Br. (Asclepiad.).
Hasseltii, Blume, l. c. = H. Hasseltii.
Kuhlii, Blume, l. c. = H. longifolia.
longifolium, Blume, l. c. = H. Rumphii.
pictum, Blume, l. c. = H. picta.
pruinosum, Blume, l. c. 30 = H. pruinosa.
puberum, Blume, l. c. 29 = H. pubera.
Rumphii, Blume, l. c. 29 = H. Rumphii.

ACANTHOSYRIS, Griseb. in Goett. Abh. xxiv. (1879) 151. *SANTALACEAE*, Benth. & Hook. f. iii. 224.
falcata, *Griseb. l. c.*—Reg. Argent.
spinescens, *Griseb. l. c.*—Reg. Argent.

ACANTHOTHECA, DC. Prod. vi. 73 (1837) = **Dimorphotheca**, Moench (Compos.).
dentata, DC. l. c. 74 = D. dentata.
integrifolia, DC. l. c. 74 = D. integrifolia.
pinnatisecta, DC. l. c. 73 = D. pinnata.

ACANTHOXANTHIUM, Fourr. in Ann. Soc. Linn. Lyon. N.S. xvii. (1869) 110 = **Xanthium**, Tourn. (Compos.).
spinosum, Fourr. l. c. 110 = X. spinosum.

ACANTHUS, Tourn. ex Linn. Syst. ed. I. (1735).
ACANTHACEAE, Benth. & Hook. f. ii. 1090.
ACANTHODUS, Rafin. Princ. Somiol. 26 (1814).
CHEILOPSIS, Moq. in Ann. Sc. Nat. Sér. I. xxvii. (1832) 230.
DILIVARIA, Juss. Gen. 103 (1789).
ZONABLEPHIS, Rafin. Sylva Tellur. 143 (1838).
arboreus, *Forsk. Fl. Aegypt. Arab.* 115.—Aegypt.
Barteri, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 37.—Afr. trop.
capensis, *Thunb. Prod. Pl. Cap.* 97.—Afr. austr.
carduaceus, *Griff. Notul.* iv. 138.—Reg. Himal.
carduifolius, Linn. f. Suppl. 294 = Blepharis carduifolia.
ciliaris, Burm. f. Fl. Ind. 139 = Blepharis boerhaviaefolia.
cristatus, Rottl. ex Nees, in DC. Prod. xi. 256 = Lepidagathis cristata.
Delilei, Spreng. Syst. iii. 819 = Blepharis edulis.
Dioscoridis, *Linn. Cent.* ii. 23 (*Am. Acad.* iv. 321).—Syria.
Doloarin, Blanco, Fl. Filip. ed. I. 487 = ilicifolius.
ebracteatus, *Vahl, Symb. Bot.* ii. 75. t. 40.—As. trop.
edulis, Forsk. Fl. Aegypt. Arab. 115 = Blepharis edulis.
furcatus, Linn. f. Suppl. 295 = Blepharis furcata.
glaber, E. Mey. ex Nees, in DC. Prod. xi. 278 = Blepharis carduifolia.
glaucus, E. Mey. ex Nees, l. c. 277 = Blepharis glauca.
glomeratus, Lam. Encyc. i. 23 = Blepharis glomerata.
hirsutus, *Boiss. Diagn.* Ser. I. iv. 86.—Oriens.
hispanicus, Hort. ex Loud. Hort. Brit. 256 = niger.
humilis, Vahl, ex Nees, in DC. Prod. xi. 274 (= Acanthodium sinuatum.)
ilicifolius, *Linn. Sp. Pl.* 639 (err. typ. 939).—As. trop.; Austral.
ilicifolius, Lour. Fl. Cochinch. 375 = ebracteatus.
imbricatus, *Edgew. in Journ. As. Soc. Beng.* xvi. (1847) 217.—Ind. or.
integrifolius, E. Mey. ex Nees, in DC. Prod. xi. 276 = Blepharis furcata.
integrifolius, Thunb. Prod. Pl. Cap. 97 = Blepharis saturaejaefolia.
Kirkii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 37.—Angola.
latifolius, Hort. ex Goeze, in Gard. Chron. (1867) 856, 903 = mollis.
leucostachyus, *Wall. Cat.* n. 2512.—Ind. or.
longibracteatus, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 79.—Burma.
longifolius, *Poir. Encyc. Suppl.* i. 88.—Europ. austr.
longipetiolatus, *Kurz, For. Fl.* ii. 241.—Burma.
lusitanicus, Hort. cf. Gard. Chron. (1867) 903 = mollis.
macer, E. Mey. ex Nees, in DC. Prod. xi. 276 = Blepharis furcata.
maderaspatanus, Vahl, ex Nees, in DC. Prod. xi. 267 = Blepharis boerhaviaefolia.
maderaspatensis, Linn. Sp. Pl. 639 (err. typ. 939) = Blepharis molluginifolia.
mollis, Graf & Noé, ex Nees, in DC. Prod. xi. 270 = longifolius.
mollis, *Linn. Sp. Pl.* 639 (err. typ. 939).—Europ. austr.

ACANTHUS:—
mollis, Riedel, ex Nees, in DC. Prod. xi. 271 = spinosus.
montanus, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 37.—Afr. trop.
mucronatus, Heyne, ex C. B. Clarke, in Hook. f. Fl. Br. Ind. iv. 517 = Lepidagathis spinosa.
niger, *Mill. Gard. Dict.* ed. VIII. n. 2.—Lusitan.
nitidus, *S. Moore, in. Journ. Bot.* xviii. (1880) 232.—Afr. trop.
pectinatus, Willd. ex Nees, in DC. Prod. xi. 274 = Blepharis edulis.
polystachyus, *Delile, in Caill. Voy. au Méroé*, 2. t. 62. f. 2.—Afr. trop.
procumbens, Herb. Madr. ex Wall. Cat. sub n. 2513 = Blepharis boerhaviaefolia.
procumbens, Thunb. Prod. Pl. Cap. 97 = Blepharis glomerata?
pungens, Herb. Madr. ex Wall. Cat. n. 2498 = Lepidagathis pungens.
Raddei, Trautv. in Act. Hort. Petrop. iii. ii. (1875) 269 = Dioscoridis.
repandus, Nees, in DC. Prod. xi. 435, sphalm. = Anthacanthus repandus.
repens, Vahl, Symb. Bot. ii. 76 = Blepharis molluginifolia.
rubens, *Rafin. Fl. Ludov.* 36.—Am. bor.
spathularis, *E. Mey. ex Nees, in. DC. Prod.* xi. 277.—Am. bor.
spinosissimus, Desf. Tabl. Hort. Par. 51 = spinosus.
spinosissimus, Host, Fl. Austr. ii. 218 = mollis.
spinosus, *Linn. Sp. Pl.* 639 (err. typ. 939).—Europ.
spinosus, Nees, in DC. Prod. xi. 271 = hirsutus, syriacus
spinulosus, Host, Fl. Austr. ii. 217 = spinosus.
syriacus, *Boiss. Diagn.* Ser. I. xi. 135.—Syria.
tetragonus, *R. Br. in Salt, Abyss. App.* 65; *Verm. Schr. ed. Nees*, iii. 249.—Abyss.
verus, *Wender. ex Steud. Nom.* ed. II. i. 11.—Hab.?
volubilis, *Wall. Pl. As. Rar.* ii. 56. t. 172.—Ind. or.

ACANTHYLLIS, Pomel, Nouv. Mat. Fl. Atl. 179 (1874) = **Anthyllis**, Riv. (Legum.).
numidica, Pomel, l. c. = Anth. numidica.
tragacanthoides, Pomel, l. c. = Astragalus armatus.

ACARNA, All. Fl. Pedem. i. 153 (1785) = **Atractylis**, Linn. (Compos.).
caespitosa, Willd. Sp. Pl. iii. 1700 = Atractylis caespitosa.
cancellata, All. Fl. Pedem. i. 153 = Atractylis cancellata.
chinensis, Bunge, Enum. Pl. Chin. Bor. 36 = Atractylis ovata.
comosa, Spreng. Syst. iii. 380 = Atractylis comosa.
cyanoides, Bess. Enum. Pl. Volh. 32 = Jurinea cyanoides.
cyanoides, Wibel, Prim. Fl. Werth. 169 = Jurinea Pollichii.
gummifera, Willd. Sp. Pl. iii. 1699 = Carlina gummifera.
lancea, Willd. l. c. 1700 = Atractylis ovata.
macrocephala, Willd. l. c. = Carlina Fontanesii.
macrophylla, Willd. l. c. = Carlina macrophylla.
mollis, Bess. Enum. Pl. Volh. 32 = Jurinea mollis.
ovata, Willd. Sp. Pl. iii. 1701 = Atractylis ovata.
rigida, Godet, ex DC. Prod. vi. 674 = Jurinea longifolia.
serratuloides, Spreng. Syst. iii. 379 = Atractylis serratuloides.

ACARNA, Hill, Veg. Syst. iv. 13 (1762) = **Cnicus**, Linn. (Compos.).
Casabonae, Hill, Hort. Kew. 59 = C. Casabonae.
leucographa, Hill, l. c. = Tyrimnus leucographus.
onopyxe, Hill, l. c. = Carduus Onopyxos.

ACARPHA, Griseb. in Goett. Abh. vi. (1853–55) 125 = **Boopis**, Juss. (Calycer.).
australis, Griseb. l. c. 126 = B. australis.

ACARPHAEA, Harv. & Gray, ex A. Gray, in Mem. Am. Acad. iv. (1849) i. (Pl. Fendl.) 98 = **Chaenactis**, DC. (Compos.).
artemesiaefolia, Harv. & Gray, l. c. = C. artemisiaefolia.

ACASTE, Salisb. in Trans. Hort. Soc. i. (1812) 322 = **Babiana**, Ker-Gawl. (Irid.).
pulchra, Salisb. l. c. 323 = B. stricta.
venusta, Salisb. l. c. 322 = B. rubrocyanea.

ACCA, Berg, in Linnaea, xxvii. (1855) 138 = **Psidium**, Linn. (Myrtac.).
domingensis, Berg, l. c. 138.—Ind. occ.
peruviana, Berg, l. c. 139.—Peruv.

ACCIA, St. Hil. Morphol. 499 (1840) = **Plukenetia**, Linn. (Euphorb.).
scandens, St. Hil. l. c. (= Fragariopsis scandens).

ACCIPITRINA, Riv. ex Rupp. Fl. Jen. ed. Hall. 81 (1745) = **Sisymbrium**, Linn. (Crucif.).

ACCOROMBONA, Endl. Gen. Suppl. i. 1427 (1841) = **Galega**, Linn. (Legum.).
tricolor, Benth. ex Walp. Rep. i. 672 = G. officinalis.

ACEDILANTHUS, Benth. & Hook. f. Gen. iii. 835 (1881) sphalm. = seq.

ACELIDANTHUS, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 94. t. 28) (1847) = **Veratrum**, Linn. (Liliac.).
anticleoides, Trautv. & Mey. l. c. = V. Maackii.

ACENTRA, Phil. Sert. Mendoc. Alt. 3 (1871) *VIOLARIEAE*.
serrata, *Phil. l. c.*—Reg. Argent.

ACER, [Tourn.] Linn. Syst. ed. I. (1735). *SAPINDACEAE*, Benth. & Hook. f. i. 409.
EUACER, Opiz, Seznam, 42 (1852).
abchasicum, *Rupr. Fl. Cauc.* 279.—Reg. Caucas.
acuminatum, *Wall. ex D. Don, Prod. Fl. Nep.* 249.—Reg. Himal.
aetnense, Hort. ex C. Koch, Dendrol. 531 = platanoides.
affine, Hoffmgg. ex Walp. Rep. i. 410 = campestre.
affine, Opiz, in Flora, vii. (1824) Beil. i. 83; xii. (1829) 547 = campestre.
Amaliae, *Orphan. ex Boiss. Fl. Orient.* i. 951.—Graecia.
amoenum, Carr. in Rev. Hortic. (1867) 280 = palmatum.
argutum, *Maxim. ex Miq. in Arch. Néerl.* ii. (1867) 475.—Japon.
austriacum, Tratt. Arch. Gew. i. t. 36 = campestre.
barbatum, Hook. Fl. Bor. Am. i. 113 = glabrum.
barbatum, *Michx. Fl. Bor. Am.* ii. 252 = saccharinum.
barbinerve, *Maxim. ex Miq. in Arch. Néerl.* ii. (1867) 476.—Japon.
bohemicum, Presl, ex Opiz, Seznam, 9 = campestre.
Boscii, *Spach, Hist. Vég. Phan.* iii. 88.—Am. bor.?
Buergerianum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 88 = palmatum.
Buzimpala, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 249 = oblongum.
caesium, *Wall. ex Brand. For. Fl.* iii. t. 21.—Reg. Himal.
californicum, D. Dietr. Syn. ii. 1283 = Negundo aceroides.
californicum, *Torr. & Gray, Fl. N. Am.* i. 250.—Calif.
Campbellii, *Hook. f. & Thoms., ex Brand. For. Fl.* 109.—Reg. Himal.
campestre, *Linn. Sp. Pl.* 1055.—Europ.; Oriens.
canadense, Duham. Arb. i. t. 12 = pensylvanicum.
capillipes, *Maxim. ex Miq. in Arch. Néerl.* ii. (1867) 477.—Japon.
carolinianum, Walt. Fl. Carol. 251 = rubrum?
carpinifolium, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1845) 154.—Japon.
cassiaefolium, *Blume, Rumphia*, iii. 193. t. 167. B. f. 2.—Java.
caudatum, *Wall. Pl. As. Rar.* ii. 4.—Reg. Himal.
cinerascens, *Boiss. Diagn.* Ser. I. vi. 29.—Persia.
circinnatum, *Pursh, Fl. Am. Sept.* i. 267.—Amer. bor.
circumlobatum, *Maxim. in Bull. Acad. Pétersb.* xii. (1867) 225.—Japon.
cissifolium, *C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 252.—Japon.
coccineum, Michx. f. Arb. Am. i. 203 = rubrum.
colchicum, Hartwiss, in Loud. Gard. Mag. (1840) 632 = pictum.
Collinsona, *Thunb. Diss. Acere*, 6.—Hab.?
collinum, Tenore, Syll. 194 = campestre.
commutatum, J. & C. Presl, Delic. Prag. 31 = monspessulanum.
cordifolium, Moench, Meth. 55 = tatarium.
coriaceum, Lodd. ex Spach, Hist. Vég. Phan. iii. 107 = polymorphum.
coriaceum, *Bosc, ex Tausch, in Flora*, xii. (1829) 550.—Hab.?
crataegifolium, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1845) 155.—Japon.

ACER:—
creticum, *Linn. Sp. Pl.* ed. II. 1497.—Oriens.
creticum, Schmidt, ex Tratt. Arch. Gew. i. t. 45 = parvifolium.
crispum, *Moench, Verz. Ausl. Baeume*, 2.—Hab.?
cultratum, Wall. Pl. As. Rar. ii. 4 = pictum.
dasycarpum, *Ehrh. Beitr.* iv. 24.—Am. bor.
Dedyle, Maxim. in Bull. Phys. Math. Acad. Sc. Pétersb. xv. (1856) 125 = spicatum.
diabolicum, *Blume, ex Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 251.—Japon.
discolor, *Maxim. in Bull. Acad. Sc. Pétersb.* xxvi. (1880) 436.—China.
dissectum, Thunb. Fl. Jap. 160 = palmatum.
distylum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1845) 154.—Japon.
Dittrichii, Opiz, Seznam, 9 = A. Pseudo-Platanus.
Douglasii, Hook. Lond. Journ. Bot. vi. (1846) 77. t. 6 = glabrum.
Drummondii, Hook. & Arn. in l. c. i. (1834) 200 = rubrum.
eriocarpon, Opiz, in Flora, vii. (1824) i. Beil. 83 = campestre.
eriocarpum, Michx. Fl. Bor. Am. ii. 253 = dasycarpum.
Fabri, *Hance, in Journ. Bot.* xxii. (1884) 76.—China.
Fieberi, *Opiz, Seznam*, 9.—Bohem.
floridanum, Hort. ex C. Koch, Dendrol. 542 = dasycarpum.
formosum, *Carr. in Rev. Hortic.* (1867) 300.—Japon.
fraxinifolium, Nutt. Gen. Am. i. 253 = Negundo aceroides.
fraxinifolium, *Rafin. New Fl. Am.* i. 48.—Am. bor.
Friederici-Guillelmi, Carr. in Rev. Hortic. (1867) 300 = palmatum.
Ginnale, Maxim. in Bull. Phys. Math. Acad. Sc. Pétersb. xv. (1856) 126 = tataricum.
glaucum, Hort. ex C. Koch, Dendrol. 543 = sanguineum.
glaucum, Marsh. Arbust. 2 = rubrum?
glabrum, *Torr. in Ann. Lyc. New York*, ii. (1828) 172.—Am. bor.
granatense, Boiss. Elench. 25 = opulifolium.
grandidentata, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 249.—Am. bor.
Heldreichii, *Boiss. & Heldr. in Boiss. Diagn.* Ser. II. v. 71.—Graecia.
heterolobum, Opiz, Natural. ix. (1825) 114 = campestre.
heterophyllum, Willd. Berl. Baumz. 10. t. 1 = creticum.
hispanicum, Pourr. in Mém. Acad. Toul. iii. (1788) 305 = opulifolium.
Hookeri, *Miq. in Arch. Néerl.* ii. (1867) 471.—Reg. Himal.
humile, Salisb. Prod. 281 = creticum.
hybridum, Bosc, Dict. Agr. v. 251 = pensylvanicum.
hypoleucum, Hort. ex C. Koch, Dendrol. 543 = rubrum.
hyrcanum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 31.—Oriens.
ibericum, Bieb. Fl. Taur. Cauc. ii. 447 = monspessulanum.
illyricum, Jacq. f. in Flora, xii. (1827) 551 = monspessulanum.
insigne, *Boiss. & Buhse, in Nouv. Mém. Mosc.* xii. (1860) 46.—Persia.
intermedium, Panč. ex Nym. Consp. 135 = monspessulanum.
isolobum, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 304.—Burma.
italum, *Lauth, de Acere*, n. 8.—Europ.; Oriens.
japonicum, *Thunb. Fl. Jap.* 161.—Japon.
javanicum, Jungh. in Hoev. & De Vr. Tijdschr. viii. (1841) 391 = niveum.
javanicum, *Burm. f. Fl. Ind.* 221.—Java.
jucundum, *Carr. in Rev. Hortic.* (1867) 380.—Japon.
laciniatum, Borckh. ex Tratt. Arch. Gew. 35 = platanoides.
lactescens, Pers. Syn. i. 417 = platanoides.
laetum, *C. A. Mey. Verz. Pfl. Cauc.* 236.—Oriens.
laevigatum, *Wall. Pl. As. Rar.* ii. 3. t. 104.—Reg. Himal.
laurifolium, D. Don, Prod. Fl. Nep. 249 = oblongum.
laurinum, *Hassk. in Hoev. & De Vr. Tijdschr.* x. (1843) 138 = niveum.
laurinum, Hassk. l. c. = niveum.
leiocarpon, Opiz, in Flora, vii. (1824) i. Beil. 82 = campestre.
leptopterum, Guss. ex Nym. Consp. 135 = opulifolium.

ACER:—
lobatum, Bosc, ex C. Koch, Dendrol. 519 = Boscii.
lobatum, *Rafin. New Fl. Am.* i. 48.
Lobelii, Tenore, Fl. Nap. v. 29. t. 199 = platanoides.
macrocarpon, Opiz, in Flora, vii. (1824) i. Beil. 82 = campestre.
macrophyllum, *Pursh, Fl. Am. Sept.* i. 267.—Am. bor.
macropterum, Vis, in Mem. Ist. Venet. ix. (1860) 175; Suppl. 28 = Visianii.
majus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 635 = A. Pseudo-Platanus.
mandshuricum, Maxim. in Bull. Acad. Sc. Pétersb. xii. (1868) 228.—Manchur.
marsicum, Guss. Pl. Rar. 375 = campestre.
Martini, Jord. Pugill. 52 = opulifolium.
marucum, Walp. Rep. i. 408 = campestre.
Maximowiczianum, *Miq. in Arch. Néerl.* ii. (1867) 478.—Japon.
melliodorum, Opiz, Seznam, 9 = A. Pseudo-Platanus.
mexicanum, *A. Gray, in Proc. Am. Acad.* v. (1861) 176.—Mexico.
micranthum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1845) 155.—Japon.
microcarpon, Masn. ex Opiz, Natural. vi. (1824) 48; in Flora, vii. (1824) i. 83 = campestre.
microphyllum, Opiz, in Flora, vii. (1824) i. Beil. 82; et Natural. vi. (1824) 48 = campestre.
molle, Opiz, l. c. 83 = campestre.
Mono, Maxim. in Bull. Phys. Math. Acad. Sc. Pétersb. xv. (1856) 126 = pictum.
monspessulanum, *Linn. Sp. Pl.* 1056.—Europ. austr.
montanum, Ait. Hort. Kew. ed. II. iii. 435 = spicatum.
montanum, Lam. Fl. Fr. ii. 553 = A. Pseudo-Platanus.
neapolitanum, Tenore, Fl. Nap. ii. 372. t. 100 = opulifolium.
neglectum, *Hoffmgg. ex Walp. Rep.* i. 410 = Hab.?
neglectum, *Lange, in Bot. Tidss.* xiii. (1882) 30.—An hybr.?
Negundo, Linn. Sp. Pl. 1056 = Negundo aceroides.
nigrum, Michx. f Arb. Am. ii. 238. t. 16 = saccharinum.
nikoense, Maxim. in Bull. Acad. Sc. Pétersb. xii. (1868) 227 = Maximowiczianum.
nikoense, *Miq. in Arch. Néerl.* ii. (1867) 478.—Japon.
niveum, *Blume, Rumphia*, iii. 193.—Java; Ind. or.
oblongum, *Wall. ex DC. Prod.* i. 593.—Reg. Himal.
obtusatum, Waldst. & Kit. ex Willd. Sp. Pl. iv. 984 = opulifolium.
obtusifolium, Sibth. & Sm. Fl. Graec. t. 361 = creticum.
Opalus, Mill. Gard. Dict. ed. VIII. n. 8 = opulifolium.
Opizii, *Ortm. ex Opiz, Seznam*, 9.—Bohem.
opulifolium, Thuil. ex Loud. Hort. Brit. Supp. ii. 606 = A. Pseudo-Platanus.
opulifolium, *Vill. Hist. Pl. Dauph.* iv. 802 = Europ.; Oriens.
orientale, Linn. Syst. ed. X. ii. 1300 = creticum.
ornatum, *Carr. in Rev. Hortic.* (1867) 300.—Japon.
orthopteron, Masn. ex Opiz, Natural. vi. (1824) 49; et Flora, vii. (1824) i. Beil. 82 = campestre.
pallidum, Ait. ex Steud. Nom. ed. II. i. 11 = dasycarpum.
palmatifidum, Tausch, ex Steud. l. c. = platanoides.
palmatisectum, Ortm. ex Čelak. Prod. Fl. Boehm. 538 = campestre.
palmatum, Rafin. New Fl. Am. i. 48 = macrophyllum.
palmatum, *Thunb. Fl. Jap.* 162.—Japon.
palmifolium, Borckh. Fl. Katzeln. 107 = saccharinum.
parviflorum, Ehrh. Beitr. iv. 25 = spicatum.
parviflorum, *Franch. & Sav. Enum. Pl. Jap.* ii. 321. Japon.
parvifolium, *Tausch, in Flora*, xii. (1829) 545.—Europ. austr.
pectinatum, Wall. Pl. As. Rar. ii. 4 = caudatum.
pennsylvanicum, Duroi, Diss. 61 = spicatum.
pensylvanicum, *Linn. Sp. Pl.* 1055.—Am. bor.
pentapomicum, *Stewart, ex Brand. For. Fl.* 111.—Reg. Himal.
pictum, *Thunb. Fl. Jap.* 162.—Japon.
pilosum, *Maxim. in Bull. Acad. Sc. Pétersb.* xxvi. (1880) 436.—As. or.
pinnatum, *Lour. Fl. Cochinch.* ii. 649.—Cochinch.
platanifolium, *Griff. Itin. Not.* 200 (= villosum?).—Reg. Himal.
platanifolium, Stokes, Bot. Mat. Med. ii. 382 = platanoides.
platanoides, *Linn. Sp. Pl.* 1055.—Europ.; Oriens.
Platanus, Burm. f. Fl. Ind. 221 (= quid?).—Ind. or.
polycarpon, Opiz, Natural. vi. (1824) 49; in Flora, vii. (1824) i. Beil. 82 = campestre.

ACER:—
polymorphum, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1845) 158 = palmatum.
polymorphum, *Spach, Hist. Vég. Phan.* iii. 107.—Oriens.
praecox, Opiz, Seznam, 9 = campestre.
procerum, Salisb. Prod. 280 = A. Pseudo-Platanus.
Pseudoplatanus, Falk, Beitr. ii. 279 = platanoides.
Pseudo-Platanus, *Linn. Sp. Pl.* 1054.—Europ.; Oriens.
pubescens, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 246.—Turkest.
purpurascens, *Franch. & Sav. Enum. Pl. Jap.* ii. 320. —Japon.
pycnanthum, *C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 250.—Japon.
quinquelobatum, Wagn. ex Opiz, in Flora, vii. I. (1824) Beil. 83 = campestre.
quinquelobum, C. Koch, Dendrol. 540 = creticum.
quinquelobum, Gilib. Fl. Lituan. ii. 198 = A. Pseudo-Platanus.
rectangulum, Dulac, Fl. Hautes-Pyr. 242 = monspessulanum.
Reginae-Amaliae, Orphan. ex Boiss. Diagn. Ser. II. i. 109 = Amaliae.
reticulatum, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 312.—Hongkong.
Reygassei, Boiss. Diagn. Ser. I. ii. 5, 71 = hyrcanum.
robustum, Opiz, Seznam, 9 = campestre?
rotundifolium, Lam. Encyc. ii. 382 = italum.
rotundum, Dulac, Fl. Hautes-Pyr. 242 = platanoides.
rubrum, Lam. Encyc. ii. 380 = dasycarpum.
rubrum, *Linn. Sp. Pl.* 1055.—Am. bor.
rubrum, Wats. Dendrol. Brit. t. 169 = rubrum.
rufinerve, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1845) 155. —Japon.
saccharinum, Linn. Sp. Pl. 1055 = dasycarpum.
saccharinum, *Wangenh. Nordamer. Holzarten* (1787) 36. t. 11. f. 26.—Am. bor.
saccharophorum, *C. Koch & Fint. Wochenschr.* (1859) 349.—Am. bor.
Saira, Hort. ex C. Koch, Dendrol. 541 = dasycarpum.
sanguineum, *Carr. in Rev. Hortic.* (1867) 280.—Japon.
sanguineum, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 176 = rubrum.
Schwedleri, Hort. ex C. Koch, Dendrol. 530 = platanoides.
Semenovii, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) I. 550.—As. temp.
sempervirens, Linn. Mant. i. 128 = heterophyllum.
sempervirens, *Tausch, in Flora*, xii. (1829) 550.—Hab.?
septemlobum, Thunb. Fl. Jap. 162 = palmatum.
serotinum, *Kit. in Linnaea*, xxxii. (1863) 552.—Europ.
sessilifolium, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1845) 158 = palmatum.
Sieboldianum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 87. —Japon.
sikkimense, *Miq. in Arch. Néerl.* ii. (1867) 471.—Reg. Himal.
spicatum, *Lam. Encyc.* ii. 381.—Am. bor.
stachyophyllum, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 694. —Reg. Himal.
sterculiaceum, Griff. Itin. Notes, 148 = pictum.
sterculiaceum, Wall. Pl. As. Rar. ii. 3. t. 105 = villosum.
striatum, Du Roi, Harbk. Baumz. i. 8 = spicatum.
suberosum, Dum. Fl. Belg. Prod. 113 = campestre.
sylvestre, Wender. in Schr. Nat. Ges. Marb. ii. (1830) 250 = campestre.
syriacum, *Boiss. & Gaill Diagn.* Ser. II. v. 72.—Syria; Cyprus.
tataricum, *Linn. Sp. Pl.* 1054.—Oriens.
tauricolum, Boiss. & Balansa, Diagn. Ser. II. v. 72 = hyrcanum.
Tauschianum, Opiz, Seznam, 9 = campestre.
tegmentosum, *Maxim. in Bull. Phys. Math. Acad. Sc. Pétersb.* xv. (1856) 125.—Reg. Amur.
ternatum, *Moç. & Sesse, ex DC. Prod.* i. 596.—Mexico.
tomentosum, Dulac, Fl. Hautes-Pyr. 242 = opulifolium.
Thomsoni, Miq. in Arch. Néerl. ii. (1867) 470 = villosum.
tomentosum, Kit. in Linnaea, xxxii. (1863) 552 = campestre.
tomentosum, Desf. Cat. Hort. Par. ed. III. 136 = criocarpum.
Trautvetteri, *Medw. ex Trautv. in Act. Hort. Petrop.* vii. (1880) 428 in obs.—Hab.?
tricuspifolium, Stokes, Bot. Mat. Med. ii. 370 = pensylvanicum.
trifidum, *Hook. & Arn. Bot. Beech. Voy.* 174.—China.
trifidum, Thunb. Fl. Jap. 163 = Lindera triloba.
trifoliatum, *Rafin. New Fl. Am.* i. 48.—Am. bor.
trifolium, Duham, Arb. i. t. 10. f. 8 = monspessulanum.

ACER:—
trilobatum, Lam. Encyc. ii. 382 = monspessulanum.
trilobum, Gilib. Fl. Lituan. ii. 198 = campestre.
trilobum, Moench, Meth. 56 = monspessulanum.
tripartitum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 248 = glabrum.
truncatum, Bunge, Enum. Pl. Chin. Bor. 10 = pictum.
ukurunduense, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 24) = spicatum.
Van-Volxemii, *Mast. in Gard. Chron.* vii. (1877) 72. —Reg. Caucas.
velutinum, Boiss. Diagn. Ser. I. vi. 28 = insigne.
villosum, J. & C. Presl, Delic. Prag. 31 = A. Pseudo-Platanus.
villosum, *Wall. Pl. As. Rar.* ii. 4.—Reg. Himal.
virens, *Thunb. Diss. Acere*, 3. (err. typ. 5).—Oriens.
virgatum, *Rafin. New Fl. Am.* i. 48.—Am. bor.
virginianum, Mill. ex Steud. Nom. ed. I. 5 = dasycarpum.
Visianii, *Nym. Consp.* 135.—Servia.
vitifolium, Opiz, ex Tausch, in Flora, xii. (1829) II. 549 = platanoides.
Wondračekii, Opiz, Seznam, 9 = campestre.

ACERANTHUS, Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1837) 349. t. 14. *BERBERIDEAE*, Benth. & Hook. f. i. 44.
diphyllus, *Morr. & Decne. l. c.* 350.—Japon.
macrophyllus, *Blume, ex C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 253.—Japon.
sagittatus, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1845) 175 = Epimedium sagittatum.
triphyllus, *C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 253.—Japon.

ACERAS, R. Br. in Ait. Hort. Kew. ed. II. v. 191 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 621.
affinis, Boiss. Fl. Orient. v. 56.—As. Min.
alpina, Steud. Nom. ed. II. i. 12 = Herminium alpinum.
angustifolia, Lindl. in Wall. Cat. n. 7061 = longicruris.
anthropomorpha, Steud. Nom. ed. II. i. 12 = anthropophora.
anthropophora, *R. Br. in Ait. Hort. Kew.* ed. II. v. 191. —Europ.
caprina, Lindl. Gen. et Sp. Orch. 282 = Orchis hircina.
densiflora, Boiss. Voy. Espagne, ii. 595 = Habenaria intacta.
Duquesnei, Reichb. f. Ic. Fl. Germ. xiii–xiv. 171 = Orchis pyramidalis.
formosa, Lindl. Gen. et Sp. Orch. 282 = Orchis formosa.
hircina, Lindl. l. c. = Orchis hircina.
intacta, Reichb. f. Ic. Fl. Germ. xiii–xiv. 2 = Habenaria intacta.
lancea, Steud. Nom. ed. II. i. 12 (= Ophrys lancea).
longibracteata, Reichb. f. Ic. Fl. Germ. xiii et xiv. 3 = Orchis longibracteata.
longicruris, *C. Wright, ex A. Gray, in Mem. Am. Acad. N. S.* vi. (1858–9) 411.—Japon.
maculata, Gren. in Gren. & Godr. Fl. Fr. iii. 282 in nota = Habenaria intacta.
pyramidalis, Reichb. f. Ic. Fl. Germ. xiii–xiv. 6 = Orchis pyramidalis.
secundiflora, Lindl. in Bot. Reg. t. 1525 = Habenaria intacta.

ACERATES, Ell. Sketch, i. 316 (1817) = **Gomphocarpus**, R. Br. (Asclepiad.).
angustifolia, Decne. in DC. Prod. viii. 522 = Asclepias stenophylla.
asperula, Decne. l. c.—Mexico.
atropurpurea, Kellogg, in Proc. Calif. Acad. (1854–59) 65; repr. 69 = G. cordifolius.
auriculata, Engelm. ex Torr. in Bot. Mex. Bound. 160. —Am. bor.
bifida, Rusby, ex A. Gray, in Proc. Am. Acad. xx. (1885) 296.—Arizona.
circinalis, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 322.—Mexico.
connivens, Decne. in DC. Prod. viii. 521 (= Anantherix connivens).
cordifolia, Benth. Pl. Hartw. 323 = G. cordifolius.
decumbens, Decne. in DC. Prod. viii. 522 = Asclepiodora decumbens.
Feayi, Chapm. in Coulter, Bot. Gazette, iii. (1878) 12. —Florida.
gomphocarpoides, Benth. Pl. Hartw. 66 = Nephradenia neriifolia.
gomphocarpoides, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 323.—Mexico.
humilis, Benth. Pl. Hartw. 291.—Mexico.

ACERATES:—
incarnata, Decne. in DC. Prod. viii. 523.—Am. bor.
lanceolata, Steud. Nom. ed. II. i. 12 = G. viridiflorus.
lanuginosa, Decne. in DC. Prod. viii. 523.—Am. bor.
latifolia, Torr. & Frém. in Frém. Rep. Exped. (1845) 317 = Asclepias cryptoceras.
longifolia, Ell. Sketch, i. 317 = Asclepias longifolius.
paniculata, Decne. in DC. Prod. viii. 521 = Asclepiodora viridis.
Schaffneri, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 386.—Mexico.
tomentosa, Torr. in Bot. Mex. Bound. 160 = G. tomentosus.
vinosa, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 387.—Mexico?
viridiflora, Ell. Sketch, 317 = G. viridiflorus.

ACERATIUM, DC. Prod. i. 519 (1824) = **Elaeocarpus**, Linn. (Tiliac.).
Calomala, Blanco, Fl. Filip. ed. II. 306 = E. nitidus.
oppositifolium, DC. Prod. i. 519 = E. oppositifolius.

ACEROTIS, Rafin. New Fl. Am. i. 49 (1836) = **Asclepias**, Tourn.

ACETOSA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Rumex**, Linn. (Polygon.).
Acetosella, Mill. Gard. Dict. ed. VIII. n. 2 = R. Acetosella.
aculeata, Chaz. in Mill. Dict. Suppl. i. 8 = R. bucephalophorus.
agrestis, Rafin. New Fl. Am. iv. 52 = R. Acetosa.
alpina, Mill. Gard. Dict. ed. VIII. n. 5 = R. montanus.
alpina, Moench, Meth. 357 = R. alpinus.
amplexicaulis, Rafin. New Fl. Am. iv. 53 = R. Acetosa.
angustata, Rafin. l. c. = R. Acetosa.
arifolia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 64 = R. arifolius.
bidentula, Rafin. New Fl. Am. iv. 52 = R. Acetosa.
bipinnata, Chaz. in Mill. Dict. Suppl. i. 8 = R. pictus.
bucephalophora, Fourr. in Ann. Soc. Linn. Lyon. N. S. xvii. (1869) 145 = R. bucephalophorus.
digyna, Mill. Gard. Dict. ed. VIII. n. 4 = Oxyria digyna.
hastata, Moench, Meth. 357 = R. Acetosella.
hastifolia, Schur, Enum. Pl. Transs. 581 = R. Acetosa.
hastulata, Rafin. New Fl. Am. iv. 52 = R. Acetosa.
heterophyla, Rafin. l. c. = R. maximus?
integrifolia, Rafin. l. c. = R. integrifolia.
intermedia, Fourr. in Ann. Soc. Linn. Lyon. N. S. xvii. (1869) 145 = R. intermedius.
Lunaria, Mill. Gard. Dict. ed. VIII. n. 6 = R. Lunaria.
luxurians, Chaz. in Mill. Dict. Suppl. i. 8 = R. luxurians.
magna, Gilib. Exercit. ii. 445 = R. Acetosa.
multifida, Chaz. in Mill. Dict. Suppl. i. 8 = R. multifidus.
officinalis, Gueldenst. ex Ledeb. Fl. Ross. iii. 510 = R. Acetosa.
olitoria, Rafin. New Fl. Am. iv. 53 = R. Acetosa.
parva, Gilib. Exercit. ii. 446 = R. Acetosella.
pratensis, Mill. Gard. Dict. ed. VIII. n. 1 = R. Acetosa.
repens, S. F. Gray, Nat. Arr. Brit. Pl. ii. 276 = R. Acetosella.
rosea, Mill. Gard. Dict. ed. VIII. n. 7 = R. roseus.
scutata, Mill. l. c. n. 3 = R. scutatus.
sterilis, Mill. l. c. n. 8 = R. Acetosella.
subalpina, Schur, Enum. Pl. Transs. 581 = R. Acetosa.
tingitana, Fourr. in Ann. Soc. Linn. Lyon. N. S. xvii. (1869) 145 = R. tingitanus.
tuberosa, Chaz. in Mill. Dict. Suppl. i. 8 = R. tuberosus.

ACETOSELLA, Rupp. Fl. Jen. ed. Hall. 127 (1745) = **Rumex**, Linn. (Polygon.).
vulgaris, Fourr. in Ann. Soc. Linn. Lyon. N. S. xvii. (1869) 145 = R. Acetosella.

ACHAEMENES, St. Lag. in Ann. Soc. Bot. Lyon. vii. (1880) 41 = **Achimenes**, P. Br. (Gesneriac.).

ACHAETA, Fourn. (Gram. Mex. 109) ex Benth. & Hook. f. Gen. iii. 1153 (1883) = **Deyeuxia**, Clar. (Gramin.).
geniculata, Fourn. l. c. ex Hemsl. Biol. Centr. Am. Bot. iii. 553.—Mexico.
plumosa, Fourn. l. c. ex Hemsl. l. c. 553.—Mexico.

ACHAETOGERON, A. Gray, in Mem. Am. Acad. N.S. iv. (1849) Pl. Fendl. 72. *COMPOSITAE*, Benth. & Hook. f. ii. 1232.
Galeottii, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 119.—Mexico.
Seemannii, *A. Gray, ex Hemsl. l. c.* 120.—Mexico.

ACHAETOGERON :—
Wislizeni, *A. Gray, in Mem. Am. Acad. N. S.* iv. (1849) (*Pl. Fendl.*) 72.—Mexico.

ACHANIA, Sw. Prod. Veg. Ind. Occ. 102 (1788) = **Malvaviscus**, Dill. (Malvac.).
ciliata, Spreng. Syst. iii. 100 = M. ciliatus.
coccinea, Salisb. Prod. 385 = M. arboreus.
concinna, Spreng. Syst. iii. 100 = M. concinnus.
cordata, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 99 = M. Balbisii.
floridana, Rafin. New Fl. Am. i. 49 = Hibiscus floridanus?
Malvaviscus, Sw. Prod. Veg. Ind. Occ. 102 = M. arboreus.
mollis, Ait. Hort. Kew. ii. 459 = M. mollis.
pilosa, Sw. Prod. Veg. Ind. Occ. 102 = M. pilosus.
Poeppigii, Spreng. Syst. iii. 100 = Hibiscus Poeppigii.
stylosa, Schrank, in Flora, ii. (1819) 449.—Bras.
tomentosa, Sterler, ex Steud. Nom. ed. II. i. 12.—Hab.?

ACHARIA, Thunb. Prod. Pl. Cap. 14 (1794). *PASSIFLOREAE*, Benth. & Hook. f. i. 814.
tragodes, *Thunb. l. c.*—Afr. austr.

ACHARITEA, Benth. in Benth. & Hook. f. Gen. ii. 1142 (1876). *VERBENACEAE*, Benth. & Hook.
tenuis, *Benth. l. c.*—Madag.

ACHARITERIUM, Bluff & Fingerh. Fl. Germ. ii. 345 (1825) = **Filago**, Linn. (Compos.).
arvense, Bluff & Fingerh. l. c. 346 = F. arvensis.

ACHASMA, Griff. Notul. iii. 426 (1851) = **Amomum**, Linn. (Scitam.).
macrocheilos, Griff. l. c. 429.—Ind. or.
megalocheilos, Griff. l. c. 426.—Ind. or.
metriocheilos, Griff. l. c. 427.—Ind. or.

ACHATOCARPUS, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 45. *AMARANTACEAE*, Benth. & Hook. f. iii. 26.
nigricans, *Triana, l. c.* 46.—N. Granat.
praecox, *Griseb. in Goett. Abh.* xxiv. (1879) 32.—Reg. Argent.
spinulosus, *Griseb. l. c.* 32.—Reg. Argent.

ACHETARIA, Cham. & Schlecht. in Linnaea, ii. (1827) 566 = **Beyrichia**, Cham. (Scrophularin.).
Sprengelii, Cham. & Schlecht. l. c. 557 = B. villosa.

ACHILLAEA, Linn. Phil. Bot. 29 (1764) = **Achillea**, Linn. (Compos.).

ACHILLEA, Linn. Syst. ed I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 419.
AGERATOIDES, Ponted. ex Linn. Class. Pl. 527 (1747).
ALITUBUS, Dulac, Fl. Hautes-Pyr. 499 (1867).
ARTHROLEPIS, Boiss. Diagn. Ser. I. xi. 14 (1849).
CONFORATA, Caesalp. ex Fourr. in Ann. Soc. Linn. Lyon. N. S. xvii. (1869) 91.
MILLEFOLIUM, Tourn. ex Adans. Fam. ii. 128 (1763).
PTARMICA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 174 (1745).
abrotanifolia, Hort. ex Steud. Nom. ed. II. i. 12 (= Pyrethrum myriophyllum).
abrotanifolia, Linn. Sp. Pl. 897 = Tanacetum millefoliatum.
abrotanoides, *Vis. Fl. Dalmat.* ii. 81. t. 10. f. 1.—Europ.
absinthifolia, [Clairv.] Man. Herb. 251 = Clavennae.
acuminata, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Sibiria.
adulterina, Leresch, ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 167 = obscura.
aegyptiaca, S. G. Gmel. It. ii. 198 = Gerberi.
aegyptiaca, Hort. ex Steud. Nom. ed. II. i. 12 = semipectinata.
aegyptiaca, Linn. Sp. Pl. 900 = Tournefortii.
aegyptiaca, Sm. Fl. Graec. ix. 71 = taygetea.
ageratifolia, *Benth. & Hook. f. Gen.* ii. 420.—Graecia.
Ageratum, *Linn. Sp. Pl.* 897.—Europ.
albicaulis, C. A. Mey. Verz. Pfl. Cauc. 76 = tenuifolia.
albida, Willd. Enum. Hort. Berol. Suppl. 61 = tenuifolia.
aleppica, *DC. Prod.* vii. 296.—Oriens.
alpina, *Linn. Sp. Pl.* 899 = Europ.
ambigua, Boiss. Fl. Orient. iii. 255 = A. Millefolium.
ambigua, Pollini, Viag. Lag. Garda, 117 = magna.

ACHILLEA :—
ambrosiaca, *Boiss. Fl. Orient.* iii. 276.—Thessal.
amoena, C. A. Mey. Verz. Pfl. Cauc. 76 = vermicularis.
anethifolia, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. (1865) I. 404 = A. Millefolium.
anglica, G. Don, in Loud. Hort. Brit. 355 = decolorans.
angustifolia, Salisb. Prod. 204 = alpina.
anthemoides, *Willd. Sp. Pl.* iii. 2200.—Hab.?
arabica, *Kotschy. in Sitzb. Acad. Wien. Math. Nat.* lii. Abth. I (1866) 255.—Arab.
arborea, Hornem. Hort. Hafn. Suppl. 102 = Gonospermum elegans.
argentea, Salisb. Prod. 204 = Clavennae.
argentea, Lam. Encyc. i. 29 = Tanacetum argenteum.
armenorum, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 269.—Armen.
asperula, Ces. & Pass. ex Nym. Consp. 362 = Anthemis asperula.
aspleniifolia, Leresch, ex Sch. Bip. in Oestr. Bot. Wochenschr. vi. (1856) 301 = Lereschei.
asplenifolia, *Vent. Jard. Cels*, t. 95.—Am. bor.?
atrata, *Linn. Sp. Pl.* 899.—Europ.
Aucheri, *Boiss. Diagn.* Ser. I. vi. 87.—Persia.
aurea, Lam. Encyc. i. 26 = Chrysanthemum achilleaefolium.
auriculata, *Willd. Sp. Pl.* iii. 2203.—Oriens.
baikalensis, *Fisch. Hort. Gorenk.* ed. II. (1812) 41.—Sibir. baikal.
bandana, Buch.-Ham. in Wall. Cat. n. 3291 = Chrysanthellum indicum.
bannatica, Kit. ex DC. Prod. vi. 29 = crithmifolia.
Barbeyana, *Heldr. & Heimerl, in Denkschr. Acad. Wien*, xlviii. (1884) 161.—Graecia.
Bardana, Buch.-Ham. ex DC. Prod. v. 631. sphalm. = *bandana*.
Barrelierii, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Europ.
biaristata, Spreng. Syst. iv. Cur. post. 303 = Anacyclus clavatus.
bicolor, *Wender. in Flora*, xii. (1829) I. *Erg.* 25.—Europ.?
bipinnata, Linn. Sp. Pl. 900 (= Pyrethrum myriophyllum).
biserrata, *Bieb. Fl. Taur. Cauc.* ii. 334.—As. Min.
Bocconii, Koch, Ind. Sem. Hort. Erlang. (1834) = alpina.
Boissieri, *Haussk. ex Boiss. Fl. Orient.* iii. 272.—Persia.
Boissieriana, Heldr. ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 162 = Barbeyana.
borealis, Bong. Veget. Ins. Sitcha, 149 = A. Millefolium.
brachyphylla, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 273.—Mesopot.
buglossis, Frivald. in Flora, xix. (1836) 433 = lingulata.
callichroa, *Boiss. Diagn.* Ser. I. vi. 86.—Persia.
camphorata, Gilib. Fl. Lituan. i. 217 = nobilis.
camtschatica, Rupr. ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 189 = sibirica.
Cancrini, *Gruner, in Bull. Soc. Nat. Mosc.* xli. (1868) II. 417.—Tauria.
capillaris, Poir. Encyc. Suppl. i. 102 = setacea.
capillata, Lapeyr. Hist. Abr. Pl. Pyr. 534 = chamaemelifolia.
capitata, Willd. Tract. Ach. 17 = Clavennae.
cartilaginea, *Ledeb. Ind. Dorp. ex Boiss. Fl. Orient.* iii. 274.—Europ.; Oriens.
chamaecyparissus, *Sch. Bip. in Flora*, xxxviii. (1855) 14.—Europ.
chamaecyparyssus, Reichb. f. Ic. Fl. Germ. xvi. 64. t. 1012 = praec.
chamaemelifolia, *Pourr. in Mém. Acad. Toul.* iii. (1788) 305.—Mont. Pyren.
chrysantha, Griseb. ex Boiss. Fl. Orient. iii. 265 = chrysocoma.
chrysocoma, *Frivald. in Flora*, xviii. (1835) 336.—Graecia.
cilicica, *J. Gay, ex Tchihat. Asie Min. Bot.* ii. 270.—As. Min.
claudiopolina, Janka, in Oestr. Bot. Wochenschr. (1854) 403 = impatiens.
claudiopolitana, Wolff, ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 185 = impatiens.
Clavennae, *Linn. Sp. Pl.* 898.—Europ.
Clusiana, *Tausch, in Flora*, iv. (1821) II. 551.—Europ.
Clusii, Sauter, in Flora, vii. (1824) I. 211 = Clusiana.
clypeolata, *Sibth. & Sm. Fl. Graec.* t. 893.—Oriens.
coarctata, Poir. Encyc. Suppl. i. 94 = compacta.

ACHILLEA :—
collina, Becker, ex Reichb. Fl. Germ. Excurs. 850 = A. Millefolium.
collina, Schur, ex Nym. Consp. 367 = setacea.
commutata, *Heimerl, in Flora*, lxvi. (1883) 385.—Europ.
compacta, Lam. Encyc. i. 27 = magna.
compacta, *Willd. Sp. Pl.* iii. 2206.—Oriens.
condensata, *Miq. Fl. Ind. Bat.* ii. 85.—Java.
conferta, *DC. Prod.* vi. 32.—Oriens.
coronopifolia, Willd. Tract. Ach. 18. n. 9. t. 1. f. 2 = tanacetifolia.
corymbifera, S. G. Gmel. It. i. 137. t. 25. f. 1 = nobilis.
crassifolia, Hort. ex Steud. Nom. ed. II. i. 13 = A. Millefolium.
cretica, *Linn. Sp. Pl.* 899.—Oriens.
crispa, Lam. ex DC. Prod. vi. 26 = asplenifolia.
cristata, Hort. ex DC. l. c. 25 = tanacetifolia.
cristata, Retz. Obs. Fasc. ii. 25 = falcata?
cristata, Willd. Tract. Ach. n. 4 et 8; Sp. Pl. iii. 2192 = alpina.
crithmifolia, Frivald. ex Hampe, in Flora, xx. (1837) I. 229 = micrantha.
crithmifolia, *Waldst. & Kit. Pl. Rar. Hung.* i. 68. t. 66.—Oriens.
crustata, *Schur, Enum. Pl. Transs.* 329.—Transylv.
cuneata, *Sch. Bip. in Linnaea*, xix. (1836) 313 = grandifolia.
cuneatiloba, *Boiss. & Buhse, in Nouv. Mém. Mosc.* xii. (1860) 117. t. 10. f. 2.—Persia.
cuneifolia, Lam. Encyc. i. 28 = A. Herba-rota.
cuspidata, Wall. Cat. n. 3230 = A. Millefolium.
cylindrica, Poir. Encyc. Suppl. i. 95 = nobilis.
damascena, DC. Prod. vii. 296 = falcata.
decipiens, Vest, in Flora, xii. (1829) I. Erg. 60 = nobilis.
decolorans, *Schrad. in Willd. Enum. Hort. Berol.* 913.—Europ.
decumbens, *Lam. Encyc.* i. 27.—Kamtschat.
dentato-serrata, *Heuffel, ex Heimerl, in Denkschr. Acad. Wien*, xlviii. (1884) 179.—Europ.
denticulata, Bess. ex Heimerl, l. c. 188 = sibirica.
dentifera, DC. Fl. Fr. Suppl. 485 = tanacetifolia.
dentifera, Reichb. Fl. Germ. Exc. 230 = A. Millefolium.
depressa, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. (1865) I. 405 = sibirica.
depressa, *Janka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 204.—Balkan.
distans, Waldst. & Kit. ex Willd. Sp. Pl. iii. 2207 = tanacetifolia.
diversifolia, Schur, Enum. Pl. Transs. 330 = crithmifolia.
djimilensis, Boiss. Fl. Orient. iii. 259 = latiloba.
dracunculoides, Desf. Cat. Hort. Par. ed. III. 180 = A. Ptarmica.
Dracunculus, Hort. ex Steud. Nom. ed. II. i. 13 = Artemisia Dracunculus?
dshimilensis, C. Koch, in Linnaea, xxiv. (1851) 324 = latiloba.
dubia, *Fisch. ex Steud. Nom.* ed. II. i. 13.—Sibir.
Dumasiana ×, *Vatke, in Oestr. Bot. Zeitschr.* xxii. (1872) 374.—Europ.
elegans, *Sch. Bip. in Flora*, xxxviii. (1855) 13.—Hispan.
erba-rota, 'Auct.' = A. Herba-rota.
eridania, Bertol. Fl. Ital. ix. 407 = nobilis.
erioclada, DC. Prod. vii. 296 = cretica.
eriophora, *DC. Prod.* vi. 31.—Persia.
Eupatorium, Bieb. Fl. Taur. Cauc. ii. 337 = filipendulina.
falcata, Lapeyr. Hist. Abr. Pl. Pyr, 534 = chamaemelifolia.
falcata, *Linn. Sp. Pl.* 897.—Oriens.
fasciculata, Burm. f. Fl. Cap. Prod. 27 = Eriocephalus umbellatus.
Feliciana ×, *Bruegg. in Jahresb. Naturf. Ges. Graub. N. Fol.* xxiii-xxiv. (1880) 105.—Europ.
filicifolia, Bieb. Fl. Taur. Cauc. ii. 338 = filipendulina.
fililoba, *Freyn, in Flora*, lxiv. (1881) 210.—Graecia.
filipendulina, *Lam. Encyc.* i. 27.—Oriens.
flabellifolia, Boiss. & Heldr. Diagn. Ser. I. xi. 27 = Tanacetum flabellifolium.
flabelliformis, Sibth. & Sm. Fl. Graec. ix. 72 = holosericea.
flava, Hort. ex Steud. Nom. ed. II. i. 13 = Chrysanthemum achilleaefolium.
flosculosa, *Desf. Tab. Hort. Par.* 105 (nomen)—Oriens.
formosa, *Sch. Bip. in Flora*, xxxviii. (1855) 13.—As. Min.
Fraasii, *Sch. Bip. l. c.* xxv. (1842) *Beibl.* i. 159.—Graecia.

ACHILLEA :—
fragilis, Balb. ex DC. Prod. vi. 23 = A. Ptarmica.
fragrantissima, *Sch. Bip. in Flora*, xxxviii. (1855) 13. Oriens; Aegypt.
fruticulosa, Willd. ex Ledeb. Fl. Ross. ii. 536 = Gerberi.
Genipi, Murr. App. Med. i. 168 = moschata.
Gerberi, Bieb. Fl. Taur. Cauc. i. 336 = nobilis.
Gerberi, *Willd. Sp. Pl.* iii. 2196.—Europ.; Caucas.
gilva, *Moench, Meth. Suppl.* 257.—Europ.?
Glavennae, (err. typ.) Schrank, Baier. Fl. ii. 409 = Clavennae
glomerata, Bieb. Fl. Taur. Cauc. iii. 585 = compacta.
goniocephala, *Boiss. & Bal. in Boiss. Diagn.* Ser. II. vi. 98.—As. Min.
gracilis, Rafin. ['Fl. texensis, 12'] Atl. Journ. 177 = A. Millefolium.
grandiflora, *Bieb. Fl. Taur. Cauc.* ii. 333.—As. Min.
grandifolia, *Frivald. in Flora*, xix. (1836), II. 433.—Oriens.
grandifolia, Bieb. Fl. Taur. Cauc. ii. 444 = A. Ptarmica.
grandis, Fisch. ex Herd, in Bull. Soc. Nat. Mosc. (1865) I. 407 = A. Ptarmica.
grata, Fenzl, in Tchihat. Asie Min. Bot. ii. 261 = odorata.
Gussonii, Ces. & Pass. ex Nym. Consp. 362 = Anthemis abrotanifolia.
Haenkeana, Tausch, in Flora, iv. (1821), 567 = A. Millefolium.
Halleri, Crantz, Inst. i. 303 = atrata.
Haussknechtiana, Aschers. in Festschr. Ges. Naturf. Fr. Berl. (1873) 243 = scq.
Haussknechtii, *Boiss. Fl. Orient.* iii. 264.—Persia.
helvetica, *Schleich. ex Gaud. Fl. Helv.* v. 368 = valesiaca?
Herba-rota, *All. Fl. Pedem.* i. 180.—Europ. merid.
Herba-rotta, Vill. Prosp. Delph. 32 = praec.
heterophylla, *Spreng, Nachr.* i. *Bot. Gart. Halle*, 9.—Sibir.
hispanica, Schrank, Pl. Rar. t. 70 = nobilis.
holosericea, *Sibth. & Sm. Fl. Graec* t. 894.—Graccia.
hybrida, *Gaud. in Koch, Syn. Fl. Germ.* ed. II. 409.—Europ.
imbricata, Aucher, ex DC. Prod. vii. 296 = aleppica.
imbricata, Vent. Jard. Cels, t. 54 = membranacea.
impatiens, *Linn. Sp. Pl.* 898.—Sibir.
impunctata, Vest, in Flora, iii. (1820) 3, 7 = moschata.
incisa, [*Clairv.*] *Man. Herb.* 251.—Europ.
inodora, Linn. Sp. Pl. 900 = Lonas inodora.
intermedia, Schleich. Cat. Pl. (1821) 5 = A. Millefolium.
ircutiana, Sch. Bip. in Flora, xxxviii. (1855) 15 = A. Ptarmica.
Jaborneggi ×, *Halacsy, in Oestr. Bot. Zeitschr.* xxvii. (1877) 45.—Europ.
Jacea, Griseb. Iter, ii. 264. ex Spicil. Fl. Rumel. ii. 216 = chrysocoma.
Jankae, *Sch. Bip. ex Nym. Consp.* 366.—Europ.
japonica, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 125 = sibirica.
kellalensis, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 263.—Persia.
Kotschyi, Boiss. Diagn. Ser. II. iii. 19 = odorata.
Kraettliana ×, *Bruegg. in Jahresb. Naturf. Ges. Graub. N. Fol.* xxiii–xxiv. (1880) 105, nomen.—Europ.
Laggeri, *Sch. Bip. in Oestr. Bot. Wochenschr.* vi. (1856) 300.—Europ.
lanata, Lam. Fl. Fr. iii. 640 = A. Millefolium.
lanata, *Spreng. Cat. Fl. Hal. ex Willd. Enum. Hort. Berol.* 915.—Hab.?
lanulosa. *Nutt. in Journ. Acad. Philad.* vii. (1834) 36 = A. Millefolium.
latifolia, Salisb. Prod. 204 = macrophylla.
latiloba, *Ledeb. ex Nordm. in Bull. Acad. Sc. Pétersb.* ii. (1837) 312.—Caucas.
Ledebourii, Heimerl, in Flora, lxvi. (1883) 389 = impatiens.
leneusis. Turcz. ex DC. Prod. vii. 295 = ircutiana.
leptophylla, *Bieb. Fl. Taur. Cauc.* ii. 335.—Tauria.
leptophylla, C. Koch, ex Nym. Consp. 367 = crithmifolia.
Lereschii, *Sch. Bip. in Oestr. Bot. Wocheuschr.* vi. (1856) 301.—Europ.
leucanthema, Pers. Syn. ii. 467 = A. Ptarmica.
ligustica, *All. Fl. Pedem.* i. 181. t. 52. f. 2.—Europ.; Oriens.
ligustica, Vis. ex Nym. Consp. 367 = nobilis.
linearis, Steud. Nom. ed. II. i. 13 = A. Ptarmica.
lingulata, *Waldst. & Kit. Pl. Rar. Hung.* i. 2.—Europ. orient.
livea, Scop. Delic. Insub. i. 6 = moschata.
lobata, *Forsk. Fl. Aegypt. Arab.* 218.—Aegypt.

ACHILLEA :—
lucida, Salisb. Prod. 204 = impatiens.
lycaonica, Boiss. & Heldr. Diagn. Ser. I. xi. 18 = conferta.
macrocephala, Rupr. Beitr. Pfl. Russ. Reiches, ii. (1845) 41 = A. Ptarmica.
macrophylla, Pill. & Mitterb. It. Poseg. 111. t. 11 = Chrysanthemum macrophyllum.
macrophylla, *Linn. Sp. Pl.* 898.—Europ.
macrophylloides, Panč. ex Nym. Consp. 366 = A. grandifolia.
magna, Haenke, in Jiras. Beob. Riesengeb. 103 = A. Millefolium.
magna, All. Fl. Pedem. i. 184 = tanacetifolia.
magna, *Linn. Sp. Pl.* ed. II. 1267.—Europ.
magna, Sibth. & Sm. Fl. Graec. ii. 194 = ligustica.
major, *Heimerl, in Denkschr. Acad. Wien*, xlviii. (1884) 157.—Europ.
marginata, Turcz. ex Ledeb. Fl. Ross. ii. 532 = A. Millefolium.
maxima, Fischer, ex Heimerl, in Denkschr. Acad. Wien, xlviii (1884) = A. Ptarmica.
membranacea, *DC. Prod.* vi. 32.—Oriens.
Michelii, *Sch. Bip. in Flora*, xxxviii. (1855) 14.—Europ.
micrantha, Bieb. Fl. Taur. Cauc. ii. 336 = pubescens.
micrantha, *Willd. Sp. Pl.* iii. 2209.—Soongar; Oriens.
microloba, DC. Prod. vi. 32 = teretifolia.
microphylla, Willd. Sp. Pl. iii. 2210 = odorata.
Millefolium, Ledeb. Fl. Alt. iv. 124 = setacea.
Millefolium, *Linn. Sp. Pl.* 899.—Europ.; As.; Am.
mollis, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 451, nomen.—Rossia.
mongolica, Fisch. ex Spreng. Nov. Prov. Hort. Hal. et Berol. (1818) i. 3 = sibirica.
monocephala, *Boiss. & Bal. Diagn.* Ser. I. v. 110.—Cilicia.
montana, *Schleich. ex DC. Prod.* vi. 21.—Europ.
monticola, Martr. Pl. Crit. du Tarn, 31 = A. Millefolium.
Morisiana, Reichb. Ic. Fl. Germ. xvi. 66. t. 1019 = A. Herba-rota.
moschata, *Jacq. Fl. Austriac.* v. 45.—Europ.
mucronata, Hort. ex Steud. Nom. ed. II. i. 13 = heterophylla.
mucronulata, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Europ.
multifida, Griseb. Spicil. Fl. Rumel. ii. 212 = atrata.
multifida, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Graccia.
multifida, Sprun. ex Nym. Consp. 367 = setacea.
multiflora, *Hook. Fl. Bor. Am.* i. 318.—Am. bor.
multiplex, Renar. Fl. Orn. 78 = A. Ptarmica.
muricata, Ces. & Pass. ex Nym. Consp. 362 = Anthemis muricata.
muschensis, C. Koch, in Linnaea, xxv. (1851) 329 = vermicularis.
myriophylla, Willd. Enum. Hort. Berol. 916 = Tanacetum millefoliatum.
nana, Lam. Fl. Fr. ed. I. ii. 154 = A. Herba-rota.
nana, *Linn. Sp. Pl.* 899.—Europ.
Neumayeri, Heldr. ex Heimerl, in Denkschr. Acad. Wien, xlviii (1884) 163 = Fraasii.
Neilreichii, Kern. in Oestr. Bot. Zeitschr. xxi. (1871) 148 = nobilis.
nitida ×, Pers. Syn. ii, 467 = A. impatiens.
nivea, *Sprun. ex Nym. Consp.* 365.—Europ.
nobilis, *Linn. Sp. Pl.* 899.—Europ.; Oriens.
nobilis, Roch. ex Nym. Consp. 367 = crithmifolia.
Novae-Graeciae, *Sch. Bip. in Flora* (1841), i. 158.—Enum.
obscura ×, *Nees, Ind. Sem. Hort. Bonn.* (1831) 1.—Europ.
occidentalis, Rafin. ex DC. Prod. vi. 24 = A. Millefolium.
ochroleuca, Eichw. Skizze, 149 = A. Millefolium.
ochroleuca, Ehrh. Beitr. vii. 166 = nobilis.
ochroleuca, Waldst. & Kit. Pl. Rar. Hung. i. 33 = pectinata.
odorata, *Linn. Syst.* ed. X. 1225.—Europ.; Oriens.
odorata, Pall. Reise, i. 379 = nobilis.
odorata, Wulf. in Jacq. Coll. i. 259 = setacea.
oligocephala, *DC. Prod.* vi. 32.—Mesopot.
oravicensis, *Kit. & Kanitz, Reliq. Kitaib.* 82, *in Verh. Zool. Bot. Ges. Wien*, xiii. (1863) 506.—Europ.
ossica, C. Koch, in Linnaea, xxiv. (1851) 323 = A. Millefolium.
oxyodonta, *Boiss. Diagn.* Ser. I. xi 15.—Persia.

ACHILLEA :—
oxylepis, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 271.—Persia.
oxyloba, *Sch. Bip. in Flora*, xxxviii. (1855) 15.—Afr. austr.
pallescens, DC. Prod. vii. 296 = grandifolia.
pallescens, *Linn. Syst.* ed. X. 1225.—Hab.?
pallida, Salisb. Prod. 204 = Tournefortii.
pannonica, Scheele, in Linnaea, xviii. (1835) 471 = Millefolium.
partheniflora, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. (1865) I. 407 = A. Ptarmica.
partheniifolia, *Desf. ex Steud. Nom.* ed. II. i. 14, nomen—Hab.?
pauciflora, Lam. Encyc. i. 26. = Tanacetum Vahlii.
pectinata, Lam. Encyc. i. 28 = odorata.
pectinata, Willd. Sp. Pl. iii. 2197.—Hungar; Tauria.
Peronini, *Boiss. Fl. Orient.* iii. 261.—As. Min.
peucedanifolia, Griseb. Spicil. Fl. Rumel. ii. 214 = grandifolia.
phrygia, *Boiss. & Bal. Diagn.* Ser. II. vi. 99.—As. Min.
pilosa, *Willd. Sp. Pl.* iii. 2196.—Hab.?
pinnata, *Reichb. f. Ic. Fl. Germ.* xvi. 64; *Sch. Bip. in Flora*, xxxviii. (1855) 14.—Europ.
polyphylla, Schleich. ex Koch, Syn. Fl. Germ. ed. II. 411 = setacea.
pseudo-tanacetifolia, Wier. ex Reichb. Fl. Germ. xvi. 69 = magna.
pseudo-pectinata, *Jauka, in Oestr. Bot. Zeitschr.* xxii. (1872) 179.—Turcia.
Ptarmica, *Linn. Sp. Pl.* 898.—Reg. temp. bor.
Ptarmica, Richards. in Frankl. 1st Journ. ed. II. App. 33 = multiflora.
ptarmicoides, Maxim. Prim. Fl. Amur. 154 = sibirica.
pubescens, Sibth. & Sm. Fl. Graec. t. 895 = micrantha.
pubescens, *Linn. Sp. Pl.* 897.—Oriens.
pubescens, Willd. Sp. Pl. iii. 2206 = odorata.
pumila, Schur, Enum. Pl. Transs. 330 = tanacetifolia.
punctata, Moench, Meth. 603 = alpina.
punctata, Tenore, ex Fenzl, in Tchihat. Asie Min. Bot. ii. 264 = odorata.
pyrenaica, Sibth. ex DC. Fl. Fr. iv. 211 = A. Ptarmica.
ramosissima, Moench, Meth. Suppl. 256 = alpina?
recurvifolia, Lapeyr. Hist. Abr. Pl. Pyr. 534 = chamaemelifolia.
Reichardtiana ×, *G. Beck, in Sitzb. Zool. Bot. Ges. Wien*, xxviii. (1878) 44.—Europ.
rigida, *Wallr. Beitr. Fl. Hercyn.* 331.—Europ.
rosea, Desf. Tabl. Hort. Par. ed. I. 105; ed. III. 181 = asplenifolia.
rosea, *Waldst. & Kit. Pl. Hung. Rar.*—Hungar.
rosmarinifolia, *Sch. Bip. in Flora*, xxxviii. (1855) 11.—Europ. occ.
rubra, Hildebr. ex Steud. Nom. ed. II. i. 11 = dentifera.
rupestris, *Huter, ex Beck, in Denkschr. Acad. Wien*, xlviii (1882) 150.—Europ.
salicifolia, Bess. in DC. Prod. vi. 23 = A. Ptarmica.
salina, Schur, Enum. Pl. Transs. 328 = setacea.
sambucifolia, Desf. Cat. Hort. Par. ed. III. 180 = Chrysanthemum macrophyllum.
sambucina, Thouin, ex DC. Prod. vi. 58 = Chrysanthemum macrophyllum.
Santolina, Falk, Beitr. ii. 243 = Gerberi.
Santolina, C. A. Mey. Verz. Pfl. Cauc. 76 = teretifolia.
Santolina, Hort. ex Willd. Sp. Pl. iii. 2200 = anthemoides.
Santolina, *Linn. Sp. Pl.* 896.—Oriens.; Afr. bor.
Santolina, Sibth. & Sm. Fl. Graec. ix. 70 = cretica.
Santolina, Willd. Tract. Ach. ex Steud. Nom. ed. I. 6 = tenuifolia.
santolinoides, *Lag. Gen. et Sp. Nov.* 30.—Hispan.
scabra, Host. Fl. Austr. ii. 512 = A. Millefolium.
Schkuhrii, Spreng. ex Nym. Consp. 367 = nobilis.
Schurii, Sch. Bip. in Oestr. Bot. Wochenschr. vi. (1856) 300 = oxyloba.
Seidlii, J. & C. Presl, Fl. Cech. 173 = A. Millefolium.
semipectinata, Desf. Cat. Hort. Par. ed. III (1829) 403 = auriculata.
senecioniflora, *Vest, in Flora*, xii. (1829) *Erg.* i. 59 = dentifera.
sericea, Janka, in Linnaea, xxx. (1859–60) 579 = compacta.
sericea, Vis. ex Reichb. Fl. Germ. Excurs. ii. 225 = Clavennae.
serrata ×, Retz. Obs. Fasc. ii. 25.—Sibir.
serrata, Sm. Engl. Bot. t. 2531 = decolorans.

ACHILLEA:—
serratifolia, Salisb. Prod. 204 = serrata.
serrulata, Hornem. Hort. Hafn. ii. 835 = A. Ptarmica.
setacea, *Waldst. & Kit. P. Rar. Hung.* i. 68. t. 66.—Europ.; Oriens.
setacea, Schwein. in Long, Exped. ii. 119 = A. Millefolium.
sibirica, *Ledeb. Ind. Sem. Hort. Dorpat*, (1811); *Fl. Ross.* ii. 528.—Reg. temp. bor.
sicula, Rafin. Préc. Dec. 40; et in Journ. Bot. ii. (1814) 271 = ligustica.
Siegfriedi ×, *Bruegg. in Jahresb. Naturf. Ges. Graub. N. Fol.* xxiii-xxiv. (1880) 58.—Europ.
silvatica, Auct. = sylvatica.
simplicicaulis, Boiss. Fl. Orient. iii. 259 = latiloba.
speciosa, Henckel, Adumbr. Pl. Hort. Hal. 1 = A. Ptarmica.
spinosa, *Schur, in Verh. Siebenb. Ver. Naturw.* x. (1859) 103.—Europ.
spinulifolia, *Fenzl, ex Tchihat. Asie Min. Bot.* ii. 268.—Taurus.
spinulosa. Schur, ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 185 = impatiens.
spithamea, *Coss. & Dur. ex Coss. in Ann. Sc. Nat.* Sér. IV. i. (1854) 239.—Algeria.
squarrosa, *Ait. Hort. Kew.* ed. I. iii. 242 (sp. dubia).—Hab.?
squarrosa, Hassk. Cat. Pl. Hort. Bogor. 101 = sibirica.
stricta, Schleich. ex Koch, Syn. Fl. Germ. ed. I. 374 = magna.
subcartilaginea, *Heimerl, in Denkschr. Acad. Wien*, xlviii. (1884) 188.—Europ.
subhirsuta, Gilib. Fl. Lituan. i. 217 = A. Millefolium.
sudetica, Opiz, in Hesperus (1813), 623, ex ej. Natural. n. vi. (1824) 49 = A. Millefolium.
sulphurea, *Boiss. Diagn.* Ser. I. xi. 17.—Syria.
sylvatica, Becker, Fl. Frankf. i. 295 = A. Millefolium.
sylvatica, Tenore, Fl. Nap. iv. 126 = ligustica.
sylvestris, S. F. Gray, Nat. Arr. ii. 458 = A. Ptarmica.
Szowitzii, Fisch. & Mey. ex DC. Prod. vi. 26 = latiloba.
talagonica, *Boiss. Diagn.* Ser. I. xi. 17.—Persia.
tanacetifolia, *All. Fl. Pedem.* i. 183 = Europ.; Oriens.
tanacetifolia, Mill. Gard. Dict. ed. VIII. n. 7 = A. Millefolium?
taurica, Bieb. Fl. Taur. Cauc. i. 336 = leptophylla.
taygetea, *Boiss. & Heldr. Diag.* Ser. I. xi. 16.—Oriens.
tenuifolia, *Lam. Encyc.* i. 26.—Oriens.
tenuifolia, Salisb. Prod. 204 = A. Millefolium.
tenuis, Schur, Enum. Pl. Transs. 329 = A. Millefolium.
teretifolia, *Willd. Sp. Pl.* iii. 2198.—Oriens.
Thomasiana, *Hall. f. ex Murith, Bot. Val.* 49.—Europ.
Thomasii, Thomas, ex Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 167 = pracc.
tomentosa, Fraas, ex Nym. Consp. 367 = holosericea.
tomentosa, Frivald. ex Nym. Consp. 368 = compacta.
tomentosa, *Linn. Sp. Pl.* 897.—Europ.; Oriens.; As. bor.
tomentosa, Pall. Ind. Taur. ex Ledeb. Fl. Ross. ii. 536 = leptophylla.
tomentosa, Pursh, Fl. Am. Sept. 563 = A. Millefolium.
Tournefortii, *DC. Prod.* vi. 28.—Graecia.
Trautmanni ×, *Stein, in Oestr. Bot. Zeitschr.* xxviii. (1878) 243, nomen; *cf. Heimerl, in Denkschr. Acad. Wien*, xlviii. (1882) 180.—Europ.
trichophylla, *Schrenk, Enum. Pl. Nov.* 48.—Soongaria.
trilingulata, Sch. Bip. ex Boiss. Fl. Orient. iii. 271 = aleppica.
tyrolensis, Wender. in Flora, ix. (1826) 353 = Clavennae.
umbellata, *Sibth. & Sm. Prodr.* ii. 192.—Graecia.
valesiaca ×, *Suter, Fl. Helvet.* ii. 199.—Europ.
velutina, Desf. Cat. Hort. Par. ed. III. 180 = compacta.
vermicularis, *Trin. in Mém. Acad. Petersb.* vi. (1818) 494. t. 13.—Oriens.
vermiculata, DC. Prod. vi. 32 = vermicularis.
versecensis, *Kit. ex Kanitz, Reliq. Kitaib.* 82. *in Verh. Zool.-bot. Ges. Wien*, xiii. (1863) 506.—Europ.
virgata, Hort. ex DC. Prod. vi. 31 = tenuifolia.
viridis, *Reichb. f. Ic. Fl. Germ.* xvi. t. 1012.—Europ.
viscosa, Lam. Fl. Fr. ii. 156 = A. Ageratum.
viscosa, *Sch. Bip. in Flora*, xxxviii. (1855) 14.—Europ.
Wilhelmsii, *C. Koch, in Linnaea*, xxiv. (1851) 328.—Oriens.
Wolffii, Schur, ex Nym. Consp. 365 = impatiens.

ACHILLEOPSIS, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) ii. 9 = **Rulingia**, R. Br. (Stercul.).
densiflora, Turcz. l. c. 10 = R. densiflora.

ACHILLIOS, St. Lag. in Ann. Soc. Bot. Lyon. vii. (1880) 118 = **Achillea**, Linn. (Compos.).
millefoliatus, St. Lag. l. c. = A. Millefolium.

ACHIMENES, Vahl, Symb. Bot. ii. 71. (1791) = **Artanema**, D. Don (Scroph.).
sesamioides, Vahl, l. c. = Art. sesamoides.

ACHIMENES, P. Br. Hist. Jamaic. 271. t. 30. f. 1. (1756). *GESNERACEAE*, Benth. & Hook. f. ii. 998.
Cyrilla, L'Hérit. Stirp. Nov. 147. t. 71 (1784).
Dolichodeira, Hanst. in Linnaea, xxvi. (1853) 205.
Eucodonia, Hanst. l. c. 201.
Guthnickia, Regel, in Flora, xxxii. (1849) 179.
Houttea, Heynh. Nom. ii. 306 (1846).
Koernickia, Regel, Ind. Sem. Hort. Petrop. (1857) 48; et Gartenfl. (1858) 309.
Locheria, Regel, in Flora, xxxi. (1848) 251.
Mandirola, Decne. in Rev. Hortic. (1848) 468.
Scheeria, Seem. in Bot. Mag. t. 4743 (1853).
Trevirana, Willd. Enum. Hort. Berol. 637 (1809).
amabilis, Decne. in Fl. des Serres, Sér. II. ii. (1857) 21. t. 1192 = Naegelia multiflora.
amoena, *Van Houtte, ex Hanst. in Linnaea*, xxvii. (1854) 713.—Mexico.
Andrieuxii, *DC. Prod.* vii. 536.—Mexico.
argyrostigma, Hook. Bot. Mag. t. 4175 = Koellikeria argyrostigma.
atrosanguinea, Lindl. in Journ. Hort. Soc. iii. (1848) 318 = foliosa.
autumnalis, *Hanst. in Linnaea*, xxvii. (1854) 724.—Guatemala.
candida, Lindl. in Journ. Hort. Soc. iii. (1848) 317 = Dicyrta candida.
cardinalis, *A. Dietr. in Allg. Gartenz.* xv. (1847) 314.—Mexico.
coccinea, *Pers. Syn.* ii. 165.—Ind. occ.
cochinchinensis, Spreng. Syst. ii. 812 (= Diceros cochinchinensis).
comifera, *DC. Prod.* vii. 536.—Am. austr.
cordata, Hort. ex Regel, in Flora, xxxii. (1849) 182 = grandiflora.
cornifera, Walp. Rep. ii. 719 = comifera.
cupreata, Hook. Bot. Mag. t. 4312 = Episcia cupreata.
Diceros, Gmel. ex Steud. Nom. ed. I. 7. 273 (= Diceros cochinchinensis).
divaricata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 9.—Peruv.
erinoides, *DC. Prod.* vii. 536.—Am. austr.
foliosa, *Morr. in Ann. Soc. Gand*, ii. (1846) 403. t. 91.—Mexico.
Georgeana, *Hort. ex Belg. Hortic.* x. (1860) 66. t. 5.—Hab.?
Ghiesbrechtiana, Hort. ex Regel, in Flora, xxxi. (1849) 182 = grandiflora.
Ghiesbrechtii, Lindl. in Journ. Hort. Soc. v. (1850) 193 = grandiflora.
gloxiniaeflora, Forkel, ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 318 = Gloxinia glabrata.
gracilis, Warcsz. ex Regel, in Flora, xxxi. (1849) 180 = Dicyrta candida.
grandiflora, *DC. Prod.* vii. 536.—Mexico.
Haageana, Otto, ex Regel, in Flora, xxxi. (1849) 182 = longiflora.
heterophylla, *DC. Prod.* vii. 536.—Bras.
hirsuta, *DC. l. c.*—Bras.
ichthyostoma, *Hanst. in Linnaea*, xxxiv. (1865-66) 434.—Bras.
igneseens, Lem. in Fl. des Serres, Sér I. iii. (1847) t. 248 = heterophylla.
incisa, *Klotzsch, ex Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 93.—Guatemala.
Jaureguia, Warscz. ex Haage, in Fl. des Serres, Sér. I. v. (1849) t. 536 = longiflora.
Kleei, Paxt. Mag. Bot. xvi. (1849) 289 = longiflora.
Knightii, *Lem. in Fl. des Serres*, Sér. I. iv. (1848) t. 420.—Guatemala.
lanata, *Hanst. in Linnaea*, xxxiv. (1865-66) 434.—Mexico.
Liebmanni, *Lem. in Fl. des Serres*, Sér. I. iii. (1847) t. 262; *Misc.* 47 B.—Hab.?
longiflora, *DC. Prod.* vii. 536.—Mexico.
maculata, Walp. Rep. ii. 719 = foliosa.
maculata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 8. t. 207 A.—Peruv.
magnifica, Lindl. in Gard. Chron. (1856) 116 = Locheria magnifica.

ACHIMENES:—
Martensiana, Walp. Rep. ii. 719 = foliosa.
misera, Lindl. in Journ. Hort. Soc. iii. (1848) 319 = Dicyrta Warscewicziana.
mollis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 9.—Chili.
multiflora, *Gardn. in Hook. Ic. Pl.* t. 468.—Bras.
naegelioides, *Van Houtte, Fl. des Serres*, Sér. II. viii. (1870) 117.—Hab.?
ocellata, *Hook. Bot. Mag.* t. 4359.—N. Granat.
panamensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 475.—Panama.
patens, *Benth. Pl. Hartw.* 67.—Mexico.
pauciflora, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 91.—Costa Rica.
pedunculata, *Benth. Pl. Hartw.* 78.—Mexico.
picta, Benth. in Hook. Bot. Mag. t. 4126 = Tydaea picta.
purpurea, Pritzel, Ind. Ic. App. 3, sphalm. = seq.
pyropaea, Gordon, in Journ. Hort. Soc. ii. (1847) 293. t. 4. f. 1 = coccinea.
repens, Herb. Madr. ex Benth. in DC. Prod. x. 387 = Limnophila repens.
rosea, *Lindl. Bot. Reg.* (1841) 65.—Guatemala.
rupestris, Gardn. in Hook. Ic. Pl. t. 480.—Bras.
scabra, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 8.—Peruv.
Scheerii, *Hemsl. Biol. Centr.-Am. Bot.* ii. 476.—Mexico.
Skinneri, Gordon, in Journ. Hort. Soc. ii. (1847) 293. t. 4. f. 2 = hirsuta.
superba, *Moritz, ex Hanst. in Linnaea*, xxxiv. (1865-66) 436.—N. Granat.
tenella, *DC. Prod.* vii. 536.—Mexico.
tenerrima, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 9.—Peruv.
urticaefolia, *Poepp. & Endl. l. c.* 8. t. 207. B.—Peruv.
urticaefolia, Warcz. ex Regel, in Flora, xxxii. (1849) 181 = Dicyrta Warszewicziana.
Warszewicziana, *Regel, in Flora*, xxxii. (1849) 181.—Am. austr.

ACHIMUS, Poir. in Dict. Sc. Nat. v. 51 (1827) = **Strebius**, Lour. (Urticac.).

ACHIRANTHES, P. Br. Hist. Jamaic. 180 (1756) = **Achyranthes**, Linn. (Amarantac.).

ACHIRIDA, Horan. Prod. Monog. Scitam. 18. t. 2. (1862) = **Canna**, Linn. (Scitam.).
iridiflora, Horan. l. c. = C. iridiflora.

ACHLAENA, Griseb. Cat. Pl. Cub. 228 (1866). *GRAMINEAE*, Benth. & Hook. f. iii. 1117.
piptostachya, *Griseb. l. c.* 229.—Cuba.

ACHLYS, DC. Syst. ii. 35 (1821). *BERBERIDEAE*, Benth. & Hook. f. i. 45.
japonica, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 61.—Japon.
triphylla, *DC. Syst.* ii. 35.—Am. bor.

ACHNATHERUM, Beauv. Agrost. 19. t. 6. f. 7 (1812) = **Stipa**, Linn. (Gramin.).
argenteum, Beauv. l. c. 146 = S. Calamagrostis.
bromoides, Beauv. l. c. 146. 20 = S. Aristella.
Calamagrostis, Beauv. l. c. 146 = S. Calamagrostis.
capense, Beauv. l. c. 146 = Danthonia papillosa.
conspicuum, Beauv. l. c. 146. 20 = Arundo conspicua.
Hallerii, Beauv. l. c. 146 = S. Calamagrostis.
lanceolatum, Beauv. l. c. 146. 20 = Calamagrostis lanceolata.
miliaceum, Beauv. l. c. 146. 20 (= Urachne parviflora).
soboliferum, Beauv. l. c. 146. 20 = Muehlenbergia sobolifera.
speciosum, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 317, nomen.—Europ.
tenuifolium, Beauv. Agrost. 146. 20 = Agrostis canina.

ACHNERIA, Beauv. Agrost. 72 (1812) = **Eriachne**, R. Br. (Gramin.).
brevifolia, Beauv. l. c. 73. 146 = E. mucronata.
capillaris, Beauv. l. c. 73. 146 = E. capillaris.
mucronata, Beauv. l. c. 70. 146 = E. mucronata.
obtusa, Beauv. l. c. 73. 146 = E. obtusa.

ACHNERIA, Munro, in Harv. Gen. Pl. Cap. ed. II. 449 (1868).
GRAMINEAE, Benth. & Hook. f. iii. 1158.

ACHNODON, Link, Hort. Berol. i. 65 (1827) = **Phleum**, Linn. (Gramin.).
arenarius, Link, l. c. = P. arenarium.
Bellardi, Link, l. c. = P. tenue.

ACHNODONTON, Beauv. Agrost. 24. t. 7. f. 5. (1812) = **Phleum**, Linn. (Gramin.).
arenarium, Trin. Gr. Unifl. 166 = P. arenarium.
Bellardi, Beauv. Agrost. 25 = P. tenue.
tenuis, Beauv. l. c. = P. tenue.

ACHNOPHORA, F. Muell. in Trans. Roy. Soc. S. Austral. vi. (1883) 32. *COMPOSITAE.*
Tatei, *F. Muell. l. c.* 33.—Austral.

ACHRAS, Linn. Gen. ed. I. 365 (1737). *SAPOTACEAE*, Benth. & Hook. f. ii. 657.
SAPOTA, *Gaertn. Fruct.* ii. 103. t. 104 (1791).
arnhemica, F. Muell. ex Benth. Fl. Austral. iv. 280 = Sideroxylon arnhemicum.
australis, R. Br. Prod. 530 = S. australe.
Balata, Aubl. Pl. Gui. i. 308 = Mimusops Kauki.
brasiliensis, Willd. ex Steud. Nom. ed. II. i. 15 = Chrysophyllum brasiliense.
Brownlessiana, F. Muell. Fragm. vii. 111 = Sideroxylon Brownlessianum.
Cainito, Ruiz & Pav. Fl. Per. iii. 18 = Lucuma Cainito.
chartacea, F. Muell. ex Benth. Fl. Austral. iv. 281 = Sideroxylon chartaceum.
costata, Endl. Prod. Fl. Ins. Norf. 49 = S. costatum.
cotinifolia, F. Muell. in Nuov. Giorn. Bot. iii. (1871) 31 = Hormogyne cotinifolia.
cuneifolia, Poir. Encyc. vi. 534 = Sideroxylon obovatum?
dissecta, Linn. f. Suppl. 210; Forst. f. Plant. Esc. 43 = Mimusops Kauki.
elengioides, Bedd. Fl. Sylv. t. 235 = Sideroxylon tomentosum.
euphlebia, F. Muell. Fragm. vii. 110 = S. euphlebium.
ferruginea, Casar. Stirp. Bras. Dec. vii. 63 = A. Sapota.
glaucescens, Willd. ex Steud. Nom. ed. II. i. 15 = Lucuma Rivicoa.
Guapeda, Casar. Stirp. Bras. Dec. vii. 61 = L. laurifolia.
Howeana, F. Muell. Fragm. ix. 72 = Sideroxylon Howeanum.
laurifolia, Casar. Stirp. Bras. Dec. vii. 62 = Lucuma Casaretti.
laurifolia, F. Muell. ex Benth. Fl. Austral. iv. 282 = Sideroxylon Richardi.
Lucuma, Blanco, Fl. Filip. ed. I. 237 = Lucuma mammosa.
Lucuma, Ruiz & Pav. Fl. Per. iii. 17 = L. obovata.
mammosa, Bonpl. ex. Miq. in Mart. Fl. Bras. vii. 64 = L. Bonplandii.
mammosa, Linn. Sp. Pl. ed. II. 469 = L. mammosa.
mammosa, Sieber, ex A. DC. in DC. Prod. viii. 204 = Mimusops Kauki.
mammosa, Tuss. Fl. Antill. iii. t. 6 = Bassia Jussiaei.
montana, Poir. Encyc. vi. 533 = Dipholis montana.
myrsinoides, A. Cunn. ex Benth. Fl. Austral. iv. 283 = Sideroxylon myrsinoides.
nigra, Poir. Encyc. vi. 532 = Dipholis nigra.
obovata, F. Muell. ex Benth. Fl. Austral. iv. 283 = Sideroxylon Brownii.
octodecemfida, Stokes, Bot. Mat. Med. ii. 294 = Mimusops Kauki.
pallida, Poir. Encyc. vi. 533 = Sideroxylon pallidum.
pentagona, Poir. l. c. 533 = S. pentagonum.
Pohlmaniana, F. Muell. Fragm. v. 184 = Sideroxylon Pohlmanum.
retusa, Dennst. Schluess. Hort. Malab. 30 = Aglaia edulis.
retusa, Poir. Encyc. vi. 533 = Bumelia retusa.
rotundifolia, Poir. l. c. 534 = B. rotundifolia.
salicifolia, Linn. Sp. Pl. ed. II = Dipholis salicifolia.
Sapota, *Linn. Sp. Pl.* ed. II. 470.—Am. austr.
scricea, *Schum. & Thonn. Beskr. Guin. Pl.* 176.—Afr. trop.
serrata, Poir. Encyc. Suppl. v. 36 = Prunus caroliniana.
sessiliflora, Poir. Encyc. vi. 531.—Ins. Borb.
tomentosa, Bedd. Forest. Man. Bot. 142 = Sideroxylon tomentosum.
venosa, Mart. Herb. Fl. Bras. 180 = Lucuma venosa.
verrucosa, Stokes, Bot. Mat. Med. ii. 294 = A. Sapota (?).
vitellina, Tuss. Fl. Antill. iv. 19 = Lucuma pauciflora.
xerocarpa, F. Muell. ex Benth. Fl. Austral. iv. 281 = Sideroxylon xerocarpum.
Zapota, Linn. Sp. Pl. ed. I. app. 1190 = A. Sapota.
Zapotilla, Jacq. ex Nutt. Sylv. Am. iii. 90 = A. Sapota.

ACHROANTHES, Rafin. in Med. Repos. New York, v. (1808) 352 = **Microstylis**, Lindl. (Orchid.).
acuminata, Rafin. New Fl. Am. i. 59 = M. monophylla.
cilifolia, Rafin. l. c. = M. monophylla.
laxiflora, Rafin. l. c. 58 = M. ophioglossoides.
obtusifolia, Rafin. l. c. = M. ophioglossoides.
unifolia, Rafin. in. Med. Repos. New York, v. (1808) 352 = M. ophioglossoides.

ACHROCHLOA, Griseb. Spicil. Fl. Rumel. ii. 446 (1845) = **Airochloa**, Link (Gramin.).

ACHROMOLAENA, Cass. in Dict. Sc. Nat. lvi. 222 (1828) = **Cassinia**, R. Br. (Compos.).
viscosa, Cass. l. c. = C. quinquefaria.

ACHROOSTACHYS, Benth. in Journ. Linn. Soc. xix. (1881) 134, err. typ. = **Athroostachys**, Benth. (Gramin.).

ACHRYSUM, A. Gray, in Hook. Kew Journ. iv. (1852) 228 = **Calocephalus**, R. Br. (Compos.).
glomeratum, A. Gray, l. c. 229 = C. multiflorus.

ACHUDEMIA, Blume, Mus. Bot. Lugd. Bat. ii. 57. t. 20. '1852' [1856]. *URTICACEAE*, Benth. & Hook. f. iii. 385.
japonica, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 241.—Japon.
javanica, *Blume, Mus. Bot. Lugd. Bat.* ii. 57. t. 20.—Java.

ACHYMUS, Vahl, ex Juss. in Dict. Sc. Nat. i. Suppl. 31 (1816) = **Streblus**, Lour. (Urticac.).
pallens, Soland. ex Blume, Mus. Bot. Lugd. Bat. ii. 79 = S. asper.

ACHYRACHAENA, Schau. Del. Sem. Hort. Vratisl. (1837) 3, ex Linnaea, xii. (1838) Litt. 87. *COMPOSITAE*, Benth. & Hook. f. ii. 396.
LEPIDOSTEPHANUS, Bartl. Ind. Sem. Hort. Goett. (1837), ex Linnaea, xii. (1838) Litt. 82.
mollis, *Schau. l. c.* 3.— Am. bor. occ.

ACHYRANTHES, Linn. Gen. ed. I. 34 (1737). *AMARANTACEAE*, Benth. & Hook. f. iii. 35.
AMORGINE, Rafin. New Fl. Am. iv. 44 (1836).
CADELARIA, Rafin. Fl. Tellur. iii. 39 (1836).
CENTROSTACHYS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 497 (1835).
CODIVALIA, Rafin. Fl. Tellur. iii. 40 (1836).
ECLOTORIPA, Rafin. l. c. (1836).
STEIREMIS, Rafin. l. c. (1836).
abyssinica, *Nees, in Del. Sem. Hort. Vratisl.* (1850) 3.—Abyss.
alba, *Eckl. & Zeyh. ex Moq. in DC. Prod.* xiii. II. 311.—Afr. austr.
alopecuroides, Lam. Encyc. i. 548 = Aerva javanica.
alternifolia, Linn. Mant. i. 50 = Digera arvensis.
alternifolia, Linn. f. Suppl. 159 = Cyathula globosa.
alternifolia, Russ. ex Wall. Cat. n. 6887 = Digera arvensis.
altissima, Jacq. Enum. Pl. Carib. 17 = Chamissoa altissima.
amaranthoides, Lam. Encyc. i. 548 = Digera arvensis.
angustifolia, *Benth. in Hook. Nig. Fl.* 492.—Afr. trop.
aquatica, *R. Br. Prod.* 417.—Ind. or.; Afr. trop.
arborescens, *R. Br. l. c.*—Ins. Norfolk.
argentea, Lam. Encyc. i. 545 = aspera.
aspera, Jacq. f. Ecl. 109. t. 74 = crispa.
aspera, Linn. Sp. Pl. 204.—Geront. trop.
aspera, Wall. Cat. n. 6924 H = bidentata.
atra, Pav. ex Moq. in DC. Prod. xiii. II. 372 = Telanthera elongata.
atropurpurea, Lam. Encyc. i. 546 = Pupalia atropurpurea.
australis, *R. Br. Prod.* 417.—Austral.
avicularis, *E. Mey. ex Moq. in DC. Prod.* xiii. II. 311.—Afr. austr.
axillaris, Willd. Enum. Hort. Berol. 270 = Alternanthera spinosa.
baccata, Pav. ex Moq. in DC. Prod. xiii. II. 250 = Chamissoa altissima.
bengalensis, *Lam. Encyc.* i. 459 (Sp. dub.).—Ind. or.
bidentata, *Blume, Bijdr.* 545.—As. trop.
borbonica, *Willd. ex Roem. et Schult. Syst.* v. 549.—Ins. Borbon.
brachiata, Linn. Mant. 50 = Nothosaerua brachiata.
brasiliana, Thouin, ex Moq. in DC. Prod. xiii. II. 360, 318 = Alternanthera procumbens.

ACHYRANTHES:—
calceolata, Russ. ex Wall. Cat. n. 6927 = Psilotrichum calceolatum.
Calea, *Ibantz, in La Natural.* iii. (1879) 79. t. 3.—Am. centr.
canescens, *R. Br. Prod.* i. 417.—Austral.
capitata, *Forsk. Fl. Aegypt. Arab.* 48.—Arab.
capitata, Moq. in DC. Prod. xiii. II. 310 = involucrata.
capituliflora, *Bert. in Colla, Hort. Ripul.* 4. t. 18.—N. Granat.
carnosa, Wall. Cat. n. 6931 = Psilotrichum trichotomum.
ciliata, Lam. Encyc. i. 549 = Digera arvensis.
conferta Kuiz, ex Seub. in Mart. Fl. Bras. v. I. 223 = Guilleminea densa.
cordata, Hochst. in Flora, xxvii (1844) Beil. 6 = Psilostachys gnaphalobrya,
corymbosa, Heyne, ex Wall. Cat. n. 6913 = Celosia polygonoides.
corymbosa, Linn. Sp. Pl. 205 = Polycarpaea corymbosa.
crispa, *Desf. ex Poir. Encyc. Suppl.* ii. 10.—Am. austr.
cylindrica, Boj. ex Moq. in DC. Prod. xiii. II. 328 = Cyathula cylindrica.
debilis, Poir. Encyc. Suppl. ii. 10 = Cyathula prostrata.
decumbens, Forsk. Fl. Aegypt. Arab. 47 = Celosia polygonoides.
diandra, *Roxb. Fl. Ind. ed. Carey*, ii. 504.—Zeylan.
dichotoma, *Linn. Mant.* 51.—Am. bor.
diffusa, Moench, Meth. 315 = Cyathula prostrata?
Digera, Poir. Encyc. Suppl. i. 2. 11 = Digera arvensis.
dioica, Pav. ex Moq. in DC. Prod. xiii. II. 350 = Alternanthera flavescens.
echinata, Retz. Obs. Fasc. ii. 12 = Pupalia lappacea.
ellipticifolia, Stokes, Bot. Mat. Med. i. 479 = aspera.
excelsa, Thou. ex Moq. in DC. Prod. xiii. II. 315 = aspera.
fasciculata, *Ehrenb. ex Schweinf. Beitr. Fl. Aethiop.* 179.—Afr. trop.
ferruginea, Roxb. Fl. Ind. ed. Carey, ii. 502 = Psilotrichum ferrugineum.
ficoidea, Lam. Encyc. i. 548 = Telanthera ficoidea.
filifolia, *Willd. ex Roem. et Schult. Syst.* v. 549.—Venezuela.
frumentacea, *Burm. f. Fl. Cap. Prod.* 7.—Afr. austr.
frutescens, Hort. ex Steud. Nom. ed. II. i. 15 = Digera arvensis.
fruticosa, Desf. Cat. Hort. Par. ed. III. 76 = aspera.
fruticosa, *Lam. Encyc.* i. 545.—Ind. or.
geminata, *Schum. & Thonn. Beskr. Guin. Pl.* 38.—Afr. trop.
geniculata, Pav. ex Moq. in DC. Prod. xiii. II. 381 = Telanthera capituliflora.
glabella, *A. Gray, in Proc. Am. Acad.* vii. (1868) 200.—Ins. Sandv.
globosa, Pers. Syn. i. 259 = Cyathula globosa.
glauca, Heyne, ex Wall. Cat. n. 6927 = Psilotrichum calceolatum.
globulifera, Boj. ex Moq. in DC. Prod. xiii. II. 329 = Cyathula globulifera.
grandifolia, *Moq. in DC. Prod.* xiii. II. 313.—Java.
halimifolia, Lam. Encyc. i. 547 = Telanthera frutescens.
Heudelotii, Moq. in DC. Prod. xiii. II. 310 = Pandiaka Heudelotii.
hirtiflora, *A. Rich. Fl. Cub. Fanerog.* ii. 175.—Cuba.
hispida, Moq. in DC. Prod. xiii. II. 312 = bidentata.
incana, *Fisch. ex Schrank, in Denkschr. Bot. Ges. Regensb.* ii. (1822) 36.—Bras.
incana, Pav. ex Moq. in DC. Prod. xiii. II. 365 = Telanthera frutescens.
incana, Roxb. Fl. Ind. i. 671; ed. Carey, ii. 495 = Aerva javanica.
indica, Mill. Gard. Dict. ed. VIII. n. 2 = aspera.
involucrata, Moq. in DC. Prod. xiii. II. 310 = Pandiaka involucrata.
javanica, *Moq. l. c.* 312.—Java.
javanica, Pers. Syn. i. 259 = Aerva javanica.
lanata, Linn. Sp. Pl. 204 = Aerva lanata.
lanceolata, Klein, in Wall. Cat. n. 6929 = bidentata.
lanceolata, Zoll. & Mor. ex Moq. in DC. Prod. xiii. II. 312 = javanica.
lappacea, Linn. Sp. Pl. 204 = Pupalia atropurpurea.
lappacea, Willd. Sp. Pl. i. 1192 = Pupalia lappacea.
leptostachya, E. Mey. in Hook. Lond. Journ. Bot. ii. (1843) 548 err. typ. 448 = alba.
linarifolia, *Sw. in Vet. Akad. Handl. Stockh.* (1825) 48.—Ins. Molucc.
Linkiana, Roem. & Schult. Syst. v. 545 = Chamissoa macrocarpa.

ACHYRANTHES:—
lupulina, Willd. ex Steud. Nom. ed. II. i. 15 = Telanthera lupulina.
mauritiana, *Moq. in DC. Prod.* xiii. II. 313.—Ins. Maurit.
mollis, Lepr. ex Seub. in Mart. Fl. Bras. v. I. 230 = Cyathula prostrata.
mollis, *Schum. & Thonn. Beskr. Guin. Pl.* 137.—Afr. trop.
Monsoniae, Pers. Syn. i. 258 = Aerva Monsoniae.
montana, Klein, ex Wall. Cat. n. 6926 = Psilotrichum nudum.
Moorcroftiana, Hook. f. Fl. Brit. Ind. v. 8. lapsu, pro Axyridi = Euotia ceratoides.
mucronata, Herb. ex Moq. in DC. Prod. xiii. II. 248 = Allmania albida.
mucronata, Lam. Encyc. i. 547 = Alternanthera Achyrantha.
muricata, Linn. Sp. Pl. ed. II. 295 = Digera arvensis.
mutica, *A. Gray, ex H. Mann, in Proc. Am. Acad.* vii. (1867) 200.—Ins. Hawaii.
natans, Buch.-Ham. ex Wall. Cat. n. 6932 = aquatica.
nervosa, *Roth, Nov. Pl. Sp.* 170.—Ind. or.
nivea, *Ait. Hort. Kew.* i. 286.—Ins. Canar.
nivea, Link, in Buch, Beschreib. Canar. 141 = Alternanthera Achyrantha.
nodiflora, Roxb. Fl. Ind. i. 678 = Allmania nodiflora.
nodosa, Bert. ex Moq. in DC. Prod. xiii. II. 352 = Alternanthera argentata.—Am. trop.
nodosa, *Vahl, in Schum. & Thonn. Beskr. Guin. Pl.* 139.—Afr. trop.
Novae-Brasiliae, Steud. Nom. ed. II. i. 15 = seq.
novae-brasilianae, Thouin, ex Roem. & Schult. Syst. v. 564 = Alternanthera procumbens.
nuda. Heyne, ex Wall. in Roxb. Fl. Ind. ed. Carey, ii. 507 = Psilotrichum nudum.
obovata, Pav. ex Moq. in DC. Prod. xiii. II. 359 = Alternanthera Achyrantha.
obovatifolia, Stokes, Bot. Mat. Med. i. 480 = aspera.
obtusifolia, Lam. Encyc. i. 545 = aspera.
orbiculata, Heyne, in Roxb. Fl. Ind. ed. Carey, ii. 507 = Pupalia lappacea.
ovata, *Ehrenb. ex Schweinf. Beitr. Fl. Aethiop.* 135.—Afr. trop.
paludosa, Bunb. in Proc. Linn. Soc. i. (1841) 109 = Telanthera philoxeroides.
paniculata, Forsk. Fl. Aegypt. Arab. 48 = Celosia caudata.
papposa, Forsk. l. c. 48 = Eurotia ceratoides.
patula, Linn. f. Suppl. 160 = Pupalia lappacea.
persica, Steud. Nom. ed. I. 8 = Aerva javanica.
picta, *Passer. in Bot. Zeit.* (1864) 31.—Bras.
pilosa, Pav. ex Moq. in DC. Prod. xiii. II. 352 = Alternanthera argentata.
piloselloides, Poit. ex Moq. l. c. 337 = Gossypianthus lanuginosus.
polygonata, Heyne, ex Wall. Cat. n. 6914 = Banalia thyrsiflora.
polygonoides, Heyne, ex Wall. l. c. n. 6926 = Psilotrichum nudum.
polygonoides, Lam. Encyc. i. 547 = Telanthera polygonoides.
polygonoides, Retz. Obs. Fasc. ii. 12 = Digera arvensis.
polygonoides, R. Br. Prod. 416 = Telanthera polygonoides.
polystachya, *Forsk. Fl. Aegypt. Arab.* 48.—Arab.
porphyristachya, Wall. Cat. n. 6925 = aspera.
porrigens, Jacq. Hort. Schoenb. iii. 54 = Telanthera porrigens.
prostrata, Linn. Sp. Pl. ed. II. 296 = Cyathula prostrata.
prostrata, Willd. Sp. Pl. i. 1194 = Cyathula globosa.
pubescens, Roth, Nov. Pl. Sp. 171 = Aerva Monsonia.
pungens, Lam. Encyc. i. 546 = Aerva Monsonia.
purpurea, Pav. ex Moq. in DC. Prod. xiii. II. 377 = Telanthera porrigens.
radicans, Cav. in Anal. Hist. Nat. Madr. iii. (1801) 27 = Alternanthera Achyrantha.
radicans, Heyne, in Roth, Nov. Pl. Sp. 170 = Aerva radicans.
repens, Ell. Sketch. i. 309 = Telanthera polygonoides.
repens, Heyne, in Roth, Nov. Pl. Sp. 167 = Cyathula prostrata.
repens, Linn. Sp. Pl. 205 = Alternanthera Achyrantha.
rivinaefolia, *Desf. Cat. Hort. Par.* ed. III. (1829) 390, 76.—Hab.?
rosea, Spreng. Syst. iv. Cur. Post. 103 = Polycarpaea violacea.

ACHYRANTHES:—
rotundifolia, Klein, ex Wall. Cat. n. 6936 = Pupalia orbiculata.
rubrofusca, Wight, Ic. v. t. 1778 = aspera.
sanguinolenta, *Linn. Sp. Pl.* ed. II. 294.—Ind. or.
sarmentosa, Link, ex Seub. in Mart. Fl. Bras. v. I. 241 = Chamissoa macrocarpa.
sarmentosa, *Vahl, Symb. Bot.* ii. 44.—Ind. or.
scandens, Hook. f. & Thoms. ex C. B. Clarke, in Journ. Linn. Soc. xv. (1877) 139 = Stilbanthus scandens.
scandens, Heyne, ex Moq. in DC. Prod. xiii. II. 348 = Iresine amarantoides.
scandens, Roxb. Fl. Ind. i. 676.; ed. Wall. ii. 503 = Aerva scandens.
scandens, Thonn. ex Moq. in DC. Prod. xiii. II. 331 = Pupalia atropurpurea.
sequax, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 506 = Cyathula tomentosa.
sericea, *Heyne, ex Wall. Cat.* n. 6390.—Ind. or.
sericea, Koen. in Roxb. Fl. Ind. iii. 675; ed. Carey, ii. 502 = Psilostachys sericea.
serpyllifolia, Poir. Encyc. Suppl. ii. 11 = Telanthera serpyllifolia.
sessilis, Steud. Nom. ed. II. i. 16 = Alternanthera sessilis.
setacea, Roth, Nov. Pl. Sp. 168 = Aerva Monsonia.
sicula, Roth, Catalect. Bot. i. 39 = aspera.
spinosa, *Desv. Hort. Divion.* (1834–36).—Hab.?
spinosa, Hornem. Hort. Hafn. i. 240 = Alternanthera spinosa.
splendens, *Mart. ex Moq. in DC. Prod.* xiii. II. 316.—Ins. Sandv.
stellata, Willd. Sp. Pl. i. 1195 = Polycarpaea stellata.
styracifolia, Lam. Encyc. i. 546 = Pupalia lappacea.
tenella, Poit. ex Moq. in DC. Prod. xiii. II. 341 = Iresine serpyllifolia.
tenuifolia, *Steud. Nom.* ed. II. i. 16 = aspera?
tenuifolia, Willd. Sp. Pl. i. 1196 = Polycarpaea tenuifolia.
ternata, Host, ex Fisch. Mey. & Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 12 = Gomphrena decumbens.
Thonningii, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 137.—Afr. trop.
thyrsiflora, Heyne, ex Wall. Cat. n. 6914 = Banalia thyrsiflora.
tomentella, *Zipp. ex Span. in Linnaea*, xv. (1841) 345.—Malaya.
tomentosa, Roth, Nov. Pl. Sp. 167 = Cyathula tomentosa.
triandra, Roxb. Fl. Ind. i. 678 = Alternanthera sessilis.
trichotoma, Perr. ex Moq. in DC. Prod. xiii. II. 322 = aquatica.
uncinata, Willd. ex Roem. & Schult. Syst. v. 554 = Cyathula achyranthoides.
uncinulata, *Schrad. Ind. Sem. Hort. Goett.* (1833) 1.—Afr. austr.
velutina, *Hook. & Arn. Bot. Beech. Voy.* 68.—Ins. Societ.
Verreauxii, Moq. in DC. Prod. xiii. II. 311 = alba.
Verschaffeltii, Lem. Illustr. Hortic. (1846) t. 409 = Iresine Herbstii.
verticillata, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1803) 26.—Afr. austr.
villosa, Blanco, Fl. Filip. ed. I. 189 = Alternanthera nodiflora.
villosa, Forsk. Fl. Aegypt. Arab. 48 = Aerva lanata.
violacea, Spreng. Syst. iv. Cur. Post. 102 = Polycarpaea violacea.
virgata, [*Desf.*] *Poir. Encyc. Suppl.* ii. 10.—Portorico.
virgata, Hort. ex Steud. Nom. ed. II. i. 16 = Celosia virgata.
virgata, Klein, ex Wall. Cat. n. 6915 = Banalia thyrsiflora.
Wightiana, Wall. Cat. n. 6928 = bidentata.

ACHYRASTRUM, Neck. Elem. i. 57 (1790) = **Hyoseris**, Linn. (Compos.).

ACHYROBACCHARIS, Sch. Bip. in Walp. Rep. ii. 952 (1843) = **Baccharis**, Linn. (Compos.).

ACHYROCLINE, Less. Syn. Comp. 332 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 305.
adoensis, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 428.—Abyss.
alata, *DC. Prod.* vi. 221.—Ecuad.
albicans, Griseb. in Goett. Abh. xxiv. (1879) 187 = satureioides.
argentina, *O. Hoffm. in Linnaea*, xliii. (1880–82) 135.—Reg. Argent.

ACHYROCLINE:—
batocana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 339.—Afr. trop.
bogotensis, *DC. Prod.* vi. 221.—N. Granat.
candicans, DC. l. c. = satureioides.
capitata, *Baker, in Mart. Fl. Bras.* vi. III. 117.—Bras.
celosioides, *DC. Prod.* vi. 221.—Peruv.
citrina, Griseb. in Goett. Abh. xxiv. (1879) 187 = satureioides.
coquimbense, *Klatt, in Linnaea*, xlii. (1878–79) 112.—Chile.
disjuncta, *Hemsl. Bot. Voy. Challenger*, i. II. 128.—Ins. Trinit.
flaccida, *DC. Prod.* vi. 220.—Bras.
flavescens, Griseb. in Goett. Abh. xix. (1874) 181 = alata.
glumacea, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 340.—Afr. trop.
hirta, *Klatt, in Linnaea*, xlii. (1878–79) 113.—N. Granat.
Hochstetteri, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 428.—Abyss.
latifolia, *Wedd. Chlor. And.* i. 148.—Boliv.
leptocephala, *DC. Prod.* vi. 220.—Madag.
luzuloides, *Vatke, in Linnaea*, xxxix. (1875) 489; *et in Oestr. Bot. Zeitschr.* (1875) 326.—Afr. trop.
madioides, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 275.—Peruv.
mathiolaefolia, DC. Prod. vi. 221 = satureioides.
mollis, Benth. Pl. Hartw. 207 = satureioides.
Moritziana, *Klatt, in Linnaea*, xlii. (1878–79) 112.—N. Granat.
pterocaula, DC. Prod. vi. 221 = alata.
rufescens, *DC. l. c.* 220.—N. Granat.
rugosa, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 419 = Stenocline chionaea.
satureioides, *DC. Prod.* vi. 220.—Am. merid.
satureioides, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 424 = Stenocline Gardneri.
Schimperi, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 428.—Abyss.
sclerochlaena, *Sch. Bip. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 341.—Afr. trop.
Steetzii, Vatke, in Oestr. Bot. Zeitschr. xxvii. (1877) 194 = batocana.
tamariscina, Schrank, in Denkschr. Akad. Muench. viii. (1821–22) 170 = Stoebe fusca.
Trianae, *Klatt, in Linnaea*, xlii. (1878–79) 113.—N. Granat.
Vargasiana, DC. Prod. vi. 220 = satureioides.
Vautheriana, DC. l. c. = alata.
ventosa, *Klatt, in Linnaea*, xlii. (1878–79) 112.—Mexico.

ACHYROCOMA, Cass. in Dict. Sc. Nat. v. 57 (1828) = **Vernonia**, Schreb. (Compos.).
tomentosa, Cass. l. c. xxvi. 21 = V. Achyrocoma.

ACHYROCOME, Schrank, in Denkschr. Akad. Muench. viii. (1821–22) 147 & 170 = **Elytropappus**, Sch. Bip. (Compos.).
ambigua, Schrank, l. c. = E. ambiguus.

ACHYRONIA, Royen, ex Linn. Gen. ed. II. 346 (1742) = **Aspalathus**, Linn. (Legumin.).

ACHYRONIA, Wendl. Obs. Bot. 39. t. 12. (1798) = **Priestleya**, DC. (Legum.).
arborea, Spreng. Syst. iii. 285, iv. Cur Post. 271, 287 = Carmichaelia australis.
villosa, Wendl. Obs. Bot. 39 = P. Thunbergii?

ACHYRONYCHIA, Torr. & Gray, in Proc. Am. Acad. vii. (1868) 330. *ILLECEBRACEAE*, Benth. & Hook. f. iii. 15.
Cooperi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 331.—N. Mexico; Calif.
Palmeri, Hook. f. in Benth. & Hook. f. Gen. iii. 15 = Parryi.
Parryi, *Hemsl. Diag. Pl. Nov.* 36.—Mexico.

ACHYROPAPPUS, Bieb. ex Fisch. in DC. Prod. vi. 563 (1838) = **Tricholepis**, DC. (Compos.).

ACHYROPAPPUS, H. B. & K. Nov. Gen. et Sp. iv. 257. t. 390 (1820) = **Schkuhria**, Roth (Compos.).
anthemoides, H. B. & K. l. c. = Bahia anthemoides.
maritimus, Poepp. ex DC. Prod. v. 657 = Bahia ambrosioides.

ACHYROPAPPUS :—
pedatus, Less. Syn. Comp. 239 = Florestina pedata.
schkuhrioides, *D. Don, ex Hook. & Arn. in Hook. Lond. Journ. Bot.* iii. (1840) 322.—Hab.?
schkuhrioides, Link & Otto, Abb. Neuer Gew. Gart. Berl. t. 30 = Schkuhria senecioides.
Woodhousei, A. Gray, in Proc. Am. Acad. vi. (1864–5) 546 = Bahia Woodhousei.

ACHYROPHORUS, Adans. Fam. ii. 112 (1763) = **Hypochoeris**, Linn. (Compos.).
acaulis, Remy, in C. Gay, Fl. Chil. iii. 448.—Chili.
albiflorus, *Sch. Bip. in Pollichia*, xvi–xvii. (1859) 52.—N. Granat.
alpicolus, Schur, Enum. Pl. Transs. 365 = H. maculata.
ambiguus, DC. Prod. vii. 93 = H. cretensis.
andinus, DC. l. c. 92 = H. andina.
angustissimus, Phil. in Linnaea, xxxiii. (1864–65) 127.—Chili.
apargioides, DC. Prod. vii. 94 = H. apargioides.
araucanus, Phil. in Linnaea, xxviii. (1856) 721.—Chili.
arenarius, DC. Prod. vii. 95 = H. arenaria.
aurantiacus, DC. l. c. 93 = H. grandiflora.
Balbisii, Hornem. Hort. Hafn. Suppl. 91 = H. Balbisii.
barbatus, Sch. Bip. in Pollichia, xvi–xvii. (1859) 48.—N. Granat.
Bipontinae, Sch. Bip. l. c. 54 = H. Gardneri.
Boissierianus, Walp. Ann. ii. 956.—Creta.
brasiliensis, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 106 = H. brasiliensis.
brasiliensis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 128 = H. Gardneri.
brevicaulis, Phil. in Linnaea, xxxiii. (1864–65) 127.—Chili.
chilensis, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 104.—Chili.
chondrilloides, A. Gray, in Proc. Am. Acad. v. (1845) 145 = H. brasiliensis.
chrysanthus, DC. Prod. vii. 94 = H. chrysantha.
ciliatus, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 128 = H. grandiflorus.
clarionoides, Remy, in C. Gay, Fl. Chil. iii. 450.—Chili.
coronopifolius, Sch. Bip. in Flora, xxxviii. (1855) 123.—Reg. Magell.
cryptocephalus, Sch. Bip. in Lechl. Berb. Am. Austr. 57.—Peruv.
elatus, Wedd. Chlor. And. i. 223 = H. elata.
discolor, Godr. in Mém. Acad. Stanislas, Sér. III. [xvii.] 1853 (1854), 407.—Hab.?
eriolaenus, Sch. Bip. in Lechl. Berb. Am. Austr. 57.—Peruv.
foliosus, Phil. Fl. Atac. 29.—Chili.
Gardneri, Sch. Bip. in Pollichia, xvi–xvii. (1859), 64 = H. Gardneri.
glabrescens, Steud. Nom. ed. II. i. 16 = H. arenaria.
glaucus, Phil. Fl. Atac. 29 = H. andina.
graminifolius, Sch. Bip. ex Walp. Rep. vi. 333 = tenuifolius.
grandidentatus, Phil. Fl. Atac. 29.—Chili.
grandiflorus, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 113.—Sibiria.
grandiflorus, Ledeb. Fl. Ross. ii. 777 = H. grandiflora.
helveticus, Hornem. Hort. Hafn. ii. 772 = H. uniflora.
helveticus, Less. Syn. Comp. 130 = H. helvetica.
hispidus, Hornem. Hort. Hafn. ii. 772 = H. hispida.
Hohenackeri, Sch. Bip. in Lechl. Berb. Am. Austr. 57.—Peruv.
Humboldtii, Sch. Bip. in Pollichia, xvi–xvii. (1859) 52.—N. Granat.
humilis, Phil. in Anales Univ. Chil. (1873) 482.—Chili.
incanus, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 100.—Am. austr.
Jussieui, Sch. Bip. l. c. 96.—Chili.
Lechleri, Sch. Bip. in Lechl. Berb. Am. Austr. 54.—Chili.
Lessingii, Sch. Bip. l. c. 98.—Chili.
maculatus, Scop. Fl. Carn. ed II. ii. 116 = H. maculata.
Meyenianus, Walp. Rep. vi. 336 = H. Meyeniana.
microcephalus, Sch. Bip. in Pollichia, xvi–xvii. (1859), 59 = H. brasiliensis.
microphyllus, Remy, in C. Gay, Fl. Chili. iii. 449.—Chili.
minimus, Hornem. Hort. Hafn. ii. 773 = H. minima.
montanus, Phil. in Linnaea, xxviii. (1856) 720.—Chili.
odoratus, Walp. Rep. vi. 336.—Chili.
oxydontus, DC. Prod. vii. 92 = H. cretensis.
palustris, Phil. in Anales Univ. Chil. (1865) ii. 317—Chili.

ACHYROPHORUS :—
petiolaris, DC. Prod. vii. 94 = H. petiolaris.
pinnatifidus, DC. l. c. 93 = H. pinnatifida.
praemorsus, Webb, ex Walp. Rep. vi. 329 = Webbii.
psychrophilus, Wedd. Chlor. And. i. 222.—Chili.
quitensis, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 120.—Ecuad.
radicatus, Scop. Fl. Carn. ed. II. ii. 117 = H. radicata.
ramosus, DC. Prod. vii. 92.—Hab.?
Robertia, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 127.—Italia.
roseus, Less. in Linnaea, v. (1830) 133. t. 2. f. 5 = Pinaropappus roseus.
sagittatus, Phil. Scrt. Mend. alt. 20.—Chili.
Scorzonerae, DC. Prod. vii. 94 = H. Scorzonerae.
Selloi, Sch. Bip. in Pollichia, xvi–xvii. (1859), 60 = H. brasiliensis.
serioloides, DC. Prod. vii. 93 = H. cretensis.
sessiliflorus, DC. l. c. 95 = H. sessiliflora.
setosus, Wedd. Chlor. And. i. 220.—Peruv.
sonchoides, DC. Prod. vii. 95 = H. sonchoides.
spathulatus, Remy, in C. Gay, Fl. Chil. iii. 447.—Chili.
spinuliger, Phil. in Linnaea, xxviii. (1856) 721.—Chili.
stenocephala, A. Gray, ex Wedd. Chlor. And. i. 221.—Peruv.
taraxacifolius, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 101 = H. radicata.
taraxacoides, Walp. Rep. vi. 336.—Peruv.
tenerifolius, Remy, in C. Gay, Fl. Chil. iii. 446.—Chili.
tenuifolius, DC. Prod. vii. 94 = H. tenuifolia.
tenuisectus, Sch. Bip. in Pollichia, xvi–xvii. (1859), 57 = H. variegata.
thrincioides, Remy, in C. Gay, Fl. Chil. iii. 445.—Chili.
trichocephalus, Sch. Bip. in Pollichia, xvi–xvii. (1859) 57 = H. variegata.
uniflorus, F. W. Schmidt, Samml. Phy. Aufz. i. (1795) 284 = H. uniflora.
variegatus, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 119 = H. variegata.
Volckmanni, Phil. in Anales Univ. Chil. (1873) 482.—Chili.
Webbii, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 97.— Ins. Falk.

ACHYROPSIS, Benth. & Hook. f. Gen. iii. 36 (1880). *AMARANTACEAE*, Benth. & Hook. f. l. c.
acicularis, *Benth. & Hook. f. l. c.*—Afr. austr.
alba, *Benth. & Hook. f. l. c.*—Afr. austr.
leptostachya, *Benth. & Hook. f. l. c.*—Afr. austr.

ACHYROSERIS, Sch. Bip. in Nov. Act. Nat. Cur. xxi. 1. (1845) 165 = **Scorzonera**, Linn. (Compos.).
macrospermum, Sch. Bip. l. c. = S. macrosperma.

ACHYROSPERMUM, Blume, Bijdr. 840 (1826). *LABIATAE*, Benth. & Hook. f. ii. 1208.
LAMPROSTACHYS, *Boj. ex Benth. in DC. Prod.* xii. 457 (1848).
SIPHOTORIS, *Boj. ex Benth. l. c.* (1848).
aethiopicum, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 56.—Afr. trop.
densiflorum, *Blume, Bijdr.* 841.—Ins. Malay.
fruticosum, *Benth. Lab. Gen. et Sp.* 643 & 745.—Madag.
philippinense, Benth. in DC. Prod. xii. 458 = densiflorum.
phlomoides, *Blume, Bijdr.* 841.—Java.
seychellarum, *Baker, Fl. Maurit.* 259.—Ins. Seychelles.
Wallichianum, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 673.—Reg. Himal.

ACHYROSTEPHUS, Kunze, ex Reichb. Consp. 212[a] (1828). (Compos.).

ACIA, Schreb. Gen. ii. 458 (1791) = **Couepia**, Aubl. (Rosac.).
amara, Willd. Sp. Pl. iii. 717 = C. guianensis.
dulcis, Willd. l. c. = C. guianensis.

ACIACHNE, Benth. in Hook. Ic. Pl. t. 1362 (1881). *GRAMINEAE*, Benth. & Hook. f. iii. 1143.
pulvinata, *Benth. l. c.*—N. Granat.

ACIANTHERA, Scheidw. in Otto & Dietr. Allgem. Gartenz. x. (1842) 292. = **Pleurothallis**, R. Br. (Orchid.).
punctata, Scheidw. l. c. = P. Acianthera.

ACIANTHUS, R. Br. Prod. 321 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 611.
atepalus, *Reichb. f. in Linnaea*, xli. (1877) 56.—N. Caled.
bifolius, R. Br. Prod. 322 = Chiloglottis diphylla.
caudatus, *R. Br. l. c.* 321.—Austral.
elegans, *Reichb. f. in Linnaea*, xli. (1877) 56.—N. Caled.
exsertus, *R. Br. Prod.* 321.—Austral.
fornicatus, *R. Br. l. c.*—Austral.
petiolatus, D. Don, Prod. Fl. Nep. 29 = Liparis nepalensis.
rivularis, *A. Cunn. in Hook. Comp. Bot. Mag.* ii. (1836) 376.—N. Zel.
Sinclairii, *Hook. f. Fl. N. Zeal.* i. 246.—N. Zel.
viridis, *Hook. f. Fl. Tasm.* ii. 372.—Austral.

ACIALYPTUS, A. Gray, Bot. U. S. Expl. Exped. i. 551. t. 67. (1854). *MYRTACEAE*, Benth. & Hook. f. i. 708.
Fullagari, *F. Muell. Fragm.* viii. 15.—Ins. D. Howe.
myrtoides, *A. Gray, Bot. U. S. Expl. Exped.* i. 551. t. 67.—Ins. Fiji.
myrtoides, Seem. in Bonplandia, ix. (1861) 206 = Seemanni.
nitida, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 186; *et in Ann. Sc. Nat.* Sér. V. iii. (1865) 227.—N. Caled.
Seemanni, *A. Gray, in Bonplandia*, x. (1862) 35.—Ins. Fiji.

ACICARPA, Raddi, Agrost. Bras. 31 (1823) = **Panicum**, Linn. (Gramin.).
sacchariflora, Raddi, l. c. t. 1. f. 4 = P. leucophaeum.

ACICARPA, R. Br. in Trans. Linn. Soc. xii. (1818) 132 = seq.

ACICARPHA, Juss. in Ann. Mus. Par. ii. (1803) 347. t. 58. *CALYCEREAE*, Benth. & Hook. f. ii. 162.
ACANTHOSPERMA, Vell. Fl. Flum. Ic. viii. t. 152 (1827).
CRYPTOCARPHA, Cass. in Bull. Soc. Philom. (1817) 34.
ECHINOLEMA, Jacq. f. ex DC. Prod. v. 3 (1836).
SOMMEA, Bory, in. Ann. Gén. Sc. Phys. vi. (1820) 92. t. 87.
bupleuroides, Less. in Linnaea, vi. (1831) 527 = Boopis bupleuroides.
crassifolia, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 402.—Bras.
lanata, *Lag. ex Pers. Syn.* ii. 488.—Mexico.
pinnatifida, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 404.—Bras.
procumbens, *Less. in Linnaea*, vi. (1831) 527.—Bras.
runcinata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 404.—Bras.
spathulata, *R. Br. in Trans. Linn. Soc.* xii. (1818) 129.—Bras.
tribuloides, *Juss. in Ann. Mus. Par.* ii. (1803) 348. t. 58.—Reg. Argent.

ACICARPHAEA, Walp. Ann. ii. 877 (1852) = **Chaenactis**, DC. (Compos.).
artemisiaefolia, Walp. l. c. = C. artemisiaefolia.

ACICLINIUM, Torr. & Gray, Fl. N. Am. ii. 231 in adnot. (1841) = **Bigelowia**, DC. (Compos.).

ACIDANDRA, Mart. ex Spreng. Gen. i. 388 (1830) = **Zollernia**, Mart. (Legumin.).

ACIDANTHERA, Hochst. in Flora, xxvii. (1844) 25. *IRIDEAE*, Benth. & Hook. f. iii. 706.
SPHAEROSPORA, Klatt, in Linnaea, xxxii. (1863) 725.
HOUTTUYNIA, Houtt. Handl. Plantenk. xii. 448. t. 85 (1780).
acquinoctialis, *Baker, in Journ. Linn. Soc.* xvi. (1877) 160.—Afr. trop.
bicolor, *Hochst. in Flora*, xxvii. (1844) 25.—Abyss.
brachystachys, *Baker, in Journ. Bot.* xiv. (1876) 338.—Afr. trop.
brevicollis, *Baker, l. c.* 339.—Afr. trop.
excapa, Baker, ex Bouché & Wittm. in Berl. Monatschr. Bef. Gartenb. xix. (1876) 15 = bulbulosa.
flexuosa, *Baker, ex Bouché & Wittm. l. c.*—Afr.

ACIDANTHERA:—
graminifolia, *Baker, in Journ. Bot.* xiv. (1876) 338.—Afr. trop.
Huttoni, *Baker, l. c.* 339.—Afr. trop.
platypetala, *Baker, l. c.*—Afr. austr.
tubulosa, *Baker, in Journ. Linn. Soc.* xvi. (1877) 160.—Afr. trop.
unicolor, *Hochst. ex Baker, l. c.*—Abyss.

ACIDOCROTON, Griseb. Fl. Brit. W. Ind. 42 (1864). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 291.
adelioides, *Griseb. l. c.*—Ind. occ.

ACIDOTON, Sw. Prod. Veg. Ind. Occ. 83 (1788). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 328.
innocuus, *Baill. Etud. Gén. Euphorb.* 402. t. 18. f. 10–11.—Ind. occ.
urens, *Sw. Prod. Veg. Ind. Occ.* 83.—Jamaica.

ACILEPIS, D. Don, Prod. Fl. Nep. 169 (1825) = **Vernonia**, Schreb. (Compos.).
cirsiifolia, Spreng. Syst. iii. 387 = Pacourina cirsiifolia.
lanata, Spreng. ex DC. Prod. vii. 36 = Dicoma tomentosa.
squarrosa, D. Don, Prod. Fl. Nep. 169 = V. teres.

ACINAX, Rafin. Fl. Tellur. iv. 53 (1836) = **Costus**, Linn. (Scitam.).

ACINETA, Lindl. Bot. Reg. (1843) Misc. 67. *ORCHIDEAE*, Benth. & Hook. f. iii. 551.
NEIPPERGIA, Morr. in Ann. Soc. Agr. Gand, v. (1849) 375. t. 282.
Barkeri, *Lindl. in Bot. Reg.* (1843) *Misc.* 68.—Mexico.
chrysantha, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 31.—Am. centr.
cryptodonta, *Reichb. f. in Bonplandia*, ii. (1854) 92.—Am. centr.
densa, *Lindl. & Paxt. Flow. Gard.* i. 91 (1850–51).—Am. centr.
erythroxantha, *Reichb. f. in Bonplandia*, ii. (1854) 92.—Hab.?
fulva, Klotzsch, in Otto & Dietr. Allg. Gartenz. (1852) 146 = Humboldtii.
glauca, Linden, ex Lindl. & Paxt. Flow. Gard. i. (1850–51) 123 = Cycnoches Pescatorei.
Hrubyana, *Reichb. f. in Gard. Chron.* N. S. xviii. (1882) II. 102.—N. Granat.
Humboldtii, *Lindl. Bot. Reg.* (1843) *Misc.* 68.—Am. centr.
longiscapa, Reichb. f. in Bot. Zeit. x. (1852) 706 = Lacaena bicolor.
Schilleriana, *Reichb. f. in Walp. Ann.* vi. 609.—Hab.?
Sella-turcica, *Reichb. f. in Bot. Zeit.* x. (1852) 705.—Am. centr.
sulcata, *Reichb. f. in Gard. Chron.* N. S. xi. (1879) I. 652.—Ecuador?
superba, Reichb. in Walp. Ann. vi. 609 = Humboldtii.
Warscewiczii, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 145 = densa.

ACINODENDRUM, Linn. Gen. ed. I. 129 (1737) = **Melastoma**, Burm.?
aurea, Rafin. Sylva Tellur. 94.—Hab.?
cymosa, Rafin. l. c. = Amphiblema cymosum?
laxiflora, Rafin. l. c.—Hab.?

ACINOLIS, Rafin. Sylva Tellur. 95 (1838) = **Melastoma**, Burm.
elliptica, Rafin. l. c. = M. ellipticum?

ACINOS, Rupp. Fl. Jen. ed. Hall. 233 (1745); Moench, Meth. 407 (1794) = **Calamintha**, Lam. (Labiat.).
acuminatus, Frivald. in Flora, xviii. (1835) 332 = C. suaveolens.
acutifolius, Schur, Enum. Pl. Transs. 530 = C. alpina.
adscendens, Moench, Meth. Suppl. 138 = C. patavina.
aestivalis, Kit. in Linnaea, xxxii. (1863) 432.—Europ.
alpinus, Moench, Meth. 407 = C. alpina.
canus, Reichb. Fl. Germ. Excurs. 327 = C. graveolens.
clinopodiifacie, Gilib. Fl. Lituan. i. 75 = C. Acinos.
corsicus, G. Don, in Sweet, Hort. Brit. ed. III. 768.—Corsica.
diffusus, Boenningh. Prod. Fl. Monast. 182.—Europ.

ACINOS:—
erectus, Frivald. in Flora, xviii. (1835) 333 = C. graveolens.
exiguus, G. Don, ex Steud. Nom. ed. II. i. 17 = C. graveolens.
grandiflorus, G. Don, ex Loud. Hort. Brit. 239 = C. grandiflora.
graveolens, Link, Enum. Hort. Berol. ii. 117 = C. graveolens.
Herba-barona, G. Don, in Loud. Hort. Brit. 239 = Thymus Herba-Barona.
heterophyllus, G. Don, l. c. = C. suaveolens.
incanus, Griseb. Spicil. Fl. Rumel. ii. 123 = C. maritima.
multiflorus, C. Koch, ex Boiss. Fl. Orient. iv. 579 = C. umbrosa.
patavinus, Pers. Syn. ii. 131. (err. typ. 151) = C. alpina.
purpurascens, Pers. l. c. = C. graveolens.
rotundifolius, Frivald. ex Walp. Rep. iii. 730 = C. alpina.
rotundifolius, Pers. Syn. ii. 131. (err. typ. 151) = C. Nepeta.
suaveolens, G. Don, ex Loud. Hort. Brit. 239 = C. suaveolens.
thymoides, Moench, Meth. 407 = C. Acinos.
transsilvanica, Schur, Enum. Pl. Transs. 529 = C. Nepeta.
villosus, Pers. Syn. ii. 131. (err. typ. 151) = C. Acinos.
vulgaris, Pers. l. c. = C. Acinos.

ACINOTUM, Reichb. Handb. 260 (1837) = **Mathiola**, R. Br. (Crucif.).

ACIOA, Aubl. Pl. Gui. ii. 698. t. 280 (1775) = **Couepia**, Aubl. (Rosac.).
amara, Steud. Nom. ed. I. 9 = C. guianensis.
Bellayana, Baill. Adansonia, vii. (1866–67) 224 = Griffonia Bellayana.
dulcis, Steud. Nom. ed. I. 9 = C. guianensis.
guianensis, Aubl. Pl. Gui. ii. 698 = C. guianensis.
Icondere, Baill. Adansonia, vii. (1866–67) 223 = Griffonia Icondere.
pallescens, Baill. l. c. 224 = Griffonia pallescens.
Smeathmanni, Baill. l. c. x. (1872) 244.—Afr. trop.

ACIOTIS, D. Don, in Mem. Wern. Soc. iv. (1823) 300. *MELASTOMACEAE*, Benth. & Hook. f. i. 744.
SPENNERA, Mart. in DC. Prod. iii. 115 (1828).
acuminifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Am. trop.
acuminifolia, Triana, l. c. = sphaeranthera, etc.
acutiflora, *Triana, l. c.* 52. pro parte.—Am. trop.
acutiflora, Triana, l. c. pro parte = laxa.
aequatorialis, *Cogn. in Mart. Fl. Bras.* xiv. III. 464.—Bras.
amazonica, *Cogn. l. c.* 461.—Bras.
annua, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 52.—Am. merid.
aquatica, *G. Don, in Loud. Hort. Brit.* 174.—Guiana.
brachybotria, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Bras.
caulialata, *Triana, l. c.* 52.—Peruv.
circaeifolia, *Triana, l. c.*—Bras.; Guiana.
dichotoma, *Cogn. in Mart. Fl. Bras.* xiv. III. 460.—Guiana.
discolor, *D. Don, in Mem. Wern. Soc.* iv. (1823) 301.—Ins. St. Vincent.
dysophylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 52.—Bras.; Peruv.
ferruginea, Triana, l. c. 51 = brachybotria.
fragilis, *Cogn. in Mart. Fl. Bras.* xiv. III. 475.—Guiana.
herbacea, *Cogn. l. c.* 468.—Bras.
indecora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 52.—Bras.; Guiana.
latifolia, *Cogn. in Mart. Fl. Bras.* xiv. III. 464.—Bras.
laxa, *Cogn. l. c.* 476.—Bras.
Levyana, *Cogn. l. c.* 460.—Nicarag.
longifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 52.—Bras.
Martiana, *Cogn. in Mart. Fl. Bras.* xiv. III. 463.—Bras.
paludosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Bras.

ACIOTIS:—
pendulifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 52.—Guiana.
polystachya, *Triana, l. c.* 52.—Bras.
purpurascens, *Triana, l. c.* 52. t. 3. f. 36.—Am. trop.
rostellata, *Triana, l. c.* 51.—Mexico.
rubricaulis, *Triana, l. c.* 52. t. 3. f. 36 b.—Bras.
Sieberi, Triana, l. c. 52 = Nepsera aquatica.
sileniflora, *Triana, l. c.* 52.—Bras.
sphaeranthera, *Cogn. in Mart. Fl. Bras.* xiv. III. 458.—Peruv.
uliginosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Venezuela.
viscosa, *Triana, l. c.* 52.—Am. trop.

ACIPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 577 = **Cambessedesia**, DC. (Melast.).
lineatum, Turcz. l. c. 578 = Pyramia pityrophylla.
rugosum, Turcz. l. c. = C. rugosa.

ACIPHYLLA, Forst. Char. Gen. 135. t. 68 (1776).—*UMBELLIFERAE*, Benth. & Hook. f. i. 912.
ANISOTOME, Hook. f. Fl. Antarct. i. 16. t. 8, 9 (1845).
CALOSCIADIUM, Endl. Gen. Suppl. iv. III. 7 (1850).
GINGIDIUM, F. Muell. in Trans. Phil. Inst. Vict. i. (1855) 103.
Colensoi, *Hook. f. Handb. N. Zeal. Fl.* 92.—N. Zel.
crenulata, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 336.—N. Zel.
Dobsoni, *Hook. f. Handb. N. Zeal. Fl.* 93.—N. Zel.
glacialis, *F. Muell. ex Benth. Fl. Austral.* iii. 375.—Austral.
Hectori, *J. Buch, in Trans. N. Z. Inst.* xiv. (1882) 346.—N. Zel.
Lyallii, *Hook. f. Handb. N. Zeal. Fl.* 92.—N. Zel.
Monroi, *Hook. f. Fl. N. Zel.* ii. 330.—N. Zel.
montana, *Armst. in Trans. N. Z. Inst.* iv. (1872) 290.—N. Zel.
procumbens, *F. Muell. ex Benth. Fl. Austral.* iii. 375.—Austral.
simplicifolia, *F. Muell. ex Benth. l. c.*—Austral.
squarrosa, *Forst. Char. Gen.* 136.—N. Zel.
Traillii, *T. Kirk, in Trans. N. Z. Inst.* xvi. (1884) 371.—N. Zel.
Traversii, *F. Muell. ex J. Buch, l. c.* vii. (1875) 335.—N. Zel.

ACIPHYLLAEA, A. Gray, in Mem. Am. Ac. N. S. iv. (1849) (Pl. Fendl.) 91 = **Hymenatherum**, Cass. (Compos.).
acerosa, A. Gray, l. c. = H. acerosum.

ACIPHYLLUM, Steud. Nom. ed. II. i. 17 (1840) = **Chorizema**, Labill. (Legum.).

ACIS, Salisb. Parad. Lond. sub t. 74 (1807) = **Leucojum**, Linn. (Amaryll.).
autumnalis, Salisb. l. c. = L. autumnale.
Broteri, Jord. & Fourr. Ic. Pl. Europ. 25 = L. trichophyllum.
grandiflora, Sweet, Brit. Fl. Gard. iii. sub t. 297 = L. trichophyllum.
hiemalis, M. Roem. Syn. Ensat. 24 = L. hyemale.
hyemalis, Sweet, Brit. Fl. Gard. iii. sub t. 297 = praec.
longifolia, M. Roem. Syn. Ensat. 25 = L. longifolium.
rosea, Sweet, Brit. Fl. Gard. t. 297 = L. hyemale.
oporantha, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 51 = L. autumnale.
pulchella, Jord. & Fourr. l. c. = L. autumnale.
roseus, Sweet, ex Loud. Hort. Brit. Supp. iii. 482 = L. roseum.
trichophylla, Sweet, Brit. Fl. Gard. iii. sub t. 297 = L. trichophyllum.

ACISANTHERA, P. Br. Hist. Jamaic. 217. t. 22. f. i. (1756). *MELASTOMACEAE*, Benth. & Hook. f. i. 739.
ANISOCENTRUM, Turcz. in Bull. Soc. Nat. Mosc. xxxv. (1862) I. 322.
DICRANANTHERA, Presl, Symb. Bot. i. 75. t. 50 (1832).
EXODICILIS, Rafin. Sylva Tellur. 97 (1838).
MIOCARPUS, Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 146.
NOTEROPHILA, Mart. Nov. Gen. et Sp. iii. 110. t. 254 (1829).
URANTHERA, Naud. in Ann. Sc. Nat Sér. III. ii. (1844) 146.

ACISANTHERA:—
adscendens, Wright, in Sauvalle, Cat. Fl. Cub. 52 = quadrata.
alata, *Cogn. in Mart. Fl. Bras.* xiv. III. 225.—Bras.
alsinaefolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 34. t. 2. f. 18 h.—Bras.
Beccabunga, Triana, l. c. 33 = brevifolia.
bivalvis, Cogn. in Mart. Fl. Bras. xiv. III. 216 = brevifolia.
Boissieriana, *Cogn. l. c.* 209.—Bras.
brevifolia, *Griseb Fl. Brit. W. Ind.* 269.—Ind. occ.; Bras.
brevifolia, Triana, in Trans. Linn. Soc. xxviii. (1871) 33 = trivalvis.
divaricata, *Cogn. in Mart. Fl. Bras.* xiv. III. 212.—Bras.
fluitans, *Cogn. l. c.* 214.—Bras.
Glazioviana, *Cogn. l. c.* 223.—Bras.
hedyotoidea, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 33.—Bras.
hedyotoidea, Triana, l. c. pro parte = Salzmanni.
inundata, *Triana, l. c.*—Bras.
limnobios, *Triana, l. c.* t. 11.—Am. trop.
pellucida, *Wright, in Griseb. Cat. Pl. Cub.* 104.—Cuba.
pulchella, *Cogn. in Mart. Fl. Bras.* xiv. III. 211.—Bras.
punctatissima, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 34. t. 11. f. 18 c.—Bras.; N. Granat.
quadrata, *Juss. ex Poir. Encyc. Suppl.* i. 111.—Jamaica.
recurva, *Griseb. in Goett. Abh.* ix. (1861) 58.—Bras.; Guiana.
recurva, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 34. t. 2. f. 18 b.—Bras.
Salzmanni, *Cogn. in Mart. Fl. Bras.* xiv. III. 209.—Bras.
tetramera, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 33.—Bras.
trivalvis, *Cogn. in Mart. Fl. Bras.* xiv. III. 217.—Bras.
variabilis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 34. t. 11. f. 18 g.—Bras.

ACISPERMUM, Neck. Elem. i. 34 (1790) = **Coreopsis**, Linn. (Compos.).

ACISTOMA. Zipp. ex Span. in Linnaea, xv. (1841) 202 = **Woodfordia**, Sm. (Lythrar.).
coccineum, Zipp. ex Span. l. c. = W. floribunda.

ACKAMA, A. Cunn. in Ann. Nat. Hist. ii. (1839) 358. *SAXIFRAGEAE*, Benth. & Hook. f. i. 653.
Muelleri, *Benth. Fl. Austral.* ii. 444.—Austral.
rosaefolia, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 358.—N. Zel.

ACLADODEA, Ruiz & Pav. Prod. 133. t. 29 (1794) = **Talisia**, Aubl. (Sapind.).
pinnata, Ruiz & Pav. Syst. Veg. 262 = T. Acladodea.

ACLEIA, DC. Prod. vi. 340 (1837) = **Senecio**, Linn. (Compos.).
belbeysia, DC. l. c. = S. belbeysius.

ACLEISANTHES, A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 259. *NYCTAGINEAE*, Benth. & Hook. f. iii. 6.
PENTACROPHYS, *A. Gray, l. c.* 261 (1853).
anisophylla, *A. Gray, l. c.*—Texas.
Berlandieri, *A. Gray, l. c.* 260.—Texas.
crassifolia, *A. Gray, l. c.*—Texas.
longiflora, *A. Gray, l. c.*—N. Mexico.
Wrightii. *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 6.—Texas.

ACLINIA, Griff. Notul. iii. 320 (1851) = **Dendrobium**, Sw. (Orchid.).

ACLISIA, E. Mey. in Presl, Rel. Haenk. i. 137. t. 25 (1827) = **Pollia**, Thunb. (Commel.).
celebica, Hassk. Commel. Ind. 49 = P. macrophylla.
Cumingiana, Hassk. l. c. 47 = P. macrophylla.
elegans, Hassk. in Miq. Pl. Jungh. 49 = P. sorzogonensis.
florida, Hort. Berol. (1837) ex Kunth, Enum. Pl. iv. 47 = Athyrocarpus pallidus.
gigantea, Hassk. Commel. Ind. 46 = P. sorzogonensis.
indica, Wight, Ic. t. 2068 = P. sorzogonensis.
sorsogonensis, Kunth, Enum. Pl. iv. 74 = P. indica.
sorzogonensis, E. Mey. in Presl, Rel. Haenk. i. 138. t. 25 = P. sorzogonensis.

ACLISIA:—
guayaquilensis, *G. Don, Gen. Syst.* iv. 461.—Peruv.
subumbellata, C. B. Clarke, Commel. et Cyrt. Beng. t. 30 = P. subumbellata.
Thomsoni, C. B. Clarke, l. c. t. 31 = Aneilema Thomsoni.
Zollingeri, Hassk. Commel. Ind. 51 = P. Zollingeri.

ACMADENIA, Bartl. & Wendl. f. Diosm. 59. t. A (1824). *RUTACEAE*, Benth. & Hook. f. i. 289.
alternifolia, *Cham. in Linnaea*, v. (1830) 52.—Afr. austr.
assimilis, *Sond. in Harv. & Sond. Fl. Cap.* i. 383.—Afr. austr.
cassiopoides, Turcz. in Bull Soc. Nat. Mosc. xxxi. (1858) I. 439 = Macrostylis dubia.
cucullata, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 383.—Afr. austr.
densifolia, *Sond. l. c.*—Afr. austr.
flaccida, *Eckl. & Zeyh. Enum.* 105.—Afr. austr.
juniperina, *Bartl. & Wendl. f. Diosm.* 61.—Afr. austr.
laevigata, *Bartl. & Wendl. f. l. c.* 64.—Afr. austr.
laevigata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 383 = assimilis.
Mundiana, *Eckl. & Zeyh. Enum.* 105.—Afr. austr.
muraltioides, Eckl. & Zeyh. l. c. 106 = juniperina.
Nivenі, *Sond. in Harv. & Sond. Fl. Cap.* i. 380.—Afr. austr.
obtusata, *Bart. & Wendl. f. Diosm.* 63.—Afr. austr.
obtusata, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. i 376 = Diosma tercfolia.
psilopetala, *Sond. in Harv. & Sond. l. c.* i. 380.—Afr. austr.
pungens, *Bartl. & Wendl. f. Diosm.* 64.—Afr. austr.
pungens, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. i. 381 = obtusata.
rosmarinifolia, *Bartl. in Linnaea*, xvii. (1844) 355.—Afr. austr.
strobilina, E. Mey. ex Harv. & Sond. Fl. Cap. i. 381 = tetragona.
tetragona, *Bartl. & Wendl. f. Diosm.* 65.—Afr. austr.
tetragona, Eckl. & Zeyh. Enum. 105 = laevigata.

ACMANTHERA, Griesb. in Mart. Fl. Bras. xii. I. 28 (1858). *MALPIGHIACEAE*, Benth. & Hook. f. i. 253.
latifolia, *Griseb. l. c.* 29.—Bras.

ACMELLA, Rich. in Pers. Syn. ii. 472 (1807) = **Spilanthes**, Jacq. (Compos.).
biflora, Spreng. Syst. iii. 591 = Wedelia biflora.
brachyglossa, Cass. in Dict. Sc. Nat. l. 258 = S. Acmella.
brasiliensis, Spreng. Syst. iii. 592 = Wedelia paludosa.
buphthalmoides, Pers. Syn. ii. 473 = Heliopsis buphthalmoides.
caulorrhiza, Delile, Cent. Pl. Afr. Méroé, 45. t. iii. f. 7 = S. Acmella.
ciliata, Cass. in Dict. Sc. Nat. xxiv. 331 = S. ciliata.
debilis, Cass. l. c. 331 = S. debilis.
fimbriata, Cass. l. c. = S. fimbriata.
flavicaulis, Rafin. New Fl. Am. i. 52.—Am. bor.
Garcini, Spreng. Syst. iii. 591 = Anvillea Garcini.
globosa, Spreng. l. c. 592 = Zaluzania globosa.
hirta, Lag. Gen. et Sp. Nov. 30 = Jaegeria hirta.
intermedia, Rich. ex Pers. Syn. ii. 472 = S. intermedia.
lanceolata, Link, ex Spreng. Syst. iii. 444 = S. Pseudo-acmella.
Linnaei, Cass. in Dict. Sc. Nat. xxiv. 330 = S. Acmella.
mauritanica, Rich. ex Pers. Syn. ii. 472 = S. Acmella.
Mutisii, Cass. in Dict. Sc. Nat. xxiv. 331 = S. Mutisii.
nudicaulis, Rafin. New Fl. Am. i. 52.—Am. bor.
Nuttaliana, Rafin. l. c. = S. repens.
occidentalis, Nutt. Gen. Am. ii. 171 = S. repens.
occidentalis, Rich. ex Pers. Syn. ii. 473 = S. Mutisii.
parvifolia, Rafin. New Fl. Am. i. 52.
repens, Rich. ex. Pers. Syn. ii. 473 = S. repens.
spilanthoides, Cass. Dict. Sc. Nat. xxiv. 329 = Wedelia carnosa.
tenella, Cass. in l. c. xxiv. 331 = S. tenella.
trilobata, Spreng. Syst. iii. 591 = Zaluzania triloba.
uliginosa, Cass. in Dict. Sc. Nat. xxiv. 331 = S. Acmella.

ACMENA, DC. in Dict. Class. Hist. Nat. xi. 401 (1826) = **Eugenia**, Linn. (Myrtac.).
bracteolata, Walp. Rep. ii. 181 = E. bracteolata.
Championi, Benth. in Hook. Kew Journ. Bot. i. (1852) 118.—Hongkong.
chinensis, Planch. Hort. Donat. 84.—China.
claviflora, Walp. Rep. ii. 181 = E. claviflora.
elliptica, G. Don, ex Steud. Nom. ed. II. i. 17 = E. Smithii.

ACMENA:—
floribunda, DC. in Dict. Class. Hist. Nat. xi 401; et Prod. iii. 262 = E. Smithii.
Gerrardi, Harv. Gen. S. Afr. Pl. ed. II. 112.—Afr. austr.
grata, Walp. Rep. ii. 181 = E. grata.
inophylla, Wight, Ic. ii. t. 623 = E. inophylla.
Kingii, G. Don, Gen. Syst. ii. 851 = E. Smithii.
lanceolata, Thw. Enum. Pl. Zeyl. 119 = E. Wightiana.
leptantha, Walp. Rep. ii. 181 = E. leptantha.
oblata, Walp. l. c. = E. oblata.
parviflora, DC. Prod. iii. 262 = E. zeylanica.
Wightiana, Walp. Rep. ii. 181 = E. Wightiana.
zeylanica, Thw. Enum. Pl. Zeyl. 118 = E. zeylanica.

ACMISPON, Rafin. Atl. Journ. 144 (1832) = **Hosackia**, Dougl. (Legumin.).
multiflorum, Rafin. New Fl. Am. i. 53 = seq.?
sericeum, Rafin. Atl. Journ. 145 = H. sericea.

ACMOSTIMA, Rafin. Sylva Tellur. 26 (1838) = **Pavetta**, Linn. (Rubiac.).
brevifolia, Rafin. l. c. = P. pentandra.
longifolium, Rafin. l. c. = P. barbata.

ACNADENA, Rafin. Sylva Tellur. 41 (1838) = **Cordia**, Linn. (Borag.).
elliptica, Rafin. l. c. 41 = C. elliptica.

ACNIDA, Linn. Diss. Chen. 32 (1750) (Am. acad. iii. 19). *AMARANTACEAE*, Benth. & Hook. f. iii. 2. MONTELIA, A. Gray, Man. Bot. N. U. States, ed. II. 369 (1856).
altissima, Michx. ex. Moq. in DC. Prod. xiii. II. 277 = rusocarpa.
altissima, Ridd. ex Moq. l. c. = tuberculata?
australis, *A. Gray, in Am. Natural.* x. (1876) 489.—Am. bor.
cannabina, Chapm. Fl. S. U. St. Suppl. 645 = australis.
cannabina, *Linn. Sp. Pl.* 1027.—Am. bor.
cuspidata, Bert. ex Spreng. Syst. iii. 903 = cannabina.
Ellioti, *Rafin. New Fl. Am.* i. 54.—Am. bor.
floridana, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 376.—Am. bor.
obtusifolia, *Rafin. New Fl. Am.* i. 54.—Am. bor.
odorata, *F. G. Dietr. ex Moq. in DC. Prod.* xiii. II. 278, nomen.—Hab.?
rhyssocarpa, Steud. Nom. ed. II. i. 17 = rusocarpa.
ruscocarpa, Willd. Sp. Pl. iv. 768 = rusocarpa.
rusocarpa, Moq. in DC. Prod. xiii. II. 277 = tuberculata.
rusocarpa, *Michx. Fl. Bor. Am.* ii. 234. t. 50.—Am. bor
salicifolia, *Rafin. in Am. Monthly Mag.* ii. (1817) 43.—Am. bor.
tamariscina, *A. Gray, in Am. Natural.* x. (1876) 489.—Am. bor.
tuberculata, *Moq. in DC. Prod.* xiii. II. 277.—Am. bor.

ACNISTUS, Schott, in Wien. Zeitschr. (1829) iv, 1180, ex Linnaea, vi. (1831) Litt. 54. *SOLANACEAE*. Benth. & Hook. f. ii. 894.
CODOCHONIA, Dun. in DC. Prod. xiii. I. 482 (1852).
LYCIOPLESIUM, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 330.
EPHAILA, HUANUCA, KOKABUS, PEDERLEA, PLICULA, TROZELIA, Rafin. Sylva Tellur. 54-55, 124 (1838).
aggregatus, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 341.—Peruv.
arborescens, *Schlecht. in Linnaea*, vii. (1832) 67.—Mexico.
Benthami, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 342.—Mexico.
breviflorus, *Sendtn. in Mart. Fl. Bras.* x. 152.—Bras.
cauliflorus, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 337 = Miersi.
cauliflorus, *Schott, in Wien. Zeitschr.* (1829) iv. 1180.—Bras.
cestroides, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 343 = Lycium cestroides.
confertiflorus, *Miers, in Ann. et Mag. Nat. Hist.* Ser. II. iv. (1849) 31.—Peruv.
ellipticus, *Hook. f. in Trans. Linn. Soc.* xx. (1850) 205.—Ins. Galap.
fasciculatus, *Benth. & Hook. f. Gen.* ii. 894.—Boliv.
floribundus, *G. Don, Gen. Syst.* iv. 461.—Peruv.
frutescens, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 299.—Porto Rico.
grandiflorus, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 344.—Venezuela.

ACNISTUS:—
guayaquilensis, *G. Don, Gen. Syst.* iv. 461.—Peruv.
Miersi, *Dun. in DC. Prod.* xiii. 1. 497.—Bras.
parviflorus, *Griseb. in Goett. Abh.* xix. (1874) 219.—Reg. Argent.
Plumieri, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 340.—Am. austr.
ramiflorus, *Miers, l. c.* 342.—Ind. occ.
sideroxyloides, *G. Don, Gen. Syst.* iv. 461.—N. Granat.
spatulatus, *G. Don, l. c.*—Peruv.
umbellatus, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 342.—Peruv.
virgatus, *Griseb. in Goett. Abh.* xxiv. (1879) 247.—Reg. Argent.

ACOELORRHAPHE, H. Wendl. in Bot. Zeit. xxxvii. (1879) 148. *PALMAE.*

ACOIDIUM, Lindl. Bot. Reg. (1837) sub t. 1951 = **Trichocentrum**, Poepp. & Endl. (Orchid.).
fuscum, Lindl. l. c. = T. fuscum.

ACOKANTHERA, G. Don, Gen. Syst. iv. 485 (1837). *APOCYNACEAE*, Benth. & Hook. f. ii. 696.
TOXICOPHLAEA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 24.
Lamarkii, *G. Don, Gen. Syst.* iv. 485.—Afr. austr.
lycioides, *G. Don, l. c.*—Afr. austr.
pubescens, *G. Don, l. c.* 486.—Afr. austr.
spectabilis, *Hook. f. Bot. Mag.* t. 6359.—Afr. austr
venenata, *G. Don, Gen. Syst.* iv. 485.—Afr. austr.

ACOMA, Benth. Bot. Voy. Sulph. 29. t. 17 (1844) = **Coreocarpus**, Benth. (Compos.).
dissectum, Benth. l. c. (= *Leptosyne dissecta*).

ACOMIS, F. Muell. Fragm. ii. 89 (1860). *COMPOSITAE*, Benth. & Hook. f. ii. 317.
macra, *F. Muell. l. c.* iv. 145.—Austral.
rutidosea, *F. Muell. l. c.* ii. 89.—Austral.

ACONCEVEIBUM, Miq. Fl. Ind. Bat. i. II. 389 (1859). *EUPHORBIACEAE.*
trinerve, *Miq. l. c.*—Malaya.

ACONITUM, Tourn. ex Linn. Syst. ed. I. (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 9.
ANTHORA, Hall. in Rupp. Fl. Jen. ed. Hall. 290 (1745).
CAMMARUM, Fourr. in. Ann. Soc. Linn. Lyon. N. S. xvi. (1868) 327.
LYCOCTONUM, Fourr. l. c. 326.
NAPELLUS, Rupp. Fl. Jen. ed. Hall. 275 (1745).
NIRBISIA, G. Don, Gen. Syst. i. 63 (1831).
abbreviatum, Langsd. ex DC. Prod. i. 61 = Fischeri.
achranthum, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 285.—Sib. altaic.
acuminatum, Reichb. Uebers. Acon. 46 = paniculatum.
acutum, Reichb. l. c. 21 = A. Napellus.
aegophonum, Reichb. l. c. 69 = A. Lycoctonum.
alatum, Wender. in Linnaea, xv. (1841) Litt. 99 = A. Napellus.
albicans, *Host, Fl. Austr.* ii. 76.—Europ.
albidum, Bernh. ex Reichb. Uebers. Acon. 31 = A. Napellus.
album, Ait. Hort. Kew. ii. 246 = variegatum.
alienum, Reichb. Uebers. Acon. 73. = A. Lycoctonum.
alpinum, Mill. Gard. Dict. ed. VIII. n. 7 = variegatum.
altigaleatum, Hayne, Arzn. Gew. xii. t. 16 = variegatum.
altissimum, Mill. Gard. Dict. ed. VIII. n. 2 = A. Lycoctonum.
ambiguum, Reichb. Monog. Acon. t. 23 = A. Napellus.
amoenum, Reichb. Uebers. Acon. 23 = A. Napellus.
ampliflorum, Reichb. l. c. 37 = A. Napellus.
angustifolium, Bernh. ex Reichb. l. c. 29 = A. Napellus.
Anthora, *Linn. Sp. Pl.* 532.—Europ.; Oriens.; As. bor.
anthorum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Anthora.
anthoroideum, DC. Syst. i. 366 = A. Anthora.
arctophonum, Reichb. Uebers. Acon. 71 = A. Lycoctonum.
arcuatum, *Maxim. Prim. Fl. Amur.* 27.—Reg. Amur.
Atees, Royle, in Journ. As. Soc. Beng. i. (1832) 459 = heterophyllum.
atlanticum, *Coss. in Bull. Soc. Fr.* xxii. (1875) 53, nomen.—Afr. bor.
atrox, Walp. Rep. i. 57 = ferox.
australe, Reichb. Uebers. Acon. 71 = A. Lycoctonum.

ACONITUM:—
austriacum, Tratt. ex Reichb. Monog. Acon. '18' (t. 71) = A. Cammarum.
autumnale, Lindl. in Journ. Hortic. Soc. ii. (1847) 77 = Fischeri.
autumnale, Reichb. Monog. Acon. t. 17. f. 2 = A. Napellus.
baicalense, *Turcz. ex Steud. Nom.* ed. II. i. 18.—Sibir. baical.
barbatum, Patr. ex Pers. Syn. ii. 83 = A. Lycoctonum.
Baumgarteni, *Schur, in Verh. Siebenb. Ver. Naturw.* iii. (1852) 84.—Europ.
Bernhardianum, Reichb. Uebers. Acon. 34 = A. Napellus.
Bernhardianum, Wallr. Sched. Crit. i. 250. t. 2 = variegatum.
Besserianum, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 58, nomen.—Podolia.
bicolor, Schult. Obs. Bot. 101 = A. Cammarum.
biflorum, *Fisch. ex DC. Syst.* i. 380.—Sibir. altaic.
boreale, Ser. ex Reichb. Uebers. Acon. 62 = A. Lycoctonum.
bracteosum, *Bercht. & Presl, Rostl.* i. *Ranunculac.* 127. nomen.—Europ.
Braunii, Reichb. Uebers. Acon. 29 = A. Napellus.
Breiterianum, Reichb. l. c. 54 = A. Cammarum.
bulbiferum, Reichb. l. c. 55 = variegatum.
caeruleum, Dulac, Fl. Hautes-Pyr. 223 = A. Napellus.
callibotryon, Reichb. Monog. Acon. 98. t. 16. f. 2 = A. Napellus.
Camarum, Schleich. Cat. (1821) 5 = paniculatum.
Cammarum, *Linn. Sp. Pl.* ed. II. 751.—Europ.
Cammarum, Jacq. Fl. Austr. v. t. 224 = variegatum.
camtschaticum, auct. = *kamtschaticum*.
Candollei, Reichb. ex Steud. Nom. ed. II. i. 18 = A. Anthora.
canescens, Schleich. ex Reichb. Fl. Germ. Excurs. 741 = A. Napellus.
Carmichaeli, Debeaux, in Act. Linn. Soc. Bord. xxxiii. (1879) 87 (Florule du Tientsin, 61) = Fischeri.
cernuum, Baumg. ex Schur, Enum. Pl. Transs. 32 = A. Cammarum.
cernuum, Wulf. ex Koelle, Spicil. 17 = paniculatum.
Chamissonianum, Reichb. Uebers. Acon. 37 = A. Napellus
chinense, Siebold, ex Paxt. Mag. v. (1838) 3, cum tab. = Fischeri.
chinense, *Sieb. ex Sieb. & Zucc. Fl. Jap. Fam. Nat.* 76 [75].—Japon.
ciliare, DC. Syst. i. 378 = villosum, volubile.
Clairvilleanum ×, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxiii.–xxiv. (1881) 82.—Europ.
Clusianum, Reichb. Monog. Acon. 13 = seq.
Clusii, Reichb. Uebers. Acon. 22 = A. Napellus.
columbianum, Nutt. in Torr. & Gray, Fl. N. Am. i. 34 = Fischeri.
commutatum, Reichb. Uebers. Acon. 36 = A. Napellus.
compactum, Reichb. l. c. 27 = A. Napellus.
confertum, Reichb. l. c. 36 = A. Napellus.
cordatum, Rafin. New Fl. Am. i. 56.—Am. bor.
cordatum, Royle, Illustr. 56 = heterophyllum.
cynoctonum, Reichb. Uebers. Acon. 73 = A. Lycoctonum.
Decandollii, Reichb. l. c. 16 = A. Anthora.
decorum, Reichb. l. c. 51 = A. Cammarum.
delphinifolium, DC. Syst. i. 380 = A. Napellus.
delphinifolium, Hort. Prag. ex Steud. Nom. ed. II. i. 18 = A. Lycoctonum.
densiflorum, Hoppe, ex Steud. Nom. ed. II. i. 18 = A. Napellus.
dissectum, D. Don, Prod. Fl. Nep. 197 = A. Napellus.
dissectum, Tausch, ex Reichb. Monog. Acon. t. 47 = A. Lycoctonum.
divergens, *Panč. Elem.* 18.—Europ.
divergens, *Rafin. New Fl. Am.* i. 56.—Am. bor.
elatum, Salisb. Prod. 375 = A. Napellus.
eminens, Koch, ex Reichb. Uebers. Acon. 35 = A. Napellus.
engadinense ×, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 82.—Europ.
eriostemum, DC. Syst. i. 377 = A. Cammarum.
eulophum, Reichb. Uebers. Acon. 15 = A. Anthora.
eustachium, Reichb. l. c. 24 = A. Napellus.
exaltatum, Bernh. ex Reichb. l. c. 51 = A. Cammarum.
excelsum, *Reichb. Monog. Acon.* t. 53.—Rossia; Sibir.
excelsum, Turcz. Cat. Baikal. n. 70 = A. Lycoctonum.
ferox, *Wall. in Ser. Mus. Helv.* i. 160.—Reg. Himal.

ACONITUM:—
ferox, Wall. Cat. n. 4721 A (Pl. As. Rar. t. 41) = A. Napellus.
firmum, Reichb. Uebers. Acon. 20 = A. Napellus.
Fischeri, *Reichb. Monog. Acon.* t. 22.—As. bor.; Am bor.
flaccidum, *Reichb. Uebers. Acon.* 39.—Sibir.
flexicaule, Hoppe & Hornsch. ex Reichb. Monog. t. 38 in syn. = paniculatum.
flexuosum, Presl, ex Reichb. l. c. t. 7 in syn. = variegatum.
flexuosum, *Rafin. New Fl. Am.* i. 55.—Am. bor.
formosum, Reichb. Uebers. Acon. 36 = A. Napellus.
fornicatum, Gilib. Fl. Lituan. ii. 285 = A. Napellus.
Funckianum, Reichb. Monog. Acon. t. 66 = A. Napellus.
Funckii, Reichb. Uebers. Acon. 28 = A. Napellus.
galeatum, Steven, ex Steud. Nom. ed. II. i. 18 = orientale.
galeriflorum, Stokes, Bot. Mat. Med. iii. 216 = A. Lycoctonum.
galectonum, Reichb. Uebers. Acon. 67 = A. Lycoctonum.
geraniifolium, *Host, Fl. Austr.* ii. 76.—Europ.
gibbiferum, Reichb. Monog. Acon. t. 19 = paniculatum.
gibbosum, Ser. Mus. Helv. i. 141. t. 15. f. 14, 15 = variegatum.
giganteum, Host, ex Steud. Nom. ed. II. i. 18 = A. Napellus.
giganteum, Zumag. Fl. Pedem. ii. 345 = A. Lycoctonum?
glabrum, DC. Syst. i. 379 = variegatum.
Gmelini, Reichb. Uebers. Acon. 63 = A. Lycoctonum.
gracile, Reichb. l. c. 55 = variegatum.
grandiflorum, [Fisch.] Hort. Gorenk. ed. I. (1808) 77 = biflorum.
grandiflorum, Pall.; cf. Reichb. Uebers. Acon. 38 = A. Napellus.
guttatum, Turcz. ex Regel, in Bull. Soc. Nat. Mosc. (1861) II. 114 = A. Napellus.
gymnandrum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 308.—China.
Halleri, Reichb. Uebers. Acon. 27 = A. Napellus.
hamatum, Hort. Berol. ex Steud. Nom. ed. II. i. 18 = variegatum.
hamatum, Reichb. in Cat. Hort. Berol. (1818); Uebers. Acon. 48 = A. Cammarum.
Hamiltonii, G. Don, Gen. Syst. i. 62 = A. Napellus.
hebegynum, DC. Syst. i. 376 = paniculatum.
heterophyllum, *Wall. Cat.* n. 4722.—Reg. Himal.
hians, *Host, Fl. Austr.* ii. 71.—Europ.
hians, Reichb. Fl. Germ. Excurs. 742 = A. Napellus.
hispidum, DC. Syst. i. 367 = A. Lycoctonum.
Hoppeanum, Reichb. Monog. Acon. t. 65 = A. Napellus.
Hoppii, Reichb. Uebers. Acon. 24 = A. Napellus.
hortense, Hoppe, ex Reichb. Monog. Acon. t. 71 in syn. = A. Cammarum.
Hosteanum, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 177 = A. Lycoctonum.
humile, Bernh. ex Reichb. Uebers. Acon. 34 = A. Napellus.
humile, Salisb. Prod. 375 = paniculatum.
illinitum, Reichb. Uebers. Acon. 54 = variegatum.
inclinatum, Sweet, Hort. Brit. ed. II. 9 = A. Anthora?
intermedium, DC. Syst. i. 374 = A. Cammarum.
intermedium, Gaud. ex Steud. Nom. ed. II. i. 18 = variegatum.
intermedium, Host, Fl. Austr. ii. 69 = A. Lycoctonum.
inunctum, Koch, ex Reichb. Monog. Acon. '6. app.' (Index) = A. Napellus.
italicum, Tratt. ex Reichb. l. c. t. 7. in syn. = variegatum.
Jacquini, Reichb. Uebers. Acon. 17 = A. Anthora.
Jacquinianum, Host, Fl. Austr. ii. 68 = A. Lycoctonum.
japonicum, *Decne. in Rev. Hortic.* (1851) 175.—Japon.
japonicum, Hort. ex Steud. Nom. ed. II. i. 18 = variegatum.
japonicum, Thunb. Fl. Jap. 231 = uncinatum.
kamtschaticum, Pall. ex Reichb. Uebers. 39 = Fischeri.
Koehleri, Reichb. Uebers. Acon. 30 = A. Napellus.
Koelleanum, Reichb. l. c. 19 = A. Napellus.
Kusnezoffii, *Reichb. Monog. Acon.* t. 21.—As. bor. or.
laciniosum, Schleich. ex Steud. Nom. ed. II. i. 18 = variegatum.
laetum, Reichb. Uebers. Acon. 21 = A. Napellus.
laeve, Royle, Illustr. 56 = A. Lycoctonum.
laevigatum, Schleich. ex Steud. Nom. ed. II. i. 18 = variegatum.
lagoctonum, Reichb. Uebers. Acon. 71 = A. Lycoctonum.
Lamarckii, Reichb. Monog. Acon. t. 40 = A. Lycoctonum.
lasiocarpum, Reichb. Uebers. Acon. 55 = variegatum.

ACONITUM:—
lasiostomum, *Reichb. Monog. Acon.* t. 49.—Ross.; Podol.
lasiostomum, Spreng. Syst. ii. 620 = orientale.
laxiflorum, Sehleich. Cat. (1821) 5 = A. Napellus.
laxum, Reichb. Illustr. Aeon. t. 13 verso = A. Napellus.
leptanthum, *Reichb. Monog. Acon.* t. 64—Rossia.
lethale, Griff. Notul. iv. 732 = palmatum.
leucanthemum, Wender. in Linnaea, v. (1830) Litt. 53 = variegatum.
leucanthum, *Reichb. Uebers. Acon.* 55.—Hab.?
Lobelianum, *Host, Fl. Austr.* ii. 74.—Europ.
Lubanskyi, Reichb. Monog. Acon. t. 20 = Fischeri.
Lubarscii, Heynh. Nom. i. 13 = praec.
luparia, Reichb. Uebers. Acon. 74 = A. Lyeoetonum.
lupicida, Reiehb. l. e. 70 = A. Lycoctonum.
luridum, *Hook. f. & Thoms. Fl. Ind.* i. 55.—Reg. Himal.
luridum, Salisb. Prod. 375 = variegatum.
Lycoctonum, Bess. Enum. Pl. Volh. 22 = orientale.
Lyeoctonum, *Linn. Sp. Pl.* 532.—Europ.; As. bor.
macranthum, Reiehb. Uebers. Acon. 51 = varicgatum.
macrophyllum, Hort. ex G. Don, in Loud. Hort. Brit. 223 = orientale.
macrorhynchum, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 83.—As. temp.
maximum, Pall. ex DC. Syst. i. 380; Prod. i. 61 = Fischeri.
medium, Schrad. Cat. Hort. Goett. ex Reiehb. Monog. Aeon. t. 71 in syn. = A. Cammarum.
meloctonum, Reichb. Uebers. Acon. 65 = A. Lyeoetonum.
Meyeri, Reichb. l. c. 33 = A. Napellus.
microphyllum, Gaud. ex Steud. Nom. ed. II. i. 19 = A. Napellus.
microstachyum, Reiehb. Uebers. Acon. 36 = A. Napellus.
Mielichhoferi, Reichb. l. e. 29 = A. Napellus.
mixtum, Reichb. l. c. 61 = variegatum.
moldavicum, Hacq. ex Reiehb. l. e. 67 = A. Lycoctonum.
molle, Reichb. l. c. 47 = paniculatum.
monanense, Schmidt, ex Reichb. Monog. Aeon. t. 51 in obs. = A. Lycoctonum.
monogynum, Forsk. Fl. Aegypt. Arab. p. xxvii = Delphinium Aconiti.
multifidum, Koeh, ex Reichb. Monog. Acon. t. 70 = A. Napellus.
multifidum, Royle, Illustr. 56 = A. Napellus.
myoctonum, Reichb. Uebers. Acon. 68 = A. Lycoetonum.
nanum, Stephan, ex Ledeb. Fl. Ross. i. 70 = biflorum.
napelloides, Sw. ex Reichb. Monog. Aeon. t. 69 in syn. = A. Napellus.
Napellus, *Linn. Sp. Pl.* 532.—Reg. bor. temp
Napellus, Pall. Reise, ii. 528 = villosum.
Napellus, S. G. Gmel. It. i. 8 = A. Lyeoctonum.
Napellus, Thunb. Fl. Jap. 231 = Fiseheri.
nasutum, Fisch. ex G. Don, Gen. Syst. i. 61 (1831) 58 = variegatum.
nasutum, Hook. Fl. Bor. Am. i. 26 = Fischeri.
neapolitanum, Tenore, Fl. Nap. iv. 327 = A. Lycoetonum.
nemorosum, Bieb. ex Reichb. Uebers. Acon. 17 = A. Anthora.
neomontanum, Baumg. Fl. Transs. ii. 100 = paniculatum.
neomontanum, Koelle, Spieil. 16 = A. Napellus.
neomontanum, Willd. Sp. Pl. ii. 1236 = A. Cammarum.
neubergense, DC. Syst. i. 373 = A. Napellus.
nitidum, Fisch. ex Steud. Nom. ed. II. i. 19 = barbatum.
nutans, Hort. Helmst. ex Steud. Nom. ed. II. i. 19 = A. Napellus.
Nuttallii, *Sweet, Hort. Brit.* ed. II. 584, nomen.—Am. bor.
obscurum, Reichb. ex S. Wats. Bibl. Ind. 2 = A. Napellus.
obtusifolium, *Host, Fl. Austr.* ii. 72.—Europ.
ochotense, Reichb. Monog. Acon. t. 18.—Sibir. or.
ochranthum, C. A. Mey. in Ledeb. Fl. Alt. ii. 285 = A. Lycoctonum.
ochroleucum, Hort. ex Steud. Nom. ed. II. i. 19 = A. Lycoctonum.
ochroleucum, Salisb. Prod. 375 = A. Anthora.
ochroleucum, Willd. Sp. Pl. ii. 1233 = orientale.
oliganthemum, Kern. in Ber. Naturw. Ver. Innsbruck, i. (1870) 119 = A. Napellus.
oligocarpum, Reichb. Uebers. Acon. 24 = A. Napellus.
oregonsense, *Rafin. New Fl. Am.* i. 57.—Oregon.
orientale, *Mill. Gard. Dict.* ed. VIII. n. 10.—Reg. Caucas.

ACONITUM:—
Ottonianum, Reichb. Uebers. Aeon. 54 = A. Cammarum.
ovatum, Lindl. in Bot. Reg. (1840) Misc. 53 = heterophyllum?
Pallasii, Reichb. Uebers. Acon. 18 = A. Anthora.
pallidum, Nutt. ex Loud. Hort. Brit. Suppl. 482, nomen = Nuttallii.
pallidum, Reiehb. Uebers. Aeon. 65 = A. Lyeoctonum.
palmatifidum, Reichb. l. c. 48 = A. Cammarum.
palmatum, *D. Don, Prod. Fl. Nep.* 196.—Reg. Himal.
panieulatum, *Lam. Fl. Fr.* iii. 646.—Europ.
paradoxum, Reichb. Monog. Acon. 76. t. 10. f. 3 = A. Napellus.
parviflorum, Reichb. Uebers. Acon. 46 = paniculatum.
parvifolium, *Host, Fl. Austr.* ii. 70.—Europ.
pauciflorum, Bertol. Misc. xxiv. p. 17. t. 6 = A. Napellus.
pauciflorum, Host, Fl. Austr. ii. 70 = A. Lycoctonum.
perniciosum, Reichb. Uebers. Acon. 68 = A. Lyeoetonum.
Phthora, Reichb. l. c. 71. = A. Lycoetonum.
plicatum, Koehler, ex Reiehb. l. e. 29 = A. Napellus.
productum, *Reichb. l. c.* t. 38 = A. Napellus.
pubescens, Moeneh, Meth. 310 = A. Napellus.
pubiceps, *Rupr. Fl. Caucas.* 42.—Reg. Caueas.
pumilum, Schur, Enum. Pl. Transs. 33 = A. Napellus.
purpureum, *Host, Fl. Austr.* ii. 74.—Europ.
pygmaeum, Steud. Nom. ed. I. 10 = A. Napellus.
pyramidale, Mill. Gard. Dict. ed. VIII. n. 6 = A. Napellus.
pyrenaicum, Hort. ex Steud. Nom. ed. II. i. 20 = leptanthum.
pyrenaicum, Koelle, Spicil. 24 = A. Lycoetonum.
pyrenaicum, Linn. Sp. Pl. 532 = A. Lyeoctonum.
pyrenaicum, Pall. Reise, ii. 316 = A. Anthora.
Raddeanum, *Regel, in Ind. Sem. Hort. Petrop.* (1861) 43; *et Fl. Ussur.* 11.—Mantehuria.
ranunculifolium, Reichb. Ie. Fl. Germ. iv. 22 = A. Lycoctonum.
ranunculoides, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 78.—Rossia.
reclinatum, A. Gray, in Hook. Lond. Journ. Bot. ii. (1843) 118 = A. Napellus.
recognitum, Reichb. Uebers. Acon. 42 = volubile.
rectum, Bernh. ex Reichb. l. c. 69 = A. Lycoctonum.
reflexum, Reichb. l. c. 43 = panieulatum.
rhynchanthum, Reichb. l. c. 56 = variegatum.
rigidum, Reichb. l. c. 36 = A. Napellus.
rostratum, Bernh. Ind. Sem. Hort. Erf. (1815); Reichb. Uebers. Acon. 56 = variegatum.
rotundifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 139 = A. Napellus.
rubellum, Sweet, Hort. Brit. ed. I. 9 = A. Napellus.
rubicundum, Fisch. ex Steud. Nom. ed. II. i. 20 = A. Lycoetonum.
sachalinense, *F. Schmidt, Reise Amurland.* 107.—Ins. Sachal.
scandens, Muhl. ex Reiehb. Uebers. Acon. 38 = uneinatum.
scandens, *Rafin. New Fl. Am.* i. 55.—Am. bor.
Schleicheri, Reiehb Uebers. Acon. 35 = A. Napellus.
Sczukini, *Turcz. in Bull. Soc. Nat. Mosc.* xiii. (1840) 61.—Sibir.
semigaleatum, Pall. ex Reichb. Uebers. Acon. 38 = A. Napellus.
septentrionale, Koelle, Spicil. 22 = A. Lyeoctonum.
sibiricum, Poir. Encyc. Suppl. i. 113 = barbatum.
sinense, Siebold, ex Lindl. & Paxt. Flow. Gard. i. 17 = Fischeri.
speciosum, Otto, ex DC. Prod. i. 60 = A. Cammarum.
spicatum, Donn, Hort. Cantab. ed. X. 209 = A. Napellus.
Sprengelii, Reichb. Uebers. Aeon. 35 = A. Cammarum.
squarrosum, Linn. ex DC. Syst. i. 368 = barbatum.
squarrosum, Koch, ex Reiehb. Uebers. Acon. 29 = A. Napellus.
Stoerkianum, Reichb. l. c. 49 = A. Cammarum.
strictissimum, Reichb. l. e. 69 = A. Lyeoctonum.
strictum, Bernh. ex DC. Syst. i. 373 = A. Napellus.
strictum, Willd. ex Reichb. Uebers. Acon. 51 = A. Lycoctonum.
taurericum, Reichb. Monog. Acon. 87. t. 12 = A. Napellus.
tauricum, Wulf. in Jacq. Coll. ii. 112 = A. Napellus.
Tauschianum, *Steud. Nom.* ed. II. i. 20.—Sibir.
tenuifolium, Reichb. Uebers. Aeon. 24 = A. Napellus.
tenuifolium, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 83 = macrorhynchum.

ACONITUM:—
tenuisectum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 65 = A. Lyeoetonum?
thelyphonum, Reichb. Uebers. Acon. 73 = A. Lycoctonum.
theriophonum, Reichb. l. c. 73 = A. Lyeoctonum.
tianschanicum, Rupr. Sert. Tianschan. 38 = A. Napellus.
tortuosum, *Willd. Enum. Hort. Berol.* 576.—Sibir.
toxicarium, Salisb. Prod. 375 = A. Lyeoctonum.
toxicum, Reichb. Uebers. Acon. 43 = paniculatum.
tragoctonum, Reiehb. l. c. 73 = A. Lycoetonum.
transilvanicum, Lereh. ex Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 165 = A. Lycoctonum.
Trattinickii, *Bess. ex Reichb. Uebers. Acon.* 75, nomen.—Hab.?
trichoearpum, *Reichb. l. c.* 29.—Hab.?
triste, Fisch. ex Steud. Nom. ed. II. i. 20 = A. Lycoctonum.
truncatum, *Rafin. New Fl. Am.* i. 55.—Am. bor.
tuberosum, *Host, Fl. Austr.* ii. 72.—Europ.
tuberosum, Patr. ex Reichb. Uebers. Acon. 16 = A. Anthora.
umbraticolum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 65 = A. Lycoctonum.
uncinatum, Hort. ex Steud. Nom. ed. II. i. 20 = variegatum.
uneinatum, *Linn. Sp. Pl.* ed. II. 750.—Am. bor.; Japon.
uncinatum, Sievers, ex Pall. Nord. Beitr. vii. 229 = biflorum.
variegatum, Hook. f. & Thoms. Fl. Ind. 56 = uncinatum.
variegatum, *Linn. Sp. Pl.* 532.—Europ.
venustum, Reichb. Uebers. Acon. 28 = A. Napellus.
versicolor, Reiehb. l. c. 50 = A. Cammarum.
versicolor, Ster. ex Ledeb. Fl. Ross. i. 66 = A. Anthora.
villosum, Maxim. Ind. Fl. Pek. in Prim. Fl. Amur. 468 = Kusnezoffi.
villosum, *Reichb. Uebers. Acon.* 39.—Sibir.
virgatum, Reichb. l. c. 28 = A. Napellus.
virosum, D. Don, Prod. Fl. Nep. 196 = A. Napellus.
volubile, Moench, Meth. Suppl. 110 = variegatum.
volubile, *Koelle, Spicil.* 21.—Sibir. altaic.
volubile, Muhl. Cat. ii. 54 = uneinatum.
vulgare, DC. Syst. i. 371 = A. Napellus.
vulparia, Reichb. Uebers. Acon. 70 = A. Lycoctonum.
Willdenowii, Reiehb. l. c. 35 = A. Napellus.
Wilemetianum, Delarb. Fl. Auv. ed. II. 499 = paniculatum.
Wulfenianum, Reichb. ex Steud. Nom. ed. II. i. 20 = A. Napellus.
zooctonum, Reichb. Uebers. Acon. 65 = A. Lycoctonum.

ACONOGONUM, Reiehb. Consp. 162 (1828) = **Polygonum**, Linn.
alpinum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 64 = P. alpinum.

ACONTIAS, Schott, Melet. i. 19 (1832) = **Xanthosoma**, Schott (Aroid.).
conspurcatus, Schott, Prod. 193 = X. conspurcatum.
cubensis, Schott, in Bonplandia, vii. (1859) 163 = X. cubense.
diversifolius, Regel, Ind. Sem. Hort. Petrop. (1865) 44; Gartenfl. (1866) 68 = X. platylobum.
hastifolius, Sehott, Melet. i. 19 = X. hastifolium.
helleborifolius, Sehott, l. c. = X. helleborifolium.
Hoffmanni, Schott, Prod. 196 = X. Hoffmanni.
luridus, G. Don, in Sweet, Hort. Brit. ed. III. 632 = Caladium luridum.
pentaphyllus, Schott, Syn. 65 = X. pentaphyllum.
platylobus, Schott, in Ann. Mus. Lugd. Bat. i. 279 = X. platylobum.
Plumieri, Schott, Syn. i. 65 = X. Plumieri.
Riedelianus, Schott, Prod. 192 = X. Riedelianum.
striatipes, Schott, l. c. = Caladium striatipes.
variegatus, Kunth, Enum. Pl. iii. 45 = X. helleborifolium.
Wendlandii, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 178 = X. Hoffmanni.

ACOPHORUM, Gaudich. ex Steud. Nom. ed. II. 20. (1840), nomen. *GRAMINEAE*?
caerulescens, *Gaudich, l. c.*—Hab.?

ACORIDIUM, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. I. (1843) 131 = **Ceratostylis**, Blume (Orchid.).
tenellum, Nees & Meyen, l. c. = C. gracilis.

ACORUS, Linn. Gen. ed. I. 104 (1737). *AROIDEAE*, Benth. & Hook. f. iii. 999.
americanus, Rafin. New Fl. Am. i. 57 = A. Calamus ?
angustifolius, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 284 = A. Calamus.
aromaticus, Gilib. Exercit. ii. 507 = A. Calamus.
Belangeri, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 284 = A. Calamus.
brasiliensium, Schott, ex Nees, in Mart. Fl. Bras. ii. 1. 170 = Scleria hirtella.
Calamus, *Linn. Sp. Pl.* 324.—Reg. bor. temp.
Calamus-aromaticus, [Clairv.] Man. Herb. 104 = A. Calamus.
Casia, Bertol. in Mem. Acad. Sci. Bolog. Ser. II. iv. (1864) 310 = A. Calamus.
cochinchinensis, *Schott, Melet.* i. 22.—Cochinch.
Commersonii, Schott, in Ann. Mus. Lugd. Bot. i. 284 = A. Calamus.
commutatus, Schott, Prod. 578 = A. Calamus.
elatus, Salisb. Prod. 263 = A. Calamus.
europaeus, Dum. Fl. Belg. 162 = A. Calamus.
flexuosus, Rafin. 'New Fl. Tex. 29'; Atl. Journ. 178 = A. Calamus.
floridanus, Rafin. New Fl. Am. i. 57 = A. Calamus.
gramineus, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 474. —Japon.
Griffithii, Schott, in Oestr. Bot. Zeitschr. (1858) 351 = A. Calamus
humilis, Salisb. Prod. 263 = A. gramineus.
nilaghirensis, Schott, in Oestr. Bot. Zeitschr. (1859) 101 = A. Calamus.
odoratus, Lam. Fl. Fr. iii. 299 = A. Calamus.
Palmita, Lichtenst. Reise, ii. 256 = Prionium Palmita.
pusillus, Siebold, in Verh. Bat. Genoot. xii. (1830) 2 = gramineus.
spurius, Schott, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 284 = A. Calamus.
Tatarinowii, Schott, in Oestr. Bot. Zeitschr. (1859) 101 = A. Calamus.
terrestris, Spreng. Syst. ii. 118 = A. Calamus.
triqueter, Turcz. ex Schott, Prod. 578 = A. Calamus.
undulatus, Stokes, Bot. Mat. Med. ii. 282 = A. Calamus.
verus, Rafin. Med. Fl. i. 26 = A. Calamus.

ACOSMIA, Benth. in Wall. Cat. n. 644 (1829) = **Gypsophila**, Linn. (Caryophyll.).
rupestris, Benth. l. c. = G. cerastioides.

ACOSMIUM, Schott, in Spreng. Syst. iv. Cur. Post. 406 (1827) = **Sweetia**, Spreng. (Legumin.).
lentiscifolium, Schott, l. c. = S. lentiscifolia.
lentiscifolium, Vog. in Linnaea, xi. (1837) 395 = S. fruticosa.

ACOSMUS, Desv. in Desf. Cat. Hort. Par. ed. III. 233 (1829) = **Aspicarpa**, Lag. (Malpig.).
pruriens, Desv. l. c. = A. hirtella.

ACOSTA, Adans. Fam. ii. 117 (1763) = **Centaurea**, Linn. (Compos.).

ACOSTA, DC. Prod. v. 90 (1836) = **Spiracantha**, H. B. & K. (Compos.).
Rolandrae, DC. l. c. = S. cornifolia.

ACOSTA, Lour. Fl. Cochinch. 276 (1790) = **Agapetes**, G. Don (Vaccin.).
spicata, Lour. l. c. = A. Acosta.

ACOSTA, Ruiz & Pav. Prod. 1 (1794) = **Moutabea**, Aubl. (Polygal.).
aculeata, Ruiz & Pav. Fl. Per. i. 5 = M. Acosta.

ACOUROA, Aubl. Pl. Gui. ii. 753. t. 301 (1775) = **Ecastaphyllum**, Rich. (Legumin.).
violacea, Aubl. l. c. (= *Geoffraea violacea*).

ACOURTIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 203 = **Perezia**, Lag. (Compos.).
formosa, DC. Prod. vii. 66 = P. fruticosa.
formosa, D. Don, in Trans. Linn. Soc. xvi. (1830) 203 = P. turbinata.
formosa, Hook. & Arn. Bot. Beech. Voy. 437 = P. rigida.
formosa, Sch. Bip. in Seem. Bot. Voy. Herald, t. 56 = P. patens.
hebeclada, DC. in Deless. Ic. Sel. iv. t. 95 = P. hebeclada.

ACOURTIA :—
macrocephala, Sch. Bip. in Seem. Bot. Voy. Herald, t. 55 = P. turbinata.
microcephala, DC. Prod. vii. 66 = P. microcephala.
moschata, DC. l. c. = P. moschata.
rigida, DC. l. c. = P. rigida.
Seemannii, Sch. Bip. in Seem. Bot. Voy. Herald, t. 54 = P. Seemannii.
turbinata, DC. Prod. vii. 66 = P. turbinata.

ACRACHNE, Wight & Arn. ex Lindl. Introd. Nat. Syst. ed. II. 381 (1836) = **Eleusine**, Gaert. (Gramin.).
eleusinoides, Wight & Arn. ex Steud. Nom. ed. II. i. 21 = E. verticillata.
verticillata, Wight & Arn.—Cf. Lindl. Introd. Nat. Syst. ed. II. 381 = E. verticillata.

ACRADENIA, Kipp. in Proc. Linn. Soc. ii. (1852) 201; et Trans. Linn. Soc. xxi. (1855) 207. t. 22. *RUTACEAE*, Benth. & Hook. f. i. 292.
Frankliniae, *Kipp. ll. cc.*—Tasmania.
zierioides, Kipp. in Proc. Linn. Soc. ii. (1852) 201 = Frankliniae.

ACRAEA, Lindl. in Benth. Pl. Hartw. 155 (1845) = **Pterichis**, Lindl. (Orchid.).
Mandonii, Reichb. f. Xenia Orch. iii. 18.—Boliv.
multiflora, Lindl. Orchid. Linden. 26.—N. Granat.
parvifolia, Lindl. in Benth. Pl. Hartw. 155.—Ecuador.
triloba, Lindl. in Ann. Nat. Hist. xv. (1845) 386.—Peruv.
Widgreni, Reichb. f. Otia Bot. Hamb. 83.—Bras.

ACRANDRA, Berg, in Linnaea, xxvii. (1854) 435; et in Mart. Fl. Bras. xiv. 1. 459 (1857) = **Campomanesia**, Ruiz & Pav. (Myrtac.).
laurifolia, Berg, in Mart. Fl. Bras. xiv. 1. 459 = C. laurifolia.
Sellowiana, Berg, l. c.—Bras.
verrucosa, Berg, l. c.—Bras.

ACRANTHERA, Arn. ex Meissn. Gen. 162 (115) (1838); et in Ann. Nat. Hist. iii. (1839) 20. *RUBIACEAE*, Benth. & Hook. f. ii. 64.
ANDROTROPIS, R. Br. in Wall. Cat. n. 8398 (1847 ?).
anamallica, *Bedd. Ic. Pl. Ind. Or.* i. 5.—Ind. or.
grandiflora, *Bedd. l. c.*—Ind. or.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 92.—Reg. Himal.
Maingayi, *Hook. f. l. c.*—Molucc.
tomentosa, *Hook. f. l. c.*—Reg. Himal.
uniflora, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 312 = Mussaenda uniflora.
zeylanica, *Arn. in Ann. Nat. Hist.* iii. (1839) 21.—Zeylan.

ACRATHERUM, Link, Hort. Berol. i. 230 (1827) = **Arundinella**, Raddi (Gramin.).
fuscatum, Nees, ex Steud. Syn. Pl. Gram. 114 = Arund. fuscata.
miliaceum, Link, Hort. Berol. i. 320 = Arund. nepalensis.
pumilum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 414 = Arund. pumila.

ACRIDOCARPUS, Guill. & Perr. Tent. Fl. Seneg. i. 123. t. 29 (1832). *MALPIGHIACEAE*, Benth. & Hook. f. i. 256.
ANOMALOPTERIS, G. Don, Gen. Syst. i. 647 (1831).
adenophorus, *A. Juss. in Arch. Mus. Par.* iii. (1843) 487.—Madag.
angolensis, *A. Juss. l. c.* 490.—Afr. trop.
argyrophyllus, *A. Juss. l. c.* 493.—Madag.
austro-caledonicus, *Baill. Adansonia*, xi. (1873-76) 248.—N. Caled.
Cavanillesii, A. Juss. in Arch. Mus. Par. iii. (1843) t. 15 = Smeathmanni.
chloropterus, *Oliver, Fl. Trop. Afr.* i. 279.—Afr. trop.
corymbosus, *Hook. f. in Hook. Ic. Pl.* t. 774.—Afr. trop.
excelsus, *A. Juss. in Arch. Mus. Par.* iii. (1843) 489. —Madag.
galphimiaefolius, *A. Juss. l. c.* 491.—Afr. trop.
guineensis, A. Juss. l. c. t. 15 = Smeathmanni.
Hirundo, *S. Moore, in Journ. Bot.* xviii. (1880) 1.—Afr. trop.
longifolius, Hook. f. in Hook. Niger Fl. 244 = Smeathmanni.
natalitius. *A. Juss. in Arch. Mus. Par.* iii. (1843) 486. —Natal.

ACRIDOCARPUS :—
orientalis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 488. —Natal.
plagiopterus, *Guill. & Perr. Tent. Fl. Seneg.* i. 123. t. 29.—Senegamb.
pruricus, *A. Juss. in Arch. Mus. Par.* iii. (1843) 492. —Natal.
Smeathmanni, *Guill. & Perr. Tent. Fl. Seneg.* i. 124. —Afr. trop.
zanzibaricus, *A. Juss. in Arch. Mus. Par.* iii. (1843) 485. t. 15.—Zanzibar.

ACRILIA, Griseb. Fl. Brit. W. Ind. 129 (1864) = **Trichilia**, Linn. (Meliac.).
Sloanei, Griseb. l. c. = T. terminalis.

ACRIOPSIS, Reinw. ex Blume, Cat. Gew. Buitenz. 97 (1823); Blume, Bijdr. 376 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 586.
densiflora, *Lindl. in Bot. Reg.* (1847) sub t. 20.—Borneo.
Griffithii, Reichb. in Bonplandia, ii. (1854) 92 = javanica.
indica, *Wight, Ic.* v. t. 1748.—Malaya.
javanica, *Reinw. ex Blume, Cat. Gew. Buitenz.* 97.; *Blume, Bijdr.* 377—Malaya.
picta, Lindl. in Bot. Reg. (1843) Misc. 69 = javanica.

ACRIULUS, Ridl. in Journ. Linn. Soc. xx. (1884) 336. *CYPERACEAE*.
griegifolius, *Ridl. l. c.*—Madag.
madagascariensis, *Ridl. l. c.*—Madag.

ACROANTHES, Rafin. in Journ. Phys. lxxxix. 261 (1819) = **Microstylis**, Lindl. (Orchid.).

ACROCARPIDIUM, Miq. in Versl. en Medel. Kon. Nederl. Inst. '1842' (1843) 198. t. 2 = **Peperomia**, Ruiz & Pav. (Piperac.).
brevipes, Benth. Bot. Voy. Sulph. 166 = P. brevipes.
cordifolium, Miq. Syst. Pip. 56 = P. cordifolia.
exile, Miq. l. c. = P. exilis.
guayaquillense, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 411 = P. guayaquilensis.
Guildingianum, Miq. l. c. 412 = P. Guildingiana.
hispidulum, Miq. Syst. Pip. 54 = P. hispidula.
Jamesonianum, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 412 = P. Jamesoniana.
majus, Miq. Syst. Pip. 60 = P. hederacea.
mexicanum, Miq. l. c. 58 = P. metapalcoensis.
molle, Miq. in Versl. en Medel. Kon. Nederl. Inst. '1842' (1843) 199 = P. mollis.
nummularifolium, Miq. l. c. 198. t. 2 = P. nummularifolia.
pulicare, Miq. l. c. 199 = P. scandens.
repens, Miq. l. c. = P. scandens.
rotundifolium, Miq. l. c. = P. circularis ?
scandens, Miq. Syst. Pip. 59 = P. scandens.
Sellowianum, Miq. l. c. 55 = P. hispidula.
serpens, Miq. in Versl. en. Medel. Kon. Nederl. Inst. '1842' (1843) 199 = Verhuellia serpens.
tenellum, Miq. l. c. = P. tenella.
urocarpum, Miq. Syst. Pip. 63 = P. urocarpa.
urospermum, Schlecht. in Linnaea, xxvii. (1854) 486 = praec. ?

ACROCARPUS, Nees, in Mart. Fl. Bras. ii. 1. 158. t. 16, 17 (1842) = **Cryptangium**, Schrad. (Cyperac.).
campestris, Nees, l. c. 160 = C. campestre.
densifolius, Nees, l. c. 159 = C. densifolium.
humilis, Nees, l. c. 161 = C. humile.
junciformis, Nees, l. c. = C. junciforme.
minarum, Nees, l. c. 159 = C. minarum.
polyphyllus, Nees, l. c. 160 = C. polyphyllum.
stellatus, Nees, ex Boeck. in Linnaea, xxxviii. (1874) 422 = C. stellatum.

ACROCARPUS, Wight, ex Arn. in Mag. Zool. & Bot. ii. (1839) 547. *LEGUMINOSAE*, Benth. & Hook. f. i. 568.
combretiflorus, *Teysm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1866) 258.—Malaya.
fraxinifolius, *Wight & Arn. in Mag. Zool. & Bot.* ii. (1839) 547.—Ind. or.
grandis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 87.—Sumatra.

ACROCENTRON, Cass. in Dict. Sc. Nat. xliv. 37. (1826) = **Centaurea**, Linn. (Compos.).
Boissieri, Webb, Iter Hisp. 32 = C. Boissieri.

ACROCENTRON:—
collinum, Cass in. Dict. Sc .Nat. xlvii. 37 = C. collina.
granatense, Webb, Iter Hisp. 32 = C. granatensis.
ornatum, Webb, l. c. = C. ornata.
tenuifolium, Cass. in Dict. Sc. Nat. l. 254 = C. rupestris.

ACROCEPHALUS, Benth. in Bot. Reg. sub tt. 1282, 1300 (1829). *LABIATAE*, Benth. & Hook. f. ii. 1173.
axillaris, *Benth. in DC. Prod.* xii. 48.—Ind. or.
Blumei, Benth. in Bot. Reg. sub t. 1300 = capitatus.
caeruleus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 135.—Afr. trop.
capitatus, *Benth. in Bot. Reg.* sub t. 1300.—Ind. or.; China.
cylindraceus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 135.—Afr. trop.
lilacinus, *Oliver, l. c.*—Afr. trop.
scariosus, Benth. in Bot. Reg. sub t. 1282 = capitatus.
verbenaceus, *Vatke, in Linnaea*, xl. (1876) 178.—Afr. trop.
villosus, *Benth. in Bot. Reg.* sub t. 1300; *Lab. Gen. et Sp.* 23.—Madag.

ACROCHAENE, Lindl. Fol. Orch. (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 507.
punctata, *Lindl. l. c.*—Ind. or.
Rimanni, *Reichb. f. in Gard. Chron.* N. S. xvii. (1882) 1. 796.—As. trop.

ACROCLINIUM, A. Gray, in Hook. Kew Journ. iv. (1852) 270 = **Helipterum**, DC. (Compos.).
multicaule, A. Gray, l. c. 271 = H. chlorocephalum.
phyllocephalum, A. Gray, l. c. = Podotheca fuscescens.
roseum, Hook. Bot. Mag. t. 4801 = H. roseum.
rubellum, A. Gray, in Hook. Kew Journ. iv. (1852) 71 = H. rubellum.

ACROCOMIA, Mart. Hist. Nat. Palm. ii. 66. tt. 56, 57 (1823). *PALMAE*, Benth. & Hook. f. iii. 943.
aculeata, Lodd. ex Mart. l. c. iii. 286 = sclerocarpa.
cubensis, Lodd. ex H. Wendl. Ind. Cycl. Pand. Cycad. 1 = lasiospatha.
fusiformis, Sweet, Hort. Brit. ed. I. 432 = lasiospatha?
glaucophylla, *Drude, in Mart. Fl. Bras.* iii. II. 392.—Bras.
globosa, *Lodd. ex Mart. Hist. Nat. Palm.* iii. 286.—Ins. S. Vincent.
guianensis, *Lodd. ex G. Don, in Loud. Hort. Brit.* 382, nomen.—Guiana.
horrida, *Lodd. ex Mart. Hist. Nat. Palm.* iii. 286.—Ins. Trinit.
intumescens, *Drude, in Mart. Fl. Bras.* iii. II. 391.—Bras.
lasiospatha, *Mart. in Palm. Orbign.* 81.—Ind. occ.
lasiospatha, Wallace, Palms Amaz. 97. t. 37 = sclerocarpa.
mexicana, *Karw. ex Mart. Hist. Nat. Palm.* iii. 285. t. 138.—Mexico.
minor, Lodd. ex G. Don, in Loud. Hort. Brit. 382 = lasiospatha?
sclerocarpa, *Mart. Hist. Nat. Palm.* ii. 66. tt. 56, 57, 100. f. 5.—Am. trop.
sphaerocarpa, Desf. Cat. Hort. Par. ed. III. 30 = sclerocarpa.
tenuifrons, *Lodd. ex Mart. Hist. Nat. Palm.* iii. 286.—Ins. S. Domingo.
Totai, *Mart. in Palm. Orbign.* 78. t. 9. f. 1.—Boliv.
vinifera, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 47.—Nicaragua.
Zapotecis, Karw. ex H. Wendl. in Kerch. Palm. 230 = mexicana.

ACROCORION, Adans. Fam. ii. 57 (1763) = **Galanthus**, Linn. (Amaryll.).

ACROCORYNE, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 316 = **Metastelma**, R. Br. (Asclep.).
caribaea, Turcz. l. c. = M. bahamense.

ACRODICLIDIUM, Nees, Laurin. Disp. Progr. 13 (1833) *LAURINEAE*, Benth. & Hook. f. iii. 154.
TRIPLOMEIA, Rafin. Sylva Tellur. 134 (1838).
anacardioides, *Spruce, ex Meissn. in DC. Prod.* xv. 1. 86.—Bras.
brasiliense, Nees, Syst. Laurin. 267 = Aydendron brasiliense.
Camara, *Schomb. in Hook. Lond. Journ. Bot.* iii. (1844) 630.—Guiana.

ACRODICLIDIUM:—
chrysophyllum, *Meissn. in DC. Prod.* xv. 1. 87.—Guiana.
dominicense, *Meissn. l. c.* 86.—Ins. Dominic.
Gardneri, *Meissn. in DC. Prod.* xv. 1. 87.—Bras.
guianense, *Nees, Syst. Laurin.* 268.—Guiana.
Ita-uba, *Meissn. in DC. Prod.* xv. 1. 86.—Bras.
jamaicense, Griseb. Fl. Brit. W. Ind. i. 280, ex parte = Wrightii.
jamaicense, *Nees, Syst. Laurin.* 270.—Jamaic.
Kunthianum, Nees, l. c. 269 = Aydendron aciphyllum.
oppositifolium, *Nees, in Linnaea*, xxi. (1868) 500.—Guiana.
salicifolium, *Griseb. Fl. Brit. W. Ind.* i. 280.—Ind. occ.
sericeum, *Griseb. l. c.* 280.—Ins. Dominic.
Sprucei, *Meissn. in DC. Prod.* xv. 1. 86.—Bras.
Wrightii, *Meissn. l. c.* 85.—Cuba.

ACRODRYON, Spreng. Syst. i. 365 (1825) = **Cephalanthus**, Linn. (Rubiac.).
angustifolius, Spreng. l. c. 336 = C. angustifolius.
orientalis, Spreng. l. c. 336 = C. orientalis.

ACROELYTRUM, Steud. in Flora, xxix. (1846) 20 = **Lophatherum**, Brongn. (Gramin.).
japonicum, Steud. l. c. 21 = L. elatum.

ACROGLOCHIN, Schrad. Cat. Hort. Goett. ex Schult. f. Mant. i. 69, 227 (1822). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 48.
BLITANTHUS, Reichb. Cat. Hort. Dresd. (1824).
LECANOCARPUS, Nees, Pl. Hort. Bonn. 4. t. 2. (1823).
chenopodioides, *Schrad. ex Schult. f. Mant.* i. 227.—Reg. Himal.; China.
persicarioides, Moq. in DC. Prod. xiii. II. 254 = chenopodioides.
Schraderiana, Schult. f. ex Moq. l. c. = chenopodioides.

ACROGLYPHE, E. Mey. Zwei Pfl. Docum. 100 (1843). = **Annesorhiza**, Cham. & Schlecht. (Umbell.).
flexuosa, E. Mey. l. c. 100 = Ann. hirsuta + montana.
hispida, Sond. in Harv. & Sond. Fl. Cap. ii. 546 = Ann. villosa.
hispidula, E. Mey. Zwei Pfl. Docum. 74 = Ann. villosa.
runcinata, E. Mey. l. c. 100.—Afr. austr.

ACROLEPIS, Schrad. in Comm. Gotting. vii. (1832) 142 = **Ficinia**, Schrad. (Cyperac.).
ferruginea, Nees, ex Boeck. in Linnaea, xxxvii. (1871–73) 91.—Afr. austr.
ramosissima, Boeck. l. c. 89 = F. ramosissima.
trichodes, Schrad. in Comm. Gotting. vii. (1832) 142 = F. trichodes.

ACROLOBUS, Klotzsch, in Verh. Akad. Berl. (1856) 236. t. 3 = **Heisteria**, Linn. (Olacin.).
parvifolia, Klotzsch, l. c. 237. t. 3 = H. parvifolia.
Schoenleinii, Klotzsch, l. c. 236. t. 3 = H. parvifolia.

ACROLASIA, Presl, Rel. Haenk. ii. 39. t. 55. (1831) = **Mentzelia**, Linn. (Loasac.).
bartonioides, Presl, l. c. 39. = M. albicaulis.
Solierii, C. Gay, Fl. Chil. ii. 430.—Chili.
squalida, Hook. f. in Trans. Linn. Soc. xx. (1851) 222 = M. aspera?

ACROLOPHUS, Cass. in Dict. Sc. Nat. l. (1827) 253 = **Centaurea**, Linn. (Compos.).
maculosus, Cass. l. c. 253 = C. maculosa.
Mierghii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 97 = C. paniculata.
paniculatus, Cass. in Dict. Sc. Nat. l. 253 = C. paniculata.
ragusinus, Webb, ex Steud. Nom. ed. II. i. 23 = C. ragusina.
rigidulus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 97 = C. paniculata.
subalbidus, Fourr. l. c. = C. maculosa.
tenuisectus, Fourr. l. c. = C. maculosa.

ACRONEMA, Falcon. ex Edgew. in Trans. Linn. Soc. xx. (1851) 51 = **Pimpinella**, Linn. (Umbellif.).
tenerum, Edgew. l. c. = P. tenera.

ACRONIA, Presl, Rel. Haenk. i. 103 (1827); Symb. Bot. ii. 9 (1858) = **Cryptostylis**, R. Br. (Orchid.).
phalangifera, Presl, ll. cc. 104. 10.—Peruv.

ACRONODA, Hassk. Cat. Hort. Bogor. Alt. 208 (1844) = seq.

ACRONODIA, Blume, Bijdr. (1825) 123 = **Elaeocarpus**, Linn. (Tiliac.).
punctata, Blume, l. c. = E. Acronodia.

ACRONOZUS, Steud. Nom. ed. II. i. 21 (1840) = praec.

ACRONYCHIA, Forst. Char. Gen. 53. t. 27 (1776). *RUTACEAE*, Benth. & Hook. f. i. 302.
CUNTO, Adans. Fam. ii. 446 (1763).
CYMINOSMA, Gaert. Fruct. i. 280. t. 58 (1788).
DORRIENIA, Dennst. Schluess. Hort. Malab. 31 (1818).
GELA, Lour. Fl. Cochinch. 232 (1790).
HUONIA, Montrous. in Mém. Acad. Lyon, x. (1860) 185.
JAMBOLANA, Adans. Fam. ii. 508 (1763).
JAMBOLIFERA, Linn. Diss. Dass. 15 (1747); Amoen. Acad. i. 398 (1749).
KOELPINIA, Scop. Introd. 231, et Ind. (1777).
SELAS, Spreng. Syst. ii. 216 (1825).
acidula, *F. Muell. Fragm.* iv. 154.—Austral.
apiculata, *Miq. Fl. Ind. Bat. Suppl.* 532.—Sumatra.
arborea, *Blume, Bijdr.* 244.—Java.
Baueri, *Schott, Fragm. Bot.* 5. t. 3.—Austral.
Cunninghamii, Hook. Bot. Mag. t. 3994 = Medicosma Cunninghamii.
Endlicheri, *Schott, Fragm. Bot.* 3. t. 2.—Ins. Norfolc.
Halmaheirae, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 245.—Malaya.
heterophylla, *A. Gray, Bot. U. S. Expl. Exped.* i. 333. t. 32.—Ins. Samoa.
Hillii, F. Muell. Fragm. i. 26 = Baueri.
imperforata, *F. Muell. l. c.*—Austral.
laevis, *Forst. Char. Gen.* 54. t. 27.—Austral.
laurifolia, *Blume, Cat. Gew. Buitenz.* 27. 63; *Bijdr.* 245.—As. trop.
laurina, F. Muell. 3rd Rep. Veg. Vict. 7; Fragm. i. 27 = laevis.
melicopoides, *F. Muell. Fragm.* v. 3.—Austral.
Minahassae, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 245.—Ins. Celebes.
oblongifolia, Endl. ex Heynh. Nom. ii. 8 = laevis.
ovata, Endl. ex Heynh. l. c. = Baueri.
pedunculata, Miq. Fl. Ind. Bat. Suppl. 532 = laurifolia.
petiolaris, *A. Gray, Bot. U. S. Expl. Exped.* i. 335. t. 33.—Ins. Fiji.
Porteri, *Hook. f. Fl. Brit. Ind.* i. 498.—Malaya.
retusa, *A. Gray, Bot. U. S. Expl. Exped.* i. 338. t. 34.—Ins. Samoa.
Richii, *A. Gray, l. c.* 336. t. 33.—Ins. Samoa.
simplicifolia, Steud. Nom. ed. II. i. 21 = Endlicheri.
tetrandra, *F. Muell. Fragm.* ix. 104.—Austral.
trifoliolata, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Ind.* ii. (1845) 585.—Java.
vestita, *F. Muell. Fragm.* iv. 155.—Austral.

ACROPERA, Lindl. Gen. et Sp. Orch. 172 (1833) = **Gongora**, Ruiz & Pav. (Orchid.).
armeniaca, Lindl. in Lindl. & Paxt. Flow. Gard. i. (1850–51) 94 = G. armeniaca.
atropurpurea, Hort. ex Heynh. Nom. ii. 8 = G. galeata.
Batemanni, Lindl. ex Reichb. f. in Gard. Chron. N. S. ii. (1874) 1. 322 = G. galeata.
citrina, Hort. ex Reichb. f. Xenia Orch. i. 51 = G. galeata.
cornuta, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 186 = G. armeniaca.
flavida, Klotzsch. l. c. xix. (1851) 217 = G. galeata.
fuscata, Hort. Belg. ex Heynh. Nom. ii. 8.—Mexico.
grandiflora, Hort. ex Pasq. Cat. Orto Bot. Napol. 3. nomen.—Hab.?
Loddigesii, Lindl. Gen. et Sp. Orch. 172 = G. galeata.
luteola, Hort. ex Heynh. Nom. ii. 8 = *fuscata*.
pallida, Hort. ex Heynh. l. c. = G. galeata.
purpurea, Hort. ex Heynh. l. c. = G. galeata.
sulphurea, Hort. ex Heynh. l. c. = *fuscata*.

ACROPETALUM, A. Juss. in Orb. Dict. vii. 617 (1849) = XEROPETALUM, Delile = **Dombeya**, Cav. (Stercul.).

ACROPHYLLUM, Benth. in Maund, Bot. ii. t. 95 (1840?). *SAXIFRAGEAE*, Benth. & Hook. f. i. 652.
CALYCOMIS, D. Don, in Edinb. N. Phil. Journ. ix. (July–Oct. 1830) 98.

ACROPHYLLUM :—
venosum, *Benth. in Maund, Bot.* ii. t. 95.—Austral.
verticillatum, Hook. Bot. Mag. t. 4050 = venosum.

ACROPHYLLUM, E. Mey. Zwei Pfl. Docum. 140 (1843) = **Kiggelaria**, DC. (Bixin.).
obtusatum, C. Muell. in Walp. Ann. vii. 233 = K. africana.
obtusum, E. Mey. Zwei Pfl. Docum. 140 = K. africana.

ACROPODIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 408 = **Aspalathus**, Linn. (Legumin.).
suffruticosum. Desv. l. c. = Asp. suffruticosa.

ACROPSELION, Spach, Hist. Vég. Phan. xiii. 163 (1846) = ACROSPELION, Bess. = **Trisetum**, Pers. (Gramin.).

ACROPTILION, Endl. Ench. 247 (1841) = seq.

ACROPTILON, Cass. in Dict. Sc. Nat. l. 464 (1827) = **Centaurea**, Linn. (Compos.).
angustifolium, Cass. l. c. 466 = C. Picris.
obtusifolium, Cass. l. c. 465 = C. Picris.
Picris, DC. Prod. vi. 662 = C. Picris.
repens, DC. l. c. 663 = C. Picris.
serratum, Cass. in Dict. Sc. Nat. l. 466 = C. Picris.
subdentatum, Cass. l. c. 465 = C. Picris.

ACROSANTHES, Eckl. & Zeyh. Enum. 328 (1837). *FICOIDEAE*, Benth. & Hook. f. i. 855.
DIDASTE, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 472 (1862).
anceps, *Sond. in Harv. & Sond. l. c.*—Afr. austr.
angustifolia, *Eckl. & Zeyh. Enum.* 329.—Afr. austr.
decandra, *Fenzl, in Ann. Wien. Mus.* ii. (1838) 270. —Afr. austr.
fistulosa, *Eckl. & Zeyh. Enum.* 328.—Afr. austr.
humifusa, *Sond. in Harv. & Sond. Fl. Cap.* ii. 472.—Afr. austr.
teretifolia, *Eckl. & Zeyh. Enum.* 329.—Afr. austr.

ACROSPELION, Bess. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 59 = **Trisetum**, Pers. (Gramin.).

ACROSPIRA, Welw. ex Baker, in Trans. Linn. Soc. Bot. Ser. II. i. (1878) 255. t. 34. *LILIACEAE*, Benth. & Hook. f. iii. 787.
asphodeloides, *Welw. l. c.*—Afr. trop.

ACROSSANTHES, Presl, Bot. Bemerk. 22 (1844) = **Vismia**, Vell. (Hypercin.).
Lhotskyanus, Presl, l. c. = V. decipiens.

ACROSTEMON, Klotzsch, in Linnaea, xii. (1838) 227 = **Grisebachia**, Klotzsch (Ericac.).
equisetoides, Klotzsch, l. c. 228.—Afr. austr.
glandulosus, Rach, l. c. xxvi. (1853–55) 790.—Afr. austr.
hirsutus, Klotzsch, l. c. xii. (1838) 228.—Afr. austr.
incanus, Klotzsch, l. c. 228 = G. incana.
incurvus, Benth. in DC. Prod. vii. 702.—Afr. austr.

ACROTAPHROS, Steud. ex A. Rich. Tent. Fl. Abyss. i. 207. t. 38 [1847] = **Ormocarpum**, Beauv. (Legum.).
bibracteata, Steud. ex A. Rich. l. c. = O. bibracteatum.

ACROTICHE, Poir. in Dict. Sc. Nat. i. Suppl. 50 (1816) = **Acrotriche**, R. Br. (Epacrid.).

ACROTOME, Benth. in Endl. Gen. 627 (1838). *LABIATAE*, Benth. & Hook. f. ii. 1206.
hispida, *Benth. in DC. Prod.* xii. 436.—Afr. austr.
inflata, *Benth. l. c.*—Afr. austr.
pallescens, *Benth. l. c.*—Afr. austr.

ACROTREMA, Jack, in Malay Misc. i. (1820) Art. V. p. 36. *DILLENIACEAE*, Benth. & Hook. f. i. 13.
Arnottianum, *Wight, Illustr. Ind. Bot.* i. 9. t. 3.—Ind. or.
bullatum, Thw. Enum. Pl. Zeyl. 20 = uniflorum.
costatum, *Jack, in Malay Misc.* i. (1820) Art. V. p. 36. —Ind. or.; Malaya.
costatum, Wall. Cat. n. 1117B = Arnottianum.
dissectum, *Thw. in Hook. Kew Journ.* viii. (1856) 242. —Zeylan.

ACROTREMA :—
Gardneri, *Thw. Enum. Pl. Zeyl.* 3.—Zeylan.
intermedium, *Thw. l. c.*—Zeylan.
lanceolatum, *Hook. Ic. Pl.* sub t. 157.—Zeylan.
lyratum, *Thw. in Hook. Kew Journ.* viii. (1856) 242. —Zeylan.
sylvaticum, *Thw. Enum. Pl. Zeyl.* 2 = uniflorum.
Thwaitesii, *Hook. f. & Thoms. in Hook. Kew Journ.* viii. (1856) 241.—Zeylan.
uniflorum, *Hook. Ic. Pl.* t. 157.—Zeylan.
Walkeri, *Wight, ex Thw. Enum. Pl. Zeyl.* 3.—Zeylan.
Wightianum, Wall. Cat. n. 3669 = costatum.

ACROTRICHE, R. Br. Prod. 547 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 614.
FROEBELIA, Regel, Gartenfl. i. (1852) 164. t. 18.
affinis, DC. Prod. vii. 757 = serrulata.
aggregata, *R. Br. Prod.* 547.—Austral.
aristata, Benth. in Enum. Pl. Hueg. 76 = Leucopogon cuspidatus.
cordata, R. Br. Prod. 548 = ovalifolia.
depressa, *R. Br. l. c.*—Austral.
divaricata, *R. Br. l. c.* 547.—Austral.
fasciculiflora, *Benth. Fl. Austral.* iv. 229.—Austral.
latifolia, A. Cunn. ex DC. Prod. vii. 757 = Leucopogon cordifolius.
Manglesii, Sond. in Lehm. Pl. Preiss. i. 326 = ramiflora.
ovalifolia, *R. Br. Prod.* 548.—Austral.
patula, Hook. f. Fl. Tasm. i. 253 = serrulata.
patula, *R. Br. Prod.* 547.—Austral.
prostrata, F. Muell. in Trans. Vict. Inst. (1855) 40 = serrulata.
ramiflora, *R. Br. Prod.* 547.—Austral.
ramiflora, Sond. in Linnaea, xxvi. (1853–55), 251 = fasciculiflora.
serrulata, *R. Br. Prod.* 547.—Austral.
subcordata, DC. Prod. vii. 757 = ovalifolia.

ACROXIS, Trin. ex Steud. Nom. ed. II. i. 22 (1840) = **Muehlenbergia**, Schreb. (Gramin.).

ACROZUS, Spreng. Syst. iv. Cur. Post. 145, 149 (1827) = **Elaeocarpus**, Linn. (Tiliac.).
punctatus, Spreng. l. c. = E. Acronodia.

ACRYPHYLLUM, Lindl. Introd. Nat. Syst. ed. II. 156 (1836) err. typ. = ARCYPHYLLUM, Ell. = **Tephrosia**, Lour. (Legum.).

ACTAEA, [Tourn.] Linn. Syst. ed. I (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 9.
CHRISTOPHORIANA, Tourn. ex Moench, Meth. 276 (1794).
acuminata, Wall. Cat. n. 4726 = spicata.
alba, *Mill, Gard. Dict.* ed. VIII. n. 2.—Am. bor.
americana, Pursh, Fl. Am. Sept. 366 = alba.
arguta, Nutt. ex Torr. & Gray, Fl. N. Am. i. 35 = spicata.
aspera, Lour. Fl. Cochinch. 332. = Tetracera sarmentosa.
brachypetala, var. *α*, DC. Syst. i. 385 = alba.
brachypetala, var. *β*, DC. l. c. = spicata.
Christophoriana, Gouan, Fl. Monsp. 152 = spicata.
Cimicifuga, Hook. Fl. Bor. Am. i. 26 = Cimicifuga elata.
Cimicifuga, Linn. Sp. Pl. 504 = Cimicifuga foetida.
cordifolia, DC. Syst. i. 383 = Cimicifuga cordifolia.
corymbosa, Stokes, Bot. Mat. Med. iii. 176 = spicata.
dahurica, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 = Cimicifuga foetida.
davurica, Franch. Pl. David. 23 = Cimicifuga davurica.
dioica, *Walt. Fl. Carol.* 152.—Carolina.
erythrocarpa, Fisch. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 20 (1835) = spicata.
frigida, Wall. Cat. n. 4725 = Cimicifuga foetida.
grandis, D. Dietr. Syn. Pl. iii. 233 = Trautvetteria palmata.
gyrostachya, Wender. Ind. Sem. Hort. Marb. (1840).—Cf. Linnaea, xv. (1841), Litt. 100 = Trautvetteria palmata.
japonica, Thunb. Fl. Jap. 221 = Cimicifuga japonica.
longipes, Spach, Hist. Vég. Phan. vii. 388 = spicata.
macrophylla, Baill. in Bull. Soc. Linn. Par. i. (1879) 224 = Anemopsis macrophylla.
macropoda, Turcz. ex Fisch. & Mey. fide Ledeb. Fl. Ross. i. 73 = Cimicifuga foetida.
monogyna, Walt. Fl. Carol. 151 = Cimicifuga racemosa.
nigra, Gaert. Mey. & Scherb. Fl. Wett. ii. 265 = spicata.

ACTAEA :—
orthostachya, Wender. Ind. Sem. Hort. Marb. (1840). —Cf. Linnaea, xv. (1841) Litt. 100 = Cimicifuga racemosa.
pachypoda, Ell. Sketch, ii. 15 = alba.
palmata, DC. Syst. i. 383 = Trautvetteria palmata.
pentacarpa, Michx. ex Torr. & Gray, Fl. N. Am. i. 39 = Cimicifuga americana.
pentagyna, Walt. Fl. Carol. 151 = Cimicifuga americana.
podocarpa, DC. Syst. i. 282 = Cimicifuga americana.
podocarpa, Schlecht. in Linnaea, vi. (1831) 583 = Cimicifuga foetida.
pterosperma, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 = Cimicifuga foetida.
racemosa, Geners. ex Steud. Nom. ed. II. i. 22 = Cimicifuga foetida.
racemosa, Gilib. Fl. Lituan. ii. 226 = spicata.
racemosa, Linn. Sp. Pl. 504 = Cimicifuga racemosa.
racemosa, Walt. ex Steud. Nom. ed. II. i. 22 = Trautvetteria palmata.
rubra, Bigel. Fl. Bost. ed. II. 211 = alba?
rubra, Willd. Enum. Hort. Berol. 561; Ledeb. Fl. Alt. ii. 275 = spicata.
simplex, Wormsk. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 = Cimicifuga foetida.
spicata, *Linn. Sp. Pl.* 504.—Reg. bor. temp.

ACTAEOGETON, Steud. Nom. ed. II. i. 22 (1840) = **Scirpus**, Linn. (Cyperac.).

ACTEGETON, Blume, Bijdr. 1143 (1826) = **Azima**, Lam. (Salvador).
sarmentosum, Blume, l. c. 1144 = A. sarmentosa.

ACTEGITON, Endl. Gen. 1089 (1840), err. typ. = praec.

ACTEPHILA, Blume, Bijdr. 581 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 269.
ANOMOSPERMUM, Dalz. in Hook. Kew Journ. iii. (1851) 228.
LITHOXYLON, Endl. Gen. 1122 (1840).
bantamensis, Miq. Fl. Ind. Bat. i. II. 356 = excelsa.
excelsa, *Muell. Arg. in Linnaea*, xxxii. (1863) 78.—Ind. or.; Malaya.
grandifolia, *Baill. Adansonia*, vi. (1865–66) 330.—Austral.
javanica, Miq. Fl. Ind. Bat. i. II. 356 = excelsa.
latifolia, *Benth. Fl. Austral.* vi. 89.—Austral.
major, *Muell. Arg. in Linnaea*, xxxii. (1863) 77.—Java.
Mooreana, *Baill. Adansonia*, vi. (1865–66) 330.—Austral.
neilgherrensis, Wight, Ic. t. 1910 = excelsa.
petiolaris, *Benth. Fl. Austral.* vi. 89.—Austral.
puberula, *Kurz, in Journ. As. Soc Beng.* xlii. (1873) II. 236.—Ins. Andaman.
rectinervis, *Kurz, in Journ. Bot.* xiii. (1875) 329.—Ins. Nicobar.
sessilifolia, *Benth. Fl. Austral.* vi. 90.—Austral.
Thomsoni, Muell. Arg. in Linnaea, xxxiv. (1863) 65 = excelsa.
zeylanica, Muell. Arg. l. c. 77 = excelsa.

ACTIMERIS, Rafin. in Am. Monthly Mag. (1819) 195 = **Actinomeris**, Nutt. (Compos.).

ACTINANTHUS, Ehrenb. in Linnaea, iv. (1829) 398. *UMBELLIFERAE*, Benth. & Hook. f. i. 903.
syriacus, *Ehrenb. l. c.*—As. min.; Syria.

ACTINEA, Juss. in Ann. Mus. Par. ii. (1803) 425. t. 61 = **Cephalophora**, Cav. (Compos.).
acaulis, Spreng. Syst. iii. 574 = Actinella acaulis.
alternifolia, Spreng. l. c. = C. heterophylla.
biternata, Spreng. l. c. = Coreopsis biternata.
chrysanthemoides, H. B. & K. Nov. Gen. et Sp. iv. 298. t. 411 = Hymenoxis chrysanthemoides.
commutata, Spreng. Syst. iii. 574 = Calea pinnatifida.
heterophylla, Juss. in Ann. Mus. Par. ii. (1803) 425 = C. heterophylla.
integrifolia, H. B. & K. Nov. Gen. et Sp. iv. 297. t. 410 = Helenium integrifolium.
integrifolia, *Torr. in Ann. Lyc. New York*, ii. (1828) 213.—Am. bor.
lanata, Steud. Nom. ed. I. 15 = Bahia lanata.
oppositifolia, Spreng. Syst. iii. 574 = Blainvillea bahiensis.

ACTINELLA, Nutt. Gen. Am. ii. 173 (1818). *COMPOSITAE*, Benth. & Hook. f. ii. 414.
PHILEOZERA, Buckl. in Proc. Ac. Sc. Philad. '1861.' (1862) 459.
PICRADENIA, Hook. Fl. Bor. Am. i. 317. t. 108 (1833).
acaulis, *Nutt. Gen. Am.* ii. 173.—Am. bor.
argentea, *A. Gray, in Mem. Am. Ac.* iv. (1849) (*Pl. Fendl.*) 100.—N. Mexico.
biennis, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 373.—Am. bor. occ.
Bigelovii, *A. Gray, Pl. Wright.* ii. 96, in adnot.—Mexico.
Brandegei, *T. C. Porter, ex A. Gray, in Proc. Am. Acad.* xiii. (1878) 373.—Am. bor. occ.
chrysanthemoides, *A. Gray, l. c.* xix. (1883) 32.—Mexico.
Cooperi, *A. Gray, l. c.* vii. (1868) 359.—Calif.
depressa, *Harv. & Gray, ex Gray, in Mem. Am. Acad.* iv. (1849) (*Pl. Fendl.*) 100.—Am. bor. occ.
glabra, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 379 = acaulis.
grandiflora, *Torr. & Gray, in Bost. Journ. Nat. Hist.* v. (1847) 110.—Am. bor. occ.
insignis, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xviii. (1882-83) 109.—Am. bor. occ.
lanata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 379 = acaulis.
lanata, Pursh, Fl. Am. Sept. ii. 560 = Eriophyllum caespitosum.
lanuginosa, Buckl. in Proc. Ac. Sc. Philad. (1862) 459 = scaposa.
leptoclada, *A. Gray, in Pacif. Rail. Rep.* iv. 107.—Am. bor. occ.
linearifolia, *Torr. & Gray, Fl. N. Am.* ii. 383.—Texas.
odorata, *A. Gray, in Mem. Am. Acad.* iv. (1849) (*Pl. Fendl.*) 101.—Am. bor. occ.
Palmeri, *A. Gray, in Proc. Am. Acad.* xix. (1883) 31.—Mexico.
Richardsonii, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 379 = biennis.
Rusbyi, *A. Gray, in Proc. Am. Acad.* xix. (1883) 33.—N. Mexico.
scaposa, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 379.—Am. bor. occ
Torreyana, Nutt. l. c. = acaulis.
Vaseyi, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 219.—Am. bor. occ.

ACTINELLA, Pers. Syn. ii. 469 (1807) = **Cephalophora**, Cav. (Compos.).
heterophylla, Pers. l. c. = C. heterophylla.

ACTINIDIA, Lindl. Introd. Nat. Syst. ed. II. 439 (1836). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 184.
KALOMIKTA, Regel, in Bull. Phys. Math. Acad. Pétersb. xv. (1857) 219.
TROCHOSTIGMA, Sieb. & Zucc. in Abh. Akad. Wiss. Muench. iii. (1843) 726. t. 2. f. 2.
acuminata, *Budisch. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 167.—Manchur.
arguta, Franch. & Sav. Enum. Pl. Jap. i. 58 = callosa.
callosa, *Lindl. Introd. Nat. Syst.* ed. II. 439.—Reg. Himal.; Japon.; Manchur.
Championi, *Benth. Fl. Hongk.* 26.—China.
chinensis, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 303.—China.
cordifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 15.—Japon.
Davidii, *Franch. Pl. Davidian.* i. 57.—China.
eriantha, *Benth. in Journ. Linn. Soc.* v. (1861) 55.—Hab.?
fulvicoma, *Hance, in Journ. Bot.* xxiii. (1885) 321.—China.
Kolomikta, Maxim. Prim. Fl. Amur. 63 = callosa.
platyphylla, *A. Gray, in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 15.—Japon.
polygama, *Franch. & Sav. Enum. Pl. Jap.* i. 59.—Japon.
rufa, Franch. & Sav. l. c. 58 = callosa.
strigosa, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 55.—Reg. Himal.
suifunensis, Budisch. ex Trautv. in Act. Hort. Petrop. viii. (1883) 167 = acuminata.
volubilis, *Franch. & Sav. Enum. Fl. Jap.* i. 59.—Japon.

ACTINOBOLE, Endl. Gen. Suppl. iii. 70 (1843) = **Gnaphalodes**, A. Gray (Compos.).

ACTINOCARPUS, R. Br. Prod. 342 (1810) = **Damasonium**, Juss. (Alismac.).
australis, Spreng. in Ersch & Gruber, Encyc. i. 348 = D. australe.
Damasonium, Sm. in Rees, Cycl. Suppl. u. 1 = D. stellatum.
europaeus, Spreng. in Ersch & Gruber, Encyc. i. (1818) 348 = D. stellatum.
maior, Bercht. & Presl, Rostl. i. Alismac. 8 = D. stellatum.
minor, R. Br. Prod. 343 = D. australe.

ACTINOCARYA, Benth. in Benth. & Hook. f. Gen. ii. 846 (1876). *BORAGINEAE*, Benth. & Hook. f. ii. 846.
tibetica, *Benth. l. c.*—Reg. Himal.

ACTINOCHLOA, Willd. ex Beauv. Agrost. 41 (1812); et ex Roem. & Schult. Syst. ii. (1817) 22. 417 = **Bouteloua**, Lag. (Gramin.).
barbata, Roem. & Schult. l. c. ii. 420 = B. barbata.
bromoides, Roem. & Schult. l. c. = B. bromoides.
ciliata, Willd. ex Beauv. Agrost. 41 = Chloris ciliata.
gracilis, Willd. ex Roem. & Schult. Syst. ii. 418 = B. oligostachya.
hirsuta, Roem. & Schult. l. c. 149 = B. hirsuta.
humilis, Willd. ex Beauv. Agrost. 41 = B. prostrata.
juncifolia, Roem. & Schult. Syst. ii. 420 = B. juncifolia.
ovata, Roem. & Schult. l. c. = B. ovata.
procumbens, Roem. & Schult. l. c. 417 = B. tenuis.
prostrata, Roem. & Schult. l. c. 419 = B. prostrata.
scorpioides, Roem. & Schult. l. c. 420 = B. scorpioides.
simplex, Roem. & Schult. l. c. 418 = B. simplex.
tenuis, Willd. ex Beauv. Agrost. 41 = B. tenuis.

ACTINOCLADUS, E. Mey. Ind. Sem. Hort. Regiom. (1846) = **Capnophyllum**, Gaertn. (Umbellif.).
cinerascens, E. Mey. l. c. = C. africanum.

ACTINOCYCLUS, Klotzsch, in Monatsb. Akad. Berl. (1857) 14 = **Pyrola**, Tourn. (Ericac.).
secundus, Klotzsch, l. c. = P. secunda.

ACTINODAPHNE, Nees, in Wall. Pl. As. Rar. ii. 68 (1831). *LAURINEAE*, Benth. & Hook. f. iii. 160.
IOZOSTE, Nees, l. c. 63 (1831).
IOZOSMENE, Lindl. Introd. Nat. Syst. ed. II. 202 (1836).
acuminata, *Meissn. in DC. Prod.* xv. 1. 211.—Japon.
angustifolia, Hook. f. & Thoms. ex Meissn. in DC. Prod. xv. 1. 218 = Hookeri.
angustifolia, *Nees, in Wall. Pl. As. Rar.* iii. 34.—Ind. or.; Malaya.
areolata, *Blume, Mus. Bot. Lugd. Bat.* i. 344.—Java.
Blumii, Nees, Syst. Laurin. 598 = Litsea ferruginea.
bornecnsis, *Meissn. in DC. Prod.* xv. 1. 213.—Borneo.
Candolleana, Meissn. l. c. 219 = specciosa.
celebica, *Miq. Fl. Ind. Bat.* i. 972.—Ins. Celebes.
chinensis, *Nees, Syst. Laurin.* 600.—China.
cochinchinensis, *Meissn. in DC. Prod.* xv. 1. 216.—Cochinch.
concolor, Blume, Mus. Bot. Lugd. Bat. i. 345 = scleroptera.
concolor, *Nees, in Wall. Pl. As. Rar.* iii. 31.—Ind. or.
confertiflora, *Meissn. in DC. Prod.* xv. 1. 216.—Reg. Himal.
corymbosa, *Blume, Mus. Bot. Lugd. Bat.* i. 343.—Sumatra.
elegans, *Thw. Enum. Pl. Zeyl.* 256.—Zeylan.
furfuracea, *Blume, Mus. Bot. Lugd. Bat.* i. 345.—Malaya.
glabra, *Blume, l. c.* i. 344.—Java.
glauca, *Nees, Syst. Laurin.* 599.—Zeylan.
glomerata, *Nees, l. c.* 597.—Java.
gracilis, *Miq. Fl. Ind. Bat. Suppl.* 368.—Sumatra.
heterophylla, Blume, Mus. Bot. Lugd. Bat. i. 342 = angustifolia.
hirsuta, *Blume, l. c.* 345.—Java.
Hookeri, *Meissn. in DC. Prod.* xv. 1. 218.—Ind. or.
javanica, *Miq. Fl. Ind. Bat.* i. 1. 969.—Java.
lanata, *Meissn. in DC. Prod.* xv. 1. 219.—Ind. or.
lancifolia, *Meissn. l. c.* 211.—Japon.
macrophylla, *Nees, Syst. Laurin.* 598.—Java.
macroptera, *Miq. Fl. Ind. Bat.* i. 1. 970.—Sumatra.

ACTINODAPHNE:—
mollis, *Blume, Mus. Bot. Lugd. Bat.* i. 343.—Sumatra.
molochina, *Nees, Syst. Laurin.* 603.—Zeylan.
moluccana, *Blume, Mus. Bot. Lugd. Bat.* i. 344.—Ins. Molucc.
Moonii, Thw. Enum. Pl. Zeyl. 256 = molochina.
multiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 230.—Ins. Fiji.
obovata, *Blume, Mus. Bot. Lugd. Bat.* i. 342.—Ind. or.
pauciflora, *Blume, l. c.*—Borneo.
pedunculata, *Meissn. in DC. Prod.* xv. 1. 211.—China.
procera, *Nees, Syst. Laurin.* 605.—Java.
pruinosa, *Nees, in Wall. Pl. As. Rar.* ii. 68.—Ind. or.
pubescens, *Blume, Mus. Bot. Lugd. Bat.* i. 345.—Malaya.
quercina, *Blume, l. c.* 344.—Java.
reticulata, *Meissn. in DC. Prod.* xvi. 1. 212.—Ind. or.
rufescens, *Blume, Mus. Bot. Lugd. Bat.* i. 343.—Malaya.
Rumphii, *Blume, l. c.* 344.—Ins. Molucc.
salicina, *Meissn. in DC. Prod.* xv. 1. 212.—Ind. or.
scleroptera, *Miq. Fl. Ind. Bat.* i. 1. 972.—Malaya.
sesquipedalis, *Hook. f. & Thoms. ex Meissn. in DC. Prod.* xv. 1. 216.—Ind. or.
Sieboldiana, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 114, 116.—Japon.
sikkimensis, *Meissn. in DC. Prod.* xvi. 1. 213.—Reg. Himal.
specciosa, *Nees, Syst. Laurin.* 602.—Zeylan.
sphaerocarpa, *Nees, l. c.* 605.—Java.
stenophylla, *Thw. Enum. Pl. Zeyl.* 256.—Zeylan.
Thwaitesii, Meissn. in DC. Prod. xv. 1. 212 = stenophylla.

ACTINODIUM, Schau. ex Schlecht. in Linnaea, x. (1835) 311. *MYRTACEAE*, Benth. & Hook. f. i. 696.
TRIPHELIA, R. Br. ex Endl. in Enum. Pl. Hueg. 48 (1837).
Cunninghami, *Schau. ex Lindl. Introd. Nat. Syst.* ed. II. 440.—Austral.
proliferum, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 17 = Cunninghami.

ACTINOLEMA, Fenzl, Pugill. Pl. Nov. Syr. 16. t. 12 (1842) = **Astrantia**, Linn. (Umbellif.).
eryngioides, Fenzl, l. c.—Oriens.
macrolema, Boiss. Fl. Orient. ii. 831.—Oriens.

ACTINOLEPIS, DC. Prod. v. 655 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 399.
HYMENOXYS, Torr. & Gray, Fl. N. Am. ii. 380 (1841).
anthemoides, *A. Gray, in Bot. Calif.* i. 378.—Calif.
coronaria, *A. Gray, in Proc. Am. Acad.* ix. (1874) 197.—Calif.
lanosa, *A. Gray, l. c.*—Utah.
Lemmoni, *A. Gray, l. c.* xvi. (1881) 101.—Calif.
multicaulis, *DC. Prod.* v. 656.—Calif.
mutica, *A. Gray, in Proc. Am. Acad.* ix. (1874) 197.—Calif.
nivea, *A. Gray, l. c.*—Calif.
tenella, *A. Gray, l. c.*—Calif.
Wallacei, *A. Gray, l. c.*—Utah.

ACTINOMERIS, Nutt. Gen. Am. ii. 181 (1818). *COMPOSITAE*, Benth. & Hook. f. ii. 379.
ACTIMERIS, Rafin. in Am. Monthly Mag. (1819) 195.
PTEROPHYTON, Cass. in Bull. Soc. Philom. (1818) 76.
RIDAN, Adans. Fam. ii. 130 (1763).
alata, Nutt. Gen. Am. ii. 181 = Verbesina occidentalis.
alba, *Torr. & Gray, Fl. N. Am.* ii. 335.—Am. bor.
alternifolia, DC. Prod. v. 572 = squarrosa.
Ellioti, *Rafin. New Fl. Am.* i. 61.—Am. bor.
helianthoides, *Nutt. Gen. Am.* ii. 181.—Am. bor.
heterophylla, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 6.—Am. bor.
longifolia, *A. Gray, in Smithson. Contrib.* v. (1853); *Pl. Wright.* ii. 89.—Arizona.
nudicaulis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 364.—Florida.
oppositifolia, DC. Prod. v. 575 = helianthoides.
oppositifolia, Fresen. Ind. Sem. Hort. Francof. (1836) = squarrosa.
ovata, *Nutt. Gen. Am.* ii. 181.—Mexico.

ACTINOMERIS :—
pauciflora, *Nutt. in Am. Journ. Sc.* v. (1822) 301.—Florida.
pedunculosa, DC. Prod. v. 576 = Verbesina Capitaneja.
procera, Steud. Nom. ed. II. i. 23 = squarrosa.
squarrosa, *Nutt. Gen. Am.* ii. 181.—Am. bor. occ.
stricta, *Hemsl. Biol. Centr. Am. Bot.* ii. 186.—Mexico.
tetragona, *DC. Prod.* v. 576.—Mexico.
tetraptera, *DC. l. c.* 575.—Mexico.
Wrightii, *A. Gray, in Mem. Am. Acad.* iv. (1849) (*Pl. Fendl.*) 85.—N. Mexico.

ACTINOMORPHE, Miq. Comm. Phyt. 102 (1840) = **Heptapleurum**, Gaertn. (Araliac.).
humilis, Miq. l. c. = H. parasiticum.

ACTINOPAPPUS, Hook. f. ex A. Gray, in Hook. Kew Journ. iv. (1852) 226 = **Rutidosis**, DC. (Compos.).
Drummondii, A. Gray, l. c. = R. Pumilio.
perpusillus, Hook. f. ex A. Gray, l. c. = R. Pumilio.

ACTINOPHORA, Wall. Cat. n. 1163 (1832) = **Schoutenia**, Korth. (Tiliac.).
fragrans, Wall. l. c. = S. ovata.

ACTINOPHYLLUM, Ruiz & Pav. Prod. 51. t. 8 (1794) = **Sciadophyllum**, P. Br. (Araliac.).
acuminatum, Pav. in Mem. Acad. Med. Madr. i. (1797) 196 = S. acuminatum.
angulatum, H. B. & K. Nov. Gen. et Sp. v. 92 = S. Humboldtianum.
angulatum, Pav. in Mem. Acad. Med. Madr. i. (1797) 194 = S. angulatum.
conicum, Pav. l. c. 195 = S. conicum.
digitatum, Wall. ex Steud. Nom. ed. II. i. 23 = S. digitatum.
divaricatum, Blume, Cat. Gew. Buitenz. 42 = S. divaricatum.
farinosum, Blume, l. c. 43 = S. tomentosum.
pedicellatum, Pav. in Mem. Acad. Med. Madr. i. (1797) 195 = S. pedicellatum.
pentandrum, Pav. l. c. 197 = S. pentandrum.

ACTINORHYTIS, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 184. *PALMAE*, Benth. & Hook. f. iii. 889.
calapparia, *H. Wendl. & Drude, l. c.*—Malaya.

ACTINOSCHOENUS, Benth. in Hook. Ic. Pl. xiv. 33. t. 1346 (1881). *CYPERACEAE*, Benth. & Hook. f. iii. 1058.
ARTHROSTYLIS, Boeck. in Linnaea, xxxvii. (1873) 523.
chinensis, *Benth. l. c.*—Hongkong.
filiformis, *Benth. l. c.*—Zeylan.
Thouarsii, *Benth. l. c.*—Madag.

ACTINOSPERMUM, Ell. Sketch, ii. 448 (1824) = **Baldwinia**, Nutt. (Compos.).
angustifolium, Torr. & Gray, Fl. N. Am. ii. 389 = B. multiflora.
multiflorum, Ell. Sketch, ii. 441 = B. multiflora.

ACTINOSPORA, Turcz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 (1835) = **Cimicifuga**, Linn. (Ranunc.).
dahurica, Turcz. ex Fisch. & Mey. l. c. = C. davurica.
frigida, Fisch. & Mey. l. c. = C. frigida.
pterosperma, Walp. Rep. i. 60 = C. davurica.
Wallichii, Meissn. Gen. Comm. 4 = C. foetida.

ACTINOSTEMA, Lindl. Veg. Kingd. 281 (1846) err. typ. = **Actinostemon**, Mart. (Euphorb.).

ACTINOSTEMMA, Griff. Pl. Cantor, 24. t. 3 (1837?). *CUCURBITACEAE*, Benth. & Hook. f. i. 838.
MITROSICYOS, Maxim. in Ann. Sc. Nat. Sér. IV. xiii. (1860) 95.
POMASTERION, Miq. Ann. Mus. Bot. Lugd. Bat. ii. (1865) 80.
japonicum, Miq. l. c. iii. 188 = racemosum.
lobatum, *Maxim. in Franch. & Sav. Enum. Pl. Jap.* i. 175.—Japon.
paniculatum, *Maxim. ex Cogn. in DC. Monog. Phan.* iii. 920.—China.
racemosum, *Maxim. ex Cogn. l. c.* 922.—China.

ACTINOSTEMMA :—
tenerum, *Griff. Pl. Cantor*, 25. t. 3.—Ind. or.
tenerum, Naud. in Ann. Sc. Nat. Sér. V. (1866) 39 = lobatum.

ACTINOSTEMON, Mart. ex Klotzsch, in Wiegm. Archiv, vii. (1841) 184. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 338.
DACTYLOSTEMON, Klotzsch, l. c. 181.
acuminatus, Baill. Etud. Gén. Euphorb. 532. t. 5 = concolor.
angustifolius, Klotzsch, ex Regel, Gartenfl. (1859) 363 = concolor.
caribaeus, *Griseb. in Goett. Abh.* vii. (1857) 168.—Bras.; Ind. occ.
concolor, Muell. Arg. in DC. Prod. xv. II. 1193 = caribaeus.
cuneatus, Baill. Adansonia, v. (1864–65) 535.—Bras.
echinatus, *Muell. Arg. in Linnaea*, xxxii. (1863) 107.—Bras.
furcatus, Klotzsch, ex Baill. Etud. Gén. Euph. 532 = caribaeus.
grandifolius, Baill. l. c. = caribaeus.
imbricatus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 216.—Bras.
Klotzschianus, Baill. Adansonia, v. (1864–65) 334 (= *Dactylostemon grandifolius*).
lanceolatus, *Saldanha, in Baill. l. c.* viii. (1867–68) 263.—Bras.
lasiocarpoides, *Baill. l. c.* v. (1864–65) 334.—Bras.
lasiocarpus, *Baill. l. c.*—Bras.
macrocarpus, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 597.—Bras.
marginatus, Klotzsch, ex Baill. Etud. Gén. Euph. 532 = caribaeus.
multiflorus, *Muell. Arg. in Linnaea*, xxxii. (1863) 111.—Bras.
oligandrus, *Baill. Adansonia*, v. (1864–65) 335.—Bras.
polymorphus, Muell. Arg. in Linnaea, xxxii. (1863) 108 = caribaeus.
sessilifolius, Klotzsch, Ind. Sem. Hort. Berol. App. (1851) 13 = caribaeus.
Sprengelii, Baill. Etud. Gén. Euph. 333 (= *Dactylostemon Klotzschii*).
trachycarpus, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 591.—Bras.
verticillatus, *Baill. Adansonia*, v. (1864–65) 334.—Bras.

ACTINOSTIGMA, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) 259 = **Seringia**, J. Gay (Sterculiac.).
lanceolatum, Turcz. l. c.—Austral.

ACTINOSTIGMA, Welw. Apont. 560 (1858) = **Symphonia**, Linn. f. (Gutlif.).
speciosum, Welw. l. c. = S. globulifera.

ACTINOSTROBUS, Miq. in Lehm. Pl. Preiss. i. 644 (1845). *CONIFERAE*, Benth. & Hook. f. iii. 425.
acuminatus, *Parl. Ind. Sem. Hort. Florent.* (1862) 25.—Austral.
pyramidalis, *Miq. in Lehm. Pl. Preiss.* i. 644.—Austral.

ACTINOTUS, Labill. Pl. Nov. Holl. i. 67. t. 92 (1804). *UMBELLIFERAE*, Benth. & Hook. f. i. 880, 1008.
ERIOCALIA, Sm. Exot. Bot. t. 78, 79 (1806).
HEMIPHUES, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 469.
HOLOTOME, Endl. Gen. 766 (1839).
bellidioides, *Benth. Fl. Austral.* iii. 369.—Austral.
Gibbonsii, *F. Muell. Fragm.* vi. 23.—Austral.
glomeratus, *Benth. Fl. Austral.* iii. 369.—Austral.
Helianthi, *Labill. Pl. Nov. Holl.* i. 67. t. 92.—Austral.
leucocephala, *Benth. in Enum. Pl. Hueg.* 56.—Austral.
minor, *DC. Prod.* iv. 83.—Austral.
Novae-Zealandiae, *Petrie, in Trans. N. Z. Inst.* xiii. (1881) 324.—N. Zel.
omnifertilis, *F. Muell. ex Benth Fl. Austral.* iii. 368.—Austral.
rhomboideus, *Benth. Fl Austral.* iii. 368.—Austral.

ACTIPSIS, Rafin. Fl. Tellur. ii. 43 (1836) = **Solidago**, Linn. (Compos.).
glomerata, Rafin. l. c. = S. glomerata.
squamosa, Rafin. l. c. = praec.?

ACTISPERMUM, Rafin. New Fl. Am. i. 61 (1836) (= ACTINOSPERMUM, Ell.) = **Baldwinia**, Nutt. (Compos.).

ACUAN, Medic. Theod. 62 (1786) = **Desmanthus**, Willd. (Legum.).
arboreum, Medic. l. c.—Hab.?
virgatum, Medic. l. c. = D. virgatus.

ACUBA, Link, Enum. Hort. Berol. ii. 386 (1822), sphalm. = **Aucuba**, Thunb. (Cornac.).

ACUBALUS, Neck. Elem. ii. 167 (1790) = **Cucubalus**, Tourn. (Caryophyll.).

ACUNA, Endl. Gen. 759 (1839) = seq.

ACUNNA, Ruiz & Pav. Prod. 69. t. 12 (1794) = **Befaria**, Mutis (Ericac.).
lanceolata, Cav. ex DC. Prod. vii. 731 = B. coarctata.
lanceolata, Ruiz & Pav. Syst. Veg. 124 = B. aestuans.
oblonga, Ruiz & Pav. l. c. 123 = B. oblonga.

ACURA, Hill, Veg. Syst. v. 52 (1769) (Compos.).

ACUROA, J. F. Gmel. Syst. 1083 (1791) = ACOUROA, Aubl. = **Ecastaphyllum**, P. Br. (Legum.).

ACUSTON, Rafin. Sylva Tellur. 131 (1838) = **Farsetia**, Desv. (Crucif.).
clypeatum, Rafin. l. c. 131 = F. clypeata.
lunaroides, Rafin. l. c. = F. lunarioides.

ACYNOS, Pers. Syn. ii. 131 (1807) (err. typ. '151'). = ACINOS, Rupp. = **Calamintha**, Moench (Labiat.).

ACYNTHA, Medic. Theodora, 76 (1786) = **Sansevieria**, Thunb. (Haemodor.).
guinensis, Medic. l. c. = S. guineensis.

ADA, Lindl. Fol. Orch. (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 566.
aurantiaca, *Lindl. l. c.*—N. Granat.

ADAMANTOGETON, Schrad. ex Nees, in Mart. Fl. Bras. ii. I. 165 (1842) = **Lagenocarpus**, Nees, (Cyper.).

ADAMBEA, Lam. Encyc. 39 (1783) = **Lagerstroemia**, Linn. (Lythrar.).
glabra, Lam. l. c. = L. Flos-reginae.
hirsuta, Lam. l. c. = L. hirsuta.

ADAMBOE, Rafin. Fl. Tellur. iv. 73 (1836) = **Argyreia**, Lour. (Convolv.).
bicolor, Rafin. l. c. = A. tiliaefolia.

ADAMIA, Wall. Tent. Fl. Nep. 46. t. 36 (1824) = **Dichroa**, Lour. (Saxifrag.).
chinensis, Gardn. & Champ. in Hook. Kew Journ. i. 311 (1849) = D. febrifuga.
cyanea, Wall. Tent. Fl. Nep. 46. t. 36 = D. febrifuga.
sylvatica, Meissn. in Walp. Rep. ii. 377 = D. febrifuga.
versicolor, Fortune, in Lindl. Journ. Hort. Soc. i. (1846) 298 = D. febrifuga.

ADAMSIA, Fisch. ex Steud. Nom. ed. I. 366, 367, in syn. (1821) = **Geum**, Linn. (Rosac.).
anemonoides, Fisch. ex Steud. l. c. 366 = G. anemonoides.
glacialis, Fisch. ex Steud. l. c. 366 = G. glaciale.
reptans, Fisch. ex Steud. l. c. 367 = G reptans.
rotundifolia, Fisch. ex Steud. l. c. = G. rotundifolium.

ADAMSIA, Willd. in Mag. Ges. Naturf. Fr. Berl. ii. (1808) 16 = **Puschkinia**, Adams (Liliac.).
scilloides, Willd. l. c. = P. scilloides.

ADANSONIA, Linn. Sp. Pl. 1190 (1753). *MALVACEAE*, Benth. & Hook. f. i. 209.
BAOBAB, Adans. Fam. ii. 399 (1763).
OPHELUS, Lour. Fl. Cochinch. 412 (1790).
Bahobab, Linn. Sp. Pl. ed. II. 960 = digitata.
Baobab, Gaert. Fruct. ii. 253. t. 135 = digitata.

ADANSONIA :—
digitata, *Linn. Sp. Pl.* 1190.—Afr. trop.; Ind. or.
Gregorii, *F. Muell. in Hook. Kew Journ.* ix. (1857) 14.—Austral.
integrifolia, Rafin. Sylva Tellur. 149 = A. Situla.
madagascaricnsis, *Baill. Adansonia*, xi. (1873-76) 251.—Madag.
microphylla, A. Rich. Fl. Cub. Fanerog. ii. 325, lapsu = Adelia microphylla.
pedunculosa, A. Rich. l. e. lapsu = Adelia pedunculosa.
Scutula, Steud. Nom. ed. II. i. 24, sphalm. = seq.
Situla, *Spreng. Syst.* iii. 124.—Afr. or.

ADAPHUS, Neck. Elem. ii. 240 (1790) = **Laurus**, Tourn.

ADATODA, Rafin. Fl. Tellur. iv. 60 (1836) = **Justicia**, Linn. (Acanth.).
arborea, Rafin. l. c. 61 = J. Adhatoda.
Betonica, Rafin. l. c. = J. Betonica.
repens, Rafin. l. c.—Hab. ?

ADELANTHUS, Endl. Gen. 1327. n. 6839; Suppl. ii. 30 (1841) = **Pyrenacantha**, Wight (Olacin.).
grandiflorus, Harv. ex Baill. Adansonia, x. (1872) 271 = P. grandiflora.
scandens, Endl. ex Baill. in DC. Prod. xvii. 18 = P. scandens.

ADELASTER, Lindl.in Proc. Roy. Hort. Soc. i. (1861) 568; Lem. in Illustr. Hortic. t. 320 = **Eranthemum** Linn. ? (Acanthac.).
albivenis, Lindl. ll. cc.—Hab. ?

ADELBERTIA, Meissn. Gen. Pl. 114 (1837) = **Merania**, Sw. (Melast.).
calophylla, Meissn. l. c. (= *Davya calophylla*).—Bras.

ADELIA, P. Br. Nat. Hist. Jamaic. 361 (1756) = **Forestiera**, Poir. (Oleac.).
acuminata, Michx. Fl. Bor. Am. ii. 225 = F. acuminata.
cassinioides, Steud. Nom. ed. II. i. 24 = F. cassinoides.
ligustrina, Michx. Fl. Bor. Am. ii. 224 = F. ligustrina.
ovata, [Desf.] Hort. Par. ex Pers. Syn. ii. 635.—Am. bor.
porulosa, Michx. Fl. Bor. Am. ii. 224 = F. porulosa.
virgata, Poir. Encyc. Suppl. i. 132 = Securinega buxifolia.

ADELIA, Linn. Syst. ed. X. (1759) 1298. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 312.
RICINELLA, Muell. Arg. in Linnaea, xxxiv. (1865) 153.
Acidoton, Blanco, Fl. Filip. ed. I. 815 = Sumbavia rottleroides.
Acidoton, *Linn. Syst.* ed. X. 1298; *Pl. Jam. Pugill.* 29.—Jamaica; Cuba.
anomala, Juss. ex Steud. Nom. ed. II. i. 24 = ovata.
anomala, Poir. Encye. Suppl. i. 132 = Erythrococca aculeata.
barbata, Blanco, Fl. Filip. ed. II. 561 = Mallotus ricinoides.
barbinervis, *Cham. & Schlecht. in Linnaea*, vi. (1831) 362.—Mexico.
Bernardia, Blanco, Fl. Filip. ed. I. 814 = Mallotus ricinoides.
Bernardia, Linn. Syst. ed. X. 1298 = Bernardia dichotoma.
caperoniaefolia, Baill. Adansonia, iv. (1864) 376 = Bernardia caperoniaefolia.
castanicarpa, Roxb. Fl. Ind. iii. 848 = Chaetocarpus castanocarpus.
celastrinea, Baill. Adansonia, iv. (1864) 375 = Bernardia axillaris.
cordifolia, *Roxb. Fl. Ind.* iii. 849.—Ins. Molucc.
cuneata, Wall. Cat. n. 7954 = Homonoia retusa.
dodecandra, Scsse, ex Cav. in Anal. Cienc. Nat. v. (1802) 254 = Bernardia interrupta.
ferruginea, Poit. ex Baill. Etud. Gén. Euph. 418 = Bernardia leprosa.
glandulosa, Blanco, Fl. Filip. ed. I. 814 = Alchornea javensis.
gracilis, Salisb. Prod. 388 = A. Acidoton.
hirsutissima. Baill. Adansonia, iv. (1864) 372 = Bernardia peduncularis.
Houlletiana, Baill. l. c. 373 = Bernardia axillaris.
javanica, *Miq. Fl. Ind. Bat.* i. II. 388.—Java.

ADELIA :—
Martii, Spreng. Syst. iii. 147 = Bernardia axillaris.
microphylla, *A. Rich. Fl. Cub. Fanerog.* ii. 209.—Cuba.
monoica, Blanco, Fl. Filip. ed. II. 561 = Mallotus moluccanus.
myrtifolia, *Vent. ex Spreng. Syst.* iii. 147.—Hab. ?
nereifolia, Heyne, ex Roth, Nov. Pl. Sp. 375 = Homonoia riparia.
oaxacana, *Hemsl. Biol. Centr. Am. Bot.* iii. 129.—Mexico.
papillaris, Blaneo, Fl. Filip. ed. II. 562 = Mallotus Zollingeri.
peduneulosa, *A. Rich. Fl. Cub. Fanerog.* ii. 210.—Cuba.
pulchella, Baill. Adansonia, iv. (1864) 374 = Bernardia pulehella.
resinosa, Blaneo, Fl. Filip. ed. II. 562 = Claoxylon Walliehianum.
retusa, Wight, Ic. t. 1869 = Homonoia retusa.
Ricinella, *Linn. Syst.* ed. X. 1298; *Pl. Jam. Pugill.* 29.—Ind. oce.
scabrida, Baill. Adansonia, iv. (1864) 373 = Bernardia axillaris.
scandens, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 350, nomen.—Malaya.
spartioides, Baill. Adansonia, v. (1865) 319 = Bernardia spartioides.
sylvestris, *Griseb. in Goett. Nachr.* (1865) 174.—Cuba.
tamanduana, Baill. Adansonia, iv. (1864) 374 = Bernardia tamanduana.
timoriana, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 350, nomen.—Ins. Timor.
triloba, *Hemsl. Biol. Centr. Am. Bot.* iii. 130.—Panama.

ADELIOIDES, R. Br. ex Benth. Fl. Austral. i. 59 (1863) = seq.

ADELIOPSIS, Benth. in Benth. & Hook. f. Gen. i. 1. Add. 436 (1862). *MENISPERMACEAE*, Benth. & Hook. f. Gen. i. 1. Add. 436; i. 964.
ADELIOIDES, R. Br. ex Benth. Fl. Austral. i. 59 (1863).
deeumbens, Benth. l. c.; et Fl. Austral. i. 59.—Austral.

ADELMANNIA, Reichb. Consp. 110 (1828) = **Borrichia**, Adans. (Compos.).

ADELOBOTRYS, DC. Prod. iii. 127 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 750.
SARMENTARIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 140.
SPHANELLOPSIS, Steud. ex Naud. l. c. 137.
XERACINA, Rafin. Sylva Tellur. 98 (1838).
adscendens, *Triana, in Journ. Bot.* v. (1867) 210.—Am. trop.
barbata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 68.—Bras.
ciliata, *Triana, l. c.* 68.—Guiana.
fuscescens, *Triana, l. c.*—N. Granat.
laxiflora, *Triana, l. c.*—Guiana.
Lindeni, *Naud. in Belg. Hortic.* xv. (1865) 98.—Bras.
rotundifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 68.—Bras.
seandens, *DC. Prod.* iii. 127.—Guiana.
scandens, Macfad. Fl. Jamaic. ii. 99 = adscendens.

ADELODA, Rafin. Fl. Tellur. iv. 61 (1836) = **Justicia**, Linn. (Acanthac.).
integra, Rafin. l. c.—Hab. ?
serrata, Rafin. l. c. = J. serrata.

ADELONEMA, Schott, Prod. Aroid. 316 (1860) = **Homalonema** Schott (Aroid.).
erythropus, Schott, l. c. 317.—Bras.

ADELONENGA, Becc. in Ann. Jard. Buitenz. ii. (1885) 82. *PALMAE*, Benth. & Hook. f. iii. 885.
Geelvinkiana, *Becc. l. c.*—Malaya.
variabilis, *Becc. l. c.*—Malaya.

ADELOSA, Blume, Mus. Bot. Lugd. Bat. i. 176 (1850). *VERBENACEAE*, Benth. & Hook. f. ii. 1153.
microphylla, *Blume, l. c.*—Madag.

ADELOSTEMMA, Hook. f. in Hook. Ic. Pl. xv. (1883) 22. *ASCLEPIADEAE.*
gracillimum, *Hook. f. l. c.*—Burma.

ADELOSTIGMA, Steetz, in Peters, Reise Mossamb. Bot. 428 (1863). *COMPOSITAE*, Benth. & Hook. f. ii. 285.
athrixioides, *Steetz, l. c.* 430.—Afr. trop.
senegalensis, *Benth. in Hook. Ic. Pl.* t. 1144.—Afr. trop.

ADENACANTHUS, Nees, in Wall. Pl. As. Rar. iii. 75 & 196 (1832) = **Strobilanthes**, Blume (Acanthac.).
acuminatus, Nees, l. c. iii. 84 = S. acuminatus.
latifolia, Nees, in DC. Prod. xi. 197.—Hab. ?

ADENACHAENA, DC. Prod. vi. 49 (1837) = **Phymaspermum**, Less. (Compos.).
leptophylla, DC. l. e. = P. leptophyllum.
parvifolia, DC. l. c. = P. parvifolium.
pubescens, DC. l. c. 49.—Afr. austr.

ADENANDRA, Willd. Enum. Hort. Berol. 256. (1809). *RUTACEAE*, Benth. & Hook. f. i. 290.
GLANDULIFOLIA, Wendl. Collect. t. 33, 37 (1808).
HAENKAEA, Usteri, Ann. Bot. vi. (1793) 117.
HAENKEA, F. W. Schmidt, Neue u. Seltne Pfl. 19 (1793); nomen prius.
OCENIA, F. G. Dietr. Lex. Gaert. Nachtr. v. 307 (1819).
OCKEA, F. G. Dietr. l. c. i. 105 (1815).
OCKENIA, Steud. Nom. ed. II. 24. (1840).
OKEA, Steud. l. e. ed. I. 562 (1821).
acuminata, Sweet, Hort. Brit. ed. I. 88 = amoena.
alba, *Roem. & Schult. Syst.* v. 451.—Afr. austr
alternifolia, *Cham. & Schlecht. in Linnaea*, v. (1830) 52.—Afr. austr.
amoena, *Bartl. & Wendl. f. Diosm.* 80.—Afr. austr.
amoena, Link, Enum. Hort. Berol. i. 239 = fragrans.
Bartlingiana, Eckl. & Zeyh. Enum. 101 = viscida.
biseriata, *E. Mey. ex Bartl. & Wendl. f. Diosm.* 75.—Afr. austr.
brachyphylla, *Schlecht. in Linnaea*, vi. (1831) 199.—Afr. austr.
calycina, *Steud. Nom.* ed. II. i. 24.—Afr. austr.
ciliata, *Sond. in Harv. & Sond. Fl. Cap.* i. 388.—Afr. austr.
cistoides, Eckl. & Zeyh. Enum. 100 = biseriata.
cordata, Link, Enum. Hort. Berol. i. 239 = Barosma serratifolia.
coriacea, Eckl. & Zeyh. Enum. 100 ex Sond. in Harv. & Sond. Fl. Cap. i. 389 = macrodenia.
coriacea, Bartl. & Wendl. f. Diosm. 75 = biseriata.
coriacea, *Lichtst. in Roem. & Schult. Syst.* v. 452.—Afr. austr.
crenata, Willd. ex Bartl. & Wendl. f. Diosm. 82 = umbellata.
cuspidata, Eckl. & Zeyh. Enum. 101 ex Sond. in Harv. & Sond. Fl. Cap. i. 385 = obtusata.
cuspidata, *E. Mey. ex Bartl. & Wendl. f. Diosm.* 87.—Afr. austr.
fragrans, *A. Gonault, in Rev. Hortic.* Sér. IV. i. (1852) 321.—Afr. austr.
fragrans, Eckl. & Zeyh. Enum. 99 = serpyllacea.
fragrans, *Roem. & Schult. Syst.* v. 451.—Afr. austr.
glandulosa, Lichtst. ex Roem. & Schult. Syst. v. 450 = umbellata.
gracilis, *Eckl. & Zeyh. Enum.* 101.—Afr. austr.
humilis, *Eckl. & Zeyh. l. c.* 99.—Afr. austr.
intermedia, Eckl. & Zeyh. l. c. 99 = marginata.
Kraussii, *Meissn. in Flora*, xxvii. (1844) 301.—Afr. austr.
lasiantha, *Sond. in Harv. & Sond. Fl. Cap.* i. 390.—Afr. austr.
linearis, A. Juss. in Mém. Mus. Par. xii. (1825) 471 = uniflora.
linifolia, *Bartl. in Linnaea*, xvii. (1843) 359.—Afr. austr.
macradenia, *Sond. in Harv. & Sond. Fl. Cap.* i. 389.—Afr. austr.
marginata, Eckl. & Zeyh. Enum. 99 = linifolia.
marginata, *Roem. & Schult. Syst.* v. 452.—Afr. austr.
mundiaefolia, *Eckl. & Zeyh. Enum.* 99.—Afr. austr.
muraltioides, *Eckl. & Zeyh. l. c.* 106.—Afr. austr.
obtusata, *Sond. in Harv. & Sond. Fl. Cap.* i. 385.—Afr. austr.
ovata, Link, Enum. Hort. Berol. i. 239 = amoena.

ADENANDRA:—
pubescens, *Sond. in Harv. & Sond. Fl. Cap.* i. 385.—Afr. austr.
rosmarinifolia, A. Juss. in Mém. Mus. Par. xii. (1825) 471 = marginata.
rotundifolia, *Eckl. & Zeyh. Enum.* 101.—Afr. austr.
rugosa, Otto, ex Steud. Nom. ed. II. i. 24 = umbellata.
Schmidtiana, *Steud. l. c.*—Afr. austr.
serpyllacea, *Bartl. in Linnaea*, xvii. (1843) 359.—Afr. austr.
serratifolia, Link, Enum. Hort. Berol. i. 239 = Barosma serratifolia.
speciosa, Link, l. c. = umbellata.
tetragona, Sweet, Hort. Suburb. 45 = Acmadenia tetragona.
trigona, *Eckl. & Zeyh. Enum.* 100.—Afr. austr.
umbellata, *Willd. Enum. Hort. Berol.* 257.—Afr. austr.
uniflora, E. Mey. & Drège, in Harv. & Sond. Fl. Cap. i. 391 = serpyllacea.
uniflora, *Willd. Enum. Hort. Berol.* 256.—Afr. austr.
villosa, Eckl. & Zeyh. Enum. p. 101 = pubescens.
villosa, Hort. ex Harv. & Sond. Fl. Cap. i. 388 = uniflora.
villosa, Lichtst. ex Roem. & Schult. Syst. v. 452 = cuspidata.
viscida, *Eckl. & Zeyh. Enum.* 101.—Afr. austr.
Wendlandiana, Eckl. & Zeyh. Enum. 100 = umbellata.

ADENANTHERA, Royen, ex Linn. Coroll. Gen. 7 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 590.
GONSII, Adans. Fam. ii. 318 (1763).
PONGELION, Adans. Fam. ii. 319 (1763).
abrosperma, *F. Muell. Fragm.* v. 30.—Austral.
aculeata, Roxb. Hort. Beng. 90; Fl. Ind. ii. 37 = Prosopis spicigera.
bicolor, *Moon, Cat. Pl. Ceyl.* 34, *ex Thw. Enum. Pl. Zeyl.* 98.—Zeylan.; Malaya.
Bonplandiana, H. B. & K. Nov. Gen. et Sp. vi. 311 = Entada polystachya.
chrysostachys, Benth. in Hook. Lond. Journ. Bot. iv. (1852) 343 = Piptadenia chrysostachys.
circinalis, DC. Prod. ii. 446 = Pithecolobium clypearia.
falcata, *Linn. Diss. Herb. Amb.* 14 (*Amoen. Acad.* iv. 124).—Ins. Molucc.
Falcataria, Linn. Sp. Pl. ed. II. 550 = falcata.
Gersenii, *Scheff. in Flora*, lii. (1869) 307.—Malaya.
glauca, *Span. in Hook. Comp. Bot. Mag.* i. (1835) 347, nomen.—Malaya.
Gogo, Blanco, Fl. Filip. ed. I. 358 = Entada scandens.
microsperma, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* xxvii. (1864) 58.—Java.
oudhensis, J. L. Stewart, ex Brand. For. Fl. Ind. 168 = Piptadenia oudhensis.
pavonina, *Linn. Sp. Pl.* 384.—As. trop.; Malaya.
polita, *Miq. Fl. Ind. Bat.* i. 47.—Malaya.
scandens, Forst. f. Prod. 33 = Entada scandens.
scandens, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 340 = Entada Adenanthera.
tetraptera, Schum. & Thonn. Beskr. Guin. Pl. 213 = Tetrapleura Thonningii.
triphysa, Dennst. Schluess. Hort. Malab. 32 = Ailanthus malabarica.

ADENANTHES, Knight, Prot. 96. (1809) = seq.

ADENANTHOS, Labill. Pl. Nov. Holl. i. 28. t. 36 (1804). *PROTEACEAE*, Benth. & Hook. f. iii. 173.
apiculata, *R. Br. Prod. Suppl.* 9.—Austral.
armata, Meissn. in DC. Prod. xiv. 313 = pungens.
barbigera, *Lindl. Swan Riv. App.* 36. n. 182.—Austral.
crenata, Willd. ex Spreng. Syst. i. 472 = cuneata.
cuneata, *Labill. Pl. Nov. Holl.* i. 28. t. 36.—Austral.
Cunninghamii, *Meissn. in Lehm. Pl. Preiss.* i. 513.—Austral.
Detmoldi, *F. Muell. Fragm.* viii. 149.—Austral.
Dobsoni, *F. Muell. l. c.* vi. 204.—Austral.
Drummondii, Meissn. in Lehm. Pl. Preiss. i. 514 = apiculata.
filifolia, *Benth. Fl. Austral.* v. 355.—Austral.
flabellifolia, [Salisb. in] Knight, Prot. 96 = cuneata.
flavidiflora, *F. Muell. Fragm.* i. 157.—Austral.
Forrestii, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 230.—Austral.
linearis, *Meissn. in DC. Prod.* xiv. 311.—Austral.
Meissneri, *Lehm. Pl. Preiss.* i. 512; ii. 248.—Austral.
obovata, *Labill. Pl. Nov. Holl.* i. 29. t. 37.—Austral.
procumbens, Meissn. in Lehm. Pl. Preiss. i. 512; ii. 248 = apiculata.
pungens, *Meissn. l. c.* i. 515.—Austral.
sericea, *Labill. Pl. Nov. Holl.* i. 29. t. 38.—Austral.
terminalis, *R. Br. in Trans. Linn. Soc.* x. (1810) 152.—Austral.
velutina, *Meissn. in DC. Prod.* xiv. 312.—Austral.
venosa, *Meissn. l. c.*—Austral.

ADENANTHUS, Roem. & Schult. Syst. iii. 22, 397 (1818) = praec.

ADENARIA, H. B. & K. Nov. Gen. et Sp. vi. 185. t. 549 (1823). *LYTHRARIEAE*, Benth. & Hook. f. i. 777.
floribunda, *H. B. & K. l. c.* 188. t. 549.—N. Granat.
griseleoides, H. B. & K. l. c. 185 = floribunda.
lanceolata, *Beurling, in Vet. Akad. Handl. Stockh.* (1854) 124.—Panama.
parvifolia, Hook. Ic. Pl. t. 166 = floribunda.
purpurata, H. B. & K. Nov. Gen. et Sp. vi. 185 = floribunda.

ADENARIUM, Rafin. in Am. Monthly Mag. (1818) 266 = **Arenaria**, Linn. (Caryophyll.).
marinum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 545 = Aren. peploides.
maritimum, Rafin. New Fl. Am. i. 62 = Aren. peploides.
peploides, Rafin. in Am. Monthly Mag. (1818) 266 = Aren. peploides.

ADENELEUTEROPHORA, Rodrig. Orch. Nov. ii. 170 (1881). *ORCHIDEAE.*
graminifolia, *Rodrig. l. c.* 171.—Bras.

ADENEMA, G. Don, Gen. Syst. iv. 201 (1837) = **Enicostema**, Blume (Gentian.).
hyssopifolium, G. Don, l. c. = E. littorale.

ADENESMA, Griseb. in DC. Prod. ix. 65 (1845) = praec.

ADENIA, Forsk. Fl. Aegypt. Arab. 77 (1775) = **Modecca**, Linn. (Passif.).
venenata, Forsk. l. c.—Arabia.

ADENIA, Torr. Fl. New York, ii. t. 122 (1843) = **Pilea**, Lindl. (Urtic.).
pumila, Torr. l. c. = P. pumila.

ADENILEIMA, Reichb. Nom. 159 (1841) = seq.

ADENILEMA, Blume, Bijdr. 1120 (1826) = **Neillia**, D. Don (Rosac.).
fallax, Blume, l. c. 1121 = N. thyrsiflora.

ADENILEMMA, Hassk. Cat. Hort. Bogor. Alt. 170 (1844) = praec.

ADENIUM, Roem. & Schult. Syst. iv. 35 & 411 (1819). *APOCYNACEAE*, Benth. & Hook. f. ii. 722.
arboreum, *Ehrenb. Symb. Phys.* 4.—Afr. trop.
Honghel, *A. DC. in DC. Prod.* viii. 412.—Afr. trop.
Honghel, Lindl. Bot. Reg. xxxii. t. 54 = obesum.
multiflorum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 279.—Afr. trop.
namaquanum, Wyley, ex Harv. Thes. Cap. ii. 11. t. 117 = Pachypodium namaquanum.
obesum, *Roem. & Schult. Syst.* iv. 411.—Arab.; Afr. trop.
speciosum, *Fenzl, in Sitzb. Acad. Wien. Math. Nat.* li. *Abth.* II. (1865) 140.—Afr. trop.

ADENOBASIUM, Presl, Symb. Bot. i. 39. t. 27 (1830) = **Sloanea**, Linn. (Tiliac.).
obtusifolium, Moric. Pl. Nouv. 83. t. 55 = S. obtusa.
salicifolium, Presl, Symb. Bot. i. 40. t. 27 = S. monosperma.

ADENOCALYMMA, Endl. Ench. 347 (1841) = seq.

ADENOCALYMNA, Mart. in Meissn. Gen. 300 (208) et in DC. Prod. ix. 199 (1840). *BIGNONIACEAE*, Benth & Hook. f. ii. 1036.
LEGUMINARIA, Bur. Monog. Bignon. 42 (1864).
MEMORA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 185.
PACHYPTERA, DC. Prod. ix. (1840) 175.
acutissimum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 395.—Bras.
alliaceum, Miers, l. c. 394 = Bignonia alliacea.
brachybotrys, *DC. Prod.* ix. 203.—Guiana.
bracteatum, *DC. l. c.* 200.—Bras.
bracteolatum, *DC. l. c.* 200.—Bolivia.
divaricatum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 390.—Bras.
comosum, *DC. Prod.* ix. 201.—Bras.
coriaceum, *DC. l. c.* 202.—Bras.
flavidum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 389.—Bras.
flavum, *Mart. ex DC. Prod.* ix. 202.—Bras.
floribundum, *DC. Prod.* ix. 201.—Bolivia.
grandifolium, *Mart. ex DC. l. c.* 199.—Bras.
Guillemini, *DC. l. c.* 202.—Bras.
inundatum, *Mart. ex DC. l. c.* 201.—Bras.
japurense, Mart. ex DC. l. c. 203.—Bras.
laevigatum, Mart. ex DC. l. c. 200 = marginatum.
lanceolatum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 390.—Bras.
longeracemosum, *Mart. ex DC. Prod.* ix. 201.—Bras.
macranthum, Mart. ex DC. l. c. 203.—Bras.
macrophyllum, *DC. l. c.* 199.—Bras.
macrostachyum, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 180.—Bras.
magnificum, *Mart. ex DC. Prod.* ix. 202.—Bras.
marginatum, *DC. l. c.* 200.—Bras.
nitidum, *Mart. ex DC. l. c.*—Bras.
prasinum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 395.—Bras.
paniculatum, *Benth. ex Miers, l. c.* 393.—Bras.
pliciflorum, *Mart. ex DC. Prod.* ix. 202.—Bras.
prehensile, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 388.—Bras.
Salzmanni, *DC. Prod.* ix. 200.—Bras.
scabriusculum, *Mart. ex DC. l. c.* 201.—Bras.
scansile, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 388.—Bras.
sepiarium, Mart. ex DC. Prod. ix. 200 = nitidum.
stipidula, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 392.—Bras.
subsessilifolium, *DC. Prod.* ix. 199.—Bras.
verruciferum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vii. (1861) 393.—Bras.

ADENOCALYX, Bert. ex H. B. & K. Nov. Gen. et Sp. vi. 330 (1823) = COULTERIA, H. B. & K. = **Caesalpinia**, Linn. (Legum.).
racemosus, Bert. ex H. B. & K. l. c. = C. mollis.
remotus, Steud. Nom. ed. II. i. 428, sphalm. = praec.

ADENOCARPUM, D. Don, ex Hook. & Arn. in Hook. Lond. Journ. Bot. iii. (1841) 318 = ADENOSPERMUM, Hook. & Arn. = **Chrysanthellum**, Rich. (Compos.).
tuberculatum, D. Don, ex Hook. & Arn. l. c. = C. indicum.

ADENOCARPUS, DC. Fl. Fr. Suppl. 549 (1815). *LEGUMINOSAE*, Benth. & Hook. f. i. 481.
anagyrifolius, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 246; xxii. (1875) 57, nomen.—Marocc.
anagyrus, *Spreng. Syst.* iii. 226.—Ins. Canar.
anisochilus, *Boiss. Diagn.* Ser. II. ii. 5.—Hispan.
benguellensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 47.—Afr. trop.
Bivonii, *Presl, Fl. Sic.* i. p. xix. = intermedius.
Boissieri, Webb, Iter Hispan. 53 = decorticans.
cebennensis, Delile, Ind. Sem. Hort. Monsp. (1838) 1 = intermedius.
commutatus, Guss. Fl. Sic. Prod. ii. 875 = intermedius.
complicatus, J. Gay, ex Gren. & Godr. Fl. Fr. i. 364 = intermedius.
decorticans, *Boiss. in Bibl. Univ. Genèv.* (Feb. 1836).—Hispan.
divaricatus, Sweet, Hort. Brit. ed. I. 112 = intermedius.
foliolosus, *DC. Fl. Fr. Suppl.* 549.—Ins. Canar.
frankenioides, Choisy, in DC. Prod. ii. 158 = anagyrus.
graecus, Griseb. Spicil. Fl. Rumel. i. 10 = intermedius.
grandiflorus, *Boiss. in Bibl. Univ. Genèv.* (Feb. 1836).—Gallia.
hispanicus, *DC. Fl. Fr. Suppl.* 549.—Hispan.
intermedius, *DC. l. c.*—Europ.; Oriens.

ADENOCARPUS :—
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 189. —Afr. trop.
parvifolius, DC. Fl. Fr. Suppl. 549 = intermedius.
speciosus, *Pomel, Nouv. Mat. Fl. Atl.* 178.—Afr. bor.
telonensis, DC. Fl. Fr. Suppl. 549 = grandiflorus?
umbellatus, *Coss. & Dur. in Ann. Sc. Nat.* Sér. IV. i. (1854) 222, nomen.—Afr. bor.
villosus, Boiss. Diagn. Ser I. ii. 14 = intermedius.
viscosus, Webb & Berth. Phyt. Canar. ii. 32 = anagyrus.
wolgensis, Spreng. Syst. iii. 226 = Calophaca wolgarica.

ADENOCAULON, Hook. Bot. Misc. i. 19. t. 15 (1830). *COMPOSITAE*, Benth. & Hook. f. ii. 344.
adhaerescens, Maxim. Prim. Fl. Amur. 152 = bicolor.
bicolor, *Hook. Bot. Misc.* i. (1830) 19. t. 15.—As. bor.; Himal.; Am. bor.
chilense, *Less. in Linnaea*, vi. (1831) 107.—Chili.
himalaicum, Edgew. in Trans. Linn. Soc. xx. (1851) 64 = bicolor.
integrifolium, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 289 = bicolor.
Lechleri, *Sch. Bip. in Flora*, xxxviii. (1855) 113.—Reg. Magell.

ADENOCHAENA, DC. Prod. vi. 680 (1837) = ADENACANTHA, DC. = **Phymospermum**, Less. (Compos.).

ADENOCHETON, Fenzl, in Flora, xxvii. (1844) 312 = **Coceulus**, Linn. (Menisperm.).
phyllanthoides, Fenzl, l. c. = C. Leaeba.

ADENOCHETUS, Baill. Etud. Gén. Euph. 653 (1858) = praec.

ADENOCHILUS, Hook. f. Fl. N. Zel. i. 246. t. 56 (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 613.
gracilis, *Hook. f. l. c.*—N. Zel.
Nortoni, *Fitzg. Austral. Orch.* i. II. 2.—Austral.

ADENOCHLAENA, Boiss. ex Baill. Etud. Gén. Euph. 472 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 307.
CENTROSTYLIS, Baill. Etud. Gén. Euph. 469 (1858).
SYMPHYLLIA, Baill. l. c. 473. t 11 (1858).
SYMPHYLLIUM, Benth. in Journ. Linn. Soc. xvii. (1878) 228.
calycina, *Bedd. For. Man. Bot.* 207.—Ind. or.
indica, *Bedd. Fl. Sylv.* ii. t. 261.—Ind. or.
leucocephala, Baill. Etud. Gén. Euph. 473 = Cephalocroton leucocephalus.
silhetensis, *Benth. in Journ. Linn. Soc.* xvii. (1880) 228.—Ind. or.
zeylanica, *Thw. Enum. Pl. Zeyl.* 270.—Zeylan.

ADENOCLINE, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 59. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 310.
DIPLOSTYLIS, Sond. in Linnaea, xxiii. (1850) 113.
PARADENOCLINE, Muell. Arg. in DC. Prod. xv. II. 1141 (1866).
acuta, *Baill. Etud. Gén. Euph.* 457.—Afr. austr.
humilis, *Turcz. in Bull. Soc. Nat. Mosc.* xvi. (1843) 61.—Afr. austr.
Mercurialis, *Turcz. l. c.* 60.—Afr. austr.
ovalifolia, *Turcz. l. c.*—Afr. austr.
pauciflora, *Turcz. l. c.* 61.—Afr. austr.
serrata, *Turcz. l. c.* xxv. (1852) II. 180.—Afr. austr.
sessiliflora, *Baill. Etud. Gén. Euph.* 457.—Afr. austr.
sessilifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xvi. (1843) 61.—Afr. austr.

ADENOCREPIS, Blume, Bijdr. 579 (1825) = **Baccaurea**, Lour. (Euphorb.).
javanica, Blume, l. c. = B. javanica.
lanceolata, Muell. Arg. in Linnaea, xxxii. (1863) 82 = B. lanceolata.
tetrandra, Baill. Etud. Gén. Euph. 601 = B. tetrandra.

ADENOCHAENA, Steud. Nom. ed. II. i. 25 (1840) sphalm. = **Adenachaena**, DC.

ADENOCYCLUS, Less. in Linnaea, iv. (1829) 337 = **Oliganthes**, Cass. (Compos.).
condensatus, Less. l. c. = O. condensata.

ADENODISCUS, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 504 = **Belotia**, A. Rich. (Tiliac.).
mexicanus, Turcz. l. c. = B. grewiaefolia.

ADENODUS, Lour. Fl. Cochinch. 294 (1790) = **Elaeocarpus**, Linn. (Tiliac.).
sylvestris, Lour. l. c. = E. sylvestris.

ADENOGRAMMA, Reichb. Iconogr. Exot. ii. 3. t. 109 (1827). *FICOIDEAE*, Benth. & Hook. f. i. 858.
STEUDELIA, Presl, Steud. Nov. Pl. Gen. (1829); et Symb. i. 3. t. 2 (1830).
diffusa, *Fenzl, in Ann. Wien. Mus.* ii. (1840) 275 (err. typ. 255).—Afr. austr.
galioides, *Fenzl, l. c.* 277.—Afr. austr.
lampocarpa, *E. Mey. ex Fenzl, in Ann. Wien. Mus.* ii. (1840) 276.—Afr. austr.
Mollugo, *Reichb. Iconogr. Exot.* ii. 3. t. 109.—Afr. austr.
oppositifolia, *Hassk. in Flora*, xxxi. (1851) 754.—Java.
physocalyx, *Fenzl, in Ann. Wien. Mus.* ii. (1840) 276.—Afr. austr.
rigida, *Sond. in Harv. & Sond. Fl. Cap.* 149.—Afr. austr.
sylvatica, *Fenzl, in Ann. Wien. Mus.* ii. (1840) 275 (err. typ. 255).—Afr. austr.

ADENOGYNA, *Rafin. New Fl. Am.* i. 62 (1836) = **Saxifraga**, Linn.
sarmentosa, Rafin. l. c. 63 = S. sarmentosa.

ADENOGYNE, Klotzsch, in Wiegm. Archiv, vii. (1841) 183 = **Sebastiania**, Spreng. (Euphorb.).
angustifolia, Klotzsch, l. c. 184 = S. angustifolia.
brachyclada, Klotzsch, l. c. = S. brachyclada.
discolor, Klotzsch, l. c. = Gymnanthes marginata.
guianensis, Klotzsch, in Schomb. Fauna & Fl. Guy. 1185 = S. guyanensis.
marginata, Klotzsch, in Wiegm. Archiv, vii. (1841) 184 = Gymnanthes marginata.
mucronata, Klotzsch, l. c. = S. Schottiana.
pachystachya, Klotzsch, l. c. = S. pachystachys.
rotundifolia, Klotzsch, l. c. = S. brevifolia.
serrata, Klotzsch, l. c. = S. serrata.

ADENOGYNE, Rafin. Fl. Tellur. ii. 67 (1836) = **Saxifraga**, Linn.

ADENOGYNUM, Reichb. f. & Zoll. in Verh. Natuurk. Ver. Nederl. Indie, i. (1856) 23; et in Linnaea, xxviii. (1856) 325 = **Chloradenia**, Baill. (Euphorb.) 312.
discolor, Reichb. f. & Zoll. ll. cc. = C. discolor.
odontophyllum, Miq. Fl. Ind. Bat. i. II. 400.—Java.

ADENOGYRUS, Klotzsch, in App. Ind. Sem. Hort. Berol. (1854) 1; et in Walp. Ann. iv. 226 (1857) = **Scolopia**, Schreb. (Bixin.).
Braunii, Klotzsch, ll. cc.—Hab.?

ADENOLEPIS, Less. in Linnaea, vi. (1831) 510 = **Cosmos**, Cav. (Compos.).
calva, Sch. Bip. in Miq. Fl. Ned. Ind. ii. 79 = Cosmos sulphureus.
pulchella, Less. in Linnaea, vi. 510 (1831).—Ins. Sandvic.

ADENOLINUM, Reichb. Handb. (1837) 306 = **Linum**, Linn.
alpicola, Reichb. l. c. 307 = L. alpinum.
alpinum, Reichb. l. c.; et Ic. Fl. Germ. xvi. 65 = L. alpinum.
angustifolium, Reichb. l. c. 307 = L. angustifolium.
austriacum, Reichb. l. c. 307; et Ic. Fl. Germ. xvi. 66 = L. austriacum.
barbatulum, Reichb. Ic. Fl. Germ. xvi. 73 = L. austriacum.
barbulatum, Schur, Enum. Pl. Transs. 127 = L. barbulatum.
Leonii, Reichb. Ic. Fl. Germ. xvi. 65 = L. Leonii.
marginatum, Reichb. Handb. 307; et Ic. Fl. Germ. xvi. 73 = L. austriacum.
montanum, Reichb. Ic. Fl. Germ. xvi. 65 = L. montanum.
nervosum, Reichb. Handb. 307 = L. nervosum.
pallescens, Reichb. l. c. = L. pallescens.

ADENOLINUM:—
perenne, Reichb. Ic. Fl. Germ. xvi. 65 = L. perenne.
provinciale, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 350 = L. alpinum.
pyrenaicum, Reichb. Handb. 307 = L. alpinum.
sibiricum, Reichb. l. c. 307 = L. sibiricum.
squamulosum, Reichb. Ic. Fl. Germ. xvi. 66 = L. squamulosum.
Tommasinii, Reichb. l. c.—Europ.

ADENONCOS, Blume, Bijdr. 381 (1825) = **Sarcochilus**, R. Br. ? (Orchid.).
virens, *Blume, l. c.*—Java.

ADENONEMA, Bunge, in Mém. Sav. Étrang. Pétersb. ii. (1836) 548; *reimpr.* Verz. Suppl. Fl. Alt. 36 = **Stellaria**, Linn. (Caryophyll.).
petraeum, Bunge, ll. cc. = S. petraea.

ADENOON, Dalz. in Hook. Kew Journ. ii. (1850) 344. *COMPOSITAE*, Benth. & Hook. f. ii. 226.
indicum, *Dalz. l. c.*—Ind. or.

ADENOPA, Rafin. Fl. Tellnr. iii. 37 (1836) = **Drosera**, Linn.
anglica, Rafin. l. c. = D. anglica.

ADENOPAPPUS, Benth. Pl. Hartw. 41 (1840). *COMPOSITAE*, Benth. & Hook f. ii. 411.
persicaefolius, *Benth. l. c.*—Mexico.

ADENOPELTIS, Bert. in Bull. Féruss. xxi. (1830), nomen; et ex A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 24. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 338.
Colliguaya, *Bert. l. c.*—Chili.

ADENOPETALUM, Klotzsch & Garcke, in Monatsb. Akad. Berl. 1859 (1860) 250; et Abh. Acad. Berl. (1860) 47 = **Euphorbia**, Linn.
boerhaaviifolium, Klotzsch & Garcke, ll. cc. 250, 47 = E. boerhavifolia.
bracteatum, Klotzsch & Garcke, ll. cc. 48 = E. scabrella.
discolor, Klotzsch & Garcke, ll. cc. 49 = E. discolor.
ellipticum, Klotzsch & Garcke, in Monatsb. Akad. Berl. 1859 (1860) 250 = E. graminea.
ellipticum, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 50 = E. saccharata.
gramineum, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 250; et in Abh. Akad. Berl. (1860) 47 = E. graminea.
Hoffmanni, Klotzsch & Garcke, ll. cc. 250, 47 = E. graminea.
irasuense, Klotzsch & Garcke, ll. cc. 250, 50 = E. graminea.
mexicanum, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 50 = E. graminea.
Oerstedii, Klotzsch & Garcke, in Monatsb. Akad. Berl. 1859) 250; et in Abh. Akad. Berl. (1860) 48 = E. graminea.
pictum, Klotzsch & Garcke, ll. cc. 250, 47.—Hab.?
pubescens, Klotzsch & Garcke, ll. cc. 250, 49 = E. graminea.
sphaerorhizum, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 250.—Hab.?
subsinuatum, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 250; et in Abh. Akad. Berl. (1860) 48 = E. graminea.

ADENOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxxi (1858) I. 416 = **Vitis**, Linn. (Ampel.).
palmatum, Turcz. l. c. 417.—Hab.?

ADENOPHAEDRA, Muell. Arg. in Mart. Fl. Bras. xi. II. 385. t. 101 (1873). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 314.
megalophylla, *Muell. Arg. l. c.* 386.—Bras.

ADENOPHORA, Fisch. in Mém. Soc. Nat. Mosc. vi. (1823) 165. *CAMPANULACEAE*, Benth. & Hook. f. ii. 563.
FLOERKEA, Spreng. Anleit. ii. II. 523 (1818).
communis, *Fisch. l. c.* 168.—Europ.
cordata, *Tausch. in Flora*, xxv. (1842) 288.—Hab.?
coronata, A. DC. Mon. Camp. 363 = marsupiiflora.
coronopifolia, *Fisch. in Mém. Soc. Nat. Mosc.* vi. (1823) 167.—As. bor.
crispata, Turcz. ex Ledeb. Fl. Ross. ii. 893 = coronata.

ADENOPHORA :—
denticulata, Fisch. in Mém. Soc. Nat. Mosc. vi. (1823) 167 = tricuspidata.
divaricata, *Franch. & Sav. Enum. Pl. Jap.* ii. 423.—Japon.
Fischeri, G. Don, in Loud. Hort. Brit. 75 = communis.
Gmelini, *Fisch. in Mém. Soc. Nat. Mosc.* vi. (1823) 167.—Dahur.
intermedia, Sweet, Hort. Brit. ed. I. 249 = communis.
intermedia, Ledeb. Ind. Sem. Hort. Dorpat. (1824) 1 = coronata.
Isabellae, *Hemsl. in Journ. Bot.* xiv. (1876) 207.—China.
Lamarkii, *Fisch. in Mém. Soc. Nat. Mosc.* vi. (1823) 168.—Sibir.
latifolia, *Fisch. l. c.*—Sibir.
liliflora, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 48 = communis.
lilifolia, Ledeb. Cat. Hort. Dorp. (1822); ex A. DC. Prod. vii. 492 = communis.
marsupiiflora, *Fisch. in Mém Soc. Nat. Mosc.* vi. (1823) 167.—Dahur.
montana, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) ii. 484.—Dahur.
nikoensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 423.—Japon.
pereskiaefolia, G. Don, in Loud. Hort. Brit. 74 = latifolia.
periplocaefolia, *A. DC. Mon. Camp.* 361.—Sibir.
Pichleri, Vis. Suppl. ex Nym. Consp. 479 = Campanula trichocalycina.
polymorpha, *Ledeb. Fl. Alt.* i. 246 in nota.—Rossia.
pomponiifolia, Fisch. in Mém. Soc. Nat. Mosc. vi. (1823) 167 = Gmelini.
Rabelaisiana, *G. Don, in Loud. Hort. Brit.* 75.—Sibir.
remotiflora, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 193.—Japon.
reticulata, *G. Don, in Loud. Hort. Brit. Suppl.* 471.—Sibir.
sinensis, *A. DC. Mon. Camp.* 354.—China.
stricta, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 192.—Japon.
stylosa, *Fisch. in Mém. Soc. Nat. Mosc.* vi. (1823) 168.—As. bor.
stylosa, Reichb. Iconogr. Exot. 32. t. 45 = communis.
suaveolens, Reichb. l. c. 22. t. 32 = communis.
tetraphylla, Fisch. in Mém. Soc. Nat. Mosc. vi. (1823) 169 = verticillata.
trachelioides, *Maxim. Prim. Fl. Amur.* 186.—China.
tricuspidata, A. DC. Mon. Camp. 355 = denticulata.
triphylla, *A. DC. l. c.* 365.—Japon.
verbenaefolia, *Bunge, ex Heynh. Nom.* ii. 9.—Japon.
verticillata, *Fisch. in Mém. Soc. Nat. Mosc.* vi. (1823) 167.—Dahur.; Japon.

ADENOPHYLLUM, Pers. Syn. ii. 458 (1807). *COMPOSITAE*, Benth. & Hook. f. ii. 408.
Dyssodia, Willd. Enum. Hort. Berol. 900 (1809).
Lebetina, Cass. in Dict. Sc. Nat. xxv. 394 (1822).
Schlechtendahlia, Willd. Sp. Pl. iii. 2125 (1803).
Willdenowa, Cav. Ic. Pl. i. 61. t. 89 (1791).
capillaceum, DC. Prod. v. 638 = coccineum.
coccineum, *Pers. Syn.* ii. 458.—Mexico.
Porophyllum, *Hemsl. Biol. Centr. Am. Bot.* ii. 218.—Mexico.
Wrightii, *A. Gray, Pl. Wright.* ii. 92.—Mexico.

ADENOPLEA, Radlk. in Bremen, Abh. viii. (1883) 406 = **Buddleia**, Houst. (Logan.).
baccata, Radlk. l. c. 407 = B. sinuata.
sinuata, Radlk. in Ber. Deutsch. Bot. Ges. ii. (1884) 261 = B. sinuata.

ADENOPLUSIA, Radlk. in Bremen Abh. viii. (1883) 461. *LOGANIACEAE.*
axillaris, *Radlk. l. c.* 462.—Madag.
Wildenowii, *Radlk. in Ber. Deutsch. Bot. Ges.* ii. (1884) 259.—Madag.

ADENOPODIA, C. Presl, Epimel. Bot. 206 (1849) = **Entada**, Adans. (Legum.).
spicata, C. Presl, l. c. 207 = E. natalensis.

ADENOPUS, Benth. in Hook. Niger Fl. 372 (1849). *CUCURBITACEAE*, Benth. & Hook. f. i. 823.
abyssinicus, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 528.—Afr. trop.

ADENOPUS :—
breviflorus, *Benth. in Hook. Niger Fl.* 372.—Afr. trop.
Cienkowskii, Schweinf. Rel. Kotschy. 45. t. 30 = Peponia Cienkowskii.
eglandulosus, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 529.—Afr. trop.
longiflorus, *Benth. in Hook. Niger Fl.* 372.—Afr. trop.

ADENORHOPIUM, Reichb. Consp. 195 (1828) = ADENOROPIUM, Pohl, = **Jatropha**, Linn. (Euphorb.).

ADENORIMA, Rafin. Fl. Tellur. iv. 112 (1836) = **Euphorbia**, Linn.
punicea, Rafin. l. c. = E. punicea.

ADENOROPIUM, Pohl, Pl. Bras. Ic. i. 12. t. 9 (1827) = **Jatropha**, Linn. (Euphorb.).
Berterii, Pohl, l. c. 14 = J. Berterii.
divaricatum, Pohl, l. c. 14 = J. divaricata.
divergens, Pohl, l. c. 15 = J. Pohliana.
elegans, Pohl, l. c. 15 = J. gossypifolia.
ellipticum, Pohl, l. c. 13. t. 9 = J. elliptica.
Forskohlii, Pohl, l. c. 15 = J. glandulosa.
glaucum, Pohl, l. c. 15 = J. glauca.
gossypifolium, Pohl, l. c. 16 = J. gossypifolia.
hernandiaefolium, Pohl, l. c. 14 = J. hernandiaefolia.
integerrimum, Pohl, l. c. 14 = J. integerrima.
Jacquini, Pohl, l. c. 15 = J. gossypifolia.
luxurians, Pohl, l. c. 16 = J. Pohliana.
Martiusii, Pohl, l. c. = J. Martiusii.
mollissimum, Pohl, l. c. 15 = J. Pohliana.
multifidum, Pohl, l. c. 16 = J. multifida.
mutabile, Pohl, l. c. 14 = J. mutabilis.
panduraefolium, Pohl, l. c. 14 = J. hastata.
peltatum, Pohl, l. c. 16 = J. peltata.
ribifolium, Pohl, l. c. 15 = J. gossypifolia.
Roxburghii, Kostel. Allg. Med.-pharm. Flora, v. 1750 = J. glandulifera.
spinosum, Pohl, Pl. Bras. Ic. i. 15 = J. spinosa.
tomentosum, Pohl, l. c. 16 = J. tomentosa.
tripartitum, Pohl, l. c. 15 = Manihot tripartita.
variegatum, Pohl, l. c. 14 = J. variegata.
villosum, Pohl, l. c. 15 = J. Pohliana.

ADENOSACHMA, A. Juss. in Orb. Dict. xi. 255 (1849) = seq.

ADENOSACME, Wall. Cat. n. 6280–82 (1832). *RUBIACEAE*, Benth. & Hook. f. ii. 69.
Lawia, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 14.
Mycetia, Reinw. in Syll. Ratisb. ii. (1828) 9.
cauliflora, *Miq. Fl. Ind. Bat.* ii. 215.—Malaya.
fasciculata, *Miq. l. c.* 216.—Java.
holotricha, *Miq. l. c.* 217.—Sumatra.
lanceolata, *Miq. l. c* 217.—Sumatra.
Lawii, *Hook. f. Fl. Brit. Ind.* iii. 96.—Ind. or.
longifolia, *Wall. Cat.* n. 6280.—Ind. or.
macrostachya, *Hook. f. Fl. Brit. Ind.* iii. 96.—Ind. or.
malayana, Wall. Cat. n. 6282 = longifolia.
nepalensis, Wall. l. c. n. 6281 = longifolia.
obovata, Miq. Fl. Ind. Bat. ii. 216 = fasciculata.
stipulata, *Hook. f. Fl. Brit. Ind.* iii. 95.—Ind. or.

ADENOSCILLA, Gren. & Godr. Fl. Fr. iii. 187 (1855) = **Scilla**, Linn. (Liliac.).
bifolia, Gren. & Godr. l. c. = S. bifolia.
nivalis, J. Gay, ex Baker, in Journ. Linn. Soc. xiii. (1873) 239 = S. bifolia.
unifolia, Texidor, in Rev. Prog. Cienc. xviii. (1869).—Cf. Bull. Soc. Bot. Fr. xvi. (1869); Bibl. 222 = S. bifolia?

ADENOSELEN, Spach. Hist. Vég. Phan. x. 23 (1841) = ADENOSOLEN, DC. = **Marasmodes**, DC. (Compos.).

ADENOSEPALUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 352 = **Hypericum**, Linn.
hirsutum, Fourr. l. c. 353 = H. hirsutum.
montanum, Fourr. l. c. = H. montanum.
nummularium, Fourr. l. c. = H. nummularium.
pulchrum, Fourr. l. c. 352 = H. pulchrum.
tomentosum, Fourr. l. c. = H. tomentosum.

ADENOSMA, R. Br. Prod. 442 (1810). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 949.
Anisanthera, Griff. Notul. iv. 100 (1854).
Pterostigma, Benth. Scroph. Ind. 20 (1835).
Spathestigma, Hook. & Arn. Bot. Beech. Voy. 204 (1836).
Brownii, *Zipp. ex Span. in Linnaea*, xv. (1841) 332, nomen.—Malaya.
caeruleum, *R. Br. Prod.* 443.—As. trop.; Austral.
camphoratum, *Hook. f. Fl. Brit. Ind.* iv. 264.—Zeylan.
capitatum, *Benth. ex Hance, in Journ. Linn. Soc.* xiii. (1873) 114.—As. trop.
chenopodiifolia, *Spreng. Syst.* ii. 829.—Ins. Societ.
chenopodioides, *Zipp. ex Span. in Linnaea*, xv. (1841) 332, nomen.—Hab.?
cuspidatum, Benth. in Wall. Cat. n. 3852 = Lindenbergia macrostachya.
fragrans, *Spreng. Syst.* ii. 829.—Ins. Societ.
grandiflorum, *Benth. ex Hance, in Journ. Linn. Soc.* xiii. (1874) 114.—China.
hirsutum, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 143.—Ins. Nicobar.
macrophyllum, *Benth. in Wall. Cat.* n. 3853.—Ind. or.
malabaricum, *Hook. f. Fl. Brit. Ind.* iv. 264.—Malabar.
microcephalum, *Hook. f. l. c.*—As. trop.
Muelleri, *Benth. Fl. Austral.* iv. 485.—Austral.
ovatum, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 263.—Malaya.
subrepens, *Benth. ex Hook. f. l. c.*—Zeylan.
triflora, Seem. Fl. Vit. 284 = Limnophila Roxburghii.
villosum, Benth. in Wall. Cat. n. 3851 = caeruleum.

ADENOSMA, Nees, in Wall. Pl. As. Rar. iii. 75 (1832) = **Cardanthera**, Buch.-Ham. (Acanthac.).
affinis, Griff. Notul. iv. 133 = Strobilanthes Acrocephalus.
africana, T. Anders. in Journ. Linn. Soc. vii. (1864) 21—Afr. trop.
balsamea, Spreng. Syst. ii. 829 = C. balsamica.
biplicata, Nees, in Wall. Pl. As. Rar. iii. 79 = C. avana.
elongata, Blume, Bijdr. 757 = Ebermaiera elongata.
Griffithii, T. Anders. in Journ. Linn. Soc. ix. (1867) 454 = C. Griffithii.
lanceolata, Blume, Bijdr. 757 = E. lanceolata.
pinnatifida, T. Anders. in Journ. Linn. Soc. ix. (1867) 455 = C. pinnatifida.
polysperma, Spreng. Syst. ii. 829 = Hygrophila Serpyllum.
spatulata, Blume, Bijdr. 737 = Ebermaiera glauca.
Thymus, Nees, in Wall. Pl. As. Rar. iii. 79 = C. balsamica.
triflora, Nees, l. c. = C. triflora.
Thwaitesii, T. Anders. in Thw. Enum. Pl. Zeyl. 224; et in Journ. Linn. Soc. ix. (1867) 455 = C. Thwaitesii.
uliginosa, Nees, in R. Br. Verm. Schrift. iii. 298 = C. uliginosa.
verticillata, Nees, in DC. Prod. xi. 69 = C. verticillata.

ADENOSOLEN, DC. Prod vi. 136 (1837) = **Marasmodes**, DC. (Compos.).
tenuifolius, DC. l. c.—Afr. austr.

ADENOSPERMUM, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 318 = **Chrysanthellum**, Rich. (Compos.).
tuberculatum, Hook. & Arn. l. c. = C. indicum.

ADENOSTEGIA, Benth. in Lindl. Syst. Nat. ed. II. 445 (1836) = **Cordylanthus**, Nutt. (Scrophul.).
rigida, Benth. l. c. = C. filifolius.

ADENOSTEMA, Desport. in Dict. Sc. Nat. i. 256 (1816) = seq.

ADENOSTEMMA, Forst. Char. Gen. 89. t. 45 (1776). *COMPOSITAE*, Benth. & Hook. f. ii. 239.
Lavenia, Sw. Prod. Veg. Ind. Occ. 112 (1788).
angustifolium, Arn. in Nov. Act. Nat. Cur. xviii. 1 (1836) 347 = viscosum.
aquaticum, D. Don, Prod. Fl. Nep. 81 = viscosum.
Berterii, *DC. Prod.* v. 110.—Ins. S. Doming.
biflorum, Less. Syn. Comp. 136 = Wedelia biflora.
brasilianum, *Cass. in Dict. Sc. Nat.* xxv. 363.—Bras.
caffrum, *DC. Prod.* v. 112.—Afr. austr.
decumbens, Don, ex Heynh. Nom. i. 19 = Beretrii.
Dregei, *DC. Prod.* v. 111.—Afr. austr.

ADENOSTEMMA :—
elatum, D. Don, Prod. Fl. Nep. 181 = viscosum.
erectum, DC. Prod. v. 113 = viscosum.
fastigiatum, DC. l. c. 111 = viscosum.
glandulosum, Benth. ex Hook. f. Fl Brit. Ind. i. 242, sphalm. (= Odontostemma) = Arenaria Benthami.
glutinosum, *DC. Prod.* v. 111.—Ins. Sandv.
Gymnocoronis, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 183 = Gymnocoronis spilanthoides.
hirsutum, *DC. Prod.* v. 113.—Java.
hirtiflorum, *Benth. Pl. Hartw.* 75.—Mexico.
lanceolatum, *Miq. Fl. Ind. Bat.* ii. 24.—Malaya.
latifolium, D. Don, Prod. Fl. Nep. 181 = viscosum.
leiocarpum, DC. in Wight, Contrib. 9 = viscosum.
macrophyllum, DC. Prod. v. 113 = viscosum.
madurense, DC. in Wight, Contrib. 9 = viscosum.
mauritianum, DC. Prod. v. 110 = viscosum.
microcephalum, DC. l. c. 111 = viscosum.
natalense, *DC. l. c.* 112.—Afr. austr.
nitidum, Spreng. Syst. ii. 357, lapsu = Gomortega nitida.
ovatum, *Miq. Fl. Ind. Bat.* ii. 25.—Malaya.
parviflorum, DC. Prod. v. 111 = viscosum.
Perrottetii, *DC. l. c.* 110.—Prom. Virid.
platyphyllum, *Cass. in Dict. Sc. Nat.* xxv. 363.—Peruv.
reticulatum, DC. in Wight, Contrib. 8 = viscosum.
rivale, Dalz. in Hook. Kew Journ. iii. (1851) 231 = viscosum.
Roylei, DC. Prod. v. 112 = viscosum.
rufescens, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 120, nomen.—Malaya.
rugosum, DC. in Deless. Ic. Sel. 4. t. 10 = viscosum.
scaevolifolium, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 186 = suffruticosum.
Schimperi, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 382.—Abyss.
suffruticosum, *Gardn. in Hook. Lond. Journ. Bot.* vi. (1867) 433.—Bras.
Swartzii, *Cass. in Dict. Sc. Nat.* xxv. 362.—Ind. occ.; Ecuador.
tinctorium, Cass. l. c. = viscosum.
triangulare, DC. Prod. v. 113 = viscosum.
viscosum, *Forst. Char. Gen.* 90.—Geront. trop.

ADENOSTEMON, Spreng. Anleit. ii. 788 (1818) = seq.

ADENOSTEMUM, Pers. Syn. i. 467 (1805) = **Gomortega**, Ruiz & Pav. (Laurin.?).
nitidum, Bert. Merc. Chil. 557; ex Am. Journ. Sc. xix. (1831) 67 = Cryptocarya Peumus.
nitidum, Pers. Syn. i. 467 = G. nitida.

ADENOSTEPHANUS, Klotzsch, in Linnaea, xv. (1841) 51 = **Euplassa**, Salisb. (Proteac.).
austro-caledonicus, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 229.—N. Caled.
bahiensis, Meissn. in Mart. Fl. Bras. v. I. 94. t. 36.—Bras.
Bleasdalii, Benth. Fl. Austral. v. 417.—Austral.
guyanensis, Meissn. in Mart. Fl. Bras. v. I. 95 = E. meridionalis.
incana, Klotzsch, in Linnaea, xv. (1841) 52.—Bras.
inaequalis, Endl. Gen. Suppl. iv. II. 87.—Bras.
laxiflora, Meissn. in Mart. Fl. Bras. v. I. 94.—Bras.
nitida, Meissn. l. c.—Bras.
organensis, Endl. Gen. Suppl. iv. II. 87.—Bras.
Sellowii, Klotzsch, in Linnaea, xv. (1841) 52.—Bras.

ADENOSTOMA, Blume, in Flora, viii. (1825) 680, nomen. *SCROPHULARINEAE.*

ADENOSTOMA, Hook. & Arn. Bot. Beech. Voy. 139, et 338. t. 30 (1832). *ROSACEAE*, Benth. & Hook. f. i. 613.
brevifolium, Nutt. ex Torr. & Gray, Fl. N. Am. i. 430 = fasciculatum.
fasciculatum, *Hook. & Arn. Bot. Beech. Voy.* 139, 338. t. 30.—Calif.
sparsifolium, *Torr. in Emory, Notes Mil. Reconnois.* 140; *Bot. Bound.* 63. t. 20.—Calif.; Mexico.

ADENOSTYLES, Cass. in Dict. Sc. Nat. i. Suppl. 59 (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 247.
ADENOSTYLIUM, Reichb. Ic. Fl. Germ. xvi. II. t. 895 (1853).

ADENOSTYLES :—
albida, *Cass. 'Bull. Soc. Philom.* 1816;' *Dict. Sc. Nat.* i. *Suppl.* 60.—Europ.
albifrons, Reichb. Fl. Germ. Excurs. 278 = albida.
Alliariae, *Kern. in Oestr. Bot. Zeitschr.* xxi. (1871) 12.—Europ.
alpina, Bluff & Fing. Comp. Fl. Germ. ii. 329 = viridis.
alpina, *Kern. in Oestr. Bot. Zeitschr.* xxi. (1871) 12.—Europ.
australis, Nym. Consp. 396 = viridis.
candidissima, *Cass. in Dict. Sc. Nat.* i. *Suppl.* 60.—Europ.
dubia, *Reichb. f. Ic. Fl. Germ.* xvi. II.—Europ.
glabra, DC. Prod. v. 203 = viridis.
hirsuta, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = albida.
hybrida, DC. Prod. v. 204 = alpina, candidissima, leucophylla, viridis.
leucophylla, DC. l. c. vii. 271 = orientalis.
leucophylla, Reichb. Fl. Germ. Excurs. 278 = candidissima.
macrophylla, Tenore, Syll. ex Nym. Consp. 396 = viridis.
Nardosmia, A. Gray, in Proc. Am. Acad. viii. (1873) 631 = Cacaliopsis Nardosmia.
nebrodensis, *Strobl, in Flora*, lxv. (1882) 196.—Sicil.
orientalis, *Boiss. Fl. Orient.* iii. 155.—Orient.
Petasites, Bluff & Fing. Comp. Fl. Germ. ii. 329 = albida.
pontica, *C. Koch, in Linnaea*, xiii. (1839) 696.—As. Min.
pyrenaica, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 64.—Hispan.
viridis, *Cass. in Dict. Sc. Nat.* i. *Suppl.* 60.—Europ.

ADENOSTYLES, Benth. & Hook. f. Gen. iii. 599 (1883) err. typ. = seq.

ADENOSTYLIS, Blume, Bijdr. 414 (1825) = **Zeuxine**, Lindl. (Orchid.).
emarginata, Blume, l. c. = Z. sulcata.
integerrima, Blume, l. c. = Z. integerrima.

ADENOSTYLIUM, Reichb. f. Ic. Fl. Germ. xvi. 2. t. 895 (1853) = **Adenostyles**, Cass. (Compos.).
dubium, Reichb. l. c. = Adenostyles dubia.
hybridum, Reichb. l. c. = Adenostyles viridis.

ADENOTHECA, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 255 = **Schizobasis**, Baker (Liliac.).
aphylla, Welw. ex Baker, l. c. = S. angolensis.

ADENOTRIAS, Jaub. & Spach, Illustr. Pl. Or. i. (1842) 76 = **Hypericum**, Linn.
Kotschyi, Jaub. & Spach, l. c. 77 (= *Triadenia Russeggeri*).—Syria.
phrygia, Jaub. & Spach, l. c. 76 = praec.

ADENOTRICHIA, Lindl. Bot. Reg. t. 1190 (1828) = **Senecio**, Linn. (Compos.).
amplexicaulis, Lindl. l. c. = S. adenotrichius.
senecioides, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 22 = S. sinuatilobus.

ADENUM, G. Don, Gen. Syst. iv. (1837) 80 = **Adenium**, Roem. & Schult. (Apocyn.).

ADESMIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 94. *LEGUMINOSAE*, Benth. & Hook. f. i. 517.
LOUDONIA, Bert. ex Hook. Bot. Misc. iii. (1833) 193.
PATAGONIUM, Schrank, in Muench. Denkschr. (1808) 93, nomen vetustius.
STREPTODESMA, A. Gray, in Proc. Am. Acad. iii. (1853) 6.
Aegiceras, *Phil. in Linnaea*, xxviii. (1856) 684.—Chili.
affinis, *Hook. f. Fl. Antarct.* 257.—Terr. Magell.
alpina, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
angulata, Hook. f. Fl. Antarct. ii. 257 = incana.
angustifolia, *Hook. & Arn. Bot. Beech. Voy.* 19.—Chili.
aphylla, *Clos, in C. Gay, Fl. Chil.* ii. 177.—Chili.
aprica, *Phil. in Anales Univ. Chil.* (1872) 697.—Chili.
arachnipes, *Clos, in C. Gay, Fl. Chil.* ii. 153.—Chili.
araucana, *Phil. in Linnaea*, xxviii. (1856) 633.—Chili.

ADESMIA :—
arborea, *Bert. Merc. Chil.* xii. 559; *cf. Bull. Féruss.* xx. (1830) 108.—Chili.
arborea, Colla, in Mem. Acc. Torin. xxxvii. (1834) 58 = glutinosa.
argentea, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
argentea, *Meyen, Reise*, i. 402, *et in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 22.—Peruv.
argyrophylla, *Phil. in Linnaea*, xxviii. (1856) 631.—Chili.
Arnottiana, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
arvensis, *Phil. in Linnaea*, xxxiii. (1864-65) 55.—Chili.
aspera, *Gill. in Hook. Bot. Misc.* iii. (1833) 190.—Chili.
atacamensis, *Phil. Fl. Atac.* 16.—Chili.
axillaris, *Phil. in Linnaea*, xxviii. (1856) 683.—Chili.
balsamica, *Bert. ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 59. t. 10.—Chili.
balsamifera, *Hook. & Arn. Bot. Beech. Voy.* 20.—Chili.
Berteroniana, Steud. Nom. ed. II. i. 27 = arborea.
bicolor, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Reg. Argent.
bijuga, *Phil. in Anales Univ. Chil.* (1884) reimpr. 3.—Chili.
boronioides, *Hook. f. Fl. Antarct.* 257.—Terr. Magell.
brachycarpa, *Phil. in Linnaea*, xxviii. (1856) 684.—Chili.
bracteata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 193.—Chili.
caespitosa, Gill. ex Hook. l. c. 191 = horrida.
calocarpa, *Phil. in Anales Univ. Chil.* (1865) II. 334.—Chili.
calycina, Vog. in Linnaea, xii. (1838) 75 = punctata.
candida, *Hook. f. Fl. Antarct.* 257.—Patag.
canescens, *Phil. in Anales Univ. Chil.* (1872) 703.—Chili.
capricornica, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = horrida.
Capricorni, *Phil. in Anales Univ. Chil.* (1862) II. 393; *et in Linnaea*, xxxiii. (1864-65) 58.—Chili.
Caragana, *Griseb. in Goett. Abh.* xix. (1874) 119.—Reg. Argent.
chillanensis, *Phil. in Anales Univ. Chil.* (1865) II. 326.—Chili.
ciliata, *Vog. in Linnaea*, xii. (1838) 74.—Bras.
cinerea, *Clos, in C. Gay, Fl. Chil.* ii. 179.—Chili.
Closii, *Phil. in Anales Univ. Chil.* (1872) 700.—Chili.
coluteoides, *Gill. in Hook. Bot. Misc.* iii. (1833) 190.—Reg. Argent.
coluteoides, *Phil. in Anales Univ. Chil.* (1872) 783.—Chili.
compacta, *Phil. in Linnaea*, xxviii. (1856) 628.—Chili.
conferta, *Hook. & Arn. Bot. Beech. Voy.* 20.—Chili.
coronilloides, *Gill. in Hook. Bot. Misc.* iii. (1833) 190.—Chili.
corymbosa, *Clos, in C. Gay, Fl. Chil.* ii. 165.—Chili.
cuneata, *Meyen, Reise*, i. 402, *et in Nov. Act. Cur.* xix. *Suppl.* i. (1843) 22.—Chili.
curvifolia, *Clos, in C. Gay, Fl. Chil.* ii. 189.—Chili.
cytisoides, *Griseb. in Goett. Abh.* xix. (1874) 119.—Reg. Argent.
Davilae, *Phil. in Anales Univ. Chil.* (1872) 699.—Chili.
decipiens, *Steud. Nom.* ed. II. i. 27.—Chili.
decumbens, *Phil. in Linnaea*, xxviii. (1856) 628.—Chili.
dendroidea, *Phil. l. c.* 683.—Chili.
dentata, *DC. in Ann. Sc. Nat.* Ser. I. iv. (1825) 95.—Chili.
dentata, Gill. ex Hook. Bot. Misc. iii. (1833) 189 = Gilliesii.
denticulata, *Clos, in C. Gay, Fl. Chil.* ii. 191.—Chili.
denticulata, *Steud. Nom.* ed. II. i. 27.—Chili.
denudata, *Phil. in Linnaea*, xxxiii. (1864-65) 56.—Chili.
descrticola, *Phil. in Anales Univ. Chil.* (1865) II. 344.—Chili.
dichotoma, *Clos, in C. Gay, Fl. Chil.* ii. 180.—Chili.
diffusa, *Bert. ex Steud. Nom.* ed. II. i. 27.—Chili.
Doniana, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
Echinus, *Presl, Symb. Bot.* ii. 14. t. 16.—Am. austr.
elata, *Clos, in C. Gay, Fl. Chil.* ii. 160.—Chili.
elegans, *Clos, l. c.* 188.—Chili.

ADESMIA:—
elongata, *Phil. in Linnaea*, xxviii. (1856) 633.—Chili.
emarginata, *Clos, in C. Gay, Fl. Chil.* ii. 187.—Chili.
eremophila, *Phil. Fl. Atac.* 15.—Chili.
erinacea, *Phil. l. c.* 16.—Chili.
exilis, *Clos, in C. Gay, Fl. Chil.* ii. 150.—Chili.
filicaulis, *Phil. in Anales Univ. Chil.* (1872) 702.—Chili.
filifolia, *Clos, in C. Gay, Fl. Chil.* ii. 157.—Chili.
filipes, *A. Gray, Bot. U. S. Expl. Exped.* i. 425.—Patag.
floribunda, *G. Don, Gen. Syst.* ii. 283.—Mexico.
frigida, *Phil. Fl. Atac.* 16.—Chili.
fruticulosa, *G. Don, Gen. Syst.* ii. 282.—Mexico.
Gayana, *Phil. in Linnaea*, xxviii. (1856) 634.—Am. austr.
genistoides, *Presl, Symb. Bot.* ii. 16.—Am. austr.
Germaini, *Phil. in Linnaea*, xxviii. (1856) 629.—Chili.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 188.—Chili.
glabriuscula, Vog. in Linnaea, x. (1836) 592 = radicans.
glandulosa, *Phil. in Anales Univ. Chil.* (1861) 47; *et in Linnaea*, xxxiii. (1864–65) 57.—Chili.
glauca, *Phil. in Linnaea*, xxviii. (1856) 682.—Chili.
glaucescens, *Phil. in Anales Univ. Chil.* (1872) 707.—Chili.
glomerata, *Auct. ex Phil. f. Cat. Pl. Chil.* 49 (sp. dub.).—Chili.
glomerula, *Clos, in C. Gay, Fl. Chil.* ii. 184.—Chili.
glutinosa, *Hook. & Arn. Bot. Beech. Voy.* 19.—Chili.
gracilis, *Meyen, ex Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 25.—Chili.
grandiflora, *Gill. in Hook. Bot. Misc.* iii. (1833) 190.—Chili.
graveolens, *Phil. Fl. Atac.* 16.—Chili.
grisea, *Hook. f. Fl. Antarct.* 257.—Patag.
hirsuta, *Phil. in Anales Univ. Chil.* (1872) 706.—Chili.
hispidula, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Peruv.
Hookeriana, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
horrida, *Gill. in Hook. l. c.*—Chili.
humifusa, *Phil. in Anales Univ. Chil.* (1872) 698.—Chili.
humilis, *Phil. in Linnaea*, xxviii. (1856) 682.—Am. austr.
Hystrix, *Phil. Fl. Atac.* 16.—Chili.
incana, *Vog. in Linnaea*, xii. (1838) 76.—Bras.
inconspicua, *Phil. l. c.* xxviii. (1856) 631.—Chili.
inflexa, *Griseb. in Goett. Abh.* xix. (1874) 121.—Reg. Argent.
Kingi, *Phil. in Anales Univ. Chil.* (1872) 706.—Chili.
Lambertiana, Gill. ex Hook. Bot. Misc. iii. (1833) 191 = trijuga.
lanata, *Hook. f. Fl. Antarct.* 256.—Terr. Magell.
Landbecki, *Phil. in Linnaea*, xxxiii. (1864–65) 59.—Chili.
latifolia, *Vog. l. c.* xii. (1838) 74.—Bras.
laxa, *Clos, in C. Gay, Fl. Chil.* ii. 162.—Chili.
leiocarpa, *Hook. & Arn, in Hook. Bot. Misc.* iii. (1833) 189.—Chili.
leptacantha, *Phil. in Anales Univ. Chil.* (1872) 705.—Chili.
longipes, *Phil. in Linnaea*, xxviii. (1856) 630.—Am. austr.
longiseta, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Chili.
lotoides, *Hook. f. Fl. Antarct.* 225.—Terr. Magell.
Loudonia, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 193.—Chili.
macrostachya, *Benth. in Mart. Fl. Bras.* xv. 1. 55.—Bras.
Meyeniana, *Phil. in Linnaea*, xxviii. (1856) 632.—Am. austr.
Meyeriana, Walp. Rep. i. 729 = Aeschynomene micrantha.
micrantha, *Phil. Fl. Atac.* 15.—Chili.
microphylla, *Hook. & Arn. Bot. Beech. Voy.* 19. t. 9.—Chili.
mimosoides, *G. Don, Gen. Syst.* ii. 282.—Mexico.
miraflorensis, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 357.—Bolivia.
monosperma, *Clos, in C. Gay, Fl. Chil.* ii. 184.—Chili.
monosperma, *Steud. Nom.* ed. II. i. 27.—Chili.
montana, *Phil. in Linnaea*, xxviii. (1856) 629.—Am. austr.

ADESMIA:—
mucronata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 189.—Chili.
multicuspis, *Clos, in C. Gay, Fl. Chil.* ii. 190.—Chili.
muricata, Bert. ex Steud. Nom. ed. II. i. 27 = decipiens.
muricata, *DC. in Ann Sc. Nat.* Sér. I. iv. (1825) 95.—Patag.; Chili.
muricata, Gill. ex Hook. Bot. Misc. iii. (1833) 188 = retrofracta.
obcordata, *Clos, in C. Gay, Fl. Chil.* ii. 199.—Chili.
obovata, *Clos, l. c.*—Chili.
obscura, *Clos, l. c.* 183.—Chili.
odontophylla, *Phil. in Anales Univ. Chil.* (1865) II. 344.—Chili.
oligophylla, *Phil. in Linnaea*, xxviii. (1856) 636.—Chili.
Onononis, *Griseb. in Lechl. Berb. Am. Austr.* 55, nomen.—Chili.
oresigena, *Phil. in Anales Univ. Chil.* (1872) 698.—Chili.
oxalidiphylla, *Phil. l. c.* (1865) II. 345.—Chili.
papposa, Bert. ex Steud. Nom. ed. II. i. 27 = angustifolia.
papposa, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Chili.
parviflora, *Clos, in C. Gay, Fl. Chil.* ii. 158.—Chili.
parvifolia, *Phil. in Linnaea*, xxviii. (1856) 683.—Am. austr.
parvula, *Phil. in Anales Univ. Chil.* (1872) 697.—Chili.
pauciflora, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 25.—Chili.
Pearcei, *Phil. in Anales Univ. Chil.* (1872) 704.—Chili.
pedicellata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 191.—Chili.
pendula, Bert. ex Steud. Nom. ed. II. i. 27 = denticulata, diffusa.
pendula, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Reg. Argent.
phylloidea, *Clos, in C. Gay, Fl. Chil.* ii. 186.—Chili.
pinifolia, *Gill. in Hook. Bot. Misc.* iii. (1833) 192.—Chili.
polyacantha, *Wedd. Chlor. And.* ii. 265.—Am. austr.
propinqua, *Clos, in C. Gay, Fl. Chil.* ii. 188.—Chili.
prostrata, *Clos, l. c.* 169.—Chili.
psilocarpa, *Phil. in Anales Univ. Chil.* (1865) II. 345.—Chili.
psoraleoides, *Vog. in Linnaea*, xii. (1838) 80.—Bras.
pugionata, *Griseb. in Goett. Abh.* xix. (1874) 120.—Reg. Argent.
pulchra, *Phil. in Linnaea*, xxviii. (1856) 632.—Am. austr.
pumila, *Hook. f. Fl. Antarct.* 255.—Patag.
punctata, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Am. austr.
pungens, *Clos, in C. Gay, Fl. Chil.* ii. 197.—Chili.
pusilla, *Phil. Fl. Atac.* 15.—Chili.
quadrijuga, *Phil. in Anales Univ. Chil.* (1872) 705.—Chili.
racemosa, Steud. Nom. ed. II. i. 27 = Aeschynomene micrantha.
radicans, *Bert. ex Steud. l. c.*—Chili.
radicans, *Clos, in C. Gay, Fl. Chil.* ii. 173.—Chili.
radicifolia, *Clos, l. c.* 169.—Chili.
ramosissima, *Phil. in Linnaea*, xxxiii. (1864–65) 56.—Am. austr.
retrofracta, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 188.—Chili.
retusa, *Griseb. in Goett. Abh.* vi. (1855) 119.—Am. austr.
rupicola, *Wedd. Chlor. And.* ii. 266.—Am. austr.
Sckickendanzii, *Griseb. in Goett. Abh.* xxiv. (1879) 104.—Reg. Argent.
Schneideri, *Phil. in Linnaea*, xxxiii. (1864–65) 60.—Am. austr.
sericea, *Griseb. in Lechl. Berb. Am. Austr.* 55, nomen.—Chili.
sessiliflora, *Phil. Fl. Atac.* 16.—Chili.
Simonsi, *Phil. in Anales Univ. Chil.* (1872) 701.—Chili.
Smithiae, Bert. ex Steud. Nom. ed. II. i. 27 = muricata.
Smithiae, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 95.—Am. austr.
spinosissima, *Meyen, Reise*, ii. 27; *et in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 24.—Chili.
stipulacea, *Clos, in C. Gay, Fl. Chil.* ii. 164.—Chili.
subterranea, *Clos, l. c.* 192.—Chili.
suffocata, *Hook. f. Fl. Antarct.* 256.—Terr. Magell.
tenella, *Hook. & Arn. Bot. Beech. Voy.* 19.—Chili.

ADESMIA:—
tomentosa, *Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 23.—Chili.
tribuloides, *Steud. Nom.* ed. II. i. 27.—Chili.
trifoliata, *Gill. in Hook. Bot. Misc.* iii. (1833) 194.—Chili.
trifoliata, *Phil. in Linnaea*, xxviii. (1856) 635.—Am. austr.
trijuga, *Gill. in Hook. Bot. Misc.* iii. (1833) 191.—Chili.
tristis, *Vog. in Linnaea*, xii. (1838) 77.—Bras.
ulicina, *Presl, Symb. Bot.* ii. 15.—Am. austr.
uspallatensis, *Gill. in Hook. Bot. Misc.* iii. (1833) 192.—Chili.
Valdesia, *Clos, in C. Gay, Fl. Chil.* ii. 174.—Chili.
verrucosa, *Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 20.—Chili.
vesicaria, *Bert. ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 58.—Chili.
villosa, *Hook. f. Fl. Antarct.* 256.—Reg. Magell.
virgata, *Bert. ex Steud. Nom.* ed. II. i. 27.—Chili.
viscida, Bert. Merc. Chil. xii. 557; Colla, in Mem. Acc. Torin. xxxvii. (1834) 59 = mucronata.
viscosa, *Gill. in Hook. Bot. Misc.* iii. (1833) 192.—Chili.
Volckmanni, *Phil. in Linnaea*, xxviii. (1856) 619.—Am. austr.
Wallichiana, Gill. ex Hook. Bot. Misc. iii. (1833) 190 = Aeschynomene aspera.

ADHADOTA, Steud. Nom. ed. II. i. 27 (1840) err. typ. = scq.

ADHATODA, Nees, in DC. Prod. xi. 384 (1847). Genus minime retinendum, species fere omnes ad *Justiciam* referendae. Vide infra.

ADHATODA, Tourn. ex Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 393; Nees, in Wall. Pl. As. Rar. iii. 102 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1112.
CARIMA, Rafin. Fl. Tellur. iv. 62 (1836).
DUVERNOYA, E. Mey. in Drège, Zwei Pfl. Docum. 150 (1843), nomen.
acuminata, *Nees, in DC. Prod.* xi. 400.—Abyss.
adenostachya, *Nees, l. c.* 408.—Bras.
amaranthoides, Nees, l. c. 400 = ceylanica.
anagalloides, *Nees, l. c.* 403.—Afr. austr.
Anselliana, Nees, l. c. 403 = Dianthera Anselliana.
arenaria, Nees, in Wall. Pl. As. Rar. iii. 103 = Justicia trinervia.
argyrostachya, Nees, l. c. 102 = Justicia argyrostachya.
aspera, *Nees, in DC. Prod.* xi. 397.—Ins. Philipp.
aurantiaca, *Regel, Ind. Sem. Hort. Petrop.* (1857) 47.—Hab.?
auriculata, *Nees, in DC. Prod.* xi. 407.—Bras.
bahiensis, *Nees, in Mart. Fl. Bras.* ix. 148.—Bras.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* 540.—Ind. or.
Betonica, Nees, in Wall. Pl. As. Rar. iii. 103 = Justicia Betonica.
boerhaaviaefolia, *Nees, in DC. Prod.* xi. 409.—Madag.
Bojeriana, *Nees, l. c.* 386.—Madag.
candicans, Nees, l. c. 396 = Dianthera candicans.
capensis, *Nees, l. c.*—Afr. austr.
carthagenensis, Nees, l. c. 403 = Beloperone violacea.
cheiranthifolia, *Nees, l. c.* 387.—Afr. austr.
ceylanica, Nees, l. c. 400 = Justicia ceylanica.
chinensis, *Benth. in Hook. Kew Journ.* v. (1853) 134.—Hongkong.
chrysotrichoma, *Nees, in DC. Prod.* xi. 408.—Bras.
ciliata, *Nees, l. c.* 396.—Bras.
congrua, *Nees, in Mart. Fl. Bras.* ix. 147.—Bras.
cuneata, *Nees, in DC. Prod.* xi. 392.—Afr. austr.
cuneifolia, *Nees, l. c.* 405.—Bras.
cydoniaefolia, *Nees, in Mart. Fl. Bras.* ix. 148.—Bras.
decussata, Nees, in DC. Prod. xi. 408 = Justicia decussata.
diffusa, Benth. in Hook. Niger Fl. 483 = Justicia insularis.
diosmophylla, *Nees, in DC. Prod.* xi. 394.—Afr. austr.
Dipteracanthus, Nees, l. c. 396 = Siphonoglossa pilosella.
divaricata, *Nees, l. c.* 391.—Afr. austr.
divergens, *Nees, l. c.* 406.—Am. merid.
diversiformis, Nees, l. c. 399 = Justicia glauca.
Eustachyana, Nees, l. c. 404 = Beloperone Eustachyana.
fasciata, *Nees, l. c.* 402.—Afr. austr.

ADHATODA:—
Felisbertiana, *Nees, in DC. Prod.* xi. 406.—Bras.
flava, *Nees, l. c.* 401.—Arab.
flexuosa, *Nees, in Mart. Fl. Bras.* ix. 148.—Bras.
formosissima, *Kotzsch, in Peters, Reise Mossamb. Bot.* 215.—Afr. trop.
furcata, Nees, in DC. Prod. xi. 398 = Justicia furcata.
genistiformis, *Nees, l. c.* 394.—Bras.
Gilliesii, *Nees, l. c.* 395.—Reg. Argent.
glandulosa, *Nees, l. c.* 405.—Bras.
glomerata, *Miq. Fl. Ind. Bat.* ii. 831.—Malaya.
holosericea, *Nees, in Mart. Fl. Bras.* ix. 148.—Bras.
Hookeriana, Nees, in DC. Prod. xi. 403 = Justicia Hookeriana.
Hypericum, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 102.—Afr. trop.
hyssopifolia, *Medic. in Act. Acad. Theod. Palat.* vi. *Phys.* (1790) 393 = Dicliptera ocymoides.
hyssopifolia, *Nees, in DC. Prod.* xi. 392.—Afr. austr.
incana, *Nees, l. c.* 393.—Afr. austr.
jalappaefolia, Buchoz, ex Steud. Nom. ed. I. 15 = Dicliptera ocymoides.
Kotschyi, *Nees, in DC. Prod.* xi. 397.—Afr. trop.
leptantha, *Nees, l. c.* 390.—Afr. austr.
leptostachya, *Nees, l. c.* 403.—Arab.
Lindeniana, *Nees, l. c.* 405.—N. Granat.
lithospermifolia, *Nees, l. c.* 398.—Peruv.
longiflora, Torr. in Bot. Mex. Bound. 125 = Siphonoglossa longiflora.
longifolia, *Nees, in DC. Prod.* xi. 404.—Ins. Mascar.
lupulina, *Nees, l. c.* 385.—Afr. austr.
Magnolia, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. 4. 6.—Malaya.
major, *Nees, in DC. Prod.* xi. 397.—Abyss.
Martiana, *Nees, l. c.* 405.—Bras.
matammensis, *Schweinf. in Verh. Zool. Bot. Ges. Wien*, xviii. (1868) 674.—Afr. trop.
microphylla, *Klotzsch, in Peters, Reise Mossamb. Bot.* 216.—Afr. trop.
minor, *Nees, in DC. Prod.* xi. 400.—Abyss.
mollissima, *Nees, l. c.* 391.—Afr. austr.
Moretiana, *Miq. Fl. Ind. Bat.* ii. 830.—Malaya.
Moritziana, *Nees, in DC. Prod.* xi. 388.—N. Granat.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 217.—Afr. trop.
natalensis, *Nees, in DC. Prod.* xi. 391.—Afr. austr.
Neesiana, Nees, l. c. 397 = Justicia Neesiana.
nilgerrhensis, Nees, in Wall. Pl. As. Rar. iii. 103 = Justicia nilgerrhensis.
nilgherrica, Nees, in DC. Prod. xi. 386 = Justicia nilgerrhensis.
nuda, *Nees, l. c.* 402.—Arab.
odora, *Nees, l. c.* 399.—Arab.
Orbigniana, *Nees, l. c.* 404.—Bolivia.
orchioides, *Nees, l. c.* 393.—Afr. austr.
origanoides, *Nees, l. c.* 407.—Ind. or.
orixensis, Nees, l. c. 400 = Justicia glauca.
pacifica, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 164 = Justicia pacifica.
palustris, *Nees, in DC. Prod.* xi. 402.—Afr. trop.
paniculata, *Benth. in Hook. Niger Fl.* 482.—Afr. trop.
patula, *Nees, in DC. Prod.* xi. 393.—Afr. austr.
periplocifolia, *Nees, l. c.* 395.—Venezuela.
petiolaris, *Nees, l. c.* 402.—Afr. trop.
pilosa, *Nees, l. c.* 405.—Bras.
plicata, *Nees, l. c.* 401.—Afr. trop.
protracta, *Nees, l. c.* 390.—Afr. austr.
pubescens, Mocuch, Meth. 431 = Gendarussa Adhatoda.
pygmaea, *Nees, in DC. Prod.* xi. 394.—Afr. austr.
quadrifaria, Nees, l. c. 396 = Justicia quadrifaria.
ramosissima, Nees, in Wall. Pl. As. Rar. iii. 103 = Justicia Betonica.
reflexiflora, *Nees, in DC. Prod.* xi. 398.—Ind. occ.
rostellaria, Nees, l. c. 397 = Justicia neglecta.
rostrata, Hochst. ex Oliver, in Trans. Linn. Soc. xxix. (1872) 131 = Asystasia Schimperi.
rostrata, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 104.—Afr. trop.
rotundifolia, *Nees, in DC. Prod.* xi. 391.—Afr. austr.
scabra, *Nees, l. c.* 407.—Venezuela.
Schimperiana, *Hochst. ex Nees, l. c.* 388.—Abyss.
sericea, Nees, fide Benth. & Hook. f. Gen. ii. 1115 = Jacobinia mexicana.
sisparensis, Benth. in Journ. Linn. Soc. ix. (1867) 518 = Rungia sisparensis.
sphaerosperma, *Nees, in DC. Prod.* xi. 409.—Ind. occ.

ADHATODA:—
spicata, Burm. ex Steud. Nom. ed. I. 15, 274 = Rungia repens.
spicata, *Nees, in DC. Prod.* xi. 408.—Malaya.
striata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 216.—Afr. trop.
suaveolens, *Nees, in DC. Prod.* xi. 401.—Arab.
subserrata, *Nees, l. c.* 409.—Ins. Philipp.
sulcata, *Nees, l. c.* 401.—Arab.
tetramera, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 301.—Porto-Rico.
thymifolia, *Nees, in DC. Prod.* xi. 392.—Afr. austr.
tranquebariensis, Nees, l. c. 399 = Justicia tranquebariensis.
trinervia, Nees, in Wall. Pl. As. Rar. iii. 103 = Justicia trinervia.
tristiflora, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 250.—Java.
tristis, *Nees, in DC. Prod.* xi. 404.—Afr. trop.
tubulosa, *Nees, l. c.* 392.—Afr. austr.
Tweediana, *Nees, l. c.* 395.—Bras.
umbrosa, *Nees, l. c.* 406.—Bras.
variegata, *Nees, l. c.* 385.—Afr. austr.
vasculosa, Nees, l. c. 407 = Justicia vasculosa.
Vasica, *Nees, in Wall. Pl. As. Rar.* iii. 103.—As. trop.
Velasquezii, *Nees, in DC. Prod.* xi. 409.—Guatemala.
ventricosa, Nees, l. c. 407 = Justicia ventricosa.
verticillata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 6.—Java.
wynaadensis, Nees, in DC. Prod. xi. 406 = Justicia wynaadensis.
Zollingeriana, Nees, l. c. xi. 396 = Justicia Zollingeriana.

ADHUNIA, Vell. Fl. Flum. 236 (1825); v. t. 118 (1827). —'Gen. dub.'
emarginata, Vell. ll. cc.—Bras.

ADIANTHUM *truncatum*, Burm. f. Fl. Ind. 235. t. 66. f. 4 = Acacia decipiens.

ADIANTUM *truncatum*, Murr. Syst. Veg. 790 = Acacia decipiens.

ADICEA, Rafin. Cat. 13 (1824) = **Urtica**, Tourn.
glabra, Rafin. l. c. = U. glabra.
pumila, Rafin. ex Torr. Fl. New York, ii. 223 = Pilea pumila.
trinervia, Rafin. ex Torr. l. c. = Pilea pumila.

ADIKE, Rafin. New Fl. Am. i. 63 (1836); Fl. Tellur. iii. 48 (1836) = **Pilea**, Lindl. (Urtic.).
allophyla, Rafin. Fl. Tellur. iii. 48 = P. trianthemoides.
glaberrima, Rafin. New Fl. Am. i. 64 = P. smilacifolia.
lanceolata, Rafin. l. c. = P. lanceolata.
pumila, Rafin. l. c. 63 = P. pumila.
rhomboidea, Rafin. l. c. 64 = P. rhombea.

ADINA, Salisb. Parad. Lond. t. 115 (1807). *RUBIACEAE*, Benth. & Hook. f. ii. 30.
aralioides, *Benth. & Hook. f. Gen.* ii. 30.—Java.
cordifolia, *Benth. & Hook. f. l. c.*—Ind. or.
globiflora, *Salisb. Parad. Lond.* t. 115 (errore *globifera*). —China.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 24.—Ind. or.
microcephala, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 40. —Afr. trop.
peduncularis, DC. Prod. iv. 349 = globiflora.
polycephala, *Benth. Fl. Hongk.* 146.—Ind. or.
racemosa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 184.—China; Japon.
rubella, *Hance, in Journ. Bot.* vi. (1868) 114.—China.
sessilifolia, *Benth. & Hook. f. Gen.* ii. 30.—Ind. or.
trichotoma, *Benth. & Hook. f. l. c.*—Malaya.

ADINANDRA, Jack, in Malay Misc. ii. n. 7. p. 49 (1822). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 182.
HAENKAEA, Usteri, Ann. Bot. vi. (1793) 117.
SAROSANTHERA, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 103. t. 16 (1840?).
acuminata, *Korth. l. c. ed. Temminck*, 109.—Sumatra.
cordata, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 114.
crenulata, *T. Anders. in Hook. f. Brit. Ind. Fl.* i. 283. —Ind. or.
cyrtopoda, Miq. Fl. Ind. Bat. Suppl. i. 478 = dumosa.

ADINANDRA:—
dasyantha, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 180 = integerrima.
dasyantha, *Korth. Verh. Nat. Gesch. Bot. ed. Temminck*, 108.—Borneo.
dumosa, *Jack, in Malay Misc.* ii. (1822) n. 7. p. 50. —Malaya; China.
dumosa, Vidal, Sinops. Filip. t. 12 = integerrima.
excelsa, *Korth. Verh. Nat. Gesch. Bot. ed. Temminck*, 109.—Borneo.
glabra, Miq. Fl. Ind. Bat. Snppl. i. 479 = dumosa.
Griffithii, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 282.—Reg. Himal.
integerrima, *T. Anders, ex Dyer, l. c.*—Malaya.
Jackiana, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 106 = dumosa.
javanica, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 143, 146.—Java.
Lamponga, *Miq. Fl. Ind. Bat. Suppl.* i. 479.—Sumatra.
lasiopetala, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 114.—Zeylan.
leiopetala, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 104.—Sumatra.
macrantha, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 421.—Java.
maculosa, *T. Anders. ex Dyer, in Hook. f. Fl. Brit. Ind.* i. 282.—Malaya.
Mannii, *Oliver, Fl. Trop. Afr.* i. 170.—Afr. trop.
Millettii, *Benth. & Hook. f. ex Hance, in Journ. Bot.* xvi. (1878) 9.—China.
phlebophylla, *Hance, l. c.* xiv. (1876) 240.—Cambodia.
Sarosanthera, *Miq. Fl. Ind. Bat.* i. ii. 477.—Borneo.
stylosa, Miq. l. c. Suppl. 478 = dumosa.
sylvestris, *Jack, in Malay Misc.* ii. (1822) n. 7 p. iii. —Sumatra.
trichocoryna, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 107 = dumosa.
villosa, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855). —Burma.

ADIPE, Rafin. Fl. Tellur. ii. 101 (1836) = **Bifrenaria**, Lindl. (Orchid.) (1843).
fulva, Rafin. l. c. = B. racemosa.
racemosa, Rafin. l. c. = B. racemosa.

ADIPERA, Rafin. Sylva Tellur. 129 (1838) = **Cassia**, Tourn. (Legum.).
Herberti, Rafin. l. c. = C. Herbertiana.

ADISA, Steud. Nom. ed. II. i. 28 (1840), sphalm. = seq.

ADISCA, Blume, Bijdr. (1825) 609 = **Mallotus**, Lour. (Euphorb.).
acuminata, Blume, l. c. 610 = M. acuminatus.
albicans, Blume, l. c. 611 = Cephalocroton albicans.
floribunda, Blume, l. c. 610 = M. floribundus.
subfalcata, Zoll. in Linnaea, xxix. (1857–58) 464 = Macaranga subfalcata.
subpeltata, Blume, Bijdr. 610 = M. subpeltatus.
timoriana, Span. Ic. n. 26; et in Linnaea, xv. (1841) 348 = M. repandus.
Zippelii, Blnme, Bijdr. 611 = M. ricinoides.

ADLERIA, Neck. Elem. ii. 642 (1790) = **Eperua**, Aubl. (Legum.).

ADLUMIA, Rafin. in Med. Repos. New York, v. (1808) 352. *PAPAVERACEAE*, Benth. & Hook. f. i. 55.
BICUCULLA, Borckh. in Roem. Arch. i. ii. 46 (1797), nomen prius.
cirrhosa, *Rafin. l. c.*—Am. bor.

ADNARIA, Rafin. Fl. Ludov. 56 (1817). *VACCINIACEAE*.
odorata, *Rafin. l. c.*—Am. bor.

ADNULA, Rafin. Fl. Tellur. ii. 87 (1836) = **Pelexia**, Poit. (Orchid.).
petiolaris, Rafin. l. c. = P. spiranthoides.

ADODENDRON, DC. Prod. vii. 725 = seq.

ADODENDRUM, Neck. Elem. i. 214 (1790) = **Rhododendron**, Reichb. (Ericac.).

ADOKETON, Rafin. Mus. N. Sc. 1; New Fl. Am. i. 65 (1836) = **Telanthera**, R. Br. (Chenop.).
saxatile, Rafin. ll. cc. = T. frutescens.
uniflorum, Rafin. New Fl. Am. i. 65.—Am. bor.

ADOLIA, Lam. Encyc. i. (1783) 44 = **Scutia**, Comm. (sec. Endl. Gen. Suppl. i. 1424) (Rhamnac.).
alba, Lam. l. c. 45 = S. indica.
rubra, Lam. l. c. = S. indica.

ADOLPHIA, Meissn. Gen. 70 (50) (1837). *RHAMNEAE*, Benth. & Hook. f. i. 384.
californica, *S. Wats. in Proc. Am. Acad.* xi. (1876) 126.—Calif.
infesta, *Meissn. Gen. Pl.* 70 (50).—Mexico.

ADONANTHE, Spach, Hist. Vég. Phan. vii. 227 (1839) = **Adonis**, Linn. (Ranunc.).
pyrenaica, Spach, l. c. 230 = Adonis pyrenaica.
sibirica, Spach, l. c. 229 = Adonis sibirica.
vernalis, Spach, l. c. 228 = Adonis vernalis.

ADONIASTRUM, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 27 = **Adonis**, Dill. (Ranunc.).
vernale, Schur, l. c. = Adonis vernalis.

ADONIGERON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 404 = **Senecio**, Linn. (Compos.).
adonidifolius, Fourr. l. c. = S. adonidifolius.

ADONIS, Dill. ex Linn. Syst. ed. I. (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 5.
ADONANTHE, Spach, Hist. Vég. Phan. vii. 227 (1839).
ADONIASTRUM, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 27.
COSMARIUM, Dulac, Fl. Hautes-Pyr. 215 (1867).
aestivalis, Bieb. Fl. Taur. Cauc. iii. 378 = autumnalis.
aestivalis, *Linn. Sp. Pl.* ed. II. 771.—Europ.; Oriens.
aestivalis, Link, ex Webb & Berth. Phyt. Canar. i. 12 = microcarpa.
aethiopica, Thunb. Prod. Pl. Cap. 94 = Knowltonia gracilis.
aleppica, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 350.—Syria.
ambigua, Gaud. Fl. Helv. iii. 518 = aestivalis.
amurensis, *Regel & Radde, in Bull. Soc. Nat. Mosc.* (1861) II. 35. t. 2. f. 12.—Manchur.
annua, Linn. Sp. Pl. 547 = aestivalis et autumnalis.
anomala, Wallr. Sched. Crit. 273 = flammea.
apennina, Linn. Sp. Pl. 548 = vernalis.
apennina, Pall. Reise, ii. 388 = wolgensis.
apennina, Schouw, ex DC. Prod. i. 25 = pyrenaica.
autumnalis, Bieb. Fl. Taur. Cauc. ii. 23 = aestivalis.
autumnalis, *Linn. Sp. Pl.* ed. II. 771.—Europ.; Oriens.
baetica, Coss. Not. Pl. Crit. ii. 25 = autumnalis.
caerulea, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 306.—China.
capensis, Linn. Sp. Pl. 548 = Knowltonia rigida.
capensis, Thunb. Prod. Pl. Cap. 94 = Knowltonia gracilis.
caudata, Stev. in Bull. Soc. Nat. Mosc. xxi. (1848) II. 271, 273 = flammea.
chaerophylla, Fisch. in DC. Prod. i. 25 = wolgensis.
Chrysocyathus, *Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind.* 15.—Reg. Himal.
citrina, DC. Syst. i. 223 = flammea.
citrina, Hoffm. Fl. Germ. i. 251 = aestivalis.
coriacea, Poir. Encyc. Suppl. i. 146 = Knowltonia rigida.
Cupaniana, Guss. Syn. Fl. Sic. ii. 36 = microcarpa.
cyllenea, *Boiss. Heldr. & Orph. Diagn.* Sér. II. v. 5.—Graecia.
daucifolia, Lam. Encyc. i. 46 = Knowltonia daucifolia.
davurica, Reichb. Ic. Crit. Cent. iv. 21. t. 321 = vernalis.
dentata, *Delile, Fl. Egypte*, 287.—Europ.; Afr. bor.
dentata, Duby, ex Nym. Consp. 4 = aestivalis.
distorta, *Tenore, Fl. Nap.* iv. 337. t. 149.—Ital.
eriocalycina, *Boiss. Fl. Orient.* i. 17.—Armenia.
erosipetala, Stokes, Bot. Mat. Med. iii. 255 = autumnalis.
filia, Linn. f. Suppl. 271 = Knowltonia daucifolia.
flammea, *Jacq. Fl. Austr.* iv. 29. t. 355.—Europ.; Oriens.
flammea, Schleich. ex Steud. Nom. ed. II. i. 28 = aestivalis.
flava, Vill. Cat. Jard. Strasb. 247 = aestivalis.

ADONIS:—
fulgens, *Hochst. in Lorent, Wanderungen*, 338.—As. Min.
gracilis, Poir. Encyc. Suppl. i. 147 = Knowltonia gracilis.
Helleborus, Crantz, Stirp. Austr. ed. I. fasc. ii. 82; ed. II. 110 = vernalis.
hirsuta, Poir. Encyc. Snppl. i. 147 = Knowltonia hirsuta.
hybrida, Wolff, ex Nym. Consp. 4 = wolgensis.
Inglisii, Royle, Illustr. Himal. Pl. 53 = aestivalis.
intermedia, Webb & Berth. Phyt. Canar. i. 121 = microcarpa.
ircutiana, Fisch. ex DC. Prod. i. 25 = vernalis.
laserpitiifolia, Poir. Encyc. Suppl. i. 147 = Knowltonia vesicatoria.
maculata, *Roth, Enum. Pl. Phan. Germ.* ii. 602.—Europ.
maculata, Wallr. Sched. Crit. 270 = aestivalis.
magellanica, Raeusch, Nom. ed. III. 162 = Hamadryas magellanica.
marginata, Bienert, ex Hook. f. Fl. Brit. Ind. i. 15 = scrobiculata.
Marschalliana, Andrz. ex Bess. Enum. Pl. Volh. 47 = wolgensis.
micrantha, DC. Syst. i. 222 = autumnalis.
microcarpa, *DC. l. c.*—Europ.
miniata, Jacq. Fl. Austr. iv. 28 = aestivalis.
palaestina, *Boiss. Diagn.* Ser. I. viii. 1.—Syria.
palustris, Raeusch, Nom. ed. III. 162 = Ranunculus sceleratus.
parviflora, *Fisch. ex DC. Prod.* i. 24.—Oriens.
parviflora, Janka, ex Nym. Consp. 4 = vernalis.
perramosa, Schur, Enum. Pl. Transs. 6 = autumnalis.
persica, Boiss. Diagn. Ser. I. vi. 4 = dentata.
phoenicea, Bercht. et Presl, Rostl. i. Ranunculac. 42 = autumnalis.
pilosa, Raeusch, Nom. ed. III. 162 = Ranunculus pensylvanicus.
polymorpha, Zumag. Fl. Pcdem. ii. 362 = aestivalis, autumnalis.
pratensis, Ledeb. Fl. Ross. i. 24 = vernalis.
Preslii, Tod. ex Nym. Consp. 4 = flammea.
pyrenaica, *DC. Fl. Fr.* v. 635.—Europ.
pyrenaica, Hook. f. & Thoms. Fl. Ind. 26 = A. Chrysocyathus.
riparia, *Rafin. New Fl. Am.* i. 66.—Am. bor.
scrobiculata, *Boiss. Diagn.* Ser. II. i. 6.—Afghan.
sibirica, *Patrin, ex Ledeb. Ind. Hort. Dorp. Suppl.* (1824) 1.—Sibir. altaic.
Sibthorpii, *Boiss. Orphan. & Heldr. in Bull. Congr. Bot. Pétersb.* 1869 (1870) 124.—Graecia.
squarrosa, *Stev. in Bull. Soc. Nat. Mosc.* xxi. (1848) II. 271, 272 = aestivalis.
vernalis, Asso, Syn. Stirp. Arag. 70 = dentata.
vernalis, *Linn. Sp. Pl.* 547.—Europ.; Graecia.
vesicatoria, Linn. f. Suppl. 272 = Knowltonia vesicatoria.
villosa, Ledeb. Ind. Hort. Dorp. Suppl. (1824) 1 = wolgensis.
wolgensis, *Stev. ex DC. Syst.* i. 545.—Europ.; Oriens; As. bor.

ADOPOGON, Neck. Elem. i. 55 (1790) = **Krigia**, Schreb. (Compos.).

ADORIUM, Rafin. Neogenyt. 3 (1825) = **Seseli**, Linn. (Umbellif.).
crassifolium, Rafin. l. c. = S. divaricatum.

ADOXA, Linn. Syst. ed. I. (1735). *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 2.
MOSCATELLA, Adans. Fam. ii. 243 (1763).
MOSCHATELLINA, [Tourn.] Hall. Enum. Stirp. Helv. i. 412 (1742).
inodora, Falc. ex Clarke, in Hook. f. Fl. Brit. Ind. iii. 2 = A. Moschatellina.
moschata, Dulac, Fl. Hautes-Pyr. 462 = A. Moschatellina.
Moschatellina, *Linn. Sp. Pl.* 367.—Europ.; As. bor.; Am. bor.
tuberosa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 493 = A. Moschatellina.

ADRASTAEA, DC. Syst. Veg. i. 424 (1818). *DILLENIACEAE*, Benth. & Hook. f. i. 15.
ADRASTEA, Spreng. Syst. ii. 360 (1825).
salicifolia, *DC. l. c.*—Austral.

ADRIANA, Gaudich. in Ann. Ser. Nat. Ser. I. v. (1825) 223. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 306.
TRACHYCARYON, Klotzsch, in Lehm. Pl. Preiss. i. 175 (1844).
acerifolia, Hook. in Mitch. Journ. Trop. Austral. 371 = glabrata.
Billardieri, Baill. Etud. Gén. Euph. Atl. 6. t. 2. f. 19 = quadripartita.
Gaudichaudi, Baill. Adansonia, vi. (1865-66) 312 = tomentosa.
glabrata, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 223.—Austral.
heterophylla, Hook. in Mitch. Journ. Trop. Austral. 371 = glabrata.
Hookeri, *Muell. Arg. in DC. Prod.* xv. II. 891.—Austral.
Klotzschii, *Muell. Arg. l. c.* 892.—Austral.
quadripartita, *Gaudich. Bot. Voy. Freyc.* 489.—Austral.
tomentosa, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 223.—Austral.

ADROMISCHUS, Lem. Jard. Fleur. ii. (1852) Misc. 59 = **Cotyledon**, Tourn. (Crassul.).
caryophyllaceus, Lem. l. c. 60 = C. caryophyllacea.
clavifolius, Lem. l. c. = C. clavifolia.
cristatus, Lem. l. c. = cristata.
cuneatus, Lem. l. c. = cuneata.
hemisphaericus, Lem. l. c. = C. hemisphaerica.
jasminiflorus, Lem. l. c.—C. jasminiflora.
maculatus, Lem. l. c. = C. maculata.
mammillaris, Lem. l. c. = C. mammillaris.
mucronatus, Lem. l. c. = C. maculata.
rhombifolius, Lem. l. c. = C. rhombifolia.
robustus, Lem. l. c. = C. triflora.

ADULPA, Endl. Gen. (1836) 119 (err. typ.) = seq.

ADUPLA. Bosc, in Jaume St. Hil. Expos. Fam. Nat. i. 65 (1805) = **Cyperus**, Linn.

ADVENTINA, Rafin. New Fl. Am. i. 67 (1836). *COMPOSITAE*.
ciliata, Rafin. l. c.—Am. bor.
parviflora, Rafin. l. c.—Am. bor.

ADYSETON, Adans. Fam. ii. 420 (1763) = **Alyssum**, Linn. (Crucif.).
alpestre, Sweet, Hort. Brit. ed. I. 467 = Alyss. alpestre.
altaicum, G. Don, in Sweet, l. c. ed. III. 30 = Alyss. Fischerianum.
argenteum, Sweet, l. c. ed. I. 467 = Alyss. argenteum.
atlanticum, G. Don, Gen. Syst. i. 178 = Alyss. atlanticum.
Bertolonii, Sweet, Hort. Brit. ed. I. 467 = Alyss. montanum.
calycinum, Scop. Fl. Carn. ed. II. ii. 13 = Alyss. calycinum.
campestre, Moench, Meth. Suppl. 87 = Alyss. campestre.
cuneifolium, Sweet, Hort. Brit. ed. I. 467 = Alyss. cuneifolium.
dentatum, G. Don, Gen. Syst. i. 179 = Camelina sativa.
diffusum, Sweet, Hort. Brit. ed. I. 467 = Alyss. montanum.
dubium, G. Don, Gen. Syst. i. 179 = Alyss. dubium.
edentulum, Link, Enum. Hort. Berol. ii. 157 = Alyss. gemonense.
Fischerianum, Sweet, Hort. Brit. ed. II. 25 = Alyss. canescens.
gemonense, Sweet, l. c. = Alyss. gemonense.
gnaphalodes, G. Don, Gen. Syst. i. 179 = Vesicaria gnaphalodes.
halimifolium, Link, Enum. Hort. Berol. ii. 157 = Alyss. halimifolium.
hyperboreum, G. Don, Gen. Syst. i. 179 = Alyss. hyberboreum.
lanigerum, G. Don, l. c. 178 = Alyss. lanigerum.
lenense, G. Don, l. c. = Alyss. lenense.
maritimum, Link, Enum. Hort. Berol. ii. 157 = Alyss. maritimum.
Marschallianum, Sweet, Hort. Brit. ed. II. 25 = Alyss. Marschallianum.
montanum, Scop. Fl. Carn. ed. II. ii. 14 = Alyss. Wulfenianum.
murale, Sweet, Hort. Brit. ed. I. 467 = Alyss. argenteum.
mutabile, Moench, Meth. 267 = Alyss. calycinum.

ADYSETON:—
nebrodense, Sweet, Hort. Brit. ed. II. 25 = Alyss. serpyllifolium.
obcordatum, G. Don, Gen. Syst. i. 179 = Abyss. obcordatum.
obtusifolium, Sweet, Hort. Brit. ed I. 467 = Alyss. alpestre.
orientale, Sweet, l. c. 467 = Alyss. orientale.
petalodes, G. Don, Gen. Syst. i. 179 = Farsetia obovata.
repens, Sweet, Hort. Brit. ed. II. 25 = Alyss. alpestre.
savranicum, G. Don, Gen. Syst. i. 178 = Alyss. alpestre.
saxatile, Sweet, Hort. Brit. ed. I. 467 = Alyss. saxatile.
serpyllifolium, Sweet, l. c. = Alyss. serpyllifolium.
sibiricum, G. Don, Gen. Syst. i. 179 = Alyss. saxatile.
spathulatum, Sweet, Hort. Brit. ed. I. 467 = Alyss. spathulatum.
spinosum, Link, Enum. Hort. Berol. ii. 157 = Alyss. spinosum.
tortuosum, Sweet, Hort. Brit. ed. I. 467 = Alyss. alpestre.
vernale, Sweet, l. c. = Alyss. rostratum.
Wulfenianum, Sweet, l. c. = Alyss. montanum.

ADYSETUM, Link, Enum. Hort. Berol. ii. 157 (1822) = praec.

AECEOCLADES, Duchartre, in Orb. Dict. ix. 170 (1849) err. typ. = OECEOCLADES, Lindl.

AECHMAEA, Brongn. in Ann. Sc. Nat. Sér. II. xv. (1841) 371 = **Aechmea**, Ruiz & Pav. (Bromel.).

AECHMANDRA, Arn. in Hook. Lond. Journ. Bot. iii. (1841) 274 = **Melothria**, Linn. (Cucurb.).
Blumeana, M. Roem. Syn. Pepon. 33 = M. Rumphiana.
conocarpa, Dalz. & Gibs. Bomb. Fl. 100 = Corallocarpus conocarpus.
deltoidea, Arn. in Hook. Lond. Journ. Bot. iii. (1841) 274 = M. zeylanica.
epigaea, Arn. l. c. = Corallocarpus epigaeus.
indica, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 626 = M. indica.
odorata, Hook. f. &. Thoms. ex Benth. Fl. Hongkong, 124 = M. odorata.
rostrata, Arn. in Hook. Lond. Journ. Bot. iii. (1841) 274 = Kedrostis rostrata.
velutina, Dalz. & Gibs. in Bomb. Fl. 200 = Corallocarpus velutina.
zeylanica, Thw. Enum. Pl. Zeyl. 125 = Cerasiocarpum zeylanicum.

AECHMANTHERA, Nees, in Wall. Pl. As. Rar. iii. 75, 87 (1837). *ACANTHACEAE*, Benth. & Hook f. ii. 1088.
gossypina, Nees, l. c. 87 = tomentosa.
leiosperma, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 429.—Ind. or.
tomentosa, *Nees, in Wall. Pl. As. Rar.* iii. 87.—Ind. or.
Wallichii, Nees, in DC. Prod. xi. 170 = tomentosa.

AECHMEA, Ruiz & Pav. Prod. 47. t. 8 (1794). *BROMELIACEAE*, Benth. & Hook. f. iii. 663.
Androlepis, Brongn. ex Houllet, in Rev. Hortic. (1870) 12.
Canistrum, E. Morr. Belg. Hortic. (1873) 257. t. 15.
Disquamia, Lem. Jard. Fleur. iii. (1852–3) t. 269.
Echinostachys, Brongn. in Planch. Hort. Donat. 25 (1854).
Eriostax, Rafin. Fl. Tellur. iv. 25 (1836).
Hohenbergia, Schult. f. Syst. vii. II. 71, 1251 (1830).
Hoplophytum, E. Morr. Belg. Hortic. (1865) 162.
Lamprococcus, Beer, Bromel. 103 (1857).
Macrochordion, De Vriese, in Jaarb. Nederl. Maat. Tuinb. (1853) 14.
Pironneava, Gaudich. Voy. Bonite, Bot. t. 63. (1839–52).
Pothuava, Gaudich. l. c. tt. 116, 117 (1839–52).
angustifolia, Pocpp. & Endl. Nov. Gen. et Sp. ii. 43 = spicata.
aquilega, *Griseb. Fl. Brit. W. Ind.* 592.—Ind. occ.
augusta, *Baker, in Journ. Bot.* xvii. (1879) 162.—Am. trop.
aurantiaca, *Baker, l. c.* 235.—Am. trop.
Barlcei, *Baker, in Gard. Chron.* xx. (1883) 102.—Honduras.
bracteata, *Griseb. Fl. Brit. W. Ind.* 592.—Ind. occ.
brasiliensis, *Regel, Gartenfl.* (1885) 258.—Bras.
brassicoides, *Baker, in Journ. Bot.* xx. (1882) 329.—Brit. Guian.

AECHMEA:—
bromeliaefolia, *Baker, ex Benth. & Hook. f. Gen.* iii. 664.—Am. trop.
Burchellii, *Baker, in Journ. Bot.* xvii. (1879) 231.—Am. trop.
caerulea, E. Morr. Cat. (1871) 1 = seq.
caerulescens, *Hort. ex Baker, in Journ. Bot.* xvii. (1879) 227.—Am. trop.
calyculata, *Baker, in Journ. Bot.* xvii. (1879) 232.—Am. trop.
capitata, Baker, l. c. 167 = exsudans.
coelestis, *E. Morr. Fl. des Serres*, Sér. II. xi. (1875) l. t. 2146.—Am. trop.
columnaris, *André, in Illustr. Hortic.* xxv. (1878) 50.—N. Granat.
comata, *Baker, in Journ. Bot.* xvii. (1879) 234.—Am. trop.
contracta, *Baker, l. c.*—Am. trop.
corallina, *Brongn. ex Beer, Bromel.* 106.—Bras.
Cornui, Carr. in Rev. Hortic. (1885) 36 = nudicaulis.
Cumingii, *Baker, in Journ. Bot.* xvii. (1879) 227.—N. Granat.
cymoso-paniculata, *Baker, l. c.* 165.—Venezuela.
dactylina, *Baker, l. c.* 161.—Panama.
dichlamydea, *Baker, l. c.* 133.—Ins. Tobago.
discolor, Morr. in Ann. Soc. Hort. Gand, ii. (1846) 175. t. 65 = fulgens.
distans, *Griseb. Fl. Brit. W. Ind.* 592.—Ind. occ.
distans, Griseb. Pl. Cub. 253 = Wrightii.
distichantha, *Lem. Jard. Fleur.* iii. (1853) t. 269.—Bras.
exsudans, *E. Morr. ex C. H. Morr. in Belg. Hortic.* xxviii. (1878) 303.—Bras.
excavata, *Baker, in Journ. Bot.* xvii. (1879) 134.—Paraguay.
fasciata, *Baker, l. c.* 231.—Am. trop.
floribunda, *Mart. ex Schult. f. Syst.* vii. II. 1271.—Bras.
fulgens, *Brongn. in Ann. Sc. Nat.* Sér. II. xv. (1841) 371.—Guiana.
Furstenbergii, E. Morr. ex C. H. Morr. in Belg. Hortic. xxviii. (1878) 252 = Streptocalyx Furstenbergi.
Glaziovii, *Baker, in Journ. Bot.* xvii. (1879) 133.—Bras.
glomerata, *Hook. f. Bot. Mag.* t. 5668.—Bras.
Hookeri, *Lem. Illustr. Hortic.* xi. (1864) *Misc.* 42.—Bras.
Hystrix, *E. Morr. ex C. H. Morr. in Belg. Hortic.* xxx. (1880) 90 & 243.—Guiana.
Jenmani, Baker, in Journ. Bot. xx. (1882) 329 = Melinonii.
Lalindei, *Linden & Rodigas, in Illustr. Hortic.* xxx. (1883) t. 481.—N. Granat.
laxiflora, *Benth. in Bot. Voy. Sulph.* 173.—Mexico.
Legrelliana, *Baker, in Journ. Bot.* xvii. (1879) 236.—Am. merid.
Lindeni, *E. Morr. ex C. Koch, Wochenschr.* viii. (1865) 398.—Bras.
lingulata, *Baker, in Journ. Bot.* xvii. (1879) 164.—Jamaic.
Luddemanniana, Brongn. ex E. Morr. Cat. Bromel. (1871) 1 = caerulescens.
macracantha, Brongn. in Illustr. Hortic. (1880) 59 = Schiedeana.
Mariae-Reginae, *H. Wendl. in Hamb. Gartenz.* xix. (1863) 32.—Am. centr.
martinicensis, *Baker, in Journ. Bot.* xvii. (1879) 132.—Ins. Martinic.
melanacantha, *De Vriese, in Tuinb. Fl.* i. (1855) 356.—Hab.?
Melinonii, *Hook. Bot. Mag.* t. 5235.—Am. trop.
Mertensii, *Schult. f. Syst.* vii. II. 1272.—Am. austr.
mexicana, *Baker, in Journ. Bot.* xvii. (1879) 165.—Mexico.
Meyeri, Baker, l. c. xviii. (1880) 16 = aquilega.
micrantha, Brongn. in Ann. Sc. Nat. Sér. II. xv. (1841) 370 = Aracococcus micranthus.
mucroniflora, *Hook. Bot. Mag.* t. 4832.—Guiana.
multiceps, *Baker, in Journ. Bot.* xviii. (1880) 49.—Am. trop.
Noettigii, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 117.—Bras.
nudicaulis, *Griseb. Fl. Brit. W. Ind.* 593.—Am. trop.
odora, *Baker, in Journ. Bot.* xvii. (1879) 226.—Am. trop.
organensis, Wawra, in Oestr. Bot. Zeitschr. xxx. (1880) 116 = floribunda.
ornata, *Baker, in Journ. Bot.* xvii. (1879) 162.—Am. trop.
Ortgiesii, *Baker, l. c.* 236.—Am. trop.

AECHMEA:—
paniculata, *Ruiz & Pav. Fl. Per.* iii. 37.—Peruv.
paniculigera, *Griseb. Fl. Brit. W. Ind.* 593.—Ind. occ.
parviflora, *Baker, in Journ. Bot.* xvii. (1879) 167.—Bras.
patentissima, *Baker, l. c.* 227.—Am. trop.
pectinata, *Baker, l. c.* 233.—Am. trop.
petropolitana, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 116.—Bras.
Pineliana, *Baker, in Journ. Bot.* xvii. (1879) 232.—Am. trop.
platynema, *Baker, l. c.* 166.—Bras.
polycephala, *Baker, l. c.* 164.—Ins. Jamaic.
pubescens, *Baker, l. c.* 135.—Nicarag.
pyramidalis, *Benth. in Bot. Voy. Sulph.* 173.—Mexico.
ramosa, *Mart. ex Schult. f. Syst.* vii. II. 1272.—Bras.
regularis, *Baker, in Journ. Bot.* xvii. (1879) 229.—Am. trop.
Schiedeana, *Schlecht. in Linnaea*, xviii. (1844) 437.—Mexico.
setigera, *Mart. ex Schult. f. Syst.* vii. II. 1273.—Bras.
spectabilis, *Brongn. ex Houllet, in Rev. Hortic.* (1875) 311.—Guatemala.
sphaerocephala, *Baker, in Journ. Bot.* xvii. (1879) 162.—Am. trop.
spicata, *Mart. ex Schult. f. Syst.* vii. II. 1273.—Bras.
suaveolens, *Knowl. & Westc. Flor. Cab.* iii. (1840) 177.—Bras.
subinermis, *Baker, in Journ. Bot.* xvii. (1879) 228.—Am. trop.
tillandsioides, *Baker, l. c.* 134.—Bras.
Veitchii, *Baker, in Bot. Mag.* t. 6329.—N. Granat.
viridis, *Baker, in Journ. Bot.* xvii. (1879) 235.—Am. austr.
vriesioides, Baker, l. c. 134 = tillandsioides.
Weilbachea, Carr. in Rev. Hortic. (1871) 171 = seq.
Weilbachii, *F. Didr. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 375.—Bras.
Wrightii, *Baker, in Journ. Bot.* xvii. (1879) 163.—Cuba.

AECHMOLEPIS, Deene. in DC. Prod. viii. (1844) 493. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 741.
rosmarinifolia, *Decne. l. c.*—Afr. trop.

AECHMOPHORA, Spreng. ex Steud. Nom. ed. II. i. (1840) 29 = **Bromus**, Linn. Gramin.
arduensis, Spreng. ex Steud. l. c. = B. arduensis.

AEDEMONE, Kotschy, in Oestr. Bot. Zeitschr. viii. (1858) 116 = **Herminiera**, Guill. & Perr. (Legumin.).
mirabilis, Kotschy, l. c. = H. Elaphroxylon.

AEDMANNIA, Spach, in Orb. Dict. i. (1841) 139, err. typ. = OEDMANNIA, Thunb. (Legum.).

AEDULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. 4. 1, nomen.

AEEGIPHILA, Sw. Prod. Veg. Ind. Occ. (1788) 31, err. typ. = **Aegiphila**, Jacq.

AEËRIDIUM, Salisb. in Trans. Hort. Soc. i. (1812) 295 = **Aërides**, Lour. (Orchid.).
odorum, Salisb. l. c. = Aërides odoratum.

AEGELATIS, Roxb. Fl. Ind. ii. 111 (1832) err. typ. = **Aegialitis**, R. Br. (Plumbag.).

AEGIALEA, Klotzsch, in Walp. Ann. ii. (1852) 1113 = **Pieris**, D. Don (Ericac.).

AEGIALINA, Schult. Mant. ii. (1824) 13, 222 = **Koeleria**, Pers. (Gramin.).
tenuis, Schult. l. c. = K. villosa.

AEGIALINITES, Presl, Bot. Bemerk. (1844) 103 = **Aegialitis**, R. Br. (Plumbag.).
annulata, Presl, l. c. = Aegialitis annulata.
rotundifolia, Presl, l. c. = Aegialitis rotundifolia.

AEGIALITIS, R. Br. Prod. (1810) 426. *PLUMBAGINEAE*, Benth. & Hook. f. ii. 624.
Aegialinites, Presl, Bot. Bemerk. (1844) 103.
annulata, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 217 = rotundifolia.
annulata, *R. Br. Prod.* i. 426.—Austral.
rotundifolia, *Roxb. Fl. Ind.* ii. 111.—Ind. or.

AEGIALITIS, Trin. Fund. Agrost. 127. t. 9. (1820) = **Koeleria**, Pers. (Gramin.).
tenuis, Trin. l. c. = K. villosa.

AEGIALOPHILA, Boiss. & Heldr. Diagn. Ser. I. x. (1849) 105 = **Centaurea**, Linn. (Compos.).
cretica, Boiss. & Heldr. l. c. 106 = C. pumila.
pumila, Boiss. l. c. 105 = C. pumila.

AEGIANILITES, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 479 (1882), err. typ. = **Aegialitis**, R. Br.

AEGIATILIS, Griff. Not. iv. 207 (1854), err. typ. = praec.

AEGICERAS, Gaertn. Fruct. i. 216. t. 46 (1788). *MYRSINEAE*, Benth. & Hook. f. ii. 648.
MALASPINAEA, Presl, Reliq. Haenk. ii. 68. t. 61 (1830).
corniculatum, Blanco, Fl. Filip. ed. I. 79 = majus.
ferreum, Blume, Bijdr. 693 = majus.
floridum, Roem. & Schult. Syst. iv. 512 = majus.
fragrans, Kon. in Kon. & Sims, Ann. Bot. i. (1805) 131 = majus.
majus, *Gaertn. Fruct.* i. 216. t. 46.—Geront. trop.
Malaspinaea, A. DC. Prod. viii. 143 = majus.
minus, A. DC. l. c. 412 = majus.
minus, *Gaertn. Fruct.* i. 216. t. 46.—Ind. or.
nigricans, A. Rich. Voy. Astrol. Bot. ii. 57. t. 21 = majus.
obcordatum, Steud. Nom. ed. II. i. 29 = majus.
obovatum, Blume, Bijdr. 693 = majus.

AEGICON, Adans. Fam. ii. 36 (1763) = **Triticum**, Linn. (Gramin.).

AEGILOPS, Linn. Coroll. Gen. 20 (1737) = **Triticum**, Linn. (Gramin.).
agropiroides, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 456.—Syria.
aromatica, Walt. Fl. Carol. 249 = Ctenium americanum.
Aucheri, Boiss. Diagn. Ser. I. v. 74 = Agropyron Aucheri.
Augeri, Steud. Syn. Pl. Gram. 355 (err. typ.) = praec.
bicornis, Jaub. & Spach, Illustr. Pl. Or. iv. 11 = T. bicorne.
biuncialis, Vis. Fl. Dalm. 1.—Europ.
brachyathera, Pomel, Nouv. Mat. Fl. Atl. 389.—Afr. bor.
caudata, Bertol. Fl. Ital. vi. 622 = Agropyron Aucheri.
caudata, C. A. Mey. Verz. Pfl. Cauc. 26 = T. Aegylops.
caudata, Balb. Elencho, 98 ; Griseb. Spicil. Fl. Rumel. ii. 425 = T. cylindricum.
caudata, Linn. Sp. Pl. 1051 = Agropyron villosum ?
ciliaris, Koen. ex Roem. & Schult. Syst. ii. 772.—Ind. or.
comosa, Sibth. & Sm. Fl. Graec. 75. t. 94.—Graecia.
connata, Steud. Syn. Pl. Gram. 356.—Graecia.
crassa, Boiss. Diagn. Ser. I. vii. 123.—Oriens.
crithodium, Steud. Syn. Pl. Gram. 355 = T. monococcum.
cylindrica, C. A. Mey. Verz. Pfl. Cauc. 26 = T. Aegylops ?
cylindrica, Host, Gram. Austr. ii. 6. t. 7 = T. cylindricum.
cylindrica, Schur, Enum. Pl. Transs. 813 = T. ovatum.
cylindrica, Sibth. & Sm. Fl. Graec. t. 95 = T. caudatum.
divaricata, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 129 = T. ovatum.
echinata, Presl, Gram. Sic. 47 = T. triunciale.
Echinus, Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 456.—Gallia ?
elongata, Lam. Fl. Fr. iii. 632 = T. triunciale.
erigens, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 131 = T. ovatum.
erratica, Jord. & Fourr. l. c. 130 = T. ovatum.
exaltata, Linn. Mant. ii. App. 575 = Ophiurus corymbosus.
fluviatilis, Blanco, Fl. Filip. ed. I. 47.—Ins. Philipp.
fragilis, Parl. Fl. Ital. i. 515 = T. fragile.
geniculata, Roth, Catalect. i. 121 = T. ovatum.
hordeiformis, Steud. Syn. Pl. Gram. 354.—Mesopotam.
Hystrix, Nutt. Gen. Am. i. 86.—Am. bor.
incurva, Linn. Sp. Pl. 1051 = Lepturus incurvatus.
incurvata, Linn. Sp. Pl. ed. II. 1490 = praec.
Kotschyi, Boiss. Diagn. Ser. I. vii. 129 = T. triunciale.
Larenti, Steud. Syn. Pl. Gram. 354 = *Lorentii*.
loliacea, Jaub. & Spach, Illustr. Pl. Or. iv. 23 = *mutica*.

AEGILOPS :—
longearistata, Steud. Syn. Pl. Gram. 356.—Guiana.
Lorentii, Hochst. in Lorent, Wanderungen, 326 = T. ovatum.
macrochaeta, Shuttl. & Huet, ex Duval, in Bull. Soc. Bot. Fr. xvi. (1869) 384 = T. ovatum.
macrura, Jaub. & Spach, Illustr. Pl. Or. iv. 21 = Agropyron Aucheri.
microstachys, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 132 = T. ovatum.
muricata, Retz. Obs. ii. 27 = Ischaemum pectinatum.
mutica, Boiss. Diagn. Ser. I. v. 73.—Oriens.
neglecta, Req. ex Bertol. Fl. Ital. i. 787 = T. triaristatum.
nigricans, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 128 = T. ovatum.
Notarisii, Clem. in Mem. Acc. Torin. Ser. II. xvi. (1857) 335 = T. ovatum.
ovata, Linn. Sp. Pl. 1050 = T. ovatum.
ovata, Roth, in Usteri, Ann. iv. (1793) 41 = T. triaristatum.
parvula, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 131 = T. ovatum.
persica, Boiss. Diagn. Ser. I. vii. 129 = T. caudatum.
platyathera, Jaub. & Spach, Illustr. Pl. Or. iv. 17.—Mesopotam.
procera, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 129 = T. ovatum.
pubiglumis, Jord. & Fourr. l. c. 131 = T. ovatum.
saccharina, Walt. Fl. Carol. 249 = Eleusine aegyptiaca.
singularis, Steud. Syn. Pl. Gram. 354.—Persia.
sicula, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 129 = T. ovatum.
speltaeformis, Jord. in Ann. Sc. Nat. Sér. IV. iv. (1855) 313 = T. ovatum.
speltoides, Tausch, in Flora, xxxix. (1837) 109 = T. bicorne.
squarrosa, Willd. Sp. Pl. iv. 944 = T. ventricosum.
squarrosa, Cav. Ic. t. 90. f. 2 = *platyathera*.
squarrosa, DC. ex Steud. Nom. ed. II. i. 29 = T. cylindricum.
squarrosa, Linn. Sp. Pl. 1051 = T. Aegylops.
subulata, Pomel, Nouv. Mat. Fl. Atl. 388.—Afr. bor.
Tauschii, Coss. Notes Crit. 69 = T. Aegylops.
triaristata, Req. ex Bertol. Fl. Ital. i. 789 = T. triunciale.
triaristata, Willd. Sp. Pl. iv. 943 = T. triaristatum.
tripsacoides, Jaub. & Spach, Illustr. Pl. Or. ii. 121. t. 200 = *mutica*.
triticoides ×, Req. ex Bertol. Fl. Ital. i. 788.—Europ.
triuncialis, Linn. Sp. Pl. 1051 = T. triunciale.
truncata, Ledeb. ex Trautv. in Act. Hort. Petrop. ix. (1884) 312, nomen.—Rossia.
uniaristata, Steud. Syn. Pl. Gram. 354.—Graecia.
uniaristata, Vis. Fl. Dalm. iii. 345.—Europ.
vagans, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 130 = T. ovatum.
ventricosa, Tausch, in Flora, xxxix. (1837) 108 = T. ventricosum.
virescens, Jord. & Fourr. Brev. Pl. Nov. Fasc. ii. 130 = T. ovatum.

AEGINETIA, Cav. in Anal. Cienc. Nat. iii. (1801) 129 = **Bouvardia**, Salisb. (Rubiac.).
capitata, R. Grah. in Edinb. N. Phil. Journ. (Jan. 1828) 389 = Coccocipselum capituliflorum.
longiflora, Cav. in Anal. Hist. Nat. iii. (1801) 130 = Houstonia longiflora.
multiflora, Cav. l. c. = Bouvardia Cavanillesii.

AEGINETIA, Linn. Syst. ed. I. (1735). *OROBANCHACEAE*, Benth. & Hook. f. ii. 982.
CENTRONIA, Blume, Bijdr. 776 (1826).
CENTRONOTA, DC. ex Meissn. Gen. 303 (213) (1840) ; et Prod. ix. 286 (1846).
GASPARINIA, Endl. Gen. 1408 (1840).
TRONICENA, Steud. Nom. ed. II. ii. 721 (1841).
abbreviata, Buch.-Ham. ex Benth. Scroph. Ind. 55 = pedunculata.
acaulis, Walp. Rep. iii. 481 = pedunculata.
Centronia, *Miq. Fl. Ind. Bat.* ii. 713.—Java.
coccinea, Juss. ex Walp. Rep. iii. 481 = Phelypaea coccinea.
indica, *Linn. Sp. Pl.* 632 ; *Roxb. Pl. Corom.* i. 63. t. 91.—As. trop.
japonica, Sieb. & Zucc. in Abh. Math. Akad. Muench. iv. III. (1845) 341 = indica.
pedunculata, *Wall. Pl. As. Rar.* iii. 13. t. 219.—As. trop.

AEGIPHILA, Jacq. Stirp. Am. t. 16 (1763) ; Obs. Bot. ii. 3. t. 27 (1767). *VERBENACEAE*, Benth. & Hook. f. ii. 1151.
AMERINA, DC. in Meissn. Gen. 278 (186) (1840).
BRUECKIA, Klotzsch & Karst. Ausw. Gew. Venez. 31. t. 10 (1848).
MANABEA, Aubl. Pl. Gui. i. 61. t. 23, 25 (1775).
OMPHALOCOCCA, Willd. ex Roem. & Schult. Syst. iii. Mant. 10 (1827).
arborescens, *Vahl, Eclog.* i. 15. t. 10.—Am. austr.
aurea, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1868) II. 218.—Cuba.
Berteriana, *Schau. in DC. Prod.* xi. 654.—N. Granat.
brachiata, *Schlecht. in Linnaea*, vi. (1831) 371.—Mexico.
brachiata, Vell. Fl. Flum. 38 ; i. t. 93 = triantha.
casseliaeformis, *Schau. in DC. Prod.* xi. 651.—Bras.
cestrifolia, *Gardn. in Hook. Lond. Journ. Bot* v. (1846) 184.—Bras.
cordata, *Poepp. ex DC. Prod.* xi. 650.—Peruv.
cornifolia, *Kunth, in Abh. Akad. Berl.* (1831) 215.—Bras.
cuspidata, *Mart. ex DC. Prod.* xi. 653.—Bras.
Deppeana, Steud. Nom. ed. II. i. 29 = brachiata.
diffusa, *Andr. Bot. Rep.* ix. t. 578.—Ind. occ.
dumosa, Salisb. Prod. 67 = martinicensis.
elata, *Sw. Prod. Veg. Ind. Occ.* 31.—Am. trop.
filipes, *Mart. & Schau. in DC. Prod.* xi. 652.—Bras.
fluminensis, *Vell. Fl. Flum.* 38 ; i. t. 95.—Bras.
foetida, *Sw. Prod. Veg. Ind. Occ.* 32.—Jamaic.
glabra, Lam. Illustr. t. 70. f. 1 = martinicensis.
glomerata, *Benth. Bot. Voy. Sulph.* 154.—Ecuador.
grandiflora, Hook. Bot. Mag. t. 4230 = Clerodendron grandiflorum.
grandifolia, Walp. Rep. iv. 121 = magnifolia.
graveolens, *Mart. & Schau. in DC. Prod.* xi. 651.—Bras.
hirta, *Casar. Nov. Stirp. Bras. Decad.* 47.—Bras.
Humboldtii, *Schau. in DC. Prod.* xi. 652.—Venezuela.
incana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 218.—Cuba.
inflexa, Vell. Fl. Flum. 38 ; i. t. 96. (Rubiacea, sec. Schau.).—Bras.
integrifolia, *Jacq. Hist. Stirp. Am.* 15. t. 173. f. 7.—N. Granat.
laeta, *H. B. & K. Nov. Gen. et Sp.* ii. 249.—N. Granat.
laevigata, *Juss. in Ann. Mus. Par.* vii. (1806) 76.—Ind. occ.
laevis, *Poepp. ex Walp. Rep.* iv. 119.—Bras.
laevis, *Willd. Sp. Pl.* i. 617.—Guiana.
lanuginosa, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 134 = obducta.
latifolia, Willd. ex Walp. Rep. iv. 120 = Mutisii.
laxiflora, *Benth. in Ann. Nat. Hist.* ii. (1839) 449.—Guiana.
Lhotskiana, *Cham. in Linnaea*, vii. (1832) 112.—Bras.
longifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 218.—N. Granat.
Luschnatii, *Schau. in DC. Prod.* xi. 651.—Bras.
lutea, Lam. Illustr. t. 70. f. 3 = laevis.
macrophylla, Desf. Cat. Hort. Paris. ed. III. 391 = magnifolia.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* ii. 251.—Am. austr.
macrophylla, Sieber, ex Schau. in DC. Prod. xi. 655 (= Rubiacea, sec. Schau.).—Ins. Trinit.
magnifolia, *Steud. Nom.* ed. II. i. 29.—Ind. occ.
Manabaea, Sw. Prod. Veg. Ind. Occ. i. 256 = laevis.
martinicensis, *Jacq. Obs. Bot.* ii. 3.—Ins. Martinic.
mediterranea, *Vell. Fl. Flum.* 38 ; i. t. 94.—Bras.
membranacea, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 219.—Am. trop.
mollis, *H. B. & K. Nov. Gen. et Sp.* ii. 250. t. 130.—Am. austr.
multiflora, *Ruiz & Pav. Fl. Per.* i. 50. t. 76.—Peruv.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* ii. 250. t. 131.—Am. austr.
Nuxia, Willd. Sp. Pl. i. 616 = Nuxia verticillata.
obducta, *Vell. Fl. Flum.* 39 ; i. t. 97.—Bras.
obovata, *Andr. Bot. Rep.* ix. 578.—Ind. occ.
oleifera, Casar. Nov. Stirp. Bras. Decad. 47 = fluminensis.
peruviana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 219.—Peruv.
pubescens, Willd. ex Steud. Nom. ed. II. i. 29 = salutaris.
punctata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 219.—Jamaic.

AEGIPHILA:—
racemosa, Vell. Fl. Flum. 37 ; i. t. 88 = cuspidata.
Riedeliana, *Schau. in DC. Prod.* xi. 649.— Bras.
salutaris, *H. B. & K. Nov. Gen. et Sp.* ii. 249.—Am. austr.
Sellowiana, *Cham. in Linnaea*, vii. (1832) 111.—Bras.
serrata, Vell. Fl. Flum. 38 ; i. t. 92 = graveolens.
splendens, *Schau. in DC. Prod.* xi. 648.—Bras.
stipulata, Vell. Fl. Flum. 37 ; i. t. 90 (Rubiacea, sec. Schau.).—Bras.
straminea, *Hoffmgg. Verz. Pfl. Nachtr.* iii. 18.—Bras.
tomentosa, *Cham. in Linnaea*, vii. (1832) 110.—Bras.
triantha, *Schau. in DC. Prod.* xi. 650.—Bras.
trifida, *Sw. Prod. Veg. Ind. Occ.* 32.—Jamaic.
umbellata, Vell. Fl. Flum. 37 ; i. t. 89. (Rubiacea, sec. Schau.).—Bras.
verrucosa, *Schau. in DC. Prod.* xi. 650.—N. Granat.
verticillata, Vell. Fl. Flum. 37 ; i. t. 91 = tomentosa.
viburnifolia, *Juss. in Ann. Mus. Par.* vii. (1806) 76. —Ins. Philipp.
villosa, *Vahl, Eclog.* i. 16.—Guiana.
virgata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 220.—Bras.
vitelliniflora, *Klotzsch, ex Walp. Rep.* iv. 123.—Hab. ?

AEGIPHYLA, Steud. Nom. ed. II. i. 29 = **Aegiphila**, Jacq.

AEGLE, Correa, in Trans. Linn. Soc. v. (1800) 22. *RUTACEAE*, Benth. & Hook. f. i. 306.
Belou, Adans. Fam. ii. 408 (1763).
decandra, *F. Villar, in Blanco, Fl. Philipp.* ed III. *Nov. App.* 38.—Ins. Philipp.
Marmelos, *Correa, in Trans. Linn. Soc.* v. (1800) 223.—Ind. or.
sepiaria, DC. Prod. i. 538 = Citrus trifoliata.

AEGLE, Dulac, Fl. Hautes-Pyr. 43 (1867) = **Posidonia**, Koen. (Naiad.).
fragilis, Dulac, l. c. = P. fragilis.

AEGOCHLOA, Benth. in Bot. Reg. sub t. 1622 (1833) = **Gilia**, Ruiz & Pav. (Polemon.).
atractyloides, Benth. l. c. = G. atractyloides.
cotulaefolia, Benth. l. c. = G. cotulaefolia.
eryngioides, Benth. l. c. = G. Navarettia.
intertexta, Benth. l. c. = G. intertexta.
pubescens, Benth. l. c. = G. cotulaefolia.
pungens, Benth. l. c. = G. squarrosa.
Torreyi, G. Don, Gen. Syst. iv. 246 = G. pungens.

AEGOMARATHRUM, Steud. Nom. ed. II. i. 30 (1840) = **Cachrys**, Tourn. (Umbell.).
crispum, Steud. l. c. = C. crispa.
pterochlaena, Steud. l. c. = C. sicula.
siculum, Mor. ex Nym. Consp. 314 = Hippomarathrum Bocconi.

AEGONYCHON, S. F. Gray, Nat. Arr. Brit. Pl. ii. 354 (1821) = **Lithospermum**, Linn. (Borag.).
arvense, S. F. Gray, l. c. = L. arvense.
repens, S. F. Gray, l. c. = L. purpureo-caeruleum.

AEGOPICRON, Giseke, Prael. 474 (1792) = AEGOPRICON, Linn. f. = **Maprounea**, Aubl. (Euphorb.).

AEGOPODION, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = seq.

AEGOPODIUM, Knaut, ex Linn. Syst. ed. I. (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 893.
Podagraria, Hall. Enum. Stirp. Helv. ii. 427 (1742).
alpestre, *Ledeb. Fl. Alt.* i. 354.—Sibir. altaic.
angelicaefolium, Salisb. Prod. 169 = Ae. Podagraria.
Carum, Wibel, Prim. Fl. Werth. 199 = Carum Carvi.
involucratum, Orphan. ex Boiss. Fl. Orient. ii. 923 = Physospermum aegopodioides.
latifolium, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 711 = Ae. Podagraria.
Podagraria, *Linn. Sp. Pl.* 265.—Europ. ; As. bor.
ternatum, Gilib. Fl. Lituan. ii. 41 = Ae. Podagraria.

AEGOPOGON, Beauv. Agrost. 122. t. 22. f. 4. (1812) partim = **Amphipogon**, R. Br. (Gramin.).
avenaceus, Beauv. l. c. = Am. strictus.
debilis, Beauv. l. c. = Am. debilis.
laguroides, Beauv. l. c. = Am. laguroides.

AEGOPOGON:—
pusillus, Spreng. Syst. i. 270 = Chloris cruciata.
strictus, Beauv. Agrost. 122 = Am. strictus.
turbinatus, Beauv. l. c. = Am. turbinatus.

AEGOPOGON, Beauv. Agrost. 122 (1812) partim ; Humb. & Bonpl. in Willd. Sp. Pl. iv. 899 (1805). *GRAMINEAE*, Benth. & Hook. f. iii. 1121.
Hymenothecium, Lag. Gen. et Sp. Nov. 4 (1816).
Schellingia, Steud. in Flora, xxxiii. (1850) 231. t. 1.
bryophilus, *Doell, in Mart. Fl. Bras.* ii. iii. 239.—Bras.
cenchroides, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 899. —Venezuela.
geminiflorus, *Humb. & Bonpl. Nov. Gen. et Sp.* iv. 133. t. 43.—Am. trop.
multisetus, Steud. Syn. Pl. Gram. 146 = cenchroides.
quinquesetus, Roem. & Schult. Syst. ii. 805 = cenchroides.
pusillus, Beauv. Agrost. 122 = geminiflorus.
rigidisctus, *Steud. Syn. Pl. Gram.* 146.—Texas.
setifer, Nees, in Linnaea, xix. (1847) 691 = cenchroides.
submuticus, *Rupr. ex Trin. in Mém. Acad. Pétersb.* Sér. VI. v. ii. (1840) 3.—Hab. ?
tenellus, Trin. Gram. Unifl. 164 = geminiflorus.
trisetus, Roem. & Schult. Syst. ii. 805 = cenchroides.
unisetus, Roem. & Schult. l. c. = geminiflorus.

AEGOPORDON, Boiss. Diagn. Scr. I. vi. (1845) 112 = **Jurinea**, Cass. (Compos.).
berardioides, Boiss. l. c. 113.—Persia.

AEGOPRICON, Linn. f. Suppl. 413 (1789) = seq.

AEGOPRICUM, Linn. Pl. Surinam. 15 (1775) ; (Amoen. Acad. viii. 263) = **Maprounea**, Aubl. (Euphorb.).
betulinum, Linn. l. c. = M. guianensis.

AEGOSERIS, Steud. Nom. ed. II. i. (1840) 30 = **Crepis**, Linn. (Compos.).
leontodontoides, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 103 = C. leontodontoides.
setosa, Fourr. l. c. = C. setosa.
suffreniana, Fourr. l. c. = C. suffreniana.

AEGOTOXICON, Molina, Sagg. Chile, ed. II. 298 (1810) = **Aextoxicon**, Ruiz & Pav. (Euphorb.).

AEGOTOXICUM, Endl. Ench. 589 (1841) = praec.

AEGYLOPS, Honck. Syn. i. 484 (1792) err. typ. = AEGILOPS, Linn. = **Triticum**, Linn. (Gramin.).

AEIPHANES, Spreng. Anleit. ii. ii. 875 (1818) err. typ. = AIPHANES, Willd. = **Martinezia**, Ruiz & Pav. (Palm.).

AELUROPUS, Trin. Fund. Agrost. 143 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1193.
Calotheca, Spreng. Anleit. ii. i. 167 (1817) ; Syst. i. 347 (1825).
Chamaedactylis, T. Nees, Gen. Fl. Germ. Monocot. i. n. 66 (1840 ?).
arabicus, *Steud. Nom.* ed. II. i. 30.—Aegypt. ; Abyss.
bombycinus, Figar. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 258 = arabicus.
brevifolius, Nees, ex Steud. Nom. ed. II. i. 30 = pubescens.
concinnus, Figar. & De Not. in Mem. Acc. Torin. Scr. II. xii. (1852) 257 = laevis.
grandiflorus, *Munro, in Cat. Kew Distrib. Griffith*, &c. 57, nomen.—Afghan.
hirsutus, *Munro, l. c.* nomen.—Ind. or.
intermedius, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) ii. 292.—As. temp.
laevis, *Trin. Fund. Agrost.* 143.—Oriens.
litoralis, Parl. Fl. Ital. i. 461 = laevis.
mucronatus, Aschers. in Schweinf. Beitr. Fl. Acthiop. 297 = arabicus
niliacus, *Steud. Nom.* ed. II. i. 30.—Aegypt.
pubescens, *Trin. ex Steud. Nom.* ed. II. i. 30.— Ind. or.
pungens, C. Koch, in Linnaea, xxi. (1848) 408 = laevis.
repens, Parl. Fl. Ital. i. 462 = villosus.
sinaicus, Figar. & De Not. in Mem. Acc. Torin. Ser. II. xii. (1852) 257 = laevis.
Smithii, *Steud. Nom.* ed. II. i 30.—Ins. Canar.
villosus, *Trin. Fund. Agrost.* 143.—Afr. bor. ; Graecia.

AENOTHERA, Lam. Encyc. iv. 550 (1796), err. typ. — **Oenothera**, Linn. (Onagr.).

AEOLANTHUS, Mart. Amoen. Bot. Monac. 4. t. 2 (1825). *LABIATAE*, Benth. & Hook. f. ii. 1176.
Orollanthus, E. Mey. Comm. Pl. Afr. Austr. ii. 230 (1837).
abyssinicus, *Hochst. ex Benth. in DC. Prod.* xii. 80.— Abyss.
ambustus, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 138.—Afr. trop.
heliotropioides, *Oliver, l. c.* 137.—Afr. trop.
Livingstonei, *C. Koch, Berl. Wochenschr.* iii. (1860) 183.—Afr. trop.
parvifolius, *Benth. in DC. Prod.* xii. 80.—Afr. austr.
pinnatifidus, *Hochst. ex Benth. l. c.*—Abyss.
pubescens, *Benth. l. c.*—Afr. trop.
repens, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 137. —Afr. trop.
suaveolens, G. Don, in Loud. Hort. Brit. 247 = suavis.
suavis, *Mart. Amoen. Bot. Monac.* 4. t. 2.—Bras. ; Afr. ?
zanzibaricus, *S. Moore, in Journ. Bot.* xv. (1877) 68. —Afr. trop.

AEOLLANTHUS, Spreng. Syst. ii. 750 (1825) = praec.

AEONIA, Lindl. Bot. Reg. sub t. 817 (1824). *ORCHIDEAE*, Benth. & Hook. f. iii. 584.
Brachystepis, Thou. ex Pritzel, Ind. Ic. 162 (1855).
Oeonia, Lindl. Orch. Scel. 14, err. typ. 1826).
alata, *A. Rich. in Ann. Sc. Nat.* Sér. II. xvi. (1841) t. 11.—Ind. or.
Auberti, *Lindl. Bot. Reg.* sub t. 817.—Madag.
brachystachya, *Lindl. Gen. et Sp. Orch.* 245.—Madag.
macrostachya, *Lindl. l. c.* 245.—Madag.
polystachya, *Benth. & Hook. f. Gen.* iii. 584.— Madag.
rosea, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 496.— Madag.

AEONIUM, Webb. & Berth. Phyt. Canar. i. 184 (1836), t. 28 = **Sempervivum**, Linn. (Crassul.).
arboreum, Webb & Berth. l. c. 185 = S. arboreum.
balsamiferum, Webb & Berth. l. c. 192 = S. balsamiferum.
barbatum, Webb & Berth. l. c. 188 = S. piliferum.
Berthelotianum, Bolle, in Bonplandia, vii. (1859) 239.
Bethencourtianum, Webb, ex Bolle, l. c. 243.—Ins. Canar.
caespitosum, Webb & Berth. Phyt. Canar. i. 191 = S. caespitosum.
canariense, Webb & Berth. l. c. i. 196 = S. canariense.
Castello-Paivae, C. Bolle, in Bonplandia, vii. (1859) 240.—Ins. Canar.
ciliatum, Webb & Berth. Phyt. Canar. i. 195 = S. ciliatum.
cruentum, Webb & Berth. l. c. 186 = S. cruentum.
cuneatum, Webb & Berth. l. c. 197 = S. cuneatum.
decorum, Webb, ex C. Bolle, in Bonplandia, vii. (1859) 241.—Ins. Canar.
frutescens, Webb & Berth. Phyt. Canar. i. 198 ? = S. frutescens.
glandulosum, Webb & Berth. l. c. 185 = S. glandulosum.
glutinosum, Webb & Berth. l. c. 185 = S. glutinosum.
Goochiae, Webb. & Berth. l. c. 190 = S. Goochiae.
gorgoneum, J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 258.—Ins. Cap. Verd.
Haworthii, Webb & Berth. Phyt. Canar. i. 193 = S. Haworthii.
holochrysum, Webb & Berth. l. c. 194 = S. holochrysum.
laxum, Webb & Berth. l. c. i. 198 ? = S. laxum.
leucoblepharum, Webb, ex A. Rich. Tent. Fl. Abyss. i. 314 = S. chrysanthum.
Lindleyi, Webb & Berth. Phyt. Canar. i. 189 = S. villosum.
manriqueorum, C. Bolle, in Bonplandia, vii. (1859) 241.—Ins. Canar.
Meyerheimii, C. Bolle, l. c. 239.— Ins. Mader.
Saundersii, C. Bolle, l. c. 241.—Ins. Canar.
Smithii, Webb & Berth. Phyt. Canar. i. 187 = S. Smithii.
strepsicladum, Webb & Berth. l. c. = S. strepsicladum.
abulaeforme, Webb & Berth. l. c. 185 = S. tabulaeforme.

AEONIUM :—
undulatum, Webb & Berth. Phyt. Canar. i. 197 = S. undulatum.
urbicum, Webb & Berth. l. c. 194 = S. urbicum.
viscatum, Webb, ex C. Bolle, in Bonplandia, vii. (1859) 241.—Ins. Canar.
Webbii, C. Bolle, l. c. 238.—Ins. Canar.
Youngianum, Webb & Berth. Phyt. Canar. i. 197 = S. Youngianum.

AËRA, Aschers. Fl. Brandenb. 830 (1864) = **Aira**, Linn. (Gramin.).

AËRANGIS, Reichb. f. in Flora, xlviii. (1865) 190 = **Angraecum**, Thou. (Orchid.).
flabellifolia, Reichb. f. l. c. 191.—Afr. trop.
Kotschyi, Reichb. f. Otia Bot. Hamb. 78 = Angr. Kotschyi.

AËRANTHES, Lindl. Bot. Reg. t. 817 (1824). *ORCHIDEAE*, iii. 576.
Arachnitis, *Lindl. Bot. Reg.* t. 817.—Ins. Mascar.
filiformis, Griseb. Fl. Brit. W. Ind. 625.—Ind. occ.
grandiflorus, *Lindl. Bot. Reg.* t. 817.—Madag.
grandiflorus, Lindl. Gen. et Sp. Orch. 243, pro parte = A. Arachnitis.
jamaicensis, Reichb. f. ex Griseb. Fl. Brit. W. Ind. 625 = Angraecum micranthum.
sesquipedalis, Lindl. Bot. Reg. sub t. 817 = Angraecum sesquipedale.

AËRANTHUS, Bartl. Ord. 57 (1830) = praec.

AËRANTHUS, Reichb. f. in Walp. Ann. vi. 899 (1861) = **Mystacidium**, Lindl. (Orchid.), &c.
aciculatus, Reichb. f. & Warm. in Otia Bot. Hamb. 91.—Bras.
arachnanthus, Reichb. f. in Flora, lxviii. (1885) 539.—Ins. Comor.
brachystachyus, Boj. Hort. Maurit. 314 = Aeonia brachystachya.
Calceolus, S. Moore, in Baker, Fl. Maurit. 353 = Angraecum Calceolus.
comorensis, Reichb. f. in Flora, lxviii. (1885) 540.—Ins. Comor.
Curnowianus, Reichb. f. in Gard. Chron. N. S. xix. (1883) i. 306.—Madag.
dentiens, Reichb. f. in Flora, lxviii. (1885) 381.—Ins. Comor.
distichus, Reichb. f. in Walp. Ann. vi. 901 = Angraecum distichum.
erythropollinius, Reichb. f. in Flora, xlviii. (1865) 190.—Afr. trop.
expansus, S. Moore, in Baker, Fl. Maurit. 351 = Angraecum expansum.
Fasciola, Reichb. f. in Walp. Ann. vi. 902 = Angraecum Fasciola.
filicornis, Reichb. f. l. c. 900 = M. filicorne.
fragrans, Reichb. f. l. c. 899 = Angraecum fragrans.
funalis, Reichb. f. l. c. 902 = Angraecum funale.
Gerrardi, Reichb. f. in Flora, l. (1867) 117.—Afr. austr.
Gladiator, Reichb. f. l. c. lxviii. (1885) 539.—Ins. Comor.
gladiifolius, Reichb. f. in Walp. Ann. vi. 900 = Angraecum gladiifolium.
gracilis, Reichb. f. in Flora, l. (1867) 117 = M. gracile.
Grandidieranus, Reichb. f. l. c. lxviii. (1885) 381.—Ins. Comor.
Guyonianus, Reichb. f. l. c. xlviii. (1865) 190 = Angraecum Guyonianum.
hispidulus, Reichb. f. in Linnaea, xli. (1877) 31.—N. Granat.
hymenanthus, Griseb. Cat. Pl. Cub. 264 = Dendrophylax hymenanthus.
intermedius, Reichb. f. & Warm. in Otia Bot. Hamb. 91.—Bras.
jamaicensis, Reichb. f. in Walp. Ann. vi. 901 = Angraecum micranthum.
Lansbergii, Reichb. f. l. c. = Angraecum Lansbergii.
Lehmanni, Reichb. f. Otia Bot. Hamb. 10.—Hab. ?
Leonii, Reichb. in Flora, lxviii. (1885) 380 = seq.
Leonis, Reichb. f. in Gard. Chron. N. S. xxiii. (1885) 726 = Angraecum Humblotii.
Lindenii, Reichb. f. in Walp. Ann. vi. 902 = Angraecum Lindenii.
macrostachys, Reichb. f. l. c. 900 = Aeonia macrostachya.

AËRANTHUS :—
Meirax, Reichb. f. in Flora, lxviii. (1885) 540.—Afr. austr.
micranthus, Reichb. f. in Walp. Ann. vi. 90 = Angraecum micranthum.
Monteverdi, Reichb. f. in Flora, xlviii. (1865) 279.—Hab. ?
muscicola, Reichb. f. l. c. 190.—Afr. trop.
neglectus, Reichb. f. & Warm. in Otia Bot. Hamb. 91.—Bras.
organensis, Reichb. f. in Walp. Ann. vi. 901 = Angraecum organense.
ornithorrhynchus, Reichb. f. l. c. 903 = Angraecum ornithorrhynchum.
pachyrhyzus, Reichb. f. in Flora, xlviii. (1865) 279.—Ind. occ.
parahybunensis, Rodrig. Orch. Nov. ii. 245.—Bras.
pectinatus, Reichb. f. in Walp. Ann. vi. 900 = Angraecum pectinatum.
phalaenophorus, Reichb. f. in Flora, lxviii. (1885) 539.—Ins. Comor.
polystachyus, Reichb. f. in Walp. Ann. vi. 901 = Aeonia polystachia.
porrectus, Reichb. f. in Flora, xlviii. (1865) 279.—Ind. occ.
pusillus, Reichb. f. l. c. l. (1867) 117 = M. pusillum.
rectus, S. Moore, in Baker, Fl. Maurit. 351 = Angraecum rectum.
rutilus, Reichb. f. in Flora, lxviii. (1885) 540.—Ins. Comor.
Sallei, Reichb. f. in Walp. Ann. vi. 902.—Ins. S. Doming.
Schiedei, Reichb. f. l. c. 901 = Campylocentron Schiedei.
sesquipedalis, Reichb. in Flora, lxviii. (1885) 381 = Angraecum sesquipedale.
spathaceus, Griseb. Cat. Pl. Cub. 264.—Cuba.
Thouarsii, S. Moore, in Baker, Fl. Maurit. 35 = Angraecum filicornu.
trifurcus, Reichb. f. in Flora, lxviii. (1885) 540.—Ins. Comor.
volucris, Reichb. f. in Walp. Ann. vi. 900 = Aeonia Auberti.
xanthopollinius, Reichb. l. c. xlviii. (1865) 190.—Afr. trop.

AËRIDES, Lour. Fl. Cochinch. 525 (1790). *ORCHIDEAE*, Benth. & Hook. f. iii. 576.
AËERIDIUM, Salisb. in Trans. Hort. Soc. i. (1812) 295.
ORXERA, Rafin. Fl. Tellur. iv. 37 (1836).
acuminatissimum, *Lindl. Gen. et Sp. Orch.* 240.—Java.
affine, Wall. Cat. n. 7316 = multiflorum.
amplexicaule, Lindl. Gen. et Sp. Orch. 239 = Sarcochilus lilacinus.
ampullaceum, Roxb. Fl. Ind. iii. 476 = Sarcolabium ampullaceum.
anceps, Lindl. Gen. et Sp. Orch. 242 = Sarcochilus anceps.
angustifolium, Lindl. l. c. 241 = Sarcochilus angustifolius.
appendiculatum, Wall. Cat. n. 7315 = Sarcanthus appendiculatus.
Arachnites, Lindl. Gen. et Sp. Orch. 238 = Sarcochilus Arachnites.
Arachnites, Sw. in Schrad. Journ. ii. (1799) 233 = Arachnanthe moschifera.
Ballantinianum, Reichb. f. in Gard. Chron. N. S. xxiv. (1885) ii. 198 = suavissimum.
Bernhardianum, *Reichb. f. l. c.* 650.—Borneo.
bicolor, Reinw. ex Lindl. Fol. Orchid. Vanda, 4 = Vanda furva.
Borassi, *Buch.-Ham. ex Sm. in Rees, Cyclop.* xxxix. n. 8.—Ind. or.
Brockessii, Heynh. Nom. ii. 11 = crispum.
Brookei, Batem. ex Lindl. in Bot. Reg. (1841) Misc. 55 = crispum.
calceolare, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 19.—Sumatra.
calceolare, Buch.-Ham. ex Sm. in Rees, Cyclop. xxxix. n. 11 = Saccolabium calceolare.
carnosum, Griff. Notul. iii. 365 = Doritis taenialis.
compressum, Lindl. Gen. et Sp. Orch. 240 = Dendrocolla compressa.
coriaceum, *Sw. in Schrad. Journ.* (1799) 234.—Madag.
cornutum, Roxb. Hort. Beng. 63; Lindl. Bot. Reg. t. 1485 = odoratum.

AËRIDES :—
costatum, '*Westk.*' *ex Steud. Nom.* ed. II. i. 30.—Ind. or.
crassifolium, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 145.—Burma.
crispum, *Lindl. in Wall. Cat.* n. 7319.—Malabar.
cylindricum, Hook. Bot. Mag. t. 4982 = A. vandarum.
cylindricum, *Lindl. in Wall. Cat.* n. 7317.—Ind. or.
decumbens, *Griff. Notul.* iii. 365; *Ic. Pl. Asiat.* t. 320.—Burma.
dasypogon, *Sm. in Rees, Cyclop.* xxxix. n. 10.—Nepal.
dasypogon, Hort. ex Batem. in Hook. Bot. Mag. sub t. 5630 = Sarcanthus erinaceus.
difforme, Wall. ex Lindl. Gen. et Sp. Orch. 242 = Ornithochilus fuscus.
diurnum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 19.—Sumatra.
elongatum, *Blume, Bijdr.* 367.—Java.
emarginatum, Lindl. Gen. et Sp. Orch. 241 = Sarcochilus emarginatus.
Emericii, *Reichb. f. in Gard. Chron.* xviii. (1882) ii. 586.—Ins. Andam.
expansum, Reichb. f. l. c. 40 = falcatum.
falcatum, *Lindl. & Paxt. Fl. Gard.* ii. (1851) 142.—Burma.
Farmeri, *Boxall, ex Naves, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 239.—Ins. Philipp.
Fenzlianum, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 282.—Ins. Philipp.
Fieldingi, *Lodd. ex E. Morr. in Belg. Hortic.* (1876) 286. t. 10.—Reg. Himal.
flavidum, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 101 = suavissimum.
Flos-aëris, Sw. in Schrad. Journ. ii. (1799) 233 = Sarcochilus Arachnites.
guttatum, Roxb. Fl. Ind. iii. 471 = Rhynchostylis retusa.
Houlletianum, *Reichb. f. in Gard. Chron.* (1872) 1194.—Cochinch.
Hystrix, *Lindl. Gen. et Sp. Orch.* 242 = Sarcochilus Hystrix.
Hystrix, Lindl. in Journ. Linn. Soc. iii. (1859) 42 = Ornithochilus fuscus.
illustre, Reichb. f. in Gard. Chron. N. S. xviii. (1882) ii. 71 = maculosum.
inflexum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 324.—Malaya.
japonicum, *Reichb. f. in Hamb. Gartenz.* xix (1863) 210.—Japon.
jucundum, *Reichb. f. l. c.* xvi. (1860) 281.—Java.
Larpentae, Hort. ex Reichb. f. in Otto et Dietr. Allg. Gartenz. xxiv. (1856) 219 = falcatum.
lasiopetala, *Willd. Sp. Pl.* iv. 130.—Ind. or.
latifolium, *Thw. Enum. Pl. Zeyl.* 429.—Zeylan.
Lawrenciae, *Reichb. f. in Gard. Chron.* xx. (1883) 460.—As. trop.
Leeanum, *Reichb. f. l. c.* xv. (1881) 656.—Hab. ?
Leonaei, *Reichb. f.*—Cf. *L'Orchidoph.* (1885) 301.—Burma.
lepidum, *Reichb. f. in Gard. Chron.* xix. (1883) 466.—Ind. or.
Lindleyanum, Wight, Ic. Pl. Ind. Or. 1677 = crispum.
Lobbii, Hort. ex Lem. Illustr. Hortic. xv. (1868) t. 559 = multiflorum.
Lobbii, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 323.—Burma.
McMorlandi, *Williams, Man. Orch.* 66; ed. VI. 105.—Ind. or.
macrostachyon, Spreng. Syst. iii. 719 = Aeonia macrostachya.
maculatum, Buch.-Ham. ex Sm. in Rees, Cyclop. xxxix. n. 9 = Vanda spathulata.
maculatum, *Llanos, Fragm.* 93.—Ins. Philipp.
maculosum, *Lindl. Bot. Reg.* xxxi. (1845) 58.—Ind. or.
margaritaceum, *Williams, Man. Orch.* 66; ed. VI. 105.—Ind. or.
marginatum, *Reichb. f. in Gard. Chron.* N. S. xxiii. (1885) i. 533.—Hab. ?
matutinum, *Blume, Bijdr.* 366.—Java.
matutinum, Willd. Sp. Pl. iv. 131 = Renanthera Arachnites.
Mendeli, Hort. ex E. Morr. in Belg. Hortic. (1876) 288 = falcatum.
minimum, Lindl. Gen. et Sp. Orch. 241 = Sarcochilus minimus.

AËRIDES :—
mitratum, *Reichb. f. in Bot. Zeit.* xxii. (1864) 415.—Ind. or.
multiflorum, *Roxb. Fl. Ind.* iii. 475.—Ind. or.
nobile, Warn. Scl. Orch. Ser. I. t. 11 = suavissimum.
obtusum, Lindl. Gen. et Sp. Orch. 239 = Sarcochilus obtusus.
odoratum, *Lour. Cochinch.* 525.—China; Ind. or.
odoratum, Reinw. ex Blume, Rumphia, iv. 53 = suavcolens.
Ortgiesianum, *Reichb. f. in Gard. Chron.* xxiii. (1885) 501.—Hab.?
pachyphyllum, *Reichb. f. l. c.* xiv. (1880) 231.—Burma.
pallidum, *Blume, Rumphia*, iv. 53.—Ins. Timor.
pallidum, Lindl. Gen. et Sp. Orch. 241 = Sarcochilus pallidus.
pallidum, Roxb. Fl. Ind. iii. 475 (= *Camarotis pallida*).
paniculatum, Ker-Gawl. Bot. Reg. iii. 220 = Vanda paniculata.
praemorsum, Willd. Sp. Pl. iv. 130.—Rhynchostylis retusa.
purpurascens, *Lindl. Gen. et Sp. Orch.* 240.—Java.
pusillum, Lindl. l. c. 241 = Sarcochilus pusillus.
quinquevulnera, *Lindl. Sert. Orch.* 30.—Malaya.
radiatum, *Roxb. Fl. Ind.* iii. 476.—Ind. or.
radicosum, *A. Rich. in Ann. Sc. Nat.* Sér. II. xv. (1841) 65. t. 1. f. c.—Ind. or.
Reichenbachii, Linden, in C. Koch & Fint. Wochenschr. (1858) 61 = suavissimum.
retusum, Sw. in Schrad. Journ. ii. (1799) 233 = Rhynchostylis retusa.
rigidum, *Buch.-Ham. ex Sm. in Rees, Cyclop.* xxxix. n. 12.—Ind. or.
Roebelenii, *Reichb. f. in Gard. Chron.* xxi. (1884) 510.—Ins. Philipp.
Rohanianum, Reichb. f. l. c. 206 = suavissimum.
roseum, Lodd. ex Paxt. Fl. Gard. t. 60 = multiflorum.
rostratum, Roxb. Fl. Ind. iii. 474 = Sarcochilus purpureus.
Ruckeri, *Hort. Donat. ex Lubbers, Cat. Pl. Rar. San Donato* (1880) 14.—Hab.?
Sanderianum, *Reichb. f. in Gard. Chron.* xxii. (1884) 134.—As. trop.
Schroederi, Hort. ex Reichb. f. in Allgem. Gartenz. (1855) 226 = maculosum.
shibatianum, *Boxall, ex Naves, in Blanco, Fl. Philipp.* ed. II. *Nov. App.* 239.—Ins. Philipp.; sphalm. ? = Thibautianum.
spicatum, D. Don, Prod. Fl. Nep. 31 = Rhynchostylis retusa.
spurium, *Lindl. Gen. et Sp. Orch.* 241.—Java.
suaveolens, *Blume, Rumphia*, iv. 53.—Java.
suaveolens, *Roxb. Fl. Ind.* iii. 473.—Ind. or.
suavissimum, Lindl. in Journ. Hort. Soc. iv. (1849) 264.—Malacca.
subulatum, Lindl. Gen. et Sp. Orch. 241 = Sarcochilus subulatus.
Sulingi, Blume, Bijdr. 367 = Renanthera Sulingi.
taeniale, Lindl. Gen. et Sp. Orch. 239 = Doritis taenialis.
teres, *Lindl. l. c.* 240.—Java.
tessellatum, Wight, in Wall. Cat. n. 7318 = Vanda Roxburghii.
testaceum, Lindl. Gen. et Sp. Orch. 238 = Vanda parviflora.
Teysmanni, *Miq. Choix Pl. Buitenz.* t. 18.—Malaya.
Thibautianum, *Reichb. f. in Gard. Chron.* (1866) 100.—Polynesia.
Thunbergii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 205.—Japon.
timoranum, *Miq. Fl. Ind. Bat.* iii. 695.—Ins. Timor.
trigonum, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 177 = multiflorum.
umbellatum, Wall. Cat. n. 1137 = Saccolabium acutifolium.
uncinatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 324.—Java.
undulatum, Sm. in Rees, Cyclop. xxxix. n. 13 = Saccolabium praemorsum.
vandarum, *Reichb. f. in Gard. Chron.* (1867) 997.—Ind. or.
Veitchi, Hort. ex E. Morr. in Belg. Hortic. (1876) 291, xxxi. (1881) 123 = multiflorum.
virens, *Lindl. Bot. Reg.* (1843) *Misc.* 41.—Hab.?
viridiflorum, *Thw. Enum. Pl. Zeyl.* 430.—Zeylan.
Wightianum, Lindl. in Wall. Cat. n. 7320 = Vanda parviflora.

AËRIDES :—
Williamsii, Warn. Scl. Orch. Ser. I. t. 21 = Fieldingi.
Wilsonianum, Hort. ex Gard. Chron. xxiii. (1885) 705 = suavissimum?
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 40.—Java.

AËRIDIUM, Pfeiff. Nom. i. 67 (1873) = AEËRIDIUM, Salisb. = **Aërides**, Lour. (Orchid.).

AËROBION, Kaempf. ex Spreng. Syst. iii. 679 (1826) = **Angraecum**, Thou. (Orchid.).
aphyllum, Spreng. l. c. 718 = Angr. aphyllum.
Calceolus, Spreng. l. c. 717 = Angr. Calceolus.
carinatum, Spreng. l. c. 718 = Eulophia virens.
caulescens, Spreng. l. c. 717 = Mystacidium caulescens.
citratum, Spreng. l. c. 718 = Angr. citratum.
crassum, Spreng. l. c. 717 = Angr. crassum.
cucullatum, Spreng. l. c. = Angr. cucullatum.
expansum, Spreng. l. c. 716 = Angr. expansum.
filicornu, Spreng. l. c. = Angr. filicornu.
fragrans, Spreng. l. c. = Angr. fragrans.
gladiifolium, Spreng. l. c. = Angr. gladiifolium.
gracile, Spreng. l. c. 717 = Angr. gracile.
implicatum, Spreng. l. c. 716 = Angr. implicatum.
inapertum, Spreng. l. c. 717 = Angr. inapertum.
maculatum, Spreng. l. c. 718 = Angr. maculatum.
multiflorum, Spreng. l. c. 717 = Angr. multiflorum.
palmiforme, Spreng. l. c. = Angr. palmiforme.
parviflorum, Spreng. l. c. 718 = Angr. parviflorum.
pectinatum, Spreng. l. c. 717 = Angr. pectinatum.
rectum, Spreng. l. c. 716 = Angr. rectum.
recurvum, Spreng. l. c. = Angr. recurvum.
striatum, Spreng. l. c. 717 = Angr. striatum.
superbum, Spreng. l. c. 718 = Angr. superbum.
triquetrum, Spreng. l. c. 717 = Angr. triquetrum.

AEROKORION, Scop. Ann. iv. Hist. Nat. 96 (1770) = AEROCORION, Adans. = **Galanthus**, Tourn. (Amaryll.).

AERUA, Juss. Gen. 88 (1789) = **Aerva**, Forsk. (Amarant.).

AERVA, Forsk. Fl. Aegypt. Arab. 170 (1775.) *AMARANTACEAE* (Benth. & Hook. f. iii. 34).
aegyptiaca, J. F. Gmel. Syst. ii. 1026 = tomentosa.
ambigua, *Moq. in DC. Prod.* xiii. II. 302.—Afr. austr.
bengalensis, Buch.-Ham. ex Wall. Cat. sub n. 6911 = scandens.
Bovii, Edgew. in Journ. Linn. Soc. vi. (1862) 206 = tomentosa.
brachiata, Mart. Beitr. Amarant. 83 = Nothosaerua brachiata.
Burchellii, Moq. in DC. Prod. xiii. II. 302 = A. leucura.
caudata, *Boj. Hort. Maurit.* 268.—Madag.?
chenopodiifolia, Boj. l. c. = Nothosaerua brachiata.
congesta, *Balf. f. ex Baker, Fl. Maurit.* 267.—Ins. Mascar.
clegans, *Moq. in DC. Prod.* xiii. II. 303.—Arab.
floribunda, Wight, Ic. Pl. t. 1776 = lanata.
glabrata, *Hook. f. Fl. Brit. Ind.* iv. 728.—Ind. or.
incana, Mart. in Nov. Act. Acad. Nat. Cur. xii. (1826) 291 = tomentosa.
javanica, Juss. in Ann. Mus. Par. xi. (1808) 131 = tomentosa.
lanata, *Juss. l. c.*—As. trop.; Arab.
leucura, *Moq. in DC. Prod.* xiii. II. 302.—Afr. austr.
microphylla, *Moq. l. c.* 301.—'Mare rubrum.'
Monsonia, *Mart. Beitr. Amarant.* 83.—Ind. or.
nervosa, *Mart. l. c.*—Ind. or.
ovata, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 174 = tomentosa.
platyphylla, Ehrenb. ex Schweinf. l. c. = tomentosa.
pubescens, *Mart. Beitr. Amarant.* 83.—Ind. or.
radicans, Mart. l. c.—Ind. or.
revoluta, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 92.—Ins. Socotr.
sanguinea, Hort. ex Moq. in DC. Prod. xiii. II. 300 = sanguinolenta.
sanguinolenta, *Blume, Bijdr.* 547.—Java.
scandens, Gamble, Trees of Darjeeling = Stilbanthus scandens.
scandens, *Wall. Cat. n.* 6911.—As. et Afr. trop.
sericea, *Moq. in DC. Prod.* xiii. II. 304.—Ins. Sandvic.
setacea, Mart. Beitr. Amarant. 83 = A. Monsonia.
suaveolens, Edgew. in Journ. Linn. Soc. vi. (1862) 207 = tomentosa.

AERVA :—
timorensis, *Moq. in DC. Prod.* xiii. II. 301.—Ins. Timor.
timoriensis, Zipp. ex Span. in Linnaea, xv. (1841) 345 = sanguinolenta.
tomentosa, *Lam. Encyc.* i. 46.—As. et Afr. trop.
toncafolia, Buch.-Ham. ex Wall. Cat. sub n. 6911 = scandens.
Tondalo, Buch.-Ham. l. c. 6909 = lanata.
trichophylla, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 174 = tomentosa.
velutina, Moq. in DC. Prod. xiii. II. 301 = Pupalia lappacea.
villosa, *Moq. l. c.* 304.—Ins. Maurit.
viridis, E. Mey. ex Moq. l. c. = lanata.
Wallichii, Moq. l. c. 300 = tomentosa.
Wightii, *Hook. f. Fl. Brit. Ind.* iv. 728.—Ind. or.

AESCHINANTHUS, Endl. Gen. 716 (1838) = **Aeschynanthus**, Jack (Gesnerac.).

AESCHINOMENE, Nocca in Usteri, Ann. Bot. v. (1793) 16 = **Aeschynomene**, Linn. (Legum.).

AESCHRION, Vell. Fl. Flum. 58; i. t. 152 (1827) = **Picraena**, Lindl. (Simarub.).
crenata, Vell. ll. cc. = P. Vellozii.

AESCHYNANTHUS, Jack, in Trans. Linn. Soc. xiv. (1823) 42. *GESNERACEAE*, Benth. & Hook. f. ii. 1013.
RHEITHROPHYLLUM, Hassk. Cat. Hort. Bogor. Alt. 123 (1844).
TRICHOSPORUM, D. Don, in Edinb. Phil. Journ. vii. (1822) 84 (nomen prius).
acuminata, G. Don, Gen. Syst. iv. 657 = Chirita acuminata.
acuminata, *Wall. Cat. n.* 6397.—Ind. or.; China.
albida, A. DC. Prod. ix. 262 = purpurascens.
albida, *Steud. Nom.* ed. II. i. 32.—Java.
alba, Hort. ex Hanst. in Mart. Fl. Bras. viii. 418 = Codonanthe Devosiana.
amoena, *C. B. Clarke, in DC. Monog. Phan.* v. 47.—Java.
Andersoni, *C. B. Clarke, Comm. et Cyrt. Beng.* t. 48.—Burma.
angustifolia, *Steud. Nom.* ed. II. i. 32.—Java.
apicidens, *Hance, in Journ. Bot.* xxi. (1883) 167.—China.
arfakensis, *C. B. Clarke, in DC. Monog. Phan.* v. 36.—N. Guin.
atropurpurea, Van Houtte, Hort. Vanhout. Fasc. ii. 42 = purpurascens.
atrosanguinea, Van Houtte, Cat. (1851) ex C. B. Clarke, in DC. Monog. v. 52.—Hab.?
Aucklandiae, *Low, Sarawak*, 382, 386.—Borneo.
Aucklandi, Hort. Low, ex Loud. Hort. Brit. Supp. iii. 484 = praec.
Beccarii, *C. B. Clarke, in DC. Monog. Phan.* v. 47.—Sumatra.
Blumei, Mor. ex C. B. Clarke, l. c. 43 = A. Lamponga.
Boschiana, De Vriese, in Morr. Ann. Soc. Agric. Bot. Gand, ii. (1846) 403 = A. Lamponga.
bracteata, Benth. Fl. Hongkong, 258 = acuminata.
bracteata, *Wall. Cat. n.* 794.—Ind. or.
brevicalyx, *Miq. Fl. Ind. Bat.* ii. 720.—Sumatra.
candida, *Henders. Cat.* (1851) *ex C. B. Clarke, in DC. Monog. Phan.* v. 52.—Hab.?
caudata, *C. B. Clarke, l. c.* 34.—Sumatra.
ceylanica, *Gard. in Calc. Journ. Nat. Hist.* vi. (1846) 474.—Zeylan.
ceylanica, Wight, Ic. t. 1347 = Perrottetii.
chinensis, Gard. & Champ. in Hook. Kew Journ. i. (1849) 328 = acuminata.
chiritoides, *C. B. Clarke, in DC. Monog. Phan.* v. 28.—Reg. Himal.
conferta, Griff. Priv. Journ. 68. nomen = Lepionotus confertus.
cordifolia, *Hook. Bot. Mag.* t. 5131.—Borneo.
cryptantha, *C. B. Clarke, in DC. Monog. Phan.* v. 50.—Borneo.
Curtisii, *C. B. Clarke, l. c.* 45.—Borneo.
discolor, T. Moore, in Paxt. Fl. Gard. iii. (1852-53) 55 = purpurascens.
elongata, *C. B. Clarke, in DC. Monog. Phan.* v. 45.—Borneo.
fulgens, *Wall. Cat. n.* 797.—Ind. or.
geminata, *Zoll. & Morr. in Nat. en Geneesk. Arch. Neerl. Ind.* ii. (1845) 574.—Java.

AESCHYNANTHUS :—
gracilis, Hort. ex Hanst. in Mart. Fl. Bras. viii. 418 = Codonanthe Devosiana.
gracilis, *Parish, ex C. B. Clarke, Comm. et Cyrt. Beng. t.* 48.—Ind. or.
grandiflora, J. Grah. Cat. Pl. Bomb. 146 = Perrottetii.
grandiflora, *Spreng. Syst.* iv. *Cur. Post.* 238.—Ind. or.
Griffithii, *R. Br. in Benn. Pl. Jav. Rar.* 115.—Ind. or.
hians, *C. B. Clarke, in DC. Monog. Phan.* v. 51.—Borneo.
Hookeri, *C. B. Clarke, l. c.* 21.—Reg. Himal.
Horsfieldii, *R. Br. in Benn. Pl. Jav. Rar.* 116.—Java.
intermedia, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 414.—Malaya.
javanica, *Hook. Bot. Mag.* t. 4503.—Java.
Kingii, *C. B. Clarke, in DC. Monog. Phan.* v. 31.—Reg. Himal.
Lamponga, *Miq. Fl. Ind. Bat. Suppl.* 563.—Sumatra.
lepidosperma, *C. B. Clarke, in DC. Monog. Phan.* v. 35.—Borneo.
leptoclada, *C. B. Clarke, l. c.* 39.—N. Guin.
levipes, *C. B. Clarke, l. c.* 28.—Assam.
Lobbiana, *Hook. Bot. Mag.* t. 4260.—Java.
longicaulis, *Wall. Cat. n.* 888.—Ind. or.
longiflora, *Blume, Bijdr.* 766.—Malaya.
longiflora, Wall. Cat. n. 795 = superba.
macrocalyx, *C. B. Clarke, in DC. Monog. Phan.* v. 46.—Sumatra.
maculata, *Lindl. Bot. Reg.* (1841) t. 28.—Ind. or.
Mannii, *Kurz, ex C. B. Clarke, in DC. Monog. Phan.* v. 29.—Ind. or.
marmorata, *F. Moore, in Paxt. Fl. Gard.* iii. (1852–53) 56.—Hab. ?
Masoniae, *Kurz, ex C. B. Clarke, l. c.* 29—Ind. or.
micrantha, *C. B. Clarke, l. c.* 27.—Reg. Himal.
microphylla, *C. B. Clarke, l. c.* 49.—Ins. Celebes.
microtricha, *C. B. Clarke, l. c.* 51.—N. Guin.
miniata, *Lindl. Bot. Reg.* N. S. xix. t. 61.—Java.
Motleyi, *C. B. Clarke, in DC. Monog. Phan.* v. 20.—Malaya.
Neesii, Zoll. & Moritzi, Syst. Verz. Zoll. 45 = pulchra.
obconica, *C. B. Clarke, in DC. Monog. Phan.* v. 50.—Malacca.
oblongifolia, G. Don, Gen. Syst. iv. 657 = Chirita acuminata.
obovata, *C. B. Clarke, in DC. Monog. Phan.* v. 47.—Java.
parasitica, C. B. Clarke, Comm. et Cyrt. Beng. t. 49 = Hookeri.
parasitica, Wall. Cat. n. 796 = grandiflora, pro parte.
parviflora, C. B. Clarke, Comm. et Cyrt. Beng. t. 47 = micrantha.
parviflora, *Spreng. Syst.* iv. *Cur. Post.* 238 = ramosissima.
parvifolia, *R. Br. in Benn. Pl. Jav.* 115.—Banca.
Paxtonii, Lindl. in Journ. Hort. Soc. iv. (1849) 79 = bracteata.
Peelii, Hook. f. & Thoms. Illustr. Himal. Pl. 17 = bracteata.
Perrottetii, *A. DC. in DC. Prod.* ix. 261.—Ind. or.
Perrottetii, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 475 = ceylanica.
philippinensis, *C. B. Clarke, in DC. Monog. Phan.* v. 39.—Ins. Philipp.
podocarpa, *C. B. Clarke, l. c.* 40.—N. Guin.
pulchella, Hort. ex Hanst. in Mart. Fl. Bras. viii. 418 = Codonanthe Devosiana.
pulchra, *G. Don, Gen. Syst.* iv. 656.—Malaya.
purpurascens, *Hassk. Cat. Pl. Hort. Bogor. Alt.* 154.—Java.
radicans, *Jack, in Trans. Linn. Soc.* xiv. (1823) 43.—Malaya.
radicans, Wall. Cat. n. 798 = Griffithii.
ramosissima, *Wall. Cat. n.* 799.—Ind. or.
Roxburghii, *Paxt. Bot. Dict. ex C. B. Clarke, in DC. Monog.* v. 52.—Hab. ?
rubiginosa, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 413.—Java.
siphonantha, *C. B. Clarke, in DC. Monog. Phan.* v. 45.—Borneo.
speciosa, *Hook. Bot. Mag.* t. 4320.—Java.
splendens, Lindl. & Paxt. Fl. Gard. iii. 169. t. 108 = speciosa.
splendida ×, *Lem. Jard. Fleur.* iii. (1853) t. 255.
superba, *C. B. Clarke, Comm. et Cyrt. Beng.* t. 46.—Ind. or.
Szcokhoffii, Hort. Berol. ex C. B. Clarke, in DC. Monog. Phan. v. 43 = pulchra.
tetraquetra, *C. B. Clarke, l. c.* 38.—Sumatra.

AESCHYNANTHUS :—
Teysmanniana, *Miq. in Bot. Zeit.* vi. (1848) 509.—Java.
tricolor, *Hook. Bot. Mag.* t. 5031.—Java.
tubiflora, *C. B. Clarke, in DC. Monog. Phan.* v. 36.—N. Guin.
verticillata, *C. B. Clarke, l. c.* 40.—N. Guin.
viridiflora, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 34.—Ins. Celebes.
volubilis, *Jack, in Trans. Linn. Soc.* xiv. (1823) 42. t. 3. f. 3.—Sumatra.
Wallichii, *Benn. Pl. Jav. Rar.* 116.—Malaya.
Wilsoni, Walker, ex C. B. Clarke, in DC. Monog. v. 26 = ceylanica.
zebrina, Van Houtte [Cat. 1851] ex Walp. Rep. vi. 521 = marmorata.
zeylanica, Walp. Ann. iii. 95 = ceylanica.
Zollingeri, *C. B. Clarke, in DC. Monog. Phan.* v. 44.—Java.

AESCHYNOMENE, Linn. Gen. ed. I. 350 (1737).
LEGUMINOSAE, Benth. & Hook. f. i. 515.
GAJATI, Rumph. ex Adans. Fam. ii. 328 (1763).
MANTODDA, Adans. Fam. ii. 508 (1763).
MACROMISCUS, Turcz. in Bull. Soc. Nat. Mosc. (1846) II. 507.
PATAGONIUM, E. Mey. Comm. Pl. Afr. Austr. 122 (1835).
ROCHEA, Scop. Introd. 296 (1777).
RUEPPELIA, A. Rich. Tent. Fl. Abyss. i. 203. t. 37 (1847).
abyssinica, Vatke, in Oest. Bot. Zeitschr. xxix. (1879) 224 = Ruppellii.
aegyptiaca, Steud. Nom. ed. II. i. 31 = Sesbania aegyptiaca.
acutangula, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 148.—Afr. trop.
americana, *Linn. Sp. Pl.* 713.—Ind. occ.; Am. trop.
aquatica, Steud. Nom. ed. II. i. 31 = aspera.
arborea, Linn. Sp. Pl. 713 = Desmodium umbellatum.
aristata, Jacq. Hort. Schoenb. i. 59 = Pictetia aristata.
aspera, *Linn. Sp. Pl.* 713.—Geront. trop.
aspera, Wall. Cat. n. 5666 = indica.
aspera, Muhl. ex Willd. Sp. Pl. iii. 1163 = hispida.
atropurpurea, *Span. in Linnaea*, xv. (1841) 192.—Ins. Timor.
bispinosa, Jacq. Ic. iii. 13 = Sesbania aculeata.
bracteosa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 150.—Afr. trop.
brasila, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 63 = Martiusii.
brasiliana, *DC. Prod.* ii. 322.—Bras.
brevifolia, *Poir. Encyc.* iv. 451.—Madag.
brevipes, *Benth. in Mart. Fl. Bras.* xv. 1. 66.—Bras.
Burmanni, *Raeusch. Nom.* ed. III. 214.—Ind. or.
cachemiriana, Cambess. in Jacquem. Voy. Bot. 40 = indica.
cannabina, Retz. Obs. Fasc. v. 26 = Sesbania aculeata.
cassioides, Desv. in Ham. Prod. Pl. Ind. Occ. 51 = Histrix.
cassioides, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 240 = Sesbania cassioides.
ciliata, Vog. in Linnaea, xii. (1838) 84 = hispida.
coccinea, Linn. f. Suppl. 330 = Sesbania grandiflora.
coluteoides, Less. & A. Rich. Voy. Astrol. Bot. ii. 87. t. 32 = Ormocarpum sennoides.
conferta, *Benth. in Ann. Nat. Hist.* iii. (1839) 433.—Guian.
crepita, *Jacq. Fragm.* 37. t. 42. f. 2.—Venezuela.
cristata, *Vatke, in Oest. Bot. Zeitschr.* xxviii. (1878) 215.—Afr. trop.
debilis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 150.—Afr. trop.
decumbens, Zipp. ex Span. in Linnaea, xv. (1841) 193 = sensitiva.
densiflora, *Benth. in Hook. Lond Journ. Bot.* ii. (1840) 56.—Guian.
dentata, Lag. Gen. et Sp. Nov. 22 = Adesmia dentata.
diffusa, Roxb. ex Wight & Arn. Prod. 215 = Sesbania procumbens.
diffusa, Klein, ex Willd. Sp. Pl. iii. 1164 = indica.
dimidiata, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 151.—Afr. trop.
divisa, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1826) 31 = americana.
Echinus, *Vog. in Linnaea*, xii. (1838) 92.—Bras.

AESCHYNOMENE :—
elegans, *Cham. & Schlecht. in Linnaea*, v. (1830) 583.—Mexico.
elongata, Salisb. Prod. 341 = Sesbania aegyptiaca.
erubescens, *E. Mey. Comm. Pl. Afr. Austr.* i. 123.—Afr. austr.
evenia, *Wright, in Sauv. Fl. Cub.* 27.—Cuba.
falcata, *DC. Prod.* ii. 322.—Bras.
fascicularis, *Cham. & Schlecht. in Linnaea*, v. (1830) 584.—Mexic.
filipes, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 415.—Madag.
filosa, *Mart. ex Benth. in Mart. Fl. Bras.* xv. 1. 61.—Bras.
fistulosa, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 259.—Porto Rico.
floribunda, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 186 = americana.
fluminensis, *Vell. Fl. Flum.* 310; vii. t. 119.—Bras.
frutescens, Poir. in Lam. Encyc. iv. 451 = Lespedeza capitata.
fruticulosa, Nees, ex G. Don, Gen. Syst. ii. 282 = Adesmia fruticulosa.
fulgida, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 149.—Afr. trop.
fusa, Steud. Nom. ed. I. 17, sphalm. = seq..
fusca, *Desf. Tabl. Cat. Hort. Par.* ed. I. 198.—Hab. ?
gangetica, Poir. in Lam. Encyc. iv. 453 = Desmodium gangeticum.
glaberrima, Poir. Encyc. Suppl. iv. 76 = hispida ?
glabrescens, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 148.—Afr. trop.
glandulosa, Benth. in Miq. Pl. Jungh. 210 = javanica.
glandulosa, Poir. Encyc. Suppl. iv. 76 = americana.
gracilis, Miq. in Linnaea, xviii. (1844) 566 = paniculata.
gracilis, Vog. l. c. xii. (1838) 891 = falcata.
grandiflora, Linn. Sp. Pl. ed. II. 1050 = Sesbania grandiflora.
guayaquilensis, *G. Don, Gen. Syst.* ii. 284.—Ecuador.
hedysaroides, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 180.—Mexic.
herbacea, *Aubl. Pl. Gui.* ii. 775.—Guian.
heterophylla, *Lour. Fl. Cochinch.* 446.—Cochinch.
Heurckeana, *Baker, in Journ. Linn. Soc.* xx. (1884) 130.—Madag.
hirsuta, Vitm. Summa Pl. iv. 270 = Burmanni.
hirsuta, DC. Prod. ii. 322 = americana.
hirta, *Lam. Illustr.* t. 629, *ex ejusd. Encyc.* iv. 450.—Ind. or.
hispida, Roxb. ex Steud. Nom. ed. II. i. 31 = Robinia hispida.
hispida, *Willd. Sp. Pl.* iii. 1163.—Am. bor.
hispidula, *H. B. & K. Nov. Gen. et Sp.* vi. 530.—N. Granat.
hispidula, Lag. Gen. et Sp. Nov. 22. n. 297 = Adesmia hispidula.
Histrix, *Poir. Encyc. Suppl.* iv. 77.—Guian.
honesta, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1826) 32 = sensitiva.
incana, G. F. W. Mey. ex DC. Prod. ii. 332 = Desmodium incanum.
incana, *Vog. in Linnaea*, xii. (1838) 90.—Bras.
indica, Burm. f. Fl. Ind. 169 = Sesbania aegyptiaca.
indica, *Linn. Sp. Pl.* 713.—Ind. or.
indica, Wall. Cat. n. 5667 = aspera.
interrupta, *Benth. in Hook. Lond. Journ. Bot.* ii. (1840) 56.—Guian.
javanica, *Miq. Fl. Ind. Bat.* i. 275.—Java.
kashmiriana, Baker, in Hook. f. Fl. Brit. Ind. i. 151 = indica.
laevis, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1845) 180.—Mexico.
Lagenaria, Lour. Fl. Cochinch. ii. 446 = aspera.
laevis, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. *Art.* iv. 6.—Malaya.
latifolia, Spreng. Syst. ii. 322 = Adesmia latifolia.
latisiliquosa, Hill, ex Steud. Nom. ed. II. i. 31 = Poiretia latisiliquosa.
laxiflora, *Boj. ex Baker, in Journ. Linn. Soc.* xx. (1884) 130.—Madag.
leptostachya, *Benth. in Mart. Fl. Bras.* xv. 1. 65.—Am. austr.
littoralis, *Vog. in Linnaea*, xii. (1838) 96.—Bras.
longifolia, Cav. Ic. iv. 8 = Piscidia longifolia.
longifolia, Orteg. Hort. Matr. Dec. 70 = Sesbania Cavanillesii.

AESCHYNOMENE:—
macropoda, DC. Prod. ii. 320 = indica.
maculata, Poir. in Lam. Encyc. iv. 452 = Desmodium gangeticum.
marginata, *Benth. in Mart. Fl. Bras.* xv. 1. 66.—Am. trop.
Martii, *Benth. l. c.* 62.—Am. trop.
Martiusii, *Steud. Nom.* ed. II. i. 31.—Bras.
mazangayana, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 415.—Madag.
micrantha, *DC. Prod.* ii. 321.—Madag.
microphylla, *Desv. ex DC. l. c.* 322.—Bras.
mimosoides, Necs, ex G. Don, Gen. Syst. ii. 282 = Adesmia mimosoides.
mimulosa, Blume, ex Miq. Fl. Ind. Bat. i. 276 = javanica.
miniata, Orteg. Hort. Matr. Dec. 28 = Piscidia punicea.
mimosifolia, *Vatke, in Oest. Bot. Zeitschr.* xxix. (1879) 224.—Afr. trop.
mollicula, *H. B. & K. Nov. Gen. et Sp.* vi. 532.—Am. austr.
moluccana, *Kostel. Allg. Med.-pharm. Flora*, iv. 1285.—Molucca.
montana, Span. in Linnaea, xv. (1841) 192 = indica.
montevidensis, *Vog. in Linnaea*, xii. (1838) 83.—Reg. Argent.
mucronata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1840) 56.—Guian.
mucronulata, *Benth. l. c.* 56.—Guian.
multiflora, *Steud. Nom.* ed. II. i. 31.—Ind. or.
obovalis, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 415.—Madag.
oligantha, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 146.—Afr. trop.
oroboides, *Benth. in Mart. Fl. Bras.* xv. 1. 64.—Am. austr.
paludosa, Roxb. Fl. Ind. iii. 333 = Sesbania aculeata.
paniculata, *Willd. ex Vog. in Linnaea*, xii. (1838) 95.—Bras.
papposa, Lag. Gen. et Sp. Nov. 23 = Adesmia papposa.
papulosa, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 146.—Afr. trop.
parviflora, *Micheli, in Kjoeb. Vidensk. Meddel.* (1875) 66.—Bras.
patagonica, Hort. ex Steud. Nom. ed. II. i. 31 = Adesmia muricata.
patula, Poir. Encyc. Suppl. iv. 78 = sensitiva.
pauciflora, *Vog. in Linnaea*, xii. (1838) 93.—Bras.
paucifolia, *Vog. l. c.* 94.—Bras.
paucijuga, DC. Prod. ii. 321 = brasiliana.
picta, Cav. Ic. iv. 7 = Sesbania aegyptiaca.
pilosa, *Poir. in Lam. Encyc.* iv. 450.—Ind. or.
platycarpa, *Benth. in Mart. Fl. Bras.* xv. 1. 63.—Peruv.
platycarpa, Michx. Fl. Bor. Am. ii. 75 = Sesbania vesicaria.
pleuronervia, *DC. Prod.* ii. 321.—Ins. S. Doming.
pluriarticulata, *G. Don, Gen. Syst.* ii. 284.—Ecuador.
podocarpa, Vog. in Linnaea, xii. (1838) 89 = falcata.
procumbens, Roxb. Fl. Ind. iii. 337 = Sesbania procumbens.
prostrata, Poir. Encyc. Suppl. iv. 76 = viscidula.
pseudoacacia, Roxb. ex DC. Prod. ii. 261 = Robinia Pseudacacia.
pseudoviscosa, Blume, ex Miq. Fl. Ind. Bat. i. 276 = javanica.
puberula, DC. Prod. ii. 321 = scoparia.
pubescens, *Poir. in Lam. Encyc.* iv. 450.—Ind. or.
pubescens, Vahl, ex DC. Prod. ii. 265 = Sesbania pubescens.
pudica, *F. G. Dietr. Lexik. Gaertn.* i. 136.—Aegypt.
pudica, *Zoll. & Morr. in Nat. en Geneesk. Arch. Neerl. Ind.* iii. (1846) 55, nomen.—Java.
pulchella, *Planch. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 149.—Afr. trop.
pulchra, *Vatke, in Oest. Bot. Zeitschr.* xxix. (1879) 250.—Afr. trop.
pumila, Linn. Sp. Pl. ed. II. 1061 = indica.
punctata, Steud. Nom. ed. I. 17 = indica.
pygmaea, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 148.—Afr. trop.
quadrata, Schum. & Thonn. Beskr. Guin. Pl. 356 = indica.
racemosa, *Vog. in Linnaea*, xii. (1838) 92.—Bras.
remota, Poir. in Lam. Encyc. iv. 452 = Desmodium Scalpe.
rostrata, *Benth. in Mart. Fl. Bras.* xv. 1. 60.—Bras.
Roxburghii, Spreng. Syst. iii. 322 = indica.
rudis, Benth. Pl. Hartw. 116 = hispidula.

AESCHYNOMENE:—
Ruppellii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 149.—Afr. trop.
scabra, *G. Don, Gen. Syst.* ii. 284.—Ecuador.
Schimperi, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 202.—Abyss.
Scoparia, *H. B. & K. Nov. Gen. et Sp.* vi. 532—Peruv.
scoparia, *Splitgerb. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 107.—Bras.
Selloi, *Vog. in Linnaea*, xii. (1838) 82.—Bras.
sensitiva, Beauv. Fl. Owar. 89. t. 53 = indica.
sensitiva, *Sw. Fl. Ind. Occ.* iii. 1276.—Am. trop.
Sesban, Jacq. Collect. ii. 283 = Sesbania aculeata.
Sesban, Linn. Sp. Pl. 714 = Sesbania aegyptiaca.
siifolia, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 151.—Afr. trop.
spicata, Poir. in Lam. Encyc. iv. 452 = Desmodium supinum.
spinulosa, Roxb. Fl. Ind. iii. 333 = Sesbania aculeata.
subviscosa, DC. Prod. ii. 321 = indica.
sulcata, H. B. & K. Nov. Gen. et Sp. vi. 530 = sensitiva.
surattensis, *Wight & Arn. Prod.* 219 = Sesbania aculeata?
Suymenta, Steud. Nom. ed. II. i. 31 = Sesbania aegyptiaca.
tecta, Vog. in Linnaea, xii. (1838) 87 = falcata.
tenuirama, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 150.—Afr. trop.
tenuis, *Griseb. Cat. Pl. Cub.* 72.—Cuba.
ternata, Spreng. ex DC. Prod. ii. 314 = Pictetia ternata.
timoriana, Span. Ic. 62; et in Linnaea, xv. (1841) 193 = sensitiva.
trachyloba, Miq. Fl. Ind. Bat. i. 276 = aspera.
tribuloides, *Baill. in Bull. Soc. Linn. Par.* i. (1884) 414.—Madag.
triflora, Poir. in Lam. Encyc. iv. 451 = Desmodium triflorum.
triquetra, St. Hil. ex Steud. Nom. ed. II. i. 32 = Desmodium triquetrum.
trisperma, *Klotzsch, in Schomb. Faun. & Fl. Guy.* 1103.—Guian.
tuberculata, *Griseb. Cat. Pl. Cub.* 72.—Cuba.
uliginosa, Roxb. Fl. Ind. iii. 334 = Sesbania aculeata.
uniflora, *E. Mey. Comm. Pl. Afr. Austr.* i. 123.—Afr. austr.
versicolor, *Wender. in Bot. Zeit.* i. (1843) 347.—Mexic.
villosa, *Poir. Encyc. Suppl.* iv. 76.—Ins. Porto-Rico.
violacea, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. *Art.* iv. 5.—Malaya.
virgata, Cav. Ic. iii. 47 = Coursetia virgata.
viscidula, *Michx. Fl. Bor. Am.* ii. 75.—Florida.
viscidula, Roxb. ex Willd. Enum. Hort. Berol. 776 = indica.
viscosa, *Vatke, in Bremen Abh.* vii. (1882) 247.—Afr. trop.
Warmingii, *Micheli, in Kjoeb. Vidensk. Meddel.* (1875) 68.—Bras.

AESCULUS, Linn. Syst. ed. II. 22 (1740). *SAPINDACEAE*, Benth. & Hook. f. i. 398.
BILLIA, Peyritsch, in Bot. Zeit. xvi. (1858) 153.
CALOTHYRSUS, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 62.
ESCULUS, Linn. Gen. ed. I. 109 (1737).
HIPPOCASTANUM, Tourn. ex Rupp. Fl. Jen. ed. Hall. 291 (1745).
ISYPUS, Rafin. Fl. Tellur. iv. 73 (1836).
MACROTHYRSUS, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 61.
NEBROPSIS, Rafin. Alsog. Am. 68 (1838).
OESCULUS, Neck. Elem. ii. 232 (1790).
OZOTIS, Rafin. Alsog. Am. 71 (1838).
PAVIA, Boerh. ex Mill. Gard. Dict. ed. VI. (1752).
PAVIANA, Rafin. Fl. Ludov. 87 (1817).
PUTZEYSIA, Planch. & Linden, in Linden, Cat. n. 22 (1857), 3.
alba, *Rafin. Alsog. Am.* 69.—Am. bor.
arguta, *Buckl. in Proc. Acad. Phil.* (1860) 443.—Am. bor.
asplenifolia, Hort. ex Loud. Encyc. Trees, 124 = A. Hippocastanum.
assamica, Griff. Notul. iv. 540 = punduana.
barbata, *Rafin. Alsog. Am.* 70.—Am. bor.
californica, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 251.—Calif.
carnea, Wats. Dendr. Brit. t. 121 = glabra.
Castanea, Gilib. Fl. Lituan. ii. 197 = A. Hippocastanum.

AESCULUS:—
chinensis, *Bunge, in Mém. Sav. étr. Pétersb.* ii. (1832–84.—China.
discolor, Pursh, Fl. Am. Sept. i. 255 = flava.
dissimilis, *Blume, Rumphia*, iii. 195.—Japon.
echinata, Muhl. Cat. Pl. Am. Sept. 38 = glabra.
flava, *Ait. Hort. Kew.* i. 494.—Am. bor.
glabra, *Willd. Enum. Hort. Berol.* 405.—Am. bor.
Hippocastanum, *Linn. Sp. Pl.* 344.—Tnrc.
humilis, Lindl. Bot. Reg. t. 1018 = Ac. Pavia?
hybrida, *DC. Hort. Monsp.* (1813) 75 = flava.
indica, *Coleb. ex Wall. Cat.* n. 1181.—Reg. Himal.
intermedia, *Steud. Nom.* ed. II. i. 32.—Am. bor.
levigata, *Rafin. Alsog. Am.* 70.—Am. bor.
livida, *Steud. Nom.* ed. II. i. 32.—Am. bor.
lutea, Wangh. in Schr. Berl. Ges. Naturf. Fr. viii. (1788) 133. t. 6 = flava.
Lyonii, Loud. Arb. Brit. i. 468 = A. Pavia.
macrostachya, Michx. Fl. Bor. Am. i. 220 = parviflora.
macrostachys, Pers. Syn. i. 403 = parviflora.
Memmingeri, Hort. ex C. Koch, Dendrol. i. 506 = A. Hippocastanum.
mexicana, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* i. 213.—Mexic.
muricata, *Rafin. Alsog. Am.* 68.—Am. bor.
neglecta, Lindl. Bot. Reg. t. 1009 = flava.
ochroleuca, *Rafin. Alsog. Am.* 68.—Am. bor.
odorata, *F. G. Dietr. Lexik. Gaertn. Nachtr.* i. 118.—Am. bor.
ohioensis, Michx. Arb. For. Am. Sept. iii. 242 = glabra.
pallida, Willd. Enum. Hort. Berol. 406 = glabra.
Parryi, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 200.—Am. bor.
parviflora, *Walt. Fl. Carol.* 128.—Am. bor.
Pavia, *Linn. Sp. Pl.* 344.—Am. bor.
procera, Salisb. Prod. 280 = A. Hippocastanum.
punduana, *Wall. Cat. n.* 1189.—Reg. Himal.
rosea, Hort. ex Loud. Encyc. Trees, 127 = glabra.
rubella, Wender. Ind. Sem. Marb. (1853) 1. et 4 = glabra?
rubescens, Tausch, in Flora, xxi. (1838) 1. Beibl. 79 = A. Pavia?
rubicunda, Lodd. Bot. Cab. t. 1242 = glabra.
rubra, Pers. Syn. i. 403 = A. Pavia.
septenata, Stokes, Bot. Mat. Med. ii. 337 = A. Hippocastanum.
turbinata, *Blume, Rumphia*, iii. 195.—Japon.
verrucosa, *Rafin. Alsog. Am.* 69.—Am. bor.
versicolor, Wender. Ind. Sem. Hort. Marb. (1853) 4 = A. Pavia?
Watsoniana, D. Dietr. Syn. ii. 1225 = glabra.
Whitleyi, Hort. ex C. Koch, Dendrol. i. 511 = A. Pavia.

AETHEILEMA, R. Br. Prod. 478 (1810) = **Phaylopsis**, Willd. (Acanth.).
alopecuroides, Spreng. Syst. ii. 826 = Lepidagathis alopecuroidea.
anisophyllum, E. Mey. ex Nees, in DC. Prod. xi. 262 = P. longifolia.
glutinosum, Stend. Nom. ed. II. i. 32 = P. parviflora.
Haenkei, Nees, in Mart. Fl. Bras. ix. 72 = Lepidagathis alopecuroidea.
imbricatum, R. Br. Prod. i. 478 = P. longifolia.
javanicum, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 129.—Java.
longifolium, Spreng. Syst. Cur. Post. 236 = P. longifolia.
micranthum, Benth. in Hook. Nig. Fl. 480.—Afr. trop.
mucronatum, Griff. Notul. iv. 137 = P. parviflora.
parviflorum, Spreng. Syst. ii. 826 = P. parviflora.
Rothii, Steud. Nom. ed. II. i. 32 = P. parviflora.
reniforme, Nees, in Wall. Pl. As. Rar. iii. 94 = P. parviflora.
rigidum, Bartl. ex Nees, in DC. Prod. xi. 243 = Barleriola solanifolia.
rupestre, Nees, l. c. 262.—Madag.

AETHEOLAENA, Cass. in Dict. Sc. Nat. xlviii. 450 (1827) = **Senecio**, Linn. (Compos.).
involucrata, Cass. l. c. 451 = S. involucratus.

AETHEONEMA, Reichb. Nom. 133 (1841) = **Gaertnera**, Lam. (Logan.).

AETHEOPAPPUS, Cass. in Dict. Sc. Nat. l. 250 (1827) = **Centaurea**, Linn. (Compos.).
Balansae, Boiss. Fl. Orient. iii. 604 = C. Balansae.
pulcherrimus, Cass. in Dict. Sc. Nat. li. 52 = C. pulcherrima.

AETHEORHIZA, Cass. in Dict. Sc. Nat. xlviii. 425 (1827) = **Crepis**, Linn. (Compos.).
bulbosa, Cass. l. c. = C. bulbosa.
montana, Willk. in Oest. Bot. Zeitschr. xxv. (1875) 110 = C. montana.

AETHEORRHIZA, Reichb. Consp. 98 (1828) = praec.

AETHERIA, Endl. Gen. 214 (1837) = **Hetaeria**, Blume (Orchid.).

AETHIONEMA, R. Br. in Ait. Hort. Kew. ed. II. iv. 80 (1812). *CRUCIFERAE*, Benth. & Hook. f. i. 88.
Diastrophis, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 35 (1835).
Disynoma, Rafin. Fl. Tellur. iii. 81 (1836).
Ethionema, Brongn. Enum. Gen. 92 (1843).
Eunomia, DC. Syst. ii. 555 (1821).
Oethionema, Knowl. & Westc. Flor. Cab. i. (1837) 41.
arabicum, Andrz. ex DC. Syst. ii. 560 = cappadocicum.
armenum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 191.—Armen.
Balansae, Boiss. Diagn. Ser. II. vi. 20 (= *Crenularia glaucescens*).
banaticum, Janka, in Linnaea, xxx. (1859–60) 558 = saxatile.
Beyrichii, *Tausch, in Flora*, xix. (1836) 411.—Persia.
Bourgaei, *Boiss. Fl. Orient.* i. 344.—Syria.
Buxbaumii, DC. Syst. ii. 560 = cappadocicum.
caespitosum, Boiss. Fl. Orient. i. 343 = Iberidella caespitosa.
campylopterum, *Boiss. l. c.* 353.—Syria.
capitatum, *Boiss. & Balansa, Diagn.* Ser. II. v. 43.—As. Min.
cappadocicum, *Spreng. Nov. Prov.* i. 2.—Oriens.
cardiophyllum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 43.—As. Min.
chloraefolium, *Boiss. Fl. Orient.* i. 344.—As. Min.
clandestinum, *Delile, in Ann. Sc. Nat.* Sér. II. vii. (1837) 285.—Hab.?
cordatum, *Boiss. Fl. Orient.* i. 350.—Oriens.
coridifolium, *DC. Syst.* ii. 561.—Oriens.
coridifolium, Hort. ex Boiss. Diagn. Ser. I. viii. 43 = pulchellum.
cristatum, *DC. Syst.* ii. 560.—Oriens.
diastrophis, *Bunge, Del. Sem. Dorp.* (1841) 7.—Armen.
elongatum, *Boiss. Diagn.* Ser. I. vi. 17.—Persia.
fimbriatum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 193.—Persia.
Fischeri, *Tausch, in Flora*, xix. (1836) 410.—Sibir.
gracile, *DC. Syst.* ii. 559.—Carniol.
graecum, *Boiss. & Sprun. Diagn.* Ser. I. vi. 16.—Graecia.
grandiflorum, *Boiss. & Hohen. l. c.* viii. 42.—Persia.
heterocarpum, *J. Gay, in Fisch. & Mey. Ind. Sem. Petrop.* iv. 1, nomen.—Syria.
heterophyllum, Boiss. Fl. Orient. i. 343 = Iberidella heterophylla.
iberideum, *Boiss. l. c.* i. 351.—Oriens.
lacerum, *Boiss. Diagn.* Ser. II. v. 45.—Cilicia.
leptocarpium, *Delile, in Ann. Sc. Nat.* Sér. II. vii. (1837) 285.—Hab.?
lignosum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 348.—As. Min.
membranaceum, *DC. Syst.* ii. 561.—Persia.
monospermum, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 80.—Hispan.
moricandianum, *Boiss. Fl. Orient.* i. 351.—As. Min.
oppositifolium, *Boiss. l. c.* 344.—Syria.
ovalifolium, *Boiss. l. c.* 351.—Oriens.
polygaloides, *DC. Syst.* ii. 562.—Ins. Scios.
polygaloides, Ledeb. Fl. Ross. i. 209 = armenum.
pulchellum, *Boiss. & Huet, Diagn.* Ser. II. v. 43.—Oriens.
pyrenaicum, *Boutigny, in Bull. Soc. Bot. Fr.* iv. (1857) 777.—Mont. Pyren.
rotundifolium, *Boiss. Fl. Orient.* i. 344.—Reg. Caucas.
rubescens, *Boiss. l. c.* 343.—As. Min.
sagittatum, Boiss. l. c. 342 = Iberidella sagittata.
salmasium, *Boiss. l. c.* 350.—Persia.
saxatile, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 80.—Europ. austr.
schistosum, *Boiss. & Kotschy, Diagn.* Ser. II. v. 42.—As. Min.
speciosum, *Boiss. & Huet, l. c.* 44.—Oriens.

AETHIONEMA:—
spicatum, *Post, in Hook. Ic. Pl.* xv. 62, t. 1478.—Syria.
stenopterum, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 192.—Persia.
stylosum, *DC. Syst.* ii. 562.—Syria.
subulatum, *Boiss. Fl. Orient.* i. 349.—As. Min.
Szowitzii, *Boiss. l. c.* 345.—Persia.
tenue, Boiss. l. c. 343 = Iberidella tenuis.
thesiifolium, *Boiss. & Heldr. Diagn.* Ser. I. viii. 44.—As. Min.
Thomasianum, *J. Gay, in Ann. Sc. Nat.* Sér. III. iv. (1845) 81.—Europ.
trinervium, Boiss. Fl. Orient. i. 342 = Iberidella trinervia.

AETHIOPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 134 = **Salvia**, Tourn. (Labiat.).
Sclarea, Fourr. l. c. = S. Sclarea.
vera, Fourr. l. c. = S. Aethiopis.

AETHONIA, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 309 = **Tolpis**, Adans. (Compos.).
filiformis, D. Don, l. c. = T. filiformis.
fruticosa, D. Don, l. c. = T. fruticosa.
lagopoda, D. Don, l. c. = T. lagopoda.

AETHUSA, Linn. Sp. Pl. 256 (1753). *UMBELLIFERAE*, Benth. & Hook. f. i. 907.
Cynapium, Riv. ex Rupp. Fl. Jen. ed. Hall. 284 (1745).
Ethusa, Linn. Syst. ed. I. (1735).
Ammi, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 260 = Ammi majus.
Bunius, Murr. Syst. Veg. ed. XIII. 236 = Carum Bunius.
capillacea, Nutt. ex DC. Prod. iv. 106 = Apiastrum angustifolium.
Cicuta, Neck. Delic. Gallo-Belg. i. 146 = A. Cynapium.
cynapioides, Bieb. Fl. Taur. Cauc. i. 227 = A. Cynapium.
Cynapium, *Linn. Sp. Pl.* 256.—Europ.; Oriens.
cynica, Dulac, Fl. Hautes-Pyr. 353 = A. Cynapium.
divaricata, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 260 = Apium divaricatum.
elata, Friedl. ex Fisch. Cat. Hort. Gorenk. (1813) 45 = A. Cynapium.
fatua, Ait. Hort. Kew. i. 352 = Carum verticillatum.
leptophylla, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 260 = Apium leptophyllum.
Meum, Murr, Syst. Veg. ed. XIII. 236 = Meum athamanticum.
micrantha, Opiz, ex Nym. Consp. 297 = A. Cynapium.
montana, Lam. Fl. Fr. iii. 649 = Carum Bunius.
mutellina, Lam. l. c. 443 = Meum mutellina.
petroselinifolia, Gilib. Fl. Lituan. ii. 25 = A. Cynapium.
pinnata, Nutt. ex DC. Prod. iv. 141 = Cynosciadium pinnatum.
segetalis, Boenn. Prod. Fl. Monast. 85 = A. Cynapium.
tenuifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 513 = A. Cynapium.
tenuifolium, Salisb. Prod. 166 = Meum athamanticum.
toxicaria, Salisb. l. c. = A. Cynapium.

AETIA, Adans., Fam. ii. 84 (1763) = **Combretum**, Linn.

AEXTOXICON, Ruiz & Pav. Prod. 131. t. 29 (1794). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 285.
Aegotoxicon, Molina, Sagg. Chile. ed. II. 298 (1810).
punctatum, *Ruiz & Pav. Syst. Veg.* 260.—Chili.

AFARCA, Rafin. Sylva Tellur. 30 (1838) = **Sageretia**, Brongn. (Rhamn.).
parviflora, Rafin. l. c. = S. Michauxii.

AFFONSEA, A. St. Hil. Voy. Diam. i. 387 (1833). *LEGUMINOSAE*, Benth. & Hook. f. i. 599.
bullata, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 77.—Bras.
comosa, Benth. l. c. 76. t. 1 = juglandifolia.
densiflora, *Benth. l. c.*—Bras.
juglandifolia, *A. St. Hil. Voy. Diam.* i. 385.—Bras.

AFZELIA, J. F. Gmel. Syst. 927 (1791) = **Seymeria**, Pursh (Scrophul.).
cassioides, J. F. Gmel. l. c. = S. tenuifolia.

AFZELIA, Sm. in Trans. Linn. Soc. iv. (1798) 221. *LEGUMINOSAE*, Benth. & Hook. f. i. 580.
Intsia, Thou. Gen. Nov. Madag. 22 (1809).
africana, *Sm. l. c.*—Afr. trop.
attenuata, *Klotzsch, in Peters, Reise Mossamb. Bot.* i. 20.—Afr. or. trop.
bijuga, *A. Gray, Bot. U. St. Expl. Exped.* i. 467.—Ins. Pacif.
bijuga, Spreng. Syst. iv. Cur. Post. 170 = Erioglossum cauliflorum.
bracteata, *Vog. ex Benth. in Hook. Ic. Pl.* t. 790.—Afr. trop.
cambodensis, *Hance, in Journ. Bot.* xiv. (1876) 258.—Cambod.
coriacea, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 275.—Ind. or.
cuanzensis, Oliver, Fl. Trop. Afr. ii. 302 = quanzensis.
discolor, *Hort. ex Steud. Nom.* ed. II. i. 33.—Afr. trop.
grandis, Hort. ex Loud. Hort. Brit. 168 = Erythrophleum guineense.
madagascariensis, Baill. Hist. Pl. ii. 112 = bijuga.
palembanica, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 275.—Ind. or.
Pancovia, DC. Prod. ii. 507 = Erioglossum cauliflorum.
Petersiana, Klotzsch, in Peters, Reise Mossamb. Bot. i. 19 = quanzensis.
quanzensis, *Welw. Apont.* 586.—Angola.
retusa, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 73.—Malacca.
rhomboidea, *Vidal, Phanerog. Cuming. Philipp.* 26, 110.—Ins. Philipp.
splendens, Hort. ex Loud. Hort. Brit. 168 = Carapa procera.

AGALINIS, Rafin. New Fl. Am. ii. 61 (1836) = **Gerardia**, Linn. (Scrophul.).
aphylla, Rafin. l. c. 65 = G. aphylla.
corymbosa, Rafin. l. c. 63 = G. Plukenetii.
fasciculata, Rafin. l. c. = G. purpurea.
filifolia, Rafin. l. c. 65 = G. filifolia.
longifolia, Rafin. l. c. 62 = G. aspera?
maritima, Rafin. l. c. = G. maritima.
microphylla, Rafin. l. c. 65.—Am. bor.
obtusifolia, Rafin. l. c. 64 = Aulaya obtusifolia?
palustris, Rafin. l. c. 62 = G. purpurea.
perennis, Rafin. l. c. 63 = G. linifolia.
Plukeneti, Rafin. l. c. = G. Plukenetii.
setacea, Rafin. l. c. 64 = G. setacea.
tenuifolia, Rafin. l. c. = G. tenuifolia.
virgata, Rafin. l. c. 62 = Seymeria virgata?

AGALLIS, Phil. in Linnaea, xxxiii. (1864–65) 12. *CRUCIFERAE*.
montana, *Phil. l. c.*—Chili.

AGALLOCHUM, Lam. Encyc. i. 48 (1783) = **Aquilaria**, Lam. (Thymel.).
officinarum, Lam. l. c. = Aquil. malaccensis.
praestantissimum, Lam. l. c. 48.—As. or.
sylvestre, Lam. l. c. 49.—Ins. Timor.

AGALLOSTACHYS, Beer, Bromel. 35 (1857) = **Bromelia**, Linn.
antiacantha, Beer, l. c. 37 = B. fastuosa.
chrysantha, Beer, l. c. 38 = B. chrysantha.
Commeliniana, Beer, l. c. 39 = B. fastuosa.
fastuosa, Beer, l. c. 36 = B. fastuosa.
laciniosus, C. Koch, Wochenschr. ii. (1859) 140 = B. laciniosa.
Pinguin, Beer, Bromel. 36 = B. Pinguin.
sylvestris, Beer, l. c. 35 = B. sylvestris.
variegata, Beer, l. c. 38 = Billbergia speciosa.

AGALMA, Miq. Fl. Ind. Bat. i. 1. 751 (1855) = **Heptapleurum**, Gaertn. (Araliac.).
aesculifolium, Seem. in Journ. Bot. ii. (1864) 298 = Brassaiopsis aculeata.
aromaticum, Seem. l. c.—Java.
elatum, Seem. l. c. = H. elatum.
glaucum, Seem. l. c. 299 = H. glaucum.
Griffithii, Seem. l. c. = H. elatum.
Horsfieldii, Seem. l. c. 298.—Java.
kavaiense, Seem. Revis. Hederac. 103 = H. kavaiense.
lucescens, Seem. in Journ. Bot. ii. (1864) 299 = H. octophyllum.
octophyllum, Seem. l. c. 298 = H. octophyllum.
racemosum, Seem. l. c. = H. racemosum.
redivivum, Seem. l. c. 299.—Borneo.

AGALMA :—
rostratum, Seem. in Journ. Bot. ii. (1864) 298 = H. rostratum.
rugosum, Miq. Fl. Ind. Bat. i. 1. 752. t. 11.—Malaya.
simillimum, Miq. l. c.—Malaya.
tomentosum, Seem. in Journ. Bot. ii. (1864) 298 = H. impressum.
vitiense, Seem. Fl. Vit. 116.—Ins. Fiji.

AGALMA, Steud. Nom. ed. II. i. 33 (1840) = **Sonchus**, Tourn. (Compos.).

AGALMANTHUS, Hombr. & Jacquinot, ex Decne. Pot. Voy. Astrol. et Zel. 78 (1845) = **Metrosideros**, Banks (Myrtac.).
umbellata, Hombr. & Jacquinot, ex Decne. l. c. = M. lucida.

AGALMYLA, Blume, Bijdr. 766 (1826). *GESNERACEAE*, Benth. & Hook. f. ii. 1014.
Orithalia, Blume, Fl. Jav. Praef. 6 (1828).
Orythia, 'Blume,' ex Endl. Gen. Suppl. i. 1408 (1841).
angustifolia, *Miq. Fl. Ind. Bat.* ii. 723.—Malaya.
asperifolia, Blume, Bijdr. 767 = Dichrotrichum asperifolium.
Beccarii, *C. B. Clarke, in DC. Monog. Phan.* v. 57.—Sumatra.
longistyla, Carr. in Rev. Hortic. (1873) 271 = staminea.
staminea, *Blume, Bijdr.* 767.—Java.
tuberculata, *Hook. f. Ic. Pl.* t. 897.—Borneo.

AGALOMA, Rafin. Fl. Tellur. iv. 116 (1836) = **Euphorbia**, Linn.
angustifolia, Rafin. l. c. 117 = E. angustifolia?
corollata, Rafin. l. c. = E. corollata.
graminifolia, Rafin. l. c. = E. granulata.

AGANIPPEA, Moç. & Sesse, ex DC. Prod. vi. 3 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 358.
Heliogenes, Benth. Pl. Hartw. 42 (1840).
bellidiflora, *Moç. & Sesse, ex DC. Prod.* vi. 3.—Mexic.
dentata, *DC. l. c.*—Mexic.

AGANISIA, Lindl. in Bot. Reg. xxv. (1839) Misc. 45; xxvi. t. 32 (1840). *ORCHIDEAE*, Benth. & Hook. f. iii. 544.
Koellensteinia, Reichb. f. in Bonplandia, ii. (1854) 17.
caerulea, *Reichb. f. in Gard. Chron.* N. S. vi. (1876) ii. 226.—Bras.
cyanea, Reichb. f. in Nov. Act. Nat. Cur. xxxv. ii. (1876) 13. t. 4 = Acacallis cyanea.
fimbriata, *Reichb. f. in Gard. Chron.* N. S. ii. (1874) ii. 452.—Guian.
lepida, *Linden & Reichb. f. in Nov. Act. Nat. Cur.* xxxv. ii. (1876) 15. t. 5.—Am. trop.
Oliveriana, *Reichb. f. in Gard. Chron.* N. S. ix. (1878) i. 558.—Bras.
pulchella, *Lindl. in Bot. Reg.* (1839) *Misc.* 45; xxvi. (1840) t. 32.—Guian.

AGANON, Rafin. Sylva Tellur. 161 (1838) = **Callicarpa**, Linn. (Verben.).
umbellata, Rafin. l. c. = C. arborea.

AGANOPE, Miq. Fl. Ind. Bat. i. 1. 151 (1855) = **Derris**, Lour. (Legumin.).
floribunda, Miq. l. c. = D. thyrsiflora.
marginata, Miq. l. c. 152 = D. marginata.
microphylla, Miq. l. c.—Java.
subavenis, Miq. l. c. Suppl. i. 299.—Sumatra.
sumatrana, Miq. l. c. 300.—Sumatra.

AGANOSMA, G. Don, Gen. Syst. iv. 77 (1837). *APOCYNACEAE*, Benth. & Hook. f. ii. 717.
Ganosma, Decne. in DC. Prod. viii. 625 (1844), sphalm.
acuminata, G. Don, l. c. = marginata.
affinis, G. Don, l. c. = Ichnocarpus ovatifolius.
Blumei, *A. DC. in DC. Prod.* viii. 432.—Java.
Blumei, Wight, Ic. t. 1305 = caryophyllata.
calycina, *A. DC. in DC. Prod.* viii. 432.—Ind. or.
caryophyllata, *G. Don, Gen. Syst.* iv. 77.—Ind. or.
concanensis, Hook. Ic. Pl. v. t. 841 = Parsonsia spiralis.
conferta, G. Don, in Loud. Encyc. Pl. Suppl. ii. 1312 = cymosa.
cymosa, *G. Don, Gen. Syst.* iv. 77.—Ind. or.

AGANOSMA :—
Doniana, Wight, Ic. t. 1306 = cymosa.
elegans, G. Don, Gen. Syst. iv. 77 = cymosa.
Edithae, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 227 = Cryptolepis elegans.
euloba, Miq. Fl. Ind. Bat. ii. 447 = acuminata.
gracilis, *Hook. f. Fl. Brit. Ind.* iii. 664.—Reg. Himal.
laevigata, J. Grah. Cat. Bomb. Pl. 113 (= Parsonsiae sp.?).—Ind. or.
laevis, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 335 = Anodendron laeve.
macrocarpa, A. DC. in DC. Prod. viii. 434 = marginata.
marginata, *G. Don, Gen. Syst.* iv. 77.—Ind. or.
Roxburghii, G. Don, l. c. = calycina.
velutina, *A. DC. in DC. Prod.* viii. 434.—Malaya.
Wallichii, G. Don, Gen. Syst. iv. 77 = calycina.

AGAPANTHUS, L'Hérit. Sert. Angl. 17 (1788). *LILIACEAE*, Benth. & Hook. f. iii. 798.
Abumon, Adans. Fam. ii. 54 (1763).
Mauhlia, Dahl, Obs. Syst. Linn. 25 (1787).
africanus, Hoffmgg. Verz. Pfl. 35 = umbellatus.
ensifolius, Willd. Sp. Pl. ii. 48 = Massonia ensifolia.
medius, Lodd. ex Steud. Nom. ed. II. i. 33 = umbellatus.
minor, Lodd. Bot. Cab. t. 42 = umbellatus.
multiflorus, Willd. Enum. Hort. Berol. 353 = umbellatus.
praecox, Willd. l. c. = umbellatus.
tuberosus, Linn. ex [DC. in] Red. Liliac. i. 6 = umbellatus.
umbellatus, *L'Hérit. Sert. Angl.* 18.—Afr. austr.
umbelliferus, Poir. Encyc. Suppl. i. 155 = umbellatus.
variegatus, Hort. ex Steud. Nom. ed. II. i. 33 = umbellatus.

AGAPATEA, Steud. in Bot. Zeit. xiv. (1856) 391 = **Distichia**, Nees & Meyen (Juncac.).
filamentosa, Buchen. in Bremen Abh. iv. (1874) 124 = D. filamentosa.
peruviana, Steud. in Bot. Zeit. xiv. (1856) 390 = D. muscoides.

AGAPETES, G. Don, Gen. Syst. iii. 862 (1834). *VACCINIACEAE*, Benth. & Hook. f. ii. 571.
Caligula, Klotzsch, in Linnaea, xxiv. (1851) 28.
Paphia, Seem. in Journ. Bot. ii. (1864) 77.
Acosta, Dun. in DC. Prod. vii. 556 = Vaccinium Acosta.
acuminata, D. Don, ex G. Don, Gen. Syst. iii. 862 = Corallobotrys acuminata.
amblyornidis, *Becc. Malesia*, i. (1878) 208.—Malaya.
angulata, *Benth. & Hook. f. Gen.* ii. 571.—Ind. or.
arborea, Dun. in DC. Prod. vii. 555 = Vaccinium Leschenaultii.
auriculata, *Benth. & Hook. f. Gen.* ii. 571.—Ind. or.
bracteata, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 448.—Ind. or.
buxifolia, *Nutt. ex Hook. f. Bot. Mag.* t. 5012.—Reg. Himal.
campanulata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 449.—Ind. or.
coriacea, G. Don, Gen. Syst. iii. 863 = Vaccinium Korthalsii.
cuneifolia, G. Don, l. c. = Vaccinium cuneifolium.
discolor, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 448.—Reg. Himal.
elliptica, G. Don, Gen. Syst. iii. 862 = Vaccinium ellipticum.
floribunda, G. Don, l. c. = Vaccinium Teysmanni.
glabra, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 444.—Ind. or.
grandiflora, *Benth. & Hook. f. Gen.* ii. 571.—Ind. or.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 449.—Reg. Himal.
hirsuta, Klotzsch, in Linnaea, xxiv. (1851) 38 = setigera.
laurifolia, G. Don, Gen. Syst. iii. 862 = Vaccinium laurifolium.
lincarifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 449.—Reg. Himal.
Lobbii, *C. B. Clarke, l. c.* 448.—Ind. or.
loranthiflora, *D. Don, ex G. Don, Gen. Syst.* iii. 862.—Ind. or.
lucida, G. Don, l. c. 863 = Vaccinium lucidum.
macrantha, *Benth. & Hook. f. Gen.* ii. 571.—Ind. or.
macrophylla, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 445.—Reg. Himal.
macrostemon, *C. B. Clarke, l. c.* 443.—Burma.
madagascariensis, *Dun. in DC. Prod.* vii. 555.—Madag.

AGAPETES :—
meliphagidum, *Becc. Malesia*, i. (1878) 208.—Malaya.
microphylla, *Jungh. in Nat. en Geneesk. Arch. Neerl. Ind.* ii. (1845) 35.—Java.
miniata, *Benth. & Hook. f. Gen.* ii. 571.—Reg. Himal.
mitrarioides, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 447.—Reg. Himal.
myrtoidea, G. Don, Gen. Syst. iii. 863 = Vaccinium myrtoides.
Myzomelae, *Becc. Malesia*, i. (1878) 209.—Malaya.
nana, *Benth. & Hook. f. Gen.* ii. 571.—Reg. Himal.
Nuttallii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 445.—Reg. Himal.
obliqua, Benth. & Hook. f. Gen. ii. 571 = setigera.
obovata, *Benth. & Hook. l. c.* 572.—Reg. Himal.
odontocera, Benth. & Hook. f. l. c. 571 = variegata.
Parishii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 445.—Ind. or.
pilifera, *Hook. f. ex C. B. Clarke, l. c.* 448.—Reg. Himal.
pulcherrima, Benth. & Hook. f. Gen. ii. 571 = variegata.
rosea, *Jungh. in Nat. en Geneesk. Arch. Neerl. Ind.* ii. (1845) 35.—Java?
salicifolia, *C. B. Clarke in Hook. f. Fl. Brit. Ind.* iii. 445.—Reg. Himal.
salicifolia, Gardn. ex Thw. Enum. Pl. Zeyl. 170 = Vaccinium Leschenaultii.
saligna, *Benth. & Hook. f. Gen.* ii. 571.—Ind. or.
serrata, G. Don, Gen. Syst. ii. 862 = Vaccinium serratum.
setigera, *D. Don, ex G. Don, l. c.*—Ind. or.
Sprengelii, G. Don, l. c. = Vaccinium Donianum.
symplocifolia, D. Don, ex G. Don, l. c. = Vaccinium Leschenaultii.
vaccinacea, Dun. in DC. Prod. vii. 554 = Vaccinium serratum.
variegata, Dun. l. c. = setigera.
varicgata, *D. Don, ex G. Don, Gen. Syst.* iii. 862.—Ind. or.
varingiaefolia, G. Don, l. c. = Vaccinium varingiaefolium.
verticillata, D. Don, ex G. Don, l. c. = setigera.
vulgaris, Jungh. in Nat. en Geneesk. Arch. Neerl. Ind. ii. (1845) 34 = Vaccinium ellipticum.
Wallichiana, Klotzsch, in Linnaea, xxiv. (1851) 38 = setigera.

AGARDHIA, Spreng. Syst. i. 4, 17 (1825) = **Qualea**, Aubl. (Vochys.).
cryptantha, Spreng. l. c. 17 = Q. cryptantha.
grandiflora, Spreng. l. c. = Q. grandiflora?

AGARISTA, DC. Prod. v. 569 (1836) = **Coreopsis**, Linn. (Compos.).
calliopsidea, DC. l. c. = C. calliopsidea.

AGARISTA, D. Don, in G. Don, Gen. Syst. iii. 837 (1834). *ERICACEAE*, Benth. & Hook. f. ii. 586.
Amechania, DC. Prod. vii. 578 (1838).
anastomosans, *G. Don, l. c.* iii. 838.—N. Granat.
bracamorensis, *G. Don, l. c.* 837.—N. Granat.
buxifolia, G. Don, l. c. = Agauria buxifolia.
chlorantha, *G. Don, l. c.* 838.—Bras.
eriophylla, *G. Don, l. c.*—Bras.
eucalyptoides, *G. Don, l. c.* 837.—Bras.
ilicifolia, *G. Don, l. c.* 838.—Peruv.
Itatiaiae, *Wawra, It. Princip. Coburg.* i. 73.—Bras.
multiflora, *G. Don, Gen. Syst.* iii. 837.—Bras.
neriifolia, *G. Don, l. c.* 838.—Bras.
nummularia, *G. Don, l. c.* 837.—Bras.
oleifolia, *G. Don, l. c.* 838.—Bras.
Pistrix, *G. Don, l. c.*—Bras.
Pohlii, *G. Don, l. c.* 837.—Bras.
pulchella, *G. Don, l. c.* 838.—Bras.
pulchra, *G. Don, l. c.* 837.—Bras.
pyrifolia, G. Don, l. c. 838 = Agauria salicifolia.
salicifolia, G. Don, l. c. 837 = Agauria salicifolia.
serrulata, *G. Don, l. c.* 838.—Bras.
Sprengelii, *G. Don, l. c.* 837.—Bras.
subrotunda, *G. Don, l. c.*—Bras.

AGASSIZIA, Chav. Monog. Antirrh. 180. t. 11 (1838) = **Galvezia**, Domb. (Scrophul.).
limensis, Chav. l. c. = G. limensis.

AGASSIZIA, Gray & Engelm. in Proc. Am. Acad. i. (1847) 46 = **Gaillardia**, Fouger. (Compos.).
suavis, A. Gray & Engelm. l. c. 50 = G. simplex.

AGASSIZIA, Spach, Hist. Vég. Phan. iv. 347 (1835) = **Oenothera**, Linn. (Onagrar.).
cheiranthifolia, Spach, l. c. 348 = O. Agassizia.

AGASTA, Miers, in Trans. Linn. Soc. Ser. II. i. 59 (1875) = **Barringtonia**, Forst. (Myrtac.).
asiatica, Miers, l. c. 61 = B. speciosa.
indica, Miers, l. c. 63 = B. speciosa.
splendida, Miers, l. c. 60 = B. speciosa.

AGASTACHYS, R. Br. in Trans. Linn. Soc. x. (1811) 158. *PROTEACEAE*, Benth. & Hook. f. iii. 175.
odorata, *R. Br. l. c.*—Tasmania.

AGASTIANIS, Rafin. in New Fl. Am. iii. 85 (1836) = **Sophora**, Linn. (Legum.).
secundiflora, Rafin. l. c. 86 = S. secundiflora.

AGASYLLIS, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 259 = **Siler**, Scop. (Umbellif.).
caucasica, Spreng. l. c. = S. caucasicum.
Galbanum, Spreng. l. c. = Peucedanum Galbanum.
latifolia, Boiss. Fl. Orient. ii. 980 = S. caucasicum.
salsa, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 259 = Peucedanum redivivum.

AGATEA, A. Gray, in Proc. Am. Acad. ii. (1852) 323 = **Agation**, Brongn. (Violar.).
violaris, A. Gray, l. c. 324 = Agation violare.

AGATEA, Rich, ex A. Gray, Bot. U. S. Expl. Exped. i. 609 (1854) = **Haplopetalum**, A. Gray (Rhizoph.).

AGATHAEA, Cass. in Bull. Soc. Philom. (1815) 175; Dict. Sc. Nat. i. Suppl. 77. t. 89 (1817) = **Felicia**, Cass. (Compos.).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 383 = F. Richardi.
amelloides, DC. Prod. v. 225 = Aster rotundifolius.
amoena, Sch. Bip. ex Walp. Rep. ii. 956 = Aster amoenus.
barbata, DC. Prod. v. 224 = Aster barbatus.
Bergeriana, DC. l. c. 223 = Aster Bergerianus.
brevifolia, DC. l. c. 225.—Afr. austr.
coelestis, Cass. Opusc. i. 3 = Aster rotundifolius.
corymbosa, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 60 = Gymnostephium corymbosum.
dentata, A. Rich. Tent. Fl. Abyss. i. 384 = F. Richardi.
diffusa, DC. Prod. v. 224 = Aster demissus.
Ecklonis, Nees, Gen. et Sp. Ast. 262 = Aster Ecklonis.
elongata, Nees, l. c. 266 = Aster elongatus.
hirta, DC. Prod. v. 224 = Aster hirtus.
hispida, DC. l. c. = Aster elongatus.
Kraussii, Sch. Bip. ex Walp. Rep. ii. 95 = Aster Kraussii.
leiocarpa, DC. Prod. v. 225 = Aster leiocarpus.
linifolia, G. Don, in Loud. Hort. Brit. 349 = Euryops linifolius.
microphylla, Cass. in Bull. Soc. Philom. (1817) 183 = Aster aethiopicus.
natalensis, Sch. Bip. ex Walp. Rep. ii. 956 = Aster natalensis.
rotundifolia, Nees, Gen. et Sp. Ast. 258 = Aster rotundifolius.
scabrida, DC. Prod. v. 225 = Aster scabridus.
serrata, Nees, Gen. et Sp. Ast. 259 = Aster serratus.
spathulata, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. x. 43 = Amellus annuus.
spathulata, Kunze, Pugill. Pl. Ined. i. 9 = Amellus hispidus.
stricta, DC. Prod. v. 224 = F. adfinis.
strigosa, Nees, Gen. et Sp. Ast. 263 = Aster strigosus.
tenera, DC. Prod. v. 224 = Aster tener.
tricolor, Nees, Gen. et Sp. Ast. 267 = Aster tricolor.
Zeyheri, Nees, l. c. 260 = Aster linifolius.

AGATHELPIS, Choisy, in Mém. Soc. Phys. Genèv. ii. (1824) 85, 89. *SELAGINEAE*, Benth. & Hook. f. ii. 1129.
aduncа, *E. Mey. Comm. Pl. Afr. Austr.* i. 252.—Afr. austr.
angustifolia, *Choisy, in Mém. Soc. Phys. Genèv.* ii. (1824) 95.—Afr. austr.
brevifolia, *E. Mey. Comm. Pl. Afr. Austr.* i. 253.—Afr. austr.
mucronata, *E. Mey. l. c.*—Afr. austr.

AGATHELPIS :—
nitida, *E. Mey. Comm. Pl. Afr. Austr.* i. 252.—Afr. austr.
parvifolia, *Choisy, in Mém. Soc. Phys. Genèv.* ii. (1824) 95.—Afr. austr.

AGATHIS, Salisb. in Trans. Linn. Soc. viii. (1807) 311. t. 15. *CONIFERAE*, Benth. & Hook. f. iii. 436.
DAMMARA, [Rumph.] Lam. Encyc. ii. 259 (1786).
australis, *Steud. Nom.* ed. II. i. 34.—N. Zel.
Dammara, Rich. Conif. 83. t. 19 = loranthifolia.
loranthifolia, *Salisb. in Trans. Linn. Soc.* viii. (1807) 312. t. 15.—Malaya.

AGATHISANTHEMUM, Klotzsch, in Peters, Reise Mossamb. Bot. 294 (1861) = **Oldenlandia**, Plum. (Rubiac.).
Bojeri, Klotzsch, l. c. = O. Bojeri.
globosum, Klotzsch, l. c. = O. globosa.
Petersii, Klotzsch, l. c. 295 = O. Bojeri.

AGATHISANTHES, Blume, Bijdr. 645 (1825) = **Nyssa**, Linn. (Cornac.).
javanica, Blume, l. c. = N. sessiliflora.

AGATHOMERIS, Delaun. in Bon Jard. (1805) ex DC. Prod. vi. 157 = **Humea**, Sm. (Compos.).

AGATHOPHORA, Bunge (Anabas. Revis. 92); in Mém. Acad. Pétersb. Sér. VII. iv. (1862) n. 11 = **Halogeton**, C. A. Mey. (Chenopod.).
alopecuroides, Bunge, l. c. = H. alopecuroides.

AGATHOPHYLLUM, Juss. Gen. 431 (1789) = **Ravensara**, Sonner. (Laurin.).
acuminatum, Willd. ex Meissn. in DC. Prod. xv. I. 110 = R. acuminata.
aromaticum, Willd. Sp. Pl. ii. 842 = R. aromatica.
cupulare, Blume, Mus. Bot. Lugd. Bat. i. 340 (= *Mespilodaphne cupularis*).
cymosum, Meissn. in DC. Prod. xv. I. 111 (= *Mespilodaphne cymosa*).
heteromorphum, Blume, Mus. Bot. Lugd. Bat. i. 340 (= *Mespilodaphne heteromorpha*).
Lindleyanum, Blume, l. c. (= *Mespilodaphne Lindleyana*.
marginatum, Blume, l. c. (= *Mespilodaphne marginata*).
Neesianum, Blume, l. c. 339 (= *Mespilodaphne mauritiana*).
obtusatum, Blume, l. c. 340 (= *Mespilodaphne heteromorpha*).
Persoonianum, Blume, l. c. 341 (= *Mespilodaphne Persooniana*).
Ravensara, Mirb. ex Steud. Nom. ed. II. 34 = R. aromatica.
retusum, Willd. ex Meissn. in DC. Prod. xv. I. 110 = R. retusa.

AGATHOPHYTON, Moq. in DC. Prod. xiii. II. 84, in syn. (1849) = seq.

AGATHOPHYTUM, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 291. t. 10. c. = **Chenopodium**, Linn.
acuminatum, Schur, Enum. Pl. Transs. 879; Index = C. acuminatum.
Bonus-Henricus, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 291 = C. Bonus-Henricus.
crassifolium, Fuss, Fl. Transs. 553 = C. crassifolium.
glaucum, Fuss, l. c. = C. glaucum.
humifusum, Kitt. Taschenb. ed. II. 264 = C. crassifolium.
rubrum, Reichb. Fl. Sax. 321 = C. rubrum.

AGATHOSMA, Willd. Enum. Hort. Berol. 259 (1809). *RUTACEAE*, Benth. & Hook. f. i. 290.
BUCCO, Wendl. Coll. t. 2. etc. (1808).
DICHOSMA, DC. ex Loud. Hort. Brit. 85 (1830).
GYMNONYCHIUM, Bartl. in Linnaea, xvii. (1843) 354. t. 11.
HARTOGIA, Linn. Syst. ed. X. 939 (1759) (nomen prius).
acerosa, Eckl. & Zeyh. Enum. 114.—Afr. austr.
acuminata, Drège, ex Walp. Ann. i. 155 = punctata.
acuminata, Willd. Enum. Hort. Berol. 260 = imbricata.
acutifolia, *Sond. in Harv. & Sond. Fl. Cap.* i. 416.—Afr. austr.

AGATHOSMA :—
adenocaulis, *Eckl. & Zeyh. Enum.* 115.—Afr. austr.
affinis, *Sond. in Harv. & Sond. Fl. Cap.* i. 405.—Afr. austr.
alaris, *Cham. in Linnaea*, v. (1830) 53.—Afr. austr.
ambigua, *Sond. in Harv. & Sond. Fl. Cap.* i. 426.—Afr. austr.
ambigua, *Sweet, Hort. Brit.* ed. I. 89.—Afr. austr.
anomala, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 408.—Afr. austr.
apiculata, E. Mey. ex Sond. l. c. 405 = apiculata.
apiculata, E. Mey. in Bartl. & Wendl. f. Diosm. 176 = Barosma apiculata.
aristata, *Presl, Bot. Bemerk.* 32.—Afr. austr.
asperifolia, *Eckl. & Zeyh. Enum.* 111.—Afr. austr.
aulonophila, Eckl. & Zeyh. l. c. 115 = commutata, virgata.
barbata, Spreng. Pugill. i. 20 = Macrostylis lanceolata.
barosmaefolia, *Eckl. & Zeyh. Enum.* 111.—Afr. austr.
barosmoides, *Sond. in Harv. & Sond. Fl. Cap.* i. 416.—Afr. austr.
Bartlingiana, Eckl. & Zeyh. Enum. 114 = A. Cerefolium, serpyllacea.
berzeliaefolia, Eckl. & Zeyh. l. c. 115 = virgata.
bifida, Bartl. & Wendl. f. Diosm. 152 = umbellata.
biophylla, Eckl. & Zeyh. Enum. 110 = hirta.
bisulca, *Bartl. & Wendl. f. Diosm.* 129.—Afr. austr.
blaerioides, *Cham. in Linnaea*, v. (1830) 55.—Afr. austr.
blaerioides, Eckl. & Zeyh. Enum. 112 = chortophila.
brevifolia, Loud. Hort. Brit. 86 = erecta.
bruniades, Loud. l. c.; G. Don, Gen. Syst. i. 787 = cuspidata.
capitata, *Sond. in Harv. & Sond. Fl. Cap.* i. 434.—Afr. austr.
cephalotes, E. Mey. ex Sond. l. c. 406 = globuliflora.
Cerefolium, *Bartl. & Wendl. f. Diosm.* 159.—Afr. austr.
chortophila, *Eckl. & Zeyh. Enum.* 116.—Afr. austr.
ciliata, *Link, Enum. Hort. Berol.* i. 238.—Afr. austr.
clavata, Hoffmgg. Verz. Pfl. 36 = obtusa.
collina, *Eckl. & Zeyh. Enum.* 110.—Afr. austr.
commutata, *Sond. in Harv. & Sond. Fl. Cap.* i. 412.—Afr. austr.
concinna, *Hoffmgg. Verz. Pfl.* 36.—Afr. austr.
cordata, Hoffmgg. l. c. = subcordata.
corymbosa, G. Don, Gen. Syst. i. 789 = villosa.
craspedota, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 415.—Afr. austr.
crassifolia, *Sond. l. c.* 438.—Afr. austr.
cuspidata, *Bartl. & Wendl. f. Diosm.* 182.—Afr. austr.
cyminoides, Eckl. & Zeyh. Enum. 117 = chortophila.
decumbens, *Eckl. & Zeyh. l. c.* 116.—Afr. austr.
dioica, Sweet, Hort. Brit. ed. I. 89 = Barosma lanceolata.
Dregeana, *Sond. in Harv. & Sond. Fl. Cap.* i. 410.—Afr. austr.
Eckloniana, *Schlecht. in Linnaea*, v. (1830) 207.—Afr. austr.
elata, *Sond. in Harv. & Sond. Fl. Cap.* i. 438.—Afr. austr.
elegans, *Cham. in Linnaea*, v. (1830) 54.—Afr. austr.
erecta, *Bartl. & Wendl. f. Diosm.* 135.—Afr. austr.
eriantha, Steud. Nom. ed. II. i. 35 = Schlechtendalii.
ericoides, *Schlecht. in Linnaea*, vi. (1831) 206.—Afr. austr.
fastigiata, *Eckl. & Zeyh. Enum.* 117.—Afr. austr.
filipetala, *Eckl. & Zeyh. l. c.* 112.—Afr. austr.
florida, *Sond. in Harv. & Sond. Fl. Cap.* i. 424.—Afr. austr.
florulenta, *Sond. l. c.* 437.—Afr. austr.
foetidissima, Hort. ex Steud. Nom. ed. II. i. 35 = Barosma foetidissima.
foliosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 421.—Afr. austr.
fraudulenta, *Sond. l. c.* 415.—Afr. austr.
geminifolia, *Eckl. & Zeyh. Enum.* 115.—Afr. austr.
Gillivrayi, *Sond. in Harv. & Sond. Fl. Cap.* i. 437.—Afr. austr.
glabra, Eckl. & Zeyh. Enum. 116 = serpyllacea.
glabrata, *Bartl. & Wendl. f. Diosm.* 165.—Afr. austr.
glabrata, E. Mey. ex C. Muell. in Walp. Ann. vii. 513 = crassifolia.
glandulosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 420.—Afr. austr.
glauca, Eckl. & Zeyh. Enum. 112 = bisulca.
globuliflora, *Presl, Bot. Bemerk.* 31.—Afr. austr.
gnidioides, *Schlecht. in Linnaea*, vi. (1831) 206.—Afr. austr.

AGATHOSMA :—
gonaquensis, *Eckl. & Zeyh. Enum.* 110.—Afr. austr.
gracilicaulis, *Sond. in Harv. & Sond. Fl. Cap.* i. 409.—Afr. austr.
graveolens, Mcissn. in Flora, xxvii. (1844) 1. 302 = collina.
hirta, *Bartl. & Wendl. f. Diosm.* 188.—Afr. austr.
hirta, [Ker-Gawl.] in Bot. Reg. t. 369 = villosa.
hirtella, *Sond. in Harv. & Sond. Fl. Cap.* i. 430.—Afr. austr.
hispida, *Bartl. & Wendl. f. Diosm.* 132.—Afr. austr.
Hookeri, *Sond. in Harv. & Sond. Fl. Cap.* i. 406.—Afr. austr.
humilis, *Sond. l. c.*—Afr. austr.
hybrida, Bartl. & Wendl. f. Diosm. 167 = rugosa.
hyponeura, Eckl. & Zeyh. Enum. 114 = serpyllacea.
imbricata, Drège, ex Walp. Ann. i. 155 = polyphylla.
imbricata, *Willd. Enum. Hort. Berol.* 259.—Afr. austr.
involucrata, *Eckl. & Zeyh. Enum.* 109.—Afr. austr.
Joubertiana, *Schlecht. in Linnaea*, vi. (1831) 207.—Afr. austr.
juncea, Hoffmgg. Verz. Pfl. 36 = imbricata.
juniperifolia, *Bartl. in Linnaea*, xvii. (1843) 376.—Afr. austr.
juniperina, *Sond. in Harv. & Sond. Fl. Cap.* i. 433.—Afr. austr.
lanceolata, Spreng. ex Bartl. & Wendl. f. Diosm. 156 = ciliata.
lancifolia, *Eckl. & Zeyh. Enum.* 109.—Afr. austr.
lasiophylla, G. Don, Gen. Syst. i. 788 = villosa.
latifolia, Loud. Hort. Brit. 85 = Barosma serratifolia.
latipetala, *Sond. in Harv. & Sond. Fl. Cap.* i. 422.—Afr. austr.
laxa, Bartl. & Wendl. f. Diosm. 162 = villosa.
lediformis, *Eckl. & Zeyh. Enum.* 109.—Afr. austr.
leptospermoides, Sond. in Harv. & Sond. Fl. Cap. i. 405 = aristata.
linifolia, *Lichtst. ex Roem. & Schult. Syst.* v. 448.—Afr. austr.
lycopodioides, *Bartl. & Wendl. f. Diosm.* 148.—Afr. austr.
marifolia, *Eckl. & Zeyh. Enum.* 113.—Afr. austr.
Martiana, *Sond. in Harv. & Sond. Fl. Cap.* i. 416.—Afr. austr.
mclaleucoides, *Sond. l. c.* 431.—Afr. austr.
Meyeriana, *Presl, Bot. Bemerk.* 32.—Afr. austr.
microphylla, *E. Mey. in Bartl. & Wendl. f. Diosm.* 173.—Afr. austr.
minuta, *Schlecht. in Linnaea*, vi. (1831) 206.—Afr. austr.
mollis, Bartl. & Wendl. f. Diosm. 168 = rugosa.
mollis, Eckl. & Zeyh. ex C. Muell. in Walp. Ann. vii. 514 = florida.
montana, *Schlecht. in Linnaea*, vi. (1831) 207.—Afr. austr.
monticola, *Sond. in Harv. & Sond. Fl. Cap.* i. 409.—Afr. austr.
mucronulata, *Sond. l. c.* 417.—Afr. austr.
Mundtii, *Cham. in Linnaea*, v. (1830) 56.—Afr. austr.
myrsinites, G. Don, Gen. Syst. i. 789 = ciliata.
nigra, *Eckl. & Zeyh. Enum.* 117.—Afr. austr.
nigromontana, *Eckl. & Zeyh. l. c.* 111.—Afr. austr.
Nivcni, *Sond. in Harv. & Sond. Fl. Cap.* i. 413.—Afr. austr.
obtusa, *Bartl. & Wendl. f. Diosm.* 169 = rugosa.
obtusata, G. Don, Gen. Syst. i. 789 = imbricata.
obtusifolia, *Presl, Bot. Bemerk.* 31.—Afr. austr.
orbicularis, *Bartl. & Wendl. f. Diosm.* 175.—Afr. austr.
Owanii, *Harv. & Sond. ex Harv. Thes. Cap.* ii. 42.—Afr. austr.
parviflora, *Bartl. & Wendl. f. Diosm.* 181.—Afr. austr.
patens, Hort. ex Barth. & Wendl. f. Diosm. 159 = A. Cerefolium.
patentissima, Eckl. & Zeyh. Enum. 114 = cuspidata.
patula, Eckl. & Zeyh. l. c. 116 = variabilis.
patula, *G. F. W. Mey. ex Bartl. & Wendl. f. Diosm.* 134.—Afr. austr.
pentachotoma, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 405.—Afr. austr.
perforata, *G. Don, Gen. Syst.* i. 789.—Afr. austr.
pilifera, *Schlecht. in Linnaea*, vi. (1831) 206.—Afr. austr.
planifolia, *Sond. in Harv. & Sond. Fl. Cap.* i. 413.—Afr. austr.
platypetala, *Eckl. & Zeyh. Enum.* 116.—Afr. austr.

AGATHOSMA :—
polyphylla, *Presl, Bot. Bemerk.* 32.—Afr. austr.
prolifera, *Bartl. & Wendl. f. Diosm.* 185.—Afr. austr.
prolifera, Eckl. & Zeyh. Enum. 114 = bisulca.
propinqua, *Sond. in Harv. & Sond. Fl. Cap.* i. 423.—Afr. austr.
pubescens, G. Don, Gen. Syst. i. 788 = villosa.
pubescens, *Sond. in Harv. & Sond. Fl. Cap.* i. 413.—Afr. austr.
pubescens, *Willd. Enum. Hort. Berol.* 259.—Afr. austr.
pubigera, *Sond. in Harv. & Sond. Fl. Cap.* i. 414.—Afr. austr.
pulchella, Link, Enum. Hort. Berol. i. 238 = Barosma pulchella.
punctata, G. Don, Gen. Syst. i. 789 = ovata.
punctata, *Sond. in Harv. & Sond. Fl. Cap.* i. 417.—Afr. austr.
recurvifolia, *Sond. l. c.* 415.—Afr. austr.
reflexa, *Link, Enum. Hort. Berol.* i. 238.—Afr. austr.
robusta, *Eckl. & Zeyh. Enum.* 110.—Afr. austr.
rubra, *Willd. & Lichtst. ex Bartl. & Wendl. f. Diosm.* 178.—Afr. austr.
rufescens, G. Don, Gen. Syst. i. 789 = rubra.
rugosa, Link, Enum. Hort. Berol. i. 238 = pubescens.
sabulosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 406.—Afr. austr.
salina, Eckl. & Zeyh. Enum. 111 = asperifolia.
Schlechtendalii, *Sond. in Harv. & Sond. Fl. Cap.* i. 407.—Afr. austr.
Schlechtendahliana, Eckl. & Zeyh. Enum. 109 = Schlechtendalii.
sedifolia, *Schlecht. in Linnaea*, vi. (1831) 206.—Afr. austr.
serpyllacea, *Lichtst. ex Roem. & Schult. Syst.* v. 447.—Afr. austr.
serruriaefolia, Eckl. & Zeyh. Enum. 109 = bifida.
speciosa, Hoffmgg. Verz. Pfl. Nachtr. i. 126 = laxa.
spicata, *Lichtst. ex G. Don, Gen. Syst.* i. 788.—Afr. austr.
spinosa, *Sond. in Harv. & Sond. Fl. Cap.* i. 417.—Afr. austr.
squamosa, *Bartl. & Wendl. f. Diosm.* 141.—Afr. austr.
stadensis, Eckl. & Zeyh. Enum. 113 = microphylla.
stenopetala, Steud. Nom. ed. II. i. 36 = filipetala.
suaveolens, Eckl. & Zeyh. Enum. 113 = A. Cerefolium.
subcordata, Hoffmgg. Verz. Pfl. 36, 163 = imbricata?
tabularis, *Sond. in Harv. & Sond. Fl. Cap.* i. 403.—Afr. austr.
tenuis, *Sond. l. c.* 423.—Afr. austr.
tenuissima, Otto, ex Bartl. & Wendl. f. Diosm. 137 = erecta.
teretifolia, G. Don, Gen. Syst. i. 787 = Diosma teretifolia.
Thunbergiana, *Bartl. & Wendl. f. Diosm.* 150.—Afr. austr.
Thunbergiana, Schlecht. in Linnaea, vi. (1831) 204 = ambigua.
Thunbergiana, Sond. in Harv. & Sond. Fl. Cap. i. 425 = platypetala.
thymifolia, Hoffmgg. Verz. Pfl. 36 = acuminata.
thymifolia, *Schlecht. in Linnaea*, vi. (1831) 205.—Afr. austr.
thuyoides, Sond. in Harv. & Sond. Fl. Cap. i. 437 = glabrata.
thyoides, G. Don, Gen. Syst. i. 787; Eckl. & Zeyh. Enum. 116 = glabrata.
trachyphylla, Eckl. & Zeyh. l. c. 110 = hispida.
umbellata, Sond. in Harv. & Sond. Fl. Cap. i. 408 = bifida.
variabilis, *Sond. l. c.* 433.—Afr. austr.
Ventenatiana, *Bartl. & Wendl. f. Diosm.* 161.—Afr. austr.
vestita, Willd. ex Roem. & Schult. Syst. v. 447 = imbricata.
villosa, *Willd. Enum. Hort. Berol.* 259.—Afr. austr.
virgata, *Bartl. & Wendl. f. Diosm.* 139.—Afr. austr.
virgata, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 412 = commutata.
virgata, Spreng. ex Bartl. & Wendl. f. Diosm. 145 = imbricata.
Wendlandiana, G. Don, Gen. Syst. i. 788 = erecta.

AGATHOTES, D. Don, in Phil. Mag. viii. (1836) 77 = **Swertia**, Linn. (Gentian.).
alata, D. Don, ex G. Don, Gen. Syst. iv. 177 = S. alata.
Chirayta, D. Don, l. c. 182 = S. Chirata.
nervosa, G. Don, l. c. 177 = S. nervosa.

AGATHYRSUS, D. Don, in Edinb. N. Phil. Journ. (1828–29) 310 = **Lactuca**, Linn. et **Sonchus**, Tourn. (Compos.).
alpinus, D. Don, l. c. = L. alpina.
cacaliaefolius, D. Don, l. c. (= *Mulgedium cacaliaefolius*).
caeruleus, D. Don, l. c. = L. alpina.
cyaneus, D. Don, l. c. = L. hastata.
floridanus, D. Don, l. c. = L. leucophaea.
leucophaeus, Beck, Bot. 170 = L. leucophaea.
macrophyllus, Beck, l. c. = L. macrophylla.
Plumieri, D. Don, in Edinb. N. Phil. Journ. (1828–29) 310 = L. Plumieri.
prenanthoides, D. Don, l. c. = L. prenanthoides.
pulchellus, D. Don, l. c. = L. pulchella.
sibiricus, D. Don, l. c. = L. sibirica.
spicatus, D. Don, l. c.—Hab.?
tataricus, D. Don, l. c. = L. tatarica.

AGATHYRUS, Rafin. New Fl. Am. iv. 84 (1836) = ENOXIS, Rafin. = **Lactuca**, Linn.? (Compos.).

AGATI, Adans. Fam. ii. 326 (1763) = **Sesbania**, Pers. (Legum.).
cannabina, Desv. Journ. Bot. i. (1813) 120 = S. aculeata.
coccinea, Desv. l. c. t. 4 = S. grandiflora.
formosa, F. Muell. Fragm. ii. 88 = S. grandiflora.
grandiflora, Desv. Journ. Bot. i. (1813) 120. t. 4 = S. grandiflora.
tomentosa, Nutt. ex A. Gray, Bot. U. St. Expl. Exped. i. 409 = S. tomentosa.
virgata, Desv. Journ. Bot. i. (1813) 120 = Coursetia virgata.

AGATION, Brongn. in Bull. Soc. Bot. Fr. viii. (1861) 79. *VIOLARIEAE*, Benth. & Hook. f. i. 118.
AGATEA, A. Gray, in Proc. Am. Acad. ii. (1852) 323. (nomen prius).
Pancheri, *Brongn. l. c.* 80.—N. Caled.
Vieillardii, *Brongn. l. c.*—N. Caled.
violare, *Brongn. l. c.*—Ins. Fiji.

AGATOPHYTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 144 = AGATHOPHYTUM, Moq. = **Chenopodium**, Linn.

AGAURIA, Benth. & Hook. f. Gen. ii. 586 (1876). *ERICACEAE*, Benth. & Hook. f. ii. 586.
AGARISTA, D. Don, in G. Don, Gen. Syst. iii. 837 (1834) (quoad spp. Africanae).
buxifolia, *Baker, Fl. Maurit.* 187, *in obs.*—Ins. Borb.
polyphylla, *Baker, in Journ. Linn. Soc.* xx. (1884) 194.—Madag.
salicifolia, *Hook. f. ex Oliver, Fl. Trop. Afr.* iii. 483.—Afr. trop.

AGAVE, Linn. Hort. Ups. 87 (1748). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 738.
ALLIBERTIA, Marion, in Rev. Hortic. Bouch. Rhone (Nov. 1882), ex Baker, in Gard. Chron. (1883) i. 176.
BONAPARTEA, Haw. Syn. Succ. 68 (1812).
LITTAEA, Tagliabue, in Bibl. Ital. i. 106 (1816).
MANFREDA, Salisb. Gen. Pl. Fragm. 78 (1866).
albicans, *Jacobi, in Hamb. Gartenz.* xxi. (1865) 256.—Mexic.
Alibertii, *Baker, in Gard. Chron.* N. S. xix. (1883) i. 176.—Mexic.?
aloides, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 265.—Hab.?
aloina, C. Koch, Wochenschr. iii. (1860) 37 = Sartorii.
altissima, Zumag. Fl. Pedem. ii. 50 = americana.
americana, *Linn. Sp. Pl.* 323.—Am. trop.
amoena, Lem. ex Jacobi, in Hamb. Gartenz. xxi. (1865) 120 = A. Scolymus.
amurensis, Jacobi, l. c. xx. (1864) 548 = xylonacantha.
ananassoides, De Jonghe & Jacobi, Nachtr. i. 32 = Regeliana.
angustifolia, Haw. Syn. Succ. 72 = rigida.
angustifolia, Hort. Par. ex Steud. Nom. ed. II. i. 34 = Furcraea tuberosa.
angustissima, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 306.—Mexic.
antillarum, Descourt. Antill. iv. 239. t. 284 = soboliifera.
applanata, *Lem. ex Jacobi, in Hamb. Gartenz.* xx. (1864) 550.—Mexic.

AGAVE :—
aspera, Jacq. Enum. Vindob. App. 307 = Furcraea cubensis.
asperrima, *Jacobi, in Hamb. Gartenz.* xx. (1864) 561. —Texas.
atrovirens, *Karw. in Salm-Dyck, Hort.* (1834) 302.—Mexic.
attenuata, *Salm-Dyck, l. c.* 303.—Mexic.
auricantha, Hort. ex Baker, in Gard. Chron. (1877) II. 40 = A. Scolymus.
australis, Steud. Nom. ed. I. 18 = Furcraea cubensis.
Banlan, *Perr. in Mém. Soc. Linn. Par.* iii. (1824) 97.—Java.
Beaucarnei, Lem. Illustr. Hortic. xi. (1864) Misc. 65 = Kerchovei.
Beauleriana, Jacobi, in Abh. Schles. Gesell. (1869) 150 = mexicana.
Bernhardi, *Jacobi, Nachtr.* i. 38.—Mexic.
Besseriana, De Smet, ex Jacobi, in Hamb. Gartenz. xxi. (1865) 155 = macroacantha.
Bonnetiana, Peacock, ex Baker, in Gard. Chron. (1877) II. 40 = ferox.
Botterii, *Baker, in Bot. Mag.* t. 6248.—Mexic.
Bouchei, *Jacobi, in Hamb. Gartenz.* xxi. (1865) 217.—Mexic.
brachystachys, *Cav. Descr.* 453.—Mexic.
bracteosa, *S. Wats. ex Engelm. in Gard. Chron.* (1882) I. 776. f. 139.—Mexic.; Calif.
Brauniana, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 216.—Mexic.
bromeliaefolia, *Salm-Dyck, Hort.* (1834) 303.—Mexic.
bulbifera, Bonpl. ex Steud. Nom. ed. II. i. 34 = mexicana.
bulbifera, *Salm-Dyck, Hort.* (1834) 8, 303.—Mexic.
caerulescens, Salm-Dyck, ex Jacobi, Monog. 38 = lophantha.
caespitosa, Tod. Hort. Bot. Panorm. (1877) 32. t. 8 = Sartorii.
californica, Jacobi, Nachtr. 47 = falcata.
Canartiana, Jacobi, in Abh. Schles. Gesell. (1869) 147 = atrovirens.
Candelabrum, Tod. Hort. Bot. Panorm. (1877) 66. t. 15 = rigida.
Cantula, Roxb. Fl. Ind. ii. 167 = vivipara.
caribaea, *Hort. Kew. ex Baker, in Gard. Chron.* (1879) II. 683.—Ind. occ.
Celsiana, Jacobi, in Hamb. Gartenz. xxi. (1865) 446 = Celsii.
Celsii, *Hook. Bot. Mag.* t. 4934.—Mexic.
chiapensis, Hort. Belg. ex Jacobi, in Hamb. Gartenz. xxii. (1866) 213 = polyacantha.
chloracantha, *Salm-Dyck, in Otto, Allg. Gartenz.* (1834) 50.—Mexic.
cinerascens, Jacobi, in Hamb. Gartenz. xx. (1864) 553 = applanata.
coarctata, Jacobi, in Abh. Schles. Gesell. (1869) 147 = atrovirens.
coccinea, *Roezl, ex Jacobi, in Hamb. Gartenz.* xxi. (1865) 61.—Mexic.
cochlearis, *Jacobi, Nachtr.* ii. 67.—Mexic.
Cohniana, Jacobi, in Hamb. Gartenz. xxii. (1866) 321 = yuccaefolia.
Commelini, Salm-Dyck, Hort. (1834) 301 = Furcraea Commelyni.
concinna, *Baker, in Gard. Chron.* (1877) II. 137.—Mexic.
concinna, Hort. Angl. ex Baker, l. c. 717 = albicans.
conduplicata, *Jacobi & Bond. in Hamb. Gartenz.* xxi. (1865) 562.—Mexic.
Consideranti, Carr. in Rev. Hortic. (Nov. 1875) 429. f. 68 = A. Victoriae-reginae.
Corderoyi, Versch. ex Baker, in Gard. Chron. (1877) II. 398.—Mexic.
crenata, Jacobi, in Hamb. Gartenz. xxii. (1866) 176 = A. Scolymus.
cubensis, Jacq. Amer. 100 = Furcraea cubensis.
cucullata, *Lem. ex Jacobi, in Hamb. Gartenz.* xxi. (1865) 124.—Mexico.
cuspidata, Hort. Saunders, ex Baker, in Gard. Chron. (1877) II. 137 = flaccida.
cyanophylla, Jacobi, in Hamb. Gartenz. xxii. (1866) 175 = mexicana.
dasylirioides, *Jacobi & Bouché, l. c.* xxi. (1865) 344.—Guatemal.
dealbata, Lem. ex Jacobi, l. c. 346 = dasylirioides.
Debaryana, Jacobi, in Abh. Schles. Gesell. (1869) 164 = pruinosa.
Decaisneana, *Jacobi, l. c.* 153.—Mexic.
deflexispina, Jacobi, Nachtr. ii. 72 = atrovirens.

AGAVE :—
Demeesteriana, Hort. ex Jacobi, Monog. 218 = miradorensis.
De-Meesteriana, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 167.—Mexic.
densiflora, *Hook. Bot. Mag.* t. 5006.—Mexic.
densiflora, Regel, Gartenfl. (1863) 274. t. 410 = rupicola.
dentata, Hort. ex Baker, in Gard. Chron. (1877) II. 748 = pruinosa.
Deserti, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 310.—Calif.
Desmetiana, Hort. ex Baker, in Gard. Chron. (1877) I. 621 = horrida.
Desmettiana, Jacobi, in Hamb. Gartenz. xxii. (1866) 217 = miradorensis.
drimiaefolia, Hort. Petrop. ex Baker, in Gard. Chron. (1877) II. 808 = undulata.
Ehrenbergiana, Baker, l. c. 620 = Ehrenbergii.
Ehrenbergii, *Jacobi, in Hamb. Gartenz.* xxi. (1865) 255.—Mexic.
echinoides, Jacobi, Nachtr. i. 48 = striata.
Ellemeetiana, *C. Koch, Wochenschr.* viii. (1865) 103.—Mexic.
elongata, Jacobi, in Hamb. Gartenz. xxi. (1865) 168 = rigida.
ensifera, Jacobi, Nachtr. 14 = univittata.
ensiformis, Hort. ex Baker, in Gard. Chron. (1877) II. 556 = striata.
excelsa, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 215.—Honduras.
expansa, Jacobi, Nachtr. ii. 26 = americana.
falcata, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 304.—Mexic.
Fenzliana, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 170. —Mexic.
ferox, *C. Koch, Wochenschr.* iii. (1860) 23.—Mexic.
filamentosa, Salm-Dyck, in Bonplandia, vii. (1859) 94 = filifera.
filifera, *Salm-Dyck, Hort.* (1834) 309.—Mexic.
flaccida, Haw. Syn. Succ. 73 = rubescens.
flaccida, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 174.—Mexic.
flavescens, Salm-Dyck, Hort. Dyck. 8 = macroacantha.
flavovirens, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 270. —Hab.?
foetida, Linn. Sp. Pl. 323 = Furcraea gigantea.
fourcroydes, Lem. Illustr. Hortic. xi. (1864) Misc. 65 = rigida.
fragrantissima, *Jacq. Enum. Vindob. App.* 309.—Hab.?
Fuerstenbergii, Jacobi, Nachtr. ii. 72 = americana.
funifera, Lem. Illustr. Hortic. xi. (1864), Misc. 65, sub t. 419 = Yucca funifera.
Funkiana, C. Koch & Bouché, in C. Koch, Wochenschr. iii. (1860) 47 = lophantha.
Galeottei, *Baker, in Gard. Chron.* (1877) II. 40.—Mexic.
geminiflora, *Ker-Gawl. in Brande, Journ. Sc.* iii. (1817) no. 3. t. 1.—Peruv.
Ghiesbrechtii, *C. Koch, Wochenschr.* v. (1862) 83, 143.—Mexic.
gigantea, D. Dietr. Syn. Pl. ii. 1192 = Furcraea gigantea.
Gilbeyi, Hort. cf. Gard. Chron. (1873) 1305 = horrida.
glabra, *Karw. in M. Roem. Syn. Ensat.* 292.—Mexic.
glaucescens, Hook. Bot. Mag. t. 5333 = attenuata.
glaucescens, *Otto, in M. Roem. Syn. Ensat.* 292.—Mexic.
Goeppertiana, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 219.—Mexic.
gracilis, Jacobi, Nachtr. ii. 66 = atrovirens.
grandidentata, Hort. Belg. ex Jacobi, in Hamb. Gartenz. xxii. (1866) 114 = horrida.
granulosa, *Scheidw. in C. Koch, Wochenschr.* iv. (1861) 286.—Mexic.
Guedeneyri, *Houllet, in Rev. Hortic.* (1875) 466.—Mexic.
Gustaviana, Hort. Corderoy, ex Baker, in Gard. Chron. N. S. viii. (1877) II. 201 = Maximiliana.
guttata, *Jacobi & Bouché, in Hamb. Gartenz*, xxi. (1865) 561.—Mexic.
Haseloffii, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 220.—Mexic.
Haworthiana, M. Roem. Syn. Ensat. 290 = mexicana?
Haynaldi, *Tod. Hort. Bot. Panorm.* i. 88. t. 24.—Mexic.?
heteracantha, Hort. ex Jacobi, in Hamb. Gartenz. xxii. (1866) 62 = lophanta.
heteracantha, *Zucc. in Nov. Act. Nat. Cur.* xvi. (1833) II. 675.—Mexic.

AGAVE :—
hexapetala, *Jacq. Enum. Pl. Carib.* 18.—Ind. occ.
Hookeri, C. Koch, Wochenschr. viii. (1865) 104 = yuccaefolia.
Hookeri, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 168.—Mexic.
Hookeri, Baker, in Bot. Mag. t. 6589 = Fenzliana.
horizontalis, *Jacobi, in Abh. Schles. Gesell.* (1869) 148, bis.—Mexic.
horrida, *Lem. ex Jacobi, in Hamb. Gartenz.* xx. (1864) 546.—Mexic.
Houlletii, Hort. Par. ex Jacobi, l. c. xxii. (1866) 268 = rigida.
Humboldtiana, *Jacobi, l. c.* 264.—Mexic.
humilis, M. Roem. Syn. Ensat. 285 = brachystachys.
hybrida, Versch. Cat. n. 12 (1868–69) = xylonacantha.
Hystrix, Hort. ex Baker, in Gard. Chron. (1877) II. 556 = striata.
inaequidens, *C. Koch, Wochenschr.* iii. (1860) 28.—Mexic.
Inghami, Versch. Cat. n. 16 (1871–72), 11 = Roezliana.
Ixtli, Karw. in Salm-Dyck, Hort. (1834) 304 = rigida.
ixtlioides, Hook. f. Bot. Mag. t. 5893 = rigida.
Jacobiana, Salm-Dyck, in Bonplandia, viii. (1860), 88 = atrovirens.
Jacquiniana, Schult. f. Syst. vii. 727 = lurida.
Karatto, *Mill. Gard. Dict.* ed. VIII. n. 6.—Mexic.
Karwinskii, Zucc. in Nov. Act. Nat. Cur. xvi. II. (1833) 675 = rigida.
Kellocki, Jacobi, in Abh. Schles. Gesell. (1869) 165 = pruinosa.
Keratto, Steud. Nom. ed. II. i. 36 = Karatto.
Keratto, Salm-Dyck, Hort. Dyck. 304 = mexicana.
Keratto, Salm-Dyck, in Bonplandia, vii. (1859) 93 = Salmdyckii.
Kerchovei, *Lem. Illustr. Hortic.* xi. (1864) 64.—Mexic.
kewensis, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 218. —Mexic.
Kochii, Jacobi, l. c. 117 = xylonacantha.
lamprochlora, Jacobi, in Abh. Schles. Gesell. (1869) 150 = Smithiana,
laticincta, *Versch. Cat.* n. 12 (1868) 2.—Mexic.
latifolia, *Karw. in M. Roem. Syn. Ensat.* 292.—Mexic.
latissima, Jacobi, in Hamb. Gartenz. xx. (1864) 551 = atrovirens.
Laurentiana, *Jacobi, l. c.* xxii. (1866) 266.—Mexic.
laxa, *Hort. Monac. ex Salm-Dyck, Hort.* (1834) 8.—Am. austr.
Lecheguilla, Torr. Bot. Mex. Bound. 213 = heteracantha.
Legrelliana, Jacobi, in Hamb. Gartenz. xxii. (1866) 266 = Ehrenbergii.?
Leguayana, Versch. Cat. n. 12 (1868) 2 = Ghiesbrechtii.
Lehmanni, Jacobi, in Abh. Schles. Gesell. (1869) 146 = atrovirens.
Lemairei, *Hort. J. Versch. ex Illustr. Hortic.* xi. (1864) *Misc.* 65.—Hab.?
lepida, D. Dietr. Syn. ii. 1192 = lurida.
Lindleyi, *Jacobi, in Abh. Schles. Gesell.* (1869) 152, bis.—Mexic.
linearis, Jacobi, in C. Koch, & Fint. Wochenschr. xii. (1869) 179 = macroacantha.
lophantha, *Schiede, in Linnaea*, iv. (1829) 582.—Mexic.
lurida, *Ait. Hort. Kew.* i. 472.—Mexic.
macrantha, *Tod. Hort. Bot. Panorm.* ii. 11. t. xxvii. —Mexic.?
macroacantha, *Zucc. in Nov. Act. Acad. Nat. Cur.* xvi. II. (1833) 676.—Mexic.
maculata, *Regel, Ind. Sem. Hort. Petrop.* (1856) 16.—Texas.
maculosa, Hook. Bot. Mag. t. 5122 = maculata.
madagascariensis, Spreng. Syst. ii. 79 = Furcraea madagascariensis.
Magni, Desf. Tabl. ed. II. 33 = mexicana.
Maigretiana, Jacobi, in Hamb. Gartenz. xxii. (1866) 121 = horrida.
Malinezii, C. Koch, Wochenschr. v. (1862) 198 = rudis.
Manguai, Desf. Tabl. Hort. Par. ed. II. 33 = mexicana.
marmorata, *Roezl, in Belg. Hortic.* (1883) 238.—Mexic.
Martiana, *C. Koch, Wochenschr.* iii. (1860) 38.—Mexic.
Maximiliana, *Hort. Saunders, ex Baker, in Gard. Chron.* (1877) II. 201.—Mexic.

AGAVE:—
megalacantha, *Hemsl. Diag. Pl. Nov. Mex.* 55.—Mexic.
melanacantha, *Lem. ex Jacobi, in Hamb. Gartenz.* xxi. (1865) 215.—Mexic.
Mescal, Ellem. ex C. Koch, Wochenschr. viii. (1865) 94 = A. Scolymus.
mexicana, *Lam. Encyc.* i. 52.—Mexic.
micracantha, *Salm-Dyck, in Bonplandia*, vii. (1859) 93.—Mexic.
Milleri, Haw. Syn. Succ. 71 = americana.
miradorensis, *Jacobi, in Abh. Schles. Gesell.* (1869) 156.—Mexic.
mitis, *Mart. Del. Sem. Hort. Monac.* (1848) 4.—Mexic.
mitraeformis, Jacobi, in Abh. Schles. Gesell. (1859) 145 = atrovirens.
Muilmanni, *Jacobi, l. c.* (1870–71) 76.—Mexic.
Newberyi, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 309.—Arizona.
Nissoni, *Baker, in Gard. Chron.* (1877) i. 528.—Mexic.
Noacki, Jacobi, in Hamb. Gartenz. xxi. (1865) 221 = Sartorii.
oblongata, *Jacobi, in Abh. Schles. Gesell.* (1869) 155.—Mexic.
obscura, *Schiede, in Linnaea*, xviii. (1844) 413.—Mexic.?
odorata, Pers. Syn. i. 380 = Furcraea cubensis.
Offoyana, *De Smet, ex Jacobi, in Hamb. Gartenz.* xxi. (1865) 214.—Havana.
oligophylla, *Baker, in Gard. Chron.* (1878) ii. 492.—Mexic.?
ornata, Jacobi, in Hamb. Gartenz. xxi. (1865) 59 = americana.
Ottonis, Jacobi, l. c. xxii. (1866) 320 = atrovirens.
Ousselghemiana, Jacobi, in Abh. Schles. Gesell. (1869) 156 = albicans.
pallida, Salisb. Prod. 247 = virginica.
pallida, *Sartorius, ex Jacobi, in Hamb. Gartenz.* xxi. (1865) 171.—Mexic.
Palmeri, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 319.—Arizona.
Parryi, *Engelm. l. c.* 311.—Arizona, Mexic.
parviflora, *Torr. Bot. Mex. Bound.* 214.—Mexic.
paucifolia, *Tod. Hort. Bot. Panorm.* i. (1877) 77.—Mexic.
Peacockii, *Croucher, in Gard. Chron.* (1873) 1400. t. 283.—Mexic.
pendula, Schnitts. in Zeits. Gartenb. Darmst. (1857) Jun. n. 21 = Sartorii.
perbella, Hort. ex Baker, in Gard. Chron. (1877) 527, 528 = xylonacantha.
perlucida, Jacobi, in Abh. Schles. Gesell. (1869) 151 = Smithiana.
picta, Salm-Dyck, in Bonplandia, vii. (1859) 88 = americana.
polianthoides, *M. Roem. Syn. Ensat.* 286.—Mexic.
polyacantha, *Haw. Rev. Pl. Succ.* 35.—Mexic.
polyanthoides, Cham. & Schlecht. in Linnaea, vi. (1831) 55 = brachystachys.
polyanthoides, *Schiede, ex Schlecht. in Linnaea*, xviii. (1844) 413.—Mexic.
polyphylla, C. Koch, Wochenschr. iii. (1860) 38 = mexicana.
Poselgerii, Salm-Dyck, in Bonplandia, vii. (1859) 92 = heteracantha.
potatorum, *Zucc. in Nov. Act. Nat. Cur.* xvi. (1833) ii. 675.—Mexic.
pruinosa, *Lem. ex Jacobi, in Hamb. Gartenz.* xxi. (1865) 449.—Mexic.
pubescens, *Regel & Ortg. in Gartenfl.* (1874) 227. t. 804.—Mexic.
pugioniformis, *Zucc. in Nov. Act. Nat. Cur.* xvi. (1833) ii. 676.—Mexic.
pulcherrima, Hort. ex C. Koch, Wochenschr. viii. (1865) 104 = Sartorii.
pulcherrima, *Otto, in M. Roem. Syn. Ensat.* 292.—Mexic.
pulchra, *Hort. Monac. in Salm-Dyck, Cat.* 8.—Am. austr.
pulverulenta, *Versch. Cat.* (1863).—Sierra Nevada.
punctata, *Hort. Berol. ex Kunth, Enum. Pl.* ii. 837.—Mexic.
quadrata, *Lem. Illustr. Hortic.* xi. (1864) *Misc.* 65.—Mexic.
ramosa, Moench, Meth. 642 = americana.
recurva, Zucc. in Abh. Akad. Muench. iv. iii. (1845) 22 = striata.

AGAVE:—
Regeliana, Hort. ex Baker, in Gard. Chron. N. S. vii. (1877) i. 621 = horrida.
Regeliana, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 214.—Mexic.
regia, *Hort. Saunders, ex Baker, in Gard. Chron.* N. S. viii. (1877) ii. 620.—Mexic.
revoluta, *Klotzsch, in Otto & Dietr. Gartenz.* viii. (1840) 274.—Mexic.
rigida, *Mill. Gard. Dict.* ed. VIII. n. 8.—Mexic.
rigidissima, Jacobi, in C. Koch & Fint. Wochenschr. xii. (1869) 179 = Kerchovei.
Roezliana, *Baker, in Gard. Chron.* N. S. vii. (1877) i. 528. f. 79.—Mexic.
Rohanii, Hort. Belg. ex Jacobi, in Hamb. Gartenz. xx. (1864) 546 = Ghiesbrechtii.
rubescens, *Salm-Dyck, Hort.* (1834) 306.—Mexic.
rubrocincta, Jacobi, in Abh. Schles. Gesell. (1869) 153 = Sartorii.
rudis, *Lem. ex Jacobi, in Hamb. Gartenz.* xxi. (1865) 216.—Mexic.
rufocincta, Jacobi, Nachtr. i. 88 = Sartorii.
Rumphii, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 121 = vivipara.
rupicola, *Regel, Gartenfl.* vii. (1858) 312.—Mexic.
Salmdyckii, *Baker, in Gard. Chron.* N. S. viii. (1877) ii. 490.—Mexic.
Salmiana, Otto, in Otto & Dietr. Allg. Gartenz. (1842) 51 = atrovirens.
Saponaria, Lindl. Bot. Reg. (1838) Misc. 76 = brachystachys.
Sartorii, *C. Koch, Wochenschr.* iii. (1860) 37.—Mexic.
Saundersii, Hook. f. Bot. Mag. t. 5493 = A. Scolymus.
scabra, Salm-Dyck, in Bonplandia, vii. (1859) 89 = Wislizeni.
schidigera, *Lem. in Illustr. Hortic.* viii. t. 289, verso (1860) ix. t. 330.—Mexic.
Schlechtendahlii, Jacobi, in Hamb. Gartenz. xx. (1864) 555 = atrovirens.
Schnittspahnii, Jacobi, l. c. xxi. (1865) 120 = A. Scolymus.
Schottii, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 305.—Mexic.
Scolymus, *Karw. in Salm-Dyck, Hort.* (1834) 307.—Mexic.
Seemanniana, *Jacobi, in Abh. Schles. Gesell.* (1869) 154.—Guatemal.
serrulata, *Karw. in Otto & Dietr. Allg. Gartenz.* (1842) 51.—Mexic.
sessiliflora, *Hemsl. Diag. Pl. Nov.* 55.—Mexic.
Shawii, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 314. t. 2–4.—Calif.
sisalana, Perrine, ex Engelm. l. c. (reimpr. 28) = rigida.
Smithiana, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 263.—Mexic.
sobolifera, *Salm-Dyck, Hort.* (1834) 307, 309.—Ind. occ.; Mexico.
spectabilis, Salisb. Prod. 247 = americana.
spectabilis, *Tod. Hort. Bot. Pan.* ii. 4. t. 25.—Mexic.?
spicata, *Cav. in Anal. Cienc. Nat.* v. (1802) 261; *Desc.* 454.—Cuba.
spicata, DC. Cat. Hort. Monsp. (1813). 74 = brachystachys.
spicata, Guss. Ind. Sem. (1825) ex Kunth, Enum. Pl. v. 831 = yuccaefolia.
spinosa, Steud. Nom. ed. II. i. 37 = Furcraea tuberosa.
splendens, *Jacobi, in Abh. Schles. Gesell.* (1870–71) 63.—Mexic.
stenophylla, Jacobi, in Hamb. Gartenz. xxii. (1866) 269 = vivipara.
striata, *Zucc. in Nov. Act. Nat. Cur.* xvi. ii. (1833) 678.—Mexic.
stricta, Salm-Dyck, in Bonplandia, vii. (1859) 94 = striata.
subfalcata, Jacobi, in C. Koch & Fint. Wochenschr. xii. (1869) 179 = macroacantha.
subinermis, *M. Roem. Syn. Ensat.* 289.—Bras.
sudburyensis, Hort. Peacock, ex Baker, in Gard. Chron. N. S. viii. (1877) ii. 137 = concinna.
Taylori ×, *Hort. cf. Gard. Chron.* N. S. ii. (1874) 60 et viii. (1877) 621. f. 125.
Taylorea ×, Hort. Veitch. in Rev. Hortic. (1877) 36 = Taylori.
thuacanensis, Karw. ex Otto in Otto & Dietr. Allg. Gartenz. (1842) 51 = atrovirens.

AGAVE:—
teoxomuliana, Karw. ex M. Roem. Syn. Ensat. 292 = bromeliaefolia.
Theometel, *Zuccagn. in Roem. Collect.* 138.—Mexic.
Thomsoniana, *Jacobi, in Hamb. Gartenz.* xxii. (1866) 262.—Mexic.
Toneliana, *Hort. Peacock, ex Baker, in Gard. Chron.* xv. (1881) i. 362.—Mexic.?
triangularis, Jacobi, in C. Koch & Fint. Wochenschr. xii. (1869) 178 = horrida.
tuberosa, Mill. Gard. Dict. ed. VIII. n. 4 = Furcraea tuberosa.
uncinata, Jacobi, in Hamb. Gartenz. xxi. (1865) 165 = polyacantha.
undulata, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 274.—Mexic.
univittata, *Haw. ex Salm-Dyck, Hort.* (1834) 308.—Mexic.
utahensis, *Engelm. in Bot. King, Exp.* 497.—Utah et Arizona.
Vandervinneni, *Lem. Illustr. Hortic.* xi. (1864), *Misc.* 64.—Mexic.
Vanderdonckii, Hort. ex Baker, in Gard. Chron. N. S. vii. (1877) i. 527 = xylonacantha.
variegata, Hort. ex Steud. Nom. ed. I. 18 = americana.
variegata, *Jacobi, in Hamb. Gartenz.* xxi. (1865) 459.—Mexic.
Vera-Crucis, Haw. Syn. Succ. 72 = seq.
Vera-cruz, Mill. Gard. Dict. ed. VIII. n. 7 = lurida.
Verschaffeltii, *Lem. ex Jacobi, in Hamb. Gartenz.* xxi. (1865) 122.—Mexic.
Victoriae-Reginae, *T. Moore, in Gard. Chron.* N. S. iv. (1875) ii. 485.—Mexic.
virginica, Hort. ex Baker, in Gard. Chron. N. S. xix. (1883) i. 176 = Alibertii.
virginica, *Linn. Sp. Pl.* 323.—Am. bor.
virginica, Mill. Gard. Dict. ed. VIII. n. 2 = americana.
viridissima, *Hort. Peacock, ex Baker, in Gard. Chron.* N. S. viii. (1877) ii. 137.—Mexic.
vivipara, Arruda, Diss. Pl. Bras. 23 = Furcraea agavephylla.
vivipara, *Linn. Sp. Pl.* 323.—Mexic.
Wallisii, *Jacobi, Nachtr.* ii. 78.—N. Granat.
Warelliana, *Hort. cf. Gard. Chron.* N. S. v. (1876) i. 603, *et Baker, l. c.* viii. (1877) ii. 264.—Mexic.
wiesenbergensis, *Wittm. in Berl. Gartenz.* (1885) 14.—Mexic.
Wislizeni, *Engelm. in Trans. Acad. Sc. St. Louis*, iii. (1875) 320.—Mexic.
xalapensis, Roezl, ex Jacobi, in Hamb. Gartenz. xxi. (1865) 61 = polyacantha.
xylacantha, Hort. cf. Gard. Chron. N. S. vii. (1877) i. 528 = xylonacantha.
xylonacantha, *Salm-Dyck, in Bonplandia*, vii. (1859) 92.—Mexico.
yuccaefolia, *DC. in Red. Lil.* t. 328, 329.—Mexic.
Yxtli, Karw. ex G. Don, in. Sweet, Hort. Brit. ed. III. 705 = rigida.
Zuccarinii, *Otto, in Allg. Gartenz.* x. (1842) 51, nomen.—Mexic.?

AGDESTIS, Moç. & Sesse, ex DC. Syst. Veg. i. 543 (1818). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 87.
clematidea, *Moç. & Sesse, l. c.*—Mexic.
teterrima, De Not. Ind. Sem. Bot. Genuens. (1855) 29 = clematidea.

AGELAEA, Soland. ex Planch. in Linnaea, xxiii. (1850) 437. *CONNARACEAE*, Benth. & Hook. f. i. 432.
Hemiandrina, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 171. t. 28.
ferruginea, Soland. ex Planch. l. c. 440 = Cnestis ferruginea.
glabrifolia, *Hance, in Journ. Bot.* xiv. (1876) 257.—Cambodia.
Koneri, *O. Hoffm. & Hildebr. ex O. Hoffm. Sert. Madag.* 15.—Madag.
Lamarckii, *Planch. in Linnaea*, xxiii. (1850) 438.—Afr. trop.
nitida, *Soland. ex Planch. l. c.* 437.—Afr. trop.
obliqua, *Baker, in Oliver, Fl. Trop. Afr.* i. 454.—Afr. trop.
pentagyna, Baill. in Bull. Soc. Linn. Par. i. (1882) 345 = Lamarckii.
pruriens, Soland. ex Planch. in Linnaea, xxiii. (1850) 440 = Cnestis corniculata.

AGELAEA :—
Thouarsiana, *Baill. Adansonia*, vii. (1866-67) 240.—Madag.
vestita, *Hook. f. Fl. Brit. Ind.* ii. 46.—Malaya.
villosa, *Soland. ex Planch. in Linnaea*, xxiii. (1850) 438.—Afr. trop.
Wallichii, *Hook. f. Fl. Brit. Ind.* ii. 47.—Malaya.

AGENIUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 447 (1836) = **Andropogon**, Linn. (Gramin.).
nutans, Nees, l. c. = A. Neesii.

AGENORA, D. Don, in Edinb. New Phil. Journ. (Mar. 1829) 310 = **Hypochoeris**, Linn. (Compos.).
laevigata, D. Don, l. c.—H. laevigata.

AGERATINA, Spach, Hist. Vég. Phan. x. 286 (1841) = **Eupatorium**, Linn. (Compos.).
ageratoides, Spach, l. c. = E. ageratoides.
aromatica, Spach, l. c. = E. aromaticum.
cordata, Spach, l. c. = E. aromaticum.

AGERATIOPSIS, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 246 (1873) = **Eupatorium**, Linn. (Compos.).
apiculata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 362 = E. apiculatum.

AGERATIUM, 'Adans.' ex Steud. Nom. ed. II. i. 37 (1841) = AGERATUM, Tourn. = **Erinus**, Linn. (Scrophul.).

AGERATIUM, Reichb. Consp. 209 (1828), err. typ. = ACERATIUM, DC. = **Elaeocarpus**, Linn. (Tiliac.).

AGERATOIDES, Ponted. ex Linn. Class. Pl. 527 (1747) = **Achillea**, Linn. (Compos.).

AGERATUM, Linn. Gen. ed. I. 247 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 241.
CAELESTINA, Cass. Dict. Sc. Nat. vi. Suppl. 8. t. 93 (1817).
CARELIA, Adans. Fam. ii. 123 (1763).
COELESTINA, Cass. in Bull. Soc. Philom. (Jan. 1817) 10.
DECACHAETA, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 462.
OXYLOBUS, Moç. ex DC. Prod. v. 115 (1836).
adscendens, *Sch. Bip. ex Benth. & Hook. f. Gen.* ii. 243.—Mexic.
albidum, *Hemsl. Biol. Centr. Am. Bot.* ii. 81.—Mexic.
album, Steud. Nom. ed. I. 18 = conyzoides.
alternifolium, *Baker, in Mart. Fl. Bras.* vi. II. 195.—Bras.
altissimum, Linn. Sp. Pl. 839 = Eupatorium ageratoides.
angustifolium, Spreng. Syst. iii. 446 = Calea angustifolia.
aquaticum, Roxb. Hort. Beng. 61; et Fl. Ind. iii. 416 = Adenostemma viscosum.
arbutifolium, *H. B. & K. Nov. Gen. et Sp.* iv. 149.—Mexic.
brachystephanum, *Regel, Gartenfl.* iii. (1854) 245. t. 108 f. c.—Caraccas.
caeruleum, Hort. ex Poir. Encyc. Suppl. i. 241 = conyzoides.
campuloclinioides, *Baker, in Mart. Fl. Bras.* vi. II. 196.—Bras.
ciliare, *Linn. Sp. Pl.* 839.—Ind. or. ?
ciliare, *Lour. Fl. Cochinch.* 484.—Cochinch.
coelestinum, Sims, Bot. Mag. t. 1730 = corymbosum.
coeruleum, Desf. Tabl. ed. I. 98 = conyzoides.
coeruleum, Sieber, ex Baker, in Mart. Fl. Bras. vi. II. 345 = Eupatorium macrophyllum.
confertum, *Baker, l. c.* 196.—Bras.
conspicuum, Hort. ex C. Koch & Fint. Wochenschr. i. (1858) 33 = Eupatorium glechonophyllum.
conyzoides, *Linn. Sp. Pl.* 839.—Reg. trop.
conyzoides, Sieber, ex Steud. Nom. ed. II. i. 37 = Eupatorium repandum.
cordifolium, Roxb. Fl. Ind. iii. 415 = conyzoides.
corymbosum, Benth. & Hook. f. Gen. ii. 242 = Agrianthus corymbosus.
corymbosum, *Zucc. ex Pers. Syn.* ii. 402.—Mexic.
domingense, Spreng. Syst. iii. 446.—S. Doming.
echioides, *Hemsl. Biol. Centr. Am. Bot.* ii. 81.—Mexic.

AGERATUM :—
febrifugum, Sesse, ex DC. Prod. v. 104 = Piqueria trinervia.
glanduliferum, *Sch. Bip. ex Benth. & Hook. f. Gen.* ii. 242.—Mexic.
guianense, Aubl. Pl. Gui. ii. 800 = Eupatorium macrophyllum.
heterolepis, *Baker, in Mart. Fl. Bras.* vi. II. 198.—Bras.
hirsutum, Poir. Encyc. Suppl. i. 242 = seq.
hirtum, Lam. Encyc. i. 54 = conyzoides.
Houstonianum, *Mill. Gard. Dict.* ed. VIII. n. 2.—Mexic.
humile, Salisb. Prod. 188 = conyzoides.
isocarphoides, *Hemsl. Biol. Centr. Am. Bot.* ii. 82.—Mexic.
Lasseauxii, Carr. in Rev. Hortic. (1870-71) 90 = Conoclinium Lasseauxii.
latifolium, Cav. Ic. iv. 33 = conyzoides.
latifolium, *Hemsl. Biol. Centr. Am. Bot.* ii. 82.—Guatemal.
lineare, Cav. Ic. iii. 3. t. 205 = Palafoxia linearis.
littorale, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 78.—Am. bor.
longifolium, *Baker, in Mart. Fl. Bras.* vi. II. 197.—Bras.
maritimum, H. B. & K. Nov. Gen. et Sp. iv. 150 = conyzoides.
matricarioides, *Less. Syn. Comp.* 155.—Cuba.
melissaefolium, *DC. Prod.* v. 109.—Bras.
mexicanum, Sims, Bot. Mag. t. 2524 = conyzoides.
mexicanum, Sweet, Brit. Fl. Gard. i. t. 89 = brachystephanum.
microcarpum, *Hemsl. Biol. Centr. Am. Bot.* ii. 82.—Costa-Rica.
microcephalum, *Hemsl. l. c.*—Mexic.
micropappum, *Baker, in Mart. Fl. Bras.* vi. II. 198.—Bras.
microphyllum, *Sch. Bip. in Seem. Bot. Voy. Herald*, 298.—Mexic.
muticum, *Griseb. Fl. Brit. W. Ind.* 356.—Jamaic.
nanum, Hort. ex Sch. Bip. in C. Koch & Fint. Wochenschr. i. (1858), Garten-Nachr. 26 = suffruticosum.
obtusifolium, Lam. Encyc. i. 54 = conyzoides.
odoratum, Vilm. Fl. Pl. Terre, ed. II. 42 = conyzoides.
paleaceum, *Hemsl. Biol. Centr. Am. Bot.* ii. 83.—Mexic.
paniculatum, Hort. ex Steud. Nom. ed. II. i. 37 = Eupatorium paniculatum.
pedatum, Ortega, Hort. Matr. 38 = Florestina pedata.
petiolatum, *Hemsl. Biol. Centr. Am. Bot.* ii. 83.—Nicarag.
Pohlianum, *Baker, in Mart. Fl. Bras.* vi. II. 197.—Bras.
punctatum, *Jacq. Hort. Schoenb.* iii. 28. t. 300.—Mexic.
punctatum, Ortega, Hort. Matr. 37 = Stevia Eupatoria.
purpureum, *Vitm. Summa Pl.* iv. 474.—Guian.
purpureum, Moç. & Sesse, ex DC. Prod. v. 122 = Stevia viscida.
quadriflorum, Blanco, Fl. Filip. ed. I. 624 = Elephantopus spicatus.
rubens, *Viv. Elench. Pl. Hort. Dinegro*, 9.—Hab. ?
salicifolium, *Hemsl. Biol. Centr. Am. Bot.* ii. 83.—Mexic.
scabriusculum, *Hemsl. l. c.*—Mexic.
scorpioideum, *Baker, in Mart. Fl. Bras.* vi. II. 197.—Am. trop.
sessilifolium, *Schau. in Linnaea*, xix. (1847) 715.—Mexic.
strictum, *Hemsl. Biol. Centr. Am. Bot.* ii. 83.—Mexic.
strictum, Sims, Bot. Mag. t. 2410 = Adenostemma viscosum.
suffruticosum, *Regel, Gartenfl.* iii. (1854) 389. t. 108.—Mexic.
tomentosum, *Hemsl. Biol. Centr. Am. Bot.* ii. 84.—Mexic.
viscosum, Ortega, Hort. Matr. 36 = Stevia lucida, salicifolia.
Wendlandii, *Hort. ex Vilm. Fl. Pl. Terre, Suppl.* 2.—Mexic.
Wrightii, *Torr. & Gray, in Proc. Am. Acad.* i. (1848) 46.—Texas.

AGERATUM, Tourn. ex Adans. Fam. ii. 210 (1763) = **Erinus**, Linn. (Scrophul.).

AGERELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 129 = **Veronica**, Linn. (Scrophul.).
acinifolia, Fourr. l. c. = V. acinifolia.
arvensis, Fourr. l. c. = V. arvensis.
verna, Fourr. l. c. = V. verna.

AGERIA, Adans. Fam. ii. 166 (1763) = **Ilex**, Linn.
acuminata, Rafin. Sylva Tellur. 46 = I. Macoucoua.
Cassena, Rafin. l. c. 47 = I. Cassine.
geminata, Rafin. l. c. 48 = I. Cassine.
heterophylla, Rafin. l. c. = I. Dahoon.
mucronata, Rafin. l. c. 47.—Hab. ?
obovata, Rafin. l. c. 48 = I. Dahoon.
opaca, Rafin. l. c. 47 = I. opaca.
ovalis, Rafin. l. c. 48 = I. ovalifolia ?
palustris, Rafin. l. c. 47 = I. Dahoon.
retusa, Rafin. l. c. 46 = I. retusa.
uniflora, Rafin. l. c. 47 = I. uniflora.

AGGERANTHUS, Wight, Ic. v. 18. t. 1737 (1852) = **Eria**, Lindl. (Orchid.).
marchantioides, Wight, l. c. = E. reticulata.

AGIALID, Adans. Fam. ii. 508 (1763) = **Balanites**. Delile (Simarub.).

AGIHALID, Alp. ex Juss. in Dict. Sc. Nat. i. 297 (1804) = praec.

AGINA, Neck. Elem. ii. 153 (1790) = **Sagina**, Linn. (Caryophyll.).

AGIRTA, Baill. Etud. Gén. Euph. 463 (1858) = **Tragia**, Linn. (Euphorb.).
Boiviniana, Baill. l. c. = T. Boiviniana.

AGLAEA, Steud. Nom. ed. I. 19 (1821) = **Melasphaerula**, Ker-Gawl. (Irid.).
graminea, Steud. l. c. = M. graminea.

AGLAIA, Allem. in Nov. Act. Nat. Cur. iv. (1770) 93 = **Cyperus**, Tourn. ?

AGLAIA, Lour. Fl. Cochinch. 173 (1790). *MELIACEAE*, Benth. & Hook. f. i. 334, 994.
AMERINA, Noronha, in Verh. Batav. Gen. v. (1790) Art. IV. 1.
ARGOPHILUM, Blanco, Fl. Filip. 186 (1837).
EUPHORA, Griff. Notul. iv. 547 (1854).
MILNEA, Roxb. Fl. Ind. ed. Wall. ii. 430 (1824).
NEMEDRA, Juss. in Mém. Mus. Par. xix. 223 (1830).
NYALELIA, Dennst. Schluess. Hort. Malab. 30 (1818).
SELBYA, M. Roem. Syn. Hesper. 89, 126 (1846).
acuminatissima, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* xvii. (1864) 42.—Malaya.
ancolana, Miq. Fl. Ind. Bat. Suppl. i. 506 = argentea.
andamanica, *Hiern, in Hook. Fl. Brit. Ind.* i. 555.—Ins. Andaman.
angustifolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 55.—Sumatra.
apiocarpa, *Hiern, in Hook. Fl. Brit. Ind.* i. 555.—Zeylan.
argentea, *Blume, Bijdr.* 170.—Java.
aspera, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* xxvii. (1864) 42.—Java.
basiphylla, *A. Gray, Bot. U. St. Expl. Exped.* i. 237.—Ins. Fiji.
batjanica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 46.—Malaya.
chittagonga, Miq. l. c. 44 = Amoora chittagonga.
cordata, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 557.—Malacca.
coriacea, *Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 57.—Borneo.
crassinervia, *Kurz, ex Hiern, in Hook. f. Fl. Brit. Ind.* i. 556.—Tenasserim.
Cumingiana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 409 = Hearnia Cumingiana.
decandra, Wall. Cat. n. 1276 = Amoora decandra.
denticulata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 410.—Luzon.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* i. 507.—Sumatra.
Dookkoo, Griff. Notul. iv. 505 (= *Lansium domesticum* ?).
edulis, *A. Gray, Bot. U. St. Expl. Exped.* i. 237.—Ind. or.; Fiji.
edulis, Hiern, in Hook. Fl. Brit. Ind. iii. 556 = undulata, minutiflora.

AGLAIA :—
edulis, Seem. Fl. Vit. 37 = basiphylla.
elacagnoidea, *Benth. Fl. Austral.* i. 383.—Austral.
elliptica, Blume, Bijdr. 171 = Hearnia elliptica.
eximia, Miq. Fl. Ind. Bat. Suppl. i. 506 = argentea.
Forstenii, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 46.—Amboina.
Ganggo, *Miq. Fl. Ind. Bat. Suppl.* i. 506.—Sumatra.
glabrata, Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. xxvii. (1864) 42 = Dysoxylum Halmaheirae.
glabriflora, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 555.—Malacca.
grandis, *Korth. in Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 56.—Borneo.
grata, Wall. ex Voigt, Hort. Suburb. Calc. 136 = Roxburghiana.
Griffithii, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 146.—Malacca.
Halmaheirae, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 58 = Dysoxylum Halmaheirae.
hexandra, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 410.—Luzon.
hypoleuca, *Miq. Fl. Ind. Bat. Suppl.* i. 507.—Sumatra.
inaequalis, Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. xxvii. (1864) 305 = Hearnia elliptica.
khasiana, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 554.—Ind. or.
Korthalsii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 42.—Sumatra.
latifolia, *Miq. l. c.*—Java.
laxiflora, *Miq. l. c.* 52.—Borneo.
lepidota, Miq. Fl. Ind. Bat. Suppl. i. 507 = Roxburghiana.
leptantha, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 51.—Sumatra.
littoralis, *Miq. l. c.* 45.—Nov. Guin.
Llanosiana, *C. DC. in DC. Monog. Phan.* i. 621.—Ins. Philipp.
longifolia, *Teijsm. & Binn. in Tijdschr. Ned. Ind.* xxvii. (1864) 42.—Java.
macrobotrys, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 409.—Luzon.
macrophylla, Teijsm. & Binn. in Tijdschr. Ned. Ind. xxvii. (1864) 42 = Dysoxylum Halmaheirae.
midnaporensis, Carey, ex Voigt, Hort. Subnrb. Calc. 136 = Roxburghiana.
minutiflora, *Bedd. Pl. Ind. Or.* i. 44.—Ind. or.
mucronulata, *C. DC. in DC. Monog. Phan.* i. 601.—Java.
multijuga, *Seem. Fl. Vit.* 37.—Ins. Fiji.
odorata, *Lour. Cochinch.* 173.—China.
odoratissima, Benth. in Hook. Lond. Journ. Bot. ii. (1845) 213 = elacagnoidea.
odoratissima, Blume, Bijdr. 171 = Diepenhorstii.
oligantha, *C. DC. in DC. Monog. Phan.* i. 603.—Ins. Philipp.
oligocarpa, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 45.—Sumatra.
oligophylla, *Miq. Fl. Ind. Bat. Suppl.* i. 507.—Sumatra.
ovata, Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. xxvii. (1864) 43 = Hearnia elliptica.
pachyphylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 57.—Sumatra.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* i. 507.—Borneo.
Pamattonis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 53.—Borneo.
paniculata, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 199.—Ind. or.
pedicellaris, *C. DC. in DC. Monog. Phan.* i. 607.—Ind. or.
pentaphylla, Kurz, in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 48 = odorata.
perviridis, *Hiern, in Hook. Fl. Brit. Ind.* i. 556.—Ind. or.
pirifera, *Hance, in Journ. Bot.* xv. (1877) 331.—China.
polyantha, Bedd. Pl. Ind. Or. i. 44 = minutiflora.
polyphylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 56.—Java.
polystachya, Wall. in Roxb. Fl. Ind. ii. 429 = Amoora Rohituka.
pycnocarpa, *Miq. Fl. Ind. Bat. Suppl.* i. 196.—Sumatra.
pyramidata, *Hance, in Journ. Bot.* xv. (1877) 331.—China.
pyrrholepis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 47.—Java.
Reinwardtii, *Miq. l. c.* 51.—Celebes.
Roxburghiana, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 147 = undulata.

AGLAIA :—
Roxburghiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 41.—As. or.
rufa, *Miq. l. c.* 49.—Sumatra.
samoënsis, *A. Gray, Bot. U. St. Expl. Exped.* i. 236.—Ins. Navigat.
sexipetala, Griff. Notul. iv. 505 = edulis.
sipamias, Miq. Fl. Ind. Bat. Suppl. i. 506 = palembanica.
Spanoghei, Blume, ex Miq. l. c. 197 = Roxburghiana.
speciosa, *Blume, Bijdr.* 171.—Java.
speciosa, Teijsm. & Binn. Cat. Hort. Bogor. 211 = argentea.
spectabilis, Hort. Calc. ex C. DC. in DC. Monog. Phan. i. 585 = Amoora spectabilis.
subgrisea, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 54.—Malaya.
submonophylla, *Miq. l. c.* 40.—Borneo.
Sulingi, *Blume, Bijdr.* 170.—Java.
Sulingi, Hassk. Cat. Hort. Bogor. 220 = Diepenhorstii.
tenuicaulis, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 556.—Malaya.
Teysmanniana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 48.—Sumatra.
tomentosa, Teijsm. & Binn. in Tijdschr. Ned. Ind. xxvii. (1864) 43 = palembanica.
trichostemon, *C. DC. in DC. Monog. Phan.* i. 608.—Borneo.
Turezaninowii, *C. DC. l. c.* 623.—Ins. Philipp.
undulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 44.—Ind. or.
Wallichii, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 555.—Ind. or.
Zippelii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 54.—Nov. Guin.

AGLAIOPSIS, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 58 (1868–69) = **Hearnia**, F. Muell. (Meliac.).
lancifolia, Miq. l. c. 59 = H. lancifolia.
glaucescens, Miq. l. c. 58 = H. glaucescens.

AGLAODENDRON, Remy, in Ann. Sc. Nat. Sér. III. xii. (1879) 175 = **Plazia**, Ruiz & Pav. (Compos.).
cheiranthifolium, Remy, l. c. = P. cheiranthifolia.

AGLAODORUM, Schott, Gen. Aroid. t. 58 (1858) = **Aglaonema**, Schott (Aroid.).
Griffithii, Schott, l. c. = Aglaonema palustre.

AGLAONEMA, Schott, Melet. i. 20 (1829). *AROIDEAE*, Benth. & Hook. f. iii. 981.
AGLAODORUM, Schott, Gen. Aroid. t. 58 (1858).
acutispathum, *N. E. Br. in Gard. Chron.* N. S. xxiv. (1885) II. 39.—China.
commutatum, Schott, Syn. Aroid. 123 = Scindapsus Cuscuaria.
Cuscuaria, Miq. Fl. Ind. Bat. iii. 217 = Scindapsus Cuscuaria.
fallax, Schott, ex Engl. in DC. Monog. ii. 439 = simplex.
gracile, Schott, in Ann. Mus. Lugd. Bat. i. 279 = pictum.
Haenkii, *Schott, in Bonplandia*, vii. (1859) 30.—Ins. Philipp.
Hookerianum, *Schott, l. c.*—Ind. or.
integrifolium, *Schott, Melet.* i. 20.—Malaya.
longicuspidatum, Schott, Prod. 304 = Schottianum.
maculatum, *Blume, Rumphia*, i. 154.—Ins. Philipp.
malaccense, Schott, in Bonplandia, vii. (1859) 30 = Schottianum.
marantifolium, *Blume, Rumphia*, i. 153. t. 66.—Malaya.
Mannii, *Hook. f. Bot. Mag.* t. 5760.—Afr. trop.
modestum, *Schott, ex Engl. in DC. Monog. Phan.* ii. 442.—Malaya.
nitidum, Kunth, Enum. Pl. iii. 56 = integrifolium.
oblongifolium, Kunth, l. c. 55 = marantifolium.
palustre, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* xxv. (1863) 399.—Malaya.
pictum, *Kunth, Enum. Pl.* iii. 55.—Ind. or.; Malaya.
princeps, Kunth, l. c. = simplex.
propinquum, Schott, in Ann. Mus. Lugd. Bat. i. 280 = Schottianum.
pygmaeum, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 161 = Chamaecladon pygmaeum.
Schottianum, *Miq. Fl. Ind. Bat.* iii. 216.—Malaya.
simplex, *Blume, in Rumphia*, i. 152. t. 65.—Java.
subundulatum, *Zoll. Syst. Verz. Ind. Archip.* 76.—Celebes.

AGLITHEIS, Rafin. Fl. Tellur. ii. 17 (1836) = **Allium**, Linn. (Liliac.).
carinata, Rafin. l. c. 18 = All. carinatum.
tatarica, Rafin. l. c. = All. tataricum.
ursina, Rafin. l. c. = All. ursinum.

AGLOTOMA, Rafin. Fl. Tellur. ii. 44 (1836) = **Aster**, Linn. (Compos.).
multiflora, Rafin. l. c. = Aster multiflorus.

AGLYCIA, Willd. ex Steud. Nom. ed. II. i. 37 (1840) = **Eriochloa**, Kunth (Gramin.).
coarctata, Willd. ex Steud. l. c. = E. polystachya.
distachya, Willd. ex Steud. l. c. = E. distachya.

AGNANTHUS, Vaill. ex Linn. Gen. ed. I. 366 (1737) = **Cornutia**, Linn. (Verben.).

AGNISTUS, G. Don, in Sweet, Hort. Brit. ed. III. 503 (1839) err. typ. = **Acnistus**, Schott (Solanac.).

AGNOSTUS, A. Cunn. in Loud. Hort. Brit. Suppl. i. 580 (1832) = **Stenocarpus**, R. Br. (Proteac.).
sinuatus, A. Cunn. l. c. = S. sinuatus.

AGNUS-CASTUS, [Tourn.] Carr. in Rev. Hortic. xlii. (1870–71) 415 = **Vitex**, Linn. (Verben.).
incisa, Carr. l. c. = V. Negundo.
Negundo, Carr. l. c. = V. Negundo.
robusta, Carr. l. c. = seq.?
vulgaris, Carr. l. c. 414 = V. Agnus-castus.

AGONANDRA, Miers, in Ann. Nat. Hist. Ser. II. viii. (1851) 172, nomen. *OLACINEAE*, Benth. & Hook f. i. 349.
brasiliensis, *Benth. & Hook. f. Gen.* i. 349.—Bras.
excelsa, *Griseb. in Goett. Abh.* xxiv. (1879) 149.—Am. austr.

AGONIS, Lindl. Swan River App. 10 (1839). *MYRTACEAE*, Benth. & Hook. f. i. 703.
BILLOTTIA, R. Br. in Journ. Soc. Geogr. i. (1832) 19.
angustifolia, Schau. in Lehm. Pl. Preiss. i. 118 = linearifolia.
conspicua, Schau. l. c. = linearifolia.
flexuosa, *Lindl. Swan River App.* 10.—Austral.
floribunda, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 20.—Austral.
glabra, *Turcz. in Bull. Acad. Pétersb.* x. (1852) 334.—Austral.
grandiflora, *Benth. Fl. Austral.* iii. 100.—Austral.
hypericifolia, *Schau. in Lehm. Pl. Preiss.* i. 117.—Austral.
juniperina, *Schau. l. c.* 118.—Austral.
linearifolia, *Lindl. Swan River App.* 10.—Austral.
marginata, *Schau. in Lehm. Pl. Preiss.* i. 117.—Austral.
obtusissima, *F. Muell. Fragm.* xi. 119.—Austral.
parviceps, *Schau. in Lehm. Pl. Preiss.* i. 119.—Austral.
Scortechiniana, *F. Muell. Fragm.* xi. 118.—Austral.
spathulata, *Schau. in Lehm. Pl. Preiss.* i. 119.—Austral.
theaeformis, Schau. l. c. ii. 223 = hypericifolia.
undulata, *Benth. Fl. Austral.* iii. 100.—Austral.

AGONOLOBUS, Reichb. Nom. 183 (1841) = **Cheiranthus**, Linn. (Crucif.).

AGONOMYRTUS, Schau. ex Reichb. Handb. 253 (1837) = **Leptospermum**, Forst.? (Myrtac.).

AGOPHYLLUM, Neck. Elem. ii. 227 (1790) = **Zygophyllum**, Linn.

AGORRHINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 126 = **Antirrhinum**, Tourn. (Scrophul.).
Oruntium, Fourr. l. c. = Ant. Orontium.

AGOSERIS, Rafin. Fl. Ludov. 58 (1817) = **Troximon**, Nutt. (Compos.).
cuspidata, Steud. Nom. ed. II. i. 37 = T. marginatum.
glaucus, Steud. l. c. = T. glaucum.
parviflora, D. Dietr. Syn. Pl. iv. 1832 = T. parviflorum.
rosea, D. Dietr. l. c. = T. roseum.
taraxacifolia, D. Dietr. l. c. = T. taraxicifolium.

AGOSTANA, Bute, ex S. F. Gray, Nat. Arr. Brit. Pl. ii. 526 (1821) = **Bupleurum**, Tourn. (Umbell.).
divaricata, S. F. Gray, l. c. 527 = B. Odontites.
tenuissima, Bute, ex S. F. Gray, l. c. 526 = B. tenuissimum.

AGRAPHIS, Link, Handb. i. 166 (1829) = **Scilla**, Linn. (Liliac.).
campanulata, Link, l. c. = S. hispanica.
cernua, Reichb. Fl. Germ. Excurs. 105 = S. cernua.
nutans, Link, Handb. i. 166 = S. nutans.
patula, Reichb. Fl. Germ. Excurs. 106 = S. hispanica.
serotina, Heynh. Nom. ii. 14 = Dipcadi serotinum.

AGRAULUS, Beauv. Agrost. 5. t. 4 (1812) = **Agrostis**, Linn. (Gramin.).
alpinus, Beauv. l. c. = Agrostis rupestris.
anemagrostoides, Trin. in Spreng. N. Entdeck. ii. 56 = Agrostis effusa.
caninus, Beauv. Agrost. 5. t. 4. f. 7 = Agrostis canina.
flavescens, Panz. ex Steud. Nom. ed. I. 19, 340 = Festuca varia.
mexicanus, Presl, Reliq. Haenk. i. 236 = Agrostis tolucensis.
montanus, Panz. ex Steud. Nom. ed. I. 19 = Agrostis alpina.
pallidus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 149 = Agrostis pallida.
setaceus, S. F. Gray, l. c. = Agrostis setacea.

AGRETTA, Eckl. Verz. 23 (1827) = **Tritonia**, Ker-Gawl. (Irid.).
crispa, Eckl. l. c. 24 = T. undulata.
grandiflora, Eckl. 23, nomen.—Afr. austr.
pallideflavens, Eckl. l. c. 24.—Afr. austr.
pentandra, Eckl. l. c. 23 = T. scillaris.
retusa, Steud. Nom. ed. II. i. 37.—Afr. austr.
stricta, Eckl. Verz. 23, nomen.—Afr. austr.

AGRIANTHUS, Mart. ex DC. Prod. v. 125 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 244.
campestris, *Mart. l. c.*—Bras.
corymbosus, *DC. l. c.* vii. 266.—Bras.
empetrifolius, *Mart. l. c.* v. 126.—Bras.

AGRICOLAEA, Schrank, in Denkschr. Akad. Muench. (1808) 98 = **Clerodendron**, Linn. (Verben.).
fragrans, Schrank, l. c. = C. fragrans.

AGRIMONIA, Tourn. ex Linn. Syst. ed. I. (1735). *ROSACEAE*, Benth. & Hook. f. i. 622.
AMONIA, Nestl. Potent. 17 (1816).
AREMONIA, Neck. Elem. ii. 100 (1790).
SPALLANZANIA, Pollini, Hort. Veron. Pl. Nov. 10 (1816).
acutifolia, Dum. Fl. Belg. Prod. 97 = odorata.
adhaerens, Gilib. Fl. Lituan. ii. 185 = A. Eupatorium.
adscendens, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 311, nomen.—Rossia.
agrimonoides, *Linn. Sp. Pl.* 448.—Europ.
americana, Luca, ex Wallr. Beitr. Bot. i. 43 = A. Eupatoria.
Blumei, *G. Don, Gen. Syst.* ii. 563.—Java; Japon.
bracteata, E. Mey. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 349 = A. Eupatoria.
bracteosa, E. Mey. Zwei Pfl. Docum. 51, 142, nomen = A. Eupatoria.
caffra, E. Mey. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 349 = A. Eupatoria.
canescens, Dum. Fl. Belg. Prod. 97 = odorata.
ceylanica, Wight, Ic. t. 224 = zeylanica.
conopsea, Tschern. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 346 = pilosa.
dahurica, Willd. ex Stev. ex DC. Prod. ii. 587 = pilosa.
davurica, Schlecht. pat. ex Ledeb. Fl. Ross. ii. 32 = praec.
decumbens, Linn. f. Suppl. 251 = Acaena latebrosa.
elata, Moench, Meth. 674 = repens.
elata, Salisb. Prod. 361 = A. Eupatoria.
Eupatoria, Michx. Fl. Am. Bor. i. 287 = parviflora.
Eupatoria, *Linn. Sp. Pl.* 448.—Reg. bor. temp.
Eupatorium, Thw. Enum. Pl. Zeyl. 102 = zeylanica.
eupatoroides, Gueldenst. ex Ledeb. Fl. Ross. ii. 31 = A. Eupatoria.
glabrata, Spreng. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 346 = A. Eupatoria.
Goctectiana, Andrz. ex C. A. Mey. l. c. = A. Eupatoria.

AGRIMONIA :—
graeca, *Boiss. & Heldr. ex Boiss. Fl. Orient.* ii. 728 = A. Eupatoria.
grandis, Andrz. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 343 = A. Eupatoria.
gryposepala, Wallr. Beitr. Bot. i. 49 = A. Eupatoria.
hirsuta, *Bong. ex C. A. Mey. in Bull. Sc. Acad. Pétersb.* x. (1842) 347.—Bras.
humilis, *Wallr. Beitr. Bot.* i. 37.—Dalmat.
incisa, *Torr. & Gray, Fl. N. Am.* i. 431.—Am. bor.
inermis, Pers. Syn. ii. 9 = agrimonoides.
javanica, Jungh. Java, i. 664, ex Miq. Fl. Ind. Bat. i. 1. 370 = Blumei.
lanata, Wall. Cat. n. 709 = A. Eupatoria.
leucantha, *Kunze, Del. Sem. Hort. Lips.* (1842).—Hab.?
major, Uspensky, in Bull. Soc. Nat. Mosc. vii. (1834) 373 = odorata?
microcarpa, Wallr. Beitr. Bot. i. 39 = A. Eupatoria.
minor, Mill. Gard. Dict. ed. VIII. n. 2 = A. Eupatoria.
nepalensis, D. Don, Prod. Fl. Nep. 229 = A. Eupatoria.
octandra, *Luce, Topogr. Nachr. Ins. Oesel*, 150.—Ins. Ossil.
odorata, *Mill. Gard. Dict.* ed. VIII. n. 3.—Europ.
officinalis, Lam. Encyc. i. 62 = A. Eupatoria.
officinarum, Lam. Fl. Fr. iii. 477 = A. Eupatoria.
parviflora, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 130. —Am. austr.
parviflora, Spreng. Syst. ii. 461 = A. Eupatoria.
pilosa, Ledeb. Ind. Hort. Dorp. Suppl. (1823) 1 = A. Eupatoria.
platycarpa, Wallr. Beitr. Bot. i. 38 = A. Eupatoria.
procera, Wallr. Beitr. Fl. Hercyn. 203; in Linnaea, xiv. (1840) 573 = odorata.
pubescens, Wallr. Beitr. Bot. i. 45 = A. Eupatoria.
pumila, Muhl. ex Rafin. Fl. Ludov. 134, nomen. —Am. bor.
repens, Eckl. & Zey. Enum. 264 = A. Eupatoria.
repens, *Linn. Syst.* ed. X. 1046.—Hispan.; As. Min.
repens, Mutel, Fl. Fr. i. 342 = odorata.
robusta, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 311 = odorata?
rostellata, Wallr. Beitr. Bot. i. 42 = A. Eupatoria.
serrifolia, Wallr. l. c. 40 = A. parviflora.
sessiliflora, Dum. Fl. Belg. Prod. 97 = odorata.
sororia, *Fisch. & Mey. ex C. A. Mey. in Bull. Sc. Acad. Pétersb.* x. (1842) 344.—Oriens.
stipularis, Dum. Fl. Belg. Prod. 97 = odorata.
striata, Michx. Fl. Bor. Am. i. 887 = A. Eupatoria.
suaveolens, Hort. ex C. A. Mey. in Bull. Sc. Acad. Pétersb. x. (1842) 344 = odorata.
suaveolens, Blume, Bijdr. 1113 = Blumei.
suaveolens, Pursh, Fl. Am. Sept. i. 336 = parviflora.
sulcata, Dulac, Fl. Hautes-Pyr. 303 = A. Eupatoria.
viscidula, Bunge, in Mém. Sav. étr. St. Pétersb. ii. 100; (Enum. Pl. Chin. Bor. 26) = A. Eupatoria.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 577 = A. Eupatoria.
zeylanica, Moon, Cat. 37 = Alchemella indica.

AGRIODENDRON, Endl. Gen. 144 (1837) = **Aloë**, Linn. (Liliac.).

AGRIOPHYLLUM, Bieb. Fl. Taur. Cauc. iii. 6 (1819). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 58.
arenarium, *Bieb. l. c.*—As. bor.
arenarium, *Bunge, in Mém. Sav. étr. Pétersb.* vii. (1851) 456 (*Lehm. Rel.* 280).—As. temp.
gobicum, *Bunge, in Bull. Acad. St. Pétersb.* xxv. (1879) 355.—Sibir.
lateriflorum, *Moq. in DC. Prod.* xiii. II. 139.—Armen.
latifolium, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1839) 170.—Oriens.
minus, *Fisch. & Mey. l. c.*—Turkest.
mongolicum, Hort. Aven. ex Moq. in DC. Prod. xiii. II. 139 = arenarium.
pungens, Link, ex A. Dietr. Sp. Pl. i. 124 = Corispermum pungens.
squarrosum, Moq. in DC. Prod. xiii. II. 139 = arenarium.
Tournefortii, Fisch. ex Moq. l. c. = lateriflorum.

AGRIPHYLLUM, Juss. Gen. 190 (1789) = **Berkheya**, Ehrh. (Compos.).
asteroides, Desf. Tabl. Hort. Par. ed. I. 109 = B. fruticosa.

AGRIPHYLLUM :—
ciliare, Desf. Cat. Hort. Par. ed. III. 186 = Collumia ciliaris.
echinatum, Desf. Tabl. Hort. Par. ed. I. 109 = B. cernua.
grandiflorum, Desf. Cat. Hort. Par. ed. II. 186 = B. grandiflora.
incanum, Desf. l. c. = B. fruticosa.
spinosum, Desf. ex Steud. Nom. ed. I. 19, 106 = B. fruticosa.

AGROCHARIS, Hochst. in Flora, xxvii. (1844) I. 19 = **Caucalis**, Linn. (Umbell.).
gracilis, Hook. f. in Journ. Linn. Soc. vi. (1860) 9 = C. melanantha.
melanantha, Hochst. in Flora, xxvii. (1844) I. 19 = C. melanantha.

AGROPHYLLUM, Neck. Elem. ii. 227 (1790) = **Zygophyllum**, Linn.

AGROPYRON, J. Gaert. in Nov. Comm. Petrop. xiv. (1770) I. 539. *GRAMINEAE* [Benth. & Hook. f. iii. 1202].
ANTHOSACHNE, Steud. Syn. Pl. Gram. 237 (1855).
BRACONOTIA, Godr. l. c. ed. I. iii. 191 (1844).
COSTIA, Willk. in Bot. Zeit. (1858) 377.
CREMOPYRUM, Schur, Enum. Pl. Transs. 807 (1866).
CRITHOPYRUM, Hort. Prag. ex Steud. Syn. Pl. Gram. 344 (1855).
ELYTRIGIA, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190.
EREMOPYRUM, Jaub. & Spach, Illustr. Pl. Or. iv. 26 (1850–53).
HAYNALDIA, Schur, Enum. Pl. Transs. 807 (1866).
HETERANTHELIUM, Hochst. ex Jaub. & Spach, Illustr. Pl. Or. iv. 24 (1850).
ROEGNERIA, C. Koch, in Linnaea, xxi. (1848) 413.
SECALIDIUM, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 91.
acutum, C. Koch, in Linnaea, xxi. (1848) 424 = repens.
acutum, *Roem. & Schult. Syst.* ii. 751.—Europ.
acutum, Reichb. ex Nym. Consp. 841 = pungens.
affine, Reichb. Ic. Fl. Germ. xi. 7 = acutum.
alpinum, Schur, Enum. Pl. Transs. 810.—Europ.
altissimum, Schur, l. c. 809 = repens.
angustifolium, Schult. Mant. ii. 412.—Sibir.
apricum, *Opiz, in Lotos*, iii. (1853) 63.—Europ.
aristatum, Bess. Enum. Pl. Volh. 41 = cristatum.
attenuatum, Roem. & Schult. Syst. ii. 751.—Ecuador.
Aucheri, *Boiss. Diagn.* Ser. I. v. 75.—Oriens.
barbatulum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 21 = seq.
barbulatum, *Schur, Enum. Pl. Transs.* 809.—Europ.
bicorne, Roem. & Schult. Syst. ii. 760 = Triticum bicorne.
biflorum, *Roem. & Schult. l. c.*—Syria.
biforme, Fig. & De Not. in Mem. Torin. Ser. II. xii. (1852) 260 = patulum.
Bourgaei, *Boiss. Fl. Orient.* v. 669.—Armen.
brachyphyllum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 663.—Persia.
brevissimum, Beauv. Agrost. 102 (sphalm. pro laevissimo) = Triticum polonicum.
bromiforme, Schur, Enum. Pl. Transs. 808 = repens.
bulbosum, *Boiss. Diagn.* Ser. I. v. 75.—Persia.
caesium, J. & C. Presl, Delic. Prag. 213 = repens.
caespitosum, *C. Koch, in Linnaea*, xxi. (1848) 424.—Armen.
Caldesii, *Goran. in Nuov. Giorn. Bot.* xii. (1880) 145. —Ital.
campestre, *Godr. & Gren. Fl. Fr.* iii. 607.—Europ.
caninum, *Beauv. Agrost.* 102.—Europ.
capillare, Beauv. l. c. = repens.
caudatum, Beauv. l. c. = villosum.
collinum, Opiz, Natural. x. (1825) 248; Seznam, 12 = repens.
concinnum, De Not. ex Parl. Fl. Ital. i. 498 = pungens.
condensatum, *Presl, Rel. Haenk.* i. 266.—Chili.
crinitum, *Link, Hort. Berol.* i. 32.—Hab.?
cristatum, *J. Gaert. in Nov. Comm. Petrop.* xiv. (1770) I. 540.—Iberia; Tauria; Sibir.
cristatum, *Beauv. Agrost.* 102; *Roem. & Schult. Syst.* ii. 758.—Hab.?
curvifolium, Lange, in Kjoeb. Vidensk. Meddel. (1860) 55 = scirpeum.
dasyanthum, *Ledeb. in Ind. Sem. Dorpat.* (1820) 3.—Rossia.

AGROPYRON :—
dasystachyum, *Scrib. in Torrey Bot. Club.* x. (1883) 78. —Am. bor.
densiflorum, *Beauv. Agrost.* 102.—Sibir.
desertorum, Schult. Mant. ii. 412 = sibiricum ?
distans, C. Koch, in Linnaea, xxi. (1848) 426 = Bourgaei.
distichum, *Beauv. Agrost.* 102.—Afr. austr.
divaricatum, *Boiss. & Bal. in Bull. Soc. Bot. Fr.* iv. (1857) 307.—Oriens.
divergens, *Nees, ex Steud. Syn. Pl. Gram.* 347.—Am. bor.
dumetorum, *Trautv. in Act. Hort. Petrop.* ix. (1884) 322.—Europ.
elatum, Opiz, Seznam, 12 = cristatum.
elongatum, Host, ex Beauv. Agrost. 102 = rigidum.
elymogenes, Arndt, ex Nym. Consp. 840 = strictum.
farctum, Boiss. Fl. Orient. v. 665 in syn. = junceum
filiforme, *Poir. ex Roem. & Schult. Syst.* ii. 760.—Ins. Borbon.
firmum, J. & C. Presl, Fl. Cech. 28 = repens.
geminatum, Schult. Mant. iii 655 = Eleusine indica.
giganteum, Roem. & Schult. Syst. ii. 753 = rigidum.
glaucum, Blanco, ex Nym. Consp. 841 = repens.
glaucum, *Roem. & Schult. Syst.* ii. 752.—Europ.
Halleri, Reichb. Fl. Germ. Excurs. 20 = Festuca Poa.
hirsutum, Hort. Berol. ex Ledeb. Fl. Alt. i. 114 = cristatum.
hispanicum, Presl, Cyp. et Gram. Sic. 49 = Festuca Poa.
hordeaceum, Boiss. Diagn. Ser. I. xiii. 67 = orientale.
imbricatum, *Roem. & Schult. Syst.* ii. 757.—Rossia.
intermedium, Beauv. Agrost. 102 = glaucum.
intermedium, Reichb. Fl. Germ. Excurs. 140 = glaucum, junceum, et obtusiflorum.
junceum, *Beauv. Agrost.* 102.—Europ.
Kotschyanum, Boiss. & Hohenh. Diagn. Ser. I. xiii. 69 = patulum.
laevissimum, Beauv. Agrost. 146 = Triticum polonicum.
lasianthum, Boiss. Diagn. Ser. I. xiii. 68 = orientale.
laxum, 'Fr.' ex Willk. & Lange, Prod. i. 109 = acutum.
lazicum, *Boiss. Fl. Orient.* v. 661.—As. Min.
ligusticum, *De Not. in Atti Ottav. Riun. Sc. Ital.* 602.—Ital.
ligusticum, Savign. in Flora, xxx. (1847) 569 = Triticum ligusticum.
littorale, *Dum. Obs. Gram. Belg.* 97.—Europ.
loliiforme, Schur, Enum. Pl. Transs. 808 = repens.
longearistatum, *Boiss. Fl. Orient.* v. 660.—Oriens.
longiglume, *Hack. ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 11.—Europ.
maritimum, Beauv. Agrost. 102 = Cutandia maritima.
mucronatum, Opiz, Seznam, 12 = glaucum.
multiflorum, Beauv. Agrost. 102 = repens.
muricatum, Eichw. Skizze, 119 = cristatum.
muricatum, *Schult. Mant.* ii. 414.—Sibir.
nubigenum, *Nees, ex Steud. Syn. Pl. Gram.* 342.—Java.
obtusiflorum, *Roem. & Schult. Syst.* ii. 753.—Batav.
obtusiusculum, *Lange, Handb.* ed. II. 48.—Dan.; Scan.
orientale, *Roem. & Schult. Syst.* ii. 757.—Oriens; Afr. bor.
panormitanum, *Parl. ex Boiss. Voy.* 680; *Pl. Rar. Sic. Fasc.* ii. 20.—Europ.; Oriens.
patulum, *Trin. Fund.* 152.—Aegypt.
pauciflorum, *Schur, in Verh. Siebenb. Ver. Naturw.* x. (1859) 77.—Europ.
pectinatum, *Beauv. Agrost.* 102.—Austral.; N. Zel.
pectiniforme, *Roem. & Schult. Syst.* ii. 758.—Tauria.
peruvianum, *Roem. & Schult. l. c.* 761.—Peruv.
pilosum, *Presl, Reliq. Haenk.* i. 267.—Mexic.
pilosum, Schur, Enum. Pl. Transs. 809 = barbulatum.
Pouzolzii, Godr. & Gren. Fl. Fr. iii. 608 = glaucum.
prostratum, Beauv. Agrost. 102 = triticeum.
pseudo-caninum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 91 = caninum.
pumilum, Beauv. Agrost. 102 = triticeum.
pungens, Reichb. ex Nym. Consp. 840 = acutum.
pungens, *Roem. & Schult. Syst.* ii. 753.—Europ.
pycnanthum, Gren. & Godr. Fl. Fr. iii. 606 = littorale.
Raegnerii, *Boiss. Fl. Orient.* v. 662.—Reg. Caucas.
repens, *Beauv. Agrost.* 102.—Reg. temp.
Richardsoni, *Schrad. in Linnaea*, xii. (1838) 467 in syn.—Am. bor.
rigidum, *Beauv. Agrost.* 102.—Europ.
Rothii, *Opiz, in Lotos*, iii. (1853) 64.—Europ.
Rouxii, *Gren. & Duval, in Mém. Soc. Emul. Doubs*, III. iv. (1860) 391.—Europ.
rupestre, *Schult. Mant.* ii. 409.—Hab.?
Savignonii, *De Not. Prosp. Fl. Ligust.* 57.—Liguria.

AGROPYRON :—
scabrum, *Beauv. Agrost.* 102.—Austral.
scirpeum, *Presl, Gram. Sic.* 49.—Sicilia.
Scribneri, *Vasey, in Bull. Torrey, Bot. Club*, x. (1883) 128.—Am. bor.
secundum, *Presl, Rel. Haenk.* i. 266.—Chili.
semicostatum, *Nees, ex Steud. Syn. Pl. Gram.* 346.—Affghan.
sepium, Beauv. Agrost. 102 = repens.
sibiricum, *Beauv. l. c.*—Europ. or.; As. occ.
squarrosum, Link, Hort. Berol. i. 32 = patulum.
stipaefolium, Czern. Consp. Fl. Charcov. 70, nomen.—Rossia.
striatum, Nees, ex Steud. Syn. Pl. Gram. 346 = Triticum striatum.
strictum, *Reichb. in Moessl. Handb.* ed. II. iii. 1812.—Europ.
strigosum, *Boiss. Fl. Orient.* v. 661.—Oriens.
subulatum, *Roem. & Schult. Syst.* ii. 761.—Aleppo.
Tauri, *Boiss. & Bal. in Bull. Soc. Bot. Fr.* iv. (1857) 307.—As. Min.
tenerum, *Vasey, in Coult. Bot. Gaz.* x. (1885) 258.—Am. bor.
tenuiculum, Steud. Nom. ed. II. i. 38 = Festuca Poa.
Tournefortii, *De Not. in Atti Ottav. Riun. Sc. Ital.* 602.—Oriens.
Tournefortii, Savign. in Flora, xxx. (1847) 569 = Aucheri.
triticeum, *J. Gaertn. in Nov. Comm. Petrop.* xiv. (1770) i. 539.—Europ.; As. bor.
truncatum, *Fuss, Fl. Transs.* 749.—Europ.
unilaterale, Beauv. Agrost. 102 = Festuca tenuiflora.
vaginans, *Beauv. Agrost.* 102.—Am. merid.
Vaillantianum, *Trautv. in Act. Hort. Petrop.* ix. (1884) 323.—Europ.
variegatum, *Roem. & Schult. Syst.* ii. 759.—Hab.?
velutinum, *Nees, in Hook. Lond. Journ. Bot.* ii. (1843) 417.—Austral.
villosum, *Link, Hort. Berol.* i. 31.—Europ.; Oriens.
violaceum, *Vasey, Gram. Un. St.* 45.—Am. bor.

AGROPYRUM, Roem. & Schult. Syst. ii. 750 (1817) = praec.

AGROSINAPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 329 = **Brassica**, Tourn. (Crucif.).
arvensis, Fourr. l. c. = B. arvensis.

AGROSTANA, Hill, Veg. Syst. v. 32 (1772) = **Bupleurum**, Linn. (Umbell.).

AGROSTEMA, Linn. Syst. ed. I. (1735) = seq.

AGROSTEMMA, Linn. Gen. ed. I. 135 (1737) = **Lychnis**, Linn. (Caryophyll.).
apetala, G. Don, Gen. Syst. i. 416 = L. apetala.
alpina, Forbes, Hort. Woburn. 104 = L. alpina.
Banksia, Meerb. Fl. Sel. 28 = L. coronaria.
brachypetala, Sweet, Hort. Brit. ed. I. 44 = L. apetala.
Bungeana, D. Don, in Sweet, Fl. Gard. Ser. II. 317 = L. Bungeana.
Coeli-rosea, Linn. Sp. Pl. 436 = L. Coeli-rosa.
Coronaria, Linn. l. c. = L. Coronaria.
corsica, G. Don, Gen. Syst. i. 416 = L. corsica.
declinis, G. Don, l. c. = L. diclinis.
decumbens, Denson, ex Loud. Hort. Brit. Suppl. i. 580 = L. decumbens.
dioica, Dufour, ex G. Don. Gen. Syst. i. 416 = L. diclinis.
fimbriata, G. Don, l. c. 417 = L. indica.
Flos-cuculi, G. Don, l. c. = L. Flos-cuculi.
Flos-jovis, Linn. Sp. Pl. 436 = L. Flos-jovis.
Flos-jovis, Pollich, Hist. Pl. Palat. i. 443 = L. coronaria.
fulgens, Doell, Rhein. Fl. 643 = L. fulgens.
Githago, Linn. Sp. Pl. ed. II. 435 = L. Githago.
Gitthago, Retz. Fl. Scand. 86 = praec.
gracilis, Boiss. Diagn. Ser. II. i. 80.—As. Min.
grandiflora, Doell, Rhein. Fl. 643 = L. coronaria.
hirsuta, Gilib. Fl. Lituan. ii. 173 = L. Githago.
hirsuta, Stokes, Bot. Mat. Med. ii. 559 = L. Githago.
inflata, G. Don, Gen. Syst. i. 417 = L. inflata.
involucrata, G. Don, l. c. 416 = L. apetala.
laeta, G. Don, l. c. = Silene laeta.
lusitanica, G. Don, l. c. 417 = L. Flos-cuculi.
nicaeensis, Pers. Syn. i. 519 = L. Githago.
nivalis, G. Don, Gen. Syst. i. 416 = L. nivalis.
parviflora, Cav. ex Steud. Nom. ed. I. 19 = Silene laeta.
perennans, Haw. ex Loud. Hort. Brit. Suppl. i. 580 = L. Densoni.
pulchra, G. Don, Gen. Syst. i. 417 = Silene laciniata.

AGROSTEMMA :—
pusilla, Desf. ex Steud. Nom. ed. II. i. 38 = L. Coeli-rosa.
pyrenaica, G. Don, Gen. Syst. i. 417 = L. pyrenaica.
saxatilis, G. Don, in Sweet, Hort. Brit. ed. III. 67 = L. saxatilis.
sibirica, G. Don, Gen. Syst. i. 416 = L. sibirica.
suecica, Maund, ex Steud. Nom. ed. II. i. 38 = L. alpina.
sylvestris, G. Don, Gen. Syst. i. 416 = L. dioica.
triflora, G. Don, l. c. 417 = L. triflora.
variegata, G. Don, l. c. 416 = L. variegata.

AGROSTICULA, Raddi, Agrost. Bras. 33. t. 1. f. 2 (1823) = **Sporobolus**, R. Br. (Gram.).
muralis, Raddi, l. c. = S. minutiflorus.

AGROSTIS, Linn. Syst. ed. I. (1735); Gen. ed. I. 19 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1149.
AGRAULUS, Beauv. Agrost. 5. t. 4 (1812).
AVENA, Hall. ex Scop. Introd. 74 (1777).
DECANDOLIA, Bast. Fl. Maine-et-Loire, 28 (1809).
NOSONEMA, Rafin. Neogenyt. 4 (1825).
PEREILEMA, Presl, Reliq. Haenk. i. 233. t. 37 (1830).
TRICHODIUM, Michx. Fl. Bor. Am. i. 41. t. 8 (1803).
VILFA, Adans. Fam. ii. 495 (1763).
abakanensis, Less. ex Trin. in Mém. Acad. Pétersb. Sér. VI. vi. 11. (1845) 325 = Michauxii.
aberrans, *Steud. Syn. Pl. Gram.* 422.—Reg. Magell.
actinoclada, F. Muell. Fragm. vi. 84 = Sporobolus actinoclados.
aculeata, Scop. Fl. Carn. ed. II. i. 62 = Crypsis aculeata.
acutiflora, Steud. Nom. ed. I. 19 = Deyeuxia acutiflora.
adscendens, Langc, in Kjoeb. Vidensk. Meddel (1860) 33 = verticillata.
aemula, R. Br. Prod. 172 = Deyeuxia Forsteri.
aenea, Spreng. Syst. Cur. Post. 337 = Sporobolus aeneus.
aenea, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1845) 332.—Sitcha.
aequata, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 413 = Deyeuxia aequata.
aequivalvis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1845) 362.—Am. bor. occ.
affinis, Kunth, Enum. Pl. i. 228 = canina.
affinis, Schult. Mant. ii. 195 = Sporobolus Muchlenbergii.
africana, Poir. Encyc. Suppl. i. 254 = Sporobolus capensis.
airaeformis, *Steud. Syn. Pl. Gram.* 172.—Chili.
airoides, *Torr. ex Spreng. Syst. Cur. Post.* 31 = Sporobolus airoides.
alba, Kunze, in Flora, xxix. (1846) 678 = Reuterii.
alba, *Linn. Sp. Pl.* 63.—Europ.; As.
albicans, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863) 91 = exarata.
albicans, Presl, ex Steud. Nom. ed. I. 19.—Hab.?
albida, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. 11. (1845) 344.—Sarepta; Rossia.
algida, [Soland. in] Phipps, Voy. 200 = Phippsia algida.
alopecuroides, Lam. Illustr. n. 812 = Polypogon monspeliensis.
alopecuroides, Link, ex Kunth, Enum. Pl. i. 228 = effusa.
alopecuroides, Vahl, ex Kunth, Enum. Pl. i. 86 = Panicum Myurus.
alpestris, Laest. ex Nym. Consp. 802 = rubra.
alpicola, *Hochst. in Flora*, xxxviii. (1855) 284.—Afr. or.
alpina, Leyss. Fl. Halen. 16 = canina.
alpina, Willd. Sp. Pl. i. 368 = rupestris.
alpina, Savi [Fl. Pisana, i. 81?], ex Steud. Nom. ed. I. 19 = vulgaris.
alpina, *Scop. Fl. Carn.* ed. II. i. 60.—Europ.
alpina, Stokes, in With. Bot. Arr. Brit. Pl. ed. II. i. 71 = sctacea.
altissima, *Tuckerm. in Am. Journ. Sc.* xlv. (1843) 44 = elata.
ambigua, Roem. & Schult. Syst. ii. 354 = alba.
anatolica, C. Koch, in Linnaea, xxi. (1848) 379 = verticillata.
andina, *Phil. in Anales Univ. Chil.* (1873) 561.—Chili.
Anemagrostis, Syme, Eng. Bot. ed. III. xi. 43 = Apera Spica-venti, et interrupta.
anemagrostoides, Trin. in Spreng. Neue Entdeck. ii. 56 = effusa.
anomala, Willd. Sp. Pl. i. 370 = elata.
antarctica, *Hook. f. Fl. Antarct.* 374. t. 132.—Reg. Antarct.
Antoniana, *Grisch. in Goett. Abh.* xxiv. (1879) 293.—Reg. Argent.
aperta, Bose, ex Trin. in Mém. Acad. Petersb. Ser. VI. vi. 11. (1845) 287 = Muehlenbergia diffusa.

AGROSTIS :—
aphanes, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 346.—N. Zembl.
aquatica, *Buckl. in Proc. Ac. Sc. Philad.* '1862.' (1863) 90.—Am. bor.
aquatica, Pourr. in Mém. Acad. Toul. iii. (1788) 306 = verticillata.
arachnoidea, Poir. Encyc. Suppl. i. 249 = Cinna arundinacea.
arachnoides, *Ell. Sketch*, i. 134.—Am. bor.
arenaria, Gouan, Illustr. 3 = Sporobolus pungens.
arenaria, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 51 = canina.
arenosa, Schur, Enum. Pl. Transs. 736 = canina.
argentea, Lam. Fl. Fr. iii. 570 = Stipa Calamagrostis.
aristata, Sincl. ex Steud. Nom. ed. II. i. 39 = alba.
aristulata, *C. Muell. in Bot. Zeit.* xiv. (1856) 349.—Ins. Falkland.
articulata, Brot. Fl. Lusit. i. 73 = Chaeturus fasciculatus.
arundinacea, Linn. Sp. Pl. 61 = Deyeuxia sylvatica.
arundinacea, *Presl, in Rel. Haenk.* i. 238.—Peruv.
arundinacea, Vill. Hist. Pl. Dauph. ii. 79 = Calamagrostis varia.
arvensis, *Phil. in Linnaea*, xxix. (1857-58) 87.—Chili.
aspera, Michx. Fl. Am. Bor. i. 53 = Sporobolus asper.
aspera, Willd. ex Steud. Nom. ed. II. i. 39 = Muehlenbergia stipoides.
asperifolia, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. II. (1845) 317 = exarata.
asperigluma, *Steud. Syn. Pl. Gram.* 422.—Chili.
asperula, *Phil. in Linnaea*, xxix. (1857-58) 87.—Chili.
atropurpurea, Hort. Hafn. ex Steud. Nom. ed. II. i. 39 = vulgaris.
atrovirens, Roem. & Schult. Syst. ii. 361 = Sporobolus atrovirens.
aucklandica, Hook. f. Fl. Antarct. i. 96 = Deschampsia caespitosa.
australis, *Lam. Encyc.* i. 57.—Europ.
australis, Linn. Mant. i. 30 = Gastridium australe.
avenacea, J. F. Gmel. Syst. i. 171 = Deyeuxia Forsteri.
avenoides, *Hook. f. Handb. New Zeal. Fl.* 330.—N. Zel.
barbata, Buckl. in Proc. Acad. Sc. Philad. '1862.' (1863) 334 = Sporobolus ramulosus.
barbata, Banks & Soland. ex Hook. f. Fl. Nov. Zel. i. 293 = Dichelachne crinita.
barbata, Pers. Syn. i. 75 = Sporobolus virginicus.
Bergiana, *Trin. Gram. Unifl.* 203.—Afr. austr.
berica, Hort. Madr. ex Steud. Nom. ed. II. i. 39 (= *Urachne parviflora*).
Berlandieri, '*Fourn. Mex. Pl. En. Gram.* 96,' *ex Hemsl. Biol. Centr. Am. Bot.* iii. 550.—Mexic.
bermudiana, Tuss. ex Kunth, Enum. Pl. i. 259 = Cynodon Dactylon.
Berteriana, Spreng. ex Steud. Nom. ed. II. 39, 143 = Arundinella martinicensis.
Biebersteiniana, Claus, Beitr. Russ. Reich. viii. 264 = trichoclada.
Billardieri, R. Br. Prod. 171 = Deyeuxia Billardieri.
biseta, Spreng. ex Steud. Nom. ed. II. i. 39, 143 = Arundinella indica.
Boeckeleri, Seubert, ex Steud. Syn. Pl. Gram. 218 (= *Gymnopogon distichophyllus*).
borbonica, *Steud. l. c.* 170.—Ins. Borbon.
borealis, Hartm. Skand. Fl. ed. IV. 23 = canina.
Bourgaei, '*Fourn. Mex. Pl. En. Gram.* 95,' *ex Hemsl. Biol. Centr. Am. Bot.* iii. 550.—Mexic.
brachyathera, *Steud. Syn. Pl. Gram.* 422.—Reg. Magell.
brachyphylla, Steud. l. c. 175 = Polypogon brachyphyllus.
brachyphylla, Willd. ex Steud. Nom. ed. II. i. 39 = Sporobolus atrovirens.
brachystachys, *Schult. Mant.* iii. 569.—Sicilia.
brasiliensis, Spreng. Nov. Prov. 45 = Reimaria acuta.
brevifolia, *Nutt. Gen. Am.* i. 44.—Am. bor.
breviglumis, *F. Muell. Census*, 133.—Austral.
brevis, Knapp, Gram. Brit. 116 = alba?
bromidioides, *Griseb. in Goett. Abh.* xxiv. (1879) 293.—Reg. Argent.
bromoides, Linn. Mant. i. 30 = Stipa Aristella.
bryoides, *Dum. Fl. Belg.* 152.—Belgium.
byzantina, *Boiss. Diagn.* Ser. I. xiii. 46.—Byzantium.
caerulescens, Poir. Encyc. Suppl. i. 258 (= *Urachne caerulescens*).
caerulescens, vide etiam coerulescens, DC.
caespitosa, Gaudich. in Freyc. Voy. Bot. 407 (= *Vilfa Gaudichaudii*).

AGROSTIS :—
caespitosa, Hoppe, ex Steud. Nom. ed. II. i. 39 = rupestris.
caespitosa, Presl, Rel. Haenk. i. 237 = Preslii.
caespitosa, Torr. in Ann. Lyc. N. York, i. (1824) 152 = Torreyi.
caespitosa, Willd. ex Steud. Nom. ed. II. i. 39 = Muehlenbergia quadridentata.
caespitosa, Salisb. Prod. 25 = Deschampsia caespitosa.
Calamagrostis, Georgi, Besch. Russ. Reich. iii. IV. 687 = Calamagrostis lanceolata.
Calamagrostis, Linn. Syst. ed. X. 872 = Stipa Calamagrostis.
Calamagrostis, Suter, Fl. Helv. i. 36 = Calamagrostis epigeios.
calamagrostoides, Regel, Ind. Sem. Hort. Petrop. (1865) 38 = Calamagrostis agrostoides.
californica, Trin. Agrost. ii. 113 = exarata?
calycina, Willd. ex Steud. Nom. ed. II. i. 39; ii. 530 = Schismus marginatus.
campestris, *Phil. in Linnaea*, xxix. (1857-58) 87.—Chili.
campyla, *Tuckerm. in Am. Journ. Sc.* Ser. II. vi. (1848) 231.—Am. bor.
canariensis, *Parl. in Webb & Berth. Phyt. Canar.* iii. 391.—Ins. Canar.
Candollei, Parl. Fl. Palerm. i. 68 = alba.
canescens, *Griseb. in Goett. Abh.* xix. (1874) 253.—Reg. Argent.
canescens, Salisb. Prod. 25 = Corynephorus canescens.
canina, *Linn. Sp. Pl.* 62.—Amphig.
canina, Ucria, Hort. Reg. Panom. 57 = pallida.
capensis, *Lam. Encyc.* i. 58.—Afr. austr.
capensis, Steud. in Flora, xii. (1829) 467 = Bergiana.
capensis, Willd. Sp. Pl. i. 372 = Sporobolus capensis.
capillaris, All. Fl. Pedem. ii. 233 = canina.
capillaris, Huds. Fl. Angl. ed. II. 27 = alba.
capillaris, Boiss. & Lange, ex Anderss. in Walp. Ann. vi. 981 = truncatula.
capillaris, Desf. Fl. Atlant. i. 69 = Reuterii.
capillaris, Bieb. Fl. Taur. Cauc. i. 55 = trichoclada.
capillaris, *Linn. Sp. Pl.* 62; *Huds. Fl. Angl.* 27.—Europ.
capillaris, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. i. 55 = trichoclada.
capillaris, Pourr. ex Nym. Consp. 801 = olivetorum.
capillaris, Thore, Chlor. 26 = elegans.
capillaris, Vill. Hist. Pl. Dauph. ii. 73 = vulgaris.
cenisia, De Not. ex Parl. Fl. Ital. i. 189 = alpina.
Carmichaeli, *Roem. & Schult. Mant.* iii. 571.—Ins. Tristan d'Acugn.
caryophyllea, Salisb. Prod. 25 = Aira caryophyllea.
castellana, *Boiss. & Reut. Diagn. Hisp.* 26.—Hispan.; Afr. bor.
chamaecalamus, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 365.—Peruv.
Chamissonis, *Trin. l. c.* 374.—Ins. Paschalis.
chilensis, *Kunze, ex Trin. l. c.* 339.—Chili.
Chinantlae, '*Fourn. Mex. Pl. En. Gram.* 96,' *ex Hemsl. Biol. Centr. Am. Bot.* iii. 550.—Mexic.
chinensis, Koen. ex Steud. Nom. ed. II. i. 39 = Leptochloa tenerrima.
chonotica, *Phil. in Linnaea*, xxix. (1857-58) 87.—Chili.
ciliaris, Steud. Syn. Pl. Gram. 116 = Arundinella anomala.
ciliata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 373.—Ind. or.
ciliata, Pall. ex Ledeb. Fl. Ross. iv. 404 = Pappophorum boreale.
ciliata, Thunb. Fl. Jap. 49 = Festuca Thunbergii.
ciliata, Willd. ex Steud. Nom. ed. II. i. 39 = Muehlenbergia ciliata.
cimicina, Poir. Encyc. Suppl. i. 258 = Panicum cimicinum.
Cinna, Lam. ex Roem. & Schult. Syst. ii. 373 = Cinna arundinacea.
Cinna, Retz. Obs. Fasc. v. 18 = Muehlenbergia mexicana.
clandestina, Spreng. Fl. Hal. Mant. 32 = Sporobolus asper.
clausa, *Phil. in Anales Univ. Chil.* (1873) 562.—Chili.
clausa, *Steud. Syn. Pl. Gram.* 169.—Mesopot.
clavata, *Trin. in Spreng. Neue Entdeck.* ii. 55.—Kamtschat.
coarctata, Ehrh. ex Hoffm. Fl. Germ. ed. II. i. 37 = alba.
coerulescens, DC. Fl. Fr. Suppl. 250 = Piptatherum coerulescens.
cognata, *Steud. Syn. Pl. Gram.* 421.—Reg. Magell.

AGROSTIS :—
colorata, Willd. ex Steud. Nom. ed. II. i. 40 = Muehlenbergia elegans.
comosa, Poir. Encyc. Suppl. i. 257 (= *Urachne parviflora*).
complanata, Ait. Hort. Kew. i. 96 = Chloris petraea.
composita, Poir. Encyc. Suppl. i. 254.—Carolina.
compressa, Willd. in Roem. & Ust. Mag. iv. (1790) XI. 39 = alba.
compressa, *Poir. Encyc. Suppl.* i. 258.—Carolina.
compressa, *Doell, in Mart. Fl. Bras.* ii. III. 27.—Bras.
compressa, Torr. Cat. Pl. N. York, 91 = Sporobolus compressus.
concinna, Tuckerm. in Journ. Am. Sc. Ser. II. xlv. (1843) 43 = canina.
condensata, Willd. ex Steud. Nom. ed. II. i. 40 = verticillata.
conferta, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. 312 = nana.
conferta, Willd. ex Steud. Nom. ed. II. i. 40 = alba.
conspicua, Roem. & Schult. Syst. ii. 364 = Arundo conspicua.
contracta, F. Muell. ex Hook. f. Fl. Tasm. ii. 116 = Deyeuxia scabra.
Cornucopiae, *Sm. ex Fraser, Monog.* 2.—Am. bor.
Cornucopiae, Lam. ex Steud. Nom. ed. II. i. 40 = laxiflora.
coromandeliana, Roxb. Fl. Ind. ed. Wall. i. 318 (= *Vilfa Roxburghiana*).—Ind. or.
coromandelina, Retz. Obs. Fasc. iv. 19 = Sporobolus coromandelinus.
corralensis, *Phil. in Linnaea*, xxxiii. (1864-65) 286.—Chili.
cremenica, Bess. ex Steud. Nom. ed. II. i. 39, 40 = alba.
crinita, Moench, Meth. 178 = Polypogon monspeliensis.
crinita, *R. Br. Prod.* i. 170.—Austral.
crinita, Rich. ex. Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 4 = Dichelachne Forsteriana.
cruciata, Linn. Pl. Jam. Pugill. ii. 7; Syst. ed. X. 873 = Chloris cruciata.
cryptandra, Torr. in Ann. Lyc. N. York, i. (1824) 151 = Sporobolus cryptandrus.
cuspidata, Willd. ex Steud. Nom. ed. II. i. 40 = Calamagrostis toluccensis.
cylindrica, *R. Br. Prod.* i. 171.—Austral.
dalmatica, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 357.—Europ.
debilis, Poir. Encyc. Suppl. i. 249 = Deyeuxia Forsteri.
debilis, Spreng. Syst. i. 262 = Muehlenbergia debilis.
decipiens, R. Br. Prod. i. 172 = Deyeuxia scabra.
decipiens, Salzm. ex Steud. Syn. Pl. Gram. 101 = Eriochloa polystachya.
decumbens, Host, Ic. Gram. iv. t. 54 = alba.
decumbens, *Link, Hort. Berol.* i. 80.—Mexic.
delicatula, *Pourr. ex Kunth, Enum. Pl.* i. 219.—Lusitan.
delicatula, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
Delislei, *Hemsl. in Bot. Challeng.* i. II. 269.—Ins. Amsterdam.
densa, Bieb. Fl. Taur. Cauc. i. 56 = verticillata.
densa, *F. Muell. Census*, 133.—Austral.
Desvauxii, *Phil. in Linnaea*, xxxiii. (1864-65) 288.—Chili.
diandra, Hornem. in Syll. Ratisb. i. (1824) 193 = dianthera.
diandra, Retz. Obs. Fasc. v. 19 = Sporobolus diander.
dianthera, *Schult. Mant.* ii. 571.—Ind. or.
diaphora, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. II. (1845) 366 = Deyeuxia quadriseta.
difficilis, *Hemsl. Bot. Challeng.* i. II. 268. t. 43.—Ins. Amsterdam.
diffusa, Banks & Soland. ex Hook. f. Fl. New Zeal. i. 298 = Deyeuxia Billardieri.
diffusa, Bess. Prim. Fl. Galic. i. 68 = verticillata.
diffusa, Host, Ic. Gram. iv. t. 55 = alba.
diffusa, Mulh. Desc. Gram. 64 = Muehlenbergia sylvatica.
digitata, Lam. Encyc. i. 59 = Paspalum digitatum.
dispar, Host, Fl. Par. ex Steud. Nom. ed. II. i. 40 (= *Urachne parviflora*).
dispar, Michx. Fl. Am. Bor. i. 52 = elata?
distans, Kunze, Del. Sem. Hort. Lips. (1847) = Deyeuxia scabra.
disticha, Schweig. ex Steud. Nom. ed. II. i. 39, 40 = alba.
distichophylla, *Phil. Fl. Atac.* 54.—Chili.
distichophylla, *Roem. & Schult. Syst.* ii. 354.—Austral.

AGROSTIS :—
divaricata, *Griseb. in Goett. Nachr.* (1868) 81.—Ind. or.
divaricata, Hoffm. Fl. Germ. ed. II. i. 37 = vulgaris.
divaricata, Hort. ex Steud. Nom. ed. II. i. 40 = Chaeturus fasciculatus.
divaricata, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 713 = alba.
domingensis, Schult. Mant. iii. 570 = Sporobolus domingensis.
Dregeana, *Steud. Syn. Pl. Gram.* 173.—Afr. austr.
dubia, DC. Fl. Fr. iii. 21 = vulgaris.
dubia, Leers, Fl. Herborn. 21 = canina.
dubia, Auct. ex Steud. Nom. ed. II. i. 40 = Chaeturus fasciculatus.
dulcis, Hort. Par. ex Poir. Encyc. Suppl. i. 250 = alba.
dulcis, Sibth. ex Kunth, Enum. Pl. i. 218 = verticillata.
dura, Willd. ex Steud. Nom. ed. II. i. 40 = Sporobolus virginicus?
Duriaei, Boiss. & Reut. ex Nym. Consp. 802 = truncatula.
Ecklonis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 364.—Afr. austr.
effusa, Lam. Encyc. i. 59 = Milium effusum.
effusa, *Spreng. Syst.* i. 260.—Europ.
elata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 317.—Am. bor.
elatior, *Steud. Syn. Pl. Gram.* 174.—N. Zel.
elegans, Salisb. Prod. 25 = A. Cornucopiae.
elegans, Roem. & Schult. Syst. ii. 362 = Humboldtiana.
elegans, *Thore, ex Loisel. in Journ. Bot.* ii. (1809) 207. t. 8.—Europ.
Elliotiana, Schult. Mant. ii. 372 = arachnoides.
elongata, Lam. Illustr. i. 162 = Sporobolus indicus.
elongata, Roth, Nov. Sp. 41 = Sporobolus orientalis.
eminens, *Griseb. in Goett. Abh.* xix. (1874) 254.—Reg. Argent.
erecta, Spreng. Syst. i. 264 = Muehlenbergia aristata.
exarata, *Trin. Gram. Unifl.* 205; *Ic. Gram.* iii. t. 27.—Am. bor.
exasperata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 352.—Chili.
exigua, *Thurb. in S. Wats. Bot. Calif.* ii. 275.—Calif.
exilis, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 399 = elegans.
expansa, Poir. ex Steud. Nom. ed. II. i. 40 = Cinna arundinacea.
falklandica, *Hook. f. Fl. Antarct.* 373.—Ins. Falkland.
fascicularis, Sinel. Hort. Gram. Wob. 154 = canina.
fasciculata, *Roem. & Schult. Syst.* ii. 362.—Ecuador.
fertilis, Steud. Syn. Pl. Gram. 170 = Sporobolus elongatus.
festucoides, Muhl. ex Kunth, Enum. Pl. i. 200 = Muehlenbergia glomerata.
festucoides, Vill. Prosp. 16 = alpina.
filifolia, *Link, in Schrad. Journ.* ii. (1799) 313.—Lusitan.
filiformis, Koen. ex Kunth, Enum. Pl. i. 261 = Cynodon elongatus.
filiformis, Spreng. Fl. Hal. Mant. 32 = Deyeuxia Forsteri?
filiformis, Vill. Hist. Pl. Dauph. ii. 78 = alpina.
filiformis, Willd. Enum. Hort. Berol. 95 = Muehlenbergia mexicana.
flabellata, *Salzm. ex Steud. Syn. Pl. Gram.* 53 = Setaria sulcata?
flava, O. F. Muell. Fl. Dan. t. 751 = alba.
flavescens, Host, Ic. Gram. iv. 52 = alpina.
flavida, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 48 = alba.
flavidula, *Steud. Syn. Pl. Gram.* 421.—Reg. Magellan.
foliata, *Hook. f. Fl. Antarct.* 95.—N. Granat.
foliosa, Hort. ex Kunth, Enum. Pl. i. 207 = Muehlenbergia mexicana.
foliosa, *Vasey, in Bull. Torrey Bot. Club*, x. (1883) 63.—Am. bor.
Fonckii, *Phil. in Linnaea*, xxxiii. (1864–65) 286.—Chili.
Forsteri, Rich. ex Roem. & Schult. Syst. ii. 359 = Deyeuxia Forsteri.
frigida, F. Muell. ex Benth. Fl. Austral. vii. 583 = Deyeuxia frigida.
frondosa, *Guss. Fl. Sic. Prod.* i. 58.—Sicilia.
frondosa, Poir. Encyc. Suppl. i. 252 = Muehlenbergia mexicana.
frondosa, Tenore, ex Spreng. Pugill. ii. 5 = vulgaris.
fulva, *Griseb. in Goett. Abh.* xxiv. (1879) 294.—Reg. Argent.
gaditana, *Nym. Syll.* 408.—Lusitan.
gallecica, Hort. ex Steud. Nom. ed. I. 20; ed. II. i. 40 = Airopsis agrostidea.

AGROSTIS :—
Gayana, *E. Desv. in C. Gay, Fl. Chil.* vi. 313.—Chili.
gelida, F. Muell. in Trans. Vict. Inst. (1855) 43 = Muelleri.
gelida, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 343.—Peruv.
geminata, *Trin. Gram. Unifl.* 207.—Unalasch.
geniculata, Lam. Fl. Fr. iii. 572 = canina.
Ghiesbreghtii, '*Fourn. Mex. Pl. En. Gram.* 97,' *ex Hemsl. Biol. Centr. Am.* iii. 551.—Mexic.
gigantea, Roth, Tent. Fl. Germ. i. 31 = alba.
glabra, *Kunth, Enum. Pl.* i. 226.—Chili.
glabra, Presl, ex Trin. in Mém. Acad. Pétersb. Sér. VI. vi. II. (1845) 342 = stricta.
glabrata, Spreng. Syst. i. 262 = Muehlenbergia glabrata.
glauca, Muhl. Desc. Gram. 76 = Deyeuxia Nuttaliana.
glauca, *Steud. Syn. Pl. Gram.* 175.—Mexic.
glaucescens, Opiz, ex Steud. Nom. ed. II. i. 40 = parvula.
glaucescens, Spreng. Syst. i. 258 = sicula.
glaucina, Bast. Fl. Maine-et-Loire, Suppl. 25 = vulgaris.
globosa, Poir. Encyc. Suppl. i. 257 = Milium globosum.
glomerata, *Kunth, Enum. Pl.* i. 219.—Peruv.
glutinosa, Fisch. ex Kunth, l. c. 138 = Mclinis minutiflora.
gracilis, Kit. ex Schult. Fl. Austr. ed. II. i. 180; et in Linnaea, xxxii. (1863) 306 = alba.
gracilis, Willd. ex Steud. Nom. ed. II. i. 40 = Muehlenbergia gracilis.
gracilis, Salisb. Prod. 25 = Koeleria cristata.
graeca, Sibth. Fl. Graec. t. 66 (= *Urachne parviflora*).
grandis, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. II. (1845) 316 = exarata.
groenlandica, *Steud. Syn. Pl. Gram.* 175.—Groenland.
Gunniana, *F. Muell. Census*, 133.—Austral.
gynostyla, *Steud. Syn. Pl. Gram.* 170.—Afr. austr.
Hegetschweileri ×, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 60.—Europ.
Helenae, *Steud. Syn. Pl. Gram.* 171.—Ins. S. Helena.
herbacea, Steud. l. c. 224 = Aira paradoxa.
Hillebrandii, Thurb. ex Boland. in Trans. Calif. Agr. Soc. (1864) 136 = acquivalvis.
hirsuta, *Linn. f. Suppl.* 108.—Teneriff.
hirtella, *Hochst. ex Steud. Syn. Pl. Gram.* 173.—Abyssin.
hispanica, Boiss. & Reut. Pugill. 120 = castellana.
hispida, Brot. ex Nym. Consp. 802 = truncatula.
hispida, Willd. Sp. Pl. i. 370 = vulgaris.
holciformis, Bieb. Fl. Taur. Cauc. i. 54 = Oryzopsis holciformis.
Hornungiana, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 46 = vulgaris.
Humboldtiana, *Steud. Nom.* ed. II. i. 40.—Quito.
humifusa, Roem. & Schult. Syst. ii. 361 = Sporobolus humifusus.
humilis, *Vasey, in Bull. Torr. Club*, x. (1883) 21.—Am. bor.
hybrida, Gaud. Agrost. i. 66 = canina.
hygrometrica, Nees, in Mart. Fl. Bras. ii. 404 (= *Bromidium hygrometricum*).
hyperborea, Laest. ex Nym. Consp. 802 = rubra.
implicata, Spreng. Syst. i. 262 = Muehlenbergia implicata.
inaequiglumis, *Griseb. in Goett. Nachr.* (1868) 80.—Ind. or.
inconspicua, *Kunze, ex C. Gay, Fl. Chil.* vi. 315.—Chili.
incurvata, Scop. Fl. Carn. ed. II. i. 62 = Lepturus incurvatus.
indica, Forsk. Fl. Aegypt. Arab. 104. n. 66 = Sporobolus coromandelinus.
indica, Linn. Sp. Pl. 63 = Sporobolus indicus.
infirma, *Buese, in Miq. Pl. Jungh.* 342.—Java.
insignita, Schur, Enum. Pl. Transs. 732 = alba.
intermedia, Balb. Elench. 85 = vulgaris.
interrupta, Linn. Syst. ed. x. 872 = Apera interrupta.
intricata, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 413 = scabra.
involuta, Muhl. Desc. Gram. 72 = Sporobolus asper.
involuta, *Poir. Encyc. Suppl.* i. 252.—Marocc.
jamaicensis, Poir. l. c. 258 = Digitaria filiformis.
Jamesoniana, *Steud. Syn. Pl. Gram.* 163 = Am. merid.
japonica, *Steud. l. c.* 171.—Japon.
javanica, Burm. ex Kunth, Enum. Pl. i. 512 = Ischaemum muticum.
juncea, Lam. Encyc. i. 60 = Sporobolus tremulus.
juncea, Michx. Fl. Am. Bor. i. 52 = Sporobolus junceus.

AGROSTIS :—
junciformis, Willd. ex Steud. Nom. ed. II. i. 41 = Deyeuxia junciformis.
Juressi, *Link, in Schrad. Journ.* ii. (1799) 312.—Lusitan.; Hispan.
Kitaibeli, Schult. Fl. Austr. ed. II. i. 178 = alba.
koelerioides, *E. Desv. in C. Gay, Fl. Chil.* t. vi. 317.—Chili.
Labillardieri, Roem. & Schult. Syst. ii. 360 = Deyeuxia Billardieri.
lachnantha, *Nees, in Ind. Sem. Hort. Bot. Vratisl.* (1834).—Afr. austr.
Laersii, J. F. Gmel. Syst. ii. 169 = canina.
lanata, H. B. & K. Nov. Gen. et Sp. i. 136 = Epicampes lanata.
lanceolata, Roem. & Schult. Syst. ii. 364 = Calamagrostis lanceolata.
Langei, Nym. Consp. 801 = vulgaris.
lanigera, *Desv. Opusc.* 61.—Bras.
lasiantha, *Phil. in Linnaea*, xxix. (1857–58) 87.—Chili.
lateriflora, Michx. Fl. Bor. Am. i. 53 = Muehlenbergia mexicana.
latifolia, Trevir. in Goeppert, Beschr. Bot. Gaert. in Breslau, 82 = Cinna arundinacea.
laxa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 147 = vulgaris.
laxa, Schreb. ex Pursh, Fl. Am. Sept. i. 61 = scabra.
laxiflora, *Poir. Encyc. Suppl.* i. 255.—Am. bor.
laxiflora, Richards. Frankl. Voy. App. 3 = scabra.
lazica, *Balansa, in Bull. Soc. Bot. Fr.* xxi. (1874) 12.—Oriens.
Lechleri, *Steud. Syn. Pl. Gram.* 422.—Chili.
Leersii, J. F. Gmel. Syst. 1495 = canina.
lendigera, Poir. Encycl. Suppl. i. 259 = Gastridium australe.
lenta, Ait. Hort. Kew. i. 96 = Panicum filiforme.
leptos, *Steud. Syn. Pl. Gram.* 169.—Louisiana.
leptostachys, *Hook. f. Fl. Antarct.* 94.—Reg. Antarct.
leptotricha, *E. Desv. in C. Gay, Fl. Chil.* vi. 316.—Chili.
Lessoniana, Steud. Nom. ed. II. i. 41 = Arundo conspicua.
ligulata, *Steud. Syn. Pl. Gram.* 173.—Austral.
limonias, *Phil. in Linnaea*, xxxiii. (1864–65) 287.—Chili.
limosa, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 82; et Oestr. Bot. Zeitschr. ix. (1859) 48 = alba.
Lindeniana, *Steud. Syn. Pl. Gram.* 171.—Colombia.
linearis, Retz. Obs. Fasc. iv. 19 = Cynodon Dactylon.
lithuanica, Bess. ex Roem. & Schult Mant. 568, in add. = vulgaris.
litigans, *Steud. Syn. Pl. Gram.* 162.—Lusitan.
littoralis, Lam. Illustr. 161 = Sporobolus virginicus.
littoralis, With. Arr. Brit. Pl. ed. III. ii. 129. t. 23 = Polypogon littoralis.
lobata, R. Br. Prod. i. 171 = Deyeuxia quadriseta.
longiaristata, Herb. Ross. ex Kunth, Enum. i. 178 = Stipa splendens.
longiflora, Willd. ex Steud. Nom. ed. II. i. 41 = Cinna arundinacea.
longifolia, *Torr. ex A. Gray, Man. Bot. N. U. St.* ed. I. 575 = Sporobolus asper.
longiseta, Hort. ex Steud. Nom. ed. I. 21, 645 = Chaeturus fasciculatus.
lusitanica, Steud. Syn. Pl. Gram. 169 = castellana.
lutescens, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 299 = Arundo conspicua.
lutosa, Poir. Encycl. Suppl. i. 249 = Polypogon littoralis.
Lyallii, *Hook. f. Fl. N. Zeal.* 297.—N. Zel.
magellanica, *Lam. Illustr.* i. 160.—Reg. Magellan.
malaccensis, Willd. ex Kunth, Enum. Pl. i. 113 = Isachne pulchella.
maritima, Lam. Encyc. i. 61; Illustr. i. 161 = alba.
maritima, With. Arr. Brit. Pl. ed. III. ii. 132 = Elymus arenarius?
Matrella, Linn. Mant. ii. 185 = Zoysia pungens.
maxima, Roxb. Fl. Ind. i. 319 = Thysanolaena Agrostis.
media, *Carmich. in Trans. Linn. Soc.* xii. (1819) 504.—Ins. Trist. da Cunha.
melanosperma, Lam. Encyc. i. 58 = *Urachne Linnaei*.
melaleuca, Trin. in Mem. Acad. St. Petersb. ii. (1832) 170 = aenea.
Mertensii, *Trin. in Linnaea*, x. (1835) 302.—Chili: Unalaschk.
mexicana, *Kunth, Enum. Pl.* i. 223.—Mexic.
mexicana, Linn. Mant. 31 = Muehlenbergia mexicana.

AGROSTIS :—
mexicana, Pers. Syn. i. 76 = Deyeuxia canadensis.
Meyeni, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 312.—Chili.
Michaelis, *Steud. Syn. Pl. Gram.* 171.—Ins. Azor.
Michauxii, Trin. Gram. Unifl. 206 = scabra.
Michauxii, Zuccagn. in Roem. Collect. 123 = Chaeturus fasciculatus.
micrantha, *Steud. Syn. Pl. Gram.* 170.—Nepal.
microcarpa, Hort. ex Steud. Nom. ed. II. i. 164; ii. 41 = Muehlenbergia debilis.
microphylla, Steud. Syn. Pl. Gram. 164 = exarata.
microsperma, Lag. Elench. 2 = Muehlenbergia debilis.
miliacea, Linn. Sp. Pl. 61 = Piptatherum miliaceum.
Milium-comosum, Poir. Encyc. Suppl. i. 257 = Piptatherum miliaceum.
minima, Linn. Sp. Pl. 63 = Mibora verna.
minima, *Lour. Fl. Cochinch.* 50.—Cochinch.
minor, F. Muell. ex Benth. Fl. Austral. vii. 582 = Deyeuxia minor.
minutiflora, Desf. ex Steud. Nom. ed. II. i. 41 = Sporobolus minutiflorus.
minutiflora, Steud. Syn. Pl. Gram. 170 (= *Vilfa indica*).
minutissima, Steud. l. c. 171 = Sporobolus raminulosus.
monandra, Hornem. Hort. Hafn. i. 72 (= *Urachne parviflora*).
monostachya, Poir. Encyc. Suppl. i. 256 = Panicum amplexicaule.
montana, Krock. Fl. Siles. i. 110 = alpina.
montana, R. Br. Prod. i. 171 = Deyeuxia montana.
montevidensis, *Spreng. ex Nees, Agrost. Bras.* 403 = Montevid.
monticola, Roem. & Schult. Syst. ii. 359 = Deyeuxia montana.
Montis-aurei, Delarbre, Fl. Auverg. 8, ex Poir. Encyc. Suppl. i. 246 = Piptatherum multiflorum.
mucronata, *Presl, Rel. Haenk.* i. 238.—Am. bor. occ.
mucronata, Spreng. Syst. i. 262 = Muehlenbergia mucronata.
Muelleri, *Benth. Fl. Austral.* vii. 576.—Austral.
Muelleri, Presl, Bot. Bemerk. 120 = verticillata.
multicaulis, *Hook. f. Fl. Antarct.* 95.—Ins. Campbell.
multicaulis, Kit. in Linnaea, xxxii. (1863) 306 = alba.
Munroana, Aitch. & Hemsl. Journ. Linn. Soc. xix. (1881) 192 = Calamagrostis Munroana.
muricata, *Kunth, Enum. Pl.* i. 219.—Peruv.
muscosa, *T. Kirk, in Trans. N. Z. Inst.* xiii. (1881) 385.—N. Zel.
Mustaphae, *Steud. Syn. Pl. Gram.* 169.—Afr. bor.
mutabilis, Knapp, Gram. Brit. 28 = alba.
mutabilis, Sibth. Fl. Oxon. 36 = setacea.
Myuros, Lam. ex Beauv. Agrost. 49 = Panicum Myurus.
nana, Delarbre, Fl. Auverg. ed. II. 664 = pumila.
nana, *Kunth, Enum. Pl.* i. 266.—Peruv.
nardifolia, *Griseb. in Goett. Abh.* xix. (1874) 252.—Reg. Argent.
nebulosa, *Boiss. & Reut. Diagn.* 26.—Hispan.
nebulosa, Bourg. ex Reut. & Lange, in Bull. Soc. Bot. Fr. xi. (1864) 47 = delicatula.
Neesii, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 361.—Afr. austr.
neglecta, Schult. Fl. Austr. ed. II. i. 179 = alba.
nemoralis, *Phil. in Linnaea*, xxx. (1859-60) 205.—Chili.
neogaea, *Steud. Syn. Pl. Gram.* 171.—N. Zembl.
nervosa, *Nees, ex Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 328.—Ind. or.
nevadensis, *Boiss. Elench.* 87.—Hispan.
nigra, With. Arr. Brit. Pl. ed. III. ii. 131 = vulgaris.
nigrescens, *Salzm. ex Steud. Nom.* ed. II. i. 41.—Bahia.
nigricans, Poir. Encyc. Suppl. i. 259 = Milium nigricans.
nitens, Guss. Fl. Sic. Prod. i. 50 = Triplachne nitens.
nivalis, F. Muell. in Trans. Vict. Inst. (1855) 43 = Deyeuxia nivalis.
nivea, Sincl. Hort. Gram. Woburn. 152 = alba?
nobilis, De Not. ex Parl. Fl. Ital. i. 186 = canina.
nootkaensis, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. II. (1845) 326 = nutkaensis.
Novae-Hollandiae, Beauv. Agrost. 148 = Deyeuxia Forsteri.
noveboracensis, Spreng. Syst. i. 260 = Epicampes lanata.
nutans, Poir. Encyc. Suppl. i. 255 = Ichnanthus pallens.
nutans, Spreng. Syst. i. 263 = Muehlenbergia gracilis.
nutkaensis, *Kunth, Enum. Pl.* i. 222.—Am. bor. occ.
obliqua, Auct. ex Trin. Gram. Unifl. 196 = alba.

AGROSTIS :—
obliqua, *Kit. in Linnaea*, xxxii. (1863) 306.—Austria.
obtusata, Auct. ex Steud. Nom. ed. II. i. 41 = Eatonia obtusata.
odorata, *Lour. Cochinch.* 64.—Cochinch.
oligantha, Roem. & Schult. Syst. ii. 372 = pauciflora.
olivetorum, *Gren. & Godr. Fl. Fr.* iii. 483.—Gallia.
oreophila, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 323.—Am. bor.
orientalis, Nees, Agrost. Bras. 393 = Sporobolus tenacissimus.
oseroensis, Scenus, Reise Istr. (1805) 50 (= *Urachne parviflora*).
ovata, Forst. f. Prod. 8 = Echinopogon ovatus.
owarensis, Roem. & Schult. Syst. ii. Mant. 199 = Sporobolus pyramidalis.
pallens, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 328.—Am. bor.
pallida, *DC. Fl. Fr. Suppl.* 251.—Europ.
pallida, Schk. Handb. 3; Nachtr. 1 = canina.
pallida, With. Arr. Brit. Pl. ed. III. ii. 128. t. 22 = alba.
palustris, Huds. Fl. Angl. 27 = alba.
panicea, Lam. Encyc. i. 58; Illustr. 811 = Gastridium australe.
panicea, Ait. Hort. Kew. i. 94 = Polypogon monspeliensis.
panicoides, Lam. Illustr. 162 = Panicum Lamarckii.
paniculata, Hort. Madrit. ex Steud. Nom. ed. II. i. 41 = Muehlenbergiae sp. sec. Trin.
papposa, Gilib. Exercit. ii. 524 = Stipa Calamagrostis.
paradisiaca, Steud. Syn. Pl. Gram. 163 = chilensis.
paradoxa, Linn. Sp. Pl. 62 (= *Urachne Linnaei*).
paradoxa, R. Br. in Ross, Voy. ed. II. ii. 192 = Colpodium latifolium.
parviflora, R. Br. Prod. 170 = scabra.
parvula, Schult. Fl. Austr. ed. II. i. 182 = Sporobolus minutus.
patagonica, *Phil. in Linnaea*, xxix. (1857-58) 87.—Patagon.
patens, *Trin. in. Mém. Acad. Pétersb.* Sér. VI. vi. II. (1845) 322.—Chili.
patula, Gaud. Agrost. Helv. i. 80 = alba.
pauciflora, *Pursh, Fl. Am. Sept.* i. 63.—Am. bor.
pauciflora, Schrad. Fl. Germ. 3 = alba.
paucinodis, *Hack. in Engl. Bot. Jahrb.* vi. (1885) 242.—Reg. Magell.
perennans, *Tuckerm. in Am. Journ. Sc.* Ser. II. xlv. (1843) 44.—Am. bor.
pernambucensis, Spreng. Syst. i. 258 = Panicum rivulare.
peruviana, Spreng. l. c. 262 = Muehlenbergia Clomena.
phleoides, *E. Desv. in C. Gay, Fl. Chil.* vi. 320.—Chili.
phleoides, Willd. ex Steud. Nom. ed. II. i. 41 = Calamagrostis coarctata.
Pickeringii, *Tuckerm. in Am. Journ. Sc.* Ser. II. xlv. (1843) 42.—Am. bor.
pilosa, Schleich. ex Gaud. Agrost. Helv. i. 75 = Calamagrostis tenella.
pilosa, *Less. in Rich. Voy. Astrol.* 134. t. 23.—N. Zel.
pilosa, Retz. Obs. Fasc. vi. 22 = Cynodon pilosus.
pilosula, *Trin. in Mém Acad. Pétersb.* Sér. VI. vi. II. (1845) 372.—Ind. or.
pisidica, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 255 = trichoclada.
planifolia, *C. Koch, in Linnaea*, xxi. (1848) 380.—Armen.
plebeia, R. Br. Prod. i. 172 = Deyeuxia plebeia.
plicata, *Lour. Fl. Cochinch.* 64.—China.
plumosa, Tenore, Prod. i. 59 = Panicum Teneriffae.
poaeformis, Willd. ex Steud. Nom. ed. II. i. 42 = Poa subuniflora.
polymorpha, Huds. Fl. Angl. ed. II. i. 31 = alba, vulgaris.
Polypogon, Salzm. ex Steud. Syn. Pl. Gram. 178 = Muehlenbergia brasiliensis.
polystachya, Bosc, ex Steud. Nom. ed. II. i. 42 = composita.
Pourretii, Spreng. Syst. i. 260 = Avena Pourretii.
Pourretii, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 290 = effusa.
praecox, Salisb. Prod. 24 = Aira praecox.
Preslii, *Kunth, Enum. Pl.* i. 225.—Peruv.
procera, A. Rich. Voy. Astrol. 125 = Arundo conspicua.
procera, *Retz. Obs.* iv. 19.—Malabar.
procumbens, Roem. & Schult. Syst. ii. 371 = alba.

AGROSTIS :—
prorepens, G. Mey. Korresp. Bl. Naturf. Ver. Riga, xxvi. 56 (quid?).—Rossia.
prostrata, *Hook. f. Fl. Antarct.* 373.—Ins. Falkland.
pseudo-arundinacea, Schleich. ex Gaud. Agrost. Helv. i. 92 = Calamagrostis varia.
pubens, Willd. ex Steud. Nom. ed. II. i. 42 = Epicampes lanata.
pubescens, S. F. Gray, Nat. Arr. Brit. Pl. ii. 148 = alba.
pubescens, H.B. & K. Nov. Gen. et Sp. i. 136 = Epicampes pubescens.
pulchella, *Guss. Fl. Sic. Prod.* i. 58.—Sicilia.
pulchella, Kunth, Gram. i. 70 = Humboldtiana.
pulchella, Loisel. ex Trin. Gram. Unifl. 205 = elegans.
pulchella, Roth, ex Roem. & Schult. Syst ii. 367 = Sporobolus pulchellus.
pumila, Linn. Mant. 31 = vulgaris.
pumila, *Steud. Syn. Pl. Gram.* 171.—Ins. Adscens.
punctata, Lam. Encyc. i. 58. n. 17 = Eriochloa polystachya.
pungens, Muhl. Desc. Gram. 72 = Zoysia pungens.
pungens, Pall. 'Ind. Taur.' ex Georgi, Beschr. iii. iv. 688 = Aeluropus litoralis.
pungens, Schreb. Graes. ii. 46 = Sporobolus pungens.
pungens, Pursh, Fl. Am. Sept. 64 = Sporobolus virginicus.
purpurascens, *Bert. ex Steud. Nom.* ed. II. i. 42.—S. Doming.
purpurascens, Sw. Prod. Veg. Ind. Occ. 25 = Sporobolus purpurascens.
purpurea, Gaud. Agrost. Helv. i. 70 = Apera Spica-venti.
pusilla, Dum. Obs. Gram. Belg. 129 = canina.
pyramidalis, *Rich. ex Steud. Syn. Pl. Gram.* 153 (= *Vilfa Richardi*).—Ind. occ.
pyramidata, Lam. Illustr. i. 161 = Sporobolus affinis.
pyrenaea, Timb. in Mém. Acad. Toul. Sér. IV. vi. (1856) 97 = alpina.
pyrenaica, Pourr. l. c. iii. (1788) 306 = rupestris.
quadridentata, Spreng. Syst. i. 263 = Muehlenbergia quadridentata.
quadrifida, Labill. Pl. Nov. Holl. i. 20 = Pentapogon Billardieri.
quadriseta, R. Br. Prod. i. 171 = Deyeuxia quadriseta.
quitensis, Spreng. Syst. i. 262 = Muehlenbergia elegans.
quitensis, Willd. ex Steud. Nom. ed. II. i. 42 = Muehlenbergia calamagrostidea.
racemosa, Michx. Fl. Bor. Am. i. 53 = Muehlenbergia glomerata.
radiata, Linn. Pl. Jam. Pugill. 7; Syst. ed. X. 873 = Chloris radiata.
ramosa, Poir. Encyc. Suppl. i. 257 = Eriochloa annulata.
ramosa, Roem. & Schult. Syst. ii. 364 = Calamagrostis lanceolata.
ramulosa, Carm. in Trans. Linn. Soc. xii. (1818) 504 = Carmichaeli.
ramulosa, Roem. & Schult. Syst. ii. 361 = Sporobolus ramulosus.
rara, R. Br. Prod. i. 171 = Dichelachne sciurea.
rata, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 3 = Dichelachne sciurea.
Ravennae, Beauv. Agrost. 148 = Erianthus Ravennae.
refracta, Moench, Meth. Suppl. 60 = verticillata.
Reinwardtii, *Miq. Fl. Ind. Bat.* iii. 750.—Malaya.
repens, Sincl. Hort. Gram. Woburn. 148 = alba.
retrofracta, Willd. Enum. Hort. Berol. 94 = Deyeuxia Forsteri.
Reuterii, *Boiss. Voy. Espagne*, ii. 645.—Hispan.
Reygeri, [Honck.] Verz. Aller Gew. Teutschl. 204 = alpina?
rigescens, *Presl, in Rel. Haenk.* i. 237.—Peruv.
rigida, Lesson, ex A. Rich. Voy. Astrol. 124 = Dichelachne rigida.
rigida, Spreng. Syst. i. 262 = Muehlenbergia rigida.
rigida, Willd. ex Steud. Nom. ed. II. i. 42 = Calamagrostis recta.
rigidula, *Steud. Syn. Pl. Gram.* 171.—Java.
rigidifolia, Poir. Encyc. Suppl. i. 257 = Panicum rigidifolium.
rivularis, Brot. Fl. Lusit. i. 75 = verticillata.
rosea, *Griseb. in Goett. Abh.* xix. (1874) 253.—Reg. Argent.
Royleana, *Trin. Agrost.* ii. 125.—Ind. or.
rubicunda, Bosc, ex Desv. Opusc. 55 = Cinna arundinacea.
rubra, Auct. ex Steud. Nom. ed. II. i. 42 = canina.

AGROSTIS :—
rubra, Huds. Fl. Angl. 26 = Gastridium australe.
rubra, *Linn. Sp. Pl.* 62.—Europ.
rubra, Wahlenb. ex Nym. Consp. 801 = vulgaris.
rudis, Roem. & Schult. Syst. ii. 360 = Deyeuxia scabra.
rupestris, *All. Fl. Pedem.* ii. 237.—Europ.; Am. bor.
rupestris, Willd. ex Kunth, Gram. i. 70 = alpina.
Ruprechtii, *Boiss. Fl. Orient.* v. 517.—Caucas.
salina, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 366 = alba.
salmantica, *Kunth, Gram.* i. 71.—Hispan.
scaberrima, *Hassk. Pl. Jav. Rar.* 26.—Java.
scabra, R. Br. Prod. 172 = Deyeuxia scabra.
scabra, *Tuckerm. in Am. Journ. Sc.* Ser. II. xlv. (1845) 44.—Am. bor.
scabra, *Willd. Sp. Pl.* i. 370.—Am. bor.
scabriglumis, Boiss. & Reut. Pugill. 125 = alba.
scabriuscula, Buckl. in Proc. As. Sc. Philad. '1862.' (1863) 90 = scabra.
Schaffneri, *Fourn. Mex. Pl. En. Gram.* 94, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 551.—Mexic.
Schiedeana, *Fourn. l. c.* 96, *ex Hemsl. l. c.*—Mexic.
Schiedeana, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 327.—Mexic.
Schimperiana, *Hochst. ex Steud. Syn. Pl. Gram.* 170.—Abyss.
Schleicheri, Jord. & Verlot, in F. Schultz, Arch. Fl. Fr. et Allem. 339, 347 = alpina.
Schottii, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 356.—Gibraltar.
Schultesii, *Kunth, Enum. Pl.* i. 228.—Pannonia.
Schweinitzii, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. (1845) 311 = elata.
sciurea, R. Br. Prod. i. 171 = Deyeuxia sciurea.
scoparia, Koen. ex Steud. Syn. Pl. Gram. 115 = Panicum semialatum.
Scouleri, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 329.—Am. bor. occ.
semibarbata, Trin. l. c. 378 = Deyeuxia Forsteri.
sepium, Jacq. Enum. Stirp. Vindob. 12 (= *Urachne parviflora*).
sericea, Banks & Soland. ex Hook. f. Fl. Nov. Zel. i. 299 = Arundo conspicua.
sericea, Ell. Sketch, i. 135 = Muehlenbergia capillaris.
serotina, Linn. Mant. 30 = Diplachne serotina.
serotina, Torr. Fl. Un. St. i. 88 = Sporobolus serotinus.
sesquiflora, *E. Desv. in C. Gay, Fl. Chil.* vi. 318.—Chili.
setacea, *Curt. Fl. Lond.* vi. t. 12.—Europ.
setacea, Vill. Hist. Pl. Dauph. ii. 76 = rupestris.
setifolia, Brot. Fl. Lusit. i. 74 = setacea.
setifolia, Willd. ex Steud. Nom. ed. II. i. 42 = tenuifolia.
setifolia, *Hook. f. Handb. New Zeal. Fl.* 329.—N. Zel.
setifolia, *Fourn. Mex. Pl. En. Gram.* 97; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 551.—Mexic.
setosa, Muhl. Descr. Gram. 68 = Muehlenbergia glomerata.
setosa, Spreng. Syst. i. 262 = Muehlenbergia debilis.
Sibthorpii, Hort. ex Steud. Nom. ed. II. i. 42 = verticillata.
sicula, Guss. ex Steud. Nom. ed. II. i. 42 = Gastridium muticum.
sicula, *Kunth, Rev. Gram.* i. 71.—Sicilia.
signata, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 47 = alba.
simensis, *Hochst. ex Steud. Syn. Pl. Gram.* 173.—Abyss.
simulans, *Hemsl. in Hook. Ic. Pl.* t. 1455.—Ins. Sta. Helena.
sinaica, Boiss. Diagn. Ser. I. xiii. 46 = alba.
sobolifera, Muhl. ex Willd. Enum. Hort. Berol. 95 = Muehlenbergia sobolifera.
Solandri, F. Muell. Veg. Chath. Isl. 60 = Deyeuxia Forsteri.
spicaeformis, Linn. f. Suppl. 108 = Perotis latifolia.
spicata, Thunb. Prod. Pl. Cap. 19 = Sporobolus capensis.
spicata, Vahl, Symb. i. 9 = Sporobolus spicatus.
Spica-venti, Linn. Sp. Pl. ed. 1. 61 = Apera Spica-venti.
Sporobolus, Spreng. Nov. Prov. 46 = Sporobolus Sprengelii.
spuria, Roem. & Schult. Syst. ii. 346 = Calamagrostis lanceolata.
stellata, Willd. Sp. Pl. i. 376 = Cynodon Dactylon.
stipata, Koel. Gram. 77 = Stipa Calamagrostis.
stipoides, Spreng. Syst. i. 263 = Muehlenbergia stipoides.
stolonifera, Host, Gram. iv. t. 56 = alba.

AGROSTIS :—
stolonifera, Leers, Fl. Herb. t. 4. f. 6 = vulgaris.
stolonifera, *Linn. Sp. Pl.* 62.—Europ.
stolonifera, Linn. ex Sm. Fl. Brit. i. 80 = verticillata.
stolonizans, Bess. in Roem. & Schult. Mant. 567, in add. = alba.
straminea, Hartm. Gram. Scand. 4 = alba.
stricta, *Buese, in Miq. Pl. Jungh.* 341.—Java.
stricta, *De Not. ex Parl. Fl. Ital.* i. 187.—Italia.
stricta, J. F. Gmel. Syst. 170 = alba.
stricta, Sincl. Hort. Gram. Woburn. 150 = canina.
stricta, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 342.—Chili.
stricta, *Willd. Sp. Pl.* i. 366.—Am. bor.
suaveolens, Blytt, in Mag. for Naturvidensk. (1837); sec. Sommerfelt, in Skand. Akad. Handl. (1837) 25 = Cinna arundinacea.
subaristata, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 192.—Afghan.
subulata, *Hook. f. Fl. Antarct.* 95.—Ins. Campbell.
sylvatica, Georgi, Beschr. Russ. Reich. iii. 687 = Calamagrostis sylvatica.
sylvatica, Huds. Fl. Angl. 28 = alba.
sylvatica, Koel. Gram. 92 = vulgaris.
sylvatica, Torr. Fl. N. Amer. i. 87 = Muehlenbergia Willdenowii.
tabularis, Steud. Nom. ed. II. i. 42 = Bergiana.
tacubayensis, *Fourn. Mex. Pl. En. Gram.* 95, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 551.—Mexic.
tarda, *Drude, in Flora*, lx. (1877) 273.—Tyrolia.
tenacissima, Linn. f. Suppl. 197 = Sporobolus indicus.
tenella, Hoffm. Fl. Germ. ed. II. i. 36 = vulgaris.
tenella, Poir. Encyc. Suppl. i. 257 = Airopsis globosa.
tenella, Willd. ex Steud. Nom. ed. II. i. 42 = Muehlenbergia tenuifolia.
tenella, Schleich. ex Gaud. Agrost. Helv. i. 75 = Calamagrostis tenella.
tenerrima, Trin. Gram. Unifl. 205 = elegans.
Tenorii, Otto, ex Loud. Hort. Brit. 26.—Italia?
tenuiflora, Willd. Sp. Pl. i. 364 = Muehlenbergia Willdenowii.
tenuiflora, Steud. Syn. Fl. Gram. 163 = perennans.
tenuifolia, Curt. Brit. Gr. 42 = canina.
tenuis, Bast. ex Roem. & Schult. Syst. ii. 277 = canina.
tenuis, Sibth. Fl. Oxon. 36 = vulgaris.
tenuis, *Vasey, in Bull. Torr. Club*, x. (1883) 21.—Calif.
tenuissima, Spreng. Syst. i 258 = Sporobolus indicus.
tenuissima, Vest, ex Steud. Nom. ed. II. i. 42 = Polypogon monspeliensis.
Thunbergii, *Steud. Syn. Pl. Gram.* 163.—Japon.
thyrsoides, Bosc, ex Steud. Nom. ed. II. i. 42 (= *Vilfa fulvescens*).
Tillierii, Willd. ex Steud. l. c. = Catabrosa aquatica.
tolucensis, *H. B. & K. Nov. Gen. et Sp.* i. 135.—Mexic.
toluccensis, Willd. ex Steud. Syn. Pl. Gram. 164 = virescens.
Torreyana, Schult. Mant. ii. 203 = compressa.
Torreyi, *Kunth, Enum. Pl.* i. 226.—Am. bor.
transilvanica, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 50 = canina.
tremula, Willd. Sp. Pl. i. 372 = Sporobolus tremulus.
triaristata, Knapp, Gram. Brit. 23 = Polypogon monspeliensis.
trichantha, *Schrank, in Regensb. Bot. Denkschr.* ii. (1818) 4.—Labrador.
trichoclada, *Griseb. in Ledeb. Fl. Ross.* iv. 439.—Oriens.
trichoides, *Roem. & Schult. Syst.* ii. 361.—Peruv.
trichopodes, Ell. Sketch, i. 135 = Muehlenbergia trichopodes.
trichotoma, Auct. ex Steud. Nom. ed. II. i. 43 = alba.
trichotoma, *Nees, ex Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 338.—Monte Video.
trichotoma, *Kit. in Linnaea*, xxxii. (1863) 306.—Austria.
tricuspidata, Hack. in Oestr. Bot. Zeitschr. xxvii. (1877) 46 = canina.
triflora, Hort. Hafn. ex Steud. Nom. ed. II. i 43 = Chaeturus fasciculatus.
trinervata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 347.—Caucas.
trisetoides, *Steud. Syn. Pl. Gram.* 172.—Chili.
tropica, *Beauv. Fl. Oware*, ii. 37.—Afr. trop.

AGROSTIS :—
truncatula, *Parl. Fl. Ital.* i. 185.—Europ.
uliginosa, *Phil. in Anales Univ. Chil.* (1865) II. 323.—Chili.
umbellata, *Colla, in Mem. Acc. Torin.* xxxix. (1836) 23.—Chili.
umbellulata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi (1845) 370.—Afr. austr.
umbrosa, *Pers. ex Trin. l. c.* 345.—Gallia.
umbrosa, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 47 = vulgaris.
vaginaeflora, Torr. in A. Gray, Man. 576 = Sporobolus vaginaeflora.
vaginata, *Steud. Syn. Pl. Gram.* 173.—Ins. Sandvic.
valentina, *Roem. & Schult. Syst.* ii. 348.—Hispan.
valvata, *Steud. Syn. Pl. Gram.* 171.—Japon.
varia, Host, Gram. iv. t. 57; Fl. Austr. i. 100 = alba.
varians, Thuil. Fl. Par. ii. 35 = canina.
varians, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 314.—Am. bor.; Chili.
velutina, Poir. ex Steud. Nom. ed. II. i. 43 = Paspalum velutinum.
ventosa, Dulac, Fl. Hautes-Pyr. 74 = Apera Spica-venti.
ventricosa, Gouan, Hort. Monsp. 39, 547 = Gastridium australe.
venusta, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 340.—Austral.
vernalis, Hort. Dorp. ex Trin. Gram. Unifl. 198 = alba.
vernalis, Poir. Encyc. Suppl. i. 259 = Milium vernale.
versicolor, Hoppe, ex Steud. Nom. ed. II. i. 43 = alpina.
versicolor, Hort. Hafn. ex Trin. Gram. Unifl. 201, 203 = vulgaris.
versicolor, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 282 = Catabrosa versicolor.
verticillata, Lam. Encyc. i. 59; Illustr. n. 831 = Andropogon squarrosus.
verticillata, Thuil. Fl. Par. ii. 36 = vulgaris.
verticillata, *Vill. Prosp.* 16.—Europ.
vestita, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 401.—Abyss.
Vidali, *Phil. in Anales Univ. Chil.* (1873) 561.—Chili.
Villarsii, Poir. Encyc. Suppl. i. 251 = verticillata.
villosa, A. Spreng. Syst. Suppl. 5 = Sporobolus ciliatus.
villosa, Poir. ex Steud. Nom. ed. II. i. 43 = Tricholaena insularis.
villosa, Reichb. ex Steud. l. c. = Sporobolus villosus.
villosa, Vill. Hist. Pl. Dauph. i. 378 = Calamagrostis sylvatica.
vinealis, [Honck.] Verz. aller Gew. Teutschl. 205 = castellana.
vinealis, With. Arr. Brit. Pl. ed. III. ii. 127 = canina.
vinealis, Schreb. Spicil. 47 = alba.
violacea, *Phil. in Anales Univ. Chil.* (1873) 560.—Chili.
violacea, Thuil. Fl. Par. ii. 35 = vulgaris.
violaceo-purpurea, Gilib. Exercit. ii. 525 = canina.
virescens, *H. B. & K. Nov. Gen. et Sp.* i. 135.—Calif.; Mexic.
virescens, Rupr. in Bull. Acad. Brux. ix. (1842) 229 = tolucensis.
virgata, Willd. ex Steud. Nom. ed. II. i. 43 = Leptochloa tenerrima.
virginica, Forsk. Fl. Aegypt. Arab. 20 = Sporobolus spicatus.
virginica, Linn. Sp. Pl. 63 = Sporobolus virginicus.
virginica, Muhl. ex Steud. Nom. ed. II. i. 43 = Sporobolus vaginaeflorus.
viridis, Gouan, Hort. Monsp. 546 = Sporobolus pungens.
viridis, *Rafin. in Am. Monthly Mag.* (1811) 356, nomen.—Am. bor.
Virletii, *Fourn. Mex. Pl. En. Gram.* 96, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 551.—Mexic.
vivipara, Bivona, ex Roem. & Schult. Syst. ii. 347 in nota = alba.
vulgaris, *With. Arr. Brit. Pl.* ed. III. ii. 132.—Reg. temp.
Wallichiana, *Steud. Syn. Pl. Gram.* 174.—Reg. Himal.
Welwitschii, Steud. l. c. 168 = Reuteri.
Wightii, *Nees, ex Steud. l. c.*—Ind. or.
Youngii, *Hook. f. Handb. New Zeal. Fl.* 330.—N. Zel.
Zenkeri, *Trin. in Mém. Acad. Pétersb.* Sér. VI. vi. (1845) 363.—Ind. or.
zeylanica, *Klein, ex Steud. Syn. Pl. Gram.* 83.—Zeylan.

AGROSTISTACHYS, Dalz. in Hook. Kew Journ. ii. (1850) 41. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 302.
SARCOCLINIUM, Wight, Ic. v. 24. t. 1887, 1888, (1852).
africana, *Muell. Arg. in Flora*, xlvii. (1864) 534.—Afr. trop.
Gaudichaudii, *Muell. Arg. in Linnaea*, xxxix. (1865–66) 144.—Ind. or.
Hookeri, *Benth. & Hook. f. Gen.* iii. 303.—Zeylan.
indica, *Dalz. in Hook. Kew Journ. Bot.* ii. (1850) 41.—Ind. or.
longifolia, *Benth. ex Hook. f. Fl. Brit. Ind.* v. 407.—Canara.
longifolia, Kurz, For. Fl. Brit. Burma, ii. 377 = Gaudichaudii.

AGROSTOCRINUM, F. Muell. Fragm. ii. 94 (1860). *LILIACEAE*, Benth. & Hook. f. iii. 792.
stypandroides, *F. Muell. l. c.* 95.—Austral.

AGROSTOMIA, Cerv. in La Naturaleza (1870) 345 = **Panicum**, Linn. (Gramin.).
aristata, Cerv. l. c.—Mexic.
barbata, Cerv. l. c. 346.—Mexic.
mutica, Cerv. l. c. 345.—Mexic.

AGROSTOPHYLLUM, Blume, Bijdr. 368. t. 53 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 517.
DIPLOCONCHIUM, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 428. t. 12.
callosum, *Reichb. f. in Seem. Fl. Vit.* 296.—Reg. Himal.
Hasseltii, *Reichb. f. in Bonplandia*, v. (1857) 41; *et in Walp. Ann.* vi. 909.—Ind. or.
javanicum, *Blume, Bijdr.* 369. t. 53.—Java.
khasiyanum, *Griff. in Calc. Journ. Nat. Hist.* iv. (1844) 376.—Reg. Himal.
longifolium, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.
megalurum, *Reichb. f. in Seem. Fl. Vit.* 296.—Ins. Samoa.
planicaule, Reichb. f. in Walp. Ann. vi. 909 = khasiyanum.

AGUAVA, Rafin. Sylva Tellur. 107 (1838) = **Myrcia**, DC. (Myrtac.).
guianensis, Rafin. l. c. = M. guianensis.
tomentosa, Rafin. l. c. = M. tomentosa.

AGYLLA, Phil. in Anales Univ. Chil. (1865) i. 643 = **Cladium**, P. Br (Cyperac.).
ensifolia, Phil. l. c. = C. scirpoideum.

AGYLOPHORA, Neck. Elem. i. 145 (1790) = **Uncaria**, Schreb. (Rubiac.).

AGYNAIA, Hassk. in Flora, xxv. (1842); Beibl. 40 = seq.

AGYNEIA, Linn. Mant. ii. 161 (1771); Vent. Jard. Cels, t. 23 (1800). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 271.
DIPLOMORPHA, Griff. Notul. iv. 479 (1854).
affinis, Kurz, ex Teijsm. & Binn. in Tijdschr. Ned. Ind. xxvii. (1864) 118 = bacciformis.
bacciformis, *A. Juss. Tent. Euph.* 24.—Ind. or.; Malay.
Berterii, Spreng. Syst. iii. 19 = Phyllanthus grandifolius.
ciliata, Wall. Cat. n. 7852 = Trigonostemon semperflorens.
coccinea, [Buch.-Ham. in] Symes, Embassy, ed. II. iii. 317 = Glochidion coccineum.
flexuosa, Heyne, ex Wall. Cat. n. 7863 = Glochidion zeylanicum.
glomerulata, *Miq. Fl. Ind. Bat. Suppl.* 447.—Sumatra.
hirsuta, *Miq. l. c.* i. ii. 368.—Malaya.
impubera, *Miq. l. c.* 367.—Malaya.
impubera, Wall. Cat. n. 7869 = Glochidion multiloculare.
impubes, Herb. Madr.? Wall. Cat. n. 7990 partim = Glochidion littorale.
impubes, Linn. Mant. ii. 296 = Phyllanthus puberus.
impubes, Vent. Jard. Cels, 23. t. 23 = bacciformis.
lanceolata, *F. G. Dietr. Lexik. Gaertn.* i. 186.—Austral.?
latifolia, Moon, Cat. 65 = Aporosa latifolia.

AGYNEIA:—
multiflora, Hassk. Cat. Hort. Bog. Alt. 240 (= Phyllanthi, sp.).—Malaya.
multiloculare, Rottl. ex Willd. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 206 = Glochidion multiloculare.
obliqua, Willd. Sp. Pl. iv. 568 = Glochidion zeylanicum.
ovata, Poir. Encyc. Suppl. i. 243; Miq. Fl. Ind. Bat. i. ii. 367 = Sauropus albicans.
phyllanthoides, Spreng. Syst. iii. 19 = bacciformis.
pinnata, Miq. Fl. Ind. Bat. i. ii. 368 = Glochidion obscurum.
pubera, Linn. Mant. ii. 296 = Phyllanthus puberus.
sinica, Miq. Fl. Ind. Bat. i. ii. 368 = Phyllanthus puberus.
tenera, Zoll. & Mor. ex Miq. Fl. Ind. Bat. i. ii. 365 = Andrachne australis.
tetrandra, Buch.-Ham. in Trans. Linn. Soc. xv. (1826) 125 = Phyllanthus Hamiltonianus.
tetrandra, Wall. Cat. n. 7852 = Trigstemon Hookerianus.

AHOUAI, Tourn. ex Adans. Fam. ii. 171 (1763) = **Thevetia**, Linn. (Apocyn.).

AICHRYSON, Webb & Berth. Phyt. Canar. i. 180. t. 27 (1840) = **Sempervivum**, Linn. (Crassul.).
Bethencourtianum, C. Bolle, in Bonplandia, vii. (1859) 243.—Ins. Canar.
Bollei, Webb, ex Bolle, l. c.—Ins. Canar.
dichotomum, Webb & Berth. Phyt. Canar. i. 181 = S. dichotomum.
pachycaulon, C. Bolle, in Bonplandia, vii. (1859) 244.—Ins. Canar.
palmense, Webb, ex C. Bolle, l. c. 243.—Ins. Canar.
Parlatorei, C. Bolle, l. c. 244.—Ins. Canar.
parviflorum, C. Bolle, l. c. 243.—Ins. Canar.
porphyrogennetos, C. Bolle, l. c.—Ins. Canar.
pulchellum, C. A. Mey. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 55.—Hab.?
punctatum, Webb & Berth. Phyt. Canar. i. 182 = S. punctatum.
pygmaeum, Webb & Berth. l. c. 184.—Ins. Canar.
radicescens, Webb & Berth. l. c. 183 = S. villosum.
sedifolium, Webb, ex C. Bolle, in Bonplandia, vii. (1859) 242.—Ins. Canar.
tortuosum, Webb & Berth. Phyt. Canar. i. 184 = S. tortuosum.
villosum, Webb & Berth. l. c. 181 = S. villosum.

AIDELUS, Spreng. Syst. iv. Cur. Post. 17 (1827) = **Veronica**, Tourn. (Scrophul.).
micranthus, Spreng. l. c.—Nepal.

AIDIA, Lour. Fl. Cochinch. 143 (1790). Gen. haud recogn. Cf. Benth. & Hook. f. ii. 29.
cochinchinensis, Lour. l. c.—Cochinch.

AIGIROS, Rafin. Alsog. Am. 42 (1838) = **Populus**, Tourn. (Salic.).
heterophylos, Rafin. l. c. = P. heterophylla.

AIKINIA, R. Br. in Wall. Pl. As. Rar. iii. 65. t. 288 (1832) = **Epithema**, Blume (Gesnerac.).
Brunonis, Wall. Pl. As. Rar. iii. 66. t. 288 = E. Brunonis.
carnosa, G. Don, Gen. Syst. iv. 665 = E. carnosum.
Horsfieldii, R. Br. in Wall. Pl. As. Rar. iii. 66 = E. difforme.

AIKINIA, Salisb. ex A. DC. Monog. Camp. 135, 141 (1830) = **Wahlenbergia**, Schrad. (Campan.).
hederacea, Salisb. ex A. DC. l. c. = W. hederacea.

AIKINIA, Wall. Pl. As. Rar. iii. t. 273. p. 46 (1832) = **Ratzeburgia**, Kunth (Gramin.).
elegans, Wall. l. c. = R. pulcherrima.

AILANTHUS, Desf. in Mém. Acad. Sc. Par. 1786. (1789) 265. t. 8. *SIMARUBEAE*, Benth. & Hook. f. i. 309.
excelsa, *Roxb. Pl. Corom.* i. t. 23.—Ind. or.
flavescens, Carr. in Rev. Hortic. (1865) 366 = Cedrela sinensis.
glandulosa, *Desf. in Mém. Acad. Sc. Par.* 1786. (1789) 265. t. 8.—China.
gracilis, Salisb. Prod. 171 = Brucea sumatrana.

AILANTHUS:—
imberbiflora, *F. Muell. Fragm.* iii. 42; xix. 44.—Austral.
integrifolia, Lam. ex Steud. Nom. ed. II. i. 43 = moluccana.
japonica, Hort. ex C. Koch, Dendrol. 569 = glandulosa.
malabarica, *DC. Prod.* ii. 89.—Malabar.
moluccana, *DC. l. c.*—Ins. Molucc.
Pongelion, J. F. Gmel. Syst. ii. 726 = glandulosa.
procera, Salisb. Prod. 171 = glandulosa.
punctata, F. Muell. Fragm. iii. 42 = Pentaceras australis.
rhodoptera, F. Muell. l. c. 43 = glandulosa.

AILANTUS, DC. Prod. ii. 88 (1825) = praec.

AILUROSCHIA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 151 = **Astragalus**, Tourn. (Legum.).
amygdaliger, Stev. l. c. = Astrag. amygdaliger.
diffusa, Stev. l. c. = Astrag. dolichophyllus.
galactites, Stev. l. c. = Astrag. galactites.
glomeratus, Stev. l. c. = Astrag. glomeratus.
hypogaeus, Stev. l. c. = Astrag. hypogaeus.
rupifraga, Stev. l. c. 152 = Astrag. rupifragus.
scabriseta, Stev. l. c. 151 = Astrag. scabrisetus.

AILLYA, De Vriese, in Nat. Verh. Haarl. x. (1854) 75 = **Goodenia**, Sm.
umbellata, De Vriese, l. c. = G. filiformis.

AIMORRA, Rafin. New Fl. Am. iv. 70 (1836). (Compos.).
acuminata, Rafin. l. c.—Am. bor.

AINSLIAEA, DC. Prod. vii. 13 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 493.
DIASPANANTHUS, Miq. in Ann. Mus. Lugd. Bat. ii. 186 (1865).
acerifolia, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 126.—Malaya.
affinis, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 187 = acerifolia.
angustifolia, *Hook. f. & Thoms. ex Clarke, in Journ. Linn. Soc.* xiv. (1875) 412.—Ind. or.
apiculata, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 126.—Japon.
aptera, *DC. Prod.* vii. 14.—Reg. Himal.
asperrima, *Sch. Bip. in Pollichia*, xx–xxi. (1863) 407.—Nepal.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 318.—Burma.
cordifolia, *Franch. & Sav. Enum. Pl. Jap.* ii. 416.—Japon.
dissecta, *Franch. & Sav. l. c.* i. 264.—Japon.
fragrans, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 236.—Hongkong.
glumacea, Sch. Bip. in Pollichia, xx–xxi. (1863) 407; Klatt, in Sitzb. Muench. Akad. (1878) 97 = angustifolia.
latifolia, *Sch. Bip. in Pollichia*, xviii–xix. (1861) 190.—Nepal.
pteropoda, *DC. Prod.* vii. 14.—Reg. Himal.
silhetensis, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 411 = pteropoda.
uniflora, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 126.—Japon.
Walkeri, *Hook. f. Bot. Mag.* t. 6225.—Hongkong.

AINSWORTHIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 343 = **Tordylium**, Linn. (Umbell.).
Carmeli, Boiss. Diagn. Ser. I. x. 44.—Palestin.
cordata, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 343 = T. cordatum.
elegans, Boiss. & Bal. Diagn. Ser. II. v. 100.—Taurus.
trachycarpa, Boiss. l. c. Ser. I. x. 44.—Oriens.

AIOLOTHECA, DC. Prod. v. 508 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 352.
parthenioides, *DC. l. c.*—Mexic.

AIOUEA, Aubl. Pl. Gui. i. 310. t. 120 (1775). *LAURINEAE*, Benth. & Hook. f. iii. 153.
APIVEA, Steud. Nom. ed. I. 25 (1821).
COLOMANDRA, Neck. Elem. ii. 142 (1790).
DOUGLASSIA, Schreb. Gen. Pl. 809 (1791).
EHRHARDIA, Scop. Introd. 107 (1777).
brasiliensis, *Meissn. in DC. Prod.* xv. i. 83.—Bras.

AIOUEA:—
densiflora, *Nees, in Linnaea*, xxi. (1848) 513.—Am. trop.
elliptica, *Meissn. in DC. Prod.* xv. 1. 83.—Am. trop.
guianensis, Nees, Syst. Laurin. 363.—(Cf. Meissn. in DC. Prod. xv. 1. 82) = seq.?
guianensis, *Aubl. Pl. Gui.* i. 310.—Guiana.
guyanensis, Griseb. ex Meissn. in DC. Prod. xv. 1. 83 = Aydendron riparium.
Malabonga, *Blanco, Fl. Filip.* ed. I. 233.—Ins. Philipp.
saligna, *Meissn. in DC. Prod.* xv. 1. 82.—Bras.
Schomburgkii, *Meissn. l. c.*—Am. trop.
tenella, Nees, in Linnaea, xxi. (1848) 512 = guianensis.
tenuiflora, Nees, ex Meissn. in DC. Prod. xv. 1. 510 (= *Mespilodaphne indecora*).
trinervis, *Meissn. l. c.* 83.—Am. trop.

AIPHANES, Willd. in Mém. Acad. Berl. (1804) 32 = **Martinezia**, Ruiz & Pav. (Palm.).
aculeata, Willd. l. c. = M. Aiphanes.
caryotaefolia, H. Wendl. in Kerch. Palm. 230 = M. caryotaefolia.
corallina, H. Wendl. l. c. = M. corallina.
elegans, H. Wendl. l. c. = M. elegans.
erinacea, H. Wendl. l. c.—N. Granat.
Lindeniana, H. Wendl. l. c. = M. Lindeniana.
Praga, H. B. & K. Nov. Gen. et Sp. i. 303 = Euterpe Praga.
truncata, H. Wendl. in Kerch. Palm. 230 = M. truncata.

AIPYANTHUS, Stev. in Bull. Soc. Nat. Mosc. (1851) 1. 599 = **Macrotomia**, DC. (Borag.).
echioides, Stev. l. c. 600 = M. echioides.

AIRA, Linn. Fl. Lapp. 27 (1737); Gen. ed. I. 335. n. 838 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1156.
ANTINORIA, Parl. Fl. Palerm. i. 92 (1845).
ASPRIS, Adans. Fam. ii. 496 (1763).
FIORINIA, Parl. Fl. Ital. i. 232 (1848).
FUSSIA, Schur, Enum. Pl. Transs. 754 (1866).
LEPTOPHOBA, Ehrh. Beitr. iv. 146 (1789).
MOLINERIA, Parl. Fl. Ital. i. 236 (1848).
PERIBALLIA, Trin. Fund. Agrost. 133 (1848).
PROINEIA, Ehrh. Beitr. iv. 149 (1789).
acutigluma, *Steud. Syn. Pl. Gram.* 221.—Chili.
aegilopsoides, *Walt. Fl. Carol.* 78.—Carolina.
aetnensis, Guss. Fl. Sic. Syn. i. 151 = Corynephorus articulatus.
aggregata, Timer. ex Bor. Fl. Centr. Fr. ed. III. ii. 702 = caryophyllea.
agrostidea, Guss. Fl. Sic. Prod. i. 61 = insularis.
agrostidea, *Loisel. Notic.* 16.—Europ. austr.
alba, Vahl, in Roem. & Schult. Syst. ii. 688 = Deschampsia alba.
alba, Wulf. ex Kunth, Gram. 110 = Scsleria elongata.
alpina, Liljebl. Utkast Sv. Fl. ed. I. 49; ed. II. 41 = Hierochloë alpina.
alpina, Linn. Sp. Pl. 65 = Deschampsia alpina.
alpina, Roth, Tent. Fl. Germ. ii. 1. 98 = Deschampsia caespitosa.
alpina, Savi, Bot. Etrusc. i. 52 = Deschampsia juncea.
alpina, Vahl, in Fl. Dan. t. 961 = atropurpurea.
alpina, Sobol. ex Rupr. in Beitr. Pfl. Russ. Reich. iv. 37 = Deschampsia flexuosa.
altaica, Trin. in Bunge, Verz. Suppl. Fl. Alt. 8 = Poa soongarica.
altissima, Lam. Fl. Fr. iii. 581 = Deschampsia caespitosa.
ambigua, Michx. Fl. Bor. Am. i. 61 = Deschampsia ambigua.
ambigua, Dc Not. in Ann. Sc. Nat. Sér. III. v. (1846) 365 = capillaris.
Andraei, Auersw. ex Andrae, in Bot. Zeitschr. xiv. (1856) 205 = Deschampsia caespitosa.
anomala, *Trin. in Linnaea*, x. (1836) 301.—Chili.
antarctica, Forst. f. Prod. 8 = Trisetum antarcticum.
antarctica, *Hook. Ic. Pl.* t. 150.—Ins. Antaret.
aquatica, Guss. Fl. Sic. Prod. i. 61 = Catabrosa ochroleuca.
aquatica, Linn. Sp. Pl. 64 = Catabrosa aquatica.
aquatica, Huds. Fl. Angl. ed. II. 34, pro parte = Glyceria distans.
arctica, Spreng. Syst. Cur. Post. 32 = Deschampsia caespitosa.

AIRA:—
arenaria, *F. W. Schmidt, in Neu. Abh. Boehm. Ges.* i. (1791) 11.—Europ.
argentea, *Lowe, in Trans. Cambr. Phil. Soc.* iv. (1831) 9.—Ins. Mader.
aristulata, *Torr. Fl. U. St.* 132.—Am. bor.
articulata, Desf. Fl. Atlant. i. 70 = Corynephorus articulatus.
arundinacea, Liljebl. Utkast Sv. Fl. ed. I. 49 = Graphephorum arundinaceum.
arundinacea, Linn. Sp. Pl. 64 = Poa tatarica.
atropurpurea, *Wahlenb. Fl. Lapp.* 37.—Europ.; Am. bor.
atrovirens, Thuil. Fl. Par. ed. II. i. 37 = Molinia caerulea.
aurea, Steud. in Flora, xii. (1829) 470 = Airopsis aurea.
australis, Raoul, Choix, 12 = Deschampsia caespitosa.
australis, Steud. Syn. Pl. Gram. 220 = Deschampsia australis.
avenacea, Spreng. Syst. i. 278 = Eriachne avenacea.
avenacea, Willd. ex Steud. Nom. ed. II. i. 44 = Trisetum palustre.
baetica, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 67 = Deschampsia baetica.
benghalensis, J. F. Gmel. Syst. i. 174 = Arundo bengalensis.
Berteroniana, *Steud. Syn. Pl. Gram.* 220.—Chili.
Biebersteinii, Steud. l. c. 219 = Deschampsia caespitosa.
bottnica, Wahlenb. Fl. Lapp. 36. t. 4 = Deschampsia caespitosa.
brasiliensis, Raddi, Agrost. Bras. 36 = Eragrostis airoides.
brasiliensis, Spreng. Syst. i. 278 = Arundinella brasiliensis.
breviaristata, Gilib. Exercit. ii. 528 = Deschampsia caespitosa.
breviculmis, Loisel. Fl. Gall. i. 57 = Corynephorus canescens.
brevifolia, Bieb. Fl. Taur. Caue. iii. 63 = Deschampsia caespitosa.
brevifolia, *Pursh, Fl. Am. Sept.* i. 76.—Missouri.
brigantiaca, Chaix, in Vill. Hist. Pl. Dauph. ii. 378 = Glyceria distans.
bulbosa, Krock. Fl. Sil. Suppl. i. 131 = Koeleria eristata?
caerulea, Linn. Sp. Pl. 63 = Molinia caerulea.
caespitosa, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 380, pro parte = Festuea fuegiana.
caespitosa, Linn. Sp. Pl. 64 = Deschampsia caespitosa.
caespitosa, Muhl. Desc. Gram. 85 = Deschampsia ambigua.
calycina, *Steud. Syn. Pl. Gram.* 220.—Peruv.
canescens, All. Fl. Pedem. ii. 244 = Corynephorus articulatus.
canescens, Host, Gram. iv. t. 36 = caryophyllea.
canescens, Linn. Sp. Pl. 65 = Corynephorus canescens.
capensis, Linn. f. Suppl. 108 = Ehrharta calycina.
capensis, Steud. in Flora, xii. (1829) 468 = Koeleria capensis.
capensis, Spreng. Syst. ii. 114 = Ehrharta villosa.
capillacea, Hort. Madr. ex Wall. Cat. sub n. 3804 = Leptochloa filiformis.
capillacea, *Lam. Illustr.* i. 177.—Am. bor.
capillaris, Guss. Fl. Sic. Prod. i. 64 = Avena corymbosa.
capillaris, *Host, Gram.* iv. 20. t. 35.—Europ.; Oriens.
capillaris, Lag. in Varied. Ciene. ii. iv. (1805) 39 = Lagascae, Airopsis agrostidea.
capillaris, Salzm. ex Steud. Nom. ed. II. i. 44 = caryophyllea.
capillaris, Savi, Fl. Pis. i. 86 = Deschampsia juncea.
caryophyllaea, Leers, Fl. Herborn. ed. II. 24 = Deschampsia flexuosa.
caryophyllea, Bertol. [ex Steud.; an Fl. Ital. i. 455] = capillaris.
caryophyllea, *Linn. Sp. Pl.* 66.—Europ.; Oriens.
catabrosoides, *Steud. Syn. Pl. Gram.* 223.—Caucas.
cespitosa, Muhl. Desc. Gram. 85 = Deschampsia ambigua.
Chamaemespilus, *Host, Fl. Austr.* ii. 8.—Europ.
chinensis, *Retz. Obs.* iii. 10.—China.
ciliaris, *Kit. in Linnaea*, xxxii. (1863) 310.—Austria.
ciliaris, Auct. ex Steud. Nom. ed. I. 23, 487 = Koeleria villosa.
ciliata, Spreng. Pugill. i. 7 = Eragrostis minutiflora.
ciliata, Spreng. Syst. i. 278 = Eriachne ciliata.

AIRA:—
clavata, Gilib. Exercit. ii. 526 = Corynephorus canescens.
clavata, Willd. ex Steud. Nom. ed. II. i. 44 = Corynephorus articulatus.
collina, Dum. Obs. Gram. Belg. 121 = Deschampsia flexuosa.
compressa, *Rafin. in Am. Monthly Mag.* (1811) 356, nomen.—Am. bor.
controversa, *Steud. Syn. Pl. Gram.* 224.—Am. bor.
complanata, Steud. Nom. ed. II. i. 44 = Chloris petraea.
corsica, Jord. Pugill. 353 = sicula.
corsica, Tausch, in Flora, xx. (1837) 102 = Deschampsia flexuosa.
corymbosa, Willd. ex Steud. Nom. ed. II. i. 44 = caryophyllea.
corymbosa, Fauché & Chaub. Fl. Pelop. 5. n. 120. t. 7. f. 1 = intermedia.
cristata, Linn. Sp. Pl. 63 = Koeleria cristata.
Cupaniana, Guss. Fl. Sic. Syn. i. 145 = caryophyllea.
cuprina, Fuss, Fl. Transs. 722 = Deschampsia flexuosa.
curta, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 701 in nota = caryophyllea.
dactyloides, Rochel, Pl. Rar. Ban. 32. f. 3 = Koeleria glauca.
danthonioides, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1830) 57 = Deschampsia calycina.
discolor, Thuil. Fl. Par. i. 39 = setacea.
distichophylla, Spreng. Syst. i. 276 = Panicum subulatum.
divaricata, Pourr. in Mém. Acad. Toul. iii. (1788) 307 = caryophyllea.
duriuscula, *Poir. Encyc. Suppl.* ii. 76.—Hab.?
Edouardi, Reut. ex Jouve, in Bull. Soc. Bot. Fr. xii. (1865) 88 = Avena corymbosa.
effusa, Spreng. Syst. i. 278 = Eriachne pallescens.
elatior, 'Hoffmgg.' ex Steud. Nom. ed. II. i. 44 = Melica Bauhini?
elatior, *Steud. Syn. Pl. Gram.* 423.—Reg. Magellan.
elegans, Willd. ex Gaud. Agrost. Helv. i. 130, 355 = capillaris.
elegans, 'Vill.' ex Steud. Nom. ed. II. i. 44 = Airopsis agrostidea.
elegantissima, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 85 = capillaris.
elodes, Brign. Fase. Pl. Foroj. 10; cf. Flora (1840), 104 = Poa nemoralis.
elongata, Hook. Fl. Bor. Am. ii. 243 = Deschampsia elongata.
elongata, Salzm. ex Steud. Nom. ed. II. i. 44 = Koeleria phleoides.
euflexuosa, Syme, Engl. Bot. ed. III. xi. 67 = Deschampsia flexuosa.
fascicularis, Steud. Syn. Pl. Gram. 219 = Corynephorus fasciculatus.
festucoides, Vill. Prosp. 16; Hist. Pl. Dauph. ii. 86 = Deschampsia alpina.
filiformis, *Koen. ex Roxb. Fl. Ind.* i. 328.—Ind. or.
flammosa, Buchoz, ex Honck. Syn. Pl. Germ. i. 521 = Deschampsia flexuosa.
flexilis, Linn. ex Gieseke, Prael. 147 = Deschampsia flexuosa.
flexuosa, Linn. Sp. Pl. 65 = Deschampsia flexuosa.
Gayana, *Steud. Syn. Pl. Gram.* 220.—Chili.
gigantea, Schrenk, ex Ledeb. Fl. Ross. iv. 446 = Muehlenbergia alpestris.
gigantea, *Steud. Syn. Pl. Gram.* 224.—Ind. occ.?
glabra, *Brot. Fl. Lusit.* 91.—Lusit.
glabrata, Brot. l. c. 92 = lendigera.
glauca, Hartm. ex Steud. Nom. ed. II. i. 44 = Deschampsia caespitosa.
glauca, Rochel, Pl. Rar. Banat. 32 = Koeleria dactyloides.
glauca, Spreng. Nachtr. Bot. Gart. Halle, i. 10 = Daetylis glauca.
globosa, Thore, in Journ. Bot. Paris, i. (1808) 197 = Airopsis globosa.
glomerata, G. Don, ex S. F. Gray, Nat. Arr. Brit. Pl. ii. 137 = Deschampsia caespitosa.
Gmelini, [Honck.] Verz. aller Gew. Teutschl. 212 = praecox?
gracilis, Trin. ex Steud. Nom. ed. II. i. 44 = Koeleria eristata.
grandiflora, Bertol. Fl. Ital. i. 436 = Koeleria setacea.
grandiflora, Steud. Syn. Pl. Gram. 220 = Deschampsia grandiflora.
grandis, Nym. Consp. 807 = Deschampsia caespitosa.

AIRA :—
Halleri, [Honck.] Verz. aller Gew. Teutschl. 212 = Trisetum rigidum.
Hartmaniana, Nym. Consp. 807 = Deschampsia caespitosa.
helodes, Steud. Nom. ed. II. i. 44 = Poa nemoralis.
hirsuta, Hall. f. ex Schleich. Cat. Nov. 5 = Koeleria hirsuta.
hispida, Spreng. Syst. i. 276 = Eriachne capillaris.
holciformis, Steud. Syn. Pl. Gram. 221 = Deschampsia holciformis.
Holcus-lanatus, Vill. Hist. Pl. Dauph. ii. 87 = Holcus lanatus.
Holcus-mollis, Vill. l. c. 88 = Holcus mollis.
Hostii, *Steud. Syn. Pl. Gram.* 221.—Bohem.
humilis, Bieb. Fl. Taur. Cauc. i. 57 = Catabrosa humilis.
hybrida, Gaud. Agrost. Helv. i. 124 = Corynephorus articulatus.
incompleta, Bosc, ex Steud. Nom. ed. II. i. 44 = Paspalum debile.
indica, Linn. Sp. Pl. ed. II. 94 = Panicum conglomeratum.
inflexa, Loisel. Fl. Gall. i. 56. t. 22 = Avena corymbosa.
insularis, *Boiss. Fl. Orient.* v. 528.—Europ.; Oriens.
intermedia, *Guss. Fl. Sic. Prod. Suppl.* i. 105.—Europ.
interrupta, Rottl. ex Steud. Syn. Pl. Gram. 55 = Panicum canaliculatum.
involucrata, Cav. Ic. i. 33. t. 44 = Deschampsia involucrata.
ischaemoides, Koen. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. (1834) 328 = Panicum antipodum.
juncea, Vill. Prosp. 16 = Deschampsia juncea.
juncifolia, Wulf. ex Kunth, Enum. i. 322 = Sesleria tenuifolia.
Kingii, Hook. f. Fl. Antarct. ii. 376. t. 136 = Deschampsia caespitosa.
koelerioides, Peyr. in Linnaea, xxx. (1859-60) 5 = Deschampsia koelerioides.
Koenigii, J. F. Gmel. Syst. i. 174 = Eragrostis minutiflora.
labradorica, *Steud. Syn. Pl. Gram.* 220.—Labrador.
laevigata, Sm. Eng. Bot. t. 2102 = Deschampsia alpina.
laevis, Brot. Fl. Lusit. i. 90 = Deschampsia laevis.
laevis, Spreng. Syst. i. 278 = Eriachne glauca.
Lagascae, *Kunth, Gram.* i. 99.—Hispan.; Corsic.
latifolia, Hook. Fl. Bor. Am. ii. 243 = Deschampsia latifolia.
latifolia, Steud. Syn. Pl. Gram. 219 = Deschampsia caespitosa.
latigluma, *Steud. l. c.* 221.—Abyss.
laxa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Orthoclada laxa.
Leersii, Kunth, Enum. Pl. i. 289 = caryophyllea.
Legei, Bor. in Soc. Angers, xxiv. (1855); et Fl. Centr. Fr. ed. III. ii. 700 = Deschampsia flexuosa.
lendigera, *Lag. in Varied. Cienc.* ii. IV. (1805) 39.—Hispan.
Lensaei, Loisel. Fl. Gall. ed. II. i. 58 = caryophyllea.
Lensii, Reichb. Fl. Germ. Excurs. 140[5] = praec.
Lenzei, Gren. & Godr. Fl. Fr. iii. 620 = praec.
litoralis, Godr. ex Nym. Consp. 807 = Deschampsia caespitosa.
macrantha, Ledeb. in Mém. Acad. Pétersb. v. (1812) 515 = Koeleria cristata.
macranthera, *Steud. Syn. Pl. Gram.* 219.—Hispan.
maderaspatana, *Nees, ex Steud. l. c.* 221.—Madras.
magellanica, *Hook. f. Fl. Antarct.* ii. 376. t. 134.—Reg. Magellan.
magellanica, Lam. ex Beauv. Agrost. 65 = Hierochloë magellanica.
major, Syme, Engl. Bot. ed. III. xi. 63.—Deschampsia caespitosa, alpina.
media, Gouan, Illustr. 3 = Deschampsia juncea.
media, Trin. in Mém. Acad. Pétersb. Sér. VI. iv. (1831) Suppl. 57 = Deschampsia caespitosa.
melicoides, Michx. Fl. Bor. Am. i. 62 = Graphephorum melicoides.
micrantha, *Steud. Syn. Pl. Gram.* 423.—Madag.
miliacea, Vill. Hist. Pl. Dauph. ii. 81 = Poa miliacea.
minuta, *Loefl. It. Hisp.* 117.—Europ.; Oriens.
minuta, Loisel. Fl. Gall. i. 45 = Airopsis agrostidea.
missurica, Spreng. Syst. ii. 578 = brevifolia.
mollis, Muhl. Descr. Gram. 82 = Eatonia purpurascens.
mollis, Schreb. Spicil. 51 = Holcus mollis.
montana, Brot. ex Nym. Consp. 808 = Deschampsia stricta.

AIRA :—
montana, Desv. Obs. Angers, 55 = setacea.
montana, Koel. Gram. 131 = Deschampsia flexuosa.
montana, *Linn. Sp. Pl.* 65.—Europ.
mucronata, Spreng. Syst. i. 276 = Eriachne brevifolia, mucronata.
multicaulis, Kunth, Enum. Pl. i. 292 = caryophyllea.
multiculmis, Dum. Obs. Gram. Belg. 121 = caryophyllea.
nana, Kunze, ex Steud. Nom. ed. II. i. 45 = praecox.
nana, Spreng. ex Steud. l. c. = caryophyllea.
nitida, Spreng. Mant. i. 32 = Eatonia purpurascens.
nitida, *Steud. Syn. Pl. Gram.* 220.—Mexic.
nivea, *Host, Fl. Austr.* ii. 8.—Europ.
Notarisiana, Steud. Syn. Pl. Gram. 221 = capillaris.
nutans, [Web. in] Wigg. Prim. Fl. Holsat. 7 = Melica nutans.
obtusata, Michx. Fl. Bor. Am. i. 62 = Eatonia obtusata.
orzyetorum, Spreng. Pugill. ii. 18 = Festuca arundinacea.
pallens, Muhl. Descr. Gram. 84 = Trisetum palustre.
pallens, *Spreng. Fl. Hal. Mant.* 36.—Hab. ?
pallescens, *Kit. in Roem. & Schult. Syst.* ii. 685.—Hungar.
paludosa, Roth, Tent. Fl. Germ. ii. 1. 96 = Deschampsia flexuosa.
paludosa, Wibel, Prim. Fl. Werth. 118 = Deschampsia caespitosa.
palustris, Pollini, Cat. Ort. Veron. (1814) 8 = Festuca arundinacea.
panicea, Willd. ex Steud. Nom. ed. II. i. 45 = Leptochloa filiformis.
paradoxa, *Steud. Syn. Pl. Gram.* 223.—Guinea.
parviflora, Thuil. Fl. Par. ed. II. i. 38 = Deschampsia caespitosa.
parvula, *Hook. f. Fl. Antarct.* ii. 377.—Fuegia.
patulipes, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 701 = caryophyllea.
pensylvanica, Spreng. in Mém. Acad. Pétersb. (1810) 299. t. 7 = Eatonia purpurascens.
petraea, Bellardi, ex Steud. Nom. ed. II. i. 45; ii. 361 = Poa nemoralis.
phleoides, Steud. Nom. ed. I. 24 = Koeleria phleoides.
pictigluma, *Steud. Syn. Pl. Gram.* 221.—Abyss.
pilosa, *Kit. in Linnaea*, xxxii. (1863) 310.—Austria.
plesiantha, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 701 = caryophyllea.
poaeformis, Willd. ex Steud. Nom. ed. II. i. 45 = Poa depauperata.
praecox, *Linn. Sp. Pl.* 65.—Europ.; Am. bor.
provincialis, Gennar. ex Nym. Consp. 814 = Avena corymbosa.
provincialis, Jord. ex Jouve, in Bull. Soc. Bot. Fr. xii. (1865) 84. t. 1 = capillaris.
pubescens, Vahl, Symb. iii. 9 = Koeleria villosa.
pulchella, Nocca & Balb. Fl. Ticin. i. 403. t. add. 1 = capillaris.
pulchella, Link, Hort. Berol. i. 130 = Avena corymbosa.
pulchella, Willd. Enum. Hort. Berol. 101 = lendigera.
pulchra, *Steud. Syn. Pl. Gram.* 220.—Chili.
pumila, Agardh, ex Roem. & Schult. Syst. ii. 578 = minuta.
pumila, Pursh, Fl. Am. Sept. i. 76 = Poa annua.
pumila, Vill. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 572 = Deschampsia juncea.
purpurea, Muhl. Descr. Gram. 85 = Uralepsis aristulata.
purpurea, Walt. Fl. Carol. 78 = Triplasis purpurea.
purpureo-argentea, Gilib. Exercit. ii. 526 = caryophyllea.
pusilla, Schur, Enum. Pl. Transs. 754 = praecox.
rara, Spreng. Syst. i. 278 = Eriachne rara.
refracta, Lag. in Varied. Cienc. ii. IV. (1805) 39 = Deschampsia refracta.
rigida, Dulac, Fl. Hautes-Pyr. 82 = caryophyllea.
Ruprechtii, Griseb. ex Hook. f. in Trans. Linn. Soc. xxiii. (1861) 346 = Avena Ruprechtii.
sabuletorum, Steud. Nom. ed. II. i. 45 = sabulonum.
sabulonum, *Labill. Sert. Austr. Caled.* 16. t. 21.—Nov. Caled.
saburralis, Jan, Elench. 2, ex Guss. Fl. Sic. Syn. i. 148 = caryophyllea.
saxatilis, Salzm. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 59 = capillaris.
scabra, *Kit. in Linnaea*, xxxii. (1863) 309.—Austr.
scabra, Hort. ex Kunth, Enum. Pl. i. 370 = Catabrosa viridula.

AIRA :—
scabra, Zuccagn. in Roem. Collect. 124 = Catabrosa scabra.
scabrosetacea, Knapp, Gram. Brit. t. 32 = Deschampsia flexuosa.
semineutra, Waldst. & Kit. in Schult. Oestr. Fl. i. 199 = Poa semineutra.
serotina, *Torr. sec. Trin. ex Steud. Nom.* ed. II. i. 45.—Am. bor.
setacea, *Huds. Fl. Angl.* ed. I. 30; *Retz. Obs.* iii. 10.—Europ.
setacea, Pourr. ex Nym. Consp. 808 = Deschampsia juncea.
setiformis, *Host, sec. Trin. ex Steud. Nom.* ed. II. i. 45.—Hab. ?
sicula, Spreng. Syst. i. 277 = Holcus tenuis.
sicula, *Tod. Ind. Sem. Pan.* (1872) 38; *in Nuov. Giorn. Bot.* v. (1873) 156.—Sicilia.
spectabilis, *Steud. Syn. Pl. Gram.* 224.—Afr. austr.
spicaeformis, *Steud. l. c.* 424.—Reg. Magell.
spicata, Linn. Sp. Pl. 63 = Trisetum subspicatum.
spicata, Linn. l. c. 64 = Panicum indicum.
splendida, Willd. ex Steud. Nom. ed. II. i. 45 = Deschampsia flexuosa.
squarrosa, Spreng. Syst. i. 278 = Eriachne squarrosa.
stolonifera, Hausm. Fl. Tir. 890, sec. Neilr. Nachtr. 25 = Deschampsia caespitosa.
stricta, Nym. Consp. 808 = Deschampsia stricta.
subaristata, Faye, ex Anderss. in Walp. Ann. vi. 992 = Deschampsia juncea.
subspicata, Linn. Syst. ed. X. 873 = Trisetum subspicatum.
subspicata, Rainv. in Gort. Fl. Belg. Foed. n. 74 = praecox.
subtriflora, Lag. in Varied. Cienc. ii. IV. 39 = Deschampsia juncea.
superbiens, *Steud. Syn. Pl. Gram.* 424.—Reg. Magell.
subulorum, D. Dietr. Syn. i. 339 err. = sabulonum.
tenera, *Kit. in Schult. Oestr. Fl.* ed. II. i. 199.—Slavonia; Banatus.
Tenorei, Fuckel, ex Willk & Lange, Prod. i. 55 = Agrostis capillaris.
Tenorii, Guss. Fl. Sic. Prod. i. 62 = Avena corymbosa.
tenuifolia, *Steud. Syn. Pl. Gram.* 223.—Chili.
tenuissima, Spreng. Syst. i. 276 = Sporobolus minutiflorus.
Todari, Tenore, ex Tod. Ind. Sem. Hort. Panorm. (1859) (Linnaea, xxx. (1859-60) 758) = caryophyllea.
triaristata, *Clarke, Trav.* ii. 452.—Palaestina.
trichodes, Spreng. Syst. i. 276 = Agrostis trichodes.
triflora, Frivald. in Flora, xix. (1836) 433 = Deschampsia caespitosa.
triflora, Ell. Sketch, i. 153 = Graphephorum Elliotti.
triflora, Roem. & Schult. Syst. ii. 680 = Deschampsia juncea.
triflora, Willd. ex Steud. Nom. ed. II. i. 45 = Corynephorus canescens.
triglumis, *Steud. Syn. Pl. Gram.* 223.—Texas.
truncata, Muhl. Descr. Gram. 83 = Eatonia obtusata.
truncata, Salzm. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 58 = caryophyllea.
uliginosa, Weihe, ex Boenn. Prod. Fl. Monast. 25 = setacea.
uniaristata, *Cav. in Anales Cienc. Nat.* vi. (1803) 148.—Hab. ?
valesiaca, Colla, Herb. Pedem. t. 94. f. 1 = Koeleria setacea.
valesiana, Bertol. Fl. Ital. i. 438 = praec.
varia, Jacq. Hort. Vindob. 15 = Sesleria caerulea.
variegata, *Steud. Syn. Pl. Gram.* 223. – Graecia.
versicolor, *Roem. & Schult. Syst.* ii. 679.—'Port. Egmont.'
vestita, *Steud. Syn. Pl. Gram.* 424.—Reg. Magell.
viatica, *Griff. Notul.* iii. 54.—Reg. Himal.
villosa, Gilib. Exercit. ii. 527 = montana.
villosa, Linn. f. Suppl. 109 = Ehrharta gigantea.
violacea, Gilib. Exercit. ii. 525 = Catabrosa aquatica.
violacea, *Steud. Syn. Pl. Gram.* 223.—Persia.
violacea, Willd. ex Steud. Nom. ed. II. i. 105 = Panicum antipodum.
vivipara, *Steud. Syn. Pl. Gram.* 222.—Groenland.
viridula, *Poir. Encyc. Suppl.* ii. 75.—Hab. ?
Wibeliana, *Sond. in Fl. Dan.* t. 2461, 2522.—Europ.
Wilhelmsii, Steud. Syn. Pl. Gram. 219 = Deschampsia caespitosa.

AIRIDIUM, Steud. Syn. Pl. Gram. 423 (1855) = **Deschampsia**, Beauv. (Gram.).
elegantulum, Steud. l. c. = D. antarctica.

AIROCHLOA, Link, Hort. Berol. i. 126 ; ii. 276 (1827) = **Koeleria**, Pers. (Gram.).
albescens, Link, l. c. i. 128 = K. albescens.
capensis, Nees, in Mart. Fl. Bras. ii. 321 = K. capensis.
caudata, Link, in Linnaea, xvii. (1843) 405 = K. crassipes.
cristata, Link, Hort. Berol. i. 127 = K. cristata.
glauca, Link, ex Schur, Enum. Pl. Transs. 751 = K. glauca.
gracilis, Link, Hort. Berol. ii. 276 = K. cristata.
hirsuta, Link, l. c. i. 129 = K. hirsuta.
hispida, Link, l. c. = K. hispida.
permollis, Nees, ex Steud. Syn. Pl. Gram. 293 = K. permollis.
pubescens, Link, Hort. Berol. i. 128 = K. cristata.
uniflora, Hochst. in Flora, xxxviii. (1855) 330.—Afr. or.
valesiaca, Link, Hort. Berol. i. 127 = K. setacea.
villosa, Link, l. c. 128 = K. villosa.

AIROPSIS, Desv. Journ. Bot. i. (1808) 200, pro parte. *GRAMINEAE*, Benth. & Hook. f. iii. 1156.
agrostidea, *DC. Fl. Fr. Suppl.* 262.—Europ. austr.
ampla, *Nees, ex Steud. Nom.* ed. II. i. 45.—Afr. austr.
aurea, *Nees, in Linnaea*, vii. (1832) 317.—Afr. austr.
brevifolia, Roem. & Schult. Syst. ii. 578 = Aira brevifolia.
Candolii, Desv. Journ. Bot. i. (1808) 200 = agrostidea.
capillaris, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 328 = Aira capillaris.
caryophyllea, Schur, l. c. = Aira caryophyllea.
Cupaniana, Lange, in Kjoeb. Vidensk. Meddel. (1860) 38 = caryophyllea.
globosa, *Desv. Journ. Bot.* i. (1808) 200.—Gall. ; Hisp.
glabra, *Desv. Opusc.* 64.—Ins. Borbon.
insularis, Nym. Syll. 411 = Aira insularis.
involucrata, Desv. Journ. Bot. i. (1808) 201 = Deschampsia involucrata.
jubata, *Griseb. in Goett. Abh.* xxiv. (1879) 292.—Reg. Argent.
lendigera, Lange, in Kjoeb. Vidensk. Meddel. (1860) 37 = Aira lendigera.
millegrana, Griseb. in Goett. Abh. xix. (1874) 252 = Eragrostis airoides.
minuta, Desv. Journ. Bot. i. (1808) 201 = Catabrosa minuta.
multiculmis, Lange, in Kjoeb. Vidensk. Meddel. (1860) 38 = Aira caryophyllea.
obtusata, Desv. Journ. Bot. i. (1808) 200 = Eatonia obtusata.
pallida, Nees, ex Steud. Nom. ed. II. i. 45 = Eriachne pallida.
peruviana, [Nees &] Meyen, Reise, i. 484 = Dissanthelium supinum.
praecox, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 328 = Aira praecox.
pulchella, Tenore, Fl. Nap. iii. 56, 102 = Avena corymbosa.
Steudelii, Nees, in Linnaea, vii. (1838) 318 = Eriachne capensis.
triseta, Nees, ex Steud. Nom. ed. II. i. 45 = Eriachne triseta.
tuberculata, Nees, ex Steud. l. c. = Eriachne tuberculata.

AITCHISONIA, Hemsl. ex Aitch. in Journ. Linn. Soc. xix. (1882) 166. t. 14. *RUBIACEAE*.
rosea, *Hemsl. l. c.*—Afghan.

AITHALES, Webb & Berth. Phyt. Canar. i. 178 (1836) = **Sedum**, Linn. (Crassul.).
caespitosa, Webb & Berth. l. c. 179, 180 = S. caespitosum.
rubens, Webb & Berth. l. c. 179 = S. rubens.

AITITARA, Endl. Gen. 254. (1837) = ATITARA, Marcgr. = **Evodia**, Forst. (Rutac.).

AITONIA, Thunb. in Phys. Sallsk. Handl. i. (1776) 166 ; Nov. Gen. 52. *SAPINDACEAE*, Benth. & Hook. f. i. 411.
AYTONIA, Linn. f. Suppl. 49. 303 (1781).
capensis, *Thunb. ll. cc.*—Afr. austr.

AITOPSIS, Rafin. Fl. Tellur. iii. 91 (1836) = **Salvia**, Tourn. (Labiat.).
foliosa, Rafin. l. c. = S. foliosa.

AIZODRABA, Fourr. in Ann. Soc. Linn. Lyon, N. S., xvi. (1868) 335 = **Draba**, Linn. (Crucif.).
aizoides, Fourr. l. c. = D. aizoides.
beugesiaca, Jord. & Fourr. l. c.—Gallia.
chrysantha, Jord. & Fourr. l. c.—Gallia.
ochroleuca, Jord. & Fourr. l. c. = D. algida.
saxigena, Fourr. l. c. = D. aizoides.
trichocarpa, Jord. & Fourr. l. c.—Gallia.

AIZOON, Linn. Gen. ed. I. 161 (1737). *FICOIDEAE*, Benth. & Hook. f. i. 854.
VESTINGIA, Hcist. Syst. (1748); Fabric. Hort Helmst. 363 (1759).
angustifolium, D. Dietr. Syn. Pl. iii. 130 = Acrosanthes angustifolia.
argenteum, Eckl. & Zeyh. Enum. 325 = rigidum.
Camforosma, *Reichb. ex Spreng. Syst. Cur. Post.* 195.—Afr. austr.
canariense, Andr. Bot. Rep t. 201 = Sesuvium pedunculatum.
canariense, *Linn. Sp. Pl.* 488.—Ins. Canar. ; Afr. bor. ; Oriens.
contaminatum, Eckl. & Zeyh. Enum. 326 = Galenia secunda.
crystallinum, Eckl. & Zeyh. l. c. 327 = Galenia crystallina.
elongatum, Eckl. & Zeyh. l. c. 326 = Galenia secunda.
fistulosum, D. Dietr. Syn. Pl. iii. 130 = Acrosanthes fistulosa.
fruticosum, Eckl. & Zeyh. Enum. 327 = Galenia affinis.
fruticosum, Linn. f. Suppl. 261 = Galenia fruticosa.
galenioides, *Fenzl, ex Harv. & Sond. Fl. Cap.* ii. 469.—Afr. austr.
glinoides, Eckl. & Zeyh. Enum. 325 = Galenia secunda.
glinoides, Linn. f. Suppl. 261.—Afr. austr.
Herniaria, Reichb. ex Steud. Nom. ed. II. i. 46 = Galenia herniariaefolia.
hirsutum, Eckl. & Zeyh. Enum. 325 = glinoides.
hispanicum, *Linn. Sp. Pl.* 488.—Hispan.
lanceolatum, *Murr. Syst. Veg.* ed. XIII. 392.—Afr austr.
microphyllum, Bartl. in Linnaea, vii. (1832) 541 = Galenia herniariaefolia.
montevidense, Sprcng. ex Rohr, in Mart. Fl. Bras. xiv. II. 310 = Sesuvium Portulacastrum.
mossamedense, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 584.—Afr. trop.
paniculatum, *Linn. Sp. Pl.* 488.—Afr. austr.
papulosum, Eckl. & Zeyh. Enum. 326 = Galenia papulosa.
perfoliatum, Linn. f. Suppl. 261 = Tetragonia Zeyheri.
procumbens, Crantz, Inst. i. 135 = canariense.
propinquum, Eckl. & Zeyh. Enum. 326 = Galenia secunda.
pubescens, Eckl. & Zeyh. l. c. = Galenia secunda.
quadrifidum, *F. Muell. Fragm.* ii. 148.—Austral.
rigidum, *Linn. f. Suppl.* 261.—Afr. austr.
sarmentosum, *Linn. f. l. c.* 260.—Afr. austr.
secundum, Linn. f. l. c. 261 = Galenia secunda.
sericeum, Fenzl, ex Harv. & Sond. Fl. Cap. ii. 470 = rigidum.
sessiliflorum, Moench, Meth. 698 = hispanicum.
spathulatum, *Eckl. & Zeyh. Enum.* 325.—Afr. austr.
squamulosum, Eckl. & Zeyh. l. c. 326 = Galenia squamulosa.
stellatum, Lam. Encyc. iii. 418 = sarmentosum.
teretifolium, D. Dietr. Syn. Pl. iii. 130 = Acrosanthes teretifolia.
tomentosum, Lam. Encyc. iii. 418 = paniculatum.
virgatum, *Welw., ex Oliver, Fl. Trop. Afr.* ii. 584.—Afr. trop.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 470.—Afr. austr.
zygophylloides, *F. Muell. Fragm.* vii. 129.—Austral.

AIZOUM, Linn. Syst. ed. I. (1755) = **Aizoon**, Linn. (Ficoid.).

AJAX, Salisb. in Trans. Hort. Soc. i. (1812) 343 = **Narcissus**, Linn. (Amaryll.).
(NOTA.—Cl. Baker *Ajaces* omnes ad *Narcissum Pseudo-Narcissum* ducit.)
abscissus, Haw. Rev. 116, 117 = N. abscissus.
albicans, Haw. Monog. 2 = N. Pseudo-Narcissus.
albus, Haw. Rev. 117 = N. moschatus.

AJAX :—
bicolor, Salisb. in Trans. Hort. Soc. i. (1812) 346 = N. Pseudo-Narcissus.
breviflos, Haw. Monog. 2. n. 6 = N. Pseudo-Narcissus.
cambricus, Haw. l. c. 3 = N. Pseudo-Narcissus.
capax, M. Roem. Syn. Ensat. 201 = N. Pseudo-Narcissus.
cernuus, Haw. Monog. 2 = N. Pseudo-Narcissus.
cuneiflorus, Salisb. in Trans. Hort. Soc. i. (1812) 343 = N. Pseudo-Narcissus.
cuneifolius, Haw. Rev. 43 = N. Pseudo-Narcissus.
cyclamineus, Haw. Monog. 2. n. 4 = N. Pseudo-Narcissus.
fenestralis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 191 = N. Pseudo-Narcissus.
festalis, Salisb. in Trans. Hort. Soc. i. (1812) 347 = N. Pseudo-Narcissus, serratus.
glaucus, M. Roem. Syn. Ensat. 199 = N. propinquus.
grandiflorus, Salisb. in Trans. Hort. Soc. i. (1812) 344 = N. major.
hexangularis, Herb. Amar. 305.—Hab.?
hispanicus, M. Roem. Syn. Ensat. 200 = N. Pseudo-Narcissus.
incomparabilis, Car. Prod. Fl. Tosc. 616 = N. incomparabilis.
lacinularis, Salisb. in Trans. Hort. Soc. i. (1812) 344 = N. propinquus.
lobularis, Haw. in Phil. Mag. (1830) 131 = N. Pseudo-Narcissus.
longiflorus, Salisb. in Trans. Hort. Soc. i. (1812) 349 = N. tortuosus.
lorifolius, Haw. Monog. 2. n. 7 = N. Pseudo-Narcissus.
lusitanicus, M. Roem. Syn. Ensat. 202 = N. triandrus.
luteus, Herb. Amar. 304, 415. t. 38. f. 33 = N. major.
major, Haw. Rev. 116 = N. major.
maximus, Haw. Monog. 3. n. 23 = N. Pseudo-Narcissus.
minimus, Haw. l. c. 1. n. 1 = N. Pseudo-Narcissus.
minor, Herb. Amar. 299, 415 = N. Pseudo-Narcissus.
moschatus, Haw. Monog. 2 = N. moschatus.
nanus, Haw. l. c. 2. n. 5 = N. Pseudo-Narcissus.
nobilis, Haw. Syn. 327 = N. Pseudo-Narcissus.
obvallaris, Haw. Rev. 120 = N. Pseudo-Narcissus.
odorus, Car. Prod. Fl. Tosc. 616 = N. odorus.
patulus, Salisb. in Trans. Hort. Soc. i. (1812) 348 = N. moschatus.
propinquus, Haw. Monog. iii. n. 22 = N. Pseudo-Narcissus.
Pseudo-narcissus, Haw. l. c. 2 = N. Pseudo-Narcissus.
pumilus, Haw. l. c. 1. n. 3 = N. Pseudo-Narcissus.
pygmaeus, M. Roem. Syn. Ensat. 201 = N. Pseudo-Narcissus.
radians, M. Roem. l. c. 193 = N. Pseudo-Narcissus.
reflexa, M. Roem. l. c. 202 = N. triandrus.
Rudbeckii, M. Roem. l. 201 = N. Pseudo-Narcissus.
rugilobus, Haw. Monog. 3 = N. Pseudo-Narcissus.
Sabiniamus, Herb. Amar. 306 = N. Pseudo-Narcissus.
serratus, Haw. Rev. 114 = N. serratus.
sexangularis, M. Roem. Syn. Ensat. 201 = N. Pseudo-Narcissus.
spurius, Haw. Syn. 327 = N. Pseudo-Narcissus.
Telamonius, Haw. l. c. 326 = N. Pseudo-Narcissus.
tortuosus, Haw. Monog. 2 = N. Pseudo-Narcissus.
tubaeflorus, Herb. Amar. 303 = N. Pseudo-Narcissus.

AJOVEA, Juss. Gen. 80 (1789) = **Aiouea**, Aubl. (Laurin.).
guianensis, Rafin. Sylva Tellur. 134 = Aiouea tenella.

AJUGA, Linn. Gen., ed. I. 167. n. 466 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1222.
ABIGA, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 85.
BUGULA, Moench, Meth. 381 (1794).
CHAMAEPITYS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 220 (1745).
MOSCHARIA, Forsk. Fl. Aegypt. Arab. 158 (1775).
PHLEBOANTHE, Tausch, in Flora, xi. (1828) 322.
acaulis, *Brocchi, in Bibliot. Ital.* xxix. 85.—Europ.
adulterina ×, *Wallr. Beitr. Fl. Hercyn.* 221.—Europ.
africana, Pers. Syn. ii. 109 = Teucrium africanum.
alpestris, Dum. Fl. Belg. 42 = genevensis.
alpina, Fries, Mant. iii. 54 = reptans.
alpina, Vill. Hist. Pl. Dauph. ii. 347 = pyramidalis.
alpina, Linn. Mant. 80 = genevensis.
argyrea, *Stapf, in Denkschr. Acad. Wien* l. (1885) 103.—Lycia.
arenosa, Stev. ex Ledeb. Fl. Ross. ii. 449 = oblongata.

AJUGA :—
astolonosa, *Schur, Enum. Pl. Transs.* 545.—Europ.
australis, *R. Br. Prod.* 503.—Austral.
Barrelieri, Tenore, Fl. Nap. t. 240. f. 2 = reptans.
bombycina, *Boiss. Fl. Orient.* iv. 803.—As. Min.
brachystemon, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1883) 192.—Reg. Himal.
bracteosa, *Benth. in Wall. Pl. As. Rar.* i. 59.—Afr. trop.; Ind. or.; As. or.
caespitosa, Schleich. ex Steud. Nom. ed. II. i. 46 = pyramidalis.
capensis, Pers. Syn. ii. 109 = Teucrium africanum.
Chamaecistus, *Ging. ex Benth. Lab. Gen. et Sp.* 698. —Persia.
Chamaepytis, Guss. Fl. Sic. Prod. ii. 66 = A. Chia.
Chamaepytis, *Schreb. Pl. Vert. Unilab.* 24.—Europ. Oriens.
Chia, *Schreb. l. c.* 25.—Oriens.
ciliata, *Bunge, Enum. Pl. Chin. Bor.* 51.—Chin. bor.
comata, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 50. —Lycia.
controversa, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, vi. (1859) 136; xiv. (1863) 44.—Gallia.
crenata, Hochst. ex Benth. in DC. Prod. xii. 597 = bracteosa.
cryptostylon, Lagrèze-Fossat, ex Nym. Consp. 567 = genevensis.
cuneatifolia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 102.—Lycia.
decumbens, Tenore, Fl. Nap. Prod. 'p. xxxiii' = acaulis.
decumbens, *Thunb. Fl. Jap.* 243.—Japon.
decumbens, D. Don, Prod. 108 = macrosperma.
densiflora, Wall. Cat. n. 2031; Benth. in Wall. Pl. As. Rar. i. 696 = bracteosa.
densiflora, Tenore, Fl. Nap. t. 239. f. 2 = reptans.
depressa, Maxim. in Bull. Acad. Pétersb. xxvii. (1883) 192 = parviflora.
diemenica, Benth. Lab. Gen. et Sp. 695 = australis.
disticha, Roxb. Fl. Ind. iii. 2 = Anisomeles malabarica.
euphrasioides, Boiss. Diagn. Ser. I. vii. 63 = A. Chamaecistus.
flaccida, *Baker, in Journ. Linn. Soc.* xx. (1884) 234.—Madag.
foliosa, Tratt. Arch. Gewaechsk. i. 25 = genevensis.
fruticosa, Roxb. Fl. Ind. iii. 1 = Anisomeles malabarica.
furcata, Link, Enum. Hort. Berol. ii. 99 = Crainotome versicolor.
genevensis, *Linn. Sp. Pl.* 561.—Europ.; Oriens.
genevensis, With. Arr. Brit. Pl. ed. III. iii. 516 = reptans.
geniculata, Maxim. in Bull. Acad. Pétersb. xxix. (1883) 189 = macrosperma.
glabra, Benth. Lab. Gen. et Sp. 700 = laevigata.
glabra, Presl, Fl. Sicul. Praef. 36 = A. Chamaepitys.
glabrata, Benth. in Wall. Cat. n. 2041 = Anisomeles ovata.
grosse-serrata, *Franch. & Sav. Enum. Pl. Jap.* ii. 467. —Japon.
Hampeana ×, *A. Br. & Vatke, in Oestr. Bot. Zeitschr.* xxii. (1872) 357.—Europ.
humilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 114.—Japon.
hybrida ×, *Kerner, in Oestr. Bot. Zeitschr.* xxiv. (1874) 382.—Austria.
Hyosciami, Wall. ex Benth. in Wall. Pl. As. Rar. i. 59, 46 = bracteosa.
incisa, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 390.—Japon.
integrifolia, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 108.—Nepal.
intermedia, Boiss. & Orph. ex Boiss. Fl. Orient. iv. 802 = A. Chia.
interrupta, Dulac, Fl. Hautes-Pyr. 407 = genevensis.
Iva, *Schreb. Pl. Vert. Unilab.* 15.—Reg. Mediterr.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 114.—Japon.
laevigata, *Boiss. Fl. Orient.* iv. 804.—Oriens.
lanata, F. Mart. ex Steud. Nom. ed. II. i. 46 = genevensis.
latifolia, Host, Fl. Austr. ii. 119 = genevensis.
latifolia, Schur, Enum. Pl. Transs. 545 = astonolosa.
Laxmanni, *Benth. Lab. Gen. et Sp.* 697.—Oriens.
lobata, *D. Don, Prod. Fl. Nep.* 108.—Reg. Himal.
lucida, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, xxii. (1868) 9.—Corsica.
lupulina, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 390.—China. bor.

AJUGA :—
lycia, *Stapf, in Denkschr. Acad. Wien*, I. (1885) 102. —Lycia.
macrosperma, *Wall. ex Benth. in Wall. Pl. As. Rar.* i. 58.—Ind. or.
mesogitana, Boiss. Diagn. Ser. I. vii. 62 = A. Chia.
mollissima, Wall. ex Steud. Nom. ed. II. i. 46 = Anisomeles ovata.
montana, Reichb. Fl. Germ. Excurs. 314 = genevensis.
moschata, Mill. ex Schreb. Pl. Vert. Unilab. 26 = A. Iva.
multiflora, *Bunge, Enum. Pl. Chin. Bor.* 51.—China bor.
nana, *Gilib. Fl. Lituan.* 323.—Europ.
Nanti, *Bor. in Mém. Soc. Acad. Maine-et-Loire*, xiv. (1863) 44.—Gallia.
oblongata, *Bieb. Fl. Taur. Cauc. Suppl.* 388.—Caucas.
ophrydis, *Burch. ex Benth. in DC. Prod.* xii. 597.—Afr. austr.
orientalis, *Linn. Sp. Pl.* 561.—Oriens.
ovata, *Wall. Cat.* n. 2036.—Burma.
palaestina, Boiss. Diagn. Ser. I. xii. 92 = A. Chia.
parviflora, *Benth. in Wall. Pl. As. Rar.* i. 59.—Reg. Himal.
pseudoiva, Rob. Cast. ex DC. Fl. Fr. v. 395 = A. Iva.
pseudo-pyramidalis, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 60.—Europ.
pumila, *Schur, Enum. Pl. Transs.* 545.—Europ.
pygmaea, *A. Gray, in Mem. Am. Acad.* N. S. VI. (1858–9) 402.—Japon.
pyramidalis, Bieb. Fl. Taur. Cauc. ii. 32 = genevensis.
pyramidalis, Huds. Fl. Ang. ed. I. 219 = reptans.
pyramidalis, *Linn. Sp. Pl.* 561.—Europ.
remota, Benth. in Wall. Pl. As. Rar. i. 59 = bracteosa.
repens, Gueldenst. ex Ledeb. Fl. Ross. iii. 447 = reptans.
repens, Roxb. Fl. Ind. iii. 3 = macrosperma.
reptans, *Linn. Sp. Pl.* 561.—Europ.
robusta, *Baker, in Journ. Linn. Soc.* xx. (1884) 235.—Madag.
rotundifolia, *Willk. & Cut. ex Willk. in Linnaea*, xxx. (1859–60) 120.—Hispan.
rugosa, Host, Fl. Austr. ii. 120 = genevensis.
rupestris, Schleich. ex Steud. Nom. ed. II. i. 46 = pyramidalis.
salicifolia, *Schreb. Fl. Vert. Unilab.* 26.—As. Min.
salicifolia, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 265 = oblongata.
scoparia, Boiss. Diagn. Ser. I. vii. 63 = A. Chamaecistus.
sikkimensis, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 114 in syn. = macrosperma.
sinuata, *R. Br. Prod.* 503.—Austral.
stolonifera, *Jeanb. & Timb. in Bull. Sc. Phys. & Nat. Toulouse*, iii. (1875–76) 577.—Europ.
Tenorii, Presl, Fl. Sicul. Praef. p. xxxvi. = acaulis.
Thomsoni, Maxim. in Bull. Acad. Pétersb. xxix. (1883) 189 = macrosperma.
tridentata, Benth. Lab. Gen. et Sp. 701 = australis.
tridactylites, Ging. ex Benth. l. c. 699 = A. Chia.
trifida, Dulac, Fl. Hautes-Pyr. 406 = A. Chamaepitys.
trifidifolia, Stokes, Bot. Mat. Med. iii. 288 = A. Chamaepitys.
vesiculifera, *Herd. in Bull. Soc. Nat. Mosc.* xli. (1868) i. 394.—As. temp.
vestita, *Boiss. Diagn.* Ser. I. vii. 62.—Graecia.
virgata, Benth. Lab. Gen. et Sp. 700 = australis.
yesoensis, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* ii. 467.—Japon.

AJUVEA, Steud. Nom. ed. I. 25 (1821) = **Aiouea**, Aubl. (Laurin.).

AKANIA, Hook. f. in Benth. & Hook. f. Gen. i. 409 (1862). *SAPINDACEAE*, Benth. & Hook. f. i. 409. Hillii, *Hook. f. l. c.*—Austral.

AKEA, Stokes, Bot. Mat. Med. ii. 353 (1812) = **Blighia**, Koen. (Sapind.).
solitaria, Stokes, I. c. 354 = B. sapida.

AKEBIA, Decne. in Arch. Mus. Par. i. (1837) 195. t. 13. *BERBERIDEAE*, Benth. & Hook. f. i. 42.
clematifolia, *Sieb. & Zucc. Fl. Jap.* i. 146.—Japon.
lobata, *Decne. in Ann. Sc. Nat.* Sér. II. xii. (1839) 107. —Japon.
quercifolia, *Sieb. & Zucc. Fl. Jap.* i. 146.—Japon.
quinata, *Decne. in Ann. Sc. Nat.* Sér. II. xii. (1839) 107.—China.

AKEESIA, Tussac, Fl. Antill. i. 66 (1808) = **Blighia**, Koen. (Sapind.).
africana, Tussac, l. c. = B. sapida.

AKENTRA, Benj. in Linnaea, xx. (1847) 319 = **Utricularia**, Linn. (Lentib.).
inflata, Benj. l. c.—Surinam.

AKLEMA, Rafin. Fl. Tellur. iv. 114 (1836) = **Euphorbia**, Linn.
nudiflora, Rafin. l. c. = E. nudiflora.

AKYLOPSIS, Lehm. Ind. Sem. Hort. Hamb. (1850) 3; cf. Otto, Hamb. Gartz. viii. (1852) 549 = **Matricaria**, Linn. (Compos.).
suaveolens, Lehm. I. c. = M. discoidea.

ALACOSPERMUM, Neck. Elem. i. 167 (1790) = **Cryptotaenia**, DC. (Umbell.).

ALAFIA, Thou. Gen. Nov. Madag. 11 (1809). *APOCYNACEAE*, Benth. & Hook. f. ii. 717.
Blastotrophe, Didrichs. in Kjoeb. Vidensk. Meddel. (1854) 192.
madagascariensis, Rich. ex A. DC. in DC. Prod. viii. 415 = Thouarsii.
pauciflora, *Radlk. in Bremen Abh.* viii. (1883) 403.—Afr. trop.
Thouarsii, *Roem. Collect.* i. 203.—Madag.

ALAGOPHYLLA, Rafin. Fl. Tellur. ii. 33 (1836) = **Gesnera**, Linn.
dasyanthes, Rafin. l. c.—Hab.?

ALAGOPTERA, Mart. Gen. Comm. 267 (1842) = ALLAGOPTERA, Nees = **Diplothemium**, Mart. (Palm.).

ALALANTIA, Corr. in Ann. Mus. Hist. Nat. Par. vi. (1804) 383, *err. typ.* = **Atalantia**, Corr. (Rutac.).

ALAMANIA, La Llave & Lex. Nov. Veg. Descr. Fasc. ii. (Orch. Opusc.) 31. *ORCHIDEAE*, Benth. & Hook. f. iii. 526.
punicea, *La Llave & Lex. l. c.*—Mexic.

ALAMANNIA, Lindl. Orch. Scel. 14 (1826) = praec.

ALANDINA, Neck. Elem. ii. 456 (1790) = **Moringa**, Juss.

ALANGIUM, Lam. Encyc. i. 174 (1783). *CORNACEAE*, Benth. & Hook. f. i. 949.
Angolam, Adans. Fam. ii. 85 (1763).
Kara-angolam, Adans. Fam. ii. 84 (1763).
acuminatum, Wight, ex Steud. Nom. ed. II. i. 47 = Lamarckii.
begoniaefolium, Baill. Hist. Pl. vi. 270 = Marlea begoniaefolia.
cordifolium, *Zoll. & Mor. ex Zoll. Syst. Verz. Ind. Archip.* 63.—Malaya.
decapetalum, Lam. Encyc. i. 174 = Lamarckii.
ebenaceum, Griff. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 742 = Marlea ebenacea.
frutescens, *Zoll. & Mor. ex Zoll. Syst. Verz. Ind. Archip.* 63.—Malaya.
glandulosum, Thw. Enum. Pl. Zeyl. 133 = Lamarckii.
hexapetalum, Lam. Encyc. i. 174 = Lamarckii.
Lamarckii, *Thw. Enum. Pl. Zeyl.* 133.—Geront. trop.
latifolium, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 741 = Lamarckii.
Mohillae, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 105. —Ins. Comor.
octopetalum, Blanco, Fl. Filip. ed. II. 310 = Lamarckii.
sundanum, Miq. Fl. Ind. Bat. i. 1. 774 = Lamarckii.
tomentosum, Lam. Encyc. i. 174 = Lamarckii.

ALANIA, Endl. Gen. Pl. 151 (1836). *LILIACEAE*, Benth. & Hook. f. iii. 797.
Allania, Meissn. Gen. 401 (304) (1842).
Cunninghami, *Steud. Nom.* ed. II. i. 47.—Austral.
Endlicheri, Kunth, Enum. Pl. iv. 644 = Cunninghami.

ALARÇONIA, DC. Prod. v. 537 (1836) = **Wyethia**, Nutt. (Compos.).
angustifolia, DC. l. c. = W. angustifolia.
helenioides, DC. l. c. = W. robusta.

ALATERNOIDES, Adans. Fam. ii. 304 (1763) = **Phylica**, Linn. (Rhamnac.).

ALATERNUS, [Tourn.] Mill. Gard. Dict. ed. VI. (1752) = **Rhamnus**, Linn.
alpinus, Moench, Meth. 344 = R. alpinus.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 3 = R. Alaternus.
balearica, Duham. ex Steud. Nom. ed. I. 685 = R. Alaternus.
cordatus, Rafin. Sylva Tellur. 27.—Hab. ?
glabra, Mill. Gard. Dict. ed. VIII. n. 2 = R. Alaternus.
glandulosus, Rafin. Sylva Tellur. 27 = R. glandulosus.
hispanicus, Hort. ex Steud. Nom. ed. I. 685 = R. Alaternus.
hybridus, Rafin. Sylva Tellur. 27 = R. hybridus.
integrifolius, Rafin. l. c. = R. integrifolius.
lanceolatus, Rafin. l. c. = R. lanceolatus.
latifolia, Mill. Gard. Dict. ed. VIII. n. 4 = R. Alaternus.
ovatus, Rafin. Sylva Tellur. 27.—Hab. ?
Phylica, Mill. Gard. Dict. ed. VIII. n. 1 = R. Alaternus.
prinoides, Rafin. Sylva Tellur. 27 = R. prinoides.
pumilus, Rafin. l. c. = R. pumilus.
rotundifolia, Hort. ex Steud. Nom. ed. I. 685 = R. Alaternus.
sempervirens, Rochl. ex Steud. Nom. ed. I. 493 = Lonicera sempervirens.
variegata, Hort. ex Steud. Nom. ed. I. 685 = R. Alaternus.

ALATHRAEA, Steud. Nom. ed. II. ii. 13, in syn. (1841) = seq.

ALATRAEA, Neck. Elem. i. 374 (1790) = **Phelypaea**, Tourn. (Orobanch.).

ALBERSIA, Kunth, Fl. Berol. ed. II. ii. 144 (1838) = **Amarantus**, Linn.
arenaria, Schur, Enum. Pl. Transs. 566 = Am. Blitum.
ascendens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 142 = Am. ascendens.
Blitum, Kunth, Fl. Berol. ed. II. ii. 144 = Am. Blitum.
caudata, Boiss. Fl. Orient. iv. 992 = Am. viridis.
deflexa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 142 = Am. deflexus.
gracilis, Webb & Berth. Phyt. Canar. iii. 287 = Am. gracilis.
livida, Kunth, Fl. Berol. ed. II. 144 = Am. Blitum.
oleracea, Kunth, l. c. = Am. Blitum.
polygama, Boiss. Fl. Orient. iv. 991 = Am. polygamus.
polygonoides, Kunth, Fl. Berol. ed. II. ii. 144 = Am. polygonoides.
polystachya, Kunth, l. c. = Am. polystachyus.
prostrata, Kunth, l. c. = Am. deflexus.

ALBERTA, E. Mey. in Linnaea, xii. (1838) 258. *RUBIACEAE*, Benth. & Hook. f. ii. 109.
laurifolia, *Baker, in Journ. Linn. Soc.* xviii. (1881) 271.—Madag.
magna, *E. Mey. in Linnaea*, xii. (1838) 258.—Afr. austr.
minor, *Baill. Adansonia*, xii. (1876–79) 247.—Madag.

ALBERTIA, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 603. *UMBELLIFERAE.*
commutata, *Regel & Schmalh. l. c.* 604.—Turkestan.
margaritifera, *Regel & Schmalh. l. c.* 605.—Turkestan.
paleacea, *Regel & Schmalh. l. c.* 606.—Turkestan.

ALBERTINIA, Spreng. Neue Entdeck. ii. 133 (1821). *COMPOSITAE*, Benth. & Hook. f. ii. 227.
SYMBLOMERIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 284.
arborea, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 236 = Vanillosmopsis arborea.
bardanoides, Mart. in DC. Prod. v. 81 = Vernonia crotonoides.
bicolor, DC. Prod. v. 81 = Eremanthus bicolor.

ALBERTINIA :—
bicolor, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 236 = Eremanthus incanus.
brasiliensis, *Spreng. Neue Entdeck.* ii. 133.—Bras.
Candolleana, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 235 = Vanillosmopsis erythropappa.
capitata, DC. Prod. v. 82 = Vanillosmopsis capitata.
Clausseni, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 15 = Vanillosmopsis erythropappa.
crotonoides, DC. Prod. v. 81 = Vernonia crotonoides.
Elaeagnus, Mart. ex DC. l. c. = Eremanthus Elaeagnus.
erythropappa, *DC. in Deless. Ic. Sel.* iv. t. 5.—Bras.
erythropappa, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 286 = Vanillosmopsis polycephala.
glomerulata, DC. Prod. v. 82 = Eremanthus glomerulatus.
gonoclados, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 144 = Piptolepis Martiana.
goyazensis, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 425 = Eremanthus goyazensis.
incana, Less. in Linnaea, iv. (1829) 342 = Eremanthus incanus.
incanescens, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 15 = Vanillosmopsis erythropappa.
mollis, Sch. Bip. ex Baker, l. c. 164 = Eremanthus mollis.
multiflora, Mart. in DC. Prod. v. 82 = Vanillosmopsis polycephala.
obtusata, Mart. l. c. 81 = Eremanthus glomerulatus.
Oleaster, Mart. ex Baker, in Mart. Fl. Bras. vi. II. 145 = Vernonia Oleaster.
pallidiseta, DC. Prod. v. 81 = Eremanthus glomerulatus.
polycephala, DC. l. c. 82 = Vanillosmopsis polycephala.
rufiseta, DC. l. c. 81 = Eremanthus glomerulatus.
saligna, Mart. in DC. l. c. 82 = Vanillosmopsis polycephala.
stellata, *Gardn. in Hook. Lond. Journ Bot.* v. (1846) 235.—Bras.
verbascifolia, Mart. in DC. Prod. v. 81 = Vernonia crotonoides.

ALBERTINIA, DC. Prod. v. 82, pro parte (1836) = **Vanillosmopsis**, Sch. Bip. (Compos.).

ALBERTISIA, Becc. Malesia, i. (1877) 161. *MENISPERMACEAE.*
papuana, *Becc. l. c.* 162 = N. Guin.

ALBIKIA, J. & C. Presl, Rel. Haenk. i. 184 et 34, 35 (1828) = **Hypolytrum**, Rich. (Cyperac.).
schoenoides, C. Presl, l. c. 185 = H. trinervium.
scirpoides, C. Presl, l. c. = H. latifolium.

ALBINA, Giseke, Prael. Linn. 199, 207 (1792) = **Alpinia**, Linn. (Scitam.).

ALBINEA, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zel. 37 (1853) = **Pleurophyllum**, Hook. f. (Compos.).
oresigenesa, Hombr. & Jacquinot, ex Decne. l. c. = P. oresigenesum.

ALBIZIA, Auct. = seq.

ALBIZZIA, Durazz. in Mag. Tosc. iii. IV. (1772) 11; F. Muell. in Journ. Bot. x. (1872) 8. *LEGUMINOSAE*, Benth. & Hook. f. i. 596.
BESENNA, A. Rich. Tent. Fl. Abyss. i. 253 (1847).
SERIANTHES, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 84 in nota.
SERICANDRA, Rafin. Sylva Tellur. 119 (1838).
ZYGIA, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 92.
acrodena, *Miq. Fl. Ind. Bat.* i. *Suppl.* 281.—Sumatra.
affinis, Fourn. in Ann. Sc. Nat. Sér. IV. xiv. (1860) 371 = amara.
altissima, Hook. f. in Hook. Niger Fl. 332 = Pithecolobium altissimum.
amara, *Boiv. in Encyc. xix. Siècle*, ii. 34.—Ind. or.
amoenissima, *F. Muell. Fragm.* viii. 165.—Austral.
angolensis, *Welw. in Oliver, Fl. Trop. Afr.* ii. 360.—Afr. trop.
anthelmintica, *Brongn. in Bull. Soc. Bot. Fr.* vii. (1860) 902.—Abyss.
auriculata, *Fourn.* in Ann. Sc. Nat. Sér. IV. xii. (1865) 400.—N. Caled.
basaltica, *Benth. Fl. Austral.* ii. 422.—Austral.

ALBIZZIA :—
Benthamiana, Blume, ex Miq. Fl. Ind. Bat. i. 30 = montana.
bigemina, F. Muell. Sel. Pl. Indust. Cult. 264 = Pithecolobium bigeminum.
Boivini, *Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 378.—Madag.
brachycalyx, *Oliver, Fl. Trop. Afr.* ii. 361.—Afr. trop.
Brownii, *Walp. Rep.* i. 928.—Afr. trop.
bubalina, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 129.—Ins. Nicob.
canescens, *Benth. Fl. Austral.* ii. 423.—Austral.
Championi, Benth. in Hook. Kew Journ. iv. (1852) 79 = Pithecolobium lucidum.
Charpentieri, *Fourn. in Bull. Soc. Bot. Fr.* xii. (1865) 400.—N. Caled.
clypearia, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 129.—Ins. Nicob.
conjugato-pinnata, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 278.—Afr. trop.
coriacea, F. Muell. Sel. Pl. Indust. Cult. 264 = Pithecolobium subcoriaceum.
coriaria, *Welw. in Oliver, Fl. Trop. Afr.* ii. 360.—Afr. trop.
cyclosperma, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = fulgens.
dulcis, F. Muell. Sel. Pl. Indust. Cult. 12 = Pithecolobium dulce.
elata, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = procera.
elegans, *Kurz, in Pegu Rep. App. B.* 47.—Burma.
elliptica, *Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 374.—Abyss.
fasciculata, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 129.—Ins. Nicob.
fastigiata, *Oliver, Fl. Trop. Afr.* ii. 361.—Afr. trop.
ferruginea, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 88.—Abyss.
floribunda, Fenzl, Reliq. Kotschy (1868). 5. t. 4 = anthelmintica.
Forbesii, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 92.—Afr. trop.
Fournieri, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 341 = Pithecolobium Fournieri.
fulgens, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 85.—Austral.; N. Caled.
glaberrima, *Benth. l. c.* 88.—Guinea.
glabrescens, *Oliver, Fl. Trop. Afr.* ii. 357.—Afr. trop.
glomeriflora, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 74.—Burma.
granulosa, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 85.—N. Caled.
Harveyi, *Fourn. in Bull. Soc. Bot. Fr.* xii. (1865) 399.—Afr. trop.
Hendersonia, F. Muell. in Journ. Bot. x. (1872) 10 = Pithecolobium Hendersonii.
heterophylla, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 300.—Burma.
hypoleuca, Oliver, Fl. Trop. Afr. ii. 356 = Harveyi.
Isembergiana, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = A. Julibrissin.
Jaubertiana, *Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 381.—Madag.
Jiringa, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 300.—Burma.
Julibrissin, *Durazz. in Mag. Tosc.* iii. IV. (1772) 11.—As.; Afr. trop.
Junghuhniana, F. Muell. in Journ. Bot. x. (1872) 9 = Pithecolobium Junghuhnianum.
latifolia, Boiv. in Encyc. xix. Siècle, ii. 33 = A. Lebbek
latisiliqua, F. Muell. Sel. Pl. Indust. Cult. 12 = Lysiloma latisiliqua.
Lebbek, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 87.—Geront. trop.
lebbekioides, *Benth. l. c.* 89 = odoratissima.
lentiscifolia, *Benth. in Trans. Linn. Soc.* xxx. (1875) 559.—N. Caled.
littoralis, *Teijsm. & Binn. in Tijdschr. Ned. Ind.* xxix. (1866) 259.—Molucc.
lomatocarpa, DC. Prod. ii. 467 = odoratissima.
lophantha, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 86.—Austral.
lucida, *Benth. l. c.*—Ind. or.; Malaya.
macrothyrsa, *Miq. Fl. Ind. Bat.* i. *Suppl.* 281.—Java.
malacophylla, Walp. Ann. ii. 457 = ferruginea.
marginata, Buch.-Ham. ex Wall. Cat. n. 5243 = stipulata.

ALBIZZIA:—
micrantha, Boiv. in Encyc. xix. Siècle, ii. 34 = odoratissima.
Milleti, *Benth. in Hook. Lond. Journ. Bot.* iii. (1846) 89.—China.
mollis, Boiv. in Encyc. xix. Siècle, ii. 33 = A. Julibrissin.
moluccana, *Miq. Fl. Ind. Bat.* i. 26.—Ins. Molucc.
monilifera, *F. Muell. in Journ. Bot.* x. (1872) 10.—Austral.
montana, *Benth. in Miq. Pl. Jungh.* 267.—Java.
mossambicensis, *Bolle, in Peters, Reise Mossamb. Bot.* i. 4.—Afr. trop.
myriophylla, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 90.—Ind. or.
nefasia, Walp. Ann. ii. 457 = Acacia abyssinica.
nellyrenza, R. Grah. in Wall. Cat. n. 5240 = amara.
Nemu, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = A. Julibrissin.
Neumanniana, *Heynh. Nom.* ii. 17.—Austral.
obovata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 558.—N. Caled.
odoratissima, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 88.—Ind. or.
paivana, *Fourn. in Ann. Sc. Nat.* Sér. IV. xv. (1861) 172.—N. Caled.
pallida, *Fourn. l. c.* xiv. (1860) 375.—Abyss.
pallida, Harv. in Harv. & Sond. Fl. Cap. ii. 284 = Harveyi.
pedicellata, *Baker, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 563.—Malacca.
Petersiana, *Oliver, Fl. Trop. Afr.* ii. 362.—Afr. trop.
polyphylla, *Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 372.—Madag.
procera, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 89.—As. trop.; Austral.
pruinosa, F. Muell. in Journ. Bot. x. (1872) 9 = Pithecolobium pruinosum.
purpurascens, Blume, ex Miq. Fl. Ind. Bat. i. 1. 28 = stipulata.
purpurea, *Boiv. ex Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 378.—Ins. Comor.
Quartiniana, *Walp. Ann.* ii. 457.—Abyss.
ramiflora, *F. Muell. Fragm.* ix. 178.—Austral.
retusa, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 90.—Ins. Philipp.
rhombifolia, *Benth. l. c.* 87.—Senegal.
rivularis, Fourn. in Ann. Soc. Sc. Nat. Sér. IV. xv. (1861) 172 = fulgens.
rosea, *Carr. in Rev. Hortic.* (1870-71) 490.—Am. bor.
rostrata, Blume, Miq. Fl. Ind. Bat. i. 24 = Pithecolobium rostratum.
rotundata, *Blume, l. c.* 20.—N. Guinea.
rubiginosa, *Miq. l. c.* 27.—Java.
rufa, *Benth. in Hook. Lond. Journ. Bot.* v. (1844) 101.—As. trop.; Java.
salajeriana, *Miq. Fl. Ind. Bat.* i. 1. 21.—Ins. Salager.
Saman, F. Muell. Sel. Pl. Indust. Cult. 12 = Pithecolobium Saman.
saponaria, *Blume, Miq. Fl. Ind. Bat.* i. 1. 19.—N. Guin.
Schimperiana, *Oliver, Fl. Trop. Afr.* ii. 359.—Afr. trop.
sericocephala, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 91 = amara.
Smithiana, Benth. l. c. i. (1842) 527 = stipulata.
speciosa, Benth. l. c. = A. Lebbek.
splendens, *Miq. Fl. Ind. Bat. Suppl.* 280.—Sumatra.
stipulata, *Boiv. in Encyc. xix. Siècle*, ii. 33.—Ind. or.
streptocarpa, *Fourn. in Bull. Soc. Bot. Fr.* xii. (1865) 399.—N. Caled.
subfalcata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 559.—N. Caled.
Sutherlandi, *F. Muell. Fragm.* vi. 22.—Austral.
tenerrima, De Vriese, in Miq. Pl. Jungh. 270 = Acacia pennata.
tengerensis, *Miq. Fl. Ind. Bat.* i. 25.—Java.
Teysmanni, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 299, 300.—Burma.
Thozetiana, *F. Muell. ex Benth. Fl. Austral.* ii. 422.—Austral.
tomentella, *Miq. Fl. Ind. Bat.* i. 1. 20.—Java.
Tozeri, F. Muell. in Journ. Bot. x. (1872) 10 = Pithecolobium grandiflorum.
umbrosa, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 86 = Calliandra umbrosa.
Vaillantii, F. Muell. Fragm. v. 9 = Archidendron Vaillantii.
versicolor, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 359.—Afr. trop.

ALBIZZIA:—
viridis, *Fourn. in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 377.—Madag.
Welwitschii, *Oliver, Fl. Trop. Afr.* ii. 362.—Afr. trop.
Wightii, Wight & Arn. Prod. 274 = amara.
Wightiana, R. Grah. in Wall. Cat. n. 5259 = amara.

ALBRADIA, D. Dietr. Syn. v. 280 (1852) = seq.

ALBRANDIA, Gaudich. in Freyc. Voy. Bot. 509 (1826) = **Streblus**, Lour. (Urtic.).
Gaudichaudi, D. Dietr. Syn. Pl. v. 280 = S. asper?
orientalis, D. Dietr. l. c. = S. asper.
spinosa, D. Dietr. l. c. = Taxotrophis zeylanica.
timorensis, D. Dietr. l. c. = Taxotrophis zeylanica.

ALBUCA, Linn. Sp. Pl. ed. II. 438 (1762). *LILIACEAE*, Benth. & Hook. f. iii. 809.
BRANCIONA, Salisb. Gen. Pl. Fragm. 36 (1866).
FALCONERA, Salisb. l. c. (1866).
NEMAULAX, Rafin. Fl. Tellur. iii. 52 (1836).
PALLASTEMA, Salisb. Gen. Pl. Fragm. 36 (1866).
VIRDIKE, Adans. Fam. ii. (19) (1763).
abyssinica, *Dryand. in Vet. Acad. Nya Handl. Stockh.* (1784) 297; *Jacq. Coll.* i. 55.—Afr. austr.
alba, Lam. Encyc. i. 76 = altissima.
altissima, *Dryand. in Vet. Acad. Nya Handl. Stockh.* (1784) 292; *Jacq. Coll.* ii. 264.—Afr. austr.
angolensis, *Welw. Apont.* 591; *et in Saund. Ref. Bot.* t. 336.—Angola.
anthericoides, Sweet, Hort. Brit. ed. I. 417 = Ornithogalum suaveolens.
aurea, *Jacq. Coll.* iv. 202.—Afr. austr.
Bainesii, *Baker, in Journ. Linn. Soc.* xiii. (1873) 290.—Afr. trop.
caudata, *Jacq. Coll.* iv. 203.—Afr. austr.
chlorantha, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 251.—Angola.
coarctata, *Dryand. in Vet. Acad. Nya Handl. Stockh.* (1784) 295.—Afr. austr.
Cooperi, *Baker, in Journ. Bot.* xii. (1874) 366.—Afr. austr.
cornuta, [DC. in] Red. Liliac. i. t. 70 = altissima.
elastica, Salisb. Prod. 239 = abyssinica.
elata, Salisb. l. c. 238 = altissima.
Elwesi, Regel, in Act. Hort. Petrop. vi. (1879) 538 = Wakefieldii.
exuviata, Ker-Gawl. in Bot. Mag. t. 871 = Urginea exuviata.
fastigiata, *Dryand. in Vet. Acad. Nya Handl. Stockh.* (1784) 296.—Afr. austr.
fibrosa, *Baker, in Gard. Chron.* (1874) ii. 386.—Afr. austr.
filifolia, Ker-Gawl. Bot. Reg. vii. t. 557 = Urginea filifolia.
flaccida, Baker, in Saund. Bot. Ref. t. 334 = Cooperi.
flaccida, *Jacq. Coll.* iv. 201.—Afr. austr.
fragrans, *Jacq. Hort. Schoenb.* i. 44. t. 84.—Afr. austr.
fugax, Ker-Gawl. Bot. Reg. iv. t. 311 = Urginea fragrans.
galeata, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 251.—Angola.
Gardeni, Hook. Bot. Mag. t. 4842 = Speirantha convallarioides.
glandulosa, *Baker, in Gard. Chron.* (1875) i. 814.—Afr. austr.; China.
juncifolia, *Baker, l. c.* (1876) i. 534.—Afr. austr.
juncifolia, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 251 = monophylla.
longifolia, *Fisch. ex Steud. Nom.* ed. II. i. 47.—Afr. austr.
lutea, Lam. Encyc. i. 76 = major, minor.
major, *Linn. Sp. Pl.* ed. II. 438.—Afr. austr.
maxima, *Burm. f. Fl. Cap. Prod.* 9.—Afr. austr.
minor, Gleditsch, in Mem. Acad. Berl. (1796) 91 = Uropetalum serotinum.
minor, *Linn. Sp. Pl.* ed. II. 438.—Afr. austr.
monophylla, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 251.—Angola.
myogaloides, *Welw. ex Baker, l. c.* 250.—Angola.
Nelsoni, *N. E. Br. in Gard. Chron.* (1880) ii. 198.—Afr. austr.
nematodes, *Sweet, Hort. Brit.* ed. II. 526.—Afr. austr.
parviflora, *Donn, Hort. Cant.* ed. II. 41.—Afr. austr.
physodes, Ker-Gawl. in Bot. Mag. t. 1046 = Urginea physodes.
polyphylla, *Baker, in Gard. Chron.* (1874) i. 471.—Afr. austr.
rigida, *Salisb. Prod.* 239.—Afr. austr.
setosa, *Jacq. Coll. Suppl.* 100. t. 14.—Afr. austr.

ALBUCA:—
Shawii, *Baker, in Journ. Bot.* xii. (1874) 367.—Afr. austr.
sordida, Salisb. Prod. 238 = major.
spiralis, *Linn. f. Suppl.* 196.—Afr. austr.
subspicata, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 251.—Angola.
tenuifolia, *Baker, in Saund. Ref. Bot.* t. 335.—Afr. austr.
viridiflora, *Jacq. Coll. Suppl.* 98.—Afr. austr.
viscosa, *Linn. f. Suppl.* 196.—Afr. austr.
vittata, Ker-Gawl. in Bot. Mag. t. 1329 = Ornithogalum vittatum.
Wakefieldii, *Baker, in Bot. Mag.* t. 6429.—Afr. trop.
Yerburyi, *Ridl. in Journ. Bot.* xxii. (1884) 370.—Aden.

ALBUCEA, Reichb. Fl. Germ. Excurs. 109 (1830) = **Ornithogalum**, Linn. (Liliac.).
chlorantha, Reichb. Ic. Fl. Germ. t. 1032 = O. nutans.
nutans, Reichb. Fl. Germ. Excurs. 110 = O. nutans.

ALBUGA, Schreb. Gen. i. 221 (1789) *err. typ.* = **Albuca**, Linn. (Liliac.).

ALBUGOIDES, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 372 = **Albuca**, Linn. (Liliac.).
erectum, Medic. l. c.—Afr. austr.?

ALCAEA, Hill, Hort. Kew. 213 (1768); Burm. f. Fl. Ind. 149 (1768) = **Althaea**, Linn. (Malrac.).

ALCAMASPINORA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 6.—Quid?

ALCANNA, Gaertn. Fruct. ii. 133. t. 110 (1791) = **Lawsonia**, Linn. (Lythrar.).
spinosa, Gaertn. l. c. = L. alba.

ALCANNA, Orph. in Bull. Congr. Bot. Pétersb. 1869 (1870) 138 = **Alkanna**, Tausch (Borag.).

ALCEA [Tourn.] Linn. Syst. ed. I. (1735) = **Althaea**, Tourn. (Malvac.).
acaulis, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 251 = Alth. acaulis.
aegyptiaca, Boiss. Diagn. Ser. II. i. 103 = Alth. lavateriflora.
africana, Lour. Fl. Cochinch. 421 = Alth. africana
apterocarpa, Boiss. Fl. Orient. i. 830 = Alth. apterocarpa.
Aucheri, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252 = Alth. Aucheri.
Calverti, Boiss. Fl. Orient. i. 832 = Alth. Calverti.
denudata, Boiss. l. c. 833.—Armenia.
digitata, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252 = Alth. lavateraeflora.
ficifolia, Linn. Sp. Pl. 687 = Alth. ficifolia.
ficifolia, Nym. Consp. 127 = Alth. rosea.
glabrata, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252 = Alth. ficifolia.
Haussknechtii, Boiss. Fl. Orient. i. 830.—Syria.
Heldreichii, Boiss. l. c. 832 = Alth. Heldreichii.
Hohenackeri, Boiss. l. c. 833 = Alth. Hohenackeri.
indica, Burm. f. Fl. Ind. 149.—Java.
kurdica, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252, 253 = Alth. kurdica.
lasiocalycina, Boiss. Fl. Orient. i. 830.—Armenia.
lavateraeflora, Boiss. l. c. 828 = Alth. lavateraeflora.
lilacina, Boiss. l. c. 831 = Alth. apterocarpa.
microchiton, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 254.—As. Min.
pallida, Bess. Enum. Pl. Volh. 28 = Alth. pallida.
palmata, Gilib. Fl. Lituan. i. 65 = Malva Alcea.
pinnatifida, Gilib. l. c. 66 = Malva moschata.
pontica, Janka, in Oestr. Bot. Zeitschr. xxiii. (1873) 201.—Turcia
procumbens, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 235. t. 17.—Hab.?
remotiflora, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 252 = Alth. remotiflora.
rhyticarpa, Trautv. ex Boiss. Fl. Orient. i. 827 = Alth. sulphurea.
rosea, Linn. Sp. Pl. 687 = Alth. rosea.
rosea, Falk, Beitr. ii. 221 = Alth. ficifolia.
rufescens, Boiss. Fl. Orient. i. 828 = Alth. rufescens.
rugosa, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 254 = ficifolia.
schirazana, Alef. l. c. 252 = Alth. kurdica.
setosa, Alef. l. c. 255 = Alth. setosa.
striata, Alef. l. c. 252 = Alth. striata.
sulphurea, Alef. l. c. = Alth. sulphurea.

ALCHEMILLA, Linn. Syst. ed. I. (1735). *ROSACEAE*, Benth. & Hook. f. i. 621.
APHANES, Linn. Gen. ed. I. 33 (1737).
PERCEPIER, Dill. ex Moench, Meth. 690 (1794).
abyssinica, *Fres. in Mus. Senkenb.* ii. (1837) 161.—Abyss.
achilleaefolia, Remy, in Ann. Sc. Nat. Sér. III. viii. (1847) 224, linea penultima = pinnata.
acutiloba, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 173 = vulgaris.
alata, Ruiz & Pav. ex Steud. Nom. ed. II. i. 47 = pinnata.
algida ×, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 64.—Europ.
alpestris, F. W. Schmidt, Fl. Boëm. iii. 88 = vulgaris.
alpina, Bieb. Fl. Taur. Cauc. i. 114 = sericea.
alpina, *Linn. Sp. Pl.* 123.—Reg. bor. temp.
ambigens, Jord. ex Nym. Consp. 238 = vulgaris.
Aphanes, Leers, Fl. Herborn. 54 = arvensis.
aphanoides, *Mutis, ex Linn. f. Suppl.* 122.—N. Granat.
argentea, Lam. Fl. Fr. iii. 303 = alpina.
arvensis, *Scop. Fl. Carn.* ed. II. i. 115.—Reg. bor. temp.
barbata, *Presl, Epim. Bot.* 199.—Peruv.
bifurcata, *Hils. & Boj ex Baill. in Bull. Soc. Linn. Par.* i. (1882) 342.—Madag.
capensis, *Thunb. Prod. Pl. Cap.* 31.—Afr. austr.
ceylanica, Hook. f. Fl. Brit. Ind. ii. 361, sphalm. = indica.
ciliata, Willd. ex Steud. Nom. ed. II. i. 47, 48 = orbiculata.
conglomerata, F. W. Schmidt, Fl. Boëm. iii. 89 = vulgaris.
conjuncta, Bab. in Mag. Nat. Hist. x. (1842) 25 = alpina.
cornucopioides, *Roem. & Schult. Syst.* iii. 471.—Hispan.
eryptantha, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 259.—Abyss.
cuneata, Gaud. Syn. Fl. Helv. 117 = pentaphyllea.
cuneifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 432 = arvensis.
dendroidea, *Zoll. in Nat. en Geneesk. Arch. Neerl. Indie*, i. (1844) 484.—Malaya.
diandra, Willd. ex Steud. Nom. ed. II. i. 48 = rupestris.
elongata, *Eckl. & Zeyh. Enum.* 264.—Afr. austr.
equisetiformis, *Trev. in Bot. Zeit.* vii. (1849) 215.—Venezuela.
erodiifolia, *Wedd. Chl. And.* ii. 247.—N. Granat.
Eupatoria, *Michx. Fl. Bor. Am.* i. 287.—Am. bor.
fissa, Schum. ex Sturm, Deutschl. Fl. Heft 56 (1830) = pyrenaica.
frigida, *Wedd. Chl. And.* ii. 245.—Boliv.
galioides, *Benth. Pl. Hartw.* 134.—N. Granat.
galioides, *Trev. in Bot. Zeit.* vii. (1849) 215.—Venezuela.
Gehecbii, *Siegfr. ex Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 70.—Europ.
glaberrima, F. W. Schmidt, Fl. Boëm. iii. 89 = vulgaris.
glabra, Suffr. in Flora, xix. (1836) 321 = vulgaris.
glabrata, Tausch, ex Steud. Nom. ed. II. i. 48 = vulgaris.
glaucescens, Wallr. in Linnaea, xiv. (1840) 549 = vulgaris.
Gunae, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xvii. (1868) 668.—Afr. trop.
helvetica ×, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 64.—Europ.
heptaphylla, Schleich. ex Steud. Nom. ed. II. i. 48 = pentaphyllea
hirsuta, *H. B. & K. Nov. Gen. et Sp.* vi. 226.—Am. austr.
hybrida, F. W. Schmidt, Fl. Boëm. iii. 89 = vulgaris.
indica, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 8.—Mont. Nilghir.; Zeylan.
intermedia, [Clairv.] Man. Herb. 43 = alpina.
latifolia, Salisb. Prod. 361 = vulgaris.
Lechleriana, *Grisch. in Goett. Abh.* xxiv. (1879) 124.—Reg. Argent.
lobata, Pall. Reise, ii. 568 = vulgaris.
lycopodioides, *Trev. in Bot. Zeit.* vii. (1849) 215.—Venezuela.
madagascariensis, *O. Hoffm. in Bremen Abh.* vii. (1882) 336.—Afr. trop.
Mandoniana, *Wedd. Chl. And.* ii. 246.—Boliv.
microcarpa, *Boiss. & Reut. Diagn. Hisp.* 11.—Hispan.
minor, Tausch, ex Host, Fl. Austr. i. 219 = pyrenaica.
minor, Huds. Fl. Angl. ed. I. 59 = vulgaris.
monandra, Stokes, Bot. Mat. Med. i. 236 = arvensis.
monandria, *Sw. Prod. Veg. Ind. Occ.* 38.—Ind. occ.
montana, F. W. Schmidt, Fl. Boëm. iii. 88 = vulgaris.

ALCHEMILLA:—
multicaulis, Tausch, ex Steud. Nom. ed. II. i. 48 = vulgaris.
multifida, *Scheele, in Flora*, xxvi. (1843) 449.—Europ.
nivalis, *H. B. & K. Nov. Gen. et Sp.* vi. 224. t. 560.—Equador.
occidentalis, Nutt. in Torr. & Gray, Fl. N. Am. i. 432 = arvensis.
orbiculata, *Ruiz & Pav. Fl. Per.* i. 68.—Peruv.
palmata, *E. Mey. in Drège, Zwei Pfl. Docum.* 163, nomen.—Afr. austr.
palmata, Gilib. Exercit. ii. 429 = vulgaris.
palmatifida, Tausch, ex Reichb. Pl. Crit. i. 6 = pyrenaica.
paniculata, F. W. Schmidt, Fl. Boëm. iii. 89 = vulgaris.
pectinata, H. B. & K. Nov. Gen. et Sp. vi. 226 = orbiculata.
pedata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 260.—Abyss.
pentaphyllea, *Linn. Sp. Pl.* 123.—Europ.
pinnata, *Ruiz & Pav. Fl. Per.* i. 69.—Peruv.
polylepis, *Wedd. Chl. And.* ii. 246.—Am. austr.
potentilloides, Hils. & Boj. ex Baill. in Bull. Soc. Linn. Par. i. (1882) 343 = madagascariensis.
pratensis, F. W. Schmidt, Fl. Boëm. iii. 88 = vulgaris.
pubescens, Bieb. Fl. Taur. Cauc. i. 114 = vulgaris.
pubescens, Willd. Hort. Berol. t. 79 = praec.?
pusilla, *Pomel, Nouv. Mat. Fl. Atl.* 159.—Afr. bor.
pyrenaica, *Dufour, in Ann. Gén. Sc. Phys.* viii. (1821) 228.—Mont. Pyren.
repens, *Presl, Epim. Bot.* 199.—Peruv.
Rothii, *Oliver, Fl. Trop. Afr.* ii. 378.—Afr. trop.
rupestris, *H. B. & K. Nov. Gen. et Sp.* vi. 224.—Quito.
Rutenbergii, *O. Hoffm. in Bremen Abh.* vii. (1882) 336.—Afr. trop.
schizophylla, *Baker, in Journ. Linn. Soc.* xx. (1884) 137.—Madag.
sericata, Reichb. Pl. Crit. i. 6, in obs. = vulgaris.
sericea, *Willd. Enum. Hort. Berol.* 171, in nota.—Oriens.
sibbaldiaefolia, *H. B. & K. Nov. Gen. et Sp.* vi. 226. t. 561.—Mexic.
stemmatophylla, *Wedd. Chl. And.* ii. 248. t. 75.—N. Granat.
subsericea, Reut. Compte-rend. Soc. Hallér. (1853–54) 20 = alpina.
sylvestris, F. W. Schmidt, Fl. Boëm. iii. 88 = vulgaris.
tenuicaulis, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 191.—Afr. trop.
triloba, Gilib. Exercit. ii. 430 = arvensis.
tripartita, *Ruiz & Pav. Fl. Per.* i. 68.—Peruv.
truncata, Tausch, ex Steud. Nom. ed. II. i. 48 = vulgaris.
velutina, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 354.—Am. bor.
venusta, *Cham. & Schlecht. in Linnaea*, v. (1830) 573.—Mexic.
villosa, *Jungh. in Nat. en Geneesk. Arch. Neerl. Indie*, ii. (1845) 46.—Java.
vulcanica, *Cham. & Schlecht. in Linnaea*, v. (1830) 573.—Mexic.
vulcanica, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie*, ii. (1845) 587.—Malaya.
vulgaris, *Linn. Sp. Pl.* 123.—Reg. bor. temp.
vulgaris, Wight, Ic. t. 229 = indica.

ALCHORNEA, Sw. Prod. Veg. Ind. Occ. 98 (1788).
EUPHORBIACEAE, Benth. & Hook. f. iii. 314.
APARISTHMIUM, Endl. Gen. 1112 (1841).
BLEEKERIA, Miq. Fl. Ind. Bat. i. II. 407 (1859).
CLADODES, Lour. Fl. Cochinch. 574 (1790).
COELEBOGYNE, J. Sm. in Trans. Linn. Soc. xviii. (1841) 509. t. 36.
DIDEROTIA, Baill. Adansonia, i. (1860–61) 274.
HERMESIA, Humb. & Bonpl. in Willd. Sp. Pl. iv. 809 (1805).
LAUTEMBERGIA, Baill. Etud. Gén. Euph. 451 (1858).
ORFILEA, Baill. l. c. 452 (1858).
PALISSYA, Baill. l. c. 502 (1858).
PSEUDOTREWIA, Miq. Fl. Ind. Bat. i. II. 414 (1859).
SCHOUSBOEA, Schum. Beskr. Guin. Pl. 449 (1827).
STIPELLARIA, Benth. in Hook. Kew Journ. vi. (1854) 2.
WETRIA, Baill. Etud. Gén. Euph. 409 (1858).

ALCHORNEA:—
acutifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 171.—Peruv.
Blumeana, *Muell. Arg. l. c.* 167.—Java.
castaneaefolia, *A. Juss. Tent. Euph.* 42.—Fl. Orinoco.
columnularis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 378.—Bras.
cordata, *Benth. in Hook. Niger Fl.* 507.—Afr. trop.
eordata, *Muell. Arg. in DC. Prod.* xv. II. 901.—Am. trop.
eordifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 170.—Afr. trop.
eoriacea, *Muell. Arg. l. c.* 168.—Madag.
cuneata, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 122 = cuneifolia.
cuneifolia, *Muell. Arg. in DC. Prod.* xv. II. 900.—Sumatra.
Davidi, *Franch. Pl. Davidian.* i. (1884) 264.—Mongolia.
discolor, *Poepp. & Endl. Nov. Gen.* iii. 19.—Bras.
Duparquetiana, *Baill. Adansonia*, xi. (1873–76) 175.—Afr. trop.
erythrosperma, *Klotzsch, ex Benth. in Hook. Kew Journ.* vi. (1854) 331.—Bras.
floribunda, *Muell. Arg. in Journ. Bot.* i. (1863) 336.—Afr. occ.
Gardneri, *Muell. Arg. in Flora*, xlvii. (1864) 435.—Bras.
glandulosa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 18. t. 221.—Bras.
glandulosa, Poit. ex Baill. Etud. Gén. Euph. 446 = latifolia.
grandiflora, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 170.—Venezuela.
grandis, *Benth. Bot. Sulph. Voy.* 164.—Am. trop.
Hilariana, *Baill. Adansonia*, v. (1864–65) 240.—Bras.
hirtella, *Benth. in Hook. Niger Fl.* 507.—Afr. trop.
ilicifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 170.—Austral.
intermedia, *Klotzsch, ex Benth. in Hook. Kew Journ.* vi. (1854) 331.—Bras.
Iricurana, *Casar. Nov. Stirp. Bras. Dec.* 24.—Bras.
javensis, Muell. Arg. in Linnaea, xxxiv. (1865–66) 170 = rugosa.
janeirensis, Casar. Nov. Stirp. Bras. Dec. 15 = nemoralis.
latifolia, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 46 = cordata.
latifolia, *Sw. Prod. Veg. Ind. Occ.* 98.—Ind. oce.
macrophylla, Mart. Herb. Fl. Bras. 271 = cordata.
madagascariensis, *Muell. Arg. in DC. Prod.* xv. II. 900.—Madag.
Martiana, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 375.—Bras.
megalophylla, *Muell. Arg. in Flora*, xlvii. (1864) 434.—N. Granat.
mollis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 168.—Ind. or.
multispicata, *Muell. Arg. l. c.* 168.—Madag.
nemoralis, *Mart. Herb. Fl. Bras.* 271.—Bras.
parviflora, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 168.—Ins. Philipp.
parvifolia, *Klotzsch, ex Benth. in Hook. Kew Journ.* vi. (1854) 331.—Bras.
parvifolia. Miq. in Linnaea, xxii. (1849) 797 = nemoralis.
platyphylla, *Muell. Arg. l. c.* xxxiv. (1865–66) 171.—Am. centr.
psilorhachis, Klotzsch, ex Benth. in Hook. Kew Journ. vi. (1854) 331 = nemoralis.
puberula, *Klotzsch, l. c.*—Bras.
pycnogyne, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 378.—Bras.
rotundifolia, Moric. ex Baill. Etud. Gén. Euph. 447 = nemoralis.
rufescens, *Franch. Pl. Davidian.* i. (1884) 265.—Mongol.
rugosa, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 170.—Ind. or.; Malaya.
scandens, *Muell. Arg. l. c.* 170.—Cochinch.
Schomburgkii, Benth. in Hook. Kew Journ. vi. (1854) 330 = Gardneri.
Schomburgkii, *Klotzsch, in Hook. Lond. Journ. Bot.* ii. (1843) 46.—Am. austr.
sidaefolia, *Baill. Etud. Gén. Euph.* 447.—Bras.
similis, *Muell. Arg. in Flora*, xlvii. (1864) 434.—Mexic.

ALCHORNEA:—
subrotunda, Baill. Etud. Gén. Euph. 447 = glandulosa.
Thozetiana, *Baill. ex Benth. Fl. Austral.* vi. 137.—Austral.
tiliaefolia, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 168.—Ind. or.
trewioides, *Muell. Arg. l. c.*—China.
triplinervia, *Muell. Arg. in DC. Prod.* xv. II. 909 = nemoralis.
villosa, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 168.—Malaya.
Zollingeri, Hassk. Cat. Hort. Bogor. Alt. 37; Retzia, 156 = villosa.

ALCHORNEOPSIS, Muell. Arg. in Linnaea, xxxiv. (1865-66) 156. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 315.
floribunda, *Muell. Arg. l. c.*—Peruv.

ALCHYMILLA, Rupp. Fl. Jen. ed. Hall. 64 (1745) = **Alchemilla**, Linn. (Rosac.).

ALCINA, Cav. Ic. i. 10. t. 15 (1791) = **Melampodium**, Linn. (Compos.).
minor, Cass. in Dict. Sc. Nat. lix. 243 = M. sericeum.
ovatifolia, Jacq. Eclog. Pl. 78 = M. divaricatum.
perfoliata, Cav. Ic. i. 11. t. 15 = M. perfoliatum.

ALCINIA, H. B. & K. Nov. Gen. et Sp. iv. 274 (1820) = praec.

ALCIOPE, DC. Prod. v. 209 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 439.
CELMISIA, Cass. in Dict. Sc. Nat. vii. 356 (1817).
lanata, *DC. Prod.* v. 210.—Afr. austr.
tabularis, *DC. l. c.*—Afr. austr.

ALDAMA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 14 (1824) = **Sclerocarpus**, Jacq. (Compos.).
dentata, La Llave, l. c. = S. dentatus.
dentata, Less. in Linnaea, v. (1830) 154 = S. Schiedeanus.
uniserialis, A. Gray, Pl. Lindh. 228 = S. uniserialis.

ALDEA, Ruiz & Pav. Prod. 19 (1794) = **Phacelia**, Juss. (Hydrophyll.).
circinnata, Willd. Enum. Suppl. 9 = P. circinnata.
magellanica, Willd. ex Steud. Nom. ed. II. i. 48 = Hydrophyllum magellanicum.
peruviana, Ruiz & Pav. Fl. Per. ii. 8. t. 114 = P. peruviana.
pinnata, Ruiz & Pav. l. c. = P. peruviana.

ALDELASTER, C. Koch, in Belg. Hortic. (1861) 324 = ADELASTER, Veitch = **Eranthemum**? Linn. (Acanth.).

ALDINA, Adans. Fam. ii. 328 (1763) = **Brya**, P. Br. (Legum.).

ALDINA, Endl. Gen. 1322 (1841). *LEGUMINOSAE*, Benth. & Hook. f. i. 560.
ALLANIA, Benth. in Hook. Lond. Journ. Bot. ii. (1840) 91.
discolor, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 12.—Bras.
heterophylla, *Benth. l. c.* 13.—Bras.
insignis, *Endl. in Walp. Rep.* i. 843.—Guiana.
latifolia, *Benth. in Mart. Fl. Bras.* xv. II. 12.—Bras.
macrophylla, *Spruce, ex Benth. l. c.* 13.—Bras.

ALDINIA, Scop. Introd. 173 (1777) = **Justicia**, Linn. (Acanth.).

ALDROVANDA, Monti, in Bonon. Sc. & Art. Inst. Comm. ii. III. 404. t. 12 (1747); Linn. Diss. Chen. 39 (1751). *DROSERACEAE*, Benth. & Hook. f. i. 663.
verticillata, Roxb. Fl. Ind. ii. 113 = vesiculosa.
vesiculosa, *Linn. Sp. Pl.* 281.—Europ.; Ind. or.

ALDUNATEA, Remy, in C. Gay, Fl. Chil. iii. 320. t. 38 (1849) = **Chaetanthera**, Ruiz & Pav. (Compos.).
chilensis, Remy, l. c. 322. t. 38. f. 1 (= *Oriastrum chilense*).

ALDUNATEA:—
gnaphalioides, Remy, in C. Gay, Fl. Chil. iii. 323 (= *Oriastrum gnaphalioides*).
lycopodioides, Remy, l. c. (= *Oriastrum gnaphalioides*).

ALECTORIDIA, A. Rich. Tent. Fl. Abyss. ii. 447. t. 99 (1851) = **Arthraxon**, Beauv. (Gramin.).
Quartiniana, A. Rich. l. c. 448 = Arth. ciliaris.

ALECTOROCTONUM, Schlecht. in Linnaea, xix. 252 (1847) = **Euphorbia**, Linn.
caracasanum, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 40 = E. caracasana.
cotinifolium, Schlecht. in Linnaea, xix. (1847) 252 = E. cotinifolia.
cotinoides, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 248 = E. cotinoides.
dilatatum, Klotzsch & Garcke, l. c. = E. dilatata.
nudiflorum, Klotzsch & Garcke, l. c. 40 = E. nudiflora.
ovatum, Schlecht. in Linnaea, xix. (1847) 252 = E. Schlechtendalii.
petiolare, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 248 = E. petiolaris.
Riedelianum, Klotzsch & Garcke, in Abh. l. c. (1860) 41 = E. cotinoides.
sanguineum, Klotzsch & Garcke, in Monatsb. l. c. (1859) 248 = E. haematodes.
scandens, Klotzsch & Garcke, l. c. = E. scandens.
Scotanum, Schlecht. in Linnaea, xix. (1847) 252 = E. Scotanum.
viride, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 248 = E. viridis.
Willdenowii, Klotzsch & Garcke, in Abh. l. c. (1860) 41 = E. cotinifolia.
Wrightii, Klotzsch & Garcke, in Monatsb. l. c. (1859) 248 = E. Wrightii.
Yavalquahuitl, Schlecht. in Linnaea, xix. (1847) 252 = E. Scotanum.

ALECTOROLOPHUS, Hall. Enum. Stirp. Helv. ii. 623 (1742); Moench, Meth. 440 (1794) = **Rhinanthus**, Linn. (Scrophul.).
alpinus, D. Dietr. Syn. Pl. iii. 553 = R. major.
alpinus, Reichb. f. Ic. Fl. Germ. xx. 66 = R. Crista-galli.
angustifolius, D. Dietr. Syn. Pl. iii. 553 = R. angustifolius.
buccalis, Heynh. Nom. ii. 18 = R. Crista-galli.
Crista-galli, Bieb. Fl. Taur. Cauc. ii. 68 = R. Crista-galli.
glaber, All. Fl. Pedem. i. 58 = R. Crista-galli.
grandiflorus, Wallr. Sched. Crit. 316 = R. Crista-galli.
hirsutus, All. Fl. Pedem. i. 58 = R. Crista-galli.
major, Reichb. Ic. Pl. Crit. viii. 13. fig. 975; Ic. Fl. Germ. xx. 65 = R. major.
minor, Dum. Fl. Belg. 33 = R. minor.
minor, Reichb. Fl. Germ. Excurs. 358 = R. Crista-galli.
parviflorus, Wallr. Sched. Crit. 318 = R. minor.
pulchella, Schum. ex Wimm. Fl. Schles. (1841) 277 = R. Crista-galli.
pulcher, Schum. ex Wimm. l. c. ed. III. (1857) 410 = R. Crista-galli.
ramosus, Schur, Enum. Pl. Transs. 880, Index = R. minor.
stenophyllus, Schur, l. c. = R. minor.
Trixago, Bieb. Fl. Taur. Cauc. ii. 68 = Bartsia Trixago.
villosus, Dum. Fl. Belg. 33 = R. Crista-galli.

ALECTRA, Thunb. Nov. Gen. 81 (1784). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 966.
GLOSSOSTYLIS, Cham. & Schlecht. in Linnaea, iii. (1828) 22.
HYMENOSPERMUM, Benth. in Wall. Cat. n. 3963 (1831).
STARBIA, Thou. Gen. Nov. Madag. 7 (1806).
abyssinica, *A. Rich. Tent. Fl. Abyss.* ii. 118.—Abyss.
asperrima, *Benth. in DC. Prod.* x. 340.—Abyss.
brasiliensis, *Benth. l. c.* 339.—Bras.
capensis, *Thunb. Prod. Pl. Cap.* 97.—Afr. austr.
cordata, Benth. in DC. Prod. x. 339 = indica.
hirsuta, *Klotzsch, in Peters, Reise Mossamb. Bot.* 221.—Afr. trop.
indica, *Benth. in DC. Prod.* x. 339.—Ind. or.
lurida, *Harv. Gen. S. Afr. Pl.* 250.—Afr. austr.

ALECTRA:—
major, *E. Mey. in Drège, Zwei Pfl. Docum.* 163.—Afr. austr.
melampyroides, *Benth. in DC. Prod.* x. 339.—Afr. austr.
minor, *E. Mey. in Drège, Zwei Pfl. Docum.* 163.—Afr. austr.
orobanchoides, *Benth. in DC. Prod.* x. 340.—Afr. austr.
parasitica, *A. Rich. Tent. Fl. Abyss.* ii. 117.—Abyss.
pedicularioides, *Baker, in Journ. Linn. Soc.* xx. (1884) 214.—Madag.
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 118.—Afr. trop.
pumila, *Benth. in DC. Prod.* x. 340.—Afr. austr.
senegalensis, *Benth. l. c.* 339.—Afr. trop.
stricta, *Benth. l. c.* 338.—Bras.
Thomsoni, *Hook. f. Fl. Brit. Ind.* iv. 297.—Ind. or.
Vogelii, *Benth. in DC. Prod.* x. 339.—Afr. trop.

ALECTRYON, Gaertn. Fruct. i. 217. t. 46 (1788) *SAPINDACEAE*, Benth. & Hook. f. i. 410.
canescens, *DC. Prod.* i. 617.—Austral.
carinatum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 119.—N. Caled.
connatum, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 299, 307 = Nephelium connatum.
coriaceum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 118.—Austral.
excelsum, *Gaert. Fruct.* i. 216. t. 46.—N. Zel.
ferrugineum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 84.—Ins. Molucc.
glabrum, *Radlk. l. c.*—Ins. Timor.
laeve, *Radlk. l. c.* 119.—Austral.
semicinereum, *Radlk. l. c.* 118.—Austral.
serratum, *Radlk. l. c.*—Austral.
sphaerococcum, *Radlk. l. c.* 119.—Celebes.
subcinereum, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* ix. (1879) 524.—Austral.
subdentatum, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 117.—Austral.
tomentosum, *Radlk. l. c.*—Austral.

ALEGRIA, Moç. & Sesse, ex DC. Prod. i. 516 (1824) = **Luehea**, Willd. (Tiliac.).
candida, Moç. & Sesse, l. c. 517 = L. platypetala..

ALEOME, Neck. Elem. iii. 68 (1790) = **Cleome**, Linn. (Cappar.).

ALEPIDEA, La Roche, Eryng. Hist. 19. t. 1 (1808). *UMBELLIFERAE*, Benth. & Hook. f. i. 878.
amatymbica, Eckl. & Zeyh. Enum. 339.—Afr. austr.
ciliaris, *La Roche, Eryng. Hist.* 19. t. 1.—Afr. austr.
cordata, E. Mey. in Drège, Zwei Pfl. Docum. 163 = ciliaris.
ferrata, Steud. Nom. ed. II. i. 48, sphalm. = *serrata*.
longifolia, E. Mey. ex Steud. l. c. = amatymbica.
peduncularis, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 320.—Abyss.
serrata, Eckl. & Zeyh. Enum. 339 = ciliaris.
Woodii, *Oliver, in Hook. Ic. Pl.* t. 1452.—Afr. austr.

ALEPYRUM, B. Br. Prod. 252, 253 (1810) = **Centrolepis**, Labill.
monogynum, Hook. f. Fl. Tasm. ii. 77. t. 138 = C. monogyna.
Muelleri, Hook. f. l. c. 78 = C. glabra.
muscoides, Hook. f. l. c. 77 = C. muscoides.
muticum, R. Br. Prod. 253 = C. mutica.
pallidum, Hook. f. Fl. Nov. Zel. i. 268 = Gaimardia pallida.
polygynum, R. Br. Prod. 253 = C. polygyna.
Pumilio, R. Br. l. c. = C. polygyna.
viride, T. Kirk, in Journ. Linn. Soc. xix. (1884) 286.—N. Zel.

ALETHRIS, Linn. Ord. Nat. (1764) = seq.

ALETRIS, Linn. Diss. Chen. 15 (1751); (Amoen. Acad. iii. 11). *HAEMODORACEAE*, Benth. & Hook. f. iii. 677.
STACHYOPOGON, Klotzsch, in Bot. Ergeb. Waldem. Reise, 49. t. 94 (1862).
alba, Michx. Fl. Bor. Am. i. 189 = farinosa.
arborea, Willd. Enum. Hort. Berol. 381 = Dracaena arborea.
aurea, *Walt. Fl. Carol.* 121.—Am. bor.

ALETRIS :—
bifolia, Burm. f. Fl. Cap. Prod. 10 = Lachenalia pendula.
capensis, Linn. Syst. ed. X. 985 = Veltheimia viridiflora.
chinensis, Lam. Encyc. i. 79 = Dracaena terminalis.
cochinchinensis, *Lour. Fl. Cochinch.* 204.—Cochinch.
dracaenoides, Roxb. ex Roem. & Schult. Syst. vii. 347.—Ind. or.
farinosa, Ker-Gawl. in Bot. Mag. t. 1418 = aurea.
farinosa, *Linn. Sp. Pl.* 319.—Am. bor.
farinosa, Thunb. Fl. Jap. 136 = japonica.
fragrans, Linn. Sp. Pl. ed. II. 456 = Dracaena fragrans.
glauca, Ait. Hort. Kew. i. 463 = Veltheimia glauca.
guineensis, Jacq. Hort. Vind. i. 63. t. 84 = Sanseviera guineensis.
hyacinthoides, Linn. Sp. Pl. ed. II. 456 = Sansevicra guinccnsis.
japonica, Houtt. Syst. xi. 486. t. 84. f. 2 = Funkia subcordata.
japonica, *Lamb. in Trans. Linn. Soc.* x. (1811) 407.—Japon.
japonica, Thunb. in Nov. Act. Ups. iii. (1780) 208 = Funkia ovata.
linguaeformis, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
littoralis, Koen. ex Steud. Nom. ed. II. i. 49 = Urginia indica ?
pumila, Ait. Hort. Kew. i. 464 (= *Tritomanthe pumila*).
pumila, Donn, Hort. Cant. ed. II. 42 = Dracaena ovata.
punicea, Labill. Pl. Nov. Holl. i. 85. t. 111 = Blandfordia marginata.
sarmentosa, Andr. Bot. Rep. t. 54 (= *Tritomanthe media*).
Uvaria, Linn. Mant. 367 = Kniphofia alooides.
zeylanica, Mill. Gard. Dict. ed. VIII. n. 4 = Sanseviera lanuginosa.

ALEURITES, Forst. Char. Gen. iii. t. 56 (1776). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 292.
Camirium, Gaertn. Fruct. ii. 194. t. 125 (1791).
Dryandra, Thunb. Fl. Jap. 13. t. 27 (1784).
Elaeococca, A. Juss. Euph. Tent. 38. t. 11 (1824).
Telopea, Soland. ex Baill. Etud. Gén. Euph. 345 (1858).
Vernicia, Lour. Fl. Cochinch. 586 (1790).
Ambinux, Pers. Syn. ii. 579 = triloba.
commutata, Geisel. Crot. Monogr. 82 = triloba.
cordata, *Steud. Nom.* ed. II. i. 49.—China; Japon.
cordifolia, Steud. l. c. = triloba.
japonica, Blume, in Miq. Ann. Mus. Bot. Lugd. Bat. iv. 120 = cordata.
laccifera, Willd. Sp. Pl. iv. 590 = Croton aromaticus.
lanceolata, Blanco, Fl. Filip. ed. I. 757 = triloba.
lobata, Blanco, l. c. 756 = triloba.
moluccana, Willd. Sp. Pl. iv. 590 = triloba.
peltata, Geisel. Crot. Monogr. 81 = Mallotus peltatus.
pentaphylla, *Wall. Cat.* n. 7959.—Ind. or.
saponaria, *Blanco, Fl. Filip.* ed. II. 519.—Ins. Philipp.
triloba, *Forst. Char. Gen.* 112.—As. trop.; Ins. Pacif.
trisperma, Blanco, Fl. Filip. ed. I. 755 = triloba.
Vernicia, Hassk. in Flora, xxv. (1842); Beibl. ii. 40 = cordata.
verniciflua, Baill. Hist. Pl. v. 116 = praec.

ALEURITIA, Spach, Hist. Vég. Phan. ix. 360 (1840) = **Primula**, Linn.
cortusoides, Spach, l. c. 361 = P. cortusoides.
farinosa, Spach, l. c. = P. farinosa.
longiflora, Spach, l. c. 362 = P. longiflora.

ALEURODENDRON, Reinw. in Blume, Cat. Gew. Buitenz. 87 (1823), nomen; Syll. Ratisb. ii. (1828) 12 = **Melochia**, Linn. (Sterculiat.).
album, Reinw. ll. cc. = M. Aleurodendron.

ALEVIA, Baill. Etud. Gén. Euph. 508 (1858) = **Bernardia**, P. Br. (Euphorb.).
leptostachya, Baill. l. c. 509 = B. interrupta.

ALEXA, Moq. in DC. Prod. xiii. II. 168 in nota (1849). *LEGUMINOSAE*, Benth. & Hook. f. i. 556.
Alexandra, Schomb. in Hook. Lond. Journ. Bot. iv. (1845) 12.

ALEXANDRA, Bunge, in Linnaea, xvii. (1843) 120. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 67.
Pterocalyx, Schenk, in Bull. Phys. Math. Acad. Pétersb. i. (1843) 361.
Lehmanni, *Bunge, in Linnaea* (1843), 120.—Soongar.

ALEXANDRA, Schomb. in Hook. Lond. Journ. Bot. iv. (1845), 12; et Barbac. (1845) 18 = **Alexa**, Moq. (Legum.).
Imperatricis, Schomb. ll. cc.—Guiana.

ALEXIA, Wight, Ic. t. 1293 (1850) = **Alyxia**, R. Br. (Apocyn.).

ALEXIS, Salisb. in Trans. Hort. Soc. i. (1812) 284 = **Amomum**, Linn. (Scitam.).
bifurca, Salisb. l. c. = A. exscapum.
grandiflora, Salisb. l. c. = A. grandiflorum.

ALEXITOXICON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 67 = **Vincetoxicum**, Moench (Asclep.).

ALFONSIA, H. B. & K. Nov. Gen. et Sp. i. 306 (1815) = **Elaeis**, Jacq. (Palm.).
oleifera, H. B. & K. l. c. 307 = E. melanococca.

ALFREDIA, Cass. in Bull. Soc. Philom. (1815) 175 = **Carduus**, Linn. (Compos.).
acantholepis, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 394 = C. acantholepis.
cernua, Cass. in Bull. Soc. Philom. (1815) 175 = C. cernuus.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 185 = Cnicus japonicus.
Karelini, Ledeb. Fl. Ross. ii. 754 = C. acantholepis.
nivea, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 395 = C. niveus.
squarrosa, 'Kostel. Select. Sem. Hort. Bot. Prag. (1838).'—Hab. ?
stenolepis, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 452 = C. stenolepis.
suaveolens, Rupr. Sert. Tiansch. 56 = C. suaveolens.
tianschanica, Rupr. l. c.—As. centr.

ALGA, [Tourn.] Lam. Fl. Fr. iii. 539 (1778) = **Zostera**, Linn. (Naiad.).
marina, Lam. l. c. = Z. marina.

ALGAROBIA, Benth. Pl. Hartw. 13 (1839) = **Prosopis**, Linn. (Legum.).
dulcis, Benth. l. c. = P. juliflora.
glandulosa, Torr. & Gray, Fl. N. Am. i. 399 = P. juliflora.
juliflora, Hcynh. Nom. ii. 18 = P. juliflora.

ALGERNONIA, Baill. in Ann. Sc. Nat. Sér. IV. ix. 198 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 339.
Tetraplandra, Baill. l. c. 200 (1858).
brasiliensis, *Baill. l. c.* 199.—Bras.
obovata, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 536.—Bras.

ALGUELAGUEN, Feuill. ex Adans. Fam. ii. 505 (1763) = **Sphaeele**, Benth. (Labiat.).

ALHAGI, Tourn. ex Adans. Fam. ii. 328 (1763); Desv. Journ. Bot. i. (1813) 120. t. 4. *LEGUMINOSAE*, Benth. & Hook. f. i. 512.
Manna, D. Don, Prod. Fl. Nep. 246 (1825).
camelorum, *Fisch. Hort. Gorenk.* ed. II. (1812) 72.—Oriens; As. centr.
graecorum, *Boiss. Diagn.* Ser. I. ix. 114.—Graecia.
karduchorum, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 555 = maurorum.
kirghisorum, Schrenk, Enum. Pl. Nov. 84 = camelorum.
mannifera, Desv. Journ. Bot. i. (1813) 120 = maurorum.
mannifera, Jaub. & Spach, Illustr. t. 401 = graecorum.
maurorum, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 397; *Desv. Journ.* i. (1813) 120. t. 4.—Oriens; Ind. or.; Syria.
napaulensium, DC. Prod. ii. 352 = maurorum.
persarum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 76 = camelorum.
Pseudalhagi, Desv. Journ. Bot. i. (1813) 120 = camelorum.
turcorum, Boiss. Diagn. Ser. I. ix. 113 = camelorum.

ALHAGIA, Reichb. Nom. 153 (1841) = praec.

ALIBERTIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 234. t. 21. *RUBIACEAE*, Benth. & Hook. f. ii. 81.
Cordiera, A. Rich. l. c. 222. t. 20 (1830).

ALIBERTIA :—
Garapatica, Karst. Fl. Colomb. i. 57. t. 28 (1858).
Gardeniola, Cham. in Linnaea, ix. (1834) 247.
Genipella, Rich. ex DC. Prod. iv. 443 (1830).
Melanopsidium, Poit. ex A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 234.
Scepseothamnus, Cham. in Linnaea, ix. (1834) 248.
Thieleodoxa, Cham. l. c. 251 (1834).
edulis, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1830) 234. t. 21.—Guiana.
hexagyna, *Karst. Fl. Colomb.* i. 57.—N. Granat.
jacquinioides, *Griseb. Cat. Pl. Cub.* 123.—Cuba.
Melloana, *Hook. f. in Trans. Linn. Soc.* xxviii. (1873) 520.—Bras.
utilis, A. Rich. in Mém. Soc. Nat. Hist. Par. v. (1830) t. 21 = edulis.

ALIBREXIA, Miers. in Hook. Lond. Journ. Bot. iv. (1845) 506 = **Dolia**, Lindl. (Convolv.).
brevifolia, Phil. in Linnaea, xxxiii. (1864-65) 209.—Chili.
Gayana, Gaudich. Bot. Voy. Bonite, Atl. t. 105.—Chili ?
incana, Phil. Fl. Atac. 45.—Chili.
revoluta, Miers, in Hook. Lond. Journ. Bot. iv. (1845) 508 = D. revoluta.
rupicola, Miers, l. c. 506 = D. tomentosa.
tomentosa, Miers, l. c. 508 = D. tomentosa.
villosa, Phil. Fl. Atac. 45.—Chili.

ALIBUM, Less. Syn. Comp. 152 (1832) = **Liabum**, Adans. (Compos.).
liaboides, Less. l. c. = L. lyratum.

ALICABON, Rafin. Sylva Tellur. 56 (1838) = **Physalis**, Linn. (Solanac.).
barbadense, Rafin. l. c. 57 = P. barbadense.
somniferum, Rafin. l. c. 56 = P. somniferum.

ALICASTRUM, P. Br. Hist. Jamaic. 372 (1756) = **Brosimum**, Sw. (Urticac.).

ALICOSTA, Dulac, Fl. Hautes-Pyr. 381 (1867) = **Bartsia**, Linn. (Scrophul.).
alpina, Dulac, l. c. = B. alpina.
spicata, Dulac, l. c. = B. aspera.

ALICTERES, Neck. Elem. ii. 304 (1790) = **Helicteres**, Linn. (Sterculiac.).

ALIFANA, Rafin. Sylva Tellur. 101 (1838) = **Brachyotum**, Triana (Melast.), etc.
canescens, Rafin. l. c. = Brachyotum canescens.
lutescens, Rafin. l. c. = Miconia lutescens.
montana, Rafin. l. c. = Conostegia montana.
striata, Rafin. l. c. = Brachyotum strigosum.

ALIFANUS, Pluk. ex Adans. Fam. ii. 234 (1763) = **Rhexia**, Linn. (Melast.).

ALINA, Adans. Fam. ii. 84 (1763).—Gen. dub.

ALIPENDULA, Neck. Elem. ii. 89 (1790) = **Spiraea**, Linn. (Rosac.).

ALIPSA, Hoffmgg. [Verz. Orchid. 20 (1840)] ex Linnaea, xvi. (1842) Litt. 228 = **Liparis**, Rich. (Orchid.).
foliosa, Hoffmgg. l. c. = L. reflexa ?

ALISETA, Rafin. New Fl. Am. ii. 22 (1836) = **Arnica** Linn. (Compos.).
plantaginea, Rafin. l. c. = A. alpina.

ALISMA, [Dill.] Linn. Syst. ed. I. (1735). *ALISMACEAE*, Benth. & Hook. f. iii. 1004.
Baldellia, Parl. Nov. Gen. Pl. Monocot. 57 (1854).
Caldesia, Parl. Fl. Ital. iii. 598 (1858).
Helanthium, Engelm. ex Benth. & Hook. f. Gen. iii. 1005 (1883).
Nectalisma, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 156.
acanthocarpum, *F. Muell. Fragm.* i. 23.—Austral.
alpestre, Coss. in Bull. Soc. Bot. Fr. xi. (1864) 333 = Echinodorus alpestris.
Ancile, Mart. ex Steud. Nom. ed. II. i. 49, nomen = Sagittaria guyanensis.
Andrieuxii, Hook. & Arn. Bot. Beech. Voy. 311 = = Echinodorus ellipticus.

ALISMA:—
angustifolium, Gilib. Fl. Lituan. ii. 224 = Echinodorus ranunculoides.
angustifolium, Hoppe, Taschenb. (1797) 13 = A. Plantago.
apetalum, Buch.-Ham. ex Wall. Cat. n. 4996 = oligococcum.
aphylla, Steud. Nom. ed. II. i. 49 = praec.
arcuatum, Micheli, in Bull. Soc. Bot. Fr. i. (1853) 312 = A. Plantago.
Berterii, Spreng. Syst. ii. 163 = Echinodorus rostratus.
Berteroanum, Balb. in Roem. & Schult. Syst. vii. 1605 = Echinodorus rostratus.
Bourgaei, Nym. Syll. 379 = Damasonium Bourgaei.
californicum, *Micheli, in A. DC. Monog. Phan.* iii. 34. —Calif.
calyophyllum, Wall. Cat. n. 4997 = parnassifolium.
canaliculatum, A. Br. & Bouché, Ind. Sem. Hort. Berol. (1867) 4 = A. Plantago.
cordifolium, Linn. Sp. Pl. 343 = Echinodorus macrophyllus, rostratus.
cordifolium, Thunb. Fl. Jap. 153 = A. Plantago.
cristatum, Wall. Cat. n. 4994 = Sagittaria guyanensis.
Damasonium, Desf. Fl. Atlant. i. 324 = Damasonium Bourgaei.
Damasonium, Linn. Sp. Pl. 343 = Damasonium stellatum.
Damasonium, Willd. Prod. Fl. Berol. 132 = parnassifolium.
difformifolium, *Steud. in Lechl. Berb. Am. Austr.* 53. —Chili.
diversifolium, Gilib. Fl. Lituan. ii. 223 = Elisma natans.
dubium, Willd. Prod. Fl. Berol. 132 = parnassifolium.
echinocarpum, Seub. in Mart. Fl. Bras. iii. 1. 105. t. 13. f. 3 = Sagittaria guyanensis.
ellipticum, Mart. in Roem. & Schult. Syst. vii. 1607 = Echinodorus ellipticus.
enneandrum, Hochst. ex Micheli, in DC. Monog. Phan. iii. 82 = Burnatia enneandra.
flavum, Linn. Sp. Pl. 343 = Limnocharis flava.
flavum, Thunb. Fl. Jap. 153 = A. Plantago.
floribundum, Seub. in Mart. Fl. Bras. iii. 1. 109 = Echinodorus grandiflorus.
Geyeri, Torr. in Nicollet, Rep. Hydrog. Mississip. (1843) 162 = A. Plantago.
glandulosum, Thw. Enum. Pl. Zeyl. 332 = oligococcum.
gramineum, C. C. Gmel. Fl. Bad. iv. 256 = A. Plantago.
graminifolium, Ehrh. ex Ledeb. Fl. Ross. iv. 40 = A. Plantago.
grandiflorum, Cham. & Schlecht. in Linnaea, ii. (1827) 152 = Echinodorus grandiflorus.
Hamiltonianum, Wall. Cat. n. 4993 = Sagittaria guyanensis.
heterophyllum, Schur, Enum. Pl. Transs. 880, Index = A. Plantago.
humile, Rich. ex Kunth, Enum. Pl. iii. 154 = Echinodorus humilis.
intermedium, *Griff. ex Voigt, Hort. Suburb. Calc.* 680.—Ind. or.
intermedium, Mart. in Roem. & Schult. Syst. vii. 1609 = Echinodorus Martii.
Kotschyi, Hochst. in Flora, xxvi. (1843) 492 = Limnophyton obtusifolium.
lanceolatum, With. Bot. Arr. Brit. Pl. ed. III. ii. 362 = A. Plantago.
Lappula, Buch.-Ham. ex Kunth, Enum. Pl. iii. 162 = Sagittaria Lappula.
latifolium, Gilib. Fl. Lituan. ii. 222 = A. Plantago.
Loeselii, Eichw. Skizze, 127 = A. Plantago.
macrophyllum, Kunth, Enum. Pl. iii. 151 = Echinodorus macrophyllus.
majus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 216 = A. Plantago.
minus, Spreng. Syst. ii. 163 = Damasonium australe.
natans, Linn. Sp. Pl. 343 = Elisma natans.
natans, Pollich, Hist. Pl. Palat. iii. 319 = A. Plantago.
nathpurense, Steud. Nom. ed. II. i. 49 = Echinodorus ellipticus.
nymphaeifolium, *Griseb. Cat. Pl. Cub.* 218.—Cuba.
obtusifolium, Thw. Enum. Pl. Zeyl. 332 = Limnophyton obtusifolium.
odoratum, Rafin. Fl. Ludov. 17 = A. Plantago.
oligococcum, *F. Muell. Fragm.* i. 23.—Austral.
paleaefolium, Kunth, Enum. Pl. iii. 152 = Echinodorus ellipticus.

ALISMA:—
paniculatum, Stokes, Bot. Mat. Med. ii. 334 = A. Plantago.
parnassifolium, Buch.-Ham. ex Wall. Cat. n. 4997 = calophyllum.
parnassifolium, *Bassi, ex Linn. Syst.* ed. XII. ii. 230.—Europ.; Ind. or.; Afr. trop.; Austral.
parviflorum, Pursh, Fl. Am. Sept. i. 252 = A. Plantago.
plantagineum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Plantago.
Plantago, *Linn. Sp. Pl.* 342 (A. Plantago-aquatica). —Reg. temp. bor.; Austral.
polyspermum, Nym. Syll. 379 = Damasonium polyspermum.
pubescens, Buch.-Ham. ex Wall. Cat. n. 4995 = Sagittaria guyanensis.
pubescens, Mart. in Roem. & Schult. Syst. vii. 1607 = Echinodorus ellipticus.
ranunculoides, All. Fl. Pedem. i. 234 = A. Plantago.
ranunculoides, Linn. Sp. Pl. 343 = Echinodorus ranunculoides.
ranunculoides, Willd. Prod. Fl. Berol. 133 = Elisma natans.
reniforme, D. Don, Prod. Fl. Nep. 22 = parnassifolium.
repens, Lam. Encyc. ii. 515 = Echinodorus ranunculoides.
roseum, Rafin. in Med. Repos. New York, v. (1808) 354 = A. Plantago.
rostratum, Nutt. in Trans. Am. Phil. Soc. v. (1837) 159 = Echinodorus rostratus.
Sagittaria, Stokes, Bot. Mat. Med. ii. 335 = Sagittaria sagittifolia.
sagittifolium, Willd. Sp. Pl. ii. 277 = Limnophyton obtusifolium.
Sprengelii, Rich. ex Kunth, Enum. Pl. iii. 154 = Echinodorus rostratus.
stellatum, Buch.-Ham. ex Wall. Cat. n. 4993 = Sagittaria guyanensis.
stellatum, Lam. Fl. Fr. iii. 77 = Damasonium stellatum.
subalatum, Mart. in Roem. & Schult. Syst. vii. 1609 = Echinodorus subalatus.
subcordatum, Rafin. in Med. Repos. New York, v. (1808) 362 = A. Plantago.
subulatum, Linn. Sp. Pl. 343 = Sagittaria natans.
subulatum, Hill, Veg. Syst. x. 33 = praec.?
taeniaefolium, *Steud. in Lechl. Berb. Am. Austr.* 53.—Chili.
tenellum, Mart. in Roem. & Schult. Syst. vii. 1600 = Echinodorus tenellus.
trinerve, *Link, ex Steud. Nom.* ed. II. i. 49.—Lusitan.
triviale, Pursh, Fl. Am. Sept. i. 252 = A. Plantago.
verticillatum, Dulac, Fl. Hautes-Pyr. 45 = A. Plantago.
virgatum, Hook. & Arn. Bot. Beech. Voy. 311 = Echinodorus virgatus.

ALISMORCHIS, Thou. Orch. Afr. t. 35 (1822) = **Calanthe**, R. Br. (Orchid.).
Centrosis, Steud. Nom. ed. II. i. 49 = C. sylvatica.

ALISSON, Vill. Prosp. 38 (1779) = **Alyssum**, Linn. (Crucif.).

ALISSUM, Neck. Delic. Gallo-Belg. i. 276 (1768) = praec.

ALITUBUS, Dulac, Fl. Hautes-Pyr. 499 (1867) = **Achillea**, Linn. (Compos.).
Millefolium, Dulac, l. c. 500 = Ach. Millefolium.
pyrenaicus, Dulac, l. c. = Ach. Ptarmica.
tomentosus, Dulac, l. c. = Ach. tomentosa.

ALIX, Comm. ex DC. Prod. v. 318 (1836) = **Psiadia**, Jacq. (Compos.).
salsifolia, Comm. ex DC. l. c. 319 = P. retusa.

ALKANNA, Tausch, in Flora, vii. (1824) 234.
BORAGINEAE, Benth. & Hook. f. ii. 857.
BAPHORHIZA, Link, Handb. ii. 578 (1829).
CAMPTOCARPUS, C. Koch, in Linnaea, xvii. (1843) 304.
CAMPYLOCARYUM, DC. Prod. x. 97 in syn. (1846).
alba, Boiss. Voy. Espagne, 747 = Nonea alba.
amplexicaulis, A. DC. in DC. Prod. x. 101 = Kotschyana.
areolata, *Boiss. Diagn.* Ser. I. iv. 45.—As. Min.
Aucheriana, A. DC. in DC. Prod. x. 102 = syriaca, rostellata.
azurea, Boiss. Pl. Or. Dec. ii. 7 = areolata.
baeotica, DC. Prod. x. 98 = graeca.
bicolor, Boiss. Pl. Or. Dec. ii. 7 = Kotschyana.
brachysolen, Boiss. l. c. = Kotschyana.

ALKANNA:—
bracteosa, *Boiss. Diagn.* Ser. I. xi. 118.—Persia.
calliensis, Heldr. ex Nym. Consp. 517 = graeca.
cappadocica, Boiss. & Bal. Diagn. Ser. II. vi. 123 = megacarpa.
controversa. *A. DC. in DC. Prod.* x. 103.—Hab.?
cordifolia, *C. Koch, in Linnaea*, xxii. (1849) 640.—Pontus.
filaginoides, DC. Prod. x. 98 = Pinardi.
frigida, *Boiss. Diagn.* Ser. I. vii. 32.—Persia.
galilaea, *Boiss. l. c.* Ser. I. xi. 118.—Palaestin.
graeca, *Boiss. & Sprun. l. c.* Ser. I. iv. 47.—Graecia.
hirsutissima, *A. DC. in DC. Prod.* x. 101.—Oriens.
incana, *Boiss. Diagn.* Ser. I. iv. 45.—As. Min.
Kotschyana, *DC. Prod.* x. 98.—Oriens.
Lehmanni, A. DC. in DC. l. c. 588 = tinctoria.
lutea, *A. DC. l. c.* 102.—Europ.
macrophylla, *Boiss. & Heldr. Diagn.* Ser. I. xi. 120.—As. Min.; Syria.
macrosiphon, *Boiss. & Heldr. l. c.* 120.—Graecia.
Mathioli, Tausch, in Flora, vii. (1824) 234 = tinctoria.
megacarpa, *A. DC. in DC. Prod.* x. 101.—Oriens.
noneiformis, Griseb. Spicil. Fl. Rumel. ii. 90 = A. Pulmonaria.
orientalis, *Boiss. Diagn.* Ser. I. iv. 46.—Oriens.
Pinardi, *Boiss. l. c.* 48.—As. Min.
primuliflora, *Griseb. Spicil. Fl. Rumel.* ii. 89.—Thracia.
Pulmonaria, *Griseb. l. c.* 90.—Macedon.
rostellata, *Boiss. Diagn.* Ser. I. xi. 119.—Taurus.
Sartoriana, *Boiss. & Heldr. l. c.* Ser. II. iii. 134.—Graecia.
scardica, *Griseb. Spicil. Fl. Rumel.* ii. 91.—Macedon.
Sieberi, *A. DC. in DC. Prod.* x. 99.—Creta.
strigosa, *Boiss. & Hohenh. in Boiss. Diagn.* Ser. I. iv. 46.—Syria.
syriaca, *Boiss. & Hohenh. Pl. Or. Nov. Dec.* ii. 7.—Syria.
tinctoria, DC. Prod. x. 99 = Sieberi.
tinctoria, *Tausch, in Flora*, vii. (1824) 234.—Oriens.
tomentosa, Griseb. Spicil. Fl. Rumel. ii. 88 in nota = Onosma mollis.
tubulosa, *Boiss. Diagn.* Ser. I. iv. 48.—As. Min.
versicolor, *Orph. in Bull. Congr. Bot. Pétersb.* 1869 (1870) 138.—Graecia.
viscidula, *Boiss. Diagn.* Ser. I. xi. 118.—Cappadoc.
Wagneri, *Bartl. ex Tchihat. Asie Min. Bot.* ii. 107, nomen.—As. Min.

ALKEKENGI, Tourn. ex Hall. Enum. Stirp. Helv. ii. 508 (1742) = **Physalis**, Linn. (Solanac.).
fissum, Moench, Meth. 472 = P. viscosa.
officinarum, Moench, Meth. Suppl. 177 = P. Alkekengi.
procumbens, Moench, Meth. 473 = P. pubescens.
pubescens, Moench, l. c. = P. peruviana.
villosa, Moench, Meth. Suppl. 178 = P. pubescens.

ALKIBIAS, Rafin. Fl. Tellur. iv. 119 (1836) (Compos.).
hastata, Rafin. l. c. (= *Linosyris hastata*).
mucronata, Rafin. l. c. = Iphione juniperifolia.
ovata, Rafin. l. c. = Chrysocoma ovata.

ALLAEANTHUS, Thw. in Hook. Kew Journ. vi. (1854) 302. t. 9. *URTICACEAE*, Benth. & Hook. f. iii. 361.
luzonicus, *Benth. & Hook. f. Gen.* iii. 361.—Ins. Phil.
zeylanicus, *Thw. in Hook. Kew Journ.* vi. (1854) 302. —Zeylan.

ALLAEOPHANIA, Thw. Enum. Pl. Zeyl. 147 (1859). *RUBIACEAE*, Benth. & Hook. f. ii. 129.
Arnottii, *Hook. f. Fl. Brit. Ind.* iii. 48.—Zeylan.
decipiens, *Thw. Enum. Pl. Zeyl.* 147.—Zeylan.

ALLAGANTHERA, Mart. Hort. Erlang. 69 (1814) = **Alternanthera**, Forsk. (Amarant.).
Forskohlei, Mart. l. c. = Alt. sessilis.

ALLAGAS, Rafin. Fl. Tellur. iv. 53. (1836) = **Halenia**, Borckh. (Gentianac.).

ALLAGOPAPPUS, Cass. in Dict. Sc. Nat. lvi. 21. (1828). *COMPOSITAE*, Benth. & Hook. f. ii. 333.
chrysopsioides, Sch. Bip. in. Schweinf. Beitr. Fl. Aethiop. 153 = Pulicaria petiolaris.
dichotomus, *Cass. in Dict. Sc. Nat.* lvi. 22.—Ins. Canar.
viscosissimus, *C. Bolle, in Bonplandia*, vii. (1859) 293.—Ins. Canar.

ALLAGOPTERA, Nees, in Neuwied, Reise nach Bras. ii. 335 (1821) = **Diplothemium**, Mart. (Palm.).
pumila, Nees, l. c. = D. maritimum.

ALLAGOSPERMA, M. Roem. Syn. Pepon. 68 (1846) = **Cayaponia**, Manso (Cucurb.).
Taynia, M. Roem. l. c. = C. Tayuya.

ALLAGOSTACHYUM, Nees, ex Steud. Nom. ed. II. i. 50 (1840) = **Poa**, Linn. (Gramin.).
lasiochloa, Nees, l. c. = Poa alternans.

ALLAMANDA, Linn. Mant. ii. 146. (1771). *APOCYNACEAE*, Benth. & Hook. f. ii. 690.
ORELIA, Aubl. Pl. Gui. i. 270. t. 106 (1775).
angustifolia, *Pohl, Pl. Bras.* i. 73. t. 59.—Bras.
Aubletii, Pohl, l. c. 75 = cathartica.
Blanchetii, *A. DC. in DC. Prod.* viii. 319.—Bras.
brasiliensis, Schott, ex Pohl, Pl. Bras. i. 73 = Schottii.
cathartica, *Linn. Mant.* ii. 214.—Am. trop.
cathartica, Schrad. in Goett. Gelehrt. Anz. (1821) 707 = Schottii.
Doniana, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 11.—Bras.
grandiflora, Lam. Encyc. iv. 601 = cathartica.
Hendersoni, *Bull, ex Dombr. Flor. Mag.* v. (1866) t. 263.—Guiana.
latifolia, Presl, Bot. Bemerk. 103 = cathartica.
Linnei, Pohl, Pl. Bras. i. 74 = cathartica.
Martii, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 12.—Bras.
neriifolia, *Hook. Bot. Mag.* t. 4594.—Hab. ?
nobilis, *T. Moore, in Gard. Chron.* (1868) 180, 918.—Bras.
ocnotheraefolia, *Pohl, Pl. Bras.* i. 71. t. 57.—Bras.
parviflora, *Presl, Bot. Bemerk.* 103.—Ins. Martin.
polyantha, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 12.—Bras.
puberula, *A. DC. in DC. Prod.* viii. 319.—Bras.
Schottii, Hook. Bot. Mag. t. 4351 = cathartica.
Schottii, *Pohl, Pl. Bras.* i. 73. t. 58.—Bras.
setulosa, *Mig. in Linnaea*, xviii. (1844) 743.—Surinam.
thevetifolia, *Muell. Arg. in Linnaea*, xxx. (1859–60) 388.—Bras.
verrucosa, Gardn. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 14 = puberula.
verticillata, *Desf. Tabl. Bot. Par.* ed. II. 93.—Hab. ?
violacea, Gardn. & Field. Sert. t. 41 = Blanchetii.
Wardleyana, Lebas, in Rev. Hortic. (1877) 20 = Hendersoni.

ALLANBLACKIA, Oliver, in Benth. & Hook. f. Gen. i. 980 (1867). *GUTTIFERAE*, Benth. & Hook. f. i. 980.
floribunda, Oliver, in Journ. Linn. Soc. x. (1869) 43.—Afr. trop.

ALLANIA, Benth. in Hook. Lond. Journ. Bot. ii. (1840) 91 = **Aldina**, Endl. (Legum.).
insignis, Benth. l. c.—Guiana.

ALLANIA, Meissn. Gen. 401 (304), (1842) = **Alania**, Endl. (Liliac.).

ALLANTOMA, Miers, in Trans. Linn. Soc. xxx. (1874) 291. *MYRTACEAE*.
aulacocarpa, *Miers, l. c.* 296.—Bras.
Burchelliana, *Miers, l. c.* 294.—Bras.
corbula, *Miers, l. c.* 297.—Bras.
cylindrica, *Miers, l. c.* 294.—Bras.
dictyocarpa, *Miers, l. c.* 297.—Bras.
fagifolia, *Miers, l. c.* 298.—Guiana.
lineata, *Miers, l. c.* 297.—Bras.
macrocarpa, *Miers, l. c.* 295.—Bras.
multiflora, *Miers, l. c.* 291.—Guiana.
scutellata, *Miers, l. c.* 296.—Bras.
subramosa, *Miers, l. c.* 292.—Guiana.
torulosa, *Miers, l. c.* 293.—Bras.

ALLARDIA, Decne. in Jacquem. Voy. Bot. 87. t. 95, 96 (1841 ?). *COMPOSITAE*, Benth. & Hook. f. ii. 424.
WALDHEIMIA, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1842) 125.
glabra, *Decne. l. c.* 88. t. 96.—Tibet.
Huegelii, Sch. Bip. in Pollichia, xx–xxi. (1863) 442 = glabra.
incana, *Klatt, in Sitzb. Akad. Muench.* (1873) 88.—Tibet.

ALLARDIA :—
lanata, Kurz, ex C. B. Clarke, Comp. Ind. 144 = tomentosa.
nivea, *Hook. f. & Thoms. ex C. B. Clarke, l. c.* 145.—Tibet. occ.
Stoliczkai, *C. B. Clarke, Comp. Ind.* 145.—Tibet. occ.
tomentosa, *Decne. in Jacquem. Voy. Bot.* 87. t. 95.—Tibet. occ.
tridactylites, Sch. Bip. in Pollichia, xx–xxi. (1863) 442 = glabra.
vestita, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 145.—Reg. Himal.

ALLARDTIA, A. Dietr. in Otto & Dietr. Allg. Gartenz. xx. (1852) 242 = **Tillandsia**, Linn. (Bromel.).
cyanea, A. Dietr. l. c. = T. cyanea.
Potockii, Antoine, in Oestr. Bot. Zeitschr. xxviii. (1878) 56 = T. utriculata.

ALLASIA, Lour. Fl. Cochinch. 84 (1790) Inc. sed.; cf. Benth. & Hook. f. i. 820.
Jobini, *Manso, Enum.* 36.—Bras.
Payos, *Lour. Fl. Cochinch.* 85.—Afr. or.

ALLAZIA, Manso, Enum. 36. (1836) = praec.

ALLELOTHECA, Steud. Syn. Pl. Gram. 117 (1855) = **Lophatherum**, Brongn. (Gramin.).
Urvillei, Steud. l. c. = L. gracile.

ALLEMANIA, Endl. Gen. 304 in nota (1837) = **Allmania**, R. Br. (Amarant.).

ALLENDEA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. i. 10 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 437.
lanceolata, *La Llave, l. c.*—Mexic.

ALLETOTHECA, Benth. & Hook, f. iii. 1191 (1883) sphalm. = ALLELOTHECA, Steud. = **Lophatherum**, Brongn. (Gramin.).

ALLIARIA, Marsh. ex Rupp. Fl. Jen. ed. Hall. 75 (1745); DC. Syst. Veg. ii. 488 = **Sisymbrium**, Linn. (Crucif.).
brachycarpa, Bieb. Fl. Taur. Cauc. Suppl. 445 = Erysimum brachycarpon.
Fuchsii, Rupr. Fl. Cauc. 85 = S. Alliaria.
Mathioli, Rupr. l. c. = S. Alliaria.
officinalis, Andrz. ex DC. Syst. Veg. ii. 489 = S. Alliaria.

ALLIBERTIA, Marion, in Rev. Hort. Bouch. Rhone, Nov. (1882); ex Baker, in Gard. Chron. N. S. xix. (1883) 176 = **Agave**, Linn. (Amaryll.).
intermedia, Marion, ex Baker, l. c. = A. Alibertii.

ALLINUM, Neck. Elem. i. 179 (1790) = **Selinum**, Linn. (Umbell.).

ALLIONIA, Loefl. Iter, 181 (1758); Linn. Syst. ed. X. 890 (1759). *NYCTAGINEAE*, Benth. & Hook. f. iii. 4.
WEDELIA, Loefl. Iter, 180 (1758).
aggregata, Spreng. Syst. i. 384 = Oxybaphus aggregatus.
albida, Walt. Fl. Carol. 84 = Oxybaphus albidus.
Cervantesii, Steud. Nom. ed. II. i. 50 = Oxybaphus Cervantesii.
cucullata, Mey. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 55 = Oxybaphus cucullatus.
decumbens, Spreng. Syst. i. 384 = Oxybaphus decumbens.
hirsuta, Pursh, Fl. Am. Sept. ii. 728 = Oxybaphus hirsutus.
incarnata, *Linn. Syst.* ed. X. 890.—Am. bor.; Ind. occ.; Am. austr.
linearis, Pursh, l. c. = Oxybaphus angustifolius.
malacoides, Benth. Bot. Voy. Sulph. 44 = incarnata.
mendocina, *Phil. Sert. Mendoc. Alt.* 41.—Chili.
nyctaginea, Michx. Fl. Bor. Am. i. 100 = Oxybaphus nyctagineus.
ovata, Pursh, Fl. Am. Sept. i. 97 = Oxybaphus floribundus.
violacea, Linn. Syst. ed. X. 890 = Oxybaphus violaceus.

ALLIUM, [Tourn.] Linn. Syst. ed. I. (1735). *LILIACEAE*, Benth. & Hook. f. iii. 802.
ANGUINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 160.

ALLIUM :—
ASCALONICUM, Renault, Fl. Dép. Orne, 33 (1800).
CEPA, Tourn. ex Linn. Gen. ed. I. 103 (1737).
CODONOPRASUM, Reichb. Consp. 66 (1828).
GEBOSCON, Rafin. Cat. 14 (1824).
MOENCHIA, Medic. in Act. Acad. Palat. vi. Phys. (1790) 343.
MOLIUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159.
MOLY, Boerh. ex Moench, Meth. 286 (1794).
NECTAROSCORDUM, Lindl. Bot. Reg. t. 1912 (1836).
OPHIOSCORODON, Wallr. Sched. Crit. 129 (1822).
PORRUM, Tourn. ex Linn. Syst. ed. I. (1735).
RHIZIRIDEUM, Fourr. l. c. 160.
SATURNIA, Maratti, Diss. Romul. et Sat. 18. t. 2. (1772).
SCHOENOPRASUM, H. B. & K. Nov. Gen. et Sp. i. 227 (1815).
SCORODON, Fourr. l. c. 160.
TRIGONEA, Parlat. in Occhio, 161 (1839).
AGLITHEIS, ENDOTIS, GETUONIS, GYNODON, KALABOTIS, KEPA, KROMON, LONCOSTEMON, MALIGA, PANSTENUM, PLEXISTENA, STELMESUS, STEMODOXIS, Rafin. Fl. Tellur. ii. 17–21 (1836).
PERILOBA, PRASKOINON, Rafin. l. c. iv. 87, 29 (1836).
BERENICE, BRISEIS, BUTOMISSA, CAMARILLA, CANIDIA, HEXONYCHIA, HYLOGETON, JULUS, MOLYZA, PHYLLODOLON, RAPHIONE, SCHOENISSA, XYLORHIZA, Salisb. Gen. Pl. Fragm. 88–92 (1866).
achaium, *Boiss. & Orph. ex Boiss. Fl. Orient.* v. 259.—Graecia.
acre, *J. & C. Presl, Delic. Prag.* 147.—Sicilia.
acuminatum, *Hook. Fl. Bor. Am.* ii. 184. t. 196.—Am. bor.
acutangulum, Ambros. Fl. Tyr. Austr. i. 565 = fallax.
acutangulum, Reichb. Ic. Crit. viii. t. 734 = montanum.
acutangulum, Schrad. Sem. Hort. Goett. 1808 = angulosum.
acutiflorum, Bové, ex Regel, in Act. Hort. Petrop. iii. II. (1875) 192 = paniculatum.
acutiflorum, Loisel. in Desv. Journ. Bot. ii. (1809) 279 = multiflorum.
acutum, Spreng. Pugill. i. 28 = A. Schoenoprasum.
adscendens, Kunth, Enum. Pl. iv. 385 = A. Ampeloprasum.
aestivum, Tineo, Pl. Rar. Sicil. 39 = sphaerocephalum.
affine, Boiss. & Heldr. Diagn. Ser. II. iv. 114 = vineale.
affine, Ledeb. Fl. Ross. iv. 166 = margaritaceum.
africanum, F. G. Dietr. Gartenlex. Nachtr. i. 160 = odoratissimum.
afrum, Kunth, Enum. Pl. iv. 455 = Ornithogalum afrum.
Aitchisoni, *Boiss. Fl. Orient.* v. 248.—Afghan.
Akaka, *Gmel. ex Roem. & Schult. Syst.* vii. 1132.—Sibir.
Akaka, Regel, in Act. Hort. Petrop. iii. (1875) 243 = karataviense.
alataviense, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) I. 448.—Turkestan.
alatum, Schreb. ex Roth, in Roem. Arch. i. III. (1798) 40 = cernuum.
Alberti, *Regel, in Act. Hort. Petrop.* v. (1877) 632.—Turkestan.
albescens, Guss. Enum. Pl. Inarim. 338 = A. Ampeloprasum.
albidum, Bieb. Fl. Taur. Cauc. iii. Suppl. 260 = flavescens.
albidum, *Fisch. Cat. Gorenk.* (1812) 10.—Oriens; Sibir.
albidum, J. & C. Presl, Delic. Prag. 146 = paniculatum.
album, Hort. ex [Delar. in] Red. Lil. v. 300 = ciliatum.
album, Santi, Viaggi, 352. t. 7 = neapolitanum.
album, Spreng. Syst. ii. 38 = pendulinum.
Alexeianum, *Regel, in Act. Hort. Petrop.* iii. (1875) 244.—Turkestan.
alibile, *A. Rich. Tent. Fl. Abyss.* ii. 330.—Abyss.
alpicolum, *Verl. in Bull. Soc. Stat. Isère*, xiv. (1872) 397.—Gallia.
alpinum, Regel, in Act. Hort. Petrop. iii. II. (1875) 181 = narcissiflorum.
alpinum, Hegetschw. Fl. Helv. 322 = A. Schoenoprasum.
altaicum, Pall. Reise, ii. 737 = fistulosum.
altissimum, *Regel, in Act. Hort. Petrop.* viii. (1884) 666.—Bokhara.
Amansii, Bor. Fl. Cent. Fr. ed. III. ii. 631. in obs. = paniculatum.
ambiguum, DC. Fl. Fr. iii. 320 = ochroleucum.
ambiguum, Sibth. & Sm. Fl. Graec. t. 327 = roseum.
amblyopetalum, Link, in Linnaea, ix. (1834) 139 = neapolitanum.

ALLIUM:—
amblyophyllum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 510.—Soongaria.
amethystinum, Tausch, in Syll. Ratisb. ii. 256 = A. Ampeloprasum.
ammophilum, Heuff. in Flora, xxviii. (1845) 241; et Verh. Zool.-Bot. Ges. Wien, viii. (1858) 210 = angulosum.
amoenum, G. Don, in Mem. Wern. Soc. vi. (1827) 75 = roseum.
Ampeloprasum, Boiss. Voy. Espagne (ii. 615) = polyanthum, baeticum.
Ampeloprasum, *Linn. Sp. Pl.* 295.—Europ.; Oriens.
Ampeloprasum, Thunb. Prod. Pl. Cap. 65 = rotundum.
amphibolum, *Ledeb. Fl. Alt.* ii. 5. t. 357.—Sibir.
amplectens, Torr. in Pacif. Rail. Rep. iv. 148 = attenuifolium.
anceps, *Kellogg, in Proc. Calif. Ac.* ii. (1863) 109. t. 32.—Calif.
Andersonii, G. Don, in Mem. Wern. Soc. vi. (1827) 59 = senescens.
andicolum, *Regel, in Act. Hort. Petrop.* iii. (1875) 211. —Peruv.
angolense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 262.—Afr. trop.
angulare, *Pall. Reise*, iii. 299.—Sibir. baic.
angulatum, Pall. l. c. 408. = praec.
angulosum, All. Fl. Pedem. ii. 158 = albidum.
angulosum, DC. Fl. Fr. iii. 222 = fallax.
angulosum, Krock. Fl. Sil. i. 524 = acutangulum, fallax, senescens.
angulosum, *Linn. Sp. Pl.* 300.—Sibir.
angulosum, Lour. Fl. Cochinch. 203 = uliginosum.
angulosum, Pursh, Fl. Am. Sept. i. 223 = reticulatum.
angustum, *G. Don, in Mem. Wern. Soc.* vi. (1827) 33. —Sibir.
anisopodium, Ledeb. Fl. Ross. iv. 183 = tenuissimum.
apenninum, Huet, ex Nym. Consp. 740 = paniculatum.
appendiculatum, Ram. ex Pers. Syn. i. 356 = suaveolens.
approximatum, Gren. & Godr. Fl. Fr. iii. 200 = sphaerocephalum.
arenarium, Linn. Sp. Pl. 297 = A. Scorodoprasum.
arenarium, Sadler, Fl. Pesth. ex Reichb. Fl. Germ. Excurs. 40 = sativum.
arenarium, [Wahlenb. in] Svensk. Bot. t. 691 = vineale.
arenarium, Thunb. [Fl. Jap. 132] = japonicum.
armenum, Boiss. & Kotschy, ex Boiss. Fl. Orient. v. 254 = flavum.
armerioides, *Boiss. Diagn.* Ser. I. vii. 116.—Assyria.
arvense, Guss.-Ind. Sem. (1825) = sphaerocephalum.
ascalonicum, *Linn. Amoen. Acad.* iv. 454.—Ubique cult.
ascalonium, Bory & Chaub. Fl. Pelop. 21 = gomphrenoides.
ascendens, Tenore, Syll. 164 = A. Ampeloprasum.
Aschersonianum, *Barbey, Herbor. au Levant*, 163. t. 4. —Oriens.
asperum, G. Don, in Mem. Wern. Soc. vi. (1827) 42 = carinatum.
atropurpureum, Waldst. & Kit. Ic. Pl. Rar. Hung. i. 16. t. 17 = stenopetalum.
atrorubens, *S. Wats. in Bot. King, Exp.* 352.—Calif.
atrosanguineum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 508.—As. med.
atroviolaceum, Boiss. Diagn. Ser. I. vii. 112 = A. Ampeloprasum.
atroviolaceum, Hornem. ex Steud. Nom. ed. II. i. 50 = sphaeroccphalum.
attenuatum, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 110. f. 33.—Calif.
attenuifolium, *Kellogg, l. c.* 110. f. 34.—Calif.
Aucheri, *Boiss. Diagn.* Ser. I. vii. 116.—Persia.
aureum, Lam. Fl. Fr. iii. 260 = A. Moly.
auriculatum, *Kunth, Enum. Pl.* iv. 418.—Ind. or.
australe, Guss. ex Kunth, l. c. 404 = paniculatum.
azureum, Bunge, in Mém. Sav. Étr. Acad. Pétersb. vii. (1851) 510; Reliq. Lehm. 334. n. 1366 = urceolatum.
azureum, Ledeb. Fl. Alt. ii. 13 = caeruleum.
Babingtonii, Borrer, in Engl. Bot. t. 2906 = A. Scordoprasum.
bachtiaricum, Boiss. Fl. Orient. v. 285 = bakhtiaricum.
Backhousianum, *Regel, Gartenfl.* (1885) 213.—Reg. Himal.
baeticum, *Boiss. Diagn.* Ser. I. vii. 113.—Hispan.
baicalense, Willd. Enum. Hort. Berol. 360 = senescens.
Bakeri, *Regel, in Act. Hort. Petrop.* iii. (1875) 141.—China; Japon.
bakhtiaricum, *Regel, l. c.* 222.—Persia.

ALLIUM:—
Balansae, *Boiss. Fl. Orient.* v. 268.—Pontus.
Bauerianum, *Baker, in Gard. Chron.* (1874) II. 131.—Aegypt.; Turcia?
Baumannianum, C. Koch, in Linnaea, xxii. (1849) 241 = rotundum.
Beckerianum, Regel, Ind. Sem. Hort. Petrop. (1860) 30 = tataricum.
Bertoloni, De Not. Rep. Fl. Lig. 414 = A. Ampeloprasum.
bidentatum, Fisch. ex Regel, in Act. Hort. Petrop. iii. (1875) 157 = tenuissimum.
Bidwelliae, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 231.—Am. bor.
bifolium, *Willd. ex Schult. f. Syst. Veg.* vii. 1117 = decipiens.
Bigelovii, *S. Wats. in Bot. King, Exped.* v. 487. t. 38. f. 8 & 9.—N. Mexic.
bisceptrum, *S. Wats. l. c.* 351 & 487. t. 37. f. 1–3.—Calif.
bisulcum, [Delar. in] Red. Lil. t. 286 = A. Stellerianum.
blandum, *Wall. Pl. As. Rar.* iii. 38. t. 260.—Tibet. occ.
Bodeanum, *Regel, in Act. Hort. Petrop.* iii. (1875) 238. —Persia.
bogdoicolum, *Regel, l. c.* vi. (1879) 530.—Turkestan.
Boissieri, Haussk. ex Regel, l. c. iii. II. (1875) 205 = frigidum.
Boissieri, Regel, l. c. 75 = Lehmannianum.
Bolanderi, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 229.—Calif.
bonariensis, *Griseb. in Goett. Abh.* xxiv. (1879) 319.—Reg. Argent.
borbonicum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 218.—Ins. Borb.
Borbasii, *Kerner, in Oestr. Bot. Zeitschr.* xxix. (1879) 39.—Europ.
Borszczowi, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 74.—Turkestan.
Boryanum, Kunth, Enum. Pl. iv. 411 = paniculatum.
bosniacum, Kumm. & Sendt. in Flora, xxxii. (1849) 762 = sphaerocephalum.
Bouddae, *Debeaux, in Actes Linn. Soc. Bord.* xxxii. (1878) 24; *Fl. du Tché-fou*, 137.—China.
brachyodon, Boiss. Diagn. Scr. I. vii. 117 = strictum.
brachystemum, [Delar. in] Red. Lil. vii. t. 374 = subhirsutum.
brahuicum, *Boiss. Fl. Orient.* v. 278.—Beluchist.
Brandegei, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 380.—Colorado.
brevicaule, Boiss. & Bal. Diagn. Ser. II. iv. 119 = paniculatum.
brevicuspis, *Boiss. & Buhse, l. c.* 114.—Persia.
brevipes, Ledeb. Fl. Ross. iv. 165 = Aucheri.
breviscapum, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 14.—Persia.
brevistylum, *S. Wats. in Bot. King, Exp.* 350.—Am. bor. occ.
Breweri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 233. —Calif.
Broteri, Kunth, Enum. Pl. iv. 454 = A. Schoenoprasum.
bucharicum, *Regel, in Act. Hort. Petrop.* viii. (1884) 660.—Bokhara.
Buhseanum, Regel, l. c. iii. II. (1875) 81 = A. Schoenoprasum.
Bungei, *Boiss. Fl. Orient.* v. 267.—Persia.
byzantinum, C. Koch, in Linnaea, xxii. (1849) 240 = A. Ampeloprasum.
caerulescens, G. Don, in Mem. Wern. Soc. vi. (1832) 34 = caeruleum.
caerulescens, *Boiss. Diagn.* Ser. II. iv. 115.—Armenia.
caeruleum, *Pall. Reise*, ii. 737.—Sibir.
caeruleum, Stschégléew, in Bull. Soc. Nat. Mosc. (1854) I. 202 = urceolatum.
caeruleum, Wall. Cat. n. 5070 = Wallichii.
caesium, *Schrenk, in Bull. Phys. Math. Acad. Pétersb.* ii. (1844) 113.—Turkestan.
caespitosum, *Siev. in Pall. Neu. Nord. Beitr.* iii. 304. —Soongaria.
calcareum, *Reut. in Compte-rend. Soc. Hallér.* (1855–56) 127.—Helvet.
calcareum, Wallr. Beitr. Fl. Hercyn. 196 = angulosum.
callidictyon, Regel, in Act. Hort. Petrop. iii. II. (1875) 74 = Borszczowi.
callidyction, *C. A. Mey. ex Kunth, Enum. Pl.* iv. 413. —Oriens.

ALLIUM:—
callimischon, Link, in Linnaea, ix. (1834) 140 = paniculatum.
callistemon, Webb, ex Regel, in Act. Hort. Petrop. iii. (1875) 189 = flavum.
calyptratum, *Boiss. Diagn.* Ser. I. xiii. 30.—Syria.
Cambiasii, De Not. Rep. Fl. Lig. 412 = rotundum.
campanulaeflorum, *Gey. ex Hook. Kew Journ.* vii. (1855) 375.—Am. bor.
campanulatum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 231.—Am. bor.
campestre, Schleich. ex Steud. Nom. ed. II. i. 51 = vineale.
canadense, Hort. ex Schult. f. Syst. vii. 1007 = Purshii.
canadense, Herbb. ex Schult. f. l. c. 1095 = mutabile.
canadense, *Linn. Sp. Pl.* 1195.—Am. bor.
canadense, Michx. Fl. Bor. Am. i. 191 = striatum.
candidissimum, Cav. Lecc. 994 = neapolitanum.
candidum, *C. Presl, Fl. Sic.* 42.—Sicilia.
canescens, Hort. ex Schult. f. Syst. vii. 1067 = glaucum.
capillare, Cav. Icones, iii. 4. t. 206 = moschatum.
capillare, Desf. Tabl. ed. I. 27 = pusillum.
capitellatum, *Boiss. Diagn.* Ser. I. vii. 118.—Persia; Affghan.
cappadocicum, *Boiss. & Balansa, ex Boiss. Fl. Orient.* v. 241.—Cappadoc.
cardiostemon, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 43.—Persia.
carduchorum, C. Koch, in Linnaea, xxii. (1849) 237 = Aucheri.
caricifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 854 = Pallasi.
caricoides, *Regel, in Act. Hort. Petrop.* vi. (1879) 532. —Turkestan.
carinatum, Habl. Taur. 152 = flavum.
carinatum, *Linn. Sp. Pl.* ed. I. 297.—Europ.
carinatum, Schulz, Fl. Starg. 86 = arenarium.
carinatum, Sm. Engl. Bot. t. 1658 = oleraceum.
Carmeli, *Boiss. Diagn.* Ser. I. xiii. 28.—Palaestin.
carneum, Tenore, Fl. Nap. 159 = roseum.
carneum, Hort. ex Schult. f. Syst. vii. 1028 = sibiricum.
carneum, Targ.-Tozz. Ist. Bot. ed. II. 1. t. 6 = ambiguum.
carneum, Willd. Enum. Hort. Berol. 359 = ascalonicum.
carolinianum, [*Delar. in*] *Red. Lil.* t. 101.—Carolina.
caspium, *Bieb. Fl. Taur. Cauc.* i. 265; iii. 260.—Ross. austr.; Turkest.; Sibir.
cassium, *Boiss. Diagn.* Ser. I. xiii. 28.—Taurus; Syria.
caucasicum, Bieb. Fl. Taur. Cauc. iii. 258 = globosum.
caucason, Turra, ex Sacc. in Atti Ist. Venet. Ser. III. ix. (1863–64) 487 = globosum.
Cepa, Banks, ex Schult. f. Syst. vii. 1008 = proliferum.
Cepa, Falk, Beitr. ii. 157 = fistulosum.
Cepa, *Linn. Sp. Pl.* ed. I. 300.—Persia; Beluchist.
Cepa, Willd. ex Ledeb. Fl. Ross. iv. 169 = galanthum.
cepaeforme, G. Don, in Mem. Wern. Soc. vi. (1827) 31 = A. Cepa.
cepaeum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 113 = A. Cepa.
ceratophyllum, Bess. ex Schult. f. Syst. vii. 1029 = fistulosum.
cernuum, Hort. ex Schult. f. l. c. 1081 = prostratum.
cernuum, *Roth, in Roem. Archiv*, i. II. 40.—Am. bor.
Chamaemoly, *Linn. Sp. Pl.* 301.—Reg. Mediterr.
Chamaemoly, Viviani, Fl. Lib. Spec. 19 = subvillosum.
chamaespathum, *Boiss. Diagn.* Ser. I. vii. 113.—Graecia.
chilense, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 218.—Chili.
chinense, G. Don, in Mem. Wern. Soc. vi. (1827) 83 = Thunbergii.
chinense, Maxim. Prim. Fl. Amur. 284 = odorum.
chionantha, Boiss. Fl. Orient. v. 272 = zebdanense.
chloroneurum, *Boiss. l. c.* 247.—Persia.
chloranthum, Boiss. Diagn. Ser. I. xiii. 33 = flavum.
chloranthum, Lall. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 246 = decipiens.
chloranthum, Schur, Enum. Pl. Transs. 672 = globosum.
Christii, *Janka, in Linnaea*, xxx. (1859–60) 604.—Europ. austr.
chrysantherum, *Boiss. & Reut. ex Boiss. Fl. Orient.* v. 280.—Assyria.
chrysanthum, *Regel, in Act. Hort. Petrop.* iii. (1875) 91. —China.
ciliare, [Delar. in] Red. Lil. vi. t. 311 = subhirsutum.

ALLIUM:—
ciliatum, C. Koch, in Linnaea, xix. (1847) 11 = sphaerocephalum.
ciliatum, Cyr. Pl. Rar. Neap. Fasc. ii. 16 = subhirsutum.
cilicicum, *Boiss. Diagn.* Ser. I. vii. 115.—As. Min.
circinnatum, *Sieber, Reise Kreta*, ii. 316.—Graecia.
cirrhosum, Raeusch. Nom. ed. III. 74 = roseum.
cirrhosum, Vand. Fasc. Pl. 11 = flavum.
clathratum, *Ledeb. Fl. Alt.* ii. 18.—Sibir. alt.
Clusianum, Bory, Fl. Pelop. 21 = circinnatum.
Clusianum, Retz. in Willd. Sp. Pl. ii. 79 = subvillosum.
colchicifolium, *Boiss. Diagn.* Ser. II. iv. 112.—Armenia.
collinum, *Dougl. ex S. Wats. in Proc. Am. Acad.* xiv. (1879) 228.—Am. bor. occ.
collinum, Guss. in Tenore, Syll. 169 = paniculatum.
coloratum, Dulac, Fl. Hautes-Pyr. 110 = A. Schoenoprasum.
coloratum, Spreng. Syst. ii. 38 = flavum.
commune, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 6; ed. II. 68 (1827) = A. Cepa.
commutatum, Guss. Enum. Pl. Inarim. 17 = rotundum.
compactum, Thuil. Fl. Par. 167 = vincale.
complanatum, Boreau, Not. xxxv. p. 12 = oleraceum.
condensatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 121.—Sibir. baical.
confertum, Fisch. Cat. (1821).—Cf. G. Don, in Mem. Wern. Soc. vi. (1827) 63 = deflexum, strictum, senescens.
confertum, Jord. & Fourr. Brev. Fl. Nov. fasc. ii. 127 = roscum.
congestum, G. Don, in Mem. Wern. Soc. vi. (1827) 66 = Stellerianum.
consanguineum, *Kunth, Enum. Pl.* iv. 431.—Reg. Himal.
consimile, Jord. ex Gren. & Godr. Fl. Fr. iii. 208 = carinatum.
contortum, Stokes, Bot. Mat. Med. ii. 229 = A. Scorodoprasum.
controversum, Costa, Suplem. Catal. 62 = pyrenaicum.
controversum, Schrad. Cat. Goett. ex Willd. Enum. Hort. Berol. 358 = sativum.
convallarifolium, Pall. ex Ledeb. Fl. Ross. iv. 184 = A. Victorialis.
Coppoleri, Tineo, Cat. Pl. Hort. Panorm. 18 = paniculatum.
corbariense, Timb. ex Nym. Consp. 737 = subhirsutum.
cornutum, *Clem. It. Dalm.* t. 1. f. 1.—Dalmatia.
cornutum, Kit. in Linnaea, xxxii. (1863) 340 = macrocephalum.
Cowani, Lindl. Bot. Reg. t. 758 = neapolitanum.
Crameri, *Aschers. & Boiss. ex Boiss. Fl. Orient.* v. 279.—Aegypt.
crinitum, Tausch, in Syll. Ratisb. ii. 255 = sphaerocephalum.
cristatum, *Boiss. Fl. Orient.* v. 237.—Thracia.
Cristophi, *Trautv. in Act. Hort. Petrop.* ix. (1884) 268.—As. temp.
croceum, *Torr. in Bot. Mex. Bound.* 218.—Calif.
Cumaria, Buch.-Ham. ex Wall. Cat. n. 5072 = A. Cepa.
Cupani, Guss. Pl. Rar. 143 = moschatum.
Cupani, *Rafin. Caratteri*, (1818) 86. n. 206.—Reg. Mediterr.
cupuliferum, *Regel, in Act. Hort. Petrop.* iii. (1875) 234.—Turkestan.
curtum, *Boiss. & Gaillard, Diagn.* Ser. II. iv. 116.—Syria; Aegypt.
Cusickii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 228.—Am. bor. occ.
cyaneum, *Regel, in Act. Hort. Petrop.*. iii. (1875) 174.—China.
Cydni, Schott & Kotschy, ex Schott, in Oestr. Bot. Wochenbl. (1854) 217 = cassium.
Cyrilli, *Tenore, Fl. Nap.* iii. 364.—Italia; Graecaic; Tauria.
danubiale, Reichb. ex Kunth, Enum. Pl. iv. 423 = senescens.
danubiale, Spreng. Fl. Hal. Mant. 38 = angulosum.
darwasicum, *Regel, in Act. Hort. Petrop.* vii. (1884) 659.—Bokhara.
Decaisnei, Presl, Bot. Bemerk. 114 = flavum.
decipiens, *Fisch. Hort. Gorenk.* ed. II. (1812) 10.—Reg. Caucas; Turkest.; Sibir.
declinatum, Reichb. Ic. Fl. Hort. Germ. v. 46. t. 477 = Stellerianum.
deflexum, Fisch. ex Kunth, Enum. Pl. iv. 428 = Stellerianum.

ALLIUM:—
deflexum, Willd. Sp. Pl. ii. 65 = lineare.
delicatulum, *Sievers, ex Willd. in Roem. & Schult. Syst.* vii. 1133.—Sibir.
densiflorum, De Not. Rep. Fl. Lig. 413 = sphaerocephalum.
denticulatum, Kit. in Linnaea, xxxii. (1863) 342 = carinatum.
dentiferum, Webb & Berth. Phyt. Canar. iii. 345. t. 234 = paniculatum.
denudatum, [Delar. in] Red. Lil. t. 357 = albidum.
Derderianum, Regel, in Act. Hort. Petrop. ii. (1875) 242 = nacmanthoides.
descendens, Koch, Syn. Fl. Germ. ed. I. 719 = vineale.
descendens, Linn. Sp. Pl. ed. I. 299 = sphaerocephalum.
descendens, Pall. Ind. Taur. ex Schult. f. Syst. vii. 1011 = rotundum.
descendens, Sibth. & Sm. Fl. Gr. iv. 15. t. 316 = margaritaceum.
Déséglisei, Bor. Not. Nov. (1853) 17; (in Bull. Soc. Indust. d'Angers).—Cf. Ann. Soc. Linn. Maine-et-Loire, i. (1857) 63 = sphaerocephalum.
deserti, Boiss. ex Regel, in Act. Hort. Petrop. iii. (1875) 76 = Boissieri.
desertorum, *Forsk. Fl. Aegypt. Arab.* 72.—Aegypt.
diaphanum, Janka, in Linnaea, xxx. (1859–60) 605 = tataricum.
dictyoprasum, Ledeb. Fl. Ross. iv. 166 = dyctioprasum.
dilutum, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 13.—Persia.
Dioscoridis, *Sibth. & Sm. Prod. Fl. Gr.* i. 222.—Europ. austr.; Oricns.
djimilense, *Boiss. ex Regel, in Act. Hort. Petrop.* iii. II. (1875) 108.—Pontus.
doloncarense, *Regel, l. c.* 113.—Sibir.
Douglasii, *Hook. Fl. Bor. Am.* ii. 184. t. 197.—Calif.
Douglasii, Wood, in Proc. Philad. Acad. (1868) 170 = unifolium, bisceptrum.
Drègeanum, Kunth, Enum. Pl. iv. 382 = A. Scorodoprasum.
Drummondi, Regel, in Act. Hort. Petrop. iii. (1875) 112 = mutabile.
Duriaeanum, Regel, l. c. 54 = A. Ampeloprasum.
Durieuanum, Walp. Ann. Bot. i. 860 = praec.
Durtaeanum, J. Gay, in Ann. Sc. Nat. Sér. III. viii. (1847) 218 = A. Ampeloprasum.
dyctioprasum, *C. A. Mey. ex Kunth, Enum. Pl.* iv. 390.—Armenia.
effusum, Boiss. Fl. Orient. v. 257 = stamineum.
clatum, *Regel, in Act. Hort. Petrop.* viii. (1884) 665.—Bokhara.
Elwesi, Regel, l. c. v. (1877) 266 = acuminatum ?
ellipticum, Wall. Cat. n. 5069 = A. Victorialis.
eminens, Gren. ex Parl. Fl. Ital. ii. 567 = sphaerocephalum.
Erdelii, Baker, in Bot. Mag. t. 6426 = orientale.
Erdelii, *Zucc. in Abh. Math. Phys. Acad. Muench.* iii. (1843) 236. t. 5.—Palaestin.
erectum, G. Don, in Mem. Wern. Soc. vi. (1827) 18 = rotundum.
ericetorum, Thore, Chlor. Land. 123 = ochroleucum, suaveolens.
eriophyllum, *Boiss. Diagn.* Scr. I. vii. 112.—Persia.
erubescens, C. Koch, in Linnaea, xxii. (1849) 242 = rudbaricum.
erythraeum, *Griseb. Spicil. Fl. Rumel.* ii. 396.—Macedon.
erythraeum, Weiss, ex Boiss. Fl. Orient. v. 253 = Weissii.
esculentum, Salisb. Prod. 235 = A. Cepa.
euosmum, Link & Otto, Ic. Pl. Rar. Hort. Berol. i. 15. t. 8 = nudicaule.
euschoenoprasum, Syme, Engl. Bot. ed. III. ix. 315 = A. Schoenoprasum.
exaltatum, Kar. & Kir. ex Ledeb. Fl. Ross. iv. 173 = obliquum.
exile, Boiss. & Orph. Diagn. Scr. II. iv. 118 = sipylum.
exsertum, Baker, in Journ. Bot. xii. (1874) 294 = Bakeri.
exsertum, *G. Don, in Mem. Wern. Soc.* vi. (1827) 17.—Rossia.
falcifolium, *Hook. & Arn. Bot. Beech. Voy.* 400.—Calif.
fallax, *Schult. f. Syst.* vii. 1072.—Europ.; As. Min.
Fedschenkoanum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 82.—Turkestan.
Fetisowi, *Regel, l. c.* v. (1877) 631.—Turkestan.
filidens, *Regel, l. c.* iii. II. (1875) 174.—Turkestan.

ALLIUM:—
filiforme, *Colla, Herb. Pedem.* t. 90. f. 3.—Italia.
fimbriatum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 232.—Calif.
Fischeri, Bess. in Roem. & Schult. Syst. vii. 1081 = Stellerianum.
Fischeri, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 161.—Sibir.
fissile, S. F. Gray, Nat. Arr. ii. 183 = A. Ascalonicum.
fissile, *Steud. Nom.* ed. II. i. 51, nomen.—As. Min.
fistulosum, *Linn. Sp. Pl.* 301.—Sibir.
flavescens, Bess. Enum. Pl. Volh. 56 = albidum.
flavescens, Pocpp. ex Kunth, Enum. Pl. iv. 459 = striatum.
flavidum, *Ledeb. Fl. Alt.* ii. 7.—Sibir.
flavum, Less. ex Ledeb. Fl. Ross. iv. 172 = delicatulum.
flavum, *Linn. Sp. Pl.* 299.—Europ.; Oriens.
flavum, Salisb. Prod. 235 = A. Moly.
flexifolium, Jord. in Gren. & Godr. Fl. Fr. iii. 208 = carinatum.
flexum, Wallst. & Kit. Pl. Rar. Hung. ii. 307 = carinatum.
flexuosum, Host, Fl. Austr. i. 422 = carinatum.
flexuosum, Urville, in Mém. Soc. Linn. Paris, i. (1822) 294 = staticaeforme.
foetidum, Willd. Enum. Suppl. 17 = flexum, carinatum.
foliosum, Clar. ex [DC. in] Red. Lil. t. 214 = A. Schoenoprasum.
Fontanesii, J. Gay, in Ann. Sc. Nat. Sér. III. viii. (1847) 209 = flavum.
fragrans, Boj. Hort. Maurit. 348 = Milla borbonica.
fragrans, Cyr. ex Tenore, Fl. Nap. iii. 364 = Cyrilli.
fragrans, Vent. Jard. Cels, 26 = Nothoscordum fragrans.
frigidum, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 34.—Graecia.
Frivaldzkyanum, Kunze, in Linnaea, xvi. (1842) 311 = margaritaceum.
fuscum, Schrad. ex Hornem. Hort. Hafn. ii. 962 = pulchellum.
fuscum, Spreng. Syst. ii. 37 = longispathum.
fuscum, Waldst. & Kit. Pl. Rar. Hung. iii. 267. t. 241 = paniculatum.
Fussii, *Kerner, in Oestr. Bot. Zeitschr.* xxviii. (1878) 155.—Austria.
galanthum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 508.—Soongaria.
gaditanum, *Perez-Lara, ex Willk. in Illustr. Fl. Hisp.* 81.—Morocco.
Gardinia, Bertero, ex Steud. Nom. ed. II. i. 664 = Brodiaea violacea.
Gasparrini, Guss. in Enum. Pl. Inarim. 337. t. 16 = A. Ampeloprasum.
Gayi, *Boiss. Fl. Orient.* v. 271.—Cilicia.
Geyeri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 227.—Calif.
Gherardi, De Not. Rep. Fl. Lig. 412 = sphaerocephalum.
giganteum, *Regel, Gartenfl.* (1883) 91.—Reg. Himal.
glandulosum, *Link & Otto, Ic. Rar.* i. 33. t. 17.—Mexic.
glaucum, [Desf.] Hort. Par. ex [DC. in] Red. Lil. iv. t. 214 = foliosum.
glaucum, Hort. Par. ex Ledeb. Fl. Ross. iv. 167 = A. Schoenoprasum.
glaucum, Schrad. ex Poir. Encyc. Suppl. i. 265 = senescens.
globosum, *Bieb. Fl. Taur. Cauc.* i. 262; iii. 257.—Europ. austr.; Reg. Cauc.; Sibir.
globosum, [*DC. in*] *Red. Lil.* iii. t. 179.—Hab. ?
glumaceum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 260.—Cataonia.
gomphrenoides, *Boiss. & Heldr. Diagn.* Ser. I. vii. 114.—Lycaonia.
gorumsense, *Boiss. Fl. Orient.* v. 233.—Cilicia.
Gouanii, G. Don, in Mem. Wern. Soc. vi. (1827) 85 = neapolitanum.
Govanianum, *Wall. Cat.* n. 5071.—Reg. Himal.
gracile, [Dryand. in] Ait. Hort. Kew. i. 429 = Nothoscordum fragrans.
graecum, J. Gay, ex Boiss. Fl. Orient. v. 271 = Gayi.
graecum, Urville, in Mem. Soc. Linn. Par. i. (1822) 293 = trifoliatum.
gramineum, C. Koch, in Linnaea, xxii. (1849) 239 = rotundum.
graminifolium, *Buch, in Abh. Acad. Berl.* xvii. (1816) 362.—Ins. Canar.
graminifolium, Loisel. Fl. Gall. i. 253 = subhirsutum.
graminifolium, Pers. Syn. i. 356 = ochroleucum.
grandiflorum, Lam. Encyc. i. 68 = narcissiflorum.

ALLIUM :—
Grazi, *Regel, in Act. Hort. Petrop.* iii. (1875) 125.—Japon.
Griffithianum, Boiss. Diagn. Ser. II. iv. 117 = rubellum.
Grimmi, *Regel, Dec. Pl. Nov.* 10.—Cf. *Act. Hort. Petrop.* viii. (1883) 272.—Turkestan.
Guicciardii, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 233 = flavum.
guttatum, Stev. in Mém. Soc. Nat. Mosc. ii. (1809) 173. t. 11 = margaritaceum.
haemanthoides, *Boiss. ex Regel, Act. Hort. Petrop.* iii. II. (1875) 240.—Persia.
haematochiton, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 227.—Calif.
Halleri, Bab. Man. Brit. Bot. ed. I. 305 = A. Scorodoprasum.
Halleri, Crantz, Inst. i. 488 = lineare.
Halleri, G. Don, in Mem. Wern. Soc. vi. (1827) 15 = A. Ampeloprasum.
Heldreichii, *Boiss. Diagn.* Ser. II. iv. 116.—Graecia.
hierochuntinum, *Boiss. Fl. Orient.* v. 244.—Palaestin.
himalayense, Regel, in Act. Hort. Petrop. iii. II. (1875) 118 = Stocksianum.
hirsutum, Lam. Fl. Fr. iii. 262 = subhirsutum.
hirsutum, *Zucc. in Abh. Bayer. Akad.* iii. (1843) 232. t. 2.—Palestin.
hirtifolium, *Boiss. Fl. Orient.* v. 281.—Persia.
hirtovaginatum, Kunth, Enum. Pl. iv. 412 = Cupani.
Hoeltzeri, *Regel, in Act. Hort. Petrop.* viii. (1884) 657; *Regel, Gartenfl.* (1884) 291.—Turkestan.
holmense, Mill. ex Kunth, Enum. Pl. iv. 384 = A. Ampeloprasum.
Hookeri, Thw. Enum. Pl. Zeyl. 339 = Wallichii.
humile, *Kunth, Enum. Pl.* iv. 442.—Ind. or.
hyalinum, *Curran, in Bull. Calif. Acad.* n. 3. (1885) 155.—Calif.
hyemale, Rafin. Fl. Ludov. 19 = fragrans.
hymenorrhizum, *Ledeb. Fl. Alt.* ii. 12.—Turkest.; Persia.
hymettium, *Boiss. & Heldr. Diagn.* Ser. II. iv. 120.—Graecia.
iliense, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) I. 452.—Soongaria.
illyricum, Jacq. Collect. iii. 273 = roseum.
illyricum, Link & Otto, ex Kunth, Enum. Pl. iv. 422 = senescens.
inaequale, Janka, in Linnaea, xxx. (1859–60) 603 = moschatum.
incarnatum, Hornem. Hort. Hafn. i. 323 = roseum.
inderiense, Fisch. ex Roem. & Schult. Syst. vii. 1133 = tartaricum.
inodorum, Ait. Hort. Kew. i. 427 = Nothoscordum fragrans.
inodorum, Willd. Sp. Pl. ii. 76 = angulosum.
insubricum, Boiss. & Reut. Cat. Sem. Jard. Genèv. (1854).—Cf. Linnaea, xxix. (1857) 721 = narcissiflorum.
intermedium, DC. Fl. Fr. vi. 318 = tataricum, paniculatum.
intermedium, G. Don, in Mem. Wern. Soc. vi. (1827) 38 = oleraceum.
Jacquemonti, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 162.—Ind. or.
Jacquemontii, Kunth, Enum. Pl. iv. 399 = rubellum.
japonicum, *Steud. Nom.* ed. II. i. 50.—Japon.
japonicum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 133.—Japon.
jeholense, *Franch. in Nouv. Arch. Mus. Hist. Nat. Par.* Sér. II. vii. (1884) 113; *Pl. Davidian.* i. (1884) 305.—China.
jenischianum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 237.—Persia.
jesdianum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 217.—Persia.
juldusicolum, *Regel, in Act. Hort. Petrop.* vi. (1879) 523.—Turkestan.
junceum, Jacquem. ex Baker, in Journ. Bot. xii. (1874) 295 = Jacquemonti.
junceum, *Sibth. & Sm. Prod.* i. 226.—Cyprus.
juncifolium, *D. Don, ex Steud. Nom.* ed. II. i. 52.—Chili.
karakense, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 76.—Turkestan.
karataviense, *Regel, l. c.* 243.—Turkestan.
Kaufmanni, *Regel, l. c.* 84.—Turkestan.
kermesinum, Reichb. Ic. Fl. Germ. x. 25 = globosum.
Kesselringi, *Regel, in Act. Hort. Petrop.* viii. (1883) 272.—Turkestan.

ALLIUM :—
Klotzschi, Regel, in Act. Hort. Petrop. iii. II. (1875) 255 = lilacinum.
Kochii, Lange, ex Nym. Consp. 736 = vinealc.
kokanicum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 104.—Turkestan.
Korolkowi, *Regel, l. c.* ii. (1873) 407.—Turkestan.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 117.—Persia.
Kunthii, *G. Don, in Mem. Wern. Soc.* vi. (1827) 82.—Mexic.
Kuschakerwiczi, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 117.—Turkestan.
laceratum, *Boiss. & Noë, Diagn.* Ser. II. iv. 112.—Persia.
lachnophyllum, *Payne, in Pal. Expl. Soc. Cat. Pl.* 125.—Palaestin.
lacteum, Sibth. & Sm. Prod. 226 = neapolitanum.
lacunosum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 231.—Calif.
lacteum, Benth. Pl. Hartw. 339 = Brodiaea lactea.
laetum, Pollini, Cat. Veron. (1812) 7 = neapolitanum.
laetum, Salisb. Prod. 235 = A. Porrum.
Lallemanti, Regel & Rach, in Ind. Sem. Hort. Petrop. (1858) 24 = decipiens.
latifolium, Gilib. Exercit. ii. 470 = ursinum.
latifolium, Jaub. & Spach, Illustr. Pl. Orient. ii. 3. t. 103 = A. Akaka.
laxiflorum, Tausch, in Flora, xii. (1829) Erg. 46 = vineale.
laxum, G. Don, in Mem. Wern. Soc. vi. (1827) 63 = angulosum.
Ledebourianum, *Schult. f. Syst.* vii. 1029.—Rossia.
Lehmannianum, *Merckl. ex Bunge, in Mém. Sav. Étr. Pétersb.* vii. (1851) 509; (*Rel. Lehm.* 333).—Turkestan.
Lemmoni, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 234.—Calif.
lepidum, Kunth, Enum. Pl. iv. 408 = paniculatum.
lepidum, Ledeb. Ic. Fl. Ross. iv. 17 = Pallasii.
leptophyllum, Schur, Enum. Pl. Transs. 674 = senescens.
leptophyllum, Wall. Cat. n. 5073 = rubellum.
Lessingii, Kar. ex Ledeb. Fl. Ross. iv. 170 = sabulosum.
leucanthum, C. Koch, in Linnaea, xxii. (1849) 240 = A. Ampeloprasum.
leucocephalum, Turcz. ex Ledeb. Fl. Ross. iv. 179 = flavidum.
Libani, *Boiss. Diagn.* Ser. I. xiii. 26.—Syria.
ligusticum, De Not. Prosp. Fl. Lig. 55 = flavum.
lilacinum, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 50.—Reg. Himal.
lilacinum, Royle, ex Regel, in Act. Hort. Petrop. iii. (1875) 89 = rubens.
liliflorum, Zeyh. Cat. Schwetz. (1818) 15 = neapolitanum.
lineare, *Linn. Sp. Pl.* 295.—Sibir.
lineare, Tenore, Fl. Nap. i. 154 = sphaerocephalum.
lineare, Willd. ex Kunth, Enum. Pl. iv. 419 = strictum.
littoreum, Bertol. Amoen. It. 241 = vineale.
littoreum, G. Don, in Mem. Wern. Soc. vi. (1827) 8 = monspessulanum.
longibulbum, Dulac, Fl. Hautes-Pyr. 110 = A. Victorialis.
longicaule, *Hornem. Hort. Hafn. Suppl.* 130.—Am. bor.
longicaule, Hort. ex Schult. f. Syst. vii. 1067 = glaucum.
longicuspis, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 45.—Kokania.
longifolium, *Spreng. Syst.* ii. 38.—Am. mer.; Mexic.
longifolium, Lindl. Bot. Reg. t. 1034 = glandulosum.
longipetiolatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = ursinum.
longisepalum, Bertol. in Nov. Comm. Acad. Bonon. v. (1842) 429 = rubellum.
longispathum, [Delar. in] Red. Lil. t. 316 = paniculatum.
longistylum, Baker, in Journ. Bot. xii. (1874) 294 = sacculiferum.
loratum, *Baker, l. c.* 290.—Tibet. occ.
lusitanicum, *Lam. Encyc.* i. 70.—Lusitan.
lusitanicum, Link, ex Regel, in Act. Hort. Petrop. iii. II. (1875) 78 = A. Schoenoprasum.
lusitanicum, [Delar. in] Red. Lil. t. 271 = angulosum.
luteum, Hort. ex Dietr. Lexic. Gart. ed. II. i. 265 = obliquum.
Macleanii, *Baker, in Bot. Mag.* t. 6707.—Affghan.
Macnabianum, *Hort. ex Regel, Gartenfl.* (1873) 261. t. 77. f. 2–3.—Hab. ?

ALLIUM :—
macranthum, *Baker, in Journ. Bot.* xii. (1874) 293.—Reg. Himal.
macrocephalum, *Kit. in Linnaea*, xxxii. (1863) 340.—Austria.
macrochaetum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 239.—Assyria.
macrorhizon, Regel, in Act. Hort. Petrop. iii. II. (1875) 154.—Turkestan.
macrorrhizum, Boiss. Diagn. Ser. I. xiii. 32 = hymenorhizum.
macrospermum, Boiss. & Kotschy, ex Boiss. Fl. Orient. v. 282 = orientale.
macrostemon, *Bunge, in Mém. Sav. Étr. Acad. Pétersb.* ii. (1835) 139.—China bor.
macrostylum, *Regel, in Act. Hort. Petrop.* vi. (1879) 296.—Tianschan.
macrum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 233.—Am. bor.
madidum, *S. Wats. l. c.* 228.—Am. bor.
magicum, Linn. Sp. Pl. ed. II. 424 = nigrum.
majale, Cyr. Pl. Rar. Neap. Fasc. i. 13. t. 4 = roseum.
margaritaceum, Heldr. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 201 = staticiformc.
margaritaceum, Moench, Meth. Suppl. 80 = A. Scorodoprasum.
margaritaceum, *Sibth. & Sm. Fl. Graec. Prod.* i. 224.—Europ.; Oriens.
marginatum, *Janka, in Termesz. Füzetek*, viii. (1884) 29.—Europ. austr.
maritimum, Rafin. Caratt. 86. n. 205 = pusillum.
maritimum, *Torr. in Rep. U. St. & Mex. Bound.* 217.—Calif.
Maximowiczi, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 153.—As. or.
medium, G. Don, in Mem. Wern. Soc. vi. (1827) 88 = triquetrum.
megalobulbon, *Regel, in Act. Hort. Petrop.* vi. (1879) 526.—Turkestan.
melanantherum, *Panč. Elem.* 64.—Europ. merid.
microcephalum, Tausch, in Flora, ii. (1819) 462 = strictum.
microcephalum, Willd. ex Kunth, Enum. Pl. iv. 425 = angulosum.
minutiflorum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 242.—Persia.
mobilense, Regel, l. c. = mutabile.
modestum, Boiss. Diagn. Ser. I. xiii. 33 = paniculatum.
mogadorense, Willd. ex Kunth, Enum. Pl. iv. 384 = A. Ampeloprasum.
Moly, Georgi, Beschreib. Russ. Reich. iii. IV. 896 = decipiens.
Moly, Griseb. & Schur, ex Regel, in Act. Hort. Petrop. iii. II. (1875) 199 = globosum.
Moly, *Linn. Sp. Pl.* 301.—Europ.
Moly, Ucria, Hort. Panorm. 153 = subhirsutum.
monadelphum, *Turcz. ex Kunth, Enum. Pl.* iv. 393.—Sibir. baical.
mongolicum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 160.—Mongolia.
monserratense, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 207 = carinatum.
monspessulanum, Gouan, Illustr. 24 = nigrum.
monspessulanum, Willd. Enum. Suppl. 16 = paniculatum.
montanum, Guss. Pl. Rar. 142 = tenuiflorum.
montanum, Reichb. Ic. Fl. Germ. x. t. 463. f. 1060 = flavum.
montanum, Tenore, Fl. Nap. iii. 367 = Cupani.
montanum, F. W. Schmidt, ex Schult. f. Syst. vii. 1072 = losum.
montanum, Sibth. & Sm. Fl. Graec. t. 319 = paniculatum.
moschatum, *Linn. Sp. Pl.* ed. I. 299.—Europ.; Oriens.
moschatum, Moris, Stirp. Sard. Elench. i. 46 = parciflorum.
moschatum, Sintenis, ex Regel, in Act. Hort. Petrop. iii. II. (1875) 199 = globosum.
moschatum, Urville, in Mém. Soc. Linn. Par. i. (1822) 293 = Cupani.
multibulbosum, Jacq. Fl. Austr. i. 9 = nigrum.
multiflorum, DC. Fl. Fr. vi. 316 = polyanthum.
multiflorum, *Desf. Fl. Atlant.* i. 288.—Algeria.
multiflorum, Kunth, Enum. Pl. iv. 387 = rotundum.
multiflorum, Regel, in Act. Hort. Petrop. iii. (1875) 61, pro parte = rudbaricum.
Murrayanum, Regel, in Gartenfl. (1873) 260. t. 770. f. 1 = acuminatum.
mutabile, *Michx. Fl. Am. Bor.* i. 195.—Georgia.

ALLIUM :—
myrianthum, Boiss. Diagn. Ser. I. v. 59 = A. Ampeloprasum.
Napus, Pall. ex Kunth, Enum. Pl. iv. 395 = A. Cepa.
narcissiflorum, *Vill. Prosp.* 18.—Europ.
narcissifolium, Lam. Fl. Fr. iii. 263 = pedemontanum.
narcissifolium, Scop. Fl. Carn. ed. II. i. 238 = senescens.
neapolitanum, *Cyr. Pl. Rar. Neap.* i. 13. t. 4.—Europ.; Oriens.
nebrodense, Guss. Prod. Fl. Sic. i. 404 = flavum.
neglectum, Wender. in Schrift. Ges. Nat. Marb. ii. (1831) 211 = A. Scorodoprasum.
nemorale, Salisb. Prod. 235 = ursinum.
Nereidum, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 244.—China.
neriniflorum, *G. Don, ex Loud. Encyc. Pl. Suppl.* ii. 1342.—China.
nerinifolium, Baker, in Journ. Bot. xii. (1874) 290 = Thunbergii.
nevadense, *S. Wats. in Bot. King, Exp.* 351, 487. t. 38. f. 1–3.—Calif.
Nevii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 231.—Calif.
nigrum, All. Fl. Pedem. ii. 158. t. 25 f. 1 = narcissiflorum.
nigrum, Bieb. Fl. Taur. Cauc. i. 266 = decipiens.
nigrum, G. Don, in Mem. Wern. Soc. vi. (1827) 89 = multibulbosum.
nigrum, *Linn. Sp. Pl.* ed. II. 430.—Europ.; Oriens.
nigrum, Sibth. & Sm. Fl. Graec. t. 323 = Bauerianum.
nipponicum, *Franch. & Sav. Enum. Pl. Jap.* ii. 527.—Japon.
nitens, Sauzé & Maill. Cat. Pl. Deux-Sèvres, 51 = vineale.
nitidulum, Fisch. ex Ledeb. Fl. Ross. iv. 171 = Pallasii.
Nitschmanni, Willd. ex Ledeb. l. c. 175 = flavum.
nivale, Jacquem. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 178 = Govanianum.
niveum, Roth, Cat. Bot. fasc. ii. 35 = subhirsutum.
Noëoanum, *Reut. ex Regel, in Act. Hort. Petrop.* iii. II. (1875) 235.—As. Min.
nudicaule, *Lehm. Cat. Hort. Hamb.* (1826).—Bras.
nutans, Herbb. quorr. ex Schult. f. Syst. vii. 1088 = cernuum.
nutans, *Linn. Sp. Pl.* 299.—Sibir.
Nuttallii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 227.—Am. bor.
obliquum, *Linn. Sp. Pl.* 296.—Sibir.
obscurum, Bieb. ex Schult. f. Syst. vii. 1003 = A. Scorodoprasum.
obtusiflorum, [DC. in] Red. Lil. t. 118 = pusillum.
obtusifolium, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 51.—Reg. Himal.
occidentale, A. Gray, in Proc. Am. Acad. vii. (1868) 390 = attenuifolium.
ochroleucum, Reichb. Ic. Fl. Germ. x. t. 498. f. 1090 = globosum.
ochroleucum, *Waldst. & Kit. Pl. Rar. Hung.* ii. 204. t. 186.—Europ.
odoratissimum, *Desf. Fl. Atlant.* i. 289. t. 83.—Afr. bor.
odorum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 854 = angulosum.
odorum, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 48 = ochroleucum.
odorum, *Linn. Mant.* 62.—Sibir.
odorum, Lour. Fl. Cochinch. 251 = Thunbergii.
odorum, Tenore, Fl. Nap. Prod. 21 = nigrum.
odorum, Thunb. Fl. Jap. 132 = Thunbergii.
oleraceum, Des Moul. Cat. Dord. 142, ex parte = dentiferum.
oleraceum, *Linn. Sp. Pl.* 299.—Europ.
oliganthum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 856 = A. Schoenoprasum.
Olivieri, *Boiss. Fl. Orient.* v. 284.—Mesopotamia.
olympicum, *Boiss. Diagn.* Ser. I. v. 58.—Thracia.
Ophioscorodon, G. Don, in Mem. Wern. Soc. vi. (1827) 5 = sativum.
Opisii, Wolfner, in Lotos, iv. (1854) 176 = triquetrum.
oreophiloides, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 114.—Turkestan.
orcophilum, *C. A. Mey. Verz. Pfl. Cauc.* 37.—Caucas.
orcoprasum, *Schrenk, in Bull. Sc. Acad. St. Pétersb.* x. (1842) 354.—Turkestan.
orientale, *Boiss. Diagn.* Ser. I. xiii. 25.—Oriens.
ornithogaloides, Walt. Fl. Carol. 121 = striatum.

ALLIUM :—
Ostrowskianum, *Regel, in Act. Hort. Petrop.* vii. (1880) 545.—Turkestan.
oviflorum, *Regel, l. c.* viii. (1884) 658.—Reg. Himal.
oxypetalum, G. Don, in Mem. Wern. Soc. vi. (1827) 37 = oleraceum.
Pallasii, *Murr. in Comm. Gott.* vi. (1775) 32. t. 3.—Sibir.
pallens, Linn. Sp. Pl. ed. II. 427 = paniculatum.
pallens, Reichb. Ic. Fl. Germ. x. t. 484. f. 1062 = flavum.
pallens, *Sibth. & Sm. Fl. Graec.* t. 317.—Graecia.
Palmeri, *S. Wats. Bot. in Rep. Geol. Expl. Fortieth Parallel*, 487. t. 27. f. 10, 11.—N. Mexico.
palustre, Pourr. ex Lag. Elench. 13 = A. Schoenoprasum.
palustre, Salisb. Prod. 235 = sibiricum.
paniculatum, All. Fl. Pedem. ii. 157 = flavum.
paniculatum, Ker-Gawl. in Bot. Mag. t. 1432 = caucasicum.
paniculatum, Vill. Hist. Pl. Dauph. ii. 354 = intermedium.
paniculatum, *Linn. Syst.* ed. X. 978.—Europ.; Oriens.
paniculatum, Sibth. & Sm. Fl. Graec. iv. t. 318 = fuscum.
paniculatum, Viv. Fl. Lib. Spec. 19 = nigrum.
pannonicum, Gromow. ex Trautv. in Act. Hort. Petrop. ix. (1884) 273, nomen.—Europ.
papillare, *Boiss. Diagn.* Ser. I. xiii. 27.—Palaestin.
paradoxum, *G. Don, in Mem. Wern. Soc.* vi. (1827) 72.—Caucas.; Persia.
parciflorum, *Viv. App. Fl. Cors. Prod.* 2.—Corsic.
Parishii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 380.—Am. bor.
Parryi, *S. Wats. l. c.* xiv. (1879) 231.—Calif.
parviflorum, Desf. Fl. Atlant. i. 290 = paniculatum.
parviflorum, Desv. Fl. Anj. 96 = sphaerocephalum.
parviflorum, *Linn. Sp. Pl.* ed. II. 427.—Europ.
parviflorum, Thuill. Fl. Paris, ed. II. i. 166 = oleraceum.
parvifolium, Crantz, Inst. i. 49 = parviflorum.
parvum, *Kellogg, in Proc. Calif. Acad.* iii. (1863) 54. f. 13.—Calif.
Paterfamilias, *Boiss. Diagn.* Ser. II. iv. 114.—Reg. Caucas.
paucibulbosum, *Haw. ex Steud. Nom.* ed. II. i. 53.—Hispan.
pauciflorum, Gren. & Godr. Fl. Fr. iii. 210 = parciflorum.
pauciflorum, Willd. ex Ledeb. Fl. Ross. iv. 169 = A. Cepa.
pedemontanum, Willd. Sp. Pl. ii. 77 = narcissiflorum.
pendulinum, Tenore, Fl. Nap. i. 168. t. 31 = triquetrum.
permixtum, Guss. Fl. Sic. Prod. i. in add. 8 = subhirsutum.
persicum, Fisch. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 43 = A. Scorodoprasum.
petiolatum, Lam. Fl. Fr. iii. 261 = ursinum.
petraeum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 511 = globosum.
phalereum, Heldr. & Sart. ex Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 233 = staticiforme.
phanerantherum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 235.—Syria.
philistaeum, Boiss. Diagn. Ser. I. xiii. 26 = Erdelii.
phrygium, *Boiss. in Bal. ex Boiss. Fl. Orient.* v. 256.—Phrygia.
phthioticum, *Boiss. & Heldr. ex Boiss. l. c.* 274.—Oriens.
pilosum, *Sibth. & Sm. Fl. Graec.* t. 321.—Archip. Graec.
pisidicum, Boiss. & Heldr. ex Boiss. Diagn. Ser. I. xiii. 34 = Cupani.
plantaginense, Willk. & Lange, Prod. Fl. Hisp. i. 211 = A. Victorialis.
plantagineum, Lam. Fl. Fr. iii. 262 = A. Victorialis.
platycaule, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 234.—Calif.
platyspathum, *Schrenk, Enum.* 7.—Turkestan.
platystemon, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 514 = oreophilum.
pleianthum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 233.—Am. bor.
Plummerae, *S. Wats. l. c.* xviii. (1882–83) 195.—Mexic.
Poeppigi, *Kunth, Enum. Pl.* iv. 456.—Chili.
polyanthum, Schult. f. Syst. vii. 1016 = rotundum.
polyphyllum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 509.—Turkestan.
polyrhizum, *Turcz. ex Regel, in Act. Hort. Petrop.* iii. (1875) 162.—Sibir. baical.

ALLIUM :—
porphyroprasum, Heldr. & Sart. ex Boiss. Fl. Orient. v. 233 = rotundum.
porraceum, S. F. Gray, Nat. Arr. ii. 181 = A. Ampeloprasum.
Porrum, Georgi, Beschreib. Russ. Reich. iii. IV. 891 = obliquum.
Porrum, *Linn. Sp. Pl.* 295.—Europ.
Potanini, *Regel, in Act. Hort. Petrop.* vi. (1879) 295.—Mongol.
praecissum, Reichb. Ic. Fl. Germ. v. 17. t. 429 = paniculatum.
pratense, Schleich. ex Kunth, Enum. Pl. iv. 382 = carinatum.
Preslianum, Schult. f. Syst. vii. 1132 = rotundum.
procerum, *Trautv. in Act. Hort. Petrop.* viii. (1884) 663; ix. (1884) 274.—Bokhara.
proliferum, Schrad. ex Willd. Enum. Hort. Berol. 358 = A. Cepa.
prostratum, Maxim. Prim. Fl. Amur. 283 = Maximowiczi.
prostratum, Trev. Ind. Sem. Vratisl. (1821) = Stellerianum.
pruinatum, Link, in Spreng. Syst. ii. 35 = monspessulanum.
Przewalskianum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 164.—China occ.
Pseudo-Cepa, Schrenk, in Bull. Sc. Acad. Pétersb. x. (1842) 355 = galanthum.
Pseudo-ochroleucum, Schur, Enum. Pl. Transs. 673 = globosum.
pulchellum, *G. Don, in Mem. Wern. Soc.* vi. (1827) 46.—Europ.; Oriens.
pulchrum, Clark, Trav. iii. 337 = nigrum.
punctulatum, Schlecht. in Linnaea, xix. (1847) 401 = A. Schoenoprasum.
purpureum, Salisb. Prod. 235 = senescens.
purpureum, Schur, Enum. Pl. Transs. 672 = carinatum.
Purshii, G. Don, in Mem. Wern. Soc. vi. (1827) 10 = vineale.
pusillum, *Cyr. ex Tenore, Cat. Neap.* (1813) 4.—Ital.
pusillum, Presl, Delic. Prag. 147 = Cupani.
pusillum, Willd. Enum. Hort. Berol. Suppl. 17 = delicatulum.
pustulosum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 243.—Cataonia.
pylium, De Not. in Linnaea, xviii. (1844) 157 = A. Ampeloprasum.
pyrenaicum, *Costa, Suplem. Catal.* 92; *Costa & Vayr. in Anal. Soc. Esp. Hist. Nat.* ix. (1880) 87.—Pyren.
Raddeanum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 155.—Sibiria.
ramosum, Georgi, Beschr. Russ. Reich. iii. IV. 892 = odorum.
ramosum, Jacq. Misc. ii. 303 = obliquum.
ramosum, *Linn. Sp. Pl.* 296.—Sibiria.
ramosum, Pall. ex Ledeb. Fl. Ross. iv. 184 = tataricum.
reflexum, F. G. Dietr. Lexic. Gaert. ed. II. i. 285 = A. Schoenoprasum.
reflexum, *Boiss. & Reut. ex Boiss. Fl. Orient.* v. 281.—Mesopot.
Regelii, *Trautv. in Act. Hort. Petrop.* ix. (1884) 275.—Transcauc.
Renari, *Regel, l. c.* vi. (1879) 521.—Turkestan.
reticulatum, Benth. Pl. Hartw. 339 = attenuifolium.
reticulatum, *Fras. ex G. Don, in Mem. Wern. Soc.* vi. (1827) 36.—Am. bor.
reticulatum, J. & C. Presl, Fl. Cech. 73 = strictum.
reticulatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Victorialis.
reticulatum, Wallr. Sched. Crit. 135 = angulosum?
Reuterianum, *Boiss. Diagn.* Ser. I. v. 60.—As. Min.
riparium, Opiz, Natural. vi. (1824) 50 = sibiricum.
robustum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 853 = atropurpurcum.
Rosenbachianum, *Regel, in Act. Hort. Petrop.* viii. (1884) 664.—Bokhara.
roseum, Krock. Fl. Siles. i. 516. t. 43 = sibiricum.
roseum, *Linn. Sp. Pl.* ed. I. 296.—Reg. Mediterr.
Rothii, *Zucc. in Abh. Bayer. Akad.* iii. (1843) 232. t. 2. f. 2.—Palaestin.
rotundifolium, Lumnitz. ex Steud. Nom. ed. II. i. 53 = rotundum.
rotundum, All. Fl. Pedem. ii. 155 = acutiflorum.
rotundum, *Linn. Sp. Pl.* ed. II 423.—Europ.; As. Min.
rotundum, Wimm. & Grab. Enum. 56 = vineale.
Roxburghi, Kunth, Enum. Pl. iv. 454 = odorum.

ALLIUM:—
rubellum, *Bieb. Fl. Taur. Cauc.* i. 264.—Europ.; Oriens; Sibir.
rubellum, Presl, Fl. Sic. 42 = rotundum.
rubellum, Regel, in Act. Hort. Petrop. iii. II. (1875) 107 = Tschulpias
rubellum, Willd. ex Ledeb. Fl. Ross. iv. 179 = globosum.
rubens, *Baker, in Journ. Bot.* xii. (1874) 293.—Reg. Himal.
rubens, Schrad. ex Willd. Enum. 360 = Stellerianum.
rubicundum, Niven, ex G. Don, in Mem. Wern. Soc. vi. (1827) 16 = rotundum.
rubicundum, *Willd. ex Kunth, Enum. Pl.* iv. 439.—Hab.?
rubrovittatum, *Boiss. & Heldr. in Boiss. Diagn.* Ser. I. xiii. 29.—Creta.
rudbaricum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 215.—Persia.
rupestre, *Bieb. Fl. Taur. Cauc.* iii. 259.—Iberia; Taur.
rupestre, Fisch. ex Schrank, in Denkschr. Regensb. Bot. Ges. ii. (1822) 38 = saxatile.
rupestre, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 260 = paniculatum.
Ruprechti, *Boiss. Fl. Orient.* v. 264.—Caucas.
ruthenicum, Steud. Nom. ed. II. i. 54 = flavum.
sabulosum, *Fisch. ex Goebel, Reise*, ii. 311.—Reg. Casp.
sabulosum, *Stev. ex Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1842) 513.—Persia; Turkestan.
sacculiferum, *Maxim. Prim. Fl. Amur.* 281.—Reg. Amur.
sairamense, *Regel, in Act. Hort. Petrop.* vi. (1879) 520.—Turkestan.
Salesovianum, Fisch. ex Regel, l. c. iii. II. (1875) 162 = Fischeri.
Sanbornii, *Wood, in Proc. Ac. Nat. Sc. Philad.* (1868) 171.—Calif.
sapidissimum, Hedw. in Roem. Collect. 76 = fistulosum.
sarawschanicum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 244.—Turkestan.
sardoum, Moris, Stirp. Sard. Elench. fasc. ii. 10 = sphaerocephalum.
sativum, *Linn. Sp. Pl.* 296.—Europ.
Savii, Parl. Fl. Ital. ii. 554 = paniculatum.
savranicum, Bess. Cat. Sem. Hort. Crem. Suppl. (1818) = globosum.
saxatile, Bieb. Fl. Taur. Cauc. i. 264 = globosum.
saxatile, Hohen. ex Boiss. Fl. Orient. v. 263 = lepidum.
saxatile, Pall. Reise, iii. 293 = fistulosum.
saxatile, Willd. ex Ledeb. Fl. Ross. iv. 181 = albidum.
scaberrimum, Serres, in Bull. Soc. Bot. Fr. iv. (1857) 439 = sphaerocephalum.
scabrellum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 215.—Persia.
scabriflorum, *Boiss. Diagn.* Ser. I. v. 60.—Cappadoc.
scabriscapum, *Boiss. l. c.* xiii. 31.—Persia.
scabrum, Gilib. Exercit. ii. 469 = oleraceum.
scaposum, *Benth. in Pl. Hartw.* 26.—Mexic.; Texas.
scariosum, Jan, ex Schult. f. Syst. vii. 1011 = rotundum.
schergianum, *Boiss. Fl. Orient.* v. 251.—Syria.
Schmidtianum, *Tausch, in Flora*, xii. (1829) *Erg.* 45.—Bohemia.
schoenoprasioides, Fries, ex Nym. Consp. 741 = A. Schoenoprasum.
schoenoprasoides, *Regel, in Act. Hort. Petrop.* v. (1877) 630.—Reg. Himal.
Schoenoprasum, *Linn. Sp. Pl.* 301.—Reg. temp. bor.
Schoenoprasum, Regel & Tiling, Fl. Ajan. 123 = Ledebourianum.
Schrenki, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 172.—Ind. or.
Schuberti, *Zucc. in Abh. Bayer. Akad.* iii. (1843) 234. t. 3. f. 1.—Oriens.
Schulleri, Stev. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 69 = Aucheri.
scilloides, Dougl. ex S. Wats. in Proc. Am. Acad. xiv. (1879) 229 = collinum.
Scordoprassum, Aschers. Fl. Brandenb. 725 = seq.
Scorodoprasum, *Linn. Sp. Pl.* 297.—Europ.; Reg. Cauc.; Syria.
scorzoneraefolium, [*DC. in*] *Red. Lil.* ii. t. 92.—Europ. austr.
segetum, *Jan, ex Schult. f. Syst.* vii. 1020.—Sicilia.
Sellovianum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 117.—Bras.; Chili.
Semenovi, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) I. 449.—Reg. Himal.
semiretschenskianum, *Regel, in Act. Hort. Petrop.* v. (1877) 630.—As. centr.

ALLIUM:—
senescens, Ker-Gawl. in Bot. Mag. t. 1150 = glaucum.
senescens, Host, Fl. Austr. i. 420 = fallax.
senescens, *Linn. Sp. Pl.* 299.—Europ.; Sibir.
senescens, Suter, Fl. Helv. i. 193 = acutangulum.
senescens, Miq. Ann. Bot. Lugd. Bot. iii. 318 = odorum.
senescens, Thunb. Fl. Jap. 132 = japonicum.
serbicum, Vis. & Panč. in Mem. Ist. Venet. xii. (1864) 479 = moschatum.
serotinum, Lapeyr. Fl. Pyr. t. 84 = ochroleucum × suaveolens.
serotinum, Schleich. ex G. Don, in Mem. Wern. Soc. vi. (1827) 63 = senescens.
serrulatum, Boiss. Fl. Orient. v. 241 = atroviolaceum.
serratum, *S. Wats. Bot. King, Exped.* 487. t. 37. f. 4, 5.—Calif.
sessiliflorum, Fisch. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 140 = senescens.
setaceum, Waldst. & Kit. Pl. Rar. Hung. i. 70. t. 68 = moschatum.
setifolium, *Schrenk, Enum.* 6.—Soongaria.
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xli. (1868) 453.—Turkestan.
sibiricum, Linn. Mant. 562 = A. Schoenoprasum.
Sibthorpianum, Schult. f. Syst. vii. 1057 = paniculatum.
siculum, Ucria, in Roem. Arch. i. 68 = Dioscoridis.
Sieberianum, Schult. f. Syst. vii. 1099 = neapolitanum.
sikkimense, *Baker, in Journ. Bot.* xii. (1874) 292.—Reg. Himal.
sinaiticum, *Boiss. Diagn.* Ser. I. xiii. 31.—Arabia.
sindjarense, *Boiss. & Hausskn. ex Regel, in Act. Hort. Petrop.* iii. II. (1875) 121.—Mesopotamia.
sinicum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. *Art.* IV. 6; ed. II. (1827) 68.—China.
sipyleum, *Boiss. Diagn.* Ser. I. v. 58.—Lydia; Ins. Chios.
sordidum, Schur, Enum. Pl. Transs. 673 = globosum.
spathaceum, Steud. ex A. Rich. Tent. Fl. Abyss. ii. 330 = subhirsutum.
speciosum, Cyr. Pl. Rar. Neap. Fasc. ii. 13 = nigrum.
spectabile, De Not. Fl. Lig. 414 = A. Ampeloprasum.
sphaerocephalum, Crome, ex Schlecht. Fl. Berol. ii. 249 = vineale.
sphaerocephalum, *Linn. Sp. Pl.* 297.—Europ.; Oriens.
spirale, Hort. Dorp. ex Kunth, Enum. Pl. iv. 422 = spurium.
spirale, Willd. Enum. Suppl. 17 = senescens, nutans.
splendens, Miq. in Ann. Mus. Lugd. Bat. iii. 154 = Bakeri.
splendens, *Willd. ex Schult. f. Syst.* vii. 1025.—Sibiria.
spurium, *G. Don, in Mem. Wern. Soc.* vi. (1827) 59 = senescens.
spurium, Hort. ex Schult. f. Syst. vii. 1067 = glaucum.
stamineum, *Boiss. Diagn.* Ser. II. iv. 119.—Oriens.
staticiforme, *Sibth. & Sm. Fl. Graec.* iv. t. 320.—Graecia.
stellatum, *Fras. ex Ker-Gawl. in Bot. Mag.* t. 1576.—Am. bor.
Stellerianum, Bess. Cat. Hort. Crem. (1824) = albidum.
Stellerianum, *Willd. Sp. Pl.* ii. 82.—Sibiria.
Stellerianum, Ledeb. Fl. Ross. iv. 179 = flavidum.
stenopetalum, *Boiss. & Kotschy, ex Regel, in Act. Hort. Petrop.* iii. II. (1875) 231.—Cilicia.
stenopetalum, Guss. Sic. Syn. i. 394 = margaritaceum.
stenophyllum, Regel, ex Herd. in Bull. Soc. Nat. Mosc. xli. (1868) II. 446 = doloncarense.
stenophyllum, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1844) 210.—Soongaria.
stenophyllum, Wall. Cat. n. 5073 = Jacquemonti.
Stevenii, Willd. ex Ledeb. Fl. Ross. iv. 176 = globosum.
stipitatum, *Regel, in Act. Hort. Petrop.* vii. (1880) 546.—As. centr.
Stocksianum, *Boiss. Diagn.* Ser. II. iv. 117.—Beluchistan.
Stoliczki, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 160.—Tibet. occ.
Stracheyi, *Baker, in Journ. Bot.* xii. (1874) 293.—Reg. Himal.
stramineum, Boiss. & Reut. Diagn. 25 = A. Moly.
stramineum, *Regel, Gartenfl.* xxv. (1876) 355.—Manchuria.
stramineum, Schur, Enum. Pl. Transs. 674 = angulosum.

ALLIUM:—
striatellum, *Lindl. in Trans. Hort. Soc.* vi. (1851) 97.—Chili.
striatum, Jacq. Coll. Suppl. 51 = Nothoscordum striatum.
strictum, *Schrad. Hort. Goett.* t. 1.—Europ.; Oriens.
suaveolens, Gaud. Fl. Helv. ii. 491 = strictum.
suaveolens, *Jacq. Coll.* ii. 305.—Europ.
suaveolens, Duby, Bot. Gall. i. 470 = ochroleucum.
subalbidum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 128 = roseum.
subalpinum, Pall. Ind. Taur. ex Kunth, Enum. Pl. iv. 448 = decipiens.
subbiflorum, *Colla, in Mem. Acc. Torin.* xxxix. (1836) 13. t. 52.—Chili.
subhirsutum, Delile, Fl. Aegypt. ex Boiss. Fl. Orient. v. 274 = neapolitanum.
subhirsutum, Desf. Fl. Atlant. 286 = subvillosum.
subhirsutum, *Linn. Sp. Pl.* 295.—Europ.; Oriens.
subhirsutum, Reichb. Fl. Germ. Excurs. 113 = subvillosum.
subhirsutum, Sieber, ex Kunth, Enum. Pl. iv. 441 = neapolitanum.
subquinqueflorum, *Boiss. Fl. Orient.* v. 264.—Armenia.
subtilissimum, *Ledeb. Ic. Fl. Alt.* ii. 22. t. 360.—Sibir. altaic.
subvillosum, Salzm. ex Schult. f. Syst. vii. 1104 = subhirsutum.
sulcatum, [DC. in] Red. Lil. t. 482 = neapolitanum.
Sulvia, Buch.-Ham. in D. Don, Prod. Fl. Nep. 53 = fragrans.
Suworowi, *Regel, in Act. Hort. Petrop.* vii. (1880) 546.—As. centr.
Synnotia, G. Don, in Mem. Wern. Soc. vi. (1827) 19 = rotundum.
Synnotianum, Schult. f. Syst. vii. 1015 = praec.
syntamanthum, C. Koch, in Linnaea, xxii. (1849) 238 = rubellum.
syriacum, Boiss. Fl. Orient. v. 232 = A. Ampeloprasum.
Szovitsi, Regel, in Act. Hort. Petrop. iii. II. (1875) 171 = strictum.
szurulense, *Lerch. ex Kanitz, in Verh. Siebenb. Ver. Naturw.* xxxiv. (1884) 55.—Europ.
talassicum, *Regel, in Act. Hort. Petrop.* v. (1877) 628.—Turkestan.
tartaricum, [Dryand. in] Ait. Hort. Kew. ed. II. ii. 233 = odorum.
tartaricum, Hort. ex Schult. f. Syst. vii. 1066 = nutans.
tartaricum, *Linn. f. Suppl.* 196.—Rossia; Sibiria; Persia.
tauricolum, Boiss. Diagn. Ser. II. iv. 118 = frigidum.
tauricum, Hort. ex Schult. f. Syst. vii. 1051 = flavum.
tauricum, Pall. ex G. Don, in Mem. Wern. Soc. vi. (1827) 50 = globosum.
tchefouense, *Debeaux, in Actes Linn. Soc. Bord.* xxxii. (1878) 25; *Fl. Tché-Fou*, 138.—China.
Tchihatschewii, *Boiss. in Tchihat. Asie Min. Bot.* ii. 551.—Armenia.
Tenorii, Spreng. Pugill. i. 29; Syst. ii. 35 = roseum.
tenue, C. Koch, in Linnaea, xxii. (1849) 238 = rubellum.
tenue, Frivald. ex Griseb. Spicil. Fl. Rumel. ii. 396 = erythraeum.
tenue, G. Don, in Mem. Wern. Soc. vi. (1827) 34 = Pallasii.
tenue, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 206.—Turkestan.
tenuiflorum, Delastre, Fl. Vienne, 420 = sphaerocephalum.
tenuiflorum, Tenore, Fl. Nap. i. 165. t. 30 = paniculatum.
tenuifolium, *Frivald. in Syll. Ratisb.* ii. 33.—Europ.
tenuifolium, Salisb. Prod. 235 = A. Schoenoprasum.
tenuifolium, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 76 = globosum.
tenuissimum, Habl. Taur. 162 = moschatum.
tenuissimum, *Linn. Sp. Pl.* 301.—Sibiria.
teretifolium, *Regel, in Act. Hort. Petrop.* v. (1877) 629.—Turkestan.
thessalum, Boiss. Fl. Orient. v. 232 = A. Ampeloprasum.
Thomsoni, *Baker, in Journ. Bot.* xii. (1874) 294.—Reg. Himal.
Thunbergii, *G. Don, in Mem. Wern. Soc.* vi. (1827) 84.—Japon; China.

ALLIUM :—
tianschanicum, Rupr. in Mém. Acad. Pétersb. Sér. III. xiv. (1869) 33. (Sert. Tiansch. 33) = hymenorhizum.
Tilingi, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 124. —Calif.
Tinei, J. & C. Presl, Delic. Prag. 145 = subhirsutum.
Tolmiei, Baker, ex Coult. Rocky Mt. Bot. 349 = tribractcatum.
tortuosum, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 76.—Europ.
tortuosum, Willd. ex Schult. f. Syst. vii. 1064 = lineare.
Tourneuxii, Boiss. Fl. Orient. v. 274 = roseum.
trachyanthum, Griseb. Spicil. Fl. Rumel. ii. 395 = rotundum.
trachypus, Boiss. & Sprun. Diagn. Ser. I. vii. 114 = sphaerocephalum.
transsilvanicum, *Lerch. ex Kanitz, in Verh. Siebenb. Ver. Naturw.* xxxiv. (1884) 54.—Europ.
Trautvetterianum, *Regel, in Act. Hort. Petrop.* viii. (1884) 661.—Bokhara.
tribractcatum, *Torr. in Pacif. Rail. Rep.* iv. 148.—Calif.
trichocnemis, *J. Gay, in Ann. Sc. Nat.* Sér. III. viii. (1847) 210.—Algeria.
tricoccum, [*Solandr. in*] *Ait. Hort. Kew.* i. 428.—Am. bor.
tricoccum, Blanco, Fl. Filip. ed. I. 239 = uliginosum.
tricorne, Desv. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 145 = angulosum.
tricorne, Hort. ex Poir. Encyc. Suppl. i. 270 = cernuum.
triflorum, *Rafin. in Med. Repos. New York*, v. (1808) 362.—Pennsylv.
trifoliatum, *Cyr. Pl. Rar. Neap. Fasc.* ii. 11.—Europ.; Orient.
tripedale, Trautv. in Act. Hort. Petrop. ii. (1873) 485 = Dioscoridis.
triquetrum, Hort. ex Schrad. in Schult. f. Syst. vii. 1078 = angulosum.
triquetrum, *Linn. Sp. Pl.* 300.—Europ.; Afr. bor.
triquetrum, Lour. Fl. Cochinch. 202 = Thunbergii.
triquetrum, Sebast. & Maur. Prod. Fl. Rom. 126 = pendulinum.
triste, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845). —Orient.
trisulcum, Hort. Genev. ex Schult. f. Syst. vii. 1067 = glaucum.
Tschonoskianum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 160.—Japon.
Tschulpias, *Regel, l. c.* 107.—Turkestan.
tuberosum, *Rottl: ex Spreng. Syst.* ii. 38.—Malabar.
tuberosum, Roxb. Hort. Beng. 24 = odorum.
tulipaefolium, Ledeb. Fl. Alt. ii. 9; Ic. t. 37 = decipiens.
turkestanicum, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 197.—Turkestan.
turtschicum, *Regel, l. c.* vi. (1879) 529.—As. centr.
tythocephalum, *Schult. f. Syst.* vii. 1133.—Sibiria.
uliginosum, G. Don, in Mem. Wern. Soc. vi. (1827) 60 = odorum.
uliginosum, Kanitz, in Linnaea, xxxii. (1863) 399 = angulosum.
uliginosum, Ledeb. Fl. Alt. ii. 16 = Ledebourianum.
umbellatum, Hall. ex Steud. Nom. ed. I. 29 = odorum.
umbilicatum, *Boiss. Diagn.* Ser. II. iv. 113.—Beluchistan.
undulatum, Fisch. ex Steud. Nom. ed. I. 29 = caspicum.
undulatum, Schousb. ex Trev. All. Sp. 8. n. 9 = nutans.
unifolium, *Kellogg, in Proc. Calif. Acad. Nat. Sc.* ii. (1863) 112. f. 35.—Calif.
uratense, *Franch. in Nouv. Arch. Mus. Hist. Nat. Par.* Sér. II. vii. (1884) 112 (*Pl. Davidian.* i. 304).—As. or.
urceolatum, *Regel, Gartenfl.* (1873) 236; *et in Act. Hort. Petrop.* ii. (1873) 405.—Turkestan.
ursinoides, G. Don, ex Sweet, Hort. Brit. ed. I. 422 = ursinum.
ursinum, *Linn. Sp. Pl.* ed. I. 300.—Europ.; As. bor.
Urvillei, Heldr. & Sart. ex Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 233 = staticiforme.
vaginatum, Pančić ex Boiss. Fl. Orient. v. 266 = Cupani.
valdense, Reut. in Comptes-rend. Soc. Hallér. (1854–56) 126 = flavum.
valdensium, Nym. Consp. 740 = flavum.

ALLIUM :—
validum, *S. Wats. in Bot. King, Exp.* 350.—Am. bor.
variegatum, *Boiss. Diagn.* Ser. I. vii. 118.—Assyria.
veratrifolium, *Griseb. ex Tchihat. Asie Min. Bot.* ii. 555.—As. Min.
vernale, Tineo, in Guss. Fl. Sic. Prod. Suppl. i. 96 = subvillosum.
veronense, Linn. Syst. ed. X. 978 = paniculatum.
veronense, Spreng. Pugill. ii. 60 = sphaerocephalum.
verrucosum, G. Don, in Mem. Wern. Soc. vi. (1827) 20 = rotundum.
versicolor, *Nem. ex Chaub, in Actes Linn. Soc. Bord.* xix. (1853) 50.—Europ.
verticillatum, *Regel, in Act. Hort. Petrop.* vi. (1875) 518.—Turkestan.
Victorialis, *Linn. Sp. Pl.* 295.—Europ.; Caucas.; Sibir.
Vincetoxicum, Pall. ex Ledeb. Fl. Ross. iv. 186 = ursinum.
vineale, Sch. Bip. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 46 = sphaerocephalum.
vineale, *Linn. Sp. Pl.* 299.—Europ.
violaceum, Gilib. Exercit. ii. 470 = arenarium.
violaceum, Wall. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 143 = Wallichii.
violaceum, Willd. Enum. Suppl. 17 = carinatum.
virescens, Guss. in Tenore, Syll. 164 = margaritaceum.
virescens, Lam. Fl. Fr. iii. 259 = oleraceum.
virens, Lam. Encyc. i. 67 = olcraceum.
viridi-album, Tineo, Cat. Pl. Hort. Panorm. (1827) 275 = sphaerocephalum.
viridulum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 855 = delicatulum.
viridulum, *Ledeb. Ic. Fl. Alt.* iii. t. 378.—Sibir. altaic.
viviparum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 852.—Sibiria.
volhynicum, Bess. Cat. Hort. Crem. Suppl. iii. 2 = strictum.
vulcanicum, Boiss. Diagn. Ser. I. xiii. 33 = rubellum.
Waldsteinianum, Schult. f. Syst. vii. 1011 = seq.
Waldsteini, G. Don, in Mem. Wern. Soc. vi. (1827) 17 = rotundum.
Wallichianum, Hort. Leichtl. ex Regel, in Act. Hort. Petrop. iii. II. (1875) 232 = acuminatum.
Wallichianum, *Steud. Nom.* ed. II. i. 53.—Reg. Himal.
Wallichii, Kunth, Enum. Pl. iv. 443 = praec.
Webbii, Clem. in Mem. Acc. Sc. Torino, Ser. II. xvi. (1857) 327. t. 8 = flavum.
Weissii, *Boiss. Fl. Orient.* v. 253.—Ins. Cyclad.
Welwitschi, *Regel, in Act. Hort. Petrop.* iii. II. (1875) 66.—Lusit.
Weschniakowi, *Regel, l. c.* vi. (1879) 531.—Turkestan.
Wiedemannianum, *Regel, l. c.* iii. II. (1875) 199.—Anatolia.
Wierzbickii, *Heufel, ex Steud. Nom.* ed. II. i. 55.—Hungar..
Willdenovi, Kunth, Enum. Pl. iv. 452 = delicatulum.
Wildii, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 232 = A. Ampeloprasum.
Winklerianum, *Regel, in Act. Hort. Petrop.* viii. (1884) 661.—Bokhara.
xanthicum, Griseb. & Schrenk, in Wiegm. Archiv, xviii. (1852) I. 358 = globosum.
zebdanense, *Boiss. & Noë, Diagn.* Ser. II. iv. 113.—Syria; Armenia.

ALLMANIA, R. Br. in Wall. Cat. n. 6890-92 (1832). *AMARANTACEAE*, Benth. & Hook. f. iii. 27.
albida, *R. Br. l. c.* n. 6891.—Ind. or.; Malaya.
aspera, Wight, Ic. t. 1770 = nodiflora.
dichotoma, Wight, l. c. t. 1771 = nodiflora.
esculenta, R. Br. in Wall. Cat. n. 6892 = nodiflora.
nodiflora, *R. Br. l. c.* n. 6890.—Ind. or.; Malaya.
procumbens, Wight, ex Hook. f. Fl. Brit. Ind. iv. 717 = nodiflora.

ALLOBIA, Rafin. Fl. Tellur. iv. 116 (1836) = **Euphorbia**, Linn.
portlandica, Rafin. l. c. = E. portlandica.

ALLOBIUM, Miers, in Ann. Nat. Hist. Ser. II. viii. (1881) 178, 179 = **Phoradendron**, Nutt. (Loranth.).

ALLOBORGIA, Steud. Nom. ed. II. i. 55 (1840) = seq.

ALLOBROGIA, Tratt. Fl. Austriac. ii. 71 (1792) = **Paradisea**, Mazz. (Liliac.).

ALLOCARPUS, H. B. & K. Nov. Gen. et Sp. iv. 291. t. 405 (1820) = **Calea**, Linn. (Compos.).
caracasanus, H. B. & K. l. c. 292.—Venezuela.
integrifolius, DC. Prod. v. 676 = C. integrifolia.
Lindenii, Sch. Bip. ex Wedd. Chlor. And. i. 74.—N. Granat.
sabazioides, Less. Syn. Comp. 242 = C. sabazioides.
scabrifolius, Hook. & Arn. Bot. Beech. Voy. 300 = C. scabrifolia.

ALLOCERATIUM, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 129, 135 = **Diptychocarpus**, Trautv. (Crucif.).
strictum, Hook. f. & Thoms. l. c. 135 = D. strictus.

ALLOCHLAMYS, Moq. in DC. Prod. xiii. II. 463 (1849) = **Pleuropetalum**, Hook. f. (Amarant.).
Darwinii, Moq. l. c. = P. Darwinii.

ALLOCHRUSA, Bunge, in Boiss. Fl. Orient. i. 559 (1867) = **Acanthophyllum**, C. A. Mey. (Caryophyll.).
Bungei, Boiss. l. c. 560 = Ac. Bungei.
persica, Boiss. l. c.—Persia.
pulchella, Bunge, ex Boiss. l. c. = *persica*.
versicolor, Boiss. l. c. 559 = Ac. versicolor.

ALLODAPE, Endl. Gen. 749 (1839) = **Lebetanthus**, Endl. (Epacrid.).
americana, Endl. l. c. = L. americanus.

ALLODAPHNE, Steud. Nom. ed. II. i. 55 (1840) = praec.

ALLOEOSPERMUM, Spreng. Anleit. ii. 901 (1818) = ALLOISPERMUM, Willd. = **Calea**, Linn. (Compos.).

ALLOHEMIA, Rafin. Sylva Tellur. 125 (1838) = **Loranthus**, Linn.
pedunculata, Rafin. l. c. = L. pedunculatus.
purpurea, Rafin. l. c. = L. occidentalis.

ALLOIANTHEROS, Steud. Syn. Pl. Gram. 217 (1855) = seq.

ALLOIATHEROS, Ell. Sketch, i. 146 (1817) = **Gymnopogon**, Beauv. (Gramin.).
ambiguus, Ell. (cf. Sketch, i. 146) ex Kunth, Enum. i. 284 = G. racemosus.
aristatus, Rafin. in Ser. Bull. Bot. i. (1830) 221 = G. racemosus.
lepturoides, Steud. Nom. ed. II. i. 55 = G. racemosus.

ALLOIOZONIUM, Kunze, in Linnaea, xvii. (1843) 572 = **Cryptostemma**, R. Br. (Compos.).
arctotideum, Kunze, l. c. = C. calendulaceum.

ALLOISPERMUM, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 139 = **Calea**, Linn. (Compos.).
divaricatum, Willd. l. c. (= *Allocarpus caracasanus*).

ALLOMIA, DC. Prod. vii. 310 (1838) = **Alomia**, H. B. & K. (Compos.).

ALLOMORPHIA, Blume, in Flora, xiv. (1831) 522. *MELASTOMACEAE*, Benth. & Hook. f. i. 752.
exigua, *Blume, l. c.* 523.—Ind. or.; Malaya.
Griffithii, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 74.—Malacca.
hispida, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) II. 53.—Burma.
ovalifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 74.—Ins. Fiji.
pauciflora, Benth. in Hook. Lond. Journ. Bot. i. (1842) 485 = Oxyspora pauciflora?
umbellulata, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 74.—Burma.

ALLOPHYLLUS, Gleditsch, Syst. 82 (1764) = seq.

ALLOPHYLUS, Linn. Fl. Zeyl. 58 (1747); Diss. Dass. 14 (1747); Am. Acad. i. 398 (1749). *SAPINDACEAE*, Benth. & Hook. f. i. 396.
TOXICODENDRUM, Gaertn. Fruct. i. 207. t. 44 (1788).
USTUBIS, Burm. f. Fl. Ind. 89. t. 32. f. i. (1768).
africanus, Beauv. Fl. Owar. ii. 54. t. 107 = zeylanicus.
amboinensis, *Blume, Rumphia*, iii. 129.—Amboina.
aporeticus, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 185.—Burma.
Blancoi, Blume, Rumphia, iii. 129 = A. Cobbe.
Cambessedei, *Blume, l. c.*—Malaya.

ALLOPHYLUS:—
celebicus, *Blume, Rumphia*, iii. 128.—Ins. Celebes.
Cobbe, *Blume, l. c.* 131.— As. or.
Cominia, Sw. Prod. Veg. Ind. Occ. 62 = S. Cominia.
Commersoni, *Blume, Rumphia*, iii. 129.—Ins. Borbon.
dimorphus, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 126.—Ins. Philipp.
filiger, *Radlk. l. c.*—Ins. Philipp.
fulvinervis, *Blume, Rumphia*, iii. 183.—Java.
integrifolius, *Blume, l. c.* 129.—Malaya.
javensis, *Blume, l. c.* 126.—Java.
lanatus, Roxb. Hort. Beng. 88 = A. Cobbe.
leptococcus, *Radlk. in Act. Congr. Bot. Amst.* 1877 (1879) 126.—Ins. Key.
ligustrinus, *Blume, Rumphia*, iii. 126.—Sumatra.
litoralis, *Blume, l. c.* 124.—Java.
nigrescens, *Blume, l. c.* 129.—Madag.
ornitrophioides, Roxb. ex Wall. Cat. n. 8055 = A. Cobbe.
Pervillei, *Blume, Rumphia*, iii. 123.—Madag.
pinnatus, Roxb. Ic. t. 1407 = Sapindus fruticosus.
racemosus, Sw. Prod. Veg. Ind. Occ. 62 = S. occidentalis.
rhusiphyllus, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 507.—Ins. Socotra.
rigidus, Sw. Prod. Veg. Ind. Occ. 62 = S. rigida.
rufescens, *Blume, Rumphia*, iii. 159.—Borneo.
rugosus, *Blume, l. c.* 138.—Java.
salignus, *Blume, l. c.* 129.—Madag.
serratus, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 185.—Burma.
sessilis, *Blume, Rumphia*, iii. 125.—Java.
sumatranus, *Blume, l. c.* 132.—Sumatra.
sundanus, Miq. Fl. Ind. Bat. i. II. 575 = A. Cobbe.
ternatus, Lour. Fl. Cochinch. 232 = A. Cobbe.
ternatus, Roxb. ex Steud. Nom. ed. II. i. 55 = S. zeylanicus.
timorensis, *Blume, Rumphia*, iii. 130.—Timor.
timorensis, Miq. Fl. Ind. Bat. ii. I. 575 = A. Cobbe.
zeylanicus, *Linn. Sp. Pl.* 348.—Himal.

ALLOPLECTUS, Mart. Nov. Gen. et Sp. iii. 53. t. 223 (1829). *GESNERACEAE*, Benth. & Hook. f. ii. 1008.
ANISOPLECTUS, CALANTHUS, CALOPLECTUS, CALYCOPLECTUS, CORYTOPLECTUS, ERYTHRANTHUS, PRIONOPLECTUS, SACCOPLECTUS, Oerst. in Danske Vid. Selsk. Skr. V. v. (1858) 117–121.
CRANTZIA, Scop. Introd. 173 (1777).
HEINTZIA, Karst. Ausw. Gew. Venez. 34. t. 11. (1848).
GLOSSALOMA, Hanst. in Linnaea, xxvi. (1853–55) 209.
LOPHIA, Desv. in Ham. Prod. Fl. Ind. Occ. 47 (1825).
LOPHALIX, Rafin. Sylva Tellur. 70 (1838).
MACROCHLAMYS, Decne. in Rev. Hortic. (1849) 243.
OROBANCHIA, Vand. Fl. Lusit. Spec. 41. f. 18, 19 (1788).
POLYSTHYSANIA, Hanst. ex Oerst. in Danske Vid. Selsk. Skr. V. v. (1858) 121.
PTERYGOLOMA, Hanst. in Linnaea, xxvi. (1853–1855) 211.
TYSSACIA, Reichb. ex Steud. Nom. ed. II. ii. 727 (1841).
VIREYA, Rafin. Specch. i. 194 (1814).
angustifolius, *DC. Prod.* vii. 546.—Bras.
bicolor, *Klotzsch, ex Hanst. in Linnaea*, xxxiv. (1865–66) 377.—Bras.
bracteatus, *Linden, Cat.* n. 5 (1850) 2.—Hab.?
calcaratus, *G. Don, Gen. Syst.* iv. 655.—Reg. Andin.
capitatus, *Hook. Bot. Mag.* t. 4452.—Am. austr.
chrysanthus, *Planch. & Linden, in Fl. des Serres*, Sér. I. viii. (1852–53) 211.—N. Granat.
circinnatus, *Mart. Nov. Gen. et Sp.* iii. 55. t. 223. f. 1.—Bras.
coccineus, *Mart. l. c.* 57.—Guiana.
concolor, Hook. Bot. Mag. t. 4371 = dichrous.
coriaceus, *Hanst. in Linnaea*, xxxiv. (1865–66) 363.—Costa Rica.
corymbosus, *Klotzsch, ex Hanst. l. c.* 371.—Peruv.
cristatus, *Mart. Nov. Gen. et Sp.* iii. 57.—Ind. occ.
darienensis, *Seem. Bot. Voy. Herald*, 187.—Darien.
dichrous, *DC. Prod.* vii. 546.—Bras.
dichrous, Hook. Bot. Mag. t. 4216 = sparsiflorus.
dimidiatus, *Benth. Pl. Hartw.* 235.—N. Granat.
eriocalyx, Galeott. ex Hook. Bot. Mag. t. 4371 = dichrous.
Forseithii, *Hanst. in Linnaea*, xxxiv. (1865–66) 375.—Am. centr.

ALLOPLECTUS:—
glaber, *DC. Prod.* vii. 546.—Mexic.
hispidus, *Mart. Nov. Gen. et Sp.* iii. 57.—Am. austr.
ichthyoderma, *Hanst. in Linnaea*, xxxiv. (1865–66) 372.—Costa Rica.
Kunthii, *G. Don, Gen. Syst.* iv. 655.—N. Granat.
lanatus, *Seem. Bot. Voy. Herald*, 186.—Panama.
macrophyllus, *Hemsl. Biol. Centr. Am. Bot.* ii. 484.—Costa Rica.
melittifolius, Mart. ex Loud. Encyc. Pl. 1402 = Tussacia pulchella.
meridensis, *Klotzsch, ex Hanst. in Linnaea*, xxxiv. (1865–66) 373.—N. Granat.
multiflorus, *Hanst. l. c.* 361.—Costa Rica.
parviflorus, *Hemsl. Biol. Centr. Am. Bot.* ii. 484.—Costa Rica.
Patrisii, *DC. Prod.* vii. 545.—Guiana.
peltatus, *Oliver, in Bot. Mag. t.* 6333.—Am. centr.
pendulus, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 6. t. 205.—Peruv.
Pinelianus, *Lem. in Fl. des Serres.* Sér. I. ii.(Août 1846) t. v.—Bras.
repens, *Hook. Bot. Mag.* t. 4250.—Ind. occ.
sanguineus, *Mart. Nov. Gen. et Sp.* iii. 57.—Am. austr.
scabridus, Lem. Jard. Fleur. i. (1851) Misc. 51 = glaber.
Schlimii, *Planch. & Linden, in Fl. des Serres*, Sér. I. viii. (1852–53) 211. t. 827.—N. Granat.
Schottii, G. Don, Gen. Syst. iv. 655 = dichrous.
semicordatus, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 5.—Peruv.
sparsiflorus, *Mart. Nov. Gen. et Sp.* iii. 55.—Bras.
speciosus, Hort. ex Hook. Bot. Mag. t. 4452 = capitatus.
speciosus, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 6.—Chili.
strigosus, *Hanst. in Linnaea*, xxxiv. (1865–66) 374.—Mexic.
tenuis, *Benth. Pl. Hartw.* 235.—Peruv.
tetragonus, *Hanst. in Linnaea*, xxxiv. (1865–66) 368.—Costa Rica.
tigrinus, *Hanst. l. c.* 366.—N. Granat.
villosus, *Hanst. in Mart. Fl. Bras.* viii. 407.—Bras.
vittatus, *André, in Illust. Hortic.* xvii. (1870) t. 13.—Peruv.
zamorensis, *Linden & André, in Illust. Hortic.* xix. (1872) 110.—N. Granat.

ALLOPLEIA, Rafin. Med. Fl. ii. 109 (1830) = **Veronica**, Tourn. (Scrophul.).
rotundifolia, Rafin. l. c. = V. rotundifolia.

ALLOPYTHION, Schott, Gen. Aroid. t. 24 (1858) = **Thomsonia**, Wall. (Aroid.).
Hookeri, Schott, l. c. = T. Hookeri.

ALLOSAMPELA, Rafin. Med. Fl. ii. 122 (1830) = **Vitis**, Linn. (Ampel.).
heterophylla, Rafin. l. c. = V. heterophylla.

ALLOSCHEMONE, Schott, Prod. Syst. Aroid. 358 (1860) = **Monstera**, Adans. (Aroid.).
Poeppigiana, Schott, l. c. = M. occidentalis.

ALLOSPERMA, Rafin. Fl. Tellur. iv. 122 (1836) = **Commelina**, Linn.
tuberosa, Rafin. l. c. = C. tuberosa.

ALLOSTIS, Rafin. Sylva Tellur. 104 (1838) = **Baeckea**, Linn. (Myrtac.).

ALLOTEROPSIS, C. Presl, Rel. Haenk. i. 343. t. 47 (1830). Gen. dub. *GRAMIN.* Benth. & Hook. f. iii. 1096.
distachya, *C. Presl, l. c.*—Calif.

ALLOTRIA, Rafin. Fl. Tellur. iii. 70 (1836) = **Commelina**, Linn.
scabra, Rafin. l. c. = C. scabra.

ALLOTROPA, A. Gray, in Pacif. Rail. Rep. Bot. iv. 81 (1855); et in Proc. Am. Acad. vii. (1868) 368. *MONOTROPEAE*, Benth. & Hook. f. ii. 607.
virgata, *Torr. & Gray, in Proc. Am. Acad. l. c.*—Calif.

ALLOUYA, Plum. ex Aubl. Pl. Gui. ii. 931 (1775) = **Maranta**, Linn. (Scitam.).

ALLOZYGIA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 309. t. 15 = **Oxyspora**, DC. (Melast.).
cernua, Naud. l. c. = O. cernua.

ALLUGHAS, Steud. Nom. ed. I. 29 (1821) = ALUGHAS, Lindl. = **Alpinia**, Linn. (Scitam.).

ALMANA, Rafin. Fl. Tellur. iv. 68 (1836) = **Gloxinia**, Linn. (Gesner.).
hirsuta, Rafin. l. c. = G. hirsuta.

ALMEIDA, Cham. in Linnaea, v. (1830) 56 = seq.

ALMEIDEA, St. Hil. in Bull. Soc. Philom. (1823) 129. *RUTACEAE*, Benth. & Hook. f. i. 284.
ARUBA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 152. t. 19, 27, 29.
acuminata, *St. Hil. ex G. Don, Gen. Syst.* i. 798.—Bras.
alba, *St. Hil. ex G. Don, l. c.*—Bras.
caerulea, *St. Hil. ex G. Don, l. c.*—Bras.
lanceolata, *St. Hil. Pl. Rem.* 144.—Bras.
lilacina, *St. Hil. in Bull. Soc. Philom.* (1823) 129.—Bras.
longifolia, *St. Hil. l. c.*—Bras.
longipes, *Presl, Bot. Bemerk.* 32.—Bras.
macropetala, *Fisch. & Mey. Sert. Petrop.* t. 10.—Bras.
myriantha, *Cham. in Linnaea*, v. (1830) 56.—Bras.
rubra, *St. Hil. in Bull. Soc. Philom.* (1823) 129.—Bras.

ALMELOVEENIA, Dennst. Schluess. Hort. Malab. 32 (1818) = **Moullava**, Adans. (Inc. Sed.).
spinosa, Dennst. l. c.—Ind. or.

ALMIDEIA, Reichb. Nom. 197 (1841) = **Almeidea**, St. Hil. (Rutac.).

ALMYRA, Salisb. in Trans. Hort. Soc. i. (1812) 336 = **Pancratium**, Linn. (Amaryll.).
illyrica, Salisb. Gen. Pl. Fragm. 109 = P. illyricum.
parviflora, Salisb. l. c. = P. parviflorum.
stellaris, Salisb. in Trans. Hort. Soc. i. (1812) 336 = P. illyricum.

ALNASTER, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 200 = **Alnus**, Tourn. (Cupul.).
fruticosus, Ledeb. Fl. Ross. iii. 655 = Alnus viridis.
viridis, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 201 = Alnus viridis.

ALNOBETULA, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853); (Enum. Pl. Trans.) 68 = **Alnus**, Tourn. (Cupul.).
viridis, Schur, l. c. = Alnus viridis.

ALNUS, [Tourn.] Linn. Syst. ed. II. (1735); Fl. Lapp. 260 (1737). *CUPULIFERAE*, Benth. & Hook. f. iii. 404.
ALNASTER, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 200.
ALNOBETULA, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858); (Enum. Pl. Trans. 68. n. 614).
CLETHROPSIS, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 201.
SEMIDOPSIS, Zumag. Fl. Pedem. i. 249 (1849).
acuminata, *H. B. & K. Nov. Gen. et Sp.* ii. 20.—Am. austr.
Alnobetula, C. Koch, Dendrol. ii. I. 625 = viridis.
alpina, Borckh. Handb. Forstb. i. 477 = viridis.
americana, Hort. ex C. Koch, Dendrol. ii. I. 636 = serrulata.
arguta, *Benth. Pl. Hartw.* 348.—Mexic.
arguta, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 205 = acuminata.
aurea, Hort. ex C. Koch, Dendrol. ii. I. 630 = glutinosa.
autumnalis, Hartig, ex Garcke, Fl. Deutschl. ed. IV. 305 = serrulata.
badensis, Lange, ex Regel, in DC. Prod. xvi. II. 188 = glutinosa.
barbata, Fries, ex Nym. Consp. 671 = pubescens.
barbata, C. A. Mey. Verz. Pfl. Cauc. 43 = glutinosa.
brembana, Rota, Prosp. Prov. Bergamasco, 79 = viridis.
canadensis, Lodd. ex Loud. Hort. Brit. 378 = serrulata.
castanaefolia, *Mirb. in Mém. Mus. Par.* xiv. (1827) 463. t. 21.—Peruv.
cerifera, Hartig, ex Regel, in DC. Prod. xvi. II. 187 = glutinosa.
communis, Desf. Tabl. Hort. Par. ed. I. 213 = glutinosa.
cordata, Desf. l. c. ed. II. 244 = cordifolia.
cordifolia, *Tenore, Fl. Nap. Prod.* 54. t. 99.—Europ.; Oriens.
corylifolia, *Kern. ex Dalla Torre, Atl. Alpenfl.* 199.—Europ.

ALNUS:—
crispa, Hort. ex Steud. Nom. ed. II. i. 55 = undulata.
crispa, *Pursh, Fl. Am. Sept.* ii. 181.—Am. bor.
cuneata, Geyer, ex Walp. Ann. i. 286 = Cercocarpus parvifolius.
denticulata, C. A. Mey. Verz. Pfl. Cauc. 43 = glutinosa.
dioica, Roxb. Fl. Ind. iii. 580 = Aporosa Roxburghii.
dubia, Req. ex Regel, in DC. Prod. xvi. II. 187 = glutinosa.
elliptica, Req. in Ann. Sc. Nat. Sér. I. v. (1825) 381 = glutinosa.
emarginata, Krock. ex Steud. Nom. ed. I. 29 = glutinosa.
februaria, Kuntze, Taschen. Fl. Leipz. 238 = glutinosa.
ferruginea, *H. B. & K. Nov. Gen. et Sp.* ii. 17.—N. Granat.
firma, *Sieb. & Zucc. in Abh. Acad. Muench.* iv. III. (1845) 230.—Japon.
fruticosa, Rupr. Beitr. Pfl. Russ. Reich. ii. 53 = viridis.
glauca, Michx. f. Arb. Am. iii. 320. t. 4. f. 2 = incana.
glutinosa, *Medic. Pfl. Anat.* 393.—Europ.; As. bor.
Harinoki, Siebold, in Verh. Gen. Bat. xii. (1830) 25 = japonica.
hirsuta, Turcz. in Bull. Soc. Nat. Mosc. (1838) 101; xxvii. (1854) I. 406 = incana.
hybrida, Reichb. Ic. Fl. Germ. xii. t. 630. f. 1292 = glutinosa, incana.
incana, *Medic. Pfl. Anat.* 393.—Reg. bor. temp.
incisa, Steud. Nom. ed. I. 29 = glutinosa.
integrifolia, Roxb. ex Steud. Nom. ed. II. i. 56 = Aporosa Roxburghii.
intermedia, Hort. ex C. Koch, Dendrol. ii. I. 631 = oblongata.
intermedia, Schrad. ex Regel, in DC. Prod. xvi. II. 189 = incana.
japonica, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1845) 230 = maritima.
japonica, Steud. Nom. ed. II. i. 55 = Betula alba.
jorullensis, Benth. Pl. Hartw. 52 = acuminata.
jorullensis, *H. B. & K. Nov. Gen. et Sp.* ii. 20.—Mexic.
laciniata, Ehrh. Beitr. iii. 22 = glutinosa.
lanuginosa, Gilib. Exercit. ii. 402 = incana.
latifolia, Desf. Cat. Hort. Par. ed. III. 352 = serrulata.
Lindeni, *Regel, in Nouv. Mém. Soc. Nat. Mosc.* xiii. (1861) 144.—As. temp.
lobata, Nym. Consp. 671 = incana.
longifolia, Bové, ex Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 348 = orientalis.
macrocarpa, Lodd. ex Loud. Hort. Brit. 378 = glutinosa.
macrocarpa, Req. ex Nym. Consp. 672 = cordifolia.
macrophylla, Desf. ex Steud. Nom. ed. I. 29; ed. II. i. 56 = serrulata.
maritima, *Nutt. Sylv. Am. Suppl.* i. 34. t. 10 (1842).—Am. bor.
microphylla, *Arvet, in Bull. Soc. Bot. Fr.* xxvi. (1879) *Bibl.* 39.—Europ.
Mirbellii, *Spach, in Ann. Sc. Nat.* Sér. II. xv. (1841) 204.—Am. austr.
Mitchelliana, M. A. Curtis, ex A. Gray, in Am. Journ. Sc. xlii. (1842) 42 = crispa.
montana ×, *Brucgg. in Jahresb. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 115.—Europ.
Morisiana, Bertol. Fl. Ital. x. 163 = glutinosa.
nana, [*Clairv.*] *Man. Herb.* 280.—Europ.
neapolitana, Savi, Tratt. Alb. Tosc. ed. II. ii. 21 = cordifolia.
nepalensis, *D. Don, Prod. Fl. Nep.* 58.—Reg. Himal.
nigra, Gilib. Exercit. ii. 401 = glutinosa.
nitens, C. Koch, in Linnaea, xxii. (1849) 334 = glutinosa.
nitida, *Endl. in Gen. Suppl.* iv. II. 20.—Reg. Himal.
obcordata, C. A. Mey. ex Steud. Nom. ed. II. i. 56, sphalm. = cordifolia.
oblongata, Kotschy, ex Regel, in DC. Prod. xvi. II. 185 = orientalis.
oblongata, Regel, Monog. 113. t. 7. f. 3–9 = maritima.
oblongata, *Willd. Sp. Pl.* iv. 335.—Europ. austr.
oblongifolia, *Torr. Bot. Mex. Bound.* 204.—N. Mexic.
oregana, Nutt. Sylv. i. 9 = rubra.
orientalis, *Decne. in Ann. Sc. Nat.* Sér. II. iv. (1835) 348.—Orient.
ovata, Lodd. Bot. Cab. t. 1141 = crispa.
oxyacanthifolia, Lodd. Cat. (1836) = glutinosa.
plicata, Hort. ex C. Koch, Dendrol. ii. I. 638 = incana.
prunifolia, Hort. ex C. Koch, l. c. 630 = glutinosa.
pubescens, Tausch, in Flora, xvii. (1834) 520 = glutinosa × incana.
pumila, *Steud. Nom.* ed. I. 29.—Am. bor.

ALNUS:—
quercifolia, Willd. Berl. Baumz. 45 = glutinosa.
repens, Wormsk. Fl. Dan. t. 2738 = viridis.
rhombifolia, *Nutt. Sylva*, i. 33.—Calif.
rotundifolia, Bertol. Fl. Ital. x. 160 = cordifolia.
rotundifolia, Stokes, Bot. Mat. Med. iv. 369 = glutinosa.
rubra, *Bongard, in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 162.—Am. bor. occ.
rubra, Desf. ex Steud. Nom. ed. I. 29 = serrulata.
rufescens, Liebm. ex Hemsl. Biol. Centr. Am. Bot. iii. 165 = acuminata.
rugosa, Hort. ex Regel, in DC. Prod. xvi. II. 188 = serrulata.
rugosa, Spreng. Syst. iii. 848 = acuminata.
serrulata, *Willd. Sp. Pl.* iv. 336.—Am. bor.
sibirica, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. (1838) 101; xxvii. (1854) I. 406 = incana.
suaveolens, Mor. ex Nym. Consp. 671 = glutinosa.
suaveolens, *Req. in Ann. Sc. Nat.* v. (1825) 381.—Europ.
subcordata, C. A. Mey. Verz. Pfl. Cauc. = cordifolia.
subrotunda, *Desf. Tabl. Hort. Par.* ed. II. 244.—Hab.?
tiliacfolia, Hort. ex C. Koch, Dendrol. ii. I. 634 = subcordata.
tenuifolia, *Nutt. Sylva*, i. 10.—Am. bor.
tirolensis, Nym. Consp. 671 = incana.
tomentosa, Hartig, ex Regel, in DC. Prod. xvi. II. 185 = orientalis.
tristis, Wormsk. ex Regel, l. c. 182 = viridis.
undulata, *Willd. Sp. Pl.* iv. 336.—Canada.
Vilmoriana, *Hort. ex Lebas, in Rev. Hortic.* (1866) 446.—Hab.?
viridis, *DC. Fl. Fr.* iii. 304; *Cham. in Linnaea*, vi. (1831) 538.—Europ.; As. bor.; Am. bor.
viridis, A. Gray, Man. Bot. N. Un. St. (1848) 424 = crispa.
vulgaris, Pers. Syn. ii. 550 = glutinosa.

ALOCASIA, Neck. Elem. iii. 289 (1790); Schott, in Schott & Endl. Melet. 18 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 975.
ENSOLENANTHE, Schott, in Bonplandia, ix. (1861) 368.
acuminata, *Schott, in Bonplandia*, vii. (1859) 28.—Ind. or.
alba, *Schott, in Oestr. Bot. Wochenbl.* ii. (1852) 59; (1854) 410.—Zeylan.
argyroneura, C. Koch, Berl. Wochenschr. (1861) 8 = Caladium Schomburgkii.
Beccarii, *Engl. in Bull. Soc. Tosc. Ortic.* iv. (1879) 300.—Malaya.
commutata, Schott, in Oestr. Bot. Wochenbl. (1854) 409 = Alocasia odora.
cucullata, *Schott, in Schott & Endl. Melet.* 18.—Ind. or.
cuprea, *C. Koch, in Berl. Wochenschr.* iv. (1861) 141; vii. (1864) 90.—Borneo.
decipiens, *Schott, in Bonplandia*, vii. (1859) 28.—Ind. or.
denudata, *Engl. in DC. Monog. Phan.* ii. 507.—Singapor.
erythraea, C. Koch, in Berl. Wochenschr. (1861) 8 = Colocasia Schomburgkii.
fallax, Schott, in Bonplandia, vii. (1859) 28 = navicularis.
fornicata, *Schott, in Oestr. Bot. Wochenbl.* (1854) 410.—Ind. or.
grandis, *Clémenc. in Rev. Hortic.* (1868) 380.—Hab.?
hetcroneura, C. Koch, in Berl. Wochenschr. (1861) 283 = Caladium Schomburgkii.
indica, *Schott. in Oestr. Bot. Wochenbl.* (1854) 410, pro parte.—Ind. or.; Malaya.
Jenningsii, Hort. Veitch. Cf. Gard. Chron. (1868) 26; in Illustr. Hortic. (1869) t. 585 = Colocasia affinis.
Johnstonii, Hort. in Gard. Chron. xiii. (1880) 759; in Illustr. Hortic. xxvii. (1880) t. 395 = Cyrtosperma Johnstonii.
Korthalsii, *Schott, in Ann. Mus. Lugd. Bat.* i. 124.—Borneo.
Liervalii, *Hort. ex Hérincq, L'Hortic. Franç.* (1869) 243.—Ins. Philipp.
lobata, *Rafin. Fl. Tellur.* iii. 64.—Hab.?
longiloba, *Miq. Fl. Ind. Bat.* iii. 207.—Malaya.
Lowii, *Hook. Bot. Mag.* t. 5376.—Borneo.
macrorhiza, *Schott, in Schott & Endl. Melet.* 18.—As. trop.; Austral.; Ins. Pacific.
Marchalli, Engl. in DC. Monog. Phan. ii. 494 = Colocasia Marchalli.
montana, *Schott, in Oestr. Bot. Wochenbl.* (1854) 410.—Ind. or.; Malaya.

ALOCASIA:—
metallica, Hook. Bot. Mag. t. 5190 = cuprea.
metallica, Schott, in Oestr. Bot. Wochenbl. (1854) 410 = indica.
navicularis, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1855) App. 2.—Reg. Himal.
odora, *C. Koch, l. c.* (1854), *App.* 5.—Ind. or.; China.
pallida, C. Koch & Bouché, l. c. = alba.
pedata, Rafin. Fl. Tellur. iii. 64 = Sauromatum pedatum.
plumbea, Van Houtte, in Fl. des Serres, Sér. II. xi. (1875) 93. t. 2206 = indica.
porphyroneura, Lem. in Illustr. Hortic. (1861) t. 297 = Caladium picturatum.
Portei, *Becc. & Engl. in DC. Monog. Phan.* ii. 645.—N. Guin.
pubera, *Schott, Syn. Aroid.* 47.—Java.
Putzeysi, *N. E. Br. in Illustr. Hortic.* xxix. (1882) t. 439, & p. 22.—Sumatra.
rapiformis, *Schott, Prod. Aroid.* 157.—Burma.
Regina, *N. E. Br.—Cf. Gard. Chron.* (1884) I. 711.—Borneo.
Roezlii, Hort. ex Gard. Chron. iii. (1875) 589 = Caladium marmoratum.
rugosa, Schott, in Oestr. Bot. Wochenbl. (1854) 410 = cucullata.
scabriuscula, *N. E. Br. in Gard. Chron.* (1879) II. 296.—Borneo.
Scdeni ×, Hort. Veitch.—Cf. Gard. Chron. (1869) 587, 615 = cuprea × Lowii.
sinuata, *N. E. Br. in Gard. Chron.* (1885) II. 678.—Malaya.
Thibautiana, *Mast. l. c.* (1878) I. 527.—Borneo.
undipes, C. Koch & Bouché, in Ind. Sem. Hort. Berol. (1854), App. 5 = Xanthosoma Jacquini.
varicgata, C. Koch & Bouché, l. c. = indica.
Veitchii, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 125 = Lowii.
virginica, Rafin. Fl. Tellur. iii. 64 = Peltandra virginica.
zebrina, *C. Koch & Veitch, in Berl. Wochenschr.* VI. (1863) 331.—Ins. Philipp.

ALOË, Tourn. ex Linn. Syst. ed. I. (1735). *LILIACEAE*, Benth. & Hook. f. iii. 776.
AGRIODENDRON, Endl. Gen. 144 (1837).
BOWIEA, Haw. in Phil. Mag. (1824) 299.
BUSIPHO, Salisb. Gen. Pl. Fragm. 76 (1866).
CATEVALA, Medic. Theodora, 67 (1786).
KUMARA, Medic. l. c. 69. t. 4 (1786).
PACHIDENDRON, Haw. Rev. Pl. Succ. 35 (1821).
PTYAS, Salisb. Gen. Pl. Fragm. 76 (1866).
RHIPIDODENDRON, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 164.
abyssinica, *Lam. Encyc.* i. 86.—Abyss.
acinacifolia, Jacq. Ecl. 49. t. 31 = Gasteria acinacifolia.
acuminata, Haw. Syn. Pl. Succ. 84 = A. humilis.
acuminata, Lam. Encyc. i. 90 = Gasteria verrucosa.
africana, *Mill. Gard. Dict.* ed. VIII. n. 4.—Afr. austr.
africana, Salm-Dyck, Cat. 31 = Salm-Dyckiana.
agavefolia, *Tod. Hort. Bot. Panorm.* i. 85. t. 23.—Afr. trop.
albicans, Haw. in Trans. Linn. Soc. vii. (1804) 18 = Haworthia albicans.
albispina, *Haw. l. c.* 22.—Afr. austr.
albocincta, Haw. Suppl. Pl. Succ. 43 = striata.
altilinea, Schult. f. Syst. vii. 638 = Haworthia altilinea.
americana, Crantz, Inst. i. 466 = Agave americana.
andongensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 263.—Afr. trop.
angolensis, *Baker, l. c.*—Afr. trop.
angulata, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 276 = Gasteria disticha.
angusta, *Schult. f. Syst.* vii. 710.—Afr. austr.
angustifolia, Salm-Dyck, Obs. 57 = Gasteria disticha.
anomala, Haw. in Trans. Linn. Soc. vii. (1804) 25 = Haworthia recurva.
arabica, *Lam. Encyc.* i. 91.—Arab.
arabica, Salm-Dyck, Cat. 27 = microstigma.
arachnoidea, Mill. Gard. Dict. ed. VIII. n. 17 = Haworthia arachnoides.
arachnoides, Thunb. Diss. Aloe, 7 = Haworthia arachnoides.
arborea, Medic. Beobacht. (1783) 305 = arborescens.
arborescens, *Mill. Gard. Dict.* ed. VIII. n. 3.—Afr. austr.

ALOË :—
aristata, *Haw. in Phil. Mag.* (Oct. 1825) 280.—Afr. austr.
aristata, *Schult. f. Syst.* vii. 638.—Afr. austr.
aspera, Haw. in Trans. Linn. Soc. vii. (1804) 6 = Apicra aspera.
asperiuscula, Schult. f. Syst. vii. 653 = Haworthia asperiuscula.
asperula, Schult. f. l. c. 635 = Haworthia asperula.
Atherstonei, *Baker, in Journ. Linn. Soc.* xviii. (1881) 170.—Afr. austr.
atrovirens, DC. Pl. Grass. t. 51 = Haworthia atrovirens.
attenuata, Haw. in Trans. Linn. Soc. vii. (1804) 11 = Haworthia attenuata.
Bainesii, *Dyer, in Gard. Chron.* (1874) i. 567.—Afr. austr.
barbadensis, Mill. Gard. Dict. ed. VIII. n. 2 = vera.
Barberae, Dyer, in Gard. Chron. (1874) i. 566 = Bainesii.
Barteri, *Baker, in Journ. Linn. Soc.* xviii. (1881) 168.—Afr. trop.
Bayfieldii, Salm-Dyck, Monog. ii. § 29. f. 14 = Gasteria Bayfieldii.
bicarinata, Spreng. Syst. ii. 70; Schult. f. Syst. vii. 652 = Apicra bicarinata.
bicolor, Schult. f. l. c. 682 = Gasteria bicolor.
Bolusii, *Baker, in Journ. Linn. Soc.* xviii. (1881) 179.—Afr. austr.
Bourea, *Schult. f. Syst.* vii. 704.—Afr. austr.
Boureana, Schult. f. l. c. 662 = Gasteria picta.
brachyphylla, Salm-Dyck, Hort. 332 = Gasteria brevifolia.
Bradlyana, Jacq. Hort. Schoenbr. iv. 11, in nota ad t. 421 = Haworthia atrovirens.
brevifolia, *Mill. Gard. Dict.* ed. VIII. in errat.—Afr. austr.
brevifolia, Haw. in Trans. Linn. Soc. vii. (1804) 23 = distans.
brevioribus, Mill. Gard. Dict. ed. VIII. n. 8, sphalm. = brevifolia.
brevis, *Schult. f. Syst.* vii. 649.—Afr. austr.
bullulata, Jacq. Fragm. 72. t. 109 = Apicra pentagona.
caesia, *Salm-Dyck, Cat.* 29.—Afr. austr.
Candelabrum, *Tod. Hort. Panorm.* (1876) 46.—Hab.?
candicans, Schult. f. Syst. vii. 681 = Gasteria candicans.
capitata, *Baker, in Journ. Linn. Soc.* xx. (1883) 272.—Madag.
carinata, Ker-Gawl. in Bot. Mag. t. 1331 = Gasteria glabra.
carinata, DC. Pl. Grass. t. 63 = Gasteria verrucosa.
carinata, Mill. Gard. Dict. ed. VIII. n. 21 = Gasteria carinata.
caroliniana, Hill, Veg. Syst. xiii. 64 = Agave virginica.
chinensis, *Steud. Nom.* ed. II. i. 56; *Baker, in Bot. Mag.* t. 6301.—China.
chloracantha, Schult. f. Syst. vii. 641 = Haworthia chloracantha.
chloroleuca, *Baker, in Gard. Chron.* (1877) ii. 38.—Afr. austr.
ciliaris, *Haw. in Phil. Mag.* (Oct. 1825) 281.—Afr. austr.
clariperla, Schult. f. Syst. vii. 646, 1712 = Haworthia attenuata.
claviflora, *Burch. Trav.* i. 272.—Afr. austr.
coarctata, Schult. f. Syst. vii. 647 = Haworthia coarctata.
Commelyni, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 282 = mitriformis.
commutata, *Tod. Hort. Bot. Panorm.* i. 75. t. 18.—Hab.?
concinna, Schult. f. Syst. vii. 653 = Haworthia viscosa.
congesta, Salm-Dyck, Monog. i. § 2. f. 1 = Apicra congesta.
consobrina, *Salm-Dyck, l. c.* § 18. f. 2.—Afr. austr.
conspurcata, Salm-Dyck, Obs. 59 = Gasteria disticha.
constricta, *Baker, in Journ. Linn. Soc.* xviii. (1881) 168.—Afr. trop.
Cooperi, *Baker, in Gard. Chron.* (1874) i. 628.—Afr. austr.
cordifolia, Schult. f. Syst. vii. 653 = Haworthia cordifolia.
crassifolia, Schult. f. l. c. 664 = Gasteria nigricans.
crassipes, *Baker, in Journ. Linn. Soc.* xviii. (1880) 162.—Afr. trop.
Croucheri, Hook. f. Bot. Mag. t. 5812 = Gasteria Croucheri.

ALOË :—
cryptopoda, *Baker, in Journ. Bot.* xxii. (1884) 52.—Afr. trop.
curta, *Spreng. Syst.* ii. 69.—Afr. austr.
cuspidata, Schult. f. Syst. vii. 639 = Haworthia cuspidata.
cylindracea, Lam. Encyc. i. 89 = Apicra spiralis, et Haworthia rigida.
cymbaefolia, Schrad. Neues Journ. ii. (1807) 17 = Haworthia cymbiformis.
cymbiformis, Haw. in Trans. Linn. Soc. vii. (1804) 8 = Haworthia cymbiformis.
decipiens, Schult. f. Syst. vii. 674 = Gasteria decipiens.
deltoidea, Hook. f. Bot. Mag. t. 6071 = Apicra deltoidea.
deltoideodonta, *Baker, in Journ. Linn. Soc.* xx. (1883) 271.—Madag.
denticulata, Schult. f. Syst. vii. 639 = Haworthia denticulata.
dependens, Steud. Nom. ed. I. 30 = pendens.
depressa, *Haw. in Trans. Linn. Soc.* vii. (1804) 16.—Afr. austr.
depressa, Salm-Dyck, ex Steud. Nom. ed. II. i. 57 = distans.
dichotoma, *Linn. f. Suppl.* 206.—Afr. austr.
dictyodes, Schult. f. Syst. vii. 663 = Gasteria retata.
distans, *Haw. Syn. Pl. Succ.* 78.—Afr. austr.
disticha, *Linn. Sp. Pl.* 321.—Afr. austr.
disticha, Mill. Gard. Dict. ed. VIII. n. 5 = saponaria.
disticha, Thunb. Diss. Aloë, n. 9 = Gasteria verrucosa.
dorsalis, Haw. in Trans. Linn. Soc. vii. (1804) 28 = lineata.
drepanophylla, *Baker, in Gard. Chron.* iii. (1875) 814.—Afr. austr.
echinata, Willd. Enum. Hort. Berol. 385 = humilis.
Ecklonis, *Salm-Dyck, Monog.* ii. § 21. f. 2.—Afr. austr.
elegans, *Tod. Ind. Sem. Panorm.* (1880) 37.—Hab.?
elongata, Murr. in Comm. Gott. ix. (1789) 191 = vera.
elongata, Salm-Dyck, Monog. § 29. f. 15 = Gasteria trigona.
ensifolia, Schult. f. Syst. vii. 681 = Gasteria acinacifolia.
erecta, *Schult. f. l. c.* 649.—Afr. austr.
erecta, Salm-Dyck, Aloë, § 6. f. 7 = Haworthia margaritifera.
excavata, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 276 = Gasteria excavata.
expansa, Haw. in Trans. Linn. Soc. vii. (1804) 8 = Haworthia rigida.
falcata, *Baker, in Journ. Linn. Soc.* xviii. (1880) 181.—Afr. austr.
fasciata, Salm-Dyck, in Schult. f. Syst. vii. 1713 = Haworthia fasciata.
ferox, *Mill. Gard. Dict.* ed. VIII. n. 22.—Afr. austr.
flabelliformis, Salisb. Prod. 246 = plicatilis.
flava, Pers. Syn. i. 378 = vera.
flavispina, Haw. in Trans. Linn. Soc. vii. (1804) 22 = mitriformis.
foetida, Crantz, Inst. i. 466 = Agave americana.
foliolosa, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Apicra foliolosa.
formosa, *Schult. f. Syst.* vii. 662.—Afr. austr.
fragrantissima, Jacq. Enum. Stirp. Vind. App. 399 = Dracaena fragrans.
frutescens, Salm-Dyck, Cat. 30 = arborescens.
fruticosa, Lam. Encyc. i. 87 = arborescens.
gasterioides, *Baker, in Journ. Linn. Soc.* xviii. (1880) 166.—Afr. austr.
glabra, Salm-Dyck, Obs. 48 = Gasteria glabra.
glabrata, Salm-Dyck, Hort. 325 = Haworthia glabrata.
glauca, *Mill. Gard. Dict.* ed. VIII. n. 16.—Afr. austr.
gracilis, *Haw. in Phil. Mag.* (Oct. 1825) 280.—Afr. austr.
granata, *Schult. f. Syst.* vii. 650.—Afr. austr.
granata, Salm-Dyck, Aloë, § 6. f. 6 = Haworthia margaritifera.
grandidentata, *Salm-Dyck, Obs.* (1822) 329.—Afr. austr.
Greenii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 165.—Afr. austr.
guineensis, Jacq. Enum. Stirp. Vind. App. 308 = Sanseviera guineensis.
guttata, Salm-Dyck, Hort. 332 = Gasteria nigricans.
Hanburiana, Naud. in Rev. Hortic. (1875) 165 = striata.
Haynaldi, *Tod. Hort. Panorm.* i. 75. t. 24.—Hab.?
hebes, Schult. f. Syst. vii. 637 = Haworthia cymbiformis.

ALOË :—
herbacea, DC. Hort. Monsp. 76 = Haworthia reticulata.
herbacea, Mill. Gard. Dict. ed. VIII. n. 18 = Haworthia atrovirens.
heteracantha, *Baker, in Journ. Linn. Soc.* xviii. (1880) 161.—Hab.?
hexapetala, *Salm-Dyck, Cat.* 28.—Hab.?
horrida, Haw. in Trans. Linn. Soc. vii. (1804) 27 = ferox.
humilis, *Mill. Abr. Gard. Dict.* ed. VI. n. 10 (1771).—Afr. austr.
hyacinthoides, Linn. Sp. Pl. 321 = Sanseviera zeylanica.
hybrida, Salm-Dyck, Cat. 7 = Haworthia hybrida.
imbricata, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Apicra spiralis.
incurva, Haw. Syn. Pl. Succ. 85 = humilis.
indica, Royle, Illust. Pl. Himal. 390 = vera.
indurata, Schult. f. Syst. vii. 654.—Afr. austr.
inermis, *Forsk. Fl. Aegypt. Arab.* 74.—Arabia.
inermis, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 161 = heteracantha.
insignis ×, *N. E. Br. in Gard. Chron.* xxiv. (1885) 40. f. 8.—Hab.?
intermedia, Haw. in Trans. Linn. Soc. vii. (1804) 12 = Gasteria verrucosa.
Kraussii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 159.—Afr. austr.
laetepunctata, Schult. f. Syst. vii. 676 = Gasteria laetepunctata.
laetevirens, Link, Enum. Hort. Berol. i. 335 = Haworthia laetevirens.
laevigata, Schult. Syst. vii. 636 = Haworthia albicans.
laevis, *Salm-Dyck, Cat.* 16.—Afr. austr.
latifolia, *Haw. Syn. Pl. Succ.* 82.—Afr. austr.
leptocaulon, Boj. Hort. Maurit. 345 = oligophylla?
leptophylla, N. E. Br. ex Baker, in Journ. Linn. Soc. xviii. (1881) 165 = latifolia.
limpida, *Schult. f. Syst.* vii. 638.—Afr. austr.
lineata, *Haw. in Trans. Linn. Soc.* vii. (1804) 18.—Afr. austr.
Lingua, Ker-Gawl. in Bot. Mag. t. 1322 = Gasteria verrucosa.
Lingua, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 277 = sp. plurim.
Lingua, Thunb. Diss. Aloë, 8 = Gasteria disticha.
linguiformis, DC. Pl. Grass. t. 68 = Gasteria sulcata.
linguiformis, Linn. f. Suppl. 206 = plicatilis.
linguiformis, Mill. Gard. Dict. ed. VIII. n. 13 = Gasteria disticha.
linguiformis, *Medic. Beobacht.* (1783) 64.—Hab.?
linita, *Schult. f. Syst.* vii. 682.—Afr. austr.
littoralis, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i (1878) 293.—Afr. trop.
littoralis, Koen. ex Baker, in Journ. Linn. Soc. xviii. (1880) 176 = vera.
lomatophylloides, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 22.—Ins. Mascar.
longiaristata, Schult. f. Syst. vi. 684 = aristata.
longifolia, Haw. Syn. Pl. Succ. 88 = Gasteria disticha.
longifolia, Lam. Encyc. i. 90 = Tritomanthe Uvaria.
longistyla, *Baker, in Journ. Linn. Soc.* xviii. (1880) 158.—Afr. austr.
Lynchii ×, *Baker, in Gard. Chron.* (1881) i. 266.
macilenta, Nichols. Dict. Gard. i. 52 = humilis.
Macowani, *Baker, in Journ. Linn. Soc.* xviii. (1881) 170.—Afr. austr.
macra, Haw. Suppl. Pl. Succ. 45, 102 = Lomatophyllum macrum.
macracantha, *Baker, in Journ. Linn. Soc.* xviii. (1880) 167.—Afr. austr.
macrocarpa, *Tod. Hort. Bot. Panorm.* 36. t. 9.—Abyss.
macroclada, *Baker, l. c.* xx. (1883) 273.—Madag.
maculata, Ker-Gawl. in Bot. Mag. t. 765 = Gasteria pulchra.
maculata, Desf. Tabl. Hort. Par. ed. I. 25 = obscura.
maculata, Forsk. Fl. Aegypt. Arab. 73 = abyssinica.
maculata, Thunb. Diss. Aloë, 8 = Gasteria maculata.
maculosa, Lam. Encyc. i. 87 = obscura, saponaria.
margaritifera, Burm. f. Fl. Cap. Prod. 10 = Haworthia margaritifera.
marginalis, DC. Pl. Grass. t. 31 = Lomatophyllum borbonicum.
marginata, Lam. Encyc. i. 89 = Haworthia albicans.
marginata, Willd. Enum. Hort. Berol. 382 = Lomatophyllum borbonicum.
marmorata, Hitchen, ex Steud. Nom. ed. II. i. 58 = formosa.

ALOË :—

micracantha, *Haw. Suppl. Pl. Succ.* 105.—Afr. austr.
microstigma, *Salm-Dyck, Monog.* ii. § 26. f. 4.—Afr. austr.
minor, Schult. f. Syst. vii. 648 = Haworthia margaritifera.
mirabilis, Haw. in Trans. Linn. Soc. vii. (1804) 9 = Haworthia mirabilis.
mitraeformis, DC. Pl. Grass. 99; Haw. Syn. Succ. 77 = distans, nobilis.
mitraeformis, Willd. Enum. Hort. Berol. 385 = mitriformis.
mitriformis, *Mill. Gard. Dict.* ed. VIII. n. 1.—Afr. austr.
mollis, Schult. Syst. vii. 665 = Gasteria mollis.
mucronata, *Schult. f. l. c.* 638.—Afr. austr.
multifaria, *Schult. f. l. c.* 635.—Afr. austr.
muricata, Haw. in Trans. Linn. Soc. vii. (1804) 25 = ferox.
muricata, Schult. Obs. 70 = glauca.
mutica, *Schult. f. Syst.* vii. 635.—Afr. austr.
myriacantha, *Schult. f. l. c.* 704.—Afr. austr.
neglecta, *Tenore, in Atti Accad. Sc. Nap.* v. 1. (1837) p. 257.—Hab.?
nigra, Schult. f. Syst. vii. 657 = Haworthia nigra.
nigricans, Haw. in Trans. Linn. Soc. vii. (1804) 13 = Gasteria nigricans.
nitens, *Baker, in Journ. Linn. Soc.* xviii. (1880) 170. —Afr. austr.
nitens, Schult. f. Syst. vii. 680 = Gasteria acinacifolia.
nitida, Salm-Dyck, Cat. 13 = Gasteria nitida.
nobilis, *Haw. Syn. Pl. Succ.* 78.—Afr. austr.
obliqua, DC. Pl. Grass. t. 91 = Gasteria pulchra.
obliqua, Haw. in Trans. Linn. Soc. vii. (1804) 14 = Gasteria maculata.
obliqua, Jacq. Hort. Schoenb. iv. 9 = Gasteria nigricans.
obscura, *Mill. Gard. Dict.* ed. VIII. n. 6.—Afr. austr.
obscura, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 275 = Gasteria excavata.
obscurevirens, *Martinati, ex Vis. Orto Bot. Padov.* 132.—Hab.?
obtusa, Schult. f. Syst. vii. 679 = Gasteria obtusa.
obtusifolia, Salm-Dyck, Obs. 62 = Gasteria obtusifolia.
officinalis, Forsk. Fl. Aegypt. Arab. 73 = vera.
oligophylla, *Baker, in Journ. Linn. Soc.* xx. (1883) 272.—Madag.
pallescens, Haw. Rev. Pl. Succ. 41 = serrulata.
pallida, Schult. f. Syst. vii. 641 = Haworthia pallida.
palmiformis, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 263.—Angola.
paniculata, Jacq. Fragm. 48. t. 68 = striata.
papillosa, Salm-Dyck, Cat. 7 = Haworthia papillosa.
parva, Schult. f. Syst. vii. 653 = Haworthia tessellata.
Peacockii, Baker, in Journ. Linn. Soc. xviii. (1880) 175 = abyssinica.
pellucens, Haw. Syn. Pl. Succ. 96 = Haworthia translucens.
pendens, *Forsk. Fl. Aegypt. Arab.* 74.—Arabia.
pentagona, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Apicra pentagona.
pentagona, *Jacq. Fragm.* 72. t. 111.—Afr. austr.
pentagona, Salm-Dyck, Monog. i. § 1. f. 4 = quinquangularis.
percrassa, Tod. Hort. Bot. Panorm. i. 81. t. 21 = abyssinica.
perfoliata, Linn. Sp. Pl. 319, 320 = sp. plurim.
perfoliata, Meyen, Reise, ii. 45 = arborescens.
Perryi, *Baker, in Journ. Linn. Soc.* xviii. (1880) 161. —Ins. Socotra.
pertusa, Haw. in Trans. Linn. Soc. vii. (1804) 25 = Crassula corymbulosa.
picta, Thunb. Diss. Aloë, 6 = obscura.
planifolia, Baker, in Saund. Refug. Bot. t. 162 = Gasteria planifolia.
planifolia, Schult. f. Syst. vii. 638 = Haworthia cymbiformis.
platylepis, *Baker, in Gard. Chron.* (1877) II. 38.—Afr. austr.
platyphylla, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 264.—Angola.
plicatilis, *Mill. Gard. Dict.* ed. VIII. n. 7.—Afr. austr.
pluridens, *Haw. in Phil. Mag.* (Oct. 1824) 299.—Afr. austr.
pluripunctata, *Schult. f. Syst.* vii. 681.—Afr. austr.
postgenita, *Schult. f. l. c.* 1714.—Afr. austr.

ALOË :—

pratensis, *Baker, in Journ. Linn. Soc.* xviii. (1880) 156. —Afr. austr.
prolifera, Haw. in Trans. Linn. Soc. vii. (1804) 16 = brevifolia.
pseudoafricana, Salm-Dyck, Cat. 31 = africana.
pseudoangulata, Salm-Dyck, l. c. 16 = Gasteria subcarinata.
pseudoferox, Salm-Dyck, l. c. 31 = ferox.
pseudonigricans, Salm-Dyck, l. c. 17 = Gasteria nigricans.
pseudorigida, Salm-Dyck, l. c. 9 & 41 = Haworthia subrigida.
pseudotortuosa, Salm-Dyck, l. c. 8 & 41 = Haworthia viscosa.
pulchra, Jacq. Hort. Schoenbr. iv. 10. t. 419 = Gasteria pulchra.
pumila, Linn. Sp. Pl. 322 = sp. plurim.
Pumilio, Jacq. Hort. Schoenb. iv. 11 = Haworthia reticulata.
punctata, Haw. in Trans. Linn. Soc. vii. (1804) 26 = variegata.
punctata, *Martinati, in Vis. Orto Bot. Padov.* 132.—Hab.?
purpurascens, *Haw. in Trans. Linn. Soc.* vii. (1804) 20.—Afr. austr.
purpurea, Lam. Encyc. i. 85 = Lomatophyllum borbonicum.
pusilla, *Schult. f. Syst.* vii. 674.—Hab.?
quinquangularis, *Schult. f. l. c.* 658.—Afr. austr.
racemosa, Lam. Encyc. i. 91 = Gasteria verrucosa.
Radula, Jacq. Hort. Schoenb. iv. 11. t. 422 = Haworthia Radula.
Radula, Ker-Gawl. in Bot. Mag. t. 1345 = Haworthia attenuata.
ramifera, *Schult. f. Syst.* vii. 636.—Afr. austr.
ramosa, Haw. in Trans. Linn. Soc. vii. (1804) 26 = purpurascens.
recurva, Haw. l. c. 10 = Haworthia recurva.
reflexa, Van Marum, ex Steud. Nom. ed. II. i. 59 = distans.
Reinwardtii, Salm-Dyck, Obs. (1821) 37 = Haworthia Reinwardtii.
repens, Schult. f. Syst. vii. 674 = Gasteria repens.
reticulata, Haw. in Trans. Linn. Soc. vii. (1804) 9 = Haworthia reticulata.
retusa, Linn. Sp. Pl. 322 = Haworthia retusa.
retusifolia, Haw. ex Steud. Nom. ed. II. i. 59 = Gasteria obtusifolia.
rhodacantha, DC. Pl. Grass. 44 = glauca.
rhodocincta, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 163 = striata.
rigida, Ker-Gawl. in Bot. Mag. t. 1337 = Haworthia tortuosa.
rigida, DC. Pl. Grass. t. 62 = Haworthia rigida.
rigida, Salisb. Prod. 245 = Kniphofia alooides.
rigida, Jacq. Fragm. t. 108 = Haworthia subrigida.
rubescens, DC. Pl. Grass. t. 15 = vera.
rufocincta, Haw. Suppl. Pl. Succ. 45 = Lomatophyllum rufocinctum.
rugosa, Salm-Dyck, in Schult. f. Syst. vii. 1712 = Haworthia rugosa.
Sabila, *Karw. ex Steud. Nom.* ed. II. i. 59.—Mexic.
Sahundra, *Boj. Hort. Maurit.* 345.—Madag.
Salm-Dyckiana, *Schult. f. in Syst.* vii. 710.—Afr. austr.
saponaria, *Haw. in Trans. Linn. Soc.* vii. (1804) 17.—Afr. austr.
scaberrima, Salm-Dyck, Hort. 332; Schult. f. Syst. vii. 1713 = Gasteria verrucosa.
scabra, Schult. f. Syst. vii. 644 = Haworthia scabra.
Schimperi, *Tod. Hort. Panorm.* (1876) 46.—Abyss.
Schmidtiana, Regel, Gartenfl. (1879) 97 = Cooperi.
Schweinfurthii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 175.—Afr. trop.
semiglabrata, Schult. f. Syst. vii. 650 = Haworthia semiglabrata.
semimargaritifera, Salm-Dyck, Cat. 6 = Haworthia margaritifera.
Serra, *DC. Pl. Grass.* t. 80.—Afr. austr.
Serra, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 282 = depressa.
serrulata, *Haw. in Trans. Linn. Soc.* vii. (1804) 18.—Afr. austr.
setosa, Schult. f. Syst. vii. 641 = Haworthia setata.
sigmoidea, *Baker, in Journ. Linn. Soc.* xviii. (1880) 177.—Afr. austr.
sinuata, Thunb. Diss. Aloë, 6 = succotrina.
sinuata, Willd. Sp. Pl. ii. 187 = purpurascens.

ALOË :—

soccotorina, Schult. f. Syst. vii. 701 = seq.
soccotrina, DC. Pl. Grass. 61. t. 85 = succotrina.
sordida, Schult. f. Syst. vii. 644 = Haworthia sordida.
specciosa, *Baker, in Journ. Linn. Soc.* xviii. (1880) 178. —Afr. austr.
spicata, *Linn. f. Suppl.* 205; *Haw. Syn. Succ.* 76.—Afr. austr.
spinulosa, Salm-Dyck, Obs. (1822) 4 = mitriformis.
spiralis, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Apicra pentagona.
spiralis, Linn. Sp. Pl. 322 = Apicra spiralis.
spirella, Salm-Dyck, Cat. 10 = Apicra pentagona.
squarrosa, *Baker, ex Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 97.—Ins. Socotr.
stenophylla, Schult. f. Syst. vii. 641 = Haworthia angustifolia.
striata, *Haw. in Trans. Linn. Soc.* vii. (1804) 18.—Afr. austr.
striatula, *Haw. in Phil. Mag.* (Oct. 1825) 281.—Afr. austr.
strigata, *Schult. f. Syst.* vii. 676.—Afr. austr.
subalbicans, Salm-Dyck, Monog. i. § 6. f. 1 = Haworthia margaritifera.
subattenuata, Salm-Dyck, in Schult. f. Syst. vii. 1712 = Haworthia subattenuata.
subcarinata, Salm-Dyck, Obs. 51 = Gasteria subcarinata.
suberecta, Haw. in Trans. Linn. Soc. vii. (1804) 16 = humilis.
subfasciata, Salm-Dyck, in Schult. f. Syst. vii. 1712 = Haworthia subfasciata.
subferox, *Spreng. Syst.* ii. 73.—Afr. austr.
subnigricans, Spreng. l. c. 71 = Gasteria nigricans.
subrigida, Schult. f. Syst. vii. 654 = Haworthia subrigida.
subtortuosa, Schult. f. l. c. = Haworthia viscosa.
subtuberculata, Haw. in Phil. Mag. (Oct. 1825) 280 = humilis.
subulata, Salm-Dyck, in Schult. f. Syst. vii. 1712 = Haworthia subulata.
subverrucosa, Salm-Dyck, Obs. 67 = Gasteria subverrucosa.
succotrina, *Lam. Encyc.* i. 85.—Afr. austr.
sulcata, Salm-Dyck, Obs. 54 = Gasteria sulcata.
supralaevis, *Haw. in Trans. Linn. Soc.* vii. (1804) 22. —Afr. austr.
tenuifolia, *Lam. Encyc.* i. 87.—Afr. austr.
tenuior, *Haw. in Phil. Mag.* (Oct. 1825) 281.—Afr. austr.
tessellata, Schult. f. Syst. vii. 635 = Haworthia tessellata.
Thraskii, *Baker, in Journ. Linn. Soc.* xviii. (1880) 180.—Afr. austr.
torquata, Schult. f. Syst. vii. 654 = Haworthia viscosa.
tortuosa, Haw. in Trans. Linn. Soc. vii. (1804) 7 = Haworthia tortuosa.
translucens, [Soland. in] Ait. Hort. Kew. ed. II. ii. 300 = Haworthia translucens.
triangularis, Lam. Encyc. i. 89 = Haworthia viscosa.
trichotoma, Colla, Hort. Ripul. 13 = saponaria.
tricolor, *Baker, in Bot. Mag.* t. 6324.—Afr. austr.
tricolor, Haw. in Trans. Linn. Soc. vii. (1804) 25 = Haworthia venosa.
trigona, Salm-Dyck, Monog. ii. § 20. f. 18 = Gasteria obtusa.
tripetala, Medic. Beobacht. (1783) 55 = plicatilis.
tristicha, Medic. Theod. 74 = Gasteria carinata.
tuberculata, Haw. in Trans. Linn. Soc. vii. (1804) 16 = humilis.
tuberculata, *Lag. Gen. et Sp. Nov.* 13.—Hab.?
turgida, Schult. f. Syst. vii. 635 = Haworthia turgida.
umbellata, DC. Pl. Grass. t. 98 = saponaria.
undata, Schult. f. Syst. vii. 677 = Gasteria subcarinata.
Uvaria, Linn. Sp. Pl. 323 = Kniphofia alooides.
vacillans, Forsk. Fl. Aegypt. Arab. 74 = vera.
variegata, Forsk. l. c. 74 = arabica.
variegata, *Linn. Sp. Pl.* 321.—Afr. austr.
venosa, Lam. Encyc. i. 89 = Haworthia venosa.
venusta, *Schult. f. Syst.* vii. 681.—Hab.?
vera, *Linn. Sp. Pl.* 320.—Reg. medit.
vera, Mill. Gard. Dict. ed. VIII. n. 15 = succotrina.
verrucosa, Mill. l. c. n. 20 = Gasteria verrucosa.
verrucula, Medic. Beobacht. (1783) 63 = prace.
virens, *Haw. in Trans. Linn. Soc.* vii. (1804) 17.—Afr. austr.
virescens, *Schult. f. Syst.* vii. 637.—Hab.?

ALOË :—
virginica, Crantz, Inst. i. 466 = Agave virginica.
viscosa, Linn. Sp. Pl. 322 = Haworthia viscosa.
vittata, *Schult. f. Syst.* vii. 662.—Afr. austr.
vivipara, Crantz, Inst. i. 466 = Agave vivipara.
vulgaris, Lam. Encyc. i. 86 = vera.
xanthacantha, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 282 = mitriformis.
yuccaefolia, A. Gray, in Proc. Am. Acad. vii. (1868) 390 = Hesperaloë yuccaefolia.
zebrina, *Baker, in Trans. Linn. Soc.* Ser. II. *Bot.* i. (1878) 264.—Afr. trop.
Zeyheri, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1880) 178 = Bainesii.
Zeyheri, Salm-Dyck, Aloë, § 29. f. 3bis = Gasteria Zeyheri.
zeylanica, Jacq. Hort. Vind. 310 = Sanseviera zeylanica.

ALOËATHEROS, Fndl. Gen. 95 (1836) = ALLOIATHEROS, Ell. = **Gymnopogon**, Beauv. ? (Gramin.).

ALOËXYLUM, Lour. Fl. Cochinch. 267 (1790) = **Aquilaria**, Lam. (Thymel.).
Agallochum, Lour. l. c. = Aq. Agallochum.
verum, Lour. l. c. = Aq. Agallochum.

ALOIDES, Boerh. ex Linn. Gen. ed. II. 253 (1742) = **Stratiotes**, Linn. (Hydrochar.).

ALOIOZONIUM, Lindl. Veg. Kingd. 713 (1847) = ALLOIOZONIUM, Kunze = **Cryptostemma**, R. Br. (Compos.).

ALOITIS, Rafin. Fl. Tellur. iii. 21 (1836) = **Gentiana** Linn.
anceps, Rafin. l. c. 22.—Hab. ?
parviflora, Rafin. l. c. 21 = G. quinqueflora.
quinqueflora, Rafin. l. c. 22 = G. quinqueflora.

ALOMIA, H. B. & K. Nov. Gen. et Sp. iv. 151, 312. t. 354 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 240.
Lycapsus, Phil. in Bot. Zeit. (1870) 499. t. 8.
Orsinia, Bertol. ex DC. Prod. v. 104 (1836).
ageratoides, *H. B. & K. Nov. Gen. et Sp.* iv. 151. t. 354.—Mexic.
alata, *Hemsl. Biol. Centr. Am. Bot.* ii. 79.—Mexic.
angustata, *Benth. ex Baker, in Mart. Fl. Bras.* vi. II. 190.—Bras.
Armani, *Baker, l. c.* 191.—Bras.
cinerea, *Benth. ex Baker, l. c.*—Bras.
fastigiata, *Benth. ex Baker, l. c.* 192.—Bras.
foliosa, *Benth. & Hook. f. Gen.* ii. 240.—Bras.
myriadenia, *Baker, in Mart. Fl. Bras.* vi. II. 192.—Bras.
Pohlii, *Baker, l. c.* 190.—Bras.
polyphylla, *Baker, l. c.* 191.—Bras.
spilanthoides, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 238 = Gymnocoronis spilanthoides.

ALONA, Lindl. Bot. Reg. (1844) t. 46. *CONVOLVULACEAE*, Benth. & Hook. f. ii. 879.
Rayeria, Gaudich. Voy. Bonite, t. 108 (1839–46).
baccata, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
balsamiflua, *Gaudich. Voy. Bonite*, t. 107.—Hab. ?
carnosa, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
coelestis, *Lindl. l. c.*—Chili.
descrticola, *Phil. Fl. Atac.* 44.—Chili.
dubia, *Phil. in Anales Univ. Chil.* (1865) ii. 350.—Chili.
ericifolia, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 500.—Chili.
floribunda, *Phil. in Anales Univ. Chil.* (1873) 525.—Chili.
glandulosa, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
longifolia, *Lindl. l. c.*—Chili.
micrantha, *Phil. Fl. Atac.* 44.—Chili.
microphylla, *Miers, in Hook. Lond. Journ. Bot.* iv. (1845) 501.—Chili.
mollis, *Phil. Fl. Atac.* 44.—Chili.
obtusa, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
pusilla, *Phil. Fl. Atac.* 45.—Chili.
revoluta, *Lindl. Bot. Reg.* (1844) sub t. 46.—Peruv.
rostrata, Lindl. l. c. = Osteocarpus rostratus.
sphaerophylla, *Phil. Fl. Atac.* 44.—Chili.
tomentosa, *Lindl. Bot. Reg.* (1844) sub t. 46.—Chili.
vernicosa, *Phil. Fl. Atac.* 44.—Chili.
xerophila, *Phil. l. c.*—Chili.

ALONSOA, Ruiz & Pav. Syst. Veg. 150 (1798). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 930.
Hemimeris, Pers. Syn. ii. 162 (1807) partim; H. B. & K. Nov. Gen. et Sp. ii. 376 (1817).

ALONSOA :—
Hemitomus, L'Hérit. ex Desf. Tabl. ed. I. 66 (1804).
Schistanthe, Kunze, in Linnaea, xvi. (1842) Litt. 109.
acutifolia, *Ruiz & Pav. Syst. Veg.* 153.—Peruv.
caulialata, *Ruiz & Pav. l. c.* 152.—Peruv.
hirsuta, *Steud. Nom.* ed. II. i. 60.—Bras.
incisifolia, *Ruiz & Pav. Syst. Veg.* 154.—Peruv.
intermedia, Lodd. Bot. Cab. t. 1456 = linearis.
linearifolia, *Steud. Nom.* ed. II. i. 60.—Peruv.
linearis, *Ruiz & Pav. Syst. Veg.* 154.—Peruv.
linifolia, *Roezl, in Belg. Hortic.* xxiv. (1874) 38.—Peruv.
Mathewsii, *Benth. in DC. Prod.* x. 250.—Peruv.
Mutisii, G. Don, Gen. Syst. iv. 513 = caulialata.
myrtifolia, *Roezl, in Belg. Hortic.* xxiv. (1874) 39.—Peruv.
parviflora, *G. Don, Gen. Syst.* iv. 513.—Venezuela.
procumbens, *Ruiz & Pav. Fl. Per.* 154.—Peruv.
quadrifolia, *G. Don, Gen. Syst.* iv. 513.—Mexic.
unilabiata, Ruiz & Pav. ex Steud. Nom. ed. I. 57, 31 = Diascia unilabiata.
urticaefolia, Steud. l. c. ed. II. i. 61 = incisifolia.
Warscewiczii, *Regel, in Gartenfl.* iii. (1854) 211. t. 91.—Peruv.

ALONZOA, Brongn. Enum. Genr. 59 (1843) = praec.

ALOPECIAS, Stev. in Bull. Soc. Nat. Mosc. iv. 266 (1832) = **Astragalus**, Tourn. (Legum.).
alopecuroides, Stev. l. c. = Astr. alopecuroides.
barbatus, Stev. l. c. 266 = Astr. barbatus.
chartaceus, Stev. l. c. xxix. (1856) II. 143 = Astr. chartaceus.
Clausii, Stev. l. c. 143 = Astr. vulpinus.
dasyanthus, Stev. l. c. iv. (1832) 266 = Astr. dasyanthus.
dasycephalus, Stev. l. c. 266 = Astr. roseus.
emarginatus, Stev. l. c. 266 = Astr. emarginatus.
Kotschyanus, Stev. l. c. xxix. (1856) II. 143 = Astr. Kotschyanus.
lanatus, Stev. l. c. iv. (1832) 266 = Astr. lanatus.
Libanotis, Stev. l. c. 266 = Astr. lanatus.
macrocephalus, Stev. l. c. 266 = Astr. macrocephalus.
narbonensis, Stev. l. c. 266 = Astr. narbonensis.
orbiculatus, Stev. l. c. xxix. (1856) II. 143 = Astr. orbiculatus.
ponticus, Stev. l. c. iv. (1832) 266 = Astr. ponticus.
roseus, Stev. l. c. xxix. (1856) II. 143 = Astr. roseus.
Sieversianus, Stev. l. c. 143 = Astr. Sieversianus.
sphaerocephalus, Stev. l. c. iv. (1832) 266 = Astr. macrocephalus.
vulpinus, Stev. l. c. 266 = Astr. vulpinus.

ALOPECURUS, Linn. Syst. ed. I. (1735); Gen. ed. I. 18 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1140.
Colobachne, Beauv. Agrost. 22. t. 6 (1812).
Tozzetia, Savi, in Mem. Soc. Ital. Sc. viii. (1798) 477.
aequalis, Sobol. Fl. Petrop. 16 = fulvus.
affinis, *Desv. Opusc.* 55.—Hab. ?
agrestis, *Linn. Sp. Pl.* ed. II. 89.—Europ.; Cauc.
agrestis, Sieber, ex Kunth, Enum. Pl. i. 24 = creticus.
alpestris, Wahlenb. ex Steud. Nom. ed. II. i. 61 = pratensis.
alpinus, *Sm. Fl. Brit.* iii. 1086.—Reg. bor. et arct.; Chili.
altissimus, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 13 = arundinaceus.
angustifolius, Sibth. & Sm. Fl. Graec. t. 64 = vaginatus.
antarcticus, Gies. in Brewst. Edinb. Cycl. x. 494 = alpinus.
antarcticus, Lechler, ex Steud. Syn. Pl. Gram. 148 = Lechleri.
antarcticus, Vahl, Symb. Bot. ii. 18 = alpinus.
anthoxantoides, *Boiss. Diagn.* Ser. I. xiii. 42.—Oriens.
aquaticus, Dum. Obs. Gram. Belg. 133 = nigricans.
aristatus, Gouan, Hort. Monsp. 37 = Polypogon monspeliensis.
aristulatus, *Michx. Fl. Bor. Am.* i. 43.—Europ.; Am. bor.
arundinaceus, *Poir. Encyc.* viii. 766.—Europ.; Am. bor.; Oriens.
arvensis, Gromow. ex Trautv. in Act. Hort. Petrop. ix. (1884) 353 = pratensis.

ALOPECURUS :—
Aucheri, Balansa, in Bull. Soc. Bot. Fr. xxi. (1874) 11 = vaginatus.
Aucheri, *Boiss. Diagn.* Sér. I. xiii. 41.—Persia.
australis, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 412 = geniculatus.
baikalensis, Turcz. ex Ledeb. Fl. Ross. iv. 462 = brachystachyus.
borealis, Trin. Fund. Agrost. 58 = alpinus.
brachyglossus, Peterm. Fl. Lips. (1844) 232 = pratensis.
brachystachyus, *Bieb. Fl. Taur. Cauc. Suppl.* 56.—Sibir.
brachystachyus, C. Koch, in Linnaea, xxi. (1848) 383 = vaginatus.
brachystylus, Peterm. Fl. Lips. (1844) 230 = pratensis.
bulbosus, With. Bot. Arr. Brit. Pl. ed. II. i. 59 = geniculatus.
bulbosus, Poir. Voy. Barb. ii. 94 = macrostachyos.
caesius, Schur, Enum. Pl. Transs. 727 in syn. = geniculatus.
cacspitosus, *Trin. Sp. Gram.* [xxi. t. 241. *sec. Pritzel, Ic. Ind.*].—Am. bor.
candicans, Salzm. ex Steud. Nom. ed. II. i. 61 = pratensis.
capensis, Eckl. ex Steud. Syn. Pl. Gram. 294 = Koeleria Alopecuros.
capensis, *Thunb. Prod. Fl. Cap.* 19.—Afr. austr.
capillaris, *Raeusch. Nom.* ed. III. 20.—Europ.
capitatus, Lam. Illustr. i. 168 = Phleum Gerardi.
carolinianus, Spreng. Nachr. i. Bot. Gart. Halle, 10 = Agrostis Cornucopiae.
carolinianus, *Walt. Fl. Carol.* 74.—Am. bor.
cassius, Boiss. Diagn. Ser. I. xiii. 41 = vaginatus.
castellanus, *Boiss. & Reut. Diagn. Pl. Hispan.* 26 = brachystachyus.
caudatus, Thunb. in Nov. Act. Ups. iii. (1780) 208 = Brylkinia caudata.
caudicans, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 707 = pratensis.
ciliatus, All. Fl. Pedem. ii. 235 = Koeleria phleoides.
ciliatus, Vitm. Summa Pl. i. 178 = Gerardi.
coerulescens, *Steud. & Hochst. ex Steud. Nom.* ed. II. i. 60, nomen.—As. Min.
colobachnoides, Trin. Fund. Agrost. 58 = brachystachyus.
creticus, *Trin. in Spreng. N. Entdeck.* ii. 45.—As. Min.; Crete.
creticus, Willk. ex Willk. & Lange, Prod. Fl. Hispan. i. 41 = agrestis.
crypsoides, *Griseb. Spicil. Fl. Rumel.* ii. 466.—Macedonia.
dasyanthus, Trautv. in Act. Hort. Petrop. v. (1877) 486 = vaginatus.
diandrus, *Griff. Notul.* iii. 11.—Ind. or.
echinatus, Thunb. Prod. Pl. Cap. 19 = Lasiochloa adscendens.
effusus, Link, ex Kunth, Enum. Pl. i. 328 = Agrostis effusa.
elongatus, *Peterm. in Flora*, xxvii. (1844) 230.—Europ.
elongatus, Poir. Encyc. Suppl. v. 495 = Polypogo elongatus.
exaltatus, *Less. in Linnaea*, ix. (1834) 207.—Sibir. ural.
fallacinus, *Gren. Fl. Mass. Adven.* (1859) 21.—Hab. ?
fasciculatus, Poir. Encyc. viii. 780 = Chaeturus fasciculatus.
foliosus, Clark, Trav. ii. 261 = vaginatus.
fulvus, Forsk. ex Steud. Nom. ed. II. i. 61 = Heleochloa alopecurioides.
fulvus, Sm. Engl. Bot. xxi. t. 1467 = geniculatus.
geniculatus, *Linn. Sp. Pl.* ed. I. 60.—Reg. bor. temp.
geniculatus, Lindh. ex Scheele, in Linnaea, xxii. (1849) 340 = Setaria polystachya.
geniculatus, Sibth. ex Steud. Nom. ed. II. i. 61 = Heleochloa alopecurioides.
Gerardi, *Vill. Fl. Delph.* 5; *Hist. Pl. Dauph.* ii. 66. t. 2.—Europ.; Oriens.
Giesseckeanus, Roem. & Schult. Mant. ii. 154 = alpinus.
glacialis, *C. Koch, in Linnaea*, xxi. (1848) 382.—Oriens.
glaucus, *Less, l. c.* ix. (1834) 206.—Sibir. ural.
glomeratus, Poir. Encyc. Suppl. v. 495 = Muehlenbergia glomerata.
gracilis, Trautv. in Act. Hort. Petrop. v. (1877) 486 = glacialis.
gracilis, *Willd. ex Trin. Agrost.* i. 16.—Carolina.
granulatus, *Beauv. Agrost.* 4, nomen.—Hab. ?

ALOPECURUS :—
hordeiformis, Linn. Sp. Pl. 60 = Pennisetum compressum.
hordeiformis, Willd. ex Steud. Nom. ed. II. i. 61 = Pennisetum orientale.
hybridus, *Wimm. Fl. Schles.* ed. III. (1857) 31.—Europ.
indicus, Murr. Syst. Veg. ed. XIII. 92 = Pennisetum Linnaei.
intermedius, *Hallier, in Bot. Zeit.* xxi. (1863) *App.* i. 6.—Ins. Heligol.
intermedius, Bock, ex Steud. Nom. ed. II. i. 61 = arundinaceus.
interruptus, Poir. Encyc. Suppl. v. 495 = Polypogon interruptus.
japonicus, *Steud. Syn. Pl. Gram.* 149.—Japon.
Konradii, Opiz, ex Mert. & Koch, in Roehl. Deutschl. Fl. i. 481 = geniculatus.
laguriformis, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 182 = brachystachyus.
laguroides, Balansa, in Bull. Soc. Bot. Fr. xxi. (1874) 11 = vaginatus.
lanatus, *Sibth. & Sm. Prod.* i. 43.—Oriens.
lasiochlamys, Boiss. Fl. Orient. v. 490 = textilis.
lasiostachyus, *Link, Hort. Berol.* i. 71.—Hispan.
latifolius, *Cav. Descr.* 37.—Ins. Tobago.
Lechleri, *Steud. Syn. Pl. Gram.* 148.—Valdivia.
litoreus, All. Fl. Pedem. ii. 235 = Koeleria villosa.
longearistatus, *Maxim. Prim. Fl. Amur.* 327.—Manchuria.
longiradiatus, Regel, ex Trautv. in Act. Hort. Petrop. ix. (1884) 354 = longearistatus.
longistylus, *Opiz, in Lotos*, ii. (1852) 228.—Bohem.
macrostachyos, *Poir. Encyc.* viii. 775.—Afr. bor.
magellanicus, Lam. Illustr. i. 168 = antarcticus.
malacostachyus, A. Gray, in Perry, Exped. Jap. ii. 328 = japonicus.
maritimus, Poir. Encyc. viii. 780 = Polypogon maritimus.
monspeliensis, Linn. Sp. Pl. 61 = Polypogon monspeliensis.
muticus, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 527 = arundinaceus.
myosuroides, Huds. Fl. Angl. 23 = agrestis.
nepalensis, *Trin. ex Steud. Syn. Pl. Gram.* 148.—Nepalia.
nigrescens, Jacq. Ecl. ii. t. 13 = arundinaceus.
nigricans, Hornem. Hort. Hafn. i. 68 = arundinaceus.
nigricans, Koch, Syn. Fl. Germ. ii. 896 = pratensis.
nigricans, Wichura, in Jahrb. Schles. Ges. (1846) 61 = hybridus,
obscurus, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 13 = pratensis.
otiporensis, *Griff. Notul.* iii. 94.—Affghan.
ovatus, Forst. Fl. Austral. 8 = alpinus?
ovatus, Knapp, Gram. Brit. 15 = alpinus.
Pallasii, Trin. Sp. Gram. t. 2 = vaginatus.
pallidus, *Dum. Obs. Gram. Belg.* 133.—Belg.
palulosus, Beauv. ex Mert. & Koch, in Roehl. Deutschl. Fl. i. 481 = geniculatus.
palulosus, Crantz, Inst. i. 368 = Polypogon monspeliensis.
paniceus, Cav. Descr. 37 = Polypogon maritimus.
paniceus, O. Muell. Fl. Dan. t. 861 = geniculatus.
paniceus, Linn. Sp. Pl. ed. II. 90; Lam. Fl. Fr. iii. 560 = Polypogon monspeliensis.
paniculatus, *Gueldenst. ex Ledeb. Fl. Ross.* iv. 465.—Reg. Caucas.
pedalis, Bosc, ex Beauv. Agrost. 4 = gracilis.
pedatus, Steud. Nom. ed. II. i. 61 = praec.
phalaroides, C. Koch, in Linnaea, xix. (1847) 5 = Gerardi.
phleiformis, Beauv. Agrost. 4 = villosus.
ponticus, C. Koch, in Linnaea, xxi. (1848) 382 = glacialis.
prutensis, Bourg. ex Lange, in Bull. Soc. Bot. Fr. xi. (1864) 47 = castellanus.
pratensis, *Linn. Sp. Pl.* 60.—Reg. bor. temp.
pratensis, Trin. Sp. Gram. t. 45 = arundinaceus.
pronus, Mitt. ex Hook. f. Steud. Fl. ed. I. 428 = geniculatus.
purpurascens, Link, in Linnaea, xvii. (1843) 400 = agrestis.
ramosus, *Poir. Encyc.* viii. 776.—Carolina.
repens, Bieb. Fl. Taur. Cauc. iii. 54 = arundinaceus.
ruthenicus, Weinm. Fl. Dan. t. 2881; et Cat. Dorpat. (1810) 10 = arundinaceus.
saccatus, *Vasey, in Coult. Bot. Gaz.* vi. (1881) 290.—Oregon.

ALOPECURUS :—
scaber, Opiz, Natural. n. ix. (1825) 166 = pratensis.
sericeus, Gaert. Fruct. i. 2. t. 1. f. 2 = pratensis.
setarioides, *Gren. in Mém. Soc. Emul. Doubs*, Sér. III. ii. (1858) 459.—Gallia.
sibiricus, Hort. Vind. ex Roem. & Schult. Syst. ii. 271 = arundinaceus.
subaristatus, Pursh, Fl. Am. Sept. i. 66 = geniculatus.
tenellus, Poir. Encyc. Suppl. v. 496 = Polypogon tenellus.
textilis, *Boiss. Diagn.* Ser. I. xiii. 40.—Oriens.
tonsus, Dum. Obs. Gram. Belg. 132 = agrestis.
transsilvanicus, Schur, Enum. Fl. Transs. 727 = brachystachyus.
trivialis, Seidl, ex Opiz, Natural. n. ix. (1825) 116 = pratensis.
turicensis, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 111.—Europ.
tuscheticus, Trautv. in Act. Hort. Petrop. ix. (1884) 354 = gracilis.
typhoides, Burm. f. Fl. Ind. 27 = Penicillaria spicata.
uralensis, *Uspenski, Ekaterinenburg.* 24.—Sibir. ural.
utriculatus, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 243.—Europ.; Oriens.
vaginatus, *Pall. in Nov. Act. Acad. Petrop.* x. (1792) 304, nomen.—Tauria; Oriens.
variegatus, *Steud. Syn. Pl. Gram.* 148.—Chili.
varius, Auct. ex Steud. Nom. ed. I. 32, 771 = Sesleria caerulea.
ventricosus, Huds. Fl. Angl. ed. II. i. 28 = Gastridium australe.
ventricosus, Pers. Syn. i. 80 = arundinaceus.
villosus, Gilib. Exercit. ii. 517 = pratensis.
villosus, *Poir. Encyc.* viii. 777.—Helvet.
Vlassovii, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. II. (1845) 42 = brachystachyus.

ALOPHIA, Herb. in Bot. Mag. sub t. 3779 (1838) = **Herbertia**, Sweet (Irid.).
Drummondiana, Herb. l. c. = H. Drummondiana.
geniculata, Klatt, in Mart. Fl. Bras. iii. 1. 517.—Bras.
linearis, Klatt, in Linnaea, xxxi. (1861–62) 558 = H. lineata.
linearis, Warm. in Kjoeb. Vidensk. Meddel. (1872) 128.—Bras.
pulchella, Benth. & Hook. f. Gen. iii. 692 = H. pulchella, lineata.
Sellowiana, Klatt, in Linnaea, xxxi. (1861–62) 557.—Bras.

ALOPHIUM, Cass. in Dict. Sc. Nat. liv. 493 (1828) = **Centaurea**, Linn. (Compos.).
tenuifolium, Cass. l. c. = C. Alophium.

ALOPHOCHLOA, Endl. Gen. 100 (1836) *sphalm.* = LOPHOCHLOA, Reichb. = **Koeleria**, Pers. (Gram.).

ALOPHYLLUS, Linn. Sp. Pl. ed. II. 496 (1762) = ALLOPHYLUS, Linn. (1747) = **Schmiedelia**, Linn. (1764) (Sapind.).

ALOPICARPUS, Neck. Elem. ii. 369 (1790) = **Paris**, Rupp. (Liliac.).

ALORANTHUS, F. S. Voigt, in Rich. Anal. Frucht, 29 (1811) *sphalm.* = **Chloranthus**, Sw.

ALOSEMIS, Rafin. Sylva Tellur. 96 (1838) = **Melastoma**, Linn.
grandiflora, Rafin. l. c.—Hab.?
villosa, Rafin. l. c.—Hab.?
zeylanica, Rafin. l. c.—Zeylan.

ALOYSIA, Ort. & Palav. ex L'Hérit. Stirp. i. 21 (1784) = **Lippia**, Linn. (Verben.).
attenuata, Walp. Rep. iv. 42.—Bras.
chamaedryfolia, Cham. in Linnaea, vii. (1832) 234 = L. chamaedryfolia
chamaedryoides, Steud. Nom. ed. II. i. 62 = L. chamaedryfolia.
citriodora, Orteg. ex Pers. Syn. ii. 139 = L. citriodora.
floribunda, Mart. & Gal. in Bull. Acad. Brux. xi. (1844) II. 320 = L. lycioides.
lycioides, Cham. in Linnaea, vii. (1832) 237 = L. lycioides.
polygalaefolia, Cham. l. c. 236 = L. polygalaefolia.
scorodonioides, Cham. l. c. 234 = L. scorodonioides.
urticoides, Cham. l. c. 238 = L. urticoides.
virgata, Juss. in Ann. Mus. Par. vii. (1806) 73 = L. urticoides.

ALPHANDIA, Baill. Adansonia, xi. (1873) 85. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 300.
furfuracea, *Baill. l. c.* 86.—N. Caled.
resinosa, *Baill. l. c.* 86.—N. Caled.

ALPHITONIA, Reissek, ex Endl. Gen. 1098 (1840). *RHAMNEAE*, Benth. & Hook. f. i. 381.
Emmenosperma, F. Muell. Fragm. iii. 63 = Emmenosperma alphitonoides.
erubescens, *Baill. Adansonia*, xi. (1874–76) 271.—N. Caled.
excelsa, *Reissek, ex Endl. Gen.* 1098.—Austral.
franguloides, *A. Gray, Bot. U. St. Expl. Exped.* i. 280.—Ins. Viti.
moluccana, *Teijsm. & Binn. Cat. Hort. Bogor.* 221.—Molucca.
xerocarpus, *Baill. Adansonia*, xi. (1873–76) 270.—N. Caled.
zizyphoides, *A. Gray, Bot. U. St. Expl. Exped.* i. 278.—Hab.?

ALPHONSEA, Hook. f. & Thoms. Fl. Ind. i. 152 (1855). *ANONACEAE*, Benth. & Hook. f. i. 29.
ceramensis, *Scheff. in Ann. Jard. Buitenz.* ii. (1885) 29.—Malaya.
elliptica, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 90.—Malaeca.
javanica, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 340; *in Flora*, liii. (1870) 242.—Java.
lutea, *Hook. f. & Thoms. Fl. Ind.* i. 153.—Ind. or.
lutea, Teijsm. & Binn. ex Scheff. in Flora, liii. (1870) 242 = javanica.
madraspatana, *Bedd. Fl. Sylv.* i. t. 76.—Ind. or.
Maingayi, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 90.—Malacea.
sclerocarpa, *Thw. Enum. Pl. Zeyl.* 11.—Zeylan.
ventricosa, *Hook. f. & Thoms. Fl. Ind.* i. 152.—Ind. or.; Malaya.
zeylanica, *Hook. f. & Thoms. l. c.* 153.—Ind. or.

ALPINIA, Linn. Gen. ed. I. 332 (1737). *SCITAMINEAE*, Benth. & Hook. f. iii. 648.
ALBINA, Giseke, Prael. Linn. 199, 207 (1792).
ALUGHAS, Linn. Fl. Zeyl. 207 (1747).
BANGLIUM, Buch.-Ham. ex Wall. Cat. sub n. 6579 (1832).
BOJERIA, Rafin. Fl. Tellur. iv. 50 (1836).
BUEKIA, Giseke, Prael. Linn. 199, 204 (1792).
CATIMBIUM, Juss. Gen. 62 (1789).
CENOLOPHON, Blume, Enum. Pl. Jav. 60 (1830).
DOXANTHES, Rafin. Fl. Tellur. iv. 51 (1836).
GALANGA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2; ed. II. 65 (1827).
HELLENIA, Willd. Sp. Pl. i. 4 (1797).
HERITIERA, Retz. Obs. vi. 17. t. 1 (1791).
KOLOWRATIA, Presl, Rel. Haenk. i. 113. t. 20 (1827).
LANGUAS, Koen. in Retz. Obs. iii. 64 (1783).
MARTENSIA, Giseke, Prael. Linn. 199, 207 (1792).
MONOCYSTIS, Lindl. Introd. Nat. Syst. ed. II. 445 (1836).
ZERUMBET, Wendl. Sert. Hannov. fasc. iv. 3. t. 19 (1798).
acrostachya, Steud. Nom. ed. II. i. 62 (= Amomi sp.). —Java.
alba, D. Dietr. Syn. Pl. i. 12 = Elettaria alba.
alba, *Rosc. in Trans. Linn. Soc.* viii. (1807) 346.—Ind. or.
alata, *A. Dietr. Sp. Pl.* i. 45.—China.
Allughas, *Rosc. in Trans. Linn. Soc.* viii. (1807) 346.—Ind. or.
Antillarum, *Roem. & Schult. Syst. Veg.* i. 20.—Ind. occ.
aquatica, *Rosc. in Trans. Linn. Soc.* viii. (1807) 346.—Ind. or.
arctiflora, *F. Muell. Fragm.* viii. 25.—Austral.
aromatica, Aubl. Pl. Gui. i. 3 = Renealmia domingensis.
auriculata, *Rosc. Scit. Pl.* t. 74.—Ins. Molucc.
Blumei, D. Dietr. Syn. Pl. i. 13 = javanica.
Boia, *Seem. Fl. Vit.* 290.—Ins. Fiji.
bracteata, Rosc. in Trans. Linn. Soc. xi. (1815) 281 = alata.
bracteata, Roxb. Fl. Ind. i. 61 = Roxburghii.
brevilabris, *Presl, Rel. Haenk.* i. 110. t. 17.—Ins. Luzon.
breviscapa, Poepp. & Endl. Nov. Gen. et Sp. ii. t. 135 = Renealmia breviscapa.
caerulea, *Benth. Fl. Austral.* vi. 265.—Austral.
calcarata, *Rosc. in Trans. Linn. Soc.* viii. (1807) 347.—Ind. or.; China.

ALPINIA:—
capitellata, *Jack, in Malay. Misc.* ii. (1822) n. VII. 4. —Malaya.
Cardamomum, Roxb. in As. Res. xi. (1810) 356 = Elettaria Cardamomum.
Cardamon-medium, Roxb. Fl. Ind. i. 72 = Amomum costatum.
caribaea, Gaert. Fruct. i. 36. t. 12 = Renealmia racemosa.
carnea, *Griff. Notul.* iii. 420.—Ind. or.
cernua, *Sims, Bot. Mag.* t. 1900.—Ind. or.
chinensis, *Rosc. in Trans. Linn. Soc.* viii. (1807) 346.—China.
coccinea, D. Dietr. Syn. Pl. i. 12 = Amomum coccineum.
comosa, Jacq. Coll. iv. 112 = Costus comosus.
conchigera, *Griff. Notul.* iii. 424.—Ind. or.
costata, Roxb. Corom. Pl. iii. 48. t. 252 = Amomum costatum.
cristata, *Griff. Notul.* iii. 421.—Ind. or.
diffissa, *Rosc. in Trans. Linn. Soc.* xi. (1815) 281.—China.
elatior, *Jack, in Malay. Misc.* ii. (1822) n. VII. 2.—Malaya.
erecta, Lodd. ex Steud. Nom. ed. II. i. 62 = calcarata.
exaltata, Roem. & Schult. Syst. i. 21. add. 563 = Renealmia exaltata.
fasciculata, Steud. Nom. ed. II. i. 62 = Amomum fasciculatum.
ferruginea, Steud. Nom. ed. II. i. 62 = mollis.
foetens, D. Dietr. Syn. Pl. i. 12 = Amomum foetens.
Galanga, *Willd. Sp. Pl.* i. 12.—As. trop.
gigantea, *Blume, Enum. Pl. Jav.* 59.—Java.
gigantea, F. Vill. in Blanco, Fl. Filip. ed. III. Nov. App. 225 = gracilis.
globosa, *Horan. Prod. Scitam.* 34.—China.
gracilis, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 316.—Ins. Philipp.
Haenkei, *Presl, Symb.* 66. t. 43.—Chili.
Hamiltoniana, *Wall. Cat.* n. 6579.—Ind. or.
hemisphaerica, D. Dietr. Syn. Pl. i. 13 = Elettaria hemisphaerica.
hirsuta, *Horan. Prod. Scitam.* 34.—Cochinch.
humilis, Vell. Fl. Flum. 1; i. t. 4 (= Renealmiae sp.).—Bras.
humilis, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 327.—Java.
involucrata, *Griff. Notul.* iii. 422.—Ind. or.
jamaicensis, Gaert. Fruct. i. 36. t. 12 = Renealmia jamaicensis.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 140.—Japon.
javanica, *Blume, Enum. Pl. Jav.* 59.—Java.
javanica, D. Dietr. Syn. Pl. i. 13 = *acrostachya*.
latifolia, Willd. ex Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 130 = Calathea latifolia.
linguiformis, Roxb. Fl. Ind. i. 73 = Elettaria linguiformis.
macrantha, Poepp. & Endl. Nov. Gen. et Sp. t. 134 = Renealmia macrantha.
macrantha, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 56.—Malaya.
maculata, *Rosc. in Trans. Linn. Soc.* viii. (1807) 347.—Ind. or.
magnifica, Rosc. Scit. t. 75 = Amomum magnificum.
malaccensis, Presl, Rel. Haenk. i. 110 = Haenkei.
malaccensis, *Rosc. in Trans. Linn. Soc.* viii. (1807) 345.—Ind. or.; Java.
media, Spreng. Syst. i. 14 = Amomum costatum.
mesana, Horan. Prod. Scitam. 27 = Zingiber macrostachyum.
minor, D. Dietr. Syn. Pl. i. 12 = Elettaria minor.
minuta, D. Dietr. l. c. = Elettaria minuta.
missionis, *Wall. Cat.* n. 6589.—Ind. or.
mollis, *Presl, Rel. Haenk.* 110.—Ins. Luzon.
multicaulis, Roem. & Schult. Syst. i. 21 = Renealmia domingensis.
mutica, *Roxb. in As. Res.* xi. (1810) 354.—Ind. or.
Neesana, *J. Grah. Cat. Bomb. Pl.* 207.—Ind. or.
Nimmonii, J. Grah. l. c. 206 = Zingiber Nimmonia.
nutans, *Rosc. in Sm. Exot. Bot.* ii. 93.—Ind. or.
occidentalis, Sw. Prod. Veg. Ind. Occ. 11 = Renealmia jamaicensis.
officinarum, *Hance, in Journ. Linn. Soc.* xiii. (1873) 6.—China.
oxyphylla, *Miq. in Journ. Bot. Néerl.* i. (1861) 93.—China.
Paco-Secora, Jacq. Fragm. 54. n. 17. t. 76 = Renealmia Paco-Secora.
pallida, D. Dietr. Syn. Pl. i. 12 = Elettaria pallida.

ALPINIA:—
paludosa, D. Dietr. Syn. Pl. i. 12 = Elettaria paludosa.
papuana, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 56.—N. Guin.
parviflora, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 316.—Ins. Philipp.
penicillata, *Rosc. in Trans. Linn. Soc.* xi. (1814) 280.—China.
phoenicea, *Kamphoevener, ex Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 154, nomen.—Ins. Nicobar.
Pininga, *D. Dietr. Syn. Pl.* i. 12.—Java.
porrecta, *Wall. Cat.* n. 6576.—Ind. or.
Presliana, Steud. Nom. ed. II. i. 62 = Renealmia racemosa.
pumila, *Hook. f. Bot. Mag.* t. 6832.—China.
punicea, Roxb. Fl. Ind. i. 71 = Elettaria punicea.
pyramidata, *Blume, Enum. Pl. Jav.* 58.—Java.
racemigera, *F. Muell. Fragm.* viii. 27.—Austral.
racemosa, Linn. Sp. Pl. ed. I. 2 = Renealmia racemosa.
racemosa, Sw. Obs. 5 = A. Antillarum.
Rafflesiana, *Wall. Cat.* n. 6575.—Ind. or.
Renealmia, Sm. in Rees, Cyclop. Suppl. xxxix. n. 14 = Renealmia exaltata.
Rheedii, *Wight, Ic.* 6. t. 2026.—Malabar.
Roscoeana, Steud. Nom. ed. II. i. 63 = alata.
Roxburghii, *Sweet, Hort. Brit.* ed. I. 390.—Ind. or.
rubra, *D. Dietr. Syn. Pl.* i. 12.—Java.
Ruitziana, Steud. Nom. ed. II. i. 63 = Renealmia Ruiziana.
serrulata, *Willd. ex Steud. l. c.*—Bras.
sessilis, Koen. in Retz. Obs. fasc. iii. 64 = Kaempferia Galanga.
Simsii, *Gasp. Observ. Boccodifalc.* 5.—Hab. ?
solaris, D. Dietr. Syn. Pl. i. 12 = Elettaria solaris.
speciosa, D. Dietr. l. c. 13 = Elettaria speciosa.
spicata, Jacq. Enum. Pl. Carib. 11 = Costus spicatus.
spicata, Presl, Rel. Haenk. i. 109 = Renealmia racemosa.
spicata, *Roxb. in As. Res.* xi. (1810) 356.—Ind. or.
spiralis, Jacq. Hort. Schoenb. i. 1 = Costus Pisonis.
stachyodes, *Hance, in Journ. Linn. Soc.* xiii. (1873) 126.—China.
striata, *Hort. Angl. ex Link, Enum.* i. 5.—Ind. or.
strobilifera, Poepp. & Endl. Nov. Gen. et Sp. t. 136 = Renealmia strobilifera.
thyrsoidea, Steud. Nom. ed. II. i. 63 = Renealmia thyrsoidea.
tomentosa, *D. Dietr. Syn. Pl.* i. 12.—Java.
tubulata, [*Rosc. in*] *Bot. Reg.* t. 777.—Ind. occ.
uviformis, *Horan. Prod. Scitam.* 35.—Malaya.
viridiflora, *Griff. Notul.* iii. 423.—Ind. or.
vitiensis, *Seem. Fl. Vit.* 290.—Ins. Fiji.
Walang, *D. Dietr. Syn. Pl.* i. 13.—Java.

ALSCHINGERA, Vis. Fl. Dalm. iii. 69 (1849) = **Physospermum**, Cusson (Umbell.).
acteaefolia, Vis. sec. Tomm. fide Nym. Consp. 315 = P. verticillatum.
verticillata, Vis. Fl. Dalm. iii. 69 = P. verticillatum.

ALSEIS, Schott, in Spreng. Syst. iv. Cur. Post. 404 (1827). *RUBIACEAE*, Benth. & Hook. f. ii. 38.
Blackiana, *Hemsl. Diag. Pl. Nov.* 30.—Panama.
floribunda, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 404.—Bras.

ALSENOSMIA, Endl. Ench. 284 (1841) sphalm. = **Alseuosmia**, A. Cunn. (Caprifol.).

ALSEODAPHNE, Nees, in Wall. Pl. As. Rar. ii. 71 (1831). *LAURINEAE*, [Benth. & Hook. f. iii. 157.]
EUPHOEBE, Blume, ex Meissn. in DC. Prod. xv. i. 58, 59 in syn. (1864).
NOTHAPHOEBE, Blume, Mus. Bot. Lugd. Bat. i. 328 (1851).
amara, Kostel. Allg. med.-pharm. Fl. ii. 484 = Nectandra cymbarum.
bancana, *Miq. Fl. Ind. Bat.* i. 915.—Banca.
breviflora, *Benth. Fl. Hongk.* 292.—China.
chinensis, *Champ. ex Meissn. in DC. Prod.* xv. i. 28.—Hongkong.
costalis, *Nees, in Wall. Pl. As. Rar.* ii. 72.—Ind. or.
Cujumary, Kostel. Allg. med.-pharm. Fl. ii. 484 = Ocotea Cujumary.
cymbarum, Kostel. l. c. = Nectandra cymbarum.
excelsa, *Blume, Mus. Bot. Lugd. Bat.* i. 332.—Java.
grandis, Kurz, For. Fl. Burma, ii. 293 = Dehaasia cuneata.

ALSEODAPHNE:—
grandis, Nees, in Wall. Pl. As. Rar. ii. 72 = Phoebe opaca.
lucida, *Nees, l. c.*—Singapore.
opifera, Kostel. Allg. med.-pharm. Fl. ii. 484 = Ocotea opifera.
polyneura, *Miq. Fl. Ind. Bat.* i. 916.—Penang.
semicarpifolia, *Nees, in Wall. Pl. As. Rar.* ii. 72.—Ind. or.

ALSEUOSMIA, A. Cunn. in Ann. Nat. Hist. ii. (1839) 209. *CAPRIFOLIACEAE*, Benth. & Hook. f. ii. 6.
FAGOIDES, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. ii. (1839) 210.
atriplicifolia, A. Cunn. l. c. 210 = Banksii.
Banksii, *A. Cunn. l. c.* 209.—N. Zel.
bupleurifolia, *A. Cunn. ex Heynh. Nom.* ii. 27.—N. Zel.
Hookeria, *Colenso, Excurs. North Isl.* 84.—N. Zel.
Ilex, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 210.—N. Zel.
ligustrifolia, A. Cunn. l. c. 209 = linariifolia.
linariifolia, *A. Cunn. l. c.*—N. Zel.
macrophylla, *A. Cunn. l. c.* 210.—N. Zel.
palaeiformis, A. Cunn. l. c. = Banksii.
pusilla, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 241.—N. Zel.
quercifolia, *A. Cunn. in. Ann. Nat. Hist.* ii. (1839) 210.—N. Zel.

ALSINANTHE, Reichb. Ic. Fl. Germ. v. (1841) 29. t. 209 = **Arenaria**, Linn. (Caryophyll.).
biflora, Reichb. l. c. 30. t. 209. f. 4939 = Ar. biflora.
stricta, Reichb. l. c. 29. t. 209. f. 4935 = Ar. stricta.
uliginosa, Reichb. l. c. t. 209. f. 4935 = Ar. stricta.

ALSINANTHEMOS, J. G. Gmel. Fl. Sib. iv. 46 (1769) = **Trientalis**, Linn. (Primul.).

ALSINANTHUS, Reichb. Handb. 298 (1837) = **Arenaria**, Linn. (Caryophyll.).
trinervius, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 347 = Ar. trinervia.

ALSINASTRUM, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 15 = **Elatine**, Linn.
callitrichoides, Rupr. Fl. Ingr. 197 = E. triandra.
galiifolia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 15 = E. Alsinastrum.
gyrospermum, Rupr. Fl. Ingr. 195 = E. Hydropiper.
orthospermum, Rupr. l. c. 196 = E. Hydropiper ?
triandrum, Rupr. l. c. 197 = E. triandra.

ALSINE, Burm. Fl. Zeyl. 13. t. 8 (1737) = **Mollugo**, Linn. (Ficoid.).

ALSINE, Tourn. ex Linn. Syst. ed. I. (1735); Gen. ed. I. 133 (1737) = **Arenaria**, Linn., **Stellaria**, Linn. &c. (Caryophyll.).

ALSINE, Scop. Fl. Carn. ed. II. 224 (1772); Wahlenb. Fl. Lapp. 127 (1812) = **Arenaria**, Linn. (Caryophyll.).
aizoides, Boiss. Diagn. Ser. I. i. 47.—Kurdistan.
alpina, Willk. in Bot. Zeit. v. (1847) 238 = Spergularia rubra.
Alsinastrum, Crantz, Inst. ii. 405 = Elatine Alsinastrum.
anatolica, Boiss. Diagn. Ser. I. viii. 97 = Ar. setacea.
apetala, Kit. ex Nym. Consp. 111 = Stellaria media.
arabica, Fisch. Mey. & Avé-Lall. Ind. Sem. Petrop. xi. 49 = Ar. tenuifolia.
arctica, Fenzl, Verbreit. Alsin. in tab. ad p. 18; et 36 = Ar. arctica.
Arduinei, Fenzl, l. c. in tab. ad p. 57 = Ar. graminifolia.
aretioides, Mert. & Koch, in Roehl. Deutsch. Fl. iii. 277 = Ar. aretioides.
aristata, [Soland. in] Russ. Aleppo, ed. II. ii. 249.—Syria.
arvensis, Crantz, Inst. ii. 408 = Spergula arvensis.
arvatica, Guss. Fl. Sic. Syn. i. 503 = Ar. mucronata.
attica, Boiss. & Sprun. Diagn. Ser. I. v. 84 = Ar. verna.
Aucheriana, Boiss. Fl. Orient. i. 678 = Ar. Aucheriana.
austriaca, Wahlenb. Fl. Lapp. 129 = Ar. austriaca.
avicularum, Lam. Fl. Fr. iii. 46 = Stellaria media.
barbata, Stokes, Bot. Mat. Med. ii. 537 = Stellaria media.

ALSINE :—
Barrelieri, Bor. in Mém. Soc. Acad. Maine-et-Loire, vi. (1859) 57.—Gallia.
Bartolotti, Huet, ex Nym. Consp. 119 = Arenaria procumbens.
banatica, Bluff & Fingh. Comp. ed. II. ii. 11. 99 = Ar. banatica.
Bauhinorum, J. Gay, ex Lamotte, Prod. i. 144 = Ar. linifolia.
bavarica, Crantz, Inst. ii. 407 = Ar. bavarica.
Biebersteinii, Rupr. Fl. Cauc. 215.—Caucas.
biflora, Wahlenb. Fl. Lapp. 128 = Ar. scandinavica.
Billardieri, Boiss. Diagn. Ser. I. viii. 95 = Ar. Meyeri.
bipartita, Gilib. Fl. Lituan. ii. 150 = Stellaria media.
brachycarpa, Boiss. Diagn. Ser. II. vi. 37 = Ar. tenuifolia.
brachypetala, Opiz, Natural. xi. (1826) 406 = Stellaria media.
brevicaulis, Sternb. ex Nym. Consp. 119 = Ar. laneeolata.
breviflora, Gilib. Fl. Lituan. ii. 150 = Ar. tenuifolia.
brevifolia, Chapm. Fl. S. Un. St. 49 = Ar. brevifolia.
brevis, Boiss. Diagn. Ser. I. viii. 96.—Persia ; Arabia.
Bufonia, Crantz, Inst. ii. 405 = Bufonia perennis.
caespitosa, Ehrh. ex Nym. Consp. 119 = Ar. verna.
campestris, Boiss. Voy. Espagne, ii. 723 (= *Minuartia campestris*).—Hispan.
campestris, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. montana.
canaliculata, Dulac, Fl. Hautes-Pyr. 246 = Ar. Cherleria.
capillipes, Boiss. ex Steud. Nom. ed. II. i. 63 = Ar. capillipes.
cappadocica, Boiss. Diagn. Ser. II. i. 88 = A. setacea.
catarctarum, Janka, in Közlem. Magyar Tudom. Akad. xii. (1876) 165.—Europ.
caucasica, Boiss. Diagn. Ser. II. i. 87 = Ar. montana.
caucasica, Rupr. Fl. Caucas. 203.—Caucas.
cerastiifolia, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. cerastiifolia.
Cherleriae, Fenzl, l. c. 37 = Stellaria petraea.
Cherleriana, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = Ar. Cherleria.
Cheveneauxii, Reut. ex Nym. Consp. 119 = Ar. recurva.
ciliata, Crantz, Inst. ii. 408.—Italia.
condensata, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. condensata.
conferta, Jord. Pugill. 35 = Ar. tenuifolia.
confusa, Heldr. & Sart. ex Nym. Consp. 118 = Ar. trichocalycina.
corymbulosa, Boiss. & Bal. Diagn. Ser. II. vi. 37 = Ar. setacea.
corymbulosa, Bor. in Mém. Soc. Acad. Maine-et-Loire, vi. (1859) 57 = Ar. Barrelieri.
costata, Fenzl, Verbreit. Alsin. 37 = Ar. costata.
Cupaniana, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 19 = Stellaria media.
decandra, Dalla Torre, Atl. Alpenfl. 78.—Europ.
decipiens, Fenzl, Pugill. Pl. Nov. Syr. 12.—Syria.
Delastrei, Bor. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 12 = Ar. Barrelieri.
diandra, Guss. Fl. Sic. Syn. i. 501 = Ar. diandra.
dianthifolia, Boiss. Diagn. Ser. I. viii. 99.—As. Min.
dichotoma, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Minuartia dichotoma.
Dillenii, Vill. Fl. Delphin. 30.—Gallia.
Drummondii, Fenzl, ex Torr. & Gray, Fl. N. Am. i. 675 = Stellaria Nuttallii.
Elatina, Crantz, Inst. ii. 405 = Elatine Hydropiper.
elegans, Fenzl, Verbreit. Alsin. tab. 18 = Ar. elegans.
elongata, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 19 = Stellaria media.
emarginata, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. emarginata.
erecta, Crantz, Inst. ii. 405 = Cerastium quaternellum.
erythrosepala, Boiss. Diagn. Ser. I. viii. 98.—As. Min.
eurytanica, Boiss. l. c. Ser. II. vi. 35.—Graecia.
extensa, Nym. Consp. 119 = Ar. procumbens.
Facchinii, Hausm. Fl. Tyr. 138.—Tyrol.
falcata, Grisch. Fl. Rumel. i. 200 = Ar. setacea.
fasciculata, Wahlenb. Fl. Lapp. 129 = Ar. fascieulata.
fastigiata, Bab. Man. ed. I. 51 = Ar. fastigiata.
formosa, Fenzl, in Flora, xxvi. (1843) 403.—Assyria.
Funkii, Jord. Pugill. 36 = Ar. fasciculata.
filiformis, Fenzl, ex Heynh. Nom. ii. 27 = Ar. picta.
Gerardi, Wahlenb. Fl. Carpat. 132 = Ar. verna.
Gieseckii, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. verna.

ALSINE :—
glabella, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 20 = Stellaria media.
glabra, A. Gray, Man. ed. II. 57 = Arenaria glabra.
glandulosa, Dulac, Fl. Hautes-Pyr. 246 = Ar. cerastiifolia.
glandulosa, Boiss. & Huet, Diagn. Ser. II. v. 61 = Ar. juniperina.
glandulosa, Mutel, ex Nym. Consp. 116 = Ar. linifolia.
globulosa, C. A. Mey. Verz. Pfl. Cauc. 219 = Ar. globulosa.
glomerata, Fenzl, Verbreit. Alsin. in tab. ad p. 46 = Ar. glomerata.
graminifolia, J. F. Gmel. Syst. 507 = Ar. Rosani.
grandiflora, Crantz, Inst. ii. 408 = Ar. grandiflora.
granuliflora, Fenzl, in Ledeb. Fl. Ross. i. 346 = Ar. setacea.
Griffithsii, Boiss. Diagn. Ser. II. i. 89 = Ar. Griffithii.
grineensis, Gren. & Godr. Fl. Fr. i. 252 = Ar. Villarsii.
groenlandica, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. groenlandica.
Gussonii, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 20 = Stellaria media.
Halophila, Turcz. in Bull. Soc. Nat. Mose. xv. (1842) 587.—Altai.
Heldreichiana, Boiss. & Heldr. in Boiss. Fl. Orient. i. 677 = Ar. juniperina.
Helmii, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. Helmii.
herniarioides, Rion, ex Nym. Consp. 119 = Ar. aretioides.
heteromalla, Bieb. Fl. Taur. Cauc. i. 350 = Ar. setacea.
heterosperma, Guss. Fl. Sic. Syn. i. 501.—Sicil.
hirsuta, Fenzl, Verbreit. Alsin. in tab. 46 = Ar. recurva.
hirta, Hartm. Handb. Skand. Fl. ed. III. 104 = Ar. verna.
hispanica, Fenzl, Verbreit. Alsin. in tab. ad p. 46 = Queria hispanica.
hispida, Crantz, Inst. ii. 408 = Ar. hispida.
hybrida, Jord. Pugill. 33 = Ar. tenuifolia.
imbricata, C. A. Mey. Verz. Pfl. Cauc. 217 = Ar. imbricata.
inamoena, C. A. Mey. l. c. 218 = Ar. imbricata.
intermedia, Boiss. Fl. Orient. i. 685.—As. Min.
intricata, Martr. Pl. Crit. Tarn, 15.—Gallia.
irregularis, Vis. ex Boiss. Fl. Orient. i. 684 = Ar. globulosa.
Jacquini, Koch, Syn. Fl. Germ. ed. II. 125 = Ar. fasciculata.
Jacquiniana, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = Ar. fasciculata.
juniperina, Wahlenb. Fl. Lapp. 129 = Ar. juniperina.
Juressi, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. Juressi.
kabylica, Pomel, Nouv. Mat. Fl. Atl. 327.—Afr. bor.
Kitaibelii, Nym. Consp. 116 = Ar. laricifolia.
lanceolata, Mert. & Koch, in Roehl. Deutsch. Fl. ed. III. ii. 275 = Ar. lanceolata.
lancifolia, Vis. Ic. Fl. Dalm. t. 34 = Ar. globulosa.
Langii, G. Reuss, Květ. Slov. (1853) 76 = Ar. laricifolia.
laricifolia, Crantz, Inst. ii. 407 = Ar. laricifolia.
laricifolia, Linn. ex Nym. Consp. 117 = Ar. setaeea.
laricina, Crantz, Inst. ii. 408 = Spergula laricina.
lateriflora, Crantz, l. c. 406 = Ar. lateriflora.
latifolia, Dum. Fl. Belg. 109.—Belg.
laxa, Jord. Pugill. 34 = Ar. tenuifolia.
leucocephala, Boiss. Diagn. Ser. I. i. 45.—As. Min.
libanotica, Boiss. l. c. viii. 98.—Libanus.
lineata, C. A. Mey. in Ledeb. Fl. Ross. = Ar. juniperina.
liniflora, Vis. Fl. Dalmat. iii. 1. 178 = Ar. liniflora.
linifolia, Borb. in Rad. Jugos. Akad. Zagrab, xxxvi. (1876) 166.—Europ. austr.
Lydia, Boiss. Diagn. Ser. II. i. 86 = Ar. tenuifolia.
macrocarpa, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. macrocarpa.
marginata, C. A. Mey. Verz. Pfl. Cauc. 217 = Spergularia media.
marginata, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Spergularia marginata.
marginata, Reichb. Fl. Germ. Excurs. 566. n. 3662 = Ar. media.
marginata, Schreb. Spicil. 31 = Spergularia pentandra.
marina, Webb & Berth. Phyt. Canar. i. 147 = Spergularia Bourgeaei.
marina, Mert. & Koch, in Roehl. Deutsch. Fl. ii. 293 = Spergularia media.

ALSINE :—
media, Linn. Sp. Pl. ed. I. 272 = Spergularia media.
media, Vell. Fl. Flum. 127 ; iii. t. 106 = Drymaria cordata.
mediterranea, Gren. Fl. Massil. Adven. (1859) 8.—Gallia.
mesogitana, Boiss. Diagn. Ser. I. i. 45 = Arenaria tenuifolia.
Meyeri, Boiss. l. c. Ser. I. viii. 96 = Arenaria globulosa.
mexicana, Bartl. ex Presl, Rel. Haenk. ii. 14 = Ar. mexicana.
Michauxii, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. stricta.
microsperma, Fenzl, ex Torr. & Gray, Fl. N. Am. i. 674 = Ar. Pitcheri.
minuta, Rohrb. in Linnaea, xxxvii. (1871–72) 246.—Chili.
Moehringia, Crantz, Inst. ii. 405 = Ar. muscosa.
moehringioides, Rohrb. in Linnaea, xxxvii. (1871–73) 247 (= *Hymenella moehringioides*).
molluginea, Lag. Gen. et Sp. Nov. (1815) 13 = Ar. molluginea.
montana, Crantz, Inst. ii. 406 = Ar. montana.
mucronata, Linn. Sp. Pl. ed. II. 389 = Ar. mueronata.
mucronata, Poir. Encye. Suppl. iv. 310 = Ar. fasciculata.
multicaulis, Crantz, Inst. ii. 406 = Ar. multicaulis.
multinervis, Boiss. Fl. Orient. i. 683.—As. Min.
Munbyi, Boiss. Diagn. Ser. II. i. 85 = Ar. Munbyi.
muscorum, Adams, ex Ledeb. Fl. Ross. i. 349 = Ar. verna.
mutabilis, Lap. ex Nym. Consp. 118 = Ar. rostrata.
nemorum, Schreb. Spieil. 30 = Stellaria nemorum.
nodosa, Crantz, Inst. ii. 408 = Sagina nodosa.
occulta, Kar. & Kir. in Bull. Soe. Nat. Mose. xv. (1842) 590 = Ar. polygonoides, artica.
octandra, Schur, Enum. Pl. Transs. 882, Index = Ar. aretioides.
Orphanides, Boiss. Diagn. Ser. II. i. 62 = Ar. tenuifolia.
pallida, Dum. Fl. Belg. 109 ; et in Bull. Soc. Bot. Belg. xii. (1873) 176.—Belg.
palustris, Kellogg, in Proc. Calif. Aead. iii. (1863) 61 = Ar. palustris.
paniculata, Fenzl, Verbreit. Alsin. 37 = Ar. paniculata.
parnassica, Boiss. & Sprun. Diagn. Ser. I. i. 46.—Graecia.
patula, A. Gray, Man. Bot. U. St. ed. II. 58 = Ar. patula.
pauciflora, Kit. ex Nym. Consp. 119 = Ar. verna.
pentandra, Crantz, Inst. ii. 408 = Spergularia pentandra.
peploides, Crantz, l. c. 406 = Ar. peploides.
Pestalozzae, Boiss. Diagn. Ser. I. viii. 99.—As. Min.
petraea, Jord. Obs. Pl. Crit. vii. (1849) 14 = Ar. rostrata.
pharnaceoides, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. filiformis.
picta, Fenzl, l. c. = Ar. filiformis.
pilifera, Turcz. in Bull. Soc. Nat. Mose. xv. (1842) 588 = Spergula larieina.
pinifolia, Fenzl, Verbreit. Alsin. in tab. ad p. 46 = Ar. pinifolia.
Pitcheri, Wood, Cl.-Book, 260 = Ar. Pitcheri.
polycarpa, Crantz, Inst. ii. 405 = Polycarpon tetraphyllum.
Preslii, Reuss, Květ. Slov. (1853) 75 = Ar. graminifolia.
procumbens, Crantz, Inst. ii. 404 = Sagina proeumbens.
procumbens, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. geniculata.
propinqua, Lange, Fl. Dan. t. 2903 = Ar. verna.
prostrata, Forsk. Fl. Aegypt. Arab. 207 = Polycarpea prostrata.
pubescens, Presl, ex. Nym. Consp. 117 = Ar. tenuifolia.
pulvinaris, Boiss. Diagn. Ser. I. i. 46.—Lydia.
Pumilio, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. arctica.
purpurascens, Bubani, in Nuov. Ann. Sc. Nat. Bolog. ix. (1843) 94.—Italia ?
purpurea, Heynh. Nom. i. 38 = Spergularia purpurea.
radicans, Guss. Fl. Sic. Syn. i. 502 = Spergularia rubra.
ramosissima, Fenzl, ex Schur, Enum. Pl. Transs. 113 = Ar. fasciculata.
recurva, Wahlenb. Fl. Helv. 87 = Ar. recurva.
repens, Vell. Fl. Flum. 127 ; iii. t. 105 = Stellaria media.

ALSINE :—
rimarum, Boiss. & Bal. in Boiss. Fl. Orient. i. 678.—Cilicia.
Rosani, Guss. Fl. Sic. Syn. i. 498 = Arenaria Rosani.
rosea, Presl, ex Nym. Consp. 119 = Ar. procumbens.
Rossii, Fenzl, Verbreit. Alsin. tab. ad p. 18 = Ar. stricta.
rostrata, Fenzl, l. c. 46 = Ar. rostrata.
rotundifolia, Stokes, Bot. Mat. Med. ii. 537 = Drymaria cordata.
Roylei, Fenzl, ex Hook. f. Fl. Brit. Ind. i. 236 = Ar. foliosa.
rubella, Schrenk, ex Ledeb. Fl. Ross. i. 349 = Ar. verna.
rubella, Wahlenb. Fl. Lapp. 128. t. 6 = Ar. verna.
rubra, Webb & Berth. Phyt. Canar. i. 148 = Spergularia canariensis.
rubra, C. A. Mey. Verz. Pfl. Cauc. 217 = Spergularia salsuginea.
rubra, Crantz, Inst. ii. 407 = Spergularia rubra.
rupestris, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. rupestris.
Sagina, Crantz, Inst. ii. 405 = Bartonia tenella.
saginoides, Crantz, l. c. 408 = Sagina Spergella.
saxatilis, Crantz, l. c. 407 = Ar. saxatilis.
saxatilis, Wahlenb. Fl. Helv. 87 = Ar. verna.
Saxifraga, Boiss. Diagn. Ser. I. i. 47 = Ar. Saxifraga.
scabra, Stokes, Bot. Mat. Med. ii. 538 = Stellaria Holostea.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 47.—Abyss.
sclerantha, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 400.—Caucas.
scleranthoides, Boiss. Diagn. Ser. II. i. 88 = Ar. setacea.
sedoides, Froel. in Koch, Syn. Fl. Germ. ed. I. 114 = Ar. sedoides.
segetalis, Linn. Sp. Pl. 272 = Spergularia segetalis.
Seguieri, Turra, ex Sacc. in Atti Ist. Venet. Ser. III. ix. (1863–64) 486.—Italia.
serpyllifolia, Crantz, Inst. ii. 406 = Ar. serpyllifolia.
setacea, Mert. & Koch, in Roehl. Deutsch. Fl. iii. 286 = Ar. setacea.
sinaica, Boiss. Diagn. Ser. I. viii. 100 = Ar. picta.
Smithii, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. globulosa.
spinuliflora, Fenzl, l. c. = Habrosia spinuliflora.
squarrosa, Fenzl, ex A. Gray, Man. Bot. U. St. ed. II. 57 = Ar. squarrosa.
striata, Crantz, Inst. ii. 407 = Ar. striata.
striata, Gren. in Mém. Soc. Doubs, (1841) 33. t. 1. f. 1 = Ar. laricifolia.
stricta, Mert. & Koch, in Roehl. Deutsch. Fl. iii. 278 = Ar. stricta.
stricta, Wahlenb. Fl. Lapp. 127 = Ar. lapponica.
subtilis, Fenzl, ex Boiss. Fl. Orient. i. 687 = Ar. tenuifolia.
subulifolia, Guss. Fl. Sic. Syn. i. 503 = Ar. mucronata.
succulenta, Delile, Fl. Egypt. 211. t. 24 = Polycarpon succulentum.
taurica, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. 319 = Ar. Villarsii.
Tchihatchewii, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 247.—As. Min.
tenella, Fuss, Fl. Transs. 112 = Ar. austriaca.
tenuifolia, Crantz, Inst. ii. 407 = Ar. tenuifolia.
tenuissima, Pomel, Nouv. Mat. Fl. Atl. 327.—Afr. bor.
tetraquetra, Crantz, Inst. ii. 406 = Ar. tetraquetra.
thymifolia, Fenzl, Verbreit. Alsin. in tab. ad p. 57 = Ar. thymifolia.
trichocalycina, Heldr. & Sart. Diagn. Ser. II. i. 87.—Ital.; Graecia.
trifolia, Baring-Gould, ex Bab. in Journ. Linn. Soc. xi. (1871) 298 = Ar. norvegica.
trinervia, Crantz, Inst. ii. 406 = Ar. trinervia.
uliginosa, Vill. Prosp. Dauph. 48 = Stellaria aquatica.
umbellata, Lam. Fl. Fr. iii. 45 = Holosteum umbellatum.
umbellifera, Boiss. Diagn. Ser. II. v. 61.—Cilicia.
unilateralis, Moench, Meth. 228 = Spergularia segetalis.
uralensis, Clerc, in Bull. Soc. Oural. iv. 105.—Ural.
velutina, Boiss. & Orphan. Diagn. Ser. II. vi. 636 = Ar. glomerata.
verna, Wahlenb. Fl. Lapp. 129; Bartl. Beitr. ii. 63 = Ar. verna.
verticillata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 6.—Java.
Villarsii, Fenzl, in Ledeb. Fl. Ross. 350 = Ar. juniperina.

ALSINE :—
Villarsii, Mert. & Koch, in Roehl. Deutsch. Fl. iii. 282 = Arenaria Villarsii.
Villarsiana, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = praec.
villosa, Heynh. Nom. i. 39 = Ar. villosa.
viscidula, Thouin, ex Nym. Consp. 117 = Ar. tenuifolia.
viscosa, Schreb. Spicil. 30 = Ar. tenuifolia.
vulgaris, Moench, Meth. 228 = Stellaria media.

ALSINELLA, Dill. ex Linn. Gen. ed. I. 118 (1737) = **Sagina**, Linn. (Caryophyll.).

ALSINELLA, Hornem. Nom. 33 (1827) = **Spergularia**, J. & C. Presl (Caryophyll.).
media, Hornem. l. c. = S. marina.

ALSINELLA, [Sw.] Summa Veg. Scand. 17 (1814) pro parte; Benth. in Wall. Cat. n. 630 (1832) = **Stellaria**, Linn. (Caryophyll.).
biflora, Sw. l. c. = S. biflora.
Wallichiana, Benth. l. c. = S. media.

ALSINELLA, [Sw.] Summa Veg. Scand. 17 (1814) pro parte; S. F. Gray, Nat. Arr. Brit. Pl. ii. 655 (1821) = **Arenaria**, Linn. (Caryophyll.).
ciliata, S. F. Gray, l. c. = Ar. ciliata.
rubella, Sw. Summa Veg. Scand. 17 = Ar. verna.
stricta, Sw. l. c. = Ar. stricta.
trinervis, S. F. Gray, Nat. Arr. ii. 655 = Ar. trinervia.
viscosa, Hartm. Fl. Dan. t. 1754 = Ar. tenuifolia.
serpyllifolia, S. F. Gray, Nat. Arr. ii. 665 = Ar. serpyllifolia.

ALSINELLA, Moench, Meth. 222 (1794) = **Cerastium**, Linn. (Caryophyll.).
erecta, Moench, l. c. = C. quaternellum.

ALSINIDENDRON, H. Mann, in Proc. Bost. Soc. Nat. Hist. x. (1866) 311. *CARYOPHYLLEAE*, Benth. & Hook. f. i. 978.
trinerve, *H. Mann*, *l. c.* 312.—Ins. Sandvic.

ALSOBIA, Hanst. in Linnaea, xxvi. (1853) 207 = **Episcia**, Mart. (Gesnerac.).
antirrhina, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 114.—Am. centr.
punctata, Hanst. in Linnaea, xxvi. (1853) 207 = E. punctata.

ALSOCYDIA, Mart. ex DC. Prod. ix. 149, &c. (1845) = **Bignonia**, Linn. &c.
acutiflora, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Lundia nitidula.
angustidens, Mart. ex DC. l. c. ix. 179 = Cuspidaria angustidens.
cordata, Mart. ex DC. l. c. 163 = Bign. cordifolia.
craterophora, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Bign. craterophora.
dichotoma, Mart. ex DC. l. c. ix. 150 = Bign. corymbifera.
erubescens. Mart. ex DC. l. c. 157 = Bign. erubescens.
firmula, Mart. ex DC. l. c. 146 = Bign. brachypoda.
glandulosa, Mart. ex DC. l. c. 149, 164 = Bign. glandulosa, lanceolata.
glutinosa, Mart. ex DC. l. c. 162 = Bign. glutinosa.
hebantha, Mart. ex Buck, Ind. DC. Prod. iii. 14 = Lundia hebantha.
hirsuta, Mart. ex DC. l. c. ix. 158 = Bign. trichoclada.
lasianthera, Mart. ex DC. l. c. 180 = Lundia cordata.
longidens, Mart. ex DC. l. c. 190 = Anemopaegma longidens.
nitidula, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Lundia nitidula.
noterophila, Mart. ex Buek, l. c. = Bign. noterophila.
tetragonocaulos, Mart. ex DC. l. c. ix. 169 = Bign. tetragonocaulos.
triplinervia, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Bign. triplinervia.
variabilis, Mart. ex DC. l. c. ix. 149 = Bign. varians.
villosa, Mart. ex Buek, Ind. DC. Prod. iii. 14 = Bign. velutina.
virginalis, Mart. ex Buek, l. c. = Lundia virginalis.

ALSODEIA, Thou. Hist. Vég. Afr. ii. 55. t. 17, 18 (1807). *VIOLARIEAE*, Benth. & Hook. f. i. 118.
CERANTHERA, Beauv. Fl. Owar. et Ben. ii. 10 (1807).

ALSODEIA :—
CERATANTHERA, Beauv. Fl. Owar. et Ben. ii. 10, in nota (1807).
CONOHORIA, Aubl. Pl. Gui. 239. t. 95 (1775).
CONORIA, Juss. Gen. 287 (1789).
CUSPA, Humb. Relat. i. 366 (1814) ex H. B. & K. Nov. Gen. et Sp. vii. 242.
DIORYKTANDRA, Hassk. Retzia, 125 (1855).
GONOHORIA, G. Don, Gen. Syst. i. 340 (1831).
JUERGENSIA, Spreng. Syst. iii. 50 (1826).
MEDUSA, Lour. Fl. Cochinch. 406 (1790).
MEDUSULA, Pers. Syn. ii. 215 (1807).
PASSALIA, Soland. ex R. Br. in Tuckey, Congo, 440 (1818).
PASSOURA, Aubl. Pl. Gui. [Suppl.] 21. t. 380 (1775).
PENTALOBA, Lour. Fl. Cochinch. 154 (1790).
PHYSIPHORA, Soland. ex R. Br. in Tuckey, Congo, 441 (1818).
PROSTHESIA, Blume, Bijdr. 866 (1826).
RIANA, Aubl. Pl. Gui. 237. t. 94 (1775).
RINOREA, Aubl. l. c. 235. t. 93 (1775).
VARECA, Roxb. Fl. Ind. i. 647 (1832).
albicaulis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 557.—N. Granat.
andina, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 366.—N. Granat.
angustifolia, *Thou. Hist. Vég. Afr.* ii. 57. t. 18. f. 2.—Madag.
arborea, *Thou. l. c.* 57. t. 17. f. 1.—Madag.
ardisiaeflora, *Welw. in Oliver, Fl. Trop. Afr.* i. 108.—Afr. trop.
Aucuparia, *Welw. l. c.* 109.—Afr. trop.
auriculata, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 319.—Madag.
bahiensis, *Moric. Pl. Nouv. Am.* 68. t. 46.—Bras.
bengalensis, *Wall. in Trans. Med. & Phys. Soc. Calc.* vii. (1846) 224.—Ind. or.
brachypetala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 558.—Afr. trop.
brevipes, *Benth. in Hook. Lond. Journ. Bot.* iv. (1842) 106.—Guiana.
Browni, Korth. in Neder. Kruid. Arch. i. (1848) 361 = zeylanica.
caledonica, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 558.—N. Caled.
calophylla, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 311.—Madag.
calycina, *Tul. l. c.* 313.—Madag.
castaneaefolia, *Spreng. Syst.* iv. *Cur. Post.* 99.—Bras.
castanioides, *Welw. in Oliver, Fl. Trop. Afr.* i. 110.—Afr. trop.
caudata, *Oliver, l. c.* 107.—Afr. trop.
cauliflora, *Oliver, l. c.* 111.—Afr. trop.
ceylanica, Auct. vide zeylanica.
chrysodasys, *Miq. Fl. Ind. Bat. Suppl.* i. 390.—Sumatra.
corylifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 559.—Java.
Cuspa, *Spreng. Syst.* iv. *Cur. Post.* 99.—Cumana.
cymulosa, *Miq. Fl. Ind. Bat. Suppl.* i. 116.—Java.
cymulosa, *Welw. in Oliver, Fl. Trop. Afr.* i. 108.—Afr. trop.
dasycaula, Miq. Fl. Ind. Bat. Suppl. i. 391 = mollis.
dasyprysis, *Miq. l. c.*—Sumatra.
decora, *Trim. in Journ. Bot.* xxiii. (1885) 203.—Zeylan.
deflexa, *Benth. Bot. Voy. Sulph.* 67.—Peruv.
dentata, *Beauv. Fl. Owar. et Ben.* ii. 11. t. 65.—Afr. trop.
disticha, *Zoll. in Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 216.—Malaya.
echinocarpa, *Korth. in Neder. Kruid. Arch.* i. (1848) 360.—Sumatra.
elliptica, *Oliver, Fl. Trop. Afr.* i. 108.—Afr. trop.
falcata, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. i. 386.—Bras.
fasciculata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 559.—Ins. Philipp.
flavescens, *Spreng. Syst.* i. 806.—Guiana.
floribunda, Moric. Pl. Nouv. Am. 70. t. 47 = castaneaefolia.
glabra, *Burgersdyke, in Miq. Pl. Jungh.* 122.—Sumatra.
Gossypium, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 367.—N. Granat.
grandis, *Miq. Fl. Ind. Bat. Suppl.* i. 391.—Sumatra.
Griffithii, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 187.—Burma.

ALSODEIA:—
guianensis, *Eichl. in Mart. Fl. Bras.* xiii. 1. 387.—Bras.
guineensis, *Spreng. Syst.* i. 806.—Afr. trop.
haplobotrys, *Hassk. in Versl. Kon. Akad. Wetensch.* iv. (1855) 137.—Java.
Horneri, *Korth. in Neder. Kruid. Arch.* i. (1848) 360.—Borneo.
Horsfieldii, *Miq. Fl. Ind. Bat.* i. ii. 117.—Java.
Horsfieldii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 559.—Java.
ilicifolia, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 12.—Afr. trop.
integrifolia, *Steud. Nom.* ed. II. i. 64.—Bras.
javanica, *Miq. Fl. Ind. Bat.* i. ii. 117.—Sumatra.
lanceolata, *Oudem. in Arch. Néerl.* ii. (1867) 196; *et in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 68.—Ind. or.
latifolia, *Thou. Hist. Vég. Afr.* ii. 57. t. 18. f. 3.—Madag.
laxiflora, Benth. in Hook. Lond. Journ. Bot. iv. (1842) 107 = guianensis.
leucoclada, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 312.—Madag.
Lindeniana, Tul. l. c. Sér. III. vii. (1847) 364 = guianensis.
longiflora, *Oudem. in Arch. Néerl.* ii. (1867) 195; *et in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. (1867) 68.—Guiana.
longipes, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 309.—Madag.
longiracemosa, *Kurz, in Journ. As. Soc. Beng.* xxxix (1870) ii. 63.—Ind. or.
macrocarpa, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 385.—Bras.
macrophylla, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 468. t. 19.—Ins. Timor.
maculata, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 316.—Madag.
Maingayi, *Hook. f. & Thoms. ex Fl. Brit. Ind.* i. 188.—Malacca.
marginata, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 127.—Nov. Granat.
Martinii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 557.—Guin.
Maximiliana, *Eichl. in Mart. Fl. Bras.* xiii. 1. 381.—Bras.
megapotamica, *Spreng. Syst.* iv. *Cur. Post.* 99.—Bras.
mollis, *Hook. f. & Thoms. ex Fl. Brit. Ind.* i. 188.—Tenasserim.
monticola, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 315.—Madag.
mutica, *Tul. l. c.* 318.—Madag.
nigricans, Boiv. ex Tul. l. c. 305 = pubescens.
obtusa, *Korth. in Neder. Kruid. Arch.* i. (1848) 359.—Borneo.
owariensis, *Spreng. Syst.* i. 806.—Afr. trop.
pallida, *Klotzsch, in Schomb. Fauna et Fl. Guy.* 1165.—Guiana.
paniculata, *Mart. Nov. Gen. et Sp.* i. 30.—Bras.
paradoxa, *Blume, ex Oudem. in Arch. Neerl.* ii. (1867) 204; *et in Miq. Ann. Mus. Lugd. Bat. Bot.* iii. 71.—Java.
pauciflora, *Thou. Hist. Vég. Afr.* ii. 57. t. 17.—Madag.
Perrini, *Spreng. Syst.* i. 807.—Am. austr.
Pervillii, *Oudem. in Arch. Néerl.* ii. (1867) 206; *et in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 72.—Madag.
physiphora, *Mart. Nov. Gen. et Sp.* i. 28.—Bras.
Piparea, Spreng. Syst. i. 807 = Casearia dentata.
prunifolia, *Spreng. l. c.*—Guiana.
pubescens, *Thou. Hist. Vég. Afr.* ii. 57. t. 18. f. 3.—Madag.
pubiflora, *Benth. in Hook. Journ. Bot.* iv. (1842) 106.—Guiana.
pugionifera, *Oudem. in Arch. Néerl.* ii. (1867) 205; *et in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 71.—Madag.
racemosa, Hook. f. & Thoms. ex Fl. Brit. Ind. i. 187 = longiracemosa.
racemosa, *Mart. & Zucc. Nov. Gen.* i. 29. t. 20.—Bras.
Regnellii, Walp. Ann. ii. 67 = Ionidium atropurpureum.
Rinorea, Spreng. Syst. i. 807 = paniculata.
Roxburghii, *Hook. f. & Thoms. ex Fl. Brit. Ind.* i. 186.—Ind. or.
rubra, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 315.—Madag.

ALSODEIA:—
rugosa, *Miq. Fl. Ind. Bat. Suppl.* i. 390.—Sumatra.
sclerocarpa, *Burgersdyke, in Miq. Pl. Jungh.* 122.—Sumatra.
semigyrata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 559.—Java.
sessilis, *Spreng. Syst.* i. 806.—Cochinch.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 402.—Ins. Socotra.
spinosa, *Boivin, ex Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 307.—Madag.
Sprucei, *Eichl. in Mart. Fl. Bras.* xiii. 1. 385.—Bras.
squamosa, *Boivin, ex Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 307.—Madag.
Storckii, *Seem. in Journ. Bot.* ii. (1864) 75.—Ins. Fiji.
subintegrifolia, *Beauv. Fl. Owar. et Ben.* ii. 11. t 66.—Afr. trop.
sylvatica, *Seem. Bot. Voy. Herald*, 79. t. 40.—Panama.
trigonocarpa, *Zoll. & Reichb. in Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 216.—Malaya.
ulmifolia, *Spreng. Syst.* i. 807.—N. Granat.
verticillata, *Boivin, ex Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 317.—Madag.
virgata, Thw. ex Hook. f. Fl. Brit. Ind. i. 189 = Scyphyllandra virgata.
viridiflora, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 310.—Madag.
Wallichiana, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 187.—Penang.
Welwitschii, *Oliver, Fl. Trop. Afr.* i. 110.—Afr. trop.
zeylanica, *Thw. Enum. Pl. Zeyl.* 21.—Ind. or.

ALSODEIOPSIS, Oliver, in Benth. & Hook. f. Gen. i. 996 (1867). *OLACINEAE*, Benth. & Hook. l. c.
Manni, *Oliver, in Journ. Linn. Soc.* x. (1869) 43.—Afr. trop.

ALSOLINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349 = **Linum**, Tourn.
maritimum, Fourr. l. c. = L. maritimum.

ALSOMITRA, M. Roem. Syn. Pepon. 117 partim (1846). *CUCURBITACEAE*, Benth. & Hook. f. i. 840.
angulata, M. Roem. l. c. 117 = Zanonia indica.
Beccariana, *Cogn. in DC. Monog. Phan.* iii. 932.—Ins. Kei.
brasiliensis, *Cogn. in Mart. Fl. Bras.* vi. iv. 115.—Bras.
capricornica, *F. Muell. Fragm.* vii. 61.—Austral.
cissoides, M. Roem. Syn. Pepon. 118 = Gynostemma pedata.
clavigera, *M. Roem. l. c.; Benth. & Hook. f. Gen.* i. 840.—Ind. or.
heterosperma, M. Roem. l. c. = Gomphogyne heterosperma.
Hookeri, *F. Muell. Fragm.* vi. 188.—Austral.
laxa, M. Roem. Syn. Pepon. 118 = Gynostemma laxa.
macrocarpa, M. Roem. l. c. 117 = Zanonia macrocarpa.
pedatifolia, *Cogn. in Mart. Fl. Bras.* vi. iv. 116.—Bras.
sarcophylla, *M. Roem. Syn. Pepon.* 118.—Burma; Siam.
Schefferiana, *Cogn. in DC. Monog. Phan.* iii. 932.—Ins. Celebes.
Stephensiana, *Cogn. l. c.* 934.—Ins. Barnardi.
timorana, *M. Roem. Syn. Pepon.* 117.—Ins. Timor.

ALSTONIA, Mutis, ex Linn. f. Suppl. 39 (1781) = **Symplocos**, Linn. (Styrac.).
ciliata, Benth. Pl. Hartw. 48 = S. ciliata.
theaeiformis, Linn. f. l. c. 264 = S. Alstonia.

ALSTONIA, Scop. Introd. 198 (1777) = **Landolphia**, R. Br. (Apocyn.).

ALSTONIA, R. Br. in Mem. Wern. Soc. i. (1809) 75.
APOCYNACEAE, Benth. & Hook. f. ii. 705.
BLABEROPUS, A. DC. Prod. viii. 410 (1844).
PALA, [Rheede] Juss. in Ann. Mus. Par. xv. (1825) 346.
acuminata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 140.—Ins. Ceram.
angustifolia, *Wall. Cat.* n. 1650.—Malaya.
angustiloba, *Miq. Fl. Ind. Bat.* ii. 438.—Malaya.
Batino, Blanco, Fl. Filip. ed. II. Suppl. 589 = macrophylla.
calophylla, *Miq. Fl. Ind. Bat.* ii. 439.—Java.

ALSTONIA:—
constricta, *F. Muell. Fragm.* i. 57.—Austral.
costata, *R. Br. in Mem. Wern. Soc.* i. (1809) 75.—Ins. Tahiti.
costata, Wall. Cat. n. 1649 = macrophylla.
costulata, Miq. Fl. Ind. Bat. Suppl. 556 = Dyera costulata.
cuneata, Wall. Cat. n. 1645 = spathulata.
Deplanchei, *Heurck & Muell. Arg. in Flora*, liii. (1870) 171.—N. Caled.
edulis, *G. Benn. in Journ. Bot.* v. (1867) 150.—N. Caled.
eximia, *Miq. Fl. Ind. Bat. Suppl.* 555.—Ins. Banca.
grandifolia, *Miq. l. c.*—Sumatra.
Kurzii, *Hook. f. Fl. Brit. Ind.* iii. 643.—Ins. Andaman.
lanceolata, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 199.—N. Caled.
Legouixiae, *Heurck & Muell. Arg. in Flora*, liii. (1870) 170.—N. Caled.
Lenormandii, *Heurck & Muell. Arg. l. c.* 172.—N. Caled.
linearis, *Benth. Fl. Austral.* iv. 314.—Austral.
longissima, *F. Muell. Papuan Plants*, 91.—N. Guin.
lucida, D. Don, Prod. Fl. Nep. 131 = Trachelospermum fragrans.
macrophylla, *Wall. Cat.* n. 1648 (*Fl. Brit. Ind.* iii. 643).—Malaya.
mollis, *Benth. Fl. Austral.* iv. 315.—Austral.
neriifolia, *D. Don, Prod. Fl. Nep.* 131.—Reg. Himal.
oleandraefolia, Lodd. ex Loud. Hort. Brit. 67 = scholaris.
ophioxyloides, *F. Muell. Fragm.* i. 57.—Austral.
plumosa, *Labill. Sert. Austr. Caled.* 28. t. 32.—N. Caled.
polyphylla, *Miq. Fl. Ind. Bot. Suppl.* 556.—Ins. Banca.
quaternata, *Heurck & Muell. Arg. in Flora*, liii. (1870) 170.—N. Caled.
Roeperi, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 201.—N. Caled.
scholaris, *R. Br. in Mem. Wern. Soc.* i. (1809) 75.—Geront. trop.
sericea, *Blume, Bijdr.* 1038.—Malaya.
spatulata, *Blume, l. c.* 1037.—Malaya.
spectabilis, *R. Br. in Mem. Wern. Soc.* i. (1809) 75.—Ins. Timor.
spectabilis, Kurz, For. Fl. Burma, ii. 183 = Kurzii.
subsessilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 140.—Amboina.
venenata, *R. Br. in Mem. Wern. Soc.* i. (1809) 77.—Ind. or.
verticillosa, *F. Muell. Fragm.* vi. 116.—Austral.
Vieillardii, *Heurck & Muell. Arg. in Flora*, liii. (1870) 171.—N. Caled.
villosa, *Blume, Bijdr.* 1038.—Java; Austral.
villosa, Seem. Fl. Vit. 161 = vitiensis.
vitiensis, *Seem. l. c.* 430.—Ins. Fiji.

ALSTROEMERIA, Linn. Diss. Alstroem. 8 (1762). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 736.
LIGTU, Adans. Fam. p. 20 (1763).
LILAVIA, Rafin. Fl. Tellur. iv. 35 (1836).
PRIOPETALON, Rafin. Fl. Tellur. iv. 34 (1836).
acutifolia, Link & Otto, Ic. Pl. Rar. Hort. Bot. Berol. 57. t. 29 = Bomarea acutifolia.
affinis, Mart. & Gal. in Bull. Acad. Brux. x. (1843) reimpr. 6 = Bomarea edulis.
albiflora, *Presl, Rel. Haenk.* i. 121. t. 22.—Chili.
amoena, Salisb. Prod. 248 = A. Pelegrina.
anceps, *Ruiz & Pav. Fl. Per.* iii. 61.—Peruv.
andina, Phil. in Linnaea, xxix. (1857–58) 69 = spathulata.
angustifolia, Herb. Amar. 96 = A. Ligtu.
araucana, *Phil. in Anales Univ. Chil.* (1873) 547.—Chili.
argento-vittata, *Lem. Illustr. Hortic.* iv. (1857), *Misc* 88; vi. (1858) t. 192.—Bras.
aurantiaca, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 205.—Chili.
aurea, *R. Grah. in Edinb. Phil. Journ.* (1838) 181.—Chili.
aurea, Meyen, Reise, i. 311 = aurantiaca.
Banksiana, M. Roem. Syn. Ensat. 259 = pulchella.
Barklayana, *Hort. ex Loud. Hort. Brit. Suppl.* ii. 608.—Hab.?
Berteroiana, *Penny, ex Loud. l. c.* = Chili.
bicolor, Hook. Exot. Fl. i. t. 95 = A. Ligtu.
bracteata, *Ruiz & Pav. Fl. Per.* iii. 61.—Peruv.
brasiliensis, *Spreng. Syst.* ii. 81.—Bras.
Bredemeyeriana, Willd. ex Kunth, Enum. Pl. ii. 808 = Bomarea multiflora.

ALSTROEMERIA :—

Burchellii, *Baker, in Journ. Bot.* xv. (1877) 262.—Bras.
Caldasiana, Hemsl. Biol. Centr. Am. Bot. iii. 336 = seq.
Caldasii, H. B. & K. Nov. Gen. et Sp. i. 283 = Bomarea Caldasiana.
caryophyllacea, *Jacq. Schoenbr.* vi. 33. t. 465.—Bras.
chilensis, *J. Cree, ex Loud. Hort. Brit. Suppl.* ii. 608, nomen.—Chili.
chilensis, *Lem. in Fl. des Serres*, Sér. I. i. (1845) 98.—Chili.
chiloënsis, *Phil. in Linnaea*, xxix. (1857–58) 71.—Ins. Chiloe.
chorillensis, *Herb. in Bot. Reg.* (1843) *Misc.* 64.—Chili.
ciliata, *Poepp. Fragm. Syn.* 6.—Chili.
citrina, Phil. in Linnaea, xxxiii. (1864–65) 264 = versicolor.
coccinea, Ruiz & Pav. Fl. Per. iii. 62 = Bomarea coccinea.
concolor, Steud. in Lechl. Berb. Am. Austr. 53 = aurantiaca.
cordifolia, *Ruiz & Pav. Fl. Per.* iii. 64.—Peruv.
crispata, *Phil. in Linnaea*, xxix. (1857–58) 70.—Chili.
crocea, *Phil. l. c.* xxxii. (1864–65) 263.—Chili.
crocea, Ruiz & Pav. Fl. Per. iii. 61 = Bomarea crocea.
Cummingiana, Herb. Amar. 96, 396 = versicolor.
Cunea, *Vell. Fl. Flum.* iii. t. 121; *Seub. in Mart. Fl. Bras.* iii. 172.—Bras.
Cunha, Vell. l. c. 131 = praec.
Curtisiana, E. Mey. ex Presl, Rel. Haenk. ii. 122 = A. Ligtu.
Curtisiana, Schult. f. Syst. vii. 735 = caryophyllacea.
declinata, Poepp. & Endl. Nov. Gen. ii. 44. t. 160 = Bomarea dispar.
densiflora, Herb. in Bot. Mag. t. 5531 = Bomarea tomentosa.
denticulata, *Ruiz & Pav. Fl. Per.* iii. 62.—Peruv.
Despuenta, *Pasq. Cat. Orto Bot. Nap.* 7.—Hab. ?
Diazi, *Phil. in Linnaea*, xxxiii. (1864) 261.—Chili.
Didierana, *Jaub. in Bull. Soc. Bot. Fr.* viii. (1861) 118.—Peruv.
distichifolia, *Ruiz & Pav. Fl. Per.* iii. 60.—Peruv.
distichophylla, Spreng. Syst. ii. 80 = Bomarea distichophylla.
dulcis, Hook. Bot. Misc. ii. (1831) 237. t. 95 = Bomarea glaucescens.
edulis, Andr. ex Poir. Encyc. Suppl. iv. 348 = Bomarea edulis.
Errembaultii ×, *Hort. ex Lem. in Fl. des Serres*, Sér. I. iii. (1847) 262.
exserens, Meyen, Reise, i. 34 = A. Ligtu.
Feuillaeana, E. Mey. ex Presl, Rel. Haenk. ii. = A. Ligtu.
filipendula, Seub. in Mart. Fl. Bras. iii. 1. 180 = caryophyllacea.
fimbriata, Ruiz & Pav. Fl. Per. iii. 63 = Bomarea fimbriata.
flava, *Phil. in Linnaea*, xxxiii. (1864–65) 263.—Chili.
floribunda, *H. B. & K. Nov. Gen. et Sp.* i. 283.—Ecuador.
Flos-Martini, [Ker-Gawl. in] Bot. Reg. t. 731 = A. Ligtu.
fluminensis, *M. Roem. Syn. Ensat.* 260.—Bras.
foliosa, *Mart. in Schult. f. Syst.* vii. 740.—Bras.
formosa, Pers. Syn. i. 381 = seq.
formosissima, Ruiz & Pav. Fl. Per. iii. 64 = Bomarea formosissima.
Gardneri, *Baker, in Journ. Bot.* xv. (1877) 261.—Bras.
Gayana, *Phil. in Linnaea*, xxix. (1857–58) 71.—Chili.
glaucescens, H. B. & K. Nov. Gen. et Sp. i. 282 = Bomarea glaucescens.
gloriosa, Cham. & Schlecht. in Linnaea, vi. (1831) 51 = Bomarea edulis.
grandifolia, *H. B. & K. Nov. Gen. et Sp.* i. 285.—Bras.
haemantha, *Ruiz & Pav. Fl. Per.* iii. 60.—Chili.
Herbertiana, M. Roem. Syn. Ensat. 250 = revoluta.
hirsuta, H. B. & K. Nov. Gen. et Sp. i. 285 = Bomarea hirsuta.
hirtella, H. B. & K. l. c. 284 = Bomarea edulis.
hirtella, *Phil. in Linnaea*, xxix. (1857–58) 70.—Chili.
Hookeri, Lodd. Bot. Cab. t. 1272 = A. Ligtu.
Hookeriana, *C. Gay, Fl. Chil.* v. 90.—Chili.
Hookeriana, Schult. f. Syst. vii. 733 = A. Ligtu.
inaequalis, *Phil. in Linnaea*, xxix. (1857–58) 67.—Chili.
inconspicua, *Phil. Anales Univ. Chil.* (1873) 546.—Chili.
inodora, *Herb. Amar.* 90. t. 2. f. 1.—Bras.
insignis, *Année, ex Hérincq, L'Hortic. Franç.* iv. (1854) 2.—Peruv.
Isabellana, *Herb. Amar.* 88, 396. t. 6. f. 4–6.—Bras.

ALSTROEMERIA :—

Jacobi, *Hort. ex Neumann, Ann. Fl. et Pomone*, (1834) t. 29.—Hab. ?
Jacquesiana, Lem. in Fl. des Serres, Sér. I. ii. (Dec. 1846) t. 182 = Bomarea Jacquesiana.
Kingii, Phil. in Anales Univ. Chil. (1873) 548 = versicolor.
Kunziana, *Klotzsch, ex Kunth, Enum. Pl.* v. 769.—Chili.
latifolia, *Ruiz & Pav. Fl. Per.* iii. 64.—Peruv.
Ligra, Crantz, Inst. i. 467 = A. Ligtu.
Ligtu, Curt. Bot. Mag. t. 125 = caryophyllacea.
Ligtu, *Linn. Diss. Alstroem.* 10.—Chili.
lineatiflora, Lindl. Bot. Reg. (1834) t. 58 = chorillensis.
lineatiflora, *Ruiz & Pav. Fl. Per.* iii. 60 t. 289.—Peruv.
linifolia, H. B. & K. Nov. Gen. et Sp. i. 282 = Bomarea linifolia.
longistaminea, *Mart. in Schult. f. Syst.* vii. 739.—Bras.
longistyla, *Schenk, in Mart. Fl. Bras.* iii. 1. 173.—Bras.
Macraeana, *Herb. Amar.* 90.—Chili.
macrocarpa, *Ruiz & Pav. Fl. Per.* iii. 63.—Peruv.
magnifica, *Herb. in Bot. Reg.* (1843) *Misc.* 64.—Chili.
Meyeniana, *J. C. Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* 1. (1843) 440.—Chili.
miniata, Mart. & Gal. in Bull. Acad. Brux. x. (1843) repr. 6 = Bomarea edulis.
montana, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 183 = Ixiolirion montanum.
monticola, *Mart. in Schult. f. Syst.* vii. 739.—Bras.
multiflora, Linn. f. Suppl. 207 = Bomarea multiflora.
mutabilis, Kunze, ex Kunth, Enum. Pl. v. 777 = aurantiaca.
Neilliana, Herb. Amar. 94 = spathulata.
Neillii, Gill. ex Hook. in Bot. Mag. t. 3105 = praec.
nemorosa, Gardn. in Bot. Mag. t. 3958 = inodora.
nivalis, Meyen, Reise, i. 315 = versicolor.
nivalis, *Phil. in Linnaea*, xxix. (1857–58) 69.—Chili.
nubigena, *Phil. l. c.* 67.—Chili.
Ochagavii, *Anneé, ex Hérincq, L'Hortic. Franç.* iv. (1854) 2.—Chili.
oculata, Lodd. Bot. Cab. t. 1851 = Bomarea Salsilla.
odorata, *Knight & Perry, ex Loud. Hort. Brit. Suppl.* III. 486.—Hab. ?
Oreas, *J. C. Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* 1. (1843) 440.—Am. austr.
ovata, Cav. Ic. i. 54. t. 76 = Bomarea edulis.
pallens, *Phil. in Linnaea*, xxxiii. (1864–65) 265.—Chili.
pallida, R. Grah. in Edinb. N. Phil. Journ. (Jul.–Oct. 1829) 344 = A. Ligtu,
parvula, *Phil. in Linnaea*, xxxiii. (1864–65) 261.—Chili.
pauciflora, Hort. ex Lem. in Fl. des Serres, Sér. I. ii. (Dec. 1846) t. 182 = Bomarea Jacquesiana.
pauciflora, *H. B. & K. Nov. Gen. et Sp.* i. 284.—N. Granat.
paupercula, *Phil. Fl. Atac.* 51.—Chili.
Pelegrina, *Linn. Diss. Alstroem.* 10; *Amoen. Acad.* vi. 247.—Peruv.; Chili.
Pelegrina, Vell. Fl. Flum. 131; iii. t. 119 = caryophyllacea.
peregrina, Ruiz & Pav. Fl. Per. iii. 51 = A. Pelegrina.
peruviana, Van-Houtte, Fl. des Serres, xxi. (1875) 51. t. 2175 = versicolor.
plantaginea, *Mart. in Schult. f. Syst.* vii. 737.—Bras.
platyphylla, *Baker, in Journ. Bot.* xv. (1877) 261.—Bras.
Pohliana, Seub. in Mart. Fl. Bras. iii. 1. 172 = A. Cunea.
Preslana, Herb. Amar. 95. t. 1. f. 48 = A. Ligtu.
Presliana, Kunth, Enum. Pl. v. 773 = praec.
psittacina, Lehm. Cat. Hort. Hamb. (1826) ex Schult. f. Syst. vii. 739 = pulchella.
puberula, Phil. in Linnaea, xxix. (1857–58) 68 = versicolor.
pulchella, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 48 = versicolor.
pulchella, *Linn. f. Suppl.* 206.—Am. austr.
pulchella, Sims, Bot. Mag. t. 2354 = haemantha.
pulchella, Salisb. Prod. 248 = A. Ligtu.
pulchra, *Sims, Bot. Mag.* t. 2421 = A. Ligtu.
purpurea, Ruiz & Pav. Fl. Per. iii. 63 = Bomarea purpurea.

ALSTROEMERIA :—

purpurea, Willd. ex Steud. Nom. ed. II. i. 61, 62 = Bomarea Caldasiana.
pygmaea, *Herb. Amar.* 100, 397. t. 8. f. 4–13.—Peruv.
quillotensis, Herb. l. c. 97. t. 2. f. 2 = A. Pelegrina.
reclinata, *Poepp. & Endl. Gen. et Sp.* ii. 44. t. 160.—Chili.
recumbens, Herb. Amar. 97. t. 3. f. 2 = versicolor.
reflexa, *M. Roem. Syn. Ensat.* 252.—Chili.
revoluta, *Ruiz & Pav. Fl. Per.* iii. 59.—Chili.
Riedelliana, *Andot, in Rev. Hortic.* Sér. II. iv. (1845–46) 207.—Bras.
rosea, Hook. Exot. Fl. t. 181 = A. Ligtu.
rosea, *Phil. Sert. Mend. Alt.* 43.—Chili.
rosea, *Ruiz & Pav. Fl. Per.* iii. 61.—Peruv.
rosmarinifolia, Benth. Pl. Hartw. 258 = Bomarea linifolia.
rubra, *Morel, in Rev. Hortic.* Sér. IV. ii. (1853) 81. f. 5.—Bras.
Salsilla, Linn. Diss. Alstroem. 10; Amoen. Acad. vi. 247 = Bomarea Salsilla.
Salsilla, Vell. Fl. Flum. 131; iii. t. 120 = Bomarea edulis.
salsilloides, *Mart. in Schult. f. Syst.* vii. 748.—Bras.
scaberula, *Baker, in Journ. Bot.* xv. (1877) 261.—Bras.
secundifolia, Ruiz & Pav. Fl. Per. iii. 60 = Bomarea secundifolia.
Sellowiana, *Seub. in Mart. Fl. Bras.* iii. 1. 173.—Bras.
sepium, Schott, ex Seub. l. c. 168 = Bomarea edulis.
sericantha, J. C. Schau. in Nov. Act. Nat. Cur. xix. Suppl. 1. (1843) 441 = spathulata.
setacea, Ruiz & Pav. Fl. Per. iii. 62 = Bomarea tomentosa.
Simsii, Spreng. Syst. ii. 80 = haemantha.
Simsii, Sweet, Hort. Brit. ed. I. 408 = A. Ligtu.
spathulata, *Presl, Rel. Haenk.* ii. 122. t. 22. f. 2.—Peruv.
stenopetala, *Phil. in Anales Univ. Chil.* (1873) 547.—Chili.
stenopetala, *Seub. in Mart. Fl. Bras.* iii. 1. 174.—Bras.
subrosulacea, *Herb. Amar.* 397. t. 5. f. 1. 2.—Chili.
tenuifolia, Herb. l. c. 94. t. 1. f. 47 = versicolor.
tigrina, Phil. in Linnaea, xxix. (1857–58) 68 = versicolor.
tigrina, Seub. in Mart. Fl. Bras. iii. 1. 174 = brasiliensis.
tomentosa, Ruiz & Pav. Fl. Per. iii. 62 = Bomarea tomentosa.
torta, H. B. & K. Nov. Gen. et Sp. i. 283 = Bomarea torta.
tricolor, Hook. Exot. Fl. t. 65 = A. Ligtu.
triflora, Griff. Notul. iii. 240 = Ixiolirion montanum.
umbellata, Meyen, Reise, i. 356 = spathulata.
uniflora, Matth. ex Kunth, Enum. Pl. v. 782 = Bomarea glaucescens.
venustula, *Phil. in Linnaea*, xxxiii. (1864–65) 260.—Chili.
versicolor, *Ruiz & Pav. Fl. Per.* iii. 59.—Chili.
violacea, *Knight & Perry, ex Loud. Hort. Brit. Suppl.* iii. 486.—Hab. ?
violacea, *Phil. Fl. Atac.* 51.—Chili.
viridiflora, *Warm. in Kjoeb. Vidensk. Meddel.* (1872) 113.—Bras.
xanthina, Phil. in Anales Univ. Chil. (1873) 546 = versicolor.
zamioides, *Baker, in Journ. Bot.* xv. (1877) 262.—Bras.

ALTENSTEINIA, H. B. & K. Nov. Gen. et Sp. i. 332. t. 72. 73 (1815). *ORCHIDEAE*, Benth. & Hook. f. iii. 592.
Aa, Reichb. f. Xen. Orchid. i. 18 (1858).
Myrosmodes, Reichb. f. l. c. i. 19. t. 8 (1858).
argyrolepis, *Benth. & Hook. f. (in Herb. Kew.)*.—Ecuad.
calceata, *Reichb. f. l. c.* iii. 19.—Peruv.
erosa, *Reichb. f. Otia Bot. Hamb.* 51.—Peruv.
fimbriata, *H. B. & K. Nov. Gen. et Sp.* i. 333. t. 72.—Ecuad.
gymnandra, *Reichb. f. Xenia Orch.* iii. 18.—Boliv.
inaequalis, *Reichb. l. c.* 19.—Boliv.
Mandonii, *Reichb. f. l. c.*—Peruv.
marginata, *Reichb. f. l. c.* 20.—Peruv.
Matthewii, *Reichb. f. l. c.* 19.—Peruv.
nubigena, *Reichb. f. l. c.* 18 in nota.—Hab. ?

ALTENSTEINIA:—
paleacea, *Kunth, Syn.* i. 325.—Ecuad.
paludosa, *Reichb. f. Xenia Orch.* iii. 19.—Boliv.
pilifera, *H. B. & K. Nov. Gen. et Sp.* i. 333. t. 73.—Ecuad.
rostrata, *Reichb. f. Xenia Orch.* iii. 18.—Ecuad.
Sceptrum, *Reichb. f. l. c.* i. 18.—Ecuad.
virescens, *Lindl. in Ann. & Mag. Nat. Hist.* xv. (1845) 385.—N. Granat.
Weddelliana, *Reichb. f. Xenia Orch.* iii. 19.—Peruv.

ALTERNANTHERA, Forsk. Fl. Aegypt. Arab. 28 (1775). *AMARANTACEAE*, Benth. & Hook. f. iii. 39.
ALLAGANTHERA, Mart. Hort. Erlang. 69 (1814).
ILLECEBRUM, Spreng. Anleit. ii. 1. 317 (1817); Syst. i. 818 (1825).
LITHOPHILA, Sw. Prod. Veg. Ind. Occ. 14 (1788); Fl. Ind. Occ. i. 47. t. 1 (1797).
LITOPHILA, Sw. Prod. l. c.
PITYRANTHUS, Mart. in Denks. Acad. Muench. v. (1817) 179.
acaulis, *Anderss. Galap. Öarnes Veget.* 164.—Ins. Galap.
Achyrantha, *R. Br. Prod.* i. 417.—Am. trop.
Achyranthes, Forsk. Fl. Aegypt. Arab. p. lxix = sessilis.
albida, *Griseb. in Goett. Abh.* xix. (1874) (*Pl. Lorent.* 36).—Reg. Argent.
amabilis, Hort. ex Lem. Illust. Hortic. xv. (1868) t. 558 = Telanthera ficoidea.
angustifolia, Moq. in DC. Prod. xiii. II. 324 = polycephala.
angustifolia, *R. Br. Prod.* i. 417.—Austral.
argentata, *Moq. in DC. Prod.* xiii. II. 352.—Porto Rico.
aurata, *Moq. l. c.* 353.—Am. austr.
axillaris, D. Dietr. Syst. i. 866 in syn. = spinosa.
Baueri, *Moq. in DC. Prod.* xiii. II. 354.—Austral.
brachiata, Hort. ex Moq. l. c. 304 = Aerva brachiata.
canescens, *H. B. & K. Nov. Gen. et Sp.* ii. 204.—Cumana.
canescens, Moq. in DC. Prod. xiii. II. 350 = Iresine canescens.
caracasana, *H. B. & K. Nov. Gen. et Sp.* ii. 205.—Venezuela.
earibaea, *Moq. in DC. Prod.* xiii. II. 354.—Ind. occ.
celosioides, *Moq. l. c.* 360.—Ins. Philipp.
ciliata, Poepp. ex Seub. in Mart. Fl. Bras. v. I. 183 = A. Achyrantha.
decipiens, *Benth. Fl. Austral.* v. 251.—Austral.
denticulata, R. Br. Prod. i. 417 = nodiflora.
denticulata, Wall. Cat. n. 6922 = sessilis.
dubia, Hort. Paris. ex Moq. in DC. Prod. xiii. II. 361 = sessilis?
dubia, H. B. & K. Nov. Gen. et Sp. ii. 209 = Tromnsdorffia canescens.
echinata, Hort. Bonn. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 43 = armata.
echinata, Sm. in Rees, Cyclop. Suppl. n. 10 = A. Achyrantha.
erecta, Reichb. in DC. Prod. iii. 375 = Polycarpaea lineariifolia.
ficoidea, R. Br. Prod. 417; Roem. & Schult. Syst. v. 555 = Telanthera ficoidea.
ficoidea, Griseb. Fl. Brit. W. Ind. 67 = Telanthera maritima.
ficoides, Beauv. Fl. Owar. ii. 66. f. 1 = denticulata.
ficoides, Reichb. ex Moq. in DC. Prod. xiii. II. 357 = sessilis.
flavescens, H. B. & K. Nov. Gen. et Sp. ii. 207 = Telanthera flavescens.
flavescens, *Moq. in DC. Prod.* xiii. II. 350.—N. Granat.
frutescens, *R. Br. ex Spreng. Syst.* i. 819 in syn.—Am. austr.
glabra, Moq. in DC. Prod. xiii. II. 361 = denticulata?
gomphrenoides, *H. B. & K. Nov. Gen. et Sp.* ii. 208.—Peruv.
herniarioides, *Beurling, in Vet. Akad. Handl. Stockh.* (1854) 143.—Panama.
iresinoides, H. B. & K. Nov. Gen. et Sp. ii. 209 = Gomphrena iresinoides.
lanuginosa, Moq. in DC. Prod. xiii. II. 359 = Cladothrix lanuginosa.
lappulacea, Schlecht. ex Moq. l. c. 329 = Cyathula globulifera.
latifolia, Moq. l. c. 351 = Iresine latifolia.
leptophylla, *Benth. Fl. Austral.* v. 251.—Austral.
leucantha, *Moq. in DC. Prod.* xiii. II. 357.—Jamaic.

ALTERNANTHERA:—
littoralis, Beauv. ex Moq. in DC. Prod. xiii. II. 365 = maritima.
linearis, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 107.—Porto Rico.
longipes, *Benth. Fl. Austral.* v. 252.—Austral.
lupulina, *H. B. & K. Nov. Gen. et Sp.* ii. 206.—Peruv.
maritima, *D. Dietr. Syn. Pl.* i. 867.—Bras.
maritima, St. Hil. Voy. Brés. ii. 43 = Telanthera maritima.
mexicana, *Moq. in DC. Prod.* xiii. II. 352.—Mexico?
muscoides. *Benth. & Hook. f. Gen.* iii. 39.—Ind. occ.
nana, F. Bauer, ex Moq. in DC. Prod. xiii. II. 341 = Iresine Brownii.
nana, *R. Br. Prod.* i. 417.—Austral.
nigriceps, Hook. Bot. Misc. ii. (1831) 237 = Telanthera elongata.
nigripes, Steud. Nom. ed. II. i. 65. (sphalm.) = praec.
nodifera, *Griseb. in Goett. Abh.* xxiv. (1879) 36.—Reg. Argent.
nodiflora, *R. Br. Prod.* i. 417.—Geront. trop.
paniculata, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 106.—Porto Rico.
paniculata, *H. B. & K. Nov. Gen. et Sp.* ii. 208.—Cumana.
paronychioides, Hort. ex Regel, Gartenfl. (1869) 102 = Telanthera Bettzichiana.
paronychioides, *St. Hil. Voy. Brés.* ii. II. 43.—Am. austr.
parviflora, *Moq. in DC. Prod.* xiii. II. 360.—Mexic.?
philoxeroides, *Griseb. in Goett. Abh.* xxiv. (1879) 36.—Reg. Argent.
pilosa, *Moq. in DC. Prod.* xiii. II. 357.—Bras.
polycephala, *Benth. Fl. Austral.* v. 251.—Austral.
polygonoides, R. Br. Prod. 417 = Telanthera polygonoides.
praelonga, St. Hil. Voy. Brés. ii. II. 409 = Telanthera praelonga.
procumbens, *Roem. & Schult. Syst.* v. 564.—Bras.
prostrata, D. Don, Prod. Fl. Nep. 76 = sessilis.
puberula, D. Dietr. Syn. Pl. i. 866 = Telanthera puberula.
pubescens, Hort. Prag. ex Moq. in DC. Prod. xiii. II. 303 = Aerva lanata.
pulchella, *H. B. & K. Nov. Gen. et Sp.* ii. 205.—Bras.
pulverulenta, *Moq. in DC. Prod.* xiii. II. 351.—Peruv.
pungens, *H. B. & K. Nov. Gen. et Sp.* ii. 206.—Bras.
purpurea, Hort. ex Pynaert, in Rev. Hortic. Belge, v. (1879) 25 = Telanthera versicolor.
radicata, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 262.—Ins. Galapag.
repens, Steud. Nom. ed. I. 34 = sessilis.
repens, Steud. l. c. ed. II. i. 65 = A. Achyrantha.
Richardi, Moq. in DC. Prod. xiii. II. 353 = Iresine interrupta.
rufa, D. Dietr. Syn. Pl. i. 866 = Telanthera rufa.
sericea, *H. B. & K. Nov. Gen. et Sp.* ii. 207.—Ecuad.
sessilis, Lem. Illustr. Hortic. (1865) t. 447 = Telanthera amoena.
sessilis, *R. Br. Prod.* i. 417.—Reg. trop.
sibirica, *Steud. Nom.* ed. II. i. 65.—Sibir.
spathulata, Lem. Illustr. Hortic. xii. (1865) t. 445 = Telanthera ficoidea.
spinosa, *Roem. & Schult. Syst.* v. 555.—Am. trop.
strigosa, *Hassk. in Hoev. & De Vriese, Tijdschr.* v. (1838-39) 259.—Malaya?
subscaposa, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 189.—Ins. Galap.
suffruticosa, *Torr. in Emory, Mex. Bound. Surv.* 181.—N. Mexico.
tenella, Colla, in Mem. Acc. Torin. xxxiii. (1829) 131. t. 9 = Telanthera polygonoides.
tenella, *Moq. in DC. Prod.* xiii. II. 355.—Hab.?
tomentella, *Seub. in Mart. Fl. Bras.* v. I. 186.—Bras.
triandra, Steud. Nom. ed. I. 33, 34 = Telanthera ficoidea.
triandra, Lam. Encyc. i. 95 = sessilis.
truxillensis, *H. B. & K. Nov. Gen. et Sp.* ii. 206.—Peruv.
versicolor, Hort. ex Regel, Gartenfl. (1869) 101 = Telanthera versicolor.
villiflora, *Scheele, in Linnaea*, xxii. (1849) 149.—Texas.
villosa, D. Dietr Syn. Pl. i. 867 = Brandesia villosa.
villosa, H. B. & K. Nov. Gen. et Sp. ii. 208 = Telanthera villosa.
villosa, *Sm. in Rees, Cycl. Suppl.* n. 9.—Hab.?
virgata, Spreng. ex Moq. in DC. Prod. xiii. II. 364 = Telanthera polygonoides.

ALTERNASEMINA, Silva Manso, Enum. 35 (1836) = **Cayaponia**, Mart. (Cucurb.).
Tayuia, Silva Manso, l. c. = C. Tayuya.

ALTHAEA, [Tourn.] Linn. Syst. ed. I (1735). *MALVACEAE*, Benth. & Hook. f. i. 260.
ALCEA, [Tourn.] Linn. Gen. ed. I. 206 (1737).
FERBERIA, Scop. Introd. 281 (1777).
acaulis, *Cav. Diss.* ii. t. 27. f. 3 (p. 93).—Syria.
aegyptiaca, Boiss. Diagn. Ser. II. i. 103 = lavateraeflora.
africana, *Lam. ex G. Don, Gen. Syst.* i. 467.—Afr. or.
ambigua, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 260 = Lavatera thuringiaca.
apterocarpa, *Fenzl, in Del. Sem. Hort. Vind.* (1858) 4.—Oriens.
arborea, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 260 = Lavatera arborea.
armeniaca, *Tenore, Ind. Sem. Hort. Neap.* (1837) 1.—Oriens.
atropurpurea, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva atropurpurea.
Aucheri, *Boiss. Diagn.* Ser. I. vi. 28.—Persia; Affghan.
borbonica, *DC. Prod.* i. 438 (= Pavonia?).—Ins. Borbon.
borealis, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva borealis.
Burchellii, DC. Prod. i. 438 = Pavonia Columella.
Calverti, *Boiss. Diagn.* Ser. II. v. 66.—Armenia.
cannabina, *Linn. Sp. Pl.* 686.—Europ.; Oriens.
caribaea, Sims, Bot. Mag. t. 1916 = rosea.
chinensis, Wall. Cat. n. 2689 = rosea.
coromandelina, Cav. Diss. ii. 93 = rosea.
corymbosa, Sw. Prod. Veg. Ind. Occ. 101 = Pavonia paniculata.
cretica, Weinm. in Syll. Ratisb. ii. (1828) 171 = rosea.
crispa, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva crispa.
digitata, Boiss. Diagn. Ser. I. viii. 106 = lavateraeflora.
ficifolia, *Cav. Diss.* ii. 92. t. 28. f. 2.—Sibir.
ficifolia, Sibth. & Sm. Fl. Graec. t. 663 = rosea.
flavo-virens, *Boiss. & Buhse, in Nouv. Mem. Soc. Nat. Mosc.* xii. (1860) 44.—Sibir. baical.
flexuosa, Sims, Bot. Mag. t. 892 = rosea.
Froloviana, *Fisch. ex G. Don, Gen. Syst.* i. 467.—Sibir.
Frutex, Hort. ex Mill. Gard. Dict. ed. VIII = Hibiscus syriacus.
garipensis, E. Mey. Zwei Pfl. Docum. 163 = Ludwigii.
gariepina, E. Mey. ex Boiss. Fl. Orient. i. 824 = praec.
glabriflora, *Ledeb. ex Sweet, Hort. Brit.* ed. II. 60.—Sibir.
Godroni, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva ambigua.
grandiflora, Scop. Del. Insub. ii. 33 = Hibiscus roseus.
Heldreichii, *Boiss. Diagn.* Ser. II. i. 103.—Thessal.
hirsuta, *Linn. Sp. Pl.* 687.—Europ.; Oriens.
hirsuta, Sieber, ex Steud. Nom. ed. II. i. 66 = Malva cretica.
hispida, Moench, Meth. 612 = hirsuta.
Hohenackeri, *Boiss. & Huet, Diagn.* Ser. II. v. 67.—Oriens.
indica, Steud. Nom. ed. II. i. 66.—Java.
Irolowiana, Steud. l. c. sphalm. = Froloviana.
kirrindensis, Boiss. Diagn. Ser. II. v. 67 = lavateraeflora.
Kotschyi, Boiss. l. c. Ser. II. i. 102 = cannabina.
Kraussiana, Buching. ex Krauss, in Flora, xxviii. (1844) 293 = Pavonia Kraussiana.
kurdica, *Schlecht. in Linnaea*, xvii. (1843) 127.—Oriens.
laevis, Moench, Meth. Suppl. 199 = Ludwigii.
lavateraeflora, *DC. Prod.* i. 437.—Oriens.
laxiflora, *A. DC. ex Schlecht. in Bot. Zeit.* vi. (1848) 341.—Hab.?
leucantha, Fisch. ex Loud. Hort. Brit. 287 = nudiflora.
lilacina, Boiss. & Kotschy, ex Boiss. Fl. Orient. i. 821 = apterocarpa.
longiflora, Boiss. & Reut. Diagn. Pl. Hispan. 9 = hirsuta.
Ludwigii, *Linn. Mant.* 98.—Oriens; Afr. bor. et austr.
marcotica, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva marcotica.
mauritiana, Alef. l. c. 260 = Malva mauritiana.
meonantha, Link, in Linnaea, ix. (1834) 586 = rosea.
mexicana, *Kunze, l. c.* xx. (1847) 51, 404.—Mexic.
microcarpa, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva parviflora.
multiflora, Reichb. ex Regel, Ind. Sem. Hort. Petrop. (1867) 36 = armeniaca.

ALTHAEA:—
mutica, J. Gay, ex Ball, in Journ. Linn. Soc. xvi. (1878) 376 = hirsuta.
narbonensis, Jacq. Coll. ii. 275 = cannabina.
narbonnensis, *Pourr. ex Cav. Diss.* ii. 94. t. 29. f. 2.—Europ.
nicaeensis, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva nicaeensis.
nudiflora, *Lindl. in Trans. Hort. Soc.* vii. (1830) 251.—Sibir. altaic.
officinalis, *Linn. Sp. Pl.* ed. I. 686.—Europ.; Oriens.
pallida, *Wald. & Kit. Pl. Rar. Hung.* t. 47.—Hungar.; Turc.
parviflora, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva parviflora.
plebeia, Schult. ex Steud. Nom. ed. II. i. 66 = Lavatera plebeia.
pulchra, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 112.—Ind. or.
racemosa, Sw. Prod. Veg. Ind. Occ. 102 = Pavonia racemosa.
remotiflora, *Boiss. & Heldr. Diagn.* Ser. I. viii. 108.—As. Min.
rhyticarpa, Trautv. in Bull. Soc. Nat. Mosc. (1860) i. 454 = sulphurea.
rosea, *Cav. Diss.* ii. 91. t. 29. f. 3.—Oriens.
rosea, Hohen. ex Boiss. Diagn. Ser. I. v. 67 = Hohenackeri.
rufescens, *Boiss. Diagn.* Ser. II. i. 102.—Syria.
setosa, *Boiss. l. c.* Ser. I. viii. 107.—Syria.
Sieberi, *Hort. ex G. Don, in Loud. Hort. Brit.* 486.—Sicilia.
sinensis, Blanco, Fl. Filip. ed. I. 552 = rosea.
sinensis, *Cav. Diss.* ii. t. 29. f. 3.—China.
speciosa, *Vis. Cat. Orto Bot. Padov.* 132.—Hab.?
spicata, Spreng. Syst. iii. 108 = Hibiscus spicatus.
striata, *DC. Prod.* i. 437.—Arab.
sublobata, Stokes, Bot. Mat. Med. iii. 530 = officinalis.
sulphurea, *Boiss. & Hohen. Diagn.* Ser. I. viii. 108.—Persia; Affghan.; Soongar.
sylvestris, *Briganti, ex Tenore, Syll.* 540.—Italia.
tabrisiana, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 44 = ficifolia.
taurinensis, C. A. Mey. Verz. Pfl. Cauc. 207 = armeniaca.
taurinensis, DC. Prod. i. 436 = officinalis.
thuringiaca, Kuntze, Taschen-Fl. Leipz. 212 = Lavatera thuringiaca.
Tournefortiana, *Tausch, in Flora*, xii. (1829) 9.—Creta.
ulmifolia, Pritzel, Ic. Ind. 50 = Sida glomerata.
verticillata, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 261 = Malva verticillata.
vulgaris, Alef. l. c. 260 = Malva vulgaris.

ALTHEA, Crantz, Stirp. Austr. 144 (1769) = praec.

ALTHENIA, Petit, in Ann. Sc. Observ. i. (1829) 451. *NAIADACEAE*, Benth. & Hook. f. iii. 1016.
Barraudonii, Duval-Jouve, in Bull. Soc. Bot. Fr. xix. (1872) p. lxxxix; Compt.-rend. lxxv. (1872) 95 = filiformis.
filiformis, *Petit, in Ann. Sc. Observ.* i. (1829) 451.—Gallia.
setacea, Delile, ex Petit, l. c. = filiformis.

ALTHERIA, Thou. Gen. Nov. Madag. 19 (1809) = **Melochia**, Linn. (Stercul.).
madagascariensis, DC. Prod. i. 493.—Madag.

ALTHINGIA, Steud. Nom. ed. II. i. 66 (1840) = seq.

ALTINGIA, [G. Don, in] Loud. Hort. Brit. 403 (1830) = **Araucaria**, Juss. (Conif.).
Cunninghamii, [G. Don,] l. c. = Ar. Cunninghamii.
excelsa, [G. Don,] l. c. = Ar. excelsa.

ALTINGIA, Noronha, in Verh. Bat. Gen. v. (1785) Art. ii. 9; ed. II. 41. *HAMAMELIDEAE*, Benth. & Hook. f. i. 669.
SEDGWICKIA, Griff. in As. Res. xix. (1836) 98. t. 15.
chinensis, Oliver, ex Hance, in Journ. Linn. Soc. xiii. (1873) 103.—China.
caerulea, Poir. Encyc. Suppl. v. 545 = Liquidambar Altingia.
excelsa, *Noronha, in Verh. Bat. Gen.* v. ed. II. 41 (1827).—Malaya.

ALTORA, Adans. Fam. ii. 356 (1763) = **Cluytia**, Linn. (Euphorb.).

ALUGHAS, Linn. Fl. Zeyl. 207 (1747) = **Alpinia**, Linn. (Scitam.).

ALVARADOA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 100. *SAPINDACEAE*, Benth. & Hook. f. i. 411.
amorphoides, *Liebm. l. c.* 101.—Mexic.
arborescens, *Griseb. Cat. Pl. Cub.* 50.—Cuba.
jamaicensis, *Benth. Pl. Hartw.* 344.—Jamaica.
mexicana, *Liebm. ex Benth. l. c.*—Mexic.

ALVARDIA, Fenzl, in Flora, xxvii. (1844) 312 = **Peucedanum**, Linn. (Umbellif.).
arborescens, Fenzl, l. c. = P. fraxinifolium.

ALVAREZIA, Pav. ex Nees, in DC. Prod. xi. 465 (1847) = **Blechum**, R. Br. (Acanth.).
distícha, Pav. ex Nees, l. c. 246 = Lophostachys Alvarezia.
parviflora, Pav. ex Nees, l. c. 466 = B. Linnaei.

ALVESIA, Welw. Apont. 587. n. 47 (1858) = **Bauhinia**, Linn. (Legum.).
bauhinioides, Welw. l. c. = B. tomentosa.

ALVESIA, Welw. in Trans. Linn. Soc. xxvii. (1869) 55. t. 19. *LABIATAE*, Benth. & Hook. f. ii. 1176.
rosmarinifolia, *Welw. l. c.*—Afr. trop.

ALVISIA, Lindl. Fol. Orch. (1859) = **Eria**, Lindl. (Orchid.).
tenuis, Lindl. l. c. = E. articulata.

ALYCIA, Willd. ex Steud. Nom. ed. II. i. 66 (1840) = **Eriochloa**, H. B. & K. (Gramin.).
coarctata, Willd. ex Steud. l. c. = E. polystachya.
distachia, Willd. ex Steud. l. c. = E. polystachya.

ALYMERIA, D. Dietr. Syn. i. in Ind. p. xiii (1839) = AYLMERIA, Mart. = **Polycarpaea**, Lam. (Caryoph.).

ALYMNIA, Neck. Elem. i. 31 (1790) = **Polymnia**, Linn. (Compos.).

ALYOGYNE, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 12 = **Fugosia**, Juss. (Malvac.).
hakeifolia, Alef. l. c. = F. hakeaefolia.

ALYPUM, Fisch. Hort. Gorenk. ed. II. (1812) 19 = **Globularia**, Tourn. (Selagin.).
monspeliensium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 142.—G. Alypum.
salicifolium, Fisch. Cat. Hort. Gorenk. ed. II. (1812) 19 = G. Alypum.
Solandri, Hort. ex Steud. Nom. ed. I. 373 = G. Alypum.

ALYSICARPUS, Neck. Elem. iii. 15 (1790); Desv. Journ. Bot i. (1813) 120. t. 4. *LEGUMINOSAE*, Benth. & Hook. f. i. 522.
FABRICIA, Scop. Introd. 307 (1777).
HALLIA, J. St. Hil. in Bull. Soc. Philom. (Dec. 1813) 192.
HEGETSCHWEILERA, Heer & Regel, Cat. Sem. Hort. Turic. (1842); ex Bot. Zeit. i. (1843) 47.
MYSICARPUS, Webb, in Hook. Niger Fl. 122 (1849) lapsu.
belgaumensis, *Wight, Ic.* t. 92.—Ind. or.
bupleurifolius, Wall. Cat. n. 5761 = rugosus.
bupleurifolius, *DC. Prod.* ii. 352.—Ind. or.
cylindricus, Desv. in Ann. Linn. Soc. Par. iv. (1826) 301 = styracifolius.
diversifolius, Wall. Cat. n. 5772 = vaginalis.
ferrugineus, *Hochst. & Steud. ex A. Rich Tent. Fl. Abyss.* i. 207.—Abyss.
glaber, E. Mey. Comm. Pl. Afr. Austr. 125 = rugosus.
glumaceus, *DC. Prod.* ii. 353.—Arab.
glumaceus, Wall. Cat. n. 5764 = rugosus.
gracilis, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 170 = bupleurifolius.
gramineus, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Ind.* iii. (1846) 59, nomen.—Malaya.
hamosus, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 171.—Ind. or.
Harnieri, Scheinf. Reliq. Kotschy, 24. t. 19 = vaginalis.
heterophyllus, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 158 = vaginalis.

ALYSICARPUS:—
Heyneanus, Wight & Arn. Prod. i. 234 = rugosus.
Hochstetteri, A. Rich. Tent. Fl. Abyss. i. 209 = rugosus.
longifolius, *Wight & Arn. Prod.* i. 233.—Ind. or.
ludens, Wall. Cat. n. 5762 = bupleurifolius, rugosus.
lupulinus, Stocks, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 159 = rugosus.
monilifer, *DC. Prod.* ii. 353.—Ind. or.; Afr. trop.
nummularifolius, DC. l. c. = vaginalis.
nummularioides, Zipp. ex Span. in Linnaea, xv. (1841) 194 = scaber.
nummularius, Baill. in Bull. Soc. Linn. Par. i. (1884) 432 = vaginalis.
obovatus, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 169 = rugosus.
paradoxus, *Boiv. ex Baill. in Bull. Soc. Linn. Par.* i. (1884) 432.—Madag.
parviflorus, Dalz. in Hook. Kew Journ. iii. (1851) 211 = Desmodium parviflorum.
pilifera, Wall. Cat. n. 5765 = rugosus.
porrectus, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 171.—Afr. trop.
pubescens, *Law, in Wight, Ic.* i. t. 250.—Ind. or.
pylifer, Steud. Nom. ed. II. i. 66 = rugosus.
quadrangularis, Edgew. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 159 = tetragonolobus.
Quartinianus, A. Rich. Tent. Fl. Abyss. i. 208 = rugosus.
racemosus, Benth. in Linnaea, xxiv. (1851) 642 = belgaumensis.
rotundifolius, Dalz. ex Hook. f. Fl. Brit. Ind. ii. 172 = Desmodium rotundifolium.
rotundifolius, Wight, ex Hook. f. l. c. 157 = hamosus.
Rubibarna, Buch.-Ham. ex Wall. Cat. n. 5771 = vaginalis, monilifera.
rugosus, *DC. Prod.* ii. 353.—Reg. trop.
rupicola, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 170.—Ind. or.
scaber, *Span. in Linnaea*, xv. (1841) 194.—Malaya.
scariosus, R. Grah. in Wall. Cat. n. 5766 = rugosus.
spicatus, Royle, ex Hook. f. Fl. Brit. Ind. ii. 159 = longifolius.
styracifolius, DC. Prod. ii. 353 = rugosus.
tetragonolobus, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 169.—Ind. or.
timorensis, *Span. in Linnaea*, xv. (1841) 194.—Ins. Timor.
trifoliatus, Stocks, ex Hook. f. Fl. Brit. Ind. ii. 158 = vaginalis.
vaginalis, *DC. Prod.* ii. 353.—Ind. or.
vaginalis, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 170 = monilifer.
vaginalis, Wall. Cat. n. 5763 = longifolius, rugosus.
varius, Wall. l. c. n. 5768 = vaginalis.
Wallichii, Wight & Arn. Prod. i. 234 = rugosus.
Zeyheri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 230.—Afr. austr.

ALYSSOIDES, Tourn. ex Adans. Fam. ii. 419 (1763) = **Vesicaria**, Lam. (Crucif.).
creticum, Medic. Phil. Bot. i. 189 = Alyssum creticum.
gemonense, Medic. l. c. = Alyssum gemonense.
leucoifolium, Medic. Pflanzeng. 63 = V. graeca.
sinuatum, Medic. Phil. Bot. i. 189 = Alyssum sinuatum.
tomentosum, Moench, Meth. 264 = Alyssum creticum.
utriculatum, Medic. Phil. Bot. 189 = V. graeca.

ALYSSOPSIS, Boiss. Diagn. Ser. I. vi. 14 (1842). *CRUCIFERAE*, Benth. & Hook. f. i. 74.
deflexa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 58.—Persia.
Drumondii, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) ii. 291 = Sisymbrium brachypodum.
Kotschyi, *Boiss. Diagn.* Ser. I. vi. 14.—Persia.
sagittata, Boiss. Fl. Orient. i. 182 = deflexum.

ALYSSOPSIS, Reichb. Nom. 181. n. 6961 (1841) = **Vesicaria**, Lam. (Crucif.).

ALYSSON, Crantz, Stirp. Austr. ed. II. 15 (1769) = seq.

ALYSSUM, Tourn. ex Linn. Syst. ed. I. (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 73.
ADUSETA, Adans. Fam. ii. 420 (1763).
ANODONTEA, G. Don, Gen. Syst. i. 180 (1831).
AURINIA, Desv. Journ. Bot. iii. (1814) 162.
BERTEROA, DC. Syst. ii. 290 (1821).

ALYSSUM:—
Glyce, Lindl. Syn. Brit. Fl. 26 (1828).
Konig, Adans. Fam. ii. 420 (1763).
Koniga, R. Br. in Denh. Clapp. & Oud. Voy. App. 214 (1826).
Lobularia, Desv. Journ. Bot. iii. (1814) 162.
Meniocus, Desv. l. c. 173.
Moenchia, Roth, Tent. Fl. Germ. i. 273 (1788).
Octadenia, R. Br. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 41 (1836).
Odontarrhena, C. A. Mey. in Ledeb. Fl. Alt. iii. 58 (1831).
Onodontea, G. Don, Gen. Syst. i. 148 (1831), sphalm.
Psilonema, C. A. Mey. in Ledeb. Fl. Alt. iii. 50 (1831).
Ptilotrichum, C. A. Mey. l. c. 64 (1831).
Schivereckia, Andrz. ex DC. Syst. ii. 300 (1821).
Stevena, Andrz. ex DC. Syst. ii. 291 (1821).
affine, Tenore, Syll. 315 = orientale.
aizoides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 153.—Armenia.
albescens, Schlecht. ex DC. Syst. ii. 309, et Prod. i. 162 = montanum.
algeriense, *Pomel, Nouv. Mat. Fl. Atl.* 232.—Afr. bor.
alpestre, Bieb. Fl. Taur. Cauc. ii. 102, Suppl. 432 = Marschallianum.
alpestre, *Linn. Mant.* 92.—Europ.
alpestre, Wulf. ex Nym. Consp. 56 = montanum.
alpestre, Salis, in Flora, xvii. (1877) II. Beibl. 77 = Robertianum.
alpestre, Sibth. & Sm. Fl. Graec. t. 623 = argenteum.
alpinum, Scop. Fl. Carn. ed. II. ii. 9 = montanum.
alsinefolium, Host, Fl. Austr. ii. 248 = calycinum.
altaicum, C. A. Mey. in Ledeb. Fl. Alt. iii. 55 = Fischerianum.
alyssoides, Linn. Syst. ed. X. 1130 = calycinum.
andinum, *Rupr. Fl. Caucas.* 103.—Reg. Caucas.
ar[c]ticum, Wormsk. ex Vahl, in Fl. Dan. t. 1520 = Vesicaria arctica.
arenarium, Loisel, Fl. Gallica, ed. I. 401 = montanum.
arenarium, *Presl, Fl. Sic.* i. 58.—Sicilia.
argenteum, *Vitm. Summ.* iv. 30.—Europ.; Oricns.
argyrophyllum, *Schott & Kotschy, in Oestr. Bot. Wochenbl.* vii. (1857) 229.—As. Min.
armenum, *Boiss. Fl. Orient.* i. 278.—Armenia.
arvaticum, Jord. Diag. i. 200 = calycinum.
atlanticum, Desf. Fl. Atlant. ii. 71. t. 149 = montanum.
Aucheri, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 156 = foliosum.
aurantiacum, *Boiss. Fl. Orient.* i. 276.—Lycia.
aurcum, *Boiss. l. c.* 286.—Oriens.
Bertholonii, Desv. Journ. Bot. iii. (1814) 172 & 185 = argenteum.
Besseri, *Stev. ex DC. Syst.* ii. 300 = podolicum.
beugesiacum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 10 = montanum.
bidentatum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 108 = Draba brachycarpa.
Bilimekii, *Willk. in Willk. & Lange, Prod.* iii. 830.—Hispan.
brachystachyum, Bieb. Fl. Taur. Cauc. iii. 434 = umbellatum.
bracteatum, Boiss. Diagn. Ser. I. xiii. 77 sphalm. = Acanthophyllum bracteatum.
bracteatum, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 18.—Persia; Armenia.
brevifolium, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 11 = montanum.
brigantiacum, Jord. & Fourr. l. c. 12 = montanum.
Bungei, *Boiss. Fl. Orient.* i. 274.—Persia.
calicinum, Neck. Delic. Gallo-Belg. i. 277 = calycinum.
callichroum, Boiss. Diagn. Ser. II. v. 34 = constellatum.
calycinum, Haenscl. ex Nym. Consp. 56 = campestre.
calycinum, *Linn. Sp. Pl.* ed. II. 908.—Europ.; Oricns.
calycinum, Pall. ex DC. Syst. ii. 318 = dasycarpum.
calyocarpum, *Rupr. Fl. Cauc.* 103, in textu.—Tauria.
campestre, Hoffm. Fl. Deutschl. ed. II. ii. 43 = calycinum.
campestre, *Linn. Syst.* ed. X. 1130.—Europ.; Oricns.
campestre, Marz. ex Poll. Fl. Ver. ii. 362 = montanum.
canariense, Del. ex Webb & Berth. Phyt. Canar. i. 91 = lybicum.
candidum, Schlecht. ex DC. Syst. ii. 308 = Marschallianum.
canescens, *DC. Syst.* ii. 322.—Sibir.; Reg. Himal.

ALYSSUM:—
cassium, *Boiss. & Heldr. Diagn.* Ser. I. viii. 34.—Syria.
castellanum, Bourg. ex Nym. Consp. 57 = serpyllifolium.
cedrorum, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 169 = hirsutum.
Cephalotes, *Boiss. Diagn.* Ser. II. i. 34.—Caria.
chalcidicum, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 175.—Turcia.
cheiranthifolium, Willd. Sp. Pl. iii. 468 = Farsetia clypeata.
cheirifolium, Berger. ex Steud. Nom. ed. I. 34 = saxatile.
chrysanthum, Boiss. Fl. Orient. i. 271 = argenteum.
ciliatum, Lam. Fl. Fr. ii. 479 = Draba aizoides.
cilicicum, *Boiss. & Bal. Diagn.* Ser. II. v. 34.—Cilicia.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 236.—Afr. bor.
clypeatum, Linn. Sp. Pl. ed. I. 651 = Farsetia clypeata.
cochlearioides, Roth, Nov. Pl. Sp. 322 = Cochlearia flava.
cochleatum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 11.—Afr. bor.
collinum, Brot. Phyt. Lusit. ii. 209. t. 180 = campestre.
collinum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 9 = montanum.
compactum, *De Not. ex Ces. Ic. Stirp. Ital.* fasc. iii. 15. t. 5.—Sicilia.
condensatum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 268.—As. Min.; Syria.
confertum, Hort. Berol. ex Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 155 = strictum.
constellatum, *Boiss. l. c.* Sér. IV. ii. (1854) 244.—Cilicia.
contemptum, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 177 = Szowitsianum.
cordatum, Spreng. ex Steud. Nom. ed. II. i. 67 = obcordatum.
cornutum, Steph. ex Ledeb. Fl. Ross. i. 139 = rostratum.
corsicum, *Duby, Bot. Gall.* ed. II. i. 34.—Corsica.
corsicum, Rob. ex Gren. & Godr. Fl. Fr. i. 117 = Robertianum.
corymbosum, *Boiss. Fl. Orient.* i. 265.—Turc.
corymbulosum, Boiss. l. c. 282 = umbellatum.
crenulatum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 33.—Syria.
cretaceum, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 109.—Sibir.
creticum, *Linn. Sp. Pl.* 661.—Creta.
cryptopetalum, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 142; *Lehm. Rel.* (1847) 142 = marginatum.
cuneifolium, Tenore, Fl. Nap. Prod. xxxvii. App.; Cat. Neap. (1815) 58 = montanum.
dagestanicum, *Rupr. Fl. Caucas.* 104.—Caucas.
dahuricum, Schlecht. ex DC. Syst. ii. 715 = canescens.
damascenum, *Boiss. & Gaill. Diagn.* Ser. II. vi. 18.—Syria.
dasycarpum, *Steph. in Willd. Sp. Pl.* iii. 469.—Oriens; Turkest.; Affghan.
dauricum, Schlecht. in DC. Syst. ii. 322, & Prod. i. 164 = canescens.
davuricum, Willd. ex Ledeb. Fl. Ross. i. 138 = Fischerianum.
decoloratum, *Pomel, Nouv. Mat. Fl. Atl.* 236.—Afr. bor.
decumbens, Herbich, Stirp. Rar. Bucov. 47 = repens.
deltoideum, Linn. Sp. Pl. ed. II. 908 = Aubrietia deltoidea.
densiflorum, Desf. Cor. Tourn. 65. t. 48 = strictum.
dentatum, Nutt. Gen. Am. ii. 63 = Draba ramosissima.
dentatum, Willd. Enum. Hort. Berol. 671 = Camelina sativa.
dicranoides, Boiss. & Huet, Diagn. Ser. II. v. 30 = Draba rigida.
diffusum, Jord. & Fourr. ex Nym. Consp. 57 = argenteum.
diffusum, Tenore, App. Hort. Neap. (1815) 58 = montanum.
Draba, Bieb. ex Willd. Enum. Hort. Berol. 672 = linifolium.
Draba, Hort. ex Webb & Berth. Phyt. Canar. i. 91 = lybicum.
Draba, Willd. Enum. Hort. Berol. 672 = hirsutum.
dubium, Bory, in Ann. Gén. Soc. Phys. iii. (1820) 4 = Lagascae.
edentulum, Waldst. & Kit. Pl. Rar. Hung. i. 95. t. 92 = gemonense.

ALYSSUM:—
elatum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 33.—As. Min.
emarginatum, *Zahl. ex Vis. Fl. Dalm.* iii. 117.—Dalmat.
erigens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 9 = montanum.
eriocarpum, Pourr. ex Steud. Nom. ed. II. i. 67 = Clypeola eriocarpa.
eriophorum, Willd. Enum. Hort. Berol. 671 = Clypeola eriophora.
eriophyllum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 273.—As. Min.
erosulum, Genn. & Pestal. ex Clem. in Mem. Acc. Sc. Torin. Ser. II. xvi. (1857) 248 = suffrutescens.
erraticum, Jord. Diag. i. 200 = calycinum.
filifolium, *Wahlenb. in Isis*, xxi. 998.—As. Min.
Fischerianum, *DC. Syst.* ii. 311.—Sibir.
flexicaule, Jord. Obs. Pl. Crit. i. (1846) 10 = montanum.
floribundum, *Boiss. Diagn.* Ser. II. v. 33.—Cilicia.
foliosum, *Bory & Chaub. Nouv. Fl. Pélop.* 42 n. 1030.—Graecia; As. Min.
fruticulosum, Steud. Nom. ed. I. 34 = Vesicaria fruticulosa.
fulvescens, *Sibth. & Sm. Prod. Fl. Graec.* ii. 13.—Graecia; As. Min.
gemonense, *Linn. Mant.* 92.—Europ.
gemonense, Roth, Tent. Fl. Germ. iii. 85 (err. glemonense) = saxatile.
globosum, Steud. Nom. ed. I. 34 = Vesicaria globosa.
glomeratum, *Burch. ex DC. Syst.* ii. 316.—Afr. austr.
Gmelini, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 8 = montanum.
gnaphalodes, DC. Syst. ii. 311 = Vesicaria gnaphalodes.
gracile, *Rafin. in Am. Monthly Mag.* (1811) 356, nomen.—Hab.?
granatense, Boiss. & Reut. Pugill. Pl. Nov. (1852) 9 = calycinum.
granatense, Nym. Consp. 57 = serpyllifolium.
halimifolium, DC. Fl. Fr. v. 594 = pyrenaicum.
halimifolium, Lap. Hist. Abr. Pl. Pyr. 371 = macrocarpum.
halimifolium, Linn. Sp. Pl. ed. I. 650 = maritimum.
halimifolium, Pall. Reise, iii. 161 = lenense.
halimifolium, *Willd. Sp. Pl.* iii. 460.—Europ.
Heideri, *Krock. Fl. Sil. Suppl.* ii. 168.—Europ.
Haussknechtii, *Boiss. Fl. Orient.* i. 269.—As. Min.
Heinzi, *Ullep. in Oestr. Bot. Zeitschr.* xxxv. (1885) 307.—Europ.
helianthemifolium, *Timb. & Jeanb. in Rev. Bot. Soc. Fr.* iii. (1884–85) 105.—Gallia.
heterophyllum, Ruiz & Pav. ex DC. Syst. ii. 542 = Lepidium perfoliatum.
heterotrichum, *Boiss. Diagn.* Ser. I. vi. 15.—Persia.
hirsutum, Bess. ex Nym. Consp. 56 = rostratum.
hirsutum, *Bieb. Fl. Taur. Cauc.* ii. 106.—Oriens.
hirsutum, Hohen. Enum. Elisabethpol. 243 = campestre.
homalocarpum, *Boiss. Fl. Orient.* i. 285.—Arabia.
horebicum, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 156 = homalocarpum.
Huetii, *Boiss. Fl. Orient.* i. 287.—As. Min.
hyperborcum, *Linn. Sp. Pl.* 651.—Am. bor.
idaeum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 35.—Creta.
illyricum, Willd. Enum. Suppl. 44 = linifolium.
imbricatum, Schlecht. ex DC. Syst. ii. 308, & Prod. i. 161 = canescens.
incanum, *Linn. Sp. Pl.* 650.—Europ.; Sibir.
Jonthlaspi, [Clairv.] Man. Herb. 216 = Clypeola Jonthlaspi.
Kotschyanum, Boiss. Fl. Orient. i. 270 = masmenacum.
Lagascae, *Janka, in Termész. Füzetek*, vii. (1883) 114.—Europ.
lanigerum, *DC. Syst.* ii. 308.—Persia.
lanuginosum, Pourr. ex DC. Syst. ii. 327 = Clypeola eriophora.
Lapeyrousianum, *Jord. Obs. Pl. Crit.* i. (1846) 3 = Mont. Pyren.
latifolium, *Vis. Fl. Dalm.* iii. 118.—Dalmat.
leiocarpum, *Pomel, Nouv. Mat. Fl. Atl.* 235.—Afr. bor.
lenense, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 110.—Sibir.
lepidotum, *Boiss. Diagn.* Ser. I. i. 73.—As. Min.
Lescurii, *A. Gray, Man. Bot. U. St.* ed. V. 72.—Am. bor.

ALYSSUM :—
leucadeum, *Guss. Pl. Rar.* 268.—Ital.
libycum, *Coss. in Bull. Soc. Bot. Fr.* iv. (1857) 467.—Afr. bor.
linearifolium, Lag. Gen. et Sp. Nov. 19 = linifolium.
linifolium, *Steph. ex Willd. Sp. Pl.* iii. 467.—Europ.; Oriens.
litorale, *Rupr. Fl. Cauc.* 103. in textu.—Littora Maris Balt.
longicaule, *Boiss. in Bibl. Genèv.* (1838) 407.—Hispan.
ludovicianum, Nutt. Gen. Am. ii. 63 = Vesicaria ludoviciana.
lunarioides, Spreng. ex DC. Syst. ii. 292 = mutabile.
lunarioides, Willd. Sp. Pl. iii. 461 = Farsetia lunarioides.
lusitanicum, Brot. ex Nym. Consp. 56 = campestre.
luteolum, *Pomel, Nouv. Mat. Fl. Atl.* 234.—Afr. bor.
macrocalyx, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 12.—Afr. bor.
macrocarpum, *DC. Syst.* ii. 321.—Gall. austr.
macropodum, *Boiss. & Bal. Diagn.* Ser. II. vi. 18.—Cappadoc.
macrostylum, *Boiss. & Huet, ex Boiss. Fl. Orient.* i. 279.—As. Min.
marginatum, *Steud. Nom.* ed. II. i. 68 (*ex Boiss. in Ann. Sc. Nat.* xvii. (1842) 157).—Arab.; Persia; Affghan.
marginatum, *Timb. & Jeanb. in Rev. Bot. Soc. Fr.* iii. (1884–85) 106.—Gallia.
maritimum, *Lam. Encyc.* i. 98.—Europ.
Marschallianum, *Andrz. ex DC. Syst.* ii. 308.—Caucas.
masmenaeum, *Boiss. Diagn.* Ser. II. v. 36.—Cappadoc.
medium, Host, Fl. Austr. ii. 244 = gemonense.
meniocoides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 158.—Mesopot.
mesopotamicum, Fenzl, in Tchichat. Asie Min. Bot. i. 311 = argenteum.
Meyerianum, Rupr. Fl. Cauc. 100, in textu = tortuosum.
micranthum, Fisch. & Mey. in Linnaea, x. (1836) 83 = campestre.
microcarpon, Vis. Fl. Dalmat. iii. 115 = corymbosum.
microcarpum, Neilr. ex Nym. Consp. 55 = gemonense.
microcephalum, Boiss. Diagn. Ser. I. xiii. 77, sphalm. = Acanthophyllum microcephalum.
micropetalum, Fisch. in DC. Syst. ii. 313 = campestre.
micropetalum, *Kit. in Linnaea*, xxxii. (1863) 497.—Europ.
microphyllum, *Steud. Nom.* ed. II. i. 68.—Sibir. alt.
minimum, Linn. Sp. Pl. 651 = maritimum.
minimum, Pall. Reise, ii. 741 = linifolium.
minimum, *Willd. Sp. Pl.* iii. 464.—Europ.; Oriens; Reg. Himal.
minutiflorum, Boiss. Diagn. Ser. I. i. 73 = alpestre.
minutulum, Schleich. ex DC. Syst. ii. 307 = alpestre.
minutum, Patr. ex DC. l. c. 322 = canescens.
minutum, *Schlecht. ex DC. l. c.* 316.—Sibir.
modestum, Boiss. Diagn. Ser. II. vi. 17 = minutum.
montanum, Brot. Fl. Lusit. ii. 558 = campestre.
montanum, *Linn. Sp. Pl.* 650.—Europ.; Oriens.
montanum, Pall. Reise, i. 252 = Fischerianum.
montanum, Patr. ex DC. Syst. ii. 322 = canescens.
mouradicum, *Boiss. & Bal. Diagn.* Ser. II. vi. 16.—As. Min.
Muellerii, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 16.—Armenia; Persia.
murale, Bieb. Fl. Taur. Cauc. ii. 103 = alpestre.
murale, Waldst. & Kit. Pl. Rar. Hung. i. 5. t. 6 = argenteum.
murcicum, Jord. ex Nym. Consp. 57 = serpyllifolium.
muricatum, *Kit. in Linnaea*, xxxii. (1863) 498.—Europ.
mutabile, *Vent. Jard. Cels*, 85.—Ital.
myagroides, All. Fl. Pedem. i. 241 = Graellsia saxatilis.
Myagrum, Wibel, Prim. Fl. Werth. 204 = Camelina sativa.
nanum, *Pomel, Nouv. Mat. Fl. Atl.* 236.—Afr. bor.
nebrodense, Tineo, Pugill. Pl. Sic. 12 = serpyllifolium.
novum, Kit. Ind. Pesth, t. 6 = alpestre.
numidicum, *Pomel, Nouv. Mat. Fl. Atl.* 377.—Afr. bor.
obcordatum, *DC. Syst.* ii. 324.—Hab.?
obliquum, Sibth. & Sm. Fl. Graec. t. 623 = mutabile.
obovatum, Turcz. in Bull. Soc. Nat. Mosc. (1837) i. 57 = sibiricum.
obtusifolium, Stev. ex DC. Syst. ii. 305 = alpestre.

ALYSSUM :—
ochroleucum, Boiss. Diagn. Ser. II. v. 36 = montanum.
Oederi, Durand, Fl. Bourg. i. 161 = Vesicaria utriculata.
orbiculare, *Timb. & Jeanb. in Rev. Bot. Soc. Fr.* iii. (1884–85) 106.—Europ.
orbiculatum, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 335.—Graecia.
orientale, *Ard. Animad. Alt.* 32.—Graecia; As. Min.
orophilum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 11 = montanum.
ovirense, *A. Kern. Sched. Fl. Exsicc.* ii. 99.—Austria.
oxycarpum, *Boiss. & Bal. Diagn.* Ser. II. v. 35.—Cappadoc.
paniculatum, Desf. Choix, 67 = Vesicaria paniculata.
paniculatum, Willd. Enum. Hort. Berol. 671 = Neslia paniculata.
parviflorum, Bieb. Fl. Taur. Cauc. iii. 434 = campestre.
parviflorum, Hochst. & Steud. ex Steud. Nom. ed. II. i. 68 = marginatum.
parviflorum, Schloss. ex Nym. Consp. 57 = calycinum.
parvulum, *Schrad. in Linnaea*, x. (1836) *Litt.* 69.—Hab.?
patulum, *Pomel, Nouv. Mat. Fl. Atl.* 238.—Afr. bor.
pedemontanum, *Rupr. Fl. Cauc.* 102, *in textu*.—Europ.
peltarioides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 158.—As. Min.
persicum, *Boiss. l. c.* 152.—Persia.
Perusianum, J. Gay, ex Gren. & Godr. Fl. Fr. i. 118 = Lapeyrousianum.
petalodes, DC. Syst. ii. 312 = Farsetia obovata.
petraeum, Andrz. ex DC. l. c. 302 = saxatile.
petraeum, Ard. Animad. Alt. 30 = gemonense.
petrophilum, Heldr. ex Nym. Consp. 56 = montanum.
Peyrousianum, Willk. & Lange, Prod. Fl. Hisp. iii. 835 = Lapeyrousianum.
phymatocarpum, Schlosser & Vukot. Syll. 156 = calycinum.
pleiospermum, Fenzl, in Russeg. Reise, Illustr. 54 = aureum.
podolicum, *Bess. Cat. Hort. Crem.* (1816) 8.—Europ.; As. Min.; Sibir.
polyodon, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 17 = campestre.
porphyreticum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 11 = montanum.
praecox, *Boiss. Fl. Orient.* i. 275.—As. Min.
psammeum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 9 = montanum.
psilocarpum, Boiss. Voy. Bot. Espagne, App. ii. 718 = calycinum.
purpureum, Lag. & Rodr. in Analcs Cienc. Nat. v. (1802) 275 = Lagascae.
pyrenaicum, *Lapeyr. Hist. Abr. Pl. Pyr.* 371. *Suppl.* 91.—Mont. Pyren.
ramosissimum, *Webb, in Giorn. Bot. Ital.* ii. (1846) 215.—Graecia.
repens, Baumg. Enum. Transs. ii. 237 = montanum.
rhodanense, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 10 = montanum.
Robertianum, *Bernard, ex Gren. & Godr. Fl. Fr.* i. 117.—Corsica.
Rochelii, Andrz. ex Reichb. Ic. Pl. Crit. xi. p. 8. n. 4273 b = montanum.
rostratum. *Stev. in Mém. Acad. Pétersb.* iii. (1809–10) 295. t. 15. f. 1.—Graecia; As. Min.
rostratum, Wierzb. ex Nym. Consp. 56 = montanum.
ruderale, Jord. Diagn. i. 198 = calycinum.
rupestre, *Tenore, Fl. Nap.* t. 60.—Europ.
rupestre, Willd. Enum. Hort. Berol. 672 = Graellsia saxatilis.
sabulosum, Jord. Diag. i. 199 = calycinum.
samariferum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 272.—As. Min.
samolifolium, Desf. Choix, 66 = Thlaspi samolifolium.
sativum, Scop. Fl. Carn. ed. II. ii. 9 = Camelina sativa.
savranicum, Andrz. ex Bess. Enum. Pl. Volh. 82 = alpestre.
saxatile, Bory & Chaub. Nouv. Fl. Pélop. 41 = orientale.
saxatile, Crantz, Stirp. Austr. 16 = gemonense.
saxatile, *Linn. Sp. Pl.* 650.—Europ.
saxatile, Sibth. & Sm. Prod. Fl. Graec. ii. 11 = orientale.
scabrum, Weinm. Cat. Hort. Dorpat. (1810) 11 = umbellatum.

ALYSSUM :—
Schlosseri, Heuff. ex Schlosser & Vukot. Syll. 156 = calycinum.
schirwanicum, Rupr. Fl. Cauc. 100 = tortuosum.
Scovitzianum, Fisch. & Mey. in Linnaea, xii. (1838) sphalm. 151 = Szovitzianum.
scutigerum, *Dur. in Expl. Sc. Algér.* t. 72. f. 4.—Afr. bor.
semidodecandra, *Schrank, Baier. Fl.* ii. 189.—Europ.
serpyllifolium, Bieb. Fl. Taur. Cauc. ii. 103 = tortuosum.
serpyllifolium, Desf. Fl. Atlant. ii. 70 = alpestre.
sibiricum, Fisch. ex DC. Syst. ii. 311 = Fischerianum.
sibiricum, Trevir. in Ges. Naturf. Fr. Berl. Mag. vii. (1811) 151 = dasycarpum.
sibiricum, *Willd. Sp. Pl.* iii. 465.—Sibir.
siculum, Jord. Diag. i. 202 = calycinum.
simplex, Rudolphi, in Schrad. Journ. ii. (1799) 290 = campestre.
sinuatum, *Linn. Sp. Pl.* 651.—Europ.; As. Min.
smyrnaeum, *C. A. Mey. in Bull. Sc. Acad. Pétersb.* vii. (1840) 132.—Smyrna.
spathulatum, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 110 = alpestre.
spathulatum, *Steph. in Willd. Sp. Pl.* iii. 465.—Sibir.
speciosum, *Pomel, Nouv. Mat. Fl. Atl.* 237.—Afr. bor.
sphacioticum, *Boiss. & Heldr. Diagn.* Scr. I. viii. 35.—Creta.
spinosum, *Linn. Sp. Pl.* 650.—Europ.
Spruneri, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 12 = montanum.
strictum, C. A. Mey. Verz. Pfl. Cauc. 137 = Szowitsianum.
strictum, *Willd. Sp. Pl.* iii. 464.—As. Min.
strigosum, [Soland. in] Russell, Aleppo, ed. II. ii. 257 = hirsutum.
strigulosum, *Amo, Fl. Iber.* vi. 571.—Hispan.
stylare, *Boiss. & Bal. ex Boiss. Fl. Orient.* i. 287.—As. Min.
styriacum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 7 = montanum.
subalpestre, Schur, Enum. Pl. Transs. 61 = argenteum.
subalpinum, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. ii. 103 = tortuosum.
sublineare, Jord. Diagn. i. 201 = calycinum.
subtumidum, Boiss. & Heldr. ex Nym. Consp. 56 = foliosum.
suffruticscens, *Boiss. Fl. Orient.* i. 275.—Syria.
superbum, L'Hérit. ex DC. Syst. ii. 288 = Farsetia suffruticosa.
Szovitsianum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 31.—Oriens.
tenuifolium, Bieb. Fl. Taur. Cauc. Suppl. 435 = cancscens.
tenuifolium, *Steph. in Willd. Sp. Pl.* iii. 460.—Ross.
tetraspermum, Bertol. in Nov. Comm. Acad. Bonon. vi. (1844) 226 = meniocoides.
tetrastemon, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 153.—As. Min.; Syria.
tortuosum, *Rupr. Fl. Caucas.* 98.—Caucas.
tortuosum, Waldst. & Kit. Pl. Rar. Hung. i. 94. t. 91 = alpestre.
transsylvanicum, Schur, Enum. Pl. Transs. 63 = montanum.
trichostachyum, *Rupr. Fl. Caucas.* 101.—Reg. Caucas.
triquetrum, Portenschl. ex DC. Syst. ii. 290 = Farsetia triquetra.
trochocarpum, Fenzl, in Tchihat. Asie Min. Bot. i. 297 = argenteum.
tuberculatum, Kit. in Linnaea, xxxii. (1863) 498 = muricatum.
turkestanicum, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 6.—Turkest.
umbellatum, *Desv. Journ. Bot.* iii. (1814) 173 & 184.—As. Min.; Syria.
utriculatum, Linn. Mant. i. 92 = Vesicaria utriculata.
utriculatum, Sibth. & Sm. Fl. Graec. t. 607 = Vesicaria Barrelieri.
vagum, Jord. Diag. i. 198 = calycinum.
vernale, Kit. ex Hornem. Hort. Hafn. 601 = montanum.
Vesicaria, Linn. Sp. Pl. ed. I. 651 = Vesicaria utriculata.
vesicarioides, Andrz. ex Steud. Nom. ed. II. i. 68 = gemonense.
Wierzbickii, Heuff. in Flora, xviii. (1835) i. 242 = rostratum.

ALYSSUM:—
Willkommii, De Roem. ex Willk. in Linnaea, xxv. (1852) 8 = granatense.
Wulfenianum, Willd. Enum. Hort. Berol. Suppl. 44 = montanum.
xanthocarpum, Boiss. in Ann. Sc. Nat. Sér. xvii. (1842) 154 = macrostylum.
xerophilum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 10 = montanum.

ALYXIA, Banks, ex R. Br. Prod. 469 (1810). *APOCYNACEAE*, Benth. & Hook. f. ii. 697.
GYNOPOGON, Forst. Char. Gen. 35. t. 18 (1776), nomen prius.
actinophylla, A. Cunn. in Bot. Mag. sub t. 3313 = verticillosa.
affinis, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 193.—N. Caled.
aromatica, Reinw. ex A. DC. in DC. Prod. viii. 346 = stellata.
bracteolosa, *Rich, ex A. Gray, in Proc. Am. Acad.* v. (1862) 332.—Ins. Fiji.
breviflora, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 195.—N. Caled.
buxifolia, *R. Br. Prod.* 470.—Austral.
calophylla, Wall. Cat. n. 1607 = Winchia calophylla.
capitellata, Benth. in Enum. Pl. Hueg. 81 = buxifolia.
ceylanica, *Wight, Ic. Pl.* t. 1293.—Zeylan.
coriacea, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 541. —Ind. or.
cuspidata, Wall. ex A. DC. in DC. Prod. viii. 351 = Melodinus orientalis.
daphnoides, *A. Cunn. in Hook. Bot. Mag.* t. 3313.—Ins. Norfolc.
disphaerocarpa, *Heurck & Muell. Arg. in Flora*, li. (1870) 169.—N. Caled.
erythrocarpa, *Vatke, in Bremen Abh.* ix. (1885) 124. —Ins. Mascar.
fascicularis, *Benth. & Hook. f. Gen.* ii. 698, *in nota ad Hunteriam.*—Reg. Himal.
Forsteri, *A. Cunn. ex Loud. Hort. Brit. Suppl.* i. 580 (1830) = A. Gynopogon.
glaucescens, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 542; G. Don, Gen. Syst. iv. 97 = Winchia calophylla.
glaucophylla, *Heurck & Muell. Arg. in Flora*, liii. (1870) 170.—N. Caled.
gracilis, *Benth. & Hook. f. Gen.* ii. 697, *in nota ad Hunteriam.*—Reg. Himal.
Gynopogon, *Roem. & Schult. Syst.* iv. 440.—Ins. Norfolc.
Gynopogon, Wall. Cat. n. 1605 = lucida.
gynopogona, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = seq.
Halmaheirae, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 140. —Malaya.
ilicifolia, *F. Muell. Fragm.* iv. 149.—Austral.
laurina, *Gaudich. in Freyc. Voy. Bot.* 451. t. 62.—Ins. Molucc.
leucogyne, *Heurck & Muell. Arg. in Flora*, liii. (1870) 170.—N. Caled.
Lindii, *F. Muell. Fragm.* viii. 46.—Ins. D. Howe.
lucida, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 540.—Malaya.
madagascariensis, *A. DC. in DC. Prod.* viii. 345.—Madag.
obovata, *Seem. Fl. Vit.* 156, *in nota.*—Ins. Pinet.
obtusifolia, *R. Br. Prod.* 470.—Austral.
odorata, *Wall. Cat.* n. 1606.—Tenasserim.
olivaeformis, *Gaudich. in Freyc. Voy. Bot.* 451.—Ins. Sandvic.
pilosa, *Miq. Fl. Ind. Bat.* ii. 408.—Java.
podocarpa, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 197.—N. Caled.
pugioniformis, A. Cunn. in Hook. Bot. Mag. sub t. 3312 = ruscifolia.
pumila, *Hook. f. Fl. Brit. Ind.* iii. 635.—Malacca.
quinata, *Miq. Fl. Ind. Bat.* ii. 407.—Sumatra.
Reinwardti, Blume, Cat. Gew. Buit. 43 = stellata.
Richardsonii, Sweet, Hort. Brit. ed. I. 273 = ruscifolia.
ruscifolia, *R. Br. Prod.* 470.—Austral.
scandens, *Roem. & Schult. Syst.* iv. 440.—Ins. Societ.
sinensis, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 334.—Hongkong.
Spanogheana, *Miq. Fl. Ind. Bat.* ii. 409.—Ins. Timor.
spicata, *R. Br. Prod.* 470.—Austral.
squamulosa, *C. Moore & F. Muell. in F. Muell. Fragm.* viii. 47.—Ins. D. Howe.

ALYXIA:—
stellata, *Roem. & Schult. Syst.* iv. 439.—As. trop.; Malaya; Ins. Pacif.
sulcata, Hook. & Arn. Bot. Beech. Voy. 90 = olivaeformis.
tetragona, R. Br. Prod. 470 = spicata.
Thozetii, *F. Muell. Fragm.* x. 103.—Austral.
thyrsiflora, *Benth. Fl. Austral.* iv. 309.—Austral.
Tisserantii, *Montr. in Mém. Acad. Lyon*, x. (1860) 233. —N. Caled.
Torresiana, *Gaudich. in Freyc. Voy. Bot.* 451.—Ins. Marian.
zeylanica, Walp. Ann. iii. 33 = ceylanica.

ALZALIA, F. G. Dietr. Lexik. Gaertn. i. 299 (1802) = **Alzatea**, Ruiz & Pav. (Celastr.).

ALZINIANA, F. G. Dietr. ex Pfeiff. Nom. i. 129 (1873) = praec.

ALZATEA, Ruiz & Pav. Prod. 40. t. 7 (1794). *CELASTRINEAE*, Benth. & Hook. f. i. 362.
mexicana, F. G. Dietr. Lexik. Gaertn. i. 299 (1802) Nachtr. = verticillata.
verticillata, *Ruiz & Pav. Fl. Per.* iii. 20. t. 241.—Peruv.

AMADEA, Adans. Fam. ii. 230 (1763) = **Androsace**, Tourn. (Primul.).

AMAGRIS, Rafin. in Préc. Somiol. 27 (1814) & Am. Monthly Mag. (1819) 190 = **Calamagrostis**, Adans. (Gramin.).

AMAIOUA, Aubl. Pl. Gui. Suppl. 13. t. 375 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 81.
EHRENBERGIA, Spreng. Neue Entdeck. ii. 129 (1821) & Syst. ii. 12 (1825).
HEXACTINA, Willd. ex Schult. f. Syst. vii. 91 (1830).
affinis, *Miq. in Linnaea*, xviii. (1844) 613.—Surinam.
africana, Spreng. Syst. ii. 126 = Coffea zanguebarica.
brasiliana, *A. Rich. in Mém. Soc. Nat. Par.* v. (1829) 251? (*Mém. Rub.* 170).—Bras.
corymbosa, *H. B. & K. Nov. Gen. et Sp.* iii. 419. t. 294.—Mexic.
edulis, *Baill. Hist. Pl.* vii. 387.—Guiana.
eriopila, *Baill. l. c.*—Guiana.
fagifolia, *Desf. in Mém. Mus. Par.* vi. (1820) 14. t. 5. —Guiana.
grandifolia, *Miq. in Linnaea*, xviii. (1844) 291.—Guiana.
guianensis, *Aubl. Pl. Gui. Suppl.* 13. t. 375.—Guiana.
hirsuta, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 25. t. 230. —Bras.
intermedia, *Mart. in Schult. f. Syst.* vii. 90.—Bras.
Laureaster, *Mart. in Flora*, xxiv. (1841); *Beibl.* ii. 84. —Bras.
longifolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 26.—Bras.
peruviana, *Desf. in Mém. Mus. Par.* vi. (1820) 16. t. 4.—Peruv.
saccifera, *Mart. in Schult. f. Syst.* vii. 91.—Bras.
surinamensis, *Steud. in Flora*, xxvi. (1843) 763.—Surinam.
utilis, Baill. in Bull. Soc. Linn. Par. i. (1879) 220 = Alibertia edulis.

AMALAGO, Rafin. Sylva Tellur. 84 (1838) = **Piper**, Linn.
antillana, Rafin. l. c. = P. officinarum.
Malamiri, Rafin. l. c. = pracc.

AMALIA, Hort. Hisp. ex Endl. Gen. 183 (1837) = **Tillandsia**, Linn. (Bromel.).
Aeris-incola, Hort. Hisp. ex Steud. Nom. ed. II. i. 69 = T. dianthoidea.

AMALIA, Reichb. Nom. 52 (1841) = **Laelia**, Lindl. (Orchid.).

AMALIAS, Hoffmgg. Preis-Verz. Orch. ex Linnaea, xvi. (1842) Litt. 228 = **Laelia**, Lindl. (Orchid.).
anceps, Hoffmgg. l. c. = Laelia anceps.

AMALOBOTRYA, Kunth, ex Meissn. in DC. Prod. xiv. 185 (1856-57) = **Symmeria**, Benth. (Polygon.).
latifolia, Kunth, ex Meissn. l. c. = S. paniculata.

AMAMELIS, Lem. in Orbign. Dict. iv. 745 (1849) = **Hamamelis**, Gron.

AMANNIA, Blume, Bijdr. 1130 (1826) = **Ammannia**, Houst. (Lythrar.).

AMANOA, Aubl. Pl. Gui. i. 256. t. 101 (1775). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 268.
acuminata, Thw. Enum. Pl. Zeyl. 428 = Cleistanthus acuminatus.
Boiviniana, *Baill. Etud. Gén. Euph.* 582.—Madag.
bracteata, Rich. ex Baill. l. c. 581 = guianensis.
bractcosa, *Planch. in Hook. Ic. Pl.* t. 797.—Afr. trop.
brasiliensis, Baill. Etud. Gén. Euph. 581. n. 4 = Gonatogyne brasiliensis.
chartacea, Baill. l. c. 582 = Cleistanthus chartaceus.
collina, Baill. l. c. = Cleistanthus collinus.
Cunninghamii, Baill. Adansonia, vi. (1865-66) 335 = Cleistanthus Cunninghamii.
Dallachiana, Baill. l. c. = Cleistanthus Dallachyanus.
divaricata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 22. t. 226.—Am. austr.
faginea, Baill. Adansonia, vi. (1865-66) 336 = Briedelia faginea.
ferruginea, Thw. Enum. Pl. Zeyl. 280 = Cleistanthus ferrugineus.
grandiflora, *Muell. Arg. in Flora*, lv. (1872) 2.—Surinam.
guianensis, *Aubl. Pl. Gui.* i. 256. t. 101.—Guiana.
indica, Thw. Enum. Pl. Zeyl. 428 = Cleistanthus robustus.
indica, Wight, Ic. t. 1911 = Cleistanthus patulus.
Leichardtii, Baill. Adansonia, vi. (1865-66) 336 = Briedelia faginea.
oblongifolia, *Muell. Arg. in Linnaea*, xxxii. (1863) 77.—Bras.
ovata, Baill. Adansonia, vi. (1865-66) 336 = Briedelia exaltata.
pallida, Thw. Enum. Pl. Zeyl. 280 = Cleistanthus pallidus.
patula, Thw. l. c. = Cleistanthus patulus.
purpurascens, *Poepp. ex Baill. Etud. Gén. Euph.* 598.—Bras.
racemosa, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 23.—Bras.
ramiflora, Poepp. & Endl. l. c. = Richeria grandis.
robusta, Thw. Enum. Pl. Zeyl. 428 = Cleistanthus robustus.
strobilacea, *Muell. Arg. in Flora*, xlvii. (1864) 515.—Afr. occ.
strobilantha, Planch. in Hook. Niger Fl. t. 47 = bracteosa.
tomentosa, Baill. Adansonia, vi. (1865-66) 336 = Briedelia tomentosa.

AMAPA, Steud. Nom. ed. I. 35 (1821) = **Carapa**, Aubl. (Meliac.).
guianensis, Steud. l. c. = C. guianensis.

AMARABOYA, Linden, ex Mast. in Gard. Chron. (1871) 944. *MELASTOMACEAE.*
amabilis, *Linden, ex Mast. l. c.*—N. Granat.
princeps, *Linden, ex Mast. l. c.*—N. Granat.
splendida, *Linden, ex Mast.* l. c.—N. Granat.

AMARACANTHUS, Steud. Nom. ed. II. i. 69 (1840) sphalm. = seq.

AMARACARPUS, Blume, Bijdr. 954 (1826). *RUBIACEAE*, Benth. & Hook. f. ii. 130.
microphyllus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 211.—Ins. Celebes.
pubescens, *Blume, Bijdr.* 954.—Java.

AMARACUS, Gleditsch, Syst. 189 (1764) = **Origanum**, Linn. (Labiat.).
cordifolius, Montbr. & Auch. ex Benth. in Ann. Sc. Nat. Sér. II. vi. (1836) 43 = O. cordifolium.
Dictamnus, Benth. Lab. Gen. et Sp. 333 = O. Dictamnus.
sipyleus, Rafin. Fl. Tellur. iii. 86 = O. sipyleus.
tomentosus, Moench, Meth. Suppl. 137 = O. Dictamnus.
Tournefortii, Benth. Lab. Gen. et Sp. 333 = O. Tournefortii.

AMARALIA, Welw. ex Benth. & Hook. f. Gen. ii. 90 (1873) = **Sherbournia**, G. Don (Rubiac.).
bignoniaeflora, Welw. l. c. ii. 91 = S. foliosa.
Doniana, Welw. l. c. = S. foliosa.

AMARANTESIA, Hort. ex Regel, Gartenfl. (1869) 102 = **Telanthera**, R. Br. (Amarant.).
brasiliensis, Hort. ex Regel, l. c. = T. Bettzichiana.

AMARANTHUS, Linn. Syst. ed. I. (1735); Gen. ed. I. 286 (1737). *AMARANTACEAE*, Benth. & Hook. f. iii. 28.
ALBERSIA, Kunth, Fl. Berol. ed. II. 144 (1838).
AMBLOGYNA, Rafin. Fl. Tellur. iii. 42 (1836).
BAJAN, Adans. Fam. ii. 506 (1763).
BLITON, Adans. l. c. (1763).
BLITUM, Linn. Gen. ed. I. 20 (1737).
DIMEIANDRA, Rafin. Neogenyt. 2 (1825).
DIMEIANTHUS, Rafin. Fl. Tellur. iii. 41 (1836).
EUXOLUS, Rafin. l. c. 42 (1836).
GLOMERARIA, Cav. Descr. 319 (1827).
MENGEA, Schau. in Nov. Act. Cur. xix. Suppl. I. (Pl. Meyen.) 405 (1843).
PENTRIUS, Rafin. Fl. Tellur. iii. 42 (1836).
PYXIDIUM, Moq. Tand. in DC. Prod. xiii. II. 262 (1849) (vox fructus desc. lapsu progenere habita).
ROEMERIA, Moench, Meth. 341 (1794).
SARRATIA, Moq. ex DC. Prod. xiii. II. 233, 268 (1849).
SCLEROPUS, Schrad. Ind. Sem. Hort. Goett. (1835).

Acroglochin, Spreng. Syst. i. 927 = Acroglochin chenopodioides.
adscendens, Auct. vide *ascendens*.
aeneus, *Bess. Cat. Hort. Volhyn.* (1816) 8.—Hab.?
aeruoides, Hochst. & Steud. ex A. Rich. Tent. Fl. Abyss. ii. 214 = Aerva lanata.
albiflorus, *Moq. in DC. Prod.* xiii. II. 263.—Hab.?
albus, *Linn. Syst.* ed. X. 1268.—Am. bor. (Europ. alien.).
albus, Robin, Voy. Louis. 374; ex Moq. in DC. Prod. xiii. II. 265 = leucanthus.
albus, Rodsch. ex F. G. Dietr. Lex. Gaertn. Nachtr. II. i. 196 = A. Blitum.
albus, Thunb. Fl. Cap. ed. Schult. i. 215 = Thunbergii.
Alopecurus, Hochst. ex A. Br. & Bouché, Ind. Sem. Hort. Berol. App. (1872) 1 = caudatus.
altissimus, Riddell, Syn. Fl. West. St. (1835) 41 = Acnida rusocarpa.
amboinicus, Buch.-Ham. in Wall. Cat. n. 6987 = gangeticus.
Anacardana, Hook. f. Fl. Brit. Ind. iv. 719 = seq.
Anardana, Buch.-Ham. in Wall. Cat. n. 6903 = paniculatus.
angustifolius, Bieb. ex Willd. Sp. Pl. iv. 381 = sylvestris.
angustifolius, Lam. Encyc. i. 115 = A. Blitum.
Arardhanus, Sweet, Hort. Brit. ed. III. 569 = paniculatus.
arctioideus, Perr. ex Moq. in DC. Prod. xiii. II. 332 = Pupalia lappacea.
ascendens, *Hornem. Hort. Hafn. Suppl.* 107.—China.
ascendens, Loisel. Not. 141 = viridis.
atropurpureus, *Roxb. Fl. Ind.* iii. 608.—Ind. or.
atrosanguineus, Hort. Lugd. (1838) ex Moq. in DC. Prod. xiii. II. 266 = hypochondriacus?
aureus, Bess. Cat. Hort. Volhyn. (1816) 8 = flavus?
aureus, *F. G. Dietr. Lexik. Gaertn.* i. 310.—Nepal.
aureus, Hort. ex Moq. in DC. Prod. xiii. II. 259 = hybridus.
bahiensis, *Mart. Herb. Fl. Bras.* n. 969.—Bahia.
Bellardi, Hort. ex Moq. in DC. Prod. xiii. II. 259 = hybridus.
Berchtholdi, Hort. ex Moq. l. c. 257 = paniculatus.
Berchtoldi, Seidl, ex Opiz, Boeheims Gew. 164; Natural. i. (1823) [2] = A. Blitum.
Berchtholdi, Hort. ex Moq. in DC. Prod. xiii. II. 259 = hybridus.
Bernhardi, Hort. ex Moq. l. c. 258 = flavus.
bicolor, Nocca, ex Willd. Sp. Pl. iv. 384 = melancholicus.
blitoides, *S. Wats. in Proc. Am. Acad.* xii. (1877) 273.—Am. bor.
blitonius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Blitum.
Blitum, Auct. ex Moq. in DC. Prod. xiii. II. 267, 274 = viridis.
Blitum, *Linn. Sp. Pl.* ed. I. 990.—Reg. temp. et trop.
Blitum, Moq. in DC. Prod. xiii. II. 263 = sylvestris.
Blitum, Reichb. Ic. ex Steud. Nom. ed. II. i. 69 = polygonoides.
brasiliensis, *Moq. in DC. Prod.* xii. II. 266.—Bras.
bullatus, Bess. ex Spreng. Syst. i. 928 = retroflexus.
californicus, *S. Wats. in Bot. Calif.* ii. 42.—Calif.
campestris, *Willd. Sp. Pl.* iv. 382.—Ind. or.
canariensis, *Bess. Cat. Hort. Volhyn.* (1816) 8.—Ins. Canar.

AMARANTHUS:—
capitatus, Cat. Hort. Turic. (1827–28).—Hab.?
Caracam, Bess. Cat. Hort. Volhyn. (1816) 8 = A. Cararu.
caracasanus, *H. B. & K. Nov. Gen. et Sp.* ii. 195.—Venezuela.
Caracu, Zucc. ex Steud. Nom. ed. II. i. 69 = A. Cararu.
Cararia, Bess. Cat. Hort. Volhyn. (1816) 8 = praec.?
Cararu, Hort. Paris. ex Moq. in DC. Prod. xiii. II. 274 = caudatus.
Cararu, *Jacq. ex Zuccag. in Roem. Collect.* 158.—Hab.?
carneus, Hort. ex Moq. in DC. Prod. xiii. II. 257 = paniculatus.
Cathecu, Hort. ex Moq. l. c. 259 = hybridus.
Caturus, *Roxb. in Herb. Madr. sec. Wall. Cat.* n 6900.—Ind. or.
caudatus, *Linn. Sp. Pl.* 990.—Oriens; Afr. trop.; Ind. or.
cauliflorus, Link, Enum. Hort. Berol. ii. 389 = Acroglochin chenopodioides.
celosioides, *H. B. & K. Nov. Gen. et Sp.* ii. 194.—N. Granat.
cernuus, Bess. in Cat. Hort. Turic. (1827–8) 'An Celosia?'
chlorostachys, Moq.-Tand. in DC. Prod. xiii. II. 259 = paniculatus.
chlorostachys, *Willd. Amarant.* 34. t. 10. f. 19.—Europ.; Oriens; Ind. or.
chlorostachys, Willk. ex Willk. & Lange, Prod. Fl. Hisp. i. 275 = retroflexus.
circinnatus, Hort. Paris. ex Poir. Encyc. Suppl. i. 311 = lividus?
coesius, *F. G. Dietr. ex Moq. in DC. Prod.* xiii. II. 266.—Hab.?
commutatus, *Kerner, in Oestr. Bot. Zeitschr* xxv. (1875) 194.—Austr.
coracanus, Mart. Hort. Erlang. 197 = caracasanus.
crassipes, Schlecht. in Linnaea, vi. (1831) 757 = polygonoides.
cristatus, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 7 = Celosia cristata.
crocatus, Bess. Cat. Hort. Volhyn. (1816) 9.—Hab.?
cruentus, Linn. Syst. ed. X. 1269 = paniculatus.
cruentus, Willd. ex Roxb. Fl. Ind. iii. 610 = caudatus.
curvifolius, Spreng. Syst. i. 928 = retroflexus.
cuspidatus, Vis. Cat. Hort. Patav. (1841) = gangeticus.
deflexus, *Linn. Mant.* ii. 295.—Europ.
Delilei, Richter & Loret, in Bull. Soc. Bot. Fr. xiii. (1866) 316 = patulus.
diacanthus, *Rafin. Fl. Ludov.* 31.—Am. bor.
diandrus, Spreng. Neue Entdeck. iii. 20 = Acroglochin chenopodioides.
diffusus, Dulac, Fl. Hautes-Pyr. 174 = A. Blitum.
dioicus, Michx. ex Moq. in DC. Prod. xiii. II. 277 = Acnida rusocarpa.
divaricatus, *Andrz. ex Lindem. Prod. Fl. Cherson.* 185.—Cherson.
dubius, Mart. Hort. Erlang. (1814) 197 = tristis.
edulis, Michx. ex Moq. in DC. Prod. xiii. II. 277 = Acnida cannabina.
emarginatus, *Salzm. ex Moq. l. c.* 274 = viridis.
enervis, *F. Muell. ex Benth. Fl. Austral.* v. 216.—Austral.
esculentus, Bess. Cat. Hort. Volhyn. (1816) 9; ex Moq. in DC. Prod. xiii. II. 266 = paniculatus.
farinaceus, Roxb. ex Moq. l. c. = paniculatus.
fasciatus, Roxb. Fl. Ind. iii. 609 = viridis.
Fatine, Mart. Hort. Erlang. 197.—Hab.?
fimbriatus, *S. Wats. in Bot. Calif.* ii. 42.—Calif.
flavescens, Hort. ex Moq. in DC. Prod. xiii. II. 259 = hybridus.
flavus, *Linn. Syst.* ed. X. 1269.—Ind. or.
flexuosus, Hort. ex Moq. in DC. Prod. xiii. II. 260 = tristis.
floridus, Benth. Bot. Voy. Sulph. 158. t. 51 = scariosus.
frumentaceus, Buch.-Ham. in Roxb. Fl. Ind. iii. 699 = paniculatus.
frutescens, Hort. ex Moq. in DC. Prod. xiii. II. 348 = Iresine amarantoides.
gangeticus, *Linn. Syst.* ed. X. 1268.—Reg. trop.
gangeticus, Wall. Cat. n. 6896 = A. Blitum.
giganteus, Bess. Cat. Hort. Volhyn. (1816) 91; F. G. Dietr. Lexik. Gaertn. i. 313.—Hab.?
giganteus, L'Hérit. ex Moq. in DC. Prod. xiii. II. 277 = Acnida cannabina.
gracilis, Desf. Tabl. Hort. Par. ed. I. 43 = caudatus.
graecizans, Auct. Hisp. ex Cut. Fl. Comp. Madr. 579 = albus.
graecizans, Linn. Sp. Pl. 990 = sylvestris.

AMARANTHUS:—
Greggii, *S. Wats. in Proc. Am. Acad.* xii. (1877) 274.—Am. bor. occ.
guadelupensis, Hort. ex Moq. in DC. Prod. xiii. II. 257 = paniculatus.
hecticus, Willd. Amaranth. 25 = hybridus.
hierichuntinus, *Vis. in Att. Ist. Ven. Sc.* Ser. III. iv. (1858–59) 139.—Syria.
Huttoni, Hort. Veitch. cf. Gard. Chron. (1872) 215 = Celosia Huttoni.
hybridus, *Linn. Sp. Pl.* 990.—Am. bor.
hybridus, Vell. Fl. Flum. x. t. 28 = flavus?
hypochondriacus, *Linn. Sp. Pl.* 991.—Am. bor. (Europ. alien.).
inamoenus, Willd. Sp. Pl. iv. 386 (1805) Amar. 14 = gangeticus.
incarnatus, Hort. ex Moq. in DC. Prod. xiii. II. 257 = paniculatus.
incomptus, Willd. Enum. Hort. Berol. 64 = gangeticus.
incurvatus, Gren. & Godr. Prosp. Fl. Fr. 8 = chlorostachys.
intermedius, Guss. ex Moq. in DC. Prod. xiii. II. 259 = chlorostachys.
interruptus, *R. Br. Prod.* 414.—Austral.
japonicus, Houtt. ex Steud. Nom. ed. I. 36 = gangeticus.
lacteus, Hort. Amstelod. (1814) 52 = Aerva sanguinolenta.
laetus, Willd. Amaranth. 28; Sp. Pl. iv. 391 (1805) = hybridus.
lanatus, Dum.-Cours. Bot. Cult. ed. I. i. 640 = Aerva lanata.
lancefolius, Roxb. Hort. Beng. 67 = gangeticus.
lanceolatus, Roxb. Fl. Ind. iii. 607 = gangeticus.
lancifolius, *Del. ex Moq. in DC. Prod.* xiii. II. 276 = interruptus.
laxiflorus, Comell. ex Poll. Fl. Veron. iii. 114 = patulus.
leptostachyus, *Benth. Fl. Austral.* v. 214.—Austral.
leucanthus, *Rafin. Fl. Ludov.* 32.—Am. bor.
leucocarpus, *S. Wats. in Proc. Am. Acad.* x. (1875) 347.—Arizona.
lineatus, R. Br. Prod. 414 = interruptus.
littoralis, Bernh. in Hort. Tur. (1813) ex Moq. in DC. Prod. xiii. II. 274 = viridis.
littoralis, Hornm. Hort. Hafn. (1815) 893 = albus.
lividus, Hort. Petrop. ex Hook. f. Fl. Brit. Ind. iv. 721 = A. Blitum.
lividus, *Linn. Sp. Pl.* 990.—Am. bor.
lividus, Roxb. Fl. Ind. iii. 605 = gangeticus.
macrocarpus, *Benth. Fl. Austral.* v. 216.—Austral.
macrocaulos, Poir. Encyc. Suppl. i. 314 = Acnida cannabina.
macrostachyus, Merat, ex Moq. in DC. Prod. xiii. II. 256 = hypochondriacus.
major, Salzm. ex Moq. in DC. Prod. xiii. II. 274.—Hab.?
mangostanus, Blanco, Fl. Filip. ed. I. 711 = melancholicus.
mangostanus, *Linn. Cent.* i. 32; *Am. Acad.* iv. 294.—Ind. or.
Mantegazzianus, *Passer. in Hort. Parm.* (1864) 4.—Am. austr.
maximus, Mill. Gard. Dict. ed. VIII. n. 5 = caudatus.
melancholicus, Linn. Sp. Pl. 989 = gangeticus.
miamiensis, *Riddell, Syn. Fl. West. St.* (1835) 41.—Am. bor.
minor, S. F. Gray, Nat. Arr. Brit. Pl. ii. 289 = A. Blitum.
minutus, Leschen. ex Moq. in DC. Prod. xiii. II. 304 = Aerva brachiata.
Mitchelli, *Benth. Fl. Austral.* v. 214.—Austral.
monstrosus, Cat. Hort. Div. (1837); Hort. Tonelle, ex Moq. in DC. Prod. xiii. II. 256 = hypochondriacus.
montevidensis, Hort. ex Moq. l. c. 257 = paniculatus.
morosus, *Reichb. Fl. Germ. Excurs.* 585.—Europ.
mucronatus, Hort. Petrop. ex Hook. f. Fl. Brit. Ind. iv. 720 = mangostanus.
mucronatus, Poir. Encyc. Suppl. i. 311 = A. Blitum.
muricatus, *Gill. ex Moq. in DC. Prod.* xiii. II. 276.—Hab.?
neglectus, Hort. ex Moq. l. c. 259 = hybridus.
nepalensis, Cat. Hort. Lugd. (1836); ex Moq. l. c. = hybridus.
obovatus, *S. Wats. in Proc. Am. Acad.* xii. (1877) 275—Am. bor.
officinalis, Gromow. ex Trautv. in Act. Hort. Petrop. ix. (1884) 139 = seq.?
oleraceus, *Linn. Sp. Pl.* 1403.—Egypt.; Ind. or.
oleraceus, Rodsch. Essequib. 24 = A. Blitum.
oleraceus, Roxb. Fl. Ind. iii. 605 = gangeticus.
olitorius, Bess. Cat. Hort. Volhyn. (1816) 9 = oleraceus?

AMARANTHUS :—
pachystachys, Reichb. ex Moq. in DC. Prod. xiii. II. 265 = parisiensis.
pallidiflorus, *F. Muell. Fragm.* i. 140.—Austral.
pallidus, Bieb. Fl. Taur. Cauc. ii. 399 = A. Blitum.
Palmeri, *S. Wats. in Proc. Am. Acad.* xii. (1877) 274. —Calif.
panieulatus, *Linn. Sp. Pl.* ed. II. 1406.—Am. bor. (in Orient. alien.).
paniculatus, Wall. Cat. n. 6904 = caudatus.
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 273; *Contrib.* 79.—Reg. Argent.
parisiensis, *Schk. Handb.* iii. 249.—Hab. ?
patulus, *Bertol. Comm. Neap.* 171. t. 12.—Am. bor. (in Europ. alien.).
pendulinus, Hort. ex Moq. in DC. Prod. xiii. II. 255 = caudatus.
pendulus, Hort. ex Moq. l. c. = caudatus.
persicarioides, Hort. ex Poir. Encyc. Suppl. i. 311 = Aerogloehin chenopodioides.
polychroa, Raeusch. Nom. ed. III. 275 = Pellionia brevifolia ?
polyflagellus, *Spreng. in Hort. Argent.* (1834) *ex Moq. in DC. Prod.* xiii. II. 266.—Hab. ?
polygamus, *Linn. Cent.* i.; *Am. Acad.* iv. 294.—Afr. trop.; Ind. or.
polygamus, Wall. Cat. n. 514 = A. Blitum.
polygamus, Thw. Enum. Pl. Zeyl. 247 = mangostanus.
polygamus, Roxb. Fl. Ind. iii. 607 = gangeticus.
polygonoides, *Linn. Pl. Jam. Pugill.* ii. 27.—Jamaic.; Zeyl.
polygonoides, Zolling. ex Moq. in DC. Prod. xiii. II. 274 = A. Blitum.
polygonoides, Roxb. Fl. Ind. iii. 721 = polygamus.
polystachyus, Buch.-Ham. in Wall. Cat. n. 6901 = viridis.
polystaehyus, *Willd. Sp. Pl.* iv. 385.—Ind. or.
Powellii, *S. Wats. in Proc. Am. Acad.* x. (1875) 347. —Am. bor.
prostratus, Balb. Mise. Bot. 44. t. 10 = deflexus.
prostratus, Bast. Fl. Maine-et-Loire, 344 = A. Blitum.
pumilus, *Rafin. in Med. Repos. New York*, v. (1808) 360.—Am. bor.
purgans, Hort. ex Moq. in DC. Prod. xiii. II. 257 = panieulatus.
pyramidalis, Noronha, in Verh. Batav. Gen. v. (1790), ed. I. Art. iv. 7 = Celosia argentea.
quitensis, *H. B. & K. Nov. Gen. et Sp.* ii. 194.—Peruv.
recurvatus, Desf. Cat. Hort. Par. ed. III. (1829) 39 = retroflexus.
retroflexus, *Linn. Sp. Pl.* 991.—Am. bor. (in Europ. alien.).
rhombeus, R. Br. Prod. 414 = interruptus.
rigidus, Sehult. ex Steud. Nom. ed. II. i. 70 = retroflexus.
rotundifolius, Herb. Par. ex Moq. in DC. Prod. xiii. II. 261 = gangeticus.
rubescens, Hort. ex Moq. l. e. 257 = panieulatus.
rubrieaulis, *Page; Hort. Angl. ex Moq. in DC. Prod.* xiii. II. 267, nomen.—Ind. or.
ruderalis, Koch, ex Moq. l. c. 274 = A. Blitum.
salicifolius, *Hort. Veitch, ex Gard. Chron.* (1871) 1550. —Ins. Philipp.
sanguineus, Linn. Sp. Pl. ed. II. 1407 = panieulatus.
sanguincus, *Vell. Fl. Flum.* x. t. 29.—Bras.
sanguinolentus, Schrad. ex Moq. in DC Prod. xiii. II. 267 = paniculatus ?
scandens, *Linn. f. Suppl.* 419.—Am. austr.
scariosus, *Benth. Bot. Voy. Sulph.* 158. t. 51.—Guatemala.
spathulatus, Desf. ex Moq. in DC. Prod. xiii. II. 276 = lineatus.
speciosus, Sims, Bot. Mag. t. 2227 = panieulatus.
spicatus, Lam. Fl. Fr. ii. 192 = retroflexus.
spicatus, Wirzen, Diss. Pl. Casan. 454 = panieulatus.
spicatus, Reichb. Fl. Germ. Excurs. 585 = patulus.
spinosus, *Linn. Sp. Pl.* 991.—Reg. trop.
spiratus, *Zipp. ex Span. in Linnaea*, xv. (1841) 345, nomen.—Malaya.
strictus, Tenore, Syll. 127 = retroflexus.
strictus, Willd. Amaranth. 27 = paniculatus.
sylvestris, Desf. Tabl. Hort. Par. ed. I. 44.—Europ.; Oriens.
tenuiflorus, Fisch. Hort. Hal. ex Moq. in DC. Prod. xiii. II. 267 = seq. ?
tenuifolius, Roxb. Fl. Ind. iii. 602 = A. Blitum.
tenuifolius, *Willd. Sp. Pl.* iv. 381.—Ind. or.
tenuifolius, Wall. Cat. n. 6893 = polygamus.
tenuis, *Benth. Fl. Austral.* v. 216.—Austral.

AMARANTHUS :—
Thunbergii, *Moq. in DC. Prod.* xiii. II. 262.—Afr. austr.
Timeroyi, Jord. ex Moq. l. e. 259 = patulus.
Torreyi, *Benth. ex S. Wats. Bot. Calif.* ii. 42.—Am. bor.
tortuosus, *Hornem. Hort. Hafn. Suppl.* 107.—Ins. S. Cruee.
tricolor, Linn. Sp. Pl. 989 = gangeticus.
tristis, Linn. l. e. = gangeticus.
tristis, Wall. Cat. n. 6905 = mangostanus.
trivialis, Rota, in Giorn. Bot. Ital. ii. II. (1846) 287 = patulus.
undulatus, Lindl. in Miteh. Trop. Austr. 102 = Mitehelli ?
undulatus, R. Br. Prod. 414 = interruptus.
ureeolatus, *Benth. in Bot. Voy. Sulph.* 158.—Guayaquil.
venulosus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 376.—Am. bor.
vernus, *Opiz, ex Moq. in DC. Prod.* xiii. II. 267, nomen. —Europ.
verticillatus, Pav. ex Moq. in DC. Prod. xiii. II. 270 = polygonoides.
violaceus, Hort. ex Moq. l. c. = panieulatus.
viridis, All. Fl. Pedem. ii. 19 = A. Blitum.
viridis, *Linn. Sp. Pl.* ed. II. 1405.—Reg. trop.
viridis, Pall. ex Steud. Nom. ed. II. i. 36 = pallidus.
viridis, Pollieh, ex Moq. in DC. Prod. xiii. II. 268 = prostratus.
Wrightii, *S. Wats. in Proc. Am. Acad.* xii. (1877) 275.—Am. bor.
zanensis, *Hornm. ex Moq. in DC. Prod.* xiii. II. 267.—Hab. ?

AMARANTUS, Linn. Syst. ed. x. 1268 (1759) = **Amaranthus**, Linn.

AMARELLA, Gilib. Fl. Lituan. i. 36 (1781) = **Gentiana**, Linn.
acuta, Rafin. Fl. Tellur. iii. 21 = G. Amarella.
brevicaulis, Rafin. l. e.—Hab. ?
gracilis, Rafin. l. e. = G. Amarella.
quadrifida, Gilib. Fl. Lituan. i. 36 = G. eampestris.
quinquefida, Gilib. l. e. = G. Amarella.
rugosa, Rafin. Fl. Tellur. iii. 21.—Hab. ?
vulgaris, Rafin. l. e. 20 = G. eampestris.

AMARENUS, C. Presl, Symb. Bot. i. 46 (1830) = **Trifolium**, Linn. (Legum.).
agrarius, C. Presl, l. c. = T. agrarium.
aureus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 362 = T. agrarium.
badius, C. Presl, l. c. = T. badium.
campestris, Fourr. l. c. = T. procumbens.
filiformis, C. Presl, l. c. = T. filiforme.
flavus, C. Presl, l. c. = T. flavum.
micranthus, Fourr. l. e. = T. filiforme.
patens, C. Presl, l. e. = T. patens.
procumbens, C. Presl, l. e. = T. proeumbens.
Schreberi, Fourr. l. e. = T. proeumbens.
Sebastiani, C. Presl, l. c. = T. Sebastiani.
spadiceus, C. Presl, l. c. = T. spadiceum.
speciosus, C. Presl, l. c. = T. speeiosum.

AMARIA, Mutis, ex Caldas, in Seman. Nuev. Gran. ii. (1810) 25 = **Bauhinia**, Linn. (Legumin.).
petiolata, Mutis, l. e. = B. petiolata.
sessilifolia, Mutis, l. e.—N. Granat.

AMARIDIUM, Hort. ex Lubbers, Cat. Pl. Rar. San Donato, 15 (1880) = **Camaridium**, Lindl. (Orehid.).

AMARORIA, A. Gray, Bot. U. St. Expl. Exped. i. 356. t. 40 (1854). *SIMARUBEAE*, Benth. & Hook. f. i. 314.
soulameoides, A. Gray, l. e.—Ins. Fiji.

AMARYLLIS, Linn. Syst. ed. I. (1735). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 727.
ATAMOSCO, Adans. Fam. ii. 57 (1763).
BELLADONNA, Sweet, Hort. Brit. ed. II. 506. (1830).
CALLICORE, Link, Handb. i. 193 (1829).
CALLIROË, Endl. Gen. 176 (1837).
acuminata, Ker-Gawl. in Bot. Reg. t. 534 = Hippeastrum rutilum.
advena, Ker-Gawl. l. e. t. 1125 = Hippeastrum advenum.
aethnensis, 'Rafin.' vide *etnensis*.
africana, Lam. Encyc. i. 124 = Lycoris aurea.
Agatha, Steud. Nom. ed. II. i. 72 = H. reticulatum.
alba, Forsk. Fl. Aegypt. Arab. 208 = Crinum album.

AMARYLLIS :—
Alberti, Lem. Illustr. Hortic. (1866) t. 498 = Hippeastrum Reginae.
alta, Steud. Nom. ed. II. i. 72 = Hippeastrum splendens.
Andersonii, Steud. l. c. i. 71 = Zephyranthes Andersoni.
andicola, Poepp. Fragm. Syn. 5 (= *Habranthus andicolus*).
angusta, Sehult. f. Syst. vii. 807 = Hippeastrum bifidum.
Aramasco, Linn. Sp. Pl. ed. II. 420 (sphalm.) = Zephyranthes Atamaseo.
arvensis, F. G. Dietr. Lexie. Gaertn. Naehtr. i. 204 = peruviana.
Atamasca, Linn. Sp. Pl. 292 = Zephyranthes Atamasco.
Atamasco, Blanco, Fl. Filip. ed. I. 254 = Hippeastrum miniatum.
atanasia, Crantz, Inst. i. 486 = Zephyranthes Atamasea.
aucta, Tratt. Thes. 45 = Nerine undulata.
aulica, Ker-Gawl. in Journ. Se. & Arts, ii. (1817) 353; et in Bot. Reg. t. 444 = Hippeastrum aulicum.
aurea, L'Hérit. Sert. Angl. 14 = Lycoris aurea.
aurea, Ruiz & Pav. Fl. Per. iii. 56 (= *Pyrolirion aureum*).
australasica, Ker-Gawl. in Bot. Reg. t. 426 = Crinum flaecidum.
australis, Spreng. Syst. ii. 54 = praec.
Bagnoldi, D. Dietr. Syn. Pl. ii. 1177 = Hippeastrum Bagnoldi.
bahiensis, DC. Pl. Rar. Jard. Genève, 7me Notiee, 18 = Hippeastrum bahiense.
Bangoldi, Steud. Nom. ed. II. i. 70, sphalm. = *Bagnoldi*.
Banksiana, Lindl. Bot. Reg. (1842), t. 11. & Append. 18 = Brunsvigia Slateriana.
Belladonna, *Linn. Sp. Pl.* 293.—Afr. austr.
Belladonna, E. Mey. ex Steud. Nom. ed. II. i. 70 = Hippeastrum equestre.
Berterii, Spreng. Syst. ii. 49 (= *Habranthus robustus*).
bicolor, Ruiz & Pav. Fl. Per. iii. 57 = Hippeastrum bieolor.
bifida, Spreng. Syst. Cur. Post. 133 = Hippeastrum bifidum.
bifolia, Aubl. Pl. Gui. i. 304 = Zephyranthes rosea.
bivaginata, Donn, Hort. Cantab. ed. VIII. 98 = Cyrtanthus pallidus.
blanda, Ker-Gawl. in Bot. Mag. t. 1450 = A. Belladonna.
brasiliensis, Andr. Bot. Rep. t. 358 = Hippeastrum equestre, Reginae.
brasiliensis, Tratt. Tabul. t. 333 = Hippeastrum rutilum.
Broussonetii, [DC. in] Red. Liliae. ii. 62 = Crinum yueeaeflorum.
bulbisperma, Burm. f. Fl. Cap. Prod. 9 = Crinum eapense.
caerulea, *Griseb. in Goett. Abh.* xxiv. (1879) 320.—Reg. Argent.
calyptrata, *Hoffmgg. in Linnaea*, xvi. (1842) *Litt.* 27. —Hab. ?
calyptrata, Ker-Gawl. Rev. 11; et in Journ. Sc. & Arts, ii. (1817) 353 = Hippeastrum calyptratum.
candida, Lindl. in Bot. Reg. t. 724 = Zephyranthes candida.
candida, Tratt. Tabul. t. 488 = Crinum giganteum.
capensis, Linn. Pl. Afr. 10 ? (Amoen. Acad. vi. 86) = Hypoxis stellatus.
capensis, Mill. Gard. Dict. ed. VIII. n. 12 = Crinum longifolium.
carinata, Spreng. Syst. Cur. Post. 152 = Zephyranthes carinata.
carnea, Sehult. f. Syst. vii. 799 = Zephyranthes rosea.
Carolinae, Sterler, Hort. Nymph. (1826) 44 = Hippeastrum retieulatum.
caspia, Willd. Sp. Pl. ii. 62 = Allium caspium.
chilensis, L'Herit. Sert. Angl. 11 = Hippeastrum chileuse.
chilensis, Ruiz & Pav. Fl. Per. iii. 56 = Hippeastrum miniatum.
chloroleuca, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 347. t. 8. f. 1 = Zephyranthes chloroleuca.
ciliaris, Linn. Sp. Pl. ed. II. 422 = Buphane ciliaris.
cinnamomea, L'Hérit. Sert. Angl. 15 = Hessea crispa.
citrina, Sibth. & Sm. Fl. Graec. iii. t. 311 = Sternbergia colchiciflora.
clavata, L'Herit. Sert. Angl. 11 = Cyrtanthus unifloras.
Clusiana, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 346 = Sternbergia colchiciflora.

AMARYLLIS:—
coccinea, Molina, Saggio Chil. ed. II. 284 = chilensis.
colchiciflora, Hort. Mosq. ex Schult. f. Syst. vii. 1135 = Bulbocodium versicolor.
colchiciflora, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 345 = Sternbergia colchiciflora.
concinna, *Morris, Fl. Conspic.* t. 44.—Hab. ?
coracina, Steud. Nom. ed. II. i. 70 = seq.
coranica, Ker-Gawl. in Bot. Reg. ii. 139 = Ammocharis falcata.
correiensis, Bury, Hexand. Pl. 9 = Hippeastrum organense.
corusca, Ker-Gawl. in Bot. Mag. fol. 1430 = Nerine curvifolia.
crispa, Jacq. Hort. Schoenb. i. 37 = Hessea crispa.
crocata, Ker-Gawl. in Bot. Reg. t. 38 = Hippeastrum rutilum.
curvifolia, Jacq. Hort. Schoenb. i. 33 = Nerine curvifolia.
Cybister, Herb. ex Fl. des Serres, v. t. 455, 456 = Hippeastrum Cybister.
cylindracea, L'Hérit. Sert. Angl. 15 = Cyrtanthus angustifolius.
cyrtanthoides, Sims, Bot. Mag. t. 2399 = Hippeastrum bicolor.
demissa, Steud. Nom. ed. II. i. 71.—Chili.
depauperata, Poepp. Fragm. Syn. 4 = Zephyranthes depauperata.
disticha, Sims, Bot. Mag. sub t. 1943 = Crinum yuccaeflorum.
disticha, Linn. f. Suppl. 195 = Buphane disticha.
Drummondii, Steud. Nom. ed. II. i. 71 = Cooperia Drummondii.
Dryades, Vell. Fl. Flum. 130; iii. t. 117 = Griffinia Dryades.
dubia, Houtt. Plantenk. xii. 181. t. 84. f. 1 = Nerine sarniensis.
dubia, Linn. Diss. Pl. Surin. n. 39; et Am. Acad. viii. 254 = Hippeastrum equestre.
elata, Jacq. Hort. Schoenb. i. 32 = Vallota purpurea.
elegans, Spreng. Pugill. ii. 59 = Hippeastrum solandraeflora.
Entreriana, Hoffmgg. in Linnaea, xliii. (1880–82) 137 = Zephyranthes mesochloa.
equestris, Ait. Hort. Kew. i. 417 = Hippeastrum equestre.
etnensis, Rafin. Caratteri, 84 = Sternbergia colchiciflora.
exigua, Schousb. Vextr. Marokko, 160 = Tapeinanthus humilis.
falcata, L'Hérit. Sert. Angl. 13 = Ammocharis falcata.
falcata, Lee, ex Steud. Nom. ed. II. i. 73, 475 = Cyrtanthus angustifolius.
flaccida, Weinm. in Flora, iii. (1820) 608 = Crinum flaccidum.
flammea, Ruiz & Pav. Fl. Per. iii. 56 (= *Pyrolirion flammeum*).—Peruv.; Chili.
flava, Pav. ex Roem. & Schult. Syst. vii. 805 = Zephyranthes flava.
flavovirens, Schott, ex Steud. Nom. ed. II. i. 71.—Cf. *fulvovirens*.
flexuosa, Jacq. Hort. Schoenb. i. 35 = Nerine flexuosa.
Forbesii, Lindl. in Trans. Hort. Soc. vi (1826) 87 = Crinum Forbesianum.
formosa, Salisb. Prod. 234 = seq.
formosissima, Linn. Sp. Pl. 293 = Sprekelia formosissima.
Fothergillia, Andr. Bot. Rep. iii. 163 = Nerine curvifolia.
Fothergillii, Poir. Encyc. Suppl. i. 320 = praec.
fulgida, Ker-Gawl. in Bot. Reg. t. 226 = Hippeastrum rutilum.
fulvovirens, Schott, in Schreib. Nachr. Oesterr. Naturf. 129 = Hippeastrum calyptratum.
Gardneri, Seub. in Mart. Fl. Bras. iii. 149 = Hippeastrum organense.
gigantea, Ait. Hort. Kew. ed. II. ii. 226 = Crinum giganteum.
gigantea, Van Marum, in Verh. Maats. Haarl. iii. (1806) 345 = Brunsvigia Josephinae.
glaucescens, Mart. in Schult. f. Syst. vii. 813 = Hippeastrum rutilum.
gracilifolia, Schult. f. l. c. 806 (= *Habranthus gracilifolius*).
gracilis, Spreng. Syst. Cur. Post. 133 = praec.
grandiflora, D. Dietr. Syn. Pl. ii. 1181 = Brunsvigia grandiflora.
grandiflora, Steud. Nom. ed. II. i. 73 = Sprekelia formosissima.

AMARYLLIS:—
guttata, Linn. Sp. Pl. 294 = Buphane ciliaris.
haemanthoidea, Tratt. Gartenpfl. 38 = Haemanthus amarylloides.
humilis, Jacq. Hort. Schoenb. i. 36 = Nerine humilis.
humilis, Link, Enum. i. 313 = Cyrtanthus uniflorus.
hyacinthina, Ker-Gawl. in Bot. Reg. t. 163 = Griffinia hyacinthina.
ignea, Lindl. Bot. Reg. t. 809 = Hippeastrum bicolor.
ignescens, Regel, Gartenfl. xxiii. (1874) 150 = Hippeastrum ignescens.
illustris, Vell. Fl. Flum. 131; iii. t. 118 = Hippeastrum psittacinum.
insignis, Ker-Gawl. Bot. Reg. t. 579 = Crinum latifolium.
intermedia, Lindl. Bot. Reg. t. 1148 (= *Habranthus intermedius*).
intermedia, Spreng. Syst. iv. Cur. Post. 133 = Griffinia intermedia.
Jacquinii, Tratt. Gartenpfl. 43 = Nerine sarniensis.
Jagus, Thomps. Bot. Displ. t. 6 = Crinum giganteum.
Josephinae, [Del.] Red. Lil. vii. 370 = Brunsvigia Josephinae.
Josephiniana, Herb. Amar. 278 = praec.
Karwinskii, Zucc. in Otto & Dietr. Allg. Gartz. ii. (1834) 245 = Sprekelia formosissima.
kermesiana, Lindl. in Bot. Reg. t. 1638 = Hippeastrum bifidum.
lateritia, *A. Dietr. in Otto & Dietr. Allg. Gartz.* xviii. (1850) 65.—Guinea.
laticoma, Ker-Gawl. in Bot. Reg. vi. 497 = Nerine lucida.
latifolia, Lam. Encyc. i. 124 = Crinum giganteum.
latifolia, L'Hérit. Sert. 14 = Crinum ornatum.
leonensis, *Van Houtte, Hort. Vanhoutt. fasc.* i. (1845) 5.—Afr. trop.
Leopoldii, Hort. Veitch. in Gard. Chron. (1870) 733 = Hippeastrum Leopoldi.
Lindleyana, Schult. f. Syst. vii. 802 = Zephyranthes carinata, Lindleyana.
linearifolia, Molina, Chile, ed. II. 284 = Hippeastrum chilense.
linearis, Willd. Sp. Pl. ii. 53 = Crinum lineare.
lineata, Colla, Hort. Rip. 3 = Hippeastrum vittatum.
lineata, Lam. Encyc. i. 123 = Crinum zeylanicum.
littoralis, Salisb. Prod. 230 = Crinum latifolium.
liturata, *Reichb. Ic.* t. 82.—Hab. ?
longa, Hort. ex Schult. f. Syst. vii. 854 = Crinum yuccaeflorum.
longiflora, Steud. Nom. ed. II. i. 70, 71 = Hippeastrum solandrifolium.
longifolia, Linn. Sp. Pl. 293 = Crinum longifolium.
longipetala, *Lem. Illustr. Hortic.* xiii. (1866) *Misc.* 78.—Hab. ?
lorifolia, Steud. Nom. ed. II. i. 71 (= *Habranthus lorifolius*).—Am. austr.
Lowii, *Morr. in Ann. Soc. Agr. Gand*, ii. (1846) 141. t. 62.—Hab. ?
lucida, D. Dietr. Syn. Pl. ii. 1181 = Nerine lucida.
lutea, Bieb. Fl. Taur. Cauc. iii. 255 = Sternbergia Fischeriana.
lutea, Linn. Sp. Pl. 292 = Sternbergia lutea.
lutea, Pav. ex Roem. & Schult. Syst. vii. 797.—Cf. *lutescens*.
lutescens, Herb. ex Kunth, Enum. Pl. v. 497 = Hippeastrum chilense.
maculata, L'Hérit. Sert. Angl. 10 (= *Habranthus maculatus*).
maranensis, Ker-Gawl. in Bot. Reg. t. 719 = Hippeastrum stylosum.
marginata, Jacq. Hort. Schoenb. i. 34 = Nerine marginata.
mesochloa, Steud. Nom. ed. II. i. 71 = Zephyranthes mesochloa.
mexicana, Hort. ex Savi, Fl. Ital. iii. 29. t. 94 = Vallota purpurea.
miniata, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 355 = Hippeastrum rutilum.
miniata, Ruiz & Pav. Fl. Per. iii. 57 = Hippeastrum miniatum.
minima, Griseb. in Goett. Abh. xxiv. (1879) 321 = Zephyranthes minima.
minima, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 350; Rev. 8 = Zephyranthes verecunda.
minor, D. Dietr. Syn. Pl. ii. 1180 = Brunsvigia minor.
minuta, H. B. & K. Nov. Gen. et Sp. i. 278 = Zephyranthes verecunda.
moluccana, Ker-Gawl. in Journ. Sc. & Arts, iii. (1817) 109 = Crinum latifolium.

AMARYLLIS:—
montana, Labill. Ic. Pl. Syr. Dec. ii. 5 = Ixiolirion montanum.
multiflora, D. Dietr. Syn. Pl. ii. 1180 = Brunsvigia gigantea.
multiflora, Tratt. Gartenpfl. 37 = Haemanthus multiflorus.
nervosa, H. B. & K. Nov. Gen. et Sp. i. 278 = Zephyranthes tubispatha.
nervosa, Poir. Encyc. Suppl. i. 321 = Brunsvigia striata.
nivea, Schult. f. Syst. vii. 799 = Zephyranthes candida.
nobilis, Salisb. Prod. 235 = Brunsvigia gigantea.
ochroleuca, Ker-Gawl. ex Schult. f. Syst. vii. 797 = Zephyranthes chloroleuca.
orientalis, Linn. Sp. Pl. 293 = Brunsvigia gigantea.
ornata, Ait. Hort. Kew. i. 418 = Crinum yuccaeflorum.
ornata, Ker-Gawl. in Bot. Mag. t. 1171 = Crinum zeylanicum.
Osbeckii, Sabine, ex Desf. Cat. ed. III. 388 = Crinum Osbeckii.
ovigera, Schult. f. Syst. vii. 840 = Peliosanthes javanica ?
pallida, [Del.] Red. Liliac. viii. 479 = A. Belladonna.
pallida, Willd. ex Schult. f. Syst. vii. 801 = Zephyranthes verecunda.
pardina, Hook. f. Bot. Mag. t. 5645 = Hippeastrum pardinum.
parvula, Seub. in Mart. Fl. Bras. iii. 145 = Zephyranthes minima.
pendula, Salisb. Prod. 231 = Cyrtanthus obliquus.
peruviana, Ker-Gawl. in Bot. Mag. sub t. 1089 (= *Pyrolirion aureum*).
phycelloides, Steud. Nom. ed. II. i. 72 = Hippeastrum phycelloides.
platypetala, Lindl. ex Bury, Hexand. Pl. 15 = Lycoris aurea.
praeclara, Sweet, Hort. Brit. ed. II. 506 = Hippeastrum reticulatum.
Princeps, Vell. Fl. Flum. 130; iii. t. 115 = Hippeastrum reticulatum.
Principis, Salm-Dyck, in Nov. Act. Nat. Cur. x. (1821) 154 = Hippeastrum reticulatum.
procera, Duchart. in Bull. Soc. Bot. Fr. x. (1863) 75 = Hippeastrum procerum.
procera, Salisb. Prod. 230 = Crinum erubescens.
propinqua, *Salisb. l. c.*—Panama.
psittacina, Ker-Gawl. in Bot. Reg. t. 199 = Hippeastrum psittacinum.
pudica, Ker-Gawl. in Journ. Sc. & Arts, ii. (1817) 348 = A. Belladonna.
pulchella, Salisb. Prod. 229 = Zephyranthes Atamasco.
pulchella, Spreng. ex Kunth, Enum. Pl. ii. 633 = Hessea crispa.
pulchra, Salisb. Prod. 233 = Nerine sarniensis.
pulverulenta, Lodd. Bot. Cab. t. 484 = Hippeastrum rutilum.
pumila, Spin. Jard. Sébast. 28; ex Spreng. Neue Entdeck. i. 252 = Zephyranthes pumila.
Pumilio, Ait. Hort. Kew. i. 415 = Cyrtanthus uniflorus.
punicea, Lam. Encyc. i. 122 = Hippeastrum equestre.
purpurea, Ait. Hort. Kew. i. 417 = Vallota purpurea.
radiata, L'Hérit. Sert. Angl. 15 = Lycoris radiata.
Radula, Jacq. Hort. Schoenb. i. 35 = Brunsvigia Radula.
Rayneri, Hook. Bot. Mag. t. 5883 = Hippeastrum procerum.
regalis, Salisb. Prod. 232 = A. Belladonna.
Reginae, Linn. Syst. Nat. ed. X. ii. 977 = Hippeastrum Reginae.
reticulata, L'Hérit. Sert. Angl. 12 = Hippeastrum reticulatum.
revoluta, L'Hérit. l. c. 13 = Crinum lineare.
riparia, Burch. ex Kunth, Enum. Pl. ii. 580 = Crinum longifolium.
robusta, A. Dietr. in Otto & Dietr. Allg. Gartz. xviii. (1850) 41 = Hippeastrum aulicum.
robusta, Sweet, ex Steud. Nom. ed. II. i. 74 (= *Habranthus robustus*).
Roezli, Regel, Gartenfl. (1874) 290. f. 809 = Hippeastrum equestre.
rosea, Lam. Encyc. i. 122 = A. Belladonna.
rosea, Spreng. Syst. iv. Cur. Post. 133 = Zephyranthes rosea.
rotundifolia, Lam. Encyc. i. 124 = Eurycles sylvestris.
Rougieri, Hort. ex Carr. in Rev. Hortic. (1882) 312 = Hippeastrum aulicum.
rutila, Ker-Gawl. Rev. 16; in Bot. Reg. i. t. 23 = Hippeastrum rutilum.

AMARYLLIS:—
rutilans, Lodd. ex Steud. Nom. ed. II. i. 74 = Hippeastrum rutilum.
sarniensis, Linn. Sp. Pl. 293 = Nerine sarniensis.
scillifolia, Hort. Angl. ex Schult. f. Syst. vii. 854 = Crinum distichum.
semibarbata, Steud. Nom. ed. II. i. 72, sphalm. (*loco* Antherici) = Bulbine bulbosa.
Slateriana, Lindl. Bot. Reg. (1844) App. 76 = Brunsvigia Slateriana.
solandraeflora, Lindl. Coll. Bot. t. 11 = Hippeastrum solandraeflorum.
spathacea, Schult. f. Syst. vii. 808 (= *Habranthus spathaceus*).—Bonaria.
speciosa, L'Hérit. Sert. Angl. 12 = Vallota purpurea.
spectabilis, Andr. Bot. Rep. t. 390 = Crinum yuccaeflorum.
spectabilis, Lodd. Bot. Cab. t. 159 = Hippeastrum Reginae.
spiralis, L'Hérit. Sert. Angl. 10 = Hessea spiralis.
staminea, Seub. in Mart. Fl. Bras. iii. 150 = Hippeastrum stylosum.
stellaris, Jacq. Hort. Schoenb. i. 37 = Hessea stellaris.
stilosa, Bury, Hexand. Pl. t. 33 = Hippeastrum stylosum.
striata, Jacq. Hort. Schoenb. i. 36 = Brunsvigia striata.
striata, Lam. Encyc. i. 125 = Hippeastrum rutilum.
striatula, Schult. f. Syst. vii. 801 = Zephyranthes verecunda.
stylosa, Sweet, Hort. Brit. ed. I. 403 = Hippeastrum stylosum.
subbarbata, Schult. f. Syst. vii. 819 = Hippeastrum rutilum.
superba, Bury, Hexand. Pl. t. 31 = Hippeastrum vittatum.
Sweetiana, Steud. Nom. ed. II. i. 72 = Hippeastrum adoenum.
Sweetii, Herb. ex Sweet, Hort. Brit. ed. II. 506 = Hippeastrum reticulatum.
sylvatica, Mart. in Schult. f. Syst. vii. 807 (= *Habranthus sylvaticus*).
Symoni, *Funck, in Journ. Hort. Prat.* Sér. II. iii. (1859) 242. t. 20.—Hab.?
tatarica, Georgi, Beschreib. Russ. Reich. iii. iv. 890 = Ixiolirion Ledebouri.
tatarica, Pall. Reise, iii. 727 = Ixiolirion tataricum.
tenella, Thunb. ex Steud. Nom. ed. II. i. 72 = Carpolyza spiralis.
toxicaria, D. Dietr. Syn. Pl. ii. 1181 = Brunsvigia toxicaria.
tubiflora, L'Hérit. Sert. Angl. 10 (= *Pyrolirion aureum*).—Peruv.
tubispatha, L'Hérit. l. c. 9 = Zephyranthes tubispatha.
Umbella, L'Hérit. l. c. 15 = Cyrtanthus obliquus.
undulata, Linn. Syst. ed. XII. 237 = Nerine undulata.
unguiculata, Mart. ex Schult. f. Syst. vii. 189 = Hippeastrum calyptratum.
vaginata, D. Dietr. Syn. Pl. ii. 1181 = Strumaria truncata.
valparadisiaca, Steud. Nom. ed. II. i. 183 (= *Habranthus pallidus*).—Chili.
variabilis, Jacq. Hort. Schoenb. iv. 14 = Crinum variabile.
venusta, Ker-Gawl. in Bot. Mag. t. 1090 = Nerine sarniensis.
verecunda, Salisb. Prod. 229 = Zephyranthes Atamasco.
verecunda, Schult. f. Syst. vii. 800 = Zephyranthes verecunda.
vernalis, Mill. Gard. Dict. ed. VIII. n. 10 = Hypoxis stellatus.
Verreauxii, *Hort.—Cf. Journ. et Flore des Jardins*, ii. (1832) t. 119.—Hab.?
versicolor, Spreng. Syst. Cur. Post. 133 (= *Habranthus versicolor*).
vittata, L'Hérit. Sert. Angl. 13 = Hippeastrum vittatum.
vivipara, Lam. Encyc. i. 123 = Crinum defixum.
yuccaeoides, Thomps. Bot. Displ. t. 12 = Crinum yuccaeflorum.
zeylanica, Linn. Sp. Pl. 293 = Crinum zeylanicum.

AMARYLLIS, Sweet, Hort. Brit. ed. II. 506 (1830) = **Hippeastrum**, Herb.

AMASONIA, Linn. f. Suppl. 48, 294 (1781). *VERBENACEAE*, Benth. & Hook. f. ii. 1147.
DIPHYSTEMA, Neck. Elem. i. 382 (1790).
DIPLOSTEMMA, 'Neck.' ex DC. Prod. vii. 306. in nota (1838) lapsu.
TALIGALEA, Aubl. Pl. Gui. ii. 625, t. 252 (1775).

AMASONIA:—
angustifolia, *Mart. & Schau. in DC. Prod.* xi. 678.—Bras.
arborea, *H. B. & K. Nov. Gen. et Sp.* ii. 253.—Venezuela.
erecta, *Linn. f. Suppl.* 294.—Guiana.
hirta, *Benth. in Ann. Nat. Hist.* ii. (1839) 451.—Guiana.
integerrima, *Spreng. Syst.* ii. 765.—N. Granat.
lasiocaulon, *Mart. & Schau. in DC. Prod.* xi. 678.—Am. austr.
punicea, *Vahl, Eclog.* ii. 51.—Ins. Trinit.
Taligalea, Auct. ex Steud. Nom. ed. II. i. 74 = erecta.
velutina, *Schau. in DC. Prod.* xi. 677.—Bras.

AMATHEA, Rafin. Fl. Tellur. iv. 65 (1836) = **Aphelandra**, R. Br. (Acanth.).
pulchra, Rafin. l. c. = Aph. pulcherrima.

AMATULA, Medic. Bot. Beobacht. 38 (1782); Malv. 106 (1787) = **Lycopersicum**, Mill. (Solan.).
flava, Medic. Malv. 106 = L. esculentum.
rubra, Medic. l. c. = L. esculentum.

AMAURIA, Benth. Bot. Voy. Sulph. 31 (1844). *COMPOSITAE*, Benth. & Hook. f. ii. 404.
dissecta, *A. Gray, Pl. Fendl.* 104.—N. Mexic.
rotundifolia, *Benth. Bot. Voy. Sulph.* 31.—Calif.

AMAXITIS, Adans. Fam. ii. 34 (1763) = **Dactylis**, Linn. (Gramin.).

AMBAIBA, Adans. l. c. 377 (1763) = **Cecropia** Loefl. (Urtic.).

AMBASSA, Steetz, in Peters, Reise Mossamb. Bot. 364 (1863) = **Vernonia**, Schreb. (Compos.).
Hochstetteri, Steetz, l. c. = V. Hochstetteri.

AMBELANIA, Aubl. Pl. Gui. i. 265. t. 104 (1775). *APOCYNACEAE*, Benth. & Hook. f. ii. 694.
BENTECA, Adans. Fam. ii. 166 (1763).
KASAILO, Dennst. Schluess. Hort. Malab. 30 (1818).
WILLUGHBEIA, Scop. in Schreb. Gen. 162 (1789).
acida, *Aubl. Pl. Gui.* i. 265. t. 104.—Guiana.
cuneata, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 17.—Bras.
laxa, *Muell. Arg. l. c.*—Am. austr.
macrophylla, *Muell. Arg. l. c.* 18.—Bras.
quadrangularis, *Muell. Arg. l. c.*—Bras.
Sagoti, *Muell. Arg. in Linnaea*, xxx. (1859–60) 389.—Guiana.
tenuiflora, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 16.—Bras.

AMBERBOA, Less. Syn. Comp. 8 (1832) = **Volutarella**, Cass. etc. (Compos.).
abyssinica, A. Rich. Tent. Fl. Abyss. i. 453 = Centaurea Hochstetteri.
Bollei, Sch. Bip. ex C. Bolle, in Bonplandia, vii. (1859) 297.—Ins. Canar.
carthamoides, DC. Prod. vi. 561 (= *Phaeopappus carthamoides*).
crupinoides, DC. l. c. 559 = V. bicolor.
decurrens, DC. l. c. 560 = Centaurea decurrens.
discolor, Lojac. in Natural. Sicil. iv. (1885) 133.—Sicil.
erucifolia, DC. Prod. vi. 559 = V. erucifolia.
glauca, Less. Syn. Comp. 8 = Centaurea moschata?
Goniocaulon, DC. Prod. vi. 558 = Goniocaulon glabrum.
Hochstetteri, Buching. ex Oliver, Fl. Trop. Afr. iii. 438 = Centaurea Hochstetteri.
incanescens, DC. Prod. vi. 561 = Centaurea integrifolia.
indica, DC. l. c. 558 = Goniocaulon glabrum.
lancifolia, DC. l. c. 661 = Centaurea lancifolia.
leucantha, Coss. Ind. Pl. Maroc. 21.—Algeria.
Lippii, DC. Prod. vi. 559 = V. Lippii.
micractis, Boiss. & Reut. Diagn. Ser. II. iii. 62.—Algeria.
moschata, Less. Syn. Comp. 8 = Centaurea moschata.
muricata, DC. Prod. vi. 559 = V. muricata.
odorata, DC. l. c. = Centaurea moschata.
Perralderiana, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 62, nomen.—Afr. bor.
phaeopappa, DC. Prod. vi. 560 = Centaurea leuzeoides.
sinaica, DC. l. c. 559 = V. bicolor.
spectabilis, DC. l. c. 561 (= *Phaeopappus spectabilis*).
Wallichiana, DC. l. c. 558 = Centaurea Wallichiana.
xanthocephala, DC. l. c. 561 (= *Psephellus xanthocephalus*).

AMBERBOI, Adans. Fam. ii. 117 (1763) = AMBERBOA.

AMBINAX, Comm. ex Juss. Gen. 389 (1789) = **Aleurites**, Forst. (Euphorb.).

AMBLACHAENIUM, Turcz. ex DC. Prod. vii. 94 (1838) = **Hypochoeris**, Linn. (Compos.).
aurantiacum, Turcz. ex DC. l. c. = H. grandiflora.

AMBLATUM, [Tourn.] G. Don, Gen. Syst. iv. 635 (1837) = ANBLATUM, Tourn. = **Lathraea**, Linn. (Orobanch.).

AMBLEIA, Spach, Hist. Vég. Phan. ix. 166 (1840), nomen = **Stachys**, Linn. (Labiat.).

AMBLIRION, Rafin. in Am. Monthly Mag. (1818) 265 = **Fritillaria**, Linn. (Liliac.).
album, Sweet, Hort. Brit. ed. II. 538 = F. alba.
camtschacense, Sweet, l. c. = F. kamtschatcensis.
grecum, Rafin. Fl. Tellur. ii. 35 = F. graecum?
lanceolatum, Sweet, l. c. = F. kamtschatcensis.
pudicum, Rafin. in Am. Monthly Mag. (1818) 265 = F. pudica.

AMBLOGYNA, Rafin. Fl. Tellur. iii. 42 (1836) = **Amaranthus**, Linn.
fimbriata, A. Gray, in Proc. Am. Acad. v. (1861) 167, 168 = Amar. fimbriatus.
polygonoides, Dalz. & Gibs. Bomb. Fl. 219 = Amar. polygamus.
polygonoides, Rafin. Fl. Tellur. iii. 42 = Amar. polygonoides.
scariosa, Moq. in DC. Prod. xiii. II. 269; A. Gray, in Proc. Am. Acad. v. (1861) 169 = Amar. scariosus.
squarrulosa, A. Gray, l. c.—Ins. Galap.
Torreyi, A. Gray, l. c. 167, 169 = Amar. Torreyi.
urceolata, Moq. in DC. Prod. xiii. II. 269; A. Gray, l. c. 168 = Amar. urceolatus.

AMBLOSTIMA, Rafin. Fl. Tellur. ii. 26 (1836) = **Schoenolirion**, Torr. (Liliac.).
albiflora, Rafin. l. c. = Sc. Elliottii.
crocea, Rafin. l. c. = Sc. croceum.
latifolia, Rafin. ex S. Wats. in Proc. Am. Acad. xiv. (1879) 242 = Sc. Elliottii.

AMBLOSTOMA, Scheidw. in Otto & Dietr. Gartenz. vi. (1838) 383. *ORCHIDEAE*, Benth. & Hook. f. iii. 522.
cernuum, *Scheidw. l. c.*—Bras.
densum, *Reichb. f. Xenia Orch.* iii. 22.—Peruv.
micranthum, *Reichb. f. in Saund. Refug. sub* t. 101.—Boliv.
tridactylum, Reichb. f. in Walp. Ann. vi. 485 = cernuum.

AMBLYACHYRUM, Hochst. ex Steud. Syn. Pl. Gram. 413 (1855) = **Apocopis**, Nees (Gramin.).
mangalorense, Hochst. ex Steud. l. c.; et in Flora, xxxix. (1856) 26 = Apoc. Wightii.

AMBLYANTHERA, Blume, Mus. Bot. Lugd. Bat. i. 50 (1849) = **Osbeckia**, Linn. (Melast.).
parvifolia, Blume, l. c. = O. parvifolia.
truncata, Blume, l. c. = O. truncata.

AMBLYANTHERA, Muell. Arg. in Mart. Fl. Bras. vi. 1. 141, t. 44, 45 (1860) = **Mandevilla**, Lindl. (Apocyn.).
acutiloba, Mucll. Arg. in Linnaea, xxx. (1859–60) 426 = Echites acutiloba.
andina, Mucll. Arg. l. c. 425 = Echites andina.
Andrieuxii, Muell. Arg. l. c. 422 = Mandevilla Andrieuxii.
angustifolia, Mucll. Arg. l. c. 430 = Echites angustifolia.
antennacea, Muell. Arg. l. c. 448 = Echites antennacea.
Benthami, Muell. Arg. l. c. 451 = Echites Benthami.
bogotensis, Muell. Arg. l. c. 452 = Echites bogotensis.
brachyloba, Muell. Arg. l. c. 423 = Echites brachyloba.
Bridgesii, Muell. Arg. l. c. 420.—Boliv.
campestris, Muell. Arg. in Mart. Fl. Bras. vi. 1. 149 = Echites campestris.
ciliata, Muell. Arg. l. c. 145 = Echites ciliata.
Claussenii, Miers, Apocyn. S. Am. 274, t. 25.—Am. austr.
convolvulacea, Muell. Arg. in Linnaea, xxx. (1859–60) 430 = M. convolvulacea.

AMBLYANTHERA :—
crassifolia, Muell. Arg. in Mart. Fl. Bras. vi. 1. 143 = Echites crassifolia.
cuiabensis, Muell. Arg. l. c. 145 = Echites cuyabensis.
cuyabensis, Muell. Arg. in Linnaea, xxx. (1859–60) 450 = praec.
Fendleri, Muell. Arg. l. c. 417.—Venezuela.
foliola, Muell. Arg. l. c. 427 = Mandevilla foliosa.
fluminensis, Muell. Arg. in Mart. Fl. Bras. vi. 1. 148 = Echites fluminensis.
funiformis, Muell. Arg. l. c. 144 = Echites funiformis.
guyanensis, Muell. Arg. in Linnaea, xxx. (1859–60) 448 = Echites guianensis.
hirsuta, Miers, Apocyn. S. Am. 185 = M. hispida ?
hispida, Muell. Arg. in Mart. Fl. Bras. vi. 1. 147 = M. hispida.
Karwinskii, Muell. Arg. in Linnaea, xxx. (1859–60) 426 = M. Karwinskii.
lasiocarpa, Muell. Arg. in Kjoeb. Vidensk. Meddel. (1869) 113 = M. hispida.
leptophylla, Muell. Arg. in Mart. Fl. Bras. vi. 1. 142 = Echites leptophylla.
macrophylla, Muell. Arg. in Linnaea, xxx. (1859–60) 449 = Echites macrophylla.
madida, Muell. Arg. in Mart. Fl. Bras. vi. 1. 150 = Echites madida.
membranacea, Muell. Arg. in Linnaea, xxx. (1859–60) 423, in nota = Echites membranacea.
mexicana, Muell. Arg. l. c. 424 = Echites mexicana.
microcalyx, Muell. Arg. l. c. 428 = Echites microcalyx.
Moritziana, Muell. Arg. l. c. 421.—N. Granat.
ovata, Miers, Apocyn. S. Am. 188.—Boliv.
oaxacana, Muell. Arg. in Linnaea, xxx. (1859–60) 447 = M. oaxacana.
palustris, Muell. Arg. in Mart. Fl. Bras. vi. 1. 145 = M. palustris.
pandurata, Muell. Arg. in Linnaea, xxx. (1859–60) 448 = Echites pandurata.
Pavonii, Muell. Arg. l. c. 450 = Echites Pavonii.
Prieurei, Muell. Arg. l. c. 448 = Echites Prieurei.
Schlimi, Muell. Arg. l. c. 419.—N. Granat.
Spruceana, Muell. Arg. in Mart. Fl. Bras. vi. 1. 143.—Bras.
suaveolens, Muell. Arg. in Linnaea, xxx. (1859–60) 447 = M. suaveolens.
subsessilis, Muell. Arg. l. c. = Echites subsessilis.
torosa, Muell. Arg. l. c. 446 = Echites torulosa.
tubiflora, Muell. Arg. l. c. 454 = Echites tubiflora.
versicolor, Muell. Arg. in Mart. Fl. Bras. vi. 1. 146 = Echites versicolor.

AMBLYANTHUS, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 83. t. 6. *MYRSINEAE*, Benth. & Hook. f. ii. 648.
glandulosus, *A. DC. l. c.*—Reg. Himal.

AMBLYCARPUM, Lem. in Orb. Dict. i. 335 (1841) = **Amblyocarpum**, Fisch. & Mey. (Compos.).

AMBLYCHLOA, Link, in Linnaea, xvii. (1843) 399 = **Sclerochloa**, Beauv. (Gramin.).
dura, Link, l. c. = S. dura.

AMBLYGLOTTIS, Blume, Bijdr. 369. t. 64 (1825) = **Calanthe**, R. Br. (Orchid.).
abbreviata, Blume, l. c. 371 = C. abbreviata.
angustifolia, Blume, l. c. 369 = C. angustifolia.
emarginata, Blume, l. c. 370 = C. emarginata.
flava, Blume, l. e. t. 64 = C. veratrifolia.
pilosa, De Vriese, ex Lindl. Fol. Orch. Calanthe, 11 = C. pilosa.
pulchra, Blume, Bijdr. 371 = C. curculigoides.
speciosa, Blume, l. e. 371 = C. speciosa.
veratrifolia, Blume, l. c. 370 = C. veratrifolia.

AMBLYGONUM, Reichb. Handb. 236 (1837) = **Polygonum**, Linn.

AMBLYLEPIS, Decne. in Orb. Dict. i. 337 (1841) = **Amblyolepis**.

AMBLYOCALYX, Benth. in Benth. & Hook. f. Gen. ii. 698 (1876). *APOCYNACEAE*, Benth. & Hook. f. ii. 698.
Beccarii, *Benth. in Hook. Ic. Pl.* t. 1179.—Borneo.

AMBLYOCARPUM, Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 30 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 336.
inuloides, *Fisch. & Mey. l. c.*—Reg. Caspica.

AMBLYOGLOSSUM, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 310 = **Tylophora**, R. Br. (Asclep.).
brevipes, Turcz. l. c. 311 = T. brevipes.
longipes, Turcz. l. c.—Malaya.

AMBLYOLEPIS, DC. Prod. v. 667 (1836) = **Helenium**, Linn. (Compos.).
setigera, DC. l. c.—Mexic.

AMBLYOPAPPUS, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 321. *COMPOSITAE*, Benth. & Hook. f. ii. 406.
Aromia, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 395.
Infantea, Remy, in C. Gay, Fl. Chil. iv. 257. t. 48 (1849).
mendocinus, *Phil. Sert. Mend. Alt.* 26.—Chili.
neo-mexicanus, *A. Gray, in Pacif. Rail. Rep.* iv. 106.—Mexic.
pusillus, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 321.—Chili.

AMBLYOPELIS, Steud. Nom. ed. II. i. 75 (1840), sphalm. = AMBLYOLEPIS, DC.

AMBLYOPOGON, Fisch. & Mey. ex DC. Prod. vi. 561 (1837) = **Centaurea**, Tourn. (Compos.).
Balansae, Boiss. Fl. Orient. iii. 604, in syn. = C. Balansae.
concinnus, Boiss. & Huet, Diagn. Ser. II. iii. 65 = C. pulcherrima.
incanescens, Fisch. & Mey. ex DC. Prod. vi. 561 = C. integrifolia.
speciosus, Boiss. Fl. Orient. iii. 604, in syn. = C. pulcherrima.

AMBLYORHINUM, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) ii. 168 = **Phyllactis**, Pers. (Valerian.).
angustifolium, Turcz. l. c. 169.—Venezuela.
grandiflorum, Turcz. l. c. 168.—Venezuela.
phylicoides, Turcz. l. c. 171.—Venezuela.
spicatum, Turcz. l. c. 170 = P. cordifolia.
triplinerve, Turcz. l. c.—Venezuela.

AMBLYSPERMA, Benth. in Enum Pl. Hueg. 67 (1837) = **Trichocline**, Cass. (Compos.).
scapigerum, Benth. l. c. = T. scapigera.

AMBLYSTIGMA, Benth. in Benth. & Hook. f. Gen. ii. 748 (1876). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 748.
cionophorum, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 366.—Reg. Argent.
hypoleucum, *Benth. in Hook. Ic. Pl.* xii. 76. t. 1188.—Bolivia.
pedunculare, *Benth. l. c.* 77. sub t. 1188.—Bolivia.

AMBLYTES, Dulac, Fl. Hautes-Pyr. 80 (1867) = **Molinia**, Scop. (Gramin.).
caerulea, Dulac, l. c. = M. caerulea.

AMBORA, Juss. Gen. 401 (1789) = **Tambourissa**, Sonn. (Monim.).
alternifolia, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 31 = T. alternifolia.
amplifolia, Tul. in Arch. Mus. Par. viii. (1855–56) 299 = T. amplifolia.
cymosa, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 31 (an Rubiacea ?).—Hab. ?
elliptica, Tul. l. c. = T. elliptica.
Ficus, Tul. l. c. 30 = T. Ficus.
fruticosa, Boiv. ex Tul. l. c. = T. purpurea.
grandifolia, R. Br. ex Baker, Fl. Maurit. 287 = T. amplifolia.
leptophylla, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 29 = T. leptophylla.
neglecta, Tul. in Arch. Mus. Par. viii. (1855–56) 432 = T. neglecta.
obovata, Tul. in Ann. Sc. Nat. Sér. IV. t. iii. (1855) 31 = T. obovata.
peltatus, R. Br. ex Baker, Fl. Maurit. 288 = T. peltata.
purpurea, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 30 = T. purpurea.
quadrifida, Poir. Encyc. vii. 565 = T. quadrifida.
racemosa, R. Br. ex Baker, Fl. Maurit. 288 = T. quadrifida.
religiosa, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 30 = T. religiosa.

AMBORA :—
sempervirens, F. G. Dietr. Lexik. Gaertn. i. 342 = T. quadrifida.
Sieberi, Tul. in Arch. Mus. Par. viii. (1855–56) 304 = T. Sieberi.
Tambourissa, Lam. ex Steud. Nom. ed. I. 37, 535 = T. quadrifida.
Tambourissa, Tul. in Arch. Mus. Par. viii. (1855–56) 303 = T. Boivinii.
tetragona, Boiv. ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 30 = T. tetragona.
tomentosa, Bory, Voy. i. 317. t. 13 = Monimia rotundifolia.
tomentosa, Sieber, ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 31 = T. Sieberi.
vestita, Tul. l. c. = T. vestita.

AMBORELLA, Baill. Hist. Pl. i. 328 in nota (1869); et Adansonia, x. 354 (1873). *MONIMIACEAE*, Benth. & Hook. f. iii. 142.
trichopoda, *Baill. l. c.*—N. Caled.

AMBRARIA, Cruse, Rub. Cap. 16. t. 1, f. 3, 4 (1825) = **Nenax**, Gaertn. (Rubiac.).
acerosa, Sond. in Harv. & Sond. Fl. Cap. 34 = N. acerosa.
glabra, Cruse, Rub. Cap. 16. t. 1 = N. acerosa.
hirta, Cruse, l. c.—Afr. austr.
microphylla, Sond. in Harv. & Sond. Fl. Cap. 34.—Afr. austr.

AMBRARIA, Heist. Syst. 11 (1748) = **Anthospermum**, Linn. (Rubiac.).
Heisteri, Willd. Sp. Pl. iv. 732 in syn. = Anth. aethiopicum.

AMBRINA, Spach, Hist. Vég. Phan. v. 295 (1836) = **Chenopodium**, Linn.
ambrosioides, Spach, l. c. 297 = C. ambrosioides.
anthelmintica, Spach, l. c. 298 = C. ambrosioides.
Botrys, Moq. Chen. Enum. 37 = C. Botrys.
carinata, Moq. l. c. 41 = C. carinatum.
chilensis, Spach, Hist. Vég. Phan. v. 297 = C. chilense.
denudata, Phil. in Linnaea, xxix. (1857–58) 37 = Roubieva multifida.
dissecta, Moq. Chen. Enum. 38 = C. bipinnatifidum.
foetida, Moq. l. c. 38 = C. Botrys.
graveolens, Moq. l. c. 41 = C. graveolens.
incisa, Moq. l. c. 36 = C. incisum.
obovata, Moq. l. c. 40 = C. obovatum.
pinnatisecta, Spach, Hist. Vég. Phan. v. 296 = Roubieva multifida.
Pumilio, Moq. Chen. Enum. 42 = C. Pumilio.
retusa, Moq. l. c. 40 = C. retusa.
spathulata, Moq. l. c. 39 = C. spathulatum.
tenuis, Moq. l. c. 42 = Roubieva multifida.
tomentosa, Moq. l. c. 37 = C. tomentosum.

AMBROMA, Linn. f. Suppl. 54, 341 (1781) = **Abroma**, Jacq. (Stercul.).
Augusta, Linn. f. l. c. 341 = Abr. fastuosum.

AMBROSIA, Bassi, in Comm. Bonon. v. (1763) 82 = **Ambrosinia**, Linn. (Aroid.).

AMBROSIA, Linn. Gen. ed. I. 287. n. 718 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 354.
abrotanifolia, *Wall. Cat.* n. 3234.—Bras.
absynthyfolia, Michx. Fl. Bor. Am. ii. 183 = artemisiaefolia.
acanthicarpa, Hook. Fl. Bor. Am. i. 309 = Franseria Hookeriana.
aptera, *DC. Prod.* v. 527.—Am. bor.
arborescens, Lam. Encyc. i. 127 = Franseria artemisioides.
arborescens, *Mill. Gard. Dict.* ed. VIII. n. 5.—Peruv.
artemisioides, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 268.—Peruv.
artemisiaefolia, Bess. Prim. Fl. Galic. ii. 390 = maritima.
artemisifolia, *Linn. Sp. Pl.* 988.—Am. bor.
bidentata, *Michx. Fl. Bor. Am.* ii. 182.—Am. bor.; Ind. occ.
cheiranthifolia, *A. Gray, in Torr. Bot. Mex. Bound,* 87.—Mexic.
chilensis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1840) 311.—Chili.
confertiflora, DC. Prod. v. 526 = Franseria tenuifolia.
coronopifolia, Torr. & Gray, Fl. N. Am. ii. 271 = psilostachya.
crithmifolia, DC. Prod. v. 525 = hispida.

AMBROSIA :—
cumanensis, *H. B. & K. Nov. Gen. et Sp.* iv. 276.—Venezuela.
elata, Salisb. Prod. 175 = artemisifolia.
elatior, Bess. Prim. Fl. Galic. ii. 389 = maritima.
elatior, Linn. Sp. Pl. 987 = artemisifolia.
fruticosa, DC. Prod. v. 525 = Franseria tenuifolia.
fruticosa, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 247 = arborescens.
glandulosa, Scheele, in Linnaea, xxii. (1849) 157 = psilostachya.
heterophylla, Muhl. ex Willd. Sp. Pl. iv. 377 = artemisifolia.
hispida, *Pursh, Fl. Am. Sept. Suppl.* 743.—Florida.
hispida, Torr. in Ann. Lyc. N. York, ii. (1828) 216 = psilostachya.
humilis, *Vell. Fl. Flum.* x. t. 25.—Bras.
incisa, F. G. Dietr. Lexik. Gaertn. i. 344 = artemisifolia.
integrifolia, Muhl. ex Willd. Sp. Pl. iii. 275 = trifida.
Lindheimeriana, Scheele, in Linnaea, xxii. (1849) 156 = psilostachya.
longistylis, A. Gray, Pl. Fendler. 79 = Franseria tenuifolia.
longistylis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 344.—Am. bor.
maritima, Ferrero, ex DC. Prod. v. 525 = crithmifolia.
maritima, *Linn. Sp. Pl.* 988.—Europ.; Orient; Afr. bor.
maritima, Vell. Fl. Flum. x. t. 26 = polystachya.
microcephala, *DC. Prod.* v. 527.—Guiana.
orobanchifera, *Meyen, Reise*, i. 444.—Peruv.
paniculata, Michx. Fl. Bor. Am. ii. 183 = artemisifolia.
peruviana, DC. Prod. v. 526 = psilostachya.
peruviana, Willd. Sp. Pl. iv. 377.—Peruv.
Pitcheri, Torr. ex Hook. Comp. Bot. Mag. i. (1835) 99 = Iva ciliata.
polystachya, *DC. Prod.* v. 526.—Bras.
psilostachya, *DC. l. c.*—Am. bor.
pumila, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 217.—Am. bor.
pusilla, Rafin. ex DC. Prod. v. 527 = artemisifolia.
scabra, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1840) 311.—Bras.
senegalensis, DC. Prod. v. 525 = maritima.
simplicifolia, Racusch. Nom. ed. III. 274 = artemisifolia.
tacorensis, *Meyen, Reise*, i. 456.—Peruv.
tenuifolia, Gren. & Godr. Fl. Fr. ii. 395 = Franseria tenuifolia.
tenuifolia, Spreng. Syst. iii. 851 = polystachya?
tomentosa, Nutt. Gen. Am. ii. 186 = Franseria discolor.
trifida, *Linn. Sp. Pl.* 987.—Am. bor.
umbellata, Moench, Meth. Suppl. 272 = maritima.
villosissima, *Forsk. Fl. Aegypt. Arab.* 160.—Arabia.

AMBROSINIA, Linn. Gen. ed. VI. 517 (1764) = *AROIDEAE*, Benth. & Hook. f. iii. 964.
AMBROSIA, Bassi, in Comm. Bonon. v. (1763) 82.
Bassii, *Linn. Gen.* ed. VI. 517.—Europ.
ciliaris, Spreng. Syst. iii. 771 = seq.
ciliata, Roxb. Pl. Corom. iii. 90 = Cryptocoryne ciliata.
maculata, Ucria, in Roem. Arch. i. 70 = A. Bassii.
nervosa, Lam. Encyc. i. 128 = A. Bassii.
reticulata, Tin. Cat. Hort. Panorm. 24, 176, sec. Guss. Fl. Sic. Syn. ii. II. 595 = A. Bassii.
retrospiralis, Roxb. Hort. Beng. 65 = Cryptocoryne retrospiralis.
Roxburghiana, Voigt, Hort. Suburb. Calc. 685 = Cryptocoryne unilocularis.
spiralis, Roxb. Hort. Beng. 65 = Cyptocoryne spiralis.
unilocularis, Roxb. Fl. Ind. iii. 493 = Cryptocoryne unilocularis.
velutina, Blume, ex Heynh. Nom. ii. 32 = A. Bassii.

AMBULI, Adans. Fam. ii. 208 (1763) = scq.

AMBULIA, Lam. Encyc. i. 128 (1783) = **Limnophila**, R. Br. (Scrophul.).
alata, Rafin. Fl. Tellur. iv. 58 = Gratiola tetragona.
aromatica, Lam. Encyc. i. 128 = L. gratissima.
rigida, Rafin. Fl. Tellur. iv. 58 = Gratiola tetragona.

AMBUYA, Rafin. Fl. Tellur. iv. 98 (1836) = **Aristolochia**, Tourn.
labiosa, Rafin. l. c. = Ar. cymbifera.

AMEBIA, Regel, Pl. Nov. Fedsch. 58 (1882) crr. typ. = **Arnebia**, Forsk. (Boragin.).

AMECARPUS, Benth. in Lindl. Veg. Kingd. 554 (1847) = **Indigofera**, Linn. (Legumin.).

AMECHANIA, DC. Prod. vii. 578 (1839) = **Agarista**, D. Don (Ericac.).
hispidula, DC. l. c. 579 (= *Leucothoë hispidula*).
subcanescens, DC. l. c. (= *Leucothoë subcanescens*).

AMELANCHIER, Medic. Phil. Bot. i. 135 (1789). *ROSACEAE*, Benth. & Hook. f. i. 628.
ARONIA, Pers. Syn. ii. 39 (1807).
PERAPHYLLUM, Nutt. in Torr. & Gray, Fl. N. Am. i. 474 (1840).
XEROMALON, Rafin. New Fl. Am. iii. 11 (1836).
alnifolia, *Nutt. in Journ. Acad. Phil.* vii. (1834) 22.—Am. bor.
asiatica, Endl. in Walp. Rep. ii. 55 = canadensis.
Bartramiana, M. Roem. Syn. Rosifl. 145 = canadensis.
Botryapium, DC. Prod. ii. 632 = canadensis.
canadensis, *Medic. Gesch.* 79; *Torr. & Gray, Fl. N. Am.* i. 473.—Am. bor.; As. or.
chinensis, Hort. ex C. Koch, Dendrol. i. 186 = Sorbus arbutifolia.
cretica, DC. Prod. ii. 632 = vulgaris.
denticulata, C. Koch, Dendrol. i. 183 = Cotoneaster denticulata.
florida, Lindl. Bot. Reg. t. 1589 = alnifolia.
grandiflora, Dougl. ex M. Roem. Syn. Rosifl. 145 = canadensis.
integrifolia, Boiss. & Hohen. Diagn. Ser. I. iii. 8 = vulgaris.
intermedia, Spach, Hist. Vég. Phan. ii. 85 = canadensis?
japonica, Hort. ex C. Koch, Dendrol. i. 179 = canadensis.
melanocarpa, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 136.—Hab.?
oblongifolia, M. Roem. Syn. Rosifl. 147 = canadensis.
oligocarpa, M. Roem. l. c. 145 = canadensis.
orbicularis, Borck. ex Steud. Nom. ed. II. i. 76 = vulgaris.
ovalis, Medic. Gesch. 78 = canadensis.
parviflora, *Boiss. Diagn.* Ser. I. iii. 9.—As. Min.
parviflora, Hort. ex Loud. Arb. Brit. ii. 877 = alnifolia.
pisidica, Boiss. & Heldr. Diagn. Ser. I. x. 2 = parviflora.
pumila, Nutt. ex Torr. & Gray, Fl. N. Am. i. 474 = alnifolia.
racemosa, Lindl. in Bot. Reg. (1847) sub. t. 38 = Exochorda grandiflora.
rotundifolia, M. Roem. Syn. Rosifl. 146 = canadensis.
sanguinea, DC. Prod. ii. 633 = canadensis?
spicata, C. Koch, Dendrol. i. 182 = canadensis?
suborbicularis, Borkh. Forstbot. ii. 1262 = vulgaris.
tomentosa, Steud. Nom. ed. II. ii. 136 in syn. = vulgaris?
vulgaris, *Moench, Meth.* 682.—Europ.; Oriens.
Wangenheimiana, M. Roem. Syn. Rosifl. 146 = canadensis.

AMELETIA, DC. in Mém. Soc. Phys. Genèv. iii. II. (1826) 82 = **Ammannia**, Houst. (Lythrar.).
acutidens, Miq. Fl. Ind. Bat. i. I. 617 = Amman. peploides.
diandra, F. Mucll. Fragm. iii. 108 = Amman. diandra.
elatinoides, Blume, Mus. Bot. Lugd. Bat. ii. 136 = Amman. elatinoides.
elongata, Blume, l. c. 135 = Amman. peploides.
floribunda, Wight, Illustr. Ind. Bot. 206 = Amman. floribunda.
indica, DC. in Mém. Soc. Hist. Nat. Genèv. iii. II. (1825) 2 & 82. t. 3 f. A. = Amman. peploides.
polystachya, Wall. Cat. n. 2094 = Amman. peploides.
rotundifolia, Wight, Ic. t. 258 = Amman. rotundifolia.
subspicata, Benth. in Hook. Kew Journ. iv. (1842) 81 = Amman. rotundifolia.
tenella, Blume, Mus. Bot. Lugd. Bat. ii. 136 = Amman. tenella.
tenuis, Wight, Ic. Pl. Ind. Or. t. 257 B = Amman. tenuis.
uliginosa, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 261 = Amman. peploides.

AMELIA, Alefeld, in Linnaea, xxviii. (1856) 25 = **Pyrola**, Tourn. (Ericac.).

AMELIA :—
media, Alef. in Linnaea, xxviii. (1856) 30 = Pyrola media.
minor, Alef. l. c. 25 = P. minor.

AMELINA, C. B. Clarke, Commel. et Cyrt. Beng. 38. t. 26 (1874) = **Aneilema**, R. Br. (Commelin.).
Wallichii, C. B. Clarke, l. c. 39. t. 26 = Aneilema acquinoctiale.

AMELLUS, P. Br. Hist. Jamaic. 317 (1756) = **Melanthera**, Rohr. (Compos.).

AMELLUS, Linn. Syst. ed. x. 1225 (1759). *COMPOSITAE*, Benth. & Hook. f. ii. 265.
HAENELIA, Walp. Rep. ii. 974 (1843).
KRAUSSIA, Sch. Bip. in Flora, xxvii. (1844) 672.
alternifolius, Roth, Catalecta, ii. 111 = strigosus.
anisatus, *Cass. in Dict. Sc. Nat.* xxxvii. 489.—Afr. austr.
annuus, *Willd. Sp. Pl.* iii. 2215.—Afr. austr.
carolinianus, Walt. Fl. Carol. 213 = Eclipta alba.
coilopodius, *DC. Prod.* v. 214.—Afr. austr.
diffusus, Forst. in Comm. Gott. ix. (1787) 39 = Chiliotrichum amelloideum.
divaricatus, Gater. Fl. Montaub. 147 = Aster Novi-Belgii.
floribundus, Willd. ex DC. Prod. v. 96 = Liabum igniarium.
flosculosus, DC. Prod. v. 214 = hispidus.
hispidus, *DC. l. c.*—Afr. austr.
Lychnitis, *Linn. Syst.* ed. X. 1255.—Afr. austr.
Lychnitis, E. Mey. ex Steud. Nom. ed. II. i. 76 = tenuifolius.
microglossus, *DC. Prod.* v. 215.—Afr. austr.
mutabilis, Gater. Fl. Montaub. 147 = Aster mutabilis.
nanus, *DC. Prod.* v. 215.—Afr. austr.
officinalis, Gater. Fl. Montaub. 147 = Aster Amellus.
pallidus, Salisb. Prod. 205 = A. Lychnitis.
pedunculatus, Orteg. ex Willd. Enum. Hort. Berol. 916 = Tridax procumbens.
rosmarinifolius, Poepp. ex DC. Prod. v. 216 = Chiliotrichum rosmarinifolium.
scabridus, *DC. l. c.* 214.—Afr. austr.
speciosus, Gater. Fl. Montaub. 146 = Aster chinensis.
spinulosus, *Pursh, Fl. Am. Sept.* ii. 564.—Am. bor.
strigosus, Less. in Linnaea, vi. (1831) 110 = annuus.
tenuifolius, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
tridactylus, *DC. Prod.* v. 215.—Afr. austr.
umbellatus, Linn. Pl. Jam. Pugill. 24 (Syst. ed. X. 1225) = Liabum Brownei.
villosus, Pursh, Fl. Am. Sept. ii. 564 = Chrysopsis villosa.

AMERIMNON, P. Br. Hist. Jamaic. 288 (1756); DC. Prod. ii. 421 (1825) = **Dalbergia**, Linn. (Legumin.).
affine, Spreng. Syst. iii. 192.—Am. austr.
Brownei, Sw. Fl. Ind. Occ. 1234 = Dalbergia Amerimnum.
Brownii, Jacq. Enum. Pl. Carib. 27.—Ind. occ.
Ebenus, Sw. Prod. Veg. Ind. Occ. 104 = Brya Ebenus.
fasciculatum, Buch.-Ham. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 109 = Millettia fruticosa.
guianense, Klotzsch, in Schomb. Faun. et Fl. Guy. 1204.—Guiana.
horridum, Dennst. Schluess. Hort. Malab. 34 (quid?).—Ind. or.
latifolium, Willd. Sp. Pl. iii. 909 = Lonchocarpus latifolius.
mimosella, Blanco, Fl. Filip. ed. I. 563 = Dalbergia lanceolaria.
obovatum, Buch.-Ham. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 246 = Derris thyrsiflora.
pallidum, Buch.-Ham. ex Baker, l. c. 108 = Millettia auriculata.
parviflorum, Spreng. Neue Entdeck. ii. 159.—Bras.
pinnatum, Jacq. Sel. Stirp. Am. 200 = Dalbergia Amerimnum.
polystachyon, Spreng. Syst. Cur. Post. 268 = Eysenhardtia amorphoides.
pubescens, Willd. Sp. Pl. iii. 909 = Pterocarpus scandens.
scandens, Willd. l. c. = Pterocarpus scandens.
secundum, Buch.-Ham. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 217 = Derris secunda.
Sieberi, Reichb. ex DC. Prod. ii. 420 = Ecastaphyllum Brownei.
strigulosum, H. B. & K. Nov. Gen. et Sp. vi. 389.—Venezuela.

AMERINA, DC. in Meissn. Gen. 278 (1840) = **Aegiphila**, Jacq. (Verben.).
ternifolia, DC. Prod. ix. 513.—N. Granat.
tomentosa, DC. l. c. = Aeg. tomentosa ?
triphylla, DC. l. c.—Afr. austr.

AMERINA, Noronha, in Verh. Batav. Gen. v. (1790) Art. iv. 1; ed. II. (1827) 64 = **Aglaia**, Lour. (Meliac.).

AMERINA, Rafin. Alsog. Amer. 15 (1838) = **Salix**, Tourn.
Balduiniana, Rafin. l. c. 16.—Am. bor.
caroliniana, Rafin. l. c. 15 = S. nigra.
glandulosa, Rafin. l. c. = S. pentandra.
Houstoniana, Rafin. l. c. = S. nigra.
missurica, Rafin. l. c. 16.—Am. bor.
nigra, Rafin. l. c. 15 = S. nigra.
tinctoria, Rafin. l. c. 15.—Am. bor.
tomentosa, Rafin. l. c. = S. Lapponum.

AMETHYSTEA, Linn. Amoen. Acad. i. 385 (1749). *LABIATAE*, Benth. & Hook. f. ii. 1220.
AMETHYSTINA, Mess. ex Hall. in Act. Upsal. 1742 (1748) 51. t. 4.
caerulea, *Linn. Sp. Pl.* 21.—Sibiria.
corymbosa, Pers. Syn. i. 24 = caerulea.
trifida, Hill, Veg. Syst. xvii. 44 = caerulea.

AMETHYSTINA, Mess. ex Hall. in Act. Upsal. 1742 (1748) 51. t. 4 = praec.

AMETRON, Rafin. Sylva Tellur. 160 (1838) = **Rubus**, Tourn. (Rosac.).
pedatum, Rafin. l. c. 161 = R. pedatus.
pyrifolium, Rafin. l. c. = R. pyrifolius.

AMHERSTIA, Wall. Pl. As. Rar. i. 1. t. 1, 2 (1830). *LEGUMINOSAE*, Benth. & Hook. f. i. 578.
nobilis, *Wall. l. c.*—Burma.

AMIANTANTHUS, Kunth, Enum. Pl. iv. 179 (1843) = AMIANTHIUM, A. Gray = **Zygadenus**, Michx. (Liliac.).

AMIANTHEMUM, Steud. Nom. ed. II. i. 76 (1840) = seq.

AMIANTHIUM, A. Gray, in Ann. Lyc. N. York, iv. (1848) 121 = **Zygadenus**, Michx. (Liliac.).
angustifolium, A. Gray, l. c. 124 = Z. angustifolius.
aspericaule, A. Gray, l. c. 126 = Melanthium aspericaule.
leimanthoides, A. Gray, l. c. 125 = Z. leimanthoides.
macrotox, Rafin. Fl. Tellur. iv. 27 = Z. muscaetoxicum.
muscaetoxicum, A. Gray, in Ann. Lyc. N. York, iv. (1848) 122 = Z. muscaetoxicum.
Nuttallii, A. Gray, l. c. 123 = Z. Nuttallii, paniculatus.

AMIANTHUM, Rafin. Fl. Tellur. iv. 27 (1836) = praec.

AMICIA, H. B. & K. Nov. Gen. et Sp. vi. 511. t. 600. (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 514.
ZYGOMERIS, Moç. & Sesse, ex DC. Prod. ii. 315 (1825).
glandulosa, *H. B. & K. l. c.*—N. Granat.
medicaginea, *Griseb. in Goett. Abh.* xxiv. (1879) 105.—Reg. Argent.
Zygomeris, *DC. Prod.* ii. 315.—Mexico.

AMICTONIS, Rafin. Sylva Tellur. 161 (1838) = **Callicarpa**, Linn. (Verbenac.).
japonica, Rafin. l. c. = C. longifolia.

AMIDA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 390 = **Madia**, Molina (Compos.).
gracilis, Nutt. l. c. = M. glomerata.
hirsuta, Nutt. l. c. = M. glomerata.

AMIDENA, Adans. Fam. ii. 470 (1763) = **Arisaema**, Mart. (Aroid.).
japonica, Rafin. Fl. Tellur. iv. 15 = Arisaema japonicum.

AMIRIS, La Llave, in La Naturaleza, vii. Apénd. 71 (1885) = **Amyris**, Linn. (Rutac.).

AMIROLA, Pers. Syn. ii. 565 (1807) = **Llagunoa**, Ruiz & Pav. (Sapindac.).
glandulosa, Hook. & Arn. Bot. Beech. Voy. 12 = Ll. glandulosa.
mollis, DC. Prod. i. 616 = Ll. mollis.
nitida, Pers. Syn. ii. 565 = Ll. nitida.
parvifolia, Steud. Nom. ed. II. i. 76 = Ll. nitida.
prunifolia, DC. Prod. i. 616 = Ll. prunifolia.

AMISCHOTOLYPE, Hassk. in Flora, xlvi. (1863) 391 = **Forrestia**, A. Rich. (Commelin.).
glabrata, Hassk. l. c. 392 = F. glabrata.
marginata, Hassk. l. c. = F. marginata.
mollissima, Hassk. l. c. = F. mollis.

AMMANELLA, Miq. Fl. Ind. Bat. i. 1. 618 (1855) = **Ammannia**, Linn. (Lythrar.).
linearis, Miq. l. c. 619 = Ammannia octandra.

AMMANNIA, [Houst.] Linn. Gen. ed. I. 337 (1737). *LYTHRARIEAE*, Benth. & Hook. f. i. 776.
AMELETIA, DC. in Mém. Soc. Phys. Genèv. iii. II. (1826) 82.
CORNELIA, Ard. Specim. ii. 9. t. 1 (1764).
CRYPTOTHECA, Blume, Bijdr. 1128 (1826).
NEXILIS, Rafin. New Fl. Am. iv. 9 (1836).
ORTEGIOIDES, Soland. ex DC. Prod. iii. 76 (1828).
NIMMOIA, Wight, in Madras Journ. Sc. vii. (1837) 311. t. 20.
ROTALA, Linn. Mant. ii. 143 (1771).
SELLOWIA, Roth, ex Roem. & Schult. Syst. v. 407 (1819).
SUFFRENIA, Bellardi, in Act. Taur. vii. (1796) 445. t. 1.
TRITHECA, Miq. Fl. Ind. Bat. i. 614 (1855).
WINTERLIA, Spreng. Syst. i. 519 (1825).
acutidens, Miq. Fl. Ind. Bat. i. 1. 617 = peploides.
aegyptiaca, Kotschy, ex Koehne, in Engl. Bot. Jahrb. i. (1881) 253, 254 = urceolata, apiculata.
aegyptiaca, Willd. Hort. Berol. t. 6 = salicifolia.
alata, *Steud. Nom.* ed. II. i. 76.—Reg. Caucas.
anagalloides, Sond. in Linnaea, xxiii. (1850) 40 = Nesaea anagalloides.
apiculata, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 254.—Afr. trop.
arenaria, H. B. & K. Nov. Gen. et Sp. vi. 190 = senegalensis.
aspera, Guill. & Perr. Tent. Fl. Seneg. i. 304 = Nesaca aspera.
attenuata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 278 = baccifera.
auriculata, Willd. Hort. Berol. i. 7. t. 7 = senegalensis.
australasica, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 41 = multiflora.
baccifera, Roth, Sp. Nov. 100 = leptopetala.
baccifera, *Linn. Sp. Pl.* 120.—Geront. calid.
baccifera, Pollini, Fl. Veron. i. 178 = salicifolia.
Boraei, Guép. Suppl. Fl. Maine-et-Loire, 39 = Lythrum nummularifolium.
borysthenica, DC. Prod. iii. 78 = Lythrum nummularifolium.
borysthenica, Kar. & Kir. in Bull. Soc. Nat. Mosc. xiv. (1841) 421 = Peplis alternifolia.
caspia, Hohen. Enum. Talysch. 133 = baccifera.
caspica, Bieb. Fl. Taur. Cauc. ii. 457; Suppl. 111 = salicifolia.
catholica, Cham. & Schlecht. in Linnaea, ii. (1827) 378 = latifolia.
catholica, Hook. & Arn. ex Secm. Bot. Voy. Herald, 284 = latifolia.
coccinea, Pers. Syn. i. 147 = octandra.
coccinea, Rottb. Pl. Hort. Univ. (Havn.) Programm. Descr. (1773) 7 = latifolia.
cordata, *Wight & Arn. Prod.* 304.—Ind. or.
crassicaulis, Guill. & Perr. Tent. Fl. Seneg. i. 303 = Nesaea polyantha.
crinipes, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 49 = Nesaea crinipes.
cryptantha, *Baker, in Journ. Linn. Soc.* xxi. (1884) 345.—Madag.
debilis, Ait. Hort. Kew. ed. I. i. 163 = baccifera.
densiflora, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 569 = salicifolia.
densiflora, Roth, Sp. Nov. 99 = pentandra.
dentifera, *A. Gray, Pl. Wright.* ii. 55.—Am. bor.
dentelloides, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) II. 76.—Bengal.
diandra, *F. Muell. Fragm.* iii. 108.—Austral.
diffusa, Willd. Enum. Hort. Berol. i. 167 = senegalensis.
dodecandra, DC. in Mém. Soc. Phys. Genèv. iii. II. (1825) 89. t. 2 = Nesaea Candollei.
elatinoides, *DC. l. c.* 92. t. 3. f. B.—Afr. trop.
elatinoides, Schimp. ex Hiern, in Oliver, Fl. Trop. Afr. ii. 468 (= *Rotala stagnina*).
elongata, Blume, Mus. Bot. Lugd. Bat. ii. 135 = peploides.

AMMANNIA :—
filiformis, Baill. Hist. Pl. vi. 438 (= *Rotala filiformis*).
filiformis, DC. in Mém. Soc. Phys. Genèv. iii. II. (1825) t. 79 = senegalensis.
fimbriata, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 569 = pentandra.
floribunda, *C. B. Clarke, l. c.* 567.—Ind. or.
floribunda, Guill. & Perr. Tent. Fl. Senceg. i. 302 = senegalensis, multiflora.
glauca, Wall. Cat. n. 2100 = salicifolia.
gracilis, Guill. & Perr. Tent. Fl. Seneg. i. 301 = senegalensis
hastata, DC. Prod. iii. 78 = latifolia.
hexandra, Wall. Cat. n. 2103 = pentandra.
Heyneana, Wall. l. c. 2104 = pentandra.
Hildebrandtii, *Koehne, in Engl. Bot. Jahrb.* i. (1881) 257.—Afr. trop.
humilis, *Michx. Fl. Bor. Am.* i. 99.—Am. bor.
hyrcanica, Fisch. ex Steud. Nom. ed. II. i. 77 = auriculata.
illecebroides, Arn. ex Koehne, in Engl. Bot. Jahrb. i. (1881) 161 (= *Rotala decussata*).
indica, Lam. Illustr. i. 311. n. 1555 = baccifera.
japonica, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 261.—Japon.
lanceolata, *Heyne, in Wall. Cat.* n. 2106.—Ind. or.
latifolia, *Linn. Sp. Pl.* 119.—Am. bor.; Ind. occ.
latifolia, Wall. Cat. n. 2096 = peploides.
latifolia, Walp. Rep. ii. 102 = sanguinolenta.
leptopetala, *Blume, Mus. Bot. Lugd. Bat.* ii. 134.—Java.
lingulata, Griseb. Cat. Pl. Cub. 106 = latifolia.
littorea, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 261.—Japon.
loandensis, Welw. ex Hiern, in Oliver, Fl. Trop. Afr. ii. 480 = Nesaea loandensis.
longipes, *Wright, in Sauv. Fl. Cub.* 53.—Ind. occ.
lythrifolia, Salisb. Prod. 65 = latifolia.
madagascariensis, Boiv. ex Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 129 = multiflora.
microcarpa, DC. in Mém. Soc. Phys. Genèv. iii. II. (1825) 93 = multiflora ?
monoflora, Blanco, Fl. Filip. ed. I. 64 = latifolia ?
multiflora, *Roxb. Hort. Beng.* 11; *Fl. Ind. ed. Carey*, i. 447.—Ind. or.
nana, Roxb. ll. cc. 11, 448 = peploides.
nana, Roxb. in Wall. Cat. n. 2105 = pentandra.
Nuttallii, A. Gray, Man. Bot. N. U. St. ed. IV. add. p. xcii = Didiplis linearis.
occidentalis, DC. Prod. iii. 78 = humilis.
octandra, *Linn. f. Suppl.* 127.—Ind. or.; Malaya.
octandra, Cham. & Schlecht. in Linnaea, ii. (1827) 376 = latifolia.
pallida, Lehm. Sem. Hort. Hamb. (1823) 3; (Linnaea, iii. (1829) 9) = latifolia.
parviflora, DC. Prod. iii. 77 = multiflora.
passerinoides, Welw. ex Hiern, in Oliver, Fl. Trop. Afr. ii. 480 = Nesaea passerinoides.
pentandra, *Roxb. Fl. Ind.* i. 448.—Geront. trop.
peploides, *Spreng. Syst.* i. 444.—Oriens; As. trop.
pinnatifida, *Linn. f. Suppl.* 127.—Java.
polystachya, Wall. Cat. n. 2094 = peploides.
Portula, Baill. in Bull. Soc. Linn. Par. i. (1874) 87 = Peplis Portula.
Prieuriana, Guill. & Perr. Tent. Fl. Seneg. i. 303 = senegalensis.
prostrata, Buch.-Ham. ex Blume, Mus. Bot. Lugd. Bat. ii. 132 = baccifera.
purpurea, Lam. Encyc. i. 131 = humilis.
pusilla, Sond. in Linnaea, xxiii. (1848) 40 = senegalensis.
pygmaea, *Kurz, in Journ. Bot.* v. (1867) 376.—Ind. or.
racemosa, Poir. Encyc. Suppl. i. 329 = arenaria.
racemosa, Roth, Catalect. fasc. iii. 25 = octandra.
ramosa, Hill, Veg. Syst. xi. 14 = humilis.
ramosior, Blanco, Fl. Filip. ed. II. 64 = peploides.
ramosior, Linn. Sp. Pl. 120 = humilis.
ramosior, Linn. Mant. ii. 332 = latifolia.
repens, Rottl. ex Mart. in Acad. Muench. Phil. vi. (1820) 150, ex DC. Prod. iii. 80 = peploides.
retusa, Koehne, in Engl. Bot. Jahrb. i. (1881) 254.—Afr. trop.
Ritchiei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 566.—Ind. or.
robusta, Heer, Ind. Sem. Hort. Turic. (1842) = latifolia.
rosea, Poir. Encyc. Suppl. i. 329 (= *Rotala mexicana*).

AMMANNIA :—
Rotala, *F. Muell. Fragm.* iii. 108.—Geront. trop.
rotundifolia, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 220.—Ind. or.; China.
rubra, Buch.-Ham. l. c. = pentandra.
sagittata, DC. Prod. iii. 80 = latifolia.
salicifolia, Hiern, in Oliver, Fl. Trop. Afr. ii. 478 = baccifera.
salicifolia, *Monti, in Comm. Bonon.* i. (1767) 112.—As. & Afr. trop.
salsuginosa, *Guill. & Perr. Tent. Fl. Seneg.* i. 302.—Afr. trop.
sanguinolenta, Cham. & Schlecht. in Linnaea, v. (1830) 568 = auriculata.
sanguinolenta, Heyne, ex Steud. Nom. ed. II. i. 77 = salicifolia.
sanguinolenta, Sw. Prod. Veg. Ind. Occ. 33 = latifolia.
sanguinolenta, Hook. & Arn. ex Scem. Bot. Voy. Herald, 284 = latifolia.
sarcophylla, Welw. ex Hiern, in Oliver, Fl. Trop. Afr. ii. 479 = Nesaea sarcophylla.
senegalensis, *Lam. Illustr.* i. 312. n. 155. t. 77. f. 2.—Geront. trop.
simpliciuscula, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 54.—Ind. or.
stylosa, Fisch. et Mey. Ind. Sem. Hort. Petrop. vii. 41 = latifolia.
subrotunda, *Wall. ex Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 55.—Ind. or.
subspicata, Benth. in Hook. Lond. Journ. Bot. i. (1843) 484 = rotundifolia.
subspicata, Hohen. ex Hook. f. Fl. Brit. Ind. ii. 568 = pentandra.
tenella, *Guill. & Perr. Tent. Fl. Seneg.* i. 297.—Afr. trop.
tenuis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 567.—Ind. or.
texana, Scheele, in Linnaea, xxi. (1848) 588 = latifolia.
triflora, R. Br. ex Benth. Fl. Austral. iii. 297 = Nesaea lanceolata.
triflora, Wall. Cat. n. 6323 = Nesaea triflora.
undulata, C. A. Mey. Ind. Sem. Hort. Petrop. ix. 56 = senegalensis.
urceolata, *Hiern, in Oliver, Fl. Trop. Afr.* ii. 478.—Afr. trop.
verticillaris, Baill. Hist. Pl. vi. 439 (= *Rotala verticillaris*).
verticillata, Boiss. Fl. Orient. ii. 743 = baccifera.
verticillata, Lam. Encyc. i. 131 = salicifolia.
verticillata, Wight & Arn. Prod. i. 304 = lanceolata.
vesicatoria, Roxb. Hort. Beng. 11; Fl. Ind. ed. Carey, i. 447 = baccifera.
viridis, Willd. ex Hornem. Hort. Hafn. i. 146 = baccifera.
Wallichii, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 84 = Hydrolythrum Wallichii.
Wormskioldii, *Fisch. et Mey. Ind. Sem. Hort. Petrop.* vii. 42.—Bras.
Wrightii, *A. Gray, Pl. Wright.* ii. 55.—Am. bor.

AMMANTHUS, Boiss. & Heldr. Diagn. Ser. I. xi. 18 (1849) = **Chrysanthemum**, Tourn. (Compos.).
ageratifolius, Boiss. & Heldr. ex Boiss. Fl. Orient. iii. 358 = C. maritimum.
filicaulis, Boiss. & Heldr. ex Boiss. Diagn. Ser. I. xi. 19.—Ins. Creta.
maritimus, Boiss. & Heldr. l. c. = C. maritimum.

AMMI, [Tourn.] Linn. Syst. ed. I. (1735); Gen. ed. I. 72 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 889.
Dasyspermum, Neck. Elem. i. 276 (1790).
Gohoria, Neck. l. c. 172 (1790).
Visnaga, Riv. ex Gaertn. Fruct. i. 92. t. 21 (1788).
acaule, Sprengel, in Roem. & Schult. Syst. vi. 530 = Chamaesciadium acaule.
anethifolium, *Lam. Encyc.* i. 132.—Orient; Sibir.
anethifolium, Ledeb. Fl. Ross. ii. 246 = Artedia squamata.
Boeberi, Hellan. ex Hoffm. Gen. Umbell. p. xviii; Hornem. Suppl. 32 = majus.
Broussonetii, DC. Prod. iv. 113 = majus?
capillaceum, *Michx. Fl. Bor. Am.* i. 164 = Discopleura capillacea.
cicutaefolium, Willd. ex Roem. & Schult. Syst. vi. 531 = majus.
cicutarium, Willd. ex Roem. & Schult. l. c.—Am. calid.

AMMI :—
copticum, Linn. Mant. 56 = Carum copticum.
costatum, Ell. Sketch Fl. Carol. i. 350 = Discopleura capillacea.
crinitum, *Guss. Pl. Rar.* 128. t. 25.—Calabr.; Sicil.
daucifolium, Buch.-Ham. in Wall. Cat. n. 7209 = Seseli daucifolium.
daucifolium, Scop. Fl. Carn. ed. II. i. 207 = Libanotis athamantoides.
daucoides, Salzm. ex DC. Prod. iv. 113 = Ammiopsis daucoides.
dilatatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Visnaga.
divaricatum, Michx. Fl. Bor. Am. i. 168 = Apium divaricatum.
elatum, Salisb. Prod. 162 = majus.
ferulaceum, Link, Enum. Hort. Berol. i. 266 = Ligusticum ferulaceum.
ferulaefolium, Lag. ex Loud. Hort. Brit. 103 = praec.?
glaucifolium, Blanco, Fl. Filip. ed. I. 213 = Carum copticum.
glaucifolium, Linn. Sp. Pl. 243 = majus.
Huntii, *H. C. Wats. in Hook. Lond. Journ. Bot.* vi. (1847) 384.—Ins. Azor.
indicum, *Buch.-Ham. in Wall. Cat.* n. 7208.—Ind. or.
intermedium, DC. Prod. iv. 113 = majus.
majus, *Linn. Sp. Pl.* ed. I. 243.—Europ.; Orient; Afr. bor.
majus, Walt. Fl. Carol. 113 = Discopleura capillacea.
meoides, Pers. Syn. i. 308 = anethifolium.
minus, *Griseb. Hort. Gotting.* (1876) 8.—Europ.
pauciradiatum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 322 = majus.
petroselinoides, Presl, ex DC. Prod. iv. 102 = Carum peregrinum.
procerum, *Lowe, Man. Fl. Mad.* 351.—Madera.
pumilum, *DC. Prod.* iv. 113.—Lusitan.
pyrenaicum, *Lapeyr. Hist. Abr. Pl. Pyr.* 145.—Pyren.
rubricaule, Hornem. Hort. Hafn. i. 272 = Discopleura capillacea.
saxatile, *Turcz. ex Bess. in Flora*, xvii. (1834) i.; *Beibl.* 13, nomen.—Rossia.
trachycarpum, Fisch. ex Boiss. Fl. Orient. ii. 855 = Aphanopleura trachysperma.
Visnaga, *Lam. Fl. Fr.* iii. 462.—Reg. Mediterr.

AMMIANTHUS, Spruce, ex Benth. & Hook. f. Gen. ii. 98 (1873) = **Retiniphyllum**, Humb. & Bonpl. (Rubiac.).

AMMIOPSIS, Boiss. Diagn. Ser. II. ii. 96 (1856). *UMBELLIFERAE*, Benth. & Hook. f. i. 927.
daucoides, *Boiss. l. c.* 97.—Mauritan.

AMMIOS, Moench, Meth. 99 (1794) = **Carum**, Tourn. (Umbellif.).
muricata, Moench, l. c. = C. copticum.

AMMOBIUM, R. Br. in Bot. Mag. t. 2459 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 316.
alatum, *R. Br. l. c.*—Austral.
craspedioides, *Benth. Fl. Austral.* iii. 584.—Austral.
plantagineum, *G. Don, in Loud. Hort. Brit.* 333.—Austral.
spathulatum, Gaudich. Voy. Freyc. Bot. 467. t. 91 = alatum.

AMMOBROMA, Torr. in Ann. Lyc. N. York, viii. (1867) 51. t. 1. *LENNOACEAE*, Benth. & Hook. f. ii. 622.
Sonorae, *Torr. l. c.* 52.—N. Mexic.

AMMOCHARIS, Herb. App. 17 (1821). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 727.
Palinetes, Salisb. Gen. Pl. Fragm. 116 (1866).
coranica, Herb. App. 17 = falcata.
falcata, *Herb. l. c.*—Afr. austr.
longifolia, M. Roem. Amar. 62 = Crinum longifolium.
Slateriana, Kunth, Enum. Pl. v. 613 = Brunsvigia Slateriana.

AMMOCHLOA, Boiss. Diagn. Ser. I. xiii. 51 (1853). *GRAMINEAE*, Benth. & Hook. f. iii. 1181.
Cephalochloa, Coss. & Dur. in Ann. Sc. Nat. Sér. IV. i. (1854) 229.
palaestina, Boiss. Diagn. Ser. I. xiii. 52 = subacaulis.

AMMOCHLOA :—
pungens, *Boiss. Diagn.* Ser. I. xiii. 52.—Afr. bor.
subacaulis, *Balansa, ex Coss. in Bull. Soc. Bot. Fr.* i. (1854) 317; iv. (1857) 399.—Syria; Afr. bor.

AMMODAUCUS, Coss. & Dur. in Bull. Soc. Bot. Fr. vi. (1859) 393 = **Daucus**, Linn. (Umbellif.).
leucotrichus, *Coss. l. c. et* xxii. (1875) 60.—Algeria.

AMMODENDRON, Fisch. ex DC. Prod. ii. 523 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 554.
caspium, Eichw. Casp.-Cauc. 7 = Eichwaldii.
Conollyi, *Bunge, ex Boiss. Fl. Orient.* ii. 627.—Turkest.
Eichwaldii, *Ledeb. & Mey. in Eichw. Casp.-Cauc.* 37. t. 33.—Reg. Casp.
Karelini, Bunge, Pl. Rel. Lehm. 69 = Conollyi.
Karelinii, *Fisch. & Mey. in Ledeb. Fl. Ross.* i. 717.—Reg. Casp.
Lehmanni, *Bunge, ex Boiss. Fl. Orient.* ii. 626.—Turkest.
persicum, *Bunge, ex Boiss. l. c.* 627.—Persia.
Sieversii, *DC. Prod.* ii. 523.—Soongaria.
Zablotskii, *Fisch. & Mey. ex Kar. in Bull. Soc. Nat. Mosc.* (1839) 150, nomen.—Rossia.

AMMODENIA, Patrin. ex J. G. Gmel. Fl. Sibir. iv. 160 (1769) = **Arenaria**, Linn. (Caryophyll.).
peploides, Rupr. in Beitr. Pfl. Russ. Reich. ii. 25 = Ar. peploides.

AMMODIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 321 = **Chrysopsis**, Nutt. (Compos.).
oregana, Nutt. l. c. = C. oregana.

AMMODYTES, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 263 = **Astragalus**, Tourn. (Legum.).
melilotoides, Stev. l. c. = Astr. melilotoides.

AMMOGETON, Schrad. Ind. Sem. Hort. Goett. (1833) 1 = **Troximon**, Nutt. (Compos.).
scorzoneraefolius, Schrad. l. c. = T. glaucum.

AMMOIDES, Adans. Fam. ii. 96 (1763) = **Carum**, Tourn. (Umbellif.).

AMMOLIRION, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 515 = **Eremurus**, Bieb. (Liliac.).
stenophyllum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 218 = E. Bungei.
Steveni, Kar. & Kir. l. c. = E. inderiensis.

AMMONALIA, Desv. ex Endl. Gen. 966, in syn. (1840) = **Arenaria**, Linn. (Caryophyll.).

AMMOPHILA, Host, Gram. Austr. iv. 24. t. 41 (1809). *GRAMINEAE*, Benth. & Hook. f. iii. 1153.
Psamma, Beauv. Agrost. 143. t. 6. f. 1. (1812).
arenaria, Link, Hort. Berol. i. 105 = arundinacea.
arundinacea, *Host, Gram. Austr.* iv. 24. t. 41.—Europ.; Am. bor.
baltica, *Link, Hort. Berol.* i. 105; *Dum. Fl. Belg.* 152.—Europ.
brevipilis, *Benth. ex Vasey, Cat. Gram. U. St.* 52.—Am. bor.
Curtissii, *Vasey, in Torrey Bot. Club*, xi. (1884) 7.—Florida.
longifolia, *Vasey, Agric. Grasses U. St.* 71.—Am. bor.

AMMORRHIZA, Ehrh. Phytophyl. (1789); Beitr. iv. 146 = **Carex**, Linn.

AMMOSELINUM, Torr. & Gray, in Pacif. Rail. Rep. ii. 165 (1855). *UMBELLIFERAE*, Benth. & Hook. f. i. 1009.
Popei, *Torr. & Gray, l. c.*—Texas.

AMMOSERIS, Endl. Gen. 500 (1838) = **Launaea**, Cass. (Compos.).
asplenifolia, D. Dietr. Syn. Pl. iv. 1319 = L. asplenifolia.
bellidifolia, D. Dietr. l. c. 1320 = L. bellidifolia.
Dregeana, D. Dietr. l. c.—Afr. austr.
nudicaulis, D. Dietr. l. c. 1319 = L. nudicaulis.
patens, D. Dietr. l. c.—Ind. or.
sarmentosus, D. Dietr. l. c. = L. pinnatifida.
surinamensis, D. Dietr. l. c.—Guiana.

AMMOSPERMA, Hook. f. in Benth. & Hook. f. Gen. i. 82 (1862). *CRUCIFERAE*, Benth. & Hook f. i. 82.
cinerea, *Hook. f. l. c.; Baill. Hist. Pl.* iii. 278.—Afr. bor.

AMMOTHAMNUS, Bunge, Enum. Pl. Lehm. 67. t. 12 (1847). *LEGUMINOSAE*, Benth. & Hook. f. i. 555.
gibbosus, *Boiss. Fl. Orient.* ii. 628.—Assyria.
Lehmanni, *Bunge, l. c.* 243.—Turkest.

AMMYRSINE, Pursh, Fl. Am. Sept. i. 301 (1814) = **Leiophyllum**, Hedw. f. (Ericac.).
buxifolia, Pursh, l. c. = L. buxifolium.
Lyoni, Sweet, Hort. Brit. ed. II. 344 = L. buxifolium.
prostrata, Sweet, ex Steud. Nom. ed. II. i. 78 = L. buxifolium.
thymifolia, Forbes, Hort. Woburn. 93 = L. buxifolium.

AMNI, Brongn. Enum. Genr. 110 (1843) err. typ. = **Ammi**, Tourn.

AMOGETON, Neck. Elem. iii. 267 (1790) = **Aponogeton**, Thunb. (Naiad.).

AMOLEIACHYRIS, Sch. Bip. in Walp. Rep. ii. 951 (1843), sphalm. = AMPHIACHYRIS, Nutt. = **Gutierrezia**, Lag. (Compos.).

AMOMIS, Berg, in Linnaea, xxvii. (1854) 416 = **Pimenta**, Lindl. (Myrtac.).
acris, Berg, l. c. 417 = P. acris.
oblongata, Berg, l. c. 421 (= P. acris?).—Ind. or. cult.; Guiana.
Pimento, Berg, l. c. 418 (= P. acris?).—Ind. occ.; Guiana.
pimentoides, Berg, l. c. 420 = P. acris.

AMOMOPHYLLUM, Engl. in Gard. Chron. (1877) i. 139 = **Spathiphyllum**, Schott (Aroid.).
floribundum, Engl. l. c. 140 = S. floribundum.
Patini, Engl. l. c. 139 = S. Patini.

AMOMUM, Linn. Mus. Cliff. 26 (1736); Gen. ed. I. 330 (1737). *SCITAMINEAE*, Benth. & Hook. f. iii. 644.
ACHASMA, Griff. Notul. iii. 426 (1851).
ALEXIS, Salisb. in Trans. Hort. Soc. i. (1812) 284.
CENOLOPHON, Horan. Prod. Monog. Scitam. 36 (1862).
DIRACODES, Blume, Enum. Pl. Jav. i. 55 (1830).
DONACODES, Blume, l. c. 54 (1830).
ETLINGERA, Giseke, Prael. Linn. 199, 209 (1792)
GEANTHUS, Reinw. in Syll. Ratisb. ii. (1828) 5.
GEOCALLIS, Horan. Prod. Monog. Scit. 33 (1862).
GREENWAYA, Giseke, Prael. Linn. 199, 206 (1792).
HORNSTEDTIA, Retz. Obs. vi. 18 (1791).
MAROGNA, Salisb. in Trans. Hort. Soc. i. (1812) 283.
MEISTERA, Giseke, Prael. Linn. 199, 205 (1792).
NICOLAIA, Horan. Prod. Monog. Scitam. 32. t. 1 (1862).
PALUDANA, Giseke, Prael. Linn. 199, 207 (1792).
PHAEOMERIA, Lindl. Introd. Nat. Syst. ed. II. 446 (1836).
STENOCHASMA, Griff. Notul. iii. 431 (1851).
TORYMENES, Salisb. in Trans. Hort. Soc. i. (1812) 283.
WURFBAINIA, Giseke, Prael. Linn. 199, 206 (1792).
ZEDOARIA, Rafin. Fl. Tellur. iv. 53 (1836).
aculeatum, *Roxb. in As. Res.* xi. (1810) 344.—Malaya.
acuminatum, *Thw. Enum. Pl. Zeyl.* 317.—Zeylan.
Afzelii, Rosc. in Trans. Linn. Soc. viii. (1807) 354 = A. Granum-paradisi.
Alpinia, Rottb. in Coll. Soc. Med. Havn. ii. (1775) 245 t. 2 = Renealmia racemosa.
angustifolium, Hanb. in Journ. Linn. Soc. xiii. (1873) 155 = Danielli.
angustifolium, Salisb. Prod. 4 = Zingiber officinale.
angustifolium, *Sonner. Voy. Indes*, iii. 276.—Madag.
aquaticum, *Raeusch. Nom.* ed. III. 1.—Zeylan.
arboreum, Lour. Fl. Cochinch. 7 = Costus Loureiri.
aromaticum, Roxb. Hort. Beng. 1; Fl. Ind. i. 44.—Ind. or.
arundinaceum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 109.—Afr. trop.
Benthamianum, *Trim. in Journ. Bot.* xxiii. (1885) 265.—Zeylan.

AMOMUM:—
bifidum, Stokes, Bot. Comment. i. 163 = Alpinia Allughas.
biflorum, *Jack, in Malay. Misc.* i. (1820) n. 1. p. 2.—Penang.
cannaecarpum, *Benth. & Hook. f. Gen.* iii. 644.—Ind. or.
capitatum, *Roxb. ex Link, Jahrb.* ii. (1811) 64.—Ind. or.
Cardamomum, Koen. in Retz. Obs. fasc. iii. 59 = seq.
Cardamon, *Linn. Sp. Pl.* 1.—Ind. or.; Malaya.
Cardamomum, Willd. Enum. Hort. Berol. Suppl. 1 = Alpinia striata.
cereum, *Hook. f. in Kew Journ. Bot.* iv. (1854) 296.—Afr. trop.
Cevuga, *Seem. Fl. Vit.* 291.—Ins. Fiji.
ciliatum, *Blume, Enum. Pl. Jav.* 49.—Java.
citratum, Pereira, Mat. Med. ii. 1137 = cereum.
Clusii, Hook. f. Bot. Mag. t. 5250 = Danielli.
Clusii, *Sm. in Rees. Cyclop.* xxxix. n. 4.—Afr. trop.
coccineum, *Benth. & Hook f. Gen.* iii. 644.—Java.
compactum, Roem. & Schult. Syst. i. 28 = A. Cardamon.
corynostachyum, *Wall. Pl. As. Rar.* i. 48 t. 58.—Ind. or.
costatum, *Benth. & Hook. f. Gen.* iii. 235.—Ind. or.
Curcuma, Jacq. Hort. Vindob. iii. 5. t. 4 = Curcuma longa.
Dallachyi, *F. Muell. Fragm.* viii. 25.—Austral.
Danielli, *Hook. f. in Kew Journ.* iv. (1854) 129. t. 5.—Afr. trop.
dealbatum, *Roxb. Hort. Beng.* 1; *Fl. Ind.* i. 43.—Ind. or.
echinatum, Willd. Sp. Pl. i. 8.—Ind. or.
elatum, Salisb. Prod. 5 = A. Granum-paradisi.
Ensal, Raeusch. Nom. ed. III. 1 = Elettaria Cardamomum.
exscapum, Sims, in Kon. & Sims, Ann. Bot. i. (1805) 549 = A. Granum-paradisi.
fasciculatum, *Benth. & Hook. f. Gen.* iii. 645.—Ind. or.
Fenzlii, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) ii. 154.—Ins. Nicobar.
floribundum, *Benth. & Hook. f. Gen.* iii. 644.—Zeylan.
foetens, *Benth. & Hook. f. l. c.*—Java.
fulviceps, *Thw. Enum. Pl. Zeyl.* 317.—Zeylan.
Galanga, Lour. Fl. Cochinch. 5 = Alpinia Galanga.
giganteum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 109.—Afr. trop.
glaberrimum, *Benth. & Hook. f. Gen.* iii. 644.—Malaya.
globosum, Lour. Fl. Cochinch. 4 = Alpinia globosa.
gracile, *Blume, Enum. Pl. Jav.* 49.—Java.
gramineum, *Wall. Cat.* n. 6558.—Ind. or.
graminifolium, *Thw. Enum. Pl. Zeyl.* 430.—Zeylan.
grandiflorum, Sm. Exot. Bot. ii. 103. t. 11 = A. Granum-paradisi.
Granum-paradisi, *Linn. Sp. Pl.* 2.—Afr. trop.
heteranthum, *Blume, Enum. Pl. Jav.* 49.—Java.
hirsutum, Lam. Illustr. i. 134. t. 3 = Costus speciosus.
hirsutum, Lour. Fl. Cochinch. 5 = Alpinia hirsuta.
hypoleucum, *Thw. Enum. Pl. Zeyl.* 318.—Zeylan.
involucratum, *Benth. & Hook. f. Gen.* iii. 644.—Zeylan.
Koenigii, *Roem. & Schult. Syst.* i. 30.—Ind. or.
latifolium, *Afzel. Remed. Guin.* i. 5.—Afr. trop.
latifolium, Salisb. Prod. 4 = Curcuma Zedoaria.
Leonurus, *Koen. ex Retz. Obs. fasc.* iii. 69.—Malacca.
limbatum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 110.—Afr. trop.
linguiforme, *Benth. & Hook. f. Gen.* iii. 644.—Ind. or.
littorale, *Koen. ex Retz. Obs. fasc.* iii. 52.—Burma.
longiscapum, *Hook. f. in Journ. Bot.* vi. (1854) 296.—Afr. trop.
macrospermum, Sm. in Rees, Cyclop. xxxix. n. 3 = Clusii.
madagascariense, Lam. Illustr. t. 2. f. 1 = angustifolium.
magnificum, *Benth. & Hook. f. Gen.* iii. 644.—Ins. Maurit.
Mannii, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 110.—Afr. trop.
marginatum, *Roxb. ex Spreng. Schrad. & Link, Jahrb.* ii. (1811) 64.—Ind. or.
masticatorium, *Thw. Enum. Pl. Zeyl.* 317.—Zeylan.
maximum, *Roxb. in As. Res.* xi. (1810) 344.—Malaya.
medium, Lour. Fl. Cochinch. 4 = Hellenia alba.
Melegueta, *Rosc. Monog. Scitam.* t. 98.—Afr. trop.
Mioga, Thunb. Fl. Jap. 14 = Zingiber Mioga.
montanum, Koen. ex Retz. Obs. fasc. iii. 51 = Zingiber Cassumunar.

AMOMUM:—
muricatum, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 59.—Ind. or.
nemorale, *Benth. & Hook. f. Gen.* iii. 644.—Zeylan.
nemorosum, Boj. Hort. Maurit. 327 = Danielli.
nigrum, Raeusch. Nom. ed. III. 1 = Alpinia Allughas.
nutans, Roem. & Schult. Syst. i. Mant. 40 = Alpinia nutans.
palustre, *Afzel. Stirp. Med. Nov.* 9.—Afr. trop.
parviflorum, Presl, Rel. Haenk. ii. 112. t. 19 = Alpinia parviflora.
petiolatum, Lam. Encyc. i. 136 = Costus spicatus.
pierardioides, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 58.—Ind. or.
pilosum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 110.—Afr. occ.
pterocarpum, *Thw. Enum. Pl. Zeyl.* 317.—Zeylan.
pulchellum, Thw. l. c. 318 = Cyphostigma pulchellum.
pyramidale, Lam. Encyc. i. 135 = Renealmia racemosa.
racemosum, Lam. Illustr. t. 2. f. 2 = A. Cardamon.
racemosum, Ruiz & Pav. Fl. Per. i. 2 = Renealmia Ruiziana.
Renealmia, Lam. Illustr. i. 5 = Renealmia exaltata.
repens, Sonner. Voy. Indes, iii. 272 = Alpinia Cardamomum
roseum, *Benth. & Hook. f. Gen.* iii. 644.—Malaya.
roseum, Roxb. Pl. Corom. 2. t. 126 = Zingiber roseum.
Rumphii, Sm. in Rees, Cyclop. xxxix. n. 15 = Elettaria musacea.
rufescens, *Benth. & Hook. f. Gen.* iii. 645.—Zeylan.
Sceptrum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 109.—Afr. trop.
Scottii, F. Muell. Fragm. viii. 24 = Elettaria Scottiana.
scyphiferum, *Koen. ex Retz. Obs. fasc.* iii. 68.—Malacca.
sericeum, Roxb. Hort. Beng. 1; Fl. Ind. i. 44 = dealbatum.
spirale, Hort. ex Steud. Nom. ed. II. i. 78 = Costus Pisonis.
spurium, J. F. Gmel. Syst. i. 6 = Zingiber Zerumbet.
spurium, *Roem. & Schult. Syst.* i. 30.—Ind. or.
striatum, Stokes, Bot. Comment. i. 159 = Renealmia sylvestris.
strobilaceum, *Sm. in Rees, Cyclop.* xxxix. n. 8.—Afr. trop.
subsericeum, *Oliver & Hanb. in Journ. Linn. Soc.* vii. (1864) 110.—Afr. trop.
subulatum, *Roxb. Pl. Corom.* t. 227; *Fl. Ind.* i. 43.—Reg. Himal.
sylvestre, Poir. Encyc. Suppl. v. 548 = Zingiber Zerumbet.
sylvestre, Sw. Prod. Veg. Ind. Occ. 11 = Renealmia sylvestris.
Taraca, Horan. Prod. 30 = Alpinia Allughas.
Taracca, Roxb. ex Rosc. Monog. Scitam. in Obs. de Amomo = praec.
thyrsoideum, Ruiz & Pav. Fl. Per. i. 2. t. 2 = Renealmia thyrsoidea.
uliginosum, *Koen. ex Retz. Obs. fasc.* iii. 56.—Ind. or.
uncinatum, Stokes, Bot. Comment. i. 161 = A. Cardamon.
villosum, *Lour. Fl. Cochinch.* i. 4.—Ind. or.; China.
vitellinum, *Lindl. in Journ. Hort. Soc.* ii. (1847) 245.—Ind. or.?
xanthioides, *Wall. Cat.* n. 6557.—Burma.
xanthorhiza, Roxb. ex Steud. Nom. ed. II. i. 78 = Zingiber Cassumunar.
Zedoaria, Berg. Mat. Med. 41 = Curcuma Zedoaria.
Zerumbet, Koen. ex Retz. Obs. fasc. iii. 55 = Curcuma Zedoaria.
Zerumbet, Linn. Sp. Pl. 1 = Zingiber Zerumbet.
Zingiber, Blanco, Fl. Filip. 2 = Zingiber Blancoi.
Zingiber, Linn. Sp. Pl. 1 = Zingiber officinale.
Zinziba, Hill, Veg. Syst. xvi. 50 = praec.

AMONIA, Nestl. Mon. Potent. 17 (1816) = **Agrimonia**, Linn. (Rosac.).
agrimonoides, Steud. Nom. ed. I. 39, 19 = Agr. Eupatoria.

AMOORA, Roxb. Pl. Corom. iii. 54. t. 258 (1819). *MELIACEAE*, Benth. & Hook. f. i. 334.
ANDERSONIA, Roxb. Hort. Beng. 87 (1814); Fl. Ind. ii. 212 (1832).
APHANAMIXIS, Blume, Bijdr. 165 (1825).
MONOSOMA, Griff. Notul. iv. 502 (1854).
NIMMOIA, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 13.

AMOORA:—
ORAOMA, Turcz. in Bull. Soc. Nat. Mosc. (1858) I. 411.
SPHAEROSACME, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 429 (1824).
amboinensis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 36.—Amboina.
Aphanamixis, *Schult. f. Syst.* vii. 1621.—Malaya.
auriculata, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 37 = cucullata.
Balanseana, *C. DC. in DC. Monog. Phan.* i. 590.—N. Caled.
borneensis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 36.—Borneo.
canarana, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 560.—Ind. or.
Championii, Benth. & Hook. f. Gen. i. 335; Hiern, in Hook. f. l. c. 562 = Pseudocarapa Championii.
chittagonga, *Hiern, in Hook. f. l. c.* 559.—Ind. or.
cucullata, *Roxb. Pl. Corom.* iii. 54. t. 258.—Ind. or.
Cumingiana, *C. DC. in DC. Monog. Phan.* i. 580.—Ins. Philipp.
decandra, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 562.—Ind. or.
dysoxyloides, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 200.—Burma.
filiciformis *Wight, Illustr. Ind. Bot.* i. 147.—Ind. or.
Ganggo, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 123 = Aglaia Ganggo.
grandifolia, Walp. Rep. i. 429; C. DC. in DC. Monog. Phan. i. 581 = A. Aphanamixis.
Korthalsii, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 36.—Borneo.
lactescens, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 200.—Burma.
lanceolata, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 560.—Malacca.
Lawii, *Benth. & Hook. f. Gen.* i. 335.—Ind. or.
laxa, Wight & Arn. Prod. 119 = cucullata.
macrophylla, Nimmo, in J. Grah. Cat. Bomb. Pl. t. 31 = A. Rohituka.
Maingayi, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 562.—Malacca.
moulmeiniana, *C. DC. in DC. Monog. Phan.* i. 584.—Burma.
nitidula, *Benth. Fl. Austral.* i. 383.—Austral.
Perrottetiana, *Steud. Nom.* ed. II. i. 89.—Ins. Philipp.
polystachya, Wight & Arn. Prod. i. 119 = A. Rohituka.
Rohituka, *Wight & Arn. l. c.*—Ind. or.
Roxburghiana, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 211 = Santiria laevigata.
rubescens, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 561.—Singapore.
rubiginosa, *Hiern, l. c.*—Malacca.
spectabilis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 37. Burma.
sumatrana, *Miq. l. c.* 35.—Sumatra.
Teysmanniana, Miq. Fl. Ind. Bat. Suppl. i. 503 = Aglaia Teysmanniana.
timorensis. *Wight & Arn. ex Steud. Nom.* ed. II. i. 78.—Ins. Timor.
tomentosa, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 211 = Santiria tomentosa.
Vieillardi, *C. DC. in DC. Monog. Phan.* i. 591.—N. Caled.

AMOORIA, Walp. Rep. i. 640 (1842), sphalm. = AMORIA, C. Presl, = **Trifolium**, Tourn. (Legum.).

AMORDICA, Neck. Elem. i. 241 (1790) = **Momordica**, Tourn. (Cucurb.).

AMOREA, Moq. in Delile, Cat. Hort. Monsp. (1844) = **Cyclolοma**, Moq. (Chenopod.).
platyphylla, Delile, l. c. = C. platyphyllum.

AMOREUXIA, Moç. & Sesse, ex DC. Prod. ii. 638 (1825). *BIXINEAE*, Benth & Hook. f. i. 124.
EURYANTHE, Cham. & Schlecht. in Linnaea, v. (1830) 224.
malvaefolia, *A. Gray, Pl. Wright.* i. 29.—Am. bor.
palmatifida, Moç. & Sesse, ex DC. Prod. ii. 638 = Schiedeana?
Schiedeana, *Planch. in Hook. Lond. Journ. Bot.* vi. (1846) 140. t. 1.—Am. bor.
Wrightii, *A. Gray, Pl. Wright.* ii. 26.—Am. bor.

AMOREUXIA, Moq. in Mém. Soc. Hist. Nat. Monsp. (1826) = **Cyclolοma**, Moq. (Chenopod.).
platyphylla, Moq. l. c. = C. platyphyllum.

AMORGINE, Rafin. New Fl. Am. iv. 44 (1836) = **Achyranthes**, Linn. (Amaranth.).
albescens, Rafin. l. c.—Am. bor.

AMORIA, C. Presl, Symb. Bot. 45 (1830) = **Trifolium**, Tourn. (Legum.).
angulata, C. Presl, l. c. 47 = T. angulatum.
Biasollettiana, C. Presl, l. c. = T. repens.
caespitosa, C. Presl, l. c. = T. Thalii.
calycina, C. Presl, l. c. 43. t. 30 = T. calycinum.
caroliniana, C. Presl, l. c. 47 = T. carolinianum.
elegans, C. Presl, l. c. = T. elegans.
hybrida, C. Presl, l. c. = T. hybridum.
isthmocarpa, C. Presl, l. c. 50 = T. Jaminianum.
macropoda, C. Presl, l. c. 51. t. 31 = T. macropodum.
Micheliana, C. Presl, l. c. 47 = T. Michelianum.
nigrescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 362 = T. Michelianum.
obcordata, C. Presl, Symb. Bot. i. 47 = T. polymorphum.
pallescens, C. Presl, l. c. = T. pallescens.
parviflora, C. Presl, l. c. = T. parviflorum.
polymorpha, C. Presl, l. c. = T. polymorphum.
repens, C. Presl, l. c. = T. repens.
reflexa, C. Presl, l. c. = T. reflexum.
Thalii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 362 = T. Thalii.
Vivianii, C. Presl, ex Steud. Nom. ed. II. i. 78 = T. caeruleum.

AMORPHA, Linn. Gen. ed. I. 229 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 492
californica, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 306.—Calif.
canescens, *Nutt. in Fras. Cat.* (1813).—Am. bor.
caroliniana, Croom, in Am. Journ. Sc. xxv. (1834) 74 = fruticosa.
coerulea, Lodd. Cat. (1830) = fruticosa.
croceo-lanata, Wats. Dendrol. t. 139 = fruticosa?
cynostachya, M. A. Curt. ex Torr. & Gray, Fl. N. Am. i. 305 (cit. falsa) = fruticosa.
elata, *Bouché, ex Schlecht. in Linnaea*, xxiv. (1851) 687.—Hab.?
elata, Hayne, Dendrol. Fl. 134 = fruticosa.
emarginata, Sweet, Hort. Brit. ed. I. 121. n. 466 = fruticosa.
fragrans, Sweet, Brit. Fl. Gard. iii. t. 241 = fruticosa.
fruticosa, *Linn. Sp. Pl.* 713.—Am. bor.
fruticosa, Torr. Bot. Mex. Bound. 53 = californica.
fruticosa, Thunb. Fl. Jap. 278 = Cladrastis amurensis.
Gaertneri, Hort. ex C. Koch, Dendrol. i. 70 = fruticosa.
Gardneri, Hort. ex C. Koch, l. c. Index, 701 = pracc.
glabra, Desf. Tabl. Hort. Par. ed. I. 192 = fruticosa.
glandulosa, *Blanco, Fl. Filip.* ed. I. 555.—Ins. Philipp.
herbacea, *Walt. Fl. Carol.* 179.—Am. bor.
herbacea, Schlecht. Ind. Sem. Hort. Hal. (1848) 8.—Cf. Linnaea, xxiv. (1851) 185 = fruticosa.
humilis, Tausch, in Flora, xxi. (1838) II. 750 = fruticosa?
laevigata, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 306.—Am. bor.
Lewisii, Lodd. ex Loud. Hort. Brit. Suppl. ii. 609 = fruticosa.
Ludwigii, Hort. ex C. Koch, Dendrol. i. 70 = fruticosa.
lutea, *Rafin. Fl. Ludov.* 105.—Am. bor.
microphylla, *Pursh, Fl. Am. Sept.* ii. 466.—Am. bor.
nana, Nutt. in Fras. Cat. (1813) = fruticosa?
nonperforata, Schk. Bot. Handb. ii. 333 = fruticosa.
ornata, Wender. Ind. Sem. Hort. Marb. (1835) = fruticosa.
paniculata, *Torr. & Gray, Fl. N. Am.* i. 306.—Am. bor.
pedalis, *Blanco, Fl. Filip.* ed. I. 553.—Ins. Philipp.
pendula, Carr. in Rev. Hortic. 1868) 340 = fruticosa?
perforata, Schk. Bot. Handb. ii. 333 = fruticosa.
pubescens, Schlecht. in Linnaea, xxiv. (1851) 691 = fruticosa.
pubescens, Willd. Berl. Baumz. 17 = herbacea.
pumila, Michx. Fl. Bor. Am. ii. 64 = herbacea.
pumila, Schlecht. Ind. Sem. Hort. Hal. (1848) 8.—Cf. Linnaea, xxiv. (1851) 185 = fruticosa.
punctata, *Rafin. New. Fl. Am.* iii. 14.—Am. bor.
Rabiae, *Lex. in La Llave & Lex. Nov. Veg. Desc. fasc.* i. 22.—Mexic.
Roemeriana, Scheele, in Linnaea, xxi. (1848) 461 = paniculata.

AMORPHA:—
tenessensis, Schult. in Kunze, Del. Sem. Hort. Lips. (1848) = fruticosa.
texana, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 452 = laevigata.
tomentosa, *Rafin. Fl. Ludov.* 105.—Am. bor.

AMORPHOCALYX, Klotzsch, in Rich. Schomb. Faun. & Fl. Guian. 1104, nomen, (1848) (Legumin.).
Roraimae, *Klotzsch, l. c.*—Guiana.

AMORPHOPHALLUS, Blume, in Batav. Diar. (1825); ex Decne. Herb. Tim. Descr. 38 (1835). *AROIDEAE*, Benth. & Hook. f. iii. 970.
BRACHYSPATHA, Schott, Syn. Aroid. 35 (1856).
CANDARUM, Reichb. ex Schott, Melet. i. 17 (1832).
CONOPHALLUS, Schott, Syn. Aroid. 34 (1856).
CORYNOPHALLUS, Schott, in Oestr. Bot. Wochenbl. (1857) 389.
HANSALIA, Schott, in Oestr. Bot. Zeitschr. (1858) 82.
HYDROSME, Schott, in Oestr. Bot. Wochenbl. (1857) 389.
KUNDA, Rafin. Fl. Tellur. ii. 82 (1836).
PROTEINOPHALLUS, Hook. f. Bot. Mag. t. 6195 (1875).
PYTHION, Mart. in Flora, xiv. (1831) 459.
TAPEINOPHALLUS, Baill. Dict. Bot. i. Ic. sine desc. (1877).
Beccarii, *Engl. in Engl. Bot. Jahrb.* i. (1881) 182.—Sumatra.
Blumei, Schott, in Ann. Mus. Lugd. Bat. i. 124 = giganteus.
bulbifer, *Blume, Rumphia*, i. 148.—Ind. or.
campanulatus, *Blume, ex Decne. Herb. Timor.* 38.—As. trop.
Chatty, *André, Illustr. Hortic.* xix. (1872) 361.—Ind. or.
commutatus, *Engl. in DC. Monog. Phan.* ii. 319.—Ind. or.
consimilis, *Blume, Rumphia*, i. 149.—Afr. trop.
decurrens, *Kunth, Enum. Pl.* iii. 581.—Ins. Philipp.
difformis, Blume, Rumphia, i. 149 = Anchomanes difformis.
dubius, *Blume, l. c.* 142.—Ind. or.; Malaya.
Fontanesii, *Kunth, Enum. Pl.* iii. 35.—Afr. trop.
Gigas, *Teijsm. & Binn. in Nat. Tijdschr. Neder. Ind.* xxiv. (1862) 329.—Sumatra.
giganteus, *Blume, Rumphia*, i. 144. t. 34.—Ind. or.; Malaya.
gracilis, *Engl. in Engl. Bot. Jahrb.* i. (1881) 183.—Sumatra.
hirsutus, *Teijsm. & Binn. in Nat. Tijdschr. Neder. Ind.* xxiv. (1862) 332.—Java.
imperialis, *Hort. cf. Gard. Chron.* (1883) I. 510.—Ins. Philipp.
Konjac, C. Koch, in Berl. Allg. Gartenz. (1858) 166 = Rivieri.
Lacourii, *Linden & André, in Illustr. Hortic.* xxv. (1878) t. 316.—Cochinch.
leonensis, *Lem. in Fl. des Serres*, Sér. I. ii. (1846) t. 161.—Afr. trop.
longistylus, *Kurz, Rep. Andam.* 50, nomen.—Ins. Andaman.
lyratus, *Kunth, Enum. Pl.* iii. 34.—Ind. or.
margaritifer, Kunth, l. c. = Plesmonium margaritiferum.
mossambicensis, *Klotzsch, ex Garcke, in Peters, Reise Mossamb. Bot.* 509.—Afr. trop.
Mulleri, *Blume, Rumphia*, i. 143.—Java.
nivosus, Lem. Illustr. Hortic. xii. (1865) t. 424 = Dracontium asperum.
palmaeformis, Dur. ex Riv. in Act. Linn. Soc. Bord. xxviii. (1869) 15; Rivière, L'Amorph. 4. 5 = Rivieri.
papillosus, *Hort. ex Rafarin, in Rev. Hortic.* (1871) 476.—Hab.?
Peyri, *cf. Gard. Chron.* (1880) II. 120.—Ins. Seychelles.
planus, *Teijsm & Binn. in Nat. Tijdschr. Neder. Ind.* xxiv. (1862) 331.—Sumatra.
punctulatus, *Blume, Rumphia*, i. 148.—Java.
Rivieri, *Dur. ex Rivière, in Act. Linn. Soc. Bord.* xxviii. (1869) 15.—Cochinch.; Japon.
sativus, *Blume, Rumphia*, i. 145.—Ins. Molucc.
spectabilis, *Engl. in DC. Monog. Phan.* ii. 316.—Java.
sylvaticus, Kunth, Enum. Pl. iii. 34 = Synanthemas silvaticus.
Titanum, *Becc. in Bull. Soc. Tosc. Ortic.* (1879) 46.—Sumatra.
tuberculiger, *Engl. in DC. Monog. Phan.* ii. 317.—Reg. Himal.
variabilis, *Blume, Rumphia*, i. 146. t. 35.—Java; Ins. Philipp.

AMORPHOPHALLUS:—
virosus, *N. E. Br. in Gard. Chron.* (1885) i. 759.—Siam.
Wallisii, Regel, Gartenfl. (1861) 322 = Dracontium asperum.
zebrinus, *Hort.—Cf. Gard. Chron.* (1865) 485.—Hab.?
zeylanicus, *Blume, Rumphia*, i. 148.—Java; Zeylan.

AMORPHOSPERMUM, F. Muell. Fragm. vii. 112 (1870) = **Lucuma**, Juss. (Sapotac.).
antilogum, F. Muell. l. c.—Austral.

AMOSA, Neck. Elem. ii. 459 (1790) = **Inga**, Plum. (Legum.).

AMOUREUXIA, C. Muell. in Walp. Ann. iv. 340 (1857), sphalm. = **Amoreuxia**, Moç. & Sesse (Bixin.).

AMPALIS, Boj. Hort. Maurit. 291 (1837). *URTICACEAE*, Benth. & Hook. f. iii. 365.
madagascariensis, *Boj. l. c.*—Madag.

AMPELANUS, Rafin. in Am. Monthly Mag. (1819) 192 = **Enslenia**, Nutt. (Asclep.).

AMPELOCERA, Klotzsch, in Linnaea, xx. (1847) 541. *URTICACEAE*, Benth. & Hook. f. iii. 353.
cubensis, *Griseb. Cat. Pl. Cub.* 57.—Ind. occ.
Ruizii, *Klotzsch, in Linnaea*, xx. 542.—Peruv.

AMPELOCISSUS, Planch. in Journ. Vigne Am. (1884) 374 = **Vitis**, Linn. (Ampelid.).
abyssinica, Planch. in Journ. Vigne Am. (1885) 24 = V. abyssinica.
acetosa, Planch. l. c. 96.—Austral.
aculeata, Planch. l. c. 95 = V. aculeata.
angolensis, Planch. l. c. 48 = V. angolensis.
arcuata, Planch. l. c. = V. arcuata.
asarifolia, Planch. l. c. 29 = V. asarifolia.
Bakeri, Planch. l. c. 30 = V. Schimperiana.
bombycina, Planch. l. c. 31 = V. bombycina.
cavicaulis, Planch. l. c. 32 = V. cavicaulis.
Chantinii, Planch. l. c. 27 = V. Chantinii.
cinnamochroa, Planch. l. c. 31.—Am. trop.
concinna, Planch. l. c. 48 = V. concinna.
dissecta, Planch. l. c. = V. dissecta.
divaricata, Planch. l. c. (1884) 375 = V. divaricata.
elephantina, Planch. l. c. 379.—Ins. Borbon.
Harmandi, Planch. l. c. 378.—Cochinch.
heracleifolia, Planch. l. c. (1885) 48 = V. heracleifolia.
indica, Planch. l. c. (1884) 375 = V. indica.
ipomeaefolia, Planch. l. c. (1885) 26 = V. ipomeaefolia.
latifolia, Planch. l. c. (1884) 374 = V. latifolia.
Lecardii, Planch. l. c. (1885) 29 = V. Lecardii.
leonensis, Planch. l. c. 30 = V. leonensis.
Leprieurii, Planch. l. c. 44 = V. multistriata.
mossambicensis, Planch. l. c. 49 = V. mossambicensis.
obtusata, Planch. l. c. 48 = V. obtusata.
platanifolia, Planch. l. c. 47 = V. platanifolia.
rugosa, Planch. l. c. (1884) 374 = V. lanata, indica.
salmonea, Planch. l. c. (1885) 31 = V. salmonea.
sarcocephala, Planch. l. c. 45 = V. sarcocephala.
Schimperiana, Planch. l. c. 26 = V. Schimperiana.
sikkimensis, Planch. l. c. (1884) 375 = V. sikkimensis.
tomentosa, Planch. l. c. 375 = V. tomentosa.

AMPELODAPHNE, Meissn. in DC. Prod. xv. i. 81 (1864). *LAURINEAE*, Benth. & Hook. f. iii. 153.
arunciflora, *Meissn. l. c.*—Bras.
dasyantha, *Meissn. l. c.*—Bras.
macrophylla, *Meissn. l. c.*—Bras.

AMPELODESMA, Beauv. Agrost. 78. t. 15. f. 11 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1178.
australe, Brongn. in Duperr. Voy. Bot. 31 = Arundo pilosa.
bicolor, *Kunth, Gram.* i. 79.—Afr. trop.
effusum, *Steud. Syn. Pl. Gram.* 195.—Afr. bor.
festucoides, Steud. l. c. = tenax.
tenax, *Link, Hort. Berol* i. 136.—Reg. Mediterr.

AMPELODESMOS, Link, Hort. Berol. i. 136 (1827) = praec.

AMPELODESMUS, Woods, in Trans. Linn. Soc. xviii. (1838) 34 = praec.

AMPELOPLIS, Rafin. Sylva Tellur. 33 (1838) = **Rhamnus**, Linn.
chinensis, Rafin. l. c. = R. theezans.

AMPELOPSIS, [Rich. in] Michx. Fl. Bor. Am. i. 159 (1803) ex parte = **Vitis**, Linn. (Ampelid.).
aconitifolia, Bunge, Enum. Pl. Chin. Bor. 12 = V. serjaniaefolia.
africana, Steud. Nom. ed. II. i. 219 = V. africana.
bipinnata, Michx. Fl. Bor. Am. i. 160 = V. arborea.
Botria, DC. Prod. i. 633 = V. africana.
brevipedunculata, Maxim. ex Trautv. in Act. Hort. Petrop. viii. (1883) 176.—As. or.
citrulloides, Lebas, in Rev. Hortic. (1875) 179.—Hab.?
cordata, Michx. Fl. Bor. Am. i. 159 = V. indivisa.
cordifolia, Rafin. Med. Fl. ii. 122 = praec.
dissecta, Carr. in Rev. Hortic. (1868) 10 = V. serjaniaefolia.
hederacea, DC. Prod. i. 633 = V. hederacea.
heptaphylla, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 450 = V. hederacea.
heptaphylla, Roem. & Schult. Syst. v. 321 = V. heptaphylla.
heterophylla, Blume, Bijdr. 194 = V. Landuk.
heterophylla, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1846) 196 = V. heterophylla.
himalayana, Royle, Illustr. Himal. Bot. 149 = V. himalayana.
hirsuta, Donn, Hort. Cantab. 166 = hederacea.
humulifolia, Bunge, Enum. Pl. Chin. Bor. 12 = V. heterophylla.
indica, Blume, Bijdr. 193 = V. indica.
japonica, Hort.—Cf. Gard. Chron. (1868) 1118. = V. inconstans.
latifolia, Tausch, in Flora, xxi. (1838) II. 738; Kostel. Allg. Med.-Pharm. Fl. iv. 1198 = V. latifolia.
lucida, Carr. in Rev. Hortic. (1868) 39 = V. serjaniaefolia.
minima, Hort.—Cf. Gard. Chron. (1873) 1469 = V. inconstans.
napiformis, Carr. in Rev. Hortic. (1870–71) 16.—China.
neilgherrensis, Wight, Spicil. Neilgh. i. 32. t. 36; et Ic. t. 965 = V. himalayana.
palmiloba, Carr. in Rev. Hortic. (1867) 451.—China.
pinnata, Roem. & Schult. Syst. v. 322 = V. pinnata.
pubescens, Schlecht. in Linnaea, x. (1836) 251 = V. pubescens.
quinquefolia, Michx. Fl. Bor. Am. i. 159. t. 162 = V. hederacea.
Regeliana, Carr. Rev. Hortic. (1866) 440.—Japon?
sempervirens, Hort.—Cf. Gard. Chron. (1882) II. 491 & 790 = V. striata.
serianaefolia, Bunge, Enum. Pl. Chin. Bor. 12 = V. serjaniaefolia.
ternata, DC. Prod. i. 633 = V. tomentosa.
tricuspidata, Sieb. & Zucc. in Abh. Akad. Muench. iv (1846) 196 = V. inconstans.
triloba, Carr. in Rev. Hortic. (1868) 39 = V. serjaniaefolia.
tripartita, Carr. l. c. = V. serjaniaefolia.
tuberosa, Carr. l. c. (1870–71) 16 = V. serjaniaefolia.
Veitchii, Hort.—Cf. Gard. Chron. (1868) 814 = V. inconstans.

AMPELOSICYOS, (AMPELOSYCIOS), Thou. Hist. Vég. Isles Afr. austr. 68. t. 22 (1807) = **Telfairia**, Hook. (Cucurb.).
scandens, Thou. l. c. = T. pedata.

AMPELYGONUM, Lindl. in Bot. Reg. (1838) Misc. 62 = **Polygonum**, Linn.
chinense, Lindl. l. c. = P. chinense.
tinctorium, Steud. Nom. ed. II. i. 79 = P. tinctorium.

AMPEREA, A. Juss. Euph. Tent. 35. t. 10 (1824) *EUPHORBIACEAE*, Benth. & Hook. f. iii. 265.
conferta, *Benth. Fl. Austral.* vi. 83.—Austral.
cuneiformis, F. Muell. ex Baill. Etud. Gén. Euph. 455 = spartioides.
ericoides, *A. Juss. Tent. Euph.* 112. t. 10.—Austral.
micrantha, *Benth. Fl. Austral.* vi. 83.—Austral.
protensa, *Nees, in Lehm. Pl. Preiss.* ii. 229.—Austral.
rosmarinifolia, Klotzsch, in Lehm. l. c. i. 176 = ericoides.
spartioides, *Brongn. in Duperr. Voy. Bot.* 226. t. 49.—Austral.

AMPEREA:—
subnuda, *Nees, in Lehm. Pl. Preiss.* ii. 229.—Austral.
volubilis, *F. Muell. ex Benth. Fl. Austral.* vi. 82.—Austral.

AMPHANIA, Banks, ex DC. in Mém. Soc. Phys. Genèv. i. (1821) 408 = **Ternstroemia**, Linn.
integrifolia, Soland. ex DC. l. c. 410 = T. elliptica.

AMPHEREPHIS, H. B. & K. Nov. Gen. et Sp. iv. 31. t. 314, 315 (1820) = **Centratherum**, Cass. (Compos.).
aristata, H. B. & K. l. c. 25 = C. punctatum.
indica, Wall. [Cat. n. 2956]; Less. in Linnaea, vi. (1831) 686 = C. reticulatum.
intermedia, Link, Abbild. 5. t. 29 = C. intermedium.
mollis, Wall. Cat. n. 2957 = C. molle.
mutica, H. B. & K. Nov. Gen. et Sp. iv. 26, 315 = C. muticum.
pilosa, Cass. ex Steud. Nom. ed. II. i. 324, in syn. = C. muticum.
psilocarpa, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 4 = Oiospermum involucratum.
pulchella, Cass. ex Steud. Nom. ed. II. i. 79 = C. pulchellum.
senegalensis, Less. in Linnaea, vi. (1831) 687 = Herderia truncata.
Wightiana, Wall. Cat. n. 2958 = Carpesium cernuum.

AMPHIACHYRIS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 313. *COMPOSITAE*, Benth. & Hook. f. ii. 250.
AMPHIPAPPUS, Torr. & Gray, in Proc. Bost. Soc. i. (1844) 210.
dracunculoides, *Nutt. l. c.*—Am. bor.
Fremontii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 633.—Calif.

AMPHIANTHUS, Torr. in Ann. Lyc. N. York, iv. (1837) 82. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 958.
pusillus, *Torr. l. c.*—Am. bor.

AMPHIBECIS, Schrank, in Syll. Ratisb. i. (1824) 86 = **Centratherum**, Cass. (Compos.).
violacea, Schrank, l. c. = C. intermedium.

AMPHIBLEMMA, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) t. 7; xv. (1851) 50. *MELASTOMACEAE*, Benth. & Hook. f. i. 754.
cymosum, *Naud. l. c.* xv. (1851) 51.—Afr. trop.
mollc, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 456; *in Trans. Linn. Soc.* xxviii. (1871) 79.—Afr. trop.
setosum, *Hook. f. ll. cc.*—Afr. trop.

AMPHIBOLIS, C. Agardh, Sp. Alg. ii. i. 474 (1822). *NAIADACEAE*, Benth. & Hook. f. iii. 1015.
GRAUMUELLERA, Reichb. Consp. 43 (1828).
antarctica, Aschers. in Linnaea, xxxv. (1867–68) 164 = Cymodocea antarctica.
bicornis, *C. Agardh, Sp. Alg.* ii. i. 474.—Austral.
zosteraefolia, C. Agardh, l. c. 475 = Cymodocea antarctica.

AMPHIBROMUS, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 420. *GRAMINEAE*, Benth. & Hook. f. iii. 1161.
fluitans, *T. Kirk, in Trans. N. Zeal. Inst.* xvi. (1884) 374. t. 28.—N. Zel.
Neesii, *Steud. Syn. Pl. Gram.* 328.—Austral.

AMPHICALEA, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 411 = **Calea**, Linn. (Compos.).
fruticosa, Gardn. l. c. 412 = C. rotundifolia.
gentianoides, Gardn. l. c. = Geissopappus gentianoides.

AMPHICALYX, Blume, Fl. Jav. Praef. p. vii (1828) = **Diplycosia** Blume (Ericac.).

AMPHICARPAEA, Ell. in Journ. Acad. Philad. i. (1818) 372 (Amphicarpa). *LEGUMINOSAE*, Benth. & Hook. f. i. 529.
CRYPTOLOBUS, Spreng. Syst. iii. 218 (1826).
FALCATA, J. F. Gmel. Syst. 1131 (1791).
LOBOMON, Rafin. New Fl. Am. i. 81 (1836).
SAVIA, Rafin. in Med. Repos. N. York, v. (1808) 352.
TETRODEA, Rafin. New Fl. Am. i. 81 (1836).
ciliata, *Rafin. l. c.* 82.—Am. bor.

AMPHICARPAEA :—
comosa, G. Don, in Loud. Hort. Brit. 314 = monoica.
cuspidata, *Rafin. New Fl. Am.* i. 83.—Am. bor.
deltifolia, Rafin. l. c. = monoica.
Edgeworthii, *Benth. in Miq. Pl. Jungh.* 231.—Reg. Himal.
Ellioti, Rafin. New Fl. Am. i. 83 = monoica.
ferruginea, Benth. in Miq. Pl. Jungh. 231 = Shuteria ferruginea.
heterophylla, Rafin. New Fl. Am. i. 82 = monoica.
monoica, *Ell. in Journ. Acad. Philad.* i. (1818) 372; *& ex Nutt. Gen. Am.* ii. 115.—Am. bor.
Pitcheri, *Torr. & Gray, Fl. N. Am.* i. 292.—Am. bor.
sarmentosa, Ell. ex Nutt. Gen. Am. ii. 114 = monoica.
trisperma, Baker, in Hook. f. Fl. Brit. Ind. ii. 181, in obs. = Edgeworthii.
villosa, Rafin. New Fl. Am. i. 82 = monoica.

AMPHICARPON, Rafin. in Am. Monthly Mag. (Jan. 1818) 175 = seq.

AMPHICARPUM, Kunth, Gram. i. 28 (1829). *GRAMINEAE*, Benth. & Hook. f. iii. 1099.
floridanum, *Chapman, Fl. S. U. St.* 572.—Florida.
Purshii, *Kunth, Gram.* i. 28.—Am. bor.

AMPHICOME, Royle, Illustr. Himal. Pl. 296. t. 72. f. 1 (1835). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1049.
arguta, *Royle, l. c.*—Reg. Himal.
Emodi, *Royle, ex Lindl. Bot. Reg.* (1838) t. 19.—Reg. Himal.

AMPHIDERRIS, Spach, Hist. Vég. Phan. x. 402 (1841) = **Orites**, R. Br. (Proteac.).

AMPHIDETES, Fourn. in Mart. Fl. Bras. vi. IV. 213 (1885). *ASCLEPIADEAE*.
laciniatus, *Fourn. l. c.* 214.—Bras.
quinquedentatus, *Fourn. l. c.* 213.—Bras.

AMPHIDONAX, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836) = **Arundo**, Linn. (Gramin.).
bengalensis, Nees, l. c. = Arundo bengalensis.

AMPHIDONAX, Nees, ex Steud. Syn. Pl. Gram. 197 = **Zenkeria**, Trin. (Gramin.).
bifaria, Nees, ex Steud. l. c.—Japon.
Heynii, Nees, ex Steud. l. c. = Z. elegans.
obtusiflora, Thw. Enum. Pl. Zeyl. 370 = Z. obtusiflora.
tenella, Wight & Arn. ex Steud. Syn. Pl. Gram. 197. —Ind. or.

AMPHIDOXA, DC. Prod. vi. 246 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 300.
filaginea, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 21.—Afr. trop.
gnaphalodes, *DC. l. c.*—Afr. austr.

AMPHIGENES, Janka, in Linnaea, xxx. (1859–60) 619 = **Festuca**, Linn. (Gramin.).
carpathica, Janka, l. c. = F. carpathica.
nutans, Janka, l. c. = F. nutans.

AMPHIGLOSSA, DC. Prod. vi. 258 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 324.
alopecuroides, *Sch. Bip. in Poltichia*, xviii–xix. (1861) 182.—Afr. austr.
callunoides, *DC. Prod.* vi. 259.—Afr. austr.
corrudaefolia, *DC. l. c.* 258.—Afr. austr.
nitidula, *DC. l. c.* 259.—Afr. austr.
perotrichoides, DC. l. c. 258 = Pterothrix perotrichoides.
tomentosa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 276. —Afr. austr.
triflora, DC. Prod. vi. 258 = Pterothrix spinescens.

AMPHIGLOTTIS, Salisb. in Trans. Hort. Soc. i. (1812) 294 = **Epidendrum**, Linn. (Orchid.).
lurida, Salisb. l. c. = E. anceps.
secunda, Salisb. l. c. = E. secundum.

AMPHILOBIUM, Loud. Hort. Brit. 242 (1830), sphalm. = **Amphilophium**, H. B. & K. (Bignon.).

AMPHILOCHIA, Mart. Nov. Gen. et Sp. i. 127. t. 77 (1824) = **Qualea**, Aubl. (Vochysiac.).

AMPHILOCHIA :—
acuminulata, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 343 = Q. cryptantha.
cordata, Mart. Nov. Gen. et Sp. i. 129 = Q. cordata.
dichotoma, Mart. l. c. 128. t. 77 = Q. dichotoma.
Lundii, Warm. in Kjoeb. Vidensk. Meddel. (1867) 27 = Q. Lundii.
marginata, Miq. in Linnaea, xxii. (1849) 796 = Q. marginata.

AMPHILOPHIUM, Kunth, in Journ. Phys. lxxxvii. (1818) 451. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1039.
glanduliferum, Mart. ex DC. Prod. ix. 193 = Vauthieri.
macrophyllum, *H. B. & K. Nov. Gen. et Sp.* iii. 149. —N. Granat.
molle, *Cham. & Schlecht. in Linnaea*, v. (1830) 120. —Mexic.
Mutisii, *H. B. & K. Nov. Gen. et Sp.* iii. 149. t. 219. —N. Granat.
paniculatum, *H. B. & K. l. c.*—Honduras.
pubescens, Spreng. Syst. ii. 836 = Bignonia fallax.
Vauthieri, *DC. Rev. Bignon.* (1838); *Prod.* ix. 193.— Bras.

AMPHINOMIA, DC. Prod. ii. 522 (1825) = **Lotononis**, Eckl. & Zeyh. (Legumin.).
decumbens, DC. l. c. = L. polycephalus.

AMPHIOLANTHUS, Griseb. Cat. Pl. Cub. 186 (1866) = **Micranthemum**, Michx. (Scrophul.).
arenaroides, Griseb. l. c. 187.—Cuba.
bryoides, Griseb. l. c. 186.—Cuba.

AMPHION, Salisb. Gen. Pl. Fragm. 66 (1866) = **Semele**, Kunth (Liliac.).

AMPHIONE, Rafin. Fl. Tellur. iv. 79 (1836) = **Ipomaea**, Linn. (Convolv.).
asarifolia, Rafin. l. c. = I. asarifolia.
chenopodia, Rafin. l. c.—Hab.?
cymosa, Rafin. l. c. = I. cymosa.
lobata, Rafin. l. c.—Hab.?
splendens, Rafin. Sylva Tellur. 130 = Argyreia splendens.
venosa, Rafin. Fl. Tellur. iv. 79 = I. venosa.

AMPHIPAPPUS, Torr. & Gray, in Proc. Bost. Soc. i. (1844) 210; Bost. Journ. Nat. Hist. v. (1847) 107 = **Amphiachyris**, Nutt. (Compos.).
Fremontii, Torr. & Gray, l. c. 108 = Amphiachyris Fremontii.

AMPHIPLEIS, Rafin. Fl. Tellur. iii. 75 (1836) = **Nicotiana**, Linn. (Solanac.).
fetida, Rafin. l. c.—Hab.?
quadrivalvis, Rafin. l. c. = N. quadrivalvis.

AMPHIPOGON, R. Br. Prod. 175 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1145.
AEGOPOGON, Beauv. Agrost. 122. t. 22. f. 4 (1812), partim.
GAMELYTHRUM, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 415.
PENTACRASPEDON, Steud. Syn. Pl. Gram. 151 (1855).
avenaceus, R. Br. Prod. 175 (1810) = strictus.
Brownei, F. Muell. Fragm. viii. 201 = strictus.
caricinus, F. Muell. in Linnaea, xxv. (1852) 445 = strictus.
cygnorum, *Nees, in Lehm. Pl. Preiss.* ii. 100.—Austral.
debilis, *R. Br. Prod.* i. 175.—Austral.
gracilis, Nees, in Lehm. Pl. Preiss. ii. 101 = strictus.
laguroides, F. Muell. Fragm. viii. 201 = cygnorum.
laguroides, *R. Br. Prod.* i. 175.—Austral.
strictus, *R. Br. l. c.*—Austral.
turbinatus, *R. Br. l. c.*—Austral.

AMPHIREPHIS, Nees & Mart. in Nov. Act. Acad. Nat. Cur. xii. (1824) 4 = AMPHIEREPHIS, H. B. & K. = **Centratherum**, Cass. (Compos.).

AMPHIRHAPIS, DC. Prod. v. 343 (1836) = **Inula**, Linn.; **Microglossa**, DC. & **Solidago**, Linn. (Compos.).
albescens, DC. Prod. v. 343 = M. albescens.
chinensis, *Sch. Bip. in Flora*, xxxv. (1852) 58 = Solidago Virga-aurea.

AMPHIRHAPIS :—
cuspidata, DC. Prod. v. 343 = Inula cuspidata.
heterotricha, DC. l. c. = I. cupatorioides.
indica, 'Less.' ex Hook. f. Fl. Brit. Ind. iii. 227, in syn. = Centratherum reticulatum.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 172.— Japon.
leiocarpa, Benth. in Hook. Lond. Journ. Bot. i. (1842) 488 = Solidago Virga-aurea.
mollis, 'Wall.' ex Hook. f. Fl. Brit. Ind. iii. 227, in syn. = Centratherum molle.
pedunculares, DC. Prod. v. 344 = Aster asperulus.
pubescens, DC. l. c. 343 = Solidago Virga-aurea.
rubricaulis, DC. l. c. = I. rubricaulis.
Wightiana, Hook. f. Fl. Brit. Ind. iii. 301 = Carpesium abrotanoides.
zeylanica, DC. Prod. vii. 279 = M. zeylanica.
Zollingeriana, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 126, nomen.—Malaya.

AMPHIRRHOX, Spreng. Syst. Cur. Post. 51 (1827). *VIOLARIEAE*, Benth. & Hook. f. i. 118.
BRADDLEYA, Vell. Fl. Flum. ii. t. 140 (1827).
SPATHULARIA, St. Hil. Pl. Rem. 317. t. 28 (1824).
latifolia, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 376.—Bras.
longifolia, *Spreng. Syst. Cur. Post.* 51.—Bras.
surinamensis, *Eichl. in Mart. Fl. Bras.* xiii. i. 377.— Bras.

AMPHISCOPIA, Nees, in Wall. Pl. As. Rar. iii. 112 (1832) = **Justicia**, Linn. (Acanth.).
aequilabris, Nees, in DC. Prod. xi. 358.—Bras.
Beyrichii, Nees, in Mart. Fl. Bras. ix. 130 = J. nummularia.
ciliata, Moricand, ex Nees, l. c.—Bras.
ciliata, Nees, in DC. Prod. xi. 359.—Reg. Argent.
comosa, Nees, l. c. 360 = J. comosa.
cuneifolia, Nees, in Mart. Fl. Bras. ix. 130 = J. cuneifolia.
inficiens, Nees, in DC. Prod. xi. 357 = J. inficiens.
lancifolia, Nees, l. c.—Peruv.
Martiana, Nees, in Mart. Fl. Bras. ix. 130.—Bras.
Middletoni, Nees, in DC. Prod. xi. 356.—Afr. trop.
montana, Nees, l. c. 359.—Bras.
Moricandiana, Nees, l. c. 357.—Bras.
Pohliana, Nees, l. c. 359.—Bras.
polystachya, Nees, l. c. 356 = J. polystachya.
retusa, Nees, l. c. 357 = J. retusa.
rosea, Nees, l. c. 358 = Eranthemum roseum?
serpens, Nees, l. c. 357 = Hypoëstes serpens?
strobilacea, Nees, l. c. 358.—Bras.
venosa, Nees, l. c. 358 = J. cristata.

AMPHISCOPIUM, St. Lag. in Ann. Soc. Bot. Lyon, vii. 119 (1880) = praec.

AMPHISTELMA, Griseb. Fl. Brit. W. Ind. 417 (1864) = **Vincetoxicum**, Moench (Asclep.).
angulatum, Fourn. in Mart. Fl. Bras. vi. IV. 226.— Bras.
aphyllum, Fourn. l. c.—Bras.
ephedroides, Griseb. Cat. Pl. Cub. 174.—Cuba.
exsertum, Griseb. in Goett. Abh. xxiv. (1879) 229.— Reg. Argent.
ferrugineum, Fourn. in Mart. Fl. Bras. vi. IV. (1885) 224.—Bras.
filiforme, Griseb. Fl. Brit. W. Ind. 418.—Ind. occ.
graminifolium, Griseb. Cat. Pl. Cub. 174.—Cuba.
leptocladon, Griseb. Fl. Brit. W. Ind. 418 = V. leptocladum.
linearifolium, Griseb. Cat. Pl. Cub. 175.—Cuba.
melanthum, Fourn. in Mart. Fl. Bras. vi. IV. 226 = V. melanthum.
parviflorum, Fourn. l. c. 224.—Bras.
Riedelii, Fourn. l. c. 225.—Bras.
salinarum, Wright, ex Griseb. Cat. Pl. Cub. 175.— Cuba.
Selloanum, Fourn. in Mart. Fl. Bras. vi. IV. 224.— Bras.
streptolobium, Fourn. l. c. 225.—Bras.
tomentosum, Fourn. l. c. 226.—Bras.

AMPHITECNA, Miers, in Trans. Linn. Soc. xxvi. (1868) 163. *BIGNONIACEAE*.
macrophylla, *Miers, ex Baill. in Rev. Hortic.* 1882 465.—Am. centr.
nigripes, *Baill. l. c.*—Am. trop.?

AMPHITHALEA, Eckl. & Zeyh. Enum. 167 (1836).
LEGUMINOSAE, Benth. & Hook. f. i. 472.
CRYPHIANTHA, Eckl. & Zeyh. Enum. 171 (1836).
EPISTEMUM, Walp. in Linnaea, xiii. (1839) 473.
INGENHOUSSIA, E. Mey. Comm. Pl. Afr. Austr. 20 (1835).
ciliaris, Eckl. & Zeyh. Enum. 169 = Coelidium ciliare.
cuneifolia, *Eckl. & Zeyh. l. c.* 167.—Afr. austr.
densa, *Eckl. & Zeyh. l. c.*—Afr. austr.
densiflora, Eckl. & Zeyh. l. c. 168 = ericaefolia.
ericaefolia, *Eckl. & Zeyh. l. c.* 169.—Afr. austr.
hilaris, Eckl. & Zeyh. l. c. = ericaefolia.
humilis, Eckl. & Zeyh. l. c. 168 = intermedia.
incurvifolia, Eckl. & Zeyh. l. c. = ericaefolia.
intermedia, *Eckl. & Zeyh. l. c.*—Afr. austr.
Kraussiana, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 65 = virgata.
micrantha, *Walp. in Linnaea*, xiii. (1839) 470.—Afr. austr.
multiflora, Eckl. & Zeyh. Enum. 168 = ericaefolia.
perplexa, Eckl. & Zeyh. l. c. 170 = Coelidium roseum.
phylicoides, *Eckl. & Zeyh. l. c.*—Afr. austr.
spinosa, Steud. Nom. ed. II. i. 80 = Coelidium spinosum.
tortilis, Steud. l. c. = Coelidium Vogelii.
verticillata, Steud. l. c. = Priestleya elliptica.
violacea, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 451.—Afr. austr.
virgata, *Eckl. & Zeyh. Enum.* 169.—Afr. austr.
Vogelii, Walp. in Linnaea, xiii. (1839) 472 = ericaefolia.
Williamsonii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 23.—Afr. austr.

AMPHIZOMA, Miers, in Trans. Linn. Soc. xxviii. (1872) 422 = **Salacia**, Linn. (Hippocrat.).
laxiflora, Miers, l. c. = S. laxiflora.
Martiana, Miers, l. c. 423 = S. Martiana.
tenuicula, Miers, l. c.—Bras.

AMPHOCHAETA, Anderss. in Vet. Akad. Hand. Stockh. (1853) 136 = **Pennisetum**, Pers. (Gramin.).
exaltata, Anderss. l. c. = P. exaltatum.

AMPHODUS, Lindl. Bot. Reg. t. 1101 (1827) = **Kennedya**, Vent. (Legum.).
ovatus, Lindl. l. c. = K. rubicunda.

AMPHORCHIS, Thou. Orch. Iles Afr. Tabl. Synopt. (1822) = **Cynorchis**, Thou. (Orchid.).
calcarata, Thou. l. c. t. 4 = C. squamosa.
colorata, Steud. Nom. ed. II. i. 80, sphalm. = praec.
inermis, Thou. Orch. Isles Afr. t. 5 = Arnottia inermis.
laxiflora, Blume, Mus. Bot. Lugd. Bat. ii. 190.—Malaya.
lilacina, Ridl. in Journ. Linn. Soc. xxi. (1885) 518.—Madag.
nilarmis, Steud. Nom. ed. II. i. 80, sphalm. = inermis.
occidentalis, Lindl. in Journ. Linn. Soc. vi. (1862) 139.—Afr. trop.

AMPHORICARPUS, Visiani, Fl. Dalm. ii. 27. t. 10. f. 2 (1844). *COMPOSITAE*, Benth. & Hook. f. ii. 465.
Neumayeri, *Visiani, Fl. Dalm.* ii. 28. t. 10. f. 2.—Dalmat.

AMPHORICARPUS, Spruce, ex Miers, in Trans. Linn. Soc. xxx. (1874) 288 = **Couratari**, Aubl. (Myrtac.).
uaupensis, Spruce, ex Miers, l. c. = C. uaupensis.

AMPHYMENIUM, H. B. & K. Nov. Gen. et Sp. vi. 380 (1823) = **Pterocarpus**, Linn. (Legum.).
cordatum, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1205, nomen.—Guiana.
floridum, Benth. in Hook. Kew Journ. ii. (1850) 234 = P. Rohrii.
latifolium, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1205, nomen.—Guiana.
laxiflorum, Benth. in Hook. Lond. Journ. Bot. ii. (1840) 65.—Bras.
laxum, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1205, nomen.—Guiana.
longifolium, Benth. in Mart. Fl. Bras. xv. 1. 269 = P. ancylocalyx.

AMPHYMENIUM :—
mediterraneum, Benth. in Ann. Wien. Mus. ii. (1838) 95 = Martia parvifolia.
nitidum, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1205, nomen.—Guiana.
orbiculatum, Benth. in Ann. Wien. Mus. ii. (1838) 95 = P. orbiculatus.
pubescens, H. B. & K. Nov. Gen. et Sp. vi. 380 = P. Amphymenium.
reticulatum, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1205, nomen.—Guiana.
Rohrii, H. B. & K. Nov. Gen. et Sp. vi. 380 = P. Rohrii.
tenuifolium, Klotzsch, in Schomb. Faun. et Fl. Guyan. 1205, nomen.—Guiana.
villosum, Benth. in Ann. Wien. Mus. ii. (1838) 95 = P. villosus.
violaceum, Walp. Rep. i. 793 = P. violaceus.

AMPOMELE, Rafin. Sylva Tellur. 160 (1838) = **Rubus**, Tourn. (Rosac.).
triphylla, Rafin. l. c. = R. trifidus.

AMSINCKIA, Lehm. Del. Sem. Hort. Hamb. (1831) 7. *BORAGINEAE*, Benth. & Hook. f. ii. 851.
BENTHAMIA, Lindl. Nat. Syst. ed. I. 241 (1830).
angustifolia, *Lehm. l. c.*—Chili.
basistaminea, *Cesati, in Atti Acc. Sc. Nap.* v. (1873) VII. 14.—Reg. Andin.
Douglasiana, A. DC. in DC. Prod. x. 118 = spectabilis.
echinata, *A. Gray, in Proc. Am. Acad.* x. (1874) 54.—Calif.
humifusa, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 371.—Reg. Argent.
intermedia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1836) 26.—Calif.
lycopsioides, *Lehm. Del. Sem. Hort. Hamb* (1831) 7.—Calif.
mexicana, *Mart. & Galeot. in Bull. Acad. Brux.* xi. (1844) reimpr. 23.—Mexic.
parviflora, Bernh. Del. Sem. Hort. Erf. (1833) = angustifolia.
spectabilis, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1836) 26.—Calif.
tessellata, *A. Gray, in Proc. Am. Acad.* x. (1874) 54.—Calif.
vernicosa, *Hook. & Arn. Bot. Beech. Voy.* 370.—Calif.

AMSONIA, Walt. Fl. Carol. 98 (1788). *APOCYNACEAE*, Benth. & Hook. f. ii. 702.
angustifolia, *Michx. Fl. Bor. Am.* i. 121.—Am. bor.
brevifolia, *A. Gray, in Proc. Am. Acad.* xii. (1877) 64.—Utah; Arizona.
ciliata, Walt. Fl. Carol. 98 = angustifolia.
elliptica, *Roem. & Schult. Syst.* iv. 432.—Japon.
latifolia, Michx. Fl. Bor. Am. i. 121 = A. Tabernaemontana.
longiflora, *Torr. Bot. Mex. Bound.* 159.—N. Mexic.
orientalis, Decne. in Jacquem. Voy. Bot. 105 = Rhazya orientalis.
Palmeri, *A. Gray, in Proc. Am. Acad.* xii. (1877) 64.—Arizona.
salicifolia, Pursh, Fl. Am. Sept. i. 184 = A. Tabernaemontana.
Tabernaemontana, *Walt. Fl. Carol.* 98.—Am. bor.
tomentosa, *Torr. & Frém. in Frém. Rep.* 1843-44 (1845) 316.—Am. bor.
tristis, Sm. in Rees, Cycl. xxxv. ad calc. art. Tabernaemontanae, n. 2 = A. Tabernaemontana.

AMSORA, Bartl. Ord. 536 (1830), sphalm. = **Amoora**, Roxb. (Meliac.).

AMURA, Schult. f. Syst. vii. p. 11, 75 (1829) = **Amoora**, Roxb. (Meliac.).

AMYDRIUM, Schott, in Miq. Ann. Mus. Lugd. Bat. i. 127 (1863). *AROIDEAE*, Benth. & Hook. f. iii. 993.
humile, *Schott, l. c.*—Borneo.

AMYGDALOPHORA, Neck. Elem. ii. 70 (1790) = **Prunus**, Tourn. (Rosac.).

AMYGDALOPSIS, Carr. in Rev. Hortic. (1862) 91 = **Prunus**, Tourn. (Rosac.).
eburnea, M. Roem. Syn. Rosifl. 16 = P. eburnea.
furcata, M. Roem. l. c. = P. eburnea.
horrida, M. Roem. l. c. 15 (= *Amygdalus horrida*).—Persia.

AMYGDALOPSIS :—
Lindleyi, Carr. in Rev. Hortic. (1862) 91; (1868) 197 = P. triloba.
lycioides, M. Roem. Syn. Rosifl. 15 (= *Amygdalus lycioides*).—As. Min.; Syria.
Scorpius, M. Roem. l. c. = P. eburnea.
virgata, Carr. in Rev. Hortic. (1872) 200 = P. virgata.

AMYGDALUS, [Tourn.] Linn. Syst. ed. I. (1735) = **Prunus**, Tourn. (Rosac.).
agrestis, Boiss. Diagn. Ser. I. x. 1.—Syria.
amara, Hayne, Arzn. iv. in nota ad t. 39. f. i. = P. Amygdalus.
amygdalina, Oken, ex M. Roem. Syn. Rosifl. 8 = P. Amygdalus.
arabica, Olivier, Voy. t. 47.—Arabia.
argentea, Lam. Encyc. i. 103 = orientalis.
Armeniaca, Dum. Fl. Belg. 91 = P. Armeniaca.
Armeniacaria, Oken, ex M. Roem. Syn. Rosifl. 10 = P. Amygdalus.
Balansae, Boiss. Diagn. Ser. II. vi. 71.—Phrygia.
Besseriana, Schott, ex Schlecht. in Abh. Naturf. Ges. Halle, ii. (1854) 1 = P. nana.
Boissierii, Carr. in Rev. Hortic. (1875) 352.—As. Min.
Bovei, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 124.—Syria.
brahuica, Boiss. Fl. Orient. ii. 645.—Beluchist.
camelliaeflora, Morr. in Belg. Hortic. viii. (1858) 97. t. 26 = P. Persica.
campaniflora, Carr. in Rev. Hortic. (1872) 129.—Hab.?
campestris, Bess. Enum. Pl. Volh. 46 = P. nana.
caryophyllacea, Hort. ex C. Koch, Dendrol. i. 85 = P. Persica.
cerasina, Oken, ex M. Roem. Syn. Rosifl. 9 = P. Amygdalus.
chinensis, Hort. ex C. Koch, Dendrol. i. 85 = P. Persica.
chinensis, Leroy, ex C. Koch, l. c. 79 = P. nana.
cochinchinensis, Lour. Fl. Cochinch. 316.—Cochinch.
collinus, Wall. Cat. n. 723 = P. Persica.
communis, Bunge, Enum. Pl. Chin. Bor. 21 = P. Persica.
communis, Linn. Sp. Pl. ed. I. 473 = P. Amygdalus.
cordifolia, Roxb. Fl. Ind. ii. 500.—China.
dasylepis, Miq. in Journ. Bot. Néerl. i. (1861) 122 = P. Persica.
dianthiflora, Hort. ex C. Koch, Dendrol. i. 85 = P. Persica.
discolor, M. Roem. Syn. Rosifl. 12 = orientalis.
dulcis, Mill. Gard. Dict. ed. VIII. n. 2 = P. Amygdalus.
eburnea, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 123.—Persia.
elaeagrifolia, Spach, l. c. 120.—Persia.
elata, Salisb. Prod. 356 = P. Amygdalus.
Fischeriana, Spach, in Ann. Sc. Nat. Sér II. xix. (1843) 125.—Hab.?
fragilis, Hell. Fl. Wirceb. i. 500 = P. Amygdalus.
frutescens, Steud. Nom. ed. II. i. 81 = (*A. fruticosa*).
fruticosa, Wender. in Bot. Zeit. (1865) 338. t. 12.—Hab.?
furcatus, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 124 = P. eburnea.
Gaertneriana, Schlecht. in Abh. Ges. Halle, ii. (1854) 19 = P. nana.
georgica, Desf. Hist. Arbr. ii. 221 = P. nana.
glandulosa, Hort. ex C. Koch, Dendrol. i. 118 = P microphylla.
glandulosa, Hook. Ic. Pl. t. 288 = P. glandulosa.
glandulosa, Leroy, ex C. Koch, Dendrol. i. 116 = P. japonica.
grandiflora, Hort. ex C. Koch, l. c. 81 = P. Amygdalus.
grandiflora, Hort. Gall. ex C. Koch, l. c. i. 80 = P. Persica.
grata, Salisb. Prod. 356 = P. Persica.
heterophylla, Hort. ex C. Koch, Dendrol. i. 81 = P. Amygdalus.
Heuckeana, Schlecht. in Abh. Ges. Halle, (1854) 1. 22.—Sibir.
horrida, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 121.—Persia.
humilis, Edgew. in Trans. Linn. Soc. xx. (1846) 44 = P. Jacquemontii.
hybrida, Dierb. ex M. Roem. Syn. Rosifl. 11 = P. Amygdalus.

AMYGDALUS :—
incana, Pall. Fl. Ross. i. 13. t. 7 = Prunus prostrata.
ispahanensis, Thouin, in Ann. Mus. Par. Sér. I. viii. (1826) 425 = P. Persica.
Kotschyi, Hohen. ex Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 117 = *eleagnifolia*.
laevis, D. Dietr. Syn. Pl. iii. 42 = P. Persica.
Ledebouriana, Schleeht. in Abh. Ges. Halle, (1854) 1 = P. nana.
leiocarpa, Boiss. Diagn. Ser. I. vi. 52.—Persia.
lycioides, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 120.—As. Min.; Syria.
microphylla, H. B. & K. Nov. Gen. et Sp. vi. 245. t. 564 = P. microphylla.
minor, Duh. Arbr. Fruit. i. t. 7.—Hab.?
nana, Linn. Sp. Pl. 473.—Europ.; Reg. Cauc.; Sibir.
Nuci-persica, Reichb. Fl. Germ. Excurs. 647 = P. Persica.
orientalis, Mill. Gard. Dict. ed. VIII. n. 4.—As. Min.; Syria.
Pallasiana, Schlecht. in Abh. Ges. Halle, (1854) 1 = P. nana.
Pallasii, Turcz. ex Steud. Nom. ed. II. i. 81 = P. pedunculata.
pedunculata, Bunge, Enum. Pl. Chin. Bor. 22 = P. triloba.
pedunculata, Pall. in Nov. Act. Petrop. vii. (1789) 355. t. 8, 9.—Sibir.
Persica, Linn. Sp. Pl. 472 = P. Persica.
persico-amygdala, Reichb. Fl. Germ. Excurs. 647 = P. Amygdalus.
persicoides, C. Koch, Dendrol. i. 84 = P. Persica.
pilosa, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 189.—China.
prostrata, Sweet, Hort. Brit. ed. II. 173 = P. prostrata.
pumila, Linn. Mant. i. 74 = P. japonica.
Reuteri, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 79 = *horrida*.
rosaeflora, Hort. ex C. Koch, Dendrol. i. 85 = P. Persica.
salicifolia, Boiss. & Bal. Diagn. Ser. II. vi. 671 = Webbii.
sativa, Mill. Gard. Dict. ed. VIII. n. 3 = P. Amygdalus.
scoparia, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 109.—Persia.
scorpius, Spach, l. c. 122 = P. eburnea.
serratifolia, Hort. ex C. Koch, Dendrol. i. 79 = P. nana.
sibirica, Lodd. ex Loud. Hort. Brit. 199.—Sibir.
sinensis, Hort. ex Steud. Nom. ed. II. i. 81 = *cochinchinensis*.
spartioides, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 108.—Syria; Persia.
spathulata, Boiss. Diagn. Ser. I. vi. 53 = P. eburnea.
spinosissima, Bunge, Rel. Lehm. 106.—Turkest.
Stocksiana, Boiss. Diagn. Ser. II. ii. 45.—Beluchist.
tomentosa, Hort. ex C. Koch, Dendrol. i. 81 = P. Amygdalus.
tomentosula, Steud. Nom. Bot. ed. II. i. 81 = P. Amygdalus.
Webbii, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 117.—Phrygia.

AMYRIS, P. Br. Hist. Jam. 208 (1756). *BURSERACEAE*, Benth. & Hook. f. i. 327.
Elemi, Adans. Fam. ii. 342 (1763).
Elemifera, Linn. Hort. Cliff. 486 (1737).
Rhinophora, Neck. Elem. ii. 229 (1790).
acuminata, Poir. Encyc. Suppl. ii. 136 = Icica acuminata.
acuminata, Roxb. Hort. Beng. 28; Fl. Ind. ii. 246 = Commiphora caudata.
Agallocha, Roxb. Hort. Beng. 28 (1814) = Commiphora Agallocha.
altissima, Willd. Sp. Pl. ii. 336 = Protium altissimum.
ambrosiaca, Linn. f. Suppl. 216 = Protium Icicariba.
ambrosiaca, Moç. & Sesse, in DC. Prod. ii. 77, 78 = Icica serrata.
ambrosiaca, Spreng. Syst. ii. 218, pro parte = Protium guianense.
ambrosiaca, Willd. Sp. ii. 335 = Protium heptaphyllum.
ambrosiaca, Vell. Fl. Flum. iv. t. 3 = Protium heptaphyllum?
anisata, Roxb. ex Steud. Nom. ed. II. i. 81 = Clausena heptaphylla.
anisata, Willd. Sp. Pl. ii. 337 = Clausena anisatum.
ariensis, Spreng. Syst. iv. Cur. Post. 149 = Elaphrium ariense.
armata, Spreng. Syst. ii. 219 = Zanthoxylon Thunbergii.
axilliflora, *Griseb. Cat. Pl. Cub.* 66.—Cuba.

AMYRIS :—
balsamifera, *Linn. Syst.* ed. X. 1000.—Jamaica.
bicolor, Willd. ex Engl. in DC. Monog. Phan. iv. 54 = Bursera bicolor.
bipinnata, Moç. & Sesse, ex DC. Prod. ii. 82 = Bursera bipinnata.
brachybotrys, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 614.—Bras.
brasiliensis, Spreng. Syst. ii. 217 = Protium brasiliense.
brasiliensis, Willd. ex Engl. in DC. Monog. Phan. iv. 64 = Protium heptaphyllum.
capensis, Spreng. Syst. ii. 218 (= *Fagarastrum capense*).
Carana, Humb. Relat. ii. 421 = Protium Carana.
caranifera, Willd. ex Engl. in DC. Monog. Phan. iv. 49 = Bursera graveolens.
Commiphora, Roxb. Fl. Ind. ii. (1824) 244 = Commiphora Agallocha.
Cataf, 'Forsk.' = *A. Kataf.*
copallifera, *Oliva, in La Naturaleza*, (1870) 40.—Mexico.
copallifera, Spreng. Syst. ii. 219 = Elaphrium copalliferum.
cymosa, Reichb. ex Steud. Nom. ed. II. i. 81 = sylvatica.
decandra, *Brotero, ex Pritz. in Bot. Zeit.* i. (1843) 561.—Bras.
decandra, Willd. Sp. Pl. ii. 336 = Icica decandra.
dentata, Blume, Bijdr. 1160 = Clausena javanica.
dentata, Willd. Sp. Pl. ii. 337 = Clausena Willdenowii.
diatrypa, *Spreng. Neue Entdeck.* iii. 48; *Syst.* ii. 217.—Ins. S. Doming.
dioica, Hort. ex Steud. Nom. ed. II. i. 81 = Sehinus dependens.
Elaphrium, Spreng. Syst. ii. 219 = Bursera glabra.
elegans, Willd. ex Engl. in DC. Monog. Phan. iv. 57 = Bursera excelsa.
elemifera, Koen. ex Blume, Mus. Bot. Lugd. Bat. i. 218 = Canarium zeylanicum.
elemifera, Linn. Syst. ed. X. 1000 = Plumieri.
enneandra, Willd. Sp. Pl. ii. 335 = Protium decandrum.
fagaroides, Spreng. Syst. iv. Cur. Post. 148 = Bursera fagaroides.
floridana, *Ware, ex Nutt. in Am. Journ. Sc.* v. (1822) 294.—Florida.
Funckiana, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 475.—Venezuela.
fruticosa, *Roxb. Hort. Beng.* 28.—Ind. or.
gilcadensis, Roxb. Hort. Beng. 28; Fl. Ind. ii. 246 = C. Berryi.
giliadensis, Linn. Diss. Opob.; Amoen. Acad. vii. 68 = Commiphora Opobalsamum.
graveolens, Buch.-Ham. ex Steud. Nom. ed. II. i. 81 = Clausena excavata.
graveolens, Spreng. Syst. iv. Cur. Post. 148 = Bursera graveolens.
guianensis, *Aubl. Pl. Gui.* i. 336.—Guiana.
guianensis, Willd. Sp. Pl. ii. 325 = Protium guianense.
heptaphylla, Roxb. Hort. Beng. 28; Fl. Ind. ii. 248 = Clausena heptaphylla.
heterophylla, Willd. Sp. Pl. ii. 335 = Protium Aracouchini.
hexandra, Ham. Prod. Fl. Ind. Occ. 34 = Dacryodes hexandra.
Hypelate, Robinson, ex Lunan, Hort. Jamaic. i. 149 = Hypelate trifoliata.
inaequalis, Spreng. Syst. ii. 218 = Clausena inaequalis.
jorullensis, Spreng. Syst. Cur. Post. 148 = Bursera jorullensis.
Kafal, Forsk. Fl. Aegypt. Arab. 80 = an seq.?
Kataf, Forsk. l. c. = Commiphora Kataf.
lanuginosa, Spreng. Syst. iv. Cur. Post. 149 = Bursera lanuginosa.
Linaloë, La Llave, in La Naturaleza, vii. (1885) Apénd. 73 = seq.
Linanoë, La Llave, ex Schlecht. in Linnaea, xvii. (1843) 252 = Bursera Aloëxylon.
lineata, *C. Wright, ex Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 177.—Ind. occ.
Lunani, Spreng. Syst. ii. 217 = maritima.
Lypelata, Steud. Nom. ed. II. i. 81, sphalm. = Hypelate trifoliata.
maritima, *Jacq. Enum. Pl. Carib.* 19.—Am. trop.
maritima, Llanos, in Mem. Acad. Madr. Ser. III. ii. iii. (1856) 503.—Ins. Philipp.?
maritima, Moç. & Sesse, ex DC. Prod. ii. 82 = A. Tecomaca.
multijuga, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 476.—Bras.

AMYRIS :—
nana, Roxb. Hort. Beng. 28 = Clausena Willdenowii.
Novae-Hollandiae, Hort. ex Steud. Nom. ed. II. i. 81 = Botryceras laurinum.
oleosa, Lam. Encyc. i. 362 = Canarium microcarpum.
Opobalsamum, Linn. Diss. Opob.; Amoen. Acad. vii. 68; Forsk. Fl. Aegypt. 79 = Commiphora Opobalsamum.
papyrifera, Caill. ex Del. Voy. Meroé, 99 = Boswellia papyrifera.
penicillata, Spreng. Syst. ii. 219 = Bursera penicillata.
pentaphylla, Roxb. Hort. Beng. 28; Fl. Ind. ii. 247 = Clausena pentaphylla.
pernambucensis, *Arruda, in Koster, Trav. Braz.* 500.—Bras.
Philippaea, Robinson, ex Lunan, Hort. Jam. i. 150 = Hypelate trifoliata.
pinnata, *H. B. & K. Nov. Gen. et Sp.* vii. 37.—Am. trop.
Plumieri, *DC. Prod.* ii. 81.—Ind. occ.
polygama, Cav. Ic. iii. 20 = Schinus dependens.
Protium, Linn. Mant. i. 65 = Protium javanicum.
pubescens, Willd. ex Schlecht. in Linnaea, xvii. (1842) 527 = Elaphrium pubescens.
punctata, Roxb. Hort. Beng. 28; Fl. Ind. ii. 251 = Clausena excavata.
Robinsonii, *DC. Prod.* ii. 82.—Jamaica.
rugosa, Willd. ex Engl. in DC. Monog. Phan. iv. 58 = Bursera jorullensis.
serratifolia, Rottl. ex A. W. Benn. in Hook. f. Fl. Brit. Ind. i. 529 = Balsamodendron Roxburghii.
simplicifolia, Roxb. Fl. Ind. ii. 244 = Atalantia Roxburghiana.
suffruticosa, Roxb. l. c. 250 = Clausena suffruticosa.
sumatrana, Roxb. Hort. Beng. 28; Fl. Ind. ii. 250 = Clausena excavata.
sylvatica, *Jacq. Am. Select.* 107.—Ind. or.
Tecomaca, *Moç. & Sesse, Fl. Mex. ined.* t. 195, *ex DC. Prod.* ii. 82.—Mexico.
terebinthifolia, *Tenore, Ind. Sem. Hort. Neap.* (1837) 1.—Bras.
thyrsiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 475.—Mexico.
tomentosa, Spreng. Syst. ii. 219 = Bursera tomentosa.
toxifera, *Linn. Syst.* ed. X. 1000.—Ind. occ.
ventricosa, Llave, ex Schlecht. in Linnaea, xvii. (1843) 246 = Bursera fagaroides.
zeylanica, Retz. Obs. fasc. iv. 25 = Canarium zeylanicum.

AMYRSIA, Rafin. Sylva Tellur. 106 (1838) = **Eugenia**, Linn. (Myrtac.).
compressa, Rafin. l. c. = E. compressa.
discolor, Rafin. l. e. = E. discolor.
foliosa, Rafin. l. e. = E. foliosa.
microphyla, Rafin. l. c. (= *Myrtus microphylla*).

ANABAENA, A. Juss. Euph. Tent. 46. t. 15 (1824) = **Plukenetia**, Linn. (Euphorb.).
tamnoides, A. Juss. l. c. t. 15 = P. tamnoides.

ANABASIS, Linn. Pl. Camts. (1750) (Amoen. Acad. ii. 347). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 72.
Borith, Adans. Fam. ii. 262 (1763).
Brachylepis, C. A. Mey. Fl. Alt. i. 370 (1829).
Fredolia, Coss. & Dur. ex Moq. & Coss. in Bull. Soc. Bot. Fr. ix. (1862) 301.
Microlepis, Eichw. Casp. Cauc. 2 (1831).
affinis, Bunge, Rel. Lehm. 480 = Arthrophytum subulifolium.
affinis, C. A. Mey. in Schrenk, Enum. Pl. Nov. Song. (1841) 10 = brevifolia.
alopecuroides, Moq. in DC. Prod. xiii. ii. 210 = Halogeton alopecuroides.
Ammodendron, C. A. Mey. in Ledeb. Fl. Alt. i. 375 = Haloxylon Ammodendron.
annua, *Bunge, Anab. Rev.* 46.—Persia.
aphylla, *Linn. Sp. Pl.* 223.—Orient; Sibir.
arbuscula, *Thunb. ex Steud. Nom.* ed. II. i. 81.—Hab.?
arctioides, *Moq. & Coss. in Bull. Soc. Bot. Fr.* ix. (1862) 301. t. 2.—Afr. bor.
articulata, Choulette, ex Pomel, Nouv. Mat. Fl. Atl. 334 = prostrata.
articulata, *Moq. in DC. Prod.* xiii. ii. 212.—Afr. bor.; Syria.
brachiata, *Fisch. & Mey. ex Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 433.—Orient; Turkest.
brachiata, Kar. & Kir. l. c. = truncata.

ANABASIS:—
brachylepis, *D. Dietr. Syn. Pl.* ii. 995.—Reg. Casp.
brevifolia, *C. A. Mey. in Ledeb. Fl. Alt.* i. 377.—Sibir. alt.
caespitosa, Steph. ex Moq. in DC. Prod. xiii. II. 216 = Nanophyton caspicum.
cinerea, Moq. l. c. 214 = Seidlitzia florida.
clavata, S. G. Gmel. It. ii. 41 = Salsola clavifolia.
conjugata, Hoffm. Hort. Mosq. (1808) No. 184 = Petrosimonia brachiata.
crassa, Moq. Chenop. Enum. 166 = articulata.
cretacea, C. A. Mey. Enum. ex Bunge, Anab. Rev. 39 = brachiata.
cretacea, *Pall. Reise*, i. 493.—Sibir.
cretica, Vitm. Summ. Pl. ii. 104 = cretacea.
Echinus, Bieb. Beschr. Laend. Casp. 152 = Noaea spinosissima.
Ehrenbergii, *Schweinf. ex Boiss. Fl. Orient.* iv. 970, in nota.—Nubia; Arab.
florida, Bieb. Fl. Taur. Cauc. i. 190 = Seidlitzia florida.
foliata, Pall. ex Bunge, Anab. Rev. 47 = Salsola foliosa.
foliosa, Pall. Reise, i. 380 = Salsola clavifolia.
foliosa, Linn. Sp. Pl. 223 = Salsola foliosa.
glomerata, Bieb. in Mém. Soc. Nat. Mosc. i. (1808) 110 = Halogeton glomeratus.
Haussknechti, *Bunge, ex Boiss. Fl. Orient.* iv. 969.—Persia.
heterophylla, Walp. Ann. i. 571 (sphalm. = seq.).
heteroptera, Jaub. & Spach, Illustr. ii. 45. t. 134 = Girgensohnia oppositiflora.
intermedia, Moq. in DC. Prod. xiii. II. 214 = phyllophora?
juncea, Bieb. ex Moq. l. c. 203 = Ofaiston monandrum.
Kareliniana, Fenzl, in Ledeb. Fl. Ross. iii. 825 = truncata.
lutea, Moq. in DC. Prod. xiii. II. 215 = setifera.
macroptera, *Moq. l. c.* 213.—Affghanist.
monandra, Schrad. Neues Journ. iii. (1809) 89 = Ofaiston pauciflorum.
multiflora, Moq. in DC. Prod. xiii. II. 212 = Haloxylon multiflorum.
oppositiflora, Bieb. in Mém. Soc. Nat. Mosc. i. (1808) 148 = Girgensohnia oppositiflora.
oppositifolia, Bieb. Beschr. Laend. Casp. App. 154 = Petrosimonia crassifolia.
oppositifolia, Schrad. Neues Journ. iii. (1809) 89 = Halogeton oppositifolius.
phyllophora, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 731.—Syria; Soongaria.
prostrata, *Pomel, Nouv. Mat. Fl. Alt.* 334.—Afr. bor.
ramosa, *Sievers, in Pall. N. Nord. Beitr.* vii. 304.—Sibiria.
setifera, *Moq. Chenop. Enum.* 164.—Aegypt.; Oriens.
Sieversii, Willd. in Roem. & Schult. Syst. vi. 223 = Nanophyton caspicum.
spinosissima, Linn. f. Suppl. 173 = Noaea spinosissima.
subulifolia, C. A. Mey. in Bull. Acad. Pétersb. Sér. IV. i. (1843) 360 = phyllophora.
tamariscifolia, Linn. Syst. ed. X. 949 = Salsola tamariscifolia.
tamariscifolia, Webb, Iter Hisp. 17 = Salsola Webbii.
tatarica, Pall. Illustr. 13. t. 8 = aphylla.
Tournefortii, Jaub. & Spach, Illustr. Pl. Or. ii. 43. t. 132 = Noaea Tournefortii.
triandra, Bieb. Beschr. Laend. Casp. App. 154 = Petrosimonia Volvox.
truncata, *Bunge, Anab. Rev.* 38.—Reg. Casp.

ANABATA, Willd. ex Roem. & Schult. Syst. iv. 60, 707 (1819). Gen. dub. *LOGAN.* Benth. & Hook. f. ii. 788.
SULZERIA, Roem. & Schult. l. c.
odorata, *Humb. ex Roem. & Schult. l. c.*—Am. austr.

ANACAMPSEROS, Linn. Syst. ed. I. (1735); Gen. ed. I. 152 (1737). *PORTULACEAE*, Benth. & Hook. f. i. 157.
RUELINGIA, Ehrh. Beitr. iii. 132 (1788).
TALINUM, Adans. Fam. ii. 245 (1763).
TELEPHIASTRUM, Medic. Phil. Bot. i. 100 (1789).
angustifolia, *Sweet, Hort. Brit.* ed. I. 170.—Afr. austr.
arachnoides, *Sims, Bot. Mag.* t. 1368.—Afr. austr.
filamentosa, *Sims, l. c.* 1367.—Afr. austr.
intermedia, Haw. ex G. Don, Gen. Syst. iii. 75 = A. Telephiastrum.

ANACAMPSEROS:—
lanceolata, *Sweet, Hort. Brit.* ed. I. 170.—Afr. austr.
lanigera, *Burch. Trav.* ii. 333.—Afr. austr.
papyracea, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 383.—Afr. austr.
polyphylla, *Sweet, Hort. Brit.* ed. I. 170.—Afr. austr.
quinaria, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 383.—Afr. austr.
rotundifolia, *Sweet, Hort. Brit.* ed. I. 170.—Afr. austr.
rubens, Sweet, l. c. 170 = arachnoides.
rufescens, Sweet, l. c. 170 = arachnoides.
Telephiastrum, *DC. Cat. Hort. Monsp.* 77.—Afr. austr.
trigona, *DC. Prod.* iii. 356.—Afr. austr.
ustulata, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 383.—Afr. austr.
varians, Sweet, Hort. Brit. ed. I. 170 = A. Telephiastrum.

ANACAMPSEROS, [Tourn.] Adans. Fam. ii. 248 (1763); Haw. Syn. Pl. Succ. 111 (1812) = **Sedum**, Linn. (Crassul.).
Aizoon, Haw. Syn. Pl. Succ. 112 = S. Aizoon.
albicans, Haw. l. c. 111 = S. albicans.
albida, Haw. ex DC. Prod. iii. 402 = S. Telephium?
alpestris, Jord. & Fourr. Brev. Nov. Pl. fasc. i. 22 = S. maximum.
angustifolia, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 168.—Gallia.
aprica, Jord. & Four. Brev. Pl. Nov. fasc. i. 19 = S. maximum.
arguta, Haw. Rev. Pl. Succ. 25 = S. argutum.
assurgens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 24 = S. maximum.
aurigerana, Jord. & Fourr. l. c. 30 = S. Telephium.
beugesiaca, Jord. & Four. l. c. 24 = S. Telephium.
Borderi, Jord. & Fourr. l. c. 30 = S. Telephium.
buxicola, Jord. & Fourr. l. c. 27 = S. Telephium.
caerulescens, Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse, ii. (1874) 70 = S. caeruleum?
Cepaea, Willd. Sp. Pl. ii. 762 = S. Cepaea.
cebennensis, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 18 = S. maximum.
chlorotica, Jord. & Fourr. l. c. 20 = S. maximum.
ciliaris, Haw. Syn. Pl. Succ. 113 = S. involucratum.
collina, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 23 = S. maximum.
conferta, Jord. & Fourr. Ic. Pl. Eur. i. 36 = S. Telephium.
convexa, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 29 = S. Telephium.
corsica, Jord. & Fourr. l. c. 17 = S. maximum.
delphinensis, Jord. & Fourr. l. c. 21 = S. maximum.
dentata, Haw. Rev. Pl. Succ. 26 = S. dentatum.
divaricata, Haw. Syn. Pl. Succ. 113 = S. divaricatum.
dumeticola, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 28 = S. Telephium.
erubescens, Jord. & Fourr. l. c. 18 = S. maximum.
formosa, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 168.—Gallia.
hybrida, Haw. Syn. Pl. Succ. 113 = S. hybridum.
Julliana, Jord. & Fourr. Ic. Pl. Eur. i. 36 = S. Telephium.
lapidicola, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 29 = S. Telephium.
linguaefolia, Hort. ex Lem. Illustr. Hortic. x. (1863) Misc. 81 = Cotyledon linguaefolia.
livida, Haw. Rev. Pl. Succ. 25 = S. lividum.
lugdunensis, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 28 = S. Telephium.
maxima, Haw. Syn. Pl. Succ. ed. Germ. 121 = S. maximum.
Millieri, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 23 = S. maximum.
minor, Jord. & Fourr. l. c. 19 = S. maximum.
monticulorum, Jord. & Fourr. l. c. 26 = S. Telephium.
Navieri, Jord. & Fourr. l. c. 27 = S. Telephium.
pachyphylla, Jord. & Fourr. l. c. 17 = S. maximum.
paucidens, Haw. Rev. Pl. Succ. 24.—Hab.?
pectinata, Haw. l. c. 24 = S. pectinatum.
populifolia, Haw. Syn. Pl. Succ. 113 = S. populifolium.
Pourretii, Jeanb. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse, iii. (1875–76) 556.—Gallia.
praecelsa, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 25 = S. Telephium.
praeruptorum, Jord. & Fourr. l. c. 20 = S. maximum.
purpurea, Haw. ex DC. Prod. iii. 402 = S. Telephium.
pycnantha, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 25 = S. Telephium.

ANACAMPSEROS:—
pyramidalis, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 169.—Gallia.
recurva, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 20 = S. maximum.
repens, Jord. & Fourr. l. c. 31 = S. Telephium.
rhodanensis, Jord. & Fourr. l. c. 24 = S. Telephium.
rubella, Jord. & Fourr. l. c. 30 = S. Telephium.
rupifraga, Jord. & Fourr. l. c. 29 = S. Telephium.
Saxifraga, Jord. & Fourr. l. c. 26 = S. Telephium.
sempervirens, Haw. Syn. Pl. Succ. 112 = S. Anacampseros.
serotina, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 22 = S. maximum.
silvatica, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 167.—Gallia.
spathulata, Hort. ex Steud. Nom. ed. II. i. 82 = S. spathulatum.
spectabilis, Jord. & Fourr. Ic. Pl. Eur. i. 37 = S. spectabile.
spuria, Haw. Rev. Pl. Succ. 25 = S. stoloniferum.
stellata, Haw. Syn. Pl. Succ. 114 = S. stellatum.
subalbida, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 27 = S. Telephium.
subrotunda, Jord. & Fourr. l. c. 19 = S. maximum.
subpyrenaica, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 169.—Gallia.
telephioides, Haw. Syn. Pl. Succ. 114 = S. telephioides.
ternata, Haw. l. c. = S. ternatum.
ternata, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 21 = S. maximum.
Thevensei, Timb. in Mém. Acad. Toulouse, Sér. VII. vi. (1874) 646.—Gallia.
triphylla, Haw. Syn. Pl. Succ. 111 = S. Telephium.
Verloti, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 22 = S. maximum.
viridula, Jord. & Fourr. l. c. 26 = S. Telephium.
vogesiaca, Jord. & Fourr. l. c. 31 = S. Telephium.
vulgaris, Haw. Syn. Pl. Succ. 112 = S. Telephium.

ANACAMPTA, Miers, Apocyn. S. Amer. 64 (1878) = **Tabernaemontana**, Linn. (Apocyn.).
acutissima, Miers, l. c. 66 = T. acutissima.
angulata, Miers, l. c. 65 = T. angulata.
congesta, Miers, l. c. = T. rubrostriolata.
hirtula, Miers, l. c. 67 = T. hirtula.
longifolia, Miers, l. c. 66 = T. longifolia.
submollis, Miers, l. c. 67 = T. submollis.

ANACAMPTIS, Rich. in Mém. Mus. Par. iv. (1818) 47, 55 = **Orchis**, Linn.
Brancifortii, Lindl. Gen. et Sp. Orch. 275 = O. Brancifortii.
condensata, C. Koch, in Linnaea, xxii. (1849) 285, in syn. = O. pyramidalis.
Durandi, Bréb. Fl. Norm. ed. II. 258 = O. pyramidalis.
pyramidalis, Rich. in Mém. Mus. Par. iv. (1818) 55 = O. pyramidalis.
quadripunctata, Lindl. Gen. et Sp. Orch. 275 = O. Brancifortii.
trichocera, C. Koch, in Linnaea, xxii. (1849) 286 = O. quadripunctata.

ANACARDIA, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = **Anacardium**, Linn.
rhinocarpa, St. Lag. l. c. = Anacardium Rhinocarpus.

ANACARDIUM, Linn. Syst. ed. I. (1735); Gen. ed. I. 129 (1737); Rottb. in Act. Hafn. ii. (1775) 252.
ANACARDIACEAE, Benth. & Hook. f. i. 420.
ACAJOU, Tourn. ex Adans. Fam. ii. 344 (1763).
ACAJUBA, Gaertn. Fruct. i. 192. t. 40 (1788).
CASSUVIUM, Lam. Encyc. i. 22 (1783).
MONODYNAMUS, Pohl, Pl. Bras. ii. 67. t. 144 (1831).
RHINOCARPUS, Bert. & Balb. ex H. B. & K. Nov. Gen. et Sp. vii. 5. t. 601 (1825).
brasiliense, *Barb. Rodr. in Rev. Engenh.* (28 Jul. 1883) repr. p. 5.—Bras.
curatellaefolium, *St. Hil. in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 272.—Bras.
dubium, Roxb. Hort. Beng. [88]; Fl. Ind. ii. 313 = Gluta Benghas.
encardium, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 5.—Malaya.
giganteum, *Hancock, ex Engl. in Mart. Fl. Bras.* xii. II. 410 (*errore Hance*).—Bras.

ANACARDIUM:—
giganteum, *Loud. ex Steud. Nom.* ed. II. i. 82, nomen. —Guiana.
humile, Engl. in Mart. Fl. Bras. xii. II. 411 = nanum.
humile, Mart. Reise, i. 314 = pumilum.
humile, *St. Hil. in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 272.—Bras.
latifolium, Lam. Encyc. i. 139 = Semecarpus Anacardium.
longifolium, Lam. l. c. = Semecarpus Anacardium.
mediterraneum, *Vell. Fl. Flum.* 163; iv. t. 46.—Bras.
nanum, *St. Hil. in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 272.—Bras.
occidentale, *Linn. Sp. Pl.* ed. I. 383.—Ind. occ.
officinale, Pritz. Ind. Ic. 59 = Semecarpus Anacardium.
officinarum, Gaertn. Fruct. i. 192 = Semecarpus Anacardium.
orientale, Auct. ex Steud. Nom. ed. II. i. 82 = Semecarpus Anacardium.
pumilum, Walp. Rep. i. 555 = nanum.
Rhinocarpus, *DC. Prod.* ii. 62.—Am. austr.
solitarium, Stokes, Bot. Mat. Med. ii. 159 = Semecarpus Anacardium.
Spruceanum, *Benth. ex Engl. in Mart. Fl. Bras.* xii. II. 410.—Bras.
subcordatum, Presl, Bot. Bemerk. 40 = curatellaefolium.
subterraneum, Liais, Climats, etc., Brésil, 604 = humile.

ANACHARIS, Rich. in Mém. Inst. ii. (1811) 62 = **Elodea**, Michx. (Hydroch.).
Alsinastrum, Bab. ex Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 74 = E. canadensis.
callitrichoides, Rich. in Mém. Inst. ii. (1811) 7. t. 2.—Bras.
canadensis, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 75 = E. canadensis.
chilensis, Planch. l. c.—Chili.
Hilariana, Casp. olim ex Casp. in Monatsb. Berl. Akad. (1857) 13 = E. Naias.
Matthewsii, Planch. in Ann. Sc. Nat. Sér. III. xi. (1849) 73.—Peruv.
Nuttallii, Planch. l. c. 74 = E. canadensis.
pomeranica, Peterm. Deutschl. Fl. (1849) 529. t. 82. fig. 650 = E. canadensis.
Tripteris, Steud. Nom. ed. II. i. 82 (*sphalm.* = Anacis, Schrank) = Coreopsis Tripteris.

ANACHEILIUM, Hoffmgg. ex Linnaea, xvi. (1842) Litt. 229 = **Epidendrum**, Linn. (Orchid.).
cochleatum, Hoffmgg. l. c. = E. cochleatum.

ANACHYRIS, Nees, in Hook. Kew Journ. ii. (1850) 103 = **Paspalum**, Linn. (Gramin.).
paspaloides, Nees, l. c. = P. malacophyllum.

ANACHYRIUM, Steud. Syn. Pl. Gram. 33 (1855) = praec.

ANACIS, Schrank, in Denkschr. Acad. Mucnch. v. (1817) 5 = **Coreopsis**, Linn. (Compos.).
Tripteris, Schrank, l. c. = C. Tripteris.

ANACOLOSA, Blume, Mus. Bot. Lugd. Bat. i. 250. t. 46 (1850). *OLACINEAE*, Benth. & Hook. f. i. 348.
crassipes, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 153.—Pegu.
densiflora, *Bedd. in Madr. Journ. Scr.* III. i. (1864) 38.—Travancor.
frutescens, *Blume, Mus. Bot. Lugd. Bat.* i. 251.—Java.
Griffithii, *Mast. in Hook. f. Fl. Brit. Ind.* i. 580.—Burma.
heptandra, *Maingay, ex Mast. l. c.* 581.—Malacca.
ilicoides, *Mast. l. c.* 580.—Reg. Himal.
Maingayi, *Mast. l. c.*—Singapore.
Pervilleana, *Baill. Adansonia*, iii. (1862-63) 119.—Ins. Masear.
puberula, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 297.—Ins. Andaman.

ANACOLUS, Griseb. in DC. Prod. ix. 70 (1845) = **Hockinia**, Gardn. (Gentian.).
Gardneri, Griseb. l. c. = H. montana.

ANACTORION, Rafin. Fl. Tellur. iv. 34 (1836) = **Synnotia**, Sweet (Irid.).
bicolor, Rafin. l. c. = S. bicolor.
galeatum, Rafin. l. c. = S. galeata.

ANACTINIA, Remy, in C. Gay, Fl. Chil. iv. 8 (1849) = **Nardophyllum**, Hook. (Compos.).
Hookeri, Remy, l. c. = N. humile.

ANACTIS, Cass. in Dict. Sc. Nat. xlvii. 510 (1827) = **Atractylis**, Linn. (Compos.).
caespitosa, Cass. l. c. l. 56 = At. caespitosa.
glabra, Rafin. Fl. Tellur. ii. 44 = Aster vimineus?
pilosa, Rafin. l. c. = Carthamus lanatus.
serratuloides, Cass. in Dict. Sc. Nat. l. 56 = At. serratuloides.

ANACYCLODON, Jungh. in Nat. en Geneesk. Arch. Néerl. Indie, ii. (1845) 49 = **Leucopogon**, R. Br. (Epacrid.).
pungens, Jungh. l. c. = L. javanicus.

ANACYCLUS, Linn. Gen. ed. I. 256 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 419.
ARTHROLEPIS, Boiss. Diagn. Scr. I. xi. 14 (1849).
CYRTOLEPIS, Less. in Linnaea, vi. (1831) 166.
HIORTHIA, Neck. Elem. i. 97 (1790).
LEUCOCYCLUS, Boiss. Diagn. Ser. I. xi. 13 (1849).
alexandrinus, *Willd. Sp. Pl.* iii. 2173.—Afr. bor.
anomalus, J. Gay, ex Boiss. Fl. Orient. iii. 283 = Anthemis palestina.
anthemoides, Lag. ex Spreng. Syst. iii. 497 = Anthemis abrotanifolia.
aristulatus, Link, ex Nym. Consp. 363 = clavatus.
aureus, Brot. Fl. Lusit. i. 363 = radiatus.
aureus, Lam. ex DC. Fl. Fr. iv. 202 = Matricaria aurea.
aureus, Linn. Mant. ii. 287 = Anthemis aurea.
australis, Sieber, ex Spreng. Syst. iii. 497 = Cotula australis.
Barrelieri, Guss. Pl. Rar. 357 = Achillea Barrelierii.
bicolor, Pers. Syn. ii. 465 = radiatus.
Candolii, Nym. Syll. 8 = clavatus.
ciliatus, Traut. in Bull. Soc. Nat. Mosc. xli. (1868) I. 461 = Anthemis ciliata.
clavatus, *Pers. Syn.* ii. 465.—Reg. Mediterr.
creticus, Linn. Sp. Pl. 892 = Anthemis cretica.
cyrtolepidioides, *Pomel, Nouv. Mat. Fl. Atl.* 54.—Afr. bor.
depressus, *Ball, in Journ. Bot.* xi. (1873) 365.—Marocco.
dissimilis, *Pomel, Nouv. Mat. Fl. Atl.* 53.—Afr. bor.
divaricatus, Cav. ex Steud. Nom. ed. i. 41 = clavatus.
formosus, *Fenzl, ex Boiss. Diag.* Ser. I. xi. 14.—Cilicia.
hirsutus, Lam. Fl. Fr. ii. 47 = valentinus.
inconstans, *Pomel, Nouv. Mat. Fl. Atl.* 52.—Afr. bor.
inflatus, *Lehm. ex Steud. Nom.* ed. II. i. 82.—Sibir.
lanuginosus, Moench, Meth. 581 = valentinus.
lincarilobus, *Boiss. & Reut. Pugill. Pl. Hispan.* 57.—Hispan.
maroccanus, *Ball, in Journ. Bot.* xi. (1873) 365.—Marocco.
mauritanicus, *Pomel, Nouv. Mat. Fl. Atl.* 55.—Afr. bor.
membranaceus, *Benth. & Hook. f. Gen.* iii. 419.—Syria.
mucronulatus, Guss. Pl. Rar. 356 = Achillea Barrelierii.
mucronulatus, Hort. ex Steud. Nom. ed. II. i. 82 = clavatus.
nigellaefolius, *Boiss. Diagn.* Ser. I. xi. 13.—Syria; Assyr.
officinarum, *Hayne, Arzneigew.* ix. t. 46.—Europ. austr.
orientalis, Linn. Sp. Pl. 892 = Anthemis montana.
pallescens, Guss. Fl. Sic. Syn. ii. 494 = radiatus.
pectinatus, Bory & Chaub. Fl. Pélop. 59. t. 30 = Anthemis tygetea.
pedunculatus, *Pers. Syn.* ii. 465.—Barbar.
prostratus, *Pomel, Nouv. Mat. Fl. Atl.* 52.—Afr. bor.
Pseudopyrethrum, *Aschers. in Bonplandia*, vi. (1858) 123.—Hab.?
pubescens, Brot. ex Link, Handb. i. 763; Reichb. Fl. Exc. ii. 225 = clavatus.
pulcher, Bess. ex DC. Prod. vi. 16 = officinarum.
purpurascens, DC. Fl. Fr. v. 481 = radiatus.
Pyrethraria, Spreng. Syst. iii. 497 = Cotula Pyrethraria.
Pyrethrum, *DC. Fl. Fr. Suppl.* 480.—Afr. bor.; Orient.
radiatus, *Loisel. Fl. Gall.* ed. I. 583; *Pers. Syn.* ii. 465.—Reg. Mediterr.
tomentosus, DC. Fl. Fr. Suppl. 481 = clavatus.
truncatus, J. Gay, in Boiss. Fl. Orient. iii. 283 = Anthemis palestina.
valentinus, *Linn. Sp. Pl.* 892.—Reg. Mediterr.

ANACYLANTHUS, Steud. Nom. ed. II. i. 82 (1840), sphalm. = **Ancylanthos**, Desf. (Rubiac.).

ANADELPHIA, Hack. in Engl. Bot. Jahrb. vi. (1885) 240 = **Andropogon**, Linn. (Gramin.).
virgata, Hack. l. c. 241 = Andr. leptocomus.

ANADENDRUM, Schott, in Bonplandia, v. (1857) 45. *AROIDEAE*, Benth. & Hook. f. iii. 991.
affine, *Schott, in Ann. Mus. Lugd. Bat.* i. 282.—Sumatra.
cordatum, *Schott, l. c.* 283.—Sumatra.
Lobbi, Schott, in Bonplandia, v. (1857) 45 = montanum.
marginatum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 116.—Ind. or.
medium, Schott, in Bonplandia, v. (1857) 45 = Epipremnum medium.
montanum, *Schott, l. c.*—Java.
punctulatum, Schott, Prod. Aroid. 393 = Monstera punctulata.
semivestitum, Schott, in Ann. Mus. Lugd. Bat. i. 283 = affine.

ANADENDRON, Wight, Illustr. Ind. Bot. ii. 164 (1850) err. typ. = **Anodendron**, A. DC. (Apocyn.).

ANADEMIA, Agardh, Aphor. 236 (1826), sphalm. = seq.

ANADENIA, R. Br. in Trans. Linn. Soc. x. (1809) 166 = **Grevillea**, R. Br. (Proteac.).
anethifolia, A. Cunn. ex Benth. Fl. Austral. v. 483 = G. anethifolia.
Aquifolium, Lindl. Swan River App. 31 = G. monticola.
Caleyi, R. Br. Prot. Nov. 16 = G. ramosissima.
filifolia, Endl. in Ann. Wien. Mus. ii. (1838) 209 = G. didymobotrya.
flexuosa, Lindl. Swan River App. 31 = G. flexuosa.
gracilis, Lindl. l. c. = G. Synapheae.
hakeoides, Lindl. l. c. 30 = Hakea undulata.
ilicifolia, R. Br. in Trans. Linn. Soc. x. (1809) 167 = G. ilicifolia.
integrifolia, Endl. in Ann. Wien. Mus. ii. (1838) 209 = G. integrifolia.
Manglesii, R. Grah. in Hook. Ic. Pl. t. 337 = Manglesia cuneata.
pulchella, R. Br. in Trans. Linn. Soc. x. (1809) 167 = G. pulchella.
Roëi, Endl. Gen. Suppl. iv. II. 83 = G. didymobotrya.
tenuiflora, Lindl. Swan River App. 31 = G. tenuiflora.
trifida, R. Br. in Trans. Linn. Soc. x. (1809) 167 = G. trifida.
triternata, A. Cunn. ex Heynh. Nom. ii. 34 = G. triternata.

ANAECTOCALYX, Triana, ex Benth. & Hook. f. Gen. i. 765 (1867). *MELASTOMACEAE*, Benth. & Hook. f. i. 765.
bracteosa, *Triana, l. c. & in Trans. Linn. Soc.* xxviii. (1871) 97.—Venezuela.

ANAECTOCHILUS, Lindl. Orch. 498 (1840) = **Anoectochilus**, Blume (Orchid.).

ANAFRENIUM, Arn. in Hook. Lond. Journ. Bot. iii. (1841) 148 = ANAPHRENIUM, E. Mey. = **Rhus**, Linn. (Anacard.).

ANAGALLIDASTRUM, Mich. ex Adans. Fam. ii. 230 (1763) = **Centunculus**, Linn. (Primul.).

ANAGALLIDIUM, Griseb. Gen. et Sp. Gent. 311 (1836) = **Swertia**, Linn. (Gentian.).
dichotoma, Griseb. l. c. 312 = S. dichotoma.
tetrapetalum, Griseb. l. c. = S. tetrapetala.

ANAGALLIS, [Tourn.] Linn. Syst. ed. I. (1735); Gen. ed. I. 43 (1737). *PRIMULACEAE*, Benth. & Hook. f. ii. 637.
EUPAREA, Banks & Soland. in Gaertn. Fruct. i. 230. t. 50 (1788).
IRASEKIA, S. F. Gray, Nat. Arr. ii. 299 (1821).
JIRASEKIA, F. W. Schmidt, Neue u. Selt. Pfl. 49 (1793).
TIRASEKIA, G. Don, in Sweet, Hort. Brit. ed. III. 563 (1839).

ANAGALLIS:—
alba, Buch.-Ham. ex Steud. Nom. ed. II. i. 83.—(Quid?) Ind. or.
alternifolia, *Cav. Ic.* vi. 3. t. 506.—Chili.
angustifolia, Salisb. Prod. 121 = linifolia.
aquatica, Erndl, ex Ledeb. Fl. Ross. iii. 31 = Samolus Valerandi.
arabica, *Duby, in DC. Prod.* viii. 69.—Arab.
arvensis, *Linn. Sp. Pl.* 148.—Europ.; As. temp.
caerulea, Lam. Fl. Fr. ed. I. ii. 285 = arvensis.
capensis, *Linn. Sp. Pl.* 149.—Afr. austr.
carnea, Schrank, Baier. Fl. i. 461 = arvensis.
Centunculus, Afzel. Veget. Suecan. (1785) 1 = Centunculus minimus.
collina, Schousb. Marok. 64 = linifolia.
crassifolia, *Thore, Chlor. Land.* 62.—Hispan.; Marocco.
filiformis, *Cham. & Schlecht. in Linnaea*, i. (1826) 225.—Bras.
foemina, Mill. Gard. Dict. ed. VIII. n. 2 = arvensis.
fruticosa, Vent. Choix, 14 = linifolia.
Gmelini, *Steud. Nom.* ed. II. i. 83.—Camtsch.
grandiflora, Andr. Repos. t. 367 = linifolia.
Huttoni, *Harv. in Proc. Dub. Univ. Zool. & Bot. Assoc.* i. (1859) 141.—Afr. austr.
indica, Sweet, Fl. Gard. t. 132 = arvensis.
Jacquemontii, Duby, in DC. Prod. viii. 69 = arvensis.
latifolia, Linn. Sp. Pl. 148 = arvensis.
linifolia, *Linn. l. c.* ed. II. 212.—Europ.
Marryattae ×, Sweet, ex G. Don, in Sweet, Hort. Brit. ed. III. 563.
mas, Vill. Hist. Pl. Dauph. ii. 461 = arvensis.
Monelli, Bieb. Fl. Taur. Cauc. i. 143 = arvensis.
Monelli, Linn. Sp. Pl. 148 = linifolia.
multangularis, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iii. 503 = Lysimachia pyramidalis.
nummularifolia, *Baker, in Journ. Linn. Soc.* xx. (1884) 196.—Madag.
orientalis, Hort. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. xi. 49 = arvensis.
ovalis, Ruiz & Pav. Fl. Per. ii. 8 = Centunculus pentandrus.
palustris, [Clairv.] Man. Herb. 59 = tenella.
parviflora, *Hoffmgg. & Link, Fl. Port.* i. 325. t. 64.—Lusitan.
parviflora, Loisel. in Mém. Soc. Linn. Par. vi. (1827) 491 = arvensis.
pedunculata, Salzm. ex Duby, in DC. Prod. viii. 72 = Centunculus pentandrus.
peploides, *Baker, in Journ. Linn. Soc.* xx. (1884) 196.—Madag.
Philipsii, Hort. ex Duby, in DC. Prod. viii. 70—linifolia.
phoenicea, Scop. Fl. Carn. ed. II. i. 139 = arvensis.
pilosa, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 83 = Lysimachia japonica.
platyphylla, *Baudo, in Exped. Alg.* t. 67.—Algeria.
pulchella, Salisb. Prod. 120 = arvensis.
pumila, Sw. Prod. Veg. Ind. Occ. 40 = Centunculus pentandrus.
punctifolia, Stokes, Bot. Mat. Med. i. 305 = arvensis.
pusilla, Salisb. Prod. 121 = Centunculus minimus.
repens, DC. Fl. Fr. v. 381 = arvensis.
repens, *Pomel, Nouv. Mat. Fl. Atl.* 126.—Algeria.
rubricaulis, Boj. ex Duby, in DC. Prod. viii. 72 (= *Micropyxis rubricaulis*).
serpens, *Hochst. ex A. DC. Prod.* viii. 668.—Abyss.
serpyllifolia, Dum. Fl. Belg. 29 = tenella.
sessilis, Salzm. ex Duby, in DC. Prod. viii. 72 = Centunculus pentandrus.
spicata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 6.—Malaya.
tenella, *Linn. Mant.* 335.—Europ.
tenuicaulis, *Baker, in Journ. Bot.* xx. (1882) 172.—Madag.
verticillata, All. Fl. Pedem. i. 87 = arvensis.
Webbiana, Penny, ex Loud. Gard. Mag. v. (1829) 471 = *Marryattae*.
Wellsiana ×, *Penny, ex Loud. Hort. Brit. Suppl.* i. 580.
Willmoreana, Don, ex Hook. Bot. Mag. t. 3380 = linifolia.

ANAGALLOIDES, Krock. Fl. Siles. ii. 1. 398 (1790) = **Vandellia**, Linn. (Scrophul.).
procumbens, Krock. l. c. = V. erecta.

ANAGANTHOS, Hook. f. in Fl. N. Zeal. i. 226 (1854) = **Australina**, Gaudich. (Urticac.).

ANAGLYPHA, DC. Prod. v. 311 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 326.
OXYLAENA, Benth. in Hook. Ic. Pl. t. 1109 (1876).
acicularis, *Benth. l. c.* xii. 9. t. 1109.—Afr. austr.
aspera, *DC. Prod.* v. 311.—Afr. austr.

ANAGYRIS, Linn. Syst. ed. I. (1735); Gen. ed. I. 219 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 465.
barbata, R. Grah. in Wall. Cat. n. 5341 = Thermopsis barbata.
cretica, *Mill. Gard. Dict.* ed. VIII. n. 2.—Ins. Archip.
foetida, *Linn. Sp. Pl.* 374.—Reg. Mediterr.; Orient.
foetida, Lour. Fl. Cochinch. 260; ed. Willd. i. 318 = sinensis.
glauca, Hort. ex Loud. Hort. Brit. 162 = foetida.
indica, Hort. ex Steud. Nom. ed. II. i. 83 = sinensis.
indica, Lindl. in Trans. Hort. Soc. vii. (1830) 245 = Piptanthus nepalensis.
inodora, *Lour. Fl. Cochinch.* 260.—Cochinch.
latifolia, *Brouss. ex Willd. Enum. Hort. Berol.* 439.—Teneriffa.
neapolitana, Tenore, Syll. 198 = foetida.
nepalensis, R. Grah. in Wall. Cat. n. 5340 = Piptanthus nepalensis.
sinensis, *Steud. ex DC. Prod.* ii. 98 ('forte ad Ormosiam referend.').—China.

ANAITIS, DC. Prod. v. 628 (1836) = **Sanvitalia**, Linn. (Compos.).
acapulcensis, DC. l. c. 629 = S. acapulcensis.

ANALECTIS, Juss. in Jaume St. Hil. Expos. ii. 362 (1805) = **Symphorema**, Roxb. (Verben.).

ANALITON, Rafin. Fl. Tellur. iii. 46 (1836) = **Rumex**, Tourn. (Polygon.).
bipinnatus, Rafin. l. c. = R. lacerus.

ANALYRIUM, E. Mey. Zwei Pfl. Docum. 163 (1843), *nomen*; Presl, in Abh. Boehm. Ges. V. iii. (1845) 505 = **Peucedanum**, Koch (Umbell.).
millefolium, E. Mey. l. c. = P. millefolium.

ANAMENIA, Vent. Malm. 22. t. 22 (1803) = **Knowltonia**, Salisb. (Ranunc.).
capensis, Hoffmgg. Verz. Pfl. Nachtr. i. 204 = K. rigida.
coriacea, Vent. Malm. 22 = K. rigida.
daucifolia, Vent. ex Cav. in Anal. Cienc. Nat. vi. (1803) 344 = K. daucifolia.
gracilis, Vent. Malm. i. 22, obs. = K. gracilis.
hirsuta, Vent. ex Cav. in Anal. Cienc. Nat. vi. (1803) 344 = K. hirsuta.
laserpitiifolia, Vent. ex Cav. l. c. = K. vesicatoria.
vesicatoria, Dum.-Cours. Bot. Cult. iv. 438 = K. vesicatoria.

ANAMIRTA, Colebr. in Trans. Linn. Soc. xiii. (1822) 52. *MENISPERMACEAE*, Benth. & Hook. f. i. 35.
Baueriana, Endl. Atakt. Bot. t. 39, 40 = A. Cocculus.
Cocculus, Wight & Arn. Prod. i. 446 = paniculata.
flavescens, Miq. Fl. Ind. Bat. i. II. 79 = A. Cocculus.
florescens, C. Muell. in Walp. Ann. vii. 63, sphalm. = flavescens.
jucunda, *Miers, Contr. Bot.* iii. 55.—Java.
lemniscata, *Miers, l. c.* 54.—Java.
Lourieri, *Pierre, Fl. For. Coch. Chin.* t. 110.—Cambodia.
luctuosa, *Miers, Contr. Bot.* iii. 55.—Java.
paniculata, *Colebr. in Trans. Linn. Soc.* xiii. (1822) 66.—Ind. or.
populifera, *Miers, Contr. Bot.* iii. 54.—Timor.
racemosa, Colebr. ex Steud. Nom. ed. II. i. 83 = Cocculus suberosus.
toxifera, Miers, Contr. Bot. iii. 53 = paniculata.

ANAMOMIS, Griseb. Fl. Brit. W. Ind. 240 (1864) = **Myrtus**, Linn.
esculenta, Griseb. l. c. in obs. = Eugenia esculenta.
ferruginea, Griseb. Cat. Pl. Cub. 91 = Eugenia ferruginea?
fragrans, Griseb. Fl. Brit. W. Ind. 240 = Myrcia fragrans?

ANAMOMIS:—
Guayabillo, Griseb. Cat. Pl. Cub. 90 = Eugenia Guayabillo.
punctata, Griseb. Fl. Brit. W. Ind. 240 = Myrcia Balbisiana.

ANAMORPHA, Karst. & Triana, in Linnaea, xxviii. (1856) 443 = **Melochia**, Linn. (Stercul.).
walthérioides, Karst. & Triana l. c. = M. lupulina.

ANANAS, Gaertn. Fruct. i. 30. t. 11 (1788) = **Bromelia**, Linn.
Pinguin, Gaertn. l. c. = B. Pinguin.

ANANAS, Tourn. ex Linn. Syst. ed. I. (1735); Mill. Gard. Dict. ed. VI. (1752); Adans. Fam. ii. 67 (1763).
BROMELIACEAE, Benth. & Hook. f. iii. 662.
ACANTHOSTACHYS, Link, Klotzsch & Otto, Ic. Pl. Rar. i. 21. t. 9. (1841).
ANANASSA, Lindl. Bot. Reg. sub t. 1068 & t. 1081 (1827).
arvensis, *Steud. Nom.* ed. II. i. 83.—Bras.
bracteatus, *Schult. f. Syst.* vii. 1286.—Bras.
debilis, Schult. f. l. c. 1287 = sativus.
glaber, Mill. Gard. Dict. ed. VIII. n. 3 (et corr.) = sativus.
lucidus, Mill. l. c. n. 4 = sativus.
macrodontes, *E. Morr. Belg. Hortic.* xxviii. (1878) 140. t. 4 et 5.—Bras.
Mordilona, Hort. Linden, ex E. Morr. in Belg. Hortic. xxix. (1879) 302 = sativus.
muricatus, Schult. f. Syst. vii. 1287 = bracteatus.
ovatus, Mill. Gard. Dict. ed. VIII. n. 1 = sativus.
Porteanus, Hort. Veitch, ex C. Koch, Wochenschr. xiv. (1871) 130 = sativus.
pyramidalis, Mill. Gard. Dict. ed. VIII. n. 2 = sativus.
Sagenaria, Schult. f. Syst. vii. 1286 = bracteatus.
sativus, *Schult. f. l. c.* 1283.—Am. trop.
semiserratus, Schult. f. l. c. 1286 = sativus.
serotinus, Mill. Gard. Dict. ed. VIII. n. 5 = sativus.
viridis, Mill. l. c. n. 6 = sativus.

ANANASSA, Lindl. Bot. Reg. sub t. 1068 et t. 1081 (1827) = **Ananas**, Tourn. (Bromel.).
bracamorensis, Linden, Cat. n. 98. (1878) 3 = Ananas sativus.
bracteata, Lindl. Bot. Reg. sub t. 1068 et t. 1081 = Ananas bracteatus.
debilis, Lindl. l. c. sub t. 1068 = Ananas sativus.
lucida, Lindl. l. c. = Ananas sativus.
Mordilona, Linden, Cat. n. 22. (1869) 2 = Ananas sativus.
muricata, D. Dietr. Syn. Pl. ii. 1088 = Ananas bracteatus.
Porteana, Carr. in Rev. Hortic. (1878) 140 = Ananas sativus.
Sagenaria, D. Dictr. Syn. Pl. ii. 1088 = Ananas bracteatus.
sativa, Lindl. Bot. Reg. sub t. 1068 = Ananas sativus.
semiserrata, D. Dictr. Syn. Pl. ii. 1088 = Ananas sativus.

ANANDRIA, Siegesb. ex Linn. Amoen. Acad. i. 243. t. 11 (1745) = **Gerbera**, Gronov. (Compos.).
Bellidiastrum, DC. Prod. vii. 40 = G. Anandria.
dimorpha, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) II. 88 = G. Anandria.
discoidea, Less. in Linnaea, v. (1830) 347. t. 3 = G. Anandria.
radiata, Less. l. c. 346. t. 4 = G. Anandria.

ANANTHERIX, Nutt. Gen. Am. i. 169 (1818) = **Asclepiodora**, A. Gray (Asclep.).
angustifolia, Rafin. Atl. Journ. 146.—Am. bor.
Bungei, G. Don, Gen. Syst. iv. 146 = Pycnostelma chinense.
connivens, A. Gray, in Proc. Am. Acad. xii. (1877) 66 = Asc. viridis.
decumbens, Nutt. in Trans. Am. Phil. Soc. v. (1837) 203 = Asc. decumbens.
grandiflora, Rafin. in New Fl. Am. iv. 59.—Am. bor.
Nuttaliana, G. Don, Gen. Syst. iv. 146 = Asc. decumbens.
ovata, Rafin. Atl. Journ. 146.—Am. bor.
paniculata, Nutt. in Trans. Am. Phil. Soc. v. (1837) 203 = Asc. viridis.
pumilis, Nutt. l. c.—Am. bor.
Torreyana, G. Don, Gen. Syst. iv. 146 = Asc. viridis.
verrucosa, Rafin. New Fl. Am. iv. 60 = Asc. viridis.
viridis, Nutt. Gen. Am. i. 169 = Asc. viridis.

ANANTHOCYCLUS, Vaill. ex Linn. Syst. ed. I. (1735) = **Cotula**, Tourn. (Compos.).

ANANTHOPUS, Rafin. Fl. Ludov. 20 (1817) = **Commelina**, Linn.
clandestinus, Rafin. l. c. 21 = C. ignorata.
cordatus, Rafin. l. c. 22 = C. Rafinesquei.
undulatus, Rafin. Fl. Tellur. iii. 70 = C. undulata.

ANAPHALIS, DC. Prod. vi. 271 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 303.
MARGARIPES, DC. ex Steud. Nom. ed. II. ii. 101 (1841).
adnata, *DC. Prod.* vi. 274.—Reg. Himal.
alata, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 478.—As. or.
arancosa, *DC. Prod.* vi. 275.—Reg. Himal.
aristata, *DC. in Wight, Contrib.* 21.—Ind. or.
Beddomei, *Hook. f. Fl. Brit. Ind.* iii. 282.—Ind. or.
brevifolia, *DC. Prod.* vi. 273.—Zeylan.
Busua, DC. l. c. 275 = araneosa.
chionantha, DC. l. c. 273 = Royleana.
cinnomomea, *C. B. Clarke, Comp. Ind.* 104.—Ind. or.; China; Japon.
contorta, *Hook. f. Fl. Brit. Ind.* iii. 284.—Reg. Himal.
cuneifolia, *Hook. f. l. c.* 280.—Reg. Himal.
cutchica, *C. B. Clarke, Comp. Ind.* 111.—Kutch.
elliptica, DC. Prod. vi. 274 = oblonga.
Falconeri, C. B. Clarke, Comp. Ind. 107 = contorta.
fruticosa, *Hook. f. Fl. Brit. Ind.* iii. 282.—Zeylan.
Griffithii, *Hook. f. l. c.* 280.—Reg. Himal.
Hancockii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 479.—China.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 282.—Reg. Himal.
japonica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 480.—Japon.
javanica, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 124.—Java.
lactea, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 479.—As. or.
leptophylla, *DC. Prod.* vi. 273.—Ind. or.
linearis, DC. l. c. 274 = marcescens?
longifolia, DC. l. c.—Java.
marcescens, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 286.—Ind. or.
margaritacea, *Benth. & Hook. f. Gen.* ii. 303.—Am. bor.
monocephala, DC. Prod. vi. 272 = nubigena.
mucronata, DC. l. c. = nubigena.
neelgerryana, *DC. l. c.*—Ind. or.
Notoniana, *DC. l. c.* 273.—Ind. or.
nubigena, *DC. l. c.* 272.—Reg. Himal.
oblonga, *DC. l. c.* 274.—Ind. or.
oligandra, *DC. l. c.* 275.—Madag.
polylepis, DC. l. c. 272 = Royleana.
pterocaulon, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 478.—As. or.
racemifera, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 310.—As. or.
Royleana, *DC. Prod.* vi. 272.—Reg. Himal.
Royleana, C. B. Clarke, Comp. Ind. 104 = Griffithii.
semidecurrens, C. B. Clarke, l. c. 109 = araneosa.
sinica, *Hance, in Journ. Bot.* xii. (1874) 261.—China.
Stoliczkai, *C. B. Clarke, Comp. Ind.* 108.—Tibet. occ.
subumbellata, *C. B. Clarke, l. c.* 104.—Reg. Himal.
tenella, DC. Prod. vi. 273 = contorta.
Thwaitesii, *C. B. Clarke, Comp. Ind.* 110.—Zeylan.
Timmua, D. Don, Prod. Fl. Nep. 174 = cinnamomea.
triplinervis, *Sims, ex C. B. Clarke, Comp. Ind.* 105.—Reg. Himal.
triplinervis, Benth. ex Hance, in Journ. Bot. xvi. (1878) 12 = Hancockii.
virgata, *Thoms. in C. B. Clarke, Comp. Ind.* 107.—Reg. Himal.
viscida, *DC. Prod.* vi. 274.—Java.
Wightiana, Thw. Enum. Pl. Zeyl. 166 = zeylanica.
Wightiana, *DC. Prod.* vi. 273.—Ind. or.
xylorhiza, *Sch. Bip. ex Hook. f. Fl. Brit. Ind.* iii. 281.—Reg. Himal.
yedoensis, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 480.—Japon.
zeylanica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 286.—Zeylan.
Zollingeri, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 54 = Antennaria saxatilis.
Zollingeriana, Sch. Bip. ex Zoll. Syst. Verz. Ind. Archip. 124 = praec.

ANAPHRAGMA, Stev. in Bull. Soc. Nat. Mosc. iv. 265 (1832) = **Astragalus**, Tourn. (Legum.).

ANAPHRENIUM, E. Mey. ex Endl. Gen. Suppl. i. 1425 (1841). *ANACARDIACEAE*, [Benth. & Hook. f. i. 419].
HEERIA, Meissn. Gen. Comm. 55 (1837).
OZOROA, Delile, in Ann. Sc. Nat. Sér. II. xx. (1843) 91. t. 1.
ROEMERIA, Thunb. Nov. Gen. Pl. ix. 130 (1798), pro parte.
abyssinicum, *Hochst. in Flora*, xxvii. (1844) 32.—Afr. trop.
argenteum, *E. Mey. Zwei Pfl. Docum.* 163.—Afr. austr.
concolor, *E. Mey. l. c.*—Afr. austr.
dispar, *E. Mey. l. c.*—Afr. austr.
longifolium, Bernh. in Flora, xxvii. (1844) 1. 349 = Protorhus longifolia.
mucronatum, *Bernh. l. c.*—Afr. austr.
mucronifolium, *Sond. in Harv. & Sond. Fl. Cap.* i. 522 = praec.
paniculosum, *Engl. in DC. Monog. Phan.* iv. 358.—Afr. austr.
pulcherrimum, *Schweinf. Fl. Aethiop. Beitr.* i. 32.—Afr. trop.

ANAPHYLLUM, Schott, Gen. Aroid. t. 83 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 996.
Wightii, *Schott, l. c.*—Ind. or.

ANAPODOPHYLLUM, Tourn. ex Moench, Meth. 277 (1794) = **Podophyllum**, Linn. (Berberid.).
peltatum, Moench, l. c. = P. peltatum.

ANARMODIUM, Schott, in Bonplandia, ix. 368 (1861) = **Dracunculus**, Schott (Aroid.).
canariense, Schott, l. c. = D. canariensis.

ANARMOSA, Miers, ex Hook. Bot. Misc. iii. (1833) 338 = **Tetilla**, DC. (Saxifrag.).
gracilis, Miers, ex Hook. l. c. = T. hydrocotylifolia.

ANARRHINUM, Desf. Fl. Atlant. ii. 51 (1800). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 933.
CARDIOTHECA, Ehrenb. ex Steud. Nom. ed. II. i. 282 (1840).
SIMBULETA, Forsk. Fl. Aegypt. Arab. 115 (1775).
abyssinicum, *Jaub. & Spach, Illustr. Pl. Or.* v. 52. t. 447.—Abyss.
arabicum, *Jaub. & Spach, l. c.* v. 50. t. 446.—Arabia.
bellidifolium, *Desf. Fl. Atlant.* ii. 51; *Chav. Monog.* 175. t. 10.—Reg. Mediterr.
bellidifolium, Fenzl, ex Jaub. & Spach, Illustr. Pl. Or. v. 54 = orientale.
brevifolium, *Coss. & Kral. in Bull. Soc. Bot. Fr.* iv. (1857) 407.—Afr. bor.
canadense, Poir. Encyc. Suppl. i. 334, in obs. = Linaria canadensis.
corsicum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 41 = bellidifolium.
crassifolium, Willd. Sp. Pl. iii. 261 = Linaria origanifolia.
Duriminium, *Pers. Syn.* ii. 159.—Lusitania.
ecalcaratum, Fenzl, ex Jaub. & Spach, Illustr. Pl. Or. v. 54 = orientale.
fruticosum, *Desf. Fl. Atlant.* ii. 52.—Afr. bor.
fruticosum, Schimp. ex Benth. in DC. Prod. x. 289 = orientale.
hirsutum, Hoffmgg. & Link, Fl. Port. 227. t. 33 = A. Duriminium.
incarnatum, Poir. Encyc. Suppl. iv. 18, in obs. = Linaria bipartita.
Linnaeanum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 41 = bellidifolium.
laxiflorum, *Boiss. Elench.* 71.—Hispan.
lusitanicum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 41 = bellidifolium.
murale, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 557 = A. Duriminium.
orientale, *Benth. in DC. Prod.* x. 289.—Oriens.
pedatum, *Desf. Fl. Atlant.* ii. 51. t. 141.—Afr. bor.
pedicellatum, *T. Anders. in Journ. Linn. Soc.* v. *Suppl.* i. (1860) 26.—Aden.
pubescens, *Fresen. in Mus. Senkenb.* i. 89.—Arab. petr.
pubescens, Hort. ex Loud. Hort. Brit. 250 = A. Duriminium.
tenellum, *Willd. Sp. Pl.* iii. 262.—Hispan.
violaceum, Dulac, Fl. Hautes-Pyr. 373 = bellidifolium.

ANARTHRIA, R. Br. Prod. 248 (1810). *RESTIACEAE*, Benth. & Hook. f. iii. 1030.
canaliculata, Nees, in Lehm. Pl. Preiss. ii. 62 = gracilis.
gracilis, *R. Br. Prod.* 249.—Austral.
gracilis, Nees, ex Benth. Fl. Austral. vii. 216 = Lepyrodia Anarthria.
grandiflora, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 50 = gracilis.
humilis, Nees, l. c. = gracilis.
ischaemoides, Nees, in Lehm. Pl. Preiss. ii. 62 = gracilis.
laevis, *R. Br. Prod.* 249.—Austral.
pauciflora, R. Br. l. c. 249 = prolifera.
polyphylla, *Nees, in Lehm. Pl. Preiss.* ii. 63.—Austral.
prolifera, *R. Br. Prod.* 249.—Austral.
scabra, *R. Br. l. c.*—Austral.

ANARTHROPHYLLUM, Benth. in Benth. & Hook. f. Gen. i. 478 (1865). *LEGUMINOSAE*, Benth. & Hook. f. i. 478.
andicolum, *Phil. f. Cat. Pl. Chil.* 52.—Chili.
Bergii, *Hieron. in Bol. Acad. Nac. Cordova*, iii. (1879) 340.—Reg. argent.
Cumingii, *Phil. f. Cat. Pl. Chil.* 52.—Chili.
desideratum, *Benth. in Benth. & Hook. f. Gen.* i. 478.—Patagon.
elegans, *Phil. f. Cat. Pl. Chil.* 52.—Chili.
juniperinum, *Phil. f. l. c.*—Chili.
rigidum, *Hieron. in Bol. Acad. Nac. Cordova*, iii. (1879) 340.—Reg. Argent.
umbellatum, *Phil. f. Cat. Pl. Chil.* 52.—Chili.

ANARTHROSYNE, E. Mey. Comm. Pl. Afr. Austr. 124 (1835) = **Pseudarthria**, Wight & Arn. Legum.
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 204 = Desmodium spirale.
cordata, Klotzsch, in Peters, Reise Mossamb. Bot. 39. t. 7 = Desmodium latifolium.
densiflora, Klotzsch, l. c. 41 = P. Hookeri.
gracilis, Klotzsch, l. c. 40 = P. confertiflora.
robusta, E. Mey. Comm. Pl. Afr. Austr. 124 = P. Hookeri.

ANARTIA, Miers, Apocyn. S. Amer. 79. (1878) = **Tabernaemontana**, Linn. (Apocyn.).
bogotensis, Miers, l. c. 82.—N. Granat.
flavescens, Miers, l. c. = Maloutia flavescens.
flavicans, Miers, l. c. = T. flavicans.
glabrata, Miers, l. c. 81 = T. oblongifolia.
Meyeri, Miers, l. c. 80 = T. Meyeri.
recurva, Miers, l. c. = T. recurva.
Wulfschlaegelii, Miers, l. c. 81 = T. Wullschlaegelii.

ANASCHOVADI, Adans. Fam. ii. 122 (1763) = **Elephantopus**, Vaill. (Compos.).

ANASSER, Juss. Gen. 150 (1789) = **Geniostoma**, Forst. (Logan.).
Borboniae, J. F. Gmel. Syst. i. 303 = G. borbonicum.
borbonica, Lam. Illustr. ii. 40 = G. borbonicum.
moluccana, Lam. l. c. = Strychnos Pseudo-quina.
Rumphii, Span. in Linnaea, xv. (1841) 325 = Pittosporum timorense.

ANASSERA, Pers. Syn. i. 265 (1805) = praec.

ANASTATICA, Linn. Gen. ed. I. 191 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 71.
HIERICONTIS, Adans. Fam. ii. 421 (1763).
IERIKONTIS, Adans. l. c. 565 (1763).
hierachuntica, Crantz, Stirp. Austr. ed. II. fasc. i. 7 = Euclidium syriacum.
hierochuntica, *Linn. Sp. Pl.* ed. II. 895.—Oriens.
littoralis, Salisb. Prod. 266 = hierochuntica.
syriaca, Linn. Sp. Pl. ed. II. 895 = Euclidium syriacum.

ANASTRABE, E. Mey. ex Benth. in Hook. Comp. Bot. Mag. ii. (1836) 54. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 937.
integerrima, *E. Mey. ex Benth. l. c.*—Afr. austr.
serrulata, *E. Mey. ex Benth. l. c.*—Afr. austr.

ANASTRAPHIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 295. *COMPOSITAE*, Benth. & Hook. f. ii. 491.
ilicifolia, *D. Don, l. c.*—Am. trop.
intertexta, *Wright, ex Griseb. Cat. Pl. Cub.* 158.—Cuba.
mantuensis, *Wright, ex Griseb. l. c.*—Cuba.
microcephala, *Griseb. l. c.*—Cuba.

ANASTREPHIA, Decne. in Orb. Dict. i. 432 (1841) = **Anastraphia**, D. Don (Compos.).

ANASTROPHEA, Wedd. in DC. Prod. xvii. 78 (1873) = **Sphaerothylax**, Bisch. (Podostem.).
abyssinica, Wedd. l. c. 79.—Abyss.

ANASTROPHUS, Schlecht. in Bot. Zeit. viii. (1850) 681 = **Paspalum**, Linn. (Gram.).
barbatus, *compressus*, *dissitiflorus*, *nematanthus*, *pectinatus*, *pellitus*, *platycaulis*, *platyculmis*, *pulchellus*, *setifolius*, Schlecht, l. c. 682, 683, omnes Paspali homonymae.

ANASYLLIS, E. Mey. Zwei Pfl. Docum. 16 (1843) = **Loxostylis**, Spreng. (Anacard.).
angustifolia, E. Mey. l. c. = L. alata.
latifolia, E. Mey. l. c. = L. alata.

ANATHALLIS, Barb. Rodrig. Orch. Nov. i. 23 (1877) = **Pleurothallis**, R. Br. (Orchid.).
densiflora, Rodrig. l. c. ii. 73.—Bras.
fasciculata, Rodrig. l. c. i. 23.—Bras.
hebesepala, Rodrig. l. c. ii. 75.—Bras.
heterophylla, Rodrig. l. c. ii. 74.—Bras.
micrantha, Rodrig. l. c. ii. 76.—Bras.
nectarifera, Rodrig. l. c. ii. 74.—Bras.
osmosperma, Rodrig. l. c. ii. 73.—Bras.
parahybunensis, Rodrig. l. c. ii. 76.—Bras.
pulvinata, Rodrig. l. c. ii. 77.—Bras.
racemosa, Rodrig. l. c. i. 24.—Bras.
secunda, Rodrig. l. c. ii. 72.—Bras.

ANATHERIX, Steud. Nom. ed. I. 41 (1821) err. typ. = ANANTHERIX, Nutt. = **Asclepiodora**, A. Gray (Asclep.).

ANATHERUM, Beauv. Agrost. 128. t. 22. f. 10, 11 (1812) = **Andropogon**, Linn. (Gram.).
abortivum, Beauv. Agrost. 169 = Panicum abortivum.
Berterianum, Spreng. ex Roem. & Schult. Syst. ii. Mant. ii. 443 = Imperata exaltata.
bicorne, Beauv. Agrost. 128 = Androp. bicornis.
brasiliense, Spreng. ex Steud. Nom. ed. II. i. 84 = Androp. leucostachyus.
caudatum, Schult. Mant. ii. 445 = Imperata exaltata.
domingense, Roem. & Schult. Syst. ii. 809 = Androp. leucostachyus.
glabrum, Schult. Mant. ii. 446 = Androp. intermedius.
holcoides, Nees, Agrost. Bras. 324 = Saccharum holcoides.
inerme, Griseb. Cat. Pl. Cub. 236 = Androp. spathiflorus.
macrurum, Griseb. Pl. Wright. Cub. ii. 534 = Androp. macrourus.
megapotamicum, Spreng. Syst. Cur. Post. 33 = Elionurus rostratus.
montanum, Roem. & Schult. Syst. Mant. ii. 445 = Androp. montanus.
muricatum, Beauv. Agrost. 150 = Androp. squarrosus.
muticum, Beauv. l. c. = Chloris petraea?
nitidum, Spreng. Syst. i. 290 = Chrysopogon fuscus.
parviflorum, Spreng. l. c. = Chrysopogon violascens.
pedunculosum, Desv. Opusc. 70.—Ind. occ.
portoricense, Spreng. l. c. = Imperata exaltata.
refractum, Beauv. Agrost. 150 = Androp. refractus.
scoparium, Desv. Opusc. 70.—Afric.
setifolium, Spreng. Syst. i. 290 = Androp. setifolium.
spathiflorum, Griseb. Cat. Pl. Cub. 236 = Androp. spathiflorus.
squarrosum, Beauv. Agrost. 128, 150 = Panicum abortivum.
tripsacoides, Spreng. Syst. i. 290 = Elionurus tripsacoides.
virgatum, Desv. Opusc. 71 = Androp. spathiflorus.
virginicum, Spreng. Pugill. ii. 16 = Androp. virginicus.

ANATROPA, Ehrenb. in Linnaea, iv. (1829) 402 = **Tetradiclis**, Stev. (Rutac.).
tenella, Ehrenb. l. c. 404 = T. salsa.

ANAUA, Miq. Fl. Ind. Bat. Suppl. 410 (1860) = **Hemicyclia**, Wight & Arn. (Euphorb.).
sumatrana, Miq. l. c. = H. sumatrana.

ANAUXANOPETALUM, Teijsm. & Binn. in Journ. Bot. Néerl. i. (1861) 368 = **Swintonia**, Griff. (Anacard.).
Schwenkii, Teijsm. & Binn. l. c. = S. Schwenkii.

ANAVINGA, Adans. Fam. ii. 448 (1763); Lam. Illustr. t. 355 = **Casearia**, Jacq. (Samyd.).
comocladifolia, Rafin. Sylva Tellur. 149 = C. comocladifolia.
fuliginosa, Blanco, Fl. Filip. ed. I. 372 = C. fuliginosa.
ilicifolia, Rafin. Sylva Tellur. 149 = C. ilicifolia.
lanceolata, Lam. Encyc. i. 148 = C. tomentosa?
ovata, Lam. l. c. = C. ovata.
parvifolia, Lam. Tabl. ii. 484 = C. parvifolia.
Samyda, Gaertn. f. Fruct. iii. 240. t. 224 = C. Samyda.

ANAXAGORAEA, Mart. Fl. Bras. xiii. i. 39 (1841) = seq.

ANAXAGOREA, St. Hil. in Bull. Soc. Philom. (1825) 91. *ANONACEAE*, Benth. & Hook. f. i. 25.
RHOPALOCARPUS, Teijsm. & Binn. ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 22 (1865).
acuminata, *St. Hil. l. c.*—Bras.
brevipes, *Benth. in Hook. Kew Journ.* v. (1853) 8.—Bras.
crassipetala, *Hemsl. Diagn. Pl. Nov.* 2.—Nicaragua.
fruticosa, *Scheff. in Flora*, lii. (1869) 302.—Malaya?
javanica, *Blume, Fl. Jav. Anon.* 66.—Java.
luzonensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 27.—Ind. or.; Ins. Philipp.
Meyeriana, *Zoll. in Linnaea*, xxix. (1857–58) 323.—Java.
phaeocarpa, *Mart. Fl. Bras.* xiii. i. 40.—Bras.
prinoides, *St. Hil. & A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 211.—Bras.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* i. 382.—Sumatra.
zeylanica, Hook. f. & Thoms. Fl. Ind. i. 144 = luzonensis.

ANAXETON, Gaertn. Fruct. ii. 406 (1791); Cass. in Dict. Sc. Nat. xxvi. (1823) 52; xxxiv. 37 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 314.
ARGYRANTHUS, Neck. Elem. i. 75 (1790).
arborescens, Less. Syn. Comp. 354 = arboreum.
arboreum, *Gaertn. Fruct.* ii. 407.—Afr. austr.
asperum, *DC. Prod.* vi. 268.—Afr. austr.
crispum, Gaertn. Fruct. ii. 407 = Helichrysum crispum.
floridum, Lam. Tabl. iii. 269 = arboreum.
foetidum, Gaertn. Fruct. ii. 406. t. 166 = Helichrysum foetidum.
hirsutum, *Less. Syn. Comp.* 356.—Afr. austr.
nudifolium, Gaertn. Fruct. ii. 407 = Helichrysum nudifolium.
nycthemerum, *Less. Syn. Comp.* 355.—Afr. austr.
recurvum, *DC. Prod.* vi. 368.—Afr. austr.
septentrionalis, Vatke, in Oestr. Bot. Zeitschr. xxvii. (1877) 194 = Gnaphalium stenophyllum.
virgatum, *DC. Prod.* vi. 268.—Afr. austr.

ANAXETON, Schrank, in Denkschr. Acad. Muench. viii. (1824) 146 et 162 = **Helipterum**, DC. (Compos.).
eximium, Schrank, l. c. = Helipterum eximium.
racemosum, Schrank, l. c. = Helichrysum mucronatum.

ANBLATUM, Tourn. ex Adans. Fam. ii. 207 (1763) = **Lathraea**, Linn. (Orobanch.).
tartaricum, Fisch. Hort. Gorenk. ed. II. 19 = Phelipaea salsa.
Tournefortii, G. Don, Gen. Syst. iv. 635 = Lathraea Squamaria.

ANCANA, F. Muell. Fragm. v. 27. t. 35 (1865) = **Unona**, Linn. f. (Anonac.).
stenopetala, F. Muell. l. c. = U. Ancana.

ANCANTHIA, Steud. Nom. ed. II. i. 85 (1840) err. typ. = seq.

ANCATHIA, DC. in Guill. Arch. Bot. ii. (1833) 331 = **Cnicus**, Tourn. (Compos.).
igniaria, DC. in Deless. Ic. Sel. iv. t. 73 = C. igniarius.

ANCHIETEA, A. St. Hil. in Ann. Sc. Nat. Sér. I. ii. (1824) 252; in Mém. Mus. Par. xi. (1824) 464; Pl. Us. t. 19 (1824); Pl. Rem. Bras. 290 (1824). *VIOLARIEAE*, Benth. & Hook. f. i. 117.
exaltata, *Eichl. in Mart. Fl. Bras.* xiii. i. 355.—Bras.
pyrifolia, G. Don, Gen. Syst. i. 340 = salutaris.
pyrifolia, *A. St. Hil. Pl. Rem. Bras.* 290.—Bras.
Roquefeilliana, Spreng. Syst. iv. Cur. Post. 98 = salutaris.

ANCHIETEA:—
salutaris, *A. St. Hil. in Ann. Sc. Nat.* Sér. I. ii. (1824) 252.—Bras.
Selloviana, *Cham. & Schlecht. in Linnaea*, ii. (1827) 148.—Bras.

ANCHIONIUM, Reichb. Consp. 181. err. typ. = **Anchonium**, DC.

ANCHOMANES, Schott, in Oestr. Bot. Wochenbl. iii. (1853) 314. *AROIDEAE*, Benth. & Hook. f. iii. 973.
difformis, Engl. in DC. Monog. Phan. ii. 304 = Hookeri.
dubius, *Schott, Prod. Aroid.* 135.—Afr. occ.
Hookeri, *Schott, in Oestr. Bot. Wochenbl.* iii. (1853) 314.—Afr. trop.

ANCHONIUM, DC. in Mém. Mus. Par. vii. (1821) 242. *CRUCIFERAE*, Benth. & Hook. f. i. 101.
Billardierii, *DC. l. c.*—Syria.
elichrysifolium, Boiss. Fl. Orient. i. 240 = Tournefortii.
Tournefortii, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 386.—Oriens.

ANCHUSA, Linn. Syst. ed. I. (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 855.
BRUNNERA, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 582.
BUGLOSSOIDES, Tausch, ex Reichb. Fl. Germ. Excurs. 344 (1831).
BUGLOSSUM, Gaertn. Fruct. i. 322. t. 67 (1788).
CARYOLOPHA, Fisch. & Trautv. Ind. Sem. Hort. Petrop. iii. 31 (1837).
PENTAGLOTTIS, Tausch, in Flora, xii. (1829) 643.
STOMOTECHIUM, Lehm. Asperif. 395 (1818).
Adami, [Pieri] in Ionios Anthol. v. 182 = sempervirens.
aegyptiaca, DC. Prod. x. 48 = flava.
aegyptiaca, Maly, ex Nym. Consp. iii. 511 = Lycopsis variegata.
affinis, *R. Br. in Salt, Voy. Abyss. App.* 62.—Abyss.
africana, *Burm. f. Prod. Fl. Cap.* 4.—Afr. austr.
Agardhii, *Lehm. Asperif.* 241.—Sibir.
aggregata, *Lehm. l. c.* 219. t. 47.—Sicil.; Oriens.
alpestris, Stev. in Trans. Linn. Soc. xi. (1815) 409. t. 32 = Nonca alpestris.
altissima, *Desf. Cat. Hort. Par.* ed. III. 397.—Hab.?
amoena, J. F. Gmel. Syst. ii. 318 = italica.
amplexicaulis, Sibth. & Sm. Prod. Fl. Graec. i. 116 = hybrida.
angustifolia, Balb. Misc. Alt. 9 = undulata.
angustifolia, DC. Fl. Fr. ed. III. iii. 632 = officinalis.
angustifolia, Georgi, It. i. 200 (= *Stenosolenium saxatile*).
angustifolia, *Linn. Sp. Pl.* 133.—Europ. austr.
angustifolia, Willd. Enum. Hort. Berol. i. 178 = ochroleuca.
angustissima, Bourg. ex Nym. Consp. 510 = undulata.
angustissima, C. Koch, in Linnaea, xxii. (1849) 633 = ochroleuca.
annua, Pall. Ind. Taur. ex A. DC. in DC. Prod. x. 48 = stylosa.
arvalis, Reichb. Fl. Germ. Excurs. 343 = officinalis.
arvensis, Bieb. Fl. Taur. Cauc. i. 123 = Lycopsis arvensis.
arvensis, Tausch, in Flora, vii. (1824) 233 = officinalis.
aspera, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 243 = ochroleuca.
asperrima, Delile, Fl. Aeg. Illustr. 55 = Arnebia hispidissima.
atlantica, *Ball, in Journ. Bot.* xi. (1873) 373.—Marocco.
Aucheri, *A. DC. in DC. Prod.* x. 49.—As. Min.
azurea, Mill. Gard. Dict. ed. VIII. n. 9 = italica.
azurea, Schur, Enum. Pl. Transs. 466 = ochroleuca.
Barrelieri, *Vitm. Summa Pl.* i. 388.—Europ.; As. Min.
Baumgarteni, Nym. Consp. iii. 510 = officinalis.
biceps, *Vest, in Flora*, iv. (1821) 148.—Ital.
bracteolata, Viv. Fl. Lib. Spec. 10. t. 4 = Alkanna tinctoria.
bulbosa, [*Pieri*] *in Ionios Anthol.* v. 182.—Corcyra.
calcarea, *Boiss. Voy. Espagne*, ii. 431. t. 123.—Hispan.
canescens, Muhl. Cat. 19 = Lithospermum canescens.
canescens, Stev. ex Ledeb. Fl. Ross. iii. 118 = ochroleuca.

ANCHUSA :—
Capellii, Moris, Stirp. Sard. Elench. ii. 6 = hybrida.
capensis, *Thunb. Prod. Pl. Cap.* 34.—Afr. austr.
Cesatiana, Fenzl, in Frieder. Reise in N. Gricch. 282 = Pulmonaria mollis.
cespitosa, *Lam. Encyc.* i. 504.—Creta.
ciliata, *Cham. in Linnaea*, iv. (1829) 440.—Bras.
ciliata, Schlecht. ex Ledeb. Fl. Ross. iii. 118 = ochroleuca.
clavata, Viv. ex DC. Prod. x. 52 = undulata.
coronensis, Schur, Enum. Pl. Transs. 466 = Barrelieri.
cretica, Mill. Gard. Diet. ed. VIII. n. 7 = Lycopsis variegata.
crispa, *Moris*, *Stirp. Sard. Elench.* iii. 9.—Sardinia.
crispa, Viviani, Fl. Cors. App. 1 = hybrida.
crispata, Presl, Fl. Sic. i. p. xxxiii = hybrida.
cruciata, Cham. in Linnaea, iv. (1829) 438 = Antiphytum cruciatum.
dasyantha, *Cham. l. c.* 437.—Bras.
deflexa, Lehm. Ind. Sem. Hort. Hamb. (1823) 17 = hispida.
divaricata, Murr. in Nov. Comm. Gotting. (1776) 25 = Nonea lutea.
Dregei, *A. DC. in DC. Prod.* x. 44.—Afr. austr.
dubia, Nocca, Hort. Ticin. t. 3 = Nonea lutea.
echinata, Lam. Encyc. i. 505 = strigosa.
echioides, Bieb. Fl. Taur. Cauc. i. 123 = Macrotomia echioides.
exasperata, Clark, Trav. ed. III. i. 625 = ochroleuca.
flava, Forsk. Fl. Aegypt. Arab. 40 = aegyptiaca.
foliosa, Lehm. ex DC. Prod. x. 53 = Myosotis foliosa.
Gmelini, Ledeb. ex Spreng. Syst. i. 562 = ochroleuca.
granatensis, *Boiss. Voy. Espagne*, ii. 430.—Hispan.
granatensis, Lange, Pugill. 191 = hybrida.
hirta, Michx. ex DC. Prod. x. 53 = Lithospermum hirtum.
hispida, *Forsk. Fl. Aegypt. Arab.* 40.—Oriens; Turkest.; Ind. bor.
hispida, Pall. ex Ledeb. Fl. Ross. iii. 120 = stylosa.
hispidissima, Sieber, ex DC. Prod. x. 94 = Arnebia hispidissima.
hyalotricha, Boiss. Pl. Or. Nov. Dec. ii. 9 = Aucheri.
hybrida, Boiss. ex DC. Prod. x. 42 = granatensis.
hybrida, *Tenore*, *Fl. Nap.* i. 65. t. 11.—Ital.; Oriens.
incana, Ledeb. Fl. Ross. iii. 117 = ochroleuca.
incarnata, Schrad. ex Steud. Nom. ed. I. 42 = officinalis.
intermedia, Schur, Enum. Pl. Transs. 466 = ochroleuca.
italica, *Retz. Obs.* i. 12.—Reg. Mediterr.
Kunthii, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 372 = Antiphytum Walpersii.
lanata, Linn. Syst. ed. X. 914 = Cynoglossum cheirifolium.
lanceolata, Willd. ex Steud. Nom. ed. II. i. 85 = Krynitzkia linifolia.
lateriflora, *Dum. Fl. Belg. Prod.* 41.—Belgia.
latifolia, Lehm. Sem. Hort. Hamb. (1821) 2 = Nonea rosea.
laxiflora, DC. Fl. Fr. iii. 631 = Borago laxiflora.
leiosperma, Bory, Fl. Pélop. t. 7. f. 3 = Nonea obtusifolia.
leptophylla, Koch, Syn. Fl. Germ. ed. II. ii. 573 = officinalis.
leptophylla, Roem. & Schult. Syst. iv. 90 = ochroleuca.
leucantha, Willd. ex DC. Prod. x. 122 = Antiphytum linifolium.
leucojifolia, Lehm. Asperif. 249 = Echinospermum barbatum.
licopsipula, Tardent, Ess. Hist. Nat. Bessar. 57 = Nonea pulla.
limbata, *Boiss. Diagn.* Ser. I. xi. 99.—As. Min.
linearifolia, Hochst. in Lorent, Wanderungen, 334 = ochroleuca.
linearifolia, Urville, in Mém. Soc. Linn. Par. i. (1822) 276 = ochroleuca.
linifolia, Lehm. Asperif. 215 = Krynitzkia linifolia.
littorea, *Moris, in Atti Ottava Riun. Ital. Scienzi*, 566.—Sardin.
longifolia, *Lam. Encyc.* i. 504.—Italia.
Lorenti, A. DC. in DC. Prod. x. 52 = ochroleuca.
lucida, Vittm. Summa Pl. i. 387 = italica.
Luschani, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 86.—Lycia.
lutea, Bieb. Fl. Taur. Cauc. i. 126 = Nonea lutea.
lutea, Cav. Ic. i. 50 = Myosotis versicolor.
Lycopsidis, Bess. ex Link, Enum. Hort. Berol. i. 166 = officinalis.

ANCHUSA :—
Lycopsidis, Retz. f. Pugill. 8 = Nonea lutea.
macrocarpa, *Boiss. & Hohen. Diagn.* Ser. I. iv. 42.—Kurdist.
macrophylla, *Desf. Fl. Atlant.* i. 157.—Marocco.
maculata, Hornem. ex Steud. Nom. ed. II. i. 85 = officinalis.
micrantha, Roem. & Schult. Syst. iv. 98 = aggregata.
microcalyx, *Vis. in Flora*, xii. (1829) 8.—Dalmat.
Milleri, *Willd. Enum. Hort. Berol.* i. 179.—Aegypt.; Arab.
moltkioides, Benth. in Royle, Illustr. Pl. Himal. t. 73. f. 1 = Mertensia moltkioides.
myosotidiflora, *Lehm. Asperif.* 234.—Sibir.; Caucas.
neglecta, *A. DC. in DC. Prod.* x. 49.—Oriens.
nigricans, Brot. Fl. Lusit. i. 298 = Nonea nigricans.
nonneoides, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 35 = undulata.
obliqua, *Vis. Ind. Sem. Hort. Patav.* (1836).—As. Min.
ochroleuca, Baumg. Fl. Transs. i. 120 = officinalis.
ochroleuca, *Bieb. Fl. Taur. Cauc.* i. 125.—Transsylv.; Oriens.
officinalis, Bieb. l. c. 124 = ochroleuca.
officinalis, Gouan, Hort. Monsp. 81; Lam. Encyc. i. 502 = italica.
officinalis, *Linn. Sp. Pl.* 133.—Europ.
officinalis, Thunb. Prod. 34; Fl. Cap. ed. Schult. 162 = riparia.
oppositifolia, H. B. & K. Nov. Gen. et Sp. iii. 91 = Krynitzkia linifolia.
orientalis, Linn. Sp. Pl. 133 = Alkanna orientalis.
orientalis, Reichb. f. Ic. Fl. Germ. xviii. 63 t. 1310 = Lycopsis orientalis.
orientalis, Pall. Reise, iii. 694 = Nonea lutea.
ovata, Lehm. Asperif. 222 = Lycopsis orientalis.
paniculata, Ait. Hort. Kew. i. 177 = italica.
parnassica, Boiss. & Orph. Diagn. Ser. II. iii. 134 = officinalis.
parviflora, Sibth. & Sm. Fl. Graec. ii. 57. t. 167 = aggregata.
parviflora, Willd. Sp. Pl. i. 759. (Sp. delenda; cf. Boiss. Fl. Or. iii. 162).—As. Min.
pauciflora, Roem. & Schult. Syst. iv. 774 = Eritrichium rupestre.
perlata, Lam. Encyc. i. 504; Illustr. n. 1820 = Lycopsis variegata.
petiolata, Hook. Bot. Mag. t. 3858 = Cynoglossum petiolatum.
picta, Bieb. Fl. Taur. Cauc. i. 127 = Nonea picta.
Pinardi, Boiss. Pl. Or. Nov. Dec. ii. 8 = hybrida.
procera, Bess. ex Link, Enum. Hort. Berol. i. 166 = ochroleuca.
pubescens, *Kit. in Rochel, Pl. Bann.* 11.—Bannat.
pulla, Bieb. Fl. Taur. Cauc. i. 125 = Nonea pulla.
pustulata, *Schur, Enum. Pl. Transs.* 465.—Transsylv.
pygmaea, H. B. & K. Nov. Gen. et Sp. iii. 92 = Krynitzkia linifolia.
racemosa, DC. Prod. x. 50 = Mertensia racemosa.
rhizochroa, Viv. Dec. n. 20 = Alkanna tinctoria.
riparia, *DC. Prod.* x. 43.—Afr. austr.
rosea, Bieb. Fl. Taur. Cauc. i. 125 = Nonea rosea.
rupestris, Roem. & Schult. Syst. iv. 101 = Eritrichium rupestre.
ryssosperma, Steud. ex A. Rich. Tent. Fl. Abyss. ii. 88 = Milleri.
sabulosa, Stev. ex Ledeb. Fl. Ross. iii. 118 = ochroleuca.
saxatilis, Pall. Reise, iii. 718 (= *Stenoselenium saxatile*).
Sellowiana, *Cham. in Linnaea*, viii. (1833) 115.—Bras.
sempervirens, *Linn. Sp. Pl.* 134.—Europ.
sericea, Roem. & Schult. Syst. iv. 774 = Eritrichium sericeum.
sikkimensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 168.—Reg. Himal.
spathulata, R. Br. ex Roem. & Schult. Syst. iv. 100 = Myosotis spathulata.
spicata, Lam. Illustr. n. 1810 = angustifolia.
spinocarpos, Forsk. Fl. Aegypt. Arab. 40 = Echinospermum spinocarpum.
Spruneri, Boiss. Diagn. Ser. I. xi. 98 = stylosa.
stoechadifolia, Cham. in Linnaea, iv. (1829) 439 = Antiphytum stoechadifolium.
stricta, *Herbich, in Flora*, xxxviii. (1855) 641.—Europ.
strigosa, *Labill. Syr. Dec.* iii. 7. t. 4.—Syria; Persia.

ANCHUSA :—
strigosa, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 246.—Syria.
stylosa, *Bieb. Fl. Taur. Cauc.* i. 123.—Graccia; As. Min.; Tauria.
subrepanda, *Kit. in Linnaea*, xxxii. (1863) 437.—Europ.
tenella, Hornem. Hort. Hafn. i. 176 = Bothriospermum tenellum.
tenuiflora, Fisch. ex Ledeb. Fl. Ross. iii. 142 = Eritrichium saxatile.
tetraquetra, Cham. in Linnaea, viii. (1833) 113 = Antiphytum tetraquetrum.
thessala, *Boiss. & Sprun. Diagn.* Ser. I. xi. 99.—Thessal.
tinctoria, Linn. Sp. Pl. ed. II. i. 192 = Alkanna tinctoria.
tinctoria, Pall. Ind. Taur. ex Lehm. Asperif. 266 = Nonea pulla.
tinctoria, Sieber, ex Steud Nom. ed. II. i. 86 = Lycopsis orientalis.
tinctoria, Woodv. Med. Bot. t. 92 = officinalis.
tomentosa, Boiss. Diagn. Ser. II. iii. 133 = ochroleuca.
tuberculata, Forsk. Fl. Aegypt. Arab. 40 = Alkanna tinctoria.
tuberosa, *H. B. & K. Nov. Gen. et Sp.* iii. 92.—Amer. austr.
undulata, *Linn. Sp. Pl.* 133.—Reg. Mediterr.
undulata, Pall. Reise, iii. 682 = ochroleuca.
undulata, Sibth. & Sm. Fl. Graec. ii. t. 165 = hybrida.
undulata, Vahl, ex DC. Prod. x. 42 = granatensis.
variegata, Lehm. Asperif. 223 = Lycopsis variegata.
ventricosa, Sibth. & Sm. Fl. Graec. ii. 58. t. 169 = Nonea alba.
ventricosa, Viv. Fl. Lib. Spec. 10. t. 6. f. 1 = Nonea Vivianii.
verrucosa, Lam. Encyc. i. 504; Illustr. 403. n. 1819 = aegyptiaca.
verrucosa, Lejeun. ex Reichb. Fl. Germ. Excurs. 342 = Lycopsis variegata.
versicolor, Schur, Enum. Pl. Transs. 466 = ochroleuca.
versicolor, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 254 = Nonea rosea.
villosa, Roem. & Schult. Syst. iv. 775 = Eritrichium villosum.
villosa, *Spreng. Syst.* i. 560.—Hab.?
virginiana, *Linn. Sp. Pl.* 133 (= Lithospermum hirtum?).—Am. bor.
virginica, Linn. Syst. ed. XII. 145 = praec.
vulgaris, Dum. Fl. Belg. 41 = sempervirens.
zeylanica, Jacq. f. Eclog. i. 47. t. 29 = Bothriospermum tenellum.
zeylanica, Vahl, ex Hornem. Hort. Hafn. i. 176 = Cynoglossum denticulatum.

ANCHUSOPSIS, Bisch. Del. Sem. Hort. Heidelb. (1852) 8 = **Lindelofia**, Lehm. (Borag.).
longiflora, Bisch. l. c. = L. spectabilis.

ANCISTROCARPHUS, A. Gray, in Proc. Am. Acad. vii. (1868) 355 = **Microplus**, Linn. (Compos.).
filagineus, A. Gray, l. c. 356.—Calif.

ANCISTROCARPUS, H. B. & K. Nov. Gen. et Sp. ii. 186. t. 122. (1817) = **Microtea**, Sw. (Phytolacc.).
hexander, J. Gay, ex Moq. in DC. Prod. xiii. II. 17, 18 = M. maypurensis.
maypurensis, H. B. & K. Nov. Gen. et Sp. ii. 186 = M. maypurensis.
maypurensis, Spreng. Neue Entdeck. i. 257 = M. Sprengelii.
Schrankii, Ledeb. Ind. Sem. Hort. Dorp. (1822) = M. maypurensis.

ANCISTROCARPUS, Oliver, in Journ. Linn. Soc. ix. (1867) 173. *TILIACEAE*, Benth. & Hook. f. i. 986.
brevispinosus, *Oliver, l. c.* 174.—Afr. trop.
densispinosus, *Oliver, l. c.*—Afr. trop.

ANCISTROCARYA, Maxim. in Bull. Acad. Petersb. xvii. (1872) 443. *BORAGINEAE*, Benth. & Hook. f. ii. 861.
japonica, *Maxim. l. c.* 444.—Japon.

ANCISTROCLADUS, Wall. Cat. n. 1052 (1832). *DIPTEROCARPEAE*, Benth. & Hook. f. i. 191.
BIGAMEA Koenig, ex Endl. Gen. 1183 (1840).

ANCISTROCLADUS:—
WORMIA, Vahl, in Skrift. Nat. Selsk. Kjobenh. vi. (1810) 104.
attenuatus, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 300.—Burma.
extensus, *Wall. Cat. n.* 1052.—Burma.
Griffithii, *Planch. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 318.—Burma.
guineensis, *Oliver, Fl. Trop. Afr.* i. 175.—Afr. trop.
Heyneanus, *Wall. Cat.* n. 7262.—Ind. or.
pinangianus, *Wall. l. c.* n. 1054.—Malaya.
sagittatus, Wall. l. c. n. 1055 = Tetramerista glabra.
stelligerus, Wall. l. c. n. 1053 = Wallichii.
Vahlii, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 325.—Zeylan.
Wallichii, *Planch. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 319.—Ind. or.

ANCISTRODESMUS, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 302 = **Microlepis**, Miq. (Melast.).
oleaefolius, Naud. l. c. 303 = M. oleaefolia.

ANCISTROLOBUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 352. t. 6 = **Cratoxylon**, Blume (Hyper.).
brevipes, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 383 = C. polyanthum.
floribundus, Turcz. l. c. 382 = C. floribundum.
formosus, Zoll. & Mor. Syst. Verz. Zoll. 25 = C. formosum.
glaucescens, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 383 = C. arborescens.
ligustrinus, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 352 = C. polyanthum.
micradenius, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 382 = C. Blancoi.

ANCISTROPHORA, A. Gray, in Mem. Am. Acad. Ser. II. vi. (1859) 457 = **Verbesina**, Linn. (Compos.).
Wrightii, A. Gray, l. c. = V. Wrightii.

ANCISTROPHYLLUM, G. Mann & H. Wendl. in Kerch. Palm. 230 (1878). *PALMAE*, Benth. & Hook. f. iii. 937.
LACCOSPERMA, G. Mann & H. Wendl. l. c. 249 (1878).
secundiflorum, *G. Mann & H. Wendl. l. c.*—Afr. trop.

ANCISTROSTIGMA, Fenzl, in Ann. Wien. Mus. ii. (1838) 293; Nov. Stirp. Dec. 85 (1838) = **Trianthema**, Linn. (Ficoid.).
cypseleoides, Fenzl, l. c. = T. cypseleoides.

ANCISTRUM, Forst. Char. Gen. 3. t. 2 (1776) = **Acaena**, Linn. (Rosac.).
Acaena, Lag. Gen. et Sp. Nov. 7 = Acaena sericea.
acorna, DC. Prod. ii. 592, sphalm. = praec.
adscendens, Poir. Encyc. Suppl. i. 347 = Acaena adscendens.
anserinaefolium, Forst. Char. Gen. 4 = Acaena Sanguisorbae.
argenteum, Poir. Encyc. Suppl. i. 346 = Acaena argentea.
barbatum, Lam. Illustr. i. 77. t. 22. f. 3 = Margyricarpus setosus.
cylindristachyum, Poir. Encyc. Suppl. i. 347 = Acaena cylindristachya.
decumbens, Gaertn. Fruct. i. 163. t. 32 = Acaena Sanguisorbae.
decumbens, Thunb. Prod. Pl. Cap. 6 = Acaena latebrosa.
diandrum, Forst. f. Prod. 10 = Acaena Sanguisorbae.
elongatum, Poir. Encyc. Suppl. i. 345 = Acaena elongata.
humile, Pers. Syn. i. 30 = Acaena adscendens.
laevigatum, Lag. Gen. et Sp. Nov. 7 = Acaena laevigata.
lappaceum, Poir. Encyc. Suppl. i. 344 = Acaena lappacea.
latebrosum, Ait. Hort. Kew. i. 16 = Acaena latebrosa.
lucidum, Ait. l. c. 15 = Acaena lucida.
magellanicum, Lam. Illustr. i. 76. t. 22. f. 2 = Acaena adscendens, laevigata.
ovalifolium, Sweet, Hort. Brit. ed. l. 148 = Acaena ovalifolia.
pinnatifidum, Poir. Encyc. Suppl. i. 345 = Acaena pinnatifida.
pumilum, Poir. l. c. 347 = Acaena pumila.
repens, Vent. Jard. Cels, 6. t. 6 = Acaena ovalifolia.
Sanguisorbae, Linn. f. Suppl. 89 = Acaena Sanguisorbae.

ANCISTRUM:—
sarmentosum, Thou. Fl. Tristan d'Acugna, 44 = Acaena Sanguisorbae.
trifidum, Poir. Encyc. Suppl. i. 346 = Acaena trifida.

ANCLYANTHUS, A. Juss. in Orb. Dict. xi. 254 (1849) = **Ancylanthos**, Desf.

ANCRUMIA, Harv. ex Baker, in Hook. Ic. Pl. t. 1227 (1877). *LILIACEAE*, Benth. & Hook. f. iii. 806.
cuspidata, *Harv. ex Baker, l. c.*—Chili.

ANCYLANTHOS, Desf. in Mém. Mus. Par. iv. (1818) 5. t. 2. *RUBIACEAE*, Benth. & Hook. f. ii. 112.
Bainesii, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 160.—Afr. trop.
cinerascens, *Welw. ex Hiern, l. c.* 159.—Afr. trop.
cistifolius, *Welw. ex Hiern, l. c.*—Afr. trop.
fulgidus, *Welw. ex Hiern, l. c.*—Afr. trop.
Monteiroi, *Oliver, in Hook. Ic. Pl.* t. 1208.—Afr. trop.
rubiginosa, *Desf. l. c.*—Angola.

ANCYLOCALYX, Tul. in Ann. Sc. Nat. Sér. II. xx. (1843) 136. t. 2 = **Pterocarpus**, Linn. (Legumin.).
acuminata, Tul. l. c. 137 = P. Ancylocalyx.

ANCYLOCLADUS, Wall. Pl. As. Rar. iii. 45 (1832) = **Willughbeia**, Roxb. (Apocyn.).

ANCYLOGYNE, Nees, in Mart. Fl. Bras. ix. 63. t. 7 (1847) = **Sanchezia**, Ruiz & Pav. (Acanth.).
capitata, Nees, in DC. Prod. xi. 222.—Peruv.
longiflora, Hook. f. Bot. Mag. t. 5588.—Ecuador.
macrocnemis, Nees, in Mart. Fl. Bras. ix. 64.—Am. trop.
munita, Nees, l. c.—Am. trop.
peruviana, Nees, in DC. Prod. xi. 222.—Peruv.

ANCYROSSEMON, Poepp. & Endl. Nov. Gen. et Sp. iii. t. 272 (1845) = seq.

ANCYROSTEMMA, Poepp. & Endl. Nov. Gen. et Sp. iii. 65 (1845) = **Sclerothrix**, Presl (Loasac.).
micranthum, Poepp. & Endl. l. c. = S. fasciculata.

ANDA, A. Juss. [in Dict. Sc. Nat. ii. 113 (1816)]; Euph. Tent. 39. t. 12 (1824) = **Joannesia**, Vell. (Euphorb.).
brasiliensis, Raddi, in Mem. Mod. xviii.; Fis. (1820) 404 = J. Princeps.
Gomesii, A. Juss. Euph. Tent. 39. t. 12 = J. Princeps.
Jomesii, Mart. in Regensb. Arch. Gew. iv. (1830) 85 = praec.
Pisonis, Mart. ex Steud. Nom. ed. II. i. 86 = J. Princeps.

ANDACA, Rafin. Fl. Tellur. iii. 82 (1836) = **Lotus**, Tourn. (Legumin.).
arabica, Rafin. l. c. = L. arabicus.

ANDERSONIA, R. Br. Prod. 553 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 617.
ANTHEROCEPHALA, DC. Prod. vii. 755 (1839).
HOMALOSTOMA, Stschegl. in Bull. Soc. Nat. Mosc. (1859) i. 22.
SPHINCTEROSTOMA, Stschegl. l. c. 21.
aristata, *Lindl. Swan Riv. App.* 25.—Austral.
brachyanthera, *F. Muell. Fragm.* vi. 61.—Austral.
brachyota, F. Muell. l. c. iv. 125 = parvifolia.
brevifolia, *Sond. in Lehm. Pl. Preiss.* i. 332.—Austral.
coerulea, *R. Br. Prod.* 554.—Austral.
colossea, *F. Muell. Fragm.* vi. 63.—Austral.
depressa, *R. Br. l. c.* 553.—Austral.
depressa, A. Cunn. ex DC. Prod. vii. 767 = squarrosa.
Fraseri, Sond. in Lehm. Pl. Preiss. i. 331 = sprengelioides.
gracilis, *DC. Prod.* vii. 767.—Austral.
grandiflora, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 21.—Austral.
heterophylla, *Sond. in Lehm. Pl. Preiss.* i. 333.—Austral.
homatostoma, *Benth. Fl. Austral.* iv. 253.—Austral.
involucrata, *Sond. in Lehm. Pl. Preiss.* i. 331.—Austral.

ANDERSONIA:—
lanuginosa, A. Cunn. ex DC. Prod. vii. 767 = depressa.
latiflora, *F. Muell. Fragm.* vi. 61.—Austral.
Lehmanniana, Sond. in Lehm. Pl. Preiss. i. 330 = sprengelioides.
macranthera, F. Muell. Fragm. iv. 124 = heterophylla.
macronema, F. Muell. l. c. viii. 51 = Sprengelia macronema.
micrantha, *R. Br. Prod.* 553.—Austral.
parvifolia, *R. Br. l. c.*—Austral.
patens, Sond. in Lehm. Pl. Preiss. i. 331 = sprengelioides.
patricia, *F. Muell. Fragm.* vi. 79.—Austral.
pauciflora, *Sond. in Lehm. Pl. Preiss.* i. 332.—Austral.
prostrata, Sond. l. c. 333 = depressa.
pubescens, Sond. l. c. 331 = sprengelioides.
setifolia, *Benth. Fl. Austral.* iv. 252.—Austral.
spirophylla, F. Muell. Fragm. vi. 62 = grandiflora.
sprengelioides, *R. Br. Prod.* 553.—Austral.
squarrosa, R. Br. Prod. 553 = depressa.
subsessilis, *Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 67.—N. Caled.
subulata, *Benth. Fl. Austral.* iv. 256.—Austral.
variegata, *Sond. in Lehm. Pl. Preiss.* i. 334.—Austral.

ANDERSONIA, Buch.-Ham. ex Wall. Cat. sub n. 4014 (1832) = **Anogeissus**, Wall. (Combret.).
acuminata, Roxb. ex Wight & Arn. Prod. 316 = Anog. acuminata.
altissima, Roxb. ex Wight & Arn. l. c. = Anog. latifolia.
lanceolata, Buch.-Ham. ex Wall. l. c. = Anog. acuminata.

ANDERSONIA, Roxb. Hort. Beng. [87] (1814); Fl. Ind. ii. 212 (1832) = **Amoora**, Roxb. (Meliac.).
cucullata, Roxb. ll. cc. = Am. cucullata.
Rohituka, Roxb. ll. cc. 87, 213 = Am. Rohituka.

ANDERSONIA, Willd. ex Roem. & Schult. Syst. v. 21 (1819) = **Gaertnera**, Lam. (Logan.).
vaginata, Willd. ex Roem. & Schult. l. c. = G. vaginata.

ANDICUS, Vell. Fl. Flum. 80 (1825) = **Joannesia**, Vell. (Euphorb.).
pentaphyllus, Vell. l. c. = J. Princeps.

ANDIRA, Lam. Encyc. i. 171 (1783). *LEGUMINOSAE*, Benth. & Hook. f. i. 550.
LUMBRICIDIA, Vell. Fl. Flum. 305 (1825).
POLTOLOBIUM, Presl, Bot. Bemerk. 63 (1844).
VOUACAPOUA, Aubl. Pl. Gui. Suppl. 9. t. 373 (1775).
acuminata, Benth. in Ann. Wien. Mus. ii. (1838) 109 = inermis.
Amazonum, *Mart. ex Benth. l. c.* 107.—Bras.
anthelminthica, *Benth. l. c.* 108.—Bras.
Araroba, *Aguiar, Mem. sobre a Araroba*, 31.—*Cf. Pharm. Journ.* Ser. III. x. (1879-80) 44.—Bras.
Aubletii, *Benth. in Ann. Wien. Mus.* ii. (1838) 108.—Guiana.
bracteosa, Benth. l. c. 107 = Amazonum.
cinerascens, Mart. Herb. Fl. Bras. n. 1156 = Tipuana macrocarpa.
cubensis, *Benth. in Journ. Linn. Soc.* iv. (1860) *Suppl.* 121.—Cuba.
cuiabensis, *Benth. in Mart. Fl. Bras.* xv. i. 295.—Bras.
excelsa, *H. B. & K. Nov. Gen. et Sp.* vi. 385.—Mexic.
fraxinifolia, *Benth. in Ann. Wien. Mus.* ii. (1838) 108.—Bras.
frondosa, *Mart. in Walp. Rep.* i. 803.—Bras.
gabonica, *Baill. in Adansonia*, vi. (1866) 219 (adnot.).—Afr. trop.
grandiflora, Guill. & Perr. Tent. Fl. Seneg. i. 254 = inermis.
Harfieldii, Leschen. in Ann. Mus. Par. xvi. (1810) t. 24 = seq.
Horsfieldii, Leschen. l. c. 481 (err. Harsfieldii) = Euchresta Horsfieldii.
humilis, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 109.—Bras.
inermis, *H. B. & K. Nov. Gen. et Sp.* vi. 385.—Ind. occ.; Bras.
laurifolia, *Benth. in Ann. Wien. Mus.* ii. (1838) 109.—Bras.

ANDIRA :—
microcarpa, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 179.—Cuba.
nitida, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 109.—Bras.
oblonga, Benth. in Journ. Linn. Soc. iv. Suppl. (1860) 121 = retusa.
ormosioides, Benth. in Ann. Wien. Mus. ii. (1838) 108 = anthelminthica.
paniculata, *Benth. l. c.* 109.—Bras.
parvifolia, *Mart. ex Benth. l. c.* 108.—Bras.
pauciflora, *Benth. l. c.* 109.—Bras.
Pisonis, Auct. ex Steud. Nom. ed. II. i. 87 = A. Aubletii.
Pisonis, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 108.—Bras.
racemosa, Lam. Encyc. i. 17 = excelsa.
retusa, *H. B. & K. Nov. Gen. et Sp.* vi. 385.—Guiana.
riparia, *H. B. & K. l. c.*—N. Granat.
rosca, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 108.—Bras.
sapindoides, *Benth. in Journ. Linn. Soc.* iv. *Suppl.* (1860) 123.—Ind. occ.
spinulosa, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 108.—Bras.
stipulacea, *Benth. l. c.* 107.—Bras.
vermifuga, *Mart. ex Benth. l. c.* 108.—Bras.

ANDISCUS, Vell. Fl. Flum. ii. t. 86 (1827) = **Andicus**, Vell.

ANDOUINIA, Reichb. Consp. 160 (1828) err. typ. = **Audouinia**, Brongn. (Bruniac.).

ANDRACHNE, Linn. Syst. ed. I. (1735). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 270.
ARACHNE, Neck. Elem. ii. 348 (1790).
ERACLISSA, Forsk. Fl. Aegypt. Arab. 208 (1775).
LEPIDANTHUS, Nutt. in Trans. Am. Phil. Soc. v. (1841) 175.
LEPTOPUS, Decne. in Jacquem. Voy. Bot. 155. t. 156 (1844).
MASCHALANTHUS, Nutt. in Trans. Am. Phil. Soc. v. (1837) 175.
PHYLLANTHIDIA, Didrichs. in Vidensk. Meddel. Kjoeb. (1857) 150.
TELEPHIOIDES, Tourn. ex Moench, Meth. Suppl. 310 (1802).
apetala, Roxb. ex Wall. Cat. n. 7956 = Bischofia javanica.
arborea, *Mill. Gard. Dict.* ed. VIII. n. 3.—Campech.
aspera, *Spreng. Syst.* iii. 884.—Oriens; Afr. trop.; Ind. or.
australis, *Zoll. & Mor. in Natur. en Geneesk. Arch. Neerl. Indie*, ii. (1845) 17.—Java.
Cadishaco, Roxb. ex Wall. Cat. n. 7877 = Cleistanthus collinus.
capensis, *Baill. in Adansonia*, iii. (1862-63) 163.—Afr. austr.
chinensis, *Bunge, Enum. Pl. Chin. Bor.* 59.—China.
colchica, *Fisch. & Mey. ex Boiss. Fl. Orient.* iv. 1137.—Caucas.
cordifolia, *Muell. Arg. in DC. Prod.* xv. II. 234.—Reg. Himal.
Decaisneana, Baill. Etud. Gén. Euphorb. 577 = cordifolia.
Decaisnei, *Benth. Fl. Austral.* vi. 88.—Austral.
Doonkyboisca, Heyne, ex Wall. Cat. n. 7879 = Bridelia retusa.
Dregeana, *Baill. Adansonia*, iii. (1862-63) 164.—Afr. austr.
elliptica, *Roth, Nov. Sp.* 364.—Ind. or.
frutescens, Ehret, in Phil. Trans. lvii. (1768) 114. t. 6 = Arbutus Andrachne.
fruticosa, Decne. ex Muell. Arg. in DC. Prod. xv. II. 235 = Decaisnei.
fruticosa, *Linn. Sp. Pl.* 1014.—China; N. Hisp.
fruticosa, Heyne, ex Hook. f. Fl. Brit. Ind. iv. 287 = Phyllanthus suberosus.
fruticulosa, *Boiss. Diagn.* Ser. I. vii. 86.—Persia.
maroccana, *Ball, in Journ. Bot.* xiii. (1875) 205.—Marocco.
microphylla, *Baill. Etud. Gén. Euph.* 577.—Peruv.
orbicularis, Muell. Arg. in DC. Prod. xv. II. 510, in syn. = seq.
orbiculata, Roth, Nov. Sp. 364 = Cleistanthus collinus.
ovalis, *Muell. Arg. in Linnaea*, xxxii. (1863) 78.—Afr. austr.

ANDRACHNE :—
ovata, Lam. ex Poir. Encyc. Suppl. i. 243, in syn. = Agyneia ovata.
phyllanthoides, Muell. Arg. in DC. Prod. xv. II. 435, in syn. = Roemeriana.
Roemeriana, *Muell. Arg. l. c.* 234.—Amer. bor.
rotundifolia, Eichw. ex C. A. Mey. Casp. 18. t. 20 = telephioides.
telephioides, *Linn. Sp. Pl.* 1014.—Reg. Mediterr.; Oriens; Ind. or.
tenera, *Miq. Fl. Ind. Bat.* i. II. 365.—Malaya.
trifoliata, Roxb. Hort. Beng. 70; Fl. Ind. iii. 728 = Bischofia javanica.

ANDRADEA, Allemão, Diss. Andrad. (1845). *NYCTAGINEAE*, Benth. & Hook. f. iii. 11.
floribunda, *Allemão, l. c.*—Bras.

ANDRASTIS, Rafin. ex Benth. in Ann. Wien. Mus. ii. (1838) 86 = **Virgilia**, Lam. (Legum.).

ANDREOSKIA, Boiss. Fl. Orient. i. 303 (1867) = **Andrzeiowskia**, Reichb. (Crucif.).

ANDREOSKIA, DC. Prod. i. 190 (1824) = **Dontostemon**, Andrz. (Crucif.).
crassifolia, Bunge, ex Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 271.—Mongolia.
dentata, Bunge, Enum. Pl. Chin. Bor. 6 = D. dentatus.
eglandulosa, DC. Prod. i. 190 = D. eglandulosus.
integrifolia, DC. l. c. = D. integrifolius.
parviflora, Turcz. Cat. Pl. Baic. n. 155. sec. Bull. Soc. Nat. Mosc. xv. (1842) 269 = Malcolmia Ledebourii.
pectinata, DC. Prod. i. 190 = D. pectinatus.

ANDREUSIA, Dun. in DC. Prod. vii. 560 (1839) = **Hornemannia**, Vahl (Vaccin.).
guadalupensis, Dun. l. c. = H. martinicensis.

ANDREUSIA, Vent. Jard. Malm. 108. t. 108 (1807) = **Myoporum**, Banks.
debilis, Vent. l. c. sub t. 108 = M. debile.
glabra, Vent. l. c. t. 108 = M. ellipticum.
salicifolia, Hort. ex Steud. Nom. ed. II. i. 87 = Freylinia cestroides.
scabra, Zeyh. ex Steud. l. c. = M. parviflorum.

ANDREWSIA, Spreng. Anleit. ii. I. 474 (1817); Syst. i. 428 (1825) = **Bartonia**, Muhl. (Gentian.).
autumnalis, Spreng. Syst. i. 162 = B. tenella.
paniculata, Bart. ex Steud. Nom. ed. II. i. 87 = B. tenella.
verna, Spreng. Syst. i. 162 = B. verna.

ANDRIALA, Decne. in Orb. Dict. i. 485 (1841) err. typ. = **Andryala**, Linn. (Compos.).

ANDRIAPETALUM, Pohl, Pl. Bras. Ic. i. 113 (1827) = **Panopsis**, Salisb. (Proteac.).
cayennense, Klotzsch, ex Meissn. in Mart. Fl. Bras. v. I. 78.—Bras.
ferrugineum, Meissn. in DC. Prod. xiv. 345.—Peruv.
multiflorum, Schott, ex Endl. Suppl. iv. II. 82.—Bras.
polystachya, Schott, ex Meissn. in DC. Prod. xiv. 345.—N. Granat.
reticulatum, Pohl, Pl. Bras. i. 115. t. 92.—Bras.
rubescens, Benth. ex Meissn. in DC. Prod. xiv. 347 = *Sprucei*.
rubescens, Pohl, Fl. Bras. i. 114. t. 91.—Bras.
sessilifolium, Benth. ex Meissn. in DC. Prod. xiv. 346 = cayennense.
sessilifolium, Klotzsch, in Linnaea, xv. (1841) 53 = P. hameliaefolia.
Sprucei, Meissn. in DC. Prod. xiv. 347.—Bras.
suaveolens, Klotzsch, in Linnaea, xx. (1847) 472.—N. Granat.
Yolombo, Posado, in Bull. Soc. Bot. Fr. xviii. (1871) 374.—N. Granat.

ANDRIEUXIA, DC. Prod. v. 559 (1836) = **Heliopsis**, Pers. (Compos.).
mexicana, DC. l. c. et in Deless. Ic. Sel. iv. t. 31 = H. buphthalmoides.
quitensis, Benth. Pl. Hartw. 206.—Ecuador.

ANDROCENTRUM, Lem. in Fl. des Serres, Sér. I. iii. (1847) 242, verso. *ACANTHACEAE*, Benth. & Hook. f. ii. 1084.
multiflorum, *Lem. l. c.*—Mexico.

ANDROCERA, Nutt. Gen. Am. i. 129 (1818) = **Solanum**, Linn.
lobata, Nutt. l. c. = S. rostratum.

ANDROCHILUS, Liebm. in Bot. Notiser, (1844) 101 = **Liparis**, Rich. (Orchid.).
campestris, Liebm. l. c. 197.—Mexico.

ANDROCOMA, Nees, in Hook. Journ. Bot. ii. (1840) 396 = **Scirpus**, Linn. (Cyperac.).
speciosa, Nees, l. c. 397 = S. giganteus.

ANDROCYMBIUM, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 21. t. 2. *LILIACEAE*, Benth. & Hook. f. iii. 822.
CYMBANTHES, Salisb. in Trans. Hort. Soc. i. (1812) 329.
ERYTHROSTICTUS, Schlecht. in Linnaea, i. (1826) 90.
PLEXINIUM, Rafin. Fl. Tellur. ii. 32 (1836).
Burchellii, *Baker, in Journ. Bot.* xii. (1874) 246.—Afr. austr.
Burkei, *Baker, l. c.*—Afr. austr.
circinatum, *Baker, in Journ. Linn. Soc.* xvii. (1880) 443.—Afr. austr.
cucullatum, *Eckl. Verz.* 7.—Afr. austr.
cuspidatum, *Baker, in Journ. Bot.* xii. (1874) 245.—Afr. austr.
Dregei, *Presl, Bot. Bemerk.* 116.—Afr. austr.
eucomoides, Sweet, Brit. Flow. Gard. t. 165 = leucanthum.
eucomoides, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 21.—Afr. austr.
leucanthum, *Willd. l. c.*—Afr. austr.
littorale, *Eckl. Verz.* 7.—Afr. austr.
longipes, *Baker, in Journ. Bot.* xii. (1874) 246.—Afr. austr.
melanthioides, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 21.—Afr. austr.
palaestinum, *Baker, in Journ. Linn. Soc.* xvii. (1880) 445.—Palaestin.
punctatum, Baker, in Journ. Bot. xii. (1874) 246 = leucanthum.
punctatum, *Baker, in Journ. Linn. Soc.* xvii. (1880) 445.—Afr. bor.
striatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 336.—Abyss.
subulatum, *Baker, in Journ. Bot.* xii. (1874) 245.—Afr. austr.
volutare, *Burch. Trav.* i. 213.—Afr. austr.
volutum, Burch. ex Baker, in Journ. Bot. xii. (1874) 245 = volutare.

ANDROGLOSSA, Benth. in Hook. Kew Journ. iv. (1852) 42 = **Sabia**, Colebr.
reticulatum, Benth. l. c. = S. limoniacea.

ANDROGRAPHIS, Wall. Pl. As. Rar. iii. 77 (1832). *ACANTHACEAE*, Benth. & Hook. f. ii. 1099.
ERIANTHERA, Nees, in Wall. Pl. As. Rar. iii. 77 (1832).
affinis, *Nees, l. c.* iii. 116.—Ind. or.
alata, *Nees, in DC. Prod.* xi. 516.—Ind. or.
Beddomei, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 506.—Ind. or.
ceylanica, Nees, in Hook. Comp. Bot. Mag. ii. (1836) 313 = macrobotrys.
ceylanica, Wight, Ic. t. 1560 = viscosula.
echioides, *Nees, in Wall. Pl. As. Rar.* iii. 117.—Ind. or.
elongata, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 502.—Ind. or.
glandulosa, *Nees, in Wall. Pl. As. Rar.* iii. 115.—Ind. or.
gracilis, Nees, in DC. Prod. xi. 516 = Wightiana.
humifusa, Wall. ex Nees, l. c. 514 = serpyllifolia.
lineata, *Nees, in Wall. Pl. As. Rar.* iii. 116.—Ind. or.
lobelioides, *Wight, Ic.* t. 1557.—Ind. or.
macrobotrys, *Nees, in DC. Prod.* xi. 516.—Ind. or.
Neesiana, Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 504 = stellulata.
Neesiana, *Wight, Ic.* t. 1561.—Ind. or.
orbiculata, Wall. ex Nees, in DC. Prod. xi. 514 = serpyllifolia.
ovata, *Benth. & Hook. f. Gen.* ii. 1100.—Ind. or.
paniculata, *Nees, in Wall. Pl. As. Rar.* iii. 116.—Ind. or.
Rothii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 506.—Ind. or.
serpyllifolia, *Wight, Ic.* t. 517.—Ind. or.

ANDROGRAPHIS:—
stellulata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 504.—Ind. or.
stenophylla, *C. B. Clarke, l. c.* 503.—Ind. or.
subspathulata, *C. B. Clarke, l. c.* 502.—Ind. or.
tenuiflora, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 502.—Ind. or.
viscidula, Steud. Nom. ed. II. i. 87 = viscosula.
viscosula, *Nees, in Wall. Pl. As. Rar.* iii. 116.—Ind. or.
Wightiana, *Arn. ex Nees, in DC. Prod.* xi. 517.—Ind. or.
Wightiana, T. Anders. in Thw. Enum. Pl. Zeyl. 232 = macrobotrys.

ANDROGYNE, Griff. Notul. iii. 279 (1851) = **Panisea**, Lindl. (Orchid.).

ANDROLEPIS, Brongn. ex Houll. in Rev. Hortic. xlii. (1870) 12 = **Aechmea**, Ruiz & Pav. (Bromel.).
Skinneri, Brongn. l. c.—Guatemala.

ANDROMACHIA, Humb. & Bonpl. Pl. Aequin. ii. 104. t. 112 (1809) = **Liabum**, Adans. (Compos.).
acaulis, H. B. & K. Nov. Gen. et Sp. iv. 98 = L. acaule.
alternifolia, Kunze, ex Steud. Nom. ed. II. i. 87 = Aplopappus chrysanthemifolius.
Deppeana, Less. in Linnaea, vi. (1831) 401 = L. Deppeanum.
excelsa, Poepp. & Endl. Nov. Gen. et Sp. iii. 44.—Peruv.
grandiflora, H. B. & K. Nov. Gen. et Sp. iv. 99 = L. grandiflorum.
hieracioides, H. B. & K. l. c. 98 = L. hieracioides.
igniaria, Humb. & Bonpl. Pl. Eq. ii. 104. t. 112 = L. igniarium.
Jussieui, Cass. in Bull. Soc. Philom. (1817) 184 = L. Jussieui.
melastomoides, H. B. & K. Nov. Gen. et Sp. iv. 101. t. 137 = L. melastomoides.
nubigenum, H. B. & K. l. c. 102 = L. nubigenum.
Poiteaui, Cass. in Bull. Soc. Philom. (1817) 184 = L. Brownei.
sessilifolia, H. B. & K. Nov. Gen. et Sp. iv. 102 = L. sessiliflorum.
solidaginea, H. B. & K. l. c. 100 = L. solidagineum.
verbascifolia, H. B. & K. l. c. = L. verbascifolium.
volubilis, H. B. & K. l. c. 102 = L. volubile.

ANDROMEDA, Linn. Syst. ed. I. (1735); Gen. ed. I. 123 (1737), pro parte. *ERICACEAE*, Benth. & Hook. f. ii. 587.
Xolisma, Rafin. in Am. Monthly Mag. (1819) 193.
acuminata, Ait. Hort. Kew. ii. 70 = Leucothoë acuminata.
acuminata, Sm. Exot. Bot. ii. 59 = Leucothoë Catesbaei.
adenothrix, Miq. Ann. Mus. Bot. Lugd. Bat. i. 31 = Gaultheria adenothrix.
alata, Herb. Dombey, ex Dun. in DC. Prod. vii. 562 (= *Proclesia alata*).—Peruv.
ambigua, *Schrad. in Goett. Gel. Anz.* i. (1821) 710.—Hab.?
americana, Hort. ex DC. Prod. vii. 607 = A. Polifolia.
anastomosans, Mutis, ex Linn. f. Suppl. 830 = Gaultheria anastomosans.
angustifolia, Pursh, Fl. Am. Sept. i. 291 = Cassandra calyculata.
arborea, Linn. Sp. Pl. ed. I. 394 = Oxydendrum arboreum.
axillaris, Lam. Encyc. i. 157 = Leucothoë axillaris.
axillaris, Michx. Fl. Bor. Am. i. 253 = Leucothoë Catesbaei.
baccata, Wangenh. Beitr. Am. 30. t. 39. f. 69 = Gaylussacia resinosa.
bracomorensis, H. B. & K. Nov. Gen. et Sp. iii. 288. t. 263 = Agarista bracomorensis.
bracteata, Cav. Ic. vi. 42 = Gaultheria bracteata, tomentosa.
bracteosa, DC. Prod. vii. 607 = Arctostaphylos tomentosa.
bryantha, Linn. Mant. 238 = Bryanthus Gmelini.
buxifolia, Lam. Encyc. i. 159 = Agauria buxifolia.
caerulea, Linn. Sp. Pl. 393 = Phyllodoce taxifolia.
calyculata, Linn. l. c. 394 = Cassandra calyculata.
campanulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 31.—Japon.
canadensis, Cels, ex Lam. Encyc. i. 158 = A. Polifolia.

ANDROMEDA:—
candida, Vell. Fl. Flum. 175; iv. t. 98 = Gaultheria candida.
capricida, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 148 = Pieris ovalifolia.
cassinefolia, Vent. Jard. Cels, 60 = Zenobia speciosa.
Catesbaei, Walt. Fl. Carol. 137 = Leucothoë Catesbaei.
cerea, Linn. f. Suppl. 238 = Vaccinium cereum.
cernua, *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 31.—Japon.
chinensis, Lodd. Bot. Cab. t. 1642 = Vaccinium bracteatum.
chlorantha, Cham. in Linnaea, viii. (1833) 508 = Leucothoë chloranthus.
ciliata, Nees, in Princ. Neuwied, Reise, i. 104 = Leucothoë ciliata.
ciliicalyx, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 30 = Menziezia ciliicalyx.
ciliosa, Boj. ex DC. Prod. vii. 603 = (*Leucothoë Bojeri*).—Madag.
coccinea, Nees, in Princ. Neuwied, Reise, i. 104 = Leucothoë coccinea.
coccinea, Schrad. in Goett. Gel. Anz. i. (1821) 709 = Gaylassacia Pseudo-vaccinium.
columellaris, Fisch. ex Herd. in Act. Hort. Petrop. i. (1871–72) 330 = Cassiope Redowskyi.
coriacea, Ait. Hort. Kew. ed. I. ii. 70 = Pieris nitida.
coriifolia, Thunb. & Billb. Pl. Bras. Dec. 1. n. 5 = Leucothoë coriifolia.
crassifolia, Pohl, Pl. Bras. ii. 34 = Leucothoë crassifolia.
crispa, Hort. Paris. ex Poir. Encyc. Suppl. i. 356 = Cassandra calyculata.
cubensis, *Griseb. Cat. Pl. Cub.* 51.—Cuba.
cupressiformis, Wall. ex D. Don, in Mem. Wern. Soc. iii. (1821) 411 = Cassiope fastigata.
cupressina, Hook. Fl. Bor. Am. ii. 38 = Cassiope Mertensiana.
Daboëcia, Linn. Syst. ed. XII. 300 = Daboëcia polifolia.
Daboëcia, Pall. Fl. Ross. ii. 57 = Phyllodoce taxifolia.
dealbata, Lindl. Bot. Reg. t. 1010 = Zenobia speciosa.
droseroides, Linn. Mant. 239 = Erica glutinosa.
elegans, Meerb. Pl. Sel. Ic. Pict. t. 27 = Leucothoë racemosa.
elliptica, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 126 = Pieris ovalifolia.
empetrifolia, Lam. Encyc. i. 155 = Pernettya empetrifolia.
ericoides, Pall. Fl. Ross. i. II. 56 = Cassiope ericoides.
eriophylla, Vell. in Vandel. Fl. Lusit. et Bras. Sp. ed. Roem. 107 = Gaultheria ferruginea.
eucalyptoides, Cham. & Schlecht. in Linnaea, i. (1826) 518 = Leucothoë eucalyptioides.
fasciculata, Sw. Prod. Veg. Ind. Occ. 73 = Lyonia fasciculata.
fastigiata, Wall. in As. Res. xiii. (1820) 394 = Cassiope fastigiata.
ferruginea, Walt. Fl. Carol. 138 = Lyonia ferruginea.
flexuosa, Moon, Cat. ex Walp. Ann. i. 480 = Gaultheria fragrantissima.
floribunda, Pursh, Fl. Am. Sept. i. 293 = Pieris floribunda.
Fontanesiana, Steud. Nom. ed. II. i. 88 = Leucothoë axillaris.
formosa, Wall. in As. Res. xiii. (1820) 395; Wall. Cat. n. 761 = Pieris formosa.
formosissima, Bart. Trav. N. & S. Car. Georg. ii. (1791) = Leucothoë acuminata.
frondosa, Pursh, Fl. Am. Sept. i. 295 = Lyonia paniculata.
glandulosa, *Wright, in Sauv. Fl. Cub.* 84.—Cuba.
glauca, Hort. ex C. Koch, Dendrol. ii. 1. 118 = Zenobia speciosa.
glaucifolia, Wender. Ind. Sem. Hort. Marb. (1827) in App. n. 2 = A. Polifolia.
glaucophylla, Link, Enum. Hort. Berol. i. 394 = A. Polifolia.
globulifera, Hort. ex DC. Prod. vii. 599 = Lyonia paniculata.
glomerata, Cav. Ic. vi. 42 = Gaultheria anastomosans.
grandiflora, Hort. ex Steud. Nom. ed. II. i. 88 = A. Polifolia.
grandiflora, Herb. Mus. Par. ex Dun. in DC. Prod. vii. 553 = Ceratostemma grandiflorum.
grandiflora, Meerb. Pl. Sel. Ic. Pict. t. 25 = Pieris Mariana.
Hahniana, *Meerb. l. c.* t. 24.—Hab.?
Halmiana, Steud. Nom. ed. I. 44; ed. II. i. 88, sphalm. = Hahniana.

ANDROMEDA:—
hirsuta, Vell. Fl. Flum. 175; iv. t. 99 = Gaultheria ferruginea.
hirta, Willd. Enum. Hort. Berol. Suppl. 23 = Leucothoë hirta.
hispidula, Mart. ex Meissn. in Mart. Fl. Bras. vii. 155 = Leucothoë Martii.
humilis, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 326 = Pernettya pumila.
hypnoides, Linn. Sp. Pl. ed. I. 393 = Cassiope hypnoides.
ilicifolia, Pers. Syn. i. 481 = Leucothoë ilicifolia.
jamaicensis, Sw. Prod. Veg. Ind. Occ. 73 = Lyonia jamaicensis.
japonica, Naves, in Blanco, Fl. Filip. ed. III. 120 = Micromelum tephrocarpum.
japonica, Thunb. Fl. Jap. 181 = Pieris japonica.
kalmiaefolia, *Hoffmgg. Verz. Pfl. Nachtr.* i. 204.—Am. bor.
katagherensis, Hook. Ic. Pl. t. 246 = Gaultheria fragrantissima.
lacustris, *Wright, in Sauv. Fl. Cub.* 85.—Cuba.
lanceolata, Desf. Cat. Hort. Par. ed. III. 136 = Leucothoë Catesbaei.
lanceolata, Wall. in As. Res. xiii. (1820) 390 = Pieris ovalifolia.
lanceolata, Vell. Fl. Flum. 175; iv. t. 97 = Leucothoë multiflora.
latifolia, *Wright, in Sauv. Fl. Cub.* 85.—Cuba.
laurina, Michx. Fl. Bor. Am. i. 253 = Leucothoë acuminata.
laxiflora, Mart. ex Meissn. in Mart. Fl. Bras. vii. 157 = Leucothoë laxiflora.
ledifolia, H. B. &. K. Proleg. p. xvi = Arctostaphylos polifolia.
ligustrina, Muhl. Cat. 44; Ell. Sketch, i. 490 = Lyonia paniculata.
littoralis, Steud. Nom. ed. II. i. 88 = Leucothoë littoralis.
longifolia, Pursh, Fl. Bor. Sept. i. 293 = Leucothoë Catesbaei.
lucida, Jacq. Coll. i. 95 = Leucothoë acuminata.
lucida, Lam. Encyc. i. 157 = Pieris nitida.
lycopodioides, Pall. Fl. Ross. i. II. 55 = Cassiope lycopodioides.
marginata, Duham. Arb. ed. nov. i. 188. t. 40 = Pieris nitida.
mariana, Jacq. Ic. Rar. iii. t. 465 = Pieris nitida.
mariana, Linn. Sp. Pl. 393 = Pieris mariana.
Mertensiana, Bong. Veget. Ins. Sitcha, 152. t. 5 = Cassiope Mertensiana.
mexicana, *Hemsl. in Biol. Centr. Am. Bot.* ii. 282.—Mexico.
montana, Buckl. in Am. Journ. Sc. xlv. (1843) 172 = Pieris floribunda.
montana, Salisb. Prod. 290 = Daboëcia polifolia.
multiflora, Pohl, Pl. Bras. ii. 33. t. 122 = Leucothoë multiflora.
musciformis, Poir. Encyc. Suppl. i. 353 = Bryanthus Gmelini.
myrsinites, *Lam. Encyc.* i. 155.—Reg. Magell.
myrtifolia, Salisb. Prod. 290 = Pieris nitida.
myrtifolia, Willd. ex DC. Prod. vii. 603 = Agarista bracamorensis.
nana, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 47.—Japon.
nereifolia, Cham. & Schlecht. in Linnaea, i. (1826) 522 = Leucothoë nereifolia.
nitida, Bartr. ex Marsh. Arbust. (1785) 8 = Pieris nitida.
nitida, Vell. Fl. Flum. 174; iv. t. 94 = Leucothoë revoluta.
nummularia, Cham. & Schlecht. in Linnaea, i. (1826) 520 = Leucothoë nummularia.
obtusa, *Wright, in Sauv. Fl. Cub.* 84.—Cuba.
octandra, Sw. Prod. Veg. Ind. Occ. 73 = Lyonia octandra.
oleaefolia, *Hoffmgg. Verz. Pfl. Nachtr.* i. 204.—Hab.?
oleifolia, Cham. in Linnaea, viii. (1833) 504 = Leucothoë oleifolia.
oleifolia, Hort. ex Steud. Nom. ed. II. i. 89 = A. Polifolia.
ovalifolia, Wall. in As. Res. xiii. (1820) 391; Cat. n. 763 = Pieris ovalifolia.
ovata, Soland. ex DC. Prod. vii. 598 = Zenobia speciosa.
paniculata, Linn. Sp. Pl. 394 = Leucothoë racemosa.
paniculata, Willd. Sp. Pl. ii. 612 = Lyonia paniculata.

ANDROMEDA:—
parabolica, Duham. Arb. ed. II. i. 191 = Lyonia parabolica.
perulata, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 31 = Enkianthus japonicus.
phillyreaefolia, Hook. Ic. Pl. t. 122 = Pieris phillyreaefolia.
pilulifera, Hort. ex DC. Prod. vii. II. 599 = Lyonia paniculata.
Pistrix, Cham. in Linnaea, viii. (1833) 508 = Leucothoë subrotunda.
Polifolia, *Linn. Sp. Pl.* 393.—Reg. bor. et arct.
polyantha, Steud. Nom. Bot. ed. II. i. 89 = Lyonia paniculata.
populifolia, Lam. Encyc. i. 159 = Leucothoë acuminata.
prostrata, Cav. Ic. vi. 43 = Pernettya microphylla.
pubescens, Poir. Encyc. Suppl. i. 355 = Lyonia rubiginosa.
pulchella, Cham. ex Meissn. in Mart. Fl. Bras. vii. 161 = Leucothoë pulchra.
pulchella, Cham. in Linnaea, viii. (1833) 509 = Leucothoë pulchella.
pulchella, Salisb. Prod. 289 = Pieris mariana.
pulchra, Vell. Fl. Flum. 174; iv. t. 96 = Gaultheria venusta.
pulchra, Cham. & Schlecht. in Linnaea, i. (1826) 521 = Leucothoë pulchra.
pulverulenta, Bartr. Trav. N. & S. Car. Georg. ii. 476 = Zenobia speciosa.
punctata, Domb. ex Dun. in DC. Prod. vii. 564 = Thibaudia punctatifolia.
pyrifolia, Thou. Obs. 14; Pers. Syn. i. 481 = Agauria salicifolia.
racemosa, Linn. Sp. Pl. 394 = Leucothoë racemosa.
racemosa, Walt. Fl. Carol. 138 = Lyonia paniculata.
recurva, Buckl. in Am. Journ. Sc. xlv. (1843) 172 = Leucothoë recurva.
Redowskii, Cham. & Schlecht. in Linnaea, i. (1826) 517 = Cassiope Redowskii.
reticulata, Walt. Fl. Carol. 137 = Leucothoë acuminata.
revoluta, Bouton, ex DC. Prod. vii. II. 603 = Leucothoë Bojeri.
revoluta, Hort. ex Steud. Nom. ed. II. i. 89 = A. Polifolia.
revoluta, Spreng. Neuc Entdeck. ii. 131 = Leucothoë revoluta.
rhomboidalis, Veill. in Duham. Arb. ed. II. i. 192 = Gaultheria ferruginea.
rigida, Humb. Itin. i. 397 = Gaultheria rigida.
rigida, Pursh, Fl. Am. Sept. i. 292 = Lyonia ferruginea.
rosmarinifolia, Gilib. Fl. Lithuan. i. 3 = A. Polifolia.
rubiginosa, Pers. Syn. i. 481 = Lyonia rubiginosa.
rupestris, Forst. f. Prod. n. 195 = Gaultheria rupestris.
salicifolia, Lam. Encyc. i. 159 = Agauria salicifolia.
secunda, Moench, Meth. 457 = A. Polifolia.
Sellowii, Steud. Nom. ed. II. i. 89 = Leucothoë pulchra.
serrata, Sm. ex DC. Prod. vii. II. 609 = Leucothoë acuminata.
serrata, Vell. Fl. Flum. 174; iv. t. 95 = Gaultheria elliptica.
serratifolia, Hort. ex DC. Prod. vii. 602 = Zenobia speciosa.
serrulata, Cham. in Linnaea, viii. (1833) 506 = Leucothoë serrulata.
sinensis, Steud. Nom. ed. II. i. 89 = Vaccinium bracteatum.
speciosa, Michx. Fl. Bor. Am. i. 256 = Zenobia speciosa.
speciosa, Willd. ex Steud. Nom. ed. II. i. 89 = Leucothoë racemosa.
spicata, Wats. Dendrol. i. t. 36 = Leucothoë racemosa.
spinulosa, Pursh, Fl. Am. Sept. i. 293 = Leucothoë Catesbaei.
squamulosa, D. Don, Prod. Fl. Nep. 149 = Pieris ovalifolia.
Stelleriana, Pall. Fl. Ross. i. II. 58 = Cassiope Stelleriana.
subrotunda, Pohl, Pl. Bras. ii. 32. t. 121 = Leucothoë subrotunda.
subsessilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 32.—Japon.
symplocifolia, Wall. Cat. n. 1522 = Vaccinium Leschenaultii.

ANDROMEDA:—
taxifolia, Pall. Fl. Ross. ii. 64. t. 72. f. 2 = Phyllodoce taxifolia.
tetragona, Linn. Sp. Pl. 393 = Cassiope tetragona.
tomentosa, Dum.-Cours. Bot. Cult. ed. II. iii. 495 = Lyonia racemosa.
trinervis, Domb. ex Dun. in DC. Prod. vii. 563 (= *Proclesia bracteata*).—Peruv.
vaccinifolia, Hort. ex C. Koch, Dendrol. ii. I. 116 = Pieris floribunda.
venulosa, DC. Prod. vii. 607 = Arctostaphylos Hookeri.
villosa, Wall. Cat. n. 762 = Pieris villosa.
Walteri, Willd. Enum. Hort. Berol. 453 = Leucothoë Catesbaei.
Watsoniana, Steud. Nom. ed. II. i. 89 = Lyonia parabolica.

ANDROMYCIA, A. Rich. Fl. Cub. Fanerog. ii. 282. t. 89 (1853). *AROIDEAE*, Benth. & Hook. f. iii. 987.
cubensis, *A. Rich. l. c.*—Cuba.

ANDROPHORANTHUS, Karst. in C. Koch & Fint. Wochenschr. (1859) 5 = **Caperonia**, St. Hil. (Euphorb.).
glandulosus, Karst. l. c. 6 = C. palustris.

ANDROPHILAX, Steud. Nom. ed. II. i. 89 (1840) = seq.

ANDROPHYLAX, Wendl. Beobacht. 38 (1798) = **Cocculus**, Linn. (Menisp.).
scandens, Wendl. l. c. = C. carolinus.

ANDROPHTHOË, Scheff. in Flora, liii. (1870) 249, err. typ. = DENDROPHTHOË, Mart. = **Loranthus**, Linn.

ANDROPHYSA, Moq. ex seipso in DC. Prod. xiii. II. 201 (1849) = **Halocharis**, Moq. (Chenop.).

ANDROPOGON, Linn. Sp. Pl. ed. I. 1045 (1753). *GRAMINEAE*, Benth. & Hook. f. iii. 1133.
AGENIUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 447 (1836).
ALLOIANTHENA, Steud. Syn. Pl. Gram. 217 (1855).
ALLOIATHEROS, Rafin. in Ser. Bull. Bot. 221 (1830).
ANADELPHIA, Hack. in Engl. Bot. Jahrb. vi. (1885) 240.
ANATHERUM, Beauv. Agrost. 128. t. 22 (1812).
ARTHROSTACHYS, Desv. Opusc. 74. t. 6 (1831).
CYMBACHNE, Retz. Obs. vi. 36, obs. ad calc. (1791).
CYMBOPOGON, Spreng. Pugill. ii. 14 (1815).
DICHANTHIUM, Willem. in Ust. Ann. Bot. xviii. 11 (1796).
DIECTOMIS, H. B. & K. Nov. Gen. et Sp. i. 193. t. 64 (1815).
DIMEIOSTEMON, Rafin. Neogenyt. 4 (1825).
DIPLASANTHUM, Desv. Opusc. 66 (1831).
DISCHANTHIUM, Kunth, Enum. Pl. i. 498 (1833).
ERIOPODIUM, Hochst. in Flora, xxix. (1846) 115, in obs.
EUKLASTAXON, Steud. in Flora, xxxiii. (1850) I. 229; Syn. Pl. Gram. 412 (1854).
EUPOGON, Desv. Opusc. 67, in obs. (1831).
EXOTHECA, Anderss. in Nov. Act. Upsal. ii. (1856) 253. t. 3.
GRAYA, Arn. ex Steud. Nom. ed. II. i. 705 (1840).
GYMNANDROPOGON, [Munro] Duthie, in Atkins. Gaz. N. W. Ind. 638 (1878 ?), nomen.
GYMNANTHELIA, Anderss. in Schweinf. Beitr. Fl. Aeth. 299 (1867).
HETEROCHLOA, Desv. Opusc. 66. t. 5. f. 3 (1831).
HOMOËATHERUM, Nees, in Hook. & Arn. Bot. Beech. Voy. 239 (1836).
HYPARRHENIA, Anderss. in Nov. Act. Upsal. ii. (1856) 231. nomen.
HYPOGYNIUM, Nees, Agrost. Bras. 364 (1829).
LENORMANDIA, Steud. in Flora, xxxiii. (1850) 229, nomen; Syn. Pl. Gram. 359 (1854).
LEPEOCERCIS, Trin. Fund. Agrost. 203 (1820).
MANDELORNA, Steud. Syn. Pl. Gram. 559 (1854).
OROPOGON, Neck. Elem. iii. 211 (1790).
PITHECURUS, Willd. ex Kunth, Enum. Pl. i. 490 (1833).
SCHIZACHYRIUM, Nees, Agrost. Bras. 331 (1829).
SCHIZOPOGON, Reichb. Consp. 52 (1828).
SPIROTHEROS, Rafin. in Ser. Bull. Bot. i. (1830) 221.

ANDROPOGON:—
VETIVERIA, Thou. ex Virey, in Journ. Phar. Sér. I. xiii. (1827) 499.
abyssinicus, *R. Br. ex Fres. in Mus. Senkenb.* ii. (1837) 146.—Abyss.
aciculatus, Willd. Sp. Pl. iv. 906 = Chrysopogon aciculatus.
aciculatus, Retz. Obs. v. 22 = Chrysopogon aciculatus.
aculeatus, Steud. Nom. ed. II. i. 90 = Chrysopogon aciculatus.
adustus, Trin. in Mém. Acad. Pétersb. Sér. VI. II. (1833) 259 = Elionurus latiflorus.
aesthenos, Steud. Syn. Pl. Gram. 376 = Ischaemum ciliare.
aethiopicus, Rupr. ex Steud. Syn. Pl. Gram. 372 = Gayanus.
afer, Bruce, Trav. v. App. 47 (Polliniae sp. ?).—Afr. bor.
affinis, J. Presl, in Rel. Haenk. i. 343 = Sorghum halepense.
affinis, *R. Br. Prod.* i. 201.—Austral.
affinis, Spreng. Nov. Prov. 5 = annulatus.
Agenium, Steud. Syn. Pl. Gram. 395 = Neesi.
agrostoides, *Speg. in Anal. Soc. Cient. Argent.* xvi. (1883) 136.—Reg. Argent.
agrostoides, Speg. Pl. Nov. Am. Austr. Dec. ii. 27; ex Anal. Soc. Cient. Argent. xvi. (1883) 15 = Chrysopogon avenaceus.
albescens, Fourn. Mex. Pl. Enum. Gram. 56; ex Hemsl. Biol. Centr. Am. iii. 530 = Chrysopogon nutans.
albidus, *Wall. Cat.* n. 8821.—Ind. or.
Alectoridia, Steud. Syn. Pl. Gram. 383 = Arthraxon ciliaris.
Allionii, DC. Fl. Fr. iii. 97 = Heteropogon contortus.
alopecuroides, Linn. Sp. Pl. 1045 = Erianthus saccharoides.
alternans, J. Presl, in Rel. Haenk. i. 342 = Chrysopogon violascens.
altissimus, Colla, in Mem. Acc. Torin. xxxix. (1836) 29. t. 60 = saccharoides.
altissimus, *Hochst. ex A. Br. in Flora*, xxiv. (1841) 277.—Abyss.
amaurus, Buese, in Miq. Pl. Jungh. 360 = Pollinia praemorsa.
ambiguus, Gennari, ex Barbey, Consp. Fl. Sard. 190 = Heteropogon hirtus.
ambiguus, Michx. Fl. Bor. Am. i. 58 = Gymnopogon racemosus.
ambiguus, Steud. Syn. Pl. Gram. 385 = exaltatus.
amethystinus, *Steud. l. c.* 371.—Afr. trop.
amplectens, *Nees, Fl. Afr. Austr.* 104.—Afr. austr.
amplexifolius, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 274 = Arthraxon ciliaris.
ampliflorus, Steud. Syn. Pl. Gram. 388 = A. Nardus.
amplus, J. Presl, in Rel. Haenk. i. 339 = glaucescens.
angustatus, Steud. Syn. i. 370 = apricus.
angustifolius, H. B. & K. Nov. Gen. et Sp. i. 184 = Trachypogon angustifolius.
angustifolius, Parl. Fl. Palerm. i. 269 = pertusus.
angustifolius, Sibth. & Sm. Prod. Fl. Graec. i. 47 = A. Ischaemum.
anias, *Llanos, Fragm.* 29.—Ins. Philipp.
annulatus, F. Muell. Fragm. viii. 123 = sericeus.
annulatus, *Forsk. Fl. Aegypt. Arab.* 173.—Afr. bor.; Ind. or.; Austral.
annulatus, F. Schmidt, Fl. Ins. Capverd. 157 = A. Ischaemum.
annulatus, Schult. Mant. ii. 453 = foveolatus.
annuus, *Hack. in Flora*, lxviii. (1885) 137.—Afr. trop.
anomalus, *Steud. Syn. Pl. Gram.* 397.—Arabia.
anthephoroides, Steud. l. c. 375 = Ischaemum anthephoroides.
anthisterioides, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 463.—Abyss.
anthisterioides, Rupr. in Bull. Acad. Brux. ix. (1842) 245 = Ruprechtii.
appendiculatus, *Nees, Fl. Afr. Austr.* 105.—Afr. austr.
apricus, *Trin. in Bull. Sc. Acad. Petersb.* i. (1836) 70.—Reg. trop.
arctatus, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 20.—Florida.
arcuatus, Steud. Syn. Pl. Gram. 374 = Ischaemum fasciculatum.
arenarius, *Hack. in Flora*, lxviii. (1885) 134.—Uruguay.

ANDROPOGON:—
argenteus, DC. Cat. Hort. Monsp. 77 = saccharoides.
argenteus, Ell. Sketch, 148 = argyreus.
argutus, Steud. Syn. Pl. Gram. 391 = Chrysopogon argutus.
argyreus, *Schult. Mant.* ii. 450.—Am. bor.
Ariani, Edgew. in Journ. Linn. Soc. vi. (1862) 208 = A. Iwarancusa.
aristatum, Poir. Eneyc. Suppl. i. 585 = earicosus.
aristidoides, Steud. Syn. Pl. Gram. 397 = Chrysopogon ciliolatus.
aristulatus, *Hochst. ex Steud. Syn. Pl. Gram.* 397 = Chrysopogon Wightianus.
aristulatus, Steud. Syn. Pl. Gram. 372 (Ischaemi sp.?).—Ins. Borbon.
armillaris, *Willd. ex Steud. Nom.* ed. II. i. 90.—Ind. or.
Arnottianus, Steud. Syn. Pl. Gram. 375 = Ischaemum rugosum.
Arnottii, *Nees, ex Steud. l. c.* 380.—Ind. or.
aromaticus, Roxb. ex Schult. Mant. 442 = muricatus.
aromaticus, Sieber, ex Schult. l. c. 481, 388 = A. Schoenanthus.
arrhenobasis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 460 = Hcteropogon arrhenobasis.
arundinaceus, Berg. Desc. Pl. Cap. 356 = Bergii.
arundinaceus, Scop. Fl. Carn. ed. II. ii. 274 = Sorghum halepense.
arundinaceus, Voigt, Hort. Suburb. Calc. 706 = Anthistiria gigantea.
asper, Kunth, Rev. Gram. i. 166 = Danthonia papillosa.
asperifolius, Hack. in Flora, lxviii. (1885) 140 = Zollingeri.
assimilis, Steud. Syn. Pl. Gram. 397 = Chrysopogon pictus.
asthenos, Steud. l. c. 376 = Ischaemum ciliare.
asthenostachys, Steud. l. c. 381 = Pollinia articulata.
astictus, *Buese, in Miq. Pl. Jungh.* 360.—Sumatra.
attenuatus, *Bertol. in Mem. Acc. Sc. Bolog.* x. (1859) 34; *Misc. Bot.* xx. 10. t. 2.—Guatemala.
Aucheri, Boiss. Diagn. Ser. I. v. 77 = Chrysopogon serrulatus.
aureo-fulvus, Steud. Syn. Pl. Gram. 373 = Pollinia Cnmingii.
aureo-villosus, Steud. Syn. Pl. Gram. 380 = Pollinia aurea.
aureus, Bory, Voy. i. 367. t. 21 = Pollinia aurea.
australis, *Spreng. Syst.* i. 287.—Austral.; Timor.
avenaceus, Berg. Desc. Pl. Cap. 356 = Bergii.
avenaceus, H. B. & K. Nov. Gen. et Sp. i. 189 = Sorghum halepense.
avenaceus, Michx. Fl. Bor. Am. i. 58 = Chrysopogon avenaceus.
barbatus, *Linn. Pl. Jam. Pugill.* 30; *Mant.* ii. 302 = Chloris barbata.
barbigerus, Steud. Syn. Pl. Gram. 376 = Ischaemum aristatum.
barbinodis, Lag. Gen. et Sp. Nov. 3 = saccharoides.
Barteri, *Hack. in Flora*, lxviii. (1885) 124.—Afr. trop.
Bellardi, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 317 = Heteropogon contortus.
bellariensis, *Hack. in Flora*, lxviii. (1885) 123.—Ind. or.
Belvisii, Desv. Opusc. 67 = argyreus.
Benthamianus, Steud. Syn. Pl. Gram. 382 = condensatus.
Bergii, *Roem. & Schult. Syst.* ii. 813.—Afr. austr.
Berteronianus, Steud. & Hochst. ex Steud. Syn. Pl. Gram. 380 = saccharoides.
Besseri, Kunth, Rev. Gram. i. 166 = Sorghum vulgare.
besukiensis, Steud. Syn. Pl. Gram. 367 = Heteropogon contortus.
biaristatus, Steud. l. c. 379 = Pollinia ciliata.
bicolor, Nees, Fl. Afr. Austr. 113 = dichrous.
bicolor, Roxb. Hort. Beng. 21; Fl. Ind. i. 272 = Sorghum vulgare.
bicornis, Benth. Pl. Hartw. 263 = condensatus.
bicornis, *Linn. Sp. Pl.* 1046.—Am. austr.
bicornis, Forsk. Fl. Aegypt. Arab. 173 = A. Schoenanthus.
bifidus, Steud. Syn. Pl. Gram. 373 = Ischaemum ciliare.
bifoveolatus, Steud. l. c. 380 = pertusus.
bilinguis, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 262 = Elionurus bilinguis.

ANDROPOGON:—
binatus, Retz. Obs. v. 21 = Spodiopogon pertnsus.
binatus, Roxb. Hort. Beng. 6; Fl. Ind. i. 255 = tenellus.
bipennatus, *Hack. in Flora*, lxviii. (1885) 142.—Afr. trop.; Madag.
bisquamulatus, Hochst. in Flora, xxvii. (1844) 245 = Gayanus.
Bladhii, Retz. Obs. ii. 27 = annulatus.
Blumii, Steud. Syn. Pl. Gram. 373 = Spodiopogon Blumii.
Boivini, *Steud. l. c.* 431.—Ins. Nossibé.
bombycinus, *R. Br. Prod.* i. 202.—Austral.
Bourgaei, *Hack. in Flora*, lxviii. (1885) 134.—Mexico.
bourouensis, Steud. Syn. Pl. Gram. 382 = Ischaemum digitatum.
brachyatherus, *Hochst. in Flora*, xxvii. (1844) 241.—Abyss.
brachystachyus, *Chapm. Fl. S. Un. St. Suppl.* 668.—Florida.
bracteatus, *Willd. Sp. Pl.* iv. 914.—Am. trop.
breviaristatus, Steud. Syn. Pl. Gram. 396 = Chrysopogon Wightianus.
brevifolius, *Sw. Prod. Veg. Ind. Occ.* 26.—Geront. trop.
brevis, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 268 = lateralis.
Brongniarti, *Steud. Nom.* ed. II. i. 90.—Ins. Molucc.
Brownii, Kunth, Rev. Gram. i. 163 = Ischaemum laxum.
Byronis, Steud. Syn. Pl. Gram. 398 = Spodiopogon Byronis.
Cabanisii, *Hack. in Flora*, lxviii. (1885) 133.—Am. bor.
caerulescens, Kunth, Enum. Pl. i. 504 = Chrysopogon violascens.
caespitosus, A. Rich. Tent. Fl. Abyss. ii. 451 = Elionurus argenteus.
caespitosus, Phil. in Linnaea, xxxiii. (1864–65) 275 = hirtiflorus.
caffrorum, Kunth, Rev. Gram. i. 165 = Sorghum vulgare.
campestris, Kunth, Enum. Pl. i. 489 = tener.
campestris, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 277.—Bras.
camporum, Trin. ex Steud. Syn. i. 378 = campestris.
canaliculatus, *Schum. Beskriv. Guin. Pl.* 52.—Guinea.
candidus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 260 = Elionurus candidus.
canescens, Kunth, Enum. Pl. i. 487 = Trachypogon polymorphus.
canus, *Steud. Nom.* ed. II. i. 90.—Mexico.
capensis, Houtt. Plant.-Syst. xii. t. 93. f. 3 = Chloris petraea.
capillaris, Heyne, ex Steud. Syn. Pl. Gram. 116 = Arundinella mutica.
capillaris, *Kunth, Rev. Gram.* i. 166.—Afr. austr.
capilliflorus, Steud. Syn. Pl. Gram. 397 = Chrysopogon violascens.
caricosus, *Linn. Sp. Pl.* ed. II. 1480.—As. trop.; Ins. Mascar.
carinatus, *Nees, Agrost. Bras.* ii. 330.—Bras.
castratus, Griff. Notul. iii. 89 = Arthraxon rude?
caucasicus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 286 = intermedius.
caudatus, Bieb. Tabl. Prov. Côte Occ. Mer Casp. 58 = Erianthus Ravennae.
ceresiaeformis, *Nees, Fl. Afr. Austr.* 109.—Afr. austr.
ceriferus, Hack. in Mart. Fl. Bras. ii. III. 281 = A. Nardus.
cernuus, Kunth, Enum. Pl. i. 501 = Sorghum halepense.
cernuus, Roxb. Fl. Ind. i. 270 = Sorghum vulgare.
chloridiformis, Gaudich. in Freyc. Voy. Bot. 412 = Dimeria pilosissima.
chlorostachys, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 260 = Elionurus rostratus.
chordatus, Steud. Syn. Pl. Gram. 398 = Spodiopogon Blumii.
chrysatherus, F. Muell. in Linnaea, xxv. (1852) 443 = sericeus.
chrysopus, Coss. in Bull. Soc. Bot. Fr. xxii. (1875) 68 = Aucheri.
chrysostachyus, *Steud. Syn. Pl. Gram.* 377.—Abyss.
ciliaris, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 260 = Elionurus latiflorus.

ANDROPOGON:—
ciliatus, Ell. Sketeh, i. 144 = Chrysopogon avenaceus.
ciliatus, Thunb. Fl. Jap. 40 = Anthistiria barbata.
ciliolatus, Steud. Syn. Pl. Gram. 396 = Chrysopogon serrulatus.
cinctus, Steud. l. c. 398 = Chrysopogon violascens.
circinnatus, Hochst. & Steud. ex Steud. l. e. 387 = laniger.
cirratus, *Hack. in Flora*, lxviii. (1885) 119.—Am. bor.
citratus, Hort. ex DC. Cat. Hort. Monsp. 78 = A. Schoenanthus.
citreus, R. Br. Prod. i. 203 = Elionurus citreus.
citriodorus, Hort. ex Desf. Tabl. ed. II. 15 = A. Schoenanthus.
citrosus, Hort. ex Steud. Nom. ed. II. i. 90 = A. Nardus.
clandestinus, Hale, ex Vasey, Gram. Un. St. 19 = Elliottii.
clandestinus, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Schoenanthus.
coeruleus, Steud. l. e. 395 = Chrysopogon serrulatus.
coesius, Nees, ex Hook. & Arn. Bot. Beech. Voy. 244 = A. Schoenanthus.
cognatus, Steud. Syn. Pl. Gram. 374 = schirensis.
coleotrichus, *Steud. l. c.* 386.—Abyss.
commutatus, Aitch. in Journ. Linn. Soc. xix. (1882) 191 = laniger.
commutatus, *Steud. Syn. Pl. Gram.* 387.—Abyss.
comoročnsis, Steud. l. c. 431 (= Ischaemi sp.).
comosus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 461 = coleotrichus.
comosus, Link, Hort. Berol. i. 239 = annulatus.
compactus, Brot. Fl. Lusit. i. 88 = Sorghum halepense.
condensatus, *H. B. & K. Nov. Gen. et Sp.* i. 188.—Amer. austr.
condensatus, Rupr. in Bull. Acad. Brux. ix. (1842) 244 = Chrysopogon Francavillanus.
condylotrichus, *Hochst. ex Steud. Syn. Pl. Gram.* 377.—Abyss.
confertiflorus, Steud. l. c. 385 = A. Nardus.
confinis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 461.—Abyss.
conjugatus, Roxb. Hort. Beng. 6; Fl. Ind. i. 260 (Polliniae sp.).—Bengal.
connatus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 464 = A. Schoenanthus.
consanguineus, *Kunth, Enum. Pl.* i. 494.—Bras.; Uruguay.
consanguineus, Trin. in Linnaca, x. (1836) 291 = condensatus.
consimilis, Steud. Syn. Pl. Gram. 394 = serratus.
contortus, Linn. Sp. Pl. ii. 1045 = Heteropogon contortus.
controversus, Steud. Syn. Pl. Gram. 391 = Sorghum halepense.
cordatifolius, Steud. l. e. 376 = Ischaemum conjugatum.
cordofanus, Hochst. in Flora, xxvii. (1844) 245 = Gayanus.
Cornucopiae, *Hack. l. c.* lxviii. (1885) 126.—Afr. trop.
corollatus, Steud. Syn. Pl. Gram. 369 = Ischaemum corollatum.
cotulifer, Thunb. Fl. Jap. 41 = Pollinia cotulifera.
courtallumensis, Steud. Syn. Pl. Gram. 377 = Apocopis Wightii.
corassipes, Steud. l. e. 375 = Ischaemum Sieboldii.
crinitus, Thunb. Fl. Jap. 40 = Pogonatherum saccharoideum.
Crupina, Kunth, Rev. Gram. i. 166 = Sorghum halepense.
cryptatherus, Steud. Syn. Pl. Gram. 376 = Ischaemum australe.
cubensis, *Hack. in Flora*, lxviii. (1885) 121.—Cuba.
curtipendulus, Steud. Nom. ed. II. i. 90 = Eutriana curtipendula.
Curtisianus, Steud. Syn. Pl. Gram. 390 = virginicus.
cuspidatus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 456 = Arthraxon cuspidatus.
cymbarius, *Linn. Mant.* ii. 303.—Afr. trop.; Ind. or.
dactyloides, Steud. Syn. Pl. Gram. 381 = Trachypogon polymorphus.
debilis, Kunth, Enum. Pl. i. 488 = brevifolius.
decolorans, H. B. & K. Nov. Gen. et Sp. i. 190 = Sorghum halepense.
decussatus, Steud. Syn. Pl. Gram. 377 = koleostachys.
demissus, *Steud. l. c.* 388.—Ind. or.
densiflorus, Steud. l. c. 386 = A. Schoenanthus.

ANDROPOGON :—

densus, Desv. in Ham. Prod. Fl. Ind. Occ. 8 = condensatus.
depauperatus, *Kunth, Rev. Gram.* i. 166.—Bengal.
depressus, Steud. Syn. Pl. Gram. 373 = Spodiopogon depressus.
Dianae, Steud. l. c. 371 = leucostachyus.
diatherus, Steud. l. c. 378 = Ischaemum latifolium.
dichroanthus, Steud. l. c. 393 = serratus.
dichroos, *Steud. l. c.* 389.—Afr. austr.
digitatus, Hochst. ex Steud. l. c. 382 = Trachypogon dactyloides.
digitatus, *Vahl, ex Trin. Fund.* 186.—Hab.?
diplandrus, *Hack. in Flora*, lxviii. (1885) 123.—Afr. trop.
dissitiflorus, Michx. Fl. Bor. Am. i. 57 = virginicus.
dissitiflorus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 267 = consanguineus.
dissolutus, Steud. Syn. Pl. Gram. 381 = Trachypogon polymorphus.
distachyos, *Linn. Sp. Pl.* 1046.—Reg. Mediterr.
distans, Nees, ex Steud. Syn. Pl. Gram. 387 = A. Nardus.
divaricatus, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. i. 99 = Aristida Humboldtiana.
divaricatus, *Linn. Sp. Pl.* 1045.—Virginia.
diversiflorus, Steud. in Zoll. Syst. Verz. 58 = Pollinia praemorsa.
domingensis, Fourn. Mex. Pl. Enum. Gram. 61; ex Hemsl. Biol. Centr. Am. iii. 530 = leucostachyus.
domingensis, Spreng. ex Steud. Nom. ed. II. i. 91 = Trachypogon polymorphus.
domingensis, Steud. l. c. 45 = leucostachyus.
Donianus, Benth. in Hook. Niger Fl. 570 = Elionurus platypus.
Dregeanus, Nees, Gram. Afr. Austr. 112 = Schimperi.
Drummondii, Steud. Syn. Pl. Gram. 393 = Sorghum vulgare.
dubitatus, Steud. l. c. 394 = Sorghum halepense.
dubius, C. Koch, in Linnaea, xxi. (1848) 443 = Sorghum halepense.
dubius, Kunth, Rev. Gram. i. 166 = Elionurus rostratus.
dulcis, *Burm. f. Fl. Ind.* 219 = Sorghum vulgare.
durus, Steud. Nom. ed. II. i. 91 = Erianthus strictus.
echinatus, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 182.—Ind. or.
echinatus, Heyne, in Steud. Nom. ed. II. i. 91 = Arthraxon lanceolatus.
echinulatus, Steud. Syn. Pl. Gram. 397 = Chrysopogon Gryllus.
elatus, Steud. l. c. 376 = Ischaemum aristatum.
elegans, J. Gay, ex Steud. l. c. 364 = Elionurus elegans.
elegantissimus, Steud. l. c. = Elionurus Royleanus.
Elliottii, *Chapm. Fl. S. Un. St.* 581.—Am. bor.
elongatus, Spreng. Syst. i. 287 = Chrysopogon elongatus.
elythrophyllus, *Steud. Syn. Pl. Gram.* 378.—Mexico.
emersus, Fourn. Mex. Pl. Enum. Gram. 58; ex Hemsl. Biol. Centr. Am. iii. 527 = saccharoides.
erianthoides, *F. Muell. Fragm.* x. 75.—Austral.
Erianthus, Link, Hort. Berol. i. 243 = Erianthus saccharoides.
eriophorus, Schoele, in Flora, xxvii. (1844) 51 = virginicus.
eriophorus, Willd. Sp. Pl. iv. 910 = laniger.
eriostachyus, J. Presl, in Rel. Haenk. i. 339 = ternatus.
erythrocoleos, *Steud. Syn. Pl. Gram.* 391.—Nepal.
Esenbeckii, Steud. l. c. 375 = Ischaemum aristatum.
eucnemis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 275.—Guinea.
eucomus, *Nees, Fl. Afr. Austr.* 104.—Afr. trop. et austr.
exaltatus, *R. Br. Prod.* i. 202.—Austral.
exaratus, *Hack. in Flora*, lxviii. (1885) 135.—Paraguay.
excavatus, Hochst. l. c. xxix. (1846) 116 = A. Schoenanthus.
exilis, *Hochst. l. c.* xxvii. (1844) 142.—Afr. trop.
exsertus, Steud. Syn. Pl. Gram. 366 = brevifolius.
falcatus, Steud. l. c. 369 = Ischaemum pectinatum.
familiaris, *Steud. l. c.* 385.—Afr. trop.
fascicularis, *Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 269.—Ind. or.
fascicularis, Thw. Enum. Pl. Zeyl. 437 = intermedius.
fasciculatus, Linn. Sp. Pl. 1047 = Chloris radiata.
fasciculatus, Steud. Syn. Pl. Gram. 382 = Ischaemum fasciculatum.
fastigiatus, *Sw. Prod. Veg. Ind. Occ.* 26.—Reg. trop.

ANDROPOGON :—

feensis, Fourn. Mex. Pl. Enum. Gram. 62; ex Hemsl. Biol. Centr. Am. iii. 527 = hirtiflorus.
ferrugineus, Hochst. ex Steud. Syn. Pl. Gram. 377 = chrysostachyus.
festucoides, J. Presl, in Rel. Haenk. i. 340 = squarrosus.
filifolius, *Steud. Syn. Pl. Gram.* 374.—Afr. austr.
filiformis, Pers. Syn. i. 103 = caricosus.
filiformis, Roxb. Hort. Beng. 6; Fl. Ind. i. 260 = Dimeria ornithopoda.
filipendulus, *Hochst. in Flora*, xxix. (1846) 115.—Afr. trop.; Zeylan.; Austral.
finitimus, *Hochst, ex A. Rich. Tent. Fl. Abyss.* ii. 465.—Abyss.; Ins. Mascar.
firmandus, Steud. Syn. Pl. Gram. 370 = Pollinia praemorsa.
firmus, Kunth, Enum. Pl. i. 486 = Heteropogon firmus.
flabelliformis, *Roxb. ex Spreng. Pugill.* ii. 9.—Ind. or.
flaccidus, A. Rich. Tent. Fl. Abyss. ii. 452 = brevifolius.
flavescens, *J. Presl, in Rel. Haenk.* i. 339.—Peruv.
flexile, Poir. Encyc. Suppl. i. 583 = scoparius.
flexuosus, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Nardus.
floridus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 265 = brevifolius.
foliatus, Steud. Syn. Pl. Gram. 389 = A. Schoenanthus.
foveolatus, *Delile, Fl. Egypte*, 160. t. 8. f. 2.—Afr. et As. trop.
fragilis, R. Br. Prod. i. 202 = brevifolius.
fragillimus, Hochst. ex Steud. Syn. Pl. Gram. 384 = exilis.
Francavilleanus, Fourn. Pl. Mex. Enum. ii. 56; ex Hemsl. Biol. Centr. Am. iii. 530 = Chrysopogon avenaceus.
fulvibarbis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 287.—Afr. trop.
fulvicomus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 463 = rufus.
fulvispica, Steud. Syn. Pl. Gram. 379 = Pollinia Cumingii.
fulvus, *Spreng. Pugill.* ii. 8.—Bengal.
furcatus, Muhl. in Willd. Sp. Pl. iv. 919 = provincialis.
fuscus, J. Presl, in Rel. Haenk. i. 342 = serratus.
Galeottii, *Fourn. Mex. Pl. Enum. Gram.* 56; *ex Hemsl. Biol. Centr. Am.* iii. 527 = Chrysopogon nutans.
galloinsulanus, *Steud. Syn. Pl. Gram.* 371.—Ins. Mascar.
garipensis, Steud. l. c. 379 = annulatus.
Gayanus, *Kunth, Rev. Gram.* i. 163.—Senegal.
geniculatus, Steud. Syn. Pl. Gram. 376 = Ischaemum aristatum.
Gerardi, Vitm. Summa Pl. vi. 16 = provincialis.
gibbus, Steud. Syn. Pl. Gram. 376 = Ischaemum aristatum.
Gidarba, *Buch.-Ham. in Wall. Cat.* n. 8797.—Ind. or.
giganteus, Fenzl, ex Steud. Syn. Pl. Gram. 386 = A. Iwarancusa.
giganteus, Hochst. in Flora, xxvii. (1844) 242 = A. Schoenanthus.
giganteus, Tenore, Fl. Nap. v. 285 = Heteropogon hirtus.
glaber, Roxb. Hort. Beng. 7; Fl. Ind. i. 271 = intermedius.
glabratus, Steud. Nom. Bot. ed. II. i. 91 = Pollinia glabrata.
glabratus, Steud. Syn. Pl. Gram. 395 = Chrysopogon Gryllus.
glabrescens, Hochst. ex Steud. l. c. 374 = polyatherus.
glabriusculus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 468.—Abyss.
glaucescens, Fourn. Pl. Mex. ii. 60; ex Hemsl. Biol. Centr. Am. iii. 527, partim = Bourgaei.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* i. 186.—Am. austr.
glaucescens, Schlecht. in Linnaea, vi. (1831) 39 = virginicus.
glaucopsis, Ell. ex Steud. Nom. ed. II. i. 91 = macrourus.
glaucopsis, Steud. Syn. Pl. Gram. 397 = montanus.
glaucus, Muhl. Descr. Uber. 278 = macrourus.
glaucus, Retz. Obs. v. 20 = Apluda mutica.
glaucus, Torr. in Ann. Lyc. N. York, i. (1824) 157 = saccharoides.
Glaziovii, *Hack. in Mart. Fl. Bras.* ii. III. 286.—Rio de Janeiro.
Goeringi, Steud. in Flora, xxix. (1846) 22 = A. Nardus.
gracilipes, *Hack. in Flora*, lxviii. (1885) 120.—Paraguay.

ANDROPOGON :—

gracilis, J. Presl, in Rel. Haenk. i. 336 = tener.
gracilis, *Spreng. Syst.* i. 284.—Ind. occ.
grandiflorus, *Hack. in Flora*, lxviii. (1885) 127.—Afr. trop.
grandis, Nees, ex Steud. Syn. Pl. Gram. 399 = A. Nardus.
Grayia, Steud. Nom. ed. II. i. 91 = pseudograya.
Griffithsiae, Steud. Syn. Pl. Gram. 375 = Ischaemum rugosum.
Grisebachii, Steud. l. c. 365 = Elionurus Royleanus.
Gryllus, Linn. Cent. Pl. ii. 33 = Chrysopogon Gryllus.
guianensis, Steud. Syn. Pl. Gram. 382 = Ischaemum guianense.
guineensis, Schum. Beskr. Guin. Pl. 51 = Gayanus.
Haenkei, J. Presl, in Rel. Haenk. i. 340 = intermedius.
halepensis, Brot. Fl. Lusit. i. 89 = Sorghum halepense.
Hallii, *Hack. in Sitzb. Acad. Wien*, lxxxix. (1884) 127.—Am. bor.
hamatulus, Hook. & Arn. Bot. Beech. Voy. 244 = A. Nardus.
hamulatus, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Nardus.
hermaphroditum, Pourr. in Mém. Acad. Toul. iii. (1788) 308 = Stipa aristella.
heteranthus, *Steud. Syn. Pl. Gram.* 370.—Hab.?
heteranthus, Trin. ex Steud. Nom. ed. II. i. 91 = bicornis.
heteroclitus, *Nees, Fl. Afr. Austr.* 115.—Ind. or.
hexastachyus, Steud. Syn. Pl. Gram. 380 = Pollinia argentea.
Hildebrandtii, *Hack. in Flora*, lxviii. (1885) 141.—Madag.
himalayensis, Steud. Syn. Pl. Gram. 377 = Apocopis Royleanus.
hirsutus, H. B. & K. Nov. Gen. et Sp. i. 186 = saccharoides.
hirtiflorus, *Kunth, Rev. Gram.* ii. 569. t. 198.—Am. trop.
hirtifolius, *J. Presl, in Rel. Haenk.* i. 338.—Mexico.
hirtinoides, Regel, in Act. Hort. Petrop. vii. (1880) 657 = Ischaemum hirtinodes.
hirtus, Linn. Sp. Pl. 1046 = Heteropogon hirtus.
hispidissimus, Steud. Syn. Pl. Gram. 367 = Heteropogon contortus.
hispidus, Willd. Sp. Pl. iv. 908 = Arundinella brasiliensis.
Hochstetteri, Steud. Syn. Pl. Gram. 384 = fastigiatus.
Hohenackeri, Hochst. ex Steud. l. c. 368 = Heteropogon contortus.
holcoides, Kunth, Rev. Gram. ii. 49. t. 160 = Saccharum holcoides.
Humboldtianus, Steud. Syn. Pl. Gram. 392 = Chrysopogon avenaceus.
humilis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 458.—Abyss.
humilis, Wright, ex Steud. Syn. Gram. i. 388 = demissus.
hypogynus, *Hack. in Mart. Fl. Bras.* ii. III. 290.—Bras.
imberbis, *Hack. in Flora*, lxviii. (1885) 119.—Paraguay.
imberbis, Steud. Syn. Pl. Gram. 375 = Ischaemum aristatum.
importunus, *Steud. l. c.* 382.—China.
incertus, Steud. l. c. 375 = Ischaemum ciliare.
incompletus, J. Presl, in Rel. Haenk. i. 342 = Chrysopogon nutans.
increscens, Steud. Syn. Pl. Gram. 396 = Chrysopogon serrulatus.
incurvatus, Koen. ex Trin. in Spreng. Neue Entdeck. ii. 91 = caricosus.
incurvatus, *Retz. Obs.* v. 21.—Ind. or.
inermis, Steud. Syn. Pl. Gram. 390 = spathiflorus.
infirmus, Steud. l. c. 369 = Ischaemum fragile.
insculptus, Anderss. in Schweinf. Beitr. Fl. Aethiop. 310 = Ischaemum laxum.
insculptus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 458 = pertusus.
insularis, Linn. Pl. Jam. Pugill. 30; Syst. ed. X. 1304 = Panicum leucophaeum.
intermedius, *R. Br. Prod.* i. 202.—Oriens; Afr. et As. trop.; Austral.
intonsus, Nees, Fl. Afr. Austr. 114 = lepidus.
intortum, Crantz, Inst. i. 398 = Sporobolus indicus.
intrans, *F. Muell. Census*, 132.—Austral.
inundatus, F. Muell. in Linnaea, xxv. (1852) 44 = intermedius.
involucratus, Koen. ex Steud. Nom. ed. II. i. 91 = Apluda aristata.
involutus, Steud. Syn. Pl. Gram. 373 = Spodiopogon angustifolius.

ANDROPOGON :—
Ischaemum, *Linn. Sp. Pl.* 1047.—Eur.; As. et Afr. trop. et temp.
Ischaemum, Thunb. Fl. Cap. i. 108 = appendiculatus.
ischyranthus, Steud. Syn. Pl. Gram. 367 = triticeus.
Iwarancusa, Boiss. Fl. Orient. v. 466 = A. Schoenanthus.
Iwarancusa, Nees, Fl. Afr. Austr. 117 = A. Nardus.
Iwarancusa, *Roxb. in Phil. Trans.* lxxx. (1790) 284.—Ind. or.; Afr. trop.
Jamesii, Torr. in Marc. Rep. (1852) 302 = saccharoides.
javanicus, *Miq. Fl. Ind. Bat.* iii. 448.—Java.
javanicus, Steud. Syn. Pl. Gram. 396 = Chrysopogon aciculatus.
jubatus, Balansa, in Bull. Soc. Bot. Fr. xix. (1872) 322 = sericeus.
junceus, *Rich. ex Steud. Nom.* ed. II. i. 91.—Ins. S. Domingo.
juncifolius, *Desv. in Ham. Prod. Fl. Ind. Occ.* 9.—Ins. S. Cruz.
Koenigii, Raeusch. Nom. ed. III. 293 = caricosus.
koleostachys, *Steud. Syn. Pl. Gram.* 376.—Ins. Mascar.
koretrostachys, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 273 = Pollinia articulata.
lachnatherus, Benth. Fl. Austral. vii. 534 = filipendulus.
laguriformis, Griseb. in Goett. Abh. xxiv. (1879) 309 = saccharoides.
laguroides, DC. Cat. Hort. Monsp. 78 = saccharoides.
lanatus, R. Br. Prod. i. 202 = exaltatus.
lanceolatus, Baker, Fl. Maur. and Seych. 414 = Arthraxon cuspidatus.
lanceolatus, Roxb. Hort. Beng. 6; Fl. Ind. i. 262 = Arthraxon lanceolatus.
lancifolius, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 271 = Arthraxon microphyllus.
laniger, *Desf. Fl. Atlant.* ii. 379.—Oriens; Ind. or.; Afr. bor.
laniger, Munro, in Journ. Linn. Soc. xix. (1882) 409 = Spodiopogon laniger.
lanuginosus, H. B. & K. Nov. Gen. et Sp. i. 187 = leucostachyus.
lasiocoleus, Steud. Syn. Pl. Gram. 383 = Arthraxon ciliaris.
lateralis, *Nees, Agrost. Bras.* ii. 329.—Am. trop.
latiflorus, Steud. Syn. Pl. Gram. 364 = Elionurus latiflorus.
latifolius, Kunth, Rev. Gram. i. 166 = Centotheca lappacea.
latifolius, Spreng. Syst. i. 286 = Ischaemum latifolium.
laxiflorus, Steud. Syn. Pl. Gram. 380 = Trachypogon polymorphus.
laxus, Roxb. Hort. Beng. 7; Fl. Ind. ed. Carey, i. 271 = Sorghum halepense.
laxus, Willd. Sp. Pl. iv. 907 = serratus.
Lechleri, Steud. ex Hack. in Mart. Fl. Bras. ii. III. 297 = condensatus.
lepidus, *Nees, Fl. Afr. Austr.* 113.—Abyss.
leptanthus, Steud. Syn. Pl. Gram. 391 = intermedius.
leptocladus, *Hack. in Flora*, lxviii. (1885) 122.—Paraguay.
leptocomus, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 264.—Sierra Leone.
leptophyllus, Trin. l. c. = tener.
leptos, Steud. Syn. Pl. Gram. 397 = tener.
leptostachyus, Benth. in Hook. Niger Fl. 571 = semiberbis.
Leschenaultianus, Decne. Herb. Timor. Descr. 29 = Pollinia Cumingii.
leucopogon, Nees, in Linnaea, xix. (1847) 694 = saccharoides.
leucostachyus, *H. B. & K. Nov. Gen. et Sp.* i. 187.—Am. trop.
Lhotskyi, Steud. Syn. Pl. Gram. i. 384 = condensatus.
lianatherus, Steud. l. c. 367 = Heteropogon insignis.
Liebmanni, *Hack. in Flora*, lxviii. (1885) 132.—Mexico.
ligularis, Kunth, Enum. Pl. i. 497 = Trachypogon polymorphus.
Lindeni, Steud. Syn. Pl. Gram. 389 = apricus.
lineatus, Steud. l. c. 369 = Ischaemum laxum.
lithophilus, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 277.—Bras.
lividus, *Thw. Enum. Pl. Zeyl.* 367.—Ind. or.
lodicularis, Steud. Syn. Pl. Gram. 375 = Ischaemum aristatum.
lolioides, Fourn. Mex. Pl. Enum. Gram. 62; ex Hemsl. Biol. Centr. Am. iii. 528 = scoparius.
longiberbis, *Hack. in Flora*, lxviii. (1885) 131.—Florida.

ANDROPOGON :—
longipes, *Hack. l. c.* 138.—Ind. or.
Louisianae, Steud. Syn. Pl. Gram. 383 = virginicus.
macer, Steud. l. c. 371.—'Mont. cerul.'
Macraei, Steud. l. c. 377 = Ischaemum ciliare.
macroglossus, Steud. l. c. 368 = Trachypogon polymorphus.
macrolepis, *Hack. in Flora*, lxviii. (1885) 125.—Afr. trop.
macrostachyus, Anderss. in Schweinf. Beitr. Fl. Aethiop. 310 = Ischaemum laxum.
macrothrix, Fourn. Mex. Pl. Enum. Gram. 60; ex Hemsl. Biol. Centr. Am. iii. 528 = Liebmanni.
macrothrix, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 270 = ternatus.
macrourus, *Michx. Fl. Bor. Am.* i. 56.—Am. bor.; Mexico.
madagascariensis, Hack. in Flora, lxviii. (1885) 136 = trichozygus.
malacophyllus, Hochst. ex Steud. Syn. Pl. Gram. 372 = Ischaemum ciliare.
malacostachyus, Fourn. Pl. Mex. Pl. Enum. Gram. 62; ex Hemsl. Biol. Centr. Am. iii. 528 = hirtiflorus.
malacostachyus, *J. Presl, in Rel. Haenk.* i. 337.—Am. centr.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 232.—Fernando Po.
marginatus, Steud. in Flora, xii. (1829) 472 = A. Nardus.
Mariannae, Steud. Syn. Pl. Gram. 382 = Ischaemum digitatum.
maritimus, Chapm. Fl. S. Un. St. Suppl. 668 = scoparius.
Martini, Roxb. Hort. Beng. 7; Fl. Ind. i. 280 = A. Schoenanthus.
Martini, Thw. Enum. Pl. Zeyl. 367 = A. Nardus.
medius, Steud. Syn. Pl. Gram. 382 = Ischaemum intermedium.
megapotamicus, Spreng. Syst. iv. Cur. Post. 33 = Trachypogon polymorphus.
melanocarpus, Ell. Sketch, i. 146 = Heteropogon acuminatus.
messanensis, Biv. ex Parl. Fl. Ital. i. 148 = Heteropogon contortus.
Meyenianus, Steud. Syn. Pl. Gram. 375 = Ischaemum aristatum.
micans, Steud. l. c. 382 = Arthraxon cuspidatus.
micranthus Kunth, Rev. Gram. i. 165 = Chrysopogon violascens.
microphyllus Trin. in Mém Acad. Pétersb. Sér. VI. ii. (1833) 275 = Arthraxon microphyllus.
microstachyus, *Desv. in Ham. Prod. Fl. Ind. Occ.* 8.—Ind. occ.
miliaceus, *Roxb. Hort. Beng.* 7; *Fl. Ind.* i. 276.—Ind. or.
miliformis, Schult. Mant. ii. 448 = praec.
Minarum, Kunth, Enum. Pl. i. 507 = Chrysopogon stipoides.
minor, Steud. Syn. Pl. Gram. 377 = Ischaemum Urvilleanum.
minutiflorus, Steud. l. c. 379 = Pollinia argentea.
mollicomus, Kunth, Rev. Gram. i. 365. t. 96 = caricosus.
mollis, Kunth, l. c. ii. 561. t. 195 = Trachypogon polymorphus.
monandrus, Roxb. Hort. Beng. 82; Fl. Ind. i. 264 = Pogonatherum saccharoideum.
monatherus, *A. Rich. Tent. Fl. Abyss.* ii. 462.—Abyss.
monostachys, Spreng. Pugill. ii. 9 = foveolatus.
monostachyus, Steud. Syn. Pl. Gram. 369 = Ischaemum aristatum.
montanus, Koen. ex Trin. in Spreng. Neue Entdeck. ii. 93 = Chrysopogon montanus.
montanus, Nees, in Nov. Act. Nat. Cur. xix. Suppl. I. (1843) 188 = micranthus.
montanus, *Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 271 = Chrysopogon pictus.
monticolus, Schult. Mant. iii. 665 = Chrysopogon montanus.
Montufari, H. B. & K. Nov. Gen. et Sp. i. 184 = Trachypogon polymorphus.
Muehlenbergianus, Schult. Mant. ii. 455 = argyreus.
multicaulis, Steud. Syn. Pl. Gram. 383 = Arthraxon microphyllus.
multinervis, Hochst. ex Steud. l. c. 374 = polyatherus.
muricatus, Retz. Obs. iii. 43 = squarrosus.
murinus, Steud. Syn. Pl. Gram. 377 = Ischaemum murinum.
muticus, Linn. Sp. Pl. ed. II. 1482 = Chloris petraea?
muticus, Steud. Syn. Pl. Gram. 374 = Ischaemum muticum.

ANDROPOGON :—
Myosurus, Hack. in Mart. Fl. Bras. ii. III. 300 = hirtiflorus.
Myosurus, *J. Presl, in Rel. Haenk.* i. 337.—Mexico.
nardoides, Nees, Fl. Afr. Austr. 116 = A. Schoenanthus.
Nardus, *Linn. Sp. Pl.* 1046.—Reg. trop.
Nardus, Trin. in Spreng. Neue Entdeck. ii. 90 = A. Schoenanthus.
Neesi, *Kunth, Enum. Pl.* i. 491.—Bras.
Neesii, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 263 = tener.
nemophyllus, Steud. Syn. Pl. Gram. 377 = Ischaemum filiforme.
nervatus, Hochst. in Flora, xxvii. (1844) 243 = A. Schoenanthus.
nervosus, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 218 = Ischaemum laxum.
Newtonii, *Hack. in Bol. Soc. Broter.* iii. (1885) 137.—Angola.
niger, Kunth, Rev. Gram. i. 164 = Sorghum vulgare.
nigritanus, Benth. in Hook. Niger Fl. 573 = squarrosus.
nitens, Schreb. ex Steud. Nom. ed. II. i. 92 = serratus.
nitidus, Kunth, Rev. Gram. i. 166 = serratus.
nodulibarbis, Hochst. ex Steud. Syn. Pl. Gram. 396 = Chrysopogon Arnottianus.
nodulosus, *Hack. in Flora*, lxviii. (1885) 116.—Afr. trop.
notopogon, Steud. Syn. Pl. Gram. 373 = Spodiopogon angustifolius.
nudus, Steud. l. c. 383 = Arthraxon ciliaris.
Nuttallii, Chapm. Fl. S. Un. St. 580 = Elionurus tripsacoides.
nutans, *Linn. Mant.* ii. 303 = Anthistiria ciliata.
nutans, Linn. Sp. Pl. 1045 = Chrysopogon nutans.
nutans, Muhl. Descr. Uber. 284 = avenaceus.
obliquiberbis, *Hack. in Flora*, lxviii. (1885) 117.—N. Caledon.
obtusifolius, Poir. Encyc. Suppl. i. 583 = brevifolius.
obtusus, Nees, in Hook. & Arn. Bot. Beech. Voy. 243 = annulatus.
obvallatus, Steud. Syn. Pl. Gram. 373 = Spodiopogon angustifolius.
odoratus, Auct. ex Steud. Nom. ed. II. i. 92 = squarrosus.
oliganthus, *Hochst. ex Steud. Syn. Pl. Gram.* 368.—Ind. or.
oligostachyus, Chapm. Fl. S. Un. St. 581 = hirtiflorus.
Olivieri, Boiss. Diagn. Ser. I. v. 76 = laniger.
orthos, Schult. Mant. ii. 455 = foveolatus.
pachnodes, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 284 = A. Schoenanthus.
paleaceus, Steud. Syn. Pl. Gram. 376 = Apocopis Royleanus.
pallidus, Kunth, Rev. Gram. i. 165 = Chrysopogon Gryllus.
paniceus, Steud. Syn. Pl. Gram. i. 375 = Ischaemum intermedium.
paniculatus, Kunth, Enum. Pl. i. 494 = condensatus.
paniculatus, Lam. Fl. Fr. iii. 633 = Chrysopogon Gryllus.
panormitanus, Parl. Fl. Ital. i. 140 = pertusus.
papillipes, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 460.—Abyss.
papillosus, *Hochst. ex A. Rich. l. c.* 457.—Afr. trop.
parviflorus, Roxb. Hort. Beng. 82; Fl. Ind. i. 277 = brevifolius.
parvispica, Steud. Syn. Pl. Gram. 397 = Chrysopogon violascens.
patentivillosus, Steud. l. c. 373 = Ischaemum ciliare.
patulus, Moench, Meth. 180 = A. Ischaemum.
pauciflorus, Salzm. ex Steud. Syn. Pl. Gram. 371 = virginicus.
pectinatus, *Steud. l. c.* 369 = Ischaemum pectinatum.
pedicellatus, Steud. l. c. 394 = serratus.
peduncularis, *Kunth, Rev. Gram.* i. 166.—Ind. or.
peduncularis, Baker, Fl. Maur. 444 = A. Alopecurus.
pendulus, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Nardus.
peninsulae, Steud. l. c. 396 = Chrysopogon Arnottianus.
pennatus, Willd. ex Steud. Nom. ed. II. i. 92 = lanuginosus.
perforatus, Trin. ex Fourn. Mex. Pl. Enum. Gram. 59, ex Hemsl. Biol. Centr. Am. iii. 529 = saccharoides.

ANDROPOGON :—
perfossus, Nees & Meyen, ex Steud. Nom. ed. II. i. 92 = intermedius.
pertusus, *Willd. Sp. Pl.* iv. 922.—Sicil.; As. et Afr. trop.; Austral.
petiolaris, Steud. Syn. Pl. Gram. 398 = Pollinia Lehmanni.
Petitianus, Balansa, in Bull. Soc. Bot. Fr. xix. (1872) 322 = obliquiberbis.
Petitianus, A. Rich. Tent. Fl. Abyss. ii. 451 = exilis.
pilifer, Steud. Syn. Pl. Gram. 373 = pilosus.
pilosus, *Klein, ex Willd. Sp. Pl.* iv. 920.—Ind. or.
pilosus, Sieber, ex Kunth, Enum. Pl. i. 514 = Ischaemum latifolium.
piptatherus, *Hack. in Mart. Fl. Bras.* ii. III. 293.—Bras.
plagiopus, Hochst. ex Steud. Syn. Pl. Gram. 374 = polyatherus.
platypus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 261 = Elionurus platypus.
plumosus, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 918 = Trachypogon polymorphus.
podotrichus, Hochst. ex Steud. Syn. Pl. Gram. 384 = Heteropogon hirtus.
poecilotrichus, *Hack. in Bol. Soc. Broter.* iii. (1885) 138.—Afr. trop.
pogonanthus, Boiss. & Bal. in Bull. Soc. Bot. Fr. v. (1858) 168 = Spodiopogon pogonanthus.
Pohlianus, *Hack. in Mart. Fl. Bras.* ii. III. 304.—Bras.
polyatherus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 455.—Abyss.
polydactylon, Linn. Amoen. Acad. v. 412; Sp. Pl. ed. II. 1483 = Chloris polydactyla.
polymorphus, Steud. Syn. Pl. Gram. 375 = Ischaemum muticum.
polyneuros, Steud. l. c. 385 = A. Schoenanthus.
polyptychos, *Steud. l. c.* 380.—Zeylan.
polystachyus, Roxb. Hort. Beng. 82; Fl. Ind. i. 265 = Heteropogon contortus.
polystichus, Hochst. ex Steud. Syn. Pl. Gram. 369 = Heteropogon acuminatus.
Preslii, Kunth, Enum. Pl. i. 489 = tener.
Princeps, A. Rich. Tent. Fl. Abyss. ii. 470 = Thelepogon elegans.
prionodes, Steud. Syn. Pl. Gram. 383 = Arthraxon lanceolatus.
procerus, F. Muell. Fragm. viii. 124 = filipendulus.
procerus, *R. Br. Prod.* i. 202.—Austral.
productus, Regel, in Bull. Acad. Pétersb. x. (1866) 378 = Pollinia tenuis.
propinquus, Kunth, Enum. Pl. i. 502 = Sorghum halepense.
prostratus, Linn. Mant. ii. 304 = Anthistiria prostrata.
provincialis, Retz. Obs. iii. 43 = Chloris radiata.
provincialis, *Lam. Encyc.* i. 376.—Europ.; Am. bor.
proximus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 464 = A. Iwarancusa.
pruinosus, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Schoenanthus.
pseudograya, *Steud. l. c.* 365.—Zeylan.; China; Madag.
pseudohirtus, Steud. in Flora, xii. (1829) 471 = A. Nardus.
pseudoischaemum, *Nees, ex Steud. Syn. Pl. Gram.* 380.—Ind. or.
pubescens, [Dryand. in] Ait. Hort. Kew, ed. I. iii. 423 = Chloris ciliata.
pubescens, Visiani, in Flora, xii. (1829) I. Erg. 3 = Heteropogon hirtus.
pubescens, Willd. ex Kunth, Enum. Pl. i. 498 = mollicomis.
pubiflorus, Fourn. Mex. Fl. Enum. Gram. 57; ex Hemsl. Biol. Centr. Am. iii. 529 = hirtifolius.
pulchellus, *D. Don, in Hook. Niger Fl.* 571.—Afr. trop.
pumilus, *Roxb. Hort. Beng.* 7; *Fl. Ind.* i. 277.—Ind. or.
punctatus, Roxb. ll. cc. 82, 268 = intermedius.
purpurascens, Muhl. ex Willd. Sp. Pl. iv. 913 = scoparius.
purpureo-sericeus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 469.—Abyss.
pusillus, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 233.—Afr. trop.
quadrivalvis, Linn. Syst. ed. XIII. 758 = Anthistiria ciliata.

ANDROPOGON :—
Quartinianus, *A. Rich. Tent. Fl. Abyss.* ii. 469.—Abyss.
quinqueplumis, Hochst. ex A. Rich. l. c. 450 = Chrysopogon serrulatus.
radicans, Lehm. Ind. Hort. Hamb. (1828); ex Kunth, Enum. Pl. i. 499 = A. Ischaemum.
racemosus, Poir. ex Steud. Nom. ed. I. 412, in syn. = ramosus.
ramosus, Blanco, Fl. Filip. ed. I. 37 = Ischaemum ciliare.
ramosus, *Forsk. Fl. Aegypt. Arab.* 173.—Aegypt.; Arab.
Ravennae, Linn. Sp. Pl. ed. II. 1481 = Erianthus Ravennae.
reconditus, Steud. Syn. Pl. Gram. 386 = Gayanus.
rectirhachis, *Fourn. Mex. Pl. Enum. Gram.* 61; *ex Hemsl. Biol. Centr. Am.* iii. 529.—Mexico.
rectus, Steud. Syn. Pl. Gram. 389 = A. Nardus.
refractus, *R. Br. Prod.* i. 202.—Ins. Molucc. et Pacific.; Austral.
relictus, Steud. Syn. Pl. Gram. 377 = Ischaemum muticum.
repens, Steud. l. c. 374 = Ischaemum muticum.
Retzii, Steud. l. c. 375 = Ischaemum ciliare.
rhizophorus, Steud. l. c. 381 = Spodiopogon albidus.
Riedelii, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 263.—Am. austr.
Robertianus, Steud. Syn. Pl. Gram. 369 = Ischaemum laxum.
rostratus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 261 = Elionurus rostratus.
rottboellioides, Steud. Syn. Pl. Gram. 382 = Rottboellia ophiuroides.
Roxburghianus, Schult. Mant. ii. 451 = Dimeria ornithopoda.
Roxburghii, Nees, ex Steud. Syn. Pl. Gram. 395 = A. Nardus.
Royleanus, Steud. l. c. 397 = Chrysopogon Gryllus.
rubens, Kunth, Rev. Gram. i. 164 = Sorghum vulgare.
rudis, Steud. Syn. Pl. Gram. 383 = Arthraxon rudis.
rufidulus, Steud. l. c. 392 = Chrysopogon avenaceus.
rufispica, *Steud. l. c.* 379.—Java.
rufus, *Kunth, Enum. Pl.* i. 492.—Afr. trop.; Bras.
rufus, Rich. ex Kunth, l. c. 473 = Saccharum cayennense.
rugosus, Steud. Syn. Pl. Gram. 376 = Ischaemum rugosum.
Ruprechtii, *Hack. in Flora*, lxviii. (1885) 126.—Afr. trop.; Mexico.
saccharatus, Kunth, Enum. Pl. i. 502 = Sorghum vulgare.
saccharatus, Roxb. Hort. Beng. 7; Fl. Ind. i. 274 = Sorghum vulgare.
saccharoides, *Sw. Prod. Veg. Ind. Occ.* 26.—Am. temp. et trop.
sanguinarius, *Schreb. Beschr. Graes.* ii. 48.—Afr. bor.
scaberrimus, Kunth, Enum. Pl. i. 504 = Chrysopogon nutans.
scabriflorus, *Rupr. ex Hack. in Mart. Fl. Bras.* ii. III. 299.—Bras.; Paraguay.
scandens, Roxb. Hort. Beng. 7; Fl. Ind. i. 262 = annulatus.
Schaffneri, Griseb. ex Fourn. Mex. Pl. Enum. Gram. 57; ex Hemsl. Biol. Centr. Am. iii. 529 = hirtifolius.
schangulensis, Rupr. ex Steud. Syn. Pl. Gram. 369 = Ischaemum laxum.
Schimperi, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 466.—Abyss.
schirensis, *Hochst. l. c.* 456.—Abyss.
Schlumbergerii, *Fourn. Mex. Pl. Enum. Gram.* 59, *ex Hemsl. Biol. Centr. Am.* iii. 529.—Mexico.
Schoenanthus, Benth. Fl. Austral. vii. 584 = A. Nardus.
Schoenanthus, *Linn. Sp. Pl.* 1046.—As. et Afr. trop. et subtrop.
Schoenanthus, Thunb. Fl. Cap. i. 108; Wall. Pl. As. Rar. t. 280 = A. Nardus.
Schottii, *Rupr. ex Hack. in Mart. Fl. Bras.* ii. III. 299.—Am. trop.
Schweinfurthii, *Hack. in Flora*, lxviii. (1885) 118.—Afr. trop.
scoparius, *Michx. Fl. Bor. Am.* i. 57.—Am. bor.; Mexico.
scoparius, J. Presl, in Rel. Haenk. i. 338 = condensatus.
scrobiculatus, Kunth, Enum. Pl. i. 507 = Heteropogon acuminatus.

ANDROPOGON :—
secundus, Kunth, Enum. Pl. i. 487 = Trachypogon polymorphus.
secundus, Willd. ex Nees, Agrost. Bras. 364 = Heteropogon contortus.
seguenensis, Steud. Syn. Pl. Gram. 367 = Heteropogon acuminatus.
segetum, Steud. l. c. 376 = Ischaemum rugosum.
Sehima, Steud. l. c. 369 = Ischaemum laxum.
semiberbis, *Kunth, Enum. Pl.* i. 489.—Afr. et Am. trop.
semisagittatus, Steud. Syn. Pl. Gram. 376 = Ischaemum semisagittatum.
sennarensis, Hochst. in Flora, xxvii. (1844) 243 = A. Iwarancusa.
sericeus, *R. Br. Prod.* i. 201.—Malaya; Austral.
serratus, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 290 = Chrysopogon violascens.
serratus, Retz. Obs. v. 21 = caricosus.
serratus, *Thunb. Fl. Jap.* 41.—As. et Afr. trop.; Austral.
serrulatus, Link, Enum. Hort. Berol. i. 241 = Arthraxon lanceolatus.
setifolius, *Kunth, Rev. Gram.* i. 166.—Afr. austr.
setosus, Griseb. Cat. Pl. Cub. 235 = Chrysopogon avenaceus.
sibiricus, Steud. Syn. Pl. Gram. 398 = Spodiopogon sibiricus.
simillimus, Steud. l. c. 373 = Ischaemum ciliare.
simplex, *Schum. Beskr. Guin. Pl.* 49.—Afr. trop.
sinaicus, Del. Ind. Sem. Hort. Monsp. (1836); et in Ann. Sc. Nat. Sér. II. vii. (1837) 285 = Heteropogon hirtus.
Smithianus, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 232.—Afr. trop.
Sorghum, Brot. Fl. Lusit. i. 88 = Sorghum vulgare.
spathaceus, Trin. Fund. Agrost. 186 = macrourus.
spathiflorus, *Kunth, Enum. Pl.* i. 496.—Bras.
speciosus, Hort. Goett. ex Steud. Nom. ed. II. i. 93 = saccharoides.
speciosus, Steud. Syn. Pl. Gram. 375 = Rottboellia speciosa.
spicatus, Steud. Nom. ed. II. i. 93 = Trachypogon polymorphus.
Sprengelii, Kunth, Rev. Gram. i. 166 = Chrysopogon montanus.
squamulatus, Hochst. in Flora, xxvii. (1844) 244 = Gayanus.
squarrosus, Koen. ex Kunth, Enum. Pl. i. 126 = Panicum abortivum et paludosum.
squarrosus, *Linn. f. Suppl.* 433.—Reg. trop.
stenophyllus, Roem. & Schult. Syst. ii. 819 = Trachypogon angustifolius.
stipaeformis, Steud. Syn. Pl. Gram. 377 = Dimeria ornithopoda.
stipoides, *Cham. & Schlecht. in Linnaea*, vi. (1831) 39 = Chrysopogon Francavillanus.
stipoides, H. B. & K. Nov. Gen. et Sp. i. 189 = Chrysopogon avenaceus.
stipoides, Kunth, Enum. Pl. i. 487 = Trachypogon polymorphus.
stipoides, Steud. Syn. Pl. Gram. 368 = Trachypogon polymorphus.
stipoides, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 288 = Chrysopogon avenaceus.
striatus, Klein, ex Willd. Sp. Pl. iv. 903 = Ischaemum laxum.
strictus, Host, Gram. ii. 2. t. 2 = Erianthus Hostii.
strictus, Roxb. Hort. Beng. 82; Fl. Ind. i. 265 = foveolatus.
subglabrescens, Steud. Syn. Pl. Gram. 393 = Sorghum vulgare.
submembranaceus, *Steud. l. c.* 431.—Ins. Mascar.
submuticus, Steud. l. c. 382 = Arthraxon ciliaris.
subrepens, Steud. l. c. 397 = montanus.
subtilis, Steud. l. c. 396 = Chrysopogon subtilis.
subulatus, J. Presl, in Rel. Haenk. i. 341 = Chrysopogon aciculatus.
subunifoveolatus, Steud. Syn. Pl. Gram. 380 = pertusus.
superciliatus, *Hack. in Engl. Bot. Jahrb.* vi. (1885) 239.—Austral.; Timor.
Tabina, Steud. in Lechl. Berb. Am. Austr. 56 = hirtiflorus.
tacazensis, Steud. Syn. Pl. Gram. 369 = Ischaemum laxum.
tahitensis, Hook. & Arn. Bot. Beech. Voy. 72 = refractus.

ANDROPOGON:—
Tamba, Hochst. ex Steud. Syn. Pl. Gram. 385 = lepidus.
tectorum, *Schum. & Thonn. Beskr. Guin. Pl.* 49.—Guinea.
tenellus, J. Presl, in Rel. Haenk. i. 335 = brevifolius.
tenellus, *Roxb. Hort. Beng.* 6; *Fl. Ind.* i. 259.—Bengal.
tener, *Kunth, Rev. Gram.* ii. 565. t. 197.—Am. bor. et austr.
tenuiculus, Steud. Syn. Pl. Gram. 371 = sericeus.
tenuifolius, Steud. l. c. 365 = Elionurus argenteus.
tenuipedicellatus, Steud. l. c. 393 = australis?
tenuirachis, Fourn. Mex. Pl. Enum. Gram. 58; ex Hemsl. Biol. Centr. Am. iii. 529 = saccharoides.
tenuis, R. Br. Prod. i. 201 = Heteropogon contortus.
ternarius, Bertol. Bot. Misc. xi. 14 = provincialis.
ternarius, Desv. Opusc. 68 = squarrosus.
ternarius, Michx. Fl. Bor. Am. i. 57 = provincialis?
ternatus, *Nees, Agrost. Bras.* ii. 326.—Am. austr.
tetrastachyus, Ell. Sketch, i. 150 = virginicus.
Thunbergii, Kunth, Rev. Gram. i. 166 = Danthonia livida.
thymiodorus, Steud. Syn. Pl. Gram. 365 = Elionurus thymiodorus.
timorensis, Steud. l. c. 376 = Spodiopogon Blumii.
tomentellus, Steud. l. c. 371 = Gayanus?
Tong-Dong, Steud. l. c. 375 = Ischaemum rugosum.
Torreyanus, Steud. Nom. ed. II. i. 93 = saccharoides.
tortus, Willd. ex Steud. Nom. ed. II. i. 92 = Trachypogon polymorphus.
trachypus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 280 = bracteatus.
trichozygus, *Baker, in Journ. Linn. Soc.* xx. (1884) 300.—Madag.
tridentatus, Hochst. in Flora, xxvii. (1844) 246 = Gayanus.
tridentatus, Roxb. Hort. Beng. 82; Fl. Ind. i. 261 = Apocopis tridentata.
Trinii, Steud. Syn. Pl. Gram. 395 = Chrysopogon serrulatus.
tripsacoides, Trin. ex Steud. Nom. ed. II. i. 92; et Syn. Pl. Gram. 364 = Elionurus tripsacoides.
trispicatus, Schult. Mant. ii. 452 = Pollinia argentea.
tristachyos, Roxb. Hort. Beng. 6; Fl. Ind. i. 261 = Pollinia argentea.
tristachyus, H. B. & K. Nov. Gen. et Sp. i. 187 = glaucescens.
triticeus, R. Br. Prod. i. 201 = Heteropogon insignis.
triticiformis, Steud. Syn. Pl. Gram. 376 = Ischaemum triticeum.
tropicus, Kunth, Enum. Pl. i. 503 = australis.
tropicus, Spreng. Syst. i. 287 = serratus.
truchmenorum, Walp. Ann. iii. 801 = Sorghum vulgare.
truncatus, Steud. l. c. 368 = Trachypogon polymorphus.
tumbackianus, *Roxb. ex Kunth, Enum. Pl.* i. 507, *in syn.*—Afr. austr.
tumbuckianus, Steud. Nom. ed. II. i. 93 = praec.
umbrosus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 467 = lepidus.
undatus, Jacq. Coll. iii. 237 = Oplismenus Jacquini.
undulatus, Pers. Syn. i. 105 = annulatus.
urceolatus, *Hack. in Flora*, lxviii. (1885) 115.—Afr. centr.
Urvilleanus, Steud. Syn. Pl. Gram. 376 = Ischaemum Urvilleanum.
usorum, Steud. l. c. 392 = Sorghum vulgare.
Vachellii, Nees, ex Hook. & Arn. Bot. Beech. Voy. 243 = intermedius.
vaginatus, Ell. Sketch, 148 = virginicus.
vaginatus, J. Presl, in Rel. Haenk. i. 336 = semiberbis.
velatus, Kunth, Enum. Pl. i. 488 = semiberbis.
venustus, *Thw. Enum. Pl. Zeyl.* 367.—Zeylan.
versicolor, Nees, ex Steud. Syn. Pl. Gram. 388 = A. Schoenanthus.
verticillatus, Boj. Hort. Maurit. i. 57 = squarrosus.
verticillatus, Roxb. Hort. Beng. 82; Fl. Ind. i. 267 = Chrysopogon verticillatus.
verticillatus, Schum. Beskr. Guin. Pl. 50 = fulvibarbis.
verticilliflorus, Steud. Syn. Pl. Gram. 393 = Sorghum halepense?
vestitus, *Steud. l. c.* 378.—Bras.
villiferus, Steud. l. c. 376 = Ischaemum australe.
villosulus, Steud. l. c. 397 = Chrysopogon violascens.
villosus, Gilib. Exercit. ii. 522 = A. Ischaemum.

ANDROPOGON:—
villosus, Lam. Fl. Fr. iii. 634 = A. Ischaemum et provincialis.
villosus, Thunb. Prod. Pl. Cap. 20 = Pollinia villosa.
vimineus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 268 = Pollinia imberbis.
violaceus, Heyne, ex Steud. Syn. Pl. Gram. 414 = Arthraxon ciliaris.
violascens, Nees, ex Steud. l. c. 396 = Chrysopogon violascens.
virens, Spreng. Syst. i. 287 = Arundinella brasiliensis.
virgatus, Desv. in Ham. Prod. Ind. Occ. 9 = spathiflorus.
virginicum, *Linn. Sp. Pl.* 1046.—Am. bor.
virginicus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 269 = leucostachyus.
Wightianus, Steud. Syn. Pl. Gram. 395 = Chrysopogon Wightianus.
Wrightii, *Hack. in Flora*, lxviii. (1885) 139.—Am. bor.
xanthoblepharis, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 281.—Afr. trop.
Wrightii, *Munro, ex Wright, in Sauv. Fl. Cub.* 201.—Cuba.
Zeae, *Roem. & Schult. Syst.* ii. 818.—N. Granat.
zeylanicus, Arn. ex Benth. Fl. Hongk. 423 = A. pseudograya.
zeylanica, Steud. Syn. Pl. Gram. 397 = Chrysopogon Arnottianus.
Zollingeri, *Steud. l. c.* 369.—Java.

ANDROSACE, [Tourn.] Linn. Syst. ed. I. (1735); Gen. ed. I. 40 (1737). *PRIMULACEAE*, Benth. & Hook. f. ii. 632.
AMADEA, Adans. Fam. ii. 230 (1763).
ARETIA, Linn. Sp. Pl. ed. I. 141 (1753).
acaulis, Hort. ex Duby, in DC. Prod. viii. 52 = septentrionalis.
acaulis, *Otto, ex Duby, l. c.* 55.—Sibiria.
acutifolia, Turcz. ex Steud. Nom. ed. II. i. 93 = A. Chamaejasme.
affinis, Biroli, in Brugnat. Giorn. Fis. II. iii. (1820) 143. t. 6 = obtusifolia.
Aizoon, *Duby, in DC. Prod.* viii. 50.—Reg. Himal.; China.
alba, *Duby, l. c.* 55.—Am. bor.
albana, *Stev. in Trans. Linn. Soc.* xi. (1815) 410. t. 34.—As. Min.; Caucas.
alismoides, Hornem. Hort. Hafn. i. 185 = lactiflora.
alpina, *Lam. Fl. Fr.* iii. 642.—Europ.
altaica, *C. Koch, in Linnaea*, xxiii. (1850) 612.—Reg. altaic.
apennina, Huet, ex Nym. Consp. 607 = alpina.
aquatica, [Clairv.] Man. Herb. 58 = Hottonia palustris.
arachnoidea, Schott, Nyman & Kotschy, Anal. Bot. 17 = villosa.
arctica, *Cham. in Linnaea*, i. (1826) 220.—Am. arct.
Aretia, D. Dietr. Syn. i. 632 = bryoides.
Aretia, Lap. Hist. Abr. Pl. Pyr. 91 = imbricata.
Aretia, *Vill. Hist. Pl. Dauph.* ii. 469.—Europ.
aretioides, Heer, ex Duby, in DC. Prod. viii. 51 = obtusifolia.
aretioides ×, *Kerner, in Oestr. Bot. Zeitschr.* xxv. (1875) 159.—Europ.
argentea, Lap. Hist. Abr. Pl. Pyr. 92 = imbricata.
arizonica, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 221.—Am. bor.
armeniaca, *Duby, in DC. Prod.* viii. 52.—As. Min.
articulata, *Schur, Enum. Pl. Transs.* 551.—Europ.
baikalensis, Fisch. ex Ledeb. Fl. Ross. iii. 19 = lactiflora.
baikalensis, Schlecht. ex Roem. & Schult. Syst. iv. 786 = ochotensis.
bidentata, C. Koch, in Linnaea, xxiii. (1850) 610 = multiscapa.
Bocconi, Hort. ex Hornem. Hort. Hafn. i. 184 = elongata.
brevifolia, Vill. Hist. Pl. Dauph. ii. 480 = septentrionalis.
brigantiaca, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 105 = carnea.
Brueggeri ×, *Jaegg. ex Bruegg. in Jahresb. Naturf. Ges. Graub. Neue Fol.* xxii. (1879) p. xxix.—Helvet.
bryoides, *DC. Fl. Fr.* iii. 440.—Helvet.
caespitosa, *Lehm. ex Spreng. in Isis*, (1817) 1288. n 162. t. 9.—Persia.
cana, *Willd. ex Steud. Nom.* ed. II. i. 93.—Am. austr.
capitata, Willd. ex Roem. & Schult. Syst. iv. 786 = A. Chamaejasme.

ANDROSACE:—
carinata, Torr. in Ann. Lyc. Nat. Hist. New York, i. (1824) 30 = A. Chamaejasme.
carnea, *Linn. Sp. Pl.* 142.—Europ.
carnosula, Duby, in DC. Prod. viii. 54 = saxifragaefolia.
Chaixi, *Gren. & Godr. Fl. Fr.* ii. 458.—Europ.; Dahuria?
Chamaejasme, Bieb. Fl. Taur. Cauc. iii. 132 = villosa.
Chamaejasme, *Willd. Sp. Pl.* i. 799.—Europ.; As. Min.; As. bor.
Chamaejasme, Wulf. in Jacq. Coll. i. 194 = obtusifolia.
Charpentieri, *Heer, in Moritzi, Graub. Verz.* 113.—Helvet.
ciliata, *DC. Fl. Fr.* iii. 441.—Europ.
commutata, *Schlecht. in Bot. Zeit.* xiv. (1856) 521.—As. temp.
cordifolia, *Wall. in As. Res.* xiii. (1820) 371.—Reg. Himal.
coronopifolia, Andr. Bot. Repos. t. 647 = lactiflora.
Croftii, *Watt, in Journ. Linn. Soc.* xx. (1884) 17.—Reg. Himal.
cylindrica, *DC. Fl. Fr.* iii. 439.—Mont. Pyren.
dasyphylla, Bunge, in Ledeb. Fl. Alt. i. 218 = villosa.
Diapensia, Vill. Hist. Pl. Dauph. ii. 472 = bryoides.
diapensioides, Lap. Hist. Abr. Pl. Pyr. 93 = pyrenaica.
elegans, Duby, in DC. Prod. viii. 55 = rotundifolia.
elongata, *Linn. Sp. Pl.* ed. II. *App.* 1668.—Europ.; Caucas.
elongata, Richards. App. Frankl. 1st Voy. 733 = septentrionalis.
elongata, Pall. ex Ledeb. Fl. Ross. iii. 21 = filiformis.
Escheri ×, *Bruegg. in Jahresb. Naturf. Ges. Graub. Neue Fol.* xxiii-xxiv. (1880) 100.—Helvet.
farinosa, Spreng. Pugill. ii. 37 = Primula farinosa.
fasciculata, Willd. ex Roem. & Schult. Syst. iv. 786 = septentrionalis.
filiformis, *Retz. Obs.* ii. 10.—Persia; As. bor.; Am. bor.
foliosa, Duby, ex Decne. in Jacquem. Voy. Bot. 142. t. 146 = sarmentosa.
foliosa, Klatt, in Linnaea, xxxii. (1863) 293 = strigilosa.
Friesii, *Trautv. in Act. Hort. Petrop.* ix. (1884) 7.—N. Zemlaya.
frutescens, Lap. Hist. Abr. Pl. Pyr. 92 = cylindrica.
geraniifolia, *Watt, in Journ. Linn. Soc.* xx. (1882) 17.—Reg. Himal.
glacialis, Hoppe, in Koch, Syn. ed. II. ii. 670 = alpina.
globifera, *Duby, in DC. Prod.* viii. 48.—Reg. Himal.
globifera, Klatt, in Linnaea, xxxii. (1863) 292 = microphylla.
globifera, Schur, Enum. Pl. Transs. 551 = villosa.
Gmelini, *Gaert. Fruct.* i. 232.—Sibir.
Halleri, [Honck.] Verz. Gew. Teutschl. 561 = carnea.
Halleri, *Linn. Sp. Pl.* 142.—Europ.
Hausmanni, *Leyb. in Flora*, xxxv. (1852) 101.—Austria.
hedreantha, *Griseb. Spicil. Fl. Rumel.* ii. 3.—Thracia.
Heerii, Hegetschw. ex Hausm. Fl. Tirol. iii. 1474 = Hausmanni.
Heerii, Koch, Syn. Fl. Germ. ed. II. 670 = bryoides.
helvetica, All. Fl. Pedem. i. 91 = bryoides.
hirtella, L. Dufour, in Act. Soc. Linn. Bord. viii. (1836) 100 = pubescens.
Hookeriana, *Klatt, in Linnaea*, xxxii. (1863) 293.—Reg. Himal.
hybrida ×, *Kerner, in Oestr. Bot. Zeitschr.* xxv. (1875) 159.—Europ.
imbricata, *Lam. Fl. Fr.* ii. 253.—Europ.
incana, Lam. Illustr. i. 432 = villosa.
incisa, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 16 = rotundifolia.
intermedia, Ledeb. Fl. Ross. iii. 20 = armeniaca.
Jacquemontii, Duby, in DC. Prod. viii. 50 = villosa.
Lachenalii, C. C. Gmel. Fl. Bad. i. 2 = carnea.
lactea, *Linn. Sp. Pl.* 142.—Europ.
lactea, Vill. Hist. Pl. Dauph. ii. 476 = obtusifolia.
lactea, Pall. Reise, ii. 618 = septentrionalis.
lactiflora, [*Fisch.*] *Hort. Gorenk.* ed. I. (1808) 30.—Sibir.
lactiflora, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 728. n. 717 = septentrionalis.
Laggeri, *Huet, in Ann. Sc. Nat.* Sér. III. xix. (1853) 254.—Europ.
lanuginosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 15.—Reg. Himal.
Lehmanni, *Wall. Cat.* n. 617.—Reg. Himal.
Lehmanniana, Spreng. in Isis, (1817) 1289 = A. Chamaejasme.
linearis, R. Grah. in Edinb. N. Phil. Journ. (Apr.-June, 1829) 179 = septentrionalis.

ANDROSACE:—
longifolia, C. Koch, in Linnaea, xxiii. (1850) 610 = albana.
longifolia, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 202.—Mongolia.
longiscapa, C. Koch, in Linnaea, xxiii. (1850) 611 = A. Chamaejasme.
longiseta, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 61 = obtusifolia.
lutea, Lam. Fl. Fr. ii. 253 = Douglasia Vitaliana.
macrantha, *Boiss. & Huet, Diagn.* Ser. II. iii. 119.—Armen.
macrocarpa, *Ledeb. ex Duby, in DC. Prod.* viii. 55, nomen.—Sibir.
Mathildae, Levier, in Nuov. Giorn. Bot. ix. (1877) 43 = alpina.
maxima, *Linn. Sp. Pl.* 141.—Europ.; Oriens; As. et Afr. bor.
microphylla, *Hook. f. Fl. Brit. Ind.* iii. 498.—Reg. Himal.; Chin.
mucronifolia, *Watt, in Journ. Linn. Soc.* xx. (1884) 17.—Reg. Himal.
multiflora, Lam. Fl. Fr. ii. 252 = septentrionalis.
multiflora, Steud. Nom. ed. II. i. 94 = imbricata.
multiscapa, *Duby, in DC. Prod.* viii. 51.—As. Min.; Syria.
muscoidea, *Duby, l. c.* 48.—Reg. Himal.
nana, Hornem. Hort. Hafn. i. 184 = elongata.
neglecta, *Cleve, in Bull. Soc. Nat. Mosc.* xlv. (1872) i. 430.—Mont. Ural.
obovata, Wall. ex Duby, in DC. Prod. viii. 35 = Primula floribunda.
obtusifolia, *All. Fl. Pedem.* i. 90. t. 46. f. 1.—Europ.
obtusifolia, Willd. ex Ledeb. Fl. Ross. iii. 18 = A. Chamaejasme.
occidentalis, *Pursh, Fl. Am. Sept.* i. 137.—Am. bor.
ochotensis, *Willd. ex Roem. & Schult. Syst.* iv. 786.—Sibir.
odoratissima, Schreb. Dec. Pl. Min. Cogn. 7. t. 4 = villosa.
olympica, Boiss. Diagn. Ser. I. iv. 37 = villosa.
orbicularis, Roem. & Schult. Syst. iv. 816, Ind. = Gmelini.
Pacheri, Leyb. in Flora, xxxvi. (1853) 585 = alpina.
parviflora, Jacq. ex Duby, in DC. Prod. viii. 54 = rotundifolia.
patens, Wright, ex A. Gray, in Mem. Am. Acad. N.S. vi. (1858–9) 401 = saxifragaefolia.
pauciflora, Vill. Hist. Pl. Dauph. ii. 477 = obtusifolia.
pedemontana, Reichb Ic. Fl. Germ. xvii. 48 = obtusifolia.
pedunculata, [*Clairv.*] *Man. Herb.* 57.—Europ.
penicillata, Schott, Nyman & Kotschy, Anal. Bot. 18 = villosa.
pennina, Gaud. Fl. Helv. ii. 108 = alpina.
pleiocarpa, Walp. Ann. iii. 7 = seq.
pleioscapa, C. Koch, in Linnaea, xxiii. (1850) 611 = armeniaca.
primulina, Spreng. Syst. iv. Cur. Post. 56 = Primula pusilla.
primuloides, D. Don, Prod. 81 = Primula pusilla.
primuloides, Duby, in DC. Prod. viii. 51 = sarmentosa.
primuloides, Moench, Meth. Suppl. 152 = Primula cortusioides.
puberula, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 105 = carnea.
pubescens, DC. Fl. Fr. iii. 438 = alpina.
pyrenaica, *Lam. Illustr.* i. 432.—Pyren.
radiata, Lehm. in Roem. & Schult. Syst. iv. 164 = filiformis.
Reverchoni, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 105 = carnea.
rosea, Jord. & Fourr. l. c. 104 = carnea.
rotundifolia, *Hardw. in As. Res.* iv. (1795) 350.—Reg. Himal.; China.
rotundifolia, Lehm. ex Roem. & Schult. Syst. iv. 165 = Gmelini.
rotundifolia, Sm. Exot. Bot. ii. 107 = saxifragaefolia.
rugosa, [Clairv.] Man. Herb. 57 = Douglasia Vitaliana.
sarmentosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 14.—Reg. Himal.
saxifragaefolia, *Bunge, Enum. Pl. Chin. Bor.* 53.—China.
Selago, *Hook. f. & Thoms. ex Watt, in Journ. Linn. Soc.* xx. (1882) 18.—Reg. Himal.
sempervivoides, *Jacquem. ex Duby, in DC. Prod.* viii. 50.—Reg. Himal.; Mongol.

ANDROSACE:—
septentrionalis, *Linn. Sp. Pl.* 142.—Europ.; Caucas.; As. et Am. bor.
spathulata, Cav. Ic. v. 56 = Samolus spathulatus.
spatulata, Steud. Nom. ed. II. i. 93 = praec.
stricta, Hartm. ex Duby, in DC. Prod. viii. 44 = Primula stricta.
strigilosa, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 10.—China.
tomentosa, Schlecht. ex Duby, in DC. Prod. viii. 47 = imbricata.
triflora, *Adans. in Mém. Soc. Nat. Mosc.* v. (1817) 89.—Sibir.
valerianoides, Lehm. ex Spreng. in Isis, (1817) 162 = albana.
villosa, Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind. iii. 498 = microphylla.
villosa, Jacq. Fl. Austr. t. 352 = A. Chamaejasme.
villosa, *Linn. Sp. Pl.* 142.—Europ.; Oriens; As. temp.
villosa, S. G. Gmel. It. ii. 196 = maxima.
Vitaliana, Lap. Hist. Abr. Pl. Pyr. 94 = Douglasia Vitaliana.
Wiedemanni, *Boiss. Fl. Orient.* iv. 16.—Anatol.
Wulfeniana, Sieber, ex Koch, Syn. Fl. Germ. ed. II. ii. 670 = hedreantha.

ANDROSAEMUM, [Tourn.] ex Adans. Fam. ii. 444 (1763) = **Hypericum**, Linn.
calycinum, Presl, ex Steud. Nom. ed. II. i. 94 = H. calycinum.
Cambessedesii, Coss. ex Nym. Consp. 131.—Ins. Balear.
foetidum, Spach, ex Willk. & Lange, Prod. iii. 589 = H. hircinum.
grandifolium, Reichb. Ic. Fl. Germ. xvi. 70.—Europ.
hircinum, Spach, Hist. Vég. Phan. v. 419 = H. hircinum.
officinale, All. Fl. Pedem. ii. 147 = H. Androsaemum.
parviflorum, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 361 = H. elatum.
pyramidale, Spach, Hist. Vég. Phan. v. 417 = H. elatum.
vulgare, Gaert. Fruct. i. 282. t. 59 = H. Androsaemum.
Webbianum, Spach, Hist. Vég. Phan. v. 418 = H. grandiflorum.
xylosteifolium, Spach, l. c. 420 = H. inodorum.

ANDROSCEPIA, Brongn. in Duperr. Voy. Coq. Bot. 77 (1829) = **Anthistiria**, Linn. (Gram.).
Anathera, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 249 = Anth. Anathera.
Barteri, Anderss. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 176 = Anth. Barteri.
gigantea, Brongn. in Duperr. Voy. Coq. Bot. 78 = Anth. gigantea.
tremula, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 247 = Anth. tremula.

ANDROSEMUM, Neck. Elem. ii. 395 (1790) = praec.

ANDROSTEMMA, Lindl. Swan Riv. App. 46 (1839) = **Conostylis**, R. Br. (Haemodor.).
junceum, Lindl. l. c. = C. Androstemma.

ANDROSTEPHIUM, Torr. in Bot. Mex. Bound. 218 (1858). *LILIACEAE*, Benth. & Hook. f. iii. 801.
breviflorum, *S. Wats. in Amer. Natural.* vii. (1873) 303.—Utah.
violaceum, *Torr. in Bot. Mex. Bound.* 219.—Texas.

ANDROSTOMA, Hook. f. Fl. Antarct. i. 44. t. 30 (1845) = **Cyathodes**, Labill. (Orchid.).
empetrifolia, Hook. f. l. c. = C. empetrifolia.

ANDROSTYLIUM, Miq. in Nat. Verh. Wet. Haarl. vii. (Stirp. Surin.) 93 (1851) = **Clusia**, Linn. (Guttif.).
Fockeanum, Miq. l. c. = C. Fockeana.

ANDROSYNE, Salisb. Gen. Pl. Fragm. 61 (1866). An **Dioscorea**, Linn. 'Genus dubium, prope Paridem.' —Cf. Benth. & Hook. f. Gen. iii. 762.
gracilis, Salisb. l. c.—Ins. Nicob.

ANDROTRICHUM, Brongn in Duperr. Voy. Coq. Bot. 177. t. 32 (1829). *CYPERACEAE*, Benth. & Hook. f. iii. 1046.
COMOSTEMUM, Nees, in Linnaea, ix. (1834) 283.
polycephalum, *Brongn. l. c.*—Bras.

ANDROTROPIS, R. Br. ex Wall. Cat. n. 8398 (1847) = **Acranthera**, Arn. (Rubiac.).
tomentosa, R. Br. ex Wall. l. c. = Acranth. tomentosa.

ANDRYALA, Linn. Gen. ed. I. 238 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 517.
ERIOPHORUS, Vaill. ex DC. Fl. Fr. iv. 36 (1808).
FORNEUM, Adans. Fam. ii. 112 (1763).
ROTHIA, Schreb. Gen. Pl. 531 (1791).
VOIGTIA, Roth, in Roem. & Ust. Mag. iv. (x.) 17 (1790).
aestivalis, *Pomel, Nouv. Mat. Fl. Atl.* 259 (err. typ. oestivalis).—Afr. bor.
Agardhii, *Haensel. ex DC. Prod.* vii. 244.—Hispan.
allochroa, *Hoffmgg. & Link, Fl. Port.* ii. 154.—Hispan.
arenaria, Boiss. & Reut. Pugill. Pl. Hispan. 71 = integrifolia.
aurea, Scop. Ann. ii. Hist. Nat. (1769) 58 = Crepis aurea.
Bourgeauii, Sch. Bip. ex Webb. & Berth. Phyt. Canar. ii. 417 = pinnatifida.
candidissima, Desf. Cat. Hort. Par. ed. III. 399 = varia.
cheiranthifolia, L'Hérit. Stirp. Nov. 35. t. 18 = varia.
cheiranthifolia, Link, ex Steud. Nom. ed. II. i. 94 = pinnatifida.
chondrilloides, *Scop. Fl. Carn.* ed. II. ii. 115.—Europ.
coronopifolia, Hoffmgg. & Link, Fl. Port. ii. 155 = pinnatifida.
corymbosa, Brot. Fl. Lusit. i. 337 = nigricans.
corymbosa, Lam. Illustr. iii. t. 657 = sinuata.
cossyrensis, Guss. Fl. Sic. Syn. ii. 407 = dentata.
crithmifolia, *Ait. Hort. Kew.* iii. 129.—Ins. Canar.
dentata, *Sibth. & Sm. Fl. Graec.* t. 811.—Reg. Mediterr.
denudata, Soland. ex Lowe, Man. Fl. Mad. i. 563 = cheiranthifolia,
diffusa, Jan. ex DC. Prod. vii. 246 = integrifolia.
dissecta, *Hoffmgg. & Link, Fl. Port.* ii. 153.—Hispan.
Ficalheana, *Daveau, in Bolet. Soc. Brot.* i. (1883) 42, 51.—Lusit.
floccosa, *Pomel, Nouv. Mat. Fl. Atl.* 3.—Afr. bor.
glandulosa, Lam. Encyc. i. 154 = varia.
gomeraea, Sch. Bip. ex Webb & Berth. Phyt. Canar. ii. 417 = pinnatifida.
incana, DC. Fl. Fr. Suppl. 445 = ragusina.
integrifolia, *Linn. Sp. Pl.* 808.—Reg. Mediterr.
integrifolia, Ucria, Hort. Panhorm. 336, ex Guss. Fl. Sic. Syn. ii. 404 = Hieracium crinitum.
laciniata, Lam. Encyc. i. 153 = ragusina.
lanata, Habl. Taur. 155 = Leontodon biscutellaefolius.
lanata, Linn. Cent. i. n. 57, in Amoen. Acad. iv. 288 = Hieracium andryaloides.
lanata, Vill. Hist. Pl. Dauph. iii. 65 = integrifolia.
lanata, Sibth. & Sm. Prod. ii. 141 = Hieracium pannosum.
laxiflora, *DC. Prod.* vii. 246.—Hispan.; Marocc.
lyrata, Pourr. in Mém. Acad. Toul. iii. (1788) 308 = ragusina.
macrocephala, Boiss. ex DC. Prod. vii. 244 = ragusina.
malacitana, Haensel. ex Willk. in Willk. & Lange, Prod. ii. 272 = laxiflora.
mogadorensis, *Coss. ex Hook. f. Bot. Mag.* t. 6010.—Marocc.
mollis, [Asso] Mant. Stirp. Arag. 175 = integrifolia.
nemausensis, Vill. Prosp. 37 = Pterotheca sancta.
nigricans, *Poir. Voy. Barb.* ii. 228.—Numidia.
nudicaulis, Lam. Encyc. i. 154 = Pterotheca sancta.
oestivalis, err. typ. *vide* aestivalis.
parviflora, Lam. Fl. Fr. ii. 117 = integrifolia.
pinnatifida, *Ait. Hort. Kew.* ed. I. iii. 129.—Ins. Canar.
pontana, Vill. Prosp. 37 = Crepis montana.
ragusina, *Linn. Sp. Pl.* ed. II. 1136.—Reg. Mediterr.
ramosissima, Boiss. Elench. 63 = ragusina.
Reboudiana, *Pomel, Nouv. Mat. Fl. Atl.* 259.—Afr. bor.
robusta, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) repr. 18.—Ins. Mader.
Rothia, *Pers. Syn.* ii. 378.—Hispan.
runcinata, Pers. l. c. = integrifolia.
sinuata, Linn. Sp. Pl. 808 = integrifolia.
sinuata, Reichb. Ic. Fl. Germ. xiv. 1436 = integrifolia.
sinuata, Willk. & Lange, Prod. ii. 271 = ragusina.
tenuifolia, *DC. Prod.* vii. 245.—Sicilia.
tenuifolia, Tenore, Fl. Neap. Syll. App. v. 38 = dentata.
tomentosa, Salisb. Prod. 182 = ragusina.

ANDRYALA:—
tomentosa, Scop. Delic. Insub. ii. 12 = varia.
undulata, J. & C. Presl, Delic. Prag. 113 = integrifolia.
uniflora, Schrank, Baier. Fl. ii. 343 = integrifolia.
varia, *Lowe, ex DC. Prod.* vii. 245.—Ins. Mader.
Webbiana, Sch. Bip. ex Webb & Berth. Phyt. Canar. iii. 417 = pinnatifida.

ANDRZEIOWSKIA, Reichb. Pl. Crit. i. 15. t. 13 (1837). *CRUCIFERAE*, Benth. & Hook. f. i. 70.
MACROCERATIUM, Reichb. Consp. 184 (1828).
Cardamine, *Reichb. l. c.*—Pontus; Syria.

ANECHITES, Griseb. Fl. Brit. W. Ind. 410 (1864). *APOCYNACEAE*, Benth. & Hook. f. ii. 707.
adglutinata, *Miers, Apocyn. S. Am.* 236.—Ind. occ.
Asperuginis, *Griseb. Fl. Brit. W. Ind.* 410.—Cuba.
circinalis, *Miers, Apocyn. S. Am.* 236.—Ind. occ.
lappulacea, *Miers, l. c.* 237.—Ins. S. Doming.
revoluta, *Miers, l. c.* 238.—Bras.
thomasiana, *Miers, l. c.* 237.—Ins. S. Thomae.

ANECIO, Neck. Elem. i. 28 (1790) = **Senecio**, Linn. (Compos.).

ANECOCHILUS, Blume, Bijdr. 411. t. 15 (1825) = **Anoectochilus**, Blume (1828) (Orchid.).

ANECTOCHILUS, Blume, Orch. Archip. Ind. 44 (1858) = **Anoectochilus**, Blume (Orchid.).

ANEILEMA, R. Br. Prod. 270 (1810). *COMMELINACEAE*, Benth. & Hook. f. iii. 849.
AMELINA, C. B. Clarke, Commel. et Cyrt. Beng. 38. t. 26 (1874).
ANILEMA, Kunth, Enum. Pl. iv. 64 (1843).
APHYLAX, Salisb. in Trans. Hort. Soc. i. (1812) 271.
BAUSCHIA, Seub. in Vidensk. Meddel. Kjoeb. 123 (1872).
DICHOESPERMUM, Wight, Ic. vi. 31. t. 2078 (1853).
DICTYOSPERMUM, Wight, l. c. 29. t. 2069 (1853).
DILASIA, Rafin. Fl. Tellur. iv. 122 (1836).
LAMPRODITHYROS, Hassk. in Peters, Mossamb. Reise, Bot. 529 (1863).
MURDANNIA, Royle, Illustr. Bot. Himal. 403. t. 95 (1839).
PILETOCARPUS, Hassk. in Flora, xlix. (1866) 212.
PRIONOSTACHYS, Hassk. l. c.
RHOPALEPHORA, Hassk. ex Mohl & Schlecht. in Bot. Zeit. (1864) 58.
STREPTYLIS, Rafin. Fl. Tellur. iv. 122 (1836).
TALIPULIA, Rafin. l. c. ii. 17 (1836).
acuminatum, Benth. Fl. Austral. vii. 85 = laxum.
acuminatum, *R. Br. Prod.* 270.—Austral.
adhaerens, Kunth, Enum. Pl. iv. 72 = aequinoctiale.
aequinoctiale, *Kunth, l. c.*—Afr. trop.
affine, R. Br. Prod. 271 = gramineum.
africanum, Beauv. Fl. Owar. ii. 57. t. 93 = Floscopa africana.
anthericoides, R. Br. Prod. 271 = gramineum.
asperum, Buch.-Ham. ex Wall. Cat. sub n. 5204 = Floscopa scandens.
beninense, *Kunth, Enum. Pl.* iv. 73.—Afr. trop.
biflorum, *R. Br. Prod.* 270.—Austral.
bracteolatum, Mart. in Flora, xxiv. (1841) II. Beibl. ii. 59 = ovato-oblongum.
brasiliense, *C. B. Clarke, in DC. Monog. Phan.* iii. 225. —Bras.
calandrinoides, *F. Muell. Fragm.* ix. 191.—Austral.
canaliculatum, Dalz. in Hook. Kew Journ. iii. (1851) 137 = spiratum.
compressum, Dalz. l. c. 138 = nudiflorum.
conspicuum, *Kunth, Enum. Pl.* iv. 69.—Ind. or.
cordatum, Buch.-Ham. ex Wall. Cat. sub n. 5223 = Commelina herbacea.
crispatum, R. Br. Prod. 270 = Pollia crispata.
crocea, Griff. Notul. iii. 235 = ochraceum.
cymosum, Kunth, Enum. Pl. iv. 69 = Floscopa scandens.
debile, Wall. Cat. n. 5215 = nudiflorum.
densiflorum, Kunth, Enum. Pl. iv. 69 = Floscopa scandens.
diandrum, Buch.-Ham. in Wall. Cat. n. 5211 = nudiflorum.
didymum, Buch.-Ham. l. c. n. 5202 = Commelina didyma.
didymum, Seem. Bot. Voy. Herald, 422 = Pollia sorzogonensis.

ANEILEMA:—
dimorphum, *Dalz. in Hook. Kew Journ.* iii. (1851) 138. —Ind. or.
dimorphum, Thw. Enum. Pl. Zeyl. 322 = zeylanicum.
divergens, *C. B. Clarke, Commel. et Cyrt. Beng.* 28. t. 16.—Ind. or.
diversifolium, Hassk. in Miq. Pl. Jungh. 142 = nudiflorum.
Dregeanum, *Kunth, Enum. Pl.* iv. 73.—Afr. austr.
Ehrenbergii, *C. B. Clarke, in DC. Monog. Phan.* iii. 229.—Afr. trop.
elatum, *Dalz. in Hook. Kew Journ.* iii. (1851) 137. —Ind. or.
elatum, Kunth, Enum. Pl. iv. 70 = herbaceum.
ensifolium, Wight, Ic. Pl. Ind. Or. t. 2072 = giganteum.
esculentum, *Wall. Cat.* n. 5208.—Ind. or.
esculentum, Wall. l. c. litt. C. (nec A. B.) = sinicum.
filiforme, Buch.-Ham. Wall. l. c. n. 5209 = vaginatum.
filipes, Mart. in Flora, xxiv. (1841) II. Beibl. ii. 59 = geniculatum.
floribundum, Hook. & Arn. Bot. Beech. Voy. 311 = Leptorhoeo filiformis.
foliosum, Hassk. in Miq. Pl. Jungh. 144 = nudiflorum.
Forskalei, *Kunth, Enum. Pl.* iv. 71.—Arab.
Gardneri, *Seub. in Mart. Fl. Bras.* iii. 259.—Bras.
giganteum, *R. Br. Prod.* 271.—Austral.
glaucum, *Thw. ex C. B. Clarke, in DC. Monog. Phan.* iii. 200.—Zeylan.
gracile, *C. B. Clarke, l. c.* 228.—Afr. occ.
gramineum, *R. Br. Prod.* 270.—Austral.
Hamiltonianum, *Wall. Cat.* n. 5222.—Assam.
herbaceum, *Wall. l. c.* n. 5223.—Ind. or.; Malaya.
herbaceum, Wall. l. c. n. 8223 = protensum.
hirtum, *A. Rich. Tent. Fl. Abyss.* ii. 343.—Abyss.
hispidum, D. Don, Prod. 45 = Floscopa scandens.
Hookeri, *C. B. Clarke, Commel. et Cyrt. Beng.* 28. t. 17.—Ind. or.
japonicum, *Kunth, Enum. Pl.* iv. 70.—Japon.
Junghunianum, Miq. Fl. Ind. Bat. iii. 528 = nudiflorum.
Keisak, *Hassk. Commel. Ind.* 32.—Japon.
Koenigii, *Wall. Cat.* n. 5214.—Ind. or.
lanceolatum, Baker, ex C. B. Clarke, in DC. Monog. Phan. iii. 227 = Schweinfurthii.
lanceolatum, *Benth. in Hook. Niger Fl.* 546.—Afr. trop.
lancifolium, Griff. Notul. iii. 236 = nudiflorum.
lanuginosum, *Wall. Cat.* n. 5221.—Ind. or.
latifolium, Wight, Ic. Pl. Ind. Or. t. 2072 = herbaceum.
laxum, R. Br. Prod. 270 = Commelina laxa.
laxiflorum, Benth. ex C. B. Clarke, in DC. Monog. Phan. iii. 226 = ovato-oblongum.
laxum, *R. Br. Prod.* 270.—Austral.
lineolatum, Kunth, Enum. Pl. iv. 69 = herbaceum.
longifolium, Hook. Exot. Fl. t. 204 = giganteum.
longifolium, Wall. Cat. n. 5213 = divergens.
loriforme, Hassk. in Miq. Pl. Jungh. 143 = nudiflorum.
Loureirii, *Hance, in Seem. Journ. Bot.* vi. (1868) 250. —China.
macrophyllum, R. Br. Prod. 270 = Pollia macrophylla.
medicum, *R. Br. l. c.* 271.—China.
melanosticum, Hance, in Seem. Journ. Bot. vii. (1869) 167 = esculentum.
micranthum, *Kunth, Enum. Pl.* iv. 70.—Ind. or.
minutum, Kunth, l. c. 661 = nudiflorum.
monadelphum, *R. Br. ex Kunth, l. c.* 70.—Java.
montanum, Thw. Enum. Pl. Zeyl. 322 = zeylanicum.
montanum, *Wight, in Wall. Cat.* n. 5203.—Ind. or.
nanum, Kunth, Enum. Pl. iv. 65 = spiratum.
nudicaule, Kunth, l. c. 67 = nudiflorum.
nudiflorum, F. Muell. Fragm. viii. 62 = giganteum.
nudiflorum, Miq. ex C. B. Clarke, in DC. Monog. Phan. iii. 206, 215 = dimorphum, Koenigii.
nudiflorum, *R. Br. Prod.* 271.—Ind. or.
nudiflorum, Wall. Cat. n. 5224 = vaginatum.
Nummularia, Miq. Fl. Ind. Bat. 535 = spiratum.
ochraceum, *Dalz. in Hook. Kew Journ.* iii. (1851) 135.—Ind. or.
oliganthum, Franch. & Sav. Enum. Pl. Jap. ii. 522 = Keisak.
ovalifolium, *Hook. f. ex C. B. Clarke, in DC. Monog. Phan.* iii. 219.—Ind. or.
ovatum, *Wall. Cat.* n. 5206.—Malaya.
ovato-oblongum, *Beauv. Fl. Owar.* ii. 71. t. 104.—Afr. trop.
paniculatum, *Wall. Cat.* n. 5216.—Ind. or.
paniculatum, Wight, Ic. Pl. Ind. Or. t. 2075 = dimorphum.

ANEILEMA:—
pauciflorum, Dalz. in Hook. Kew Journ. iii. (1851) 136 = vaginatum.
pauciflorum, *Wight, Ic. Pl. Ind. Or.* t. 2077.—Ind. or.
pedunculosum, *C. B. Clarke, in DC. Monog. Phan.* iii. 228.—Afr. austr.
Petersii, *C. B. Clarke, l. c.* 225.—Afr. trop.
pilosum, Wall. Cat. n. 5219 = Commelina pilosa.
poaeoides, *Seub. in Mart. Fl. Bras.* iii. 259.—Bras.
protensum, *Wall. Cat.* n. 5218.—Ind. or.
protensum, Thw. Enum. Pl. Zeyl. 322 = zeylanicum.
radicans, D. Don, Prod. Fl. Nep. 45 = nudiflorum.
reniforme, Buch.-Ham. in Wall. Cat. n. 5205 = Pollia subumbellata.
rivulare, A. Rich. Tent. Fl. Abyss. ii. 342 = Floscopa rivularis.
scaberrimum, Kunth, Enum. Pl. iv. 69 = protensum.
scapiflorum, *Kostel. Allg. Med.-Pharm. Fl.* i. 127.—Ind. or.
scapiflorum, Hook. f. & Thoms. ex C. B. Clarke, in DC. Monog. Phan. iii. 203 = divergens.
Schomburgkianum, *Kunth, Enum. Pl.* iv. 661.—Bras.
Schweinfurthii, *C. B. Clarke, in DC. Monog. Phan.* iii. 227.—Afr. trop.
sclerocarpum, *F. Muell. Fragm.* viii. 61.—Austral.
secundum, Wight, Ic. Pl. Ind. Or. t. 2074 = sinicum.
secundiflorum, Kunth, Enum. Pl. iv. 69 = Pollia sorzogonensis.
semiteres, Dalz. in Hook. Kew Journ. iii. (1851) 138 = paniculatum.
sepalosum, *C. B. Clarke, in DC. Monog. Phan.* iii. 202. —Afr. trop.
serotinum, Don, ex C. B. Clarke, l. c. 200 = scapiflorum.
siliculosum, F. Muell. Fragm. viii. 61 = acuminatum.
siliculosum, *R. Br. Prod.* 270.—Austral.
simplex, Kunth, Enum. Pl. iv. 71 = sinicum.
sinicum, Herb. Berol. ex C. B. Clarke, in DC. Monog. Phan. iii. 211 = nudiflorum.
sinicum, [*Lindl. in*] *Bot. Reg.* t. 659.—China.
spectabile, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 77 = Loureirii.
spicatum, Wall. Cat. n. 5217 = Loureirii.
spiratum, *R. Br. Prod.* 271.—Ind. or.
Tacazzeanum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 343 = aequinoctiale.
Tacazzeanum, Baker, in Trans. Linn. Soc. xxix. (1872) 163 = lanceolatum.
terminale, Wight, Ic. Pl. Ind. Or. t. 2076 = nudiflorum.
Thomsoni, *C. B. Clarke, in Journ. Linn. Soc.* xv. (1877) 121.—Reg. Himal.
trichocoleum, Schau. in Nov. Act. Acad. Leop. Carol. xix. (1843) Suppl. i. 448 (Pl. Meyen.) = nudiflorum.
triquetrum, *Wall. Cat.* n. 5220.—Ind. or.
tuberosum, Buch.-Ham. in Wall. Cat. n. 5207 = scapiflorum.
umbrosum, Kunth, Enum. Pl. iv. 71 = ovato-oblongum.
vaginatum, *R. Br. Prod.* 271.—Ind. or.
versicolor, *Dalz. in Hook. Kew Journ.* iii. (1851) 136. —Ind. or.
violaceum, F. Muell. ex C. B. Clarke, in DC. Monog. Phan. iii. 205 = gramineum.
vitiense, *Seem. Fl. Vit.* 314. t. 96.—Ins. Fiji.
Welwitschii, *C. B. Clarke, in DC. Monog. Phan.* iii. 229.—Angola.
zeylanicum, *C. B. Clarke, l. c.* 204.—Zeylan.

ANELASMA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 42 = **Abuta**, Aubl. (Menisp.).
concolor, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 260 = Ab. concolor.
Gardnerianum, Miers, l. c. Ser. II. vii. (1851) 43 = Ab. concolor.
guianense, Miers, l. c. = Ab. concolor.
intaminatum, Miers, l. c. Ser. III. xiv. (1864) 260.—Bras.
jamaicense, Miers, ex Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 50 = Pachygone domingensis.
laurifolium, Sagot, ex Benth. l. c. = Ab. concolor.
laurifolium, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 260.—Ins. S. Thomae.
Martianum, Miers, l. c. = Ab. concolor.
minutiflorum, Sagot, ex Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 50 = Pachygone domingensis.

ANELASMA :—
pallidum, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 260 = Ab. concolor.
Sellowianum, Miers, l. c.—Bras.
Spruceanum, Miers, ex Benth. in Hook. Kew Jour . iii. (1851) 116 = Ab. concolor.
strumosum, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 260.—Bras.
urophyllum, Miers, l. c.—Bras.

ANEMAGROSTIS, Trin. Fund. Agrost. 128 (1820) = **Apera**, Adans. (Gram.).
interrupta, Trin. l. c. 129 = Ap. interrupta.
Neesiana, Wight & Arn. ex Steud. Syn. Pl. Gram. 115 = Panicum semialatum.
Spica-venti, Trin. Fund. Agrost. 129 = Ap. Spica-venti.
tenella, Wight, ex Steud. Syn. Pl. Gram. 115 = Arundinella tenella.

ANEMANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 323 = **Anemone**, Tourn. (Ranunc.).
nemorosus, Fourr. l. c. = Anemone nemorosa.
ranunculoides, Fourr. l. c. = Anemone ranunculoides.

ANEMARRHENA, Bunge, in Mém. Acad. Pétersb. ii. (1831) 140 (Enum. Pl. Chin. Bor. 66). *LILIACEAE*, Benth. & Hook. f. iii. 782.
asphodeloides, *Bunge, ll. cc.*—China.

ANEMIA, Nutt. in Ann. Nat. Hist. Ser. I. i. (1838) 136 = **Houttuynia**, Thunb. (Piperac.).
californica, Nutt. l. c. = H. californica.

ANEMIOPSIS, Endl. Gen. Suppl. I. 1374 (1841) = ANEMOPSIS, Hook. & Arn.

ANEMITIS, Rafin. Fl. Tellur. iii. 87 (1836) = **Phlomis**, Tourn. (Labiat.).
rigida, Rafin. l. c. = P. Herba-venti.

ANEMONANTHEA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 724 (1821) = **Anemone**, Linn. (Ranunc.).
apennina, S. F. Gray, l. c. 724 = Anemone apennina.
nemorosa, S. F. Gray, l. c. 725 = Anemone nemorosa.
ranunculoides, S. F. Gray, l. c. = Anemone ranunculoides.

ANEMONE, Linn. Syst. ed. I. (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 4.
ANEMANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 323.
ANEMONANTHEA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 724 (1821).
ANEMONELLA, Spach, Hist. Vég. Phan. vii. 240 (1839).
BARNEOUDIA, C. Gay, Fl. Chil. i. 29. t. 1 (1845).
FLAMMARA, Hill, Veg. Syst. xvi. 37 (1770).
HARTIANA, Rafin. Neogenyt. 2 (1825).
HEPATICA, Dill. ex Linn. Syst. ed. I (1735).
HOMALOCARPUS, Schur, Enum. Pl. Transsyl. 3 (1866).
ISOPYRUM, Adans. Fam. ii. 460 (1763).
ORIBA, Adans. l. c. 459 (1763).
PREONANTHUS, Ehrh. Beitr. iv. 149 (1789).
PULSATILLA, [Tourn.] Linn. Syst. ed. I (1735).
SYNDESMON, Hoffmgg. in Flora, xv. (1832) Intell. 34.
acanthifolia, *Rich. Ind. Sem. Hort. Patav.* (1841).—Hab. ?
aconitifolia, Michx. Fl. Bor. Am. i. 320 = dichotoma.
aconitifolia, Turcz. in Bull. Soc. Nat. Mosc. (1840) 61 = narcissiflora.
acutiloba, *Laws. Ranunc. Canad.* 30.—Am. bor.
acutipetala, Schlcich. Cat. Pl. Helv. ed. III. 6 = A. Pulsatilla.
acquinoctialis, *Poepp. Fragm. Syn.* 27.—Peruv.
affinis, G. Don, Gen. Syst. i. 21 = pratensis.
alba, Gilib. Fl. Lituan. ii. 275 = nemorosa.
alba, Juss. in Ann. Mus. Par. iii. (1804) 248. t. 20. f. 1 = sylvestris.
alba, Goaty & Pons, in Bull. Soc. Bot. Fr. xxx. (1883) p. lxxix = coronaria.
albana, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 264.—Oriens ; As. bor.
alborosea, Gilib. Fl. Lituan. ii. 276 = vernalis.
alchemillaefolia, *E. Mey. ex Pritz. in Linnaea*, xv. (1841) 758.—Afr. austr.

ANEMONE :—
alpina, *Linn. Sp. Pl.* 539.—Europ. ; Caucas. ; Am. bor.
alpina, Scop. Fl. Carn. ed. II. i. 384 = baldensis.
altaica, *Fisch. ex Ledeb. Fl. Alt.* ii. 362.—Sibir. altaic.
ambigua, Turcz. ex Pritz. in Linnaea, xv. (1841) 601 = albana.
angulosa, *Lam. Encyc.* i. 169.—Europ. or.
anomala, *Rafin. Fl. Ludov.* 82 (quid ?).—Am. bor.
antucensis, *Poepp. Fragm. Syn.* 27.—Chili.
apennina, Auct. Orient. ex Boiss. Fl. Orient. i. 13 = blanda.
apennina, *Linn. Sp. Pl.* 541.—Europ.
apiifolia, Scop. Fl. Carn. ed. II. i. 385 = alpina.
arborea, Hort. ex Steud. Nom. ed. II. i. 95 = capensis.
arctica, Fisch. in Linnaea, vi. (1831) 574 = Richardsoni.
argentea, Raim. ex Wedd. Chlor. Andina, ii. 305 = Ranunculus Raimondii.
armena, *Boiss. Fl. Orient.* i. 10.—As. Min.
baicalensis, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 40.—As. temp.
baldensis, G. Don, Gen. Syst. i. 18 = multifida.
baldensis, Lam. Encyc. i. 614 = alpina.
baldensis, *Linn. Mant.* i. 78.—Europ.
barbulata, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) n. VII. 149.—China.
Bauhini, *Tausch, in Flora*, xxv. (1842) i. 69.—Europ.
Berlandieri, Pritz. in Linnaea, xv. (1841) 628 = decapetala.
bicolor, Poepp. ex Pritz. l. c. 624 = decapetala.
biflora, *DC. Syst.* i. 201.—Oriens ; Reg. Himal.
bilobata, Phil. f. Cat. Pl. Vasc. Chil. 5 = deeapetala.
blanda, *Schott & Kotschy, in Oestr. Bot. Wochenbl.* iv. (1854) 129.—Oriens.
Bogenhardiana, *Pritz. in Linnaea*, xv. (1841) 591.—Europ.
borealis, Richards. in Frankl. 1st Journ. ed. II. App. 22 = parviflora.
Bungeana, *Pritz. in Linnaea*, xv. (1841) 603.—Sibiria.
Burseriana, Scop. Fl. Carn. ed. II. i. 385 = alpina.
caelestina, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 4.—China.
caerulea, *DC. Syst.* i. 203.—Sibir.
caerulea, Lam. Fl. Fr. iii. 318 = appenina.
caffra, *Harv. Gen. S. Afr. Pl.* 9.—Afr. austr.
Campanella, Fisch. ex Ledeb. Fl. Ross. i. 22 = albana.
campaniflora, Stokes, Bot. Mat. Med. iii. 247 = pratensis.
canadensis, Linn. Syst. ed. XII. iii. App. 231 = dichotoma.
capensis, *Lam. Encyc.* i. 164.—Afr. austr.
caroliniana, Torr. & Gray, Fl. N. Am. i. 12, partim = heterophylla.
caroliniana, Walt. Fl. Carol. 156 = decapetala.
caucasica, Willd. ex Rupr. Fl. Caucas. 14 = blanda.
cernua, *Thunb. Fl. Jap.* 238.—Japon.
chilensis, Spreng. ex Eichl. in Mart. Fl. Bras. xiii. i. 152 = decapetala.
chinensis, *Bunge, Enum. Pl. Chin.* 2.—China.
coccinea, Jord. Diag. i. 57 = coronaria.
collina, Salisb. Prod. 371 = A. Pulsatilla.
Commersoniana, DC. in Deless. Ic. i. 4 = multifida.
coronaria, *Linn. Sp. Pl.* 539.—Reg. Mediterr. ; Oriens.
coronarioides, Hanry, Prod. Hist. Nat. Var, 142 = coronaria.
crassifolia, *Hook. Ic. Pl.* t. 257.—Tasmania.
cuneata, Schlecht. in Linnaea, vi. (1831) 574 = parviflora.
cuneifolia, Juss. in Ann. Mus. Par. iii. (1804) 248. t. 21 = parviflora.
curta, Wall. Cat. n. 4690 = Clematis montana.
cyanea, Risso, Fl. Nice, 7 = coronaria.
cylindrica, *A. Gray, in Ann. Lyc. N. York*, iii. (1828–36) 221.—Am. bor.
cynosurus, Griff. Notul. iv. 735 = Ranunculus falcatus.
dacica, Freyn, ex Borb. in Just, Bot. Jahresb. (1874) 1076, *err. typ.* = Arabis dacica.
dahurica, *Fisch. ex DC. Prod.* i. 17.—As. temp.
debilis, *Fisch. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 274 [bis].—Sibir.
decapetala, J. F. Gmel. Syst. 871 = Geum rivale.
decapetala, *Arduini, Animad. alt.* 27.—Am. bor. et austr.
decaphilla, Krock, Fl. Sil. ii. 1. 235 = Geum rivale.
deltoidea, *Dougl. in Hook. Fl. Bor. Amer.* i. 6. t. 3. f. A.—Am. bor. occ.
demissa, *Hook. f. & Thoms. Fl. Ind.* i. 23.—Reg. Himal.

ANEMONE :—
dichotoma, *Linn. Sp. Pl.* 540.—As. bor. ; Am. bor.
discolor, Royle, Illustr. Bot. Himal. 52 = obtusiloba.
dodecaphilla, Krock, Fl. Sil. ii. 1. 235 = Geum rivale.
Drummondii, *S. Wats. Bot. Calif.* ii. 424.—Calif.
dubia, Bell. App. Fl. Pedem. 232 = narcissiflora.
dubia, Wall. Cat. n. 4697 = rivularis.
elegans, Decne. in Rev. Hortic. (1852) 41. cum ic. = japonica.
elongata, *D. Don, Prod. Fl. Nep.* 194.—Reg. Himal.
eranthoides, *Regel, in Act. Hort. Pétrop.* viii. (1884) 691.—As. temp.
exigua, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 306.—China.
Falconeri, *Thoms. in Hook. Ic. Pl.* t. 899.—Reg. Himal.
Fanninii, *Harv. Gen. S. Afr. Pl.* ed. II. 2, nomen.—Natal.
fasciculata, Linn. Sp. Pl. 542 = narcissiflora.
fasciculata, Vahl, Symb. Bot. iii. 74 = umbellata.
Fischeriana, *DC. Prod.* i. 20.—Sibiria.
flaccida, *Fr. Schmidt, Reisen Amurl.* 103.—Japon. China.
flava, Gilib. Fl. Lituan. ii. 274 = ranunculoides.
flavescens, Zucc. in Flora, ix. (1826) I. 371 = patens.
flemmensis, Scop. Ann. ii. Hist. Nat. (1796) 54 = alpina.
formosa, *Clark, Trav.* ii. 145.—As. Min.
fragifera, Wulf. in Jacq. Misc. ii. 55 = baldensis.
fulgens, *J. Gay, in DC. Prod.* i. 18.—Reg. Mediterr.
fumariaefolia, Juss. in Ann. Mus. Par. iii. (1804) 247 = triternata.
geraniifolia, Wall. Cat. n. 4693 = rivularis.
gigantea, Raim. ex Wedd. Chlor. And. ii. 304 = Ranunculus gigantea.
glacialis, [Clairv.] Man. Herb. 180 = Halleri.
Glazioviana, *Urb. in Linnaea*, xliii. (1880–82) 25.—Bras.
globosa, Nutt. ex Pritz. in Linnaea, xv. (1841) 673 = multifida.
Gmeliniana, *Presl, ex Pritz. l. c.* 697.—Sibiria.
Gortschakowii, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 131 = biflora.
Govaniana, Lindl. Bot. Reg. (1844) Misc. 45 = polyanthes.
Govaniana, Wall. Cat. n. 4688 = obtusiloba.
gracilis, *Fr. Schmidt, Reisen Amurl.* 102.—Japon.
grandiflora, Hoppe, ex Reichb. Fl. Germ. Excurs. 732 = alpina.
grassensis, Goaty & Pons, in Bull. Soc. Bot. Fr. xxx. (1883) p. lxxviii = coronaria.
Grayi, *Behr. ex Kellogg, in Bull. Calif. Acad.* n. 1. (1884) 5.—Calif.
Griffithii, *Hook. f. & Thoms. Fl. Ind.* i. 24.—Reg. Himal.
groenlandica, O. F. Muell. in Fl. Dan. t. 566 = Coptis trifolia.
Hackelii, Steud. Nom. ed. I. 48 = Halleri.
Halleri, *All. Fl. Pedem.* ii. 170.—Europ.
helleboriflora, Richter, ex Pritz. in Linnaea, xv. (1841) 598 = pratensis.
helleborifolia, *DC. Syst.* i. 211.—Am. austr.
Hepatica, *Linn. Sp. Pl.* 538.—Europ. ; Oriens ; Am. bor.
hepaticifolia, *Hook. Ic. Pl.* t. 1.—Chili.
heterophylla, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 12.—Am. bor.
hirsuta, Gilib. Fl. Lituan. ii. 276 = sylvestris.
hirsuta, Moench, Meth. Suppl. 105 = virginiana.
hispida, Wall. Cat. n. 4694 = rivularis.
hortensis, Linn. Sp. Pl. 540 = stellata.
hortensis, Thore, Chl. Land. 238 = fulgens.
Hudsoniana, Richards. in Frankl. 1st Journ. ed. II. App. 22 = multifida.
hybrida, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 74 = japonica.
hybrida, Miq. ex Pritz. in Linnaea, xv. (1841) 573 = Halleri.
incisa, Ledeb. Ind. Hort. Dorp. Suppl. (1824) 1 = caerulea.
indeterminata, Banks, ex Walp. Rep. i. 23, sphalm. = parviflora.
integrifolia, *Humb. & Bonpl. ex DC. Syst.* i. 217.—Am. merid.
intermedia, G. Don, Gen. Syst. i. 21 = patens.
intermedia, Hoppe, ex Steud. Nom. ed. I. 48, 49 = montana.
intermedia, Schult. Obs. 101 = A. Pulsatilla.

ANEMONE:—
intermedia, Winkl. ex Pritz. in Linnaea, xv. (1841) 652 = A. nemorosa.
irregularis, Lam. Encyc. i. 167 = dichotoma.
isopyroides, *Juss. in Ann. Mus. Par.* iii. (1804) 249. —Sibir.
Jamesoni, *Hook. Ic. Pl.* t. 670.—Ecuador.
Jankae, *F. Schultz, in Flora*, xxxix. (1856) 205.—Transsyl.
japonica, *Sieb. & Zucc. Fl. Jap.* i. 15.—Japon.
lancifolia, Pursh, Fl. Am. Sept. ii. 386 = nemorosa.
lanigera, C. Gay, Fl. Chil. i. 22 = multifida.
latifolia, Bell. ex Re, in Mem. Acc. Sc. Torin. xxxiii. (1829) 233 = fulgens.
Laxmanni, Steud. Nom. ed. II. i. 96 = dichotoma.
lepida, Jord. Diag. i. 59 = hortensis.
lineariloba, *Fisch. ex Pritz. in Linnaea*, xv. (1841) 697.—Kamtschatka.
linearis, Schleich. ex Pritz. l. c. 661 = caerulea.
lobata, Pers. Syn. ii. 96 = palmata.
longipetala, Schleich. ex Steud. Nom. ed. I. 48, 49 = patens.
longiscapa, Wall. Cat. n. 4691 = polyanthes.
ludoviciana, Nutt. Gen. Am. ii. 20 = patens.
lutea, Lam. Fl. Fr. iii. 318 = ranunculoides.
macrorrhiza, Domb. ex Eichl. in Mart. Fl. Bras. xiii. 1. 151 = decapetala.
malvifolia, Linn. Sp. Pl. 1197 = palmata.
mexicana, *H. B. & K. Nov. Gen. et Sp.* v. 33.—Mexic.
micrantha, Hort. ex Steud. Nom. ed. II. i. 96 = alpina.
micrantha, Klotzsch, in Bot. Ergeb. Waldem. Reise, 133 = obtusiloba.
millefoliata, Bertol. Amoen. 574 = alpina.
minima, DC. Syst. i. 206 = nemorosa.
minuta, *Laxm. ex Sievers, in Pall. Neu. Nord. Beitr.* vii. 153.—Sibir.
mollis, Wall. Cat. n. 4689 = obtusiloba.
montana, *Hoppe, ex Sturm, Fl. Deutschl. Heft* xc. n. 2.—Europ.; Caucas.
Mouansii, Hanry, ex Arduino, Fl. Alp. Marit. 12 = coronaria.
multifida, *Poir. Encyc. Suppl.* i. 364.—Am. bor. et austr.
myrrhidifolia, Vill. Prosp. 50 = alpina.
narcissiflora, *Linn. Sp. Pl.* 542.—Europ.; Orient; As. bor.; Am. bor.
narcissiflora, Hook. & Arn. Bot. Beech. Voy. 121 = multifida.
nemorosa, *Linn. Sp. Pl.* 541.—Europ.; As. bor.; Am. bor.
nemorosa, Schang. in Pall. Neu. Nord. Beitr. vi. 34 = altaica.
nemorosa-alba, Crantz, Stirp. Austr. ed. I. fasc. ii. 101; ed. II. 121 = nemorosa.
nemorosa-lutea, Crantz, l. c. = ranunculoides.
nikoënsis, *Maxim. in Bull. Acad. Pétersb.* xviii. (1875) 275.—Japon.
nobilis, Jord. Diag. i. 57 = coronaria.
Nuttalliana, DC. Syst. i. 193 = patens.
Nuttallii, Nutt. in Journ. Acad. Soc. Nat. Phil. v. (1825) 158 = patens.
obsoleta, Hort. ex Steud. Nom. ed. II. i. 98 = pratensis.
obtusiloba, *D. Don, Prod.* 194.—Reg. Himal.
obtusiloba, Lindl. Bot. Reg. xxx. (1844) t. 65 = scaposa.
occidentalis, *S. Wats. in Proc. Am. Acad.* xi. (1876) 121. —Calif.
ochotensis, Fisch. Hort. Gorenk. ed. II. (1812) 47 = sylvestris.
ochroleuca ×, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 77.—Helvet.
octopetala, *Patrin, ex Pritz. in Linnaea*, xv. (1841) 697, nomen.—Hab.?
Oenanthe, Ucria, in Roem. Arch. i. 69 = coronaria.
palmata, *Linn. Sp. Pl.* 538.—Reg. Mediterr.
parviflora, *Michx. Fl. Bor. Am.* i. 319.—Am. bor.
patens, Hoppe, in Sturm, Deutschl. Fl. Heft xlvi. 1 = Halleri.
patens, *Linn. Sp. Pl.* 538.—Europ.; Am. bor.
Pavoniana, *Boiss. Diagn.* Ser. II. i. 6; *et ex Leresche & Levier, in Journ. Bot.* xvii. (1879) 196.—Iberia.
pavonina, Lam. Encyc. i. 166 = hortensis.
pedata, Rafin. in Med. Repos. New York, v. (1808) 361 = nemorosa.
pennsylvanica, Linn. Mant. 247 = dichotoma.
pentaphylla, Hook. ex Pritz. in Linnaea, xv. (1841) 657 = nemorosa.
pinnatifida, Dulac, Fl. Hautes-Pyr. 214 = Halleri.
pisciensis, Sism. Agr. Tosc. 223 (1801) = A. Pulsatilla.

ANEMONE:—
Pittonii ×, *Glowacki, in Verh. Zool.-Bot. Ges. Wien*, xix. (1869) 901.—Europ.
polyanthes, *D. Don, Prod. Fl. Nep.* 194.—Reg. Himal.
praecox, Salisb. Prod. 371 = A. Hepatica.
praestabilis, Jord. Diagn. i. 58 = coronaria.
pratensis, C. Koch, in Linnaea, xv. (1841) 248 = montana.
pratensis, *Linn. Sp. Pl.* 539.—Europ.
pratensis, Pall. ex Pritz. in Linnaea, xv. (1841) 637 = sylvestris.
pratensis, Sibth. Fl. Ox. 169; With. Arr. Brit. Pl. ed. III. 498 = A. Pulsatilla.
propinqua, G. Don, Gen. Syst. i. 21 = pratensis.
pubescens, Domb. ex Steud. Nom. ed. I. 458, in syn. = Ranunculus Krapfia.
Pulsatilla, Bieb. Fl. Taur. Cauc. ii. 18 = montana.
Pulsatilla, *Linn. Sp. Pl.* 539.—Europ.
Pulsatilla, Sturm, Fl. Deutschl. i. Heft lxxxix. 12 = pratensis.
punica, Sism. Agr. Tosc. 223 = A. Pulsatilla.
pusilla, DC. Syst. i. 197 = coronaria.
pusilla, Gaertn. ex Steud. Nom. ed. I. 49 = Geum anemonoides.
pygmaea, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 90 = apennina.
pyrenaica, Pall. ex Pritz. in Linnaea, xv. (1841) 640 = apennina.
quinquefolia, Linn. Sp. Pl. 541 = nemorosa.
Raddeana, *Regel, in Bull. Soc. Nat. Mosc.* xxxiv. (1861) ii. 16.—Reg. Amur.
ranunculiflora, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = ranunculoides.
ranunculiformis, St. Lag. l. c. 67 = ranunculoides.
ranunculoides, *Linn. Sp. Pl.* 541.—Europ.; Caucas.; Sibir.
ranunculoides, Richards. in Frankl. 1st Journ. App. 12 = Richardsoni.
ranunculoidi-nemorosa, Kunze, in Fl. Dan. t. 2723 = nemorosa.
reflexa, Ledeb. ex Spreng. Syst. ii. 662 = caerulea.
reflexa, *Steph. in Willd. Sp. Pl.* ii. 1282.—Sibir.
Regina, Risso, Fl. Nice, 6 = coronaria?
Richardsoni, *Hook. Fl. Bor. Am.* i. 6. t. 4.—Am. arct.
rigida, *C. Gay, Fl. Chil.* i. 25.—Chili.
Rissoana, Jord. Diag. i. 58 = coronaria.
rivularis, *Buch.-Ham. ex DC. Syst.* i. 211.—Ind. or.
rivularis, Wall. Cat. n. 4692 = elongata.
rosea, Hanry, ex Nym. Consp. 3 = coronaria.
Rossii, *S. Moore, in Journ. Linn. Soc.* xvii. (1880) 376 t. 16.—China.
rubra, Lam. Encyc. i. 163 = A. Pulsatilla.
rupestris, Jacquem. Bot. Voy. 5 = rupicola.
rupestris, *Wall. Cat.* n. 4696.—Reg. Himal.
rupicola, *Camb. in Jacquem. Bot. Voy.* iv. 5.—Reg. Himal.
Salesovii, Fisch. ex Pritz. in Linnaea, xv. (1841) 655 = altaica.
sanguinea, Pursh, ex Pritz. l. c. 672 = multifida.
scaposa, Edgew. in Trans. Linn. Soc. xx. (1851) 27 = polyanthes.
Sellowii, *Pritz. in Linnaea*, xv. (1841) 667.—Bras.
sibirica, *Linn. Sp. Pl.* 541.—Sibir.
silvestris, vide sylvestris.
slavica, *Reuss, Bericht. Oestr. Litt.* i. 82.—Europ.
sordida, Schur, Enum. Pl. Transs. 3 = sylvestris.
speciosa, Adams, ex Pritz. in Linnaea, xv. (1841) 685 = narcissiflora.
speciosa, *Rupr. Fl. Caucas.* 12.—Caucas.
sphenophylla, *Poepp. Fragm. Syn.* 27.—Chili.
stellata, Lam. Encyc. i. 166 = hortensis.
stolonifera, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 225.—Japon.
sulphurea, All. Fl. Pedem. ii. 170 = vernalis.
sulphurea, Linn. Mant. 78 = alpina.
sumatrana, *De Vriese, in Miq. Pl. Jungh.* 76.—Sumatra.
sylvestris, *Linn. Sp. Pl.* 540.—Europ.; Caucas.
sylvestris, Vill. Hist. Pl. Dauph. iii. 726 = A. Pulsatilla.
tenella, Banks, ex Pritz. in Linnaea, xv. (1841) 632 = parviflora.
tenella, Pursh, Fl. Am. Sept. ii. 386 = decapetala.
tenuifolia, *DC. Syst.* i. 196.—Afr. austr.
tenuifolia, Schleich. Cat. Pl. Helv. ed. IV. 7 = Bogenhardiana.
tetrasepala, *Royle, Illustr. Bot. Himal.* 53.—Reg. Himal.
thalictroides, *Linn. Sp. Pl.* 542.—Am. bor.

ANEMONE:—
Thomsoni, *Oliver, in Journ. Linn. Soc.* xxi. (1885) 397.—Afr. trop.
transylvanica, *Heuff. in Verh. Zool.-Bot. Ver. Wien*, (1858) 42.—Europ.
trifolia, DC. Prod. i. 20 = reflexa.
trifolia, *Linn. Sp. Pl.* 540.—Europ.; Am. bor.
trifoliata, Georgi, Beschr. Russ. Reich. iii. iv. 1058 = reflexa.
triloba, Stokes, Bot. Mat. Med. iii. 243 = A. Hepatica.
trilobata, Juss. in Ann. Mus. Par. iii. (1804) 247. t. 1 = decapetala.
trilobata, Pers. Syn. ii. 97 = parviflora.
triternata, *Vahl, Symb. Bot.* iii. 74. t. 65.—Am. austr.
trullifolia, *Hook. f. & Thoms. Fl. Ind.* i. 22.—Reg. Himal.
Tschernaewi, *Regel, in Act. Hort. Petrop.* viii. (1884) 690.—As. temp.
udensis, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 6).—Manchur.
umbellata, Lam. Fl. Fr. iii. 322; Willd. Sp. Pl. ii. 1284 = narcissiflora.
umbrosa, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 361.—Sibir.
uniflora, Pursh, ex Steud. Nom. ed. II. i. 96 = Thalictrum anemonoides.
uralensis, Fisch. ex DC. Prod. i. 19 = caerulea, baldensis.
Vahlii, Hornem. Fl. Dan. t. 2176 = Richardsoni.
variata, Jord. ex Nym. Consp. 3 = hortensis.
Ventreana, Hanry, ex Arduino, Fl. Alp. Marit. 12 = coronaria.
vernalis, *Linn. Sp. Pl.* 538.—Europ.
versicolor, Jord. Pugill. (1852) 1 = coronaria.
versicolor, Salisb. Prod. 371 = hortensis.
villosa, Royle, Illustr. Bot. Himal. 52 = polyanthes.
virginiana, *Linn. Sp. Pl.* 540.—Am. bor.
vitifolia, *Buch.-Ham. ex DC. Syst.* i. 210.—Reg. Himal.
Wahlenbergii, *Szontagh, ex Nym. Consp.* 2.—Europ.
Wallichiana, Royle, Illustr. Bot. Himal. 52 = albana.
Walteri, *Pursh, Fl. Am. Sept.* ii. 387.—Am. bor.
Wightiana, *Hook. Ic. Pl.* ii. t. 176.—Ind. or.
Wightiana, Wall. Cat. n. 4697 = rivularis.
Wolfgangiana, *Bess. ex Koch, in Roehl. Deutschl. Fl.* iv. 106.—Europ.

ANEMONELLA, Spach, Hist. Vég. Phan. vii. 239 (1839) = **Anemone**, Linn. (Ranunc.).
thalictroides, Spach, l. c. 240 = Anemone thalictroides.

ANEMONOPSIS, Pritz. Ind. Ic. 71, sphalm. = ANEMOPSIS, Hook.

ANEMONOPSIS, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1846) 181. *RANUNCULACEAE*, Benth. & Hook. f. i. 8.
XAVERIA, Endl. Gen. Suppl. iv. 30 (1847).
macrophylla, *Sieb. & Zucc. l. c.* 182.—Japon.

ANEMONOSPERMUM, Commers. ex Steud. Nom. ed. II. i. 450, in syn. (1840) = **Cryptostemma**, R. Br. (Compos.).

ANEMOPAEGMA, Mart. ex DC. Prod. ix. 187 (1845). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1037.
acutifolium, *DC. Prod.* ix. 188.—Bras.
album, *Mart. ex DC. l. c.*—Bras.
citrinum, *Mart. ex DC. l. c.* 189.—Bras.
clematideum, *Griseb. in Goett. Abh.* xix. (1874) 222. —Reg. Argent.
floridum, *Mart. ex DC. Prod.* ix. 188.—Bras.
glaucum, *Mart. ex DC. l. c.*—Bras.; Peruv.?
laeve, *DC. l. c.* 189.—Bras.
lanceaefolium, *DC. l. c.* 188.—Bras.
longidentatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = seq.
longidens, *DC. Prod.* ix. 190.—Bras.
mirandum, *Mart. ex DC. l. c.* 187.—Bras.
orbiculatum, *DC. l. c.* 190.—Panama.
Poeppigii, *DC. l. c.*—Am. austr.
prostratum, *DC. l. c.* 189.—Bras.
quinquedentatum, *Mart. ex DC. l. c.* 190.—Bras.
racemosum, *Mart. ex DC. l. c.* 189.—Bras.
scabriusculum, *Mart. ex DC. l. c.* 188.—Bras.
sessilifolium, Mart. ex DC. l. c. 187 = mirandum.
triplinervium, *Mart. ex DC. l. c.* 188.—Bras.
Vargasianum, *DC. l. c.* 190.—Bras.
velutinum, *Mart. ex DC. l. c.* 189.—Bras.

ANEMOPSIS, [Hook.] in Ann. Nat. Hist. i. (1838) 136, in nota; Hook. & Arn. Bot. Beech. Voy. 390. t. 92 = **Houttuynia**, Thunb. (Piperac.).
Berlanderi, C. DC. in Linnaea, xxxvii. (1871–73) 333 = H. californica.
californica, Hook. & Arn. Bot. Beech. Voy. 390. t. 92 = H. californica.
Ludovici-Salvatoris, Willk. in Regel, Gartenfl. xxvi. (1877) 228.—Calif.

ANEPSA, Rafin. Fl. Tellur. ii. 31 (1836) = **Veratrum**, Tourn. (Liliac.).
carinata, graminifolia, latifolia, spicata, Rafin. l. c.—(Spp. ignotae).

ANEPSIAS, Schott, Gen. Aroid. t. 73 (1858) = **Rhodospatha**, Poepp. (Aroid.).
Moritzianus, Schott, l. c.—Venezuela.

ANERINCLEISTUS, Korth. in Verh. Nat. Gesch. ed. Temminck, 250. t. 68 (1842). *MELASTOMACEAE*, Benth. & Hook. f. i. 753.
Griffithii, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 75.—Ind. or.
Helferi, *Hook. f. ex Triana, l. c.*—Ind. or.
hirsutus, *Korth. in Verh. Nat. Gesch.* 250. t. 68.—Sumatra.

ANERMA, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 170 (1842) = **Scleria**, Berg. (Cyperac.).
hispidula, Schrad. ex Nees, l. c. = S. hispidula.

ANESLEA, Reichb. Consp. 46 (1828) = ANNESLEA, Andr. = **Euryale**, Salisb. (Nymph.).

ANESORHIZA, Endl. Gen. 773 (1839), sphalm. = **Annesorhiza**, Cham. & Schlecht. (Umbell.).

ANETANTHUS, Hiern, ex Benth. & Hook. f. Gen. ii. 1025 (1876). *GESNERACEAE*, Benth. & Hook. f. ii. 1025.
alatus, *Benth. & Hook. f. Gen.* ii. 1025.—Bras.
gracilis, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 93.—Bras.
parviflora, *Benth. & Hook. f. Gen.* ii. 1025.—Mexic.
villosa, *Benth. & Hook. l. c.*—Bras.

ANETHUM, Tourn. ex Linn. Gen. ed. I. 82 (1737) = **Peucedanum**, Linn. (Umbell.).
arvense, Bess. Ind. Sem. Hort. Crem. (1820) = Carum Ridolfia.
arvense, Salisb. Prod. 168 = Peucedanum graveolens.
capense, Thunb. in Hoffm. Phytogr. Blaett. i. (1803) 26 = Carum capense.
chryseum, *Boiss. & Heldr. Diagn.* Ser. I. x. 32.—Hispan.
cymbocarpum, DC. Prod. iv. 186 = Cymbocarpum anethoides.
dulce, DC. Cat. Hort. Monsp. 78 = Foeniculum vulgare.
erythraeum, DC. Prod. iv. 186 = Cymbocarpum erythraeum.
Foeniculum, Linn. Sp. Pl. 263 = Foeniculum vulgare.
Foeniculum, Thunb. Prod. Pl. Cap. 51 = Carum capense.
graveolens, Linn. Sp. Pl. 263 = Peucedanum graveolens.
graveolens, Ucria, Hort. Pan. 138 = Carum Ridolfia.
minus, Gouan, Illustr. 20 = Foeniculum vulgare.
Panmori, Roxb. Fl. Ind. ii. 94 = Foeniculum vulgare.
Panmorium, Roxb. ex Flem. in As. Res. xi. (1810) 156 = praec.
Pastinaca, Wibel, Prim. Fl. Werth. 146 = Peucedanum Pastinaca.
pinnatum, Ruiz & Pav. ex Urban, in Mart. Fl. Bras. xi. 1. 342 = Apium Ammi.
piperitum, Ucria, in Roem. Arch. i. 68 = Foeniculum piperitum.
pusillum, Hort. ex Steud. Nom. ed. II. i. 97 = Carum Ridolfia.
rupestre, Salisb. Prod. 168 = Foeniculum vulgare.
segetum, Linn. Mant. 219 = Carum Ridolfia.
Sowa, Roxb. ex Flem. in As. Res. xi. (1810) 156 = Peucedanum graveolens.
trifoliatum, Roxb. Hort. Beng. 86; Fl. Ind. ii. 96 = Pimpinella Heyneana.

ANETIA, Endl. Gen. 923 (1839) = **Byrsanthus**, Guill. (Samyd.).
Byrsanthus, Steud. Nom. ed. II. i. 97 = B. Brownii.

ANEULOPHUS, Benth. in Benth. & Hook. f. Gen. i. 244 (1862). *LINEAE*.
CAUCANTHUS, Forsk. Fl. Aegypt. Arab. 91 (1775).
africanus, *Benth. l. c.*—Afr. trop.

ANEURISCUS, Presl, Symb. Bot. i. 71. t. 48 (1832) = **Symphonia**, Linn. (Guttif.).
Aubletii, Presl, l. c. 72 = S. globulifera.
exserens, Presl, l. c. t. 48 = S. globulifera.

ANGADENIA, Miers, Apocyn. S. Amer. 173 (1878). *APOCYNACEAE*.
almadensis, *Miers, l. c.* 179.—Bras.
amazonica, Miers, l. c. 175 (= *Anisolobus amazonicus*).
Berterii, *Miers, l. c.* 180.—Ins. S. Doming.
cognata, Miers, l. c. 176 (= *Anisolobus cognatus*).
coriacea, *Miers, l. c.* 177.—Guiana.
cubensis, *Miers, l. c.* 182.—Cuba.
Cururu, Miers, l. c. 175 (= *Anisolobus Cururu*).
elegans, *Miers, l. c.* 178.—Reg. Amazon.
elliptica, *Miers, l. c.* 180.—Bras.
geminata, Miers, l. c. 178 = Odontadenia geminata.
grandifolia, *Miers, l. c.* 175.—Reg. Amazon.
havanensis, *Miers, l. c.* 181.—Cuba.
hypoglauca, Miers, l. c. 173 = Odontadenia hypoglauca.
latifolia, *Miers, l. c.* 176.—Bras.
Lindeniana, *Miers, l. c.* 180.—Cuba.
majuscula, *Miers, l. c.* 174.—Reg. Amazon.
nitida, Miers, l. c. 177 = Odontadenia nitida.
pandurata, Miers, l. c. 182 = Echites pandurata.
Poeppigii, Miers, l. c. 179 = Odontadenia Poeppigii.
Prieurii, *Miers, l. c.* 182.—Guiana.
pruinosa, *Miers, l. c.* 177.—Bras.
reticulata, *Miers, l. c.* 179.—Bras.
Sagraei, Miers, l. c. 181 = Echites Sagraei.
Sprucei, Miers, l. c. 176 (= *Anisolobus Sprucei*).
sylvestris, Miers, l. c. 174 = Odontadenia sylvestris.
Valenzuelana, Miers, l. c. 181 = Echites Valenzuelana.

ANGEJA, Vandelli, Fasc. Pl. 13 (1771) = Quid?
Vandellii, Steud. Nom. ed. II. i. 97.—Bras.

ANGELANDRA, Endl. Gen. Suppl. iii. 69 (1843) = **Engelmannia**, Torr. & Gray (Compos.).
pinnatifida, Endl. ex Walp. Rep. ii. 976 = E. pinnatifida.

ANGELANDRA, Endl. Gen. Suppl. iv. III. 91 (1850) = **Croton**, Linn. (Euphorb.).
elliptica, Baill. Etud. Gén. Euph. 379 = C. ellipticum.

ANGELESIA, Korth. in Ned. Kruidk. Arch. iii. (1855) 384 = **Trichocarya**, Miq. (Rosac.).
splendens, Korth. l. c. = T. splendens.

ANGELICA, [Riv.] Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 916.
ANGELOPHYLLUM, Rupr. Rev. Umb. Kamtsch. 8 (1859).
CALLISACE, Fisch. in Hoffm. Gen. Umb. ed. II. 170 (1816).
EUSTYLIS, Hook. f. Fl. N. Zel. t. 19 (1853).
GINGIDIUM, Forst. Char. Gen. 41. t. 21 (1776).
GOMPHOPETALUM, Turcz. in Bull. Soc. Nat. Mosc. (1841) 537.
OSTERICUM, Hoffm. Gen. Umb. 162 (1814).
alpina, Krock. ex Steud. Nom. ed. I. 49 = sylvestris.
angustifolia, Hoffm. Gen. Umb. 162 = Peucedanum angustifolium.
anomala, Avé-Lall. in Ind. Sem. Hort. Petrop. ix. 57 = sylvestris.
aquilegifolia, Lam. Fl. Fr. iii. 452 = Siler trilobum.
Archangelica, Linn. Sp. Pl. 250 = Archangelica officinalis.
arguta, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 620.—Am. bor.
atropurpurea, Linn. Sp. Pl. ed. I. 251 = Archangelica atropurpurea.
bracteata, Roxb. in Beats. Tracts, App. 297 = Sium heleneanum.
Breweri, *A. Gray, in Proc. Am. Acad.* vii. (1868) 348.—Calif.
Candollei, Wall. Cat. n. 582 = Selinum Candollii.
Carvifolia, Vill. Prosp. 25; Spreng. Umbellif. Prod. 16 = Selinum Carvifolia.
chacrophyllea, *Lotterm. ex Eysenh. Diss. Accur. Pl. Pruss.* 12.—Livon.
citriodora, *Hance, in Journ. Bot.* ix. (1871) 131.—China.

ANGELICA:—
controversa, *Zumag. Fl. Pedem.* ii. 11.—Italia.
Curtisii, *Buckl. in Am. Journ. Sc.* xlv. (1843) 173.—Am. bor.
dahurica, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 35.—As. or.
Dawsoni, *S. Wats. in Proc. Am. Acad.* xx. (1885) 369.—Am. bor.
decipiens, *Hook. f. Handb. New Zeal. Fl.* 98.—N. Zel.
Dieffenbachii, *Benth. & Hook. f. Gen.* i. 916.—N. Zel.
diversicolor, Dulac, Fl. Hautes-Pyr. 345 = Razulii.
dura, C. Koch, in Linnaea, xvi. (1842) 357 (= *Tommasinia purpurascens*).
ebulifolia, Lap. Hist. Abr. Pl. Pyr. 156 = sylvestris.
Fischeri, Spreng. in Roem. & Schult. Syst. vi. 605 = Silaus longifolius.
flavescens, Hoffm. Gen. Umb. 162 = sylvestris.
Florenti, *Franch. & Sav. ex Maxim. in Bull. Acad. Pétersb.* xix. (1874) 274.—Japon.
geniculata, *Hook. f. Handb. New Zeal. Fl.* 98.—N. Zel.
genuflexa, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 620.—Am. bor.
Gingidium, *Hook. f. Handb. New Zeal. Fl.* 97.—N. Zel.
glauca, *Edgew. in Trans. Linn. Soc.* xx. (1846) 53.—Reg. Himal.
Gmelini, Wormsk. ex DC. Prod. iv. 170 = Archangelica Gmelini.
gracilis, *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) 222.—China.
graveolens, Steud. Nom. ed. II. ii. 555 in syn. = Peucedanum graveolens.
grosseserrata, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 275.—China bor.
hakonensis, *Maxim. l. c.* 277.—Japon.
heteroearpa, *Lloyd, in Bull. Soc. Bot. Fr.* vi. (1859) 709.—Hispan.
hirsuta, Muhl. Cat. 30 = Archangelica hirsuta.
inaequalis, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 184.—Japon.
integrifolia, *Walt. Fl. Carol.* 115.—Carolina.
intermedia, Schult. ex Steud. Nom. ed. I. 49 = Archangelica officinalis.
japonica, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–9) 390.—Japon.
Kiusiana, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 284.—Japon.
laevigata, Fisch. Cat. Hort. Gorenk. (1812) 45 = Peucedanum latifolium.
laevis, *J. Gay, in Ann. Sc. Nat.* Sér. II. vi. (1836) 221 (err. typ. 121).—Hispan.
lancifolia, *Spreng. Syst.* i. 916.—Hab.?
leporina, *S. Wats. in Proc. Am. Acad.* xii. (1877) 252.—Am. bor.
Levisticum, All. Fl. Pedem. ii. 10 = Levisticum officinale.
linearifloba, *A. Gray, in Proc. Am. Acad.* vii. (1868) 347.—Calif.
litoralis, Fries, Fl. Halland. 51 = Archangelica officinalis.
lobata, *Walt. Fl. Carol.* 115.—Am. bor.
lucida, *Linn. Sp. Pl.* 251.—Canada.
Lyallii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 374.—Am. bor.
macrophylla, Schur, Enum. Pl. Transs. 262 = sylvestris.
maculata, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 72.—Ins. Kuril.
major, Gilib. Fl. Lituan. ii. 24 = Archangelica officinalis.
major, *Lag. Gen. et Sp. Nov.* 13.—Hispan.
Maximowiczii, *Benth. ex Maxim. in Bull. Acad. Pétersb.* xix. (1874) 274.—Sibir.
mexicana, *Vatke, App. Sem. Hort. Berol.* (1876) *App.* 2. n. 12.—Mexico.
minor, Gilib. Fl. Lituan. ii. 23 = sylvestris.
Miqueliana, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 276.—Japon.
mongolica, *Franch. in Nouv. Arch. Mus. Par.* Sér. II. vi. (1883) 21; *Pl. Davidian.* i. (1884) 141.—Mongol.
montana, Brot. Fl. Lusit. i. 426 = sylvestris.
nemorosa, Tenore, Fl. Nap. t. 231; Syll. 561 = sylvestris.
norvegica, *Nym. Syll. Suppl.* 26.—Norveg.
officinalis, Bernh. Syst. Verz. Erf. 170 = Peucedanum Ostruthium.

ANGELICA:—
officinalis, Moench, Meth. 81 = Archangelica officinalis.
pachycarpa, *Lange, Descr. Ic. Ill.* 7. t. 9.—Hispan.
pachyptera, Avé-Lall. Ind. Sem. Hort. Petrop. ix. 58 = sylvestris.
paludapifolia, Lam. Fl. Fr. iii. 451 = Levisticum officinale.
palustris, *Hoffm. Gen. Umb.* 162.—Europ.
paniculata, Lam. Encyc. i. 172 = Trochiscanthes nodiflorus.
pinnata, *S. Wats. in King, Rep.* v. 126.—Am. bor.
polyclada, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 86.—Japon.
polymorpha, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 185.—Japon.
pratensis, Bieb. in Fisch. Hort. Gorenk. (1812), ex Bess. Enum. Pl. Volh. 94 = palustris.
pratensis, J. & C. Presl, Fl. Cech. 61 = sylvestris.
pubescens, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 34.—Japon.
procera, Salisb. Prod. 164 = Archangelica officinalis.
pyrenaea, Spreng. Umb. 62 = Selinum pyrenaeum.
Razulii, *Gouan, Illustr.* 13. t. 6.—Pyren.
Razulii, All. Fl. Pedem. ii. 10 = sylvestris.
refracta, *F. Schmidt, Reisen Amurl.* 138.—Ins. Sachalin.
Reuteri, *Boiss. Diagn.* Ser. II. ii. 87.—Hispan.
rosaefolia, *Hook. Ic. Pl.* t. 581.—N. Zel.
ruthenica, Schott, ex Ledeb. Fl. Ross. ii. 296 = sylvestris.
sachalinensis, *Maxim. Prim. Fl. Amur.* 127.—As. or.
sativa, Mill. Gard. Dict. ed. VIII. n. 1 = Archangelica officinalis.
saxatilis, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 729.—Sibir.
scabra, Petit, in Ann. Sc. Obs. i. 99. t. 3 = Xatardia scabra.
scotica, Lam. Encyc. i. 173 = Ligusticum scoticum.
Sieboldi, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 61 (fide Maxim. = A. pubescens, Kiusiana et dahurica).—Japon.
songorica, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 590.—Kokania.
Strattoniana, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 164.—Affghan.
sylvestris, *Linn. Sp. Pl.* 251.—Europ.; Oriens; As. bor.
ternata, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 32.—As. temp.
tomentosa, *S. Wats. in Proc. Am. Acad.* xi. (1876) 141.—Calif.; Am. bor.
Tournefortiana, Cusson, ex Steud. Nom. ed. I. 50 = Razulii.
Traversii, Hook. f. Handb. N. Zeal. Fl. 729 = Aciphylla Traversii.
triloba, Desf. ex Steud. Nom. ed. I. 464, 50 = Siler trilobum.
triquinata, Michx. Fl. Bor. Am. i. 167 = Archangelica atropurpurea.
triquinata, Turcz. ex Trautv. & Mey. in Middend. Reise (Fl. Ochot. 44) = Physolophium saxatile.
ursina, *Regel, in Act. Hort. Petrop.* v. (1877) 590.—Kamtsch.
verticillaris, Linn. Mant. 217 = Peucedanum verticillare.
verticillata, *Geyer, ex Hook. Lond. Journ. Bot.* vi. (1847) 233.—Am. bor.
villosa, Lag. Gen. et Sp. Nov. 12 = sylvestris.
Wheeleri, *S. Wats. in Am. Natural.* vii. (1873) 301.—Amer. bor.

ANGELINA, Pohl, ex Tul. in Arch. Mus. Nat. Par. viii. (1885) 363 = **Siparuna**, Aubl. (Monim.).
divergentifolia, Pohl, ex Tul. l. c. = S. guianensis.

ANGELOCARPA, Rupr. in Mém. Acad. Sc. Pétersb. Sér. VII xiv. (1869) n. iv. (Scrt. Tiansch.) p. 48. *UMBELLIFERAE.*
brevicaulis, *Rupr. l. c.*—As. centr.

ANGELONIA, Humb. & Bonpl. Pl. Aequin. ii. 92. t. 108 (1809). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 930.
PHYLACANTHUS, Benth. Scroph. Ind. 1 (1835).
PHYSIDIUM, Schrad. in Goett. Gel. Anz. i. (1821) 714.
SCHELVERIA, Nees & Mart. in Flora, iv. (1821) 299.

ANGELONIA:—
THYLACANTHA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1822) 45.
angustifolia, *Benth. in DC. Prod.* x. 254.—Mexic.
arguta, *Benth. l. c.* 252.—Bras.
biflora, *Benth. l. c.* 254.—Bras.
bisaccata, *Benth. l. c.* 252.—Bras.
Blanchetii, *Benth. l. c.* 253.—Bras.
campestris, *Nees & Mart. in Flora*, iv. (1821) i. 299.—Bras.
ciliata, Gardn. ex Hook. Bot. Mag. sub t. 3848 = cornigera.
ciliolata, Benth. in DC. Prod. x. 252 = Hookeriana.
Coffreana, Moric. ex Benth. in DC. Prod. x. 252 = procumbens.
cornigera, *Hook. Bot. Mag.* t. 3848.—Bras.
crassifolia, *Benth. in DC. Prod.* x. 252.—Bras.
eriostachys, *Benth. l. c.* 254.—Bras.
Gardneri, *Hook. Bot. Mag.* t. 3754.—Bras.
goyayensis, *Benth. in DC. Prod.* x. 253.—Bras.
grandiflora, *C. Morr. in Ann. Soc. Hort. Gand*, iii. (1847) 93. t. 119.—Bras.
hirta, *Cham. in Linnaea*, viii. (1833) 27.—Bras.
Hookeriana, *Gardn. ex Benth. in DC. Prod.* x. 251.—Bras.
integerrima, *Spreng. Syst.* iv. *Cur. Post.* 235.—Bras.
Leandri, *J. Kickx, in Bull. Acad. Belg.* vi. i. (1839) 309.—Bras.
linearifolia, Moric. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 242 = Blanchetii.
lobelioides, Mart. ex J. A. Schmidt, l. c. 246 = cornigera.
micrantha, *Benth. in DC. Prod.* x. 252.—Bras.
minor, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* v. 33.—Bras.
pilosella, *J. Kickx, in Bull. Acad. Belg.* vi. i. (1839) 309.—Cuba.
pratensis, *Gardn. ex Benth. in DC. Prod.* x. 253.—Bras.
procumbens, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1822) 45.—Bras.
pubescens, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 15.—Bras.
salicariaefolia, *Humb. & Bonpl. Pl. Aequin.* ii. 92. t. 108.—Am. austr.
serrata, *Benth. in DC. Prod.* x. 253.—Bras.
spectabilis, Moric. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 243 = eriostachys.
tomentosa, *Moric. ex Benth. in DC. Prod.* x. 254.—Bras.
veronicoides, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 240 = procumbens.
viscosa, Mart. ex J. A. Schmidt, l. c. 244 = tomentosa.

ANGELOPHYLLUM, Rupr. in Beitr. Pfl. Russ. Reich. xi. (Rev. Umb. Kamtsch.) 8 = **Angelica**, Linn. (Umbell.).
dahuricum, Rupr. l. c. 9 = Angelica sylvestris.
ursinum, Rupr. l. c. 8 = Angelica ursina.

ANGELOPOGON, Poepp. ex Poepp. & Endl. Gen. et (Sp. i. 1 (1835) = **Myzodendron**, Banks & Soland. Santal.).

ANGERVILLA, Neck. Elem. i. 351 (1790) = **Stemodia**, Linn. (Scrophul.).

ANGIANTHUS, Wendl. Coll. ii. 31. t. 48 (1809). *COMPOSITAE*, Benth. & Hook. f. ii. 319.
CEPHALOSORUS, A. Gray, in Hook. Kew Journ. iii. (1851) 152.
CHRYSOCORYNE, Endl. in Bot. Zeit. i. (1843) 457.
CROSSOLEPSIS, Benth. in Enum. Pl. Hueg. 61 (1837).
CYLINDROSORUS, Benth. l. c. 62 (1837).
DITHYROSTEGIA, A. Gray, in Hook. Kew Journ. iii. (1851) 100.
EPITRICHE, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 74.
ERIOCLADIUM, Lindl. Swan Riv. App. 24 (1839).
GAMOZYGIS, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) ii. 75.
HYALOCHLAMYS, A. Gray, in Hook. Kew Journ. iii. (1851) 101.
OGGEROSTYLUS, Cass. in Dict. Sc. Nat. xlix. 221 (1827).
OXEROSTYLUS, Steud. Nom. ed. II. ii. 242 (1841), sphalm.

ANGIANTHUS;—
PHYLLOCALYMMA, Benth. in Enum. Pl. Hueg. 61 (1837).
PIPTOSTEMMA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 191.
PLEUROPAPPUS, F. Muell. in Trans. Vict. Inst. (1855) 37.
POGONOLEPIS, Steetz, in Lehm. Pl. Preiss. i. 440 (1845).
SILOXERUS, Labill. Pl. Nov. Holl. ii. 57. t. 209 (1806), nomen prius.
SKIRRHOPHORUS, DC. ex Lindl. Introd. Nat. Syst. ed. II. 260 (1836).
STYLONCERUS, Spreng. Syst. iii. 356, 451 (1826).
amplexicaulis, *Benth. Fl. Austral.* iii. 568.—Austral.
brachypappus, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 44.—Austral.
codonopappus, *F. Muell. Fragm.* ix. 2.—Austral.
Cunninghamii, *Benth. Fl. Austral.* iii. 565.—Austral.
demissus, *Benth. l. c.* 567.—Austral.
Drummondii, *Benth. l. c.* 566.—Austral.
eriocephalus, *Benth. l. c.* 567.—Austral.
flavescens, Steetz, in Lehm. Pl. Preiss. i. 438 = tomentosus.
globifer, *Benth. Fl. Austral.* iii. 567.—Austral.
humifusus, *Benth. l. c.* 563.—Austral.
microcephalus, *Benth. l. c.* 566.—Austral.
micropoides, *Benth. l. c.* 565.—Austral.
Milnei, *Benth. l. c.* 564.—Austral.
myosuroides, *Benth. l. c.* 563.—Austral.
phyllocephalus, *Benth. l. c.* 565.—Austral.
platycephalus, *Benth. l. c.* 566.—Austral.
pleuropappus, *Benth. l. c.* 563.—Austral.
plumiger, *Benth. l. c.* 568.—Austral.
Preissianus, *Benth. l. c.* 566.—Austral.
pusillus, *Benth. l. c.* 564.—Austral.
pygmaeus, *Benth. l. c.* 567.—Austral.
strictus, *Benth. l. c.* 568.—Austral.
tenellus, *Benth. l. c.* 564.—Austral.
tomentosus, *Wendl. Coll.* ii. 31. t. 48 (1809).—Austral.

ANGIOPETALUM, Reinw. in Syll. Ratisb. ii. (1828) 7. *MYRSINEAE.*
punctatum, *Reinw. l. c.*—Java.

ANGKALANTHUS, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 88. *ACANTHACEAE.*
oligophylla, *Balf. f. l. c.* 89.—Socotra.

ANGOLAEA, Wedd. in DC. Prod. xvii. 300 (1873). *PODOSTEMACEAE*, Benth. & Hook. f. iii. 111.
fluitans, *Wedd. l. c.*—Afr. trop.

ANGOLAM, Adans. Fam. ii. 85 (1763) = **Alangium**, Linn. (Cornac.).

ANGOLAMIA, Scop. Introd. 107 (1777) = praec.

ANGOPHORA, Cav. Ic. iv. 21. t. 338 (1797). *MYRTACEAE*, Benth. & Hook. f. i. 707.
cordifolia, *Cav. l. c.*—Austral.
floribunda, Sweet, Hort. Brit. ed. II. 209 = Eugenia Smithii, Ventenatii.
intermedia, *DC. Prod.* iii. 222.—Austral.
lanceolata, *Cav. Ic.* iv. 22. t. 337.—Austral.
subvelutina, *F. Muell. Fragm.* i. 31.—Austral.
Woodsiana, *F. M. Bailey, in Proc. Linn. Soc. N. S. W.* vi. (1881) 143.—Austral.

ANGORCHIS, Nees, in R. Br. Verm. Schrift. ii. 423, in adnot. (1826) = **Angraecum**, Bory (Orchid.).

ANGOSTURA, Roem. & Schult. Syst. iv. 188 (1819) = **Galipea**, Aubl. (Rutac.).
Cupare, Roem. & Schult. l. c. = G. trifoliata.

ANGOSTYLES, Benth. in Hook. Kew Journ. vi. (1854) 328. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 327.
longifolia, *Benth. l. c.*—Bras.

ANGRAECUM, Bory, Voy. i. 359. t. 19 (1804). *ORCHIDEAE*, Benth. & Hook. f. iii. 583.
AËRANGIS, Reichb. f. in Flora, xlviii. (1865) 190.
AËROBION, Kaempf. ex Spreng. Syst. iii. 679 (1826).
ANGORCHIS, Nees, in R. Br. Verm. Schrift. ii. 423 in adnot. (1826).

ANGRAECUM:—
Listrostachys, Reichb. f. in Bot. Zeit. x. (1852) 930.
Microcaelia, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 285 (1851).
Plectrelminthus, Rafin. Fl. Tellur. iv. 42 (1836).
alcicorne, *Reichb. f. Otia Bot. Hamb.* 117.—Afr. trop.
aphyllum, *Thou. Orch. Afr.* t. 73.—Ins. Masc.
apiculatum, *Hook. Bot. Mag.* t. 4159.—Afr. trop.
arachnopus, *Reichb. f. in Bonplandia*, ii. (1854) 93.—Afr. trop.
arcuatum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 204.—Afr. austr.
arcuatum, Lindl. ex Hook. f. in Bot. Mag. t. 5589 = Chailluanum.
armeniacum, *Lindl. in Bot. Reg.* (1839) *Misc.* 67.—Afr. trop.
articulatum, *Reichb. f. in Gard. Chron.* (1872) 73.—Madag.
ashantense, *Lindl. in Bot. Reg.* (1843) *Misc.* 56.—Afr. trop.
bicaudatum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 205.—Afr. austr.
bilobum, *Lindl. Bot. Reg.* (1840) *Misc.* 69.—Afr. occ.
Boutoni, *Reichb. f. Otia Bot. Hamb.* 117.—Ins. Comor.
bracteosum, *Balf. f. & S. Moore, in Journ. Bot.* xiv. (1876) 293.—Ins. Borbon.
brevifolium, *Lindl. Bot. Reg.* (1840) sub t. 68.—Bras.
Brongniartianum, *Reichb. f. ex Linden, Pescatoria*, i. t. 16.—Ins. Mascar.
Burchellii, *Reichb. f. in Flora*, l. (1867) 117.—Afr. austr.
Calceolus, *Thou. Orch. Afr.* t. 78.—Ins. Mascar.
campyloplectron, *Reichb. f. in Bonplandia*, iii. (1855) 226.—Hab.?
capense, Lindl. Gen. et Sp. Orch. 248 = Mystacidium filicorne.
capitatum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 137.—Afr. trop.
carinatum, Kostel. Allg. Med.-pharm. Fl. i. 250 = Eulophia virens.
carpophorum, *Thou. Orch. Afr.* t. 76.—Afr. trop.
caudatum, *Lindl. Bot. Reg.* t. 1844.—Afr. trop.
caulescens, Thou. Orch. Afr. t. 75 = Mystacidium caulescens.
Chailluanum, *Hook. f. Bot. Mag.* t. 5589.—Afr. trop.
Chiloschistae, *Reichb. f. in Linnaea*, xx. (1847) 678.—Natal.
Christyanum, *Reichb. f. in Gard. Chron.* (1880) i. 806.—Hab.?
citratum, *Thou. Orch. Afr.* t. 61.—Madag.
clandestinum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 206.—Afr. trop.
clavigerum, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 485.—Madag.
conchiferum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 205.—Afr. austr.
cornutum, *Reichb. f. in Flora*, lxviii. (1885) 538.—Ins. Comor.
Cowanii, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 484.—Madag.
crassum, *Thou. Orch. Afr.* tt. 70 et 71.—Madag.
cryptodon, *Reichb. f. in Gard. Chron.* (1883) i. 307.—Madag.
cucullatum, *Thou. Orch. Afr.* t. 48.—Ins. Mascar.
culiciferum, *Reichb. f. in Flora*, lxviii. (1885) 538.—Ins. Comor.
descendens, *Reichb. f. in Gard. Chron.* (1882) i. 558.—Hab.?
distichum, *Lindl. Bot. Reg.* t. 1781.—Afr. trop.
eburneum, Lindl. l. c. (1832) xviii. 1522 = superbum.
eburneum, *Bory, Voy.* i. 359. t. 19.—Ins. Borbon.
elatum, Thou. Orch. Afr. t. 79 = Cryptopus elatus.
Ellisii, *Reichb. f. in Flora*, lv. (1872) 278.—Madag.
expansum, *Thou. Orch. Afr.* t. 57.—Ins. Mascar.
falcatum, *Lindl. Coll. Bot.* 15.—China.
fasciola, *Lindl. Bot. Reg.* xxvi. (1840) sub t. 68.—Demerara.
fastuosum, *Reichb. f. in Gard. Chron.* (1881) ii. 748, 844.—Madag.
filicorne, *Thou. Orch. Afr.* t. 52.—Ins. Mascar.; Madag.
filiforme, *Lindl. Gen. et Sp. Orch.* 248.—Ind. occ.
floribundum, *Reichb. f. in Gard. Chron.* (1885) i. 787; *in Flora*, lxviii. (1885) 380.—Ins. Comor.
fragrans, *Thou. Orch. Afr.* t. 54.—Ins. Mascar.
funale, Lindl. in Gard. Chron. (1846) 135 = Dendrophylax funalis.

ANGRAECUM:—
fuscatum, *Reichb. f. in Gard. Chron.* (1882) ii. 488.—Madag.
Galeandrae, *Reichb. f. in Flora*, xlviii. (1865) 189.—Afr. occ.
Gilpinae, *Reichb. f. & S. Moore, in Journ. Linn. Soc.* xvi. (1877) 206.—Madag.
gladiifolium, *Thou. Orch. Afr.* t. 53.—Ins. Mascar.; Madag.
globulosum, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 285 = Guyonianum.
glomeratum, *Ridl. in Gard. Chron.* (1885) ii. 678.—Afr. trop.
gracile, *Thou. Orch. Afr.* t. 77.—Ins. Mascar.
Grantii, Bateman, ex Baker, in Trans. Linn. Soc. xxix. (1875) 153 = Kotschyi.
Guyonianum, *Reichb. f. in Linnaea*, xxii. (1849) 865.—Abyss.
Hildebrandtii, *Reichb. f. in Gard. Chron.* (1878) i. 725.—Afr. trop.
Humblotii, *Reichb. f. in Flora*, lxviii. (1885) 381.—Ins. Comor.
hyaloides, *Reichb. f. in Gard. Chron.* (1880) i. 264.—Madag.
ichneumoneum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 136.—Afr. occ.
imbricatum, *Lindl. l. c.* 137.—Afr. occ.
implicatum, *Thou. Orch. Afr.* t. 58.—Madag.
inapertum, Thou. l. c. t. 50 = Mystacidium inapertum.
infundibulare, *Lindl. in Journ. Linn. Soc.* vi. (1862) 136.—Afr. occ.
jamaicense, *Reichb. f. & Wullschl. in Walp. Ann.* vi. 901.—Jamaic.
japonicum, *Linden, Illustr. Hortic.* xxviii. (1881) 64.—Japon.
Kotschyi, *Reichb. f. in Gard. Chron.* (1880) ii. 456.—Afr. trop.
Lansbergii, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 316.—Caracas.
Leonis, Hort.—Cf. Gard. Chron. (1885) ii. 80–81 = Aëranthus Leonis.
lepidotum, *Reichb. f. l. c.* (1880) i. 806, in obs.—Hab.?
Lindenii, *Lindl. l. c.* (1846) 135.—Cuba.
luridum, *Lindl. ex Steud. Nom.* ed. II. i. 98.—Afr. trop.
maculatum, Lindl. Coll. 15 = Eulophia maculata.
maxillarioides, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 479.—Madag.
megalorrhizum, *Reichb. f. Otia Bot. Hamb.* 117.—Afr. trop.
micranthum, *Lindl. Bot. Reg.* xxi. t. 1772.—Afr. trop.
modestum, *Hook. f. Bot. Mag.* t. 6693.—Madag.
monodon, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 102.—Afr. occ.
monophyllum, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 58 = Eulophia monophylla.
multiflorum, Thou. Orch. Afr. t. 74 = Mystacidium caulescens.
Mystacidii, *Reichb. f. in Linnaea*, xx. (1847) 677.—Natal.
odoratissimum, *Reichb. f. in Bonplandia*, iv. (1856) 326.—Hab.?
organense, *Reichb. f. in Walp. Ann.* vi. 901.—Bras.
ornithorhynchum, *Lindl. Bot. Reg.* xxvi. (1840) sub t. 68.—Bras.
palmatum, Thou. Orch. Afr. t. 69 = palmiforme.
palmiforme, *Thou. l. c.* 68.—Ins. Mascar.
parviflorum, *Thou. l. c.* t. 60.—Ins. Mascar.
parvulum, *Ayres, ex Baker, Fl. Maurit.* 357.—Ins. Mascar.
pectinatum, *Thou. Orch. Afr.* t. 51.—Ins. Mascar.; Madag.
pellucidum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 134.—Afr. occ.
pertusum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 205.—Afr. occ.
Pescatorianum, *Lindl. in Journ. Hort. Soc.* iv. (1849) 263.—Ins. Borbon.
physophorum, *Reichb. f. Otia Bot. Hamb.* (1881) 77 et 449.—Ins. Mascar.
Poeppigii, *Reichb. f. in Linnaea*, xxii. (1849) 858.—Cuba.
polystachyum, *Lindl. Bot. Reg.* xxvi. (1840) sub t. 68.—Peruv.
polystachyum, *A. Rich. in Mém. Hist. Nat. Par.* iv. (1828) 66 = Aëonia polystachya.
pugioniforme, *Klotzsch, Ind. Sem. Hort. Berol. App.* (1851) 12.—Venezuela.

ANGRAECUM:—
pusillum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 205.—Afr. austr.
pygmaeum, *Linden, Illustr. Hortic.* xxviii. (1881) 64.—Japon.
ramosum, *Thou. Orch. Afr.* t. 59.—Ins. Mascar.
rectum, *Thou. l. c.* 55.—Ins. Mascar.
recurvum, *Thou. l. c.* t. 56.—Ins. Mascar.
rhipsalisocium, *Reichb. f. in Flora*, xlviii. (1865) 189.—Afr. occ.
Rohrii, *Reichb. f. Otia Bot. Hamb.* 117.—Abyss.
rostellare, *Reichb. f. in Flora*, lxviii. (1885) 380; *et in Gard. Chron.* (1885) i. 726.—Ins. Comor.
rostratum, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 485.—Madag.
Rutenbergianum, *Kraentz. in Bremen Abh.* vii. (1882) 257.—Afr. trop.
sacciferum, *Lindl. in Hook. Comp. Bot. Mag.* ii. (1836) 205.—Afr. austr.
Schiedei, Reichb. f. in Linnaea, xxii. (1849) 857 = Campylocentron Schiedei.
Schimperianum, *Reichb. f. in Walp. Ann.* iii. 573.—Abyss.
Scottianum, *Reichb. f. in Gard. Chron.* (1878) ii. 556.—Ins. Comor.
Sedeni, Hort. ex Nichols. Dict. Gard. i. 79 = arcuatum.
Sellowii, *Reichb. f. in Linnaea*, xxii. (1849) 857.—Bras.
sesquipedale, *Thou. Orch. Afr.* tt. 66 et 67.—Madag.
spathulatum, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 478.—Madag.
striatum, *Thou. Orch. Afr.* t. 72.—Ins. Mascar.
subulatum, *Lindl. in. Hook. Comp. Bot. Mag.* ii. (1836) 206.—Afr. occ.
superbum, *Thou. Orch. Afr.* tt. 62 et 64.—Madag.
tenue, *Lindl. Bot. Reg.* xxvi. (1840) sub t. 68.—Bras.
teretifolium, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 484.—Madag.
tridens, *Lindl. l. c.* vi. (1862) 135.—Afr. trop.
tridentatum, *Harv. Thes. Cap.* ii. 6.—Afr. austr.
triquetrum, *Thou. Orch. Afr.* t. 49.—Ins. Mascar.
vagans, *Lindl. in Journ. Linn. Soc.* vi. (1862) 135.—Afr. occ.
virens, Lindl. Bot. Reg. (1847) t. 19 = eburneum.
vesicatum, *Lindl. l. c.* (1843) *Misc.* 6.—Afr. trop.
Weigelti, *Reichb. f. in Linnaea*, xxii. (1849) 857.—Guiana.
xylopus, *Reichb. f. in Flora*, lxviii. (1885) 538.—Ins. Comor.
yuccaefolium, *Boj. Hort. Maurit.* 318.—Ins. Maurit.
zeylanicum, *Lindl. in Journ. Linn. Soc.* iii. (1859) 40.—Ind. or.

ANGUILLARIA, R. Br. Prod. 273 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 824.
australis, F. Muell. Fragm. vii. 74 = dioica.
biglandulosa, R. Br. Prodr. 273 = dioica.
capensis, Spreng. Syst. ii. 146 = Androcymbium lencanthum.
caricina, *Schult. f. Syst.* vii. 1539.—Ind. or.
cochinchinensis, *Spreng. Syst.* ii. 147.—Cochinch.
densiflora, *Benth. Fl. Austral.* vii. 29.—Austral.
dioica, *R. Br. Prod.* 273.—Austral.
Heyneana, Wall. Cat. n. 5086 = Iphigenia indica.
indica, R. Br. Prod. 273 = Iphigenia indica.
monantha, Endl. in Lehm. Pl. Preiss. ii. 45 = dioica.
pygmaea, Endl. l. c. = Wurmbea pygmaea.
racemosa, Schult. f. Syst. vii. 1539 = Iphigenia indica.
tenella, Endl. in Lehm. Pl. Preiss. ii. 45 = Wurmbea tenella.
uniflora, R. Br. Prod. 273 = dioica.

ANGUILLARIA, Gaert. Fruct. i. 372 (1788) = Ardisia. Sw. (Myrsin.).
bahamensis, Gaertn. l. c. 373 = Ar. excelsa.
Barthesia, Lam. Illustr. ii. 109 = Ar. Barthesia.
chrysophylla, Lam. l. c. = Ar. serrulata.
coriacea, Poir. Encyc. vii. 685 = Ar. coriacea.
crenulata, Poir. l. c. 688 = Parathesis crenulata.
excelsa, Lam. Illustr. ii. 110 = Ar. excelsa.
Icacorea, Lam. l. c. 109 = Ar. acuminata.
lateriflora, Lam. l. c. = Ar. lateriflora.
latifolia, Steud. Nom. ed. II. i. 98 = Ar. laurifolia.
laurifolia, Lam. Illustr. ii. 109 = Ar. laurifolia.
parasitica, Lam. l. c. = Grammadenia parasitica.
pyramidalis, Cav. Ic. vi. 1 = Ar. pyramidalis.
serrata, Cav. l. c. 2 = Ar. Cavanillesii.
serrulata, Poir. Encyc. vii. 686 = Ar. serrulata.
solanacea, Poir. l. c. 688 = Ar. humilis.

ANGUILLARIA :—
tinifolia, Lam. Illustr. ii. 109 = Ardisia tinifolia.
zeylanica, Gaertn. Fruct. i. 373. t. 77. f. 1 = Ar. Moonii.

ANGUINA, Mich. ex Linn. Syst. ed. I. (1735) = **Trichosanthes**, Linn. (Cucurb.).

ANGUINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. 160 (1869) = **Allium**, Tourn. (Liliac.).
Victorialis, Fourr. l. c. = Allium Victorialis.

ANGULOA, Ruiz & Pav. Prod. Fl. Per. 118. t. 26 (1794). *ORCHIDEAE*, Benth. & Hook. f. iii. 548.
Clowesii, *Lindl. Bot. Reg.* (1844) *Misc.* 26.—Venezuela.
dubia, *Reichb. f. in Gard. Chron.* (1882) I. 764.—N. Granat.
eburnea, *Hort. ex Nichols. Dict. Gard.* i. 79.—N. Granat.
grandiflora, H. B. & K. Nov. Gen. et Sp. i. 345 = Stanhopea Bucephalus.
Hernandezii, Kunth, Syn. i. 332 = Stanhopea tigrina.
Hohenlohii, *Morr. in Belg. Hortic.* iii. (1853) 201. t. 31.—N. Granat.
lurida, Link, in Preuss. Gart. Ver. i. 289. t. 6 = Catasetum luridum.
media ×, *Reichb. f. in Gard. Chron.* (1881) II. 38.
purpurea, *Linden, in Illustr. Hortic.* xxviii. (1881) t. 427.—Venezuela.
Ruckeri, *Lindl. Bot. Reg.* (1846) t. 41.—N. Granat.
squalida, *Poepp. & Endl. Nov. Gen. et Sp.* i. 74.—Peruv.
superba, H. B. & K. Nov. Gen. et Sp. i. 343. t. 93 = Acineta Humboldtii.
uniflora, *Ruiz & Pav. Syst.* 228.—Peruv.
virginalis, Hort.—Cf. Gard. Chron. (1851) 392 = uniflora.

ANGURIA, [Tourn.] Linn. Syst. ed. I. (1735). *CUCURBITACEAE*, Benth. & Hook. f. i. 833.
aculeolata, Schlecht. in Linnaea, xxiv. (1851) 748 = Wilbrandia verticillata.
affinis, Schlecht. l. c. 760 = pedata.
Arrabidae, Schlecht. l. c. 736 = Gurania multiflora.
bignoniacea, Poepp. & Endl. Nov. Gen. et Sp. ii. 53 = triphylla, Gurania cissoides.
Boissieriana, *Cogn. in DC. Monog. Phan.* iii. 673.—Peruv.
capitata, Poepp. & Endl. Nov. Gen. et Sp. ii. 52 = Gurania capitata.
cissoides, Benth. in Hook. Kew Journ. ii. (1850) 242 = Gurania cissoides.
Citrullus, Mill. Gard. Dict. ed. VIII. add. = Citrullus vulgaris.
dentata, Schlecht. in Linnaea, xxiv. (1851) 713 = pedata.
dubia, Hook. & Arn. Bot. Beech. Voy. 292 = Cayaponia attenuata.
eriantha, Poepp. & Endl. Nov. Gen. et Sp. ii. 52. t. 169 = Gurania eriantha.
eriantha, Seem. Bot. Voy. Herald, 127 = Gurania Seemanniana.
Gaudichaudiana, Schlecht. in Linnaea, xxiv. (1851) 753 = Wilbrandia verticillata.
glomerata, Eggers, Fl. St. Croix & Virgin. Isl. 55 = Corallocarpus glomeratus.
grandiflora, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1877).—Bras.
guianensis, Klotzsch, in Schomb. Reis. Brit. Guian. iii. 988 = Gurania guianensis.
heterophylla, Willd. ex Schlecht. in Linnaea, xxiv. (1851) 772 = Gurania cissoides.
integrifolia, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1824) 9.—Bras.
indica, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 6.—Malaya.
Jacquiniana, Schlecht. in Linnaea, xxiv. (1851) 708 = trilobata.
Kegeliana, Schlecht. l. c. 776 = Gurania Kegeliana.
Kunthiana, *Schlecht. l. c.* 762.—Bras.
laciniosa, Schlecht. l. c. 755 = Apodanthera laciniosa.
leptantha, Schlecht. l. c. 779 = Helmontia leptantha.
lobata, *Linn. Amoen. Acad.* viii. 263.—Guian.
longipedunculata, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1877).—Mexic.
Makoyana, Lem. in Fl. des Serres, Sér. I. iii. (1847) 222 = Gurania Makoyana.

ANGURIA :—
Meyeniana, Schlecht. in Linnaea, xxiv. (1851) 750 = Wilbrandia verticillata.
multiflora, Miq. in Linnaea, xviii. (1846) 366 = Gurania multiflora.
obtusiloba, Schlecht. in Linnaea, xxiv. (1851) 744 = (Vitis sp.).
Ottoniana, *Schlecht. l. c.* 758.—Cuba.
pallida, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1877) 10, 22.—Ecuador.
pedata, *Jacq. Enum. Pl. Carib.* 3.—Ins. S. Doming.
pedata, Willd. ex Cogn. in DC. Monog. Phan. iii. 828 = Cyclanthera leptostachya.
pedatisecta, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 10 = Cyclanthera pedata.
pedisecta, *Nees & Mart. l. c.* = Apodanthera pedisecta.
Plumieriana, *Schlecht. in Linnaea*, xxiv. (1851) 709.—Ins. Martinic.
Poeppigiana, Schlecht. l. c. 769 = Dieudonnaea rhizantha.
polyanthos, Klotzsch, in Schomb. Reis. Brit. Guian. iii. 988 = Gurania Klotzschiana, G. subumbellata.
polyphyllos, Schlecht. in Linnaea, xxiv. (1851) 712 = pedata.
rhizantha, Poepp. & Endl. Nov. Gen. et Sp. ii. 52. t. 171 = Dieudonnaea rhizantha.
rosea, H. B. & K. Nov. Gen. et Sp. ii. 122 = trilobata.
Schomburgkiana, *Schlecht. in Linnaea*, xxiv. (1851) 765.—Guiana.
Sellowiana, Schlecht. l. c. 746 = Gurania Sellowiana.
Sellowiana, Wawra, Bot. Maximil. Reise, i. 57. t. 52 = Gurania Wawraei.
sinuata, Benth. in Hook. Kew Journ. ii. (1850) 242 = Gurania sinuata.
speciosa, Poepp. & Endl. Nov. Gen. et Sp. ii. 51. t. 169 = Gurania speciosa.
spinulosa, Poepp. & Endl. l. c. 52. t. 170 = Gurania spinulosa.
subumbellata, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) (Stirp. Surin. Select.) 103 = Gurania subumbellata.
ternata, *M. Roem. Syn. Pepon.* i. 26.—Bras.
trifoliata, Aubl. Pl. Guian. ii. 842 = Gurania cissoides.
trifoliata, *Linn. Sp. Pl.* ed. II. 1376.—Ins. S. Doming.
trifoliata, Vell. Fl. Flum. x. t. 2 = ternata.
trilobata, *Jacq. Enum. Pl. Carib.* 31.—N. Granat.
trilobata, Vell. Fl. Flum. x. t. 1 = Gurania multiflora.
triphylla, Klotzsch, in Schomb. Reis. Brit. Guian. iii. 988 = Schomburgkiana.
triphylla, *Miq. in Linnaea*, xix. (1847) 136.—Guian.
umbrosa, *H. B. & K. Nov. Gen. et Sp.* ii. 121.—Mexic.
Wageneriana, Schlecht. in Linnaea, xxiv. (1851) 785 = Gurania Wageneriana.
Warmingiana, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxvii. (1877) 21.—Bras.
Warmingii, *Cogn. l. c.*—Bras.
Warscewicziana, Hort. Belg. ex Cogn. l. c. 18 = seq.
Warscewiczii, *Hook. f. in Bot. Mag.* t. 5304.—Mexic.

ANGUSTINEA, A. Gray, Bot. U. St. Expl. Exped. i. 593, in syn. (1854) = AUGUSTINEA, St. Hil. & Naud. = **Miconia**, Ruiz & Pav. (Melast.).

ANHALONIUM, Lem. Cact. Nov. Gen. et Sp. 1. (1839) = **Mammillaria**, Haw. (Cact.).
areolosum, Lem. Illustr. Hortic. vi. (1859) Misc. 35.—Mexic.
elongatum, Salm-Dyck, Cact. Hort. Dyck. (1845) 15 = M. elongata.
Engelmanni, Lem. Cact. 42 = M. fissurata.
fissipedum, Monv. Cat. ex Forst. Cact. Handb. ed. II. i. 232 = sulcatum.
fissuratum, Engelm. Cact. Mex. Bound. 75 = M. fissurata.
Kotchubeyi, Lem. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 5 = Mammillaria sulcata.
Kotschoubeyanum, Lem. Cact. 41; Illustr. Hortic. xvi. (1869) Misc. 72.—Mexic.
prismaticum, Lem. Cact. Nov. Gen. et Sp. 1 = M. prismatica.
pulvilligerum, Lem. Illustr. Hortic. xvi. (1869) 72; Hort. Monv. i. 275 = Mammillaria elongata.

ANHALONIUM :—
retusum, Salm-Dyck, Cact. Hort. Dyck. (1845) 15 = Mammillaria prismatica.
sulcatum, Salm-Dyck, l. c. ed. II. 5, 78.—Mexic.
Williamsii, Lem. ex Forst. Handb. Cact. ed. II. i. 233 = Echinocactus Williamsii.

ANIA, Lindl. in Wall. Cat. n. 3740–41; et Gen. et Sp. Orch. 129 (1831) = **Tainia**, Blume (Orchid.).
angustifolia, Lindl. in Wall. Cat. n. 3740.—Ind. or.; China.
bicornis, Lindl. in Bot. Reg. (1842) Misc. 37 = T. bicornis.
latifolia, Lindl. in Wall. Cat. n. 3741 = T. latifolia.
latifolia, Wight, Ic. Pl. iii. 914 = T. bicornis?
maculata, Thw. Enum. Pl. Zeyl. 301 = (Chrysoglossi sp.).—Zeylan.

ANIBA, Aubl. Pl. Gui. i. 327 (1775) = **Ocotea**, Aubl. (Laurin.).
guianensis, Aubl. l. c. = O. commutata.

ANICTOCLEA, Nimmo, in J. Grah. Cat. Pl. Bomb. 252 (1839) = **Tetrameles**, R. Br. (Datisc.).
Grahamiana, Nimmo, l. c. = T. nudiflora.

ANIDRUM, Neck. Elem. i. 188 (1790) = **Bifora**, Hoffm. (Umbell.).

ANIGOSIA, Salisb. in Trans. Hort. Soc. i. (1812) 327 = **Anigozanthos**, Labill. (Haemodor.).
flavida, Salisb. l. c. = Anigozanthos flavida.

ANIGOZANTHOS, Labill. Voy. i. 409. t. 22 (1798). *HAEMODORACEAE*, Benth. & Hook. f. iii. 676.
ANIGOSIA, Salisb. in Trans. Hort. Soc. i. (1812) 327.
SCHWAEGRICHENIA, Spreng. Syst. ii. 26 (1825).
bicolor, *Endl. in Lehm. Pl. Preiss.* ii 26.—Austral.
coccinea, Paxt. Bot. Mag. v. 271 = flavida.
flavescens, Steud. Nom. ed. II. i. 98 = flavida.
flavida, [*DC. in*] *Red. Lil.* t. 176.—Austral.
fuliginosa, Hook. Bot. Mag. t. 4291 = Macropodia fumosa.
grandiflora, Salisb. Parad. Lond. t. 97 = flavida.
humilis, *Lindl. Swan Riv. App.* 46. t. 6 B.—Austral.
Manglesii, *D. Don, in Sweet, Brit. Fl. Gard.* Ser. II. t. 265.—Austral.
Manglesii, Maund, Botanist, t. 67 = flavida.
minima, Lehm. Pl. Preiss. ii. 274 = Preissii.
Preissii, *Endl. in Lehm. l. c.* 26.—Austral.
pulcherrima, *Hook. Bot. Mag.* t. 4180.—Austral.
rufa, *Labill. Voy.* i. 409. t. 22.—Austral.
tyrianthina, Hook. Bot. Mag. t. 4507 = rufa.
viridis, *Endl. in Lehm. Pl. Preiss.* ii. 25.—Austral.

ANILEMA, Kunth, Enum. Pl. iv. 64 (1843) = **Aneilema**, R. Br. (Commel.).

ANIOTUM, Soland. ex Endl. Gen. 332 (1837) = **Inocarpus**, Forst. (Legum.).

ANISACANTHA, R. Br. Prod. 410 (1810). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 62.
bicornis, F. Muell. Fragm. vii. 14 = Sclerolaena bicornis.
bicuspis, *F. Muell. in Trans. Vict. Inst.* (1855) 133.—Austral.
Birchii, *F. Muell. Fragm.* viii. 163.—Austral.
brevicuspis, *F. Muell. l. c.* iv. 150.—Austral.
diacantha, Nees, in Lehm. Pl. Preiss. i. 635 = Sclerolaena diacantha.
divaricata, *R. Br. Prod.* 410.—Austral.
Drummondii, *Benth. Fl. Austral.* v. 199.—Austral.
echinopsila, *F. Muell. Fragm.* vii. 14.—Austral.
erinacea, *Moq. in DC. Prod.* xiii. II. 122.—Austral.
glabra, *F. Muell. ex Benth. Fl. Austral.* v. 200.—Austral.
gracilicuspis, *F. Muell. Fragm.* ii. 170.—Austral.
incana, A. Cunn. ex Moq. in DC. Prod. xiii. II. 122 = muricata.
kentropsidea, F. Muell. in Trans. Vict. Inst. (1855) 133 = Sclerolaena diacantha.
lanicuspis, F. Muell. Fragm. ii. 170 = Sclerolaena lanicuspis.
muricata, *Moq. Chenop. Enum.* 84.—Austral.
quinquecuspis, Muell. in Trans. Vict. Inst. (1885) 134 = muricata.
tricuspis, *F. Muell. l. c.* 133.—Austral.

ANISACANTHUS, Nees, in Linnaea, xvi. (1842) 307. *ACANTHACEAE*, Benth. & Hook. f. ii. 1117.
BIRNBAUMIA, Kostel. Ind. Pl. Hort. Prag. (1844) 21.
Greggii, *A. Gray, Synop. Fl. N. Am.* ii. 328.—Mexic.
juncea, *Hemsl. Biol. Centr. Am. Bot.* 522.—Mexic.
pumilus, *Nees, in DC. Prod.* xi. 445.—Mexic.
Thurberi, *A. Gray, Synop. Fl. N. Am.* ii. 328.—Am. bor.
virgularis, *Nees, in DC. Prod.* xi. 445.—Mexic.
Wrightii, *A. Gray, Synop. Fl. N. Am.* ii. 328.—Am. bor.

ANISACTIS, Dulac, Fl. Hautes-Pyr. 347 (1867) = **Carum**, Linn. (Umbell.).
segetalis, Dulac, l. c. = C. Petroselinum.

ANISADENIA, Wall. Cat. n. 1510 (1830). *LINEAE*, Benth. & Hook. f. i. 243.
khasyana, Griff. Notul. iv. 534 = saxatilis.
pubescens, *Griff. l. c.* 535.—Reg. Himal.
saxatilis, *Wall. Cat.* n. 1510.—Reg. Himal.

ANISANDRA, Bartl. in Lehm. Pl. Preiss. i. 361 (1845) = **Microcorys**, R. Br. (Labiat.).
glabra, Bartl. l. c. = M. glabra.

ANISANDRA, Planch. ex Oliver, Fl. Trop. Afr. i. 348 in syn. (1868) = **Ptychopetalum**, Benth. (Olacin.).
nigrescens, Planch. l. c. = P. anceps.

ANISANTHA, C. Koch, in Linnaea, xxi. (1848) 394 = **Bromus**, Linn. (Gramin.).
pontica, C. Koch, l. c. = B. tectorum.

ANISANTHERA, Griff. Notul. iv. 100 (1854) = **Adenosma**, R. Br. (Scrophul.).

ANISANTHERA, Rafin. Fl. Tellur. iii. 80 (1836) = **Caccinia**, Savi (Borag.).
ciliata, Rafin. l. c. = C. Rauwolfii.

ANISANTHERA, Rafin. Fl. Tellur. ii. 60 (1836) = **Crotalaria**, Tourn. (Legum.).
hastata, Rafin. l. c. 61 = C. verrucosa.
versicolor, Rafin. l. c. 60 = C. verrucosa.

ANISANTHUS, Sweet, Hort. Brit. ed. I. 396 (1827) = **Antholyza**, Linn. (Irid.).
abyssinicus, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 344 = Antholyza abyssinica.
caryophyllaceus, Klatt, l. c. = Watsonia humilis.
Cunonia, Sweet, Hort. Brit. ed. I. 397 = Anth. Cunonia.
huillensis, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 344.—Afr. trop.
quadrangularis, Sweet, Hort. Brit. ed. I. 397 = Anth. quadrangularis.
saccatus, Klatt, in Linnaea, xxxv. (1867–68) 300 = Anth. saccata
splendens, Sweet, Brit. Flow. Gard. Ser. II. i. t. 84 = Anth. caffra.

ANISANTHUS, Willd. ex Roem. & Schult. Syst. v. p. xiv (1819) = **Symphoricarpos**, Juss. (Caprifol.).
microphyllus, Willd. ex Roem. & Schult. l. c. 223 = S. microphyllus.

ANISEIA, Choisy (Conv. Or. 99) in Mém. Soc. Genèv. vi. (1833) 481; viii. (1838) 65 = **Ipomoea**, Linn. (Convolv.).
Afzelii, G. Don, Gen. Syst. iv. 295 = Hewittia bicolor.
arifolia, Meissn. in Mart. Fl. Bras. vii. 319.—Bras.
aurea, Kellogg, in Proc. Calif. Acad. v. (1873) 83.—Calif.
barlerioides, Choisy, Conv. Or. 102 = I. barlerioides.
biflora, Choisy, l. c. 101 = I. biflora.
bracteata, Hassk. Pl. Rar. Jav. 516 = Hewittia bicolor.
calycina, Choisy, Conv. Or. 100 = I. calycina.
calystegioides, Choisy, in DC. Prod. ix. 431 = I. crassipes.
carnea, Griseb. Fl. Brit. W. Ind. 471, *sphalm.* = cernua.
cernua, Moric. N. Pl. Am. t. 38.—Bras.
costata, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 367.—Chili.
diversifolia, Walp. l. c.—Chili.

ANISEIA:—
emarginata, Hassk. Cat. Hort. Bog. Alt. 139 = Ipomoea martinicensis.
ensifolia, Choisy, Conv. Rar. 144 = I. martinicensis.
fulvicaulis, Hochst. ex Choisy, in DC. Prod. ix. 431.—Abyss.
gracillima, Choisy, l. c. 430.—Bras.
hastata, Meissn. in Mart. Fl. Bras. vii. 319 = I. phylloneura.
heterantha, Choisy, in DC. Prod. ix. 430.—Bras.
heterophylla, Meissn. in Mart. Fl. Bras. vii. 321 = I. heterophylla.
martinicensis, Choisy, Conv. Rar. 144 = I. martinicensis.
Medium, Choisy, Conv. Or. 100 = I. denticulata.
nitens, Choisy, Conv. Rar. 145.—Bras.
salicifolia, Choisy, l. c. 144 = I. martinicensis.
tomentosa, Meissn. in Mart. Fl. Bras. vii. 321 = I. martinicensis.
uniflora, Choisy, Conv. Or. 101 = I. martinicensis.
Velloziana, Choisy, in DC. Prod. ix. 430.—Bras.

ANISEION, St. Lag. in Ann. Soc. Bot. Lyon, vii. 119 (1880) = praec.

ANISEPTA, Rafin. Cat. 13 (1824) = **Croton**, Linn. (Euphorb.).
graveolens, Rafin. l. c.—Am. bor.

ANISOCALYX, Hance, in Walp. Ann. iii. 195 (1853) = **Herpestis**, Gaertn. (Scrophul.).
limnanthiflorus, Hance, l. c. = H. Monniera.

ANISOCARPUS, Nutt. in. Trans. Am. Phil. Soc. Ser. II. vii. (1841) 388 = **Madia**, Molina (Compos.).
Bolanderi, A. Gray, in Proc. Am. Acad. vii. (1868) 360 = M. Bolanderi.
madioides, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 388 = M. Nuttallii.

ANISOCENTRA, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 593 = **Tropaeolum**, Linn. (Geran.).
cardiopetala, Turcz. l. c. = T. speciosum.

ANISOCENTRUM, Turcz. in Bull. Soc. Nat. Mosc. (1862) i. 322 = **Acisanthera**, P. Br. (Melast.).
Gardneri, Turcz. l. c. 323 = Ac. punctatissima.

ANISOCHAETA, DC. Prod. v. 109 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 493.
mikanioides, *DC. l. c.* 110.—Afr. austr.

ANISOCHILUS, Wall. Pl. As. Rar. ii. 18 (1831). *LABIATAE*, Benth. & Hook. f. ii. 1177.
adenanthus, *Dalz. & Gibs. Bomb. Fl.* 206.—Ind. or.
albidus, Wight, Ic. iv. t. 1435 = dysophylloides.
carnosus, *Wall. Pl. As. Rar.* ii. 18.—Ind. or.
crassus, Benth. in DC. Prod. xii. 81 = carnosus.
decussatus, Dalz. & Gibs. Bomb. Fl. 206 = eriocephalus.
dysophylloides, *Benth. in Wall. Pl. As. Rar.* ii. 19.—Ind. or.
eriocephalus, *Benth. in DC. Prod.* xii. 81.—Ind. or.
euneurus, *Miq. Fl. Ind. Bat.* ii. 957.—Java.
glaber, Schrad. Ind. Sem. Hort. Gott. (1833) 1 = carnosus.
pallidus, *Wall. Pl. As. Rar.* ii. 18.—Ind. or.
paniculatus, *Benth. in DC. Prod.* xii. 82.—Ind. or.
plantagineus, *Hook. f. Fl. Brit. Ind.* iv. 628.—Ind. or.
polystachys, *Benth. in Wall. Pl. As. Rar.* ii. 19.—Ind. or.
purpureus, Wight, Ic. iv. t. 1435 = dysophylloides.
robustus, *Hook. f. Fl. Brit. Ind.* iv. 629.—Ind. or.
rupestris, Wight, ex Hook. f. Fl. Brit. Ind. iv. 627 = carnosus.
scaber, *Benth. in DC. Prod.* xii. 81.—Ind. or.
sericeus, *Benth. l. c.* 82.—Ind. or.
sinensis, *Hance, in Journ. Bot.* xxiii. (1885) 327.—China.
suffruticosus, *Wight, Ic.* iv. t. 1437.—Ind. or.
suffruticosus, Thw. Enum. Pl. Zeyl. 238 = paniculatus?
verticillatus, *Hook. f. Fl. Brit. Ind.* iv. 629.—Ind. or.
Wightii, *Hook. f. l. c.* 628.—Ind. or.

ANISOCOMA, Torr. & Gray, in Bost. Journ. Nat. Hist. v. (1844) 111. t. 13. *COMPOSITAE*, Benth. & Hook. f. ii. 518.
PTEROSTEPHANUS, Kellogg, in Proc. Calif. Acad. iii. (1863) 20. f. 4.
acaulis, *Torr. & Gray, l. c.*—Calif.

ANISODENS, Dulac, Fl. Hautes-Pyr. 465 (1867) = **Scabiosa**, Tourn. (Dipsac.).
arvensis, Dulac, l. c. = S. arvensis.
longifolius, Dulac, l. c. = S. sylvatica.
sylvaticus, Dulac, l. c. = S. sylvatica.

ANISODERIS, Cass. in Dict. Sc. Nat. xlviii. 429 (1827) = **Crepis**, Linn. (Compos.).
foetida, Fisch. & Mey. Ind. Sem. Petrop. iv. 32 = C. foetida.
rhoeadifolia, Fisch. & Mey. l. c. = C. foetida.
rubra, Cass. in Dict. Sc. Nat. xlviii. 429 = C. rubra.

ANISODONTEA, Presl, Bot. Bemerk. 18 (1844) (in Abh. Boehm. Ges. Fol. V. iii.) = **Sphaeralcea**, St. Hil. (Malvac.).
Dregeana, Presl, l. c. = S. Dregeana.

ANISODUS, Link & Otto, Ic. Sel. Pl. 77. t. 35 (1824) = **Scopola**, Jacq. (Solanac.).
humilis, Hook. f. ex C. B. Clarke, in Fl. Brit. Ind. iv. 242 = Mandragora caulescens.
luridus, Link & Otto, Ic. Sel. Pl. 77 = S. lurida.
stramonifolius, G. Don, in Loud. Hort. Brit. 61 = S. lurida.

ANISOLEPIS, Steetz, in Lehm. Pl. Preiss. i. 446 (1845) = **Helipterum**, DC. (Compos.).
Pyrethrum, Steetz, l. c. 447 = H. Pyrethrum.

ANISOLOBUS, A. DC. Prod. viii. 395 (1844) = **Odontadenia**, Benth. (Apocyn.).
amazonicus, Muell. Arg. in Mart. Fl. Bras. vi. i. 114.—Bras.
anomalus, Heurck & Muell. Arg. Obs. Bot. 160.—Peruv.
cognatus, Muell. Arg. in Mart. Fl. Bras. vi. i. 113.—Bras.
Cururu, Muell. Arg. l. c. 112.—Bras.
distinctus, Miers, Apocyn. S. Am. 169.—Am. austr.
Fockei, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) (Stirp. Surin. Sel.) 159.—Guian.
gracilipes, Muell. Arg. in Mart. Fl. Bras. vi. i. 115.—Bras.
hebecarpus, Muell. Arg. l. c. 111.—Bras.
Hostmanni, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) (Stirp. Surin. Sel.) 159.—Surinam.
Kappleri, Miq. in Linnaea, xviii. (1844) 737 = *Cururu*.
oblongus, Miers, Apocyn. S. Am. 169.—Am. austr.
Perrottetii, A. DC. in DC. Prod. viii. 395.—Bras.
Pohlianus, Miers, Apocyn. S. Am. 171.—Am. austr.
psidiifolius, Miers, l. c. 172.—Am. austr.
pulcherrimus, Miers, l. c. 171.—Am. austr.
puncticulosa, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) (Stirp. Surin. Sel.) 158.—Guian.
rubidulus, Miers, Apocyn. S. Am. 173.—Am. austr.
Salzmanni, A. DC. in DC. Prod. viii. 395.—Bras.
Sprucei, Muell. Arg. in Mart. Fl. Bras. vi. i. 114.—Bras.
Stadelmeyeri, Muell. Arg. l. c. 113.—Bras.
Zuccarinianus, Miers, Apocyn. S. Am. 171.—Am. austr.

ANISOLOTUS, Bernh. Sem. Hort. Erf. (1837); Linnaea, xii. (1838) Litt. 75 = **Hosackia**, Dougl. (Legum.).
anthylloides, Bernh. ll. cc. = H. subpinnata.
Poeppigiana, Bernh. ll. cc.—Chili.
Wrangeliana, Bernh. ll. cc. = H. subpinnata.

ANISOMALLON, Baill. Adansonia, xi. (1874) 197. *OLACINEAE*.
clusiaefolium, *Baill. l. c.*—N. Caled.

ANISOMELES, R. Br. Prod. 503 (1810). *LABIATAE*, Benth. & Hook. f. ii. 1207.
albiflora, *Miq. Fl. Ind. Bat.* ii. 976.—Malaya.
australis, *Spreng. Syst. iv. Cur. Post.* 226.—Austral.
candicans, *Benth. in Wall. Pl. As. Rar.* i. 59.—Ind. or.
cuneata, *Jacq. Ecl. Pl.* 127.—Ind. or.
disticha, Heyne, ex Roth, Nov. Pl. Sp. 254 = ovata.
furcata, Loud. Hort. Brit. 232 = Craniotome versicolor.
glabrata, Benth. in Wall. Cat. n. 2041 = ovata.
Heyneana, *Benth. in Wall. Pl. As. Rar.* i. 59.—Ind. or.

ANISOMELES:—
inodora, *R. Br. Prod.* 503.—Austral.
intermedia, Wight, in Benth. Lab. Gen. et Sp. 703 = malabarica.
malabarica, *R. Br. ex Sims, Bot. Mag.* t. 2071.—Ind. or.; Ins. Masc.
mollissima, Wall. Cat. n. 2039 = ovata.
moschata, *R. Br. Prod.* 503.—Austral.
nepalensis, Spreng. Syst. ii. 706 = Craniotome versicolor.
ovata, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 364.—Ind. or.; Malaya.
salviaefolia, *R. Br. Prod.* 503.—Austral.

ANISOMERIA, D. Don, in Edinb. N. Phil. Journ. xiii. (1832) 238. *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 85.
coriacea, *D. Don, l. c.*—Chili.
drastica, *Moq. in DC. Prod.* xiii. II. 25.—Chili.
fruticosa, *Phil. in Linnaea*, xxix. (1857-8) 38.—Chili.
littoralis, *Moq. in DC. Prod.* xiii. II. 25.—Chili.

ANISOMERIS, Presl, Symb. Bot. ii. 5. t. 54 (1833) = **Chomelia**, Jacq. (Rubiac.).
diversifolia, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 211.—Bras.
spinosa, Presl, Symb. Bot. ii. 5. t. 54 = C. Anisomeris.

ANISOMETROS, Hassk. in Flora, xx. (1847) 602 = **Pimpinella**, Linn. (Umbell.).
alpina, Hassk. l. c. = P. javana.

ANISONEMA, A. Juss. Euph. Tent. 19. t. 4 (1824) = **Phyllanthus**, Linn. (Euphorb.).
dubium, Blume, Bijdr. 589 = P. reticulatus.
eglandulosum, Decne. Herb. Tim. Descr. 154 = P. reticulatus.
glaucinum, Miq. Fl. Ind. Bat. Suppl. 449.—Sumatr.
hypoleucum, Miq. l. c.—Sumatr.
intermedium, Decne. in Nouv. Ann. Mus. Par. iii. (1831) 482 = P. reticulatus.
jamaicense, Griseb. Fl. Brit. W. Ind. 716 = P. jamaicensis.
multiflorum, Wight, Ic. t. 1899 = P. reticulatus.
reticulatum, A. Juss. Euph. Tent. t. 4. f. 11 = P. reticulatus.
Zollingeri, Miq. Fl. Ind. Bat. i. II. 375 = P. reticulatus.

ANISOPAPPUS, Hook. & Arn. Bot. Beech. Voy. 196 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 339.
africanus, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 369.—Afr. trop.
chinensis, *Hook. & Arn. Bot. Beech. Voy.* 196.—China.

ANISOPETALA, Walp. Rep. iv. 228 (1848) = **Pelargonium**, L'Hérit. (Geran.).
crispa, Walp. l. c. = P. crispum.
purpurascentia, Walp. l. c. = P. purpurascens.
Radulae, Walp. l. c. = P. Radula, etc.
tomentosa, Walp. l. c. = P. tomentosa.

ANISOPETALUM, Hook. Exot. Fl. t. 149 (1825) = **Bulbophyllum**, Thou. (Orchid.).
Careyanum, Hook. l. c. = B. Careyanum.

ANISOPHYLLEA, R. Br. ex Sabine, in Trans. Hort. Soc. v. (1824) 446. *RHIZOPHOREAE*, Benth. & Hook. f. i. 683.
ANISOPHYLLUM, G. Don, ex Hook. Niger Fl. 342, 575 (1849).
TETRACARPAEA, Benth. in Journ. Linn. Soc. iii. (1859) 72, sphalm.
TETRACRYPTA, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 314.
Beccariana, *Baill. Adansonia*, xi. (1875) 311.—Borneo.
disticha, Baill. l. c. = trapezoidalis.
Gaudichaudiana, *Baill. l. c.*—Am. trop.
grandis, *Benth. in Journ. Linn. Soc.* i. (1857) 79, in nota.—Penang.
grandifolia, G. Henslow, in Hook. f. Fl. Brit. Ind. ii. 442 = grandis.
Griffithii, *Oliver, in Linn. Trans. Soc.* xxiii. (1862) 460.—Malacca.
laurina, *R. Br. ex Sabine, in Trans. Hort. Soc.* v. (1824) 446.—Afr. trop.

ANISOPHYLLEA:—
rhomboidea, *Baill. Adansonia*, xi. 1875) 310.—Borneo.
trapezoidalis, *Baill. l. c.* iii. (1862) 4—Singapur.
zeylanica, *Benth. in Hook. Niger Fl.* 343, 575.—Zeylan.

ANISOPHYLLUM, Boiv. ex Baill. Adansonia, i. (1860–61) 153 = **Croton**, Linn. (Euphorb.).
acutifolium, Boiv. ex Baill. l. c. = C. muricatus.
scutelligerum, Boiv. ex Baill. l. c. 154 = C. Paycrianus.

ANISOPHYLLUM, G. Don, ex Benth. in Hook. Niger Fl. 342, 575 (1849) = **Anisophyllea**, R. Br. (Rhizoph.).

ANISOPHYLLUM, Haw. Syn. Pl. Succ. 159 (1812) = **Euphorbia**, Linn.
aegypticum, Schweinf. Beitr. Fl. Acthiop. 34 = E. aegyptiaca.
affine, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 29 = E. Balbisii.
amoenum, Klotzsch & Garcke, l. c. 33 = E. adenoptera.
arabicum, Schweinf. Beitr. Aethiop. 34 = E. arabica.
arenarium, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 24 = E. zygophylloides.
Atoto, Klotzsch & Garcke, l. c. 37 = E. Atoto.
bahiense, Klotzsch & Garcke, l. c. 33 = E. bahiensis.
Berterianum, Klotzsch & Garcke, l. c. 32 = E. Berteriana.
Besseri, Klotzsch & Garcke, l. c. 30 = E. Besseri.
Burmannianum, Klotzsch & Garcke, l. c. 25 = E. thymifolia.
caecorum, Klotzsch & Garcke, l. c. 38 = E. caecorum.
californicum, Klotzsch & Garcke, l. c. 36 = E. Deppeana.
centunculoides, Klotzsch & Garcke, l. c. 38 = E. centunculoides.
Chamaesyce, Haw. Syn. Pl. Succ. 160 = E. Chamaesyce.
Chamissonis, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 22 = E. Chamissonis.
cheirolepis, Klotzsch & Garcke, l. c. 21 = E. cheirolepis.
congenerum, Klotzsch & Garcke, l. c. 22 = E. congenera.
convolvuloides, Klotzsch & Garcke, l. c. 34 = E. convolvuloides.
cordatum, Klotzsch & Garcke, l. c. 38 = E. cordata.
crassipes, Klotzsch & Garcke, l. c. = E. macropus.
densiflorum, Klotzsch & Garcke, l. c. 28 = E. adenoptera.
dentatum, Haw. Syn. Pl. Succ. 162 = E. dentata.
dioicum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 31 = E. dioica.
emarginatum, Klotzsch & Garcke, l. c. 24 = E. emarginata.
Fendleri, Klotzsch & Garcke, l. c. 26 = E. Fendleri.
flexuosum, Klotzsch & Garcke, l. c. 35 = E. flexuosa.
Forskalii, Klotzsch & Garcke, l. c. 25 = E. Forskalii.
Geyeri, Klotzsch & Garcke, l. c. 23 = E. Geyeri.
glabratum, Klotzsch & Garcke, l. c. 35 = E. glabrata.
glaucophyllum, Klotzsch & Garcke, l. c. = E. glaucophylla.
golianum, Klotzsch & Garcke, l. c. 29 = E. goliana.
granulatum, Schweinf. Beitr. Fl. Aethiop. 34 = E. granulata.
Humboldtii, Haw. Syn. Pl. Succ. 160 = E. graminea.
humifusum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 21 = E. humifusa.
humistratum, Klotzsch & Garcke, l. c. 26 = E. humistrata.
hypericifolium, Haw. Syn. Pl. Succ. 161 = E. hypericifolia.
hyssopifolium, Haw. l. c. = E. hyssopifolia.
hyssopifolium, Boiss. in DC. Prod. xv. II. 23 = E. hypericifolia.
inaequale, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 27 = E. adenoptera.
inaequilaterum, Klotzsch & Garcke, l. c. 22 = E. inaequilatera.
indicum, Schweinf. Beitr. Fl. Aethiop. 34 = E. hypericifolia.
Ipecacuanha, Haw. Syn. Pl. Succ. 164 = E. Ipecacuanha.

ANISOPHYLLUM:—
laevigatum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 35 = E. laevigata.
lasiocarpum, Klotzsch & Garcke, l. c. = E. lasiocarpa.
leiospermum, Klotzsch & Garcke, l. c. 23 = E. obliqua.
leucanthum, Klotzsch & Garcke, l. c. 27 = E. leucantha.
Lindenianum, Klotzsch & Garcke, l. c. 37 = E. Lindeniana.
lineare, Klotzsch & Garcke, l. c. 38 = E. linearis.
macropus, Klotzsch & Garcke, l. c. 33 = E. macropus.
maculatum, Haw. Syn. Pl. Succ. 162 = E. maculata.
melanadenium, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 23 = E. melanadenia.
Meyenianum, Klotzsch & Garcke, l. c. 25 = E. Meyeniana.
mossambicense, Klotzsch & Garcke, l. c. 30 = E. mozambicensis.
multiforme, Klotzsch & Garcke, l. c. 37 = E. multiformis.
Mundii, Klotzsch & Garcke, l. c. 25 = E. parvifolia.
Nagleri, Klotzsch & Garcke, l. c. 31 = E. Nagleri.
nanum, Klotzsch & Garcke, l. c. 32 = E. chamaerrhodos.
nodosum, Klotzsch & Garcke, l. c. 22 = E. clusiaefolia.
novo-mexicanum, Klotzsch & Garcke, l. c. 31.—N. Mexic.
ocymoideum, Haw. Syn. Pl. Succ. 161 = E. ocymoidea.
orbiculatum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 24 = E. orbiculata.
ovalifolium, Klotzsch & Garcke, l. c. 26 = E. ovalifolia.
Peplis, Haw. Syn. Pl. Succ. 159 = E. Peplis.
pilulifera, Haw. l. c. 162 = E. pilulifera.
polycnemoides, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 34 = E. polycnemoides.
polygonifolium, Haw. Syn. Pl. Succ. 160 = polygonifolia.
prostratum, Haw. l. c. 163 = E. prostrata.
ramosissimum, Klotzsch & Garcke, in Abh. Akad. Berl. 1859 (1860) 30 = E. Garckeana.
rhytispermum, Klotzsch & Garcke, l. c. 34 = E. rhytisperma.
roseum, Haw. Syn. Pl. Succ. 163 = E. rosea.
Ruizianum, Klotzsch & Garcke, l. c. 37 = E. Ruiziana.
scordifolium, Klotzsch & Garcke, l. c. 35 = E. scordifolia.
Selloi, Klotzsch & Garcke, l. c. 32 = E. Selloi.
senile, Klotzsch & Garcke, l. c. 28 = E. stictospora.
serpens, Klotzsch & Garcke, l. c. 23 = E. serpens.
setigerum, Klotzsch & Garcke, l. c. 29 = E. setigera.
tenuiflorum, Klotzsch & Garcke, l. c. 26 = E. velligera.
tettense, Klotzsch & Garcke, l. c. 34 = E. tettensis.
thymifolium, Haw. Syn. Pl. Succ. 160 = E. thymifolia.
Vahlii, Klotzsch & Garcke, l. c. 36 = E. Vahlii.
velleriflorum, Klotzsch & Garcke, l. c. 28 = E. velleriflora.
virgatum, Klotzsch & Garcke, l. c. 36 = E. Hookeri.

ANISOPHYLLUM, Jacq. Select. Am. 283 (1763) (Quid?).
pinnatum, Jacq. l. c.—Ins. Baru.

ANISOPLECTUS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 120 (Gesn. Centr. Am. 46) = **Alloplectus**, Mart. (Gesnerac.).
pendulus, Oerst. ll. cc. = Allop. pendulus.

ANISOPLEURA, Fenzl, in Flora, xxvi. (1843) 459 = **Prangos**, Lindl. (Umbell.).
crenata, Fenzl, l. c.—Mesopotam.

ANISOPOGON, R. Br. Prod. 176 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1162.
avenaceus, *R. Br. l. c.*—Austral.
capensis, *Nees, Gram. Afr. Austr.* 265.—Afr. austr.

ANISOPTERA, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 65. t. 6 (1839). *DIPTEROCARPEAE*, Benth. & Hook. f. i. 192.
ANTHEROTRICHE, Turcz. in Bull. Soc. Nat. Mosc. (1846) II. 505.
bantamensis, Hassk. Retzia, i. 140 = Vatica bantamensis.
costata, *Korth. Verh. Nat. Gesch. Bot.* 65. t. 6.—Borneo.
glabra, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 61.—Burma.

ANISOPTERA :—
Guiso, *A. DC. in DC. Prod.* xvi. ii. 616.—Ins. Philipp.
lanceolata, Walp. Ann. i. 113 = thurifera.
Mangachapoi, A. DC. in DC. Prod. xvi. II. 616 = Vatica Mangachapoi.
marginata, *Korth. Verh. Nat. Gesch. Bot.* 66. t. 6.—Borneo.
melanoxylon, Hook. f. in Trans. Linn. Soc. xxiii. (1860) 160 = Vatica mclanoxylon.
oblonga, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 301.—Burma.
odorata, Kurz, in Flora, lv. (1872) 190 = Vatica grandiflora.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* i. 485.—Sumatr.
plagata, *Blume, Mus. Bot. Lugd. Bat.* ii. 42.—Ins. Philipp.
polyandra, *Blume, l. c.* t. 6.—N. Guin.
thurifera, *Blume, l. c.*—Ins. Philipp.

ANISORA, Rafin. Sylva Tellur. 74 (1838) = **Helicteres**, Linn. (Stercul.).
angulata et *Murri*, Rafin. l. c.—Hab. ?

ANISORAMPHUS, DC. Prod. vii. 251 (1838) = **Crepis**, Linn. (Compos.).
hypochaerideus, DC. l. c.—Afr. austr.
hypochaeroides, Hook. f. in Journ. Linn. Soc. vii. (1864) 204 = C. Hookeriana.

ANISOSCIADIUM, DC. Coll. Mém. v. 63. t. 15 (1829); Prod. iv. 234 (1830) = **Echinophora**, Linn. (Umbell.).
orientale, DC. Prod. l. c.—Persia.

ANISOSPERMA, Manso, Enum. Subst. Bras. 38 (1836). *CUCURBITACEAE*, Benth. & Hook. f. i. 820.
Passiflora, *Manso, l. c.*—Bras.

ANISOSTACHYA, Nees, in DC. Prod. xi. 368, 730 (1847) = **Justicia**, Linn. (Acanth.).
Bojeri, Nees, in DC. Prod. xi. 368, 730.—Madag.
Commersoni, T. Anders. in Journ. Linn. Soc. vii. (1864) 46 = J. haplostachya.
velutina, Nees, in DC. Prod. xi. 730.—Madag.

ANISOSTEMON, Turcz. in Bull. Soc. Nat Mosc. xx. (1847) I. 152 = **Connarus**, Linn.
trifoliatus, Turcz. l. c. = C. polyanthus.

ANISOSTICHUS, Bur. Monog. Bign. 43. t. 6 (1864) = **Bignonia**, Tourn.
capreolata, Bur. l. c. = B. caprcolata.

ANISOSTICTUS, Benth. & Hook. f. Gen. ii. 1033 (1883), lapsu = ANISOSTICHUS, Bur. = **Bignonia**, Tourn.

ANISOTES, Nees, in DC. Prod. xi. 424 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1111.
CALASIAS, Rafin. Fl. Tellur. iv. 64 (1836), nomen prius.
diversifolius, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 88.—Ins. Socotr.
trisulcus, *Nees, in DC. Prod.* xi. 424.—Arab.

ANISOTES, Lindl. Introd. Nat. Syst. ed. II. 100, 441 (1836) = **Lythrum**, Linn.
anomalus, Lindl. l. c. 441 = L. anomalum.
Hilarianus, Meissn. Gen. Comm. 84 = L. anomalum.

ANISOSTICTE, Bartl. Ord. 294 (1830) = **Capparis**, Linn. (C. frondosa, Jacq.).

ANISOTOMA, Fenzl, in Linnaea, xvii. (1843) 330. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 781.
ANISOTOMARIA, Presl, Bot. Bemerk. 103 (1844).
DECACERAS, Harv. Thes. Cap. ii. 9. t. 114 (1863).
LOPHOSTEPHUS, Harv. l. c. 113 (1863).
Arnottii, *Benth. & Hook. f. Gen.* ii. 781.—Afr. austr.
cordifolia, *Fenzl, in Linnaea*, xvii. (1843) 331.—Afr. austr.

ANISOTOMARIA, Presl, Bot. Bemerk. 103 (1844) = **Anisotoma**, Fenzl (Asclep.).
mollis, Presl, l. c. = Anisotoma cordifolia.

ANISOTOME, Hook. f. Fl. Antarct. 16. t. 8, 9 (1844) = **Aciphylla**, Forst. (Umbell.).
antipoda, Hook. f. Fl. Antarct. 17.—Ins. Antarct.
aromatica, Hook. f. Fl. N. Zeal. i. 89.—N. Zel.
geniculata, Hook. f. l. c. 90 = Angelica geniculata.
Gingidium, Hook. f. l. c. 89 = Angelica Gingidium.
intermedia, Hook. f. l. c. = Ligusticum intermedium.
latifolia, Hook. f. Bot. Antarct. 16 = Ligusticum latifolium.
Lyallii, Hook. f. Fl. N. Zeal. i. 88 = Ac. Lyallii.
rosaefolia, Hook. f. l. c. 90 = Angelica rosaefolia.

ANISTYLIS, Rafin. Neogenyt. 4 (1825) = **Liparis**, Rich. (Orchid.).
Convallaria, Rafin. l. c. = L. liliifolia.
lutea, Rafin. l. c. = L. Loeselii.

ANISUM, Riv. ex Linn. Gen. ed. I. 83 (1737) = **Pimpinella**, Linn. (Umbell.).
caffrum, Eckl. & Zeyh. Enum. 341 = P. caffra.
officinale, DC. Prod. iv. 122 = P. Anisum.
officinarum, Moench, Meth. 100 = P. Anisum.
stadense, Eckl. & Zeyh. Enum. 341 = P. stadensis.
vulgare, Gaertn. Fruct. i. 23 = P. Anisum.

ANKYLOBUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 148 = **Astragalus**, Tourn. (Legum.).
caspicus, Stev. l. c. = A. caspius.
contortuplicatus, Stev. l. c. = A. contortuplicatus.
cruciatus, Stev. l. c. = A. cruciatus.
hamosus, Stev. l. c. = A. hamosus.
reticulatus, Stev. l. c. = A. reticulatus.

ANKYROPETALUM, Fenzl, in Bot. Zeit. i. (1843) 393 = **Gypsophila**, Linn. (Caryophyll.).
arsusianum, Kotschy, in Boiss. Fl. Orient. i. 533.—Syria.
coelesyriacum, Boiss. Diagn. Ser. I. viii. 59.—Oriens.
gypsophiloides, Fenzl, in Bot. Zeit. i. (1843) 393.—Mesopot.
Reuteri, Boiss. & Haussk. in Boiss. Fl. Orient. i. 533.—Oriens.

ANNESLEA, W. Hook. in Salisb. Parad. Lond. t. 64 (1807) = ANNESLIA, Salisb.

ANNESLEA, Roxb. ex Andr. Bot. Rep. t. 618 (1810); Fl. Ind. ii. 573 (1832) = **Euryale**, Salisb. (Nymph.).
amazonica, Presl, Bot. Bemerk. 103 = Victoria regia.
brasiliana, Presl, l. c. = Victoria regia.
spinosa, Andr. Bot. Rep. x. t. 618 = E. ferox.

ANNESLEA, Wall. Pl. As. Rar. i. 5. t. 5 (1829). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 182.
CALLOSMIA, Presl, Bot. Bemerk. 103 (1844).
crassipes, *Hook. ex Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 121.—Malaya.
fragrans, *Wall. Pl. As. Rar.* i. 5. t. 5.—Burma.
montieola, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 59.—Burma.

ANNESLEIA, = ANNESLEA, W. Hook. = **Calliandra**, Benth. (Legum.).

ANNESLIA, Salisb. Parad. Lond. t. 64 (1807) = **Calliandra**, Benth. (1840) (Legum.).
falcifolia, Salisb. l. c. in textu = C. Houstoni.
grandiflora, Salisb. l. c., t. 64 = C. Houstoni.
Houstoni, Sweet, Hort. Brit. ed. II. 163 = C. Houstoni.

ANNESORHIZA, Cham. & Schlecht. in Linnaea, i. (1826) 398. *UMBELLIFERAE*, Benth. & Hook. f. i. 913.
ACROGLYPHE, E. Mey. Zwei Pfl. Docum. 100 (1843).
GLIA, Sond. in Harv. & Sond. Fl. Cap. ii. 547 (1862).
STENOSEMIS, E. Mey. Zwei Pfl. Docum. 141, 148 (1843).
capensis *Cham. & Schlecht. l. c.* 399.—Afr. austr.
elata, *Eckl. & Zeyh. Enum.* 345.—Afr. austr.
filicaulis, *Eckl. & Zeyh. l. c.*—Afr. austr.
hirsuta, *Eckl. & Zeyh. l. c.*—Afr. austr.
macrocarpa, *Eckl. & Zeyh. l. c.*—Afr. austr.
montana, *Eckl. & Zeyh. l. c.* 344.—Afr. austr.
spuria, *Eckl. & Zeyh. l. c.* 345.—Afr. austr.
villosa, *Sond. in Harv. & Sond. Fl. Cap.* ii. 546.—Afr. austr.

ANNONA, Linn. Hort. Cliff. 222 (1737); Forsk. Fl. Aegypt. Arab. 102 = **Anona**, Linn.

ANNULARIA, Hochst. in Flora, xxiv. (1841) 670 = **Voacanga**, Thou. (Apocyn.).
natalensis, Hochst. l. c. 671 = V. Dregei.

ANOCHEILE, Hoffmgg. ex Reichb. Nom. 235 (1841) = **Epidendrum**, Linn. (Orchid.).

ANODA, Cav. Diss. i. 38. t. 10. f. 3 (1785). *MALVACEAE*, Benth. & Hook. f. i. 202.
CAVANILLEA, Medic. Malv. 19 (1787).
acerifolia, Cav. in Anal. Cienc. Nat. vi. (1803) 336; DC. Prod. i. 459 = hastata.
angustifolia, *Spreng. Nachtr.* i. *Bot. Gart. Halle*, 11.—Hab. ?
arenariaeflora, Hort. ex Steud. Nom. ed. II. i. 100 = crenatiflora.
brachyantha, Reichb. Hort. Bot. t. 34 = triangularis.
crenatiflora, Ortcg. Hort. Matr. Dec. 96 = parviflora ?
cristata, Schlecht. in Linnaea, xi. (1837) 210 = lavateroides.
Dilleniana, Cav. Diss. i. 40. t. 11. f. 1 = cristata.
fernandeziana, *Steud. in Flora*, xxxix. (1856) 437.—Ins. Juan Fern.
hastata, *Cav. Diss.* i. 39. t. 10. f. 3.—Mexic.; Ind. occ.
hirsuta, *Phil. in Linnaea*, xxviii. (1856) 675.—Chili.
incarnata, *H. B. & K. Nov. Gen. et Sp.* v. 266.—Mexic.
incarnata, Spreng. ex Steud. Nom. ed. II. i. 100 = triangularis.
lanceolata, *Hook. & Arn. Bot. Beech. Voy.* 411.—Mexic.
lavateroides, *Medic. Malv.* 19.—Mexic.
Ochsenii, Phil. in Linnaea, xxviii. (1856) 613 = Abutilon Ochsenii.
Ortegae, Spreng. Nachtr. i. Bot. Gart. Halle, 11 = crenatiflora.
ovata, Meyen, Reise, ii. 139 = Sida fallax.
parviflora, *Cav. Ic.* v. 19. t. 431.—Mexic.
pentaschista, *A. Gray, Pl. Wright.* ii. 22.—Nov. Mexic.
populifolia, *Phil. in Linnaea*, xxviii. (1856) 313.—Chili.
pubescens, *Schlecht. in Linnaea*, xi. (1837) 218.—Mexic.
punicea, *Lag. Gen. et Sp. Nov.* t. 21.—Mexic.
reticulata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 368.—Am. bor.
strictiflora, *Steud. in Flora*, xxxix. (1856) 437.—Chili.
triangularis, *DC. Prod.* i. 459.—Mexic.
triloba, Cav. Diss. i. 39. t. 10. f. 3 = lavateroides.
Wrightii, *A. Gray, Pl. Wright.* ii. 22.—Nov. Mexic.

ANODENDRON, A. DC. Prod. viii. 443 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 719.
Candolleanum, *Wight, Ic.* iv. t. 1309.—Malaya.
coriaccum, *Miq. Fl. Ind. Bat.* ii. 455.—Java.
inflatum, *Hassk. in Flora*, xxviii. (1845) 269.—Java.
laeve, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* i. 315.—Japon.
moluccanum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 140.—Malaya.
paniculatum, *A. DC. in DC. Prod.* viii. 444.—Ind. or.; Malaya.
pauciflorum, *Hook. f. Fl. Brit. Ind.* iii. 669.—Penang.
rhinosporum, *Thw. Enum. Pl. Zeyl.* 194.—Zeylan.
rubescens, Teijsm. & Binn. Cat. Hort. Bogor. 127 = Candollcanum.
tenuiflorum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 140.—Sumatr.

ANODIA, Hassk. in Flora, xxv. (1842) II. Beibl. 38, in nota = **Anoda**, Cav. (Malvac.).

ANODISCUS, Benth. in Benth. & Hook. f. Gen. ii. 998 (1876). *GESNERACEAE*, Benth. & Hook. f. ii. 998.
peruvianus, *Benth. in Hook. Ic. Pl.* t. 1199.—Peruv.

ANODONTEA, Sweet, Hort. Brit. ed. I. 467 (1827) = **Alyssum**, Tourn. (Crucif.).
arenaria, G. Don, Gen. Syst. i. 180 = Al. montanum ?

ANODONTEA :—
Aucheri, D. Dietr. Syn. Pl. iii. 643 = Alyssum foliosum.
canescens, Sweet, Hort. Brit. ed II. 25 = Alyssum canescens.
dasycarpa, Sweet, l. c. = Alyssum dasycarpum.
edentula, Sweet, l. c. ed 1. 467 = Alyssum gemonense.
halimifolia, Sweet, l. c. = Alyssum halimifolium.
macrocarpa, Sweet, l. c. = Alyssum macrocarpum.
marginata, D. Dietr. Syn. Pl. iii. 643 = Alyssum marginatum.
meniocoides, D. Dietr. l. c. = Alyssum meniocoides.
peltarioides, D. Dietr. l. c. = Alyssum peltarioides.
pyrenaica, G. Don, Gen. Syst. i. 180 = Alyssum pyrenaicum.
rupestris, Sweet, Hort Brit. ed. I. 467 = Alyssum rupestris.
spinosa, Sweet, l. c. = Alyssum spinosum.
tenuifolia, G. Don, Gen. Syst. i. 180 = Alyssum tenuifolium.
xanthocarpa, D. Dietr. Syn. Pl. iii. 643 = Alyssum macrostylum.

ANODOPETALUM, A. Cunn. ex Endl. Gen. 818 (1839). *SAXIFRAGEAE*, Benth. & Hook. f. i. 652.
biglandulosum, *A. Cunn. ex Hook. f. Fl. Tasm.* i. 148. —Austral.
glandulosum, Baill. Hist. Pl. iii. 378 = biglandulosum.

ANOECTOCHILUS, Blume, Bijdr. 411. t. 15 (1825) (Anecochilus). *ORCHIDEAE*, Benth. & Hook. f. iii. 598.
albo-lineatus, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 141 = brevilabris.
albo-marginatus, *Lour. Encyc. Pl. Suppl.* ii. 1539.—Cochinch.
argenteus, Hort. ex Loud. Hort. Brit. Suppl. iii. 608 = Physurus argenteus.
argyroneurus, *C. Koch & Lauche, in Berl. Allg. Gartenz.* xxv. (1857) 4.—Java.
aureus, Hort. ex C. Koch & Lauche, l. c. = setaceus.
brevilabris, *Lindl. Gen. et Sp. Orch.* 499.—Ind. or.
Bulleni, *Low, ex Morr. in Belg. Hortic.* xii. (1862) 10. —Borneo.
crispus, *Lindl. in Journ. Linn. Soc.* i. (1857) 180.—Ind. or.
Croesus ×, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 424.—Hab. ?
Dawsonianus, Low, ex Reichb. f. in Gard. Chron. (1868) 1038 = Haemaria discolor.
Dominii ×.—*Cf. Burbridge, in Gard. Chron.* (1871) 44.
elatus, *Lindl. in Journ. Linn. Soc.* i. (1857) 178.—Ind. or.
elongatus, *Miq. Fl. Ind. Bat.* iii. 734.—Java.
flavescens, Blume, Bijdr. 412 = Odontochilus flavescens.
flavus, Benth. & Hook f. Gen. iii. 598 = Odontochilus flavus.
Frederici-Augusti, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 177, 424.—Hab. ?
glaber, *Miq. Fl. Ind. Bat.* iii. 733.—Malaya.
gracilis, *Blume, Bijdr.* 413.—Malaya.
grandiflorus, Lindl. in Journ. Linn. Soc. i. (1857) 179 = Odontochilus grandiflorus.
Hasseltii, *Miq. Fl. Ind. Bat.* iii. 734.—Java.
intermedius, *Hort. ex Loud. Encyc. Pl. Suppl.* ii. 1492. —Ind. or.
intermedius, *Reichb. f. in Hamb. Gartenz.* xvi. (1860) 177.—Hab. ?
Jaubertii, Gaudich. in Voy. Bonite, t. 100 = Odontochilus Jaubertii.
lanceolatus, Lindl. Gen. et Sp. Orch. 499 = Odontochilus lanceolatus.
latomaculatus, Hort. ex Blume, Coll. des Orch. i. 50 = Roxburghii.
Lobbianus, Planch. in Fl. des Serres, Sér. I. v. (1849) 519 = Roxburghii.
longiflorus, Reichb. f. in Seem. Fl. Vit. 294 = Odontochilus longiflorus.
Lowii, Hort. ex Lond. Encyc. Pl. Suppl. ii. 1492 = Dossinia marmorata.
luteus, Lindl. in Journ. Linn. Soc. i. (1857) 179 = Odontochilus lanceolatus.
neglectus, *Blume, Orch. Arch.* 47 *in nota*.—Zeylan. ?
occultus, *Miq. Fl. Ned. Ind.* iii. 734.—Java.
Ordianus, Hort.—Cf. Gard. Chron. (1869) 1236 = Haemaria discolor.
Ortgiesi, *Hort. ex Van-Geert, Cat.* n. 70 (1873).—N. Granat.
Nevilleana, *Low, ex C. Morr. in Belg. Hortic.* xii. (1862) 10.—Hab. ?

ANOECTOCHILUS :—
pictus, Hort. ex Loud. Hort. Brit. Suppl. iii. 608 = Physurus pictus.
pubescens, *Blume, Bijdr.* 412.—Java.
regalis, *Blume, Orch. Arch. Ind.* 46.—Zeylan.
regalis, Low, ex C. Morr. in Belg. Hortic. xii. (1862) 4 = setaceus.
Reinwardtii, *Blume, Orch. Arch. Ind.* 48.—Malaya.
Roxburghii, *Lindl. in Wall. Cat.* n. 7387.—Ind. or.
Ruckeri, *Low, ex C. Morr. in Belg. Hortic.* xii. (1862) 10.—Borneo.
Rumphii, *Low, ex C. Morr. l. c.* 5.—Malaya.
sandwicensis, *Lindl. Gen. et Sp. Orch.* 500.—Ind. Sandv.
setaceus, *Blume, Bijdr.* 412.—Java.
setaceus, Lindl. Bot. Reg. t. 2010 = regalis.
setaceo-pictus, *C. Koch & Lauche, in Berl. Allg. Gartenz.* xxv. (1857) 4.—Hab. ?
spicatus, *Miq. Fl. Ind. Bat.* iii. 734.—Borneo.
striatus, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1492 = Zeuxine regia.
uniflorus, *Miq. Fl. Ind. Bat.* iii. 733.—Sumatra.
Veitchianus, Hort. ex Blume, Coll. des Orch. i. 119 = Macodes petola.
xanthophyllus, Planch. Fl. des Serres, Sér. I. v. (1849) 519 = Frederici-Augusti.

ANOGEISSUS, Wall. Cat. n. 4014 (1832) *COMBRETACEAE*, Benth. & Hook. f. i. 687.
ANDERSONIA, Buch.-Ham. ex Wall Cat. l. c. (1832).
acuminata, *Wall. Cat.* n. 4014.—Ind. or.
hirta, Wall. Cat. n. 4016 = acuminata.
latifolia, *Wall. Cat.* n. 4015.—Ind. or.
leiocarpa, *Guill. & Perr. Tent. Fl. Seneg.* i. 280. t. 65. —Afr. trop.
myrtifolia, Wall. Cat. n. 4017 = pendula.
pendula, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 171.—Ind. or.
phillyreaefolia, *Heurck & Muell. Arg. Obs. Bot.* 209. —Burma.

ANOGRA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 339 = **Oenothera**, Linn. (Onagr.).
Douglasiana, Spach, l. c. 340 = O. albicaulis.
Nuttalliana, Spach, l. c. = O. albicaulis.
pinnatifida, Spach, l. c. 341 = O. albicaulis.

ANOGYNA, Nees, in Hook. Journ. Bot. ii. (1840) 395 = **Lagenocarpus**, Nees (Cyperac.).
tremula, Nees, l. c. = L. tremula.

ANOIGANTHUS, Baker, in Journ. Bot. xvi. (1878) 76. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 722.
breviflorus, *Baker, l. c.*—Natal.
luteus, Baker, l. c. 77 = breviflorus.

ANOMA, Lour. Fl. Cochinch. 278 (1790) = **Moringa**, Juss.
cochinchinensis, Lour. l. c. 280 (Caesalpinaeae sp. ?).
Moringa, Lour. l. c. 279 = M. pterygosperma.
Morunga, Lour. l. c. = M. pterygosperma.

ANOMALANTHUS, Klotzsch, in Linnaea, xii. (1838) 238 = **Simocheilus**, Klotzsch (Ericac.).
discolor, Klotzsch, l. c. 239.—Afr. austr.
scoparius, Klotzsch, l. c.—Afr. austr.

ANOMALOPTERIS, G. Don, Gen. Syst. i. 647 (1831) = **Acridocarpus**, Guill. & Perr. (Malpig.).
longifolia, G. Don, l. c. = Ac. Smeathmanni.
obovata, G. Don, l. c. = Ac. plagiopterus.
spicata, G. Don, l. c. = Ac. Smeathmanni.

ANOMALOSTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 162. t. 28 (1861) = **Cleome**, Linn. (Capparid.).
bororensis, Klotzsch, l. c.—Afr. trop.

ANOMALOTIS, Steud. Syn. Pl. Gram. 198 (1855) = **Trisetaria**, Forsk. (Gramin.).
quinqueseta, Steud. l. c. = T. quinqueseta.

ANOMANTHODIA, Hook. f. in Benth. & Hook. Gen. ii. 87 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 87.
auriculata, *Hook. f. l. c.*—Malaya.

ANOMANTIA, Rafin. ex DC. Prod. v. 617 (1836) = **Verbesina**, Linn. (Compos.). (V. virginica, Linn.)

ANOMATHECA, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 227 = **Lapeirousia**, Pourr. (Irid.).
angolensis, Baker, in Journ. Bot. xiv. (1876) 337.—Afr. austr.
calamifolia, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 355.—Afr. austr.
cruenta, Lindl. Bot. Reg. t. 1369.—Afr. austr.
grandiflora, Baker, in Journ. Bot. xiv. (1876) 337.—Afr. austr.
juncea, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 227 = L. juncea.
xanthospila, Ker-Gawl. ex Spreng. Syst. i. 154.—Afr. austr.

ANOMAZA, Laws. ex Salisb. in Trans. Hort. Soc. i. (1812) 323 = **Lapeirousia**, Pourr. (Irid.).
excisa, Laws. Cat. 2 ; ex Salisb. l. c. = L. juncea.

ANOMIANTHUS, Zoll. in Linnaea, xxix. (1857–58) 324. *ANONACEAE*, Benth. & Hook. f. i. 27.
heterocarpus, *Zoll. l. c.*—Ins. Bantam.

ANOMOCARPUS, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 351 ; Contrib. ii. 28 = **Calicera**, Cav.
axillaris, Miers, ll. cc. 352 ; 28.—Chili.
eryngioides, Miers, ll. cc. 353 ; 29 = C. eryngioides.
leucanthemus, Miers, ll. cc. 355 ; 31.—Chili.
pulvinatus, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 354 = C. pulvinata.
subsessiliflorus, Miers, l. c. 352 ; Contrib. ii. 29 = C. sessiliflora.
tenuifolius, Miers, ll. cc. 356 ; 33 = A. tenuifolia.
tenuis, Miers, ll. cc. 356 ; 32.—Chili.

ANOMOCHLOA, Brongn. in Ann. Sc. Nat. Sér. III. xvi. (1851) 368. t. 23. *GRAMINEAE*, Benth. & Hook. f. iii. 1111.
macrantoidea, Pritz. Ind. Ic. 74 = marantoidea.
marantoidea, *Brongn. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 369.—Bras.

ANOMORHEGMIA, Meissn. Gen. i. 303 (1840) = **Stauranthera**, Benth. (Gesner.).
Miquelii, Meissn. l. c. ii. 212 = S. ecalcarata.

ANOMOSANTHES, Blume, Rumphia, iii. 151, in obs. (1847) = **Hemigyrosa**, Blume (Sapind.).
deficiens, Blume, l. c. = H. deficiens.

ANOMOSPERMUM, Dalz. in Hook. Kew Journ. iii. (1851) 228 = **Actephila**, Blume (Euphorb.).
excelsum, Dalz. l. c. = Actephila excelsa.

ANOMOSPERMUM, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 39. *MENISPERMACEAE*, Benth. & Hook. f. i. 35.
ELISSARRHENA, Miers, l. c. Ser. III. xiii. (1864) 124.
axilliflorum, *Griseb. Cat. Pl. Cub.* 4.—Cuba.
grandifolium, *Eichl. in Flora*, xlvii. (1864) 388.—Bras.
Hostmanni, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiv. (1864) 103 = Schomburgkii.
japurense, *Eichl. in Flora*, xlvii. (1864) 388.—Bras.
lucidum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xiv. (1864) 103.—Guian.
nitidum, Miers, l. c. = Schomburgkii.
oblongatum, *Miers, l. c.*—Bras.
ovatum, *Miers, l. c.*—Bras.
reticulatum, *Eichl. in Flora*, xlvii. (1864) 388.—Bras.
Schomburgkii, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. vii. (1851) 102; Ser. III. xiv. (1864) 103.—Guian.

ANOMOSTEPHIUM, DC. Prod. v. 560 (1836) = **Aspilia**, Thou. (Compos.).
angustifolium, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 290 = Aspilia foliacea.
buphthalmoides, DC. in Deless. Ic. Sel. iv. t. 32.—Guadalup.
ellipticum, DC. Prod. v. 560 = Aspilia elliptica.
foliosum, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 290 = Aspilia foliosa.
oblongifolium, DC. Prod. v. 560 = Aspilia foliacea.
ovatifolium, DC. l. c. = Aspilia ovalifolia.

ANONA, Linn. Syst. ed. I. (1735); Gen. ed. I. 158 (1737). *ANONACEAE*, Benth. & Hook. f. i. 27.
ANNONA, Linn. Hort. Cliff. 222 (1737).
acutiflora, *Mart. Fl. Bras.* xiii. 1. 10.—Bras.
aethiopica, Steud. Nom. ed. II. ii. 737 = Xylopia aethiopica.
afrieana, *Linn. Sp. Pl.* 537.—Hab.?
amara, *Raeusch. Nom.* ed. III. 161.—Guiana.
Ambotay, *Aubl. Pl. Gui.* i. 616. t. 249.—Am. trop.
amplexieaulis, *Lam. Encyc.* ii. 127.—Madag.; Ins. Masear.
antioquensis, *Linden, Cat.* n. 87 (1871).—N. Granat.
arenaria, Sehum. & Thonn. Beskr. Guin. Pl. 257 = senegalensis.
aromatica, Boj. Hort. Maurit. 6 = Xylopia Richardi.
asiatica, Heyne, ex Wall. Cat. sub n. 6475 = Unona pannosa.
asiatiea, *Linn. Sp. Pl.* 537.—Zeylan.
asiatica, Vahl, Symb. Bot. iii. 73 = Forskahlii.
atabapensis, *H. B. & K. Nov. Gen. et Sp.* v. 58.—Guian.
australis, *St. Hil. Fl. Bras. Mer.* i. 33.—Bras.
axilliflora, *DC. Prod.* i. 86.—Guian.
Barteri, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 477.—Afr. trop.
bibracteata, Hook. Ic. Pl. iv. t. 328 = Guatteria bibraeteata.
Bonplandiana, H. B. & K. Nov. Gen. et Sp. v. 58 = murieata.
bullata, *A. Rich. Ess. Fl. Cub.* 31.—Cuba.
caeans, *Warm. in Kjoeb. Vidensk. Meddel.* (1873) 155.—Bras.
casearilloides, *Griseb. Cat. Pl. Cub.* 2.—Cuba.
cauliflora, *Mart. Fl. Bras.* xiii. 1. 16.—Bras.
Cherimolia, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. trop.
eherimolioides, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 27.—N. Granat.
eheriquensis, *Linden, Cat.* n. 87 (1871).—N. Granat.
chrysocarpa, Lepr. in Guill. & Perr. Tent. Fl. Seneg. i. 6 = palustris.
chrysopetala, *Steud. in Flora*, xxvi. (1843) 754.—Guian.
chrysophylla, *Boj. in Ann. Sc. Nat.* xx. (1830) 53.—Ins. Masear.
cinerea, *Dunal, Monog. Anon.* 71. t. 8.—Ind. oec.
colorata, *Vell. Fl. Flum.* 239; v. t. 129.—Bras.
conica, *Ruiz & Pav. Fl. Per.* v. t. 490; *ex G. Don, Gen. Syst.* i. 88.—Peruv.
eonophylla, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 27.—N. Granat.
coriaeea, *Mart. Fl. Bras.* xiii. 1. 6.—Bras.
eornifolia, *St. Hil. Fl. Bras. Mer.* i. 33.—Bras.
crassifolia, *Mart. Fl. Bras.* xiii. 1. 6.—Bras.
crotonifolia, *Mart. l. c.* 46.—Bras.
densieoma, *Mart. l. c.* 9.—Bras.
depressa, *Baill. in Adansonia*, viii. (1868) 267.—Mexie.
dioiea, *St. Hil. Fl. Bras. Mer.* i. 34.—Bras.
distincta, Raeusch. Nom. ed. III. 160 = Forskahlii.
distincta, Willem. in Ust. Ann. Bot. xviii. 42 = Guatteria Willemetiana.
dodecapetala, Lam. Eneye. ii. 127 = Talauma Plumieri.
dolabripetala, Raddi, in Mem. Mod. xviii. Fis. (1820) 394 = Rollinia longifolia.
eehinata, *Dunal, Monog. Anon.* p. 68. t. 4.—Guian.
exalbida, Vell. Fl. Flum. 240; v. t. 131 = Rollinia exalbida.
exeelsa, *H. B. & K. Nov. Gen. et Sp.* v. 59.—Mexic.
exsucca, Dunal, Monog. Anon. 77 = Rollinia exsueea.
fagifolia, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 132.—Bras.
foetida, *Mart. Fl. Bras.* xiii. 1. 16.—Bras.
Forskahlii, *DC. Syst.* i. 472.—Aegypt.
furfuraeea, *St. Hil. Fl. Bras. Mer.* i. 34.—Bras.
glabra, Forsk. Fl. Aegypt. Arab. 102 = Forskahlii.
glabra, *Linn. Sp. Pl.* 537.—Hab.?
glauca, *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* iv. (1824) 31.—Guinea.
globiflora, *Schlecht. in Linnaea*, x. (1836) 235.—Mexic.
grandiflora, *Lam. Encyc.* ii. 126.—Ins. Mascar.; Madag.
grandiflora, Bart. Trav. 20 & 171. t. 2 = Asimina grandiflora.
grandifolia, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 131.—Bras.
haematantha, *Miq. in Linnaea*, xxii. (1849) 465.—Guian.
hexapetala, Linn. f. Suppl. 270 = Artabotrys odoratissima.

ANONA:—
Hostmanni, *Steud. in Flora*, xxvi. (1843) ii. 754.—Guian.
Humboldtiana, H. B. & K. Nov. Gen. et Sp. v. 56 = seq.
Humboldtii, *Dunal, Monog. Anon.* 64. t. 3.—Cumana.
humilis, *Benth. in Hook. Kew Journ.* iii. (1851) 113.—Bras.
hypoglauea, *Mart. Fl. Bras.* xiii. 1. 13.—Bras.
involuerata, *Baill. Adansonia*, viii. (1868) 265.—Mexic.
ionophylla, *Triana, & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 27.—N. Granat.
laevigata, *Mart. Fl. Bras.* xiii. 1. 17.—Bras.
laevis, *H. B. & K. Nov. Gen. et Sp.* v. 60.—Guian.; Venezuela.
lanceolata, Willd. ex Steud. Nom. ed. II. i. 100 = Guatteria maypurensis.
lasiocalyx, *Mart. Fl. Bras.* xiii. 1. 16.—Bras.
laurifolia, *Dunal, Monog. Anon.* 65.—Ind. oec.
lepidota, *Miq. in Linnaea*, xix. (1847) 129.—Guian.
Liebmanniana, *Baill. Adansonia*, viii. (1868) 266.—Mexie.
longifolia, Aubl. Pl. Gui. i. 615. t. 248 = Duguetia longifolia.
madagascariensis, Pers. Syn. ii. 95 = grandiflora.
Manirote, *H. B. & K. Nov. Gen. et Sp.* v. 59.—Guian.
Mannii, *Oliver, in Hook. Ic. Pl.* t. 1010.—Afr. trop.
Maregravii, *Mart. Fl. Bras.* xiii. 1. 5.—Bras.
mexieana, *Lodd. ex G. Don, in Loud. Hort. Brit.* 227.—Mexie.
micrantha, *Bert. ex Spreng. Syst.* ii. 640.—Ins. S. Doming.
microcarpa, Jacq. Fragm. 40. t. 44 = Diospyros Cargillia.
mieroearpa, *Ruiz & Pav. Fl. Peruv.* v. t. 487; *ex G. Don, Gen. Syst.* i. 90.—Peruv.
montana, *Macfad. Fl. Jamaic.* i. 7.—Jamaie.
montieola, *Mart. Fl. Bras.* xiii. 1. 7.—Bras.
mucosa, Jaeq. Obs. i. 16 = Rollinia mueosa.
muricata, *Linn. Sp. Pl.* 536.—Am. trop.
muscosa, Aubl. Pl. Gui. i. 618 = Rollinia mueosa.
Myristica, Gaertn. Fruet. ii. 194. t. 125 = Monodora Myristiea.
nitida, *Mart. Fl. Bras.* xiii. 1. 9.—Bras.
nitida, Willd. ex Steud. Nom. ed. II. i. 101 = Guatteria maypurensis.
obovata, Willd. Sp. Pl. ii. 1269 = Asimina grandiflora.
obtusiflora, Tuss. Fl. Antill. t. 28 = Rollinia mueosa.
obtusifolia, DC. Prod. i. 84 = Rollinia mueosa.
paludosa, *Aubl. Pl. Gui.* i. 611. t. 246.—Guian.
palustris, *Linn. Sp. Pl.* ed. II. 757.—Am. et Afr. trop.
Pavonii, *G. Don, Gen. Syst.* i. 89.—Peruv.
peduncularis, *Steud. in Flora*, xxvi. (1843) ii. 754.—Guian.
pendula, Salisb. Prod. 380 = Asimina triloba.
Perrottetii, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 197.—Senegal.
peruviana, Humb. & Bonpl. ex Dunal, Monog. Anon. 67 = uliginosa.
phaeoelados, *Mart. Fl. Bras.* xiii. 1. 46.—Bras.
Pisonis, *Mart. l. c.* 5.—Bras.
polyearpa, *DC. Syst.* i. 499.—Afr. trop.
Porpetae, Boiv. ex Baill. in Bull. Soc. Linn. Par. i. (1882) 341 = senegalensis.
pubescens, Salisb. Prod. 380 = A. Chermolia.
punetata, *Aubl. Pl. Gui.* i. 614. t. 247.—Guian.
punieifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 26.—N. Granat.
purpurea, *Moç. & Sesse, ex Dunal, Monog. Anon.* 64.—Mexie.
pygmaea, Bartr. Trav. 18. t. 1 = Asimina pygmaea.
pyriformis, *Boj. Hort. Maurit.* 5; *Baker, Fl. Maurit.* 3.—Ins. Mascar.
quinducnsis, *H. B. & K. Nov. Gen. et Sp.* v. 60.—N. Granat.
retieulata, *Linn. Sp. Pl.* 537.—Am. trop.
reticulata, Sieber, ex A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 200 = Rollinia mueosa.
reticulata, Vell. Fl. Flum. 239; v. t. 130 = Pisonis.
rhizantha, *Eichl. in Eichl. Jahrb. Berl.* ii. (1883) 322.—Bras.
rhombipetala, *Ruiz & Pav. Fl. Per.* v. t. 489; *ex G. Don, Gen. Syst.* i. 87.—Peruv.
riparia, *H. B. & K. Nov. Gen. et Sp.* v. 59.—Peruv.
rufa, *Presl, Rel. Haenk.* ii. 75.—Am. trop.
rufescens, 'Aubl.' ex Steud. Nom. ed. I. 51 (lapsu pro 'Abuta rufescens').—Guian.

ANONA:—
Salzmanni, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 197.—Bras.
senegalensis, *Pers. Syn.* ii. 95.—Afr. trop.; Guian.
sericea, *Dunal, Monog. Anon.* 69.—Guian.
sessiliflora, *Benth. in Hook. Kew Journ.* v. (1853) 8.—Bras.
silvestris, Vell. Fl. Flum. 239; v. t. 128 = Rollinia sylvatiea.
sinensis, *Raeusch. Nom.* ed. III. 160.—China.
sphaeroearpa, *Splitg. in Hoeven & De Vriese, Tijdschr.* ix. (1842) 96.—Guian.
spineseens, *Mart. Fl. Bras.* xiii. 1. 11.—Bras.
squamosa, Delile, Ill. Aegypt. 17 = Forskahlii.
squamosa, *Linn. Sp. Pl.* 537.—Ind. oee.
squamosa, Vell. Fl. Flum. 239; v. t. 127 = obtusiflora.
surriffa, *Roxb. Hort. Beng.* 43, nomen.—Ind. or.
sylvatica, St. Hil. Fl. Bras. Mer. i. 32 = Rollinia sylvatiea.
sylvestris, Auet.—Cf. *silvestris*.
tenuiflora, *Mart. Fl. Bras.* xiii. 1. 10.—Bras.
tenuifolia, *A. DC. in Mém. Soc. Phys. Genèv.* v. (1832) 198.—Guian.
triloba, Linn. Sp. Pl. 537 = Asimina triloba.
tripetala, Ait. Hort. Kew. ii. 252 = A. Cherimolia.
tuberosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 7.—Malaya.
uliginosa, *H. B. & K. Nov. Gen. et Sp.* v. 56.—Peruv.
uncinata, Lam. Encye. ii. 127 = Artabotrys odoratissima.
uniflora, Dunal, Monog. Anon. 76 = Duguetia Candollei.
velutina, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 132.—Bras.
vepertonum, *Mart. Fl. Bras.* xiii. 1. 16.—Bras.
xestropetala, Spreng. Syst. ii. 641 = Rollinia longifolia.
xylopifolia, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 132.—Bras.

ANONIS, Tourn. ex Seop. Fl. Carn. ed. II. ii. 54 (1772) = **Ononis**, Linn. (Legum.).
americana, Aubl. Pl. Gui. ii. 763 = Stylosanthes proeumbens, viscosa.
arvensis, Lam. Fl. Fr. ii. 611 = O. arvensis.
fruticosa, Lam. l. e. 612 = O. frutieosa.
minutissima, Lam. l. c. 608 = O. minutissima.
Natrix, Seop. Fl. Carn. ed. II. ii. 55 = O. Natrix.
pinguis, Lam. Fl. Fr. ii. 609 = O. Natrix.
pusilla, Lam. l. c. 610 = O. reclinata?
reclinata, Lam. l. e. 611 = O. reclinata.
rotundifolia, Lam. l. e. 612 = O. rotundifolia.
scariosa, Moench, Meth. 157 = O. mitissima.
spinosa, Seop. Fl. Carn. ed. II. ii. 54 = O. spinosa.
variegata, Lam. Fl. Fr. ii. 608 = O. variegata.
viscosa, Lam. l. e. 609 = O. viseosa.

ANONYCHIUM, Sehweinf. Reliq. Kotschy. 7. t. 7 (1868) = **Prosopis**, Linn. (Legum.).
lanceolatum, Sehweinf. l. e. = P. oblonga.

ANONYMOS, Walt. Fl. Carol. (1788) (passim pro omnibus dubiis).
aquatica, Walt. l. e. 37 = Limnanthemum trachyspermum.
aquatica, Walt. l. e. 230 = Planera aquatica.
aquatica, Walt. l. c. 109 = Saururus cernuus.
aquatica, Walt. l. c. 137 = Nesaea verticillata.
asteroides, Michx. ex DC. Prod. v. 301 = Boltonia asteroides.
bracteata, Walt. Fl. Carol. 181 = Zornia tetraphylla.
capitata, Walt. l. c. 69 = Burmannia capitata.
carolinensis, Walt. l. c. 60 = Elytraria virgata.
carolinensis, Walt. l. c. 91 = Lithospermum hirtum.
carolinensis, Walt. l. e. 164 = Verbena Aubletia.
carolinensis, Walt. l. c. 168 = Ruellia carolinensis.
carolinensis, Walt. l. c. 188 = Amphicarpaea monoica.
cassioides, Walt. l. c. 171 = Seymeria tenuifolia.
ciliata, Walt. l. c. 197 = Liatris graminifolia?
corymbosus, Walt. l. c. 139.—Am. bor.
dioscoroides, Croom, in Am. Journ. Sc. xxix. (1835) 165 = Croomia paueiflora.
erecta, Walt. Fl. Carol. 170 = Houstonia caerulea.
flavus, Walt. l. c. = Gerardia flava.
frondosus, Walt. l. c. 139.—Am. bor.

ANONYMOS:—
frutescens, Walt. Fl. Carol. 186 = Wistaria speciosa.
graminifolia, Walt. l. c. 197 = Liatris graminifolia?
ligustrinus, Walt. l. c. 139.—Am. bor.
odoratissima, Walt. l. c. 198 = Trilisia odoratissima.
paniculata, Walt. l. c. = Trilisia paniculata.
pedicularis, Walt. l. c. 170 = Gerardia?
petiolata, Walt. l. c. 108 = Mitreola petiolata.
pilosa, Walt. l. c. 197 = Liatris pycnostachya.
pinnatifida, S. Wats. Bibl. Ind. 247, lapsu = seq.
pinnatus, Walt. Fl. Carol. 103 = Petalostemon corymbosus.
procumbens, Walt. l. c. 86 = Anotis rotundifolia.
pudica, Walt. l. c. 123 = Uvularia perfoliata.
purpurea, Walt. l. c. 170 = Gerardia purpurea.
quaternata, Walt. l. c. 246 = Gerardiae sp.
quinata, Walt. l. c. = Gerardiae sp.
ramosa, Walt. l. c. 198 = Liatris scariosa.
repens, Walt. l. c. 110 = Dichondra repens.
repens, Walt. l. c. 160 = Mitchella repens.
rotundifolia, Walt. l. c. 181 = Crotalaria ovalis.
sagittalis, Walt. l. c. = Crotalaria sagittalis.
sempervirens, Walt. l. c. 98 = Gelsemium sempervirens.
sessilifolia, Walt. l. c. 108 = Mitreola petiolata.
setacea, Walt. l. c. 170 = Gerardia setacea?
tinctoria, Walt. l. c. 68 = Lachnanthes tinctoria.
umbrosa, Walt. l. c. 63 = Micranthemum orbiculatum.
uniflora, Walt. l. c. 198 = Carphephorus bellidifolius?

ANOOSPERMA, Kunze, in Bot. Zeit. i. (1843) 661, *err. typ.* = **Oncosperma**, Blume (Palm.).

ANOPLANTHUS, Endl. Gen. 727 (1839) = **Phelypaea**, Tourn. et **Aphyllon**, Mitch. (Orobanch.).
Biebersteinii, Reut. in DC. Prod. xi. 42 = P. coccinea.
coccineus, Walp. Rep. iii. 481 = P. coccinea.
comosus, Walp. l. c. 480 = Aphyllon comosum.
fasciculatus, Walp. l. c. = Aphyllon fasciculatum.
Tournefortii, Reut. in DC. Prod. xi. 42 = P. coccinea.
uniflorus, Endl. Ic. Gen. t. 72 = Aphyllon uniflorum.

ANOPLOCARYUM, Ledeb. Fl. Ross. iii. 154 (1847) = **Eritrichium**, Schrad. (Borag.).
compressum, Ledeb. l. c.—Rossia.

ANOPLON, Reichb. Consp. 212 b (1828) = **Phelypaea**, Tourn. et **Aphyllon**, Mitch. (Orobanch.).
Biebersteinii, C. A. Mey. Verz. Pfl. Cauc. 104 = Phelypaea coccinea.
biflorum, G. Don, Gen. Syst. iv. 634 = Aphyllon uniflorum.
fasciculatum, G. Don, l. c. 633 = Aphyllon fasciculatum.
Tournefortii, G. Don, l. c. = Phelypaea coccinea.

ANOPLOPHYTUM, Beer, in Flora, xxxvii. (1854) 346 = **Tillandsia**, Linn. (Bromel.).
aëranthos, Beer, Bromel. 40 = T. dianthoidea.
amoenum, E. Morr. ex C. Morr. in Belg. Hortic. xxxiii. (1883) 265. t. 17 = T. pulchra.
Benthamianum, Beer, Bromel. 263 = T. Benthamiana.
bicolor, Beer, l. c. 41 = T. stricta.
calothyrsum, Beer, l. c. 263 = T. calothyrsus.
dianthoideum, Beer, l. c. 41 = T. dianthoidea.
didistichum, E. Morr. ex C. Morr. in Belg. Hortic. xxxi. (1881) 164 = T. Lorentziana.
Duratii, Beer, Bromel. 42 = T. Duratii.
flexuosum, Beer, l. c. 43 = T. flexuosa.
geminiflorum, E. Morr. ex C. Morr. in Belg. Hortic. xxix. (1879) 226, & xxx. 191. t. 111 = T. geminiflora.
guianense, Beer, Bromel. 44.—Guian.
incanum, E. Morr. ex C. Morr. in Belg. Hortic. xxxi. (1881) 209. t. 11 = T. Gardneri.
lineare, Beer, Bromel. 42 = T. linearis.
longebracteatum, Beer, l. c. 264 = T. purpurea.
pulchellum, Beer, l. c. 41 = T. pulchra.
Rollissoni, E. Morr. ex C. Morr. in Belg. Hortic. xxix. (1879) 275.—Bras.
roseum, Beer, Bromel. 40 = T. dianthoidea.
rubidum, Beer, l. c. = T. geminiflora.
setaceum, Beer, l. c. 265 = T. compressum?
stramineum, C. Koch, in Ind. Sem. Hort. Berol. (1856) 7 = T. straminea.
Sprengelianum, Beer, Bromel. 266 = T. Sprengelianum.

ANOPLOPHYTUM:—
strictum, Beer, Bromel. 39 = Tillandsia stricta.
strobilanthum, Beer, l. c. 42 = T. strobilantha.
vestitum, Beer, l. c. 266 = T. Benthamiana.
violaceum, Beer, l. c. = T. violacea.
vittatum, Beer, l. c. 43 = T. vittata.

ANOPTERUS, Labill. Pl. Nov. Holl. i. 85. t. 112 (1804). *SAXIFRAGEAE*, Benth. & Hook. f. i. 648.
glandulosus, *Labill. l. c.* 86.—Tasman.
Macleayanus, *F. Muell. in Journ. Pharm. Soc. Vict.* (1858) 43.—Austral.

ANOSMIA, Bernh. in Linnaea, vii. (1832) 608 = **Smyrnium**, Linn. (Umbell.).
idaea, Bernh. l. c. 609 = S. apiifolium.

ANOSPORUM, Nees, in Linnaea, ix. (1835) 287 = **Cyperus**, Linn.
Cephalotes, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 84 = C. Cephalotes.
Colymbetes, Boeck. in Bot. Zeit. xxvii. (1869) 26 = C. Colymbetes.
cubense, Boeck. l. c. = Scirpus cubensis.
monocephalum, Nees, in Linnaea, ix. (1835) 287 = C. Cephalotes.
nudicaule, Boeck. in Bot. Zeit. xxvii. (1869) 26 = C. nudicaulis.
pallidum, Boeck. in Linnaea, xxxvi. (1869–70) 412 = C. platystylis.

ANOTIS, DC. Prod. iv. 431 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 59.
affinis, Benth. & Hook. f. Gen. ii. 59 = Leschenaultiana.
boerhaavioides, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 165.—China.
calycina, *Hook. f. Fl. Brit. Ind.* iii. 73.—Ind. or.
capitata, Korth. in Ned. Kruid. Arch. ii. (1851) 151 = Hedyotis stipulata?
carnosa, *Benth. & Hook. f. Gen.* ii. 59.—Ind. or.
Cervantesii, DC. Prod. iv. 432 = Hedyotis Cervantesii.
ciliolosa, G. Don, Gen. Syst. iii. 535 = Houstonia purpurea.
coerulea, G. Don, l. c. = Houstonia caerulea.
conferta, DC. Prod. iv. 431 = Mallostoma conferta.
decipiens, *Hook. f. Fl. Brit. Ind.* iii. 72.—Ind. or.
dentelloides, Steud. Nom. ed. II. i. 101 = Oldenlandia dentelloides.
ericoides, DC. Prod. iv. 433 = Hedyotis ericoides.
filiformis, DC. l. c. 431 = Mallostoma filiformis.
Finlaysoniana, *Benth. & Hook. f. Gen.* ii. 60.—Sumatr.
foetida, *Benth. & Hook. f. Gen.* ii. 59.—Ind. or.
gentianoides, Walp. Rep. iv. 230 = Houstonia caerulea.
gracilis, *Hook. f. Fl. Brit. Ind.* iii. 71.—Reg. Himal.
hypnoides, DC. Prod. iv. 432 = Hedyotis hypnoides.
ingrata, *Benth. & Hook. f. Gen.* ii. 59; *et Hook. f. Fl. Brit. Ind.* iii. 71.—Reg. Himal.
juniperifolia, DC. Prod. iv. 432 = Mallostoma juniperifolia.
lanceaefolia, Benth. & Hook. f. Gen. ii. 59 = lancifolia.
lanceolata, DC. Prod. iv. 433 = Houstonia purpurea.
lancifolia, *Hook. f. Fl. Brit. Ind.* iii. 73.—Ind. or.
laricifolia, DC. Prod. iv. 432 = Mallostoma laricifolia.
Lechleriana, Schlecht. in Linnaea, xxviii. (1856) 488 = Oldenlandia Lechleriana.
Leschenaultiana, *Benth. & Hook. f. Gen.* ii. 59; *et Hook. f. Fl. Brit. Ind.* iii. 72.—Ind. or.
longiflora, Benth. Pl. Hartw. 23 = Houstonia triflora.
longifolia, G. Don, Gen. Syst. iii. 535 = Houstonia purpurea.
microphylla, DC. Prod. iv. 432 = Hedyotis microphylla.
mitrasacmoides, *Benth. & Hook. f. Gen.* ii. 60.—Austral.
monosperma, *Benth. & Hook. f. l. c.* 59; *et Hook. f. Fl. Brit. Ind.* iii. 75.—Ind. or.
Montholoni, *Hook. f. Fl. Brit. Ind.* iii. 73.—Ind. or.
nummularia, *Benth. & Hook. f. Gen.* ii. 60; *et Hook. f. Fl. Brit. Ind.* iii. 75.—Zeylan.
nummulariformis, Benth. & Hook. f. l. c. = praec.
oxyphylla, *Hook. f. l. c.* 72.—Ind. or.

ANOTIS:—
pilifera, *Schlecht. in Linnaea*, xxviii. (1856) 487.—Peruv.
pterospora, *Benth. & Hook. f. Gen.* ii. 60.—Austral.
purpurea, G. Don, Gen. Syst. iii. 535 = Houstonia purpurea.
quadrilocularis, *Benth. & Hook. f. Gen.* ii. 59; *et Hook. f. Fl. Brit. Ind.* iii. 74.—Ind. or.
Rheedii, *Benth. & Hook. f. l. c.*; *et Hook. f. l. c.* 73.—Ind. or.
Richardiana, *Benth. & Hook. f. l. c.*; *et Hook. f. l. c.* 75.—Ind. or.
Ritchiei, *Hook. f. Fl. Brit. Ind.* iii. 74.—Ind. or.
rotundifolia, *DC. Prod.* iv. 433.—Am. bor.
Salzmanni, DC. l. c. = Oldenlandia Salzmanni.
serpens, *DC. l. c.*—Am. austr.
serpyllifolia, G. Don, Gen. Syst. iii. 535 = Houstonia serpyllifolia.
setosa, DC. Prod. iv. 432 = Mallostoma setosa.
tenella, G. Don, Gen. Syst. iii. 535 = Houstonia tenella.
Thwaitesiana, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 165.—China.
thymifolia, DC. Prod. iv. 432 = Mallostoma thymifolia.
urophylla, *Benth. & Hook. f. Gen.* ii. 59; *et Hook. f. Fl. Brit. Ind.* iii. 72.—Ind. or.
Wightiana, *Benth. & Hook. f. l. c.* 60; *et Hook. f. l. c.* 75.—Ind. or.

ANPLECTRUM, A. Gray, Bot. U. St. Expl. Exped. i. 597 (1854). *MELASTOMACEAE*, Benth. & Hook. f. i. 758.
APLECTRUM, Blume, in Flora, xiv. (1831) 502.
annulatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 84.—Penang.
assamicum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 546.—Reg. Himal.
barbatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 84.—Burma.
biligulatum, *Triana, l. c.* 85.—Sumatr.
confine, *Blume, Mus. Bot. Lugd. Bat.* i. 38.—Sumatr.
cyanocarpum, Triana, in Trans. Linn. Soc. xxviii. (1871) 84 = glaucum.
divaricatum, *Triana, l. c.*—Ind. or.
glaucum, *Triana, l. c.*—Malaya.
Korthalsii, *Triana, l. c.* 85.—Borneo.
latifolium, *Triana, l. c.*—Borneo.
ligulatum, *Triana, l. c.*—Java.
myrtiforme, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 305 = Medinilla myrtiformis.
myrtifolium, Miq. Fl. Ind. Bat. i. 1. 555 = Medinilla myrtiformis.
nodosum, *Blume, Mus. Bot. Lugd. Bat.* i. 37.—Sumatr.
nodosum, *Blume, l. c.* 37.—Sumatr.
ovalifolium, A. Gray, Bot. U. St. Expl. Exped. i. 597 = Allomorphia ovalifolia.
pallens, *Blume, Mus. Bot. Lugd. Bat.* i. 38.—Malaya.
parviflorum, Benth. Fl. Hongk. 116 = Blastus cochinchinensis.
reformatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 85.—Borneo.
rostratum, *Blume, in Flora*, xiv. (1831) 502.—Malaya.
stipulare, *Blume, l. c.* 503.—Java.
viminale, *Blume, l. c.* 502.—Malaya.

ANQUETILIA, Decne. in Jacquem. Voy. Bot. 161. t. 161 (1848) = **Skimmia**, Thunb. (Rutac.).
Laureola, Decne. l. c. = S. Laureola.

ANREDERA, Juss. Gen. 84 (1789). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 78, 1219.
BERIESA, Steud. Nom. ed. II. i. 199 (1840).
CLAIRISIA, Benth. & Hook. f. Gen. iii. 78 (1880).
CLARISIA, Abat, in Act. Soc. Med. Sev. x. 418; ex Spreng. Gen. i. 202 (1830).
SIEBERA, Presl, in Oken, Isis, xxi. (1828) 275.
Cumingii, Hassk. in Flora, xlviii. (1865) 401 = spicata.
scandens, Moq. in DC. Prod. xiii. II. 230 = spicata.
spicata, *Pers. Syn.* i. 297.—Am. trop.; Ins. Philipp.
vesicaria, Gaertn. f. Fruct. iii. 176. t. 213 = spicata.
vesiculosa, Poir. Encyc. Suppl. i. 391 = spicata.

ANSELLIA, Lindl. Bot. Reg. xxx. (1844) sub t. 12. *ORCHIDEAE*, Benth. & Hook. f. iii. 537.
africana, *Lindl. l. c.*—Afr. trop.
gigantea, *Reichb. f. in Linnaea*, xx. (1847) 673.—Natal.

ANSERINA, Dum. Fl. Belg. 21 (1827) = **Chenopodium**, Linn.
Bonus-Henricus, Dum. l. c. = C. Bonus-Henricus.

ANSONIA, Bert. ex Hemsl. Bot. Voy. Challenger, i. III. 53 (1884) = **Lactoris**, Phil. (Piperac.).
nodulosa, Bert. ex Hemsl. l. c. = L. fernandezia.

ANSONIA, Rafin. New Fl. Am. iv. (1836) 58 = **Amsonia**, Walt. (Apocyn.).
elliptica, Rafin. l. c.—Am. bor.
latifolia, Rafin. l. c. = Amsonia Tabernaemontana.
salicifolia, Rafin. l. c. = Amsonia Tabernaemontana.
tenuifolia, Rafin. l. c.—Am. bor.

ANSTRUTHERIA, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 344. t. 4 = **Weihea**, Spreng. (Rhizoph.).
africana, Benth. in Journ. Linn. Soc. iii. (1859) 78 = W. africana.
ceylanica, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 345 = W. ceylanica.
madagascariensis, Benth. in Journ. Linn. Soc. iii. (1859) 78 = W. madagascariensis.
zeylanica, Henslow, in Hook. f. Fl. Brit. Ind. ii. 441 = W. ceylanica.

ANTACANTHUS, Rich. ex DC. Prod. iv. 484 (1830) = **Scolosanthus**, Vahl (Rubiac.).

ANTAGONIA, Griseb. in Goett. Abh. xix. (1874) 144 = **Cayaponia**, Manso (Cucurb.).
citrullifolia, Griseb. l. c. 145 = C. citrullifolia.
Sandia, Griseb. in Lorentz, Veg. del Nordest. Prov. Entre-Rios, 69 = C. Sandia.

ANTAUREA, Neck. Elem. i. 70 (1790) = **Centaurea**, Linn. (Compos.).

ANTELAEA, Gaertn. Fruct. i. 277. t. 68 (1788). Genus inc. scd.
canescens, *Cels, ex Heynh. Nom.* ii. 38.—Hab. ?
javanica, *Gaertn. Fruct.* i. 277.—Java.

ANTENNARIA, Gaertn. Fruct. ii. 410. t. 167 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 301.
DISYNANTHUS, Rafin. in Am. Monthly Mag. (1818) 268.
alpina, *Gaertn. l. c.*—Reg. bor.
andina, *A. Gray, in Proc. Am. Acad.* v. (1861) 138.—Am. austr.
aretioides, *A. Gray, l. c.*—Peruv.
argentea, *Benth. Pl. Hartw.* 319.—Calif.
carpathica, *R. Br. in Trans. Linn. Soc.* xii. (1818) 123.—Reg. temp. bor.
chilensis, *Remy, in C. Gay, Fl. Chil.* iv. 235.—Chili.
cinnamomea, DC. Prod. vi. 270 = Anaphalis cinnamomea.
contorta, D. Don, in Bot. Reg. t. 605 = Anaphalis contorta.
dimorpha, *Torr. & Gray, Fl. N. Am.* ii. 431.—Am. bor.
dioica, *Gaertn. Fruct.* ii. 410. t. 167. f. 3.—Europ.; As. bor.
flagellaris, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 212.—Am. bor.
Geyeri, *A. Gray, Pl. Fendl.* 107.—Am. bor.
hyperborea, D. Don, in Engl. Bot. Suppl. t. 2640 = dioica.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 178 = Anaphalis margaritacea.
japonica, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 126; Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 200 = praec. ?
javanica, *DC. Prod.* vi. 271.—Java.
labradorica, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 406 = alpina.
leontopodina, DC. Prod. vi. 269 = Leontopodium sibiricum.
Leontopodium, Gaertn. Fruct. ii. 410 = Leontopodium alpinum.
leucophylla, Holl. & Reichb. in Flora, xiii. (1830) 382 = Helichrysum obconicum.
lincarifolia, *Wedd. Chlor. And.* i. 150.—Bolivia.
luzuloides, *Torr. & Gray, Fl. N. Am.* ii. 430.—Am. bor. occ.
magellanica, *Sch. Bip. in Flora*, xxxviii. (1855) 117.—Reg. Magell.
margaritacea, R. Br. in Trans. Linn. Soc. xii. (1817) 123 = Anaphalis margaritacea.

ANTENNARIA:—
Meredithae, F. Muell. in Papers Roy. Soc. Tasm. (1870) 15 = Leontopodium Meredithae.
microcephala, *A. Gray, in Proc. Am. Acad.* x. (1874) 74.—Calif.
monocephala, DC. Prod. vi. 269 = alpina.
monoica, *Wedd. Chlor. And.* i. 150.—N. Granat.
montana, S. F. Gray, Nat. Arr. ii. 458 = dioica.
muscoides, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 100.—Reg. Himal.
nana, *Hook. f. & Thoms. ex C. B. Clarke, l. c.*—Reg. Himal.
nivalis, Tenore, ex Nym. Consp. 383 = Leontopodium alpinum.
nubigena, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 45 = Raoulia catipes.
parvifolia, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 406 = dioica.
Planchoni, F. Muell. Census, 80 = Raoulia Planchoni.
plantaginea, *R. Br. in Trans. Linn. Soc.* xii. (1817) 123.—Am. bor.
plantaginifolia, Hook. Fl. Bor. Am. i. 330 = plantaginea.
racemosa, *Hook. l. c.*—Am. bor.
rubicunda, C. Koch, in Linnaea, xvii. (1843) 49 = Helichrysum arenarium.
saxatilis, *DC. Prod.* vi. 271.—Java.
semidecurrens, DC. l. c. = Anaphalis araneosa.
sordida, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 124.—Java.
Steetziana, *Turcz. in Bull. Soc. Nat. Mosc.* (1857) I. Add. p. xxxix.—Dahuria.
stenophylla, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 213.—Am. bor.
triplinervis, Sims, Bot. Mag. t. 2468 = Anaphalis triplinervis.
uniceps, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 105.—Austral.
villosissima, D. Don, Prod. Fl. Nep. 175 = Anaphalis araneosa.
viscida, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 124.—Malaya.

ANTENORON, Rafin. Fl. Ludov. 28 (1817) = **Polygonum**, Linn.
racemosum, Rafin. l. c. = P. virginianum.

ANTEPHORA, Steud. Syn. Pl. Gram. 111 (1855) = **Anthephora**, Schreb. (Gram.).

ANTERISCIUM, Meyen, Reise, i. 315 (1834) sphalm. = **Asteriscium**, Cham. & Schlecht. (Umbellif.).

ANTHACANTHA, Lem. Illustr. Hortic. (1858) Misc. 64 = **Euphorbia**, Linn.
Desmetiana, Lem. l. c. = E. hexagona.

ANTHACANTHUS, Nees, in DC. Prod. xi. 460 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1097.
CRATEOLA, Rafin. Fl. Tellur. iv. 64 (1836).
OPLONIA, Rafin. l. c.
acicularis, *Nees, l. c.*—Jamaica.
armatus, *Nees, l. c.*—Jamaica.
bispinosus, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 525.—Ind. occ.
cuneatus, *Nees, in DC. Prod.* xi. 461.—Jamaica.
cuneifolius, Nees, l. c. 432 = cuneatus.
emarginatus, *Nees, l. c.* 461.—Ins. S. Thom.
jamaicensis, *Griseb. Fl. Brit. W. Ind.* 457.—Jamaica.
lycioides, *Nees, in DC. Prod.* xi. 461.—Ind. occ.
microphyllus, *Nees, l. c.*—Ind. occ.
purpurascens, *Griseb. Cat. Pl. Cub.* 198.—Cuba.
repandus, *Nees, in DC. Prod.* xi. 462.—Ins. Tanna.
sinuatus, *Nees, l. c.*—Ins. Tanna.
spinosus, *Nees, l. c.* 460.—Am. austr.
Sprengelii, *Nees, l. c.* 461.—Ins. S. Doming.
tetrastichus, *Griseb. Cat. Pl. Cub.* 197.—Cuba.
vincoides, *Nees, in DC. Prod.* xi. 461.—Madag.

ANTHACANTHUS, Rich. ex DC. Prod. iv. 484 (1830) = **Scolosanthus**, Vahl (Rubiac.).

ANTHACTINIA, Bory, in Ann. Gén. Sc. Phys. ii. (1819) 139 = **Passiflora**, Linn.
Horsfieldii, M. Roem. Syn. Pepon. 191 = P. Horsfieldii.
longipes, Bory, in Ann. Gén. Sc. Phys. ii. (1819) 139 = P. longipes.

ANTHACTINIA:—
moluccana, M. Roem. Syn. Pepon. 190 = Passiflora moluccana.
nepalensis, M. Roem. l. c. = P. nepalensis.
penangiana, M. Roem. l. c. 192 = Modecca nicobarica.
singaporiana, M. Roem. l. c. = Modecca singaporiana.
sumatrana, M. Roem. l. c. 191 = P. sumatrana.
timorensis, M. Roem. l. c. = P. moluccana.
Walkerii, M. Roem. l. c. = P. suberosa.

ANTHADENIA, Lem. Hort. Vanhoutt. fasc. i. 4 (1845); Fl. des Serres, ii. (Avr. 1846) 10. t. 6 = **Sesamum**, Linn. (Pedalin.).
sesamoides, Lem. ll. cc.—Afr. trop.

ANTHAENANTIA, Beauv. Agrost. 48. t. 10. f. 7 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1098.
ANTHENANTIA, R. Br. in Flind. Voy. ii. App. III. 582 (1814).
AULAXANTHUS, Ell. Sketch Bot. Carol. i. 102 (1817).
AULAXIA, Nutt. Gen. Am. i. 47 (1818).
LEPTOCORYPHIUM, Nees, Agrost. Bras. 83 (1824).
lanata, *Benth. in Journ. Linn. Soc.* xix. (1882) 39.—Mexic.
rufa, *Schult. Mant.* ii. 258.—Am. bor.
villosa, *Beauv. Agrost.* 48. t. 10. f. 7.—Am. bor.

ANTHAERIUM, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 179 *err. typ.* = **Anthurium**, Schott (Aroid.).

ANTHALOGEA, Rafin. New Fl. Am. iv. 88 (1836) = **Polygala**, Linn.
polygama, Rafin. l. c. = P. rubella.

ANTHANEMA, Rafin. Fl. Tellur. iv. 90 (1836) = **Cuscuta**, Linn. (Convolv.).
capitata, Rafin. l. c. = C. capitata.
paradoxa, Rafin. l. c. = C. glomerata.
verticillata, Rafin. l. c. = C. glomerata.

ANTHANOTIS, Rafin. Fl. Ludov. 52, 149 (1817) = **Asclepiodora**, A. Gray (Asclep.).
pedicellata, Rafin. l. c. 52 = Podostigma pubescens.
procumbens, Rafin. l. c. 149.—Am. bor.
viridis, Rafin. l. c. 52 = Asc. viridis.

ANTHEIDOSORUS, A. Gray, in Hook. Kew Journ. iii. (1851) 173 = **Myriocephalus**, Benth. (Compos.).
gracilis, A. Gray, l. c. = M. gracilis.

ANTHEILEMA, Rafin. Fl. Tellur. iv. 66 (1836) = **Phaylopsis**, Willd. (Acanth.).
imbricata, Rafin. l. c. = P. longifolia.
paniculata, Rafin. l. c. = Ruellia paniculata.

ANTHEISCHIMA, Korth. Verh. Nat. Gesch. Bot. ed. Temminck, 137 t. 27 (1840) = **Gordonia**, Ellis (Ternstroem.).
excelsa, Korth. l. c. = G. excelsa.

ANTHELIA, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 127 (1863–64). *AROIDEAE*, Benth. & Hook. f. iii. 963.
nobilis, *Schott, l. c.*—Celebes.

ANTHELIS, Rafin. Chl. Aetn. 9 (1813) = **Helianthemum**, Tourn. (Cistin.).
albidus, Rafin. l. c.—Italia.
arenaria, Rafin. New Fl. Am. iii. 31.—Am. bor.
glaucus, Rafin. Chl. Aetn. 9 = H. glaucum.
glutinosus, Rafin. l. c. = H. glutinosum.
laevipes, Rafin. l. c. = H. laevipes.
podanisia, Rafin. New Fl. Am. iii. 30.—Am. bor.

ANTHELMINTHICA, P. Br. Hist. Jamaic. 156 (1756) = **Spigelia**, Linn. (Logan.).

ANTHEMA, Medic. Malv. 42 (1787) = **Lavatera**, Tourn. (Malvac.).
arborea, Medic. l. c. = L. arborea.
cretica, Medic. l. c. = L. cretica.
scabra, Moench, Meth. 612 = L. cretica.
Tenoreana, Presl, Fl. Sicul. i. 181 = L. cretica.

ANTHEMIOPSIS, Boj. ex DC. Prod. v. 547, 548 (1836) = **Wedelia**, Jacq. (Compos.).
elongata, Boj. l. c. 548 = W. elongata.
macrophylla, Boj. l. c. 547.—Afr. trop.

ANTHEMIS, Mich. ex Linn. Syst. ed. I. (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 420.
CHAMAEMELUM, Tourn. ex Adans. Fam. ii. 128 (1763).
CHAMOMILLA, Godr. Fl. Lorr. ii. 19 (1843).
COTA, J. Gay, in Guss. Fl. Sic. Syn. ii. 866 (1844).
LEPIDOPHORUM, Neck. Elem. i. 14 (1790).
LEUCANTHEMUM, Adans. Fam. ii. 127 (1763).
LYONNETIA, Cass. in Dict. Sc. Nat. l. 106 (1827).
MARCELIA, Cass. l. c. xxxiv. 107 (1825).
MARUTA, S. F. Gray, Nat. Arr. ii. 456 (1821).
MATRICARIA, Tourn. ex Adans. Fam. ii. 127 (1763).
ORMENIS, Cass. in Dict. Sc. Nat. xxix. 180 (1823).
PERIDERAEA, Webb, Iter Hisp. 37 (1838).
RHETINOLEPIS, Coss. in Bull. Soc. Bot. Fr. iii. 707 (1856).

abrotanifolia, *Guss. Fl. Sic. Syn.* ii. 490.—Italia; Lusitan.
abrotanifolia, Willk. in Bot. Zeit. v. (1847) 858 = montana.
absinthifolia, *Boiss. & Sprun. Diagn.* Ser. I. iv. 8.—Caria.
abyssinica, J. Gay, ex A. Rich. Tent. Fl. Abyss. i. 418 = A. Cotula.
abyssinica, *Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop.* 142, nomen.—Abyss.
aciphylla, *Boiss. Diagn.* Ser. I. iv. 7.—As. Min.
adonidifolia, *Boiss. Fl. Orient.* iii. 317.—Armenia.
aetnensis, *Schouw, in Spreng. Syst.* iii. 595.—Mont. Aetna.
afra, Burm. f. Fl. Cap. Prod. 27 = Osmites dentata.
afra, *Linn. Syst.* ed. X. 1223.—Afr. austr.
ageratifolia, Sibth. & Sm. Fl. Graec. ix. 68. t. 887 = Achillea ageratifolia.
agrestis, *Presl, Fl. Sic.* p. xxix, nomen.—Sicilia.
agrestis, Wallr. Sched. Crit. 484 = arvensis.
aizoides, Boiss. & Orph. ex Hook. f. Bot. Mag. t. 6117 = Achillea ageratifolia.
Aizoon, Griseb. Spicil. Fl. Rumel. ii. 210 = Achillea ageratifolia.
albida, *Boiss. Diagn.* Ser. I. iv. 10.—As. Min.
alpina, Gouan, Fl. Monsp. 370 = montana.
alpina, *Linn. Cent.* ii. 31. n. 195.—Europ.
alpina, Moric. ex Nym. Consp. 360 = carpatica.
altaica, *Fisch. Hort. Gorenk.* ed. II. (1812) 41.—Sibir. alt.
altissima, Boiss. Diagn. Ser. I. xi. 12 = coelopoda.
altissima, Bell. in Mém. Acad. Torin. v. (1790–91) 245 = peregrina.
altissima, Guss. ex Nym. Consp. 359 = brachycentros.
altissima, *Linn. Sp. Pl.* 893.—Europ.; Oriens.
americana, Linn. l. c. 895 = Chrysanthellum procumbens.
americana, Mutis, in Linn. f. Suppl. 378 = Spilanthes Mutisii.
ammophila, *Boiss. & Heldr. Diagn.* Ser. I. xi. 9.—Pamphyl.
Anacyclus, J. Gay, ex Boiss. Fl. Orient. iii. 299 = cretica.
anatolica, *Boiss. Diagn.* Ser. I. xi. 10.—As. Min.
anemonifolia, Boj. ex Steud. Nom. ed. II. i. 102 = Epallage anemonifolia.
anglica, Spreng. Syst. iii. 594 = arvensis.
apennina, *Tausch, in Syll. Ratisb.* ii. 250.—Europ.
apiifolia, R. Br. in Bot. Reg. t. 527 = parthenoides.
arabica, Linn. Sp. Pl. 896 = Cladanthus arabicus.
arabica, Viv. Fl. Lib. Spec. 56 = Cyrtolepis alexandrina.
arenicola, *Boiss. Fl. Orient.* iii. 306.—Cilicia.
argaea, Boiss. & Bal. Diagn. Ser. II. vi. 97 = ptarmicaeformis.
artemisiaefolia, Willd. Sp. Pl. iii. 2184 = Chrysanthemum sinense.
artemisifolia, Willd. in Ges. Naturf. Fr. Neue Schr. iii. (1801) 431 = praec.?
arvensis, *Linn. Sp. Pl.* ed. I. 894.—Europ.; Oricns.
arvensis, Pall. ex Bieb. Fl. Taur. Cauc. ii. 330 = altissima.
asperula, *Bertol. Fl. Ital.* ix. 368.—Corsica.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 289.—Afr. bor.
atrata, *Schur, ex Heimerl. in Denkschr. Acad. Wien*, xlviii. (1884) 137.—Europ. austr.
Aucheri, Boiss. Diagn. Ser. I. xiii. 77, sphalm. = Achillea Aucheri.
aurea, *Brot. Fl. Lusit.* i. 394.—Europ. austr.
aurea, DC. Prod. vi. 7 = nobilis.
aurea, Webb, ex Nym. Consp. 375 = Matricaria aurea.
auriculata, *Boiss. Diagn.* Ser. I. iv. 5.—As. Min.
australis, Sibth. & Sm. Fl. Graec. ix. 67 = cypria.
australis, Willd. Sp. Pl. iii. 2177 = incrassata.

ANTHEMIS:—
austriaca, *Jacq. Fl. Austr.* v. 22.—Europ.; Oriens.
Barrelieri, Tenore, Fl. Nap. ii. 245 = Achillea Barrelieri.
Baumgarteniana, Schur, Enum. Pl. Transs. 334 = montana.
Bellidiastrum, Linn. Syst. ed. X. 1223 = Osmites Bellidiastrum.
biaristata, DC. Fl. Fr. iv. 204 = Anacyclus clavatus.
Biebersteiniana, *C. Koch, in Linnaea*, xxiv. (1851) 320.—Oriens.
Blancheana, *Boiss. Fl. Orient.* iii. 292.—Syria.
Bollei ×, Asch. ex Nym. Consp. ii. 363 = A. Cotula.
Bourgei, *Boiss. & Reut. Pugill. Pl. Hispan.* 56.—Hispan.
Boveana, *J. Gay, in Expl. Sc. Algér.* t. 60. f. 2.—Algeria.
Bovei, Willk. & Lange Prod. ii. 91 = praec.
Brachmanni, *Boiss. & Heldr. in Boiss. Diagn.* Ser. I. vi. 84.—Graecia.
brachycentros, *J. Gay, in Koch, Syn. Fl. Germ.* ed. II. 414.—Graecia; As. Min.
brachyglossa, C. Koch, in Linnaea, xxiv. (1851) 318 = tinctoria.
buphthalmoides, Jacq. Hort. Schoenb. ii. 13 = Heliopsis buphthalmoides.
byzantina, C. Koch, in Linnaea, xxiv. (1851) 321 = montana.
caespitosa, Herbich, in Flora, xl. (1857) 509 = tenuifolia.
cairica, Vis. Fl. Quaed. Aegypt. 36. t. 6.—Aegypt.; Arab.
callichroa, Boiss. Diagn. Scr. I. xiii. 77, sphalm. = Achillea callichroa.
camptocarpa, Boiss. Fl. Orient. iii. 319 = tripolitana.
candicans, *Boiss. l. c.* 293.—Lycia.
candidissima, *Willd. ex Spreng. Syst.* iii. 593.—Oriens; Lycia.
canescens, *Brot. Fl. Lusit.* i. 395.—Lusit.
canescens, Tenore, ex Nym. Consp. 359 = brachycentros.
carnea, *Boiss. Diagn.* Ser. I. iv. 4.—As. Min.
carpathica, *Willd. Sp. Pl.* iii. 2179.—Europ.; As. Min.
Cassia, *Boiss. Diagn.* Ser. I. xi. 10.—Graecia.
caucasica, Adam, in Weber & Mohr, Beitr. i. 71 = Chrysanthemum caucasicum.
caucasica, Henckel, in Donnersm. Pl. Select. H. Halens. 9 = tinctoria.
caucasica, Hornem. Hort. Hafn. 832 = Triumfetti.
caucasica, C. A. Mey. Verz. Pfl. Cauc. 75 = iberica.
Chamomilla, Tenore, Syll. 440 = brachycentros.
Chamomilla, *Willd. Enum. Hort. Berol.* 910.—'Europ. austr.'
Chamomilla-romana, Crantz, Inst. i. 300 = nobilis.
chamomilloides, Steven, ex Bieb. Fl. Taur. Cauc. ii. 331 = candidissima.
chia, *Linn. Sp. Pl.* 894.—Graecia; As. Min.
chilensis, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 247.—Chili.
chronia, Nym. Syll. Suppl. 2 = montana.
chrysantha, *J. Gay, in Expl. Sc. Algér.* t. 60. f. 2.—Afr. bor.
chrysantha, Schur, Enum. Pl. Transs. i. 331 = tinctoria.
chrysocephala, Boiss. & Reut. Diagn. Ser. II. iii. 25 = montana.
chrysoleuca, *J. Gay, ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. 50.—Algeria.
ciliata, *Boiss. Fl. Orient.* iii. 312.—Reg. Caucas.
cinerea, *Panč. Elem.* 39.—Europ. austr.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 50.—Afr. bor.
clavata, Desf. Fl. Atlant. ii. 287 = Anacyclus clavatus.
clavata, *Guss. Fl. Sic. Syn.* ii. 865.—Sicilia.
coarctata, Sibth. & Sm. Fl. Graec. ix. 68 = tinctoria.
coelopoda, *Boiss. Diagn.* Ser. I. xi. 12.—As. Min.; Syria.
collina, Jord. Cat. Jard. Dijon, (1848) 18 = montana.
concinna, Guss. Fl. Sic. Syn. ii. 870 = intermedia.
confusa, *Pomel, Nouv. Mat. Fl. Atl.* 49.—Afr. bor.
coronopifolia, *Willd. Sp. Pl.* iii. 2176.—Hispan.
Cornucopiae, *Boiss. Diagn.* Scr. I. xi. 8.—Palaest.
corymbosa, Haenke, in Jacq. Coll. ii. 73 = Achillea atrata.
corymbulosa, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 316.—Mesopot.
Cossoniana, Reichb. f. Ic. Fl. Germ. xvi. 63. t. 1000 = austriaca.
cossyrensis, Guss. Fl. Sic. Syn. ii. 870 = secundiramea.
Cota, Koch, Syn. Fl. Germ. 414 = brachycentros.
Cota, *Linn. Sp. Pl.* 893.—Europ. austr.
Cota, Savi, ex DC. Prod. vi. 11 = austriaca.

ANTHEMIS:—
Cota, Sibth. & Sm. Fl. Graec. t. 880 = mixta.
Cotula, Blanco, Fl. Filip. ed. I. 638 = Eclipta erecta.
Cotula, *Linn. Sp. Pl.* 894.—Europ.; Afr. bor.; Oriens.
Cotula-foetida, Crantz, Inst. i. 300 = A. Cotula.
cotuloides, Rafin. ex DC. Prod. vi. 13 = A. Cotula.
crassifolia, Humb. ex Steud. Nom. ed. II. i. 102 = Ivа chciranthifolia.
crassipes, Boiss. Diagn. Scr. I. xi. 9 = hyalina.
cretica, *Linn. Sp. Pl.* 895.—Reg. Mediterr.
cronia, Boiss. & Heldr. Diagn. Ser. II. iii. 22 = montana.
Cupaniana, Tod. ex Nym. Consp. 360 = montana.
cypria, *Boiss. Fl. Orient.* iii. 300.—Ins. Cyprus.
cyrenaica, *Coss. in Bull. Soc. Bot. Fr.* xix. (1872) 82.—As. Min.
dalmatica, *Scheele, in Linnaea*, xviii. (1844) 464.—Europ. austr.
damascena, *Boiss. & Gaill. in Boiss. Fl. Orient.* iii. 315.—Syria.
dentata, Boj. ex DC. Prod. vi. 4 = Epallage dentata.
dentata, DC. Prod. vi. 291, in syn. = Osmites anthemoides.
dentata, Moench, Meth. 606 = austriaca.
deserti, *Boiss. Fl. Orient.* iii. 305.—Arab.
diffusa, Salzm. ex DC. Prod. vi. 5 = arvensis.
discoidea, *Spreng. Nachtr.* i. *Bot. Gart. Halle*, 11.—Hab.?
discoidea, *Willd. Sp. Pl.* iii. 2187 = tinctoria.
divergens, *Boiss. Fl. Orient.* iii. 300.—Cappadoc.
drepanensis, Huet, ex Nym. Consp. 361 = maritima.
dubia, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 380 = arvensis.
Duriaei, Delile, Ind. Sem. Hort. Monsp. (1843) ex Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 50 = chrysoleuca.
erucaefolia, *J. & C. Presl, Delic. Prag.* 102.—Sicilia.
euxina, *Boiss. Fl. Orient.* iii. 282.—Reg. Caucas.
fallax, Willd. Enum. Hort. Berol. Suppl. 60 = fuscata.
fimbriata, *Boiss. Fl. Orient.* iii. 306.—Cilicia.
foetida, Lam. Fl. Fr. ii. 164 = A. Cotula.
fruticosa, C. Sm. ex DC. Prod. vi. 14 = Gonospermum revolutum.
fruticosa, *Linn. Cent. Pl.* ii. 31. n. 196; *Amoen. Acad.* iv. 330.—Afr. trop.
fruticulosa, *Bieb. Fl. Taur. Cauc.* ii. 329.—Reg. Caucas.
fumariaefolia, *Boiss. Fl. Orient.* iii. 304.—Cappadoc.
fumarioides, *Hochst. in Lorent, Wanderungen*, 330.—Armcn.
fungosa, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 318.—Persia.
fuscata, *Brot. Fl. Lusit.* i. 394.—Reg. Mediterr. occ.
Garcini, Burm. f. Fl. Ind. 183 = Anvillea Garcini.
Gayana, *Boiss. Fl. Orient.* iii. 313.—Persia; Beluchist.
Gemmellari, *Tineo, Pl. Rar. Sicil.* 47.—Mont. Etna.
Gerardiana, Jord. Obs. Pl. Crit. vii. (1849) 31 = montana.
glabra, Jacquem. ex Decne. in Jacquem. Voy. Bot. iv. 88 = Allardia glabra.
globosa, Orteg. Hort, Matr. 46 = Zaluzania globosa.
granatensis, Boiss. Elench. 60 = arvensis.
grandiflora, Host, Fl. Austr. ii. 506 = carpatica.
grandiflora, Jacquem. Voy. Bot. iv. 87 = Allardia tomentosa.
grandiflora, *Des Moul. in Act. Linn. Soc. Bord.* xx. (1855) 556.—Gallia.
grandiflora, Ramatuelle, in Journ. Hist. Nat. Par. ii. (1792) 234 = Chrysanthemum sinense.
granulata, *Pomel, Nouv. Mat. Fl. Atl.* 290.—Afr. bor.
graveolens, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) (Sert. Fl. Trans.) 40 = carpatica.
graveolens, *Boiss. Fl. Orient.* iii. 295.—Lycia.
Guicciardii, Heldr. & Sart. ex Boiss. Fl. Orient. iii. 308 = peregrina.
halimifolia, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 284.—Algeria.
halophila, *Boiss. & Bal. in Boiss. Fl. Orient.* iii. 285.—Cilicia.
Haussknechtii, *Boiss. & Reut. l. c.* 310.—Syria; Mesopot.
Haynaldi, Janka, in Oestr. Bot. Wochenbl. vi. (1856) 1 = arvensis.
hebronica, *Boiss. & Kotschy, Diagn.* Ser. II. v. 108.—Palaest.
hemisphaerica, Herbich, Stirp. Rar. Bucov. 59 = macrantha.
hemistephana, *Boiss. Fl. Orient.* iii. 314.—Persia.

ANTHEMIS :—
heracleotica, Boiss. & Heldr. ex Nym. Consp. 362 = peregrina.
heterophylla, *Ball, in Journ. Linn. Soc.* xvi. (1878) 507.—Morocco.
heterophylla, *Wallr. Beitr. Fl. Hercyn.* 281.—Europ.
hirpina, Tenore, ex Guss. Fl. Sic. Syn. ii. 868 = montana.
hispanica, Zuccagni, in Roem. Collect. 155 = mixta.
hondensis, Willd. ex Steud. Nom. ed. II. i. 102 = Wedelia acapulcensis.
hungarica, Hornem. Hort. Hafn. Suppl. 101 = austriaca.
hyalina, *DC. Prod.* vi. 4.—Syria; Mesopot.
hydruntina, *Groves, in Journ. Linn. Soc.* xxi. (1885) 533.—Ital.
iberica, *Bieb. Fl. Taur. Cauc.* ii. 328.—Reg. Caucas.
incisa, *Viviani, Fl. Ital.* 9.—Italia.
inconspicua, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xxxviii. (1865) i. 409 = Matricaria discoidea.
incrassata, Link, Enum. Hort. Berol. ii. 345 = Anacyclus clavatus.
incrassata, Loisel. Not. 129 = arvensis.
indurata, *Delile, Fl. Egypte*, 274. t. 47. f. 3.—Aegypt.
inodora, Gilib. Fl. Lituan. i. 223 = arvensis.
intermedia, *Guss. Fl. Sic. Syn.* ii. 491.—Sicil.
italica, *Linn. Syst.* ed. X. 1223.—Ital.
kahirica, Boiss. Fl. Orient. iii. 316 = cairica.
Kitaibelii, DC. Prod. vi. 7 = carpatica.
Kitaibelii, Spreng. Syst. iii. 592 = montana.
Kotschyana, *Boiss. Diagn.* Ser. II. iii. 23.—Cilicia.
lanuginosa, Moench, Meth. 606 = Anacyclus radiatus.
laxa, *Spreng. Fl. Hal. Mant.* 52.—Hab.?
leucantha, Linn. Pl. Rar. Afr. 23 = Osmites anthemoides.
leucanthemifolia, *Boiss. & Blanch. Diagn.* Ser. II. iii. 20.—Syria.
leucanthemoides, *Boiss. l. c.* Ser. I. iv. 9.—Smyrna.
leucocephala, Schur, Enum. Pl. Transs. 334 = montana.
libanotica, DC. Prod. vi. 9 = A. chia.
Lindleyi, Nym. Syll. 7 = montana.
lithuanica, Bess. ex DC. Prod. vi. 13 = A. Cotula.
littoralis, Roj.-Clem. Ens. Vit. And. 307 = maritima.
lutescens, Cerv. in La Llave et Lex. Nov. Veg. Desc. fasc. i. 30 = Zaluzania angusta.
lyonnetioides, *Boiss. Fl. Orient.* iii. 286.—Syria.
macedonica, *Boiss. & Orph. Diagn.* Ser. II. vi. 97.—Macedon.
macrantha, *Heuff. in Flora*, xvi. (1833) 362.—Austria.
maritima, Urville, Enum. Pl. Orient. 114 = secundiramea.
maritima, *Linn. Sp. Pl.* 893.—Reg. Mediterr.
maritima, Pall. ex Bieb. Fl. Taur. Cauc. ii. 330 = ruthenica.
maritima, Sibth. & Sm. Fl. Graec. ix. 64 = tinctoria.
Marschalliana, Willd. Sp. Pl. iii. 2187 = Bieberstein iana.
Melampodia, Spreng. Syst. iii. 592 = seq.
melampodina, *Delile, Fl. Egypte*, 268. t. 45.—Aegypt.; Arab.
melanacme, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 315.—Mesopot.
melanoloma, Trautv. in Bull. Soc. Nat. Mosc. xli. (1868) i. 461 = rigescens.
membranacea, Labill. Ic. Pl. Syr. dec. iii. 14 = Achillea membranacea.
Micheliana, Bertol. Cat. Sem. Hort. Bonon. (1838); et Fl Ital. ix. 364 = maritima.
Micheliana, Guss. Fl. Sic. Syn. ii. ii. 868, obs. = montana.
micrantha, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 300.—Assyria.
microsperma, *Boiss. & Kotschy, Diagn.* Ser. II. v. 108.—Aegypt.; Arab.
millefolia, Linn. Sp. Pl. 896 = Chrysanthemum millefoliatum.
millefoliata, Pall. Reise Statthalt. Russ. Reich. i. 87, 129 = pracc.
minuta, *Lehm. ex Steud. Nom.* ed. II. i. 103.—Hab.?
misella, 'H. S.' ex Nym. Consp. 361 = parnesia.
mixta, *Linn. Sp. Pl.* 894.—Reg. Mediterr.
monantha, Willd. Sp. Pl. iii. 2187 = tinctoria.
monilicostata, *Pomel, Nouv. Mat. Fl. Atl.* 41.—Afr. bor.
montana, Koch, Syn. Fl. Germ. ed. II. 415 = carpatica.
montana, Ledeb. Fl. Ross. ii. 523 = sterilis.
montana, *Linn. Sp. Pl.* ed. II. 1261.—Europ.; Oriens.

ANTHEMIS :—
montana, Sibth. & Sm. Prod. Fl. Graec. ii. 190 = carpatica.
mucronulata, C. Koch, in Linnaea, xxiv. (1851) 319 = Chrysanthemum caucasicum.
mucronulata, Bertol. Amoen. 46 = Achillea Barrelierii.
mucronulata, Hort. ex Steud. Nom. ed. II. i. 103 = Anacyclus clavatus.
mucronulata, Reichb. Fl. Germ. Excurs. 227 = carpatica.
multicaulis, Boiss. & Heldr. in Boiss. Fl. Orient. iii. 301 = arvensis.
multicaulis, *Janka, in Oestr. Bot. Wochenbl.* vi. (1856) 363.—Europ. austr.
murieata, *Guss. Fl. Sic. Syn.* ii. 490.—Sicil.
mysorensis, Herb. Madr. ex DC. Prod. v. 551 = Guizotia oleifera.
Neilreichii, Ortm. in Flora, xxxv. (1852) 676 = ruthenica.
nicaeënsis, Willd. Sp. Pl. iii. 2182 = arvensis.
nigrescens, Vest, in Flora, iii. (1820) 8 = austriaca.
nigricans, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 365 = rigescens.
nobilis, *Linn. Sp. Pl.* 894.—Europ.
occidentalis, Willd. Sp. Pl. iii. 2185 = Spilanthes Mutisii.
odontostephana, *Boiss. Diagn.* Ser. I. vi. 85.—Persia; Ind. or.
odorata, Ait. Hort. Kew. iii. 238 = Ursinia odorata.
odorata, Lam. Fl. Fr. ii. 163 = nobilis.
oppositifolia, Lam. Encyc. i. 576 = Spilanthes Mutisii.
ormenioides, Boiss. Diagn. Ser. I. iv. 5 = Wiedemanniana.
ovatifolia, Orteg. Hort. Matr. 73 = Heliopsis buphthalmoides.
oxylepis, *Boiss. Fl. Orient.* iii. 282.—Cilicia.
oxyloba, Schur, Enum. Pl. Transs. 884, sphalm. = Achillea oxyloba.
paecilolepis, Boiss. Fl. Orient. iii. 288 = Kotschyana.
palestina, *Reut. ex Boiss. l. c.* 283.—As. Min.; Syria.
pallescens, Heldr. ex Nym. Consp. ii. 359 = tinctoria.
pamphylica, *Boiss. & Heldr. Diagn.* Ser. I. xi. 8.—As. Min.
parnassica, Nym. Syll. Suppl. 1 = tinctoria.
Parnassii, Boiss. & Heldr. ex Nym. Consp. 359 = tinctoria.
parnesia, *Boiss. & Heldr. in Boiss. Fl. Orient.* iii. 305.—Graecia.
parthenioides, *Benth. ex Hornem. Hort. Hafn.* ii. 833.—Hab.?
pauciloba, *Boiss. Diagn.* Ser. I. vi. 83.—Syria; Mesopot.
pectinata, *Boiss. & Reut. l. c.* Ser. I. iv. 6.—As. Min.
peduncularis, *Benth. & Hook. f. Gen.* ii. 422.—Afr. bor.
pedunculata, Bory & Chaub. Fl. Pélop. 59 = Anacyclus clavatus.
pedunculata, Desf. Fl. Atl. ii. 288 = Anacyclus pedunculatus.
pentelica, Boiss. ex Nym. Consp. 360 = montana.
peregrina, Bieb. Fl. Taur. Cauc. ii. 331 = candidissima.
peregrina, *Linn. Syst. Nat.* ed. X. 1223.—Ital.; Graecia; As. Min.
persepolitana, *Boiss. Diagn.* Ser. I. xi. 7.—Persia.
persica, *Boiss. l. c.* Ser. I. vi. 85.—Persia.
Pestalozzae, *Boiss. l. c.* Ser. I. xi. 12.—Lycia.
petraea, Tenore, Fl. Nap. ii. 246 = carpatica.
philistea, *Boiss. Fl. Orient.* iii. 302.—Palestin.
piscinalis, *Durieu, ex Munby, Fl. Algér.* 95, nomen.—Algeria.
platyglossa, C. Koch, in Linnaea, xxiv. (1851) 319 = rigescens.
plebeia, *Boiss. & Noë, in Boiss. Fl. Orient.* iii. 313.—Kurdist.
poecilolepis, Boiss. Diagn. Ser. II. iii. 24 = Kotschyana.
pontica, Urville, Enum. Pl. Orient. 114 = ruthenica.
pontica, Sibth. & Sm. Fl. Graec. ix. 66 = tricolor.
pontica, Willd. Sp. Pl. iii. 2184 = montana.
praecox, Link, ex Willd. l. c. 2182 = fuscata.
prolifera, Pers. Syn. ii. 467 = Cladanthus arabicus.
pseudo-atrata, Schur, Enum. Pl. Transs. 327 = tenuifolia.
pseudocota, Visiani, Fl. Dalm. ii. 78 = brachycentros.
pseudocotula, *Boiss. Diagn.* Ser. I. vi. 86.—Oriens.
psorosperma, Tenore, Viag. Abruz. 128 = A. Cotula.

ANTHEMIS :—
ptarmicaefolia, Adam, in Weber & Mohr, Beitr. i. 71 = Achillea grandiflora.
ptarmicaeformis, *C. Koch, in Linnaea*, xxiv. (1851) 319.—As. Min.
pterygantha, C. Koch, l. c. xvii. (1843) 44 = candidissima.
pubescens, Willd. Sp. Pl. iii. 2177 = Anacyclus clavatus.
punctata, Guss. ex Nym. Consp. 360 = montana.
punctata, *Vahl, Symb. Bot.* ii. 91. t. 46.—Atlas; Sicilia.
purpurascens, Nym. Consp. 363 = Anacyclus radiatus.
pygmaea, Bertol. in Nov. Comm. Acad. Bonon. vi. (1843) 237 (Misc. Bot. ii. 23) = Matricaria praecox.
pygmaea, *Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop.* 142, nomen.—Afr. trop.
pyrenaica, *Miégev. in Bull. Soc. Bot. Fr.* xii. (1865) 397.—Mont. Pyren.
pyrenaica, Sch. Bip. in Oestr. Bot. Wochenbl. iv. (1854) 164 = austriaca.
pyrethriformis, Schur, Enum. Pl. Transs. 335 = carpatica.
Pyrethrum, Gouan, Fl. Monsp. 370 = montana.
Pyrethrum, Linn. Sp. Pl. 895 = Anacyclus Pyrethrum.
ramosa, Link, ex Spreng. Syst. iii. 593 = A. Cotula.
Raseheyana, *Boiss. Diagn.* Ser. I. xi. 8.—Syria.
reflectens, Reichb. f. Ic. Fl. Germ. xvi. 61. t. 1006 = arvensis.
reflexa, Moench, Meth. 605 = altissima.
repanda, *Linn. Sp. Pl.* 895.—Lusitan.
repens, Walt. Fl. Carol. 211 = Spilanthes repens.
Requienii, Sch. Bip. ex Nym. Consp. 361 = ruthenica.
retusa, Delile, Fl. Aeg. Illustr. 74 = cairica.
retusa, Link, ex Spreng. Syst. iii. 593 = ruthenica.
retusa, Tenore, ex Nym. Consp. 359 = brachycentros.
Reuteriana, Boiss. Diagn. Ser. I. iv. 8 = smyrnaea.
revoluta, Chr. Sm. ex Buch, in Berl. Acad. Abh. 1816-17 (1819) 370 = Gonospermum revolutum.
rhodensis, *Boiss. Diagn.* Ser. I. xi. 11.—Ins. Rhodus.
rigescens, *Willd. Hort. Berol.* i. t. 62.—Reg. Caucas.
rosea, Jacquem. Voy. Bot. iv. 88 = Allardia glabra.
rosea, *Sibth. & Sm. Fl. Graec.* t. 887.—Cyprus.
rotata, *Boiss. Fl. Orient.* iii. 318.—Aegypt.; Arab.
Rudolphiana, Adam, in Weber & Mohr, Beitr. i. 72 = Biebersteiniana.
rupestris, Boj. ex Steud. Nom. ed. II. i. 103 = Epallage rupestris.
ruthenica, *Bieb. Fl. Taur. Cauc.* ii. 330. n. 1772.—Europ. austr. or.; Reg. Caucas.
ruthenica, Sch. Bip. ex Schur, Enum. Pl. Transs. 332 = austriaca.
salvifolia, Boj. ex DC. Prod. v. 3 = Epallage salvifolia.
santolinoides, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 284.—Algeria.
saxatilis, Baumg. ex Schur, Enum. Pl. Transs. 334 = Baumgarteniana.
saxatilis, Bieb. Fl. Taur. Cauc. iii. 581 = sterilis.
saxatilis, DC. ex Willd. Enum. Hort. Berol. 910 = montana.
scaposa, J. Gay, in Boiss. Fl. Orient. iii. 313 = Gayana.
scariosa, Boiss. Diag. Ser. I. xi. 7 = persepolitana.
scariosa, *DC. Prod.* vi. 4.—Syria; Mesopot.
scariosa, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 263.—Syria.
schizostephana, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 314.—Kurdist.
Schurii, Heimerl, in Denkschr. Acad. Wien, xlviii. (1884) 137 = Achillea oxyloba.
secundiramea, *Bivona, Sicul. Pl. Cent.* ii. 10.—Sicil.; Aegypt.
segetalis, Tenore, Fl. Neap. Syll. App. v. 54 = brachycentros.
sessilis, Salisb. Prod. 204 = Cladanthus arabicus.
Sibthorpii, *Griseb. Spicil. Fl. Rumel.* ii. 210.—Graecia.
sicula, Guss. Fl. Sic. Syn. ii. 866 = arvensis.
sinuata, Cerv. in La Llave et Lex. Nov. Veg. Desc. fasc. i. 30 = Zaluzania triloba.
Sismondaeana, *Clem. in Mem. Accad. Sc. Torino*, Ser. II. xvi. (1857) 297 (*Sert. Orient.* 61).—Byzant.
smyrnaea, *Boiss. Diagn.* Ser. I. iv. 7.—Smyrna.
sphacelata, Willk. in Bot. Zeit. v. (1847) 858 = Anacyclus clavatus.

ANTHEMIS :—
spbacelata, *J. & C. Presl, Delic. Prag.* 103.—Sicilia.
Spruneri, Boiss. & Heldr. Diagn. Ser. II. iii. 24 = montana.
sterilis, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) II. 379.—Tauria.
stiparum, *Pomel, Nouv. Mat. Fl. Atl.* 48.—Afr. bor.
stipulacea, Moench, Meth. Suppl. 258 = Chrysanthemum sinense.
styriaca, Vest, in Syll. Ratisb. i. 12 = carpatica.
suaveolens, *Schwaegr. ex Steud. Nom.* ed. II. i. 103.—Hab.?
subulicola, *Pomel, Nouv. Mat. Fl. Atl.* 49.—Afr. bor.
suffruticosa, *Hornem. Hort. Hafn. Suppl.* 101.—Hab.?
sulphurea ×, Wall. ex Nym. Consp. 363 = A. Cotula.
Tatrae, *Kit. ex Schult. Fl. Austr.* ii. 534.—Europ.
taygetea, *Boiss. & Heldr. Diagn.* Ser. I. vi. 83.—Graecia.
tenella, *Host, Fl. Austr.* ii. 507.—Europ.
tenuifolia, *Schur, in Verh. Siebenb. Ver. Naturw.* ii. (1851) 171.—Transsylvan.
tenuiloba, Boiss. Diagn. Ser. II. iii. 25 = montana.
tenuisecta, *Ball, in Journ. Bot.* xi. (1873) 365.—Marocco.
tenuisecta, *Boiss. Fl. Orient.* iii. 301.—Armenia.
tenuisecta, *Pomel, Nouv. Mat. Fl. Atl.* 50.—Algeria.
tenuispherica, *Herbich, Stirp. Rar. Bucov.* 59.—Austria.
tigrensis, *J. Gay, ex A. Rich. Tent. Fl. Abyss.* i. 418.—Abyss.
tinctoria, *Linn. Sp. Pl.* 896.—Europ.; Oriens.
tomentosa, Gouan, ex Boiss. Fl. Orient. iii. 321 = Anacyclus clavatus.
tomentosa, *Linn. Sp. Pl.* 893.—As. Min.
tomentosa, O. F. Muell. in Fl. Dan. xii. t. 685 = arvensis.
tomentosa, Linn. ex Boiss. Fl. Orient. iii. 291 = montana.
tomentosa, Willd. Sp. Pl. iii. 2176 = peregrina.
tricolor, *Boiss. Fl. Orient.* iii. 288.—Cyprus.
triloba, Orteg. Hort. Matr. 72 = Zaluzania triloba.
trinervia, *Raeusch. Nom.* ed. III. 249.—Am. merid.
tripolitana, *Boiss. & Blanch. Diagn.* Ser. II. iii. 22.—Syria.
Triumfetti, All. Fl. Pedem. i. 187 = austriaca.
Trotzkiana, *Claus, in Beitr. Pflanz. Russ. Reich.* viii. 299.—Turkestan.
tubaeformis, Guss. Fl. Sic. Syn. ii. 870 = Anacyclus clavatus.
tuberculata, *Boiss. Elench.* 59. n. 107.—Hispan.; Afr. bor.
tubicina, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 319.—Mesopot.
ubensis, *Pomel, Nouv. Mat. Fl. Atl.* 289.—Afr. bor.
umbilicata, *Boiss. & Huet, Diagn.* Ser. II. iii. 24.—Armenia.
Urvillaeana, Sch. Bip. ex Nym. Consp. ii. 362 = tomentosa.
Urvillaei, Nym. Syll. 7 = cretica.
valentina, Linn. Sp. Pl. 895 = Anacyclus radiatus.
viridis, Blanco, Fl. Filip. ed. II. 436 = Eclipta erecta.
Visianii, Weiss, ex Boiss. Fl. Orient. iii. 311 = cbia.
vulgaris, Fl. Dan. ex Steud. Nom. ed. I. 53 = Matricaria inodora.
vulgaris, Linn. ex Steud. Nom. l. c. = Matricaria Chamomilla.
Wiedemanniana, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1835) 27.—As. Min.

ANTHEPHORA, Schreb. Beschr. Graes. ii. 105. t. 44 (1779). *GRAMINEAE*, Benth. & Hook. f. iii. 1122.
ANTEPHORA, Steud. Syn. Pl. Gram. 111 (1855).
HYPUDAEURUS, Reicbb. Nom. 37 (1841).
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 389 = Hochstetteri.
axilliflora, Steud. Syn. Pl. Gram. 111 = Buchloë dactyloides.
Belangeri, Steud. l. c. = Hilaria cenchroides.
cuspidata, *Anderss. Enum. Pl. Ins. Galap.* 141.—Ins. Galap.
elegans, Rupr. ex Steud. Syn. Pl. Gram. 111 = Hochstetteri.
elegans, *Schreb. Beschr. Graes.* ii. 105. t. 44.—Mexico.

ANTHEPHORA :—
Hochstetteri, *Nees, in Flora*, xxvii. (1844) 249.—Abyss.
Kotschyi, Hochst. ex Steud. Syn. Pl. Gram. 111 = Hochstetteri.
persica, *Boiss. Diagn.* Ser. I. xiii. 44.—Persia.
pubescens, Nees, Fl. Afr. Austr. i. 74 = Cenchrus pubescens.
punctulata, *Steud. Syn. Pl. Gram.* i. 111.—Mexico.
villosa, Spreng. Neue Entdeck. iii. 14 = elegans.

ANTHERICLIS, Rafin. in Am. Monthly Mag. (1819) 195; Journ. Phys. lxxxix. (1819) 261 = **Tipularia**, Nutt. (Orchid.).

ANTHERICUM, Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 788.
DILANTHES, Salisb. Gen. Pl. Fragm. 70 (1866).
HESPERANTHES, S. Wats. in Proc. Am. Acad. xiv. (1879) 241.
LILIAGO, Salisb. Gen. Pl. Fragm. 70 (1866).
PESSULARIA, Salisb. l. c. (1866).
PHALANGANTHUS, Sebrank, ex Haw. Syn. Pl. Succ. 67 (1819).
PHALANGIUM, Juss. Gen. 52 (1789).
TRACHYANDRA, Kunth, Enum. Pl. iv. 573 (1843).
BLEPHANTHERA, ENDOGONA, LICINIA, OBSITILA, TRACHINEMA, Rafin. Fl. Tellur. ii. 27, 59; iii. 57.
adenanthera, Forst. f. Prod. 24 = Dianella ensifolia.
affine, *Baker, in Journ. Bot.* x. (1872) 138.—Afr. austr.
Aitoni, Baker, in Journ. Linn. Soc. xv. (1876) (= Bulbinella, Gen. Pl.) 294.—Afr. austr.
albucoides, Ait. Hort. Kew. i. 449 = Ornithogalum suaveolens.
aloifolium, Salisb. Prod. 250 = Bulbine alooides.
alooides, Linn. Sp. Pl. 311 = Bulbine alooides.
altissimum, Mill. Gard. Dict. ed. VIII. n. 8 = Bulbine longiscapa.
amoenum, Salisb. Prod. 250 = A. Liliago.
anceps, *Baker, in Journ. Linn. Soc.* xv. (1876) 305.—Afr. austr.
anceps, Soland. ex Baker, l. c. 363 = Tricoryne anceps.
andongense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 257.—Afr. trop.
angulicaule, *Baker, in Journ. Linn. Soc.* xv. (1876) 305.—Afr. austr.
angustifolium, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 332.—Abyss.
annuum, Linn. Sp. Pl. 311 = Bulbine annua.
annuum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 203 = Asphodelus tenuifolius.
aphyllum, *Forsk. Fl. Aegypt. Arab.* 209.—Arabia.
arenarium, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 259.—Afr. trop.
asperatum, *Baker, in Journ. Bot.* x. (1872) 138.—Afr. austr.
asphodeloides, Ait. Hort. Kew. i. 450 = Bulbine longiscapa.
asphodeloides, Linn. Sp. Pl. 311 = Bulbine aspbodeloides.
asphodelum, *Forsk. Fl. Aegypt. Arab.* 209.—Ins. Rhodus.
auratum, *Fisch. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 281.—Soongaria.
autumnale, Scop. Fl. Carn. ed. II. i. 247 = Scilla autumnalis.
baeticum, Steud. Nom. ed. II. i. 103 = A. Liliago.
benguellense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 257.—Angola.
bicolor, Desf. Fl. Atl. i. 304 = Morgagnia bicolor.
bifolium, Scop. Fl. Carn. ed. II. i. 247 = Scilla bifolia.
bipedunculatum, Jacq. Coll. Suppl. 88 = triflorum.
blepharophoron, Schult. f. Syst. vii. 461 = ciliatum.
boeticum, Boiss. Voy. Espagne, ii. 619. t. 172 = A. Liliago.
brasiliense, *Baker, in Journ. Linn. Soc.* xv. (1876) 306.—Bras.
Brehmeanum, *Baker, in Journ. Bot.* x. (1872) 139.—Afr. austr.
brevicaule, *Baker, in Journ. Linn. Soc.* xv. (1876) 298.—Afr. austr.
brevifolium, *Thunb. Prod. Pl. Cap.* 62.—Afr. austr.
bulbosum, R. Br. Prod. 275 = Bulbine bulbosa.

ANTHERICUM :—
Burkei, Baker, in Journ. Bot. x. (1872) 140 = Bulbinella Burkei.
caeruleum, Ruiz & Pav. Fl. Per. iii. 67. t. 299 = Pasitbea caerulea.
calyculatum, Georgi, It. i. 206 = Tofieldia cernua.
calyculatum, Linn. Sp. Pl. 311 = Tofieldia calyculata.
calyculatum, Pall. Reise, iii. 33 = Tofieldia palustris.
calyptrocarpum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 258.—Afr. trop.
Cameroni, *Baker, in Journ. Linn. Soc.* xv. (1876) 96, 314.—Afr. trop.
canaliculatum, *Ait. Hort. Kew.* i. 448.—Afr. austr.
capillare, Spreng. Syst. ii. 83; Schult. f. Syst. vii. 457 = Bulbinella capillaris.
carnosum, Baker, in Journ. Linn. Soc. xv. (1876) 296 (= Bulbinellae sp. Gen. Pl.).—Hab.?
Cauda-felis, Linn. f. Suppl. 202 = Bulbinella caudata.
caudatum, Thunb. Prod. Pl. Cap. 63 = Bulbinella caudata.
caulescens, *Baker, in Journ. Linn. Soc.* xv. (1876) 303.—Afr. trop.
cernuum, *Hornem. Hort. Hafn.* i. 332.—Hab.?
Chamaemoly, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 332 = Chlorophytum tetraphyllum.
ciliatum, *Linn. f. Suppl.* 202.—Afr. austr.
ciliatum, Schult. f. Syst. vii. 466 = Humboldtii.
ciliatum, *Spreng. l. c.* ii. 84.—Venezuela.
ciliolatum, Baker, in Journ. Bot. x. (1872) 137 = Bulbinella ciliolata.
cirrhatum, Forst. f. Prod. n. 148 = Arthropodium cirrhatum.
coarctatum, Ruiz & Pav. Fl. Per. iii. 67 = Excremis ramosa.
colubrinum, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 256 = Dasystachys colubrina.
comosum, Thunb. Prod. Pl. Cap. 63 = Chlorophytum Sternbergianum.
contortum, *Linn. f. Suppl.* 202.—Afr. austr.
Cooperi, *Baker, in Journ. Linn. Soc.* xv. (1876) 304.—Natal.
corymbosum, *Baker, in Journ. Bot.* xv. (1877) 71.—Afr. trop.
costatum, *Andr. Bot. Rep.* t. 298.—Afr. austr.
crispum, Thunb. Prod. Pl. Cap. 63 = Chlorophytum crispum.
croceum, *Schult. f. Syst.* vii. 476.—Georgia.
dendroides, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 249 = Asphodeline prolifera.
dissitiflorum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 257.—Angola.
divaricatum, Jacq. Hort. Schoenb. iv. 7 = revolutum.
Dorsetii, Hort. ex Baker, in Journ. Linn. Soc. xv. (1876) 301 = ramosum.
Dregeanum, *Baker, in Journ. Bot.* x. (1872) 139.—Afr. austr.
Drimiopsis, *Baker, in Journ. Linn. Soc.* xv. (1876) 301.—Afr. trop.
eccremorrhizum, *Ruiz & Pav. Fl. Per.* iii. 67.—Peruv.
ecbeandioides, *Baker, in Bot. Mag.* cxi. (1885) t. 6809.—Mexic.?
elatum, Ait. Hort. Kew. i. 448 = Chlorophytum elatum.
elongatum, Willd. Sp. Pl. 136 = Jacquinianum.
ensatum, Banks & Soland. ex Hook. f. Fl. N. Zeal. i. 255 = Dianella intermedia.
ensiforme, *Vell. Fl. Flum.* 143; iii. t. 162.—Bras.
ericetorum, Berger. Fl. Pyr. ii. 168 = Simethis bicolor.
esculentum, Spreng. Syst. ii. 84 = Camassia esculenta.
exuviatum, Jacq. Coll. Suppl. 89 = Urginea exuviata.
falcatum, *Linn. f. Suppl.* 202.—Afr. austr.
falcatum, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 256 = Dasystachys falcata.
fasciculatum, *Baker, in Journ. Linn. Soc.* xv. (1876) 316.—Afr. austr.
fastigiatum, Steud. Nom. ed. I. 53 = Chlorophytum elatum.
favosum, Thunb. Prod. Pl. Cap. 63 = Bulbine favosa.
filifolium, Jacq. Coll. Suppl. 93 = Urginea filifolia.
filifolium, Thunb. Prod. Pl. Cap. 62 = Urginea nematodes.
filiforme, Ait. Hort. Kew. i. 451 = Bulbinella triquetra.

ANTHERICUM :—
filiforme, Thunb. Prod. Pl. Cap. 62 = Jacquinianum.
fimbriatum, Steud. Nom. ed. I. 35 = Dichopogon Sieberianus?
fimbriatum, Thunb. Prod. Pl. Cap. 63 = longifolium.
flagelliforme, Baker, in Journ. Bot. x. (1872) 140 = Schizobasis flagelliformis.
flavescens, *Schult. f. Syst.* vii. 1692.—Mexic.
flexifolium, Jacq. Coll. Suppl. 93 = Jacquinianum.
flexifolium, *Linn. f. Suppl.* 201.—Afr. austr.
flexuosum, R. Br. ex Benth. Fl. Austral. vii. 50 = Corynotheca dichotoma.
floribundum, Ait. Hort. Kew. i. 447 = Bulbinella latifolia.
fragrans, Jacq. Hort. Schoenb. i. 45 = Urginea fragrans.
frutescens, Linn. Sp. Pl. 310 = Bulbine frutescens.
fruticosum, Salisb. Prod. 250 = pracc.
fugax, Moris, Elench. i. 46 = Urginea fugax.
Gerrardi, *Baker, in Journ. Bot.* x. (1872) 137.—Afr. austr.
glaucum, *Ruiz & Pav. Fl. Per.* iii. 68.—Peruv.
Gmelinianum, Schult. f. Syst. vii. 481 = Melanthium sibiricum.
gracile, Baker, in Journ. Bot. x. (1872) 137 = Bulbinella gracilis.
gracile, Salisb. Prod. 249 = ramosum.
gracillimum, *Regel, in Act. Hort. Petrop.* vi. (1879) 289.—Hab. ?
graecum, Linn. Sp. Pl. ed. II. 444 = Lloydia graeca.
gramineum, *Vell. Fl. Flum.* 143; iii. t. 161.—Bras.
gramineum, *Baker, in Journ. Linn. Soc.* xv. (1876) 302.—Madag.
graminifolium, Willd. Sp. Pl. ii. 139 = undulatum.
grandiflorum, Salisb. Prod. 250 = Paradisea Liliastrum.
graminiformc, Dulac, Fl. Hautes-Pyr. 115 = Paradisea Liliastrum.
Grantii, *Baker, in Trans. Linn. Soc.* xxix. (1875) 160. t. 103 B.—Afr. trop.
graptophyllum, *Baker, in Gard. Chron.* (1882) i. 460.—Ins. Socotra.
Heynei, Rottl. ex Baker, in Journ. Linn. Soc. xv. (1876) 323 = Chlorophytum Heyneanum.
hirsutum, *Thunb. Prod. Pl. Cap.* 63.—Afr. austr.
hispidum, *Linn. Sp. Pl.* ed. II. 446.—Afr. austr.
Hookeri, Colenso, ex Hook. f. Handb. New Zeal. Fl. 286 (= Bulbinellae sp. Gen. Pl.).—N. Zel.
Humboldtii, *Hemsl. Biol. Centr. Am. Bot.* iii. 374.—Mexic.
humile, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 331.—Abyss.
hyacinthoides, Willd. ex Kunth, Enum. Pl. iv. 336 = Scilla indica.
inconspicuum, *Baker, in Journ. Bot.* xv. (1877) 71.—Afr. trop.
incurvum, Thunb. Prod. Pl. Cap. 62 = Bulbine rostrata.
intermedium, Willk. in Willk. & Lange, Prod. Fl. Hisp. i. 202 = A. Liliago.
intricatum, Baker, in Journ. Bot. x. (1872) 140 = Schizobasis intricata.
involucratum, *Baker, in Journ. Linn. Soc.* xv. (1876) 311.—Afr. austr.
Jacquinianum, *Schult. f. Syst.* vii. 462.—Afr. austr.
japonicum, *Thunb. Fl. Jap.* 138.—China; Japon.
lagopus, Thunb. Prod. Pl. Cap. 63 = Bulbine longiscapa.
latifolium, Linn. f. Suppl. 202 = Bulbinella latifolia.
latifolium, Schult. f. Syst. vii. 477 = simplex.
latifolium, Banks & Soland. ex Kunth, Enum. Pl. iv. 622 = Arthropodium cirrhatum.
leptophyllum, *Baker, in Journ. Linn. Soc.* xv. (1876) 317.—Mexic.
Liliago, *Linn. Sp. Pl.* 310.—Europ.; Afr. bor.
Liliastrum, Linn. l. c. ed. II. 445 = Paradisea Liliastrum.
limosum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 257.—Afr. trop.
longepedunculatum, *Steud. ex Schult. f. Syst.* vii. 457, 1692.—Afr. austr.
longifolium, A. Rich. Tent. Fl. Abyss. ii. 333 = Chlorophytum longifolium.
longifolium, *Jacq. Coll. Suppl.* 92.—Afr. austr.
longipes, *N. E. Br. in Gard. Chron.* (1882) ii. 297.—Bras.
longiscapum, Jacq. Coll. Suppl. 84 = Bulbine longiscapa.

ANTHERICUM :—
longistylum, *Baker, in Journ. Linn. Soc.* xv. (1876) 305.—Afr. austr.
Macowani, *Baker, l. c.* 309.—Afr. austr.
macrophyllum, A. Rich. Tent. Fl. Abyss. ii. 334 = Chlorophytum macrophyllum.
Makoyanum, *Regel, Gartenfl.* (1880) 36.—Hab.?
marginatum, Thunb. Prod. Pl. Cap. 63 = Urginea marginata.
Mattiazzi, Vand. Fasc. Pl. 12 = Simethis bicolor.
mesembryanthemoides, Baker, in Journ. Bot. x. (1872) 136 = Bulbine mesembryanthemoides.
milleflorum, [DC. in] Red. Lil. t. 58 = Arthropodium paniculatum.
minus, Steud. Nom. ed. I. 53 = Arthropodium minus.
molle, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 259.—Angola.
monophyllum, *Baker, in Journ. Bot.* xvi. (1878) 324.—Afr. trop.
muricatum, *Linn. f. Suppl.* 202.—Afr. austr.
nanum, *Baker, in Journ. Linn. Soc.* xv. (1876) 305.—Mexic.
nematodes, Schult. f. Syst. vii. 472 = Urginea nematodes.
nepalense, Spreng. Syst. iv. Cur. Post. 135 = Chlorophytum nepalense.
nidulans, *Baker, in Journ. Linn. Soc.* xv. (1876) 314.—Afr. trop.
Nimmonii, J. Grah. Cat. Pl. Bomb. 220 = Chlorophytum glaucum.
niveum, Spreng. Syst. ii. 84; Schult. f. in Roem. & Schult. Syst. vii. 480 = Chlorophytum tuberosum.
non-ramosum, Gilib. Exercit. ii. 465 = A. Liliago.
nubicum, *Baker, in Journ. Linn. Soc.* xv. (1876) 301.—Afr. trop.
Nuttallianum, *Schult. f. Syst.* vii. 477.—Am. bor.
nutans, Jacq. Coll. Suppl. 86 = Bulbine nutans.
nutans, *Thunb. Prod. Pl. Cap.* 63.—Afr. austr.
Oatesii, *Baker, in Journ. Bot.* xvi. (1878) 324.—Afr. trop.
orchideum, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 258.—Afr. trop.
ornithogaloides, Baker, in Journ. Linn. Soc. xv. (1876) 294 = Bulbinella ornithogaloides.
ornithogaloides, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 332 = Chlorophytum tuberosum.
ossifragum, Georgi, Beschreib. Russ. Reich. iii. iv. 907 = Tofieldia cernua.
ossifragum, Linn. Sp. Pl. 311 = Narthecium ossifragum.
pachyphyllum, *Baker, in Journ. Linn. Soc.* xv. (1876) 304.—Afr. austr.
palustre, Salisb. Prod. 250 = Narthecium ossifragum.
paniculatum, Andr. Bot. Rep. t. 395 = Arthropodium paniculatum.
paradoxum, *Schult. f. Syst.* vii. 459.—Afr. austr.
Parkeri, *Baker, in Journ. Bot.* xx. (1882) 269.—Madag.
parviflorum, Benth. Fl. Hongk. 373 = Chlorophytum laxum.
parvum, *N. E. Br. in Gard. Chron.* (1880) ii. 588.—Bras.
pauciflorum, Thunb. Prod. Pl. Cap. 63 = triflorum.
pendulum, Hornem. Hort. Hafn. i. 333 = Arthropodium paniculatum.
peronatum, Baker, in Journ. Bot. x. (1872) 137 = Bulbinella peronata.
peruvianum, Willd. ex Kunth, Enum. Pl. iv. 596 = Humboldtii.
physodes, Jacq. Coll. Suppl. 94; Ic. t. 418 = Urginea physodes.
pilosum, Jacq. l. c. 87; Ic. t. 416 = Ornithogalum hispidum.
planifolium, Linn. Mant. ii. 224 = Simethis bicolor.
pleiostachyum, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 255 = Dasystachys pleiostachya.
plumosum, Ruiz & Pav. Fl. Per. iii. 68 = Bottionea thysanotoides.
pomeridianum, Ker-Gawl. in Bot. Reg. t. 564 = Chlorogalum pomeridianum.
praemorsum, Jacq. Coll. Suppl. 85 = Bulbine praemorsa.
Pseudo-asphodelus, Jacq. Enum. Vind. 59, 233 = Tofieldia calyculata.
pterocaulon, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 258.—Angola.
pubescens, *Baker, in Journ. Linn. Soc.* xv. (1876) 309.—Afr. austr.

ANTHERICUM :—
pubirhachis, *Baker, l. c.* 302.—Afr. trop.
pudicum, *Baker, l. c.* 308.—Afr. austr.
pugioniformc, Jacq. Coll. Suppl. 83 = Bulbine pugioniformis.
pulchellum, *Baker, in Journ. Bot.* x. (1872) 140.—Afr. austr.
pusillum, Jacq. Coll. Suppl. 95 = Urginea pusilla.
pyrenicarpum, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 259.—Afr. trop.
Quamash, Steud. Nom. ed. 1. 54 = Camassia esculenta.
ramosissimum, *Schult. f. Syst.* vii. 469.—Mexic.
ramosum, Steud. Nom. ed. 1. 54 = Melasphaerula graminifolia.
ramosum, *Linn. Sp. Pl.* 310.—Europ.
reflexum, Cav. Ic. iii. 21 = Echeandia terniflora.
revolutum, *Linn. Sp. Pl.* 310.—Afr. austr.
rigidum, *Baker, in Journ. Bot.* x. (1872) 141.—Afr. austr.
Rossii, *Hook. f. Handb. New Zeal. Fl.* 285 (= Bulbinellae sp. Gen. Pl.).—N. Zel.
rostratum, Jacq. Coll. Suppl. 82 = Bulbine rostrata.
Saltii, *Baker, in Journ. Linn. Soc.* xv. (1876) 309.—Abyss.
sardoum, Muell. ex Kunth, Enum. Pl. iv. 335 = Urginea fugax.
scabrellum, *Baker, in Journ. Linn. Soc.* xv. (1876) 318.—Mexic.
scabrum, *Linn. f. Suppl.* 202.—Afr. austr.
Schultesii, *Baker, in Journ. Bot.* x. (1872) 140.—Afr. austr.
scilliflorum, *Eckl. ex Steud. Nom.* ed. II. i. 248, in syn.; *et ex Baker, in Journ. Linn. Soc.* xv. (1876) 298.—Afr. austr.
scilloides, *Schult. f. Syst.* vii. 477.—Reg. Argent.
semibarbatum, Hook. Bot. Mag. t. 3129 = Bulbine bulbosa.
semibarbatum, R. Br. Prod. 275 = Bulbine semibarbata.
serotinum, Linn. Sp. Pl. ed. II. 444 = Lloydia alpina.
sessiliflorum, Steud. Nom. ed. II. i. 104 = scilliflorum.
setosum, Willd. ex Schult. f. Syst. vii. i. 473 = Bulbinella robusta.
sibiricum, *Hort. Erf. ex Steud. Nom.* ed. II. i. 104.—Sibir.
simplex, *Willd. ex Kunth, Enum. Pl.* iv. 596.—Reg. Colomb.
Skinneri, *Baker, in Journ. Linn. Soc.* xv. (1876) 315.—Am. trop.
sphacelatum, *Baker, l. c.* 303.—Afr. trop.
spirale, Linn. Mant. ii. 224 = Eriospermum spirale.
spiratum, Thunb. Prod. Pl. Cap. 62 = Urginea filifolia.
squameum, Linn. f. Suppl. 202 = hispidum.
steuocarpum, *Baker, in Journ. Linn. Soc.* xv. (1876) 317.—Mexic.
Sternbergianum, Schult. f. Syst. vii. 1693 = Chlorophytum Sternbergianum.
strictum, Steud. ex Steud. Nom. ed. I. 54 = Dichopogon strictus.
subpetiolatum, *Baker, in Journ. Linn. Soc.* xv. (1876) 302.—Afr. trop.
subtrigynum, Jacq. Coll. iii. 273 = Amianthium muscaetoxicum.
succulentum, Salisb. Prod. 250 = Bulbine asphodeloides.
sulphureum, Waldst. & Kit. Pl. Rar. Hung. i. 98 = Ornithogalum pyrenaicum.
superpositum, *Baker, in Journ. Bot.* xvi. (1878) 324.—Afr. trop.
tenellum, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 256.—Afr. trop.
thrysoideum, *Baker, in Journ. Bot.* x. (1872) 139.—Afr. austr.
Torreyi, *Baker, in Journ. Linn. Soc.* xv. (1877) 317.—Mexic.
triflorum, *Ait. Hort. Kew.* i. 448.—Afr. austr.
trinervium, Viviani, Fl. Lib. Spec. 21. t. 8. f. 2 = Lloydia alpina.
tripedale, *Baker, in Journ. Bot.* xx. (1882) 269.—Madag.
triquetrum, Linn. f. Suppl. 202 = Bulbinella triquetra.
tuberosum, Roxb. Pl. Corom. ii. 20. t. 138 = Chlorophytum tuberosum.
undulatum, *Jacq. Coll. Suppl.* 87.—Afr. austr.
undulatum, Thunb. Prod. Fl. Cap. 63 = hispidum.
uniflorum, *Roxb. Hort. Beng.* 87; *Fl. Ind.* ii. 149.—Ind. or.

ANTHERICUM :—
ustulatum, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 258.—Afr. trop.
variegatum, Hort. ex Fl. Mag. N. S. t. 152 = Chlorophytum elatum.
vermiculare, Soland. ex Baker, in Journ. Linn. Soc. xv. (1876) 308 = longepedunculatum.
vespertinum, Jacq. Hort. Schoenb. i. 44 = falcatum.
vestitum, *Baker, in Journ. Linn. Soc.* xv. (1876) 307.—Mexic.
villosum, Labill. Ic. Pl. Syr. Dec. v. 14. t. 9 = Gagea bohemica.
viscosum, *Baker, in Journ. Bot.* x. (1872) 141.—Afr. austr.
Williamsii, Hort. ex Gard. Chron. (1875) i. 490 = Chlorophytum elatum.
yedoënse, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* ii. 529.—Japon.
zanguebaricum, *Baker, in Journ. Linn. Soc.* xv. (1876) 302.—Afr. trop.
Zeyheri, *Baker, in Journ. Bot.* x. (1872) 140.—Afr. austr.

ANTHEROCEPHALA, DC. Prod. vii. 755 (1839) = **Andersonia**, R. Br. (Epacrid.).
Drummondii, DC. l. c. = And. aristata.

ANTHEROCERAS, Bert. in Am. Journ. Sc. xix. (1831) 301 = **Leucocoryne**, Lindl. (Liliac.).
odorata, Bert. (Merc. Chil. 561) in Am. Journ. Sc. xix. (1831) 301.—Chili.
ornithogaloides, Bert. ll. cc.—Chili.

ANTHEROSPERMA, Poir. ex Steud. Nom. ed. II. i. 105 (1840), err. typ. = **Atherosperma**, Labill. (Monim.).

ANTHEROTOMA, Hook. f. in Benth. & Hook. f. Gen. i. 745 (1867). *MELASTOMACEAE*, Benth. & Hook. f. i. 745.
Afzelii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 444.—Afr. trop.
Naudini, *Hook. f. in Benth. & Hook. f. l. c.*—Madag.

ANTHEROTRICHE, Turcz. in Bull. Soc. Nat. Mosc. (1846) ii. 505 = **Anisoptera**, Korth. (Dipterocarp.).
lanceolata, Turcz. l. c. 506 = Anisopt. thurifera.

ANTHERURA, Lour. Fl. Cochinch. 144 (1790). *RUBIACEAE*, Benth. & Hook. f. ii. 29.
rubra, *Lour. l. c.*—Cochinch.; Molucc.

ANTHERYLIUM, Rohr & Vahl, in Skr. Naturh. Selsk. Hafn. ii. (1792) i. 211. t. 8. *LYTHRARIEAE*, Benth. & Hook. f. i. 782.
floribundum, Spreng. Syst. ii. 474 = Adenaria floribunda.
grisleoides, Spreng. l. c. = Adenaria floribunda.
nudiflorum, *Hemsl. Diagn. Pl. Nov.* 13.—Mexic.
purpuratum, Spreng. l. c. = Adenaria floribunda.
Rohrii, *Vahl, in Skr. Naturh. Selsk. Hafn.* ii. (1792) i. 211.—Ind. occ.

ANTHERYTA, Rafin. Sylva Tellur. 95 (1838) = **Tetrazygia**, Rich. (Melast.).
granulosa, Rafin. l. c. 96 = T. granulosa.

ANTHESTERIA, Spreng. Anleit. ii. i. 79 (1817) = **Anthistiria**, Linn. f. (Gramin.).

ANTHILLIS, Neck. Delic. Gallo-Belg. ii. 303 (1768) = **Anthyllis**, Linn. (Legum.).

ANTHIPSIMUS, Rafin. in Journ. Phys. lxxxix. (1819) 105 = **Panicum**, Linn.? (Gramin.).
gonopodos, Rafin. l. c.—Am. bor.

ANTHIRRINUM, Moench, Meth. 521 (1794) = **Antirrhinum**, Tourn. (Scroph.).

ANTHISTIRIA, Linn. f. Nov. Gram. Gen. 35 (1779) (Amoen. Acad. x. 38). *GRAMINEAE*, Benth. & Hook. f. iii. 1136.
Androscepia, Brongn. in Duperr. Voy. Coq. Bot. 77 (1829).
Aristaria, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 296.

ANTHISTIRIA :—
Exotheca, Anderss. in Nov. Act. Upsal. Ser. III. i. (1856) 253.
Germainia, Balansa & Poitrasson, in Bull. Soc. Hist. Nat. Toul. vii. (1873) 344.
Heterelytron, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 294.
Iseilema, Anderss. in Nov. Act. Soc. Upsal. Ser. III. i. (1856) 250.
Perobachne, Presl, Rel. Haenk. i. 348. t. 48 (1830).
Themeda, Forsk. Fl. Aegypt. Arab. 178 (1775).
abyssinica, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 449.—Abyss.
anathera, *Nees, in Steud. Syn. Pl. Gram.* 402.—Ind. or.
andropogonoides, Steud. l. c. = Andropogon bracteatus.
argentea, Nees, Gram. Afr. Austr. 127 = imberbis.
arguens, *Willd. Sp. Pl.* iv. 901.—Ins. Molucc.
arguens, Nees, Fl. Afr. Austr. 124 = imberbis.
arguta, *Nees, ex Steud. Syn. Pl. Gram.* 401.—Ind. or.; Burma.
arundinacea, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 115 = arguens.
arundinacea, Roxb. Hort. Beng. 6; Fl. Ind. i. 251 = gigantea.
australis, R. Br. Prod. i. 200 = imberbis.
avenacea, F. Muell. Fragm. v. 206 = gigantea.
barbata, Desf. in Journ. Phys. xl. 294. t. 2 = ciliata.
Barteri, *Munro, ex Oliver, in Trans. Linn. Soc.* xxix. (1875) 176.—Afr. trop.
basisericea, F. Muell. Fragm. v. 207 = gigantea.
Bladhii, *Wight, in Wall. Cat.* n. 8770.—Ind. or.
brachyantha, Boiss. Diagn. Ser. I. xiii. 71 = imberbis.
caespitosa, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 241 = imberbis.
caudata, Nees, ex Hook. & Arn. Bot. Beech. Voy. 245 = gigantea.
ciliata, Cav. ex Spreng. Syst. i. 291 = australis.
ciliata, *Linn. f. Diss. Nov. Gram. Gen.* 35.—Ind. or.; Afr.; Jamaic.
ciliata, Retz. Obs. fasc. iii. 11 = imberbis.
cimicina, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 182 = prostrata.
cuspidata, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 242 = imberbis.
cymbaria, *Roxb. Hort. Beng.* 6; *Fl. Ind.* i. 251.—Ind. or.
depauperata, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 243 = imberbis.
Desfontainii, Kunth, Rev. Gram. i. 161 = imberbis.
dissoluta, Nees, ex Steud. Syn. Pl. Gram. 400 (est Andropoginis sp. ign.).—Afr. trop.
fasciculata, Thw. Enum. Pl. Zeyl. 366 = Andropogon filipendulus.
flosculosa, F. Muell. Fragm. x. 75 = gigantea?
foliosa, H. B. & K. Nov. Gen. et Sp. i. 191 = Cymbopogon Humboldtii.
Forskalii, Kunth, Rev. Gram. i. 162 = imberbis.
frondosa, R. Br. Prod. i. 200 = arguens.
gigantea, *Cav. Ic.* v. 35.—Ins. Philipp.
glabrifolia, Desv. Opusc. 71 (= 'ciliata, Sw.').—Ind. occ.
glandulosa, Wight, ex Steud. Syn. Pl. Gram. 400 = prostrata.
glauca, Desf. Fl. Atl. ii. 380. t. 254 = imberbis.
heteroclita, Roxb. Hort. Beng. 6; Fl. Ind. i. 249 = Andropogon heteroclitus.
hispida, Thunb. Fl. Cap. i. 403 = imberbis.
Hookeri, Griseb. in Goett. Nachr. (1868) 91 = gigantea.
Humboldtii, Nees, Agrost. Bras. 369 = Cymbopogon Humboldtii.
imberbis, *Retz. Obs.* iii. 11.—Afr. austr.
japonica, Willd. Sp. Pl. iv. 901 = imberbis.
Junghuhniana, Nees, ex Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 233 = arguens.
latifolia, Anderss. in Peters, Reise Mossamb. Bot. 562 = Andropogon lepidus.
laxa, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 243 = imberbis.
Linneana, Steud. Syn. Pl. Gram. 401 = prostrata.
membranacea, *Lindl. in Mitch. Trop. Austr.* 88.—Austral.
multiplex, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 449 = Andropogon multiplex.
mutica, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 117 = gigantea.
paleacea, *Ball, in Journ. Linn. Soc.* xvi. (1878) 734.—Marocco.

ANTHISTIRIA :—
pilifera, Steud. Syn. Pl. Gram. 400 = arguens.
pilosa, Presl, Rel. Haenk. i. 348 = Cymbopogon Humboldtii.
polystachya, Roxb. Hort. Beng. 6; Fl. Ind. i. 248 = imberbis.
prostrata, *Willd. Sp. Pl.* iv. 901.—Ind. or.
pseudo-cymbaria, Steud. Syn. Pl. Gram. 399 = Cymbopogon elegans.
puberula, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 240 = imberbis.
punctata, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 448 = imberbis.
quinqueplex, Steud. Syn. Pl. Gram. 400 = Cymbopogon elegans.
ramosa, Thunb. ex Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 247 = Apluda gigantea.
reflexa, H. B. & K. Nov. Gen. et Sp. i. 191 = Cymbopogon Humboldtii.
scandens, Roxb. Hort. Beng. 6; Fl. Ind. i. 248 = ciliata.
setosa, Thunb. ex Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 233 = arguens.
subglabra, Buese, in Miq. Pl. Jungh. 363 = imberbis.
subsericans, Nees, ex Steud. Syn. Pl. Gram. 401 = gigantea.
syriaca, Boiss. Diagn. Ser. I. xiii. 72 = imberbis.
tortilis, Presl, Rel. Haenk. i. 347 = Andropogon Nardus.
tremula, *Nees, ex Steud. Syn. Pl. Gram.* 401.—Ind. or.
villosa, Lam. Illustr. t. 841. f. 3 = gigantea.
vulcanica, *Steud. Syn. Pl. Gram.* 400.—Ins. Santorin.
vulpina, Anderss. in Nov. Act. Upsal. Ser. III. ii. (1856) 245 = gigantea.
Wightii, Nees, ex Steud. Syn. Pl. Gram. 400 = prostrata.

ANTHOBOLUS, R. Br. Prod. 357 (1810). *SANTALACEAE*, Benth. & Hook. f. iii. 230.
cxocarpoides, *F. Muell. Fragm.* ix. 3.—Austral.
filifolius, *R. Br. Prod.* 357.—Austral.
fovculatus, *F. Muell. Fragm.* i. 212.—Austral.
leptomeroides, *F. Muell. l. c.* 21.—Austral.
triqueter, A. Cunn. ex A. DC. in DC. Prod. xiv. 687 = filifolius.
triqueter, *R. Br. Prod.* 357.—Austral.

ANTHOCEPHALUS, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 237, excl. syn.; Korth. Verh. Nat. Gesch. 154. t. 48 (1840?). *RUBIACEAE*, Benth. & Hook. f. ii. 29.
Cephalidium, A. Rich. l. c. 290.
Cadamba, Miq. Fl. Ind. Bat. ii. 135 = morindaefolius.
chinensis, Hassk. in Flora, xxviii. (1845) 231 = Cephalanthus orientalis.
chinensis, Walp. Rep. ii. 491 = Nauclea purpurea.
indicus, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 238 = Nauclea purpurea.
morindaefolius, *Korth. Verh. Nat. Gesch.* 154. t. 48; Obs. Naucl. Ind. 20.—Sumatr.; Ind. or.

ANTHOCERAS, Baker, in Journ. Linn. Soc. xi. 374 (1871) = ANTHEROCERAS, Bert. = **Leucocoryne**, Lindl. (Liliac.).

ANTHOCERASTES, A. Gray, in Hook. Kew Journ. iv. (1852) 225 = **Toxanthus**, Turcz. (Compos.).
Drummondii, A. Gray, l. c. = T. perpusillus.
Muelleri, Sond. in Linnaea, xxv. (1852) 480 = T. Muelleri.

ANTHOCERCIS, Labill. Nov. Holl. Pl. ii. 19. t. 158 (1806). *SOLANACEAE*, Benth. & Hook. f. ii. 912.
Cyphanthera, Miers, in Ann. Nat. Hist. Ser. II. xi. (1853) 376.
Eadesia, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 71.
albicans, *A. Cunn. in Field, New South Wales, App.* 335. t. 2.—Austral.
amblyantha, F. Muell. Fragm. i. 179 = myosotidea.
angustifolia, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 21.—Austral.
anisantha, *Endl. in Ann. Wien. Mus.* ii. (1838) 201.—Austral.

ANTHOCERCIS:—
arborea, *F. Muell. Fragm.* i. 212.—Austral.
Eadesii, *F. Muell. l. c.* ii. 139.—Austral.
fasciculata, *F. Muell. l. c.* i. 122.—Austral.
genistoides, *Miers, Illustr. S. Am. Pl.* ii. *App.* 27. t. 83 D.—Austral.
glabella, Miers, l. c. = viscosa.
gracilis, *Benth. in DC. Prod.* x. 192.—Austral.
Hopwoodii, *F. Muell. Fragm.* ii. 138.—Austral.
ilicifolia, Hook. Bot. Mag. sub t. 2961 = viscosa.
intricata, *F. Muell. Fragm.* i. 211.—Austral.
Leichhardtii, *F. Muell. l. c.* vi. 142.—Austral.
littorea, Endl. Iconogr. t. 68 = viscosa.
littorea, *Labill. Pl. N. Holl.* ii. 19. t. 158.—Austral.
microphylla, *F. Muell. Fragm.* i. 179.—Austral.
myosotidea, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 20.—Austral.
Odgersii, *F. Muell. Fragm.* x. 19.—Austral.
racemosa, *F. Muell. l. c.* i. 211.—Austral.
scabrella, *Benth. in DC. Prod.* x. 192.—Austral.
spinescens, F. Muell. Fragm. i. 122 = genistoides.
tasmanica, *Hook. f. Fl. Tasm.* i. 289. t. 92.—Austral.
viscosa, *R. Br. Prod.* 448.—Austral.

ANTHOCEROS dissectus, Mont. ex Wedd. in DC. Prod. xvii. 62 (= *Neolacis dissecta*) = **Apinagia**, Tul. (Podost.).

ANTHOCHLAMYS, Fenzl, ex Endl. Gen. 300 (1837). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 58.
PELTISPERMUM, Moq. Chenop. Enum. 102 (1840).
polygaloides, [*Fenzl*?] *Moq. l. c.* 102.—Reg. Caucas.

ANTHOCHORTUS, Endl. Gen. 121 (1837); Kunth, Enum. Pl. iii. 485 (1841) = ANTOCHORTUS, Nees = **Willdenowia**, Thunb. (Restiac.).

ANTHOCHYTRUM, Reichb. f. Ic. Fl. Germ. xix. 39. t. 1432 (1859) = **Crepis**, Vaill. (Compos.).
alpinum, Reichb. f. l. c. 39. t. 81 = C. alpina.

ANTHOCLEISTA, Afzel. ex R. Br. in Tuckey, Congo, App. 449 (1818). *LOGANIACEAE*, Benth. & Hook. f. ii. 795.
madagascariensis, *Baker, in Journ. Bot.* xx. (1882) 173.—Madag.
macrophylla, *G. Don, in Loud. Hort. Brit.* 471.—Afr. trop.
nobilis, *G. Don, Gen. Syst.* iv. 68.—Afr. trop.
scandens, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 16.—Afr. trop.
Vogelii, *Planch. in Hook. Ic. Pl.* t. 793.—Afr. trop.

ANTHOCOMA, Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie, ii. (1845) 569 = **Phlomis**, Linn.? (Labiat.).
flavescens, Zoll. & Mor. l. c. 570.—Malaya.

ANTHOCOMETES, Nees, in DC. Prod. xi. 311 (1847) = **Monothecium**, Hochst. (Acanth.).
aristatus, Nees, l. c. 312 = M. aristatum.

ANTHODENDRON, Reichb. in Moessl. Handb. i. p. xl. 244, 308 (1827) = **Rhododendron**, Linn. (Ericac.).
flavum, Reichb. l. c. 309 = R. flavum.
nudiflorum, Reichb. l. c. = R. nudiflorum.
ponticum, Reichb. Fl. Germ. Excurs. 416 = R. ponticum.
viscosum, Reichb. in Moessl. Handb. ed. II. i. 424 = R. viscosum.

ANTHODISCUS, Endl. Gen. 1091 in syn. (1841) = ANTHODON, Ruiz & Pav. = **Salacia**, Linn. (Hippocrat.).

ANTHODISCUS, G. F. W. Mey. Prim. Fl. Esseq. 193 (1818). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 181.
guianensis, *G. F. W. Mey. l. c.* = Guiana.
peruanus, *Baill. Adansonia*, x. (1872) 241.—Peruv.
trifoliatus, *G. F. W. Mey. Prim. Fl. Esseq.* (1818) 194.—Guiana.

ANTHODON, Ruiz & Pav. Fl. Per. i. 45. t. 74 (1798) = **Salacia**, Linn. (Hippocrat.).
apiculatum, Miers, ex Peyr. in Mart. Fl. Bras. xi. t. 137 = Hippocratea megacarpa.
crassifolium, Mart. in Schult. Mant. i. 348 = S. crassifolia.

ANTHODON:—
decussatum, Ruiz & Pav. Fl. Per. i. 45. t. 74. f. 2 = Hippocratea decussata.
ellipticum, Mart. in Schult. Mant. i. 348 = Salacia elliptica.
glomeratum, Mart. l. c. = S. glomerata.
grandiflorum, Benth. in Hook. Kew Journ. iv. (1852) 10 = S. grandiflora.
grandifolium, Mart. in Schult. Mant. i. 348 = S. grandifolia.
laevigatum, Steud. Nom. ed. II. i. 105 = S. laevigata.
laxiflorum, Benth. in Hook. Kew Journ. iv. (1852) 10 = S. laxiflora.
micranthum, Mart. in Schult. Mant. i. 348 = S. micrantha.
oblongifolium, Mart. l. c. = S. elliptica.
paniculatum, Mart. l. c. = S. paniculata.
Selloanum, Schult. Mant. iii. Add. II. 411.—Bras.
trinerve, Schult. l. c.—Bras.
undulatum, Mart. in Schult. Mant. i. 348 = S. arborea.
verrucosum, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 172.—Cuba.

ANTHODUS, Mart. ex Roem. & Schult. Syst. Mant. i. 253 (1822) = ANTHODON, Ruiz & Pav. = **Salacia**, Linn. (Hippocrat.).

ANTHOGONIUM, Wall. ex Lindl. Introd. Nat. Syst. ed. II. 341 (1836), nomen. *ORCHIDEAE*, Benth. & Hook. f. iii. 515.
gracile, *Wall. Cat.* n. 7398.—Reg. Himal.
Griffithii, Reichb. f. in Bonplandia, ii. (1854) 90 = gracile.

ANTHOLOMA, Labill. Voy. Rech. Pérouse, ii. 235. t. 41 (1800). *TILIACEAE*, Benth. & Hook. f. i. 239.
Billardieri, *Vieill. in Bull. Soc. Linn. Normand.* ix. (1865) 335.—N. Caled.
montana, *Labill. Voy. Rech. Pérouse*, 236. t. 41.—N. Caled.

ANTHOLYZA, Linn. Syst. ed. I. (1735); Gen. ed. I. 10 (1737). *IRIDEAE*, Benth. & Hook. f. iii. 710.
ANISANTHUS, Sweet, Hort. Brit. ed. II. 500 (1830); Brit. Fl. Gard. Ser. II. t. 84 (1838).
CUNONIA, Mill. Gard. Dict. ed. VIII. (1768).
GLADIOLUS, Gaertn. Fruct. i. 31. t. 11. f. 4. (1788).
HOMOGLOSSUM, Salisb. in Trans. Hort. Soc. i. (1812) 325.
PETAMENES, Salisb. l. c. 324.
abbreviata, Pers. Syn. i. 42 = quadrangularis.
abyssinica, *A. Brongn. in A. Rich. Tent. Fl. Abyss.* ii. 306.—Abyss.
aethiopica, *Linn. Syst.* ed. X. 863.—Afr. austr.
aletroides, Burm. f. Fl. Cap. Prod. 1 = Watsonia aletroides.
bicolor, Gasp. ex Tenore, Cat. Orto Nap. (1845) 78; fide E. Morr. in Belg. Hortic. ii. (1852) 145 = aethiopica.
brevifolia, Hort. ex Steud. Nom. ed. II. i. 105 = Gladiolus brevifolius.
brevifolia, Eckl. ex Ball, in Journ. Linn. Soc. xvi. (1878) 179 = caffra.
caffra, *Banks, ex Ker-Gawl. in Konig & Sims, Ann. Bot.* i. (1805) 232.—Afr. austr.
caryophyllacea, Burm. f. Fl. Cap. Prod. 1 = Watsonia humilis.
caryophyllea, Vahl, Enum. Pl. ii. 123 = Watsonia humilis.
cepacea, *Linn. Mant.* ii. 176.—Afr. austr.
coccinea, Spreng. Neue Entdeck. i. 253 = A. Cunonia.
Cunonia, *Linn. Sp. Pl.* 37.—Afr. austr.
fimbriata, Klatt, in Linnaea, xxxv. (1867–68) 299 = Babiana fimbriata.
floribunda, Salisb. in Trans. Hort. Soc. i. (1812) 324 = aethiopica.
fucata, *Baker, in Journ. Linn. Soc.* xvi. (1878) 180.—Afr. austr.
fulgens, Andr. Bot. Rep. t. 192 = Watsonia Meriana.
hirsuta, Klatt, in Linnaea, xxxv. (1867–68) 379 = Gladiolus sericeo-villosus.
hirsuta, Lam. Encyc. ii. 211 = Gladiolus mollis.
hirta, Pers. Syn. i. 42 = Babiana Thunbergii.
huillensis, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 270.—Afr. trop.
hypogaea, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 345.—Afr. austr.

ANTHOLYZA:—
intermedia, *Baker, in Journ. Linn. Soc.* xvi. (1880, nomen.—Afr. austr.
lucidior, *Linn. f. Suppl.* 96.—Afr. austr.
lucidior, Vahl, Enum. Pl. ii. 122 = Watsonia plantaginea.
Ludwigii, Eckl. Verz. 34 = aethiopica?
marginata, Page, ex Steud. Nom. ed. II. i. 106 = Watsonia marginata.
Maura, Linn. Mant. ii. 175 = Witsenia Maura.
Meriana, Blanco, Fl. Filip. ed. I. 24 = Sisyrinchium palmifolium.
Meriana, Linn. Sp. Pl. ed. II. 54 = Watsonia Meriana.
Merianella, Curt. Bot. Mag. t. 441 = Watsonia aletroides.
Merianella, *Murr. Syst.* ed. XIII. 77.—Afr. austr.
Meriania, Linn. Syst. ed. X. 863 = Watsonia Meriana.
montana, [Ker-Gawl. in] Konig & Sims, Ann. Bot. i. (1805) 233 = Gladiolus montanus.
nemorosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 346.—Afr. austr.
nervosa, *Thunb. Prod. Pl. Cap.* 7.—Afr. austr.
orchidiflora, Klatt, in Linnaea, xxxii. (1863) 733 = Gladiolus arenarius.
paniculata, *Klatt, in Linnaea*, xxxv. (1867–68) 379.—Afr. austr.
plicata, Linn. f. Suppl. 96 = Babiana Thunbergii.
praealta, [Del. in] Red. Liliac. vii. t. 387 = aethiopica.
quadrangularis, *Burm. f. Fl. Cap. Prod.* 1.—Afr. austr.
quinquenervia, *Schrank, Comm. Irid. Cap.* 56.—Afr. austr.
ramosa, Klatt, in Linnaea, xxxii. (1863) 734 = Gladiolus montanus.
revoluta, *Burm. f. Fl. Cap. Prod.* 1.—Afr. austr.
ringens, Andr. Bot. Rep. t. 32 = aethiopica.
ringens, Linn. Sp. Pl. 37 = Babiana ringens.
rupestris, Eckl. ex Baker, in Journ. Linn. Soc. xvi. (1877) 179 = caffra.
saccata, *Baker, in Journ. Linn. Soc.* xvi. (1877) 180.—Afr. austr.
spicata, Andr. Bot. Rep. t. 56 = Watsonia brevifolia.
spicata, *Brehm, ex Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 345.—Afr. austr.
spicata, Mill. Gard. Dict. ed. VIII. n. 2 = Gladiolus Milleri.
splendens, *Steud. Nom.* ed. II. i. 106.—Afr. austr.
striata, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 346.—Afr. austr.
tubulosa, Andr. Bot. Rep. t. 174 = Watsonia aletroides.
vittigera, Salisb. in Trans. Hort. Soc. i. (1812) 324 = aethiopica.

ANTHOMELES, M. Roem. Syn. (Rosiflorae) 102 & 140 (1847) = **Crataegus**, Linn. (Rosac.).
aestivalis, M. Roem. l. c. 141 = Cr. aestivalis.
Douglasii, M. Roem. l. c. 140 = Cr. Douglasii.
flava, M. Roem. l. c. 142 = Cr. flava.
flexispina, M. Roem. l. c. = Cr. parvifolia.
glandulosa, M. Roem. l. c. 141 = Cr. flava.
opaca, M. Roem. l. c. = Cr. aestivalis.
rotundifolia, M. Roem. l. c. 140 = Cr. coccinea.
sanguinea, M. Roem. l. c. = Cr. sanguinea.
stipulosa, M. Roem. l. c. 142 = Cr. stipulosa.
subserrata, M. Roem. l. c. = Cr. subserrata.
turbinata, M. Roem. l. c. = Cr. flava.

ANTHONOTA, Beauv. Fl. Owar. et Ben. i. 70. t. 42 (1805) = **Macrolobium**, Schreb. (Legum.).
crassifolia, Baill. Adansonia, vi. (1864–65) 179.—Afr. trop.
elliptica, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 430.—Am. merid.
macrophylla, Beauv. Fl. Owar. et Ben. i. 70. t. 42 = M. Palisoti.

ANTHOPHYLLUM, Steud. Syn. Pl. Cyp. 160 (1855) = **Scirpus**, Linn. (Cyper.).
Urvillei, Steud. l. c. = S. frondosus.

ANTHOPOGON, Neck. Elem. ii. 12 (1790) = **Gentiana** Linn.
barbata, Rafin. Fl. Tellur. iii. 25 = G. barbata.
ciliata, Rafin. l. c. = G. ciliata.
crinita, Rafin. l. c. = G. crinita.
detonsa, Rafin. l. c. = G. serrata.
virgata, Rafin. l. c.—Am. bor.

ANTHOPOGON, Nutt. Gen. Am. i. 81 (1818) = **Gymnopogon**, Beauv. (Gram.).
albescens, Rafin. New Fl. Am. iv. 91.—Am. bor.

ANTHOPOGON :—
brevifolium, Rafin. New Fl. Am. iv. 91 = Gymnopogon brevifolius.
incarnatum, Rafin. l. c. 90.—Am. bor.
lepturoides, Nutt. Gen. Am. i. 82 = G. racemosus.

ANTHOPTERUS, Hook. Ic. Pl. t. 243 (1840). *VACCINIACEAE*, Benth. & Hook. f. ii. 568.
calycinus, Spruce, ex Benth. & Hook. f. Gen. ii. 570 = Ceratostema calycinum.
longiflorus, Benth. ex Benth. & Hook. f. l. c. = Ceratostema longiflorum, *Benth. & Hook. f.*
mucronatus, *Benth. Pl. Hartw.* 221.—Reg. Colomb.
racemosus, *Hook. Ic. Pl.* iii. t. 243.—Peruv.
Wardii, *Ball, in Hook. Ic. Pl.* t. 1465.—Reg. Colomb.

ANTHORA, Hall. in Rupp. Fl. Jen. 290 (1745) ; DC. Prod. i. 56 = **Aconitum**, Tourn. (Ranunc.).
saxatilis, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 326 = Ac. Anthora.
versicolor, Stev. ex Ser. in DC. Prod. i. 57 = Ac. Anthora.

ANTHOSACHNE, Steud. Syn. Pl. Gram. 237 (1855) = **Agropyron**, J. Gaertn. (Gramin.).
australasica, Steud. l. c. = Agrop. scabrum.

ANTOSCHMIDTIA, Steud. Syn. Pl. Gram. 199 (1855) = **Schmidtia**, Steud. (Gramin.).

ANTHOSCIADIUM, Fenzl, in Endl. Gen. Suppl. iv. III. 9 (1850) = **Selinum**, Linn. (Umbell.).

ANTHOSPERMUM, Linn. Gen. ed. I. 302 (1737). *RUBIACEAE*, Benth. & Hook. f. ii. 140.
CHRYSOSPERMUM, Reichb. Consp. 166 (1828) ?
aethiopicum, *Linn. Sp. Pl.* 1058.—Afr. austr.
ambrosiacum, Moench, Meth. Suppl. 269 = aethiopicum.
aromaticum, Salisb. Prod. 59 = aethiopicum.
asperuloides, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 11.—Fernando Po.
Bergianum, *Cruse, Rub. Cap.* 8.—Afr. austr.
Burkei, *Sond. in Harv. & Sond. Fl. Cap.* iii. 29.—Afr. austr.
calycophyllum, *Sond. l. c.* 31.—Afr. austr.
ciliare, *Linn. Sp. Pl.* ed. II. 1521.—Afr. austr.
ciliare, Thunb. ex Sond. in Harv. & Sond. Fl. Cap. iii. 30 = hirtum.
confertum, *Cruse, in Walp. Rep.* ii. 462.—Afr. austr.
Crocyllis, *Sond. in Harv. & Sond. Fl. Cap.* iii. 32.—Afr. austr.
Dregei, *Sond. l. c.* 29.—Afr. austr.
Ecklonis, *Sond. l. c.* 32.—Afr. austr.
emirnense, *Baker, in Journ. Bot.* xx. (1882) 139.—Madag.
ferrugineum, *Eckl. & Zeyh. Enum.* 366.—Afr. austr.
foetidum, Eckl. & Zeyh. ex Harv. & Sond. Fl. Cap. iii. 30 = lanceolatum.
galioides, Reichb. ex Spreng. Syst. iv. Cur. Post. 338 = ciliare.
Galopina, Thunb. Prod. Pl. Cap. 32 = Galopina circaeoides.
hedyotideum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 30.—Afr. austr.
herbaceum, Linn. f. Suppl. 440 = lanceolatum.
hirsutum, DC. Prod. vi. 580 = hirtum.
hirtum, *Cruse, Rub. Cap.* 11.—Afr. austr.
hispidulnm, *E. Mey. ex Harv. & Sond. Fl. Cap.* iii. 29.—Afr. austr.
lanceolatum, Sieber, ex Harv. & Sond. l. c. 30 = hirtum.
lanceolatum, *Thunb. Prod. Fl. Cap.* 32.—Afr. austr.
latifolium, E. Mey. ex Harv. & Sond. l. c. 30 = lanccolatum.
Lichtensteinii, *Cruse, Rub. Cap.* 9, 15.—Afr. austr.
muriculatum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 345.—Abyss.
nodosum, E. Mey. ex Harv. & Sond. Fl. Cap. iii. 30 = lanceolatum.
pachyrrhizum, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 229.—Afr. trop.
paniculatum, *Cruse, Rub. Cap.* 9, 15.—Afr. austr.
plicatum, Hilsenb. & Boj. ex Baker, in Journ. Linn. Soc. xx. (1884) 143 = Myosurandra moschata.
polyacanthum, *Baker, l. c.* 171.—Madag.

ANTHOSPERMUM :—
prostratum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 28.—Afr. austr.
pumilum, *Sond. l. c.* 31.—Afr. austr.
rigidum, *Eckl. & Zeyh. Enum.* 367.—Afr. austr.
rubiaceum, Reichb. ex Spreng. Syst. iv. Cur. Post. 338 = hirtum.
scabrum, Thunb. Prod. Pl. Cap. 32 = Carpacoce scabra.
spathulatum, Spreng. Neue Entdeck. iii. 45 = ciliare.
spermacoceum, Reichb. ex Spreng. Syst. iv. Cur. Post. 338 = Carpacoce Spermacoce.
thymoides, *Baker, in Journ. Linn. Soc.* xx. (1884) 171.—Madag.
tricostatum, *Sond. in Harv. & Sond. Fl. Cap.* iii. 28.—Afr. austr.

ANTHOSTEMA, A. Juss. Euph. Tent. 56. t. 18 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 261.
madagascariense, *Baill. Étud. Gén. Euph.* 60, 544.—Madag.
senegalensis, *Juss. Euph. Tent.* 117.—Afr. trop.

ANTHOTIUM, R. Br. Prod. 582 (1810). *GOODENOVIEAE*, Benth. & Hook. f. ii. 537.
glabrum, Poir. in Dict. Sc. Nat. ii. Suppl. 80 = humile.
humile, *R. Br. Prod.* 582.—Austral.
rubriflorum, *F. Muell. ex Benth. Fl. Austral.* iv. 45.—Austral.

ANTHOTROCHE, Endl. Nov. Stirp. Dec. 6 (1839). *SOLANACEAE*, Benth. & Hook. f. ii. 912.
Blackii, *F. Muell. Fragm.* viii. 232.—Austral.
pannosa, *Endl. Nov. Stirp. Dec.* 7.—Austral.
Walcottii, *F. Muell. Fragm.* i. 123.—Austral.

ANTHOXANTHUM, Linn. Gen. ed. I. 18 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1138.
XANTHANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, viii. 189 (1881).
XANTHONANTHOS, St. Lag. l. c. vii. (1880) 119; viii. 189 (1881).
aculeatum, Linn. f. Suppl. 89 = Crypsis aculeata.
alpinum, Schur, Enum. Pl. Transs. 725 = odoratum.
amarum, *Brot. Phyt. Lusit.* 11. t. 4. f. 1.—Lusitan.
angustifolium, Plan.-Gir. Ensaya Fl. Gallega, 398 = Puelii.
aristatum, *Boiss. Voy. Espagne*, 638.—Hispan.
aristatum, Bor. Fl. Centr. Fr. ed. II. 576 = Puelii.
asperum, Mann, ex Opiz, Natural. vii. (1824) 51 = odoratum.
avenaceum, Pourr. ex Lange, in Kjoeb. Vidensk. Meddel. (1860) 26 = amarum.
avenaceum, *Retz. Obs.* 3, 8.—Ind. or.
Carrenianum, Parl. Pl. Nov. 37 = aristatum.
crinitum, Linn. f. Suppl. 90 = Dichelachne crinita.
giganteum, Walt. Fl. Carol. 65 = Erianthus saccharoides.
glaucum, Vahl, ex Roem. & Schult. Syst. i. 288 = amarum.
gracile, *Bivon. Stirp. Rar. Sicil.* i. [13] t. 1. f. 2.—Reg. Mediterr.
indicum, Linn. Sp. Pl. 28 = Perotis latifolia.
lasianthum, *Salzm. ex Ball, in Journ. Linn. Soc.* xvi. (1878) 709.—Marocco.
Lloydii, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 697 = aristatum.
myrthense, Tod. ex Nym. Consp. 790 = Puellii.
odoratum, *Linn. Sp. Pl.* 28.—Europ.; As. bor.; Afr. bor.
ovatum, *Lag. Elench.* 2. n. 15.—Hispan.; Marocco.
paniculatum, Linn. Sp. Pl. 28 = Festuca spadicea.
Puelii, *Lecoq & Lamotte, Cat. Pl. Auver.* 385.—Europ.
pulchellum, D. Dietr. Syn. Pl. i. 625, err. typ. 69 = pulcherrimum.
pulcherrimum, *Lour. Fl. Cochinch.* 36.—Cochinch.
Sommierianum, Ricei, in Nuovo Giorn. Bot. xiii. (1881) 135 = aristatum.
spiculaeforme, Auct. ex Steud. Nom. ed. II. i. 106 = Perotis latifolia.
villosum, Dum. Obs. Gram. Belg. 129 = odoratum.

ANTHRISCUS, Bernh. Syst. Verz. Erf. 113 (1800). *UMBELLIFERAE*, Benth. & Hook. f. i. 899.

ANTHRISCUS :—
abortiva, Jord. Obs. Pl. Crit. vii. 28 = alpestris.
africana, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 195.—Afr. occ.
alpestris, *Wimm. & Grab. Fl. Silesiae*, i. 289.—Europ.
alpina, Jord. Obs. Pl. Crit. vii. 29 = sylvestris.
anatolica, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 59 = nemorosa.
capensis, Spreng. in Roem. & Schult. Syst. vi. 526 = Carum hispidum.
Caucalis, Bieb. Fl. Taur. Cauc. i. 230 = vulgaris.
Cerefolium, *Hoffm. Gen. Umb.* 41, 47. t. 1.—Europ.; Oriens; As. bor.
chaerophyllus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Cerefolium.
Cicutaria, Duby, in DC. Bot. Gall. i. 239 = alpestris.
dubia, Kabath, Fl. Gleiw. 80 = alpestris.
elatior, Bess. Enum. Pl. Volh. 55 = sylvestris.
fumarioides, *Spreng. Umb. Prod.* 27.—Europ.
gracilis, Hampe, in Flora, xxv. (1842) 1. 64 = tenerrima.
heterosantha, Schur, Enum. Pl. Transs. 273 = alpestris.
humilis, Bess. Enum. Pl. Volh. 13 = alpestris.
intermedia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858) 31 = sylvestris.
Kotschyi, *Fenzl, ex Boiss. Diagn.* Ser. II. v. 102.—Oriens.
laevigata, Griseb. Spicil. Fl. Rumel. i. 367 = sylvestris.
lamprocarpa, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 59.—Syria; Cilic.
longirostris, Bertol. Fl. Ital. iii. 197 = A. Cerefolium.
macrocarpa, Boiss. & Heldr. Diagn. Ser. II. ii. 100 = nemorosa.
mollis, *Boiss. & Reut. l. c.* 99.—Afr. bor.
neglecta, Boiss. & Reut. ex Lange, Ind. Sem. Haun. (1854) 22; Losc. & Pard. Ser. Inconf. Arag. 46 = vulgaris.
nemorosa, Baker & S. Moore, in Journ. Linn. Soc. xvii. (1879) 382 = sylvestris.
nemorosa, *Spreng. Umb. Prod.* 27.—Enrop.; Oriens.
nitens, *Bernh. in Linnaea*, xii. (1838) *Litt.* 76 = Ruthenia.
nitida, *Garcke, Fl. Deutschl.* ed. VII. 180.—Europ.
nodosa, Pers. Syn. i. 320 = Chaerophyllum nodosum.
odoratissima, *Gueldenst. Reisen durch Russl.* i. 190.—Rossia.
pilosa, Schur, Enum. Pl. Transs. 272 = sylvestris.
Prescottii, *Veesenm. in Beitr. Pfl. Russ. Reich.* ix. 84.—Rossia.
procera, Bess. Enum. Pl. Volh. 13 = sylvestris.
rivularis, Dolliner, ex Vukot. in Rad. Jugos. Akad. Zagrab, xxxix. (1877) 209 = nemorosa.
rupicola, Godet, Fl. Jura, i. 300 = alpestris.
Ruprechti, *Boiss. Fl. Orient.* ii. 914.—Reg. Caucas.
sativa, Bess. Enum. Pl. Volh. 13 = A. Cerefolium.
scandens, Steud. Nom. ed. II. i. 107, sphalm. = seq.
Scandix, Bieb. Fl. Taur. Cauc. i. 231 = Chaerophyllum nodosum.
sicula, *DC. Prod.* iv. 223.—Sicil.
sicula, Vis. ex Nym. Consp. 301 = fumarioides.
sylvestris, *Hoffm. Gen. Umb.* 40, 46. t. 1. f. 19.—Europ.; Reg. Caucas.; As. bor.
taurica, *Fisch. ex Loud. Hort. Brit.* 101.—Tauria.
tenerrima, *Boiss. & Sprun. in Ann. Sc. Nat.* Sér. III. ii. (1869) 60.—Graecia; As. Min.
torquata, Duby, in DC. Bot. Gall. i. 239 = alpestris.
trachysperma, Reichb. ex Nym. Consp. 301 = A. Cerefolium.
trichosperma, Pers. Syn. i. 320 = Caucalis trichosperma.
trichosperma, Spreng. in Roem. & Schult. Syst. vi. 525 = A. Cerefolium.
tuberculata, *Spreng. in Roem. & Schult. Syst.* vi. 527.—Hab.?
villosa, Boiss. Fl. Orient. ii. 911 = nemorosa.
vulgaris, *Bernh. Syst. Verz. Erf.* 168.—Europ.; Oriens.

ANTHROCEPHALUS, [Schlecht. in] Linnaea, xiv. (1840) Litt. 179, sphalm. = **Anthocephalus**, A. Rich. (Rubiac.).

ANTHROPODIUM, Sims, in Bot. Mag. Ind. (1816) 5, err. typ. = **Arthropodium**, R. Br. (Liliac.).

ANTHROSTYLIS, D. Dietr. Syn. Pl. i. 129, 207 (1839), err. typ. = **Arthrostylis**, R. Br. (Cyper.).

ANTHURIUM, Schott, in Wien. Zeitschr. f. Kunst, iii. 828 (1829). *AROIDEAE*, Benth. & Hook. f. iii. 998. PLEUROSPA, Rafin. Fl. Tellur. iv. 8 (1836). STREPSANTHERA, Rafin. l. c. 13 (1836).

acaule, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 83. t. 294.—Peruv.

acaule, *Schott, Melet.* i. 22.—Ind. occ.

acuminatum, Schott, in Oestr. Bot. Wochenbl. v. (1855) 66 = gracile.

aduncum, *Schott, Prod. Aroid.* 478.—Bras.

aemulum, *Schott, in Bonplandia*, vii. (1859) 165.—Mexic.

aeranthe, *Hort. ex Baker, in Saund. Ref.* t. 278.—Am. trop.

affine, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 82.—Bras.

agnatum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181.—Nicaragua.

albo-costatum, *Miq. Hort. Amstelod.* (1853).—Hab.?

alienatum, *Schott, Prod. Aroid.* 507.—Peruv.

alienigenum, Schott, ex Engl. in DC. Monog. Phan. ii. 130 = Olfersianum.

amoenum, *Kunth, Ind. Sem. Hort. Berol.* (1848) 11.—Peruv.

amplum, Kunth, Enum. Pl. iii. 77 = Hookeri.

andegavense ×, Hort. cf. Gard. Chron. (1883) i. 510.

Andersonii, *Schott, in Oestr. Bot. Wochenbl.* (1857) 325.—Ind. occ.

andicola, *Liebm. in Kjoeb. Vidensk. Meddel.* (1849) 22.—Mexic.

Andraeanum, *Linden, Illustr. Hortic.* (1877) 43. t. 271.—Reg. Colomb.

angustatum, *Kunth, Enum. Pl.* iii. 79.—Venezuela.

Appunianum, *Schott, Prod. Aroid.* 512.—Venezuela.

araliaefolium, *Regel, in Ind. Sem. Hort. Petrop.* (1869) 10; *Gartenfl.* (1870) 98. t. 648.—'N. Calcd.'

assimile, Schott, in Oestr. Bot. Wochenbl. (1855) 82 = Harrisii.

attenuatum, Miq. Cat. Hort. Amstel. (1853) = acaule.

Aubletii, Kunth, Enum. Pl. iii. 81 = pentaphyllum.

augustinum, *C. Koch & Lauche, Ind. Sem. Hort. Berol.* (1855) 7 = trinervium.

bahiense, N. E. Br. ex Engl. in DC. Monog. Phan. ii. 638 = bellum.

Bakeri, *Hook. f. Bot. Mag.* t. 6261.—Am. centr.

Belangeri, Engl. in Engl. Bot. Jahrb. i. (1881) 480 = gracile.

bellum, *Schott, in Oestr. Bot. Zeitschr.* (1859) 100.—Bras.

Beyrichianum, Kunth, Enum. Pl. iii. 69 = Harrisii.

Binoti, Linden, in Regel, Gartenfl. (1872) t. 723 = bellum.

bogotense, *Schott, in Oestr. Bot. Wochenbl.* (1857) 302.—Reg. Colomb.

bombacifolium, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 182.—Costa Rica.

Boucheanum, *C. Koch, Ind. Sem. Hort. Berol.* (1855) *App.* 7.—Venezuela.

brachygonatum, *Schott, Prod. Aroid.* 468.—Mexic.

brachyspathum, C. Koch & Bouché, in Koch, Allg. Gartenz. (1857) 233 = trinerve.

Bredemeyeri, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 269.—Bras.

breviscapum, *Kunth, Enum. Pl.* iii. 78; *Poepp. & Endl. Nov. Gen. et Sp.* iii. 84.—Peruv.

Brownii, *Mast. in Gard. Chron.* (1876) ii. 744.—Reg. Colomb.

candidum, Bull, Cat. (1875) 3. t. 11 = brevirostre.

cardiophyllum, C. Koch & August, Ind. Sem. Hort. Berol. (1854) App. 9 = nymphacifolium.

carinatum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 275.—Reg. Colomb.

cartilagineum, *Kunth, Enum. Pl.* iii. 79.—Venezuela.

caucanum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 274.—Reg. Colomb.

caudatum, Kunth, Enum. Pl. iii. 80 = undatum.

Chamissonis, Schott, in Oestr. Bot. Wochenbl. (1857) 269 = Gaudichaudianum.

chelseiensis ×, N. E. Br. in Gard. Chron. (1885) ii. 650.

clavigerum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 84.—Peruv.

cochleatum, Moritz, ex Schott, Prod. Aroid. 499 = nymphacifolium.

colocasiaefolium, *Devans. in Rev. Hortic.* (1879) 452.—Am. trop.?

commutatum, *Schott, in Bonplandia*, x. (1862) 148.—Hab.?

comtum, *Schott. l. c.* 87.—Bras.

ANTHURIUM:—

concinnatum, Schott, Prod. Aroid. 522 = bogotense.

concinnum, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 301 = Guildingii.

consanguineum, Kunth, Enum. Pl. iii. 71 = Harrisii.

consimile, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 294.—Peruv.

consobrinum, *Schott, l. c.* v. (1855) 66.—Nicaragua.

contemtum, Schott, ex Engl. in DC. Monog. Phan. ii. 129 = Harrisii.

corallinum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 84.—Peruv.

cordato-sagittatum, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 100 = rubrinervium.

cordatum, *C. Koch & H. Sello, Ind. Sem. Hort. Berol.* (1853) *App.* 5.—Venezuela.

cordatum, G. Don, in Sweet, Hort. Brit. ed. III. 633 = Pothos cordata.

cordatum, Schott, Prod. Aroid. 491 = Lhotzkyanum.

cordifolium, Hort. ex Gard. Chron. (1865) 934 = grande.

cordifolium, *Kunth, Enum. Pl.* iii. 76.—Costa Rica.

coriaceum, *G. Don, in Sweet, Hort. Brit.* ed. III. 633.—Bras.

costatum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1853) *App.* 6.—Venezuela.

crassifolium, *N. E. Br. in Gard. Chron.* (1883) i. 10.—Reg. Colomb.

crassinervium, Hort. ex Engl. in DC. Monog. Phan. ii. 138 = acaule.

crassinervium, *Schott, Melet.* i. 22.—Panama.

crenatum, Kunth, Enum. Pl. iii. 75 = acaule.

cristallinum, *Linden & André, in Linden, Cat.* (1873) n. 90. t. 128.—Peruv.

cucullatum, Koch & Sello, Ind. Sem. Hort. Berol. (1853) App. 6 = andicola.

cuprcum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 278.—Reg. Colomb.

cuspidatum, *Mast. in Gard. Chron.* (1875) i. 428.—Reg. Colomb.

cuspidifolium, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.—Costa Rica.

cymatophyllum, *C. Koch & Sello, in C. Koch, Wochenschr.* xi. (1868) 276.—Bras.

Dechardi, André, in Illustr. Hortic. xxiv. (1877) t. 269 = Spathiphyllum cannaeforme.

decurrens, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 83. t. 293.—Peruv.

denudatum, Engl. in Engl. Bot. Jahrb. vi. (1885) 280.—Reg. Colomb.

Devansayanum ×, *Hort.—Cf. Gard. Chron.* (1883) i. 510.

digitatum, *G. Don, in Sweet, Hort. Brit.* ed. III. 633.—Peruv.

dolosum, Schott, in Oestr. Bot. Zeitschr. viii. (1858) 179 = scandens.

Dombeyanum, *Brongn. ex Engl. in DC. Monog. Phan.* ii. 146.—Peruv.

dominicense, *Schott, in Oestr. Bot. Zeitschr.* (1858) 350.—Ind. occ.

egregium, Schott, Prod. Aroid. 475 = ellipticum.

elatum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1853) App. 5 = pentaphyllum.

elegans, *Engl. in Engl. Bot. Jahrb.* i. (1881) 482.—Ins. Martinica.

ellipticum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1853) App. 6.—Venezuela.

emarginatum, *Baker, in Saund. Refug.* t. 274.—Am. trop.

eminens, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 273.—Peruv.

erythropodum, Miq. Hort. Amstelod. (1853) 8 = Harrisii.

fallax, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 309 = Guildingii.

Fendleri, *Schott, Prod. Aroid.* 468.—Venezuela.

Ferrierense ×.—Cf. Gard. Chron. (1883) ii. 758.

fissum, *C. Koch, in Ender, Ind. Aroid.* 102.—Reg. Colomb.

flavescens, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 83.—Peruv.

flavidum, *N. E. Br. in Gard. Chron.* (1885) ii. 651.—Reg. Colomb.

flexile, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.—Costa Rica.

flexuosum, *Kunth, Enum. Pl.* iii. 82.—Venezuela.

floribundum, Linden & André, in Illustr. Hortic. (1874) 22. t. 159 = Spathiphyllum floribundum.

ANTHURIUM:—

Fontanesii, *Schott, in Bonplandia*, x. (1862) 347.—Venezuela.

formosum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181.—Costa Rica.

Frascri, *Engl. in DC. Monog. Phan.* ii. 112.—Ecuador.

fraternum, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 293 = Lindenianum.

Friedrichsthalii, Schott, l. c. v. (1855) 65 = Rudgeanum.

fucatum, Schott, l. c. viii. (1858) 387 = nymphacifolium.

funiferum, Klotzsch & Karst. ex Engl. in DC. Monog. Phan. ii. 111 = Bredemeyeri.

Galeottianum, *C. Koch, in Ind. Sem. Hort. Berol.* (1855) *App.* 5.—Bras.

Galeottii, *C. Koch, l. c.* (1854).—Mexic.

Gaudichaudianum, *Kunth, Enum. Pl.* iii. 74.—Bras.

Geitnerianum, *A. Regel, in Ind. Sem. Hort. Petrop.* (1866) 88.—Am. trop.

Ghiesbrechtii, Linden, ex Schott, Prod. Aroid. 538 = podophyllum.

gladiifolium, *Schott, in Journ. Bot.* i. (1863) 5.—Bras.

glaucescens, Kunth, Enum. Pl. iii. 73 = coriaceum.

glaucum, Schott, Melet. i. 22 = coriaceum.

Glaziovii, *Hook. f. Bot. Mag.* t. 6833.—Bras.

gracile, *Lindl. Bot. Reg.* t. 1635.—Guiana.

grande, *Hort. Bull, ex Gard. Chron.* (1865) 934.—Am. austr.

grandifolium, *Kunth, Enum. Pl.* iii. 77.—Reg. Colomb.

grossum, Schott, in Oestr. Bot. Zeitschr. (1859) 101 = pentaphyllum.

Guildingii, *Schott, in Oestr. Bot. Wochenbl.* (1857) 301.—Ind. occ.

Gustavi, *Regel, Gartenfl.* (1878) 324, nomen; (1882) 67.—Reg. Colomb.

gymnopus, *Griseb. Cat. Pl. Cub.* 219.—Cuba occ.

Hahnii, *Engl. in Engl. Bot. Jahrb.* i. (1881) 481.—Ins. Martinica.

Harrisii, *G. Don, in Sweet, Hort. Brit.* ed. III. 633.—Bras.

helleborifolium, Hort. ex Engl. in DC. Monog. Phan. ii. 205 = variabile.

helleborifolium, *Schott, in Bonplandia*, x. (1862) 148.—Hab.?

Hoffmanni, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181.—Costa Rica.

Hoffmannsegii, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 325.—Bras.

Holtonianum, *Schott, l. c.* 317.—Reg. Colomb.

Hookeri, *Kunth, Enum. Pl.* iii. 74.—Ind. occ.

Hookeri, Schott, Prod. Aroid. 472 = tetragonum.

Huegelii, Schott, in Oestr. Bot. Wochenbl. v. (1855) 83 = Hookeri.

Humboldtianum, Kunth, Enum. Pl. iii. 78 = rubrinervium.

Humboldtianum, Schott, Prod. Aroid. 524 = polyrrhizon.

humile, Schott, l. c. 442 = amoenum.

hybridum, *Hort. ex Engl. in DC. Monog. Phan.* ii. 191.—Hab.?

hygrophilum, *Engl. in Bot. Jahrb.* vi. (1885) 276.—Reg. Colomb.

ianthinopodum, Schott, ex Engl. in Mart. Fl. Bras. iii. ii. 89 = Harrisii.

illepidum, Schott, in Oestr. Bot. Zeitschr. (1859) 100 = longifolium.

imperiale, *Miq. ex Schott, Prod. Aroid.* 465.—Hab.?

inamoenum, Schott. l. c. 504 = nymphacifolium.

inconditum, Schott, in Oestr. Bot. Zeitschr. (1858) 181 = gracile.

inconspicuum, *N. E. Br. in Gard. Chron.* (1885) i. 787.—Bras.

indecorum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 350.—Peruv.

indecorum, Schott, ex Engl. in DC. Monog. Phan. ii. 205 = variabile.

insigne, *Mast. in Gard. Chron.* (1878) i. 430.—Reg. Colomb.

intermedium, Kunth, Enum. Pl. iii. 70 = Harrisii.

Isertianum, Schott, Prod. Aroid. 493 = Guildingii.

Jilekii, Schott, in Bonplandia, x. (1862) 5 = Harrisii.

Kalbreyeri, *Hort. ex Gard. Chron.* (1881) ii. 117.—N. Granat.?

Karwinskyi, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 101; *Prod. Aroid.* 553.—Am. cent.

ANTHURIUM:—
Kastelskii, *Schott, in Oestr. Bot. Zeitschr.* (1859) 100. —Chili.
Kellermannii, *Schubert, ex Kellerm. in Wien. Illustr. Gart. Zeit.* ix. (1884) 6.—Hab.?
Kunthianum, *Liebm. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 372.—Mexic.
Kunthii, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 84.—Peruv.
lactiflorum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 277. —Reg. Colomb.
lanceolatum, *Kunth, Enum. Pl.* iii. 71.—Am. austr.
lanceolatum, G. Don, in Sweet, Hort. Brit. ed. III. 634 = Willdenowii.
lancifolium, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180. —Costa Rica.
Langsdorffii, *Schott, Prod. Aroid.* 458.—Bras.
lapathifolium, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 309 = ochranthum.
Laucheanum, *C. Koch, in Allgem. Gartenz.* (1857) 191.—Bras.
Lechlerianum, *Schott, Prod. Aroid.* 534.—Peruv.
leptostachyum, Schott, in Oestr. Bot. Wochenbl. v. (1855) 66 = Olfersianum.
leucocarpum, Schott, l. c. (1857) 53 = scandens.
leuconeurum, *Lem. in Illustr. Hortic.* ix. (1862) t. 314. —Am. trop.
Lhotzkyanum, *Schott, Prod. Aroid.* 491.—Bras.
Libonianum, Lindl. & Regel, in Gartenfl. (1867) 291. t. 558 = Gaudichaudianum.
Liebmanni, *Schott, in Bonplandia*, vii. (1859) 165.—Mexic.
Lindenianum, Hort. ex Engl. in DC. Monog. Phan. ii. 176 = nymphaeifolium.
Lindenianum, *C. Koch & August, in C. Koch, Allg. Gartenz.* (1857) 234.—Reg. Colomb.
Lindigii, Hérincq, L'Hortic. Franç. (1866) 17; N. E. Br. in Gard. Chron. (1879) ii. 554 = Lindenianum.
linguifolium, Hort. ex Engl. in DC. Monog. Phan. ii. 124 = Geitnerianum.
longifolium, Hort. ex Engl. l. c. 129 = Harrisii.
longifolium, *G. Don, in Sweet, Hort. Brit.* ed. III. 633; *Kunth, Enum. Pl.* iii. 69.—Bras.
longispathum, *Schott, Prod. Aroid.* 520.—Costa Rica.
lucidum, *Kunth, Enum. Pl.* iii. 73.—Bras.
Luschnathianum, Kunth, l. c. = Olfersianum.
macilentum, Schott, in Bonplandia, vii. (1859) 165 = gracile.
Macleani, *Schott, in Oestr. Bot. Zeitschr.* (1858) 350. —Peruv.
macrolobium ×, *Hort. Bull, in Gard. Chron.* (1883) i. 404.—Hab.?
macrophyllum, G. Don, in Sweet, Hort. Brit. ed. III. 633 = Pothos macrophylla.
macrophyllum. Schott, in Oestr. Bot. Wochenbl. v. (1855) 83 = Selloum.
macrospadix, *Lem. Illustr. Hortic.* ix. (1862) *Misc.* 20. —Guiana.
magnificum, *Linden, in Belg. Hortic.* xv. (1865) 98.—Reg. Colomb.
magnificum, Hort. ex Gard. Chron. (1865) 934 = grande.
Malyi, *Maximil. (Archd. Austr.) ex Schott, in Bonplandia*, x. (1862) 322.—Bras.
mandiocanum, Schott, Prod. Aroid. 454 = Harrisii.
margaritaceum, Baker, in Saund. Refug. t. 280 = Hoffmanni.
Martianum, *C. Koch & Kolb, in C. Koch, Wochenschr.* xi. (1868) 276.—Guiana.
Martini, *Schott, in Oestr. Bot. Wochenbl.* vii. (1857) 325.—Guiana.
martinicense, *Engl. in Engl. Bot. Jahrb.* i. (1881) 480. —Ins. Martinica.
Maximiliani, *Schott, in Bonplandia*, x. (1862) 5.—Bras.
membranuliferum, Schott & Ohlend. ex Engl. in DC. Monog. Phan. ii. 196 = podophyllum.
metallicum, *Linden, ex Schott, Prod. Aroid.* 506.—Hab.?
mexicanum, *Engl. in DC. Monog. Phan.* ii. 105.—Mexic.
mexicanum, Liebm. in Kjoeb. Vidensk. Meddel. (1849) 21 = Schlechtendalii.
microphyllum, *G. Don, in Sweet, Hort. Brit.* ed. III. 633.—Bras.
microspadix, Schott, in Oestr. Bot. Zeitschr. (1858) 180 = porrectum.
microstachyum, *Kunth, Enum. Pl.* iii. 72.—Reg. Colomb.
Miquelianum, *C. Koch & August, Ind. Sem. Hort. Berol.* (1855) 5.—Bras.
montanum, *Hemsl. Diag. Pl. Nov.* 36.—Guatemala.

ANTHURIUM:—
myosuroides, *Endl. ex Kunth, Enum. Pl.* iii. 72.—Reg. Colomb.
neglectum, Miq. in Cat. Hort. Amstel. (1853) = Hookeri.
nitidum, *Benth. Pl. Hartw.* 255.—Reg. Colomb.
nymphaefolium, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1853) *App.* 6.—Venezuela.
obliquatum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 387.—Venezuela.
oblongatum, *Schott, ex Engl. in Mart. Fl. Bras.* iii. ii. 290 = Olfersianum.
obtusifolium, *G. Don, in Sweet, Hort. Brit.* ed. III. 633; *Kunth, Enum. Pl.* 79.—Barbadoes.
obtusilobum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181.—Costa Rica.
ochranthum, *C. Koch, Ind. Sem. Hort. Berol.* (1853) *App.* 6.—Am. trop.
Oerstedtianum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.—Costa Rica.
Olfersianum, *Kunth, Enum. Pl.* iii. 72.—Bras.
ornatum, Schott, in Oestr. Bot. Wochenbl. (1857) 294 = nymphaeifolium.
Ottonianum, *Kunth, Enum. Pl.* iii. 82.—Bras.
oxybelium, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 310 = nitidum.
oxycarpum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 83.—Peruv.
pachiraefolium, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 273.—Bras.
pallens, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.—Costa Rica.
palmatifidum, *Van Houtte, ex Schott, Prod. Aroid.* 557.—Hab.?
palmatum, *G. Don, in Sweet, Hort. Brit.* ed. III. 633; *Kunth, Enum. Pl.* iii. 80.—Ins. Dominic.
panduraeforme, *Schott, Prod. Aroid.* 536.—Costa Rica.
panduratum, *Mart. ex Schott, in Oestr. Bot. Wochenbl.* (1855) 273.—Bras.
panduratum, Schott, l. c. = panduraeforme.
parvispathum, *Hemsl. Biol. Cent. Am. Bot.* iii. 432. —Guatemala.
Patini, R. Hogg, in Gard. Chron. (1875) i. 524 = Spathiphyllum Patini.
pedatifidum, Regel & Linden, Petrop. (1865) 44; et in Gartenfl. (1866) 66. t. 501 = pedato-radiatum.
pedato-radiatum, *Schott, in Bonplandia*, vii. (1859) 337.—Mexic.
pedatum, *Endl. ex Kunth, Enum. Pl.* iii. 79.—N. Granat.
peltatum, Poepp. & Endl. Nov. Gen. et Sp. iii. 83. t. 294 = Homalonema peltata.
pentaphyllum, *G. Don, in Sweet, Hort. Brit.* ed. III. 633; *Kunth, Enum. Pl.* iii. 81.—Bras.
podophyllum, *Kunth, l. c.* 80.—Mexic.
Poiteanum, Kunth, l. c. 68 = gracile.
Poiteauanum, Schott, Prod. Aroid. 453 = gracile.
polyrrhizon, *C. Koch & August, Ind. Sem. Hort. Berol.* (1855) *App.* 7.—Venezuela.
polytomum, Schott, in Bonplandia, vii. (1859) 337 = podophyllum.
popayense, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 274. —Reg. Colomb.
porrectum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180. —Costa Rica.
protensum, *Schott, l. c.* 181.—Am. centr.
pseudopodophyllum, Schott, in Bonplandia, vii. (1859) 338 = podophyllum.
pulchellum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 273. —Reg. Colomb.
Purdieanum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 53.—Reg. Colomb.
quindiuense, Schott, l. c. 293 = fraternum.
quinquenervium, *Kunth, Enum. Pl.* iii. 76.—Reg. Colomb.
quinquevulnerum, Schott, ex Engl. in Mart. Fl. Bras. iii. ii. 100 = undatum.
radicans, *C. Koch & A. Haage, Ind. Sem. Hort. Berol.* (1854) *App.* 10.—Bras.
recusatum, *Schott, Prod. Aroid.* 467.—Cuba.
reflexum, *Hort. Par. ex Regel, Gartenfl.* (1866) 259.—Am. trop.
reflexum, G. Don, in Sweet, Hort. Brit. ed. III. 633 = coriaceum.
regale, *Linden, in Belg. Hortic.* xvi. (1866) 200.—Am. trop.
Regnellianum, *Engl. in Mart. Fl. Bras.* iii. ii. 96.—Bras.

ANTHURIUM:—
repandum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 317. —Reg. Colomb.
reptans, *G. Don, in Sweet, Hort. Brit.* ed. III. 633.—Bras.
reticulatum, *Benth. Pl. Hartw.* 255.—Peruv.
rigidulum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180 —Costa Rica.
Roezlii, *Regel, in Gartenfl.* (1872) 331.—Reg. Colomb.
Rothschildianum ×, *Hort.—Cf. Gard. Chron.* (1883) i. 510.
rubescens, *G. Don, in Sweet, Hort. Brit.* ed. III. 634.—Bras.
rubidum, Schott, ex Engl. in DC. Monog. Phan. ii. 129 = Harrisii.
rubricaule, Kunth, Enum. Pl. iii. 39 = Harrisii.
rubrinervium, *G. Don, in Sweet, Hort. Brit.* ed. III. 633. —Venezuela.
Rudgeanum, *Schott, in Oestr. Bot. Wochenbl.* v. (1855) 65.—Bras.
rugosum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 389. —Venezuela.
sagittaria, Linden, ex Schott, Prod. Aroid. 537 = panduraeforme.
sagittatum, G. Don, in Sweet, Hort. Brit. ed. III. 633 = rubrinervium.
Salviniae, *Hemsl. Diag. Pl. Nov.* 36.—Guatemala.
sanguineum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 279. —Reg. Colomb.
Saundersii, *Hook. f. Bot. Mag.* t. 6218.—Bras.
scandens, *Engl. in Mart. Fl. Bras.* iii. ii. 78.—Bras.
Scherzerianum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 53.—Guatemala.
Schlechtendalii, *Kunth, Enum. Pl.* iii. 75.—Mexic.
scolopendrinum, Kunth, l. c. 68 = gracile.
scolopendroides, Hort. ex Engl. in DC. Monog. Phan. ii. 129 = Harrisii.
Selloum, *C. Koch, Ind. Sem. Hort. Berol.* (1854).—Ins. S. Juan.
Sellowianum, *Kunth, Enum. Pl.* iii. 70.—Bras.
signatum, *C. Koch & L. Mathieu, Ind. Sem. Hort. Berol.* (1855) 8.—Reg. Colomb.
sinuatum, *Benth. ex Schott, in Oestr. Bot. Wochenbl.* (1857) 318.—Bras.
smilaciforme, C. Koch, Ind. Sem. Hort. Berol. (1855) 8 = undatum.
solitarium, Schott, Prod. Aroid. 478 = affine.
Sonderianum, Schott, in Bonplandia, vi. (1858) 372 = pentaphyllum.
sororium, *Schott, Prod. Aroid.* 522.—Peruv.
spathiphyllum, *N. E. Br. in Gard. Chron.* (1877) i. 652. —Hab.?
spectabile, *Hérincq, L'Hortic. Franç.* (1866) 17.—N. Granat.
spectabile, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181. —Costa Rica.
splendidum, *Hort. Bull, ex Illustr. Hortic.* xxx. (1883) t. 510. (*Cf. Gard. Chron.* (1883) i. 381).—Reg. Colomb.
stipitatum, *Benth. Pl. Hartw.* 255.—Reg. Colomb.
striatum, *C. Koch & L. Mathieu, Ind. Sem. Hort. Berol.* (1855) 7.—Peruv.
strictum, *N. E. Br. in DC. Monog. Phan.* ii. 638. —Paraguay.
subcordatum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 181.—Guatemala.
subhastatum, *Schott, l. c.* 350.—Reg. Colomb.
subsagittatum, *Kunth, Enum. Pl.* iii. 79.—Venezuela.
subsignatum, *Schott, in Bonplandia*, ix. (1862) 368.—Costa Rica.
subtriangulare, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 279.—Reg. Colomb.
subtrilobum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 198.—Venezuela.
surinamensis, *Miq. in Nat. Verh. Wet. Haarl.* vii. (1851) 211.—Surinam.
Swartzianum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 238.—Ind. occ.
tapinostachyum, *Schott, in Oestr. Bot. Zeitschr.* (1858) 180.—Costa Rica.
tetragonum, *Hook. ex Schott, Prod. Aroid.* 475.—Hab.?
tolimense, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 277. —Reg. Colomb.
trifidum, Oliver, in Hook. f. Bot. Mag. t. 6339 = signatum.
trilobum, Hort. Lindl. in Illustr. Hortic. xxiv. (1877) t. 283 = signatum.
trinerve, *Miq. in Linnaea*, xvii. (1843) 67.—Bras.
trinervium, *Kunth, Enum. Pl.* iii. 76.—Bras.

ANTHURIUM:—
triphyllum, *Brongn. ex Schott, Prod. Aroid.* 548.—Bolivia.
truncatulum, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 275.—Reg. Colomb.
umbrosum, *Liebm. in Kjoeb. Vidensk. Meddel.* (1849) 21.—Mexic.
undatum, *Schott, Melet.* i. 22.—Bras.
undulatum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1854) App. 107 = Harrisii.
undulifolium, C. Koch, ex Ender, Ind. Aroid. 17 = undatum.
Urvilleanum, Schott, Prod. Aroid. 459 = Harrisii.
variabile, *Kunth, Enum. Pl.* iii. 81.—Bras.
varians, Miq. Hort. Amstel. (1853) = aeaule.
Veitcheum, Hort. in Rev. Hortic. (1877) 37 = seq.
Veitchi, *Mast. in Gard. Chron.* (1876) 775.—Reg. Colomb.
Vellozianum, Schott, Prod. Aroid. 482 = Olfersianum.
venosum, *Griseb. Cat. Pl. Cub.* 219.—Cuba.
violaceum, Schott, Melet. i. 22 = seandens.
violascens, Schott, in Oestr. Bot. Wochenbl. (1854) 89 = costatum.
viride, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1855) 6 = Sellowianum.
virgosum, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 100.—Bras.
Wagenerianum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1853) App. 7.—Venezuela.
Wallisii, *Mast. in Gard. Chron.* (1875) 429.—Reg. Colomb.
Waluiewi, *Regel, in Act. Hort. Petrop.* vi. (1879) 290.—Venezuela.
Warocqueanum, *Moore, Florist & Pomol.* (1878) 101.—Reg. Colomb.
Warscewiczii, C. Koch, Ind. Sem. Hort. Berol. (1855) App. 4; et in Ender, Ind. Aroid. 17 = sinuatum.
Wendlandii, *Schott, in Oestr. Bot. Zeitschr.* (1858) 182.—Costa Rica.
Willdenowii, Kunth, Enum. Pl. iii. 71 = lanceolatum.
Williamsii, Hort. ex Engl. in DC. Monog. Phan. ii. 117 = Scherzerianum.

ANTHYLLIS, Riv. ex Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 488.
ACANTHYLLIS, Pomel, Nouv. Mat. Fl. Atl. 179 (1874).
BARBA-JOVIS, Tourn. ex Adans. Fam. ii. 321 (1763).
CORNICINA, Boiss. Voy. Bot. Espagne, 162 (1843).
DORYCNOPSIS, Boiss. l. c. 163 (1843).
FAKELOBA, Rafin. Sylva Tellur. 69 (1838).
PHYSANTHYLLIS, Boiss. Voy. Bot. Espagne, t. 162 (1843).
POGONITIS, Reichb. Handb. 226 (1837).
TRIPODION, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 348.
VULNERARIA, Tourn. ex Hall. Enum. Stirp. Helv. ii. 569 (1742).
ZENOPOGON, Link, Handb. ii. 481 (1831).
affinis, Britting. ex Koch, in Rochl. Deutschl, Fl. ed. III. v. 124 = A. Vulneraria.
alpestris, Kit. ex Schult. Oestr. Fl. ed. II. ii. 317 = A. Vulneraria.
alpina, G. Don, in Loud. Hort. Brit. 283 = A. Vulneraria.
argentea, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 408 = Cytisopsis dorycnifolia.
argentea, Salisb. Prod. 332 = A. Barba-Jovis.
arundana, Boiss. & Reut. Pugill. Pl. Nov. (1852) 35 = A. Vulneraria.
Aspalathi, Ser. in DC. Prod. ii. 169 = A. Hermanniae.
asphaltoides, Linn. Cent. Pl. ii.; Amoen. Acad. iv. 326 = Aspalathus anthylloides.
aurea, *Welden. ex Host, Fl. Austr.* ii. 319.—Graecia; Dalmat.
balearica, *Coss. ex Mares & Vigin. Cat. Pl. Balear.* 75.—Ins. Balear.
Barba-Jovis, *Linn. Sp. Pl.* 720.—Reg. Mediterr.
bicephalos, Gilib. Fl. Lituan. ii. 96 = A. Vulneraria.
bicolor, Bertol. Fl. Ital. vii. 401 = A. Vulneraria.
bidentata, *Munby, Fl. Alger.* 76.—Algeria.
biflora, [*Soland. in*] *Russ. Aleppo.* ed. II. ii. 259.—Syria.
calcicola, Schur, Enum. Pl. Transs. 150 = A. Vulneraria.
capensis, Lam. Encyc. i. 205 = Lebeckia cytisoides.
carpatica, *Pantoč. in Magyar Növént. Lapok*, vi. (1882) 162.—Europ.

ANTHYLLIS:—
chilensis, Bert. ex Steud. Nom. ed. II. i. 107 = Adesmia Loudonia.
chilensis, Ser. in DC. Prod. ii. 171 = Hosackia subpinnata.
cicerifolia, *Pourr. ex Colm. in Anal. Soc. Hist. Nat.* i. (1872) 88.—Hispan.
collina, Salisb. Prod. 332 = A. Vulneraria.
cornicina, *Linn. Sp. Pl.* 719.—Hispan.
cornicina, Pall. ex Georgi, Beschr. Russ. Reich. Nachtr. 294 = Medicago cretacea.
cornicina, Poir. Voy. Barb. ii. 211 = hamosa.
cretica, Lam. Encyc. i. 204 = Ebenus cretica.
cuneata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 246 = Lespedeza criocarpa.
cytisoides, *Linn. Sp. Pl.* 720.—Europ.
Dillenii, Schult. ex Steud. Nom. ed. I. 56 = A. Vulneraria.
Erinacea, Linn. Sp. Pl. 720 = Erinacea pungens.
flava, Gouan, Herbor. 173, nomen = A. Vulneraria.
Genistae, *Dufour, ex DC. Prod.* ii. 169.—Hispan.
genistoides, Dufour, in Bull. Soc. Bot. Fr. vii. (1860) 324 = A. Genistae.
Gerardi, *Linn. Mant.* 100.—Reg. Mediterr. occ.
glaucescens, Kit. in Linnaea, xxxii. (1863) 612 = alpestris.
gracilis, Salisb. Prod. 332 = A. cytisoides.
hamosa, *Desf. Fl. Atlant.* ii. 151.—Hispan.; Afr. bor.
Hermanniae, *Linn. Sp. Pl.* 720.—Reg. Mediterr.; Oriens.
heterophylla, *Linn. l. c.* ed. II. 1013.—Hispan.
hispida, Boiss. & Reut. Pugill. Pl. Nov. (1852) 36 = A. Vulneraria.
Hystrix, Sweet, Hort. Brit. ed. I. 114, nomen = Erinacea pungens?
incisa, Willd. Sp. Pl. iii. 1017 = Cicer ervoides.
indica, *Lour. Fl. Cochinch.* 429.—Cochineh.
involucrata, Linn. Mant. 265 = Lotononis involucrata.
italica, Hort. ex Loud. Hort. Brit. Suppl. ii. 610 = A. Vulneraria.
Jacquini, *A. Kern. in Zeitschr. Ferdinand. Tirol.* III. xv. (1870) 287.—Europ.
leguminosa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 597 = A. Vulneraria.
Lemanniana, *Lowe, in Hook. Kew Journ.* viii. (1856) 291.—Ins. Madera.
linifolia, Linn. Mant. 265 = Aspalathus linearifolia.
lotoides, *Linn. Sp. Pl.* 720.—Hispan.
macrocarpa, Walp. Rep. i. 634 = A. Vulneraria.
macrocephala, Wender. in Linnaea, v. (1830) Litt. 54 = A. Vulneraria.
maritima, Schweig. in Fl. Dan. t. 2728 = A. Vulneraria.
monocephalos, Gilib. Fl. Lituan. ii. 97 = A. Vulneraria.
montana, *Linn. Sp. Pl.* 719.—Europ.
numidica, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 396, nomen.—Algeria.
onobrychioides, *Cav. Ic.* ii. 40. t. 150.—Hispan.
onoides, Burm. f. Fl. Cap. Prod. 21 = Ononis anthylloides.
pallida, Opiz, ex Nym. Consp. 164 = A. Vulneraria.
pentaphylla, *Steud. Nom.* ed. II. i. 107.—Sibir.
podocephala, *Boiss. Elench.* 34.—Hispan.
polycephala, *Desf. Fl. Atlant.* ii. 150. t. 195.—Atlas.
polycephala, Reichb. Fl. Germ. Excurs. 516 = aurea.
polyphylla, Kit. ex DC. Prod. ii. 170 = A. Vulneraria.
pseudo-cytisus, Walp. Rep. ii. 846 = Cytisopsis dorycnifolia.
pubescens, Stokes, Bot. Mat. Med. iii. 517 = A. Vulneraria.
quinquefiora, Linn. f. Suppl. 325 = Aspalathus araneosa, parviflora.
Ramburei, *Boiss. Elench.* 35.—Hispan.
rosea, Willk. ex Rodr. in Anal. Soc. Esp. Hist. Nat. viii. (1879) 53 = balearica.
rubicunda, Wender. ex Steud. Nom. ed. II. i. 107 = A. Vulneraria.
rubra, Gouan, Herbor. 173, nomen = A. Vulneraria.
rupestris, Coss. Notes Crit. 155.—Hispan.
rustica, Mill. Gard. Dict. ed. VIII. n. 3 = A. Vulneraria.
rusticana, Wender. in Linnaea, v. (1830) Litt. 54 = A. Vulneraria.
sanguinea, Schur, Enum. Pl. Transs. 150, in syn. = A. Vulneraria.
sericea, *Lag. Gen. et Sp. Nov.* 22.—Hispan.
sericea, Willd. Sp. Pl. iii. 1014 = Ebenus Sibthorpii.
splendens, Willd. l. c. 1018 = A. Barba-Jovis.
Sprunerі, Heldr. ex Nym. Consp. 164 = A. Vulneraria.

ANTHYLLIS:—
subsimplex, *Pomel, Nouv. Mat. Fl. Atl.* 320.—Algeria.
tejedensis, *Boiss. Elench.* 35.—Hispan.; Marocc.
tetraphylla, *Linn. Sp. Pl.* 719.—Reg. Mediterr.
Tournefortii, Schult. ex Steud. Nom. ed. I. 56 = A. Vulneraria.
tragacanthoides, Desf. Fl. Atlant. ii. 150 = Astragalus armatus.
tragacanthoides, Labill. Ic. Pl. Syr. Dec. ii. 16 = Astragalus coluteoides.
tricolor, *Vukot. in Rad. Jugos. Akad. Zagrab*, xxxiv. (1876).—Europ.
variegata, Boiss. Diagn. Ser. II. v. 81 = A. Vulneraria.
visciflora, *Linn. f. Suppl.* 335.—Afr. austr.
Vulneraria, *Linn. Sp. Pl.* 719.—Europ.; Orient; Abyss.
vulnerarioides, *Bonj. ex Reichb. Fl. Germ. Excurs.* 515.—Ital.
Webbiana, Webb, ex Walp. Rep. i. 63 = tejedensis.
Webbiana, Hook. Bot. Mag. t. 3284 = A. Vulneraria.
Weldeniana, Reichb. Fl. Germ. Excurs. 515 = A. Vulneraria.

ANTIARIS, Lesch. in Ann. Mus. Par. xvi. (1810) 478. t. 22. *URTICACEAE*, Benth. & Hook. f. iii. 371.
ANTSCHAR, Horsf. in Verh. Bat. Gen. vii. (1814) 8.
IPO, Pers. Syn. ii. 566 (1807).
LEPURANDRA, J. Grah. Cat. Pl. Bomb. 193 (1839).
Bennettii, *Seem. in Bonplandia*, x. (1862) 3.—Ins. Fiji.
dubia, Span. ex Hook. Comp. Bot. Mag. i. (1835) 310 = toxicaria.
innoxia, Blume, Rumphia, 172. t. 54 = toxicaria.
macrophylla, *R. Br. in Flind. Voy.* ii. 602. t. 5.—Austral.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* i. 424.—Sumatra.
rufa, *Miq. l. c.* 423.—Sumatra.
saccidora, Dalz. in Hook. Kew Journ. iii. (1851) 232 = toxicaria.
toxicaria, *Lesch. in Ann. Mus. Par.* xvi. (1810) 478.—Ind. or.; Malaya.
zeylanica, Seem. in Bonplandia, x. (1862) 4, in nota; et Fl. Vit. 253 = toxicaria.

ANTICHARIS, Endl. Nov. Stirp. Dec. 22; Gen. 682 (1839); Iconogr. t. 93. *SCROPHULARINEAE*. Benth. & Hook. f. ii. 928.
DORATANTHERA, Benth. in Endl. Gen. 685 (1839).
MEISARRHENA, R. Br. in Salt, Abyssin. App. 63 (1818).
arabica, *Endl. Nov. Stirp. Dec.* 23.—Arabia.
arabica, Hochst. ex Benth. in DC. Prod. x. 347 = linearis.
glandulosa, *Aschers. in Monatsb. Acad. Berl.* (1866) 880.—Aegypt.; Arab.; Ind. or.
linearis, *Hochst. ex Aschers. l. c.* 882.—Afr. trop.; Arab.; Ind. or.

ANTICHLOA, Steud. Nom. ed. II. i. 108 (1840) = ACTINOCHLOA, Willd. = **Bouteloua**, Lag. (Gram.).

ANTICHORUS, Linn. Mant. i. 9 (1767) = **Corchorus**. Linn. (Tiliac.).
depressus, Linn. l. c. 64 = C. Antichorus.

ANTICLEA, Kunth, Enum. Pl. iv. 191 (1843) = **Zygadenus**, Michx. (Liliac.).
Fremonti, Torr. in Pacif. Rail. Rep. iv. 144 = Z. Fremonti.
glauca, Kunth, Enum. Pl. iv. 192 = Z. glaucus.
mexicana, Kunth, l. c. 193 = Z. mexicanus.
Nuttallii, Torr. in Pacif. Rail. Rep. iv. 144 = Z. Nuttallii.
sibirica, Kunth, Enum. Pl. iv. 191 = Z. sibiricus.
volcanica, Baker, in Journ. Linn. Soc. xvii. (1880) 482 = Z. volcanicus.

ANTICORYNE, Turcz. in Bull. Math.-Phys. Acad. Petersb. x. (1852) 332 = **Baeckea**, Linn. (Myrtac.).
diosmoides, Turcz. l. c. = B. ovalifolia.

ANTIDAPHNE, Poepp. & Endl. Nov. Gen. et Sp. iii. 70. t. 199 (1838). *LORANTHACEAE*, Benth. & Hook. f. iii. 216.
viscoidea, *Poepp. & Endl. l. c.*—Peruv.

ANTIDESMA, Burm. ex Linn. Diss. de Anandr. 6 (1745); Amoen. Acad. i. 249 (1749). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 284.
BESTRAM, Adans. Fam. ii. 354 (1763).
MINUTALIA, Fenzl, in Flora, xxvii. (1844) 312.
RHYTIS, Lour. Fl. Cochinch. 660 (1790).
STILAGO, Linn. Mant. i. 16 (1767).
acidum, Retz. Obs. v. 30 = A. Ghesaembilla.
acuminatum, *Wight, Ic.* t. 1991.—Ind. or.
Alexiteria, Gaertn. Fruct. i. 188. t. 39 = A. Menasu?
Alexiteria, *Linn. Sp. Pl.* 1027.—Ind. or.
Alexiteria, Willd. Sp. Pl. iv. 762 = nigricans.
alexiterium, Presl, Epim. Bot. 234 = leptocladum.
alnifolium, *Hook. Ic. Pl.* t. 481.—Afr. austr.
amboinense, Miq. Ann. Mus. Bot. Lugd. Bat. i. 218 = stipulare.
attenuatum, Wall. Cat. n. 7286 = velutinosum.
auritum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 203.—Java.
barbatum, Presl, Epim. Bot. 233 = rostratum.
bicolor, Hassk. Cat. Hort. Bog. Alt. 81 = Excaecaria bicolor.
bifrons, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 229 = venosum.
Blumei, Tul. l. c. 211 = tetrandrum.
Boivinianum, Baill. Adansonia, ii. (1861–62) 45 = venosum.
Boutoni, *Baker, Fl. Maurit.* 306.—Ins. Mascar.
Bunius, *Spreng. Syst.* i. 826.—Ind. or.; Malaya.
Bunius, Wall. Cat. n. 7282 A. = nigricans.
celebicum, *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 218.—Ins. Celebes.
ciliatum, Presl, Epim. Bot. 234 = A. Bunius.
comptum, *Tul. in Ann. Sc. Nat.* Sér. xv. (1851) 190.—Ind. or.
cordifolium, Presl, Epim. Bot. 235 = A. Bunius.
coriaceum, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 204 = fallax.
Cumingii, *Muell. Arg. in DC. Prod.* xv. II. 249.—Ins. Philipp.
cuspidatum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 67.—Ind. or.
Dallachyanum, *Baill. Adansonia*, vi. (1865–66) 337.—Austral.
diandrum, *Heyne, ex Roth, Nov. Pl. Spec.* 369.—Ind. or.
diandrum, *Spreng. Syst.* i. 826; *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 196.—Ind. or.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 467.—Sumatra.
digitaliforme, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 191.—Ins. Philipp.
diversifolium, *Miq. Fl. Ind. Bat. Suppl.* 468.—Sumatra.
erostre, *F. Muell. ex Benth. Fl. Austral.* vi. 87.—Austral.
erythrocarpum, *Muell. Arg. in DC. Prod.* xv. II. 258.—Java.
erythroxyloides, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 208.—Madag.
excavatum, *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 218.—Ins. Celebes.
fallax, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 68.—Malaya.
filiforme, Blume, Bijdr. 1124 = Bennettia filiformis.
flexuosum, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 225 = nigricans.
floribundum, Tul. l. c. 189 = A. Bunius.
frutescens, *Jack, in Malay. Misc.* ii. (1890) n. VII. 91.—Sumatra.
fruticosum, *Muell. Arg. in DC. Prod.* xv. II. 259.—Cochinch.
fruticulosum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 237.—Pegu.
Ghesaembilla, *Gaertn. Fruct.* i. 189. t. 39.—Ind. or.; Malaya; Austral.; Afr. trop.
glabrum, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 188 = A. Bunius.
Grossularia, Raeusch. Nom. ed. III. 287 = Embelia Ribes.
guineensis, G. Don, ex Hook. Niger Fl. 515, nomen = Uapaca guineensis.
heterophyllum, *Blume, Bijdr.* 1123.—Java.
heterophyllum, Presl, Epim. Bot. 234 = leptocladum.
japonicum, *Sieb. & Zucc. in Abh. Baier. Akad.* iv. III. (1846) 212.—China; Japon.
laciniatum, *Muell. Arg. in Flora*, xlvii. (1864) 529.—Afr. occ.
lancaefolium, *Boj. Hort. Maurit.* 289.—Ins. Mascar.
lanceolarium, Wall. Cat. n. 7284 = diandrum.
lanceolarium, Moritzi, Syst. Verz. Pfl. Zoll. 73 = lanceolatum.
lanceolatum, Dalz. & Gibs. Bomb. Fl. 237 = A. Menasu.
lanceolatum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 195.—Ind. or.
leptobotryum, *Muell. Arg. in Flora*, xlvii. (1864) 529.—Afr. trop.
leptocladum, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 199 = Moritzii.
leucopodum, *Miq. Fl. Ind. Bat. Suppl.* 465.—Sumatra.
littorale, *Blume, Bijdr.* 1123.—Java.
Lobbianum, *Muell. Arg. in DC. Prod.* xv. II. 254.—Java.
longifolium, *Boj. Hort. Maurit.* 289.—Ins. Mascar.
longifolium, Decne. ex Baill. Adansonia, v. (1865) 349, nomen = Richeria grandis.
lunatum, Miq. Fl. Ind. Bat. Suppl. 467 = Aporosa lunata.
macrophyllum, *Wall. ex Voigt, Hort. Suburb. Calc.* 295, nomen.—Ind. or.
madagascariense, *Lam. Encyc.* i. 206.—Madag.
Mannianum, Muell. Arg. in Flora, xlvii. (1864) 519 = Thecacoris Manniana.
martabanicum, *Presl, Epim. Bot.* 232.—Burma.
membranaceum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 68.—Afr. occ.
Menasu, *Miq. ex Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 215.—Ind. or.
Menasu, Kurz, For. Fl. ii. 360 = martabanicum.
minus, *Blume, Bijdr.* 1123.—Java.
minus, Wall. Cat. n. 7288 = coriaceum.
molle, Wall. Cat. n. 7287; Muell. Arg. in Linnaea, xxxiv. (1865–66) 67 = velutinosum.
montanum, *Blume, Bijdr.* 1124.—Java.
montanum, Thw. Enum. Pl. Zeyl. 289 = pyrifolium.
Moritzii, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 67.—Ind. or.; Malaya.
natalensis, Harv. Thes. Cap. ii. 45 = venosum.
nervosum, Wall. Cat. n. 7289 = Gironniera nervosa.
neurocarpum, *Miq. Fl. Ind. Bat. Suppl.* 466.—Sumatra.
nigricans, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 225.—Ind. or.
nitidum, Tul. l. c. 193 = leptocladum.
oblongatum, *Muell. Arg. in DC. Prod.* xv. II. 254.—Ind. or.
oblongifolium, *Blume, Bijdr.* 1125.—Java.
ovalifolium, *Zipp. ex Span. in Linnaea*, xv. (1841) 350.—Ins. Timor.
pacificum, *Muell. Arg. in DC. Prod.* xv. II. 254.—Ins. Pacif.
palembanicum, *Miq. Fl. Ind. Bat. Suppl.* i. 465.—Sumatra.
paniculatum, Roxb. ex Willd. Sp. Pl. iv. 764 = A. Ghesaembilla.
parvifolium, *F. Muell. Fragm.* iv. 86.—Austral.
persimile, *Kurz, in Journ. Bot.* xiii. (1875) 330.—Ins. Nicobar.
petiolare, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 207.—Madag.
platyphyllum, *H. Mann, in Proc. Am. Acad.* vii. (1867) 202.—Hawaii.
pleuricum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 213.—Ins. Philipp.
puberum, *Zipp. ex Span. in Linnaea*, xv. (1841) 350.—Ins. Timor.
pubescens, Moritzi, Verz. Pfl. Zolling. 73 = Moritzii.
pubescens, Roxb. Pl. Corom. ii. 35. t. 167 = A. Ghaesembilla.
puncticulatum, *Miq. Fl. Ind. Bat. Suppl.* 468.—Sumatra.
pyrifolium, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 68.—Zeylan.
refractum, Muell. Arg. in DC. Prod. xv. II. 257 = acuminatum.
retusum, *Zipp. ex Span. in Linnaea*, xv. (1841) 350, nomen.—Ins. Timor.
rhamnoides, *Brongn. ex Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 217.—Hab.?
Ribes, Raeusch. Nom. ed. III. 287 = Embelia Ribes.
rostratum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 218.—Ins. Philipp.
rotatum, Muell. Arg. in DC. Prod. xv. II. 256 = cuspidatum.
rotundifolium, *Boj. Hort. Maurit.* 289.—Ins. Mascar.
Roxburghii, *Wall. Cat.* n. 7283; *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 234.—Ind. or.
rufescens, Tul. in Ann. Sc. Nat. Sér. III. xv. (1851) 231 = venosum.
rugosum, *Wall. ex Voigt, Hort. Suburb. Calc.* 295, nomen.—Ind. or.
Rumphii, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 238.—Amboina.
salaccense, Zoll. & Moritzi, Syst. Verz. Zoll. 74 = tetrandrum.
salicifolium, *Miq. Fl. Ind. Bat. Suppl.* i. 467.—Malaya.
salicifolium, Presl, Epim. Bot. 233 = rostratum.
scandens, *Lour. Fl. Cochinch.* ii. 617.—China.
Schultzii, *Benth. Fl. Austral.* vi. 86.—Austral.
simile, Muell. Arg. in DC. Prod. xv. II. 259 = acuminatum.
sinuatum, *Benth. Fl. Austral.* vi. 87.—Austral.
sphaerocarpum, *Muell. Arg. in DC. Prod.* xv. II. 255.—Ins. Pacif.
spicatum, Blanco, Fl. Filip. ed. I. 794 = A. Ghesaembilla.
stenopetalum, Mucll. Arg. in Flora, xlvii. (1864) 520 = Thecacoris stenopetala.
Stilago, *Poir. Encyc. Suppl.* i. 403.—Hab.?
stipulare, *Blume, Bijdr.* 1125.—Ins. Molucc.
sylvestre, Lam. Encyc. i. 207 = A. Ghesaembilla et A. Bunius.
sylvestre, Wall. Cat. n. 7281 = diandrum.
tetrandum, *Blume, Bijdr.* 1124.—Java.
Thwaitesianum, Muell. Arg. in DC. Prod. xv. II. 263 = A. Bunius.
tomentosum, *Blume, Bijdr.* 1126.—Java.
tomentosum, Blume, ex Miq. Fl. Ind. Bat. i. II. 427 = velutinosum.
tomentosum, Voigt, Hort. Suburb. Calc. 294; Wight, Ic. t. 767–768 = Roxburghii.
triplinervium, Spreng. Neue Entdeck. ii. 116 = Alchornea triplinervia.
Tulasneanum, *Baill. Etud. Gén. Euph.* 602.—Ins. Mascar.?
velutinosum, *Blume, Bijdr.* 1125.—Burma; Malaya.
velutinosum, Miq. Fl. Ind. Bat. i. II. 428 = molle.
velutinum, *Tul. in Ann. Sc. Nat.* Sér. III. xv. (1851) 223.—Burma.
venosum, *E. Mey. ex Tul. l. c.* 232.—Afr. austr.
vestitum, Presl, Epim. Bot. 232 = A. Ghesaembilla.
Vogelianum, *Muell. Arg. in Flora*, xlvii. (1864) 529.—Afr. trop.
Wallichianum, Presl, Epim. Bot. 235 = diandrum.
zeylanicum, *Lam. Encyc.* i. 207; *Buch.-Ham. in Wall. Cat.* n. 8572.—Zeylan.
zeylanicum, Presl, Epim. Bot. 234 = pleuricum.

ANTIGONA, Vell. Fl. Flum. 186; iv. t. 145 (1827) = **Casearia**, Jacq. (Samyd.).
serrata, Vell. ll. cc. 187. t. 145 = C. lasiophylla.

ANTIGONON, Endl. Gen. 310 (1837). *POLYGONACEAE*. Benth. & Hook. f. iii. 103.
cinerascens, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 14.—Mexic.
cordatum, *Mart. & Gal. l. c.*—Mexic.
guatemalense, *Meissn. in DC. Prod.* xiv. 184.—Guatemala.
insigne, Mast. in Gard. Chron. (1877) i. 780 = guatemalense.
leptopus, *Hook. & Arn. Bot. Beech. Voy.* 308. t. 69.—Am. austr.
platypus, *Hook. & Arn. l. c.* 309.—Mexic.

ANTINORIA, Parl. Fl. Palerm. i. 92 (1845) = **Aira** Linn. (Gram.).
agrostidea, Parl. l. c. 95 = Aira agrostidea.
insularis, Parl. l. c. 94 = Aira insularis.

ANTIPHYLLA, Rafin. New Fl. Am. iv. 16 (1836) = **Melochia**, Dill. (Stercul.).
serrata, Rafin. l. c. = M. serrata.

ANTIPHYLLA, Haw. Enum. Saxif. 43 (1821) = **Saxifraga**, Linn.
arctica, Haw. l. c. 45 = S. muscoides.
biflora, Haw. l. c. 44 = S. biflora.
caerulea, Haw. l. c. 43 = S. oppositifolia.
oppositifolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 386 = S. oppositifolia.
retusa, Haw. Enum. Saxif. 44 = S. retusa.
spathulata, Haw. l. c. = S. spathulata.

ANTIPHYTUM, DC. ex Meissn. Gen. 280 (1840).
BORAGINEAE, Benth. & Hook. f. ii. 859.
cruciatum, *DC. Prod.* x. 121.—Bras.
floribundum, A. Gray, in Proc. Am. Acad. x. (1874) 55 = Krynitzkia floribunda.
fruticosum, Moç. ex DC. Prod. x. 121 = mexicanum.
heliotropioides, A. DC. l. c. 122 = Krynitzkia heliotropioides.
linifolium, DC. l. c. = Krynitzkia linifolia.
mexicanum, *DC. l. c.*—Mexic.
Parryi, S. Wats. in Proc. Am. Acad. xviii. (1882-83) 122 = Krynitzkia Parryi.
staechadifolium, *DC. Prod.* x. 121.—Bras.
tetraquetrum, *DC. l. c.* 122.—Bras.
Walpersii, *A. DC. l. c.*—Peruv.

ANTIRRHOEA, Comm. ex Juss. Gen. 204 (1789) (Antirhea). *RUBIACEAE* [Benth. & Hook. f. ii. 100].
GUETTARDELLA, Champ. in Hook. Kew Journ. iv. 197 (1852).
PITTONIOTIS, Griseb. in Bonplandia, vi. (1858) 8.
STENOSTOMUM, Gaertn. f. Fruct. iii. 69 (1805).
STURMIA, Gaertn. l. c. t. 192 (1805).
aristata, *Benth. & Hook. f. Gen.* ii. 100.—Ind. occ.
bifurcata, *Benth. & Hook. l. c.*—Ind. occ.
borbonica, J. F. Gmel. Syst. i. 244 = Quivisia Sieberi.
dichotoma, *Hemsl. Biol. Centr. Am. Bot.* ii. 42.—Mexic.
dioca, Bory, ex DC. Prod. iv. 459 = verticillata.
frangulacea, *DC. Prod.* iv. 459.—Ins. Mascar.
lucida, *Benth. & Hook. f. Gen.* ii. 100.—Ind. occ.
Lostaeana, Comm. ex DC. Prod. iv. 459 = borbonica.
philippinensis, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 312.—Ins. Philipp.
protracta, *Hemsl. Biol. Centr. Am. Bot.* ii. 42.—Mexic.
rotundata, *Benth. & Hook. f. Gen.* ii. 100.—Cuba.
strigosa, *Korth. in Neder. Kruid. Arch.* ii. II. (1851) 217.—Sumatr.
tenuifolia, *F. Muell. ex Benth. Fl. Austral.* iii. 418.—Austral.
trichantha, *Hemsl. Biol. Centr. Am. Bot.* ii. 42.—Panama.
verticillata, DC. Prod. iv. 459 = borbonica.
viscosa, *Benth. & Hook. f. Gen.* ii. 100.—Ind. occ.

ANTIRRHINUM, Tourn. ex Linn. Syst. ed. I (1735).
SCROPHULARINEAE, Benth. & Hook. f. ii. 934.
AGORRHINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 126.
ASARINA, Tourn. ex Mill. Gard. Dict. ed. VIII (1768).
GAMBELIA, Nutt. in Journ. Acad. Phil. Ser. II. i. (1847) 149.
ORONTIUM, Pers. Syn. ii. 158 (1807).
SAIROCARPUS, Nutt. ex A. DC. in DC. Prod. x. 592 (1846).
acutangulum, Tenore, Fl. Nap. ii. 47. t. 59 = Linaria Cymbalaria.
Adamii, Willd. ex Ledeb. Fl. Ross. iii. 207 = Linaria macroura.
aegyptiacum, *Forsk. Fl. Aegypt. Arab.* 112.—Arab.
aegyptiacum, Linn. Sp. Pl. ed. I. 613 = Linaria aegyptiaca.
aequitrilobum, Viv. Fl. Lib. Spec. 68; Fl. Cors. Sp. Nov. 10 = Linaria aequitriloba.
aerugineum, Gouan, Illustr. 38 = Linaria tristis.
albescens, Medic. Beobacht. (1783) 136 = Linaria aegyptiaca.
albifrons, Sibth. & Sm. Prod. i. 432 = Linaria albifrons.
album, Lam. Fl. Fr. ii. 345 = Linaria chalepensis.
alpinum, Linn. Sp. Pl. 615 = Linaria alpina.
alsinaefolium, Viv. Fl. Lib. Spec. 67; Fl. Cors. Sp. Nov. 10 = Linaria alsinaefolia.
ambiguum, *Lange, in Kjoeb. Vidensk. Meddel.* (1863) 37 (*Pugill.* 203).—Hispan.
amethysteum, Vent. in Lam. Encyc. iv. 353 = Linaria amethystea.
amethystinum, Steud. Nom. ed. II. i. 108 = Linaria amethystea.
angustifolium, Poir. Encyc. Suppl. iv. 27 = siculum.
angustifolium, Urv. in Mém. Soc. Linn. Par. i. (1822) 329 = tortuosum?
angustissimum, Loisel. Not. Suppl. 167 = Linaria italica.
aparinoides, Willd. Sp. Pl. iii. 247 = Linaria stricta.
aphyllum, Thunb. Prod. Pl. Cap. 105 = Utricularia capensis.
aquaticum, *Lour. Fl. Cochinch.* ii. 384 (Utricularia sp.?).—Cochinch.

ANTIRRHINUM:—
arenarium, Poir. Encyc. Suppl. iv. 26 = Linaria arenaria.
arvense, Linn. Sp. Pl. 614 = Linaria arvensis.
asarifolium, Salisb. Prod. 98 = seq.
Asarina, *Linn. Sp. Pl.* 612.—Italia.
asarinum, Lam. Fl. Fr. ii. 348 = A. Orontium.
assurgens, Bianca, ex Caruel, in Parl. Fl. Ital. vi. 656 = siculum.
auriculatum, Linn. Fl. Fr. ii. 339 = Linaria Elatine.
barbatum, *Thunb. Prod. Pl. Cap.* 105.—Afr. austr.
Barrelieri, *Bor. Graines Rec. Jard. Bot. Angers* (1854).—Hispan.
Bauhinii, Gaud. Fl. Helv. iii. 154 = Linaria italica.
bellidifolium, Linn. Sp. Pl. 617 = Anarrhinum bellidifolium.
bicorne, Linn. Pl. Rar. Afr. 11 (Amoen. Acad. vi. 88) = Nemesia bicornis.
bipartitum, Vent. Jard. Cels, t. 82 = Linaria bipartita.
bipunctatum, Cav. Ic. i. t. 33. f. 1 = Linaria amethystea.
bipunctatum, Linn. Sp. Pl. ed. I. 614 = Linaria glauca.
bipunctatum, Thuill. Fl. Paris, ed. II. 311 = Linaria supina.
breviflorum, Gilib. Fl. Lituan. i. 137 = A. Orontium.
Breweri, *A. Gray, in Proc. Am. Acad.* vii. (1868) 374.—Calif.
Broussonetii, Poir. Encyc. Suppl. iv. 23 = Linaria multipunctata.
caesium, Lag. ex Pers. Syn. ii. 157 = Linaria tristis?
calycinum, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 256.—Syria.
calycinum, Vent. in Lam. Encyc. iv. 365 = A. Orontium.
canadense, Linn. Sp. Pl. 618 = Linaria canadensis.
capense, *Burm. f. Fl. Cap. Prod.* 16.—Afr. austr.
capense, Thunb. Prod. Pl. Cap. 105 = Nemesia linearis.
capitatum, C. & J. Presl, Delic. Prag. 74 = sempervirens.
capitellatum, Vent. in Lam. Encyc. iv. 353 = Linaria glauca.
caprarium, Caruel, in Parl. Fl. Ital. vi. 362 = Linaria capraria.
caucasicum, Willd. ex Ledeb. Fl. Ross. iii. 205 = Linaria spuria.
chalepense, Linn. Sp. Pl. 617 = Linaria chalepensis.
Charidemi, *Lange, in Kjoeb. Vidensk. Meddel.* (1881) 99.—Lusit.
chytrospermum, *A. Gray, in Proc. Am. Acad.* xii. (1877) 81.—Arizona.
cimbalaria, Neck. Delic. Gallo-Belg. i. 267 = Linaria Cymbalaria.
cirrhosum, Linn. Mant. ii. 249 = Linaria cirrhosa.
comintanum, Blanco, Fl. Filip. ed. I. 502 = Hygrophila salicifolia.
commune, Lam. Fl. Fr. ii. 340 = Linaria vulgaris.
compositum, *Lojac. in Natural. Sicil.* ii. (1883) 297.—Sicilia.
confertiflorum, Benth. in DC. Prod. x. 592 = Mohavea viscida.
confertum, Jan, ex Benth. l. c. 278 = Linaria striata.
Cooperi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 376.—Calif.
cordatum, Buch, ex Chav. Monog. 181 = Linaria lanigera.
cornutum, *Benth. Pl. Hartw.* 328.—Calif.
cornutum, Durand, in Pacif. Rail. Rep. v. 11. t. 10 = leptaleum.
corsicum, *Vitm. Summa*, iii. 495.—Corsica.
Coulterianum, *Benth. in DC. Prod.* x. 592.—Calif.
craniolare, Stokes, Bot. Mat. Med. iii. 383 = A. Orontium.
crassifolium, Cav. Ic. ii. 11 = Linaria origanifolia.
cyathiferum, *Benth. Bot. Voy. Sulph.* 40. t. 19.—Calif.
Cymbalaria, Gueldenst. ex Ledeb. Fl. Ross. iii. 205 = Linaria Elatine.
Cymbalaria, Linn. Sp. Pl. ed. I. 612 = Linaria Cymbalaria.
Cymbalaria, Sieber, ex Benth. in DC. Prod. x. 267 = Linaria acutangula.
dalmaticum, Gueldenst. ex Ledeb. Fl. Ross. iii. 209 = Linaria genistifolia.
dalmaticum, Linn. Sp. Pl. ed. I. 616 = Linaria dalmatica.
delawariana, *Roxb. Hort. Beng.* 45, nomen.—Am. bor.?
dentatum, Poir. Encyc. Suppl. iv. 22 = Nemesia monomotapensis.

ANTIRRHINUM:—
dentatum, Vahl, Symb. Bot. ii. 66 = Linaria scariosa.
diffusum, Bernh. ex Steud. Nom. ed. II. i. 108 = latifolium.
diffusum, Brot. Phyt. t. 141 = Linaria multicaulis.
diminutum, *Pomel, Nouv. Mat. Fl. Atl.* 100.—Algeria.
dubium, Vill. Hist. Pl. Dauph. ii. 437 = Linaria supina.
Duriminium, Brot. Fl. Lusit. i. 197 = Anarrhinum Duriminium.
ecaudatum, *Lucé, Topog. Nachr. Oesel*, 218.—Ross.
Elatine, Linn. Sp. Pl. 612 = Linaria Elatine.
elatiniflorum, Wahlenb. in Isis, xxi. 989 = Linaria Elatine.
elatinioides, Tenore, Syll. 48. f. 2 = Linaria Elatine.
elatinoides, Willd. Sp. Pl. iii. 235 = Linaria elatinoides.
elegans, Pers. Syn. ii. 156 = Linaria amethystea.
elegans, Tenore, Viag. Abruz. 175 = A. Orontium.
filiforme, Poir. Encyc. Suppl. iv. 27 = Linaria rubrifolia.
filipes, *A. Gray, in Bot. Ives, Report*, 19.—Colorado.
flavum, Poir. Voy. Barb. ii. 191 = Linaria flava.
flexuosum, *Pomel, Nouv. Mat. Fl. Atl.* 100.—Algeria.
flexuosum, Willd. Sp. Pl. iii. 250 = Linaria flexuosa.
fragrans, *Pall. Reise Statth. Russ. Reich.* i. 111, 129.—Reg. Casp.
fruticans, Thunb. Prod. Pl. Cap. 105 = Nemesia fruticans.
fruticosum, Willd. Sp. Pl. iii. 237 = Linaria fruticosa.
fugax, Boiss. Diagn. Ser. II. iii. 160 = Linaria persica.
gallioides, Vent. in Lam. Encyc. iv. 351 = Linaria striata.
genistaefolium, 'Vill.' ex Benth. in DC. Prod. x. 272 = Linaria italica.
genistifolium, Lapeyr. Hist. Abr. Pl. Pyr. 354 = Linaria vulgaris.
genistifolium, Linn. Sp. Pl. 616 = Linaria genistifolia.
genistifolium, Vill. Hist. Pl. Dauph. ii. 440 = Linaria italica.
gibbosum, Wall. Pl. As. Rar. ii. 44 = A. Orontium.
glandulosum, Lejeune, Fl. Spa, 320 = Linaria vulgaris.
glandulosum, *Lindl. Bot. Reg.* t. 1893.—Calif.
glaucophyllum, Brot. Phyt. Lusit. 138 = Linaria glaucophylla.
glaucum, Lapeyr. Hist. Abr. Pl. Pyr. 352 = Linaria supina.
glaucum, Linn. Cent. I. n. 43; in Amoen. Ac. iv. 277 = Linaria glauca.
glaucum, Stocks, in Wight, Ic. iv. t. 1459 = Schweinfurthia sphaerocarpa.
glaucum, Thore, Chl. Land. 265 = Linaria thymifolia.
glutinosum, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 82.—Hispan.
gracile, Pers. Syn. ii. 156 = Linaria Pelisseriana.
gracile, Reut. Ind. Sem. Hort. Genev. (1852) 4; ex Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 582 = A. Orontium.
graecum, Bory & Chaub. Fl. Pélop. 175. t. 21 = Linaria graeca.
grandiflorum, Stokes, Bot. Mat. Med. iii. 384 = majus.
gratissimum, Roxb. ex Hook. f. Fl. Brit. Ind. iv. 266 = Limnophila Roxburghii.
Haelava, Forsk. Fl. Aegypt. Arab. 111 = Linaria Haelava.
hederaceum, Lam. Fl. Fr. ii. 338 = Linaria Cymbalaria.
hederaefolium, Salisb. Prod. 96 = Linaria Cymbalaria.
hepaticifolium, Poir. Encyc. Suppl. iv. 19 = Linaria hepaticaefolia.
heterophyllum, Schousb. Vextr. Marokko. 194; Willd. Sp. Pl. iii. 234 = Linaria heterophylla.
hexandrum, Forsk. Fl. Aegypt. Arab. 43 = Vandellia crustacea.
hirtum, Linn. Sp. Pl. ed. I. 616 = Linaria hirta.
hispanicum, *Chav. Monog.* 83.—Hispan.
Huetii, Reut. in Ann. Sc. Nat. Ser. IV. ii. (1854) 380 = sempervirens.
humile, Salisb. Prod. 98 = A. Orontium.
hybridum, *Benth. Cat. Pl. Pyren.* 60.—Mont. Pyren.
incarnatum, Vent. in Lam. Encyc. iv. 364 = Linaria bipartita.
indicum, Royle, ex Ball, in Journ. Linn. Soc. xvi. (1878) 595 = A. Orontium.
intermedium, Debeaux, in Bull. Soc. Bot. Fr. xx. (1873) 13 = latifolium.
intermedium, Hort. ex Steud. Nom. ed. II. i. 108 = striatum.

ANTIRRHINUM:—
intricatum, *Ball, in Journ. Bot.* xiii. (1875) 173.—Marocco.
italicum, Mill. Gard. Dict. ed. VIII. n. 5 = majus.
jamaicense, Fisch. ex Steud. Nom. ed. II. i. 108 = siculum.
junceum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 377.—Calif.
junceum, Linn. Cent. i.; Amoen. Acad. iv. 277; Vent. in Lam. Encyc. iv. 352 = Linaria spartea.
junceum, Pall. Reise, ii. 332 = Linaria Loeselii.
Kelloggii, *Greene, in Torrey Bot. Club*, x. (1883) 126.—Calif.
Kingii, *S. Wats. in Bot. King, Exp.* 215. t. 21.—Utah.
laevigatum, Herb. Banks, ex Benth. in DC. Prod. x. 258 = Diascia parviflora.
lagopodioides, Linn. f. Suppl. 279 = Linaria saxatilis.
lanigerum, Willd. Sp. Pl. iii. 233 = Linaria lanigera.
latifolium, Bourg. ex Willk. & Lange, Prod. ii. 584 = hispanicum.
latifolium, *Mill. Gard. Dict.* ed. VIII. n. 4; *DC. Fl. Fr.* v. 411.—Europ.
latifolium, Willd. Sp. Pl. iii. 238 = Linaria latifolia.
laxiflorum, Willd. l. c. 242 = Linaria laxiflora.
leptaleum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 373.—Calif.
Linaria, Linn. Sp. Pl. 616 = Linaria vulgaris.
linarioides, Linn. l. c. ed. II. 853 = Linaria vulgaris.
linifolium, Linn. l. c. 858 = Linaria linifolia.
linifolium, Pall. Ind. Taur. 161 = Linaria macroura.
Linkianum, Boiss. & Reut. Diagn. Ser. II. iii. 160 = latifolium.
linogriseum, Brot. Phyt. 135 = Linaria bipartita.
littorale, Bernh. in DC. Fl. Fr. v. 410; Poir. Encyc. Suppl. iv. 20 = Linaria littoralis.
longicorne, Thunb. Prod. Pl. Cap. 105 = Diascia Thunbergiana.
lusitanicum, Vent. in Lam. Encyc. iv. 361 = Linaria tristis.
lusitanicum, Link, ex Steud. Nom. ed. II. i. 109 = Linaria pedunculata.
macrocarpum, Soland. in Ait. Hort. Kew. ii. 335 = Nemesia chamaedrifolia.
macrourum, Bieb. Fl. Taur. Cauc. ii. 75 = Linaria macroura.
maculatum, Moç. & Sesse, ex DC. Prod. vii. 534 = Achimenes foliosa.
majus, *Linn. Sp. Pl.* 617.—Reg. Mediterr.
marginatum, Desf. in Act. Soc. Hist. Nat. Par. i. (1794) 36. t. 7 = Linaria tristis.
maritimum, Poir. Encyc. Suppl. iv. 23 = Linaria supina.
massiliense, Poir. ex Steud. Nom. ed. II. i. 57, 482 = Linaria maritima.
maurandioides, *A. Gray, in Proc. Am. Acad.* vi. (1868) 376.—Texas.
medium, Hort. ex Steud. Nom. ed. II. i. 109 = striatum.
meionanthum, Brot. Phyt. t. 126 = sempervirens.
meonanthum, Hoffmgg. & Link, Fl. Port. i. 261. t. 51 = sempervirens.
micranthum, Cav. Ic. i. 51 = Linaria parviflora.
microcarpum, *Pomel, Nouv. Mat. Fl. Atl.* 99.—Algeria.
minus, Linn. Sp. Pl. 617 = Linaria viscida.
molle, Blanco, Fl. Filip. ed. I. 503 = Hygrophila undulata.
molle, Boiss. Voy. Espagne, 449 = hispanicum.
molle, *Linn. Sp. Pl.* 1198.—Mont. Pyren.
molle, St. Amans, Bouq. Pyren. n. 154 = sempervirens.
monspessulanum, Georgi, Beschr. Russ. Reich. iii. iv. 1106 = Linaria Loeselii.
monspessulanum, Linn. Sp. Pl. 616 = Linaria striata.
montevidense, Mart. ex Colla, in Mem. Acc. Torino, xxxv. (1831) 178 = majus.
multibracteatum, Brot. ex Steud. Nom. ed. II. i. 109 = variabile.
multicaule, Linn. Sp. Pl. 615 = Linaria multicaulis.
multicaule, Tenore, Fl. Nap. Prod. 36 = Linaria stricta.
multipunctatum, Brot. Fl. Lusit. i. 195 = Linaria amethystea.
murale, Salisb. Prod. 98 = majus.
muralis, Pers. Syn. ii. 155 = Linaria Cymbalaria.
neglectum, Spreng. Neue Entdeck. iii. 160 = Linaria triphylla.
nicaeense, Risso, Fl. Nice, 347 = latifolium.
Nevinianum, *A. Gray, in Coult. Bot. Gazette*, ix. (1884) 54.—Calif.

ANTIRRHINUM:—
Nuttallanum, *Benth. in DC. Prod.* x. 592.—Calif.
ochroleucum, Salisb. Prod. 98 = Linaria vulgaris.
ocimifolium, Pourr. ex Spreng. Syst. ii. 793 = Linaria ocimifolia.
odoratissimum, Gueldenst. ex Ledeb. Fl. Ross. iii. 208 = Linaria Loeselii.
odorum, Bieb. Fl. Taur. Cauc. ii. 75 = Linaria Loeselii.
oppositiflorum, Poir. Encyc. Suppl. iv. 23 = Linaria villosa.
orbiculare, *Medic. Beobacht.* (1788) 242.—Hab. ?
orchidiflorum, [Desf.] ex Pers. Syn. ii. 155 = Linaria bipartita.
Orcuttianum, *A. Gray, in Coult. Bot. Gazette*, ix. (1884) 53.—Calif.
origanifolium, Gouan, Hort. Monsp. 301 = Linaria rubrifolia.
origanifolium, Linn. Sp. Pl. 615 = Linaria origanifolia.
Orontium, *Linn. Sp. Pl.* 617.—Europ.; As. Min.; Afr. bor.
ottomanum, *Janka, in Termész. Füzetek*, iv. (1880) 297.—Turcia.
Osyris, Cyrill. Pl. Neap. Nov. Fasc. ii. 21 = Linaria chalepensis.
pallidiflorum, Lam. Fl. Fr. ii. 341 = Linaria genistifolia.
pallidum, Tenore, Fl. Nap. i. 323 = Linaria pilosa.
papilionaceum, Linn. Mant. 86 = Schweinfurthia sphaerocarpa.
parviflorum, Jacq. Coll. iv. 204 = Linaria arvensis.
parviflorum, Willd. Sp. Pl. iii. 245 = Linaria parviflora.
patens, Thunb. Prod. Pl. Cap. 105 = Nemesia patens.
pauciflorum, Steud. Nom. ed. II. i. 109 = Linaria multicaulis.
pauciflorum, Poepp. ex Chav. Monog. 149 = Linaria canadensis.
pedunculatum, Linn. Sp. Pl. ed. II. 857 = Linaria pedunculata.
Pelisserianum, Linn. l. c. ed. I. 615 = Linaria Pelisseriana.
Pelisserianum, Mout.-Font. ex Chav. Monog. 155 = Linaria gracilis.
pilosum, Bory, Fl. Pélop. 39 = Linaria microcalyx.
pilosum, Jacq. Obs. Bot. ii. 28 = Linaria pilosa.
pinifolium, Poir. Voy. Barb. ii. 193 = Linaria reticulata.
pinnatum, Linn. f. Suppl. 280 = Nemesia pinnata.
polygalaefolium, Brot. Phyt. t. 138 = Linaria supina.
polygalaefolium, Poir. Encyc. Suppl. iv. 21 = Linaria italica.
polygalaefolium, Willd. ex Steud. Nom. ed. II. i. 109 = Linaria tristis.
polygonifolium, Poir. Encyc. Suppl. iv. 26 = Linaria amethystea.
porcinum, *Lour. Fl. Cochinch.* ii. 384.—Cochinch.
praecox, Brot. Phyt. t. 131 = Linaria spartea.
prostratum, Cyrill. ex Chav. Monog. 120 = Linaria reflexa.
pterospermum, A. Rich. Tent. Fl. Abyss. ii. 115 = Schweinfurthia pterosperma.
pubescens, Pers. Syn. ii. 157 = Linaria arenaria.
pubescens, Tenore, Fl. Nap. Prod. 36 = Linaria pallida.
pumilum, Poir. ex Steud. Nom. ed. II. i. 109 = Linaria flava.
purpureum, Dulac, Fl. Hautes-Pyr. 374 = A. Orontium.
purpureum, Linn. Sp. Pl. ed. I. 613 = Linaria purpurea.
pygmaeum, *Gouan, Illustr.* 37.—Europ. austr.
pyramidale, Vent. in Lam. Encyc. iv. 360 = Linaria pyramidata.
pyramidatum, Benth. in DC. Prod. x. 272 = praec.
pyrenaicum, Ram. ex DC. Fl. Fr. iii. 581 = Linaria supina.
quadrifolium, *G. Shaw, Cimel. Phys.* 84. t. 44.—Hab. ?
quinquelobatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Asarina.
quinquelobum, Stokes, Bot. Mat. Med. iii. 385 = Linaria Cymbalaria.
ramosissimum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 254.—Algeria.
reflexum, Linn. Syst. ed. X. 1112 = Linaria reflexa.
repens, Linn. Sp. Pl. 614 = Linaria striata.
reticulatum, Sm. Ic. Fict. [2] = Linaria reticulata.
Rhodium, Boiss. ex Benth. in DC. Prod. x. 291 = majus.
rhytidospermum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 27 = Linaria persica.
romanum, Seb. & Maur. ex Nym. Consp. 536 = tortuosum.
rosmarinifolium, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 6.—Malaya.

ANTIRRHINUM:—
rotundifolium, Gilib. Fl. Lituan. i. 138 = Linaria spuria.
rubrifolium, Poir. Encyc. Suppl. v. 737 = Linaria rubrifolia.
rupestre, Boiss. & Reut. Pugill. Pl. Nov. (1852) 82 = glutinosum.
rupestre, Bourg. ex Willk. & Lange, Prod. ii. 585 = hispanicum.
ruscinonense, Debeaux, in Bull. Soc. Bot. Fr. xx. (1873) 14 = Barrelieri.
rytidocarpum, G. Don, Gen. Syst. iv. 516 = Linaria persica.
rytidospermum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 27 = Linaria persica.
sagittatum, Poir. Encyc. Suppl. iv. 19 = Linaria sagittata.
saphyrinum, Brot. Fl. Lusit. i. 197 = Linaria amethystea.
saphyrinum, Sieber, ex Steud. Nom. ed. II. i. 109 = Linaria Pelisseriana.
saxatile, Bonani, Nann. Prod. 69 = Linaria arenaria.
saxatile, Linn. Cent. Pl. i. 16 = Linaria saxatilis.
scabrum, Thunb. Prod. Pl. Cap. 105 = Nemesia chamaedrifolia.
scariosum, Vent. in Lam. Encyc. iv. 349 = Linaria scariosa.
sempervirens, *Lapeyr. Fl. Pyr.* i. 7. t. 4; *Hist. Abr. Pl. Pyr.* 354.—Mont. Pyren.
serpyllifolium, Pourr. ex Chav. Monog. 93 = Linaria villosa.
siculum, *Mill. Gard. Dict.* ed. VIII. n. 6; *Ucria, in Roem. Arch.* i. 69.—Sicilia.
simplex, *Link, in Schrad. Neues Journ.* ii. (1799) 300.—Gallia.
simplex, Willd. Sp. Pl. iii. 243 = Linaria arvensis.
sparteum, Linn. Sp. Pl. ed. I. 1197 = Linaria spartea.
sparteum, Viviani, Fl. Lib. Spec. 33 = Linaria viscosa.
spartum, Linn. Cent. i. 17, postea *sparteum*.
speciosum, Donn, Hort. Cant. ed. IV. 139 = Linaria bipartita.
speciosum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 376.—Calif.
spinescens, Viviani, Fl. Lib. Spec. 32. t. 27. f. 2 = Linaria aegyptiaca.
spurium, Brot. ex Steud. Nom. ed. II. i. 109 = Linaria reticulata.
spurium, Linn. Sp. Pl. 613 = Linaria spuria.
striatum, Lam. Fl. Fr. ii. 343 = Linaria striata.
striatum, *Schrank, ex Steud. Nom.* ed. I. 57.—Europ.
strictum, *A. Gray, in Proc. Am. Acad.* vii. (1868) 375.—Calif.
strictum, Sibth. & Sm. Fl. Graec. vi. 75. t. 594 = Linaria Sibthorpiana.
strumosum, Banks, ex Benth. in DC. Prod. x. 260 = Nemesia strumosa.
subalpinum, Brot. Fl. Lusit. i. 196 = Linaria alpina.
subcordatum, *A. Gray, in Proc. Am. Acad.* xx. (1885) 306.—Calif.
subsessile, *A. Gray, in Coult. Bot. Gazette*, ix. (1884) 53.—Calif.
subulatum, Cyrill. ex Chav. Monog. 136 = Linaria Loeselii.
supinum, Brot. Fl. Lusit. i. 191 = Linaria reticulata.
supinum, Linn. Sp. Pl. 615 = Linaria supina.
supinum, Pall. Reise, ii. 522 = Linaria Loeselii.
supinum, Sibth. &. Sm. Fl. Graec. vi. 75. t. 595 = Linaria genistifolia.
tenellum, Cav. Ic. ii. 61. t. 180. f. 1 = Linaria tenella.
tenue, Viviani, Fl. Lib. Spec. 33. t. 16. f. 5–6; Spreng. Syst. ii. 495 = Linaria viscosa.
Thuillierii, Poir. Encyc. Suppl. iv. 26 = Linaria supina.
thymifolium, Vahl, Symb. Bot. ii. 67 = Linaria thymifolia.
tortuosum, *Bosc, ex Lam. Encyc.* iv. 365.—Reg. Mediterr. occ.
Tournefortii, Poir. Encyc. Suppl. iv. 24 = Linaria Tournefortii.
tricolor, Hort. ex Chav. Monog. 118 = Linaria triphylla.
tricolor, Pourr. ex Chav. l. c. 172 = Linaria amethystea.
triornithophorum, Linn. Sp. Pl. 613 = Linaria triornithophora.
triphyllum, Cav. Ic. ii. 61. t. 179 = Linaria Cavanillesii.
triphyllum, Linn. Sp. Pl. 613 = Linaria triphylla.
triste, Linn. l. c. = Linaria tristis.
uniflorum, Moç. &. Sesse, ex DC. Prod. vii. 532-3 = Gesneria uniflora.

ANTIRRHINUM :—
unilabiatum, Linn. f. Suppl. 279 = Diascia unilabiata.
vagans, *A. Gray, in Proc. Am. Acad.* vii. (1868) 375. —Calif.
variable, *Link, ex Steud. Nom.* ed. I. 57.—Lusit.
versicolor, Linn. Sp. Pl. 866 = Linaria versicolor.
vexillo-calyculatum, *Kellogg, in Proc. Calif. Acad.* i. ed. II. (1873) 7, 26.—Calif.
villosum, Lap. Fl. Pyren. 353 = Linaria origanifolia.
villosum, Linn. Sp. Pl. ed. II. 852 = Linaria villosa.
violaceum, Salisb. Prod. 97 = Linaria Pelisseriana.
Virga, *A. Gray, in Proc. Am. Acad.* vii. (1868) 373 — Calif.
virgatulum, Brot. Fl. Lusit. i. 193 = Linaria spartea.
virgatum, Poir. Voy. Barb. ii. 192 = Linaria virgata.
viscosum, Curt. Bot. Mag. t. 368 = Linaria hirta.
viscosum, Linn. Cent. ii. 21 = Linaria viscosa.

ANTISOLA, Rafin. Sylva Tellur. 97 (1838) = **Miconia**, Ruiz & Pav. (Melast.).
racemosa, Rafin. l. c. = M. racemosa.

ANTISTHIRIA, Pers. Syst. 127 (1797), err. typ. = **Anthistiria**, Linn. (Gramin.).

ANTISTROPHE, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 84. *MYRSINEAE*, Benth. & Hook. f. ii. 647.
oxyantha, *A. DC. l. c.*—Reg. Himal.
serratifolia, *Hook. f. in Benth. & Hook. f. Gen.* ii. 647. —Ind. or.

ANTITAXIS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 37; Ser. III. xx. (1867) 13. *MENISPERMACEAE*, Benth. & Hook. f. Gen. i. 33, 959, iii. 256.
ealocarpa, *Kurz, in Journ. Bot.* xiii. (1875) 324.—Ins. Nicob.
cauliflora, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xx. (1867) 14.—Java.
fascieulata, *Miers, l. c.*—Malaya.
longifolia, *Miers, l. c.*—Ins. Timor.
lucida, *Miers, l. c.*—Java.

ANTITHRIXIA, DC. Prod. vi. 277 (1837). *COMPOSITAE*, Benth. & Hook. f. ii. 329.
abyssinica, *Benth. & Hook. f. Gen.* ii. 329.—Afr. trop.
angustifolia, *Oliver & Hiern, in Oliver, Fl. Afr. Trop.* iii. 356.—Afr. trop.
flavicoma, *DC. Prod.* vi. 278.—Afr. austr.

ANTITRAGUS, Gaertn. Fruet. ii. 7. t. 80 (1791) = **Crypsis**, Ait. (Gram.).
aculeatus, Gaertn. l. c. = C. aculeata.

ANTIZOMA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 41; Ser. III. xvii. (1866) 266 = **Cissampelos**, Linn. (Menisp.).
angustifolia, Miers, ex Harv. & Sond. Fl. Cap. i. 11 = C. angustifolia.
Burchelliana, Miers, l. c. = C. calcarifera.
calcarifera, Miers, l. c. 12 = C. calcarifera.
Harveyana, Miers, l. c. = C. angustifolia.
lycioides, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xvii. (1866) 266.—Afr. austr.
Miersiana, Harv. in Harv. & Sond. Fl. Cap. i. 13.— Afr. austr.

ANTOCHORTUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 451 (1836) = **Willdenowia**, Thunb. (Restiac.).
Ecklonii, Nees, l. c. = W. striata?

ANTODON, Neck. Elem. i. 58 (1790) = **Leontodon**, Linn. (Compos.).

ANTOMMARCHIA, Colla, Hort. Ripul. App. ii. 345 (1826) = **Correa**, Sm. (Rutac.).
rubra, Colla, l. c. 25. t. 72 = C. speciosa.
speciosa, Schlecht. in Linnaea, iv. (1829) Litt. 56 = C. speciosa.

ANTONIA, R. Br. in Wall. Pl. As. Rar. iii. 65 (1832) = **Rhynchoglossum**, Blume (Gesner.).
obliqua, R. Br. l. c. = R. obliquum.

ANTONIA, Pohl, Pl. Bras. Ic. ii. 13. t. 109 (1831). *LOGANIACEAE*, Benth. & Hook. f. ii. 796.

ANTONIA :—
Griffithii, Wight, Illustr. Ind. Bot. ii. t. 156 b = Norrisia malacensis.
ovata, *Pohl, Pl. Bras.* ii. 13. t. 109.—Bras.
pilosa, Hook. Ic. Pl. t. 64 = ovata.
pubescens, Bong. in Mém. Acad. Pétersb. Sér. VI. iii. (1835) 2. t. 1 = ovata.

ANTONIANA, Tuss. Fl. Antill. 2. t. 16 (1818) ex Griseb. Fl. Brit. W. Ind. 338 = **Faramea**, Aubl. (Rubiac.).
laurifolia, Tuss. l. c. = F. odoratissima.

ANTOPETITIA, A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 261 = **Ornithopus**, Linn. (Legum.).
abyssinica, A. Rich. l. c. 262 = O. coriandrinus.
cancellata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 210 = Helminthocarpon abyssinicum.

ANTOPHYLAX, Poir. in Dict. Sc. Nat. ii. Suppl. 95 (1816), err. typ. = ANDROPHYLAX, Wendl. = **Cocculus**, DC. (Menisp.).

ANTOSCHMIDTIA, Steud. Syn. Gram. 199 in syn. Schmidtiae (1855) = **Schmidtia**, Steud. (Gram.).
quinqueseta, Boiss. Fl. Orient. v. 559 = S. quinqueseta.

ANTRIBA, Rafin. Sylva Tellur. 126 (1838) = **Loranthus**, Linn.
budleoides, Rafin. l. c. = L. Scurrula.

ANTROLEPIS, Welw. Apont. 578 (1858) = **Ascolepis**, Nees (Cyper.).
anthemiflora, Welw. l. c. = A. anthemiflora.
elata, Welw. l. c. = A. elata.
leucantha, Welw. l. c. = A. protea.
santolina, Welw. l. c. = A. protea.
sulphurea, Welw. l. c. = A. protea.

ANTROSPERMUM, Sch. Bip. in Flora, xxvii. (1844) 773 = **Venidium**, Less. (Compos.).
floribundum, Passer. Ind. Sem. Parm. (1859).—Hab.?
Kraussii, Sch. Bip. in Flora, xxvii. (1844) 773 = V. Kraussii.

ANTSCHAR, Horsf. in Verh. Bat. Gen. vii. (1814) 8 = **Antiaris**, Leseh. (Urtic.).

ANTURA, Forsk. Fl. Aegypt. Arab. 63 (1775) = **Carissa**, Linn. (Apocyn.).
hadiensis, J. F. Gmel. Syst. 405 = C. edulis.

ANUBIAS, Schott, in Oestr. Bot. Wochenbl. (1857) 398. *AROIDEAE*, Benth. & Hook. f. iii. 975.
Afzelii, *Schott, l. c.* 399.—Afr. occ.
Barteri, *Schott, Prod. Aroid.* 159.—Afr. occ.
heterophylla, *Engl. in DC. Monog. Phan.* ii. 435.— Afr. trop.

ANURUS, Presl, in Weitenw. Beitr. ii. (1837) 24 = **Lathyrus**, Linn. (Legum.).
linifolius, Presl, l. c. = L. Nissolia.
Nissolia, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 366 = L. Nissolia.

ANVILLEA, DC. Prod. v. 487 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 338.
SYCODIUM, Pomel, Nouv. Fl. Atl. 39 (1874).
Garcini, *DC. Prod.* v. 487.—Aegypt.; Persia.
radiata, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 471, *nomen*.—Algeria.

ANYCHIA, Michx. Fl. Bor. Am. i. 112 (1803). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 16.
BUINALIS, Rafin. New Fl. Am. iv. 40 (1836).
Argyrocoma, Michx. Fl. Bor. Am. i. 114 = Paronychia Argyrocoma.
Baldwinii, Torr. & Gray, Fl. N. Am. i. 172 = Paronychia Baldwinii.
canadensis, *Ell. Sketch*, i. 307.—Am. bor.
capillacea, *DC. Prod.* iii. 369.—Am. bor.
conferta, *Rafin. Atl. Journ.* (1832) 16.—Am. bor.
dichotoma, *Michx. Fl. Bor. Am.* i. 113.—Am. bor.
divaricata, *Rafin. New Fl. Am.* iv. 42.—Am. bor.
fastigiata, *Rafin. Atl. Journ.* (1832) 16.—Am. bor.
Herniariae, Pers. Syn. i. 261 = Illecebrum herniarioides.
herniarioides, Ell. Sketch, i. 308 = Herniaria americana.
herniarioides, Michx. Fl. Bor. Am. i. 113 = Paronychia herniarioides.

ANYCHIA :—
lateralis, *Rafin. Atl. Journ.* (1832) 16.—Am. bor.
nudiflora, *Rafin. New Fl. Am.* iv. 43.—Am. bor.
polygonoides, *Rafin. Atl. Journ.* (1832) 16.—Am. bor.

AOPLA, Lindl. Bot. Reg. sub t. 1701 (1835) = **Habenaria**, Linn.
reniformis, Lindl. l. c. = H. reniformis.

AOTUS, Sm. in Kon. & Sims, Ann. Bot. i. (1805) 504. *LEGUMINOSAE*, Benth. & Hook. f. i. 470.
carinata, *Meissn. in Lehm. Pl. Preiss.* ii. 215.—Austral.
cordifolia, *Benth. in Enum. Pl. Hueg.* 96.—Austral.
cordifolia, Lindl. & Paxt. Fl. Gard. i. 76 = Pultenaea ternata.
dillwynioides, Meissn. in Lehm. Pl. Preiss. i. 60 = Dillwynia floribunda.
Drummondii, *T. Moore, in Gard. Comp.* (1852) 97.— Austral.
ericoides, G. Don, Gen. Syst. ii. 120 = villosa.
ferruginea, Labill. Pl. Nov. Holl. i. t. 132 = villosa.
genistoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 268.—Austral.
gracilis, Loud. Hort. Brit. Suppl. ii. 610 = gracillima.
gracillima, *Meissn. in Lehm. Pl. Preiss.* i. 59.— Austral.
intermedia, Meissn. l. c. 60 = gracillima.
lanigera, *A. Cunn. ex Benth. in Ann. Wien. Mus.* ii. (1838) 78.—Austral.
mollis, *Benth. in Mitch. Journ. Trop. Austral.* 236.— Austral.
passerinoides, *Meissn. in Lehm. Pl. Preiss.* i. 61.— Austral.
phylicoides, *F. Muell. ex Benth. Fl. Austral.* ii. 92.— Austral.
Preissii, *Meissn. in Lehm. Pl. Preiss.* ii. 214.—Austral.
procumbens, Meissn. l. c. i. 60 = Preissii.
Tietkensii, *F. Muell. Fragm.* x. 33.—Austral.
villosa, *Sm. in Kon. & Sims, Ann. Bot.* i. (1805) 504. —Austral.
virgata, *DC. Prod.* ii. 108.—Austral.
Wuerthii, Regel, in Bot. Zeit. ix. (1851) 596 = Dillwynia floribunda.

APABUTA, Griseb. Ges. Abhandl. 280 (1880), nomen. *MENISPERMACEAE*.

APACTIS, Thunb. Nov. Gen. 66 (1783) = **Roydsia**, Roxb. (Capparid.).
japonica, Thunb. l. c.—Japon.

APALANTHE, Planch. in Ann. & Mag. Nat. Hist. Ser. II. i. (1848) 87 = **Elodea**, Michx. (Hydroch.).
granatensis, Planch. l. c. = E. granatensis.
guyanensis, Planch. l. c. = E. guyanensis.
Schweinitzii, Planch. l. c. = E. canadensis.

APALANTUS, 'Adans. Fam. ii. p. (19)' fide Pfeiff. Nom. i. 225 (1870) = **Hapalanthus**, Jacq.

APALATOA, Aubl. Pl. Gui. i. 382. t. 147 (1775) = **Crudia**, Schreb. (Legumin.).
aromatica, Lam. Encyc. iv. 559 = C. aromatica.
bantamensis, Baill. Hist. Pl. ii. 104.—Java.
guianensis, Lam. Encyc. i. 208 = C. Apalatoa.
spicata, Aubl. Pl. Gui. i. 383 = C. Apalatoa.

APALOCHLAMYS, Cass. in Dict. Sc. Nat. lvi. 223 (1828) = **Cassinia**, R. Br. (Compos.).
Billardierii, DC. Prod. vi. 157 = C. spectabilis.
Endlicheri, DC. l. c. = C. spectabilis.
Kerrii, DC. l. c. = C. spectabilis.

APALOPTERA, Nutt. ex A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 319 = **Abronia**, Juss. (Nyctag.).
annua, Nutt. ex Hook. Kew Journ. v. (1853) 261 = Ab. cycloptera.

APALUS, DC. Prod. v. 507 (1836) = **Blennosperma**, Less. (Compos.).
anthemifolius, DC. l. c. = B. chilense.

APAMA, Lam. Encyc. i. 91 (1783) = **Bragantia**, Lour. (Aristoloch.).
laurifolia, Rafin. Sylva Tellur. 13 = B. Wallichii.
siliquosa, Lam. Encyc. i. 91 = B. Wallichii.

APARGIA, Scop. Fl. Carn. ed. II. ii. 113 (1772) = **Leontodon**, Linn. (Compos.).
alpina, Willd. Sp. Pl. iii. 1547 = L. pyrenaicus.
angustifolia, Schleich. ex DC. Prod. vii. 102 = L. tenuiflorus.
annua, Visiani, Pl. Aeg. 38. t. 6 = L. hispidulus.
arenaria, Moretti, ex DC. Prod. vii. 103 = L. Berinii.
aspera, Sprun. ex Nym. Consp. 468 = L. graecus.
aspera, Waldst. & Kit. Pl. Rar. Hung. ii. 114 = L. asperus.
aurantiaca, Tenore, Syll. 392 = L. pyrenaicus.
aurea, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 282 (Hoppe, ex DC. Prod. vii. 167) = Crepis aurea.
autumnalis, Hoffm. Deutschl. Fl. ed. II. ii. 113 = L. autumnalis.
Berinii, Jan, ex DC. Prod. vii. 103 = L. Berinii.
borealis, Bongard, Veget. Ins. Sitcha, 146 = L. borealis.
Brumati, Schied. ex Steud. Nom. ed. II. i. 110 = L. hastilis.
bulbosa, Balb. in Mem. Acad. Tor. (1808) 224 = L. tuberosus.
canescens, Sieber, ex DC. Prod. vii. 103 = L. Berinii.
caucasica, Bieb. Fl. Taur. Cauc. ii. 247 = L. hastilis.
caucasica, Reichb. in Moessl. Handb. ed. II. ii. 1373 = L. asperus.
chillensis, H. B. & K. Nov. Gen. et Sp. iv. 3 = L. chillensis.
cichoracea, Tenore, Prod. 46; Fl. Nap. ii. 166 = L. cichoraceus.
ciliata, Presl, Fl. Sicul. p. xxxi = L. Preslii.
communis, Spenn. Fl. Friburg. ii. 558 = L. hispidus.
coronopifolia, Willd. Sp. Pl. iii. 1553 = Picris radicata.
crispa, Willd. l. c. 1551 = L. crispus.
crocea, Less. in Linnaea, ix. (1834) 152. t. 182.—Sibir. ural.
crocea, Pollin. Fl. Veron. ii. 565. t. 2. f. 8 = L. pyrenaicus.
danubialis, Scop. Fl. Carn. ed. II. ii. 114 = L. hastilis.
dubia, Willd. Sp. Pl. iii. 1549 = L. hastilis.
ericetorum, Richt. [Taschenb. ed. II. (1830)] = L. hastilis.
fasciculata, Bivona, Stirp. Rar. Sicil. ii. [11] t. 2 = L. cichoraceus.
Gouani, Schleich. Cat. Pl. Helv. ed. IV. 7 = L. hirtus.
guestphalica, Boenning. ex Reichb. in DC. Prod. vii. 102 = L. hastilis.
Haenkei, Sternb. ex Steud. Nom. ed. II. i. 110 = L. pyrenaicus.
hastilis, Host, Syn. 423 = L. hastilis.
hastilis, Viviani, Fl. Lib. Spec. 50. t. 26. f. 1 = L. Salzmannia.
heterophylla, Moench, Meth. 543 = L. hastilis.
hieracioides, Willd. Sp. Pl. iii. 1554 = Picris hieracioides.
hirta, Geners. ex Steud. Nom. ed. II. i. 110; ii. 23 = L. pyrenaicus.
hirta, Steud. l. c. i. 110 = L. hispidus.
hirta, Scop. Fl. Carn. ed. II. ii. 114 = L. hirtus.
hispanica, Bieb. Fl. Taur. Cauc. ii. 248 = L. asperus.
hispanica, Willd. Sp. Pl. iii. 1552 = L. hispanicus.
hispida, Bory, in Ann. Sc. Gén. Phys. iii. (1820) 5 = L. Boryi.
hispida, Hoffm. Deutschl. Fl. ed. II. ii. 113 = L. hispidus.
hispida, Willd. Sp. Pl. iii. 1553 = L. hastilis.
hyoseridiflora, Saut. ex Schur, Enum. Pl. Transs. 359 = L. caucasicus.
hyoseridifolia, Less. Syn. Comp. 132 = Crepis hyoseridifolia.
hyoseroides, Sieber, in Spreng. Syst. iv. 664 = L. creticus.
hyoseroides, Vest, in Flora, iii. (1820) 7 = L. hirtus?
incana, Mart. Fl. Mosq. 135 = L. hastilis.
incana, Scop. Fl. Carn. ed. II. ii. 114 = L. incanus.
incrassata, Moench, Meth. Suppl. 220 = L. hispidus.
intermedia, Schult. ex Steud. Nom. ed. II. i. 110 = L. hastilis.
laevis, Moench, Meth. 541 = L. hirtus.
livida, Moench, l. c. 543 = L. hastilis.
lucida, Tenore, Cat. Hort. Neap. App. (1819) 71 = L. lucidus.
lyrata, Tenore, Fl. Nap. Prod. 46 = L. tuberosus.
macrorrhiza, Guenth. ex Spreng. Syst. iii. 665 = L. macrorrhizus.
media, Host, Fl. Austr. ii. 404 = L. pyrenaicus.
obtusa, Kit. in Linnaea, xxxii. (1863) 405.—Europ. austr.
pratensis, Link, Handb. i. 791 = L. autumnalis.
pyrenaica, Sternb. & Hoppe, ex Steud. Nom. ed. I. 58 = L. alpinus.

APARGIA :—
rigens, Mart. ex Reichb. in Moessl. Handb. ed. II. ii. 1377 = Picris hieracioides.
Rosani, Tenore, Fl. Nap. App. v. 26 = Leontodon Villarsii.
salina, Gaert. Meyer & Scherb. Fl. Wett. iii. 146 = L. crispus.
salsa, Schur, Enum. Pl. Transs. 884, Index = Taraxacum palustre.
saxatilis, Tenore, Cat. Hort. Neap. (1819) 59 = L. crispus.
scabra, Nocca, ex Steud. Nom. ed. II. i. 110 = Picris asplenioides.
serotina, Jan, ex Steud. l. c. = Hypochoeris radicata.
setosa, Moench, Meth. Suppl. 219 = Crepis setosa.
sicula, Guss. Fl. Sic. Syn. ii. 399 = L. hastilis.
sinuata, Kit. in Linnaea, xxxii. (1863) 404.—Europ. austr.
strigosa, Bieb. Fl. Taur. Cauc. ii. 249 = L. asperrimus.
sudetica, Link, ex Spreng. Syst. iii. 664 = L. hispidus.
Taraxaci, Hornem. in Fl. Dan. t. 1523 = L. autumnalis.
Taraxaci, Willd. Sp. Pl. iii. 1550 = L. Taraxaci.
taraxaciflora, Viv. Fl. Lib. Spec. 50. t. 22. f. 1 = Picris taraxaciflora.
tenuiflora, Gaud. Fl. Helv. vi. 362 = L. alpinus.
tergestina, Hoppe, ex DC. Prod. vii. 103 = L. crispus.
tuberosa, Willd. Sp. Pl. iii. 1549 = L. tuberosus.
umbellata, Schrank, in Denkschr. Regensb. Bot. Ges. ii. (1818) 66 = Picris hieracioides.
variegata, Willd. Sp. Pl. iii. 1552 = Hypochoeris variegata.
verna, Salzm. in Delile, Sem. Hort. Monsp. (1836) = L. Salzmanni.
Villarsii, Willd. Sp. Pl. iii. 1553 = L. Villarsii.

APARGIDIUM, Torr. & Gray, Fl. N. Am. ii. 474 (1841) = **Leontodon**, Linn. (Compos.).
boreale, Torr. & Gray, l. c. = L. borealis.

APARINANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 397 = **Galium**, Tourn. (Rubiac.).
muralis, Fourr. l. c. = G. murale.
verticillatus, Fourr. l. c. = G. verticillatum.

APARINE, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Galium**, Tourn. (Rubiac.).
borealis, Hill, Herb. Brit. ii. 151 = G. boreale.
hispida, Moench, Meth. 640 = G. Aparine.
latifolia, Moench, l. c. = G. Cruciata.
minima, All. ex DC. Prod. iv. 610 = G. murale.
pariscensis, Hill, Herb. Brit. ii. 150 = G. parisiense.
spuria, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 397 = G. spurium.
tenera, Fourr. l. c. = G. spurium.
tricornis, Fourr. l. c. = G. tricorne.
verrucosa, Moench, Meth. 640 = G. saccharatum.
viscidiflora, Rafin. Ann. Nat. (1820) 14.—Am. bor.
Vaillantii, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 397 = G. spurium.
vulgaris, Hill, Herb. Brit. ii. 149 = G. Aparine.

APARINELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 397 = **Galium**, Tourn. (Rubiac.).
decipiens, Fourr. l. c. = G. parisiense.
divaricata, Fourr. l. c. = G. divaricatum.
litigiosa, Fourr. l. c. = G. parisiense.
parisiensis, Fourr. l. c. = G. parisiense.
ruricola, Fourr. l. c. = G. parisiense.
setacea, Fourr. l. c. = G. setaceum.
tenuicaulis, Fourr. l. c. = G. divaricatum.

APARISTHMIUM, Endl. Gen. 1112 (1840) = **Alchornea**, Sw. (Euphorb.).
cordatum, Baill. Adansonia, v. (1863-64) 307 = Al. cordata.
javense, Hassk. Hort. Bogor. 235 = Al. rugosa.
Spruceanum, Baill. Adansonia, v. (1863-64) 307 = Al. cordata.
sumatranum, Reichb. f. & Zoll. in Verh. Nat. Ver. Nederl. Ind. i. (1856) 28 = Al. villosa.

APATANTHUS, Viv. Fl. Lib. Spec. 54. t. 7. f. 3 (1824) = **Hieracium**, Linn. (Compos.).
crinitus, Viv. l. c. = H. crinitum.

APATELIA, DC. Mém. Ternstr. 34. t. 8 (1822) = **Saurauja**, Willd. (Ternstroem.).

APATELIA :—
biserrata, DC. Prod. i. 526 = S. biserrata.
glabrata, DC. Prod. i. 526 = Saurauja serrata.
lanceolata, DC. l. c. = S. lanceolata.
scabra, G. Don, Gen. Syst. i. 569 = S. excelsa.
tomentosa, G. Don, l. c. 568 = S. tomentosa.

APATEMONE, Schott, Gen. Aroid. t. 57 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 984.
Motleyana, *Schott, l. c.*—Borneo.

APATITIA, Desv. in Ham. Prod. Pl. Ind. Occ. 42 (1825) = **Bellucia**, Neck. (Meliac.).
blakeoides, Desv. l. c. = B. Aubletii.

APATURIA, Lindl. Gen. et Sp. Orch. 130 (1831) = **Pachystoma**, Blume (Orchid.).
chinensis, Lindl. l. c. 131 = P. chinense.
Lindleyana, Wight, Ic. Pl. Ind. v. 1662 = P. senile.
montana, Lindl. in Wall. Cat. n. 3738 = P. senile.
senilis, Lindl. l. c. n. 3739 = P. senile.
Smithiana, Lindl. l. c. n. 3737 = P. senile.

APEGIA, Neck. Elem. i. 251 (1790) = **Ceropegia**, Linn. (Asclep.).

APEIBA, Aubl. Pl. Gui. i. 537. t. 213 ad 216 (1775). *TILIACEAE*, Benth. & Hook. f. i. 237.
ANBLETIA, Schreb. Gen. 353 (1789).
OXYTANDRUM, Neck. Elem. ii. 255 (1790).
aspera, *Aubl. l. c.* i. 545. t. 216.—Guiana.
australis, A. Rich. Fl. Nov. Zel. t. 34 = Entelea arborescens.
Cimbalaria, *Arruda, in Koster, Trav. Braz.* 488.—Bras.
discolor, *G. Don, Gen. Syst.* i. 555.—Bras.
echinata, *Gaertn. Fruct.* ii. 189. t. 121.—Hab.?
emarginata, Lam. Encyc. i. 210 = Sloanea emarginata.
glabra, *Aubl. Pl. Gui.* i. 541. t. 214.—Guiana.
hirsuta, Lam. Encyc. i. 208.—A. Tibourbou.
hispida, Gaertn. Fruct. ii. t. 121 = A. Petoumo.
hypoleuca, *Steud. in Flora*, xxvi. (1843) II. 755.—Guiana.
laevis, *G. Don, Gen. Syst.* i. 555.—Guiana.
laevis, Sw. Prod. Veg. Ind. Occ. 83 = glabra.
levis, Aubl. Pl. Gui. t. 214 = glabra.
membranacea, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) *Suppl.* ii. 61.—Venezuel.
Petoumo, *Aubl. Pl. Gui.* i. 543. t. 215.—Guiana.
Tibourbou, *Aubl. l. c.* 538. t. 213.—Guiana; Venezuel.
ulmifolia, *H. B. & K. Nov. Gen. et Sp.* v. 347.—Venezuel.

APEMON, Rafin. Fl. Tellur. ii. 11 (1836) = **Datura**, Linn. (Solan.).
crassicaule, Rafin. l. c. = D. ceratocaula.

APENTOSTERA, Rafin. New Fl. Am. ii. 72 (1836) = **Penstemon**, Mitch. et **Chelone**, Linn. (Scrophul.).
secundiflora, Rafin. l. c. = P. Menziesii.
triflora, Rafin. l. c. 73 = Chelone nemorosa.

APELLA, Scop. Introd. 107 (1777) = APPELLA, Adans. = **Laurus**, Linn.

APENULA, Neck. Elem. i. 234 (1790) = **Specularia**, Heist. (Campan.).

APERA, Adans. Fam. ii. 495 (1763). *GRAMINEAE*, Benth. & Hook. f. iii. 1152.
ANEMAGROSTIS, Trin. Fund. Agrost. 128 (1820).
arundinacea, Hook. f. Fl. N. Zeal. 295. t. 67 (= Stipae sp.).—N. Zel.
aspera, Beauv. Agrost. 151 = Sporobolus asper.
crinita, Beauv. l. c. 31 = Dichelachne crinita.
effusa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 148 = A. Spica-venti.
interrupta, *Beauv.* Agrost. 31.—Rossia austr.; Oriens.
palustris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 148 = Agrostis alba.
procumbens, Auct. ex Steud. Nom. ed. II. i. 111 = Agrostis alba, vulgaris.
purpurea, Beauv. Agrost. 31; et Reichb. Ic. Fl. Germ. i. 32 = A. Spica-venti.
Spica-venti, *Beauv. l. c.*—Europ.; Oriens.
tenuiflora, Beauv. l. c. 151 = Muehlenbergia Willdenowii.

APERIPHRACTA, Nees, ex Meissn. in DC. Prod. xv. 1. 111 (1864) = **Ocotea**, Aubl. (Laurin.).

APERIPHRACTA:—
cuneata, Nees, ex Meissn. in DC. Prod. xv. 1. 119 (=*Oreodaphne cuneata*).—Peruv.
Martiana, Nees, l. c. 117=Ocotea rufo-tomentosa.
Minarum, Nees, l. c. 140=O. umbrosa.
velutina, Nees, l. c. 132=O. velutina.

APERULA, Blume, Mus. Bot. Lugd. Bat. i. 365 (1851) =**Lindera**, Thunb. (Laurin.).
assamica, Meissn. in DC. Prod. xvi. 240=L. assamica.
citriodora, Blume, Mus. Bot. Lugd. Bat. i. 366.—Japon.
confusa, Meissn. in DC. Prod. xv. 1. 243.—Java.
cuspidata, Blume, Mus. Bot. Lugd. Bat. i. 367.—Ins. Celebes.
densiflora, Meissn. in DC. Prod. xv. 1. 242.—Java.
insignis, Blume, in Mus. Bot. Lugd. Bat. i. 367.—Sumatr.
lanceolata, Blume, l. c.—Java.
lucida, Blume, l. c. 366.—Java.
macrophylla, Blume, l. c. 367.—Java.
Neesiana, Blume, l. c. 366=L. Neesiana.
oxyphylla, Blume, l. c.=L. oxyphylla.
piporicarpa, Meissn. in DC. Prod. xv. 1. 242.—Sumatr.
polyantha, Blume, Mus. Bot. Lugd. Bat. i. 366.—Java.
puberula, Blume, l. c. 367.—Java.
reticulata, Blume, l. c.=L. reticulata.
salicifolia, Blume, l. c.—Java.

APERULA, Gled. in Mém. Acad. Berl. (1751) 127, sphalm.=**Asperula**, Linn. (Rubiac.).

APETAHIA, Baill. in Bull. Soc. Linn. Par. i. (1882) 311. *LOBELIACEAE.*
raiateensis, *Baill. l. c.*—Ins. Pacif.

APETALON, Wight, Ic. v. 22. t. 1758 (1852)=**Didymoplexis**, Griff. (Orchid.).
minutum, Wight, l. c.=D. pallens.

APHACA, [Tourn.] Linn. Gen. ed. I. 217 (1737)= **Lathyrus**, Linn. (Legum.).
disperma, Alef. in Oestr. Bot. Zeitschr. xi. (1861) 282.—Aegypt.
marmorata, Alef. l. c.=L. polyanthus.
pseudophaca, Alef. in Bonplandia, ix. (1861) 139=L. pseudoaphaca.
vulgaris, C. Presl, in Weitenw. Beitr. ii. (1837) 24= L. Aphaca.

APHAENANDRA, Miq. Fl. Ind. Bat. ii. 341 (1856). *RUBIACEAE.*
sumatrana, *Miq. l. c.*—Sumatr.

APHAEREMA, Miers, in Proc. Hort. Soc. iii. (1863) 294. *SAMYDACEAE*, Benth. & Hook. f. i. 799.
spicata, *Miers, l. c.*—Bras.

APHANACTIS, Wedd. Chlor. And. i. 142. t. 37 A (1855). *COMPOSITAE*, Benth. & Hook. f. ii. 360.
Jamesoniana, *Wedd. l. c.*—Ecuador.

APHANAMIXIS, Blume, Bijdr. 165 (1825)=**Amoora**, Roxb. (Meliac.).
Blumei, Span. Ic. 14; et in Linnaea, xv. (1841) 182= Am. timorensis.
grandiflora, Blume, Bijdr. 165=Am. grandifolia.
Perrottetiana, A. Juss. Mém. Mél. 108=Am. Perrottetiana.
rubiginosa, Griff. ex C. DC. in DC. Monog. Phan. i. 585=Am. rubiginosa.
timorensis, A. Juss. Mém. Mél. 107=Am. timorensis.

APHANANTHE, Link, Enum. Hort. Berol. i. 383 (1821)=**Microtea**, Sw. (Phytolacc.).
celosioides, Link, l. c.=M. maypurensis.

APHANANTHE, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 265. *URTICACEAE*, Benth. & Hook. f. iii. 355.
Homoioceltis, Blume, Mus. Bot. Lugd. Bat. ii. 64. t. 34 (1852) [1856].
aspera, *Planch. in DC. Prod.* xvii. 208.—Japon.
cuspidata, Planch. l. c. 209=Girounniera reticulata.
philippinensis, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 337.—Austral.
rectinervis, *Planch. in DC. Prod.* xvii. 208.—Austral.

APHANANTHEMUM, Steud. Nom. ed. II. i. 111 (1840)=**Helianthemum**, Tourn. (Cistin.).
salicifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 340=H. salicifolium.

APHANES, Linn. Gen. ed. I. 33 (1737)=**Alchemilla**, Linn. (Rosac.).
alata, Steud. Nom. ed. I. 58=Al. pinnata.
arvensis, Linn. Sp. Pl. ed. I. 123=Al. arvensis.
cornucopioides, Lag. Gen. et Sp. Nov. 99. n. 7=Al. cornucopioides.
monogyna, [Pieri] in Ionios Anthol. iii. (1834) 694= Al. arvensis.
orbiculata, Pers. Syn. i. 150.=Al. orbiculata.
pinnata, Pers. l. c.=Al. pinnata.
triloba, Gilib. Exercit. ii. 430=Al. arvensis.
tripartita, Pers. Syn. i. 150=Al. tripartita.
vulgaris, Schur, Enum. Pl. Transs. 884=Al. arvensis.

APHANIA, Blume, Bijdr. 236 (1825) = **Sapindus**, Linn.
bifoliata, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 238=S. bifoliatus.
cuspidata, Radlk. l. c.=S. cuspidatus.
Danura, Radlk. l. c.=S. Danura.
longipes, Radlk. Sapind. Holland-Ind. Nachtr. 68.—Malaya.
microcarpa, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 238=S. microcarpus.
montana, Blume, Bijdr. 236=S. montanus.
paucijuga, Radlk. Sapind. Holland-Ind. Nachtr. 68.—Malaya.
rubra, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 238=S. attenuatus.
senegalensis, Radlk. l. c.=S. abyssinicus.
sphaerococca, Radlk. in Act. Congr. Bot. Amst. 1877 (1879) 91.—N. Guinea.

APHANISMA, Nutt. ex Moq. in DC. Prod. xiii. ii. 54 (1849). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 50.
Cryptanthus, Nutt. ex Moq. l. c.
blitoides, *Nutt. ex Moq. l. c.*—Calif.

APHANOCALYX, Oliver, in Hook. Ic. Pl. xi. 53. t. 1066 (1870). *LEGUMINOSAE.*
cynometroides, *Oliver, l. c.*—Afr. occ.

APHANOCHILUS, Benth. in Bot. Reg. sub t. 1282 (1829)=**Elsholtzia**, Willd. (Labiat.).
blandus, Benth. l. c.=E. blanda.
eriostachyus, Benth. l. c.=E. eriostachya.
flavus, Benth. l. c.=E. flava.
foetens, Benth. l. c.=E. incisa.
incisus, Benth. l. c.=E. incisa.
paniculatus, Benth. l. c.=E. incisa.
pilosus, Benth. l. c.=E. pilosa.
polystachyus, Benth. l. c.=E. polystachya.

APHANODON, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 110=**Henriettella**, Naud. (Melast.).

APHANOMYRTUS, Miq. Fl. Ind. Bat. i. 1. 480 (1855). *LAURIN.* gen. dub. (=**Baeckia**?).
rostrata, *Miq. l. c.* 481.—Malaya.

APHANOMYXIS, DC. Prod. vii. ii. 766, in nota (1839) =APHANOMIXIS, Blume = **Amoora**, Roxb. (Meliac.).

APHANOPAPPUS, Endl. Gen. Suppl. ii. 43 (1842)= **Lipochaeta**, DC. (Compos.).
Nuttallii, Walp. Rep. ii. 620=L. micrantha.

APHANOPETALUM, Endl. Gen. 818 (1839). *SAXIFRAGEAE*, Benth. & Hook. f. i. 650.
Platypetelea, J. Drum. in Hook. Kew Journ. vii. (1855) 55.
occidentale, *F. Muell. Fragm.* i. 228.—Austral.
resinosum, *Endl. Nov. Stirp. Dec.* 35.—Austral.

APHANOPLEURA, Boiss. Fl. Orient. ii. 855 (1872). *UMBELLIFERAE.*
trachysperma, *Boiss. l. c.*—Armenia.

APHANOSTEMMA, St. Hil. Fl. Bras. Mer. i. 12 (1825) =**Ranunculus**, Linn.
apiifolia, St. Hil. l. c. 13=R. apiifolius.

APHANOSTEPHUS, DC. Prod. v. 310 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 262.
Keerlia, DC. Prod. v. 309 (1836), ex parte.
Leucopsidium, DC. Prod. vi. 43 (1837).
arizonicus, A. Gray, in Proc. Am. Acad. xvi. (1881) 81=humilis.
arkansanus, *A. Gray, Pl. Wright.* i. 93.—Texas.
humilis, *A. Gray, l. c.*—Texas.
pilosus, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 457=ramosissimus.
ramosissimus, *DC. Prod.* v. 310.—Texas.
ramosus, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 81. —Mexic.
Riddelii, Torr. & Gray, Fl. N. Am. ii. 189=ramosissimus.

APHANTOCHAETA, A. Gray, in Pacif. Rail. Rep. iv. 99 (1856). *COMPOSITAE*, Benth. & Hook. f. ii. 251.
exilis, *A. Gray, l. c.* 100.—Calif.

APHARICA, Schlecht. in Linnaea, vi. (1831) Litt. 78= APHANIA, Blume=**Sapindus**, Linn.

APHELANDRA, R. Br. Prod. 475 (1810). *ACANTHACEAE*, Benth. & Hook. f. ii. 1102.
Amathea, Rafin. Fl. Tellur. iv. 65 (1836).
Hemisandra, Scheidw. in Bull. Acad. Brux. ix. 1. (1842) 22.
Hemitome, Nees, in DC. Prod. xi. 295 (1847).
Hydromestus, Scheidw. in Otto & Dietr. Gartenz. x. (1842) 285.
Lagochilium, Nees, in Mart. Fl. Bras. ix. 85. t. 10 (1847).
Poecilocnemis, Mart. ex Nees, in DC. Prod. xi. 295 (1847).
Strobilorhachis, Link, Klotzsch & Otto, Ic. Pl. Rar. 117. t. 48 (1839).
Synandra, Schrad. in Goett. Gel. Anz. i. (1821) 715.
acanthifolia, *Hook. Ic. Pl.* t. 113.—Peruv.
Acanthus, *Nees, in DC. Prod.* xi. 302.—Ecuador.
acutifolia, *Nees, l. c.* 299.—Ind. occ.
atroviirens, *N. E. Br. in Illustr. Hortic.* xxxi. (1884) t. 527.—Bras.
aurantiaca, *Lindl. Bot. Reg.* 31. t. 12.—Mexic.
bullata, *H. Wendl. in Hamb. Gartenz.* xix. (1863) 30. —Am. centr.
carduifolia, Hook. Ic. Pl. t. 718=formosa.
chartacea, *T. Anders. in Journ. Bot.* ii. (1864) 289.— Peruv.
Chamissoniana, *Nees, in Mart. Fl. Bras.* ix. 90.— Bras.
Clava, *Wawra, in Oestr. Bot. Zeitschr.* xii. (1862) 174. —Bras.
cristata, *R. Br. Prod.* 475, in obs.—Ind. occ.
cristata, H. B. & K. Nov. Gen. et Sp. ii. 236= pectinata.
cristata, Lindl. Bot. Reg. t. 1477=tetragona.
Dehnhardtii, *Tenore, Ind. Sem. Hort. Neap.* (1829) 14.—Hab. ?
Deppeana, *Cham. & Schlecht. in Linnaea*, v. (1830) 96.—Mexic.
Fascinator, *Linden & André, Illustr. Hortic.* xxi. (1874) t. 164.—Am. trop.
flava, *Nees, in DC. Prod.* xi. 297.—Reg. Colomb.
formosa, *Nees, l. c.* 301.—Peruv.
fulgens, *Decne. in Rev. Hortic.* Ser. III. i. (1847) 21.— Mexic.
fulgens, *Hort. ex Regel & Rach, Gartenfl.* (1858) 342=tenuiflora.
glabrata, *Willd. ex Nees, in DC. Prod.* xi. 296.—Am. merid.
Ghiesbreghtiana, Hort. ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 339=Jacobinia Ghiesbreghtiana.
Haenkeana, *Nees, Pl. Haenk. ex DC. Prod.* xi. 298.— Am. centr.
Hartwegiana, *Nees, in Benth. Pl. Hartw.* 236.—Reg. Colomb.
Hieronymi, *Grisch. in Goett. Abh.* xxiv. (1879) 260.— Reg. Argent.
hirsuta, Mart. ex Nees, in DC. Prod. xi. 417=Beloperone hirsuta.
hydromestus, *Hemsl. Biol. Centr. Am. Bot.* ii. 512. — Mexic.
ignea, Nees, ex Steud. Nom. ed. II. i. 111=Stenandrium igneum.

APHELANDRA:—
illustrata.—*Cf. Gard. Chron.* v. (1876) 603.—Bras.
lateritia, *Loud. Encyc. Pl.* 1539.—Guatemala.
Leopoldii, Van Houtte, in Fl. des Serres, ix. Sér. I. (1853) 85 = squarrosa.
Liboniana, *Linden, ex Hook. f. Bot. Mag.* 5463.—Am. trop.
longiracemosa, Hort. ex Lindl. in Paxt. Fl. Gard. ii. 78 = Thyrsacanthus callistachys.
longiscapa, Hort. ex Lem. in Fl. des Serres, Sér. I. iii. (1847) 240 = Thyrsacanthus Lemairianus.
lutea, *Nees, in DC. Prod.* xi. 300.—Bras.
lyrata, *Nees, l. c.* 302.—Am. austr.
macrostachya, *Nees, in Mart. Fl. Bras.* ix. 88.—Bras.
Margaritae, *E. Morr. in Belg. Hortic.* xxxiii. (1883) 315. t. 19.—Bras.
marginata, *Nees & Morr. in Nov. Act. Nat. Cur.* xi. (1823) i. 52.—Bras.
mediaurata, *Cogn. & March. Pl. Ornem.* i. (1873) t. 30.—Bras.
melgariensis, *Nees, in DC. Prod.* xi. 297.—Reg. Colomb.
micans, *Moritz. ex Vatke, Ind. Sem. Hort. Berol.* (1876) *App.* 1.—Venezuela.
monticola, Mart. ex Nees, in DC. Prod. xi. 364 = Jacobinia monticola.
mucronata, *Nees, in DC. Prod.* xi. 301.—Peruv.
nemoralis, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 90.—Bras.
nervosa, *T. Anders. in Journ. Bot.* ii. (1864) 289.—Peruv.
nitens, *Hook. f. Bot. Mag.* t. 5741.—Reg. Colomb.
nitida, Nees & Mart. in Nov. Act. Nat. Cur. xi. 1. (1823) 57 = Geissomeria nitida.
nobilis, *Hort.—Cf. Gard. Chron.* (1867) 461.—Hab.?
nuda, *Nees, in Mart. Fl. Bras.* ix. 89.—Bras.
oostachya, *Wawra, in Oestr. Bot. Zeitschr.* xii. (1862) 172.—Bras.
ornata, *T. Anders. in Journ. Bot.* ii. (1864) 289.—Bras.
pectinata, *Willd. ex Nees, in DC. Prod.* xi. 297.—Am. austr.
Porteana, *Morel, in Fl. des Serres*, Sér. I. x. (1854–55) 41. t. 984.—Bras.
prismatica, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 78.—Am. centr.
pulcherrima, *H. B. & K. Nov. Gen. et Sp. Pl.* ii. 236.—Reg. Colomb.
pulcherrima, Willd. ex Nees, in DC. Prod. xi. 295 = tetragona.
pumila, *Nichols. Dict. Gard.* i. 91.—Bras.
punctata, *Hort.—Cf. Gard. Chron.* (1881) II. 86; *Illustr. Hortic.* xxix. (1882) t. 457.—Am. merid.
quercifolia, *Nees, in DC. Prod.* xi. 302.—Reg. Colomb.
repanda, *Nees, l. c.* 728.—Peruv.
Roezlii, Carr. in Rev. Hortic. (1872) 100 = aurantiaca.
runcinata, *Klotzsch, ex Nees, in DC. Prod.* xi. 302.—Reg. Colomb.
Schiedeana, *Cham. & Schlecht. in Linnaea*, v. (1830) 95.—Mexic.
sciophila, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 91.—Bras.
Sinclairiana, *Nees, in Benth. Bot. Voy. Sulph.* 146. t. 47.—Am. centr.
Sinitzini, *Linden, Cat.* n. 95 (1876); *Rev. Hortic.* (1876) 237.—Peruv.
squarrosa, *Nees, in Mart. Fl. Bras.* ix. 89.—Bras.
stephanophysa, *Nees, l. c.* 90.—Am. austr.
sulphurea, *Hook. f. Bot. Mag.* t. 5951.—Ecuador.
tarapotensis, Linden, ex Rafarin, in Rev. Hortic. (1877) 38 = Sinitzini.
tenuiflora, *Regel & Rach, Gartenfl.* (1858) 342.—Hab.?
tetragona, *Nees, in DC. Prod.* xi. 295.—Reg. Colomb.
tridentata, *Hemsl. Biol. Centr. Am. Bot.* ii. 513.—Costa Rica.
variegata, *Morel, Fl. des Serres*, Sér. I. x. (1854–55) 33. t. 981.—Hab.?
verticillata, *Nees, in DC. Prod.* xi. 281.—Mexic.

APHELANDROS, St. Lag. in Ann. Soc. Bot. Lyon, vii. 119 (1880) = **Aphelandra**, R. Br. (Acanth.).

APHELEXIS, D. Don, in Mem. Wern. Soc. v. (1826) 546 = **Helichrysum**, Tourn. (Compos.).
adhaerens, Boj. ex DC. Prod. vi. 217.—Madag.
Candollei, Boj. ex DC. l. c.—Madag.
ericoides, Sweet, Hort. Brit. ed. I. 223 = Helichrysum ericoides.

APHELEXIS:—
fasciculata, D. Don, in Mem. Wern. Soc. v. (1826) 546 = H. sesamoides.
filiformis, D. Don, l. c. 547 = H. sesamoides.
heterophylla, G. Don, in Sweet, Hort. Brit. ed. III. 378 = Helipterum heterophyllum.
humilis, D. Don, in Mem. Wern. Soc. v. (1826) 547 = H. humile.
hypnoides, DC. Prod. vi. 218.—Madag.
incana, G. Don, in Sweet, Hort. Brit. ed. III. 379 = Helipterum incanum.
lycopodiodes, Boj. in DC. Prod. vi. 217.—Madag.
macrantha, Paxt. Fl. Mag. xv. 269 = H. humile.
selaginifolia, DC. Prod. vi. 217.—Madag.
sesamoides, D. Don, in Mem. Wern. Soc. v. (1826) 546 = H. sesamoides.

APHELIA, R. Br. Prod. 251 (1810). *CENTROLEPIDEAE*, Benth. & Hook. f. iii. 1026.
BRIZULA, Hieron. in Abh. Naturf. Ges. Halle, xii. (1873) 206, 208 (Centrolep. 92, 94).
Brizula, *F. Muell. Fragm.* v. 203.—Austral.
cyperoides, *R. Br. Prod.* i. 252.—Austral.
Drummondii, *Benth. Fl. Austral.* vii. 201.—Austral.
gracilis, *Sond. in Linnaea*, xxviii. (1855) 227.—Austral.
Gunnii, Hook. f. Fl. Tasm. ii. 75. t. 138 = gracilis.
monogyna, Hieron. in Abh. Naturf. Ges. Halle, xii. (1873) 208 (Centrolep. 94) = Centrolepis monogyna.
nutans, *Hook. f. ex Benth. Fl. Austral.* vii. 200.—Austral.
Pumilio, *F. Muell. in Linnaea*, xxviii. (1855) 226.—Austral.

APHILLANTHES, Neck. Elem. iii. 199 (1790) = **Aphyllanthes**, Tourn. (Liliac.).

APHLOIA, Benn. Pl. Jav. Rar. 192 (1838) in adnot. *BIXINEAE*, Benth. & Hook. f. i. 126.
NEUMANNIA, A. Rich. Fl. Cub. 96, in adnot. (1845).
integrifolia, Benn. Pl. Jav. Rar. 192 in adnot. = mauritiana.
madagascariensis, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 273.—Madag.
mauritiana, *Baker, Fl. Maurit.* 12.—Ins. Mascar.
theaeformis, Benn. Pl. Jav. Rar. 192, in adnot. = mauritiana.

APHOMA, Rafin. Fl. Tellur. ii. 31 (1836) = **Iphigenia**, Kunth (Liliac.).
angustiflora, Rafin. l. c. 32 = I. indica.
cuneata, Rafin. l. c. = I. indica.

APHOMONIX, Rafin. Fl. Tellur. ii. 65 (1836) = **Saxifraga**, Tourn.
hederacea, Rafin. l. c. = S. sarmentosa.

APHONINA, Neck. Elem. iii. 237 (1790) = **Pariana**, Aubl. (Gramin.).

APHORA, Neck. Elem. iii. 50 (1790) = **Virgilia**, Lam. (Legum.).

APHORA, Nutt. in Trans. Am. Phil. Soc. Ser. II. v. (1837) 174 = **Argithamnia**, P. Br. (Euphorb.).
Blodgettii, Torr. ex Chapm. Fl. S. U. St. 408.—Florida.
catamariensis, Griseb. in Goett. Abh. xxiv. (1879) 58 = Arg. catamariensis.
Drummondii, Engelm. & Gray, Pl. Lindh. i. 25.—Texas.
humilis, Engelm. & Gray, l. c. 54 = Arg. humilis.
humilis, Torr. Bot. Mex. Bound. 196 = Arg. neomexicana.
laevis, A. Gray, ex Torr. l. c. = Arg. laevis.
lanceolata, Engelm. & Gray, Pl. Lindh. i. 25 = Arg. lanceolata.
mercurialina, Nutt. in Trans. Am. Phil. Soc. N. S. v. (1837) 174 = Arg. mercurialina.
pilosissima, Torr. in Bot. Mex. Bound. 197 = Arg. mercurialina.
serrata, Torr. l. c. = Arg. serrata.

APHRAGMIA, Nees, in Lindl. Nat. Syst. ed. II. 444 (1836) = **Ruellia**, Linn. (Acanth.).
Haenkei, Nees, l. c. = R. albicaulis.
rotundifolia, Nees, ex Benth. Bot. Voy. Sulph. 146 = R. floribunda.

APHRAGMUS, Andrz. ex DC. Prod. i. 209 (1824) = **Braya**, Sternb. & Hoppe, (Crucif.).
Eschscholtziana, Andrz. ex DC. l. c.—Ins. Aleutian.

APHYLAX, Salisb. in Trans. Hort. Soc. i. (1812) 271 = **Aneilema**, R. Br. (Commelin.).
spiralis, Salisb. l. c. = An. spiratum.

APHYLLANTHES, Tourn. ex Linn. Gen. ed. I. 357 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 795.
juncea, Salisb. Parad. Lond. t. 9 = monspeliensis.
monspeliensis, *Linn. Sp. Pl.* 294.—Reg. Mediterr.

APHYLLEIA, Champ. in Calc. Journ. Nat. Hist. vii. (1847) 468 = **Sciaphila**, Blume (Triurid.).
erubescens, Champ. l. c. = S. erubescens.

APHYLLOCAULON, Lag. Amen. Nat. i. 38 (1811) = **Gerbera**, Gronov. (Compos.).

APHYLLOCLADUS, Wedd. Chlor. And. i. 11. t. 3 (1855) = **Hyalis**, D. Don (Compos.).
decussatus, Hieron. Ic. Pl. Argent. 28. t. 5. f. 1.—Reg. Argent.
spartioides, Wedd. l. c.—Bolivia.

APHYLLON, Mitch. in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) 221. *OROBANCHACEAE*, Benth. & Hook. f. ii. 983.
ANOPLON, Reichb. Consp. 212[b] (1828).
GYMNOCAULIS, Nutt. ex Benth. & Hook. f. Gen. ii. 983 (1876).
LOXANTHES, Rafin. Neogenyt. 3 (1825).
MYZORRHIZA, Phil. in Linnaea, xxix. (1857–58) 36.
THALESIA, Rafin. in Am. Monthly Mag. (1818) 267.
californicum, *A. Gray, in Bot. Calif.* i. 584.—Am. bor. occ.
comosum, *A. Gray, l. c.*—Am. bor. occ.
Cooperi, *A. Gray, in Proc. Am. Acad.* xx. (1885) 307.—Am. bor. occ.
Dugesii, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 132.—Mexic.
fasciculatum, *Torr. & Gray, in A. Gray, Man. Bot. U. St.* ed. I. 281.—Am. bor.
ludovicianum, *A. Gray, in Bot. Calif.* i. 585.—Am. bor.
multiflorum, *A. Gray, l. c.*—N. Mexic.
pinetorum, *A. Gray, l. c.*—Am. bor. occ.
tuberosum, *A. Gray, l. c.*—Calif.
uniflorum, *A. Gray, Man. Bot. U. St.* ed. I. 290.—Am. bor.

APHYLLORCHIS, Blume, 'Bijdr. t. 77 (1825)' (nobis latet). *ORCHIDEAE*, Benth. & Hook. f. iii. 606.
Hasseltii, *Blume, Mus. Bot. Lugd. Bat.* i. 31.—Java.
montana, *Reichb. f. in Linnaea*, xli. (1877) 57.—Ind. or.
pallida, *Blume, Bijdr.* t. 77; *Mus. Bot. Lugd. Bat.* i. 30.—Malaya.
Spiculaea, *Reichb. f. in Linnaea*, xli. (1877) 58.—Borneo.

APHYTEIA, Linn. Diss. Aphyt. 7 (1776); (Amoen. Acad. viii. 310. t. 7) (1785) = **Hydnora**, Thunb. (Cytin.).
Acharii, Steud. Nom. ed. II. i. 111, sphalm. (Cf. Thunb. Fl. Cap. ed. Schult. 499) = H. africana.
Hydnora, Linn. f. Diss. 10 = H. africana.
multiceps, Burch. Trav. i. 213, in nota = Cytinus dioicus.
triceps, Steud. Nom. ed. II. i. 111 = H. triceps.

APIASTRUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 643 (1840). *UMBELLIFERAE*, Benth. & Hook. f. i. 888.
angustifolium, *Nutt. l. c.* 644.—Calif.
latifolium, Nutt. l. c. = angustifolium.

APICRA, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 167. *LILIACEAE*, Benth. & Hook. f. iii. 776.
APRICA, D. Dietr. Syn. ii. 1097 (1840).
CATEVALA, Medic. Theod. 67 (1786).
albicans, Willd. l. c. 271 = Haworthia albicans.
anomala, Willd. l. c. 273 = Haworthia recurva
arachnoides, Willd. l. c. 168 = Haworthia arachnoides.
aspera, *Willd. l. c.* 274.—Afr. austr.

APICRA:—
asperula, Haw. ex Steud. Nom. ed. II. i. 111 = Haworthia asperula.
atrovirens, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 168 = Haworthia atrovirens.
attenuata, Willd. l. c. 270 = Haworthia attenuata.
bicarinata, *Haw. Suppl.* 63.—Afr. austr.
bullulata, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 273 = pentagona.
congesta, *Baker, in Journ. Linn. Soc.* xviii. (1880) 218. —Afr. austr.
cymbaefolia, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 271 = Haworthia cymbiformis.
deltoidea, *Baker, in Journ. Linn. Soc.* xviii. (1880) 217. —Afr. austr.
expansa, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 272 = Haworthia rigida.
fasciata, Willd. l. c. = Haworthia fasciata.
foliolosa, *Willd. l. c.* 274.—Afr. austr.
granata, Willd. l. c. 269 = Haworthia margaritifera.
imbricata, Willd. l. c. 273 = spiralis.
margaritifera, Willd. l. c. 269 = Haworthia margaritifera.
maxima, Steud. Nom. ed. I. 59; ed. II. i. 112 = Haworthia margaritifera.
minor, Steud. ll. cc. = Haworthia margaritifera.
mirabilis, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 269 = Haworthia mirabilis.
nigra, Haw. in Phil. Mag. (1824) 302 = Haworthia nigra.
patula, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 272 = Haworthia rigida.
pentagona, *Willd. l. c.* 273.—Afr. austr.
pseudo-rigida, Haw. Suppl. 62 = Haworthia subrigida.
Pumilio, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 271 = Haworthia reticulata.
quinquangularis, Sweet, Hort. Brit. ed. II. 521 = Aloë quinquangularis.
Radula, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 270 = Haworthia Radula.
recurva, Willd. l. c. = Haworthia recurva.
reticulata, Willd. l. c. 272 = Haworthia reticulata.
retusa, Willd. l. c. 271 = Haworthia retusa.
rigida, Willd. l. c. 272 = Haworthia subrigida.
spiralis, *Baker, in Journ. Linn. Soc.* xviii. (1880) 217. —Afr. austr.
spiralis, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 273 = pentagona.
spirella, Willd. ex Haw. Suppl. Pl. Succ. 64 = pentagona.
tortuosa, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 274 = Haworthia viscosa.
translucens, Willd. l. c. 168 = Haworthia translucens.
tricolor, Willd. l. c. 271 = Haworthia venosa.
viscosa, Willd. l. c. 274 = Haworthia viscosa.

APILIA, Rafin. Alsog. Am. 39 (1838) = **Fraxinus**, Tourn. (Oleac.).
laciniata, Rafin. l. c. = F. excelsior, var.?
macrophyla, Rafin. l. c. = F. macrophylla.
pendula, Rafin. l. c. = F. excelsior.

APINAGIA, Tul. in Ann. Sc. Nat. Sér. III. xi. (1849) 97. *PODOSTOMACEAE*, Benth. & Hook. f. iii. 111.
BLANDOWIA, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 100. t. 4. f. 2, nomen princeps.
MONOSTYLIS, Tul. Podost. Monog. 201 (1852).
NEOLACIS, Wedd. in DC. Prod. xvii. 59 (1873).
divaricata, *Tul. & Wedd. ex Tul. l. c.*—Bras.
fucoides, *Tul. l. c.* 98.—Bras.
Gardneriana, *Tul. l. c.*—Bras.
membranacea, *Tul. l. c.* 99.—Bras.
Preissii, *Tul. in Arch. Mus. Par.* vi. (1852) 108.—Italia.
psyllophora, *Tul. & Wedd. ex Tul. in Ann. Sc. Nat.* Sér. III. xi. (1849) 97.—Bras.
pusilla, *Tul. l. c.* 99.—Guiana.
pygmaea, *Tul. l. c.*—Bras.
Riedelii, *Tul. l. c.* 98.—Bras.
ruppioides, *Tul. l. c.*—Reg. Colomb.
striata, *Griseb. ex Wedd. in DC. Prod.* xvii. 64.—Peruv.
Willdenowii, *Tul. in Arch. Mus. Par.* vi. (1852) 106. —Peruv.; Chili.

APINELLA, Neck. Elem. i. 191 (1790) = **Trinia**, Hoffm. (Umbellif.).

APINUS, Neck. Elem. iii. 369 (1790) = **Pinus**, Tourn. (Conif.).

APIOCARPUS, Montrous. in Mém. Acad. Lyon, x. (1860) 190 = **Akania**, Hook.? (Sapindac.).
Moguini, Montrous. l. c. 191.—N. Caled.

APIOPETALUM, Baill. Adansonia, xii. (1878) 133. *ARALIACEAE*.
glabratum, *Baill. l. c.* 134.—N. Caled.
velutinum, *Baill. l. c.*—N. Caled.

APIOS, Moench, Meth. 165 (1794). *LEGUMINOSAE*, Benth. & Hook. f. i. 532.
BRADLEA, Adans. Fam. ii. 324 (1763).
CYRTOTROPIS, Wall. Pl. As. Rar. i. 49. t. 62 (1830).
americana, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 355.—Am. bor.
carnea, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 188.—Reg. Himal.
chinensis, Spreng. Syst. iii. 310 = Wisteria chinensis.
Fortunei, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 396.—China; Japon.
frutescens, Pursh, Fl. Am. Sept. ii. 474 = Wisteria frutescens.
tuberosa, *Moench, Meth.* 165.—Am. bor.

APIOSPERMUM, Klotzsch, in Abh. Akad. Berl. (1853) 351, 352 = **Pistia**, Linn. (Aroid.).
obcordatum, Klotzsch, l. c. = P. Stratiotes.

APIROPHORUM, Neck. Elem. ii. 72 (1790) = **Pyrus**, Tourn. (Rosac.).

APISTA, Blume, Bijdr. 296 (1825) = **Podochilus**, Blume (Orchid.).
tenuis, Blume, l. c. = P. tenuis.

APIUM, [Tourn.] Linn. Syst. ed. I. (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 888.
CELERI, Adans. Fam. ii. 498 (1763).
CICLOSPERMUM, Lag. Amen. Acad. Madr. ii. 101 (1811).
CRITAMUS, Hoffm. Gen. Umb. 182 (1814).
CYCLOSPERMUM, DC. Prod. iv. 105, in syn. (1830).
HELODIUM, Dum. Fl. Belg. 77 (1827).
HELOSCIA, Dum. Anal. Fam. 34 (1829).
HELOSCIADIUM, Koch, in Nov. Act. Nat. Cur. xii. 1. (1825) 125.
LEPTOCAULIS, Nutt. ex DC. Prod. iv. 107 (1830).
MAUCHARTIA, Neck. Elem. i. 172 (1790).
SELINON, Adans. Fam. ii. p. '(23)' 603 (1763).
SIUMIS, Rafin. New Fl. Am. iv. 31 (1836).
SPERMOLEPIS, Rafin. Neogenyt. 2 (1825).
Ammi, Crantz, Stirp. Austr. ed. I. fasc. iii. 109; ed. II. 217 = Ammi majus.
Ammi, *Urban, in Mart. Fl. Bras.* xi. 1. 341.—Bras.
Amomum, Stokes, Bot. Mat. Med. ii. 156 = Sison Amomum.
Anisum, Crantz, Cl. Umb. Emend. 101 = Pimpinella Anisum.
australe, Thou. Fl. Trist. d'Acugn. 43 = prostratum.
bipinnatum, Walt. Fl. Carol. 115 = Selinum canadense.
biternatum, Stokes, Bot. Mat. Med. ii. 150 = Aegopodium Podagraria.
Carsonii, *Benth. & Hook. f. Gen.* i. 888.—Am. bor.
Carvi, Crantz, Stirp. Austr. ed. I. fasc. iii. 110; ed. II. 218 = Carum Carvi.
Celleri, Gaertn. Fruct. i. t. 22 = graveolens.
chilense, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 353.—Chili.
Commersoni, *DC. Prod.* iv. 101.—Hab.?
crassipes, *Reichb. f. Ic. Fl. Germ.* xxi. 9.—Hispan.
crispum, Mill. Gard. Dict. ed. VIII. n. 2 = Carum Petroselinum.
decumbens, Eckl. & Zeyh. Enum. 340 = graveolens.
divaricatum, *Benth. & Hook. f. ex S. Wats. Bibl. Ind. N. Am. Bot.* 412.—Am. bor.
dulce, Mill. Gard. Dict. ed. VIII. n. 5 = graveolens.
echinatum, *Benth. & Hook. f. ex S. Wats. Bibl. Ind. N. Am. Bot.* 412.—Am. bor.
filiforme, Hook. Ic. t. 819 = prostratum.
flexuosum, *Phil. in Linnaea*, xxxiii. (1864–65) 91.—Chili.
fractophyllum, *Hornem. Hort. Hafn. Suppl.* 128.—Hab.?
glaucescens, *H. B. & K. Nov. Gen. et Sp.* v. 17.—Reg. Colomb.
glaucum, Crantz, Cl. Umb. Emend. 100 = Trinia vulgaris.
gracile, *Benth. & Hook. f. Gen.* i. 888.—Chili.

APIUM:—
graveolens, Cham. in Linnaea, i. (1826) 388 = prostratum.
gravcolens, *Linn. Sp. Pl.* 264.—Europ.; Orient; Ind. bor. occ.; Calif.
humile, *Benth. & Hook. f. Gen.* i. 888.—Afr. austr.
inundatum, *Reichb. f. Ic. Fl. Germ.* xxi. 9.—Europ.
involucratum, Roxb. ex Flem. in As. Res. xi. (1810) 157 = Carum Roxburghianum.
junceum, Stokes, Bot. Mat. Med. ii. 155 = Carum segetum.
laetum, Salisb. Prod. 169 = Carum Petroselinum.
latifolium, Mill. Gard. Dict. ed. VIII. n. 3 = Carum Petrosclinum.
latifolium, Poir. Encyc. iv. 369 = Carum peregrinum.
leptophyllum, *F. Muell. ex Benth. Fl. Austral.* iii. 372. —Austral.
lineare, *Benth. & Hook. f. Gen.* i. 888.—Am. bor.
lobatum, Gilib. Fl. Lituan. ii. 41 = graveolens.
lusitanicum, Mill. Gard. Dict. ed. VIII. n. 7 = graveolens.
maritimum, Salisb. Prod. 169 = graveolens.
montanum, *H. B. & K. Nov. Gen. et Sp.* v. 17.—Reg. Colomb.
nodiflorum, *Reichb. f. Ic. Fl. Germ.* xxi. 10.—Europ.
patens, *S. Wats. Bibl. Ind. N. Am. Bot.* 413.—Am. bor.
peregrinum, Crantz, Cl. Umb. Emend. 101 = Carum peregrinum.
petraeum, Crantz, l. c. 104 = Ammi majus.
Petroselinum, Cham. in Linnaea, i. (1826) 389 = prostratum.
Petroselinum, Linn. Sp. Pl. ed. I. 264 = Carum Petroselinum.
Popei, *A. Gray, in Proc. Am. Acad.* vii. (1868) 343.—Am. bor.
prostratum, *Labill. Relat.* i. 141; *Vent. Jard. Malm.* 81.—Austral.; Reg. Antarct.
pumilum, Crantz, Stirp. Austr. ed. I. fasc. iii. 103; ed. II. 213 = Trinia vulgaris.
ranunculifolium, *H. B. & K. Nov. Gen. et Sp.* v. 18. t. 421.—Reg. Colomb.
rapaceum, Mill. Gard. Dict. ed. VIII. n. 6 = graveolens.
repens, *Reichb. f. Ic. Fl. Germ.* xxi. 10.—Europ.
romanum, Zuccagni, in Roem. Collect. 135 = Carum Petroselinum.
segetum, Dum. Fl. Belg. 71 = Carum segetum.
Sium, Crantz, Stirp. Austr. ed. I. fasc. iii. 106; ed. II. 215 = Sium erectum.
ternatum, Pall. ex Steud. Nom. ed. II. i. 112 = Archangelica Gmelini.
ternatum, Schlecht. ex Roem. & Schult. Syst. vi. 431 = Ligusticum scoticum.
ternatum, Siebold, in Verh. Batav. Gen. xii. (1830) 46 = Ligusticum acutilobum.
trifoliatum, Wight & Arn. Prod. i. 368, sphalm. = Pimpinella Heyneana.
tuberosum, Steud. Nom. ed. I. 59 = Carum Petroselinum
Visnaga, Crantz. Cl. Umb. Emend. 104 = Ammi Visnaga.
vulgare, Lam. Fl. Fr. iii. 444 = Carum Petroselinum.

APLACTIS, Rafin. Fl. Tellur. ii. 42 (1836) = **Solidago**, Tourn. (Compos.).
paniculata, Rafin. l. c. = S. pauciflosculosa.

APLARINA, Rafin. New Fl. Am. iv. 99 (1836) = **Euphorbia**, Tourn.
microphyla, Rafin. l. c. 100 = E. thymifolia.
prostrata, Rafin. l. c. 99 = E. prostrata.

APLECTRA, Rafin. Cat. 13 (1824) = **Aplectrum**, Nutt. (Orchid.).
elatior, Rafin. l. c. = Aplectrum hyemale.

APLECTROCAPNOS, Boiss. & Reut. Diagn. Ser. I. v. 79 (1844) = **Sarcocapnos**, DC. (Papav.).
boetica, Boiss. & Reut. l. c. = S. baetica.
integrifolia, Boiss. Diagn. Ser. II. i. 13 = S. baetica.

APLECTRUM, Torr. Comp. Fl. St. 322 (1826). *ORCHIDEAE*, Benth. & Hook. f. iii. 497.
APLECTRA, Rafin. Cat. 13 (1824).
hyemale, *Torr. l. c.*—Am. bor.

APLECTRUM, Blume, in Flora, xiv. (1831) 502 = **Anplectrum**, A. Gray (Melast.).
confine, Blume, Mus. Bot. Lugd. Bat. i. 38 = Anpl. confine
myrtiforme, Naud. in Ann. Sc. Nat. Ser. III. xv. (1851) 305 = Medinilla myrtiformis.
nodosum, Blume, Mus. Bot. Lugd. Bat. i. 37 = Anpl nodosum.

APLECTRUM :—
ovalifolium, Naud. ex A. Gray, Bot. U. St. Expl. Exp. i. 597 = Allomorpha ovalifolia.
pallens, Blume, Mus. Bot. Lugd. Bat. i. 38 = Anplectrum pallens.
rostratum, Blume, in Flora, xiv. (1831) 502 = Anpl. rostratum.
stipulare, Blume, l. c. 503 = Anpl. stipulare.
viminale, Blume, l. c. 502 = Anpl. viminale.

APLEURA, Phil. in Linnaea, xxxiii. (1864-65) 89. *UMBELLIFERAE*, Benth. & Hook. f. i. 875.
nucamentacea, *Phil. l. c.* 90.—Chili.

APLEXIA, Rafin. Neogenyt. 4 (1825) = **Leersia**, Sw. (Gram.).
virgata, Rafin. in Ser. Bull. Bot. 220 = L. virginica.
virginica, Rafin. Neogenyt. 4 = L. virginica.

APLILIA, Rafin. New Fl. Am. iii. 93 (1836) = **Fraxinus**, Tourn. (Oleac.).
laciniata, Rafin. l. c. = F. excelsior.
macrophyla, Rafin. l. c. = F. excelsior.

APLINA, Rafin. Fl. Tellur. iv. 119 (1836) = **Staehelina**, Linn. (Compos.).
fuscata, Rafin. l. c. = S. unifloscula.

APLOCA, Neck. Elem. i. 255 (1790) = **Periploca**, Tourn. (Asclep.).

APLOCARYA, Lindl. Bot. Reg. (1844) sub t. 46 = **Dolia**, Lindl. (Convolv.).
divaricata, Lindl. l. c.—Chili.

APLOCHLAMIS, Steud. Nom. ed. II. i. 309, in syn. (1840) sphalm. = APALOCHLAMYS, Cass. = **Cassinia**, R. Br. (Compos.).

APLOLEIA, Rafin. Fl. Tellur. ii. 17 (1836) = **Tradescantia**, Linn. (Commelin.).
diffusa, Rafin. l. c.—Hab. ?

APLOLOPHIUM, Cham. in Linnaea, vii. (1832) 556. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1038.
HAPLOLOPHIUM, Endl. Gen. 712 (1839).
Blanchetii, *DC. Prod.* ix. 192.—Bras.
bracteatum, *Cham. in Linnaea*, vii. (1832) 556.—Bras.
Lundii, *DC. Prod.* ix. 192.—Bras.

APLOPAPPUS, Cass. in Dict. Sc. Nat. lvi. 168 (1828). *COMPOSITAE*, Benth. & Hook. f. ii. 253.
ACAMPHOPAPPUS, A. Gray, in Proc. Am. Acad. viii. (1873) 634.
CHROILEMA, Bernh. Del. Sem. Hort. Erf. (1840).
CROILEMA, Heynh. Nom. ii. 171 (1846).
CROPTILON, Rafin. Fl. Tellur. ii. 47 (1836).
ERIOCARPUM, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 320.
HAPLOPAPPUS, Endl. Gen. 385 (1838).
HOMOPAPPUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 330.
HOOREBEKIA, Corneliss. in Mussch. Hort. Gand. 120 (1817), nomen princeps.
ISOPAPPUS, Torr. & Gray, Fl. N. Am. ii. 239 (1841).
PRIONOPSIS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 329.
PYRROCOMA, Hook. Fl. Bor. Am. i. 306. t. 107 (1833).
SIDERANTHUS, Fras. ex Sweet, Hort. Brit. ed. I. 227 (1826).
STENOTUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 334.
acaulis, *A. Gray, in Proc. Am. Acad.* vii. (1868) 353.—Calif.
aceros us, *Bert. ex Phil. in Linnaea*, xxviii. (1856) 724.—Chili.
acuminatus, DC. Prod. v. 348 = Leucopsis sericeus.
alpigenus, Torr. & Gray, Fl. N. Am. ii. 241 = Aster pulchellus.
alternifolius, Steud. Nom. ed. II. i. 87, sphalm. = chrysanthemifolius.
anthylloides, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 261.—Chili.
apargioides, *A. Gray, in Proc. Am. Acad.* vii. (1868) 354.—Calif.
araucanus, *Phil. in Anal. Univ. Chil.* (1873) 491.—Chili.
arbutoides, *Remy, in C. Gay, Fl. Chil.* iv. 53.—Chili.
arenarius, *Benth. Bot. Voy. Sulph.* 24 (Asteris sp. ?).—Calif.
argenteus, *Steud. Nom.* ed. II. i. 112.—Chili.
armerioides, *A. Gray, Color. Exp. Surv. Bot.* (1860) 16.—N. Mexic.
aureus, *A. Gray, Pl. Fendl.* 76.—Texas.
baccharoides, Benth. Bot. Voy. Sulph. 24 = Solidago occidentalis.
Baylahuen, *Remy, in C. Gay, Fl. Chil.* iv. 42.—Chili.
bellidioides, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 112 = Erigeron pappochroma.
Berberidis, *A. Gray, Syn. Fl. N. Am.* i. II. 126.—Calif.
Berterii, *DC. Prod.* v. 348.—Chili.
blephariphyllus, A. Gray, Pl. Wright. i. 97 = Aster gymnocephalus.
Bloomeri, *A. Gray, in Proc. Am. Acad.* vi. (1865) 541.—Am. bor.
Brandegei, *A. Gray, Syn. Fl. N. Am.* i. II. 132.—Am. bor.
bustillosianus, *Remy, in C. Gay, Fl. Chil.* iv. 49.—Chili.
caespitosa, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 330.—Chili.
calendulaceus, *Griseb. in Goett. Abh.* xxiv. (1879) 179.—Reg. Argent.
canescens, DC. Prod. v. 349 = Chrysopsis villosa.
carthamoides, *A. Gray, in Proc. Acad. Sc. Philad.* (1863) 65.—Am. bor.
cervinus, *Rothrock, in Wheeler, Geogr. Surv.* 40. t. 6.—Am. bor.
cervinus, S. Wats. in Am. Natural. vii. (1873) 301 = nanus.
Chamissonis, *DC. Prod.* v. 347.—Chili.
chrysanthemifolius, *DC. l. c.* 348.—Chili.
ciliatus, *DC. l. c.* 346.—Am. bor.
coronopifolius, *DC. l. c.* 347.—Chili.
Coulteri, Harv. & Gray, ex A. Gray, Pl. Fendl. 75 = spinulosus.
croceus, *A. Gray, in Proc. Acad. Sc. Philad.* '1863' (1864) 65.—Am. bor.
cuneatus, *A. Gray, in Proc. Am. Acad.* viii. (1873) 635.—Calif.
cuneifolius, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 330.—Chili.
decurrens, *Remy, in C. Gay, Fl. Chil.* iv. 58.—Chili.
densifolius, *Remy, l. c.* 53.—Chili.
diffusus, DC. Prod. v. 349 = Leucopsis diffusa.
Diplopappus, *Remy, in C. Gay, Fl. Chil.* iv. 53.—Chili.
discoideus, DC. Prod. v. 350 = Bigelovia veneta.
divaricatus, *A. Gray, in Pacif. Rail. Rep.* iv. 99.—Am. bor.
Doellianus, *Phil. in Anal. Univ. Chil.* (1873) 489.—Chili.
ericoides, *DC. Prod.* v. 347.—Calif.
filifolius, DC. Prod. vii. 280 = Erigeron filifolius ?
florifer, Hook. & Arn. Bot. Beech. Voy. 351 = Townsendia florifera.
foliosus, *DC. Prod.* v. 346.—Chili.
Fremonti, *A. Gray, in Proc. Acad. Sc. Philad.* (1863) 65.—Am. bor.
glabratus, *Phil. in Linnaea*, xxviii. (1856) 727.—Chili.
glutinosus, Cass. ex DC. Prod. v. 347 = coronopifolius, pulchellus.
glutinosus, *DC. Prod.* v. 347.—Chili.
gracilis, *A. Gray, Pl. Fendl.* 76.—Texas.
gramineus, Benth. Pl. Hartw. 65 (err. typ. 61) = Chrysopsis graminifolia.
Greenei, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 80.—Am. bor.
grindelioides, *DC. Prod.* v. 348.—Chili.
Gunnii, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 111. = Erigeron pappochroma.
gymnocephalus, DC. Prod. v. 346 = Aster gymnocephalus.
Haenkei, DC. l. c. 349 = Corethrogyne filaginifolia.
Hallii, *A. Gray, in Proc. Am. Acad.* viii. (1872) 389.—Oregon.
hirtellus, *Phil. in Linnaea*, xxviii. (1856) 726.—Chili.
hirtus, *A. Gray, Syn. Fl. N. Am.* i. II. 127.—Am. bor.
hispidulus, *DC. Prod.* v. 348.—Chili.
Hookerianus, *A. Gray, Syn. Fl. N. Am.* i. II. 131.—Texas.
Hookerianus, *DC. Prod.* vii. 279.—Chili.
Hookerianus, A. Gray, Pl. Lindh. i. 40 = divaricatus.
Hoorebekia, *DC. Prod.* v. 346.—Chili.
Howellii, *A. Gray, Syn. Fl. N. Am.* i. II. 446.—Oregon.
hypoleucus, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 177 = Bigelovia hypoleuca.
ilicifolius, *Remy, in C. Gay, Fl. Chil.* iv. 55.—Chili.
illinitus, *Phil. in Linnaea*, xxviii. (1856) 727.—Chili.
integrifolius, *T. C. Porter, ex A. Gray, in Proc. Am. Acad.* xvi. (1881) 79.—Am. bor.
inuloides, Torr. & Gray, Fl. N. Am. ii. 241 = uniflorus.
junceus, Greene, in Bull. Acad. Calif. i. (1884) 190 = spinulosus.
lanatus, *Hook. f. in Trans. Linn. Soc.* xx. (1847) 215.—Ins. Galapag.
lanceolatus, *Torr. & Gray, Fl. N. Am.* ii. 241.—Calif.
lanuginosus, *A. Gray, Bot. U. St. Expl. Exped.* ii. 347.—Am. bor.
laricifolius, *A. Gray, Pl. Wright.* ii. 80.—N. Mexic.
Lastarrianus, *Remy, in C. Gay, Fl. Chil.* iv. 45.—Chili.
linearifolius, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 457 = Baccharis texana ?.
linearifolius, *DC. Prod.* v. 347.—Calif.
Lyallii, *A. Gray, in Proc. Acad. Sc. Philad.* '1863' (1864) 64.—Am. bor.
Macraeanus, *A. Gray, in Proc. Am. Acad.* v. (1861) 122.—Chili.
macrocephalus, *DC. Prod.* v. 348.—Chili.
Macronema, *A. Gray, in Proc. Am. Acad.* vi. (1864) 542.—Am. bor.
marginalis, *Phil. in Linnaea*, xxxiii. (1864-65) 139.—Chili.
marginatus, *Griseb. Cat. Pl. Cub.* 149.—Cuba.
medicinalis, *Phil. in Linnaea*, xxviii. (1856) 729.—Chili.
Menziesii, Torr. & Gray, Fl. N. Am. ii. 241 = Bigelovia Menziesii.
Meyenii, *Walp. in Nov. Act. Cur.* xix. *Suppl.* i. (1843) 261.—Chili.
mollis, A. Gray, in Proc. Am. Acad. xvi. (1881) 80 = Greenei.
monactis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 1.—Calif.
mucronatus, *Hook. & Arn. Bot. Beech. Voy.* 147.—Chili.
multicaulis, *A. Gray, in Am. Natural.* viii. (1874) 213.—Am. bor.
nanus, *Eaton, in Bot. King, Exped.* 159.—Am. bor.
nevadensis, *Kellogg, in Proc. Calif. Acad.* iii. (1863) 9.—Calif.
Nuttallii, *Torr. & Gray, Fl. N. Am.* ii. 242.—Am. bor.
ochagavianus, *Phil. in Linnaea*, xxviii. (1856) 725.—Chili.
Orcuttii, *A. Gray, in Proc. Am. Acad.* xx. (1885) 297.—Am. bor.
Palmeri, *A. Gray, l. c.* xi. (1876) 74.—Calif.
paniculatus, A. Gray, l. c. vii. (1868) 354 = racemosus.
paniculatus, *Phil. in Anal. Univ. Chil.* (1873) 490.—Chili.
pappochroma, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 111 = Erigeron pappochroma.
Parryi, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 239.—Am. bor.
parvifolius, *A. Gray, in Proc. Am. Acad.* v. (1861) 122.—Chili.
patagonicus, *Phil. in Linnaea*, xxviii. (1856) 726.—Chili.
paucidentatus, *Phil. in Anal. Univ. Chil.* (1873) 492.—Chili.
pectinatus, *Phil. in Linnaea*, xxxiii. (1864-65) 140.—Chili.
pedunculosus, *Remy, in C. Gay, Fl. Chil.* iv. 51.—Chili.
phyllocephalus, DC. Prod. v. 347 = rubiginosus.
pinifolius, *A. Gray, in Proc. Am. Acad.* viii. (1873) 636.—Calif.
pinnatifidus, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 330.—Am. bor.
Poeppigianus, *A. Gray, in Proc. Am. Acad.* v. (1862) 121.—Chili.
polyphyllus, *Phil. in Linnaea*, xxx. (1859-60) 192.—Chili.
prunelloides, *DC. Prod.* v. 346.—Chili.
pulchellus, *DC. l. c.* 347.—Chili.
pygmaeus, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 239.—Am. bor.
racemosus, *Torr. in Sitgreave, Rep.* 162.—Am. bor.
radicans, *Remy, in C. Gay, Fl. Chil.* iv. 45.—Chili.
ramulosus, DC. Prod. v. 350 = Baccharis ramulosa.
reflexus, *Phil. in Linnaea*, xxviii. (1856) 728.—Chili.
Remyanus, *Wedd. Chlor. And.* i. 210.—Chili.
Rengifoanus, *Remy, in C. Gay, Fl. Chil.* iv. 43.—Chili.
resinosus, A. Gray, Bot. U. St. Expl. Exped. ii. 346 = nanus.
reticulatus, *Phil. in Linnaea*, xxviii. (1856) 725.—Chili.

APLOPAPPUS:—
rigidus, *Phil. Fl. Atac.* 30.—Chili.
rubiginosus, *Torr. & Gray, Fl. N. Am.* ii. 240.—Am. bor.
scaposus, DC. Prod. v. 350 = Leucopsis scaposa.
scaposus, *Remy, in C. Gay, Fl. Chil.* iv. 50.—Chili.
scrobiculatus, *DC. Prod.* v. 348.—Chili.
sericeus, *DC. l. c.* 349.—Chili.
sericeus, Phil. in Linnaea, xxviii. (1856) 724 = Poeppigianus.
sphaerocephala, *Harv. & Gray, ex A. Gray, Pl. Fendl.* 76.—Calif.
spinulosus, *Phil. in Anal. Univ. Chil.* (1873) 490.—Chili.
spinulosus, *DC. Prod.* v. 347.—Am. bor.
squarrosus, *Hook. & Arn. Bot. Beech. Voy.* 146.—Calif.
stellatus, Hook. f. in Lond. Journ. Bot. vi. (1847) 112 = Erigeron pappochroma.
stelliger, *Remy, in C. Gay, Fl. Chil.* iv. 52.—Chili.
stenophyllus, *A. Gray, Bot. U. St. Expl. Exped.* ii. 347.—Calif.
stoloniferus, *DC. Prod.* v. 349.—Mexic.
suffruticosus, Eaton, in Bot. King, Exped. 159 = Watsoni.
suffruticosus, *A. Gray, in Proc. Am. Acad.* vi. (1865) 542.—Calif.
tasmanicus, Hook. f. in Lond. Journ. Bot. vi. (1847) 110 = Erigeron pappochroma.
tenuicaulis, Eaton, in Bot. King, Exped. 160 = lanceolatus.
tortifolius, Torr. & Gray, in Bost. Journ. Nat. Hist. v. (1845) 109 = Aster tortifolius.
uncinatus, *Phil. in Linnaea*, xxviii. (1856) 728.—Chili.
uniflorus, *Torr. & Gray, Fl. N. Am.* ii. 241.—Am. bor.
velutinus, *Remy, in C. Gay, Fl. Chil.* iv. 57.—Chili.
villosus, *Phil. in Linnaea*, xxviii. (1856) 725.—Chili.
virgatus, *Phil. in Anal. Univ. Chil.* (1873) 491.—Chili.
Watsoni, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 79.—Am. bor.
Whitneyi, *A. Gray, l. c.* vii. (1868) 353.—Calif.

APLOPHYLLUM, Cass. in Dict. Sc. Nat. xxxiii. 472 (1824) = **Mutisia**, Linn. (Compos.).
decurrens, Cass. l. c. = M. decurrens.
ilicifolium, Cass. l. c. = M. ilicifolia.
linearifolium, Cass. l. c. = M. linearifolia.

APLOPHYLLUM, A. Juss. in Mém. Mus. Par. xii. (1825) 464 = **Ruta**, Linn.
acutifolium, G. Don, Gen. Syst. i. 780 = R. acutifolia.
arabicum, Boiss. Diagn. Ser. I. viii. 127 = R. tuberculata.
arbusculum, Franch. in Revoil, Çomali, 23.—Afr. trop.
armenum, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 179 = R. suaveolens?
Besseri, Spach, l. c. = R. Besseri.
Biebersteinii, Spach, l. c. 178 = R. Buxbaumii.
Blanchei, Boiss. Fl. Orient. i. 937.—Orient.
Boissieianum, Viv. & Panč. in Mem. Ist. Venet. xv. (1870) 14 = R. patavina.
Bourgaei, Boiss. Fl. Orient. i. 928.—Armenia.
Broussonetianum, Coss. in Bull. Soc. Bot. Fr. xx. (1873) 244; xxii. (1875) 56, nomen.—Afr. bor.
Buhsei, Boiss. Fl. Orient. i. 926.—Persia.
Bungei, Trautv. in Act. Hort. Petrop. viii. (1883) 183.—Turkestan.
Buxbaumii, G. Don, Gen. Syst. i. 780 = R. Buxbaumii.
Buxbaumii, Schur, Enum. Pl. Transs. 140 = R. suaveolens.
canaliculatum, Boiss. Diagn. Ser. I. viii. 128.—Persia.
Candolleanum, Spach, ex Jaub. & Spach, Illustr. Pl. Or. t. 270 = R. glabra.
cappadocicum, Spach, ex Jaub. & Spach, l. c. t. 263.—Cappadoc.
Chesneyanum, Boiss. Fl. Orient. i. 941.—Mesopot.
ciliatum, Griseb. Spicil. Fl. Rumel. i. 130 = R. suaveolens.
cilicicum, Boiss. Fl. Orient. i. 927.—Cilicia.
congestum, Jaub. & Spach, Illustr. Pl. Or. iii. 77 = R. suaveolens.
consanguineum, Bunge, ex Boiss. Fl. Orient. i. 941.—Persia.
cordatum, G. Don, Gen. Syst. i. 780 = R. cordata.
coronatum, Griseb. Spicil. Fl. Rumel. i. 129 = R. patavina.
corymbulosum, Boiss. Diagn. Ser. II. i. 116 = R. Buxbaumii.

APLOPHYLLUM:—
crenulatum, Boiss. Diagn. Ser. II. i. 117.—Beluchist.
dahuricum, Juss. in Mém. Mus. Par. xii. (1825) 464 = Ruta dahurica.
davuricum, Ledeb. Fl. Ross. i. 492 = praec.
dichotomum, G. Don, Gen. Syst. i. 780 = R. dichotoma.
dichotomum, Griseb. Spicil. Fl. Rumel. i. 131 = R. Buxbaumii.
elatum, Fisch. & Mey. ex Kar. in Bull. Soc. Nat. Mosc. (1839) 149.—Turcomania.
erythraeum, Boiss. Diagn. Ser. II. i. 114.—Afghan.
filifolium, Spach, ex Jaub. & Spach, Illustr. Pl. Or. t. 265.—Persia.
flexuosum, Boiss. Diagn. Ser. II. i. 118.—Afghan.
fruticulosum, G. Don, Gen. Syst. i. 780 = R. fruticulosa.
furfuraceum, Bunge, in Boiss. Fl. Orient. i. 934.—Persia.
glaberrimum, Bunge, l. c. 930.—Persia.
glabrum, G. Don, Gen. Syst. i. 780 = R. glabra.
Griffithianum, Boiss. Diagn. Ser. II. i. 115.—Afghan.
halepense, Spach, in Jaub. & Spach, Illustr. Pl. Or. t. 266 = R. Buxbaumii.
hirsutum, Regel & Schmalh. ex Regel, Pl. Nov. Fedsch. 17.—As. temp. or.
hispanicum, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 176 = R. linifolia.
hispidulum, Bunge, in Boiss. Fl. Orient. i. 933.—Persia.
Kotschyi, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 186.—Persia.
Kowalenskyi, Stschég. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 467.—Reg. Caucas.
lasianthum, Bunge, in Schrift. Naturf. Ver. Riga, i. (1847) 206; et Pl. Lehm. 239.—Turkest.
latifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 398.—As. cent.
lineare, G. Don, Gen. Syst. i. 780 = R. linearis.
linifolium, G. Don, l. c. = R. linifolia.
linifolium, Reichb. ex Nym. Consp. 142 = R. patavina.
linifolium, Schur, Enum. Pl. Transs. 140 = R. suaveolens.
longifolium, Boiss. Diagn. Ser. I. viii. 127.—Palaestin.
mesopotamicum, Boiss. l. c. Ser. II. i. 116 = R. Buxbaumii.
myrtifolium, Boiss. l. c. Ser. I. i. 63.—As. Min.
obtusifolium, Ledeb. Fl. Ross. i. 490 = R. obtusifolia.
patavinum, G. Don, Gen. Syst. i. 780 = R. patavina.
pedicellatum, Bunge, ex Boiss. Fl. Orient. i. 925.—Persia.
perforatum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 397 = R. acutifolia.
pilosum, Franch. in Ann. Sc. Nat. Sér. VI. xv. (1883) 248, t. 13.—Turkest.
pilosum, Szczegl. ex Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 273.—Persia.
propinquum, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 189.—Mesopot.
ptilostylum, Spach, ex Jaub. & Spach, Illustr. Pl. Or. t. 267.—Persia.
pubescens, Boiss. Voy. Bot. Espagne, i. 371 = R. linifolia.
pubescens, G. Don, Gen. Syst. i. 780 = R. pubescens.
pumilum, Boiss. Diagn. Ser. I. i. 62.—As. Min.
Reuteri, Boiss. l. c. I. v. 87.—Caria.
robustum, Bunge, in Schrift. Naturf. Ver. Riga, i. (1847) 208, et Pl. Lehm. 249.—Persia.
rosmarinifolium, G. Don, Gen. Syst. i. 781 = R. rosmarinifolia.
Sieversii, Fisch. in Schrenk, Enum. Pl. Nov. 89 = R. acutifolia.
spathulatum, Reichb. ex Nym. Consp. 143 = R. Buxbaumii.
Stocksianum, Boiss. Diagn. Ser. II. i. 117.—Beluchist.
suaveolens, G. Don, Gen. Syst. i. 780 = R. suaveolens.
suaveolens, Sweet, Hort. Brit. ed. III. 126 = R. Buxbaumii.
subcordatum, Bunge, in Boiss. Fl. Orient. i. 932.—Persia.
sulphureum, Boiss. Diagn. Ser. I. v. 87.—As. Min.
sylvaticum, Boiss. l. c. viii. 126.—Syria.
tauricum, Spach, in Ann. Sc. Nat. Sér. III. xi. (1849) 178 = R. taurica.
Tchihatchewi, Boiss. Diagn. Ser. II. vi. 41 = *sulphureum*.
telephioides, Boiss. Fl. Orient. i. 929.—Cappadoc.
tenue, Boiss. l. c. 932.—Armenia.
thesioides, G. Don, Gen. Syst. i. 780 = R. thesioides.

APLOPHYLLUM:—
trichostylum, Bunge, in Boiss. Fl. Orient. i. 941.—Persia.
tuberculatum, Juss. in Mém. Mus. Par. xii. (1825) 528. t. 17 = R. tuberculata.
versicolor, Fisch. & Mey. in Bull. Phys. Math. Akad. Pétersb. iii. (1845) 307 = *lasianthum*.
villosulum, Boiss. & Haussk. in Boiss. Fl. Orient. i. 936.—Syria.
villosum, G. Don, Gen. Syst. i. 780 = R. villosa.
virgatum, Spach, ex Jaub. & Spach, Illustr. Pl. Or. t. 267.—Persia.
vulcanicum, Boiss. & Heldr. Diagn. Ser. I. viii. 126.—Lycaonia.

APLOSTELLIS, Thou. Orch. Iles Afr. t. 24 (1822) = **Pogonia**, Juss. (Orchid.).
ambigua, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 36 = P. Thouarsii.
simplex, Thou. Orch. Isles Afr. t. 24 = P. Thouarsii.
truncata, Lindl. Gen. et Sp. Orch. 411.—Ins. Mascar.?

APLOSTEMON, Rafin. in Journ. Phys. lxxxix. (1819) 105 = **Scirpus**, Linn. etc. (Cyper.).
atropurpureum, Rafin. l. c. = Eleocharis atropurpurea.
bracteatum, Rafin. l. c. = S. bracteatus.
compressum, Rafin. l. c. = S. Caricis.
polytrichoides, Rafin. l. c. = Fimbristylis polytrichoides.
triquetrum, Rafin. l. c. = Fimbristylis argentea.

APLOSTYLIS, Rafin. Fl. Tellur. iv. 91 (1836) = **Cuscuta**, Linn. (Convolv.).
lupuliformis, Rafin. l. c. = C. lupuliformis.

APLOTAXIS, DC. in Guill. Archiv. Bot. ii. (1833) 330 = **Saussurea**, DC. (Compos.).
albescens, DC. Prod. vi. 540 = S. albescens.
andryaloides, DC. l. c. 542 = S. sorocephala.
auriculata, DC. l. c. 541 = S. hypoleuca.
australasica, F. Muell. Fragm. i. 36 = S. affinis.
Bungei, DC. Prod. vi. 539 = S. affinis.
Bungei, Less. ex Ledeb. Fl. Ross. ii. 753 = Leuzea salina.
caespitosa, DC. Prod. vi. 539 = S. taraxacifolia.
candicans, DC. l. c. 540 = S. candicans.
Candolleana, DC. l. c. 541 = S. Candolleana.
carthamoides, Ham. in DC. l. c. 540 = S. affinis.
cirsoides, DC. l. c. = Cnicus arvensis.
deltoidea, DC. l. c. 541 = S. deltoidea.
denticulata, DC. l. c. 539 = S. denticulata.
discolor, DC. l. c. 541 = S. piptathera.
fastuosa, Decne. in Jacquem. Voy. Bot. 97. t. 105 = S. denticulata.
foliosa, Edgew. in Trans. Linn. Soc. xx. (1846) 77 = S. foliosa.
Frolovii, DC. Prod. vi. 538 = S. Frolovii.
gnaphalodes, Royle, Illustr. Bot. Himal. 251. t. 59. f. 1 = S. sorocephala.
gossypina, DC. Prod. vi. 541 = S. gossypiphora.
involucrata, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 389.—As. temp.
Jacea, Klotzsch, in Bot. Ergeb. Waldem. Reise, 79. t. 79 = S. Jacea.
Lappa, Decne. in Jacquem. Voy. Bot. 96. t. 104 = S. Lappa.
leontodontoides, DC. Coll. Mém. ix. 29. t. 10 (Prod. vi. 539) = S. Kunthiana.
multicaulis, DC. in Deless. Ic. Sel. iv. t. 68 = S. affinis.
nepalensis, DC. l. c. t. 67 = S. Eriostemon.
nivea, DC. Prod. vi. 541 = S. deltoidea.
obvallata, DC. in Deless. Ic. Sel. iv. t. 69 = S. obvallata.
pungens, DC. Prod. vi. 539 = Cnicus arvensis.
Roylei, DC. Prod. vi. 538 = S. Roylei.
scaposa, Edgew. in Trans. Linn. Soc. xx. (1846) 76 = S. candicans.
Simpsoniana, Field. & Gardn. Sert. Pl. t. 26 = S. andryaloides.
sorocephala, Schrenk, Enum. Pl. Nov. (1841) 43 = S. sorocephala.
taraxacifolia, DC. Prod. vi. 532 = S. taraxacifolia.
uniflora, DC. l. c. 539 = S. uniflora.

APLOTHECA, Mart. ex Cham. in Linnaea, v. (1830) 91 = **Froelichia**, Moench. (Amarant.).
interrupta, Mart. ex Cham. l. c. = F. interrupta.

APLUDA, Linn. Sp. Pl. ed. I. 82 (1753). *GRAMINEAE*, Benth. & Hook. f. iii. 1137.
CALAMINA, Beauv. Agrost. 128. t. 23. f. 1 (1812).
DIECTOMIS, Beauv. l. c. 132. t. 23. f. 5 (1812).
aristata, Linn. Cent. ii. 7; Sp. Pl. ed. II. 1487 = mutica.
ciliata, *Anderss. in Oefver. Vet. Acad. Stockh.* (1855) 177.—Ind. or.
communis, Nees & Arn. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 194 = mutica.
Cumingii, Buese, ex De Vriese, Pl. Ind. Bat. Reinw. 105 = mutica.
digitata, Linn. f. Suppl. 434 = Polytoca bracteata.
distachya, Beauv. Agrost. 133, 151 = Andropogon distachyus.
geniculata, Roxb. Hort. Beng. 8; Fl. Ind. i. 327 = mutica.
gigantea, Spreng. Syst. i. 290 = mutica.
glauca, Schreb. Graes. ii. 99 = mutica.
Gryllus, Beauv. Agrost. 133 = mutica.
Gryllus, Presl, Cyp. & Gram. Sic. 55 = Chrysopogon Gryllus.
humilis, Kunth, Enum. Pl. i. 517 = mutica.
imberbis, Steud. Nom. ed. II. i. 251, sphalm. = Anthistiria imberbis.
inermis, Regel, in Act. Hort. Petrop. vii. (1880) 658 = mutica.
Kobila, Buch.-Ham. ex Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 194 = mutica.
microstachya, Nees, l. c. 193 = mutica.
mucronata, Steud. Syn. Pl. Gram. 404 = mutica.
mutica, *Linn. Sp. Pl.* 82.—As. trop.; Polynes.; Austral.
pedicellata, Buese, ex De Vriese, Pl. Ind. Bat. Reinw. 105 = mutica.
rostrata, Arn. & Nees in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 194 = mutica.
scabra, Anderss. in Oefver. Vet. Acad. Stockh. (1855) 179 = mutica.
scirpoides, Walt. Fl. Carol. 250 (pl. dubia).—Carolina.
villosa, Schreb. Graes. ii. 92 = mutica.
Zeugites, Linn. Syst. ed. X. 1306; Pl. Jam. Pugill. 30 = Zeugites americana.
Zeygites, Aubl. Pl. Gui. ii. 933 = Ichnanthus pallens.

APOBALLIS, Schott, in Oestr. Bot. Zeitschr. (1858) 318 = **Schismatoglottis**, Zoll. (Aroid.).
neglecta, Schott, l. c. 317 = S. rupestris.

APOCHORIS, Duby, in DC. Prod. viii. 67 (1844). *PRIMULACEAE*, Benth. & Hook. f. ii. 635.
pentapetala, *Duby*, *l. c.*—China.

APOCOPIS, Nees, in Proc. Linn. Soc. i. (1841) 93. *GRAMINEAE*, Benth. & Hook. f. iii. 1128.
AMBLYACHYRUM, Hochst. ex Steud. Syn. Pl. Gram. 413 (1885).
Beckettii, Thw. ex Trimen, Cat. Pl. Ceylon, 107 = Wightii.
Royleana, *Nees, in Proc. Linn. Soc.* i. (1841) 94.—Ind. or.
tridentata, *Benth. in Journ. Linn. Soc.* xix. (1882) 67.—Ind. or.
Wightii, *Nees, ex Steud. Syn. Pl. Gram.* 377.—Ind. or.; China.
Wrightii, Munro, in Proc. Am. Acad. iv. (1864–65) 363 = Wightii.

APOCOPSIS, Meissn. Gen. 429 (328) (1843) = **Apocopis**, Necs (Gram.).

APOCYNUM, [Tourn.] Linn. Syst. ed. I. (1735); Gen. cd. I. 65 (1737). *APOCYNACEAE*, Benth. & Hook. f. ii. 716.
QUIRIVELIA, Poir. Encyc. vi. 42 (1804).
Acouci, Aubl. Pl. Gui. i. 274 = Forsteronia Acouci.
africanum, *Lour. Fl. Cochinch.* i. 168 (Quid ?).—Zanzib.
agglomeratum, *Poir. Encyc. Suppl.* i. 407.—Ins. S. Doming.
alterniflorum, Lour. Fl. Cochinch. i. 168 = Strophanthus alterniflorus.
androsaemifolium, *Linn. Sp. Pl.* 213.—Am. bor.
bursiflorum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 5.—Malaya.
canariense, *Lam. Encyc.* i. 212.—Ins. Canar.
cannabinum, *Linn. Sp. Pl.* 213.—Am. bor.
citrifolium, Descourt. Fl. Med. Antill. iii. 180. t. 191 = Echites citrina.
compressum, Moench, Meth. 464 = venetum.
APOCYNUM:—
cordatum, Mill. Gard. Dict. ed. VIII. n. 10 = Rhabdadenia cordata.
cordatum, Thunb. Prod. Pl. Cap. 47 = Astephanus cordatus.
cordifolium, Tenore, ex Steud. Nom. ed. II. i. 113 = praec.?
cotinifolium, *Lam. Encyc.* i. 215.—Java.
crassifolium, Salisb. Prod. 149 = Ichnocarpus frutescens.
cumanense, Willd. ex Roem. & Schult. Syst. vi. 796 = Metaplexis fimbriatum.
erectum, Vell. Fl. Flum. 123; iii. t. 87 = Barjonia racemosa.
filiforme, Thunb. Prod. Pl. Cap. 47 = Eustegia filiformis.
floristratum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 5 = Strophanthus dichotomus.
foetidum, Burm. f. Fl. Ind. 71 = Paederia foetida.
frutescens, Afzel. Remcd. Guin. Coll. iv. 28 = Ichnocarpus Afzelii.
frutescens, Linn. Sp. Pl. 213 = Ichnocarpus frutescens.
hastatum, Thunb. Prod. Pl. Cap. 47 = Eustegia hastata.
Hendersonii, *Hook. f. in Hume & Henders. Lahore to Yarkand*, 327.—Turkest.
hypericifolium, Ait. Hort. Kew. ed. I. i. 304 = cannabinum.
indicum, Lam. Encyc. i. 214 = reticulatum.
Juventas, Lour. Fl. Cochinch. 167 = Tylophora ovata.
lanceolatum, Thunb. Prod. Pl. Cap. 47 = Astephanus lanceolatus.
lineare, Linn. f. Suppl. 169 = Astephanus linearis.
maculatum, Descourt. Fl. Med. Antill. iii. 177. t. 190 = Echites maculata.
minutum, Linn. f. Suppl. i. 169 = Eustegia hastata.
mucronatum, Blanco, Fl. Filip. cd. I. 852 = Toxocarpus gracilis.
muscipulum, Moench, Meth. 464 = androsaemifolium.
nervosum, *Mill. Gard. Dict.* ed. VIII. n. 9.—Vera Crux.
obliquum, Mill. l. c. n. 8 = Echites umbellata.
odoratissimum, *Lour. ex Pritz. Ic. Ind.* 83.—Malaya.
orixense, Rottler, ex Hook. f. Fl. Brit. Ind. iv. 6 = Cryptolepis elegans.
paniculatum, Lam. Encyc. i. 214 = Forsteronia Acouci.
pictum, *Schrenk, in Bull. Phys. Math. Acad. Pétersb.* ii. (1844) 115.—Soongaria.
piscatorium, Dougl. ex A. DC. in DC. Prod. viii. 439 = cannabinum.
platyanthum, Salzm. ex A. DC. l. c. 594 = Philibertia cynanchoides.
pubescens, R. Br. in Mem. Wern. Soc. i. (1809) 68 = cannabinum.
purpureum, *Tausch, in Flora*, xix. (1836) II. 385.—Am. bor.
rectinerve, Salzm. ex Decne. in DC. Prod. viii. 603 = Blepharodon pallidum.
reticulatum, Herb. Madras, ex Wall. Cat. n. 8247 = Decalepis Hamiltonii.
reticulatum, *Linn. Sp. Pl.* 214.—Ind. or.
reticulatum, Lour. Fl. Cochinch. 208 = Parsonsia Helicandra.
salicifolium, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 406 = Gomphocarpus fruticosus.
salicifolium, *Willd. ex Roem. & Schult. Syst.* iv. 796.—Ins. S. Doming.
scandens, Mill. Gard. Dict. ed. VIII. n. 5 = Echites asperuginis.
sibiricum, Jacq. Hort. Vindob. iii. 37 = cannabinum.
sibiricum, Pall. ex Roem. & Schult. Syst. iv. 405 = venetum.
soleare, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 5.—Malaya.
speciosissimum, Mill. Gard. Dict. ed. VIII. n. 4 (= *Haemadictyon suberectum*).
stellatum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 5.—Malaya.
stylospermum, *Noronha, l. c.*—Malaya.
syriacum, S. G. Gmel. Reise, ii. 198, 257 = Calotropis procera.
tiliaefolium, Lam. Encyc. i. 214 = Dregea volubilis.
triflorum, Linn. f. Suppl. 169 = Astephanus triflorus.
umbellatum, Aubl. Pl. Gui. i. 275 = Thenardia umbellata.
umbellatum, Salzm. ex Decne. in DC. Prod. viii. 585 = Oxypetalum densiflorum.
venetum, A. DC. Prod. viii. 440 = cannabinum.
venetum, *Linn. Sp. Pl.* 213.—Reg. Mediterr.; Oriens; As. bor.; Ind. or.
APOCYNUM:—
villosum, *Mill. Gard. Dict.* ed. VIII. n. 11.—Vera Crux.
viminale, Bassi, in Comm. Bonon. (1753) ex Roem. & Schult. Syst. vi. 113 = Sarcostemma viminale.
vincaefolium, *Burm. f. Fl. Ind.* 71.—Ind. or.
volubile, Vell. Fl. Flum. 123; iii. t. 88 = Araujia calycina.

APODANTHERA, Arn. in Hook. Journ. Bot. iii. 274 (1841). *CUCURBITACEAE*, Benth. & Hook. f. i. 834.
argentea, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 42.—Bras.
aspera, *Cogn. l. c.* 43.—Mexic.
biflora, *Cogn. l. c.*—Ecuador.
Buraeavi, *Cogn. l. c.* 44.—Mexic.
Galeottii, *Cogn. l. c.* 45.—Mexic.
gracilis, *Benth. Bot. Voy. Sulph.* 99 = Melothria fluminensis.
laciniosa, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 12.—Bras.
Mandonii, *Cogn. l. c.* 41.—Bolivia.
Mathewsi, *Arn. in Hook. Journ. Bot.* iii. (1841) 274.—Peruv.
mucronata, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 40.—Peruv.
pedisecta, *Cogn. l. c.* 17.—Bras.
scabra, *Cogn. in Bull. Acad. Belg.* Sér. II. xlix. (1880) 192.—Ecuador.
smilacifolia, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) 42.—Bras.
undulata, *A. Gray, Pl. Wright.* ii. 60.—N. Mexic.

APODANTHES, Poit. in Ann. Sc. Nat. Sér. I. iii. (1824) 422. t. 26. *CYTINACEAE*, Benth. & Hook. f. iii. 118.
FROSTIA, Bertero, ex Guill. in Ann. Sc. Nat. Sér. II. ii. (1834) 19, in nota.
PILOSTYLES, Guill. in Ann. Sc. Nat. Sér. II. ii. (1834) 21. t. 1.
SARNA, Karst. in Nov. Act. Nat. Cur. xxvi. (1858) 920. t. 65.
Berterii, *Gardn. in Hook. Ic. Pl.* t. 655.—Chili.
Blanchetii, *Gardn. l. c.* t. 655 b.—Bras.
Calliandrae, *Gardn. l. c.* t. 644.—Bras.
Cascariae, *Poit. in Ann. Sc. Nat.* Sér. I. iii. (1824) 422. t. 26.—Am. austr.
Flacourtiae, *Karst. in Linnaea*, xxviii. (1856) 413.—Venezuela.

APODOCEPHALA, Baker, in Journ. Linn. Soc. xxi. (1885) 417. *COMPOSITAE*.
pauciflora, *Baker, l. c.*—Madag.

APODOLIRION, Baker, in Journ. Bot. xvi. (1878) 74. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 722.
Bolusii, *Baker, l. c.* 75.—Afr. austr.
Buchanani, *Baker, l. c.*—Natal.
lanceolatum, *Benth. & Hook. f. Gen.* iii. 722.—Afr. austr.
Mackenii, *Baker, in Journ. Bot.* xvi. (1878) 75.—Natal.

APODOSTACHYS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 557 = **Ercilla**, A. Juss. (Phytolac.).
densiflora, Turcz. l. c. = E. volubilis.

APODYNOMENE, E. Mey. Comm. Pl. Afr. Austr. 111 (1835) = **Tephrosia**, Pers. (Legum.).
aemula, E. Mey. l. c. 113 = T. acmula.
diffusa, E. Mey. l. c. = T. diffusa.
grandiflora, E. Mey. l. c. 111 = T. grandiflora.
macropoda, E. Mey. l. c. 112 = T. macropoda.
Meyeri, Presl, Bot. Bemerk. 57 = T. grandiflora.

APODYTES, E. Mey. ex Bernh. in Linnaea, xii. (1838) 136; et ex Arn. in Hook. Journ. Bot. iii. (1841) 155. *OLACINEAE*, Benth. & Hook. f. i. 351.
NOTHAPODYTES, Blume, Mus. Bot. Lugd. Bat. i. 248 (1850).
RHAPHIOSTYLIS, Planch. ex Benth. in Hook. Niger Fl. 259 (1849).
acutifolia, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 92 = dimidiata.
andamanica, *Kurz, Andam. Rep.* 33.—Ins. Andam.
Beddomei, *Mast. in Hook. f. Fl. Brit. Ind.* i. 588.—Ind. or.
beninensis, *Hook. f. in Hook. Ic. Pl.* t. 778.—Afr. trop.

APODYTES :—
Benthamiana, *Wight, Ic.* t. 1153.—Ind. or.
brachystylis, *F. Muell. Fragm.* ix. 149.—Austral.
dimidiata, *E. Mey. ex Bernh. in Linnaea*, xii. (1838) 136.—Afr. austr.
Gardneriana, Miers, in Ann. & Mag. Nat. Hist. Ser. II. ix. (1852) 389 = Benthamiana.
grandifolia, *Benth. & Hook. f. Gen.* i. 352.—Madag.
mauritiana, *Benth. & Hook. f. l. c.*—Ins. Mascar.

APOGANDRUM, Neck. Elem. i. 216 (1790) = **Erica**, Linn.

APOGETON, Schrad. ex Steud. Nom. ed. II. i. 114 (1840) = **Aponogeton**, Linn. f. (Naiad.).

APOGON, Ell. Sketch, ii. 267 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 509.
LAXANON, Rafin. New Fl. Am. ii. 16 (1836).
gracilis, *DC. Prod.* vii. 79.—Am. bor.
humilis, *Ell. Sketch*, ii. 267.—Am. bor.
humilis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 424 = gracilis.
lyratum, Nutt. in Journ. Acad. Philad. vii. (1834) 71 = humilis.
Wrightii, *A. Gray, Syn. Fl. N. Am.* i. II. 411.—Texas.

APOGONIA, Fourn. Gram. Mex. 63 ex Benth. & Hook. f. Gen. iii. 1130 (1883) = **Rottboellia**, Linn. (Gramin.).
glabrata, Fourn. ex Hemsl. Biol. Centr. Am. Bot. iii. 521 = Andropogon imberbis.
ramosa, Fourn. ex Hemsl. l. c. = R. aurita.

APOLANESIA, Reichb. Nom. 154 (1841) = **Apoplanesia**, Presl (Legum.).

APOLEPSIS, Hassk. Cat. Hort. Bogor. Alt. 150 (1844) = **Lepidagathis**, Willd. (Acanth.).
dispar, Hassk. l. c. = L. repanda.
repanda, Nees, in DC. Prod. xi. 261 = L. repanda.

APOLLINARIS, Linn. Syst. ed. I. (1735) = **Nicotiana**, Linn. ? (Solan.).

APOLLONIAS, Nees, Laurin. Expos. 10 (1833). *LAURINEAE*, Benth. & Hook. f. iii. 152.
Arnottii, *Nees, Syst. Laurin.* 670.—Ind. or.
barbusana, A. Br. in Verh. Befoerd. Gartenb. Preuss. xxi. (1852) 10 = canariensis.
canariensis, *Nees, Syst. Laurin.* 696.—Ind. or.
zeylanica, Thw. Enum. Pl. Zeyl. 253 = Beilschmiedia oppositifolia.

APOMAEA, Neck. Elem. ii. 24 (1790) = **Ipomoea**, Linn. (Convolv.).

APOMOEA, Steud. Nom. ed. II. i. 114 (1840) = praec.

APONOA, Rafin. Sylva Tellur. 84 (1838) = **Columnea**, Linn. (Gesner.).
repens, Rafin. l. c. = C. scandens ?

APONOGETON, Linn. f. Suppl. 32 (1781). *NAIADACEAE*, Benth. & Hook. f. iii. 1013.
AMOGETON, Neck. Elem. iii. 267 (1790).
APOGETON, Schrad. ex Steud. Nom. ed. II. i. 114 (1840).
HYDROGETON, Pers. Syn. i. 400 (1805).
LIMNOGETON, Edgew. in Lindl. Veg. Kingd. 210 (1847).
OUVIRANDRA, Thou. in Roem. Collect. 196 (1809); Gen. Nov. Madag. 2 (1810).
SPATHIUM, Edgew. in Journ. As. Soc. Beng. (1842) 145.
abyssinicum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 351.—Abyss.
angustifolium, *Ait. Hort. Kew.* i. 495.—Afr. austr.
Bernerianum, *Hook. f. in Benth. & Hook. f. Gen.* iii. 1014.—Madag.
crinifolium, *Lehm. ex Steud. Nom.* ed. II. i. 114.—Afr. austr.
crispum, *Thunb. Nov. Gen.* 73.—Zeylan.
desertorum, *Zeyh. ex Steud. Nom.* ed. II. i. 114.—Afr. austr.
distachys, Edgew. in Hook. Lond. Journ. Bot. iii. (1844) 402 = distachyum.
distachyum, *Thunb. Nov. Gen.* 74.—Afr. austr.
echinatum, *Roxb. Hort. Beng.* [87]; *et Fl. Ind. ed. Carey*, ii. 210.—Ind. or.
elongatum, *F. Muell. ex Benth. Fl. Austral.* vii. 188.—Austral.
fenestrale, *Hook. f. in Benth. & Hook. f. Gen.* iii. 1014.—Madag.
Holubii, *Oliver, in Hook. Ic. Pl.* t. 1470.—Afr. trop.
junceum, *Lehm. ex Steud. Nom.* ed. II. i. 114.—Afr. austr.
leptostachyum, *E. Mey. ex Baker, in Trans. Linn. Soc.* xxix. (1875) 158.—Afr. austr.
lineare, Vahl, Symb. Bot. iii. 51 = monostachyon.
microphyllum, *Roxb. Hort. Beng.* 26; *Fl. Ind. ed. Carey*, ii. 211.—Ind. or.
monostachys, Edgew. in Hook. Lond. Journ. Bot. iii. (1844) 402 = scq.
monostachyon, *Linn. f. Suppl.* 214.—Ind. or.; China.
natalense, *Oliver, in Hook. Ic. Pl.* t. 1471.—Afr. austr.
Rehmanni, *Oliver, l. c.* 1471 B.—Afr. trop.
spathaceum, *E. Mey. in Drège, Zwei Pfl. Docum.* 165, *nomen*.—Afr. austr.
quadrangulare, *Baker, in Journ. Linn. Soc.* xviii. (1881) 279.—Madag.
ulvaceum, *Baker, l. c.*—Madag.
undulatum, Roxb. Hort. Beng. 26; Fl. Ind. ed. Carey, ii. 211 = crispum.
vallisnerioides, *Baker, in Trans. Linn. Soc.* xxix. (1875) 158.—Afr. trop.
zosteraefolium, *Schrad. ex Steud. Nom.* ed. II. i. 114 (an lapsu = Potamogeton zosteraefolius ? Schum.).—Am. bor.

APOPHRAGMA, Griseb. Gen. et Sp. Gent. 163 (1839) = **Curtia**, Cham. & Schlecht. (Gentian.).
tenuifolium, Griseb. l. c.—Guiana.

APOPHYLLUM, F. Muell. in Hook. Kew Journ. ix. (1857) 306. *CAPPARIDEAE*, Benth. & Hook. f. i. 109.
anomalum, *F. Muell. l. c.* 307.—Austral.

APOPLANESIA, Presl, Symb. Bot. i. 63. t. 41 (1831). *LEGUMINOSAE*, Benth. & Hook. f. i. 492.
paniculata, *Presl, l. c.*—Mexic.

APOPLEUMON, Rafin. Fl. Tellur. iv. 72 (1836) = **Ipomoea**, Linn. (Convolv.).
bignonioides, Rafin. l. c. = I. bignonioides.

APORANTHUS, Bromf. Fl. Vect. 117 (1856) = **Trigonella**, Linn. (Legum.).
Trifoliastrum, Bromf. l. c. = T. ornithopodioides.

APORETIA, Walp. Ann. ii. 217 (1851-52) = seq.

APORETICA, Forst. Char. Gen. 131. t. 66 (1776) = **Schmidelia**, Linn. (Sapind.).
gemella, Blanco, Fl. Filip. ed. II. 203 = S. Cobbe.
gemella, DC. Prod. i. 610 = S. cochinchinensis.
penicillata, Blanco, Fl. Filip. ed. I. 291 = S. Cobbe.
pinnata, Forst. f. Prod. n. 392 = S. Cobbe.
pinnata, Hook. & Arn. Bot. Beech. Voy. 61 = Guioa lentiscifolia.
ternata, Forst. Char. 132 = S. Cobbe.

APOROCACTUS, Lem. Illustr. Hortic. vii. (1860) Misc. 67 = **Cereus**, Linn. (Cact.).
Baumanni, Lem. l. c. = C. Baumanni.
colubrinus, Lem. l. c. = C. colubrinus.
flagelliformis, Lem. l. c. = C. flagelliformis.
flagriformis, Lem. Cact. 58 = C. flagriformis.

APOROSA, Blume, Bijdr. 511 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 282.
LEIOCARPUS, Blume, Bijdr. 581 (1825).
LEPIDOSTACHYS, Wall. Cat. n. 6816 (1832).
SCEPA, Lindl. Introd. Nat. Syst. ed. II. 441 (1836).
TETRACTINOSTIGMA, Hassk. Hort. Bogor. Descr. 55 (1855).
acuminata, *Thw. Enum. Pl. Zeyl.* 288.—Ind. or.
affinis, Baill. Etud. Gén. Euph. 645 = Lindleyana.
arborea, *Muell. Arg. in DC. Prod.* xv. II. 470.—Java.
arborescens, *Muell. Arg. l. c.*—Java.
aurita, *Baill. Etud. Gén. Euph.* 645.—Ins. Philipp.
aurita, Miq. Fl. Ind. Bat. i. II. 431 = microcalyx.
australiana, *F. Muell. Census*, 20.—Austral.
Cumingiana, Baill. Etud. Gén. Euph. 645 = microcalyx.
dioica, Muell. Arg. in DC. Prod. xv. II. 472 = Roxburghii.
ficifolia, *Baill. Adansonia*, xi. (1873-76) 177.—Malaya; Cochinch.
frutescens, *Blume, Bijdr.* 514.—Java.
frutescens, Benth. Fl. Hongk. 317 = microcalyx.
fruticosa, *Muell. Arg. in DC. Prod.* xv. II. 475.—Java.
fusiformis, *Thw. Enum. Pl. Zeyl.* 288.—Zeylan.
glabrifolia, *Kurz, in Journ. Bot.* xiii. (1875) 330.—Ins. Nicobar.
lanceolata, Kurz, For. Fl. Burma, ii. 363 = Planchoniana.
lanceolata, *Thw. Enum. Pl. Zeyl.* 288.—Zeylan.
latifolia, *Thw. l. c.*—Zeylan.
leptostachya, Benth. Fl. Hongk. 317 = microcalyx.
Lindleyana, *Baill. Etud. Gén. Euph.* 645 (Lyndleyana).—Ind. or.
lunatum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 239.—Malaya.
macrophylla, *Muell. Arg. in DC. Prod.* xv. II. 470.—Burma.
microcalyx, *Hassk. in Bull. Soc. Bot. France*, vi. (1859) 714.—China; Austral.; Malaya.
microstachya, *Muell. Arg. in DC. Prod.* xv. II. 474.—Burma.
Miquelliana, *Muell. Arg. l. c.*—Sumatra.
oblonga, *Muell. Arg. in Linnaea*, xxxii. (1863) 78.—Burma.
Planchoniana, *Baill. Etud. Gén. Euph.* 645.—Burma.
quadrilocularis, *Muell. Arg. in DC. Prod.* xv. II. 476.—Sumatra.
Roxburghii, *Baill. Etud. Gén. Euph.* 645.—Ind. or.
sphaerocarpa, Muell. Arg. in Flora, xlvii. (1864) 519 = Lindleyana.
tetrapleura, *Hance, in Journ. Bot.* xiv. (1876) 260.—Cambodia.
Thwaitesi, Baill. Etud. Gén. Euph. 645 = fusiformis.
Tulasneana, Baill. l. c. = microstachya.
villosa, *Baill. l. c.*—Burma; Cochinch.
villosula, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 239.—Burma.

APORRHIZA, Radlk. in Sitzb. Math. Phys. Acad. Muench. viii. (1878) 338. *SAPINDACEAE*.
paniculata, *Radlk. l. c.* 339.—Afr. trop.

APORUM, Blume, Bijdr. 334. t. 39 (1825) = **Dendrobium**, Sw. (Orchid.).
acinaciforme, Griff. in Calc. Journ. Nat. Hist. v. (1845) 370 = D. acinaciforme.
anceps, Lindl. in Wall. Cat. n. 2020 = D. Nathanielis.
anceps, Lindl. Gen. et Sp. Orch. 71 = D. anceps.
concinnum, Lindl. ex Reichb. f. in Trans. Linn. Soc. xxx. (1874) 149 = D. atropurpureum.
cuspidatum, Wall. ex Lindl. in Bot. Reg. (1841) Misc. 2 = D. Nathanielis.
incrassatum, Blume, Bijdr. 334 = D. incrassatum.
incrassatum, Brongn. ex Lindl. in Journ. Linn. Soc. iii. (1859) 60 = Epidendrum aporoides ?
indivisum, Blume, Bijdr. 334. t. 39 = D. indivisum.
indivisum, Lindl. Gen. et Sp. Orch. 70 = D. Leonis.
Jenkinsii, Griff. in Calc. Journ. Nat. Hist. v. (1845) 367 = D. parciflorum.
Leonis, Lindl. in Bot. Reg. (1840) Misc. 59 = D. Leonis.
lobatum, Blume, Bijdr. 334 = D. lobatum.
micranthum, Griff. in Calc. Journ. Nat. Hist. iv. (1844) 375. t. 16. f. 7 = D. Serra.
Roxburghii, Griff. l. c. v. (1845) 370 = D. Roxburghii.
Serra, Lindl. in Wall. Cat. n. 2021 = D. Serra.
sinuatum, Lindl. in Bot. Reg. (1841) Misc. 1 = D. sinuatum.
subteres, Griff. in Calc. Journ. Nat. Hist. v. (1845) 370 = D. acerosum.

APOSERIS, Neck. Elem. i. 57 (1790) = **Hyoseris**, Linn. (Compos.).
foetida, Cass. in Dict. Sc. Nat. xlviii. 427 = H. lucida.
foetida, Less. Syn. Comp. 128 = H. foetida.
lucida, Less. l. c. = H. lucida.

APOSTASIA, Blume, Bijdr. 428. t. 5 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 635.
MESODACTYLIS, Wall. Pl. As. Rar. i. 74, in obs. (1830).

APOSTASIA :—
NEUMAYERA, Rcichb. f. in Flora, lv. (1872) 278.
NIEMEYERA, F. Muell. Fragm. vi. 96 (1867).
Brunonis, Griff. Notul. iii. 243 = nuda.
Lobbii, *Reichb. f. in Flora*, lv. (1872) 278.—Borneo.
nuda, *R. Br. in Wall. Pl. As. Rar.* i. 76. t. 85.—Ind. or.; Malaya.
odorata, *Blume, Bijdr.* 423.—Java.
stylidioides, *Reichb. f. in Flora*, lv. (1872) 278.—Austral.
Wallichii, *R. Br. in Wall. Pl. As. Rar.* i. 75. t. 84.—Ind. or.; Malaya.

APOTERIUM, Blume, Bijdr. 218 (1825) = **Calophyllum**, Linn. (Guttif.).
Sulatri, Blume, l. c. = C. spectabile.

APOXYANTHERA, Hochst. in Flora, xxvi. (1843) 78 = **Raphionacme**, Harv. (Asclep.).
pubescens, Hochst. l. c. = R. pubescens.

APOZIA, Willd. ex Steud. Nom. ed. II. i. 114 (1840) = **Micromeria**, Benth. (Labiat.).
chamaedryoides, Willd. ex Steud. l. c. = M. Brownei.
herniarioides, Willd. ex Steud. l. c. = Herpestis herniarioides.
serpyllacea, Willd. ex Steud. l. c. = M. Brownei.

APPELLA, Adans. Fam. ii. 84 (1763) = **Laurus**, Linn.

APPENDICULA, Blume, Bijdr. 297. t. 40 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 588.
CONCHOCHILUS, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 146.
METACHILUM, Lindl. Gen. et Sp. Orch. 74 (1830).
SCOLIOCHILUS, Reichb. f. Xenia Orch. ii. 118, in obs. (1872).
alba, *Blume, Bijdr.* 299.—Java.
anceps, *Blume, l. c.*—Java.
angustifolia, *Blume, l. c.* 301.—Java.
bifaria, *Lindl. ex Benth. in Hook. Kew Journ.* vii. (1855) 35.—Ind. or.; China.
bracteosa, *Reichb. f. in Seem. Fl. Vit.* 299.—Ins. Fiji.
buxifolia, *Blume, Bijdr.* 300.—Java.
callosa, *Blume, l. c.* 303.—Burma; Malaya.
carnosa, *Blume, l. c.* 300.—Java.
Chalmersiana, *F. Muell. in Wing, South. Sc. Record*, N. S. (1. May 1885).—N. Guin.
congenera, *Blume, Bijdr.* 303.—Java.
cornuta, *Blume, l. c.* 302.—Java.
cristata, *Blume, l. c.* 298.—Java.
elegans, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.
graminifolia, *Teijsm. & Binn. in Tijdschr. Neder. Ind.* xxiv. (1862) 322.—Java.
Hasseltii, *Blume, Bijdr.* 304.—Burma.
Hasseltii, Wight, Ic. t. 1748 = Agrostophyllum khasyanum.
Kuhlii, *Reichb. f. Xenia Orch.* ii. 116.—Java.
Lewisii, *Griff. in Calc. Journ. Nat. Hist.* iv. (1844) 378.—Malaya.
longifolia, *Blume, Bijdr.* 304.—Java.
membranacea, *Teijsm. & Binn. in Kruid. Arch.* iii. (1855) 399.—Java.
micrantha, Lindl. in Ann. Nat. Hist. xv. (1845) 386.—Ins. Philipp.
monoceras, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.
muricata, *Teijsm. & Binn. in Tijdschr. Neder. Ind.* xxiv. (1862) 322.—Java.
pauciflora, *Blume, Bijdr.* 300.—Java.
pendula, *Blume, l. c.* 298.—Java.
penicillata, *Blume, Rumphia*, iv. 46.—N. Guin.
peruligera, *Reichb. f. in Bonplandia*, v. (1857) 41.—Java.
purpurascens, *Blume, Bijdr.* 302.—Java.
ramosa, *Blume, l. c.* 299.—Java.
reduplicata, Par. & Reichb. f. Otia Bot. Hamb. 45 = bifaria.
reflexa, *Blume, Bijdr.* 301.—Java.
rhodiola, *Reichb. f. Xenia Orchid.* ii. 118.—Java.
sororia, *Miq. Fl. Ind. Bat.* iii. 703.—Java.
stipulata, Griff. Notul. iii. 358 = callosa.
teres, Griff. l. c. 359 = Ceratostylis teres.
tomentella, *Zoll. in Nat. en Geneesk. Arch. Neerl. Indie*, i. (1844) 403.—Java.
torta, *Blume, Bijdr.* 303.—Malaya.

APPENDICULA :—
tricornis, *Reichb. f. in Bonplandia*, v. (1857) 41.—Hab.?
undulata, *Blume, Bijdr.* 301.—Java.
Vieillardi, *Reichb. f. in Linnaea*, xli. (1877) 76.—N. Caled.
viridiflora, *Teijsm. & Binn. in Tijdschr. Neder. Ind.* xxiv. (1862) 321.—Java.
xytriophora, *Reichb. f. in Seem. Fl. Vit.* 299.—Malaya.

APPENDICULARIA, DC. Prod. iii. 114 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 741.
thymifolia, *DC. l. c.*—Guiana.

APPUNIA, Hook. f. in Benth. & Hook. f. Gen. ii. 120 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 120.
BELLYNKIA, Muell. Arg. in Flora, xlviii. (1875) 465.
angulata, *Baill. Hist. Pl.* vii. 415.—Bras.
tenuiflora, *Benth. & Hook. f. Gen.* ii. 120.—Guiana.

APRADUS, Adans. Fam. ii. 102 (1763) = **Arctopus**, Linn. (Umbell.).

APRELLA, Steud. Nom. ed. II. i. 152 (1840), err. typ. = ASPRELLA, Schrcb. = **Leersia**, Sw. (Gram.).

APREVALIA, Baill. in Bull. Soc. Linn. Par. i. (1884) 428. *LEGUMINOSAE.*
floribunda, *Baill. l. c.*—Madag.

APTANDRA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 201. *OLACINEAE*, Benth. & Hook. f. i. 345.
Benthamiana, Miers, l. c. Ser. III. iv. (1859) 360 = liriosmoides.
liriosmoides, *Spruce, ex Miers, l. c.*—Bras.
Spruceana, *Miers, l. c.* Ser. II. viii. (1851) 202.—Bras.
tubicina, *Benth. ex Miers, l. c.* Ser. III. iv. (1859) 360.—Bras.

APTERIA, Nutt. in Journ. Acad. Sc. Philad. vii. (1834) 64. t. 9. *BURMANNIACEAE*, Benth. & Hook. f. iii. 458.
NEMITIS, Rafin. Fl. Tellur. iv. 33 (1836).
STEMOPTERA, Miers, in Proc. Linn. Soc. i. (1840) 62.
hymenanthera, Miq. in Verh. Nat. Wet. Haarl. vii. (1851) 216. t. 64 = setacea.
lilacina, Miers, in Trans. Linn. Soc. xviii. (1841) 546. t. 37 = setacea.
orobanchioides, Hook. Ic. Pl. t. 254 = Dictyostega orobanchioides.
setacea, *Nutt. in Journ. Acad. Sc. Philad.* vii. (1834) 64. t. 9.—Am. bor.

APTERANTHES, Mik. in Nov. Act. Nat. Cur. xvii. (1835) 593. t. 41 = **Boucerosia**, Wight & Arn. (Asclep.).
cylindrica, Decne. in Ann. Sc. Nat. Sér. V. xiii. (1870–1) 406.—Abyssin.
Gussoneana, Mik. in Nov. Act. Nat. Cur. xvii. (1835) 594. t. 41 = B. Gussoniana.
tessellata, Decne. in Ann. Sc. Nat. Sér. V. xiii. (1870–1) 406 = Echidnopsis cereiformis.

APTEROCARYON, Opiz, in Lotos, v. (1855) 258 = **Betula**, Linn. (Cupulif.).
Michauxii, Opiz, l. c. 259 = B. Michauxii.

APTERON, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 300. *RHAMNEAE.*
lanceolatum, *Kurz, l. c.* 301.—Tenasserim.

APTEUXIS, Griff. Notul. iv. 672 (1854) = **Pternandra**, Jack (Melast.).
trinervis, Griff. l. c. = P. caerulescens.

APTOSIMUM, Burch. Trav. i. 219 (1824). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 927.
CHILOSTIGMA, Hochst. in Flora, xxiv. (1841) 372.
OHLENDORFIA, Lehm. Ind. Sem. Hort. Hamb. (1835).
abietinum, *Burch. Trav.* i. 308.—Afr. austr.
depressum, *Burch. l. c.* i. 260.—Afr. austr.
eriocephalum, *E. Mey. ex Benth. in Bot. Reg.* sub t. 1882.—Afr. austr.
indivisum, *Burch. Trav.* i. 219.—Afr. austr.
procumbens, Burch. ex Steud. Nom. ed. II. i. 114 = depressum.

APTOSIMUM :—
pumilum, *Benth. in DC. Prod.* x. 345.—Afr. austr.
tragacanthoides, *E. Mey. ex Benth. in Bot. Reg.* sub t. 1882.—Afr. austr.
viscosum, *Benth. l. c.*—Afr. austr.

APTOTHECA, Miers, Apocyn. S. Am. 150 (1878). *APOCYNACEAE.*
corylifolia, *Miers. l. c.*—Cuba.

APULEIA, Gaertn. Fruct. ii. 439. t. 171 (1791) = **Berkheya**, Ehrh. (Compos.).
asteroides, Gaertn. l. c. 440 = B. fruticosa.
atractyloides, Less. Syn. Comp. 64 (= *Stobaea atractyloides*).
fruticosa, Bartl. Sem. Goett. (1835).—Cf. Linnaea, xii. (1838) Litt. 80 = B. fruticosa.
fruticosa, Gaertn. Fruct. ii. 440 = B. grandiflora.
heterophylla, Less. Syn. Comp. 65 (= *Stobaea heterophylla*).
rigida, Gaertn. Fruct. ii. 439. t. 171 (= *Stobaea Gaertneri*).
Zeyheri, Less. Syn. Comp. 64 = B. Zeyheri.

APULEIA, Mart. Herb. Fl. Bras. 123 (1837). *LEGUMINOSAE*, Benth. & Hook. f. i. 574.
ZENKERIA, Arn. in Mag. Zool. & Bot. ii. (1838) 548.
molaris, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 177.—Bras.
praecox, *Mart. Herb. Fl. Bras.* 123.—Bras.

AQUARTIA, Jacq. Enum. Pl. Carib. i. 12 (1760) = **Solanum**, Linn.
aculeata, Jacq. l. c. = S. Aquartia.
microphylla, Lam. Illustr. n. 1604. t. 82. f. 2 = S. microphyllum.
microphylla, Willd. Sp. Pl. i. 622.—Ins. S. Doming.
solanacea, Willd. ex Steud. Nom. ed. II. i. 114.—N. Granat.
tomentosa, Lam. Illustr. n. 1603. t. 82. f. 1 = S. Aquartia.

AQUIFOLIUM, Tourn. ex Hall. Enum. Stirp. Helv. ii. 467 (1742) = **Ilex**, Linn.
croceum, Rafin. Sylva Tellur. 44 = I. Aquifolium?
ferox, Mill. ex Rafin. l. c. = I. Aquifolium.
heterophyllum, Rafin. l. c. = I. Aquifolium.
Ilex, Scop. Fl. Carn. cd. II. i. 116 = I. Aquifolium.
japonicum, Rafin. Sylva Tellur. 44 = Berberis japonica.
lanceolatum, Rafin. l. c.—Hab.?
planifolium, Rafin. l. c.—Hab.?
spinosum, Lam. Fl. Fr. iii. 652 = I. Aquifolium.
undulatum, Rafin. Sylva Tellur. 44 = I. Aquifolium.
vulgare, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128 = I. Aquifolium.

AQUILARIA, Lam. Encyc. ii. 610 (1783). *THYMELAEACEAE*, Benth. & Hook. f. iii. 200.
AGALLOCHUM, Lam. Encyc. i. 47 (1783).
OPHIOSPERMUM, Rcichb. Consp. 82 (1828).
OPHISPERMUM, Lour. Fl. Cochinch. 280 (1790).
Agallocha, *Roxb. Hort. Beng.* 33; *Fl. Ind.* ii. 422.—Ind. or.; Burma.
bancana, Miq. Fl. Ind. Bat. Suppl. 355 = Gonostylus Miquelianus.
chinensis, Spreng. Syst. ii. 356 = A. Ophispermum.
grandiflora, *Benth. Fl. Hongk.* 297.—China.
macrophylla, Miq. Fl. Ind. Bat. Suppl. 356 = Gonostylus Miquelianus.
malaccensis, Benth. in Hook. Kew Journ. v. (1853) 195 = grandiflora.
malaccensis, *Lam. Encyc.* i. 49.—Malaya.
microcarpa, *Baill. Adansonia*, xi. (1873–76) 304.—Borneo.
Ophispermum, *Poir. in Dict. Sc. Nat.* xviii. 161.—China.
ovata, Cav. Diss. vii. 377 = malaccensis.
pentandra, *Blanco, Fl. Filip.* ed. I. 373.—Ins. Philipp.
secundaria, *DC. Prod.* ii. 59.—Ins. Molucc.

AQUILEGIA, [Tourn.] Linn. Syst. ed. I. (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 8.
advena, *Regel, in Ind. Sem. Hort. Petrop.* (1856) 32.—Hab.?
aggericola, Jord. Diagn. i. 87 = vulgaris.
alpina, Deless. Ic. t. 48 = jucunda.
alpina, Georgi, It. i. 220 = glandulosa.
alpina, Lam. Encyc. i. 150 = pyrenaica.

AQUILEGIA :—
alpina, *Linn. Sp. Pl.* 533.—Europ. ; Sibir.
Amaliae, *Heldr. in Boiss. Diagn.* Ser. II. i. 11.—Thessal.
anemonoides, Willd. in Ges. Nat. Fr. Berl. Mag. v. (1811) 401 t. 9. f. 6 = Isopyrum grandiflorum.
aragonensis, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 966.—Hispan.
arbascensis, *Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse*, ii. (1874) 449 ; (*Massiv d'Arbas*, 85).—Europ.
arctica, Lond. Hort. Brit. 610 = formosa.
atrata, Koch, in Flora, xiii. (1830) i. 118 = vulgaris.
atropurpurea, Miq. in Ann. Mus. Lugd. Bat. iii. 8 = Buergeriana.
atropurpurea, Willd. Enum. Hort. Berol. 577 = viridiflora.
aurea, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 174 = chrysantha.
Bauhini, Schott, in Verh. Bot. Zool. Ver. Wien, iii. (1853) 128 = Einseleana.
Bernardi, Gren. & Godr. Fl. Fr. i. 45 = vulgaris.
Bertolonii, *Schott, in Verh. Bot. Zool. Ver. Wien*, iii. (1853) 127.—Europ.
bicolor, Ehrb. Beitr. viii. 146 = sibirica.
blanda ×, Lem. Illustr. Hortic. (1857) t. 146 = vulgaris.
brachyceras, Turcz. ex Fisch. & Mey. in Maund, Bot. Gard. n. 755 = leptoceras.
Braunii, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10, nomen.—Europ.
brevistyla, *Hook. Fl. Bor. Am.* i. 24.—Am. bor.
Buergeriana, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1846) 183.—Japon.
caerulea, *James, in Long, Exped. Rocky Mount.* ii. 345.—Am. bor. occ.
californica, *A. Gray, in Proc. Am. Acad.* vii. (1868) 328.—Am. bor. occ.
californica, Hartw. in Gard. Chron. (1854) 836 = formosa.
campyloceutra, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10.—Europ.
canadensis, Hook. Fl. Bor. Am. i. 24 = formosa.
canadensis, *Linn. Sp. Pl.* 533.—Am. bor.
canadensis, Pall. Reise, iii. 230 = viridiflora.
caucasica, Ledeb. ex Rupr. Fl. Cauc. 32 = olympica.
chrysantha, *A. Gray, in Proc. Am. Acad.* viii. (1873) 621.—N. Mexie.
coerulea, Auct. = caerulea.
collina, Jord. Diag. i. 84 = vulgaris.
concolor, Fisch. ex Steud. Nom. ed. II. i. 115 = vulgaris.
confusa, Rota, Prosp. Fl. Bergamo, 99 = Einseleana.
corniculata, Vill. Cat. Hort. Strasb. 250 = vulgaris.
cornuta, Gilib. Fl. Lituan. ii. 286 = vulgaris.
corsica, Soleir. ex Nym. Consp. 18 = vulgaris.
cyclophylla, Jeaub. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse, iii. (1875–76) 555 ; (Massif Laurenti, 139) = pyrenaica.
dahurica, Patr. in Deless. Ic. Sel. t. 49 = viridiflora.
dichroa, *Freyn, in Flora*, xliii. (1880) 26.—Europ.
dioica, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10, nomen.—Europ.
discolor, Steud. Nom. ed. II. i. 115 = glandulosa.
discolor, *Levier & Leresche, in Journ. Bot.* xvii. (1879) 197.—Hispan.
dumeticola, Jord. Diag. i. 86 = vulgaris.
Ebneri, Zimm. ex Nym. Consp. 18 = vulgaris.
ecalcarata, Hort. ex Steud. Nom. ed. II. i. 115 = vulgaris.
Einseleana, *Fr. Schultz, Arch. Fl. Fr. et Allem.* 135.—Europ.
elata, Ledeb. Ind. Hort. Dorp. Suppl. (1824) 41 = vulgaris.
elegans, Pope, ex Steud. Nom. ed. II. i. 115 = vulgaris.
elegans, Salisb. Prod. 374 = canadensis.
eximia, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10, nomen.—Europ.
eximia, Van Houtte, ex Planch. in Fl. des Serres, Sér. II. ii. (1857) 13. t. 1188 = formosa.
flabellata, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1846) 183.—Japon.
flava, Lam. Encyc. i. 149 = viridiflora.
flavescens, *S. Wats. in Bot. King, Exp.* 10.—Calif.
flaviflora, Tenney, in Am. Natural. i. (1868) 388 = canadensis.
formosa, *Fisch. in DC. Prod.* i. 50.—Kamtsch.
formosa, Hook. f. Bot. Mag. t. 6552 = Hookeri.
fragrans, *Benth. in Maund, Botanist*, iv. t. 181.—Reg. Himal.
Gaertneri, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10, nomen.—Europ.
Garnieriana, Sweet, Brit. Fl. Gard. Ser. II. v. t. 103 = sibiriea.
Gebleri, *Bess. ex Turcz. in Flora*, xvii. (1834) i. *Beibl.* 5.—Europ.
glandulosa, *Fisch. Hort. Gorenk.* ed. II. (1812) 48 ; *et ex Link, Enum. Hort. Berol.* ii. 84.—Sibir. altaic.
glandulosa, Miq. in Ann. Mus. Lugd. Bat. iii. 8 = flabellata.
glauca, *Lindl. Bot. Reg.* xxvi. (1840) t. 46.—Reg. Himal.
glaucescens, Baker, in Gard. Chron. (1878) II. 203 = vulgaris.
glaucophylla, Steud. in Flora, xxxix. (1856) 407 = vulgaris.
grandiflora, Patr. ex DC. Syst. i. 337 = alpina.
grandiflora, Schangin, in Pall. N. Nord. Beitr. vi. 49 = glandulosa.
grata, *Maly, ex Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 8, nomen.—Europ.
Haenkeana, Koch, Syn. Fl. Germ. 23 = vulgaris.
Haynaldi, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 9.—Europ.
hirsuta, *Timb.—Cf. Bull. Soc. Bot. Fr.* xxix. (1882), *Bibl.* 47.—Gallia merid.
hispanica, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10.—Hispan.
Hookeri, *Borbas, in Természs. Füzetek*, viii. (1884) 311.—Am. bor.
Huteri, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 9.—Europ.
hybrida, Sims, in Bot. Mag. t. 1221 = vulgaris.
inversa, Mill. Gard. Diet. ed. VIII. n. 3 = vulgaris.
Jonesii, *Parry, in Am. Natural.* viii. (1874) 211.—Am. bor.
jucunda, Fisch. & Lallem. Ind. Sem. Hort. Petrop. vi. (1840) 43 = glandulosa.
jucunda, Fisch. & Mey. ex Ledeb. Fl. Ross. i. 736 = vulgaris.
kanaoriensis, Jaequem. ex Camb. in Jaequem. Voy. Bot. 7. t. 5 = Moorcroftiana.
Karelini, Baker, in Gard. Chron. (1878) II. 76 = vulgaris.
Kareliniana, *C. A. Mey. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 53.—Hab. ?
Kitaibelii, *Schott, in Verh. Bot. Zool. Ver. Wien*, iii. (1853) 129.—Armenia.
lactiflora, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 374.—Sibir.
leptoceras, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 33.—Sibir.
leptoceras, Nutt. in Journ. Ac. Philad. vii. (1834) 9 = caerulea.
longisepala, *Zimm. ex Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 9, nomen.—Europ.
longissima, *A. Gray, ex S. Wats. in. Proc. Am. Acad.* xvii. (1881–82) 317.—Am. bor.
lutea, *Lam. Encyc.* i. 149.—Hab. ?
lutescens, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10, nomen.—Europ.
macrantha, Hook. & Arn. Bot. Beech. Voy. 317. t. 72 = caerulea.
macrocentra, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10, nomen.—Europ.
Magnolii, Loret, in Loret & Barraud. Fl. Montp. ii. 796 = Kitaibelii.
mexicana, Hook. Bot. Mag. t. 3919 = Skinneri.
minuta, Steph. ex Ledeb. Fl. Ross. i. 53 = Isopyrum grandiflorum.
mollis, *Jeanb. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse*, iii. (1875–76) 555.—Gallia.
montana, Sternb. in Regensb. Denkschr. ii. (1818) 60 = alpina.
Moorcroftiana, *Wall. Cat.* n. 1713.—Reg. Himal.
Moorensteniana, Steud. Nom. ed. II. i. 115, sphalm. = pracc.
nemoralis, *Jord. Diag.* i. 83.—Gallia.
nevadensis, *Boiss. & Reut. Cat. Hort. Genev.* (1854) 4 ; *et in Ann. Sc. Nat.* Sér. IV. ii. (1854) 380.—Hispan.
nigricans, Baumg. Enum. Stirp. Transs. ii. 104 = vulgaris.
nivalis, Falc. ex Baker, in Gard. Chron. (1878) II. 76 = glauca.
nivea, Baumg. ex Baker, in Gard. Chron. (1878) II. 76 = vulgaris.
olympica, *Boiss. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 360.—Oriens.
orthantha, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10, nomen.—Europ.
Ottonis, *Orph. ex Boiss. Diagn.* Ser. II. i. 11.—Graecia.
oxysepala, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 10).—As. or.
paraplesia, Schur, Enum. Pl. Transs. 28 = vulgaris.
parviflora, *Ledeb. in Mém. Acad. Pétersb.* v. (1814) 544.—Sibir.
platysepala, Reichb. Ic. Fl. Germ. t. 4730 = vulgaris.
plena, Hort. ex Steud. Nom. ed. II. i. 115 = vulgaris.
praecox, Jord. Diag. i. 85 = vulgaris.
pubiflora, *Wall. Cat.* n. 4714.—Ind. or.
pycnotricha, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10.—Europ.
pyrenaica, *DC. Fl. Fr.* iv. 912.—Pyren. ; Appenin.
pyrenaica, Koch, Syn. Fl. Germ. ed. II. 24 = Einseleana.
pyrenaica, Reichb. Ic. Fl. Germ. t. 4732 = Bertolonii.
Reuteri, Boiss. Diagn. Ser. II. i. 10 = Bertolonii.
Reuteriana, Reichb. f. ex Nym. Consp. 18 = alpina.
ruscinonensis, *Timb. & Jeanb. in Bull. Soc. Sc. Nat. Toulouse*, iii. (1875–76) 554.—Gallia.
Schinneri, Port. in Rev. Hortic. Sér. II. iv. (1845–46) 185 = Skinneri.
Schottii, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10.—Europ.
sibirica, Don, ex Steud. Nom. ed. II. i. 115 = vulgaris.
sibirica, *Lam. Encyc.* i. 150.—Sibir.
sibirica, Schur, ex Nym. Consp. 18 = glandulosa.
silvestris, Neck. Delic. Gallo-Belg. i. 234 = vulgaris.
sinensis, *Heynh. Nom.* ii. 41.—China.
Skinneri, *Hook. Bot. Mag.* t. 3919.—Mexie.
speciosa, DC. Syst. i. 336 = sibirica.
speciosa, Timb. in Bull. Soc. Hist. Nat. Toulouse, iv. (1870) 155 = arbascensis.
spectabilis, Lem. in Illustr. Hortic. xi. (1864) t. 403 = sibiriea.
stellata, Hort. ex Steud. Nom. ed. I. 61 ; ed. II. i. 115 = vulgaris.
stenopetala, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 8, nomen.—Europ.
Sternbergii, Reichb. Fl. Germ. Exeurs. 749 = alpina.
subalpina, Bor. Fl. Centr. Fr. ed. III. ii. 24 = vulgaris.
subscaposa, Borbas, in Magyar Tudom. Akad. xii. n. VI. (1882) 18 = vulgaris.
sulphurea, *Zimm. ex Borbas, l. c.* 9.—Europ.
sylvestris, vide *silvestris*.
sylvestris, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 4 = vulgaris.
Szaboi, *Borbas, in Magyar Tudom. Akad.* xii. n. VI. (1882) 10, nomen.—Europ.
thalictrifolia, Schott & Kotschy, in Verh. Bot. Zool. Ver. Wien, iii. (1853) 130 = Einseleana.
thalictroides, Schlecht. in Linnaea, vi. (1831) 581 = parviflora.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 94 ; iv. (1853) 4 = vulgaris.
truncata, Fisch. & Mey. Ind. Sem. Hort. Petrop. (1843) Suppl. 8 = formosa.
variegata, Moench, Meth. 311 = canadensis.
versicolor, Salisb. Prod. 374 = vulgaris.
viridiflora, *Pall. in Act. Petrop.* (1779) 260. t. 11.—Sibir.
viscosa, Gouan, Fl. Monsp. 267 = Kitaibelii.
viscosa, Reichb. Fl. Germ. Excurs. 748 = Einseleana.
viscosa, Suffr. ex Reichb. l. c. 749 = pyrenaica.
volubilis, *Maack, ex Trautv. in Act. Hort. Petrop.* viii. (1883) 54.—Manchuria.
vulgaris, *Linn. Sp. Pl.* 533.—Europ. ; Oriens.
vulgaris, Richards. App. Frankl. Journ. ed. II. 21 = brevistyla.
vulgaris, Thunb. Fl. Jap. 232 = flabellata.
Wittmanniana, Stev. ex Fisch. & Mey. & Ave-Lall. Ind. Sem. Hort. Petrop. xi. Suppl. 15 = olympica.

AQUILICIA, Linn. Mant. ii. 211 (1771) = **Leea**, Linn. (Ampel.).
Ottilis, Gaertn. Fruct. i. 275 = L. sambucina.
sambucina, Linn. Mant. 211 = L. sambucina.

ARABIDAEA.—Cf. Rev. Hortic. (1868) 154 = **Arrabidaea**, DC. (Bignon.).

ARABIDIUM, Spach, Hist. Vég. Phan. vi. 436 (1838) = **Arabis**, Linn. (Crucif.).
alpestre, Spach, l. c. 438 = Arabis alpina.
alpinum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = Arabis alpina.
saxaticolum, Fourr. l. c. = Arabis saxatilis.

ARABIDOPSIS, Schur, Enum. Pl. Transs. 55 (1866) = **Sisymbrium**, Linn. (Crucif.).
pinnatifida, Rupr. Fl. Caucas. 86 = Braya pinnatifida.
Thaliana, Schur, Enum. Pl. Transs. 55 = S. Thalianum.

ARABIS, Adans. Fam. ii. 422 (1763) = **Iberis**, Dill. (Crucif.).

ARABIS, Linn. Gen. ed. I. 198 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 69.
ABASICARPON, Andrz. ex Reichb. in Moessl. Handb. ii. 1150 (1828).
ARABIDIUM, Spach, Hist. Vég. Phan. vi. 436 (1838).
ARABISA, Reichb. Handb. 260 (1841).
CAULOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332.
LOMASPORA, DC. ex Steud. Nom. ed. II. ii. 65 (1841).
PILOSELLA, Kostel. Hort. Prag. 104 (1844).
PSILARABIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332.
STEVENIA, Adams & Fisch. in Mém. Soc. Nat. Mosc. v. 84 (1820).
TURRITIS, Tourn. ex Linn. Syst. ed. I. (1737).
accedens, Jord. Diag. i. 115 = hirsuta.
aizoides, *Bonato, Cat. Pl. Hort. Patav. (sec. Steud. Nom.* ed. II. i. 115).—Hab. ?
alaschanica, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 421.—Mongolia.
albida, *Stev. in Fisch. Cat. Hort. Gorenk.* 51.—Reg. Mediterr.; Oriens.
Alliaria, Bernh. Syst. Verz. Erf. 195 = Sisymbrium Alliaria.
Allionii, *DC. Fl. Fr.* iv. 676.—Pedemont.
alpestris, *Schleich. ex Reichb. Ic. Fl. Germ.* ii. 13. n. 4338 b.—Helvetia.
alpina, Georgi, Beschr. Russ. Reich. iii. IV. 1137 = albida.
alpina, Krock. ex Steud. Nom. ed. II. i. 115 = sagittata.
alpina, *Linn. Sp. Pl.* 664.—Europ.; Am. bor.
alpina, Lucé, Fl. Osil. 228 = Gerardi.
amara, Banks, ex DC. Syst. ii. 246 = Cardamine rhomboidea.
ambigua, DC. Syst. ii. 231 = petraea.
amplexicaulis, *Edgew. in Trans. Linn. Soc.* xx. (1851) 31.—Reg. Himal.
andicola, H. B. & K. Nov. Gen. et Sp. v. 81 = Sisymbrium hispidulum.
androsacea, *Fenzl, Pugill. Pl. Nov. Syr.* 13.—As. Min.
angustifolia, Lam. Encyc. i. 220 = Hutchinsia alpina.
apennina, *Tausch, in Syll. Ratisb.* ii. (1828) 244.—Italia.
arcuata, Dulac, Fl. Hautes-Pyr. 200 = Sisymbrium Thalianum.
arcuata, *A. Gray, in Proc. Am. Acad.* vi. 187.—Calif.
arcuata, Shuttlw. ex Godet, Fl. Jura, i. (1852) 38 = alpestris.
arenaria, *Schrank, Fl. Monac.* iii. 256.—Europ.
arenosa, Cham. & Schlecht. in Linnaea, i. (1826) 17 = kamtschatica.
arenosa, *Scop. Fl. Carn.* ed. II. ii. 32.—Europ.
arvensis, Bernh. Syst. Verz. Erf. 195 = Conringia austriaca.
arvensis, *Edgew. in Trans. Linn. Soc.* xx. (1851) 32.—Ind. bor. occ.
aspera, All. Ped. Auct. 18. t. 2. f. 2 = auriculata.
attenuata, Royle, ex Hook. f. in Journ. Linn. Soc. vi. (1862) 143 = tibetica.
aubrietioides, *Boiss. Fl. Orient.* i. 175.—Cilicia.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 52.—As. Min.; Syria.
auriculata, *Lam. Encyc.* i. 219.—Europ.; Oriens.
Balansae, *Boiss. & Reut. Diagn.* Sér. II. v. 17.—Algeria.
Barbarea, Bernh. Syst. Verz. Erf. 195 = Barbarea vulgaris.
bellidifolia, Crantz, Stirp. Austr. ed. I. fasc. i. 42; ed. II. 44 = pumila.
bellidifolia, *Jacq. Obs. Bot.* i. 22.—Europ.
bellidifolia, Scop. Fl. Carn. ed. II. ii. 31 = Cardamine alpina.

ARABIS:—
bellidioides, Lam. Encyc. i. 220 = Cardamine alpina.
bijuga, *Watt, in Journ. Linn. Soc.* xviii. (1881) 378.—Reg. Himal.
Billardieri, DC. Syst. ii. 218 = albida.
blepharophylla, *Hook. & Arn. Bot. Beech. Voy.* 321.—Calif.
Boissieri, Griseb. Spicil. Fl. Rumel. i. 248 = drabaeformis.
borealis, Andrz. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 33 = hirsuta.
Boryi, Boiss. Voy. Bot. Espagne, ii. 26. t. 5 = Braya pinnatifida.
brachycarpa, *Rupr. Fl. Cauc.* 73, *in textu*.—Reg. Caucas.
brachypoda, Boiss. Elench. 8 = parvula.
bracteata, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 291.—Bolivia.
brassicaeformis, *Wallr. Sched. Crit.* 359.—Europ.
Breutelii, *Lange, Consp. Fl. Groenl.* i. 50.—Groenl.
brevifolia, DC. Syst. ii. 218 = albida.
Breweri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 123.—Calif.
bryoides, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 55.—Graecia.
bulbosa, Muhl. Cat. 63; Spreng. Fl. Hal. Mant. 46 = Cardamine rhomboidea.
bursifolia, Lam. Fl. Fr. ii. 511 = Sisymbrium bursifolium.
cadmea, *Boiss. Diagn.* Ser. I. viii. 21.—Caria.
caerulea, *Haenke, in Jacq. Collect.* ii. 56.—Europ.
caespitosa, *Schleich. Cat. Pl. Helv.* ed. IV. 8.—Helvet.
camtschatica, Willd. ex DC. Syst. ii. 242 = Stelleri.
canadensis, *Linn. Sp. Pl.* 665.—Am. bor.
canadensis, Mill. Gard. Dict. ed. VIII. n. 6 = Diplotaxis muralis.
canescens, Brocchi, in Bibliot. Ital. xxix. 90 = alpina.
canescens, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 83.—Am. bor.
cantabrica, *Leresche & Levier, in Journ. Bot.* xvii. (1879) 197.—Hispan.
capensis, Linn. Syst. ed. X. 1135; Burm. f. Prod. Fl. Cap. 18 = Heliophila pusilla.
Cardamines, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 292.—Austral.
carduchorum, *Boiss. Fl. Orient.* i. 173.—Armenia.
carpetana, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 822.—Hispan.
caucasica, Willd. Enum. Hort. Berol. Suppl. 45 = albida.
cebennensis, *DC. Syst.* ii. 234.—Gallia.
cenisia, Reut. Cat. Jard. Bot. Genèv. (1853) 4 = alpestris.
chinensis, Rottl. ex Wight, Illustr. i. sub t. 12 = Lepidium sativum.
ciliaris, Willd. Enum. Hort. Berol. 684 = pumila.
ciliata, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 107.—Europ.
ciliata, Auct. ex Nym. Consp. i. 33 = alpestris.
ciliata, Tausch, Ind. Pl. Sudet. (sec. Steud. Nom. ed. II. i. 117) = hirsuta.
cilicica, Boiss. & Bal. Diagn. Ser. II. v. 17 = ionocalyx.
Clusiana, Schrank, Fl. Monac. ii. 125 = alpina.
coerulea, vide caerulea.
colchaguensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 293.—Chili.
collina, *Tenore, Prod. Fl. Nap.* 39.—Italia.
collisparsa, Jord. Diagn. i. 116 = hirsuta.
colorata, *Tausch, in Flora*, xix. (1836) II. 405.—Europ.
Commersonii, *DC. Syst.* ii. 228.—Reg. Argent.
conferta, Willd. ex Reichb. Fl. Germ. Excurs. 680 = hirsuta.
conringioides, *Ball, in Journ. Bot.* xi. (1873) 297.—Marocc.
constricta, Griseb. Spicil. Fl. Rumel. i. 249 = hirsuta.
contracta, *Celak. Prod. Fl. Boehm.* 155.—Bohem.
contracta, Spenn. Fl. Friburg. iii. 925 = hirsuta.
cordata, Boiss. in Ann. Sc. Nat. Sér. II. xviii. (1842) 51 = laxa.
Costae, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 820.—Hispan.
Crantziana, Ehrh. Beitr. v. 177 = petraea.
crassifolia, *Ledeb. in Baier. Bot. Ges. Denkschr.* iii. (1841) 107.—Rossia.
cremocarpa, Boiss. & Bal. Diagn. Ser. II. v. 16 = laxa.
crepidiopoda, Griseb. ex. Pant. in Oestr. Bot. Zeitschr. xxiii. (1873) 268 = pseudoturritis.

ARABIS:—
cretica, Boiss. & Heldr. Diagn. Ser. I. viii. 20 = muralis.
crispata, Willd. Enum. Hort. Berol. 684 = alpina.
croatica, Schott, ex Nyman & Kotschy, Analect. Bot. 44 = neglecta.
cuneifolia, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 17 = alpina.
curtisiliqua, DC. Syst. ii. 223 = hirsuta.
Cusickii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 363.—Am. bor.
dacica, Heuff. ex Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) I. 307 = Halleri.
dasycarpa, Andrz. ex DC. Syst. ii. 244 = auriculata.
declinata, Schrad. Ind. Sem. Hort. Goett. (1831); (ex Linnaea, viii. (1833) Litt. 22) = Holboellii.
decumbens, *Ball, in Journ. Bot.* xi. (1873) 297.—Marocco.
deflexa, *Boiss. Fl. Orient.* i. 175.—Lycia.
dentata, [Clairv]. Man. Herb. 223 = Braya pinnatifida.
dentata, *Torr. & Gray, Fl. N. Am.* i. 80.—Am. bor.
docica, Fuss, in Verh. Siebenb. Ver. Naturw. xiii. (1862) 150, sphalm. = dacica.
Douglassii, Torr. in Torr. & Gray, Fl. N. Am. i. 83, in syn. (cum cit. falsa) = Cardamine rotundifolia.
drabaeformis, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 55.—Bithyn.
drabaeformis, Griseb. Spicil. Fl. Rumel. i. 248 = bryoides.
Dregeana, *Harv. & Sond. Fl. Cap.* i. 22.—Afr. austr.
Drummondii, *A. Gray, in Proc. Am. Acad.* vi. 187.—Am. bor.
elongata, Salisb. Prod. 272 = A. Turrita.
eriocarpa, Schur, Enum. Pl. Transs. 46 = A. Turrita.
eriopoda, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 63.—Sibir. or.
erubescens, *Ball, in Journ. Bot.* xi. (1873) 297.—Marocco.
erysimoides, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 145.—Soongaria.
Eschscholtziana, Andrz. ex Ledeb. Fl. Ross. i. 118 = hirsuta.
falcata, Michx. Fl. Am. Bor. i. 31 = canadensis.
falcata, *A. Rich. Tent. Fl. Abyss.* i. 17.—Abyss.
falklandica, Benth. & Hook. f. Gen. i. 69 = macloviana.
farinacea, *Rupr. Fl. Caucas.* 71.—Caucas.
fastigiata, Hook. f. Fl. N. Zel. ii. 324 = Cardamine fastigiata.
flagellosa, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 72.—Japon.
flaviflora, *Bunge, in Mém. Acad. Pétersb.* Sér. VI. vii. (1859) 582.—Armenia.
flexuosa, Hort. Berol. ex Reichb. Fl. Germ. Excurs. 679 = serpyllifolia.
floribunda, Schur, Enum. Pl. Transs. 44 = petraea.
foliosa, *Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1862) 160.—Reg. Himal.
fruticulosa, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 19.—Sibir. alt.
furcata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 362.—Am. bor.
Gerardi, *Bess. in Koch, Syn. Fl. Germ.* ed. I. 38.—Europ.; Reg. Cauc.
gigantea, Hook. Ic. Pl. t. 259 = Cardamine stylosa.
glabra, Bernh. Syst. Verz. Erf. 195 = perfoliata.
glandulosa, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 146.—Tibet.; Soongar.
glareosa, Schur, in Verh. Siebenb. Ver. Naturw. i. (1850) 106; iv. (1853) 59 = arenosa.
glastifolia, *Reichb. Ic. Fl. Germ.* xii. 13.—Europ.
Gmelini, Willd. ex Ledeb. Fl. Ross. i. 131 = Parrya macrocarpa.
gracilescens, Jord. Diag. i. 119 = hirsuta.
gracilis, Willd. ex Ledeb. Fl. Ross. i. 185 = Sisymbrium mollissimum.
gracilis, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 158 = Sisymbrium Wallichii.
grandiflora, Linn. Amoen. Acad. ii. 279 = Parrya macrocarpa.
grandiflora, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 142 = alpina.
Halleri, *Linn. Sp. Pl.* ed. II. 929.—Europ.
Halleri, Lucé, Fl. Osil. 228 = arcnosa.
hastulata, Bert. ex DC. Prod. i. 150 = Cardamine resedifolia.
Heliophila, DC. Syst. ii. 237 = Farsetia Jacquemonti.
hesperidoides, A. Gray, Man. Bot. U. St. ed. V. 68 = Thelypodium pinnatifidum.

ARABIS:—
heteromalla, Schrad. Ind. Sem. Hort. Goett. ex Linnaea, viii. (1833) Litt. 22 = Holboellii.
heterophylla, Forst. ex DC. Syst. ii. 265 = Cardamine hirsuta.
heterophylla, Nutt. ex Torr. & Gray, Fl. N. Am. i. 81 = laevigata.
heterophylla, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 59.—Austria.
himalaica, *Edgew. in Trans. Linn. Soc.* xx. (1851) 31.—Reg. Himal.
hirsuta, DC. Fl. Fr. iv. 675 = ciliata.
hirsuta, *Scop. Fl. Carn.* ed. II. ii. 30.—Europ.; Oriens; Am. bor.
hirsuta, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 142 = alpina.
hirta, Lam. Encyc. i. 220 = stricta.
hirtella, Jord. Diag. i. 120 = hirsuta.
hispida, Soland. in Ait. Hort. Kew. ed. I. ii. 400 = stricta.
hispida, Murr. Syst. Veg. ed. XIII. 501 (cf. Linn. f. Suppl. 298) = petraea.
hispida, Moench, Meth. 258 = pendula.
Holboellii, *Hornem. Fl. Dan.* t. 1879.—Am. bor.
Holboelliana, Hort. Hafn. ex Trevir. in Flora, xvi. (1833) 124 = praec.
Hookeri, *Lange, Consp. Fl. Groenl.* i. 50.—Groenl.
Hornungiana, *Schur, Enum. Pl. Transs.* 43.—Transsylv.
Huetii, *Trautv. in Act. Hort. Petrop.* ii. (1878) 496.—Reg. Caucas.
humilis, Schleich. ex DC. Syst. ii. 224 = muralis.
hybrida, Reut. Suppl. Cat. Pl. Genèv. 8 = alpestris.
idanensis, Jord. Diag. i. 117 = hirsuta.
incana, Moench, Meth. 257 = alpina.
incana, *Roth, Catalect. Bot.* i. 79.—Europ.
incana, Willd. Enum. ex Steud. Nom. ed. II. i. 116 = muralis.
incana, Willd. Herb. ex Steud. l. c. = collina.
incanescens, Munro, Pl. Agra, 6, ex Hook, f. & Thoms. in Journ. Linn. Soc. v. (1861) 148 = Farsetia Hamiltonii.
incarnata, *Lamb. ex DC. Syst.* ii. 210.—Sibir.
integrifolia, *Lapeyr. Hist. Abr. Pl. Pyr.* 385. *Suppl.* 93.—Europ.
intermedia, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 92.—Europ.
intermedia, Hoppe, ex Steud. Nom. ed. II. i. 116 = bellidifolia.
ionocalyx, *Boiss. & Heldr. Diagn.* Ser. I. viii. 20.—As. Min.
japonica, *Regel, Gartenfl.* (1863) 308. t. 414.—Japon.
japonica, A. Gray, in Mem. Am. Acad. N. S. vi. (1858) 381 = Arabis Stelleri.
jugicola, Jord. Diag. i. 121 = bellidifolia.
kamtschatica, Auct. vide *camtschatica*, Willd.
kamtschatica, *Fisch. ex DC. Syst.* ii. 231.—Kamtschatca.
kamtschatica, Pall. ex Ledeb. Fl. Ross. i. 185 = Sisymbrium mollissimum.
kanawarensis, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 142 = amplexicaulis.
Kochii, Jord. Diag. i. 112 = Gerardi.
kokanika, *Regel, Pl. Nov. Fedsch.* 6.—Turkestan.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 232.—Turkestan.
lactea, Stev. ex Rupr. Fl. Cauc. 289 = mollis.
laevigata, *DC. Syst.* ii. 237.—Am. bor.
laevigata, Hook. Fl. Bor. Am. i. 43 = Drummondii?
lanata, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 248.—Peruv.
lasioloba, *Link, Enum. Hort. Berol.* ii. 163.—Mexico.
lateripendens, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = A. Turrita.
latifolia, *Dur. in Expl. Sc. Algér.* t. 72.—Algeria.
latifolia, Willd. ex Ledeb. Fl. Ross. i. 131 = Parrya Linnaeana.
laxa, *Sibth. & Sm. Prod. Fl. Graec.* ii. 28.—As. Min.
leptocarpa, Wall. Cat. n. 4784 = Sisymbrium Wallichii.
leptocarpaea, Fisch. ex Reichb. Fl. Germ. Excurs. 680 = alpina.
Lerchenfeldiana, Schur, Enum. Pl. Transs. 42 = alpina.
levigata, Poir. Encyc. Suppl. i. 411 = laevigata.
lilacina, Schrad. Ind. Sem. Hort. Goett. vii. (1832) 42 = Holboellii.
longesiliqua, *C. & J. Presl, Delic. Prag.* 16.—Sicilia.

ARABIS:—
longifolia, DC. Syst. ii. 219 = albida.
longirostris, *S. Wats. in Bot. King, Exp.* 17.—Calif.
longisiliqua, Wallr. Sched. Crit. 359 = A. Turrita.
lucens, Dulac, Fl. Hautes-Pyr. 202 = bellidifolia.
lucida, *Linn. f. Suppl.* 298.—Europ.
ludoviciana, *C. A. Mey. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* ix. 60.—Am. bor.
lusitanica, *Boiss. Diagn.* Ser. II. i. 20.—Lusitan.
Lyallii, *S. Wats. in Proc. Am. Acad.* xi. (1876) 122.—Am. bor. occ.
lyraefolia, DC. Syst. ii. 244 = canadensis.
lyrata, *Linn. Sp. Pl.* 665.—Am. bor.
macloviana, *Hook. Ic. Pl.* t. 498.—Ins. Falkland.
macrocarpa, Torr. Mex. Bound. Rep. 32 = perfoliata.
madonia, *C. & J. Presl, Delic. Prag.* 15.—Sicilia.
major, S. F. Gray, Nat. Arr. Brit. Pl. ii. 676 = A. Turrita.
Marschalliana, Hort. ex Steud. Nom. ed. II. i. 116 = hirsuta.
mexicana, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 319.—Mexico.
microphylla, Nutt. ex Torr. & Gray, Fl. N. Am. i. 82 = Drummondii?
minor, Hort. Berol. ex DC. Syst. ii. 226 = incana.
mollis, Rafin. in Am. Monthly Mag. ii. (1817) 43 = canadensis.
mollis, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 270.—Reg. Caucas.
montana, Bernh. ex DC. Syst. ii. 225 = stricta.
montana, Lam. Encyc. i. 219 = hirsuta.
Montbretiana, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 53.—Oriens.
monticola, Jord. Diag. i. 107 = alpina.
moschata, Hort. ex Reichb. Fl. Germ. Excurs. 680 = incana.
multicaulis, Bell. ex DC. Syst. ii. 233 = Halleri.
multiflora, *DC. Syst.* ii. 243.—Mont. Pyren.
multijuga, *Borb. in Linnaea*, xli. (1877) 604.—Europ.
muralis, *Bertol. Rar. Ital. Pl. Decad.* ii. 37.—Graecia.
muralis, Salisb. Prod. 272 = Sisymbrium Thalianum.
muricola, Jord. Diag. i. 125 = muralis.
Nasturtium, [Clairv.] Man. Herb. 223 = Nasturtium officinale.
neglecta, *Schult. Oestr. Fl.* ii. 248.—Mont. Carpath.
nemoralis, Steud. Nom. ed. II. i. 117 = hirsuta.
nemorensis, C. A. Mey. Verz. Pfl. Cauc. 179 = Gerardi.
nemorensis, Reichb. Fl. Germ. Excurs. 681 = hirsuta.
nepetaefolia, *Boiss. Fl. Orient.* i. 177.—Persia.
nivalis, Guss. Pl. Rar. 389 = serpyllifolia.
nivalis, Spreng. Syst. ii. 893 = Macropodium nivale.
Nordmanniana, Rupr. Fl. Cauc. 72 = mollis.
nova, Vill. Prosp. 39 = saxatilis.
nuda, Bélang. ex Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 54 = Sisymbrium nudum.
nudicaulis, DC. Syst. ii. 240 = Parrya macrocarpa.
nudiuscula, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 22.—Afr. austr.
nutans, Moench, Meth. 258 = pumila.
nutans, Sims, Bot. Mag. t. 2219 = Allionii.
oblongata, *Wender. Hort. Marb.* (1824) ex Steud. Nom. ed. II. i. 117.—Hab.?
obtusifolia, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 58, 6 = alpina.
ochroleuca, *Boiss. & Heldr. Diagn.* Ser. II. i. 21.—Graecia.
ochroleuca, Lam. Encyc. i. 218 = A. Turrita.
olympica, Sibth. ex DC. Syst. ii. 219 = albida.
ovata, Poir. Encyc. Suppl. v. 557 = hirsuta.
ovirensis, Wulf. in Jacq. Coll. i. 196 = Halleri.
owaihiensis, *Cham. & Schlecht. in Linnaea*, i. (1826) 17.—Ins. Sandvic.
oxyota, DC. Syst. ii. 236 = pendula.
pachyrhiza, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 144.—Soongaria.
pangiensis, *Watt, in Journ. Linn. Soc.* xviii. (1881) 378.—Reg. Himal.
parviflora, Rafin. in Am. Monthly Mag. ii. (1817) 43 = Sisymbrium Thalianum.
parvula, *Dufour, in DC. Syst.* ii. 228.—Europ. austr.
patens, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 142 = amplexicaulis.
patens, *Sulliv. in Am. Journ. Sc.* i. (1819) 42, 49.—Am. bor.

ARABIS:—
Patriniana, DC. Syst. ii. 236 = pendula.
patula, *Host, Fl. Austr.* ii. 271.—Austria.
patula, Torr. Bot. Mex. Bound. 32 = Holboellii.
patula, Weinm. Cat. Hort. Dorp. (1819) 18 = auriculata.
pauciflora, Garcke, Fl. Deutschl. ed. IV. 22 = brassicaeformis.
pedemontana, Boiss. Diagn. Ser. I. i. 69 = cebennensis.
pelligera, Fenzl, in Kotschy, Reise Cilic. Taur. 391 = ionocalyx.
pendula, Lachen. ex Reichb. Fl. Germ. Excurs. 681 = A. Turrita.
pendula, *Linn. Sp. Pl.* ed. I. 665.—Sibir.; Am. bor.
pendula, Stell. ex DC. Syst. ii. 242 = Stelleri.
perfoliata, *Lam. Encyc.* i. 219.—Europ.; Sibir.; Reg. Himal.
permixta, Jord. Diag. i. 110 = Gerardi.
petiolaris, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 187.—Texas.
petiolata, Bieb. Fl. Taur. Cauc. ii. 126 = Sisymbrium Alliaria.
petraea, *Lam. Encyc.* i. 221.—Reg. bor. temp.
petricola, Jord. Diag. i. 121 = hirsuta.
petrogena, Kern. in Oestr. Bot. Zeitschr. xiii. (1863) 141 = arenosa.
Piaseczkii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 420.—China.
pinnatifida, Lam. Encyc. i. 221 = Braya pinnatifida.
planisiliqua, Pers. ex Reichb. Ic. Fl. Germ. xii. 13 = Gerardi.
platysperma, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 519.—Calif.
praecox, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 684 = procurrens.
procera, Jord. Diag. i. 114 = Gerardi.
procurrens, *Waldst. & Kit. Pl. Rar. Hung.* ii. 154. t. 144.—Hungar.; Banat.; Thracia.
propera, Jord. Diag. i. 118 = hirsuta.
propinqua, Jord. l. c. = hirsuta.
pseudoturritis, *Boiss. & Heldr. Diagn.* Ser. II. i. 20.—Graecia.
pterosperma, Edgew. in Trans. Linn. Soc. xx. (1851) 33 = alpina.
puberula, Nutt. ex Torr. & Gray, Fl. N. Am. i. 82 canescens.
pubescens, *Poir. Encyc. Suppl.* i. 413.—Algeria.
pubicalyx, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 72.—Japon.
pubigera, Jord. Diag. i. 115 = Gerardi.
pumila, *Jacq. Fl. Austr.* iii. 44.—Europ.
purpurascens, Presl, Fl. Sicul. i. 50 = longisiliqua.
purpurea, *Sibth. & Sm. Fl. Graec.* vii. t. 643.—Ins. Cyprus.
ramosa, Lam. Fl. Fr. ii. 510 = Sisymbrium Thalianum.
recta, Vill. Hist. Pl. Dauph. iii. 319 = auriculata.
Reichenbachii, Syme, Eng. Bot. ed. III. i. 168 = hirsuta.
reparda, *S. Wats. in Proc. Am. Acad.* xi. (1876) 122.—Calif.
reptans, Lam. Encyc. i. 222 = Draba caroliniana?
resediflora, H. B. & K. Nov. Gen. et Sp. v. 81 = Nasturtium arabiforme.
resedifolia, Lam. Fl. Fr. ii. 511 = Cardamine resedifolia.
retrofracta, R. Grah. in Edinb. Phil. Journ. (1827) 7 = Holboellii.
Retziana, Beurl. ex Nym. Consp. i. 33 = hirsuta.
rhaetica, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 85.—Europ.
rhomboidea, Pers. Syn. ii. 204 = Cardamine rhomboidea.
rigida, Dulac, Fl. Hautes-Pyr. 200 = brassicaeformis.
rigida, Wallr. Beitr. Fl. Hercyn. 229 = auriculata.
rigidula, Jord. Diag. i. 108 = Gerardi.
Roemeriana, Andrz. ex DC. Syst. ii. 228 = parvula.
rosea, DC. Syst. ii. 215 = muralis.
rosella, Jord. Diag. i. 126 = muralis.
rotundifolia, Rafin. in Am. Monthly Mag. ii. (1817) 43 = Draba caroliniana.
rubricaulis, Jord. Diag. i. 113 = Gerardi.
rugosa, Moench, Meth. 259 = A. Turrita.
runcinata, Lam. Encyc. i. 222 = petraea.
runcinata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 319.—Am. bor.
rupestris, Nutt. ex Torr. & Gray, Fl. N. Am. i. 81 = hirsuta.
sabauda, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 87.—Europ.

ARABIS:—
sagittata, DC. Fl. Fr. vi. 592 = hirsuta.
saxatilis, *All. Fl. Pedem.* i. 268.—Europ.
saxeticola, Jord. Diag. i. 106 = alpina.
saxicola, *Edgew. in Trans. Linn. Soc.* xx. (1846) 33.—Reg. Himal.
saxigena, Jord. Diag. i. 127 = muralis.
scabra, All. Fl. Pedem. i. 268 = serpyllifolia pumila.
scabra, Gilib. Fl. Lituan. ii. 61 = Sisymbrium Thalianum.
scapigera, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 54 = Sisymbrium nudum.
Schievereckiana, *Andrz. in DC. Prod.* i. 145.—Tyrol.
Scopoliana, *Boiss. ex Nym. Consp.* i. 35.—Europ.
segetalis, Schur, Enum. Pl. Transs. 45 = arenosa.
serpyllifolia, Hoppe, ex DC. Syst. ii. 241 = vochinensis.
serpyllifolia, *Vill. Prosp.* 39.—Europ.
serrata, *Franch. & Sav. Enum. Pl. Jap.* i. 33.—Japon.
sibirica, *Hort. ex Steud. Nom.* ed. II. i. 117.—Sibir.
sicula, *Huet, ex Reut. Cat. Grain.* (1857).—Sicilia.
sicula, Stev. ex Nym. Suppl. 32 = albida.
Sieberi, *Tausch, in Flora*, xix. (1836) II. 405.—Austr.
siliculosa, Lam. Encyc. i. 221 = Parrya macrocarpa.
simplicifolia, Schur, Enum. Pl. Transs. 43 = petraea.
sinaica, Boiss. Diagn. Ser. I. viii. 21 = auriculata.
sinaica, *Steud. Nom.* ed. II. i. 117, nom.—Arab.
sinuata, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 64.—Sibir.
sisymbroides, Hook. ex Hook. f. in Trans. Linn. Soc. xxiii. (1861) 314 = petraea.
Sophia, Bernh. Syst. Verz. Erf. 196 = Sisymbrium Sophia.
Soyeri, *Reut. & Huet, ex Huet, in Ann. Sc. Nat.* Sér. III. xix. (1853) 251.—Mont. Pyren.
sparsiflora, Nutt. ex Torr. & Gray, Fl. N. Am. i. 81 = Holboellii.
spathulata, *DC. Syst.* ii. 227.—Am. austr.
spathulata, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 81.—Am. bor.
Stelleri, *DC. Syst.* ii. 242.—As. bor. or.
stellulata, Desv. & Berthol. in Desv. Journ. Bot. ii. (1813) 76 = pumila.
stenocarpa, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 4 = saxatilis.
stenopetala, *Willd. Enum. Hort. Berol. Suppl.* 46.—Hab.?
Steveniana, *Rupr. Fl. Cauc.* 289.—Reg. Cauc.
stolonifera, Clairv. ex DC. Syst. ii. 233 = Halleri.
stolonifera, *Hornem. Hort. Hafn.* ii. 618.—Hab.?
stricta, Costa, Ampliac. 22 = Costae.
stricta, *Huds. Fl. Angl.* 292.—Europ.; Am. bor.
stricta, Sieber, ex Tausch, in Flora, xix. (1836) II. 406 = alpestris.
subcoriacea, Gren. ex Nym. Consp. 34 = bellidifolia.
subnivalis, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxv. (1882) 88.—Europ.
subnitens, Jord. Diag. i. 124 = alpestris.
subpinnatifida, *S. Wats. in Proc. Am. Acad.* xx. (1885) 353.—Am. bor. occ.
sudetica, Tausch, in Flora, xix. (1836) II. 407 = hirsuta.
suecica, Fries, Summa Veg. 30, 147 = Sisymbrium suecicum.
suffrutescens, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 362.—Am. bor. occ.
sulphurea, *Boiss. Fl. Orient.* i. 173.—Persia.
supina, Lam. Fl. Fr. ii. 512 = Sisymbrium supinum.
sylvestris, Scop. Fl. Carn. ed. II. ii. 32 = Brassica Cheiranthos.
taraxacifolia, T. Anders. in Hook. f. Fl. Brit. Ind. i. 136 = arvensis.
tatarica, Pall. Reise, ii. 676 = Parrya macrocarpa.
tenella, *Guss. & Tenore, Fl. Neap. Syll. App.* v. 20.—Lucania.
tenella, Host, Fl. Austr. ii. 273 = Halleri.
Tenorii, Huet, ex Nym. Consp. 34 = albida.
Thaliana, Crantz, Stirp. Austr. ed. II. 39 = petraea.
Thaliana, Linn. Sp. Pl. 665 = Sisymbrium Thalianum.
Thomsoni, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 145 = tibetica.
thyrsoidea, Sibth. & Sm. Fl. Graec. t. 642 = albida.
tibetica, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 143.—Reg. Himal.
toxophylla, Bieb. Fl. Taur. Cauc. iii. 448 = Sisymbrium toxophyllum.
transsilvanica, Schur, Enum. Pl. Transs. 43 = petraea.

ARABIS:—
trichopoda, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 63.—Rossia.
tuberosa, Muhl. ex Pers. Syn. ii. 204 = Cardamine rhomboidea.
Turczaninowii, *Ledeb. Fl. Ross.* i. 123.—Sibir.
Turrita, *Linn. Sp. Pl.* 665.—Europ.; Oriens.
Turritis, [Clairv.] Man. Herb. 223 = A. Turrita.
umbrosa, Crantz, Stirp. Austr. ed. I. fasc. i. 43; ed. II. 41 = A. Turrita.
umbrosa, Turcz. ex Steud. Nom. ed. I. ii. 118 = petraea.
undulata, Link, Enum. Hort. Berol. ii. 161 = alpina.
uralensis, Willd. ex Ledeb. Fl. Ross. i. 120 = petraea.
verna, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 105.—Reg. Mediterr.
vesula, Jord. Diag. i. 123 = alpestris.
violacea, Moench, Meth. 259 = verna.
virescens, Jord. Diag. i. 109 = Gerardi.
virginica, Poir. Encyc. Suppl. i. 413 = Cardamine hirsuta.
viscosa, DC. Syst. ii. 216 = albida.
vochinensis, *Spreng. Pugill.* i. 46. n. 81.—Austria.
wolgensis, Willd. ex Ledeb. Fl. Ross. i. 185 = Sisymbrium toxophyllum.
yokoscensis, *Franch. & Sav. Enum. Pl. Jap.* i. 34.—Japon.
Zeyheriana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 292.—Afr. austr.

ARABISA, Reichb. Handb. 260 (1837) = **Arabis**, Linn. (Crucif.).

ARACHIDNA, Plum. ex Moench, Meth. 121 (1794) = **Arachis**, Linn. (Legum.).
hypogaea, Moench, l. c. 122 = Arachis hypogaea.
quadrifolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 7.—Malaya.

ARACHIS, Linn. Syst. ed. I. (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 518.
ARACHIDNA, Moench, Meth. 121 (1794).
MUNDUBI, Adans. Fam. ii. 323 (1763).
africana, Burm. f. Fl. Cap. Prod. 22 = Voandzeia subterranea.
africana, Lour. Fl. Cochinch. ii. 430 = hypogaea.
americana, Tenore, in Mem. Ist. Incorr. i. (1811) 48 = hypogaea.
aprica, Walt. Fl. Carol. 183 = Stylosanthes elatior.
asiatica, Lour. Fl. Cochinch. ii. 430 = hypogaea.
fruticosa, Retz. Obs. Fasc. v. 26 = Stylosanthes mucronata.
glabrata, *Benth. in Trans. Linn. Soc.* xviii. (1841) 159.—Bras.
hypogaea, *Linn. Sp. Pl.* 741.—Reg. trop. (cult.).
marginata, *Gardn. in Hook. Ic. Pl.* t. 500.—Bras.
prostrata, *Benth. in Trans. Linn. Soc.* xviii. (1841) 159.—Bras.
pusilla, *Benth. l. c.*—Bras.
tuberosa, *Benth. l. c.*—Bras.
villosa, *Benth. l. c.*—Am. austr.

ARACHNANTHE, Blume, Rumphia, iv. 55. t. 196 (1828). *ORCHIDEAE*, Benth. & Hook. f. iii. 572.
ARACHNIS, Blume, Bijdr. 365 (1825).
ARMODORUM, Breda, Orch. Kuhl. & Hass. (1827).
ARRHYNCHIUM, Lindl. in Paxt. Fl. Gard. i. 142 (1851).
ESMERALDA, Reichb. f. Xenia Orch. ii. 38 (1874).
bilinguis, *Benth. in Journ. Linn. Soc.* xviii. (1881) 332.—Ind. or.
Cathcartii, *Benth. & Hook. f. Gen.* iii. 573.—Reg. Himal.
Lowei, *Benth. & Hook. f. l. c.*—Borneo.
moschifera, *Blume, Rumphia*, iv. 55.—Malaya.
Sulingi, *Benth. in Journ. Linn. Soc.* xviii. (1881) 332.—Malaya.

ARACHNE, Neck. Elem. ii. 348 (1790) = **Andrachne**, Linn. (Euphorb.).

ARACHNIMORPHA, Desv. in Ham. Prod. Fl. Ind. Occ. 28 (1825) = **Rondeletia**, Linn. (Rubiac.).
incana, Ham. l. c. = R. incana.

ARACHNIS, Blume, Bijdr. 365 (1825) = **Arachnanthe**, Blume (Orchid.).
moschifera, Blume, l. c. = Arachnanthe moschifera.

ARACHNITES, F. W. Schmidt, Fl. Boëm. i. 74 (1793) = **Ophrys**, Linn. (Orchid.).
alpina, F. W. Schmidt, l. c. = Herminium alpinum.
anthropophora, F. W. Schmidt, l. c. 77 = Aceras anthropophora.
apifera, Hoffm. Deutschl. Fl. ed. II. ii. 180 = O. apifera.
atrata, Tod. Hort. Panor. ii. t. 28 = O. aranifera.
Benoitii, Tod. l. c.—Hab.?
Bertolonii, Tod. Orch. Sicul. 79 = O. Bertolonii.
Biancae, Tod. l. c. 83 = O. Biancae.
bombyliflora, Tod. l. c. 91 = O. bombyliflora.
fuciflora, F. W. Schmidt, Fl. Boëm. i. 76 = O. Arachnites.
fusca, Tod. l. c. 98 = O. fusca.
Insengae, Tod. Nouv. Gen. 12 = O. Inzengae.
lunulata, Tod. Orch. Sicul. 77 = O. lunulata.
lutea, Tod. l. c. 95 = O. fusca.
Monorchis, Hoffm. Deutschl. Fl. ed. II. ii. 179 = Herminium Monorchis.
musciflora, F. W. Schmidt, Fl. Boëm. i. 75 = O. muscifera.
oxyrhynchus, Tod. Orch. Sicul. 81 = O. Arachnites.
pallida, Tod. l. c. 100 = O. pallida.
Speculum, Tod. l. c. 93 = O. Speculum.
tenthredinifera, Tod. l. c. 85 = O. tenthredinifera.

ARACHNITIS, Phil. in Bot. Zeit. 217 (1864). *BURMANNIACEAE*, Benth. & Hook. f. iii. 460.
uniflora, *Phil. l. c.*—Chili.

ARACHNOPOGON, Berg, ex Haberl, Cat. Pesth.; ex Steud. Nom. ed. II. i. 118 (1840), nomen.
heterospermum, Berg, l. c.—Hab.?

ARACHNOSPERMUM, Berg (sec. Steud. Nom. ed. II. i. 118) = **Hypochaeris**, Linn. (Compos.).

ARACHNOSPERMUM, F. W. Schmidt, Samml. Phys. Aufs. i. 274 (1795) = **Scorzonera**, Linn. (Compos.).
laciniatum, F. W. Schmidt, l. c. = S. laciniata.
resedifolium, F. W. Schmidt, l. c. = praec.?

ARACHNOTHRIX, Walp. Ann. ii. 776 (1852) = seq.

ARACHNOTHRYX, Planch. in Fl. des Serres, Sér. I. v. (1849) sub t. 442 = **Rondeletia**, Linn. (Rubiac.).
buddleioides, Planch. l. c. = R. buddleioides.
discolor, Planch. l. c. = R. discolor.
elongata, Planch. l. c. = R. elongata.
hispidula, Griseb. Fl. Brit. W. Ind. 325.—Ind. occ.
laniflora, Planch. in Fl. des Serres, Sér. I. v. (1849) sub t. 442 = R. laniflora.
leucophylla, Planch. l. c. = R. lencophylla.
reflexa, Planch. l. c. = R. reflexa.
rosea, Linden, Hort. Linden. (1859) 3.—N. Granat.

ARACHUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 360 = **Vicia**, Tourn. (Legum.).
vicioides, Medic. l. c. = V. bithynica.

ARACIUM, Neck. Elem. i. 49 (1790) = **Crepis**, Linn. (Compos.).
alpinum, Monn. Ess. Hierac. 73 = C. alpina.
aureum, D. Dietr. Syn. Pl. iv. 1329 = C. aurea.
chondrilloides, D. Dietr. l. c. 1330 = C. Jacquini.
cydoniaefolium, D. Dietr. l. c. = Hieracium cydoniaefolium.
Froelichianum, D. Dietr. l. c. = C. incarnata.
Gmelini, D. Dietr. l. c. = C. Gmelini.
incarnatum, D. Dietr. l. c. 1329 = C. incarnata.
laevigatum, Miq. Fl. Ind. Bat. ii. 111 = C. laevigata.
lyratum, Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) i. 214 = C. lyrata.
mauritianum, D. Dietr. Syn. Pl. iv. 1330 = C. mauritianum.
multicaule, D. Dietr. l. c. = C. multicaulis.
paludosum, Monn. Ess. Hierac. 73 = C. paludosa.
Pancicii, Sch. Bip. in Pollichia, xxii–iv. (1866) 320.—Europ.
praemorsum, D. Dietr. Syn. Pl. iv. 1329 = C. praemorsa.
prenanthoides, D. Dietr. l. c. 1330 = Hieracium prenanthoides.
pubescens, C. Koch, in Linnaea, xxiii. (1850) 681.—As. Min.?
pygmaeum, Miq. Fl. Ind. Bat. ii. 111 = C. nudicaulis.
succisaefolium, D. Dietr. Syn. Pl. iv. 1330 = C. succisaefolia.

ARAEOCOCCUS, Brongn. in Ann. Sc. Nat. Sér. II. xv. (1841) 370. *BROMELIACEAE*, Benth. & Hook. f. iii. 664.
mieranthus, *Brongn. l. c.*—Guiana.

ARAGUS, Steud. Nom. ed. II. i. 118 (1840) = ARAGALLUS, Neck. = **Astragalus**, Tourn. (Legumin.).

ARALIA, Tourn. ex Linn. Syst. ed. I. (1735); Gen. ed. I. 88 (1737). *ARALIACEAE*, Benth. & Hook. f. i. 936.
AURELIANA, Lafit. ex Catesb. Nat. Hist. Carol. App. 16 (1754).
BLAKWELLIA, Scop. Introd. 326 (1777).
DIMORPHANTHUS, Miq. Comm. Phyt. 95 (1840).
GYNAPTEINA, Spach, Hist. Vég. Phan. viii. 113 (1839).
MOTHERWELLIA, F. Muell. Fragm. vii. 107 (1870).
NALAGU, Adans. Fam. ii. 445 (1763).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 336 = Heptapleurum abyssinicum.
acerifolia, Willd. ex Roem. & Schult. Syst. vi. 699 = Oreopanax acerifolium.
aculeata, Ham. ex Seem. in Journ. Bot. vi. (1868) 136 = Brassaiopsis aculeata.
acutifolia, Willd. ex Roem. & Schult. Syst. vi. 697 = Brassaiopsis aculeata.
angularis, Willd. l. c. 698 = Oreopanax angulare.
arborea, Linn. Pl. Jam. Pugill. ii. 11; Syst. ed. X. 967 = Dendropanax arboreum.
arborea, Vell. Fl. Flum. 129; iii. t. 110 = Gilibertia brasiliensis.
argentata, H. B. & K. Nov. Gen. et Sp. v. 4 = Oreopanax argentatum.
argentea, *Linden, Cat.* n. 18. p. 7.—Hab.?
armata, *Seem. in Journ. Bot.* vi. (1868) 134.—Ind. or.
aromatica, Blume, Bijdr. 271 (= *Agalma aromaticum*). —Java.
avicenniaefolia, H. B. & K. Nov. Gen. et Sp. v. 2. t. 2 = Oreopanax avicenniaefolium.
Bastardiana, Decne. Bot. Vénus, 24 = Panax Bastardianum.
bipinnata, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 83 = montana.
bipinnata, Blanco, Fl. Filip. ed. I. 222 = javanica.
bipinnatifida, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 722.—Reg. Himal.
brevifolia, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 74.—Mexic.
caehemiriea, *Decne. in Jacquem. Voy. Bot.* 72. t. 81.—Reg. Himal.
californiea, *S. Wats. in Proc. Am. Acad.* xi. (1876) 144. —Calif.
calyculata, Zoll. & Moritzi, Syst. Verz. Zoll. 41 = Macropanax oreophilum.
canescens, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 94 = spinosa.
capitata, Jacq. Enum. Pl. Carib. 18 = Oreopanax capitatum.
capitulata, *Jungh. & De Vriese, in Ann. Sc. Nat.* Sér. III. vi. (1846) 116.—Sumatra.
catalpaefolia, Willd. ex Roem. & Schult. Syst. vi. 697 = Orcopanax capitatum.
cheirophylla, Spreng. Syst. i. 953 = Orcopanax eheirophyllum.
chinensis, Blume, Bijdr. 870 = dasyphylla.
chinensis, Linn. Sp. Pl. 273 = spinosa.
eissifolia, *Griff. ex Seem. in Journ. Bot.* vi. (1868) 134. —Reg. Himal.
cochleata, Lam. Encyc. i. 224 = Panax eochleatum.
cordata, *Thunb. Fl. Jap.* 127.—Japon.
crassifolia, Soland. ex A. Cunn. in Ann. Nat. Hist. ii. (1839) 214 = Pseudopanax erassifolium.
crassinervia, H. B. & K. Nov. Gen. et Sp. v. 5 = Oreopanax crassinervium.
cumanensis, H. B. & K. l. e. 3 = Oreopanax eumanense.
dasyphylla, *Miq. Fl. Ind. Bat.* i. 1. 751.—Java.
Decaisneana, Hance, in Journ. Bot. iv. (1866) 172 = spinosa.
deeomposita, *Reinw. ex De Vriese, Pl. Ind. Bat. Reinw.* 84.—Java.
digitata, Roxb. Hort. Beng. 22; Fl. Ind. ii. 107 = Heptapleurum venulosum.
digitata, Willd. ex Roem. & Schult. Syst. vi. 701 = Oreopanax xalapense.
discolor, H. B. & K. Nov. Gen. et Sp. v. 4 = Oreopanax discolor.
disperma, Blume, Bijdr. 872 = Maeropanax oreophilum.

ARALIA :—
dubia, Spreng. Syst. iv. Cur. Post. 125 = Trevesia palmata.
Echinops, Cham. & Schlecht. in Linnaea, v. (1880) 409 = Oreopanax Echinops.
edulis, Sieb. & Zucc. Fl. Jap. i. 57. t. 25 = eordata.
elata, Seem. in Journ. Bot. vi. (1868) 134 = spinosa.
elegantissima, *Veitch, in Gard. Chron.* (1873) 782.—Hab.?
erinacea, Hook. in Brewst. Edinb. Journ. Sc. vi. (1827) 64 = Fatsia horrida.
Faribuki, *Sieb. ex Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 16.—Japon.
farinosa, Delile, ex Ferr. & Galin. Voy. Abyss. iii. 135 = Panax pinnatum.
ferox, *Miq. Fl. Ind. Bat.* i. 1. 750.—Malaya.
ferruginea, Willd. ex Roem. & Schult. Syst. vi. 701 = Sciadophyllum ferrugineum.
filieifolia, *Chr. Moore, ex Fourn. in Illustr. Hortic.* xxiii. (1876) 72. t. 240.—Ins. Paeif.
Finlaysoniana, *Seem. in Journ. Bot.* vi. (1868) 134.—Ind. or.
floribunda, H. B. & K. Nov. Gen. et Sp. v. 6. t. 416 = Oreopanax floribundum.
foliolosa, *Seem. in Journ. Bot.* vi. (1868) 134.—Reg. Himal.
fragrans, G. Don. ex Loud. Hort. Brit. 112 = Heteropanax fragrans.
Gemma, *Linden.—Cf. Gard. Chron.* (1880) II. 150.—N. Caled.
georgica, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 8.—Am. bor.
Ghiesbrechtii, Verseh. ex Morr. in Belg. Hortie. xiii. (1863) 3 = Oreopanax xalapense.
Ginseng, *Baill. Hist. Pl.* vii. 197.—China.
glomerulata, Blume, Bijdr. 873 = Brassaiopsis speeiosa.
gracillima, Linden, in Rev. Hortic. (1877) 38 = Veitchii.
grandis, Miq. Comm. Phyt. 95 = elata.
Guilfoylei, *Cogn. & March. Pl. Ornem.* ii. (1874). t. 58.—Ins. Pacif.
heptaphylla, Willd. ex Spreng. Syst. i. 952 = Sciadophyllum ferrugineum.
heterophylla, A. Cunn. ex Hook. Ic. Pl. t. 583–584 = Pseudopanax crassifolium.
heterophylla, Mutis, ex H. B. & K. Nov. Gen. et Sp. v. 1 = Oreopanax Mutisianum.
heterophylla, Willd. ex Roem. & Schult. Syst. vi. 698 = Oreopanax acerifolium.
hispida, *Vent. Jard. Cels*, t. 41; *Michx. Fl. Bor. Am.* i. 185.—Am. bor.
Humboldtiana, Roem. & Schult. Syst. vi. 699 = Oreopanax floribundum.
humilis, *Cav. Ic.* iv. 7. t. 313.—N. Mexic.
hypoleuca, Presl, Epim. Bot. 250 = spinosa.
incisa, Willd. ex Roem. & Schult. Syst. vi. 699 = Oreopanax floribundum.
japonica, Hort. ex Seem. in Journ. Bot. vi. (1868) 135 = Leroana.
japonica, Thunb. Fl. Jap. 128 = Fatsia japonica.
jatrophaefolia, H. B. & K. Nov. Gen. et Sp. v. 6 = Oreopanax jatrophaefolium.
javanica, *Miq. Fl. Ind. Bat.* i. 1. 749.—Java.
Kleinii, Steud. Nom. ed. II. i. 118 = Miquelia Kleinii.
laetevirens, C. Gay, Fl. Chil. iii. 151 (= *Cheirodendron laetevirens*).—Chili.
lappaefolia, *Raeusch. Nom.* ed. III. 88; *et Retz. Obs.* vi. 27.—Ind. or.
Laribuki, *Keiske, ex Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 16.—Japon.
latifolia, Wight & Arn. Prod. 376 = Polyscias pinnata.
Leroana, *C. Koch, Wochenschr.* vii. (1864) 369.—Am. bor.
Lessoni, Hook. f. Fl. Nov. Zel. 96 = Pseudopanax Lessoni.
longifolia, *D. Dietr. Syn. Pl.* ii. 1035.—Java.
longifolia, Reinw. ex App. Hort. Spaarn. Berg. 2 = Brassaia littorea.
lucescens, Blume, Bijdr. 872 (= Heptapleuri sp.).—Java.
lucida, *Hassk. Cat. Hort. Bogor. Alt.* 164.—Malaya.
lutescens, Auct. ex Steud. Nom. ed. II. err. typ. = luecscens.
Lyallii, *T. Kirk, in Trans. N. Z. Inst.* xvii. (1885) 222, 295 = (Pseudopanacis sp.?).—N. Zel.
macrophylla, A. Cunn. in Sweet, Hort. Brit. Suppl. i. 501 = Meryta latifolia.
macrophylla, Lindl. in Bot. Reg. (1844) Mise. 73 = eachemirica.

ARALIA :—
malabarica, *Bedd. Fl. Sylv.* ii. 121.—Ind. or.
mandschurica, Seem. in Journ. Bot. vi. (1868) 134 = spinosa.
mandshurica, *Rupr. in Bull. Acad. Pétersb.* xv. (1856) 134.—As. or.
Maralia, Roem. & Schult. Syst. vi. 704 = Panax Maralia.
Maximowiczii, Van Houtte, Fl. des Serres, Sér. II. xx. (1874) 39. t. 2067 = Acanthopanax ricinifolium.
micans, Willd. ex Roem. & Schult. Syst. vi. 701 = Panax splendens.
Mitsde, Siebold, in Linnaea, xxvi. (1853–55) 754 = Dendropanax japonicum.
monogynia, Vell. Fl. Flum. 129; iii. t. 111 = Gilibertia monogyna.
montana, *Blume, Bijdr.* 870.—Java.
Moorei, F. Muell. Fragm. ii. 108 = Heptapleurum venulosum.
Motherwelli, *F. Muell. l. c.* vii. 107.—Austral.
Muehlenbergiana, Roem. & Schult. Syst. vi. 704 = hispida.
multiflora, Pohl, ex DC. Prod. iv. 262 = Oreopanax capitatum.
Mutisiana, H. B. & K. Nov. Gen. et Sp. v. 1 = Oreopanax Mutisianum.
nobilis, *Hort. Williams.—Cf. Gard. Chron.* (1878) II. 56; (1880) I. 759.—Hab.?
nodosa, Blume, Bijdr. 873 = Polyseias nodosa.
nudicaulis, Blume, l. c. 870 = eordata.
nudicaulis, *Linn. Sp. Pl.* 274.—Am. bor.
nutans, *Franch. & Sav. Enum. Pl. Jap.* ii. 376.—Japon.
obtusiloba, H. B. & K. Nov. Gen. et Sp. v. 5 = Oreopanax obtusilobum.
occidentalis, Schlecht. ex Ledeb. Fl. Ross. ii. 375 = Fatsia horrida.
octonata, Stokes, Bot. Mat. Med. ii. 181 = Heptapleurum octophyllum.
octophylla, Blanco, Fl. Filip. ed. II. 158 = Heptapleurum rigidum.
octophylla, Lour. Fl. Cochineh. i. 187 = Heptapleurum octophyllum.
palmata. Lam. Eneye. i. 224 = Trevesia Zippeliana.
palmata, Lour. Fl. Cochinch. 187 = Brassaiopsis Hainla.
palmata, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 82 = Trevesia sundaica.
palmata, Willd. ex Roem. & Schult. Syst. vi. 699 = Oreopanax eheirophyllum.
paniculata, Phil. in Linnaea, xxviii. (1856) 694 = Panax valdiviense.
papyrifera, Hook. Kew Journ. iv. (1852) 50 = Fatsia papyrifera.
parvifolia, *Panch. & Sebert, Not. Bois. Nouv. Calèd.* 204.—N. Caled.
pendula, Blanco, Fl. Filip. ed. I. 223 = Polyscias nodosa.
pennata, Poir. Eneyc. Suppl. i. 418, sphalm. = Trevesia palmata.
pentaphylla, Thunb. Fl. Jap. 128 = Acanthopanax spinosum.
pergamacea, Blume, Bijdr. 873 = Heptapleurum pergamaeeum.
pinnata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 335.—Abyss.
pinnatifida, Jungh. & De Vriese, in Ann. Sc. Nat. Sér. III. vi. (1846) 115 = Aralidium pinnatifidum.
Planchoniana, Hanee, in Journ. Bot. iv. (1866) 172 = Acanthopanax spinosum.
platanifolia, H. B. & K. Nov. Gen. et Sp. v. 6 = Oreopanax platanifolium.
polaris, Hombr. & Jaequinot, ex Decne. Bot. Voy. Astrol. et Zel. 55 = Stilbocarpa polaris.
Polyscias, Spreng. ex Seem. in Journ. Bot. vi. (1868) 138 = Polyscias umbellata.
Pseudo-ginseng, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 721.—Reg. Himal.
pubescens, DC. Cat. Hort. Monsp. (1813) 80 = humilis?
quinduensis, H. B. & K. Nov. Gen. et Sp. v. 8 = Sciodaphyllum quinduense.
quinquefolia, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 105.—Am. bor.
racemosa, *Linn. Sp. Pl.* 273.—Am. bor.
ramiflora, Pohl, ex March. in Mart. Fl. Bras. xi. 1. 250 = Gilibertia cuneata.
Regeliana, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 73.—Mexico.
Reinwardtiana, Seem. in Journ. Bot. vi. (1868) 138 = Trevesia sundaica.

ARALIA:—
reticulata, Willd. ex Roem. & Schult. Syst. vi. 699 = Oreopanax reticulata.
rigida, Blume, Bijdr. 874 = Heptapleurum rigidum.
rugosa, Blume, l. c. 872 (= Heptapleuri sp.).—Java.
salicifolia, *Vent. ex Seem. in Journ. Bot.* vi. (1868). 138.—Hab.?
scabra, Presl, ex DC. Prod. iv. 258 = pubescens.
scandens, Poir. Encyc. Suppl. i. 419 = Brassaiopsis Hainla.
Schefflera, Spreng. Pl. Pugill. i. 28 = Schefflera digitata.
Sciodaphyllum, Sw. Prod. Veg. Ind. Occ. 55 = Sciodaphyllum Brownei.
septemnervia, H. B. & K. Nov. Gen. et Sp. v. 3 = Oreopanax capitatum.
Sieboldii, Hort. ex C. Koch, Dendrol. i. 678 = Fatsia japonica.
simillima, Blume, Bijdr. 871 (= Heptapleuri sp.).—Java.
soratensis, *March. in Bull. Acad. Belg.* Sér. II. xlvii. (1879) 75.—Bolivia.
spectabilis, Hort. Bull.—Cf. Gard. Chron. (1876) i. 603 = filieifolia.
spinosa, *Linn. Sp. Pl.* 273.—Am. bor.
tarchonanthifolia, Willd. ex Roem. & Schult. Syst. vi. 698 = Oreopanax avicenniaefolium.
tenuifolia, *Pancher, ex Baill. Adansonia*, x. (1871-72) 372.—N. Caled.
Thibautii, *Versch. ex Morr. in Belg. Hortic.* xiii. (1863). —Mexic.
Thomsonii, *Seem. in Journ. Bot.* vi. (1868) 134.—Reg. Himal.
trifolia, Banks & Soland. ex Walp. Ann. v. 84 = Pseudopanax crassifolium.
trifolia, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 105.—Am. bor.
trifoliata, Meyen, Reise, ii. 332 = Acanthopanax aculeatum.
trigyna, Gaudich. Voy. t. 98 (= Panacis sp.).—Hab.?
triphylla, Poir. Encyc. Suppl. i. 418 = trifolia.
tripinnata, Blanco, Fl. Filip. ed I. 223 = Panax fruticosum.
turbacensis, H. B. & K. Nov. Gen. et Sp. v. 3 = Oreopanax turbacense.
umbellata, Pav. ex Seem. in Journ. Bot. vi. (1868) 138 = Gilibertia umbellata.
umbellata, Pohl, ex March. in Mart. Fl. Bras. xi. i. 250 = Gilibertia cuneata.
umbellifera, Lam. Encyc. i. 225 = Osmoxylon amboinense.
umbraculifera, Roxb. Hort. Beng. 22; Fl. Ind. iii. 108 = Polyscias nodosa.
urticaefolia, *Blume, ex Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 9.—Hab.?
valdiviensis, C. Gay, Fl. Chil. iii. 152 = Panax valdiviense.
Veitchii, *Hort. ex T. Moore, in Flor. & Pom.* (1874) 5. —Hab.?
vitiensis, A. Gray, Bot. U. St. Expl. Exped. i. 715 = Schefflera vitiensis.
Wallichiana, *D. Dietr. Syn. Pl.* ii. 1035.—Ind. or.
xalapensis, H. B. & K. Nov. Gen. et Sp. v. 8. = Oreopanax xalapense.

ARALIDIUM, Miq. Fl. Ind. Bat. i. i. 762. t. 13 (1855). *ARALIACEAE*, Benth. & Hook. f. i. 936.
dentatum, *Miq. Fl. Ind. Bat. Suppl.* 340.—Sumatra.
pinnatifidum, *Miq. l. c.* i. i. 763. t. 13.—Malacca.

ARALIOPSIS, Kurz, in Andam. Rep. 39 (1870) = **Brassaiopsis**, Decne. & Planch. (Araliac.).
andamanica, Kurz, l. c. = B. palmata.

ARAPABACA, Adans. Fam. ii. 225 (1763) = **Spigelia**, Linn. (Logan.).

ARARIBA, Mart. in Sitzb. Acad. Muench. (1860) 323 = **Sickingia**, Willd. (Rubiae.).
alba, Mart. l. c. 324.—Bras.
rubra, Mart. l. e.—Bras.

ARAROCARPUS, Scheff. in Ann. Jard. Buitenz. ii. (1885) 10. *ANONACEAE*.
velutinus, *Scheff. l. c.*—Java.

ARASCHCOOLIA, Sch. Bip. Pl. Kotsch. Nub. Exs. n. 101 (1873); ex Benth. & Hook. f. Gen. ii. 337 = **Geigeria**, Griessel. (Compos.).
acaulis, Sch. Bip. ex Oliver, Fl. Trop. Afr. iii. 368 = G. aeaulis.

ARAUCARIA, Juss. Gen. Pl. 413 (1789). *CONIFERAE*, Benth. & Hook. f. iii. 437.
ALTINGIA, Loud. Hort. Brit. 403 (1830).
COLUMBEA, Salisb. in Trans. Linn. Soc. viii. (1807) 317.
DOMBEYA, Lam. Encyc. ii. 301. t. 828 (1786).
EUTACTA, Link, in Linnaea. xv. (1841) 543.
EUTASSA, Salisb. in Trans. Linn. Soc. viii. (1807) 316.
QUADRIFARIA, Manetti, ex Gord. Pinet. Suppl. 14 (1862).
albospica, Hort. ex Carr. Conif. ed. II. 611 = excelsa.
araucana, C. Koch, Dendrol. ii. ii. 206 = imbricata.
Balansae, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. xiii. (1870–71) 351.—N. Caled.
Bibbiani, Hort. Ital. ex Gord. Pinet. 24 = brasiliana.
Bidwillii, *Hook. Lond. Journ. Bot.* ii. (1843) 503. t. 18, 19.—Austral.
brasiliana, *A. Rich. in Dict. Class. Hist. Nat.* i. 512.—Bras.
brasiliensis, Loud. Hort. Brit. 403 = praec.
chilensis, Mirb. in Mém. Mus. Par. xiii. (1825) 49 = imbricata.
columnaris, Hook. in Bot. Mag. t. 4635 = Cookii.
Cookii, *R. Br. ex D. Don, in Trans. Linn. Soc.* xviii. (1841) 164.—N. Caled.
Cunninghamii, *Sweet, Hort. Brit.* ed. II. 475.—Austral.
Dombeyi, A. Rich. Conif. 86. t. 20 = imbricata.
elegans, Hort. ex Carr. Conif. ed. I. 415 = brasiliana.
excelsa, *R. Br. in Ait. Hort. Kew.* ed. II. v. 412.—Austral.; Ins. Norf.
glauca, Antoine, Conif. 105. t. 44 = Cunninghami.
Goldieana, *Hort. ex T. Moore, Flor. & Pom.* (1877) 39. —N. Caled.
gracilis, Hort. ex Carr. Conif. ed. I. 415 = brasiliana.
imbricata, *Pav. in Mem. Acad. Med. Madr.* i. (1797) 199.—Chili.
intermedia, *R. Br. ex Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 54.—N. Caled.
lanceolata, Hort. ex Carr. Conif. ed. I. 171 = Cunninghamia sinensis.
Lindleyana, Van Houtte, ex Gord. Pinet. 24 = brasiliana.
montana, *Brongn. & Gris, in Ann. Sc. Nat.* Sér. V. xiii. (1870–71) 358.—N. Caled.
Muelleri, *Brongn. & Gris, l. c.* 362.—N. Caled.
Ridolfiana, Savi f. in Atti Firenz. (1841) 458 et 783; et Giorn. Bot. Ital. ii. (1846) fasc. i. 52 = brasiliana.
Rulei, *F. Muell. ex Lindl. in Gard. Chron.* (1861) 868. —N. Caled.
Saviana, *Parl. in Bull. Soc. Bot. Fr.* viii. (1861) 89.—Bolivia.
subulata, *Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 55.—N. Caled.

ARAUCASIA, Benth. & Hook. f. Gen. iii. 919 (1883), *sphalm.* = ARAUSIACA, Blume = **Orania**, Zipp. (Palm.).

ARAUJIA, Brot. in Trans. Linn. Soc. xii. (1818) 62. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 751.
LAGENIA, Fourn. in Mart. Fl. Bras. vi. iv. 293 (1885).
PENTAPHRAGMA, Zucc. ex Reichb. Consp. 131 (1828).
PHYSIANTHUS, Mart. Nov. Gen. et Sp. i. 53. t. 32 (1824).
SCHUBERTIA, Mart. l. c. 55. t. 33 (1824).
albens, G. Don, Gen. Syst. iv. 149 = sericifera.
angustifolia, *Steud. Nom.* ed. II. i. 119.—Bras.
calycina, Decne. in DC. Prod. viii. 534 = sericifera.
fusca, Griseb. ex Lorentz, in Goett. Abh. xxiv. (1879) 232 = megapotamica.
hortorum, *Fourn. in Mart. Fl. Bras.* vi. iv. 293.—Bras.
megapotamica, *G. Don, Gen. Syst.* iv. 149.—Bras.
sericifera, *Brot. in Trans. Linn. Soc.* xii. (1818) 62.—Peruv.
subhastata, *Fourn. in Mart. Fl. Bras.* vi. iv. 293.—Bras.
undulata, Visiani, Hort. Patav. (1832) 134 = hortorum.

ARAUSIACA, Blume, Rumphia, in praef. p. 8. (1836) tt. 119, 122 = **Orania**, Zipp. (Palm.).
excelsa, Blume, l. c. = O. regalis.

ARBUTUS, [Tourn.] Linn. Syst. ed. I (1735). *ERICACEAE*, Benth. & Hook. f. ii. 581.
UNEDO, Hoffmgg. & Link, Fl. Port. i. 415 (1809).
acadiensis, Linn. Sp. Pl. 395 (planta obscura).—Am. bor.

ARBUTUS:—
acerba, Gilib. Fl. Lituan. i. 5 = Arctostaphylos Uva-ursi.
alpina, Linn. Sp. Pl. 395 = Arctostaphylos alpina.
Andrachne, *Linn. Syst.* ed. X. 1024. n. 488.—Reg. Mediterr. or.
andrachnoides ×, *Link, Enum. Hort. Berol.* i. 395.—Hab.?
arguta, Zucc. ex Steud. Nom. ed. II. i. 120, 119 = Arctostaphylos arguta.
buxifolia, Stokes, Bot. Mat. Med. ii. 509 = Arctostaphylos Uva-ursi.
Callicarpa, Bueh, Canar. 146 = canariensis.
canadensis, *Poir. ex Steud. Nom.* ed. I. 62.—Canada.
canariensis, *Duham. Arb.* ed. II. i. 80.—Ins. Canar.
cassinifolia, *Hort. ex Steud. Nom.* ed. II. i. 119.—Ins. Madera.
coriacea, Blume, Cat. Gew. Buitenz. 67 = Vaccinium Korthalsii.
crispa, Hoffmgg. Verz. Pfl. Nachtr. i. (1824) 204 = A. Unedo.
Croomii, Hort.—Cf. Gard. Chron. (1884) ii. 492 = A. Unedo.
densiflora, Benth. Pl. Hartw. 15 = glandulosa.
densiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 280. t. 260. —Mexic.
discolor, Hook. Ic. Pl. t. 29 = Arctostaphylos arguta.
empetrifolia, Pers. Syn. i. 483 = Pernettya empetrifolia.
ferruginea, *Linn. f. Suppl.* 238.—Reg. Colomb.
filiformis, Lam. Encyc. i. 228 = Chiogenes serpyllifolia.
floribunda, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 534.—Mexic.
furiens, *Hook. Bot. Beech. Voy.* 33.—Chili.
glaberrima, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 6.—Malaya.
glandulosa, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 533.—Mexic.
herpetica, Coleb. ex Roxb. Hort. Beng. [90]; Fl. Ind. ii. 412 = Pieris ovalifolia.
hybrida, Ker-Gawl. in Bot. Reg. t. 619 = andrachnoides.
integrifolia, Lam. Encyc. i. 227 = A. Andrachne.
integrifolia, Sieber, ex Klotzsch, in Linnaea, xxiv. (1851) 71 = Sieberi.
integrifolia, Sims, Bot. Mag. t. 2319 = A. Unedo.
intermedia, Heldr. ex Nym. Consp. 490 = A. Unedo.
lanceolata, Lam. Encyc. i. 227 = Prunus caroliniana.
laurifolia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 151 = Gaultheria fragrantissima.
laurifolia, Linn. f. Suppl. 238 (Cf. A. Gray, Syn. Fl. N. Am. ii. i. 396) = A. Unedo.
laurifolia, Lindl. Bot. Reg. (1839) t. 67 = xalapensis.
laurina, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 535.—Mexic.
leucocarpa, Poepp. & Kunze, ex DC. Prod. vii. 586 = Pernettya leucocarpa.
longifolia, Andr. Bot. Rep. x. 664 = canariensis.
lucida, Hort. ex Steud. Nom. ed. II. i. 119 = A. Andrachne.
macrophylla, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 534.—Mexic.
Madrono, Humb. Nouv. Espagne, ii. 454, ex H. B. & K. Nov. Gen. et Sp. iii. 282 = Arctostaphylos pungens.
Menziesii, *Pursh, Fl. Bor. Am.* i. 282.—Am. bor.; Mexic.
microphylla, Forst. f. in Comm. Gotting. ix. (1789) 32 = Pernettya microphylla.
microphylla, Hort.—Cf. Gard. Chron. (1884) ii. 49 = A. Unedo.
Milleri, *Sweet, Hort. Brit.* ed. II. 331.—Oriens.
mollis, *H. B. & K. Nov. Gen. et Sp.* iii. 281. t. 260.—Mexic.
mucronata, Linn. f. Suppl. 239 = Pernettya mucronata.
myrtifolia, Willd. ex Steud. Nom. ed. II. i. 120, 119 = Arctostaphylos pungens.
myrtilloides, Humb. ex Steud. Nom. ed. II. i. 119 = Vaccinium stamineum.
myrtilloides, Poepp. ex DC. Prod. vii. 593 = Pernettya myrtilloides.
nuda, Steud. Nom. ed. II. i. 119 = Arctostaphylos pungens.
obtusifolia, *Rafin. Fl. Ludov.* 55.—Am. bor.
officinalis, Boiss. Fl. Orient. iii. 967 = Arctostaphylos Uva-ursi.
ovata, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 533.—Mexic.
paniculata, *Mart. & Gal. l. c.* 535.—Mexic.

ARBUTUS:—
petiolaris, *H. B. & K. Nov. Gen. et Sp.* iii. 281.—Mexic.
phyllireaefolia, Pers. Syn. i. 483 = Pernettya phyllircaefolia.
pilosa, R. Grah. in Bot. Mag. t. 3177 = Pernettya pilosa.
procera, Dougl. ex Lindl. Bot. Reg. t. 1753 = Menziesii.
procera, Soland. ex DC. Prod. vii. 581 = canariensis.
procumbens, *Kluk, ex Bess. Fl. Galic.* ii. 355.—Galieia.
procumbens, Salisb. Prod. 289 = Arctostaphylos Uva-ursi.
prostrata, Cav. Ic. vi. 43. t. 562 = Pernettya parvifolia.
prunifolia, *Klotzsch, in Linnaea*, xxiv. (1851) 73.—Mexic.
pumila, Forst. f. in Comm. Gotting. ix. (1789) 32 = Pernettya empetrifolia.
punctata, *Hook. & Arn. Bot. Beech. Voy.* 33.—Calif.
pungens, Hook. & Arn. l. c. 144 = Aretostaphylos Hookeri.
rigida, Banks, ex Hook. f. Fl. Antaret. ii. 326 = Pernettya mucronata.
rigida, Bert. ex Colla, in Mem. Aec. Sc. Torin. xxxviii. (1835) 120 = Pernettya rigida.
rigida, Willd. ex Steud. Nom. ed. II. i. 120, 119 = Arctostaphylos pungens.
rubescens, Bertol. Fl. Guatimal. 20. t. 7 = Arctostaphylos rubescens.
salicifolia, *Cels, ex Hoffmgg. Verz. Pfl. Nachtr.* i. (1824) 204.—Oriens.
serpyllifolia, Lam. Encyc. i. 225 = Gaultheria antipoda.
serratifolia, Lodd. Bot. Cab. t. 580 = A. Andraehne.
serratifolia, Salisb. Prod. 288 = A. Unedo.
sibiriea, *Forbes, Hort. Woburn.* 95, nomen.—Sibir.
Sieberi, Klotzsch, in Linnaea, xxiv. (1851) 71 = A. Andrachne.
spinulosa, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 532.—Mexic.
texana, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 460 = Menziesii.
thymifolia, Ait. Hort. Kew. ii. 72 = Chiogenes serpyllifolia.
tomentosa, Pursh, Fl. Am. Sept. i. 282 = Aretostaphylos pungens.
turbinata, Pers. ex Reichb. Fl. Germ. Excurs. i. 416 = A. Unedo.
Unedo, *Linn. Sp. Pl.* 395.—Reg. Mediterr.; Hibern.
Uva-ursi, Linn. l. c. = Arctostaphylos Uva-ursi.
varians, *Benth. Pl. Hartw.* 77.—Mexic.
vernalis, Poepp. & Endl. Nov. Gen. i. 26. t. 42 = Gaultheria vernalis.
villosa, Willd. ex Klotzsch, in Linnaea, xxiv. (1851) 72 = glandulosa.
xalapensis, Andr. ex Hook. Ic. Pl. t. 27 = Arctostaphylos oaxacana.
xalapensis, *H. B. & K. Nov. Gen. et Sp.* iii. 279.—Mexic.
xanthocarpa, *Wangen. Beitr.* 30.—Am. bor.

ARCANGELISIA, Bece. Malesia, i. (1877) 145. *MENISPERMACEAE.*
inclyta, *Becc. l. c.* 147.—Java; Celebes.
lemniscata, *Becc. l. c.*—N. Guin.

ARCAULA, Rafin. Alsog. Am. 30 (1838) = **Quercus**, Tourn. (Cupul.).
spicata, Rafin. l. c. = Q. spicata.

ARCEUTHOBIUM, Bieb. Fl. Taur. Canc. Suppl. 629 (1819). *LORANTHACEAE*, Benth. & Hook. f. iii. 213.
RAZOUMOWSKIA, Hoffm. ex Bieb. l. c. (1819).
abietinum, Engelm. in Proc. Am. Acad. viii. (1873) 401 = occidentale.
americanum, Nutt. ex A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 214 = A. Oxyeedri.
campylopodum, *A. Gray, l. c.*—Am. bor.
capense, J. S. Presl, Rostl. ii. 28 = Viscum capense.
cryptopodum, A. Gray, in Bost. Journ. Nat. Hist. vi. (1850) 214 = vaginatum.
divaricatum, *Engelm. in Wheeler, Rep.* vi. 253.—Am. bor.
Douglasii, *Engelm. l. c.*—Am. bor.

ARCEUTHOBIUM:—
gracile, *Engelm. in A. Gray, Pl. Fendl.* 59.—Am. bor.
minutum, *Engelm. in Bull. Torrey Bot. Club*, ii. (1871) 43, 47.—Am. bor.
occidentale, *Engelm. ex S. Wats. in Bot. Calif.* ii. 107.—Calif.
opuntioides, J. Presl, Rostl. ii. 28 = Dendrophthora opuntioides.
Oxycedri, *Bieb. Fl. Taur. Canc. Suppl.* 629.—Europ.; Am. bor.
robustum, *Engelm. in A. Gray, Pl. Fendl.* 59.—Am. bor.
vaginatum, *J. Presl, Rostl.* ii. 28.—Am. bor.

ARCEUTHOBIUM, Griseb. Fl. Brit. W. Ind. 314 (1864) = **Dendrophthora**, Eichl. (Loranth.).
Bonaniae, Griseb. Cat. Pl. Cub. 121 = D. Bonaniae.
cupressoides, Griseb. Fl. Brit. W. Ind. 315 = D. cupressoides.
domingense, Griseb. Cat. Pl. Cub. 121 = D. domingensis.
Epiviscum, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 192 = D. Epiviscum.
glaucum, Griseb. Fl. Brit. W. Ind. 315 = D. glaucum.
gracile, Griseb. l. c. = D. gracilis.
Mancinellae, Wright, ex Griseb. Cat. Pl. Cub. 121 = D. Mancinellae.

ARCEUTHOS, Ant. & Kotschy, in Oestr. Bot. Wochenbl. (1854) 249 = **Juniperus**, Linn. (Conif.).
drupacea, Ant. & Kotschy, l. c. = J. drupacea.

ARCHANGELICA, Hoffm. Gen. Umb. ed. I. 161 (1814); ed. II. 166 (1816). *UMBELLIFERAE*, Benth. & Hook. f. i. 917.
COELOPLEURUM, Ledeb. Fl. Ross. ii. 361 (1844).
CZERNIAJEVIA, Turcz. in Bull. Soc. Nat. Mosc. (1838) 93.
atropurpurea, *Hoffm. Gen. Umb.* 162.—Am. bor.
brevicaulis, *Reichb. f. in Journ. Bot.* xiv. (1876) 46.—As. temp.
commutata, *C. A. Mey. ex Rupr. in Beitr. Pfl. Russ. Reich.* xi. 13.—Kamtschatka.
decurrens, Ledeb. Fl. Alt. i. 316 = officinalis.
dentata, *Chapm. in Torr. & Gray, Fl. N. Am.* i. 622.—Florida.
discoearpa, *Fries, in Bot. Notiser*, (1857) 50.—Europ.
Gmelini, *DC. Prod.* iv. 170.—Am. bor.; As. bor. oce.
hirsuta, *Torr. & Gray, Fl. N. Am.* i. 622.—Am. bor.
inaequalis, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* i. 188.—Japon.
Keiskei, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 62.—Japon.
littoralis, *Agardh, ex DC. Prod.* iv. 170.—Europ. aret.
major, Lag. Gen. et Sp. Nov. 13 = littoralis?
norwegica, Rupr. in Beitr. Pfl. Russ. Reich. ii. 37 = officinalis.
officinalis, *Hoffm. Gen. Umb.* 162.—Europ.; Sibir.; Reg. Himal.
officinalis, Hook. Fl. Bor. Am. i. 267 = Gmelini.
peregrina, Nutt. ex Torr. & Gray, Fl. N. Am. i. 622 = Gmelini.
Roylei, Lindl. in Royle, Illustr. Bot. Himal. 232 = officinalis.
sativa, Bess. Enum. 13 = officinalis.
slavica, Reuss, Květ. Slov. (1853) 184 = officinalis.
spuria, Wahlenb. Fl. Casp. 84, in obs. = littoralis.

ARCHEMORA, DC. Prod. iv. 188 (1830) = **Peucedanum**, Linn. (Umbell.).
ambigua, DC. l. c. = P. ternatum.
denticulata, DC. l. c. = P. ternatum.
Fendleri, A. Gray, Pl. Fendl. 56.—Texas.
rigida, DC. Mém. Ombell. 52; Prod. iv. 188 = P. ternatum.
ternata, Nutt. ex Torr. & Gray, Fl. N. Am. i. 631 = P. ternatum.
tricuspidata, DC. Prod. iv. 188 = P. ternatum.

ARCHERIA, Hook. f. Fl. Tasm. i. 262. t. 80, 81. *EPACRIDEAE*, Benth. & Hook. f. ii. 616.
eriocarpa, *Hook. f. l. c.* 263. t. 80 B.—Tasmania.
hirtella, *Hook. f. l. c.* t. 81.—Tasmania.
minor, Hook. f. l. c. 264 = serpyllifolia.

ARCHERIA:—
racemosa, *Hook. f. Handb. New Zeal. Fl.* 180.—N. Zel.
serpyllifolia, *Hook. f. Fl. Tasm.* i. 263. t. 80 B.—Tasmania.
Traversii, *Hook. f. Handb. New Zeal. Fl.* 180.—N. Zel.

ARCHIDENDRON, F. Muell. Fragm. v. 59 (1865). *LEGUMINOSAE*, Benth. & Hook. f. i. 1004.
Lucyi, *F. Muell. l. c.* vi. 201.—Austral.
Vaillantii, *F. Muell. l. c.* v. 60.—Austral.

ARCHIMEDEA, Leandr. Sacram. ex A. St. Hil. in Ann. Sc. Nat. Sér. II. vii. (1837) 32 = **Lophophytum**, Schott & Endl. (Balanoph.).
pyramidalis, Leandr. Sacram. l. c. = L. mirabile.

ARCHITAEA, Mart. Nov. Gen. et Sp. i. 116. t. 73 (1824) = **Archytaea**, Mart. (Ternstr.).

ARCHONTOPHOENIX, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 182. *PALMAE*, Benth. & Hook. f. iii. 889.
Alexandrae, *H. Wendl. & Drude, l. c.* 212.—Austral.
Cunninghamii, *H. Wendl. & Drude, l. c.* 214.—Austral.
Veitchii, *H. Wendl. & Drude, l. c.*—Hab.?

ARCHYTAEA, Mart. Nov. Gen. et Sp. i. 117 (1824). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 188.
PLOIARUM, Korth. Verh. Nat. Gesch. Bot. 135. t. 25 (1842).
multiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 363.—Guiana.
sessilis, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 406.—Malaya.
triflora, *Mart. Nov. Gen. et Sp.* i. 117.—Bras.
Vahlii, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 163.—Malaya.

ARCTAGROSTIS, Griseb. in Ledeb. Fl. Ross. iv. 434 (1853). *GRAMINEAE*, Benth. & Hook. f. iii. 1150.
latifolia, *Griseb. l. c.*—Reg. Arct.

ARCTIO (et ARCTION), Lam. Encyc. i. 235 (1783) = **Berardia**, Vill. (Compos.).
lanuginosa, Lam. l. c. = B. subacaulis.

ARCTIODRACON, A. Gray, in Mem. Am. Acad. Ser. II. vi. (1858–9) 408 = **Lysichitum**, Schott (Aroid.).
camtschaticum, A. Gray, l. c. 409 = L. camtschatense.
japonicum, A. Gray, l. c. = L. camtschatense.

ARCTIUM, Lam. Fl. Fr. ii. 70 (1778) = **Berardia**, Vill. (Compos.).
lanuginosum, Lam. l. c. = B. subacaulis.

ARCTIUM, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 466.
LAPPA, Juss. Gen. 173 (1789).
adhaerens, Gilib. Fl. Lituan. i. 182 = A. Lappa.
Bardana, Willd. Sp. Pl. iii. 1632 = A. Lappa.
Carduelis, Linn. Mant. i. 108 = Carduus arctioides.
conglomeratum, Schur, ex Nym. Syll. Suppl. 4 = minus.
euminus, Syme, Engl. Bot. ed. III. v. 26 = minus.
grandiflorum, Desf. Cat. Hort. Par. ed. III. 154 = majus.
intermedium, Bab. in Ann. & Mag. Nat. Hist. Ser. II. xvii. (1856) 374 = majus.
intermedium, Lange, Fl. Dan. t. 2663 = nemorosum.
Lappa, Kalm, Fl. Fennic. n. 449; Sm. Engl. Bot. t. 1228 = minus.
Lappa, Willd. Sp. Pl. iii. 1631 = majus.
Lappa, *Linn. Sp. Pl.* 816 (= majus, minus, &c.).—Europ.
majus, *Bernh. Syst. Verz. Erf.* 154.—Europ.
minus, *Bernh. l. c.*—Europ.
montanum, Schweig. ex Steud. Nom. ed. II. i. 120 = minus.
nemorosum, *Lejeune, in Reichb. Ic. Fl. Germ.* xv. 81.—Europ.
Personata, Linn. Sp. Pl. 816 = Carduus Personata.

ARCTIUM:—
Personata, Pall. Reise, iii. 439 (planta ignota).—Davuria.
pubens, Bab. in Ann. & Mag. Nat. Hist. Ser. II. xvii. (1856) 376 = nemorosum.
ruderale, Salisb. Prod. 184 = A. Lappa.
tomentosum, Mill. Gard. Dict. ed. VIII. n. 3; Bernh. Syst. Verz. Erf. 154 = majus.

ARCTOCALYX, Fenzl, in Otto & Dietr. Allg. Gartenz. xvi. 305 (1848) = **Solenophora**, Benth. (Gesner.).
Endlicherianus, C. Heller, ex Fenzl, l. c. 307 = S. Endlicheriana.
insignis, Fenzl, l. c. 306 = S. insignis.

ARCTOCARPUS, Blanco, Fl. Filip. ed. I. 666 (1837) = **Artocarpus**, Forst. (Urtic.).

ARCTOGERON, DC. Prod. v. 260 (1836) = **Aster**, Linn. (Compos.).
gramineum, DC. l. c. 261.—Sibir.

ARCTOMECON, Torr. & Frém. in Frém. Rep. 312. t. 2 (1845). *PAPAVERACEAE*, Benth. & Hook. f. i. 52.
californicum, *Torr. & Frém. l. c.*—Calif.

ARCTOPHILA, Rupr. in Beitr. Pfl. Russ. Reich. ii. 62 (1845) = **Poa**, Linn. (Gram.).
fulva, Nym. Consp. 830 = P. fulva.
Laestadii, Rupr. in Beitr. Pfl. Russ. Reich. ii. 62 = P. Lacstadii.
pendulina, Nym. Consp. 830 = P. Laestadii.

ARCTOPUS, Linn. Coroll. Gen. 21 (1737); Hort. Cliff. 495 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 879.
Apradus, Adans. Fam. ii. 102, 509 (1763).
Dregei, *Sond. in Harv. & Sond. Fl. Cap.* ii. 565.—Afr. austr.
echinatus, *Linn. Sp. Pl.* 1058.—Afr. austr.
monacanthus, *Carmich. ex Harv. & Sond. Fl. Cap.* ii. 565.—Afr. austr.

ARCTOSTAPHYLOS, Adans. Fam. ii. 165 (1763). *ERICACEAE*, Benth. & Hook. f. ii. 581.
Comarostaphylis, Zucc. in Abh. Akad. Muench. ii. (1837) 331.
Daphnidostaphylis, Klotzsch, in Linnaea, xxiv. (1851) 78.
Mairania, Neck. Elem. i. 219 (1790).
Uva-ursi, Tourn. ex Moench, Meth. 470 (1794).
Xerobotrys, Nutt. in Trans. Am. Phil. Scr. II. viii (1843) 267.
Xylococcus, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1843) 258.
acuta, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1843) 258 = Hookeri?
alpina, Payot.—Cf. Bull. Soc. Bot. Fr. xxix. (1882) Bibl. 43, lapsu = *angustifolia*.
alpina, *Spreng. Syst.* ii. 287.—Am. bor. et arct.
Andersonii, *A. Gray, in Proc. Am. Acad.* ix. (1876) 83.—Calif.
angustifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 278.—Mexic.
angustifolia, Payot, Fl. du Mont Blanc, Phan. 174 = A. Uva-ursi.
arbutoides, *Hemsl. Biol. Centr. Am. Bot.* ii. 278.—Guatemala.
arguta, *Zucc. in Linnaea*, xii. (1838) *Litt.* 86; *DC. Prod.* vii. 585.—Mexic.
attenuata, *Hemsl. Biol. Centr. Am. Bot.* ii. 278.—Mexic.
bicolor, *A. Gray, in Proc. Am. Acad.* vii. (1868) 366.—Calif.
Clevelandi, A. Gray, l. c. xii. (1877) 61 = bicolor.
cordifolia, Lindl. Bot. Reg. sub t. 1791 = tomentosa.
discolor, DC. Prod. vii. 585 = arguta.
diversifolia, *Parry, ex A. Gray, Syn. Fl. N. Am.* ii. 1. 397.—Calif.
glauca, *Lindl. Bot. Reg.* sub t. 1791.—Calif.
glauca, S. Wats. in Bot. King, Exped. 210 = pungens.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* iii. 278.—Mexic.
Hartwegiana, *Hemsl. Biol. Centr. Am. Bot.* ii. 278.—Mexic.
Hookeri, *G. Don, Gen. Syst.* iii. 836.—Calif.
latifolia, Walp. Rep. ii. 726 = seq.

ARCTOSTAPHYLOS:—
ledifolia, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 536.—Mexic.
longiiolia, *Benth. Pl. Hartw.* 44.—Mexic.
mucronata, *Hemsl. Biol. Centr. Am. Bot.* ii. 279.—Mexic.
mucronifera, *DC. Prod.* vii. 585.—Mexic.
nevadensis, *A. Gray, Syn. Fl. N. Am.* ii. 1. 27.—Calif.
nitida, Benth. Pl. Hartw. 66 = arguta.
nummularia, *A. Gray, in Proc. Am. Acad.* vii. (1868) 366.—Calif.
oaxacana, Benth. Pl. Hartw. 45 = Hartwegiana.
oaxacana, *DC. Prod.* vii. 585.—Mexic.
officinalis, Wimm. & Grab. Fl. Siles. i. 391 = A. Uva-ursi.
oppositifolia, *Parry, in Proc. Davenport Acad. Nat. Sc.* iv. (1884) 37.—Calif.
polifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 277. t. 258.—Mexic.
polifolia, Torr. Bot. Mex. Bound. 108 = oppositifolia.
procera, Lindl. ex Pritz. Ind. Ic. 90, sphalm. = tomentosa.
procumbens, Patze, Mey. & Elk. Fl. Preuss. 188 = A. Uva-ursi.
pumila, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. viii. (1843) 266.—Calif.
pungens, A. Gray, in Bot. Calif. i. 453, partim = Hookeri.
pungens, *H. B. & K. Nov. Gen. et Sp.* iii. 278. t. 259.—Calif.; Mexic.
rubescens, *Hemsl. Biol. Centr. Am. Bot.* ii. 279.—Guatemala.
salicifolia, Parry, in Proc. Davenport Acad. iv. (1884) 37 = oppositifolia.
spinulosa, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 537 = arguta.
tomentosa, *Lindl. Bot. Reg.* t. 1791.—Calif.
Uva-ursi, *Spreng. Syst.* ii. 287.—Reg. bor. et arct.
Veatchii, Kellogg, in Proc. Calif. Acad. ii. (1863) 19 = bicolor.

ARCTOTHECA, Wendl. Beobacht. 41; et Hort. Herrenh. 8. t. 6 (1798). *COMPOSITAE*, Benth. & Hook. f. ii. 458.
grandiflora, Drège, ex Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 93 = Arctotis Dregei.
grandiflora, Schrad. in Ind. Sem. Hort. Goett. (1832) 1 = repens.
hirta, *Schrad. ex Steud. Nom.* ed. II. i. 120.—Afr. austr.
repens, *Wendl. Beobacht.* 41; *et Hort. Herrenh.* 8. t. 6.—Afr. austr.

ARCTOTIS, Linn. Gen. ed. I. 261 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 458.
Anemonospermos, Adans. Fam. ii. 127 (1763).
Lycotis, Hoffmgg. Verz. Pfl. Nachtr. ii. 99 (1826).
Odontoptera, Cass. in Dict. Sc. Nat. xxxv. 396 (1825).
Steganotus, Cass. l. c. (1825).
abrotanifolia, Steud. Nom. ed. II. i. 120 = Ursinia abrotanifolia.
acaulis, E. Mey. ex DC. Prod. vi. 486 = canescens.
acaulis, Jacq. Hort. Schoenb. ii. 16. t. 158 = adpressa.
acaulis, *Linn. Sp. Pl.* ed. II. 1306.—Afr. austr.
adpressa, *DC. Prod.* vi. 485.—Afr. austr.
adpressa, Eckl. ex Harv. & Sond. Fl. Cap. iii. 465 = Haplocarpha lyrata.
aenea, *Jacq. f. Eclog.* 1. t. 52.—Afr. austr.
amoena, *Jacq. Fragm.* 75.—Afr. austr.
amplexicaulis, *Less. Syn. Comp.* 25.—Afr. austr.
angustifolia, Jacq. Hort. Schoenb. ii. 21 = aspera.
angustifolia, *Linn. Sp. Pl.* 923.—Afr. austr.
anthemoides, Linn. Syst. ed. X. 1234 = Ursinia anthemoides.
anthemoides, Thunb. ex Harv. & Sond. Fl. Cap. iii. 152 = Ursinia chrysanthemoides.
arborescens, *Jacq. Hort. Schoenb.* ii. 23.—Afr. austr.
arborescens, Willd. ex Harv. & Sond. Fl. Cap. iii. 453 = aspera.
argentea, Ait. Hort. Kew. iii. 273 = linearis.
argentea, *Thunb. Diss. Arct.* 4 et 7.—Afr. austr.
aspera, *Linn. Sp. Pl.* 922.—Afr. austr.
asplenifolia, Burm. f. Fl. Cap. Prod. 28 = Gazania pinnata.
aurea, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
aureola, Ker-Gawl. in Bot. Reg. i. t. 32 = aspera.
auriculata, Jacq. Hort. Pl. Schoenb. 2. t. 169 = aspera.
bellidifolia, *Berg. Desc. Pl. Cap.* 318.—Afr. austr.

ARCTOTIS:—
bicolor, Willd. Enum. Hort. Berol. 936 = aspera.
breviscapa, Thunb. Diss. Arct. 12 = leptorhiza.
Calendula, Linn. Sp. Pl. 922 = Cryptostemma calendulaceum.
calendulacea, Linn. Syst. ed. XII. 578 = Cryptostemma calendulaceum.
calendulacea, Thunb. ex Harv. & Sond. Fl. Cap. iii. 463 = Venidium hirsutum.
campanulata, *DC. Prod.* vi. 485.—Afr. austr.
candida, *Thunb. Diss. Arct.* 14.—Afr. austr.
canescens, *DC. Prod.* vi. 485.—Afr. austr.
carduifolia, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
caroliniana, *Walt. Fl. Carol.* 216.—Carolina.
caulescens, Thunb. Diss. Arct. 11 = aspera.
cernua, Thunb. l. c. 17 = Ursinia anthemoides.
Cineraria, *Jacq. Hort. Schoenb.* ii. 24.—Afr. austr.
coronopifolia, Salisb. Prod. 210 = Ursinia crithmifolia.
corruscans, DC. Prod. vi. 496 = Cryptostemma calendulaceum.
crithmifolia, Harv. in Harv. & Sond. Fl. Cap. iii. 144 = Ursinia crithmifolia.
crithmoides, Berg. Desc. Pl. Cap. 326 = Ursinia crithmifolia.
crithmoides, Jacq. Hort. Schoenb. ii. 15 = Ursinia pilifera.
cuneata, DC. Prod. vi. 489 = stoechadifolia.
cuprea, Hort. ex Steud. Nom. ed. II. i. 120 = aenea.
cuprea, Jacq. Hort. Schoenb. ii. 25 = aspera.
decumbens, Jacq. l. c. iii. 68. t. 381 = angustifolia.
decumbens, Thunb. Diss. Arct. 9 = stoechadifolia.
decurrens, Jacq. Hort. Schoenb. ii. 20. t. 165 = angustifolia.
dentata, Linn. Pl. Afr. Rar. 25; Amoen. Acad. vi. 107 = Ursinia dentata.
dentata, Thunb. Diss. Arct. 17 (= *Sphenogyne discolor*).—Afr. austr.
denudata, Thunb. l. c. 14 = laevis.
diffusa, *Thunb. l. c.* 9.—Afr. austr.
Dregei, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 93.—Afr. austr.
echinata, DC. Prod. vi. 486 = Landtia hirsuta.
elatior, *Jacq. Hort. Schoenb.* ii. 23.—Afr. austr.
elatior, Willd. ex DC. Prod. v. 488 = amplexicaulis.
elegans, Thunb. Diss. Arct. 14 = Chrysanthemum indicum.
elongata, *Thunb. l. c.* 9.—Afr. austr.
fastuosa, *Jacq. Hort. Schoenb.* ii. 20. t. 166.—Afr. austr.
flaccida, *Jacq. l. c.* 19.—Afr. austr.
flammea, *Jacq. f. Fragm.* 41.—Afr. austr.
foeniculacea, Jacq. Hort. Schoenb. ii. 15 = Ursinia foeniculacea.
formosa, Thunb. Diss. Arct. 11 = aspera.
gigantea, *A. Rich. Tent. Fl. Abyss.* i. 449.—Abyss.
glabrata, Jacq. Hort. Schoenb. ii. 25 = laevis.
glandulosa, *Thunb. Diss. Arct.* 8.—Afr. austr.
glaucophylla, Jacq. Hort. Schoenb. ii. 22 = candida.
glutinosa, Sims, Bot. Mag. t. 1343 = Dimorphotheca cuneata.
grandiflora, Ait. Hort. Kew. iii. 272 = acaulis.
grandiflora, Jacq. Hort. Schoenb. iii. t. 378 = laevis.
grandis, Thunb. Diss. Arct. 8 = stoechadifolia.
hispidella, Juss. ex DC. Prod. vii. 258 = Hispidella hispanica.
humilis, Salisb. Prod. 210 = acaulis.
hypochondriaca, Willd. Sp. Pl. iii. 2343 = Cryptostemma calendulaceum.
incisa, Thunb. Diss. Arct. 10 = aspera.
interrupta, Thunb. l. c. 12 = Arctotheca repens.
Jacobaea, Sch. Bip. in Flora, xxvii. (1844) 772 = aspera.
Kraussii, Sch. Bip. l. c. 771 = angustifolia.
laciniata, *Lam. Encyc.* i. 237.—Afr. austr.
laevis, *Thunb. Diss. Arct.* 10.—Afr. austr.
lanata, Thunb. l. c. 11 = Haplocarpha Thunbergii.
lancea, Thunb. ex DC. Prod. vi. 503 = Didelta tomentosum.
lanceolata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 455.—Afr. austr.
Leichtliniana, *Lynch, in Gard. Chron.* (1885) II. 38.—Afr. austr.
leiocarpa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 451.—Afr. austr.
leptorhiza, *DC. Prod.* vi. 486.—Afr. austr.
leucantha, *Hoffmgg. Verz. Pfl.* 40.—Hab.?
leucanthemoides, *Jacq. Hort. Schoenb.* ii. 19. t. 164.—Afr. austr.

ARCTOTIS:—
linearis, *Thunb. Diss. Arct.* 7.—Afr. austr.
lyrata, Willd. Sp. Pl. iii. 2356 = aspera.
maculata, Jacq. Hort. Schoenb. iii. 67 = aspera.
Massoniana, Spreng. Syst. iv. Cur. Post. 304 = laevis.
melanocycla, Willd. Enum. Hort. Berol. 937 = aspera.
micrantha, Thunb. ex DC. Prod. vi. 493 = Venidium decurrens.
muricata, Thunb. Diss. Arct. 10 = bellidifolia.
nodosa, Thunb. l. c. 15 = Chrysanthemum nodosum.
nudicaulis, Thunb. l. c. 16 (= *Sphenogyne nudicaulis*).—Afr. austr.
odorata, Steud. Nom. ed. II. i. 121 = Ursinia odorata.
oocephala, *DC. Prod.* vi. 486.—Afr. austr.
paleacea, Linn. Pl. Afr. Rar. 25; Amoen. Acad. vi. 107 = Ursinia crithmifolia.
paleacea, Thunb. Diss. Arct. 16 = Ursinia paleacea.
paniculata, Jacq. Hort. Schoenb. iii. 68 = bellidifolia.
pannucea, *Jacq. l. c.* iv. 21.—Afr. austr.
paradoxa, Ait. Hort. Kew. iii. 373 = Ursinia anthemoides.
paradoxa, Linn. Cent. ii. 32; Amoen. Acad. iv. 330 = Ursinia paradoxa.
paradoxa, Thunb. Diss. Arct. 17 = Ursinia chrysanthemoides.
pectinata, Thunb. l. c. 14 = Ursinia dentata.
petiolata, *Thunb. l. c.* 10.—Afr. austr.
pilifera, Thunb. l. c. 18 = Ursinia pilifera.
piligera, Berg. Desc. Pl. Cap. 325 = Ursinia dentata.
pinnata, Thunb. Diss. Arct. 16 = Ursinia filiformis.
pinnatifida, *Thunb. l. c.* 7.—Afr. austr.
plantaginea, Linn. Sp. Pl. ed. II. 1306 = Venidium semipapposum.
plantaginifolia, Salisb. Prod. 210 = Venidium semipapposum.
populifolia, Berg. Desc. Pl. Cap. 323 = Cryptostemma niveum.
prostrata, Salisb. Prod. 210 = Arctotheca repens.
punctata, Thunb. Diss. Arct. 15 (= *Sphenogyne punctata*).—Afr. austr.
purpurea, *Willd. Enum. ex Steud. Nom.* ed. II. i. 121.—Afr. austr.
pusilla, *DC. Prod.* vi. 489.—Afr. austr.
pygmaea, *A. Rich. Tent. Fl. Abyss.* i. 448.—Abyss.
radicans, Linn. Mant. ii. 479, 480 = angustifolia.
repens, Jacq. Hort. Schoenb. iii. 31 = Arctotheca repens.
reptans, *Jacq. l. c.* 69. t. 382.—Afr. austr.
revoluta, *Jacq. l. c.* ii. 24.—Afr. austr.
rigida, Burm. f. Fl. Cap. Prod. 28 = Gazania Pavonia.
rosea, Jacq. Hort. Schoenb. ii. 18 = stoechadifolia.
runcinata, *Hoffmgg. Verz. Pfl.* 40.—Hab. ?
scabra, Thunb. Diss. Arct. 9 = Venidium semipapposum.
scapigera, Thunb. l. c. 12 = acaulis.
scariosa, Ait. Hort. Kew. iii. 274 = Ursinia scariosa.
sericea, Thunb. Diss. Arct. 18 (= *Sphenogyne sericea*).—Afr. austr.
serrata, Thunb. l. c. 8 = Ursinia serrata.
spathulata, Thunb. ex Spreng. Syst. iii. 628 = Venidium perfoliatum.
speciosa, Jacq. Hort. Schoenb. ii. 18 = acaulis.
speciosa, Salisb. Prod. 209 = Cryptostemma calendulaceum.
spinulosa, Jacq. Hort. Schoenb. ii. 21 = fastuosa.
squarrosa, Jacq. l. c. 25 = laevis.
staticefolia, Poir. Encyc. Suppl. i. 438 = Gazania subulata.
stoechadifolia, *Berg. Desc. Pl. Cap.* 324.—Afr. austr.
sulphurea, Gaertn. Fruct. ii. 439 = Cryptostemma hypochondriacum.
sulphurea, Salisb. Prod. 210 = aspera.
superba, Linn. Cent. ii. 32; Sp. Pl. ed. II. 1305 = Cryptostemma calendulaceum.
tenuifolia, Linn. Mant. 288 = Ursinia tenuifolia.
tenuifolia, Poir. Encyc. Suppl. i. 439 = Dimorphotheca graminifolia.
tenuifolia, Salisb. Prod. 210 = Ursinia scariosa.
tomentosa, Thunb. ex DC. Prod. vi. 492 = Venidium semipapposum.
tricolor, Jacq. Hort. Schoenb. ii. 16 = acaulis.
tricolor, Willd. ex DC. Prod. vi. 486 = elongata.
trifida, Thunb. Diss. Arct. 15 (= *Sphenogyne trifida*).—Afr. austr.
trifurcata, Burm. f. Fl. Cap. Prod. 28 = Ursinia crithmifolia.
tristis, Linn. Sp. Pl. 922 = Cryptostemma calendulaceum.
undulata, Gaertn. Fruct. ii. 438 = laciniata.
undulata, Jacq. Hort. Schoenb. ii. 16 = acaulis.

ARCTOTIS:—
undulata, Thunb. Diss. Arct. 13 = aspera.
venidioides, *DC. Prod.* vi. 489.—Afr. austr.
verbascifolia, *Harv. in Harv. & Sond. Fl. Cap.* iii. 452.—Afr. austr.
virgata, *Jacq. Hort. Schoenb.* iii. 32.—Afr. austr.
vitellina, *Hoffmgg. Verz. Pfl.* 40.—Hab. ?

ARCYNOSPERMUM, Turcz. in Bull. Soc. Nat. Mosc. (1858) i. 191. *MALVACEAE* (gen. dub.), Benth. & Hook. f. Gen. i. 199.
nodiflorum, *Turcz. l. c.*—Mexic.

ARCYPHYLLUM, Ell. in Journ. Acad. Philad. i. (1818) 371 = **Rhynchosia**, Lour. (Legum.).
caribacum, Ell. l. c. 372 = R. caribaea.
capitatum, Benth. in Linnaea, xxii. (1849) 525 = R. Clausseni.
difforme, Ell. in Journ. Acad. Phil. i. (1818) 372 = R. tomentosa.
erectum, Ell. l. c. = R. tomentosa.
leucophyllum, Benth. in Linnaea, xxii. (1849) 525 = R. leucophylla.
reticulatum, Ell. in Journ. Acad. Philad. i. (1818) 372 = R. reticulata.
rhombifolium, Ell. l. c. = R. minima.
simplicifolium, Ell. l. c. 371 = R. tomentosa.

ARCYTHOPHYLLUM, Willd. ex Schlecht. in Linnaea, xxviii. (1856) 491 = seq.

ARCYTOPHYLLUM, [Willd.] Roem. & Schult. Mant. iii. 5 (1827) = **Mallostoma**, Karst. (Rubiac.).
blaerioides, [Willd. ex] Roem. & Schult. l. c. = M. caracasana.
nitidum, Schlecht. in Linnaea, xxviii. (1856) 492 = M. nitida.
setosum, Schlecht. l. c. = M. setosa.

ARDERNIA, Salisb. Gen. Pl. Fragm. 35 (1866) = **Ornithogalum**, Linn. (Liliac.).
fuscata, Salisb. l. c. = O. fuscatum.

ARDINGHELIA, Comm. ex A. Juss. Tent. Euph. 21 (1824) = **Phyllanthus**, Linn. (Euphorb.).

ARDISIA, Gaertn. Fruct. ii. 78. t. 94 (1791) = **Cyathodes**, Labill. (Epacrid.).
acerosa, Gaertn. l. c. = C. acerosa.
lanceolata, Gaertn. f. l. c. iii. 148. t. 210.—Austral.
montana, Gaertn. f. l. c. 149.—Austral.
procumbens, Gaertn. f. l. c. 149 = Melichrus rotatus.
ruficarpa, Gaertn. f. l. c. 148. t. 210.—Austral.

ARDISIA, Sw. Prod. Veg. Ind. Occ. 3. 48 (1788); Fl. Ind. Occ. i. 467. t. 10 (1797). *MYRSINEAE*, Benth. & Hook. f. ii. 645.
ANGUILLARIA, Gaertn. Fruct. i. 372 (1788).
BADULA, Juss. Gen. 420 (1789).
BLADHIA, Thunb. Nov. Gen. i. 6 (1781), *nomen prius*.
BARTHESIA, Comm. ex A. DC. in Trans. Linn. Soc. xvii. (1837) 112.
CLIMACANDRA, Miq. Pl. Jungh. i. 199 (1852).
GALIZIOLA, Rafin. Sylva Tellur. 167 (1838).
ICACOREA, Aubl. Pl. Gui. ii. Suppl. i. t. 368 (1775).
KATOUTHEKA, Adans. Fam. ii. 159 (1763).
MILNEA, Rafin. Sylva Tellur. 167 (1838).
MONOPORUS, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 91.
PICKERINGIA, Nutt. in Journ. Acad. Sc. Philad. vii. (1834) 95.
PURKINJIA, Presl, Symb. Bot. ii. 17. t. 64 (1833).
PYRGUS, Lour. Fl. Cochinch. i. 120 (1790).
STYLOGYNE, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 91.
WEDELIA, Steud. Nom. ed. II. i. 133 (1840).
VEDELA, Adans. Fam. ii. 502 (1763).
acuminata, Spreng. Syst. i. 661 = lateriflora.
acuminata, *Willd. Sp. Pl.* i. 1062.—Ind. occ.
adenanthera, Miq. in Mart. Fl. Bras. x. 285 = Parathesis adenanthera.
affinis, Blume, ex A. DC. in DC. Prod. viii. 96 = Myrsine affinis.
alternata, A. DC. in Ann. Sc. Nat. Sér. II. ii. (1841) 296, sphalm. = oxyphylla.
ambigua, *Mart. Herb. Fl. Bras.* 303.—Bras.

ARDISIA:—
amboinensis, *Scheff. Comm. Myrsin.* 75.—Amboin.
Amherstiana, A. DC. in Trans. Linn. Soc. xvii. (1834) 120 = reflexa.
amplexicaulis, *Bedd. Ic. Pl. Ind. Or.* i. 37.—Ind. or.
anceps, Blume, Bijdr. 685 = Blumii.
anceps, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 280 = colorata.
andamanica, *Kurz, For. Fl. Burma*, ii. 108.—Malaya.
angustifolia, *A. DC. in DC. Prod.* viii. 134.—Ins. Philipp.
arborescens, Wall. Cat. n. 2289 = humilis.
arguta, *H. B. & K. Nov. Gen. et Sp.* iii. 247.—Am. austr.
attenuata, Wall. Cat. n. 2286 = oxyphylla.
avenis, Blume, Bijdr. 691 = Myrsine capitellata.
bahamensis, A. DC. in Trans. Linn. Soc. xvii. (1834) 128 = excelsa.
Barthesia, *Steud. Nom.* ed. I. 64.—Zeylan.
Basaal, Roem. & Schult. Syst. iv. 517 = Embelia robusta.
bhotanica, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 528.—Reg. Himal.
bipinnata, *Baker, in Journ. Linn. Soc.* xx. (1884) 201.—Madag.
Blumii, A. DC. in Trans. Linn. Soc. xvii. (1834) 117 = colorata.
Boissieri, A. DC. in DC. Prod. viii. 129 = lanceolata.
borneensis, *Scheff. Comm. Myrsin.* 79.—Borneo.
bracteata, *Baker, in Oliver, Fl. Trop. Afr.* iii. 495.—Afr. trop.
bracteosa, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 127.—Mexic.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 67.—Burma.
brasiliensis, *Spreng. Syst.* i. 662.—Bras.
breviflora, *A. DC. in DC. Prod.* viii. 122.—Peruv.
brevipedata, *F. Muell. Fragm.* vi. 163.—Austral.
bumelioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 516.—Cuba.
calothyrsa, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 371; *et in Flora*, liii. (1870) 254.—Sumatra.
calycosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 292.—Nicaragua.
canaliculata, *Lodd. Bot. Cab.* t. 1083.—Reg. Colomb.
capitata, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 331.—Ins. Fiji.
capollina, *Moç. & Sesse, ex A. DC. in Trans. Linn. Soc.* xvii. (1834) 116.—Mexic.
caribaea, *A. DC. ex Miq. in Mart. Fl. Bras.* x. 289, in adnot.—Ins. Dominica.
cauliflora, *Mart. & Miq. in Mart. Fl. Bras.* x. 291.—Bras.
Cavanillesii, *Roem. & Schult. Syst.* iv. 515.—Ins. Philipp.
celebica, *Scheff. Comm. Myrsin.* 83.—Ins. Celebes.
cestrifolia, *H. B. & K. Nov. Gen. et Sp.* vii. 213.—Reg. Colomb.
chinensis, *Benth. Fl. Hongk.* 207.—China.
chrysophylloides, Miq. Fl. Ind. Bat. Suppl. 574 = tuberculata.
clusioides, *Griseb. Fl. Brit. W. Ind.* 396.—Ind. occ.
coccinea, *Jungh. in Hoev. & De Vriese, Tijdschr.* vii. (1840) 304 ('104' err. typ.).—Java.
colorata, Link, Enum. Hort. Berol. i. 194 = paniculata.
colorata, *Roxb. Hort. Beng.* 16; *Fl. Ind.* i. 581.—Ind. or.
complanata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 280 = colorata.
compressa, *H. B. & K. Nov. Gen. et Sp.* iii. 245.—Reg. Colomb.
confinis, *Miq. Fl. Ind. Bat. Suppl.* 576.—Sumatra.
coriacea, Ledru, ex Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 105 = Ternstroemia brevipes.
coriacea, *Sw. Prod. Veg. Ind. Occ.* 48.—Ind. occ.
courtallensis, *Wight, Ic.* t. 1215.—Ind. or.
crassa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 518.—Malaya.
crenata, *Sims, in Bot. Mag.* t. 1950.—China.
crenata, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 583.—Malaya; China.
crenulata, Lodd. Bot. Cab. t. 2 = crenata.
crenulata, Vent. Choix, t. 5 = Parathesis crenulata.
crispa, A. DC. in Trans. Linn. Soc. xvii. (1834) 124 = crenata.
cubana, *A. DC. l. c.*—Cuba.
Cumingiana, *A. DC. in DC. Prod.* viii. 136.—Ins. Philipp.
cuspidata, *Benth. Bot. Voy. Sulph.* 123.—Ins. Cocos.

ARDISIA :—
cybianthoides, *Miq. Fl. Ind. Bat.* ii. 1028.—Malaya.
cymosa, *Baker, in Oliver, Fl. Trop. Afr.* iii. 495.—Afr. trop.
cymosa, *Blume, Bijdr.* 689.—Java.
decipiens, *A. DC. in DC. Prod.* viii. 125.—Peruv.
decus-montis, *Miq. Pl. Jungh.* 193.—Java.
demissa, *Miq. Fl. Ind. Bat.* ii. 1018.—Malaya.
densa, Miq. l. c. Suppl. 575 = crenata.
denticulata, *Blume, Bijdr.* 691.—Java.
depressa, *C. B. Clarke, in Hook. Fl. Brit. Ind.* iii. 522.—Reg. Himal.
disticha, A. DC. in DC. Prod. viii. 129 = Pimelandra disticha.
divaricata, *Willd. ex Roem. & Schult. Syst.* iv. 804.—Hab. ?
divergens, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 583.—Malaya.
divergens, Thw. Enum. Pl. Zeyl. 174 pro parte = Gardneri.
Doca, Buch.-Ham. ex A. DC. in DC. Prod. viii. 126 = paniculata.
Doma, Wall. ex A. DC. in Trans. Linn. Soc. xvii. (1834) 118 = humilis.
domingensis, *Willd. ex Roem. & Schult. Syst.* iv. 803.—Ins. S. Domingo.
dubia, *Scheff. Comm. Myrsin.* 67.—Malaya.
durifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 214.—N. Granat.
elegans, Andr. Bot. Repos. t. 623 = lentiginosa.
elliptica, Bedd. For. Man. 138 = humilis.
elliptica, *Thunb. Nov. Gen.* viii. 119.—Zeylan.
erythroxyloides, *Thou. ex Roem. & Schult. Syst.* iv. 804.—Madag.
escallonioides, *Schlecht. & Cham. in Linnaea*, vi. (1831) 393.—Mexic.
esculenta, *Pav. ex A. DC. in Trans. Linn. Soc.* xvii. (1834) 128.—Am. austr.
eugeniaefolia, Kurz, For. Fl. Burma, ii. 111 = Pimelandra Griffithii.
eugeniaefolia, Wall. Cat. n. 2276 = Pimelandra eugeniaefolia.
excelsa, *Ait. Hort. Kew.* i. 261.—Ins. Baham.; Madcr.
eximia, *Miq. Pl. Jungh.* 196.—Java.
ferruginea, *H. B. & K. Nov. Gen. et Sp.* iii. 244.—N. Granat.
fertilis, *Miq. Fl. Ind. Bat. Suppl.* 576.—Sumatra.
floribunda, *Roem. & Schult. Syst.* iv. 804.—Madag.
floribunda, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 272 = neriifolia.
floribunda, *Willd. ex Steud. Nom.* ed. II. i. 122, *nomen.*—Madag.
foetida, *Willd. ex Roem & Schult. Syst.* iv. 803.—Am. austr.
formosana, Rolfe, in Journ. Bot. xx. (1882) 358 = Sieboldii.
Forstenii, *Scheff. Comm. Myrsin.* 61.—Malaya.
frangulaefolia, *Zipp. et Scheff. l. c.* 82.—Malaya.
fuliginosa, *Blume, Bijdr.* 692.—Java.
fusca, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 127. t. 2.—Costa Rica.
fusco-pilosa, *Baker, in Journ. Linn. Soc.* xx. (1884) 200.—Madag.
Gardneri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 521.—Zeylan.
glabra, A. DC. in Trans. Linn. Soc. xvii. (1834) 123 = japonica.
glabrata, Blume, Bijdr. 692 = villosa.
glandulosa, Blume, l. c. 690 = crenata.
glandulosa, Roxb. Hort. Beng. 16; Fl. Ind. i. 584 = Amblyanthus glandulosus.
glandulosa-marginata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 128.—Costa Rica.
gracilis, *Miq. in Mart. Fl. Bras.* x. 287.—Bras.
grandiflora, Wall. Cat. n. 2272 = colorata.
grandifolia, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 122.—Burma.
grandis, *Seem. in Bonplandia*, ix. (1861) 259; *Fl. Vit.* 150.—Ins. Fiji.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 523.—Reg. Himal.
guadalupensis, *Duchass. ex Griseb. in Goett. Abh.* vii. (1857) 237.—Ins. Dominic.
Hamiltonii, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 126.—Ind. occ.
Hasseltii, Blume, ex Scheff. Comm. Myrsin. 86. = lanceolata.

ARDISIA :—
Helferiana, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 86.—Burma.
hortorum, *Maxim. in Gartenfl.* (1865) 363. t. 491.—Japon.
Hostmanni, *Miq. in Mart. Fl. Bras.* x. 288.—Guian.
humilis, Blume, Bijdr. 687 = javanica.
humilis, *Vahl, Symb. Bot.* iii. 40.—Ind. or.; Malaya.
Hymenandra, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 282 = Hymenandra Wallichii.
Icara, *Buch.-Ham. in Wall. Cat.* n. 2264.—Ind. or.; Burma.
insularis, *Baker, Fl. Maurit.* 190.—Maurit.
involucrata, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 68.—Reg. Himal.
irasuensis, *Oerst.* in *Kjoeb. Vidensk. Meddel.* (1861) 124.—Costa Rica.
jacquinioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 517.—Ind. occ.
jambosioides, Miq. Pl. Jungh. 196 = crassa.
japonica, *Blume, Bijdr.* 690.—China; Japon.
javanica, *A. DC. in DC. Prod.* viii. 130.—Java.
Junghuhniana, *Miq. Pl. Jungh.* 192.—Sumatra.
Keenani, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 526.—Reg. Himal.
khasiana, *C. B. Clarke, l. c.* 527.—Reg. Himal.
Korthalsiana, Scheff. Comm. Myrsin. 57 = Pimelandra Wallichii.
Kurzii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 525.—Burma.
laevigata, *Blume, Bijdr.* 690.—Java.
laevis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 125.—Costa Rica.
lamponga, *Miq. Fl. Ind. Bat. Suppl.* 577.—Sumatra.
lanceolata, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 583.—Malaya.
lateriflora, *Sw. Prod. Veg. Ind. Occ.* 48.—Ind. occ.
latifolia, Roem. & Schult. Syst. iv. 514 = laurifolia.
latifolia, Sieber, ex A. DC. in Trans. Linn. Soc. xvii. (1834) 113 = guadalupensis.
latipes, *Mart. Herb. Fl. Bras.* n. 570.—Bras.
laurifolia, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 116.—Ind. occ.
lentiginosa, Ker-Gawl. in Bot. Reg. t. 533 = crenata.
lepidota, *H. B. & K. Nov. Gen. et Sp.* iii. 247.—N. Granat.
leptantha, *Miq. in Mart. Fl. Bras.* x. 286.—Bras.
Leschenaultii, A. DC. in DC. Prod. viii. 133 = lanceolata.
Lhotskya, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 127.—Bras.
Liebmannii, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 129. t. 2.—Mexico.
Lindleyana, D. Dietr. Syst. i. 617 = crenata.
linearifolia, *Miq. Fl. Ind. Bat. Suppl.* 577.—Sumatra.
lineata, *Roem. & Schult. Syst.* iv. 804.—Am. austr.
littoralis, Andr. Bot. Rep. t. 630 = humilis.
Lobbiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 529.—Ind. or.
longifolia, *Mart. ex Miq. in Mart. Fl. Bras.* x. 290.—Bras.
longipes, *Baker, in Journ. Linn. Soc.* xxi. (1885) 421.—Madag.
lurida, *Blume, Bijdr.* 687.—Java.
luzonensis, *Presl, Rel. Haenk.* ii. 65.—Ins. Philipp.
macrocalyx, *Scheff. Comm. Myrsin.* 91.—Borneo.
macrocarpa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 277.—Reg. Himal.
macrophylla, *Reinw. ex Blume, Bijdr.* 691.—Java.
macrophylla, Wall. Cat. n. 2290 = grandifolia.
macroscypha, *Baker, in Journ. Linn. Soc.* xxi. (1885) 420.—Madag.
maculata, *Poit. ex A. DC. in Trans. Linn. Soc.* xvii. (1834) 116.—Ins. S. Doming.
mamillata, *Hance, in Journ. Bot.* xxii. (1884) 290.—China.
marginata, *Blume, Bijdr.* 688.—Java.
Martiana, *Miq. in Mart. Fl. Bras.* x. 285.—Bras.
melanosticta, Schlecht. in Linnaea, viii. (1833) 526 = Parathesis melanosticta.
membranacea, *Wall. Cat.* n. 2288; *ex A. DC. Prod.* viii. 134.—Reg. Himal.
metallica, *N. E. Br. in Illustr. Hortic.* xxviii. (1881) t. 421.—Sumatra.
micrantha, *H. B. & K. Nov. Gen. et Sp.* iii. 246.—Venezuela.
microphylla, *Roem. & Schult. Syst.* iv. 804.—Madag.
Miqueliana, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 367; *et in Flora*, liii. (1870) 253.—Malaya.

ARDISIA :—
missionis, *Wall. Cat.* n. 6524; *et ex A. DC. Prod.* viii. 130.—Zeylan.
mollis, Blume, Bijdr. 689 = villosa.
Moonii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 523.—Zeylan.
mucronata, Blume, Bijdr. 685 = lanceolata.
multiflora, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 516.—Ind. occ.
multiflora, Willd. ex Roem. & Schult. Syst. iv. 804 = revoluta.
myriantha, *Baker, in Journ. Linn. Soc.* xxi. (1885) 419.—Madag.
myristicaefolia, *Blume, ex Scheff. Comm. Myrsin.* 63.—Malaya.
nana, Buch.-Ham. ex A. DC. in Trans. Linn. Soc. xvii. (1834) 118 = humilis.
neriifolia, A. DC. Prod. viii. 127, pro parte = andamanica.
neriifolia, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) ii. 225 = depressa.
neriifolia, *Wall. Cat.* n. 2278.—Reg. Himal.
nicaraguensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 123.—Nicaragua.
nigrescens, *Oerst. l. c.* 130. t. 2.—Mexic.
nigricans, *Miq. in Mart. Fl. Bras.* x. 289.—Bras.
nitidula, *Baker, in Journ. Linn. Soc.* xx. (1884) 200.—Madag.
nigro-punctata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 127. t. 2.—Costa Rica.
nutans, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 121.—Java.
oblonga, A. DC. l. c. = oxyphylla.
obovata, Blume, Bijdr. 688 = humilis.
obovata, Ham. Prod. Fl. Ind. Occ. 26 = Hamiltonii.
odontophylla, Lindl. Bot. Reg. t. 1892 = chinensis.
odontophylla, *Wall. Cat.* n. 2279.—Reg. Himal.
oleracea, Buch.-Ham. ex A. DC. in Trans. Linn. Soc. xvii. (1834) 118 = humilis.
oligantha, *Baker, in Journ. Linn. Soc.* xxi. (1885) 419.—Madag.
Oliveri, *Mast. in Gard. Chron.* (1877) ii. 680.—Costa Rica.
Opegrapha, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 126.—Costa Rica.
orinocensis, *H. B. & K. Nov. Gen. et Sp.* iii. 244.—Fluv. Orinoco.
ovata, Thunb. Nov. Gen. viii. 119 = colorata.
oxyantha, Wall. Cat. n. 2275 = Antistrophe oxyantha.
oxyphylla, *Wall. l. c.* n. 2291; *A. DC. Prod.* viii. 130.—Burma; Malaya.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 575.—Sumatra.
paniculata, A. DC. in DC. Prod. viii. 139 = A. Samara.
paniculata, *Roxb. Hort. Beng.* 16; *Fl. Ind.* i. 580.—Ind. or.
parasitica, Sw. Prod. Veg. Ind. Occ. 48 = Grammadenia parasitica.
parvifolia, *Humb. ex Willd. sec. Roem. & Schult. Syst.* iv. 804.—Peruv.
pauciflora, Benth. Fl. Hongk. 206 = pentagona.
pauciflora, *Heyne, ex Roxb. Fl. Ind. ed. Carey*, ii. 279.—Ind. or.
pavonina, Buch.-Ham. ex A. DC. Prod. viii. 136 = odontophylla.
pedunculata, Boj. ex Baker, in Journ. Linn. Soc. xx. (1884) 203 = Oncostemum pedicellatum.
pedunculosa, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 279.—Reg. Himal.
pellucida, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 130. t. 2.—Mexic.
pentagona, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 124.—Java.
pergamacea, Blume, ex A. DC. in DC. Prod. viii. 85 = Embelia pergamacea.
Perrottetiana, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 95.—Ins. Philipp.
peruviana, *Miq. in Mart. Fl. Bras.* x. 290.—Peruv.
petocalyx, *Scheff. Comm. Myrsin.* 91.—Sumatra.
philippinensis, *A. DC. in DC. Prod.* viii. 128.—Ins. Philipp.
Pickeringia, *Torr. & Gray, ex DC. Prod.* viii. 124.—Florida.
plagioneura, *Scheff. Comm. Myrsin.* 85.—Malaya.
polycarpa, Wall. Cat. n. 2285 = colorata.
polycephala, *Wall. l. c.* n. 2293; *A. DC. Prod.* viii. 131.—Burma.
polycephala, Wight, Illustr. t. 145 = humilis.
polygama, *A. DC. in DC. Prod.* viii. 138.—Molucca.

ARDISIA :—
polyneura, Miq. Pl. Jungh. 198 = colorata.
polysticta, Miq. Fl. Ind. Bat. Suppl. 576 = crenata.
porosa *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 522. —Malaya.
primulaefolia, *Gardn. & Champ. in Hook. Kew Journ.* i. (1849) 324.—China.
procumbens, Gaertn. f. Fruct. iii. t. 210 = Melichrus rotatus.
propinqua, *H. B. & K. Nov. Gen. et Sp.* vii. 213.—Venezuela.
pseudo-Icacorea, *Miq. in Mart. Fl. Bras.* x. 283.—Bras.
pseudojambosa, *F. Muell. Fragm.* iv. 81.—Austral.
pterocaulis, *Miq. Pl. Jungh.* 194.—Sumatra.
pteropoda, *Miq. Fl. Ind. Bat.* ii. 1025.—Malaya.
pubicalyx, *Miq. l. c. Suppl.* 575.—Sumatra.
pumila, Blume, Bijdr. 688 = Labisia pothoina.
punctata, Blume, l. c. 687 = nutans.
punctata, Jack, in Roxb. Fl. Ind. ed. Carey, ii. 275 = divergens.
punctata, Lindl. Bot. Reg. t. 827 = crenata.
punctulosa, D. Dietr. Syn. Pl. i. 615 = divergens.
purpurea, Reinw. in Blume, Bijdr. 684 = lanceolata.
pusilla, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 126. —Japon.
pyramidalis, *Pers. Syn.* i. 233; *A. DC. in Trans. Linn. Soc.* xvii. (1834) 123.—Ind. occ.
pyramidalis, Roth, Nov. Pl. Sp. 123 = colorata.
pyrgina, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Pyrgus.
Pyrgus, *Roem. & Schult. Syst.* iv. 518.—China.
pyrifolia, *Willd. ex Roem. & Schult. l. c.* 804.—Madag.
quinquangularis, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 95.—Assam.
quinquegona, Blume, Bijdr. 689 = pentagona.
racemosa, *Spreng. Syst.* i. 661.—Bras.
ramiflora, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 132.—Costa Rica.
reclinata, *Scheff. Comm. Myrsin.* 69.—Malaya.
reflexa, *Wall. Cat.* n. 2282; *A. DC. Prod.* viii. 132.—Burma.
revoluta, *H. B. & K. Nov. Gen. et Sp.* iii. 246.—Mexico.
rhomboidea, *Wight, Ic. Pl.* t. 1213.—Ind. or.
rhynchocarpa, *Scheff. Comm. Myrsin.* 68.—Malaya.
rhynchophylla, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 529.—Reg. Himal.
rigida, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 87.—Burma.
rostrata, Hassk. in Flora, li. (1868) 26 = humilis.
Rothii, A. DC. in Trans. Linn. Soc. xvii. (1834) 117 = colorata.
Roxburghiana, D. Dietr. Syn. Pl. i. 617 = Amblyanthus glandulosus.
rubiginosa, *Miq. Pl. Jungh.* 192.—Sumatra.
salicifolia, A. DC. in DC. Prod. viii. 129 = humilis.
Samara, *A. DC. ex Buek, Index*, iii. praef. 8.—Ind. or.
sanguinolenta, Blume, Bijdr. 685 = colorata.
sanguinolenta, Wall. Cat. n. 2287 = Wallichii.
scandens, *Blume, Bijdr.* 686.—Java.
Schomburgkiana, *Miq. in Mart. Fl. Bras.* x. 289.—Bras.
scrobiculata, *Seem. Fl. Vit.* 150.—Ins. Fiji.
semicrenata, Mart. Herb. Fl. Bras. 571 = acuminata.
semidentata, *Miq. Pl. Jungh.* 191.—Java.
serrata, Pers. Syn. i. 233 = Cavanillesii.
serratifolia, Bedd. in Madr. Journ. Ser. III. i. (1864) 51 = Antistrophe serratifolia.
serrulata, Kurz, in Journ. As. Soc. Beng. xlii. (1873) ii. 87 = A. Icara.
serrulata, *Sw. Prod. Veg. Ind. Occ.* 48.—Ind. occ.
sessilis, *Scheff. Comm. Myrsin.* 59.—Malaya.
Sieberi, *Baker, Fl. Maurit.* 190.—Maurit.
Sieboldi, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iii. 190. —Japon.
sinensis, Hort. ex Steud. Nom. ed. II. i. 122 = A. Pyrgus.
solanacea, Roxb. Hort. Beng. 16; Fl. Ind. i. 580 = humilis.
Spanoghei, *Scheff. Comm. Myrsin.* 62.—Malaya.
speciosa, Blume, Bijdr. 684 = lanceolata.
sphenobasis, *Scheff. Comm. Myrsin.* 65.—Malaya.
spicata, Wall. Cat. n. 2273 = Labisia pothoina.
squamulosa, *Presl, Rel. Haenk.* ii. 65.—Ins. Luzon.
Storckii, *Seem. Fl. Vit.* 150.—Ins. Fiji.
stylosa, *Miq. Pl. Jungh.* 196.—Java.
subcuneata, *Miq. Fl. Ind. Bat.* ii. 1028.—Malaya.
sumatrana, *Miq. Pl. Jungh.* 195.—Sumatra.
sumbavana, *Miq. Fl. Ind. Bat.* ii. 1027.—Malaya.

ARDISIA :—
surinamensis, *Miq. in Mart. Fl. Bras.* x. 288.—Guiana.
tavoyana, A. DC. in Trans. Linn. Soc. xvii. (1834) 125 = villosa.
synneura, *Scheff. Comm. Myrsin.* 80.—Malaya.
tenuiflora, Kollmann, ex A. DC. Prod. viii. 127 = pauciflora.
tenuifolia, *Blume, Bijdr.* 686.—Java.
tenuiramis, *Miq. Pl. Jungh.* 198.—Java.
ternatensis, *Scheff. Comm. Myrsin.* 58.—Ins. Ternate.
tetrandra, H. B. & K. Nov. Gen. et Sp. iii. 243 = Cybianthus Humboldtii.
Teysmanniana, *Scheff. in Tijdschr. Nederl. Ind.* xxxi. (1870) 368; *et in Flora*, liii. (1870) 253.—Malaya.
thyrsiflora, D. Don, Prod. Fl. Nep. 148 = neriifolia.
tinifolia, *Sw. Prod. Veg. Ind. Occ.* 48.—Jamaica.
tomentosa, *Presl, Rel. Haenk.* ii. 66.—Ins. Luzon.
Tsjeriam-Cottam, Roem. & Schult. Syst. iv. 518 = Embelia robusta.
tuberculata, *Wall. Cat.* n. 2274.—Malaya.
turbacensis, *H. B. & K. Nov. Gen. et Sp.* iii. 245.—N. Granat.
umbellata, *Baker, in Journ. Linn. Soc.* xxi. (1885) 420.—Madag.
umbellata, Roxb. Hort. Beng. 16; Fl. Ind. i. 582 = humilis.
umbrosa, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie*, ii. (1845) 575.—Java.
undulata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 524. —Reg. Himal.
verrucosa, *Presl, Rel. Haenk.* ii. 65.—Ins. Philipp.
vestita, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 274 = villosa.
villosa, *Roxb. Hort. Beng.* 85; *Fl. Ind.* i. 582.—Malaya.
virens, *Kurz, For. Fl.* ii. 575.—Reg. Himal.; China.
viridiflora, *Blume, ex A. DC. in DC. Prod.* viii. 88.—Malaya.
vitiensis, *Seem. Fl. Vit.* 150.—Ins. Fiji.
volubilis, Wall. ex Steud. Nom. ed. II. i. 122 = colorata.
Wallichii, *A. DC. in Trans. Linn. Soc.* xvii. (1834) 123. —Burma.
Wallichii, Thw. Enum. Pl. Zeyl. 174 = Moonii.
Wightiana, Wall. Cat. n. 2330 = humilis.
xylosteoides, *Griseb. Fl. Brit. W. Ind.* 395.—Jamaica.
zeylanica, *C. B. Clarke, in Hook. Fl. Brit. Ind.* iii. 520. —Zeylan.
Zollingeri, A. DC. in DC. Prod. viii. 670 = colorata.

ARDISIANDRA, Hook. f. in Journ. Linn. Soc. vii. (1864) 205. t. 1. *PRIMULACEAE*, Benth. & Hook. f. ii. 633.
sibthorpioides, *Hook. f. l. c.*—Afr. trop. occ.

ARDUINA, Adans. Fam. ii. 499 (1763) = **Kundmannia**, Scop. (Umbell.).

ARDUINA, Mill. Ic. ii. t. 300 (1759); Linn. Mant. i. 7 (1767) = **Carissa**, Linn. (Apocyn.).
acuminata, E. Mey. Comm. Pl. Afr. Austr. 191 = C. acuminata.
bispinosa, Linn. Mant. i. 52 = C. bispinosa.
bispinosa, Lodd. Bot. Cab. t. 387 = C. ferox.
edulis, Spreng. Syst. i. 669 = C. edulis.
erythrocarpa, Eckl. in S. Afr. Journ. i. (1830) 372 = C. erythrocarpa.
ferox, E. Mey. Comm. Pl. Afr. Austr. 191 = C. ferox.
grandiflora, E. Mey. l. c. 190 = C. grandiflora.
haematocarpa, Eckl. in S. Afr. Journ. i. (1830) 372 = C. haematocarpa.
macrocarpa, Eckl. l. c. = C. macrocarpa.

ARECA, Linn. Sp. Pl. ed. I. 1189 (1753). *PALMAE*, Benth. & Hook. f. iii. 883.
ARECCA, Linn. Fl. Zeyl. 186 (1747).
alba, Bory, Voy. i. 306 = Dictyosperma album.
Alicae, *W. Hill, ex F. Muell. in Gartenfl.* (1879) 199. —Austral.
angulosa, *Giseke, Linn. Praelect.* 80.—Ind. or.
arundinacea, *Becc. Malesia*, i. (1877) 23.—Borneo.
augusta, Kurz, in Journ. Bot. xiii. (1875) 331 = Ptychoraphis augusta.
aurea, Hort. in Fl. des Serres, Sér. II. vii. (1867–68) 43 = Dictyosperma aureum.
Bacaba, *Arruda, in Koster, Trav. Bras.* 490.—Pernambuc.

ARECA :—
Banksii, A. Cunn. ex Kunth, Enum. Pl. iii. 185 = Rhopalostylis Baueri.
Baueri, Hook. f. ex Lem. Illustr. Hortic. xv. (1868) 575 = Rhopalostylis Baueri.
borbonica, Hort. ex Kunth, Enum. Pl. iii. 186 = Dictyosperma album.
borneensis, *Becc. Malesia*, i. (1877) 22.—Borneo.
Calapparia, Blume, Rumphia, ii. t. 100. f. 2 = Actinorhytis Calapparia.
Catechu, *Linn. Sp. Pl.* 1189.—Ind. or.; Malaya.
Cathecu, Burm. f. Fl. Ind. 241 = A. Catechu.
cincta, Walp. Ann. v. 808 = Acanthophoenix crinita.
cocoides, Griff. in Calc. Journ. Nat. Hist. iv. (1845) 454 = Actinorhytis Calapparia.
communis, Zipp. ex Blume, Rumphia, ii. 73, 119. t. 128 = Drymophloeus communis.
concinna, *Thw. Enum. Pl. Zeyl.* 328.—Zeylan.
cornuta, *Giseke, Linn. Praelect.* 81.—Ind. or.
coronata, Blume, in Mart. Hist. Nat. Palm. 179 = Pinanga coronata.
costata, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 200 (= Pinangae sp.).—Ins. Nicobar.
crinita, Bory, Voy. i. 307 = Acanthophoenix crinita.
cuneifolia, Stokes, Bot. Mat. Med. ii. 318 = humilis.
curvata, Griff. Ic. Pl. As. t. 248 = Pinanga disticha.
curvata, Griff. Notul. iii. 164 = Pinanga paradoxa.
Dicksonii, Roxb. Hort Beng. [105]; Fl. Ind. iii. 616 = Pinanga Dicksonii.
disticha, Roxb. ll. cc. 620 = Pinanga disticha.
elaeocarpa, Reinw. ex Kunth, Enum. Pl. iii. 195 = Drymophloeus oliviaeformis.
erythrocarpa, H. Wendl. ex Kerch. Palm. 231 = erythropoda.
erythropoda, Miq. in Journ. Bot. Néerl. i. (1861) 6 = Cyrtostachys Renda.
Faufel, Gaertn. Fruct. i. 19. t. 7. fig. 2 = A. Catechu.
furcata, *Becc. Malesia*, i. (1877) 23.—Borneo.
furfuracea, Hort. ex H. Wendl. in Kerch. Palm. 231 = Dictyosperma furfuraceum.
gigantea, Hort. l. c. 254 = Pinanga ternatensis.
glandiformis, *Lam. Encyc.* i. 241.—Ins. Molucc.
globulifera, Lam. Encyc. 895 = Pinanga Kuhlii.
gracilis, Roxb. Hort. Beng. 68; Fl. Ind. iii. 619 = Pinanga gracilis.
gracilis, Thou. ex Kunth, Enum. Pl. iii. 188 = Dypsis pinnatifrons.
gracilis, Giseke, Linn. Praelect. 80 = Drymophloeus appendiculata.
haematocarpon, Griff. Notul. iii. 165 = Pinanga malaiana.
hexasticha, Kurz, in Journ. As. Soc. Beng. xliii. (1874) ii. 201 = Pinanga hexasticha.
himalayana, Griff. ex H. Wendl. in Kerch. Palm. 231 = A. Catechu.
horrida, Griff. in Calc. Journ. Nat. Hist. v. (1845) 465 = Oncosperma horrida.
hortensis, Lour. Fl. Cochinch. 568, in obs. = A. Catechu.
humilis, Blanco, ex H. Wendl. in Kerch. Palm. 231 = triandra ?
humilis, Roxb. ex H. Wendl. l. c. = Pinanga disticha.
humilis, Willd. Sp. Pl. iv. 595 = Drymophloeus saxatilis.
jobiensis, *Becc. Malesia*, i. (1877) 21.—N. Guin.
lactea, Hort. Bogor. ex Miq. Fl. Ind. Bat. iii. 10 = Dictyosperma album.
lansiformis, *Giseke, Linn. Praelect.* 81.—Ind. or.
latisecta, Scheff. in Tijdschr. Nederl. Ind. xxxii. (1871) 168 = Pinanga latisecta.
laxa, *Buch.-Ham. in Mem. Wern. Soc.* v. (1824) 309. —Ins. Andam.
litoralis, Blume, Rumphia, ii. 123, in obs. = Drymophloeus communis.
lutescens, Bory, Voy. ii. 296 = Chrysalidocarpus lutescens.
macrocalyx, *Zipp. ex Blume, Rumphia*, ii. 75. t. 101, 160, 163 D.—N. Guinea.
madagascarica, *Stadtmann, ex Bojer, Hort. Maurit.* 305. —Madag.
madagascariensis, *Mart. Hist. Nat. Palm.* 179.—Madag.
malaiana, Griff. in Calc. Journ. Nat. Hist. v. (1845) 457 = Pinanga malaiana.
minuta, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 146. —Borneo.
monostachya, Mart. Hist. Nat. Palm. 178. t. 135. f. 4 = Bacularia monostachya.
nagensis, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 453. Ind. or.

ARECA :—
Nenga, Blume, in Mart. Hist. Nat. Palm. 179 = Nenga Wendlandiana.
Nibung, Griff. ex H. Wendl. in Kerch. Palm. 231 = Oncosperma filamentosa.
Nibung, Mart. Hist. Nat. Palm. 173. t. 150. f. 4 = Oncosperma horrida.
nigra, Giseke, ex H. Wendl. in Kerch. Palm. 231 = A. Catechu.
nobilis, Hort.—Cf. Gard. Chron. (1868) 349 = Nephrosperma Van-Houtteana.
Normanbyi, *F. Muell. Fragm.* viii. 235.—Austral.
oleracea, Jacq. Am. Select. 278 = Oreodoxa oleracea.
olivaeformis, Giseke, in Ord. Nat. Pl. 79 = Drymophloeus olivaeformis.
orizaeformis, Gaertn. Fruct. i. 20 = Pinanga Kuhlii.
oviformis, *Giseke, Linn. Praelect.* 81.—Ind. or.
oxycarpa, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1868) n. v. 1.—Malaya.
paniculata, Scheff. ex H. Wendl. in Kerch. Palm. 231 = Mischophloeus paniculata.
paradoxa, Griff. in Calc. Journ. Nat. Hist. v. (1845) 463 = Pinanga paradoxa.
parva, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 6.—Malaya.
polystachys, *Miq. ex H. Wendl. in Kerch. Palm.* 232. —Malaya.
propria, Hort. Bogor. ex Miq. Fl. Ind. Bat. iii. 10 = Dictyosperma album.
procera, Zipp. ex Blume, Rumphia, ii. 95 = Kentia procera.
pumila, Blume, in Mart. Hist. Nat. Palm. 177. t. 153. fig. 1–3 = Nenga Wendlandiana.
punicea, Zipp. ex Blume, Rumphia, ii. 72. t. 122 = Drymophloeus puniceus.
purpurea, *Hort. in Illustr. Hortic.* xxiv. (1877) t. 298. —Maurit.
rubra, Bory, Voy. i. 306 = Acanthophoenix rubra.
rubra, Hort. ex H. Wendl. in Kerch. Palm. 232 = Dictyosperma rubrum.
sanguinea, Zipp. ex Blume, Rumphia, ii. 72 = Drymophloeus puniceus.
sapida, Soland. ex Forst. f. Pl. Esc. 66 = Rhopalostylis sapida.
saxatilis, Burm. f. Fl. Ind. 42 = Drymophloeus saxatilis.
spicata, Lam. Encyc. i. 241 = Calyptrocalyx spicatus.
spinosa, Hasselt & Kunth, Enum. Pl. iii. 185 = Oncosperma filamentosum.
sylvestris, Lour. Fl. Cochinch. ii. 568 = glandiformis.
tenella, *Becc. Malesia*, i. (1877) 22.—Borneo.
tigillaria, Jack, in Malay. Misc. ii. (1820) n. VII. p. 88 = Oncosperma filamentosum.
triandra, *Roxb. Hort. Beng.* 68; *Fl. Ind.* iii. 617.—Ind. or.
umbrosa, *Link, Enum. Hort. Berol.* ii. 403, nomen.—Hab.?
vaginata, Giseke, Linn. Praelect. 78 = Drymophloeus appendiculatus.
Verschaffelti, Hort. ex Lem. in Illustr. Hortic. xiii. (1866) t. 462-3 = Hyophorbe Verschaffelti.
vestiaria, *Giseke, Linn. Praelect.* 78.—Hab.?
Wallichiana, Mart. Hist. Nat. Palm. 178 = Iguanura Wallichiana.
Wendlandiana, Scheff. ex H. Wendl. in Kerch. Palm. 332 = Nenga Wendlandiana.

ARELINA, Neck. Elem. i. 83 (1790) = **Berkheya**, Ehrh. (Compos.).

AREMONIA, Neck. Elem. ii. 100 (1790) = **Agrimonia**, Linn. (Rosac.).
agrimonoides, DC. Prod. ii. 588 = Agrimonia Eupatoria.

ARENARIA, Adans. Fam. ii. 256 (1763) = **Spergula**, Linn. (Caryophyll.).

ARENARIA, Rupp. ex Linn. Syst. ed. I. (1735). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 149.
ADENARIUM, Rafin. in Am. Monthly Mag. (1818) 266.
ALSINANTHUS, Reichb. Handb. 298 (1837).
ALSINE, Scop. Fl. Carn. ed. I. 224 (1772).
ALSINELLA, [Sw.] Summa Veg. Scand. 17 (1814), pro parte; S. F. Gray, Nat. Arr. Brit. Pl. ii. 655 (1821).

ARENARIA :—
AMMODENIA, Patrin, ex S. Gmel. Fl. Sibir. iv. 160 (1769).
BIGELONIA, (Bigelowia) Rafin. in Journ. Phys. lxxxix. (1819) 289.
BRACHYSTEMMA, D. Don, Prod. Fl. Nep. 216 (1828).
CHERLERIA, Hall. It. Helv. ii. n. 79. t. 1 (1740).
CHETROPIS, Rafin. Fl. Tellur. iii. 80 (1836).
DOLOPHRAGMA, Fenzl, in Ann. Wien. Mus. i. (1836) 63. t. 7.
EREMOGONE, Fenzl, Verbr. Alsin. 13. t. ad p. 18 (1833).
EUTHALIA, Rupr. Fl. Caucas. 220 (1869).
FACCHINIA, Reichb. Ic. Fl. Germ. v. 29 (1841).
GOUFFEIA, Robill. & Cast. in DC. Fl. Fr. v. 609 (1815).
GRENIERA, J. Gay, in Ann. Sc. Nat. Sér. III. iv. 27 (1845).
GYPSOPHYTUM, Adans. Fam. ii. 256 (1763).
HALIANTHUS, Fries, Fl. Halland. 75 (1817).
HONKENYA, Ehrh. Beitr. ii. 180 (1788).
LEPTOPHYLLUM, Ehrh. l. c. iv. 147 (1789).
LEPYRODICLIS, Fenzl, in Endl. Gen. 966 (1840).
MERCKIA, Fisch. ex Cham. & Schlecht. in Linnaea, i. (1826) 59.
MERKIA, Reichb. Handb. 298 (1837).
MINUARTIA, Linn. Sp. Pl. 89 (1753).
MOEHRINGIA, Linn. Syst. ed. II. 22 (1740).
NEUMAYERA, Reichb. Ic. Fl. Germ. v. 30 (1841).
ODONTOSTEMMA, Benth. in Wall. Cat. n. 645 (1828); G. Don, Gen. Syst. i. 449 (1831).
PETTERA, Reichb. Ic. Fl. Germ. v. 33. t. 220 (1841); Nom. 205 (1841).
PLINTHINE, Reichb. Handb. 298 (1837).
RHODALSINE, J. Gay, in Ann. Sc. Nat. Sér. III. iv. 25 (1845).
SABULINA, Reichb. Fl. Germ. Excurs. 785 (1832).
SIEBERA, Hoppe, in Flora, ii. (1819) 24.
SOMERAUERA, Hoppe, l. c. 26 (1819).
STROPHIUM, Dulac, Fl. Hautes-Pyr. 247 (1867).
TRYPHANE, Reichb. Nom. 205 (1841).
WIERZBICKIA, Reichb. l. c. (1841).
WILHELMSIA, Reichb. Consp. 206 (1828).
XERALSINE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 347.
abietina, J. & C. Presl, Delic. Prag. i. 63 = grandiflora.
acerosa, *Boiss. & Heldr. Diagn.* Ser. I. viii. 103.—As. Min.
achalensis, *Griseb. in Goett. Abh.* xxiv. (1879) 26.—Reg. Argent.
acicularis, Fisch. ex DC. Prod. i. 403 = juniperina.
aculeata, *S. Wats. in Bot. King, Exp.* 40.—Am. bor.
adenotrichia, Bong. ex Ledeb. Fl. Ross. i. 291 = Gypsophila petraea.
affinis, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 198.—Hab.?
africana, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 184. —Afr. trop.
agglomerata, Poir. Encyc. Suppl. v. 6 = glomerata.
aggregata, Boiss. Elench. 23, in obs. = erinacea.
aggregata, Loisel. Fl. Gall. i. 317 = tetraquetra.
airaefolia, Fisch. ex DC. Prod. i. 402 = capillaris.
algarbiensis, *Welw. ex Willk. Ic. Pl. Hispan.* i. 93.—Lusitan.
Alpamarcae, *A. Gray, Bot. U. St. Expl. Exped.* i. 116. —Peruv.
alpicola, Tenore, Fl. Nap. iv. 224 = graminifolia.
alpina, Porter, Fl. Colorad. 14 = obtusa.
alsinoides, Raddi, Brev. Observ. 13 = Stellaria uliginosa.
alsinoides, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 201 = lanuginosa.
altaica, Fisch. ex DC. Prod. i. 404 = arctica.
amabilis, Bory, in Ann. Gén. Sc. Phys. iii. (1820) 5 = tetraquetra.
andicola, *Gill. in Hook. Bot. Misc.* iii. (1833) 148.—Chili.
andina, *Rohrb. in Linnaea*, xxxvii. (1871–72) 255.—Bolivia.
androsacea, Fisch. ex DC. Prod. i. 402 = rigida.
apetala, Vill. Prosp. 48 = biflora.
aphanantha, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 293.—Peruv.
arabica, Hort. Berol. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. xi. 49 = tenuifolia.
arctica, *Stev. ex DC. Prod.* i. 404.—Reg. bor. et arct.
Arduini, Vis. Stirp. Dalm. 8. t. 3. f. 2 = graminifolia.

ARENARIA :—
aretioides, *Portenschl. ex J. Gay, in Ann. Sc. Nat.* Sér. I. iii. (1824) 34, nota.—Europ.
aristata, Rafin. Caratteri, n. 187 = setacea.
armeniaca, *Boiss. Diagn.* Ser. I. i. 48.—Armenia.
Armeriastrum, *Boiss. Elench.* 22.—Hispan.
armerina, *Bory, in Ann. Gén. Sc. Phys.* iii. (1820) 5. —Hispan.
arvatica, Presl, Fl. Sicul. 163 = mucronata.
arvensis, Wallr. Sched. Crit. 200 = Spergula arvensis.
Aucheriana, *Boiss. Diagn.* Ser. I. i. 50.—Persia.
austriaca, All. Fl. Pedem. ii. 112. t. 64. f. 2 = Villarsii.
austriaca, *Jacq. Fl. Austr.* iii. 39. t. 270.—Europ.
baetica, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1877) 365 = emarginata.
baicalensis, Steud. Nom. ed. II. i. 127 = uliginosa.
Balansae, *Boiss. Fl. Orient.* i. 700.—As. Min.; Persia.
baldensis, Reichb. ex Nym. Consp. 113 = bavarica.
balearica, *Linn. Syst. Nat.* ed. XII. App. 230.—Ins. Balear.; Corsic.
banatica, *Heuff. in Flora*, xxi. (1838) 359.—Hungaria.
Barrelieri, Vill. Hist. Pl. Dauph. iv. 634 = tenuifolia.
Bartolloti, *Tineo, Pl. Rar. Sic. Pugill.* i. 10.—Sicilia.
bavarica, *Linn. Cent.* ii. 17; *Amoen. Acad.* iv. 315.—Europ.
bavarica, *Pall. Reise*, ii. 522.—Rossia.
Benthamii, *Edgew. & Hook. f. in Hook. f. Fl. Brit. Ind.* i. 242.—Reg. Himal.
Benthamii, *Fenzl, ex Torr. & Gray, Fl. N. Am.* i. 675. —N. Mexic.
Berteroana, Phil. in Linnaea, xxviii. (1856) 673 = Spergularia villosa?
Biebersteinii, Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 202 = graminifolia.
biflora, Griseb. Spicil. Fl. Rumel. i. 203 = rotundifolia.
biflora, *Linn. Mant.* 74.—Europ.
bisulca, *Fenzl, ex Rohrb. in Linnaea*, xxxvii. (1871–73) 248.—Peruv.
blepharophylla, *Boiss. Fl. Orient.* i. 693.—Armenia.
Bocconi, Soleir. ex Steud. Nom. ed. II. i. 123, 125 = Spergularia media.
bonariensis, *Gill. in Hook. Bot. Misc.* iii. (1833) 148. —Reg. Argent.
Bourgaeana, Coss. ex Willk. in Linnaea, xl. (1876) 117 = incrassata.
Bourgaei, *Hemsl. Diagn. Pl. Nov.* 21.—Mexic.
brevicaulis, Sternb. in Spreng. Pugill. i. 31 = lanceolata.
breviflora, Gilib. Fl. Lituan. ii. 155 = serpyllifolia.
brevifolia, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 180. —Am. bor.
Brotherana, *Trautv. in Act. Hort. Petrop.* viii. (1883) 151.—Rossia.
bryoides, Fisch. ex DC. Prod. i. 404 = arctica.
bryoides, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 201.—Mexic.
buxifolia, Poir. Encyc. vi. 362 = lateriflora.
caerulescens, Rudolph, ex Ledeb. Fl. Ross. i. 291 = Gypsophila petraea.
caespitosa, Ehrh. Beitr. v. 177 = verna.
cacspitosa, *Phil. in Linnaea*, xxviii. (1856) 673.—Chili.
caespitosa, Salisb. Prod. 299 = balearica.
caespitosa, J. Vahl, ex Nym. Consp. 121 = Sagina nivalis.
californica, *Brewer, in Bot. Calif.* i. 69.—Calif.
calycantha, Ledeb. in Mém. Acad. Sc. Pétersb. v. (1812) 534 = Stellaria borealis.
calycina, Haensel. ex Willk. & Lange, Prod. iii. 603 = Sagina subulata.
calycina, Hort. ex Ledeb. Fl. Ross. i. 343 = tenuifolia.
calycina, *Poir. Voy. Barb.* ii. 167.—Afr. bor.
calyculata, *Poir. Encyc. Suppl.* v. 7.—Hungaria?
campestris, All. Fl. Pedem. ii. 114 = Spergularia rubra.
canadensis, Pers. Syn. i. 504 = Spergularia rubra.
canescens, *Vahl, ex Hornem. Hort. Hafn.* i. 964.—Hab.?
cantabrica, Amo, Fl. Iber. vi. 140 = serpyllifolia.
capensis, *Steud. Nom.* ed. II. i. 124.—Afr. austr.
capillacea, All. Fl. Pedem. ii. 365 = grandiflora.
capillacea, Willd. ex Ledeb. Fl. Ross. i. 367 = capillaris.
capillaris, *Poir. Encyc.* vi. 380.—Sibir.; Am. bor.
capillipes, *Boiss. Voy. Bot. Espagne*, 98. t. 27.—Hispan.; Afr. bor.
capitata, *Lam. Fl. Fr.* iii. 39.—Hisp.; Lusit.
caricifolia, *Boiss. in Tchihat. Asie Min. Bot.* i. 235.—Armenia.
caroliniana, Walt. Fl. Carol. 141 = squarrosa.
cassia, *Boiss. Diagn.* Ser. I. viii. 101.—Syria.

ARENARIA:—
caucasica, Adams, ex Ledeb. Fl. Ross. i. 354 = pinifolia.
Cephalotes, *Bieb. Fl. Taur. Cauc.* i. 346.—Tauria.
cerastiiflora, Spreng. Syst. v. 62 Ind. = cerastiifolia.
cerastiifolia, *Ramond, ex DC. Fl. Fr.* v. 783.—Pyren.
cerastioides, Pers. Syn. i. 502 = purpurascens.
cerastioides, Poir. Voy. Barb. ii. 166 = spathulata.
Cervianae, *Cham. & Schlecht. in Linnaea*, i. (1826) 52.—Chili.
Cherleria, *Hook. f. Stud. Fl.* ed. I. 60.—Europ.
Cherleriae, Fisch. ex DC. Prod. i. 409 = Stellaria petraea.
cherlerifolia, G. Don, in Loud. Hort. Brit. 478 = hirta.
cherlerioides, Vill. Prosp. 48 = lanceolata.
chilensis, *DC. in Mém. Soc. Phys. Genèv.* vi. (1833) 222.—Chili.
ciliaris, *Losc. Trat. Pl. Aragon.* i. 69, 79.—Hispan.
ciliata, *Linn. Sp. Pl.* 425.—Europ.; Am. bor.
ciliata, Sibth. & Sm. Fl. Graec. t. 438 = cretica.
ciliolata, *Edgew. & Hook. f. in Hook. f. Fl. Brit. Ind.* i. 240.—Reg. Himal.
cinerea, *DC. Fl. Fr.* v. 611.—Gallia.
clandestina, *Portenschl. Pl. Dalm.* 13.—Dalmatia.
coerulescens, Auct. vide caerulescens.
collina, Turcz. ex Ledeb. Fl. Ross. i. 349 = verna.
colorata, Turcz. ex Steud. Nom. ed. II. i. 124 = capillaris.
condensata, C. & J. Presl, Delic. Prag. i. 62 = recurva.
conferta, *Boiss. Diagn.* Ser. I. i. 51.—Graecia.
conferta, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 293.—Bolivia.
congesta, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 178.—Calif.
conica, *Boiss. Voy. Bot. Espagne*, ii. 98. t. 31.—Hispan.
conimbricensis, *Brot. Fl. Lusit.* ii. 200.—Lusitan.; Hispan.
conimbricensis, J. Gay, ex Gren. & Godr. Fl. Fr. i. 260 = controversa.
connata, *Rafin. Atl. Journ.* 151.—Am. bor.
controversa, *Boiss. Pugill. Pl. Nov. Hisp.* 100.—Hispan.; Gallia.
corsica, Steud. Nom. ed. II. i. 124 = balearica.
costata, *Bunge, ex Ledeb. Fl. Alt.* ii. 170.—Sibir.
cretica, *Spreng. Syst.* ii. 396.—Graecia; Cyprus; Creta.
cucubaloides, *Sm. Pl. Ic. Ined.* i. 17. t. 17.—Armenia.
dahurica, Fisch. ex DC. Prod. i. 402 = juncea.
davurica, Spreng. Syst. ii. 398 = Stellaria petraea.
debilis, *Hook. f. Fl. Brit. Ind.* i. 242.—Reg. Himal.
debilis, Turcz. ex Ledeb. Fl. Ross. i. 348 = verna.
decussata, Salisb. Prod. 299 = tetraquetra.
decussata, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 212.—Mexic.
deflexa, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 277 = graveolens.
densa, Kit. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 200 = gracilis.
densissima, *Wall. Cat.* n. 640.—Reg. Himal.
denticulata, *Phil. Fl. Atac.* 10.—Chili.
depauperata, *C. Gay, Fl. Chil.* i. 270.—Chili.
diandra, Guss. Fl. Sic. Prod. i. 515 = Spergularia diandra.
dianthoides, Hort. ex Ledeb. Fl. Ross. i. 365 = gypsophiloides.
dianthoides, *Sm. Pl. Ic. Ined.* i. 16. t. 16.—Armen.; Persia.
dichotoma, *Krock. Fl. Sil.* ii. 1. 55.—Silesia.
dichotoma, Moench, Meth. 224 = tenuifolia.
dicranoides, *H. B. & K. Nov. Gen. et Sp.* vi. 34.—Mexic.
diffusa, Ell. Sketch, i. 519 = lanuginosa.
digyna, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 201 = scopulorum.
divaricata, Adams, ex Ser. in DC. Prod. i. 405 = verna.
Douglasii, *Fenzl, ex Torr. & Gray, Fl. N. Am.* i. 674.—Calif.
drypidea, *Boiss. Diagn.* Ser. I. i. 49.—As. Min.
dubia, *Suter, Fl. Helv.* i. 266.—Helvet.
echinata, *Poir. Encyc.* vi. 377.—Europ. austr.
elegans, Cham. & Schlecht. in Linnaea, i. (1826) 57 = stricta.
elongata, Balbis, ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 200 = austriaca, grandiflora.
emarginata, *Brot. Fl. Lusit.* ii. 202.—Lusitan.; Maroceo.
erinacea, *Boiss. Voy. Bot. Espagne*, ii. 103.—Hispan.
erythrosepala, Boiss. Diagn. Ser. I. viii. 98 = setacea.

ARENARIA:—
extensa, *Dufour, in Ann. Sc. Gén. Phys.* vii. (1820) 291.—Hispan.
fasciculata, *Forsk. Fl. Aegypt. Arab.* 211.—Arabia.
fasciculata, *Gouan, Illustr.* 30.—Europ.
fasciculata, Hohen. Enum. Pl. Elisabethpol. 232 = sclerantha.
fasciculata, Pall. Ind. Taur.; Bieb. Casp. App. 209 = glomerata.
fasciculata, Sibth. & Sm. Fl. Graec. v. 30 = globulosa.
fastigiata, *Phil. in Linnaea*, xxviii. (1856) 673.—Chili.
fastigiata, Sm. Eng. Bot. xx. t. 1744 = rostrata.
fastigiata, *Schur, Enum. Pl. Transs.* 112.—Transsylv.
Fendleri, *A. Gray, Pl. Fendl.* 13.—Am. bor.
festucoides, *Benth. in Royle, Illustr. Bot. Himal.* 81. t. 21. f. 3.—Reg. Himal.
filicaulis, Fenzl, in Griseb. Spicil. Fl. Rumel. i. 203 = graveolens.
filifolia, Bieb. Fl. Taur. Cauc. i. 344 = graminifolia.
filifolia, Forsk. Fl. Aegypt. Arab. 211 = sctacea.
filiformis, Labill. Ic. Pl. Syr. dec. iv. 8 = picta.
fimbriata, Bieb. ex Ledeb. Fl. Ross. i. 352 = laricina.
fimbriata, Salzm. ex Kindberg, Monog. Lepigon. 22 = Spergularia fimbriata.
flaccida, [Clairv.] Man. Herb. 149 = Villarsii.
flaccida, Froel. ex Reichb. Fl. Germ. Excurs. 790 = lanceolata.
flaccida, Roxb. Hort. Beng. 34; Fl. Ind. ii. 447 = Spergula pentandra.
flaccida, Schleich. Cat. Pl. Helv. ed. III. 46 = verna.
flexuosa, *Rafin. Atl. Journ.* 151.—Am. bor.
floribunda, *C. Gay, Fl. Chil.* i. 269.—Chili.
foliosa, *Royle, ex Edgew. & Hook. f. in Fl. Brit. Ind.* i. 236.—Reg. Himal.
formosa, Fisch. ex DC. Prod. i. 402 = capillaris.
Franklinii, *Dougl. ex Hook. Fl. Bor. Am.* i. 101.—Am. bor.
frigida, *Rupr. Fl. Caucas.* 202.—Caucas.
frutescens, Kit. ex Schult. Oestr. Fl. ed. II. i. 667 = sctacea.
galioides, *Ser. in DC. Prod.* i. 410.—Hab.?
geniculata, *Poir. Voy. Barb.* ii. 166.—Reg. Mediterr.
Gerardi, Willd. Sp. Pl. ii. 729 = verna.
Giesekii, Hornem. Fl. Dan. t. 1518 = verna.
glabra, Balb. Misc. 21 = Villarsii.
glabra, Ell. Sketch, i. 520 = Stellaria uniflora.
glabra, *Michx. Fl. Bor. Am.* i. 274.—Am. bor.
glabra, Torr. Fl. U. St. 455 = groenlandica.
glabrata, Cham. & Schlecht. in Linnaea, i. (1826) 56 = verna.
glanduligera, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 240.—Reg. Himal.
glandulosa, *Jacq. Hort. Schoenb.* iii. 56.—Afr. austr.
globiflora, *Wall. Cat.* n. 639.—Reg. Himal.
globosa, Spreng. Syst. ii. 400 = globulosa.
globulosa, *Labill. Ic. Pl. Syr. dec.* iv. 6.—Oriens.
glomerata, *Bieb. Fl. Taur. Cauc.* i. 350.—Tauria.
glutinosa, Bieb. l. c. 344 = Holosteum umbellatum.
glutinosa, Boiss. in Ann. Sc. Nat. Sér. IV. ii. (1854) 247. t. 2 = Ledebouriana.
glutinosa, Willd. ex Ledeb. Fl. Ross. i. 353 = imbricata.
Gmelini, Bong. ex Ledeb. l. c. 351 = Villarsii.
Gmelini, Fisch. ex DC. Prod. i. 402 = Gypsophila Gmelini.
gorgonea, *J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 278.—Ins. Cap. Verd.
gothica, *Fries, Mant.* ii. 33.—Europ.
Gouffeia, Puel, in Duch. Rev. Bot. i. (Avr. 1846) 450 = controversa.
Gouffeia, *Chaub. Fl. Pélop.* n. 701. t. 24. f. 2.—Gall. occ.
gracilis, *Kit. ex Schult. Pl. Rar. Hung.* iii. 305.—Croat.; Dalmat.
graeca, Boiss. ex Nym. Consp. 114 = graveolens.
graminea, Auct. ex Steud. Nom. ed. II. i. 124 = grandiflora.
graminea, *C. A. Mey. Verz. Pfl. Cauc.* 220.—Reg. Caucas.
graminifolia, *Arduin. Animad. Alt.* 25.—Italia; Dalmat.
graminifolia, *Schrad. Hort. Goett.* i. 5.—Hungar.; Reg. Cauc.; Sibiria.
grandiflora, Coss. ex Willk. in Linnaea, xl. (1876) 117 = incrassata.
grandiflora, *Linn. Syst.* ed. X. 1034.—Europ.
grandiflora, Pall. Reise, iii. 34 = arctica.
grandis, H. B. & K. Nov. Gen. et Sp. vi. 30 = Spergularia grandis.

ARENARIA:—
graveolens, *Schreb. in Nov. Act. Nat. Cur.* iii. (1767) 478.—Graecia; As. Min.; Syria; Arab.
Griffithii, *Boiss. Fl. Orient.* i. 697.—Tibet. occ.; Afghan.
grineensis, Thomas, Suppl. Cat. (1842) = Villarsii.
groenlandica, *Spreng. Syst.* ii. 402.—Am. arct.
Guicciardii, *Heldr. in Boiss. Diagn.* Scr. II. v. 60.—Graecia.
Guilielmi-Waldemarii, Klotzsch, in Bot. Ergeb. Waldem. Reise, 142 = festucoides.
gypsophiloides, *Linn. Mant.* i. 71.—As. Min.; Persia.
gypsophiloides, Stev. ex Ledeb. Fl. Ross. i. 365 = cucubaloides.
gypsophiloides, Willd. ex Ledeb. l. c. 366 = juncea.
Haenkeana, Bartl. ex Presl, Rel. Haenk. ii. 15 = lateriflora.
Halleri, Raeusch. Nom. ed. III. 130 = tenuifolia.
Halleri, Vitm. Summa, iii. 109 = biflora.
halophila, Bunge, ex Ledeb. Fl. Alt. ii. 163 = Spergularia media.
Helmii, *Fisch. ex DC. Prod.* i. 404.—Sibir. ural.
herniarifolia, Desf. Fl. Atlant. i. 358 = geniculata.
heteromalla, Pers. Syn. i. 504 = setacea.
heterosperma, Guss. Fl. Sic. Syn. i. 501 = Spergularia media.
hirsuta, Bieb. Fl. Taur. Cauc. i. 349 = recurva.
hirta, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = cretica.
hirta, Worsmsk. in Fl. Dan. t. 1646 = verna.
hispanica, Spreng. Syst. ii. 396 = spathulata.
hispida, *Linn. Sp. Pl.* 425.—Europ.
hispida, Salzm. ex Presl, Fl. Sicul. 164 = Salzmanni.
Holostea, *Bieb. Fl. Taur. Cauc.* i. 345.—Reg. Caucas.
Holostea, Beaupr. ex Ser. in DC. Prod. i. 402 = rigida.
holosteoides, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 241.—Reg. Himal.
Hookeri, Nutt. ex Torr. & Gray, Fl. N. Am. i. 178 = Franklinii.
Howellii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 354.—Oregon.
humifusa, Wahlenb. Fl. Lapp. 129 = ciliata.
Huteri, *Kern. in Oestr. Bot. Zeitschr.* xxii. (1872) 368.—Europ.
hybrida, Vill. Prosp. 48 = tenuifolia.
imbricata, Banks, ex Steud. Nom. ed. II. i. 124, 125 = glabra.
imbricata, *Bieb. Fl. Taur. Cauc.* i. 344.—Reg. Caucas.
imbricata, Lag. Rodr. in Anal. Cienc. Nat. v. 1802 278 = tetraquetra.
imbricata, Rafin. in Med. Repos. N. York, v. (1808) 361 = squarrosa.
inamoena, Steud. Nom. ed. II. i. 125 = imbricata.
incrassata, *Lange, Pl. Nov. Hisp.* fasc. i. 3. t. 4. f. 2.—Hispan.
intricata, Dufour, ex Ser. in DC. Prod. i. 410 = montana.
intricata, *J. Gay, ex Willk. & Lange, Prod. Fl. Hisp.* iii. 617, in syn.—Hispan.
isaurica, *Boiss. Fl. Orient.* i. 695.—As. Min.
Jamesoniana, *Rohrb. in Linnaea*, xxxvii. (1871–72) 267.—Ecuador.
juncea, *Bieb. Fl. Taur. Cauc.* iii. 309.—Tauria.
juniperina, *Linn. Mant.* 72.—Oriens; Turkest.
juniperina, Pursh, Fl. Am. Sept. 319 = verna.
juniperina, Vill. Hist. Pl. Dauph. iii. 624 = grandiflora.
Jnressi, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 212.—Hispan.
Jussiei, *St. Hil. Fl. Bras. Med.* ii. 174.—Bras.
kansuensis, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 428.—China.
kashmirica, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 236.—Reg. Himal.
Koriniana, *Fisch. ex Ledeb. Fl. Ross.* i. 363.—Rossia.
Kotschyana, Fenzl, in Russegg. Reise, ii. 930 = tmolea.
lanceolata, *All. Fl. Pedem.* ii. 114.—Europ.
lanuginosa, *Rohrb. in Mart. Fl. Bras.* xiv. ii. 274.—Am. bor. et austr.
lapponica, *Spreng. Syst.* ii. 402.—Lappon.
laricifolia, Bieb. Fl. Taur. Cauc. i. 347 = pinifolia.
laricifolia, Lapeyr. (Abr. i. 255) ex Steud. Nom. ed. II. i. 124 = grandiflora.
laricifolia, *Linn. Sp. Pl.* 424.—Europ.
laricifolia, Pall. Ind. Taur. (51) ex Steud. Nom. ed. I. 66; ed. II. i. 127 = setacea.
laricifolia, Pursh, Fl. Am. Sept. i. 319 = arctica.
laricina, Cham. & Schlecht. in Linnaea, i. (1826) 57 = Spergula laricina.
lateriflora, *Linn. Sp. Pl.* 423.—Sibir.; Am. bor.

ARENARIA:—
lateriflora, Darl. Fl. Cestr. 277 = Stellaria borealis.
laxa, Fisch. ex Ser. in DC. Prod. i. 404 = grandiflora.
Laxmanni, Fisch. ex Ser. l. c. 402 = capillaris.
Ledebouriana, *Fenzl, in Russegg. Reise*, ii. 931.—Armenia.
leptoclados, Guss. Fl. Sic. Syn. ii. 284 = serpyllifolia.
leptoclados, *Boiss. Fl. Orient.* i. 701.—Europ.; Afr. bor.; Oriens.
leptophylla, *Cham. & Schlecht. in Linnaea*, v. (1830) 233.—Mexic.
Lessertiana, *Fenzl, in Russegg. Reise*, ii. 932.—Persia.
lesurina, *Loret, Herb. Loz. et Prost.* 13.—Gall. mer.
libanotica, Kotschy, in Boiss. Fl. Orient. i. 699 = setacea.
Liebmanniana, *Rohrb. in Linnaea*, xxxvii. (1871-72) 274.—Bras.
ligericina, Lec. & Lamotte, Cat. Pl. Cent. Fr. 104 = lesurina.
lignosa, *Phil. Fl. Atac.* 10.—Chili.
linearifolia, Poir. Encyc. vi. 366 = montana.
lineata, *C. A. Mey. ex Ledeb. Fl. Ross.* i. 351.—Rossia.
liniflora, Jacq. Fl. Austr. v. 22 = verna.
liniflora, Linn. Sp. Pl. ed. II. 607 = laricifolia.
liniflora, Willd. ex Spreng. Syst. ii. 399 = grandiflora.
littoralis, *Phil. in Linnaea*, xxviii. (1856) 673.—Chili.
littoralis, Salisb. Prod. 299 = peploides.
Lloydii, Jord. Pugill. 37 = serpyllifolia.
longifolia, DC. Prod. i. 402 = graminifolia.
longifolia, Bieb. Fl. Taur. Cauc. i. 345 = saxatilis.
Loscosii, *Texid. in Losc. Trat. Pl. Aragon*, ii. 28.—Hispan.
lychnidea, *Bieb. Fl. Taur. Cauc.* i. 347.—Reg. Caucas.
lychnidea, Willd. ex Ledeb. Fl. Ross. i. 367 = capillaris.
lycopodioides, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 212.—Mexic.
macradenia, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 367.—Am. bor.
macrocalyx, *Tausch, in Syll. Ratisb.* ii. (1828) 243.—Syria.
macrocarpa, Hornem. Hort. Hafn. i. 964 = canescens.
macrocarpa, Kit. ex Ser. in DC. Prod. i. 404 = laricifolia.
macrocarpa, *Pursh, Fl. Am. Sept.* i. 316.—Am. bor.
macrophylla, *Hook. Fl. Bor. Am.* i. 102.—As. or.; Am. bor. occ.
macrorrhiza, Guir. ex Willk. & Lange, Prod. Fl. Hisp. iii. 166 = Spergularia media.
macrorrhiza, Requien, ex Loisel. Nouv. Not. 22 = Spergularia macrorrhiza.
macrosepala, *Boiss. Diagn.* Ser. I. i. 52.—As. Min.
macrotheca, Hornem. ex Cham. & Schlecht. in Linnaea, i. (1826) 53 = Spergularia macrotheca.
Mandoniana, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 294.—Reg. And.
marginata, DC. Fl. Fr. v. 793 = Spergularia marginata.
marginata, *Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 212.—Sibir. or.
marina, All. Fl. Pedem. ii. 114 = Spergularia media.
marina, Pall. Reise, iii. 603 = Spergularia marina.
marina, Roth, Tent. Fl. Germ. i. 189 = Spergularia salina.
maritima, Steud. Nom. ed. II. i. 125 = Spergularia media.
Marschlinsii, Koch, in Flora, xxiv. (1841) ii. 509 = serpyllifolia.
media, DC. Prod. i. 401 = Spergularia marina.
media, Linn. Sp. Pl. ed. II. 606 = Spergularia media.
media, Poll. ex Steud. Nom. ed. II. ii. 617, in syn. = Spergula pentandra.
mediterranea, Ledeb. ex Link, Enum. Hort. Berol. i. 431 = mucronata.
melandryoides, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 241.—Reg. Himal.
mexicana, *Steud. Nom.* ed. II. i. 125.—Mexic.
Meyeri, *Edgew. & Hook. f. in Fl. Brit. Ind.* i. 236.—Oriens; Ind. bor. occ.
Meyeri, *Fenzl, ex Ledeb. Fl. Ross.* i. 368.—As. temp.
Michauxii, Hook. f. in Trans. Linn. Soc. xxiii. (1861) 287, 322 = stricta.
microphylla, *Phil. in Linnaea*, xxviii. (1856) 673.—Peruv.
minuta, *C. Gay, Fl. Chil.* i. 273.—Chili.
mixta, Lapeyr. Hist. Abr. Pl. Pyr. 255 = grandiflora.
modesta, *Dufour, in Ann. Sc. Gén. Phys.* vii. (1820) 291.—Hispan.

ARENARIA:—
molluginea, *Ser. in DC. Prod.* i. 400.—Mexic.
molluginifolia, *Pourr. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 201.—Hispan.
montana, *Linn. Cent.* i. 12; *Amoen. Acad.* iv. 272.—Gall.; Hisp.
montana, Sieber, ex Steud. Nom. ed. II. i. 125 = macrocalyx.
monticola, Buckl. in Proc. Acad. Sc. Philad. '1861.' (1862) 449 = Benthamii.
monticola, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 238.—Reg. Himal.
Moritzii, *Bruegg. in Zeitschr. Ferdinand. Tirol.* III. ix. (1860) 117.—Europ.
mucronata, DC. Fl. Fr. iv. 791 = rostrata.
mucronata, *Linn. Sp. Pl.* 424.—Ital.; Gall.
mucronata, Sibth. & Sm. Fl. Graec. t. 294 = tenuifolia.
multicaulis, Linn. Amoen. Acad. i. 162 = ciliata.
multicaulis, *Phil. in Linnaea*, xxviii. (1856) 673.—Chili.
Munbyi, *Munby, Cat. Pl. Algér.* 6.—Algeria.
muralis, Sieber, ex Spreng. Syst. ii. 397 = graveolens.
musciformis, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 150.—N. Granat.
musciformis, *Wall. Cat.* n. 6101.—Tibet.
muscipula, *Fenzl, Verbr. Alsin.* anal. tab. iii.—Afr. bor.
muscoides, *H. B. & K. Nov. Gen. et Sp.* vi. 34.—Am. austr.
muscorum, Adams, sec. Fisch. ex DC. Prod. i. 407 = uliginosa.
muscorum, *Fisch. ex DC. l. c.* 409.—Kamtschatka.
muscosa, *Med. in Act. Acad. Theod. Palat.* iii. *Phys.* (1775) 202. t. 12.—Europ.
mutabilis, Lapeyr. Hist. Abr. Pl. Pyr. 256 = fasciculata.
nana, *Boiss. & Heldr. Diagn.* Ser. I. viii. 103.—Creta.
nana, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 201 = muscoides.
nardifolia, Ledeb. ex Hook. Fl. Bor. Am. i. 98 (1830) = capillaris.
neelgherrensis, Wight & Arn. Prod. i. 43.—Reg. Himal.; Mont. Nilgiri.
nemorosa, H. B. & K. Nov. Gen. et Sp. vi. 35 = lanuginosa.
nemorosa, Steud. Nom. ed. II. i. 126. 127 = trinervia.
nepalensis, *Spreng. Syst. Cur. Post.* 181.—Nepal.
nervosa, Lam. Fl. Fr. iii. 36 = trinervia.
nevadensis, Boiss. Diagn. Ser. II. i. 90 = conferta.
nilgherrensis, Fenzl, var. α, γ, ex Boiss. Fl. Orient. i. 700, in syn. = Balansae.
nitida, *Rohrb. in Linnaea*, xxxvii. (1871-73) 249.—Peruv.
nodosa, Bory & Chaub. Fl. Pélop. 28. t. 15 = juniperina.
nodosa, Wallr. Sched. Crit. 200, in obs. = Sagina nodosa.
norvegica, *Gunn. Fl. Norv.* ii. 1100. t. 9. f. 7-9; *ex Hornem. in Fl. Dan.* t. 1269.—Scandin.
nudiuscula, Bertol. Bot. Misc. ii. 222. t. 8 = picta.
obtusa, All. Fl. Pedem. ii. 114 = polygonoides.
obtusa, *Torr. in Ann. Lyc. New York*, ii. (1828) 170.—Am. bor.
obtusiflora, *Kunze, in Flora*, xxix. (1846) 632.—Hispan.
occulta, Fisch. ex DC. Prod. i. 408 = polygonoides, arctica.
oligosperma, *C. Gay, Fl. Chil.* i. 272.—Chili.
orbicularis, *Vis. Fl. Dalm.* iii. 180.—Europ.
orbiculata, *Royle, ex Hook. f. Fl. Brit. Ind.* i. 240.—Reg. Himal.
Orbignyana, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 293.—Bolivia.
oreophila, *Hook. f. Fl. Brit. Ind.* i. 238.—Tibet.
otitoides, *Adams, ex Fisch. in DC. Prod.* i. 402.—Sibir.
oxypetala, *Sibth. & Sm. Fl. Graec. Prod.* i. 303.—Graccia; Creta.
palustris, *C. Gay, Fl. Chil.* i. 271.—Chili.
palustris, *S. Wats. Bot. Calif.* i. 70.—Calif.
pamphylica, *Boiss. & Heldr. Diagn.* Ser. I. viii. 102.—As. Min.
paniculata, *Bunge, in Ledeb. Fl. Alt.* ii. 168.—Sibir. alt.
papillosa, Steud. Nom. ed. II. i. 126 = graveolens.
paradoxa, Bartl. ex Presl, Rel. Haenk. ii. 15 = lanuginosa.
parviflora, *Benth. Pl. Hartw.* 163.—N. Granat.

ARENARIA:—
patagonica, *Phil. in Linnaea*, xxviii. (1856) 671.—Patagonia.
patula, Martr. Fl. Tarn, 107 = serpyllifolia.
patula, *Michx. Fl. Bor. Am.* i. 273.—Am. bor.
pauciflora, *Kit. in Linnaea*, xxxii. (1863) 510.—Hungar.
pedunculosa, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 294.—Bolivia.
pendula, *Waldst. & Kit. Pl. Rar.* i. 90. t. 87.—Banat.; Thracia.
pensylvanica, Muhl. Ind. Fl. Lanc. 169 = lateriflora?
pentagyna, *Moç. & Sesse, ex DC. Prod.* i. 395.—Mexic.
pentandra, Dufour, in Ann. Sc. Gén. Phys. vii. (1820) 292 = tenuifolia.
pentandra, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 429.—Soongaria.
pentandra, *Turcz. in Flora*, xvii. (1834) i. *Beibl.* 8.—Sibir. baical.
pentandra, Waldr. Sched. Crit. 206 = Spergula pentandra.
peploides, Lapeyr. Hist. Abr. Pl. Pyr. 251 = Polycarpon peploides.
peploides, *Linn. Sp. Pl.* 423.—Reg. bor. temp. et arct.
persica, Boiss. Diagn. Ser. I. i. 49 = Lessertiana.
peruviana, Poepp. ex Rohrb. in Mart. Fl. Bras. xiv. ii. 275 = lanuginosa.
Peyritschii, *Rohrb. in Linnaea*, xxxvii. (1871-73) 264.—Mexic.
pharnaceoides, Ser. in DC. Prod. i. 408 = picta.
physodes, *Fisch. ex DC. l. c.* 413.—Am. bor.
picta, *Sibth. & Sm. Fl. Graec. Prod.* i. 304.—As. Min.; Syria.
pilifera, Fisch. ex Ledeb. Fl. Ross. i. 352 = Spergula laricina.
pinifolia, *Bieb. Fl. Taur. Cauc.* i. 348.—Reg. Caucas.
Pissisi, *Phil. in Linnaea*, xxxiii. (1864-65) 20.—Chili.
Pitcheri, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 180.—Am. bor.
plantaginea, Mert. & Koch, in Roehl. Fl. Deutschl. iii. 273, in syn. = trinervia.
Poeppigiana, *Rohrb. in Linnaea*, xxxvii. (1871-72) 265.—Peruv.
polycnemifolia, *Boiss. Diagn.* Ser. I. i. 48.—Armenia; Persia.
polygonoides, *Wulf. in Jacq. Coll.* i. 241. t. 15.—Europ.; As. bor.
polytrichoides, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 237.—Reg. Himal.
Pomeli, *Munby, in Bull. Soc. Bot. Fr.* xi. (1864) 45.—Algeria.
Ponae, Reichb. Ic. Pl. Germ. ii. 26 = bavarica.
portulacea, Lam. Fl. Fr. iii. 38 = peploides.
Preslii, Nym. Consp. 113 = graminifolia.
procera, Spreng. ex Hornem. Hort. Hafn. 424 = graminifolia.
procumbens, Bove, ex Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 277 = graveolens.
procumbens, *Rupr. Fl. Cauc.* 202.—Caucas.
procumbens, Vahl, Symb. Bot. i. 50. t. 33 = geniculata.
propinqua, Richards. in Frankl. 1st Journ. ed. II. App. 17 = verna.
prostrata, Ser. in DC. Prod. i. 400 = Polycarpaea prostrata.
Przewalskii, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 428.—China.
pseudarmeriastrum, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 46.—Gallia.
pubescens, Presl, ex Steud. Nom. ed. II. i. 126 = pubigera.
pubescens, *Redowski, ex Ledeb. Fl. Ross.* i. 349.—Rossia.
pubescens, Steud. Nom. ed. I. 66 = Spergularia media.
pubescens, Urville, in Mém. Soc. Linn. Par. i. (1822) 306 = graveolens.
pubigera, *Steud. Nom.* ed. II. i. 126.—Sicilia.
pulchella, Hoffm. Hort. Mosq. 4 = Gypsophila elegans.
pulchra, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 212.—Armenia.
pulvinata, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 238.—Reg. Himal.
Pumilio, R. Br. in Parry, 2nd Voy. App. 391 = arctica.
pungens, *Clem. in Lag. Gen. et Sp. Nov.* 15.—Hispan.; Marocco.

ARENARIA:—
pungens, Steph. ex Ser. in DC. Prod. i. 403 = subulata.
pungens, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 179.—Am. bor. occ.
purpurascens, *Ramond, ex DC. Fl. Fr.* iv. 785.—Pyren.
purpurea, Pers. Syn. i. 504 = Spergularia purpurea.
purpurea, Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 211 = rubicunda.
Purshiana, Ser. in DC. Prod. i. 414 = Stellaria humifusa.
pusilla, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 367.—Am. bor.
pycnophylla, *Rohrb. in Linnaea*, xxxvii. (1871-72) 250.—Bolivia.
pyrenaica, Hort. ex Steud. Nom. ed. II. i. 126 = montana.
quadrivalvis, R. Br. in Parry, 1st Voy. App. 271 = verna.
querioides, Pourr. ex Willk. in Bot. Zeit. v. (1847) 239 = capitata.
racemosa, *Willk. l. c.*—Hispan.
Raddiana, Ser. in DC. Prod. i. 412 = Stellaria uliginosa.
radians, *Benth. Pl. Hartw.* 163.—N. Granat.
radicans, Hort. ex Steud. Nom. ed II. i. 126 = pendula.
radicans, Spreng. Syst. ii. 400 = Spergularia rubra.
Rafinesquiana, Ser. in DC. Prod. i. 409 = squarrosa.
Ramondi, Poir. Encyc. Suppl. v. 3 = cerastiifolia.
ramosissima, Bieb. ex Steud. Nom. ed. I. 67 = setacea.
ramosissima, *Willd. Enum. Hort. Berol. Suppl.* 24.—Hungar.
recurva, *All. Fl. Pedem.* ii. 113. t. 89. f. 3.—Europ. Oriens.
Redowskii, *Cham. & Schlecht. in Linnaea*, i. (1826) 58.—As. or.
repens, *Ser. in DC. Prod.* i. 410.—Hab.?
reptans, *Hemsl. Diag. Pl. Nov.* 22.—Mexic.
retusa, *Boiss. Voy. Bot. Espagne*, 99. t. 27.—Hispan.
rhodia, Boiss. Diagn. Ser. I. i. 52 = oxypetala.
rigida, *Bieb. Fl. Taur. Cauc.* i. 346.—Tauria.
rivularis, *Phil. Fl. Atac.* 10.—Chili.
Rohrbachiana, *Garcke, in Linnaea*, xxxvii. (1871–72) 311.—Mexic.
Rosani, *Tenore, Prod. Fl. Nap.* 26.—Ital.; Dalmat.
rosea, Presl, Fl. Sicul. 165 = geniculata.
rosea, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 365 = emarginata.
Rossii, R. Br. in App. Parry, Voy. 13 = stricta.
Roylei, Fenzl, ex Klotzsch, in Bot. Ergeb. Waldem. Reise, 142 = festucoides.
rostrata, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 481 = laricifolia.
rostrata, *Pers. Syn.* i. 504.—Europ. alp.
rotundifolia, *Bieb. Fl. Taur. Cauc.* i. 343.—Graecia; Reg. Cauc.; As. Min.
rubella, Sm. Engl. Bot. Suppl. t. 2538 = verna.
rubicunda, *Spreng. Syst.* ii. 399.—Sibir.
rubra, Linn. Sp. Pl. 423 = Spergularia rubra.
rupestris, *Labill. Pl. Syr. dec.* iv. 8. t. 4. f. 1.—Liban.
rupicola, Hook. f. Fl. Brit. Ind. i. 243 err. typ. = seq.
rupifraga, Fenzl, ex Ledeb. Fl. Ross. i. 780 = Thylacospermum rupifragum.
ruscifolia, *Poir. Encyc.* vi. 365.—Canada.
ruscifolia, Requien, in Guer. Vaucl. ed. II. 254 = cinerea.
sabulinea, *Griseb. in Russegg. Reise*, ii. 333.—Mesopot.
sajanensis, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 200.—Sibir. alt.
salina, Ser. in DC. Prod. i. 401 = Spergularia marina.
salsuginea, Bunge, ex Ledeb. Fl. Alt. ii. 163 = Spergularia diandra.
Salzmanni, *Presl, Fl. Sicul.* 163.—Sicil.
sancta, *Janka, in Termész. Füzetek*, vi. (1882) 165.—Europ. austr.
saponarioides, *Boiss. & Bal. Diagn.* Ser. II. vi. 35.—Phrygia.
saxatilis, Bout. ex Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 621 = obtusiflora.
saxatilis, Auct. ex Steud. Nom. ed. II. i. 127 = setacea.
saxatilis, C. C. Gmel. Fl. Bad. ii. 267 = verna.
saxatilis, *Linn. Sp. Pl.* 423.—Europ.; Sibir.
saxatilis, Vill. Fl. Delphin. 43 = rostrata.
Saxifraga, *Frivald. in Flora*, xix. (1836) ii. 434.—Rumelia.
saxosa, *A. Gray, Pl. Wright.* ii. 18.—N. Mexic.
scabra, *Poir. Encyc.* vi. 377.—Europ.
scandinavica, Spreng. Syst. ii. 399 = Stellaria biflora.
scario-a, *Boiss. in Tchihat. Asie Min. Bot.* i. 234.—Armenia.
scopulorum, *H. B. & K. Nov. Gen. et Sp.* vi. 31.—Peruv.
sedoides, *Froel. ex Koch, Syn. Fl. Germ.* ed. I. 114.—Tyrol.

ARENARIA:—
segetalis, Lam. Fl. Fr. iii. 43 = Spergularia segetalis.
sericea, Ser. in DC. Prod. i. 414 = purpurea.
serpens, Fisch. ex Ser. l. c. 404 = arctica.
serpens, *H. B. & K. Nov. Gen. et Sp.* vi. 32.—Ecuador.
serpyllifolia, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 619 = nevadensis.
serpylliifolia, *Linn. Sp. Pl.* 423.—Europ.; As. temp.; Ind. or.
serpyllifolia, Guss. Fl. Sic. Syn. i. 495 = leptoelados.
serpylloides, *C. Gay, Fl. Chil.* i. 271.—Chili.
setacea, Fisch. ex Ser. in DC. Prod. i. 402 = lychnidea.
setacea, Adams, ex Ser. l. c. 403 = capillaris.
setacea, *Thuill. Fl. Par.* ed. II. 220.—Reg. Mediterr.; Orient; Sibir.
setosa, Willd. ex Ledeb. Fl. Ross. i. 368 = capillaris.
sibirica, Pers. Syn. i. 504 = capillaris.
sipylea, Boiss. Diagn. Ser. I. i. 51 = graveolens.
sitchensis, D. Dietr. Syn. Pl. ii. 1565 = peploides.
Smithii, Steud. Nom. ed. II. i. 127 = globulosa.
soratensis, *Rohrb. in Linnaea*, xxxvii. (1871–72) 266.—Bolivia.
sordida, *Chaub. Fl. Pélop.* n. 699. t. 24. f. 1.—Graecia.
spathulata, *Desf. Fl. Atlant.* i. 358.—Hispan.; Afr. bor.
Spergula, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 293 = Spergularia media.
sperguloides, *Rafin. Atl. Journ.* 153.—Am. bor.
sperguloides, *Trevir. ex Steud. Nom.* ed. II. i. 127.—Chili.
sphaerocarpa, Martr. Fl. Tarn, 107 = patula.
sphaerocarpa, Tenore, Syll. Fl. Neap. iv. 222 = serpyllifolia.
spinuliflora, Ser. in DC. Prod. i. 406 = Habrosia spinuliflora.
squarrosa, *Michx. Fl. Bor. Am.* i. 273.—Am. bor.
stellarioides, Pers. Syn. i. 503 = austriaca.
stellarioides, *Willd. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 209.—Caucas.
stenocarpa, *Phil. Fl. Atac.* 10.—Chili.
stenopetala, Turcz. Cat. Baikal. n. 257, in Bull. Soc. Nat. Mosc. (1838) 89 = arctica.
stenophylla, Ledeb. Ind. Hort. Dorp.; Fl. Alt. ii. 165 = graminifolia.
Steveniana, *Boiss. Fl. Orient.* i. 692.—Armenia.
stolonifera, Vill. ex DC. Prod. i. 404 = grandiflora.
Stracheyi, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 240.—Tibet.
striata, Linn. Cent. ii. 17 = capillacea, cephaelotes, laricifolia, recurva, saxatilis, verna.
striata, *Vill. Hist. Pl. Dauph.* iv. 630. t. 47.—Gallia.
stricta, *Michx. Fl. Bor. Am.* i. 274.—Am. bor.
stygia, Boiss. & Heldr. Diagn. Ser. II. i. 91 = cretica.
subulata, *Ser. in DC. Prod.* i. 403.—Sibir.
subulifolia, Presl, Fl. Sicul. 162 = mucronata.
succulenta, Ser. in DC. Prod. i. 400 = Polycarpon suculentum.
sulcata, Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 212 = hirta.
Szowitsii, *Boiss. Fl. Orient.* i. 695.—Reg. Caucas.
tenella, *Kit. in Schult. Oestr. Fl.* ed. II. i. 662.—Hungar.
tenella, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 179.—Am. bor.
tenella, Turcz. ex Steud. Nom. ed. II. i. 127 = stricta.
tenera, *Edgew. in Hook. f. Fl. Brit. Ind.* i. 242.—Reg. Himal.
tenuiflora, *Schrad. ex Steud. Nom.* ed. II. i. 127.—Hab.?
tenuifolia, *Linn. Sp. Pl.* 424.—Europ.
tenuis, *J. Gay, in Walp. Ann.* vii. 300.—Hispan.
teres, Kit. ex Steud. Nom. ed. I. 67 = setacea.
teretifolia, *Phil. Fl. Atac.* 10.—Chili.
tetragyna, *Willd. ex Schlecht. Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 201.—Am. austr.
tetraquetra, *Linn. Sp. Pl.* 423.—Gallia.
tetrastieha, *Boiss. Diagn.* Ser. I. i. 50.—Persia.
thesiifolia, *Froel. ex Reichb. Ic. Fl. Germ.* v. t. 219.—Alp. Algoviae.
thymifolia, Pursh, Fl. Am. Sept. i. 317 = Stellaria humifusa.
thymifolia, *Sibth. & Sm. Fl. Graec. Prod.* i. 304.—Creta.
tmolea, *Boiss. Diagn.* Ser. I. i. 50.—As. Min.
tomentosa, *Willk. in Linnaea*, xxv. (1852) 15.—Hispan.
tremula, *Boiss. Diagn.* Ser. I. viii. 101.—Syria.
triandra, Schrank, Hort. Monac. t. 31 = mucronata.

ARENARIA:—
trichocalycina, *Tenore & Guss. ex Boiss. Fl. Orient.* i. 681.—Graecia.
trichotoma, Boiss. Diagn. Ser. II. v. 60 = Ledebouriana.
trichotoma, *Royle, ex Hook. f. Fl. Brit. Ind.* i. 235.—Reg. Himal.
triflora, Linn. Mant. 240 = grandiflora.
triflora, Vill. Prosp. 48 = Villarsii.
trinervia, *Linn. Sp. Pl.* 423.—Europ.
tschuktschorum, *Regel, in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 254.—Sibir.
ucranica, 'Spreng. in Ersch.' sec. Steud. Nom. ed. II. i. 124, 127 = graminifolia.
uliginosa, Schleich. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag. vii. (1813) 207 = stricta.
umbellata, [Clairv.] Man. Herb. 149 = Holosteum umbellatum.
umbellata, Sibth. & Sm. Fl. Graec. v. 28 = Holosteum tenerrimum.
umbellata, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 252.—As. Min.
umbrosa, *Bunge, in Ledeb. Fl. Alt.* ii. 173.—Sibir. alt.
uniflora, *Luce, Topog. Nachr. Oesel*, 141.—Ins. Ossil.
uniflora, Poir. Encyc. vi. 375 = recurva.
unilateralis, Steud. Nom. ed. II. i. 127 = segetalis.
uralensis, *Pall. ex Spreng. Syst.* ii. 396.—Sibir. ural.
varia, Dulac, Fl. Hautes-Pyr. 249 = serpyllifolia.
valentina, *Boiss. Diagn.* Ser. II. i. 90.—Europ.
verna, Georgi, Beschr. Russ. Reich. iii. iv. 979 = setacea.
verna, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 756 = Sagina subulata.
verna, *Linn. Mant.* 72.—Europ.; Am. bor.
verticillata, Willd. Sp. Pl. ii. 725 = Acanthophyllum verticillatum.
Villarsii, *Balb. ex Schlecht. in Ges. Naturf. Fr. Berl. Mag.* vii. (1813) 199.—Europ. med.
villosa, *Ledeb. in Mém. Acad. Pétersb.* v. (1812) 534.—Sibir. or.
villosa, Wulf. in Jacq. Coll. iv. 319 = Wulfenii.
violaeea, *Ledeb. in Mém. Acad. Pétersb.* v. (1812) 533.—Sibir.
viscida, Loisel. Not. 68 = serpyllifolia.
viscidula, Thuill. Fl. Par. i. 219 = tenuifolia.
viscosa, Fisch. ex Ser. in DC. Prod. i. 403 = subulata.
viscosa, Pers. Syn. i. 504 = tenuifolia.
viscosa, Pourr. ex Webb & Berth. Phyt. Canar. i. 149 = procumbens.
vulcanorum, *Maxim. in Franch. & Sav. Enum. Pl. Jap.* i. 52.—Japon.
vulgaris, Bernh. Syst. Verz. Erf. 259 = Stellaria media.
Wallichiana, Ser. in DC. Prod. i. 411 = serpyllifolia.
Wulfenii, *Steud. Nom.* ed. II. i. 128.—Alp. Carniol.

ARENBERGIA, Mart. & Gal. in Bull. Acad. Brux. xi. i. (1844) 373 = **Eustoma**, Salisb. (Gentian.).
glauca, Mart. & Gal. l. c. 374 = E. exaltatum.

ARENGA, Labill. in Mém. Inst. Par. iv. (1801) 269.
PALMAE, Benth. & Hook. f. iii. 917.
GOMUTUS, Correa, in Ann. Mus. Par. ix. (1807) 288. t. 24. f. 1.
SAGUERUS, Adans. Fam. ii. 24 (1763).
Griffithi, Seem. ex H. Wendl. in Kerch. Palm. 232 = saccharifera.
javanica, *Hort. ex H. Wendl. l. c.*—Java.
obtusifolia, *Mart. Hist. Nat. Palm.* 191. t. 147, 148.—Ind. or.; Malaya.
saccharifera, *Labill in Mém. Inst. Par.* iv. (1801) 269.—Malaya.
Westerhoutii, Griff. in Calc. Journ. Nat. Hist. v. (1845) 474 = obtusifolia.
Wightii, *Griff. l. c.* 475.—Ind. or.

ARETHUSA, Gronov. Fl. Virg. 184 (1748). *ORCHIDEAE*, Benth. & Hook. f. iii. 614.
CRYBE, Lindl. Introd. Nat. Syst. ed. II. 446 (1836).
EPIPACTIS, Adans. Fam. ii. 70 (1763).
ORCHIDION, Mitchell, in Act. Nat. Cur. viii. (1748) 218, App.
alaris, Raeusch. Nom. ed. III. 262 = Pterygodium catholicum.
alata, Poir. Encyc. Suppl. i. 443 = Pterostylis praecox.
bengalensis, Herb. Calc. ex Hemsl. in Journ. Linn. Soc. xx. (1884) 311 = Didymoplexis pallens.
biplumata, Linn. f. Suppl. 405 = Pipinnula bonariensis.
bulbosa, *Linn. Sp. Pl.* 950.—Am. bor.

ARETHUSA:—
capensis, Linn. Diss. Pl. Rar. Afr. 28; Amoen. Acad. vi. 111 = Disperis capensis.
catenata, Sm. Exot. Bot. ii. 89 = Caladenia carnea.
ciliaris, Linn. f. Suppl. 405 = Bartholina pectinata.
cordata, Poir. Encyc. Suppl. i. 444 = Disperis cordata.
crispa, Thunb. Prod. Pl. Cap. 3 = Corycium crispum.
cucullata, Poir. Encyc. Suppl. i. 444 = Disperis cucullata.
divaricata, Linn. Sp. Pl. 951 = Pogonia divaricata.
ecristata, Griff. Notul. iii. 378 = Didymoplexis pallens.
foliosa, *Walt. Fl. Carol.* 222.—Am. bor.
gentianoides, Sw. Fl. Ind. Occ. iii. 1436 = Pogonia gentianoides.
glutinosa, Blanco, Fl. Filip. ed. I. 641 = Geodorum semicristatum.
japonica, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858–9) 409.—Japon.
lutea, Gaudich. in Ann. Sc. Nat. Sér. I. v. (1825) 101 = Chloraea Gaudichaudii.
medeoloides, Pursh, Fl. Am. Sept. ii. 591 = Pogonia verticillata.
ophioglossoides, Linn. Sp. Pl. 951 = Pogonia ophioglossoides.
parviflora, Michx. Fl. Am. Bor. ii. 160 = Pogonia pendula.
pendula, Willd. Sp. Pl. iv. 82 = Pogonia pendula.
petraea, *Pers. Syn.* ii. 512.—Afr. trop.
picta, Anders. in Trans. Soc. Arts, xxv. (1807) 199 = Spiranthes picta.
plicata, Andr. Bot. Rep. v. 321 = Pogonia plicata.
racemosa, Walt. Fl. Carol. 222 = Ponthieva glandulosa.
rosea, *Benth. in Journ. Linn. Soc.* xviii. (1881) 348.—Mexic.
secunda, Thunb. Prod. Pl. Cap. 3 = Disperis secunda.
simplex, Thou. Orch. Afr. t. 24 = Pogonia Thouarsii.
spicata, Walt. Fl. Carol. 222 = Aplectrum hyemale.
tigridiaefolia, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 14.—Mexic.
trianthophoros, Sw. in Vet. Akad. Nya Handl. Stockh. (1800) 230 = Pogonia pendula.
verticillata, Muehl. ex Willd. Sp. Pl. iv. 81 = Pogonia verticillata.
villosa, Linn. f. Suppl. 405 = Disperis villosa.

ARETIA, Hall. Enum. Stirp. Helv. ii. 485 (1742); Linn. Sp. Pl. ed. I. 141 (1753) = **Androsace**, Linn. (Primul.).
alba, Rafin. Fl. Ludov. 34 = Andr. alba.
alpina, Linn. Sp. Pl. 141 = Andr. alpina.
alpina, Wulf. in Jacq. Fl. Austr. 36 = Andr. hedreantha.
argentea, Gaertn. f. Fruct. iii. 107 = Andr. imbricata.
baldensis, Turra, Fl. Ital. Prod. 63.—Italia.
brevis, Hegetschw. Fl. Schw. 190 = Andr. alpina.
brutia, Nym. Consp. 607 = Andr. hedreantha.
bryoides, Loisel. Fl. Gall. ed. I. 111 = Andr. bryoides.
cana, Willd. ex Steud. Nom. ed. II. i. 129, 527 = Draba aretioides.
Charpentieri, Heer, in Hegetschw. Fl. Schw. 987 = Andr. imbricata.
ciliata, Loisel. Fl. Gall. ed. I. 112 = Andr. alpina.
cylindrica, Loisel. l. c. 111 = Andr. cylindrica.
glacialis, Schleich. Cat. Pl. Helv. ed. IV. 8 = Andr. alpina.
Halleri, Linn. Sp. Pl. ed. II. 204, in syn. = Andr. carnea.
Hausmanni, Nym. Consp. 607 = Andr. Hausmanni.
hedraeantha, Nym. l. c. = Andr. hedreantha.
Heerii, Hegetschw. Fl. Schw. 188 = Andr. Hausmanni.
helvetica, Georgi, Beschreib. Russ. Reich. iii. IV. 760 = Andr. villosa.
helvetica, Murr. Syst. ed. XIII. 162 = Andr. bryoides.
hirtella, Nym. Consp. 607 = Andr. alpina.
longiflora, Fisch. ex A. DC. in DC. Prod. viii. 668 = Dionysia odora.
multiflora, Vand. Fasc. Pl. 8.—Lusit.?
pennina, Murith, Bot. Val. 53 = Andr. alpina.
pubescens, Loisel. Fl. Gall. ed. I. 111 = Andr. alpina.
pyrenaica, Loisel. l. c. = Andr. pyrenaica.
rubra, Schleich. ex Steud. Nom. ed. II. i. 128 = Andr. hedreantha.
tomentosa, Schleich. Cat. Pl. Helv. ed. III. 8 = Andr. imbricata, bryoides.
Vandeli, Turra, Fl. Ital. Prod. 63.—Italia.

ARETIA:—
Vitaliana, Murr. Syst. ed. XIII. 162 = Douglasia Vitaliana.
Wulfeniana, Sieber, ex Nym. Consp. 607 = Andr. hedreantha.

ARETIA, Link, Handb. ii. 411 (1829) = **Primula**, Linn.
auriculata, Link, l. c. = P. Auricula.
carniolica, Link, l. c. 412 = P. carniolica.
ciliata, Link, l. c. = P. ciliata.
crenata, Link, l. c. 411 = P. marginata?
Floerkeana, Link, l. c. 413 = P. Floerkeana.
glaucescens, Link, l. c. 412 = P. calycina.
glutinosa, Link, l. c. 413 = P. glutinosa.
integrifolia, Link, l. c. 412 = P. integrifolia.
minima, Link, l. c. 413 = P. minima.
Palinuri, Link, l. c. 411 = P. Palinuri.
spectabilis, Link, l. c. 412 = P. spectabilis.
truncata, Link, l. c. 413 = P. minima.
villosa, Link, l. c. 411 = P. villosa.

ARETIASTRUM, Spach, Hist. Vég. Phan. x. 304 (1841) = **Valeriana**, Linn.

ARGAN, Dryand. in Trans. Linn. Soc. ii. (1794) 225 = seq.

ARGANIA, Roem. & Schult. Syst. iv. 46 (1819).
SAPOTACEAE, Benth. & Hook. f. ii. 655.
VERLANGIA, Neck. Elem. ii. 125 (1790); Rafin. Sylva Tellur. 33 (1838).
Sideroxylon, *Roem. & Schult. l. c.* 502.—Marocco.

ARGELASIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 357 = **Genista**, Tourn. (Legum.).
corsica, Fourr. l. c. = G. corsica.
Scorpius, Fourr. l. c. = G. Scorpius.

ARGELIA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 331. t. 11, G = **Solenostemma**, Hayne (Asclep.).
Delilii, Decne. l. c. = S. Argel.

ARGEMONE, Tourn. ex Linn. Syst. ed. I. (1735).
PAPAVERACEAE, Benth. & Hook. f. i. 52.
ECHTRUS, Lour. Fl. Cochinch. 344 (1790).
alba, *Lestib. Bot. Belg.* ed. II. iii. 11. 133 (= seq.?).
alba, Rafin. Fl. Ludov. 83 = mexicana.
albiflora, Hornem. Hort. Hafn. 469 = mexicana.
armeniaca, Linn. Sp. Pl. 509 = Papaver armeniacum.
Barclayana, *Penny, ex Loud. Gard. Mag.* vi. (1830) 115.—Mexic.
cambrica, Desf. in Dict. Sc. Nat. ii. 481 = Meconopsis cambrica.
fruticosa, *Thurber, ex A. Gray, in Mem. Am. Acad.* N. S. v. (1855) (*Pl. Nov. Thurb.*) 306.—Mexic.
Georgiana, Croom, in Am. Journ. Sc. Ser., I. xxv. (1834) 75 = mexicana.
grandiflora, Sweet, Brit. Flow. Gard. iii. t. 226 = mexicana.
hispida, *A. Gray, Pl. Fendl.* 5.—Am. bor.
Hunnemanni, Otto, in Otto & Dictr. Allg. Gartenz. i. (1833) 298 = Hunnemannia fumariaefolia?
intermedia, Sweet, Hort. Brit. ed. II. 585 = Barclayana.
mexicana, *Linn. Sp. Pl.* 508.—Am. bor.; Mexic.
mucronata, Dum.-Cours. ex Steud. Nom. ed. II. i. 128 = mexicana.
munita, Dur. & Hilg. in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 37 = hispida.
ochroleuca, *Sweet, Brit. Fl. Gard.* iii. t. 242.—Mexic.
platyceras, *Link & Otto, Ic.* ii. 43.—Mexic.
pyrenaica, Linn. Sp. Pl. 509 = Papaver pyrenaicum.
rosea, *Hook. Bot. Misc.* ii. (1831) 207.—Chili.
sexvalvis, Stokes, Bot. Mat. Med. iii. 195 = mexicana.
spinosa, Moench, Meth. 227 = mexicana.
sulphurea, Sweet, ex Loud. Hort. Brit. 216 = ochroleuca.
versicolor, Salisb. Prod. 376 = mexicana.

ARGENOPE, Salisb. Gen. Pl. Frag. 102 (1866) = **Narcissus**, Linn. (Amaryll.).
serotina, Salisb. l. c. = N. serotinus.

ARGENTINA, Lam. Fl. Fr. iii. 118 (1778) = **Potentilla**, Tourn. (Rosac.).
rubra, Lam. l. c. 120 = P. palustris.
rupestris, Lam. l. c. = P. rupestris.
supina, Lam. l. c. 119 = P. supina.
vulgaris, Lam. l. c. = P. Anserina.

ARGITHAMNIA, Sw. Prod. Veg. Ind. Occ. 39 (1788), emend. in ej. Fl. i. 335 (1797). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 303.
APHORA, Nutt. in Trans. Am. Phil. Soc. Ser. II. v. 174 (1837).
ARGOTHAMNIA, Spreng. Syst. iii. 847 (1826).
ARGYROTHAMNIA, Muell. Arg. in Linnaea, xxxiv. (1865) 65.
ARGYTHAMNIA, P. Br. Hist. Jamaic. 338 (1756).
CHIROPETALUM, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 21.
CHLOROCAULON, Klotzsch, in Endl. Gen. Suppl. iv. III. 89 (1847).
DESFONTAENA, Vell. Fl. Flum. 95 (1825).
DITAXIS, A. Juss. Euph. Tent. 27. t. 7 (1824).
PHILYRA, Klotzsch, in Wiegm. Archiv, vii. (1841) 199.
SEROPHYTON, Benth. Bot. Voy. Sulph. 52 (1844).
STENONIA, Didrichs. in Kjoeb. Vidensk. Meddel. (1857) 146.
adenophora, *A. Gray, in Proc. Am. Acad.* viii. (1870) 294.—Mexic.
anisotricha, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 314.—Bras.
aphoroides, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 146.—Texas.
Berteroana, *Muell. Arg. l. c.* 151.—Chili.
boliviensis, *Muell. Arg. l. c.* 149.—Chili.
brasiliensis, *Muell. Arg. l. c.* 144.—Bras.
breviramea, *Muell. Arg. l. c.* 146.—Bolivia.
calycina, *Muell. Arg. in Flora*, lv. (1872) 10.—Ind. occ.
candicans, *Sw. Prod. Veg. Ind. Occ.* 39.—Ind. occ.
canescens, *Phil. f. Cat. Pl. Vasc. Chil.* 262.—Chili.
cantonensis, *Hance, in Journ. Bot.* xvi. (1878) 14.—China.
catamarcensis, *Hieron. in Bull. Acad. Nac. Cordova*, iv. (1881) 449.—Reg. Argent.
desertorum, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 310.—Bras.
dioica, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 145.—Bras.
fasciculata, *Muell. Arg. l. c.* 146.—Ind. occ.
Fendleri, *Muell. Arg. l. c.* 145.—Venezuela.
foliosa, *Muell. Arg. l. c.* 150.—Bras.
Gardneri, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 310.—Bras.
guatemalensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 145.—Mexic.
gymnadenia, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 316.—Bras.
herbacea, *Spreng. Syst.* iii. 848.—Am. bor.
heterantha, *Muell. Arg. in DC. Prod.* xv. II. 735.—Mexic.
humilis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 147.—Mexic.
illimaniensis, *Muell. Arg. l. c.* 146.—Mexic.
laevis, *Muell. Arg. l. c.* 147.—Mexic.
lanceolata, *Muell. Arg. l. c.* 148.—Calif.
lancifolia, *Muell. Arg. l. c.* 145.—Ind. occ.
lineata, *Baill. Adansonia*, iv. (1863–64) 288.—Bras.
melochiaeflora, *Muell. Arg. in DC. Prod.* xv. II. 736.—Bras.
mercurialina, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 148.—Texas.
mollis, *Baill. Adansonia*, iv. (1863–64) 289.—Bras.
montevidensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 147.—Reg. Argent.
neo-mexicana, *Muell. Arg. l. c.*—N. Mexic.
pauciflora, *Mirb. ex Steud. Nom.* ed. II. i. 128.—Am. austr.
Pavoniana, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 149.—Peruv.
pilosissima, *Muell. Arg. l. c.* 148.—Texas.
quinquecuspidata, *Muell. Arg. l. c.* 150.—Peruv.
Ruiziana, *Muell. Arg. l. c.* 151.—Peruv.
Schiedeana, *Muell. Arg. l. c.* 150.—Mexic.
sericea, *Griseb. Fl. Brit. W. Ind.* 44.—Ind. occ.
sericophylla, *A. Gray, in Bot. Calif.* ii. 70.—Calif.
serrata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 147.—Calif.
Simoniana, *Muell. Arg. l. c.* 145.—Bras.
sponiella, *Muell. Arg. l. c.* 148.—Peruv.
tricocca, *Muell. Arg. l. c.* 150.—Bras.
tricuspidata, *Muell. Arg. l. c.*—Chili.
tuberculata, *Muell. Arg. l. c.* 144.—China.

ARGOLASIA, Juss. Gen. 60 (1789) = **Lanaria**, Ait. (Haemodor.).
aurea, Pursh, Fl. Am. Sept. i. 224, in syn. = Lophiola aurea.
capensis, J. F. Gmel. Syst. ii. 560 = L. plumosa.
lanata, Lam. ex Poir. Encyc. Suppl. i. 448 = L. plumosa.
plumosa, Juss. Gen. ed. Usteri, 67 = L. plumosa.

ARGOPHILUM, Blanco, Fl. Filip. 186 (1837) = **Aglaia**, Lour. (Meliac.).
pinnatum, *Blanco, l. c.* 186 = Aglaia angustifolia.

ARGOPHYLLUM, Blanco, Fl. Filip. p. lxv (1837) = praec.

ARGOPHYLLUM, Forst. Char. Gen. 29. t. 15 (1776). *SAXIFRAGEAE*, Benth. & Hook. f. i. 646.
ellipticum, *Labill. Sert. Aust. Caled.* 39. t. 41.—N. Caled.
Lejourdani, *F. Mucll. Fragm.* iv. 33.—Austral.
nitidum, *Forst. Char. Gen.* 30.—N. Caled.
sericeum, Poir. Encyc. Suppl. i. 449 = nitidum.

ARGORIPS, Rafin. Alsog. Am. 13 (1838) = **Salix**, Tourn.
alba, Rafin. l. c. = S. alba.
cerulea, Rafin. l. c. = S. alba.

ARGOSTEMMA, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 324 (1824). *RUBIACEAE*, Benth. & Hook. f. ii. 54.
POMANGIUM, Reinw. in Syll. Ratisb. ii. (1828) 10.
acutum, Wall. ex Hook. f. Fl. Brit. Ind. iii. 43 = verticillatum.
angustifolium, *Miq. Fl. Ind. Bat.* ii. 163.—Malaya.
begoniaccum, *Miq. l. c.* 348.—Malaya.
borragineum, *Blume, in DC. Prod.* iv. 417.—Java.
calycinum, R. Br. ex Benn. Pl. Jav. Rar. i. 97 = Neurocalyx Wightii.
coenosciadicum, *Suring. in Arch. Néerl.* v. (1870) 116.—Java.
connatum, Dalz. in Hook. Kew Journ. iii. (1851) 345 = courtallense.
courtallense, *Arn. in Ann. Nat. Hist.* iii. (1839) 22.—Ind. or.
Elatostemma, *Hook. f. Fl. Brit. Ind.* iii. 45.—Malaya.
fulvum, *Zoll. & Moritzi, Syst. Verz. Zoll.* 61.—Java.
glaberrimum, Dalz. in Hook. Kew Journ. iii. (1851) 345 = verticillatum.
humile, *Benn. Pl. Jav. Rar.* i. 94.—Reg. Himal.
inaequale, *Benn. l. c.* 96.—Sumatra.
inaequilaterum, *Benn. l. c.* 95.—Java.
Junghuhnii, *Miq. Fl. Ind. Bat.* ii. 164.—Malaya.
khasianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 43.—Reg. Himal.
Korthalsii, *Miq. Fl. Ind. Bat.* ii. 165.—Malaya.
Kurzii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 45.—Burma.
laeve, *Benn. Pl. Jav. Rar.* i. 96.—Java.
laevigatum, *Benn. l. c.*—Java.
Lobbii, *Hook. f. Fl. Brit. Ind.* iii. 44.—Burma.
longifolium, *Benn. Pl. Jav. Rar.* i. 97.—Sumatra.
macrosepalum, *Miq. Fl. Ind. Bat.* ii. 349.—Malaya.
microcnide, *Miq. l. c. Suppl.* 539.—Sumatra.
montanum, *Blume, in DC. Prod.* iv. 418.—Java.
Necsianum, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) [*Pl. Meyen.*] 349.—Ins. Philipp.
ncurocalyx, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 229.—Malaya.
oligantha, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) ii. 131.—Ins. Andam.
ophirense, *Maing. ex Hook. f. Fl. Brit. Ind.* iii. 45.—Malaya.
Parishii, *Hook. f. l. c.* 44.—Burma.
parvifolium, *Benn. Pl. Jav. Rar.* i. 96.—Malaya.
pauciflorum, *Blume, in DC. Prod.* iv. 418.—Java.
pedunculosum, *Miq. Fl. Ind. Bat.* ii. 349.—Malaya.
pictum, *Korth. in Neder. Kruid. Arch.* ii. ii. (1851) 122.—Sumatra.
pictum, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 327.—Penang.
pulchrum, *Korth. in Neder. Kruid. Arch.* ii. ii. (1851) 122.—Sumatra.
pumilum, *Benn. Pl. Jav. Rar.* i. 95.—Ind. or.; Afr. trop.
rostratum, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 326.—Reg. Himal.
sarmentosum, *Wall. l. c.* 324.—Reg. Himal.

ARGOSTEMMA:—
soneriloides, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 310.—Ins. Andaman.
spinulosum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 46.—Singapore.
subinaequale, *Benn. Pl. Jav. Rar.* i. 96.—Java.
tavoyanum, *Wall. ex Benn. l. c.* 94.—Burma.
Teysmannianum, *Miq. Fl. Ind. Bat.* ii. 349.—Java.
timorense, *Benn. Pl. Jav. Rar.* i. 95.—Timor.
uniflorum, *Blume, in DC. Prod.* iv. 418.—Java.
unifolium, *Benn. Pl. Jav. Rar.* i. 94.—Penang.
unifolium, Kurz, in Journ. As. Soc. xlvi. (1877) ii. 131 = Kurzii.
verticillatum, Wall. Cat. n. 8394. partim = khasianum.
verticillatum, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 325.—Ind. or.
Wallichii, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. 350 (1843) [Pl. Meyen.].—Ins. Philipp.

ARGOTHAMNIA, Spreng. Syst. iii. 847 (1826) = **Argithamnia**, Sw. (Euphorb.).

ARGUZIA, Amm. ex Steud. Nom. ed. I. 68 (1821); Stev. in Bull. Soc. Nat. Mosc. (1851) i. 558 = **Tournefortia**, Linn. (Borag.).
cimmerica, Stev. l. c. 560 = T. sibirica.
Messerschmidia, Stev. l. c. = T. sibirica.
montana, Steud. Nom. ed. I. 68 = T. Arguzia.
repens, Rafin. Sylva Tellur. 167.—Sibir.?
rosmarinifolia, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 559 = T. sibirica.

ARGYLIA, D. Don, in Edinb. Phil. Journ. ix. (1823) 260. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1043.
adscendens, *DC. Prod.* ix. 235.—Chili.
australis, *Phil. in Anal. Univ. Chil.* (1873) 512.—Chili.
Bustillosii, *Phil. in Linnaea*, xxix. (1857–58) 13.—Chili.
canescens, *D. Don, in Edinb. Phil. Journ.* (*Apr.–June* 1829) 88.—Chili.
chrysantha, *Phil. in Linnaea*, xxxiii. (1864–65) 180.—Chili.
cremophila, *Phil. l. c.*—Chili.
Feuillei, *DC. Prod.* ix. 235.—Chili.
geranioides, *DC. l. c.*—Chili.
glabriuscula, *Phil. in Linnaea*, xxxiii. (1864–65) 179.—Chili.
glutinosa, *Phil. Fl. Atac.* 36.—Chili.
huidobriana, *Clos, in C. Gay, Fl. Chil.* iv. 411.—Chili.
incana, *Phil. Fl. Atac.* 36.—Chili.
lutea, *Phil. in Anal. Univ. Chil.* (1873) 511.—Chili.
potentillaefolia, *DC. Prod.* ix. 235.—Chili.
puberula, *DC. l. c.*—Chili.
radiata, *D. Don, in Edinb. Phil. Journ.* ix. (1823) 260.—Peruv.
radiata, Endl. Ic. n. 4112 = canescens.
tomentosa, *Phil. Fl. Atac.* 36.—Chili.
trifoliata, DC. Prod. ix. 237 = uspallatensis.
uspallatensis, *DC. l. c.* 236.—Chili.
villosa, *Phil. in Anal. Univ. Chil.* (1873) 512.—Chili.
viridis, *Phil. in Linnaea*, xxxiii. (1864–65) 181.—Chili.

ARGYRA, Noronha, ex Baill. Adansonia, i. (1860–61) 162 = **Croton**, Linn. (Euphorb.).
fasciculata, Noronha, ex Baill. l. c. = C. Noronhae.

ARGYRANTHEMUM, Webb, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 258 (1836–50) = **Chrysanthemum**, Linn. (Compos.).
anethifolium, Webb, l. c. 265–267 = C. anethifolium.
coronopifolium, Webb, l. c. 272 = C. coronopifolium.
crithmifolium, C. Bolle, Ind. Sem. Hort. Berol. App. (1861) 5 = C. anethifolium.
dissectum, Lowe, Man. Fl. Mad. i. 464 = C. dissectum.
foeniculaceum, Webb, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 262 = C. anethifolium.
frutescens, Sch. Bip. l. c. 264 = C. frutescens.
gracile, Sch. Bip. l. c. 261 = C. gracile.
haematomma, Lowe, Man. Fl. Mad. i. 463 = C. haematomma.
jacobaeifolium, Webb, in Webb & Berth. Phyt. Canar. ii. 251.—Ins. Canar.
ochroleucum, Webb, l. c. 259 = C. ochroleucum.

ARGYRANTHEMUM:—
pinnatifidum, Webb, in Webb & Berth. Phyt. Canar. ii. 269 = C. pinnatifidum.
Webbii, Sch. Bip. in Webb & Berth. l. c. = C. Webbii.

ARGYRANTHUS, Neck. Elem. i. 75 (1790) = **Helipterum**, DC. (Compos.).

ARGYREIA, Lour. Fl. Cochinch. 134 (1790). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 869.
ADAMBOË, Rafin. Fl. Tellur. iv. 73 (1836).
ARGYREON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120.
SAMUDRA, Rafin. Fl. Tellur. iv. 72 (1836).
abyssinica, *Choisy, in DC. Prod.* ix. 332.—Abyss.
acuta, Benth. Fl. Hongk. 237, in syn. = Championi.
acuta, *Lour. Fl. Cochinch.* 135.—China; Ind. or.
aggregata, Arn. ex Choisy, Conv. Or. 45 = Lettsomia aggregata.
alata, *Montrous. in Mém. Acad. Lyon*, x. (1860) 236.—N. Caled.
alulata, Miq. Fl. Ind. Bat. ii. 587 = Ipomoea Turpethum.
ampla, Choisy, Conv. Or. 38 = Roxburghii.
arborea, *Lour. Fl. Cochinch.* 135.—China.
argentata, *Miq. Fl. Ind. Bat.* ii. 588.—Malaya.
argentea, *Arn. ex Choisy, Conv. Or.* 36.—Ind. or.
barbigera, Choisy, l. c. 42 = Lettsomia barbigera.
Bona-nox, Sweet, Hort. Brit. ed. I. 289 = Rivea hypocrateriformis.
bracteata, *Choisy, Conv. Or.* 39.—Ind. or.
capitata, Arn. ex Choisy, l. c. 41 = Lettsomia strigosa.
capitata, Brand. For. Fl. 343 = Lettsomia Thomsoni.
capitata, Kurz, For. Fl. Burm. ii. 216 = Lettsomia peguensis.
Championi, Benth. Fl. Hongk. 236 = Lettsomia Championi.
Choisyana, *Regel & Koern. in Ind. Sem. Hort. Petrop.* (1858) 40.—Hab.?
Choisyana, *Wight, ex Hook. f. in Fl. Brit. Ind.* iv. 190.—Ind. or.
courtallensis, Wight, ex Choisy, in DC. Prod. ix. 329 = malabarica.
cuneata, *Ker-Gawl. in Bot. Reg.* t. 661.—Ind. or.
cymosa, *Sweet, Hort. Brit.* ed. I. 289.—Ind. or.; Malaya.
Daltoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 190.—Ind. or.
elliptica, Arn. ex Choisy, Conv. Or. 35 = Lettsomia elliptica.
festiva, Wall. Pl. As. Rar. 68. t. 76.—China.
floribunda, *Wall. ex Voigt, Hort. Suburb. Calc.* 352, *nomen*.—Ind. or.
fulgens, *Choisy, Conv. Or.* 33.—Ind. or.
glabra, *Choisy, ex Zoll. Syst. Verz. Ind. Archip.* 128, 130.—Malaya.
Guichenotii, *Choisy, Conv. Or.* 41.—Malaya.
hancorniaefolia, Gardn. ex Thw. Enum. Pl. Zeyl. 210 = Lettsomia hancorniaefolia.
hirsuta, *Wight & Arn. in Nov. Act. Nat. Cur.* xviii. (1836) i. 356.—Ind. or.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 185.—Reg. Himal.
involucrata, *C. B. Clarke, l. c.* 187.—Ind. or.
lanceolata, *Choisy, Conv. Or.* 39.—Burma; Malaya.
laurifolia, Voigt, Hort. Suburb. Calc. 351 = Lettsomia elliptica.
Lawii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 190.—Ind. or.
Leschenaultii, *Choisy, Conv. Or.* 31.—Ind. or.
Leschenaultii, Thw. Enum. Pl. Zeyl. 209 = sericea.
malabarica, *Arn. ex Choisy, Conv. Or.* 38.—Ind. or.; Malaya.
megapotamica, *Griseb. in Goett. Abh.* xxiv. 1879) 263.—Reg. Argent.
mollis, *Choisy, Conv. Or.* 38.—Ind. or.; Malaya.
multiflora, Voigt, Hort. Suburb. Calc. 352 = Roxburghii.
nellygherya, *Choisy, Conv. Or.* 32.—Ind. or.
nervosa, *Boj. Hort. Maurit.* 224.—Ins. Mascar.
nitida, *Choisy, Conv. Or.* 40.—Ind. or.; Ins. Philip.
oblonga, *Benth. Bot. Voy. Sulph.* 133.—Mexic.
obtecta, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 186.—Burma; Malaya.
obtusifolia, Lour. Fl. Cochinch. 135 = obtecta.
ornata, Sweet, Hort. Brit. ed. I. 289 = Rivea ornata.

ARGYREIA:—
osyrensis, Choisy, in DC. Prod. ix. 334 = Lettsomia aggregata.
pallida, *Choisy, Conv. Or.* 34.—Burma.
pilosa, *Wight & Arn. in Nov. Act. Nat. Cur.* xviii. (1836) i. 356.—Ind. or.
pomacea, *Choisy, Conv. Or.* 32.—Ind. or.
populifolia, *Choisy, l. c.*—Zeylan.
populifolia, Choisy, in DC. Prod. ix. 329 = Hookeri.
Reinwardtiana, *Miq. Fl. Ind. Bat.* ii. 589.—Malaya.
Roxburghii, *Arn. ex Choisy, Conv. Or.* 37.—Ind. or.
rubicunda, Choisy, l. c. 44 = Lettsomia rubicunda.
sericea, *Dalz. & Gib. Bomb. Fl.* 169.—Ind. or.
sericea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = argentea.
setosa, Arn. ex Choisy, Conv. Or. 43 = Lettsomia setosa.
speciosa, *Boj. Hort. Maurit.* 224.—Maurit.
speciosa, *Sweet, Hort. Brit.* ed. I. 289.—Ind. or.; Malaya.
splendens, *Sweet, l. c.*—Ind. or.
tiliaefolia, *Wight, Ic.* iv. i. 12. t. 1358 (sphalm. *tillaefolia*). Ind. or.
tomentosa, *Choisy, Conv. Or.* 44.—Burma.
uniflora, Sweet, Hort. Brit. ed. I. 289 = Rivea hypocrateriformis.
venusta, Choisy, Conv. Or. 37 = argentea.
Wallichii, *Choisy, l. c.* 39.—Ind. or.
zeylanica, Kurz, For. Fl. Brit. Burma, ii. 215, excl. syn. = Lettsomia Kurzii.
zeylanica, Voigt, Hort. Suburb. Calc. 351 = pomacea.

ARGYRELLA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 300. t. 6 = **Dissotis**, Benth. (Melast.).
canescens, Harv. Gen. S. Afr. Pl. ed. II. 113 = D. incana.
incana, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 300. t. 6. f. 7 = D. incana.
phaeotricha, Naud. l. c. = D. phaeotricha.

ARGYREON, St. Lag. in Ann. Soc. Bot. Lyon, vii. 120 (1880) = **Argyreia**, Lour. (Convolv.).

ARGYREXIAS, Rafin. Fl. Tellur. iv. 86 (1836) = **Echium**, Tourn. (Borag.).
candicans, Rafin, l. c. = E. candicans.

ARGYROCHAETA, Cav. Ic. iv. 54. t. 378 (1791) = **Parthenium**, Linn. (Compos.).
bipinnatifida, Cav. l. c. = P. Hysterophorum.

ARGYROCOMA, Rafin. New Fl. Am. iv. 41 (1836) = **Anychia**, Rich. (Illecebr.).
imbricata, Rafin. l. c. 43 = Anychia dichotoma.

ARGYROCOME, Gaertn. Fruct. ii. 410 (1791); Schrank, in Denkschr. Acad. Wiss. Muench. viii. (1824) 146 et 163 = **Helipterum**, DC. (Compos.).
apiculata, Cass. in Dict. Sc. Nat. xxxix. 40 = Helichrysum apiculatum.
bracteata, Cass. l. c. = Helichrysum rutilans?
ericoides, Poir. Tabl. Encyc. iii. 270 = Helichrysum ericoides.
ferruginea, Schrank, in Denkschr. Acad. Wiss. Muench. viii. (1824) 163 = Helip. speciosissimum.
imbricata, Buek, Ind. DC. Prod. ii. 15 = Metalasia uniflora.
retorta, Cass. in Dict. Sc. Nat. xxxiv. 39 = Helichrysum stoloniferum.
retorta, Gaertn. Fruct. ii. t. 167 = Helichrysum retortum.
seminula, Schrank, in Denkschr. Acad. Wiss. Muench. viii. (1824) 164 = Helip. speciosissimum.
semipapposum, Cass. in Dict. Sc. Nat. xxxix. 40 = Helichrysum semipapposum.
Staehelina, Schrank, in Denkschr. Acad. Wiss. Muench. viii. (1824) 165 = Helichrysum squamosum.
vermiculata, Poir. Tabl. Encyc. iii. 270 = Metalasia uniflora.
vulnerata, Schrank, in Denkschr. Acad. Wiss. Muench. viii. (1824) 164 = H. canescens.

ARGYRODENDRON, Klotzsch, in Peters, Reise Mossamb. Bot. 100 (1861) = **Croton**, Linn. (Euphorb.).
bicolor, Klotzsch, l. c. 102 = C. pulchellus.
ovatum, Boiv. ex Baill. Adansonia, i. (1861) 152 = C. Bernieri.
Petersii, Klotzsch, in Peters, Reise Mossamb. Bot. i. 101 (= Combreti sp.).—Afr. trop.

ARGYRODENDRON, F. Muell. Fragm. i. 2 (1858) = **Tarrietia**, Blume (Stercul.).
trifoliolatum, F. Muell. l. c. = T. Argyrodendron.

ARGYROGLOTTIS, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 83 = **Helichrysum**, Vaill. (Compos.).
turbinata, Turcz. l. c. 84 = H. Argyroglottis.

ARGYROLOBIUM, Eckl. & Zeyh. Enum. 184 (1835). *LEGUMINOSAE*, Benth. & Hook. f. i. 480.
CHAMAECYTISUS, Vis. Fl. Dalm. 272. t. 55 (1850).
CHASMONE, E. Mey. Comm. Pl. Afr. Austr. 71 (1836).
DIOLOTUS, Tausch, in Flora, xxv. (1842) 284.
GAMOCHILUM, Walp. in Linnaea, xiii. (1839) 509.
TEPHROTHAMNUS, Sweet, Hort. Brit. ed. II. 126 (1830), nomen prius.
TRICHASMA, Walp. in Linnaea, xiii. (1839) 510.
abyssinicum, *Jaub. & Spach, Illustr.* i. 115.—Abyss.
aequinoctiale, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 46.—Afr. trop.
Andrewsianum, *Steud. Nom.* ed. II. i. 129.—Afr. austr.
angustifolium, *Eckl. & Zeyh. Enum.* 188.—Afr. austr.
arabicum, *Jaub. & Spach, Illustr.* i. 115; *Boiss. Diagn.* Ser. I. ii. 13.—Arab.
argenteum, *Eckl. & Zeyh. Enum.* 184.—Afr. austr.
argenteum, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 464 = Linnacanum.
ascendens, *Walp. in Linnaea*, xiii. (1839) 507.—Afr. austr.
baptisioides, *Walp. l. c.* 506.—Afr. austr.
barbatum, *Walp. Rep.* ii. 845.—Afr. austr.
biflorum, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
Bottae, *Jaub. & Spach, Illustr.* i. 115.—Arab.
calycinum, Ball, in Journ. Linn. Soc. xvi. (1878) 396, in obs. = Linnaeanum.
calycinum, *Jaub. & Spach, Illustr.* i. 115.—Dalmat.; Reg. Caucas.; Persia.
candicans, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
collinum, *Eckl. & Zeyh. l. c.*—Afr. austr.
comanthum, Vog. ex Steud. Nom. ed. II. i. 129 = baptisioides.
connatum, *Harv. in Harv. & Sond. Fl. Cap.* 72.—Afr. austr.
crassifolium, *Eckl. & Zeyh. Enum.* 184.—Afr. austr.
crinitum, *Walp. in Linnaea*, xiii. (1839) 506.—Afr. austr.
crotalarioides, *Jaub. & Spach, Illustr.* i. 114, 116.—Syria.
dasycarpum, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. ii. 46 = A. Schimperianum.
divaricatum, Jaub. & Spach, Illustr. i. 116 = flaccidum.
dorycnoides, *Baker, in Oliver, Fl. Trop. Afr.* ii. 45.—Afr. trop.
emirnense, *Baker, in Journ. Linn. Soc.* xx. (1883) 125.—Madag.
fallax, Ball, in Journ. Bot. xi. (1873) 302 = Linnaeanum.
filiforme, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
flaccidum, *Jaub. & Spach, Illustr.* i. 115.—Ind. or.
goodioides, *Walp. Rep.* ii. 845.—Afr. austr.
gracile, Fenzl, in Flora, xxvi. (1843) i. 394 = crotalaroides.
grandiflorum, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 29.—Afr. bor.
incanum, Eckl. & Zeyh. Enum. 185 = crassifolium.
involucratum, *Harv. in Harv. & Sond. Fl. Cap.* 75.—Afr. austr.
Kotschyi, *Boiss. Diagn.* Ser. I. vi. 32.—Persia.
Krebsianum, *Presl, Bot. Bemerk.* 139.—Afr. austr.
lanceolatum, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
Linnaeanum, *Walp. in Linnaea*, xiii. (1839) 508.—Reg. Mediterr.
longifolium, *Walp. Rep.* ii. 844.—Afr. austr.
lotoides, *Bunge, ex Trautv. in Act. Hort. Petrop.* ii. (1873) 519.—Armenia.
lotoides, *Harv. in Harv. & Sond. Fl. Cap.* 595.—Afr. austr.
microphyllum, *Ball, in Journ. Bot.* xi. (1873) 303.—Marocco.
modestum, Hochst. in Flora, xxvii. (1844) i. 100 = abyssinicum.
molle, *Eckl. & Zeyh. Enum.* 187.—Afr. austr.
obcordatum, *Steud. Nom.* ed. II. i. 130.—Afr. austr.
obovatum, *Eckl. & Zeyh. Enum.* 185.—Afr. austr.
obsoletum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 70.—Afr. austr.
ornithopodioides, Jaub. & Spach, Illustr. i. 116 = roseum.
patens, *Eckl. & Zeyh. Enum.* 185.—Afr. austr.

ARGYROLOBIUM:—
pauciflorum, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
pedunculare, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 348.—Madag.
petiolare, *Steud. Nom.* ed. II. i. 130.—Afr. austr.
Petitianum, *A. Rich. Tent. Fl. Abyss.* i. 163.—Abyss.
pilosum, *Harv. in Harv. & Sond. Fl. Cap.* 71.—Afr. austr.
polyphyllum, *Eckl. & Zeyh. Enum.* 184.—Afr. austr.
prostratum, *Boiss. Diagn.* Ser. I. ii. 13.—Assyria.
pumilum, *Eckl. & Zeyh. Enum.* 185.—Afr. austr.
ramosissimum, *Baker, in Oliver, Fl. Trop. Afr.* ii. 46.—Afr. trop.
remotum, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 162.—Abyss.
roseum, *Jaub. & Spach, Illustr.* i. 116.—Persia; Beluchist.
rupestre, *Walp. in Linnaea*, xiii. (1839) 508.—Afr. austr.
Saharae, *Pomel, Nouv. Mat. Fl. Atl.* 177.—Algeria.
Sandersoni, *Harv. in Harv. & Sond. Fl. Cap.* ii. 594.—Afr. austr.
Schimperianum, *Hochst. in Flora*, xxvii. (1844) i. 100.—Afr. trop.
sericeum, *Eckl. & Zeyh. Enum.* 184.—Afr. austr.
sessilifolium, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 175.—Turcia.
speciosum, *Eckl. & Zeyh. Enum.* 187.—Afr. austr.
splendens, *Walp. Rep.* ii. 845.—Afr. austr.
stenophyllum, *Boiss. Fl. Orient.* ii. 32.—Afghan.
stipulaceum, *Ball, in Journ. Bot.* xi. (1873) 302.—Marocco.
stipulaceum, *Eckl. & Zeyh. Enum.* 187.—Afr. austr.
strictum, *Steud. Nom.* ed. II. i. 130.—Afr. austr.
Sutherlandi, *Harv. in Harv. & Sond. Fl. Cap.* 594.—Afr. austr.
syriacum, Boiss. Diagn. Ser. I. ii. 13 = crotalarioides.
tenue, *Walp. Rep.* i. 632.—Afr. austr.
trigonelloides, *Jaub. & Spach, Illustr.* i. 116, 118.—Persia.
tuberosum, *Eckl. & Zeyh. Enum.* 188.—Afr. austr.
umbellatum, *Vog. ex Steud. Nom.* ed. II. i. 130.—Afr. austr.
uniflorum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 72.—Afr. austr.
uniflorum, *Jaub. & Spach, Illustr.* i. 115.—Oriens.
velutinum, *Eckl. & Zeyh. Enum.* 186.—Afr. austr.
venustum, *Eckl. & Zeyh. l. c.* 185.—Afr. austr.
virgatum, *Baker, in Oliver, Fl. Trop. Afr.* ii. 46.—Afr. trop.

ARGYROPHANES, Schlecht. in Linnaea, xx. (1847) 596 = **Helichrysum**, Vaill. (Compos.).
Behrii, Schlecht. l. c. = H. Baxteri.

ARGYROPHYLLUM, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 150 (1873) = **Soaresia**, Sch. Bip. (Compos.).
ovali-ellipticum, Pohl, ex Baker, l. c. = S. velutina.

ARGYROPHYTON, Hook. Comp. Bot. Mag. ii. (1836) 163 = **Argyroxiphium**, DC. (Compos.).
Douglasii, Hook. l. c. = Argyroxiph. sandwicense.

ARGYROPSIS, M. Roem. Syn. Ensat. 125 (1847) = **Zephyranthes**, Herb. (Amaryll.).
candida, M. Roem. l. c. = Z. candida.

ARGYRORCHIS, Blume, Orch. Archip. Ind. 120. t. 31, 36 (1858). *ORCHIDEAE*, Benth. & Hook. f. iii. 506.
javanica, *Blume, l. c.*—Java.

ARGYROTHAMNIA, Muell. Arg. in Linnaea, xxxiv. (1865) 144 = **Argithamnia**, Sw. (Euphorb.).

ARGYROXIPHIUM, DC. Prod. v. 668; et Mém. Comp. t. 8 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 393.
ARGYROPHYTON, Hook. Comp. Bot. Mag. ii. (1836) 163.
macrocephalum, *A. Gray, in Proc. Am. Acad.* ii. (1852) 160.—Ins. Sandvic.
sandwicense, *DC. Prod.* v. 668.—Ins. Sandvic.

ARGYTAMNIA, Duchesne, in Dict. Sc. Nat. iv. 98 (1804) = **Argithamnia**, Sw. (Euphorb.).

ARGYTHAMNIA, P. Br. Hist. Jamaic. 338 (1756), sphalm. = **Argithamnia**, Sw. (Euphorb.).

ARHYNCHIUM, Lindl. & Paxt. Flow. Gard. i. (1850–51) 142 = **Arachnanthe**, Blume (Orchid.).

ARIA, Jacq. f. Cat. Hort. Vindob. (1826) ex Host, Fl. Austr. ii. 7 (1831) = **Pyrus**, Linn. (Rosac.).
alnifolia, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 166 = P. alnifolia.
alpina, M. J. Roem. Syn. iii. Rosifl. 124 = P. alpina.
ambigua, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 165.—Hab.?
arguta, M. J. Roem. Syn. iii. Rosifl. 129 = P. intermedia.
betulaefolia, M. J. Roem. l. c. 124 = P. betulaefolia.
Chamaemespilus, Host, Fl. Austr. ii. 8 = P. Chamaemespilus.
crenata, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 162 = P. crenata.
cuspidata, M. J. Roem. Syn. iii. Rosifl. 125 = P. cuspidata.
Decaisneana, Lavall. Arbor. Segrez. 61.—Hab.?
decipiens, M. J. Roem. Syn. iii. Rosifl. 129 = P. rotundifolia.
edulis, Guimp. ex Lavall. Arbor. Segrez. 63 = *Decaisneana*.
edulis, M. J. Roem. Syn. iii. Rosifl. 124 = P. edulis.
flabellifolia, M. J. Roem. l. c. 127 = Crataegus corymbosa.
graeca, M. J. Roem. l. c. = P. Aria.
Hostii, Carr. Rev. Hortic. (1877) 210 = P. Hostii.
Hostii, Jacq. f. Cat. Hort. Vindob. (1826) ex Host, Fl. Austr. ii. 8 = P. intermedia.
intermedia, Schur, Enum. Pl. Transs. 207 = P. intermedia.
japonica, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 164 = P. japonica.
kamaonensis, M. J. Roem. Syn. iii. Rosifl. 130 = P. lanata.
kumaonensis, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 162 = praec.
lanata, Decne. l. c. 163 = P. lanata.
latifolia, M. J. Roem. Syn. iii. Rosifl. 128 = P. rotundifolia.
Mougeoti, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 378 = P. Aria.
nivea, Host, Fl. Austr. ii. 8 = P. Aria.
obtusata, M. J. Roem. Syn. iii. Rosifl. 125 = P. obtusata.
rotundifolia, M. J. Roem. l. c. 129 = P. rotundifolia.
scandica, M. J. Roem. l. c. 127 = P. intermedia.
semilobata, M. J. Roem. l. c. 129 = P. rotundifolia.
Szovitsii, Decne. in Nouv. Arch. Mus. Par. Sér. I. x. (1874) 165.—Hab.?
tiliaefolia, Decne. l. c. 166 = P. alnifolia.
vestita, M. J. Roem. Syn. iii. Rosifl. 125 = P. vestita.

ARIETINUM, Beck, Bot. N. and Midl. U. St. 352 (1833) = **Cypripedium**, Linn. (Orchid.).
americanum, Beck, l. c. = C. Arietinum.

ARILLARIA, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 70 = **Ormosia**, Jack (Legum.).
robusta, Kurz, l. c. 71 = O. robusta.

ARILLASTRUM, Panch. ex Brongn. & Gris, in Bull. Soc. Bot. Fr. xiv. (1867) 253 = **Spermolepis**, Brongn. & Gris (Myrtac.).
gummiferum, Panch. l. c. = S. gummiferum.

ARINEMIA, Rafin. Sylva Tellur. 49 (1838) = **Ilex**, Tourn.
lanceolata, Rafin. l. c. = I. lanceolata.

ARIOCARPUS, Scheidw. in Bull. Acad. Brux. v. (1838) 491 = **Mammillaria**, Haw. (Cactac.).
retusus, Scheidw. l. c. 492. t. 1 = M. prismatica.

ARIODENDRON, Meissn. Gen. Comm. 301 ('1836-43'), sphalm. = AGRIODENDRON, Endl. = **Aloë**, Linn. (Liliac.).

ARIONA, Pers. Syn. i. 212 (1805) = **Arjona**, Comm. (Santal.).

ARIOPSIS, J. Grah. Cat. Bomb. Pl. 252 (1839). *AROIDEAE*, Benth. & Hook. f. iii. 973.
peltata, *J. Grah. l. c.*—Reg. Himal.
protanthera, N. E. Br. in Rep. Roy. Gard. Kew, (1877) 51 = peltata.

ARISACONTIS, Schott, in Bonplandia, v. (1857) 129. *AROIDEAE*? Benth. & Hook. f. iii. 963.
Chamissonis, *Schott, l. c.*—Ins. Radack.

ARISAEMA, Mart. in Flora, xiv. (1831) II. 459. *AROIDEAE*, Benth. & Hook. f. iii. 965.
AMIDENA, Rafin. Fl. Tellur. iv. 15 (1836).
abbreviatum, *Schott, in Oestr. Bot. Zeitschr.* (1857) 382.—Reg. Himal.
affine, Schott, in Bonplandia, vii. (1859) 27 = concinnum.
album, *N. E. Br. in. Journ. Linn. Soc.* xviii. (1882) 247.—Siam.
alienatum, Schott, in Bonplandia, vii. (1859) 26 = concinnum.
amplissimum, *Blume, Rumphia*, i. 110.—Japon.
amurense, *Maxim. Prim. Fl. Amur.* 264.—Reg. Amur.
angustatum, *Franch. & Sav. Enum. Pl. Jap.* ii. 507.—Japon.
atrorubens, *Blume, Rumphia*, i. 97.—Am. bor.
Boscii, Blume, l. c. 104 = A. Dracontium.
Bottae, *Schott, Prod. Aroid.* 42.—Arabia.
brasilianum, Blume, Rumphia, i. 97 = atrorubens.
caudatum, *Engl. in DC. Monog. Phan.* ii. 559.—Ind. or.
cochinchinense, Blume, Rumphia, i. 107 = Pinella Wawrae.
commutatum, Schott, in Bonplandia, vii. (1859) 26 = tortuosum.
concinnum, *Schott, l. c.* 27.—Reg. Himal.
consanguineum, Schott, l. c. = erubescens.
cornutum, *Schott, l. c.*—Reg. Himal.
costatum, *Mart. in Flora*, xiv. (1831) II. 459.—Reg. Himal.
Cumingii, *Schott, Syn. Aroid.* 28.—Ins. Philipp.
curvatum, Hook. Bot. Mag. t. 5931 = tortuosum.
curvatum, *Kunth, Enum. Pl.* iii. 20.—Reg. Himal.
cuspidatum, *Engl. in DC. Monog. Phan.* ii. 536.—Penang.
decipiens, *Schott, in Oestr. Bot. Zeitschr.* (1857) 373.—Java.
dolosum, Schott, in Bonplandia, vii. (1859) 26 = intermedium.
Dracontium, *Schott, Meletem.* i. 17.—Am. bor.
echinatum, *Schott, l. c.*—Reg. Himal.
eminens, Schott, in Oestr. Bot. Zeitschr. (1857) 357 = speciosum.
enneaphyllum, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 352.—Abyss.
erubescens, *Schott, Meletem.* i. 17.—Reg. Himal.
exile, *Schott, in Bonplandia*, vii. (1859) 26.—Reg. Himal.
fallax, Schott, in Miq. Ann. Mus. Lugd. Bat. i. 123 = filiforme.
filicaudatum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1882) 253.—Zeylan.
filiforme, *Blume, Rumphia*, i. 102. t. 28.—Java.
filiforme, Thw. Enum. Pl. Zeyl. 334 = neglectum.
fimbriatum, *Mast. in Gard. Chron.* (1884) II. 680.—Ins. Philipp.
flavum, *Schott, Prod. Aroid.* 40.—Arabia.
Franchetianum, *Engl. Bot. Jahrb.* i. (1881) 487.—Tibet.
fraternum, Schott, in Bonplandia, vii. (1859) 26 = Leschenaultii.
galeatum, *N. E. Br. in Gard. Chron.* (1879) II. 102.—Reg. Himal.
gracile, Kunth, Enum. Pl. iii. 21 = Typhonium gracile.
Griffithii, *Schott, Syn. Aroid.* 26.—Reg. Himal.
hastatum, Blume, Rumphia, i. 96 = atrorubens.
helleborifolium, Schott, Syn. Aroid. 29 = tortuosum.
heptaphyllum, *Blume, Rumphia*, i. 109.—Ind. or.
heterophyllum, Blume, l. c. 110 = Thunbergii.
Hookerianum, Schott, in Oestr. Bot. Wochenbl. (1857) 334 = Griffithii.
Hookeri, Schott, Gen. Aroid. 6. f. 11–19 = Griffithii.
Huegelii, Schott, Syn. Aroid. 27 = Leschenaultii.
inclusum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1881) 249.—Java.
intermedium, *Blume, Rumphia*, i. 102.—Reg. Himal.
Jacquemontii, *Blume, l. c.* 95.—Reg. Himal.
japonicum, *Blume, l. c.* 106.—Japon.
laminatum, Benth. Fl. Hongk. 342 = penicillatum.
laminatum, *Blume, Rumphia*, i. 99. t. 27.—China.
latisectum, Blume, l. c. 110 = japonicum.
Leschenaultii, *Blume, l. c.* 93.—Ind. or.
lobatum, *Engl. Bot. Jahrb.* i. (1881) 487.—Tibet.
Loureiri, Blume, Rumphia, i. 108 = Pinellia tuberifera.
macrospathum, *Benth. Pl. Hartw.* 52.—Mexic.

ARISAEMA:—
macrourum, Kunth, Enum. Pl. iii. 644 = Pinellia tuberifera.
Makoyanum, Kunth, in Ind. Sem. Hort. Berol. (1845) 9 = filiforme.
mirabile, Schott, in Oestr. Bot. Zeitschr. (1857) 366 = speciosum.
Murrayi, Hook. Bot. Mag. t. 4388 = Jacquemontii.
neglectum, *Schott, in Bonplandia*, vii. (1859) 26.—Zeylan.
nepenthoides, *Mart. in Schott, Meletem.* i. 17.—Reg. Himal.
ochraceum, *Schott, in Bonplandia*, vii. (1859) 27.—Reg. Himal.
ornatum, *Miq. Fl. Ind. Bat.* iii. 220.—Sumatra.
papillosum, Steud. ex Schott, Prod. Aroid. 46 = Leschenaultii.
penicillatum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1881) 248. t. 5.—Hongkong.
pentaphyllum, *Schott, Meletem.* i. 17.—Ind. or.; China.
Plukenetii, Blume, Rumphia, i. 110 = A. Dracontium.
praecox, De Vriese, ex C. Koch, Allg. Gartenz. (1857) 85 = ringens.
propinquum, Schott, in Oestr. Bot. Zeitschr. (1857) 333 = intermedium.
pulchrum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1881) 252. t. 6.—Ind. or.
pumilum, Blume, Rumphia, i. 107 = trilobatum.
Pythonium, Blume, l. c. 108 = Zomicarpa Pythonium.
quinatum, *Schott, Syn. Aroid.* 28.—Am. bor.
ringens, *Schott, Meletem.* i. 17.—Japon.
Roxburghii, Kunth, Enum. Pl. iii. 18 = cuspidatum.
Schimperianum, *Schott, in Bonplandia*, vii. (1859) 27.—Abyss.
serotinum, Miq. ex Franch. & Sav. Enum. Pl. Jap. ii. 4 = Sierotium.
serratum, Schott, Meletem. i. 17 = japonicum.
Sieboldii, De Vriese, ex C. Koch, Allg. Gartenz. (1857) 85 = ringens.
Sierotium, *Siebold, ex Regel, in Ind. Sem. Hort. Petrop.* (1865) 44.—Japon.
sikokianum, *Franch. & Sav. Enum. Pl. Jap.* ii. 507.—Japon.
speciosum, *Mart. in Schott, Meletem.* i. 17.—Reg. Himal.
Steudelii, *Schott, in Bonplandia*, vii. (1859) 26.—Ind. or.
stictopodum, Miq. Fl. Ind. Bat. iii. 219 = filiforme.
Stracheyanum, Schott, in Oestr. Bot. Zeitschr. (1857) 333 = intermedium.
Tatarinowii, *Schott, in Bonplandia*, vii. (1859) 27.—China.
Thunbergii, *Blume, Rumphia*, i. 105.—Japon.
tortuosum, *Schott, Meletem.* i. 17.—Reg. Himal.
tortuosum, Steud. ex Engl. in DC. Monog. Phan. ii. 544 = Steudelii.
tripartitum, *Engl. l. c.* 538.—Japon.
triphyllum, *Schott, Meletem.* i. 17.—Am. bor.
utile, *Hook. f. ex Engl. in DC. Monog. Phan.* ii. 537.—Reg. Himal.
verrucosum, *Schott, in Oestr. Bot. Wochenbl.* 1857 341.—Reg. Himal.
vituperatum, Schott, in Bonplandia, vii. (1859) 28 = erubescens.
Wightii, Hook. f. Bot. Mag. t. 5507 = neglectum.
Wightii, Schott, in Bonplandia, vii. (1859) 26 = Jacquemontii.
zebrinum, Hort. ex Nichols. Dict. Gard. i. 112 = = triphyllum.

ARISARON, Adans. Fam. ii. 470 (1763) = **Arisarum**, Tourn. (Aroid.).

ARISARUM, Hall. in Rupp. Fl. Jen. ed. II. 251 (1745) = **Calla**, Rich. (Aroid.).

ARISARUM, [Tourn.] Targ. Toz. in Anal. Mus. Fis. Fir. ii. II. (1810) 67. *AROIDEAE*, Benth. & Hook. f. iii. 965.
ARISARON, Adans. Fam. ii. 470 (1763).
BALMISA, Lag. Gen. et Sp. Nov. 17 (1816).
HOMAIDA, Rafin. Fl. Tellur. iii. 63 (1836).
aspergillum, Dun. in Mem. Acad. Montp. Sect. Sc. i. (1847) 8 = simorrhinum.
australe, Rich. in Guill. Arch. i. (1838) 20 = vulgare.
azoricum, Schott, in Oestr. Bot. Wochenbl. (1857) 190 = vulgare.
Balansae, Schott, l. c. = vulgare.
Clusii, Schott, Prod. Aroid. 28 = vulgare.

ARISARUM:—
crassifolium, Schott, in Bonplandia, ix. (1861) 369 = vulgare.
Forbesii, Schott, in Oestr. Bot. Wochenbl. (1857) 190 = vulgare.
hastatum, *Pomel, Nouv. Mat. Fl. Atl.* 390.—Algeria.
italicum, Rafin. Fl. Tellur. iii. 63 = Arum italicum.
Jacquini, Schott, Prod. Aroid. 22 = vulgare.
Libani, Schott, l. c. 21 = vulgare.
maculatum, Rafin. Fl. Tellur. iii. 63 = Arum maculatum.
oligocarpum, *Pomel, Nouv. Mat. Fl. Atl.* 390.—Algeria.
ovatum, Rafin. Fl. Tellur. iii. 63 = Lagenandra toxicaria.
pictum, Rafin. l. c. = Arum pictum.
proboscideum, *Savi, Osserv. Div. Piante*, 6.—Italia.
serpentrium, Rafin. Fl. Tellur. iii. 63 = Arisarum vulgare.
Sibthorpii, Schott, Prod. Aroid. 21 = vulgare.
simorrhinum, *Dur. in Duch. Rev. Bot.* i. (1845–46) 360.—Algeria.
subalpinum, Kotschy, ex Engl. in DC. Monog. Phan. ii. 563 = vulgare.
subexertum, Webb & Berth. Phyt. Canar. iii. 293 = vulgare.
tingitanum, Schott, Prod. Aroid. 22 = vulgare.
Veslingii, Schott, l. c. 20 = vulgare.
vulgare, *Targ. Toz. in Anal. Mus. Fis. Fir.* ii. II. (1810) 67.—Reg. Mediterr.

ARISTAEA, A. Rich. in Orb. Dict. ii. 136 (1842) = **Aristea**, Soland. (Irid.).

ARISTARIA, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 296 = **Anthistiria**, Linn. (Gramin.).
barbata, Jungh. l. c. 297 = Anth. arguens.

ARISTEA, [Soland. in] Ait. Hort. Kew. i. 67 (1789); iii. 506. *IRIDEAE*, Benth. & Hook. f. iii. 701.
CLEANTHE, Salisb. in Trans. Hort. Soc. i. (1812) 312.
GENLISIA, Reichb. Consp. 60 (1828).
NIVENIA, Vent. Dec. Gen. Nov. 5 (1808).
WREDOWIA, Eckl. Verz. 16 (1827).
africana, *Hoffmgg. Verz. Pfl.* 40.—Afr. trop.
alata, *Baker, in Journ. Linn. Soc.* xxi. (1885) 405.—Afr. trop.
anceps, *Eckl. ex Klatt, in Linnaea*, xxxiv. (1865–66) 550.—Afr. austr.
angolensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 270.—Afr. trop.
angustifolia, *Baker, in Journ. Linn. Soc.* xx. (1884) 269.—Madag.
bracteata, Pers. Syn. i. 41 = capitata.
capitata, *Ker-Gawl. in Bot. Mag.* t. 605.—Afr. austr.
cladocarpa, *Baker, in Journ. Linn. Soc.* xx. (1884) 268.—Madag.
coerulea, Vahl, Enum. ii. 124 = capitata.
compressa, *Buching. ex Baker, in Journ. Linn. Soc.* xvi. (1878) 111.—Natal.
cyanea, [*Soland. in*] *Ait. Hort. Kew.* i. 67.—Afr. austr.
dichotoma, Eckl. ex Klatt, in Linnaea, xxxiv. (1865–66) 551 = Eckloni.
dichotoma, *Ker-Gawl. Gen. Irid.* 13.—Afr. austr.
diffusa, *Eckl. Verz.* 16, nomen.—Afr. austr.
Eckloni, *Baker, in Journ. Linn. Soc.* xvi. (1878) 112.—Afr. austr.
eriophora, Pers. Syn. i. 41 = cyanea.
fruticosa, *Pers. l. c.*—Afr. austr.
glauca, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 381.—Afr. austr.
intermedia, Eckl. Verz. 16 = dichotoma.
juncifolia, *Eckl. l. c.; Baker, in Journ. Bot.* xiv. (1876) 267.—Afr. austr.
Kitchingii, *Baker, in Journ. Linn. Soc.* xx. (1884) 269.—Madag.
lacera, Stokes, Bot. Comment. i. 224 = cyanea.
lugens, Hort. ex Steud. Nom. ed. II. i. 130 = melaleuca.
madagascariensis, *Baker, in Journ. Bot.* xiv. (1876) 267.—Madag.
major, Andr. Bot. Rep. t. 160 = capitata.
melaleuca, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 236.—Afr. austr.
pusilla, *Ker-Gawl. l. c.*—Afr. austr.
racemosa, *Baker, in Journ. Bot.* xiv. (1876) 267.—Afr. austr.
repens, *A. Dietr. Sp. Pl.* ii. 516.—Afr. austr.
schizolaena, *Harv. ex Baker, in Journ. Bot.* xiv. (1876) 267.—Afr. austr.

ARISTEA:—
spathacea, Spreng. Syst. i. 158 = Bobartia spathacea.
spicata, Pers. Syn. i. 41 = capitata.
spiralis, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 236.—Afr. austr.
torulosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 382.—Afr. austr.
umbellata, *Spreng. Syst.* i. 158.—Afr. austr.
Wredowia, *Steud. Nom.* ed. II. i. 130.—Afr. austr.

ARISTEGA, Miers, Contrib. Bot. iii. 375 (1871). *MENISPERMACEAE.*
laevifolia, *Miers, l. c.* 376. t. 51.—Ind. or.

ARISTELLA, Bertol. Fl. Ital. i. 690 (1833) = **Stipa**, Linn. (Gram.).
bromoides, Bertol. l. c. = S. Aristella.
longiflora, Regel, in Act. Hort. Petrop. vii. (1880) 645.—As. temp.

ARISTIDA, Linn. Sp. Pl. ed. I. 82 (1753). *GRAMINEAE*, Benth. & Hook. f. iii. 1140.
ARTHRATHERUM, Beauv. Agrost. 32. t. 8. f. 8 (1812).
CHAETARIA, Beauv. l. c. 30. t. 8. f. 5, 6 (1812).
CURTOPOGON, Beauv. l. c. 32. t. 8. f. 7 (1812).
KIELBOUL, Adans. Fam. ii. 31 (1763).
MOULINSIA, Rafin. in Ser. Bull. Bot. i. (1830) 221.
ORTACHNE, Nees, in Seem. Bot. Voy. Herald, 225 (1852–57).
SCHISTACHNE, Figar. & De Not. in Mem. Accad. Torin. Scr. II. xii. (1852) 252.
STIPAGROSTIS, Nees, in Linnaea, vii. 290 (1832).
STREPTACHNE, H. B. & K. Nov. Gen. et Sp. i. 124. t. 40 (1815).
TRIXOSTIS, Rafin. in Ser. Bull. Bot. i. (1830) 221.
abyssinica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 134 = coerulescens.
acutiflora, *Trin. & Rupr. l. c.* 167.—Aegypt.; Arabia.
adoënsis, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 370.—Abyss.
Adscensionis, *Linn. Sp. Pl.* 82.—Reg. temp. et calid.
Adscensionis, Walt. Fl. Carol. 74 = oligantha.
Adscensionis, *Sw. Obs.* 40 = cognata.
aequiramea, *Schlecht. in Linnaea*, xxii. (1849) 343.—Texas.
aethiopica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 134 = Adscensionis.
affinis, Kunth, Rev. Gram. i. 61 = purpurascens.
americana, Linn. f. Pl. Jam. Pugill. 7; Syst. ed. X. 879 = Bouteloua bromoides, juncifolia.
amplissima, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 155.—Afr. trop.
Antillarum, *Poir. Encyc. Suppl.* i. 451.—Ind. occ.
Antoniana, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
arabica, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 135 = Adscensionis.
arenaria, *Gaudich. in Freyc. Voy. Bot.* 407.—Austral.
arundinacea, *Linn. Mant.* 146.—Ind. or.
avenaria, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 68 = setifolia.
barbata, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 532.—Mexic.
barbicollis, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 152.—Abyss.
basiramea, *Vasey, in Coult. Bot. Gaz.* ix. (1884) 76.—Am. bor.
Behriana, *F. Muell. in Trans. Vict. Inst.* i. (1855) 44.—Austral.
Beyrichiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 104.—Am. centr.
bipartita, *Steud. Syn. Pl. Gram.* 140.—Afr. austr.
brachypoda, *Tausch, in Flora*, xix. (1836) 506.—Aegypt.; Arabia.
brachyptera, *Coss. & Bal. in Bull. Soc. Bot. Fr.* v. (1858) 169.—Afr. bor.
brevifolia, *Steud. Nom.* ed. II. i. 130.—Afr. austr.
breviseta, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863) 92 = purpurea.
bromoides, *H. B. & K. Nov. Gen. et Sp.* i. 122.—Am. bor. occ.; Ecuador.
bromoides, Salzm. ex Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 118 = longifolia.
caffra, *Bory, Voy.* ii. 376.—Afr. trop.
californica, *Thurb. ex Boland. in Trans. Calif. Agr. Soc.* (1864) 134.—Calif.

ARISTIDA:—
caloptila, Boiss. Fl. Orient. v. 497 = paradisea.
calycina, *R. Br. Prod.* i. 173.—Austral.
canariensis, Willd. Enum. Hort. Berol. 99 = Adscensionis.
capensis, Savi, in Mem. Modcn. (1837) 198 = plumosa.
capensis, *Thunb. Prod. Pl. Cap.* 19.—Afr. austr.
capillacea, Cav. Ic. v. 43. t. 468. f. 1 = Cumingiana.
capillacea, *Lam. Illustr.* i. 156.—Ind. or.; Am. austr.
capillaris, Kunth, Enum. Pl. i. 188, in syn. = Cumingiana.
caudatum, *Anderss. Enum. Pl. Ins. Galap.* 144.—Ins. Galap.
chaetophylla, *Steud. Syn. Pl. Gram.* 420.—Madag.
chapadensis, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 68.—Bras.
chinensis, *Munro, in Proc. Am. Acad.* iv. (1864–65) 363.—China.
Chloris, Salzm. ex Doell, in Mart. Fl. Bras. ii. III. 83 = Gymnopogon foliosus.
chrysopila, *Steud. Nom.* ed. II. i. 131.—Arabia.
ciliata, *Desf. in Schrad. N. Journ.* iii. (1809) 255.—Afr. bor. austr.; Arabia.
ciliata, Steud. & Hochst. ex Steud. Nom. ed. II. i. 131 = hirtigluma.
coarctata, H. B. & K. Nov. Gen. et Sp. i. 122 = setifolia.
coarctata, Lichtst. ex Kunth, Enum. Pl. i. 195 = congesta.
coerulescens, Desf. Fl. Atlant. i. 109 = Adscensionis.
coerulescens, Sieber, ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 88 = Sieberiana.
coerulescens, Hochst. ex Steud. Syn. Pl. Gram. 139 = adoënsis.
cognata, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 127 = purpurascens, stricta.
compacta, *Anderss. Enum. Pl. Ins. Galap.* 145.—Ins. Galap.
complanata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 85.—Bras.
concinna, *Sond. ex J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 140.—Afr. austr.
condensata, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 19.—Am. bor.
congesta, *Roem. & Schult. Syst.* ii. 401.—Afr. austr.
contorta, F. Muell. in Trans. Vict. Inst. i. (1855) 44 = arenaria.
crinita, Steud. Nom. ed. II. i. 131 = Stipa crinita.
crinita, *J. Presl, in Rel. Haenk.* i. 223.—Peruv.
Cumingiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 141.—Ins. Philipp.
curtiseta, A. Gray, in Proc. Acad. Sc. Philad. '1862' (1863) 334 = purpurea.
curvata, *Nees, ex A. Rich. Tent. Fl. Abyss.* ii. 392.—Abyss.; Afr. austr.
curvifolia, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 533.—Mexic.
cyanantha, *Steud. Syn. Pl. Gram.* 141.—Afghan.
decorata, *Steud. l. c.* 421.—Palaestin.
delicatula, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 393.—Abyss.
densispica, *Steud. Syn. Pl. Gram.* 139.—Afr. trop.
depressa, Retz. Obs. iv. 22 = Adscensionis.
depressa, Trin. Gram. Unifl. 175 = setacea.
desmantha, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 109.—Texas.
dichotoma, *Michx. Fl. Bor. Am.* i. 41.—Am. bor.
diffusa, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 68 = vestita.
dispersa, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 129.—Mexic.
divaricata, *Humb. & Bonpl. ex Willd. Enum. Hort. Berol.* 99.—Mexic.
divaricata, Jacq. Eclog. 7. t. 6 = coerulescens.
divaricata, Lag. Elench. 3 = spadicea.
divulsa, *Anderss. Enum. Pl. Ins. Galap.* 143.—Ins. Galap.
domingensis, Kunth, Rev. Gram. i. 62 = Andropogon hirtiflorus.
Dregeana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 169.—Afr. austr.
Ehrenbergii, *Trin. & Rupr. Stipac.* 136.—Arabia.
elatior, Cav. Ic. vi. 65. t. 589 = Adscensionis.
elegans, Rudge, Fl. Guian. 22. t. 30 = capillacea.
Elliotiana, *Steud. Syn. Pl. Gram.* 133.—Carolina.
elliptica, *Kunth, Enum. Pl.* i. 193.—Bras.

ARISTIDA:—
fasciculata, *Torr. in Ann. Lyc. N. York*, i. (1824) 154. —Canada.
Fendleriana, *Steud. Syn. Pl. Gram.* 420.—N. Mexic.
festucoides, *Poir. Encyc. Suppl.* i. 453.—Afr. trop.
filipendula, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863) 93 = purpurea.
flaccida, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 117.—Bras.
flexuosa, *Fourn. Mex. Pl. Enum. Gram.* 77, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 533.—Mexic.
floccosa, Coss. & Dur. in Expl. Sc. Algér. i. 82 = plumosa.
floridana, *Vasey, Gram. U. St.* 21.—Am. bor.
foenicularis, *Edgew. in Journ. As. Soc. Beng.* xxi. (1852) 160, 183.—Ind. or.
Forskaehlei, Tausch, in Flora, xix. (1836) 506 = plumosa.
Freycinetii, Steud. Nom. ed. II. i. 131 = Stipa crinita.
funicularis, *Trin. ex Steud. l. c.*—Nubia.
funiculata, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 159.—Afr. bor.; Arab.; Beluchist.
furcata, Poit. ex Roem. & Schult. Syst. ii. 711 = Bouteloua bromoides.
Gardneriana, Steud. Syn. Pl. Gram. 137 = setifolia.
geminata, Willd. ex Steud. Nom. ed. II. i. 131 = Gymnopogon foliosa.
geminiflora, *Fourn. Mex. Pl. Enum. Gram.* 77, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 533.—Mexic.
geminiflora, Steud. Syn. Pl. Gram. 144 = geminifolia.
geminifolia, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 169.—Afr. austr.
geniculata, *Rafin. Am. Monthly Mag.* (1817), *ex Trin. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 179.—Am. bor.
Geyeriana, Steud. Syn. Pl. Gram. 133 = purpurascens.
gibbosa, *Kunth, Enum. Pl.* i. 189.—Bras.
gigantea, Linn. f. Suppl. 113 = Adscensionis.
glaberrima, *Steud. Syn. Pl. Gram.* 135.—Chili.
glauca, Steud. l. c. = longiramea.
gossypina, *Bosc, ex Beauv. Agrost.* 30, nomen; *Roem. & Schult. Syst.* ii. 391.—Carolina.
gracilis, *Ell. Sketch*, i. 141.—Am. bor.
gracillima, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 173.—Afr. trop.
Grisebachiana, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 533.—Mexic.
guineensis, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 137.—Guinea.
gyrans, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 18.—Am. bor.
Heymanni, *Regel, in Act. Hort. Petrop.* vii. (1880) 649. —As. centr.
hirtigluma, *Steud. Nom.* ed. II. i. 131; *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 171.—Aegypt.; Syria; Arabia.
hordeacea, *Hochst. ex Steud. Syn. Pl. Gram.* 142 = Steudeliana.
hordeacea, *Kunth, Rev. Gram.* ii. t. 173.—Afr. trop.
Humboldtiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 118.—Mexic.
humilis, *H. B. & K. Nov. Gen. et Sp.* i. 121.—Venezuela.
hygrometrica, *R. Br. Prod.* 174.—Austral.
Hystrix, *Linn. f. Suppl.* 113.—Afr. austr.
implexa, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 68. —Bras.
interrupta, *Cav. Ic.* v. 45. t. 471. f. 2.—Mexic.
Jacquiniana, *Tausch, in Flora*, xix. (1836) 508.—Am. austr.
jorullensis, *Kunth, Rev. Gram.* i. 62.—Mexico.
junciformis, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 143.—Afr. austr.
Karwinskiana, *Trin. & Rupr. l. c.* 121.—Mexic.
Kotschyi, Hochst. ex Steud. Syn. Pl. Gram. 142 = funiculata.
Kunthiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1849) 151.—Afr. trop.
laevis, *Kunth, Enum. Pl.* i. 192.—Bras.
Lamarckii, Steud. Nom. ed. I. 69 = amplissima.
lanata, Forsk. Fl. Aegypt. Arab. 25 = plumosa.
lanata, *Poir. Encyc. Suppl.* i. 453.—Am. bor.
lanosa, *Muhl. Descr. Uber.* 174.—Am. bor.
lanuginosa, Bosc, ex Trin. in Mém. Acad. Pétersb. Sér. VI. iv. 11. (1838) 46 = stricta
laxa, *Cav. Ic.* v. 44. t. 470.—Ins. Philipp.; Am. austr.

ARISTIDA:—
laxa, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 85 = flaccida.
laxa, Willd. ex Trin. & Rupr. l. c. vii. (1849) 130 = dispersa.
leiocalycina, *Trin. & Rupr. l. c.* 161.—Guinea.
leptopoda, *Benth. Fl. Austral.* vii. 562.—Austral.
Liebmannii, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 534.—Mexic.
longeradiata, *Steud. Syn. Pl. Gram.* 140.—Afr. trop.
longespica, *Poir. Encyc. Suppl.* i. 452.—Am. bor.
longifolia, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 84.—Bras.
longiramea, *J. Presl, in Rel. Haenk.* i. 224.—Mexic.
longiseta, *Steud. Syn. Pl. Gram.* 420.—N. Mexic.
lutescens, *Steud. l. c.* 145.—Afr. austr.
luzoniensis, *Cav. Ic.* v. 45. t. 470.—Ins. Philipp.
macranthera, Hochst. ex Boiss. Fl. Orient. v. 493 = funiculata.
macrathera, A. Rich. Tent. Fl. Abyss. ii. 393 = praec.
macrochaeta, *Steud. Syn. Pl. Gram.* 134.—Am. bor.
macrochloa, *Hochst. in Flora*, xxxviii. (1855) 200.—Afr. or.
maritima, *Steud. Syn. Pl. Gram.* 137.—Guadel.
mauritiana, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 392.—Abyss.
mauritiana, Kunth, Rev. Gram. i. 265. t. 44 = Adscensionis.
meccana, *Hochst. ex Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 152.—Arabia.
megapotamica, *Spreng. Syst. Cur. Post.* 31.—Bras.
mendocina, *Phil. Sert. Mend. Alt.* 47.—Chili.
micropoda, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 107.—Am. bor.
minuta, Poit. ex Roem. & Schult. Syst. ii. 711 (= *Eutriana Ledebourii*).—Ins. S. Doming.
modatica, *Steud. Syn. Pl. Gram.* 139.—Abyss.
muehlenbergioides, *Fourn. Mex. Pl. Enum. Gram.* 79, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 534.—Mexic.
murina, *Cav. Ic.* v. 44. t. 469.—Reg. Argent.
mutabilis, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 150.—Afr. trop.
namaquensis, *Trin. & Rupr. l. c.* 174.—Afr. austr.
nana, *Steud. Nom.* ed. II. i. 131.—Chili.
Necsiana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 113.—Bras.
nigrescens, *J. Presl, in Rel. Haenk.* i. 223.—Mexic.
nutans, *Ehrenb. & Hempr. ex Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 135.—Arabia.
nutans, Hort. Berol. ex Steud. Nom. ed. II. i. 131 = Adscensionis.
obtusa, *Delile, Fl. Egypte*, 175. t. 13. f. 2.—Afr. bor. et austr.; Arab.
oligantha, *Michx. Fl. Br. Am.* i. 41.—Am. bor.
orizabensis, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 534.—Chili.
pallens, Nutt. Gen. Am. i. 57 = oligantha.
pallens, *Cav. Ic.* v. 43. t. 468.—Chili.
Palmeri, *Vasey, in Torrey Bot. Club*, x. (1883) 42.—Am. bor.
pallida, *Steud. Syn. Pl. Gram.* 143.—Nubia.
palustris, *Vasey, Cat. Gram. U. St.* 35.—Am. bor.
paniculata, *Forsk. Fl. Aegypt. Arab.* 25.—Arabia.
papposa, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 173.—Abyss.
paradisea, *Edgew. in Journ. As. Soc. Beng.* xvi. (1847) 11. 1219.—Persia; Afghan.; Aegypt.; Arab.
paradoxa, J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins. 140 = funiculata.
parviflora, Steud. Syn. Pl. Gram. 140 = vagans.
pauciflora, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863) 92 = oligantha.
pennata, *Trin. in Mém. Acad. Pétersb.* vi. (1815) 488. —Persia; Turkestan.
perennis, *Panz. ex Trin. & Rupr. l. c.* Sér. VI. vii. (1849) 104.—Carolina.
piligens, Burch. ex Spreng. Syst. i. 269 = capensis.
piligera, Burch. ex Roem. & Schult. Syst. ii. Mant. 579 = capensis.
pilosa, *Labill. Sert. N. Caled.* 12. t. 17.—N. Caled.
plumosa, Desf. Fl. Atlant. i. 109 = ciliata.
plumosa, *Linn. Sp. Pl.* ed. II. 1666.—Reg. Mediterr.; Persia; Arab.
pogonoptila, *Boiss. Fl. Orient.* v. 496.—Beluchistan.
prodigiosa, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 80.—Afr. trop.
proxima, *Steud. Syn. Pl. Gram.* 145.—Afr. austr.

ARISTIDA:—
pseudo-hystrix, *Steud. Syn. Pl. Gram.* 142.—Afr. austr.
pumila, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 85 = Adscensionis.
pungens, *Desf. Fl. Atlant.* i. 109. t. 35.—Afr. bor.; Turkestan.; Sibiria.
pungens, Sieber, ex Trin. Gram. Unifl. Diss. i. 181 = vulnerans.
pungens, Steud. Syn. Pl. Gram. 145 = scoparia.
purpurascens, *Poir. Encyc. Suppl.* i. 452.—Am. bor.; Jamaic.
purpurea, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 145.—Am. bor.
pusilla, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 140.—Afr. austr.
quinqueseta, *Steud. Syn. Pl. Gram.* 420.—Maurit.
racemosa, Muhl. Desc. Uber. 172 = purpurascens.
racemosa, *Spreng. Mant.* i. 35.—Ins. S. Helen.
Raddiana, *Savi, in Mem. Moden.* (1837) 198.—Aegypt.
ramosa, *R. Br. Prod.* i. 173.—Austral.
ramosissima, *Engelm. ex A. Gray, Man. Bot. U. St.* ed. V. 618.—Am. bor.
recurvata, *H. B. & K. Nov. Gen. et Sp.* i. 123.—Am. austr.
refracta, *Griseb. Cat. Pl. Cub.* 228.—Cuba.
repens, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 87.—Ins. Galap.
Riedeliana, *Trin. & Rupr. l. c.* Sér. VI. vii. (1849) 113. —Panama.
rigescens, *Roem. & Schult. Syst.* ii. 400.—Ind. or.
rigida, *Cav. Ic.* v. 44. t. 469.—Ins. Philipp.
rigida, Roth, Nov. Pl. Sp. 42 = rigescens.
riparia, *Trin. in Bull. Acad. Pétersb.* i. (1836) 68. —Bras.
Roemeriana, *Scheele, in Linnaea*, xxii. (1849) 343.—Texas.
Royleana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 160.—Ind. or.
rufescens, *Steud. Syn. Pl. Gram.* 421.—Ins. Mascar.
sabulosa, Kunth, Enum. Pl. i. 197 = setifolia.
Sanctae-Luciae, *Trin. Diss.* ii. 25.—Bras.
scabra, *Kunth, Rev. Gram.* i. 62.—Mexico.
Schaffneri, *Fourn. Mex. Pl. Enum. Gram.* 78, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 534.—Mexic.
Schiediana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 120.—Mexic.
Schimperi, Hochst. & Steud ex Steud. Syn. Pl. Gram. 143 = ciliata.
Schweinfurthii, *Boiss. Fl. Orient.* v. 493.—Aegypt.; Arab.
scoparia, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 176.—Aegypt.
secunda, Ledeb. ex Steud. Nom. ed. II. i. 132 (= *Eutriana Ledebourii*).—Ins. S. Doming.
sericea, Ehrenb. ex Boiss. Fl. Orient. v. 497 = paradisea.
setacea, Hort. ex Steud. Nom. ed. II. i. 132 = Adscensionis.
setacea, *Retz. Obs.* iv. 22.—Ind. or.
setacea, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 84 = Adscensionis.
setifolia, *H. B. & K. Nov. Gen. et Sp.* i. 122.—Mexic; Bras.
Sieberiana, *Trin. in Spreng. Neue Entdeck.* ii. 61.—Afr. bor.; Creta; Syria.
similis, *Steud. Syn. Pl. Gram.* 420.—Madag.
simpliciflora, *Chapm. in Coult. Bot. Gaz.* iii. (1878) 18. —Am. bor.
simplicissima, *Steud. Syn. Pl. Gram.* 139.—Ins. Borbon.
sorzogonensis, *J. Presl, in Rel. Haenk.* i. 224.—Ins. Luzon.
spadicea, *H. B. & K. Nov. Gen. et Sp.* i. 123.—Am. austr.
spadicea, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. 11. (1838) 43 = tincta.
spiciformis, *Ell. Sketch.* i. 141.—Am. bor.
spicigera, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 136.—Arab.
squarrosa, *Trin. in Spreng. Neue Entdeck.* ii. 62.—Am. bor.
Steudeliana, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 155.—Abyss.
stipacea, Ehrenb. & Hempr. ex Boiss. Fl. Orient. v. 493 = funiculata.
stipaeformis, Hochst. ex Steud. Syn. Pl. Gram. 143 = pallida.
stipiformis, *Lam. Illustr.* i. 157.—Afr. trop.
stipoides, *R. Br. Prod.* i. 174.—Austral.
stricta, *Michx. Fl. Bor. Am.* i. 41.—Am. bor. et austr.

ARISTIDA :—
stricta, Muhl. Desc. Gram. 174 = spiciformis.
stricta, Auct. ex Steud. Nom. ed. II. i. 132 = virgata.
stricta, Steud. Syn. Pl. Gram. 133 = purpurascens.
subacaulis, *Steud. Nom.* ed. II. i. 132.—Afr. austr.
subaequans, *Doell, in Mart. Fl. Bras.* ii. III. 19.—Bras.
subbiflora, *Steud. Syn. Pl. Gram.* 138.—Ins. Guadeloupe.
subrecurvata, *Beauv. Agrost.* 152, nomen.—Hab. ?
subspicata, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 125.—Ins. Galap.
Swartziana, Steud. Syn. Pl. Gram. 137 = purpurascens.
Teneriffae, *Steud. l. c.* 420.—Ins. Teneriffa.
tenuiflora, *Steud. l. c.* 138.—Afr. trop.
tenuis, *Hochst. in Flora*, xxxviii. (1855) 200.—Abyss.
tenuis, *Kunth, Rev. Gram.* i. 62.—Cumana; Mexic.
ternipes, *Cav. Ic.* v. 46.—Panama.
Thonningii, *Trin. & Rupr. in Mém. Acad. Pétersb.* Sér. VI. vii. (1849) 137.—Afr. trop.
tincta, *Trin. & Rupr. l. c.* 111.—Guiana.
torta, *Kunth, Enum. Pl.* i. 190.—Bras.
trichodes, *Nees, in Hook. Kew Journ.* ii. (1850) 101.—Ins. Philipp.
tripilis, Thunb. Fl. Ceyl. Diss. 2, ex Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 147 = Hystrix.
tuberculosa, *Nutt. Gen. Am.* i. 57.—Am. bor.
unilateralis, Willd. ex Steud. Nom. ed. II. i. 132 = Bouteloua aristioides.
uniplumis, *Lichtst. in Roem. & Schult. Syst.* ii. 401.—Afr. austr.
Urvillei, *Steud. Syn. Pl. Gram.* 141.—Ins. Falkland.
vagans, *Cav. Ic.* v. 45. t. 471.—Austral.
vestita, *Thunb. Prod. Pl. Cap.* 19.—Afr. austr.
virgata, *Trin. in Spreng. Neue Entdeck.* ii. 60.—Am. bor.
Virleti, *Fourn. Mex. Pl. Enum. Gram.* 76, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 535.—Mexic.
vulgaris, Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 131 = Adscensionis.
vulnerans, Trin. & Rupr. l. c. 175 = pungens.
vulpioides, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 251.—China.
Zeyheri, *Steud. Nom.* ed. II. i. 132.—Afr. austr.
Zittelii, *Aschers. in Verh. Bot. Ver. Brand.* xxi. (1880) 70.—Aegypt.

ARISTIDIUM, Lindl. Veg. Kingd. 116 (1847) = **Bouteloua**, Lag. (Gramin.).

ARISTOMENIA, Vell. Fl. Flum. 345 (1825) viii. t. 84 = **Stiffia**, Mikan (Compos.).
fruticosa, Vell. l. c. 346; viii. t. 84 = S. chrysantha.

ARISTOLOCHIA, Tourn. ex Linn. Syst. ed. I. (1735).
ARISTOLOCHIACEAE, Benth. & Hook. f. iii. 123; 1220.
AMBUYA, Rafin. Fl. Tellur. iv. 98 (1836).
DICTYANTHES, Rafin. in Loud. Gard. Mag. viii. (1832) 247.
DIGLOSSELIS, Rafin. Fl. Tellur. iv. 98 (1836).
EINOMEIA, Rafin. Med. Fl. i. 62 (1828).
ENDODECA, Rafin. l. c. (1828).
GLOSSULA, Reichb. Handb. 173 (1837).
GUACO, Liebm. in Forhandl. Skandin. Naturf. 1844 (1847) 203.
HEXAPLECTRIS, Rafin. Fl. Tellur. iv. 97 (1836).
HOCQUARTIA, Dum. Comm. Bot. 30 (1822).
HOWARDIA, Klotzsch, in Monatsb. Acad. Berl. (1859) 607.
ISIPHIA, Rafin. Med. Fl. ii. 232 (1830).
ISOTREMA, Rafin. in Am. Monthly Mag. (1819) 195.
PISTOLOCHIA, Rafin. Fl. Tellur. iv. 98 (1836).
PLAGISTRA, Rafin. l. c. (1836).
PSOPHIZA, Rafin. l. c. 99. (1836).
PTERIPHIS, Rafin. l. c. (1836).
SIPHISIA, Rafin. Med. Fl. i. 62 (1828).
TROPEXA, Rafin. Fl. Tellur. iv. 98 (1836).
abbreviata, Mart. ex Mast. in Mart. Fl. Bras. iv. II. 109 = cymbifera.
abyssinica, Klotzsch, in Monatsb. Akad. Berl. (1859) 598 = bracteata.
acuminata, *Lam. Encyc.* i. 254.—Ind. or.
acuminata, Roxb. Hort. Beng. 64; Fl. Ind. iii. 489 = Roxburghiana.
acutifolia, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 54.—Fl. Amazon.
aethiopica, *Welw. Apont.* 548.—Afr. trop.
albida, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 75.—Afr. occ.
altissima, *Desf. Fl. Atlant.* ii. 324. t. 249.—Reg. Mediterr. or.
anatolica, Boiss. Fl. Orient. iv. 1080 = maurorum.
anguicida, *Jacq. Enum. Pl. Carib.* 30.—Am. austr.
anguicida, Pav. ex Duch. in DC. Prod. xv. I. 443 = velutina.
anguicida, Sieber, ex Duch. l. c. 479 = acuminata.
angulata, *Boj. ex Duch. l. c.* 497.—Madag.
angulisans, Michx. Fl. Bor. Am. ii. 258 = tomentosa.
angulosa, Wall. ex Duch. in DC. Prod. xv. I. 480 = Roxburghiana.
angustifolia, *Boj. Hort. Maurit.* 277.—Ins. Maurit.
angustifolia, *Cham. in Linnaea*, vii. (1832) 211.—Bras.
antihysterica, *Mart. ex Duch. in DC. Prod.* xv. I. 477.—Bras.
appendiculata, Vell. Fl. Flum. ix. t. 98 = macroura.
arborescens, *Linn. Sp. Pl.* 960.—Am.
arcuata, *Mast. in Mart. Fl. Bras.* iv. II. 101.—Am. trop.
arenicola, *Hance, in Journ. Bot.* xiv. (1876) 261.—Cambodia.
argentina, *Griseb. in Goett. Abh.* xix. (1874) 156.—Reg. Argent.
arkasana, *Lodd. Cat.* ed. XIV. (1826) 37.—Am. bor.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 136.—Algeria.
attica, Orphan. ex Duch. in DC. Prod. xv. I. 487 = longa.
Aucherii, Jaub. & Spach, Illustr. i. 174. t. 99 = maurorum.
aurantiaca, *Duch. in DC. Prod.* xv. I. 475.—Venezuela.
auricularia, *Boiss. Diagn.* Ser. I. v. 49.—As. Min.
aurita, *Duch. in DC. Prod.* xv. I. 482.—Madag.
baetica, *Linn. Sp. Pl.* 961.—Europ.
barbata, *Jacq. Coll.* iii. 221.—Venezuela.
Baueri, Duch. in DC. Prod. xv. I. 484 = Thozetii.
Bernieri, *Duch. l. c.* 482.—Madag.
biflora, Willd. ex Duch. l. c. 457 = maxima.
bilabiata, *Linn. Sp. Pl.* ed. II. 1361.—Ins. S. Doming.
Billardieri, *Jaub. & Spach, Illustr.* i. 175. t. 100.—Syria; Cilicia.
bilobata, *Linn. Sp. Pl.* 960.—Ins. S. Doming.
birostris, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 60.—Bras.
Bodamae, *Dingl. in Flora*, xlvi. (1883) 301.—Thracia.
Bonplandi, Tenore, Cat. Sem. Hort. Neap. (1842) 12 = fimbriata.
Bottae, Jaub. & Spach, Illustr. i. 173. t. 98 = maurorum.
brachyura, *Duch. in DC. Prod.* xv. I. 446.—Ins. S. Doming.
bracteata, *Retz. Obs.* v. 29.—Ind. or.
bracteolata, Lam. Encyc. i. 258 = bracteata.
bracteosa, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 37.—Mexic.
Braniciana, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 155, nomen.—Hab. ?
brasiliensis, *Mart. & Zucc. Nov. Gen. et Sp.* i. 77.—Bras.
brevipes, *Benth. Pl. Hartw.* 15.—Mexic.
Bridgesii, *Duch. in DC. Prod.* xv. I. 463.—Chili.
Bruguieri, Jaub. & Spach, Illustr. ii. 40. t. 129 = Olivieri.
Burchellii, *Mast. in Mart. Fl. Bras.* iv. II. 90.—Am. trop.
californica, *Torr. in Pacif. Rail. Rep.* iv. 128.—Calif.
campanaeformis, DC. ex Duch. in DC. Prod. xv. I. 432 = Holostylis reniformis.
caracasana, Spreng. Syst. iii. 753 = trilobata.
caudata, Herb. Berol. ex Duch. in DC. Prod. xv. I. 446 = brachyura.
caudata, Booth, ex Lindl. in Bot. Reg. t. 1453 = macroura.
caudata, *Jacq. Enum. Pl. Carib.* 30.—Ins. S. Doming.
caudata, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 155 (*Contrib.* 48).—Reg. Argent.
Chamissonis, *Duch. in DC. Prod.* xv. I. 462.—Bras.
chilensis, *Miers, Trav. Chil.* ii. 531.—Chili.
chiquitensis, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 49.—Bolivia.
ciliata, Hook. Bot. Mag. t. 3756 = fimbriata.
ciliosa, Benth. in Maund, Bot. t. 90 = fimbriata.
Claussenii, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 57.—Bras.
clavidenia, *Griseb. Cat. Pl. Cub.* 115.—Cuba.
Clematitis, *Linn. Sp. Pl.* 962.—Europ.; Reg. Caucas.; As. Min.
clypeata, *Linden & André, Illustr. Hortic.* xvii. (1870) t. 40.—N. Granat.
conduplicata, Poit. ex Duch. in DC. Prod. xv. I. 460 = oblongata.
conferta, *Mill. Gard. Dict.* ed. VIII. n. 11.—Mexic.
constricta, *Griseb. in Goett. Abh.* vii. (1857) 225.—Reg. Argent.
contorta, *Bunge, Enum. Pl. Chin. Bor.* 58.—China bor.
cordiflora, *Mutis, ex H. B. & K. Nov. Gen. et Sp.* ii. 149.—N. Granat.
cordigera, *Willd. ex Klotzsch, in Monatsb. Akad. Berl.* (1859) 621.—Bras.
coriacea, *Rafin. Atl. Journ.* 146.—Hab. ?
cornuta, *Mast. in Mart. Fl. Bras.* iv. II. 97.—Am. trop.
costaricensis, *Duch. in DC. Prod.* xv. I. 450.—Costa Rica.
crenata, Ehrenb. ex Duch. l. c. 478 = bracteata.
cretica, *Lam. Encyc.* i. 225.—Creta.
cubensis, *Linden, Cat.* n. 11. *Suppl.* (1856) 12.—Cuba.
cucullata, Pohl, ex Duch. in DC. Prod. xv. I. 432 = Holostylis reniformis.
cyclochilia, Duch. in Ann. Sc. Nat. Sér. IV. ii. (1854) 45 = passifloraefolia.
cymbifera, *Mart. & Zucc. Nov. Gen. et Sp.* i. 76. t. 49.—Bras.
cynanchifolia, *Mart. & Zucc. l. c.* 77. t. 51.—Bras.
debilis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 197.—Japon.
deltantha, *F. Muell. Fragm.* vi. 179.—Austral.
deltoidea, *H. B. & K. Nov. Gen. et Sp.* ii. 146. t. 112.—Bras.
dictyantha, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 40.—Venezuela.
Duchartrei, André, Le Mouv. Hortic. (1867) 61; Mast. in Gard. Chron. (1868) 516 = Ruiziana.
durior, *Hill, Veg. Syst.* xxi. 57.—Am. bor.
ecaudata, DC. ex Duch. in DC. Prod. xv. I. 446 = dictyantha.
egensis, Poepp. ex Duch. l. c. 465 = acutifolia.
Ehrenbergiana, *Cham. in Linnaea*, vii. (1832) 210.—Ins. S. Doming.
elegans, *Mast. in Gard. Chron.* (1885) II. 301.—Bras.
elliptica, Duch. in Ann. Sc. Nat. Sér. IV. ii. (1854) 51 = tigrina.
emarginata, *Duch. in DC. Prod.* xv. I. 467.—Bras.
emarginata, Willd. ex Duch. l. c. 466 = nummulariifolia.
erecta, *Linn. Sp. Pl.* ed. II. 1362.—Vera Crux.
eriantha, *Mart. & Zucc. Nov. Gen. et Sp.* i. 78. t. 53.—Bras.
Eschscholtzii, Ledeb. ex Duch. in DC. Prod. xv. I. 481 = A. Tagala.
eurystoma, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 41.—Bras.
fasciculata, Lam. Fl. Fr. iii. 387 = A. Pistolochia.
filipendula, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 70.—Bras.
fimbriata, *Cham. in Linnaea*, vii. (1832) 210. t. 6.—Bras.
flexuosa, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 36. Mexic.
floribunda, *Lem. Illustr. Hortic.* (1868) t. 568.—Bras.
fluminensis, Buek, in DC. Prod. Index, iv. 28 = odora.
foetens, *Lindl. Bot. Reg.* t. 1824.—Ind. occ.
foetida, *H. B. & K. Nov. Gen. et Sp.* ii. 147. t. 114.—Mexic.
Fontanesii, Boiss. & Reut. Pugill. Pl. Hispan. 108 = longa.
Fordiana, *Hemsl. in Journ. Bot.* xxiii. (1885) 286.—China.
fragrantissima, *Ruiz, in Mem. Bejuc. Estrell.* (1805) 46.—Peruv.
frutescens, Marsh. Arb. Am. 24 = macrophylla.
galeata, *Mart. & Zucc. Nov. Gen. et Sp.* i. 76. t. 50.—Bras.
galeata, Moritzi, Syst. Verz. 70 = cymbifera.
Galcotti, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 44.—Mexic.

ARISTOLOCHIA :—
Gardneri, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 69.—Bras.
Gaudichaudii, *Duch. l. c.* 72.—Ins. Rawak.
geminiflora, H. B. & K. Nov. Gen. et Sp. ii. 148. t. 117 = maxima.
gibbosa, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 53. —Mexic.
Giberti, *Hook. Bot. Mag.* t. 5345.—Reg. Argent.
gigantea, Hook. l. c. 4221 = grandiflora.
gigantea, *Mart. & Zucc. Nov. Gen. et Sp.* i. 75. t. 48. —Bras.
Gigas, Lindl. in Bot. Reg. (1842) Misc. 51. t. 60 = grandiflora.
glandulosa, *Kickx, in Bull. Acad. Brux.* déc. vii. (1839) n. 11.—Cuba.
glauca, Desf. Fl. Atlant. ii. t. 250 = baetica.
glauca, Salisb. Prod. 215 = subglauca.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* ii. 147. t. 115.—N. Granat.
Glaziovii, *Mast. in Mart. Fl. Bras.* iv. II. 90.—Am. trop.
Goldieana, *Hook. f. in Trans. Linn. Soc.* xxv. (1865) 185. t. 14.—Afr. trop.
Goudotii, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 66.—N. Granat.
gracilis, *Duch. in DC. Prod.* xv. I. 460.—Am. austr.
grandiflora, *Arruda, in Koster, Trav. Braz.* 499.—Pernamb.
grandiflora, Gomez, Obs. Bot. Med. ii. (1803) 14. t. 3; et in Mem. Ac. Lisboa, iii. (1812) Mem. dos Corresp. 64 = cymbifera.
grandiflora, *Sw. Prod. Veg. Ind. Occ.* 126.—Jamaica.
grandifolia, Salisb. Prod. 215 = macrophylla.
Griffithii, *Hook. f. & Thoms. ex Duch. in DC. Prod.* xv. I. 437.—Reg. Himal.
hastata, H. B. & K. Nov. Gen. et Sp. ii. 148 = pentandra.
hastata, *Jack, in Malay. Misc.* ii. (1822) n. VII. 6.—Malaya.
hastata, Klotzsch, in Monatsb. Acad. Berl. (1859) 597 = Zollingeri.
hastata, Nutt. Gen. Am. ii. 200 = A. Serpentaria.
hians, *Willd. in Mém. Soc. Nat. Mosc.* ii. (1809) 100. —Venezuela.
Hilariana, *Duch. in DC. Prod.* xv. I. 451.—Bras.
hirsuta, Linn. Syst. ed. XII. 601 = hirta.
hirsuta, Muhl. Desc. Uber. 18 = tomentosa.
hirta, Buek, in DC. Prod. Index, iv. 28 = torta.
hirta, Georgi, Beschreib. Russ. Reich. iii. v. 1275 = pontica.
hirta, *Linn. Sp. Pl.* 961.—As. Min.
hispida, *Pohl, ex Duch. in DC. Prod.* xv. I. 498.—Bras.
iberica, *Fisch. & Mey. ex Duch. in DC. Prod.* xv. I. 493.—Reg. Caucas.
imbricata, *Mast. in Journ. Linn. Soc.* xiv. (1875) 494. —Ins. Philipp.
incisa, *Duch. in DC. Prod.* xv. I. 490.—Anatolia.
indica, *Linn. Sp. Pl.* 960.—Ind. or.
infesta, Salisb. Prod. 215 = A. Clematitis.
inflata, *H. B. & K. Nov. Gen. et Sp.* ii. 145. t. 111.—N. Granat.
insignis, *Versch. ex Gard. Chron.* (1868) 349.—Bras.
Ipemi, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 153 (*Contrib.* 46).—Paraguay.
Jackii, Steud. Nom. ed. II. i. 141 = hastata.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 136.—Japon.
Kaempferi, *Willd. Sp. Pl.* iv. 152.—Japon.
Karwinskii, *Duch. in DC. Prod.* xv. I. 442.—Mexic.
Kotschyi, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 237 = bracteata.
labiata, Willd. in Mém. Soc. Nat. Mosc. ii. (1809) 101. t. 6 = brasiliensis.
labiosa, Ker-Gawl. in Bot. Reg. (1822) t. 689 = cymbifera.
lanceolata, Wight, Ic. v. t. 1858 = indica.
Leprieurii, *Duch. in DC. Prod.* xv. I. 451.—Guiana.
leuconeura, *Linden, in Belg. Hortic.* viii. (1858) 165. —Am. austr.
Lindeniana, *Duch. in DC. Prod.* xv. I. 453.—Cuba.
lincarifolia, *Wright, ex Griseb. Cat. Pl. Cub.* 115.—Cuba.
lineata, *Duch. in Rev. Hortic.* Sér. IV. iii. (1854) 284. t. 15.—Japon.
littoralis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 155 (*Contrib.* 47).—Reg. Argent.

ARISTOLOCHIA :—
longa, Georgi, Beschr. Russ. Reich. iii. v. 1274 = A. Clematitis.
longa, *Linn. Sp. Pl.* 961.—Reg. Mediterr.
longa, Thunb. Fl. Japon. 144 = debilis.
longa, Woodv. Med. Bot. ii. t. 107 = baetica.
longiflora, *Engelm. & A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 259.—Texas.
longifolia, *Champ. ex Benth. in Hook. Kew Journ.* vi. (1854) 116.—Hongkong.
longifolia, Roxb. Hort. Beng. [102]; Fl. Ind. iii. 490 = moluccana.
lutea, Desf. Choix, 13 t. 8 = pallida.
lutescens, *Duch. in DC. Prod.* xv. I. 461.—Bras.
macradenia, Hook. Bot. Mag. t. 4467 = glandulosa.
macrocarpa, *Duch. in DC. Prod.* xv. I. 497.—Afr. occ.
macroglossa, Jaub. & Spach, Illustr. ii. 38. t. 127 = Tournefortii.
macrophylla, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 68.—Guiana.
macrophylla, Lam. Encyc. i. 252 = A. Sipho.
macrota, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 43. —Guiana.
macroura, *Gomez, in Mem. Ac. Lisboa*, iii. (1812) *Mem. dos Corresp.* 77.—Bras.
Mannii, *Hook. f. in Trans. Linn. Soc.* xxv. (1865) 186. —Afr. trop.
marsupiiflora, *Schrad. in Goett. Gel. Anz.* i. (1821) 719.—Hab.?
Mathewsii, *Duch. in DC. Prod.* xv. I. 497.—Peruv.
mauritiana, Pers. Syn. ii. 527 = bracteata.
maurorum, Klotzsch, in Monatsb. Acad. Berl. (1859) 598 = bracteata.
maurorum, *Linn. Sp. Pl.* ed. II. 1363.—As. Min.; Persia.
maxima, Cham. in Linnaea, vii. (1832) 208 = Chamissonis.
maxima, *Jacq. Enum. Pl. Carib.* 30.—Ind. occ.
maysorensis, Fisch. ex Duch. in DC. Prod. xv. I. 479 = indica.
melastoma, *Manso, ex Duch. l. c.* 461.—Bras.
mexicana, *Kostel. Allg. Med.-Pharm. Fl.* ii. 467.—Mexic.
mexicana, A. Dietr. Syn. v. 196 = maxima.
micrantha, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 35. t. 5.—Mexic.
microphylla, *Willd. Enum. Hort. Berol. Suppl.* 68.—Hab.?
microstoma, *Boiss. & Sprun. Diagn.* Ser. I. v. 50.—Graecia.
microura, Pohl, ex Duch. in DC. Prod. xv. I. 496 = setosa.
mollissima, *Hance, in Journ. Bot.* xvii. (1879) 300.—China.
moluccana, *Duch. in DC. Prod.* xv. I. 438.—Ins. Molucc.
moschata, Wedd. ex Duch. l. c. 475 = odoratissima.
multiflora, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 73.—Ins. Comoro.
multinervis, *Pomel, Nouv. Mat. Fl. Atl.* 137.—Algeria.
nervosa, *Duch. in DC. Prod.* xv. I. 480.—Ind. or.
nummularifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 145.—Venezuela.
oblonga, Vell. Fl. Flum. ix. t. 99 = rumicifolia.
oblongata, *Jacq. Hort. Schoenb.* ii. 29.—Hab.?
obtusata, *Sw. Prod. Veg. Ind. Occ.* 126.—Ind. occ.
odora, *Steud. Nom.* ed. II. i. 133.—Bras.
odoratissima, Benth. Pl. Hartw. 82 = gibbosa.
odoratissima, *Linn. Sp. Pl.* ed. II. 1362.—Jamaica.
odoratissima, Vell. Fl. Flum. ix. t. 97 = odora.
officinalis, T. Nees, Pl. Offic. t. 144 = A. Serpentaria.
Olivieri, *Colleg. ex Boiss. Diagn.* Ser. I. v. 50.—Mesopot.
orbicularis, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 59.—Bras.
orbiculata, Vell. Fl. Flum. ix. t. 96 = cymbifera.
ornithocephala, Hook. Bot. Mag. t. 4120 = brasiliensis.
Ottonis, Klotzsch, ex Duch. in DC. Prod. xv. I. 476 = pandurata.
ovalifolia, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 50.—Mexic.
paecilantha, *Boiss. Diagn.* Ser. I. xii. 104.—Syria.
pallida, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 656 = longa.
pallida, *Willd. Sp. Pl.* iv. 162.—Europ.; As. Min.
pandurata, *Jacq. Hort. Schoenb.* iv. 49. t. 407.—Venezuela.

ARISTOLOCHIA :—
pandurata, Wall. ex Duch. in DC. Prod. xv. I. 479 = indica.
panduriformis, Willd. Sp. Pl. iv. 152 = pandurata.
pannosa, *Mast. in Journ. Linn. Soc.* xxiv. (1875) 493. —Peruv.
papillaris, *Mast. in Mart. Fl. Bras.* iv. II. 100.—Am. trop.
paraguariensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 156 (*Contrib.* 48).—Paraguay.
paramaribensis, *Duch. in DC. Prod.* xv. I. 496.—Bras.
pardina, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 47. —Mexic.
parviflora, *Griseb. in Goett. Abh.* xxiv. (1879) 148.—Reg. Argent.
parvifolia, *Sibth. & Sm. Fl. Graec. Prod.* ii. 222.—Syria; Cyprus; Graecia?
parvifolia, Sprun. ex Duch. in DC. Prod. xv. I. 485 = microstoma.
passifloraefolia, *A. Rich. Fl. Cub. Fanerog.* ii. 195.—Cuba.
paucinervis, *Pomel, Nouv. Mat. Fl. Atl.* 136.—Algeria.
Pavoniana, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 54.—Mexic.
Pearcei, *Mast. in Journ. Linn. Soc.* xiv. (1875) 493.—Peruv.
Pearcei, *Phil. in Ann. Univ. Chil.* i. (1861) 68; *Linnaea*, xxxiii. (1865) 231.—Chili.
peltata, *Linn. Sp. Pl.* 960.—Ins. S. Doming.
peltata, Sw. Obs. 341 = reniformis.
pentandra, *Jacq. Enum. Pl. Carib.* 30.—Cuba.
Petersiana, *Klotzsch, in Monatsb. Akad. Berl.* (1859) 599.—Afr. trop.
picta, Karst. Ausw. Schoen. Pfl. Venez. ii. 34 = pandurata.
pilosa, *H. B. & K. Nov. Gen. et Sp.* ii. 146. t. 113.—Ecuador.
piperifolia, *Griff. Itin. Bot.* 19.—Reg. Himal.
Pistolochia, Aucher, ex Duch. in DC. Prod. xv. I. 491 = incisa.
Pistolochia, *Linn. Sp. Pl.* ed. II. 962.—Reg. Mediterr. occ.
platanifolia, *Duch. in DC. Prod.* xv. I. 437.—Mexic.
platyloba, *Garcke, in Linnaea*, xxii. (1849) 69.—Guiana.
podocarpa, *A. Bertol. in Nov. Comm. Acad. Sc. Bon.* iv. (1840) 437.—Guatem.
Pohliana, *Duch. in DC. Prod.* xv. I. 496.—Bras.
polyrrhizos, *Spreng. Syst.* iv. *Cur. Post.* 310.—Am. bor.
pontica, *Lam. Encyc.* i. 255.—As. Min.
praevenosa, *F. Muell. Fragm.* ii. 166.—Austral.
promissa, *Mast. in Gard. Chron.* (1879) I. 494.—Afr. trop.
prostrata, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 63.—Bolivia.
pubera, F. Baner, ex Duch. in DC. Prod. xv. I. 484 = Thozetii.
pubera, *R. Br. Prod.* 349.—Austral.
puberula, F. Muell. ex Duch. in DC. Prod. xv. I. 484 = pubera.
pubescens, Hort. ex C. Koch, Dendrol. ii. I. 395 = tomentosa.
pubescens, *Page, ex Steud. Nom.* ed. II. i. 133.—Am. bor.
pubescens, *Willd. ex Duch. in DC. Prod.* xv. I. 450.—Bras.
punctata, Balbis, ex Duch. l. c. 446 = brachyura.
punctata, *Lam. Encyc.* i. 253.—Ins. S. Doming.
pusilla, Pohl, ex Duch. in DC. Prod. xv. I. 466 = Clausseni.
quadriflora, *Gueldenst. Reisen Russl.* i. 421.—Reg. Caucas.
Raja, *Mart. & Zucc. Nov. Gen. et Sp.* i. 75.—Bras.
recurvilabra, *Hance, in Journ. Bot.* xi. (1873) 75.—China.
refracta, Burch. ex Mast. in Mart. Fl. Bras. iv. II. 93 = Chamissonis.
reniformis, Vell. Fl. Flum. ix. t. 100 = A. Raja.
reniformis, *Willd. Sp. Pl.* iv. 153.—Ins. S. Doming.
repens, *Mill. Gard. Dict.* ed. VIII. n. 12.—Mexic.
reticulata, Holton, ex Duch. in DC. Prod. xv. I. 477 = maxima.
reticulata, *Nutt. in Trans. Am. Phil. Soc.* N.S. v. (1837) 162.—Am. bor.
reticulata, Seem. Bot. Voy. Herald, 193 = Mathewsii.
rigida, *Duch. in DC. Prod.* xv. I. 495.—Afr. austr.
ringens, Link & Otto, Ic. Pl. Sel. i. 33. t. 14 = brasiliensis.

ARISTOLOCHIA :—
ringens, *Vahl, Symb. Bot.* iii. 99.—Ind. or.
rotunda, Bové, ex Duch. in DC. Prod. xv. i. 487 = longa.
rotunda, Georgi, Beschr. Russ. Reich. iii. v. 1274 = A. Clematitis.
rotunda, *Linn. Sp. Pl.* 962.—Reg. Mediterr.
rotunda, Moç. & Sesse, ex Duch. in DC. Prod. xv. i. 442 = bracteosa.
Roxburghiana, *Klotzsch, in Monatsb. Akad. Berl.* (1859) 596.—Ind. or.; Malaya.
rubromarginata, Fisch. ex Duch. in DC. Prod. xv. i. 451 = Leprieurii.
rufomarginata, Leprieur, ex Duch. l. c. = Leprieurii.
rugosa, Lam. Encyc. i. 252 = obtusata.
Ruiziana, *Duch. in DC. Prod.* xv. i. 476.—Peruv.
rumicifolia, *Mart. & Zucc. Nov. Gen. et Sp.* i. 79.—Bras.
rumicifolia, R. Schomb. ex Duch. in DC. Prod. xv. i. 447 = dictyantha.
Rumphii, *Kostel. Allg. Med.-Pharm. Fl.* ii. 465.—Ins. Banda.
saccata, Hook. f. & Thoms. ex Klotzsch, in Monatsb. Akad. Berl. (1859) 603 = seq.
saccata, *Wall. Pl. As. Rar.* ii. 2. t. 103.—Reg. Himal.
sagittata, *Muhl. ex Duch. in DC. Prod.* xv. i. 434 = A. Serpentaria.
scabrida, Boiss. ex Duch. in DC. Prod. xv. i. 494 = hirta.
scabridula, Boiss. Diagn. Ser. I. xii. 105 = paecilantha.
scandens, *Mill. Gard. Dict.* ed. VIII. n. 10.—Hispan.
Sellowiana, Duch. in DC. Prod. xv. i. 458 = triangularis.
sempervirens, *Forsk. Fl. Aegypt. Arab.* 156.—Arabia.
sempervirens, *Linn. Sp. Pl.* 961.—Creta.
sempervirens, Orph. ex Nym. Consp. 645 = altissima.
sepicola, *Mast. in Mart. Fl. Bras.* iv. ii. 100.—Am. trop.
sericea, *Benth. Pl. Hartw.* 81.—Mexic.
sericea, *Blanco, Fl. Filip.* 283.—Ins. Philipp.
Serpentaria, *Linn. Sp. Pl.* 961.—Am. bor.
sessiflora, DC. ex Duch. in DC. Prod. xv. i. 480 = nervosa.
sessilifolia, *Duch. l. c.* 464.—Bras.
setigera, Klotzsch, in Monatsb. Acad. Berl. (1859) 617 = chilensis.
setosa, *Duch. in DC. Prod.* xv. i. 496.—Bras.
sicula, *Tineo, in Guss. Fl. Sic. Syn.* ii. 878.—Sicilia.
sinarum, *Lindl. in Gard. Chron.* (1859) 708.—China.
Sipho, *L'Hérit. Stirp. Nov.* 13.—Am. bor.
smilacina, *Duch. in DC. Prod.* xv. i. 459.—Bras.
somaliensis, *Oliver, in Hook. Ic. Pl.* t. 1273.—Afr. trop.
Soyauxiana, *Oliver, l. c.* t. 1410.—Afr. trop.
spathulata, *Duch. in DC. Prod.* xv. i. 448.—Cuba.
Sprucei, *Mast. in Mart. Fl. Bras.* iv. ii. 88.—Am. trop.
strictiflora, Duch. in DC. Prod. xv. i. 484 = pubera.
subclausa, *S. Wats. in Proc. Am. Acad.* xx. (1885) 372.—Mexic.
subcordata, Willd. ex Duch. in DC. Prod. xv. i. 468 = deltoidea.
subglauca, Lam. Encyc. i. 257; Brot. Fl. Lusit. ii. 593 = baetica.
subsagittata, Blanco, Fl. Filip. ed. I. 283 = A. Tagala.
surinamensis, Miq. in Linnaea, xviii. (1844) 246 = macrota.
surinamensis, *Willd. Sp. Pl.* iv. 151.—Guiana.
Tagala, *Cham. in Linnaea*, vii. (1832) 207.—Ins. Luzon.
taliscana, *Hook. & Arn. Bot. Beech. Voy.* 309.—Mexic.
tamnifolia, *Duch. in DC. Prod.* xv. i. 448.—Bras.
tapetotricha, Lem. Illustr. Hortic. iii. (1856) Misc. 22 = N. macroura.
tenera, *Pohl, ex Duch. in DC. Prod.* xv. i. 466.—Granat.
theriaca, *Mart. ex Duch. l. c.* 455.—Bras.
Thozetii, *F. Muell. Fragm.* ii. 167.—Austral.
Thwaitesii, *Hook. Bot. Mag.* t. 4918.—China.
tigrina, *A. Rich. Fl. Cub. Fanerog.* ii. 194.—Cuba.
timorensis, *Decne. Herb. Tim. Descr.* 41.—Ins. Timor.
tomentosa, *Sims, Bot. Mag.* t. 1369.—Am. bor.
torta, Willd. ex Duch. in DC. Prod. xv. i. 458 = inflata.
Tournefortii, *Jaub. & Spach, Illustr.* ii. 38. t. 128.—As. Min.; Ins. Chios.
triactina, *Hook. f. in Trans. Linn. Soc.* xxv. (1865) 186.—Afr. trop.

ARISTOLOCHIA :—
Trianaei, *Duch. in DC. Prod.* xv. i. 462.—N. Granat.
triangularis, *Cham. in Linnaea*, vii. (1832) 209. t. 6.—Bras.
tricaudata, *Lem. Illustr. Hortic.* xii. (1865) *Misc.* 49; xiii. (1866) t. 523.—Mexic.
trichostoma, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 190.—Cuba.
trifida, Lam. Encyc. i. 251 = trilobata.
triloba, Salisb. Prod. 214 = trilobata.
trilobata, Lam. Encyc. i. 251 = surinamensis.
trilobata, Lindl. Bot. Reg. t. 1399 = macroura.
trilobata, *Linn. Sp. Pl.* 960.—Ind. occ.
tripteris, *Rafin. Fl. Ludov.* 24.—Am. bor.
trulliformis, *Mast. in Mart. Fl. Bras.* iv. ii. 101.—Am. trop.
truncata, *Field. & Gardn. Sert. Pl.* t. 44.—Peruv.
turbacensis, *H. B. & K. Nov. Gen. et Sp.* ii. 149.—Am. bor.
Uhdeana, *Duch. in DC. Prod.* xv. i. 465.—Mexic.
undata, Moench, Meth. 719 = sempervirens.
ungulifolia, *Mast. in Journ. Linn. Soc.* xiv. (1875) 494.—Borneo.
utriculata, Seem. Bot. Voy. Herald, 193 = Mathewsii.
valentina, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 39.—Mexic.
variifolia, *Duch. l. c.* 65.—Mexic.
velutina, *Duch. l. c.* 39.—Mexic.
veraguensis, *Duch. in DC. Prod.* xv. i. 458.—Am. centr.
Wageneriana, *Schlecht. in Linnaea*, xxvi. (1853) 631.—N. Granat.
Warmingii, *Mast. in Mart. Fl. Bras.* iv. ii. 109.—Am. trop.
Weddellii, *Duch. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 62.—Bras.
Westlandi, *Hemsl. in Journ. Bot.* xxiii. (1885) 286.—China.
Wrightii, Seem. Bot. Voy. Herald, 331. t. 72 = brevipes.
Zollingeri, *Miq. Fl. Ind. Bat.* i. 1066.—Java.

ARISTOTELA, Adans. Fam. ii. 125 (1763) = **Othonna**, Linn. (Compos.).

ARISTOTELA, J. F. Gmel. Syst. 751 (1791) = **Aristotelia**, L'Hérit. (Tiliac.).

ARISTOTELEA, Lour. Fl. Cochinch. 522 (1790) = **Spiranthes**, Rich. (Orchid.).
spiralis, Lour. l. c. = S. australis.

ARISTOTELEA, Spreng. Syst. ii. 450 (1825) = **Aristotelia**, L'Hérit. (Tiliac.).

ARISTOTELIA, L'Hérit. Stirp. Nov. 31. t. 16 (1784). *TILIACEAE*, Benth. & Hook. f. i. 239.
BEAUMARIA, Deless. ex Steud. Nom. ed. II. i. 192 (1840).
FRIESIA, DC. Prod. i. 520 (1824).
australasica, *F. Muell. Fragm.* ii. 79.—Austral.
Braithwaitei, *F. Muell. in Wing, South. Sc. Record*, i. (1881) 149.—N. Hebrid.
Colensoi, *Hook. f. Handb. New Zeal. Fl.* 33.—N. Zel.
erecta, *J. Buch. in Trans. N. Z. Inst.* iii. (1871) 209.—N. Zel.
fruticosa, *Hook. f. Fl. N. Zeal.* i. 34.—N. Zel.
glabra, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. ii. (1868) 44; *et Contrib. Bot.* ii. 180.—Am. merid.
glandulosa, Ruiz & Pav. Syst Veg. 125 = A. Macqui.
lucida, Salisb. Prod. 345 = A. Macqui.
Macqui, *L'Hérit. Stirp. Nov.* 31. t. 16.—Chili.
megalosperma, *F. Muell. Fragm.* ix. 84.—Austral.
peduncularis, *Hook. f. Fl. Tasm.* i. 52.—Tasmania.
racemosa, *Hook. f. Fl. N. Zeal.* i. 33.—N. Zel.

ARIVELA, Rafin. Sylva Tellur. 110 (1838) = **Cleome**, Linn. (Capparid.).
viscosa, Rafin. l. c. = C. viscosa.

ARIVONA, Steud. Nom. ed. II. i. 130 (1840) = **Arjona**, Comm. (Santal.).

ARJONA, Comm. ex Cav. Ic. iv. 57. t. 383 (1797). *SANTALACEAE*, Benth. & Hook. f. iii. 220.
ARIVONA, Steud. Nom. ed. II. i. 130 (1840).
ARJOONA, Endl. Gen. 325 (1838).
andina, *Phil. in Anal. Univ. Chil.* ii. (1862) 405; *et in Linnaea*, xxxiii. (1864–65) 232.—Chili.

ARJONA :—
appressa, *Phil. in Linnaea*, xxxiii. (1864–65) 233.—Chili.
linearis, *Miers, in Journ. Linn. Soc.* xvii. (1880) 133.—Reg. Argent.
longifolia, *Phil. in Anal. Univ. Chil.* ii. (1862) 405.—Chili.
minima, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 30.—Chili.
patagonica, *Homb. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél.* 22.—Patagon.
pungens, *Phil. in Anal. Univ. Chil.* (1873) 537.—Chili.
pnsilla, *Hook. f. Fl. Antarct.* ii. 342.—Patagon.
rigida, *Miers, in Journ. Linn. Soc.* xvii. (1880) 132.—Reg. Argent.
ruscifolia, *Poepp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) (*Pl. Meyen.*) 112.—Chili.
tuberosa, *Cav. Ic.* iv. 57. t. 383.—Am. merid.
tuberosa, Phil. in Linnaea, xxxiii. (1864–65) 231 = rigida.

ARKEZOSTIS, Rafin. New Fl. Am. iv. 100 (1836) = **Cayaponia**, Manso (Cucurb.).
quinqueloba, Rafin. l. c. = C. Boykinii.

ARKOPODA, Rafin. Fl. Tellur. iii. 73 (1836) = **Reseda**, Tourn.
Luteola, Rafin. l. c. = R. Luteola.

ARMANIA, Bert. ex DC. Prod. v. 576 (1836) = **Encelia**, Adans. (Compos.).
fruticulosa, Bert. l. c.—Am. austr.

ARMENIACA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Prunus**, Tourn. (Rosac.).
amarella, Schur, Enum. Pl. Transs. 178 = P. amarella.
atropurpurea, Loisel. in Duham. ed. Nov. v. 172. t. 552 = P. dasycarpa.
berricoccia, Delarb. Fl. Auv. ed. II. 326 = P. Armeniaca.
brigantiaca, Pers. Syn. ii. 36 = P. brigantiaca.
dasycarpa, Pers. l. c. = P. dasycarpa.
Davidiana, Carr. in Rev. Hortic. (1879) 237.—China.
duracina, Dierb. ex Steud. Nom. ed. II. i. 133 = P. amarella.
epirotica, Gaertn. Mey. & Scherb. Fl. Wett. ii. 167 = P. Armeniaca.
fusca, Turp. & Poit. in Nouv. Duham. Arb. Fruit. i. 123. t. 60 = P. dasycarpa.
longifolia, Desc. Fl. Antill. ii. 108 = P. occidentalis.
lucida, May, in Rev. Hortic. (1875) 179 = P. lucida.
Mume, De Vriese, Tuinb. Fl. i. (1855) 1. t. 1 = P. Mume.
nigra, Pers. Syn. ii. 36 = P. dasycarpa.
pedunculata, Hort. ex Loud. Hort. Brit. Suppl. ii. 610 = P. pedunculata.
persicifolia, Loisel. in Duham. ed. Nov. v. 172. t. 552 = P. dasycarpa.
sibirica, Pers. Syn. ii. 36 = P. sibirica.
verrucosa, May, in Rev. Hortic. (1875) 299.—As. Min.
trichocarpa, M. Roem. Syn. Rosifl. 34 = P. tomentosa.
vulgaris, Lam. Encyc. i. 2 = P. Armeniaca.

ARMENIASTRUM, Lem. Jard. Fleur. iv. (1854) Misc. 76 = **Espadaea**, A. Rich. (Verben.).
apiculatum, Lem. l. c. 78 = E. apiculata.

ARMERIA, Linn. Syst. ed. I. (1735); Willd. Enum. Hort. Berol. 333 (1809). *PLUMBAGINEAE*, Benth. & Hook. f. ii. 626.
POLYANTHEMUM, Medic. in Staatsw. Vorles. Churpf. Phys. Oek. Ges. i. (1791) 228.
alliacea, Ebel & Hoffmgg. in Hoffmgg. & Link, Fl. Port. i. 441 = plantaginea.
alliacea, Griseb. Spicil. Fl. Rumel. ii. 296 = cariensis.
alliacea, Mutel, Fl. Fr. iii. 86. f. 405 = plantaginea.
alliacea, *Roem. & Schult. Syst.* vi. 773.—Reg. Mediterr. occ.
alliacea, Schousb. Vext. Marok. 146 = baetica.
alliacea, Webb, ex Ball, in Journ. Linn. Soc. xvi. (1878) 560 = mauritanica.
allioides, *Boiss. Voy. Bot. Espagne*, ii. 524.—Hispan.
allioides, Willk. ex Willk. & Lange, Prod. Fl. Hisp. ii. 366 = longearistata.
alpina, Ebel, Armeriae Gen. Diss. (1840) 28 = Halleri.
alpina, Frivald. ex Griseb. Spicil. Fl. Rumel. ii. 296 = cariensis.

ARMERIA :—
alpina, Guss. Fl. Sic. Prod. 378 = nebrodensis.
alpina, Tenore, Syll. 160, ex Boiss. in DC. Prod. xii. 685 = majellensis.
alpina, Turcz. in Bull. Soc. Nat. Mosc. (1838) 99; xxv. (1852) II. 399 = sibirica.
alpina, *Willd. Enum. Hort. Berol.* 333.—Europ.
andicola, C. Gay, ex Boiss. in DC. Prod. xii. 682 = andina.
andina, *Poepp. ex Boiss. l. c.*—Chili.
androsacea, Boiss. l. c. 679 = vulgaris.
arctica, *Wallr. Beitr.* i. 193.—Unalashka.
arcuata, Welw. ex Boiss. & Reut. Pugill. Pl. Nov. 101 = boetica.
arenaria, Schult. in Roem. & Schult. Syst. vi. 771 = maritima.
argyrocephala, *Wallr. Beitr.* i. 206.—Europ.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 134.—Algeria.
australis, *Boiss. Voy. Bot. Espagne*, ii. 526.—Hispan.
berlengensis, *Daveau, in Bol. Soc. Broter.* ii. (1884) 24.—Lusitan.
boetica, *Boiss. Voy. Bot. Espagne*, ii. 749.—Hispan.
Boissieriana, *Coss. Notes Crit.* 44.—Hispan.
Bourgaei, Boiss. ex Nym. Consp. 615 = Duriaei.
brachyphylla, *Boiss. in DC. Prod.* xii. 682.—Chili.
brevifolia, Kunze, ex Boiss. l. c. = chilensis.
bupleuroides, Gren. & Godr. Fl. Fr. ii. 736 = plantaginea.
caespitosa, *Boiss. in DC. Prod.* xii. 679.—Hispan.; Lusitan.
campestris, Wallr. Beitr. i. 203 = maritima.
canescens, *Boiss. in DC. Prod.* xii. 686.—Dalmatia.
cantabrica, *Boiss. & Reut. ex Willk. & Lange, Prod. Fl. Hisp.* ii. 366.—Hispan.
cariensis, *Boiss. in DC. Prod.* xii. 677.—As. Min.
Cephalotes, Hook. Bot. Mag. t. 4128 = mauritanica.
Cephalotes, Hoffmgg. & Link, Fl. Port. i. 440 = latifolia.
Cephalotes, Schousb. Vext. Marok. 146 = plantaginea.
Cephalotus, Boiss. in DC. Prod. xii. 675 = pinifolia.
chilensis, *Boiss. l. c.* 681.—Chili.
Choulettiana, *Pomel, Nouv. Mat. Fl. Atl.* 135.—Algeria.
cinerea, *Welw. ex Boiss. & Reut. Pugill. Pl. Nov.* 101.—Lusitan.
curvifolia, *Colla, in Mem. Acc. Torin.* xxxix. (1835) 5.—Hab.?
curvifolia, Bert. in Mere. Chil. 563; et Bull. Férussac, xx. (1830) 108 = chilensis.
denticulata, *Bertol. Amoen. Ital.* 77.—Italia.
denticulata, Portensch. Pl. Dalm. t. 6 = canescens.
dianthoides, Hornem. & Spreng. ex Boiss. in DC. Prod. xii. 683 = plantaginea.
Duriaei, *Boiss. l. c.* 684.—Hispan.
duriuscula, Bab. in Ann. Nat. Hist. Ser. II. iii. (1849) 436 = pubescens.
ebracteata, *Pomel, Nouv. Mat. Fl. Atl.* 132.—Algeria.
elongata, *Hoffm. Deutschl. Fl.* i. 150.—Europ.
elongata, Koch, in Mert. & Koch, Roehl. Deutschl. Fl. ed. III. ii. 487 = maritima.
eriophila, *Willk. in Bol. Soc. Broter.* ii. (1884) 145.—Lusitan.
expansa, Wallr. Beitr. i. 197 = pubescens.
fasciculata, *Willd. Enum. Hort. Berol.* i. 334.—Hispan.; Sardin.
fibrosa, *Pomel, Nouv. Mat. Fl. Atl.* 133.—Algeria.
filicaulis, *Boiss. Voy. Bot. Espagne*, ii. 527. t. 154.—Hispan.
formosa, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 848 = Statice Pseudo-armeria.
formosa, Heynh. Nom. ii. 45 = mauritanica.
fruticosa, *Lodd. Bot. Cab.* t. 1587.—Europ. austr.
gaditana, *Boiss. in DC. Prod.* xii. 675.—Hispan.; Maroeco.
glauca, Wallr. Arm. 215 = boctica.
glaucescens, *Desf. Cat. Hort. Par.* ed. III. 79.—Hab.?
globosa, Link, ex Boiss. in DC. Prod. xii. 684 = latifolia.
gracilis, Tenore, Syll. 158 = nebrodensis.
grandiflora, Hort. Schwetz. ex Boiss. in DC. Prod. xii. 684 = latifolia.
Gussonei, *Boiss. in DC. Prod.* xii. 687.—Sicil.
Halleri, Wallr. Beitr. i. 194 = alpina.
hetcrophylla, Wallr. l. c. 188 = nebrodensis.
hirta, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 369 = pubigera.
hirta, Willd. Enum. Hort. Berol. 333 = boctica.
humilis, Link, Handb. ii. 263 = caespitosa.

ARMERIA :—
intermedia, Link, ex Boiss. in DC. Prod. xii. 681 = pubescens.
juncea, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 324.—Gall. merid.
juncea, Wallr. Beitr. i. 212 = allioides.
juniperifolia, Ebel, Armeriae Gen. Diss. (1840) 30 = filicaulis.
juniperifolia, J. Gay, ex Boiss. in DC. Prod. xii. 684 = Duriaei.
juniperifolia, Koch, in Flora, vi. (1823) 711 = leucocephala.
juniperifolia, Willd. ex Hoffmgg. & Link, Fl. Port. i. 442 = caespitosa.
Kochii, Boiss. in DC. Prod. xii. 686 = leucocephala.
labradorica, *Wallr. Beitr.* i. 185.—Groenl.; Labrador.
lachnolepis, *Pomel, Nouv. Mat. Fl. Atl.* 134.—Algeria.
Langei, *Boiss. ex Lange, in Kjoeb. Vidensk. Meddel.* (1861) 59.—Hispan.
latifolia, Moris, Fl. Sard. Elench. fasc. iii. 10 = Morisii.
latifolia, *Willd. Enum. Hort. Berol.* 334.—Lusitan.
leucantha, Salzm. ex Boiss. Voy. Bot. Espagne, ii. 525 = allioides.
leucocephala, *Koch, in Flora*, vi. (1823) 712.—Corsica.
littoralis, Boiss. in DC. Prod. xii. 676 = Welwitschii.
littoralis, *Hoffmgg. & Link, Fl. Port.* t. i. 441.—Lusitan.
littoralis, Willd. Enum. Hort. Berol. 333 = boctica.
longcaristata, *Boiss. & Reut. Pugill. Pl. Nov.* 103.—Hispan.
longearistata, Bourg. in Bull. Soc. Bot. Fr. ii. (1855) 643 = plantaginea.
longifolia, *Desf. Cat. Hort. Par.* ed. III. 79, nomen.—Hab.?
lusitanica, Link, ex Boiss. in DC. Prod. xii. 675 = pinifolia.
macloviana, *Cham. in Linnaea*, vi. (1831) 567.—Ins. Falkland.
macrophylla, *Boiss. & Reut. Pugill. Pl. Nov.* 100.—Hispan.
macropoda, *Boiss. in DC. Prod.* xii. 688.—Italia.
maculata, Poepp. ex Boiss. l. c. 682 = andina.
maderensis, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1833) 12.—Madera.
majellensis, *Boiss. in DC. Prod.* xii. 685.—Italia.
maritima, Girard, ex Boiss. l. c. 680 = ruscinonensis.
maritima, *Willd. Enum. Hort. Berol.* 333.—Europ.
mauritanica, *Wallr. Beitr.* i. 217.—Afr. bor.
meridionalis, Poepp. ex Boiss. in DC. Prod. xii. 682 = chilensis.
montana, G. Don, ex Loud. Hort. Brit. 114 = maritima.
montana, Soleir. ex Wallr. Beitr. i. 196 = leucocephala.
montana, Wallr. l. c. 210 = plantaginea.
Morisii, *Boiss. in DC. Prod.* xii. 687.—Sicil.
Muelleri, Huet, in Ann. Sc. Nat. Sér. III. xix. (1853) 255 = alpina.
multiceps, Wallr. Beitr. i. 196 = leucocephala.
nebrodensis, *Guss. Fl. Sic. Syn.* i. 366.—Italia.
neglecta, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 324.—Lusitan.
Orphanidis, Boiss. Diagn. Ser. II. iv. 71 = majellensis.
pinifolia, Bourg. ex Nym. Consp. 614 = macrophylla.
pinifolia, *Hoffmgg. & Link, Fl. Port.* i. 437.—Lusitan.
planifolia, Nym. Consp. 616. n. 35 = pubescens.
plantaginea, Boiss. Voy. Bot. Espagne, 526 = boctica.
plantaginea, *Willd. Enum. Hort. Berol.* 334.—Europ.
plantaginea, Webb, It. Hisp. 18 = latifolia.
praecox, Jord. in Bull. Soc. Dauph. fasc. iii. 79 = plantaginea.
pseudoarmeria, Brot. Fl. Lusit. i. 488 = plantaginea.
pseud-Armeria, Murr. Syst. Veg. ed. XIV. 300 = latifolia.
pubescens, Heyne, ex Steud. Nom. ed. II. i. 133 = maritima.
pubescens, *Link*, '*in Repert. Nat. Cur. Berol.* i. 180.'—Europ.
pubigera, Boiss. in DC. Prod. xii. 678 = maritima.
pubinervis, *Boiss. l. c.* 688.—Gallia.
pungens, *Hoffmgg. & Link, Fl. Port.* i. 439.—Lusitan.
purpurea, Koch, in Flora, vi. (1823) 710 = elongata.
rhenana, Gremli, Exc. Fl. Schw. ed. IV. 350 = maritima.
rigida, Wallr. Beitr. i. 199 = plantaginea.
rumelica, Boiss. in DC. Prod. xii. 677 = cariensis.
ruscinonensis, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 323.—Hispan.

ARMERIA :—
sabulosa, Jord. ex Bor. Fl. Centr. Fr. ed. III. ii. 537 = plantaginea.
sanguinolenta, *Wallr. Beitr.* i. 207.—Am. bor.
sardoa, *Spreng. Syst. Cur. Post.* 127.—Sardin.; Corsic.
scabra, Kunze, ex Boiss. in DC. Prod. xii. 683 = andina.
scabra, *Pall. in Willd. ex Roem. & Schult. Syst.* vi. 776.—As. bor.
scabriuscula, Kunze, ex Boiss. in DC. Prod. xii. 683 = andina.
scorzoneraefolia, Tenore, Syll. 160 = macropoda.
scorzoneraefolia, Willd. Enum. Hort. Berol. 334 = plantaginea.
scorzoneraefolia, Link, in Schrad. Journ. iii. (1800) 60 = latifolia.
scorzonerifolia, Balb. & Nocca, Fl. Ticin. t. 6 = plantaginea.
scorzonerifolia, Friv. ex Nym. Consp. 615 = rumelica.
setacea, Delile, ex Nym. l. c. 616 = juncea.
seticeps, Reichb. Ic. Fl. Germ. v. 54 = plantaginea.
sibirica, *Turcz. ex Boiss. in DC. Prod.* xii. 678.—Sibiria.
sicula, Heldr. ex Boiss. l. c. 685 = nebrodensis.
simplex, *Pomel, Nouv. Mat. Fl. Atl.* 132.—Algeria.
Soleirolii, Duby, Bot. Gall. Suppl. 1032 = leucocephala.
spinulosa, *Boiss. in DC. Prod.* xii. 675.—Barbaria.
splendens, *Boiss. Voy. Bot. Espagne*, ii. 749.—Hispan.
stenophylla, Girard, in Ann. Sc. Nat. Sér. III. ii. (1844) 323 = plantaginea.
tenuis, Balb. ex Boiss. in DC. Prod. xii. 685 = sardoa.
tenuis, Wallr. Beitr. i. 184 = filicaulis.
thessala, Boiss. Diagn. Ser. II. iv. 70 = cariensis.
Thomasii, Nym. Consp. 615 = leucocephala.
tingitana, Boiss. & Reut. Pugill. Pl. Nov. 102 = gaditana.
trigonoides, Hort. Berol. ex Ebel, Armeriae Gen. Diss. (1840) 34 = plantaginea.
undulata, Boiss. in DC. Prod. xii. 685 = argyrocephala.
velutina, *Welw. ex Boiss. & Reut. Pugill. Pl. Nov.* 100.—Lusitan.
vestita, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 366 = longearistata.
villosa, *Girard, in Ann. Sc. Nat.* Sér. III. ii. (1844) 323.—Hispan.
vulgaris, Moris, Elench. ex Nym. Consp. 615 = sardoa.
vulgaris, Vis. Fl. Dalmat. ii. 6 = canescens.
vulgaris, Willd. Enum. Hort. Berol. 333 = elongata.
Welwitschii, *Boiss. in DC. Prod.* xii. 676.—Hispan.
Willkommii, *Henriques, in Bol. Soc. Broter.* iii. (1884) 199, 224.—Hispan.

ARMERIASTRUM, Lindl. Veg. Kingd. 641 (1847) = **Acantholimon**, Boiss. (Plumbag.).

ARMODORUM, Breda, Orch. Kuhl. & Hass. (1827) = **Arachnanthe**, Blume (Orchid.).
distichum, Breda, l. c. [t. 6] = Arachnanthe Sulingi.

ARMORACIA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 426 (1800); Bernh. Syst. Verz. Erf. '258' (1800) = **Cochlearia**, Linn. (Crucif.).
amphibia, Peterm. Fl. Lips. Excurs. 477 = Nasturtium amphibium.
aquatica, Kostel. Allg. Med.-Pharm. Fl. v. 1571 = Nasturtium amphibium.
austriaca, Wimm. Fl. Schles. (1841) 30 = Nasturtium austriacum.
lapathifolia, Gilib. Fl. Lituan. ii. 53 = C. Armoracia.
macrocarpa, Baumg. Enum. Stirp. Transsylv. ii. 240 = C. macrocarpa.
rustica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1858) 8 = seq.
rusticana, Gaertn. Mey. & Scherb. Fl. Wett. ii. 426 = C. Armoracia.
sativa, Bernh. Syst. Verz. Erf. 191 = C. Armoracia.
valdiviana, Phil. in Linnaea, xxvni. (1856) 671.—Chili.

ARNEBIA, Forsk. Fl. Aegypt. Arab. 62 (1775).
BORAGINEAE, Benth. & Hook. f. ii. 862.
DIOCLEA, Spreng. Syst. i. 502 (1825).
MENEGHINIA, Endl. Gen. 648 (1839).
STENOSOLENIUM, Turcz. in Bull. Soc. Nat. Mosc. (1840) 253.

ARNEBIA :—
Strobila, G. Don, Gen. Syst. iv. 327 (1837).
Toxostigma, A. Rich. Tent. Fl. Abyss. ii. 86. t. 77 (1851).
Bungei, *Boiss. Fl. Orient.* iv. 215.—Persia.
calycina, Stev. in Bull. Soc. Nat. Mosc. (1851) 598 = cornuta.
Cephalotes, A. DC. in DC. Prod. x. 96 = Macrotomia Cephalotes.
cornuta, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. (1835) 22.—Oriens; Turkest.
decumbens, *Coss. & Kralik, in Bull. Soc. Bot. Fr.* iv. (1857) 398, 402.—Oriens.
densiflora, Ledeb. Fl. Ross. iii. 140 = Macrotomia Cephalotes.
echioides, A. DC. in DC. Prod. x. 96 = Macrotomia echioides.
endochroma, Aitch. in Journ. Linn. Soc. xviii. (1882) 81 = Macrotomia perennis.
fimbriata, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 507.—As. centr.
fimbriopetala, *Stocks, in Hook. Kew Journ.* iii. (1851) 180.—Persia; Beluchist.
flavescens, Boiss. Diagn. Ser. I. xi. 117 = linearifolia.
Griffithii, *Boiss. l. c.* Ser. II. iii. 135.—Ind. bor. occ.; Affghan.
guttata, *Bunge, Ind. Sem. Hort. Dorpat.* (1840) 7.—Sibir.
hispidissima, *DC. Prod.* x. 94.—Afr. trop.; Arab.; Ind. bor. occ.
linearifolia, *DC. l. c.* 95.—Aegypt.; Syria; Persia.
longiflora, C. Koch, in Linnaea, xxii. (1849) 640 = Macrotomia echioides.
minima, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 30.—Persia.
obovata, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 407 (*Lehm. Rel. Bot.* 231).—Turkest.; Ind. or.
Olgae, *Regel, Pl. Nov. Fedsch.* 58.—As. centr.
perennis, A. DC. in DC. Prod. x. 95 = Macrotomia perennis.
Sewerzowi, *Regel, Pl. Nov. Fedsch.* 58.—As. centr.
tetrastigma, *Forsk. Fl. Aegypt. Arab.* 62.—Arab.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 176.—Tibet. occ.
tibetana, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 189.—Tibet. occ.
tinctoria, *Forsk. Fl. Aegypt. Arab.* 63.—Aegypt.; Arab.
tingens, A. DC. in DC. Prod. x. 96 = Macrotomia perennis.
Viviani, Coss. & Dur. ex Coss. in Ann. Sc. Nat. Sér. IV. i. (1854) 240 = cornuta.

ARNEDINA, Reichb. Nom. 'n. 1984' (1841) = **Arundina**, Blume (Orchid.).

ARNICA, Rupp. ex Linn. Syst. ed. I. (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 440.
alpina, *Olin & Ladau, Diss.* 11.—Europ.; Am. bor.
alpina, Salisb. Prod. 201 = Doronicum scorpioides.
alpina, Willd. ex Steud. Nom. ed. I. 70 = montana.
altaica, *Turcz. in Bull. S c. Nat. Mosc.* xx. (1847) ii. 12.—Mont. Altai.
amplexicaulis, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 408.—Am. bor. occ.
amplexicaulis, Wall. ex DC. Prod. vi. 314 = Senecio amplexicaulis.
angustifolia, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 238 = Parryi.
angustifolia, Turcz. ex Ledeb. Fl. Ross. ii. 622 = montana.
angustifolia, Vahl, Fl. Dan. t. 1524 = alpina.
auriculata, Willd. ex Ledeb. Fl. Ross. ii. 632 = Senecio caucasicus.
austriaca, Hoppe, ex Schur, Enum. Pl. Transs. 340 = Doronicum austriacum.
Bellidiastrum, All. Fl. Pedem. i. 205 = Aster Bellidiastrum.
brevicaulis, *Rafin. New Fl. Am.* ii. 23.—Am. bor.
caffra, *Linn. Syst.* ed. X. 1219.—Afr. austr.
Chamissonis, *Less. in Linnaea*, vi. (1831) 238.—Am. bor.; Unalaschk.
ciliata, Thunb. Fl. Jap. 318 = Hypochoeris grandiflorus.
Claytoni, Pursh, Fl. Am. Sept. ii. 527 = nudicaulis.
Clusii, All. Fl. Pedem. i. 205. t. 17. f. 1 = Doronicum Clusii.

ARNICA :—
cordata, E. Mey. ex DC. Prod. vii. 16 = Gerbera tomentosa.
cordata, Thunb. Prod. Pl. Cap. 154 = Gerbera cordata.
cordata, Wulf. in Roem. Arch. iii. iii. 408 = Doronicum caucasicum.
cordifolia, *Hook. Fl. Bor. Am.* i. 331.—Am. bor.
coronopifolia, Linn. Sp. Pl. 885 = Gerbera coronopifolia.
corsica, Loisel. Fl. Gall. 576. t. 20 = Doronicum corsicum.
crenata, Thunb. Prod. Pl. Cap. 154 = Mairia crenata.
crocea, Linn. Sp. Pl. 885 = Gerbera Burmanni.
discoides, *Benth. Pl. Hartw.* 319.—Calif.
Doronicum, Jacq. Fl. Austr. i. 57 = Senecio Doronicum.
Doronicum, Pursh, Fl. Am. Sept. ii. 528 = Senecio Pseudo-arnica.
floccosa, Bert. ex DC. Prod. vi. 357 = Senecio arachnoideus.
foliosa, *Nutt. in Trans. Am. Phil. Soc.* N.S. vii. (1841) 407.—Am. bor.
fulgens, Pursh, Fl. Am. Sept. ii. 527 = alpina.
Gerbera, Burm. f. Fl. Cap. Prod. 27 = Gerbera ferruginea.
Gerbera, Linn. Sp. Pl. 885 = Gerbera asplenifolia.
glacialis, Wulf. in Jacq. Coll. i. 230; et Ic. Rar. iii. 15. t. 586 = Doronicum glaciale.
grandis, Thunb. Prod. Pl. Cap. 154 = Alciope lanata.
helvetica, G. Don, ex Loud. Hort. Brit. 351 = montana.
hirsuta, Forsk. Fl. Aegypt. Arab. 151 = Gerbera hirsuta.
incana, Pers. Syn. ii. 452 = Trichocline incana.
intermedia, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 203.—Sibir.
inuloides, Vahl, Symb. Bot. ii. 91 = Heterolepis decipiens.
japonica, Thunb. Fl. Jap. 319 = Senecio japonicus.
lanata, Thunb. Prod. Pl. Cap. 154 = Alciope lanata.
lanceolata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 407 = Chamissonis.
lanigera, Tenore, Prod. Fl. Neap. (1811) 49 = Senecio arachnoideus.
latifolia, *Bongard, Vég. Isl. Sitch.* 29.—Am. bor. occ.
linifolia, Wall. Cat. n. 3285 = Senecio linifolius.
longifolia, *Eaton, in Bot. King, Exped.* 186.—Am. bor. occ.
macrophylla, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 408 = cordifolia.
maritima, Linn. Sp. Pl. 884 = Senecio Pseudo-arnica.
Menziesii, Hook. Fl. Bor. Am. i. 331 = latifolia.
microphylla, Walp. Rep. ii. 652, err. typ. = *macrophylla*.
Mollii, *Schrank, Prim. Fl. Salisb.* 213.—Europ.
mollis, Hook. Fl. Bor. Am. i. 331 = Chamissonis.
montana, Hook. l. c. 330 (pro parte) = foliosa.
montana, *Linn. Sp. Pl.* 884.—Europ.; As. bor.
nevadensis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 55.—Calif.
nivea, Wall. Cat. n. 3143 = Gerbera nivea.
nudicaulis, *Ell. Sketch*, ii. 333.—Am. bor.
obtusifolia, *Less. in Linnaea*, vi. (1831) 236.—Unalaschk.
obtusifolia, Wall. ex DC. Prod. vi. 314 = Senecio arnicoides.
operina, Rich. & Less. Fl. Nov. Zel. 231 = seq.
oporina, Forst. f. Prod. 56 = Olearia operina.
palmata, Thunb. Fl. Jap. 319 = Senecio syneilesis.
Parryi, *A. Gray, in Am. Nat.* viii. (1874) 213.—Am. bor. occ.
parviflora, *A. Gray, in Proc. Am. Acad.* vii. (1868) 363.—Calif.
peruviana, Pers. Syn. ii. 452 = Werneria rigida.
petiolata, *Schur, Enum. Pl. Transs.* 342.—Europ.
Pilosella, Burm. ex DC. Prod. vii. 17 (= *an* Arctotidea? —*Cf. DC. in loc.*).—Afr. austr.
piloselloides, Linn. Pl. Afr. Rar. 22. n. 73 (1760) = Gerbera piloselloides.
plantaginea, *Pursh, Fl. Am. Sept.* ii. 527.—Am. arct.
plantaginisfolia, Gilib. Fl. Lituan. i. 210 = montana.
pyrolaefolia, Pers. Syn. ii. 454 = Gerbera Burmanni.
rotundifolia, Willd. Sp. Pl. iii. 2110 = Bellium rotundifolium.
sachalinensis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 55.—Ins. Sachal.
scorpioides, Adams, ex Ledeb. Fl. Ross. ii. 632 = Senecio caucasicus.

ARNICA :—
scorpioides, *Jacq. Austr.* iv. 26. t. 349.—Europ.
scorpioides, Linn. Sp. Pl. 884 = Doronicum grandiflorum.
senecionoides, Fenzl, ex Boiss. Fl. Orient. iii. 396 = Senecio cilicius.
serrata, Thunb. Fl. Cap. ed. Schult. 669 = Gerbera ferruginea.
sinuata, Thunb. Prod. Pl. Cap. 154 = Gerbera ferruginea.
spinulosa, Pers. Syn. ii. 452 = Alciope tabularis.
styriaca, Vill. Prosp. 32 = Doronicum Clusii.
tabularis, Thunb. Prod. Pl. Cap. 154 = Alciope tabularis.
tussilaginea, Burm. f. Fl. Ind. 182 = Senecio Kaempferi.
unalaschkensis, *Less. in Linnaea*, vi. (1831) 238.—Ins. Aleut.; Unalaschk.
viscosa, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 374.—Calif.
Wulfeniana, Pollini, in Giorn. Fis. Pav. ix. (1826) 185 = Doronicum Columnae.

ARNOCRINIUM, Endl. & Lehm. ex Lehm. Pl. Preiss. ii. 41 (1846). *LILIACEAE*, Benth. & Hook. f. iii. 795.
Drummondii, *Endl. ex Lehm. l. c.*—Austral.
glabrum, *Baker, in Journ. Linn. Soc.* xvii. (1880) 416.—Austral.
Preissii, *Lehm. Pl. Preiss.* ii. 42.—Austral.

ARNOGLOSSUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 292 (1821) = **Plantago**, Tourn.
alpinum, S. F. Gray, l. c. 294 = P. alpina.
gramineum, S. F. Gray, l. c. 293 = P. maritima.
incanum, S. F. Gray, l. c. 292 = P. media.
lanceolatum, S. F. Gray, l. c. 293 = P. lanceolata.
maritimum, S. F. Gray, l. c. = P. maritima.
subulatum, S. F. Gray, l. c. 294 = P. Coronopus.

ARNOGLOSSUM, Rafin. Fl. Ludov. 64 (1817). *COMPOSITAE*.
plantagineum, Rafin. l. c. 65 (quid?).—Am. bor.

ARNOLDIA, Blume, Bijdr. 868 (1826) = **Weinmannia**, Linn. (Saxifr.).
fraxinifolia, Blume, in Flora, xli. (1858) 254 = W. fraxinifolia, *Miq.*
heterophylla, Blume, Bijdr. 868 = W. Blumei, *A. Gray*.
pinnata, Blume, l. c. = W. Blumei, *Planch.*

ARNOLDIA, Cass. in Dict. Sc. Nat. xxx. 330 (1824) = **Dimorphotheca**, Moench (Compos.).
aurea, Cass. l. c. 331 = D. chrysanthemifolia.
chrysanthemifolia, Steud. Nom. ed. II. i. 256 = D. chrysanthemifolia.
Tragus, Steud. Nom. ed. II. i. 256 = D. Tragus.
Ventenatiana, Presl, Repert. 192, in nota = D. chrysanthemifolia.
viscosa, Steud. Nom. ed. II. i. 256 = D. cuneata.

ARNOPOGON, Willd. Sp. Pl. iii. 1496 (1803) = **Urospermum**, Scop. (Compos.).
asper, Willd. l. c. 1497 = U. picroides.
capensis, Willd. l. c. = U. picroides.
Dalechampii, Willd. l. c. 1496 = U. Dalechampii.
picroides, Willd. l. c. = U. picroides.

ARNOSERIS, Gaertn. Fruct. ii. 355. t. 157 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 507.
foetida, Dum. Fl. Belg. 63 = Crepis foetida.
gracilis, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 301 = Apogon gracilis.
humilis, Sch. Bip. l. c. = Apogon humilis.
minima, Dum. Fl. Belg. 63 = pusilla.
pusilla, *Gaertn. Fruct.* ii. 355. t. 157.—Europ.

ARNOTTIA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 29. *ORCHIDEAE*, Benth. & Hook. f. iii. 622.
inermis, *S. Moore, in Baker, Fl. Maurit.* 339.—Ins. Mascar.
mauritiana, *A. Rich. in Mém. Soc. Hist. Nat. Par.* iv. (1828) 30.—Ins. Mascar.

AROMADENDRON, Blume, Bijdr. 10 (1825) = **Talauma**, Juss. (Magnol.).
elegans, Blume, l. c. 10 = T. elegans.
glaucum, Korth. ex Miq. Fl. Ind. Bat. i. 2. 16 = T. glauca.

AROMADENDRON, Andr. ex Steud. Nom. ed. II. i. 134 (1840) et AROMADENDRUM, Andr. ex Steud. l. c. 599, in syn. (1840) = **Eucalyptus**, L'Hérit. (Myrtac.).

AROMIA, Nutt. in Trans. Am. Phil. Soc. Scr. II. vii. (1841) 395 = **Amblyopappus**, Hook. (Compos.).
tenuifolia, Nutt. l. c. 396 = Amblyop. pusillus.

ARON, Adans. Fam. ii. 470 (1763).
dracunculum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = Dracunculus vulgaris.
colocasium, St. Lag. l. c. = Colocasia antiquorum.

ARONGANA, Choisy, in DC. Prod. i. 541 (1824) = **Haronga**, Thou. (Hyperic.).

ARONIA, Mitchell, in Act. Acad. Nat. Cur. viii. (1748) 206, App. = **Orontium**, Linn. (Aroid.).

ARONIA, Medic. Phil. Bot. i. 140 (1789) = **Osteomeles**, Lindl. (Rosac.).

ARONIA, Pers. Syn. ii. 39 (1807) = **Amelanchier**, Lindl. (Rosac.).
affinis, Neum. ex Tausch, in Flora, xxi. (1838) i. Beibl. i. 77 = Amel. canadensis.
alnifolia, Nutt. Gen. Am. i. 306 = Amel. alnifolia.
Amelanchier, Dum. Fl. Belg. 92 = Amel. vulgaris.
arborea, Barton, Comp. Fl. Phil. i. 228 = Amel. canadensis.
arbutifolia, Medic. Phil. Bot. i. 140 = Pyrus arbutifolia.
aria-chamaemespilus, Reichb. Fl. Germ. Excurs. 630 = Pyrus Chamaemespilus.
asiatica, Sieb. & Zucc. Fl. Jap. i. 87. t. 42 = Amel. canadensis.
Botryapium, Pers. Syn. ii. 39 = Amel. canadensis.
Chamaemespilus, Pers. l. c. = Pyrus Chamaemespilus.
cordata, Rafin. Med. Fl. ii. 196 = Amel. canadensis.
cretica, Pers. l. c. 40 = Amel. vulgaris.
densiflora, Spach, Hist. Vég. Phan. ii. 88 = Pyrus arbutifolia.
depressa, Spach, l. c. 90 = Pyrus arbutifolia.
floribunda, Spach, l. c. 89 = Pyrus arbutifolia.
floribunda, Sweet, Hort. Brit. ed. I. 485 = Pyrus spectabilis.
glabrescens, Spach, Hist. Vég. Phan. ii. 89 = Pyrus arbutifolia.
grandifolia, Sweet, Hort. Brit. ed. II. 177 = Pyrus arbutifolia.
Hostii, Carr. in Rev. Hortic. (1873) 470 = Pyrus intermedia.
japonica, Hort. ex C. Koch, Dendrol. i. 179 = Pyrus japonica.
latifolia, Riddell, Suppl. Cat. Pl. Ohio, 24 = Amel. canadensis ?
melanocarpa, Nutt. ex Ell. Sketch, i. 556 = Pyrus arbutifolia.
nivea, Neum. ex Tausch, in Flora, xxi. (1838) 714 = Amel. canadensis.
ovalis, Pers. Syn. ii. 39 = Amel. canadensis.
petraea, Wallr. Beitr. Fl. Hercyn. 206.—Europ.
praecox, Neum. ex Tausch, in Flora, xxi. (1838) 715 = Amel. canadensis.
pubens, Spach, Hist. Vég. Phan. ii. 90 = Pyrus arbutifolia.
pumila, M. Roem. Syn. iii. Rosifl. 161 = Pyrus arbutifolia.
pyrifolia, Lam. Encyc. i. 83 = Pyrus arbutifolia.
racemosa, Schur, Enum. Pl. Transs. 207 = Amel. cretica.
rotundifolia, Pers. Syn. ii. 39 = Amel. vulgaris.
sanguinea, Nutt. Gen. Am. i. 306 = Amel. canadensis.
sorbifolia, Spach, Hist. Vég. Phan. ii. 87 = Pyrus spuria.
spuria, M. Roem. Syn. iii. Rosifl. 158 = Pyrus spuria.
subcordata, Rafin. ex DC. Prod. ii. 632 = Amel. canadensis.
tomentosa, Wallr. Beitr. Fl. Hercyn. 206 = Amel. vulgaris ?
Watsoniana, M. Roem. Syn. iii. Rosifl. 159 = Pyrus pinnatifida.

ARONICUM, Neck. Elem. i. 27 (1790) = **Doronicum**, Linn. (Compos.).
altaicum, DC. Prod. vi. 320 = D. altaicum.

ARONICUM :—
Bauhini, Reichb. f. Ic. Fl. Germ. xvi. 32 = D. hirsutum.
carpathicum, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 137 = D. grandiflorum.
Clusii, Koch, Syn. ed. I. 382 = D. hirsutum.
cordatum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 41 = D. cordatum.
corsicum, DC. Prod. vi. 319 = D. corsicum.
Doronicum, Reichb. Fl. Germ. Excurs. 233 = D. Clusii.
glaciale, Reichb. l. c. 234 = D. glaciale.
latifolium, Reichb. l. c. = D. grandiflorum.
scorpioides, DC. Prod. vi. 319; Koch, Syn. ed. I. 382 = D. grandiflorum.
scorpioideum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = D. grandiflorum.
viscosum, Freyn & Gautier, in Bull. Soc. Bot. Fr. xxviii. (1881) 50.—Europ.

AROSMA, Rafin. Fl. Tellur. iii. 66 (1836) = **Caladium**, Vent. (Aroid.).
acutifolia, Rafin. l. c. = C. bicolor ?
obtusifolia, Rafin. l. c. = C. bicolor ?

AROTON, Neck. Elem. ii. 336 (1790) = **Croton**, Linn. (Euphorb.).

AROUNA, Aubl. Pl. Gui. i. 16. t. 5 (1775) = **Dialium**, Linn. (Legumin.).
divaricatum, Willd. Sp. Pl. i. 156 = D. divaricatum.
guianensis, Aubl. Pl. Gui. i. 16 = D. divaricatum.

ARPITIUM, Neck. Elem. i. 168 (1790) = **Ligusticum**, Linn. (Umbell.).
simplex, Sweet, Hort. Brit. ed. II. 591 = L. simplex.

ARPOPHYLLUM, Llave & Lex. Nov. Veg. Descr. fasc. ii. (Orch. Opusc.) 19 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 492.
alpinum, *Lindl. in Benth. Pl. Hartw.* 93 (1842).—Mexic.
cardinale, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 282.—Mexic.
giganteum, *Hartw. ex Lindl. in Ann. Nat. Hist.* iv. (1840) 384.—Mexico.
medium, *Reichb. f. Beitr. Orch. Centr. Am.* 89.—Guatemala.
spicatum, *Llave & Lex. Nov. Veg. Descr.* fasc. ii. (*Orch. Opusc.*). 20.—Mexico.
squarrosum, Hort. Donat. ex Lubbers, Cat. Pl. Rar. San Donato (1880) 15 = spicatum.

ARRABIDAEA, DC. in Bibl. Univ. Genèv. xvii. (1838) 126 (Rev. Bign. 10). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1032.
Chasmia, Schott, in Spreng. Syst. Cur. Post. 409 (1827).
Vasconcellia, Mart. in Flora, xxiv. (1841) ii. Beibl. 12 (Herb. Fl. Bras. 252).
acutifolia, *DC. Prod.* ix. 185.—Bras.
Agnus-castus, *DC. l. c.* 183.—Bras.
argentea, *Wawra, in Oestr. Bot. Zeitschr.* (1879) 216.—Bras.
Blanchetii, DC. Prod. ix. 186 = Bignonia Blanchetii.
candicans, *DC. l. c.* 185.—Guiana.
Chica, Verl. in Rev. Hortic. (1866) 154 = Bignonia Chica.
Clausseni, *DC. l. c.* 186.—Bras.
conjugata, *Mart. in Flora*, xxiv. (1841) ii. *Beibl.* 46.—Bras.
corchoroides, *DC. Prod.* ix. 183.—Bras.
cordifolia, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1158.—Guiana.
desertorum, *DC. Prod.* ix. 185.—Bras.
fasciculata, *DC. l. c.*—Bras.
florida, *DC. l. c.* 184.—Bras.
obovata, *DC. l. c.* 185.—Bras.
Orbignyana, *DC. l. c.* 184.—Bolivia.
paniculata, *Seem. in Trans. Linn. Soc.* xxiii. (1862) 14.—Ins. Martinic.
platyphylla, *DC. Prod.* ix. 186.—Bras.
Pleei, *DC. l. c.* 184.—Venezuela.
pruinosa, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1158.—Guiana.
rosea, *DC. Prod.* ix. 185.—Bras.
rubrinervis, *Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 192.—Bras.
Salzmanni, *DC. Rev. Bign.* 10. (1838).—Bras.

ARRABIDAEA :—
Sego, DC. in Bibl. Univ. Genèv. xvii. (1838) 126; et Prod. ix. 183 = A. Agnus-castus.
Sieberi, *DC. l. c.* 186.—Bras.
subincana, DC. l. c. 184 = Bignonia subincana.
tuberculata, *DC. l. c.*—Bras.
virescens, *DC. l. c.*—Bras.

ARRABIDAEA, Steud. Nom. ed. II. i. 135 (1840) = **Cormonema**, Reissek (Rhamn.).
spinosa, Steud. l. c. = C. spinosum.

ARRACACHA, DC. in Bibl. Univ. Genèv. xl. (1829) 74; et Prod. iv. 243 (1830) = seq.

ARRACACIA, Bancr. in Jamaic. Journ. iv. (1826) 18. *UMBELLIFERAE*, Benth. & Hook. f. i. 884.
Arracacha, DC. in Bibl. Univ. Genèv. xl. (1829) 74; et Prod. iv. 243 (1830).
Bancroftia, Billb. in Linneska Samfund. Handl. i. 1832 (1833) 39.
Deweya, Torr. & Gray, Fl. N. Am. i. 641 (1840).
Pentacrypta, Lehm. in Linnaea, v. (1830) 380. t. 5.
Velaea, DC. Prod. iv. 230 (1830).
acuminata, *Benth. Pl. Hartw.* 187.—Mexico.
atropurpurea, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* i. 564.—Mexico.
decumbens, *Benth. & Hook. f. l. c.*—Mexico.
esculenta, DC. Prod. iv. 244 = xanthorrhiza.
glaucescens, *Benth. Pl. Hartw.* 187.—Mexico.
moschata, *DC. Prod.* iv. 244.—Am. merid.
tolucensis, *Hemsl. Biol. Centr. Am. Bot.* i. 564.—Mexico.
xanthorrhiza, *Bancr. in Jamaic. Journ.* iv. (1826) 18.—Jamaica.

ARRHENACHNE, Cass. in Dict. Sc. Nat. lii. 253 (1828) = **Baccharis**, Linn. (Compos.).
juncea, Cass. l. c. = B. juncea.

ARRHENATHERUM, Beauv. Agrost. 55. t. 11. f. 5 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1161.
americanum, Beauv. l. c. 56 = Avena pensylvanica.
asperum, Opiz, in Lotos, iii. (1853) 65 = avenaceum.
avenaceum, *Beauv. Agrost.* 55. t. 11. f. 5.—Europ.
avenaceum, Boiss. Voy. Bot. Espagne, ii. 657 = erianthum.
biaristatum, Peterm. Fl. Lips. Excurs. 106 = avenaceum.
bulbosum, Kunze, ex Anderss. in Walp. Ann. vi. 1003 = erianthum.
bulbosum, Presl, Cyp. et Gram. Sic. 29 = avenaceum.
cechicum, Opiz, in Lotos, iii. (1853) 66 = avenaceum.
elatius, Beauv. Agrost. 56 = avenaceum.
erianthum, *Boiss. & Reut. Diag. Pl. Hispan.* 121.—Lusit.; Alg.
exserens, Opiz, in Lotos, iii. (1853) 65 = avenaceum.
kentuckense, *Torr. ex Steud. Nom.* ed. II. i. 135.—Am. bor.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 122.—Oriens.
longifolium, Dulac, Fl. Hautes-Pyr. 78 = pallens.
palaestinum, Boiss. Diagn. Ser. I. xiii. 51 = avenaceum.
pallens, *Link, Hort. Berol.* i. 124.—Lusit.; Hisp.
pennsylvanicum, Torr. Fl. N. Am. i. 130 = Avena pensylvanica.
precatorium, Beauv. Agrost. 56 = avenaceum.
rivulare, *Opiz, in Lotos*, iii. (1853) 65.—Europ.
rupestre, *Dum. Obs. Gram. Belg.* 146.—Europ.
Thorei, Desm. Cat. Dord. (1840) = pallens.
zavadilianum, Opiz, in Lotos, iii. (1853) 66 = avenaceum.

ARRHOSTOXYLON, Mart. ex Nees, in Mart. Fl. Bras. ix. 57. t. 6 et 209 (1847) = seq.

ARRHOSTOXYLUM, Mart. ex Nees, in DC. Prod. xi. 209 (1847) = **Ruellia**, Linn. Acanth.).
achimeniflorum, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 131 = R. achimeniflorum.
acutangulum, Mart. ex Nees, in Mart. Fl. Bras. ix. 58 = R. acutangula.
affine, Nees, in Mart. Fl. Bras. ix. 59.—Bras.
album, Nees, l. c. 60 = Ruellia alba.
amplexicaule, Nees, l. c. 59.—Bras.
ciliatiflorum, Nees, in DC. Prod. xi. 216 = R. ciliatiflora.

ARRHOSTOXYLUM:—
coccineum, Nees, in Mart. Fl. Bras. ix. 63 = R. coccinea.
costatum, Nees, l. c. 61 = R. costata.
curviflorum, Nees, l. c. 62 = R. curviflora.
formosum, Nees, in DC. Prod. xi. 215 = R. formosa.
fulgidum, Nees, in Mart. Fl. Bras. ix. 59 = R. fulgida.
glabrum, Nees, l. c. 58.—Bras.
Haenkeanum, Nees, in DC. Prod. xi. 211.—Peruv.
Hartwegianum, Nees, l. c. 209.—N. Granat.
inflatum, Nees, l. c. 215 = R. inflata.
jussieuoides, Nees, l. c. 212 = R. jussieuoides.
laxum, Nees, in Mart. Fl. Bras. ix. 57.—Bras.
longiflorum, Nees, l. c. 62 = seq.
longifolium, Nees, in DC. Prod. xi. 214 = R. longifolia.
microphyllum, Nees, in Mart. Fl. Bras. ix. 61 = Calophanes microphylla?
nitidum, Nees, l. c. 59.—Bras.
pedunculosum, Nees, in DC. Prod. xi. 211 = R. pedunculosa.
repens, Mart. Fl. Bras. ix. 60.—Bras.
roseum, Nees, l. c. 61.—Bras.
rubrum, Nees, in DC. Prod. xi. 209 = R. rubra.
silvaccola, Nees, in Mart. Fl. Bras. ix. 60 = R. formosa.
splendidulum, Nees, l. c.—Bras.
stemonacanthoides, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 130 = R. stemonacanthoides.
subsessile, Nees, in Mart. Fl. Bras. ix. 58.—Bras.
terminale, Nees, in DC. Prod. xi. 214.—Peruv.
violaceum, Nees, l. c. 216 = R. violacea.
Willdenovianum, Nees, l. c. 209.—Ind. occ.

ARRHYNCHIUM, Lindl. in Paxt. Flow. Gard. i. 142 (1850–51) = **Arachnanthe**, Blume (Orchid.).
labrosum, Lindl. & Paxt. l. c. = Arachn. bilinguis.

ARROSTIA, Rafin. Caratteri, 75 (1810) = **Gypsophila**, Linn. (Caryophyll.).
altissima, Rafin. Fl. Tellur. ii. 54 = G. altissima.
dichotoma, Rafin. Caratteri, 75 = G. Arrostii.
paniculata, Rafin. Fl. Tellur. ii. 53 = G. paniculata.

ARROWSMITHIA, DC. Prod. vii. 254 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 329.
styphelioides, *DC. l. c.*—Afr. austr.

ARROZIA, Schrad. ex Kunth, Enum, Pl. i. 11 (1833) = **Luziola**, Juss. (Gramin.).
micrantha, Schrad. l. c. (= *Caryochloa brasiliensis*). —Bras.

ARRUDEA, A. St. Hil. & Camb. Fl. Bras. Mer. i. 318. t. 66 (1827) = **Clusia**, Linn. (Guttif.).
bicolor, Benth. in Hook. Kew Journ. iii. (1851) 166 = C. bicolor.
clusioides, A. St. Hil. et Camb. Fl. Bras. Mer. i. 319.—Bras.
purpurea, Splitg. in Hoev. & De Vriese, Tijdschr. ix. (1842) 103.—Guiana.

ARSACE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 113 = **Erica**, Linn.
arborea, Fourr. l. c. = E. arborea.

ARSIS, Lour. Fl. Cochinch. 335 (1790) = **Grewia**, Linn. (Tiliac.).
rugosa, Lour. l. c. = G. Microcos.

ARTABOTRYS, R. Br. in Bot. Reg. t. 423 (1820). *ANONACEAE*, Benth. & Hook. f. i. 24.
ROPALOPETALUM, Griff. Notul. iv. 16 (1845).
Blumei, Benth. Fl. Hongk. 10 = honkongensis.
Blumei, *Hook. f. & Thoms. Fl. Ind.* i. 128.—Java.
brachypetalus, *Benth. in Trans. Linn. Soc.* xxiii. (1862) 467.—Afr.
burmanicus, *DC. in Mém. Soc. Phys. Genèv.* v. (1832) 212.—Burma.
caudatus, *Wall. Cat.* n. 6417.—Reg. Himal.
crassifolius, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 54.—Burma; Malaya.
Cumingianus, *Vidal, Phan. Cuming. Philipp.* 169.—Ins. Philipp.
densiflorus, *Wall. ex Voigt, Hort. Suburb. Calc.* 16, nomen.—Ind. or.
hamatus, Benth. in Hook. Kew Journ. iii. (1851) 257 = honkongensis.

ARTABOTRYS:—
hamatus, Blume, Fl. Jav. Anon. 60. t. 29 = odoratissimus.
Hildebrandtii, *O. Hoffm. Sert. Pl. Madag.* 3.—Madag.
honkongensis, *Hance, in Journ. Bot.* viii. (1870) 71.—Hongkong.
inodorus, *Zipp. in Ann. Mus. Bot. Lugd. Bat.* ii. 41.—N. Guinea.
intermedius, *Hassk. Cat. Pl. Hort. Bogor.* 175.—Java.
Kurzii, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 54.—Burma.
macrophyllus, *Hook. f. in Hook. Niger Fl.* 207.—Afr. trop.
madagascariensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 42.—Madag.
Maingayi, *Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 55.—Malacca.
malayanus, Griff. Notul. iv. 713 = Xylopia ferruginea.
odoratissimus, Blume, Fl. Jav. Anon. 59 = honkongensis.
odoratissimus, *R. Br. in Bot. Reg.* t. 423.—Ind. or.; Malaya.
parviflorus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 41.—Sumatr.
pleurocarpus, *Maing. ex Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 54.—Malacca.
polygynus, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 41.—Borneo.
siamensis, *Miq. l. c.* 42.—Siam.
speciosus, *Kurz, ex Hook. f. & Thoms. in Hook. f. Fl. Brit. Ind.* i. 55.—Ins. Andam.
suaveolens, *Blume, Fl. Jav. Anon.* 62.—Ind. or.; Malaya.
sumatranus, *Miq. Fl. Ind. Bat. Suppl.* i. 375.—Sumatr.
Thomsoni, *Oliver, Fl. Trop. Afr.* i. 29.—Afr. trop.
unguiculatus, *Wall. Cat.* n. 6415 F.—Ind. or.
uncatus, Baill. Hist. Pl. i. 232 = odoratissimus.
zeylanicus, *Hook. f. & Thoms. Fl. Ind.* i. 128.—Ind. or.

ARTANEMA, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 234 (1829). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 953.
ACHIMENES, Vahl, Symb. Bot. ii. 71 (1791).
DICEROS, Pers. Syn. ii. 164 (1807).
angustifolium, *Benth. Scroph. Ind.* 39.—Singapore.
fimbriatum, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 234.—Austral.
sesamoides, *Benth. Scroph. Ind.* 39.—Ind. or.

ARTANTHE, Miq. Comm. Phyt. 40. t. 7, 8 (1840) = **Piper**, Linn.
abutilifolia, Miq. Syst. Pip. 385 = P. abutilifolium.
abutiloides, Miq. l. c. 384 = P. abutiloides.
acutifolia, Miq. l. c. 428 = P. acutifolium.
adenandra, Miq. l. c. 515 = P. adenandrum.
adenophora, Miq. l. c. 514 = P. adenophorum.
adenophylla, Miq. in Linnaea, xx. (1847) 170 = P. colubrinum.
aduncа, Miq. Comm. Phyt. 49. t. 4 = P. aduncum.
aequalis, Miq. Syst. Pip. 511 = P. aequale.
aequirophylla, Miq. l. c. 391 = P. cardiophyllum.
affinis, Miq. in Linnaea, xviii. (1844) 228 = P. affine.
alaris, Miq. Syst. Pip. 406 = P. alare.
albida, Miq. l. c. 415 = P. albidum.
alnoides, Miq. l. c. 462 = P. alnoides.
alpina, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 451 = P. alpinum.
alveolata, Miq. Syst. Pip. 406 = P. alveolatum.
amazonica, Miq. in Linnaea, xx. (1847) 146 = P. amazonicum.
ampla, Miq. Syst. Pip. 501 = P. Rohrii.
amplectens, Miq. in Linnaea, xx. (1847) 154 = P. amplectens.
andicola, Miq. Syst. Pip. 415 = P. andicolum.
angusta, Miq. l. c. 398 = P. angustum.
angustifolia, Miq. l. c. 528 = P. consanguineum.
anonaefolia, Miq. l. c. 504 = P. anonaefolium.
apiculata, Klotzsch, in Hook. Journ. Bot. iv. (1842) 321 = P. anonaefolium.
appendiculata, Benth. Pl. Hartw. 254 = P. appendiculatum.
arcuata, Miq. Syst. Pip. 533 = P. Opizianum.
areolata, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 454 = P. areolatum.
aspera, Miq. l. c. 456 = P. hirsutum.

ARTANTHE:—
asperifolia, Miq. Syst. Pip. 441 = Piper hirsutum.
asperiuscula, Miq. l. c. 464 = P. hirsutum.
auriculata, Miq. in Linnaea, xx. (1847) 167 = P. auriculatum.
aurita, Miq. Syst. Pip. 400 = P. auritum.
avellana, Miq. l. c. 478 = P. avellanum.
baccans, Miq. in Linnaea, xx. (1847) 166 = P. baccans.
bahiensis, Presl, Epim. Bot. 227 = P. bahiense.
barbata, Miq. Syst. Pip. 414 = P. barbatum.
Barclayana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 460 = P. hirsutum.
Bartlingiana, Miq. Syst. Pip. 510 = P. Bartlingianum.
Beecheyana, Miq. l. c. 403 = P. ellipticum.
begoniaefolia, Miq. l. c. = P. begoniaefolium.
Benthamiana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 447 = P. Benthamianum.
berbicensis, Miq. Syst. Pip. 500 = P. Hostmannianum.
Blanchetii, Miq. l. c. 473 = P. Blanchetii.
Botrytis, Miq. l. c. 533 = P. Botrytis.
brachypoda, Benth. Bot. Voy. Sulph. 167. t. 54 = P. brachypodon.
brachystachya, Miq. ex C. DC. in DC. Prod. xvi. 1. 296 = P. brachystachyum.
brasiliensis, Miq. in Linnaea, xx. (1847) 140.—Bras.
Bredemeyeri, Miq. Syst. Pip. 429 = P. Bredemeyeri.
brevispicata, Miq. l. c. 508 = P. brevispicatum.
Brongniartii, Miq. l. c. 486 = P. Brongniartii.
Brunoniana, Miq. l. c. 509 = P. lanceolatum.
caladiifolia, Miq. l. c. 387 = P. caladiifolium.
caldasiana, Miq. in Verh. Kon. Akad. Wetensch. II. v. (1871) 237; et in Arch. Néerl. vi. (1871) 175.—Bras.
calocoma, Miq. in Seem. Bot. Voy. Herald, 199 = P. calocoma.
Cambessedei, Miq. in Linnaca, xx. (1847) 161 = P. Cambessedesii.
caracasana, Miq. Syst. Pip. 398 = P. caracasanum.
cardiophylla, Presl, Epim. Bot. 225 = P. cardiophyllum.
caripensis, Miq. Syst. Pip. 522 = P. caripense.
casapiensis, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 463 = P. casapiense.
cassinoides, Miq. Syst. Pip. 490 = P. cassinoides.
catalpaefolia, Miq. in Versl. en Medel. Kon. Nederl. Inst. '1842' (1843) 197 = P. marginatum.
caudata, Miq. l. c. 196 = P. marginatum.
cearensis, Miq. Syst. Pip. 456 = P. angustifolium.
celtidifolia, Miq. l. c. 452 = P. celtidifolium.
cernua, Presl, Epim. Bot. 225 = P. cernuum.
Chamissonis, Miq. Syst. Pip. 457 = P. Chamissonis.
chavicoides, Miq. in Linnaea, xx. (1847) 135 = P. chavicoides.
chelseana, Miq. Syst. Pip. 533 = P. tomentosum.
chimonanthifolia, Miq. l. c. 466 = P. chimonanthifolium.
Churumayu, Miq. l. c. 481 = P. Churumayu.
cinnamomifolia, Miq. l. c. 419.—Peruv
Clausseniana, Miq. in Linnaea, xx. (1847) 153 = P. Claussenianum.
coccoloboides, Miq. Syst. Pip. 397 = P. coccoloboides.
colubrina, Miq. l. c. 512 = P. colubrinum.
concinna, Miq. in Linnaea, xx. (1847) 151 = P. concinnum.
consanguinea, Miq. Syst. Pip. 528 = P. consanguineum.
controversa, Miq. l. c. 448 = P. hispidum.
cornifolia, Miq. l. c. 479 = P. cornifolium.
coruscans, Miq. l. c. 411 = P. coruscans.
corylifolia, Klotzsch, in Hook. Journ. Bot. iv. (1842) 321 = P. dilatatum.
corylifolia, Miq. Syst. Pip. 474 = P. dilatatum.
crassispicata, Miq. l. c. 521 = P. crassispicatum.
crocata, Miq. l. c. 429 = P. crocatum.
cumanensis, Miq. l. c. 484 = P. cumanense.
curvata, Miq. l. c. 522 = P. curvatum.
cuspidata, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 468 = P. pseudo-peruvianum.
dasyoura, Miq. l. c. 452 = P. dasyourum.
dasypoda, Miq. l. c. 458 = P. dasypodum.
decipiens, Miq. Syst. Pip. 462 = P. decipiens.
decurrens, Miq. l. c. 402 = P. tuberculatum.
decurrens, Fourn. in Illustr. Hortic. xxiii. (1876) t. 239 = P. decurrens.
demerarana, Miq. in Hook. Lond. Journ. Bot iv. (1845) 464 = P. demeraranum.
denudans, Miq. in Linnaea, xx. (1847) 172 = P. denudans.
dimidiata, Presl, Epim. Bot. 225 = P. dimidiatum.
diospyrifolia, Miq. Syst. Pip. 477 = P. diospyrifolium.
divergens, Presl, Epim. Bot. 228 = P. divergens.
dubia, Miq. Syst. Pip. 535 = P. chilense.

ARTANTHE :—
Durvilleana, Miq. Syst. Pip. 507 = P. anonaefolium.
elongata, Miq. l. c. 434 = P. angustifolium.
enckeoides, Miq. l. c. 412 = P. crassinervium.
Endlicheriana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 447 = P. formosum.
eriopoda, Miq. l. c. 459 = P. eriopodon.
eucalyptifolia, Miq. Syst. Pip. 505 = P. eucalyptifolium.
excavata, Miq. l. c. 534 = P. excavatum.
eximia, Miq. l. c. 393 = P. obliquum.
exserens, Miq. in Linnaea, xx. (1847) 148 = P. crassinervium.
flagellaris, Miq. Syst. Pip. 393 = P. nobile.
flexicaulis, Miq. l. c. 533 = P. flexicaule.
flexuosa, Miq. l. c. 454 = P. flexuosum.
formosa, Miq. l. c. 386 = P. formosum.
fuliginea, Miq. l. c. 455 = P. fuligineum.
fuscescens, Miq. in Linnaea, xx. (1847) 144 = P. estrellense.
Gabrieliana, Miq. Syst. Pip. 476 = P. Gabrielanum.
Galeottii, Miq. l. c. 451 = P. aduncum.
Gaudichaudiana, Miq. l. c. 469 = P. Gaudichaudianum.
geniculata, Griseb. Cat. Pl. Cub. 65 = P. rigidum.
geniculata, Miq. in Versl. en Medel. Kon. Nederl. Inst. '1842' (1843) 194 = P. geniculatum.
glabella, Miq. Syst. Pip. 518 = P. glabellum.
glabrata, Miq. l. c. 481 = P. cumanense.
glabrescens, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 461 = P. glabrescens.
gracilis, Miq. Syst. Pip. 448 = P. gracile.
grandifolia, Miq. l. c. 408 = P. grandifolium.
granulosa, Miq. l. c. 435 = P. angustifolium.
guianensis, Klotzsch, in Hook. Journ. Bot. iv. (1842) 322 = P. guianense.
Guilleminiana, Miq. Syst. Pip. 409 = P. crassinervium.
Haenkeana, Miq. in Linnaea, xx. (1847) 145 = P. Paulianum.
halesiaefolia, Miq. Syst. Pip. 419 = P. halesiaefolium.
Hartwegiana, Benth. Pl. Hartw. 254 = P. Hartwegianum.
heterophylla, Miq. Syst. Pip. 502 = P. heterophyllum.
hexagyna, Miq. l. c. 412 = P. Betle.
Hilariana, Miq. l. c. 421 = P. Hilarianum.
hirsuta, Miq. l. c. 446 = P. hirsutum.
hirtula, Presl, Epim. Bot. 226 = P. hirtulum.
hispida, Miq. Comm. Phyt. 50. t. viii. f. A = P. asperifolia.
Hookeriana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 446 = P. Hookerianum.
Hostmanniana, Miq. l. c. 465 = P. Hostmannianum.
hymenophylla, Miq. Syst. Pip. 532 = P. Kegelii.
imperialis, Miq. in Seem. Bot. Voy. Herald, 198. t. 38 = P. imperiale.
insignis, Miq. Syst. Pip. 394 = P. insigne.
Jacquemontiana, Miq. l. c. 463 = P. citrifolium.
jalapensis, Miq. l. c. 444 = P. jalapense.
jamaicensis, Griseb. Fl. Brit. W. Ind. 170 = P. geniculatum.
japurensis, Miq. in Linnaea, xx. (1847) 168 = P. japurense.
javitensis, Miq. Syst. Pip. 401 = P. javitense.
Jussiacana, Miq. l. c. 420 = P. Jussiaeanum.
Karwinskiana, Miq. l. c. 476 = P. Karwinskianum.
Kegeliana, Miq. in Linnaea, xxii. (1849) 77 = P. Kegelianum.
Kunthiana, Miq. Syst. Pip. 453 = P. hirsutum.
lacunosa, Miq. l. c. 427 = P. lacunosum.
lanceaefolia, Miq. l. c. 433 = P. lanceaefolium.
lanceata, Miq. l. c. 526 = P. lancifolium.
lapathifolium, Miq. l. c. 401 = P. lapathifolium.
laurifolia, Miq. l. c. 520 = P. laurifolium.
Lehmanniana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 466 = P. salicariaefolium.
lentaginoides, Miq. Syst. Pip. 520 = P. rivinoides.
Leprieurii, Miq. l. c. 525.—Guiana.
leptura, Miq. l. c. 523 = P. lepturum.
Lessertiana, Miq. l. c. 405 = P. lapathifolium.
leucophylla, Miq. l. c. 460 = P. verbascifolium.
Lhotzkyana, Miq. l. c. 417 = P. Lhotzkyanum.
Lindeniana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 453 = P. Artanthe.
lineata, Miq. Syst. Pip. 459 = P. lineatum.
linguifolia, Miq. l. c. 533 = P. linguaefolium.
Lucaeana, Miq. l. c. 523 = P. Lucaeanum.
Luschnathiana, Miq. l. c. 494 = P. geniculatum.

ARTANTHE :—
macrophylla, Griseb. Fl. Brit. W. Ind. 171 = P. geniculatum.
macrophylla, Miq. Syst. Pip. 423 = P. macrophyllum.
macrorhyncha, Miq. in Linnaea, xx. (1847) 173 = P. macrorhynchon.
macrostachya, Presl, Epim. Bot. 227 = P. macrostachyum.
magnifica, Linden, ex Fourn. in Illustr. Hortic. xxiii. (1876) t. 239 = P. decurrens.
magnifica, Miq. Syst. Pip. 391 = P. obliquum.
malacophylla, Presl, Epim. Bot. 227 = P. malacophyllum.
marginata, Miq. Syst. Pip. 381 = P. marginatum.
Martiana, Miq. l. c. 422 = P. Martianum.
Martinicae, Miq. l. c. 413.—Ins. Martinica.
Maximiliana, Miq. in Linnaea, xx. (1847) 169 = P. juliflorum.
melanosticta, Miq. Syst. Pip. 404 = P. melanostictum.
mexicana, Miq. l. c. 458 = P. mexicanum.
Meyeniana, Klotzsch, ex Miq. l. c. 524 = P. lepturum.
Meyeri, Miq. l. c. 454 = P. divaricatum.
microstachya, Miq. l. c. 527 = P. microstachyon.
Miersiana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 449 = P. Miersinum.
Mikaniana, Miq. Syst. Pip. 383 = P. Parthenium.
mitis, Miq. l. c. 407 = P. mite.
modesta, Miq. l. c. 517 = P. modestum.
mollicoma, Miq. l. c. 438 = P. mollicomum.
mollis, Miq. in Linnaea, xx. (1847) 165 = P. mollipilosum.
Moritziana, Miq. l. c. xviii. (1844) 714 = P. Moritzianum.
nitida, Miq. Comm. Phyt. 47. t. vii = P. nitidum.
nitidula, Miq. Syst. Pip. 534 = P. nitidulum.
nubigena, Miq. l. c. 432 = P. crocatum.
obesa, Miq. l. c. 470 = P. scabrum.
oblonga, Miq. l. c. 480 = P. oblongum.
oblongata, Miq. l. c. 535 = P. oblongatum.
oblongifolia, Klotzsch, in Hook. Journ. Bot. iv. (1842) 322 = P. oblongifolium.
obovata, Miq. Syst. Pip. 508 = P. colubrinum.
obtusa, Miq. l. c. 416 = P. fluminense.
obumbrata, Miq. in Linnaea, xx. (1847) 142 = P. obumbratum.
Olfersiana, Klotzsch, in Hook. Journ. Bot. iv. (1842) 321 = P. hirsutum.
Opizii, Miq. Syst. Pip. 440 = P. asperifolium.
opulifolia, Miq. in Linnaea, xx. (1847) 160 = P. Cambessedesii.
paraënsis, Miq. l. c. 156 = P. paraënse.
Parkeriana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 46 = P. Parkerianum.
patens, Miq. Syst. Pip. 534 = P. nervosum.
patula, Miq. l. c. 554 = P. patens.
Pavoni, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 450 = P. Pavoni.
pedunculariis, Miq. Syst. Pip. 531 = P. pedunculare.
persicariaefolia, Miq. l. c. 499 = P. persicariaefolium.
peruviana, Miq. l. c. 482 = P. pseudo-peruvianum.
phthinotricha, Miq. l. c. 527 = P. phthinotrichon.
phytolaccaefolia, Miq. l. c. 534 = P. phytolaccaefolium.
pilosiuscula, Miq. l. c. 476 = P. pilosiusculum.
plantaginea, Miq. l. c. 389 = P. plantagineum.
Poeppigii, Miq. l. c. 437 = P. pseudo-mollicomum.
polyneura, Miq. l. c. 504 = P. indecorum.
populifolia, Miq. l. c. 418 = P. populifolium.
pothifolia, Miq. l. c. 399 = P. pothifolium.
pothomorpha, Miq. l. c. 403 = P. Pothomorphe.
prunifolia, Miq. l. c. 423 = P. prunifolium.
pseudo-churumayu, Miq. l. c. 407 = P. pseudo-churumayu.
psilostachya, Miq. l. c. 448 = P. psilostachyum.
punctata, Miq. l. c. 522 = P. punctatum.
pyrifolia, Miq. in Linnaea, xx. (1847) 159 = P. Mastersianum.
Radula, Miq. l. c. 426 = P. Bredemeyeri.
ramiflora, Miq. ex C. DC. in DC. Prod. xvi. 1. 288 = P. Hostmannianum.
Regnellii, Miq. in Linnaea, xxii. (1849) 574 = P. Regnellii.
renitens, Miq. in Mart. Fl. Bras. iv. 1. 221.—Am. trop.
reticulosa, Miq. Syst. Pip. 554 = P. reticulosum.
rhododendrifolia, Miq. l. c. 506 = P. anonaefolium.
rhynchostachya, Miq. l. c. 526 = P. rhynchostachyum.
richardiaefolia, Miq. l. c. 395 = P. richardiaefolium.
riparia, Miq. l. c. 409 = P. grandifolium.
rivinoides, Miq. l. c. 519 = P. rivinoides.

ARTANTHE :—
Rudgeana, Miq. in Linnaea, xxii. (1849) 77 = P. Rudgeanum.
rudis, Miq. Syst. Pip. 467 = P. rude.
rugosa, Miq. l. c. 488 = P. Rohrii.
Ruiziana, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 444 = P. pangoënse.
rumicifolia, Miq. in Linnaea, xx. (1847) 141 = P. rumicifolium.
salicariaefolia, Miq. Syst. Pip. 463 = P. salicariaefolium.
salicifolia, Miq. l. c. 533 = P. salicifolium.
saliunca, Miq. l. c. 528 = P. salicinum.
salviaefolia, Miq. l. c. 430 = P. salviaefolium.
sancta, Miq. in Linnaea, xviii. (1844) 714 = P. sanctum.
saururoides, Miq. Syst. Pip. 414 = P. saururoides.
scabra, Miq. l. c. 447 = P. hirsutum.
scabrata, Presl, Epim. Bot. 226.—Mexico.
scabrifolia, Miq. Syst. Pip. 461 = P. scabrifolium.
scabriuscula, Miq. l. c. 447 = P. scabriusculum.
Schachii, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 448 = P. Schachii.
Schlechtendalii, Miq. Syst. Pip. 386 = P. reticulatum.
Schomburgkii, Klotzsch, in Hook. Journ. Bot. iv. (1842) 322.—Guiana.
Schottii, Miq. in Linnaea, xx. (1847) 157 = P. Schottii.
scopulorum, Miq. Syst. Pip. 396 = P. scopulorum.
scutata, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 445 = P. subscutatum.
secunda, Miq. Syst. Pip. 516 = P. secundum.
Seemanniana, Miq. in Seem. Bot. Voy. Herald, 199. t. 39 = P. auritum.
Sellowiana, Miq. Syst. Pip. 416 = P. Sellovianum.
septuplinervia, Miq. in Seem. Bot. Voy. Herald, 199. t. 40 = P. septuplinervium.
sororia, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 443 = P. sororium.
spectabilis, Miq. in Linnaea, xx. (1847) 138 = P. cernuum.
staminea, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 461 = P. geniculatum.
stellipila, Miq. in Linnaea, xx. (1847) 163 = P. stellipilum.
stupposa, Miq. Syst. Pip. 430 = P. crocatum.
subaequalis, Presl, Epim. Bot. 226 = P. subaequale.
subauriculata, Miq. in Linnaea, xx. (1847) 150 = P. subauriculatum.
superba, Miq. l. c. 137 = P. obliquum.
Swartziana, Miq. Syst. Pip. 490 = P. Swartzianum.
tectoniaefolium, Miq. l. c. 396 = P. tectoniaefolium.
tejucensis, Miq. l. c. 496 = P. tejucense.
tenuicuspis, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 457 = P. tenuicuspe.
tiliaefolia, Miq. Syst. Pip. 388 = P. tiliaefolium.
trichoneura, Miq. l. c. 483.—Guiana.
trichostachya, Miq. l. c. 488 = P. citrifolium.
tricuspis, Miq. ex Seem. Bot. Voy. Herald, 200. t. 42.—Panama.
trinervis, Miq. l. c. 200. t. 41 = P. trinerve.
tuberculata, Miq. Syst. Pip. 497 = P. tuberculatum.
tumida, Miq. l. c. 465 = P. tumidum.
ulmifolia, Miq. l. c. 472 = P. dilatatum.
umbrosa, Miq. l. c. 470 = P. umbrosum.
variegata, Miq. l. c. 485 = P. variegatum.
Velloziana, Miq. l. c. 436 = P. aduncum.
velutina, Miq. l. c. 471 = P. velutinum.
verbascifolia, Miq. l. c. 431 = P. verbascifolium.
verrucosa, Griseb. Fl. Brit. W. Ind. 171 = P. geniculatum.
Warakabacoura, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 469 = P. Warakaboura.
Wydleriana, Miq. Syst. Pip. 487 = P. cuspidatum.
xestophylla, Miq. l. c. 491 = P. geniculatum.
xylopioides, Miq. l. c. 492 = P. geniculatum.
xylosteoides, Miq. l. c. 422 = P. xylosteoides.

ARTEDIA, Linn. Gen. ed. I. 71 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 927.
muricata, Linn. Sp. Pl. 242 = Daucus muricatus.
squamata, *Linn. l. c.*—Orient.
squamata, Pall. in Nov. Act. Petrop. x. (1797) 308 = Sanicula crithmifolia.

ARTEMISIA, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 435.
ABROTANUM, [Tourn.] Neck. Elem. i. 98 (1790).
ABSINTHIUM, Tourn. ex Linn. Syst. ed. I (1735).

ARTEMISIA:—
Oligosporus, Cass. in Bull. Soc. Philom. (1817) 33.
Picrothamnus, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 417.
abrotanifolium, Salisb. Prod. 191 = A. Abrotanum.
abrotanifolium, *Siev. in Pall. N. Nord. Beitr.* vii. 309. —Rossia.
abrotanoides, Jacq. ex DC. Prod. vi. 106 = vestita.
abrotanoides, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 399 = californica.
Abrotanum, Habl. Taur. 128 = maritima.
Abrotanum, *Linn. Sp. Pl.* 845.—Europ.; As. temp.
Abrotanum, Savi, Mat. Med. t. 38 = camphorata.
Abrotanum, S. G. Gmel. It. i. 21 = procera.
absinthia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Absinthium.
Absinthii, Heyne, ex Hook. f. Fl. Brit. Ind. iii. 329 = pallens.
absinthioides, Poir. Encyc. Suppl. i. 461 = Siversiana.
Absinthium, *Linn. Sp. Pl.* 848.—Europ.
abyssinica, *Sch. Bip. in A. Rich. Tent. Fl. Abyss.* i. 420. —Afr. trop.
acetica, Jacquem. ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 86 = maritima.
achilleaefolia, Tenore, Fl. Nap. 190 = camphorata.
achilleoides, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 195 = Tanacetum achilleoides.
Adamsii, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 27.—As. bor. or.
aemula, Steph. ex Bess. l. c. 22 = rutaefolia.
aethiopica, Linn. Sp. Pl. 845 = A. Herba-alba.
affinis, Hassk. in Flora, xxv. (1842) ii. Beibl. 22 = vulgaris.
affinis, *Steud. Nom.* ed. II. i. 136.—Sibir.
afra, *Jacq. Hort. Schoenb.* iv. 34. t. 467.—Afr. austr.
alba, Habl. Taur. 144 = austriaca.
alba, Pall. Reise, i. 502 = maritima.
alba, Turra, Fl. Ital. Prod. 67 = camphorata.
albida, Ledeb. Fl. Alt. iv. 85 = maritima.
albida, Willd. ex Ledeb. Fl. Ross. ii. 566 = inodora.
Allionii, Nym. Consp. 377 = nana.
alpina, Pall. ex Willd. Sp. Pl. iii. 1824 = lanata.
alpina, DC. ex Steud. Nom. ed. II. i. 136 = campestris.
alpina, Sch. Bip. ex Willk. in Flora, xxxiv. (1851) 746 = lanata.
altaica, Desf. Cat. Hort. Paris. (1829) 160 = pontica.
altissima, Ehrh. ex DC. Prod. vi. 108 = procera.
ambigua, Linn. Sp. Pl. ed. II. 1190 = Trichogyne laricifolia.
ambigua, Thunb. Prod. Pl. Cap. 147 = Scriphium vermiculatum.
Ammaniana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 60.—Sibir.
amorpha, *Rafin. Fl. Ludov.* 69.—Louisiana.
amygdalina, *Decne. in Jacquem. Voy. Bot.* 92. t. 100.— Reg. Himal.
andicola, Don, ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 325 = mendozana.
androsacea, Seem. Bot. Voy. Herald, 34. t. 6 = senjavinensis.
anethifolia, Fisch. ex DC. Prod. vi. 108 = procera.
anethifolia, *Weber, ex Stechm. Artem.* 29.—Sibir.
anglica, Hort. ex DC. Prod. vi. 103 = maritima.
angustifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 450 = A. Abrotanum.
annua, *Linn. Sp. Pl.* 847.—Oriens; As. bor.
annua, Pall. Reise, ii. 523 = sacrorum.
anomala, *S. Moore, in Journ. Bot.* xiii. (1875) 227.— China.
anthriscifolia, *Presl, ex Steud. Nom.* ed. II. i. 136.— Hab ?
apetala, Hort. Pest. ex Steud. l. c. = vulgaris.
apiacea, *Hance, in Walp. Ann.* ii. 895.—China.
aprica, Gebl. ex DC. Prod. vi. 125 = frigida.
aquatica, *Lour. Fl. Cochinch.* ii. 490.—China.
aragonensis, Lam. Encyc. ii. 269 = A. Herba-alba.
arborescens, *Linn. Sp. Pl.* ed. II. 1188.—Reg. Mediterr.
arbuscula, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 398.—Am. bor. occ.
arctica, Bess. ex Hook. Fl. Bor. Am. i. 323 = Richardsoniana.
arctica, Less. in Linnaea, vi. (1831) 213 = norvegica.
arenaria, *DC. Prod.* vi. 94.—Reg. Caucas.
argentea, Buch, Beschr. Canar. 148, 164 = canariensis.
argentea, *L'Hérit. Sert. Angl.* 22.—Sicil.; Madera.
argentea, Seb. & Maur. ex Willk. & Lange, Prod. Fl. Hisp. ii. 68 = arborescens.
argyrophylla, *Ledeb. Fl. Alt.* iv. 166.—Sibir.

ARTEMISIA:—
armeniaca, *Lam. Encyc.* i. 263.—Armenia.
armeniaca, Willd. ex Ledeb. Fl. Ross. ii. 592 = Tournefortiana.
Assoana, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 69 = lanata.
arragonensis, Auct. vide *aragonensis*.
Athanasia, Bess. in Bull. Soc. Nat. Mosc. vii. (1834) 24 = Tanacetum fruticulosum.
atlantica, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ix. (1862) 298.—Algeria.
atrata, Lam. Encyc. i. 253 = spicata.
atropurpurea, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. (1867) i. 231 = arctica.
Aucheri, *Boiss. Fl. Orient.* iii. 367.—Persia.
australis, Ehrh. ex DC. Prod. vi. 120 = biennis.
australis, *Less. in Linnaea*, vi. (1831) 522.—Ins. Sandvic.
austriaca, *Jacq. Fl. Austr.* i. 61.—Europ. or.; Oriens; As. bor.
baicalensis, Willd. ex DC. Prod. vi. 107 = sacrorum.
Balsamita, Willd. Enum. Hort. Berol. Suppl. 57 = pontica.
bargusinensis, Spreng. Syst. iii. 493 = borealis.
Barrelieri, *Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 87. —Hispan.
basilica, Webb, ex Willk. & Lange, Prod. Fl. Hisp. ii. 70 = granatensis.
Baumgarteni, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 73.—Europ. austr. or.
Besseriana, Ledeb. Fl. Ross. ii. 590 = lagocephala.
Biasolettiana, Visiani, Ind. Sem. Hort. Patav. (1836) = camphorata.
biennis, *Willd. Phyt.* 11. n. 39.—Am. bor.; Sibir.; Reg. Himal.
biflora, *Spreng. Nachr.* i. *Bot. Gart. Halle*, 12.—Hab.?
Bigelovii, *A. Gray, in Pacif. Rail. Rep.* iv. 110.—Am. bor.
Billardieriana, Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 81 = A. Herba-alba.
blepharolepis, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 340 (*Lehm. Reliq. Bot.* 164).—As. centr.
Bocconi, All. Fl. Pedem. i. 169 = nana, spicata.
Bolanderi, *A. Gray, in Proc. Am. Acad.* xix. (1883) 49.—Calif.
borealis, *Pall. Reise*, iii. 755.—Am. bor. et arct.; Sibir.
Boschniakiana, DC. Prod. vi. 104 = maritima.
brachyloba, *Franch. Pl. Davidian.* i. 171.—Mongol.
brevifolia, Wall. Cat. n. 3306 = maritima.
caerulescens, *Linn. Sp. Pl.* 848.—Reg. Mediterr.
caespitosa, Bess. in Hook. Fl. Bor. Am. i. 324 = Richardsoniana.
caespitosa, *Ledeb. Fl. Alt.* iv. 80.—Sibir. alt.
californica, *Less. in Linnaea*, vi. (1831) 523.—Calif.
Campbellii, *Hook. f. & Thoms. ex Clarke, Comp. Ind.* 164.—Reg. Himal.
campestris, Auct. Hisp. ex Willk. & Lange, Prod. Fl. Hisp. ii. 72 = glutinosa.
campestris, Ledeb. Fl. Alt. iv. 86 = commutata.
campestris, *Linn. Sp. Pl.* 846.—Europ.; As. bor.; Oriens; Ind. or.
campestris, Pursh, Fl. Am. Sept. ii. 522 = canadensis?
campestris, Scop. ex Steud. Nom. ed. II. i. 136 = camphorata.
campestris, Turcz. ex DC. Prod. vi. 96 = Ledebouriana.
camphorata, *Vill. Presp.* 31.—Europ. austr.
camtschatica, Schlecht. ex Ledeb. Fl. Ross. ii. 568 = borealis.
cana, *Pursh, Fl. Am. Sept.* 621.—Am. bor.
canadensis, *Michx. Fl. Bor. Am.* ii. 129.—Am. bor.; As. bor. or.
canariensis, *Less. in Linnaea*, vi. (1831) 217.—Ins. Canar.
canescens, Willd. Sp. Pl. iii. 1843 = armeniaca.
cannabina, Jacquem. ex Bess. in DC. Prod. vi. 97 = desertorum.
capillaris, Miq. in Ann. Mus. Bot. Lugd. Bat. ii. 175 = scoparia.
capillaris, *Thunb. Fl. Jap.* 309.—Japon.
capillifolia, Fisch. ex DC. Prod. vi. 122 = subviscosa.
capillifolia, Lam. Encyc. i. 266 = Eupatorium foeniculaceum.
carnifolia, Buch.-Ham. ex Roxb. Fl. Ind. iii. 422 = seq.
caruifolia, *Roxb. Hort. Beng.* 61.—Ind. or.; China.
carvifolia, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 81 = caruifolia.
caucasica, *Willd. Sp. Pl.* iii. 1823.—Reg. Caucas.
caudata, *Michx. Fl. Bor. Am.* ii. 129.—Am. bor.

ARTEMISIA:—
centiflora, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 493.—Mongolia.
cephalantha, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xl. (1867) i. 227 = glomerata.
cernua, Dufour, ex Willk. & Lange, Prod. Fl. Hisp. ii. 71 = hispanica.
cernua, Nutt. Gen. Am. ii. 143 = dracunculoides.
cernuiflora, Dufour, ex Willk. & Lange, Prod. Fl. Hisp. ii. 71 = hispanica.
chamaemelifolia, *Vill. Prosp.* 32.—Gall.; Reg. Caucas.; Persia.
Chamissoniana, Bess. ex Hook. Fl. Bor. Am. i. 324 = norvegica.
chamomillifolia, Parrot, in Engelh. & Parrot, Reise Krym. Kauk. ii. 131 = chamaemelifolia?
Chiajeana, *Kunze, ex Rich. Bot. Méd.* ii. 612; *ex Steud. Nom.* ed. II. i. 137.—Oriens?
chinensis, Bess. in Linnaea, xv. (1841) 92 = lagocephala.
chinensis, Burm. ex DC. Prod. vi. 125 = mollis.
chinensis, *Linn. Sp. Pl.* 849.—China; Sibir.
chinensis, Pursh, Fl. Am. Sept. ii. 521 = Stelleriana.
chrysantha, Fisch. ex DC. Prod. vi. 125 = frigida.
chrysantha, Steph. ex Ledeb. Fl. Ross. ii. 597 = rupestris.
Chrysanthemum, Russ. ex DC. Prod. vi. 140 = Sphaeromorphaea Russeliana.
Cina, *Berg, Darstell.* iv. t. 29c.—Oriens.
cinerea, Hort. Par. ex DC. Prod. vi. 122 = camphorata.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 55.—Algeria.
coarctata, J. H. Fors. in Linn. Inst. Skrift. (reimpr. 12) = vulgaris.
coarctata, Turcz. ex Ledeb. Fl. Ross. ii. 567 = campestris.
coerulescens, Pall. Reise, ii. 285 = sericea.
coerulescens, Presc. ex Ledeb. Fl. Ross. ii. 564 = Ammaniana.
collina, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 39 = campestris.
columbiensis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 398 = cana.
Columnae, Tenore, Ind. Sem. Hort. Neap. (1829) 14 = camphorata.
commutata, Bess. in Bull. Soc. Nat. Mosc. viii. (1835) 70 = campestris.
compacta, Fisch. ex DC. Prod. vi. 102 = maritima.
Contra, Linn. Mant. 282 = Sieberi.
Contra, Willd. ex Spreng. Syst. iii. 494 = maritima.
controversa, Jan, ex DC. Prod. vi. 126 = Siversiana.
Copa, *Phil. Fl. Atac.* 33.—Chili.
corymbosa, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 74 = glomerata.
corymbosa, Lam. Encyc. i. 265 = camphorata.
crithmifolia, *Linn. Sp. Pl.* 846.—Lusit.; Gall.
crithmifolia, Sievers, in Pall. N. Nord. Beitr. vii. 333 = glabella.
cuneifolia, DC. Prod. vi. 126 = parviflora.
cuneifolia, Schecle, in Linnaea, xxii. (1849) 162 = mexicana?
curilensis, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 67 = Meyeriana.
curilensis, Spreng. Syst. iii. 489 = leontopodioides.
Cyna, Pall. Reise, iii. 630 = maritima.
Czekanowskiana, *Trautv. in Act. Hort. Petrop.* v. (1877) 72.—Sibir.
davurica, *Vitm. Summa*, v. 12.—Dahuria.
Decaisnei, *F. W. Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 329.—Ind. or.
Deliliana, Bess. Enum. Pl. Volh. Suppl. 89 = monosperma.
delphinensis, Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 19 = Mutellina.
densiflora, Viv. App. Alt. Cors. ii. 4. t. 2 = gallica.
dentata, Willd. Sp. Pl. iii. 1826 = rupestris.
depressa, Turcz. ex DC. Prod. vi. 107 = Adamsii.
desertorum, Bess. ex Hook. Fl. Bor. Am. i. 325 = canadensis.
desertorum, *Spreng. Syst.* iii. 490.—As. centr.; Sibir.; Reg. Himal.
discolor, *Dougl. ex DC. Prod.* vi. 109.—Am. bor. occ.
discolor, Torr. & Gray, in Pacif. Rail. Rep. ii. 126 = franserioides.
Douglasiana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 60 = ludoviciana.
dracunculoides, *Pursh, Fl. Am. Sept.* 742.—Am. bor.
Dracunculus, *Linn. Sp. Pl.* 849.—Europ. or.; Oriens; Reg. Himal.

ARTEMISIA:—
Dracunculus, Pursh, Fl. Am. Sept. ii. 521 = draeunculoides.
dubia, Wall. Cat. n. 3307 = vulgaris.
Eehegarayi, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 42.—Reg. Argent.
elegans, Fisch. ex Ledeb. Fl. Ross. ii. 579 = proeera.
elegans, Jan, ex Bess. in Bull. Soc. Nat. Mose. ix. (1836) 104 = glaeialis.
elegans, Roxb. Hort. Beng. 61; Fl. Ind. iii. 421 = seoparia.
elegans, Salisb. Prod. 191 = arboreseens.
eranthema, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 339 (*Lehm. Reliq. Bot.* 163).—Turkest.
eriantha, Tenore, Ind. Sem. Hort. Neap. (1830) 14 = Baumgarteni.
erioearpa, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 336 (*Lehm. Reliq. Bot.* 160). — Persia; Turkestan.
eriophora, Schleeht. ex Ledeb. Fl. Ross. ii. 586 = vulgaris.
eriopoda, *Bunge, Enum. Chin. Bor.* 37.—Chin. bor.
Eschscholtziana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 24.—Ins. Sandvie.
exilis, Fisch. ex DC. Prod. vi. 119 = annua.
Falconeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 328.—Tibet. oee.
fasciculata, *Bieb. Fl. Taur. Cauc.* ii. 293.—Reg. Caueas.; Persia.
fasciculata, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. iii. 318 = Tanacetum artemisioides.
filaginoidea, Steehm. Artem. 29 = vallesiaea.
filifolia, *Torr. in Ann. Lyc. N. York*, ii. (1828) 211.—Am. bor.
Fischeriana, Bess. in Nouv. Mém. Soc. Nat. Mose. iii. (1834) 21 = ealifornica.
foeniculacea, Stev. ex DC. Prod. vi. 108 = proeera.
foetida, Jaequem. ex Bess. in Bull. Soc. Nat. Mose. ix. (1836) 96 = desertorum.
foliosa, Nutt. in Trans. Am. Phil. Soe. Ser. II. vii. (1841) 397 = ealifornica.
Fontanesiana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 52.—Hab.?
fragrans, Eiehw. Casp. Caue. 6 = Sieberi.
fragrans, Willd. Sp. Pl. iii. 1835 = maritima.
franserioides, *Greene, in Bull. Torrey Club*, x. (1883) 42.—Am. bor. oec.
frigida, Eiehw. Casp. Caue. 31 = austriaea.
frigida, *Willd. Sp. Pl.* iii. 1838.—Sibir.; Am. aret.
fruteseens, *Pall. Reise Statth. Russ. Reich.* i. 94, 177. —Rossia.
fruticans, Comm. ex DC. Prod. vi. 261 = Stoebe paniculata.
fruticosa, Willd. ex DC. l. e. 94 = arenaria.
furcata, Bieb. Fl. Taur. Cauc. iii. 567 = trifureata?
gallica, Schur, Enum. Pl. Transs. 324 = maritima.
gallica, *Willd. Sp. Pl.* iii. 1824.—Gall.; Hisp.
garganica, Tenore, ex DC. Prod. vi. 122 = camphorata.
Gayana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 35. —Hispan.
gelida, Ledeb. in Mém. Acad. Pétersb. v. (1805) 565 = borealis.
Genipi, Steehm. Artem. 17 = spieata.
gilveseens, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 175.—Japon.
glabella, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 441.—Soongaria.
glabra, Jaequem. ex Bess. in DC. Prod. vi. 97 = desertorum.
glabrata, DC. in Wight, Contrib. 20 = parviflora.
glabrescens, Stokes, Bot. Mat. Med. iv. 188 = campestris.
glacialis, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 70 = granatensis.
glacialis, Vitm. Sagg. 44 = lanata.
glacialis, Herb. Mosc. ex Bess. in Nouv. Mém. Soc. Nat. Mose. iii. (1834) 67 = Triniana.
glacialis, *Linn. Sp. Pl.* ed. II. 1187.—Europ. austr. med.
glacialis, Wull. in Jacq. Coll. ii. 118 = Mutellina.
glacialis, Schang. in Pall. N. Nord. Beitr. vi. 93 = commutata.
glauca, Bieb. ex Ledeb. Fl. Ross. ii. 564 = Ammaniana.
glauca, *Pall. ex Willd. Sp. Pl.* iii. 1831.—Reg. Himal.; Sibir.
globularia, Cham. ex Bess. in Nouv. Mém. Soc. Nat. Mose. iii (1834) 64 = norvegiea.
ARTEMISIA:—
globularia, Ledeb. ex Maxim. in Bull. Acad. Pétersb. xvii. (1872) 435 = glomerata.
glomerata, Hook. & Arn. Bot. Beech. Voy. 125 = senjavinensis.
glomerata, *Ledeb. in Mém. Acad. Pétersb.* v. (1805) 564.—Am. oec. et As. or. aret.
glomerata, Sieber, ex Spreng. Syst. iii. 489 = Sieberi.
glomerulata, *Sieber, ex Spreng. Syst.* iii. 489 = Sieberi.
glutinosa, Boiss. ex Walp. Rep. vi. 214 = variabilis.
glutinosa, *J. Gay, in DC. Prod.* vi. 95.—Gallia; Ital.
Gmelini, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 28 = saerorum.
Gmelini, *Stechm. Artem.* 30.—Sibir.
Gmelini, Willd. ex Bess. in Nouv. Mém. Soc. Nat. Mose. iii. (1834) 67 = Lagopus.
gnaphalioides, Bess. in DC. Prod. vi. 105 = mendozana.
gnaphalodes, Nutt. Gen. Am. ii. 143 = ludovieiana.
gorgonum, *Webb, in Hook. Niger Fl.* 142.—Ins. Cap. Verd.
gracilis, L'Hérit. ex DC. Prod. vi. 100 = seoparia.
granatensis, *Boiss. ex DC. l. c.* 298.—Hispan.
grandiflora, *DC. l. c.* 122.—Sibir.
grandiflora, Fisch. ex Herd. in Bull. Soc. Nat. Mose. xl. (1867) i. 217 = pontica.
grandiflora, Hoffm. Deutsehl. Fl. ed. II. ii. 136 = rupestris.
grata, Heyne, ex Steud. Nom. ed. II. i. 137 = Roxburghiana?
grata, Wall. Cat. n. 3294; Comp. n. 404 = Roxburghiana.
Griffithiana, Boiss. Diagn. Ser. II. iii. 31 = maeroeephala.
groenlandica, Wormsk. in Fl. Dan. t. 1585 = borealis.
Halimodendron, Ledeb. in Hook. f. Fl. Brit. Ind. iii. 321 = salsoloides.
Halodendron, *Turcz. in DC. Prod.* vi. 94.—Dahur.; Mongol.
Haussknechtii, *Boiss. Fl. Orient.* iii. 374.—Persia.
helvetica, Sehleieh. Cat. Pl. Helv. ed. III. 8 = nana.
hemisphaerica, Roxb. Hort. Beng. 61; Fl. Ind. iii. 422 = Cotula hemisphaeriea.
Herba-alba, *Asso, Syn. Arag.* 117.—Hispan.; Maroec.; Oriens.
herbacea, Clarke, ex DC. Prod. vi. 108 = proeera.
herbaeea, *Ehrh. in Willd. Sp. Pl.* iii. 1828.—Sibir.
herbacea, Hort. Wratisl. ex Bess. in Nouv. Mém. Soe. Nat. Mose. iii. (1834) 30 = proeera.
hermaphrodita, Kit. ex Ledeb. Fl. Ross. ii. 566 = campestris.
heterophylla, Bess. in Nouv. Mém. Soe. Nat. Mose. iii. (1834) 74 = Riehardsoniana?
heterophylla, Nutt. in Trans. Am. Phil. Soe. N. S. vii. (1841) 400 = vulgaris.
Heyneana, Wall. Cat. n. 3295 = vulgaris.
hirsuta, Rottl. in Spreng. Syst. iii. 490 = Cyathoeline lyrata.
hispanica, Jaeq. Coll. i. 98 = biennis.
hispaniea, *Lam. Encyc.* i. 263.—Hispan.
hispanica, Steehm. ex Bess. in Bull. Nat. Soe. Mose. ix. (1836) 114 = vulgaris.
hololeuea, *Bieb. ex Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 46.—Ukrania.
hololeuca, Boiss. & Haussk. in Boiss. Fl. Orient. iii. 374 = Haussknechtii.
holoserieea, Ledeb. Fl. Alt. iv. 65 = serieea.
Hookeriana, Bess. in Nouv. Mém. Nat. Soe. Mose. iii. (1834) 39 = ludovieiana.
humilis, Bieb. Casp. App. 83 = maritima.
humilis, Mill. Gard. Dict. ed. VIII. n. 7 = A. Abrotanum.
humilis, Wulf. in Jaeq. Coll. iv. 295 = eamphorata.
hypoleuca, Edgew. in Trans. Linn. Soc. xx. (1846) 71 = Roxburghiana.
hyrcana, Spreng. Syst. iii. 434 = annua.
iberica, *Boiss. ex Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 120.—Reg. Cauc.
igniaria, Maxim. Prim. Fl. Amur. 161 = vulgaris.
incanescens, Jord. in Ann. Soc. Agric. Lyon; ex Gren. & Godr. Fl. Fr. ii. 127 = camphorata.
incompta, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 400 = discolor.
inconspicua, Spreng. Nachr. i. Bot. Gart. Halle, 12 = biennis.
inculta, Delile, Fl. Egypte, 264 = A. Herba-alba.
ARTEMISIA:—
inculta, Sieber, ex DC. Prod. vi. 95 = monosperma.
indica, Willd. Sp. Pl. iii. 1846 = vulgaris.
infirma, S. G. Gmel. ex Ledeb. Fl. Ross. ii. 566 = eampestris.
inodora, Bieb. Fl. Taur. Caue. ii. 295 = eampestris.
inodora, Hook. & Arn. Bot. Beech. Voy. 150 = draeuneuloides.
inodora, Miil. Gard. Diet. ed. VIII. n. 16 = A. Absinthium.
inodora, Willd. Enum. Hort. Berol. 864 = A. Draeuneulus.
insipida, *Schur, Enum. Pl. Transs.* 322.—Transsylv.
insipida, *Vill. Prosp.* 32.—Gallia.
integrifolia, *Falk, Beitr.* ii. 239.—Reg. Casp.
integrifolia, Linn. Sp. Pl. 848 = vulgaris.
integrifolia, Pursh, Fl. Am. Sept. 520 = ludovieiana.
integrifolia, Riehards. App. Frankl. Journ. ed. II. 30 = Purshiana.
intermedia, Host, Fl. Austr. ii. 160 = eamphorata
intricata, *Franch. Pl. Davidian.* i. 170.—Mongol.
involucrata, Turez. ex Ledeb. Fl. Ross. ii. 598 = frigida.
italica, Bieb. ex Ledeb. l. e. 566 = eampestris.
italica, *Stechm. Artem.* 28.—Italia.
Jacquemontiana, Bess. in Bull. Soc. Nat. Mose. (1835) 59 = desertorum.
Jacquini, Raeusch. Nom. ed. III. 237 = biennis
japonica, *Thunb. Fl. Jap.* 310.—Japon.
japonica, F. Schmidt, Fl. Sachal. 149 = Keiskeana.
jenisea, Willd. ex Ledeb. Fl. Ross. ii. 598 = seq.
jeniscensis, Willd. ex Spreng. Syst. iii. 489 = frigida.
judaiea, *Linn. Fl. Palaest. in Amoen. Acad.* iv. 463; *et Mant.* 281.—Aegypt.; Arab.
judaica, Lour. Fl. Cochineh. 489 = Loureiri.
juneea, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 383.—Soongaria.
Jussieana, *J. Gay, ex Bess. l. c.* viii. (1835) 28.—Hispan.
kamtschatica, Willd. ex Bess. l. e. 67 = leontopodioides.
Keiskeana, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 176.—Japon.
Koelreuteriana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 16.—Hab.?
kohatiea, *Klatt, in Sitzb. Akad. Muench.* (1878) 91.—Ind. or.
Kotzebuensis, Bess. in Nouv. Mém. Soe. Nat. Mose. iii. (1834) 80 = Chrysanthemum bipinnatum.
Krascheninnikoviana, Bess. l. e. 41 = maerobotrys, latifolia.
Kruhsiana, Bess. l. e. 22 = lagoeephala.
kulbadiea, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 120.—Reg. Casp.
kurilensis, Walp. Rep. vi. 217 = A. Lagopus.
laeiniata, *Willd. Sp. Pl.* iii. 1843.—Reg. Himal.; Sibir.
lactiflora, Wall. ex DC. Prod. vi. 115 = vulgaris.
lagoeephala, *Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 22.—Sibir.
lagopina, St. Lag. in Ann. Soe. Bot. Lyon, vii. (1880) 119 = seq.
Lagopus, *Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 69.—Sibir.
lanata, Koeh, Syn. Fl. Germ. ed. II. 402 = nitida.
lanata, Lam. Eneye. i. 267 = variabilis.
lanata, *Willd. Sp. Pl.* iii. 1823.—Hisp.; Ital.; Taur.
lanuginosa, Poir. Eneye. Suppl. i. 462 = lanata.
laricifolia, S. G. Gmel. It. ii. 3 = maritima.
lasiocline, Boiss. Diagn. Ser. I. vi. 92 = splendens.
latifolia, Ledeb. in Mém. Acad. Petersb. v. (1805) 569 = laciniata.
lavandulaefolia, DC. Prod. vi. 110 = vulgaris.
lavandulifolia, Salisb. Prod. 191 = caerulescens.
Ledebouriana, *Bess. in Bull. Soc. Nat. Mosc.* viii. (1835) 37.—Sibir.
lednicensis, Rochel, ex Spreng. Pugill. ii. 77 = eampestris.
Lehmanniana, *Bunge, in Mem. Sav. Etr. Petersb.* vii. (1851) 340 (*Lehm. Reliq. Bot.* 164).—Afghan.; Turkest.
leontopodioides, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 63 = glomerata.
leptophylla, *D. Don, Prod. Fl. Nep.* 183 (sp. dubia).—Nepal.
leptostachya, DC. Prod. vi. 113 = vulgaris.
Lercheana, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 441 = maritima.

ARTEMISIA:—
Lercheana, Presc. ex Bess. in Bull. Soc. Nat. Mosc. viii. (1835) 33 = arenaria.
Lercheana, Weber, ex Stechm. Artem. 24 = fragrans.
Lerchii, Boeber, ex Georgi, Beschr. Russ. Reich. iii. IV. 1235 = maritima.
Lessingiana, *Bess. in Linnaea*, xv. (1841) 90.—Sibir.
leucanthemifolia, Desf. ex DC. Prod. vi. 98 = desertorum.
leucodes, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 106.—Soongor.
leucophylla, Turcz. ex C. B. Clarke, Comp. Ind. 162 = vulgaris.
Lewisii, Torr. & Gray, Fl. N. Am. ii. 417 (nom. dclend.).
Libanotis, Bieb. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 42 = macrobotrys.
Lindheimeriana, Scheele, in Linnaea, xxii. (1849) 163 = mexicana.
Lindleyana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 35.—Am. bor. occ.
linearis, Gueldenst. ex Ledeb. Fl. Ross. ii. 560 = salsoloides.
Lippii, Jan, ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 79 = A. Herba-alba.
lithophila, Turcz. ex DC. Prod. vi. 122 = lagocephala.
littoralis, Retz. Obs. v. 28 = Epaltes australis.
Lobelii, All. Fl. Pedem. i. 166 = camphorata.
lobulifolia, *Boiss. Fl. Orient.* iii. 364.—Affghan.
longepedunculata, Rudolphi, ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 77 = norvegica.
longifolia, *Nutt. Gen. Am.* ii. 142.—Am. bor. occ.
Loureiri, *Kostel. Allg. Med. Pharm. Fl.* ii. 669.—China.
ludoviciana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 38 = discolor.
ludoviciana, *Nutt. Gen. Am.* ii. 143.—Am. bor. occ.
macra, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 364.—Cataonia.
macrantha, Ledeb. in Mém. Acad. Pétersb. v. (1805) 573 = sacrorum.
macrobotrys, Ledeb. Fl. Alt. iv. 73 = laciniata.
macrocephala, *Jacquem. ex Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 28.—Affghan.; Tibet.; As. centr.
macrophylla, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 41 = laciniata.
macrorhiza, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 73.—Sibir.
maderaspatana, Linn. Sp. Pl. 849 = Grangea maderaspatana.
magellanica, *Sch. Bip. in Flora*, xxxviii. (1855) 116.—Reg. Magell.
maracandica, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 345 (Lehm. Rel. Bot. 169) = fragrans.
maritima, *Linn. Sp. Pl.* 846.—Europ.; Reg. Caucas.; Sibir.
maritima, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 74 = gallica.
maroccana, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 61.—Marocco.
Marschalliana, Spreng. Syst. iii. 496 = campestris.
matricarioides, Less. in Linnaea, vi. (1831) 210 = Matricaria discoidea.
megacephala, *Rupr. Sert. Tiansch.* 52.—As. centr.
melanolepis, *Boiss. Diagn.* Ser. I. xi. 27.—Persia.
mendozana, *DC. Prod.* vi. 105.—Am. austr.
Mertensiana, Wallr. Sched. Crit. 466. t. 4 = laciniata.
Messerschmidiana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 27 = sacrorum.
Messerschmidii, Weber, ex Stechm. Artem. 19 = maritima.
mexicana, *Willd. ex Spreng. Syst.* iii. 490.—Mexico.
Meyeriana, Bess. in Linnaea, xv. (1841) 96 = A. Lagopus.
Michauxiana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 71 = discolor.
Michauxii, Bess. ex Hook. Fl. Bor. Am. i. 324 = praec.
millefoliata, Bieb. ex DC. Prod. vi. 107 = sacrorum.
minima, Linn. Sp. Pl. 849 = Centipeda orbicularis.
minor, *Jacquem. ex Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 22.—Tibet.
mollis, *J. Gay, in DC. Prod.* vi. 125.—Hab.?
mollissima, *D. Don, Prod. Fl. Nep.* 182 (sp. ignot.).—Nepal.
molluccana, *Roxb. Hort. Beng.* [101]; *Fl. Ind.* iii. 417.—Ins. Molucc.

ARTEMISIA:—
mongolica, Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 53 = vulgaris.
monogyna, Waldst. & Kit. Pl. Rar. i. 177. t. 75 = fragrans.
monosperma, *Delile, Fl. Egypte*, 263.—Syria; Aegypt.; Arab.
monostachya, *Bunge, ex Maxim. Fl. Amur.* 482.—Mongolia.
monspeliensis, Nym. Consp. 377 = glutinosa.
montana, Schlecht. ex Ledeb. Fl. Ross. ii. 591 = Richardsoniana.
montevidensis, *Spreng. Syst.* iii. 487.—Reg. Argent.
monticola, C. Koch, in Linnaea, xxiv. (1851) 347 = caucasica.
Moorcroftiana, *Wall. Cat. Comp.* n. 406.—Himal. bor. occ.
Moxa, *DC. Prod.* vi. 121.—China.
multicaulis, Ledeb. Fl. Alt. iv. 60 = ancthifolia.
mutabilis, Salisb. Prod. 191 = A. Santonicum.
Mutellina, S. G. Gmel. It. i. 159 = splendens.
Mutellina, *Vill. Prosp.* 31.—Europ. med.
myriantha, Wall. Cat. n. 3297 = vulgaris.
nana, *Gaud. Fl. Helv.* v. 231.—Helvet.
naronitana, Visiani, ex DC. Prod. vi. 108 = procera.
neapolitana, Tenore, Prod. Fl. Nap. App. v. 28 = variabilis.
neglecta, Spreng. in Willd. Enum. Hort. Berol. ii. 860 = desertorum.
nepalensis, *Nees, in Flora*, xiv. (1831) 290.—Ind. or.
nilotica, Linn. Cent. i. 27; Sp. Pl. ed. II. 1188 = Cotula anthemoides.
nitens, Stev. ex Ledeb. Fl. Ross. ii. 595 = sericea.
nitida, *Bertol. Mant.* 53.—Italia.
nitrosa, *Weber, ex Stechm. Artem.* 24.—Sibir.
nivea, Redowsk. ex Willd. Enum. Hort. Berol. 863 = austriaca.
norica, Leyb. in Flora, xxxvii. (1854) 369 = nana.
norvegica, *Fries, Novit. Suec.* ed. I. (1817) 56.—Norveg.
nubigena, Wall. Cat. n. 3233 = Tanacetum nubigenum.
nutans, Fras. ex Pursh, Fl. Am. Sept. 742 = dracunculoides.
nutans, Willd. Sp. Pl. iii. 1831 = maritima,
Nuttalliana, Bess. ex Hook. Fl. Bor. Am. i. 326 = dracunculoides.
obtusiloba, *Ledeb. Fl. Alt.* iv. 68.—Sibir. alt.
occitanica, Salzm. ex DC. Prod. vi. 96 = campestris.
odoratissima, *Desf. Fl. Atlant.* ii. 263.—Algeria.
officinalis, Gater. Fl. Montaub. 144 = vulgaris.
officinalis, *Gueldenst. ex Ledeb. Fl. Ross.* ii. 600.—Rossia.
ohotensis, Steph. ex Ledeb. l. c. 597 = rupestris.
oligantha, Miégev. in Bull. Soc. Bot. Fr. xviii. (1871) 368 = Mutellina.
Oliveriana, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 342 (Rel. Lehm. 150) = fragrans.
Oliveriana, J. Gay, ex Bess. in DC. Prod. vi. 101 = A. Herba-alba.
Ontina, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 294 = A. Herba-alba.
orientalis, Herb. Madr. ex DC. Prod. vi. 120 = pallens.
orientalis, Willd. Sp. Pl. iii. 1836 = austriaca.
pachystachya, DC. Prod. vi. 114 = pycnocephala.
pacifica, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 401 = canadensis.
Pallasiana, *Fisch. ex Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 61.—Rossia.
Pallasii, Spreng. Syst. iii. 494 = maritima.
Pallasii, Willd. ex Spreng. l. c. 493 = borealis.
pallens, *Wall. ex DC. Prod.* vi. 120.—Ind. or.
pallida, Salisb. Prod. 191 = pontica.
palmata, Lam. Encyc. i. 268 = caerulescens.
palmata, Lap. Hist. Abr. Pl. Pyr. 504 = gallica.
palmata, Steph. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 22 = rutaefolia.
Palmeri, *A. Gray, in Proc. Am. Acad.* xi. (1876) 79.—Calif.
palustris, *Linn. Sp. Pl.* 846.—Dahur.; Mongol.
paniculaeformis, DC. Prod. vi. 107 = vulgaris.
paniculata, *Lam. Encyc.* i. 265.—Ital.; Gall.
paniculata, Rottl. ex Wight, Contrib. 12 = Cyathocline lyrata.
paniculata, Roxb. Hort. Beng. 61; Fl. Ind. iii. 418 = vulgaris.

ARTEMISIA:—
paniculata, Sch. Bip. ex Willk. in Flora, xxxiv. (1851) 747 = variabilis.
Parishii, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 220.—Calif.
Parryi, *A. Gray, l. c.* vii. (1868) 361.—N. Mexico.
parviflora, *Buch.-Ham. ex Roxb. Hort. Beng.* 61; *Fl. Ind.* iii. 420.—Ind. or.; Burma.
parviflora, Wight, Cat. n. 1460 = vulgaris.
pauciflora, Bieb. Fl. Taur. Cauc. ii. 290 = arenaria.
pauciflora, Georgi, Beschr. Russ. Reich. Nachtr. 306 = maritima.
pauciflora, Spreng. Syst. iii. 491 = desertorum.
pectinata, *Pall. Reise*, iii. 255.—Dahur.; Mongol.
pedatifida, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 399.—Am. bor.
pedemontana, Balb. ex Loisel. Not. 126 = lanata.
peduncularis, Bieb. Fl. Taur. Cauc. iii. 566 = splendens.
pedunculata, Steud. Nom. ed. II. 72 = splendens.
pedunculosa, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 176.—Japon.
pendula, Salisb. Prod. 191 = A. Absinthium.
pendula, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 39 = fragrans.
persica, *Boiss. Diagn.* Ser. I. vi. 91.—Persia; Affghan.; As. centr.
petraea, Turcz. ex Ledeb. Fl. Ross. ii. 598 = frigida.
petrosa, Jan, ex DC. Prod. vi. 118 = Baumgarteni.
peucedanifolia, Juss. ex DC. Prod. vi. 99 = canadensis.
pinnatifida, Jacquem. ex DC. Prod. vi. 120 = Tournefortiana.
piperita, Pall. ex Ledeb. Fl. Alt. iv. 88 = scoparia.
plattensis, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 397 = filifolia.
plumosa, Fisch. ex Bess. in DC. Prod. vi. 119 = annua.
pontica, Burm. f. Prod. 26 = afra.
pontica, *Linn. Sp. Pl.* 847.—Europ.; Reg. Caucas.; Soongaria.
potentillaefolia, Fisch. ex Spreng. Syst. iii. 494 = armeniaca.
potentilloides, A. Gray, in Proc. Am. Acad. vi. (1864) 551 = Tanacetum potentilloides.
Prescottiana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 72.—Am. bor. occ.
procera, Lap. Hist. Abr. Pl. Pyr. 503 = variabilis.
procera, *Willd. Sp. Pl.* iii. 1818.—Europ. austr. or.; As. Min.; Sibir.
procumbens, Schrad. ex DC. Cat. Hort. Monsp. 80 = frigida.
pseudo-pontica, *Schur, Enum. Pl. Transs.* 323.—Europ.
pubescens, *Ledeb. in Mém. Acad. Pétersb.* v. (1805) 568.—Sibir. or.
pulchella, Gmel. ex DC. Prod. vi. 102 = maritima.
pumila, Hort. ex Link, Enum. Hort. Berol. ii. 316 = frigida.
pumila, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 640 = Lindleyana.
punctata, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 43 = latifolia, macrobotrys.
purpurascens, Jacquem. ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 60 = Roxburghiana.
Pursbiana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 59.—Am. bor.
pusilla, *Luce, Topog. Nachr. Oesel*, 272.—Europ. bor.
pycnocephala, *DC. Prod.* vi. 99.—Sibir.
pycnorrhiza, *Ledeb. Fl. Alt.* iv. 79.—Sibir. alt.
pycnostachya, Torr. & Gray, Fl. N. Am. 422 = pycnocephala.
pyromacha, Viviani, Fl. Lib. Spec. 54. t. 13. f. 5 = hispanica.
quadripedalis, Gilib. Fl. Lituan. i. 174 = vulgaris.
quinqueloba, *Trautv. in Bull. Soc. Nat. Mosc.* xxxix. (1866) 1. 348.—As. temp.
racemifera, Steph. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 28 = sacrorum.
racemosa, *Kit. in Linnaea*, xxxii. (1863) 386.—Europ.
racemosa, Miégev. in Bull. Soc. Bot. Fr. xii. (1865) 341; xviii. (1871) 367 = spicata.
radicans, C. Sm. ex DC. Prod. vi. 102 = reptans.
ramosa, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 148.—Ins. Canar.
ramosa, Lag. ex Willk. & Lange, Prod. Fl. Hisp. ii. 71 = hispanica.

ARTEMISIA :—
Redowskyi, Ledeb. in Mém. Acad. Pétersb. v. (1805) 574 = A. Dracunculus.
repens, Pall. ex Willd. Sp. Pl. iii. 1840 = austriaca.
reptans, *C. Sm. ex Link, in Buch, Beschr. Canar. Ins.* 148.—Ins. Canar.
revoluta, Edgew. in Trans. Linn. Soc. xx. (1846) 72 = Roxburghiana.
rhodantha, *Rupr. Sert. Tiansch.* 52.—As. centr.
Richardsoniana, A. Gray, in Am. Journ. Sc. Scr. II. xxxiii. (1862) 239 = norvegiea.
Richardsoniana, *Hook. Fl. Bor. Am.* 323.—Am. arct.
rigida, *A. Gray, in Proc. Am. Acad.* xix. (1883) 49.—Am. bor. occ.
riparia, Buch.-Ham. ex DC. Prod. vi. 119 = carvifolia.
Rothrockii, *A. Gray, in Bot. Calif.* i. 618.—Calif.
Roxburghiana, *Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 57.—Reg. Himal.
Royleana, *DC. Prod.* vi. 115 (sp. dubia).—Ind. bor. occ.
rubella, Moench, Meth. 572 = caerulescens.
rubriflora, Turcz. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1836) 54 = vulgaris.
ruderalis, Salisb. Prod. 191 = vulgaris.
rupestris, All. Fl. Pedem. i. 169 = Mutellina.
rupestris, Asso, Syn. Arag. 117 = lanata.
rupestris, Lag. & Rodr. in Anal. Cienc. v. (1802) 284 = granatensis.
rupestris, Lam. Encyc. i. 262 = splendens.
rupestris, *Linn. Sp. Pl.* 847 ; *It. Goth.* 285.—Oeland. ; Saxon. ; Sibir.
rupestris, O. F. Muell. in Fl. Dan. t. 801 = norvegica.
rupestris, Scop. Fl. Carn. ed. II. ii. 146 = camphorata.
rupestris, Vill. Hist. Pl. Dauph. iii. 246 = spicata.
rutacea, Stev. ex DC. Prod. vi. 106 = rutaefolia.
rutaefolia, *Steph. ex Spreng. Syst.* iii. 488.—Sibir. or.
sabulosa, Stev. ex DC. Prod. vi. 108 = procera.
sabulosa, Willd. ex Ledeb. Fl. Ross. ii. 561 = arenaria.
sachaliensis, Tiles. ex Bess. in Bull. Soc. Nat. Mosc. viii. (1835) 48 = capillaris.
sacrorum, *Ledeb. in Mém. Acad. Pétersb.* v. (1805) 571.—Rossia ; Tibet. ; Sibir.
Saharae, *Pomel, Nouv. Mat. Fl. Atl.* 56.—Algeria.
saligna, Tenore, Prod. Fl. Nap. Suppl. i. 63 = variabilis.
salina, Bess. Enum. Pl. Volh. 76 = maritima.
salina, *Willd. Sp. Pl.* ii. 1834.—Europ.
salsa, F. Martini, Gesch. Nat. vii. 283 = nitrosa.
salsoloides, *Willd. Sp. Pl.* iii. 1832.—Reg. Caucas. ; Tibet. ; Sibir.
samamisica, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 50 = vulgaris.
santolinaefolia, Turcz. ex Bess. l. c. 87 = sacrorum.
Santonica, Lam. Encyc. i. 268 = caerulescens.
Santonica, Pursh, Fl. Am. Sept. 521 (sp. obscura).—Am. bor.
Santonica, Woodw. Med. Bot. t. 123 = suaveolens.
Santonicum, *Linn. Sp. Pl.* 845.—As. bor.
Santonicum, Sievers, ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 112 = campestris.
saxatilis, Reichb. Fl. Germ. Excurs. 202 = incanescens.
saxatilis, Waldst. & Kit. in Willd. Sp. Pl. iii. 1830 = camphorata.
saxorum, F. Schmidt, Reisen Amurl. 50, *err. typ.* = sacrorum.
scariosa, Rich. ex Bess. in DC. Prod. vi. 126 = Sieversiana.
Schimperi, *Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop.* 142, nomen.—Afr. trop.
Schlagintweitiana, *Klatt, in Sitzb. Acad. Muench.* (1878) 90.—Turkestan.
Schmidtiana, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 439.—Japon.
Schrenkiana, *Ledeb. Fl. Ross.* ii. 575.—Sibir.
scopaeformis, *Ledeb. l. c.*—Sibir.
scoparia, *Waldst. & Kit. Pl. Rar. Hung.* i. 66. t. 65.—Europ. ; Orient ; Ind. or.
scopulorum, *A. Gray, in Proc. Ac. Sc. Philad.* '1863' (1864) 66.—Am. bor. occ.
selengensis, Turcz. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 50 = vulgaris.
semavinensis, Bess. l. c. 65 = senjavinensis.
Semsek, *Forsk. Fl. Aegypt. Arab.* 218.—Arabia.
Senecionis, *Jacquem. ex Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 75.—Ind. or.
senjavinensis, *Bess l. c.* 64.—Am. occ. et As. or. arct.
sericea, *Weber, in Stechm. Artem.* 16. n. 5.—Sibir.

ARTEMISIA :—
sericophylla, Rupr. in Beitr. Pfl. Russ. Reich. ii. 41 = borealis.
Seriphium, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 71 = hispanica.
Seriphium, Wallr. Sched. Crit. 458 = maritima.
serotina, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 340 (*Lehm. Reliq. Bot.* 165).—Turkestan.
serrata, *Nutt. Gen. Am.* ii. 142.—Am. bor.
sibirica, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 429 = Tanacetum sibiricum.
Sieberi, Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 80 = A. Herba-alba.
sipontina, Tenore, Fl. Nap. v. 215 = caerulescens.
Siversiana, *Ehrh. ex Willd. Sp. Pl.* iii. 1845.—Rossia ; As. bor. ; Reg. Himal.
sogdiana, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 343 (*Lehm. Reliq. Bot.* 167).—Turkestan.
soongarica, *Schrenk, Enum. Pl. Nov.* 49 (1871).—Soongaria ; Turkest.
spicata, *Wulf. ex Jacq. Fl. Austr.* v. 46.—Europ.
spicigera, C. Koch, in Linnaea, xxiv. (1851) 345 = fragrans.
spinescens, *Eaton, in Bot. King, Exp.* 180. t. 19.—Am. bor. occ.
spithamaea, Pursh, Fl. Am. Sept. 522 = borealis.
splendens, *Willd. Sp. Pl.* iii. 1822.—As. Min. ; Reg. Caucas.
Stechmanniana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 35 = chamaemelifolia.
Stelleri, Stev. ex Ledeb. Fl. Ross. ii. 568 = borealis.
Stelleriana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 79.—Am. bor. ; Graecia ; Kamtschatka.
sternutatoria, Roxb. Hort. Beng. 61 ; Fl. Ind. iii. 423 = Centipeda orbicularis.
Steveniana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 64 = glomerata.
Stewartii, C. B. Clarke, Comp. Ind. 163 = annua.
Stracheyi, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 164.—Tibet. occ.
stricta, *Edgew. in Trans. Linn. Soc.* xx. (1846) 73.—Reg. Himal.
stricta, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xl. (1867) i. 204 = desertorum.
stricta, Heyne, ex DC. Prod. vi. 374 = Cyathocline lyrata.
suaveolens, Fisch. in DC. l. c. 119 = annua.
suaveolens, *Lam. Encyc.* i. 269.—Europ.
suavis, *Jord. Ind. Sem. Hort. Divion.* (1848) 18.—Gallia.
subcanescens, Willd. Enum. Hort. Berol. 861 = camphorata.
subspinescens, Boiss. Diagn. Ser. I. vi. 91 = persica.
subulata, Willd. ex Steud. Nom. ed. II. i. 140 = arenaria.
subulosa, Willd. ex DC. Prod. vi. 94 = arenaria.
subviscosa, *Bess. in Bull. Soc. Nat. Mosc.* ix. (1836) 24.—Sibir.
sueculenta, *Ledeb. Fl. Alt.* iv. 81.—Sibir.
supina, Stev. ex DC. Prod. vi. 112 = hololeuca.
sylvatica, *Maxim. Prim. Fl. Amur.* 161.—Mandchur.
sylvatica, Schlecht. ex Ledeb. Fl. Ross. ii. 581 = laciniata.
tanacetifolia, All. Fl. Pedem. i. 166 = spicata.
tanacetifolia, Georgi, It. i. 229 = sacrorum.
tanacetifolia, *Linn. Sp. Pl.* 848.—Sibir.
tanacetoides, Less. in Linnaea, vi. (1831) 210 = Matricaria multiflora.
tatarica, *Weber, ex Stechm. Artem.* 19.—Tataria.
taurica, *Bieb. Fl. Taur. Cauc.* ii. 291.—Taur.
taurica, Less. in Linnaea, x. (1835-36) 170 = Lessingiana.
taurica, Willd. Sp. Pl. iii. 1837 = fragrans.
tenella, *Weber, ex Stechm. Artem.* 26.—Hab. ?
tenuiflora, Jacquem. ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 62 = Tanacetum tenuifolium.
tenuifolia, Adams, in DC. Prod. vi. 107 = Adamsii.
tenuifolia, Ehrh. ex Ledeb. Fl. Ross. ii. 567 = commutata.
tenuifolia, Moench, Meth. 571 = pontiea.
tenuifolia, Willd. Sp. Pl. iii. 1819 = Eupatorium foeniculaceum.
tenuissima, Spreng. ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 30 = proeera.
Thunbergiana, Maxim. in Bull. Acad. Pétersb. xvii. (1872) 432 = apiacea.
thuscula, *Cav. in Anal. Cienc. Nat.* iv. (1801) 86.—Ins. Teneriffa.

ARTEMISIA :—
tibetica, Hook. f. Fl. Brit. Ind. iii. 329 = minor.
tifliensis, Adams, ex Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 62 = faseiculata.
Tilesii, Ledeb. in Mém. Acad. Pétersb. v. (1805) 568 = vulgaris.
tomentella, *Trautv. in Bull. Soc. Nat. Mosc.* xxxix. (1866) i. 351.—As. centr.
Tournefortiana, *Reichb. Ic. Exot.* i. 6. t. 5.—Orient ; Reg. Himal.
transsilvanica, Schur, Enum. Pl. Transs. 321 = Baumgarteni.
Trautvetteriana, *Bess. in Mém. Sav. Etr. Pétersb.* iv. (1845) 464. t. 5.—Orient ; Afghan.
trichophylla, Wall. Cat. n. 3305 = scoparia.
tridentata, *Nutt. in Trans. Am. Phil. Soc.* Scr. II. vii. (1841) 398.—Am. bor. occ.
trifida, *Nutt. l. c.*—Am. bor. occ.
trifida, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 196 = Tanaecetum trifidum.
trifurcata, Desf. Cat. Hort. Par. ed. III. (1829) 161 et 399 = Fontanesiana.
trifureata, *Steph. ex Spreng. Syst.* iii. 488.—Soongaria.
Triniana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 67.—Sibir.
Tschernieviana, Bess. in Bull. Soc. Nat. Mosc. viii. (1835) 33 = arenaria.
Turczaninoviana, *Bess. in Nouv. Mém. Soc. Nat. Mosc.* iii. (1834) 23.—Afghan.
turkestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 311.—Turkestan.
ucranica, Bess. ex Nym. Consp. 378 = Haussknechtii.
umbelliformis, Linn. Encyc. i. 262 = Mutellina.
umbrosa, Turcz. ex DC. Prod. vi. 113 = sclengensis.
uralensis, *Spreng. ex Bess. in Flora*, xvii. (1834) i. *Beibl.* 16, nomen.—Sibir.
Vahliana, *Kostel. Allg. Med.-Pharm. Fl.* ii. 698.—Persia.
valentina, Lam. Encyc. i. 269 = A. Herba-alba.
vallesiaca, All. Fl. Pedem. i. 169 = seq.
vallesiana, *Lam. Encyc.* i. 269.—Helv. austr. ; Ital.
variabilis, *Tenore, Fl. Neap. Prod. App.* v. 28.—Neap. ; Hisp.
Verlotorum, Lamotte, in Mém. Assoc. Franç. Cong. Clerm.-Ferr. (1876) 511 = vulgaris.
vermiculata, Linn. Mant. 291 = Stoebe artemisioides.
vermiculata, Schang. ex DC. Prod. vi. 99 = borealis.
vestita, *Wall. Cat.* n. 3301.—Reg. Himal. ; China.
Villarsii, Gren. & Godr. Fl. Fr. ii. 130 = Baumgarteni.
violacea, Desf. Tabl. ed. II. 112 = vulgaris.
violacea, Ledeb. in Mém. Acad. Pétersb. v. (1805) 567 = borealis.
virens, Moench, Meth. Suppl. 233 = vulgaris.
virgata, *Gueldenst. ex Ledeb. Fl. Ross.* ii. 600.—Ucrania.
virgata, Richards. in Frankl. Journ. App. 306 = frigida.
viridifolia, Ledeb. Ind. Sem. Hort. Dorpat. (1823) 2 = rupestris.
viridis, Willd. ex DC. Prod. vi. 124 = rupestris.
viscosa, DC. l. c. 95 = glutinosa.
vulgaris, Burm. f. Fl. Ind. 177 = indica.
vulgaris, *Linn. Sp. Pl.* 848.—Reg. temp. bor.
Wadei, Edgew. in Trans. Linn. Soc. xx. (1846) 72 = annua.
Wallichiana, Bess. in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 69 = Roxburghiana.
Willdenowii, Dietr. ex Bess. in Bull. Soc. Nat. Mosc. ix. (1836) 115 = camphorata.
Wrightii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 48.—Mexico.
Wulfenii, Schleich. Cat. Pl. Helv. ed. IV. 8 = Mutellina.
Zollingeriana, *Sch. Bip. in Zoll. Syst. Vers. Ind. Archip.* 126.—Malaya.

ARTENEMA, G. Don, in Sweet, Hort. Brit. ed. III. 516 (1839) = **Artanema**, D. Don (Scrophul.).

ARTHOSTEMA, Neck. Elem. ii. 280 (1790) = **Gnetum**, Linn.

ARTHRATHERUM, Beauv. Agrost. 32. t. 8. f. 8 (1812) = **Aristida**, Linn. (Gramin.).
arenarium, Nees, in Lehm. Pl. Preiss. ii. 98 = Aristida arenaria.
brachyatherum, Coss. & Dur. Expl. Algér Glum. 291 = Aristida acutiflora.

ARTHRATHERUM :—
brevifolium, Nees, Fl. Afr. Austr. i. 183 = Aristida brevifolia.
caloptilum, Jaub. & Spach, Illustr. iv. 54 = Aristida paradisea.
capense, Nees, Fl. Afr. Austr. 177 = Aristida capensis.
ciliatum, Nees, in Linnaea, vii. (1832) 287 = Aristida ciliata.
comosum, J. Gay, ex Kunth, Enum. Pl. i. 194 = Aristida stipiformis.
elatum, Boiss. & Buhse, Diag. Ser. II. iv. 128 = Aristida pogonoptila.
hirtiglume, Jaub. & Spach, Illustr. iv. 52 = Aristida hirtigluma.
hygrometricum, Beauv. Agrost. 33 = Aristida hygrometrica.
Hystrix, Nees, in Linnaea, vii. (1832) 287 = Aristida vestita.
Kotschyi, Hochst. ex Trin. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 159 = Aristida funiculata.
lutescens, Nees, Fl. Afr. Austr. i. 179 = Aristida lutescens.
megapotamicum, Nees, ex Trin. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 124 = Aristida implexa.
namaquense, Nees, Fl. Afr. Austr. i. 185 = Aristida namaquensis.
obtusum, Nees, l. c. 179 = Aristida obtusa.
pauciflorum, Boiss. & Buhse, Diag. Ser. II. iv. 127.—Persia.
plumosum, Nees, Fl. Afr. Austr. i. 182 = Aristida plumosa.
pogonoptilum, Jaub. & Spach, Illustr. iv. 56 = Aristida pogonoptila.
pungens, Beauv. Agrost. 33 = Aristida pungens.
rigidum, Schult. Mant. ii. 212 = Aristida rigescens.
Schimperi, Nees, Fl. Afr. Austr. 178 = Aristida ciliata.
stipoides, Beauv. Agrost. 33 = Aristida stipoides.
subacaule, Nees, Fl. Afr. Austr. 180 = Aristida subacaulis.
uniplume, Nees, l. c. 181 = Aristida uniplumis.
vestitum, Nees, l. c. 174 = Aristida vestita.
Zeyheri, Nees, l. c. 177 = Aristida Zeyheri.

ARTHRAXON, Beauv. Agrost. 111. t. 11. f. 6 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1128.
ALECTORIDIA, A. Rich. Tent. Fl. Abyss. ii. 447. t. 99 (1851).
BATRATHERUM, Nees, in Edinb. N. Phil. Journ. xviii. (1835) 180.
LASIOLYTRUM, Steud. in Flora, xxix. (1846) 18.
LUCAEA, Kunth, Rev. Gram. ii. 480. t. 159 (1830).
PLEUROPLITIS, Trin. Fund. Agrost. 174 (1820).
PSILOPOGON, Hochst. ex Reichb. Nom. 26 (1841).
ciliaris, *Beauv. l. c.*—Oriens; As. et Afr. trop.; Austral.
coloratus, Hochst. in Flora, xxxix. (1856) 188 = ciliaris.
cuspidatus, *Hochst. l. c.*—Ind. or.
echinatus, Hochst. l. c. = lanceolatus.
gracilis, Hochst. l. c. = ciliaris.
japonicum, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 288 = ciliaris.
Junghuhnii, *Hochst. in Flora*, xxxix. (1856) 189.—Java.
lanceolatus, *Hochst. l. c.* 188.—Ind. or.; Abyss.; Japon.
lancifolius, Hochst. l. c. = microphyllus.
Langsdorfianus, Hochst. l. c. = ciliaris.
lasiocoleos, Hochst. l. c. 189 = ciliaris.
major, Hochst. l. c. 188 = ciliaris.
micans, *Hochst. l. c.*—Ind. or.
microphyllus, *Hochst. l. c.*—Ind. or.
minor, Hochst. l. c. = microphyllus
nudus, Hochst. l. c. = ciliaris.
plumbeus, Hochst. l. c. 189 = ciliaris.
rudis, *Hochst. l. c.* 188 = cuspidatus.
Schimperi, Hochst. l. c. 189 = ciliaris.
Schmidtii, Hochst. l. c. = ciliaris.
serrulatus, Hochst. l. c. 188 = lanceolatus.
submuticus, Hochst. l. c. = ciliaris.
violaceus, Hochst. l. c. = ciliaris.

ARTHROCARPUM, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 510. *LEGUMINOSAE*.
gracile, Balf. f. l. c. 511.—Ins. Socotra.

ARTHROCHILUS, F. Muell. Fragm. i. 42 (1858) = **Drakaea**, Lindl. (Orchid.).
irritabilis, F. Muell. l. c. 43 = D. irritabilis.

ARTHROCHORTUS, Lowe, in Hook. Kew Journ. viii. (1856) 301 = **Lolium**, Linn. (Gram.).
loliaceus, Lowe, l. c.—Ins. Dezsertas (Madera).

ARTHROCLIANTHUS, Baill. Adansonia, ix. (1870) 296. *LEGUMINOSAE*.
sanguineus, *Baill. l. c.* 297.—N. Caled.

ARTHROCNEMUM, Moq. Chenop. Enum. 111. (1840), partim. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 65.
ambiguum, *Moq. Chenop. Enum.* 112.—Austral.
arbuscula, Moq. l. c. 113 = Salicornia arbuscula.
australasicum, F. Muell. ex Ung. Sternb. in Atti Congr. Bot. Firenze, 1874 (1876) 303 = Salicornia quinqueflora.
Belangerianum, Moq. Chenop. Enum. 112 = Halocnemum caspicum.
bidens, Nees, in Lehm. Pl. Preiss. i. 633 = Salicornia bidens.
caspicum, Moq. in DC. Prod. xiii. II.150 = Halocnemum caspicum.
ciliolatum, *Bunge, in Linnaea*, xxviii. (1856) 573, nomen.—As. Min.?
coralloides, *Losc. & Pard. Ser. Imperf. Pl. Arag.* 90; ed. II. 351.—Hispan.
fruticosum, Moq. Chenop. Enum. 111 = Salicornia fruticosa.
fruticosum, Torr. in Stansbury, Rep. 394 = Spirostachys occidentalis.
glaucum, *Delile, Fl. Egypt.* 69; *Ung. Sternb. in Atti Congr. Bot. Firenze*, 1874 (1876) 283.—Graecia; Afr. trop.; Ind. or.
halocnemoides, Nees, in Lehm. Pl. Preiss. i. 632 = Salicornia arbuscula.
indicum, *Moq. Chenop. Enum.* 113.—Ind. or.; Afr. trop.
indicum, Thw. Enum. Pl. Zeyl. 246 = Salicornia brachiata.
macrostachyum, *Bunge, ex Ung. Sternb. Vers. Syst. Salicorn.* 67.—Reg. Mediterr.
macrostachyum, Mor. & Delporte, in Ind. Sem. Hort. Bot. Taur. (1854) 35 = glaucum.
macrostachyum, Torr. Bot. Mex. Bound. Surv. 184 = Spirostachys occidentalis.
triandrum, F. Muell. Fragm. i. 139 = Salicornia robusta.

ARTHROLEPIS, Boiss. Diagn. Scr. I. xi. 14 (1849) = **Achillea**, Vaill. (Compos.).
imbricata, Boiss. l. c. 15 = Ach. membranacea.
membranacea, Boiss. l. c. = Ach. membranacea.

ARTHROLOBIUM, Reichb. Consp. 151 (1828) = ARTROLOBIUM, Desv. = **Ornithopus**, Linn. (Legum.).
ebracteatum, Auct. = O. ebracteatus.

ARTHROLOBUS, Andrz. ex DC. Syst. ii. 430 (1821) = **Rapistrum**, Crantz (Crucif.).
perennis, Andrz. ex DC. l. c. 431 = R. perenne.
rugosus, Andrz. ex DC. l. c. 432 = R. rugosum.

ARTHROLOBUS, Stev. ex DC. Syst. ii. 579 (1821) = **Sterigma**, DC. (Crucif.).
torulosus, Stev. ex DC. l. c. 580 = S. torulosum.

ARTHROMISCHUS, Thw. Enum. Pl. Zeyl. 47 (1858) = **Paramignya**, Wight (Rutac.).
armatus, Thw. l. c. = P. armata.

ARTHROPHYLLUM, Blume, Bijdr. 878 (1826). *ARALIACEAE*, Benth. & Hook. f. i. 944.
MORMORAPHIS, Jack, ex Wall. Cat. n. 4931 (1831).
alternifolium, Maing. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 734 = pinnatum.
Blumeanum, Zoll. & Moritzi, Syst. Verz. Pfl. Zoll. 41 = diversifolium.
ceylanicum, *Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 27.—Zeylan.?
dilatatum, *Miq. Fl. Ind. Bat.* i. 1. 768.—Java.
diversifolium, *Blume, Bijdr.* 879.—Malaya.
ellipticum, Blume, l. c. = diversifolium.
javanicum, Blume, l. c. = diversifolium.
ovalifolium, Jungh. & De Vriese, in Ann. Sc. Nat. Sér. III. vi. (1846) 117 = diversifolium.
pinnatum, *Maing. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 734.—Malaya.
reticulatum, *Blume, ex Miq. in Ann. Mus. Bot. Lugd. Bat.* i. 27.—Malaya.
serratifolium, *Miq. Fl. Ind. Bat. Suppl.* 341.—Ins. Banca.

ARTHROPHYLLUM, Boj. Hort. Maurit. 221 (1837) = **Phyllarthron**, DC. (Bignon.).
comorense, Boj. l. c. = P. comorense.
madagascariense, Boj. l. c.—Madag.

ARTHROPHYTUM, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 211. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 70.
Lehmannianum, *Bunge, Revis. Anab.* 46 = subulifolium.
subulifolium, *Schrenk, l. c.*—Turkestan.; Soongaria.

ARTHROPODIUM, R. Br. Prod. 276 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 790.
appendiculatum, *Avé-Lall. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* xi. 52.—Austral.
caespitosum, Spreng. Syst. ii. 87 = Stypandra caespitosa.
candidum, *Raoul, Choix Pl. Nouv. Zél.* 14.—N. Zel.
capillipes, *Endl. in Lehm. Pl. Preiss.* ii. 34.—Austral.
cirrhatum, *R. Br. in Bot. Mag.* t. 2350.—N. Zel.
dianellaccum, *F. Muell. Fragm.* x. 65.—Austral.
elongatum, DC. Cat. Hort. Monsp. 81, in obs. = paniculatum.
fimbriatum, R. Br. Prod. 276 = Dichopogon Sieberianus.
glaucum, Spreng. Syst. ii. 87 = Stypandra glauca.
hispidum, Spreng. l. c. = Anthericum hispidum.
imbricatum, Spreng. l. c. = Stypandra glauca.
laxum, Hook. f. Fl. Tasm. ii. 51 = Dichopogon strictus.
laxum, *Sieber, ex Schult. f. Syst.* vii. 441.—Austral.
Lindleyi, Kunth, Enum. Pl. iv. 621 = paniculatum.
minus, Lindl. Bot. Reg. t. 866 = paniculatum.
minus, *R. Br. Prod.* 276.—Austral.
muricatum, Spreng. Syst. ii. 87 = Anthericum muricatum.
neo-caledonicum, *Baker, in Journ. Linn. Soc.* xv. (1877) 352.—N. Caled.
paniculatum, *R. Br. Prod.* 276.—Austral.
pendulum, *DC. Cat. Hort. Monsp.* 80 = paniculatum.
Preissii, *Endl. in Lehm. Pl. Preiss.* ii. 35.—Austral.
scabrum, Spreng. Syst. ii. 87 = Agrostocrinum stipandroides.
strictum, R. Br. Prod. 276 = Dichopogon strictus.
umbellatum, Spreng. Syst. ii. 87 = Stypandra umbellata.

ARTHROPOGON, Nees, Agrost. Bras. 319 (1829). *GRAMINEAE*, Benth. & Hook. f. iii. 1119.
stipitatus, *Hackel, in Sitzb. Acad. Wien*, lxxxix. (1884) 125.—Cuba.
villosus, *Nees, Agrost. Bras.* 319.—Bras.

ARTHROSOLEN, C. A. Mey. in Bull. Phys-Math. Acad. Pétersb. i. (1843) 359. *THYMELAEACEAE*, Benth. & Hook. f. iii. 193.
anthylloides, C. A. Mey. l. c. = Lasiosiphon anthylloides.
calocephalus, *C. A. Mey. l. c.*—Afr. austr.
chrysanthus, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 165.—Afr. trop.
glaucescens, *Oliver, in Journ. Linn. Soc.* xv. (1871) 96.—Afr. trop.
gymnostachys, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 359.—Afr. austr.
inconspicuus, *Meissn. in DC. Prod.* xiv. 560.—Afr. austr.
laxus, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 359.—Afr. austr.
ornatus, *Meissn. in DC. Prod.* xiv. 559.—Afr. austr.
pimelioides, *Meissn. l. c.* 560.—Natal.
polycephalus, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 359.—Afr. austr.
sericocephalus, *Meissn. in DC. Prod.* xiv. 561.—Afr. austr.
somalensis, *Franch. in Revoil, Çomali*, 62. t. vi.—Somali.
spicatus, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* i. (1843) 359.—Afr. austr.
subspicatus, Endl. Gen. Suppl. iv. II. 63 = Wikstroemia subspicata.

ARTHROSPRION, Hassk. Retzia, i. 212 (1855) = **Acacia**, Willd. (Legum.).
stipulatum, Hassk. l. c. = Acacia concinna.

ARTHROSTACHYA, Link, Hort. Berol. i. 151 (1827) = **Gaudinia**, Beauv. (Gram.).
coarctata, Link, l. c. = G. geminiflora.

ARTHROSTACHYS, Desv. Opusc. 74 (1831) = **Andropogon**, Linn. (Gramin.).
gracilis, Desv. l. c.—Ind. or.

ARTHROSTEMA, Ruiz & Pav. Fl. Per. iv. t. 326 (1802). *MELASTOMACEAE*, Benth. & Hook. f. i. 740.
HETERONOMA, DC. Prod. iii. 122 (1828).
MELANIUM, Rich. ex DC. l. c. 135 (1828).
alatum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 35.—Venezuela.
alpestre, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 357 = Pterolepis alpestris.
angusturense, DC. Prod. iii. 135 = Pterolepis glomerata.
Aubletii, DC. l. c. 137 = Comolia Aubletii.
Bonplandii, DC. l. c. 138 = Monochaetum Bonplandii.
brachyandrum, Cham. in Linnaea, ix. (1834) 454 = Pterolepis glomerata, Tibouchina versicolor.
calcaratum, DC. Prod. iii. 138 = Monochaetum calcaratum.
campanulare, DC. l. c. 136 = Brachyotum campanulare.
campanulare, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 35.—Am. trop.
capitatum, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1850) 361.—Ind. occ.
cataphractum, Naud. l. c. 360 = Pterolepis cataphracta.
caulialatum, Ruiz & Pav. Fl. Per. iv. t. 327 = Aciotis caulialata.
cerastiifolium, DC. ex Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 122 = Tibouchina cerastifolia.
ciliatum, *Ruiz & Pav. Fl. Per.* iv. t. 326.—Peruv.
clidemioides, Miq. in Linnaea, xxii. (1849) 541 = Pterolepis clidemioides.
cubense, A. Rich. Ess. Fl. Cub. 550 = Chaetolepis cubensis.
Deppeanum, Schlecht. & Cham. in Linnaea, v. (1830) 566 = Monochaetum Deppeanum.
dicranantherum, DC. Prod. iii. 138 = Monochaetum dicranantherum.
exiguum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 359 = Pterolepis exigua.
filiforme, Naud. l. c. 358 = Pterolepis filiformis.
floribundum, Schlecht. in Linnaea, xiii. (1839) 431 = Monochaetum rivulare.
fragile, *Lindl. in Journ. Hort. Soc.* iii. (1848) 74.—Mexic.
glomeratum, Cham. in Linnaea, (1834) ix. 454 = Pterolepis glomerata.
Goudotianum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 359 = Pterolepis Goudotiana.
herbaceum, DC. Prod. iii. 137 = Tibouchina herbacea.
heterostemon, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 360 = Pterolepis filiformis.
hirsutissimum, DC. Prod. iii. 137 = Tibouchina herbacea.
hirsutissimum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 357 = Pterolepis pauciflora.
ladanoides, DC. Prod. iii. 136 = Pterolepis pusilla.
lanceolatum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 186.—Cuba.
latifolium, *D. Don, in Mem. Wern. Soc.* iv. (1823) 299.—Guiana.
lineatum, D. Don, l. c. = Monochaetum lineatum.
lutescens, *DC. Prod.* iii. 136 = Brachyotum lutescens.
macranthera, Miq. in Linnaea, xxii. (1849) 540 = Tibouchina sebastianopolitana.
Martinsianum, DC. Prod. iii. 137 = Tibouchina sebastianopolitana.
multiflorum, DC. l. c. 138 = Monochaetum multiflorum.
multiflorum, D. Don, in Mem. Wern. Soc. iv. (1823) 299 = Monochaetum dicranantherum.
myrtoideum, DC. Prod. iii. 138 = Monochaetum myrtoideum.
nitidum, R. Grah. in Edinb. N. Phil. Journ. xii. (1382) 186 = Tibouchina nitida.
nummularioides, DC. Prod. iii. 137 = Comolia nummularioides.
paniculatum, D. Don, in Mem. Wern. Soc. iv. (1823) 299 = Oxyspora paniculata.
pauciflorum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 357 = Pterolepis pauciflora.
piloselloides, DC. Prod. iii. 136 = Castratella piloselloides.
polygonoides, DC. l. c. 137 = Pterolepis polygonoides.
pumilum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 357 = Pterolepis pusilla.
pusillum, Naud. l. c. 361 = Pterolepis pusilla.

ARTHROSTEMA:—
quinquenerve, DC. Prod. iii. 136 = Brachyotum quinquenerve.
rosmarinifolium, DC. l. c. = Brachyotum rosmarinifolium.
Salzmannii, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 356 = Pterolepis Salzmannii.
Schiedeanum, Steud. Nom. ed. II. i. 141 (= *Pleroma longiflorum*?).—Mexic.
uruguayense, Cham. in Linnaea, ix. (1834) 453 = Tibouchina nitida.
versicolor, DC. Prod. iii. 137 = Tibouchina versicolor.
villosum, DC. l. c. = Comolia villosa.
villosum, Mart. Herb. Fl. Bras. n. 499 = Comolia ovalifolia.
volubile, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 35.—N. Granat.
Weddellianum, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 356 = Pterolepis Weddelliana.

ARTHROSTEMMA, DC. Prod. iii. 134 (1828) pro parte = **Brachyotum**, Triana (Melast.).

ARTHROSTEMMA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850) 355 = **Pterolepis**, Miq. (Melast.).

ARTHROSTYGMA, Steud. Nom. ed. II. i. 141 (1840) = **Petrophila**, R. Br. (Proteac.).

ARTHROSTYLIDIUM, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 117. *GRAMINEAE*, Benth. & Hook. f. Gen. iii. 1208.
AULONEMIA, Goudot, in Ann. Sc. Nat. Sér. III. v. (1846) 75. t. 5.
Burchellii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 43.—Bras.
capillifolium, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 531.—Ind. occ.
cubense, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 118. t. 4. f. 13.—Cuba.
excelsum, *Griseb. Fl. Brit. W. Ind.* 529.—Ind. occ.
fimbriatum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 531.—Ind. occ.
Haenkei, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 117. t. 3. f. 12.—Peruv.
leptophyllum, *Doell, in Mart. Fl. Bras.* ii. III. 175.—Bras.
longiflorum, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 41.—Venezuela.
maculatum, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 118. t. 5.—Peruv.
pubescens, *Rupr. l. c.* 119. t. 4. f. 14.—Ind. occ.
racemiflorum, *Steud. Syn. Pl. Gram.* 336.—Mexic.
Schomburgkii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 41.—Guiana.
Trinii, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 119. t. 4. f. 15.—Bras.

ARTHROSTYLIS, R. Br. Prod. 229 (1810). *CYPERACEAE*, Benth. & Hook. f. iii. 1058.
aphylla, Boeck. in Linnaea, xxxvii. (1871–73) 524 = Actinoschoenus Thouarsii.
aphylla, *R. Br. Prod.* 229.—Austral.
chinensis, Benth. Fl. Hongk. 397 = Actinoschoenus chinensis.
filiformis, Thw. Enum. Pl. Zeyl. 352 = Actinoschoenus filiformis.
Thouarsii, Kunth, Enum. Pl. ii. 284 = Actinoschoenus Thouarsii.

ARTHROSTYLIS, Boeck. in Linnaea, xxxvii. (1871–73) 523 = **Actinoschoenus**, Benth. (Cyperac.).
aphylla, Boeck. l. c. 524 = Act. Thouarsii.
filiformis, Boeck. l. c. = Act. filiformis.

ARTHROTAXIS, Endl. Gen. Suppl. i. 1372 (1841) err. typ. = **Athrotaxis**, D. Don (Conif.).

ARTHROTHAMNUS, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 251 = **Euphorbia**, Linn.
Bergii, Klotzsch & Garcke, in Abh. Akad. Berl. (1860) 63.—Afr. austr.
brachiatus, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 251; et Abh. Akad. Berl. (1860) 62 = E. brachiata.
Burmanni, Klotzsch & Garcke, ll. cc. = E. Burmanni.
cymosus, Klotzsch & Garcke, ll. cc. 251, 63 = E. nudiflora.

ARTHROTHAMNUS:—
densiflorus, Klotzsch & Garcke, in Monatsb. Akad. Berl. (1859) 251; et Abh. Akad. Berl. (1860) 62.—Afr. austr.
Ecklonii, Klotzsch & Garcke, ll. cc. 251, 63.—Afr. austr.
scopiformis, Klotzsch & Garcke, ll. cc. = E. scopiformis.
Tirucalli, Klotzsch & Garcke, ll. cc. 251, 62 = E. Tirucalli.

ARTHROTRICHUM, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 500; et Edinb. N. Phil. Journ. N. S. xvii. (Jan.–Apr. 1863) 234 = **Trichinium**, R. Br. (Amarant.).
calostachyum, F. Muell. ll. cc. = T. calostachyum.

ARTHROZAMIA, Reichb. Consp. 40 (1828) = **Encephalartos**, Lehm. (Cycad.).

ARTOCARPUS, Forst. Char. Gen. 101. t. 51 (1776). *URTICACEAE*, Benth. & Hook. f. iii. 376.
DURIO, Adans. Fam. ii. 399 (1763).
POLYPHEMA, Lour. Fl. Cochinch. 546 (1790).
RADERMACHIA, Thunb. in Handl. Vet. Akad. Stockh. (1776) 252.
RIMA, Sonner. Voy. Nouv. Guin. 99. t. 57 ad 60 (1776).
SITODIUM, Gaertn. Fruct. i. 344. t. 71, 72 (1791).
angustifolia, *Roxb. Hort. Beng.* 66, nomen.—Ins. Malay.
anisophylla, *Miq. Fl. Ind. Bat. Suppl.* 422.—Sumatra.
benghalensis, Roxb. ex Wall. Cat. sub n. 4655 = A. Lakoocha.
bifaria, Wall. ex Miq. Ann. Bot. Mus. Lugd. Bat. iii. 213 = Sloetia Sideroxylon.
biformis, *Miq. Fl. Ind. Bat. Suppl.* 419.—Sumatra.
Blumei, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 111.—Java.
brasiliensis, *Gomez, in Mem. Ac. Lisboa*, iii. (1812) *Mem. dos Corresp.* 84.—Bras.
calophylla, *Kurz, For. Fl.* ii. 431.—Tenasserim.
Camansi, *Blanco, Fl. Filip.* ed. I. 670.—Ins. Philipp.
canarana, *Miq. in Verh. Nederl. Inst.* III. v. (1852) (*Anal. Bot. Ind.*) 20.—Ind. or.
Cannoni, *Hort. Bull, ex Van Houtte, in Fl. des Serres*, xxi. (1875) 131.—Ins. Soc.
cerifera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 212.—Java.
Chama, Buch.-Ham. ex Wall. Cat. sub n. 4657 = A. Chaplasha.
Champeden, Spreng. Syst. iii. 804 = A. Polyphema.
Chaplasha, *Roxb. Hort. Beng.* 66; *et Fl. Ind.* iii. 525.—Ind. or.
communis, Forst. Char. Gen. 100 = incisa?
Cumingiana, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 119.—Ins. Philipp.
cuspidata, Griff. Notul. iv. 400 = rigida.
Dadak, *Miq. Fl. Ind. Bat. Suppl.* 420.—Sumatra.
dasyphylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 212.—Ins. Celebes.
dimorphophylla, *Miq. Fl. Ind. Bat. Suppl.* 417.—Ins. Banca.
echinata, Roxb. Hort. Beng. 66; et Fl. Ind. iii. 527 = rigida.
elastica, *Reinw. in Blume, Bijdr.* 481.—Borneo.
elongata, Miq. Fl. Ind. Bat. Suppl. 419 = Sloetia Sideroxylon.
erythrocarpa, Korth. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 213 = A. Dadak.
erythrocarpa, Teijsm. ex Miq. l. c. 212 = dasyphylla.
exculpta, *Hort.—Cf. Gard. Chron.* (1878) II. 84.—Hab.?
Finlaysoniana, *Wall. Cat.* n. 4662.—Malaya.
glauca, *Blume, Bijdr.* 483.—Java.
glaucescens, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 120.—Java.
Gomeziana, *Wall. ex Tréc. l. c.* 118.—Ind. or.
heterophylla, Lam. Encyc. iii. 209 = integrifolia.
hirsuta, *Lam. Encyc.* iii. 211.—Ind. or.
hirsutissima, *Kurz, in Tijdschr. Nederl. Ind.* xxvii. (1864) 182.—Ins. Banca.
hypargyrea, *Hance, ex Benth. Fl. Hongk.* 325.—Hongkong.
incisifolia, Stokes, Bot. Mat. Med. iv. 331 = incisa.
incisa, *Linn. f. Suppl.* 411.—Malaya; Ins. Pacif.
integrifolia, *Linn. f. l. c.* 412.—Ind. or.; Malaya.
Jaca, Lam. Encyc. iii. 209 = integrifolia.
Kemando, *Miq. Fl. Ind. Bat. Suppl.* 418.—Sumatra.

ARTOCARPUS:—
Kertau, *Zoll. Syst. Verz. Ind. Archip.* 89, 95.—Malaya.
laciniata, *Hort.—Cf. Gard. Chron.* (1875) II. 159; *et Rev. Hortic.* (1877) 38.—Ins. Pacif.
Lacucha, Roxb. Hort. Beng. 66, nomen = A. Lakoocha.
laevis, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 124.—Java.
Lakoocha, *Roxb. Fl. Ind.* iii. 524.—Ind. or.; Malaya.
lamellosa, Blanco, Fl. Filip. ed. I. 667 = nitida.
lanceaefolia, *Roxb. Hort. Beng.* [103]; *Fl. Ind.* iii. 527.—Malaya.
lanceolata, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 121.—Manila.
Limpato, *Miq. Fl. Ind. Bat. Suppl.* 421.—Sumatra.
madagascariensis, *Boj. Hort. Maurit.* 290.—Madag.
mariannensis, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 114.—Ins. Mariann.
maxima, Blanco, Fl. Filip. ed. I. 669 = integrifolia.
mollis, *Miq. Fl. Ind. Bat. Suppl.* 420.—Sumatra.
mollis, Wall. Cat. n. 4661 = A. Lakoocha.
Nanca, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 7.—Malaya.
nitida, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 119.—Ins. Philipp.
nobilis, *Thw. Enum. Pl. Zeyl.* 282.—Zeylan.
odoratissima, *Blanco, Fl. Filip.* ed. I. 671.—Ins. Philipp.
ovata, Blanco, l. c. 666 = Cumingiana.
ovata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 7.—Malaya.
palembanica, *Miq. Fl. Ind. Bat. Suppl.* 424.—Java.
parviflora, *Voigt, in Syll. Ratisb.* ii. (1828) 53.—Hab.?
peduncularis, *Kurz, in Journ. Bot.* xiii. (1875) 331.—Ins. Nicobar.
petiolaris, *Miq. Fl. Ind. Bat. Suppl.* 422.—Sumatra.
philippensis, *Lam. Encyc.* iii. 210.—Ins. Philipp.
pilosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 7.—Malaya.
pilosa, Reinw. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 211 = A. Polyphema.
Polyphema, *Pers. Syn.* ii. 531.—Malaya.
pomiformis, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 400 = Gomeziana.
Ponga, Dennst. Schluessel, 30 = Broussonetia integrifolia.
pubescens, *Blume, Bijdr.* 481.—Java.
pubescens, Moon, Cat. Pl. Ceyl. 61 = nobilis.
pubescens, Willd. Sp. Pl. iv. 189 = integrifolia.
reticulata, Heyne, ex Wall. Cat. sub n. 4655 = A. Lakoocha.
reticulata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 213.—Ins. Celebes.
Riedelii, *Miq. l. c.*—Ins. Celebes.
rigida, *Blume, Bijdr.* 482.—Burma; Malaya.
rigida, Wall. Cat. n. 4658 A = Gomeziana.
Rima, Blanco, Fl. Filip. ed. I. 671 = incisa.
rufa, *Miq. Fl. Ind. Bat. Suppl.* 423.—Malaya.
rufescens, *Kurz, For. Fl.* ii. 431.—Ind. or.
rufescens, *Miq. Fl. Ind. Bat. Suppl.* 420.—Malaya.
serrata, Roxb. Hort. Beng. 66, nomen.—Ind. or.
Tampang, Miq. Fl. Ind. Bat. Suppl. 421 = rufescens, *Miq.*
tephrophylla, *Miq. l. c.* 422.—Sumatra.
Teysmanni, *Miq. l. c.* 418.—Sumatra.
tylophylla, *Miq. Fl. Ind. Bat.* i. II. 289.—Malaya.
varians, *Miq. Fl. Ind. Bat. Suppl.* 417.—Sumatra.
venenosa, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie*, ii. (1845) 213.—Java.
Vriesiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 212.—Ins. Celebes.
Zollingeriana, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 90, 95.—Malaya.

ARTROLOBIUM, Desv. Journ. Bot. i. (1813) 121. t. 4 = **Coronilla**, Linn., **Ornithopus**, Linn., etc. (Legumin.).
creticum, Desv. l. c. = Coronilla cretica.
durum, Desv. l. c. = Ornithopus durus.
ebracteatum, Desv. l. c. = Ornithopus ebracteatus.
micranthum, Benth. in Mart. Fl. Bras. xv. I. 51.—Am. trop.
repandum, DC. Prod. ii. 311 = Ornithopus repandus.
scorpioides, Desv. Journ. Bot. i. (1813) 121 = Coronilla scorpioides.
Spasskii, Kalen. in Bull. Soc. Nat. Mosc. xviii. (1845) I. 233.—Tauria.
tauricum, Kalen. l. c.—Tauria.

ARUANA, Burm. f. Ind. Alt. (sign. G, verso) (1769) = **Myristica**, Linn.
silvestris, Burm. f. l. c. = M. moschata.

ARUBA, Aubl. Pl. Gui. i. 293. t. 115 (1775) = **Simaba**, Aubl. (Simarub.).
guianensis, Aubl. l. c.—Guiana.

ARUBA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 172. t. 19, 27, 29 = **Almeidea**, St. Hil. (Rutac.).
acuminata, Nees & Mart. l. c. 175 = Almeidea acuminata.
alba, Nees & Mart. l. c. 174 = Almeidea alba.
coerulea, Nees & Mart. l. c. = Almeidea caerulea.
myriacantha, Nees & Mart. ex Steud. Nom. ed. II. i. 142 = Almeidea myriantha.

ARUM, [Tourn.] Linn. Syst. ed. I. (1735), partim.
AROIDEAE, Benth. & Hook. f. Gen. iii. 967.
GYMNOMESIUM, Schott, in Oestr. Bot. Wochenbl. (1855) 17.
abyssinicum, A. Rich. Tent. Fl. Abyss. ii. 352 = Sauromatum abyssinicum.
aculeatum, Steud. Nom. ed. I. 73 = Caladium aculeatum.
alatum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 4.—Malaya.
albinervium, Kotschy, ex Engl. in DC. Monog. Phan. ii. 590 = hygrophilum.
albispathum, Hort. ex Engl. l. c. 591 = italicum.
albispathum, Stev. ex Ledeb. Fl. Ross. iv. 9 = orientale.
alpinum, Schott & Kotschy, in Bot. Zeit. ix. (1851) 285 = maculatum.
alternans, Engl. in DC. Monog. Phan. ii. 372 = Philodendron crassinervium.
alternum, Vell. Fl. Flum. ix. t. 116 = Philodendron crassinervium.
amphibium, Vell. l. c. 112 = Philodendron amphibium.
aquaticum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 5.—Malaya.
angulatum, *Griff. Notul.* iii. 143.—Ind. or.
aphyllum, Hook. in W. Gray, Western Afr. 386. t. A = Amorphophalli sp.
arborescens, Linn. Sp. Pl. 967 = Montrichardia arborescens.
Arisarum, Linn. l. c. 966 = Arisarum vulgare.
Arisarum, Lour. Fl. Cochinch. 655 = Typhonium cochinchinense.
atrorubens, Ait. Hort. Kew. iii. 315 = Arisaema atrorubens.
atrorubens, Spreng. Syst. iii. 769 = Pinellia tuberifera.
auritum, Linn. Syst. ed. X. 1251; Pl. Jam. Pugill. 26 = Syngonium auritum.
auritum, Vell. Fl. Flum. ix. t. 113 = Syngonium Vellozianum.
balearicum, Buchoz, Dec. Pl. viii. 11 = pictum.
Besserianum, Schott, in Oestr. Bot. Zeitschr. (1858) 349 = maculatum.
betifolium, *Mill. Gard. Dict.* ed. VIII. n. 12.—Am. bor.
bicolor, Ait. Hort. Kew. iii. 316 = Caladium bicolor.
bubalinum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 4.—Malaya.
bulbiferum, Roxb. Hort. Beng. 65; Sims, Bot. Mag. t. 2072 = Amorphophallus bulbifera.
bulbiferum, Salisb. Prod. 260 = Pinellia tuberifera.
bulbosum, Pers. ex Kunth, Enum. Pl. iii. 54 = Pinellia tuberifera?
byzantinum, Schott, in Oestr. Bot. Wochenbl. (1857) 212 = italicum.
caeruleo-punctatum, Zipp. ex Kunth, Enum. Pl. iii. 33 = Amorphophallus variabilis.
calyptrale, Salisb. Prod. 261 = Arisarum vulgare.
campanulatum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 509 = Amorphophallus campanulatus.
canariense, Webb & Berth. Phyt. Canar. iii. 293 = italicum.
cannaefolium, Linn. f. Pl. Surin. n. 123 = Spathiphyllum cannaeforme?
carratracense, Haensel. in Bot. Zeit. iv. (1846) 313 = Biarum Bovei.
catavericum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 4.—Malaya.
ciliatum, Hort. ex Loud. Encyc. Suppl. ii. 1508 = Cryptocoryne ciliata.

ARUM:—
cilicicum, Kotschy, ex Engl. in DC. Monog. Phan. ii. 589 = rupicolum.
clavatum, Desf. Cat. Hort. Par. ed. III. 385 = Sauromatum pedatum.
Colocasia, Linn. Sp. Pl. 965 = Colocasia antiquorum.
colocasioides, Desf. Cat. Hort. Par. ed. III. 7 et 385 = Colocasia antiquorum.
concinnatum, Schott, Ic. Aroid. t. 39 = italicum.
conophalloides, Kotschy, ex Schott, Prod. Aroid. 97 = orientale.
consobrinum, *Schott, l. c.* 87.—Reg. Cauc.
cordatum, Vell. Fl. Flum. ix. t. 111 = Philodendron cordatum.
cordifolium, Bory, Voy. i. 376; ii. 67 = Alocasia indica.
corsicum, Loisel. Fl. Gall. ii. 617 = pictum.
costatum, Wall. Tent. Fl. Nep. t. 19 = Arisaema costatum.
crassipes, Boiss. Diagn. Ser. I. xiii. 9 = Helicophyllum crassipes.
crenatum, Wight, in Hook. Bot. Misc. ii. (1831) 100 = Theriophonum crenatum.
creticum, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 9.—Creta.
crinitum, Ait. Hort. Kew. iii. 314 = Helicodiceros muscivorus.
crudele, Salisb. Prod. 261 = Dieffenbachia Seguine.
cucullatum, Lour. Fl. Cochinch. ii. 536 = Alocasia cucullata.
curvatum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 506 = Arisaema curvatum.
cuspidatum, Blume, Cat. Gew. Buitenz. 101 = Typhonium cuspidatum.
cuspidatum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 506 = Arisaema cuspidatum.
cylindraceum, Gasp. in Guss. Fl. Sic. Syn. ii. II. 597 = italicum.
cyprium, Schott, in Bonplandia, x. (1861) 368 = Dioscoridis.
decurrens, Blanco, Fl. Filip. ed. I. 656 = Amorphophallus decurrens.
detruncatum, *C. A. Mey. ex Schott, Prod. Aroid.* 8.—Pers. bor.
Dioscoridis, *Sibth. & Sm. Fl. Graec. Prod.* ii. 245.—Graecia; As. Min.
discolor, *Voigt, in Syll. Ratisb.* ii. 50.—Hab.?
divaricatum, Dulac, Fl. Hautes-Pyr. 46 = italicum.
divaricatum, Linn. Sp. Pl. 966 = Typhonium divaricatum.
diversifolium, Blume, Cat. Gew. Buitenz. 102 = Typhonium divaricatum.
Dracontium, Linn. Sp. Pl. 964 = Arisaema Dracontium.
Dracontium, Lour. Fl. Cochinch. 533 = Pinellia Wawrae.
Dracontium, Thunb. Fl. Jap. 233 = Arisaema Thunbergii.
Dracontium, Vell. Fl. Flum. ix. t. 103 = Staurostigma Luschnathianum.
Dracunculus, Linn. Sp. Pl. 964 = Dracunculus vulgaris.
Dracunculus, Thunb. Fl. Jap. 233 = Arisaema japonicum.
echinatum, Wall. Pl. As. Rar. ii. 30. t. 136 = Arisaema echinatum.
Ehrenbergii, *Schott, in Oestr. Bot. Zeitschr.* viii. (1858) 386.—Syria.
elongatum, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) II. 67 = orientale.
erubescens, Wall. Pl. As. Rar. ii. 30. t. 135 = Arisaema erubescens.
esculentum, Linn. Sp. Pl. 965 = Colocasia antiquorum.
exsertum, Salisb. Prod. 260 = A. Dracontium.
filiforme, Reinw. in Blume, Cat. Gew. Buitenz. 103 = Arisaema filiforme.
flagelliferum, *Griff. Notul.* iii. 144.—Ind. or.
flagelliforme, Roxb. Hort. Beng. 65; Lodd. Bot. Cab. t. 396 = Typhonium cuspidatum.
flavum, Forsk. Fl. Aegypt. Arab. 157 = Arisaema flavum.
foetidum, Salisb. Prod. 261 = italicum.
fornicatum, Roth, Pl. Ind. Or. 362 = Pinellia tuberifera.
fornicatum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 501 = Alocasia fornicata.
fornicatum, Wight, Ic. iii. t. 789 = Alocasia decipiens.
fugax, Salisb. Prod. 260 = venosum.
gracile, Roxb. Hort. Beng. 65; Fl. Ind. iii. 505 = Typhonium gracile.
gracile, Unverr. ex Schott, Prod. Aroid. 91 = maculatum.
gramineum, Lam. Encyc. iii. 30 = Biarum gramineum.

ARUM:—
gramineum, [Soland. in] Russ. Aleppo, ed. II. ii. 264 = Biarum Russellianum.
grandifolium, Jacq. Hort. Schoenbr. t. 189 = Philodendrum grandifolium.
gratum, Schott, Syn. Aroid. 11 = orientale.
Griffithii, *Schott, l. c.* 15.—Afghanist.
guellekense, Kotschy, ex Engl. in DC. Monog. Phan. ii. 589 = rupicolum.
guttatum, Salisb. Prod. 260 = Dracunculus vulgaris.
guttatum, Wall. Pl. As. Rar. ii. 10. t. 115 = Sauromatum guttatum.
hederaceum, Jacq. Enum. Pl. Carib. 31 = Philodendron Hoffmannii.
hederaceum, Willd. Sp. Pl. iv. 486 = Philodendron hederaceum.
Heldreichii, Orph. ex Boiss. Fl. Orient. v. 40 = maculatum.
helleborifolium, Jacq. Coll. iii. 217 = Xanthosoma helleborifolium.
heterophyllum, Aucher, ex Blume, Rumphia, i. 121 = Helicophyllum Rauwolffii.
hygrophilum, *Boiss. Diagn.* Ser. I. xiii. 8.—Syria.
immaculatum, Reichb. Ic. Fl. Germ. vii. 6; Schott, Prod. Aroid. 94 = italicum.
incomptum, Schott, l. c. 89 = orientale.
incurvatum, Lam. Fl. Fr. iii. 538 = Arisarum vulgare.
indicum, Lour. Fl. Cochinch. ii. 536 = Colocasia indica.
integrifolium, Link, Enum. Hort. Berol. ii. 394 = Aglaonema integrifolium.
intermedium, Schur, ex Schott, Prod. Aroid. 91 = maculatum.
intortum, [Soland. in] Russ. Aleppo, ed. II. ii. 264 = Helicophyllum Rauwolffii.
italicum, *Mill. Gard. Dict.* ed. VIII. n. 2.—Europ.
Jacquemontii, *Blume, Rumphia*, i. 118.—As. trop.
Korolkowi, *Regel, in Act. Hort. Petrop.* ii. (1873) 407. —Turkestan.
Kotschyi, *Boiss. & Hohen. Diagn.* Ser. I. xiii. 8.—Pers. bor.
laceratum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4 = seq.?
lacerum, Jacq. Hort. Schoenb. iv. 35 = Philodendron lacerum.
lanceolatum, Vell. Fl. Flum. ix. t. 114 = Philodendron crassinervium.
Liepoldii, Schott, Prod. Aroid. 77 = Dioscoridis.
lingulatum, *Linn. Pl. Jam. Pugill.* 26; *Syst.* ed. X. 1251.—Jamaica.
liniferum, Arruda, in Koster, Trav. Braz. 483 = Montrichardia linifera.
lividum, Salisb. Prod. 260 = Colocasia antiquorum.
longicyrrhum, Schott, Syn. Aroid. 14 = hygrophilum.
longispathum, Reichb. Ic. Fl. Germ. vii. 5 = orientale.
lyratum, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 508 = Amorphophallus lyratus.
macrorrhizum, Linn. Sp. Pl. 965 = Alocasia macrorrhiza.
macrourum, *Bunge, Enum. Pl. Chin. Bor.* 67.—China bor.
maculatum, All. Fl. Pedem. ii. 228 = italicum.
maculatum, Gueldenst. ex Ledeb. Fl. Ross. iv. 9 = orientale.
maculatum, *Linn. Sp. Pl.* 966.—Europ.
Malyi, Schott, Prod. Aroid. 93 = maculatum.
margaritiferum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 512 = Plesmonium margaritiferum.
maximum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 5.—Malaya.
marmoratum, Schott, Prod. Aroid. 85 = italicum.
melanopus, *Boiss. Fl. Orient.* v. 40.—Oriens.
minutum, Willd. Sp. Pl. iv. 484 = Theriophonum crenatum.
montanum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 497 = Colocasia montana.
mucronatum, Lam. Encyc. iii. 12 = Alocasia macrorrhiza.
mucronatum, Spreng. Syst. iii. 769 = Xanthosoma mucronatum.
Mulleri, Zipp. ex Blume, Rumphia, i. 143 = Amorphophallus Mulleri.
Murrayi, J. Grah. Cat. Bomb. Pl. 229 = Arisaema Jacquemontii.
muscivorum, Linn. f. Suppl. 410 = Helicodiceros muscivorus.
nepenthoides, Wall. Tent. Fl. Nep. 26. t. 18 = Arisaema nepenthoides.
Nickellii, Schott, Prod. Aroid. 85 = italicum.

ARUM:—
nigrum, *Schott, in Oestr. Bot. Wochenbl.* (1857) 212.—Europ.
nigrum, Vell. Fl. Flum. ix. t. 107 = Xanthosoma violaceum.
Nordmanni, Schott, Syn. Aroid. 12 = orientale.
novum, *Linn. f. Pl. Surin.* n. 122.—Guiana.
numidicum, Schott, Syn. Aroid. 10 = italicum.
nymphaeifolium, Roxb. Hort. Beng. 65; Fl. Ind. iii. 495 = Colocasia antiquorum.
oblongum, Vell. Fl. Flum. ix. t. 115 = Philodendron oblongum.
obtusilobum, *Link, Enum. Hort. Berol.* ii. 393.—Hab.?
odoratum, Heynh. Nom. ii. 46 = seq.
odorum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 499 = Alocasia odora.
orientale, *Bieb. Fl. Taur. Cauc.* ii. 207.—Taur.; Caucas.
orientale, Vis. ex Nym. Consp. 754 = nigrum.
orixense, R. Br. Prod. 336 = Typhonium Brownii.
orixense, Roxb. ex Andr. Bot. Repos. t. 356 = Typhonium trilobatum.
ovatum, Linn. Sp. Pl. 967 = Lagenandra toxicaria.
palaestinum, *Boiss. Diagn.* Ser. I. xiii. 6.—Oriens.
pedatum, Fisch. ex Spreng. Syst. iii. 769 = Sauromatum guttatum.
pedatum, Willd. Enum. Hort. Berol. Suppl. 54 = Sauromatum pedatum.
pellucidum, Fulchir, ex Kunth, Enum. Pl. iii. 42 = Caladium bicolor.
peltatum, Lam. Encyc. iii. 13 = Caladium esculentum.
pentaphyllum, Linn. Sp. Pl. 964 = Arisaema pentaphyllum.
pentaphyllum, Vell. Fl. Flum. ix. t. 104 = Xanthosoma pentaphyllum.
peregrinum, Linn. Sp. Pl. 966 = Alocasia macrorrhiza.
Petteri, Schott, Syn. Aroid. 12 = orientale.
philistaeum, *Kotschy, ex Schott, Prod. Aroid.* 79.—Palaestina.
phrygium, *Boiss. ex Tchihat. Asie Min. Bot.* ii. 648.—As. Min.
pictum, *Linn. f. Suppl.* 410.—Ins. balear.; Corsic.
pictum, Pett. Bot. Wegw. (1832); ex Schott, Syn. Aroid. 13 = orientale.
pinnatifidum, Jacq. Hort. Schoenb. ii. 31 = Philodendron pinnatifidum.
pinnatifidum, Vell. Fl. Flum. ix. t. 110 = Philodendron bipinnatifidum.
polymorphum, *Buckl. in Am. Journ. Sc.* xlv. (1843) 173.—Am. bor.
polyphyllum, *Link, Enum. Hort. Berol.* ii. 393.—Am. austr.
ponticum, Schott, in Bonplandia, x. (1862) 148 = italicum.
proboscideum, Linn. Sp. Pl. 966 = Arisarum proboscidcum.
ptychiurum, Zipp. ex Kunth, Enum. Pl. iii. 26 = Typhonium cuspidatum.
pulchrum, Salisb. Prod. 260 = Caladium bicolor.
pumilum, Kotschy, ex Engl. in DC. Monog. Phan. ii. 586 = philistaeum.
pumilum, Lam. Encyc. iii. 8 = Typhonium trilobatum.
punctatum, Desf. Cat. Hort. Par. ed. III. 385 = Alocasia indica.
punctulatum, Zipp. ex Kunth, Enum. Pl. iii. 34 = Amorphophallus bulbifer.
purpureum, *Thunb. Fl. Jav.* 20.—Japon.
pyrenaeum, Dufour, in Lap. Hist. Abr. Pl. Pyr. Suppl. 143 = maculatum.
Pythonium, Mart. Auswahl, 18 = Zomicarpa Pythonium.
quinatum, Nutt. Gen. Am. ii. 222 = Arisaema quinatum.
ramosum, *Link, Enum. Hort. Berol.* ii. 394.—Hab.?
rapiforme, Roxb. Hort. Beng. 65; Fl. Ind. iii. 497 = Alocasia rapiformis.
Rauwolffii, Blume, Rumphia, i. 122 = Helicophyllum Rauwolffii.
regnium, Rodsch. 'in Bald. Diar. xxviii. 32'; ex G. F. W. Mey. Esseq. 273 = Dieffenbachia Seguine.
ringens, *Thunb. in Trans. Linn. Soc.* ii. (1794) 337.—Japon.
Rumphii, Gaudich. in Freyc. Voy. Bot. 127. t. 34 = Amorphophallus campanulatus.
rupicola, *Boiss. Diagn.* Ser. I. xiii. 7.—Syria.
sagittaefolium, Hort. ex Steud. Nom. ed. II. i. 143 = Caladium hastifolium.

ARUM:—
sagittaefolium, Linn. Sp. Pl. 966 = Xanthosoma sagittaefolium.
sagittatum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 4.—Malaya.
sagittifolium, Link, Enum. Hort. Berol. ii. 393 = Xanthosoma hastifolium.
sagittifolium, Rodsch. Observ. 30 = Urospatha sagittifolia.
Sazensoo, Burger, ex Kunth, Enum. Pl. iii. 19 = Arisaema japonicum.
Seguine, Jacq. Enum. Pl. Carib. 31 = Dieffenbachia Seguine.
Seguinum, Linn. Sp. Pl. ed. II. 1371 = Dieffenbachia Seguine.
serratum, Thunb. in Trans. Linn. Soc. ii. (1794) 338 = Arisaema japonicum.
sessiliflorum, *Roxb. Hort. Beng.* 65; *Fl. Ind.* iii. 507. —Ind. or.
simlense, Hort. ex Engl. in DC. Monog. Phan. ii. 571 = Sauromatum venosum.
speciosum, Wall. Tent. Fl. Nep. 29. t. 20 = Arisaema speciosum.
spectabile, Schott, in Oestr. Bot. Wochenbl. (1857) 175 = Dioscoridis.
spectabile, Zipp. ex Kunth, Enum. Pl. iii. 34 = Amorphophallus bulbifer
spiculatum, Blume, Rumphia, i. 121 = Helicophyllum crassipes.
spirale, Retz. Obs. i. 30 = Cryptocoryne spiralis.
spirale, Salisb. Prod. 259 = Helicodiceros crinitus.
stercorarium, Zipp. ex Kunth, Enum. Pl. iii. 33 = Amorphophallus variabilis.
strepsiceros, *Sieber, ex Steud. Nom.* ed. II. i. 143.—Palaestina.
subulatum, Desf. Cat. Hort. Par. ed. III. 7 et 385 = Pinellia tuberifera.
sylvaticum, Roxb. Hort. Beng. [103]; Fl. Ind. iii. 511 = Synantherias silvaticus.
syriacum, Blume, Rumphia, i. 119 = Dioscoridis.
syriacum, Spreng. Syst. iii. 768 = Biarum Russellianum.
tenuifolium, Linn. Sp. Pl. 967 = Biarum tenuifolium.
ternatum, Thunb. Fl. Jap. 233 = Pinellia tuberifera.
tortuosum, Wall. Pl. As. Rar. ii. 10 = Arisaema tortuosum.
trapezuntinum, Schott, ex Engl. in DC. Monog. Phan. ii. 592 = italicum.
trifidum, Desf. Cat. Hort. Par. ed. III. 7 et 385 = Hydrosme Fontanesii.
trifoliatum, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 4.—Malaya.
trilobatum, Linn. Sp. Pl. 965 = Typhonium divaricatum.
trilobatum, Thunb. Fl. Jap. 234 = Pinellia tripartita.
tripartitum, Jacq. Hort. Schoenb. ii. 33. t. 190 = Philodendron tripartitum.
triphyllum, Houtt. Plantenk. ii. 11. 184 = Pinellia tuberifera.
triphyllum, Linn. Sp. Pl. 965 = Arisaema atrorubens.
triphyllum, Thunb. Fl. Jap. 233 = Arisaema ringens.
variolatum, Schott, Prod. Aroid. 81 = Xanthosoma violaceum.
venosum, Ait. Hort. Kew. iii. 315 = Sauromatum guttatum.
venosum, Balb. ex Engl. in DC. Monog. Phan. ii. 487 = Xanthosoma helleborifolium.
ventricosum, *Hoffmgg. Vers. Pfl. Nachtr.* ii. 25.—Bras.
vermitoxicum, Vell. Fl. Flum. ix. t. 108 = Caladium bicolor.
vernale, Salisb. Prod. 261 = maculatum.
virginicum, Linn. Sp. Pl. 966 = Peltandra virginica.
viviparum, Roxb. Hort. Beng. 65; Fl. Ind. iii. 496 = Remusatia vivipara.
virescens, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 6. —Persia.
vittatum, Salisb. Prod. 260 = Arisaema atrorubens.
vulgare, Lam. Fl. Fr. iii. 537 = maculatum.
xanthorrhizon, Jacq. Hort. Schoenbr. ii. 32 = Xanthosoma sagittifolium.
Zelebori, Schott, Prod. Aroid. 94 = maculatum.

ARUNA, Schreb. Gen. i. 26 (1789) = AROUNA, Aubl. = **Dialium**, Linn. (Legumin.).

ARUNCUS, Linn. Syst. ed. I. (1735); Gen. ed. I. 310 (1737) = **Spiraea**, Tourn. (Rosac.).
americanus, Rafin. Sylva Tellur. 152 = S. Aruncus.
astilboides, Maxim. in Act. Hort. Petrop. vi. (1879) 171 = S. Aruncus.
sylvester, Kostel. Ind. Hort. Prag. 15 = S. Aruncus.
vulgaris, Rafin. Sylva Tellur. 152 = S. Aruncus.

ARUNDINA, Blume, Bijdr. 401 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 521.
DILOCHIA, Lindl. Gen. et Sp. Orch. 38 (1830).
affinis, Griff. Notul. iii. 330 = chinensis.
bambusifolia, *Lindl. in Wall. Cat.* n. 3751.—Ind. or.; Malaya.
chinensis, *Blume, Bijdr.* 402.—Reg. Himal.; China.
densa, *Lindl. in Bot. Reg.* (1842) *Misc.* 25.—Singapore.
Meyenii, *Reichb. f. in Linnaea*, xxv. (1852) 227.—China.
minor, *Lindl. Gen. et Sp. Orch.* 125.—Zeylan.
pentandra, *Reichb. f. Xenia Orch.* ii. 11.—Java.
Philippi, Reichb. f. in Linnaea, xxv. (1852) 227 = chinensis.
pulchella, *Teijsm. & Binn. in Nederl. Kruid. Arch.* iii. (1855) 400.—China.
pulchra, *Miq. in Journ. Bot. Néerl.* i. (1861) 90.—China.
speciosa, Blume, Bijdr. 401. t. 73 = densiflora.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 37.—Ins. Pacif.
Wallichii, *Reichb. f. Xenia Orch.* ii. 13. t. 105.—Malaya.

ARUNDINARIA, Michx. Fl. Bor. Am. i. 73 (1803). *GRAMINEAE*, Benth. & Hook. f. iii. 1207.
LUDOLFIA, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 320.
MACRONAX, Rafin. in Med. Repos. New York, v. (1808) 352.
MIEGIA, Pers. Syn. i. 101 (1805).
THAMNOCALAMUS, Munro, in Trans. Linn. Soc. xxvi. (1868) 33.
TRIGLOSSUM, Fisch. Cat. Jard. Gorenk. (1812).
acuminata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 25.—Mexic.
amplissima, *Nees, in Linnaea*, ix. (1834) 479.—Bras.
aristulata, *Doell, in Mart. Fl. Bras.* ii. III. 165.—Bras.
attenuata, *Doell, l. c.* 170.—Bras.
bambusina, Trin. Fund. Agrost. 97 = macrosperma.
callosa, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 30.—Reg. Himal.
debilis, *Thw. Enum. Pl. Zeyl.* 375.—Zeylan.
decalvata, *Doell, in Mart. Fl. Bras.* ii. III. 170.—Bras.
densifolia, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 32.—Zeylan.
distans, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 621.—Bras.
elegans, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 249.—Burma.
falcata, *Nees, in Linnaea*, ix. (1834) 478.—Reg. Himal.
flexuosa, *Hance, in Journ. Bot.* xiv. (1876) 339.—China.
floribunda, *Thw. Enum. Pl. Zeyl.* 375.—Zeylan.
Fortunei, Fenzi, in Gard. Chron. (1876) II. 773 = Simoni.
Fortunei, *Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 897.—Japon.
gigantea, *Nutt. Gen. Am.* i. 39.—Am. bor.
glaucescens, *Beauv. Agrost.* 144, 152.—Ind. or.; China?
Griffithiana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 20.—Reg. Himal.
Hindsii, *Munro, l. c.* 31.—Hongkong.
hirsuta, *Munro, l. c.* 30.—Reg. Himal.
hispida, Steud. Syn. Pl. Gram. 335 = Wightiana.
Hookeriana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 29.—Reg. Himal.
intermedia, *Munro, l. c.* 28.—Reg. Himal.
interrupta, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 620 = falcata.
japonica, *Sieb. & Zucc. ex Steud. Syn. Pl. Gram.* 334.—Japon.
khasiana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 28.—Reg. Himal.
kurilensis, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* viii. (1850) 121.—Ins. Kurile.

ARUNDINARIA:—
longifolia, *Fourn. Mex. Pl. Enum. Gram.* 131, *ex Hemsl. Biol. Centr. Am. Bot.* iii. 586.—Mexic.
longiramea, Munro, in Trans. Linn. Soc. xxvi. (1868) 19 = sinica.
macrosperma, *Michx. Fl. Bor. Am.* i. 74.—Am. bor.
macrostachya, *Nees, in Linnaea*, ix. (1834) 481.—Bras.
Maximowiczii, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 783 = Simoni.
Metake, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 284 = japonica.
microphylla, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 32.—Reg. Himal.
moliniformis, Hochst. ex Munro, l. c. 19 = Wightiana.
multiflora, *Doell, in Mart. Fl. Bras.* ii. III. 166.—Bras.
parviflora, Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 619 = Arthrostylidium Trinii.
pinifolia, Nees, in Mart. Fl. Bras. ii. 525 = Chusquea pinifolia.
procera, Wall. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 34 = spathiflora.
racemosa, *Munro, l. c.* 17.—Reg. Himal.
radiata, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 115. t. 15. f. 9.—Bras.
Schomburgkii, Benn. in Trans. Linn. Soc. xviii. (1841) 562 = Arthrostylidium Schomburgkii.
Simoni, *A. & C. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 774.—China.
sinica, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 235.—China.
spathiflora, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. II. (1835) 617.—Reg. Himal.
stolonifera, Kurz, Cat. Hort. Bot. Calcut. 79 = Phyllostachys nigra.
suberecta, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 32.—Reg. Himal.
tecta, *Muhl. Desc. Uber.* 191.—Am. bor.
tessellata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 31.—Afr. austr.
Trianae, *Munro, l. c.* 25.—N. Granat.
Trinii, Hack. in Mart. Fl. Bras. ii. III. 174 = Arthrostylidium Trinii.
utilis, Cleghorn, in Journ. Agr. Soc. Ind. xiii. (1865) 388 = falcata.
verticillata, *Nees, in Mart. Fl. Bras.* ii. III. 523.—Bras.
Walkeriana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 21.—Zeylan.
Wightiana, *Nees, in Linnaea*, ix. (1834) 482.—Ind. or.
Wightiana, Thw. Enum. Pl. Zeyl. 444 = Walkeriana.
Wightii, Benth. Fl. Hongk. 434 = sinica.

ARUNDINELLA, Raddi, Agrost. Bras. 37. t. 1 (1823). *GRAMINEAE*, Benth. & Hook. f. iii. 1118.
ACRATHERUM, Link, Hort. Berol. i. 230 (1834).
BRANDTIA, Kunth, Rev. Gram. ii. 511. t. 170 (1830).
CALAMOCHLOË, Reichb. Consp. 52 (1828).
GOLDBACHIA, Trin. in Spreng. Neue Entdeck. ii. 81 (1821).
RIEDELIA, Trin. ex Kunth, Enum. Pl. i. 515 (1833).
THYSANACHNE, Presl, Thysan. Nov. Pl. Gen. (1829); et Rel. Haenk. i. 252 (1830).
Acratherum, *Nees, ex Steud. Nom.* ed. II. i. 143.—Ind. or.
agrostoides, Trin. Ic. xxiii. t. 265 = holcoides.
anomala, *Steud. Syn. Pl. Gram.* 116.—Japon.
auletica, Rupr. in Bull. Acad. Brux. ix. (1842) II. 242 = Deppeana.
avenacea, *Munro, ex Thw. Enum. Pl. Zeyl.* 362.—Zeylan.
brasiliensis, *Raddi, Agrost. Bras.* 37.—Bras.
ciliata, *Nees, ex Miq. in Verh. Nederl. Inst.* III. iv. (1851) 30.—Ind. or.
crinita, *Trin. in Linnaea*, x. (1836) 299.—Peruv.
cubensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 533.—Ind. occ.
Deppeana, *Nees, in Bonplandia*, iii. (1855) 84.—Mexic.
Ecklonii, Nees, Fl. Afr. Austr. 80 = nepalensis.
elegantula, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 233.—Afr. occ.
flammida, *Trin. Gram. Sp.* xxiii. t. 267.—Bras.
fuscata, *Nees, ex Buese, in Miq. Pl. Jungh.* 359.—Ind. or.
glabra, *Hook. & Arn. Bot. Beech. Voy.* 237.—China.

ARUNDINELLA:—
Goeringii, *Steud. Syn. Pl. Gram.* 116.—Japon.
hirsuta, Hochst. ex Steud. l. c. 115 = setifera.
hirsuta, *Nees, ex Steud. l. c.*—Reg. Himal.
holcoides, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 71 = Brandtia holcoides.
indica, *Trin. ex Steud. Nom.* ed. II. i. 143.—Ind. or.
khasiana, *Nees, ex Steud. Syn. Pl. Gram.* 115.—Reg. Himal.
latifolia, Fourn. Mex. Pl. Enum. Gram. 54; ex Hemsl. Biol. Centr. Am. Bot. iii. 515 = Deppeana.
martinicensis, *Trin. Gram. Panic.* ii. 62.—Ins. Martinic.
mesophylla, *Nees, ex Steud. Syn. Pl. Gram.* 115.—Ind. or.
Metzii, *Hochst. ex Miq. in Verh. Nederl. Inst.* III. iv. (1851) 31.—Ind. or.
Mikani, Nees, Agrost. Bras. 465 = brasiliensis.
mutica, *Nees, ex Steud. Syn. Pl. Gram* 116.—Ind. or.
nepalensis, *Trin. Sp. Gram.* t. 268.—Ind. or.
nervosa, Nees, ex Hook. & Arn. Bot. Beech. Voy. 237 = Panicum semialatum.
pallida, *Nees, Agrost. Bras.* 465.—Mexic.
peruviana, *Steud. Syn. Pl. Gram.* 115.—Peruv.
phragmitoides, *Griseb. Cat. Pl. Cub.* 234.—Cuba.
pilosa, Hochst. ex Miq. in Verh. Nederl. Inst. III. iv. (1851) 30 = ciliata.
pumila, *Steud. Syn. Pl. Gram.* 114.—Abyss.
purpurea, *Hochst. ex Steud. l. c.* 115.—Ind. or.
rigida, *Nees, Fl. Afr. Austr.* 80.—Afr. austr.
Schultzii, *Benth. Fl. Austral.* vii. 545.—Austral.
scoparia, Fourn. Mex. Pl. Enum. Gram. 55; ex Hemsl. Biol. Centr. Am. Bot. iii. 515 = Deppeana.
setifera, *Steud. Syn. Pl. Gram.* 115.—Ind. or.
setosa, *Trin. Gram. Panic.* 65.—Nepal.
stipoides, *Hack. in Sitzb. Acad. Wien*, lxxxix. (1884) 123.—Madag.
stricta, *Nees, in Hook. Kew Journ.* ii. (1850) 102.—Hongkong.
tenella, *Nees & Wight, ex Steud. Nom.* ed. II. i. 143.—Ind. or.
tricuspidula, *Buese, in Miq. Pl. Jungh.* 360.—Java.
villosa, *Arn. ex Steud. Syn. Pl. Gram.* 115.—Zeylan.
Wallichii, *Nees, ex Steud. l. c.* 114.—Ind. or.
Zollingeri, *Steud. l. c.* 115.—Java.

ARUNDO, Beauv. Agrost. 60. t. 13. f. 2 (1812) = **Phragmites**, Trin. (Gram.).

ARUNDO, Tourn. ex Linn. Gen. ed. I. 19 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1179.
AMPHIDONAX, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836).
DONACIUM, Frics, Bot. Notiser, (1843) 131.
DONAX, Beauv. Agrost. 77 (1812), pro parte.
SCOLOCHLOA, Mert. & Koch, in Roehl. Deutschl. Fl. i. 528 (1823).
acutiflora, Pollini, Fl. Veron. i. 88 = Deyeuxia varia.
acutiflora, Schrad. Fl. Germ. i. 217 = Deyeuxia acutiflora.
aegyptia, Delile, Fl. Aeg. Illustr. 4 = A. Donax.
aegyptiaca, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 984 = praec.
aggerum, *Kit. in Linnaea*, xxxii. (1863) 307.—Europ.
agrestis, Lour. Fl. Cochinch. 57 = Bambusa agrestis.
Agrostis, Lap. ex Willk. & Lange, Prod. Fl. Hisp. i. 88 = Molinia caerulea.
Agrostis, Scop. Fl. Carn. ed. II. i. 88 = Deyeuxia acutiflora.
Agrostis, [Weber, in] Wigg. Prim. Fl. Holsat. 10 = Deyeuxia acutiflora?
agrostoides, Pursh, Fl. Am. Sept. i. 86 = Deyeuxia canadensis.
airaeformis, *Raddi, Obs. Madera*, 10.—Madera.
airoides, *Poir. Encyc.* vi. 270.—Am. bor.
Alopecuros, Gaudich. ex Urv. in Mém. Soc. Linn. Par. iv. (1826) 602 = Festuca Alopecuros.
alpestris, Hall. ex Roem. & Schult. Syst. ii. 503 = Deyeuxia Halleriana.
alpina, [Honck.] Verz. Gew. Teutschl. 321; Schult. Oestr. Fl. ed. II. i. 189 = Calamagrostis tenella.
alpina, Vill. ex Kunth, Enum. Pl. i. 238 = Deyeuxia Halleriana.
altissima, Benth. Cat. Pl. Pyr. 62 = Phragmites communis.
Ampelodesmos, Cyrill. Pl. Rar. Neap. Fasc. ii. 30 = Ampelodesma tenax.
annulata, Roem. & Schult. Syst. ii. 503 = Calamagrostis littorea.

ARUNDO :—
antarctica, Urv. in Mém. Soc. Linn. Par. iv. (1826) 602 = Festuca antarctica.
arborea, Mill. Gard. Dict. ed. VIII. n. 5 = Bambusa arundinacea.
arenaria, Gorter, Fl. Ingr. 18 = Deyeuxia neglecta.
arenaria, Linn. Sp. Pl. 82 = Ammophila arundinacea.
argentea, [*Clairv.*] *Man. Herb.* 16. —Helvet.
australis, Cav. ex Roem. & Schult. Syst. ii. 511 = Phragmites australis.
australis, A. Rich. ex Steud. Nom. ed. II. i. 143, 250 = conspicua.
baltica, Fluegg. ex Schrad. Fl. Germ. 223. t. 5. f. 4 = Ammophila baltica.
Bambos, Burm. ex Steud. Nom. ed. II. i. 143 = Bambusa agrestis.
Bambos, Linn. Sp. Pl. 81 = Bambusa arundinacea.
Bambu, Lour. Fl. Cochinch. 56 = praec.
bengalensis, Boj. Hort. Maurit. 368 = Phragmites communis.
bengalensis, *Retz. Obs.* v. 20.—Bengal.
bicolor, *Desf. Fl. Atlant.* i. 107; *Poir. Voy. Barb.* ii. 104.—Afr. bor.
Biebersteiniana, Roem. & Schult. Syst. ii. Mant. 285 = Calamagrostis littorea.
bifaria, *Retz. Obs.* iv. 22.—Ind. or.
biflora, Lam. Illustr. i. 196 = Ampelodesmos tenax.
Boivini, *Steud. Syn. Pl. Gram.* 423.—Madag.
brevipilis, Torr. Fl. U. St. 95 = Ammophila brevipilis.
Calamagrostis, Bieb. Fl. Taur. Cauc. i. 78; ii. 456 = Calamagrostis epigeios.
Calamagrostis, Hall. f. in Roem. Arch. i. II. 10 = Deyeuxia Halleriana.
Calamagrostis, Leyss. Fl. Halens. 28 = Deyeuxia sylvatica.
Calamagrostis, Linn. Sp. Pl. 82 = Calamagrostis lanceolata.
Calamaria, Salisb. Prod. 24 = Calamagrostis lanceolata.
calycina, Willd. ex Steud. Nom. ed. II. i. 144 = nitida.
canadensis, Michx. Fl. Am. Bor. i. 73 = Deyeuxia canadensis.
canadensis, Steud. Nom. ed. II. i. 144 = Deyeuxia Nuttalliana.
canescens, [Weber, in] Wigg. Prim. Fl. Holsat. 10 = Calamagrostis epigeios, lanceolata.
canila, Molina, ex Steud. Syn. Pl. Gram. 336 = Chusquea Cumingii.
capillata, *Nem. ex Chaub. in Actes Linn. Soc. Bord.* xix (1853) 44.—Hab.?
ciliata, [*Clairv.*] *Man. Herb.* 16.—Helvet.
cinnoides, Muhl. Desc. Uber. 187 = Deyeuxia Nuttalliana.
Clarionis, Loisel. Fl. Gall. 719 = Deyeuxia Halleriana.
coarctata, Poir. Encyc. Suppl. iv. 704 = Deyeuxia coarctata.
coarctata, Torr. Fl. U. St. 94 = Deyeuxia Nuttalliana.
collina, Tenore, Fl. Nap. t. 108 = Pliniana.
colorata, Ait. Hort. Kew. i. 116 = Phalaris arundinacea.
confinis, Willd. Enum. Hort. Berol. 127 = Calamagrostis confinis.
conspicua, *Forst. f. Prod.* 9.—N. Zel.
contracta, Gilib. Exercit. ii. 542 = Calamagrostis lanceolata.
cuspidata, Schult. Mant. iii. 604 = Calamagrostis cuspidata.
cylindrica, J. Gay, ex Steud. Nom. ed. II. i. 144 = Deyeuxia acutiflora.
dioeca, Spreng. Syst. i. 361 = Gynerium argenteum.
dioica, *Lour. Fl. Cochinch.* 70.—Cochinch.
Donax, Georgi, Beschreib. Russ. Reich. iii. IV. 705 = Arundinaria kurilensis.
Donax, *Linn. Sp. Pl.* 81.—Reg. Mediterr.; Oriens.
effusa, C. C. Gmel. Fl. Bad. 262 = Calamagrostis littorea.
effusa, Poir. Encyc. Suppl. iv. 706 = Deyeuxia effusa.
Egmontiana, Roem. & Schult. Syst. ii. 511 = Phragmites Egmontianus.
epigeios, Forsk. ex Steud. Nom. ed. II. i. 144 = Imperata arundinacea.
epigeios, Huds. Fl. Ang. ed. I. 43; ed. II. 54 = Calamagrostis lanceolata.
epigeios, Linn. Sp. Pl. 81 = Calamagrostis epigeios.
eriantha, Poir. Encyc. Suppl. iv. 705 = Deyeuxia eriantha.
excelsa, Salisb. Prod. 24 = Bambusa arundinacea.
farcta, Aubl. Pl. Gui. i. 52 = Calamagrostis farcta.

ARUNDO :—
fastuosa, Willd. ex Steud. Nom. ed. II. i. 144 = Gynerium saccharoides.
Fax, Lour. Fl. Cochinch. 58 = Melocanna humilis.
festucacea, Willd. Enum. Hort. Berol. 126 = Graphephorum arundinaceum.
festucoides, Desf. Fl. Atlant. i. 108 = Ampelodesmos tenax.
filiformis, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 117.—Java.
fissa, Willd. ex Steud. Nom. ed. II. i. 144 = Deyeuxia canadensis.
flexuosa, Brongn. in Duperr. Voy. Bot. 29. t. 4 = Phragmites flexuosus.
flexuosa, [Clairv.] Man. Herb. 16 = Aira flexuosa.
fulvida, *F. Buch. in Trans. N. Z. Inst.* vi. (1874) 242. —N. Zel.
Gayana, *E. Desv. in C. Gay, Fl. Chil.* vi. 326.—Valdivia.
gigantea, Walt. Fl. Carol. 81 = Arundinaria macrosperma.
glauca, Bieb. Fl. Taur. Cauc. i. 79 = Calamagrostis littorea.
glauca, Hornem. Hort. Hafn. 74 = Deyeuxia Nuttalliana.
graeca, Link, in Linnaea, ix. (1834) 136 = Phragmites communis.
groenlandica, Schrank, in Regensb. Denkschr. i. II. (1818) 8 = Calamagrostis groenlandica.
Halleri, Willd. Prod. Fl. Berl. 60 = Stipa Calamagrostis.
Halleriana, Gaud. Agrost. Helv. i. 97 = Deyeuxia Halleriana.
Henslowiana, *Hook. & Arn. Bot. Beech. Voy.* 248.—China.
Hostii, Roem. & Schult. Syst. ii. 508 = Deyeuxia varia.
humilis, Roem. & Schult. l. c. 510 = Calamagrostis lanceolata, tenella.
hyperborea, Trin. ex Steud. Syn. Pl. Gram. 236 (= *Dupontia psilothantha*).
indica, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 7.—Malaya.
intermedia, C. C. Gmel. Fl. Bad. i. 266 = Calamagrostis epigeios.
isiaca, Delile, Fl. Aeg Illustr. 82 = Phragmites communis.
junciformis, Poir. Encyc. Suppl. iv. 705 = Deyeuxia junciformis.
Kakao, Steud. Syn. Pl. Gram. 194 = conspicua?
Karka, *Retz. Obs.* iv. 21; *Roxb. Hort. Beng.* 8; *Fl. Ind.* i. 348.—Ind. or.
Kila, Spreng. ex Steud. Nom. ed. II. i. 144 = Gynerium argenteum.
laevis, Steud. Nom. ed. II. i. 144 = Poa laevis.
Langsdorfii, Link, Enum. Hort. Berol. i. 74 = Deyeuxia purpurea.
lapponica, Wahlenb. Fl. Lapp. 27. t. 1 = Deyeuxia lapponica.
latifolia, Salisb. Prod. 24 = A. Donax.
laxa, Roem. & Schult. Syst. ii. 502 = Calamagrostis littorea.
Leersii, Oett. Syst. Verz. Oberlaus. 17 = Calamagrostis lanceolata.
Leersii, J. Koch, Handb. 49 = Stipa Calamagrostis.
levis, Poir. Encyc. Suppl. iv. 707 = Poa laevis.
ligulata, *Poir. l. c.* 706 = Deyeuxia ligulata.
lithuanica, Schult. Mant. ii. 602 = Calamagrostis lanceolata.
litorea, Schrad. Fl. Germ. 212 = Calamagrostis littorea.
littoralis, Steud. Nom. ed. II. i. 144 = Ammophila arundinacea.
madagascariensis, *Kunth, Rev. Gram.* i. 273. t. 48.—Madag.
maritima, *Agardh, in Flora*, vii. (1824) 561.—Scandin.
mauritanica, Poir. Voy. Barb. ii. 104; Desf. Fl. Atlant. i. 106 = Pliniana.
maxima, Forsk. Fl. Aegypt. Arab. 24 = Phragmites communis.
maxima, Lour. Fl. Cochinch. 58 = Bambusa maxima.
micrantha, Lam. Illustr. i. 196 = Pliniana.
minutiflora, *Brongn. in Duperr. Voy. Bot.* 29.—Ins. Molucc.
mitis, Lour. Fl. Cochinch. 57 = Bambusa mitis.
montana, [Clairv.] Man. Herb. 15 = Deyeuxia montana.
montana, Roem. & Schult. Syst. ii. 508 = Deyeuxia sylvatica.
multiplex, Lour. Fl. Cochinch. 58 = Gigantochloa verticillata.

ARUNDO :—
Naga, Koen. ex Steud. Nom. ed. II. i. 144 = Phragmites australis.
Naya, Koen. ex Steud. l. c. ii. 321 = praec.
neglecta, Ehrh. Beitr. vi. 84, 137 = Deyeuxia neglecta.
nemorosa, Schult. Oestr. Fl. ed. II. i. 187 = Deyeuxia varia.
nitida, *H. B. & K. Nov. Gen. et Sp.* i. 149. t. 688.—Ecuador.
occidentalis, *Sieber, ex Schult. Mant.* ii. 289.—Ins. Martin.
orientalis, Mill. Gard. Dict. ed. VIII. n. 6 = Saccharum fuscum.
pallens, *Muhl. ex Steud. Nom.* ed. II. i. 144.—Am. bor.
palustris, Salisb. Prod. 24 = Phragmites communis.
penicillata, Labill. Nov. Holl. Pl. i. 26 = Danthonia racemosa.
Phragmites, Linn. Sp. Pl. 81 = Phragmites communis.
picta, Kunze, ex Steud. Nom. ed. II. i. 144 = Poa picta.
pilosa, *Urv. in Mém. Soc. Linn. Par.* iv. (1826) 603.—Ins. Falkland.
piscatoria, *Lour. Fl. Cochinch.* 55.—Cochinch.
planifolia, Poir. Encyc. Suppl. iv. 707 = Deyeuxia planifolia.
plebica, Poir. l. c. = Deyeuxia plebeia.
Pliniana, *Turra, Farset.* 11; *Fl. Ital. Prod.* 63.—Reg. Mediterr.
Plinii, Turra, Farset. 11 = Pliniana.
plumosa, Schult. Mant. 604 = Stipa papposa.
poaeformis, Labill. Nov. Holl. Pl. i. 27. t. 35 = Poa Billardieri.
poaeoides, Poir. Encyc. Suppl. iv. 705 = Poa subuniflora.
Pseudophragmites, Hall. f. in Roem. Arch. i. II. 11 = Calamagrostis littorea.
Pseudophragmites, Lejeune, Fl. Spa, 64 = Phragmites communis.
Pseudophragmites, Schrad. Fl. Germ. 213 = Deyeuxia Halleriana.
Pseudophragmites, Wahlenb. Fl. Lapp. 28 = Calamagrostis lapponica.
pungens, Auct. ex Steud. Nom. ed. II. i. 144 = Phragmites communis.
purpurascens, Schult. Mant. iii. 603 = Deyeuxia purpurascens.
purpurea, Trin. in Spreng. Neue Entdeck. ii. 52 = Deyeuxia purpurea.
pygmaea, *Moretti, ex Pollini, in Brugnat. Giorn. Fis.* ix. (1816) 22.—Italia.
pygmaea, Spreng. Pugill. i. 9 = Trisetum hispanicum.
Quila, Molina, Sagg. Chil. 279 = Chusquea Fernandeziana.
Quila, Poir. Encyc. vi. 274 = Chusquea Quila.
quitensis, Poir. l. c. Suppl. iv. 704 = Muehlenbergia calamagrostidea.
Ragamowskii, Hort.—Cf. Gard. Chron. (1876) II. 847 = Bambusa tessellata.
ramosa, Schult. Oestr. Fl. ed. II. i. 185 = Calamagrostis lanceolata.
recta, Poir. Encyc. Suppl. iv. 706 = Deyeuxia recta.
Reynaudiana, *Kunth, Rev. Gram.* i. 275. t. 49.—Pegu.
rigida, Poir. Encyc. Suppl. iv. 705 = Deyeuxia rigida.
riparia, Salisb. Prod. 24 = Phalaris arundinacea.
riparia, Willd. ex Steud. Nom. ed. II. i. 144 = Calamagrostis lanceolata.
rivularis, Auct. ex Steud. Nom. ed. II. i. 144 = Phragmites communis.
Roxburghii, Kunth, Rev. Gram. i. 79 = A. Karka.
Rugi, *Molina, Sagg. Chil.* 279.—Chili.
Sabuli, *Kit. in Linnaea*, xxxii. 1863 306.—Europ.
saccharoides, Poir. Encyc. Suppl. iv. 705 = Gynerium saccharoides.
sagittata, Pers. Syn. i. 102 = Gynerium saccharoides.
sativa, Lam. Fl. Fr. iii. 616 = A. Donax.
Schleicheriana, Bess. Fl. Galic. i. 69 = Calamagrostis lanceolata.
Schwabii, Schult. Mant. iii. 603 = Gastridium australe.
scriptoria, Linn. Amoen. Ac. iv. 150 = A. Donax.
sedenensis, Loisel. Fl. Gall. 719 = Deyeuxia varia.
Selloana, Schult. Mant. iii. 605 = Gynerium argenteum.
semiannularis, Labill. Nov. Holl. Pl. i. 26. t. 33 = Danthonia semiannularis.
semifarcta, Hort. Ticin. ex Steud. Nom. ed. II. i. 144 = Pliniana.
speciosa, Schrad. Fl. Germ. i. 219. t. 5. f. 8 = Stipa Calamagrostis.

ARUNDO:—
splendida, Schkr. ex Steud. Nom. ed. II. i. 144 = Deyeuxia acutiflora.
stricta, Gilib. Exercit. ii. 542 = Calamagrostis epigeios.
stricta, Hofft. Cat. Kursk. 7 = Deyeuxia neglecta.
stricta, Spreng. Neue Entdeck. i. 247 = Deyeuxia Nuttalliana.
stricta, Timm, in Siemss. Meklenb. Mag. ii. 235, ex Schrad. Fl. Germ. 215 = Deyeuxia stricta.
strigosa, Wahlenb. Fl. Lapp. 29. t. 2 = Deyeuxia strigosa.
subulata, J. Gay, ex Reichb. Fl. Germ. Excurs. 27 = Deyeuxia acutiflora.
sylvatica, Schrad. Fl. Germ. i. 218. t. 4. f. 7 = Deyeuxia sylvatica.
Tabacaria, Lour. Fl. Cochinch. 58 = Bambusa Tabacaria.
tecta, Blanco, Fl. Filip. ed. I. 48 = A. Karka.
tecta, Walt. Fl. Carol. 81 = Arundinaria tecta.
tenax, Vahl, Symb. Bot. ii. 25 = Ampelodesmos tenax.
tenella, Schrad. Fl. Germ. 220 = Calamagrostis tenella.
tenella, Spreng. Pugill. ii. 6 = Muehlenbergia tenuifolia.
tenuifolia, Poir. Encyc. Suppl. iv. 704 = Muehlenbergia tenuifolia.
tolucensis, Poir. l. c. 705 = Deyeuxia tolucensis.
triodioides, Trin. Sp. Gram. t. 351 = Schedonorus littoralis.
valdiviana, *Molina, Saggio Chil.* 279.—Chili.
Vallatoria, S. F. Gray, Nat. Arr. Brit. Pl. ii. 128 = Phragmites communis.
varia, Schrad. Fl. Germ. 216 = Deyeuxia acutiflora.
versicolor, Mill. Gard. Dict. ed. VIII. n. 3 = A. Donax.
villosa, Auct. ex Schult. Mant. iii. 287 = Calamagrostis sylvatica.
villosissima, *Forsk. ex Steud. Nom.* ed. II. i. 145.—Reg. Byzant.
vilnensis, Schult. Mant. iii. 602 = Calamagrostis lanceolata.
viridescens, Steud. Syn. Pl. Gram. 192 = seq.
viridiflavescens, Poir. Encyc. vi. 271 = Deycuxia viridiflavescens.
vulgaris, Lam. Fl. Fr. iii. 615 = Phragmites communis.
vulnerans, Gilib. Exercit. ii. 541 = Phragmites communis.
Webbiana, *Steud. Syn. Pl. Gram.* 194.—Afr. austr.
Wilhelmsii, Ledeb. in Mém. Acad. Pétersb. vi. (1814) 593 = Trisetum rigidum.
Zollingeri, *Buese, in Miq. Pl. Jungh.* 243.—Java.

ARUNGANA, Pers. Syn. ii. 91 (1807) = **Haronga**, Thou. [1809] (Hyperic.).
crenata, Pers. l. c. = H. crenata.
mollusca, Pers. l. c. = H. mollusca.
paniculata, Pers. l. c. = H. madagascariensis.

ARUNIA, Pers. Syst. 251 (1797), *sphalm.* = **Brunia**, Linn.

ARVERSIA, Cambess. in St. Hil. Fl. Bras. Mer. ii. 184. t. 112 (1829) = **Polycarpon**, Linn. (Caryophyll.).
depressa, Klotzsch, in Peters, Reise Mossamb. Bot. i. 140 = P. depressum.
frankenoides, Cambess. in St. Hil. Fl. Bras. Mer. ii. 184 = P. depressum.
Loefflingii, Walp. Rep. i. 263 = P. depressum.

ARVIELA, Salisb. Gen. Pl. Fragm. 135 (1866) = **Zephyranthes**, Herb. (Amaryll.).
tubispatha, Salisb. l. c. = Z. tubispatha.

ARYTERA, Blume, Rumphia, iii. 170 (1847) = **Ratonia**, DC. (Sapind.).
angustifolia, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 552.—Hab.?
arcuata, Radlk. l. c. 554.—N. Caled.
brachyphylla, Radlk. l. c. 552.—N. Guin.
Brackenridgei, Radlk. l. c. 555.—Ins. Fiji.
chartacea, Radlk. l. c. 553.—Hab.?
distylis, Radlk. l. c.—Austral.
divaricata, F. Muell. in Trans. Phil. Inst. Victoria, iii. (1859) 25 = Nephelium divaricatum.
foveolata, F. Muell. l. c. = Nephelium foveolatum.
gigantosperma, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 674.—Sumatra.
Karang, Miq. Fl. Ind. Bat. Suppl. i. 510.—Sumatra.

ARYTERA:—
Leichhardtii, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 553.—Austral.
lepidota, Radlk. l. c. 555.—N. Caled.
littoralis, Blume, Rumphia, iii. 170.—Sumatra.
macrocarpa, Miq. Fl. Ind. Bat. Suppl. i. 510 = Triomma malaccensis.
microphylla, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 553.—Austral.
montana, Blume, Rumphia, iii. 171.—Malaya.
morocarpa, Miq. Fl. Ind. Bat. Suppl. i. 510.—Sumatra.
oligolepis, Radlk. in Sitzb. Math.-Phys. Acad. Muench. ix. (1879) 555.—Ins. Navig.
O'Shanesiana, Radlk. l. c. 554.—Austral.
pachyphylla, Radlk. l. c.—N. Caled.
rufescens, Radlk. l. c. 552.—Ins. Philipp.
semiglauca, F. Muell. in Trans. Phil. Inst. Victoria, iii. (1859) 25 = Cupania semiglauca.
Silaka, Miq. Fl. Ind. Bat. Suppl. i. 510.—Sumatra.

ASACARA, Rafin. Neogenyt. 2 (1825) = **Gleditschia**, Linn. (Legum.).
aquatica, Rafin. l. c. = G. monosperma.

ASAEMIA, Harv. in Harv. & Sond. Fl. Cap. iii. 187 (1864), in syn. *COMPOSITAE*, Benth. & Hook. f. ii. 433.
axillaris, *Harv. l. c.*—Afr. austr.

ASAGRAEA, Baill. Adansonia, ix. (1870) 232. *LEGUMINOSAE*.
spinosa, *Baill. l. c.* 233.—Am. bor. occ.

ASAGRAEA, Lindl. Bot. Reg. t. 33 (1839) = **Schoenocaulon**, A. Gray (Liliac.).
caracasana, Ernst, in Journ. Bot. (1871) 91 = S. officinale.
caricifolia, Kunth, Enum. Pl. iv. 666 = S. officinale.
officinalis, Lindl. in Bot. Reg. (1839) t. 33 = S. officinale.
tenuifolia, Kunth, Enum. Pl. iv. 700 = S. officinale.

ASAPHES, DC. Prod. ii. 90 (1825) = **Toddalia**, Juss. (Rutac.).
undulata, DC. l. c. = T. lanceolata.

ASAPHES, Spreng. Syst. iv. Cur. Post. 222 (1827) = **Morina**, Tourn. (Dipsac.).
nepalensis, Spreng. l. c. 232 = M. persica.

ASARCA, Poepp. ex Lindl. in Quart. Journ. R. Instit. i. (1827) 52 = **Chloraea**, Lindl. (Orchid.).
acutiflora, Poepp. & Endl. Nov. Gen. et Sp. ii. 14. t. 120, A.—Chili.
acutifolia, Steud. Nom. ed. II. i. 145, sphalm. = praec.
alaris, Lindl. Gen. et Sp. Orch. 408 = C. alaris.
arauacana, Phil. in Linnaea, xxix. (1857–58) 56.—Chili.
aurantiaca, Lindl. in Hook. Journ. Bot. i. (1834) 4 = C. aurantiaca.
Berteri, Reichb. f. in Linnaea, xix. (1847) 377.—Chili.
bidentata, Poepp. & Endl. Nov. Gen. et Sp. ii. 15. t. 121.—Chili.
brachychila, Phil. in Anal. Univ. Chil. (1865) II. 332.—Chili.
chrysostachya, Phil. in Linnaea, xxix. (1857–58) 55.—Chili.
Commersonii, Hook. f. Fl. Antarct. ii. 351 = C. Commersonii.
glandulifera, Poepp. & Endl. Nov. Gen. et Sp. ii. 14. t. 120. B = C. volucris.
Kingii, Hook. f. Fl. Antarct. ii. 351.—Reg. Magell.
leucantha, Poepp. & Endl. Nov. Gen. et Sp. ii. 13. t. 119.—Chili.
maculata, Steud. Nom. ed. II. i. 145 = seq.
maculosa, Poepp. Fragm. Syn. Pl. Chil. 20 = C. galeata.
odoratissima, Poepp. & Endl. Nov. Gen. et Sp. ii. 13. t. 118.—Chili.
parviflora, Poepp. Fragm. Syn. Pl. Chil. 20.—Chili.
platyantha, Reichb. f. in Linnaea, xxii. (1849) 862.—Chili.
sinuata, Lindl. Gen. et Sp. Orch. 408 = C. longibracteata.
speciosa. Lindl. in Quart. Journ. R. Instit. i. (1827) 52 = C. Pavonii.

ASARCA:—
spectabilis, Phil. in Linnaea, xxix. (1857–58) 55 = Chloraea spectabilis.
sulphurea, Phil. l. c. 57.—Chili.
verrucosa, A. Rich. ex C. Gay, Fl. Chil. v. 465. t. 66 = C. verrucosa.

ASARINA, Tourn. ex Mill. Gard. Dict. ed. VIII. (1768) = **Antirrhinum**, Tourn. (Scrophul.).
cordifolia, Moench, Meth. Suppl. 172 = Ant. Asarina.
erecta, Mill. Gard. Dict. ed. VIII. n. 2 = 'Penstemon,' Mitch.?—Am. bor.
Lobelii, Quer, Fl. Esp. iii. 115. t. 30 = Ant. Asarina.
procumbens, Mill. Gard. Dict. ed. VIII. n. 1 = Ant. Asarina.

ASARUM, [Tourn.] Linn. Syst. ed. I (1735). *ARISTOLOCHIACEAE*, Benth. & Hook. f. iii. 122.
HETEROTROPA, Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 314. t. 10.
HEXASTYLIS, Rafin. Neogenyt. 3 (1825).
albivenium, *Regel, Gartenfl.* xiii. (1864) 295. t. 440.—Japon.
arifolium, *Michx. Fl. Bor. Am.* i. 279.—Am. bor.
Blumei, *Duch. in DC. Prod.* xv. 1. 427.—Japon.
canadense, *Linn. Sp. Pl.* 442.—Am. bor.
caulescens, *Maxim. in Bull. Acad. Pétersb.* xvii. (1872) 162.—Japon.
carolinianum, Walt. Fl. Carol. 143 = canadense.
caudatum, *Lindl. Bot. Reg.* sub t. 1399.—Calif.
caudigerum, *Hance, in Journ. Bot.* xix. (1881) 142.—China.
elegans, *Duch. in DC. Prod.* xv. 1. 426.—Japon.
europaeum, *Linn. Sp. Pl.* 442.—Europ.; As. Min.; Sibir.
furcatum, *Rafin. New Fl. Am.* ii. 20.—Am. bor.
grandifolium, *Lodd. Cab. ex Loud. Hort. Brit.* 187.—Am. bor.
grandiflorum, *Klotzsch, in Monatsb. Acad. Berl.* (1859) 586.—Am. bor.
grandiflorum, Steud. Nom. ed. II. i. 145 = grandifolium.
Hartwegii, *S. Wats. in Proc. Am. Acad.* x. (1875) 346.—Calif.
heterotropoides, F. Schmidt, Reisen Amurl. 171 = Sieboldii.
himalaycum, *Hook. f. & Thoms. ex Klotzsch, in Monatsb. Akad. Berl.* (1859) 585.—Reg. Himal.
Hookeri, Field. & Gardn. Sert. t. 32 = caudatum.
Hypocistis, Linn. Sp. Pl. 442 = Cytinus Hypocistis.
ibericum, Stev. ex Ledeb. Fl. Ross. iii. 553 = europaeum.
latifolium, Salisb. Prod. 344 = canadense.
Lemmoni, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 294.—Calif.
leucodictyon, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 135 = albivenium.
lucidum, Salisb. Prod. 344 = europaeum.
maculosum, Stokes, Bot. Mat. Med. iii. 7 = virginicum.
medium, Rafin. New Fl. Am. ii. 20 = arifolium?
officinale, Moench, Meth. 292 = europaeum.
parviflorum, Regel, Ind. Sem. Hort. Petrop. (1866) 89 = elegans.
parvifolium, Rafin. New Fl. Am. ii. 20 = virginicum.
renifolium, Stokes, Bot. Mat. Med. iii. 5 = europacum.
reniforme, Gilib. Exercit. ii. 424 = europaeum.
rotundifolium, Rafin. New Fl. Am. ii. 21 = virginicum.
Sieboldii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 134.—Japon.
Thunbergii, *A. Br. Ind. Sem. Hort. Berol.* (1861) *App.* 13.—Japon.
variegatum, *A. Br. & Bouché, l. c.* 12.—Japon.
villosum, Muhl. ex Duch. in DC. Prod. xv. 1. 424 = canadense.
virginianum, Crantz, Inst. i. 134 = seq.
virginicum, *Linn. Sp. Pl.* 442.—Am. bor.
virginicum, Thunb. Fl. Jap. 190 = albivenium.
virginicum, Walt. Fl. Carol. 143 = arifolium.
viridiflorum, Regel, Ind. Sem. Hort. Petrop. (1869) 83 = variegatum.

ASCALEA, Hill, Veg. Syst. iv. 14 (1762) = **Carduus**, Linn. (Compos.).
lanceata, Hill, Hort. Kew. 60 = Cnicus lanceolatus.
nutans, Hill, l. c. = Card. nutans.

ASCALONICUM, Renault, Fl. Dép. Orne, 33 (1800) = **Allium**, Tourn. (Liliac.).
sativum, Renault, l. c. = All. Ascalonicum.

ASCALONICUM :—
Schoenoprasum, Renault, Fl. Dép. Orne, 33 (1800) = Allium Schoenoprasum.
Scorodoprasum, Renault, l. c. = All. Scorodoprasum.

ASCANICA, Crantz, Inst. ii. 313 (1766) = **Patagonula**, Linn. (Borag.).
patagonica, Crantz, l. c. = P. americana.

ASCARICIDA, Cass. in Dict. Sc. Nat. iii. Suppl. 38 (1816) = **Vernonia**, Schreb. (Compos.).
adoënsis, Steetz, in Peters, Reise Mossamb. Bot. 358 = V. adoënsis.
anthelminthica, Sweet, Hort. Brit. ed. I. 219 = V. anthelminthica.
Buchingeri, Steetz, in Peters, Reise Mossamb. Bot. 359 = V. Buchingeri.
indica, Cass. in Dict. Sc. Nat. iii. Suppl. 38 = V. anthelminthica.
Kotschyana, Steetz, in Peters, Reise Mossamb. Bot. 359 = V. Kotschyana.
mossambiquensis, Steetz, l. c. 358 = V. mossambiquensis.
Richardi, Steetz, l. c. = V. adoënsis.
triplinervia, G. Don, in Loud. Hort. Brit. 333 = Baccharis triplinervis.

ASCARIDIA, Reichb. Nom. 99 (1841) = praec.

ASCARINA, Forst. Char. Gen. 117. t. 59 (1776). *CHLORANTHACEAE*, Benth. & Hook. f. iii. 134.
lanceolata, *Hook. f. in Journ. Linn. Soc.* i. (1856) 127.—Ins. Paeif.
lucida, *Hook. f. Fl. N. Zel.* i. 228.—N. Zel.
lurida, Solms, in DC. Prod. xvi. 1. 478 = rubricaulis.
polystachya, *Forst. Char. Gen.* 59, 117. t. 59.—Ins. Societ.
rubra, Poir. Encyc. ix. 475 = Myrica sapida.
rubricaulis, *Solms, in DC. Prod.* xvi. 1. 478.—N. Caled.
serrata, Blume, Enum. Pl. Jav. i. 79 = Chloranthus brachystachyus.

ASCHAMIA, Salisb. Gen. Pl. Fragm. 134 (1866) = **Hippeastrum**, Herb. (Amaryll.).
equestris, Salisb. l. c. = H. equestre.
psittacina, Salisb. l. c. = H. psittacinum.
Reginae, Salisb. l. c. = H. Reginae.

ASCHENBORNIA, Schau. in Linnaea, xix. (1847) 716. *COMPOSITAE*, Benth. & Hook. f. ii. 241.
heteropoda, *Schau. l. c.*—Mexic.

ASCHENFELDTIA, F. Muell. ex Meissn. in DC. Prod. xiv. 515 (1856) = **Pimelea**, Banks & Soland. (Thymel.).
pimeleoides, F. Muell. ex Meissn. l. c. = P. microcephala.

ASCIADIUM, Griseb. Cat. Pl. Cub. 118 (1866). *UMBELLIFERAE*.
coronopifolium, *Griseb. l. c.*—Cuba.

ASCIUM, Schreb. Gen. 358 (1789) = **Norantea**, Aubl. (Ternstr.).
adamantium, Steud. Nom. ed. II. i. 145 = N. adamantium.
anomalum, Sprcng. Syst. iv. Cur. Post. 207 = N. anomala.
Aubletii, Spreng. Syst. ii. 599 = N. guianensis.
Berterii, Sprcng. l. c. = N. Berterii.
goyazense, Steud. Nom. ed. II. i. 145 = N. goyazensis.
japurense, Steud. l. c. = N. japurensis.
Norantea, Racusch. Nom. ed. III. 156 = N. guianensis.
paraënse, Steud. Nom. ed. II. i. 145 = N. paraënsis.
Selloi, Sprcng. Syst. ii. 599 = N. cuneifolia, brasiliensis.
violaceum, Vahl, Ecl. Am. ii. 41. t. 20 = N. guianensis.

ASCLEIA, Rafin. Sylva Tellur. 53 (1838) (= **Lycium** ? Linn.) (Solan.).
mexicana, Rafin. l. c. (= L. capsulare).—Mexic.

ASCLEPIAS, Linn. Gen. ed. I. 64 (1737). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 754.
ACEROTIS, Rafin. New Fl. Am. i 49 (1836).

ASCLEPIAS:—
CRASSA, Rupp. Fl. Jen. ed. Hall. 26 (1745).
DASSOVIA, Neck. Elem. i. 257 (1790).
OLIGORON, Rafin. New Fl. Am. iv. 60 (1836).
OTANEMA, Rafin. l. c. 61 (1836).
OTARIA, Kunth, in H. B. & K. Nov. Gen. et Sp. iii. 192 (1818).
acanthocarpus, Gmel. ex Decne. in DC. Prod. viii. 571 = Gonolobus hirsutus.
aceratoides, *M. Curt. in Am. Journ. Sc.* Scr. II. vii. (1849) 407.—Am. bor.
acida, Roxb. Hort. Beng. 20; Fl. Ind. ii. 31 = Sarcostemma brevistigma.
acornuta, *Kellogg, in Proc. Calif. Acad.* i. (1865) 54. —Calif.
acuminata, Pursh, Fl. Am. Sept. i. 182 = rubra.
acuminata, Roxb. Hort. Beng. 21; Fl. Ind. ii. 55 = Gymnema acuminatum.
acuminata, Willd. ex Roem. & Schult. Syst. vi. 82, 145 = Klaprothiana.
aequicornu, *Fourn. in Mart. Fl. Bras.* vi. IV. 201.—Bras.
ageratoides, *M. Curt. in Am. Journ. Sc.* Ser. II. vii. (1849) 407.—Am. centr.
alba, Cav. Ic. iv. 5 = Sarcostemma Dombeyanum ?
alba, Mill. Gard. Dict. ed. VIII. n. 1; Lam. Fl. Fr. ii. 301 = Vincetoxicum officinale.
albiflora, *Rafin. New Fl. Am.* iv. 62.—Am. bor.
albo-aurantiaca, Hort. ex Jaeq. Fragm. 86 = nivea.
alexicaca, Jacq. Coll. ii. 101 = Hoya pendula.
alexitoxica, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = Vincetoxicum officinale.
alpina, Vell. Fl. Flum. 116; iii. t. 57 = Oxypetalum megapotamicum.
alticola, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 371.—Mexic.
altissima, *Jacq. Enum. Pl. Carib.* 17.—Ind. occ.
amoena, Brongn. in Ann. Sc. Nat. Sér. I. xxiv. (1831) 275. t. 13 = incarnata.
amoena, Linn. Sp. Pl. 214 = purpurascens.
amplexicaulis, *Michx. Fl. Bor. Am.* i. 115.—Am. bor.
angulosa, *Regel, Cat. Pl. Hort. Aksakov.* 13.—Hab. ?
angustifolia, Ell. Sketch, i. 385 = Michauxii.
angustifolia, Hort. Berol. ex Roem. & Schult. Syst. vi. 70 = praec.
angustifolia, *Schweig. Enum. Pl. Hort. Bot. Regiomont.* (1812) 13.—Mexic.
angustissima, *Anderss. Om Galap. Oarnes Veget.* 196. —Ins. Galapag.
annularia, Roxb. Hort. Beng. 20; Fl. Ind. ii. 37 = Holostemma Rheedianum.
aphylla, Boj. ex Decne. in DC. Prod. viii. 546 = Decanema Bojerianum.
aphylla, Forsk. Fl. Aegypt. Arab. 50 = Sarcostemma Forskohlianum.
aphylla, Roxb. ex Wight, Contrib. 60 = Sarcostemma brevistigma.
aphylla, Thunb. Prod. Pl. Cap. 47 = Sarcostemma aphyllum.
Apocinum, Gater. Desc. Pl. Montaub. 58 = syriaca.
arborea, Salisb. Prod. 150 = seq.
arborescens, Linn. Mant. 216 = Gomphocarpus arborescens.
arenaria, *Torr. in Bot. Mex. Bound.* 162.—Am. bor. occ.
argentata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 6.—Malaya.
armata, *Spreng. Syst.* i. 849.—N. Granat.
arvensis, Vell. Fl. Flum. 117; iii. t. 63 = Oxypetalum campestre.
asthmatica, Linn. f. Suppl. 171 = Tylophora asthmatica.
aurantiaca, Salisb. Prod. 150 = curassavica.
auriculata, *H. B. & K. Nov. Gen. et Sp.* iii. 191. t. 228. —Mexic.
axillaris, *Vell. Fl. Flum.* 115; iii. t. 51.—Bras.
barjoniaefolia, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 372.—Bolivia.
bicolor, Moench, Meth. 717 = curassavica.
bidentata, *Hemsl. Biol. Centr. Am. Bot.* ii. 322.—Mexic.
Blanchetii, *Fourn. in Mart. Fl. Bras.* vi. IV. 201.—Bras.
brachystephana, *Engelm. ex Torr. in Bot. Mex. Bound.* 163.—Am. bor. occ.
boliviensis, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1883) 381.—Bolivia.
bracteolata, *Fourn. in Mart. Fl. Bras.* vi. IV. 200.—Bras.

ASCLEPIAS:—
brevicornu, Schecle, in Linnaea, xxi. (1848) 756 = Asclepiodora decumbens.
Bridgesii, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 383.—Bolivia.
campestris, *Decne. in DC. Prod.* viii. 566.—Bras.
campestris, Vell. Fl. Flum. iii. t. 63 = Oxypetalum campestre.
cancellata, *Burm. f. Fl. Cap. Prod.* 7.—Afr. austr.
candida, *Vell. Fl. Flum.* 118; iii. t. 65.—Bras.
canescens, Willd. in Ges. Naturf. Fr. Neu. Schr. iii. (1801) 418 = Cynanchum glaucum.
canina, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 6.—Malaya.
carnosa, Blanco, Fl. Filip. ed. I. 208 = Hoya multiflora.
carnosa, Linn. f. Suppl. 170 = Hoya carnosa.
ciliata, Murr. ex Decne. in DC. Prod. viii. 572 = Xysmalobium undulatum.
cinerea, *Walt. Fl. Carol.* 105.—Am. bor.
cirrosa, *Ehrenb. ex Schweinf. Beitr. Fl. Aethiop.* 132.—Afr. trop.
citrifolia, Hook. & Arn. in Hook. Journ. Bot. i. (1834) 295 = campestris.
citrifolia, Jacq. Coll. ii. 290 = variegata.
clausa, Jacq. Enum. Pl. Carib. 17 = Sarcostemma glaucum.
clusiaefolia, *Horsf. ex Roem. & Schult. Syst.* vi. 82.—Java.
communis, Vell. Fl. Flum. 115; iii. t. 53 = Oxypetalum Banksii.
compressa, Moench, Meth. 717 = purpurascens.
connivens, Baldw. in Ell. Sketch, i. 320 = Asclepiodora viridis.
consanguinea, *Kunze, in Linnaea*, xx. (1847) 26.—Hab. ?
contorta, Forsk. Fl. Aegypt. Arab. p. cviii. n. 188 = Sarcostemma Forskohlianum.
convolvulacea, Heyne, ex Decne. in DC. Prod. viii. 572 = Holostemma Rheedianum.
convolvulacea, Willd. Sp. Pl. i. 1269 = Daemia angolensis.
cordata, Burm. f. Fl. Ind. 72 = Pergularia minor.
cordata, Forsk. Fl. Aegypt. Arab. 49 = Daemia cordata.
cordata, Walt. Fl. Carol. 105 = rubra ?
cordata, Vell. Fl. Flum. 116; iii. t. 56 = Roulinia fluminensis.
Cornuti, Decne. in DC. Prod. viii. 564 = syriaca.
Coulteri, *A. Gray, in Proc. Am. Acad.* xii. (1877) 71. —Mexic.
crassifolia, Hort. Par. ex Decne. in DC. Prod. viii. 572 = Gomphocarpus arborescens.
crassifolia, Linn. ex Decne. l. c. = Gomphocarpus fruticosus.
crispa, Berg. Desc. Pl. Cap. 75 = Gomphocarpus crispus.
cryptoceras, *S. Wats. in Bot. King, Exp.* 283. t. 28.—Am. bor. occ.
cubensis, *Wender. in Bot. Zeit.* i. (1843) 830.—Cuba.
curassavica, Griseb. in Goett. Abh. xix. (1874) 159 = campestris.
curassavica, *Linn. Sp. Pl.* 215; *Willd. Sp. Pl.* i. 1266. —Am. austr.
curassavica, Lour. Fl. Cochinch. 170 = Toxocarpus Wightianus.
Curtissii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 85. —Am. bor.
Curupi, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 384.—Paraguay.
cutca, *Rafin. in Am. Monthly Mag.* (1817) 359.—Am. bor.
cynanchoides, *Humb. & Bonpl. ex Roem. & Schult. Syst.* vi. 86.—Venezuela.
Daemia, Blanco, Fl. Filip. ed. I. 208 = Gymnema tingens.
Daemia, Forsk. Fl. Aegypt. Arab. 51 = Daemia extensa.
dasypus, *Rafin. Atl. Journ.* 152.—Am. bor.
davurica, Willd. Sp. Pl. i. 1272 = Cynanchum boreum.
debilis, *Michx. Fl. Am. Bor.* i. 116.—Am. bor.
decumbens, Linn. Sp. Pl. 216 = tuberosa.
densa, *Jacq. Enum. Pl. Carib.* 17.—Ind. occ.
dichotoma, Rottl. in Wall. Cat. n. 8200 = Tylophora tenuis.
Douglasii, Hook. Fl. Bor. Am. ii. 53. t. 97 = speciosa.
echinata, Herb. Madr. ex Hook. f. Fl. Brit. Ind. iv. 35 = Marsdenia tenacissima.
echinata, Roxb. Hort. Beng. 20; Fl. Ind. ii. 44 = Daemia extensa.

ASCLEPIAS:—
echinata, Decne. in DC. Prod. viii. 544 = Daemia angolensis.
ecornuta, Kellogg, ex A. Gray, Syn. Fl. N. Am. ii. 1. 100 = Gomphocarpus cordifolius.
elata, Benth. Pl. Hartw. 290 = glaucescens.
elegans, *Vell. Fl. Flum.* 116; iii. t. 55.—Bras.
eriocarpa, *Benth. Pl. Hartw.* 323.—Calif.
eriocarpa, Torr. in Pacif. Rail. Rep. iv. 128 = vestita.
erosa, *Torr. in Bot. Mex. Bound.* 162.—Calif.
euphorbiaefolia, *Engelm. in Proc. Am. Acad.* xvi. (1881) 104.—Mexic.
exaltata, Linn. Amoen. Acad. iii. 404 = syriaca.
exaltata, Link, ex Spreng. Syst. i. 848 = Asclepiodora viridis.
exaltata, Muhl. ex Bigel. Fl. Bost. 102 = phytolaccoides.
fascicularis, Decne. in DC. Prod. viii. 509 = mexicana.
fasciculata, Hemsl. Biol. Centr. Am. Bot. ii. 324, sphalm. = pracc.
Feayi, *Chapm. ex A. Gray, in Proc. Am. Acad.* xii. (1876) 72.—Am. bor. trop.
ferruginea, *Humb. & Bonpl. ex Roem. & Schult. Syst.* vi. 83.—Venezuela.
filiformis, Jacq. Enum. Pl. Carib. 17 = Sarcostemma Jacquini.
filiformis, Linn. f. Suppl. 169 (= *Lagarinthus tenuis*).—Ind. occ.; Afr. austr.
floridana, Lam. Encyc. i. 284 = Gomphocarpus longifolia.
foetida, Cav. Ic. ii. 45. t. 158 = Roulinia Jacquini.
Forskohlei, *Roem. & Schult. Syst.* vi. 85.—Arab.
fragrans, *Decne. in DC. Prod.* viii. 571.—Guatemala.
fragrans, *Rafin. in Med. Repos. N. York*, v. (1808) 354.—Am. bor.
Frémontii, *Torr. in Pacif. Rail. Rep.* vi. 87.—Calif.
fruticosa, Linn. Sp. Pl. 216 = Gomphocarpus fruticosus.
fusca, Lour. Fl. Cochinch. 170 = Vincetoxicum medium.
fuscata, Willd. Enum. Hort. Berol. Suppl. 13 = praec.
Galeotti, *Fourn. in Bull. Soc. Bot. Fr.* xiv. (1867) 250.—Mexic.
galioides, H. B. & K. Nov. Gen. et Sp. iii. 148 = verticillata.
geminata, Roxb. Hort. Beng. 20; Fl. Ind. ii. 45 = Gymnema sylvestre.
gigantea, Linn. Sp. Pl. 214 = Calotropis procera.
gigantea, Willd. Sp. Pl. i. 1264 = Calotropis gigantea.
glabra, Forsk. Fl. Aegypt. Arab. 51 = Doemia glabra.
glabra, Mill. Gard. Dict. ed. VIII. n. 12 = Gomphocarpus fruticosus.
glaucescens, *H. B. & K. Nov. Gen. et Sp.* iii. 290. t. 227.—N. Mexic.
globosa, Stokes, Bot. Mat. Med. ii. 4 = syriaca.
gonocarpos, J. F. Gmel. Syst. 446 = Gonolobus carolinensis.
graminea, *Lam. Encyc.* i. 283.—Ind. or.
grandiflora, Linn. f. Suppl. 170 = Gomphocarpus grandiflorus.
grandiflora, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 379.—Mexic.
grandifolia, Bertol. in Mem. Acc. Sc. Bolog. iii. (1851) 189 (Misc. Bot. xii. 47) = syriaca.
Greeniana, *Nutt. ex Loud. Hort. Brit. Suppl.* iii. 493.—Am. bor.
Hallii, *A. Gray, in Proc. Am. Acad.* xii. (1877) 69.—Colorado.
hastata, Bunge, in Mém. Sav. Etr. Acad. Pétersb. ii. (1835) 117 = Cynanchum Bungei.
herbacea, Roxb. Hort. Beng. 21; Fl. Ind. ii. 50 = Calotropis Acia.
hispida, Vell. Fl. Flum. 115; iii. t. 52 = Ditassa Guillemiana.
humistrata, Walt. Fl. Carol. 105 = amplexicaulis.
hybrida, Michx. Fl. Am. Bor. i. 115 = variegata.
illinoënsis, Michx. ex Steud. Nom. ed. II. i. 146 = syriaca.
incarnata, *Linn. Sp. Pl.* 215.—Am. bor.
incarnata, Walt. Fl. Carol. 106 = Gomphocarpus longifolius.
involucrata, *Engelm. ex Torr. in Bot. Mex. Bound.* 163.—N. Mexic.
Jacquiniana, Spreng. Syst. iv. Cur. Post. 341 = Sarcostemma Jacquini.
Jamesii, *Torr. Bot. Mex. Bound.* 162.—Am. bor.
javanica, Burm. ex Decne. in DC. Prod. viii. 628 = Leptadenia reticulata.

ASCLEPIAS:—
Klaprothiana, *Roem. & Schult. Syst.* vi. 82.—Am. bor.
lactifera, Linn. Sp. Pl. 216 = Gymnema lactiferum.
lanceolata, Ives, in Am. Journ. Sc. i. (1819) 252 = Gomphocarpus viridiflorus.
lanceolata, Walt. Fl. Carol. 105 = paupercula.
lancifolia, Steud. Nom. ed. I. 77 = rubra.
laniflora, Forsk. Fl. Aegypt. Arab. 51 = Kanahia laniflora.
Langsdorffii, *Fourn. in Mart. Fl. Bras.* vi. iv. 203.—Bras.
lanuginosa, *H. B. & K. Nov. Gen. et Sp.* iii. 193.—Mexic.
lanuginosa, Nutt. Gen. Am. i. 168 (= *Acerates lanuginosa*).—Am. bor.
latifolia, *Rafin. Atl. Journ.* 146.—Am. bor.
laurifolia, Michx. Fl. Bor. Am. i. 117 = rubra.
laurifolia, Roxb. Hort. Beng. 20; Fl. Ind. ii. 49 = Genianthus laurifolius.
laxiflora, *Decne. in DC. Prod.* viii. 571.—Mexic.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 85.—Arizona.
leucophylla, Engelm. in Am. Natural. ix. (1875) 348 = erosa.
Linaria, *Cav. Ic.* i. 42. t. 57.—Mexic.
Lindheimeri, Engelm. & A. Gray, Pl. Lindh. 250 = longicornu.
Lindleyi, Gill. ex Steud. Nom. ed. II. i. 145, 146 = variegata.
linearifolia, Pav. ex Decne. in DC. Prod. viii. 539 = Sarcostemma Dombeyanum.
linearis, *Scheele, in Linnaea*, xxi. (1848) 758 = verticillata.
linifolia, H. B. & K. Nov. Gen. et Sp. iii. 190 = angustifolia, virgata?
linifolia, *Lag. ex Spreng. Syst.* i. 847.—Mexic.
longicornis, *Kellogg, in Proc. Calif. Acad.* i. ('1855') 65.—Calif.
longicornu, *Benth. in Pl. Hartw.* 24.—Calif.
longiflora, Jacq. Enum. Pl. Carib. 17 (= *Schubertia longiflora*).—Ind. occ.
longifolia, *Michx. Fl. Bor. Am.* i. 116.—Am. bor.
longipetala, Scheele, in Linnaea, xxi. (1848) 757 = Asclepiodora viridis.
longistigma, Roxb. Hort. Beng. 85; Fl. Ind. ii. 46 = Toxocarpus Roxburghii.
lutea, *Rafin. in Med. Repos. N. York*, v. (1808) 361.—Am. bor.
lutea, Mill. Gard. Dict. ed. VIII. n. 3 = Vincetoxicum luteum.
macrantha, *Hochst. ex Decne. in DC. Prod.* viii. 558.—Abyss.
macrophylla, *Humb. & Bonpl. in Roem. & Schult. Syst.* vi. 86.—Cumana.
macrophylla, Nutt. in Journ. Acad. Philad. Ser. II. i. (1847) 180 = mexicana.
macrotis, *Torr. Bot. Mex. Bound.* 164. t. 45.—N. Mexic.
margaritacea, *Hoffmgg. ex Roem. & Schult. Syst.* vi. 86.—Bras.
marginata, Decne. in DC. Prod. viii. 566 = candida.
maritima, Jacq. Enum. Pl. Carib. 17 = Ibatia maritima.
maritima, *Pall. Reise*, iii. 557.—Rossia austr.
maritima, Rafin. ex Decne. in DC. Prod. viii. 567 = incarnata.
Meadii, *Torr. ex A. Gray, Man. Bot. U. St.* ed. II. *Add.* 704.—Am. bor.
media, Hort. ex Steud. Nom. ed. II. i. 146 = Vincetoxicum medium.
megalotis, *Rafin. New Fl. Am.* iv. 61.—Am. bor.
melantha, *Decne. in DC. Prod.* viii. 570.—Mexic.
mellodora, St. Hil. Pl. Rem. Bras. 227 = nervosa.
mexicana, *Cav. Ic.* i. 42. t. 58 = Calif.; Mexic.
Michauxii, *Decne. in DC. Prod.* viii. 569.—Am. bor.
micrantha, *Roxb. Hort. Beng.* 21; *Fl. Ind.* ii. 50.—Ind. or.
microphylla, Heyne, ex Roth, Nov. Sp. Pl. 177 = seq.?
microphylla, Roxb. Hort. Beng. 85; Fl. Ind. ii. 35 = Pentatropis microphylla.
monantha, Vell. Fl. Flum. 115; iii. t. 54 = Oxypetalum monanthum.
montana, Roxb. Hort. Beng. 85; Fl. Ind. ii. 45 = Gymnema tingens.
montevidensis, *Spreng. ex Fourn. in Mart. Fl. Bras.* vi. iv. 202.—Reg. Argent.
mucronata, Thunb. Prod. Pl. Cap. 47 (= *Metaplexis mucronata*).—Afr. austr.

ASCLEPIAS:—
multicaulis, Vell. Fl. Flum. 117; iii. t. 60 = Oxypetalum campestre.
multinervis, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 383.—Paraguay.
muricata, Schum. & Thonn. Beskr. Guin. Pl. 153 = Doemia angolensis.
neglecta, *Hemsl. Biol. Centr. Am. Bot.* ii. 325.—Mexic.
nervosa, *Decne. in DC. Prod.* viii. 568.—Bras.
nigra, Linn. Sp. Pl. 216 = Vincetoxicum nigrum.
nitida, Vell. Fl. Flum. 116; iii. t. 59 = Oxypetalum umbellatum.
nivalis, J. F. Gmel. Syst. 444 = seq.
nivea, Forsk. Fl. Aegypt. Arab. 51 = Forskohlei.
nivea, *Linn. Sp. Pl.* 215.—Ind. occ.
nivea, Sims, Bot. Mag. t. 1181 = phytolaccoides.
nuda, *Thonn. & Schum. Beskr. Guin. Pl.* 155.—Afr. trop.
nummularia, *Torr. Bot. Mex. Bound.* 163. t. 45.—Am. bor.
nutans, Muhl. ex Steud. Nom. ed. II. i. 146 (= *Acerates viridiflorus*).—Am. bor.
Nuttalliana, A. Gray, Man. Bot. U. St. ed. II. 352, 704 = ovalifolia.
Nuttalliana, Torr. in Ann. Lyc. N. York, ii. (1828) 218 (= *Acerates lanuginosa*).—Am. bor.
nyctaginifolia, *A. Gray, in Proc. Am. Acad.* xii. (1877) 69.—Calif.
obovata, *Ell. Sketch*, 321.—Am. bor.
obtusifolia, Hort. Prag. ex Kunze, in Linnaea, xx. (1847) 27 = consanguinea.
obtusifolia, *Michx. Fl. Bor. Am.* i. 115.—Am. bor.
odoratissima, Roxb. Hort. Beng. 20; Fl. Ind. ii. 47 = Pergularia minor.
oenotheroides, Cham. & Schlecht. in Linnaea, v. (1830) 123 = longicornu.
otarioides, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 373.—Mexic.
ovalifolia, *Decne. in DC. Prod.* viii. 567.—Am. bor.
ovata, Auct. ex Steud. Nom. ed. II. i. 146 = Vincetoxicum officinale.
ovata, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1842) 363.—Mexic.
pallida, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 383.—Paraguay.
pallida, Roxb. Hort. Beng. 20; Fl. Ind. ii. 48 = Pergularia pallida.
paniculatum, *Bunge, Enum. Pl. Chin. Bor.* 43.—China.
parasitica, Roxb. Hort. Beng. 20; Fl. Ind. ii. 42 = Hoya parasitica.
parviflora, Ait. Hort. Kew. i. 307 = perennis.
parviflora, Leconte, ex Decne. in DC. Prod. viii. 570 = verticillata.
parviflora, Roem. & Schult. Syst. vi. 82, sphalm. = parvifolia.
parvifolia, Decne. in DC. Prod. viii. 568 = perennis.
parvifolia, *Roem. & Schult. Syst.* iv. 801, ind.—Am. austr.
patula, Auct. ex Decne. in DC. Prod. viii. 536 = Calotropis procera.
paupercula, *Michx. Fl. Bor. Am.* i. 118.—Am. bor.
pedicellata, Walt. Fl. Carol. 106 = Podostigma pubescens.
pellucida, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 381.—Mexic.
pendula, Roxb. Hort. Beng. 85; Fl. Ind. ii. 36 = Hoya pendula.
peregrina, Blanco, Fl. Filip. ed. I. 207 = Sarcolobus carinatus.
perennis, *Walt. Fl. Carol.* 107.—Am. bor.
periplocaefolia, Nutt. Gen. Am. i. 167 = rubra.
phytolaccoides, *Lyon, ex Pursh, Fl. Am. Sept.* i. 180.—Am. bor.
pilosa, Vell. Fl. Flum. 117; iii. t. 62 = Oxypetalum foliosum.
pinifolia, Greene, in Torrey Bot. Club, viii. (1881) 5 = A. Linaria.
planiflora, Jacq. Enum. Pl. Carib. 17 = Gonolobus planiflorus.
plicata, *Heyne, ex Roth, Nov. Pl. Sp.* 176.—Ind. or.
polystachya, Walt. Fl. Carol. 107 = rubra?
pratensis, *Benth. Pl. Hartw.* 45.—Mexic.
Princeps, *Bartl. Ind. Sem. Hort. Gott.* (1839); *ex Kunze, in Linnaea*, xx. (1847) 27.—Am. bor.
procera, Ait. Hort. Kew. i. 305 = Calotropis procera.
procumbens, *Rafin. Fl. Ludov.* 51.—Am. bor.

ASCLEPIAS:—
pseudosarsa, Roxb. Hort. Beng. 20; Fl. Ind. ii. 39 = Hemidesmus indicus.
puberula, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 104.—Mexic.
pubescens, Linn. Mant. 215 = Gomphocarpus arborescens.
pubescens, Moench, Meth. 716 = syriaca.
pubigera, Dum. Fl. Belg. 52 = syriaca.
pulchella, Roxb. Hort. Beng. 21; Fl. Ind. ii. 54 = Raphistemma pulchellum.
pulchella, Salisb. Prod. 150 = perennis.
pulchra, Ehrh. ex Willd. Enum. Hort. Berol. 278 = incarnata.
pulchra, Vell. Fl. Flum. 116; iii. t. 58 = Oxypetalum pulchrum.
purpurascens, Georgi, Beschr. Russ. Reich. iii. iv. 813 = Vincetoxicum purpurascens.
purpurascens, *Linn. Sp. Pl.* 214.—Am. bor.
purpurascens, Walt. Fl. Carol. 105 = obtusifolia.
purpurea, Pall. Reise, iii. 260 (= *Cynoctonum roseum*).—Rossia.
quadrifolia, *Jacq. Obs. Bot.* ii. 8.—Am. bor.
quinquedentata, *A. Gray, in Proc. Am. Acad.* xii. (1877) 71.—N. Mexic.
quinqueflora, *Vell. Fl. Flum.* 118; iii. t. 68.—Bras.
racemosa, Jacq. Enum. Pl. Carib. 17 = Roulinia Jacquini.
racemosa, Roxb. Hort. Beng. 20; Fl. Ind. ii. 32 = Cosmostigma racemosum.
radians, Forsk. Fl. Aegypt. Arab. 49 = Steinheilia radians.
ramosa, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 373.—Bolivia.
repanda, *Forsk. ex Steud. Nom.* ed. I. 76; ed. II. i. 146 = Gomphocarpus crispus.
revoluta, *Rafin. Fl. Ludov.* 51.—Am. bor.
rosea, *H. B. & K. Nov. Gen. et Sp.* iii. 189.—Mexic.
rosea, Roxb. Hort. Beng. 20; Fl. Ind. ii. 40 = Oxystelma esculentum.
rotundifolia, *Mill. Gard. Dict.* ed. VIII. n. 15.—Afr. austr.
rotundifolia, *Rafin. Atl. Journ.* 152.—Hab.?
rubra, *Linn. Sp. Pl.* 217.—Am. bor.
rubra, Sievers, in Pall. N. Nord. Beitr. vii. 329 = = Apocynum Venetum.
rubricaulis, *H. B. & K. Nov. Gen. et Sp.* iii. 148.—Mexic.
salicifolia, *Lodd. Bot. Cab.* t. 272.—Mexic.
scandens, Beauv. Fl. Owar. i. 93. t. 56 = Daemia angolensis.
scandens, Mill. Gard. Dict. ed. VIII. n. 19 = Sarcostemma Swartzianum.
Schaffneri, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 103.—Mexic.
Selloana, *Fourn. in Mart. Fl. Bras.* vi. iv. 202.—Bras.
Serpentaria, *Rafin. Fl. Ludov.* 52.—Am. bor.
setosa, *Benth. Pl. Hartw.* 24.—Mexic.
setosa, Forsk. Fl. Aegypt. Arab. 51 = Gomphocarpus setosus.
sibirica, *Linn. Sp. Pl.* 217.—Sibir.
sibirica, Murr. in Comm. Gotting. (1779) 23 = Vincetoxicum sibiricum.
similis, *Hemsl. Biol. Centr. Am. Bot.* ii. 326.—Mexic.
singularis, *Vell. Fl. Flum.* 117; iii. t. 61.—Bras.
sinuosa, *Burm. f. Fl. Cap. Prod.* 7.—Afr. austr.
speciosa, *Torr. in Ann. Lyc. N. York*, ii. (1828) 218.—Am. bor. occ.
spinosa, *Vell. Fl. Flum.* 118; iii. t. 67.—Bras.
spiraliflora, Stokes, Bot. Mat. Med. ii. 6 = seq.
spiralis, Forsk. Fl. Aegypt. Arab. 49 = Pentatropis spiralis.
stellata, Burm. ex Decne. in DC. Prod. viii. 573 = Hoya multiflora.
stenophylla, *A. Gray, in Proc. Am. Acad.* xii. (1876) 72.—Am. bor.
stipitacea, Forsk. Fl. Aegypt. Arab. 51 = Sarcostemma stipitaceum.
suaveolens, Leconte, ex Decne. in DC. Prod. viii. 569 = Michauxii.
suberosa, Roxb. Hort. Beng. 85; Fl. Ind. ii. 38 = Leptadenia reticulata.
subulata, *Decne. in DC. Prod.* viii. 571.—Calif.
subulata, Forsk. Fl. Aegypt. Arab. p. cviii. n. 193. t. 7 = Caralluma subulata.
Sullivantii, *Engelm. ex A. Gray, Man. Bot.* ed. I. 366.—Am. bor.
Sullivantii, Torr. Bot. Mex. Bound. 163 = glaucescens.

ASCLEPIAS:—
Sussuela, Roxb. Hort. Beng. 20; Fl. Ind. ii. 31 = Hoya imperialis.
syriaca, Blanco, Fl. Filip. ed. I. 204 = cnrassavica.
syriaca, *Linn. Sp. Pl.* ed. II. 214.—Am. bor.
tenacissima, Roxb. Pl. Corom. iii. 35. t. 240 = Marsdenia tenacissima.
tenuis, Herb. Heyne, ex Hook. f. Fl. Brit. Ind. iv. 41 = Tylophora capparidifolia.
tenuissima, Roxb. Hort. Beng. 20; Fl. Ind. ii. 41 = Tylophora tenuissima.
tetrapetala, Dennst. Schlness. Hort. Malab. 35 = Pentatropis microphylla.
tinctoria, Roxb. Hort. Beng. 20; Fl. Ind. ii. 43 = Marsdenia tinctoria.
tingens, Buch.-Ham. ex Roxb. Hort. Beng. 21; Fl. Ind. ii. 53 = Gymnema tingens.
tomentosa, Herb. Madr. ex Hook. f. Fl. Brit. Ind. iv. 35 = Marsdenia tenacissima.
tomentosa, *Ell. Sketch*, i. 320.—Am. bor.
tomentosa, Thonn. ex Decne. in DC. Prod. viii. 628 = Leptadenia lancifolia.
toxicaria, *Salisb. Prod.* 150 = Vincetoxicum officinale.
truncata, *Willd. Enum. Hort. Berol. Suppl.* 13.—Hab.?
truncata, Roxb. ex Decne. in DC. Prod. viii. 529 = Cynanchum alatum.
tuberosa, *Linn. Sp. Pl.* 217.—Am. bor.
tuberosa, Hort. ex Kunze, in Linnaea, xx. (1847) 27.—Hab.?
tuberosa, Roxb. Fl. Ind. ii. 38 = Leptadenia reticulata.
tunicata, Roxb. Hort. Beng. 20; l. c. 35 = Cynanchum pauciflorum.
tunicata, Hort. Calc. ex Hook. f. Fl. Brit. Ind. iv. 45 = Tylophora asthmatica.
umbellata, *Vell. Fl. Flum.* 118; iii. t. 66.—Bras.
uncialis, *Greene, in Coulter, Bot. Gaz.* v. (1880) 64.—Am. bor. occ.
undulata, Jacq. Enum. Pl. Carib. 17 = Xysmalobium undulatum.
undulata, Linn. Sp. Pl. 214 = Pachycarpus crispus.
Vanilla, Rafin. in Am. Monthly Mag. iv. (1818) 39 = quadrifolia.
variegata, *Linn. Sp. Pl.* 215.—Am. bor.
verecunda, Salisb. Prod. 150 = incarnata.
verticillata, *Linn. Sp. Pl.* 217.—Am. bor.
vestita, *Hook. & Arn. in Bot. Beech. Voy.* 363.—Calif.
vicinalis, Mart. Fl. Bras. vi. iv. 235 = Sarcostemma barbatum.
villosa, Balb. in Mem. Acc. Torin. vii. (1802–3) 386 = Lachnostoma Balbisii.
villosa, Mill. Gard. Dict. ed. VIII. n. 14 = Gomphocarpus fruticosus.
villosa, *Rafin. in Med. Repos. N. York*, v. (1808) 353, nomen.—Am. bor.
viminalis, Steud. Nom. ed. II. i. 146 = Sarcostemma viminale.
viminalis, Sw. Prod. Veg. Ind. Occ. 53 = Philibertia viminalis.
Vincetoxicum, Linn. Sp. Pl. 216 = Vincetoxicum officinale.
virgata, Balb. in Cat. Hort. Taur. (1813) 14 = angustifolia.
viridiflora, Rafin. in Med. Repos. N. York, v. (1808) 360; Pursh, Fl. Am. Sept. i. 181 = Gomphocarpus viridiflorus.
viridiflora, Roxb. ex Decne. in DC. Prod. viii. 573 = Hoya viridiflora.
viridis, Walt. Fl. Carol. 107 = Asclepiodora viridis.
viridula, *Chapm. Fl. S. U. St.* 363.—Am. bor. trop.
Virletii, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 378.—Mexic.
volubilis, Herb. Madr. ex Hook. f. Fl. Brit. Ind. iv. 63 = Leptadenia reticulata.
volubilis, Linn. f. Suppl. 170 = Dregea volubilis.
volubilis, Domb. ex Decne. in DC. Prod. viii. 539 = Sarcostemma Dombeyanum.
vomitoria, Koen. ex Hook. f. Fl. Brit. Ind. iv. 45 = Tylophora asthmatica.
Weddellii, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 381.—Bolivia.
Wrightii, Greene, in Proc. Am. Acad. xvi. (1881) 102 = nyctaginifolia.
zeylanica, Burm. ex Decne. in DC. Prod. viii. 628 = Leptadenia reticulata.

ASCLEPIODORA, A. Gray, in Proc. Am. Acad. xii. (1876) 66. *ASCLEPIADEAE.*
ANANTHERIX, Nutt. Gen. Am. i. 169 (1818).
ANTHANOTIS, Rafin. Fl. Ludov. 52, 149 (1817), nomen prius.
asperula, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 369.—Mexic.
circinalis, Fourn. l. c. = Asclepias brachystephana.
decumbens, *A. Gray, in Proc. Am. Acad.* xii. (1876) 67.—Am. bor.
gomphocarpoides, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 369.—Mexic.
viridis, *A. Gray, in Proc. Am. Acad.* xii. (1876) 66.—Am. bor.

ASCOCHILUS, Blume, Fl. Jav. Pracf. p. vi. (1828) = **Geodorum**, Jack (Orchid.).

ASCOLEPIS, Nees, ex Steud. Syn. Pl. Cyp. 105 (1855). *CYPERACEAE*, Benth. & Hook. f. iii. 1054.
ANTROLEPIS, Welw. Apont. Phyto-Geograph. in Annaes do Consel. Ultramarin. (1858) 578.
PLATYLEPIS, Kunth, Enum. Pl. ii. 269 (1835).
PTERACHNE, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 62, in nota (1842).
PTEROGYNE, Schrad. ex Nees, in Mart. l. c. (1842).
anthemiflora, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 77.—Angola.
capensis, *Ridl. in Trans. Linn. Soc.* Ser. II. Bot. ii. (1884) 164.—Afr. austr. et trop.
elata, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 79.—Angola.
eriocaulides, *Nees, ex Steud. Syn. Pl. Cyp.* 105.—Abyss.
kyllingioides, Steud. l. c. = Lipocarpha microcephala.
protea, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 75.—Afr. trop.
pusilla, *Ridl. in Trans. Linn. Soc.* Ser. II. Bot. ii. (1884) 164.—Afr. occ.
speciosa, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 78.—Angola.
tenuior, Steud. Syn. Pl. Cyp. 105 = Scirpus squarrosus.

ASCRA, Schott, in Spreng. Syst. iv. Cur. Post. 407 (1827) = **Banara**, Aubl. (Samyd.).
brasiliensis, Schott, l. c. = B. brasiliensis.

ASCYRUM, Linn. Gen. ed. I. 231 (1737). *HYPERICINEAE*, Benth. & Hook. f. i. 164.
HYPERICOIDES, Adans. Fam. ii. 443 (1763).
ISOPHYLLUM, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367.
amplexicaule, *Michx. Fl. Bor. Am.* ii. 77.—Am. bor.
calicinum, Poir. Tabl. Encyc. iii. 199 = Hypericum calycinum.
coriaceum, Moench, Meth. 130 = Hypericum calycinum.
cruciatum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69 = A. Crux-Andreae.
Crux-Andreae, Linn. Sp. Pl. 787 = Hypericum mutilum.
Crux-Andreae, *Linn. Sp. Pl.* ed. II. 1107.—Am. bor.; Ind. occ.
cubense, *Griseb. Cat. Pl. Cub.* 40.—Cuba.
filicaule, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 252.—Reg. Himal.
glandulosum, Moench, Meth. Suppl. 42 = Hypericum balearicum.
grandiflorum, Rafin. Fl. Ludov. 87 = stans.
helianthemifolium, Spach, Hist. Veg. Phan. v. 460 = A. Crux-Andreae.
humifusum, Labill. Nov. Holl. Pl. ii. 33. t. 175 = Hypericum japonicum.
hypericifolium, Sw. ex Rafin. Sylva Tellur. 116 = A. Hypericoides.
Hypericoides, *Linn. Sp. Pl.* 788 & ed. II. 1108 pro max. parte.—Ind. occ.
Hypericoides, Marsh. Arbust. 14 = stans.
grandiflorum, Rafin. Fl. Ludov. 87 = stans.
involutum, Labill. Nov. Holl. Pl. ii. 32. t. 174 = Hypericum gramineum.
linifolium, Spach, Hist. Veg. Phan. v. 459 = A. Crux-Andreae.
Michauxii, Spach, l. c. 460 = A. Crux-Andreae.
microsepalum, Torr. & Gray, Fl. N. Am. i. 157 = Hypericum microsepalum.
monogynum, Moench, Meth. Suppl. 42 = Hypericum monogynum.
montanum, Rafin. Sylva Tellur. 116 = A. Hypericoides.

ASCYRUM:—
multicaule, Michx. ex Willd. Sp. Pl. iii. 1472 = A. Crux-Andreae.
nummularifolium, Banks, ex Steud. Nom. ed. I. 77 = pumilum.
oblongifolium, Spach, Hist. Vég. Phan. v. 461 = A. Crux-Andreae.
pauciflorum, Nutt. Gen. Am. ii. 15 = pumilum.
Plumierii, Bertol. in Mem. Acc. Sc. Bolog. iv. (1853) 77 (Misc. Bot. xiii. 19) = A. Crux-Andreae.
pumilum, *Michx. Fl. Bor. Am.* ii. 77.—Am. bor.
sibiricum, Poir. Tabl. Encyc. iii. 200 = Hypericum Ascyrum.
simplex, Zeyh. ex Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 389 = stans.
spathulatum, Spach, Hist. Vég. Phan. v. 462 = A. Crux-Andreae.
stans, *Michx. Fl. Bor. Am.* ii. 77.—Am. bor.
tetragonum, Moench, Meth. 130 = Hypericum Ascyron.
villosum, Linn. Sp. Pl. 788 = Hypericum setosum.

ASCYUM, Vahl, ex DC. Prod. i. 566 (1824) = ASCIUM, Schreb. = **Norantea**, Aubl. (Ternstr.).

ASEMEIA, Rafin. New Fl. Am. iv. 88 (1836) = **Polygala**, Linn.
alba, Rafin. l. c. = P. alba.
carnea, Rafin. l. c.—Am. bor.
leptopsis, Rafin. l. c.—Am. bor.
rosea, Rafin. l. c. = P. Senega.

ASEMNANTHA, Hook. f. in Benth. & Hook. f. Gen. ii. 106 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 106.
pubescens, *Hook. f. l. c.* 107.—Mexic.

ASEPHANANTHES, Bory, ex DC. Prod. iii. 323 (1828) = **Passiflora**, Linn.

ASICARIA, Neck. Elem. ii. 211 (1790) = **Polygonum**, Tourn.

ASIMIA, H. B. & K. Nov. Gen. et Sp. v. 61 (1821) = **Asimina**, Adans. (Anonac.).

ASIMINA, Adans. Fam. ii. 365 (1763). *ANONACEAE*, Benth. & Hook. f. i. 24.
ORCHIDOCARPUM, Michx. Fl. Bor. Am. i. 329 (1803).
aurantiaca, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 25 = Orophea aurantiaca.
Blainii, *Griseb. Cat. Pl. Cub.* 3.—Cuba.
campechiana, *H. B. & K. Nov. Gen. et Sp.* v. 61.—Mexic.
campaniflora, Spach, Hist. Vég. Phan. vii. 528 = triloba.
conoidea, Spach, l. c. 530 = triloba?
cuneata, Shuttlw. ex Chapm. Fl. S. U. St. 15 = grandiflora.
glabra, Hort. ex C. Koch, Dendrol. i. 384 = triloba.
grandiflora, *Dun. Monog. Anon.* 84. t. 11.—Am. bor.
grandiflora, Hort. ex Spach, Hist. Vég. Phan. v. 530 = *conoidea*.
lanceolata, Hook. f. in Journ. Linn. Soc. i. (1857) 129, *err. typ.* = Ascarina lanceolata.
neglecta, *Griseb. Cat. Pl. Cub.* 3.—Cuba.
parviflora, *Dun. Monog. Anon.* 82. t. 9.—Am. bor.
pygmaea, *Dun. l. c.* 84. t. 10.—Am. bor.
reticulata, *Shuttlw. ex Chapm. Fl. S. U. St.* 15.—Florida.
rhombifolia, *Griseb. Cat. Pl. Cub.* 3.—Ind. occ.
secundiflora, Shuttlw. ex Chapm. Fl. S. U. St. 15 = pygmaea.
triloba, *Dun. Monog. Anon.* 83.—Am. bor.

ASIPHONIA, Griff. in Trans. Linn. Soc. xix. (1845) 333. t. 37 = **Bragantia**, Lour. (Aristoloch.).
piperiformis, Griff. l. c. = B. corymbosa.

ASKIDIOSPERMA, Steud. in Flora, xxxiii. (1850) 229, nomen; et Syn. Pl. Cyp. ii. 257 (1855). *RESTIACEAE*, Benth. & Hook. f. iii. 1032.
capitatum, *Steud. ll. cc.*—Afr. austr.

ASKOFAKE, Rafin. Fl. Tellur. iv. 108 (1836) = **Utricularia**, Linn. (Lentib.).
recurva, Rafin. l. c. = U. recurva.

ASKOLAME, Rafin. Fl. Tellur. ii. 11 (1836) = **Milla**, Cav. (Liliac.).
biflora, Rafin. l. c. 12 = M. biflora.

ASPALATHUS, Linn. Sp. Pl. 711 (1753). *LEGUMINOSAE*, Benth. & Hook. f. i. 478.
ACROPODIUM, Desv. in Ann. Sc. Nat. ix. (1826) 408.
CYPHOCALYX, HETEROLATHUS, LAPASANTHUS, PACHYRAPHEA, PARASPALATHUS, PLAGIOSTIGMA, PSILOLEPUS, STREPTOSEMA, et TRINCURIA, Presl, in Abh. Boehm. Ges. Wiss. Folge V. iii. 556–562 (Bot. Bemerk. 126–132 (1844).
DIALLOSPERMA, Rafin. Sylva Tellur. 69 (1838).
ERIOCYLAX, Neck. Elem. iii. 25 (1790).
NEFRAKIS, Rafin. Sylva Tellur. 69 (1838).
SARCOCALYX, Walp. in Linnaea, xiii. (1839) 479.
SARCOPHYLLUS, Thunb. in Schrad. Journ. i. (1799) 323.
SCALIGERA, Adans. Fam. ii. 373 (1763).
SEMETOR, Rafin. Sylva Tellur. 69 (1838).
abietina, E. Mey. Comm. Pl. Afr. Austr. 48 = aciphylla.
abietina, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
acanthes, *Eckl. & Zeyh. Enum.* 215.—Afr. austr.
acanthophylla, Eckl. & Zeyh. l. c. 221 = Chamissonis.
acicularis, *E. Mey. Comm. Pl. Afr. Austr.* 46.—Afr. austr.
aciphylla, *Harv. in Harv. & Sond. Fl. Cap.* ii. 129.—Afr. austr.
aculeata, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
aculeata, Willd. ex Walp. in Linnaea, xiii. (1839) 486 = tridentata.
acuminata, *Lam. Encyc.* i. 287.—Afr. austr.
acuminata, Thunb. Prod. Pl. Cap. 125 = ambigua.
adelphea, *Eckl. & Zeyh. Enum.* 212.—Afr. austr.
adscendens, Steud. Nom. ed. II. i. 147 = elongata.
aemula, *E. Mey. Comm. Pl. Afr. Austr.* 42.—Afr. austr.
affinis, Eckl. & Zeyh. Enum. 212 = microdon.
affinis, E. Mey. Comm. Pl. Afr. Austr. 60 = pinguis.
affinis, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
Agardhiana, *DC. Prod.* ii. 143.—Afr. austr.
albanensis, Eckl. & Zeyh. Enum. 211 = frankenioides.
albens, E. Mey. ex Benth. in Hook. Lond. Journ. Bot. vii. (1848) 648 = exilis.
albens, *Linn. Mant.* ii. 261; *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
albens, Willd. ex Walp. in Linnaea, xiii. (1839) 499 = armata.
albiflora, Eckl. & Zeyh. Enum. 207 = thymifolia.
alopecuroides, E. Mey. Comm. Pl. Afr. Austr. 52 = setacea.
Alopecurus, *Burch. ex Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 628, (err. typ. 618).—Afr. austr.
alpina, Eckl. & Zeyh. Enum. 218 = rubro-fusca.
alternifolia, *Harv. in Harv. & Sond. Fl. Cap.* ii. 141.—Afr. austr.
alternifolia, *Spreng. Syst.* iii. 187.—Afr. austr.
ambigua, *DC. Prod.* ii. 143.—Afr. austr.
angustissima, *E. Mey. Comm. Pl. Afr. Austr.* 44.—Afr. austr.
anthylloides, Berg. Desc. Pl. Cap. 211 = lotoides.
anthylloides, *Linn. Sp. Pl.* ed. II. 1002.—Afr. austr.
appendiculata, E Mey. in Linnaea, vii. (1832) 157 = ciliaris.
arachnoidea, Otto, ex Steud. Nom. ed. II. i. 147, 149 = setacea.
arachnoidea, *Walp. in Linnaea*, xiii. (1839) 497.—Afr. austr.
araneosa, *Linn. Sp. Pl.* 712.—Afr. austr.
arborea, *Lour. Fl. Cochinch.* 431.—Cochinch.
argentea, Eckl. & Zeyh. Enum. 203 = aemula.
argentea, *Linn. Sp. Pl.* 713.—Afr. austr.
argentea, J. F. Gmel. Syst. 1093 = sericea.
argyraea, *DC. Prod.* ii. 139.—Afr. austr.
arida, *E. Mey. in Linnaea*, vii. (1832) 156.—Afr. austr.
armata, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
ascendens, E. Mey. Comm. Pl. Afr. Austr. 41 = elongata.
asparagoides, *Linn. f. Suppl.* 321.—Afr. austr.
astroides, Willd. ex Walp. in Linnaea, xiii. (1839) 503 = racemosa.
astroites, *Linn. Sp. Pl.* 711.—Afr. austr.
astroites, Thunb. Prod. Pl. Cap. 128 = racemosa.
aulonogena, Eckl. & Zeyh. Enum. 216 = ciliaris.
batodes, Eckl. & Zeyh. l. c. 215 = aciphylla.
Benthamii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 111.—Afr. austr.

ASPALATHUS:—
bicolor, Eckl. & Zeyh. Enum. 205 = retroflexa.
biflora, E. Mey. Comm. Pl. Afr. Austr. 64 = pedunculata.
bracteata, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
Burchelliana, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 623.—Afr. austr.
calcarata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 116.—Afr. austr.
callosa, *Linn. Sp. Pl.* 713.—Afr. austr.
callosa, Sims, Bot. Mag. t. 2329 = araneosa.
canaliculata, E. Mey. Comm. Pl. Afr. Austr. 44 = stenophylla.
candicans, *Ait. Hort. Kew.* ed. II. iv. 264.—Afr. austr.
canescens, *Linn. Mant.* ii. 262.—Afr. austr.
capillaris, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 653.—Afr. austr.
capitata, *Linn. Pl. Afr. Rar.* 14; *Amoen. Acad.* vi. 92.—Afr. austr.
capitella, Burch. ex Benth. in Hook. Lond. Journ. Bot. vii. (1848) 597 = inops.
carmaënsis, Eckl. & Zeyh. Enum. 1404 = thymifolia.
carnosa, *Berg. Descr. Pl. Cap.* 206.—Afr. austr.
carnosa, Eckl. & Zeyh. Enum. 220 = pachyloba.
carnosa, Linn. Mant. 261 = sarcantha.
Cephalotes, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 615 = Forbesii.
Cephalotes, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
cerrhantha, Eckl. & Zeyh. Enum. 208 = spicata.
Chamissonis, *Vog. in Linnaea*, x. (1835–36) 597.—Afr. austr.
Chenopoda, *Linn. Sp. Pl.* 711.—Afr. austr.
chortophila, Eckl. & Zeyh. Enum. 211 = frankenioides.
ciliaris, Sims, Bot. Mag. t. 2233 = A. Chenopoda.
ciliaris, *Linn. Mant.* 262.—Afr. austr.
cinerascens, *E. Mey. Comm. Pl. Afr. Austr.* 54.—Afr. austr.
cinerea, Thunb. Prod. Pl. Cap. 125 = cytisoides.
coerulescens, E. Mey. Comm. Pl. Afr. Austr. 54 = Lebeckia microphylla.
cognata, Presl, Bot. Bemerk. 126 = corymbosa.
collina, *Eckl. & Zeyh. Enum.* 220.—Afr. austr.
comosa, Eckl. & Zeyh. l. c. 209 = A. Alopecurus.
comosa, E. Mey. Comm. Pl. Afr. Austr. 56 = Benthamii.
comosa, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
comosa, Willd. ex Walp. in Linnaea, xiii. (1839) 490 = capitata.
conferta, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 596 = exigua.
corrudaefolia, Berg. Descr. Pl. Cap. 207 = genistoides, pungens.
corrudaefolia, DC. Prod. ii. 139 = rigescens.
corymbosa, *E. Mey. in Linnaea*, vii. (1832) 159.—Afr. austr.
costulata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 641.—Afr. austr.
crassifolia, *Andr. Bot. Rep.* t. 351.—Afr. austr.
cretica, Linn. Sp. Pl. 712 = Anthyllis Hermanniae.
cuneata, D. Don, Prod. Fl. Nep. 246 = Lespedeza sericea.
cuneata, *E. Mey. Comm. Pl. Afr. Austr.* 37.—Afr. austr.
cymbaeformis, DC. Prod. ii. 140 = uniflora.
cytisoides, *Lam. Encyc.* i. 292.—Afr. austr.
dasyantha, *Eckl. & Zeyh. Enum.* 201.—Afr. austr.
deciduifolia, Eckl. & Zeyh. l. c. 210 = nigra.
densifolia, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 632.—Afr. austr.
diffusa, *Eckl. & Zeyh. Enum.* 208.—Afr. austr.
divaricata, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
divergens, E. Mey. Comm. Pl. Afr. Austr. 45 = praec.
divergens, Willd. ex Walp. in Linnaea, xiii. (1839) 493 = microphylla
Dorycnium, Lam. Encyc. i. 290 = Dorycnium suffruticosum.
Dregeana, Walp. in Linnaea, xiii. (1839) 486 = incurva.
dubia, E. Mey. in Linnaea, vii. (1832) 157 = ciliaris.
dubia, Willd. ex Walp. in Linnaea, xiii. (1839) 504 = Amphithalea ericaefolia.
Ebenus, Linn. Syst. ed. X. 1158 = Brya Ebenus.
echinata, E. Mey. Comm. Pl. Afr. Austr. 51 = rigescens.
echinata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 119 = A. Hystrix.
Ecklonis, Vog. ex Walp. in Linnaea, xiii. (1839) 489 = ciliaris.

ASPALATHUS:—
elongata, *Eckl. & Zeyh. Enum.* 202.—Afr. austr.
elongata, E. Mey. Comm. Pl. Afr. Austr. 63 = incurva.
ericaefolia, Berg. Desc. Pl. Cap. 205 = uniflora.
ericifolia, *Linn. Sp. Pl.* 711.—Afr. austr.
ericaefolia, Willd. ex Walp. in Linnaea, xiii. (1839) 495 = thymifolia.
ericoides, E. Mey. in Linnaea, vii. (1832) 160 = ericifolia.
erinaeea, *Lam. Encyc.* i. 291.—Afr. austr.
eriophylla, *Walp. in Linnaea*, xiii. (1839) 499.—Afr. austr.
erythrodes, *Eckl. & Zeyh. Enum.* 200.—Afr. austr.
exigua, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 597 = marginata.
exigua, *Eckl. & Zeyh. Enum.* 198.—Afr. austr.
exilis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 136.—Afr. austr.
falcata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 655.—Afr. austr.
ferox, *Harv. in Harv. & Sond. Fl. Cap.* ii. 138.—Afr. austr.
ferruginea, *Banks, ex Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 607.—Afr. austr.
filicaulis, *Eckl. & Zeyh. Enum.* 204.—Afr. austr.
filifolia, E. Mey. in Linnaea, vii. (1832) 158 = abietina.
flavescens, Presl, Bot. Bemerk. 126 = spinosa.
flexuosa, Thunb. Prod. Pl. Cap. 127 = mollis.
flexuosa, Willd. ex Walp. in Linnaea, xiii. (1839) 495 = ericifolia.
floribunda, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 636.—Afr. austr.
Forbesii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 114.—Afr. austr.
fornicata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 646.—Afr. austr.
frankenioides, *DC. Prod.* ii. 139.—Afr. austr.
frankenioides, Eckl. & Zeyh. Enum. 207 = thymifolia.
frutescens, Vog. ex Walp. in Linnaea, xiii. (1839) 487 = pinea.
fusca, Thunb. Prod. Pl. Cap. 125; Fl. Cap. ed. Schult. 574 = callosa.
galeata, *E. Mey. Comm. Pl. Afr. Austr.* 49.—Afr. austr.
galioides, Berg. Desc. Pl. Cap. 210 = retroflexa.
galioides, *Linn. Mant.* ii. 260; *Thunb. Diss. Aspal.* 19.—Afr. austr.
galioides, Sieber, ex Walp. in Linnaea, xiii. (1839) 502 = divaricata.
galioides, Willd. ex Walp. l. c. 493, 502 = pedunculata.
garipensis, E. Mey. Comm. Pl. Afr. Austr. 44.—Afr. austr.
genistoides, Eckl. & Zeyh. Enum. 215 = laricifolia.
genistoides, *Linn. Mant.* 261.—Afr. austr.
genistoides, Thunb. Prod. Pl. Cap. 128 = pungens.
Gillii, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 627 = setacea.
glauca, Eckl. & Zeyh. Enum. 221 = obtusata.
globosa, *Andr. Bot. Rep.* t. 510.—Afr. austr.
globosa, Hort. ex Steud. Nom. ed. II. i. 148 = nigra.
globosa, Eckl. & Zeyh. Enum. 209 = spicata.
globulosa, E. Mey. in Linnaea, vii. (1832) 159 = nigra.
glomerata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 623.—Afr. austr.
glomerata, Linn. f. Suppl. 321 = capitata.
grandiflora, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 632.—Afr. austr.
heterophylla, *E. Mey. Comm. Pl. Afr. Austr.* 40.—Afr. austr.
heterophylla, Linn. f. Suppl. 321; Thunb. Prod. Pl. Cap. 126 = lotoides.
hiatuum, Eckl. & Zeyh. Enum. 212 = frankenioides.
hilaris, Eckl. & Zeyh. l. c. 214 = laricifolia.
hirta, *E. Mey. in Linnaea*, vii. (1832) 156.—Afr. austr.
hispida, Sieber, ex Presl, Bot. Bemerk. 131 = miiflora.
hispida, Thunb. Prod. Pl. Cap. 127 = thymifolia.
holosericea, E. Mey. Comm. Pl. Afr. Austr. 37 = Buchcuroedera holosericea.
horrida, Eckl. & Zeyh. Enum. 221 = spinosa.
Hystrix, Eckl. & Zeyh. l. c. 219 = Willdenowiana.
Hystrix, *Linn. f. Suppl.* 322.—Afr. austr.
incomta, E. Mey. ex Benth. in Hook. Lond. Journ. Bot. vii. (1848) 643 = lactea.
incomta, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
incurva, Eckl. & Zeyh. Enum. 208 = comosa.
incurva, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.

ASPALATHUS:—
incurvifolia, *Vog. ex Walp. in Linnaea*, xiii. (1839) 497.—Afr. austr.
indica, Linn. Sp. Pl. 712 = Indigofera aspalathoides.
iniquua, Eckl. & Zeyh. Enum. 212 = adelphea.
inops, *Eckl. & Zeyh. l. c.* 197.—Afr. austr.
intermedia, Eckl. & Zeyh. l. c. 211 = frankenioides.
involucrata, E. Mey. Comm. Pl. Afr. Austr. 38 = undulata.
involucrata, Pappe, ex Harv. & Sond. Fl. Cap. ii. 114 = nigra.
Jacobaea, *E. Mey. Comm. Pl. Afr. Austr.* 41.—Afr. austr.
Joubertiana, Eckl. & Zeyh. Enum. 214 = canescens.
juniperina, Thunb. Prod. Pl. Cap. 128 = galioides.
kannaënsis, Eckl. & Zeyh. Enum. 207 = ericifolia.
Kraussiana, Meissn. in Hook. Lond. Journ. Bot. ii. (1848) 69 = anthylloides.
lactea, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
lanata, *E. Mey. Comm. Pl. Afr. Austr.* 64.—Afr. austr.
lanceolata, *E. Mey. l. c.* 37.—Afr. austr.
laricifolia, *Berg. Desc. Pl. Cap.* 204.—Afr. austr.
laricifolia, Lam. Encyc. i. 287 = leptophylla.
laricina, DC. Prod. ii. 141 = laricifolia.
laricina, Steud. Nom. ed. II. i. 149 = tridentata.
laxata, *Linn. Mant.* 263.—Afr. austr.
lepida, *E. Mey. Comm. Pl. Afr. Austr.* 58.—Afr. austr.
leptocoma, Eckl. & Zeyh. Enum. 205 = divaricata.
leptophylla, *Eckl. & Zeyh. l. c.* 219.—Afr. austr.
leptothria, Eckl. & Zeyh. l. c. 213 = frankenioides.
leucocephala, E. Mey. Comm. Pl. Afr. Austr. 41 = virgata.
leucophaea, *Harv. in Harv. & Sond. Fl. Cap.* ii. 112.—Afr. austr.
linearifolia, *DC. Prod.* ii. 142.—Afr. austr.
linifolia, E. Mey. in Linnaea, vii. (1832) 162 = linearifolia.
linifolia, Steud. in Flora, xiii. (1830) 543 = heterophylla.
longifolia, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 624.—Afr. austr.
longipes, *Harv. in Harv. & Sond. Fl. Cap.* ii. 141.—Afr. austr.
lotoides, E. Mey. Comm. Pl. Afr. Austr. 42 = Meyeri.
lotoides, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
lotoides, Willd. ex Walp. in Linnaea, xiii. (1839) 486, 503 = tridentata.
macrantha, *Harv. in Harv. & Sond. Fl. Cap.* ii. 117.—Afr. austr.
macrocarpa, *Eckl. & Zeyh. Enum.* 203.—Afr. austr.
macrosepala, *Steud. Nom.* ed. II. i. 149.—Afr. austr.
marginalis, *Eckl. & Zeyh. Enum.* 213.—Afr. austr.
marginata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 103.—Afr. austr.
melanoides, Eckl. & Zeyh. Enum. 210 = nigra.
Meyeri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 106.—Afr. austr.
Meyeriana, Eckl. & Zeyh. Enum. 218 = ciliaris.
micrantha, E. Mey. in Linnaea, vii. (1832) 161 = thymifolia.
microcarpa, *Burch. ex DC. Prod.* ii. 139.—Afr. austr.
microdon, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 642.—Afr. austr.
microphylla, DC. Prod. ii. 143 = divaricata.
microphylla, *Steud, in Flora*, xiii. (1830) II. 545.—Afr. austr.
minutifolia, Vog. ex Walp. in Linnaea, xiii. (1839) 500 = pinguis.
mollis, *Lam. Encyc.* i. 290.—Afr. austr.
mucronata, Linn. f. Suppl. 320 = Viborgia armata.
multiflora, Thunb. Prod. Pl. Cap. 128 = thymifolia.
multiflora, Sieber, ex Walp. in Linnaea, xiii. (1839) 495 = ericifolia.
Mundiana, Eckl. & Zeyh. Enum. 220 = verrucosa.
muraltioides, Eckl. & Zeyh. l. c. 209 = mollis.
myrtillifolia, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 597.—Afr. austr.
neanthes, Eckl. & Zeyh. Enum. 213 = canescens.
nervosa, *E. Mey. Comm. Pl. Afr. Austr.* 62.—Afr. austr.
nigra, *Linn. Mant.* 262.—Afr. austr.
nigrescens, E. Mey. in Linnaea, vii. (1832) 159 = nigra.
nivea, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
nivea, Willd. ex Walp. in Linnaea, xiii. (1839) 486 = purpurea.
nodosa, *Vog. ex Walp. l. c.* 496.—Afr. austr.

ASPALATHUS:—
nudiflora, *Harv. in Harv. & Sond. Fl. Cap.* ii. 140.—Afr. austr.
obtusata, *Thunb. Prod. Pl. Cap.* 125.—Afr. austr.
opaca, *Eckl. & Zeyh. Enum.* 217.—Afr. austr.
orbiculata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 595.—Afr. austr.
oresigena, Eckl. & Zeyh. Enum. 217 = ciliaris.
orientalis, *Linn. Mant.* 263.—Oriens.
pachyloba, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 638.—Afr. austr.
pallens, Eckl. & Zeyh. Enum. 210 = nigra.
pallescens, *Eckl. & Zeyh. l. c.* 218.—Afr. austr.
papillosa, Eckl. & Zeyh. l. c. 217 = ciliaris.
Pappeana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 117.—Afr. austr.
parviflora, *Berg. Desc. Pl. Cap.* 208; *Thunb. Prod. Pl. Cap.* 122.—Afr. austr.
pedicellata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 135.—Afr. austr.
pedunculata, E. Mey. ex Walp. Ann. ii. 338 = bracteata, capillaris.
pedunculata, *L'Hérit. Sert. Angl.* t. 26.—Afr. austr.
persica, Burm. f. Fl. Ind. 155 = Lotus Gareini.
pilosa, Eckl. & Zeyh. Enum. 202 = tridentata.
pilosa, *Linn. Mant.* 263.—Afr. austr.
pilosa, Sieber, ex Walp. in Linnaea, xiii. (1839) 488 = arancosa.
pinea, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
pinguis, Eckl. & Zeyh. Enum. 220 = arida.
pinguis, E. Mey. Comm. Pl. Afr. Austr. 60 = affinis.
pinguis, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
pinnata, *Linn. Pl. Rar. Afr.* 14; *Amoen. Acad.* vi. 92.—Afr. austr.
Plukenetiana, Eckl. & Zeyh. Enum. 200 = rugosa.
poliotes, Eckl. & Zeyh. l. c. 213 = frankenioides.
polyantha, *Walp. in Linnaea*, xiii. (1839) 485.—Afr. austr.
polycephala, *E. Mey. Comm. Pl. Afr. Austr.* 39.—Afr. austr.
Priori, *Harv. in Harv. & Sond. Fl. Cap.* ii. 127.—Afr. austr.
procumbens, *Eckl. & Zeyh. Enum.* 202.—Afr. austr.
procumbens, E. Mey. in Linnaea, vii. (1832) 162 = lotoides.
propinqua, *E. Mey. Comm. Pl. Afr. Austr.* 53.—Afr. austr.
prostrata, *Eckl. & Zeyh. Enum.* 206.—Afr. austr.
psoraleoides, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 598.—Afr. austr.
pulchella, E. Mey. Comm. Pl. Afr. Austr. 38 = Buchenroedera tenuifolia.
pungens, *Thunb. Prod. Pl. Cap.* 129.—Afr. austr.
purpurascens, E. Mey. Comm. Pl. Afr. Austr. 44 = purpurea.
purpurea, *Eckl. & Zeyh. Enum.* 201.—Afr. austr.
quinquefolia, Linn. Pl. Rar. Afr. 14; Amoen. Acad. vi. 92 = lotoides.
quinquefolia, Thunb. Fl. Cap. 575 = virgata.
racemosa, *E. Mey. in Linnaea*, vii. (1832) 157.—Afr. austr.
ramulosa, E. Mey. l. c. 163 = aemula.
recurva, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 642.—Afr. austr.
remota, *Eckl. & Zeyh. Enum.* 218.—Afr. austr.
retroflexa, Eckl. & Zeyh. l. c. 204 = abietina.
retroflexa, *Linn. Sp. Pl.* 712.—Afr. austr.
rigescens, *E. Mey. Comm. Pl. Afr. Austr.* 52.—Afr. austr.
rostrata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 631.—Afr. austr.
rubens, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
rubescens, Eckl. & Zeyh. Enum. 213 = adelphea.
rubrofusca, *Eckl. & Zeyh. l. c.* 216.—Afr. austr.
rugosa, *Thunb. Prod. Pl. Cap.* 125.—Afr. austr.
sanguinea, Eckl. & Zeyh. Enum. 212 = vermiculata.
sanguinea, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
sarcantha, *Vog. ex Walp. in Linnaea*, xiii. (1839) 489.—Afr. austr.
sarcodes, *Vog. ex Walp. l. c.* 480.—Afr. austr.
scaphoides, Eckl. & Zeyh. Enum. 206 = uniflora.
Schlliana, *Presl, Bot. Bemerk.* 129.—Afr. austr.
secunda, E. Mey. Comm. Pl. Afr. Austr. i. 47 = pungens.
securifolia, *Eckl. & Zeyh. Enum.* 198.—Afr. austr.
sericantha, E. Mey. Comm. Pl. Afr. Austr. 49 = laricifolia.
sericea, *Berg. Desc. Pl. Cap.* 212.—Afr. austr.

ASPALATHUS:—
sericea, Eckl. & Zeyh. Enum. 202 = Jacobaea.
sericea, Lam. Encyc. i. 288 = nivea.
sericea, *Linn. f. Suppl.* 321.—Afr. austr.
sericea, Thunb. Prod. Pl. Cap. 125 = Argyrolobium obsoletum.
setacea, *Eckl. & Zeyh. Enum.* 216.—Afr. austr.
Simsiana, Eckl. & Zeyh. l. c. 200 = araneosa.
spathulata, Eckl. & Zeyh. l. c. 198 = securifolia.
speciosa, *Steud. in Flora*, xiii. (1830) 544.—Afr. austr.
spicata, *Thunb. Prod. Pl. Cap.* 127.—Afr. austr.
spinescens, DC. Prod. ii. 138 = arida.
spinescens, Eckl. & Zeyh. Enum. 221 = divaricata.
spinescens, *Thunb. Prod. Pl. Cap.* 129.—Afr. austr.
spiniflora, L'Hérit. ex Steud. Nom. ed. II. i. 107, 149 = Anthyllis Hermanniae.
spinosa, *Linn. Sp. Pl.* ed. II. 1000.—Afr. austr.
squamosa, Thunb. Fl. Cap. ed. Schult. 581 = pedunculata.
squarrosa, Thunb. Prod. Pl. Cap. 128 = pedunculata.
stachyera, Eckl. & Zeyh. Enum. 202 = lotoides.
staurantha, Eckl. & Zeyh. l. c. = argentea.
stellaris, *Eckl. & Zeyh. l. c.* 197.—Afr. austr.
stenophylla, *Eckl. & Zeyh. l. c.*—Afr. austr.
Steudeliana, B[rongn.] in Bull. Féruss. xxvi. (1831) 278 = microphylla.
stricta, Steud. in Flora, xiii. (1830) II. 543 = elongata.
suaveolens, *Eckl. & Zeyh. Enum.* 199.—Afr. austr.
subinermis, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 643.—Afr. austr.
subtingens, Eckl. & Zeyh. Enum. 212 = adelphea.
subulata, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
succulenta, E. Mey. in Linnaea, vii. (1832) 159 = verrucosa.
suffruticosa, *DC. Prod.* ii. 144.—Afr. austr.
sulcata, Vog. ex Walp. in Linnaea, xiii. (1839) 490 = erythrodes.
tenuifolia, *DC. Prod.* ii. 143.—Afr. austr.
teres, *Eckl. & Zeyh. Enum.* 215.—Afr. austr.
thymifolia, Lam. Encyc. i. 288 = canescens.
thymifolia, *Linn. Sp. Pl.* 711.—Afr. austr.
thymifolia, Thunb. Prod. Pl. Cap. 127 = comosa.
tomentosa, E. Mey. Comm. Pl. Afr. Austr. 55 = frankenioides.
trichodes, E. Mey. l. c. 38 = Buchenroedera trichodes.
tridentata, Berg. ex Walp. in Linnaea, xiii. (1839) 482 = pilosa.
tridentata, Eckl. & Zeyh. Enum. 201 = ferruginea.
tridentata, *Linn. Sp. Pl.* 712.—Afr. austr.
trigona, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
triquetra, *Thunb. l. c.* 127.—Afr. austr.
truncata, *Eckl. & Zeyh. Enum.* 197.—Afr. austr.
tuberculata, *Walp. in Linnaea*, xiii. (1839) 497.—Afr. austr.
tylodes, Eckl. & Zeyh. Enum. 200 = callosa.
ulicina, *Eckl. & Zeyh. l. c.* 205.—Afr. austr.
undulata, *Eckl. & Zeyh. l. c.* 199.—Afr. austr.
uniflora, Eckl. & Zeyh. l. c. 219 = macrantha.
uniflora, *Linn. Sp. Pl.* 712.—Afr. austr.
uniflora, Thunb. Diss. Aspal. 20 = Willdenowiana.
varians, Eckl. & Zeyh. Enum. 209 = ericifolia.
variegata, *Eckl. & Zeyh. l. c.* 201.—Afr. austr.
venosa, *E. Mey. Comm. Pl. Afr. Austr.* 39.—Afr. austr.
vermiculata, *Lam. Encyc.* i. 288.—Afr. austr.
verrucosa, Eckl. & Zeyh. Enum. 220 = tuberculata.
verrucosa, *Linn. Sp. Pl.* 712.—Afr. austr.
verrucosa, Willd. ex Walp. in Linnaea, xiii. (1839) 486 = Willdenowiana.
versicolor, E. Mey. Comm. Pl. Afr. Austr. 48 = collina.
villosa, *Thunb. Prod. Pl. Cap.* 125.—Afr. austr.
viminea, E. Mey. Comm. Pl. Afr. Austr. 38 = Buchenroedera viminea.
virgata, *Thunb. Prod. Pl. Cap.* 126.—Afr. austr.
volvularis, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 7.—Malaya.
vulnerans, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 621 = hirta.
vulnerans, *Thunb. Prod. Pl. Cap.* 128.—Afr. austr.
Willdenowiana, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 630.—Afr. austr.
Wurmbeana, *E. Mey. Comm. Pl. Afr. Austr.* 58.—Afr. austr.

ASPALATUS, St. Hil. in Dict. Sc. Nat. iii. 211 (1804) = **Aspalathus**, Linn. (Legum.).

ASPALTHIUM, Medic. Phil. Bot. i. 211 (1789) = ASPHALTHIUM, Medic. = **Psoralea**, Linn. (Legum.).

ASPARAGOPSIS, Kunth, in Abh. Akad. Berl. (1842) 35 = **Asparagus**, Tourn. (Liliac.).
abyssinica, Kunth, Enum. Pl. v. 101 = Asparagus racemosus.
acerosa, Kunth, l. c. 102 = Asparagus acerosus.
adscendens, Kunth, l. c. = Asparagus adscendens.
aethiopica, Kunth, l. c. 95 = Asparagus aethiopicus.
alba, Kunth, l. c. 84 = Asparagus albus.
arborescens, Kunth, l. c. 79 = Asparagus arborescens.
ascendens, Baker, in Journ. Linn. Soc. xiv. (1875) 624 = Asparagus adscendens.
Brownei, Kunth, Enum. Pl. v. 103 = Asparagus racemosus.
consanguinea, Kunth, l. c. 76 = Asparagus consanguineus.
Curilla, Kunth, l. c. 102 = Asparagus Curillus.
Decaisnei, Kunth, l. c. 103 = Asparagus racemosus.
decumbens, Kunth, l. c. 77 = Asparagus crispus.
densiflora, Kunth, l. c. 96 = Asparagus sarmentosus.
denudata, Kunth, l. c. 82 = Asparagus denudatus.
Dregii, Kunth, l. c. 84 = Asparagus Thunbergianus.
flagellaris, Kunth, l. c. 103 = Asparagus flagellaris.
floribunda, Kunth, l. c. 98 = Asparagus racemosus.
grandiflora, Kunth, l. c. 80 = Asparagus umbellatus.
Hohenackerii, Kunth, l. c. 100 = Asparagus racemosus.
humilis, Hassk. in Versl. en Med. Akad. Wetensch. v. (1857) 93.—Japon.
javanica, Kunth, Enum. Pl. v. 100 = Asparagus racemosus.
juniperina, Kunth, l. c. 85 = Asparagus africanus.
Krebsiana, Kunth, l. c. 93 = Asparagus racemosus.
Lamarckii, Kunth, l. c. 87 = Asparagus africanus.
lancea, Kunth, l. c. 104 = Asparagus sarmentosus.
Lowei, Baker, in Journ. Linn. Soc. xiv. (1875) 611 = Asparagus umbellatus.
microrhaphis, Kunth, Enum. Pl. v. 83 = Asparagus microrhaphis.
minutiflora, Kunth, l. c. 89 = Asparagus minutiflorus.
Niveniana, Kunth, l. c. 88 = Asparagus africanus.
passerinoides, Kunth, l. c. 90 = Asparagus capensis.
retrofracta, Kunth, l. c. 88 = Asparagus retrofractus.
retrofracta, Schweinf. ex Baker, in Journ. Linn. Soc. xiv. (1875) 623 = Asparagus racemosus.
rubricaulis, Kunth, Enum. Pl. v. 87 = Asparagus rubricaulis.
sarmentosa, Dalz. & Gibs. Bomb. Fl. 246 = Asparagus racemosus.
sarmentosa, Kunth, Enum. Pl. v. 97 = Asparagus sarmentosus.
scandens, Kunth, l. c. 78 = Asparagus scandens.
Schlechtendalii, Kunth, l. c. 90 = Asparagus Thunbergianus.
scoparia, Kunth, l. c. 80 = Asparagus asiaticus.
setacea, Kunth, l. c. 82 = Asparagus declinatus.
sinica, Miq. in Journ. Bot. Néerl. i. (1861) 90 (= Asparagus lucidus ?).—Hongkong.
spinescens, Kunth, Enum. Pl. v. 93 = Asparagus stipulaceus.
stipulacea, Kunth, l. c. 91 = Asparagus stipulaceus.
suaveolens, Baker, in Journ. Linn. Soc. xiv. (1875) 617 = Asparagus suaveolens.
subquadrangularis, Kunth, Enum. Pl. v. 94 = Asparagus racemosus.
Thunbergii, Kunth, l. c. 85 = Asparagus Thunbergianus.
triacantha, Kunth, l. c. 91 = Asparagus stipulaceus.
umbellata, Kunth, l. c. 79 = Asparagus umbellatus.
umbellulata, Kunth, l. c. 81 = Asparagus umbellulatus.
Willdenowii, Kunth, l. c. 86 = Asparagus asiaticus.
Zeyheri, Kunth, l. c. 92 = Asparagus stipulaceus.

ASPARAGUS, Tourn. ex Linn. Syst. ed. I. (1735); Gen. ed. I. 93 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 765.
ASPARAGOPSIS, Kunth, in Abh. Akad. Berl. (1842) 35.
ELIDE, Medic. Phil. Bot. ii. 71 (1791).
HECATRIS, Salisb. Gen. Pl. Fragm. 66 (1866).
MYRSIPHYLLUM, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 25.
abyssinicus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 319.—Abyssin.
abyssinicus, Schweinf. ex Baker, in Journ. Linn. Soc. xiv. (1875) 616 = Schweinfurthii.

ASPARAGUS:—
acerosus, *Roxb. Hort. Beng.* 24; *Fl. Ind.* ii. 150.—Ind. or.; Burma.
acerosus, *Thunb. ex Schult. f. Syst.* vii. 337.—Afr. austr.
acutifolius, *Linn. Sp. Pl.* 314.—Reg. Mediterr.
adscendens, *Roxb. Hort. Beng.* 24; *Fl. Ind.* ii. 153.—Reg. Himal.; Afghan.
aethiopicus, *Linn. Mant.* 1.—Afr. austr.
aetnensis, Tornab. in Atti Acc. Gioenia, Ser. II. xii. (1856) 345 = acutifolius.
africanus, *Lam. Encyc.* i. 295.—Afr. austr.
Agul, *Forsk. Fl. Aegypt. Arab.* p. lxv. *ex Schult. f. Syst.* vii. 337.—Arabia.
albus, *Linn. Sp. Pl.* 313.—Reg. Mediterr. occ.
albus, Thunb. Prod. Pl. Cap. 66 = Thunbergianus.
altilis, Aschers. Fl. Brandenb. 730 = officinalis.
altissimus, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 287.—Algeria.
amarus, DC. Cat Hort. Monsp. 81 = scaber.
ambiguus, De Not. Rep. Fl. Lig. 401 = acutifolius.
angolensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 254.—Afr. trop.
aphyllus, *Linn. Sp. Pl.* 314.—Reg Mediterr.
arborescens, *Willd. ex Schult. f. Syst.* vii. 337.—Ins. Teneriffa.
ascendens, Auct. vide adscendens.
asiaticus, *Linn. Sp. Pl.* 313.—Ind. or.; Afr. trop. et austr.
asiaticus, Wight, Ic. t. 2055 = subalatus.
benguellensis, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 253.—Angola.
brachyphyllus, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 78.—Reg. Danub.; Soongaria; Afghan.
Breslerianus, Schult. f. Syst. vii. 323 = maritimus.
brevifolius, *Boiss. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 602.—Lycia.
brevifolius, Tornab. in Atti Acc. Gioenia, Ser. II. xii. (1856) 359 = acutifolius.
Broussonetii, *Jacq. Eclog.* 3. t. 40.—Afr. bor.
Burchellii, *Baker, in Journ. Linn. Soc.* xiv. (1875) 618.—Afr. austr.
Burkei, *Baker, l. c.* 607.—Afr. austr.
Buxbaumius, Gueldenst. ex Ledeb. Fl. Ross. iv. 199 = verticillatus.
capensis, *Linn. Sp. Pl.* 314.—Afr. austr.
capitatus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 607.—Ind. bor. occ.; Beluchist.
caspius, Hohen. Enum. Pl. Talusch. 24 = officinalis.
caspius, *Hort. Vind. ex Schult. f. Syst.* vii. 322.—Reg. Casp.
Chesneyi, Baker, in Journ. Linn. Soc. xiv. (1875) 603 = filifolius.
collinus, Jan, ex Parl. Fl. Ital. iii. 14 = tenuifolius.
collinus, Schur, in Oestr. Bot. Zeitschr. x. (1860) 355 = officinalis.
commutatus, Tenore, Fl. Nap. iii. 374 = acutifolius.
consanguineus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 615.—Afr. austr.
Cooperi, *Baker, in Gard. Chron.* (1874) i. 818.—Afr. austr.
Corruda, Scop. Fl. Carn. ed. II. i. 248 = acutifolius.
crispus, Boj. Hort. Maurit. 350 = umbellulatus.
crispus, *Lam. Encyc.* i. 295.—Afr. austr.
Curillus, *Buch.-Ham. ex Roxb.* 24; *Fl. Ind.* ii. 152.—Reg. Himal.
Curillus, Wall. Cat. n. 5155E = filicinus.
cuscutoides, *Burch. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 606.—Afr. austr.
davuricus, *Fisch. ex Link, Enum. Hort. Berol.* i. 348.—Davuria.
declinatus, Eckl. & Zeyh. ex Baker, in Journ. Linn. Soc. xiv. (1875) 615 = consanguineus.
declinatus, *Linn. Sp. Pl.* 313.—Afr. austr.
declinatus, Sieber, ex Baker, in Journ. Linn. Soc. xiv. (1875) 621 = retrofractus.
declinatus, Schousb. Vextr. Marok. 173 = foeniculaceus.
decumbens, Jacq. Hort. Schoenb. i. 51 = crispus.
deflexus, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 254.—Angola.
densus, *Soland. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 616.—Afr. austr.
denudatus, *Baker, l. c.* 606.—Afr. austr.
dependens, Schlecht. ex Kunth, Enum. Pl. v. 85 = juniperinus.
dependens, Thunb. Prod. Pl. Cap. 66 = africanus.
dichotomus, Brouss. ex Webb & Berth. Phyt. Canar. iii. 327 = umbellatus.

ASPARAGUS :—
divaricatus, Rottl. ex Baker, in Journ. Linn. Soc. xiv. (1875) 618 = asiaticus.
divaricatus, *Royle, Illustr. Bot. Himal.* i. 393.—Ind. or.
Draco, Linn. Sp. Pl. ed. II. 451 = Dracaena Draco.
Dregeanus, *Hort.—Cf. Gard. Chron.* (1877) II. 598.—Afr. austr.
drepanophyllus, *Welw. ex. Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 254.—Afr. trop.
dubius, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 363 = racemosus.
dumosus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 608.—Sindh.
Eckloni, *Baker, l. c.* 615.—Afr. austr.
equisetifolius, Ledru, ex Webb & Berth. Phyt. Canar. iii. 328 = arborescens.
equisetoides, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 253.—Angola.
erectus, *Thunb. Prod. Pl. Cap.* 65.—Afr. austr.
esculentus, Salisb. Prod. 252 = officinalis.
euphraticus, Kunth, Enum. Pl. v. 73 = filifolius.
exaltatus, C. Sm. & Bueh, ex Webb & Berth. Phyt. Canar. iii. 326 = scoparius.
exuvialis, *Burch. Trav.* i. 432.—Afr. austr.
falcatus, Benth. in Hook. Lond. Journ. Bot. i. (1842) 493 = lueidus.
falcatus, *Linn. Sp. Pl.* 313.—Afr. trop. & austr.; Zeylan.
falcatus, Thunb. Fl. Jap. 139 = lucidus.
fasciculatus, R. Br. Prod. 281 = racemosus, acerosus.
fasciculatus, *Thunb. Fl. Cap.* ii. 329.—Afr. austr.
filicinus, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 49.—Reg. Himal.; Burma.
filifolius, *Bertol. Nov. Comm. Acad. Bonon.* v. (1842) 430 (*Misc. Bot.* i. 20).—Mesopot.
flagellaris, *Baker, in Journ. Linn. Soc.* xiv. (1875) 614.—Senegambia.
flexuosus, Thunb. Prod. Pl. Cap. 66 = crispus.
floribundus, Rottl. ex Baker, in Journ. Linn. Soc. xiv. (1875) 611 = Rottleri.
floribundus, *Royle, Illustr. Bot. Himal.* i. 393.—Ind. or.
foeniculaceus, *Lowe, in Journ. Linn. Soc.* v. (1864) 44.—Marocco.
fragilis, Burch. ex Baker, in Journ. Linn. Soc. xiv. (1875) 606 = denudatus.
Gaudichaudianus, *Kunth, Enum. Pl.* v. 71.—Cochinch.
gibbus, Bunge, Enum. Pl. Chin. Bor. 65 = davuricus.
glycycarpus, Kunth, Enum. Pl. v. 61 = davuricus.
gonocladus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 627.—Ind. or.
gracilis, *Royle, Illustr. Bot. Himal.* i. 393.—Reg. Himal.
gracilis, Salisb. Prod. 252 = asiaticus.
graminifolius, Linn. Sp. Pl. ed. II. 450 = Dracaena graminifolia.
grandiflorus, Willd. ex Webb & Berth. Phyt. Canar. iii. 327 = umbellatus.
Griffithii, *Baker, in Journ. Linn. Soc.* xiv. (1875) 604.—Afghan.
hedecarpus, Andr. ex Baker, l. c. 599 = caspius, officinalis.
horridus, Linn. f. Suppl. 203 = stipularis.
horridus, *Murr. Syst.* ed. XIII. 274.—Hab.?
hortensis, Mill. ex Baker, in Journ. Linn. Soc. xiv. (1875) 598 = officinalis.
inarimensis, Tornab. in Atti Acc. Gioenia, Ser. II. xii. (1856) 368 = acutifolius.
inderiensis, Blume, ex Ledeb. Fl. Alt. ii. 44 = maritimus.
insularis, Hanec, in Ann. Sc. Nat. Sér. V. v. (1866) 245 = lucidus.
irregularis, *Baker, in Journ. Linn. Soc.* xiv. (1875) 620.—Afr. austr.
Jacquemonti, *Baker, l. c.* 615.—Ind. or.
Krausii, *Baker, l. c.* 628.—Afr. austr.
laetus, Salisb. Prod. 253 = acthiopicus.
laevissimus, *Steud. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 623.—Ind. or.
lanceus, *Thunb. Prod. Pl. Cap.* 66.—Afr. austr.
laricinus, *Burch. Trav.* i. 537.—Afr. austr.
lignosus, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
littoralis, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) II. 92 = scaber.
longiflorus, *Franch. Pl. Davidian.* i. 300.—As. or.
longifolius, *Fisch. ex Steud. Nom.* ed. II. i. 150.—Sibiria.

ASPARAGUS :—
Lowei, *Kunth, Enum. Pl.* v. 73.—Madera.
Lownei, *Baker, in Journ. Linn. Soc.* xiv. (1875) 601.—Palaestin.
lucidus, *Lindl. Bot. Reg.* (1844) *Misc.* 29.—Japon.
Macowani, *Baker, in Journ. Linn. Soc.* xiv. (1875) 609.—Afr. austr.
madagascariensis, *Baker, l. c.* xvi. (1878) 206.—Madag.
marinus, Reichb. Ic. Fl. Germ. t. 520 = scaber.
maritimus, Mill. Gard. Dict. ed. VIII. n. 2 = scaber.
maritimus, *Pall. Reise*, ii. 329.—Rossia; Reg. Cauc.; Persia; Sibiria.
mauritianus, Lam. ex Desf. Tabl. ed. I. 20 = declinatus.
medeoloides, *Thunb. Prod. Pl. Cap.* 66.—Afr. austr.
micranthus, Lindl. Swan Riv. App. 58 = Corynotheca dichotoma.
micranthus, Sieb. & Zucc. ex Baker, in Journ. Linn. Soc. xiv. (1875) 604 = schoberioides.
microphyllus, Burch. ex Baker, l. c. 619 = africanus.
microraphis, *Baker, l. c.* 612.—Afr. austr.
minutiflorus, *Baker, l. c.* 616.—Afr. austr.
mitis, *A. Rich. Tent. Fl. Abyss.* ii. 319.—Abyss.
monophyllus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 604.—Beluchist.
mozambicus, *Kunth, Enum. Pl.* v. 72.—Afr. trop.
multiflorus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 610.—Afr. austr.
neglectus, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 750 = trichophyllus.
Nelsoni, *Baker, in Journ. Linn. Soc.* xiv. (1875) 617.—Afr. austr.
nepalensis, *Baker, l. c.* 622.—Reg. Himal.
Nivcnianus, *Schult. f. Syst.* vii. 331.—Afr. austr.
nodosus, *Soland. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 608.—Afr. austr.
officinalis, *Linn. Sp. Pl.* 313.—Europ.; Reg. Caucas.; Sibir.
oligoclonos, *Maxim. Prim. Fl. Amur.* 286.—Reg. Amur.
oligophyllus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 604.—As. temp.
orientalis, *Bess. Cat. Hort. Crem.* (1814).—Hab.?
oxyacanthus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 625.—Afr. austr.
oxycarpus, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) II. 92 = officinalis.
palaestinus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 602.—Palaestin.
pallidus, Salisb. Prod. 252 = albus.
paragus, Gueldenst. ex Ledeb. Fl. Ross. iv. 197 = officinalis.
parviflorus, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 130.—Sibir. baical.
Pastorianus, Webb & Berth. Phyt. Canar. iii. 329. t. 229 = albus.
Pauli-Guilelmi, *Solms, in Schweinf. Beitr. Fl. Aethiop.* 203.—Afr. trop.
pectinatus, [DC. in] Red. Lil. t. 407 = scandens.
penduliflorus, Zipp. ex Span. in Linnaea, xv. (1841) 476 = dubius.
persicus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 603.—Persia.
Petersianus, *Kunth, Enum. Pl.* v. 72.—Afr. trop.?
Petitianus, *A. Rich. Tent. Fl. Abyss.* ii. 320.—Abyss.
phyllacanthus, Lam. Encyc. i. 296 = aphyllus.
pilosus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 610.—Afr. trop.
planiuscula, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
plocamoides, Webb, in Steud. Nom. ed. II. i. 150 = scoparius.
plumosus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 613.—Afr. austr.
polyphyllus, Stev. ex Ledeb. Fl. Ross. iv. 198 = officinalis.
prostratus, Dum. Fl. Belg. 138 = officinalis.
psilurus, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 253.—Angola.
puberulus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 618.—Afr. austr.
pubescens, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 254.—Angola.
purpurascens, Bieb. ex Baker, in Journ. Linn. Soc. xiv. (1875) 602 = trichophyllus.
racemosus, *Willd. Sp. Pl.* ii. 152.—Ind. or.; Afr. trop.; Austral.
ramosissima, *Baker, in Gard. Chron.* (1874) II. 6.—Afr. austr.
retrofractus, *Forsk. Fl. Aegypt. Arab.* 73.—Arabia.

ASPARAGUS :—
retrofractus, *Linn. Sp. Pl.* 313.—Afr. austr.
retrofractus, C. Sm. ex Webb & Berth. Phyt. Canar. iii. 328 = arborescens.
retrofractus, *Schousb. Vextr. Marok.* 174 (sp. dubia).—Marocco.
rivalis, *Burch. ex Schult. f. Syst.* vii. 337.—Afr. austr.
Rottleri, *Baker, in Journ. Linn. Soc.* xiv. (1875) 611.—Ind. or.
ruber, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
rubricaulis, *Baker, in Journ. Linn. Soc.* xiv. (1875) 616.—Ind. or.
rubicundus, Berg. Desc. Pl. Cap. 88 = africanus, capensis, stipulaceus.
sarmentosus, Heyne, in Wall. Cat. n. 5154 = volubilis.
sarmentosus, *Linn. Sp. Pl.* 314.—Ind. or.
sativus, Mill. Gard. Dict. ed. VIII. Add. = officinalis.
scaber, *Brign. Fasc. Pl. Forojul.* 22.—Reg. Mediterr. et Caucas.
scaber, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 11 = umbellatus.
scaberulus, *A. Rich. Tent. Fl. Abyss.* ii. 320.—Abyss.
scandens, Gueldenst. ex Ledeb. Fl. Ross. iv. 199 = verticillatus.
scandens, *Thunb. Prod. Pl. Cap.* 66.—Afr. austr.
schoberioides, *Kunth, Enum. Pl.* v. 70.—Japon.
Schweinfurthii, *Baker, in Journ. Linn. Soc.* xiv. (1875) 616.—Afr. trop.
scoparius, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 11.—Marocco; Ins. Madera.
Sieboldii, *Maxim. Prim. Fl. Amur.* 287.—Japon.
simulans, *Baker, in Journ. Linn. Soc.* xiv. (1875) 609.—Madag.
sinensis, Noisset, ex Baker, l. c. 607 = crispus.
Smithianus, *Webb, ex Steud. Nom.* ed. II. i. 150, nomen.—Ins. Teneriff.
spinescens, *Steud. in Schult. f. Syst.* vii. 334.—Afr. austr.
squarrosus, *J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 165.—Ins. Cap. Verd.
stachyoides, *Spreng. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 624.—Afr. austr.
stellatus, *Baker, l. c.* 612.—Afr. austr.
stipulaceus, *Lam. Encyc.* i. 296.—Afr. austr.
stipularis, *Forsk. Fl. Aegypt. Arab.* 72.—Reg. Mediterr.; Ins. Canar.
striatus, *Thunb. Prod. Pl. Cap.* 65.—Afr. austr.
suaveolens, *Burch. Trav.* i. 226.—Afr. austr.
subalatus, *Steud. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 614 (*err.* subulatus).—Ind. or.
subulatus, *Thunb. Prod. Pl. Cap.* 66.—Afr. austr.
sylvaticus, Burch. ex Baker, in Saund. Ref. Bot. t. 214 = virgatus.
sylvaticus, Waldst. & Kit. Pl. Rar. Hung. iii. 223 = tenuifolius.
Tauscheri, Fisch. ex Ledeb. Fl. Alt. ii. 44 = maritimus.
tenuifolius, Gilib. Exercit. ii. 459 = officinalis.
tenuifolius, *Lam. Encyc.* i. 294.—Europ. austr.
terminalis, Linn. Sp. Pl. ed. II. 450 = Cordyline terminalis.
tetragonus, Bresler, Diss. n. 15 = racemosus.
Thunbergianus, *Schult. f. Syst.* vii. 333.—Afr. austr.
triacanthus, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
tricarinatus, C. Koch, in Linnaea, xxii. (1849) 266 = verticillatus.
tricarinatus, [*Delile, in*] *Red. Lil.* t. 451.—Hab.?
trichophyllus, *Bunge, Enum. Pl. Chin. Bor.* 65.—Afghan.; As. med.; Chin. bor.
umbellatus, *Link, in Buch, Beschr. Canar. Ins.* 140.—Ins. Canar.
umbellatus, *Royle, Illustr. Bot. Himal.* i. 393.—Ind. or.
umbellulatus, *Bresler, Diss.* n. 24.—Ins. Maurit.
umbricola, *Kit. ex Kanitz, Reliq. Kitaib.* 121; *in Verh. Zool. Bot. Ges. Wien*, xiii. (1863) 545.—Europ.
undulatus, *Thunb. Prod. Pl. Cap.* 66.—Afr. austr.
vaginellatus, *Boj. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 612.—Madag.
verticillaris, Lam. Encyc. i. 297 = verticillatus.
verticillatus, *Linn. Sp. Pl.* ed. II. 450.—Ross. austr.; Orient.
virgatus, *Baker, in Saund. Ref. Bot.* t. 214.—Afr. austr.
volubilis, Georgi, Beschreib. Russ. Reich. Nachtr. 268 = verticillatus.
volubilis, Buch.-Ham. ex Wall. Cat. n. 5154 = adscendens.
volubilis, *Thunb. Prod. Pl. Cap.* 66.—Afr. austr.

ASPARAGUS :—
vulgaris, Gueldenst. ex Ledeb. Fl. Ross. iv. 197 = officinalis.
Wrightii, *A. Gray, in Mem. Am. Acad.* N. S. vi. (1858-59) 413.—Japon.
zanzibaricus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 614.—Afr. trop.

ASPASIA, Lindl. Gen. et Sp. Orch. 139 (1833). *ORCHIDEAE*, Benth. & Hook. f. iii. 560.
TROPHIANTHUS, Scheidw. in Otto & Dietr. Allg. Gartenz. xii. (1844) 218.
Biberiana, Reichb. f. in Walp. Ann. vi. 849 = seq.
Bibriana, *Reichb. f. in Bonplandia*, ii. (1854) 90.—Hab.?
epidendroides, *Lindl. in Hook. Journ. Bot.* i. (1834) 6.—Panama.
fragrans, Klotzsch, Ind. Sem. Hort. Berol. (1853) 12 = epidendroides.
lunata, *Lindl. Bot. Reg.* xxii. (1836) sub t. 1907.—Bras.
lyrata, [*Reichb. f. in*] *Bot. Zeit.* x. (1852) 721.—Bras.
papilionacea, *Reichb. f. in Gard. Chron.* (1876) II. 100.—Costa Rica.
principissa, *Reichb. f. in Bot. Zeit.* x. (1852) 637.—Veraguas.
psittacina, *Reichb. f. in Gard. Chron.* x. (1878) II. 684.—Ecuad.
variegata, *Lindl. Bot. Reg.* xxii. t. 1907.—Am. austr.

ASPASIA, E. Mey. ex Pfeiff. Nom. i. 295 (1873) = **Stachys**, Linn. (Labiat.).

ASPASIA, Salisb. Gen. Pl. Fragm. 34 (1866) = **Ornithogalum**, Linn. (Liliac.).
aurea, Salisb. l. c. = O. thyrsoides.
flavescens, Salisb. l. c. = O. thyrsoides.
lactea, Salisb. l. c. = O. lacteum.
revoluta, Salisb. l. c. = O. revolutum.
thyrsoides, Salisb. l. c. = O. thyrsoides.

ASPEGRENIA, Poepp. & Endl. Nov. Gen. et Sp. ii. 12. t. 116 (1837) = **Octomeria**, R. Br. (Orchid.).
scirpoidea, Poepp. & Endl. l. c. = O. scirpoidea.

ASPELINA, Cass. in Dict. Sc. Nat. xli. 166 (1826) = **Senecio**, Linn. (Compos.).
nivea, Cass. l. c. = S. Aspelina.

ASPERA, Columna, ex Moench, Meth. 641 (1794) = **Galium**, Tourn. (Rubiac.).
muralis, Lowe, Man. Fl. Mad. i. 399 = G. murale.
nutans, Moench, Meth. 641 = G. murale.

ASPERELLA, Humb. in Roem. & Ust. Mag. iii. VII. 5 (1790). *GRAMINEAE*, Benth. & Hook. f. iii. 1207 (Asprella).
GYMNOSTICHUM, Schreb. Beschr. Graes. ii. 127. t. 47 (1810).
HYSTRIX, Moench, Meth. 294 (1794).
Hystrix, *Humb. l. c.*—Oriens.

ASPERUGO, [Tourn.] Linn. Syst. ed. I. (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 853.
aegyptiaca, Linn. Sp. Pl. ed. II. 198 = Anchusa flava.
alba, [Pieri] in Ionios Anthol. v. 192 = procumbens.
divaricata, Murr. in Nov. Comm. Gotting. (1776) 25. t. 2 = Nonea lutea.
procumbens, *Linn. Sp. Pl.* 138.—Europ.; Arab.; Oriens; As. bor.
vulgaris, Dum.-Cours. Bot. Cult. ii. 167 = procumbens.
verrucosa, Boiss. Fl. Orient. iv. 159, in syn. = Anchusa flava.

ASPERULA, Linn. Syst. ed. I (1735). *RUBIACEAE*, Benth. & Hook. f. ii. 150.
LEPTUNIS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 396, in textu.
BLEPHAROSTEMMA, CHLOROSTEMMA, CYNANCHICA, et GALIOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 398.
acerosa, C. Koch, in Linnaea, xix. (1847) 31 = liliaciflora.
affinis, Boiss. & Huet, Diagn. Ser. II. ii. 110 = cynanchica.
algerica, Pers. Syn. i. 124 = hirsuta.
Allionii, Baumg. Enum. i. 80 = hexaphylla.
alpigena, Schur, Enum. Pl. Transs. 277 = cynanchica.
alpina, Bieb. Fl. Taur. Cauc. Suppl. 103 = cynanchica.
Aparine, *Bess. Fl. Galic.* i. 114.—Europ. austr. occ.; Oriens; Sibiria.

ASPERULA :—
arcadiensis, *Sims, Bot. Mag.* t. 2146.—Graecia.
arenicola, Reut. Cat. Pl. Genève, ed. II. 101 = cynanchica.
aristata, Griseb. Spicil. Fl. Rumel. ii. 165 = longiflora.
aristata, *Linn. f. Suppl.* 120.—Europ.; Oriens.
arundana, Ball, ex Nym. Consp. ii. 334 = pendula.
arvensis, *Linn. Sp. Pl.* 103.—Europ.; Oriens.
aspera, *Boiss. Fl. Orient.* iii. 28.—Reg. Cauc.; Persia.
asperrima, *Boiss. Elench.* 56.—Hispan.
Athoa, Boiss. Diagn. Ser. I. x. 62 = suberosa.
azurea, Jaub. & Spach, Illustr. i. 151. t. 82 = orientalis.
Boissieri, Steud. ex Boiss. Voy. Bot. Espagne, 737 = effusa.
Boissieriana, Steud. Nom. ed. II. i. 151 = praec.
brachyantha, *Boiss. Diagn.* Ser. II. vi. 66.—Persia.
breviflora, *Boiss. l. c.* Ser. I. x. 63.—Syria.
brevifolia, *Vent. Jard. Cels*, t. 63.—As. Min.
brunnea, Boiss. Fl. Orient. iii. 32 = brevifolia.
bryoides, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 106.—Lycia.
calabrica, Linn. f. Suppl. 120 = Putoria calabrica.
canescens, Vis. Fl. Dalmat. iii. 11. t. 25 = aristata.
capitata, Kit. ex Reichb. f. Ic. Fl. Germ. xvii. t. 1182 = hexaphylla.
cespitosa, Boiss. Fl. Orient. iii. 39 = liliaciflora.
chlorantha, *Boiss. & Heldr. Diagn.* Ser. II. vi. 90.—Graecia.
ciliaris, DC. Prod. iv. 582 = stylosa.
ciliata, Moench, Meth. 484 = Putoria calabrica.
ciliata, *Rochel, Banat.* t. ix. f. 22; *Regel, Ind. Sem. Hort. Petrop.* (1864) 21.—Bannat.
collina, Salisb. Prod. 58 = cynanchica.
commutata, Presl, Fl. Sic. Prod. 62 = longiflora.
conferta, *Hook. f. in Lond. Journ. Bot.* vi. (1847) 463 [bis].—Austral.
consanguinea, Klotzsch, in Bot. Ergeb. Waldem. Reise, 87 = Galium triflorum.
crassifolia, Linn. Mant. 37 = tomentosa.
cretacea, Willd. ex Roem. & Schult. Syst. iii. 529 = cynanchica.
cruciata, *Vell. Fl. Flum.* 44; i. t. 114.—Bras.
cyanea, *Vell. l. c.* 44; i. t. 113.—Bras.
cynanchica, *Linn. Sp. Pl.* 104.—Europ.; As. Min.
Danilevskiana, *Basiner, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 202.—Turkestan.
debilis, Ledeb. Ind. Hort. Dorp. Suppl. (1824) 2 = humifusa.
deficiens, *Viviani, in Ann. Stor. Nat.* iv. 231.—Corsica.
denudata, Vahl, ex Willk. & Lange, Prod. Fl. Hisp. ii. 301 = hirsuta.
digyna, Dufour, in Ann. Sc. Gén. Phyt. vii. (1820) 295 = aristata.
divaricata, *Bischoff, Del. Sem. Hort. Heidelb.* (1848) 4.—Hab.?
divergens, Boiss. Diagn. Ser. II. vi. 90 = graveolens.
dubia, Willd. ex Roem. & Schult. Syst. iii. Mant. 189 = arvensis.
effusa, *Boiss. Voy. Bot. Espagne*, 280.—Hispan.
elongata, *Schrenk, Enum. Pl. Nov.* 58.—Soongaria.
eriocarpa, Boiss. Elench. 57 = asperrima.
fallax, Ehrh. ex Steud. Nom. ed. II. i. 151 = laevigata.
fasciculata, Boiss. Diagn. Ser. I. x. 60 = stricta.
flaccida, Tenore, Fl. Nap. t. 110 = longiflora.
foetida, Pers. Syn. i. 524 = Putoria calabrica.
fragillima, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 36.—Persia.
fragrantissima, *Armst. in Trans. N. Z. Inst.* xiv. (1882) 359.—N. Zel.
galioidea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 13; et viii. (1881) 161 = galioides.
galioides, *Bieb. Fl. Taur. Cauc.* i. 101.—Europ.; Reg. Cauc.
galioides, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 312 = Galium murcicum.
galioides, Jacquem. Journ. ex Hook. f. Fl. Brit. Ind. iii. 205 = Galium triflorum.
geminifolia, *F. Muell. Fragm.* v. 147.—Austral.
glaberrima, Bunge, ex Nym. Consp. 333 = involucrata.
glabra, C. Koch, in Linnaea, xix. (1847) 31 = involucrata.
glauca, Bess. Enum. Pl. Volh. 7 = galioides.
glomerata, *Griseb. Spicil. Fl. Rumel.* ii. 166.—Oriens.
gracilis, *C. A. Mey. Verz. Pfl. Cauc.* 154.—Reg. Caucas.
graveolens, *Bieb. ex Bess. in Roem. & Schult. Syst.* iii. *Mant.* 376.—Tauria; As. Min.
Gunnii, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 463 [bis].—Austral.

ASPERULA :—
Gussonii, *Boiss. Diagn.* Ser. I. x. 63.—Sicil.
Haussknechtii, *Boiss. Pl. Or. Nov. Dec.* i. 1.—Assyria.
hexaphylla, *All. Fl. Pedem.* i. 12. t. 77. f. 3.—Ital.; Hungar.
hexaphylla, Georgi, Beschr. Russ. Reich. iii. IV. 728 = cretacea.
hirsuta, *Desf. Fl. Atlant.* i. 127.—Lusit.; Afr. bor.
hirta, *Ramond, in Bull. Soc. Philom.* (1800) 131. t. 9.—Mont. Pyren.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 87 = Galium triflorum.
humifusa, *Bess. Cat. Hort. Crem.* (1811) *Suppl.* iii. 4; *Bieb. Fl. Taur. Cauc. Suppl.* 105.—Podol.; Oriens.
incana, Bory & Chaub. Nouv. Fl. Pélop. 10 = taygetea.
incana, *Sibth. & Sm. Prod. Fl. Graec.* i. 88.—Creta.
involucrata, *Wahlenb. in Isis*, xxi. (1828) 971.—As. Min.
Jordani, Perr. & Song. ex Nym. Consp. 334 = longiflora.
Kotschyana, *Boiss. Fl. Orient.* iii. 27.—Kurdistan.
laevigata, DC. Prod. iv. 582, partim = involucrata.
laevigata, *Linn. Mant.* 38.—Europ. austr.
laxiflora, *Boiss. in Bal. ex Boiss. Fl. Orient.* iii. 37.—Pontus.
libanotica, *Boiss. Diagn.* Ser. I. x. 59.—Syria.
liliaciflora, *Boiss. l. c.* iii. 33.—As. Min.
littoralis, *Sibth. & Sm. Prod. Fl. Graec.* i. 88.—Pontus.
longiflora, Noë, ex Nym. Consp. 333 = cynanchica.
longiflora, *Waldst. & Kit. Pl. Rar. Hung.* ii. 162. t. 150.—Europ.
longifolia, Sibth. & Sm. Fl. Graec. Prod. i. 87 = Galium longifolium.
lutea, *Sibth. & Sm. l. c.* 88.—Graecia.
Lycia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 105.—Lycia.
macroclada, Huet, in Ann. Sc. Nat. Sér. III. xix. (1853) 253 = cynanchica.
macrorrhiza, Hoffmgg. & Link, Fl. Port. ii. 41. t. 84. = aristata.
maritima, Tenore, ex Nym. Consp. 334 = tomentosa.
matrisylva, Gilib. Exercit. Bot. Univ. Viln. 5 = odorata.
microphylla, *Boiss. Diagn.* Ser. I. iii. 34.—Persia.
minima, *Hook. f. in Lond. Journ. Bot.* vi. (1847) 464 [bis].—Tasmania.
minor, S. F. Gray, Nat. Arr. Brit. Pl. ii. 481 = cynanchica.
mollis, Boiss. Diagn. Ser. I. iii. 31 = arcadiensis.
molluginoides, *Reichb. Fl. Germ. Excurs.* 206.—Reg. Caucas.
montana, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 151 = cynanchica.
mucronata, *Otto, ex Steud. Nom.* ed. II. i. 151.—Hab.?
multiflora, Lapeyr. Hist. Abr. Pl. Pyr. 62 = cynanchica.
Mungieri, Boiss. & Heldr. Diagn. Ser. I. x. 61 = lutea.
muscosa, *Boiss. & Heldr. l. c.* Ser. II. ii. 109.—Thessalia.
mutica, Boiss. Fl. Orient. iii. 43 = suavis.
neglecta, Moris, Elench. Sard. ii. 4 = pumila.
neglecta, *Guss. Pl. Rar.* 70. t. 13. f. 2.—Italia.
Neilreichii, *Beck, in Verh. Zool.-Bot. Ges. Wien*, xxxii. (1883) 182.—Europ.
nemoralis, Salisb. Prod. 58 = taurina.
nitens, Guss. Pl. Rar. 70. t. 13. f. 2 = neglecta.
nitida, *Sibth. & Sm. Fl. Graec. Prod.* i. 89.—As. Min.
odora, Salisb. Prod. 57 = odorata.
odorata, *Linn. Sp. Pl.* 103.—Europ.; Oriens.
oligantha, F. Muell. ex Miq. in Nederl. Kruidk. Arch. iv. (1859) 111, 112 = Rubia syrticola.
oppositifolia, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 42.—As. centr.
orientalis, *Boiss. & Hohen. Diagn.* Ser. I. iii. 30.—As. Min.; Reg. Caucas.
ovalifolia, Boiss. & Huet, Diagn. Ser. II. ii. 113 = gracilis.
palustris, Tenore, Fl. Nap. 111 = tinctoria.
paniculata, Boiss. Elench. 56 = effusa.
paniculata, *Bunge, in Ledeb. Fl. Alt.* i. 140.—Sibir. alt.
papillosa, Kern. ex Nym. Consp. 325 = Galium aristatum.
papillosa, *Lange, in Kjoeb. Vidensk. Meddel.* 111 (*Pugill.* ii. 161).—Hispan.
pendula, *Boiss. Elench.* 55.—Hispan.

ASPERULA:—
perpusilla, *Hook. Fl. Nov. Zel.* 114.—N. Zel.
Pestalozzae, Boiss. Diagn. Ser. II. ii. 113 = graveolens.
Platygalium, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 284.—Manchur.
pontica, *Boiss. Fl. Orient.* iii. 39.—Pontus.
prostrata, *C. Koch, in Linnaea*, xxiv. (1851) 461.—Armenia.
pulvinaris, Nym. Consp. ii. 334 = lutea.
pumila, *Moris, in Mem. Acc. Torin.* xxxviii. (1835) 26.—Sardinia.
pusilla, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 464 [bis] = Gunnii.
pycnantha, Boiss. Diagn. Ser. I. x. 64 = humifusa.
pyrenaica, Linn. Sp. Pl. 104 = cynanchica.
repens, Brot. Fl. Lusit. i. 152 = hirsuta.
rigida, *Sibth. & Sm. Fl. Graec. Prod.* i. 89.—Creta.
rivalis, Sibth. & Sm. l. c. 87 = A. Aparine.
rotundifolia, Linn. Mant. ii. 230 = laevigata.
rubeola, Gratel. in Bull. Soc. Nat. Hist. Bord. i. (1826) 311 = cynanchica.
rubeola, Lam. Fl. Fr. iii. 375 = cynanchica, tinctoria.
rubioides, *Schur, Enum. Pl. Transs.* 276.—Europ.
rumelica, Boiss. Diagn. Ser. II. ii. 113 = graveolens.
rupestris, Reichb. Fl. Germ. Excurs. 205 = scutellaris.
rupicola, Jord. Pugill. 76 = cynanchica.
saxatilis, Lam. Illustr. n. 139 = nitida, pyrenaica.
scabra, Link, Enum. Hort. Berol. i. 134 = tomentosa?
scabra, Moris, Elench. Sard. ii. 4 = pumila.
scabra, J. & C. Presl, Delic. Prag. 124 = aristata.
scoparia, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 463 [bis].—Austral.
scutellaris, *Vis. Sem. Rar. Hort. Patav.* (1836).—Albania.
seticornis, *Boiss. Diagn.* Ser. I. vi. 65.—Persia.
setosa, *Jaub. & Spach, Illustr.* i. 152. t. 82. B.—Syria; Persia.
setulosa, Boiss. Diagn. Ser. I. x. 61 = littoralis.
sherardioides, *Jaub. & Spach, Illustr.* i. 153.—Persia.
sinaica, Delile, ex Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 268 = Galium sinaicum.
squarrosa, C. Koch, in Linnaea, xxiv. (1851) 461 = glomerata.
Staliana, *Vis. Fl. Dalmat.* iii. 11.—Dalmatia.
stricta, *Boiss. Diagn.* Ser. I. iii. 33.—Graecia; As. Min.
strictissima, Schur, Enum. Pl. Transs. 277 = galioides.
stylosa, *Boiss. Diagn.* Ser. I. iii. 31.—Persia.
suaveolens, Schrad. ex DC. Prod. iv. 583 = longiflora.
suavis, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* ix. 62.—Anatolia.
subalpina, Schur, Enum. Pl. Transs. 277 = cynanchica.
suberosa, Guss. Fl. Sic. Syn. i. 180 = Gussonii.
suberosa, *Sibth. & Sm. Fl. Graec. Prod.* i. 89.—Graecia.
subsimplex, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 463 [bis].—Tasmania.
subvelutina, DC. Prod. iv. 585 = Galium leiophyllum.
suffruticosa, *Boiss. & Heldr. Diagn.* Ser. II. ii. 111.—Graecia.
supina, Bieb. Fl. Taur. Cauc. i. 101 = cynanchica.
taurina, *Linn. Sp. Pl.* 103.—Europ.; Orient.
taygetea, *Boiss. & Heldr. Diagn.* Ser. I. x. 60.—Graecia.
Tenori, Wallr. ex Roem. & Schult. Syst. iii. Mant. 189, in adn. = tomentosa.
tenuiflora, Jord. Pugill. 77 = cynanchica.
tenuifolia, *Boiss. Diagn.* Ser. I. iii. 32.—As. Min.
tenuissima, *C. Koch, in Linnaea*, xxiv. (1851) 462.—Orient.
thessala, Boiss. & Heldr. Diagn. Ser. II. ii. 111 = longiflora.
tinctoria, *Linn. Sp. Pl.* 104.—Europ.
tomentosa, *Tenore, Prod. Fl. Neap.* p. xii.—Ins. Caprac.
Tournefortii, *Sieber, Reise Kreta*, i. 239; ii. 316.—Creta.
triandra, Gilib. Fl. Lituan. i. 14 = tinctoria.
trichodes, *J. Gay, in DC. Prod.* iv. 582.—Persia.
trinervia, Lam. Fl. Fr. iii. 376 = taurina.
tubiflora, Gueldst. ex Steud. Nom. ed. I. 80; ed. II. i. 151 = molluginoides.
tyraica, Bess. Enum. Pl. Volh. 41 = galioides.
umbellata, Willd. ex DC. Prod. iv. 586 = hexaphylla.

ASPHALATHUS, Burm. f. Fl. Ind. 155; et Fl. Cap. Prod. 21 (1768) = **Aspalathus**, Linn. (Legum.).

ASPHALTHIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 380 = **Psoralea**, Linn. (Legum.).

ASPHALTHIUM:—
bituminosum, Medic. Phil. Bot. i. 211 = Psoralea bituminosa.
ellipticum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 381 = seq.?
frutescens, Medic. l. c. 380 = P. bituminosa.
herbaceum, Medic. l. c. 381 = praec.?
nigrum, Medic. l. c. = praec.?
palaestinum, Medic. l. c. = P. palaestina.

ASPHALTIUM, Fourr. in Ann. Soc. Lyon, N. S. xvi. (1868) 365 = ASPHALTHIUM, Medic. = **Psoralea**, Linn. (Legum.).

ASPHODELINE, Reichb. Fl. Germ. Excurs. 116 (1830). *LILIACEAE*, Benth. & Hook. f. iii. 783.
DORYDIUM, Salisb. Gen. Pl. Fragm. 72 (1866).
HEROION, Rafin. Fl. Tellur. iv. 12 (1836).
ambigua, *J. Gay, ex Baker, in Journ. Linn. Soc.* xv. (1877) 276.—Oriens.
Balansae, *J. Gay, ex Baker, l. c.* 277.—Cilicia.
brevicaulis, *J. Gay, ex Baker, l. c.* 276.—As. Min.; Mesopot.
capillaris, *Kunth, Enum. Pl.* iv. 562.—Hab.?
cretica, Boiss. ex Baker, in Journ. Linn. Soc. xv. (1877) 276 = brevicaulis.
cretica, Vis. Fl. Dalm. i. 152 = liburnica.
damascena, *Baker, in Journ. Linn. Soc.* xv. (1877) 276.—Syria.
globifera, *J. Gay, ex Baker, l. c.* 274.—Syria.
isthmocarpa, *J. Gay, ex Baker, l. c.* 278.—As. Min.
liburnica, Reichb. Fl. Germ. Excurs. 116.—Europ. austr. or.
lutea, *Reichb. l. c.*—Reg. Mediterr. et Caucas.; As. Min.
Morisiana, Clem. in Mem. Acc. Torin. Ser. II. xvi. (1857) 326 = taurica.
parviflora, *Baker, in Journ. Linn. Soc.* xv. (1877) 276.—Byzantium.
prismatocarpa, *J. Gay, ex Boiss. Fl. Orient.* v. 320.—Cappadoc.
prolifera, *Kunth, Enum. Pl.* iv. 563.—Armenia; Reg. Cauc.
rigidiflora, *Baker, in Journ. Linn. Soc.* xv. (1877) 277.—As. Min.
Szovitsii, C. Koch, in Linnaea, xxii. (1849) 261 = tenuior.
taurica, *Kunth, Enum. Pl.* iv. 561.—Orient.
tenuior, *Ledeb. Fl. Ross.* iv. 193.—Orient; Reg. Caucas.; Persia.

ASPHODELOIDES, Moench, Meth. 634 (1794) = **Asphodelus**, Linn. (Liliac.).
ramosa, Moench, l. c. = Asphodelus ramosus.

ASPHODELOPSIS, Steud. ex Baker, in Journ. Linn. Soc. xv. (1876) 321 = **Chlorophytum**, Ker-Gawl.
arangadinensis, Steud. ex Baker, l. c. 332 = C. attenuatum.

ASPHODELUS, [Tourn.] Linn. Syst. ed. I (1735). *LILIACEAE*, Benth. & Hook. f. iii. 782.
ASPHODELOIDES, Moench, Meth. 634 (1794).
CLAUSONIA, Pomel, Mat. Fl. Atl. 1 (1874).
GETHOSYNE, Salisb. Gen. Pl. Fragm. 72 (1866).
OPHIOPRASON, Salisb. l. c. (1866).
VERINEA, Pomel, Mat. Fl. Atl. 1 (1874).
abyssinicus, Hort. ex Schult. f. Syst. vii. 500 = Albuca abyssinica.
acaulis, *Desf. Fl. Atlant.* i. 302. t. 89.—Barbaria.
aestivus, Brigant. Stirp. Rar. i. 5 = Asphodeline liburnica.
aestivus, Brot. Fl. Lusit. i. 525 = microcarpus.
aestivus, Reichb. Pl. Crit. t. 451 = tenuifolius.
affinis, Parl. Fl. Ital. ii. 596 = microcarpus.
africanus, Lam. Encyc. i. 300 = Albuca abyssinica.
africanus, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 730 = ramosus.
albus, Boiss. Voy. Bot. Espagne, 618 = cerasiferus.
albus, Mill. Gard. Dict. ed. VIII. n. 3 = ramosus.
albus, T. Nees, Gen. iv. t. 20 = fistulosus.
albus, *Willd. Sp. Pl.* ii. 133.—Europ. austr.
altaicus, Pall. in Act. Acad. Petrop. (1779) II. 258. 10 = Eremurus altaicus.
ambigens, *Jord. in Bull. Soc. Bot. Fr.* vii. (1860) 735.—Gallia.

ASPHODELUS:—
apiocarpus, Hoffmgg. ex Kunth, Enum. Pl. iv. 557 = aestivus.
approximatus, Gren. & Godr. Fl. Fr. iii. 223 = fistulosus.
Arrondeaui, Lloyd, Fl. Ouest Fr. ed. III. 317 = albus.
asiaticus, *Haw. Syn. Pl. Succ.* 332.—Asia.
Audibertii, Requien, ex Schult. f. Syst. vii. 1695 = ramosus.
australis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 124 = ramosus.
bracteatus, Willd. ex Schult. f. Syst. vii. 484 = Asphodeline taurica.
brevicaulis, Bertol. in Nov. Comm. Acad. Bonon. v. (1842) 430 (Misc. Bot. i. 20) = Asphodeline brevicaulis.
canariensis, C. Sm. & Buch, in Abh. Acad. Berl. (1816–17) 362, 368 = intermedius.
capensis, Burm. f. Fl. Cap. Prod. 10 (= Cyanella?).—Afr. austr.
capensis, Linn. Syst. ed. X. 982 = Chlorophytum elatum.
capillaris, [DC. in] Red. Lil. t. 380 = Asphodeline liburnica.
cerasiferus, *J. Gay, in Bull. Soc. Bot. Fr.* iv. (1857) 610; *et in Ann. Sc. Nat.* Sér. IV. vii. (1857) 127.—Reg. Mediterr. occ.
Chamaemoly, Hochst. ex Baker, in Journ. Linn. Soc. xv. (1877) 328 = Chlorophytum tetraphyllum.
Chambeironi, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 736 = ramosus.
clavatus, Roxb. Hort. Beng. 24; Fl. Ind. ii. 148 = tenuifolius.
clavosus, Don, ex Steud. Nom. ed. II. i. 152, sphalm. = clavatus.
collinus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 123 = ramosus.
comosus, Houtt. Plantenk. xii. 336. t. 83 = Eucomis punctata.
comosus, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 737 = ramosus.
corsicus, Jord.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 739 = ramosus.
creticus, Lam. Encyc. i. 200 = Asphodeline liburnica.
crinipes, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 729 = albus.
damascenus, Boiss. Diagn. Ser. I. xiii. 22 = Asphodeline damascena.
delphinensis, Gren. & Godr. in Mém. Soc. Emul. Doubs, Sér. II. vi. (1855) 13 = albus.
Déséglisei, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 121 = ramosus.
fistulosus, *Linn. Sp. Pl.* 309.—Reg. Mediterr.; Oriens.
fuscatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 123 = ramosus.
glaucus, Aucher, ex Boiss. Fl. Orient. v. 323 = Eremurus cappadocicus.
glaucescens, Jord.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 739 = ramosus.
gramineus, Mill. ex Steud. Nom. ed. I. 80, 273; ed. II. i. 152 = Melasphaerula graminea.
grandiflorus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 123 = ramosus.
inderiensis, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 257 = Eremurus inderiensis.
infestus, Parl. Fl. Ital. ii. 600 = microcarpus.
intermedius, Hornem. Hort. Hafn. Suppl. 131 = fistulosus.
latifolius, Breyn. ex Schult. f. Syst. vii. 485 = Wachendorfia paniculata.
liburnicus, Scop. Fl. Carn. ed. II. i. 245 = Asphodeline liburnica.
littoralis, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 729 = albus.
luteus, Linn. Sp. Pl. 309 = Asphodeline lutea.
macrocarpus, Parl. Fl. Ital. ii. 604 = albus.
major, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 204 = albus.
micranthus, Coss. & Kralik, ex Coss. in Bull. Soc. Bot. Fr. iv. (1857) 497 = viscidulus.
microcarpus, Reichb. Fl. Germ. Excurs. 117; Ic. Fl. Germ. ix. 30. n. 1116 = tenuifolius.
microcarpus, *Viviani, Fl. Cors.* 5.—Reg. Mediterr.; Ins. Canar.
Morisianus, *Parl. Fl. Ital.* ii. 596.—Italia.
neglectus, Schult. f. Syst. vii. 488 = albus.
nervosus, *Pomel, Nouv. Mat. Fl. Atl.* 385.—Afr. bor.
occidentalis, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 731 = albus.

ASPHODELUS:—
olbiensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 122 = ramosus.
Ozanoni, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 733 = ramosus.
parviflorus, Wight, Ic. t. 2062 = fistulosus.
pendulinus, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 399, 497.—Algeria; Arab.
persicus, Fisch. & Mey. ex Boiss. Fl. Orient. v. 317 = Asphodeline tenuior.
persicus, Jaub. & Spach, Illustr. ii. 2. t. 102 = Eremurus persicus.
pratensis, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. i. 203 = microcarpus.
procerus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 124 = ramosus.
proliferus, Bieb. Fl. Taur. Cauc. i. 280 = Asphodeline prolifera.
proterophyllus, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 737 = ramosus.
pyrenaicus, Jord. l. c. 732 = albus.
racemosus, Link, Enum. Hort. Berol. i. 328 = ramosus.
ramosus, Colm. ex Willk. & Lange, Prod. Fl. Hisp. i. 203 = microcarpus.
ramosus, Gouan, ex Willk. & Lange, l. c. 204 = cerasiferus.
ramosus, Lapeyr. Hist. Abr. Pl. Pyr. 188 = neglectus.
ramosus, *Linn. Sp. Pl.* ed. I. 310.—Europ. austr.
ramosus, Linn. l. c. ed. II. 444 = microcarpus, cerasiferus.
ramosus, Murr. in Nov. Comm. Gotting. (1776) 37. t. 71 = albus.
ramosus, Sturm, Deutschl. Fl. Heft vi. t. 6 = fistulosus.
refractus, Boiss. Diagn. Ser. I. xiii. (1854) 23 = pendulinus.
regius, Hort. ex Heynh. Nom. ii. 48 = Eremurus spectabilis.
repens, *Pomel, Nouv. Mat. Fl. Atl.* 256.—Afr. bor.
rigidifolius, Boiss. & Heldr. Diagn. Ser. I. xiii. 22 = Asphodeline rigidifolia.
Rouxii, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 738 = ramosus.
serotinus, Sibth. ex Jacq. Eclog. 113 = Asphodeline liburnica.
sibiricus, Hort. ex Schult. f. Syst. vii. 483 = Asphodeline lutea.
sibiricus, Sievers, in Pall. N. Nord. Beitr. vii. 257 = Eremurus altaicus.
simplex, Pers. Syn. i. 367 = albus.
sphaerocarpus, Gren. & Godr. Fl. Fr. iii. 223 = albus.
spicatus, Desf. ex Baker, in Journ. Linn. Soc. xv. (1877) 271 = ramosus.
stoechadensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 122 = ramosus.
strictus, *Burm. f. Fl. Cap. Prod.* 10.—Afr. austr.
subalpinus, Gren. & Godr. Fl. Fr. iii. 224 = albus.
Szovitsii, C. Koch, in Linnaea, xxii. (1849) 261 = Asphodeline tenuior.
tardiflorus, Jord. in Bull. Soc. Bot. Fr. vii. (1860) 736 = ramosus.
tauricus, Lodd. Bot. Cab. t. 1102 = Asphodeline tenuior.
tauricus, Pall. in Nov. Act. Petrop. x. 309; Bieb. Fl. Taur. Cauc. i. 279 = Asphodeline taurica.
tenuiflora, C. Koch, in Linnaea, xxii. (1849) 261 = Asphodeline tenuior.
tenuifolius, *Cav. in Anal. Cienc. Nat.* iii. (1801) 46.—Ind. or.
tenuior, Fisch. Cat. Hort. Gorenk. (1812) 9 = Asphodeline tenuior.
validus, Jord.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 740 = ramosus.
Villarsii, Verl. ex Jord. Billotia, 107, 109; et in Bull. Soc. Stat. Isère, Sér. II. iv. (1860) 151 = ramosus.
virgatus, Jord.—Cf. Bull. Soc. Bot. Fr. vii. (1860) 738 = ramosus.
viscidulus, *Boiss. Diagn.* Ser. I. vii. 118.—Afr. bor.; Arab.

ASPICARIA, D. Dietr. Syn. i. 2, 27 (1839) = seq.

ASPICARPA, Rich. in Mém. Mus. Par. ii. (1815) 398. t. 1; Lag. Gen. et Sp. Nov. 1 (1816). *MALPIGHIACEAE*, Benth. & Hook. f. i. 261.
Acosmus, Desv. in Desf. Cat. Hort. Par. (1829) 233.
Hartwegiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 598.—Mexic.

ASPICARPA:—
hirtella, *Rich. in Mém. Mus. Par.* ii. (1815) 399. t. 13.—Mexic.
hyssopifolia, *A. Gray*, in Bost. Journ. Nat. Hist. vi. (1850) 167.—Texas.
longipes, *A. Gray, Pl. Wright.* i. 37.—N. Mexic.
sericea, *Griseb. in Goett. Abh.* xxiv. (1879) 68.—Reg. Argent.
urens, Lag. Gen. et Sp. Nov. i. 1. n. 5 = hirtella.

ASPIDALIS, Gaertn. Fruct. ii. t. 171 (1791) = **Berkheya**, Ehrh. (Compos.).
araneosa, Gaertn. l. c. = B. cernua.

ASPIDANDRA, Hassk. in Versl. en Med. Akad. Wetensch. iv. (1855) 141. *EUPHORBIACEAE*? 'Gen. dub.'
fragrans, *Hassk. l. c.*—Java.

ASPIDANTHERA, Benth. in Hook. Journ. Bot. iii. (1841) 217 = **Ferdinandusa**, Pohl (Rubiac.).
rudgeoides, Benth. l. c.—Bras.

ASPIDEIUM, Zollik. ex DC. Prod. vii. 150 (1838) = **Chondrilla**, Tourn. (Compos.).

ASPIDISTRA, Ker-Gawl. in Bot. Reg. t. 628 (1823). *LILIACEAE*, Benth. & Hook. f. iii. 772.
Macrogyne, Link & Otto, Ic. Pl. Sel. t. 31 (1823).
Plectogyne, Link, in Otto & Dietr. Allg. Gartenz. (1834) 266.
Porpax, Salisb. Gen. Pl. Fragm. 9 (1866).
elatior, *Blume, in Hoev. & De Vriese, Tijdschr.* i. (1834) 76. t. 4.—Japon.
lurida, *Ker-Gawl. in Bot. Reg.* t. 628.—China.
lurida, Sieber, ex Presl, in Flora, xii. (1829) 568 = Wachendorfia hirsuta.
punctata, Lindl. in Bot. Reg. t. 977 = lurida.
variegata, Regel, in Flora, xxxiii. (1850) 354 = elatior.

ASPIDOCARPUS, Neck. Elem. ii. 123 (1790) = **Rhamnus**, Tourn.

ASPIDOCARYA, Hook. f. & Thoms. Fl. Ind. i. 180 (1855). *MENISPERMACEAE*, Benth. & Hook. f. i. 33.
hirsuta, *Becc. Malesia*, i. (1877) 136.—Malaya.
uvifera, *Hook. f. & Thoms. Fl. Ind.* i. 180.—Reg. Himal.

ASPIDOGLOSSUM, E. Mey. Comm. Pl. Afr. Austr. 200 (1837) = **Schizoglossum**, E. Mey. (Asclep.).
biflorum, E. Mey. l. c. 201.—Afr. austr.
fasciculare, E. Mey. l. c. 200.—Afr. austr.
heterophyllum, E. Mey. l. c.—Afr. austr.

ASPIDOPTERYS, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 266. *MALPIGHIACEAE*, Benth. & Hook. f. i. 259.
canarensis, *Dalz. in Hook. Kew Journ.* iii. (1851) 37.—Ind. or.
concava, *A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 266.—Burma.
cordata, *A. Juss. l. c.* 267.—Ind. or.
elliptica, *A. Juss. l. c.* 266.—Ind. or.
glabriuscula, A. Juss. l. c. 267 = Roxburghiana.
glomerata, Wight, Ic. t. 1986 = canarensis.
Helferiana, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) ii. 137.—Burma.
hirsuta, *A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 267.—Burma.
lanuginosa, A. Juss. l. c. = nutans.
nutans, A. Juss. l. c. = Wallichii.
nutans, *Hook. f. Fl. Brit. Ind.* i. 421.—Reg. Himal.
oxyphylla, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 267 = Roxburghiana.
rotundifolia, A. Juss. l. c. = nutans.
Roxburghiana, *A. Juss. l. c.*—Ind. or.
tomentosa, *A. Juss. l. c.*—Java.
Wallichii, *Hook. f. Fl. Brit. Ind.* i. 421.—Reg. Himal.

ASPIDOSPERMA, Mart. & Zucc. Nov. Gen. et Sp. i. 57. t. 34 ad 36 (1824). *APOCYNACEAE*, Benth. & Hook. f. ii. 702.
Macaglia, Rich. ex Vahl, in Skrivt. Nat. Selsk. Kiobenh. vi. (1810) 107.
Peltospermum, DC. in Bibl. Univ. Genèv. xvii. (1838) 133.

ASPIDOSPERMA:—
anomalum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 61.—Bras.
argenteum, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 104.—Bras.
australe, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 58.—Bras.
bicolor, *Mart. Nov. Gen. et Sp.* i. 60.—Bras.
camporum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 49.—Bras.
condylocarpon, *Muell. Arg. l. c.* 55.—Bras.
cylindrocarpum, *Muell. Arg. l. c.* 54.—Bras.
dasycarpon, *A. DC. in DC. Prod.* viii. 396.—Bras.
decipiens, *Muell. Arg. in Linnaea*, xxx. (1859–60) 398.—Venezuela.
desmanthum, *Benth. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 52.—Bras.
discolor, *A. DC. in DC. Prod.* viii. 398.—Bras.
dispermum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 60.—Bras.
eburneum, *Allem. ex Saldanha, in Ann. Sc. Nat.* Sér. V. xix. (1874) 213.—Bras.
excelsum, *Benth. in Hook. Journ. Bot.* iii. (1841) 245.—Guiana.
Francisii, *A. DC. in DC. Prod.* viii. 398.—Guiana.
Gardneri, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 47.—Bras.
Gomezianum, *A. DC. in DC. Prod.* viii. 397.—Bras.
Hilarianum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 50.—Bras.
lagoënse, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 102.—Bras.
latisiliqua, *A. DC. in DC. Prod.* viii. 676.—Guiana.
leucomelanum, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 105.—Bras.
Lhotzkianum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 60.—Bras.
macrocarpon, *Mart. Nov. Gen. et Sp.* i. 59.—Bras.
macrophyllum, *Muell. Arg. in Linnaea*, xxx. (1859–60) 397.—Guiana.
Martii, *Silva Manso, in Mart. Herb. Fl. Bras.* 163.—Bras.
megalocarpon, *Muell. Arg. in Linnaea*, xxx. (1859–60) 400.—Mexic.
melanocalyx, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 52.—Bras.
molle, Mart. Herb. Fl. Bras. 162 = pyrifolium.
multiflorum, *A. DC. in DC. Prod.* viii. 397.—Guiana.
nitidum, *Benth. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 59.—Bras.
nobile, *Muell. Arg. l. c.* 44.—Bras.
oblongum, *A. DC. in DC. Prod.* viii. 399.—Guiana.
obscurum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 48.—Bras.
olivaceum, *Muell. Arg. l. c.* 57.—Bras.
pachypterum, *Muell. Arg. l. c.* 51.—Bras.
pallidiflorum, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 101.—Bras.
parvifolium, *A. DC. in DC. Prod.* viii. 398.—Bras.
platyphyllum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 46.—Bras.
polyneuron, *Muell. Arg. l. c.* 57.—Bras.
Pohlianum, *Muell. Arg. l. c.* 44.—Bras.
populifolium, *A. DC. in DC. Prod.* viii. 397.—Bras.
pyricollum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 58.—Bras.
pyrifolium, *Mart. Nov. Gen. et Sp.* i. 60.—Bras.
Quebracho-blanco, *Schlecht. in Bot. Zeit.* xix. (1861) 137.—Am. austr.
Quebracho-colorado, *Schlecht. l. c.* 139.—Am. austr.
ramiflorum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 55.—Bras.
refractum, *Mart. Nov. Gen. et Sp.* i. 60.—Bras.
Riedelii, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 56.—Bras.
Sellowii, *Muell. Arg. l. c.*—Bras.
sessiliflorum, *Muell. Arg. in Linnaea*, xxx. (1859–60) 399.—Ins. Trinit.
Spruceanum, *Benth. ex Muell. Arg. in Mart. Fl. Bras.* vi. 1. 52.—Bras.
subincanum, *Mart. Herb. Fl. Bras.* n. 262; *ex A. DC. in DC. Prod.* viii. 397.—Bras.
tomentosum, *Mart. Nov. Gen. et Sp.* i. 58. t. 34.—Bras.
Vargasii, *A. DC. in DC. Prod.* viii. 399.—Venezuela.
velutinum, Fisch. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 45 = tomentosum.
verbascifolium, *Muell. Arg. l. c.* 46.—Bras.

ASPIDOSPERMA:—
venosum, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 103.—Bras.
verruculosum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 51.—Bras.
Warmingii, *Muell. Arg. in Kjoeb. Vidensk. Meddel.* (1869) 101.—Bras.

ASPIDOSTIGMA, Hochst. in Flora, xxvii. (1844) 18 = **Toddalia**, Juss. (Rutac.).
acuminatum, Hochst. l. c. = T. nobilis.

ASPILIA, Thou. Gen. Nov. Madag. 12 (1806). *COMPOSITAE*, Benth. & Hook. f. ii. 371.
ANOMOSTEPHIUM, DC. Prod. v. 560 (1836).
CORONOCARPUS, Schum. & Thonn. Beskr. Guin. Pl. 393 (1827).
DIPTEROTHECA, Sch. Bip. in Flora, xxv. (1842) 434.
HARPEPHORA, Endl. Gen. Suppl. i. 1382 (1841).
WIRTGENIA, Sch. Bip. in Flora, xxv. (1842) 435.
abyssinica, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 379.—Afr. trop.
abyssinica, *Vatke, in Linnaea*, xxxix. (1875) 495.—Abyss.
angustifolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 380.—Afr. trop.
anomala, *Baker, in Mart. Fl. Bras.* vi. III. 204.—Bras.
aristata, *Griseb. in Goett. Abh.* xxiv. (1879) 191.—Reg. Argent.
asperrima, *Baker, in Mart. Fl. Bras.* vi. iii. 200.—Bras.
attenuata, *Baker, l. c.* 199.—Bras.
aurantiaca, *Griseb. in Goett. Abh.* xix. (1874) 183.—Reg. Argent.
Baroni, *Baker, in Journ. Linn. Soc.* xx. (1884) 188.—Madag.
Bojeri, *DC. Prod.* v. 561.—Madag.
buphthalmiflora, *Griseb. in Goett. Abh.* xix. (1874) 136.—Reg. Argent.
Burchellii, *Baker, in Mart. Fl. Bras.* vi. III. 194.—Bras.
calendulacea, Griseb. in Goett. Abh. xxiv. (1879) 191 = buphthalmiflora.
Clausseniana, *Baker, in Mart. Fl. Bras.* vi. III. 202.—Bras.
ecliptaefolia, *Baker, l. c.* 197.—Bras.
elegans, *Benth. & Hook. f. Gen.* ii. 372.—Bras.
elliptica, *Baker, in Mart. Fl. Bras.* vi. III. 192.—Bras.
floribunda, *Baker, l. c.* 198.—Bras.
foliacea, *Baker, l. c.* 193.—Bras.
foliosa, *Benth. & Hook. f. Gen.* ii. 372.—Bras.
fruticosa, *Baker, in Mart. Fl. Bras.* vi. III. 204.—Bras.
Gayana, *Benth. & Hook. f. Gen.* ii. 372.—Afr. trop.
glabra, *Benth. & Hook. f. l. c.*—Bras.
gracilis, *Baker, in Mart. Fl. Bras.* vi. III. 198.—Bras.
helianthoides, *Benth. & Hook. f. Gen.* ii. 372.—Afr. trop.
hirsuta, *Benth. & Hook. f. l. c.*—Bras.
hispidula, *Baker, in Mart. Fl. Bras.* vi. III. 201.—Bras.
Kotschyana, *Benth. & Hook. f. Gen.* ii. 372.—Afr. trop.
laevissima, *Baker, in Mart. Fl. Bras.* vi. III. 202.—Bras.
latifolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 379.—Afr. trop.
linearifolia, *Baker, in Mart. Fl. Bras.* vi. III. 192.—Bras.
linearifolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 380.—Afr. trop.
Martii, *Baker, in Mart. Fl. Bras.* vi. III. 195.—Bras.
mexicana, *Benth. & Hook. f. Gen.* ii. 372.—Mexic.
multiflora, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 378.—Afr. trop.
oblonga, *Baker, in Mart. Fl. Bras.* vi. III. 198.—Bras.
ovalifolia, *Baker, l. c.* 199.—Bras.
pascaloides, *Griseb. in Goett. Abh.* xxiv. (1879) 191.—Reg. Argent.
phyllostachya, *Baker, in Mart. Fl. Bras.* vi. III. 201.—Bras.
podophylla, *Baker, l. c.* 200.—Bras.
Pohlii, *Baker, l. c.* 201.—Bras.
Prieureana, *Benth. & Hook. f. Gen.* ii. 372.—Afr. trop.
procumbens, *Baker, in Mart. Fl. Bras.* vi. III. 194.—Bras.
pusilla, *Baker, l. c.* 193.—Bras.
reflexa, *Baker, l. c.* 196.—Bras.
reticulata, *Baker, l. c.* 202.—Bras.
Riedelii, *Baker, l. c.* 196.—Bras.

ASPILIA:—
rudis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 380.—Afr. trop.
Schimperi, *Oliver & Hiern, l. c.* 379.—Afr. trop.
serrulata, *Baker, in Mart. Fl. Bras.* vi. III. 204.—Bras.
setosa, *Griseb. in Goett. Abh.* xxiv. (1879) 192.—Reg. Argent.
silphioides, *Benth. & Hook. f. Gen.* ii. 372.—Bras.
Smithiana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 380.—Afr. trop.
squarrosa, *Baker, in Mart. Fl. Bras.* vi. III. 203.—Bras.
strigosa, *Benth. & Hook. f. Gen.* ii. 372.—Mexic.
subalpestris, *Baker, in Mart. Fl. Bras.* vi. III. 202.—Bras.
subpetiolata, *Baker, l. c.* 203.—Bras.
Thouarsii, *DC. Prod.* v. 561.—Madag.
tomentosa, *Baker, in Mart. Fl. Bras.* vi. III. 199.—Bras.
Warmingii, *Baker, l. c.* 192.—Bras.
wedeliaeformis, *Vatke, in Oestr. Bot. Zeitschr.* xxvii. (1877) 197.—Afr. trop.

ASPILOBIUM, Soland. ex A. Cunn. in Ann. Nat. Hist. (Sept. 1838) 45 = **Geniostoma**, Forst. (Logan.).

ASPILOTUM, S[oland.] ex Steud. Nom. ed. II. i. 669, 152 (1840) = **Geniostoma**, Forst. (Logan.).
laevigatum, S[oland.] ex Steud. l. c. = G. ligustrifolium.

ASPITIUM, Neck. ex Steud. Nom. ed. II. ii. 10 (1841) = **Laserpitium**, Linn. (Umbell.).

ASPLA, Reichb. Nom. 50 (1841) err. typ. = AOPLA, Lindl. = **Habenaria**, Linn. (Orchid.).

ASPRELLA, Host, Gram. Austr. iv. 17. t. 29 (1809) = **Psilurus**, Trin. (Gram.).
nardiformis, Host, l. c. = P. nardoides.

ASPRELLA, Willd. Enum. Hort. Berol. 132 (1809) = **Asperella**, Humb. (Gram.).

ASPRELLA, Schreb. Gen. 45 (1789) = **Leersia**, Sw. (Gram.).
australis, Roem. & Schult. Syst. ii. 267 = L. hexandra.
brasiliensis, Roem. & Schult. Mant. 153 = L. brasiliensis.
digitata, Lam. Illustr. i. 167 = L. digitata.
echidnea, Rafin. in Am. Monthly Mag. (1819) 190.—Am. bor.
flexuosa, Dulac, Fl. Hautes-Pyr. 71 = L. oryzoides.
hexandra, Roem. & Schult. Mant. 551 = L. hexandra.
Hystrix, Link, Enum. Hort. Berol. 100 = Asperella Hystrix.
imbricata, Roem. & Schult. Syst. ii. 268 = L. virginica.
lenticularis, Roem. & Schult. l. c. 267 = L. lenticularis.
major, Fres. ex Steud. Nom. ed. II. i. 150 = Asperella Hystrix.
mexicana, Roem. & Schult. Mant. 551 = L. hexandra.
monandra, Roem. & Schult. l. c. = L. monandra.
oryzoides, Lam. Illustr. i. 167; Beauv. Agrost. 2. t. 4. f. 2 = L. oryzoides.
ovata, Roem. & Schult. Syst. ii. 267 = L. lenticularis.
purpurea, Boj. Hort. Maurit. 376 = L. hexandra.
sibirica, Trautv. in Act. Hort. Petrop. v. (1877) 132.—Sibir.
virginica, Roem. & Schult. Syst. ii. 266 = L. virginica.

ASPRIS, Adans. Fam. ii. 496 (1763) = **Aira**, Linn. (Gramin.) (= Aira praecox).

ASSA, Houtt. Plantenk. v. 275 (1776) = **Tetracera**, Linn. (Dillen.).
exotica, F. G. Gmel. Syst. 839 = T. Assa.
indica, [Christm. & Panz. in] Houtt. Pflanz. Syst. iv. 40 = T. Assa.

ASSARACUS, Haw. Monog. Narciss. 4; in Sweet, Brit. Flow. Gard. iv. (1838) = **Narcissus**, Linn. (Amaryll.).
capax, Haw. l. c. = N. triandrus.
reflexus, Haw. l. c. = N. triandrus.

ASSONIA, Cav. Diss. iii. 120. t. 42 (1786) = **Dombeya**, Cav. (Stercul.).
Erythroxylon, Sims, Bot. Mag. sub t. 1000 = Trochetia Erythroxylon.
populnea, Cav. Diss. iii. 120. t. 42. f. 3 = D. populnea.
viburnoides, DC. Prod. i. 498.—Ins. Borbon.

ASTARTEA, DC. in Dict. Class. xi. 400 (1826); Prod. ii. 210 (1828). *MYRTACEAE*, Benth. & Hook. f. i. 702.
ambigua, *F. Muell. Fragm.* ii. 32.—Austral.
aspera, Schau. in Linnaea, xvii. (1843) 242 = fascicularis.
clavulata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 333.—Austral.
corniculata, Schau. in Lehm. Pl. Preiss. i. 113 = fascicularis.
Endlicheriana, Schau. in Linnaea, xvii. (1843) 242 = fascicularis.
fascicularis, *DC. Prod.* iii. 210.—Austral.
glomerulosa, Schau. in Linnaea, xvii. (1843) 242 = fascicularis.
intratropica, *F. Muell. Fragm.* i. 83.—Austral.
laricifolia, Schau. in Linnaea, xvii. (1843) 242 = fascicularis.
leptophylla, Schau. l. c. = fascicularis.
muricata, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 334.—Austral.
scoparia, Schau. in Linnaea, xvii. (1843) 242 = fascicularis.

ASTELIA, Banks & Soland. ex R. Br. Prod. 291 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 781.
FUNCKIA, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 19.
HAMELINIA, A. Rich. Voy. Astrolob.; Fl. Nouv. Zél. 158. t. 24 (1832).
albicans, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 252.—N. Zel.
alpina, *R. Br. Prod.* 291.—Austral.
Banksii, *A. Cunn. in Hook. Comp. Bot. Mag.* ii. (1836) 374.—N. Zel.
Cunninghamii, *Hook. f. Fl. N. Zel.* i. 259.—N. Zel.
fragrans, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 333.—N. Zel.
grandis, *Hook. f. ex T. Kirk, in Trans. N. Z. Inst.* i. (1868) 93.—N. Zel.
Hookeriana, T. Kirk, l. c. iv. (1872) 244 = Cunninghamii.
linearis, *Hook. f. Fl. Antarct.* 76.—Ins. Antarct.
Menziesiana, *Sm. in Rees, Cycl.* xxxix. *App.*; *Schult. f. Syst.* vii. 1506.—Ins. Sandvic.
microsperma, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 251.—N. Zel.
montana, *Seem. Fl. Vit.* 313.—Ins. Fiji.
nervosa, *Banks & Soland. ex Hook. f. Fl. N. Zel.* i. 260.—N. Zel.
polyneuron, *Colenso, in Trans. N. Z. Inst.* xiv. (1882) 233.—N. Zel.
psychrocharis, F. Muell. in Hook. Kew Journ. viii. (1856) 332 = alpina.
pumila, *R. Br. Prod.* 291.—Terr. Magell.
Ricardi, Kunth, Enum. Pl. iii. 365 = Banksii.
sericea, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 261 = Banksii.
Solandri, *A. Cunn. in Hook. Comp. Bot. Mag.* ii. (1836) 374.—N. Zel.
spicata, *Colenso, in Trans. N. Z. Inst.* xiv. (1882) 335.—N. Zel.
stylosa, F. Muell. in Hook. f. Fl. Tasm. ii. 61 = Milligania stylosa.
trinervia, *T. Kirk, in Trans. N. Z. Inst.* i. (1868) 93.—N. Zel.
veratroides, *Gaudich. in Freyc. Voy. Bot.* 420. t. 31.—Ins. Sandvic.
Waialealia, *Wawra, in Flora*, lviii. (1875) 243.—Ins. Sandvic.

ASTELMA, R. Br. in Bot. Reg. sub t. 532 (1820) = **Helichrysum**, Vaill. et **Helipterum**, DC. (Compos.).
canescens, D. Don, in Mem. Wern. Soc. v. (1824) 545 = Helipterum canescens.
crassifolium, D. Don, l. c. 543 = Helipterum ferrugineum.
eximium, R. Br. in Bot. Reg. sub t. 532 = Helipterum eximium.
fragrans, Sweet, Hort. Brit. ed. I. 222 = Helichrysum fragrans.
fruticans, Ker-Gawl. Bot. Reg. t. 726 = Helichrysum fruticans.
imbricatum, D. Don, in Mem. Wern. Soc. v. (1824) 544 = Helichrysum imbricatum.
lineare, D. Don, l. c. = Helichrysum ericaefolium.
milleflorum, D. Don, l. c. 542 = Helipterum phlomoides.

ASTELMA :—
modestum, Sieber, ex Hook. Bot. Mag. t. 2710 = Helipterum gnaphaloides.
proteoides, D. Don, in Mem. Wern. Soc. v. (1824) 543. —Afr. austr.
retortum, D. Don, ex Steud. Nom. ed. II. i. 152 = Helichrysum retortum.
speciosissimum, D. Don, in Mem. Soc. Wern. v. (1824) 544 = Helipterum speciosissimum.
spirale, G. Don, in Sweet, Hort. Brit. ed. III. 378 = Helipterum variegatum.
Staehelina, D. Don, in Mem. Wern. Soc. v. (1824) 543 = Helipterum speciosissimum.
variegatum, D. Don, l. c. 544 = Helipterum variegatum.
virgatum, G. Don, in Sweet, Hort. Brit. ed. III. 378 = Helipterum virgatum.

ASTEMMA, Less. Syn. Comp. 216 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 356.
dubium, *Less. l. c.*—Ecuador.

ASTEMON, Regel, Ind. Sem. Hort. Petrop. (1860) 38 = **Bystropogon**, L'Hérit. (Labiat.).
graveolens, Regel, l. c. (*an* Hyptis suaveolens ?).—Bolivia.

ASTEPHANANTHES, Bory, in Ann. Gén. Sc. Phys. ii. (1819) 138 = **Passiflora**, Linn.
bilobata, Bory, l. c. = P. bilobata.

ASTEPHANUS, R. Br. in Mem. Wern. Soc. i. (1809) 54. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 747.
HAEMAX, E. Mey. Comm. Pl. Afr. Austr. 223 (1837).
TYLODONTIA, Griseb. Cat. Pl. Cub. 175 (1866).
arenarius, *Decne. in DC. Prod.* viii. 507.—Madag.
badius, *E. Mey. Comm. Pl. Afr. Austr.* 224.—Afr. austr.
Berterii, *Spreng. Syst.* i. 855.—N. Granat.
cernuus, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 342.—Ins. Pemba.
cordatus, *R. Br. in Mem. Wern. Soc.* i. (1809) 54.—Afr. austr.
cordifolius, *Phil. in Anal. Univ. Chil.* ii. (1862) 399 ; *et Linnaea*, xxxiii. (1864-65) 175.—Chili.
cubensis, *H. B. & K. Nov. Gen. et Sp.* iii. 206. t. 237. —Cuba.
Dregei, *D. Dietr. Syn. Pl.* ii. 909.—Afr. austr.
dumosus, D. Dietr. l. c. = Rhyssolobium dumosum.
frutescens, *E. Mey. Comm. Pl. Afr. Austr.* 223.—Afr. austr.
fruticosus, Steud. Nom. ed. II. i. 153 = frutescens.
Gardneri, *Fourn. in Mart. Fl. Bras.* vi. IV. 199.—Bras.
geminiflorus, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 342.—Chili.
lanceolatus, *R. Br. in Mem. Wern. Soc.* i. (1809) 54. —Afr. austr.
linearis, *R. Br. l. c.*—Afr. austr.
marginatus, *Decne. in DC. Prod.* viii. 508.—Afr. austr.
Massoni, *Roem. & Schult. Syst.* vi. 124.—Afr. austr.
mitophorus, *Griseb. in Goett. Abh.* xix. (1874) 205.—Reg. Argent.
mucronatus, *Schlecht. in Linnaea*, viii. (1833) 518.—Mexic.
nigrescens, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1881) 367.—Mexic.
ovalifolius, *A. Rich. Fl. Cub. Fanerog.* ii. 95.—Cuba.
ovatus, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 342.—Madag.
pauciflorus, *E. Mey. Comm. Pl. Afr. Austr.* 224.—Afr. austr.
peruvianus, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 367.—Peruv.
recurvatus, *Klotzsch, in Peters, Reise Mossamb. Bot.* 274.—Afr. trop.
Schimperi, *Vatke, in Oestr. Bot. Zeitschr.* xxvi. (1876) 145.—Afr. trop.
streptocarpus, *Fourn. in Ann. Sc. Nat.* Sér. VI. xiv. (1882) 367.—Paraguay.
triflorus, *R. Br. in Mem. Wern. Soc.* i. (1809) 54.—Afr. austr.
urceolatus, *Griseb. Cat. Pl. Cub.* 173.—Cuba.
utahensis, *Engelm. in Am. Natural.* ix. (1875) 349.—Am. bor. occ.
Zeyheri, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 314.—Afr. austr.

ASTER, Tourn. ex Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 271.
ACTIPSIS, AGLOTOMA, ANACTIS, DIPLACTIS, FIMBRISTIMA, GALACTELLA (GALATELLA), KALIMARES, LEIACHENIS, MESOLIGUS, TRIPOLION, VENATRIS, VIRGULUS, Rafin. Fl. Tellur. ii. 43–47 (1836).
ARCTOGERON, DC. Prod. v. 260 (1836).
BELLIDIASTRUM, Cass. in Dict. Sc. Nat. xxxvii. 458 (1835).
BIOTIA, DC. Prod. v. 264 (1836).
CALIMERIS, Nees, Gen. et Sp. Aster. 225 (1833).
CRINITA, Moench, Meth. 578 (1794).
CRINITARIA, Cass. in Dict. Sc. Nat. xxxvii. 475 (1825).
DIETERIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 300.
DIPLOPAPPUS, Cass. in Bull. Soc. Philom. (1817) 137 ; et in Dict. Sc. Nat. xiii. 308 (1819).
DOELLINGERIA, Nees, Gen. et Sp. Aster. 177 (1833).
EUCEPHALUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 298.
EURYBIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 464 (1821).
EXATHAMNUS, D. Don, ex Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 46.
GALATEA, Cass. in Bull. Soc. Philom. (1818) 165 ; et in Dict. Sc. Nat. xviii. 56 (1820).
HELEASTRUM, DC. Prod. v. 263 (1836).
HERSILEA, Klotzsch, Bot. Ergeb. Waldem. Reise Bot. 83 (1862).
HETEROCHAETA, DC. Prod. v. 282 (1836).
HOMOSTYLIUM, Nees, in Linnaea, xviii. (1844) 513.
KALIMERIS, Cass. in Dict. Sc. Nat. xxxvii. 464 (1825).
LINOSYRIS, Cass. l. c. 460, 476 (1825).
MACHAERANTHERA, Nees, Gen. et Sp. Aster. 224 (1833).
MARGARITA, Gaud. Fl. Helv. v. 335 (1829).
NOTICASTRUM, DC. Prod. v. 279 (1836).
PINARDIA, Neck. Elem. i. 5 (1790).
PSYCHROGETON, Boiss. Fl. Orient. iii. 156 (1875).
RHINACTINA, Less. in Linnaea, vi. (1831) 119.
SYMPHYOTRICHUM, Nees, Gen. et Sp. Aster. 135 (1833).
TRIPOLIUM, Nees, l. c. 152 (1833).
TURCZANINOWIA, DC. Prod. v. 257 (1836).
VIRGARIA, Rafin. ex DC. Prod. v. 243 (1836).
XYLORHIZA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 297.
abbreviatus, Nees, Syn. Aster. 16 = patulus.
abyssinicus, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 383 = Felicia Richardi.
acaulis, *Wedd. Chlor. And.* i. 189.—Peruv.
Acmellus, Pall. Reise, iii. 320 = A. Amellus.
acris, S. G. Gmel. It. i. 172. t. 38. f. 2 = dracunculoides.
acris, Lapeyr. Hist. Abr. Pl. Pyr. 518 = salignus.
acris, Leyss. Fl. Halens. 215 = A. Amellus.
acris, *Linn. Sp. Pl.* ed. II. 1228.—Europ. austr. ; Caucas.
acris, Willd. Sp. Pl. iii. 2023 = trinervis.
aculeatus, Labill. Nov. Holl. Pl. ii. 52. t. 200 = Olearia ramulosa.
acuminatus, *Michx. Fl. Bor. Am.* ii. 109.—Am. bor.
acuminatus, Nees, Gen. et Sp. Aster. 60 = patulus.
adenolasius, F. Muell. Fragm. v. 67 = Olearia adenolasia.
adenophorus, F. Muell. l. c. 78 = Olearia adenophora.
adfinis, Less. Syn. Comp. 174 = Felicia adfinis.
adnatus, *Nutt. in Journ. Acad. Philad.* vii. (1834) 82. —Am. bor.
adscendens, *Lindl. in DC. Prod.* v. 231.—Am. bor.
adulterinus, Willd. Enum. Hort. Berol. ii. 884 = Novi-Belgii.
aestivus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 203 = longifolius.
aestivus, A. Gray, Man. Bot. U. St. ed. V. 233 partim = junceus.
aestivus, Rothrock, in Wheeler, Rep. 364 = occidentalis.
aethiopicus, *Burm. f. Fl. Cap. Prod.* 27.—Afr. austr.
ageratoides, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 154 = trinervius.
Aitonii, *Forbes, Hort. Woburn.* 188.—Hab. ?
alatus, Aikin, in Eaton & J. Wright, Man. Bot. 146 = patens.
albescens, Wall. Cat. n. 2974 = Microglossa albescens.

ASTER :—
albus, Eaton & J. Wright, Man. Bot. 147 = ptarmicoides.
albus, Willd. ex Spreng. Syst. iii. 528 = A. Amellus.
alpicola, F. Muell. Fragm. v. 70 = Olearia alpicola.
alpigenus, A. Gray, in Proc. Am. Acad. viii. (1872) 389 = pulchellus.
alpinus, Gueldenst. ex Ledeb. Fl. Ross. ii. 476 = A. Amellus.
alpinus, *Linn. Sp. Pl.* 872.—Europ. ; Oriens ; As. bor.
altaicus, *Willd. Enum. Hort. Berol.* 881.—Persia ; Reg. Himal. ; As. bor.
altissimus, *Mill. Gard. Dict.* ed. VIII. n. 20.—Am. bor.
altissimus, Moench, Meth. 601 = Novae-Angliae.
alwartensis, Lodd. Cat. (1807) ex Sims, Bot. Mag. t. 2321 = Erigeron pulchellus.
alyssoides, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 198.—Mongol.
amabilis, *Wender. ex Nees, Gen. et Sp. Aster.* 282.—Hab. ?
ambiguus, Bernh. ex Nees, l. c. 138 = macrophyllus.
amelloides, Hoffm. in Goett. Gel. Anz. xx. (1800) 1325 = A. Amellus.
Amellus, Aitch. in Journ. Linn. Soc. xix. (1882) 168 = pseudamellus.
Amellus, *Linn. Sp. Pl.* 873.—Europ. ; As. Min. ; Reg. Caucas.
amethystinus, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 294.—Am. bor.
Ammani, Lindl. ex DC. Prod. v. 229 = sibiricus.
amoenus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 84.—Afr. austr.
amoenus, Lam. Encyc. i. 306, 307 = puniceus.
amplexicaulis, Lam. l. c. 304 = Novae-Angliae.
amplexicaulis, Muhl. in Willd. Sp. Pl. iii. 2046 = laevis.
amplexicaulis, Michx. Fl. Bor. Am. ii. 114 = patens.
amplus, *Lindl. in DC. Prod.* v. 236.—Am. arct.
amygdalinus, Lam. Encyc. i. 305 = umbellatus.
Andersoni, *A. Gray, in Proc. Am. Acad.* vii. (1868) 352.—Calif.
Andersonii, *E. Forst. ex Forbes, Hort. Woburn.* 188.—Am. bor.
andinus, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 290.—Am. bor.
angustifolius, Jacq. Hort. Schoenb. iii. 63 = Felicia angustifolia.
angustifolius, Lindl. in Royle, Illustr. Bot. Himal. 251. t. 58 = altaicus.
angustissimus, Tausch, in Flora, xi. (1828) 487 = Heteropappus hispidus.
angustus, Nees, Syn. Aster. 26 = paniculatus.
angustus, Schur, Enum. Pl. Transs. 306 = Kayserianus.
angustus, *Torr. & Gray, Fl. N. Am.* ii. 162.—As. bor. ; Am. bor.
annectens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 78.—Afr. austr.
annuus, Linn. Sp. Pl. 875 = Erigeron annuus.
anomalus, *Engelm. in Torr. & Gray, Fl. N. Am.* ii. 503.—Am. bor.
Ansif, *Mart. in Denkschr. Muench. Akad.* vi. (1820) 195.—Arabia.
antirrhinifolius, Salisb. Prod. 197 = Cymbalariae.
aragonensis, *Asso, Syn.* 121. t. 8. f. 2 ; *Lam. Encyc.* i. 302.—Hispan.
aragonensis, Bout. ex Willk. & Lange, Prod. Fl. Hisp. ii. 36 = Willkommii.
ardense, *Forbes, Hort. Woburn.* 188.—Hab. ?
arenarioides, *Eaton, ex A. Gray, in Proc. Am. Acad.* viii. (1873) 647.—Am. bor. occ.
argenteus, Michx. Fl. Bor. Am. ii. 111 = sericeus.
argophyllus, Labill. Nov. Holl. Pl. ii. 52. t. 201 = Olearia argophylla.
argunensis, DC. Prod. v. 227 = flaccidus.
argutus, R. Br. ex Benth. Fl. Austral. iii. 488 = Olearia arguta.
argutus, Nees, Gen. et Sp. Aster. 69 = Novi-Belgii.
Arnottii, Nees, ex Torr. & Gray, Fl. N. Am. ii. 115 = patens.
arragonensis, Auct. = aragonensis.
artemisiaeflorus, Poir. Encyc. Suppl i. 500 = Tradescanti.
artemisioides, F. Muell. Fragm. v. 65 = Olearia tubuliflora.
ascendens, Hook. Fl. Bor. Am. ii. 8 = adscendens.
asper, Buch.-Ham. ex DC. Prod. v. 277 = asperrimus.
asper, *Nees, Gen. et Sp. Aster.* 84.—Am. bor.
asperrimus, *Less. in Linnaea*, v. (1830) 143.—Mexic.
asperrimus, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 293 = grandiflorus.

ASTER :—
asperrimus, Wall. Cat. n. 2970 = trinervis.
asperugineus, Eaton, in Bot. King, Exp. 142 = Erigeron asperugineus.
asperulus, Torr. & Gray, Fl. N. Am. ii. 120 = undulatus.
asperulus, *Wall. Cat.* n. 2968, *ex Nees, Gen. et Sp. Aster.* 281.—Reg. Himal.
asterotrichus, F. Muell. Fragm. v. 79 = Olearia asterotricha.
attenuatus, Lindl. ex Hook. Comp. Bot. Mag. i. (1835) 97 = virgatus.
atticus, Pall. Reise Stath. Russ. Reich. i. 345 = A. Amellus ?
aurantius, Linn. Sp. Pl. 877 = Dyssodia appendiculata.
aureus, D. Don, Prod. Fl. Nep. 178 = Solidago aurea.
auriculatus, Schloss. ex Nym. Consp. 387 = salignus.
auritus, Lindl. in DC. Prod. v. 232 = patens.
australis, Hort. ex DC. l. c. 270 = Olearia ramulosa.
autumnalis, Wender. ex Nees, sec. Torr. & Gray, Fl. N. Am. ii. 119 = undulatus.
axillaris, F. Muell. Fragm. v. 64 = Olearia axillaris.
azureus, *Lindl. ex Hook. Comp. Bot. Mag.* i. (1835) 98. —Am. bor.
baccharoides, *Steetz, in Seem. Bot. Voy. Herald*, 385. —China.
Baldwinii, Torr. & Gray, Fl. N. Am. ii. 127 = undulatus.
Balli, *F. Muell. Fragm.* viii. 143.—Austral.
barbatus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 81.— Afr. austr.
Behrii, Schlecht. in Linnaea, xxi. (1848) 446 = Vittadinia australis.
Bellidiastrum, *Nees, ex Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 258.—Chili.
Bellidiastrum, *Scop. Fl. Carn.* ed. II. ii. 168.—Europ.
bellidiflorus, Nees, Syn. Aster. 21 = leucanthemus.
bellidiflorus, Steud. Nom. ed. II. i. 153 = spectabilis.
bellidiflorus, Willd. Enum. Hort. Berol. 886 = paniculatus.
bellidioides, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 177 = Erigeron bellidioides.
bellidioides, *Baker, in Mart. Fl. Bras.* vi. III. 24.— Bras.
benghalensis, Heyne, ex DC. Prod. v. 292 = Erigeron asteroides.
Benthami, *F. Muell. Census*, 78.—Austral.
Benthamii, Steetz, in Scem. Bot. Voy. Herald, 385 = trinervius.
berberifolius, A. Cunn. ex DC. Prod. v. 268 = Olearia myrsinoides.
Bergerianus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 78. —Afr. austr.
bessarabicus, Bernh. ex Reichb. Fl. Germ. Excurs. 246 = A. Amellus.
bicolor, Dietr. ex DC. Prod. v. 239 = longifolius.
bicolor, Nees, Syn. Aster. 25; Gen. et Sp. Aster. 283 = Solidago bicolor.
biennis, Ledeb. in Ind. Sem. Hort. Dorpat. Suppl. i. (1811) = hispidus.
biennis, Nutt. Gen. Am. ii. 155 = canescens.
biennis, Torr. in Ann. Lyc. N. York, i. (1824) 211 = commutatus, oblongifolius.
bifidus, Nees, Syn. Aster. 18 = dracunculoides.
biflorus, Michx. Fl. Bor. Am. ii. 114 = Radula.
biflorus, Presc. ex DC. Prod. v. 231 = sibiricus.
bifrons, All. Fl. Pedem. i. 197 = Inula bifrons.
bifrons, Lindl. in DC. Prod. v. 243 = diffusus.
Bigelovii, *A. Gray, in Pacif. Rail. Rep.* iv. 97. t. 10.— Am. bor. occ.
blandus, Pursh, Fl. Am. Sept. ii. 555 = puniceus.
blepharophyllus, *A. Gray, Pl. Wright.* ii. 77.—Mexic.
Bloomeri, A. Gray, in Proc. Am. Acad. vi. (1864–65) 539 = campestris.
bonariensis, Spreng. Syst. iii. 528 = Erigeron glaucum.
borealis, Provanch. Fl. Canad. i. 308 = junceus.
Borreri, *Forbes, Hort. Woburn.* 188.—Hab. ?
bostoniensis, Hort. ex Baker, in Gard. Chron. (1884) II. 744 = amethystinus.
botryanthus, *C. Koch, in Linnaea*, xxiii. (1850) 701 (= foliaceus ?).—Oriens.
Bowiei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 73.—Afr. austr.
brachycomoides, F. Muell. Fragm. v. 86 = Vittadinia brachycomoides.

ASTER :—
bracteatus, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. xxxviii. (1865) I. 387 = tataricus.
bracteolatus, Nutt. in Trans. Am. Phil. Soc. Scr. II. vii. (1841) 293 = Hallii.
breviflorus, *Phil. in Linnaea*, xxxiii. (1864–65) 131.— Chili.
brevipes, *Benth. Fl. Hongk.* 175.—China.
britannicus, All. Fl. Pedem. i. 197 = Inula britannica.
brumalis, Nees, Gen. et Sp. Aster. 70 = Novi-Belgii.
Bubonium, Scop. Fl. Carn. ed. II. ii. 173 = Inula spiraeifolia.
bupleurifolius, Hort. Monsp. ex Nees, Gen. et Sp. Aster. 286 = laevis.
bupleuroides, *Forbes, Hort. Woburn.* 188.—Hab. ?
Burchellii, Hook. f. Ic. Pl. sub t. 1056 = Commidendron Burchellii.
Burkei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 76.— Afr. austr.
cabulicus, Lindl. Bot. Reg. (1843) Misc. 62 = Microglossa albescens.
caespitosus, Hort. ex Lindl. Bot. Reg. xix. t. 1571 = Novi-Belgii.
caespitosus, *E. Forst. ex Forbes, Hort. Woburn.* 188.— Hab. ?
caffrorum, Less. Syn. Comp. 179 = Felicia caffrornm.
calcareus, *F. Muell. Census*, 78.—Austral.
calendulaefolius, *Loisel. Herb. Amat.* iv. 274.—Hab. ?
californicus, Less. in Linnaea, vi. (1831) 33 = Erigeron glaucus.
campestris, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 293.—Am. bor. occ.
camporum, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 79.—Bras.
Canbyi, Vasey, ex A. Gray, Syn. Fl. N. Am. i. 193 = foliaceus.
Candollii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 80.— Afr. austr.
canescens, *Pursh, Fl. Am. Sept.* ii. 547.—Am. bor. occ.
canus, *Waldst. & Kit. Pl. Rar. Hung.* i. 28.—Europ.
capensis, Less. Syn. Comp. 168 = rotundifolius.
Capusi, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 304.—Turkest.
caricifolius, H. B. & K. Nov. Gen. et Sp. iv. 92. t. 333 = pauciflorus.
carneus, Fries, ex Baker, in Gard. Chron. (1884) II. 680 = paniculatus.
carneus, *Nees, Syn. Aster.* 26.—Am. bor.
carneus, Torr. & Gray, partim, ex Baker, in Gard. Chron. (1884) II. 680 = salicifolius.
carnosus, *A. Gray, ex Hemsl. in Biol. Centr. Am. Bot.* ii. 120.—Am. bor. occ.
carnosus, Gilib. Fl. Lituan. i. 214 = A. Tripolium.
carolinianus, *Walt. Fl. Carol.* 208.—Am. bor.
caryophyllus, Steud. Nom. ed. II. i. 154, sphalm. = *argophyllus*.
cassiarabicus, *Maund. Bot. Gard.* vi. n. 672.—Rossia.
Cassiniae, F. Muell. Fragm. v. 68 = Olearia Cassiniae.
catalaunicus, *Willk. & Costa, in Linnaea*, xxx. (1859–60) 105.—Hispan.
caubulicus, Hook. Bot. Mag. lxxiii. (1847) App. 34 = Microglossa albescens.
caucasicus, *Willd. Sp. Pl.* iii. 2020.—Reg. Caucas.
Celmisia, F. Muell. Fragm. v. 84 = Celmisia longifolia.
chamaedrifolius, Lam. ex DC. Prod. vii. 13 = Printzia Bergii.
Chamissonis, *A. Gray, in Torr. Bot. U. St. Expl. Exped.* 341.—Calif.
Chapmani, *Torr. & Gray, Fl. N. Am.* ii. 161.—Am. bor.
Charieis, Less. Syn. Comp. 171 = strigosus.
chilensis, Nees, Gen. et Sp. Aster. 123 = Chamissonis.
chinensis, Linn. Sp. Pl. 877 = Callistephus hortensis.
chrysanthemifolius, Hort. ex Steud. Nom. ed. II. i. 154 = filifolius.
chrysanthemoides, Willd. ex Spreng. Syst. iii. 538 = tanacetifolius.
chrysocomifolius, Hort. ex Steud. Nom. ed. II. i. 154 = Nolletia chrysocomoides.
chrysocomoides, Desf. Arb. i. 232 = Nolletia chrysocomoides.
chrysocomoides, Turcz. ex DC. Prod. vii. 272 = alpinus.
chrysophyllus, A. Cunn. ex DC. l. c. v. 266 = Olearia chrysophylla.
ciliaris, Thunb. ex DC. l. c. 218 = Mairia taxifolia.
ciliatus, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 295 = phyllolepis.
ciliatus, *Walt. Fl. Carol.* 209 (sp. dub.).—Am. bor.

ASTER :—
ciliatus, Muhl. in Willd. Sp. Pl. iii. 2027 = multiflorus.
ciliolatus, Lindl. in DC. Prod. v. 235 = Lindleyanus.
coerulescens, DC. l. c. 235 = salicifolius.
Collinsii, Nutt. in Journ. Acad. Philad. vii. (1834) 82 = Sericocarpus tortifolius.
collinus, Salisb. Prod. 198 = A. Amellus.
coloradoënsis, *A. Gray, in Proc. Am. Acad.* xi. (1876) 76.—Colorado.
commutatus, *A. Gray, Syn. Fl. N. Am.* i. II. 185.— Am. bor. occ.
concinnus, Colla, Hort. Ripul. App. iii. 22. t. 12 = Novae-Angliae.
concinnus, Salisb. Prod. 197 = fruticosus.
concinnus, *Willd. Enum. Hort. Berol.* 884.—Am. bor.
concolor, *Linn. Sp. Pl.* 874.—Am. bor.
confertus, Desf. Cat. Hort. Par. ed. III. 401 = puniceus.
confertus, Nees, Syn. Aster. 21 = versicolor.
conocephalus, F. Muell. Fragm. v. 67 = Olearia conocephala.
consanguineus, Ledeb. Fl. Ross. ii. 473 = peregrinus.
conspicuus, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) 144 = tataricus.
conspicuus, *Lindl. in DC. Prod.* v. 230.— Am. bor. occ.
Conyzae, Griessel. Kleine Bot. Schrift. 122 = Inula vulgaris.
conyzoides, Desf. Cat. Hort. Par. ed. III. 175 = Felicia angustifolia.
conyzoides, Willd. Sp. Pl. iii. 2043 = Sericocarpus conyzoides.
cordifolius, Lam. Encyc. i. 307 = paniculatus.
cordifolius, *Linn. Sp. Pl.* 875.—Am. bor.
cordifolius, Michx. Fl. Bor. Am. ii. 112 = corymbosus.
cordifolius, Muhl. ex Willd. Sp. Pl. iii. 2035 = sagittifolius.
coriaceus, Forst. f. Prod. 56. n. 297 = Celmisia coriacea.
coriaceus, Hort. ex DC. Prod. v. 276 = fruticosus.
coridifolius, Michx. ex Willd. Sp. Pl. iii. 2028 = dumosus.
cornifolius, Muhl. ex Willd. l. c. 2039 = infirmus.
Cornuti, Mill. ex Nees, in DC. Prod. v. 234 = abbreviatus.
Cornuti, Wendl. ex Nees, Gen. et Sp. Aster. 58 = patulus.
corymbosus, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 207. —Am. bor.
corymbosus, Moench, Meth. Suppl. 250 = Inula germanica.
crassifolius, *F. W. Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 326.—N. Granat.
crenatus, Less. Syn. Comp. 166 = Mairia crenata.
crinitus, Linn. Pl. Rar. Afr. 21 = Athrixia heterophylla.
crinitus, Thunb. Prod. Pl. Cap. 160 = Athrixia capensis.
crispus, Forsk. Fl. Aegypt. Arab. 150 = Pulicaria crispa.
crocatus, *Bertol. Fl. Guatimal.* 34.—Guatemala.
crocifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 89.—N. Granat.
cuneatus, A. Cunn. ex Steud. Nom. ed. II. i. 154 = Olearia stellulata.
Cunninghami, *F. Muell. Census*, 78.—Austral.
Curtisii, *Torr. & Gray, Fl. N. Am.* ii. 110.—Am. bor.
Cusickii, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 99. —Am. bor. occ.
cyaneus, Hoffm. Phyt. Blaett. i. (1803) 71 = laevis.
cyanodiscus, F. Muell. Fragm. v. 82 = Olearia ramosissima.
cydoniaefolius, A. Cunn. ex DC. Prod. v. 267 = Olearia cydoniaefolia.
cylleneus, Orphan. ex Nym. Consp. 387 = alpinus.
Cymbalariae, *Ait. Hort. Kew.* ii. 197; *Thunb. Prod. Pl. Cap.* 160.—Afr. austr.
dahuricus, *Benth. ex Baker, in Gard. Chron.* (1885) I. 208.—Sibir.
debilis, Boj. ex DC. Prod. vi. 276 = Athrixia debilis.
decorus, Desf. ex DC. l. c. v. 246 = laevis.
decumbens, *Baker, in Mart. Fl. Bras.* vi. III. 24.— Bras.
decurrens, A. Cunn. ex DC. Prod. v. 269 = Olearia decurrens.
decurrens, Moench, Meth. Suppl. 251 = Inula bifrons.
deflexus, Moench, l. c. 250 = acris.
deflexus, *Tausch, in Flora*, xxii. (1839) 77.—Hab. ?
demissus, *Forbes, Hort. Woburn.* 188.—Hab. ?

ASTER :—
demissus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 79. —Afr. austr.
dentatus, Andr. Bot. Rep. t. 61 = Olearia dentata.
dentatus, Thunb. Prod. Pl. Cap. 159 = Felicia fragilis.
denudatus, Nutt. in Trans. in Am. Phil. Soc. Ser. II. vii. (1841) 292 = adscendens.
depressus, Kit. in Linnaea, xxxii. (1863) 373 = A. Tripolium.
desertorum, *Tchertk. in Bull. Soc. Nat. Mosc.* vi. (1833) 15.—Rossia.
dichotomus, Ell. Sketch, ii. 366 = reticulatus.
diffusus, DC. Prod. v. 242, partim = vimineus.
diffusus, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 205. —Am. bor.
dimorphophyllus, *Franch. & Sav. Enum. Pl. Jap.* i. 224.—Japon.
diplostephioides, *Benth. & Hook. f. Gen.* ii. 272.—Reg. Himal.
discoideus, Boiss. & Reut. Pugill. 54 = Willkommii.
discoideus, Ell. Sketch, ii. 358 = Solidago discoidea.
discoideus, *Sond. ex Harv. & Sond. Fl. Cap.* iii. 76.— Afr. austr.
distortus, Turcz. ex Avé-Lall. in Ind. Sem. Hort. Petrop. viii. 53. n. 546 = altaicus.
divaricatus, Lam. Encyc. i. 305 = acuminatus.
divaricatus, Linn. Sp. Pl. 873 = corymbosus.
divaricatus, *Nutt. in Torr. & Gray, Fl. N. Am.* ii. 163.—Am. bor. et austr.
divaricatus, Rafin. ex DC. Prod. v. 241 = divergens.
divaricatus, Spreng. Syst. iii. 529 = umbellatus.
divaricatus, Torr. & Gray, Fl. N. Am. ii. 163 = exilis.
divergens, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 205 = diffusus.
diversifolius, Michx. Fl. Bor. Am. ii. 113 = undulatus.
diversifolius, Spreng. Syst. iii. 533 = heterophyllus.
Donii, *E. Forst. ex Forbes, Hort. Woburn.* 188.— Hab. ?
Douglasii, Eaton, Bot. King, Exp. 141 = foliaceus.
Douglasii, *Lindl. in DC. Prod.* v. 239.—Am. bor. occ.
dracunculoides, Bess. Enum. Pl. Volh. 33 = acris.
dracunculoides, *Lam. Encyc.* i. 303.—Europ. austr.; As. bor.
dracunculoides, Willd. Sp. Pl. iii. 2050 = paniculatus.
Dregei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 74.—Afr. austr.
Drummondii, *Lindl. ex Hook. Comp. Bot. Mag.* i. (1835) 97.—Am. bor.
dumosus, DC. Prod. v. 241 = multiflorus.
dumosus, Hoffm. Phytogr. Blaett. i. (1803) 68 = ericoides.
dumosus, *Linn. Sp. Pl.* 873.—Am. bor.
dumosus, Nees, Syn. Aster. 31 = ericoides.
Durandii, Nutt. ex Durand, in Pacif. Rail. Rep. v. 8 = Menziesii.
dysentericus, Scop. Fl. Carn. ed. II. ii. 171 = Pulicaria dysenterica.
echinatus, Less. Syn. Comp. 177 = Felicia echinata.
Ecklonis, *Less. l. c.* 170.—Afr. austr.
Ehrenbergii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 302.—Mexic.
elaeagnifolius, A. Cunn. ex DC. Prod. v. 267 = Olearia furfuracea.
elaeophilus, A. Cunn. l. c. 269 = Olearia heterophila.
elatus, Bert. ex Steud. Nom. ed. II. i. 154 = corymbosus.
elegans, Hort. Par. ex A. Gray, Syn. Fl. N. Am. i. II. 184 = concinnus.
elegans, *Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind.* 44.—Sikkim.
elegans, Nees, Syn. Aster. 20 = A. Amellus.
elegans, *Torr. & Gray, Fl. N. Am.* ii. 159.—Am. bor. occ.
elegans, Willd. Sp. Pl. iii. 2042 = spectabilis.
Elliottii, *Torr. & Gray, Fl. N. Am.* ii. 140.—Am. bor.
ellipticus, A. Cunn. ex DC. Prod. v. 271 = Olearia elliptica.
elodes, Torr. & Gray. Fl. N. Am. ii. 136 = Novi-Belgii.
elongatus, *Thunb. Prod. Pl. Cap.* 160.—Afr. austr.
elongifolius, A. Cunn. in Ann. Nat. Hist. ii. (1839) 127, sphalm. = *elaeagnifolius*.
eminens, Lindl. in Bot. Reg. t. 1614 = Novi-Belgii.
eminens, Nees, Gen. et Sp. Aster. 87 = junceus, paniculatus.

ASTER :—
eminens, Willd. Enum. Hort. Berol. 886 = salicifolius.
Engelmannii, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 238.—Am. bor. occ.
ensifolius, Scop. Fl. Carn. ed. II. ii. 174 = Inula ensifolia.
erectus, *Forbes, Hort. Woburn.* 188.—Hab. ?
eremophilus, *Bunge, Verz. Suppl. Fl. Alt.* 103.—Sibir. altaic.
ericaefolius, Forsk. Fl. Aegypt. Arab. 150 = Athrixia ericaefolia.
ericaefolius, *Rothr. in Coulter, Bot. Gaz.* ii. (1877) 70. —Mexic.; Texas.
ericoides, Lam. Encyc. i. 304 = multiflorus.
ericoides, *Linn. Sp. Pl.* 875.—Am. bor.
erigeroides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 74. —Afr. austr.
erigeroides, Hook. & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 48 = Erigeron fruticosus.
erubescens, Sieber, ex Benth. Fl. Austral. iii. 471 = Olearia myrsinoides.
eryngiifolius, *Torr. & Gray, Fl. N. Am.* ii. 502.— Am. bor.
espenbergensis, Nees, Gen. et Sp. Aster. 36 = sibiricus.
exasperatus, Link, Enum. Hort. Berol. ii. 328 = Olearia ramulosa.
exiguifolius, F. Muell. Fragm. v. 67 = Olearia exiguifolia.
exilifolius, F. Muell. l. c. 69 = Olearia exilifolia.
exilis, *Ell. Sketch*, ii. 344.—Am. bor.
exscapus, Richards. App. Frankl. Journ. ed. II. 32 = Townsendia sericea.
expansus, Nees, Syn. Aster. 23 = laevis.
Exsul, Lindl. Swan Riv. Pl. App. 24 = Olearia rudis.
falcatus, Eaton, in Bot. King, Exped. 140, ex parte = adscendens.
falcatus, *Lindl. in DC. Prod.* v. 241.—Am. arct.
fascicularis, E. Mey. ex DC. l. c. 220 = Felicia muricata.
fastigiatus, *Fisch. in Mém. Soc. Nat. Mosc.* iii. (1812) 74.—Sibir.
fastigiatus, Ledeb. Ic. Pl. Fl. Ross. t. 161 = Hauptii.
Fendleri, *A. Gray, Pl. Fendl.* 66.—Am. bor.
Ferresii, F. Muell. Fragm. v. 75 = Olearia Ferresii.
ferrugineus, Edgew. in Trans. Linn. Soc. xx. (1846) 64 = Microglossa albescens.
ferrugineus, Wendl. in Flora, ii. (1819) 676 = Olearia dentata.
ficoidea, *Harv. in Harv. & Sond. Fl. Cap.* iii. 76.— Afr. austr.
filaginifolius, Hook. & Arn. Bot. Beech. Voy. 146 = Corethrogyne filaginifolia.
filifolius, *Vent. Jard. Malm.* 82.—Afr. austr.
filifolius, Willd. ex Nees, Gen. et Sp. Astcr. 210 = Felicia muricata.
filiformis, Eckl. ex DC. Prod. v. 218 = Mairia taxifolia.
filiformis, Steud. Nom. ed. II. i. 154 = Felicia muricata.
firmus, *Nees, Syn. Aster.* 25.—Am. bor.
Fischeri, *E. Forst. ex Forbes, Hort. Woburn.* 188.— Hab. ?
Fischerianus, *DC. Prod.* v. 247.—Sibir.
flaccidus, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 599.—Sibir.
flexuosus, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 73 = Fischerianus.
flexuosus, Nutt. Gen. Am. ii. 154 = tenuifolius.
floribundus, *Nutt. l. c.* 158.—Am. bor.
floribundus, Willd. Sp. Pl. iii. 2048 = Novi-Belgii.
florulentus, F. Muell. Fragm. v. 82 = Olearia floribunda.
fluminensis, *Vell. Fl. Flum.* viii. t. 123.—Bras.
foliaceus, *Lindl. in DC. Prod.* v. 228.—Am. bor. occ.
foliolosus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 202 = vimineus.
foliolosus, Ell. Sketch, ii. 345 = dumosus.
foliosus, Pers. Syn. ii. 443 = dumosus.
formosissimus, *Hort. ex Heynh. Nom.* ii. 51. nomen; *Vilm. Fl. Pl. Terre*, ed. I. 98.—Hab. ?
Forsteri, *Borr. ex Forbes, Hort. Woburn.* 188.— Hab. ?
fragilis, Lindl. in Bot. Reg. t. 1537 = dumosus.
fragilis, Jacq. ex Steud. Nom. ed. I. 82 = linariifolius.
fragilis, Nees, Gen. et Sp. Aster. 101 = vimineus.
fragilis, *Schloss. & Vukot. Syll. Fl. Croat.* 57.— Croatia.
fragilis, Willd. Sp. Pl. iii. 2051 = Tradescanti.

ASTER :—
Fremonti, *A. Gray, Syn. Fl. N. Am.* i. II. 191.—Am. bor. occ.
frondosus, *Torr. & Gray, Fl. N. Am.* ii. 165.—Am. bor. occ.
frutetorum, Wimm. ex Nym. Consp. ii. 387 = salignus.
fruticosus, *Linn. Sp. Pl.* 872.—Afr. austr.
fruticulosus, Sims, Bot. Mag. t. 2286 = pluriflorus.
fruticulosus, Burm. f. Prod. 27 = Felicia angustifolia.
fruticulosus, Krebs, ex DC. Prod. v. 276 = filifolius.
fruticulosus, Willd. Sp. Pl. iii. 2018, pro parte = obtusatus.
fruticulosus, Murr. Syst. Veg. ed. XIII. 633 = fruticosus.
fugax, Brot. Fl. Lusit. 385 = aragonensis.
furfuraceus, A. Rich. & Lesson, Fl. Nov. Zel. 216 = Olearia furfuracea.
Gardneri, *Baker, in Mart. Fl. Bras.* vi. III. 25.—Bras.
Garibaldii, Bruegg. in Verh. Schweiz. Naturf. Ges. Samaden, 1863 (1864) 229 = alpinus.
Gayanus, *DC. Prod.* v. 227.—Chili.
Gebleri, Bess. ex DC. l. c. 259 = altaicus.
Garlachii, *Hance, in Journ. Bot.* xviii. (1880) 262.— China.
giganteus, Hort. ex Vilm. Fl. Pl. Terre, ed I 928 = Vernonia noveboracensis.
Gilliesii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 49.—Chili.
glabellus, Nees, Syn. Aster. 31 = ericoides.
glaber, E. Mey. ex Nees, Gen. et Sp. Aster. 68 = vimineus.
glaber, *Mill. Gard. Dict.* ed. VIII. n. 17.—Am. bor.
glabriusculus, Torr. & Gray, Fl. N. Am. ii. 159 = xylorrhiza.
glacialis, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 291.—Am. bor.
glandulosus, Labill. Nov. Holl. Pl. ii. 50. t. 197 = Olearia glandulosa.
glaucescens, Wender. ex Nees, Gen. et Sp. Aster. 133 = laevis.
glaucus, Nees, Syn. Aster. 23 = laevis.
glaucus, *Torr. & Gray, Fl. N. Am.* ii. 159.—Am. bor. occ.
glomeratus, Bernh. ex DC. Prod. v. 265 = macrophyllus.
Glehni, *F. Schmidt, Reisen Amurl.* 146.—As. or.
glutescens, F. Muell. Fragm. v. 77 = Olearia glutinosa.
glutinosus, Cav. Ic. ii. 53 = Grindelia glutinosa.
glutinosus, Roxb. in Beats. App. 303 = Commidendron rugosum.
Gmelini, Ledeb. in Denkschr. Baier. Bot. Ges. iii. (1841) 77 = Senecio Gmelini.
Gmelini, Tausch, in Flora, xi. (1828) 486 = Heteropappus hispidus.
Gmelini, Tausch, l. c. xxii. (1839) I. 78 = altaicus.
gracilentus, Banks, ex DC. Prod. v. 327 = Chrysopsis villosa.
gracilentus, Torr. & Gray, Fl. N. Am. ii. 166 = virgatus.
gracilis, *Nutt. Gen. Am.* ii. 158.—Am. bor.
graminifolius, Pursh, Fl. Am. Sept. ii. 545 = Erigeron hyssopifolius.
grandiflorus, *Hoffm. Phyt. Blaett.* i. (1803) 65.— Hab. ?
grandiflorus, *Linn. Sp. Pl.* 877.—Am. bor.
grandiflorus, Nutt. Gen. Am. ii. 156 = paludosus.
graveolens, Nutt. in Journ. Ac. Nat. Sc. Phil. ii. (1821) 116 = oblongifolius.
gravis, F. Muell. Fragm. v. 82 = Olearia gravis.
Greenii, Nees, ex Torr. & Gray, Fl. N. Am. ii. 158 = nemoralis.
Greenii, Torr. & Gray, l. c. 134 = salicifolius.
gummiferus, Hook. f. Ic. Pl. t. 1056 = Commidendron spurium.
gymnocephalus, *A. Gray, in Proc. Am. Acad.* xv. (1880) 32.—Texas; Mexic.
Haenkei, *Nees, Gen. et Sp. Aster.* 63.—Chili.
Hallii, *A. Gray, in Proc. Am. Acad.* viii. (1872) 388 = Am. bor. occ.
Hauptii, *Ledeb. Fl. Alt.* iv. 100.—Sibir.
Hauptii, Turcz. ex DC. Prod. v. 256 = dahuricus.
Haydeni, Porter, in Hayden, Geol. Rep. 1871 (1872) 485 = pulchellus.
hebecladus, DC. Prod. v. 242 = multiflorus.
Helenium, Scop. Fl. Carn. ed. II. ii. 171 = Inula Helenium.
heleophilus, F. Muell. Fragm. v. 66 = Olearia heleophila.

ASTER:—
Hersilei, *Klatt, Comp. Herb. Schlag. Hochas.* 368.—As. temp.
Herveyi, *A. Gray, Man. Bot. U. St.* ed. V. 230.—Am. bor.
hesperius, *A. Gray, Syn. Fl. N. Am.* i. II. 192.—Am. bor. occ.
heterochaeta, C. B. Clarke, Comp. Ind. 44 = Heterochaeta asteroides.
heteromallus, Wender. in Schrift. Marb. ii. (1831) VI. 54 = undulatus.
heterophyllus, Nees, Syn. Aster. 15 = paniculatus.
heterophyllus, Thunb. Prod. Pl. Cap. 160 = Athrixia heterophylla.
heterophyllus, Willd. Sp. Pl. iii. 55 = cordifolius.
heterophyllus, *Schloss. & Vukot. Syll. Fl. Croat.* 55.—Croatia.
Heynei, F. Muell. Fragm. v. 86 = Olearia xerophila.
hiemalis, Nees, Gen. et Sp. Aster. 77 = longifolius.
himalaicus, *C. B. Clarke, Comp. Ind.* 43.—Reg. Himal.
hirsuticaulis, Lindl. in DC. Prod. v. 242 = diffusus.
hirsutus, Host, Fl. Austr. ii. 485 = alpinus.
hirsutus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 80.—Afr. austr.
hirtellus, Lindl. in DC. Prod. v. 233 = sagittifolius.
hirtus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 82.—Afr. austr.
hirtus, *C. Koch, in Linnaea*, xxiii. (1850) 701.—Armenia.
hirtus, Scop. Fl. Carn. ed. II. ii. 173 = Inula hirta.
hirtus, Thunb. Prod. Pl. Cap. 159 = Felicia angustifolia.
hispidus, *Baker, in Saund. Ref. Bot.* t. 342.—Afr. austr.
hispidus, Lam. Encyc. i. 307 = puniceus.
hispidus, Thunb. Fl. Jap. 315 = Heteropappus hispidus.
holosericeus, Forst. f. Prod. 56. n. 296 = Celmisia holosericea.
homolepis, F. Muell. Fragm. v. 65 = Olearia homolepis.
Hookerii, *Forbes, Hort. Woburn.* 189.—Hab.?
horizontalis, Desf. Cat. Hort. Par. ed. III. 402 = diffusus.
Huegelii, F. Muell. Fragm. v. 79 = Olearia ciliata.
humilis, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 125 = Vittadinia australis.
humilis, Willd. Sp. Pl. iii. 2038 = infirmus.
hungaricus, Poir. Encyc. Suppl. i. 496 = salignus.
Hunnemanni, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab.?
hybernus, *E. Forst. ex Forbes, l. c.*—Hab.?
hydrabadensis, Wall. Cat. n. 2973 = Pulicaria Wightiana.
hygrophilus, A. Cunn. ex DC. Prod. v. 269 = Olearia hygrophila.
hyssopifolius, Berg. Desc. Pl. Cap. 287 = Felicia angustifolia.
hyssopifolius, Cav. Ic. t. 232 = sedifolius.
hyssopifolius, Linn. Mant. 114 = aeris.
ibericus, Stev. ex Bieb. Fl. Taur. Cauc. ii. 311 = A. Amellus.
ignoratus, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 11.—Ind. or.
ilicifolius, A. Cunn. ex DC. Prod. v. 268 = Olearia myrsinoides.
illitus, F. Muell. Fragm. v. 76 = Olearia elliptica.
imbricatus, *Linn. Pl. Rar. Afr.* 21.—Afr. austr.
imbricatus, Walp. Rep. ii. 574 = arenarioides.
inaequalis, *Hill, Hort. Kew.* 16.—Am. bor.
incanus, *A. Gray, in Bot. Calif.* i. 322.—Calif.; Mexic.
incisus, *Fisch. in Mém. Soc. Nat. Mosc.* iii. (1812) 76.—Sibir.
inconspicuus, *Less. in Linnaea*, v. (1830) 143.—Mexic.
indicus, Heyne, in Wall. Cat. n. 2971 = Pulicaria Wightiana.
indicus, Linn. Sp. Pl. 876 = Boltonia indica.
infirmus, *Michx. Fl. Bor. Am.* ii. 109.—Am. bor.
inflexus, *Desf. Tabl.* ed. II. 122.—Hab.?
integrifolius, *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884). 224.—China.
integrifolius, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 291.—Am. bor. occ.
intermedius, Turcz. ex DC. Prod. vii. 272 = flaccidus.
intybaceus, Kunth & Bouché, Ind. Sem. Hort. Berol. (1845) 11 = tataricus.
inulaefolius, *Boiss. & Bal. ex Boiss. Fl. Orient.* iii. 159.—Reg. Caucas.

ASTER:—
inuloides, D. Don, Prod. Fl. Nep. 178 = Erigeron multiradiatus.
inuloides, Fisch. ex DC. Prod. v. 231 = sibiricus.
iodochrous, F. Muell. Fragm. v. 81 = Olearia iodochroa.
ircutianus, DC. Prod. v. 229 = tataricus.
italicus, *Hill, Veg. Syst.* ii. 88.—Italia.
jamaicensis, *Less. in Linnaea*, v. (1830) 144.—Mexic.; Ind. occ.
japonicus, *Less. ex Nees, Gen. et Sp. Aster.* 33, 34.—Japon.
junceus, *Ait. Hort. Kew.* iii. 204.—Am. bor.
junceus, Nees, Syn. Aster. 26 = tenuifolius.
Kayserianus, *Schur, Enum. Pl. Transs.* 306.—Europ.
Kingii, *Eaton, Bot. King, Exp.* 141. t. 16.—Am. bor. occ.
Kraussii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 84.—Afr. austr.
Kumleini, Fries, ex A. Gray, Syn. Fl. N. Am. i. II. 179 = oblongifolius.
lacerus, Lindl. ex DC. Prod. v. 231 = sibiricus.
lacunarum, *Aitch. & Hemsl. in Journ. Linn. Soc* xix. (1882) 168.—Afghan.
laetus, Fisch. ex Ledeb. Fl. Ross. ii. 479 = punctatus.
laevigatus, Hook. Bot. Mag. t. 2995 = laevis.
laevigatus, Lam. Encyc. i. 307 = Novi-Belgii.
laevigatus, Pursh, Fl. Am. Sept. ii. 553 = longifolius.
laevigatus, Willd. Sp. Pl. iii. 2046 = versicolor.
laevis, Lindl. in Bot. Reg. t. 1500 = versicolor.
laevis, *Linn. Sp. Pl.* 876.—Am. bor.
laevis, Vahl, ex DC. Prod. v. 238 = longifolius.
Laka, *C. B. Clarke, Comp. Ind.* 49.—Reg. Himal.
Lamarckianus, Nees, Gen. et Sp. Aster. 100 = paniculatus.
lanceolatus, Willd. Sp. Pl. iii. 2050 = paniculatus.
lanceolatus, Nutt. Gen. Am. ii. 158 = salicifolius.
lanuginosus, *Wendl. Beytr.* ii. 9.—Afr. austr.
lasiocarpus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 76.—Afr. austr.
lateralis, Hort. Bonn. ex Nees, Gen. et Sp. Aster. 74 = longifolius.
latifolius, Desf. Cat. Hort. Par. ed. III. 176 = macrophyllus.
latifolius, Banks, ex Steud. Nom. ed. II. i. 155 = acuminatus.
latifolius, Mill. Gard. Dict. ed. VIII. n. 26 = trinervis?
lautureanus, *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) 224.—China.
lavandulaeeus, *Willd. ex DC. Prod.* v. 274.—Mexic.
laxifolius, Lindl. in Hook. Fl. Bor. Am. ii. 10 = junceus.
laxifolius, Nees, Gen. et Sp. Aster. 69 = longifolius.
laxus, Torr. & Gray, Fl. N. Am. ii. 134 = Novi-Belgii.
laxus, Willd. Enum. Hort. Berol. ii. 857 = paniculatus.
ledifolius, A. Cunn. ex DC. Prod. v. 269 = Olearia ledifolia.
ledifolius, Pursh, Fl. Am. Sept. ii. 544 = nemoralis.
ledophyllus, A. Gray, in Proc. Am. Acad. xvi. (1880) 98 = Engelmanni.
leiocarpus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 84.—Afr. austr.
Lehmanni, F. Muell. Fragm. v. 66 = Olearia heleophila.
leiophyllus, *Franch. & Sav. Enum. Pl. Jap.* i. 223.—Japon.
leiophyllus, Schloss. & Vukot. Syll. Fl. Croat. 56 = salignus.
Lemmoni, *A. Gray, Syn. Fl. N. Am.* i. II. 199.—Arizona.
lepidophyllus, Pers. Syn. ii. 442 = Olearia lepidophylla.
leptophyllus, *Desf. Tabl.* ed. II. 268; *DC. Prod.* v. 256.—Am. bor.
leptophyllus, *Fisch. Hort. Gorenk.* ed. II. (1812) 39, nomen.—Sibir.
leptophyllus, Hort. Par. ex DC. Prod. v. 242 = ericoides.
Lerchenfeldianus, *Schur, Enum. Pl. Transs.* 306.—Hungar.
leucanthemus, Desf. Cat. Hort. Par. ed. III. 401 = Tradescanti.
leucanthemus, *Rafin. in Med. Repos. N. York*, v. (1808) 359.—Am. bor.
liatroides, Muhl. ex DC. Prod. v. 230 = surculosus.
lilaeinus, *Hoffmgg. Verz. Pfl.* 41.—Hab.?
Lima, *Lindl. in DC. Prod.* v. 230.—Mexic.
linariifolius, *Linn. Sp. Pl.* 874.—Am. bor.
Lindeni, *Sch. Bip. in Seem. Bot. Voy. Herald*, 302.—Mexic.

ASTER:—
Lindleyanus, *Torr. & Gray, Fl. N. Am.* ii. 122.—Am. bor.
linifolius, Burm. f. Prod. 27 = Senecio rosmarinifolius.
linifolius, *Harv. in Harv. & Sond. Fl. Cap.* iii. 81.—Afr. austr.
linifolius, Linn. Sp. Pl. 874 = aeris.
linifolius, Torr. & Gray, Fl. N. Am. ii. 162 = subulatus.
linoides, Hort. Gaybac. ex DC. Prod. v. 221 = Felicia caffrorum.
Linosyris, *Bernh. Syst. Verz. Erf.* 151.—Europ.
liratus, Sims, Bot. Mag. t. 1509 = Olearia stellulata.
lithospermifolius, Desf. Tabl. ed. II. 268; DC. Prod. v. 259 = altaicus?
litoralis, *Phil. in Anal. Univ. Chil.* (1873) 486.—Chili.
lividus, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab.?
longicaulis, Desf. ex DC. Prod. v. 253 = A. Tripolium.
longifolius, *Desf. Cat. Hort. Par.* ed. III. 400.—Hab.?
longifolius, A. Gray, Man. Bot. U. St. ed. V. 233 = Novi-Belgii.
longifolius, *Lam. Encyc.* i. 306.—Am. bor.
longifolius, Nees, Syn. Aster. 26 = laxifolius.
longipes, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 78 = Leucopsis scaposa.
lucidus, Hort. Kew. ex Loud. Hort. Brit. 347 = laevis.
lucidus, Wender. in Flora, xii. (1829) I. Erg. 25 = puniceus.
lucidus, *Moench, Meth. Suppl.* 252.—Hab.?
lusitanicus, Brot. Phyt. 63. t. 29 = aragonensis.
lutescens, Torr. & Gray, Fl. N. Am. ii. 160 = ptarmicoides.
luxurians, Spreng. Syst. iii. 538; Nees, Gen. et Sp. Aster. 83 = Novi-Belgii.
lyratus, Nees, Gen. et Sp. Aster. 188, in syn. = Olearia stellulata.
Maackii, *Regel, in Mém. Acad. Pétersb.* Sér. VII. iv. (1861) n. IV. 81.—Japon.
Macleaii, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab.?
macrophyllus, *Linn. Sp. Pl.* ed. II. ii. 1232.—Am. bor.
macrorrhizus, A. Cunn. ex DC. Prod. v. 260 = Vittadinia macrorrhiza.
macrorrhizus, *Thunb. Prod. Pl. Cap.* 159.—Afr. austr.
magellanicus, Lam. Illustr. iii. 259. t. 681 = Perezia lactucoides.
magellanicus, Spreng. Syst. iii. 526 = Chiliotrichium amelloideum.
magniflorus, F. Muell. Fragm. v. 80 = Olearia magniflora.
marginatus, *H. B. & K. Nov. Gen. et Sp.* iv. 91.—N. Granat.
maritimus, Salisb. Prod. 198 = A. Tripolium.
Martii, *Baker, in Mart. Fl. Bras.* vi. III. 23.—Bras.
marylandicus, Michx. Fl. Bor. Am. ii. 108 = Sericocarpus conyzoides.
maximus, Less. Syn. Comp. 182 = Erigeron maximus.
megalophyllus, F. Muell. Fragm. v. 70 = Olearia megalophylla.
megapotamicus, *Spreng. Syst.* iii. 526.—Bras.
Menziesii, *Lindl. in DC. Prod.* v. 243.—Am. bor. occ.
mesopotamicus, Steud. Nom. ed. I. 82 = megapotamicus.
mesopotamicus, *Spreng. Nachr.* i. *Bot. Gart. Halle*, 13.—Hab.?
mespilifolius, Less. Syn. Comp. 180 = Nidorella mespilifolia.
microcephalus, *Franch. & Sav. Enum. Pl. Jap.* i. 223.—Japon.
microphyllus, Labill. Nov. Holl. Pl. ii. 51 = Olearia lepidophylla.
microphyllus, Torr. ex Lindl. in DC. Prod. v. 244 = adnatus.
microphyllus, Vent. Jard. Malm. sub t. 83 = Olearia ramulosa.
microspermus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 72.—Afr. austr.
Milleri, *Forbes, Hort. Woburn.* 189.—Hab.?
mirabilis, *Torr. & Gray, Fl. N. Am.* ii. 165.—Am. bor.
miser, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 205 = Tradescanti.
miser, Lam. Encyc. i. 308 = virgatus.
miser, Linn. Sp. Pl. 877 (delend.—Cf. A. Gray, Syn. Fl. i. II. 172).

ASTER :—
miser, Nutt. Gen. Am. ii. 158 = diffusus.
Mitchellii, *F. Muell. Fragm.* v. 78.—Austral.
modestus, *Lindl. in DC. Prod.* v. 231.—Am. bor. occ.
molliusculus, *Wall. Cat.* n. 2972.—Reg. Himal.
monanthus, *Vell. Fl. Flum.* viii. t. 121.—Bras.
mongolicus, *Franch. Pl. Davidian.* i. 161.—As. or.
monocephalus, Boiss. in Bal. ex Boiss. Fl. Orient. iii. 160 = caucasicus.
montanus, All. Fl. Pedem. i. 195 = Inula montana.
montanus, Nutt. Gen. Am. ii. 155 = sericeus.
montanus, R. Br. in Richards. App. Frankl. Journ. ed. I. 749 = sibiricus.
montevidensis, Griseb. in Goett. Abh. xix. (1874) 170 = Leucopsis diffusa.
monticolus, Lindl. ex DC. Prod. v. 287 = Erigeron monticolus.
Mooneyi, *F. Muell. Fragm.* viii. 144.—Austral.
moranensis, *H. B. & K. Nov. Gen. et Sp.* iv. 93.—Mexic.
Muehlenbergii, Tausch, in Flora, xxii. (1839) 77, in adnot. = amplexicaulis.
Muelleri, *F. Muell. Census*, 78.—Austral.
multiceps, Lindl. in DC. Prod. v. 237 = oblongifolius.
multiflorus, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 116.—Am. bor.; Mexic.
multiflorus, Bertol. Misc. Bot. v. t. 5 = dumosus.
multiflorus, Nutt. Gen. Am. ii. 155 = vimineus.
multiradiatus, Wall. Cat. n. 2969 = Erigeron multiradiatus.
muricatus, F. Muell. Fragm. v. 66 = Olearia muricata.
muricatus, Thunb. Prod. Pl. Cap. 159 = Felicia muricata.
mutabilis, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 205 = Novi-Belgii.
mutabilis, *Vell. Fl. Flum.* viii. t. 122.—Bras.
mutabilis, *Linn. Sp. Pl.* 876 (sp. obscura).—Am. bor.
mutabilis, Willd. Sp. Pl. iii. 2045 = versicolor.
mutatus, Torr. & Gray, Fl. N. Am. ii. 142 = modestus.
myriantus, *Carr. in Rev. Hortic.* (1866) 439.—Hab.?
myrsinoides, Labill. Nov. Holl. Pl. ii. 53. t. 202 = Olearia myrsinoides.
myrtifolius, Willd. Enum. Hort. Berol. Suppl. 59 = miser.
namaquanus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 79.—Afr. austr.
nanus, E. Mey. ex DC. Prod. v. 218 = Mairia microcephala.
natalensis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 81.—Afr. austr.
Neesii, Sch. Bip. in Flora, xxxix. (1856) 354 = Chamissonis.
nemoralis, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 198.—Am. bor.
Nernstii, F. Muell. Fragm. v. 81 = Olearia Nernstii.
nervosus, *Mill. Gard. Dict.* ed. VIII. n. 23.—Am. bor.
nitidulus, *DC. in Wight, Contrib.* 9.—Ind. or.
nivalis, Adam, in Weber & Mohr, Beitr. i. 67 = alpinus.
Noëanus, Sch. Bip. ex Nym. Consp. 387 = A. Amellus.
notosericeus, *Griseb. in Goett. Abh.* xxiv. (1879) 178.—Reg. Argent.
Novae-Angliae, *Linn. Sp. Pl.* 875.—Am. bor.
Novae-Scotiae, *Forbes, Hort. Woburn.* 189; *Steud. Nom.* ed. II. i. 156.—N. Scotia.
Novi-Belgii, *Linn. Sp. Pl.* 877.—Am. bor.
nudicaulis, Lam. Encyc. i. 308 = Lagenophora Commersonii.
nudiflorus, Nutt. Gen. Am. ii. 157 = A. Radula.
Nuttallii, Torr. & Gray, Fl. N. Am. ii. 126 = adscendens.
obcordatus, F. Muell. Fragm. v. 69 = Olearia obcordata.
obliquus, DC. Prod. v. 237 = carneus.
obliquus, Nees, Syn. Aster. 27 = salicifolius.
oblongifolius, *Nutt. Gen. Am.* ii. 156.—Am. bor.
obovatus, Ell. Sketch, ii. 368 = reticulatus.
obovatus, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 95.—Sibir. altaic.
obtusatus, *Thunb. Prod. Pl. Cap.* 159.—Afr. austr.
occidentalis, *Torr. & Gray, Fl. N. Am.* ii. 164.—Am. bor. occ.
ochraceus, H. B. & K. Nov. Gen. et Sp. iv. 94 = Tetramolopium ochraceum.
Oculus-Christi, Cav. ex Steud. Nom. ed. II. i. 156 = Inula Oculus-Christi.
odontophyllus, Wall. Cat. n. 3285 = Senecio linifolius.
odorus, *All. Fl. Pedem.* i. 197.—Italia.
officinalis, All. l. c. 194 = Inula Helenium.

ASTER :—
oliganthemus, *F. Muell. Census*, 78.—Austral.
onustus, Nees, Gen. et Sp. Aster. 122 = Novi-Belgii.
oolentangiensis, Riddell, Syn. Fl. W. States, 55 = azureus.
oppositifolius, Linn. Amoen. Acad. v. 369 = Cymbalariae.
orarius, F. Muell. Fragm. v. 78 = Olearia glutinosa.
oreganus, *Torr. & Gray. Fl. N. Am.* ii. 163.—Am. bor. occ.
orientalis, S. G. Gmel. It. i. 155. t. 36. f. 1 = Inula britannica.
ornatus, Hort. Angl. ex Steud. Nom. ed. II. i. 156 = Herveyi.
Ottonis, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab.?
pallens, Willd. Enum. Hort. Berol. Suppl. 58 = patulus.
pallidulus, B. C. Vog. in Trew, Pl. Rar. t. 28 = cordifolius.
Palmeri, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 209.—Texas.
paludosus, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 201.—Am. bor.
palustris, Lam. Fl. Fr. ii. 143 = A. Tripolium.
panduratus, *Nees, ex Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 258.—China.
paniculatus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 207 = Lindleyanus.
paniculatus, *Lam. Encyc.* i. 306.—Am. bor.
paniculatus, *Mill. Gard. Dict.* ed. VIII. n. 24.—Hab.?
paniculatus, Muhl. in Ell. Sketch, ii. 365 = sagittifolius.
paniculatus, Nutt. Gen. Am. ii. 156 = undulatus.
paniculatus, Willd. Sp. Pl. iii. 2035 = cordifolius.
pannonicus, Jacq. Hort. Vindob. i. 3 = A. Tripolium.
pannosus, F. Muell. Fragm. v. 83 = Olearia pannosa.
Pappei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 79.—Afr. austr.
papposus, Willd. ex Spreng. Syst. iii. 525 = Printzia Bergii.
Parryi, *A. Gray, in Am. Natural.* viii. (1874) 212.—Am. bor. occ.
parviflorus, *A. Gray, in Bot. Calif.* i. 322.—Mexic.
parviflorus, Darlingt. Fl. Cestr. 446 = diffusus.
parviflorus, Nees, Syn. Aster. 29 = Tradescanti.
patens, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 201.—Am. bor.
patentissimus, Lindl. in DC. Prod. v. 232 = patens.
Pattersoni, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 272.—Colorado.
patulus, *Lam. Encyc.* i. 308.—Am. bor.
paucidentatus, F. Muell. Fragm. v. 66 = Olearia paucidentata.
pauciflorus, Lindl. ex DC. Prod. v. 231 = sibiricus.
pauciflorus, Martens, in Bull. Acad. Brux. viii. (1841) 67 = ericoides.
pauciflorus, *Nutt. Gen. Am.* ii. 154.—Am. bor.; Mexic.
pedunculatus, Wall. Cat. n. 2967 = asperulus.
pellitus, *H. B. & K. Nov. Gen. et Sp.* iv. 91.—N. Granat.
pendulus, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 205 = diffusus.
pensilvanicus, Poir. Encyc. Suppl. i. 498 = laevis.
peregrinus, Less. in Linnaea, vi. (1831) 123 = foliaceus.
peregrinus, *Pursh, Fl. Am. Sept.* ii. 556.—As. or.; Am. bor. arct.
perezioides, Less. Syn. Comp. 167 = Mairia perezioides.
persoonioides, A. Cunn. in DC. Prod. v. 267 = Olearia persoonioides.
pesthiensis, *DC. Prod.* v. 237.—Hungar.
petiolatus, *Harv. Thes. Cap.* ii. 35. t. 154.—Afr. austr.
philadelphicus, *Bart. Fl. Phil. Prod.* 81.—Am. bor.
phlogifolius, Muhl. in Willd. Sp. Pl. iii. 2034 = patens.
phlogopappus, Labill. Nov. Holl. Pl. ii. 49. t. 495 = Olearia stellulata.
phlogotrichus, Spreng. Syst. iii. 525 = Olearia stellulata.
phylicoides, H. B. & K. Nov. Gen. et Sp. iv. 98 = Tetramolopium phylicoides.
phyllolepis, *Torr. & Gray, Pl. N. Am.* ii. 113.—Am. bor.
pichinchensis, Willd. ex Nees, Gen. et Sp. Aster. 203 = Tetramolopium tenerrimum.
pilosus, Willd. Sp. Pl. iii. 2025 = ericoides.

ASTER :—
pimeleoides, A. Cunn. in DC. Prod. v. 268 = Olearia pimeleoides.
pinifolius, F. Muell. Fragm. v. 71 = Olearia pinifolia.
pinifolius, Nees, Syn. Aster. 29 = polyphyllus.
pinnatus, Cav. Ic. iii. 6. t. 212 = Dyssodia pubescens.
plantagineus, Poepp. ex DC. Prod. v. 418 = Senecio plantagineus.
plantaginifolius, Nutt. ex Nees, Gen. et Sp. Aster. 299 = Sericocarpus conyzoides.
pluriflorus, *G. Don, in Loud. Hort. Brit.* 346.—Afr. austr.
Pohlii, *Baker, in Mart. Fl. Bras.* vi. III. 23.—Bras.
polifolius, *Linn. Sp. Pl.* ed. II. 1224 = Printzia Bergii.
politus, Nees, Syn. Aster. 23 = laevis.
polyphyllus, Moench, Meth. Suppl. 253 = macrophyllus.
polyphyllus, Pursh, ex DC. Prod. v. 245 = tenuifolius.
polyphyllus, *Willd. Enum. Hort. Berol.* 888.—Am. bor.
Porteri, *A. Gray, in Proc. Am. Acad.* xvi. (1880) 99.—Am. bor. occ.
potosinus, *A. Gray, l. c.* xv. (1880) 32.—Mexic.
pradensis, *Phil. in Linnaea*, xxviii. (1856) 729.—Chili.
praealtus, Torr. & Gray, Fl. N. Am. ii. 134 = Novi-Belgii.
praealtus, Poir. Encyc. Suppl. i. 493 = salicifolius.
praecox, Lindl. in Hook. Fl. Bor. Am. ii. 9 = Lindleyanus.
praecox, Mill. Gard. Dict. ed. VIII. n. 19 = pyrenaeus.
praecox, Willd. Enum. Hort. Berol. Suppl. 58 = patulus.
pratensis, *Rafin. Fl. Ludov.* 67.—Am. bor.
Preissii, F. Muell. Fragm. v. 66 = Olearia heleophila.
prenanthoides, *Muhl. in Willd. Sp. Pl.* iii. 2046.—Am. bor.
Prescottii, Lindl. in DC. Prod. v. 231 = sibiricus.
procumbens, *Mill. Gard. Dict.* ed. VIII. n. 32.—Mexic.
propinquus, A. Cunn. ex DC. Prod. v. 270 = Olearia ramulosa.
pseudamellus, Wender. in Schrift. Marb. ii. (18 1) 226 = amelloides.
pseudamellus, *Hook. f. Fl. Brit. Ind.* iii. 249.—Ind. or.
pseudoamellus, DC. Fl. Fr. Suppl. v. 469 = A. Amellus.
pseudodiffusus, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab.?
pseudodumosus, *E. Forst. ex Forbes, l. c.*—Hab.?
ptarmicoides, *Torr. & Gray, Fl. N. Am.* ii. 160.—Am. bor.
pubescens, Hornem. Hort. Hafn. Suppl. 98 = heterophyllus.
pubescens, Lam. Encyc. i. 307 = tataricus.
pubescens, Moench, Meth. 600 = Inula squarrosa.
pubescens, Nees, Syn. Aster. 16 = sagittifolius.
pulchellus, *Eaton, in Bot. King, Exp.* 143. t. 16.—Am. bor. occ.
pulchellus, Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 255 = alpinus.
pulchellus, Willd. Sp. Pl. iii. 2019 = Erigeron pulchellus.
pulchellus, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. xxxviii. (1865) i. 376 = flaccidus.
pulcherrimus, Lodd. Bot. Cab. i. t. 6 = linariifolius.
Pulicaria, All. Fl. Pedem. i. 197 = Pulicaria vulgaris.
pulicarius, Scop. Fl. Carn. ed. II. ii. 172 = Pulicaria vulgaris.
pulverulentus, Hoffmgg. Verz. Pfl. Nachtr. i. (1824) 165 = Olearia stellulata.
pumilus, Banks & Soland. ex Hook. f. Fl. N. Zeal. i. 125 = Vittadinia australis.
pumilus, Fisch. Cat. Hort. Gorenk. (1808) 62; ed II. (1812) 39 = altaicus.
punctatus, Lapeyr. Hist. Abr. Pl. Pyr. 513 = Jasione tuberosa.
punctatus, Waldst. & Kit. Pl. Rar. Hung. ii. t. 109 = acris.
puniceus, Ell. Sketch, ii. 255 = Elliottii.
puniceus, *Linn. Sp. Pl.* 875.—Am. bor.
purpurascens, *Sch. Bip. in Seem. Bot. Voy. Herald*, 156.—Mexic.
purpuratus, Nees, Gen. et Sp. Aster. 118 = virgatus.
purpureo-caerulescens, *Hill, Hort. Kew.* 17.—Am. bor.
purpureus, Gueldenst. ex Ledeb. Fl. Ross. ii. 477 = A. Amellus?
purpureus, *Jacq. ex Steud. Nom.* ed. I. 83.—Hab.?

ASTER :—
pusillus, *Hornem. Hort. Hafn.* ii. 812.—Hab. ?
pygmaeus, *Lindl. in DC. Prod.* v. 228.—Am. arct.
pyrenaeus, *DC. Fl. Fr.* iv. 146.—Mont. Pyren.
pyrenaicus, Pourr. in Mém. Acad. Toul. iii. (1788) 308 = Erigeron uniflorus.
pyrrhopappus, *Boiss. Fl. Orient.* iii. 158.—Afghan.
quercifolius, Less. ex Walp. Rep. vi. 123 = Erigeron quercifolius.
quercifolius, Sieber, ex DC. Prod. v. 272 = Olearia stellulata.
quitensis, *Willd. ex Spreng. Syst.* iii. 533.—Ecuador.
racemosus, *Ell. Sketch*, ii. 348.—Am. bor.
Radula, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 210.—Am. bor.
Radula, Less. in Linnaea, vi. (1831) 125 = Chamissonis.
radulinus, *A. Gray, in Proc. Am. Acad.* viii. (1872) 388.—Am. bor. occ.
ramosissimus, Mill. Gard. Dict. ed. VIII. n. 21 = ericoides.
ramosus, *G. Don, in Loud. Hort. Brit.* 346.—Am. bor.
ramulosus, Labill. Nov. Holl. Pl. ii. 51 = Olearia ramulosa.
ramulosus, Lindl. in DC. Prod. v. 243 = falcatus.
ramulosus, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 293 = adscendens.
ranunculoides, *Moench, Meth. Suppl.* 254.—Hab. ?
recurvatus, Gucnth. ex Nees, Gen. et Sp. Aster. 97 = bellidiflorus.
recurvatus, Nees, l. c. 93 = tenuifolius.
recurvatus, Pursh, Fl. Am. Sept. ii. 556 = Tradescanti.
recurvatus, Spreng. in Schrad. Journ. Bot. (1800) 195 = squarrulosus.
recurvatus, Willd. Sp. Pl. iii. 2047 = diffusus.
reflexus, Linn. Sp. Pl. 877 = Felicia reflexa.
regalis, Salisb. Prod. 198 = Callistephus hortensis.
Regnellii, *Baker, in Mart. Fl. Bras.* vi. III. 21.—Bras.
repens, *H. B. & K. Nov. Gen. et Sp.* iv. 90. t. 332.—N. Granat.
reticulatus, *Pursh, Fl. Am. Sept.* ii. 548.—Am. bor.
retroflexus, A. Cunn. ex Hook. f. Fl. N. Zeal. i. 148 = Senecio glastifolius.
retroflexus, *Lindl. in DC. Prod.* v. 242.—Am. bor.
Reversii, Hort. ex Decne. in Rev. Hortic. Sér. IV. i. (1852) 181 = ericoides.
revolutus, *F. Muell. Census*, 78.—Austral.
rhomboideus, Lindl. ex DC. Prod. v. 230 = tataricus.
Richardsonii, Spreng. Syst. iii. 528 = sibiricus.
rigidulus, Desf. Tabl. ed. II. 122 = salicifolius.
rigidulus, Nees, Syn. Aster. 29 = leucanthemus, patulus.
rigidus, Linn. Sp. Pl. 874 (ex parte) = linariifolius.
rigidus, Moench, Meth. Suppl. 251 = Inula salicina.
riparius, *H. B. & K. Nov. Gen. et Sp.* iv. 92.—Arizona; Mexic.
riparius, Nees, Gen. et Sp. Aster. 280 = pesthiensis.
rivularis, *Cass. ex Loud. Hort. Brit.* 347.—Am. bor.
rivularis, Less. in Linnaea, v. (1830) 143 = Erigeron repens.
robustus, Hort. Bonn. ex Nees, Gen. et Sp. Aster. 282 = strictus.
robustus, Hort. ex Nees, l. c. 300 = salicifolius.
roseus, *Bieb. ex Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 245.—Reg. Caucas.
roseus, Desf. Cat. Hort. Par. ed. III. 401 = Novae-Angliae.
rosmarinifolius, A. Cunn. ex DC. Prod. v. 268 = Olearia rosmarinifolia.
rosmarinifolius, Spreng. Syst. iii. 526 = obtusatus.
rotundifolius, *Thunb. Prod. Pl. Cap.* 160.—Afr. austr.
Roxburghii, Hook. f. in Hook. Ic. Pl. xi. 45 = Commidendron robustum.
ruber, *Forbes, Hort. Woburn.* 189.—Hab. ?
rubicundus, *Forbes, l. c.*—Hab. ?
rubricaulis, Lam. Encyc. i. 305 = laevis.
rufibarbis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 76.—Afr. austr.
rugulosus, *Maxim. in Bull. Acad. Pétersb.* xv. (1871) 226, 220.—Japon.
rupestris, H. B. & K. Nov. Gen. et Sp. iv. 94 = Tetramolopium rupestre.
rutaeodorus, A. Cunn. ex DC. Prod. v. 269 = Olearia glandulosa.
sagittifolius, Ell. Sketch, ii. 262 = undulatus.

ASTER :—
sagittifolius, *Wedem. ex Willd. Sp. Pl.* iii. 2035.—Am. bor.
salicifolius, *Ait. Hort. Kew.* iii. 203; *Lam. Encyc.* i. 306.—Am. bor.
salicifolius, Nees, Syn. Astcr. 26 = longifolius.
salicifolius, Richards. App. Frankl. Journ. ed. I. 748 = junceus.
salicifolius, Scholl. Fl. Barb. Suppl. 328 = paniculatus.
salicinus, Nees, Gen. et Sp. Aster. 171, in syn. = salignus.
salicinus, Scop. Fl. Carn. ed. II. ii. 172 = Inula salicina.
salignus, Willd. Sp. Pl. iii. 2040 = paniculatus.
salinus, Schrad. Hort. Gotting. 9 = A. Tripolium.
salsuginosus, Hook. Fl. Bor. Am. ii. 6, partim = peregrinus.
salsuginosus, Less. in Linnaea, vi. (1831) 124 = sibiricus.
salsuginosus, Richards. ex R. Br. in App. Frankl. Journ. ed. I. 749 = Erigeron salsuginosus.
sanguineus, Willd. ex Spreng. Syst. iii. 533 = paniculatus.
satureiaefolius, Banks, in Reliq. Houston. t. 19 = Pectis satureioides.
Sayianus, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 294 = modestus.
scaber, Ell. Sketch, ii. 363 = undulatus.
scaber, *Moench, Meth. Suppl.* 253.—Hab. ?
scaber, *Thunb. Fl. Jap.* 316.—Japon.
scaberrimus, *Less. in Linnaea*, v. (1830) 143.—Mexic.
scabcrulus, *Miq. in Journ. Bot. Néerl.* i. (1861) 100.—China.
scabridus, *E. Mey. ex DC. Prod.* v. 225.—Afr. austr.
scabridus, Hook. f. & Thoms. ex C. B. Clarke, Comp. Ind. 47 = trinervius.
scabris, Mill. Gard. Dict. ed. VIII. n. 16 = alpinus.
scabrosus, Bertol. Misc. Bot. vi. 29 = Sericocarpus tortifolius.
scandens, Jacq. f. ex Spreng. Syst. iii. 524 = carolinianus.
scapigerus, *Ledeb. Fl. Ross.* ii. 472.—Sibiria.
scaposus, Klatt, in Sitzb. Akad. Muench. (1878) 84 = molliusculus.
scepusiensis, Kit. in Linnaea, xxxii. (1863) 373 = A. Amellus.
Schreberi, Nees, Syn. Aster. 16 = macrophyllus.
scoparius, DC. Prod. v. 242 = multiflorus.
scoparius, Nees, Syn. Aster. 28 = diffusus, Tradescauti.
scopulorum, *A. Gray, in Proc. Am. Acad.* xvi. (1880) 98.—Am. bor. occ.
scorpioides, Scop. Fl. Carn. ed. II. ii. 169 = Doronicum scorpioides.
scorzonerifolius, *Regel, Gartenfl.* (1873) 1; *et in Act. Hort. Petrop.* ii. (1873) 308.—Mexic.
secundiflorus, Desf. Tabl. ed. II. 268 = vimineus.
secundirameus, Zeyh. ex Sch. Bip. Tanacet. 20 = Matricaria nigellaefolia.
sedifolius, Linn. Sp. Pl. 874 = acris.
seliger, Nees, in Neuwied, Reise Nord. Am. ii. 439 = diffusus.
sericeus, *Vent. Jard. Cels*, t. 33.—Am. bor.
serotinus, Willd. Sp. Pl. iii. 2049 = Novi-Belgii.
serratifolius, Schloss. & Vukot. Syll. Fl. Croat. 57 = paniculatus.
serratus, Lag. ex DC. Prod. v. 315 = Grindelia inuloides.
serratus, *Thunb. Prod. Pl. Cap.* 160.—Afr. austr.
serrulatus, *Harv. in Harv. & Sond. Fl. Cap.* iii. 73.—Afr. austr.
sessilifolius, *Fisch. ex Steud. Nom.* ed. II. i. 157.—Am. bor.
setosus, *Baker, in Mart. Fl. Bras.* vi. III. 22.—Bras.
shastensis, *A. Gray, in Bot. Calif.* i. 322.—Calif.
Shortii, *Lindl. ex Hook. Fl. Bor. Am.* ii. 9; *Riddell, Syn. Pl. W. States*, 55.—Am. bor.
sibiricus, Lam. Encyc. i. 305 = pyrenaeus.
sibiricus, *Linn. Sp. Pl.* 872.—Sibir.; Am. bor. occ.
sibiricus, Turcz. ex Torr. & Gray, Fl. N. Am. ii. 107 = montanus.
Siemssenii, F. Muell. Fragm. v. 71 = Olearia viscidula.
sikkimmensis, *Hook. f. & Thoms. in Bot. Mag.* t. 4557.—Ind. or.
silenifolius, *Turcz. in DC. Prod.* v. 227.—Sibiria.
simplex, Eaton, in Bot. King, Exped. 140 = oreganus.

ASTER :—
simplex, Willd. Enum. Hort. Berol. 887 = paniculatus.
simulans, *Harv. in Harv. & Sond. Fl. Cap.* iii. 73.—Afr. austr.
sinuatus, A. Cunn. ex DC. Prod. v. 272 = Olearia quercifolia.
Solandri, *E. Forst. ex Forbes, Hort. Woburn.* 189.—Hab. ?
solidagineus, Michx. ex Willd. Sp. Pl. iii. 2024 = Sericocarpus solidagincus.
solidaginoides, Pers. Syn. ii. 443 = Sericocarpus solidagineus.
Sonderi, F. Muell. Fragm. v. 83 = Olearia grandiflora.
Sonorae, A. Gray, Pl. Wright. ii. 76 = riparius.
sparsiflorus, Michx. Fl. Bor. Am. ii. 112 = dumosus.
sparsiflorus, Pursh, Fl. Am. Sept. ii. 547 = tenuifolius.
spartioides, C. B. Clarke, Comp. Ind. 46 = altaicus.
spathularis, Brouss. ex DC. Prod. v. 315 = Grindelia inuloides.
spathulatus, Hort. ex Steud. Nom. ed. II. i. 157 = Inula salicina.
spathulatus, Lag. ex Nees, Gen. et Sp. Aster. 283 = Grindelia inuloides.
spathulatus, *Lindl. in DC. Prod.* v. 231.—Am. arct.
spathulifolius, *Maxim. in Bull. Acad. Pétersb.* xvi. (1871) 216.—Japon.
speciosus, Hornem. Hort. Hafn. ii. 816 = spectabilis.
spectabilis, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 209.—Am. bor.
spectabilis, Hook. & Arn. Bot. Beech. Voy. 146 = Chammissonis.
spectabilis, Willd. Sp. Pl. iii. 2048 = Novi-Belgii.
spinosus, *Benth. Pl. Hartw.* 20.—Am. bor.; Mexic.
spinulosus, *Chapm. Fl. S. U. St.* 199.—Florida.
spurius, Willd. Sp. Pl. iii. 2032 = Novae-Angliae.
squarrosulus, Nees, Gen. et Sp. Aster. 86 = longifolius.
squarrosus, All. Fl. Pedem. i. 196 = Inula spiraeifolia.
squarrosus, Banks, ex Steud. Nom. ed. II. i. 157 = linariifolius.
squarrosus, *Walt. Fl. Carol.* 209.—Carolina.
Steetzii, F. Muell. Fragm. v. 66 = Olearia strigosa.
stellulatus, Labill. Nov. Holl. Pl. ii. 50. t. 196 = Olearia stellulata.
stenomeres, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 209.—Am. bor. occ.
stenophyllus, Lindl. in DC. Prod. v. 242 = salicifolius.
Stracheyi, *Hook. f. Fl. Brit. Ind.* iii. 250.—Reg. Himal.
striatus, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 233.—China.
strictus, Poir. Encyc. Suppl. i. 498 = paniculatus.
strictus, Pursh, Fl. Am. Sept. ii. 556 = A. Radula.
strictus, Wender. in Schrift. Marb. ii. (1839) VI. 54 = dracunculoides.
strigosus, *Lichtenst. ex Harv. & Sond. Fl. Cap.* iii. 83.—Afr. austr.
strigosus, Thunb. Prod. Pl. Cap. 159 = Amellus annuus.
Stuartii, F. Muell. Fragm. v. 76 = Olearia Stuartii.
subasper, Lindl. ex Hook. Comp. Bot. Mag. i. (1835) 97 = salicifolius.
subcymosus, Bernh. ex Nees, Gen. et Sp. Aster. 138 = macrophyllus.
subspicatus, Nees, l. c. 74 = Douglasii.
subulatus, Hort. ex Steud. Nom. ed. II. i. 157 = ericoides.
subulatus, *Michx. Fl. Bor. Am.* ii. 111.—Am. bor.
succulentus, Gilib. Exercit. i. 174 = A. Tripolium.
surculosus, *Michx. Fl. Bor. Am.* ii. 113.—Am. bor.
tanacetifolius, *H. B. & K. Nov. Gen. et Sp.* iv. 95.—Calif.; Mexic.
tardiflorus, *Linn. Sp. Pl.* ed. II. 1231.—Am. bor.
tardiflorus, Willd. Sp. Pl. iii. 2049 = Novi-Belgii.
tataricus, *Linn. f. Suppl.* 373.—Tataria.
tataricus, Turcz. in Bull. Soc. Nat. Mosc. (1838) 94 = sibiricus.
taxifolius, Linn. Pl. Afr. 22; Amoen. Acad. vi. 102 = Mairia taxifolia.
tenellus, Linn. ll. cc. = Felicia fragilis.
tenellus, *Thunb. Fl. Cap.* 686.—Afr. austr.
tener, *Harv. in Harv. & Sond. Fl. Cap.* iii. 80.—Afr. austr.
tenerrimus, Less. in Linnaea, iv. (1829) 120 = Tetramolopium tenerrimum.

ASTER:—
tenue, *Kellogg, in Proc. Calif. Acad.* vii. (1877) 114.—Calif.
tenuiculus, *Bart. Fl. Phil. Prod.* 81.—Am. bor.
tenuifolius, Ell. Sketch, ii. 347 = vimineus.
tenuifolius, *Linn. Sp. Pl.* 873.—Am. bor.
tenuifolius, Nees, Syn. Aster. 119, partim = polyphyllus.
tenuifolius, Willd. Sp. Pl. iii. 2026 = ericoides.
tenuis, *Less. ex Ledeb. Fl. Ross.* ii. 483.—Rossia.
teretifolius, F. Muell. Fragm. v. 77 = Olcaria teretifolia.
Thomsoni, *C. B. Clarke, Comp. Ind.* 48.—Reg. Himal.
thyrsiflorus, Hoffm. Phyt. Blactt. i. (1803) 83 = Novi-Belgii.
tibeticus, *Hook. f. Fl. Brit. Ind.* iii. 251.—Tibet. occ.
Tilesii, Wichstr. in Nya Handl. Stockh. (1822) 13 = peregrinus.
tinctorius, Wallr. Beitr. Fl. Hercyn. 271; et in Linnaea, xiv. (1840) 641 = A. Amellus.
tomentellus, Hook. & Arn. Bot. Beech. Voy. 146 = Corethrogyne filaginifolia.
tomentosus, Schrad. in Wendl. Sert. Hann. 8. t. 24 = Olearia dentata.
tortifolius, *A. Gray, in Proc. Am. Acad.* vii. (1868) 353.—Am. bor. occ.
tortifolius, Michx. Fl. Bor. Am. ii. 109 = Sericocarpus tortifolius.
Townshendii, Hook. f. Bot. Mag. t. 6430 = Bigelovii.
Tradescanti, Hoffm. Phyt. Blaett. i. (1803) 68. t. A. f. 2 = patulus.
Tradescanti, *Linn. Sp. Pl.* 876.—Am. bor.
Tradescanti, Michx. Fl. Bor. Am. ii. 118 = diffusus.
Tradescanti, Nees, Syn. Aster. 28 = paniculatus.
trianthus, *Vell. Fl. Flum.* viii. t. 120.—Bras.
tricephalus, *C. B. Clarke, Comp. Ind.* 43.—Reg. Himal.
tricolor, *Spreng. in Flora*, xiii. (1830) 1. *Beil.* 15, nomen.—Afr. austr.
trifurcatus, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 317 = Senecio trifurcatus.
trilineatus, *Sch. Bip. ex Klatt, in Leopoldina*, xx. (1884) 91.—Mexic.
trinervifolius, Less. in Linnaea, ix. (1834) 183 = paniculatus.
trinervis, *Desf. Tabl.* ed. II. 122.—Gallia.
trinervis, Ruiz & Pav. ex DC. Prod. vii. 22 (= *Diazeuxis trinervis*).—Peruv.
trinervius, Gilib. Fl. Lithuan. iii. 214 = A. Amellus.
trinervius, *Roxb. Hort. Beng.* 61; *Fl. Ind.* iii. 433.—Reg. Himal.
Tripolium, *Linn. Sp. Pl.* 872.—Europ.; Afr. bor.; Oriens; As. bor.
Tripolium, Walt. Fl. Carol. 209 = tenuifolius.
tubuliflorus, F. Muell. Fragm. v. 65 = Olearia tubuliflora.
turbinatus, *S. Moore, in Journ. Bot.* xvi. (1878) 132.—China.
turbinellus, *Lindl. ex Hook. Comp. Bot. Mag.* i. (1835) 98.—Am. bor.
Turczaninowii, F. Muell. Fragm. v. 67 = Olearia imbricata.
turkestanicus, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 303.—Turkestan.
umbellatus, *Mill. Gard. Dict.* ed. VIII. n. 22.—Am. bor.
unalaschkensis, Less. in Linnaea, vi. (1831) 122 = peregrinus.
unalaschkensis, Less. ex Bong. Veg. Isl. Sitcha, 148 = salsuginosus.
unctuosus, Steud. Nom. ed. II. i. 157 = Novi-Belgii.
undulatus, Ell. Sketch, ii. 361 = patens.
undulatus, *Linn. Sp. Pl.* 875.—Am. bor.
undulatus, Moench, Meth. 600 = Inula britannica.
uniflorus, Michx. Fl. Bor. Am. ii. 110 = nemoralis.
uniflorus, Moench, Meth. 602 = Novi-Belgii.
urophyllus, Lindl. in DC. Prod. v. 233 = sagittifolius.
vagans, *E. Forst. ex Forbes, Hort. Woburn.* 190.—Hab. ?
Vaillantii, All. Fl. Pedem. i. 196 = Inula Vaillantii.
Vahlii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 49.—Chili.
vernicosus, F. Muell. Fragm. v. 67 = Olearia passerinoides.
vernus, Linn. Sp. Pl. 876 = Erigeron nudicaulis.
versicolor, Hort. ex Steud. Nom. ed. II i. 157 = mutabilis.

STER:—
versicolor, *Willd. Sp. Pl.* iii. 2045.—Am. bor.
villosus, *Sch. Bip. in F. Schultz, Arch. Fl.* 130.—Europ.
villosus, Michx. Fl. Bor. Am. ii. 113 = ericoides.
villosus, Thunb. Prod. Pl. Cap. 159 = Felicia angustifolia.
vimineus, *Lam. Encyc.* i. 306.—Am. bor.
vimineus, Nees, Gen. et Sp. Aster. 68 = tardiflorus.
vimineus, Willd. Sp. Pl. iii. 2046 = virgatus.
virgatus, *Ell. Sketch*, ii. 353.—Am. bor.
virgatus, Moench, Meth. Suppl. 250 = tenuifolius.
virgatus, Nees, Syn. Aster. 27 = salicifolius.
virgineus, Nees, l. c. 22 = longifolius.
viscosus, Labill. Nov. Holl. Pl. ii. 53 = Olearia viscosa.
Walpersianus, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 259.—China.
Warmingii, *Baker, in Mart. Fl. Bras.* vi. III. 23.—Bras.
Watsoni, *A. Gray, Syn. Fl. N. Am.* i. II. 201.—Am. bor. occ.
Willdenovii, *E. Forst. ex Forbes, Hort. Woburn.* 190.—Hab. ?
Willkommii, *Sch. Bip. ex Willk. in Flora*, xxxiv. (1851) 742.—Hispan.
Wolfii, Favrat, in Bull. Soc. Murith. vii-viii. (1879) 4 = alpinus.
Wrightii, *A. Gray, Pl. Wright.* ii. 75.—Mexic.; Texas.
xerophilus, F. Muell. Fragm. v. 76 = Olearia xerophila.
xylorrhiza, *Torr. & Gray, Fl. N. Am.* ii. 158.—Am. bor. occ.
Zeyheri, Less. Syn. Comp. 177 = Felicia Zeyheri.

ASTERACANTHA, Nees, in Wall. Pl. As. Rar. iii. 75 (1832) = **Hygrophila**, R. Br. (Acanth.).
auriculata, Nees, in DC. Prod. xi. 248 = H. spinosa.
longifolia, Nees, in Wall. Pl. As. Rar. iii. 90 = H. spinosa.
macracantha, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 146 = H. spinosa.

ASTERANDRA, Klotzsch, in Wiegm. Archiv, vii. (1841) 200 = **Phyllanthus**, Linn. (Euphorb.).
cornifolia, Klotzsch, l. c. = P. grandifolius.
Sellowiana, Klotzsch, l. c. = P. grandifolius.

ASTERANTHA, Reichb. Handb. 190 (1837) = ASTERACANTHA, Nees, = **Hygrophila**, R. Br. (Acanth.).

ASTERANTHEMUM, Kunth, Enum. Pl. v. 151 (1850) = **Smilacina**, Desf. (Liliac.).
dahuricum, Kunth, l. c. 153 = S. dahurica.
trifoliatum, Kunth, l. c. = S. trifolia.
vulgare, Kunth, l. c. 152 = S. stellata.

ASTERANTHERA, Hanst. in Linnaea, xxvi. (1853) 211. *GESNERACEAE*, Benth. & Hook. f. ii. 1011.
chiloënsis, *Hanst. l. c.*—Chiloe.
ovata, *Hanst. l. c.* xxxiv. (1865–66) 417.—Chili.

ASTERANTHOS, Desf. in Mém. Mus. Par. vi.(1820) 9. t. 3. *MYRTACEAE*, Benth. & Hook. f. i. 724.
ASTERANTHUS, Spreng. Syst. ii. 568 (1825).
brasiliensis, *Desf. l. c.*—Bras.

ASTERANTHUS, Endl. Gen. 293 (1837) = ASTRANTHUS, Lour. = **Homalium**, Jacq. (Samyd.).

ASTERIAS, Borckh. in Roem. Arch. i. 1. (1796) 25 = **Gentiana**, Linn.
hybrida, G. Don, Gen. Syst. iv. 184 = G. lutea.
lutea, Borckh. in Roem. Arch. i. 1. (1796) 25 = G. lutea.

ASTERIASTIGMA, Bedd. Forest. Man. Bot. 236 (1873). *BIXINEAE.*
macrocarpa, *Bedd. l. c.*—Ind. or.

ASTERIDEA, Lindl. Swan Riv. App. 24 (1839) = **Athrixia**, Ker-Gawl. (Compos.).
gracilis, A. Gray, in Hook. Kew Journ. iv. (1852) 274 = Athr. gracilis.
multiceps, A. Gray, l. c. = Athr. multiceps.
puberulenta, Lindl. Swan Riv. App. 24 = Athr. australis.
stricta, A. Gray, in Hook. Kew Journ. iv. (1852) 275 = Athr. stricta.

ASTERIDIUM, Engelm. ex Walp. Rep. ii. 958 (1843) = **Chaetopappa**, DC. (Compos.).

ASTERINGA, E. Mey. ex DC. Prod. vi. 138 (1837) = **Pentzia**, Thunb. (Compos.).

ASTERIOIDES, Ponted. ex Linn. Class. Pl. 530 (1747) = **Buphthalmus**, Linn. (Compos.).

ASTERIPHOLIS, Ponted. ex Linn. Class. Pl. 530 (1747) = **Aster**, Linn. (Compos.).

ASTERISCIUM, Cham. & Schlecht. in Linnaea, i. (1826) 254. t. 5. *UMBELLIFERAE*, Benth. & Hook. f. i. 877.
BUSTILLOSIA, Clos, in C. Gay, Fl. Chil. iii. 106. t. 32 (1847).
CASSIDOCARPUS, Presl, ex DC. Prod. iv. 82 (1830).
DIPTERYGIA, Presl, l. c. 80 (1830).
EREMOCHARIS, Phil. Fl. Atacam. 25. t. 2 B (1860).
GYMNOPHYTON, Clos, in C. Gay, Fl. Chil. iii. 102. t. 32 (1847).
TRITAENICUM, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) 1. 169.
aemocarpon, *Clos, in C. Gay, Fl. Chil.* iii. 101.—Chili.
anethoides, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) 1. 170.—Chili.
chilense, *Cham. & Schlecht. in Linnaea*, i. (1826) 254. t. 5.—Chili.
flexuosum, *Hemsl. Diag. Pl. Nov.* 16.—Mexic.
isatidicarpum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 349.—Chili.
oemocarpum, Walp. Ann. i. 978 = aemocarpon.
Poeppigii, DC. Prod. iv. 82 = chilense.
polycephalum, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 332.—Chili.
pozoides, *Clos, in C. Gay, Fl. Chil.* iii. 100.—Chili.
ramosissimum, *Phil. in Linnaea*, xxxiii. (1864–65) 90.—Chili.
robustum, *Baill. Hist. Pl.* vii. 145.—Chili.
verrucosum, Meyen, Reise, i. 315 = chilense.

ASTERISCUS, Moench, Meth. 592 (1794) = **Odontospermum**, Neck. (Compos.).
aquaticus, Less. Syn. Comp. 210 = O. aquaticum.
aureus, Lange, Pugill. ii. 118.—Hispan.
brachiatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 60 = O. maritimum.
Daltoni, Walp. Ann. ii. 844 = O. Daltoni.
flosculosus, Decne. in Ann. Sc. Nat. Sér. II. iv. (1835) 202 = Anvillea Garcini.
graveolens, Less. Syn. Comp. 210 = O. graveolens.
imbricatus, DC. Prod. v. 487 = O. imbricatum.
lanuginosus, Aucher, ex DC. l. c. vii. 287 = O. lanuginosum.
littoralis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 60 = O. maritimum.
maritimus, Less. Syn. Comp. 210 = O. maritimum.
mauritanicus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 61 = O. maritimum.
nanus, Nym. Consp. 391 = O. aquaticum.
odorus, DC. Prod. v. 486 = O. odorum.
pygmaeus, Coss. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 471.—Afr. bor.; Palaest.; Beluchist.
Schimperi, Boiss. Fl. Orient. iii. 180.—Arab.
sericeus, DC. Prod. v. 486 = O. sericeum.
sessilis, Moench, Meth. 592 = O. aquaticum.
Smithii, Walp. Ann. ii. 844 = O. Smithii.
Vogelii, Walp. l. c. 845 = O. Vogelii.

ASTERISCUS, Tourn. ex Sch. Bip. in Webb, Phytogr. Canar. ii. 229 (1835–60) = **Pallenis**, Cass. (Compos.).
spinosus, Sch. Bip. l. c. 230 = P. spinosa.

ASTERISCUS, Reichb. Consp. 144 (1828) = **Asteriscium**, Cham. (Umbell.).

ASTEROCARPUS, Eckl. & Zeyh. Enum. 122 (1835) = **Pterocelastrus**, Meissn. (Celastrin.).
arboreus, Eckl. & Zeyh. l. c. 123 = P. variabilis.
Burmanni, Eckl. & Zeyh. l. c. 122 = P. tricuspidatus.
litoralis, Eckl. & Zeyh. l. c. = P. litoralis.
nervosus, Eckl. & Zeyh. l. c. 123 = P. variabilis.
rostratus, Eckl. & Zeyh. l. c. = P. rostratus.
stenopterus, Eckl. & Zeyh. l. c. = P. stenopterus.
tetrapterus, Eckl. & Zeyh. l. c. = P. variabilis.
tricuspidatus, Eckl. & Zeyh. l. c. 124 = P. rostratus.
typicus, Eckl. & Zeyh. l. c. 122 = P. tricuspidatus.

ASTEROCARPUS, Reichb. Handb. 261 (1837); Gren. & Godr. Fl. Fr. i. 190 = **Astrocarpus**, Neck. (Resed.).

ASTEROCEPHALUS, [Vaill.] Adans. Fam. ii. 152 (1763) = **Scabiosa**, Linn. (Dipsac.).
acutiflorus, Reichb. Fl. Germ. Excurs. 194 = S. maritima.
africanus, Spreng. Syst. i. 380 = S. eretiea.
agrestis, Spreng. l. e. 383 = S. gramontia.
altissimus, Spreng. l. e. = S. africana.
ambiguus, Reichb. Fl. Germ. Excurs. 194 = S. maritima.
amoenus, Spreng. Syst. i. 381 = S. amoena.
arenarius, Vis. Pl. Aeg. 6. t. 1 = S. arenaria.
argenteus, Spreng. Syst. i. 382 = S. ueranica.
atropurpureus, Reichb. Ic. Fl. Germ. xii. t. 1363 = S. maritima.
atropurpureus, Spreng. l. c. 381 = S. atropurpurea.
bannaticus, Spreng. Syst. i. 383 = S. banatica.
Biebersteinii, Sweet, Hort. Brit. ed. I. 205 = S. mierantha.
bipinnatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 58 = S. oehroleuea.
brachiatus, Reichb. Fl. Germ. Excurs. 194 = S. brachiata.
brevicomus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 57 = S. rotata.
canescens, Lag. Gen. et Sp. Nov. 8, in obs.; Spreng. Syst. i. 380 = S. suaveolens.
caucasicus, Spreng. l. e. 381 = S. eaucasiea.
cochinchinensis, Spreng. l. c. 380 = S. eochinchinensis.
Columbaria, Wallr. Sched. Crit. 48 = S. Columbaria.
Columnae, Reichb. Fl. Germ. Excurs. 195 = S. holoserieea.
crenatus, Spreng. Syst. i. 382 = S. erenata.
creticus, Spreng. l. e. 380 = S. eretiea.
daucoides, Spreng. l. e. 383 = S. daueoides.
dentatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 57 = S. stellata.
dichotomus, Lag. Gen. et Sp. Nov. 8 = S. diehotoma.
diffusus, Spreng. Syst. iv. Cur. Post. 38 = S. diffusa.
divaricatus, Lag. ex Steud. Nom. ed. I. 84; ed. II. i. 158 = S. sieula.
eburneus, Spreng. Syst. i. 382 = S. ueranica.
elegans, Lag. Gen. et Sp. Nov. 8 = S. eaueasiea.
flavescens, Sehur, in Verh. Siebenb. Ver. Naturw. x. (1859) 61 = S. oehroleuea.
graminifolius, Spreng. Syst. i. 380 = S. graminifolia.
gramuntius, Spreng. l. e. i. 383 = S. gramontia.
grandiflorus, Sweet, Hort. Brit. ed. I. 205 = S. maritima.
Hladnikianus, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 171 = S. lueida.
holosericeus, Spreng. Syst. i. 381 = S. holoserieea.
holosericeus, Webb, Iter, 40 = S. sphaeiotica.
induratus, Spreng. Syst. i. 380 = S. africana.
integrifolius, Lag. ex Steud. Nom. ed. II. i. 158 = S. integrifolia.
intermedius, Lag. Gen. et Sp. Nov. 8 = S. lusitanica.
isetensis, Spreng. Syst. i. 383 = S. isetensis.
legionensis, Spreng. l. c. 379 = S. legionensis.
limonifolius, Spreng. l. e. 380 = S. limonifolia.
lobatus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 57 = S. rotata.
lucidus, Spreng. Syst. i. 380 = S. nitens.
lyratus, Spreng. l. c. 381 = S. sieula.
maritimus, Spreng. l. e. 382 = S. maritima.
Massoni, Sweet, Hort. Brit. ed. I. 205 = S. nitens.
micranthus, Spreng. Syst. i. 381 = S. mierantha.
mollissimus, Spreng. l. e. 383 = S. Columbaria.
monspeliensis, Zumag. Fl. Pedem. i. 182; Jord. & Fourr. Ic. Pl. Europ. i. 47 = S. monspeliensis.
montanus, Schur, Enum. Pl. Transs. 299 = S. oehroleuca.
ochroleucus, Wallr. Sehed. Crit. 50; Spreng. Syst. i. 383 = S. isetensis.
palaestinus, Spreng. l. e. 380 = S. palaestina.
paucisetus, Loud. Hort. Brit. Suppl. ii. 581 = S. gramontia.
pectinatus, Lag. Gen. & Sp. Nov. 8 = S. peetinata.
pilosus, Lag. l. c. = S. ucraniea.
polytomus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 58 = S. rotata.
prolifer, Spreng. Syst. i. 380 = S. prolifera.
pseudo-banaticus, Schur, Enum. Pl. Transs. 300 = S. lucida.
pumilus, Spreng. Syst. i. 382 = S. Columbaria.

ASTEROCEPHALUS:—
pyrenaicus, Sweet, Hort. Brit. ed. I. 205 = S. Columbaria.
rotatus, Spreng. Syst. i. 381 = S. rotata.
rutaefolius, Sweet, Hort. Brit. ed. I. 205 = S. rutaefolia.
saxatilis, Spreng. Syst. i. 379 = S. saxatilis.
Scopolii, Reichb. Fl. Germ. Excurs. 195 = S. Scopolii.
sericeus, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 59 = S. graminifolia.
setifer, Spreng. Syst. i. 381 = S. setifera.
siculus, Spreng. l. c. = S. sieula.
silenifolius, Spreng. l. c. 380 = S. brachiata.
simplex, Spreng. l. e. 383 = S. monspeliensis.
sphacioticus, Spreng. l. e. 382 = S. sphaeiotiea.
stellatus, Reichb. Fl. Germ. Excurs. 194; et Ic. Fl. Germ. xii. f. 1370 = S. stellata.
stellatus, Spreng. Syst. i. 381 = S. monspeliensis.
strictus, Spreng. l. e. 380 = S. lueida.
suaveolens, Lag. Gen. & Sp. Nov. 8, in obs.; Wallr. Sehed. Crit. 51 = S. suaveolens.
Succisa, Wallr. Sched. Crit. 52 = S. Succisa.
tomentosus, Spreng. Syst. i. 381 = S. Succisa.
triniaefolius, Vis. in Mem. Ist. Venet. xxi. (1880–82) 482 = S. triniaefolia.
ucranicus, Spreng. Syst. i. 382 = S. ueranica.
urceolatus, Spreng. l. e. = S. rutaefolia.
vestinus, Faeeh. ex Reichb. Ic. Fl. Germ. xii. 20. n. 1373 = S. vestina.
virescens, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 59 = S. graminifolia.
Webbianus, Spreng. Syst. i. 381 = S. oehroleuca.
Wulfenii, Reichb. Fl. Germ. Excurs. 194, in syn. n. 1156 = S. ueranica.

ASTEROCHAETE, Nees, in Linnaea, ix. (1834) 300. *CYPERACEAE*, Benth. & Hook. f. Gen. iii. 1062.
angustifolia, *Nees, l. c.*—Afr. austr.
arundinacea, *Kunth, Enum. Pl.* ii. 312.—N. Caled.
capitellata, *Nees, in Linnaea*, ix. (1834) 300; x. (1836) 194.—Afr. austr.
elongata, *Kunth, Enum. Pl.* ii. 312.—Ins. Maurit.
glomerata, *Nees, in Linnaea*, ix. (1834) 300; x. (1836) 194—Afr. austr.
Ludwigii, *Hochst. in Flora*, xxviii. (1845) 759, in nota. —Afr. austr.
nitens, *Kunth, Enum. Pl.* ii. 313.—Ins. Maurit.
tenuis, *Kunth, l. c.* 312.—Afr. austr.

ASTEROCHITON, Turez. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 138 = **Thomasia**, J. Gay (Stereul.).
pygmaeus, Turez. l. c. 139 = T. pygmaea.

ASTEROCHLAENA, Gareke, in Bot. Zeit. viii. (1850) 666 = **Pavonia**, Cav. (Malvac.).
cuspidata, Gareke, l. e. 668.—Peruv.

ASTEROCYTISUS, Sehur, ex Fuss, Fl. Transs. 154 (1866) = **Genista**, Linn. (Legumin.).
radiatus, Sehur, ex Fuss, l. c. = G. radiata.

ASTEROGEUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 294 (1821) = **Plantago**, Tourn.
laciniatum, S. F. Gray, l. c. = P. Coronopus.

ASTEROGYNE, H. Wendl. ex Benth. & Hook. f. Gen. iii. 914 (1883). *PALMAE*, Benth. & Hook. f. iii. 914.

ASTEROLASIA, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 9; et Hook. Kew Journ. viii. (1856) 34. *RUTACEAE*, Benth. & Hook. f. i. 294, 990.
PLEURANDROPSIS, Baill. Adansonia, x. (1872) 305.
UROCARPUS, J. Drumm. in Hook. Kew Journ. vii. (1855) 54.
buxifolia, *Benth. Fl. Austral.* i. 351.—Austral.
chorilaenoides, F. Muell. in Trans. Vict. Inst. i. (1855) 116 = Mierocybe pauciflora.
correifolia, *Benth. Fl. Austral.* i. 350.—Austral.
grandiflora, *Benth. l. c.* 352.—Austral.
mollis, *Benth. l. c.* 351.—Austral.
Muelleri, *Benth. l. c.* 350.—Austral.
pallida, *Benth. l. c.* 352.—Austral.
phebalioides, *Benth. l. c.*—Austral.
phebalioides, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 10 = pleurandroides.
pleurandroides, *F. Muell. ex Benth. Fl. Austral.* i. 351. —Austral.
squamuligera, *Benth. l. c.* 352.—Austral.

ASTEROLASIA:—
trymalioides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 10.—Austral.

ASTEROLINION, Brongn. Enum. Gen. 69 (1843) = seq.

ASTEROLINON, Hoffmgg. & Link, Fl. Port. i. 332 (1820). *PRIMULACEAE*, Benth. & Hook. f. ii. 636.
PELLETIERA, A. St. Hil. in Mém. Mus. Par. ix. (1822) 365.
adoënse, *Kunze, in Linnaea*, xx. (1847) 37.—Abyss.
Linum-stellatum, Duby, in DC. Prod. viii. 68 = stellatum.
lysimachioideum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 64 = A. stellatum.
serpyllifolium, *Ball, in Journ. Linn. Soc.* xxi. (1884) 226.—Patagon.
stellatum, *Hoffmgg. & Link, Fl. Port.* i. 333.—Lusitan.
trinum, *Baudo, in Ann. Sc. Nat.* Sér. II. xx. (1843) 350. —Chili.

ASTEROMAEA, DC. Prod. v. 302 (1836) = seq.

ASTEROMOEA, Blume, Bijdr. 901 (1826) = **Boltonia**, L'Hérit. (Compos.).
indica, Blume, l. c. = B. indica.
pekinensis, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 225 = B. pekinensis.

ASTEROMYRTUS, Schau. in Linnaea, xvii. (1843) 242 = **Melaleuca**, Linn. (Myrtac.).
Gaertneri, Schau. l. c. 243 = M. angustifolia.

ASTEROPEA, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 79 = seq.

ASTEROPEIA, Thou. Hist. Vég. Afr. 51. t. 15. (1807). *SAMYDACEAE*, Benth. & Hook. f. i. 801.
amblyocarpa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 81.—Madag.
densiflora, *Baker, in Journ. Bot.* xx. (1882) 49.—Madag.
multiflora, *Thou. Hist. Vég. Afr.* 51. t. 15.—Madag.

ASTEROPHYLLUM, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 1077 (1829) = **Asperula**, Linn. (Rubiac.), &c.
Aparine, Schimp. & Spenn. l. c. = Galium Aparine.
Asperula, Schimp. & Spenn. l. c. = Asperula odorata.
galioides, Schimp. & Spenn. l. c. in obs. = Asperula galioides.
Galium, Schimp. & Spenn. l. c. = Galium verum.
Scherardianum, Schimp. & Spenn. l. c. = Sherardia arvensis.
sylvaticum, Schimp. & Spenn. l. c. in obs. = Asperula odorata.
tinctorium, Schimp. & Spenn. l. c. = Asperula tinctoria.

ASTEROPSIS, Less. Syn. Comp. 188 (1832) = **Podocoma**, Cass. (Compos.).
macrocephala, Less. l. c.—Bras.

ASTEROPTERUS, Adans. Fam. ii. 124 (1763); Gaertn. Fruct. ii. 460. t. 173 (1791) = **Leysera**, Linn. (Compos.).
Callicornia, Gaertn. Fruct. ii. 460 = L. Callicornia.
ceruleus, Rafin. Fl. Tellur. ii. 48 = an praec.?

ASTEROPUS, Schult. Mant. iii. 15 (1827) = ASTROPUS, Spreng. = **Waltheria**, Linn. (Stereul.).

ASTEROSCHOENUS, Nees, in Mart. Fl. Bras. ii. 1. 124 (1842) = **Rynchospora**, Vahl (Cyperac.).
consanguineus, Nees, l. c. 125 = R. consanguinea.
rigidus, Nees, l. c. 126 = R. rigida.
speciosus, Nees, l. c. 125 = R. speciosa.

ASTEROSPERMA, Less. Syn. Comp. 389 (1832) = **Felicia**, Cass. (Compos.).
chrysocomoides, Less. l. c.—Afr. austr.

ASTEROSTEMMA, Decne. in Ann. Sc. Nat. Ser. II. ix. (1838) 271. t. 10. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 775.
repandum, *Decne. l. c.*—Java.

ASTEROSTIGMA, Fisch. & Mey. in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 148; Schott, in Oestr. Bot. Wochenbl. (1852) 67 = **Staurostigma**, Scheidw. (Aroid.).
colubrinum, Schott, in Bonplandia, x. (1862) 86 = S. concinnum.
concinnum, Schott, in Oestr. Bot. Wochenbl. (1852) 67 = S. concinnum.
Langsdorffii, C. Koch, in Ind. Sem. Hort. Berol. (1854) 8 = S. concinnum.
lineolatum, Schott, in Bonplandia, x. (1862) 86 = S. concinnum.
Luschnathianum, Schott, Syn. Aroid. 126 = S. Luschnathianum.
Pavonii, Schott, Prod. Aroid. 339 = S. Pavoni.
Tweedianum, Schott, in Oestr. Bot. Zeitschr. (1859) 39 = S. Tweedianum.
Vellozianum, Schott, Syn. Aroid. 126 = S. Luschnathianum.
vermitoxicum, Griseb. in Goett. Abh. xix. (1874) 247 = S. vermitoxicum.
zebrina, Linden, in Belg. Hortic. xv. (1865) 98.—Bras.

ASTEROSTOMA, Blume, Mus. Bot. Lugd. Bat. i. 59 (1849) = **Osbeckia**, Linn. (Melast.).
aspera, Blume, l. c. = O. aspera.
decandra, Blume, l. c. = O. decandra.
decumbens, Blume, l. c. = O. decumbens.
Kleinii, Blume, l. c. = O. aspera.
multiflora, Blume, l. c. = O. multiflora.
octandra, Blume, l. c. = O. octandra.
repens, Blume, l. c. = Melastoma repens.
rotundifolia, Blume, l. c. = O. rotundifolia.
virgata, Blume, l. c. = O. virgata.
Walkeri, Blume, l. c. = O. Walkeri.
Wightiana, Blume, l. c. = O. octandra.

ASTEROTHRIX, Cass. in Dict. Sc. Nat. xlviii. 434 (1827) = **Leontodon**, Linn. (Compos.).
asperrima, Cass. l. c. = L. asperrimus.
crispa, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 100 = L. crispus.
hispanica, DC. Prod. vii. 127 = L. hispanicus.

ASTEROTRICHION, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. 19. t. 8 (1841) = **Plagianthus**, Forst. (Malvac.).
sidoides, Klotzsch, l. c. = P. sidoides.

ASTHENOCHLOA, Buese, in Miq. Pl. Jungh. 367 (1854). *GRAMINEAE*, Benth. & Hook. f. Gen. iii. 1096.
tenera, *Buese, l. c.* 368.—Java.

ASTHOTHECA, Miers, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 254 = **Clusia**, Linn. (Guttif.).

ASTIANTHUS, D. Don, in Edinb. Phil. Journ. ix. (1843) 262 = **Tecoma**, Juss. (Bignon.).
longifolius, D. Don, l. c. = T. viminalis.

ASTILBE, Buch.-Ham. in D. Don, Prod. Fl. Nep. 210 (1825). *SAXIFRAGEAE*, Benth. & Hook. f. i. 634.
HOTEIA, Morr. & Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 316. t. 11.
Aruncus, Trev. in Bot. Zeit. xiii. (1855) 819 = Spiraea Aruncus.
barbata, Hort.—Cf. Gard. Chron. (1871) 547 = japonica.
chinensis, *Franch. & Sav. Enum. Pl. Jap.* i. 144.—China; Japon.
decandra, *D. Don, Prod. Fl. Nep.* 211.—Am. bor.
japonica, *A. Gray, in Hook. Lond. Journ. Bot.* ii. (1843) 124.—Japon.
odontophylla, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 96 = chinensis.
podophylla, Baill. Hist. Pl. iii. 332 = Rodgersia podophylla.
rivularis, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 210.—Reg. Himal.
rubra, *Hook. f. & Thoms. ex Hook. Bot. Mag.* t. 4959.—Reg. Himal.
speciosa, Jungh. Java, i. (1853) 250 = rivularis.
Stoliczkai, *Kurz, ex Seem. in Journ. Bot.* v. (1867) 240 (Spiraeae sp. ?).—Himal. bor. occ.
Thunbergii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 96.—Japon.

ASTIRIA, Lindl. in Bot. Reg. (1844) Misc. 27. t. 49. *STERCULIACEAE*, Benth. & Hook. f. i. 221.
rosea, *Lindl. l. c.*—Maurit.

ASTOMA, DC. Coll. Mém. v. 71. t. 17 (1829); et Prod. iv. 249 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 885.
ASTOMAEA, Reichb. Handb. 218 (1837).
seselifolium, *DC. ll. cc.*—Syria.

ASTORGANTHUS, Endl. Cat. Hort. Vindob. ii. 196 (1843) = **Melicope**, Forst. (Rutac.).
Huegelii, Endl. l. c. = M. simplex.

ASTRADELPHUS, Rémy, in Ann. Sc. Nat. Sér. III. xii. (1849) 185 = **Erigeron**, Linn. (Compos.).
calbucanus, Phil. in Anal. Univ. Chil. (1873) 485.—Chili.
chilensis, Rémy, in Ann. Sc. Nat. Sér. III. xii. (1849) 185 = E. chilensis.

ASTRAEA, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = **Croton**, Linn. (Euphorb.).
divaricatum, Klotzsch, l. c. = C. Klotzschii.
diversifolia, Klotzsch, l. c. = C. Klotzschii.
glandulifera, Klotzsch, ex Wawra, Bot. Ergeb. Maxim. Bras. 31. t. 42 = C. Klotzschii.
Jatropha, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = C. carnosus.
Klotzschii, Didr. in Kjoeb. Vidensk. Meddel. (1857) 137 = C. Klotzschii.
lobatum, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = C. lobatus.
Manihot, Klotzsch, l. c. = C. lobatus.
palmata, Klotzsch, l. c. = C. lobatus.
Paulina, Didr. in Kjoeb. Vidensk. Meddel. (1857) 138 = C. Paulinus.
prunifolia, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = C. Klotzschii.
Seemanni, Klotzsch, in Seem. Bot. Voy. Herald, 103 = C. lobatus.
tomentosum, Klotzsch, in Wiegm. Archiv, vii. (1841) 194 = C. comosus.

ASTRAEA, Schau. in Linnaea, xvii. (1843) 238 = **Thryptomene**, Endl. (Myrtac.).
saxicola, Schau. l. c. 239 = T. saxicola.

ASTRAGALOIDES, Adans. Fam. ii. 323 (1763) = **Astragalus**, Tourn. (Legum.).
Alopecurus, Moench, Meth. 168 = Astragalus alopecuroides.
alpina, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 373 = Astragalus alpinus.
australis, Medic. l. c. = Astragalus australis.
bicolor, Moench, Meth. Suppl. 57 = Astragalus uliginosus.
Cicera, Moench, Meth. 168 = Astragalus Cicer.
frigida, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 373 = Astragalus frigidus.
Glaux, Moench, Meth. 168 = Astragalus Glaux.
syphilitica, Moench, l. c. = Astragalus exscapus.

ASTRAGALUS, Tourn. ex Linn. Gen. ed. I. 215 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 506.
AILUROSCHIA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 151.
ALOPECIAS, Stev. l. c. 143 (1856).
AMMODYTES, Stev. l. c. iv. (1832) 263.
ANKYLOBUS, Stev. l. c. xxix. (1856) II. 148.
ARAGALUS, Neck. Elem. iii. 12 (1790).
ARAGUS, Steud. Nom. ed. II. i. 118 (1840).
AULOSEMA, Walp. Rep. i. 694 (1842).
CARYOLOBIUM, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 266.
CHONDROCARPUS, Stev. l. c. 268 (1832).
CRACCINA, Stev. l. c. xxix. (1856) II. 144.
CYMBICARPUS, Stev. l. c. iv. (1832) 266.
CYSTIUM, Stev. l. c. xxix. (1856) II. 147.
DIDYMOPELTA, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 669.
DIPELTA, Regel & Schmalh. l. c. 578 (1877).
DIPLOTHECA, Hochst. in Flora, xxix. (1846) 595.
EROPHACA, Boiss. Voy. Bot. Espagne, 176 (1839-45).
EUILUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 149.
EUPREPIA, Stev. l. c. iv. (1832) 268.
FEIDANTHUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 148.
GLANDULA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 374.
GLOTTIS, Medic. l. c. 377 (1787).
GLYCYPHYLLA, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 266.
HAMARIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 364.
HAMOSA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 376.
HEDYPHYLLA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 142.
HOMALOBUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 350 (1840).
HYPOGLOTTIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 364.
KENTROPHYTA, Nutt. ex Torr. & Gray, Fl. N. Am. i. 353 (1840).
KIRCHNERA, Opiz, in Lotos, viii. (1858) 65.
MACROSEMA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 145.
MEDYPHYLLA, Opiz, in Lotos, viii. (1858) 65.
MYOBROMA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 150.
ONIX, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 378.
PEDINA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 144.
PHACA, Roy. ex Linn. Coroll. 13 (1737).
PHILAMMOS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 146.
PHYSONDRA, Rafin. Atl. Journ. 145 (1832).
PICRAENA, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 263.
PODOCHREA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 364.
PODOLOTUS, Royle, Illustr. Bot. Himal. 198 (1835).
PROSELIAS, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 268.
PSYCHRIDIUM, Stev. l. c. 263 (1832).
RYSODIUM, Stev. l. c. (1832).
SEWERZOWIA, Regel & Schmalh. in Act. Hort. Petrop. v. (1877) 580.
SOLENOTUS, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 143.
SPIESIA, Neck. Elem. iii. 13 (1790).
STELLA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 377.
THIUM, Steud. Nom. ed. I. 833 (1821).
TIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 373.
TRIQUETRA, Medic. l. c. 376 (1787).
XEROPHYSA, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 150.
abakanensis, Less. ex Bunge, Astrag. ii. 22 = multicaulis.
abbreviatus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) II. 343.—As. temp.
aboriginorum, *Richards. in Frankl. 1st Journ. App.* 746.—Am. bor. arct.
abyssinicus, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 193.—Abyss.
acanthostachys, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 326 = Roussaeanus.
acaulis, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 132.—Reg. Himal.
acicularis, *Bunge, Astrag.* i. 90; ii. 161.—Cappadoc.
acinaciferus, *Boiss. Diagn.* Ser. I. ix. 72.—Arab.
acmonotrichus, *Fenzl, Pugill. Pl. Nov. Syr.* 5.—Cilicia.
acmophyllus, *Bunge, Astrag.* i. 93; ii. 166.—Cappadoc.
acmotrichus, Bunge, l. c. i. 109; ii. 189 = acmonotrichus.
acutiflorus, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 120 = Falconeri.
acutifolius, *Bunge, Astrag.* i. 130; ii. 227.—Persia.
acutifolius, DC. ex Steud. Nom. ed. I. 88 = A. Stella.
acutifolius, *Schleich. Cat. Pl. Helv.* ed. IV. 9.—Europ.
acutirostris, *S. Wats. in Proc. Am. Acad.* xx. (1885) 360.—Calif.
acutus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
adesmiaefolius, *Benth. ex Bunge, l. c.* i. 4; ii. 2.—Tibet. occ.
adpressus, Ehrenb. ex Walp. Ann. iv. 502 = gummifer.
adscendens, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 317.—Persia.

ASTRAGALUS:—
adsurgens, *Pall. Astrag.* 40. n. 44. t. 31.—Sibir.; Am. bor.
adsurgens, Torr. in Stansbury, Rep. 385 = iodanthus.
adsurgens, Willd. ex Steud. Nom. ed. II. i. 159 = virgatus.
adunciformis, *Boiss. Diagn.* Ser. I. ii. 42.—As. Min.
aduncus, Bunge, Astrag. i. 102; ii. 179 = usturtensis.
aduncus, Bieb. Fl. Taur. Cauc. ii. 195 = Bungeanus.
aduncus, Hohen. Talysch. 107 = Cephalotes.
aduncus, *Willd. Sp. Pl.* iii. 1269.—Reg. Caucas.
adustus, *Bunge, Astrag.* i. 80; ii. 134.—Kurdistan.
Aegiceras, *Willd. Enum. Hort. Berol. Suppl.* 52 = Ins. Chio.
aegobromus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 74.—Persia.
aegyptiacus, Mill. Gard. Dict. ed. VIII. n. 11 = annularis.
aegyptiacus, Spreng. Syst. iii. 298 = kahiricus.
aeluropus, *Bunge, Astrag.* i. 89; ii. 157.—Reg. Caucas.; Persia.
aesicarius, *Pall. Reise*, iii. 751.—Sibir.
affghanus, *Boiss. Fl. Orient.* ii. 1095 (emend.).—Beluchist.
affinis, *Steud. Nom.* ed. II. i. 159.—Chili.
africanus, *Bunge, Astrag.* i. 61; ii. 104.—Algeria.
Agraniotii, *Orph. ex Boiss. Diagn.* Ser. II. ii. 29.—Laconia.
agrestis, Dougl. ex Hook. Fl. Bor. Am. i. 148 = hypoglottis.
aintabicus, *Boiss. Diagn.* Ser. I. ix. 86.—Syria; Mesopotam.
Aitchisoni, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 121.—Ind. bor. occ.
Ajfreidii, *Aitch. & Baker, in Journ. Linn. Soc.* xix. (1882) 157.—Afghanist.
ajubensis, *Bunge, Astrag.* i. 61; ii. 102.—Persia.
akkensis, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 57, nomen.—Afr. bor.
aksuensis, *Bunge, Astrag.* i. 25; ii. 30.—Sibir. altaic.
alaschanus, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 31.—Mongolia.
alatavicus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 344.—Alatau.
albens, *Greene, in Bull. Calif. Acad.* i. ('1885') 156.—Calif.
Alberjilla, *Steud. Nom.* ed. II. i. 159.—Chili.
Alberti, *Bunge, in Act. Hort. Petrop.* vii. (1880) 375.—Turkestan.
albicans, *Bong. Verz. Pfl. Saisang-Nor*, 21. t. 2.—Soongaria.
albicaulis, *DC. Astrag.* 132. n. 59. t. 21.—Reg. Caucas.
albicaulis, Ledeb. Fl. Ross. i. 628 = medius.
albidus, Waldst. & Kit. Pl. Rar. Hung. i. 39. t. 40 = vesicarius.
aleppicus, *Boiss. Diagn.* Ser. I. ii. 58.—Syria; Mesopotam.
alexandrinus, *Boiss. l. c.* ix. 75.—Aegypt.; Arab.; Syria.
algarbiensis, *Coss. ex Bunge, Astrag.* i. 9; ii. 6.—Hispan.
alienus, *A. Gray, Bot. U. St. Expl. Exped.* i. 418.—Pernv.
alindanus, *Boiss. Diagn.* Ser. I. ii. 85.—Caria.
allochorus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 366.—Am. bor.
Alopecias, C. Koch, ex Bunge, Astrag. ii. 97 = maximus.
Alopecias, *Pall. Astrag.* 12. t. 9.—Persia; Turkestan.
alopecuroides, Ledeb. Fl. Ross. i. 633. n. 88 = maximus (ex parte).
alopecuroides, *Linn. Sp. Pl.* 755.—Sibiria; Helv.; Kamtschatka.
alopecuroides, Pall. Astrag. 9. t. 7 = vulpinus.
alopecuroideus, Lam. Fl. Fr. ii. 636 = narbonnensis.
alopecuros, *Pall. ex DC. Astrag.* 11. t. 8.—As. med.
Alpamarcae, *A. Gray, Bot. U. St. Expl. Exped.* i. 417.—Peruv.
alpinus, Ledeb. Fl. Ross. i. 601 = arcticus.
alpinus, *Linn. Sp. Pl.* i. 760.—Reg. bor. et arct.
alpinus, Vill. Hist. Pl. Dauph. i. 303 = A. Onobrychis.
altaicus, *Bunge, Astrag.* i. 37; ii. 46.—Mont. Altai.
altaicus, Pall. Astrag. 56 = Oxytropis brevirostra.
alyssoides, Kotschy, ex Walp. Ann. ii. 374 = elbrusensis.
alyssoides, *Lam. Encyc.* i. 317.—Armenia.

ASTRAGALUS:—
amalecitanus, *Boiss. Diagn.* Ser. I. ix. 46.—Palaestin.
amanus, *Boiss. Fl. Orient.* ii. 290.—Syria.
amarus, *Pall. Astrag.* 8. t. 6.—Reg. Casp.
amatus, *Clos, in C. Gay, Fl. Chil.* ii. 115.—Chili.
ambiguus, *Bunge, Astrag.* i. 86; ii. 150.—Syria.
ambiguus, Pall. Astrag. 54 = Oxytropis ambigua.
amblolepis, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 372.—Kurdistan.
Amherstianus, *Benth. in Royle, Illustr. Bot. Himal.* 199.—Reg. Himal.
ammocryptus, Boiss. Diagn. Ser. I. ix. 59. = tribuloides.
Ammodendron, *Bunge, Astrag.* i. 130; ii. 228.—As. med.
Ammodytes, *Pall. Voy. Ed. Gall. App.* n. 372. t. 83.—Sibir. or.
ammophilus, Bieb. ex Bess. in Flora, xv. (1832) II. Beibl. 33 = virgatus.
ammophilus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 335.—Persia; Soongaria.
ammotrophus, *Bunge, Astrag.* i. 112; ii. 192.—Turkestan.
amoenus, *Fenzl, Pugill. Pl. Nov. Syr.* 4.—Cilicia.
amphilogus, *Bunge, Astrag.* i. 90; ii. 161.—Anatolia.
amphioxys, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 366.—Am. bor.
ampullarius, *S. Wats. in Am. Natural.* vii. (1873) 300.—Calif.
ampullatus, Pall. Reise, i. 750 = Oxytropis ampullata.
ampullifer, Griseb. Spicil. Fl. Rumel. i. 58 = physocalyx.
amygdaliger, Less. in Linnaea, ix. (1834) 176 = testiculatus.
amygdalinus, *Bunge, Astrag. Fedsch.* 228.—As. temp.
anacamptus, *Bunge, Astrag.* i. 110; ii. 190.—Persia.
anacardius, *Bunge, l. c.* i. 61; ii. 105.—Persia.
anatolicus, *Boiss. Diagn.* Ser. I. ii. 77.—Anatolia.
ancistrocarpus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 463.—Mesopotam.
ancistron, *Pomel, Nouv. Mat. Fl. Atl.* 186.—Algeria.
ancocarpus, *Pomel, l. c.* 187.—Algeria.
ancyleus, *Boiss. Fl. Orient.* ii. 240.—Cappadoc.
andalanicus, *Boiss. & Haussk. ex Boiss. l. c.* 345.—Persia.
Andersonii, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 524.—Calif.
Andrachne, *Bunge, Astrag.* i. 89; ii. 158.—Syria; Mesopotam.
andrachnaefolius, *Fenzl, Pugill. Pl. Nov. Syr.* 4.—As. Min.; Syria.
anemophilus, *Greene, in Bull. Calif. Acad.* i. (1885) 186.—Calif.
anfractuosus, Bunge, Astrag. i. 125; ii. 218 = subulatus.
angarensis, *Turcz. ex Bunge, l. c.* i. 126; ii. 222.—Sibir. baical.
angorensis, Bunge, l. c. ii. 176 = leucocyancus.
angulosus, *DC. Astrag.* 234. t. 45.—Syria.
angustatus, Boiss. Diagn. Ser. I. ix. 47 = ruscifolius.
angustatus, *Bunge, Astrag.* i. 125; ii. 220.—Persia.
angustiflorus, *C. Koch, in Linnaea*, xv. (1840) 720.—Armenia; Persia.
angustifolius, *Lam. Encyc.* i. 321.—Graecia; As. Min.
angustissimus, Bunge, Astrag. i. 135; ii. 232 = oxypetalus.
anisacanthus, *Boiss. Diagn.* Ser. I. ix. 103.—Afghan.
anisacanthus, Bunge, Astrag. ii. 128 = jubatus.
anisomerus, *Bunge, Astrag. Fedsch.* 228.—As. temp.
ankylotus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 27.—Reg. Casp.
ankylotus, Stschegl. in Bull. Soc. Nat. Mosc. (1854) I. 162 = commixtus.
annularis, *Forsk. Fl. Aegypt. Arab.* 139.—Aegypt.; Syria.
annuus, *DC. Astrag.* 127. n. 25.—Hab.?
anomalus, *Bunge, Astrag.* i. 35; ii. 62.—Tibet. occ.
anserinaefolius, *Boiss. Diagn.* Ser. I. ii. 76.—Persia.
antabicus, *Boiss. l. c.* ix. 86.—Anatolia.
anthylloides, *Lam. Encyc.* i. 320.—As. Min.
anthylloides, Pall. Astrag. 15 = inflatus.
Antilibani, *Bunge, Astrag.* i. 55; ii. 90.—Syria.
Antiselli, *A. Gray, in Bot. Calif.* i. 152.—Calif.
antoninus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 343.—Am. bor. occ.
Apollinens, *Boiss. Diagn.* Ser. II. ii. 27.—Graecia.
apricus, *Bunge, Astrag.* i. 40; ii. 58.—Persia.

ASTRAGALUS:—
arabicus, Ehrenb. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 121 = prolixus.
arabicus, Ehrenb. ex Bunge, Astrag. i. 9; ii. 6 = fatinensis.
arabicus, *Kotschy, in Sitzb. Acad. Wien, Math.-Nat. lii. Abth.* I. (1866) 264.—Arabia.
arborescens, Bunge, Astrag. ii. 225 = paucijugus.
Arbuscula, *Pall. Astrag.* 19. n. 24. t. 17.—Sibir.
Arbuscula, Willd. ex Ledeb. Fl. Ross. i. 633 = fruticosus.
arcticus, *Bunge, Astrag.* i. 23; ii. 27.—Europ. arct.
arcticus, Spreng. Syst. iv. Cur. Post. 288 = Oxytropis uralensis.
arcticus, *Willd. Enum. Hort. Berol. Suppl.* 51.—Hab.?
arcuatus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 407.—As. Min.
arcuatus, Steph. ex Ledeb. Fl. Ross. i. 649 = scaberrimus.
arenarius, Lapeyr. Hist. Abr. Pl. Pyr. 429 = baionnensis.
arenarius, *Linn. Sp. Pl.* 759.—Dania; Suecia.
arenarius, O. F. Muell. in Fl. Dan. t. 614 = danicus.
arenarius, Pall. Astrag. t. 34 = hypoglottis.
arenicola, *Pomel, Nouv. Mat. Fl. Atl.* 184.—Algeria.
arenosus, S. G. Gmel. ex Pall. Astrag. 25 = hyrcanus.
arequipensis, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* I. (1843) 17.—Peruv.
argacus, *Boiss. & Bal. Diagn.* Ser. II. vi. 53.—Cappadoc.
arganaticus, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 23.—As. temp.
argentatus, Pall. Astrag. 60. t. 48 = Oxytropis argentata.
argenteus, Bertol. ex Vis. in Flora, xii. (1829) I. Erg. 18 = Muelleri.
argophyllus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 331 = glareosus.
arguricus, *Bunge, Astrag.* i. 103; ii. 181.—Mont. Ararat.
argutensis, *Bunge, l. c.* i. 98; ii. 170.—Armenia.
argyrophyllus, *Boiss. & Gaill. ex Boiss. Fl. Orient.* ii. 358.—Syria.
argyrostachyus, *Boiss. Diagn.* Ser. I. vi. 41.—Persia.
argyrothamnus, *Boiss. l. c.* ix. 89.—Syria.
aridus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 223.—Am. bor. occ.
aristatus, *L'Hérit. Stirp. Nov.* 170.—Europ. austr.
aristatus, Sieber, ex Steud. Nom. ed. II. i. 159 = angustifolia.
aristatus, Sibth. & Sm. Fl. Graec. viii. 25. t. 734 = Parnassi.
arizonicus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 398.—Arizona.
arkalycensis, *Bunge, Astrag.* i. 139; ii. 238.—As. temp.
armatus, *Willd. Sp. Pl.* iii. 1330.—Numidia.
armeniacus, *Boiss. Diagn.* Ser. I. ii. 86.—Armenia.
armeriastrum, Decne. ex Boiss. Fl. Orient. ii. 387 = cephalanthus.
arnacantha, *Bieb. Fl. Taur. Cauc.* ii. 205.—Reg. Caucas.
arnoceras, Bunge, Astrag. i. 13; ii. 14 = brachyceras.
arpilobus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 336.—As. temp.
arrectus, *A. Gray, in Proc. Am. Acad.* viii. (1870) 289.—Am. bor.
Arthu-Schottii, A. Gray, l. c. vi. (1864) 209 = Coulteri.
artipes, *A. Gray, l. c.* xiii. (1878) 370.—Am. bor.
asaphes, *Bunge, Astrag. Fedsch.* 251.—As. temp.
asciocalyx, *Bunge, Astrag.* i. 137; ii. 234.—As. Min.
ascophorus, Fisch. ex Bunge, l. c. ii. 60 = brachystachys.
askius, *Bunge, l. c.* i. 120; ii. 208.—Persia.
aspadanus, *Bunge, l. c.* i. 63; ii. 107.—Persia.
asper, *Jacq. Ic. Rar.* i. 13. t. 152.—Europ. austr.; Reg. Caucas.
asperulus, Dufour, in Ann. Sc. Gén. vii. (1820) 295 = A. epiglottis.
Asterias, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 267 = cruciatus.
astraboides, *Pomel, Nouv. Mat. Fl. Atl.* 186.—Algeria.
atlanticus, *Ball, in Journ. Bot.* xi. (1873) 306.—Afr. bor. occ.
atratus, *S. Wats. in Bot. King, Exp.* 69.—Am. bor. occ.
atropatanus, *Bunge, Astrag.* i. 82; ii. 138.—Persia.

ASTRAGALUS :—

atropatanus, Fisch. ex Bunge, Astrag. ii. 90 = pulchellus.
atropurpureus, *Boiss. & Heldr. Diagn.* Ser. I. ix. 54.—As. Min.
atticus, Nym. Consp. 190 = Wulfenii.
Aucheri, *Boiss. Diagn.* Ser. I. ii. 46.—Armenia.
Aucherianus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 398. = ptilodes.
auganus, *Bunge, Astrag.* i. 36 ; ii. 45.—Afghanist.
aulacolobus, *Boiss. Diagn.* Ser. I. ix. 64.—Syria ; Mesopotam.
aureus, *Willd. in Mém. Acad. Berl.* (1794–95) 29. t. 1.—Oriens.
Austinae, *A. Gray, Bot. Calif.* i. 156.—Calif.
australis, *Lam. Fl. Fr.* ii. 637 ; *Bunge, Astrag.* i. 22 ; ii. 24.—Mont. Pyren. ; Hispan.
austriacus, Delarb. Fl. Auv. ed. II. 466 = hamosus.
austriacus, *Linn. Sp. Pl.* 1070.—Europ. ; Reg. Caucas. ; Sibiria.
austriacus, Pall. ex Steud. Nom. ed. II. i. 159 = versicolor.
austriacus, Thore, ex DC. Prod. ii. 283 = baionnensis.
bachtiaricus, *Bunge, Astrag.* i. 38 ; ii. 53.—Persia.
bactrianus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 354.—Turkestan.
baeticus, *Linn. Sp. Pl.* ed. II. 1068.—Hispan. ; Sicil.
baghensis, *Bunge, Astrag.* i. 80 ; ii. 135.—Persia.
baibutensis, *Bunge, l. c.* i. 91 ; ii. 162.—Armenia.
baicalensis, *Bunge, l. c.* i. 139 ; ii. 238.—As. centr.
baicalensis, Pall. Astrag. 64. t. 52 = Oxytropis coerulea.
baicalia, Pall. Astrag. 93. t. 77. f. 1 = Oxytropis baicalia.
baionnensis, *Loisel. Fl. Gall.* ed. I. 474.—Gallia occ.
bakaliensis, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 268 (*Pl. Lehm.* 92. t. 14).—Persia.
bakuensis, *Bunge, Astrag.* i. 37 ; ii. 45.—Reg. Casp. ; Ind. bor. occ.
Balansae, *Boiss. Diagn.* Ser. II. vi. 59.—Cappadoc.
banaticus, Rochel, ex Nym. Consp. 189 = chlorocarpus.
Barba-Aronis, Ehrenb. ex Bunge, Astrag. ii. 65 = Sieberi.
Barba-Jovis, *DC. Astrag.* 188.—As. Min.
Barba-Jovis, Fenzl, ex Bunge, Astrag. ii. 149 = Fenzlii.
Barba-Mosis, *Ehrenb. ex Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 486.—Mont. Sinai.
barbatus, *Lam. Encyc.* i. 314.—Armenia.
barbatus, Sibth. & Sm. Fl. Graec. Prod. ii. 88 = Sibthorpianus.
Barrelieri, Dufour, in Ann. Sc. Gén. vii. (1820) 297 = incanus.
basianicus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 361.—Mesopotam.
bayonnensis, Auct. vide baionnensis.
Beckerianus, *Trautv. in Act. Hort. Petrop.* iv. (1876) 363.—Reg. Caucas.
Beckwithii, *Torr. & Gray, in Pacif. Rail. Rep.* iii. (1854) 120.—Am. bor. occ.
Behen, Bertol. in Nov. Comm. Bonon. vi. (1844) 233 (Misc. Bot. ii. 19) = Russelii.
Belangerianus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) 484 = leioclados.
Benthamianus, *Gill. in Hook. Bot. Misc.* iii. (1833) 187.—Chili.
Bergii, *Hieron. in Bol. Acad. Nac. Cordova*, iii. (1879) 343.—Reg. Argent.
Berteri, *Colla, in Mem. Acc. Torin.* xxxvii. (1834) 57.—Chili.
berytheus, *Boiss. & Blanche, Diagn.* Ser. II. ii. 33.—Syria.
berytius, *Bunge, Astrag.* i. 50 ; ii. 78.—Cataonia.
bethlehemiticus, *Boiss. Diagn.* Ser. I. ix. 85.—Palaestin.
bhotanensis, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 126.—Reg. Himal.
bicarinatus, Steud. Nom. ed. II. i. 159 = Gueldenstaedtia pauciflora.
bicolor, Desf. Cat. Hort. Par. ed. III. 473 = pictus.
bicolor, *Lam. Encyc.* i. 317.—Armenia.
bicolor, Steud. Nom. ed. II. i. 159 = uliginosus.
bicristatus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 75.—Calif.
bicuspis, *Boiss. Fl. Orient.* 1094 (emend.).—As. Min.
bicuspis, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 406.—Tibet. occ.
bidentatus, *H. B. & K. Nov. Gen. et Sp.* vi. 493.—Am. calid.

ASTRAGALUS :—

bidentatus, Ehrenb. ex Bunge, Astrag. ii. 17 = hispidulus.
Biebersteinii, *Bunge, l. c.* i. 114 ; ii. 194.—Reg. Caucas.
Bienerti, *Bunge, l. c.* i. 80 ; ii. 133.—Persia.
bifidus, Turcz. ex Ledeb. Fl. Ross. i. 605 = multicaulis.
biflorus, Linn. Mant. 273 = Crotalaria biflora.
biflorus, Pall. Reise, iii. 206 = Gueldenstaedtia pauciflora.
biflorus, *Viv. Fl. Lyb. Spec.* 44. t. 20.—Afr. bor.
Bigelovii, *A. Gray, Pl. Wright.* ii. 42.—Am. bor. ; Mexic.
bingoellianus, Kotschy & Boiss. ex Boiss. Fl. Orient. ii. 357 = muschianus.
biovulatus, *Bunge, Astrag.* i. 11 ; ii. 11.—Persia.
biserratus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = A. Biserrula.
Biserrula, *Bunge, Astrag.* i. 11 ; ii. 9.—Persia.
bisulcatus, *A. Gray, in Pacif. Rail. Rep.* xii. 42. t. 1.—Am. bor. occ.
bludhistanus, *Bunge, Astrag.* i. 76 ; ii. 129.—Beluchist.
Bodeanus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 435.—Persia.
boeticus, *Linn. Sp. Pl.* 758.—Reg. Mediterr.
bohemicus, Mayer, in Abh. Boehm. Ges. iii. (1787) 314. t. 1 = virgatus.
Boissieri, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 324.—Hispan.
Boissieri, Bunge, Astrag. ii. 132, pro parte = creticus.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 337.—Calif.
bombycinus, *Boiss. Diagn.* Ser. I. ii. 50.—Aegypt. ; Arab. ; Persia.
Bonanni, J. & C. Presl, Delic. Prag. (1822) 42 = leucophaeus.
Bonannianus, Bunge, Astrag. ii. 242 = depressus.
bounacanthus, *Boiss. Fl. Orient.* ii. 370.—Lycia.
bounophilus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 99.—Persia.
Bourgaeanus, *Coss. Notes Crit.* 160.—Hispan.
Bourgovii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 227.—Am. bor. occ.
brachybius, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 267 = hamosus.
brachybotrys, *Bunge, Del. Sem. Dorp.* (1839) 8 ; *et in Linnaea*, xiv. (1840) *Litt.* 116.—Sibir. alt.
brachycalyx, *Fisch. ex Boiss. Diagn.* Ser. I. ix. 90 ; *et in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 350.—Kurdist.
brachycarpus, *Bieb. Fl. Taur. Cauc.* ii. 201.—Reg. Caucas.
brachycentrus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 337.—Persia.
brachyceras, C. Koch, ex Bunge, Astrag. ii. 7 = tribuloides.
brachyceras, *Ledeb. Cat. Hort. Dorp.* (1822) 3.—Syria ; Persia ; Tauria.
brachycladus, *Boiss. Fl. Orient.* ii. 393.—Persia.
brachylobus, *DC. Prod.* ii. 285.—Reg. Casp.
brachyodontus, *Boiss. Diagn.* Ser. I. ii. 46.—Persia.
brachyphyllus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 394 = condensatus.
brachypodus, Boiss. Diagn. Ser. I. ii. 69 = lagurus.
brachypterus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 397.—Oriens.
brachypus, *Schrenk, Enum. Pl. Nov.* 79.—Soongaria.
brachystachys, *DC. Prod.* ii. 303.—Syria ; Mesopotam.
brachytropis, *C. A. Mey. Verz. Pfl. Cauc.* 140.—Reg. Caucas.
Brackenbridgei, *A. Gray, Bot. U. St. Expl. Exped.* i. 416.—Peruv.
bracteolatus, DC. Astrag. 208 = angustifolius.
bracteosus, *Boiss. & Noë, Diagn.* Ser. II. ii. 31.—Anatolia.
bracteosus, C. A. Mey. ex Bunge, Astrag. ii. 185 = hyalolepis.
bracteosus, Klotzsch, in Bot. Ergeb. Waldem. Reise, 160 = melanostachys.
Brandegei, *Porter & Coult. Fl. Col.* 24.—Am. bor. occ.
brahuicus, *Bunge, Astrag.* i. 119, 205.—Beluchist.
brazoënsis, *Buckl. in Proc. Acad. Sc. Philad.* '1861' (1862) 452.—Texas.
brevicarinatus, DC. Astrag. 241. t. 49 = Gueldenstaedtia pauciflora.
breviflorus, DC. l. c. 194. t. 31 = eriocephalus.
breviflorus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 358 = polyanthus.

ASTRAGALUS :—

breviflorus, Fenzl, ex Bunge, Astrag. ii. 149 = Fenzlii.
brevifolius, *Ledeb. Fl. Alt.* iii. 334.—Sibir. alt.
brevipes, *Bunge, Astrag.* i. 99 ; ii. 172.—Persia.
brevirostris, Poir. Encyc. Suppl. i. 528 = Oxytropis brevirostra.
Breweri, *A. Gray, in Proc. Acad. Calif.* iii. (1863) 103.—Calif.
Bruguieri, *Boiss. Diagn.* Ser. I. ix. 102.—Persia.
Buceras, *Willd. Enum. Hort. Berol. Suppl.* 51.—Ins. Maurit.
buchtormensis, *Pall. Astrag.* 176. t. 62.—Sibiria.
buchtormensis, Willd. ex Ledeb. Fl. Ross. i. 585 = Oxytropis Pallasii.
Buhseanus, *Bunge, Astrag.* i. 120 ; ii. 207.—Persia.
Bulla, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 438 = cephalanthus.
bullarius, Pall. Reise, iii. 213 = Oxytropis mixotricha.
Bungeanus, *Boiss. Fl. Orient.* ii. 436.—Reg. Caucas.
Burkeanus, *Benth. ex Harv. Thes. Cap.* i. 52. t. lxxxii.—Afr. austr.
Bustillosii, *Clos, in C. Gay, Fl. Chil.* ii. 117.—Chili.
byzantinus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) 379 = prusianus.
cabulicus, Boiss. Diagn. Ser. I. ix. 48 = Oxytropis cabulica.
cachinalensis, *Phil. Fl. Atac.* 15.—Chili.
cadmicus, *Boiss. Diagn.* Ser. I. ii. 44.—As. Min.
caelestis, *Boiss. Fl. Orient.* ii. 470.—Reg. Caucas.
caeruleus, Pall. Reise, iii. 293 = Oxytropis caerulea.
caeruleus, *Hort. Par. ex Bunge, Astrag.* i. 103 ; ii. 184.—Zeylan.
caespitosus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 230.—Am. bor. occ.
caespitosus, Pall. Astrag. 70. t. 57 = Oxytropis caespitosa.
calabricus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 402.—Calabria.
californicus, *Greene, in Bull. Calif. Acad.* n. 3 (1885) 157.—Calif.
callichrous, *Boiss. Diagn.* Ser. I. ix. 62.—Palaestin.
calliphysa, *Bunge, Astrag.* i. 76 ; ii. 128.—Persia.
callistachys, *Buhse, in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 463.—Persia.
calophyllus, *Boiss. & Heldr. Diagn.* Ser. II. ii. 26.—Anatolia.
calvescens, *Bunge, Astrag.* i. 42 ; ii. 65.—Persia.
calycinus, *Bieb. Fl. Taur. Cauc.* ii. 199.—Reg. Caucas.
calycosus, *Torr. ex S. Wats. in Bot. King, Exp.* 66.—Calif.
camelorum, *C. & W. Barbey, Herb. Levant*, 131. t. 3.—Aegypt.
campestris, *A. Gray, in Proc. Am. Acad.* vi. (1864) 229.—Am. bor. occ.
campestris, Linn. Sp. Pl. 761 = Oxytropis campestris.
camporum, *Benth. ex Bunge, Astrag.* ii. 3.—Afghanist.
camptoceras, *Bunge, l. c.* i. 12 ; ii. 12.—Cappadocia.
campylanthus, *Boiss. Diagn.* Ser. I. vi. 42.—Persia.
campylorhynchus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 23.—As. Min. ; Persia ; Soongaria.
campylosema, *Boiss. Diagn.* Ser. I. ii. 82.—Cilicia.
campylotrichus, *Bunge, Astrag. Fedsch.* 207.—As. temp.
canadensis, *Linn. Sp. Pl.* 757.—Am. bor.
canaliculatus, Willd. Enum. Hort. Berol. Suppl. 52 = scorpioides.
cancellatus, *Bunge, Astrag.* ii. 178.—Persia.
candicans, *Greene, in Bull. Calif. Acad.* n. 3 (1885) 156.—Calif.
candicans, Pall. Astrag. 61. t. 49 = Oxytropis candicans.
candidissimus, *Ledeb. Fl. Alt.* iii. 309.—Sibir. alt. ; Calif.
Candolleanus, *Boiss. Diagn.* Ser. I. ii. 80.—Persia.
Candolleanus, Royle, Illustr. Bot. Himal. 199 = Royleanus.
canescens, *Bunge, Astrag.* ii. 174.—Armenia.
canescens, DC. Astrag. 114. t. 16 = onobrychoides.
canescens, Soland. ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 34 = Solandri.
canescens, *Kit. in Linnaea*, xxxii. (1863) 625.—Hungar.
caniculatus, Willd. Enum. Hort. Berol. Suppl. 52 = scorpioides.
cano-ater, *Bunge, Astrag.* i. 50 ; ii. 80.—Cappadoc.
canus, *Bunge, l. c.* ii. 180.—Persia.
canus, Willd. ex Ledeb. Fl. Ross. i. 606 = multicaulis.
capillipes, *Fisch. ex Bunge, Astrag.* i. 20 ; ii. 21.—China.

ASTRAGALUS :—

capitatus, Lam. Fl. Fr. ii. 640 = A. Glaux.
capitatus, Linn. Sp. Pl. 755 = emarginatus.
Capito, *Boiss. & Hohen. Diagn.* Ser. I. ix. 40.—Persia.
eappadoeicus, *Boiss. Diagn.* Ser. I. ii. 74.—As. Min.
caprinus, Guss. Fl. Sic. Syn. ii. 315 = Huetii.
eaprinus, *Linn. Sp. Pl.* ed. II. 1071.—Barbaria ; Oriens.
caprinus, Pall. Reise, ii. 329 = dasyanthus.
Caraganae, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1838) 344.—Reg. Caucas. ; Persia.
carduchorum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 310.—Persia.
eariensis, *Boiss. Diagn.* Ser. I. ix. 56.—As. Min.
carinalis, Benth. ex Bunge, Astrag. i. 23 ; ii. 28 = himalayanus.
carnosus, Nutt. Gen. Am. ii. 100 = caryocarpus.
carnosus, Pursh, Fl. Am. Sept. ii. 740 = Sophora sericea.
carolinianus, Linn. Sp. Pl. 757 = canadensis.
earyocarpus, *Ker-Gawl. in Bot. Reg.* t. 176.—Am. bor.
caryocarpus, Torr. in Ann. Lyc. N. York, ii. (1828) 179 = plattensis.
earyolobus, *Bunge, Astrag.* i. 37.—Persia.
casapaltensis, *Ball, in Journ. Linn. Soc.* xxii. (1885) 36.—And. Peruv.
Casei, *A. Gray, in Bot. Calif.* i. 154.—Calif.
easchmirensis, *Bunge, Astrag.* i. 30 ; ii. 34.—Kashmir.
easpicus, *Bieb. Fl. Taur. Cauc.* ii. 204.—Reg. Casp. ; As. temp.
caspius, Hohen. ex Ledeb. Fl. Ross. i. 641 = praec.
eastanaeformis, *S. Wats. in Proc. Am. Acad.* xx. (1885) 361.—Arizona.
eastellanus, *Bunge, Astrag.* i. 9 ; ii. 5.—Hispan.
eatacamptus, *Bunge, l. c.* ii. 191.—Persia.
catalinensis, Nutt. in Journ. Acad. Sc. Phil. N. S. i. (1847) 152 = didymocarpus.
cataonicus, Boiss. Fl. Orient. ii. 247 = affghanus.
eataonicus, *Bunge, Astrag.* i. 107 ; ii. 188.—Cataonia.
caucasicus, Bieb. Fl. Taur. Cauc. ii. 205 = Marschallianus.
eaucasicus, *Pall. Astrag.* 2. t. 2.—Reg. Caucas.
caudatus, Pall. l. c. 62. t. 50 = Oxytropis songorica.
eauliculosus, *Boiss. Diagn.* Ser. II. ii. 25.—Armenia.
eauquenensis, *Phil. in Anal. Univ. Chil.* (1872) 691.—Chili.
Cedreti, *Boiss. Diagn.* Ser. I. ix. 39.—Mont. Libanus.
ecphalanthus, *DC. Astrag.* 169. t. 38.—Persia.
cephalanthus, Hohen. ex Bunge, Astrag. ii. 117 = Hohenackeri.
cephalanthus, Kotschy, ex Bunge, l. c. ii. 120 = schirasicus.
cephalonicus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 424 = aristatus.
ecphalonicus, *Presl, Bot. Bemerk.* 57.—Cephalonia.
Cephalotes, *Pall. Astrag.* 29. t. 24.—Reg. Caucas.
Cephalotes, [Soland. in] Russell, Aleppo, ed II. ii. 260 A. Andraehne.
cerasinus, *Baker, in Journ. Linn. Soc.* xviii. (1878) 47.—Afghan.
cerasocrenus, *Bunge, Astrag.* i. 93 ; ii. 167.—Persia.
ceratoides, *Bieb. Fl. Taur. Cauc.* iii. 492.—Sibir. alt.
ceratoides, Bunge, Enum. Alt. 71 = stenoceras.
Cervini, Thomas, ex Steud. Nom. ed. II. i. 160 = lapponica.
chaborasicus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 437.—Mesopotam.
chaetodon, *Bunge, in Mém. Sav. Etr. Acad. Pétersb.* vii. (1851) 272 (*Pl. Lehm.* 96).—Turkestan.
chaetodon, Torr. ex A. Gray, in Proc. Am. Acad. vi. (1866) 194 = Spaldingii.
chaetolobus, *Bunge, in Mém. Sav. Etr. Acad. Pétersb.* ii. (1835) 592.—Sibir. alt.
chaetopodus, *Bunge, Astrag.* ii. 208.—Persia.
chalaranthus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 390.—Persia.
Chamaelcuce, *A. Gray, in Ives, Rep. Bot.* 10.—Am. bor. occ.
Chardini, *Boiss. Diagn.* Ser. I. ii. 74.—Persia.
chartaceus, *Ledeb. Fl. Ross.* i. 634.—Persia.
chartostegius, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 310.—Oriens ; Persia.
Chaubardi, Bunge, Astrag. i. 118 ; ii. 202 = Wulfeni.
chinensis, *Linn. f. Decad.* i. t. 3.—China.
chionobius, *Bunge, Astrag.* ii. 74.—Persia.
chionophilus, *Boiss. & Heldr. Diagn.* Ser. I. ix. 52.—As. Min.
chionophyllus, Kotschy, ex Bunge, Astrag. ii. 76 = depressus.

ASTRAGALUS :—

chius, Boiss. & Orph. ex Boiss. Fl. Orient. ii. 370 = trojanus.
chiwensis, *Bunge, Astrag. Fedsch.* 247.—Khiva.
chloranthus, Pall. Astrag. 30. t. 25 = asper
chlorocarpus, Griseb. Spicil. Fl. Rumel. i. 50 = A. Onobrychis.
chlorocyaneus, *Boiss. & Reut. Diagn.* Ser. I. ix. 56.—Hispan. ; Afr. bor. occ.
chlorodontus, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 29.—As. temp.
chlorophaeus, Bunge, Astrag. i. 50 ; ii. 79 = oxytropifolius.
chlorosphaerus, *Boiss. & Noë, Diagn.* Ser. II. ii. 28.—Oriens.
chlorostachys, Hook. f. & Thoms. ex Bunge, Astrag. ii. 28 = carinalis.
chlorostachys, *Lindl. in Trans. Hort. Soc.* vii. (1830) 249.—Reg. Himal.
chlorostegius, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 413.—Mesopotam.
choicus, Bunge, Astrag. i. 119 ; ii. 204 = latifolius.
chorassanicus, *Bunge, l. c.* i. 83 ; ii. 145.—Persia.
chordorrhizus, *Fisch. ex Bunge, l. c.* i. 52 ; ii. 86.—Reg. Caucas.
chorinensis, *Bunge, l. c.* i. 22 ; ii. 24.—Sibir.
Christianus, Aucher, ex Bunge, l. c. 36 = diptherolobus.
Christianus, Heldr. ex Bunge, l. c. ii. 38 = pisidicus.
Christianus, *Linn. Sp. Pl.* 755.—As. Min. ; Syria.
Christianus, Russell, ex Bunge, Astrag. ii. 36 = alepicus.
Christianus, Pinard, ex Bunge, l. c. ii. 38 = columnaris.
Christianus, Siev. in Pall. N. Nord. Beitr. vii. 293 = Sieversianus.
Christianus, Sibth. & Sm. Fl. Graec. Prod. ii. 85 = graecus.
Christianus, Vahl, Symb. Bot. i. 57 = tomentosus.
chromolepis, Boiss. & Hohen. Diagn. Ser. I. ix. 84 = aureus.
chrysanthus, *Boiss. & Hohen. l. c.* 73.—Persia.
chrysochlorus, Boiss. & Kotschy, in Boiss. l. c. Ser. II. v. 86 = pannosus.
chrysomallus, *Bunge, Astrag. Fedsch.* 310.—As. temp.
chrysophyllus, *Boiss. Diagn.* Ser. I. ix. 38.—Syria.
chrysopterus, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 32.—China.
chrysostachys, *Boiss. Diagn.* Ser. I. ii. 69.—Persia.
chrysotrichus, *Boiss. l. c.* 75.—Persia.
chtonoecephalus, *Boiss. & Bal. Diagn.* Ser. II. vi. 57.—Cappadoc.
Cicer, *Linn. Sp. Pl.* 757.—Europ. austr. ; Reg. Caucas. ; Sibiria.
cicerellus, *Boiss. & Bal. Diagn.* Ser. II. vi. 53.—Cappadoc.
cicerifolius, *Royle, ex Bunge, Astrag.* i. 44 ; ii. 70.—Reg. Himal. ; Tibet. occ.
cicerifolius, Royle, ex Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 404 = scariosus.
cicerinus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = A. Cicer.
cilicius, *Boiss. Diagn.* Ser. I. ii. 59.—Cilicia.
ciliolatus, *Benth. ex Bunge, Astrag.* i. 27 ; ii. 33.—Reg. Himal.
cinerascens, DC. ex Steud. Nom. ed. II. i. 160 = rupifragus, testiculatus.
cinereus, Bess. Enum. Pl. Volh. 29 = pallescens.
cinereus, DC. Astrag. 178 = reduncus.
cinereus, Turcz. Fl. Baic.-Dah. i. 322 = multicaulis.
cinereus, *Willd. Sp. Pl.* iii. 1315.—Armenia.
citrinus, *Bunge, Astrag.* i. 34 ; ii. 42.—Persia.
clandestinus, Roth, ex Spreng. Syst. iii. 300 = cymbiformis.
Clausii, C. A. Mey. in Goebel, Reise, ii. 265. t. 5 = vulpinus.
clavatus, *DC. Astrag.* 230. t. 44.—Armenia.
clavatus, Ledeb. Fl. Ross. i. 648 = brachycarpus.
Clevelandi, *Greene, in Torrey Bot. Club.* ix. (1882) 121.—Am. bor.
Closianus, *Phil. in Anal. Univ. Chil.* (1872) 691.—Chili.
Clusii, *Boiss. Diagn.* Ser. I. ix. 101.—Hispan.
Clusii, *Pollini, ex Steud. Nom.* ed. I. 85 ; ed. II. i. 160.—Italia.
coarctatus, *Trautv. in Act. Hort. Petrop.* iv. (1876) 365.—Armenia.
cobrensis, *A. Gray, Pl. Wright.* ii. 43, in adnot.—Am. bor. occ.

ASTRAGALUS :—

cognatus, *C. A. Mey. in Schrenk, Enum. Pl. Nov.* 81.—As. centr.
collinus, *Boiss. Fl. Orient.* ii. 438.—As. Min.
collinus, *Dougl. ex Hook. Fl. Bor. Am.* 141, in syn.—Am. bor. occ.
eolumnaris, *Boiss. Diagn.* Ser. I. ix. 70.—As. Min.
eolutcocarpus, *Boiss. l. c.* 65.—Tibet. occ. ; Afghanist.
eoluteoides, *Willd. in Mém. Acad. Berl.* (1794-95) 27.—Syria.
commixtus, *Bunge, in Arb. Nat. Ver. Riga*, i. (1847) 246.—As. Min. ; Persia ; Turkest.
comosus, *Bunge, Astrag.* i. 56 ; ii. 93.—Persia.
compactus, Fisch. ex Bunge, in Bull. Soc. Nat. Mosc. (1853) II. 368 = zachlensis, strictifolius.
compactus, *Lam. Encyc.* i. 322.—Armenia.
compactus, C. A. Mey. Verz. Pfl. Cauc. 143 = Meyeri.
complanatus, *R. Br. ex Bunge, Astrag.* ii. 1.—Chin. bor.
eomplicatus, *Gill. in Hook. Bot. Misc.* iii. (1833) 187.—Chili.
compressus, *Ledeb. Fl. Alt.* iii. 304.—Sibir. alt.
concinnus, *Benth. ex Bunge, Astrag.* ii. 50.—Beluchist.
concretus, Benth. in Royle, Illustr. Bot. Himal. 199 = vicioides.
condensatus, *Ledeb. Fl. Ross.* i. 639.—Reg. Caucas.
conduplicatus, *Bertol. in Nov. Comm. Bonon.* vi. (1844) 231 (*Misc. Bot.* ii. 17).—Syria ; Mesopotam.
confertiflorus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 368.—Am. bor.
confertus, Benth. ex Bunge, Astrag. ii. 27 = hippocrepidis.
confusus, *Bunge, l. c.* ii. 216.—Persia.
Congdoni, *S. Wats. in Proc. Am. Acad.* xx. (1885) 360.—Calif.
congestus, *Baker, ex Aitch. in Journ. Linn. Soc.* xix. (1882) 158.—Ind. or.
conjunctus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 371.—Am. bor.
consanguineus, *Bong. & C. A. Mey. Verz. Pfl. Saisang-Nor*, 22. t. 3.—China.
eontortuplicatus, *Linn. Sp. Pl.* 758.—Europ. austr. occ. ; Sibiria ; Ind. or.
contortuplicatus, Sibth. & Sm. Fl. Graec. viii. 21. t. 729 = cruciatus.
Cooperi, *A. Gray, Man. Bot. U. St.* ed. II. 98.—Am. bor.
eordatus, *Bunge, Astrag.* i. 69 ; ii. 114.—Persia.
eoriaceus, *Hemsl. Biol. Centr. Am. Bot.* i. 263.—Mexic.
corniculatus, *Bieb. Fl. Taur. Cauc.* iii. 492.—Bessarab. ; Podol.
corniculatus, Turcz. Cat. Fl. Baic. 356 = angarensis.
cornutus, *Bunge, Astrag.* i. 76 ; ii. 130.—Afghanist.
cornutus, Pall. Reise, i. 499 = vimineus.
Coronilla, *Bunge, Astrag.* ii. 10.—Persia.
corrugatus, *Bertol. Amoen. Ital.* 38.—Aegypt. ; Arab. ; Pers. ; Afghanist.
corydalinus, *Bunge, in Act. Hort. Petrop.* iii. (1875) 101.—Turkestan.
Cossonii, Bunge, Astrag. i. 118 ; ii. 192 = chlorocyaneus.
eostatus, *Bunge, l. c.* i. 35 ; ii. 61.—Afghanist.
Coulteri, *Benth. Pl. Hartw.* 307.—Calif.
Cracea, *DC. Astrag.* 101. t. 9.—Peruv.
crassicarpus, Fras. Cat. ex DC. Prod. ii. 287 = caryocarpus.
crassicaulis, R. Grah. in Wall. Cat. n. 5932 = Caragana crassicaulis.
erassinervius, *Boiss. Diagn.* Ser. II. v. 88.—Kurdistan.
crassipes, Fras. ex Steud. Nom. ed. II. i. 160 = caryocarpus.
crassispinus, *Bunge, Astrag.* i. 132 ; ii. 230.—Afghanist.
crassus, *Bunge, l. c.* i. 42 ; ii. 63.—Persia.
crenatus, Schult. Obs. 186 = corrugatus.
crenophilus, *Boiss. Diagn.* Ser. I. ix. 90.—Kurdistan.
eretaceus, *Boiss. l. c.* Ser. II. v. 84.—Syria.
cretaceus, Pall. in Nov. Act. Petrop. x. (1776) 56 = tauricus.
cretensis, Pall. Astrag. 5 = caspicus.
creticus, Boiss. Voy. Bot. Espagne, 181 = Boissieri.
creticus, DC. ex Boiss. Diagn. Ser. I. ix. 82 = cruentiflorus.
creticus, *Lam. Encyc.* i. 321.—Graecia.
creticus, Pinard, ex Bunge, Astrag. ii. 160 = ptiloides.
creticus, Willd. in Mem. Acad. Berl. (1794-95) 23 = echinoides.

ASTRAGALUS:—
creticus, Sibth. Fl. Graec. viii. 26. t. 736 = prusianus.
criacanthus, Stev. ex Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 331 = arnacantha.
crinitus, *Boiss. Diagn.* Ser. I. ii. 55.—Armenia.
cristatus, Gouan, Illustr. 50 = pentaglottis.
Cristophi, *Trautv. in Act. Hort. Petrop.* viii. (1884) 378.—Rossia.
Crotalariae, A. Gray, in Proc. Am. Acad. vi. (1864) 216 = oocarpus.
cruciatus, *Link, Enum. Hort. Berol.* ii. 256.—As. Min.; Syria.
cruciatus, Auct. ex Bunge, Astrag. ii. 8 = pseudostella.
cruciatus, Auct. quib. ex Boiss. Fl. Orient. ii. 226, in nota = polyactinus.
cruentiflorus, *Boiss. Diagn.* Ser. I. ix. 82.—Syria.
cruentus, Balb. Cat. Taur. App. i. 8 = corrugatus.
Cruikschankii, *Griseb. in Goett. Abh.* xxiv. (1879) 103.—Reg. Argent.
cryptanthus, *Wedd. Chlor. And.* ii. 259.—Am. austr.
cryptocarpos, *DC. Astrag.* 187.—Armenia.
cucullaris, *Boiss. Diagn.* Ser. I. vi. 39.—Mesopotam.
cumanus, Bieb. ex Ledeb. Fl. Ross. i. 607 = lasioglottis.
cumanus, Willd. ex Steud. Nom. ed. II. i. 160, 159 = aduncus.
cuneifolius, *Bunge, Astrag.* i. 44; ii. 68.—Afghan.
curtipes, *A. Gray, in Proc. Calif. Acad.* iii. (1863) 103.—Am. bor. occ.
curviflorus, *Boiss. Diagn.* Ser. I. ii. 67.—Persia.
curvirostris, *Boiss. l. c.* Ser. I. vi. 38.—Oriens.
Cuscutae, *Bunge, Astrag.* i. 120; ii. 207.—Reg. Caucas.
Cusickii, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 370.—Am. bor.
cuspidatus, Bunge, Astrag. i. 117; ii. 198 = schizopterus.
cyaneus, *A. Gray, Pl. Fendl.* 34.—Am. bor. occ.
cygneus, Fisch. ex Bunge, Astrag. ii. 190 = subbijugus.
cylindraceus, *DC. Astrag.* 223.—Armenia.
cylindraceus, C. A. Mey. Verz. Pfl. Cauc. 143 = macrourus.
cylleneus, *Boiss. & Heldr. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 355.—Graecia.
cymbaecarpus, *Brot. Fl. Lusit.* ii. 167.—Lusit.
cymbiformis, Willd. ex Brot. Phyt. 59 = cymbaecarpus.
cymbostegis, *Bunge, Astrag.* i. 85; ii. 14.—Cataonia.
cyprius, *Boiss. Fl. Orient.* ii. 469.—Ins. Cyprus.
cyrenaicus, *Coss. in Bull. Soc. Bot. Fr.* xix. (1872) 80.—Cyren.
cyrtobasis, *Bunge, ex Boiss. Fl. Orient.* ii. 497.—Turkestan.
cyrtoides, A. Gray, in Proc. Am. Acad. vi. (1864) 201 = Gibbsii.
cysticalyx, *Ledeb. Fl. Ross.* i. 643.—Rossia.
cytisoides, *Bunge, Astrag.* i. 128; ii. 224.—Turkestan.
dacicus, *Heuff. in Verh. Zool. Bot. Ges. Wien*, (1858) 93.—Europ.
dactylocarpus, *Boiss. Diagn.* Ser. I. ii. 60.—Mesopotam.
daenensis, *Boiss. l. c.* Ser. I. vi. 35.—Persia.
daguricus, Pall. Astrag. 88. t. 72 = Oxytropis prostrata.
dahuricus, *DC. Prod.* ii. 285.—Dahuria; China.
dahuricus, Patrin, ex DC. l. c. 277 = Oxytropis dissitiflora.
dalguricus, Willd. Sp. Pl. iii. 1301 = Oxytropis prostrata.
dalmaticus, Bunge, ex Nym. Consp. 190 = Soyeri.
Daltonianus, Bunge, Astrag. i. 44; ii. 69 = polyacanthus.
damascenus, *Boiss. & Gaill. Diagn.* Ser. II. vi. 56.—Syria.
damascenus, DC. Prod. ii. 298 = coluteoides.
danicus, *Retz. Obs.* iii. 41.—Europ.
Darumbium, *C. Gay, Fl. Chil.* ii. 112.—Chili.
dasyanthus, *Pall. Astrag.* 79. t. 65; *Reise*, iii. 749.—Hungar.; Sibir.; Reg. Cauc.
dasycephalus, Bess. ex Stev. in Bull. Soc. Nat. Mosc. (1832) 264 = roseus.
dasyglottis, Fisch. ex DC. Prod. ii. 282 = A. Hypoglottis.
dasyglottis, Pall. ex Steud. Nom. ed. II. i. 160 = pentaglottis.
dasyphyllos, Pall. Astrag. 91. t. 75 = Oxytropis lanata.
dauricus, C. Koch, in Linnaea, xv. (1841) 719 = Stevenianus.
dauricus, Auct. = dahuricus.

ASTRAGALUS:—
davuricus, Ledeb. Ic. Fl. Ross. t. 318 = dahuricus.
davuricus, Patrin, ex Ledeb. Fl. Ross. i. 597 = Oxytropis dissitiflora.
dealbatus, Ehrenb. ex Bunge, Astrag. ii. 164 = Roussaeanus.
dealbatus, Pall. Astrag. t. 23. f. 1 = albicaulis.
debilis, *A. Gray, in Proc. Acad. Sc. Philad.* 1863 (1864) 60.—Am. bor.
decemjugus, Bunge, Astrag. i. 43; ii. 67 = psilacanthus.
decipiens, Bunge, l. c. i. 107; ii. 188 = Sibthorpianus.
declinatus, Salisb. Prod. 337 = monspessulanus.
declinatus, *Willd. Sp. Pl.* iii. 1294.—Armenia.
decumbens, Boiss. Fl. Orient. ii. 429 = leucocyaneus.
decumbens, *A. Gray, in Proc. Am. Acad.* vi. (1864) 229.—Am. bor. occ.
decurrens, *Boiss. Diagn.* Ser. I. vi. 40.—Mesopotam.
deflexus, Pall. in Act. Acad. Petrop. ii. (1779) 268. t. 15 = Oxytropis deflexa.
deinacanthus, *Boiss. Diagn.* Ser. I. ix. 76.—Syria.
deluensis, Bunge, Astrag. i. 55; ii. 91 = griseus.
demavendicus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 70.—Persia.
demissus, Boiss. & Heldr. Diagn. Ser. I. ix. 50 = amoenus.
dendridium, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 427 = lycioides.
dendroides, *Kar. & Kir. l. c.* xv. (1842) 339.—Dahuria.
densiflorus, *Kar. & Kir. l. c.* 329.—Reg. Himal.; Sibir.
densifolius, *Lam. Encyc.* i. 317.—Armenia.
densifolius, Torr. in Pacif. Rail. Rep. vii. 10 = Menziesii.
denudatus, Kotschy, ex Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 335 = myriacanthus.
denudatus, Stev. l. c. 328 = microcephalus.
depauperatus, *Ledeb. Fl. Alt.* iii. 314.—Sibir. alt.
dependens, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 471.—China occ.
depressus, Bertol. in Nov. Comm. Bonon. vi. (1844) 232 (Misc. ii. 232) = cruciatus.
depressus, *Linn. Cent. Pl.* ii. 29; *Amoen. Acad.* iv. 327.—Europ. austr.; Oriens.
depressus, Pall. Reise, i. 239 = diffusus.
depressus, Sibth. & Sm. Fl. Graec. viii. 24. t. 733 = leucophaeus.
derbendicus, *Bunge, Astrag.* i. 40; ii. 55.—Persia.
desertorum, *Lehm. ex Spreng. Syst.* iii. 300.—Deserta Asiae.
dianthus, *Bunge, Astrag. Fedsch.* 283.—As. centr.
diaphanus, Dougl. ex Hook. Fl. Bor. Am. i. 151 = lentiginosus.
dichopterus, Pall. Astrag. t. 39 = austriacus.
dictyocarpus, *Boiss. Diagn.* Ser. I. ii. 84.—Syria.
dictyolobus, *C. A. Mey. ex Bunge, Astrag.* i. 60; ii. 98.—Persia.
dictyophysus, *Reut. ex Bunge, l. c.* i. 74; ii. 124.—Pontus.
dicystis, *Bunge, Astrag. Fedsch.* 225.—As. centr.
didymocarpus, *Hook. & Arn. Bot. Beech. Voy.* 334.—Calif.
didymophysus, *Bunge, Astrag.* i. 113; ii. 193.—Persia.
diffusus, Willd. Sp. Pl. iii. 1321 = dolichophyllus.
dilectus, *Phil. in Anal. Univ. Chil.* (1872) 690.—Chili.
dilutus, *Bunge, Del. Sem. Dorpat.* (1840) 7.—*Cf. Linnaea*, xv. (1841) *Litt.* 85.—Oriens.
diopogon, *Bunge, Astrag.* i. 70; ii. 117.—Beluchist.
dioriticus, Schott, ex Boiss. Fl. Orient. ii. 503 = Oxytropis dioritica.
dipelta, *Bunge, in Act. Hort. Petrop.* vii. (1880) 368.—Turkestan.
diphacus, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 342.—Am. bor. occ.
diphtherites, *Fenzl, in Flora*, xxvi. (1843) 396.—Mesopotam.
diphtherolobus, *Bunge, Astrag.* i. 31; ii. 36.—Syria.
diphyllus, Lam. Encyc. i. 330 = Oxytropis microphylla.
diphysus, *A. Gray, Pl. Fendl.* 34.—Am. bor. occ.
dipodurus, *Bunge, Astrag.* i. 88; ii. 156.—Syria.
dipsaceus, *Bunge, l. c.* i. 59; ii. 97.—Anatolia.
discolor, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 33.—Mongolia.
dispermus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 365.—Am. bor.
dissitiflorus, *Bunge, in Mém. Acad. Pétersb.* Sér. VI. vii. (1859) 585.—As. temp.

ASTRAGALUS:—
distans, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 128 = macropterus.
distans, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 472.—Persia.
distans, *A. Gray, Bot. U. St. Expl. Exped.* i. 412.—Patagonia.
distortus, *Torr. & Gray, Fl. N. Am.* i. 333.—Am. bor.
divaricatus, *Bunge, Astrag.* i. 93; ii. 166.—Persia.
diversifolius, A. Gray, in Proc. Am. Acad. vi. (1864) 230 = junceus.
Dodtii, *Phil. in Linnaea*, xxxiii. (1864-65) 48.—Am. austr.
dolichophyllus, *Pall. Astrag.* 84. n. 90. t. 68.—Reg. Caucas.
dolius, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 340.—Kurdistan.
Dombeyi, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) 483.—Peruv.
Donianus, *DC. Prod.* ii. 283.—Nepal.
dorcoceras, *Bunge, Astrag.* i. 13; ii. 14.—Cilicia; Mesopot.
dorychnoides, Scop. Delic. Insub. ii. 104 = lusitanicus.
dorycnioides, Dougl. ex G. Don, Gen. Syst. ii. 258 = succumbens.
Douglasii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 215.—Am. bor. occ.
drepanocarpus, Hochst. ex Bunge, Astrag. i. 14; ii. 16 = gyzensis.
drepanolobus, *A. Gray, in Proc. Am. Acad.* xix. (1883) 75.—Am. bor. occ.
drepanophorus, *Griseb. in Goett. Abh.* xxiv. (1879) 102.—Reg. Argent.
Drummondii, *Dougl. in Hook. Fl. Bor. Am.* i. 153.—Am. bor.
drupaceus, *Orphan. ex Boiss. Diagn.* Ser. II. ii. 32.—Graecia.
drusorum, *Boiss. Diagn.* Ser. I. ix. 78.—Syria.
dubius, DC. Astrag. 167. n. 115 = buchtormensis.
dyctiocarpus, Boiss. Diagn. Ser. I. ii. 84, sphalm. = dictyocarpus.
ebenoides, *Boiss. l. c.* Ser. I. ii. 70.—Persia.
ecbatanus, *Bunge, Astrag.* i. 71; ii. 118.—Persia.
Echegarayi, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 22.—Reg. Argent.
Echidna, *Bunge, Astrag.* i. 82; ii. 143.—Persia.
echinatus, Murr. Hort. Bot. Stirp. 222 = pentaglottis.
echinoides, L'Hérit. Stirp. Nov. 170 = angustifolius.
Echinops, *Aucher, ex Boiss. Diagn.* Ser. I. ii. 57.—Syria.
Echinus, Boiss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 280 = Boissieri.
Echinus, *DC. Astrag.* 197. t. 34.—Syria.
Echinus, C. A. Mey. Verz. Pfl. Cauc. 143 = caspius.
echioides, Willd. Sp. Pl. iii. 1326 = angustifolius.
edulis, *Dur. ex Coss. in Ann. Sc. Nat.* Sér. IV. i. (1854) 222.—Afr. bor.
effusus, *Bunge, Astrag.* i. 102; ii. 179.—Persia.
Ehrenbergii, *Bunge, l. c.* i. 61; ii. 102.—Syria.
elatus, *Boiss. & Bal. Diagn.* Ser. II. vi. 57.—Cappadoc.
elbrusensis, *Boiss. l. c.* Ser. I. ix. 45.—Persia.
elegans, *Bunge, Astrag.* i. 55; ii. 89.—Persia.
Elliotii, D. Dietr. Syn. Pl. iv. 1080 = obcordatus.
ellipsoideus, Bong. & Mey. Suppl. Alt. 84, partim = kurtschumensis.
ellipsoideus, *Ledeb. Fl. Alt.* iii. 319.—Sibir. alt.
elmaluensis, *Bunge, Astrag.* i. 121; ii. 211.—Lycia.
elongatus, *Willd. Sp. Pl.* iii. 1314.—As. Min.
elymaiticus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 346.—Persia.
emarginatus, *DC. Astrag.* 150.—Oriens.
emarginatus, *Labill. Ic. Pl. Syr. Dec.* i. 19.—Syria.
Emodi, *Steud. Nom.* ed. II. i. 161.—Reg. Himal.
enoplus, *Trautv. in Act. Hort. Petrop.* iv. (1876) 364.—Reg. Caucas.
entomophyllus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 259.—Persia.
ephippium, *Pomel, Nouv. Mat. Fl. Atl.* 183.—Algeria.
epiglossus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = epiglottis.
epiglottis, *Linn. Sp. Pl.* 759.—Reg. Mediterr.
epiglottoides, Willk. in Bot. Zeit. iii. (1845) 428 = A. Epiglottis.
episcopus, *S. Wats. in Proc. Am. Acad.* x. (1875) 346.—Am. bor. occ.
eremophilus, *Boiss. Diagn.* Ser. I. ii. 54.—Aegypt.; Arab.; Beluchist.

ASTRAGALUS:—
eremospartoides, *Regel, in Act. Hort. Petrop.* iii. (1875) 103.—Turkestan.
eremothamnus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 334.—Soongaria.
eriacmus, *Boiss. Fl. Orient.* ii. 345.—Cappadoc.
erianthus, Fisch. ex Boiss. Fl. Orient. ii. 328 = crenophilus.
erianthus, Willd. Sp. Pl. iii. 1332 = gummifer.
erinaceus, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 325.—Persia.
erinaceus, Presl, Bot. Bemerk. 57 = angustifolius.
eriocalyx, *Bunge, Astrag.* i. 86; ii. 150.—Cilicia.
eriocarpus, DC. Astrag. 237. t. 47 = mollis.
eriocarpus, *S. Wats. in Bot. King, Exp.* 71 & 440.—Am. bor. occ.
eriocaulos, DC. Astrag. 188 = microcephalus.
eriocephalus, Balansa, ex Bunge, Astrag. ii. 161 = acicularis.
eriocephalus, Waldst. & Kit. Pl. Rar. Hung. i. 45. t. 46 = dasyanthus.
eriocephalus, *Willd. Sp. Pl.* iii. 1333.—Armenia.
erioceras, *Fisch. & Mey. in Ledeb. Fl. Ross.* i. 626.—As. med.
eriolobus, Bunge, in Mém. Sav. Etr. Acad. Pétersb. ii. (1835) 591 = depauperatus.
erionotus, *Benth. ex Bunge, Astrag.* i. 33; ii. 40.—Afghan.
eriophyllus, *Boiss. Diagn.* Ser. I. ix. 39.—As. Min.
eriopodus, *Boiss. l. c.* Ser. I. ii. 48.—Persia.
eriosphaerus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 347.—Persia.
eriostylus, *Boiss. & Haussk. ex Boiss. l. c.* 316.—Persia.
erpocaulos, Boiss. Diagn. Ser. I. ix. 58 = tribuloides.
ervoides, *Hook. & Arn. Bot. Beech. Voy.* 417.—Mexic.
ervoides, Turcz. in Bull. Soc. Nat. Mosc. (1838) 90 = miniatus.
erythrolepidius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = erythrolepis.
erythrolepis, *Boiss. Diagn.* Ser. I. ii. 65.—Persia.
erythrotaenius, *Boiss. l. c.* vi. 39.—Mesopotam.
eschkerensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 328.—Persia.
eubrychioides, *Boiss. Diagn.* Ser. I. ix. 41.—Cappadoc.
eucephalus, *Boiss. l. c.* ii. 56.—Syria.
euoplus, *Trautv. in Act. Hort. Petrop.* iv. (1876) 364.—As. temp.
exaltatus, Bunge, Astrag. i. 25 = secundus.
eximius, *Bunge, Astrag. Fedsch.* 244.—As. centr.
expansus, Boiss. Diagn. Ser. I. ix. 43 = Kotschyanus.
exscapus, *Linn. Mant.* 275.—Europ.
exscapus, Auct. Transs.—Cf. Nym. Consp. 193 = pubiflorus.
fabaceus, DC. Prod. ii. 302 = utriger.
fabaceus, *Bieb. Fl. Taur. Cauc.* iii 496.—Reg. Caucas.
fabaceus, Jan, ex Nym. Consp. 193 = Huetii.
falcatus, Desf. Fl. Atlant. ii. 188 = falciformis.
falcatus, *Lam. Encyc.* i. 310.—Reg. Caucas.; Sibir.
falciformis, *Desf. Emend. Atl. ex DC. Astrag.* 176.—Algeria.
falcinellus, *Boiss. Diagn.* Ser. I. ix. 63.—Aegypt.
Falconeri, *Bunge, Astrag.* i. 4; ii. 2.—Tibet. occ.
fallax, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 340 = mesoleios.
fallax, *S. Wats. in Proc. Am. Acad.* xx. (1885) 362.—Mexic.
farctus, *Bunge, in Mem. Sav. Etr. Pétersb.* vii. (1851) 276 (*Pl. Lehm.* 100).—Turkestan.
fasciculatus, *Bertol. in Nov. Comm. Bonon.* vi. (1844) 23.—Hab. ?
fasciculifolius, *Boiss. Diagn.* Ser. I. ii. 73.—Persia.
fastidiosus, *Greene, in Bull. Calif. Acad.* n. 4 (1885) 186.—Calif.
fatmensis, *Hochst. ex Bunge, Astrag.* ii. 16.—Arabia; Beluchist.
Fendleri, *A. Gray, Pl. Wright.* ii. 44.—N. Mexic.
Fenzlii, *Boiss. Diagn.* Ser. II. v. 89.—Cilicia.
ferrugineus, *C. Gay, Fl. Chil.* ii. 114.—Chili.
filagineus, Boiss. Diagn. Ser. I. ix. 89 = gossypinus.
filamentosus, *Bunge, Astrag.* i. 39; ii. 51.—Kurdist.
filicaulis, *Fisch. & Mey. Enum. Pl. Turc. et Pers.* 256.—Reg. Casp.; Turkest.; Afghan.
filifolius, A. Gray, in Pacif. Rail. Rep. iv. 80 = pictus.
filifolius, *Clos, in C. Gay, Fl. Chil.* ii. 111.—Chili.
filiformis, Poir. Encyc. Suppl. i. 528 = tauricus.
filipes, *Torr. ex A. Gray, in Proc. Am. Acad.* vi. (1866) 226.—Am. bor. occ.
finitimus, *Bunge, Astrag.* i. 61; ii. 102.—Persia.

ASTRAGALUS:—
Fischeri, Buhse, in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 429 = mucronifolius.
Fischerianus, Bunge, Astrag. i. 90; ii. 160 = pilodes.
flaccidus, *Bieb. Fl. Taur.* ii. 196. n. 1484.—Reg. Caucas.
flaccidus, Ledeb. Fl. Ross. i. 607 = onobrychioides.
flavescens, *Boiss. Diagn.* Ser. I. ii. 64.—As. Min.
flavus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 335.—Am. bor.
flexuosus, *Dougl. ex Hook. Fl. Bor. Am.* i. 141, in syn.—Am. bor.
flexus, *Fisch. in Bull. Phys.-Math. Acad. Pétersb.* iii. (1844) 307, in nota.—Persia; Beluchist.; Soongaria.
floccosus, *Boiss. Diagn.* Ser. I. ii. 61.—Persia.
floribundus, Pall. Astrag. 47. t. 37 = Oxytropis floribunda.
floridus, *Benth. ex Bunge, Astrag.* i. 24; ii. 28.—Reg. Himal.
floridus, Scop. Delic. Insub. ii. 108 = Christianus.
florulentus, *Boiss. & Haussk. Fl. Orient.* ii. 317.—Persia.
fodinarum, *Boiss. & Noë, ex Bunge, Astrag.* i. 121; ii. 209.—Armenia.
foetidus, Vill. Prosp. 42 = Oxytropis foetida.
foliolosus, *Bunge, Astrag.* i. 74; ii. 125.—Anatolia.
follicularis, *Pall. Astrag.* 19. t. 14.—As. centr.
Fontanesii, *Coss. & Dur. ex Bunge, Astrag.* i. 75; ii. 126.—Algeria.
Forskahlei, *Boiss. Diagn.* Ser. I. ix. 101.—Aegypt.; Arab.; Syria.
fragiferus, *Bunge, Astrag.* i. 72; ii. 119.—Persia.
fragiformis, Willd. Sp. Pl. iii. 1261 = inflatus.
fragrans, *Willd. l. c.* 1294.—Orient.
Fraxinella, *Bunge, Astrag.* i. 39; ii. 52.—Persia.
fraxinifolius, *DC. Astrag.* 142. n. 71.—Armenia.
Frémontii, Torr. in Pacif Rail. Rep. iv. 80 = malacus, lentiginosus.
Fresenii, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 266.—Arab.
Frickii, *Bunge, Astrag.* i. 117; ii. 201.—Reg. Caucas.
frigidus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 219.—Am. bor.
Friwaldskianus, Boiss. Diagn. Ser. I. ii. 73 = physocalyx.
frutescens, Hort. ex Kunth, Ind. Hort. Berol. (1845) 12 = Brongniartia robinioides.
fruticosus, Asso, Oryctogn. Arag. t. 2. f. 3 = Clusii.
fruticosus, Forsk. Fl. Aegypt. Arab. 139 = tomentosus.
fruticosus, Steud. Nom. ed. II. i. 161 = Hedysarum fruticosum.
fruticosus, *Pall. Astrag.* 21. n. 24. t. 19.—Sibir. or.
fruticosus, Ledeb. Fl. Alt. iii. 303 = vimineus.
gaeobotrys, *Boiss. & Bal. in Boiss. Diagn.* Ser. II. vi. 52.—Cappadoc.
Gaillardoti, *Boiss. Fl. Orient.* ii. 375.—Syria.
galactites, Ledeb. Fl. Ross. i. 656, ex parte = brevifolius.
galactites, *Pall. Astrag.* 85. t. 65.—Sibir.; Dahur.
galegiformis, Hohen. Enum. Pl. Elisabethpol. 250 = A. Glycyphyllos.
galegiformis, *Linn. Sp. Pl.* 756.—Reg. Caucas.
galegiformis, Pall. Reise, ii. 192 = uliginosus.
galegoides, Nutt. Gen. Am. ii. 100 = racemosus.
Garbancillo, *Cav. Ic.* i. 59, in textu.—Peruv.
Garbanzillo, Pers. Syn. ii. 336 = praec.
Garvanzillo, Cav. Ic. i. t. 85 = Garbancillo.
Gebleri, *Fisch. ex Bong. Verz. Pfl. Saisang-Nor*, 24. t. 4.—Soongaria.
geminanus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 330.—Persia.
geminiflorus, *Humb. & Bonpl. Pl. Aequin.* i. 126.—Ecuador.
genargenteus, Moris, Stirp. Sard. Elench. 11 = sirinicus.
geniculatus, *Desf. Fl. Atlant.* ii. 186. t. 205.—Algeria.
genistoides, *Boiss. Diagn.* Ser. I. ix. 97.—Afghanist.
geobotrys, *Boiss. l. c.* Ser. II. vi. 52.—Phrygia.
geocyamus, *Boiss. Fl. Orient.* ii. 474.—Cappadoc.
georgicus, Boiss. Diagn. Ser. I. ii. 42 = Cephalotes.
Gerardi, *Janka, in Termész. Füzetek*, viii. (1884) 310.—Europ.
Gerardianus, R. Grah. in Wall. Cat. n. 5938 = Caragana Gerardiana.
gerensis, *Boiss. Diagn.* Ser. I. ix. 71.—Persia.
gerensis, Boiss. & Buhse, ex Bunge, Astrag. ii. 64 = spinescens.
Germanii, *Phil. in Linnaea*, xxviii. (1856) 621.—Am. austr.

ASTRAGALUS:—
Geyeri, *A. Gray, in Proc. Am. Acad.* vi. (1864) 214.—Am. bor. occ.
ghilanicus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 329.—Persia.
gibberosus, Bertol. in Nov. Comm. Bonon. vi. (1844) 231 (Misc. Bot. ii. 17) = Chesneya rhytidosperma.
gibbosus, DC. Prod. ii. 300 = Ammothamnus gibbosus.
Gibbsii, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 161. f. 50.—Calif.
giganteus, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 370.—Am. bor.
gilensis, *Greene, in Torrey Bot. Club*, viii. (1881) 97.—N. Mexic.
Gilliesii, *Phil. in Anal. Univ. Chil.* (1862) II. 393.—Am. austr.
gilvus, *Boiss. Diagn.* Ser. I. ix. 71.—Caria.
glaber, DC. Astrag. 118 = fragrans.
glaber, *Michx. Fl. Bor. Am.* ii. 66.—Am. bor.
glaber, Lam. Encyc. i. 525 = Oxytropis glabra.
glabrifolius, *Bunge, Astrag.* i. 93; ii. 167.—Afghanist.
glabriusculus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 204.—Am. bor. occ.
gladiatus, *Boiss. Diagn.* Ser. I. ii. 45.—As. Min.; Syria.
glanduliferus, *Debeaux, in Act. Linn. Soc. Bord.* xxxiii. (1879) 39.—China.
glareosus, *Dougl. ex Hook. Fl. Bor. Am.* i. 151.—Am. bor. occ.
glaucacanthos, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 481.—Persia.
glauciformis, *Pomel, Nouv. Mat. Fl. Atl.* 183.—Algeria.
glaucoides, S. G. Gmel. It. i. 127. t. 21 = Coronilla varia.
glaucophyllus, *Bunge, Astrag.* i. 118; ii. 200.—Anatolia.
glaucus, Bieb. Fl. Taur. Cauc. ii. 203 = vesicarius.
Glaux, *Linn. Sp. Pl.* 759.—Hispan.; Afr. bor. occ.
Glaux, Pall. Reise, ii. 464 = A. Hypoglottis.
gliciphillos, Neck. Delic. Gallo-Belg. ii. (1768) 312 = A. Glyciphyllos.
globiceps, *Bunge, in Act. Hort. Petrop.* vii. (1880) 372.—As. centr.
globiflorus, *Boiss. Diagn.* Ser. I. ix. 81.—Persia.
globigerus, C. Koch, in Linnaea, xix. (1847) 64 = hirsutus.
globosus, *Vahl, Symb. Bot.* i. 60.—Armenia.
glomeratus, *Ledeb. Fl. Alt.* iii. 327.—Sibir. alt.
glumaceus, *Boiss. Diagn.* Ser. I. ii. 69.—Persia.
glycyphylloides, *DC. Prod.* ii. 292.—Reg. Caucas.
Glycyphyllos, *Linn. Sp. Pl.* 758.—Europ.; Reg. Caucas.; Sibiria.
Gmelini, *Scop. Delic. Insub.* ii. 109.—Europ.
Gmelini, Steud. Nom. ed. II. i. 161 = Laxmanni.
gomboeformis, *Pomel, Nouv. Mat. Fl. Atl.* 187.—Algeria.
Gombo, *Coss. & Dur. ex Bunge, Astrag.* i. 42; ii. 63.—Algeria.
Gompholobium, *Benth. ex Bunge, l. c.* i. 39; ii. 50.—Beluchist.
goniatus, Nutt. ex Torr. & Gray, Fl. N. Am. i. 330 = hypoglottis.
gossypinus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 373.—Kurdist.; Persia.
Govanianus, R. Grah. in Wall. Cat. n. 5925 = chlorostachys.
gracilentus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 223.—Am. bor. occ.
gracilipes, *Benth. ex Bunge, Astrag.* i. 15; ii. 17.—Tibet.
gracilis, *Nutt. Gen. Am.* ii. 100.—Am. bor.
gracilis, Stev. ex Ledeb. Fl. Ross. i. 609 = Stevenianus.
gracilis, Torr. in Ann. Lyc. N. York, ii. (1828) 179 = microlobus.
graecus, *Boiss. Diagn.* Ser. I. ii. 57.—Graecia.
Grahami, Steud. Nom. ed. II. i. 161 = vicioides.
Grahamianus, Fisch. in Bull. Soc. Nat. Mosc. (1853) II. 406 = Jacquemontii.
Grahamianus, Royle, Illustr. Bot. Himal. 199. t. 36. f. 2 = polyacanthus
grammocalyx, *Boiss. & Hohen. Diagn.* Ser. I. ix. 52.—Persia.
granatensis, Lam. ex Steud. Nom. ed. I. 86; ed. II. i. 161 = A. Poterium.
granatensis, Lange, in Kjoeb. Vidensk. Meddel. (1865) 177 (Pugill. 372) = A. Glaux.
grandiflorus, *Bunge, Astrag. Fedsch.* 229.—As. centr.

ASTRAGALUS:—
grandiflorus, Linn. Sp. Pl. 761 = Hedysarum grandiflorum.
grandiflorus, Pall. Astrag. 57. t. 46 = Oxytropis grandiflora.
grandiflorus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 370.—Am. bor.
Grantii, *Bunge, Astrag.* i. 83; ii. 50.—Afghanist.
graveolens, *Buch.-Ham. in Wall. Cat.* n. 5929.—Reg. Himal.
Grayii, *Parry, ex S. Wats. in Am. Natural.* viii. (1874) 212.—Am. bor. occ.
Greenei, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 105.—N. Mexic.
Greggii, *S. Wats. l. c.* xvii. (1881–82) 343.—Am. bor. occ.
Griffithii, *Benth. ex Bunge, Astrag.* ii. 192.—Afghanist.
griseus, *Boiss. Diagn.* Ser. I. vi. 36.—Persia.
gryphus, *Coss. & Dur. ex Bunge, Astrag.* i. 12; ii. 13.—Algeria.
guatemalensis, *Hemsl. Biol. Centr. Am. Bot.* i. 264. Guatemala.
gudrunensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 462.—Assyria.
Guldenstaedtiae, *Bunge, Astrag.* i. 119; ii. 203.—Persia.
gummifer, *Labill. in Journ. Phys.* (1790) 46.—Syria.
guttatus, [*Soland. in*] *Russell, Aleppo.* ed. II. ii. 260.—Aleppo.
gymnolobus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 426.—Armenia.
gymnopodus, *Boiss. Diagn.* Ser. I. ix. 66.—Afghanist.
gypsophilus, *Rouy, in Bull. Soc. Bot. Fr.* xxxi. (1884) 278, nomen.—Gallia.
gyzensis, *Delile, Fl. Egypt. Suppl.* 64.—Afr. bor.; Arab.; Persia; Soongaria.
Haarbachii, *Sprun. in Boiss. Diagn.* Ser. I. ii. 50.—Graecia.
haematocarpus, *Bunge, Astrag.* i. 118; ii. 201.—Reg. Caucas.
halepensis, Fisch. ex Boiss. Fl. Orient. ii. 274 = alepicus.
halicacabus, *Lam. Encyc.* i. 220.—Armenia.
halicacabus, Ledeb. Fl. Ross. i. 643 = mesites.
Halleri, All. Fl. Pedem. i. 343 = Oxytropis foetida.
Hallii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 224.—Am. bor. occ.
Halodendron, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 21.—As. temp.
hamatus, Lam. ex Willk. & Lange, Prod. Fl. Hisp. iii. 271 = A. Stella.
hamosus, Balansa, ex Bunge, Astrag. ii. 17 = commixtus.
hamosus, Kotschy, ex Bunge, Astrag. ii. 14 = arnoceras.
hamosus, *Linn. Sp. Pl.* 758.—Reg. Mediterr.; Transcauc.; Ind. bor. occ
hamosus, Pall. Reise, iii. 653 = reduncus, trimestris.
Hancockii, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 471.—China bor.
harpilobus, Boiss. Fl. Orient. ii. 234, in syn. = gyzensis.
Hartwegi, *Benth. Pl. Hartw.* 10. n. 53.—Am. bor. occ.
hasbeyanus, *Boiss. Diagn.* Ser. I. ix. 77.—Syria.
Hauarensis, Boiss. l. c. Ser. I. ix. 63 = gyzensis.
Hausknechtii, *Bunge, Astrag.* i. 49; ii. 77.—Cataonia.
Haydenianus, *A. Gray, in Brandegee, Fl. S. W. Col.* 235.—Am. bor. occ.
hedysaroides, Linn. Sp. Pl. 756 = Hedysarum obscurum.
hedysaroides, Sievers, in Pall. N. Nord. Beitr. vii. 265 = A. Onobrychis.
hedysaroides, Willd. Sp. Pl. iii. 1264 = macrostachys.
heldaviensis, Lerch. ex Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 19 = A. Cicer.
Heldreichii, *Boiss. Diagn.* Ser. I. ix. 43.—As. Min.
hellenicus, *Boiss. Fl. Orient.* ii. 292.—Graecia.
Helleri, *Fenzl, in Bonplandia*, viii. (1860) 56.—Mexic.
Helmii, *Fisch. ex DC. Prod.* ii. 301.—Mont. Ural.
helminthocarpus, Vill. Prosp. 42 = depressus.
hemiphaca, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 328.—Sibir.
Hemsleyi, *Aitch. & Baker, ex Aitch. in Journ. Linn. Soc.* xix. (1882) 158.—Afghanist.
Hendersoni, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 120.—Tibet. occ.
heratensis, *Bunge, Astrag.* i. 83; ii. 145.—Afghanist.
hermoneus, *Boiss. Diagn.* Ser. I. ix. 94.—Syria.
heterodoxus, *Bunge, Astrag.* i. 53; ii. 88.—Persia.

ASTRAGALUS:—
Heydei, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 118.—Tibet. occ.
hians, Jacq. Coll. i. 87 = Oxytropis deflexa.
hilaris, *Bunge, Astrag.* i. 89; ii. 158.—Syria.
himalayanus, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 160.—Reg. Himal.
himalayensis, Hort. Par. ex Bunge, Astrag. ii. 32 = chlorostachys.
himalayensis, Jacquem. ex Heynh. Nom. ii. 52 = seq.
himalensis, Jacquem. ex Baker, in Hook. Fl. Brit. Ind. ii. 128 = chlorostachys.
Hinkei, Sadl. ex Bunge, Astrag. ii. 217 = argenteus.
hippocrepidis, Benth. ex Bunge, l. c. i. 5; ii. 3.—Afghanist.
hippoglossus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = hypoglottis.
hirsutissimus, *DC. Astrag.* 119. t. 19.—Syria.
hirsutus, *Vahl, Symb. Bot.* i. 59.—As. Min.
hirtellus, Ledeb. Fl. Ross. i. 606 = viciaefolius.
hirticalyx, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* ii. 380.—Kurdistan.
hirtulus, Ledeb. Fl. Ross. i. 606 = viciaefolius.
hirtus, *Bunge, Astrag.* i. 54; ii. 87.—Persia.
hispanicus, *Coss. ex Bunge, l. c.* i. 135; ii. 231.—Hispan.
hispidulus, *DC. Astrag.* 132. t. 13.—Arabia.
hispidulus, Michx. ex Steud. Nom. ed. I. 86 = gummifer.
hispidus, Labill. Ic. Pl. Syr. Dec. i. 18 = barbatus.
Hoantchy, *Franch. Pl. Davidian.* i. 86.—As. or.
Hohenackeri, *Boiss. Diagn.* Ser. I. ii. 70.—Reg. Caucas.
holargyreus, Bunge, in Mem. Sav. Etr. Pétersb. vii. (1851) 273 (Pl. Lehm. 97) = Schrenkianus.
hololeucus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 59.—Persia.
holopsilus, *Bunge, Astrag.* i. 56; ii. 93.—Persia.
holopterus, *Turcz. ex Bunge, l. c.* i. 98; ii. 170.—Sibiria.
holosemius, *Bunge, l. c.* i. 55; ii. 89.—Persia.
Hookerianus, D. Dietr. Syn. Pl. iv. 1086 = ervoides.
Hookerianus, *A. Gray, Proc. Am. Acad.* vi. (1864) 215.—Am. bor. occ.
horizontalis, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 406 = Arbuscula.
Hornii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 398.—Am. bor. occ.
horridus, *Boiss. Diagn.* Ser. I. ii. 66.—Persia.
horridus, Willd. ex Bunge, Astrag. ii. 135 = arnacantha.
Hosackiae, *Greene, in Bull. Calif. Acad.* i. n. 3 (1885) 157.—Calif.
hosackioides, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 123.—Reg. Himal.
hostilis, *Boiss. Fl. Orient.* ii. 306.—Beluchist.
Howelli, *A. Gray, in Proc. Am. Acad.* xv. (1880) 46.—Am. bor.
Huetii, *Bunge, Astrag.* i. 37; ii. 49.—Sicilia.
Humboldtii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 195.—Am. bor.
humifusus, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 119 = subumbellatus.
humifusus, Willd. Sp. Pl. iii. 1286 = viciaefolius.
humilis, *Bieb. Fl. Taur. Cauc.* ii. 203.—Reg. Caucas.
humilis, Geyer, ex Hook. Lond. Journ. Bot. vi. (1847) 211 = Shortianus.
humilis, Willd. ex Boiss. Diagn. Ser. I. ix. 83 = Lamarckii.
humillimus, *A. Gray, in Brandegee, Fl. S. W. Col.* 235.—Am. bor. occ.
humistratus, *A. Gray, Pl. Wright.* ii. 43.—Am. bor. occ.
hyalolepis, *Bunge, Astrag.* i. 105; ii. 185.—Reg. Caucas.
hybernus, Ehrenb. ex Bunge, l. c. ii. 81 = hirsutissimus.
hybridus, S. G. Gmel. It. i. t. 18 = flaccida.
hybridus, Willd. ex Ledeb. Fl. Ross. i. 628 = vesicarius.
hymenocalyx, *Boiss. Diagn.* Ser. I. ii. 55.—Persia.
hymenocarpus, Fisch. ex Bunge, Astrag. ii. 95 = chartaceus.
hymenochlaenus, Fisch. ex Bunge, l. c. i. 38; ii. 51 = angustiflorus.
hymenocystis, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 449.—Persia.
hymenostegis, *Fisch. & Mey. l. c.* 448.—Persia.

ASTRAGALUS:—
hypogaeus, *Ledeb. Fl. Ross. Alt.* t. 95; *Fl. Alt.* iii. 329.—Sibir. alt.
hypoglottis, Brot. Phytogr. t. 60 = pentaglottis.
hypoglottis, *Linn. Mant.* 274.—Reg. bor. temp.
hypoglottis, Desf. Fl. Atlant. ii. 183 = A. Glaux.
hypoglottis, DC. Astrag. 118 = danicus.
hypoglottoides, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 124.—Tibet. occ.
hypoleucus, *Schau. in Linnaea*, xx. (1847) 747.—Mexic.
hypoxylus, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 192.—Am. bor. occ.
hypsogeton, *Bunge, Astrag.* i. 82; ii. 140.—Persia.
hyrcanus, Kar. in Bull. Soc. Nat. Mosc. (1839) 152 = A. Ammodendron.
hyrcanus, *Pall. Astrag.* 25. t. 22.—Reg. Casp.
Hystrix, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 403.—Persia.
ibicinus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 293.—Persia.
ictericus, *Dingl. in Flora*, lxiv. (1881) 381.—Thracia.
idaeus, *Bunge, Astrag.* i. 107; ii. 187.—Ins. Creta.
iliensis, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 20.—As. temp.
illyricus, [Bernh.] Sel. Sem. Hort. Erf. (1836).—Cf. Linnaea, xii. (1838) Litt. 74 = Wulfenii.
immersus, *Baker, ex Aitch. in Journ. Linn. Soc.* xviii. (1881) 45.—Afghanist.
impressus, Ehrenb. ex Bunge, Astrag. ii. 19 = peregrinus.
inarius, Pall. Astrag. 94. t. 77. f. 2 = Oxytropis inaria.
incanus, Chaub. ex Bunge, Astrag. ii. 202 = Chaubarti.
incanus, Gilib. Fl. Lithuan. ii. 112 = arenarius.
incanus, *Linn. Syst.* ed. X. 1175.—Gall. austr.
incanus, Pall. Astrag. iii. t. 84 = brachycarpus.
incanus, Sibth. & Sm. Fl. Graec. t. 732 = Spruneri.
incanus, Siev. ex Ledeb. Fl. Ross. i. 592 = Oxytropis argentata.
incanus, Wulf. in Jacq. Coll. i. 320 = Wulfeni.
incertus, *Ledeb. Fl. Ross.* i. 647.—Reg. Caucas.; Armenia.
inconspicuus, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 126.—Reg. Himal.
incurvus, *Desf. Fl. Atlant.* ii. 182. t. 103.—Hispan.; Afr. bor. occ.
incurvus, Reichb. Fl. Germ. Excurs. 512 = Wulfeni.
inderiensis, *Claus, in Beitr. Pfl. Russ. Reich.* viii. 64.—Rossia.
indigoferoides, Horan. ex Bunge, Astrag. ii. 21 = capillipes.
ineptus, A. Gray, in Proc. Am. Acad. vi. (1864–65) 525 = lentiginosus.
infestus, *Boiss. Diagn.* Ser. I. ix. 95.—Afghanist.
infirmus, *Steud. Nom.* ed. II. i. 161.—Chili.
inflatus, *DC. Astrag.* 90. n. 13.—Desert. Kirgh.
inflatus, *Gill. ex Steud. Nom.* ed. II. i. 161.—Chili.
inflexus, *Dougl. ex G. Don, Gen. Syst.* ii. 256.—Am. bor. occ.
insularis, *Kellogg, in Bull. Calif. Acad.* n. i. (1884) 6.—Calif.
intarrensis, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 258.—Turkest.
intermedius, Boiss. Diagn. Ser. I. ii. 53 = commixtus.
intermedius, *Host, Fl. Austr.* ii. 361.—Europ.
intermedius, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 340.—Dahuria.
interruptus, Benth. ex Bunge, Astrag. ii. 10 = ammophilus.
iodanthus, *S. Wats. in Bot. King, Exp.* 70.—Am. bor. occ.
iodotropis, *Boiss. & Hohen. Diagn.* Ser. I. ix. 100.—Persia.
iranicus, Bunge, Astrag. i. 56; ii. 92 = mollis.
ischredensis, *Bunge, l. c.* i. 37; ii. 47.—Persia.
isetensis, Hort. ex Willd. Enum. Hort. Berol. 783 = falcatus, falciformis.
isopetalus, Boiss. Diagn. Ser. I. ix. 67 = kahiricus.
ispahanicus, *Boiss. l. c.* Ser. I. ii. 52.—Persia.
ispirensis, *Boiss. & Huet, l. c.* Ser. II. ii. 25.—Armenia.
Jacquemontii, Bunge, Astrag. i. 44; ii. 68 = polyacanthus.
janthinus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 328.—Persia.
Jaubertianus, *Boiss. Fl. Orient.* ii. 24.—Phrygia.
Jaubertianus, Bunge, Astrag. i. 90; ii. 160 = macroptilus.

ASTRAGALUS:—
jejunus, *S. Wats. in Bot. King, Exp.* 73, 442.—Am. bor. occ.
jeniseensis, Less. ex Bunge, Astrag. ii. 183 = semibiloculares.
jesdianus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 61.—Persia.
Jessenii, *Bunge, Astrag.* i. 61; ii. 101.—Persia.
jodostachys, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 60.—Persia.
Johannis, *Boiss. Diagn.* Ser. I. vi. 41.—Persia.
jubatus, *Boiss. l. c.* ix. 104.—Persia; Afghan.
junceus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 230.—Am. bor. occ.
junceus, Ledeb. in Spreng. Syst. iii. 297 = Stevenianus.
juvenalis, *Delile, Ind. Sem. Hort. Monsp.* (1836) 22.—Aegypt.
kadschorensis, *Bunge, Astrag.* i. 103; ii. 182.—Reg. Caucas.
kahiricus, *DC. Prod.* ii. 292.—Aegypt.; Arab.; Persia; Afghan.
kapherrianus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 446 = persicus.
karabaghensis, *Bunge, Astrag.* i. 93; ii. 165.—Persia.
karakugensis, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 232.—As. Min.
karamasicus, Boiss. & Bal. Diagn. Ser. II. vi. 50 = orthodontus.
Karelini, Fisch. in Bull. Soc. Nat. Mosc. (1839) 152 = commixtus.
karputanus, *Boiss. Diagn.* Ser. II. ii. 26.—Armenia.
karsianus, *Bunge, Astrag.* ii. 115.—Armenia.
kashmirensis, Bunge, l. c. ii. 246, Index = caschmirensis.
kellalensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 388.—Persia.
kentrodes, *Buhse, in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 412.—Persia.
kentrophyta, *A. Gray, in Proc. Acad. Philad.* (1863) 60.—Am. bor. occ.
keratensis, *Bunge, Astrag.* i. 73; ii. 122.—Persia.
Kessleri, *Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) I. 496.—Soongaria.
Keyserlingii, *Bunge, Astrag.* i. 82; ii. 139.—Persia.
khasianus, *Benth. ex Bunge, l. c.* i. 27; ii. 32.—Reg. Himal.
kirghisicus, Stschegl. in Bull. Soc. Nat. Mosc. xxvii. (1854) I. 161 = tribuloides.
kirrindicus, *Boiss. Diagn.* Ser. II. ii. 32.—Persia.
koburensis, *Bunge, Astrag.* i. 116; ii. 196.—Mongolia.
kohistanus, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 254.—Turkest.
kohrudicus, *Bunge, Astrag.* i. 67; ii. 109.—Persia.
kokandensis, *Bunge, Astrag. Fedsch.* 216.—As. temp.
kongrensis, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 120.—Him. bor. occ.
Korolkowii, *Bunge, Astrag. Fedsch.* 230.—As. temp.
koschukensis, *Boiss. Fl. Orient.* ii. 492.—Afghan.
Kotschyanus, *Boiss. Diagn.* Ser. I. ii. 44.—Syria; Mesopotam.
Kotschyanus, Fisch. ex Bunge, Astrag. ii. 19 = tuberculosus.
Krauscanus, *Regel, in Act. Hort. Petrop.* iii. (1875) 104.—Turkestan.
kuhistanus, *Bunge, Astrag.* i. 82; ii. 141.—Persia.
kuldshensis, *Bunge, in Act. Hort. Petrop.* vii. (1880) 374.—Turkestan.
kuramensis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 46.—Afghan.
kurdicus, *Boiss. Diagn.* Ser. I. ix. 84.—Kurdistan; Syria.
kurrumensis, *Bunge, in Act. Hort. Petrop.* vii. (1880) 370.—Afghan.
kurtschumensis, *Bunge, Astrag.* i. 139; ii. 236.—Soongaria.
labradoricus, DC. Prod. ii. 287 = alpinus.
lacteus, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. ii. 31.—Graecia.
lactiflorus, Ledeb. Fl. Alt. iii. 333 = testiculatus.
lactus, *Bunge, Astrag.* i. 34; ii. 41.—Afghan.
lagocephalus, *Fisch. & Mey. ex Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 197.—As. temp.
lagonyx, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 364.—Persia.
lagopodioides, Vahl, Symb. Bot. i. 64 = lagopoides.
lagopoides, Bunge, Astrag. i. 67; ii. 111 = uraniolimneus.
lagopoides, *Lam. Encyc.* i. 322.—Armenia.
Lagowskii, *Trautv. in Bull. Phys.-Math. Acad. Pétersb.* xvi. (1858) 324.—Armenia.
laguroides, *Pall. Reise*, iii. 750.—Sibir.
Lagurus, Eichw. Casp. Cauc. 631 = persicus.
Lagurus, Pall. Astrag. 18. t. 16 = laguroides.
Lagurus, *Willd. in Mém. Acad. Berl.* (1794–95) 28.—Reg. Caucas.; Persia.
Lamarckii, *Boiss. Diagn.* Ser. I. ix. 82.—As. Min.
Lambertii, Spreng. Syst. iii. 308 = Oxytropis Lambertii.
lanatus, *Labill. Ic. Pl. Rar. Syr.* i. 21. t. 11.—Syria.
lanatus, Pall. Astrag. 108. t. 81 = Oxytropis Pallasii.
lancearius, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 370.—Am. bor.
lanceolatus, *Bunge, Astrag.* i. 35; ii. 60.—Afghan.
lanceolatus, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 246 = stipulatus.
lanigerus, *Desf. Fl. Atlant.* ii. 181. t. 202.—Afr. bor.
lanigerus, Viv. Fl. Lib. Spec. 44 = alexandrinus.
lanuginosus, *Clos, in C. Gay, Fl. Chil.* ii. 121.—Chili.
lanuginosus, Desc. Fl. Pitt. Antill. Ind. viii. 53 = Sophora tomentosa.
lanuginosus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 409.—As. temp.
laricus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 61.—Persia.
lasianthus, C. A. Mey. in Bong. & Mey. Suppl. Fl. Alt. n. 92 = lasiopetalus.
lasiocaulos, Bunge, Astrag. i. 83; ii. 144 = pachyacanthus.
lasiocladus, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 135 = bicuspis.
lasioglottis, *Bieb. Fl. Taur. Cauc.* iii. 486.—Reg. Caucas.
lasiopetalus, *Bunge, in Del. Sem. Dorp.* (1839).—*Cf. Linnaea* xiv. (1840) *Litt.* 117.—Soongaria.
lasiophyllus, *Ledeb. Fl. Ross.* i. 627.—Oriens.
lasiosemius, *Boiss. Diagn.* Ser. I. ix. 96.—Afghan.
lasiostylus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 367.—Turkestan.
latebracteatus, Boiss. in Tchihat. Asie Min. Bot. i. 54. = macrochlamys.
lateritius, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 339.—Kurdistan.
latifolius, DC. Astrag. 235 = Candolleanus.
latifolius, *Lam. Encyc.* i. 319.—Armen.; Persia.
laxiflorus, Boiss. Fl. Orient. ii. 413 = bracteosus.
laxiflorus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 390 = psilodontius.
Laxmanni, Jacq. Hort. Vind. iii. 22. t. 37 = adsurgens.
Laxmanni, Nutt. Gen. Am. ii. 99 = striatus.
laxus, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 269 = remotifolius.
Layneae, *Greene, in Bull. Calif. Acad.* i. n. 3. (1885) 156.—Calif.
Lehmannianus, *Bunge, in Arb. Nat. Ver. Riga*, i. (1847) 252.—Turkestan.
leiocalyx, Benth. ex Bunge, Astrag. i. 36; ii. 49 = laetus.
leiocarpus, Shuttlew. in Mag. Zool. & Bot. ii. (1838) 520 = exscapus.
leioclados, *Boiss. Diagn.* Ser. I. ii. 62.—Persia.
leiolobus, *Bunge, Astrag.* i. 14; ii. 15.—Persia.
leiophysa, *Bunge, in Act. Hort. Petrop.* iii. (1875) 104.—Turkestan.
Lemmoni, *A. Gray, in Proc. Am. Acad.* viii. (1873) 626.—Am. bor. occ.
lentiformis, *A. Gray, in Bot. Calif.* i. 156.—Calif.
lentiginosus, *Dougl. ex Hook. Fl. Bor. Am.* i. 151.—Am. bor. occ.
leontinus, Wahlenb. Fl. Lapp. 191. t. 12. f. 4 = oroboides.
leontinus, *Wulf. in Jacq. Misc.* ii. 59.—Austr.; Helv.
leontonyx, Bunge, Astrag. i. 89; ii. 157 = oleaefolius.
lepidanthus, *Boiss. Diagn.* Ser. I. ix. 85.—Syria.
leporinus, *Boiss. l. c.* Ser. II. ii. 30.—Armenia.
lepsensis, *Bunge, Astrag.* i. 25; ii. 29.—Soongaria.
leptacanthus, Benth. ex Bunge, l. c. i. 44; ii. 69 = leptocentrus.
leptacanthus, *Boiss. & Buhse, in Bull. Soc. Nat. Mosc.* (1853) II. 432.—Persia.
leptaleus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 220.—Am. bor. occ.
leptocarpus, *Torr. & Gray, Fl. N. Am.* i. 334.—Texas.
leptocaulis, *Ledeb. Fl. Alt.* iii. 296.—Sibir. altaic.
leptocentrus, *Bunge, Astrag.* i. 44; ii. 69.—Reg. Himal.
leptodendron, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 345 = caspius.
leptodermus, *Bunge, Astrag. Fedsch.* 210.—As. centr.
leptophyllus, *Desf. Fl. Atlant.* ii. 188. t. 207.—Algeria.
leptophyllus, Nutt. in Journ. Acad. Philad. vii. (1834) 18 = stenophyllus.
leptophyllus, Pall. Reise, iii. 749 = Oxytropis leptophylla.
leptopus, *Benth. ex A. Gray, Pl. Wright.* ii. 43.—Mexic.
leptostachys, Pall. Astrag. 50. t. 40 = sulcatus.
leptostachys, Siev. in Willd. Sp. Pl. iii. 1265 = puberulus.
leptothalamus, Boiss. ex Bunge, Astrag. i. 135; ii. 232 = seq.
leptothamnus, *Boiss. Fl. Orient.* ii. 480.—Cilicia.
leptus, *Boiss. l. c.* 305.—Afghan.
lessertioides, *Benth. ex Bunge, Astrag.* i. 24; ii. 28.—Reg. Himal.
leucacanthus, *Boiss. Diagn.* Ser. I. ix. 93.—Aegypt.
leucanthus, Pall. Astrag. 59. t. 47 = Oxytropis leucantha.
leucocephalus, *R. Grah. in Wall. Cat.* n. 5923.—Reg. Himal.
leucocladus, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) 27.—As. temp.
leucocyaneus, *Griseb. Spicil. Fl. Rumel.* i. 50.—As. Min.
leucolachnus, *Boiss. Fl. Orient.* ii. 357.—Syria.
leucomallus, *Phil. in Linnaea*, xxx. (1859–60) 186.—Am. austr.
leucomelas, *Bunge, Astrag.* i. 32; ii. 38.—Afghan.
leucophaeus, *Sm. in Trans. Linn. Soc.* i. (1791) 252.—Sicilia.
leucophyllus, Hook. Lond. Journ. Bot. vi. (1847) 211 = arrestus.
leucophyllus, *Torr. & Gray, Fl. N. Am.* i. 336.—Am. bor. occ.
leucophyllus, Willd. Sp. Pl. iii. 1331 = angustifolius.
leucopsis, *Torr. Bot. Mex. Bound.* 56. t. 16.—Calif.
leucoptilus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 320.—Kurdistan.
leucospermus, Bunge, in Arb. Nat. Ver. Riga, i. (1847) 250 = A. Alopecias.
libanoticus, Boiss. Diagn. Ser. II. v. 87 = A. Echinus.
Libanotis, Willd. Sp. Pl. iii. 1308 = lanatus.
lignosus, *DC. Astrag.* 175. t. 39.—Sibiria.
lilacinus, *Boiss. Diagn.* Ser. I. ix. 42.—Persia.
Lindheimeri, *Engelm. ex A. Gray, Pl. Wright.* i. 52.—Am. bor. occ.
linearifolius, Pers. Syn. ii. 336 = A. Onobrychis.
linearius, Willd. Sp. Pl. iii. 1302 = Oxytropis inaria.
lineatus, *Lam. Encyc.* i. 314.—Armenia.
Listoniae, *Boiss. Diagn.* Ser. I. ix. 37.—Anatolia.
lithophilus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 344.—Soongar.
litoreus, *Phil. in Linnaea*, xxxiii. (1864–65) 47.—Chili.
litostachyus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 350.—Oriens; Kurdistan.
lividus, Monb. ex Bunge, Astrag. ii. 19 = tuberculosus.
lobophorus, *Boiss. Diagn.* Ser. II. ii. 31.—Persia.
lonchocarpus, *Torr. in Pacif. Rail. Rep.* iv. 80.—Am. bor. occ.
longebracteatus, Steud. Nom. ed. II. i. 162 = sanguinolentus.
longicaulis, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 129.—Kashmir.
longicaulis, *Pomel, Nouv. Mat. Fl. Atl.* 185.—Algeria.
longicuspidatus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = longicuspis.
longiscuspis, *Bunge, Astrag.* i. 121; ii. 211.—Reg. Caucas.
longiflorus, Delile, Fl. Egypte, 256. t. 39 = kahiricus.
longiflorus, Gilib. Fl. Lithuan. ii. 111 = A. Onobrychis.
longiflorus, *Pall. Astrag.* 73. t. 60.—Reg. Casp.
longifolius, *Lam. Encyc.* i. 322.—Armenia.
longipes, Boiss. & Heldr. ex Boiss. Fl. Orient. ii. 480 = Apollineus.

ASTRAGALUS:—
longipes, Steud. Nom. ed. II. i. 162 = rupifragus.
longipes, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 405 = macropterus.
longirostrus, Poir. Encyc. Suppl. i. 528 = Oxytropis longirostra.
longistylus, *Bunge, Astrag.* i. 81 ; ii. 137.—Persia.
lotiflorus, *Hook. Fl. Bor. Am.* i. 152.—Am. bor. occ.
lotoides, Lam. Encyc. i. 316 = sinicus.
lunatus, Gilib. Fl. Lithuan. ii. 110 = glycyphyllos.
lunatus, *Pall. Astrag.* 44.—Persia.
lupulinus, *Pall. Reise*, iii. 750.—Sibiria.
lusitanicus, *Lam. Encyc.* i. 312.—Reg. Mediterr. ; As. Min.
luteo-caeruleus, *Baker, in Journ. Linn. Soc.* xviii. (1878) 47.—Afghanist.
luxurians, *Bunge, Astrag.* i. 23 ; ii. 25.—Sibir.
Lyalli, *A. Gray, in Proc. Am. Acad.* vi. (1864) 195.—Am. bor. occ.
lycioides, *Boiss. Diagn.* Ser. I. ii. 66.—Persia.
lycius, *Boiss. l. c.* ix. 45.—Lycia.
lydius, Boiss. l. c. ii. 43 = leucocyaneus.
lydius, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 393 = amphilogus.
macranthus, Willd. ex Ledeb. Fl. Ross. i. 650 = brachycarpus.
macrobotrys, *Bunge, Astrag.* i. 129 ; ii. 226.—Persia.
macrocarpus, *DC. Astrag.* 143. t. 28.—Syria.
macrocarpus, Pall. in Nov. Act. Petrop. x. (1776) 56 = subulatus.
macrocephalus, *Willd. Sp. Pl.* iii. 1260.—As. Min. ; Syria.
macroceras, C. A. Mey. ex Bong. Verz. Pfl. Saisang-Nor, 24 = macrolobus.
macrochlamys, *Boiss. ex Bunge, Astrag.* i. 49 ; ii. 77.—As. Min.
macrocladus, *Bunge, Astrag. Fedsch.* 293.—As. centr.
macrodon, *A. Gray, in Proc. Am. Acad.* vi. (1864) 216.—Am. bor. occ.
macrolacis, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 63 = stenolepis.
macrolobus, *Bieb. Fl. Taur. Cauc.* iii. 493.—Sibir. altaic.
macrolobus, Bunge, Enum. Alt. 70 = macroceras.
macrolobus, Ledeb. Fl. Alt. iii. 307 = eriolobus, chaetolobus.
macronyx, *Bunge, Astrag. Fedsch.* 236.—As. centr.
macropelmatus, *Bunge, Astrag.* i. 36 ; ii. 43.—Persia.
macropetalus, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1844) 307.—As. temp.
macropodius, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 352 = aureus.
macropterus, *DC. Prod.* ii. 283.—Tibet. occ. ; Sibir.
macroptilus, *Boiss. ex Bunge, Astrag.* i. 90 ; ii. 160.—Phrygia.
macropus, Boiss. Diagn. Ser. II. vi. 51 = macroscepus.
macropus, *Bunge, in Arb. Nat. Ver. Riga*, i. (1847) 238.—As. med.
macrorrhizus, *Cav. Ic.* ii. 28. t. 133.—Hispania.
macrorrhizus, D. Don, Prod. Fl. Nep. 245 = Donianus.
macroscepus, *Boiss. in Bunge, Astrag.* i. 49 ; ii. 77.—Cappadocia.
macrosemius, *Boiss. & Hohen. Diagn.* Ser. I. ix. 93.—Persia.
macrostachys, *DC. Astrag.* 141. t. 15.—Armenia.
macrotropis, *Bunge, Astrag.* i. 127 ; ii. 223.—Soongaria.
macrourus, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1838) 346.—Reg. Caucas. ; Persia.
maculatus, *Bunge, Astrag.* i. 119 ; ii. 206.—Persia.
maculatus, Lam. Encyc. i. 316 = annularis.
Maddenianus, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 127.—Reg. Himal.
majellensis, Huet, ex Nym. Consp. 193 = australis.
malacensis, Salzm. ex Bunge, Astrag. ii. 12 = sesameus.
malacophyllus, *Benth. ex Bunge, l. c.* i. 36 ; ii. 61.—Reg. Himal.
malacophyllus, Hort. ex G. Don, in Loud. Hort. Brit. 307 = galegiformis.
malacotrichus, Fisch. & Kotschy, ex Bunge, Astrag. ii. 164 = Roussaeanus.
malacus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 336.—Am. bor. occ.
mareoticus, *Delile, Fl. Egypt.* 257. t. 39. f. 3.—Aegypt.
maritimus, *Moris, Fl. Sard.* i. 523.—Sardinia.
maroniensis, *Dingl. in Flora*, lxiv. (1881) 382.—Thracia.

ASTRAGALUS:—
maroniticus, Boiss. & Balansa, Diagn. Ser. II. v. 58 = drusorum.
marosportensis, Lerch. ex Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 19 = dasyanthus.
Marschallianus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 328.—Oriens ; Reg. Caucas.
marusiensis, Lerch. ex Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 19 = arenarius.
masanderanus, *Bunge, Astrag.* i. 114 ; ii. 193.—Persia.
massiliensis, *Lam. Encyc.* i. 320.—Gallia.
Matthewsii, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 192.—Am. bor.
mauritanicus, *Coss. in Bull. Soc. Bot. Fr.* iii. (1856) 673.—Algeria.
mauritanicus, *Stev. in Bull. Soc. Nat. Mosc.* iv. (1832) 267.—Algeria.
Maximowiczii, *Trautv. in Act. Hort. Petrop.* ix. (1884) 379.—Rossia.
maximus, Lam. ex Steud. Nom. ed. II. i. 162 = A. Onobrychis.
maximus, *Willd. Sp. Pl.* iii. 1258.—Reg. Caucas. ; Armenia.
Maxwellii, Royle, Illustr. Bot. Himal. 199 = tibetanus ?
medicagineus, Boiss. Diagn. Ser. I. ix. 42 = Kotschyanus.
medius, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1843) 196.—Soongaria.
medullaris, Boiss. Diagn. Ser. I. ix. 66 = graveolens.
megacarpus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 215.—Am. bor. occ.
megalanthus, *DC. Astrag.* 131. t. 20.—Sibir.
megalocystis, *Bunge, l. c.* ii. 122.—Persia.
megalomerus, *Bunge, in Act. Hort. Petrop.* vii. (1880) 379.—As. centr.
megalotropis, *C. A. Mey. ex Bunge, Astrag.* i. 60 ; ii. 99.—Reg. Caucas.
melaleucus, *Bunge, Astrag.* ii. 100.—Persia.
melanocalyx, *Boiss. in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 59.—Persia.
melanocarpus, *Bunge, Astrag.* i. 51 ; ii. 81.—Oriens.
melanocarpus, *Hook. Fl. Bor. Am.* i. 150.—Am. bor.
melanocarpus, Richards. App. Frankl. Journ. ed. II. 87 = missouriensis.
melanocephalus, *Boiss. Diagn.* Ser. I. ix. 50.—As. Min.
melanodon, *Boiss. l. c.* Ser. I. vi. 36.—Persia.
melanophrurius, *Boiss. Fl. Orient.* ii. 276.—Cappadoc.
melanostachys, *Benth. ex Bunge, Astrag.* i. 21 ; ii. 22.—Reg. Himal.
melanotrichus, Boiss. ex Bunge, l. c. ii. 188 = cataonicus.
melanotrichus, *Ledeb. Fl. Alt.* iii. 323. — Sibir. altaic.
melilotoides, *Pall. Voy.* t. 33 ; *Desc.* iii. 748.—Sibir. ; China.
melinus, *Boiss. Fl. Orient.* ii. 446.—Cappadocia.
melitenensis, *Boiss. Diagn.* Ser. I. ii. 43.—Cappadocia.
membranaceus, *Bunge, Astrag.* i. 25 ; ii. 30.—As. temp.
membranaceus, Moench, Meth. 167 = trimestris.
Menziesii, *A. Gray, in Proc. Am. Acad.* vi. (1864) 217.—Calif.
Mercklinii, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 70.—Persia.
meridionalis, *Bunge, Astrag.* i. 61 ; ii. 104.—Persia.
meschedensis, *Bunge, l. c.* i. 82 ; ii. 140.—Persia.
mesites, *Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 66.—Armenia.
mesogitanus, *Boiss. Diagn.* Ser. I. ii. 41.—As. Min.
mesoleios, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 63 = radkanensis.
mesoleios, *Boiss. & Hohen. Diagn.* Ser. I. ix. 91.—Persia.
mesopotamicus, *Boiss. l. c.* ii. 68.—Mesopotam.
mesopterus, *Griseb. Spicil. Fl. Rumel.* i. 49.—Macedon.
mexicanus, A. Gray, Pl. Lindh. 176 = plattensis.
mexicanus, *A. DC. Pl. Nov. Hort. Genèv.* v. 17.—Am. bor.
Meyeri, *Boiss. Diagn.* Ser. I. ix. 88.—Armenia.
Michauxianus, *Boiss. l. c.* ii. 62.—Kurdistan.
micracme, *Boiss. & Noë, l. c.* Ser. II. v. 87.—Armenia.
micrancistrus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 475.—Oriens ; Assyria.
micranthellus, *Wedd. Chlor. And.* i. 262.—Peruv.
micranthus, *Desv. Journ. Bot.* iii. (1814) 78.—Hab. ?

ASTRAGALUS:—
micranthus, Nutt. in Journ. Acad. Philad. ii. (1821) 122 = Nuttallianus.
micranthus, Pall. ex Ledeb. Fl. Ross. i. 637 = striatellus.
microcarpus, *DC. Astrag.* 104.—Sibir.
microcephalus, *Willd. Sp. Pl.* iii. 1332.—As. Min.
microcystis, *Bunge, Astrag. Fedsch.* 306.—As. centr.
microcystis, *A. Gray, in Proc. Am. Acad.* vi. (1864) 220.—Am. bor. occ.
microdontus, *Baker, in Journ. Linn. Soc.* xviii. (1881) 46.—Afghan.
microlobus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 203.—Am. bor.
microphyllus, Bess. ex Steud. Nom. ed. II. i. 162 = A. Onobrychis.
microphyllus, Georgi, Beschr. Russ. Reich. Nachtr. 296 = adsurgens.
microphyllus, *Linn. Sp. Pl.* 757.—Sibir.
microphyllus, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 134 = cicerifolius.
microphyllus, Pall. Astrag. 92. t. 76 = Oxytropis microphylla.
microphyllus, Schuebl. & Martens, Fl. Wuertemb. 474 = A. Cicer.
microphysa, *Boiss. Diagn.* Ser. I. ii. 71.—Persia.
micropterus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 396.—Anatolia.
microrchis, *Barbey, in Bull. Soc. Vaud. Sc. Nat.* xxi. (1883) 95.—Helvetia.
microthamnus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 393.—Assyria.
miniatus, *Bunge, Astrag.* i. 98 ; ii. 169.—Mongolia.
minimus, DC. Astrag. 64 = alpinus.
minimus, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 18.—Peruv.
minor, *Clos, in C. Gay, Fl. Chil.* ii. 107.—Chili.
minutissimus, *Wedd. Chlor. And.* ii. 257.—Peruv.
minutus, Boiss. Diagn. Ser. I. ix. 58 = tribuloides.
minutus, Buhse, ex Bunge, in Mém. Acad. Pétersb. Sér. VII. xv. (1869) n. 1. 11 = biovulatus.
mirpoureanus, Camb. in Jacquem. Voy. Bot. 39. t. 46 = Gueldenstaedtia mirpourensis.
miser, *Dougl. in Hook. Fl. Bor. Am.* i. 152.—Am. bor. occ.
missouriensis, *Nutt. Gen. Am.* ii. 99.—Am. bor.
Mitchellianus, *Boiss. Diagn.* Ser. I. ix. 80.—Phrygia.
mixtus, Bieb. Cent. Pl. Rar. i. t. 26 = vesicarius.
modestus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 75.—Persia.
modestus, *Wedd. Chlor. And.* i. 263.—Am. austr.
Moellendorffii, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 472.—China bor.
mogollonicus, *Greene, in Torrey Bot. Club*, viii. (1881) 97.—Calif.
mohavensis, *S. Wats. in Proc. Am. Acad.* xx. (1885) 361.—Am. bor. occ.
mokiacensis, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 367.—Am. bor. occ.
moldavicus, Steud. Nom. ed. I. 87 = A. Onobrychis.
mollis, *Bieb. Fl. Taur. Cauc.* iii. 495.—Reg. Caucas. Persia.
mollissimus, Presl, Bot. Bemerk. 57 = bethlemiticus.
mollissimus, *Torr. in Ann. Lyc. N. York*, ii. (1828) 178.—Am. bor. occ.
monadelphus, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 32.—China.
monanthemus, *Boiss. Diagn.* Ser. I. ii. 78.—Persia.
mongholicus, *Bunge, Astrag.* i. 25 ; ii. 30.—Mongolia.
monophyllus, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 473.—Mongolia.
monspessulanus, Griseb. Spicil. Fl. Rumel. i. 60 = Wulfeni.
monspessulanus, *Linn. Sp. Pl.* 761.—Europ. austr. ; Reg. Cauc.
montanus, Bieb. Fl. Taur. Cauc. ii. 197 = Oxytropis cyanea.
montanus, Lapeyr. Hist. Abr. Pl. Pyr. 429 = Oxytropis pyrenaica.
montanus, Linn. Sp. Pl. 760 = Oxytropis montana.
montanus, Pall. Reise, iii. 199 = Oxytropis uralensis.
montanus, Brot. ex Steud. Nom. ed. I. 87 = macrorrhizus.
montanus, Jacq. Fl. Austr. 131 = alpinus.
monticola, *Phil. in Linnaea*, xxxiii. (1864–65) 47.—Chili.

ASTRAGALUS:—
Moorcroftianus, R. Grah. in Wall. Cat. n. 5933 = Sophora Moorcroftiana.
Mortoni, *Nutt. in Journ. Acad. Phil.* vii. (1834) 19. —Am. bor. occ.
mossulensis, *Bunge, Astrag.* i. 101; ii. 177.—Persia; Mesopotam.
mucidus, *Bunge, ex Boiss. Fl. Orient.* ii. 279.—Turkestan.
muciferus, *Bunge, Astrag. Fedsch.* 223.—As centr.
mucronatus, *DC. Astrag.* 140. t. 25.—Armenia.
mucronifolius, *Boiss. Diagn.* Scr. I. ii. 68.—Persia.
Muelleri, Steud. & Hochst. in Flora, x. (1827) 72 = A. Onobrychis.
mugodsharicus, Bunge, Astrag. i. 124; ii. 216 = mugosaricus.
mugosaricus, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 236.—Transural.
multicaulis, *Ledeb. Fl. Alt.* iii. 295.—Sibir. altaic. et baical.
multicaulis, Nutt. ex Torr. & Gray, Fl. N. Am. i. 335 = pubentissimus.
multicaulis, Turcz. in Bull. Soc. Nat. Mosc. (1838) 90 = bifidus.
multiceps, *Wall. Cat.* n. 5237; *Bunge, Astrag.* ii. 69. —Reg. Himal.
multiflorus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 226.—Am. bor. occ.
multijugus, *DC. Prod.* ii. 301.—Mesopotam.
Munroi, *Benth. ex Bunge, Astrag.* i. 30; ii. 35.—Reg. Himal.; Kashgar.
muradicus, *Bunge, l. c.* i. 90; ii. 161.—Phrygia.
muricatus, Pall. Astrag. 89. t. 73 = Oxytropis muricata.
murinus, *Boiss. Diagn.* Scr. I. ii. 72.—Persia.
muschianus, *Kotschy & Boiss. ex Boiss. Fl. Orient.* ii. 357.—Armenia.
mutabilis, Boiss. Fl. Orient. ii. 448 = paecilanthus.
mutabilis, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 125 = strictus.
myriacanthus, *Boiss. Diagn.* Ser. I. ix. 91.—Persia.
myriophyllus, *Bunge, Astrag. Fedsch.* 232.—As. centr.
myriophyllus, Pall. Astrag. 87. t. 71 = Oxytropis myriophylla.
nanodes, *Bunge, in Act. Hort. Petrop.* vii. (1880) 377. —Turkest.
nanus, *DC. Astrag.* 114. t. 17.—Cilicia; Syria.
narbonensis, *Gouan, Illustr.* 49.—Gallia; Hispan.
narbonensis, Pall. Astrag. 13. t. 10 = ponticus.
narbonensis, C. A. Mey. Verz. Pfl. Cauc. 143 = megalotropis.
nebrodensis, Presl, ex Steud. Nom. ed. II. i. 162 = Boissieri.
neglectus, *Fisch. ex Steud. l. c.*—Sibir.
nematodes, *Bunge, ex Boiss. Fl. Orient.* ii. 425.—Turkestan.
nervistipulus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 384.—Kurdistan.
neurocarpus, *Boiss. Diagn.* Ser. I. ii. 59.—Syria.
neurophyllus, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 260.—Turkest.
nevadensis, *Boiss. Diagn.* Scr. I. ii. 63.—Hispan.
Newberryi, *A. Gray, in Proc. Am. Acad.* xii. (1877) 55.—Utah.
nicharensis, *Bunge, Astrag.* i. 39; ii. 50.—Beluchist.
nigrescens, A. Gray, in Am. Journ. Sc. Ser. II. xxxiii. (1862) 410 = multiflorus.
nigrescens, Nutt. in Journ. Acad. Philad. Ser. II. i. (1847) 153 = didymocarpus.
nigrescens, Pall. Astrag. 65. t. 53 = Oxytropis nigrescens.
nigrostriatus, C. Koch, in Linnaea, xv. (1841) 719 = ornithopodioides.
nitens, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 61 = viridis.
nitens, *Boiss. & Heldr. Diagn.* Ser. I. ix. 51.—As. Min.
nitens, *Host, Fl. Austr.* ii. 362.—Europ.
nitidus, Dougl. ex Hook. Fl. Bor. Am. i. 149 = adsurgens.
nivalis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 341.—Tibet. occ.; Soongaria.
niveus, Coq. Monbr. ex Bunge, Astrag. ii. 189 = pelliger.
Noaëanus, *Boiss. Diagn.* Ser. II. ii. 24.—Armenia.
nodiflorus, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) II. 147.—Rossia.
Nortonii, *Nutt. in Journ. Acad. Phil.* vii. (1834) 19. —Am. bor.
norvegicus, *Grauer, Dec.* 13.—Norvegia.

ASTRAGALUS:—
nothoxys, *A. Gray, in Proc. Am. Acad.* vi. (1864) 232. —Am. bor. occ.
nubigenus, D. Don, Prod. Fl. Nep. 245 = Caragana crassicaulis.
nuciferus, *Bunge, Astrag. Fedsch.* 223.—Cappadoc.; Mesopotam.
nucleiferus, *Boiss. Diagn.* Ser. I. ii. 83.—Oricns.
nudatus, Bunge, Astrag. i. 85; ii. 147 = kurdicus.
nudus, *Clos, in C. Gay, Fl. Chil.* ii. 115 (1846).—Chili.
nudus, *S. Wats. in Bot. King, Exp.* 74, 443.—Am. bor. occ.
nummularioides, *Desf. Fl. Atlant. Emend. ex DC. Astrag.* 233.—Numidia.
nummularius, *DC. Astrag.* 221. t. 2. f. 2.—Ins. Creta.
nummularius, Desf. Fl. Atlant. ii. 182 = nummularioides.
nummularius, Lam. Encyc. i. 317 = pinctorum.
nummularius, C. A. Mey. ex Bunge, Astrag. ii. 57 = talyschensis.
nurensis, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 58.—Persia.
Nuttallianus, *DC. Prod.* ii. 289.—Am. bor. occ.
obcordatus, Boiss. Diagn. Ser. I. vi. 38 = anacardius.
obcordatus, *Ell. Sketch*, ii. 227.—Am. bor.
oblongifolius, *Clos, in C. Gay, Fl. Chil.* ii. 109.—Chili.
obscurus, *S. Wats. in Bot. King, Exp.* 69, 437.—Am. bor. occ.
obtusifolius, *DC. Prod.* ii. 295.—Assyria.
obvallatus, Ehrenb. ex Bunge, Astrag. ii. 149 = cruentiflorus.
obvelatus, Ehrenb. ex Bunge, l. c. ii. 163 = drusorum.
ochreatus, *Bunge, l. c.* i. 127; ii. 224.—Persia.
Ochrias, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 33.—Mongolia.
ochrobius, *Bunge, Astrag.* i. 82; ii. 141.—Persia.
ochroleucus, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 57.—Marocco.
ochroleucus, Gilib. Fl. Lituan. ii. 111 = pilosus.
ocrochlorus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 83.—Persia.
odessanus, Bess. Enum. Pl. Volh. 30 = fruticosus.
odoratus, D. Dietr. Syn. Pl. iv. 1080 in syn. = obcordatus.
odoratus, *Lam. Encyc.* i. 311.—Reg. Caucas.; Persia; Soongaria.
oleifolius, *DC. Astrag.* 192.—As. Min.
oleifolius, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) 362 = aeluropus.
Olgae, *Bunge, Astrag. Fedsch.* 223.—As. centr.
Olgini, Bunge, Astrag. i. 130; ii. 177 = mossulensis.
oliganthus, Horan. ex Bunge, l. c. ii. 1 = complanatus.
oliganthus, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 337 = reticulatus.
oligophyllus, *Boiss. Diagn.* Ser. I. ii. 47.—Persia.
oligophyllus, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 197 = unijugus.
Olivieri, Klotzsch, ex Bunge, Astrag. i. 69; ii. 115 = mesopotamicus.
olopterus, DC. Prod. ii. 284 = austriacus.
olopterus, Turcz. ex Steud. Nom. ed. II. i. 162, 164 = versicolor.
olympicus, Bunge, Astrag. ii. 162 = baibutensis.
olympicus, Lam. ex Walp. Ann. iv. 514 = angustifolius.
olympicus, Pall. Astrag. 4. t. 4 = angustifolius.
oncotrichus, *Bunge, Astrag.* ii. 11.—Persia.
onobrychioides, Bieb. Fl. Taur. Cauc. iii. 486 = A. Cephalotes.
onobrychioides, Bunge, Astrag. ii. 183 = Bungeanus.
Onobrychis, *Linn. Sp. Pl.* 760.—Europ. austr.; Oriens; Sibir.
Onobrychis, Poll. Hist. Pl. Palat. ii. 327 = hypoglottis.
oocarpus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 213. —Am. bor. occ.
ooccphalus, *Boiss. Diagn.* Ser. I. ii. 56.—Syria.
oophorus, *S. Wats. in Bot. King, Exp.* 73, 441.—Am. bor. occ.
ophiocarpus, *Benth. ex Boiss. Fl. Orient.* ii. 224.—Persia; Tibet. occ.
oplites, Benth. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 134 = cicerifolius.
orbicularis, Walp. Rep. i. 702 = seq.
orbiculatus, *Ledeb. Fl. Ross. Alt.* t. 290; *Fl. Alt.* iii. 311.—Afghan.; Turkest.; Sibir. altaic.
Orbignyanus, *Wedd. Chlor. And.* ii. 260.—Bolivia.
Orcuttianus, *S. Wats. in Proc. Am. Acad.* xx. (1885) 361.—Calif.

ASTRAGALUS:—
oreades, *C. A. Mey. Verz. Pfl. Cauc.* 141.—Reg. Caucas.
oreganus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 335. —Am. bor. occ.
orientalis, Pers. ex DC. Prod. ii. 295 = macrocephalus.
ornithopodioides, *Lam. Encyc.* i. 312.—As. Min.; Persia.
oroboides, *Hornem. in Fl. Dan.* t. 1396.—Am. bor.
orthanthus, *A. Gray, in Proc. Am. Acad.* vi. (1864) 195.—Mexic.
orthocarpus, *Boiss. Diagn.* Ser. I. ix. 63.—Afghan.
orthocarpus, Dougl. ex A. Gray, in Proc. Am. Acad. vi. (1864) 196 = canadensis.
orthodontus, Bunge, Astrag. i. 101; ii. 176 = karamasicus.
ortholobus, *Bunge, l. c.* i. 127; ii. 223.—Mont. Ural.
otiporensis, *Boiss. Fl. Orient.* ii. 271.—Afghan.
ourmitanensis, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 256.—Turkest.
ovalis, *Boiss. & Bal. Diagn.* Ser. II. vi. 52.—Cappadoc.
ovallensis, *Clos, in C. Gay, Fl. Chil.* ii. 116.—Chili.
ovatus, *DC. Astrag.* 212. n. 107.—Armenia.
ovigerus, *Boiss. Diagn.* Ser. I. ii. 67.—Persia.
ovinus, *Boiss. l. c.* 76.—Armenia.
Owerini, *Bunge, Astrag.* i. 105; ii. 186.—Reg. Caucas.
oxyglottis, *Stev. in Bieb. Fl. Taur. Cauc.* ii. 192.—Reg. Caucas.; Persia; Turcoman.
oxyodon, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 125.—Tibet. occ.
oxypetalus, *Bunge, Astrag. Fedsch.* 273.—As. centr.
oxyphyllus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 375.—Syria.
oxyphyllus, Pall. Astrag. 90. t. 74 = Oxytropis oxyphylla.
oxyphysus, *A. Gray, in Proc. Calif. Acad.* iii. (1863) 103.—Am. bor. occ.
oxyrhynchus, *Hemsl. Biol. Centr. Am. Bot.* i. 265.—Mexic.
oxysemius, *Bunge, Astrag.* i. 118; ii. 202.—Anatolia.
oxytropifolius, *Boiss. Diagn.* Ser. I. ix. 37.—As. Min.
oxytropoides, Bunge, in Mém. Acad. Pétersb. Sér. VI. vii. (1859) 585; Sér. VII. xv. (1869) 185 = incertus.
pachyacanthus, *Bunge, Astrag.* i. 82; ii. 143.—Persia.
pachycarpus, Torr. & Gray, Fl. N. Am. i. 332 = caryocarpus.
pachypus, *Greene, in Bull. Calif. Acad.* i. n. 3. (1885) 157.—Calif.
pachystachys, *Bunge, Astrag.* i. 80; ii. 134.—Persia.
paecilanthus, *Boiss. & Heldr. Diagn.* Ser. II. vi. 54.—Phrygia.
Pallasii, *Fisch. Hort. Gorenk.* (1813) 71; *et ex DC. Prod.* ii. 305 (1825).—Deserta Casp.
Pallasii, *Spreng. Fl. Hal. Mant.* (1807) 48.—Rossia.
pallescens, *Bieb. Fl. Taur. Cauc.* iii. 489.—Bessarab.; Podol.
pallidus, Boiss. Fl. Orient. ii. 372 = drusorum.
Palliseri, *A. Gray, in Proc. Am. Acad.* vi. (1864) 227. —Am. bor.
Palmeri, *A. Gray, l. c.* vii. (1868) 398.—Am. bor. occ.
pamphylicus, *Boiss. Fl. Orient.* ii. 239.—Cilicia.
Pancicii, *Heuff. in Flora*, xxxvi. (1853) 621.—Europ.
panduratus, *Bunge, Astrag.* i. 59; ii. 95.—Anatolia.
pannonicus, Schult. Oestr. Fl. ed. II. ii. 335 = dasyanthus.
pannosus, *Fenzl, in Tchihat. Asie Min. Bot.* i. 2.—Cilicia.
paphlagonicus, Bunge, Astrag. i. 121; ii. 210 = atropurpurcus.
paradoxus, *Bunge, l. c.* i. 64; ii. 108.—Persia.
paralipomenus, *Bunge, l. c.* i. 81; ii. 136.—Persia.
paralurges, *Bunge, l. c.* i. 67; ii. 111.—Persia.
paraplesius, *Bunge, l. c.* i. 46; ii. 73.—Persia.
Parishii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 75. —Am. bor.
Parnassi, *Boiss. Diagn.* Ser. I. ix. 80.—Graecia; Oriens.
Parnassi, Heldr. ex Nym. Consp. 192 = aristatus, cephalonicus.
parrowianus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 320.—Persia.
Parryi, C. Anders. Cat. Pl. Nev. 120 = malacus.
Parryi, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 410.—Am. bor. occ.
parviceps, *Bunge, Astrag.* i. 91; ii. 162.—As. Min.
parviflorus, Desf. ex Spreng. Syst. iii. 300, in syn. = racemosus.
parviflorus, Lam. Encyc. i. 310 = Oxytropis deflexa.

ASTRAGALUS :—
parvifolius, Nutt. ex A. Gray, in Proc. Am. Acad. vi. (1864) 202 = gracilis.
parvus, *Hemsl. Biol. Centr. Am. Bot.* i. 266.—Mexic.
Pastellianus, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 95 = vesicarius.
Pattersoni, *A. Gray, in Brandegee, Fl. S. W. Col.* 285.—Am. bor. occ.
pauciflorus, A. Gray, in Proc. Acad. Philad. (1863) 60 = leptaleus.
pauciflorus, Hook. in Lond. Journ. Bot. vi. (1849) 210 = alpinus.
pauciflorus, *Hook. Fl. Bor. Am.* i. 149.—Am. bor. occ.
pauciflorus, Pall. Astrag. 81. t. 66 = Gueldenstaedtia pauciflora.
paucijugus, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 196.—Turkestan.
pauper, *Bunge, Astrag. Fedsch.* 224.—As. centr.
pectinatus, *Dougl. ex Hook. Fl. Bor. Am.* i. 142.—Am. bor.
pectinatus, *Boiss. Diagn.* Ser. I. ii. 54.—Syria.
peduncularis, *Royle, Illustr. Bot. Himal.* 199.—Tibet. occ.
Pehuenches, *Niederl. in Roca, Exped. Rio Negro*, ii. 206.—Patagonia.
pelliger, *Fenzl, Pugill. Pl. Nov. Syr.* 5.—Cilicia.
pellitus, *Bunge, Astrag.* i. 33; ii. 39.—Persia.
penduliflorus, Lam. Fl. Fr. ii. 636 = alpinus.
pendulus, DC. Astrag. 232 = elongatus.
pendulus, Bunge, Astrag. ii. 211 = caelestis.
pennatus, *Bunge, l. c.* ii. 148.—Cataonia.
pentaglottis, *Linn. Mant.* 274.—Reg. Mediterr.
pentapetaloides, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 274 = flexus.
peregrinus, Fisch. & Mey. ex Bunge, Astrag. ii. 34 = A. Caraganae.
peregrinus, *Vahl, Symb. Bot.* i. 57.—Aegypt.; Palaestin.
peristereus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 351.—Persia.
permiensis, C. A. Mey. ex Rupr. Fl. Bor. Ural. ii. 82. t. 11 = Helmii.
perpusillus, Bertol. in Nov. Comm. Bonon. vi. (1844) 230 = tribuloides.
persepolitanus, Boiss. Diagn. Ser. I. ix. 60 = ammophilus.
persicus, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 3.—Reg. Casp.; Persia.
peruvianus, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 18.—Peruv.
Pestalozzae, Boiss. Diagn. Ser. I. ix. 41 = cadmicus.
petraeus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 333.—Mont. Alatav.
petropylensis, *Bunge, Astrag.* i. 139; ii. 237.—Sibir.
phacaeformis, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. 590; *Verz. Suppl. Fl. Alt.* 90.—Sibir. altaic.
phlomoides, *Boiss. Diagn.* Ser. I. ii. 56.—Persia.
physaloides, *DC. Prod.* ii. 300.—Anatolia vel Syria.
physocalyx, *Fisch. in Bull. Sc. Acad. Pétersb.* ii. (1837) 74.—Thracia.
physocalyx, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 409 = cysticalyx.
physocarpus, Ledeb. Fl. Alt. iii. 335 = physodes.
physodes, *Linn. Sp. Pl.* 760.—Reg. Casp.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 415.—Peruv.
pictus, Boiss. & Gaill. Diagn. Ser. II. vi. 55 = conduplicatus.
pictus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 214.—Am. bor. occ.
pictus, *Steud. Nom.* ed. II. i. 163.—Chili.
piestolobus, *Bunge, Astrag.* i. 38; ii. 52.—Persia.
pilosus, Linn. Sp. Pl. 756 = Oxytropis pilosa.
Pinardi, *Boiss. Diagn.* Ser. I. ix. 56.—As. Min.
pinetorum, *Boiss. l. c.* ii. 77.—As. Min.; Syria.
piptocephalus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 376.—Persia.
pisidicus, *Boiss. & Heldr. Diagn.* Ser. I. ix. 70.—Pisidia.
Pistacia, Rost. ex DC. Prod. ii. 305 = elongatus.
placens, *Clos, in C. Gay, Fl. Chil.* ii. 109.—Chili.
plattensis, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 332.—Am. bor.
platyphyllus, Fisch. ex Bunge, Astrag. ii. 205 = choicus.
platyphyllus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 345.—Soongar.
platyraphis, *Fisch. ex Bunge, Astrag.* i. 37; ii. 48.—Syria; Kurdist.; Mesopot.
platysematus, *Bunge, Astrag.* i. 120; ii. 205.—Persia.

ASTRAGALUS :—
platytropis, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 526.—Am. bor. occ.
plebejus, *Boiss. Diagn.* Ser. I. vi. 37.—Persia.
plumosus, DC. ex Bunge, Astrag. ii. 148 = Lamarckii.
plumosus, Sieber, ex Bunge, l. c. ii. 59 = bethlemiticus.
plumosus, *Willd. Sp. Pl.* iii. 1333.—Galatia.
podocarpus, *C. A. Mey. Verz. Pfl. Cauc.* 142.—Reg. Caucas.; Persia.
podocarpus, R. Grah. in Wall. Cat. n. 5930 = A. Emodi.
podocarpus, *Buch.-Ham. in Wall. l. c.*—Ind. or.
podolobus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 49.—Persia.
podosphaerus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 255.—Kurdistan.
poecilanthus, Bunge, Astrag. ii. 188 = paecilanthus.
polaris, *Benth. in Trans. Linn. Soc.* xxiii. (1861) 323.—Am. arct. occ.
polemius, *Boiss. Fl. Orient.* ii. 306.—Bcluchist.
polemoniacus, *Bunge, Astrag.* i. 135; ii. 233.—Anatolia.
poliotes, *Bunge, Astrag.* i. 116; ii. 197.—Sibir. altaic.
polyacanthus, *Royle, Illustr. Bot. Himal.* 199.—Reg. Himal.
polyacanthus, Wall. Cat. n. 5934 = Caragana polyacantha.
polyactinus, *Boiss. Fl. Orient.* ii. 226, in nota.—Hispan.; Afr. bor.
polyanthus, *Bunge, Astrag.* i. 93; ii. 165.—Persia.
polybotrys, *Boiss. Diagn.* Ser. I. ix. 64.—Afghan.
polycephalus, Tenore, ex Bunge, Astrag. ii. 95 = ponticus.
polyceras, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 332.—Soongar.
Polygala, Pall. Astrag. n. 116. t. 83 = sanguinolentus.
polyphyllus, *Bunge, Astrag.* i. 35; ii. 59.—Reg. Caucas.
polyphyllus, Willd. Sp. Pl. iii. 1300 = Oxytropis microphylla.
ponticus, *Pall. Astrag.* 14. t. 11.—Reg. Caucas.; As. Min.
porphyraemus, *Boiss. Fl. Orient.* ii. 367.—Cappadoc.
porphyrobaphis, *Fisch. in Bull. Soc. Nat. Mosc.* (1853) II. 462.—Persia.
porphyrolepis, Fisch. l. c. 378 = radkanensis.
porrectus, *S. Wats. in Bot. King, Exp.* 75, 411.—Am. bor. occ.
Poterium, DC. Astrag. 160 = Clusii.
Poterium, Pall. Astrag. 1. t. 1 = arnacantha.
Poterium, *Vahl, Symb. Bot.* i. 63.—Hispan.
praecox, Baumg. Enum. Stirp. Transs. ii. 362 = Wulfeni.
praetermissus, *Ball, in Journ. Bot.* xi. (1873) 306.—Marocco.
Preussii, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 222.—Am. bor. occ.
procerus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 464.—Persia.
procerus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 369.—Am. bor.
procumbens, *Hook. & Arn. Bot. Beech. Voy.* 18.—Chili.
procumbens, Mill. Gard. Dict. ed. VIII. n. 18 = pentaglottis.
procumbens, *S. Wats. in Proc. Am. Acad.* xx. (1885) 361.—N. Mexic.
prolixus, *Sieber, ex Bunge, Astrag.* i. 9; ii. 6.—Afr. bor.; Ind. bor. occ.
prostratus, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. (1838) 90 = adsurgens.
prostratus, *Hook. & Arn. Bot. Beech. Voy.* 18.—Chili.
prostratus, Scop. Delic. Insub. ii. 106 = Oxytropis prostrata.
prusianus, Boiss. Diagn. Ser. I. ix. 88 = Boissieri.
Przewalskii, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 32.—China.
pseudaustralis, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* xi. *Suppl.* 16.—Sibir. altaic.
pseudocaspius, *Fisch. in Bull. Soc. Nat. Mosc.* (1853) II. 347.—Anatolia.
pseudocicer, Opiz, in Flora, iv. (1821) 758 = A. Cicer.
pseudonobrychis, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 238, nomen.—Hab. ?
pseudopentaglottis, Boiss. & Noë, ex Boiss. Fl. Orient. ii. 226 = triradiatus.
pseudostella, *Delile, Fl. Egypt. Suppl.* 64.—Graecia; Aegypt.; Arab.

ASTRAGALUS :—
pseudotragacantha, Pall. Astrag. 3. t. 3 = aureus.
pseudotragacantha, Urv. in Mém. Linn. Soc. Par. i. (1822) 348 = thracicus.
psilacanthus, *Boiss. Diagn.* Ser. I. ix. 97.—Afghan.
psilacanthus, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) 407 = tenuispinus.
psilacmos, *Bunge, Astrag.* i. 86; ii. 151.—Syria.
psilocentros, Fisch. in Bull. Soc. Nat. Mosc. (1853) II. 405 = polyacanthus.
psilodontius, *Boiss. Diagn.* Ser. I. ix. 86.—Syria.
psiloglottis, Stev. in DC. Prod. ii. 288 = oxyglottis.
psilopterus, Bunge, Astrag. i. 45; ii. 71 = genistoides.
psilopus, Schrenk, in Bull. Sc. Acad. Pétersb. x. (1842) 254 = spartioides.
psilostylus, *Bunge, Astrag.* i. 87; ii. 152.—Persia.
psoraleoides, Boiss. ex Bunge, l. c. ii. 235 = asciocalyx.
psoralioides, *Willd. Sp. Pl.* iii. 1309.—Cappadoc.
psoraloides, Lam. Encyc. i. 317; DC. Astrag. 145. t. 18 = canescens.
pterocarpus, *S. Wats. in Bot. King, Exp.* 71.—Am. bor. occ.
pterocephalus, *Bunge, Astrag. Fedsch.* 252.—As. centr.
pterostylis, *DC. Prod.* ii. 294.—Austral.
ptilocephalus, *Baker, ex Aitch. in Journ. Linn. Soc.* xviii. (1881) 47.—Afghan.
ptilodes, *Boiss. Diagn.* Ser. I. ix. 81.—As. Min.
ptychophyllus, *Boiss. l. c.* vi. 42.—Persia.
pubentissimus, *Torr. & Gray, Fl. N. Am.* i. 693.—Am. bor. occ.
puberulus, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 405. n. 240 = macropterus.
puberulus, *Ledeb. Fl. Alt.* iii. 299.—Sibir. altaic.
pubescens, Schrank, Prim. Sal. 2 = frigidus.
pubiflorus, *DC. Astrag.* 173.—Sibir.
pubigerus, Pall. ex Trautv. in Act. Hort. Petrop. viii. (1883) 238 = pubiflorus.
pugioniferus, *Fisch. ex Bunge, Astrag.* i. 125; ii. 219.—Macedonia.
pugniformis, L'Hérit. Stirp. Nov. 170 = longifolius.
pugniformis, Willd. Sp. Pl. iii. 1334 = oleaefolius.
pulchellus, *Boiss. Diagn.* Ser. I. ii. 86.—Persia.
pulchellus, Salisb. Prod. 337 = danicus.
Pulsiferi, *A. Gray, in Proc. Am. Acad.* x. (1874) 69.—Am. bor. occ.
pulvinatus, *Bunge, Astrag.* i. 82; ii. 142.—Persia.
Pumilio, Pall. Astrag. 67. t. 55 = Oxytropis Pumilio.
punctatus, *Bunge, Astrag.* i. 120; ii. 209.—Persia.
pungens, Willd. Sp. Pl. iii. 1325 = angustifolius.
purpurascens, *Bunge, Astrag.* i. 35; ii. 42.—Afghan.
purpurascens, Jacq. f. ex Steud. Nom. ed. II. i. 163 = A. Onobrychis.
purpureus, Lam. Encyc. i. 314 = hypoglottis.
Purshii, *Dougl. ex G. Don, Gen. Syst.* ii. 271.—Am. bor. occ.
pusillus, *Vog. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 19.—Peruv.
pycnocephalus, *Fisch. in Bull. Soc. Nat. Mosc.* (1853) II. 371.—Kurdistan.
pycnocladus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 336.—Persia.
pycnolobus, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 594.—Sibir. altaic.
pycnophyllus, Kotschy, ex Bunge, Astrag. ii. 138 = mesoleios.
pycnophyllus, Stev. in Mém. Soc. Nat. Mosc. iv. (1814) 57 = microcephalus.
pycnorrhizus, *Wall. Cat.* n. 5927.—Reg. Himal.
pycnostachyus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 527.—Am. bor. occ.
pygmaeus, DC. ex Bunge, Astrag. ii. 86 = nanus.
pygmaeus, Pall. Astrag. 66. t. 54 = Oxytropis nigrescens.
pyramidulus, Salzm. ex Bunge, Astrag. ii. 5 = epiglottis.
pyrrhotrichus, *Boiss. Diagn.* Ser. I. ix. 73.—Afghan.
quadrisulcatus, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 248.—Persia; Afghan.; Turcoman.
quinquefoliolatus, *Bunge, Astrag.* i. 119; ii. 204.—Persia.
Quisqualis, *Bunge, Astrag. Fedsch.* 222.—As. centr.
racemosus, *Pursh, Fl. Am. Sept.* ii. *Suppl.* 740.—Am. bor.
racemulosus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 391.—Persia.
Raddeanus, Regel, Ind. Sem. Hort. Petrop. (1865) 40 = flaccidus.
radians, *Pomel, Nouv. Mat. Fl. Atl.* 185.—Algeria.
radiatus, Steud. Nom. ed. II. i. 163 = Sieberi.

ASTRAGALUS:—
radiatus, *Ehrenb. ex Bunge, Astrag.* i. 11; ii. 8.—Aegypt.; Arab.
radiatus, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 267 = A. Stella.
radicans, *Hornem. Hort. Hafn.* ii. 708.—Hab.?
radicatus, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 268 = Sieberi.
radiciflorus, Steud. Nom. ed. II. i. 163 = testiculatus.
radkanensis, *Bunge, Astrag.* i. 81; ii. 138.—Persia.
ramosissimus, *Scop. Delic. Insub.* ii. 105.—Sibir.
raphiodontus, *Boiss. Diagn.* Ser. I. ix. 96.—Afghan.
rariflorus, *Ledeb. Fl. Alt.* iii. 313.—Soongaria.
Rattani, *A. Gray, in Proc. Am. Acad.* xix. (1883) 75.—Calif.
Rauwolfii, Vahl, Symb. Bot. i. 63 = Russelii.
Reboudii, *Coss. ex Bunge, Astrag.* i. 37; ii. 49.—Afr. bor.
recognitus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 452.—Persia.
recurvus, *Greene, in Bull. Calif. Acad.* i. n. 3. (1885) 155.—Calif.
reduncus, *Pall. Astrag.* 109. t. 82.—Reg. Caucas.
reflexistipulus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 44.—Japon.
reflexus, *Torr. & Gray, Fl. N. Am.* i. 334.—Am. bor. occ.
refractus, Boiss. & Buhse, in Boiss. Diagn. Ser. II. ix. 57 = Buhseanus.
refractus, *C. A. Mey. Verz. Pfl. Cauc.* 144.—Reg. Caucas.
Regelii, *Trautv. in Act. Hort. Petrop.* ii. (1873) 472.—Armenia.
Reinii, *Ball, in Journ. Linn. Soc.* xvi. (1878) 432.—Marocco.
remotiflorus, *Boiss. Diagn.* Ser. I. ii. 72.—Persia.
remotifolius, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 1095 (emend.).—Persia.
remotijugus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 74.—Persia.
reptans, *Willd. Hort. Berol.* ii. 88. t. 88.—Mexic.
resupinatus, Bieb. Fl. Taur. Cauc. iii. 493 = fragrans.
retamocarpus, *Boiss. & Hohen. ex Boiss. Diagn.* Ser. I. ix. 68.—Persia.
reticulatus, *Bieb. Fl. Taur. Cauc.* iii. 491.—Reg. Caucas.; Afghan.
retroflexus, Pall. Astrag. 33. t. 27 = Oxytropis deflexa.
retusus, DC. Astrag. 205. t. 36 = coluteoides.
retusus, Willd. in Mém. Acad. Berl. (1794–95) 24. t. 2 = angustifolius.
Reuterianus, *Boiss. Diagn.* Ser. I. ii. 71.—Persia.
reventus, *A. Gray, in Proc. Am. Acad.* xv. (1880) 46.—Am. bor.
rhacodes, *Bunge, Astrag. Fedsch.* 308.—As. temp.
rhizanthus, Boiss. Diagn. Ser. I. ii. 83 = cariensis.
rhizanthus, *Royle, Illustr. Bot. Himal.* 199.—Reg. Himal.
rhizocephalus, *Baker, ex Aitch. in Journ. Linn. Soc.* xviii. (1878) 46.—Ind. or.
rhodochrous, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 825.—Kurdistan.
rhodosemius, *Boiss. & Haussk. l. c.* 363.—Persia.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 414.—Peruv.
rigidulus, *Benth. ex Bunge, Astrag.* ii. 25.—Reg. Himal.
rigidus, Boiss. & Bal. Diagn. Ser. II. vi. 55 = melanocephalus.
Robbinsii, *A. Gray, Man. Bot. U. St.* ed. II. 98.—Am. bor.
Robbinsii, Matthew, Pl. Acad. 15 = alpinus.
robustus, *Bunge, Astrag.* i. 122; ii. 212.—Persia.
Rochelianus, Heuff, ex Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) I. 293 = A. Onobrychis.
roseus, *Ledeb. Fl. Alt.* iii. 330.—Sibir. altaic.
Rostani, Boiss. Diagn. Ser. I. ii. 45 = densifolius, chrysophyllus.
rostratus, *C. A. Mey. Verz. Pfl. Cauc.* 144.—Reg. Caucas.
rotundifolius, Presl, ex Steud. Nom. ed. II. i. 163 = glycyphyllus.
rotundifolius, Royle, Illustr. Bot. Himal. 199 = graveolens.
rotundifolius, Willd. Sp. Pl. iii. 1317 = nummularioides.
Roussacanus, *Boiss. Diagn.* Ser. I. ii. 61.—Syria; Mesopotam.
Royleanus, Bunge, Astrag. i. 30; ii. 34 = Candollcanus.
Royleanus, D. Dietr. Syn. Pl. iv. 1090 = strobiliferus, *Royle*.
rubriflorus, *Bunge, Astrag.* i. 67; ii. 109.—Persia.

ASTRAGALUS:—
rubrostriatus, *Bunge, Astrag.* i. 68; ii. 113.—Persia.
rudbaricus, Bunge, l. c. i. 123; ii. 215 = confusus.
rugosus, *Fisch. ex Bunge, l. c.* ii. 53.—Kurdistan.
rumelicus, Bunge, l. c. 81; ii. 137 = veluchensis.
runcinatus, *Boiss. Fl. Orient.* ii. 368.—Cappadoc.
rupicola, Ehrenb. ex Bunge, Astrag. ii. 19 = peregrinus.
rupifragus, *Pall. Astrag.* 86. t. 70.—Reg. Caucas.; Ural.
Ruprechti, *Bunge, Astrag.* i. 101; ii. 174.—Reg. Caucas.
Rusbyi, *Greene, in Bull. Calif. Acad.* i. n. 1 (1884) 8.—Calif.
ruscifolius, *Boiss. Diagn.* Ser. I. ii. 47.—Persia.
Russelii, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 260. t. 13.—Syria; Mesopotam.
rytidocarpus, *Ledeb. Fl. Alt.* iii. 315.—Sibir. altaic.
rytilobus, *Bunge, Astrag. Fedsch.* 209.—As. temp.
sabuletorum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 408 = arkalycensis.
sabuletorum, *Ledeb. Fl. Alt.* iii. 321.—Sibir. altaic.
sabulorum, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 368.—Am. bor.
saccatus, *Boiss. Diagn.* Ser. I. ix. 53.—Persia.
saccharensis, Ehrenb. ex Bunge, Astrag. ii. 16 = gyzensis.
saccocalyx, *Schrenk, Enum. Pl. Nov.* 83.—Soongaria.
sachalinensis, *Bunge, Astrag.* i. 23; ii. 26.—Ins. Sachalin.
saganlugensis, *Trautv. in Bull. Phys.-Math. Acad. Pétersb.* xvi. (1858) 323.—Armenia.
Saharae, *Pomel, Nouv. Mat. Fl. Atl.* 184.—Algeria.
Sahendi, *Buhse, in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 415.—Persia.
sakkarensis, Ehrenb. ex Boiss. Fl. Orient. ii. 234 = gyzensis.
salatavicus, *Bunge, Astrag.* i. 117; ii. 201.—Reg. Caucas.
salsugineus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 341.—Soongaria.
salsula, *Scop. Delic. Insub.* ii. 104.—Sibir.
salvatoris, Willk. in Willk. & Lange, Prod. Fl. Hisp. iii. 281 = massiliensis.
samamensis, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 68.—Persia.
sanctus, *Boiss. Diagn.* Ser. I. ix. 47.—Syria.
sanguinolentus, *Bieb. Casp.* 190.—Reg. Caucas.
saratagius, *Bunge, Astrag. Fedsch.* 269.—As. temp.
scaberrimus, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 91.—China bor.
scabrifolius, *Boiss. Diagn.* Ser. I. ii. 81.—Armenia.
scabrisetus, *Bong. Verz. Pfl. Saisang-Nor*, n. 89. t. 5.—Soongaria.
scaposus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 366.—Am. bor.
scariosus, Benth. ex Bunge, Astrag. i. 44; ii. 71 = polyacanthus.
schahrudensis, *Bunge, l. c.* i. 61; ii. 103.—Persia.
Schanginianus, *Pall. Astrag.* 77. t. 73.—Sibiria.
Schelichowii, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 68.—Ochotsk.
Schimperi, *Boiss. Diagn.* Ser. I. ii. 53.—Arabia.
schirasicus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 457.—Persia.
schistocalyx, *Bunge, Astrag.* i. 46; ii. 75.—Persia.
schistosus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 49.—Persia.
schizopterus, *Boiss. l. c.* ii. 82.—Cilicia; Syria.
Schmalhauseni, *Bunge, in Act. Hort. Petrop.* vii. (1880) 369.—Turkestan.
Schottianus, *Boiss. Diagn.* Ser. II. v. 89.—Cilicia; Cappadoc.
Schrenkianus, *Fisch. & Mey. in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 197.—Turkestan.
schurabicus, *Bunge, Astrag.* i. 76; ii. 129.—Persia.
sciadophorus, *Franch. Pl. Davidian.* i. 84.—As. or.
sciurcus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 98.—Persia.
sclerocarpus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 225.—Am. bor. occ.
sclerocladus, *Bunge, Astrag.* i. 46; ii. 75.—Persia.
scleropodius, *Ledeb. Fl. Alt.* iii. 326; *Fl. Ross. Alt.* t. 295.—Sibir. altaic.
scleroxylon, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 241.—Turkestan.
scopaeformis, Ledeb. Fl. Ross. i. 615 = tauricus.
scoparius, *Schrenk, Enum. Pl. Nov.* 82.—As. temp.
scopulorum, *Porter & Coult. Fl. Col.* 24.—Colorado.

ASTRAGALUS:—
scorpioides, *Pourr. ex Willd. Sp. Pl.* iii. 1280.—Hispan.
scorpius, *Boiss. Diagn.* Ser. I. ix. 103.—Persia.
scorpius, Buhse, ex Bunge, Astrag. ii. 127 = glaucacanthus.
Scorpiurus, Bunge, l. c. i. 5; ii. 3 = hippocrepidis.
Scottianus, Stocks, ex Bunge, l. c. ii. 31 = medullaris.
scrobiculatus, Boiss. Fl. Orient. ii. 490 = gymnolobus.
secundus, *DC. Astrag.* 128. n. 55.—Sibir. austr.
secundus, Michx. Fl. Bor. Am. ii. 67 = alpinus.
segetalis, Boiss. Fl. Orient. ii. 277 = neurocarpus.
Segethi, *Phil. in Linnaea*, xxxiii. (1864–65) 47.—Chili.
scidabadensis, *Bunge, Astrag.* i. 68; ii. 113.—Persia.
Seidlitzii, *Bunge, l. c.* i. 40; ii. 56.—Persia.
Semenovii, *Bunge, in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 22.—As. temp.
semibilocularis, *DC. Astrag.* 170.—Sibiria.
semibilocularis, *Fisch. ex Bunge, Astrag.* i. 103; ii. 183.—Sibir. altaic.
semipellitus, *Bunge, l. c.* i. 83; ii. 144.—Persia.
sempervirens, Lam. Encyc. i. 320 = aristatus.
senganensis, *Bunge, Astrag.* i. 82; ii. 142.—Persia.
serbicus, Panč. ex Boiss. Fl. Orient. ii. 268 = glycyphylloides.
sericeus, DC. Astrag. 148 = globosus.
sericeus, Lam. Fl. Fr. ii. 645 = versicarius, A. Oxytropis, sordida.
sericeus, Willd. Enum. Hort. Berol. Suppl. 51 = hamosus.
sericoleucus, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 410.—Am. bor. occ.
sericophyllus, *Griseb. Spicil. Fl. Rumel.* i. 53.—Graecia.
sericostachyus, *Stocks, in Hook. Kew Journ.* iv. (1852) 146.—Beluchist.
serotinus, *A. Gray, in Pacif. Rail. Rep.* xii. 18, 51.—Am. bor. occ.
sesameus, *Linn. Sp. Pl.* 759.—Reg. Mediterr.
sesameus, Pall. Astrag. 98 = oxyglottis.
sesamoides, *Boiss. Diagn.* Ser. I. ix. 59.—Afghan.
seshanoides, Benth. in Royle, Illustr. Bot. Himal. 199 = trichocarpus.
sesquiflorus, *S. Wats. Proc. Am. Acad.* x. (1875) 346.—Utah.
setiferus, DC. Prod. ii. 296 = Cornulaca tragacanthoides.
setosus, Pall. Astrag. 55. t. 44 = Oxytropis setosa.
setulosus, *Boiss. & Bal. Diagn.* II. vi. 50.—Cappadoc.
Sewersowii, Boiss. Fl. Orient. ii. 289 = seq.
Sewertzowii, *Bunge, Astrag.* i. 38; ii. 54.—Turkest.
seytunensis, *Bunge, l. c.* i. 87; ii. 154.—Syria.
Shortianus, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 331.—Am. bor. occ.
Shortianus, S. Wats. Pl. Wheeler, 7 = cyaneus.
sibiricus, *Scop. Delic. Insub.* ii. 104.—Sibiria.
Sibthorpianus, *Boiss. Diagn.* Ser. I. ii 84.—Bithynia.
siculus, '*Bivon. Giorn. Palerm.* n. 4.'—Sicilia.
siculus, Gasp. ex Nym. Consp. 192 = calabricus.
siculus, Schimp. ex Nym. l. c. = cephalonicus.
Sieberi, *DC. Prod.* ii. 295.—Aegypt.; Arab.
Sieversianus, *Pall. Astrag.* 15. t. 12.—Persia; Soongaria.
Sieversianus, Steph. ex Ledeb. Fl. Ross. i. 592 = Oxytropis argentata.
Sieversii, Coq. Monb. ex Bunge, Astrag. ii. 36 = diphtherolobus.
sigmoideus, *Bunge, l. c.* i. 123; ii. 214.—Anatolia.
sikkimensis, *Benth. ex Bunge, l. c.* i. 23; ii. 25.—Reg. Himal.
siliquosus, *Boiss. Diagn.* Ser. I. ii. 51.—Persia.
siliquosus, Boiss. l. c. vi. 40 = lobophorus.
silvicolus, Pritz. Ic. Ind. 123 = Oxytropis sylvatica.
simplicifolius, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 231.—Am. bor. occ.
sinaicus, Boiss. Diagn. Ser. I. ix. 57 = pseudostella.
singarensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 263.—Mesopotam.
sinicus, *Linn. Mant.* i. 103.—China.
sinicus, Thunb. Fl. Jap. 290 = lotoides.
sirinicus, *Tenore, Prod. Fl. Neap. App.* v. 25 (1826)'; *Fl. Neap.* v. 132. t. 171.—Italia.
sisyrodytes, *Bunge, Astrag.* i. 33; ii. 40.—Turkestan.
sitiens, *Bunge, l. c.* i. 126; ii. 220.—Persia.
sivasicus, *Bunge, l. c.* i. 93; ii. 166.—Armenia.
skythropos, *Bunge, ex Maxim. in Bull. Acad. Petersb.* xxiv. (1878) 31.—China.
soficus, *Bunge, Astrag.* i. 81; ii. 136.—Persia.
sogdianus, *Bunge, Astrag. Fedsch.* 268.—As. temp.

ASTRAGALUS :—

Solandri, *Lowe, in Hook. Kew Journ.* viii. (1856) 294. —Marocco ; Madera.
somcheticus, C. Koch, in Linnaea, xv. (1841) 720 = sanguinolentus.
songaricus, Pall. Astrag. 63. t. 51 = Oxytropis songarica.
Sonorae, A. Gray, Pl. Wright. ii. 44 = arizonicus.
sordidus, Willd. Sp. Pl. iii. 1313 = Oxytropis campestris.
Soyeri, *Buching. ex Bunge, Astrag.* i. 118 ; ii. 199.—Dalmat.
Spachianus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 67.—Persia.
Spaldingii, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 524.—Am. bor. occ.
sparsiflorus, *A. Gray, in Proc. Acad. Philad.* (1863) 60.—Am. bor. occ.
sparsus, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 267.—Arab.
spartioides, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 330.—Soongar.
speciosus, *Boiss. & Hohen. Diagn.* Ser. I. ix. 67.—Persia.
speirocarpus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 225.—Am. bor. occ.
sphaeranthus, *Boiss. Diagn.* Ser. I. vi. 37.—Persia.
sphaerocalyx, *Ledeb. Fl. Ross.* i. 647.—Mont. Ararat.
sphaerocarpus, *Clos, in C. Gay, Fl. Chil.* ii. 117.—Chili.
sphaerocarpus, *Desf. ex Steud. Nom.* ed. II. i. 163.—Chili.
sphaerocephalus, *Trautv. in Act. Hort. Petrop.* viii. (1883) 241.—Reg. Caucas.
sphaerocystis, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 273 (*Lehm. Rel. Bot.* (1847) 97).—As. centr.
sphaerophysa, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 338.—Soongar.
spicatus, Burm. ex Guill. & Perr. Tent. Fl. Seneg. 182 = Indigofera hirsuta.
spicatus, *Pall. Reise*, ii. 742.—Soongar.
spicatus, Nutt. in Torr. & Gray, Fl. N. Am. i. 336 = Mortoni.
spinellus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 315.—Persia.
spinescens, *Bunge, Astrag.* i. 42 ; ii. 64.—Persia.
spinosissimus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = A. Tragacantha.
spinosissimus, *Wall. Cat.* n. 5935.—Ind. or.
Spruneri, *Boiss. Diagn.* Ser. I. ii. 79.—Graecia.
squalidus, *Boiss. l. c.* Ser. II. ii. 29.—As. Min.
squamulosus, Poir. Encyc. Suppl. i. 528 = Oxytropis squamulosa.
squarrosus, *Bunge, Astrag.* i. 129 ; ii. 226.—Persia.
Ssahendi, Boiss. Fl. Orient. ii. 311 = Sahendi.
Stella, Bieb. Fl. Taur. Cauc. ii. 191 = cruciatus.
Stella, *Gouan, Illustr.* 50.—Gall. austr. ; Afr. bor.
stellatus, Lam. Fl. Fr. ii. 641 = A. Stella.
stenanthus, *Bunge, Astrag. Fedsch.* 274.—As. temp.
stenoceras, *C. A. Mey. in Bong. Verz. Pfl. Saisang-Nor*, 24.—Soongar.
stenocystis, *Bunge, Astrag. Fedsch.* 301.—As. temp.
stenolepis, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 387.—Persia.
stenolobus, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 593 = ceratoides.
stenophyllus, *Torr. & Gray, Fl. N. Am.* i. 329.—Am. bor.
stenorhachis, Fisch. in Bull. Soc. Nat. Mosc. xxvi. (1853) II. 388 = andrachnaefolius.
stenosemius, *Boiss. Diagn.* Ser. II. ii. 27.—Armenia.
stenostegius, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 314.—Kurdistan.
Stevenianus, *DC. Prod.* ii. 285.—Reg. Caucas. ; Persia.
Stewartii, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 129.—Ind. bor. occ.
stipitatus, *Benth. ex Boiss. Fl. Orient.* ii. 305.—Afghanist.
stipitatus, Sims, Bot. Mag. t. 2380 = stipulatus.
stipularis, Sims, ex D. Don, l. c. l. Index = seq.
stipulatus, *D. Don, ex Sims, l. c.* t. 2380.—Reg. Himal.
stipulatus, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 131 = graveolens.
Stocksii, *Benth. ex Bunge, Astrag.* i. 6 ; ii. 4.—Afghan.
stolzenburgensis, Lerch. ex Schur, Enum. Pl. Transs. 164 = eriocephalus.

ASTRAGALUS :—

stramineus, *Boiss. Diagn.* Ser. II. v. 85.—Syria.
streptopus, *Greene, in Bull. Calif. Acad.* i. n. 3 (1885) 155.—Calif.
striatellus, *Bieb. Fl. Taur. Cauc.* ii. 189.—Reg. Caucas.
striatus, Nutt. in Torr. & Gray, Fl. N. Am. i. 330 = adsurgens.
strictifolius, *Boiss. Diagn.* Ser. I. ix. 78.—Kurdistan. ; Persia.
strictispinus, *Boiss. Fl. Orient.* ii. 366.—Lycia.
strictissimus, Sievers, ex Pall. Astrag. 16 = folliculares.
strictus, *R. Grah. in Wall. Cat.* n. 5924.—Reg. Himal.
strictus, Siev. ex Fisch. in DC. Prod. ii. 301 = folliculares.
strigillosus, *Bunge, Astrag.* i. 102 ; ii. 178.—Anatolia.
strigulosus, *H. B. & K. Nov. Gen. et Sp.* vi. 494.—Mexic.
strobiliferus, *Lindl. in Bot. Reg.* (1849) *Misc.* 39.—Armenia.
strobiliferus, *Royle, Illustr. Bot. Himal.* 199.—Reg. Himal.
stromatodes, *Bunge, Astrag.* i. 85 ; ii. 148.—Syria.
subacaulis, Bunge, l. c. i. 101 ; ii. 176 = karputanus.
subalpinus, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 69.—Persia.
subbarbellatus, *Bunge, Astrag. Fedsch.* 227.—Turkestan.
subbiflorus, Lag. ex DC. Prod. ii. 289 = scorpioides.
subbijugus, *Ledeb. Fl. Ross.* i. 627.—Turkestan.
subcaulescens, Ledeb. Fl. Ross. i. 644 = Ruprechti, hyalolepis.
subcaulescens, *Royle, Illustr. Bot. Himal.* 199.—Ind. or.
subcinereus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 366.—Am. bor.
subcompressus, *A. Gray, in Brandegee, Fl. S. W. Col.* 234.—Am. bor. occ.
suberosus, [Soland. in] Russell, Aleppo, ed. II. ii. 260 = tuberculosa.
submitis, *Boiss. & Hohen. Diagn.* Ser. I. ix. 100.—Persia.
subsecundus, *Boiss. & Hohen. l. c.* 54.—Persia.
subulatus, Desf. Fl. Atlant. ii. 186 = annularis.
subulatus, Koch, in Flora, ex Steud. Nom. ed. II. i. 159, in syn. = argenteus.
subulatus, *Pall. ex Bieb. Fl. Taur. Cauc.* ii. 193.—Reg. Caucas. ; Tibet. ; Afghan. ; Sibir.
subuliferus, *Boiss. & Kotschy, ex Bunge, Astrag.* i. 31 ; ii. 37.—Cilicia.
subuliformis, DC. Astrag. 107 = subulatus.
subumbellatus, Klotzsch, in Bot. Ergeb. Waldem. Reise, 159 = hippocrepidis.
succulentus, Richards. in Frankl. 1st Journ. App. 746 = caryocarpus.
succumbens, *Dougl. ex G. Don, Gen. Syst.* ii. 263.—Am. bor. occ.
succumbens, Torr. & Gray, in Pacif. Rail. Rep. ii. 163 = Parryi.
suffalcatus, *Bunge, Astrag.* i. 125 ; ii. 219.—Afghan.
suffruticosus, *DC. Astrag.* 82.—Sibir.
suffruticosus, Pall. Reise, i. 440 = fruticosus.
suidunensis, *Bunge, in Act. Hort. Petrop.* vii. (1880) 378.—Turkestan.
sulcatus, Lam. Fl. Fr. ii. 639 = austriacus.
sulcatus, *Linn. Sp. Pl.* 756.—Sibir.
sulfureus, *Bunge, Astrag.* i. 29 ; ii. 33.—Afghan.
sultanensis, Bunge, l. c. i. 17 ; ii. 19 = bombycinus.
superbus, *Bunge, l. c.* i. 62 ; ii. 105.—Persia.
supinus, *C. A. Mey. ex Bunge, l. c.* i. 51 ; ii. 82.—Persia.
surugensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 402.—Mesopotam.
susianus, *Boiss. Diagn.* Ser. I. ii. 64.—Persia.
susianus, Kotschy, ex Bunge, Astrag. ii. 119 = acutus.
sylvaticus, Willd. Sp. Pl. iii. 1300 = Oxytropis sylvatica.
sylvicolus, Pall. Astrag. 95 = Oxytropis sylvatica.
syriacus, *Linn. Sp. Pl.* 759.—Syria.
syriacus, Pall. ex Ledeb. Fl. Alt. iii. 295 = Laxmanni.
syriacus, Linn. Herb. ex Bunge, in Boiss. Fl. Orient. ii. 479 = vimineus.
syrtschensis, *Bunge, Astrag.* i. 46 ; ii. 74.—Persia.
Szovitsii, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 473.—Armenia.

ASTRAGALUS :—

tabrizianus, *Buhse, in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 445.—Persia.
taeniolepis, Boiss. & Hohen. Diagn. Ser. I. ix. 99 = persicus.
talagonicus, *Boiss. & Hohen. l. c.* 87.—Persia.
talasseris, *Boiss. & Bal. l. c.* Ser. II. vi. 58.—Cappadoc.
talyschensis, *Bunge, Astrag.* i. 40 ; ii. 57.—Reg. Caucas.
tanaiticus, *C. Koch, in Linnaea*, xxiv. (1851) 94.—Oriens.
tarijensis, *Wedd. Chlor. And.* ii. 262.—Bolivia.
taschkendicus, *Bunge, Astrag. Fedsch.* 280.—As. temp.
tataricus, *Franch. Pl. Davidian.* i. 87.—Mongolia.
tauricolus, *Boiss. Diagn.* Ser. II. v. 86.—As. Min.
tauricus, Claus, in Goebel, Reise, ii. 269 = scopaeformis.
tauricus, *Pall. Astrag.* 48. t. 38.—Reg. Caucas. ; Ural. ; Soongar.
Tavernieri, *Boiss. & Diagn.* Ser. I. ii. 78.—Persia.
tegetarius, *S. Wats. in Bot. King, Exp.* 76, 444.—Am. bor. occ.
teheranicus, *Boiss. & Hohen. in Boiss. Diagn.* Ser. I. ix. 44.—Persia.
tenax, *Bunge, Astrag.* i. 67 ; ii. 110.—Persia.
tenellus, *Bunge, l. c.* i. 120 ; ii. 206.—Persia.
tenellus, Pursh, Fl. Am. Sept. ii. 473 = multiflorus.
tenellus, Willd. ex Steud. Nom. ed. II. i. 164 = strigulosus.
tener, *A. Gray, in Proc. Am. Acad.* vi. (1864) 206.—Calif.
tennesseensis, A. Gray, ex Chapm. Fl. S. U. St. 98 = plattensis.
tenuicaulis, *Benth. ex Bunge, Astrag.* i. 23 ; ii. 28.—Reg. Himal.
tenuifolius, *Desf. Fl. Atlant.* ii. 186.—Algeria.
tenuifolius, Georgi, It. i. 226 = versicolor.
tenuifolius, Linn. Sp. Pl. ed. II. 1065 = A. Onobrychis.
tenuirugis, *Boiss. Diagn.* Ser. I. ix. 61.—Arab.
tenuis, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 90 (*Cat. Baikal.* n. 354).—Sibir. baical.
tenuispinus, Bunge, Astrag. i. 44 ; ii. 70 = cicerifolius.
tephrodes, *A. Gray, Pl. Wright.* ii. 45.—N. Mexic.
tephrolobus, *Bunge, Astrag.* i. 127 ; ii. 222.—Sibir. altaic.
tephrosioides, *Boiss. Diagn*, Ser. I. ix. 69.—Afghan.
teres, Lam. Encyc. i. 524 = Oxytropis floribunda.
terminalis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 370.—Am. bor. occ.
testiculatus, *Pall. Astrag.* 82. t. 67.—Reg. Caucas. ; Soongar. ; Sibir.
tetragonocarpus, *Boiss. Diagn.* Ser. I. ii. 52.—Persia.
tetrapterus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 369.—Am. bor. occ.
tetrastichus, *Bunge, Astrag. Fedsch.* 279.—As. temp.
thessalus, Boiss. Diagn. Ser. I. ii. 80 = Spruneri.
Thompsonae, *S. Wats. in Proc. Am. Acad.* x. (1875) 345.—Am. bor. occ.
Thomsonianus, Benth. ex Bunge, Astrag. ii. 234 = nivalis.
thracicus, *Griseb. Spicil. Pl. Rumel.* i. 55.—Thracia.
Thurberi, *A. Gray, in Mem. Acad. Am.* N. S. v. (1855) 312.—Arizona.
tianschanicus, *Bunge, in Rupr. Sert. Tiansch.* 43.—As. temp.
tibetanus, *Benth. ex Bunge, Astrag.* i. 52 ; ii. 85.—Afghanist.
tigreensis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 193 = venosus.
Tigridis, *Boiss. Diagn.* Ser. I. ix. 55.—Mesopotam.
timuranus, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 256.—Turkest.
tmoleus, *Boiss. Diagn.* Ser. I. ix. 79.—Lydia.
toeniolepis, Boiss. & Hohen. ex Boiss. l. c. 99 = persicus.
tokatensis, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 370.—Anatolia.
tomentosus, *Lam. Encyc.* i. 312.—Aegypt.
torrentum, *Bunge, Astrag.* i. 40 ; ii. 55.—Persia.
tortuosus, *DC. Prod.* ii. 299.—Mesopotam.
Tournefortii, *Boiss. Diagn.* Ser. I. ix. 92.—Cappadoc.
trachyacanthos, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 381.—Persia.
trachytrichus, *Bunge, Astrag.* i. 101 ; ii. 173.—Armenia.
Tragacantha, Ucria, Hort. Panor. 316 = siculus.
Tragacantha, Habl. Taur. 112 = arnacantha.

ASTRAGALUS:—
Tragacantha, *Linn. Sp. Pl.* 762.—Reg. Medit. or.
Tragacantha, Pall. Astrag. t. 4. f. 1. Pl. 2 = massiliensis.
Tragacantha, Vill. Hist. Pl. Dauph. iii. 470 = aristatus.
tragacanthoides, Hohen. Enum. Pl. Talysch. 3 = talyschensis.
tragacanthoides, Lam. ex Steud. Nom. ed. II. i. 164 = declinatus.
tragacanthoides, Linn. Sp. Pl. 762 = longiflorus.
tragacanthoides, Willd. Sp. Pl. iii. 1323 = lanigerus.
tragacanthoides, Pall. Reise, i. 148 = diffusus.
tragacanthus, Lam. Fl. Fr. ii. 642 = A. Tragacantha.
tragoides, Linn. Syst. ed. X. 1175 = longiflorus.
transoxanus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 338.—Turkestan.
transsilvanicus, *Janka, in Termész. Füzetek*, viii. (1884) 307.—Europ.
Trautvetteri, *Bunge, Astrag.* i. 39; ii. 54.—Soongaria.
triangularis, *DC. Prod.* ii. 291.—Hab. ?
tribulifolius, *Benth. ex Bunge, Astrag.* i. 4; ii. 2.—Tibet. occ.
tribuliformis, Hochst. ex Boiss. Fl. Orient. iii. 225 = tribuloides.
tribuloides, *Delile, Fl. Aegypt. Illustr.* 70.—Afr. bor.; Oriens.
tribuloides, Kotschy, ex Bunge, Astrag. ii. 10 = ammophilus.
tricarinatus, *A. Gray, in Proc. Am. Acad.* xii. (1877) 56.—Am. bor.
trichocalyx, Nutt. ex Torr. & Gray, Fl. N. Am. i. 332 = mexicanus.
trichocalyx, *Trautv. in Act. Hort. Petrop.* iv. (1876) 362.—Reg. Caucas.
trichocarpus, *R. Grah. in Wall. Cat.* n. 5926.—Reg. Himal.
trichocarpus, Young, Fl. Texas, 228 = Nuttallianus.
trichobolus, *DC. Prod.* ii. 299.—Persia.
trichopodus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 218.—Am. bor. occ.
trichopterus, *Boiss. Fl. Orient.* ii. 292.—Syria.
trichostigma, *Bunge, Astrag.* i. 40; ii. 57.—Anatolia.
tricolor, *Bunge, l. c.* i. 139; ii. 237.—Soongar.
tridactylicus, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 527.—Am. bor. occ.
triflorus, *A. Gray, Pl. Wright.* ii. 45.—Am. bor. occ.
trifoliatus, *Phil. in Linnaea*, xxviii. (1856) 681.—Chili.
trifoliolatus, *Boiss. Diagn.* Ser. I. ii. 48.—Syria.
trigonelloides, *Boiss. l. c.* 49.—Armenia.
trigonocarpus, *Bunge, Astrag.* i. 22; ii. 23.—As. centr.
trigonus, *DC. Astrag.* 186.—Aegypt.; Arab.
trigonus, Sieber, ex DC. Prod. ii. 295 = Sieberi.
trimestris, Bieb. Fl. Taur. Cauc. ii. 190 = brachyceras.
trimestris, *Linn. Sp. Pl.* 761.—Aegypt.
trimestris, Pall. ex Ledeb. Fl. Ross. i. 368 = hamosus.
trimorphus, Viv. Fl. Lyb. Spec. 44. t. 10 = annularis.
triphyllus, Pall. Astrag. 68. t. 56. f. a = Oxytropis triphylla.
triphyllus, *Pursh, Fl. Am. Sept.* ii. 740.—Am. bor.
triquetrus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 367.—Am. bor.
triradiatus, *Bunge, Astrag.* i. 10; ii. 7.—Cilicia; Mesopotam.
tristis, Nutt. in Torr. & Gray, Fl. N. Am. i. 336 = Mortoni.
troglodytus, *S. Wats. in Proc. Am. Acad.* xx. (1885) 362.—Arizona.
trojanus, *Stev. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 401.—Graecia; As. Min.; Oriens.
tschujensis, *Bunge, Astrag.* i. 22; ii. 24.—Sibir. altaic.
tuberculosus, *DC. Astrag.* 33. t. 22.—Syria; Mesopotam.
tumidus, Bieb. Fl. Taur. Cauc. ii. 198 = fabaccus.
tumidus, DC. Astrag. 165 = Russelii.
tumidus, Willd. in Mém. Acad. Berl. (1794-95) 26 = Clusii.
tunetanus, *Willd. Sp. Pl.* iii. 1277.—Algeria.
turbinatus, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 251.—Persia; Turkestan.
turcomanicus, *Bunge, Astrag.* i. 130; ii. 227.—Tureom.
Turczaninowii, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 342.—Turkestan.
turkestanus, *Bunge, ex Boiss. Fl. Orient.* ii. 278.—Turkestan.
tymphresteus, *Boiss. & Sprun. Diagn.* Ser. I. ii. 63.—Graecia.

ASTRAGALUS:—
tyrolensis, Sieber, ex Steud. Nom. ed. II. i. 164 = Oxytropis campestris.
ulachanensis, *Franch. Pl. Davidian.* i. 87.—Mongolia.
uliginosus, Bieb. Fl. Taur. Cauc. ii. 188 = glycyphylloides.
uliginosus, *Linn. Sp. Pl.* 757.—Sibiria.
umbellatus, *Bunge, Astrag.* i. 24; ii. 29.—N. Zemlaia.
umbellatus, Pers. Syn. ii. 335 = lotoides.
uncatus, *Linn. Sp. Pl.* ed. II. 1072.—Aleppo.
uncatus, Linn. Herb. ex Boiss. Fl. Orient. ii. 234 = trimestris.
uncinatus, *Bertol. Fl. Ital.* viii. 54.—Ins. Corsica.
uncinatus, Moench, Meth. 166 = baeticus.
uncinatus, *Pomel, Nouv. Mat. Fl. Atl.* 322.—Algeria.
uniflorus, *DC. Astrag.* 243. t. 50.—Peruv.
unifoliatus, *Bunge, in Arb. Naturf. Ver. Riga*, i. (1847) 231.—Turkestan.
unifultus, *L'Hérit. Stirp. Nov.* 168.—Peruv.; Chili.
unijugus, *Bunge, Astrag.* i. 130; ii. 228.—Soongaria.
unilateralis, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 404.—Sibir. altaic.
uralensis, Linn. Fl. Suec. 257 = Oxytropis campestris.
uralensis, Vahl, in Fl. Dan. t. 1041 = Oxytropis sordida.
uraniolimneus, *Boiss. Fl. Orient.* ii. 380.—Armenia.
urmiensis, *Bunge, Astrag.* i. 37; ii. 44.—Persia.
ursinus, *A. Gray, in Proc. Am. Acad.* xiii. (1878) 367.—Am. bor.
ustiurtensis, *Bunge, Astrag.* i. 102; ii. 179.—Reg. Casp.
utahensis, *Torr. & Gray, in Pacif. Rail. Rep.* ii. (1854) 120.—Am. bor. occ.
utriger, Friv. ex Bunge, Astrag. ii. 108 = physocalyx.
utriger, Boiss. & Buhse, ex Boiss. Fl. Orient. ii. 281 = maeropelmatus.
utriger, *Pall. Astrag.* 75. t. 61.—Tauria.
utriger, C. A. Mey. Verz. Pfl. Cauc. 143 = bakuensis.
utriger, Willd. ex Ledeb. Fl. Ross. i. 654 = pubiflorus.
vaccarum, *A. Gray, Pl. Wright.* ii. 43.—Am. bor. occ.
vaginans, *DC. Astrag.* 210. t. 37.—Cilicia.
vaginatus, *Pall. Astrag.* 46. t. 36.—Sibiria.
vaginatus, Richards. in Frankl. 1st Journ. App. 746 = Hallii.
valesiacus, Thomas, ex Koch, Syn. Fl. Germ. ed. I. 182 = Oxytropis lapponica.
Vanillae, *Boiss. Diagn.* Ser. I. ii. 60.—Persia.
variabilis, *Bunge, ex Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 33.—Mongolia.
variegatus, *Franch. in Ann. Sc. Nat.* Sér. VI. xv. (1883) 259.—Turkest.
varius, S. G. Gmel. It. i. 116. t. 17 = virgatus.
Vaseyi, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 370.—Am. bor. occ.
vegetus, *Bunge, Astrag.* i. 102; ii. 181.—Persia.
vegliensis, Sadl. Fl. Pest. i. 199 in nota = Muelleri.
veluchensis, *Boiss. Fl. Orient.* ii. 319.—Graecia.
velutinus, Sieber, ex Steud. Nom. ed. II. i. 164 = Oxytropis Halleri.
venosus, Aucher, ex Bunge, Astrag. ii. 217 = Aucheri.
venosus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 193.—Abyss.
venosus, Ehrenb. ex Bunge, Astrag. ii. 229 = angustifolius.
ventorum, *A. Gray, ex S. Wats. in Am. Natural.* viii. (1874) 212.—Am. bor. occ.
venulosus, *Boiss. Diagn.* Ser. I. ii. 75.—Cappadoc.
vernus, *Georgi, Bemerk. Reise Russ. Reich.* i. 226.—Rossia.
verrucosus, *Moris, Stirp. Sard. Elench.* 12.—Sardinia.
versicolor, C. Koch, in Linnaea, xix. (1847) 65 = Sibthorpianus.
versicolor, *Pall. Astrag.* 45. t. 35.—Sibiria or.
verticillaris, *Bunge, Astrag.* i. 53; ii. 88.—Afghanist.
verticillaris, Linn. Mant. ii. 275 = Oxytropis myriophylla.
verticillaris, Georgi, Beschr. Russ. Reich. iii. IV. 1185 = Oxytropis verticillaris.
verticillaris, *Scop. Delic. Insub.* ii. 107.—Sibiria.
verus, DC. ex Bunge, Astrag. ii. 164 = Roussaeanus.
verus, *Olivier, Voy.* iii. t. 44.—Persia.
vesicarius, Lam. Fl. Fr. ii. 637 = A. Cicer.
vesicarius, *Linn. Sp. Pl.* 760.—Europ. austr.; Tauria.
vesicarius, Pall. Reise, iii. App. 751 = Oxytropis caespitosa.
vesicarius, Sicv. in Pall. N. Nord. Beitr. vii. 274 = inflatus.
vesiculosus, *Clos, in C. Gay, Fl. Chil.* ii. 120.—Chili.

ASTRAGALUS:—
vesiculosus, Jacquem. ex Baker, in Hook. f. Fl. Brit. Ind. iii. 138 = mollis.
vestitus, *Boiss. & Heldr. Diagn.* Ser. I. ix. 98.—Graecia.
vestitus, *S. Wats. Bibliog. Ind.* 202.—Calif.
vexillaris, *Boiss. Diagn.* Ser. I. ii. 49.—Mesopotam.
viciaefolius, *DC. Astrag.* 119.—Reg. Caucas.; As. Min.
vicioides, *R. Grah. in Wall. Cat.* n. 5931.—Reg. Himal.
vicioides, Ledeb. Fl. Alt. iii. 301 = macropterus.
villosissimus, *Bunge, Astrag.* i. 129; ii. 225.—Turkestan.
villosus, Gueldenst. ex Ledeb. Fl. Ross. i. 654 = pubiflorus.
villosus, *Michx. Fl. Bor. Am.* ii. 67.—Am. bor.
vimineus, Pall. Astrag. 24. t. 21 = frutieosus.
virescens, Ait. Hort. Kew. iii. 73 = falcatus.
virgatus, *Pall. Astrag.* 20. t. 18.—Hung.; Taur.
virgatus, Reichb. Fl. Germ. Exeurs. 510, 868 = Muelleri.
viridis, *Bunge, Astrag.* i. 135; ii. 231.—Armenia.
viscosus, Chaix, ex Vill. Prosp. 42 = Oxytropis campestris.
Volekmanni, *Phil. in Anal. Univ. Chil.* (1861) I. 46.—Chili.
Vulnerariae, *DC. Prod.* ii. 300.—As. Min.
vulnerarioides, All. Fl. Pedem. i. 343 = Anthyllis Vulneraria.
vulpinus, Ledeb. Fl. Alt. iii. 318 = lagocephalus.
vulpinus, *Willd. Sp. Pl.* iii. 1259.—Turkestan.
Wagneri, *Bartl. ex Bunge, Astrag.* ii. 125.—Persia.
Wardii, *A. Gray, in Proc. Am. Acad.* xii. (1877) 55.—Utah.
urtoënsis, *Boiss. & Kotschy, ex Bunge, Astrag.* i. 87; ii. 153.—Armenia.
Webberi, *A. Gray, in Bot. Calif.* i. 154.—Calif.
Webbianus, *R. Grah. in Wall. Cat.* n. 5936.—Reg. Himal.
Weidmannianus, *Fisch. in Bull. Soc. Nat. Mosc.* xxvi. (1853) II. 395.—Anatolia.
Whitneyi, *R. Gray, in Proc. Am. Acad.* vi. (1864–65) 526.—Calif.
Wilhelmsii, *Fisch. ex Ledeb. Fl. Ross.* i. 603 = resupinatus.
wingatanus, *S. Wats. in Proc. Am. Acad.* xviii. 1882–83) 192.—Am. bor. occ.
wolgensis, *Bunge, Astrag.* i. 36; ii. 47.—Rossia.
Wrightii, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 176.—Texas.
Wulfeni, *Koch, Syn. Fl. Germ.* ed. II. 207.—Dalmatia; Graecia.
xanthomelas, *Bunge, Astrag. Fedsch.* 314.—As. temp.
xanthotrichos, *Ledeb. Fl. Alt.* iii. 324.—Sibir. altaic.
xerophilus, *Ledeb. Fl. Ross.* i. 607.—Mont. Ararat.
Xiphidium, *Bunge, Astrag.* i. 25; ii. 218.—Reg. Caucas.
xiphocarpus, *Benth. ex Bunge, l. c.* i. 27; ii. 32.—Reg. Himal.
xylorrhizus, *Bunge, Astrag. Fedsch.* 267.—As. temp.
yuralicus, *Boiss. Fl. Orient.* ii. 366.—Cappadocia.
zacharensis, *Bunge, Astrag.* i. 23; ii. 27.—Syria.
zachlensis, *Bunge, l. c.* i. 88; ii. 154.—Syria.
zagrosicus, *Boiss. & Haussk. ex Boiss Fl. Orient.* ii. 286.—Persia.
zanskarensis, *Benth. ex Bunge, Astrag.* i. 43; ii. 67.—Reg. Himal.
zerdanus, *Boiss. Diagn.* Ser. I. ii. 51.—Persia.
zeylanicus, Ledeb. ex Bunge, Astrag. ii. 184 = cacruleus.
Zohrabi, Bunge, l. c. i. 68; ii. 112 = lagopoides.

ASTRANTHIUM, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 312 = **Bellis**, Linn. Compos. .
integrifolia, Nutt. l. c. = B. integrifolia.

ASTRANTHUS, Lour. Fl. Cochinch. 221 (1790) = **Homalium**, Jacq. (Samyd.).
cochinchinensis, Lour. l. c. 222 = H. fagifolium.
foetida, Wall. ex Mast. in Hook. f. Fl. Brit. Ind. ii. 598 = H. foetidum.

ASTRANTIA, [Tourn.] Linn. Syst. ed. I. (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 879.
ACTINOLEMA, Fenzl, Pugill. Pl. Syr. 66. t. 12 (1842).
alpina, Stur, ex Nym. Consp. 318 = gracilis.
alpina, [Clairv.] Man. Herb. 78 = minor.
bavarica, Schult. in Flora, xli. (1858) 161 = gracilis.

ASTRANTIA:—
Biebersteinii, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 28.—Reg. Caucas.
candida, Mill. Gard. Dict. ed. VIII. n. 2 = major.
carinthiaca, Hoppe, ex Koch, Syn. Fl. Germ. ed. I. 281 = major.
carniolica, *Wulf. in Jacq. Fl. Austr.* v. 31.—Carinth.
carniolica, Koch, Syn. Fl. Germ. ed. I. 280 = gracilis.
caucasica, Willd. ex Spreng. Umb. 17 = major.
ciliaris, *Linn. f. Suppl.* 177.—Afr. austr.
croatica, *Tomm. ex Vukot. in Rad. Jugos. Akad. Zagrab*, xliv. (1878) 216.—Croatia.
Diapensia, Scop. Fl. Carn. ed. II. i. 186 = Sanicula europaea.
digitata, Moench, Meth. 94 = minor.
dondiaefolia, Schur, Enum. Pl. Transs. 245 = gracilis.
elatior, Frivald. in Flora, xix. (1836) 434 = major.
Epipactis, Scop. Fl. Carn. ed. II. i. 185 = Hacquetia Epipactis.
europaea, Linn. ex Rehm. Notiz. Veg. Gest. Schwarz. Meer. 63 = major?
gracilis, *Bartl. in Linnaea*, xv. (1841) *Litt.* 93.—Carniolia.
gracilis, Schur, in Oestr. Bot. Zeitschr. (1858) 22 = carniolica.
grandiflora, Poepp. ex DC. Prod. iv. 668 (= *Pozoa coriacea*).—Reg. Andin.
helleborifolia, *Salisb. Prod.* 159.—Reg. Caucas.
heterophylla, Willd. in Ges. Naturf. Fr. Berl. Neue Schr. iii. (1801) 419 = helleborifolia.
intermedia, Bieb. Fl. Taur. Cauc. Suppl. 194 = major.
major, Bieb. l. c. n. 509 = Biebersteinii.
major, *Linn. Sp. Pl.* 235.—Reg. Caucas.
maxima, Pall. in Nov. Act. Petrop. vii. (1793) 357. t. 11 = helleborifolia.
microphylla, Schur, Enum. Pl. Transs. 245 = gracilis.
minor, *Linn. Sp. Pl.* 235.—Europ.
minor, Scop. Fl. Carn. ed. II. i. 187 = carniolica.
minor, Vitm. Summa Veg. ii. 124 = pauciflora.
montana, [Clairv.] Man. Herb. 78 = major.
neglecta, *C. Koch & Bouché, in App. Sem. Hort. Berol.* (1853) 22.—Europ.
nigra, Scop. Fl. Carn. ed. II. i. 188 = major.
pallida, Presl, ex Koch, Syn. Fl. Germ. ed. I. 280; ed. II. 309 = major.
pauciflora, *Bertol. in Desv. Journ. Bot.* ii. (1813) 76.—Italia.
ranunculifolia, *Reichb. f. Ic. Fl. Germ.* xxi. 3.—Europ.
saniculaefolia, Salisb. Prod. 159 = major.
speciosa, Hort. ex Heynh. Nom. ii. 53 = helleborifolia.
transsilvanica, Schur, Enum. Pl. Transs. 245 = carniolica.
tridentata, *Parrot, Reise Krym Kauk.* ii. 130, nomen.—Reg. Caucas.
trifida, Hoffm. Umb. ed. II. 1. praef. 8 in nota = major.

ASTRAPAEA, Lindl. Coll. Bot. t. 14 (1822) = **Dombeya**, Cav. (Stercul.).
penduliflora, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 90 = D. Wallichii.
tiliaefolia, Sweet, Hort. Brit. ed. I. 58 = D. acutangula.
viscosa, Sweet, l. c. = D. cannabina.
Wallichii, Lindl. Coll. Bot. t. 14 = D. Wallichii.

ASTREBLA, F. Muell. [Fragm. x. 76 (1876), nomen] ex Benth. Fl. Austral. vii. 602 (1878). *GRAMINEAE*, Benth. & Hook. f. iii. 1170.
elymoides, *F. Muell. ex F. M. Bailey, Grass. Queensl.* i.—Austral.
pectinata, *F. Muell. ex Benth. Fl. Austral.* vii. 602.—Austral.
triticoides, *F. Muell. l. c.*—Austral.

ASTREPHIA, Dufr. Hist. Valer. 50 (1811), pro parte. *VALERIANEAE*, Benth. & Hook. f. ii. 153.
Oligacoce, Willd. ex DC. Prod. iv. 632 (1830).
carnosa, Dufr. l. c. 51 = Valeriana carnosa.
chaerophylloides, *DC. Prod.* iv. 629.—Peruv.
chinensis, *Dufr. Hist. Valer.* 51.—China.
coarctata, Dufr. l. c. = Valeriana coarctata.
crispa, *Dufr. l. c.*—Chili.
hyalinorrhiza, Dufr. l. c. 52 = Valeriana hyalinorrhiza.
interrupta, Dufr. l. c. 41 = Valeriana interrupta.
laciniata, Dufr. l. c. 56 = chaerophylloides.
laxa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 364.—Chili.
lobata, *Hook. & Arn. l. c.*—Chili.

ASTREPHIA:—
mexicana, Hook. & Arn. Bot. Beech. Voy. Suppl. 431 = Phyllactis mexicana.
mexicana, Seem. Bot. Voy. Herald, 297 = Valeriana mexicana.
oblongifolia, Dufr. Hist. Valer. 60 = Valeriana oblongifolia.
polystachya, Dufr. l. c. 51 = Valerianopsis polystachya.
pratensis, Benth. Pl. Hartw. 39 = Phyllactis pratensis.
serrata, Dufr. Hist. Valer. 50 = Valeriana serrata.

ASTROCARPUS, Neck. Elem. ii. 243 (1790). *RESEDACEAE*, Benth. & Hook. f. i. 111.
Sesamella, Reichb. Consp. 186 (1828).
abyssinicus, Hochst. ex Oliver, Fl. Trop. Afr. i. 103 = Caylusea abyssinica.
Clusii, *J. Gay, in F. Schultz, Arch. Fl. Fr. et Allem.* 33.—Hispan.
cochlearifolius, *Nym. in Ofver. Vet. Akad. Stockh.* xviii. (1861) 191.—Lusitan.
interruptus, Bor. in Mém. Soc. Acad. Maine-et-Loire, viii. (1860) 7 = sesamoides.
minor, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 85 (*Pugill.* 280. t. 4. f. 1).—Europ.
purpurascens, Rafin. Fl. Tellur. iii. 73 = Clusii.
sesamoides, Colm. ex Willk. & Lange, Prod. Fl. Hisp. iii. 899 = Clusii.
sesamoides, *DC. ex Duby, Bot. Gall.* i. 67.—Europ.
spathulaefolius, *Revell, ex Bor. in Mém. Soc. Acad. Maine-et-Loire*, i. (1857) 87.—Gallia.
suffruticosus, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 83 (*Pugill.* 278. t. 4).—Hispan.

ASTROCARYUM, G. F. W. Mey. Prim. Fl. Esseq. 265 (1818). *PALMAE*, Benth. & Hook. f. iii. 942.
Toxophoenix, Schott, Nachr. K. K. Oester. Naturf. in Brasil. Anh. ii. (1822).
acanthopodium, *Barb.-Rodr. Enum. Palm. Nov.* 20.—Bras.
acaule, *Mart. Hist. Nat. Palm.* ii. 78. t. 24. 63. f. 5.—Bras.
aculeatum, Barb.-Rodr. Enum. Palm. Nov. 20 = Rodriguesii.
aculeatum, *G. F. W. Mey. Prim. Fl. Esseq.* 266.—Guiana.
aculeatum, Hort. ex H. Wendl. in Kerch. Palm. 230 = Acrocomia sclerocarpa.
aculeatum, Wallace, Palm Trees Amaz. 111. t. 43 = minus?
argentenm, *Hort. Bull.—Cf. Gard. Chron.* (1875) i. 491.—Hab.?
aureum, *Griseb. & H. Wendl. in Griseb. Fl. Brit. W. Ind.* 521.—Ind. occ.
Awarra, De Vriese, in Jaarb. Maatsch. (1848) 12 = aculeatum.
Ayri, *Mart. Hist. Nat. Palm.* ii. 71. t. 59ª.—Bras.
Borsigianum, C. Koch, ex Regel, Gartenfl. (1861) 29 = Stevensonia grandifolia.
campestre, *Mart. Hist. Nat. Palm.* ii. 79. t. 63. f. 4.—Bras.
candescens, *Barb.-Rodr. Enum. Palm. Nov.* 22.—Bras.
Chichon, Linden, Illustr. Hortic. xxviii. (1881) 15 = mexicanum.
Chonta, *Mart. in Palmet. Orb.* 84. t. 4. f. 1. 2.—Peruv.
farinosum, *Barb.-Rodr. Enum. Palm. Nov.* 21.—Bras.
flexuosum, *Hort. ex H. Wendl. in Kerch. Palm.* 232.—Hab.?
guianense, *Splitg. ex Mart. Hist. Nat. Palm.* iii. 323.—Guiana.
gynacanthum, *Mart. l. c.* ii. 71. t. 60, 61.—Bras.
gynacanthum, Wallace, Palm Trees Amaz. 103. t. 39 = A. Munbaca?
Huaimi, *Mart. in Palmet. Orb.* 86. t. 13. fig. 3.—Bolivia.
humile, *Wallace, Palm Trees Amaz.* 115. t. 45.—Bras.
Jauari, *Mart. Hist. Nat. Palm.* ii. 76. t. 52 et 65. f. 1.—Bras.
Jucuma, Linden, Illustr. Hortic. xxviii. (1881) 15, sphalm. = A. Tucuma.
Malybo, *Karst. in Linnaea*, xxviii. (1856) 245.—N. Granat.
mexicanum, *Liebm. ex Mart. Hist. Nat. Palm.* iii. 323.—Mexic.

ASTROCARYUM:—
minus, *Trail, in Journ. Bot.* xv. (1877) 78.—Bras.
Munbaca, *Mart. Hist. Nat. Palm.* ii. 74.—Bras.
Murumura, *Mart. l. c.* 70. t. 58, 59.—Bras.
Murumuru, *Wallace, Palm Trees Amaz.* 101. t. 38.—Bras.
panamense, *Linden, Illustr. Hortic.* xxxviii. (1881) 15.—N. Granat.
Paramaca, *Mart. in Palmet. Orb.* 88.—Guiana.
plicatum, *Drude, in Mart. Fl. Bras.* iii. II. 375.—Guiana; Bras.
Princeps, *Barb.-Rodr. Enum. Palm. Nov.* 22.—Bras.
pygmaeum, *Drude, in Mart. Fl. Bras.* iii. II. 385.—Bras.
pumilum, *Hort. ex H. Wendl. in Kerch. Palm.* 232.—Hab.?
Rodriguesii, *Trail, in Journ. Bot.* xv. (1877) 79.—Bras.
rostratum, *Hook. Bot. Mag.* t. 4773.—Bras.
sclerocarpum, Hort. ex H. Wendl. in Kerch. Palm. 232 = Acrocomia sclerocarpa.
sclerophyllum, *Drude, in Mart. Fl. Bras.* iii. II. 377.—Bras.
segregatum, *Drude, l. c.* 382.—Bras.
tenuifolium, *Linden, Illustr. Hortic.* xxviii. (1881) 15.—Bras.
Tucuma, *Mart. Hist. Nat. Palm.* ii. 77. t. 65. f. 2.—Bras.
tucumoides, *Drude, in Mart. Fl. Bras.* iii. II. 381.—Bras.
vulgare, *Mart. Hist. Nat. Palm.* ii. 74. t. 62, 63.—Bras.
Warszewiczii, *Karst. in C. Koch & Fint. Wochenschr.* (1858) 297.—Am. austr.
Weddellii, *Drude, in Mart. Fl. Bras.* iii. II. 383.—Bras.

ASTROCEPHALUS, Rafin. Fl. Tellur. iv. 94 (1836) = ASTEROCEPHALUS, Vaill. = **Scabiosa**, Tourn. (Dipsac.).
ciliata, integrifolia, rigida, sylvatica, tatarica, uralensis, Rafin. l. c. = Scabiosae totidem homonymae.

ASTROCOCCUS, Benth. in Hook. Kew Journ. vi. (1854) 327. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 326.
coriaceus, *Baill. in Adansonia*, v. 308.—Bras.
cornutus, *Benth. in Hook. Kew Journ.* vi. (1854) 327.—Bras.

ASTROCOMA, Neck. Elem. i. 112 (1790) = **Staavia**, Dahl (Bruniac.).

ASTRODENDRUM, Dennst. Schluess. Hort. Malab. 30 (1818) = **Sterculia**, Linn.
malabaricum, Dennst. l. c. = S. guttata.

ASTROGLOSSUS, Reichb. f. ex Benth. & Hook. f. Gen. iii. 588 (1883) = **Trichoceros**, H. B. & K. (Orchid.).

ASTROGYNE, Benth. Pl. Hartw. 14 (1839) = **Croton**, Linn. (Euphorb.).
crotonoides, Benth. l. c. = C. gracilis.

ASTROGYNE, Wall. ex Laws. in Hook. f. Fl. Brit. Ind. i. (1875) 629 = **Siphonodon**, Griff. (Celastr.).
coriacea, Wall. ex Laws. l. c. = S. celastrineus.

ASTROLOBIUM, DC. Prod. ii. 311 (1825) = **Ornithopus**, Linn. et **Coronilla**, Linn. (Legumin.).
creticum, DC. l. c. 310 = C. cretica.
durum, DC. l. c. 311 = O. durus.
ebracteatum, DC. l. c. = O. ebracteatum.
repandum, DC. l. c. = C. repanda.
scorpioides, DC. l. c. = C. scorpioides.

ASTROLOMA, R. Br. Prod. 538 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 611.
Mesotricha, Stschegl. in Bull. Soc. Nat. Mosc. (1859) i. 6.
Pentataphrus, Schlecht. in Linnaea, xx. (1847) 618.
Stenanthera, R. Br. Prod. 538 (1810).
Stomarrhena, DC. Prod. vii. 738 (1838).
Ventenatia, Cav. Ic. iv. 79 (1797).
Vintenatia, Cav. l. c. 28 (1797).
Baxteri, *A. Cunn. ex DC. Prod.* vii. 739.—Austral.
Candolleanum, *Sond. in Lehm. Pl. Preiss.* i. 302.—Austral.
compactum, *R. Br. Prod.* 538.—Austral.

ASTROLOMA:—
conostephioides, *F. Muell. ex Benth. Fl. Austral.* iv. 158.—Austral.
cuneifolium, Sond. in Lehm. Pl. Preiss. i. 299 = compactum.
denticulatum, R. Br. Prod. 538 = humifusum.
dilatatum, Sond. in Lehm. Pl. Preiss. i. 298 = longiflorum.
discolor, Sond. l. c. = longiflorum.
divaricatum, *Sond. l. c.* 299.—Austral.
Drummondii, *Sond. l. c.*—Austral.
foliosum, *Sond. l. c.* 297.—Austral.
glaucescens, Sond. l. c. 298 = macrocalyx.
hirsutum, Stschegl. in Bull. Soc. Nat. Mosc. (1859) i. 7 = Drummondii.
humifusum, *R. Br. Prod.* 538.—Austral.
juniperinum, F. Muell. Fragm. vi. 36 = microphyllum.
lasionemum, F. Muell. l. c. 40 = stomarrhena.
latifolium, Sond. in Lehm. Pl. Preiss. i. 302 = tectum.
longiflorum, *Sond. l. c.* 297.—Austral.
macrocalyx, *Sond. l. c.* 301.—Austral.
marginatum, Sond. l. c. 299 = divaricatum.
microcalyx, *Sond. l. c.* 298.—Austral.
microdonta, *F. Muell. ex Benth. Fl. Austral.* iv. 155.—Austral.
microphyllum, *Stschegl. in Bull. Soc. Nat. Mosc.* (1859) i. 7.—Austral.
pallidum, *R. Br. Prod.* 538.—Austral.
pallidum, Sond. in Linnaea, xxvi. (1853–55) 246 = humifusum.
pinifolium, *Benth. Fl. Austral.* iv. 159.—Austral.
prostratum, *R. Br. Prod.* 538.—Austral.
pungens, Stschegl. in Bull. Soc. Nat. Mosc. (1859) i. 8 = divaricatum.
splendens, Planch. in Fl. des Serres, Sér. I. x. (1854–55) 129. t. 1018 = divaricatum.
stomarrhena, *Sond. in Lehm. Pl. Preiss.* i. 301.—Austral.
tectum, *R. Br. Prod.* 538.—Austral.
xerophyllum, *Sond. in Lehm. Pl. Preiss.* i. 301.—Austral.

ASTRONIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1; Blume, Bijdr. 1080 (1826). *MELASTOMACEAE*, Benth. & Hook. f. i. 771.
Astronidium, A. Gray, Bot. U. St. Expl. Exped. i. 582. t. 72 (1854).
Lomanodia, Rafin. Sylva Tellur. 97 (1838).
Naudinia, Decne. ex Triana, in Seem. Fl. Vit. 86 (1866).
concolor, Zipp. ex Steud. Nom. ed. II. i. 165 = macrophylla.
confertiflora, *A. Gray, Bot. U. St. Expl. Exped.* i. 579.—Ins. Fiji.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 174.—Ins. Philipp.
Forsteri, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 258.—Ins. Tahiti.
fraterna, *A. Gray, Bot. U. St. Expl. Exped.* i. 576. t. 72.—Ins. Tahiti.
fraterna, Seem. Fl. Vit. 85, 86 = Forsteri.
glabra, Triana, in Trans. Linn. Soc. xxviii. (1871) 152 = Forsteri.
intermedia, Blume, Mus. Bot. Lugd. Bat. i. 9 = spectabilis.
macrophylla, *Blume, Bijdr.* 1080.—Malaya.
ovalifolia, *Naud. ex Decne. Bot. Voy. Vénus*, 27.—Ins. Tahiti.
papetaria, *Blume, Bijdr.* 267.—Ins. Molucc.
parviflora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 152.—Ins. Fiji.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 577. t. 72.—Ins. Fiji.
pileolata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 152.—Ins. Tahiti.
robusta, *Seem. Fl. Vit.* 86.—Ins. Fiji.
Rolfei, *Vidal, Phan. Cuming. Philipp.* 174.—Ins. Philipp.
samoënsis, *S. Moore, in Journ. Bot.* xviii. (1880) 3.—Ins. Samoa.
smilacifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 152.—Malaya.
spectabilis, *Blume, Bijdr.* 1080.—Java.
spectabilis, Zipp. ex Steud. Nom. ed. II. i. 165 = papetaria.
Storckii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 152.—Ins. Fiji.

ASTRONIA:—
subcordata, *A. Gray, Bot. U. St. Expl. Exped.* i. 580.—Ins. Samoa.
tomentosa, *Seem. Fl. Vit.* 86.—Ins. Fiji.

ASTRONIDIUM, A. Gray, Bot. U. St. Expl. Exped. i. 581. t. 72 (1854) = **Astronia**, Blume (Melast.).
parviflorum, A. Gray, l. c. 582. t. 72 = Astronia parviflora.
Storckii, Seem. Fl. Vit. 87 = Astronia Storckii.

ASTRONIUM, Jacq. Enum. Pl. Carib. 10. *ANACARDIACEAE*, Benth. & Hook. f. i. 423.
Balansae, *Engl. in Bot. Jahrb.* i. (1881) 45.—Paraguay.
Candollei, *Engl. l. c.*—Paraguay.
concinnum, *Schott, in Spreng. Syst. Cur. Post.* 404.—Bras.
fraxinifolium, *Schott, in Spreng. l. c.*—Bras.
gracile, *Engl. Bot. Jahrb.* i. (1881) 75.—Bras.
graveolens, *Jacq. Enum. Pl. Carib.* 33.—N. Granat.
insigne, March. Rev. Anac. 177 = Parishia insignis.
juglandifolium, Griseb. in Goett. Abh. xxiv. (1879) 94 = A. Urundeuva.
Libonianum, *J. Linden, Cat.* n. 17. (1862) 7.—Bras.
macrocalyx, *Engl. Bot. Jahrb.* i. (1881) 75.—Bras.
obliquum, *Griseb. Fl. Brit. W. Ind.* 176.—Ind. occ.
Planchonianum, *Engl. in Mart. Fl. Bras.* xii. ii. 399.—Am. trop.
Urundeuva, *Engl. Bot. Jahrb.* i. (1881) 45.—Bras.

ASTROPANAX, Seem. in Journ. Bot. iii. 176 (1865) = **Heptapleurum**, Gaertn. (Araliac.).
abyssinicum, Seem. l. c. 177 = H. abyssinicum.
Baikiei, Seem. l. c. = H. Baikiei.
Barteri, Seem. l. c. = H. Barteri.
elatum, Seem. l. c. = H. elatum.
Manni, Seem. l. c. = H. Mannii.

ASTROPETALUM, Griff. Notul. iv. 411. t. 565 (1854) = **Swintonia**, Griff. (Anacard.).

ASTROPHEA, Reichb. Consp. 132 (1828) = **Passiflora**, Linn.
emarginata, M. Roem. Syn. Pepon. 152 = P. emarginata.
glaberrima, Klotzsch, in Schomb. Fauna & Fl. Guy. 1168, nomen.—Guiana.
glauca, M. Roem. Syn. Pepon. 151 = P. arborea.
ovata, M. Roem. l. c. = P. ovata.

ASTROPHIA, Nutt. in Torr. & Gray, Fl. N. Am. i. 278 (1838) = **Lathyrus**, Linn. (Legum.).
littoralis, Nutt. l. c. = L. littoralis.

ASTROPHYLLUM, Torr. & Gray, in Bot. Pope Exped. (Pacif. Rail. Rep. ii.) 161 (1855). *RUTACEAE*, Benth. & Hook. f. i. 296.
dumosum, *Torr. & Gray, l. c.*—Am. bor. occ.

ASTROPHYTUM, Lem. Cact. Gen. Nov. Gen. et Sp. 3 (1839) = **Echinocactus**, Link & Otto (Cact.).
Asterias, Lem. Cact. 50 = E. Asterias.
myriostigma, Lem. Cact. Gen. Nov. et Sp. 4 = E. myriostigma.
prismaticum, Lem. Cact. 50 = E. myriostigma.

ASTROPUS, Spreng. Neue Entdeck. iii. 64 (1822) = **Waltheria**, Linn. (Stercul.).
tomentosus, Spreng. l. c. = W. Astropus.

ASTROSCHOENUS, Lindl. Veg. Kingd. 119 (1847) = **Asteroschoenus**, Nees (Cyperac.).

ASTROSTEMMA, Benth. in Hook. Ic. Pl. xiv. 7. t. 1311 (1880). *ASCLEPIADEAE*.
spartioides, *Benth. l. c.*—Borneo.

ASTROTRICHA, DC. Coll. Mém. v. 29 t. 5 (1829); Prod. iv. 74 (1830). *ARALIACEAE*, Benth. & Hook. f. i. 937.
asperifolia, F. Muell. ex Klatt, in Linnaea, xxix. (1858) 709 = ledifolia.
floccosa, *DC. Coll. Mém.* v. 30. t. 5.—Austral.
Hamptoni, *F. Muell. Fragm.* vi. 125.—Austral.
hoveoides, *A. Cunn. ex Benth. Enum. Pl. Huegel.* 55.—Austral.
latifolia, *Benth. l. c.* 54.—Austral.

ASTROTRICHA:—
ledifolia, *DC. Coll. Mém.* v. 30. t. 6.—Austral.
linearis, *A. Cunn. ex Benth. Enum. Pl. Huegel.* 55.—Austral.
longifolia, *Benth. l. c.*—Austral.
pterocarpa, *Benth. Fl. Austral.* iii. 379.—Austral.

ASTYDAMIA, DC. Coll. Mém. v. 53 (1829); Prod. iv. 190 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 909.
canariensis, *DC. Prod.* iv. 190.—Ins. Teneriff.
latifolia, Baill. Hist. Pl. vii. 208 = canariensis.

ASTYLIS, Wight, Ic. t. 1992 (1853) = **Hemicyclia**, Wight & Arn. (Euphorb.).
venusta, Wight, l. c. = H. venusta.

ASTYLUS, Dulac, Fl. Hautes-Pyr. 188 (1867) = **Hutchinsia**, R. Br. (Crucif.).
alpinus, Dulac, l. c. = H. alpina.
petraea, Dulac, l. c. = H. petraea.

ASTYRIA, Lindl. Veg. Kingd. 364 (1847) = **Astiria**, Lindl. (Stercul.).

ASYNEUMA, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 335 = **Phyteuma**, Linn. (Campan.).
canescens, Griseb. & Schenk, l. c. = P. canescens.

ASYSTASIA, Blume, Bijdr. 796 (1826). *ACANTHACEAE*, Benth & Hook. f. ii. 1094.
Dicentranthera, T. Anders. in Journ. Linn. Soc. vii. (1863) 52.
Henfreya, Lindl. Bot. Reg. (1847) t. 31.
Intrusaria, Rafin. Fl. Tellur. iv. 66 (1836).
Mackaya, Harv. Thes. Cap. i. 8. t. 13 (1859).
Ramusia, E. Mey. in Drège, Zwei Pfl. Docum. 215 (1843).
acuminata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 201.—Afr. trop.
africana, *Hort. Bogor. ex T. Anders. in Journ. Agri.-Hort. Soc. Ind.* N. S. i. (1868) 287.—Afr. trop.
atroviridis, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 526.—Reg. Himal.
Blumei, Nees, in DC. Prod. xi. 167 = intrusa.
Bojeriana, Nees, l. c. 166 = coromandeliana.
calycina, Benth. in Hook. Niger Fl. 478 = Strobilanthes Helictus.
capensis, Nees, in Linnaea, xv. (1841) 356 = coromandeliana.
Charmian, *S. Moore, in Journ. Bot.* xviii. (1880) 38.—Afr. trop.
chelonoides, *Nees, in Wall. Pl. As. Rar.* iii. 89.—Ind. or.
chinensis, *S. Moore, in Journ. Bot.* xiii. (1875) 228.—China.
comorensis, Boj. ex Nees, in DC. Prod. xi. 166 = coromandeliana.
coromandeliana, *Nees, in Wall. Pl. As. Rar.* iii. 89.—Ind. or.; Malaya; Afr. trop. et austr.
coromandeliana, Wight, Ic. t. 1506 = crispata.
crispata, *Benth. in Linnaea*, xxiv. (1851) 647.—Ind. or.
denticulata, Nees, in Wall. Pl. As. Rar. iii. 89 = Strobilanthes denticulatus.
floribunda, *Klotzsch, in Peters, Reise Mossamb. Bot.* 200.—Afr. trop.
formosa, Hort. Calc. ex T. Anders. in Journ. Agri.-Hort. Soc. Ind. N. S. i. (1868) 270 = Ruellia formosa.
gangetica, T. Anders. in Thw. Enum. Pl. Zeyl. 235 = coromandeliana.
intrusa, *Blume, Bijdr.* 796.—Malaya.
intrusa, Nees, in DC. Prod. xi. 166 = coromandeliana.
Kunthiana, *Nees, in Wall. Pl. As. Rar.* iii. 89.—Penang.
lanceolata, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 524.—Burma.
Lawiana, *Dalz. in Hook. Kew Journ.* iv. (1852) 344.—Ind. or.
macrocarpa, *Nees, in Wall. Pl. As. Rar.* iii. 89.—Reg. Himal.
multiflora, *Klotzsch, in Peters, Reise Mossamb. Bot.* 203.—Afr. trop.
mysurensis, T. Anders. in Journ. Linn. Soc. ix. (1867) 524 = Lawiana.
Neesiana, *Nees, in Wall. Pl. As. Rar.* iii. 89.—Ind. or.; Burma.
nemorum, Nees, l. c. 90 = intrusa.
Panichanga, Nees, l. c. = Strobilanthes Panichanga.

ASYSTASIA:—
Parishii, T. Anders. in Journ. Linn. Soc. ix. (1867) 526 = Eranthemum Parishii.
pauciflora, Nees, in Wall. Pl. As. Rar. iii. 90 = Codonacanthus pauciflorus.
plumbaginea, Nees, l. c. 89 = coromandeliana.
podostachys, *Klotzsch, in Peters, Reise Mossamb. Bot.* 199.—Afr. trop.
pubescens, *Klotzsch, l. c.* 202.—Afr. trop.
quarterna, *Nees, in DC. Prod.* xi. 166.—Afr. trop.
querimbensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 204.—Afr. trop.
rostrata, Solms, in Schweinf. Fl. Aethiop. 242 = Schimperi.
scabrida, *Klotzsch, in Peters, Reise Mossamb. Bot.* 202.—Afr. trop.
scandens, *Hook. Bot. Mag.* t. 4449.—Afr. trop.
Schimperi, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 40, 53.—Abyss.
subhastata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 200.—Afr. trop.
thyrsacanthus, T. Anders. in Journ. Linn. Soc. ix. (1867) 525 = Eranthemum indicum.
travancorica, *Bedd. Ic. Pl. Ind. Or.* i. 39.—Ind. or.
violacea, *Dalz. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 494.—Ind. or.
violacea, Dalz. in Hook. Kew Journ. ii. (1850) 139 = coromandeliana.
Vogeliana, *Benth. in Hook. Niger Fl.* 479.—Afr. trop.
Welwitschii, *S. Moore, in Journ. Bot.* xviii. (1880) 308. t. 213.—Afr. trop.

ATACCA, Lem. Jard. Fleur. ii. (1852) t. 186, 187 = seq.

ATACCIA, Presl, Rel. Haenk. 149 (1828) = **Tacca**, Forst.
aspera, Kunth, Enum. Pl. v. 464 = T. integrifolia.
cristata, Kunth, l. c. 466 = T. cristata.
integrifolia, Presl, Rel. Haenk. 149 = T. integrifolia.
laevis, Kunth, Enum. Pl. v. 466 = T. laevis.
lancaefolia, Kunth, l. c. 465 = T. lancaefolia.

ATADINUS, Rafin. Sylva Tellur. 30 (1838) = **Rhamnus**, Linn.
alpinus, Rafin. l. c. = R. alpinus.

ATAENIA, Endl. Gen. Suppl. v. 5 (1850) = ATENIA, Hook. & Arn. = **Carum**, Linn. (Umbell.).

ATALANTA, Nutt. Gen. Am. ii. 73 (1818) = **Cleome**, Linn. (Cappar.).
serrulata, Nutt. l. c. = C. serrulata.

ATALANTHUS, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 311 = **Sonchus**, Linn. (Compos.).
angustifolius, Pomel, Nouv. Mat. Fl. Atl. 8 (= *Zollikoferia angustifolia*).—Afr. bor.
arboreus, Sweet, Hort. Brit. ed. II. 273 = S. arboreus.
Candolleanus, Pomel, Nouv. Mat. Fl. Atl. 8 (= *Zollikoferia Candolleana*).—Afr. bor.
divaricatus, Pomel, l. c. 7 = S. divaricatus.
longilobus, Pomel, l. c. (= *Zollikoferia longiloba*).—Afr. bor.
nudicaulis, Pomel, l. c. = Launaea nudicaulis.
pinnatus, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 311 = S. leptocephalus.
quercifolius, Pomel, l. c. 8 = S. quercifolius.
resedifolius, Pomel, l. c. (= *Zollikoferia resedifolia*).—Afr. bor.
spinosus, D. Don, in Edinb. N. Phil. Journ. (Mar. 1829) 311 = Launaea spinosa.

ATALANTIA, Correa, in Ann. Mus. Par. vi. 383 (1805). *RUTACEAE*, Benth. & Hook. f. i. 305.
CHILOCALYX, Turcz. in Bull. Soc. Nat. Mosc. (1863) i. 588.
HELIE, M. Roem. Syn. Hesper. 42 (1846).
LAMPETIA, M. Roem. l. c. (1846).
MALNAREGAM, Adans. Fam. ii. 344 (1763).
MEROPE, M. Roem. Syn. Hesper. 44 (1846).
RISSOA, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 324.
SCLEROSTYLIS, Blume, Bijdr. 133 (1825).
SEVERINIA, Tenore, Ind. Sem. Hort. Neap. (1840).
bilocularis, Wall. Cat. n. 6356 = Limonia bilocularis.
buxifolia, *Oliver, in Benth. Fl. Hongk.* 51.—China.

ATALANTIA:—
buxifolia, Oliver, in Journ. Linn. Soc. v. Suppl. ii. (1861) 26 = Limonia bilocularis.
capitellata, Lindl. in Wall. Cat. n. 6355 = racemosa.
carissoides, Wall. l. c. n. 6354 = monophylla.
caudata, *Hook. f. Fl. Brit. Ind.* i. 513.—Reg. Himal.
ceylanica, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 25.—Ind. or.
citrifolia, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 69.—Burma.
floribunda, Wight, Ic. iv. t. 1611 = monophylla.
glauca, *Benth. & Hook. f. Gen.* i. 305.—Austral.
Hindsii, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 26.—China.
longispina, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 295.—Burma.
Loureiriana, *M. Roem. Syn. Hesper.* 37.—Cochinch.
macrophylla, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) ii. 136.—Ins. Nicobar.
missionis, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 25.—Ind. or.
monophylla, Benth. in Hook. Lond. Journ. Bot. i. (1842) 483 = Hindsii.
monophylla, *DC. Prod.* i. 535.—Ind. or.
monophylla, Hook. & Arn. Bot. Beech. Voy. 172 = buxifolia.
nitida, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 25.—Malaya.
parvifolia, M. Roem. Syn. Hesper. 37 = racemosa.
platystigma, Wight, Illustr. Ind. Bot. i. 108 = monophylla.
puberula, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 211.—Siam.
racemosa, *Wight & Arn. Prod.* 91; *et in Hook. Journ. Bot.* i. (1834) 64.—Ind. or.
recurva, *Benth. Fl. Austral.* i. 370.—Austral.
Roxburghiana, *Hook. f. Fl. Brit. Ind.* i. 513.—Malaya.
Roxburghii, Oliver, in Journ. Linn. Soc. v. Suppl. ii. 25 = caudata.
trimera, *Oliver, l. c.*—Malaya.
umbellata, M. Roem. Syn. Hesper. 37 = monophylla.

ATALAYA, Blume, Rumphia, iii. 186 (1847). *SAPINDACEAE*, Benth. & Hook. f. i. 401.
annularis, *Blume, Rumphia*, iii. 186.—Malaya.
australis, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 327 = Sapindus australis.
bijuga, Span. Ic. n. 19; in Linnaea, xv. (1841) 180 = Cupania salicifolia.
cochinchinensis, *Blume, Rumphia*, iii. 186.—Cochinch.
coriacea, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 326.—Ins. Dom. Howe.
hemiglauca, *F. Muell. ex Benth. Fl. Austral.* i. 463.—Austral.
multiflora, *Benth. l. c.*—Austral.
salicifolia, *Blume, Rumphia*, iii. 186.—Timor.; Austral.
variifolia, *F. Muell. ex Benth. Fl. Austral.* i. 463.—Austral.

ATAMASCO, Rafin. Neogenyt. 3 (1825) = **Amaryllis**, Linn.

ATAMISQUEA, Miers, Trav. Chili, ii. 529 (1826). *CAPPARIDEAE*, Benth. & Hook. f. i. 109.
emarginata, *Miers, l. c.*—Chili.

ATAMOSCO, Adans. Fam. ii. 57 = **Amaryllis**, Linn.

ATAMOSKO, Adans. Fam. ii. 523 = praec.

ATASITES, Neck. Elem. i. 7 (1790) = **Gerbera**, Gronov. (Compos.).

ATAXIA, R. Br. Chlor. Melv. in App. Parry, Voy. 292 (1824) = **Hierochloë**, S. G. Gmel. (Gramin.).
Ecklonis, Nees, ex Trin. in Mém. Acad. Pétersb. v. (1840) 77 = H. Ecklonii.
Hookeri, Griseb. in Goett. Nachr. (1868) 77.—Ind. or.
Horsfieldii, Kunth, Gram. i. 22; R. Br. in Benn. Pl. Jav. Rar. 8. t. 3.—Java.
mexicana, Rupr. in Bull. Acad. Brux. ix. (1842) 233 = H. mexicana.
tenuis, Trin. in Mém. Acad. Pétersb. v. (1840) 78.—Afr. austr.
Tongo, Nees, ex Trin. l. c. = H. Tongo.

ATE, Lindl. Gen. et Sp. Orch. 326 (1835) = **Habenaria**, Willd. (Orchid.).
acuminata, Thw. Enum. Pl. Zeyl. 309 = H. acuminata.
virens, Lindl. Gen. et Sp. Orch. 327 = H. barbata.

ATECOSA, Rafin. Fl. Tellur. iii. 46 (1836) = **Rumex**, Linn. (Polygon.).

ATELANDRA, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 289. t. 10. *MYRSINEAE*.
laurina, *Bello, l. c.*—Porto Rico.
obtusifolia, *Bello, l. c.*—Porto Rico.

ATELANDRA, Lindl. Swan Riv. App. 40. t. 5 (1839) = **Hemigenia**, R. Br. (Labiat.).
incana, Lindl. l. c. = H. incana.
polystachya, Lindl. l. c. 40 = H. incana.

ATELEA, A. Rich. Pl. Vasc. Cub. 454 (1845) = **Ateleia**, Moç. & Sessé (Legumin.).

ATELANTHERA, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 129. *CRUCIFERAE*, Benth. & Hook. f. i. 68.
perpusilla, *Hook. f. & Thoms. l. c.* 138.—Tibet. occ.

ATELEIA, Moç. & Sessé, ex DC. Mém. Lég. 395. t. 57 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 558.
apetala, *Griseb. Cat. Pl. Cub.* 80.—Cuba.
cubensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 180.—Cuba.
Glazioveana, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 306.—Bras.
gummifera, *D. Dietr. Syn. Pl.* iv. 1219.—Ins. S. Doming.
microcarpa, *D. Dietr. l. c.*—Am. austr.
peltaria, *D. Dietr. l. c.*—Afr. austr.
pterocarpa, *Moç. & Sessé, ex DC. Lég. Mém.* 394.—Mexic.

ATELESTE, Sond. in Linnaea, xxiii. (1850) 9, 10 = **Aberia**, Hochst. (Bixin.).
tristis, Sond. l. c. 9 = Aberia tristis.
Zeyheri, Sond. l. c. 10 = Aberia Zeyheri.

ATELIANTHUS, Nutt. ex Benth. in DC. Prod. x. 455, in nota (1846) = **Synthyris**, Benth. (Scrophul.).
veronicoides, Nutt. ex Benth. l. c. = S. plantaginea.

ATENIA, Hook. & Arn. Bot. Beech. Voy. 349 (1840) = **Carum**, Linn. (Umbell.).
Gairdneri, Hook. & Arn. l. c. = C. Gairdneri.

ATERAMNUS, P. Br. Hist. Jamaic. 339 (1756); 'nom. dclend.' Benth. & Hook. f. Gen. iii. 256.

ATHALMUS, Neck. Elem. i. 20 (1790) = **Pallenis**, Cass. (Compos.).

ATHAMANTA, Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 900.
KILLINGA, Adans. Fam. ii. 498 (1763).
LIBANOTIS, Hall, Enum. Stirp. Helv. ii. 450 (1742).
PETRO-CARVI, Tausch, in Flora, xvii. (1834) 355.
TRIBULA, Hill, Veg. Syst. vi. 10 (1764).
TURBITH, Tausch, in Flora, xvii. (1834) 343.
achilleifolia, Wall. Cat. n. 568 = Pimpinella achilleifolia.
Ajowan, Wall. Cat. n. 572 = Carum copticum.
alata, Bieb. Fl. Taur. Cauc. i. 214 = Ligusticum alatum.
alpina, Stokes, Bot. Mat. Med. ii. 79 = cretensis.
ambigua, *Retz. Obs.* iii. 28.—Hab.?
anethifolia, Wall. Cat. n. 569 = Carum anethifolium.
annua, Linn. Sp. Pl. 245 = cretensis.
arctica, *Nym. Syll.* 155.—Europ.
aurea, *Vis. ex Reichb. Ic. Fl. Germ.* xxi. t. 1935.—Europ.
canescens, DC. Prod. iv. 155 (certe non Ath. sp. ex Boiss. Fl. Orient. ii. 971).—Oriens.
capensis, Burm. f. Fl. Cap. Prod. 7 = Ligusticum capense.
capillacea, Lam. Encyc. i. 325 = cretensis.
caruifolia, [Web. in] Wigg. Prim. Fl. Hols. 27 = Selinum Carvifolia.
carvifolia, *Steph. ex Roem. & Schult. Syst.* vi. 496.—Rossia.
carvifolia, Willd. ex Ledeb. Fl. Ross. ii. 272 = Seseli Hippomarathrum.
carvifolia, Willd. ex Ledeb. l. c. 314 = Peucedanum baikalense.
caucasica, Hort. Par. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. iii. 39 = Seseli athamantoides.

ATHAMANTA:—
Cervaria, Linn. Syst. ed. X. 956 = Peucedanum Cervaria.
cervariaefolia, *DC. Prod.* iv. 155.—Ins. Teneriff.
cervariaefolia, Schrad. ex Spreng. Pugill. ii. 56 = Seseli buchtormensis.
cervariaefolia, Viv. ex Steud. Nom. ed. II. i. 166 = Lecokia cretica.
chiliosciadia, *Boiss. & Heldr. Diagn.* Ser. II. ii. 86.—Graecia.
chinensis, Linn. Sp. Pl. 245 = Selinum Monnieri.
coarctata, Ag. ex Nym. Consp. 296 = Seseli granatense.
collina, Salisb. Prod. 163 = Seseli Libanotis.
compacta, Ledeb. Fl. Alt. i. 327 = Schultzia compacta.
condensata, Linn. Sp. Pl. 1195 = Seseli condensatum.
condensata, Pall. ex Steud. Nom. ed. II. i. 166 = Pallasiana.
cretensis, *Linn. Sp. Pl.* 245.—Europ. austr.
cretensis, Schrank, Baier. Fl. i. 545 = Matthioli.
cretensis, Ucria, Hort. Panorm. 127 = sicula.
crinita, Ledeb. Fl. Alt. i. 326 = Schultzia crinita.
crithmifolia, Juss. ex DC. Prod. iv. 145 = Seseli gummiferum.
crithmoides, Lapeyr. Hist. Abr. Pl. Pyr. 148 = Seseli Monnieri.
daucifolia, Host, Fl. Austr. i. 362 = Seseli Monnieri.
daucoides, J. F. Gmel. Syst. 477 = Seseli Libanotis.
decussata, Gilib. Fl. Lituan. ii. 18 = Peucedanum Cervaria.
densa, *Boiss. & Orph. Diagn.* Ser. II. v. 98.—Graecia.
denudata, Fisch. ex Hornem. Hort. Hafn. Suppl. 32 (= *Cenolophium Fischeri*).—Rossia.
depressa, D. Don, Prod. Fl. Nep. 184 = Cortia Lindleyi.
diffusa, Wall. Cat. n. 570 = Seseli indicum.
divaricata, Gilib. Fl. Lituan. ii. 17 = Peucedanum Oreoselinum.
divaricifolia, Stokes, Bot. Mat. Med. ii. 78 = praec.
flavescens, Vis. Orto Bot. Padov. (1842) 135 = Portenschlagia ramosissima.
flexuosa, Juss. ex DC. Prod. iv. 179 = Peucedanum palustre.
Galatta, Gmel. ex DC. Prod. iv. 230 = Pleurospermum Golaka.
Golaka, Hacq. Pl. Carn. t. 5 = Pleurospermum Golaka.
gigantea, *D. Don, Prod. Fl. Nep.* 184 (Sp. dubia).—Nepal.
Haynaldi, Borb. & Uechtr. in Termész. Füzetek, i. (1877) 30. t. 6, 7 = Matthioli.
hungarica, Borb. ex Nym. Consp. 294 = Matthioli.
incana, Steph. ex Willd. Sp. Pl. i. 1402 = Seseli graveolens.
lasiantha, Willd. ex Roem. & Schult. Syst. vi. 494 = Pimpinella villosa.
lateriflora, *Eckl. & Zeyh. Enum.* 348.—Afr. austr.
latifolia, Viv. Fl. Ital. 22 = Peucedanum Cervaria.
leucospermum, Poir. Encyc. Suppl. i. 532 = Seseli leucospermum.
Libanotis, Linn. Sp. Pl. 244 = Seseli Libanotis.
lucida, *Retz. Obs.* iii. 29.—Hab.?
macedonica, *Spreng. in Roem. & Schult. Syst.* vi. 491.—Graecia; Afr. bor.
Mathioli, DC. Fl. Fr. iv. 318 = cretensis.
Matthioli, *Wulf. in Jacq. Coll.* i. 211.—Eur. austr. or.
maxima, Choisy, in DC. Prod. iv. 155 = cervariaefolia.
media, *Nym. Syll.* 154.—Europ.
Meum, Linn. Sp. Pl. 245 = Meum athamanticum.
Meum, Pall. Reise, ii. 28 = Schultzia crinita.
monstrosa, Steph. ex Roem. & Schult. Syst. Veg. vi. 495 = Schultzia compacta.
multiflora, Sibth. & Sm. Fl. Graec. iii. 69. t. 276 = Carum multiflorum.
mutellinoides, Lam. Encyc. i. 325 = cretensis.
Oreoselinum, Huds. Fl. Angl. 115 = Seseli Libanotis.
Oreoselinum, Linn. Sp. Pl. 244 = Peucedanum Oreoselinum.
Pallasiana, *Vest, ex Steud. Nom.* ed. II. i. 166.—Sibiria.
panacifolia, Spreng. Sp. Umb. 140 = Magydaris panacina, tomentosa.
pisana, Savi, in Ust. Ann. Bot. xxi. 7 = Peucedanum palustre.
pubescens, Retz. Obs. iii. 28 = Seseli Libanotis.
pyrenaica, Jacq. Hort. Vindob. ii. 93 = Seseli Monnieri.
ramosissima, Link & Hoffmgg. Fl. Port. 405. t. 106 = Seseli tortuosum.

ATHAMANTA:—
ramosissima, Portschl. in Roem. & Schult. Syst. vi. 496 = Portenschlagia ramosissima.
Reichenbachii, *Nym. Syll.* 154.—Europ.
rigida, Hornem. Hort. Hafn. 960 = Seseli buchtormensis.
rigida, *Roth, Enum. Pl. Phan. Germ.* i. 907.—Europ.
Riviniana, Host, ex Steud. Nom. ed. II. i. 166 = Seseli Libanotis.
Roxburghiana, Wall. Cat. n. 571 = Carum Roxburghianum.
rupestris, Reichb. Fl. Germ. Excurs. 470 = Matthioli.
rupestris, Vill. Hist. Pl. Dauph. ii. 648 = cretensis.
scabra, *Burm. f. Prod. Fl. Cap.* 7.—Afr. austr.
sibirica, Linn. Sp. Pl. 244 = Seseli sibiricum.
sicula, *Linn. l. c.*—Ital.; Afr. bor.
sicula, Ucria, Hort. Panor. 137 = Kundmannia sicula.
stricta, *Ledeb. ex Steud. Nom.* ed. II. i. 166.—Sibiria.
sylvestris, [Web. in] Wigg. Prim. Fl. Hols. 26 = Angelica sylvestris.
tenuifolia, Pall. ex Roem. & Schult. Syst. vi. 495 = Seseli tenuifolium.
tenuifolia, Willd. ex Spreng. in Roem. & Schult. l. c. = Peucedanum salinum.
teres, *D. Don, Prod. Fl. Nep.* 185 (Sp. dub.).—Nepal.
tortuosa, Spreng. Syst. i. 900 (= *Deverra tortuosa*).—Aegypt.
Turbith, Brot. Phytog. Lusit. 200. t. 169 = Seseli tortuosum.
verticillata, Portschl. Enum. Pl. Dalm. 14. t. 4 = Portenschlagia ramosissima.
verticillata, Sibth. & Sm. Fl. Graec. iii. 68. t. 275 = Carum graecum.

ATHAMUS, Neck. Elem. i. 85 (1790) = **Carlina**, Linn. (Compos.).

ATHANASIA, Linn. Sp. Pl. ed. II. 1180 (1762).
COMPOSITAE, Benth. & Hook. f. ii. 416.
BEMBYCODIUM, Kunze, in Linnaea, xvi. (1842) 316.
HOLOPHYLLUM, Less. Syn. Comp. 262 (1833).
HYMENOLEPIS, Cass. in Bull. Soc. Philom. (1817) 138; et in Dict. Sc. Nat. xxii. 315 (1821).
METAGNANTHUS, Endl. Gen. 438 (1838).
MORYSIA, Cass. in Dict. Sc. Nat. xxxiii. 59 (1824).
PRISTOCARPHA, E. Mey. ex DC. Prod. vi. 86 (1837).
SAINTMORYSIA, Endl. Gen. 438 (1838).
acerosa, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
annua, Linn. Sp. Pl. ed. II. 1182 = Lonas inodora.
aspera, *Thunb. Prod. Pl. Cap.* 145.—Afr. austr.
canescens, *Thunb. l. c.* 144.—Afr. austr.
capitata, Berg. Desc. Pl. Cap. 237 = scabra.
capitata, *Linn. Pl. Rar. Afr.* n. 50; *Amoen. Acad.* vi. 97.—Afr. austr.
cinerea, Linn. f. Suppl. 361 = tomentosa.
coronopifolia, *Harv. in Harv. & Sond. Fl. Cap.* iii. 191.—Natal.
crenata, *Linn. Sp. Pl.* 1180.—Afr. austr.
crithmifolia, *Linn. l. c.* 1181.—Afr. austr.
cuneifolia, *Lam. Encyc.* i. 327.—Afr. austr.
cuneiformis, *DC. Prod.* vi. 89.—Afr. austr.
cupressiformis, Comm. ex DC. Prod. v. 314 = Lepidophyllum cupressiforme.
dentata, *Linn. Sp. Pl.* ed. II. 1181.—Afr. austr.
dentata, Linn. f. Suppl. 361 = serrata.
dimorpha, *DC. Prod.* vi. 87.—Afr. austr.
Dregeana, *Harv. in Harv. & Sond. Fl. Cap.* iii. 191.—Afr. austr.
ebracteata, *E. Mey. ex DC. Prod.* vi. 87.—Afr. austr.
eriopoda, *DC. l. c.* 89.—Afr. austr.
fasciculata, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
filiformis, *Linn. f. Suppl.* 361.—Afr. austr.
flabellifera, Salisb. Prod. 189 = Pentzia flabelliformis.
flexuosa, E. Mey. ex DC. Prod. vi. 91 = microcephala.
flexuosa, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
genistaefolia, Linn. Mant. 464 = Relhania genistaefolia.
glabra, Thunb. Prod. Pl. Cap. 144 = capitata, microphylla.
glabrescens, *DC. Prod.* vi. 88.—Afr. austr.

ATHANASIA:—
glutinosa, E. Mey. ex DC. Prod. vi. 288 = Relhania genistaefolia.
graminifolia, Walt. Fl. Carol. 200 = Marshallia angustifolia.
hastata, Walt. l. c. 201 = Melanthera hastata.
hirsuta, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
imbricata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 196.—Afr. austr.
incisa, *Harv. l. c.* 190.—Afr. austr.
indica, Roxb. Hort. Beng. 61; Fl. Ind. iii. 417 = Goniocaulon glabrum.
indivisa, *Harv. in Harv. & Sond. Fl. Cap.* iii. 190.—Afr. austr.
juncea, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
laevigata, Linn. Pl. Afr. n. 50; Amoen. Acad. vi. 51 = aspera.
lanuginosa, Cav. Ic. i. 2 = canescens.
leucoclada, *Harv. in Harv. & Sond. Fl. Cap.* iii. 191.—Afr. austr.
linifolia, *Harv. l. c.* 198.—Afr. austr.
linifolia, Burm. f. Fl. Cap. Prod. 23 = longifolia.
longifolia, *Lam. Encyc.* i. 326.—Afr. austr.
maritima, Linn. Sp. Pl. ed. II. 1182 = Diotis maritima.
microcephala, D. Dietr. Syn. Pl. iv. 1401 = linifolia.
microphylla, *DC. Prod.* vi. 87.—Afr. austr.
Mundtii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 196.—Afr. austr.
obovata, Walt. Fl. Carol. 201 = Marshallia lanceolata.
pachycephala, *DC. Prod.* vi. 88.—Afr. austr.
palmatifida, *DC. l. c.* 89.—Afr. austr.
paniculata, Walt. Fl. Carol. 201 = Actinomeris alba.
parviflora, *Ait. Hort. Kew.* iii. 167.—Afr. austr.
parviflora, *Thunb. Prod. Pl. Cap.* 145.—Afr. austr.
pauciflora, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
pectinata, *Linn. f. Suppl.* 361.—Afr. austr.
pinnata, E. Mey. ex DC. Prod. vi. 85 = incisa.
pinnata, *Linn. f. Suppl.* 361.—Afr. austr.
pubescens, *Linn. Sp. Pl.* ed. II. 1182.—Afr. austr.
pubescens, Thunb. Prod. Pl. Cap. 144 = rugulosa.
pumila, Linn. f. Suppl. 362 = Relhania pedunculata.
punctata, Berg. Desc. Pl. Cap. 238 = capitata.
punctata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 190.—Afr. austr.
punctata, Thunb. Prod. Pl. Cap. 144 = capitata.
quinquedentata, *Thunb. Prod. Pl. Cap.* 145.—Afr. austr.
recurvifolia, Salisb. Prod. 189 = dentata.
rotundifolia, *DC. Prod.* vi. 87.—Afr. austr.
rugulosa, *E. Mey. ex DC. l. c.*—Afr. austr.
salicifolia, *Dietr. ex Steud. Nom.* ed. I. 91.—Afr. austr.
scabra, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
scariosa, *DC. Prod.* vi. 88.—Afr. austr.
schizolepis, *Harv. in Harv. & Sond. Fl. Cap.* iii. 190.—Afr. austr.
serrata, *Vitm. Summa Veg.* iv. 487.—Afr. austr.
sertulifera, *DC. Prod.* vi. 87.—Afr. austr.
sessiliflora, Linn. f. Suppl. 362 = Relhania sessiliflora.
spathulata, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
squarrosa, Linn. Pl. Rar. Afr. n. 52; Amoen. Acad. vi. 98 = Relhania squarrosa.
tanacetifolia, *Roth, Catalecta*, ii. 104.—Afr. austr.
tomentosa, *Thunb. Prod. Pl. Cap.* 144.—Afr. austr.
tricuspis, Poir. Encyc. Suppl. i. 533 = trifurcata.
tridentata, Salisb. Prod. 189 = trifurcata.
trifurcata, *Linn. Sp. Pl.* ed. II. 1181.—Afr. austr.
trinervia, Walt. Fl. Carol. 201 = Marshallia latifolia
uniflora, Linn. f. Suppl. 362 = Relhania cuneata.
velutina, *D. Dietr. Syn. Pl.* iv. 1401.—Afr. austr.
virgata, *Jacq. Hort. Schoenb.* ii. 11. t. 148.—Afr. austr.

ATHECIA, Gaertn. Fruct. i. 141 (1788) = **Forstera** Linn. f. ? (Stylid.).
glabra, Gaertn. l. c. = F. sedifolia ?

ATHENAEA, Adans. Fam. ii. 121 (1763) = **Spargano phorus**, Vaill. (Compos.).

ATHENAEA, Schreb. Gen. 259 (1789) = **Casearia**, Jacq. (Samyd.).

ATHENAEA, Sendt. in Mart. Fl. Bras. x. 133 (1846) *SOLANACEAE*, Benth. & Hook. f. ii. 890.

ATHENAEA :—
LARNAX, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iv. (1849) 37.
WITHERINGIA, Miers, l. c. iii. (1849) 145.
anonacea, *Sendt. in Mart. Fl. Bras.* x. 137. t. 18.—Bras.
hirsuta, *Sendt. l. c.* 137.—Bras.
Martiana, *Sendt. l. c.* 136.—Bras.
micrantha, *Sendt. l. c.* 135.—Bras.
picta, *Sendt. l. c.* 134.—Bras.
pogogena, *Sendt. l. c.* 135.—Bras.
Pohliana, *Sendt. l. c.* 136.—Bras.
Schottiana, *Sendt. l. c.* 135.—Bras.
xalapensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 422.—Mexic.

ATHENANTHIA, Kunth, in Mém. Mus. Par. ii. (1815) 71 = **Anthaenantia**, Beauv. (Gramin.).

ATHEOLAENA, Reichb. Consp. 106 (1828) = ATHEOLAENA, Cass. = **Senecio**, Linn. (Compos.).

ATHERANDRA, Decne. in DC. Prod. viii. 497 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 743.
acuminata, *Decne. l. c.*—Java.
acutifolia, *Decne. l. c.*—Malaya.
cuspidata, Blume, Mus. Bot. Lugd. Bat. i. 147 = acutifolia.
pubescens, Blume, l. c. = acutifolia.
Wallichii, Benth. & Hook. f. Gen. ii. 744 = Atherolepsis Wallichii.

ATHERANTHERA, Mast. in Oliver, Fl. Trop. Afr. ii. 519 (1871); Trans. Linn. Soc. xxvii. (1871) 640. *PASSIFLOREAE*.
paniculata, *Mast. in Oliver, Fl. Trop. Afr.* ii. 519.—Afr. trop.
Welwitschii, Mast. in Trans. Linn. Soc. xxvii. (1871) 640 = paniculata.

ATHERNOTUS, Dulac, Fl. Hautes-Pyr. 74 (1867) = **Calamagrostis**, Adans. (Gramin.).
epigeios, Dulac, l. c. = C. epigeios.
lanceolatus, Dulac, l. c. 75 = C. lanceolatus.
pyramidalis, Dulac, l. c = Deyeuxia sylvatica.

ATHEROCEPHALA, DC. Prod. vii. 755 (1839) = **Andersonia**, R. Br. (Epacrid.).
Drummondii, DC. l. c. = Andersonia aristata.

ATHEROLEPSIS, Hook. f. in Hook. Ic. Pl. xv. 26. t. 1433 (1883). *ASCLEPIADEAE*.
Wallichii, *Hook. f. l. c.*—Burma.

ATHEROPHORA, Willd. ex Steud. Nom. ed. II. i. 167 (1840) = **Aegopogon**, Humb. & Bonpl. (Gramin.).
secunda, Willd. l. c. = Aegopogon geminiflorus.

ATHEROPOGON, Muhl. ex Willd. Sp. Pl. iv. 937 (1805) = **Bouteloua**, Lag. (Gramin.).
acuminatus, Fourn. Mex. Pl. En. Gram. 139; ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
affinis, Fourn. ex Benth. & Hook. f. Gen. iii. 1169 = B. disticha?
americanus, Fourn. Mex. Pl. En. Gram. 140; ex Hemsl. Biol. Centr. Am. Bot. iii. 561 = B. bromoides.
Antillarum, Spreng. Syst. i. 294 = Aristida Antillarum.
apludoides, Willd. Sp. Pl. iv. 937 = B. racemosa.
aristidoides, Roem. & Schult. Syst. ii. 415 = B. aristidoides.
bromoides, Roem. & Schult. l. c. = B. bromoides.
chondrosioides, Roem. & Schult. l. c. 416 = B. ovata.
curtipendulus, Fourn. Mex. Pl. En. Gram. 138; ex Hemsl. Biol. Centr. Am. Bot. iii. 563 = B. racemosa.
distichus, Spreng. Syst. i. 294 = B. disticha.
domingensis, Spreng. l. c. 293 (= *Eutriana juncea*).—Ins. S. Doming.
filiformis, Fourn. Mex. Pl. En. Gram. 140; ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
gracilis, Spreng. Syst. i. 293 = B. oligostachya.
hirtus, Spreng. l. c. = B. hirsuta.
humilis, Spreng. l. c. = B. prostrata.
juncifolius, Spreng. l. c. 294 = B. juncifolia.
medius, Fourn. Mex. Pl. En. Gram. 139; ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
oligostachyus, Nutt. Gen. Am. i. 78 = B. oligostachya.

ATHEROPOGON :—
Pentarrhaphis, Spreng. Syst. i. 294 (= *Eutriana Pentarrhaphis*).—Mexic.
polymorphus, Fourn. Mex. Pl. En. Gram. 141; ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
procumbens, Jacq. Eclog. Gram. 12 = B. tenuis.
radicosus, Fourn. Mex. Pl. En. Gram. 140; ex Hemsl. Biol. Centr. Am. Bot. iii. 561 = B. bromoides.
repens, Roem. & Schult. Syst. ii. 416 = B. bromoides.
scorpioides, Spreng. Syst. i. 293 = B. scorpioides.
stolonifer, Fourn. Mex. Pl. En. Gram. 140; ex Hemsl. Biol. Centr. Am. Bot. iii. 565.—Mexic.
Triaena, Spreng. Syst. i. 293 = B. Triathera.
villosus, Steud. Nom. ed. II. i. 167 (= *Eutriana villosa*).—Bras.

ATHEROSPERMA, Labill. Nov. Holl. Pl. ii. 74. t. 224 (1806). *MONIMIACEAE*, Benth. & Hook. f. iii. 144.
integrifolium, A. Cunn. ex A. DC. in DC. Prod. xvi. II. 676 = mos:l atum.
micranthum, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 46 = Daphnandra micrantha.
moschatum, *Labill. Nov. Holl. Pl.* ii. 74. t. 224.—Austral.
Novae-Zelandiae, *Hook. f. Handb. New Zeal. Fl.* 240.—N. Zel.
repandulum, *F. Muell. Fragm.* x. 105.—Austral.
Sassafras, A. Cunn. ex A. DC. in DC. Prod. xvi. II. 676 = Doryphora Sassafras.
sempervirens, *Baill. Adansonia*, ix. (1868) 116; *Hist. Pl.* i. 322.—Chili.

ATHEROSTEMON, Blume, Mus. Bot. Lugd. Bat. i. 126 (1849). *ASCLEPIADEAE*.
javensis, *Blume, l. c.*—Burma; Java.

ATHERSTONEA, Pappe, Sylva Cap. ed. II. 29 (1862) = **Strychnos**, Linn. (Logan.).
decussata, Pappe, l. c. = S. Atherstonei.

ATHERURUS, Blume, Rumphia, i. 136 (1835) = **Pinella**, Tenore (Aroid.).
ternatus, Blume, l. c. = P. tuberifera.
tripartitus, Blume, l. c. 137 = P. tripartita.

ATHESIANDRA, Miers, ex Benth. & Hook. f. Gen. i. 346 (1862) = **Ptychopetalum**, Benth. (Olacin.).
anceps, Miers, ex Oliver, Fl. Trop. Afr. i. 348 = P. anceps.

ATHLIANTHUS, Endl. Gen. Suppl. ii. 63 (1842) = **Justicia**, Linn. (Acanth.).
spiciflorus, Heynh. Nom. ii. 54.—Afr. trop.

ATHRIXIA, Ker-Gawl. in Bot. Reg. t. 681 (1823). *COMPOSITAE*, Benth. & Hook. f. ii. 328.
ASTERIDIA, Lindl. Swan Riv. App. 24 (1839).
CHRYSODISCUS, Steetz, in Lehm. Pl. Preiss. i. 460 (1845).
KLENZEA, Sch. Bip. ex Steud. Nom. ed. II. i. 847 (1840); et in Walp. Rep. ii. 973 (1843), partim.
TRICHOSTEGIA, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 81.
angustissima, *DC. Prod.* vi. 277.—Afr. austr.
australis, *Steetz, in Lehm. Pl. Preiss.* i. 482.—Austral.
capensis, *Ker-Gawl. in Bot. Reg.* t. 681.—Afr. austr.
Chatopodo, *F. Muell. Fragm.* x. 56.—Austral.
debilis, *DC. Prod.* vi. 276.—Madag.
elata, *Sond. in Linnaea*, xxiii. (1850) 67.—Afr. austr.
ericaefolia, *DC. Prod.* vi. 277.—Arabia.
fontana, *Mac Owan, in Journ. Linn. Soc.* xviii. (1881) 391.—Afr. austr.
Gerrardi, *Harv. in Harv. & Sond. Fl. Cap.* iii. 292.—Afr. austr.
glandulosa, *Kunth, Ind. Sem. Hort. Berol.* (1848) 15.—Afr. austr.
gracilis, *Benth. Fl. Austral.* iii. 599.—Austral.
heterophylla, *Less. Syn. Comp.* 365.—Afr. austr.
multiceps, *Benth. Fl. Austral.* iii. 599.—Austral.
phylicoides, *DC. Prod.* vi. 277.—Afr. austr.
rosmarinifolia, Oliver & Hiern, in Oliver, Fl. Trop. Afr. iii. 355.—Afr. trop.
sessilifolia, *DC. Prod.* vi. 277.—Afr. austr.
stricta, *Benth. Fl. Austral.* iii. 600.—Austral.
tenella, *Benth. l. c.*—Austral.
tomentosa, Less. Syn. Comp. 364 = Pterothrix tomentosa.

ATHRODACTYLIS, Forst. Char. Gen. 149 (1776) = **Pandanus**, Linn.
spinosa, Forst. l. c. 150 = P. odoratissimus.

ATHROISMA, DC. in Guill. Arch. Bot. ii. (1833) 516. *COMPOSITAE*, Benth. & Hook. f. ii. 296.
laciniatum, *DC. l. c.* 517.—Ind. or.
viscida, Zoll. & Mor. in Nat. en Geneesk. Arch. Nerl. Indie, ii. (1845) 232 = laciniatum.

ATHROISMA, Griff. Notul. iv. 477 (1854) = **Trigonostemon**, Blume (Euphorb.).
dentatum, Griff. l. c. 478 = T. longifolius.
serratum, Griff. l. c. 477 = T. heteranthus.

ATHRONIA, Neck. Elem. i. 32 (1790) = **Spilanthes**, Jacq. (Compos.).

ATHROOSTACHYS, Benth. ex Benth. & Hook. f. Gen. iii. 1208 (1883). *GRAMINEAE*, Benth. & Hook. f. iii. 1208.
capitata, *Benth. l. c.* 1209.—Bras.

ATHROTAXIS, D. Don, in Trans. Linn. Soc. xviii. (1839) 171. t. 13, 14. *CONIFERAE*, Benth. & Hook. f. iii. 430.
ARTHROTAXIS, Endl. Gen. Suppl. i. 1372 (1841).
alpina, Van Houtte, ex Gord. Pinet. 31 = selaginoides.
araucarioides, Brongn. ex Parl. in DC. Prod. xvi. II. 496 = Dacrydium araucarioides.
cupressoides, *D. Don, in Trans. Linn. Soc.* xviii. (1839) 173. t. 13.—Tasmania.
Doniana, Henk. & Hochst. Nadelh. 221 = laxifolia.
Gunneana, *Hook. ex Carr. Conif.* ed. II. 207.—Tasmania.
imbricata, Hort. ex Carr. l. c. 204 = selaginoides.
imbricata, Maule, ex Gord. Pinet. Suppl. 16 = cupressioides.
Joucadan, Hort. ex Carr. Conif. ed. II. 55 = Juniperus gracilis.
laxifolia, *Hook. Ic. Pl.* t. 573.—Tasmania.
selaginoides, *D. Don, in Trans. Linn. Soc.* xviii. (1839) 172. t. 14.—Tasmania.
tetragona, Hook. Ic. Pl. t. 560 = Microcachrys tetragona.

ATHRUPHYLLUM, Lour. Fl. Cochinch. 120 (1790) = **Myrsine**, Linn.
lineare, Lour. l. c. = M. Athruphyllum.

ATHYMALUS, Neck. Elem. ii. 353 (1790) = **Euphorbia**, Linn.

ATHYROCARPUS, Schlecht. in Linnaea, xxvi. (1853-55) 454. *COMMELINACEAE*, Benth. & Hook. f. iii. 847.
PHAEOSPHAERION, Hassk. in Flora, (1866) 212.
leiocarpus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 386.—Mexic.
pallidus, Hassk. ex C. B. Clarke, in DC. Monog. Phan. iii. 151 = Commelina pallida.
pallidus, *Schlecht. in Linnaea*, xxvi. (1853-55) 454.—Am. trop.
persicariaefolius, *Hemsl. Biol. Centr. Am. Bot.* iii. 386.—Panama.

ATHYRUS, Neck. Elem. iii. 39 (1790) = **Lathyrus**, Linn. (Legumin.).

ATHYSANUS, Greene, in Bull. Calif. Acad. i. (1885) 72. *CRUCIFERAE*.
pusillus, *Greene, l. c.*—Calif.

ATIMETA, Schott, Gen. Aroid. t. 71 (1858) = **Rodospatha**, Poepp. (Aroid.).
filamentosa, Reissek, ex Peyr. Aroid. Maxim. 28. t. 19, 20.—Bras.
Martii, Schott, Prod. Aroid. 349 = R. oblongata.
Videniana, Schott, Gen. Aroid. t. 71 = R. oblongata.

ATIRSITA, Rafin. New Fl. Am. iv. 36 (1836) = **Eryngium**, Tourn. (Umbell.).
pumila, Rafin. l. c. 37 = E. gracile.

ATITARA, Marcgr. ex Juss. in Dict. Sc. Nat. iii. 277 (1816) = **Evodia**, Forst. (Rutac.).

ATKINSONIA, F. Muell. Fragm. v. 34 (1865) = **Loranthus**, Linn.
ligustrina, F. Muell. l. c. v. 34; vi. 252 = L. epigaeus.

ATLANTIA, Kurz, For. Fl. Brit. Burma, i. 194 (1877) = **Atalantia**, Correa (Rutac.).

ATOCION, Adans. Fam. ii. 254 (1763) = **Silene**, Linn. (Caryophyll.).
Armeria, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344 = S. Armeria.

ATOLARIA, Neck. Elem. iii. 31 (1790) = **Crotalaria**, Linn. (Legum.).

ATOMOSTYLIS, Steud. in Flora, xxxiii. (1850) 229; Syn. Pl. Cyp. 315 (1855) = **Cyperus**, Linn.
cyperiformis, Steud. Syn. Pl. Cyp. 315 = C. nudicaulis.
flavescens, Steud. in Flora, xxxiii. (1850) 229. = C. nudicaulis.

ATOSSA, Alef. in Bonplandia, ix. (1861) 100 = **Vicia**, Linn. (Legum.).
Clusii, Alef. l. c. 101 = V. oroboides.
sepium, Alef. l. c. = V. sepium.

ATRACTYLIA, Reichb. Nom. 'n. 3416' (1841) = seq.

ATRACTYLIS, Linn. Gen. ed. I. 246 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 465.
ACARNA, All. Fl. Pedem. i. 153 (1785).
ANACTIS, Cass. in Dict. Sc. Nat. xlvii. 510 (1827).
ATRACTYLODES, DC. Prod. vii. 48 (1838).
CHAMAELEON, Cass. in Dict. Sc. Nat. xlvii. 509 (1827).
CIRSELLIUM, Gaertn. Fruct. ii. 454. t. 163 (1791).
CROCODYLUS, 'Adans.' ex Steud. Nom. ed. II. i. 167, in syn. (1840).
SPADACTIS, Cass. in Dict. Sc. Nat. xlvii. 510 (1827).
acaulis, Pers. Syn. ii. 382, in syn. = gummifera.
amplexicaulis, Poir. Encyc. vi. 30 (= Stobaeae sp.).—Afr. austr.
angustifolia, Houtt. Plantenk. x. 518 = Berkheya lanceolata.
armena, Poir. Encyc. Suppl. iv. 637 (= *Onobroma armenum*).—Armen.; Persia.
caespitosa, *Desf. Fl. Atlant.* ii. 254. t. 225.—Afr. bor. occ.
caespitosa, Viv. Fl. Lib. Spec. 52 = cancellata.
cancellata, *Linn. Sp. Pl.* 830.—Reg. Mediterr.; Orient.
canescens, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 519 = cancellata.
chinensis, DC. Prod. vi. 549 = ovata.
ciliaris, *Linn. Syst.* ed. X. 1202.—'Aethiopia.'
citrina, Coss. & Kralik, in Bull. Soc. Bot. Fr. iv. (1857) 471 = flava.
comosa, *Sieber, ex Cass. in Dict. Sc. Nat.* l. 58.—Syria.
cuncata, *Boiss. Fl. Orient.* iii. 454.—Afghanist.
diffusa, *Coss. in Bull. Soc. Bot. Fr.* iv. (1857) 396, nomen.—Algeria.
echinata, *Pomel, Nouv. Mat. Fl. Atl.* 274.—Algeria.
flava, Coss. & Dur.—Cf. Bull. Soc. Bot. Fr. iv. (1857) 471 = citrina.
flava, *Desf. Fl. Atlant.* ii. 254. t. 262.—Afr. bor.; Arab.; Beluchist.
fruticosa, Linn. Sp. Pl. 829 = Berkheya fruticosa.
Fusus-agrestis, Gaertn. Fruct. ii. 381. t. 161 = Carthamus lanatus.
gummifera, *Linn. Sp. Pl.* 829.—Reg. Mediterr.
gummifera, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 519 = Carlina racemosa.
humilis, *Linn. Sp. Pl.* 829.—Europ. austr. occ.; Marocco.
lanata, Scop. Fl. Carn. ed. II. ii. 134 = Carthamus lanatus.
lancea, Thunb. Fl. Jap. 306 = ovata.
leucocephala, Moench, Meth. 553 = Carthamus lanatus.
leucophaea, Gaertn. Fruct. ii. 321 = Carthamus leucocaulis.
macrocephala, *Desf. Fl. Atlant.* ii. 253.—Afr. bor. occ.
macrophylla, *Desf. l. c.* 255.—Algeria.
mexicana, Linn. f. Suppl. 350 = Lycoseris mexicana.
microcephala, Coss. & Dur. in Ann. Sc. Nat. Sér. IV. i. (1854) 240 = serratuloides.
oppositifolia, Linn. Mant. ii. 477 = Berkheya grandiflora.
ovata, *Thunb. Fl. Jap.* 306.—Japon.
persica, Boiss. Diagn. Ser. I. vi. 110 = Thevenotia persica.
phaeolepis, *Pomel, Nouv. Mat. Fl. Atl.* 273.—Algeria.
pilosa, Moench, Meth. 553 = Carthamus lanatus.

ATRACTYLIS:—
polycephala, *Coss. in Bull. Soc. Bot. Fr.* iv. (1857) 481, nomen.—Algeria.
Preauxiana, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 350.—Ins. Canar.
prolifera, *Boiss. Diagn.* Ser. I. x. 96.—Palaestina.
purpurata, Linn. ex Sm. Ic. fasc. iii. 65 = seq.
purpurea, Linn. f. Suppl. 349 = Onoseris purpurea.
radiciflora, *DC. Prod.* vi. 551.—Gall. austr.
rigida, Poir. Encyc. Suppl. iv. 637 = Carthamus rigidus.
scabra, *Boiss. Diagn.* Ser. I. vi. 109.—Persia.
serrata, *Pomel, Nouv. Mat. Fl. Atl.* 20.—Afr. bor.
serratuloides, *Sieber, ex Cass. in Dict. Sc. Nat.* l. 58.—Algeria; Palaest.
speciosa, DC. in Deless. Ic. Sel. iv. t. 71 = comosa.
triplinervia, Vahl, ex Steud. Nom. ed II. i. 168; ii. 83 = Lycoseris triplinervia.

ATRACTYLODES, DC. Prod. vii. 48 (1838) = **Atractylis**, Linn. (Compos.).
lancea, DC. l. c. = Atractylis ovata.
lyrata, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 193 = Atractylis ovata.
ovata, DC. Prod. vii. 48 = Atractylis ovata.

ATRAGENA, Linn. Syst. ed. I. (1735) = seq.

ATRAGENE, Linn. Diss. Dass. 21 (1747); Amoen. Acad. i. 132 = **Clematis**, Linn. (Ranunc.).
alpina, Linn. Sp. Pl. 542; Rev. Hortic. (1855) 323. t. 17. f. 1 = C. alpina.
alpina, Pall. Fl. Ross. ii. 69 = C. verticillaris.
americana, Sims, Bot. Mag. t. 887 = C. verticillaris.
austriaca, Scop. Fl. Carn. ed. II. i. 387 = C. alpina.
austriaca, Wender. in Linnaea, xi. (1837) 648, in obs. = C. Wenderothii.
balearica, Pers. Syn. ii. 98 = C. cirrhosa.
capensis, Burm. f. Fl. Cap. Prod. 16, pro parte = Anemone tenuifolia.
capensis, Linn. Sp. Pl. 543 = Anemone capensis.
ceylanica, Steud. Nom. ed. II. i. 168, vide *zeylanica*.
cirrhosa, Pers. Syn. ii. 98 = C. cirrhosa.
Clematides, Crantz, Stirp. Austr. ed. I. fasc. ii. 111; ed. II. 127 = C. alpina.
columbiana, Nutt. in Journ. Acad. Phil. vii. (1834) 7 = C. verticillaris.
cordata, Wender. ex Steud. Nom. ed. II. i. 168 = C. Wenderothii.
florida, Pers. Syn. ii. 98 = C. florida.
indica, Desf. Tabl. ed. I. 123 = C. florida.
indica, Heyne, ex Steud. Nom. ed. II. i. 168, 379 = C. Gouriana.
japonica, Thunb. Fl. Jap. 238 = C. polypetala.
japonica, Wight, ex Steud. Nom. ed. II. i. 168 = C. Gouriana.
lobata, Llanos, Fragm. 73 = Naravelia zeylanica.
macropetala, Ledeb. Fl. Alt. ii. 376 = C. macropetala.
occidentalis, Hornem. Hort. Hafn. ii. 520 = C. occidentalis.
ochotensis, Pall. Fl. Ross. i. II. 69 = C. ochotensis.
patens, Deene. in Belg. Hortic. (1853) 16 = C. patens.
platysepala, Trautv. & C. A. Mey. in Middend. Reise, (Fl. Ochot. 5).—As. bor. occ.
polygama, Jacq. Am. Select. 171 = C. dominica.
polypetala, Thunb. Fl. Jap. 239 = Anemone japonica.
sibirica, Linn. Sp. ed. I. 343 = alpina.
speciosa, *Weinm. in Bull. Soc. Nat. Mosc.* (1850) i. 538.—Rossia.
tenuifolia, Linn. f. Suppl. 270 = Anemone tenuifolia.
tenuis, Thunb. Fl. Jap. 239 = Anemone tenuifolia.
triternata, DC. Syst. i. 150 = C. triternata.
Wenderothii, Schlecht. in Linnaea, xi. (1837) 650 = C. Wenderothii.
zeylanica, Blanco, Fl. Filip. ed. I. 461 = Naravelia laurifolia.
zeylanica, Linn. Sp. Pl. 542 = Naravelia zeylanica.

ATRAPHAXIS, Linn. Syst. ed. I (1735). *POLYGONACEAE*, Benth. & Hook. f. iii. 96.
PEDALIUM, Adans. Fam. ii. 277 (1763).
PERSICARIA, Neck. Elem. ii. 210 (1790).
TRAGOPYRUM, Bieb. Fl. Taur. Cauc. iii. 284 (1819).
afghanica, *Meissn. in DC. Prod.* xiv. 76.—Afghanist.
angustifolia, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15, 21. t. 115.—Armenia.
aphylla, *Pall. Reise*, i. 386.—Rossia.

ATRAPHAXIS:—
Aucherii, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15, 20. t. 114.—Persia.
Billardieri, *Jaub. & Spach, l. c.* 14, 17. t. 111.—Graecia; As. Min.; Syria.
buxifolia, *Jaub. & Spach, l. c.* 14.—Reg. Caucas.
Calverti, Boiss. Diagn. Ser. II. iv. 76 = spinosa.
candida, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 1021.—Persia.
canescens, *Bunge, Ind. Sem. Hort. Dorpat.* (1839) 3.—As. bor.
compacta, *Ledeb. Fl. Alt.* ii. 55.—Turkestan.
crassifolia, Agardh, in Schult. f. Syst. vii. 1382 = spinosa.
cuneifolia, *Poir. Encyc. Suppl.* i. 535.—Am. bor.
davurica, Jaub. & Spach, Illustr. Pl. Or. ii. 15 = Laxmanni.
decipiens, Jaub. & Spach, l. c. = Laxmanni.
densiflora, C. Koch, in Linnaea, xxii. (1849) 212 = spinosa.
dioica, Bosc, ex Meissn. in DC. Prod. xiv. 80 = Polygonella parvifolia.
Fischeri, Jaub. & Spach, Illustr. Pl. Or. ii. 12 = spinosa.
frutescens, C. Koch, Dendrol. ii. 1. 369 = Laxmanni.
graeca, Meissn. in DC. Prod. xiv. 77 = Billardieri.
grandiflora, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15, 19. t. 113.—Armenia; Cappadoc.
inermis, Mill. Gard. Dict. ed. VIII. n. 2 = undulata.
Karelini, Jaub. & Spach, Illustr. Pl. Or. ii. 12 = spinosa.
laetevirens, *Jaub. & Spach, l. c.* 14.—Soongaria.
lanceolata, *Bunge, in Mém. Sav. Etr. Pétersb.* vii (1851) 483 (*Lehm. Rel.* 307).—As. temp.
lanceolata, Meissn. in DC. Prod. xiv. 78 = Laxmanni.
Laxmanni, *Agardh, in Schult. f. Syst.* vii. 1382.—Sibir.
micrantha, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15.—Sibiria.
pungens, *Jaub. & Spach, l. c.* 14.—Soongaria.
pyrifolia, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 483 (*Lehm. Rel.* 307).—Turkestan.
replicata, Lam. Encyc. i. 329 = spinosa.
sinaica, Fresen. in Mus. Senkenberg. i. 74; Jaub. & Spach, Illustr. Pl. Or. ii. 12 = spinosa.
spinosa, Eichw. Casp. Cauc. 7 = compacta.
spinosa, Laxm. ex Agardh, in Schult. f. Syst. vii. 1382 = crassifolia, Laxmanni.
spinosa, *Linn. Sp. Pl.* 475.—Orient.
suaedaefolia, *Jaub. & Spach, Illustr. Pl. Or.* ii. 15.—Persia.
Tournefortii, *Jaub. & Spach, l. c.* 14, 18. t. 112.—Armenia.
undulata, Linn. Sp. Pl. 333 = Polygonum atraphaxoides.
undulata, *Willd. Sp. Pl.* ii. 249.—Afr. austr.
variabilis, Jaub. & Spach, Illustr. Pl. Or. ii. 11, 13, 16. t. 110 = Billardieri.

ATRATEGIA, Bedd. ex Hook. f. Fl. Brit. Ind. i. 74 (1872) = **Gonothalamus**, Blume (Anonac.).
Thomsoni, Bedd. ex Hook. f. l. c. 75 = G. cardiopetalus.

ATREMA, DC. Coll. Mém. v. 71. t. 18 (1829); et Prod. iv. 250 (1830) = **Bifora**, Hoffm. (Umbellif.).
americanum, DC. Coll. Mém. v. 71. t. 18 = B. americana.
testiculatum, Miq. Fl. Ind. Bat. i. 1. 744 = B. testiculata.

ATRICHOSERIS, A. Gray, Syn. Fl. N. Am. i. ii. 410 (1884). *COMPOSITAE*.
platyphylla, *A. Gray, l. c.*—Am. bor. occ.

ATRIPLEX, [Tourn.] Linn. Syst. ed. I (1735); Gen. 317 (1737). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 53.
ENDOLEPIS, Torr. ex A. Gray, in Pacif. Rail. Rep. xii. 47. t. 3 (1860).
HALIMUS, Wallr. Sched. Crit. 117 (1822).
OBIONE, Gaertn. Fruct. ii. 198. t. 126 (1791).
PHYLLOTHECA, Nutt. ex Moq. in DC. Prod. xiii. ii. 98 (1849).
PTEROCARYA, Nutt. ex Moq. l. c. 106 (1849).
PTEROCHITON, Torr. in Frem. Rep. 318 (1845).
SCHIZOTHECA, Lindl. Veg. Kingd. 513 (1847).
THELEOPHYTON, Moq. in DC. Prod. xiii. ii. 115 (1849).

ATRIPLEX:—
abbreviata, Opiz, in Flora, vii. (1824) 1. Beil. 81 = laciniata.
acanthocarpa, *S. Wats. in Proc. Am. Acad.* ix. (1874) 117.—Texas.
acuminata, Bieb. Fl. Taur. Cauc. iii. 641 = hortensis.
acuminata, Waldst. & Kit. Pl. Rar. Hung. ii. 107. t. 103 = nitens.
agrestis, Schur, Enum. Pl. Transs. 575 = patula.
alaskensis, *S. Wats. in Proc. Am. Acad.* ix. (1874) 108.—Am. bor. occ.
alba, Crantz, Inst. i. 206 = Chenopodium album.
alba, Salisb. Prod. 152 = Exomis atriplicioides.
alba, Scop. Delic. Insub. ii. 16. t. 8 = rosea.
albicans, [Soland. in] Ait. Hort. Kew. ed. I. iii. 430 = Exomis atriplicioides.
albicans, Bess. Prim. Fl. Galic. i. 194 = rosea.
albicans, Willd. Sp. Pl. iv. 962 = laciniata.
Aldamae, *Griseb. Cat. Pl. Cub.* 282.—Cuba.
alexandrina, *Boiss. Fl. Orient.* iv. 914.—Aegypt.
amarantoides, Gmel. ex Moq. in DC. Prod. xiii. II. 116 = Axyris amaranthoides.
amblyostegia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 416.—As. centr.
ambrosioides, Crantz, Inst. i. 207 = Chenopodium ambrosioides.
angulata, *Benth. Fl. Austral.* v. 174.—Austral.
angustifolia, Sm. Fl. Brit. 1092 = patula.
anthelmintica, Crantz, Inst. i. 207 = Chenopodium ambrosoides.
arabicum, Ehrenb. ex Boiss. Fl. Orient. iv. 917 = farinosa.
arenaria, *Nutt. Gen. Am.* i. 198.—Am. bor.
arenaria, Tineo, Cat. Hort. Panorm. 276; Woods, in Bab. Man. ed. III. 271 = laciniata.
argentea, *Nutt. Gen. Am.* i. 198; *ex S. Wats. Rev. Chenop.* 117.—Am. bor. occ.
argentea, Pall. ex Steud. Nom. ed. II. i. 168 = nitens.
argentea, Schrad. ex Moq. in DC. Prod. xiii. II. 92 = rosea.
aristata, Crantz, Inst. i. 208 = Chenopodium aristatum.
Assoi, *Dufour, in Bull. Soc. Bot. Fr.* vii. (1860) 432.—Hispan.
astracanica, Balb. Hort. Turic. (1813) = laciniata.
astracanica, *Lag. Gen. et Sp. Nov.* (1816) 3.—As. temp.
astrachanica, Ledeb. Fl. Ross. iii. 719 = laciniata.
atacamensis, *Phil. Fl. Atac.* 48.—Chili.
Aucherii, Moq. Chenop. Enum. 51 = nitens.
australasica, Moq. l. c. 59 = patula.
axillaris, Tenore, Fl. Nap. Prod. i. 54 = rosea.
Babingtonii, Woods, in Bab. Man. ed. III. 270 = patula.
Barclayana, *D. Dietr. Syn. Pl.* v. 537.—Am. bor. occ.
Belangeri, *D. Dietr. l. c.* 535.—Persia; Afghanist.
benghalensis, Lam. Encyc. i. 276 = hortensis.
Berlandieri, Moq. Chenop. Enum. 65 = canescens.
Besseriana, Roem. & Schult. Syst. vi. 284 = rosea.
Billardieri, *Hook. f. Fl. N. Zel.* i. 215.—Austral.; N. Zel.
Bocconi, Guss. Fl. Sic. Syn. ii. 588 = laciniata.
Bonus-Henricus, Crantz, Inst. i. 206 = Chenopodium Bonus-Henricus.
Botrys, Crantz, l. c. 207 = Chenopodium Botrys.
bracteosa, Trautv. in Act. Hort. Petrop. i. n. 1 (1870) 17 = dimorphostegia.
bracteosa, *S. Wats. in Proc. Am. Acad.* ix. (1874) 115.—Calif.
Breweri, *S. Wats. l. c.* 119.—Calif.
Bunburyana, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 274.—Austral.
californica, *Moq. in DC. Prod.* xiii. II. 98.—Calif.
calotheca, *Fries, Novit. Fl. Suec. Mant.* iii. 164.—Suecia.
campanulata, *Benth. Fl. Austral.* v. 177.—Austral.
campestris, Koch & Ziz, Cat. Pl. Palat. 24 = tatarica.
cana, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 306; *Moq. Chenop. Enum.* 61.—Sibiria.
candicans, *Link, ex Steud. Nom.* ed. II. i. 168.—Lusitan.
canescens, *James, Cat.* 178.—Am. bor. occ.
canescens, Nutt. Gen. Am. i. 197 = Nuttallii.
capensis, *Moq. Chenop. Enum.* 63.—Afr. austr.

ATRIPLEX:—
Chanicowii, *Bunge, ex Boiss. Fl. Orient.* iv. 913.—Persia.
chilensis, *Colla, in Mem. Acc. Torin.* xxxix. (1836) 7. t. 49.—Chili.
chrystallina, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 279 = Billardieri.
cinerea, *Poir. Encyc. Suppl.* i. 471.—Austral.; N. Zel.
confertifolia, *S. Wats. in Proc. Am. Acad.* ix. (1874) 119.—Am. bor. occ.
coriacea, *Forsk. Fl. Aegypt. Arab.* 175.—Aegypt.
coriacea, Raddi, ex Moq. in DC. Prod. xiii. II. 108 = glauca.
coronata, *S. Wats. in Proc. Am. Acad.* ix. (1874) 114.—Calif.
Coulteri, *D. Dietr. Syn. Pl.* v. 537.—Calif.
crassifolia, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 309.—Turkestan.; Ind. or.; Sibir.
crispa, *D. Dietr. Syn. Pl.* v. 536.—Persia.
cristata, Heyne, ex Moq. in DC. Prod. xiii. II. 99; Koen. ex Moq. l. c. 109 = repens.
eristata, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 959.—N. Granat.
crystallina, Moq. l. e. 105, 116 = Billardieri.
crystallina, *Ehrenb. ex Boiss. Fl. Orient.* iv. 915.—Aegypt.
decumbens, Roem. & Schult. Syst. vi. 289 = prostrata.
decumbens, *S. Wats. in Proc. Am. Acad.* xii. (1877) 275.—Calif.
deltoidea, Bab. Man. ed. I. 253 = hastata.
denticulata, Moq. in DC. Prod. xiii. II. 97 = semibaccata.
deserticola, *Phil. Fl. Atac.* 47.—Chili.
diffusa, Tenore, Prod. Fl. Nap. i. 57 = laciniata.
dimorphostegia, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 438.—Soongar.
dioica, Rafin. in Am. Monthly Mag. (1818) 176 = Purshiana.
Drummondii, *Moq. in DC. Prod.* xiii. II. 102.—Austral.
clachophylla, *F. Muell. Fragm.* vii. 8.—Austral.
elaeagnoides, Moq. Chenop. Enum. 65 = cinerea.
elegans, *D. Dietr. Syn. Pl.* v. 537.—Mexic.
elongata, Guss. Fl. Sic. Syn. ii. 592 = hastata.
Endolepis, *S. Wats. in Proc. Am. Acad.* ix. (1874) 110.—Am. bor.
erecta, Huds. Fl. Angl. ed. I. 376 = patula.
exilifolia, *F. Muell. Fragm.* vii. 9.—Austral.
expansa, *S. Wats. in Proc. Am. Acad.* ix. (1874) 116.—Am. bor. occ.
farinosa, Dum. Fl. Belg. 20 = laciniata.
farinosa, Moq. Chenop. Enum. 55 = Verreauxii.
farinosa, *Forsk. Fl. Aegypt. Arab.* 123.—Aegypt.; Arab.
fasciculata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 377.—Am. bor.
fissivalvis, *F. Muell. Fragm.* ix. 123.—Austral.
Flabellum, *Bunge, ex Boiss. Fl. Orient.* iv. 912.—Persia; Turkestan.
flavescens, Dum. Fl. Belg. 20 (Quid ?).—Belgium.
flexuosa, *Moq. Chenop. Enum.* 57.—Afr. austr.
foliolosa, Link, in Schrad. Journ. (1800) 1. 54 = rosea.
foliolosa, *Phil. in Bot. Zeit.* xxviii. (1870) 500; *Anal. Univ. Chil.* (1875) 192.—Chili.
foliosa, Guss. ex Nym. Consp. 628 = rosea.
fruticosa, Nutt. ex Moq. in DC. Prod. xiii. II. 112 = canescens.
Gardneri, D. Dietr. Syn. Pl. v. 537 = Gordoni.
gigantea, Poir. ex Moq. in DC. Prod. xiii. II. 94 = hastata.
glauca, Brot. Fl. Lusit. i. 472 = candicans.
glauca, Crantz, Inst. i. 207 = Chenopodium glaucum.
glauca, Pall. ex Roem. & Schult. Syst. v. 277 = verrucifera.
glauca, Pall. Reise, i. 494 = cana.
glauca, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 645.—laciniata.
glauca, *Linn. Cent.* i. 33.—Europ.
glomulifera, *Nees, in Lehm. Pl. Preiss.* i. 634.—Austral.
Gmelini, *C. A. Mey. in Bong. Veg. Sitch.* 42.—As. bor. or.
Gordoni, *Hook. Kew Journ.* v. (1853) 261.—Am. bor.
gracilis, Nutt. ex Moq. in DC. Prod. xiii. II. 95 = hastata.
graeca, Sibth. & Sm. Fl. Graec. x. t. 963 = tatarica.
graeca, Willd. Sp. Pl. iv. 958 = hastata.

ATRIPLEX:—
Graeci, Tineo, Cat. Hort. Panorm. 277 = rosea.
Greggii, *S. Wats. in Proc. Am. Acad.* ix. (1874) 118.—Mexic.
Griffithii, *Moq. in DC. Prod.* xiii. II. 102.—Persia; Afghanist.
halimoides, *Lindl. in Mitch. Three Exp.* i. 285.—Austral.
halimoides, Rafin. in Am. Monthly Mag. (1818) 176 = hastata.
halimoides, *Tineo, Cat. Hort. Panorm.* 277.—Hab.?
Halimus, R. Br. Prod. 406 = cinerea.
Halimus, *Linn. Sp. Pl.* 1052.—Afr. bor. et austr.
Halimus, Nees, in Lehm. Pl. Preiss. i. 633 = cinerea, isatidea.
Halimus, Pall. Reise, ii. 477 = cana.
Halimus, Ritter, ex Moq. in DC. Prod. xiii. II. 110 (= *Obione crispa*).—Ins. S. Doming.
hastaefolia, Salisb. Prod. 153 = hastata.
hastata, Geners. ex Steud. Nom. ed. II. i. 169 = laciniata.
hastata, Koch, Syn. Fl. Germ. ed. II. 702 = calotheca.
hastata, *Linn. Sp. Pl.* 1053.—Europ.; As. bor.
hastilifolia, Reichb. ex Steud. Nom. ed. II. i. 169 = patula.
Hermannii, Villem. Phytogr. iii. 1222 = nitens.
heterantha, Wight, Ic. v. t. 1787 = hortensis.
heterophylla, Nutt. ex Moq. in DC. Prod. xiii. II. 112 = canescens.
heterosperma, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 448 (*Lehm. Rel.* 272).—As. centr.
hispanica, *S. G. Gmel. It.* iv. 40.—Ross. austr.
holocarpa, *F. Muell. Rep. Babb. Exp.* 19.—Austral.
hortensis, *Linn. Sp. Pl.* 1053.—Cosmopol. (cult.).
humilis, *F. Muell. Fragm.* iv. 48.—Austral.
hybrida, Crantz, Inst. i. 207 = Chenopodium hybridum?
hymenelytra, *S. Wats. in Proc. Am. Acad.* ix. (1874) 119.—Am. bor. occ.
hymenotheca, *Moq. in DC. Prod.* xiii. II. 101.—Austral.
hypoleuca, Nees, in Lehm. Pl. Preiss. i. 633 = cinerea.
imbricata, *D. Dietr. Syn. Pl.* v. 536.—Bolivia.
incana, *F. G. Dietr. Vollst. Gaertn. Lexik. Nachtr.* i. 418.—Hispan.
incisa, Bieb. Fl. Taur. Cauc. iii. 641 = tatariea.
incisa, Hohen. ex Ledeb. Fl. Ross. iii. 720 = laciniata.
incrassata, *F. Muell. Rep. Babb. Exp.* 20.—Austral.
inflata, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 75 = halimoides.
integrifolia, Krock. ex Steud. Nom. ed. II. i. 169 = triangularis.
isatidea, *Moq. Chenop. Enum.* 63.—Austral.
julacea, *S. Wats. in Proc. Am. Acad.* xx. (1885) 370.—Calif.
Kataf, Ehrenb. ex Boiss. Fl. Orient. iv. 916 = A. Halimus.
Koenigii, Wall. Cat. n. 6951 = repens.
Kunthiana, *D. Dietr. Syn. Pl.* v. 536.—Am. austr.
lacera, Desf. Cat. Hort. Par. ed. III. 389 = calotheca.
laciniata, Aitch. Cat. Pl. Panjab 125 = crassifolia.
laciniata, Bieb. Fl. Taur. Cauc. iii. 641 = rosea.
laciniata, *Linn. Sp. Pl.* 1053.—Europ.; Am. bor.
laciniata, Pursh, Fl. Am. Sept. i. 199 = Purshiana.
laciniata, Schkuhr, Handb. t. 349 = hastata.
laevis, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 60.—Sibir.
Lampa, *Gill. ex Moq. in DC. Prod.* xiii. II. 110.—Chili.
lasiantha, Boiss. Diagn. Ser. I. xii. 95 = tatarica.
latifolia, Wahlenb. Fl. Suec. ii. 660 = hastata.
Lehmanniana, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 451 = turcomanica.
lenticularis, *C. A. Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* (1838) 99, nomen.—As. temp.
lenticularis, Tausch, in Flora, xxiii. (1840) 1. 16 = Tauschii.
lentiformis, *S. Wats. in Proc. Am. Acad.* ix. (1874) 118.—Am. bor. occ.
leptocarpa, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 74.—Austral.
leptoclada, Boiss. Diagn. Ser. II. iv. 74 = turcomanica.
leucoclada, *Boiss. Diagn.* Ser. I. xii. 95.—Aegypt.; Arab.
leucophylla, *D. Dietr. Syn. Pl.* v. 536.—Calif.
limbata, *Benth. Fl. Austral.* v. 178.—Austral.
limosa, *Schur, Enum. Pl. Transs.* 576.—Europ.
Lindleyi, Moq. in DC. Prod. xiii. II. 100 = halimoides.

ATRIPLEX:—
linifolia, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 958.—Mexic.
littoralis, *Linn. Sp. Pl.* 1054.—Reg. bor. temp.
longipes, Drejer & Fries, Summa Veg. Scand. 201 = hastata.
lucida, Desf. ex Mert. & Koch, in Rochl. Fl. Deutschl. ii. 311 = nitens.
lucida, Moench, Meth. Suppl. 121 = sibirica.
lunulata, Del. ex Moq. in DC. Prod. xiii. II. 99 = mollis.
macrodira, Guss. Fl. Sic. Syn. ii. 592 = patula.
macrotheca, Hort. ex Moq. in DC. Prod. xiii. II. 94 = calotheca.
marina, Linn. Mant. ii. 300 = littoralis.
maritima, Crantz, Inst. i. 208 = Suaeda maritima.
maritima, Pall. Reise, ii. 289 = littoralis.
mauritanica, *Boiss. & Reut. Pugill. Pl. Nov.* 106.—Hispan.
micrantha, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 738 = nitens.
micrantha, C. A. Mey. in Ledeb. Fl. Alt. iv. 308 = hastata.
micrantha, *Moq. Chenop. Enum.* 62.—Sibir.
microcarpa, *D. Dietr. Syn. Pl.* v. 536.—Calif.
microcarpa, *Benth. Fl. Austral.* v. 176.—Austral.
microcarpa, Waldst. & Kit. Pl. Rar. Hung. iii. 278. t. 250 = hastata.
microphylla, D. Dietr. Syn. Pl. v. 536 = parvifolia, H. B. & K.
microphylla, Willd. Sp. Pl. iv. 958 = Chenopodium microphyllum.
microphylla, *Phil. Fl. Atac.* 48.—Chili.
microphylla, Schur, Enum. Pl. Transs. 578 = laciniata.
microsperma, Bab. Man. ed. I. 253 = hastata.
microsperma, Waldst. & Kit. Pl. Rar. Hung. iii. 278. t. 250 = hastata.
microtheca, Fries, Corp. Fl. Succ. 133 = hastata.
microtheca, Moq. in DC. Prod. xiii. II. 91 = hortensis.
mollis, *Desf. Fl. Atlant.* ii. 391.—Algeria.
mollis, Uspenski, Jekaterinenburg. 106 = verrucifera.
Moneta, *Bunge, ex Boiss. Fl. Orient.* iv. 912.—Persia; Afghanist.
monilifera, *S. Wats. in Proc. Am. Acad.* ix. (1874) 111.—Am. bor.
monoica, Moench, Meth. 353 = rosea.
montevidensis, *Spreng. Syst.* iii. 918.—Reg. Argent.
Moquiniana, *Webb, ex Moq. in DC. Prod.* xiii. II. 97.—Austral.
mucronata, Rafin. in Am. Monthly Mag. (1818) 176 = arenaria.
mucronata, *Phil. Fl. Atac.* 47.—Chili.
Muelleri, *Benth. Fl. Austral.* v. 175.—Austral.
multifida, Crantz, Inst. i. 207 = Roubieva multifida.
multifida, Desf. Tabl. ed. II. 51 = calotheca.
muralis, Crantz, Inst. i. 206 = Chenopodium murale.
muricata, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 959.—Am. austr.
nemorensis, Schur, Enum. Pl. Transs. 575 = patula.
nitens, *Schkuhr, Handb.* iii. (1803) 541.—Europ.; Orient.
nudicaule, *Boguslaw, in Erman, Archiv*, vi. 65.—Rossia.
nummularia, *Lindl. in Mitch. Journ. Trop. Austr.* (1848) 64.—Austral.
Nuttalii, *S. Wats. in Proc. Am. Acad.* ix. (1874) 116.—Am. bor. occ.
oahuensis, *Meyen, Reise*, ii. 127.—Ins. Sandvic.
oblongifolia, Host, Fl. Austr. i. 320 = hastata.
oblongifolia, Waldst. & Kit. Pl. Rar. Hung. iii. 278. t. 211 = patula.
obovata, *Moq. Chenop. Enum.* 61.—Mexic.
obtusa, *Willd. Sp. Pl.* iv. 964.—Sibiria.
obtusata, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 42.—Hispan.
occidentalis, D. Dietr. Syn. Pl. v. 537 = canescens.
ocymifolia, Viv. Eg. 23 = coriacea.
odorata, Pers. Syn. i. 293 = Exonis albicans.
Olivieri, Moq. Chenop. Enum. 52 = tatarica.
oppositifolia, DC. Rapp. i. 12 = hastata.
oppositifolia, *Vill. Prosp.* 21.—Gallia.
oppositifolia, *S. Wats. in Proc. Am. Acad.* ix. (1874) 118.—Mexic.
orbicularis, *S. Wats. l. c.* xvii. (1881–82) 377.—Am. bor.
palaestina, *Boiss. Diagn.* Ser. I. xii. 96.—Syria; Assyria.
Palmeri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 146.—Calif.
paludosa, *R. Br. Prod.* 406.—Austral.
paludosa, Nees, in Lehm. Pl. Preiss. i. 633 = Drummondii.
pamparum, *Griseb. in Goett. Abh.* xix. (1874) 69.—Reg. Argent.
Parishii, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 377.—Am. bor.
Parryi, *S. Wats. l. c.* 378.—Am. bor.
parviflora, Steud. Nom. ed. II. i. 169 = scq.
parvifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 192.—Mexic.
parvifolia, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 16.—Marocco; Ins. Canar.
patagonica, *D. Dietr. Syn. Pl.* v. 536.—Patagonia.
patula, *Linn. Sp. Pl.* 1053—Reg. temp. bor. et austr.
patula, Sm. Fl. Brit. 1091; Engl. Fl. iv. 257 = hastata.
pedunculata, *Linn. Cent. Pl.* i. 33.—Europ.; As. bor.; Orient.
persica, *Boiss. Diagn.* Ser. I. xii. 96.—Persia.
peruviana, *Moq. in DC. Prod.* xiii. II. 102.—Peruv.
Pharaonis, Hort. Avenion. ex Moq. l. c. 91 = microtheca.
philippica, *Weinm. in Syll. Ratisb.* ii. (1828) 29.—Ins. Philipp.
phyllostegia, *S. Wats. in Proc. Am. Acad.* ix. (1874) 108.—Am. bor. occ.
Piqueres, Lag. Pl. Barrill. 67 = laciniata.
platysepala, Guss. Fl. Sic. Syn. ii. 590 = hastata.
plebeja, *Carmich. in Trans. Linn. Soc.* xii. (1819) 508.—Ins. Trist. d'Acunha.
polycarpa *S. Wats. in Proc. Am. Acad.* ix. (1874) 117.—Calif.
polygama, *Sessé, in Lag. Gen. et Sp. Nov.* 12.—Mexic.
polymorpha, Coss. Germ. et Wedd. Cat. Pl. Par. 108 = hastata, patula.
polysperma, Crantz, Inst. i. 207 = Chenopodium polyspermum.
polysperma, Tenore, Prod. Fl. Nap. i. 57 = rosea.
portulacoides, *Linn. Sp. Pl.* 1053.—Europ.; Syria; Afr. bor.
portulacoides, Pall. Ind. Taur. 59 = verrucifera.
Powellii, *S. Wats. in Proc. Am. Acad.* ix. (1874) 114.—Arizona.
procumbens, Jundz. Fl. Lituan. 78 = patula.
procumbens, *Less. in Linnaea*, ix. (1834) 203.—Sibir. alt.
prostrata, Bouch. Fl. Abbev. 76; Bab. Man. ed. I. 252 = hastata.
prostrata, *R. Br. Prod.* i. 406.—Austral.
prostrata, Moq. in DC. Prod. xiii. II. 99 = cinerea.
prostrata, *Phil. in Anal. Univ. Chil.* (1873) 535.—Chili.
pruinosa, Sieber, ex Moq. in DC. Prod. xiii. II. 93 = laciniata.
pruinosa, Sieber, ex Boiss. Fl. Orient. iv. 910 = tatarica.
Pseudo-nitens, *Schur, Enum. Pl. Transs.* 574.—Hungaria.
Pumilio, *R. Br. Prod.* 406.—Austral.
pungens, *Trautv. in Bull. Soc. Nat. Mosc.* xli. (1867) II. 53.—Sibiria.
Purshiana, *Moq. Chenop. Enum.* 55.—Am. bor.
pusilla, *S. Wats. in Proc. Am. Acad.* ix. (1874) 110.—Am. bor. occ.
ramosissima, Nutt. ex Moq. in DC. Prod. xiii. II. 111 = microcarpa.
recta, Schur, Enum. Pl. Transs. 575 et Ind. = patula.
recurva, Urv. in Mém. Soc. Linn. Par. i. (1822) 284 = tatarica.
reniformis, R. Br. Prod. 406 = paludosa.
reniformis, F. Muell. Fragm. vii. 9 = stipitata.
repens, Aitch. Cat. Pl. Panjab, 125 = Stocksii.
repens, *Roth, Nov. Pl. Sp.* 377.—Ind. or.
retusa, *C. Gay, Fl. Chil.* v. 240.—Chili.
rhagodioides, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 74.—Austral.
rosea, Bab. Man. Brit. ed. I. 253 = crassifolia.
rosea, *Linn. Sp. Pl.* ed. II. 1493.—Europ.; As. Min.; Afr. bor.
rosea, F. Muell. Fragm. vii. 9 = Muelleri.
rotundifolia, *Dombey, ex Moq. Chenop. Enum.* 70.—Peruv.
rotundifolia, Boiss. Fl. Orient. iv. 915 = crystallina.
ruberrima, Hort. ex Moq. in DC. Prod. xiii. II. 91 = hortensis.
rubra, Crantz, Inst. i. 206 = hortensis.
ruderalis, Wallr. Sched. Crit. 115 = hastata.
saccaria, *S. Wats. in Proc. Am. Acad.* ix. (1874) 112.—Utah.
Sackii, Rostc. & Schum. Fl. Sedin. 401. t. 1 = hastata.
sagittata, Borckh. in Reinsch. Mag. 477 = nitens.
salicifolia, Lag. ex Moq. Chenop. Enum. 74 = polygama.
salicina, Pall. Reise, i. 495 = littoralis.
salina, Sievers, in Pall. N. Nord. Beitr. vii. 327 = littoralis.
salina, Desf. in Journ. Bot. i. (1813) 48 = patula.
scoparia, Crantz, Inst. i. 208 = Kochia scoparia.
scmibaccata, *R. Br. Prod.* 406.—Austral.
semibaccata, Moq. in DC. Prod. xiii. II. 97 = spongiosa.
semibracteata, Steud. Nom. ed. II. i. 169, sphalm. = scmibaccata.
serotina, Crantz, Inst. i. 206 = Chenopodium scrotinum.
serpyllifolia, *Bunge, in Act. Hort. Petrop.* v. (1877) 642.—Sibiria; Turkest.
serrata, Huds. Fl. Angl. ed. I. 377 = littoralis.
sibirica, *Linn. Sp. Pl.* ed. II. 1493.—Sibiria.
sicula, *Ucria, in Roem. Arch.* i. 70.—Sicilia.
sinuata, Hoffm. Deutschl. Fl. ed. II. ii. 277 = laciniata.
Smithii, Syme, Eng. Bot. ed. III. viii. 32 = hastata.
spectabilis, Ehrh. ex Moq. in DC. Prod. xiii. II. 91 = hortensis.
spicata, Stokes, Bot. Mat. Med. ii. 24 = Anredera scandens?
spicata, *S. Wats. in Proc. Am. Acad.* ix. (1874) 108.—Calif.
spinacifolia, Stokes, Bot. Mat. Med. ii. 23 = hastata.
spinosa, *D. Dietr. Syn. Pl.* v. 536.—Am. centr.
spongiosa, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 74.—Austral.
stipitata, *Benth. Fl. Austral.* v. 168.—Austral.
stipitata, Westerl. in Linnaea, xl. (1876) 165 = calotheca.
Stocksii, *Boiss. Diagn.* Ser. II. iv. 73.—Scinde.
stylosa, *Viv. Pl. Eg.* 23. f. 21.—Lybia.
subopposita, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 63.—Europ.
Suckleyana, *S. Wats. in Proc. Am. Acad.* ix. (1874) 111.—Am. bor.
sulcata, Michx. ex Roem. & Schult. Syst. vi. 290 = littoralis.
tartarica, Crantz, Inst. i. 205 = tatarica.
tatarica, Auct. fid. Boiss. Fl. Orient. iv. 909 = patula.
tatarica, *Linn. Sp. Pl.* 1053.—As. centr.
tatarica, Schkuhr, Handb. t. 349 = oblongifolia.
Tauschii, *D. Dietr. Syn. Pl.* v. 538.—Hab.?
tetrandra, Torr. ex Moq. in DC. Prod. xiii. II. 93 = Purshiana.
texana, *S. Wats. in Proc. Am. Acad.* ix. (1874) 113.—Texas.
thunbergiaefolia, *Boiss. Fl. Orient.* iv. 911.—Babylonia.
Thunbergiana, Roem. & Schult. Syst. vi. 281 = laciniata.
Tornabeni, Tineo, in Guss. Fl. Sic. Syn. ii. 589 = laciniata.
Torreyana, *Rafin. Atl. Journ.* 146.—Am. bor.
Torreyi, *S. Wats. in Proc. Am. Acad.* ix. (1874) 118.—Am. bor. occ.
transsilvanica, Schur, Enum. Pl. Transs. 577 = hastata.
triangularis, Willd. Sp. Pl. iv. 963 = hastata.
truncata, *A. Gray, in Proc. Am. Acad.* viii. 1873) 398.—Am. bor. occ.
turcomanica, *Fisch. & Mey. ex Karel. in Bull. Soc. Nat. Mosc.* (1839) II. 169.—Persia.
turcomanica, Turcz. ex Ledeb. Fl. Ross. iii. 720 = laciniata.
undulata, *D. Dietr. Syn. Pl.* v. 536.—Patagon.
urbica, Crantz, Inst. i. 206 = Chenopodium urbicum.
velutinella, *F. Muell. Rep. Babb. Exp.* 20.—Austral.
veneta, Moq. Chenop. Enum. 54 = microtheca.
veneta, Willd. Sp. Pl. iv. 962 = tatarica.
Verreauxii, *Moq. in DC. Prod.* xiii. II. 98.—Afr. austr.
verrucifera, *Bieb. Fl. Taur. Cauc.* ii. 441.—Reg. Caucas.; Persia.
verticillata, Lag. Gen. et Sp. Nov. 12 = rosea.
vesicaria, *Heward, ex Benth. Fl. Austral.* v. 172.—Austral.
virgata, Roth, Nov. Sp. Pl. 377 = hortensis.
virgata, Scop. Delic. Insub. ii. 14; Hoffm. Deutschl. Fl. ed. II. ii. 277 = patula.

ATRIPLEX :—
viridis, Crantz, Inst. i. 207 = Chenopodium album.
viridis, Hort. ex Moq. in DC. Prod. xiii. II. 90 = nitens.
Vulvaria, Crantz, Inst. i. 207 = Chenopodium Vulvaria.
Wolfii, *S. Wats. in Proc. Am. Acad.* ix. (1874) 112.—Am. bor.
Wrightii, *S. Wats. l. c.* 113.—N. Mexic.
zosteraefolia, *S. Wats. l. c.* 109.—Am. bor.

ATROPA, Linn. Gen. ed. I. 49 (1737). *SOLANACEAE*, Benth. & Hook. f. ii. 900.
BELLADONA, Adans. Fam. ii. 219 (1763).
acaulis, Stokes, Bot. Mat. Med. i. 396 = Mandragora officinarum.
acuminata, Royle, Illustr. Bot. Himal. 279 = A. Belladonna.
ambigua, Salisb. Prod. 132 = Scopola carniolica.
arborea, Willd. ex Dun. in DC. Prod. xiii. I. 497 = Acnistus arborescens.
arborescens, Linn. Cent. ii. 10 (Amoen. Acad. iv. 307) = Acnistus arborescens.
arborescens, Roem. & Schult. Syst. iv. 685 = Acnistus Plumieri.
arenaria, Willd. ex Roem. & Schult. Syst. iv. 686 = Nectouxia formosa.
aristata, Poir. Encyc. Suppl. i. 609 = Withania aristata.
aspera, Ruiz & Pav. Fl. Peruv. ii. 45 = Hebecladus asperus.
baetica, *Willk. in Linnaea*, xxv. (1852) 50.—Hispan.
Belladonna, *Linn. Sp. Pl.* 181.—Europ.; Orient; Ind. or.
bicolor, Ruiz & Pav. Fl. Peruv. ii. 45 = Hebecladus bicolor.
biflora, Ruiz & Pav. l. c. = Hebecladus biflorus.
contorta, Pers. Syn. i. 219 = Saracha contorta.
daturaefolia, Thore, Chlor. Land. 74 = Nicandra physaloides.
dentata, Pers. Syn. i. 219 = Saracha dentata.
dependens, Mathews, ex Hook. Ic. Pl. t. 107 = Salpichroa dependens.
erecta, Hornem. Hort. Hafn. 957 = Hebecladus biflorus.
erecta, Zuccagni, in Roem. Collect. 129 = Saracha Zuccagniana.
flexuosa, Willd. ex Roem. & Schult. iv. 686; Dun. in DC. Prod. xiii. 465 = Hebecladus viridiflorus.
frutescens, Linn. Sp. Pl. 182 = Withania frutescens.
glandulosa, Hook. Bot. Misc. ii. (1831) 230 = Salpichroa glandulosum.
gymnosperma, Schmid. Ic. 67. t. 18 = Nolana prostrata.
herbacea, *Mill. Gard. Dict.* ed. VIII. n. 3.—Campechia.
hirsuta, Meyen, Reise, i. 466 = Salpichroa hirsutum.
hirtella, *Spreng. Syst.* i. 699.—Bras.
humifusa, Goüan, Hort. Monsp. 106 = Nolana prostrata.
humilis, Salisb. Prod. 132 = Mandragora officinarum.
lethalis, Salisb. l. c. = A. Belladonna.
lutescens, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 241 = A. Belladonna.
Mandragora, Linn. Syst. ed. X. 933 = Mandragora officinarum.
Mandragora, Sibth. & Sm. Fl. Graec. iii. 26. t. 232 = Mandragora autumnalis.
origanifolia, Desf. Cat. Hort. Par. ed. III. 396 = Physalis curasavica, origanifolia.
physalodes, Linn. Sp. Pl. 181 = Nicandra physaloides.
physaloides, Georgi, Beschr. Russ. Reich. Nachtr. 261 = Scopola physaloides.
physaloides, Linn. Syst. ed. XII. 171 = *physalodes*.
plicata, Roth, Catalecta, ii. 24 = Saracha procumbens.
procumbens, Cav. Ic. i. 53 = Saracha procumbens.
punctata, Pers. Syn. i. 219. = Saracha punctata.
ramosissima, Mathews, ex Dunal, in DC. Prod. xiii. I. 473 = Salpichroa ramosissimum.
reflexa, Walp. Rep. iii. 103, in syn. = seq.
revoluta, F. G. Dietr. Vollst. Gaertn. Lexik. Nachtr. i. 422 = Hebecladus umbellatus.
rhomboidea, Gill. & Hook. in Hook. Bot. Misc. i. (1830) 135 = Salpichroa rhomboideum.
Rothii, Poir. Encyc. Suppl. i. 610 = Hebecladus umbellatus.
sideroxyloides, Willd. ex Roem. & Schult. Syst. iv. 486 = Acnistus arborescens.

ATROPA :—
solanacea, All. ex Steud. Nom. ed. II. i. 170 = Acnistus arborescens.
solanacea, Linn. Mant. ii. 205 = Solanum aggregatum.
spinosa, Meyen, Reise, i. 466 = Dunalia lycioides
umbellata, Ruiz & Pav. Fl. Peruv. ii. 44 = Hebecladus umbellatus.
villosa, Zuccagni, in Roem. Collect. 130 = Hebecladus umbellatus.
viridiflora, H. B. & K. Nov. Gen. et Sp. iii. 11 = Hebecladus viridiflorus.

ATROPIS, Rupr. Fl. Samoj. [Beitr. Fl. Russ. Reich. ii.] 64 (1845) = **Glyceria**, R. Br. (Gramin.).
angustata, Griseb. in Ledeb. Fl. Ross. iv. 390 = G. distans.
californica, Munro, ex A. Gray, in Proc. Acad. Sc. Philad. '1862' (1863) 336.—Am. bor. occ.
capillaris, Schur, Enum. Pl. Transs. 779 = G. distans.
carinata, Griseb. in Goett. Abh. xxiv. (1879) 291.—Reg. Argent.
convoluta, Griseb. in Ledeb. Fl. Ross. iv. 389 = G. distans.
distans, Rupr. Fl. Samoj. 64; Griseb. in Ledeb. Fl. Ross. iv. 388 = G. distans.
festucaeformis, Schur, Enum. Pl. Transs. 780 = G. maritima.
intermedia, Schur, l. c. 779 = G. maritima.
magellanica, E. Desv. in C. Gay, Fl. Chil. vi. 394 = G. magellanica.
maritima, Griseb. in Ledeb. Fl. Ross. iv. 389 = G. maritima.
pauciflora, Thurb. in S. Wats. Bot. Calif. ii. 310 = Poa pauciflora.
procumbens, Thurb. l. c. 309.—Am. bor.
Pseudo-maritima, Schur, Enum. Pl. Transs. 780 = G. maritima.
pumila, T. Kirk, in Gard. Chron. (1882) I. 261; Trans. N. Z. Inst. xiv. (1882) 379.—N. Zel.
scabrella, Thurb. in S. Wats. Bot. Calif. ii. 310 = Poa scabrella.
tenuiflora, Griseb. in Ledeb. Fl. Ross. iv. 389 = G. distans.
tenuifolia, Thurb. in S. Wats. Bot. Calif. ii. 310 = G. tenuifolia.
transsilvanica, Schur, Enum. Pl. Transs. 889, Index = G. maritima.

ATRUTEGIA, *Bedd. in Madr. Journ.* Ser. III. i. (1864) 37. f. 1. *ANONACEAE*, Benth. & Hook. f. Gen. i. 957.
wynadensis, *Bedd. l. c.*—Ind. or.

ATTALEA, H. B. & K. Nov. Gen. et Sp. i. 309 (1815). *PALMAE*, Benth. & Hook. f. iii. 947.
acaulis, *Hort. ex H. Wendl. Cycl. Pand. Cycad.* 4.—Hab.?
agrestis, Barb.-Rodr. Enum. Palm. Nov. 42 = microcarpa.
amygdalina, *H. B. & K. Nov. Gen. et Sp.* i. 310. t. 95.—Am. austr.
blepharopus, *Mart. in Palmet. Orb.* 116. t. 5. f. 2.—Bolivia.
butyrosa, Lodd. ex H. Wendl. in Kerch. Palm. 232 = humilis.
Cephalotus, Poepp. ex Mart. in Palmet. Orb. 119 = Scheelea Cephalotes.
Cohune, *Mart. l. c.* 121; *et Hist. Nat. Palm.* iii. 300. t. 167.—Honduras.
compta, *Mart. Hist. Nat. Palm.* ii. 137. t. 41, 75, 97.—Bras.
coronata, *Lodd. ex H. Wendl. Cycl. Pand. Cycad.* 5.—Hab.?
excelsa, *Mart. Hist. Nat. Palm.* ii. 138. t. 96. f. 3.—Bras.
exigua, *Drude, in Mart. Fl. Bras.* iii. II. 439.—Bras.
funifera, *Mart. Hist. Nat. Palm.* ii. 136. t. 95, 96. f. 4.—Bras.
gomphococca, *Mart. l. c.* 301.—Am. austr.
grandis, *Hort. ex H. Wendl. in Kerch. Palm.* 233.—Hab.?
Humboldtiana, *Spruce, Palm. Amaz.* 163.—Am. trop.
humilis, *Mart. in Spreng. Syst.* ii. 624.—Bras.
limbata, *Seem. ex H. Wendl. in Kerch. Palm.* 233.—Hab.?
macrocarpa, *Linden, Illustr. Hortic.* xxviii. (1881) 15.—Bras.

ATTALEA :—
Magdalenae, *Linden, Illustr. Hortic.* xxviii. (1881) 15.—N. Granat.
Manaca, *Linden, l. c.*—N. Granat.
maracaibensis, *Mart. in Palmet. Orb.* 124.—Peruv.
Maripa, Mart. l. c. 123 = Maximiliana Maripa.
microcarpa, *Mart. l. c.* 125.—Bras.
monosperma, Barb.-Rodr. Enum. Palm. Nov. 42 = spectabilis.
nucifera, Karst. in Linnaea, xxviii. (1856) 255 = amygdalina.
phalerata, *Mart. in Spreng. Syst.* ii. 624.—Bras.
Pixuna, Barb.-Rodr. Enum. Palm. Nov. 43 = spectabilis.
Princeps, Mart. in Palmet. Orb. 113. t. 4. f. 2 = Scheelea Princeps.
purpurea, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—N. Granat.
puruënsis, *Linden, l. c.*—Fl. Amazon.
racemosa, Spruce, Palm. Amaz. 166 = Orbignia racemosa.
Rossii, *Lodd. ex Loud. Hort. Brit.* 387.—Bras.
rostrata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 50.—Costa Rica.
speciosa, *Mart. Hist. Nat. Palm.* ii. 138.—Bras.
spectabilis, *Mart. l. c.* 136. t. 96. f. 1, 2.—Bras.
spinosa, *Meyen, Reise*, i. 469.—Peruv.
Tiasse, *Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Fl. Amazon.
transitiva, *Barb.-Rodr. Prot. App.* 49.—Bras.
venatorum, Mart. Hist. Nat. Palm. iii. 325 = Maximiliana venatorum.

ATTRACTILIS, Hall. ex Scop. Introd. (1777) 124 = **Atractylis**, Linn. (Compos.).

ATULANDRA, Rafin. Sylva Tellur. 31 (1838) = **Rhamnus**, Linn.
arragonensis, Rafin. l. c. = R. lycioides.
valentina, Rafin. l. c. = R. pumila.

ATUNA, Rafin. Sylva Tellur. 153 (1838) = **Heritiera**, Dryand. (Stercul.).
racemosa, Rafin. l. c. = H. littoralis.

ATYLOSIA, Wight & Arn. Prod. 257 (1834). *LEGUMINOSAE*, Benth. & Hook. f. i. 542.
CANTHAROSPERMUM, Wight & Arn. Prod. 255 (1834).
albicans, *Benth. in Miq. Pl. Jungh.* 243.—Ind. or.
barbata, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 216.—Ind. or.; Malaya.
calycina, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 186.—Ind. or.
candicans, Kurz, l. c. = Rhynchosia arvensis.
Candollei, *Wight & Arn. Prod.* i. 257.—Ind. or.
cinerea, *F. Muell. ex Benth. Fl. Austral.* ii. 264.—Austral.
circinnalis, Benth. in Miq. Pl. Jungh. 244 = Dunbaria circinalis.
elongata, *Benth. l. c.* 243.—Ind. or.
geminiflora, *Dalz. in Journ. Linn. Soc.* xiii. (1873) 185.—Ind. or.
glandulosa, Dalz. l. c. = mollis.
goënsis, Dalz. l. c. 186 = barbata.
grandiflora, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 214.—Reg. Himal.
grandifolia, *F. Muell. ex Benth. Fl. Austral.* ii. 264.—Austral.
kulnensis, *Dalz. in Journ. Linn. Soc.* xiii. (1873) 186.—Ind. or.
Lawii, *Dalz. l. c.*—Ind. or.
Lawii, Wight, Ic. i. t. 93 = lineata.
lineata, *Wight & Arn. Prod.* i. 258.—Ind. or.
major, Wight & Arn. l. c. 257 = Candollei.
marmorata, *Banks, ex Benth. Fl. Austral.* ii. 263.—Austral.
mollis, *Benth. in Miq. Pl. Jungh.* 243.—Ind. or.; Malaya.
nivea, *Benth. l. c.*—Burma.
platycarpa, *Benth. l. c.*—Reg. Himal.
pluriflora, *F. Muell. ex Benth. Fl. Austral.* ii. 264.—Austral.
punctata, Dalz. in Journ. Linn. Soc. xiii. (1873) 186 = Dunbaria conspersa.
reticulata, *Benth. Fl. Austral.* ii. 263.—Austral.
rostrata, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 216.—Ind. or.
rugosa, *Wight & Arn. Prod.* i. 257.—Ind. or.

ATYLOSIA:—
scarabaeoides, *Benth. in Miq. Pl. Jungh.* 242.—Austral.
sericea, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 213.—Ind. or.
subrhombea, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 51 = Dunbaria subrhombea.
villosa, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 214.—Reg. Himal.

ATYLUS, Salisb. Parad. Lond. sub t. 67 (1807) = **Petrophila**, R. Br. (Proteac.).
anemonifolia, Salisb. l. c. = Isopogon anemonifolius.
fucifolia, Salisb. l. c. = P. pulchella.

AUBENTONIA, Domb. ex Steud. Nom. ed. I. 93 (1821) = **Waltheria**, Linn. (Stercul.).
tomentosa, Domb. ex Steud. l. c. = W. ovata.

AUBERTIA, Bory, Voy. i. 356. t. 18 (1804) = **Zanthoxylum**, Linn.? (Rutac.).
Bajarnardia, Buch.-Ham. in Wall. Cat. sub n. 7118 = Z. Bajarnardia.
borbonica, Bory, Voy. i. 356. t. 18 (an Enodiae sp.?—Cf. Benth. & Hook. f. Gen. i. 296) = Z. Aubertia.
piperita, Buch.-Ham. ex Wall. Cat. sub n. 7117 = Z. Hamiltonianum.
Timbor, Buch.-Ham. ex Wall. l. c. n. 7116 = Z. acanthopodium.
tomentosa, Bory, ex Steud. Nom. ed. II. i. 171 = Z. Aubertia.

AUBERTIA, Chapel. ex Baill. Adansonia, i. (1861) 162, 166 = **Croton**, Linn. (Euphorb.).
argentea, Chapel. ex Baill. l. c. 162 = C. Noronhae.
glandulosa, Chapel. ex Baill. l. c. 166 = C. Chapelieri.

AUBION, Rafin. Sylva Tellur. 110 (1838) = **Cleome**, Linn. (Capparid.).
Chelidoni, Rafin. l. c. = C. Chelidonii.

AUBLETIA, Gaertn. Fruct. i. 379. t. 78. f. 2 (1788) = **Sonneratia**, Linn. f. (Lythrar.).
caseolaris, Gaertn. l. c. = S. acida.

AUBLETIA, Lour. Fl. Cochinch. 283 (1790) = **Paliurus**, Tourn. (Rhamnac.).
ramosissima, Lour. l. c. = P. ramosissimus.

AUBLETIA, Neck. Elem. i. 356 (1790) = **Ruellia**, Linn. (Acanth.).

AUBLETIA, Rich. in Pers. Syn. ii. 638 (1807) = **Monnieria**, Linn. (Rutac.).
trifoliata, Pers. l. c. = M. trifolia.

AUBLETIA, Schreb. Gen. 353 (1789) = **Apeiba**, Aubl. (Tiliac.).
aspera, Willd. Sp. Pl. ii. 1156 = Apeiba aspera.
discolor, Spreng. Neue Entdeck. ii. 166 = Apeiba discolor.
laevis, Willd. Sp. Pl. ii. 1156 = Apeiba glabra.
Petouma, Willd. l. c. = Apeiba Petouma.
Tibourbou, Willd. l. c. = Apeiba Tibourbou.
Tiburbu, Sw. Fl. Ind. Occ. ii. 944 = praec.

AUBRIETIA, Adans. Fam. ii. 420 (1763) (Aubrieta); DC. in Mém. Mus. Par. vii. (1821) 232; Syst. ii. 293. *CRUCIFERAE*, Benth. & Hook. f. Gen. i. 73.
canescens, Boiss. Fl. Orient. i. 252 = deltoidea.
Columnae, *Guss. Pl. Rar.* 266.—Ital.
croatica, *Schott, Nyman & Kotschy, Analect. Bot.* 47.—Croatia.
deltoidea, *DC. Syst.* ii. 294.—Italia; Graecia; As. Min.
diffusa, Spreng. Syst. ii. 871 = Heliophila diffusa.
edentula, *Boiss. Fl. Orient.* i. 254.—Kurdistan.
erubescens, *Griseb. Spicil. Fl. Rumel.* i. 268.—Graecia.
floribunda, Spach, Hist. Vég. Phan. vi. 469 = deltoidea.
gracilis, *Sprun. ex Boiss. Diagn.* Ser. I. i. 74.—Graecia.
graeca, Griseb. Spicil. Fl. Rumel. i. 268 = deltoidea.
hesperidiflora, G. Don, ex Loud. Hort. Brit. 257 = deltoidea.
integrifolia, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 35 = deltoidea.
intermedia, *Heldr. & Orph. ex Boiss. Diagn.* Ser. II. i. 36.—Graecia.

AUBRIETIA:—
Kotschyi, *Boiss. & Hohen. Diagn.* Ser. I. viii. 32.—Persia.
italica, Boiss. Fl. Orient. i. 252, in nota = Columnae.
lepidioides, Spreng. Syst. ii. 871 = Heliophila diffusa.
libanotica, *Boiss. & Hohen. Diagn.* Ser. I. viii. 32.—Mont. Libanus.
olympica, *Boiss. Fl. Orient.* i. 251.—Bithynia.
parviflora, *Boiss. Diagn.* Ser. I. vi. 14.—Persia.
Pinardi, *Boiss. l. c.* Ser. II. i. 37.—As. Min.
purpurea, DC. Syst. ii. 294 = Arabis purpurea.

AUBRYA, Baill. Adansonia, ii. 265 (1861-62). *HUMIRIACEAE*, Benth. & Hook. f. i. 988.
gabonensis, *Baill. l. c.* 266.—Afr. trop.

AUCHERA, DC. Coll. Mém. ix. t. 11; Prod. vi. 557 (1837) = **Cousinia**, Cass. (Compos.).
floribunda, Regel, Cat. Pl. Hort. Aksakov. 13.—Hab.?
orientalis, DC. Coll. Mém. ix. t. 11 = C. orientalis.

AUCKLANDIA, Falc. in Trans. Linn. Soc. xix. (1845) 23 = **Saussurea**, DC. (Compos.).
Costus, Falc. l. c. = S. Lappa.

AUCUBA, Thunb. Nov. Gen. 61 (1783). *CORNACEAE*, Benth. & Hook. f. i. 950.
EUBASIS, Salisb. Prod. 68 (1796).
chinensis, *Benth. Fl. Hongk.* 138.—China.
himalaica, *Hook. f. & Thoms. Illustr. Himal. Pl.* t. 12.—Reg. Himal.
japonica, *Thunb. Nov. Gen.* 61.—Japon.

AUCUBA, Cham. in Linnaea, v. (1830) 56, in syn. = ARUBA, Nees & Mart. = **Raputia**, Aubl. (Rutac.).

AUCUBAEPHYLLUM, Ahlburg, in Bot. Zeit. xxxvi. (1878) 113. *CORNACEAE*.
lioukiense, *Ahlburg, l. c.*—Japon.

AUCUPARIA, Riv. ex Rupp. Fl. Jen. ed. Haller. 140 (1745) = **Pyrus**, Linn. (Rosac.).
pinnata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 378 = P. Aucuparia.
sylvestris, Medic. Gesch. 86 = P. Aucuparia.

AUDIBERTIA, Benth. in Bot. Reg. sub t. 1282 (1829) = **Mentha**, Linn. (Labiat.).
parviflora, Nym. Consp. 597 = M. Requieni.
pusilla, Benth. in Bot. Reg. sub t. 1282 = M. Requieni.

AUDIBERTIA, Benth. in Bot. Reg. t. 1469 (1831). *LABIATAE*, Benth. & Hook. f. ii. 1197.
capitata, *A. Gray, in Proc. Am. Acad.* vii. (1868) 387.—Calif.
Clevelandi, *A. Gray, l. c.* x. (1874) 76.—Calif.
decurrens, Nutt. ex Benth. in DC. Prod. xii. 359 = grandiflora.
Dorrii, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 190.—Calif.
grandiflora, *Benth. Lab. Gen. et Sp.* 312.—Calif.
humilis, *Benth. l. c.* 313.—Calif.
incana, *Benth. in Bot. Reg.* t. 1469.—Am. bor. occ.
nivea, *Benth. Lab. Gen. et Sp.* 313.—Calif.
Palmeri, *A. Gray, in Bot. Calif.* i. 601.—Calif.
polystachya, *Benth. Lab. Gen. et Sp.* 314.—Calif.
spinulosa, Nutt. ex Benth. in DC. Prod. xii. 359 = stachyoides.
stachyoides, *Benth. Lab. Gen. et Sp.* 313.—Calif.
Vaseyi, *T. C. Porter, in Coult. Bot. Gazette*, vi. (1881) 207.—Calif.

AUDOUINIA, Brongn. in Ann. Sc. Nat. viii. (1826) 384. t. 38. *BRUNIACEAE*, Benth. & Hook. f. i. 673.
PAVINDA, Thunb. ex Bartl. Ord. 374 (1830).
capitata, *Brongn. l. c.*—Afr. austr.

AUGANTHUS, Link, Handb. i. 415 (1829) = **Primula**, Linn.
praenitens, Link, l. c. = P. sinensis.

AUGEA, Thunb. ex Retz. Obs. v. 3 (1789) = **Lanaria**, Ait. (Haemod.).
lanata, Thunb. ex Retz. l. c. = L. plumosa.

AUGEA, Thunb. Prod. Pl. Cap. 80 (1794); et Nov. Gen. 132 (1798). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 265.
capensis, *Thunb. ll. cc.* 80 et 133.—Afr. austr.

AUGIA, Lour. Fl. Cochinch. 337 (1790). 'Genus ignotum,' Benth. & Hook. f. Gen. i. 418.
sinensis, Lour. l. c. (= *Calophyllum Augia* ex Steud. Nom. ed. II. i. 171).—China.

AUGUSTA, Leandr. Sacram. in Denkschr. Akad. Muench. vii. (1819) 235. t. 14 = **Stifftia**, Mikan (Compos.).
grandiflora, Leandr. Sacram. l. c. = S. chrysantha.
parviflora, Leandr. Sacram. l. c. = S. parviflora.

AUGUSTA, Pohl, Pl. Bras. Ic. ii. 1. t. 101–105. *RUBIACEAE*, Benth. & Hook. f. ii. 51.
BONIFACIA, Manso, ex Steud. Nom. ed. II. i. 216 (1840).
SCHREIBERSIA, Pohl, in Flora, viii. (1825) 183.
attenuata, *Pohl, Pl. Bras. Ic.* ii. t. 104.—Bras.
glaucescens, *Pohl, l. c.* t. 105.—Bras.
lanceolata, *Pohl, l. c.* t. 101.—Bras.
oblongifolia, *Pohl, l. c.* t. 103.—Bras.
parvifolia, *Pohl, l. c.* t. 102.—Bras.

AUGUSTEA, DC. Prod. iv. 404 (1830) = **Augusta**, Pohl (Rubiac.).

AUGUSTIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 124; et Abh. Akad. Berl. (1854) 80. t. 8. B. = **Begonia**, Linn.
caffra, Klotzsch, ll. cc. 124 & 81 = B. caffra.
Dregei, Klotzsch, ll. cc. 124 & 80 = B. Dregei.
natalensis, Klotzsch, ll. cc. 181 & 80 = B. natalensis.
suffruticosa, Klotzsch, ll. cc. 124 & 82 = B. suffruticosa.

AUGUSTINEA, Karst. in Linnaea, xxviii. (1856) 395 = **Bactris**, Jacq. (Palm.).
balanoidea, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 39 = B. balanoidea.
major, Karst. in Linnaea, xxviii. (1856) 395 = B. major.
ovata, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 38 = B. ovata.

AUGUSTINEA, St. Hil. & Naud. in Ann. Sc. Nat. Sér. III. ii. (1844) 145 = **Miconia**, Ruiz & Pav. (Melast.).
speciosa, St. Hil. & Naud. l. c. = M. speciosa.

AULACIA, Lour. Fl. Cochinch. 273 (1790) = **Clausena**, Burm. f. (Rutac.).
falcata, Lour. Fl. Cochinch. 273 (= *Cookia falcata*).—Cochinch.
falcifolia, Stokes, Bot. Mat. Med. ii. 482 = praec.
punctata, Raeusch. Nom. ed. III. 119 = Cl. Wampi.

AULACIDIUM, Rich. ex DC. Prod. iii. 112 (1828) = **Macrocentrum**, Hook. f. (Melast.).
cristatum, Rich. ex DC. l. c. 113 = M. cristatum.
fasciculatum, Rich. ex DC. l. c. = M. fasciculatum.

AULACINTHUS, E. Mey. Comm. Pl. Afr. Austr. 155 (1835) = **Lotononis**, Eckl. & Zeyh. (Legum.).
gracilis, E. Mey. l. c. = L. gracilis.
rigidus, E. Mey. l. c. = L. rigidus.

AULACOCALYX, Hook. f. in Hook. Ic. Pl. t. 1126 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 109.
jasminiflora, *Hook. f. l. c.*—Afr. trop.

AULACOCARPUS, Berg, in Linnaea, xxvii. ('1854') 345. *MYRTACEAE*, Benth. & Hook. f. i. 720.
crassifolius, *Berg, l. c.* 346.—Ins. Gorgona.
quadrangularis, *Griseb. Fl. Brit. W. Ind.* 239.—Ind. occ.
Sellowianus, *Berg, in Linnaea*, xxvii. ('1854') 346.—Bras.
Wrightii, *Griseb. Cat. Pl. Cub.* 90.—Cuba.

AULACODISCUS, Hook. f. in Benth. & Hook. f. Gen. ii. 71 (1873) (*non Ehrenb. quae Crypt.*). *RUBIACEAE*, Benth. & Hook. f. ii. 71.
premnoides, *Hook. f. l. c.*—Malaya.

AULACOPHYLLUM, Regel, Gartenfl. (1876) 141 = **Zamia**, Linn. (Cycad.).
Lindeni, Regel, l. c. = Z. Lindeni.
montanum, Regel, l. c. = Z. montana.
Ortgiesi, Regel, l. c. = Z. Chigua.
Roezli, Regel, l. c. = Z. pseudo-parasitica.
Skinneri, Regel, l. c. 143 = Z. Skinneri.
Wallisii, Regel, l. c. = Z. Wallisii.

AULACORHYNCHUS, Nees, in Linnaea, ix. 305 (1834); ib. x. 198 (1835). *CYPERACEAE*, Benth. & Hook. f. Gen. iii. 1043.
crinifloras, Steud. Nom. ed. II. i. 171 = seq.
crinifolius, *Nees, in Linnaea*, ix. 305; et x. 199.—Afr. austr.

AULACOSPERMUM, Ledeb. Fl. Alt. iv. 344 (1833) = **Pleurospermum**, Hoffm. (Umbell.).
anomalum, Ledeb. l. c. 335.—Mont. Altai.
cuneatum, Ledeb. l. c.—Mont. Altai.
multifidum, Meinsh. in Linnaea, xxx. (1859–60) 515. —Sibir.
simplex, Rupr. Sert. Tiansch. 49 = P. stellatum.
tenuilobum, Meinsh. in Linnaea, xxx. (1859–60) 515. —Sibir.

AULACOSTIGMA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 149 = **Rhynchotheca**, Ruiz & Pav. (Geran.).
inerme, Turcz. l. c. 150.—Reg. Andina.

AULAX, Berg. Descr. Pl. Cap. 33 (1767). *PROTEACEAE*, Benth. & Hook. f. iii. 168.
cneorifolia, *Knight, Prot.* 15.—Afr. austr.
pinifolia, *Berg. Descr. Pl. Cap.* 33.—Afr. austr.
umbellata, R. Br. in Trans. Linn. Soc. x. (1810) 50 = cneorifolia.

AULAXANTHUS, Ell. Sketch, i. 102 (1817) = **Anthaenantia**, Beauv. (Gramin.).
ciliatus, Ell. l. c. = Anthaenantia villosa.
rufus, Ell. l. c. t. 6 = Anthaenantia rufa.

AULAXIA, Nutt. Gen. Am. i. 47 (1818) = **Anthaenantia**, Beauv. (Gramin.).
ciliata, Nutt. l. c. = Anthaenantia villosa.
rufa, Nutt. l. c. = Anthaenantia rufa.

AULAXIS, Haw. Enum. Saxif. 46 (1821) = **Saxifraga**, Linn.
micranthifolia, Haw. l. c. 47 = S. erosa.
nuda, Haw. l. c. = S. erosa.

AULAXIS, Steud. Nom. ed. II. i. 171 (1840) = AULAYIA, Nutt. = **Anthaenantia**, Beauv. (Gramin.).

AULAYA, Harv. Gen. Pl. Afr. Austr. 249 (1838) = **Harveya**, Hook. (Scrophul.).
capensis, Harv. l. c. 250 = H. capensis.
coccinea, Harv. Thes. Cap. i. 23.—Afr. austr.
grandiflora, Benth. in DC. Prod. x. 523.—Afr. austr.
obtusifolia, Benth. l. c. = H. obtusifolia.
pauciflora, Benth. l. c. 524.—Afr. austr.
purpurea, Benth. l. c. 523 = H. purpurea.
scarlatina, Benth. l. c. 524.—Afr. austr.
squamosa, Harv. Gen. Pl. Afr. Austr. 250 = H. squamosa.

AULEYA, D. Dietr. Syn. Pl. iii. 539 (1843) = praec.

AULICA, Rafin. Fl. Tellur. iv. 10 (1836) = **Hippeastrum**, Herb. (Amaryll.).
latifolia, Rafin. l. c. = H. aulicum.
platypetala, Rafin. l. c. = H. aulicum.
striata, Rafin. l. c. = H. aulicum.

AULIPHAS, Rafin. Sylva Tellur. 95 (1838) = **Miconia**, Ruiz & Pav. (Melast.).
ligustrina, Rafin. l. c. = M. ligustrioides.

AULIZA, Salisb. in Trans. Hort. Soc. i. (1812) 294 = **Epidendrum**, Linn. (Orchid.).
ciliaris, Salisb. l. c. = E. ciliare.

AULOMYRCIA, Berg, in Linnaea, xxvii. (1854) 35 = **Myrcia**, DC. (Myrtac.).
Abrantea, Berg, l. c. xxx. (1859–60) 659.—Bras.
acetosans, Berg, l. c. 662.—Ind. occ.
acrantha, Berg, in Mart. Fl. Bras. xiv. i. 71.—Bras.
acutata, Berg, l. c. = M. acutata.
acutifolia, Berg, l. c. 89.—Bras.
Aethusa, Berg, l. c. 112.—Bras.
alagoënsis, Berg, l. c. 120 = M. alagoënsis.
albo-tomentosa, Berg, in Linnaea, xxvii. (1854) 62 = M. albo-tomentosa.
alloiota, Berg, l. c. 54.—Bras.
alternifolia, Berg, l. c. 72 = M. alternifolia.
amazonica, Berg, l. c. 41 = M. amazonica.

AULOMYRCIA:—
amethystina, Berg, in Mart. Fl. Bras. xiv. i. 108.—Bras.
andromedoides, Berg, in Linnaea, xxvii. (1854) 40 = Myrcia andromedoides.
androsaemoides, Berg, l. c. xxx. (1859–60) 661.—Guiana.
angustifolia, Berg, in Mart. Fl. Bras. xiv. i. 135.—Bras.
anomala, Berg, l. c. 96 = M. anomala.
apiocarpa, Berg, l. c. 64.—Bras.
atropilosa, Berg, l. c. 101.—Bras.
atrovirens, Berg, l. c. 121.—Bras.
aureo-lanata, Berg, in Linnaea, xxvii. (1854) 52.—Bras.
badia, Berg, in Mart. Fl. Bras. xiv. i. 547.—Bras.
bahiensis, Berg, l. c. 113.—Bras.
bella, Berg, l. c. 71 = M. bella.
bicarinata, Berg, l. c. 118.—Bras.
bicudoënsis, Berg, l. c. 557.—Bras.
biformis, Berg, l. c. 141.—Bras.
Billardiana, Berg, in Linnaea, xxvii. (1854) 51 = M. Billardiana.
bimarginata, Berg, in Mart. Fl. Bras. xiv. i. 115.—Bras.
Blanchetiana, Berg, l. c. 65.—Bras.
bombycina, Berg, l. c. 66.—Bras.
Botrys, Berg, l. c. 116.—Bras.
bracteata, Berg, l. c. 554 = M. bracteata.
breviramis, Berg, l. c. 66.—Bras.
bullata, Berg, l. c. 96 = M. bullata.
buxifolia, Berg, l. c. 80 = M. buxifolia.
buxizans, Berg, in Linnaea, xxx. (1859–60) 664.—Guiana.
caerulescens, Berg, in Mart. Fl. Bras. xiv. i. 80.—Bras.
caesia, Berg, l. c. 83.—Bras.
calyptranthoides, Berg, l. c. 67.—Bras.
Camaraeana, Berg, l. c. 83 = M. Camaraeana.
Cambessedeana, Berg, in Linnaea, xxvii. (1854) 40 = M. Cambessedeana.
campestris, Berg, l. c. 74 = M. campestris.
capitata, Berg, in Mart. Fl. Bras. xiv. i. 554 = M. capitata.
capivarhyensis, Berg, in Linnaea, xxvii. (1854) 36 = M. capivarhyensis.
caracasana, Berg, l. c. 53 = M. caracasana.
cardiophylla, Berg, in Mart. Fl. Bras. xiv. i. 103 = M. cardiophylla.
cassinoides, Berg, in Linnaea, xxvii. (1854) 74 = M. cassinoides.
castrensis, Berg, in Mart. Fl. Bras. xiv. i. 111.—Bras.
chapadensis, Berg, l. c. 554.—Bras.
chilensis, Berg, in Linnaea, xxvii. (1854) 38 = M. chilensis.
chrysophylla, Berg, in Mart. Fl. Bras. xiv. i. 125.—Bras.
Clausseniana, Berg, l. c. 118.—Bras.
collina, Berg, in Linnaea, xxvii. (1854) 40.—Bras.
comosa, Berg, in Mart. Fl. Bras. xiv. i. 133.—Bras.
conduplicata, Berg, in Linnaea, xxvii. (1854) 76.—Guiana.
confusa, Berg, l. c. 56.—Bras.
coriacea, Berg, l. c. 70 = M. coriacea.
corymbiflora, Berg, l. c. 74.—Bras.
corymbosa, Berg, l. c. 41 = M. corymbosa.
costata, Berg, in Mart. Fl. Bras. xiv. i. 79 = M. costata.
Coumeta, Berg, in Linnaea, xxvii. (1854) 60 = M. Coumeta.
crassicaulis, Berg, l. c. 74 = M. crassicaulis.
crassifolia, Berg, in Mart. Fl. Bras. xiv. i. 128.—Bras.
crenulata, Berg, l. c. 141.—Bras.
cuneata, Berg, in Linnaea, xxvii. (1854) 72.—Venezuela.
cuprea, Berg, in Mart. Fl. Bras. xiv. i. 77.—Bras.
curatellaefolia, Berg, in Linnaea, xxvii. (1854) 55 = M. curatellaefolia.
cymosa, Berg, in Mart. Fl. Bras. xiv. i. 552.—Bras.
daphnoides, Berg, in Linnaea, xxvii. (1854) 77 = M. daphnoides.
dealbata, Berg, l. c. 61 = M. dealbata.
debilis, Berg, l. c. 62 = M. debilis.
decrescens, Berg, in Mart. Fl. Bras. xiv. i. 135.—Bras.
densa, Berg, in Linnaea, xxvii. (1854) 37 = Calyptranthes densa.
desertorum, Berg, in Mart. Fl. Bras. xiv. i. 556.—Bras.
detergens, Berg, in Linnaea, xxvii. (1854) 46 = M. detergens.

AULOMYRCIA:—
diaphana, Berg, in Mart. Fl. Bras. xiv. i. 82.—Bras.
dichroma, Berg, in Linnaea, xxvii. (1854) 65.—Guiana.
dictyophleba, Berg, in Mart. Fl. Bras. xiv. i. 560.—Bras.
dictyophylla, Berg, l. c. 72.—Bras.
dimorpha, Berg, l. c. 101.—Bras.
divaricata, Berg, in Linnaea, xxvii. (1854) 58 = M. divaricata.
Doniana, Berg, in Mart. Fl. Bras. xiv. i. 516 = M. Doniana.
dumosa, Berg, in Linnaea, xxx. (1859–60) 656.—Guiana.
edulis, Berg, l. c. 657.—Guiana.
egensis, Berg, in Mart. Fl. Bras. xiv. i. 99.—Bras.
elaeodendra, Berg, in Linnaea, xxvii. (1854) 75 = M. elaeodendra.
emarginata, Berg, in Mart. Fl. Bras. xiv. i. 134.—Bras.
eumecophylla, Berg, l. c. 98.—Bras.
exsucca, Berg, in Linnaea, xxvii. (1854) 79 = M. exsucca.
Fenzliana, Berg, l. c. 63 = M. Fenzliana.
ferruginea, Berg, in Mart. Fl. Bras. xiv. i. 552 = M. ferruginea.
fragilis, Berg, l. c. 117.—Bras.
Gardneriana, Berg, l. c. 129 = M. Gardneriana.
garopabensis, Berg, in Linnaea, xxvii. (1854) 47 = M. garopabensis.
Gaudichaudiana, Berg, in Mart. Fl. Bras. xiv. i. 88.—Bras.
gigantea, Berg, l. c. 548.—Bras.
glabra, Berg, l. c. 119.—Bras.
glandulosa, Berg, in Linnaea, xxvii. (1854) 79.—Bras.
glauca, Berg, l. c. 50 = M. glauca.
glaucescens, Berg, in Mart. Fl. Bras. xiv. i. 81.—Bras.
goyazensis, Berg, l. c. 85 = M. goyazensis.
grandiflora, Berg, l. c. 113.—Bras.
grandifolia, Berg, l. c. 97 = M. grandifolia.
hepatica, Berg, l. c. 132.—Bras.
hirtiflora, Berg, in Linnaea, xxvii. (1854) 40 = M. hirtiflora.
holosericea, Berg, in Mart. Fl. Bras. xiv. i. 99.—Bras.
Hostmanniana, Berg, in Linnaea, xxvii. (1854) 42.—Guiana.
hypericoides, Berg, l. c. 62 = M. hypericoides.
imbricata, Berg, l. c. 61 = M. imbricata.
insularis, Berg, in Mart. Fl. Bras. xiv. i. 98 = M. insularis.
intermedia, Berg, l. c. 107.—Bras.
Jacquiniana, Berg, in Linnaea, xxvii. (1854) 69 = M. coriacea.
jequitinhonhensis, Berg, in Mart. Fl. Bras. xiv. i. 137. —Bras.
Klotzschiana, Berg, l. c. 111 = M. Klotzschiana.
laciniosa, Berg, l. c. 545.—Bras.
lancea, Berg, l. c. 88.—Bras.
lancifolia, Berg, in Linnaea, xxx. (1859–60) 658.—Guiana.
Langsdorffii, Berg, in Mart. Fl. Bras. xiv. i. 550 = M. Langsdorffii.
lanuginosa, Berg, l. c. 102 = M. lanuginosa.
laricina, Berg, l. c. 61.—Bras.
Laruottiana, Berg, in Linnaea, xxvii. (1854) 53 = M. Laruottiana.
Laureola, Berg, in Mart. Fl. Bras. xiv. i. 110.—Bras.
lauriflora, Berg, in Linnaea, xxvii. (1854) 64 = M. lauriflora.
laurifolia, Berg, l. c. = M. laurifolia.
laxiflora, Berg, in Mart. Fl. Bras. xiv. i. 114 = M. laxiflora.
leptoclada, Berg, in Linnaea, xxvii. (1854) 40 = M. leptoclada.
Leucadendron, Berg, l. c. 64 = M. Leucadendron.
leucantha, Berg, in Mart. Fl. Bras. xiv. i. 93.—Bras.
linearifolia, Berg, in Linnaea, xxvii. (1854) 35 = M. linearifolia.
lineata, Berg, in Mart. Fl. Bras. xiv. i. 68.—Bras.
Lingua, Berg, in Linnaea, xxvii. (1854) 77.—Bras.
linguaeformis, Berg, in Mart. Fl. Bras. xiv. i. 125.—Bras.
longipes, Berg, in Linnaea, xxvii. (1854) 55.—Bras.
lucida, Berg, in Mart. Fl. Bras. xiv. i. 118.—Bras.
macrocarpa, Berg, in Linnaea, xxvii. (1854) 41 = M. macrocarpa.
macrophylla, Berg, in Mart. Fl. Bras. xiv. i. 99 = M. macrophylla.
Mansoni, Berg, l. c. 121.—Bras.

AULOMYRCIA :—
Maraguana, Berg, in Mart. Fl. Bras. xiv. 1. 120 = Myrcia Maraguana.
maritima, Berg, l. c. 553.—Bras.
Martiana, Berg, l. c. 138 = M. Martiana.
Mathewsiana, Berg, in Linnaea, xxvii. (1854) 45.—Peruv.
micrantha, Berg, in Mart. Fl. Bras. xiv. 1. 516.—Bras.
microcarpa, Berg, l. c. 81 = M. microcarpa.
micropetala, Berg, in Linnaea, xxvii. (1854) 56.—Bras.
montana, Berg, l. c. 36 = M. montana.
mugiensis, Berg, l. c. = M. mugiensis.
multiflora, Berg, l. c. 47 = M. multiflora.
mutabilis, Berg, in Mart. Fl. Bras. xiv. 1. 70.—Bras.
myrtillifolia, Berg, in Linnaea, xxvii. (1854) 36 = M. myrtillifolia.
Ncesiana, Berg, l. c. 36 = M. Ncesiana.
nigrescens, Berg, l. c. 71 = M. nigrescens.
nigricans, Berg, in Mart. Fl. Bras. xiv. 1. 101.—Bras.
nigro-punctata, Berg, l. c. 116.—Bras.
nivea, Berg, in Linnaea, xxvii. (1854) 61 = M. nivea.
oblongata, Berg, l. c. 37 = M. oblongata.
obovata, Berg, l. c. 72.—Bras.
obscura, Berg, in Mart. Fl. Bras. xiv. 1. 132.—Bras.
obtecta, Berg, l. c. 117.—Bras.
obtusa, Berg, in Linnaea, xxvii. (1854) 66 = M. obtusa.
obtusata, Berg, l. c. 39.—Bras.
orthophylla, Berg, in Mart. Fl. Bras. xiv. 1. 549.—Bras.
Ottonis, Berg, in Linnaea, xxvii. (1854) 55.—Guiana.
ovalifolia, Berg, in Mart. Fl. Bras. xiv. 1. 81.—Bras.
ovalis, Berg, l. c. 107.—Bras.
pachyclada, Berg, l. c. 133.—Bras.
pallens, Berg, in Linnaea, xxvii. (1854) 73 = M. pallens.
pallida, Berg, in Mart. Fl. Bras. xiv. 1. 87.—Bras.
paniculariS, Berg, l. c. 84.—Bras.
paniculata, Berg, in Linnaea, xxvii. (1854) 50.—Bras.
paraënsis, Berg, in Mart. Fl. Bras. xiv. 1. 76.—Bras.
parnahibensis, Berg, l. c. 135.—Bras.
pauciflora, Berg, l. c. 560 = M. pauciflora.
perforata, Berg, l. c. 83.—Bras.
pertusa, Berg, in Linnaea, xxvii. (1854) 47 = M. pertusa.
piauhiensis, Berg, in Mart. Fl. Bras. xiv. 1. 87.—Bras.
pilantha, Berg, l. c. 555.—Bras.
pinifolia, Berg, in Linnaea, xxvii. (1854) 35 = M. pinifolia.
pirarensis, Berg, l. c. 41.—Guiana.
platyclada, Berg, in Mart. Fl. Bras. xiv. 1. 545 = M. platyclada.
plumbea, Berg, l. c. 142.—Bras.
Poeppigiana, Berg, in Linnaea, xxvii. (1854) 73.—Bras.
Pohliana, Berg, in Mart. Fl. Bras. xiv. 1. 72.—Bras.
polyantha, Berg, l. c. 82 = M. Kunthiana.
polymorpha, Berg, in Linnaea, xxvii. (1854) 46.—Bras.
pruinosa, Berg, in Mart. Fl. Bras. xiv. 1. 114.—Bras.
prunifolia, Berg, in Linnaea, xxvii. (1854) 55 = M. prunifolia.
puberula, Berg, l. c. 54 = M. puberula.
pubiflora, Berg, l. c. 40 = M. pubiflora.
pulchra, Berg, in Mart. Fl. Bras. xiv. 1. 68.—Bras.
punctata, Berg, l. c. 546 = M. punctata.
pusilla, Berg, l. c. 140.—Bras.
pyrifolia, Berg, in Linnaea, xxvii. (1854) 44.—Bras.
racemosa, Berg, l. c. 52.—Bras.
ramuliflora, Berg, in Mart. Fl. Bras. xiv. 1. 64.—Bras.
ramulosa, Berg, in Linnaea, xxvii. (1854) 36 = M. ramulosa.
Regeliana, Berg, in Mart. Fl. Bras. xiv. 1. 557.—Bras.
reticulata, Berg, l. c. 108 = M. reticulata.
reticulosa, Berg, in Linnaea, xxvii. (1854) 62 = M. reticulosa.
retusa, Berg, in Mart. Fl. Bras. xiv. 1. 142.—Bras.
Richardiana, Berg, in Linnaea, xxx. (1859–60) 664.—Bras.
Riedeliana, Berg, in Mart. Fl. Bras. xiv. 1. 551.—Bras.
roraimensis, Berg, in Linnaea, xxvii. (1854) 68.—Guiana.
rorida, Berg, in Mart. Fl. Bras. xiv. 1. 552.—Bras.
rostrata, Berg, l. c. 544 = M. rostrata.
rosulans, Berg, in Linnaea, xxvii. (1854) 54.—Bras.
rotundifolia, Berg, in Mart. Fl. Bras. xiv. 1. 123.—Bras.

AULOMYRCIA :—
rubella, Berg, in Linnaea, xxvii. (1854) 40 = M. rubella.
rufa, Berg, in Mart. Fl. Bras. xiv. 1. 65.—Bras.
rufipes, Berg, in Linnaea, xxvii. (1854) 77 = M. rufipes.
rugosa, Berg, in Mart. Fl. Bras. xiv. 1. 110.—Bras.
Sagraea, Berg, in Linnaea, xxx. (1859–60) 655.—Cuba.
salicifolia, Berg, l. c. xxvii. (1854) 78 = M. salicifolia.
Salzmanni, Berg, in Mart. Fl. Bras. xiv. 1. 116.—Bras.
Schomburgkiana, Berg, in Linnaea, xxvii. (1854) 75.—Bras.
Schrankiana, Berg, l. c. 79 = M. Schrankiana.
scrobiculata, Berg, in Mart. Fl. Bras. xiv. 1. 137 = M. scrobiculata.
Sonderiana, Berg, l. c. 110.—Bras.
spathulata, Berg, l. c. 91.—Bras.
sphaerocarpa, Berg, in Linnaea, xxvii. (1854) 51 = M. sphaerocarpa.
sphenoides, Berg, in Mart. Fl. Bras. xiv. 1. 63.—Bras.
Spruceana, Berg, l. c. 76.—Bras.
stenophylla, Berg, in Linnaea, xxx. (1859–60) 654.—Paraguay.
stictophylla, Berg, in Mart. Fl. Bras. xiv. 1. 67.—Bras.
stricta, Berg, l. c. 548.—Bras.
suaveolens, Berg, in Linnaea, xxvii. (1854) 78 = M. suaveolens.
subalpestris, Berg, l. c. 73 = M. subalpestris.
subavenia, Berg, in Mart. Fl. Bras. xiv. 1. 69.—Bras.
subcordata, Berg, in Linnaea, xxvii. (1854) 62 = M. subcordata.
subobliqua, Berg, l. c. 57.—Guiana.
subverticillaris, Berg, l. c. 73.—Bras.
suffruticosa, Berg, in Mart. Fl. Bras. xiv. 1. 136.—Bras.
supraaxillaris, Berg, l. c. 69.—Bras.
surinamensis, Berg, in Linnaea, xxvii. (1854) 64 = M. surinamensis.
tenuifolia, Berg, in Mart. Fl. Bras. xiv. 1. 68.—Bras.
ternifolia, Berg, l. c. 134.—Bras.
thyrsiflora, Berg, in Linnaea, xxvii. (1854) 74.—Bras.
torta, Berg, l. c. 78 = M. torta.
tortuosa, Berg, in Mart. Fl. Bras. xiv. 1. 558.—Bras.
trichantha, Wawra, in Oestr. Bot. Zeitschr. xxix. (1879) 215.—Bras.
triflora, Berg, in Linnaea, xxvii. (1854) 79 = M. triflora.
trifolia, Berg, in Mart. Fl. Bras. xiv. 1. 107.—Bras.
undulata, Berg, l. c. 89.—Bras.
vacciniifolia, Berg, l. c. 140.—Bras.
variabilis, Berg, in Linnaea, xxvii. (1854) 62 = M. variabilis.
Vautheriana, Berg, l. c. xxx. (1859–60) 655 = M. Vauthieriana.
velhensis, Berg, in Mart. Fl. Bras. xiv. 1. 560.—Bras.
venulosa, Berg, in Linnaea, xxvii. (1854) 63 = M. venulosa.
vestita, Berg, l. c. 74 = M. vestita.
virgata, Berg, in Mart. Fl. Bras. xiv. 1. 136 = M. virgata.
Widgreniana, Berg, l. c. 70.—Bras.
Wullschlaegeliana, Berg, in Linnaea, xxvii. (1854) 48.—Guiana.

AULONEMIA, Goudot, in Ann. Sc. Nat. Sér. III. v. (1846) 76. t. 5 = **Arthrostylidium**, Rupr. (Gramin.).
Gueko, Goudot, l. c.—N. Granat.

AULONIX, Rafin. Sylva Tellur. 25 (1838) = **Cytisus**, Linn. (Legumin.).
biflorus, Rafin. l. c. = C. biflorus.

AULOSEMA, Walp. Rep. i. 694 (1842) = **Astragalus**, Linn. (Legumin.).
eximium, Walp. l. c. = Astr. hypogaeus.

AURANTIUM, Tourn. ex Mill. Gard. Dict. ed. VI (1752) = **Citrus**, Linn. (Rutac.).
acre, Mill. l. c. ed. VIII. n. 1 = C. Aurantium.
decumana, Mill. l. c. n. 4 = C. decumana.
humile, Mill. l. c. n. 5 = C. humilis.
myrtifolium, Descourt. Fl. Med. Antill. iii. 308. t. 219 = C. Aurantium.
orientale, Mill. Gard. Dict. ed. VIII. n. 3 = C. Aurantium.
silvestre, Pritz. Ic. Ind. 127 = C. Aurantium.
sinense, Mill. Gard. Dict. ed. VIII. n. 2 = C. Aurantium.

AURELIA, Cass. Bull. Soc. Philom. (1815) 175; et in Dict. Sc. Nat. xxxvii. 468 (1816) = **Grindelia**, Willd. (Compos.).
amplexicaulis, Cass. l. c. = G. squarrosa.
decurrens, Cass. l. c. = G. glutinosa.

AURELIA :—
glutinosa, Cass.—Cf. Bull. Soc. Philom. (1815) 175 = Grindelia glutinosa.
squarrosa, Cass. ex Steud. Nom. ed. II. i. 171 = G. squarrosa.

AURELIA, J. Gay, in Ann. Sc. Nat. Sér. IV. x. (1858) 95 = **Narcissus**, Tourn. (Amaryll.).
Broussonetii, J. Gay, l. c. 96 = N. Bronssonetii.

AURELIANA, Lafit. ex Catesb. Nat. Hist. Carol. App. 16 (1754) = **Aralia**, Linn.

AURELIANA, Sendt. in Mart. Fl. Bras. x. 138. t. 19. (1846) = **Bassovia**, Aubl. (Solan.).
fasciculata, Sendt. l. c. 140 = B. fasciculata.
glomuliflora, Sendt. l. c. 138.—Bras.
lucida, Sendt. l. c. 139.—Bras.
tomentosa, Sendt. l. c. 140 = B. tomentosa.
velutina, Sendt. l. c. = B. velutina.

AUREOLARIA, Rafin. New Fl. Am. ii. 58 (1836) = **Gerardia**, Linn. (Scrophul.).
cuneifolia, Rafin. l. c. 59 = Herpestis pubescens?
glauca, Rafin. l. c. 60 = G. quercifolia.
levigata, Rafin. l. c. 59 = G. laevigata.
reticulata, Rafin. l. c.—Am. bor.
rupestris, Rafin. l. c. 60 = G. rupestris.
villosa, Rafin. l. c. 59 = G. flava.

AURICULA, Tourn. ex Spach, Hist. Vég. Phan. ix. 356 (1840) = **Primula**, Tourn.
crenata, Spach, l. c. 358 = P. marginata.
glutinosa, Spach, l. c. 359 = P. glutinosa.
hortensis, Spach, l. c. 357 = P. Auricula.
integrifolia, Spach, l. c. 359 = P. integrifolia.
minima, Spach, l. c. 360 = P. minima.
Palinuri, Spach, l. c. 357 = P. Palinuri.
villosa, Spach, l. c. 358 = P. minima.

AURILA, Noronha, in Verh. Batav. Gen. v. (1790) ed. l. Art. IV. 1, nomen (Quid?).

AURINIA, Desv. Jonrn. Bot. iii. (1813) 162 = **Alyssum**, Linn. (Crucif.).
capensis, Desv. l. c. = Heliophila diffnsa.
corymbosa, Griseb. Spicil. Fl. Rumel. i. 271 = Al. corymbosum.
edentula, Fisch. Mey. & Trantv. Ind. Sem. Hort. Petrop. iii. 4 = Al. gemonense.
gemonensis, Griseb. Spicil. Fl. Rnmel. i. 272 = Al. gemonense.
media, Schur, Enum. Pl. Transs. 61 = Al. gemonense.
obcordata, Desv. Journ. Bot. iii. (1813) 162 = Al. obcordatum.
orientalis, Griseb. Spicil. Fl. Rnmel. i. 272 = Al. orientale.
petraeum, Fuss, Fl. Transs. 64 = Al. gemonense.
saxatilis, Desv. Journ. Bot. iii. (1813) 162 = Al. saxatile.
saxatilis, Griseb. Spicil. Fl. Rumel. i. 273 = Al. orientale.
sinuata Griseb. l. c. 271 = Al. sinuatum.

AUROTA, Rafin. Fl. Tellur. iii. 61 (1836) = **Curculigo**, Gaertn. (Amaryll.).
latifolia, Rafin. l. c. = C. latifolia.

AURORA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 1 = **Quisqualis**, Linn. (Combret.).

AUSTERIUM, Poit. ex DC. Prod. ii. 385 (1825) = **Rhynchosia**, Lour. (Legumin.).

AUSTRALINA, Gaudich. in Freyc. Voy. Bot. 505 (1826). *URTICACEAE*, Benth. & Hook. f. iii. 394.
ANAGANTHOS, Hook. f. Fl. N. Zel. i. 226 (1854).
DIDYMOTOCA, E. Mey. Zwei Pfl. Docum. 178 (1848). nomen.
acuminata, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 212.—Afr. austr.
capensis, *Wedd. l. c.*—Afr. austr.
flaccida, Wedd. in Arch. Mus. Par. viii. (1855–56) 546. t. 20 C. = Schimperiana.
integrifolia, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 212.—Afr. austr.
Muelleri, *Wedd. in Arch. Mus. Par.* viii. (1855–56) 545.—Austral.
Novae-Zelandiae, Hook. f. Fl. N. Zel. i. 226 = pusilla.

AUSTRALINA:—
pusilla, *Gaudich. in Freyc. Voy. Bot.* 505.—Austral.; N. Zel.
Schimperiana, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 212.—Abyss.
tasmanica, Hook. f. Fl. N. Zel. i. 226 = pusilla.

AUSTROBUXUS, Miq. Fl. Ind. Bat. Suppl. 444 (1860) = **Sarcococca**, Lindl. (Euphorb.).
nitidus, Miq. l. c.—Sumatra.

AUTOGENES, Rafin. Fl. Tellur. iv. 20 (1836) = **Narcissus**, Tourn. (Amaryll.).
angustifolius, Rafin. l. c. = N. poëticus.
biflorus, Rafin. l. c. = N. biflorus.
poëticus, Rafin. l. c. = N. poëticus.
tenuior, Rafin. l. c. = N. gracilis.

AUXEMMA, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 23. t. 5. *BORAGINEAE*, Benth. & Hook. f. ii. 840.
Gardneriana, *Miers, l. c.* 24. t. 5.—Bras.

AVELLANITA, Phil. in Linnaea, xxxiii. (1864–65) 237. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 289.
Bustillosi, *Phil. l. c.* 238.—Chili.

AVELLINIA, Parl. Pl. Nov. 59 (1842). *GRAMINEAE*, Benth. & Hook. f. iii. 1184.
Michelii, *Parl. l. c.*—Italia.
Savi, *Parl. l. c.* 61.—Italia.
scabriuscula, Nym. Consp. 815 = Koeleria scabriuscula.
tenuicula, *Nym. l. c.*—Hispan.

AVENA, Hall. ex Scop. Introd. 74 (1777) = **Agrostis**, Linn. (Gramin.).

AVENA, Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1160.
ELICTOTRICHON, Bess. ex Andrz. Rys Botan. i. (1823) 9.
HELICOTRICHUM, Bess. in Reichb. Fl. Germ. Excurs. 140⁶. n. 352 (1832).
HEUFFELIA, Schur, Enum. Pl. Transs. 760 (1866).
abyssinica, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 415.—Abyss.
adsurgens, *Schur, Enum. Pl. Transs.* 762.—Hungar.
affinis, Bernh. ex Steud. Nom. ed. II. i. 171, 173 = sterilis.
agraria, Brot. Fl. Lusit. i. 105 = strigosa.
agraria-mutica, Brot. l. c. 106 = strigosa.
agraria-sesquialtera, Brot. l. c. = strigosa.
agropyroides, *Boiss. Diagn.* Ser. I. xiii. 50.—Graecia.
agrostidea, Fries, Mant. iii. 3 = Trisetum agrostideum.
agrostoides, Griseb. Spicil. Fl. Rumel. ii. 454 = Trisetum myrianthum.
airoides, Koel. Gram. 208 = Trisetum snbspicatum.
airoides, Pourr. ex Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 290 = Ponrretii.
alba, *Vahl, Symb. Bot.* ii. 24.—Gall. austr.; Afr. bor.
albinervis, *Boiss. Voy. Bot. Espagne*, ii. 656.—Hispan.
Alopecuros, Roth, Catalecta, iii. 20 = Trisetum Alopecuros.
alpestris, DC. Fl. Fr. vi. 260 = Trisetum pratense.
alpestris, Host, Gram. Austr. iii. 27. t. 39 = T. alpestre.
alpina, [Honck.] Verz. Gew. Teutschl. 302 = Aira montana?
alpina, Latour. Chlor. Lugd. 3 = Schenzeri.
alpina, Sm. in Trans. Linn. Soc. x. (1811) 335 = pratensis.
alpina, Trin. Fund. Agrost. 157 = Deschampsia alpina.
alta, Cav. ex Roem. & Schult. Syst. ii. 691 = strigosa.
ambigua, Schoenb. ex Nym. Consp. 810 = fatua.
amethystina, *DC. Fl. Fr.* iii. 56; vi. 260.—Gallia.
andropogoides, *Steud. in Flora*, xii. (1829) II. 486.—Afr. austr.
anglica, Hort. ex Roem. & Schult. Syst. ii. 669 = sativa.
antarctica, Roem. & Schult. l. c. 676 = Trisetum antarcticnm.
antarctica, Thunb. Prod. Pl. Cap. 22 = Danthonia antarctica.

AVENA:—
arduensis, Lej. ex Steud. Nom. ed. II. i. 171 = strigosa.
arenaria, Spreng. Syst. i. 333 = Trisetum arenarium.
argaea, *Boiss. Fl. Orient.* v. 546.—Cappadoc.
argentoideum, Schur, in Oestr. Bot. Zeitschr. x. (1860) 74 = Trisetum alpestre.
argentea, Willd. Enum. Hort. Berol. 125 = Trisetum argenteum.
ariguensis, *Steud. Syn. Pl. Gram.* 233.—Chili.
aristidoides, Steud. in Flora, xii. (1829) II. 481. n. 23 = Danthonia Thunbergii.
aristidoides, *Thunb. Prod. Pl. Cap.* 22.—Afr. austr.
arundinacea, Delile, Fl. Egypte, 171. t. 12 = Danthonia Forskohlei.
arvensis, Salisb. Prod. 23 = Bromus arvensis.
aspera, *Munro, ex Thw. Enum. Pl. Zeyl.* 372.—Zeylan.
atheranthera, Presl, Cyp. et Gram. Sic. 30 = barbata.
atropurpurea, Link, Hort. Berol. i. 119 = Aira atropurpurea.
aurata, All. Fl. Pedem. ii. 255 = setacea, Agrostis alpina.
australis, Parl. Fl. Ital. i. 285 = pratensis.
barbata, *Brot. Fl. Lusit.* i. 108.—Reg. Mediterr.; As. Min.; Arab.
barbata, *Pott, ex Link, in Schrad. Journ.* ii. (1799) 315.—Lusitan.
baregensis, Nym. Consp. 813 = Trisetum agrostideum.
Banmgartenii, *Steud. Syn. Pl. Gram.* 233.—Hungar.
Bellardi, *Colla, Herb. Pedem.* t. 93. f. 2.—Italia.
benghalensis, *Schwaegr. ex Steud. Nom.* ed. I. 94.—Ind. or.
Besseri, *Ledeb. Fl. Ross.* iv. 415.—Ross. med.
bifida, *Beauv. Agrost.* 153, nomen.—Hab.?
bipartita, Link, Hort. Berol. i. 113 = Danthonia semiannularis.
Blavii, Aschers. & Janka.—Cf. Aschers. & Kan. Cat. Coron. Serb. 9 = compressa.
borbonia, *Steud. Syn. Pl. Gram.* 234.—Ins. Borbon.
brevifolia, Host, Gram. iii. t. 40 = Trisetum distichophyllum.
brevifolia, Spreng. Syst. i. 334 = Trisetnm rigidum.
brevis, *Roth, Tent. Fl. Germ.* i. 40.—Europ.
bromoides, *Gouan, Hort. Monsp.* 52.—Reg. Mediterr.
bromoides, Mert. & Koch, in Roehl. Fl. Deutschl. ed. III. i. 570 = pratensis.
Brownei, Spreng. Syst. i. 336 = Danthonia pallida.
Bruhnsiana, *Gruner, in Bull. Soc. Nat. Mosc.* xli. (1867) II. 458.—Reg. Casp.
bulbosa, Willd. in Ges. Naturf. Fr. Berl. Neue Schr. ii. (1799) 116. t. 5. f. 1 = Arrhenatherum avenaceum.
Burnoufii, Nym. Syll. 413 = Trisetum Burnoufii.
byzantina, C. Koch, in Linnaea, xxi. (1848) 392 = fatua.
caespitosa, Kuntze, Taschen-Fl. Leipz. 45 = Deschampsia caespitosa.
calicina, Vill. Hist. Pl. Dauph. ii. 148 = Danthonia provincialis.
callosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxix. (1856) I. 36.—Sibir.
calycina, Lam. Illustr. i. 200 = Danthonia calycina.
calycina, Willd. Sp. Pl. i. 453, in syn. = Danthonia provincialis.
canariensis, *Nees, ex Steud. Nom.* ed. II. i. 171.—Ins. Teneriff.
Candollei, Nym. Consp. 812 = Trisetum pratense.
canescens, [Weber, in] Wigg. Prim. Fl. Hols. 9 = Corynephorus canescens.
cantabrica, Lag. in Herrer. Agr. 141 = filifolia.
capensis, Burm. f. Fl. Cap. Prod. 3 = Anthistiria ciliata.
capensis, Linn. f. Suppl. 112 = Aristida capensis.
capensis, Steud. in Flora, xii. (1829) 481 = Danthonia papillosa.
capillacea, *Beauv. Agrost.* 153, nomen.—Hab.?
capillaris, *Mert. & Koch, in Roehl. Deutschl. Fl.* ed. III. i. 573.—Europ.
Carmeli, *Boiss. Diagn.* Ser. I. xiii. 50.—Syria.
caroliniana, Walt. Fl. Carol. 81 = Trisetum palnstre.
carpatica, Host, Gram. Austr. iv. 18. t. 31 = Trisetum carpathicum.
caryophyllea, *Sibth. & Sm. Fl. Graec. Prod.* i. 67.—Archip. Graec.
caryophyllea, [Weber, in] Wigg. Prim. Fl. Hols. 10 = Aira caryophyllea.
Cavanillesii, Hort. ex Roem. & Schult. Syst. ii. 691 = strigosa.

AVENA:—
Cavanillesii, Koch, Syn. Fl. Germ. ed. I. 797 = Trisetnm Cavanillesii.
Cavanillesii, Lag. in Varied. Cienc. ii. IV. (1805) 39 = Stipa arenaria.
cernua, Kunth, Enum. Pl. i. 306 = Trisetum cernunm.
chinensis, Fisch. ex Roem. & Schult. Syst. ii. 669 = nuda.
chlorantha, Link, in Linnaea, xvii. (1843) 401 = Trisetum aureum.
ciliaris, Kit. ex Schult. Oestr. Fl. ed. II. i. 268 = Trisetum carpathicum.
cinerea, Auct. ex Roem. & Schult. Syst. ii. 669 = sativa.
clauda, *Dur. in Duch. Rev. Bot.* i. (1845–46) 360.—Graecia; As. Min.; Afr. bor.
coarctata, Desf. Cat. Hort. Par. ed. III. (1829) 22 = Gaudinia geminiflora.
colorata, Steud. in Flora, xii. (1829) II. 481. n. 11 = Danthonia tortuosa.
compacta, *Boiss. & Heldr. Diagn.* Ser. I. vii. 122.—Graecia; Servia.
compressa, *Heuff. in Flora*, xviii. (1835) 244.—Hungar.
condensata, Link, Enum. Hort. Berol. i. 82 = Trisetum aureum, *Tenore*.
convoluta, *Presl, Cyp. et Gram. Sic.* 31.—Europ. austr. or.
coquimbensis, Roem. & Schult. Syst. ii. 677 = Hierochloë utriculata.
coronensis, *Schur, Enum. Pl. Transs.* 763.—Transsylv.
corymbosa, *Nym. Consp.* 814.—Corsica.
cristata, Auct. ex Roem. & Schult. Syst. ii. 758 = Agropyron cristatum.
Cupaniana, *Nym. Syll.* 414.—Ital.; Creta.
cuspidata, Willd. ex Kunth, Enum. Pl. i. 239 = Calamagrostis cuspidata.
daenensis, Boiss. Diagn. Ser. I. vii. 123 = Trisetum rigidum.
decora, *Janka, in Termész. Füzetek*, viii. (1884) 28.—Europ.
depauperata, Sieber, ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 68 = Danthonia provincialis.
desertorum, *Less. in Linnaea*, ix. (1834) 208.—Sibir.
deyeuxioides, H. B. & K. Nov. Gen. et Sp. i. 147 = Deyeuxia triflora.
dispermis, Mill. Gard. Dict. ed. VIII. n. 1 = sativa.
distans, *Schur, in Oestr. Bot. Zeitschr.* xx. (1870) 22.—Europ.
disticha, Lam. Encyc. i. 330 = Trisetum rigidum.
distichophylla, Host, Gram. Austr. ii. 39. t. 53 = Trisetum argenteum.
distichophylla, Vill. Prosp. 16; Pl. Dauph. ii. 144 = Trisetum distichophyllum, rigidum.
dubia, Leers, Fl. Herborn. 41 = Gaudinia tenuis.
Dufourei, Nym. Consp. 813 = Trisetum Dufourei.
dura, Salisb. Prod. 22 = Festuca duriuscula.
elata, *Forsk. Fl. Aegypt. Arab.* 19.—Constantinop.
elata, Salisb. Prod. 23 = Arrhenatherum avenaceum.
elatior, Linn. Sp. Pl. 79 = Arrhenatherum elatius.
elegans, Aschers. in Verh. Bot. Ver. Brand. iii.–iv. (1861–62) 279 = capillaris.
elephantina, Spreng. ex Steud. Nom. ed. II. i. 172 = andropogoides.
elephantina, Thunb. Prod. Pl. Cap. 23 = Danthonia elephantina.
elongata, H. B. & K. Nov. Gen. et Sp. i. 148 = Trisetum elongatum.
eriantha, *Dur. in Duch. Rev. Bot.* i. (1845–46) 360.—Afr. bor.
eriantha, *Hack. in Oestr. Bot. Zeitschr.* xxvii. (1877) 122.—Hispan.
fallax, De Not. Ind. Sem. Hort. Reg. Genuens. 24 = sempervirens.
fallax, Pollini, Fl. Vernon. iii. 773 = pubescens.
fallax, Roem. & Schult. Syst. ii. 692 = montana.
fallax, Tenore, Fl. Nap. iii. 96 = convoluta.
fatua, *Linn. Sp. Pl.* 80.—Europ.; Oriens; As. bor.
fatua, Schreb. Beschr. Graes. 109. t. 15 = sterilis.
fatua, Viv. Fl. Lib. Spec. 6 = barbata.
fertilis, All. Auct. Pedem. 45 = Gaudinia tenuis.
festucaeformis, *Hochst. in Flora*, xxxviii. (1855) 275.—Afr. or.
filifolia, *Lag. Gen. & Sp. Nov.* 4.—Hispan.
filiformis, Forst. f. Prod. 9 = Deyeuxia Forsteri.
filiformis, Labill. Nov. Holl. Pl. i. 24. t. 31 = Deyenxia Billardieri.
flava, Hort. ex Roem. & Schult. Syst. ii. 669 = sativa.

AVENA:—
flavescens, Bout. ex Willk. & Lange, Prod. Fl. Hisp. i. 72 = Trisetum gallecicum.
flavescens, Linn. Sp. Pl. 80 = Trisetum pratense.
flexuosa, Mert. & Koch, in Roehl. Deutschl. Fl. ed. III. i. 570 = Deschampsia flexuosa.
flexuosa, *Schrank, in Denkschr. Reg. Bot. Ges.* ii. 7.—Labrador.
Forskolei, Vahl, Symb. Bot. ii. 25 = Danthonia Forskolei.
Forsteri, Kunth, Rev. Gram. i. 104 = Deyeuxia Forsteri.
fragilis, Linn. Sp. Pl. 80 = Gaudinia fragilis.
Freita, Orteg. [ex Spreng.] Bot. Gart. Halle, 14 = strigosa.
fusca, Ard. in Saggi Acc. Padov. ii. (1789) t. 4 = sativa.
fusca, Kit. ex Schult. Fl. Oestr. ed. II. i. 268 = Trisetum carpathicum.
fusca, *Schur, Enum. Pl. Transs.* 756.—Hungar.
fuscoflora, Schur, l. c. = fusca.
gallecica, Nym. Syll. Suppl. 71 = Trisetum gallecicum.
geminiflora, Kunth, Rev. Gram. 103, 299. t. 61 = Gaudinia geminiflora.
Georgiana, Roem. & Schult. Syst. ii. 669 = sativa.
georgica, Boiss. olim ex Fl. Orient. v. 540 = Ventenata macra.
georgica, Zuccag. in Roem. Collect. 126 = sativa.
gigantea, Salisb. Prod. 23 = Bromus giganteus.
glabra, C. Koch, in Linnaea, xix. (1847) 5 = pubescens.
glabrata, Brot. ex Steud. Nom. ed II. i. 172 = corymbosa.
glacialis, Boiss. Elench. 87; Voy. Bot. Espagne, ii. 654 = Trisetum glaciale.
glauca, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 20 = Scheuzeri.
glomerata, Steud. in Flora, xii. (1829) II. 483 = Danthonia curvifolia.
glumosa, *Ell. Sketch*, 11.—Am. bor.
glumosa, Michx. Fl. Bor. Am. i. 72 = Danthonia spicata.
gracilis, Moris, Stirp. Sard. Elench. fasc. i. 50 = Trisetum Morisii.
Heldreichii, Parl. Fl. Ital. i. 275 = convoluta.
heteromalla, Moench, Meth. 195 = orientalis.
hexantha, Steud. in Flora, xii. (1829) II. 487. n. 87 = Danthonia stricta.
hirsuta, Moench, Meth. Suppl. 64; Roth, Catalecta, iii. 19 = barbata.
hirta, *Schrad. in Goett. Gel. Anz.* (1821) III. 2075.—Afr. austr.
hirtifolia, Boiss. Diagn. Ser. II. iv. 128 = pubescens.
hirtula, Lag. Gen. et Sp. Nov. 4 = barbata.
hispanica, Ard. in Saggi Acc. Padov. ii. (1789) 112 = sativa.
hispanica, Hort. ex Roem. & Schult. Syst. ii. 691 = strigosa.
hispanica, Lange, in Kjoeb. Vidensk. Meddel. (1860) 41 = Arrhenatherum erianthum.
hispida, Linn. f. Suppl. 111 = Tristachya leucothrix.
hispida, Nym. Syll. Suppl. 71 = Trisetum hispidum.
hybrida, Peterm. ex Reichb. Fl. Saxon. 17 = fatua.
Hoppeana, *Scheele, in Flora*, xxvii. (1844) 57.—Istria.
Hostii, Boiss. & Reut. Pugill. 121 = Parlatorii.
Hugeninii, *De Not. ex Steud. Syn. Pl. Gram.* 425.—Italia.
hungarica, *Luce, Topogr. Nachr. Oesel*, 20.—Hungar.
hydrophila, F. Muell. ex Hook. f. Fl. Tasm. i. 121 = Amphibromus Neesii.
intermedia, Lindgren, in Bot. Notiser (1841), 151 = fatua.
intermedia, Guss. Fl. Sic. Prod. Suppl. 16 = corymbosa.
involucrata, Schrad. in Goett. Gel. Anz. (1821) III. 2075 = Danthonia involucrata.
involuta, Presl, ex Steud. Nom. ed. II. i. 172 = sempervirens.
japonica, Steud. Syn. Pl. Gram. 281 = hirsuta.
Junghuhnii, *Buese, in Miq. Pl. Jungh.* 345.—Java.
Koenigii, Spreng. Syst. i. 336 = Aristida arundinacea.
Kotschyi, *Steud. Syn. Pl. Gram.* 235.—Persia.
lachnantha, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 227.—Afr. trop.
laconica, Nym. Syll. Suppl. 71 = Trisetum laconicum.
lacta, Salisb. Prod. 22 = Brachypodium pinnatum.
laevigata, Schur, in Oestr. Bot. Zeitschr. x. (1860) 72 = pubescens.
laevis, Hack. l. c. xxvii. (1877) 46 = albinervis

AVENA:—
lanata, Koel. Gram. 303 = Holcus lanatus.
lanata, Schrad. in Goett. Gel. Anz. (1821) III. 2075 = Danthonia lanata.
lanuginosa, Gilib. Exercit. ii. 539 = fatua.
lasiantha, Link, in Linnaea, ix. (1834) 135 = Bromus intermedius.
latifolia, Host, Gram. Austr. iv. 19. t. 32 = planiculmis.
lendigera, Salisb. Prod. 23 = Gastridium australe.
leonina, Steud. in Flora, xii. (1829) II. 484 = Danthonia antarctica ?
leptostachys, *Hook. f. Fl. Antarct.* 378.—Reg. Magellan.
levis, Nym. Consp. 811 = albinervis.
lodunensis, Delastr. in 'Cat. Hort. Pict. (1835)' = sulcata.
Loeflingiana, Cav. Ic. i. t. 45. f. 1 = Trisetum Gaudinianum.
Loefflingiana, Geuns. ex Roem. & Schult. Syst. ii. 370 = Aira praecox.
Loeflingiana, Balb. Misc. Bot. i. 12 = Trisetum Cavanillesii.
Loeflingiana, Linn. Sp. Pl. 79 = Trisetum hispanicum.
Loeflingiana, Lam. Encyc. i. 332 = Trisetum paniccum.
longifolia, Req. in DC. & Duby, Bot. Gall. i. 514 = pratensis.
longifolia, Spreng. Syst. i. 336 = Danthonia longifolia.
longifolia, Thore, Prom. Golf. fasc. 92 = Arrhenatherum pallens.
longiglumis, *Dur. in Duch. Rev. Bot.* i. (1845-46) 359.—Afr. bor.
lucida, Bertol. Fl. Ital. i. 701 = pubescens.
ludoviciana, Dur. in Bull. Soc. Linn. Bord. xx. (1855) i. 41 = sterilis.
lupulina, Steud. in Flora, xii. (1829) II. 486 = Danthonia lanata.
lupulina, Thunb. Prod. Pl. Cap. 23 = Danthonia lupulina.
lutea, *Linn. f. Suppl.* 112.—Ins. Martinic.
macilenta, Guss. Fl. Sic. Prod. Suppl. i. 29 = Koeleria macilenta.
macra, Hohen. Enum. Pl. Elisabethpol. 214 = rigida.
macra, Stev. in Bieb. Fl. Taur. Cauc. i. 77 = Ventenata macra.
macrocalycina, Steud. in Flora, xii. (1829) II. 482 = Danthonia macrantha.
macrocarpa, Moench, Meth. 196 = sterilis.
macrostachya, *Balansa & Dur. in Bull. Soc. Bot. Fr.* i. (1854) 318.—Algeria.
magellanica, *Beauv. Agrost.* 153, nomen.—Reg. Magellan.
Mandoniana, Coss. & Bal. in Bull. Soc. Bot. Fr. xv. (1868) 185, nomen = marginata ?
marginata, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) 529.—Ins. Madera.
maxima, *Presl, Fl. Sicul.* p. xliv.—Sicilia.
mediolanensis, Balansa & De Not. in Comoll. Fl. Comens. i. 147 = Trisetum myrianthum.
Michelii, Guss. Fl. Sic. Syn. 151 = Koeleria macilenta.
mollis, Koel. Gram. 300 = Holcus mollis.
mollis, Michx. Fl. Am. Bor. i. 72 = Trisetum subspicatum.
mollis, Salisb. Prod. 23 = Bromus mollis.
montana, Brot. Fl. Lusit. i. 109 = Arrhenatherum Thorei.
montana, Roem. & Schult. Syst. ii. 674, in syn. = pratensis.
montana, *Vill. Fl. Dauph.* ii. 151.—Mont. Pyren.
montana, [Weber, in] Wigg. Prim. Fl. Holsat. 9 = Deschampsia flexuosa.
multiculmis, Dum. Agrost. 121 = Aira caryophyllea.
muralis, Salisb. Prod. 22 = Festuca Myurus.
muricata, Spreng. Neue Entdeck. i. 247 = Aristida capensis, Danthonia papillosa.
mutica, *Krock. Fl. Sil. Suppl.* i. 187.—Enrop.
myriantha, Bertol. Fl. Ital. i. 722 = Trisetum myrianthum.
mysorensis, Spreng. Syst. i. 337.—Ind. or.
nana, Kunze, ex Steud. Nom. ed. II. i. 172, 45 = Aira praecox.
Neesii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 229.—Afr. trop.
neglecta, Bertol. Fl. Ital. i. 713 = Trisetum Gaudinianum.
neglecta, Savi, Fl. Pis. t. 1. f. 4 = Trisetum paniceum.
nemoralis, Salisb. Prod. 23 = Bromus asper.
nervosa, Lam. Illustr. n. 1115 = strigosa.
nervosa, R. Br. Prod. i. 178 = Amphibromus Neesii.

AVENA:—
Neumeyeriana, Vis. Fl. Dalm. iii. 339 = compacta.
nigra, *Wallr. Beitr. Fl. Hercyn.* 174; *et in Linnaea*, xiv. (1840) 544.—Europ.
nitida, Desf. Fl. Atlant. i. 102 = Trisetum nitidum.
nodosa, Cullum, It. Scan. 251 = Arrhenatherum avenaceum.
Noeana, Nym. Syll. Suppl. 71 = Trisetum aurenm, *Tenore*.
Notarisii, Parl. Fl. Ital. i. 279; ex Steud. Syn. Pl. Gram. 425 = sempervirens.
nuda, *Linn. Amoen. Acad.* iii. 401.—Europ. austr.
nuda, Thuill. Fl. Par. ed. II. i. 49 = brevis.
nutans, St. Lag. Rech. Hist.—Cf. Bull. Soc. Bot. Fr. xxxi. (1884) Bibl. 201 = sterilis.
nutkaensis, *J. Presl, in Rel. Haenk.* i. 254.—Sinus Nootka.
occidentalis, *Dur. Cat. Gr. Bord.* (1864) 25.—*Cf. Bull. Soc. Bot. Fr.* xii. (1865) *Bibl.* 78.—Ins. Ferro.
odorata, DC. Fl. Fr. iii. 42 = Hierochloë australis.
odorata, Pers. Syst. i. 100 = Hierochloë borealis.
oligostachya, *Munro, ex Aitch. in Journ. Linn. Soc.* xviii. (1881) 108; xix. 193. t. 30.—Afghanist.
orientalis, *Schreb. Spicil.* 52.—Europ. austr.; Oriens.
ovata, C. C. Gmel. ex Kunth, Enum. Pl. i. 295 = Trisetum ovatum.
ovina, Salisb. Prod. 22 = Festuca ovina.
palaestina, *Steud. Syn. Pl. Gram.* 425.—Palaestina.
pallens, Link, in Schrad. Journ. ii. (1799) 314 = Arrhenatherum pallens.
pallida, Salisb. Prod. 24 = Holcus lanatus.
pallida, Thunb. Prod. Pl. Cap. 22 = Danthonia pallescens.
palustris, Michx. Fl. Bor. Am. i. 72 = Trisetum palustre.
panicea, Hort. ex Roem. & Schult. Syst. ii. 625, 660 = Koeleria phleoides.
panicea, Lam. Illustr. i. 202 = Trisetum paniceum.
papillosa, Schrad. in Goett. Gel. Anz. (1821) III. 2075 = Danthonia papillosa.
papillosa, Steud. in Flora, xii. (1829) II. 434 = Danthonia scabra.
paradensis, *Kit. ex Kanitz, in Verh. Zool. Bot. Ges. Wien*, xiii. (1863) 543 (*Reliq. Kitaib.* 119).—Europ.
paradoxa, Willd. ex Kunth, Enum. Pl. i. 188 = Aristida dichotoma.
Parlatorii, *Woods, ex Dalla Torre, Atl. Alpenfl.* 228.—Italia.
parviflora, Desf. Fl. Atlant. i. 98 = Trisetum parviflorum.
parviflora, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 289 = Trisetum micranthum.
pauciflora, Auct. ex Roem. & Schult. Syst. ii. 660 = Trisetum paniceum.
pauperula, *Phil. in Linnaea*, xxix. (1857-58) 94.—Chili.
pendula, Gilib. Exercit. 539 = sativa.
penicillata, Willd. ex Stend. Nom. ed. II. i. 172; ii. 714 = Trisetum rigidum.
pennsylvanica, Muhl. Descr. Uber. 185 = Arrhenatherum kentuckense.
pensylvanica, Forsk. Fl. Aegypt. Arab. 23 = Danthonia Forskolei.
pensylvanica, *Linn. Sp. Pl.* 79.—Am. bor.
pensylvanica, Presl, Cyp. et Gram. Sic. 30 = maxima.
persica, *Steud. Syn. Pl. Gram.* 230.—Persia.
peruviana, Hort. ex Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 290 = Trisetum paniceum.
phleoides, Urv. in Mém. Soc. Linn. Par. iv. (1826) 601 = Trisetum phleoides.
pilosa, Aucher, ex Dur. in Duch. Rev. Bot. i. (1845-46) 860 = clauda.
pilosa, *Bieb. Fl. Taur. Cauc.* iii. 84.—As. Min.; Syria; Afr. bor.
pilosa, *J. Presl, in Rel. Haenk.* i. 253.—Chili.
pilosa, *Scop. Fl. Carn.* ed. II. i. 86.—Carniolia.
planiculmis, *Schrad. Fl. Germ.* i. 881. t. 6. f. 2.—Europ.; As. bor.
planiculmis, Sm. Engl. Bot. t. 2141 = pratensis.
podolica, Pascal. ex Zuccagni, in Roem. Collect. 126 = sativa.
Pourretii, *Willd. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 290. t. 8.—Hispan.
praecox, *Beauv. Agrost.* 89.—Lusitan.
praecusta, Reichb. Fl. Germ. Excurs. 140 = pratensis.
pratensis, Boiss. Voy. Bot. Espagne, ii. 656 = bromoides.
pratensis, Brot. Fl. Lusit. ii. 110 = sulcata.

AVENA:—
pratensis, Gouan, 'Hort. 53'; Kunth, Enum. Pl. i. 300 = pubescens.
pratensis, *Linn. Sp. Pl.* 80.—Europ.; As. bor.
pratensis, Sibth. & Sm. Fl. Graec. Prod. i. 67 = caryophyllea.
precatoria, Thuill. Fl. Par. ed. II. 58 = Arrhenatherum avenaceum.
Preslii, *Kunth, Enum. Pl.* i. 304.—Chili.
provincialis, Nym. Syll. Suppl. 71 = capillaris.
provincialis, Poir. Encyc. Suppl. i. 541 = Danthonia provincialis.
pseudolucida, Schur, in Oestr. Bot. Zeitschr. x. (1860) 72 = pubescens.
pseudoviolacea, *Kern. ex Dalla Torre, Atl. Alpenfl.* 228.—Europ. alp.
puberula, Guss. in Tenore, Prod. Fl. Nap. App. v. = Koeleria macilenta.
pubescens, *Huds. Fl. Angl.* 42.—Europ.; As. bor.
pulchella, Beauv. Agrost. 154, nomen = Aira lendigera.
pumila, Desf. Fl. Atlant. i. 103 = Trisetum pumilum.
pumila, Urv. in Mém. Soc. Linn. Par. i. (1822) 267 = Trisetum aureum, *Tenore*.
purpurascens, DC. Cat. Hort. Monsp. 82 = Trisetum pratense.
purpurascens, De Not. Ind. Sem. Genuen. (1847); Ann. Sc. Nat. Sér. III. ix. (1848) 323 = Trisetum alpestre.
purpurea, *Gueldenst. ex Ledeb. Fl. Ross.* iv. 419.—Rossia.
purpurea, Linn. f. Suppl. 112 = Danthonia purpurea.
pusilla, [Weber, in] Wigg. Prim. Fl. Hols. 9 = Aira praecox.
quadridentula, *Doell, in Mart. Fl. Bras.* ii. III. 100.—Bras.
quadriseta, Labill. Nov. Holl. Pl. i. 25. t. 32 = Deyeuxia quadriseta.
quinqueseta, Steud. in Flora, xii. (1829) II. 485 = Trisetum Steudelii.
racemosa, *Thuill. Fl. Par.* ed. II. 59.—Gallia.
Radula, *Franch. & Sav. Enum. Pl. Jap.* ii. 174.—Japon.
redolens, Pers. Syn. i. 100 = Hierochloë redolens.
rigida, Bieb. Fl. Taur. Cauc. i. 77 = Trisetum rigidum.
rigida, *Steud. in Flora*, xii. (1829) II. 482.—Afr. austr.
Rotae, De Not. ex Parl. Fl. Ital. i. 264 = Trisetum argenteum.
rubra, Zuccagni, in Roem. Collect. 126 = sativa.
rupestris, J. F. Gmel. Syst. ? 94 = Agrostis rupestris.
Ruprechtii, *Ledeb. Fl. Ross.* iv. 418.—Rossia arct.
sativa, *Linn. Sp. Pl.* 79.—Terr. temp. (cult.).
scabriuscula, Lag. in Varied. Cienc. ii. IV. (1805) 212 = Koeleria scabriuscula.
scabrivalvis, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 67.—Chili.
Schelliana, *Hack. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 342.—Hab.?
Scheuchzeri, *All. Fl. Pedem.* ii. 255.—Europ.; Reg. Caucas.
secalina, Salisb. Prod. 22 = Bromus secalinus.
secunda, Salisb. l. c. = Arrhenatherum avenaceum.
sedenensis, DC. Fl. Fr. iii. 719 = montana.
segetalis, Bianca, ex Nym. Consp. 810 = barbata.
sempervirens, Bory & Chaub. Fl. Pélop. 7. n. 179 = Heldreichii.
sempervirens, Host, Gram. iii. 28. t. 41 = Parlatorii.
sempervirens, Bess. Enum. Pl. Volh. 6 = Besseri.
sempervirens, Lapeyr. ex Benth. Cat. Pl. Pyr. 65 = montana.
sempervirens, Ledeb. Fl. Alt. i. 89 = desertorum.
sempervirens, *Vill. Prosp.* 17.—Europ.; As. bor.
sesquiflora, *Ledeb. Fl. Ross.* iv. 419.—As. or.
sesquitertia, Bieb. Fl. Taur. Cauc. i. 77 = Trisetum rigidum.
sesquitertia, Comol. Prod. 18 = lucida.
sesquitertia, Host, Syn. Fl. Austr. 60 = Trisetum alpestre.
sesquitertia, Hort. ex Steud. Nom. ed. I. 95 = barbata.
sesquitertia, Linn. Mant. i. 34 = pubescens.
sesquitertia, Tenore, ex Steud. Nom. ed. II. i. 173 = sempervirens.
sesquitertia, Willd. Sp. Pl. i. 448 = Trisetum argenteum.
setacea, Baumg. ex Schur, Enum. Pl. Transs. 735 = Agrostis rupestris.
setacea, Baumg. l. c. 753 = Deschampsia flexuosa.
setacea, Parl. Fl. Ital. i. 133 = Parlatorii.

AVENA:—
setacea, Muhl. ex Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 87 = Aristida dichotoma.
setacea, *Vill. Prosp.* 16.—Europ. austr.
setifolia, Brot. Fl. Lusit. 108 = Arrhenatherum pallens.
sibirica, Linn. Sp. Pl. 79 = Stipa sibirica.
Sibthorpii, Nym. Syll. 412 = bromoides.
sicula, Spreng. Syst. i. 335 = Trisetum aureum, *Tenore*.
Smithii, *Porter, ex A. Gray, Man. Bot. U. St.* ed. V. 640.—Am. bor.
spicaeformis, Auct. ex Roem. & Schult. Syst. ii. 690 = Danthonia spicaeformis.
spicata, All. ex Kunth, Enum. Pl. i. 312 = Triodia decumbens, Danthonia provincialis.
spicata, Ell. Sketch, i. 174 = Danthonia sericea.
spicata, Gilib. Exercit. 540 = pratensis.
spicata, Linn. Sp. Pl. 80 = Trisetum subspicatum.
spicata, Willd. Sp. Pl. i. 453, partim = Danthonia provincialis.
splendens, Boiss. Elench. 88 = albinervis.
splendens, Auct. Gall. ex Steud. Nom. ed. II. i. 173 = pratensis.
splendens, Guss. Fl. Sic. Prod. i. 126 = Trisetum pratense.
squarrosa, *Schrank, in Denkschr. Regensb. Ges.* i. II. (1818) 7.—Labrador.
sterilis, Delile, ex Boiss. Fl. Orient. v. 543 = fatua.
sterilis, *Linn. Sp. Pl.* ed. II. 118.—Reg. Mediterr.; Oriens.
stipaeformis, *Linn. Mant.* 34.—Afr. austr.
stipoides, Willd. ex Steud. Nom. ed. II. i. 173; ii. 146 = Oryzopsis fimbriata.
striata, Lam. Encyc. i. 332 = sempervirens.
striata, Trin. in Mém. Acad. Pétersb. Sér. VI. iv. II. (1838) 28 = callosa.
striata, *Michx. Fl. Bor. Am.* i. 73.—Am. bor.
striata, *Schousb. ex Willk. & Lange, Prod. Fl. Hisp.* i. 60.—Hispan.
striata, Vis. Fl. Dalm. i. 70 = convoluta.
stricta, Host, Fl. Austr. i. 127 = Danthonia provincialis.
strigosa, *Schreb. Spic.* 52.—Europ.; Reg. Casp.
strigosa, Vogler, Schediasma Bot. 22 = Gaudinia tenuis.
subalpestris, Hartm. Handb. ed. III. ex ej. ed. IV. 29 = Trisetum agrostideum.
subcylindrica, Ehrenb. ex Boiss. Fl. Orient. v. 536 = Trisetum glumaceum.
subspicata, [Clairv.] Man. Herb. 17 = Trisetum subspicatum.
subulata, Lam. Illustr. n. 1113 = setacea.
subvillosa, Schur, Enum. Pl. Transs. 763 = convoluta.
sulcata, *J. Gay, in Delastre, Fl. Vienne*, 477. t. 4.—Hispan.
sylvatica, Salisb. Prod. 24 = Holcus mollis.
symphicarpa, Trin. ex Steud. Nom. ed. II. i. 173 = hirta.
syriaca, Boiss. & Bal. ex Boiss. Fl. Orient. v. 542 = sterilis.
tatarica, Ard. in Saggi Acc. Padov. ii. (1789) 101. t. 1 = orientalis.
taygetana, *Steud. Syn. Pl. Gram.* 233.—Graecia.
Tenorii, Nym. Consp. 814 = corymbosa.
tenuiflora, *Bertol. ex Reichb. Ic. Fl. Germ.* i. 41. n. 1688.—Italia.
tenuis, Moench, Meth. 195 = Ventenata avenacea.
Thorei, DC. & Duby, Bot. Gall. i. 514 = Arrhenatherum pallens.
tolucensis, H. B. & K. Nov. Gen. et Sp. i. 148 = Trisetum toluccense.
triaristata, Vill. Prosp. 17 = Ventenata avenacea.
trichophylla, C. Koch, in Linnaea, xxi. (1848) 393 = fatua.
trichopodia, *J. Presl, in Rel. Haenk.* i. 254.—Mexic.
triseta, Thunb. Prod. Pl. Cap. 22 = Danthonia Thunbergii.
trisperma, Auct. ex Roem. & Schult. Syst. ii. 669 = sativa.
triticoides, *Lindl. in Mitch. Journ. Austral.* 365.—Austral.
truncata, Dulac, Fl. Hautes-Pyr. 83 = bromoides.
tuberosa, Gilib. Exercit. ii. 538 = Arrhenatherum avenaceum.
uniflora, *Parl. Pl. Nov.* 84.—Ins. Canar.
unilateralis, Brouss. ex Roem. & Schult. Syst. ii. 669 = orientalis.
valesiaca, Nym. Syll. 414 = Trisetum Cavanillesii.

AVENA:—
varia, Schur, in Oestr. Bot. Zeitschr. x. (1860) 75 = Trisetum carpathicum.
velutina, *Nym. Consp.* 812 = Trisetum velutinum.
ventricosa, *Balansa, in Bull. Soc. Bot. Fr.* i. (1854) 14.—Algeria.
versicolor, Baumg. ex Steud. Syn. Pl. Gram. 233 = Baumgartenii.
versicolor, Vill. Prosp. 17 = Scheuzeri.
versicolor, St. Am. Fl. Agen. 48 = sulcata.
vilis, *Wallr. Beitr. Fl. Hercyn.* 173; *et in Linnaea*, xiv. (1840) 543.—German.
villosa, Bertol. Exc. de Re Herb. 6. n. 3 = Trisetum villosum.
villosa, Lag. ex Roem. & Schult. Syst. ii. 670 = hirsuta.
virescens, *Regel, in Act. Hort. Petrop.* vii. (1880) 635.—Turkestan.
viridis, H. B. & K. Nov. Gen. et Sp. i. 147 = Deyeuxia viridis, *Fourn.*
Wiestii, *Steud. Syn. Pl. Gram.* 231.—Aegypt.; Arab.; Persia.
Wilhelmsii, Spreng. Syst. i. 333 = Trisetum rigidum.

AVENELLA, Koch, ex Steud. Nom. ed. II. i. 173 (1840); Parl. Fl. Ital. i. 244 (1848) = **Deschampsia**, Beauv. (Gramin.).
cuprina, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 85 = D. flexuosa.
flexuosa, Parl. Fl. Ital. i. 246 = D. flexuosa.
rubra, Jord. ex Nym. Consp. 808 = D. flexuosa.
uliginosa, Parl. Fl. Ital. i. 246 = D. uliginosa.

AVERRHOA, Linn. Syst. ed. I (1735). *GERANIACEAE*, Benth. & Hook. f. i. 277.
CARAMBOLA, Adans. Fam. ii. 508 (1763).
acida, Linn. Sp. Pl. 428 = Phyllanthus Cicca.
acutangula, Stokes, Bot. Mat. Med. ii. 543 = A. Carambola.
Bilimbi, *Linn. Sp. Pl.* 428.—Ind. or.; China.
Carambola, *Linn. Sp. Pl.* 428.—Ind. or.; China.
frondosa, Salisb. Prod. 318 = Phyllanthus Cicca.
obtusangula, Stokes, Bot. Mat. Med. ii. 542 = A. Bilimbi.
pentandra, Blanco, Fl. Filip. ed. I. 392 = Connaropsis philippica.
sinica, Hance, in Walp. Ann. ii. 241 = Rourea microphylla.

AVERRHOIDIUM, Baill. Adansonia, xi. 244 (1874). *SAPINDACEAE*.
Gardnerianum, *Baill. l. c.*—Bras.

AVERSIA, G. Don, Gen. Syst. iii. 92 (1834) = ARVERSIA, Cambess. = **Polycarpon**, Linn. (Caryophyll.).

AVICENNIA, Linn. Syst. ed. I (1735); Gen. ed. I. 27 (1737). *VERBENACEAE*, Benth. & Hook. f. ii. 1160.
BONTIA, Linn. Gen. ed. I. 180 (1737).
DONATIA, Loefl. Iter, 193 (1758).
HALODENDRUM, Thou. Gen. Nov. Madag. 8 (1806).
RACKA, J. F. Gmel. Syst. 245 (1791).
SCEURA, Forsk. Fl. Aegypt. Arab. 37 (1778).
UPATA, Adans. Fam. ii. 201 (1763).
africana, Beauv. Fl. Owar. i. 80. t. 47 = officinalis.
alba, Blume, Bijdr. 821 = officinalis.
elliptica, Thunb. Pl. Bras. Dec. iii. 37 = officinalis.
floridana, Rafin. Atl. Journ. 148 = officinalis.
intermedia, Griff. in Trans. Linn. Soc. xx. (1846) 6 = officinalis.
Lamarckiana, Presl, Bot. Bemerk. 99 = officinalis.
Meyeri, Miq. in Linnaea, xviii. (1844) 262 = officinalis.
nitida, *Jacq. Enum. Pl. Carib.* 25.—Am. trop.; Afr. trop.
oblongifolia, Nutt. ex Chapm. Fl. S. U. St. 310 = officinalis.
obovata, Griff. Notul. iv. 189 = officinalis.
Oepata, Buch.-Ham. in Trans. Linn. Soc. xvii. (1837) 221 = officinalis.
officinalis, *Linn. Sp. Pl.* 110.—Littora tropica.
resinifera, Forst. f. Pl. Escul. 72 = officinalis.
tomentosa, Jacq. Enum. Pl. Carib. 25 = officinalis.

AVICEPS, Lindl. Gen. et Sp. Orch. 345 (1838) = **Satyrium**, Sw. (Orchid.).
pumila, Lindl. l. c. 346 = S. pumilum.

AVICULARIA, Steud. Nom. ed. II. i. 174 (1840) = **Polygonum**, Tourn.
indica, Didrichs. in Bot. Notiscr, (1850) 187 = P. plebeium.
maritima, Didrichs. l. c. = P. maritimum.
vulgaris, Didrichs. l. c. = P. aviculare.

AVIUNCULUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 367 = **Coronilla**, Linn. (Legum.).
scorpioides, Fourr. l. c. = C. scorpioides.

AVORNELA, Rafin. Sylva Tellur. 24 (1838) = **Genista**, Linn. (Legum.).
canariensis, Rafin. l. c. = G. canariensis.
candicans, Rafin. l. c. = Cytisus candicans.
linifolia, Rafin. l. c. = Cytisus linifolia.

AWAYUS, Rafin. Sylva Tellur. 151 (1838) = **Spiraea**, Linn. (Rosac.).
japonicus, Rafin. l. c. 151 = S. japonica.

AXANTHES, Blume, Bijdr. 1002 (1826) = **Urophyllum**, Wall. (Rubiac.).
arborea, Blume, l. c. 1003 = U. glabrum.
Blumeana, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 145 = U. Blumeanum.
ceylanica, Wight, l. c. 146 = U. zeylanicum.
corymbosa, Blume, Bijdr. 1003 = U. corymbosum.
elliptica, Wight, in Calc. Journ. Nat. Hist. vii. (1847) 144 = U. ellipticum.
enneandra, Wight, l. c. 144 = Aulacodiscus premnoides.
Griffithiana, Wight, l. c. 147 = U. Griffithianum.
hirsuta, Wight, l. c. 148 = U. hirsutum.
longifolium, Wight, l. c. 145 = U. longifolium.
macrophylla, Blume, Bijdr. 1002 = U. macrophyllum.
philippensis, Cham. & Schlecht. ex Steud. Nom. ed. II. i. 174.—Ins. Luzon.
strigosa, Blume, Bijdr. 1003 = U. strigosum.
timoriensis, DC. Prod. iv. 441.—Ins. Timor.
tomentosa, Blume, ex DC. l. c. 440 = U. tomentosum.
zeylanica, Wight, Ic. t. 1163 = U. zeylanicum.

AXANTHOPSIS, Korth. in Nederl. Kruidk. Arch. ii. (1851) 195 = **Urophyllum**, Wall. (Rubiac.).
elongata, Korth. l. c. = U. elongatum.

AXENFELDIA, Baill. Etud. Gén. Euph. 419 (1858) = **Mallotus**, Lour. (Euphorb.).
intermedia, Baill. l. c. = M. muricatus.

AXIA, Lour. Fl. Cochinch. 35 (1790). *VALERIANACEAE*?
cochinchinensis, *Lour. l. c.* 36.—Cochinch.

AXILLARIA, Rafin. in Am. Monthly Mag. (1818) 266 = **Polygonatum**, Adans. (Liliac.).

AXINAEA, Ruiz & Pav. Prod. 68 (1794). *MELASTOMACEAE*, Benth. & Hook. f. i. 749.
CHASTENAEA, DC. Prod. iii. 102 (1828).
bracteata, Zipp. ex Blume, in Flora, (1831) 519 = Medinilla bracteata.
crassinoda, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 69.—Peruv.
dependens, *Ruiz & Pav. Fl. Per.* iv. t. 411.—Peruv.
floribunda, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 69.—Ecuador.
glandulosa, *Ruiz & Pav. Fl. Per.* iv. t. 412.—Peruv.
grandifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 69.—Venezuela.
lanceolata, *Ruiz & Pav. Fl. Per.* iv. 409; *Syst.* i. 122.—Peruv.
lepidota, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 69.—Ecuador.
macrophylla, *Triana, l. c.*—Ecuador.
Merianiae, *Triana, l. c.*—Venezuela.
muricata, D. Don, in Mem. Wern. Soc. iv. (1823) 321 = Miconia muricata.
purpurea, Ruiz & Pav. Fl. Per. iv. 410; Syst. i. 122 = Meriania radula.
Radula, Benth. Pl. Hartw. 130 = Meriania Radula.
sclerophylla, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 68.—Ecuador.
scutigera, *Triana, l. c.* 69.—N. Granat.
sessilifolia, *Triana, l. c.* 68.—Peruv.

AXINEA, Juss. in Dict. Sc. Nat. xxix. (1823) 508 = praec.

AXINANDRA, Thw. in Hook. Kew Journ. vi. (1854) 66. t. 1. C. *LYTHRARIEAE*, Benth. & Hook. f. i. 784.
alata, *Baill. Adansonia*, xii. (1876) 86.—Malaya.
Beccariana, *Baill. l. c.* 85.—Malaya.
coriacea, *Baill. l. c.*—Malaya.
Maingayi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 581.—Malacca.
zeylanica, *Thw. in Hook. Kew Journ.* vi. (1854) 67.—Zeylan.

AXINANTHERA, Karst. in Linnaea, xxx. (1859–60) 157 = **Bellucia**, Neck. (Melast.).
Bellucia, Karst. l. c. = B. Axinanthera.
macrophylla, Karst. l. c. = B. superba?

AXINIPHYLLUM, Benth. in Hook. Ic. Pl. xii. 16. t. 1118 (1872). *COMPOSITAE*, Benth. & Hook. f. ii. 362.
corymbosum, *Benth. l. c.*—Mexic.
tomentosum, *Benth. l. c.* sub t. 1118.—Mexic.

AXIRIS, Linn. Diss. Chen. 43 (1751) = **Axyris**, Linn. (Chenop.).

AXOLOPHA, Alef. in Oestr. Bot. Zeitschr. xii. (1862) 258 = **Lavatera**, Linn. (Malvac.).
aegyptia, Alef. l. c. 259 = Malva aegyptia.
althaeoides, Alef. l. c. = Malva althaeoides.
hirsuta, Alef. l. c. = Althaea hirsuta.
Ludwigii, Alef. l. c. = Althaea Ludwigii.
maritima, Alef. l. c. = L. maritima.
Wigandi, Alef. l. c. = L. maritima.

AXOLUS, Rafin. Sylva Tellur. 61 (1838) = **Cephalanthus**, Linn. (Rubiac.).
angustifolius, Rafin. l. c. = C. angustifolius.

AXONOPUS, Beauv. Agrost. 12 (1812) = **Paspalum**, Linn. (Gramin.).
aureus, Beauv. l. c. = P. aureum.
cimicinus, Beauv. l. c. = Panicum cimicinum.
compressus, Beauv. l. c. = P. compressum.
corymbosus, Schult. Mant ii. 177 = Panicum corymbosum.
digitatus, Beauv. Agrost. 12 = P. digitatum.
paniceus, Beauv. l. c. = Panicum filiforme.
Poiretii, Roem. & Schult. Syst. ii. 318 = P. compressum.
savannarum, Schlecht. in Linnaea, xxvi. (1853–55) 132 = P. savannarum.

AXONOTECHIUM, Fenzl, in Ann. Wien. Mus. i. (1836) 354 = **Orygia**, Forsk. (Ficoid.).
trianthemoides, Fenzl, l. c. 355 = O. decumbens.

AXYRIS, Linn. Syst. ed. VI. Add. (1748); Amoen. Acad. 24 (1751). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 56.
amaranthoides, *Linn. Sp. Pl.* 979.—As. bor.
ceratoides, Linn. l. c. = Eurotia ceratoides.
hybrida, Linn. l. c. 980 = amaranthoides.
Moorcroftiana, R. Br. in Wall. Cat. n. 6950 = Eurotia ceratoides.
pentandra, *Jacq. Am. Select.* 244.—Cuba.
prostrata, Linn. Sp. Pl. 980 = amaranthoides?
sphaerosperma, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. (1839) 46.—Mont. Altai.
triflora, Moench, Meth. Suppl. 120 = amaranthoides.
villosa, *Moq. in DC. Prod.* xiii. II. 117.—Afghan.

AYAPANA, Spach, Hist. Vég. Phan. x. 290 (1841) = **Eupatorium**, Linn. (Compos.).
officinalis, Spach, l. c. = E. Ayapana.

AYDENDRON, Nees, in Linnaea, viii. (1833) 36. *LAURINEAE*, Benth. & Hook. f. iii. 153.
ENDLICHERIA, Nees, in Linnaea, viii. (1833) 37.
GOEPPERTIA, Nees, Syst. Laurin. 365 (1836).
SCHAUERA, Nees, in Lindl. Nat. Syst. ed. II. 202 in adnot. (1836).
aciphyllum, *Nees, in Linnaea*, xxi. (1848) 496.—Guiana.
affinis, *Meissn. in DC. Prod.* xv. I. 92.—Bras.
amazonicum, *Meissn. l. c.* 89.—Bras.
argenteum, *Griseb. Fl. Brit. W. Ind.* 285.—Ind. occ.
bracteatum, *Nees, Syst. Laurin.* 256.—Ins. S. Vincent.

AYDENDRON:—
brasiliense, *Meissn. in Mart. Fl. Bras.* v. II. 180.—Bras.
Cannella, *Meissn. in DC. Prod.* xv. I. 90.—Guiana.
cayennense, *Meissn. l. c.* 95.—Guiana.
citrifolium, Griseb. Fl. Brit. W. Ind. 284 = trinitatis.
citrifolium, *Nees, Syst. Laurin.* 257.—Bras.
cubense, *A. Rich. Fl. Cub. Faner.* ii. 187.—Cuba.
Cujumary, *Nees, Syst. Laurin.* 247.—Bras.
curviramcum, *Meissn. in DC. Prod.* xv. I. 190.—Guiana.
desertorum, *Nees, Syst. Laurin.* 260.—Bras.
firmulum, *Nees, in Linnaea*, viii. (1836) 36.—Bras.
floribundum, *Meissn. in DC. Prod.* xv. I. 88.—Bras.
Gardneri, *Meissn. l. c.* 89.—Bras.
glaucum, *Klotzsch & Karst. ex Meissn. l. c.* 90.—N. Granat.
goyazense, *Meissn. l. c.* 94.—Bras.
Hostmannianum, *Nees, in Linnaea*, xxi. (1848) 499.—Guiana.
intermedium, *Meissn. in DC. Prod.* xv. I. 89.—Bras.
laevigatum, *Meissn. l. c.* 88.—Bras.
Laurel, *Nees, Syst. Laurin.* 249.—Venezuela.
macrophyllum, *Meissn. in DC. Prod.* xv. I. 92.—N. Granat.
microbotryum, *Nees, Syst. Laurin.* 254.—Ins. Portorico.
Muca, *Nees, in Linnaea*, xxi. (1848) 498.—Peruv.
nitidum, *Meissn. in DC. Prod.* xv. I. 95.—Bras.
obtusifolium, *Nees, Syst. Laurin.* 258.—Am. austr.
pachycarpum, *Meissn. in DC. Prod.* xv. I. 95.—Guiana.
pallidum, *Meissn. l. c.* 93.—Bras.
panurense, *Meissn. l. c.* 89.—Bras.
parviflorum, *Meissn. l. c.* 90.—Bras.
permolle, *Nees, Syst. Laurin.* 246.—Bras.
piauhyense, *Meissn. in DC. Prod.* xv. I. 93.—Bras.
riparium, *Nees, in Linnaea*, xxi. (1848) 497.—Guiana.
robustum, *Klotzsch & Karst. ex Meissn. in DC. Prod.* xv. I. 90.—N. Granat.
salicifolium, *Nees, Syst. Laurin.* 252.—Ind. occ.; Guian.
Sellowii, *Meissn. in DC. Prod.* xv. I. 88.—Bras.
sericeum, *Griseb. Fl. Brit. W. Ind.* 285.—Ind. occ.
suaveolens, *Nees, in Linnaea*, viii. (1833) 37.—Bras.
tenellum, *Meissn. in DC. Prod.* xv. I. 93.—Bras.
trinitatis, *Meissn. l. c.* 91.—Ins. Trinit.
verrucosum, *Nees, Syst. Laurin.* 259.—Bras.
verticillatum, *Meissn. in DC. Prod.* xv. I. 91.—Ins. S. Vincent.

AYENIA, Loefl. It. 199 (1758). *STERCULIACEAE*, Benth. & Hook. f. i. 225.
CYBIOSTIGMA, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 155.
DAYENIA, Mill. Gard. Dict. ed. VII. (1759).
JUNGIA, Loefl. It. 199 (1758).
abatilifolia, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 569 = magna.
acalyphifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 50.—Reg. Argent.
angustifolia, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 34.—Bras.
cardiopetala, Sesse, ex DC. Prod. i. 488 = magna.
cordifolia, *Sesse, ex DC. l. c.*—Mexic.
cordobensis, *Hieron. Ic. Pl. Argent.* 50.—Reg. Argent.
euphrasiaefolia, *Griseb. Cat. Pl. Cub.* 29.—Cuba.
hirta, *A. St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 34.—Bras.
inermis, Medic. ex Steud. Nom. ed. II. i. 174 = pusilla.
laevigata, *Sw. Prod. Veg. Ind. Occ.* 97.—Ins. Jamaic.
lingulata, *Griseb. in Goett. Abh.* xxiv. (1879) 50.—Reg. Argent.
magna, *Linn. Syst.* ed. X. 1247.—Mexic.
mexicana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 569.—Mexic.
microphylla, *A. Gray, in Smithson. Contr.* iii. (1852) 24.—N. Mexic.
ovata, *Hemsl. Diag. Pl. Nov.* 4.—Mexic.
pusilla, *Linn. Syst.* ed. X. 1247.—Ins. Carib.; Peruv.
rotundifolia, *Hemsl. Diag. Pl. Nov.* 4.—Mexic.
sidaefolia, Hemsl. Biol. Centr. Am. Bot. i. 135 = mexicana.
sidaefolia, DC. Prod. i. 488 = tomentosa.

AYENIA:—
sidaeformis, Loefl. It. 257 = tomentosa.
stipularis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 334.—N. Granat.
tomentosa, *Linn. Syst.* ed. X. 1247.—Venezuela.

AYLANTUS, Juss. Gen. 373 (1789) = **Ailanthus**, Desf. (Simarub.).

AYLMERIA, Mart. Amaranth. 68; et in Nov. Act. Nat. Cur. xiii. (1826) 276 = **Polycarpaea**, Lam. (Caryophyll.).
rosea, Mart. ll. cc. = P. violacea.
violacea, Mart. ll. cc. = P. violacea.
Zollingerii, Fenzl, in Zoll. Syst. Verz. Ind. Archip. 141.—Malaya.

AYPARIA, Rafin. Sylva Tellur. 154 (1838) = **Symplocos**, Linn. (Styrac.).
crenata, Rafin. l. c. (= *Dicalix cochinchinensis*, Lour.).—Cochinch.

AYTONIA, Linn. f. Suppl. 49, 303 (1781) = **Aitonia**, Thunb. (Sapind.).

AZADIRACHTA, A. Juss. in Mém. Mus. Par. xix. (1830) 220 (Mél. 68) (1830) = **Melia**, Linn.
indica, A. Juss. ll. cc. 221, 69 = Melia Azadirachta.

AZALEA, Gaertn. Fruct. i. 301. t. 63 (1788), pro parte = **Loiseleuria**, Desv. (Ericac.).

AZALEA, Linn. Syst. ed. I (1735) = **Rhododendron**, Linn. (Ericac.).
alba, Blume, ex DC. Prod. vii. 721 = R. album.
alba, Sweet, in Ann. Fromont, i. (1829) 104 = R. ledifolium.
amoena, Lindl. in Paxt. Flow. Gard. iii. (1852) t. 89 = R. indicum.
arborea, Bart. ex Steud. Nom. ed. II. i. 174 = R. arborescens.
arborea, Linn. Sp. Pl. 150 = R. ponticum.
arborescens, Pursh, Fl. Am. Sept. i. 153 = R. arborescens.
aurantiaca, Dietr. Darst. Vorz. Zierpfl. Heft i. 5. t. 1 = R. calendulaceum.
Bealii, Fortune, ex C. Koch, Dendrol. ii. II. 175 = R. sublanceolatum.
bicolor, Pursh, Fl. Am. Sept. i. 153 = R. nudiflorum.
bullata, Forst. ex Steud. Nom. ed. II. i. 174 = Lebetanthus americana.
Burgeri, Miq. Ann. Mus. Bot. Lugd. Bat. i. 34 = R. indicum.
calendulacea, Hook. & Arn. Bot. Beech. Voy. 361 = R. occidentale.
calendulacea, Michx. Fl. Bor. Am. i. 151 = R. calendulaceum.
californica, Torr. & Gray, ex Durand, in Journ. Acad. Nat. Sc. Philad. Ser. II. iii. (1855) 94 = R. californicum.
calycina, Lem. Jard. Fl. iii. (1853) t. 257 = R. indicum.
canescens, Michx. Fl. Bor. Am. i. 150 = R. nudiflorum.
celebica, Blume, ex DC. Prod. viii. 725 = R. malayanum.
chinensis, Steud. Nom. ed. II. i. 175 = R. sinense.
crispiflora, Hook. Bot. Mag. t. 4726 = R. indicum.
crocea, Hoffmgg. Verz. Pfl. Nachtr. iii. 22 = R. calendulaceum?
dahurica, C. Koch, Dendrol. ii. II. 181 = R. dauricum.
Danielsiana, Paxt. Mag. Bot. i. 129 = R. indicum.
decumbens, DC. Prod. vii. 718 = R. indicum.
Farrerae, C. Koch, Dendrol. ii. II. 178 = R. Farrerae.
ferruginosa, Pall. Fl. Ross. ii. t. 70 = R. Anthopogon.
fimbriata, Dum.-Cours. Bot. Cult. ed. II. iii. 337.—China.
flammea, Pritz. Ic. Ind. 131 = R. calendulaceum.
flava, Hoffmgg. Verz. Pfl. Nachtr. ii. 62 = R. calendulaceum?
fragrans, Adams, in Mém. Acad. Pétersb. ii. (1808) 332 = R. Anthopogon.
fragrans, Rafin. Ann. Nat. 12 = R. arborescens.
fulva, Michx. ex Lam. Journ. Hist. Nat. i. (1792) 410 = R. nudiflorum.
gandavensis, Hort. ex C. Koch, Dendrol. ii. II. 185 = R. ponticum.
glauca, Lam. Illustr. 493. t. 110 = R. viscosum.
glauca, Meerb. Pl. Sel. Ic. t. 6 = R. glaucum?

AZALEA:—
hispida, Pursh, Fl. Am. Sept. i. 154 = R. viscosum.
indica-alba, Lindl. in Bot. Reg. t. 811 (1824) = R. ledifolium.
indica, Linn. Sp. Pl. 150 = R. indicum.
japonica, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 400.—Japon.
javanica, Blume, ex DC. Prod. vii. 721 = R. javanicum.
Jodogawa, Hassk. Cat. Hort. Bog. Alt. 162 = R. Keiskei.
lapponica, Linn. Sp. Pl. 151 = R. lapponicum.
lapponica, Pall. Fl. Ross. ii. 52. t. 70. f. 1 = R. Anthopogon.
ledifolia, Hook. Bot. Mag. t. 2901 = R. ledifolium.
lilliflora, Poit. in Rev. Hortic. (1869) 299 = R. ledifolium.
linariaefolia, Carr. l. c. (1872) 100 = seq.?
linearifolia, Hook. f. Bot. Mag. t. 5769 = R. linearifolium.
lucida, Meerb. Pl. Sel. Ic. 7 = R. lucidum.
Ludoviciae, Herb. in Bot. Reg. xxxi. 60.—Hab.?
lugens, Dum.-Cours. Bot. Cult. ed. II. iii. 337.—China.
lutea, Linn. Sp. Pl. 150 = R. nudiflorum.
macrantha, Bunge, Enum. Pl. Chin. Bor. 41 = R. indicum.
mollis, Blume, Bijdr. 853 = R. molle.
Morterii, Sweet, ex Loud. Hort. Brit. Suppl. iii. 496 = R. calendulaceum.
Mortieriana, Hort. ex Spae, in Ann. Gand. ii. (1846) 325 = R. calendulaceum.
mucronata, Blume, Bijdr. 853 = R. ledifolium.
multiflora, Turcz. ex DC. Prod. vii. 715 = R. Anthopogon.
myrtifolia, Champ. in Hook. Bot. Mag. lxxvii. (1851) sub t. 4609 = R. ovatum.
narcissiflora, Fortune, ex Planch. in Rev. Hortic. Sér. IV. iii. (1854) 67 = R. ledifolium.
nitida, Pursh, Fl. Am. Sept. i. 153 = R. nitidum.
nivea, Dehnh. Cat. Pl. Hort. Camald. ed. II. 17.—Hab.?
nudiflora, Linn. Sp. Pl. ed. II. 214 = R. nudiflorum.
obtusa, Lindl. in Journ. Hort. Soc. i. (1846) 152 = R. indicum.
occidentalis, Torr. & Gray, ex Torr. in Pacif. Rail. Rep. iv. 116 = R. occidentale.
ovata, Lindl. in Journ. Hort. Soc. i. (1846) 149 = R. lapponicum.
pallida, Turcz. in Bull. Soc. Nat. Mosc. (1838) 96; xxi. (1848) II. 498.—Mont. Baical.
periclymena, Pers. Syn. i. 213 = R. nudiflorum.
periclymenoides, Michx. Fl. Bor. Am. i. 151 = R. nudiflorum.
phaenicea, Hort. ex Planch. in Fl. des Serres, Sér. I. ix. (1853–54) 79 = R. puniceum.
pilosa, Michx. in Lam. Journ. Nat. Hist. i. (1792) 410 = Menziesia globularis.
pontica, Linn. Sp. Pl. 150 = R. ponticum.
procumbens, Linn. l. c. 151 = Loiseleuria procumbens.
prolifera, Poit. in Ann. Fromont, i. (1829) 104 = R. Simsii.
punctata, Lour. Fl. Cochinch. 113; ed. Willd. i. 139 = R. Loureirianum.
punicea, Sweet, in Ann. Fromont, i. (1829) 104 = R. sublanceolatum.
purpurea, Dum.-Cours. Bot. Cult. ed. II. iii. 337.—Ind. or.
ramentacea, Lindl. Journ. Hort. Soc. iv. (1849) 291 = R. indicum.
Rawsonii, Paxt. Mag. Bot. ii. 121.—Hab.?
retusa, Blume, ex Steud. Nom. ed. II. i. 175 = R. retusum.
rosmarinifolia, Burm. f. Fl. Ind. 43 = R. ledifolium.
rubra, Meerb. Pl. Sel. Ic. 8.—Hab.?
scabra, Lodd. ex A. Gray, Syn. Fl. N. Am. ii. II. 41 = R. viscosum.
serpyllifolia, A. Gray, in Perry, Jap. Exped. ii. 315 = R. serpyllifolium.
Seymouri, Lindl. in Bot. Reg. t. 1975 = R. Seymouri.
sinensis, Lodd. Bot. Cab. t. 885 = R. sinense.
speciosa, Willd. Enum. Hort. Berol. Suppl. 10 = R. speciosum.
squamata, Lindl. in Journ. Hort. Soc. i. (1849) 152 = R. Farrerae.
tomentosa, Duroi, ex Steud. Nom. ed. I. 97 = Loiseleuria procumbens.
tubiflora, Blume, ex DC. Prod. vii. 725 = R. malayanum.

AZALEA:—
verticillata, Bart. ex Lodd. Cab. t. 1632.—Am. bor.
virens, Dum.-Cours. Bot. Cult. ed. II. iii. 335 = R. viscosum.
viscosa, Linn. Sp. Pl. 151 = R. viscosum.
vittata, Dum.-Cours. Bot. Cult. ed. II. iii. 337 = R. vittatum.
vittato-punctata, Lem. Illustr. Hortic. (1854) t. 20 = praec.?

AZALTEA, Walp. Rep. i. 539 (1842) = **Alzatea**, Ruiz & Pav. (Celastr.).

AZAMARA, Hochst. ex Reichb. Nom. 145 (1841) = **Schmidelia**, Linn. (Sapind.).
trifoliata, Hochst. ex Baker, in Oliver, Fl. Trop. Afr. 421 = S. africana.

AZANZA, Alef. in Bot. Zeit. xix. (1861) 298 = **Thespesia**, Corr. (Malvac.).
acuminata, Alef. l. c. 299 = T. populnea.
Lampas, Alef. l. c. 297 = T. Lampas.
Zollingeri, Alef. l. c. 298.—Java.

AZANZA, Moç. & Sessé, ex DC. Prod. i. 453 (1824) = **Hibiscus**, Linn. (Malvac.).
insignis, Moç. & Sessé, l. c. 454 = H. Azanza.

AZAOLA, Blanco, Fl. Filip. 402 (1837) = **Payena**, A. DC. (1844). (Sapot.).
Betis, Blanco, l. c. = P. Betis.
Leerii, Teijsm. & Binn. in Tijdschr. Nederl. Ind. vi. (1854) 116 = P. Leerii.

AZARA, Ruiz & Pav. Prod. 79. t. 36 (1799). *BIXINEAE*, Benth. & Hook. f. i. 127.
MYRTOPHYLLUM, Turcz. in Bull. Soc. Nat. Mosc. (1863) I. 604.
SILENIA, Bert. in Merc. Chil. (1829) ex Bull. Féruss. xx. (1830) 108.
alpina, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 30. t. 167.—Chili.
Berteroniana, *Steud. in Flora*, xxxix. (1856) 420.—Chili.
celastrina, *D. Don, in Edinb. N. Phil. Journ.* x. (Dec. 1830) 119.—Chili.
chiloënsis, *Hook. f. Fl. Antarct.* 243.—Chili.
dentata, *Ruiz & Pav. Syst.* 138.—Chili.
dubia, *Steud. in Flora*, xxxix. (1856) 418.—Chili.
Fernandesiana, C. Gay, Fl. Chil. i. 196 = serrata.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 144.—Chili.
hirtella, *Miq. in Linnaea*, xxv. (1852) 650.—Chili.
integrifolia, Bert. ex Steud. in Flora, xxxix. (1856) 420 = Berteroniana.
integrifolia, Miq. ex Steud. l. c. = Lechleriana.
integrifolia, *Ruiz & Pav. Syst.* 138.—Chili.
intermedia, *C. Gay, Fl. Chil.* i. 194.—Chili.
lanceolata, *Hook. f. Fl. Antarct.* 243.—Chili.
Lechleriana, *Steud. in Flora*, xxxix. (1856) 420.—Chili.
Lilen, Bert. in Merc. Chil. (1829).—Cf. Bull. Férussac, xx. (1830) 108 = celastrina, sparsiflora.
microphylla, *Hook. f. Fl. Antarct.* 244.—Chili.
salicifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 20.—Reg. Argent.
serrata, Bert. ex C. Muell. in Walp. Ann. vii. 226 = dubia.
serrata, *Ruiz & Pav. Prod.* 79. t. 36; *Syst.* 137.—Chili.
sparsiflora, *Steud. Nom.* ed. II. i. 175.—Chili.
tomentosa, *Bert. ex Steud. in Flora*, xxxix. (1856) 419.—Chili.
umbellata, *Phil. in Linnaea*, xxviii. (1856) 610.—Chili.
umbellata, Presl, Rel. Haenk. ii. 92 = Laetia apetala.
Valdiviae, *Lechler, ex Steud. in Flora*, xxxix. (1856) 420.—Chili.

AZAROLUS, Borkh. Handb. Forstb. ii. 1224 (1803) = **Pyrus**, Linn. (Rosac.).
alpina, Borkh. l. c. 1234 = P. alpina.
arbutifolia, Borkh. l. c. 1225 = P. arbutifolia.
Aria, Borkh. l. c. 1229 = P. Aria.
Chamaemespilus, Borkh. l. c. 1227 = P. Chamaemespilus.
crataegoides, Borkh. l. c. 1253 = Crataegus Azarolus.
heterophylla, Borkh. l. c. 1248.—Hab.?
hybrida, Borkh. l. c. 1239 = P. rotundifolia.
intermedia, Borkh. l. c. 1232 = P. intermedia.

AZAROLUS :—
maroccana, M. Roem. Syn. Rosifl. 133 = Crataegus Azarolus.
Oliveriana, M. Roem. l. c. = Crataegus pentagyna.
pinnatifida, Borkh. Handb. Forstb. ii. 1245 = Pyrus pinnatifida.
platyphylla, M. Roem. Syn. Rosifl. 133 = Crataegus pentagyna.
Pollvilleriana, Borkh. Handb. Forstb. ii. 1251 = P. Pollveria.
pubescens, M. Roem. Syn. Rosifl. 134 = Crataegus pubescens.
torminalis, Borkh. Handb. Forstb. ii. 1235 = P. torminalis.

AZEDARA, Rafin. Fl. Ludov. 135 (1817) = **Melia**, Linn.
speciosa, Rafin. l. c. = M. Azedarach.

AZEDARAC, Adans. Fam. ii. 342 (1763) = seq.

AZEDARACH, Tourn. ex Linn. Gen. ed. I. 127 (1737) = **Melia**, Linn.
Commelini, Medic. Bot. Beobacht. (1782) 164 = M. Azedarach.
deleteria, Medic. Malv. 115 = M. Azadirachta.
edule, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 5 = Sandoricum indicum.
fraxinifolia, Moench, Meth. Suppl. 58 = M. Azadirachta.
nigrum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 5.—Malaya.
odoratum, Noronha, l. c. = M. Azedarach.
ramiflorum, Noronha, l. c. = Epicharis cauliflora.
recisum, Noronha, l. c.—Malaya.

AZEREDIA, Arruda, ex Allemão, Desenb. cum Ic. (1846). *BIXINEAE*, Benth. & Hook. f. i. 125.
pernambucana, *Allemão, l. c.*—Bras.

AZIMA, Lam. Encyc. i. 343 (1789). *SALVADORACEAE*, Benth. & Hook. f. ii. 681.
ACTEGETON, Blume, Bijdr. 1143 (1826).
KANDENA, Rafin. Sylva Tellur. 138 (1838).
MONETIA, L'Hérit. Stirp. Nov. i. t. 1 (1784).
angustifolia, *A. DC. in DC. Prod.* xvii. 29.—Madag.
diacantha, Lam. Encyc. i. 343 = Plectronia parviflora.
nova, *Blanco, Fl. Filip.* ed. I. 68.—Ins. Philipp.
nova, J. F. Gmel. Syst. 262 = tetracantha.
sarmentosa, *Benth. & Hook. f. Gen.* ii. 681.—Burma; Malaya.
sarmentosa, Vill. ex Vidal, Sinops. t. 65. f. C = tetracantha ?
scandens, Baill. Hist. Pl. vi. 13 = sarmentosa.
tetracantha, *Lam. Encyc.* i. 343.—Ind. or.

AZOPHORA, Neck. Elem. ii. 361 (1790) = **Rhizophora**, Linn.

AZORELLA, Lam. Encyc. i. 344 (1783). *UMBELLIFERAE*, Benth. & Hook. f. i. 874.
BOLAX, Comm. ex Juss. Gen. 226 (1789).
CHAMITIS, Banks, ex Gaertn. Fruct. i. 94. t. 22 (1788).
DICHOPETALUM, F. Muell. in Hook. Kew Journ. vii. (1855) 378. t. 11.
FRAGOSA, Ruiz & Pav. Prod. Fl. Per. 43. t. 34 (1794).
MICROSCIADIUM, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 468.
OSCHATZIA, Walp. Ann. i. 340 (1849).
PECTOPHYTUM, H. B. & K. Nov. Gen. et Sp. v. 28. t. 425 (1821).
POZOA, Lag. Gen. et Sp. Nov. 13 (1816).
apoda, *A. Gray, Bot. U. St. Expl. Exped.* i. 699.—Chili.
aretioides, *Willd. ex DC. Prod.* iv. 77.—Ecuador.
biloba, *Wedd. Chlor. And.* ii. 196.—Bolivia.
bolacina, *Clos, in C. Gay, Fl. Chil.* iii. 80.—Chili.
bryoides, *Phil. in Linnaea*, xxviii. (1856) 695.—Chili.
caespitosa, *Cav. Ic.* v. 57. t. 484.—Chili.
caespitosa, Vahl, Symb. Bot. iii. 48 = glebaria.
Chamitis, Pers. Syn. i. 303 = filamentosa.
eladorrhiza, *Pers. l. c.*—Peruv.
compressa, Labill. Nov. Holl. Pl. i. 75. t. 101 = Siebera compressa.
corymbosa, *Pers. Syn.* i. 305.—Peruv.
crassifolia, Pers. l. c. 303 = caespitosa.

AZORELLA:—
crenata, *Pers. Syn.* 302.—Peruv.
cuneifolia, *F. Muell. ex Benth. Fl. Austral.* iii. 365.—Austral.
daucoides, Urv. in Mém. Soc. Linn. Par. iv. (1826) 613 = Orcomyrrhis andicola.
depauperata, *Phil. Fl. Atac.* 24.—Chili.
diapensioides, *A. Gray, Bot. U. St. Expl. Exped.* i. 702.—Peruv.
dichopetala, *Benth. Fl. Austral.* iii. 365.—Austral.
diversifolia, *Clos, in C. Gay, Fl. Chil.* iii. 86.—Chili.
filamentosa, *Lam. Encyc.* i. 344.—Terr. Magellan.
Gayana, *Phil. in Linnaea*, xxviii. (1856) 653.—Chili.
Gillicsii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 347.—Chili.
glabra, *Wedd. Chlor. And.* 67.—Bolivia.
glebaria, *A. Gray, Bot. U. St. Expl. Exped.* i. 701.—Chili.
gummifera, Poir. Encyc. Suppl. i. 551 = caespitosa.
Hookeriana, Clos, in C. Gay, Fl. Chil. iii. 81 = caespitosa.
incisa, *Wedd. Chlor. And.* ii. 197.—Bolivia.
integrifolia, Raeusch. Nom. ed. III. 84 = filamentosa.
lanceolata, Labill. Nov. Holl. Pl. i. 74. t. 69 = Siebera Billardieri.
linearifolia, Cav. Ic. v. 57 = Siebera lincarifolia.
lycopodioides, *Gaudich. in Ann. Sc. Nat.* Sér. I. v. (1825) 105. t. 3.—Reg. Magellan.
madreporica, *Clos, in C. Gay, Fl. Chil.* iii. 79.—Chili.
monantha, *Clos, l. c.*—Chili.
Muelleri, *Benth. Fl. Austral.* iii. 364.—Austral.
multifida, *Pers. Syn.* i. 303.—Peruv.
museoides, *Phil. in Linnaea*, xxviii. (1856) 695.—Chili.
nervosa, *Phil. l. c.* 696.—Chili.
ovata, Labill. Nov. Holl. Pl. i. 74. t. 100 = Siebera Billardieri.
pedunculata, *Willd. ex DC. Prod.* iv. 78.—Ecuador.
piligera, *Phil. in Linnaea*, xxviii. (1856) 652.—Chili.
pulvinata, *Wedd. Chlor. And.* ii. 194.—Ecuador.
Ranunculus, *Urv. in Mém. Soc. Linn. Par.* iv. (1826) 613.—Reg. Magell.
reniformis, Pers. Syn. i. 303.—Peruv.
Saxifraga, *Benth. Fl. Austral.* iii. 365.—Austral.
Selago, *Hook. f. Fl. Antarct.* 284.—Terr. Magell.; Ins. Kerguelen.
sessiliflora, *Phil. in Linnaea*, xxviii. (1856) 652.—Chili.
spinosa, *Pers. Syn.* i. 303.—Chili.
triscuspidata, Lam. Illustr. t. 189 = caespitosa.
triscuspidata, Poir. Encyc. Suppl. i. 551 = lycopodioides.
trifoliata, Walp. Ann. v. 62 = trifoliolata.
trifoliolata, *Hook. f. Fl. N. Zel.* i. (1853) t. 18.—N. Zel.
trifoliolata, *Clos, in C. Gay, Fl. Chil.* iii. 85. t. 30. f. 2.—Chili.
trifuscata, Hook. Ic. Pl. t. 539 = seq.
trifurcata, *Pers. Syn.* i. 303.—Reg. Magellan.
utriculata, *Griseb. in Goett. Abh.* vi. (1854) 122.—Reg. Magellan.

AZURINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Serophul.).
aphylla, Fourr. l. c. 128 = V. aphylla.

B

BABACTES, DC. in Meissn. Gen. 302; Comm. 211 (1840); et Prod ix. 260 (1845) = **Chirita**, Buch.-Ham. (Gesnerac.).
oblongifolia, DC. l. e. = C. acuminata.

BABBAGIA, F. Muell. Rep. Babb. Exped. 21 (1858). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 61.
aeroptera, *F. Muell. & Tate, in Trans. Roy. Soc. S. Austral.* vi. (1883) 108.—Austral.
dipterocarpa, *F. Muell. Rep. Babb. Exped.* 21.—Austral.
pentaptera, *F. Muell. & Tate, in Trans. Roy. Soc. S. Austral.* vi. (1883) 108.—Austral.
scleroptera, *F. Muell. in Wing, South Sc. Record*, (Nov. 1885).—Austral.

BABIANA, Ker-Gawl. in Bot. Mag. t. 576 (1802) *IRIDEAE*, Benth. & Hook. f. iii. 706.
ACASTE, Salisb. in Trans. Hort. Soc. i. (1812) 322.
angustifolia, *Sweet, Hort. Brit.* ed. I. 396; *Eckl. Verz.* 31.—Afr. austr.
atrocyanea, *Eckl. Verz.* 33.—Afr. austr.
atrodeltoidea, *Eckl. l. c.*—Afr. austr.
Bainesii, *Baker, in Journ. Bot.* xiv. (1876) 335.—Afr. austr.
caesia, *Eckl. Verz.* 32.—Afr. austr.
caeruleseens, *Eckl. l. c.*—Afr. austr.
cuneifolia, *Baker, in Journ. Bot.* xiv. (1876) 335.—Afr. austr.
densiflora, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 348.—Afr. austr.
disticha, *Ker-Gawl. in Bot. Mag.* t. 626.—Afr. austr.
Dregei, *Baker, in Journ. Bot.* xiv. (1876) 336.—Afr. austr.
Ecklonii, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 348.—Afr. austr.
fimbriata, *Baker, in Journ. Linn. Soc.* xvi. (1878) 166.—Afr. austr.
flabellifolia, *Harv. ex Klatt, in Linnaea*, xxxv. (1867–68) 380.—Afr. austr.
flavocaesia, *Eckl. Verz.* 32.—Afr. austr.
fragrans, *Eckl. l. c.* 31.—Afr. austr.
Gladiola, *Spreng. ex Steud. Nom.* ed. II. i. 176.—Afr. austr.
hypogaea, *Burch. Trav.* ii. 589.—Afr. austr.
lilacina, *Eckl. Verz.* 31.—Afr. austr.
lineolata, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 347.—Afr. austr.
maculata, *Klatt, l. c.* 349.—Afr. austr.
mucronata, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 234.—Afr. austr.
multiflora, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 351.—Afr. austr.
nana, *Spreng. Syst.* i. 156.—Afr. austr.
nervosa, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 234 = Antholyza nervosa.
obtusifolia, *Ker-Gawl. l. c.*—Ins. Mascar.
ochroleuca, *Sweet, Hort. Brit.* ed. I. 396.—Afr. austr.
parviflora, *Brehm, ex Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 350.—Afr. austr.
plicata, *Ker-Gawl. in Bot. Mag.* t. 576.—Afr. austr.
punctata, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 348.—Afr. austr.
punicea, *Eckl. Verz.* 32.—Afr. austr.
purpurea, Ker-Gawl. in Bot. Mag. sub t. 1019 = villosa.
pygmaea, *Spreng. ex Steud. Nom.* ed. II. i. 176.—Afr. austr.
pygmaea, Baker, in Journ. Linn. Soc. xvi. (1878) 165 = nana.
quadripartita, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 350.—Afr. austr.
reflexa, *Eckl. Verz.* 33.—Afr. austr.
reflexa, Ker-Gawl. Gen. Irid. 154 = plicata.
ringens, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 233.—Afr. austr.
rosea, *Eckl. Verz.* 32.—Afr. austr.
rubro-coerulea, Pritz. Ind. Ic. 134 = seq.
rubrocyanea, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 234.—Afr. austr.
sambucina, *Ker-Gawl. l. c.*—Afr. austr.
scabra, *Brehm, ex Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 349.—Afr. austr.
secunda, *Ker-Gawl. Gen. Irid.* 154.—Afr. austr.
socotrana, *Hook. f. Bot. Mag.* t. 6585.—Ins. Socotr.
spathacea, *Ker-Gawl. in Bot. Mag.* t. 638.—Afr. austr.
Sprengelii, Baker, in Journ. Linn. Soc. xvi. (1878) 165 = pygmaea.
stellata, *Eckl. Verz.* 32.—Afr. austr.
strieta, *Ker-Gawl. in Bot. Mag.* t. 621, 637.—Afr. austr.
sulphurea, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 234.—Afr. austr.
tenuiflora, *Sweet, Hort. Brit.* ed. I. 396, nomen.—Afr. austr.
Thunbergii, *Ker-Gawl. in Kon. & Sims, Ann. Bot.* i. (1805) 233.—Afr. austr.
tubata, *Sweet, Hort. Brit.* ed. I. 396; *Eckl. Verz.* 33.—Afr. austr.
tubiflora, *Ker-Gawl. in Bot. Mag.* t. 630.—Afr. austr.
tubulosa, Ker-Gawl. Gen. Irid. 154 = tubiflora.
undulato-venosa, *Klatt, in Abh. Naturf. Ges. Halle*, xii. (1882) 350.—Afr. austr.

BABIANA :—
villosa, *Ker-Gawl. in Bot. Mag.* t. 583.—Afr. austr.
villosula, *Ker-Gawl. ex Steud. Nom.* ed. II. i. 176.—Afr. austr.

BABINGTONIA, Lindl. in Bot. Reg. (1842) t. 10 = **Baeckea**, Linn. (Myrtac.).
Camphorosmae, Lindl. l. c. = Baeckea Camphorosmae.
corynophylla, F. Muell. Fragm. iv. 74 = Baeckea corynophylla.
crispiflora, F. Muell. l. c. = Baeckea crispiflora.
densifolia, F. Muell. l. c. = Baeckea densifolia.
gracilis, F. Muell. l. c. = Baeckea polyandra.
novo-anglica, F. Muell. l. c. = Baeckea densifolia.
ovalifolia, F. Muell. l. c. = Baeckea ovalifolia.
pentagonantha, F. Muell. l. c. = Baeckea pentagonantha.
pentandra, F. Muell. l. c. = Baeckea pentandra.
Preissiana, F. Muell. l. c. = Baeckea pentandra.
robusta, F. Muell. l. c. = Baeckea robusta.
subcuneata, F. Muell. l. c. = Baeckea subcuneata.
virgata, F. Muell. l. c. = Baeckea virgata.

BABIRON, Rafin. New Fl. Am. iv. 23 (1836) = **Daucus**, Tourn. (Umbell.), &c.
dichotomum, Rafin. l. c. 24 = Apium divaricatum.
divaricatum, Rafin. l. c. = Apium divaricatum.
pusillum, Rafin. l. c. 23 = D. pusillus.

BACA, Rafin. Princ. Somiol. 25 (1814) = **Boea**, Commers. (Gesnerac.).

BACASIA, Ruiz & Pav. Prod. Fl. Per. 105. t. 22 (1794) = **Barnadesia**, Mutis (Compos.).
corymbosa, Ruiz & Pav. Syst. Veg. 189 = Barn. corymbosa.
lanceolata, Ruiz & Pav. ex DC. Prod. vii. 3 = Barn. lanceolata.
spinosa, Ruiz & Pav. Syst. Veg. 188 = Chuquiraga spinosa.

BACCAUREA, Lour. Fl. Cochinch. 651 (1790).
EUPHORBIACEAE, Benth. & Hook. f. iii. 283.
ADENOCREPIS, Blume, Bijdr. 579 (1825).
CALYPTROON, Miq. Fl. Ind. Bat. Suppl. 471 (1860).
COCCOMELIA, Reinw. in Syll. Ratisb. ii. (1828) 5.
HEDYCARPUS, Jack, in Trans. Linn. Soc. xiv. (1825) 118.
MICROSEPALA, Miq. Fl. Ind. Bat. Suppl. 444 (1860).
PIERARDIA, Roxb. Hort. Beng. 28 (1814); Fl. Ind. ii. 254 (1832).
acuminata, *Muell. Arg. in DC. Prod.* xv. II. 463.—Sumatra.
affinis, Muell. Arg. l. c. 459 = parviflora.
Barteri, *Muell. Arg. l. c.* 464.—Afr. trop. occ.
borneensis, *Muell. Arg. l. c.* 460.—Borneo.
bracteata, *Muell. Arg. l. c.* 466.—Malaya.
cauliflora, *Lour. Fl. Cochinch.* 661.—Cochinch.
costulata, *Muell. Arg. in DC. Prod.* xv. II. 464.—Sumatra.
courtallensis, *Muell. Arg. l. c.* 459.—Ind. or.
dasystachya, *Muell. Arg. l. c.* 458.—Java.
deflexa, *Muell. Arg. l. c.* 462.—Java.
dulcis, *Muell. Arg. l. c.* 460.—Sumatra.
flaccida, *Muell. Arg. l. c.* 459.—Burma.
Griffoniana, *Muell. Arg. l. c.* 464.—Afr. occ.
javanica, *Muell. Arg. l. c.* 465.—Java.
lanceolata, *Muell. Arg. l. c.* 457.—Malaya.
macrocarpa, *Muell. Arg. l. c.* 459.—Malaya.
macrophylla, *Muell. Arg. l. c.* 464.—Malaya.
macrostachya, Wight, Ic. Pl. t. 1913 = sapida.
minutiflora, *Muell. Arg. in DC. Prod.* xv. II. 463.—Java.
Motleyana, *Muell. Arg. l. c.* 461.—Malaya.
parviflora, *Muell. Arg. l. c.* 462.—Burma; Malaya.
propinqua, Muell. Arg. l. c. 463 = sapida.
pubera, *Muell. Arg. l. c.* 458.—Sumatra.
pyrrhadasya, *Muell. Arg. l. c.* 462.—Sumatra.
racemosa, *Muell. Arg. l. c.* 461.—Java.
ramiflora, *Lour. Fl. Cochinch.* 661.—Cochinch.
sapida, Bedd. Fl. Sylv. t. 280 = courtallensis.
sapida, *Muell. Arg. in DC. Prod.* xv. II. 459.—Ind. or.; Malaya.
Seemanni, *Muell. Arg. l. c.* 462.—Ins. Pacif.
stylaris, *Muell. Arg. l. c.* 465.—Ins. Pacif.
sumatrana, *Muell. Arg. l. c.* 466.—Sumatra.
sylvestris, *Lour. Fl. Cochinch.* 662.—Cochinch.
taitensis, *Muell. Arg. in DC. Prod.* xv. II. 463.—Ins. Pacif.
tetrandra, *Muell. Arg. l. c.* 465.—Ins. Philipp.
Wilkesiana, *Muell. Arg. l. c.* 461.—Ins. Pacif.

BACCHARIOIDES, Linn. Fl. Zeyl. 196 (1747) = seq.

BACCHARIS, Linn. Hort. Cliff. 404 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 286.
ACHYROBACCHARIS, Sch. Bip. in Walp. Rep. ii. 952 (1843).
ARRHENACHNE, Cass. in Dict. Sc. Nat. lii. 253 (1828).
MOLINA, Ruiz & Pav. Prod. Fl. Per. 111. t. 24 (1794).
PINGRAEA, Cass. in Dict. Sc. Nat. xli. 57 (1826).
POLYPAPPUS, Less. in Linnaea, iv. (1829) 314.
PTEROCLADIS, Lamb. ex G. Don, in Sweet, Hort. Brit. ed. III. 349 (1839).
SERGILUS, Gaertn. Fruct. ii. 409. t. 174 (1791).
STEPHANANTHUS, Lehm. Ind. Sem. Hort. Hamb. (1826).—Cf. Flora, x. (1827) 96.
TURSENIA, Cass. in Dict. Sc. Nat. xxxvii. 480 (1825).
absinthioides, Hook. & Arn. Bot. Beech. Voy. 57 = Tessaria absinthioides.
acerosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 67.—Ecuador.
acutiloba, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 87 = incisa.
adnata, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 870 = Pluchea subdecurrens.
adscendens, *Pers. Syn.* ii. 424.—Peruv.
aegyptiaca, Forsk. ex DC. Prod. v. 450 = Pluchea Dioscoridis.
affinis, DC. l. c. 413 = subopposita.
affinis, Spreng. ex Baker, in Mart. Fl. Bras. vi. III. 94 = retusa.
Alamanni, DC. Prod. v. 402 = glutinosa.
alata, Hort. Berol. ex Loud. Hort. Brit. Suppl. ii. 612 = genistelloides.
alaternoides, *H. B. & K. Nov. Gen. et Sp.* iv. 51.—Peruv.
alaternoides, Poepp. ex DC. Prod. v. 410 = Poeppigiana.
albida, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 41.—Reg. Argent.
alnifolia, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 264.—Peruv.
alpestris, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 122.—Bras.
alpestris, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 64 = rufescens.
alpina, *H. B. & K. Nov. Gen. et Sp.* iv. 48.—Peruv.
amplexicaulis, Wall. ex DC. Prod. v. 439 = Laggera flava.
amygdalina, Griseb. in Goett. Abh. xix. (1874) 175 = oxyodonta.
angulata, *Griseb. l. c.* xxiv. (1879) 180.—Reg. Argent.
angustifolia, Desf. Cat. Hort. Par. ed. III. 163 = B. Pingraea.
angustifolia, *Michx. Fl. Bor. Am.* ii. 125.—Am. bor.
anomala, *DC. Prod.* v. 403.—Bras.
aphylla, *DC. l. c.* 424.—Bras.
apiculata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 74 = trinervis.
arborea, *Linn. Mant.* 284.—Ins. Johann. (Afr. or.).
arborescens, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 412 = arborea.
arbutifolia, *Vahl, Symb. Bot.* iii. 97.—Peruv.
arctostaphyloides, *Baker, in Mart. Fl. Bras.* vi. III. 88.—Bras.
arenaria, *Baker, l. c.* 62.—Bras.
aretioides, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
aretioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 67.—Ecuador.
arguta, Gill. ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 35 = racemosa.
arguta, Pers. Syn. ii. 423 = Psiadia trinervia.
artemisioides, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 41.—Reg. Argent.
articulata, *Pers. Syn.* ii. 425.—Bras.
arunca, Schrank, ex DC. Prod. v. 381 = Conyza Gouani.
asperifolia, Benth. Fl. Hartw. 86 = Conyza asperifolia.
asperula, *Schau. in Linnaea*, xix. (1847) 724.—Mexic.
assuensis, *H. B. & K. Nov. Gen. et Sp.* iv. 67.—Ecuador.
asteroides, Colla, in Mem. Acc. Torin. xxxviii. (1835) 14. t. 25. f. 1 (= *Tripolium conspicuum*?).—Chili.
attenuata, *D. Don, ex Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 34.—Uruguay.
auriculata, Wall. ex DC. Prod. v. 442 = Blumea hieracifolia.
avicenniaefolia, *DC. l. c.* 414.—Bras.

BACCHARIS :—
axillaris, *DC. Prod.* v. 407.—Bras.
axillaris, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 94 = halimimorpha.
bahiensis, *Baker, l. c.* 88.—Bahia.
Bairdii, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 30.—Uruguay.
Baldwinii, Griseb. in Goett. Abh. xxiv. (1879) 181 = tridentata.
Baldwinii, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 28 = rufescens.
balsamifera, *Benth. Pl. Hartw.* 202.—N. Granat.
balsamifera, Stokes, Bot. Mat. Med. iv. 204 = Blumea balsamifera.
banksiaefolia, Bert. in Merc. Chil. (1829).—Cf. Bull. Férussac, xx. (1830) 108 = Tessaria absinthioides.
berberidifolia, Walp. in Linnaea, xiv. (1840) 314 = brachylaenoides.
berberifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 67.—Ecuador.
Bezanilleana, *Remy, in C. Gay, Fl. Chil.* iv. 98.—Chili.
bifrons, *Baker, in Mart. Fl. Bras.* vi. III. 54.—Bras.
Bigelovii, *A. Gray, in Torr. Bot. Mex. Bound.* 84.—Am. bor.; Mexic.
Blanchetiana, DC. Prod. vii. 281 = Moquinia flavescens.
bogotensis, *H. B. & K. Nov. Gen. et Sp.* iv. 61.—N. Granat.
brachylaenoides, *DC. Prod.* v. 421.—Bras.
brachyphylla, *A. Gray, Pl. Wright.* ii. 83.—Am. bor. occ.
bracteata, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 35 = dracunculifolia.
bracteolata, *DC. Prod.* v. 418.—Hab.?
brasiliana, Linn. Sp. Pl. ed. II. 1205 = Vernonia scabra.
brevifolia, *DC. Prod.* v. 409.—Bras.
breviseta, DC. l. c. 402 = serrulata.
bryoides, *Pers. Syn.* ii. 425.—Reg. Magellan.
buddleoides, *H. B. & K. Nov. Gen. et Sp.* iv. 64.—Peruv.
bupleuroides, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 86 = ligustrina.
Burchellii, *Baker, in Mart. Fl. Bras.* vi. III. 44.—Bras.
buxifolia, *Pers. Syn.* ii. 424.—Peruv.
cacalioides, *Spreng. Nachr. i. Bot. Gart. Halle*, 14.—Hab.?
caerulescens, DC. Prod. v. 402 = glutinosa.
caespitosa, *Pers. Syn.* ii. 425.—Peruv.
calistemioides, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 265.—Chili.
calliprinos, *Griseb. in Goett. Abh.* xix. (1874) 177.—Reg. Argent.
calvescens, *DC. Prod.* v. 413.—Bras.
camporum, *DC. l. c.* 399.—Bras.
Candolleana, *Steud. Nom.* ed. II. i. 177.—Peruv.
canescens, *Spreng. Syst.* iii. 466.—Reg. Argent.
capitata, *H. B. & K. Nov. Gen. et Sp.* iv. 53.—Am. austr.
caprariaefolia, *DC. Prod.* v. 416.—Bras.
cassinefolia, *DC. l. c.* 412.—Bras.
cassinoides, DC. l. c. = cassinefolia.
cataphracta, *Spreng. Syst.* iii. 462.—Peruv.
Cephalotes, *DC. Prod.* v. 421.—Bras.
Chilco, *H. B. & K. Nov. Gen. et Sp.* iv. 56.—N. Granat.
chilensis, Weinm. in Syll. Ratisb. ii. (1828) 21 = Podanthus ovatifolius.
chiloënsis, Sch. Bip. in Bonplandia, iv. (1856) 55 = B. Chilquilla.
Chilquilla, *DC. Prod.* v. 419.—Chili.
chinensis, Lour. Fl. Cochinch. ii. 494 = Blumea chinensis.
ciliata, *Gard. in Hook. Lond. Journ. Bot.* iv. (1845) 122.—Bras.
cinerea, *DC. Prod.* v. 400.—Bras.
cinnamomifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 65.—Ecuador.
Clausseni, *Baker, in Mart. Fl. Bras.* vi. III. 99.—Bras.
Clausseni, Sch. Bip. ex Baker, l. c. 82 = cassinefolia.
cochinchinensis, *Spreng. Syst.* iii. 466.—Cochinch.
cognata, *DC. Prod.* v. 413.—Bras.
collina, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 75 = Lundii.
concava, *Pers. Syn.* ii. 425.—Peruv.; Chili.
conferta, *H. B. & K. Nov. Gen. et Sp.* iv. 55.—Mexic.
confertifolia, Colla, in Mem. Acc. Torin. xxxviii. (1835) 15. t. 25. f. 2 = B. Chilquilla.

BACCHARIS :—

congesta, *DC. Prod.* v. 410.—Mexic.
consanguinea, DC. l. c. = pilularis.
conyzoides, *DC. l. c.* 403.—Bras.
coridifolia, *DC. l. c.* 422.—Bras.
Corydalis, DC. ex Sch. Bip. in Linnaea, xxx. (1859–60) 181 = cassinefolia.
corymbosa, *Meyen, Reise*, i. 311.—Chili.
corymbosa, *Pers. Syn.* ii. 424.—Peruv.
crenulata, *Spreng. Syst.* iii. 465.—Bras.
crispa, Spreng. l. c. 466 = genistelloides.
cultrata, *Baker, in Mart. Fl. Bras.* vi. III. 93.—Bras.
cuneata, DC. Prod. v. 408 = conferta.
cuneifolia, DC. l. c. 406 = magellanica.
cuneifolia, Moench, Meth. 574 = halimifolia.
cuneifolia, Wendl. ex Steud. Nom. ed. II. i. 177, 179 = Wendlandii.
curvifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 87 = rufescens.
cupressiformis, Pers. Syn. ii. 425 = Lepidophyllum cupressiforme.
cylindrica, *DC. Prod.* v. 426 = genistelloides.
daphnoides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 34 = cassinefolia.
Darwinii, *Hook. & Arn. l. c.* 39.—Patagon.
deltoidea, *Baker, in Mart. Fl. Bras.* vi. III. 90.—Reg. Argent.
densiflora, *Wedd. Chlor. And.* i. 175.—Bolivia.
denticulata, *D. Don, in DC. Prod.* vii. 282.—Patagon.
dentifolia, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 66 = recurvata.
dentosa, Mart. ex Baker, l. c. 91 = subopposita.
depauperata, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 120 = serrulata.
dependens, *Pers. Syn.* ii. 424.—Peruv.
dioica, *Vahl, Symb. Bot.* iii. 98.—Ins. Carib.
Dioscoridis, Bové, ex DC. Prod. v. 382 = Conyza aegyptiaca.
Dioscoridis, Linn. Cent. Pl. i. 27; Amoen. Acad. iv. 289 = Pluchea hirsuta.
diptera, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 38 = articulata.
discolor, *Baker, l. c.* 48.—Bras.
discolor, Sch. Bip. ex Baker, l. c. 54 = B. Lychnophora.
disticha, Sch. Bip. ex Baker, l. c. 53 = calvescens.
divergens, *DC. Prod.* v. 400.—Mexic.
diversifolia, Spreng. ex Sch. Bip. in Walp. Rep. ii. 958 = Erigeron Sprengelii.
Doniana, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 26.—Bras.
Douglasii, *DC. Prod.* v. 400.—Calif.
dracunculifolia, *DC. l. c.* 421.—Bras.
effusa, *Griseb. in Goett. Abh.* xix. (1874) 177.—Reg. Argent.
elaeagnoides, *Steud. ex Sch. Bip. in Linnaea*, xxx. (1859–60) 181.—Bras.
elacoides, *Remy, in C. Gay, Fl. Chil.* iv. 88.—Chili.
elegans, *H. B. & K. Nov. Gen. et Sp.* iv. 60. t. 324.—Mexic.
elliptica, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 83.—Bras.
emarginata, *Pers. Syn.* ii. 424.—Peruv.
Emoryi, *A. Gray, in Torr. Bot. Mex. Bound.* 83.—Am. bor. occ.
ericoides, Pers. Syn. ii. 425 = Liabum ericoides.
erigeroides, *DC. Prod.* v. 418.—Bras.
crioclada, *DC. l. c.* 415.—Bras.
erioptera, Benth. in Ann. Nat. Hist. ii. (1839) 441 = Pterocaulon virgatum.
eupatorioides, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 22.—Chili.
farinosa, Spreng. in Ersch & Grub. Allgem. Encyc. vii. (1821) 27 = glutinosa.
fasciculosa, *Klatt, in Leopoldina*, xx. (1884) 91.—Mexic.
fastigiata, *Baker, in Mart. Fl. Bras.* vi. III. 39.—Bras.
ferruginea, *Pers. Syn.* ii. 425.—Peruv.
Fevillei, *DC. Prod.* v. 403.—Peruv.
filifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 69 = polyphylla.
flabellata, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 28.—Chili.
flexuosa, *Baker, in Mart. Fl. Bras.* vi. III. 83.—Bras.
flocculosa, Mart. ex Baker, l. c. 53 = calvescens.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iv. 64.—N. Granat.
floribunda, *Phil. in Linnaea*, xxxiii. (1864–65) 145.—Chili.

BACCHARIS :—

foetida, Linn. Sp. Pl. 861 = Pluchea camphorata.
foliosa, *Gill. ex Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 30.—Chili.
frigida, *H. B. & K. Nov. Gen. et Sp.* iv. 49.—Peruv.
fuchsiaefolia, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 88 = orgyalis.
fuliginea, H. B. & K. Nov. Gen. et Sp. iv. 68 = Bigelovia fuliginea.
fusca, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 67.—Ecuador.
Gaudichaudiana, DC. Prod. v. 424 = articulata.
genistelloides, *Pers. Syn.* ii. 425.—Peruv.
genistelloides, Poepp. ex DC. Prod. v. 425 = sagittalis.
genistifolia, *DC. l. c.* 423.—Bras.
Gibertii, *Baker, in Mart. Fl. Bras.* vi. III. 52.—Bras.
Gillesii, *A. Gray, in Proc. Am. Acad.* v. (1861) 123.—Chili.
glabra, *Spreng. Syst.* iii. 466.—Reg. Argent.
glauca, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 263.—Chili.
Glaziovii, *Baker, in Mart. Fl. Bras.* vi. III. 44.—Bras.
glomeruliflora, *Pers. Syn.* ii. 423.—Am. bor.; Ins. Bermud.
glutinosa, Hook. & Arn. Bot. Beech. Voy. 31 = Douglasii?
glutinosa, *Pers. Syn.* ii. 425.—Am. bor. et austr. occ.
gnaphalioides, *DC. Prod.* v. 415.—Bras.
gnaphalioides, Spreng. Syst. iii. 461 = Lucilia acutifolia.
gnidiifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 61.—Ecuador.
gracilis, *DC. Prod.* v. 423.—Bras.
grandiflora, *H. B. & K. Nov. Gen. et Sp.* iv. 58.—Ecuador.
gratissima, Blume, ex DC. Prod. vi. 447 = Blumea appendiculata.
graveolens, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
grindeliaefolia, *Wedd. Chlor. And.* i. 177.—Peruv.
grisea, *Baker, in Mart. Fl. Bras.* vi. III. 65.—Bras.
Grisebachii, *Hieron. in Bol. Acad. Nac. Córdova*, iv. (1881) 36.—Reg. Argent.
guianensis, *Schomb. Fauna & Flora Guy.* 1135.—Guiana.
Haenkei, DC. Prod. v. 401 = Douglasii.
halimifolia, *Linn. Sp. Pl.* 860.—Am. bor.; Ind. occ.
halimifolia, Moench, Meth. Suppl. 235 = Conyza ivaefolia.
halimimorpha, *DC. Prod.* v. 411.—Bras.
hambatensis, *H. B. & K. Nov. Gen. et Sp.* iv. 62.—Ecuador.
Havardi, *A. Gray, Syn. Fl. N. Am.* i. II. 224.—Texas.
helichrysoides, *DC. Prod.* v. 415.—Bras.
hemiprionodes, *Buek, Ind. Gen. DC. Prod.* i. p. v.—Bras.
heterophylla, *H. B. & K. Nov. Gen. et Sp.* iv. 62.—Mexic.
heterophylla, Spreng. Syst. iii. 464, pro parte = sebastianopolitana.
hieracifolia, Lam. Encyc. i. 347 = Conyza Gouani.
hieraciifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 129.—Mexic.
hirta, *DC. Prod.* v. 405.—Bras.
hirtella, *DC. l. c.* 418.—Mexic.
Hookeriana, DC. Prod. v. 414 = Aplopappus Hookerianus.
humifusa, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 311 = Melalema humifusa.
humifusa, *H. B. & K. Nov. Gen. et Sp.* iv. 48. t. 322.—Ecuador.
humilis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 92.—Bras.
Huydobriana, Remy, in C. Gay, Fl. Chil. iv. 90 = B. Pingraea
hypericifolia, *Baker, in Mart. Fl. Bras.* vi. III. 69.—Bras.
hyssopifolia, E. Mey. ex DC. Prod. v. 323 = Nidorella foliosa.
ilicifolia, Lam. Encyc. i. 345 = Brachylaena dentata.
Ilinissae, *Benth. Pl. Hartw.* 202.—N. Granat.
ilinita, *DC. Prod.* v. 412.—Bras.
inamoena, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 81 = trinervis.
incarum, Wedd. Chlor. And. i. 170 = microphylla.
incisa, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 29.—Reg. Argent.
indica, Linn. Sp. Pl. 861 = Pluchea indica.
intermedia, *DC. Prod.* v. 411.—Chili.

BACCHARIS :—

intermixta, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 84.—Bras.
iresinoides, *H. B. & K. Nov. Gen. et Sp.* iv. 63.—N. Granat.
ivaefolia, Blanco, Fl. Filip. ed. I. 627 = Conyza viscidula.
ivaefolia, Linn. Sp. Pl. 860 = Conyza ivaefolia.
juncea, *Desf. Cat. Hort. Par.* ed. III. 163.—Bras.
junciformis, *DC. Prod.* v. 426.—Bras.
lacrymans, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 94 = retusa.
laevis, *Spreng. Syst.* iii. 466.—Cochinch.
lanceolata, *H. B. & K. Nov. Gen. et Sp.* iv. 63.—Peruv.
lancifolia, *Less. in Linnaea*, ix. (1834) 266.—Mexic.
lanuginosa, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 82 = helichrysoides.
lateralis, *Baker, in Mart. Fl. Bras.* vi. III. 100.—Bras.
latifolia, *Pers. Syn.* ii. 424.—Peruv.
laxa, *Gardn. in Hook. Lond. Journ. Bot* iv. (1845) 121.—Bras.
Lechleri, *Sch. Bip. in Lechl. Berb. Am. Austr.* 53, nomen.—Chili.
ledifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 50.—Peruv.
lepidophylla, DC. Prod. v. 427 = Helichrysum baccharoides.
leptocephala, DC. l. c. 413 = rufescens.
leptophylla, DC. l. c. 423 = rufescens.
leptospermoides, DC. l. c. 421 = dracunculifolia.
Lessingiana, *DC. l. c.* 414.—Bras.
leucanthemifolia, Burm. f. ex DC. l. c. 387 = Conyza pinnatilobata.
leucolepis, Sch. Bip. in Linnaea, xxii. (1849) 570 = helichrysoides.
leucopappa, DC. Prod. v. 415 = helichrysoides.
ligustrina, *DC. l. c.* 421.—Bras.
limnophila, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 60 = ligustrina.
linearifolia, Pers. Syn. ii. 425 = Psiadia trinervia.
linearis, *Pers. l. c.*—Chili.
linearis, Poepp. ex Baker, in Mart. Fl. Bras. vi. III. 59 = serrulata.
lingulata, Kunze, in Linnaea, vi. (1831) 147 = rosmarinifolia.
linifolia, *DC. Prod.* v. 420.—Mexic.
linifolia, Meyen, Reise, i. 311 = rosmarinifolia.
linifolia, *Phil. in Linnaea*, xxxiii. (1864–65) 145.—Chili.
longifolia, *DC. Prod.* v. 402.—Mexic.
longipes, *Kunze, ex DC. l. c.* 401.—Chili.
loxensis, *Benth. Pl. Hartw.* 135.—Peruv.
lucida, Meyen, Reise, i. 460 = Vernonia phylicaeformis.
Lundii, *DC. Prod.* v. 404.—Bras.
Lychnophora, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 85.—Bras.
lycioides, *Remy, in C. Gay, Fl. Chil.* iv. 99.—Chili.
lycopodioides, Pers. Syn. ii. 425 = Eriotrix juniperifolia.
Macraei, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 32.—Chili.
macrantha, *H. B. & K. Nov. Gen. et Sp.* iv. 54.—Ecuador.
macrodonta, *DC. Prod.* v. 416.—Bras.
madagascariensis, Pers. Syn. ii. 423 = Psiadia madagascariensis.
magellanica, *Pers. l. c.* 425.—Reg. Magell.
Mandonii, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 533, nomen.—Bolivia.
marcetiaefolia, *Benth. Pl. Hartw.* 202.—Reg. Granat.
marginalis, *DC. Prod.* v. 402.—Peruv.; Chili.
maritima, *Baker, in Mart. Fl. Bras.* vi. III. 72.—Bras.
maxima, *Baker, l. c.* 80.—Bras.
medullosa, DC. Prod. v. 405 = serrulata.
megacephala, D. Don, ex Baker, in Mart. Fl. Bras. vi. III. 79 = triplinervis.
megapotamica, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 38 = rufescens.
megapotamica, *Spreng. Syst.* iii. 461.—Bras.
melastomaefolia, Hook. & Arn. in Hook. Journ. Bot iii. (1841) 25 = oxyodonta.
mesoneura, DC. Prod. v. 412 = retusa.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 295.—Peruv.
micrantha, *H. B. & K. Nov. Gen. et Sp.* iv. 61.—Mexic.
microcephala, *DC. Prod.* v. 425.—Bras.
microdonta, *DC. l. c.* 416.—Bras.

BACCHARIS:—
microphylla, DC. Prod. v. 406 = Candolleana.
microphylla, *H. B. & K. Nov. Gen. et Sp.* iv. 53.—N. Granat.
micropoda, *Baker, in Mart. Fl. Bras.* vi. III. 88.—Bras.
microptera, *Baker, l. c.* 42.—Bras.
milleflora, DC. Prod. v. 426 = genistelloides.
minutiflora, *Mart. ex Baker, in Mart. Fl. Bras.* vi. III. 69.—Bras.
mollis, *H. B. & K. Nov. Gen. et Sp.* iv. 58.—Ecuador.
monantha, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 69 = minutiflora.
montana, DC. Prod. v. 404 = macrodonta.
montana, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 78 = Schultzii.
montevidensis, Sch. Bip. ex Baker, l. c. 57 = coridifolia.
montevidensis, Spreng. Syst. iii. 460 = Vernonia nitidula.
mucronata, Hook. & Arn. Bot. Beech. Voy. 30 = Aplopappus Hookerianus.
mucronata, *H. B. & K. Nov. Gen. et Sp.* iv. 60.—Mexic.
Muelleri, *Baker, in Mart. Fl. Bras.* vi. III. 61.—Bras.
multiflora, *H. B. & K. Nov. Gen. et Sp.* iv. 59.—Mexic.
multisulcata, *Baker, in Mart. Fl. Bras.* vi. III. 45.—Bras.
myricaefolia, DC. Prod. v. 405 = semiserrata.
myriocephala, *Baker, in Mart. Fl. Bras.* vi. III. 93.—Bras.
myriocephala, DC. Prod. v. 426 = genistelloides.
myrsinites, *Pers. Syn.* ii. 424. n. 25.—Ins. S. Doming.
myrsinoides, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 32 = brevifolia.
myrtilloides, *Griseb. in Goett. Abh.* xix. (1874) 176.—Reg. Argent.
nana, *D. Don, ex Baker, in Mart. Fl. Bras.* vi. III. 56.—Patagon.
Neaei, *DC. Prod.* v. 411.—Am. merid.
neriifolia, Linn. Sp. Pl. 860 = Brachylaena nereifolia.
nervosa, *DC. Prod.* v. 399.—Ind. occ.
nitida, *Pers. Syn.* ii. 425.—Peruv.
nitida, Wall. Cat. n. 3043 = Blumea chinensis.
nivalis, *Sch. Bip. in Bonplandia*, iv. (1856) 55, nomen.—Chili.
notosergila, *Griseb. in Goett. Abh.* xxiv. (1879) 183.—Reg. Argent.
nuda, DC. Prod. v. 424 = aphylla.
oblongifolia, *Pers. Syn.* ii. 424; *Spreng. Syst.* iii. 460.—Chili.
obovata, *DC. Prod.* v. 408.—Peruv.
obovata, Hook. & Arn. Bot. Beech. Voy. 30 = umbelliformis.
obtusifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 51.—Peruv.
ochracea, *Spreng. Syst.* iii. 460.—Reg. Argent.
odorata, *H. B. & K. Nov. Gen. et Sp.* iv. 52.—N. Granat.
odorata, Spreng. in Ersch & Grub. Allgem. Encyc. vii. (1821) 27 = cassinefolia.
oleifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 86 = calvescens.
oligantha, DC. ex Thoms. in Journ. Linn. Soc. ix. (1867) 343 = Catamixis baccharoides.
oligophylla, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 23 = Pohlii.
opuntioides, *Mart. ex Baker, l. c.* 39.—Bras.
Orbignyana, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 327.—Chiquitos.
organensis, *Baker, in Mart. Fl. Bras.* vi. III. 74.—Bras.
orgyalis, *DC. Prod.* v. 416.—Bras.
orizabaĕnsis, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 130.—Mexic.
orizabaĕnsis, Klatt, in Leopoldina, xx. (1884) 91 = resinosa.
oronocensis, *DC. Prod.* v. 422.—Peruv.
ovalis, Pers. Syn. ii. 424 = Pluchea ovalis.
ovata, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 22.—Chili.
ovata, Sieber, ex DC. Prod. v. 450 = Pluchea ovalis.
oxyodonta, *DC. l. c.* 404.—Bras.
oxyphylla, *DC. l. c.* 400.—Peruv.
paniculata, *DC. l. c.* 420.—Chili.
parviflora, *Less. in Linnaea*, v. (1830) 146.—Mexic.
parviflora, Pers. Syn. ii. 425 = marginalis.
parviflora, Poir. Encyc. Suppl. i. 556 = bracteolata.
parvifolia, *DC. Prod.* v. 424.—Peruv.

BACCHARIS:—
patagonica, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 29.—Reg. Magellan.
patens, *Baker, in Mart. Fl. Bras.* vi. III. 52.—Bras.
paucidentata, DC. Prod. v. 420 = rufescens.
paucidentata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 71 = dracunculifolia.
paucifloscula, *DC. Prod.* v. 413.—Bras.
pedalis, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 64 = rufescens.
pedicellata, *DC. Prod.* v. 407.—Chili.
pentaptera, *DC. l. c.* 425.—Bras.
Pentlandii, *DC. l. c.* 416.—Bolivia.
pentziaefolia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 96.—Bras.
petiolata, *DC. Prod.* v. 418.—Chili.
phagnaloides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 59 = microdonta.
philippensis, *H. B. & K. Nov. Gen. et Sp.* iv. 55.—Ecuador.
phlogopappa, *DC. Prod.* v. 402.—Am. austr.
phylicaefolia, *DC. l. c.* 415.—Bras.
phylicaeformis, Meyen, Reise, ii. 31 = Vernonia phylicaeformis.
phylicoides, *H. B. & K. Nov. Gen. et Sp.* iv. 49.—Peruv.
phyteumoides, *DC. Prod.* v. 425.—Bras.
pilularis, *DC. l. c.* 407.—Am. bor. occ.
Pingraea, *DC. Prod.* v. 420.—Chili.
Pingraea, Nutt. in Trans. Am. Phil. Soc. vii. (1841) 337 = glutinosa.
pinnatifida, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 327.—Patagon.
platensis, *Spreng. Syst.* iii. 465.—Reg. Argent.
platypoda, *DC. Prod.* v. 409.—Bras.
Plummerae, *A. Gray, in Proc. Am. Acad.* xv. (1880) 48.—Calif.
Poeppigiana, *DC. Prod.* v. 410.—Chili.
polifolia, *Griseb. in Goett. Abh.* xix. (1874) 176.—Reg. Argent.
polyantha, *H. B. & K. Nov. Gen. et Sp.* iv. 64.—Peruv.?; Mont. Vid.?
polyantha, Spreng. Syst. ii. 464 = helichrysoides.
polycephala, *Wedd. Chlor. And.* i. 173.—Bolivia.
polycephala, Sch. Bip. in Linnaea, xxx. (1859–60) 181 = vernonioides.
polygalaefolia, *H. B. & K. Nov. Gen. et Sp.* iv. 49.—Peruv.
polygona, *Baker, in Mart. Fl. Bras.* vi. III. 46.—Bras.
polylepis, Sch. Bip. ex Baker, l. c. 98 = subcapitata.
polyneura, Sch. Bip. ex Baker, l. c. 100 = reticularia.
polyphylla, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 87.—Bras.
polyphylla, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 69 = minutiflora.
polyptera, *DC. Prod.* v. 426.—Bras.
potosina, *A. Gray, in Proc. Am. Acad.* xv. (1880) 33.—Mexic.
prenanthoides, *Baker, in Mart. Fl. Bras.* vi. III. 84.—Bras.
prinoides, *H. B. & K. Nov. Gen. et Sp.* iv. 56.—Reg. Amazon.
prostrata, Baldw. in Hook. Journ. Bot. iii. (1841) 28 = Baldwinii.
prostrata, *Pers. Syn.* ii. 424.—Peruv.
prunifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 63.—Peruv.
psilocalyx, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 48 = erioclada.
ptarmicaefolia, DC. Prod. v. 419 = Bigelovii?
Pterocaulon, Boj. ex DC. l. c. 455 = Pterocaulon Bojeri.
pteronioides, *DC. l. c.* 410.—N. Mexic.
puberula, *DC. l. c.* 401.—Bras.
pulchella, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 532, nomen.—Bolivia.
pulverulenta, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 327.—Peruv.
punctigera, DC. Prod. v. 404 = Lundii.
punctulata, DC. l. c. 405 = oxyodonta.
pyramidalis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 120.—Bras.
quadrangularis, Meyen, Reise, i. 460 = Lepidophyllum Meyeni.
quadrinervia, *Pers. Syn.* ii. 423.—Am. austr.
quinquenervis, *DC. Prod.* v. 399.—Bras.
quitensis, *H. B. & K. Nov. Gen. et Sp.* iv. 57.—Ecuador.

BACCHARIS:—
racemosa, *DC. Prod.* v. 401.—Chili.
radicans, DC. l. c. 416 = gnaphalioides.
Radin, *Phil. in Linnaea*, xxxiii. (1864–65) 146.—Chili.
ramiflora, *A. Gray, in Proc. Am. Acad.* xv. (1880) 33.—Mexic.
ramosissima, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 84 = retusa.
ramulosa, A. Gray, Pl. Thurb. 301 = pteronioides.
rapunculoides, E. Mey. ex DC. Prod. v. 323 = Nidorella resedifolia.
recurvata, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 86.—Bras.
refracta, *Burch. ex Baker, in Mart. Fl. Bras.* vi. III. 64.—Bras.
Regnellii, *Sch. Bip. in Linnaea*, xxii. (1849) 571.—Bras.
resediflora, Less. ex Baker, in Mart. Fl. Bras. vi. III. 58 = erigeroides.
resiniflua, *Steud. & Hochst. ex DC. Prod.* vii. 282.—Arab.
resinosa, Hook. & Arn. Bot. Beech. Voy. 31 = concava.
resinosa, *H. B. & K. Nov. Gen. et Sp.* iv. 52.—Am. austr.
retamoides, *Phil. Sert. Mend. Alt.* 22.—Chili.
reticularia, *DC. Prod.* v. 409.—Bras.
reticulata, Pers. Syn. ii. 425 = genistelloides.
retusa, *DC. Prod.* v. 412.—Bras.
revoluta, *H. B. & K. Nov. Gen. et Sp.* iv. 50.—Peruv.
rhetinodes, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 265.—Chili.
rhexioides, *H. B. & K. Nov. Gen. et Sp.* iv. 66.—Peruv.; Bras.
rhomboidalis, *Remy, in C. Gay, Fl. Chil.* iv. 99.—Chili.
Riedelii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 82 = cassinefolia.
rigida, Hook. & Arn. Bot. Beech. Voy. 57 = oblongifolia.
riparia, *H. B. & K. Nov. Gen. et Sp.* iv. 65.—Ecuador.
riparia, Poepp. ex DC. Prod. v. 418 = sessilifolia.
rivularis, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 83 = macrodonta.
Rohriana, Spreng. in Ersch & Grub. Allgem. Encyc. vii. (1821) 27 = dioica.
Roraimae, *Schomb. Fauna et Fl. Guy.* 1078, nomen.—Guiana.
rosmarinifolia, *Hook. & Arn. Bot. Beech. Voy.* 30.—Chili.
rotundifolia, *Spreng. Syst.* iii. 465.—Reg. Argent.
rufescens, *Spreng. l. c.* iii. 464.—Reg. Argent.
rufidula, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 81 = vernonioides.
rugosa, Willd. ex DC. Prod. v. 39 = Vernonia patens.
rupicola, *H. B. & K. Nov. Gen. et Sp.* iv. 52.—N. Granat.
sagittalis, *DC. Prod.* v. 425.—Chili.
salicifolia, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 337 = salicina.
salicifolia, *Pers. Syn.* ii. 425.—Peruv.
salicina, Rothr. in Wheeler, Rep. vi. 156 = Emoryi.
salicina, *Torr. & Gray, Fl. N. Am.* ii. 258.—Am. bor. occ.
saligna, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 60 = ligustrina.
saltensis, *Baker, l. c.* 93.—Bras.
Salvia, Lour. Fl. Cochinch. ii. 494 = Blumea balsamifera.
Salzmannii, *DC. Prod.* v. 409.—Bras.
sarophora, Phil. in Anal. Univ. Chil. i. (1861) 52; et in Linnaea, xxxiii. (1864–65) 147 = Heterothalamus spartioides.
sarothroides, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 211.—Calif.
scabra, *Pers. Syn.* ii. 424.—Ecuador.
scandens, *Less. in Linnaea*, v. (1830) 146.—Mexic.
scandens, *Pers. Syn.* ii. 424.—Peruv.
Schiedeana, *Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 83.—Bras.
Schlechtendalii, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 50 = tarchonanthoides.
Schomburgkii, *Baker, l. c.* 89.—Guiana.
Schultzii, *Baker, l. c.* 78.—Bras.
scolopendra, *Hook. Ic. Pl.* i. t. 68.—Ecuador.
scoparia, *Sw. Fl. Ind. Occ.* iii. 1339.—Jamaica.
scoparioides, *Griseb. Cat. Pl. Cub.* 150.—Cuba.
sculpta, *Griseb. in Goett. Abh.* xix. (1874) 175.—Reg. Argent.

BACCHARIS:—
sebastianopolitana, *Baker, in Mart. Fl. Bras.* vi. III. 65.—Bras.
Seemanni, *A. Gray, in Proc. Am. Acad.* xv. (1880) 33.—Mexic.
Selloi, *Baker, in Mart. Fl. Bras.* vi. III. 68.—Bras.
semiserrata, DC. Prod. v. 404. n. 38 = hemiprionodes.
semiserrata, DC. l. c. 419. n. 149 = Sternbergiana.
senegalensis, Pers. Syn. ii. 424 = Vernonia senegalensis.
senicula, Mart. in Flora, xxi. (1838) II. Beibl. 61 = cassinefolia.
sergiloides, *A. Gray, in Pacif. Rail. Rep.* iv. 101.—Am. bor. occ.
serraefolia, *DC. Prod.* v. 403.—Mexic.
serratifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 59.—Mexic.; Ind. occ.
serrulata, *Pers. Syn.* ii. 423.—Bras.
sessiliflora, Michx. Fl. Bor. Am. ii. 135 = glomeruliflora.
sessiliflora, *Vahl, Symb. Bot.* iii. 97.—Bras.
sessilifolia, *DC. Prod.* v. 418.—Chili.
sinensis, *Lour. Fl. Cochinch.* ii. 604.—Cochinch.
sinuata, *H. B. & K. Nov. Gen. et Sp.* iv. 53.—Peruv.
Solierii, *Remy, in C. Gay, Fl. Chil.* iv. 98.—Chili.
Solisi, *Phil. in Anal. Univ. Chil.* (1865) ii. 330.—Chili.
sordescens, *DC. Prod.* v. 405.—Mexic.
spartea, *Benth. Bot. Voy. Sulph.* 114.—Am. centr.
spartioides, C. Gay, Fl. Chil. iv. 102 = Heterothalamus spartioides.
spathulata, *Schau. in Linnaea*, xix. (1847) 724.—Mexic.
spathulata, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 326.—Peruv.
speciosa, *DC. Prod.* v. 399.—Ins. Guadalup.
sphaerocephala, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 25.—Chili.
spiropoda, F. Muell. Fragm. i. 34 = Pluchea baccharoides.
squarrosa, *Baker, in Mart. Fl. Bras.* vi. III. 50.—Bras.
squarrosa, *H. B. & K. Nov. Gen. et Sp.* iv. 67.—Mexic.
Steetzii, *Anderss. Galap. Oarnes Veg.* 177.—Ins. Galap.
stenocephala, *Baker, in Mart. Fl. Bras.* vi. III. 39.—Bras.
stenoptera, Sch. Bip. in Linnaea, xxxiv. (1865–66) 533 = subalata.
Sternbergiana, *Steud. Nom.* ed. II. i. 179.—Peruv.
striata, Ruiz & Pav. Syst. 207 = Fevillei.
Stroebeliana, *Ces. in Atti Soc. Acc. Sc. Nap.* v. (1873) VII. 12.—Europ.
stylosa, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 120.—Bras.
subalata, *Wedd. Chlor. And.* i. 174.—Bolivia.
subcapitata, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 85.—Bras.
subdentata, *DC. Prod.* v. 408.—Bras.
subopposita, *DC. l. c.* 413.—Bras.
subpenninervis, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 532, nomen.—Bolivia.
subscandens, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 76 = conyzoides.
subspathulata, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 87 = rufescens.
subulata, *D. Don, in Hook. Journ. Bot.* iii. (1841) 39.—Patagon.
sulcata, *DC. Prod.* v. 419.—Mexic.
syncephala, Sch. Bip. in Bonplandia, iv. (1856) 54; et ex Baker, in Mart. Fl. Bras. vi. III. 99 = platypoda.
tarchonanthoides, *DC. Prod.* v. 414.—Bras.
teindalensis, *H. B. & K. Nov. Gen. et Sp.* iv. 417.—N. Granat.
tenella, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 42.—Patagon.
tenuifolia, *Linn. Sp. Pl.* 860.—'Afr.'
tenuifolia, DC. Prod. v. 423 = rufescens.
teucrifolia, Hook. & Arn. ex Baker, in Mart. Fl. Bras. vi. III. 77 = anomala.
texana, *A. Gray, Pl. Fendl.* 75.—Texas.
thesioides, *H. B. & K. Nov. Gen. et Sp.* iv. 61.—Mexic.; Arizona.
Thomasii, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 326.—Mexic.
thymifolia, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 38.—Reg. Argent.
thymifolia, Miers, in Proc. Hort. Soc. (1863) 349 = Weirii.
thyoides, *Pers. Syn.* ii. 425.—Peruv.
Tola, *Phil. Fl. Atac.* 30.—Chili.
tomentosa, Steud. Nom. ed. II. i. 179 = Moquinia polymorpha.

BACCHARIS:—
tomentosa, *Pers. Syn.* ii. 424.—Peruv.
trichoclada, *DC. Prod.* v. 400.—Mexic.
trichotoma, *Klatt, in Leopoldina*, xx. (1884) 91.—Mexic.
tricuneata, Pers. Syn. ii. 424 = cuneata.
tridentata, Gaudich. Fl. Mal. 15 = magellanica.
tridentata, Poepp. ex DC. Prod. v. 411 = concava.
tridentata, *Vahl, Symb. Bot.* iii. 98.—Bras.
trifurcata, Trevir. in Nov. Act. Nat. Cur. xiii. (1826) 201 = Conyza stricta.
trimera, DC. Prod. v. 425 = genistelloides.
trinervis, *Pers. Syn.* ii. 423.—Bras.
triplinervia, DC. Prod. v. 404 = oxyodonta.
triplinervis, *Baker, in Mart. Fl. Bras.* vi. III. 79.—Bras.
triptera, Mart. Mat. Med. Bras. 43 = genistelloides.
tripterix, Poepp. ex Baker, in Mart. Fl. Bras. vi. III. 42 = sagittalis.
truncata, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 82.—Bras.
truncata, *Phil. in Linnaea*, xxxiii. (1864) 148.—Chili.
tucumanensis, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 34.—Reg. Argent.
Tweediei, *Hook. & Arn. l. c.* 33 = subopposita.
ulicina, *Hook. & Arn. l. c.* 38.—Patagon.
ulmifolia, *Burm. f. Fl. Cap. Prod.* 26 = Conyza incisa.
umbelliformis, *DC. Prod.* v. 410.—Chili.
uncinella, *DC. l. c.* 415.—Bras.
uniflora, *Pers. Syn.* ii. 425.—Peruv.
Urvilleana, *Brongn. Bot. Voy. Coq.* t. 61.—Reg. Magell.
ustulata, *Benth. Pl. Hartw.* 135.—N. Granat.
vaccinioides, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 121.—Bras.
vaccinoides, *H. B. & K. Nov. Gen. et Sp.* iv. 50.—Mexic.
Vahlii, DC. Prod. v. 411 = dioica.
valdiviana, *Phil. in Linnaea*, xxviii. (1856) 738.—Chili.
varians, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 84 = rufescens.
Vauthieri, *DC. Prod.* v. 409.—Bras.
velutina, DC. l. c. 415 = ochracea.
veneta, H. B. & K. Nov. Gen. et Sp. iv. 68 = Bigelovia veneta.
venosa, *Pers. Syn.* ii. 425.—Peruv.
venulosa, DC. Prod. v. 421 = brachylaenoides.
venusta, *H. B. & K. Nov. Gen. et Sp.* iv. 66.—N. Granat.
verbenaefolia, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 26 = hirta.
vernicosa, *Hook. & Arn. in Hook. l. c.* 28.—Uruguay.
vernonioides, *DC. Prod.* v. 422.—Bras.
villosa, *Vahl, Symb. Bot.* iii. 98.—Arabia.
viminea, *DC. Prod.* v. 400.—Calif.
vincaefolia, *Baker, in Mart. Fl. Bras.* vi. III. 83.—Reg. Argent.
virgata, Boj. ex DC. Prod. v. 385 = Conyza virgata.
viscosa, Hook. & Arn. Bot. Beech. Voy. 352 = Douglasii.
viscosa, Lam. Encyc. i. 345 = Psiadia integerrima.
viscosa, Walt. Fl. Carol. 202 = Pluchea bifrons.
vismioides, DC. Prod. v. 412 = vernonioides.
Volckmanni, *Phil. in Linnaea*, xxxiii. (1864–65) 147.—Chili.
volubilis, *H. B. & K. Nov. Gen. et Sp.* iv. 417.—Peruv.
vulneraria, *Baker, in Mart. Fl. Bras.* vi. III. 75.—Bras.
Walpersiana, D. Dietr. Syn. Pl. iv. 1447 = brachylaenoides.
Weddellii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 63 = rufescens.
Weirii, *Baker, l. c.* 67.—Bras.
Wendlandii, *Steud. Nom.* ed. II. i. 179.—Afr. austr.
Wrightii, *A. Gray, Pl. Wright.* i. 101.—Am. bor. occ.
Wrightii, Sch. Bip. in Seem. Bot. Voy. Herald, 303 = Seemanni.
xalapensis, H. B. & K. Nov. Gen. et Sp. iv. 56 = conferta.
xerophila, Mart. in Flora, xxiv. (1841) II. Beibl. 11 = sessiliflora.
xiphophylla, *Baker, in Mart. Fl. Bras.* vi. III. 68.—Bras.

BACCHAROIDES, Moench, Meth. 578 (1794) = **Vernonia**, Schreb. (Compos.).
anthelmintica, Moench, l. c. = V. anthelmintica.

BACKHOUSIA, Hook. & Harv. in Bot. Mag. t. 4133 (1845). *MYRTACEAE*, Benth. & Hook. f. i. 711.
angustifolia, *F. Muell. Fragm.* i. 79.—Austral.

BACKHOUSIA:—
citriodora, *F. Muell. Fragm.* i. 78.—Austral.
myrtifolia, *Hook. & Harv. in Bot. Mag.* t. 4133.—Austral.
riparia, Hook. & Harv. l. c. sub t. 4133 = myrtifolia.
sciadophora, *F. Muell. Fragm.* ii. 26, 171.—Austral.

BACONIA, DC. in Ann. Mus. Par. ix. (1807) 219 = **Pavetta**, Linn. (Rubiac.).
corymbosa, DC. l. c. = P. Baconia.
montana, Hook. f. in Journ. Linn. Soc. vii. (1864) 196 = P. Hookeriana.

BACOPA, Aubl. Pl. Gui. i. 128. t. 49 (1775). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 952.
aquatica, *Aubl. l. c.* 129. t. 49.—Guiana.
grandiflora, Mart. ex Benth. in Ann. Nat. Hist. ii. (1839) 443 = aquatica.

BACTRIS, Jacq. Stirp. Am. 271. t. 256 (1763). *PALMAE*, Benth. & Hook. f. iii. 941.
AUGUSTINEA, Karst. in Linnaea, xxvii. (1856) 395.
GUILIELMA, Mart. Palm. Fam. 21 (1824); Hist. Nat. Palm. ii. 81. tt. 66, 67 (1824).
PYRENOGLYPHIS, Karst. Fl. Columb. ii. 141. t. 174 (1869).
acanthocarpa, *Mart. Hist. Nat. Palm.* ii. 92. t. 70, 71. f. 2.—Bras.
acanthocarpoides, Barb.-Rodr. Enum. Palm. Nov. 31 = acanthocarpa.
acanthocnemis, Mart. in Palmet. Orb. 67 = simplicifrons.
acanthophylla, *Mart. l. c.* 70.—Ins. Porto Rico.
acanthospatha, *Trail, ex Drude, in Mart. Fl. Bras.* iii. II. 354.—Bras.
actinoneura, *Drude & Trail, ex Drude, in Mart. l. c.* 344.—Bras.
aculeifera, *Drude, l. c.* 352 in obs.—Bras.
acuminata, *Liebm. ex Mart. Hist. Nat. Palm.* iii. 321. Mexic.
arenaria, Barb.-Rodr. Enum. Palm. Nov. 29 = simplicifrons.
aristata, *Mart. Hist. Nat. Palm.* ii. 97. t. 73 a. f. 5.—Bras.
armata, Barb.-Rodr. Enum. Palm. Nov. 27 = chaetospatha.
arundinacea, *Trail, in Journ. Bot.* xv. (1877) 10.—Reg. Amazon.
Aubletiana, *Trail, l. c.* xiv. (1876) 372.—Guiana.
baculifera, *Karw. ex Mart. Hist. Nat. Palm.* iii. 322.—Mexic.
balanoides, *H. Wendl. in Kerch. Palm.* 233.—Costa Rica.
balanophora, *Spruce, in Journ. Linn. Soc.* xi. (1871) 146, 153.—Am. trop.
bicuspidata, *Spruce, l. c.* 146, 152.—Am. trop.
bidentula, *Spruce, l. c.* 146, 151.—Am. trop.
bifida, *Mart. Hist. Nat. Palm.* ii. 105. t. 73 c.—Bras.
bifida, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 44 = Oerstediana.
brevifolia, Spruce, in Journ. Linn. Soc. xi. (1871) 144, 147 = simplicifrons.
Brongniartii, *Mart. in Palmet. Orb.* 59. t. 7. f. 2.—Bolivia.
campestris, *Poepp. in Mart. Hist. Nat. Palm.* ii. 146.—Bras.
capillacea, *Trail, in Journ. Bot.* xv. (1877) 5.—Bras.
caracasana, *Lodd. ex Linden, Illustr. Hortic.* xxviii. (1881) 16.—Venezuela.
Caravallana, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Bras.
caribaea, *Karst. in Linnaea*, xxviii. (1856) 403.—Ind. occ.
carolensis, Spruce, in Journ. Linn. Soc. xi. (1871) 145, 149 = simplicifrons.
caryotaefolia, *Mart. Hist. Nat. Palm.* ii. 106. t. 74. f. 3, 4.—Bras.
caryotifolia, *Drude, in Mart. Fl. Bras.* iii. II. 337.—Bras.
Catel, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Venezuela.
chactophylla, *Mart. in Palmet. Orb.* 71.—Bras.
chactorhachis, Mart. l. c. 61 = major.
chaetospatha, *Mart. Hist. Nat. Palm.* ii. 147.—Bras.
chloracantha, *Poepp. in Mart. l. c.* 145.—Bras.
ciliata, *Mart. l. c.* 95. t. 71. f. 3.—Peruv.
concinna, *Mart. l. c.* 99. t. 72. f. 4, 5, 6.—Bras.
confluens, Linden & H. Wendl. in Linnaea, xxviii. (1856) 347 = acanthospatha.

BACTRIS:—
Constanciae, *Barb.-Rodr. Enum. Palm. Nov.* 37.—Bras.
Corossilla, *Karst. in Linnaea*, xxviii. (1856) 407.—Venezuela.
Corazillo, Hort. ex H. Wendl. Cycl. Pand. Cycad. 5 = praec.?
Cruegeriana, *Griseb. & Wendl. ex Griseb. Fl. Brit. W. Ind.* 520.—Ind. occ.
cucullata, *H. Wendl. in Kerch. Palm.* 233, nomen.—Hab.?
Cuesco, *Crueg. ex Griseb. Fl. Brit. W. Ind.* 520.—Ind. occ.
Cuesco, *Engl. in Linnaea*, xxxiii. (1864–65) 665.—N. Granat.
Curuena, Trail, in Journ. Bot. xv. (1877) 48 = socialis.
cuspidata, *Mart. Hist. Nat. Palm.* ii. 101. t. 73 B.—Bras.
Cuvaro, *Karst, in Linnaea*, xxviii. (1856) 406.—Venezuela.
cyagroides, Barb.-Rodr. Enum. Palm. Nov. 33 = syagroides.
elatior, *Wallace, Palm Trees Amaz.* 81. t. xxx.—Bras.
elegans, *Hort. ex H. Wendl. in Kerch. Palm.* 233.—Hab.?
elegans, *Trail, in Journ. Bot.* xv. (1877) 3.—Bras.
ericetina, Barb.-Rodr. Enum. Palm. Nov. 26 = bifida.
erosa, *Mart. Hist. Nat. Palm.* ii. 281.—Ind. occ.
eumorpha, *Trail, in Journ. Bot.* xv. (1877) 9. t. 184.—Bras.
exaltata, Barb.-Rodr. Enum. Palm. Nov. 32 = major?
faucium, *Mart. in Palmet. Orb.* 60. t. 6. f. 2.—Bolivia.
fissifrons, *Mart. Hist. Nat. Palm.* ii. 103. t. 73. b. f. 3, 4.—Bras.
flavispina, Hort. Belg. ex Heynh. Nom. ii. 57 = pallidispina.
floccosa, Spruce, in Journ. Linn. Soc. xi. (1871) 146, 151 = cuspidata.
fusca, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 43.—Costa Rica.
Gasipaës, *H. B. & K. Nov. Gen. et Sp.* i. 302.—N. Granat.
Gaviona, Trail, in Journ. Bot. xv. (1877) 48 = major.
geonomoides, *Drude, in Mart. Fl. Bras.* iii. II. 325.—Bras.
glandulosa, *Oerst. L'Amér. Centr.* 15. t. 8. f. 1–15.—Bras.
glaucescens, *Drude, in Mart. Fl. Bras.* iii. II. 345.—Bras.
Glazioviana, *Drude, l. c.* 348.—Bras.
globosa, Gaertn. Fruct. i. 22 = Acrocomia sclerocarpa.
gracilis, Barb.-Rodr. Enum. Palm. Nov. 27 = simplicifrons.
granariuscarpa, *Barb.-Rodr. l. c.* 37.—Bras.
granatensis, *H. Wendl. ex Linden, Illustr. Hortic.* xxviii. (1881) 16.—N. Granat.
guineensis, Steud. Nom. ed. I. 98, 99 = minor.
hirta, *Mart. Hist. Nat. Palm.* ii. 105. t. 73. C. f. 3.—Bras.
horrida, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 41.—Nicaragua.
hylophila, Spruce, in Journ. Linn. Soc. xi. (1871) 146, 152 = pectinata.
incommoda, *Trail, in Journ. Bot.* xv. (1877) 42.—Bras.
inermis, *Trail, ex Barb.-Rodr. Enum. Palm. Nov.* 30.—Bras.
infesta, *Mart. in Palmet. Orb.* 54. t. 7. f. 1.—Bolivia.
integrifolia, Wallace, Palm Trees Amaz. 91. t. xxxv = fissifrons.
interruptepinnata, *Barb.-Rodr. Enum. Palm. Nov.* 37.—Bras.
inundata, *Mart. in Palmet. Orb.* 58. t. 7. f. 3.—Bras.
juruënsis, *Trail, in Journ. Bot.* xv. (1877) 40.—Bras.
leucantha, *H. Wendl. in Linnaea*, xxviii. (1856) 345.—N. Granat.
Liboniana, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Bras.
linearifolia, Barb.-Rodr. Enum. Palm. Nov. 31 = pectinata.
littoralis, *Barb.-Rodr. l. c.* 36.—Bras.
longifrons, *Mart. Hist. Nat. Palm.* ii. 106.—Reg. Amazon.
longipes, *Poepp. in Mart. l. c.* 145. t. 73.—Bras.
Macanilla, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Venezuela.

BACTRIS:—
macracantha, *Mart. Hist. Nat. Palm.* ii. 95. t. 73.—Bras.
macroacantha, *Mart. l. c.*—Bras.
macrocarpa, *Wallace, Palm Trees Amaz.* 85. t. xxxii.—Bras.
major, *Jacq. Stirp. Am.* 280. t. 171. f. 2.—N. Granat.
Maraja, *Mart. Hist. Nat. Palm.* ii. 93. t. 71. f. 1.—Bras.
Maraja-açu, *Barb.-Rodr. Enum. Palm. Nov.* 36.—Bras.
Maraja-y, Barb.-Rodr. l. c. ed. I. 29 = cuspidata.
martineziaefolia, Hort. ex H. Wendl. in Kerch. Palm. 234 = Martinezia corallina.
mexicana, *Mart. in Palmet. Orb.* 65.—Mexic.
microcarpa, Spruce, in Journ. Linn. Soc. xi. (1871) 146, 153 = hylophila.
microspatha, Barb.-Rodr. Enum. Palm. Nov. 26 = cuspidata.
minax, *Miq. in Verh. Nat. Wet. Haarl.* vii. (1851) 207.—Guiana.
minima, Gaertn. Fruct. ii. 269. t. 139. f. 5 = Martinezia corallina.
minor, Gaertn. l. c. i. t. 9. f. 1 = Acrocomia lasiospatha?
minor, *Jacq. Stirp. Am.* 279. t. 171. f. 1.— Bras.; N. Granat.
mitis, Mart. Hist. Nat. Palm. ii. 102 = cuspidata.
monticola, *Barb.-Rodr. Enum. Palm. Nov.* 34.—Bras.
negrensis, Spruce, in Journ. Linn. Soc. xi. (1871) 145, 147 = simplicifrons.
nemorosa, *Barb.-Rodr. Enum. Palm. Nov.* 32.—Bras.
Neromanni, *Voigt, in Syll. Ratisb.* ii. (1828) 51.—Hab.?
obovata, *H. Wendl. in Kerch. Palm.* 234.—Hab.?
Oerstediana, *Trail, in Journ. Bot.* xv. (1877) 43.—Costa Rica.
oligocarpa, *Barb.-Rodr. Enum. Palm. Nov.* 28.—Bras.
ovata, *H. Wendl. in Kerch. Palm.* 234.—Nicaragua.
ovata, Stokes, Bot. Mat. Med. iv. 394 = major.
pallidispina, *Mart. in Palmet. Orb.* 62.—Guiana.
palustris, Barb.-Rodr. Enum. Palm. Nov. 36 = bidentula.
paraënsis, *Splitg. ex H. Wendl. in Kerch. Palm.* 234.—Surinam.
paucijuga, *Barb.-Rodr. Enum. Palm. Nov.* 34.—Bras.
Pavoniana, *Mart. in Palmet. Orb.* 70.—Porto Rico.
pectinata, *Mart. Hist. Nat. Palm.* ii. 98. t. 60, 73. A. f. 1, 2.—Bras.
pilosa, *Karst. in Linnaea*, xxviii. (1856) 405.—N. Granat.
Piritu, *Wendl. ex H. Wendl. in Kerch. Palm.* 234.—N. Granat.
Piranga, *Trail, in Journ. Bot.* xv. (1877) 41.—Bras.
piscatorum, *Wedd. ex Drude, in Mart. Fl. Bras.* iii. II. 354.—Bras.
Plumeriana, *Mart. in Palmet. Orb.* 64.—Ind. occ.
praemorsa, *Poepp. ex Mart. l. c.* 66.—Peruv.
pulchra, *Trail, in Journ. Bot.* xv. (1877) 4. t. 184. f. 2.—Bras.
Puyamo, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Reg. Granat.
riparia, *Mart. Hist. Nat. Palm.* ii. 97. t. 71. f. 4.—Bras.
rivularis, *Barb.-Rodr. Enum. Palm. Nov.* 36.—Bras.
rotunda, Stokes, Bot. Mat. Med. iv. 394 = minor.
Sanctae-Paulae, *Engl. in Linnaea* (1864–65), 667.—N. Granat.
sciophila, *Miq. in Verh. Nat. Wet. Haarl.* vii. (1851) 208.—Guiana.
setipinnata, Barb.-Rodr. Enum. Palm. Nov. 32 = hylophila.
setosa, *Mart. Hist. Nat. Palm.* ii. 94. t. 72. f. 1–3.—Bras.
setulosa, *Karst. in Linnaea*, xxviii. (1856) 408.—Venezuela.
simplicifrons, *Mart. Hist. Nat. Palm.* ii. 103. t. 73. C. f. 1–2.—Bras.
simplicifrons, Spruce, in Journ. Linn. Soc. xi. (1869) 148 = geonomoides.
socialis, Mart. in Palmet. Orb. 56. t. 14. f. 2 = major.
speciosa, *Karst. in Linnaea*, xxviii. (1856) 402.—N. Granat.
sphaerocarpa, *Trail, in Journ. Bot.* xv. (1877) 8.—Amazon.
subglobosa, *H. Wendl. in Kerch. Palm.* 234.—Salvador.
syagroides, *Trail, in Journ. Bot.* xv. (1877) 76.—Bras.
sylvatica, *Barb.-Rodr. Enum. Palm. Nov.* 30.—Bras.

BACTRIS:—
tenera, *H. Wendl. in Kerch. Palm.* 234.—Hab.?
tenerrima, *Mart. ex Drude, in Mart. Fl. Bras.* iii. II. 328.—Bras.
tenuis, Wallace, Palm Trees Amaz. 87. t. xxxiii. = cuspidata.
tomentosa, *Mart. Hist. Nat. Palm.* ii. 100. t. 73. A. f. 3, 4.— Bras.
Trailiana, *Barb.-Rodr. Enum. Palm. Nov.* 27.—Bras.
trichospatha, *Trail, in Journ. Bot.* xv. (1877) 41, 42.—Bras.
turbinata, Spruce, in Journ. Linn. Soc. xi. (1871) 146, 152 = pectinata.
turbinocarpa, *Barb.-Rodr. Enum. Palm. Nov.* 33.—Bras.
uaupensis, Spruce, in Journ. Linn. Soc. xi. (1871) 145, 150 = cuspidata.
umbraticola, *Barb.-Rodr. Enum. Palm. Nov.* 34.—Bras.
umbrosa, Barb.-Rodr. l. c. 29 = trichospatha.
utilis, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* iii. 413.—Costa Rica.
varinensis, *Hort. ex H. Wendl. in Kerch. Palm.* 234.—Hab.?
villosa, *H. Wendl. ex Hemsl. Biol. Centr. Am. Bot.* iii. 413.—Costa Rica.
vulgaris, *Barb.-Rodr. Enum. Palm. Nov.* (1879) *App.* 42.—Bras.
xanthocarpa, Barb.-Rodr. l. c. 30 = simplicifrons?

BACTYRILOBIUM, Willd. Enum. Hort. Berol. 439 (1809) = **Cassia**, Linn. (Legumin.).
biflorum, Willd. ex Steud. Nom. ed. II. i. 179 = C. biflora.
ferrugineum, Schrad. in Goett. Gel. Anz. (1821) I. 713 = C. ferruginea.
Fistula, Willd. Enum. Hort. Berol. 440 = C. Fistula.
grande, Hornem. Hort. Hafn. i. 135 = C. grandis.
latifolium, Steud. Nom. ed. I. 41 = Anagyris latifolia.
molle, Schrad. in Goett. Gel. Anz. (1821) I. 713 = C. grandis.

BACULARIA, F. Muell. Fragm. vii. 103 (1870). *PALMAE*, Benth. & Hook. f. iii. 903.
LINOSPADIX, H. Wendl. & Drude, in Linnaea, xxxix. (1874) 177.
minor, *F. Muell. l. c.* xi. 58.—Austral.
monostachya, *F. Muell. l. c.* vii. 103; xi. 58.—Austral.

BADAMIA, Gaertn. Fruct. ii. 90. t. 97 (1791) = **Terminalia**, Linn. (Combret.).
Commersoni, Gaertn. l. c. = T. Catappa.

BADAROA, Bert. ex Steud. Nom. ed. II. i. 179 (1840) = seq.

BADEROA, Bert. ex Hook. Bot. Misc. iii. (1833) 324 = **Sicyos**, Linn. (Cucurb.).
arenaria, Bert. ex Steud. Nom. ed. II. i. 179 = seq.
bryoniaefolia, Bert. ex Hook. l. c. = S. bryoniaefolius.

BADIERA, DC. Prod. i. 334 (1824). *POLYGALEAE*, Benth. & Hook. f. i. 137.
acuminata, *DC. l. c.* 335.—Mexic.
Berteriana, *Spreng. Syst.* iii. 172.—Ins. S. Doming.
divaricata, *DC. Prod.* i. 335.—Bras.
diversifolia, *DC. l. c.* 334.—Jamaica.
domingensis, *DC. l. c.*—Ins. S. Doming.
Penaea, *DC. l. c.* 335.—Am. trop.

BADIERA, Hassk. Cat. Pl. Hort. Bogor. Alt. 227 (1844) = **Polygala**, Linn.
pulchra, Hassk. l. c. = P. venenosa.
venenosa, Hassk. l. c. = P. venenosa.

BADULA, Juss. Gen. 420 (1789) = **Ardisia**, Sw. (Myrsin.).
angustifolia, A. DC. in Trans. Linn. Soc. xvii. (1837) 114 = Embelia angustifolia.
arborea, Thou. ex A. DC. in DC. Prod. viii. 84 = Embelia arborea.
Barthesia, A. DC. in Trans. Linn. Soc. xvii. (1837) 113 = A. Barthesia.
Barthesia, Boj. Hort. Maurit. 196 = A. insularis.
borbonica, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 90.—Ins. Borbon.
caribaea, A. DC. l. c. 89 = A. caribaea.

BADULA :—
crassa, A. DC. in Trans. Linn. Soc. xvii. (1837) 114 = Ardisia insularis.
cybianthoides, A. DC. in DC. Prod. viii. 107 = Cybianthus philippinensis.
divaricata, Thou. ex A. DC. l. c. 89 = Oncostemum divaricatum.
excelsa, Thou. ex A. DC. l. c. 113 = (*Monoporus paludosus*).—Madag.
insularis, A. DC. in Trans. Linn. Soc. xvii. (1837) 113 = A. insularis.
iteoides, Benth. Pl. Hartw. 217.—N. Granat.
Lamarkiana, A. DC. in DC. Prod. viii. 110 = A. laurifolia.
lanceolata, Boj. ex A. DC. l. c. 90 = Oncostemum Bojeranum.
lateriflora, Griseb. in Goett. Abh. vii. (1857) 237 = A. laurifolia.
laurifolia, Boj. ex A. DC. in DC. Prod. viii. 108.—Madag.
Lhotskyana, A. DC. l. c. 111 = A. Lhotskyana.
Mamcicillo, Schlecht. in Linnaea, xxvi. (1853–55) 671.—N. Granat.
micrantha, Boj. Hort. Maurit. 196 = A. insularis.
micrantha, A. DC. in Trans. Linn. Soc. xvii. (1837) 112.—Ins. Maurit.
multiflora, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 89 = A. insularis.
myrtifolia, Thou. ex A. DC. in DC. Prod. viii. 90 = Oncostemum myrtifolium.
nemorosa, Thou. ex A. DC. l. c. = Oncostemum nemorosum.
nigricans, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 89.—Bras.
ovalifolia, A. DC. in Trans. Linn. Soc. xvii. (1837) 114.—Ins. Maurit.
paludosa, Boj. ex A. DC. in DC. Prod. viii. 113 (= *Monosporus paludosus*).—Madag.
pauciflora, Boj. ex A. DC. l. c. 90 = Oncostemum pauciflorum.
peruviana, A. DC. l. c. 112 = A. nigricans.
philippinensis, A. DC. l. c. 108 = A. subcuneata.
reticulata, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 89 = A. insularis.
scandens, Boj. Hort. Maurit. 196 = Embelia micrantha.
Schomburgkiana, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 89.—Guiana.
Sieberi, A. DC. in Trans. Linn. Soc. xvii. (1837) 113. t. 5 = A. guadalupensis.

BADUSA, A. Gray, in Proc. Am. Acad. iv. 308 (1860). *RUBIACEAE*, Benth. & Hook. f. ii. 42.
corymbiflora, *A. Gray, l. c.*—Ins. Pacif.

BADUSA Barthesia, Steud. Nom. ed. II. i. 98, in syn. (1840), sphalm. = BADULA, Juss. = Ardisia Barthesia.

BAEA, Comm. ex Juss. Gen. 121 (1789) = **Boea**, Comm. (Gesner.).

BAECKEA, Burm. f. Prod. Fl. Cap. 12 (1768) = **Brunia**, Linn.
africana, Burm. f. l. c. = Br. pinifolia.
cordata, Burm. f. l. c. = Br. racemosa.
lancifolia, Eckl. & Zeyh. Enum. 140 = Br. racemosa.
laurifolia, Sond. in Harv. & Sond. Fl. Cap. ii. 315, in syn. = Br. racemosa.
racemosa, Eckl. & Zeyh. Enum. 140 = Br. racemosa.
thyrsiflora, Harv. & Sond. Fl. Cap. ii. 604 = Br. pinifolia.
thyrsophora, Eckl. & Zeyh. Enum. 140 = Br. pinifolia.

BAECKEA, Linn. Sp. Pl. 358 (1753). *MYRTACEAE*, Benth. & Hook. f. i. 701.
Allostis, Rafin. Sylva Tellur. 104 (1838).
Babingtonia, Lindl. in Bot. Reg. (1842) t. 10.
Camphoromyrtus, Schau. in Linnaea, xvii. (1843) 240.
Cyathostemon, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 331.
Ericomyrtus, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 154.
Euryomyrtus, Schau. in Linnaea, xvii. (1843) 239.
Harmogia, Schau. in Linnaea, xvii. (1843) 238.
Imbricaria, Sm. in Trans. Linn. Soc. iii. (1797) 257.

BAECKEA :—
Jungia, Gaertn. Fruct. i. 175. t. 35 (1788).
Mollia, J. F. Gmel. Syst. 420 (1791).
Murrinea, Rafin. Sylva Tellur. 104 (1838).
Neuhofia, Stokes, Bot. Mat. Med. i. 439 (1812).
Oxymyrrhine, Schau. in Linnaea, xvii. (1843) 240.
Rinzia, Schau. l. c. xvii. 239 (1843).
Schidiomyrtus, Schau. l. c. 237 (1843).
Tetrapora, Schau. l. c. 238 (1843).
Tetraspora, Miq. in Nederl. Kruidk. Arch iv. (1859) 150.
Tjongina, Adans. Fam. ii. 234 (1763).
Triplarina, Rafin. Sylva Tellur. 104 (1838).
affinis, Endl. in Enum. Pl. Hueg. 51 = Astartea fasciculatis.
affinis, Hook. Ic. Pl. t. 284 c = diffusa.
alpina, Lindl. in Mitch. Three Exped. ii. 178 = diffusa.
arbuscula, *R. Br. ex Benth. Fl. Austral.* iii. 79.—Austral.
astartioides, *Benth. l. c.* 80.—Austral.
Behrii, *F. Mucll. Fragm.* iv. 68.—Austral.
Blacketii, *F. Muell. l. c.* viii. 181.—Austral.
brevifolia, *DC. Prod.* iii. 230.—Austral.
calycina, Lindl. in Mitch. Three Exped. ii. 190 = Thryptomene Mitchelliana.
camphorata, *R. Br. in Bot. Mag.* t. 2694.—Austral.
Camphorosmae, *Endl. in Enum. Pl. Hueg.* 51.—Austral.
capitata, *F. Muell. Census*, 54.—Austral.
carnulosa, Sieber, in Spreng. Syst. iv. Cur. Post. 149 = brevifolia.
chinensis, Gaertn. Fruct. i. 157 = frutescens.
ciliata, *F. Muell. Census*, 54.—Austral.
cochinchinensis, *Blume, Mus. Bot. Lugd. Bat.* i. 69.—Cochinch.
corymbulosa, *Benth. Fl. Austral.* iii. 87.—Austral.
corynophylla, *F. Muell. Fragm.* iv. 72.—Austral.
crassifolia, *Lindl. in Mitch. Three Exped.* ii. 115.—Austral.
crenatifolia, *F. Muell. Fragm.* iv. 70.—Austral.
crenulata, *DC. Prod.* iii. 230.—Austral.
crispiflora, *F. Muell. Fragm.* iv. 72.—Austral.
cryptandroides, *F. Muell. l. c.* x. 29.—Austral.
Cumingeana, Schau. in Walp. Rep. ii. 920 = frutescens.
Cunninghamii, *Benth. Fl. Austral.* ii. 82.—Austral.
densifolia, *Sm. in Trans. Linn. Soc.* iii. (1797) 260.—Austral.
diffusa, *Sieber, ex DC. Prod.* iii. 230.—Austral.
dimorphandra, *F. Muell. ex Benth. Fl. Austral.* iii. 74.—Austral.
diosmifolia, *Rudge, in Trans. Linn. Soc.* viii. (1807) 298. t. 13.—Austral.
diosmoides, Sieber, ex DC. Prod. iii. 230 = crenulata.
Drummondii, *Benth. Fl. Austral.* iii. 75.—Austral.
ericaea, *F. Muell. Fragm.* i. 31.—Austral.
ericoides, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 184.—N. Caled.
ericoides, *Schlecht. in Bot. Zeit.* iv. (1846) 724.—Austral.
fasciculata, Sieber, in Spreng. Syst. iv. Cur. Post. 149 = densifolia.
floribunda, *Benth. Fl. Austral.* iii. 87.—Austral.
frutescens, *Linn. Sp. Pl.* 358.—Malaya; China.
fumana, *F. Muell. Fragm.* iv. 68.—Austral.
gracilis, *A. Cunn. in Field, New South Wales*, 349.—Austral.
grandiflora, *Benth. Fl. Austral.* iii. 89.—Austral.
Gunniana, *Schau. ex Walp. Rep.* ii. 920.—Austral.
involucrata, Endl. in Enum. Pl. Hueg. 170 = Scholtzia obovata.
laxiflora, *F. Muell. Census*, 54.—Austral.
leptocaulis, *Hook. f. in Hook. Ic. Pl.* t. 298.—Tasmania.
leptantha, *F. Mucll. Census*, 54.—Austral.
leptocalyx, F. Muell. Fragm. i. 30 = Micromyrtus leptocalyx.
linifolia, *Rudge, in Trans. Linn. Soc.* viii. (1807) 297. Austral.
micrantha, DC. Prod. iii. 230 = Thryptomene baeckeacea.
micrantha, Hook. f. in Hook. Ic. Pl. t. 309 = Gunniana.
microphylla, A. Cunn. ex Schau. in Linnaea, xvii. (1843) 238 = Harmogia Cunninghami.
microphylla, Sieber, ex Spreng. Syst. iv. Cur. Post. 149 = Micromyrtus microphylla.
nclitrioides, *Seem. in Journ. Bot.* ii. (1864) 74.—N. Caled.

BAECKEA :—
nova-anglica, F. Muell. Fragm. iv. 71 = densifolia.
obovata, DC. Prod. iii. 230 = Scholtzia obovata.
obtusifolia, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 185.—N. Caled.
ochropetala, *F. Muell. Fragm.* x. 29.—Austral.
oligandra, *F. Muell. Census*, 54.—Austral.
oligomera, *Radlk. in Ber. Deutsch. Bot. Ges.* ii. (1884) 264.—Austral.
ovalifolia, *F. Muell. Fragm.* iv. 72.—Austral.
oxycoccoides, *Benth. Fl. Austral.* iii. 75.—Austral.
pachyphylla, *Benth. l. c.* 85.—Austral.
parviflora, *F. Muell. Census*, 54.—Austral.
parvula, DC. Prod. iii. 229 = virgata.
pentagonantha, *F. Muell. Fragm.* iv. 73.—Austral.
pentandra, *F. Muell. l. c.* ii. 31; iv. 72.—Austral.
phylicoides, A. Cunn. ex Schau. in Walp. Rep. ii. 921 = Kunzea peduncularis.
pinifolia, *DC. Prod.* iii. 229.—N. Caled.
platystemona, *Benth. Fl. Austral.* iii. 74.—Austral.
plicata, F. Muell. Fragm. i. 30 = Micromyrtus microphylla.
polyandra, *F. Muell. l. c.* iv. 72.—Austral.
polystemonea, *F. Muell. l. c.* ii. 124.—Austral.
prostrata, Hook. f. Ic. Pl. t. 284 b = diffusa.
pulchella, *DC. Prod.* iii. 230.—Austral.
pygmaea, *R. Br. ex Benth. Fl. Austral.* iii. 86.—Austral.
ramosissima, *A. Cunn. in Field, New South Wales*, 249.—Austral.
robusta, *F. Muell. Fragm.* iv. 72.—Anstral.
saxicola, *A. Cunn. ex Hook. Bot. Mag.* t. 3160.—Austral.
scholleraefolia, *Lchm. Pl. Preiss.* ii. 369.—Austral.
serpillifolia, *F. Muell. Fragm.* x. 30.—Austral.
sinensis, Gaertn. Fruct. i. t. 31 = frutescens.
speciosa, A. Cunn. ex Schau. in Linnaea, xvii. (1843) 241 = Hypocalymma strictum.
spathulata, *F. Muell. Census*, 54.—Anstral.
spinosa, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 149.—Austral.
stenophylla, *F. Muell. Fragm.* i. 13.—Austral.
subcuneata, *F. Muell. l. c.* iv. 73.—Austral.
sumatrana, *Blume, Mus. Bot. Lugd. Bat.* 69.—Sumatra.
teretifolia, *F. Mucll. Census*, 54.—Austral.
tetragona, *F. Muell. ex Benth. Fl. Austral.* iii. 77.—Austral.
thymifolia, Hook. f. Ic. Pl. t. 284 a = diffusa.
trichophylla, Sieber, ex Spreng. Syst. iv. Cur. Post. 149 = linifolia.
uberiflora, *F. Muell. Census*, 53.—Austral.
umbellata, F. Muell. Fragm. iv. 69 = virgata.
umbellifera, *F. Muell. Census*, 54.—Austral.
unicinella, *Benth. Fl. Austral.* iii. 84.—Austral.
utilis, F. Muell. ex Miq. in Neder. Kruidk. Arch. iv. (1859) 150 = Gunniana.
virgata, *Andr. Bot. Rep.* 598.—N. Caled.

BAEICA, C. B. Clarke, in DC. Monog. Phan. v. 134 (1883) = **Boeica**, C. B. Clarke (Gesner.).

BAEOBOTRYS, Forst. Char. Gen. 21. t. 11 (1776) = **Maesa**, Forsk. (Myrsin.).
acuminata, Spreng. Syst. in Ind. v. 89 = M. ramentacea.
acuminata, Wall. Cat. n. 2321 = Gymnosporia acuminata.
argentea, Wall. l. c. n. 2316 = Maesa argentea.
Bogota, Buch.-Ham. ex A. DC. in DC. Prod. viii. 81 = M. macrophylla.
dubia, Wall. Cat. n. 2317 = M. dubia.
fragrans, Wall. l. c. n. 2322 = M. ramentacea.
fulvinervis, Kollm. ex A. DC. in DC. Prod. viii. 82 = M. mollis.
glabra, Roxb. Hort. Beng. 16; Fl. Ind. i. 560 = M. ramentacea.
indica, Roxb. ll. cc. 16, 557 = M. indica.
indica, Wall. Cat. n. 2318. A & B = M. Chisia.
lanceolata, Blume, Bijdr. 865 = M. Blumii.
lanceolata, Spreng. Syst. iv. Cur. Post. 83 = M. ramentacea.
lanceolata, Vahl, Symb. Bot. i. 19 = M. lanceolata.
latifolia, Blume, Bijdr. 865 = M. indica.
lucida, Wall. Cat. n. 2323.—Ind. or.
lutea, Roxb. ex Steud. Nom. ed. II. i. 180 = M. lutea.
macrophylla, Wall. Cat. n. 2325 = M. macrophylla.
missionis, Wall. l. c. n. 6523 = M. ramentacea?

BAEOBOTRYS:—
mollis, Blume, Bijdr. 865 = Maesa mollis.
mollissima, Blume, l. c. 866 = M. mollissima.
Morsha, Buch.-Ham. ex A. DC. in DC. Prod. viii. 80 = M. indica.
muscosa, Reinw. ex Blume, Bijdr. 866 = M. membranacea.
nemoralis, Forst. Char. Gen. 22 = M. nemoralis.
nemoralis, Roxb. Hort. Beng. 85; Fl. Ind. i. 559 = M. indica.
ovata, Wall. Cat. n. 2324 = M. ramentacea.
paniculata, Wall. l. c. n. 2320 = M. paniculata.
picta, Hochst. ex Walp. Ann. iii. 964 = M. lanceolata.
pubescens, Lodd. ex Loud. Hort. Brit. 71.—M. pubescens.
ramentacea, Roxb. Hort. Beng. 16; Fl. Ind. i. 558 = M. ramentacea.
rufescens, E. Mey. ex A. DC. in DC. Prod. viii. 81 = M. lanceolata.
tetrandra, Roxb. Hort. Beng. 85; Fl. Ind. i. 560 = M. tetrandra.
tomentosa, Buch.-Ham. ex Spreng. Syst. iv. Cur. Post. 83 = M. macrophylla.
virgata, Blume, Bijdr. 866 = M. indica.

BAEOMETRA, Salisb. in Trans. Hort. Soc. i. (1812) 330. *LILIACEAE*, Benth. & Hook. f. iii. 823.
JANIA, Schult. Syst. vii. II. p. xcviii. 1528 (1830).
KOLBEA, Schlecht. in Linnaea, i. (1826) 80.
columellaris, *Salisb. l. c.*—Afr. austr.

BAEOTERPE, Salisb. Gen. Pl. Fragm. 17 (1866) = **Hyacinthus**, Linn. (Liliac.).
brevifolia, Salisb. l. c. 18 = H. brevifolius.
corymbosa, Salisb. l. c. = H. corymbosus.

BAEOTHRYON, Ehrh. Beitr. iv. 147 (1789) = **Scirpus**, Linn. (Cyperac.).
caespitosum, A. Dietr. Sp. Pl. ii. 89 = S. caespitosus.
confervoides, A. Dietr. l. c. 94 = S. confervoides.
crinitum, A. Dietr. l. c. = Ficinia filiformis.
festucoides, A. Dietr. l. c. = Bulbostylis festucoides.
fistulosum, A. Dietr. l. c. = Eleocharis fistulosa.
flavescens, A. Dietr. l. c. 91 = Eleocharis ocreata.
Halleri, T. Nees, Gen. Germ. Ic. ii. 17 = S. pauciflorus.
Hystrix, A. Dietr. Sp. Pl. ii. 92 = Hypodiscus aristatus.
nanum, A. Dietr. l. c. 91 = S. nanus.
pauciflorum, A. Dietr. l. c. 90 = S. pauciflorus.
pusillum, A. Dietr. l. c. 92 = S. pusillus.
pygmaeum, A. Dietr. l. c. = Eleocharis Chaetaria.
retroflexum, A. Dietr. l. c. 93 = Eleocharis Chaetaria.
tenuiculum, A. Dietr. l. c. = Eleocharis Rothiana.
trichodes, A. Dietr. l. c. = Eleocharis trichodes.
variegatum, A. Dietr. l. c. 92 = Eleocharis variegata.

BAERIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 29 (1835). *COMPOSITAE*, Benth. & Hook. f. ii. 399.
DICHAETA, Nutt. in Trans. Am. Phil. Soc. N. S. VII. (1841) 383.
PTILOMERIS, Nutt. in Am. Phil. Trans. N. S. vii. (1841) 381.
affinis, *A. Gray, in Proc. Am. Acad.* xix. (1883) 23.—Calif.
anthemoides, *A. Gray, l. c.* 23.—Calif.
carnosa, *Greene, in Bull. Torrey Club*, x. (1883) 86.—Calif.
chrysostoma, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1835) 29.—Calif.
Clevelandi, *A. Gray, in Proc. Am. Acad.* xix. (1883) 22.—Calif.
coronaria, *A. Gray, l. c.* 23.—Calif.
curta, *A. Gray, l. c.* 21.—Calif.
debilis, *Greene, ex A. Gray, Syn. Fl. N. Am.* i. II. 325.—Calif.
Fremontii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 358; ix. (1874) 196.—Calif.
gracilis, *A. Gray, l. c.* ix. (1874) 196.—Calif.; Arizon.
leptalea, *A. Gray, Syn. Fl. N. Am.* i. II. 325.—Calif.
macrantha, *A. Gray, in Proc. Am. Acad.* xix. (1883) 21.—Calif.
maritima, *A. Gray, l. c.* ix. (1874) 196; *Bot. Calif.* i. 376.—Calif.
mutica, *A. Gray, in Proc. Am. Acad.* xix. (1883) 23.—Calif.
Palmeri, *A. Gray, in Bot. Calif.* i. 376.—Calif.

BAERIA:—
platycarpha, *A. Gray, in Proc. Am. Acad.* ix. (1874) 196.—Calif.
tenella, *A. Gray, l. c.* xix. (1883) 23.—Calif.
tenerrima, *A. Gray, l. c.* ix. (1874) 196.—Calif.
uliginosa, *A. Gray, l. c.*—Calif.

BAEUMERTA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 419 (1800) = **Nasturtium**, R. Br. (Crucif.).
Nasturtium, Gaertn. Mey. & Scherb. l. c. 467 = N. officinale.

BAGALATTA, Roxb. ex Reichb. Consp. 86 (1828) = **Tiliacora**, Colebr. (Menisperm.) (T. racemosa).

BAGASSA, Aubl. Pl. Gui. Suppl. 15. t. 376 (1775). *URTICACEAE*, Benth. & Hook. f. iii. 362.
LAUREA, Gaudich. Voy. Bonite, t. 88 (1826).
LOUREA, Kunth, in Linnaea, xiii. (1839) 562.
guianensis, *Aubl. l. c.*—Guiana.
Sagotiana, *Bur. ex Benth. & Hook. f. Gen.* iii. 362.—Guiana.

BAGNISIA, Becc. Malesia, i. 249. t. 12 (1878). *BURMANNIACEAE*, Benth. & Hook. f. iii. 459.
SARCOSIPHON, Blume, Mus. Bot. Lugd. Bat. i. 65. t. 18 (1849).
crocea, *Becc. l. c.*—Malaya.

BAHEL, Adans. Fam. ii. 210 (1763) = **Hygrophila**, R. Br. (Acanth.) (H. spinosa).

BAHIA, Lag. Gen. et Sp. Nov. 30 (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 402.
STYLESIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 377.
TRICHOPHYLLUM, Nutt. Gen. Am. ii. 166 (1818).
VIRLETIA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 402 (1873).
absinthifolia, *Benth. Pl. Hartw.* 18.—Texas; Mexic.
achilleoides, DC. Prod. v. 657 = Eriophyllum caespitosum.
ambigua, A. Gray, in Bot. Calif. i. 382 = Eriophyllum ambiguum.
ambrosioides, *Lag. Gen. et Sp. Nov.* 28, 30.—Chili.
angustifolia, DC. Prod. v. 656 = Calea angustifolia.
anthemoides, *A. Gray, in Proc. Am. Acad.* xv. (1880) 40.—Mexic.
arachnoidea, Fisch. & Avé-Lall. in Ind. Sem. Hort. Petrop. ix. 63 = Eriophyllum caespitosum.
artemisiaefolia, Less. in Linnaea, iv. (1829) 160 = Eriophyllum staechadifolium.
Bigelovii, *A. Gray, in Torr. Bot. Mex. Bound.* 96.—Texas.
biternata, *A. Gray, Pl. Wright.* ii. 95.—Texas.
chrysanthemoides, *A. Gray, in Proc. Am. Acad.* xix. (1883) 28.—Am. bor. occ.
confertiflora, DC. Prod. v. 657 = Eriophyllum confertiflorum.
cuneata, Kellogg, in Proc. Calif. Acad. v. (1873) 49 = Eriophyllum caespitosum.
dealbata, A. Gray, Pl. Fendl. 99 = absinthifolia.
Gilliesii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 28.—Reg. Argent.
gracilis, Hook. & Arn. Bot. Beech. Voy. 353 = Eriophyllum gracile.
integrifolia, DC. Prod. v. 656 = Eriophyllum caespitosum.
lanata, DC. l. c. 657 = Eriophyllum caespitosnm.
latifolia, Benth. Bot. Voy. Sulph. 30 = Eriophyllum caespitosnm.
latifolia, *Lindl. in Journ. Hort. Soc.* viii. (1853) 319.—Calif.
leucophylla, DC. Prod. v. 657 = Eriophyllnm caespitosum.
leucophylla, Eaton, Bot. King, Exped. 17, pro parte = Eriophyllum Watsoni.
multiflora, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 373 = Eriophyllum caespitosum.
neo-mexicana, *A. Gray, in Proc. Am. Acad.* xix. (1883) 27.—N. Mexic.
nepetaefolia, A. Gray, l. c. v. (1861) 184 = sinnata.
nudicaulis, *A. Gray, l. c.* xix. (1883) 27.—Am. bor. occ.
oblongifolia, *A. Gray, l. c.*—Colorado.
oppositifolia, *DC. Prod.* v. 656.—Am. bor. occ.
Palmeri, *A. Gray, Syn. Fl. N. Am.* ii. I. 452.—Calif.

BAHIA:—
parviflora, A. Gray, in Bot. Calif. i. 382 = Eriophyllum ambiguum.
pedata, *A. Gray, Pl. Wright.* i. 123.—Texas.
resinosa, DC. Prod. vi. 678 = Galinsoga resinosa.
rubella, A. Gray, in Torr. Bot. Mex. Bound. 95 = Eriophyllum Wallacci.
schkuhrioides, *A. Gray, in Proc. Am. Acad.* xix. (1883) 27.—Mexic.
sinuata, *Less. in Linnaea*, vi. (1831) 160.—Mexic.
staechadifolia, DC. Prod. v. 656 = Eriophyllum staechadifolium.
tenuifolia, *DC. l. c.* 657.—Calif.
trifida, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 374 = Eriophyllum confertiflorum.
trolliifolia, *DC. Prod.* v. 657.—Mexic.
Wallacei, A. Gray, in Pacif. Rail. Rep. iv. 105 = Eriospermum Wallacei.
Wallacei, A. Gray, in Proc. Nat. Hist. Bost. vii. (1859-61) 145 = Eriospermum ambiguum.
Woodhousii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 28.—N. Mexic.

BAHIOPSIS, Kellogg, in Proc. Calif. Acad. Nat. Soc. ii. (1863) 35 = **Viguiera**, H. B. & K. (Compos.).
lanata, Kellogg, l. c. = V. nivea?

BAIKAEA, Benth. in Benth. & Hook. f. Gen. i. 581 (1865). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
insignis, *Benth. in Trans. Linn. Soc.* xxv. (1866) 314. t. 41.—Afr. trop.
minor, *Oliver, Fl. Trop. Afr.* ii. 309.—Afr. trop.

BAILEYA, Harv. & Gray, ex Torr. in Emory, Notes Mil. Reconnois. 144 (1848). *COMPOSITAE*, Benth. & Hook. f. ii. 423.
multiradiata, *Harv. & Gray, ex Torr. l. c.*—Am. bor. occ.
pauciradiata, *Harv. & Gray, ex A. Gray, Pl. Fendl.* 105.—Calif.
pleniradiata, Harv. & Gray, ex A. Gray, l. c. = multiradiata.

BAILLIERIA, Aubl. Pl. Gui. ii. 804. t. 317 (1775) = **Clibadium**, Linn. (Compos.).
aspera, Aubl. l. c. = C. surinamense.
Barbasco, H. B. & K. Nov. Gen. et Sp. iv. 289 = C. Barbasco.
caroliniana, Spreng. Syst. iii. 625 = Iva imbricata.
commelinoides, Less. Syn. Comp. 215 = Ichthyothere Cunabi.
erosa, Poir. Encyc. Suppl. i. 563 = C. erosum.
nereifolia, H. B. & K. Nov. Gen. et Sp. iv. 289 = C. nereifolium.
pedunculosa, Poir. Encyc. Suppl. i. 563 = C. pedunculosum.
sylvestris, Aubl. Pl. Gui. ii. 807 = C. surinamense.
terebinthacea, Poir. Encyc. Suppl. i. 563 = C. terebinthaceum.

BAILLONIA, Bocq. ex Baill. Adansonia, ii. (1861-62) 251. t. 7. *VERBENACEAE*, Benth. & Hook. f. ii. 1143.
DIOSTEA, Miers, in Trans. Linn. Soc. xxvii. (1870) 103. t. 28.
amabilis, *Bocq. ex Baill. l. c.* 252.—Paraguay.
juncea, *Benth. & Hook. f. Gen.* ii. 1144.—Chili.

BAIMO, Rafin. Fl. Tellur. iv. 27 (1836) = **Fritillaria**, Tourn. (Liliac.).
cirrhosa, Rafin. l. c. = F. japonica.

BAISSEA, A. DC. Prod. viii. 424 (1844). *APOCYNACEAE*, Benth. & Hook. f. ii. 719.
CLEGHORNIA, Wight, Ic. t. 1310, 1312 (1850).
acuminata, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 663.—Zeylan.
leonensis, *Benth. in Hook. Niger Fl.* 452.—Afr. trop.
malaccensis, *Hook. f. Fl. Brit. Ind.* iii. 663.—Malacca.
multiflora, *A. DC. in DC. Prod.* viii. 424.—Afr. trop.

BAITARIA, Ruiz & Pav. Prod. 63. t. 36 (1797) = **Calandrinia**, H. B. & K. (Portulac.).
acaulis, *Ruiz & Pav. Fl. Per.* iv. t. 403; *Syst. Veg.* 111.—Peruv.

BAJAN, Adans. Fam. ii. 506 (1763) = **Amaranthus**, Linn.

BAKERIA, Seem. Journ. Bot. ii. (1864) 248 = **Pierandra**, A. Gray (Araliac.).
vitiensis, Seem. l. c. = P. viticnsis.

BALAKA, Becc. in Ann. Jard. Buitenz. ii. (1885) 91 in obs. *PALMAE*.
perbrevis, *Becc. l. c.*—Ins. Fiji.
Seemanni, *Becc. l. c.*—Ins. Fiji.

BALANAULAX, Rafin. Alsog. Am. 28 (1838) = **Quercus**, Tourn. (Cupul.).
molucca, Rafin. l. c. = Q. molucca.
Rumphi, Rafin. l. c. = Q. molucca?

BALANGHAS, [Burm.] Rafin. Sylva Tellur. 72 (1838) = **Sterculia**, Linn.
rubiginosa, Rafin. l. c. = S. rubiginosa.
Telabo, Rafin. l. c. = S. Balanghas.

BALANGUE, Gaertn. Fruct. ii. 495. t. 180. f. 3 (1791). *JASMINEAE*?
Gaertneri, *DC. Prod.* viii. 316.—Madag.

BALANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 1 (Quid?).

BALANITES, Delile, Fl. Egypte, 221. t. 28. f. 1. (1813). *SIMARUBEAE*, Benth. & Hook. f. i. 314.
AGIALID, Adans. Fam. ii. 508 (1763).
aegyptiaea, *Delile, l. c.*—Afr. bor.; Arab.; Palaestin.
aegyptica, Wall. Cat. n. 6855 = Roxburghii.
ferox, G. Don, Gen. Syst. i. 774 = aegyptiaca.
Roxburghii, *Planch. in Ann. Sc. Nat.* Sér. IV. iv. (1854) 258. t. 2.—Ind. or.

BALANOCARPUS, Bedd. Forest. Man. Bot. 236 [bis] (1873). *DIPTEROCARPEAE*.
erosa, *Bedd. l. c.* 237.—Ind. or.
utilis, *Bedd. l. c.*—Ind. or.

BALANOPHORA, Forst. Char. Gen. t. 50 (1776). *BALANOPHOREAE*, Benth. & Hook. f. iii. 235.
CYNOPSOLE, Endl. Gen. 74 (1838).
SARCOCORDYLIS, Wall. Cat. n. 7249 (1831?).
SCYNOPSOLE, Reichb. Handb. 164 (1837).
abbreviata, *Blume, Enum. Pl. Jav.* i. 87.—Java.
affinis, Griff. in Proc. Linn. Soc. i. (1849) 220; Trans. Linn. Soc. xx. (1851) 94. t. 4 = dioica.
alutacea, Jungh. in Nov. Act. Nat. Cur. xviii. Suppl. i. (1841) 205 = abbreviata.
alveolata, Griff. in Proc. Linn. Soc. i. (1849) 220; Trans. Linn. Soc. xx. (1851) 94 = dioica.
Burmanniana, Griff. ll. cc. 219 et 93 = dioica.
capensis, Eckl. & Zeyh. ex Eichl. in DC. Prod. xvii. 125 = Mystropetalon Thomii.
dioica, *R. Br. in Wall. Cat.* n. 7246.—Ind. or.; Burma.
dioica, Unger, in Ann. Wien. Mus. ii. (1838) t. 2. f. 1, 2 = elongata.
elongata, *Blume, Enum. Pl. Jav.* i. 87.—Java.
elongata, Schott & Endl. Melet. 13, partim = dioica.
fungosa, *Forst. Char. Gen.* 99. t. 50.—Austral.; N. Hebrid.
gigantea, Wall. Cat. n. 7249 = globosa.
globosa, *Jungh. in Nov. Act. Nat. Cur.* xviii. *Suppl.* i. (1841) 210.—Burma; Java.
Harlandi, *Hook. f. in Trans. Linn. Soc.* xxii. (1859) 426.—Hongkong.
Hildebrandtii, *Reichb. f. in Journ. Bot.* xiv. (1876) 45.—Ins. Comoro.
indica, *Wall. Cat.* n. 7247.—Ind. or.
involucrata, *Hook. f. & Thoms. in Trans. Linn. Soc.* xxii. (1859) 30.—Reg. Himal.
Lowii, *Hook. f. l. c.* 426. t. 75 C.—Borneo.
maxima, Jungh. in Nov. Act. Nat. Cur. xviii. Suppl. i. (1841) 209. t. 1 = elongata.
picta, Griff. in Proc. Linn. Soc. i. (1849) 220; Trans. Linn. Soc. xx. (1851) 94 = dioica.
picta, Miq. ex Eichl. in DC. Prod. xvii. 146 = indica.
polyandra, *Griff. in Trans. Linn. Soc.* xx. (1851) 94.—Reg. Himal.
reflexa, *Becc. in Att. Soc. Ital. Sc. Nat. Milano*, xi. (1868) 197.—Borneo.
Thwaitesii, *Eichl. in DC. Prod.* xvii. 146, in nota.—Zeylan.
typhina, Wall. Cat. n. 7248 = dioica.

BALANOPLIS, Rafin. Alsog. Amer. 29 (1838) = **Quercus**, Tourn. (Cupul.).
serrata, Rafin. l. c. 30 = Q. cuspidata.
tribuloides, Rafin. l. c. 29 = Castanopsis tribuloides.

BALANOPS, Baill. Adansonia, x. (1871) 117, 337. *BALANOPSEAE*, Benth. & Hook. f. iii. 341, 1223.
australiana, *F. Muell. Fragm.* x. 114.—Austral.
Balansae, *Baill. Adansonia*, x. (1872) 338.—N. Caled.
microstachya, *Baill. l. c.* 340.—N. Caled.
oliviformis, *Baill. l. c.* 337.—N. Caled.
Pancheri, *Baill. l. c.* 118.—N. Caled.
Theophrasta, *Baill. l. c.* 338.—N. Caled.
Vieillardi, *Baill. l. c.* 118.—N. Caled.

BALANOPSIS, Rafin. Sylva Tellur. 134 (1838) = **Ocotea**, Aubl. (Laurin.).
acuta, Rafin. l. c. (= *Mespilodaphne cupularis*).
Cassia, Rafin. l. c. = Litsaca zeylanica?
cupulata, Rafin. l. c. (= *Mespilodaphne cupularis*).
elliptica, Rafin. l. c.—Hab.?

BALANOPTERIS, Gaertn. Fruct. ii. 94. t. 98 (1791) = **Heritiera**, Ait. (Sterculiac.).
minor, Gaertn. l. c. t. 98. f. 2 = H. Fomes.
Tothila, Gaertn. l. c. p. 94. t. 99 = H. littoralis.

BALANOSTREBLUS, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 247. t. 19. *URTICACEAE*, Benth. & Hook. f. iii. 377.
ilicifolia, *Kurz, l. c.* 248.—Chittagong; Burma; Penang.

BALANSAEA, Boiss. & Reut. Pugill. Pl. Nov. 49 (1852) = **Chaerophyllum**, Linn. (Umbellif.).
Fontanesii, Boiss. & Reut. l. c. 50.—Hispan.
glaberrima, Lange, Pugill. iv. 238 = Daucus glaberrimus.

BALANTIUM, Desv. ex Ham. Prod. Fl. Ind. Occ. 34 (1825) = **Parinarium**, Juss. (Rutac.).
cordifolium, Desv. l. c.—Ind. occ.?

BALARDIA, Cambess. in A. St. Hil. Fl. Bras. Mer. ii. 180. t. 111 (1829) = **Spergularia**, Presl (Caryophyll.).
platensis, Cambess. l. c. = S. platensis.

BALAUSTION, Hook. Ic. Pl. t. 852 (1852). *MYRTACEAE*, Benth. & Hook. f. i. 702.
CHEYNIA, J. Drumm. in Hook. Kew Journ. vii. (1855) 56.
PUNICELLA, Turcz. in Bull. Acad. Pétersb. x. (1852) 333.
pulcherrimum, *Hook. Ic. Pl.* t. 852.—Austral.

BALBISIA, Cav. in Anal. Cienc. Nat. vii. (1804) 62. t. 46. *GERANIACEAE*, Benth. & Hook. f. i. 276.
CRUKSHANKSIA, Hook. & Arn. in Hook. Bot. Misc. ii. (1831) 211. t. 90.
LEDOCARPON, Desv. in Mém. Mus. Par. vi. (1818) 250.
Meyeniana, *Klotzsch, in Linnaea*, x. (1836) 432.—Pernv.
pedunculatis, *D. Don, in Edinb. N. Phil. Journ.* xi. (1831) 277.—Chili.
verticillata, *Cav. in Anal. Cienc. Nat.* vii. (1804) 62. t. 46.—Chili.

BALBISIA, DC. in Guill. Archiv. Bot. ii. (1833) 333 = **Rhetinodendron**, Meissn. (Compos.).
Berterii, DC. l. c. = R. Berteroi.

BALBISIA, Willd. Sp. Pl. iii. 2214 (1803) = **Tridax**, Linn. (Compos.).
Caledoniae, Spreng. Fl. Hal. Mant. 52 = Olearia viscosa.
canescens, Rich. ex Pers. Syn. ii. 470 = T. procumbens.
divaricata, Cass. Opusc. iii. 91; ex DC. Prod. v. 679 = T. procumbens.
elongata, Willd. Sp. Pl. iii. 2214 = T. procumbens.
pedunculata, Hoffmgg. Verz. Pfl. (1824) 228 = T. procumbens.
suffruticosa, Dietr. ex DC. Prod. v. 267 = Olearia viscosa.

BALBOA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 106 = **Teplirosia**, Pers. (Legumin.).
diversifolia, Liebm. l. c. 107 = T. madrensis.

BALBOA, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 252. *GUTTIFERAE*, Benth. & Hook. f. i. 172.
membranacea, *Planch. & Triana, l. c.*—N. Granat.

BALDELLIA, Parl. Nuov. Gen. e Spec. 57 (1854) = **Alisma**, Linn.
ranunculoides, Parl. l. c. 58 = A. ranunculoides.

BALDINGERA, Dennst. Schluess. Hort. Malab. 31 (1818) = **Premna**, Linn. (Verben.).
glandulosa, Dennst. l. c. = P. longifolia.

BALDINGERA, Gaertn. Mey. & Scherb. Fl. Wett. i. 43 (1799) = **Phalaris**, Linn. (Gramin.).
arundinacea, Dum. Obs. Gram. Belg. 130 = P. arundinacea.
colorata, Gaertn. Mey. & Scherb. Fl. Wett. i. 96 = P. arundinacea.

BALDINGERIA, Neck. Elem. i. 88? (1790) = **Cotula**, Linn. (Compos.).

BALDINGERIA, F. W. Schmidt, Samml. Phys. Aufs. i. 277 (1795) = **Leontodon**, Linn. (Compos.).
pygmaea, F. W. Schmidt, l. c. = L. hirtum?
urens, F. W. Schmidt, l. c. = Hypochoeris cretensis.

BALDWINIA, Nutt. Gen. Am. ii. 175 (1818) (Balduina). *COMPOSITAE*, Benth. & Hook. f. ii. 391.
ACTINOSPERMUM, Ell. Sketch, ii. 448 (1817).
ACTISPERMUM, Rafin. New Fl. Am. i. 61 (1836).
ENDORIMA, Rafin. in Am. Monthly Mag. (1819) 195.
MNESITEON, Rafin. Fl. Ludov. 67 (1817).
bicolor, *Rafin. New Fl. Am.* iv. 73.—Am. bor.
lutea, *Rafin. l. c.*—Am. bor.
multiflora, *Nutt. Gen. Am.* ii. 175.—Am. bor.
uniflora, *Nutt. l. c.*—Am. bor.

BALDWINIA, Rafin. in Am. Monthly Mag. (1818) 267 = **Passiflora**, Linn.
peltata, Rafin. l. c. = P. peltata.

BALENDASIA, Rafin. Fl. Tellur. iv. 105 (1836) = **Passerina**, Linn. (Thymel.).
ericoides, Rafin. l. c. = P. ericoides.

BALENERDIA, Comm. ex Steud. Nom. ed. I. 99 (1821 = **Nanodea**, Banks (Santal.).

BALESSAM, Bruce, Trav. v. t. 25 (1790) = **Balsamodendron**, Kunth, sec. Endl. Gen. 136 (Burserac.).

BALFOURIA, R. Br. in Mem. Wern. Soc. i. (1810) 70. = **Wrightia**, R. Br. (Apocyn.).
saligna, R. Br. Prod. 467 = W. saligna.

BALFOURODENDRON, Mello, ex Oliver, in Hook. Ic. Pl. xiii. 3. tt. 1203–04 (1877). *RUTACEAE*.
eburneum, *Mello, l. c.*—Bras.

BALFURIA, Reichb. Consp. 134 (1828) = BALFOURIA, R. Br. = **Wrightia**, R. Br. (Apocyn.).

BALINGAYUM, Blanco, Fl. Filip. ed. I. 187 (1837) = **Erythropalum**, Blume (Olacin.).
decumbens, Blanco, l. c.—Ins. Philipp.

BALIOSPERMUM, Blume, Bijdr. 603 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 324.
angulare, Decne. ex Baill. Etud. Gén. Euph. 395 = axillare.
axillare, *Blume, Bijdr.* 604.—Ind. or.; Malaya.
calycinum, *Muell. Arg. in Flora*, xlvii. (1864) 470.—Reg. Himal.
indicum, Decne. in Jacquem. Voy. Bot. 154. t. 155 = axillare.
micranthum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 215.—Reg. Himal.
montanum, Muell. Arg. in DC. Prod. xv. II. 1125 = axillare.
Moritzianum, Baill. Etud. Gén. Euph. 395 = axillare.
polyandrum, Wight, Ic. t. 1885 = axillare.
reidioides, *Kurz, in Flora*, lviii. (1875) 32.—Siam.
sinuatum, *Muell. Arg. in Flora*, xlvii. (1864) 470.—Reg. Himal.

BALLARDIA, Montr. in Mém. Acad. Lyon, x. (1860) 204 = **Cloëzia**, Brongn. (Myrtac.).
elegans, Montr. l. c. 205.—N. Caled.

BALLARION, Rafin. in Am. Monthly Mag. (1818) 266 = **Stellaria**, Linn. (Caryophyll.).

BALLELA, Rafin. Fl. Tellur. iv. 70 (1836) = **Campanula** ?, Linn.
repens, Rafin. l. c. = C. repens.

BALLEXERDA, Comm. ex A. DC. Prod. xiv. 675 (1857) = **Nanodea**, Banks (Santal.).

BALLIERIA, Juss. Gen. 188 (1789) = BAILLIERIA, Anbl. = **Clibadium**, Linn. (Compos.).

BALLIMON, Rafin. New Fl. Am. iv. 28 (1836) = **Daucus**, Tourn. (Umbellif.).
maritimum, Rafin. l. c. = D. maritimus.
muricatulum, Rafin. l. c. = D. muricatus.

BALLOCHIA, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 86. *ACANTHACEAE.*
amoena, *Balf. f. l. c.* 87.—Ins. Socotr.
atro-virgata, *Balf. f. l. c.*— Ins. Socotr.
rotundifolia, *Balf. f. l. c.*—Ins. Socotr.

BALLOSPORUM, Salisb. Gen. Pl. Fragm. 142 (1866) = **Gladiolus**, Linn. (Irid.).
segetum, Salisb. l. c. = G. segetum.

BALLOTA, Linn. Gen. ed. I. 175 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1212.
ACANTHOPRASIUM, Spenn, in T. Nees, Gen. Fl. Germ. Gamop. i. (1843).
BERINGERIA, Neck. Elem. i. 312 (1790).
PSEUDODICTAMNUS, Tourn. ex Adans. Fam. ii. 191 (1763).
acetabulosa, *Benth. Lab. Gen. et Sp.* 595.—Graecia; As. Min.
africana, *Benth. l. c.* 594.—Afr. austr.
africana, Colm. Apuntes, 121 = hirsuta.
alba, *Linn. Fl. Suec.* ed. II. 206. n. 530.—Europ.
ampliata, Willd. ex Steud. Nom. ed. II. i. 181 = nigra.
arabica, Hochst. & Steud. ex Steud. l. c. = Leucas urticaefolia.
aristata, Reichb. Fl. Germ. Excurs. 325 = nigra.
armena, Boiss. Diagn. Ser. II. iv. 54 = rotundifolia.
Aucheri, *Boiss. l. c.* Ser. I. v. 39.—Kurdist.; Persia.
borealis, Schweig. in Koenigsb. Arch. Naturw. i. (1812) 214 = alba.
bracteosa, Ball, in Journ. Linn. Soc. xvi. (1878) 629 = nigra.
bullata, *Pomel, Nouv. Mat. Fl. Atl.* 116.—Algeria.
cinerea, D. Don, Prod. Fl. Nep. 111 = Roylea elegans.
damascena, *Boiss. Diagn.* Ser. I. xii. 87.—Syria; Aegypt.
dictamnifolia, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = pseudodictamnus.
disticha, Linn. Mant. 83 = Anisomeles ovata.
disticha, Rodsch. Obs. 35 = Hyptis pectinata.
foetida, Lam. Fl. Fr. ii. 381 = nigra.
Forskahlei, *Benth. Lab. Gen. et Sp.* 599.—Arabia.
frutescens, Coss. ex Nym. Consp. 581 = spinosa.
glandulifera, Trautv. in Act. Hort. Petrop. iii. II. (1875) 270 = rotundifolia.
Hildebrandtii, *Vatke & Kurtz, in Oestr. Bot. Zeitschr.* xxv. (1875) 95.—Afr. or.
hirsuta, *Benth. Lab. Gen. et Sp.* 595.—Hispan.; Afr. bor.
hirsuta, Schult. ex Steud. Nom. ed. II. i. 181 = nigra.
hirta, Hort. Berol. ex Steud. l. c. = nigra.
hispanica, *Benth. Lab. Gen. et Sp.* 597.—Europ. austr.
hispanica, Willk. & Lange, Prod. Fl. Hisp. ii. 446, in syn. = hirsuta.
integrifolia, *Benth. Lab. Gen. et Sp.* 599.—Ins. Cyprus.
italica, Benth. in DC. Prod. xii. 519 = rupestris.
lanata, Linn. Sp. Pl. 582 = Leonurus lanatus.
lanata, Willd. ex Ledeb. Fl. Ross. iii. 439 = Eremostachys laciniata.
larendana, *Boiss. & Heldr. Diagn.* Ser. I. xii. 88.—As. Min.
limbata, Benth. Lab. Gen. et Sp. 743 = Otostegia limbata.
macrodonta, *Boiss. & Bal. Diagn.* Ser. II. iv. 54.—Cappadoc.
mauritanica, Steud. Nom. ed. II. i. 181 = seq.
mauritiana, Pers. Syn. ii. 126 = Anisomeles ovata.
microphylla, Benth. Lab. Gen. et Sp. 596 = Otostegia Schimperi.
mollissima, *Benth. l. c.* 595.—Hispan.
nigra, *Linn. Sp. Pl.* 582.—Europ.; Oriens; Afr. bor.
obliqua, Benth. Lab. Gen. et Sp. 596 = saxatilis.
orbicularis, Lag. ex Willk. & Lange, Prod. Fl. Hisp. ii. 446 = hirsuta.

BALLOTA :—
orientalis, C. & J. Presl, Delic. Prag. 80 (Sp. dubia negligenda, Boiss.).—As. Min.
persica, Benth. Lab. Gen. et Sp. 598 = Otostegia macrophylla.
pilosa, Lour. Fl. Cochinch. 364 = Leucas Benthamiana.
pseudodictamnus, *Benth. Lab. Gen. et Sp.* 594.—Graecia; Creta.
rotundifolia, *C. Koch, in Linnaea*, xxi. (1848) 697.—Armenia.
royleoides, *Benth. in DC. Prod.* xii. 520.—Arab.
rubra, Schrad. ex Steud. Nom. ed. II. i. 182 = nigra.
ruderalis, Sw. in Palmstr. Sv. Bot. t. 389 (1809) = nigra.
rugosa, Benth. Lab. Gen. et Sp. 596 = saxatilis.
rupestris, *Vis. Fl. Dalm.* ii. 216.—Dalmat.
Russeliana, Benth. Lab. Gen. et Sp. 597 = saxatilis.
saxatilis, Guss. Fl. Sic. Syn. ii. 82 = rupestris.
saxatilis, *Sieber, ex Benth. Lab. Gen. et Sp.* 596.—Syria; As. Min.
Schimperi, Benth. in DC. Prod. xii. 519 = Otostegia Schimperi.
sepium, Paulet, in Pers. Syn. ii. 125 = nigra.
silvestris, Hoffmgg. & Link, ex Willk. & Lange, Prod. Fl. Hisp. ii. 446 = alba.
sordida, Salisb. Prod. 83 = nigra.
spinosa, *Link, Handb.* 475.—Gallia.
stachydiformis, *Hochst. in Flora*, xxvii. (1844) 100.—Abyss.
suaveolens, Linn. Syst. ed. X. 1100; Pl. Jam. Pugill. 15 = Hyptis suaveolens.
suaveolens, Rodsch. Obs. 35 = Hyptis pectinata.
undulata, *Benth. Lab. Gen. et Sp.* 595.—Syria; Arab.
urticaefolia, Ortim. ex Reichb. Ic. Bot. viii. 30. t. 774 = nigra.
verticillata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 8.—Malaya.
vulgaris, Hoffmgg. & Link, Fl. Port. t. 115 = nigra.

BALMEDA, Nocca, Hort. Ticin. 21 (1804) = **Grewia**, Linn. (Tiliac.).
corylifolia, Nocca, l. c. = G. corylifolia.

BALMISA, Lag. Gen. et Sp. Nov. 17 (1816) = **Arisarum**, Targ.-Tozz. (Aroid.).
vulgaris, Lag. l. c. = A. vulgare.

BALOGHIA, Endl. Prod. Fl. Norf. 84 (1833). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 300.
STEIGERIA, Muell. Arg. in Linnaea, xxxiv. (1865) 215.
alternifolia, *Baill. Adansonia*, ii. (1861–62) 216.—N. Caled.
carunculata, *Baill. l. c.* 215.—N. Caled.
lucida, *Endl. Prod. Fl. Norf.* 84.—Ins. Norfolc.
Pancheri, *Baill. Adansonia*, ii. (1861–62) 214.—Austral.; N. Caled.

BALOSKION, Rafin. Fl. Tellur. iv. 32 (1836) = **Restio**, Linn.
dichotomum, Rafin. l. c. = R. quadrifarius.

BALSAMARIA, Lour. Fl. Cochinch. ii. 469 (1790) = **Calophyllum**, Linn. (Guttif.).
Inophyllum, Lour. l. c. 470 = C. Inophyllum.

BALSAMEA, Gled. in Berl. Ges. Naturf. Fr. Schr. iii. (1782) 127 = BALSAMODENDRUM, Kunth = **Commiphora**, Jacq. (sec. Engler) (Burser.).
abyssinica, Engl. in Bot. Jahrb. i. (1881) 41 = C. abyssinica.
africana, Baill. Hist. Pl. v. 295 = C. africana.
Agallocha, Baill. l. c. = C. Agallocha.
Berryi, Engl. in Bot. Jahrb. i. (1881) 41 = C. Berryi.
capensis, Engl. l. c. 42 = C. capensis.
caudata, Engl. l. c. = C. candata.
edulis, Baill. in Bull. Soc. Linn. Par. i. (1877) 122 = C. edulis.
Harveyi, Engl. in Bot. Jahrb. i. (1881) 42 = C. Harveyi.
Hildebrandtii, Engl. l. c. = C. Hildebrandtii.
Kotschyi, Engl. l. c. 41 = C. africana.
madagascariensis, Engl. l. c. 42 = C. grandifolia.
meccanensis, Gled. in Berl. Ges. Naturf. Fr. Schr. iii. (1782) 127 = C. Opobalsamum.
mollis, Engl. in Bot. Jahrb. i. (1881) 42 = C. mollis.
Mukul, Baill. Hist. Pl. v. 295 = C. Mukul.
Myrrha, Baill. l. c. 294 = C. Myrrha.
pedunculata, Engl. in Bot. Jahrb. i. (1881) 42 = C. pedunculata.
pilosa, Engl. l. c. 41 = C. pilosa.

BALSAMEA :—
Roxburghii, Engl. in Bot. Jahrb. i. (1881) 41 = C. Agallocha.
Schimperi, Engl. l. c. = C. Schimperi.
Stocksiana, Engl. l. c. = C. Stocksiana.
zanzibarica, Baill. in Adansonia, xi. (1874) 180 = C. zanzibarica.

BALSAMIFLUA, Griff. Notul. iv. 382 (1854) = **Populus**, Linn. (Salicin.).
deltoides, Griff. l. c. = P. euphratica.

BALSAMINA, Tourn. ex Scop. Fl. Carn. ed. II. ii. 183 (1772); Gaertn. Fruct. ii. 151. t. 113 (1791) = **Impatiens**, Linn. (Geraniac.).
angustifolia, Blume, Bijdr. 239 = I. Balsamina.
bifida, DC. Prod. i. 686 = I. flaccida.
capensis, DC. l. c. = I. capensis.
carnea, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 7.—Malaya.
chinensis, DC. Prod. i. 686 = I. chinensis.
coccinea, DC. l. c. 685 = I. Balsamina.
cochleata, DC. l. c. 686 = I. cochleata.
communis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 7.—Malaya.
comorensis, Boj. ex Baker, in Journ. Linn. Soc. xx. (1884) 114 = I. comorensis.
cornuta, DC. Prod. i. 686 = I. Balsamina.
cristata, Ser. Cat. Gr. Jard. Lyon, (1853) = I. scabrida?
fasciculata, DC. Prod. i. 686 = I. chinensis.
foeminea, Gaertn. Fruct. ii. 113 = I. Balsamina.
fruticosa, Leschen. ex Steud. Nom. ed. II. i. 182 = I. fruticosa.
fulva, Ser. Fl. Jard. iii. 453 = I. fulva.
glandulifera, Ser. Cat. Gr. Jard. Lyon, (1853) = I. glandulifera.
glandulifera, Lyall, ex Baker, in Journ. Linn. Soc. xx. (1884) 113 = I. Lyallii.
heterophylla, G. Don, Gen. Syst. i. 749 = I. chinensis.
hirsuta, Blume, Bijdr. 240 = I. hirsuta.
hortensis, Desp. in Dict. Sc. Nat. iii. 485 = I. Balsamina.
javensis, Blume, Bijdr. 240 = I. javensis.
lacca, Medic. Malv. 71 = I. Balsamina.
latifolia, DC. Prod. i. 686 = I. latifolia.
Leschenaultii, DC. l. c. = I. Leschenaultii.
longicornis, Ser. Fl. Jard. iii. 452 = I. leptoceras?
lutea, Delarb. Fl. Auv. ed. II. 504 = I. Noli-tangere.
macrochila, Ser. Fl. Jard. iii. 454 = I. Roylei.
madagascariensis, DC. Prod. i. 686 = I. madagascariensis.
Mastersiana, Paxt. Bot. Mag. vi. 75 = I. Mastersiana.
micrantha, Blume, Bijdr. 240 = I. leptoceras.
minor, DC. Prod. i. 686 = I. Kleinii.
minutiflora, Span. in Linnaea, xv. (1841) 185 = I. Balsamina.
mollis, G. Don, Gen. Syst. i. 749 = I. Balsamina.
mutila, DC. Prod. i. 686 = I. mutila.
mysorensis, DC. l. c. = I. mysorensis.
Noli-tangere, Scop. Fl. Carn. ed. II. ii. 184 = I. Noli-tangere.
odorata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 203 = I. Balsamina.
oppositifolia, DC. Prod. i. 686 = I. oppositifolia.
parviflora, Ser. Cat. Gr. Jard. Lyon, (1853) = I. parviflora?
puberula, G. Don, Gen. Syst. i. 749 = I. puberula.
racemosa, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 203 = I. Balsamina.
repens, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 8.—Malaya.
rosmarinifolia, DC. Prod. i. 686 = I. oppositifolia.
Roylei, Ser. Fl. Jard. iii. 455. t. 7 = I. Roylei.
salicifolia, Boj. ex Baker, in Journ. Linn. Soc. xx. (1884) 115 = I. salicifolia.
scabriuscula, G. Don, Gen. Syst. i. 748 = I. scabriuscula.
setacea, Coleb. ex Hook. Exot. Fl. ii. t. 137 = I. chinensis.
Tilo, DC. Prod. i. 686 = I. tripetala?
tricornis, Ser. Fl. Jard. iii. 452 = I. scabrida.

BALSAMITA, Desf. in Act. Soc. Hist. Nat. Par. i. (1792) t. 1 = **Chrysanthemum**, Linn. (Compos.).
ageratifolia, Desf. l. c. 2 = C. flosculosum.
annua, DC. Fl. Fr. iv. 187 = Tanacetum annuum.
Audibertii, Req. in Ann. Sc. Nat. Sér. I. v. (1825) 382 = Tanacetum Audiberti.
corymbosa, Salzm. in Flora, iv. (1821) 112 = C. flosculosum.
dentata, Wall. Cat. n. 326 = Myriactis Wallichii.

BALSAMITA :—
flabelliformis, Pers. Syn. ii. 408 = Pentzia flabelliformis.
grandiflora, Desf. in Act. Soc. Hist. Nat. Par. i. (1792) 1 (= *Plagius grandiflorus*).—Afr. bor.
major, Desf. l. c. 8 = C. Balsamita.
multifida, Roxas Clem. Ensay. Vid, 292 = Tanacetum annuum.
suaveolens, Pers. Syn. ii. 408 = C. Balsamita.
tridentata, Delile, Fl. Egypte, 273 = Cotula tridentata.
virgata, Desf. in Act. Soc. Hist. Nat. Par. i. (1792) 2 = C. discoideum.
vulgaris, Willd. Sp. Pl. iii. 1802 = Chrysanthemum Balsamita.

BALSAMOCARPON, Clos, in C. Gay, Fl. Chil. ii. 226. t. 20 (1846) = **Caesalpinia**, Linn. (Legumin.).
brevifolium, Clos, l. c.—Chili.

BALSAMODENDRON, DC. Prod. ii. 76 (1825) = seq.

BALSAMODENDRUM, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 348 = **Commiphora**, Jacq. (Burser.).
abyssinicum, Engl. in Bot. Jahrb. i. (1881) 41 = C. abyssinica.
africanum, Arn. in Ann. Nat. Hist. iii. (1839) 87 = C. africana.
Agallocha, Wight & Arn. l. c. 86 = C. Agallocha.
Arnottianum, March. in Adansonia, vii. (1866–67) 266 = C. pubescens.
Berryanum, Voigt, Hort. Suburb. Calc. 149 = C. Berryi.
Berryi, Arn. in Ann. Nat. Hist. iii. (1839) 86 = C. Berryi.
capense, Sond. in Harv. & Sond. Fl. Cap. i. 526 = C. capense.
caudatum, March. in Adansonia, vii. (1866–67) 266 = C. caudata.
Ehrenbergianum, O. Berg, in Bot. Zeit. xx. (1862) 163 = C. Opobalsamum.
gileadense, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 349 = C. Opobalsamum.
habessinica, O. Berg, in Bot. Zeit. xx. (1862) 161 = C. abyssinica.
Kafal, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 349 = C. abyssinica.
Kataf, Kunth, l. c. = C. Kataf.
Kotschyi, O. Berg, in Bot. Zeit. xx. (1862) 162 = C. africanum.
madagascariense, March. in Adansonia, viii. (1867–68) 71. t. 6 = C. grandifolia.
molle, Oliver, Fl. Trop. Afr. i. 326 = C. mollis.
Mukul, Hook. ex Stocks, in Hook. Kew Journ. i. (1849) 259 = C. Mukul.
Myrrha, T. Nees, Pl. Officin. t. 182 = C. Myrrha.
Opobalsamum, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 348 = C. Opobalsamum.
parvifolium, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 506.—Ins. Socotra.
pedunculatum, Kotschy & Peyr. Pl. Tinn. 11. t. 5; et Schweinf. Beitr. Fl. Aeth. 234 = C. pedunculata.
planifrons, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 506.—Ins. Socotra.
Playfairii, Hook. f. ex Oliver, Fl. Trop. Afr. i. 326 = C. Myrrha.
pubescens, Stocks, in Journ. As. Soc. Bomb. ii. (1848) 395. t. 25 = C. Stocksiana.
Roxburghianum, March. ex Baill. Adansonia, vii. (1866–67) 266 = C. caudata.
Roxburghii, Arn. in Ann. Nat. Hist. iii. (1839) 86 = C. Agallocha.
Roxburghii, Stocks, in Journ. As. Soc. Bomb. ii. (1848) 391 = C. Mukul.
Schimperi, O. Berg, in Bot. Zeit. xx. (1862) 162 = C. Schimperi.
socotranum, Balf. f. in Proc. Roy. Soc. Edinb. xi. (1882) 505.—Ins. Socotr.
Wightii, Arn. in Ann. Nat. Hist. iii. (1839) 86 = C. Mukul.
zeylanicum, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 349 = Canarium zeylanicum.

BALSAMONA, Vand. Fasc. Pl. 15 (1771) = **Cuphea**, P. Br. (Lythrar.).
Pinto, Vand. l. c. = C. Balsamona.

BALSAMOPHLEOS, O. Berg, in Bot. Zeit. xx. (1862) 163 = **Commiphora**, Jacq. (Burser.).
Kataf, O. Berg, l. c. = C. Kataf.

BALSAMORHIZA, Hook. Fl. Bor. Am. i. 310 (1833). *COMPOSITAE*, Benth. & Hook. f. ii. 366.
Espeletia, Nutt. in Journ. Acad. Philad. vii. (1834) 37. t. 4.
Bolanderi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 356.—Calif.
Careyana, *A. Gray, Pl. Fendl.* 81.—Am. bor. occ.
deltoidea, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 351.—Am. bor. occ.
glabrescens, Benth. Pl. Hartw. 317 = deltoidea.
helianthoides, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 351 = sagittata.
hirsuta, *Nutt. l. c.* 349.—Am. bor. occ.
Hookeri, *Nutt. l. c.*—Am. bor. occ.
incana, Nutt. l. c. 350 = Hookeri.
macrophylla, *Nutt. l. c.*—Am. bor. occ.
sagittata, *Nutt. l. c.*—Am. bor. occ.
silphioides, Geyer, ex Hook. Lond. Journ. Bot. vi. (1847) 244 = Silphium laeve.
terebinthacea, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 349.—Am. bor. occ.

BALSAMUS, Stackh. De Liban. 11 (1814) = **Commiphora**, Jacq. (Burser.).
Kafal, Stackh. l. c. 12 = C. abyssinica.
libanotus, Stackh. l. c. 11 = C. Opobalsamum?
meccanensis, Stackh. l. c. 12 = C. Opobalsamum.
Theophrasti, Stackh. l. c. 11 = C. Opobalsamum.

BALTIMORA, Linn. Mant. 158 (1771). *COMPOSITAE*, Benth. & Hook. f. ii. 348.
Fougeria, Moench, Meth. Suppl. 243 (1802).
Fougerouxia, Cass. in Dict. Sc. Nat. liv. 461 (1829).
Niebuhria, Scop. Introd. 134 (1777).
Scolospermum, Less. in Linnaea, v. (1830) 152.
Timanthea, Salisb. Prod. 208 (1796).
alata, Meerb. Afb. t. 23 = recta?
alba, Pers. Syn. ii. 489 = recta.
recta, *Linn. Mant.* 288.—Mexic.
scabra, Pohl, ex Baker, in Mart. Fl. Bras. vi. iii. 258 = Calea hymenolepis.
scolospermum, *Steetz, in Seem. Bot. Voy. Herald*, 154.—Panama.
trinervata, Moench, Meth. 592 = recta.

BAMBOS, Retz. Obs. v. 24 (1789) = **Bambusa**, Schreb. (Gramin.).
arundinacea, Retz. Obs. v. 24 = Bambusa arundinacea.
Arundo, Soland. Fl. Ins. Ocean. Pacif. 217 = Schizostachyum glaucifolium.
quinqueflora, Stokes, Bot. Mat. Med. ii. 286 = Bambusa arundinacea.
stricta, Roxb. Pl. Corom. i. 58. t. 80 (non text.) = Oxytenanthera Thwaitesii, et Dendrocalamus strictus.
verticillata, Poir. Encyc. viii. 703 = Gigantochloa verticillata.

BAMBUS, J. F. Gmel. Syst. 579 (1791) = seq.

BAMBUSA, Schreb. Gen. Pl. 236 (1789). *GRAMINEAE*, Benth. & Hook. f. iii. 1210.
Bambos, Retz. Obs. v. 24 (1789).
Bambus, J. F. Gmel. Syst. 579 (1791).
Beesha, Kunth, Syn. i. 253 (1822).
Ischurochloa, Buese, in Miq. Pl. Jungh. 389 (1854).
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 439 = Oxytenanthera abyssinica.
affinis, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 93.—Burma.
agrestis, *Poir. Encyc.* viii. 704.—Ind. or.; Cochinch.
alpina, Bory, Voy. i. 310. t. 12 = Nastus borbonicus.
amahussana, *Lindl. in Penny Cyclop.* iii. 357 (1835).—Amboina.
amplexifolia, Schult. f. Syst. vii. 1348 = Guadua amplexifolia.
andamanica, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 88.—Ins. Andam.
angulata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 94.—China.
Apus, *Schult. f. Syst.* vii. ii. 1353.—Java.
argentea-striata, *Regel, Gartenfl.* (1865) 363.—Japon.
aristata, *Lodd. & Lindl. in Penny Cyc.* iii. (1835) 357.—Ind. or.
arundinacea, Humb. & Bonpl. Pl. Aequin. i. 71 = Nastus borbonicus.

BAMBUSA :—
arundinacea, Ait. Hort. Kew. ed. II. ii. 316 = Thouarsii.
arundinacea, Wight, ex Steud. Nom. ed. II. i. 183 = orientalis.
arundinacea, *Willd. Sp. Pl.* ii. 245.—Ind. or.
Arundo, Nees, in Linnaea, ix. (1834) 471 = arundinacea.
Arundo, Wight, ex Steud. Nom. ed. II. i. 183 = orientalis.
aspera, *Schult. f. Syst.* vii. ii. 1352.—Amboina.
atra, *Lindl. in Penny Cyc.* iii. (1835) 357.—Amboina.
attenuata, Thw. Enum. Pl. Zeyl. 375 = Teinostachyum attenuatum.
aurea, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 716 = Phyllostachys aurea.
aurea, *Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 285.—Japon.
aureo-striata, *Regel, Gartenfl.* (1865) 362.—Japon.
auriculata, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 86 = vulgaris.
baccifera, Roxb. Hort. Beng. 25; Fl. Ind. ii. 197 = Melocanna bambusoides.
Balcooa, *Roxb. ll. cc.* 25, 196.—Ind. or.
barbata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* i. (1835) 627 = Nastus barbatus.
Beecheyana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 108.—China.
bifolia, Siebold, ex Munro, l. c. 36 = Phyllostachys bambusoides.
Bitung, *Schult. f. Syst.* vii. ii. 1354.—Java.
Blancoi, *Steud. Syn. Pl. Gram.* 331.—Ins. Philipp.
Blumeana, Hook. & Arn. Bot. Beech. Voy. 254 = tuldoides.
Blumeana, *Schult. f. Syst.* vii. ii. 1343.—Java.
Brandisii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 109.—Burma.
breviflora, *Munro, l. c.* 96.—China.
caesia, Sieb. & Zucc. ex Munro, l. c. 89 = nana.
calostachya, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 250.—Burma.
Cantori, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 111.—China.
capensis, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 144.—Afr. austr.
capitata, Trin. l. c. Sc. Nat. i. (1835) 626 = Guadua capitata.
capitata, Wall. & Griff. in Wall. Cat. n. 8913 = Cephalostachyum capitatum.
capitata, *Willd. ex Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 43.—Madag.
China, *Franch. & Sav. Enum. Pl. Jap.* ii. 607.—Japon.
Chusque, Poir. Encyc. Suppl. v. 494 = Chusquea scandens.
cornuta, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 113.—Java.
crinita, Thoms. ex Munro, l. c. 157 = nutans.
critica, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 250.—Pegu.
dichotoma, Donn, Hort. Cantab. ed. IV. 78 = Phyllostachys nigra.
diffusa, Blanco, Fl. Filip. ed. I. 269 = Guadua distorta.
distorta, *Nees, in Linnaea*, ix. (1834) 470.—Bras.
diversistachya, Munro, in Trans. Linn. Soc. xxvi. (1868) 126 = Gigantochloa heterostachya.
dumetorum, Hance, ex Walp. Ann. iii. 781 = Schizostachyum dumetorum.
Duquilioi, Hort. ex Carr. in Rev. Hortic. (1869) 292 = Phyllostachys Quilioi.
edulis, *Carr. l. c.* (1866) 380.—China.
elegantissima, Hassk. Pl. Jav. 42 = Beesha elegantissima.
excelsa, Miq. Fl. Ind. Bat. iii. 418 = maxima.
falcata, Hort. ex Vilm. Fl. Pl. Terre. ed. II. 98 = Arundinaria falcata.
Falconeri, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 95.—Reg. Himal.
Fax, Poir. Encyc. viii. 704 = Melocanna humilis.
fera, *Miq. Fl. Ind. Bat.* iii. 418.—Amboina.
fistulosa, Royle, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 95 = Falconeri.
flagellifera, Griff. ex Munro, l. c. 150 = Dendrocalamus flagellifer.
flexuosa, *Munro, l. c.* 101.—China.
flexuosa, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 758 = Phyllostachys flexuosa.

BAMBUSA:—
floribunda, *Zoll. & Mor. ex Steud. Syn. Pl. Gram.* 330. —Java.
floribunda, Munro, in Trans. Linn. Soc. xxvi. (1868) 34 = Thamnocalamus Falconeri.
Fortunei, *Van Houtte, Fl. des Serres*, xv. (1863) t. 1535.—Japon.
fragilis, Spruce, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 78 = Guadua latifolia.
gigantea, Wall. Cat. Bot. Gard. Calc. 79 = Dendrocalamus giganteus.
glauca, Blume, ex Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 162 = Arundinaria glaucescens.
glauca, Lodd. ex Lindl. in Penny Cyclop. iii. 357 (1835) = nana.
glaucescens, Siebold, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 89 = nana.
glaucifolia, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 147 = Schizostachyum glaucifolium.
globifera, *Griseb. in Goett. Nachr.* (1868) 72.—Ind. or.
glomerata, Royle, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 147 = Dendrocalamus strictus.
gracilis, *Hort. ex A. & C. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 682.—Hab.?
gracilis, Wall. Cat. n. 5033 = Oxytenanthera nigro-ciliata.
Griffithiana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 99.—Burma.
Guadua, Humb. & Bonpl. Pl. Aequin. 168. t. 20 = Guadua angustifolia.
Helferi, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 114. —Burma.
heterocycla, *Carr. in Rev. Hortic.* (1878) 354.—Hab.?
hexandra, Roxb. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 147 = Dendrocalamus strictus.
Hookeri, *A. & C. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 642.—Hab.?
Horsfieldii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 115.—Java.
humilis, Reichb. ex Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 140 = vulgaris.
khasiana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 97. —Reg. Himal.
kumasasa, Zoll. ex Steud. Syn. Pl. Gram. 331 = Phyllostachys kumasaca.
latiflora, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 250.—Ind. or.
latifolia, Humb. & Bonpl. Pl. Aequin. i. 68. t. 21 = Guadua latifolia.
levis, *Blanco, Fl. Filip.* ed. I. 272.—Ins. Philipp.
Lima, *Blanco, l. c.* 271.—Ins. Philipp.
lineata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 118. —Ins. Ternate.
longinodis, *Miq. Fl. Ind. Bat.* iii. 418.—Malaya.
longispatha, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 250.—Burma.
Luçoniae, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 115.—Ins. Philipp.
Lumampao, *Blanco, Fl. Filip.* ed. I. 272.—Ins. Philipp.
Mac-Clellandi, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 114.—Burma.
macra, Wall. ex Munro, l. c. 34 = Thamnocalamns spathiflorus.
macroculmis, *A. Rivière, in Bull. Soc. Acclim.* Sér. III. v. (1878) 624.—Hab.?
Macala, Buch.-Ham. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 91 = Tulda.
madagascariensis, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 631 = vulgaris.
marginata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 114.—Burma.
Mastersii, *Munro, l. c.* 113.—Ind. or.
maxima, Buch.-Ham. in Wall. Cat. n. 5039 = Dendrocalamus Hamiltonii.
maxima, *Poir. Encyc.* viii. 704.—Amboin.; China.
Maximowiczii, Hort. ex Munro, in Gard. Chron. (1876) II. 774 = Arundinaria Fortunei.
Metake, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 284 = Arundinaria japonica.
microphylla, Griff. Journ. i. 242, 259 (1847) = Arundinaria microphylla.
mitis, Blanco, Fl. Filip. ed. I. 271 = Blancoi.
mitis, Hort. Paris. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 18 = Arundinaria japonica.
mitis, *Poir. Encyc.* viii. 704.—China.
mitis, Hort. ex Carr. in. Rev. Hortic. (1866) 380 = edulis.

BAMBUSA:—
monogyna, *Blanco, Fl. Filip.* ed. I. 268.—Ins. Philipp.
monogynia, Griff. Notul. iii. 63 = Dendrocalamus Hamiltoni.
multiplex, *Raeusch. Nom.* ed. III. 103; *Schult. f. Syst.* vii. II. 1350.—Cochinch.
nana, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 199.—Japon.
Neesiana, Arn. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 103 = arundinacea.
nigra, Lodd. Cat. (1823) ex Loud. Hort. Brit. 124 = Phyllostachys nigra.
nigricans, Hort. ex Steud. Nom. ed. II. i. 183 = Phyllostachys nigra.
nigro-ciliata, Buese, in Miq. Pl. Jungh. 389 = Oxytenanthera nigro-ciliata.
nutans, *Wall. ex Munro, in Trans. Linn. Soc.* xxvi. (1868) 92.—Reg. Himal.
Oldhami, *Munro, l. c.* 109.—China.
orientalis, *Nees, in Linnaea*, ix. (1834) 475.—Ind. or.
pallida, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 97. —Ind. or.
paniculata, Willd. ex Stend. Nom. ed. II. i. 183 = Nastus borbonicus?
parviflora, Schult. f. Syst. vii. II. 1350 = Arthrostylidium maculatum.
picta, *Lindl. in Penny Cyclop.* iii. 357 (1835).—Ins. Ceram.
picta, Sieb. & Zucc. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 111 = Fortunei.
polymorpha, *Munro, l. c.* 98.—Burma.
prava, *Lindl. in Penny Cyclop.* iii. 357 (1835).—Amboina.
pseudoarundinacea, Steud. Syn. Pl. Gram. 330 = Gigantochloa verticillata.
puberula, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 285 = Phyllostachys nigra.
pubescens, *Doell, in Mart. Fl. Bras.* ii. III. 189.—Bras.
pubescens, Lodd. ex Lindl. Penny Cyclop. iii. 357 (1835) = Dendrocalamus strictus.
pungens, Blanco, Fl. Filip. ed. I. 270 = arundinacea.
pygmaea, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 286.—Japon.
quadrangularis, *Fenzi, in Bull. Soc. Tosc. Ort.* v. (1880) 401.—Japon.
Quilioi, Hort. ex A. & C. Rivière, in Bull. Soc. Acclim. Sér. III. v. (1878) 697 = Phyllostachys Quilioi.
Ragamowskii, Hort.—Cf. Gard. Chron. (1876) II. 847 = tessellata.
regia, *Thoms. Cat. Hort. Bot. Calc.* 79; *ex Munro, in Trans. Linn. Soc.* xxvi. (1868) 116.—Ind. or.
reticulata, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 148 = Phyllostachys bambusoides.
Ritchcyi, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 113.—Ind. or.
Rumphiana, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) II. 86.—Ind. or.
ruscifolia, Siebold, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 157 = Phyllostachys kumasaca.
scandens, Blume, ex Nees, in Flora, vii. (1824) 291 = Dendrocalamus Tjankorreh.
Schimperiana, Steud. Syn. Pl. Gram. 330 = Oxytenanthera abyssinica.
schizostachyoides, Kurz, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 134 = Melocanna Kurzii.
scriptoria, Dennst. Schluess. Hort. Malab. 31 = Melocanna bambusoides.
senamensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 606.—Japon.
siamensis, Kurz, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 116 = regia.
Sieberi, Griseb. Fl. Brit. W. Ind. 528 = vulgaris.
Simoni, Carr. in Rev. Hortic. (1866) 380 = Arundinaria Simoni.
spinosa, Nees, in Linnaea, ix. (1834) 475 = arundinacea.
spinosa, Blume, ex Nees, in Flora, viii. (1825) 580 = Blumeana.
spinosa, *Roxb. Hort. Beng.* 25; *Buch.-Ham. in Trans. Linn. Soc.* xiii. (1822) 480.—Ind. or.
sterilis, Kurz, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 285 = nana.
striata, *Lodd. ex Lindl. Penny Cyclop.* iii. 357 (1835). —China.
stricta, Roxb. Hort. Beng. 25; Fl. Ind. ii. 193 = Oxytenanthera Thwaitesii et Dendrocalamus strictus.
stridula, *Moon. Cat.* 26.—Zeylan.
sulfurea, Carr. in Rev. Hortic. (1873) 379 = Phyllostachys sulphurea.

BAMBUSA:—
surinamensis, Rupr. in Mém. Acad. Pétersb. Sér. VI. v. (1839) 49 = vulgaris.
tabacaria, *Poir. Encyc.* viii. 705.—Amboina; Java.
Tagoara, Nees, Agrost. Bras. 532 = Guadua Tagoara.
Teba, *Miq. Fl. Ind. Bat.* iii. 418.—Malaya; China.
tenuis, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 119. —Ins. Macass.
teres, *Buch.-Ham. in Wall. Cat.* n. 5026 B.—Ind. or.
tessellata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 110.—China.
textoria, *Blanco, Fl. Filip.* ed. I. 270.—Ins. Philipp.
Thouarsii, Kunth, Not. Gen. Bambusa, 3 = vulgaris.
trigyna, Roxb. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 91 = Tulda.
Trinii, Nees, in Linnaea, ix. (1834) 469 = Gnadua Trinii.
Tulda, Benth. Fl. Hongk. 434 = tuldoides.
Tulda, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 193.—Ind. or.; Burma.
tuldoides, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 93. —China.
variegata, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 285 = Fortunei.
vasaria, Buch.-Ham. in Wall. Cat. n. 5025 = Balcooa.
vasaria, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 122. —Malaya.
verticillata, Benth. Fl. Hongkong. 434 = Dendrocalamus latiflorus.
verticillata, Rottler, ex Munro, in Trans. Linn. Soc. xxvi. (1868) 147 = Dendrocalamus strictus.
verticillata, Hook. & Arn. Bot. Beech. Voy. 254 = Beecheyana.
verticillata, Willd. Sp. Pl. ii. 245 = Gigantochloa verticillata.
villosula, *Kurz, For. Fl. Brit. Burma*, ii. 553.—Burma.
violascens, Carr. in Rev. Hortic. (1869) 292 = Phyllostachys violascens.
virgata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1835) 624 = Guadua virgata.
viridi-glaucescens, Carr. in Rev. Hortic. (1869) 292 = nana.
viridi-striata, Regel, Ind. Sem. Hort. Petrop. (1866) 77 = striata.
vulgaris, Nees, Fl. Afr. Austr. 462 = Balcooa.
vulgaris, *Schrad. in Wendl. Coll. Pl.* ii. 26. t. 47. —Mexic.
Wightii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 111.—Ind. or.

BAMIA, [Linn.] R. Br. ex Wall. Pl. Asiat. Rar. i. 39 (1830) = **Hibiscus**, Tourn. (Malvac.).
Abelmoschus, R. Br. ex Wall. Cat. n. 1915 = H. Abelmoschus.
angulosa, Wall. Cat. n. 1927 = H. angulosus.
betulifolia, Wall. l. c. 1918 = H. Abelmoschus.
cancellata, Wall. l. c. 1920 = H. cancellatus.
chinensis, Wall. l. c. 1916 = H. Abelmoschus?
crinita, Wall. l. c. 1922 = H. cancellatus.
fusiformis, Wall. l. c. 1921 = H. cancellatus.
magnifica, Wall. l. c. 1919 = H. hostilis.
Manihot, Wall. l. c. 1926 = H. Manihot.
multiformis, Wall. l. c. 1917 = H. Abelmoschus.
pungens, Wall. l. c. 1924 = H. pungens.
rugosa, Wall. l. c. 1923 = H. rugosus.
tetraphylla, Wall. l. c. 1925 = H. tetraphyllus.

BAMMIA, Rupp. Fl. Jen. ed. Hall. 39 (1745) = praec.

BANALIA, Moq. in DC. Prod. xiii. II. 278 (1849). *AMARANTACEAE*, Benth. & Hook. f. iii. 27.
brasiliana, Moq. l. c. (= Chamissoae sp.?).—Bras.
occidentalis, Moq. l. c. 279 = Nitrophila occidentalis.
thyrsiflora, *Moq. l. c.* 278.—Ind. or.

BANARA, Aubl. Pl. Gui. i. 547. t. 217 (1775). *SAMYDACEAE*, Benth. & Hook. f. i. 798.
ASCRA, Schott, in Spreng. Syst. iv. Cur. Post. 407 (1827).
BOCA, Vell. Fl. Flum. 232 (1825); v. t. 113 (1827).
CHRISTANNIA, Presl, Rel. Haenk. ii. 91. t. 67 (1830).
PINEDA, Ruiz & Pav. Prod. 76. t. 14 (1794).
XYLADENIUS, Desv. in Ham. Prod. Fl. Ind. Occ. 41 (1825).
brasiliensis, *Benth. in Journ. Linn. Soc.* v. (1861). *Suppl.* ii. 93.—Bras.

BANARA:—
brevifolia, Blanco, Fl. Filip. ed. I. 426 = Scolopia dasyanthera.
dioica, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 94—Mexic.
domingensis, *Benth. l. c.*—Ins. S. Doming.
fagifolia, Vahl, Symb. Bot. iii. 65 = guianensis.
glaberrima, *Wright, ex Griseb. Cat. Pl. Cub.* 8.—Cuba.
glauca, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 92 = Kuhlia glauca.
grandiflora, *Spruce, ex Benth. l. c.* 93.—Peruv.
guianensis, *Aubl. Pl. Gui.* i. 548. t. 217.—Guiana; Bras.
ibaguensis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 291.—N. Granat.
incana, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 94.—Peruv.
iaxiflora, *Benth. l. c.* 91.—Peruv.
mexicana, *A. Gray, in Proc. Am. Acad.* v. (1861) 174.—Mexic.
mollis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 290.—N. Granat.
nitida, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) *Suppl.* ii. 93.—Peruv.
parviflora, *Benth. l. c.* 91.—Bras.
pubescens, *Spruce, ex Benth. l. c.* 92.—Ecuador.
racemosa, *Blanco, Fl. Filip.* ed. I. 425.—Ins. Philipp.
reticulata, *Griseb. Cat. Pl. Cub.* 8.—Cuba.
Roxburghii, *Spreng. Syst.* ii. 479 (omnino dubia).—Ind. or.
tomentosa, *Clos, in Ann. Sc. Nat.* Sér. IV. viii. (1857) 240.—Bras.
ulmifolia, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 91 = Kuhlia ulmifolia.
Vellozii, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 331.—Bras.

BANCROFFTIA, Steud. Nom. ed. II. i. 183 (1840), = seq.

BANCROFTIA, Billb. in Linneska Samfund. Handl. i. 1832 (1833) 39 = **Arracacia**, Bancr. (Umbell.).
moschata, Billb. l. c. = A. moschata.
xanthorrhiza, Billb. l. c. = A. xanthorrhiza.

BANCROFTIA, Macfad. Fl. Jamaic. i. 112 (1837) = **Tovaria**, Ruiz & Pav. (Capparid.).
diffusa, Macfad. l. c. = T. pendula.

BANDEIRAEA, Welw. ex Benth. & Hook. f. Gen. i. 577 (1865). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
Griffonia, Baill. in Adansonia, vi. (1865) 188. t. 2.
simplicifolia, *Benth. in Trans. Linn. Soc.* xxv. (1866) 306, in nota.—Afr. trop.
speciosa, *Welw. ex Benth. l. c.* 306. t. 40.—Afr. trop.
tenuiflora, *Benth. l. c.* 307.—Afr. trop.

BANDEREIA, Baill. Hist. Pl. ii. 120 (1870) = **Bandeiraea**, Welw. (Legum.).

BANDURA, Burm. Thes. Zeyl. 42. t. 17 (1737) = **Nepenthes**, Linn.

BANFFYA, Baumg. Enum. Stirp. Transs. i. 385 (1816) = **Gypsophila**, Linn. (Caryophyll.).
petraea, Baumg. l. c. = G. transsylvanica.

BANGLIUM, Buch.-Ham. ex Wall. Cat. sub n. 6579 (1832) = **Alpinia**, Linn. (Scitam.).
sulphureum, Buch.-Ham. l. c. = A. Hamiltoniana.

BANIA, Becc. Malesia, i. (1877) 162. *MENISPERMACEAE*.
thyrsiflora, *Becc. l. c.*—N. Guinea.

BANISTERIA, Linn. Syst. ed. II. 23 (1740). *MALPIGHIACEAE*, Benth. & Hook. f. i. 257.
Anomalopteris, G. Don, Gen. Syst. i. 647 (1831).
Holopetalon, Reichb. Nom. 207 (1841).
acanthocarpa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 394.—Am. austr.
aculeata, *Mill. Gard. Dict.* ed. VIII. n. 6.—Mexic.
adamantium, *Mart. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 406.—Bras.
adenopoda, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 40.—Bras.

BANISTERIA:—
albicana, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 131.—Bras.
albida, Willd. ex Steud. Nom. ed. II. i. 183 = Gaudichaudia albida.
allophila, Sieber, ex A. Juss. in Arch. Mus. Par. iii. (1843) 356 = Brachypterys borealis.
amoena, Mart. ex A. Juss. l. c. 571 = Hiraea Gaudichaudiana.
angulata, Vell. Fl. Flum. 191; iv. t. 160 = Stigmaphyllon affine.
angulosa, Linn. Sp. Pl. 427 = Stigmaphyllon angulosum.
angustifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 38.—Bras.
anisandra, Griseb. in Linnaea, xiii. (1839) 190 = Clausseniana.
anisandra, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 47.—Bras.
antifebrilis, *Ruiz, ex Griseb. in Linnaea*, xxii. (1849) 15.—Peruv.
appendiculata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 413.—Ins. S. Vincent.
appendiculata, Lamb. ex DC. Prod. i. 592 = Heteropteris appendiculata.
argentea, *Spreng. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 393.—Am. austr.
argyrophylla, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 45.—Bras.
atrosanguinea, *A. Juss. in Arch. Mus. Par.* iii. (1843) 389.—Bolivia.
aurea, Vell. Fl. Flum. 192; iv. t. 167 = Gardneriana.
auriculata, Cav. Diss. ix. 428. t. 255 = Stigmaphyllon auriculatum.
benghalensis, Linn. Sp. Pl. 427 = Hiptage Madablota.
Benthamiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 390.—Peruv.
bicolor, *Poepp. ex A. Juss. l. c.* 393.—N. Granat.
Billbergiana, *Beurling, in Vet. Akad. Handl. Stockh.* (1854) 118.—Panama.
Blanchetiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 420.—Am. austr.
bogotensis, Bonpl. ex Triana & Planch. Prod. Fl. N. Granat. i. 318 = Stigmaphyllon bogotense.
bonariensis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 157.—Reg. Argent.
brachiata, Linn. Sp. Pl. 428 = Heteropteris brachiata.
brachiata, Spreng. Syst. ii. 387 = Heteropteris platyptera.
brachiata, Willd. ex A. Juss. in Arch. Mus. Par. iii. (1843) 356 = Brachypterys borealis.
brachyptera, DC. Prod. i. 591 = Brachypterys borealis.
bracteata, DC. l. c. 590 = Tetrapteris inaequalis.
brevipes, Sesse, ex DC. Prod. i. 591 = Gaudichaudia brevipes.
Caapi, *Spruce, ex Griseb. in Mart. Fl. Bras.* xii. III. 43.—Bras.
caerulea, Lam. Encyc. i. 367 = Heteropteris caerulea.
Calcitrapa, Desv. in Ham. Prod. Fl. Ind. Occ. 40 = Brachypterys borealis.
calocarpa, *Miq. in Linnaea*, xviii. (1844) 53.—Guiana.
campestris, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 36.—Bras.
chrysophylla, Lam. Encyc. i. 368 = Heteropteris chrysophylla.
ciliata, Lam. l. c. 369 = Stigmaphyllon ciliatum.
cinerascens, *Griseb. in Linnaea*, xxii. (1849) 18.—Guiana.
Clausseniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 422.—Bras.
coccinea, *Vell. Fl. Flum.* 191. iv. t. 161.—Bras.
comans, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 406 = stellaris.
constricta, *Griseb. in Mart. Fl. Bras.* xii. III. 46.—Bras.
convolvulifolia, Cav. Diss. ix. 428. t. 256 = Stigmaphyllon convolvulifolium.
cordata, Vell. Fl. Flum. 191; iv. t. 163 = rotundifolia.
cordistipula, *Mart. Herb. Fl. Bras.* 556 = Peixotoa cordistipula.
coriacea, *G. Don, ex Griseb. in Mart. Fl. Bras.* xii. III. 94.—Bras.
cornifolia, *Spreng. Syst.* ii. 388.—N. Granat.
corymbosa, *Griseb. in Mart. Fl. Bras.* xii. III. 42, in nota.—Bras.
corymbosa, Griseb. in Linnaea, xxii. (1849) 18 = cinerascens.
cristata, *Griseb. l. c.* 16.—Guiana.

BANISTERIA:—
crotonifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 36.—Bras.
cujabensis, Griseb. in Linnaea, xiii. (1839) 194 = Hiraca cujabensis.
deformis, Desv. in Ham. Prod. Fl. Ind. Occ. 40 = Stigmaphyllon angulosum.
dichotoma, Linn. Sp. Pl. 427 = Stigmaphyllon rotundifolium.
dichotoma, DC. Prod. i. 588 = Stigmaphyllon convolvulifolium.
divaricata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 412.—Guiana.
diversifolia, Bert. ex DC. Prod. i. 588 = variifolia.
diversifolia, *H. B. & K. Nov. Gen. et Sp.* v. 159.—Cuba.
eglandulata, Vell. Fl. Flum. 190; iv. t. 155 = Tetrapteris crebriflora.
eglandulosa, DC. Prod. i. 590 = Heteropteris suberosa.
eglandulosa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 410.—Bras.
elegans, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 322.—N. Granat.
elliptica, H. B. & K. Nov. Gen. et Sp. v. 161 = Stigmaphyllon ellipticum.
emarginata, *Cav. Diss.* ix. 425. t. 249.—Am. austr.
erianthera, *A. Juss. in Arch. Mus. Par.* iii. (1843) 411.—Bras.
fagifolia, DC. Prod. i. 590 = Hiraea fagifolia.
ferruginea, *Cav. Diss.* ix. 424. t. 248.—Bras.; Guian.
ferruginea, Seem. Bot. Voy. Herald, 92 partim = Tetrapteris Seemanni.
fimbriata, *Vell. Fl. Flum.* 192; iv. t. 165.—Bras.
fimbriata, Griseb. in Linnaea, xiii. (1839) 191 = nigrescens.
Fischeriana, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1858) 48.—Cuba.
floribunda, *DC. Prod.* i. 589.—Bras.
fruticosa, Vell. Fl. Flum. 192; iv. t. 164 = Heteropteris umbellata.
fulgens, DC. Prod. i. 590 = Stigmaphyllon puberum.
fulgens, G. F. W. Mey. Prim. Fl. Esseq. 181 = ferruginea.
fulgens, *Linn. Sp. Pl.* 427.—Ins. Guadalup.
fulgens, Lam. Encyc. i. 368 = Stigmaphyllon fulgens.
Gardneriana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 420.—Bras.
glandulosa, Pav. ex A. Juss. l. c. 478 = Heteropteris rubiginosa.
glauca, Desf. Cat. Hort. Par. ed. III. 406 = Stigmaphyllon ciliatum.
glaucophylla, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 319 = Verrucularia glaucophylla.
grata, *Griseb. in Mart. Fl. Bras.* xii. III. 49.—Bras.
heterophylla, Willd. Sp. Pl. ii. 742 = Stigmaphyllon fulgens.
heterostyla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 408.—N. Granat.
hexandra, Vell. Fl. Flum. 188; iv. t. 149 = Schwannia elegans.
Humboldtiana, DC. Prod. i. 588 = Stigmaphyllon Humboldtianum.
hypericifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 42.—Bras.
intermedia, A. Juss. in Arch. Mus. Par. iii. (1843) 404 = stellaris.
jasminellum, *A. Juss. l. c.* 409.—N. Granat.
Kraussiana, *Hochst. in Flora*, xxvii. (1844) i. 296.—Afr. austr.
laevifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 38.—Bras.
lancifolia, West, ex Schlecht. in Linnaea, iv. (1829) 81 = Heteropteris laurifolia.
latifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 40.—Bras.
laurifolia, Linn. Sp. Pl. ed. II. 611 = Heteropteris laurifolia.
laurifolia, Rich. ex A. Juss. in Arch. Mus. Par. iii. (1843) 380 = Stigmaphyllon periplocifolium.
laurifolia, Sieber, ex A. Juss. l. c. 294 = Byrsonima spicata.
ledifolia, *H. B. & K. Nov. Gen. et Sp.* v. 159.—Cuba.
leiocarpa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 395.—Peruv.
leona, *Cav. Diss.* ix. 424 (sp. dubia).—Afr. trop.
leptocarpa, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 130.—Guiana.

BANISTERIA:—
lobulata, *E. Mey. in Nov. Act. Nat. Cur.* xii. II. (1825) 813.—Guiana.
longifolia, Sw. Prod. Veg. Ind. Occ. 75 = Heteropteris platyptera.
lucida, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 109.—Guiana.
lupuloides, Linn. Sp. Pl. 427 = Gouania domingensis.
lutea, *Ruiz, ex Griseb. in Linnaea*, xxii. (1849) 15.—Peruv.
macradena, *DC. Prod.* i. 590.—Guiana.
macrocarpa, Pers. Syn. i. 507 = Heteropteris platyptera.
macrophylla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 418.—Bras.
macrostachia, Vell. Fl. Flum. 189; iv. t. 152 = Clausseniana.
magnoliaefolia, Desv. ex Ham. Prod. Fl. Ind. Occ. 40 = Heteropteris platyptera.
maracabensis, A. Juss. in Arch. Mus. Par. iii. (1843) 418 = cornifolia.
maritima, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Brachypterys borealis.
Martiniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 413.—Guiana.
mediterranea, Vell. Fl. Flum. 191; iv. t. 162 = Janusia muricata.
megacarpos, Vell. l. c. 189; iv. t. 189 = Stigmaphyllon tomentosum.
megaphylla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 399.—Bras.
membranifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 39.—Bras.
metallicolor, *A. Juss. l. c.* 46.—Bras.
microphylla, *Jacq. Obs.* iii. 7. t. 56.—Carolin.
Minarum, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 403 = schizoptera.
monoptera, Vell. Fl. Flum. 189; iv. t. 153 = Heteropteris chrysophylla.
montana, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 37 = campestris.
mucronata, DC. Prod. i. 589 = Stigmaphyllon ellipticum.
multialata, Vell. Fl. Flum. 189; iv. t. 154 = Tetrapteris punctulata.
multiflora, Boj. ex A. Juss. in Arch. Mus. Par. iii. (1843) 424 = Sphedamnocarpus madagascariensis.
multiflora, DC. Prod. i. 589 = Heteropteris platyptera.
multifoliolata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 402.—Bras.
muricata, *Cav. Diss.* ix. 423. t. 243. f. 2.—Peruv.
mutabilis, Vell. Fl. Flum. 193; iv. t. 168 = Thryallis brachystachys.
nigrescens, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 44.—Bras.
nitens, Griseb. in Linnaea, xiii. (1839) 198 = argyrophylla.
nitida, Vell. Fl. Flum. 188; iv. t. 148 = Stigmaphyllon ciliatum.
nitida, Lam. Encyc. i. 369 = Heteropteris nitida.
nitrosiodora, *Griseb. in Goett. Abh.* xxiv. (1879) 65.—Reg. Argent.
nummifera, *A. Juss. in Arch. Mus. Par.* iii. (1843) 391.—Bras.
ochnaefolia, Mart. ex A. Juss. l. c. 464 = Heteropteris byrsonimaefolia.
orinoccensis, *H. B. & K. Nov. Gen. et Sp.* v. 162.—Venezuela.
ovata; Cav. Diss. ix. 429. t. 257. f. 1 = Brachypterys borealis.
oxyclada, *A. Juss. in Arch. Mus. Par.* iii. (1843) 396.—Bras.
Pagua, Steud. Nom. ed. II. i. 184 = Heteropteris syringaefolia.
palmata, *Cav. Diss.* ix. 430. t. 257. f. 2.—Am. austr.
paniculata, DC. Prod. i. 591 = Heteropteris Beccheyana.
parviflora, *A. Juss. in Arch. Mus. Par.* iii. (1843) 416.—Bras.
parvifolia, Vent. Choix, 51 = Heteropteris parvifolia.
patens, Griseb. in Linnaea, xiii. (1839) 200 = Heteropteris patens.
pauciflora, *H. B. & K. Nov. Gen. et Sp.* v. 159.—Cuba.
periplocaefolia, *Desf. Tabl.* ed. II. 160 = Stigmaphyllon retusum.
picta, H. B. & K. Nov. Gen. et Sp. v. 160 = Brachypterys borealis.

BANISTERIA:—
platyptera, *Griseb. in Linnaea*, xxii. (1849) 17.—Venezuela.
podadena, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) I. 392.—Cuba.
polita, *Miq. in Verh. Nat. Wet. Haarl.* vii. (1851) 80. t. 21.—Guiana.
praecox, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 133.—Bras.
Pragua, Vell. Fl. Flum. 190; iv. t. 138 = Heteropteris syringaefolia.
pruinosa, *Mart. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 392.—Bras.
pubera, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 109 = Stigmaphyllon puberum.
pubiflora, DC. Prod. i. 591 = Heteropteris laurifolia.
pubipetala, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 41. t. 169.—Bras.
purpurea, Linn. Sp. Pl. 427 = Heteropteris purpurea.
Quapara, *Aubl. Pl. Gui.* i. 464. t. 186.—Guiana.
Riedeliana, *Regel, Ind. Sem. Hort. Petrop.* (1855) 17.—Cuba.
rigida, *A. Juss. in Arch. Mus. Par.* iii. (1843) 416.—Bras.
sagittata, *Cav. Diss.* ix. 430. t. 257.—Ins. S. Doming.
salicifolia, *DC. Prod.* i. 590.—Bras.
schizoptera, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 43.—Bras.
Schlimii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 584.—N. Granat.
Schomburgkiana, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 129.—Guiana.
schwannioides, *Griseb. in Mart. Fl. Bras.* xii. III. 54.—Bras.
scutellata, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 131.—Bras.
Sellowiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 419.—Bras.
sepium, *Mart. ex A. Juss. l. c.* 414.—Bras.
sericea, Cav. Diss. ix. 429. t. 258 = Heteropteris sericea.
sericea, Dombey, ex A. Juss. in Arch. Mus. Par. iii. (1843) 394 = acanthocarpa.
sinemariensis, Aubl. Pl. Gui. i. 462. t. 185 = Stigmaphyllon puberum.
sinuata, *DC. Prod.* i. 588.—Guiana.
solitaria, Vell. Fl. Flum. 189; iv. t. 151 = Hiraea Salzmanniana.
Spixiana, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 561 = Heteropteris suberosa.
splendens, DC. Prod. i. 588 = Stigmaphyllon fulgens.
Spruceana, *Griseb. in Mart. Fl. Bras.* xii. III. 45.—Bras.
stellaris, *Griseb. in Linnaea*, xiii. (1839) 192.—Bras.
subcordata, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 336 = adenopoda.
suberosa, Willd. ex Steud. Nom. ed. II. i. 184 = Heteropteris suberosa.
tenuis, Lindl. in Bot. Reg. (1838) Misc. 72 = Heteropteris umbellata.
terminalis, *Vell. Fl. Flum.* 192; iv. t. 166.—Bras.
ternata, Vell. l. c. 190; iv. t. 159 = trifoliata.
ternata, Sesse, ex DC. Prod. i. 591 = Stigmaphyllon ellipticum.
tetraptera, Sonner. Voy. Indes, iii. 270 = Hiptage Madablota.
thyrsoidea, Griseb. in Linnaea, xiii. (1839) 199 = Heteropteris thyrsoidea.
tiliaefolia, H. B. & K. Nov. Gen. et Sp. v. 162 = Stigmaphyllon Humboldtianum.
tiliaefolia, *Vent. Choix*, 50. t. 50.—Java.
timoriensis, *DC. Prod.* i. 588.—Ins. Timor.
tomentosa, Desf. Tabl. ed. II. 160 = emarginata.
tomentosa, Salzm. ex A. Juss. in Arch. Mus. Par. iii. (1843) 383 = Stigmaphyllon rotundifolium.
tomentosa, Schlecht. in Linnaea, x. (1833) 244 = Heteropteris Beecheyana.
triflora, Vell. Fl. Flum. 190; iv. t. 156 = crotonifolia.
trifoliata, *Steud. Nom.* ed. II. i. 184.—Bras.
tristellocarpa, Boj. ex A. Juss. in Arch. Mus. Par. iii. (1843) 498 = Tristellatcia Bojeriana.
umbellata, Sieber, ex A. Juss. l. c. 356 = Brachypterys borealis.
umbellulata, *DC. Prod.* i. 588.—Ins. S. Doming.
umbellulata, Sieber, ex A. Juss. in Arch. Mus. Par. iii. (1843) 42 = Byrsonima trinitensis.
unialata, Vell. Fl. Flum. 190; iv. t. 157 = B. ferruginea.
unicapsularis, Lam. Encyc. i. 367 = Hiptage Madablota.

BANISTERIA:—
varia, Spreng. Syst. ii. 386 = variifolia.
variifolia, *DC. Prod.* i. 588.—N. Granat.
velutina, A. Juss. in Arch. Mus. Par. iii. (1843) 401 = crotonifolia.
vernoniaefolia, *Mart. ex A. Juss. l. c.* 398.—Bras.
verruculosa, *Sw. Ann. Bot. Posth.* 31.—Ind. occ.
virgultosa, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 414 = pubipctala.
vitifolia, Sesse, ex DC. Prod. i. 588 = variifolia.
volubilis, Endl. ex Hcynh. Nom. ii. 59 = Hiraea Simsiana.
zanzibarica, Boj. ex Steud. Nom. ed. II. i. 184 = Acridocarpus zanzibaricus.

BANIUM, Ces. ex Boiss. Fl. Orient. ii. 882 (1872) = **Carum**, Linn. (Umbell.).
ligusticifolium, Ces. ex Boiss. l. c. = C. multiflorum.

BANJOLEA, Bowdich, Madeira, 258 (1825). *ACANTHACEAE*? (Genus omnino dub.).
violacea, *Bowdich, l. c.*—Madera.

BANKESIA, Bruce, Trav. ed. II. vii. 181 (1805) = **Brayera**, Kunth (Rosac.).

BANKSEA, Koen. in Retz. Obs. iii. 75 (1783) = **Costus**, Linn. (Scitamin.).
speciosa, Koen. l. c. = C. speciosus.

BANKSIA, Bruce, Trav. ed. II. vii. tt. 22, 23 = **Brayera**, Kunth (Rosac.).
abyssinica, 'Bruce', ex Steud. Nom. ed. II. i. 184 = Brayera anthelmintica.

BANKSIA, Domb. ex DC. Prod. iii. 83 (1828) = **Cuphea**, P. Br. (Lythrar.).
ciliata, Domb. ex DC. l. c. 88 = C. microphylla.
glutinosa, Domb. ex DC. l. c. = C. cordata.

BANKSIA, Forst. Char. Gen. 7. t. 4 (1776) = **Pimelea**, Banks (Thymel.).
Gnidia, Forst. l. c. 8 = P. Gnidia.
pilosa, Forst. f. in Nov. Act. Upsal. iii. (1780) 174 = P. pilosa.
prostrata, Forst. Char. Gen. 8 = P. prostrata.
tomentosa, Forst. l. c. = P. pilosa.

BANKSIA, Linn. f. Suppl. 15, 126 (1781). *PROTEACEAE*, Benth. & Hook. f. iii. 184.
ISOSTYLIS, R. Br. ex Steud. Nom. ed. II. i. 829 (1840).
aemula, *R. Br. in Trans. Linn. Soc.* x. (1811) 209.—Austral.
aemula, Sieber, ex Meissn. in DC. Prod. xiv. 461 = serrata.
aquifolia, Lindl. Swan Riv. App. 34 = ilicifolia.
asplenifolia, Salisb. Prod. 51 = integrifolia.
asplenifolia, Knight, Prot. 113 = marcescens.
attenuata, *R. Br. in Trans. Linn. Soc.* x. (1811) 209.—Austral.
australis, R. Br. l. c. 205 = marginata.
barbigera, Meissn. in Lehm. Pl. Preiss. ii. 264 = Goodii.
Baueri, *R. Br. Prot. Nov.* 35.—Austral.
Baxteri, *R. Br. l. c.* 36.—Austral.
blechnifolia, F. Muell. Fragm. iv. 108 = repens.
Brownii, *Baxter, ex R. Br. Prot. Nov.* 37.—Austral.
Caleyi, *R. Br. l. c.* 35.—Austral.
Candolleana, *Meissn. in Hook. Kew Journ.* vii. (1855) 118.—Austral.
coccinea, *R. Br. in Trans. Linn. Soc.* x. (1811) 207.—Austral.
collina, *R. Br. l. c.* 204.—Austral.
compar, R. Br. l. c. 207 = integrifolia.
conchifera, Gaertn. Fruct. i. 221. t. 48 = serrata.
cuneifolia, Hoffm. in Roem. & Schult. Syst. iii. Mant. 379 = integrifolia.
cuneifolia, *Wendl. f. ex Hoffmgg. Verz. Pfl. Nachtr.* ii. 64.—Austral.
Cunninghamii, Sieber, in Spreng. Syst. iv. Cur. Post. 47 = collina.
cylindrostachya, Lindl. Swan Riv. App. 34 = attenuata.
dactyloides, Gaertn. Fruct. i. 221. t. 47. f. 2 = Hakea dactyloides.
dentata, *Linn. f. Suppl.* 127.—Austral.
dentata, Wendl. Hort. Herrenh. t. 8 = serrata.
denticulata, Dum.-Cours. Bot. Cult. ed. II. vii. 108 = spinulosa.

BANKSIA:—
depressa, R. Br. in Trans. Linn. Soc. x. (1811) 205 = marginata.
dilleniaefolia, Knight, Prot. 113 = latifolia.
dryandroides, *Baxter, in Sweet, Fl. Austral.* t. 56.—Austral.
elatior, R. Br. in Trans. Linn. Soc. x. (1811) 209 = aemula.
elegans, *Meissn. in Hook. Kew Journ.* vii. (1855) 119.—Austral.
erieaefolia, *Linn. f. Suppl.* 127.—Austral.
fagifolia, Hoffm. ex Roem. & Schult. Syst. iii. Mant. 379 = latifolia.
ferrea, Vent. ex Spreng. Syst. i. 485 = marginata.
gibbosa, Sm. in White, Voy. 224. t. 22. f. 2 = Hakea gibbosa.
glauca, Cav. in Anal. Hist. Nat. i. (1799) 230 = integrifolia.
Goodii, *R. Br. Prot. Nov.* 36.—Austral.
grandidentata, Dum.-Cours. Bot. Cult. ed. II. vii. 108 = speciosa.
grandis, *Willd. Sp. Pl.* i. 535.—Austral.
Guentheri, *Regel, Ind. Sem. Hort. Petrop.* (1857) 37; *Gartenfl.* (1858) 50.—Austral.
Gunnii, Meissn. in Hook. Kew Journ. iv. (1852) 210 = marginata.
heterophylla, Hort. ex Meissn. in DC. Prod. xiv. 466 = Hakea suaveolens.
Hookeri, Drumm. in Comp. Bot. Mag. (1848) 1 = Solandri.
Hookeriana, *Meissn. in Hook. Kew Journ.* vii. (1855) 119.—Austral.
Huegelii, *R. Br. ex Endl. Gen. Suppl.* iv. II. 89.—Austral.
hypoleuca, Hoffmgg. Verz. Pfl. Nachtr. ii. 66 = marginata.
ilicifolia, *R. Br. in Trans. Linn. Soc.* x. (1811) 211.—Austral. ?
insularis, *R. Br. l. c.* 206 = marginata.
integerrima, *Dum.-Cours. Bot. Cult.* ed. II. ii. 421; vii. 108.—Austral. ?
integrifolia, Labill. ex Meissn. in DC. Prod. xiv. 456 = marginata.
integrifolia, *Linn. f. Suppl.* 127.—Austral.
integrifolia, Schleeht. in Linnaea, xx. (1847) 584 = australis.
laevigata, *Meissn. in Hook. Kew Journ.* iv. (1852) 210.—Austral.
latifolia, *R. Br. in Trans. Linn. Soc.* x. (1811) 208.—Austral.
ledifolia, A. Cunn. ex Meissn. in DC. Prod. xiv. 454 = Cunninghamii.
Lemanniana, *Meissn. in Hook. Kew Journ.* iv. (1852) 210.—Austral.
Lindleyana, *Meissn. l. c.* vii. (1855) 120.—Austral.
littoralis, Lindl. in Bot. Reg. t. 1363 = eollina.
littoralis, *R. Br. in Trans. Linn. Soc.* x. (1811) 204.—Austral.
longifolia, D. Dietr. Syn. i. 538 = latifolia.
longifolia, Hort. ex Steud. Nom. ed. II. i. 184 = oblongifolia.
longifolia, Don, ex F. G. Dietr. Vollst. Gaertn. Lexik. ii. 150 = serrata.
macrophylla, Link, Enum. Hort. Berol. i. 116 = integrifolia.
marcescens, Bonpl. Jard. Malm. 116. t. 48 = marginata.
marcescens, Hook. Bot. Mag. t. 2803 = australis.
marceseens, *R. Br. in Trans. Linn. Soc.* x. (1811) 208.—Austral.
marginata, *Cav. in Anal. Hist. Nat.* i. (1799) 227.—Austral.
marginata, *R. Br. in Trans. Linn. Soc.* x. (1811) 204.—Austral.
media, Hook. f. Fl. Tasm. i. 329 = serrata.
media, *R. Br. Prot. Nov.* 35.—Austral.
Meissneri, *Lehm. Pl. Preiss.* i. 582.—Austral.
Menziesii, *R. Br. Prot. Nov.* 36.—Austral.
microstachya, Cav. in Anal. Hist. Nat. i. (1799) 224 = marginata.
mimosoides, *Hort. Cantabr. ex F. G. Dietr. Vollst. Gaertn. Lexik.* ii. 150.—Austral.
mitis, Knight, Prot. 112 = serrata.
museuliformis, *Gaertn. Fruct.* i. 221.—Austral.
nivea, Labill. Voy. i. 411. t. 24 = Dryandra nivea.
nutans, *R. Br. in Trans. Linn. Soc.* x. (1811) 203.—Austral.
oblongifolia, *Cav. Anal. Hist. Nat.* i. (1799) 225.—Austral.
oblongifolia, Lodd. Bot. Cab. t. 241 = marginata.

BANKSIA:—
oecidentalis, *R. Br. in Trans. Linn. Soc.* x. (1811) 204.—Austral.
oleaefolia, Cav. in Anal. Hist. Nat. i. (1799) 228. t. 14 = integrifolia.
oleaefolia, Salisb. Prod. 51 = Hakea dactyloides.
ornata, *F. Muell. ex Meissn. in Linnaea*, xxvi. (1853–55) 352.—Austral.
paludosa, R. Br. in Trans. Linn. Soc. x. (1811) 207 = integrifolia.
patula, R. Br. l. e. 205 = marginata.
petiolaris, *F. Muell. Fragm.* iv. 109.—Austral.
phylicaefolia, Salisb. Prod. 51 = erieaefolia.
pinifolia, Meissn. in Hook. Kew Journ. vii. (1855) 118 = sphaerocarpa.
pinifolia, Salisb. Prod. 51 = Hakea gibbosa.
pinnata, Hort. Cels, ex Roem. & Schult. Syst. iii. 444 = Hakea pinnata.
pinnatisecta, F. Muell. Fragm. vii. 58 = repens.
polypodiifolia, Knight, Prot. 113 = repens.
praemorsa, Andr. Bot. Rep. t. 258 = marceseens.
praemorsa, Hort. ex Dum.-Cours. Bot. Cult. vii. ed. II. 107 = marginata.
prionophylla, F. Muell. 1st Gen. Rep. 17 = eollina.
prionotes, *Lindl. Swan River App.* 34.—Austral.
procumbens, Hort. ex Dum.-Cours. Bot. Cult. ed. II. vii. 107 = dentata.
prostrata, *R. Br. Prot. Nov.* 36.—Austral.
pubeseens, *Donn, ex Roem. & Schult. Syst.* iii. 444.—Austral.
pulchella, *R. Br. in Trans. Linn. Soc.* x. (1811) 202.—Austral.
pyriformis, Gaertn. Fruet. i. 220. t. 47 = Xylomelum pyriforme.
quereifolia, *R. Br. in Trans. Linn. Soc.* x. (1811) 210.—Austral.
repens, *Labill. Voy.* i. 412. t. 23.—Austral.
reticulata, Hoffm. in Roem. & Schult. Syst. Mant. iii. 379 = integrifolia.
retieulata, *Wendl. f. ex Hoffmgg. Verz. Pfl. Nachtr.* ii. 67.—Austral.
Robur, Cav. in Anal. Hist. Nat. i. (1799) 226 = latifolia.
salicifolia, Cav. l. c. 230 = integrifolia.
Seeptrum, *Meissn. in Hook. Kew Journ.* vii. (1855) 120.—Austral.
serraefolia, Knight, Prot. 112 = aemula.
serrata, *Linn. f. Suppl.* 126.—Austral.
serrata, Cav. Ic. vi. 27. t. 540 = aemula.
serratifolia, Salisb. Prod. 51 = aemula.
Solandri, *R. Br. Prot. Nov.* 36.—Austral.
speciosa, Lindl. in Bot. Reg. t. 1728 = Vietoriae.
speeiosa, *R. Br. in Trans. Linn. Soc.* x. (1811) 210.—Austral.
sphaeroearpa, *R. Br. l. c.* 203.—Austral.
spicata, Gaertn. Fruet. i. 221. t. 48 = integrifolia.
spinulosa, *Sm. New Holl. Pl.* i. 13. t. 4.—Austral.
tenuifolia, Salisb. Prod. 50 = Hakea aeieularis.
teretifolia, Salisb. l. e. 51 = Hakea pugioniformis.
tomentosa, *F. Muell. in Hook. Kew Journ.* viii. (1856) 327.—Austral.
tomentosa, Hort. ex Meissn. in DC. Prod. xiv. 454 = littoralis.
trieuspis, *Meissn. in Hook. Kew Journ.* vii. (1855) 118.—Austral.
tridentata, *Meissn. l. c.* 120.—Austral.
uncigera, Knight, Prot. 112 = latifolia.
undulata, Lindl. in Bot. Reg. t. 1316 = serrata.
verticillata, *R. Br. in Trans. Linn. Soc.* x. (1811) 207.—Austral.
Vietoriae, *Meissn. in Hook. Kew Journ.* vii. (1855) 119.—Austral.

BANNISTERIA, Linn. Gen. ed. I. 377 (1737) = **Banisteria**, Linn. (Malpig.).

BANNISTERIOIDES, Linn. Fl. Zeyl. 192 (1747) = **Pellia**, Gaertn. (Inc. sed.).

BAOBAB, Mill. Gard. Diet. ed. VI. 175 (1752) = **Adansonia**, Linn. (Malvac.).

BAPHIA, Afzel. ex Lodd. Bot. Cab. t. 367 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 553.
BRACTEOLARIA, Hoehst. in Flora, xxiv. (1841) II. 638.
CARPOLOBIA, G. Don, Gen. Syst. i. 370 (1831).
DELARIA, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 406.

BAPHIA:—
angolensis, *Welw. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 249.—Afr. trop.
haematoxylon, Hoehst. ex Hook. Niger Fl. 321 = nitida.
Heudelotiana, *Baill. Adansonia*, vi. (1866) 216.—Afr. trop.
Kirkii, *Baker, in Oliver, Fl. Trop. Afr.* ii. 250.—Afr. trop.
laurifolia, Baill. Adansonia, vi. (1866) 213 = pubescens.
leptostemma, Baill. l. c. 214 = nitida.
maxima, *Baker, in Oliver, Fl. Trop. Afr.* ii. 250.—Afr. trop.
nitida, *Lodd. Bot. Cab.* t. 367.—Afr. trop.
pilosa, *Baill. Adansonia*, vi. (1866) 216.—Afr. trop.
polygalacea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 248.—Afr. trop.
pubeseens, *Hook. f. in Hook. Niger Fl.* 320.—Afr. trop.
pyrifolia, Baill. in Bull. Soc. Linn. Par. i. (1885) 445 = polygalaeea.
racemosa, *Hochst. in Flora*, xxiv. (1841) II. 638.—Afr. trop.
spathaeca, *Hook. f. in Hook. Niger Fl.* 320.—Afr. trop.

BAPHIOPSIS, Benth. ex Baker, in Oliver, Fl. Trop. Afr. ii. 256 (1871). *LEGUMINOSAE*.
parviflora, *Benth. ex Baker, l. c.*—Afr. trop.

BAPHORHIZA, Link, Handb. i. 578 (1829) = **Alkanna**, Tausch (Borag.).
tinctoria, Link, l. c. = A. tinetoria.

BAPTISIA, Vent. Dec. Gen. Nov. 9 (1808). *LEGUMINOSAE*, Benth. & Hook. f. i. 466.
CROTALOPSIS, Michx. ex DC. Prod. ii. 100, in syn. (1825).
EAPLOSIA, LASINIA, PERICAULON, RIPASIA, Rafin. New Fl. Am. ii. 48–51 (1836).
alba, Hook. Bot. Mag. t. 1177 = leucantha.
alba, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 6.—Am. bor.
albiflora, *Rafin. New Fl. Am.* ii. 47.—Am. bor.
aurieulata, *Sweet, Hort. Brit.* ed. II. 123.—Am. bor.
australis, Hort. ex Lehm. in Nov. Act. Nat. Cur. xiv. (1829) 803 = minor.
australis, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 6.—Am. bor.
bracteata, Muhl. ex Ell. Sketch, i. 469 = leueophaea.
caerulea, Eaton & Wright, Man. N. Am. Bot. 154 = australis.
ealycosa, *Engelm. in Coult. Bot. Gaz.* iii. (1878) 65.—Florida.
confusa, Sweet, ex G. Don, Gen. Syst. ii. 113 = australis ?
exaltata, Sweet, Brit. Flow. Gard. i. t. 97 = australis.
fraxinifolia, Nutt. ex Torr. & Gray, Fl. N. Am. i. 387, in obs. = Thermopsis fraxinifolia.
lanceolata, *Ell. Sketch*, i. 467.—Am. bor.
Leeontii, *Torr. & Gray, Fl. N. Am.* i. 386.—Florida.
leucantha, *Torr. & Gray, l. c.* 385.—Am. bor.
leucophaea, *Nutt. Gen. Am.* i. 282.—Am. bor.
lupinoides, *Burbidge, in Gard. Chron.* (1884) I. 827.—Am. bor. ?
megaearpa, *Chapm. ex Torr. & Gray, Fl. N. Am.* i. 386.—Florida.
microphylla, *Nutt. in Journ. Acad. Philad.* vii. (1834) 97.—Florida.
minor, Lehm. in Nov. Act. Nat. Cur. xiv. (1829) 803 = australis.
mollis, DC. Prod. ii. 100 = Thermopsis mollis.
mollis, Nutt. Gen. Am. i. 281 = Thermopsis fraxinifolia.
nepalensis, Hook. Exot. Fl. t. 131 = Piptanthus nepalensis.
perfoliata, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 5.—Am. bor.
retusa, *Rafin. New Fl. Am.* ii. 47.—Am. bor.
Serenae, *M. Curt. in Am. Journ. Sc.* Ser. II. vii. (1849) 406.—Am. bor.
simplicifolia, *Croom, in Am. Journ. Sc.* xxv. (1834) 74.—Florida.
sphaerocarpa, *Nutt. in Journ. Acad. Philad.* vii. (1834) 97.—Texas.
stipulacea, Ravenel, in Elliot Soc. Nat. Hist. i. (1856) t. 2 = microphylla.

BAPTISIA:—
sulphurea, *Engelm. in Coult. Bot. Gaz.* iii. (1878) 65.—Am. bor.
texana, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 452 = australis?
tinctoria, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 6.—Am. bor.
uniflora, Spreng. Syst. ii. 347 = lanceolata.
uniflora, *Nutt. Gen. Am.* i. 281, nomen = praec.?
versicolor, Lodd. Bot. Cab. t. 1144 = australis?
versicolor, *Rafin. New Fl. Am.* iv. 107, nomen.—Am. bor.
villosa, *Nutt. Gen. Am.* i. 281.—Carolina.

BAPTISTONIA, Barb.-Rodr. Orch. Nov. i. 95 (1877). *ORCHIDEAE.*
echinata, *Barb.-Rodr. l. c.*—Bras.

BARANDA, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 502 = **Barringtonia**, Forst. (Myrtac.).
angatensis, Llanos, l. c. = B. macrostachya.

BARATRANTHUS, Miq. Fl. Ind. Bat. i. 1. 834 (1855) = **Loranthus**, Linn.
axanthus, Miq. l. c.—Sumatra.

BARATTIA, A. Gray & Engelm. in Proc. Am. Acad. i. (1848) 48 = **Encelia**, Adans. (Compos.).
calva, A. Gray & Engelm. l. c. = E. calva.

BARAULTIA, Spreng. Syst. ii. 313 (1825) = **Barraldeia**, Thou.

BARBACENIA, Vand. Fl. Lusit. & Bras. Spec. 21. t. 1 (1796). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 740.
CAMPDERIA, H. B. & K. Nov. Gen. et Sp. vii. 155 (1825).
PLEUROSTIMA, Rafin. Fl. Tellur. ii. 97 (1836).
RADIA, A. Rich. in Kunth, Syn. Pl. Aequin. i. (1822).
VISNEA, Steud. ex Endl. Gen. 173 (1836).
Alexandrinae, *Schousb. ex Hook. Lond. Journ. Bot.* iv. (1845) 13.—Guiana.
bicolor, *Mart. Nov. Gen. et Sp.* i. 19. t. 13.—Bras.
brasiliensis, Willd. Sp. Pl. ii. 227 = Vandellii.
coccinea, *Mart. in Schult. f. Syst.* vii. 286.—Bras.
ensifolia, *Mart. l. c.*—Bras.
exscapa, *Mart. Nov. Gen. et Sp.* i. 21. t. 14.—Bras.
flava, *Mart. ex Steud. Nom.* ed. II. i. 185.—Bras.
Gardneri, *Seub. in Mart. Fl. Bras.* iii. 1. 70. t. 8. f. 5.—Bras.
glauca, *Mart. in Schult. f. Syst.* vii. 238.—Bras.
gracilis, Hort. ex Baker, in Journ. Linn. Soc. xviii. (1881) 239 = Dasylirion acrotrichum.
ignea, *Mart. in Schult. f. Syst.* vii. 285.—Bras.
longiflora, *Mart. Nov. Gen. et Sp.* i. 19. t. 12.—Bras.
luzulaefolia, *Mart. in Schult. f. Syst.* vii. 285.—Bras.
macrantha, *Lem. Jard. Fl.* iv. (1854) t. 390.—Bras.
purpurea, *Hook. Bot. Mag.* t. 2777.—Bras.
Rogierii ×, Hort. ex Moore & Ayres, Mag. Bot. ii. (1850) 209.—Bras.
rubro-virens, *Mart. Nov. Gen. et Sp.* i. 20. t. 14. f. 1.—Bras.
schidigera, *Lem. Jard. Fl.* ii. (1852) t. 198.—Bras.
squamata, *Lindl. in Bot. Reg.* (1843) *Misc.* 64.—Bras.
tomentosa, *Mart. Nov. Gen. et Sp.* i. 18. t. 11.—Bras.
tricolor, *Mart. l. c.* 18. t. 10.—Bras.
Vandelli, Schult. f. Syst. vii. 289 = glauca.
Vandellii, *Pohl, ex Seub. in Mart. Fl. Bras.* iii. 1. 72.—Bras.

BARBA-JOVIS, Adans. Fam. ii. 321 (1763) = **Anthyllis**, Linn. (Legumin.).
argyrophylla, Moench, Meth. 110 = A. Barba-Jovis.
linearifolia, Moench, Meth. Suppl. 38 = A. Hermanniae.
montana, Moench, l. c. 37 = A. montana.
tragacanthoides, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 347 = Astragalus armatus?

BARBAREA, Scop. Fl. Carn. ed. I. 522 (1760) = **Dentaria**, Linn. (Crucif.).

BARBAREA, R. Br. in Ait. Hort. Kew. ed. II. iv. 109 (1812). *CRUCIFERAE.* Benth. & Hook. f. i. 68.
CAMPE, Dulac, Fl. Hautes-Pyr. 199 (1867).

BARBAREA:—
alpina, *C. Koch, in Linnaea*, xix. (1847) 319.—As. Min.
altaica, Andrz. ex Steud. Nom. ed. II. i. 185 = arcuata.
arcuata, Andrz. ex DC. Prod. i. 141 = taurica.
arcuata, *Reichb. in Flora*, v. (1822) 296.—Europ.; As. Min.; As. bor.
augustana, Boiss. Diagn. Ser. I. i. 69 = vulgaris.
australis, Hook. f. Fl. N. Zel. i. 14 = vulgaris.
australis, Jord. Diagn. i. 103 = praecox.
brachycarpa, *Boiss. Diagn.* Ser. I. v. 80.—As. Min.
bracteosa, *Guss. Fl. Sic. Prod.* ii. 257.—Sicilia.
brevicaulis, Jord. Diagn. i. 104 = rupicola.
brevistyla, Jord. l. c. 102 = praecox.
cilicica, Schott & Kotschy, ex Boiss. Fl. Orient. i. 185 = minor.
ciliifera, *Steud. Nom.* ed. II. i. 185, nomen.—Graecia.
elata, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 140.—Reg. Himal.
erysimoides, Schur, Enum. Pl. Transs. 41 = praecox.
heterophylla, *Willk. in Bot. Zeit.* v. (1847) 217.—Hispan.
hirsuta, Weihe, in Flora, xiii. (1830) 257 = vulgaris.
humilis, *Presl, Fl. Sic.* i. 45.—Sicilia.
iberica, DC. Syst. ii. 208 = arcuata.
integrifolia, *DC. l. c.*—As. Min.
intermedia, Bor. Fl. Centr. Fr. ed. I. ii. 48 (1840) = vulgaris.
Kayserii, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 57 = vulgaris.
Linnaei, Spenn. Fl. Friburg. iii. 929 = praecox, vulgaris, taurica, &c.
longisiliqua, Jord. Diagn. i. 103 = praecox.
lyrata, Aschers. in Fl. Prov. Brandenb. i. 35 = vulgaris.
minor, *C. Koch, in Linnaea*, xix. (1847) 55.—As. Min.; Syria.
oligosperma, *C. Koch, l. c.* 319.—Armenia.
orthoceras, *Ledeb. Hort. Dorp.* (1824).—Sibir. altaic.
palustris, Hegetschw. Fl. Schw. 654 = stricta.
parviflora, Fries, Novit. ed. II. 207 = stricta.
patula, Fries, Nov. Mant. iii. 76 = praecox.
perennis, *Pomel, Nouv. Mat. Fl. Atl.* 375.—Algeria.
planisiliqua, *C. A. Mey. in Middend. Reise (Fl. Ochot.* 14).—Sibir.
plantaginea, *DC. Syst.* ii. 208.—As. Min.; Persia.
praecox, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 109.—Europ.
prostrata, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 348 = sicula.
pyrenaica, Jord. Diagn. i. 349 = vulgaris.
rivularis, Martr. Fl. Tarn, 44 = vulgaris.
rivularis, *Panč. Elem.* 14.—Servia.
rupestris, Steud. Nom. ed. II. i. 185 = vulgaris.
rupicola, *Moris, Stirp. Sard. Elench.* i. 55.—Sardinia.
sicula, Gren. & Godr. Fl. Fr. i. 92 = vulgaris.
sicula, *C. & J. Presl, Delic. Prag.* 17.—Ital.; As. Min.
stolonifera, *Pomel, Nouv. Mat. Fl. Atl.* 375.—Algeria.
stricta, Fellm. Pl. Vasc. Lapp. 6 = orthoceras.
stricta, *Andrz. in Bess. Enum. Pl. Volh.* 72.—Ross. med.
stricta, Willk. in Flora, xxxiv. (1851) 583 = vulgaris.
sylvestris, Jord. Diagn. i. 100 = vulgaris.
taurica, DC. Syst. ii. 207 = arcuata.
verna, Aschers. Fl. Prov. Brandenb. i. 36 = praecox.
vicina, Martr. Fl. Tarn, 45, in obs. = vulgaris.
vulgaris, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 109.—Europ.; As. temp.

BARBELLINA, Cass. in Dict. Sc. Nat. l. 440 (1827) = **Staehelina**, Linn. (Compos.).
sericea, Cass. l. c. = S. arborescens.

BARBERETTA, Harv. Gen. S. Afr. Pl. ed. II. 377 (1868). *HAEMODORACEAE*, Benth. & Hook. f. iii. 674.
aurea, *Harv. l. c.*—Afr. austr.

BARBERINA, Vell. Fl. Flum. 235 (1825); Ic. v. t. 117 (1827) = **Symplocos**, Linn. (Styrac.).
antillana, Miers, in Journ. Linn. Soc. xvii. (1880) 294.—Ind. occ.
celastrinea, Miers, l. c. = S. celastrinea.
crenata, Miers, l. c.—Bras.
cubensis, Miers, l. c. = S. cubensis.
estrellensis, Miers, l. c. 293 = S. estrellensis.
hirsuta, Vell. Fl. Flum. 235; v. t. 117 = S. hirsuta.
lanceolata, Mart. Herb. Fl. Bras. n. 723 = S. lanceolata.
phaeoclados, Mart. l. c. 724 = S. phaeoclados.

BARBERINA:—
ramentacea, Miers, in Journ. Linn. Soc. xvii. (1880) 294 = S. ramentacea.
revoluta, Mart. Herb. Fl. Bras. n. 832 = S. revoluta.
rhamnifolia, Miers, in Journ. Linn. Soc. xvii. (1880) 293 = S. rhamnifolia.
Spruceana, Miers, l. c. 295.—Peruv.
tetrandra, Mart. Syst. Mat. Med. Bras. 49 = S. tetrandra.

BARBEUIA, Thou. Gen. Nov. Madag. 6 (1806). *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 85.
madagascariensis, *Steud. Nom.* ed. II. i. 186.—Madag.

BARBIERA, Spreng. Gen. ii. 587 (1831) = seq.

BARBIERIA, DC. Mém. Lég. 241. t. 39 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 495.
maynensis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 58.—Peruv.
pinnata, Baill. Hist. Pl. ii. 263 = polyphylla.
polyphylla, *DC. Mém. Lég.* vi. 249. t. 39.—Porto Rico.

BARBILUS, P. Br. Hist. Jamaic. 216 (1756) = **Trichilia**, Linn. (Meliac.).

BARBULA, Lour. Fl. Cochinch. 366 (1790) = **Caryopteris**, Bunge (Verben.).
sinensis, Lour. l. c. = C. incana.

BARBYLUS, Juss. Gen. 430 (1789) = **Trichilia**, Linn. (Meliac.).
Brownii, J. F. Gmel. Syst. 611 = T. hirta?
jamaicensis, DC. Prod. ii. 91 = T. hirta?

BARCELLA, Drude, in Mart. Fl. Bras. iii. 459. t. 106. (1881) = **Elaeis**, Jacq. (Palm.).
odora, Drude, l. c. = E. odora.

BARCENA, Dugès, in Rev. Cienc. Mexic. Dec. i. (1879) 8, cum tab.; et in La Naturaleza, iv. (1879) 281. *RHAMNEAE.*
guanajuatensis, *Dugès, ll. cc.*—Mexic.

BARCENIA, Dugès, in La Naturaleza, iv. (1879) 283 = praec.

BARCKHAUSENIA, Menke, in Flora, xxvii. (1854) 723 = seq.

BARCKHAUSIA, DC. Fl. Fr. vi. 449 (1815) = BARKHAUSIA, Moench = **Crepis**, Linn. (Compos.).

BARCLAYA, Wall. in Trans. Linn. Soc. xv. (1827) 442. t. 18. *NYMPHAEACEAE*, Benth. & Hook. f. i. 47.
longifolia, *Wall. l. c.*—Burma.
Mottleyi, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 157. t. 21. f. 1–5.—Borneo.
oblonga, Wall. Cat. n. 599 = longifolia.

BARDANA, Hill, Veg. Syst. iv. 28 (1762) = **Arctium**, Linn. (Compos.).
Arctium, Hill, Hort. Kew. 64 = A. majus.
Lappa, Hill, Herb. Brit. i. 79 = A. majus.
minor, Hill, Hort. Kew. 64 = A. minus.

BARERIA, Juss. in Dict. Sc. Nat. iv. 64 (1804) = BARRERIA, Scop. = **Poraqueiba**, Aubl. (Olacin.).

BARGEMONTIA, Gaudich. Voy. Bonite, t. 8. ('1839–46'). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 880.
peruviana, *Gaudich. l. c.*—Peruv.

BARHAMIA, Klotzsch, in Seem. Bot. Voy. Herald, 104 (1852?) = **Croton**, Linn. (Euphorb.).
essequiboënsis, Klotzsch, l. c. = C. populifolius.
hispida, Klotzsch, l. c. = C. populifolius.
macrostachya, Klotzsch, l. c. = C. populifolius.
multispicata, Klotzsch, l. c. = C. urticaefolius.
ovalifolia, Klotzsch, l. c. = C. ovalifolius.
panamensis, Klotzsch, l. c. = C. populifolius.
urticaefolia, Klotzsch, l. c. = C. urticaefolius.

BARJONIA, Decne. in DC. Prod. viii. 512 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 771.
chloraefolia, *Decne. l. c.*—Bras.
eymosa, *Fourn. in Mart. Fl. Bras.* vi. iv. 206.—Bras.
deltoidea, Deene. ex Fourn. l. c. 205 = racemosa.
linearis, *Decne. in DC. Prod.* viii. 512.—Bras.
obtusifolia, *Fourn. in Mart. Fl. Bras.* vi. iv. 206.—Bras.
racemosa, *Decne. in DC. Prod.* viii. 512.—Bras.
Warmingii, *Fourn. in Mart. Fl. Bras.* vi. iv. 206.—Bras.

BARKANIA, Ehrenb. [ex Reichb. Consp. 43 (1828)] in Abh. Acad. Berl. (1832) 429 = **Halophila**, Thou. (Hydroch.).
bullata, Ehrenb. in Abh. Acad. Berl. (1832) 429 = H. bullata.
punctata, Ehrenb. Symb. 5, ex Pritz. Ic. Ind. Suppl. 42 = H. ovata.
stipulacea, Zanard. in Mem. Ist. Venet. vii. (1857) 225 = H. stipulacea.

BARKERIA, Knowles & Weste. Flor. Cab. ii. (1838) 7. t. 49 = **Epidendrum**, Linn. (Orehid.).
Barkeriola, Reichb. f. in Gard. Chron. (1884) II. 616 = E. Barkeriola.
cyclotella, Reichb. f. l. e. (1880) I. 72 = E. cyclotellum.
elegans, Knowles & Weste. Flor. Cab. ii. (1838) 7 = E. elegans.
Lindleyana, Batem. ex Lindl. in Bot. Reg. (1842) Mise. 2 = E. Lindleyanum.
melanocaulon, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 22 = E. melanoeaulon.
Skinneri, Paxt. Mag. Bot. xv. (1849) 1 = E. Skinneri.
spectabilis, Batem. in Lindl. Bot. Reg. (1842) Misc. 43 = E. spectabile.
Vanneriana, Reichb. f. in Gard. Chron. (1885) II. 678 = E. Vannerianum.

BARKHAUSENIA, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 69 = BORCKHAUSENIA, Gaert. Mey. & Scherb. = **Corydalis**, Vent. (Papav.).

BARKHAUSIA, Moench, Meth. 537 (1794) = **Crepis**, Linn. (Compos.).
aculeata, DC. Prod. vii. 159 = C. aculeata.
adenothrix, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 465 = C. Rueppellii.
albida, Cass. in Dict. Se. Nat. xxvi. 62 = C. albida.
alpina, DC. Fl. Fr. ed. III. iv. 41 = C. alpina.
amplexicaulis, Coss. & Dur. ex Coss. in Ann. Sc. Nat. Sér. IV. i. (1854) 232.—Algeria.
amplexifolia, Godr. in Mém. Acad. Montp. (Sect. Méd.) i. (1853) 437 = C. amplexifolia.
amygdalina, Lag. ex Steud. Nom. ed. II. i. 186 = C. foetida.
apargioides, Spreng. Syst. iii. 652 = Chondrilla Peltidium.
arenaria, Pomel, Nouv. Mat. Fl. Atl. 4 = C. arenaria?
aspera, Reichb. in Moessl. Handb. ed. II. ii. 1405 = C. aspera.
auriculifolia, Reichb. l. e. 1410, in nota = C. incarnata?
Balbisiana, DC. Prod. vii. 155 = C. bursifolia.
balearica, Costa, Barcin. (1861) 6 = C. balearica.
bellidifolia, DC. Fl. Fr. v. 449 = C. bellidifolia.
Bivoniana, Reichb. in Moessl. Handb. ed. II. ii. 1408.—Europ.
bursifolia, Spreng. Syst. iii. 653 = C. bursifolia.
byzantina, DC. Prod. vii. 158 = C. foetida.
caespitosa, Moris, Fl. Sard. 92 = C. caespitosa.
Candollei, Spreng. Syst. iii. 657 = C. foetida.
canescens, Spreng. l. c. 653, sub n. 16 = C. bursifolia.
carbonaria, Seh. Bip. in Mus. Senek. iii. 54 = C. carbonaria.
caroliniana, Nutt. Gen. Am. ii. 126 = Pyrrhopappus carolinianus.
cerinna, Reichb. in Moessl. Handb. ed. II. ii. 1406 = C. neglecta.
chondrilloides, Bluff & Fingerh. Comp. ii. 304 = C. foeniculacea.
chaetocephala, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 381; (Lehm. Reliq. Bot. 208).—Turkest.
Clausonis, Pomel, Nouv. Mat. Fl. Atl. 1.—Algeria.

BARKHAUSIA:—
comata, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 27 = C. comata.
cretica, Boiss. Diagn. Ser. I. xi. 53 = C. cretica.
decumbens, Gren. & Godr. Fl. Fr. ii. 332 = C. decumbens.
divaricata, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 26 = C. divaricata, *Lowe*.
divaricata, Schur, Enum. Pl. Transs. 373 = C. foetida.
dubia, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 27.—Ins. Madera.
echioides, Schur, Enum. Pl. Transs. 373 = C. foetida.
elata, Koeh, in Linnaea, xxiii. (1850) 684 = C. alpina.
elegans, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 435 = C. elegans.
erucaefolia, Gren. & Godr. Fl. Fr. ii. 331 = C. erucaefolia.
flexuosa, DC. Prod. vii. 156 = C. nana.
floribunda, Pomel, Nouv. Mat. Fl. Atl. 260 = C. floribunda.
foetida, F. W. Schmidt, Samml. Phys. Aufs. i. (1795) 283 = C. foetida.
glandulosa, Presl, Fl. Sic. p. xxxi = C. foetida.
gracilis, Dum. Fl. Belg. 61.—Belgium.
graveolens, Link, Enum. Hort. Berol. ii. 290 = C. foetida.
Haensleri, Boiss. ex DC. Prod. vii. 153 = C. taraxacifolia.
heterocarpa, Boiss. Voy. Espagne, ii. 742 = C. taraxacifolia.
hiemalis, Spreng. Syst. iii. 652 = C. taraxacifolia.
hieracioides, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 27 = C. hieracioides.
hirsuta, Pomel, Nouv. Mat. Fl. Atl. 260 = C. hirsuta.
hirta, C. Koch, in Linnaea, xxiii. (1850) 688 = C. foetida.
hispida, Link, Enum. Hort. Berol. ii. 290 = C. setosa.
hyemalis, Bivona, Stirp. Rar. Sicil. i. [17] t. 2 = C. taraxacifolia.
interrupta, Reichb. in Moessl. Handb. ed. II. ii. 1407, in nota = C. Sieberi.
intybacea, Spreng. Syst. iii. 653 = C. taraxacifolia.
juvenalis, Delile, Ind. Sem. Hort. Monsp. (1856) 23 = C. juvenalis.
Kotschyana, Boiss. Diagn. Ser. I. vii. 13 = C. Kotschyana.
Kralikii, Pomel, Nouv. Mat. Fl. Atl. 5 = C. Kralikii.
laciniata, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 25 = C. laciniata.
Leontodon, DC. Fl. Fr. ed. III. n. 2950 = C. leontodontoides.
leontodontoides, Spreng. Syst. iii. 653 = C. leontodontoides.
Lessingii, Hook. & Arn. Bot. Beech. Voy. i. 145 = Troximon apargioides.
leucocephala, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 385; (Lehm. Reliq. Bot. 209).—As. centr.
leucorhodia, Reichb. in Moessl. Handb. ed. II. ii. 1409 = C. bicolor.
macrocephala, Willk. in Bot. Zeit. v. (1847) 860 = C. albida.
macrophylla, Spreng. Syst. iii. 652 = C. macrophylla.
Marschallii, C. A. Mey. Verz. Pfl. Cauc. 57 = C. Marschallii.
melanocephala, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 385; (Lehm. Reliq. Bot. 209).—As. centr.
mucronata, Bertol. Fl. Ital. viii. 515 = C. alpestris.
muricata, Spreng. Syst. iii. 651 = C. aspera.
nana, DC. Prod. vii. 156 = C. nana.
nicaeensis, Link, ex Spreng. Syst. iii. 653 = C. taraxacifolia.
nigricans, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 278 = C. nigricans.
nuda, Presl, ex Spreng. Syst. iii. 653 = C. leontodontoides.
nudiflora, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 278 = C. aspera.
numidica, Pomel, Nouv. Mat. Fl. Atl. 261 = C. numidica.
paludosa, Baumg. ex DC. Prod. vii. 170 = C. paludosa.
pingnis, Reichb. in Moessl. Handb. ed. II. ii. 1411 = C. foetida.
porrifolia, DC. Prod. vii. 157 = Pterotheca bifida.
praecox, Reichb. in Moessl. Handb. ed. II. ii. 1409 = C. taraxacifolia.
prostrata, Dum. Fl. Belg. Prod. 61 = C. foetida.
purpurea, Bivona, Stirp. Rar. Man. iv. 22 = Lagoseris purpurea.

BARKHAUSIA:—
radicata, Godr. in Mém. Acad. Montp. (Sect. Méd.). i. (1853) 436 = Hypochoeris radicata.
raphanifolia, Chev. ex Steud. Nom. ed. II. i. 187 = C. taraxacifolia.
raphanifolia, Spreng. Syst. iii. 653 = C. raphanifolia.
recognita, DC. Prod. vii. 154 = C. recognita.
reflexa, C. & J. Presl, Delic. Prag. 111 = C. taraxacifolia.
repens, Spreng. Syst. iii. 652.—China.
rhoeadifolia, Bieb. Fl. Taur. Cauc. iii. 538 = C. foetida.
rodigioides, Sch. Bip. ex Tchihat. Asie Min. Bot. ii. 374 = C. foetida.
Roylei, DC. Prod. vii. 157 = C. bifida.
rubra, Moench, Meth. 537 = C. rubra.
sardoa, Spreng. Syst. iv. Cur. Post. 304.—Sardin.
scapigera, Kunze, ex DC. Prod. vii. 152 (= *Macrorhynchus Poeppigii*).—Chili.
scariosa, Moench, Meth. 537 = C. alpina.
scariosa, Reichb. Fl. Germ. Exc. i. 256 = C. vesicaria.
Schimperi, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 466 = C. Schimperi.
Schultzii, Hochst. ex A. Rich. l. e. 464 = C. Schultzii.
senecioides, Spreng. Syst. iii. 652 = C. senecioides.
setosa, DC. Fl. Fr. ed. III. v. 44 = C. setosa.
spathulata, Spreng. Syst. iii. 651 = C. spathulata.
Sprengelii, Steud. Nom. ed. II. i. 187 = Troximon apargioides.
stricta, Reichb. in Moessl. Handb. ed. II. ii. 1406 = C. neglecta.
Suffreniana, DC. Cat. Hort. Monsp. 83 = C. Suffreniana.
taraxacifolia, DC. Fl. Fr. iv. 43 = C. taraxacifolia.
taraxacifolia, Spreng. Syst. iii. 653 = C. taraxacoides.
taraxacoides, Reichb. in Moessl. Handb. ed. II. ii. 1408; DC. Prod. vii. 155 = C. taraxacoides.
tenella, Benth. in Hook. Lond. Journ. Bot. i. (1842) 488 = Lactuca versicolor.
tenerrima, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 464 = C. carbonaria.
tenerrima, Tenore, in Ann. St. Nat. Bolog. iv. 252 = C. leontodontoides.
tenuifolia, DC. Prod. vii. 155 = C. tenuifolia.
triangularis, C. Koch, in Linnaea, xxiii. (1850) 687 = C. foetida.
versicolor, Spreng. Syst. iii. 651 = Lactuca versicolor.
vesicaria, Spreng. l. e. 652 = C. vesicaria.
zazinthia, Margot. & Reut. Fl. Zanth. 62 = C. foetida

BARKHUSENIA, Hoppe, in Flora, xii. (1829) 512 = BARKHAUSIA, Moench = **Crepis**, Linn. (Compos.).

BARKLYA, F. Muell. in Journ. Linn. Soc. iii. (1859) 158; Fragm. i. 109 (1859). *LEGUMINOSAE*, Benth. & Hook. f. i. 559.
syringifolia, *F. Muell. ll. cc.*— Austral.

BARLAEA, Reichb. f. in Linnaea, xli. (1877) 54 = **Habenaria**, Willd. (Orchid.).
calcarata, Reichb. f. l. e. = H. calcarata.

BARLERIA, Linn. Gen. ed. I. 869 (1737). *ACANTHACEAE*, Benth. & Hook. f. ii. 1091.
BARLERIACANTHUS, BARLERIANTHUS, BARLERIOPSIS, BARLERIOSIPHON, BARLERITES, DICRANACANTHUS, PRIONITIS et PSEUDOBARLERIA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 133-137.
BARRELIERA, J. F. Gmel. Syst. 959 (1791).
CRABBEA, Harv. Gen. Pl. Cap. ed. I. 276 (1838).
SOUBEYRANIA, Neck. Elem. i. 365 (1790).
WAHABIA, Fenzl, in Flora, xxvii. (1844) 312.
acanthoides, *Vahl, Symb. Bot.* i. 47.—Arab.; Aegypt.; Ind. bor. oce.
acanthophora, *Nees, in DC. Prod.* xi. 726.—Afr. trop.
aculeata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 85.—Ins. Socotr.
acuminata, Wight, Ic. t. 450 = tomentosa.
adelensis, *Delile, in Rochet, Sec. Voy. Choa*, 341.—Afr. trop.
alata, *S. Moore, in Journ. Bot.* xviii. (1880) 266.—Afr. trop.
alba, Lodd. Bot. Cab. iv. t. 360 = cristata.
anomala, Spreng. Syst. ii. 825 = Bravaisia floribunda.
arabica, *Belang. ex Nees, in DC. Prod.* xi. 725.—Arabia.
argentea, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 86.—Ins. Socotr.

BARLERIA :—
Arnottiana, *Nees, in Hook. Comp. Bot. Mag.* ii. (1836) 311.—Zeylan. ; Burma.
Aucheriana, Nees, in DC. Prod. xi. 234 = Hochstetteri.
auriculata, Schum. Beskr. Guin. Pl. 285 = Hygrophila spinosa.
axillaris, Steud. Nom. ed. II. i. 187 = Lepidagathis trinervis ?
barbata, E. Mey. Zwei Pfl. Docum. 152 = Gueinzii.
Beddomei, T. Anders. ex Bedd. Ic. Ind. Or. i. 62 = Lawii.
bispinosa, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 144 = B. Prionitis.
bispinosa, *Vahl, Symb. Bot.* i. 46.—Arab.
bispinosa, Nees, in Wall. Pl. As. Rar. iii. 94 = mysorensis.
bituminosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 8.—Malaya.
Boivinii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 31.—Madag.
bracteata, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 490 = nitida.
brasiliensis, *Spreng. Neue Entdeck.* iii. 26.—Bras.
brevispina, R. Br. in Salt, Abyss. App. 64 = B. Prionitis.
Burchelliana, Nees, in DC. Prod. xi. 235 = macrostegia.
Burkeana, Sond. in Linnaea, xxiii. (1850) 92 = macrostegia.
buxifolia, *Linn. Sp. Pl.* 636.—Ind. or.
buxifolia, Roxb. Hort. Beng. 45 ; Fl. Ind. iii. 37 = obovata.
buxifolia, Wall. Cat. n. 2504 = mysorensis.
caerulea, Herb. Lamb. ex Nees, in DC. Prod. xi. 228 = cristata.
caerulea, Roxb. Hort. Beng. 45 ; Fl. Ind. iii. 39 = strigosa.
caerulea, Wall. Cat. n. 2509, partim = polytricha.
candida, *Nees, in DC. Prod.* xi. 240.—Oriens.
capitata, *Klotzsch, in Peters, Reise Mossamb. Bot.* 210.—Afr. trop.
cardiocalyx, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 105.—Afr. trop.
Carruthersiana, *S. Moore, in Journ. Bot.* xviii. (1880) 270.—Afr. trop.
chegosa, Herb. Madr. ex Wall. Cat. n. 2506 = cristata.
ciliata, E. Mey. ex Nees, in DC. Prod. xi. 230 = Meyeriana.
ciliata, Roxb. Hort. Beng. 45 ; Fl. Ind. iii. 38 = cristata.
coccinea, Linn. Sp. Pl. 637 = Ruellia coccinea.
consanguinea, *Klotzsch, in Peters, Reise Mossamb. Bot.* 206.—Afr. trop.
conspicua, *Nees, in Wall. Pl. As. Rar.* iii. 92.—Ins. Penang.
cordifolia, Hochst. ex T. Anders. in Journ. Linn. Soc. vii. (1864) 29 = parviflora.
cornigera, Véry, ex Nees, in DC. Prod. xi. 248 = Hygrophila spinosa.
courtallica, *Nees, in Wight, Ic.* iv. t. 1529.—Ind. or.
cristata, Lam. Encyc. i. 380 = noctiflora.
cristata, *Linn. Sp. Pl.* 636.—Ind. or. ; Burma.
cuspidata, *Heyne, ex Nees, in Wall. Pl. As. Rar.* iii. 93.—Ind. or.
cyanea, *S. Moore, in Journ. Bot.* xviii. (1880) 265.—Angola.
damarensis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 29.—Afr. austr. trop.
Decaisneana, *Nees, in DC. Prod.* xi. 230.—Madag.
diacantha, Hochst. ex Nees, l. c. 238 = B. Prionitis.
diandra, Heyne, ex Nees, in Wall. Pl. As. Rar. iii. 92 = polytricha ?
diandra, Hochst. & Steud. ex Nees, in DC. Prod. xi. 231 = Hochstetteri.
diandra, Schlecht. ex Nees, l. c. = obtusa.
dianthera, Herb. Madr. ex Wall. Cat. n. 2391 = Peristrophe montana.
dichotoma, Roxb. Hort. Beng. 45 ; Fl. Ind. iii. 39 = cristata.
diffusa, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 518 = Lepidagathis diffusa.
discolor, *Nees, in Benth. Bot. Voy. Sulph.* 146.—Mexic.
echinata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = B. Prionitis.
elata, Dalz. in Hook. Kew Journ. iii. (1851) 227 = involucrata.
eranthemoides, R. Br. in Salt, Abyss. App. 64 = acanthoides.
fasciculata, Russ. ex Wall. Cat. n. 2509 = polytricha ?

BARLERIA :—
fissiflora, *Boj. ex Nees, in DC. Prod.* xi. 238.—Madag.
flava, *Jacq. Ecl.* 67. t. 46.—Arab.
gentianoides, Desf. ex Nees, in DC. Prod. xi. 224 = flava.
Gibsoni, *Dalz. in Hook. Kew Journ.* ii. (1850) 339.—Ind. or.
glabra, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 8.—Malaya.
glabrata, Vahl, ex Nees, in DC. Prod. xi. 248 = Hygrophila spinosa.
glandulosa, Hochst. ex Nees, l. c. 243 = Lepidagathis glandulosa.
glauca, T. Anders. in Journ. Linn. Soc. ix. (1867) 493 = Gibsoni.
grandiflora, *Dalz. in Hook. Kew Journ.* ii. (1850) 339.—Ind. or.
grandiflora, *R. Br. in Salt, Abyss. App.* 64.—Abyssin.
grandifolia, R. Br. ex T. Anders. in Journ. Linn. Soc. vii. (1864) 30, sphalm. = grandiflora.
grandis, Hochst. ex Nees, in DC. Prod. xi. 233 = grandiflora.
Grantii, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 127.—Afr. trop.
Gueinzii, *Sond. in Linnaea*, xxiii. (1850) 91.—Afr. austr.
halimoides, Nees, in DC. Prod. xi. 231 = Petalidium halimoides.
Harnierii, *Solms, in Schweinf. Beitr. Fl. Aethiop.* 106.—Afr. trop.
hexacantha, Bertol. in Ann. Stor. Nat. Bolog. fasc. ix. 445 ; Disq. Pl. Nov. 3 (1832) = longifolia.
hexacantha, Moris, in Mem. Acc. Torin. xxxvi. (1833) 200. t. 7 = Hygrophila spinosa.
Hildebrandtii, *S. Moore, in Journ. Bot.* xv. (1877) 69.—Afr. trop.
hirsuta, Herb. Madr. ex Wall. Cat. n. 2503 = tomentosa.
hirsuta, Jacq. Obs. Bot. ii. 7 = Gonzalea spicata.
hirsuta, Nees, Wall. Pl. As. Rar. iii. 91 ; Cat. n. 7142 = polytricha.
Hochstetteri, *Nees, ex DC. Prod.* xi. 231.—Afr. bor. ; Arab. ; Scind.
hypocrateriformis, Hochst. ex T. Anders. in Journ. Linn. Soc. vii. (1863) 28 = B. Prionitis.
Hystrix, Linn. Mant. i. 89 = B. Prionitis.
ilicina, *E. Mey. ex T. Anders. in Journ. Linn. Soc.* vii. (1864) 28.—Afr. austr.
inaequalis, Benth. in Flora, xxxii. (1849) 557 = courtallica.
inaequalis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 149 = Phaylopsis longifolia.
indica, Linn. ex T. Anders. in Journ. Linn. Soc. vii. (1864) 115 = cristata.
involucrata, *Nees, in Wall. Pl. As. Rar.* iii. 92.—Ind. or.
irritans, *Nees, in Linnaea*, xv. (1841) 359.—Afr. austr.
Kirkii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 30.—Afr. trop.
laciniata, Wall. Pl. As. Rar. iii. 91 ; Cat. n. 7144 = cristata.
lactea, Desf. ex Steud. Nom. ed. II. i. 187 = cristata.
lancifolia, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 28.—Afr. austr. trop.
Lawii, *T. Anders. l. c.* ix. (1867) 492.—Ind. or.
Lichtensteiniana, *Nees, in DC. Prod.* xi. 235.—Afr. austr.
limnogeton, *S. Moore, in Journ. Linn. Soc.* xv. (1877) 95.—Ins. Celebes.
longiflora, *Linn. f. Suppl.* 290.—Ind. or.
longifolia, T. Anders. in Journ. Linn. Soc. ix. (1867) 491 = longiflora.
longifolia, Linn. Cent. Pl. ii. 22 ; Amoen. Acad. iv. 320 = Hygrophila spinosa.
lupulina, *Lindl. in Bot. Reg.* t. 1483.—Ind. or. ; Ins. Maurit.
Mackenii, *Hook. f. Bot. Mag.* t. 5866.—Afr. austr.
Macraei, *Arn. in Hook. Comp. Bot. Mag.* ii. (1836) 311.—Ind. or.
macracantha, R. Br. in Salt, Abyss. App. 64 = Hygrophila spinosa ?
macrolemma, *Peyr. in Sitzb. Acad. Wien. Math.-Nat.* xxxviii. (1860) 574.—Afr. trop.
macrophylla, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 489 = strigosa.
macrostachya, Boj. Hort. Maurit. 260 = lupulina.
macrostegia, *Nees, in DC. Prod.* xi. 235.—Afr. austr.

BARLERIA :—
marginata, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 127.—Afr. trop.
Meyeriana, *Nees, in DC. Prod.* xi. 230.—Afr. austr.
micans, *Nees, in Benth. Bot. Voy. Sulph.* 146.—N. Granat.
mitis, *Ker-Gawl. in Bot. Reg.* t. 191.—Ind. or.
mollis, *R. Br. in Salt, Abyss. App.* 64.—Abyssin.
monostachya, *Boj. in Hook. Journ. Bot.* i. (1834) 356.—Ins. Maurit.
montana, *Herb. Madr. ex Nees, in Wall. Pl. As. Rar.* iii. 92.—Ind. or.
montana, Wight & Nees, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 487, pro parte = Gibsoni.
multiflora, *Hassk. Cat. Fl. Bogor. Alt.* 149.—Hab. ?
mysorensis, *Heyne, in Roth, Nov. Sp. Pl.* 313.—Ind. or.
napalensis, Nees, in Wall. Pl. As. Rar. iii. 91 = cristata.
nitida, *Jacq. ex Steud. Nom.* ed. II. i. 187.—Am. austr.
nitida, *Nees, in Wall. Pl. As. Rar.* iii. 91.—Ind. or.
noctiflora, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 145 = acanthoides.
noctiflora, *Linn. f. Suppl.* 290.—Ind. or. ; Arab.
nuda, Nees, in Wall. Pl. As. Rar. iii. 92 = cristata.
nutans, *Nees, in DC. Prod.* xi. 227.—Zeylan.
obovata, Buch.-Ham. in Trans. Linn. Soc. xiv. (1825) 293 = mysorensis.
obtusa, *Nees, in Linnaea*, xv. (1841) 358.—Afr. austr.
oenotheroides, Dum.-Cours. Bot. Cult. Cat. vii. 7 = flava.
opaca, *Nees, in DC. Prod.* xi. 230.—Afr. occ.
orbicularis, *Hochst. ex Anders, in Journ. Linn. Soc.* vii. (1864) 29.—Abyssin.
ovata, *E. Mey. ex Nees, in DC. Prod.* xi. 230.—Afr. austr.
paniculata, Herb. Madr. ex Nees, in Wall. Pl. As. Rar. iii. 92 = tomentosa.
papillosa, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 31.—Afr. austr.
parviflora, *R. Br. in Salt, Abyss. App.* 65.—Abyssin.
pentandra, Arn. ex Nees, in DC. Prod. xi. 232 = involucrata et Arnottiana.
pilosa, *Heyne, ex Nees, in Wall. Pl. As. Rar.* iii. 93.—Ind. or.
polyneura, *S. Moore, in Journ. Bot.* xviii. (1880) 266.—Afr. trop.
polystachya, Hook. ex Nees, in DC. Prod. xi. 226 = strigosa.
polytricha, Boj. ex Nees, l. c. 286 (= *Polythrix stenandrium*).—Madag.
polytricha, *Wall. Pl. As. Rar.* i. 72. t. 82.—Ind. or.
prinoides, Hort. Monsp. ex Nees, in DC. Prod. xi. 229 = cristata.
Prionitis, *Linn. Sp. Pl.* 636.—As. et Afr. trop.
procumbens, *Lour. Fl. Cochinch.* 377.—China.
pubiflora, Benth. in Flora, xxxii. (1849) 558 = B. Prionitis.
pumila, Hochst. ex Nees, in DC. Prod. xi. 372 = Justicia simplex.
pungens, Heyne, ex Nees, in Wall. Pl. As. Rar. iii. 91 = polytricha ?
pungens, *Linn. f. Suppl.* 290.—Afr. austr.
pungens, Thunb. Prod. Pl. Cap. 98 = irritans.
pungens, Wall. Cat. n. 2498 = Lepidagathis pungens.
purpurea, Lodd. Bot. Cab. iv. t. 344 = montana.
pyramidalis, Lam. ex Nees, in DC. Prod. 243 = Blechum Brownei.
quadrispinosa, Stokes, Bot. Mat. Med. iii. 411 = B. Prionitis.
querimbensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 205.—Afr. trop.
racemosa, Herb. Madr. ex Wall. Cat. n. 2503 = tomentosa.
ramosa, Buch.-Ham. in Trans. Linn. Soc. xiv. (1825) 295 = noctiflora.
repens, *Nees, in DC. Prod.* xi. 230.—Afr. austr.
rhynchocarpa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 205.—Afr. trop.
Richardiana, *Nees, in DC. Prod.* xi. 225.—Madag.
rigida, Spreng. in Zeyh. ex Schlecht. in Linnaea, xiv. (1840) 304 = Pachypodium tomentosum.
rigida, *Willd. ex Nees, in DC. Prod.* xi. 242.—Afr. austr.
rubra, *Buch.-Ham. in Trans. Linn. Soc.* xiv. (1825) 296 (sp. dub.).—Ind. or.
ruellioides, Rottl. ex Nees, in DC. Prod. xi. 240 = nodiflora.

BARLERIA :—
ruellioides, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 30.—Afr. trop.
salicifolia, *S. Moore, in Journ. Bot.* xviii. (1880) 268.—Afr. trop.
satureioides, *Griseb. Cat. Pl. Cub.* 195.—Cuba.
senegalensis, Nees, in DC. Prod. xi. 224 = flava.
senensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 209.—Afr. trop.
sepalosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 487.—Ind. or.
sericea, T. Anders. in Journ. Linn. Soc. vii. (1864) 115 = longiflora.
solanifolia, Linn. Sp. Pl. 637 = Barleriola solanifolia.
somalensis, *Franch. in Révoil, Çomali*, 51.—Afr. or.
spicata, Roxb. Hort. Beng. 45 = B. Prionitis.
Spina-ceylanica, Nees, in DC. Prod. xi. 242 = mysorensis.
spinicyma, *Nees, l. c.* 240.—Arabia.
spinosa, Hook. ex Nees, l. c. 247 = Hygrophila spinosa.
spinosa, Wall. Cat. n. 2497 = Lepidagathis spinosa.
spinulosa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 208.—Afr. trop.
squarrosa, *Klotzsch, l. c.* 207.—Afr. trop.
stellato-tomentosa, *S. Moore, in Journ. Bot.* xviii. (1880) 268.—Afr. trop.
stenophylla, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 95.—Burma.
stimulans, *E. Mey. ex Nees, in DC. Prod.* xi. 241.—Afr. austr.
Stocksii, *T. Anders. in Journ. Linn. Soc.* ix. (1867) 493.—Ind. or.
strigosa, *Willd. Sp. Pl.* iii. 379.—Ind. or.
terminalis, Nees, in DC. Prod. xi. 225 = strigosa.
tetracantha, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 85.—Ins. Socotr.
tomentosa, *Roth, Nov. Sp. Pl.* 314.—Ind. or.
tomentosa, Herb. Madr. ex Wall. Cat. n. 2503 = acuminata.
triacantha, Hochst. ex Nees, in DC. Prod. xi. 240 = acanthoides.
trinervis, Herb. Ham. ex Nees, l. c. 243 = Lepidagathis trinervis.
trispinosa, *Vahl, Symb. Bot.* i. 46.—Arab.
uitenagensis, Hochst. ex Nees, in DC. Prod. xi. 231 = obtusa.
unilateralis, Buch.-Ham. ex Wall. Cat. n. 2509 = polytricha.
ventricosa, *Hochst. ex Nees, in DC. Prod.* xi. 230.—Abyss.
venulosa, Nees, l. c. 228 = cristata.
vestita, *T. Anders. in Thw. Enum. Pl. Zeyl.* 230.—Zeylan.
villosa, *S. Moore, in Journ. Bot.* xviii. (1880) 267.—Afr. trop.
violascens, *S. Moore, l. c.* 265.—Afr. trop.
Welwitschii, *S. Moore, l. c.* 267.—Afr. trop.

BARLERIACANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 136 = **Barleria**, Linn. (Acanth.).
acanthoides, Oerst. l. c. = Barleria acanthoides.
noctiflora, Oerst. l. c. = Barleria noctiflora.

BARLERIANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 136 = **Barleria**, Linn. (Acanth.).
grandis, Oerst. l. c. = Barleria grandiflora.

BARLERIOLA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 136. *ACANTHACEAE*, Benth. & Hook. f. ii. 1093.
solanifolia, *Oerst. l. c.*—Ins. S. Doming.

BARLERIOPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 133 = **Barleria**, Linn. (Acanth.).
discolor, Oerst. l. c. 134 = Barleria discolor.
glandulosa, Oerst. l. c. = Barleria micans.
micans, Oerst. l. c. = Barleria micans.

BARLERIOSIPHON, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 136 = **Barleria**, Linn. (Acanth.).
longiflora, Oerst. l. c. = Barleria longiflora.

BARLERITES, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 137 = **Barleria**, Linn. (Acanth.).
Hochstetteri, Oerst. l. c. = Barleria Hochstetteri.

BARLIA, Parl. Nouv. Gen. Piant. Monocot. 5 (1858); Fl. Ital. iii. 445 (1858) = **Orchis**, Linn.
longibracteata, Parl. l. c. 6; Fl. Ital. iii. 447 = O. longibracteata.

BARNADESIA, Mutis, in Linn. f. Suppl. 55 (1781). *COMPOSITAE*, Benth. & Hook. f. ii. 484.
BACAZIA, Ruiz & Pav. Prod. Fl. Per. 105. t. 22 (1794).
DIACANTHA, Less. in Linnaea, v. (1830) 243.
DOLICHOSTYLIS, Cass. in Dict. Sc. Nat. lvi. 139 (1828).
FULCALDEA, Poir. Encyc. Suppl. v. 375 (1817).
PENTHEA, Spach, Hist. Vég. Phan. x. 9 (1841).
RHODACTINIA, Gardn. in Hook. Lond. Journ. vi. (1847) 449.
TURPINIA, Humb. & Bonpl. Pl. Aequin. i. 113. t. 33 (1808).
VOIGTIA, Spreng. Syst. iii. 367 (1826).
XENOPHONTA, Vell. Fl. Flum. 346 (1825).
arborea, *H. B. & K. Nov. Gen. et Sp.* iv. 16.—Ecuador.
arborea, Seh. Bip. in Linnaea, xxii. (1849) 569 = rosea.
berberoides, *Sch. Bip. in Lechl. Berb. Am. Austr.* 57.—Peruv.
corymbosa, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 278.—Peruv.
dianthiflora, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 365 = rosea.
divaricata, *Griseb. in Goett. Abh.* xxiv. (1879) 209.—Reg. Argent.
Dombeyana, *Less. in Linnaea*, v. (1830) 246.—Peruv.
flavescens, *Meyen, Reise*, i 307.—Chili.
glochidiata, Mart. ex Baker, in Mart. Fl. Bras. vi. III. 361 = synacantha.
hirsuta, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 527, nomen.—Bolivia.
lanata, *Meyen, Reise*, i. 347.—Chili.
lanceolata, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 277.—Peruv.
laxa, D. Don, l. c. 280 = rosea.
media, *D. Don, l. c.* 276.—Peruv.
odorata, *Griseb. in Goett. Abh.* xxiv. (1879) 210.—Reg. Argent.
parviflora, *Spruce, ex Benth. & Hook. f. Gen.* ii. 485.—Ecuador.
polyacantha, *Wedd. Chlor. And.* i. 13.—Bolivia.
reticulata, *D. Don, in Trans. Linn. Soc.* xvi. (1833) 277.—Peruv.
rosea, *Lindl. in Bot. Reg.* (1843) t. 29.—Am. austr.
spinosa, *Linn. f. Suppl.* 348.—Peruv.; N. Granat.
spinosa, Willd. ex DC. Prod. vii. 3 = arborea.
ulicina, Hook. & Arn. Bot. Beech. Voy. 29 = Chuquiraga ulicina.

BARNARDIA, Lindl. in Bot. Reg. t. 1029 (1826) = **Scilla**, Linn. (Liliac.).
indica, Wight, Ic. t. 2041 = S. indica.
japonica, Schult. f. Syst. vii. 555 = S. japonica.
scilloides, Lindl. in Bot. Reg. t. 1029 = S. chinensis.

BARNEOUDIA, C. Gay, Fl. Chil. i. 29. t. 1 (1845) = **Anemone**, Linn. (Ranunc.).
chilensis, C. Gay, l. c.—Chili.
Domeykoana, Leybold, in Anal. Univ. Chil. (1858) 159; et in Flora, xlii. (1859) 242.—Chili.
major, Phil. in Linnaea, xxviii. (1856) 609.—Chili.

BAROLA, Adans. Fam. ii. 344 (1763) = **Trichilia**, Linn. (Meliac.).

BAROLLAEA, Neck. Elem. ii. 322 (1790) = **Caryocar**, Linn. (Ternstroem.).

BARONIA, Baker, in Journ. Bot. xx. (1882) 67. *ANACARDIACEAE*.
Taratana, *Baker, l. c.*—Madag.

BAROSMA, Willd. Enum. Hort. Berol. 257 (1809). *RUTACEAE*, Benth. & Hook. f. i. 290.
BARYOSMA, Roem. & Schult. Syst. v. 448 (1819).
BUCCO, Wendl. Collect. i. 13 (1808).
PARAPETALIFERA, Wendl. Collect. t. 15, 34 (1808).
acutata, *Sond. in Harv. & Sond. Fl. Cap.* i. 395.—Afr. austr.
alpina, *Eckl. & Zeyh. Enum.* 102.—Afr. austr.
angustifolia, *Bartl. & Wendl. f. Diosm.* 116.—Afr. austr.
apiculata, *Eckl. & Zeyh. Enum.* 104.—Afr. austr.
betulina, *Bartl. & Wendl. f. Diosm.* 102.—Afr. austr.
bruniades, Hoffmgg. Verz. Pfl. Nachtr. i. 128 = Agathosma cuspidata.
crenata, *Eckl. & Zeyh. Enum.* 102.—Afr. austr.

BAROSMA :—
crenata, *Sweet, Hort. Brit.* ed. I. 89.—Afr. austr.
crenulata, *Hook. Bot. Mag.* t. 3413.—Afr. austr.
dioica, *Bartl. & Wendl. f. Diosm.* 114.—Afr. austr.
Eckloniana, *Bartl. in Linnaea*, xvii. (1843) 363.—Afr. austr.
foetidissima, *Bartl. & Wendl. f. Diosm.* 118.—Afr. austr.
gnidioides, *Eckl. & Zeyh. Enum.* 104.—Afr. austr.
graveolens, *G. Don, Gen. Syst.* i. 786.—Afr. austr.
Kraussiana, *Meissn. in Flora*, xxvii. (1844) I. 303.—Afr. austr.
lanceolata, *Sond. in Harv. & Sond. Fl. Cap.* i. 397.—Afr. austr.
latifolia, *Roem. & Schult. Syst.* v. 449.—Afr. austr.
microcarpa, *Sond. in Harv. & Sond. Fl. Cap.* i. 398.—Afr. austr.
mucronata, *Meissn. in Flora*, xxvii. (1844) I. 303.—Afr. austr.
Niveni, *Sond. in Harv. & Sond. Fl. Cap.* i. 398.—Afr. austr.
oblonga, *Bartl. & Wendl. f. Diosm.* 112.—Afr. austr.
odorata, Willd. Enum. Hort. Berol. 257 = serratifolia.
orbicularis, Hort. ex Sweet, Hort. Brit. ed. I. 89 = betulina.
ovata, *Bartl. & Wendl. f. Diosm.* 109.—Afr. austr.
pauciflora, *Eckl. & Zeyh. Enum.* 103.—Afr. austr.
pulchella, *Bartl. & Wendl. f. Diosm.* 107.—Afr. austr.
pulchra, *Cham. in Linnaea*, v. (1830) 53.—Afr. austr.
pungens, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 397.—Afr. austr.
scoparia, *Eckl. & Zeyh. Enum.* 103.—Afr. austr.
serratifolia, *Willd. Enum. Hort. Berol.* 257.—Afr. austr.
sertuliflora, *Delponte, in Mem. Acc. Torino*, Ser. II. xiv. (1854) 408. t. 9.—Hab. ?
ternata, *Eckl. & Zeyh. Enum.* 103.—Afr. austr.
trichopodis, *Bartl. & Wendl. f. ex Juss. in Mém. Mus. Par.* xii. (1825) 474.—Afr. austr.
venusta, *Eckl. & Zeyh. Enum.* 102.—Afr. austr.

BARRALDEIA, Thou. Gen. Nov. Madag. 24 (1809) = **Carallia**, Roxb. (Rhizoph.).
madagascariensis, Spreng. Syst. ii. 313 (Baraultia).—Madag.

BARRATTIA, A. Gray, in Proc. Am. Acad. i. (1848) 48 = BARATTIA, A. Gray & Engelm. = **Enceelia**, Adans. (Compos.).

BARRELIERA, J. F. Gmel. Syst. 959 (1791) = **Barleria**, Linn. (Acanth.).

BARRERIA, Linn. Gen. ed. II. 131 (1742); (et BARRERA, Linn. Sp. Pl. ed. II. 393 (1762)).
capensis, Linn. Sp. Pl. 274.—Afr. austr.

BARRERIA, Willd. Sp. Pl. i. 1145 (1798) = **Poraqueiba**, Aubl. (Olacineae).
gujanensis, Raeusch. Nom. ed. III. 259, nomen = P. guianensis.
theobromaefolia, Willd. Sp. Pl. i. 1145 = P. guianensis.

BARRINGTONIA, Forst. Char. Gen. 75. t. 38 (1776). *MYRTACEAE*, Benth. & Hook. f. i. 720.
AGASTA, Miers, in Trans. Linn. Soc. Ser. II. Bot. i. 59 (1875).
BARANDA, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 502.
BOTRYOROPIS, Presl, Epim. Bot. 220 (1849).
BUTONICA, Juss. Gen. 326 (1789).
DOXOMMA, Miers, in Trans. Linn. Soc. Ser. II. Bot. i. (1875) 98.
MEGADENDRON, Miers, l. c. 109 (1875).
MENICHEA, Sonner. Voy. N. Guin. 133 (1776).
METEORUS, Lour. Fl. Cochinch. 410 (1790).
MITRARIA, J. F. Gmel. Syst. 799 (1791).
STRAVADIUM, Juss. Gen. 326 (1789).
acuminata, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 206.—Borneo.
acutangula, *Gaertn. Fruct.* ii. 97. t. 101.—Ins. Molucc.
alata, Wall. Cat. n. 3633 = conoidea.
alba, *Blume, in Fl. des Serres*, Ser. I. vii. (1851-52) 23.—Ins. Molucc.
alba, Kostel. Allg. Med.-Pharm. Fl. iv. 1536 = racemosa.
asiatica, Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 131; xlvi. (1877) II. 70 = speciosa.
angusta, *Kurz, l. c.* xlii. (1873) II. 233.—Burma.
Butonica, *Forst. Char. Gen.* t. 38.—Ins. Pacif.

BARRINGTONIA :—
calyptrata, R. Br. ex Benth. Fl. Austral. iii. 288, in nota = speciosa ?
Careya, *F. Muell. Fragm.* v. 183.—Austral.
ceylanica, *Gardn. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 508.—Zeylan.
coccinea, Kostel. Allg. Med.-Pharm. Fl. iv. 1536 = excelsa.
conoidea, *Griff. Notul.* iv. 656 ; *Ic. Pl. As.* t. 635, 636. f. 1.—Burma ; Malacca.
costata, *Miq. Fl. Ind. Bat.* i. 489.—Archip. Sunda.
cylindrostachya, Griff. Notul. iv. 655 = macrostachya.
edulis, *Seem. Fl. Vit.* 82.—Ins. Fiji.
elongata, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 206. —Java.
excelsa, *Blume, Bijdr.* 1097.—Malaya.
Helferi, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 509. —Burma.
Horsfieldii, *Miq. Fl. Ind. Bat.* i. 489.—Java.
inclyta, Miers, in Trans. Linn. Soc. Ser. II. Bot. i. (1876) 71. t. 14 = racemosa.
insignis, *Miq. Fl. Ind. Bat.* i. 488.—Malaya.
intermedia, *Vieill. in Bull. Soc. Linn. Normand.* x. (1866) 98.—N. Caled.
luzoniensis, *Vidal, Phan. Cuming. Philipp.* 13, 48, 113. —Ins. Philipp.
macrocarpa, *Hassk. Cat. Hort. Bogor. Alt.* 263.—Java.
macrophylla, Miq. Fl. Ind. Bat. i. 491 = speciosa.
macrostachya, *Kurz, For. Fl. Brit. Burma*, i. 498 ; *et in Journ. As. Soc. Beng.* xlvi. (1877) II. 71.—Burma.
Montrouzieri, *Vieill. in Bull. Soc. Linn. Normand.* x. (1866) 100.—N. Caled.
neocaledonica, *Vieill. l. c.* 99.—N. Caled.
nitida, *Miq. Fl. Ind. Bat.* i. 490.—Java.
pendula, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 71.—Burma.
pterocarpa, *Kurz, l. c.* xlii. (1873) II. 234.—Burma.
pubescens, Miers, in Trans. Linn. Soc. Ser. II. Bot. i. (1876) 83 = acutangula.
racemosa, *Roxb. Hort. Beng.* 52 ; *Fl. Ind.* ii. 634.— Ind. or. ; Malaya ; Polynes.
Reinwardtii, *Miq. Fl. Ind. Bat.* i. 488.—Java.
reticulata, *Miq. l. c.* 490.—Borneo.
rigida, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 510 = Symplocos rigida.
rubra, Blume, in Fl. des Serres, Sér. I. vii. (1851–52) 23 = racemosa.
samoënsis, *A. Gray, Bot. U. St. Expl. Exped.* i. 508.— Ins. Fiji.
sarcostachys, Miq. Fl. Ind. Bat. i. 490 = macrostachya.
Senequli, *Jardin, in Bull. Soc. Linn. Normand.* Sér. II. ix. (1875) 305.—N. Caled.
serrata, *Miq. Fl. Ind. Bat.* i. 488.—Java.
speciosa, Linn. f. Suppl. 312 = B. Butonica.
speciosa, Wall. Cat. n. 3632 B = racemosa.
spicata, *Blume, Bijdr.* 1097.—Java.
Stravadium, Blanco, Fl. Filip. ed. I. 533 = racemosa.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* 315.—Sumatra.
timorensis, *Blume, in Van Houtte, Fl. des Serres*, Sér. I. vii. (1851–52) 23.—Ins. Timor.
Vriesei, *Teijsm. & Binn. in Tijdschr. Neêrl. Ind.* ii. (1851) 308.—Java.

BARROËTIA, A. Gray, in Proc. Am. Acad. xv. (1880) 29. *COMPOSITAE.*
Pavonii, *A. Gray, l. c.* xvii. (1881–82) 206.—Mexic.
setosa, *A. Gray, l. c.* xv. (1880) 29.—Mexic.
subuligera, *A. Gray, l. c.*—Mexic.

BARROTIA, Gaudich. Bot. Voy. Bonite, t. 13 (1852 ?) = **Pandanus**, Linn.
altissima, Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 277 = P. Minda.
aragoënsis, Brongn. l. c. 278 = P. aragoënsis.
Balansae, Brongn. l. c. 281 = P. Balansae.
decumbens, Brongn. l. c. 285 = P. decumbens.
diodon, Gaudich. Bot. Voy. Bonite, t. 13. f. 9–14 = P. furcatus.
Gaudichaudii, Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 264 = P. tetrodon.
macrocarpa, Brongn. l. c. 279 = P. macrocarpus.
monodon, Gaudich. Bot. Voy. Bonite, t. 13. f. 15–24 = P. monodon.
Pancheri, Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 283 = P. Pancheri.
sphaerocephala, Brongn. l. c. 284 = P. sphaerocephalus.
tetrodon, Gaudich. Bot. Voy. Bonite, t. 13. f. 1–8 = P. tetrodon.

BARROWIA, Decne. in DC. Prod. viii. 629 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 779.
jasminiflora, *Decne. l. c.* 630.—Afr. austr.

BARTERIA, Hook. f. in Journ. Linn. Soc. v. (1861) 14. t. 2. *PASSIFLOREAE*, Benth. & Hook. f. i. 812.
fistulosa, *Mast. in Oliver, Fl. Trop. Afr.* ii. 511.—Afr. trop.
nigritana, *Hook. f. in Journ. Linn. Soc.* v. (1861) 15. t. 2.—Afr. trop.

BARTHEA, Hook. f. in Benth. & Hook. f. Gen. i. 751 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
chinensis, *Hook. f. l. c.*—Hongkong.

BARTHESIA, Comm. ex A. DC. in Trans. Linn Soc. xvii. (1837) 112 = **Ardisia**, Sw. (Myrsin.).

BARTHOLINA, R. Br. in Ait. Hort. Kew. ed. II. v. 194 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 623.
LATHRISIA, Sw. Annot. Bot. 48 (1829).
Burmanniana, Ker-Gawl. in Brand. Journ. Sc. iv. (1818) 204. t. 5 = pectinata.
Ethelae, *Bolus, in Journ. Linn. Soc.* xx. (1884) 472.— Afr. austr.
Lindleyana, Reichb. f. Otia Bot. Hamb. 119 = pectinata.
pectinata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 194.— Afr. austr.

BARTHOLLESIA, Manso, Enum. 45 (1836) = **Bertholletia**, Humb. & Bonpl. (Myrtac.).

BARTHRATHERUM, Anderss. in Walp. Ann. iii. 799 (1853) = BATHRATHERUM, Hochst. = **Arthraxon**, Beauv. (Gramin.).

BARTLETTIA, A. Gray, in Mem. Am. Acad. Ser. II. v. (1855) (Pl. Nov. Thurb.) 323. *COMPOSITAE*, Benth. & Hook. f. ii. 441.
scaposa, *A. Gray, l. c.*—Mexic.

BARTLINGIA, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 373 = **Pultenaea**, Sm. (Legumin.).
obovata, Brongn. l. c. = P. obovata.

BARTLINGIA, F. Muell. ex Benth. Fl. Austral. vii. 63 (1878) = **Laxmannia**, R. Br. (Liliac.).
brachyphylla, F. Muell. Census, 119 = L. brachyphylla.
gracilis, F. Muell. Fragm. vii. 88 = L. gracilis.
grandiflora, F. Muell. Census, 118 = L. grandiflora.
minor, F. Muell. l. c. = L. ramosa.
ramosa, F. Muell. Fragm. vii. 88 = L. ramosa.
sessiliflora, F. Muell. in Papers Roy. Soc. Tasm. (1877) 116 = L. sessiliflora.
sessilis, F. Muell. Census, 119 = L. sessilis.
squarrosa, F. Muell. l. c. 118 = L. squarrosa.

BARTLINGIA, Reichb. in Flora, vii. (1824) 241 = **Plocama**, Ait. (Rubiac.).
scoparia, Reichb. l. c. = P. pendula.

BARTOLINA, Adans. Fam. ii. 124 (1763) = **Tridax**, Linn. (Compos.).

BARTONIA, Muhl. ex Willd. in Ges. Naturf. Freunde, Neue Schr. iii. (1801) 444. *GENTIANEAE*, Benth. & Hook. f. ii. 818.
ANDREWSIA, Spreng. Syst. i. 428 (1825).
CENTAURELLA, Michx. Fl. Bor. Am. i. 97. t. 12 (1803).
CENTAURIUM, Pers. Syn. i. 137 (1805).
NUTTALLA, Rafin. in Am. Monthly Mag. (1818) 175.
pubescens, Rafin. in Med. Repos. New York, v. (1808) 352 = verna ?
tenella, *Muhl. ex Willd. in Ges. Naturf. Freunde, Neue Schr.* iii. (1801) 445.—Am. bor.
uniflora, Rafin. in Med. Repos. New York, v. (1808) 352 = verna ?
verna, *Muhl. ex A. Gray, in Chapm. Fl. S. U. St.* 357. —Am. bor.

BARTONIA, Sims, Bot. Mag. t. 1487 (1804) = **Mentzelia**, Linn. (Loaseae).
albescens, Gill. & Arn. in Edinb. Geog. Journ. ii. (1831) 273.—Chili.
albicaulis, Dougl. ex Hook. Fl. Bor. Am. i. 222 = M. albicaulis.
aurea, Lindl. in Bot. Reg. t. 1831 = M. Lindleyi.

BARTONIA :—
decapetala, Sims, Bot. Mag. t. 1487 = Mentzelia ornata.
laevicaulis, Dougl. ex Hook. Fl. Bor. Am. i. 221. t. 69 = M. laevicaulis.
micrantha, Hook. & Arn. Bot. Beech. Voy. 343 = M. micrantha.
micrantha, Hook. in Lond. Journ. Bot. vi. (1847) 227 = M. dispersa.
multiflora, Nutt. in Journ. Acad. Sc. Phil. N. S. i. (1847) 180 = M. multiflora.
nuda, Pursh, Fl. Am. Sept. 328, 749 = M. nuda.
ornata, Hook. Lond. Journ. Bot. vi. (1847) 226 = M. laevicaulis.
ornata, Pursh, Fl. Am. Sept. 327 = M. ornata.
parviflora, Dougl. ex Hook. Fl. Bor. Am. i. 221 = M. laevicaulis.
pumila, Nutt. ex Torr. & Gray, Fl. N. Am. i. 535 = M. pumila.
sinuata, Presl, Rel. Haenk. ii. 38 = *albescens*.
Wrightii, Walp. Ann. ii. 656 = M. Wrightii.

BARTRAMIA, Linn. Diss. Dass. 17 (1747) ; Fl. Zeyl. 77 (1747) ; Gaertn. Fruct. ii. 137. t. 111 (1791) = **Triumfetta**, Linn. (Tiliac.).
angulata, Lam. Ill. t. 430. f. 2 = T. rhomboidea.
crispifolia, Stokes, Bot. Mat. Med. iii. 14 = T. rhomboidea.
glandulosa, Lam. Ill. t. 400. f. 1 = T. rhomboidea.
indica, Linn. Sp. Pl. 389 = T. rhomboidea.
Lappago, Gaertn. Fruct. ii. 137 t. 111 = T. rhomboidea.
rhombifolia, Stokes, Bot. Mat. Med. iii. 15 = T. rhomboidea.

BARTRAMIA, Salisb. Prod. 99 (1796) = **Penstemon**, Mitch. (Scrophular.).
concinna, Salisb. l. c. = P. pubescens.
pulchella, Salisb. l. c. = P. laevigata.

BARTSCHIA, Linn. Syst. ed. II. 26 (1740) = seq.

BARTSIA, Linn. Hort. Cliff. 325 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 977.
ALICOSTA, Dulac, Fl. Hautes-Pyr. 381 (1867).
BELLARDIA, All. Fl. Pedem. i. 61 (1785).
EUFRAGIA, Griseb. Spicil. Fl. Rumel. ii. 13 (1844).
LASIOPERA, Hoffmgg. & Link, Fl. Port. i. 298 (1809).
ODONTITES, [Riv.] Hall. Enum. Stirp. Helv. ii. 627 (1742).
PARENTUCELLIA, Viv. Fl. Lib. Spec. 31. t. 21. f. 3 (1824).
STAEHELINIA, Hall. Enum. Stirp. Helv. ii. 624 (1742).
TRIXAGO, Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 4.
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 134 = B. Trixago.
acuminata, Pursh, Fl. Am. Sept. ii. 429 = Castilleja pallida.
alpina, *Linn. Sp. Pl.* 602.—Europ. ; Am. bor.
aspera, *Willk. & Lange, Fl. Hisp.* ii. 614, obs.—Hispan. ; Marocc.
bicolor, DC. Ic. Rar. t. 10 = B. Trixago.
biloba, *Wedd. Chlor. And.* ii. 123.—Peruv.
breviflora, *Benth. in DC. Prod.* x. 545.—Calif.
canescens, *Wedd. Chlor. And.* ii. 123.—Peruv.
capensis, Spreng. Syst. ii. 773 = B. Trixago.
carnea, *Griseb. Spicil. Fl. Rumel.* ii. 12.—Macedon.
chilensis, *Benth. in DC. Prod.* x. 547.—Chili.
ciliolata, *Wedd. Chlor. And.* ii. 123.—Bolivia.
coccinea, Linn. Sp. Pl. 602 = Castilleja coccinea.
crenoloba, *Wedd. Chlor. And.* ii. 124.—Bolivia.
decurva, *Hochst. ex Benth. in DC. Prod.* x. 545.— Abyssin.
densiflora, *Benth. l. c.* 548.—Peruv.
diffusa, *Benth. l. c.* 546.—Peruv.
elongata, *Wedd. Chlor. And.* ii. 127.—Am. merid.
euphrasioides, *Wedd. l. c.* 130.—Ecuador.
Fagonii, Lapeyr. Hist. Abr. Pl. Pyr. 345 = aspera.
filiformis, *Wedd. Chlor. And.* ii. 126.—Peruv.
Garideli, Ball, in Journ. Linn. Soc. xvi. (1878) 602 = purpurea.
glauca, Poir. ex Steud. Nom. ed. II. i. 189 = Lagotis glauca.
gracilis, *Benth. in DC. Prod.* x. 548.—N. Granat.
granatensis, Ball, in Journ. Linn. Soc. xvi. (1878) 602 = purpurea.
grandiflora, Spreng. in Ersch & Grub. Allgem. Encyc. vii. (1821) 461 = Castilleja sessiliflora.
Gymnandra, Linn. f. Suppl. 278 = Lagotis glauca.
Gymnandra, Pursh, Fl. Am. Sept. ii. 430 = Synthyris rubra ?

BARTSIA:—
heterophylla, *Wedd. Chlor. And.* ii. 129.—N. Granat.
hispida, *Benth. in DC. Prod.* x. 547.—Peruv.
humilis, Lapeyr. Hist. Abr. Pl. Pyr. 344 = Euphrasia minima.
imbricata, Lapeyr. l. c. = Euphrasia officinalis.
inaequalis, *Benth. in DC. Prod.* x. 547.—Peruv.; N. Granat.
integrifolia, *Wedd. Chlor. And.* ii. 131.—Peruv.
lanceolata, *Reichb. f. Ic. Fl. Germ.* xx. 57. t. 107.—Europ.
laniflora, *Benth. in DC. Prod.* x. 546.—N. Granat.
laticrenata, Benth. l. c. 547 = santolinaefolia.
latifolia, *Sibth. & Sm. Fl. Graec.* vi. 69. t. 586.—Reg. Mediterr.; Reg. Cauc.; Persia.
laxiflora, *Benth. in DC. Prod.* x. 547.—N. Granat.
longiflora, *Hochst. ex Benth. l. c.* 545.—Abyss.
lutea, *Reichb. f. Ic. Fl. Germ.* xx. 56.—Europ.; As. Min.
maxima, Gren. & Godr. Fl. Fr. ii. 611, in syn. = viscosa.
maxima, Pers. Syn. ii. 151 = B. Trixago.
melampyroides, *Benth. in DC. Prod.* x. 548.—Ecuador.
Meyeniana, *Benth. l. c.* 546.—Peruv.
mollis, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. lviii. (1883). i. 397 = Veronica ciliata.
mutica, *Benth. l. c.* 548.—Peruv.
Odontites, *Huds. Fl. Angl.* 268.—Europ.; Reg. Mediterr.
orthocarpiflora, *Benth. in DC. Prod.* x. 545.—Peruv.
pallida, Linn. Sp. Pl. 602 = Castilleja pallida.
parviflora, Charpent. ex Benth. in DC. Prod. x. 544 = alpina.
parviflora, Thom. Cat. Pl. Suisses (1837) 6 = alpina.
parvifolia, *Benth in DC. Prod.* x. 545.—N. Granat.
patens, *Benth. l. c.* 546.—Peruv.
pedicularoides, *Benth. l. c.*—Ecuador.
peruviana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 400.—Peruv.
pumila, *Benth. in DC. Prod.* x. 546.—Peruv.
purpurea, *Ball, in Journ. Linn. Soc.* xvi. (1878) 602.—Hispan.; Afr. bor.
purpurea, Duby, Bot. Gall. 354 = latifolia.
rhinanthoides, Hochst. ex Benth. in DC. Prod. x. 543 = B. Trixago.
rubricoma, Pall. Reise, iii. 34 = Castilleja pallida.
santolinaefolia, *Benth. in DC. Prod.* x. 547.—Mexic.
scabra, *Spreng. Syst.* ii. 773.—Afr. austr.
scordifolia, *Steud. Nom.* ed. II. i. 189.—Ins. Mader.
serotina, Bertol. Amoen. Ital. 33 = B. Odontites.
sibirica, Pall. Reise, ii. 123 = Castilleja pallida.
spicata, Ramond, in Bull. Soc. Philom. ii. (1799–1801) = aspera.
stricta, *Benth. in DC. Prod.* x. 547.—Ecuador.
subinclusa, *Benth. l. c.*—Peruv.
subinclusa, Griseb. ex Wedd. Chlor. And. ii. 126 = filiformis.
tenuifolia, Pursh, Fl. Am. Sept. ii. 429 = Orthocarpus tenuifolius.
trichophylla, *Wedd. Chlor. And.* ii. 122.—Peruv.
trifida, Spreng. Syst. ii. 773 = Bungea trifida.
Trixago, *Linn. Sp. Pl.* 602.—Europ.; Afr. bor.
verna, *Reichb. f. Ic. Fl. Germ.* xx. 57. t. 107.—Europ.
versicolor, Pers. Syn. ii. 151 = B. Trixago.
viscosa, *Linn. Sp. Pl.* 602.—Europ.; Reg. Mediterr.

BARYA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 122; et in Abh. Akad. Berl. (1854) 22. t. 2, B = **Begonia**, Linn.
monadelpha, Klotzsch, ll. cc. 122, 23 = Begonia monadelpha.

BARYOSMA, Gaertn. Fruct. ii. 73. t. 93 (1791) = **Dipteryx**, Schreb. (Legumin.).
oppositifolium, Pers. Syn. ii. 278 = D. oppositifolia.
Tongo, Gaertn. Fruct. ii. t. 93 = D. odorata.

BARYOSMA, Roem. & Schult. Syst. v. 448 (1819) = **Barosma**, Willd. (Rutac.).

BARYSOMA, Bunge, Del. Sem. Hort. Dorpat. (1839) 7; et in Linnaea, xiv. (1840) Litt. 117 = **Heracleum**, Linn. (Umbellif.).
villosum, Bunge, ll. cc. = H. villosum.

BARYXYLUM, Lour. Fl. Cochinch. 266 (1790). *LEGUMINOSAE*?—Cf. Benth. & Hook. f. i. 464.
rufum, *Lour. l. c.*—China.

BASALTOGETON, Salisb. Gen. Pl. Fragm. 27 (1866) = **Scilla**, Linn. (Liliac.).
peruvianum, Salisb. l. c. = S. peruviana.

BASANACANTHA, Hook. f. in Benth. & Hook. f. Gen. ii. 82 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
armata, *Hook. f. l. c.* 83.—Am. trop.
dioica, *Hook. f. l. c.*—N. Granat.
hebecarpa, *Hook. f. l. c.*—Am. trop.
Humboldtiana, *Hook. f. l. c.*—Am. trop.
monantha, *Hook. f. l. c.*—Guatemala.
reticulata, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 98.—Mexic.
tetracantha, *Hook. f. in Benth. & Hook. f. Gen.* ii. 83.—Am. trop.

BASANANTHE, Peyr. in Bot. Zeit. xvii. (1859) 101; et in Sitzb. Acad. Wien. Math. Nat. xxxviii. (1860) 569. *PASSIFLOREAE*, Benth. & Hook. f. i. 812.
littoralis, Peyr. ll. cc.—Afr. trop.
Nummularia, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 28. t. 9.—Angola.

BASELA, Linn. Gen. ed. VI. (Vindob. reimpr.) 151 (1767) = seq.

BASELLA, [Rheede] Linn. Diss. Dass. 12 (1747); Amoen. Acad. i. 396. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 76.
GANDOLA, Rumph. ex Linn. Sp. Pl. ed. II. 390 (1762).
alba, Linn. Sp. Pl. 272 = rubra.
cananifolia, Buch.-Ham. ex Wall. Cat. n. 6961 = rubra.
cordifolia, Lam. Encyc. i. 382 = rubra.
crassifolia, Salisb. Prod. 153 = rubra.
diffusa, Ruiz & Pav. ex Moq. in DC. Prod. xiii. II. 226 (= *Tandonia diffusa*).—Peruv.
filiformis, Ruiz & Pav. ex Moq. l. c. 227 (= *Tandonia filiformis*).—Peruv.
Hookeriana, Moq. ex Moq. l. c. 225 = Tournonia Hookeriana.
japonica, Burm. f. Fl. Ind. 76. t. 39. f. 4 = rubra.
lucida, Linn. Syst. ed. X. 966. n. 343 = rubra.
marginata, H. B. & K. Nov. Gen. et Sp. ii. 189 (= *Tandonia marginata*).—Ecuador.
nigra, Lour. Fl. Cochinch. i. 183 = rubra.
obovata, H. B. & K. Nov. Gen. et Sp. ii. 189 (= *Tandonia obovata*).—N. Granat.
obtusifolia, Willd. ex Moq. in DC. Prod. xiii. II. 227 (= *Tandonia obovata*).—N. Granat.
ramosa, Jacq. f. in Spreng. Syst. i. 950 = rubra.
rubra, *Linn. Sp. Pl.* 272.—Reg. trop.
trifida, Willd. in Roem. & Schult. Syst. vi. 695 (= *Tandonia marginata*).—Ecuador.
tuberosa, H. B. & K. Nov. Gen. et Sp. ii. 189 = Ullucus tuberosus.
vesicaria, Lam, Illustr. t. 215. f. 1 = Anredera spicata.
volubilis, Salisb. Prod. 153 = rubra.

BASILAEA, Juss. ex Lam. Encyc. i. 382 (1783); et Gen. 52 (1789) = **Eucomis**, L'Hérit. (1788) (Liliac.).
bifolia, Mirb. Hist. Nat. Pl. viii. (1804) 339 = Whiteheadia bifolia.
coronata, Lam. Encyc. i. 382 = E. regia.
nana, Mirb. Hist. Nat. Pl. viii. (1804) 339 = E. nana.
punctata, Mirb. l. c.; Lam. Illustr. t. 289 = E. punctata.
regia, Mirb. l. c. 338 = E. regia.
undulata, Mirb. l. c. = E. undulata.

BASILEOPHYTA, F. Muell. First. Gen. Rep. 16 (1853) = **Fieldia**, A. Cunn. (Gesneriac.).
Friderici-Augusti, F. Muell. l. c. = F. australis.

BASILICUM, Moench, Meth. Suppl. 143 (1802) = **Ocimum**, Tourn. (Labiat.).
polystachion, Moench, l. c. = O. Basilicum.

BASILLAEA, R. Hedw. Gen. 239 (1809) = BASILAEA, Juss. = **Eucomis**, L'Hérit. (Liliac.).

BASILIMA, Rafin. New Fl. Am. iii. 75 (1836) = **Spiraea**, Linn. (Rosac.).
pygmea, Rafin. l. c. 76.—Am. bor.
sorbifolia, Rafin. l. c. 75 = S. sorbifolia?

BASISTEMON, Turcz. in Bull. Soc. Nat. Mosc. (1863) 214. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 1244.
bogotensis, *Turcz. l. c.*—N. Granat.
peruviana, *Benth. & Hook. f. Gen.* ii. 1245.—Peruv.

BASKERVILLA, Lindl. Gen. et Sp. Orch. 505 (1840). *ORCHIDEAE*, Benth. & Hook. f. iii. 597.
assurgens, *Lindl. l. c.*—Peruv.

BASONCA, Rafin. Fl. Tellur. iv. 68 (1836) = **Martynia**, Linn. (Pedalin.).
longiflora, Rafin. l. c. = M. longiflora.

BASSELINIA, Vieill. in Bull. Soc. Linn. Normand. Sér. II. vi. 1872 (1873) 230. *PALMAE*.
Deplanchei, *Brongn. & Gris, l. c.* 232, nomen.—N. Caled.
gracilis, *Brongn. & Gris, l. c.* 231, nomen.—N. Caled.
kanaliensis, *Vieill. l. c.* 232, nomen.—N. Caled.
Lenormandii, *Vieill. l. c.* nomen.—N. Caled.
Pancheri, *Brongn. & Gris, l. c.* nomen.—N. Caled.

BASSIA, All. Misc. Taur. iii. 177. t. 4 (1766) = **Chenolea**, Thunb. (1781) (Chenopod.).
astracantha, F. Muell. Census, 140 = C. astrocarpa.
astrocarpa, F. Muell. Fragm. xii. 12 = C. astrocarpa.
bicornis, F. Muell. Census, 30 = Sclerolaena bicornis.
bicuspis, F. Muell. l. c. = Anisacantha bicuspis.
biflora, F. Muell. l. c. = Sclerolaena biflora.
Birchii, F. Muell. l. c. = Anisacantha Birchii.
brevicuspis, F. Muell. l. c. = Anisacantha brevicuspis.
carnosa, F. Muell. l. c. = C. carnosa.
Cornishiana, F. Muell. in Melb. Chem. & Drug. (1885).—Austral.
Dallachyana, F. Muell. Census, 30 = C. Dallachyana.
diacantha, F. Muell. l. c. = Sclerochlaena diacantha.
divaricata, F. Muell. l. c. = Anisacantha divaricata.
Drummondii, F. Muell. l. c. = Anisacantha Drummondii.
echinopsila, F. Muell. l. c. = Anisacantha echinopsila.
enchylaenoides, F. Muell. l. c. = C. enchylaenoides.
Erskineana, F. Muell. in Melb. Chem. & Drug. (Apr. 1885).—N. Guinea.
eurotioides, F. Muell. Census, 30 = C. eurotioides.
Forrestiana, F. Muell. Fragm. xii. 12 = C. Forrestiana.
glabra, F. Muell. Census, 30 = Anisacantha glabra.
hirsuta, Aschers. ex Nym. Consp. 625 = C. hirsuta.
lanicuspis, F. Muell. Census, 30 = Sclerochlaena lanicuspis.
micrantha, F. Muell. l. c. = Enchylaena micrantha.
Muelleri, F. Muell. l. c. = C. Muelleri.
muricata, All. Misc. Taur. iii. t. 4 = C. hirsuta.
paradoxa, F. Muell. Census, 30 = Sclerolaena paradoxa.
quinquecuspis, F. Muell. l. c. = Anisacantha muricata.
salsuginosa, F. Muell. l. c. = Threlkeldia salsuginosa.
sclerolaenoides, F. Muell. l. c. = C. sclerolaenoides.
tricornis, F. Muell. l. c. = C. tricornis.
tridens, F. Muell. Fragm. xii. 12 = C. tridens.
uniflora, F. Muell. Census, 30 = Sclerolaena uniflora.

BASSIA, Koenig, ex Linn. Mant. ii. App. 555 (1771). *SAPOTACEAE*, Benth. & Hook. f. ii. 658.
DASYAULUS, Thw. Enum. Pl. Zeyl. 175 (1860).
ILLIPE, F. Muell. Extra-trop. Pl. Am. ed. [5] (1884) 181.
MADHUCA, J. F. Gmel. Syst. 799 (1791).
albescens, Griseb. Cat. Pl. Cub. 164 (= Labourdonnaisiae sp.).—Cuba occ.
amicorum, *A. Gray, in Proc. Am. Acad.* v. (1861–62) 327.—Ins. Pacif.
argentea, *De Vriese. Pl. Reinw.* 62.—Malacca.
Balem, *Miq. Fl. Ind. Bat. Suppl.* i. 582.—Sumatra.
Blancoi, *A. DC. in DC. Prod.* viii. 199.—Ins. Philipp.
butyracea, *Roxb. in As. Res.* viii. (1805) 477.—Ind. or.
caloneura, *Kurz, Andaman. Rep.* 141.—Ins. Andaman.
canaliculata, Bedd. Forest. Man. Bot. 141 = Dichopsis canaliculata.
Cocco, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 34.—N. Guinea.
cuneata, *Blume. Bijdr.* 675.—Malaya.
dubia, *Gaertn. Fruct.* ii. 104. t. 104.—Amboin.
elliptica, Dalz. in Hook. Kew Journ. iii. (1851) 36 = Dichopsis elliptica.
elongata, *Miq. Fl. Ind. Bat. Suppl.* 582.—Sumatra.
fulva, *Bedd. Forest. Man. Bot.* 140.—Ind. or.

BASSIA :—
galactoxyla, F. Muell. Fragm. vi. 27 = Lucuma galactoxylon.
grandis, Bedd. Fl. Sylv. ii. t. 254 = Dichopsis grandis.
hypoleuca, Miq. Fl. Ind. Bat. Suppl. 582 = Dichopsis obovata.
insignis, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* xii. (1882) 309.—Ind. or.
Jussaei, Griseb. Fl. Brit. W. Ind. 402 (sphalm.—Cf. Achras mammosa).—Ind. occ.
Korthalsii, *Pierre, ex Burck, in Ann. Jard. Buitenz.* v. (1885) 45.—Sumatra.
Krantzii, *Hance, in Journ. Bot.* xiv. (1876) 260.—Cambodia.
laevifolia, Bedd. Forest. Man. Bot. 141 = Dichopsis laevifolia.
lanceolata, Bedd. I. c. = Dichopsis lanceolata.
lanceolata, *A. DC. in DC. Prod.* viii. 199.—Ins. Philipp.
latifolia, *Roxb. Pl. Corom.* i. 20. t. 19.—Ind. or.
Lobbii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 546.—Burma.
lorgifolia, *Linn. Mant.* ii. *App.* 563.—Ind. or.
malabarica, *Bedd. Forest. Man. Bot.* 140.—Ind. or.
microphylla, *Hook. Ic. Pl.* i. t. 74.—Zeylan.
Moonii, *Bedd. Forest. Man. Bot.* 140.—Ind. or.
Mottleyana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 546.—Malaya.
neriifolia, *Bedd. Forest. Man. Bot.* 140.—Ind. or.
obovata, *Forst. f. Prod.* n. 200.—Ins. Tanna.
oleifera, *A. DC. in DC. Prod.* viii. 199.—Ins. Philipp.
pallida, *Burck, in Ann. Jard. Buitenz.* v. (1885) 44.—Sumatra.
Parkii, G. Don, Gen. Syst. iv. 36 = Butyrospermum Parkii.
parvifolia, A. DC. in DC. Prod. viii. 198 = microphylla.
pauciflora, Bedd. Forest. Man. Bot. 141 = Dichopsis pauciflora.
petiolaris, *Bedd. l. c.* 140.—Ind. or.
polyantha, Wall. Cat. n. 4166 = Dichopsis polyantha.
rubiginosa, Bedd. Forest. Man. Bot. 141 = Dichopsis rubiginosa.
sericea, *Blume, Bijdr.* 674.—Malaya.
villosa, Wall. Cat. n. 4165 = latifolia.
Wightiana, Bedd. Forest. Man. Bot. 141 = Isonandra Wightiana.

BASSOVIA, Aubl. Pl. Gui. i. 217. t. 85 (1775). *SOLANACEAE*, Benth. & Hook. f. ii. 892.
AURELIANA, Sendt. in Mart. Fl. Bras. x. 138. t. 19 (1846).
WITHERINGIA, L'Hérit. Sert. Angl. i. f. 1 (1788).
brachypoda, *Dun. in DC. Prod.* xiii. 409.—Peruv.
cornuta, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 59.—Bras.
crassiflora, *Dun. in DC. Prod.* xiii. 409.—Bras.
fasciculata, *Dun. l. c.* 408.—Bras.
Gardneri, *Dun. l. c.* 409.—Bras.
glomuliflora, *Dun. l. c.* 405.—Bras.
hebepoda, *Dun. l. c.* 407 = lucida?
laevis, *Dun. l. c.* 408.—Java.
leptopoda, *Dun. l. c.* 411.—Bras.
lucida, *Dun. l. c.* 406.—Guatemala.
pyraster, *Dun. l. c.* 408.—Bras.
Richardi, *Dun. l. c.* 406.—Guiana.
silvatica, Blanco, Fl. Filip. ed. II. 95 = Maesa indica.
solanacea, *Benth. & Hook. f. Gen.* ii. 891.—Hab.?
Spina-alba, *Griseb. in Goett. Abhandl.* xxiv. (1879) 248.—Am. bor.
sylvatica, *Aubl. Pl. Gui.* i. 217.—Guiana.
tomentosa, *Dun. in DC. Prod.* xiii. 409.—Bras.
velutina, *Dun. l. c.* 410.—Bras.
Wallichii, Dun. l. c. 409 = Solanum crassipetalum.

BASTARDIA, H. B. & K. Nov. Gen. et Sp. v. 254. t. 472 (1821). *MALVACEAE*, Benth. & Hook. f. i. 203.
BOGENHARDIA, Reichb. Nom. 200 (1841).
angulata, Guill. & Perr. Tent. Fl. Seneg. i. 65 = Abutilon intermedium.
aristata, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 200 = bivalvis.
bivalvis, *H. B. & K. Nov. Gen. et Sp.* v. 255, in nota.—Jamaica.
crispa, St. Hil. Fl. Bras. Mer. i. 194, in nota = Abutilon crispum.
foetida, Sweet, Hort. Brit. ed. II. 64 = Abutilon foetidum.
guayquilensis, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 200 = viscosa.
hirsutiflora, Presl, Reliq. Haenk. ii. 112 = viscosa.

BASTARDIA :—
hirsutissima, Walp. Rep. i. 327 = praec.
nemoralis, St. Hil. Fl. Bras. Mer. i. 195 = Abutilon crispum.
parvifolia, H. B. & K. Nov. Gen. & Sp. v. 255 = viscosa.
spinifex, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 186 = bivalvis.
viscosa, *H. B. & K. Nov. Gen. et Sp.* v. 256.—Mexic.

BASTERA, J. F. Gmel. Syst. 1271 (1791) = BASTERIA, Houtt. = **Berkheya**, Ehrh. (Compos.).

BASTERIA, Mill. Gard. Dict. ed. VII (1759) = **Calycanthus**, Linn.
Calycanthus, Mill. l. c. ed. VIII = C. floridus.

BASTERIA, Houtt. Nat. Hist. Pl. vi. 158 (1776) = **Berkheya**, Ehrh. (Compos.).
aculeata, Houtt. l. c. = Berkheya obovata.

BASTIA, Steud. Nom. ed. II. i. 238, in syn. (1840), sphalm. = BUSTIA, Adans. = **Buphthalmum**, Linn. (Compos.).

BATANTHES, Rafin. Atl. Journ. 145 (1832) = **Gilia**, Ruiz & Pav. (Polemon.).
aggregata, Rafin. l. c. = G. aggregata.
longiflora, Rafin. l. c. = Collomia longiflora.
pungens, Rafin. l. c. = G. pungens.

BATATAS, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 434 ; (Conv. Or. 52) = **Ipomoea**, Linn. (Convolv.).
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 64 = I. abyssinica.
acetosaefolia, Choisy, Conv. Rar. 124 = I. carnosa.
auriculata, Hochst. ex Choisy, in DC. Prod. ix. 350 = I. cordofana.
betacea, Lindl. in Bot. Reg. (1839) Misc. 93 ; (1840) t. 56.—Guiana.
bignonioides, G. Don, Gen. Syst. iv. 261 = I. bignonioides.
bombycina, Choisy, in DC. Prod. ix. 340 = I. bombycina.
bonariensis, Lindl. in Bot. Reg. (1838) Misc. 55 = I. ficifolia.
Cavanillesii, G. Don, Gen. Syst. iv. 262 = I. Cavanillesii.
Choisyana, Wight, Ic. t. 491 = Argyreia Choisyana.
cissoides, Choisy, Conv. Or. 55 = I. cissoides.
crassicaulis, Benth. Bot. Voy. Sulph. 134.—Guayaquil.
edulis, Choisy, Conv. Or. 53 = I. Batatas.
glaberrima, Hassk. Retzia, i. 65.—Hab.?
glabra, Benth. in Hook. Lond. Journ. Bot. v. (1846) 352 = I. glabra.
glaucifolia, G. Don, Gen. Syst. iv. 262 = I. glaucifolia.
heterophylla, G. Don, l. c. 261 = I. heterophylla.
incurva, Benth. in Hook. Niger Fl. 464 = I. incurva?
insignis, G. Don, in Sweet, Hort. Brit. ed. III. 482 = I. insignis.
Jalapa, Choisy, Conv. Rar. 125 = I. Jalapa.
littoralis, Choisy, l. c. 124 = I. carnosa.
Loureirii, G. Don, Gen. Syst. iv. 262 = I. tuberosa.
macrorhizos, G. Don, l. c. 261 = I. macrorhizos.
maritima, Boj. Hort. Maurit. 225 = I. biloba.
multiflora, Boj. ex Choisy, in DC. Prod. ix. 371 = I. staphylina.
paniculata, Choisy, Conv. Or. 54 = I. digitata.
Papirin, G. Don, Gen. Syst. iv. 261 = I. pubescens.
pareiraefolia, Choisy, Conv. Rar. 123 = I. pareiraefolia.
pentaphylla, Choisy, Conv. Or. 54 = I. pentaphylla.
pulchella, Boj. Hort. Maurit. 226 = I. pulchella.
quinquefolia, Choisy, Conv. Rar. 127 = I. quinquefolia.
senegalensis, G. Don, Gen. Syst. iv. 261 = I. palmata.
sinuatus, Guss. Enum. Pl. Ins. Inar. (1854) 210 = I. stolonifera.
subtriloba, G. Don, Gen. Syst. iv. 261 = I. pubescens.
ternata, G. Don, l. c. 262 = I. ternata.
tomentosa, Choisy, in DC. Prod. ix. 337 = I. tomentosa.
triloba, Choisy, Conv. Rar. 127 = I. triloba.
tuberosa, Boj. Hort. Maurit. 226 = I. tuberosa.
venosa, Boj. l. c. = I. cairica.
venosa, G. Don, Gen. Syst. iv. 261 = I. venosa.
villosa, Choisy, in DC. Prod. ix. 337 = I. villosa.
viscida, Voigt, Hort. Suburb. Calc. 354.—Ind. or.?
Wallii, Morr. in Ann. Gand. ii. (1846) 285 = I. Wallii.
Willdenowii, G. Don, Gen. Syst. iv. 261 = I. heterophylla.
xanthorhiza, Boj. Hort. Maurit. 225 = I. Batatas.

BATEMANIA, Endl. Gen. 197 (1837) = seq.

BATEMANNIA, Lindl. in Bot. Reg. t. 1714 (1834). *ORCHIDEAE*, Benth. & Hook. f. iii. 546.
apiculata, *Reichb. f. in Linnaea*, xli. (1877) 109.—N. Granat.
armillata, *Reichb. f. in Gard. Chron.* (1878) i. 780.—Hab.?
Beaumontii, *Reichb. f. in Walp. Ann.* iii. 544.—Bras.
Burtii, Endr. & Reichb. f. in Gard. Chron. (1872) 1099 = Zygopetalum Burtii.
Colleyi, *Lindl. Bot. Reg.* xx. (1835) t. 1714.—Guiana.
fimbriata, *Linden & Reichb. f. in Bonplandia*, ii. (1854) 280.—N. Granat.
grandiflora, Reichb. f. l. c. iv. (1856) 323 = Zygopetalum grandiflorum.
Gustavi, *Reichb. f. in Linnaea*, xli. (1877) 108.—N. Granat.
Lalindei, *Linden, Cat.* (1873) n. 90.—N. Granat.
lepida, *Reichb. f. in Gard. Chron.* (1878) i. 588.—Bras.
Meleagris, *Reichb. f. in Bonplandia*, iii. (1855) 217.—Bras.
Wallisii, *Reichb. f. in Gard. Chron.* (1880) i. 776.—N. Granat.

BATERIUM, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xiii. (1864) 124 = **Haematocarpus**, Miers (1867) (Menisp.).
validum, Miers, l. c. = H. Thomsoni.

BATESIA, Spruce, in Benth. & Hook. f. Gen. i. 563 (1865). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
floribunda, *Spruce, ex Benth. in Trans. Linn. Soc.* xxv. (1866) 303. t. 37.—Bras. bor.

BATHYSA, Presl, in Abh. Boehm. Ges. Folge V. iii. (1845) 514 ; (Bot. Bemerk. 84). *RUBIACEAE* Benth. & Hook. f. ii. 49.
SCHOENLEINIA, Klotzsch, in Hayne, Arzneigew. xiv. in nota ad t. 15 (1846).
VOIGTIA, Klotzsch, l. c. (1846).
stipulata, *Presl, l. c.*—Bras.

BATINDUM, Rafin. Sylva Tellur. 81 (1838) = **Oftia**, Adans. (Myopor.).
jasmineum, Rafin. l. c. 82 = O. africana.

BATIS, Linn. Syst. ed. X. 1380. n. 1172 (1759). *BATIDEAE*, Benth. & Hook. f. iii. 88.
americana, *Linn. Mant.* ii. 499.—Am. bor.
aurantiaca, Wall. ex Thw. Enum. Pl. Zeyl. 263 = Plecospermum spinosum.
californica, Torr. in Smithson. Contrib. vi. (1854) 8 = maritima.
fruticosa, Roxb. Hort. Beng. [71] ; Fl. Ind. iii. 763 = Cudrania javanensis et C. fruticosa.
hermaphrodita, *Blanco, Fl. Filip.* ed. I. 791.—Ins. Philipp.
maritima, *Linn. Syst.* ed. X. 1380.—Jamaica.
spinosa, Roxb. Hort. Beng. [71] ; Fl. Ind. iii. 762 = Plecospermum spinosum.
vermiculata, Hook. Fl. Bor. Am. ii. 128 = Sarcobatus vermiculatus.

BATOCARPUS, Karst. Fl. Columb. ii. 67. t. 134 (1862?). *URTICACEAE*, Benth. & Hook. f. iii. 376.
orinocensis, *Karst. l. c.*—Venezuela.

BATOCYDIA, Mart. ex DC. Prod. ix. 143 (1845) = **Bignonia**, Linn.
capreolata, Mart. l. c. 146 = Bignonia capreolata.
cuneifolia, Mart. l. c. 157 = Bignonia cuneifolia.
exoleta, Mart. l. c. 148 = Bignonia exoleta.
Unguis, Mart. l. c. 146 = Bignonia Unguis-cati.

BATODENDRON, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1843) 261 = **Vaccinium**, Linn.
arboreum, Nutt. l. c. = V. arboreum.

BATRACHIUM, S. F. Gray, Nat. Arr. Brit. Pl. ii. 720 (1821) = **Ranunculus**, Tourn.
admixtum, Nyl. in Nyl. & Sael. Herb. Mus. Fenn. 35 = R. confervioides.
aquatile, Wimm. Fl. Schles. (1841) 8 = R. aquatilis.
aspergillifolium, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 218 = R. trichophyllus.

BATRACHIUM :—
Bachii, Wirtg. ex F. Schultz, Arch. Fl. Fr. et Allem. i. (1844) 292 = Ranunculus fluitans.
Baudotii, F. Schultz, l. c. 71 = R. Baudotii.
bipontinum, F. Schultz, ex Gren. & Godr. Fl. Fr. i. 24 = R. trichophyllus.
biternatum, Bereht. & Presl, Rostl. i. Ranunc. 49 = R. biternatus.
caespitosum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 722 = R. circinatus.
caespitosum, F. Schultz, Arch. Fl. Fr. et Allem. i. (1844) 71 = R. trichophyllus.
capillaceum, Bercht. & Presl, Rostl. i. Ranunc. 49 = R. trichophyllus.
carinatum, Schur, in Verh. Naturf. Ver. Brucnn, xv. II. (1877) 28 = R. Baudotii.
circinatoides, Arv. in Soc. Ech. Dauph. iv. (1877) 105 = R. aquatilis ?
circinatum, Spach, Hist. Vég. Phan. vii. 201 = R. circinatus.
coenosum, Nym. in Bot. Notiser (1852), 100 = R. cocnosus.
confervoides, Fries, in Bot. Notiser (1845), 121 = R. coenosus.
confusum, Garcke, Fl. Deutschl. ed. V. 8 = R. Petiveri.
confusum, Le Jolis, in Mém Soc. Sc. Nat. Cherb. vii. (1860) 257 = R. confusus.
dichotomum, Schmalh. ex Trautv. in Act. Hort. Petrop. ix. (1884) 363 = R. aquatilis.
divaricatum, Wimm. Fl. Schles. (1841) 10 = R. divaricatus.
diversifolium, Nym. Consp. 15 = R. diversifolius.
Drouetii, Nym. in Bot. Notiser (1852) 98 = R. Drouetii.
elongatum, F. Schultz, in Billotia, i. 113 = R. peltatus.
eradicatum, Lacst. ex Nym. Consp. 15 = R. confervoides.
flaccidum, Rupr. Fl. Cauc. 15 = R. aquatilis.
floribundum, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 216 = R. floribundus.
fluitans, Wimm. Fl. Schles. (1841) 9 = R. fluitans.
fluviatile, S. F. Gray, Nat. Arr. ii. 722 = R. fluviatilis.
Friesii, Nym. Consp. 15 = R. heterophyllus.
Godronii, Nym. l. c. 16 = R. diversifolius.
hederaceum, S. F. Gray, Nat. Arr. ii. 721 = R. hederaceus.
heterophyllum, S. F. Gray, l. c. = R. heterophyllus.
heterophyllum, Lange, Pugill. 251 = R. peltatus.
hololeucum, Garcke, Fl. Deutschl. ed. IV. 7 = R. hololeucus.
intermedium, Nym. Syll. 175 = R. intermedius.
Langei, F. Schultz, ex Nym. Consp. 16 = R. tripartitus.
leiospermum, Hartm. f. Skand. Fl. ed. XIII. 165 = R. hederaceus.
Lenormandii, F. Schultz, Arch. Fl. Fr. et Allem. i. (1844) 70 = R. Lenormandi.
longirostre, F. Schultz, l. c. 71 = R. longirostris.
lutarium, Revel, in Act. Linn. Soc. Bord. xxv. (1864) 413. t. 4 = R. Lenormandi.
luteolum, Revel.—Cf. Bull. Soc. Bot. Fr. xii. (1865) Bibliog. 258 = R. heterophyllus.
lutulentum, Nym. Syll. Suppl. 29 = R. trichophyllus.
marinum, Fries, Nov. Mant. iii. 51 = R. Baudotii.
minimum, Schur, in Verh. Naturf. Ver. Brucnn, xv. II. (1877) 28 = R. trichophyllus.
obtusiflorum, S. F. Gray, Nat. Arr. ii. 721 = R. tripartitus.
ololeucos, [Bossch] Prod. Fl. Bat. 7 = R. hololeucus.
omoiophyllum, Nym. Consp. 16 = R. hederaceus.
pantothrix, S. F. Gray, Nat. Arr. ii. 722 = R. trichophyllus.
paucistamineum, F. Schultz, Arch. Fl. Fr. et Allem. (1844) 71 = R. trichophyllus.
pectinatum, Nym. Consp. 15 = R. trichophyllus.
peltatum, Bercht. & Presl, Rostl. i. Ranunc. 49 = R. peltatum.
penicillatum, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 216 = R. peltatus.
Petiveri, F. Schultz, Arch. Fl. Fr. et Allem. i. (1844) 71 = R. Petiveri.
peucedanifolium, Dum. Fl. Belg. 127 = R. fluitans.
pseudofluitans, Nym. Consp. 16 = R. peltatus.
pumilum, Nym. in Bot. Notiser (1852), 98 = R. fluitans.
radians, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 215 = R. trichophyllus.

BATRACHIUM :—
radians, Revel, in Act. Linn. Soc. Bord. xix. (1853) 120 = Ranunculus radians.
rhipiphyllum, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 215 = R. diversifolius.
rigidum, Dum. Fl. Belg. 127 = R. trichophyllus.
Rionii, Nym. in Bot. Notiser (1850) = R. sceleratus.
tenellum, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 211 = R. tenellus.
trichophyllum, [Bossch] Prod. Fl. Bat. 7 ; Des Moul. in Act. Soc. Linn. Bord. xx. (1859) 456 = R. trichophyllus.
trinacrium, Nym. Consp. 16 = R. tripartitus.
tripartitum, S. F. Gray, Nat. Arr. ii. 721 = R. tripartitus.
triphyllos, Dum. in Bull. Soc. Bot. Belg. ii. (1863) 214 = R. tripartitus.
truncatum, Dum. l. c. 215 = R. peltatus.
villosum, F. Schultz, ex Gren. & Godr. Fl. Fr. i. 24 = R. trichophyllus.

BATRATHERUM, Nees, in Edinb. N. Phil. Journ. xviii. (1835) 180 = **Arthraxon**, Beauv. (Gram.).
cuspidatum, Hochst. in Flora, xxxix. (1856) 178 = A. cuspidatus.
echinatum, Nees, in Edinb. N. Phil. Journ. xviii. (1835) 181 = A. laneeolatus.
lanceolatum, Nees, l. c. = A. lanceolatus.
micans, Nees, l. c. 182 = A. cuspidatus.
molle, Nees, l. c. 181 = A. microphyllus.
nudum, Nees, ex Steud. Syn. Pl. Gram. 383 = A. ciliaris.
plumbeum, [Munro, ex] Duthie, in Atkinson, Gazetteer, 640 = A. ciliaris.
rude, Nees, ex Steud. Syn. Pl. Gram. 383 = A. rudis.
Schimperi, Nees, ex Hochst. in Flora, xxxix. (1856) 179 = A. microphyllus.
serrulatum, Hoehst. ex A. Rich. Tent. Fl. Abyss. ii. 458 = A. lanceolatus.
submuticum, Nees, ex Steud. Syn. Pl. Gram. 382 = A. ciliaris.

BATSCHIA, J. F. Gmel. Syst. ii. 315 (1791) = **Lithospermum**, Linn. (Boragin.).
albiflora, Rafin. New Fl. Am. iv. 19.—Am. bor.
canescens, Michx. Fl. Am. Bor. i. 130. t. 14 = L. canescens.
caroliniana, Roem. & Schult. Syst. iv. 52 = L. hirtum.
caroliniensis, J. F. Gmel. Syst. i. 315 = L. hirtum.
conspicua, R. Br. in Richards. App. Frankl. Journ. 732 = L. canescens.
decumbens, Nutt. Gen. Am. i. 114 = L. angustifolium.
disticha, G. Don, Gen. Syst. iv. 326 = L. distichum.
Gmelini, Michx. Fl. Am. Bor. i. 130 = L. hirtum.
longiflora, Pursh, Fl. Am. Sept. i. 132 = L. angustifolium.
pilosa, G. Don, Gen. Syst. iv. 372 = L. pilosum.
sericea, Roem. & Schult. Syst. iv. 743 = L. canescens.
Torreyi, G. Don, Gen. Syst. iv. 372 = L. pilosum.

BATSCHIA, Moench, Meth. 567 (1794) = **Eupatorium**, Linn. (Compos.).
nivea, Moench, l. c. = E. ageratoides.

BATSCHIA, Mutis, ex Thunb. in Nov. Act. Ups. v. (1792) 120. t. 2 = **Abuta**, Aubl. (Menisp.).
conferta, Miers, in Seem. Bot. Voy. Herald, i. 76 = A. Seemanni.
conferta, DC. Prod. i. 103 = A. spicata.
racemosa, Thunb. in Nov. Act. Ups. v. (1792) 122 = A. racemosa.
spicata, Thunb. l. c. = A. spicata.

BATSCHIA, Vahl, Symb. Bot. iii. 39. t. 56 (1794) = **Humboldtia**, Vahl (Legum.).
laurifolia, Vahl, l. c. = H. laurifolia.

BATTATA, Hill, Hort. Kew. 148 (1768) = **Solanum**, Linn.
tuberosa, Hill, l. c. = S. tuberosum.

BAUCHEA, Fourn. ex Benth. in Journ. Linn. Soc. xix. (1881) 88. *GRAMINEAE*, Benth. & Hook. f. iii. 1149.
Karwinskii, *Fourn. Mex. Pl. En. Gram.* 87 ; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 550.—Mexic.

BAUDINIA, Lesch. ex DC. Prod. iii. 211 (1828) = **Calothamnus**, Labill. (Myrtac.).

BAUDINIA, Lesch. ex DC. Prod. vii. 511 (1839) = **Scaevola**, Linn. (Goodenov.).
humilis, Lesch. ex DC. l. c. 511 = S. striata.

BAUDOUINIA, Baill. Adansonia, vi. (1866) 193. t. 5. *LEGUMINOSAE*, Benth. & Hook. f. i. 1003.
fluggeiformis, *Baill. l. c.* viii. (1868) 201.—Madag.
sollyaeformis, *Baill. l. c.* vi. (1866) 194.—Madag.

BAUERA, Banks, ex Andr. Bot. Rep. t. 198 (1793). *SAXIFRAGEAE*, Benth. & Hook. f. i. 655.
Billardieri, D. Don, in Edinb. N. Phil. Journ. ix. (1830) 96 = rubioides.
capitata, *Ser. in DC. Prod.* iv. 13.—Austral.
galioides, Sieber, in Reichb. Ic. Exot. t. 77 = rubioides.
humilis, Sweet, Hort. Suburb. 124 = rubioides.
juncea, Walp. Rep. ii. 375 = Tetratheca juncea.
microphylla, Sieber, ex DC. Prod. iv. 13 = rubioides.
rubiaefolia, Salisb. in Kon. & Sims, Ann. Bot. i. (1805) 514. t. 10 = rubioides.
rubioides, *Andr. Bot. Rep.* t. 198.—Austral.
sessiliflora, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 41.—Austral.

BAUHINIA, Linn. Gen. ed. I. 126 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 575.
ALVESIA, Welw. Apont. 587. n. 47 (1858).
AMARIA, Mutis, ex Caldas, in Seman. Nucv. Gran. II. (1810) 25.
BINARIA, Rafin. Sylva Tellur. 122 (1838).
CANSENIA, Rafin. l. c. (1838).
CASPAREA, H. B. & K. Nov. Gen. et Sp. vi. 317 (1823).
CAULOTRETUS, Rich. ex Spreng. Syst. iv. Cur. Post. 406 (1827).
ELAYUNA, Rafin. Sylva Tellur. 145 (1838).
LACARA, Spreng. Neue Entdeck. iii. 56 (1822).
LASIOBEMA, Miq. Fl. Ind. Bat. i. 71 (1855).
LOCELLARIA, Welw. Apont. 588 (1858).
MANDARUS, Rafin. Sylva Tellur. 122 (1838).
MONOTELES, Rafin. l. c. (1838).
PAULETIA, Cav. Ic. v. 5. t. 409, 410 (1799).
PERLEBIA, Mart. Reise, i. 555 (1828).
PHANERA, Lour. Fl. Cochinch. 37 (1790).
PILEOSTIGMA, Hochst. in Flora, xxix. (1846) 598.
SCHNELLA, Raddi, in Mem. Mod. xviii. Fis. (1820) 411.
TELESTRIA, Rafin. Sylva Tellur. 122 (1838).
abyssinica, A. Rich. Tent. Fl. Abyss. i. 253 = retieulata.
acida, Reinw. in Flora, xxxi. (1848) 578 = malabarica.
aculeata, *Linn. Sp. Pl.* 374.—Am. austr.
aculeata, Vell. Fl. Flum. 171 ; iv. t. 83 = forficata.
acuminata, Bruce, Trav. Abyss. App. 167. t. 18 = B. Farec.
acuminata, *Linn. Sp. Pl.* 376.—Ind. or.; Malaya; China.
acuminata, Vell. Fl. Flum. 171 ; iv. t. 85.—Bras.
Açuruana, *Moric. Pl. Nouv. Rar. Am.* 77. t. 51.—Bras.
Adansoniana, *Guill. & Perr. Tent. Fl. Seneg.* i. 265.—Afr. trop.
aegopoda, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69, 120 = B. Pes-caprae.
affinis, *Vog. in Linnaea*, x. (1836) 594.—Bras.
Aimouta, Steud. Nom. ed. II. i. 191 (err. typ. 291), vide *Outimouti*.
alba, Buch.-Ham. ex Wall. Cat. sub n. 5796 = variegata.
albicans, *Vog. in Linnaea*, xiii. (1839) 304.—Bras.
americana, *Delaun. Herb. Amat.* t. 315.—Am. austr.
ampla, *Span. Ic.* 29 ; *in Linnaea*, xv. (1841) 202.—Timor.
anatomica, *Link, Enum. Hort. Berol.* i. 405.—Am. austr.
Andrieuxii, *Hemsl. Diag. Pl. Nov.* 48.—Mexic.
anguina, *Roxb. Hort. Beng.* 31 ; *Fl. Ind.* ii. 328.—Ind. or.; Malaya.
angulosa, *Vog. in Linnaea*, xiii. (1839) 312.—Bras.
armata, *Otto, Pl. Bras.* 25.—Bras.
articulata, Oliver, Fl. Trop. Afr. ii. 290, err. typ. = reticulata ?
aurantiaca, *Boj. in Ann. Sc. Nat.* Ser. II. iv. (1835) 264.—Madag.
aurita, *Ait. Hort. Kew.* ii. 48.—Jamaica.

BAUHINIA:—
Bacuria, Buch.-Ham. ex Wall. Cat. sub n. 5793 = malabarica.
bahiensis, Bong. in Mém. Acad. Pétersb. Sér. VI. iv. (1836) 114 = Langsdorffiana.
Baueriana, Steud. Nom. ed. II. i. 191 (err. typ. 291) = malabarica.
Benthami, Bedd. Forest. Man. Bot. 93; Ic. Pl. Ind. Or. i. 22 = phoenicea.
Benzoin, *Kotschy, ex Schweinf. Reliq. Kotsch.* (1868) 12. t. 11.—Afr. trop.
bicolor, *D. Dietr. Syn. Pl.* ii. 1476.—Bras.
bicuspidata, *Benth. in Mart. Fl. Bras.* xv. II. 193.—Am. trop.
bidentata, *Jack, in Malay. Misc.* ii. (1822) n. VII. p. 76.—Malaya.
binata, Blanco, Fl. Filip. ed. I. 331 = Blancoi.
binata, Naves, ex Villar, in Blanco, Fl. Phil. ed. III. Nov. App. 72 = tomentosa.
Blancoi, *Benth. Fl. Austral.* ii. 296, in nota; *Baker, in Hook. f. Fl. Brit. Ind.* ii. 278.—Siam; Ins. Philipp.
Bongardi, *Steud. Nom.* ed. II. i. 191 (err. typ. 291).—Bras.
Bowkeri, *Harv. in Harv. & Sond. Fl. Cap.* ii. 596.—Afr. austr.
brachycarpa, *Wall. Cat.* n. 5786.—Burma.
brachystachya, Walp. Rep. i. 852 = cumanensis.
bracteata, *R. Grah. in Wall. Cat.* n. 5802.—Burma; Siam.
brasiliensis, *Vog. in Linnaea*, xiii. (1839) 302.—Bras.
Bredemeyeri, *Vog. l. c.*—Venezuela.
breviloba, *Benth. in Mart. Fl. Bras.* xv. II. 197.—Bras.
brevipes, Vog. in Linnaea, xiii. (1839) 307 = Bongardi.
Buchanani, Desv. in Ann. Sc. Nat. Sér. I. ix. (1829) 430 = diphylla.
Burchellii, *Benth. in Mart. Fl. Bras.* xv. II. 186.—Bras.
Burkeana, Benth. ex Harv. & Sond. Fl. Cap. ii. 275 = esculenta.
Cancara, Buch.-Ham. ex Steud. Nom. ed. II. i. 191 (err. typ. 291) = variegata.
candicans, *Benth. in Mart. Fl. Bras.* xv. II. 201.—Bras.
candida, Ait. Hort. Kew. ii. (1789) 49 = acuminata.
candida, Roxb. Hort. Beng. [31]; Fl. Ind. ii. 318 = variegata.
Carronii, *F. Muell. in Trans. Phil. Inst. Victoria*, iii. (1858) 49.—Austral.
castrata, *Blanco, Fl. Filip.* ed. I. 331.—Ins. Philipp.
castrata, *Hassk. in Flora*, xxv. (1842) *Beibl.* II. 54.—Java.
Championi, *Benth. Fl. Hongkong.* 99.—Hongkong.
cheilantha, *D. Dietr. Syn. Pl.* ii. 1476.—Bras.
chinensis, Vog. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 42 = variegata.
choriophylla, Vog. in Linnaea, xiii. (1839) 307 = rufa.
chrysophylla, Vahl, ex DC. Prod. ii. 517 = macrophylla.
chrysophylla, Vog. in Linnaea, xiii. (1839) 313 = splendens.
cinnamomea, *DC. Prod.* ii. 517.—Guiana.
cissoides, *Welw. ex Oliver, Fl. Trop. Afr.* ii. 287.—Afr. trop.
coccinea, *DC. Prod.* ii. 516.—Cochinch.
columbiensis, Vog. in Linnaea, xiii. (1839) 313 = cumanensis.
compacta, *Hort. ex Regel, Cat. Pl. Aksakov.* 15.—Hab.?
confertiflora, *Benth. in Mart. Fl. Bras.* xv. II. 206.—Bras.
cordata, Vog. in Linnaea, xi. (1837) 308 = rufa.
cordifolia, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 332 = emarginata.
corniculata, *Benth. in Mart. Fl. Bras.* xv. II. 198.—Bras.
cornifolia, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 278.—Ins. Penang.
coromandeliana, DC. Mém. Leg. xiii. 486; Prod. ii. 515 = purpurea.
coronata, *Benth. in Mart. Fl. Bras.* xv. II. 209.—Bras.
corymbosa, *Roxb. Hort. Beng.* [31]; *Fl. Ind.* ii. 329.—China.
cucullata, *Desv. Journ. Bot.* i. (1814) 74 (= variegata?).—Ind. or.
cumanensis, *H. B. & K. Nov. Gen. et Sp.* vi. 321.—Venezuela.

BAUHINIA:—
Cumingiana, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App* 73.—Ins. Philipp.
Cunninghamii, *Benth. Fl. Austral.* ii. 295.—Austral.
cupulata, *Benth. in Mart. Fl. Bras.* xv. II. 188.—Bras.
curvula, *Benth. l. c.* 194.—Bras.
cuyabensis, *Steud. Nom.* ed. II. i. 191 (err. typ. 291).—Bras.
debilis, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 149.—Java.
dipetala, *Hemsl. Diag. Pl. Nov.* 48.—Mexic.
diphylla, *Buch.-Ham. in Symes, Embassy*, ed. II. iii. 311.—Burma; Malaya.
diptera, *Blume, ex Miq. in Verh. Nederl. Inst.* III. iii. (1850) 12.—Malaya.
divaricata, Hort. ex Steud. Nom. ed. II. i. 191 (err. typ. 291) = forficata.
divaricata, Lam. Encyc. i. 389 = Lamarckiana.
divaricata, *Linn. Sp. Pl.* 374.—Am. austr.
divergens, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 282.—Burma.
dodecandra, *D. Dietr. Syn. Pl.* ii. 1475.—Bras.
dubia, *G. Don, Gen. Syst.* ii. 463.—Bras.
dubia, Vog. in Linnaea, xiii. (1839) 314 = coronata.
dumosa, *Benth. in Mart. Fl. Bras.* xv. II. 194.—Bras.
elongata, *Korth. Verh. Nat. Gesch. Bot.* (1839–42) 89.—Borneo.
emarginata, *Jack, in Malay. Misc.* ii. (1822) n. VII. p. 75.—Sumatra.
emarginata, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
emarginata, Wall. Cat. n. 5792 = retusa.
eriophora, *Tenore, Ind. Sem. Hort. Neap.* (1842) 12.—Hab.?
excelsa, Blume, ex Korth. Verh. Nat. Gesch. Bot. 90 = ferruginea.
esculenta, *Burch. Trav.* ii. 589.—Afr. austr.
excisa, *Hemsl. Biol. Centr. Am. Bot.* i. 337.—Panama.
eximia, *Miq. Fl. Ind. Bat.* i. I. 75.—Java.
Farec, *Desv. Journ. Bot.* i. (1814) 74.—Abyssin.
fassoglensis, *Kotschy, ex Schweinf. Reliq. Kotsch.* (1868) 14. t. 12.—Afr. trop.
ferruginea, D. Dietr. Syn. Pl. ii. 1476 = Bongardi.
ferruginea, *Roxb. Hort. Beng.* 90; *Fl. Ind.* ii. 331.—Malaya.
Finlaysoniana, *R. Grah. in Wall. Cat.* n. 5801.—Malaya.
flexuosa, *Moric. Pl. Nouv. Am.* 80. t. 53.—Bras.
floribunda, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 430.—Bras.
forficata, *Link, Enum. Hort. Berol.* i. 404.—Bras.
foveolata, Dalz. in Journ. Linn. Soc. xiii. (1873) 188 = Lawii?
fulva, *Korth. Verh. Nat. Gesch. Bot.* (1839–42) 91.—Java.
furcata, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 429.—Am. bor.
fusconervis, *D. Dietr. Syn. Pl.* ii. 1476.—Bras.
Gardneri, *Benth. in Mart. Fl. Bras.* xv. II. 186.—Bras.
garipensis, *E. Mey. Comm. Pl. Afr. Austr.* i. 162.—Afr. austr.
geminata, *Vog. in Linnaea*, xiii. (1839) 305.—Bras.
Gilesii, *F. Muell. & Bailey, in Wing, South Sc. Record*, ii. (1882) 151.—Austral.
glabra, *Jacq. Enum. Pl. Carib.* 20.—N. Granat.
glabrifolia, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 281.—Burma.
glandulosa, *DC. Prod.* ii. 513.—N. Barcel.
glauca, *Wall. Cat.* n. 5785; *Benth. Fl. Hongkong.* 99.—Burma; Malaya; China.
glaucescens, *DC. Prod.* ii. 515.—Venezuela.
Grahami, Steud. Nom. ed. II. i. 191 (err. typ. 291) = rufa.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 373.—Madag.
grandiflora, *Juss. in Poir. Encyc. Suppl.* i. 600.—Peruv.
grandiflora, Blanco, Fl. Filip. ed. I. 332 = variegata.
grandifolia, *D. Dietr. Syn. Pl.* ii. 1475.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 377. t. 145.—Guiana.
heterandra, *Benth. in Mart. Fl. Bras.* xv. II. 196.—Bras.
heterophylla, *Kunth, Mim.* t. 46.—Venezuela.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 107.—Afr. trop.

BAUHINIA:—
hirsuta, *Vog. in Linnaea*, xiii. (1839) 300.—Bras.
hirsuta, Weinm. in Ratisb. Syll. ii. (1828) 9 = acuminata.
hirta, *Steud. Nom.* ed. II. i. 191 (err. typ. 291).—Bras.
holophylla, *Steud. l. c.; D. Dietr. Syn. Pl.* ed. II. 1476.—Bras.
Hookeri, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 51.—Austral.
Hostmanniana, *Miq. in Verh. Nat. Wet. Haarl.* vii. (1851) 10.—Guiana.
Humblotiana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 365.—Madag.
hymenaeaefolia, *Triana, ex Hemsl. Biol. Centr. Am. Bot.* i. 337.—Panama.
indica, *Lodd. ex Loud. Hort. Brit.* 165.—Ind. or.
inermis, Forsk. Fl. Aegypt. Arab. 85 = reticulata, tomentosa.
inermis, *Pers. Syn.* i. 455.—Am. austr.
integerrima, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 197.—Bras.
integrifolia, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 331.—Malaya.
intermedia, Vog. in Linnaea, xi. (1837) 309 = rufa.
involucellata, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 72.—Burma.
isopetala, *Griff. Notul.* iv. 451.—Burma.
japonica, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 401.—Japon.
Kappleri, *Sagot, in Ann. Sc. Nat.* Sér. VI. xiii. (1882) 317.—Guiana.
khasiana, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 281.—Ind. or.
Kirkii, *Oliver, Fl. Trop. Afr.* ii. 288.—Afr. trop.
Kockiana, *Korth. Verh. Nat. Gesch. Bot.* (1839–42) 87.—Sumatra.
Krugii, *Urban, in Berl. Deutsch. Bot. Ges.* iii. (1885) 83.—Porto-Rico.
Kunthiana, *Vog. in Linnaea*, xiii. (1839) 312.—Am. austr.
Lamarckiana, *DC. Prod.* ii. 512.—Am. austr.
Langsdorffiana, *Bong. in Mém. Acad. Pétersb.* iv. (1836) 109.—Bras.
latifolia, *Cav. Ic.* v. 4. t. 405.—Mexic.
latisiliqua, *Cav. l. c.* 5.—Ins. Philipp.
Lawii, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 277.—Ind. or.
Leichhardtii, *F. Muell. in Trans. Phil. Inst. Victoria*, iii. (1859) 50.—Austral.
leiopetala, *Benth. in Mart. Fl. Bras.* xv. II. 209.—Bras.
leptopetala, *Moç. & Sessé, ex DC. Prod.* ii. 513.—Mexic.
Lingua, *DC. l. c.* 516.—Malaya.
longicuspis, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 185.—Bras.
longiflora, D. Dietr. Syn. Pl. ii. 1475 = forficata.
longifolia, *D. Dietr. l. c.* 1476.—Bras.
longipetala, *Walp. Rep.* i. 852.—Guiana.
longistyla, Benth. in Mart. Fl. Bras. xv. II. 185 = holophylla.
lucida, Wall. Cat. n. 5779 A = emarginata.
lunaria, *Cav. Ic.* v. 4. t. 407.—Mexic.
lunaroides, *A. Gray, in S. Wats. Bibliog. Ind. N. Am. Bot.* i. 205.—N. Mexic.
macrantha, *Oliver, Fl. Trop. Afr.* ii. 289.—Afr. trop.
macranthera, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 338.—Mexic.
macrophylla, *Poir. Encyc. Suppl.* i. 600.—Am. austr.
macrostachya, *Benth. in Hook. Journ. Bot.* ii. (1840) 96.—Guiana.
macrostachya, *Wall. Cat.* n. 5774.—Ind. or.
maculata, *Tenore, Cat. Hort. Neap.* (1845) 79.—Hab.?
madagascariensis, *Desv. Journ. Bot.* i. (1814) 74.—Madag.
malabarica, *Roxb. Hort. Beng.* [31]; *Fl. Ind.* ii. 321.—Ind. or.; Burma.
marginata, *D. Dietr. Syn. Pl.* ii. 1475.—Bras.
Maximiliani, *Benth. in Mart. Fl. Bras.* xv. II. 203.—Am. trop.
megalandra, *Griseb. Fl. Brit. W. Ind.* 213.—Ind. occ.
membranacea, *Benth. in Mart. Fl. Bras.* xv. II. 187.—Bras.
mexicana, *Vog. in Linnaea*, xiii. (1839) 299.—Mexic.
microphylla, *Vog. l. c.* 301.—Am. austr.

BAUHINIA :—
mollis, *D. Dietr. Syn. Pl.* ii. 1475.—Bras.
mollissima, Wall. Cat. n. 5782 = elongata.
Mombassae, Vatke, in Oestr. Bot. Zeitschr. xxx. (1880) 279.—Afr. trop.
monandra, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 73.—Burma.
mucora, C. Bolle, in Peters, Reise Mossamb. Bot. i. 22 = tomentosa.
multinervia, *DC. Prod.* ii. 515.—Venezuela.
natalensis, *Oliver, in Hook. Bot. Mag.* t. 6086.—Afr. austr.
nervosa, *Wall. Cat.* n. 5777.—Ind. or.
nitida, *Benth. in Mart. Fl. Bras.* xv. II. 184.—Bras.
notophila, *Griseb. in Goett. Abh.* xxiv. (1879) 116.—Reg. Argent.
obtusata, *Vog. in Linnaea*, xiii. (1839) 305.—Bras.
odoratissima, *Moric. Pl. Nouv. Am.* 78. t. 52.—Bras.
odoratissima, *Newm. ex G. Don, Gen. Syst.* ii. 461.—Bras.
Olfersiana, *Vog. in Linnaea*, xiii. (1839) 306.—Bras.
ornata, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 72.—Burma.
Outimouta, Aubl. Pl. Gui. i. 375. t. 144 = guianensis, macrophylla.
ovata, *Vog. in Linnaea*, xiii. (1839) 300.—Bras.
panamensis, Spreng. Syst. ii. 384 = B. Pauletia.
parviflora, Vahl, Symb. Bot. iii. 55 = racemosa.
parvifolia, Hochst. ex Field. & Gardn. Sert. t. 10 = rufescens.
parvifolia, Scem. Bot. Voy. Herald, 113 = B. Pauletia.
parvifolia, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 257.—China.
Pauletia, *Pers. Syn.* i. 455.—Panama.
pentandra, *Vog. ex D. Dietr. Syn. Pl.* ii. 1475.—Bras.
Persichii, *F. Muell. in Wing, South Sc. Record* (Feb. 1885), 25.—Austral.
Pervilleana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 374.—Madag.
Pes-caprae, *Cav. Ic.* v. 3. t. 404.—Mexic.
Petersiana, *C. Bolle, in Peters, Reise Mossamb. Bot.* i. 24.—Afr. trop.
petiolata, *Triana, ex Hook. Bot. Mag.* t. 6277.—N. Granat.
phoenicea, *Heyne, in Wall. Cat.* n. 5800.—Ind. or.
picta, *DC. Prod.* ii. 515.—N. Granat.
pinnata, *Blanco, Fl. Filip.* 334.—Ins. Philipp.
piperifolia, Roxb. Fl. Ind. ii. 327 = anguina?
platisiliqua, Steud. Nom. ed. II. i. 192 = reticulata.
platycalyx, *Benth. in Mart. Fl. Bras.* xv. II. 207.—Bras.
platypetala, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 339.—Mexic.
platyphylla, *Benth. in Mart. Fl. Bras.* xv. II. 185.—Bras.
platyphylla, Zipp. ex Span. in Linnaea, xv. (1841) 201 = purpurea.
platysiliqua, Perr. ex DC. Prod. ii. 515 = reticulata.
Poitcauana, *Vog. in Linnaea*, xiii. (1839) 309.—Guiana.
polyantha, *Vog. l. c.* 310.—Bras.
polyacantha, Steud. Nom. ed. II. i. 192 = polyantha.
polycarpa, *Wall. Cat.* n. 5787.—Burma.
porosa, *Boivin, ex Baill. in Bull. Soc. Linn. Par.* i. (1883) 373.—Madag.
porrecta, *Sw. Prod. Veg. Ind. Occ.* 66.—Ind. occ.
Pottsii, G. Don, Gen. Syst. ii. 462 = ferruginea.
pruinosa, *Voy. in Linnaea*, xiii. (1839) 301.—Bras.
pubescens, DC. Lég. Mém. xiii. 483 = aculeata.
pulchella, *Benth. in Mart. Fl. Bras.* xv. II. 190.—Bras.
punctata, *Bolle, in Peters, Reise Mossamb. Bot.* i. 23.—Afr. trop.
punctata, *Burch. ex Benth. in Mart. Fl. Bras.* xv. II. 211.—Bras.
purpurea, DC. ex Walp. Ann. ii. 448 = malabarica.
purpurea, *Linn. Sp. Pl.* 375.—Ind. or.; Burma; China.
purpurea, Wall. Cat. n. 5797 = acuminata, variegata.
pyrrhancura, *Korth. Verh. Nat. Gesch. Bot.* (1839-42) 88.—Sumatra.
pyrrhocarpa, Hochst. in Flora, xxvii. (1844) 99 = reticulata.
racemifera, *Desf. Journ. Bot.* iii. (1814) 74.—Am. austr.
racemosa, *Lam. Encyc.* i. 390.—Ind. or.; Malaya; China.
racemosa, Vahl, Symb. Bot. iii. 56. t. 62 = Vahlii.
Raddiana, *Bong. in Mém. Acad. Pétersb.* Sér. VI. iv. (1836) 111.—Bras.

BAUHINIA :—
radiata, *Vell. Fl. Flum.* 170; iv. t. 81.—Bras.
ramosissima, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 339.—Mexic.
reniformis, Royle, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 277 = malabarica.
reticulata, *DC. Lég. Mém.* xiii. 484.—Senegal.; Aegypt.
retusa, Poir. Encyc. Suppl. i. 699 = Lamarckiana, purpurea.
retusa, *Roxb. Hort. Beng.* [31]; *Fl. Ind.* ii. 322.—Reg. Himal.
rhodacantha, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 429.—Bras.
Richardiana, *DC. Prod.* ii. 517.—Guiana.
Riedeliana, *Bong. in Mém. Acad. Pétersb.* Sér. VI. iv. (1836) 113.—Bras.
Riedelii, *Baker, in Journ. Linn. Soc.* xv. (1877) 98.—Celebes.
riparia, Mart. ex Benth. in Mart. Fl. Bras. xv. II. 190 = pulchella.
rosea, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 72.—Burma.
rosea, *Miq. in Linnaea*, xviii. (1844) 588.—Surinam.
rotundata, *Mill. Gard. Dict.* ed. VIII. n. 7.—N. Granat.
rotundifolia, Pers. Syn. i. 455 = subrotundifolia.
Roxburghiana, Voigt, Hort. Suburb. Calc. 254 = retusa.
rubescens, Pers. Syn. i. 455 = rufescens.
rubiginosa, *Bong. in Mém. Acad. Pétersb.* Sér. VI. iv. (1836) 112.—Bras.
rufa, *R. Grah. in Wall. Cat.* n. 5798.—Ind. or.
rufa, *Steud. Nom.* ed. II. i. 192; *et D. Dietr. Syn. Pl.* ii. 1476.—Bras.
rufescens, *Lam. Encyc.* i. 391.—Afr. austr. et trop.
ruficarpa, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 430 = phoenicea.
Rutenbergiana, *Vatke, in Bremen Abh.* vii. (1882) 249.—Afr. trop.
rutilans, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 206.—Bras.
scandens, Blanco, Fl. Filip. ed. I. 332 = Cumingiana.
scandens, Burm. f. Fl. Ind. 94 = corymbosa.
scandens, Linn. Sp. Pl. 374 = anguina.
scandens, Roxb. Hort. Beng. [31]; Fl. Ind. ii. 326 = macrostachya, racemosa.
Schlechtendaliana, *Mart. & Gal. in Bull. Brux.* x. (1843) II. 308.—Mexic.
Sellowiana, Vog. in Linnaea, xiii. (1839) 306 = cujabensis.
semibifida, *Roxb. Hort. Beng.* [31]; *Fl. Ind.* ii. 330.—Burma; Malaya.
Serpae, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 20.—Afr. trop.
smilacifolia, *Burch. ex Benth. in Mart. Fl. Bras.* xv. II. 183.—Bras.
smilacina, *Steud. Nom.* ed. II. i. 192.—Bras.
spathacea, *Sesse, ex DC. Prod.* ii. 512.—Mexic.
speciosa, Roxb. in Wall. Cat. n. 5791 = tomentosa.
speciosa, Vog. in Linnaea, xiii. (1839) 314 = superba.
spicata, Koen. ex Wall. Cat. n. 5789 = racemosa.
spicata, Rottl. ex Spreng. Fl. Hal. Mant. 39; Syst. ii. 333 = racemosa.
spicata, *Vog. in Linnaea*, xiii. (1839) 310.—Hab.?
spinosa, Poir. Encyc. Suppl. i. 599 = B. Pauletia.
splendens, *H. B. & K. Nov. Gen. et Sp.* vi. 319.—Am. austr.
Sprucei, *Benth. in Mart. Fl. Bras.* xv. II. 205.—Bras.
stipularis, *Korth. ex Verh. Nat. Gesch. Bot.* (1839-42) 92.—Sumatra.
suavcolens, *H. B. & K. Nov. Gen. et Sp.* vi. 321.—Am. austr.
subclavata, *Benth. in Mart. Fl. Bras.* xv. II. 188.—Bras.
subrotundifolia, *Cav. Ic.* v. 4. t. 406.—Mexic.
superba, *Steud. Nom.* ed. II. i. 192.—Guiana.
tamarindacea, Delile, Voy. Méroé, 34 = reticulata.
Teysmanniana, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 19.—N. Guinea.
Thonningii, Schum. Beskr. Guin. Pl. 203 = reticulata.
tenella, *Benth. in Mart. Fl. Bras.* xv. II. 195.—Bras.
tenuis, Span. Ic. 36; et in Linnaea, xv. (1841) 202 = racemosa.
timoriana, Steud. Nom. ed. II. i. 192 = seq.
timorensis, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 466 = racemosa.
tomentosa, *Linn. Sp. Pl.* 375.—As. et Afr. trop.
tomentosa, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 72 = acuminata.

BAUHINIA :—
tomentosa, Vell. Fl. Flum. 171; iv. t. 84 = Langsdorffiana.
tomentosa, Wall. Cat. n. 5790. F & G = malabarica.
triandra, Roxb. Hort. Beng. [31]; Fl. Ind. ii. 320 = purpurea.
unguicularis, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* i. 340.—Mexic.
unguiculata, *Baker, in Hook. f. Fl. Brit. Ind.* ii. 277.—Siam.
Ungula, Jacq. Fragm. 22. t. 15 = aculeata.
Ungula, Willd. ex Steud. Nom. ed. II. i. 192 = Bredemeyeri.
ungulata, *Linn. Sp. Pl.* 374.—'Amer.'
uruguayensis, *Benth. in Mart. Fl. Bras.* xv. II. 209.—Bras.
Vahlii, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 72 = Cumingiana.
Vahlii, *Wight & Arn. Prod.* i. 297.—Ind. or.
variegata, *Linn. Sp. Pl.* 375.—Ind. or.; Burma; China.
velutina, *Wall. Cat.* n. 5781.—Burma.
verrucosa, *Vog. in Linnaea*, xiii. (1839) 311.—Bras.
versicolor, *Bertol. Hort. Bonon. Pl. Nov.* i. 7.—Mexic.
viridescens, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 429.—Ind. or.
Vogeliana, Steud. Nom. ed. II. i. 192 = spicata.
Welwitschii, *Oliver, Fl. Trop. Afr.* ii. 287.—Afr. trop.
Williamsii, *F. Muell. Descr. Pl. Papuan.* iv. 61.—N. Guinea.
xanthoneura, Korth. ex Miq. Fl. Ind. Bat. i. 63 = pyrrhaneura.

BAUKEA, Vatke, in Linnaea, xliii. (1880-82) 104. *LEGUMINOSAE.*
insignis, *Vatke, l. c.*—Madag.
maxima, Baill. in Bull. Soc. Linn. Par. i. (1883) 383 = insignis.

BAUMANNIA, DC. in Mém. Soc. Phys. Genèv. iv. (1833) 583 (Not. Pl. Rar. Jard. Genèv. vi. 1. t. 1, 25, = **Damnacanthus**, Gaertn. (Rubiac.).
geminiflora, DC. ll. cc. = D. indicus.

BAUMANNIA, Spach, Hist. Vég. Phan. iv. 351 (1835, = **Oenothera**, Linn. (Onagrar.).
Douglasiana, Spach, l. c. 352 = O. albicaulis.
Nuttalliana, Spach, l. c. = O. albicaulis.
pinnatifida, Spach, l. c. 353 = O. pinnatifida.

BAUMEA, Gaudich. in Freyc. Voy. Bot. 416. t. 29 (1826) = **Cladium**, P. Br. (Cyperac.).
acuminata, Regel, Cat. Pl. Hort. Aksakov. 15.—Hab.?
arthrophylla, Boeck. in Linnaea, xxxviii. (1874) 242 = C. arthrophyllum.
Brownii, Boeck. l. c. = C. glomeratum.
crassa, Thw. Enum. Pl. Zeyl. 353 = C. riparium.
Deplanchei, Boeck. in Flora, lxi. (1878) 143.—N. Caled.
falcata, Nees, in Hook. Kew Journ. vi. (1854) 29.—Ins. Philipp.
flexuosa, Boeck. in Bremen Abh. vii. (1880) 39.—Madag.
glomerata, Gaudich. in Freyc. Voy. Bot. 416. t. 29.—Ins. Molucc.
iridifolia, Boeck. in Linnaea, xxxviii. (1874) 244.—Madag.
laxa, Boeck. l. c. 245 = C. laxum.
loculata, Boeck. l. c. 243 = C. articulatum.
longifolia, Boeck. l. c. 244 = C. Filum.
mariscoides, Gaudich. in Freyc. Voy. Bot. 417 = C. mariscoides.
Meyenii, Kunth, Enum. Pl. ii. 314.—Ins. Sandvic.
Preissii, Nees, in Lehm. Pl. Preiss. ii. 75 = C. latissimum.
riparia, Boeck. in Linnaea, xxxviii. (1874) 246 = C. riparium.
rubiginosa, Boeck. l. c. 241 = C. glomeratum.
schoenoides, Boeck. l. c. 246 = C. schoenoides.

BAUMGARTENIA, Spreng. Anleit. ii. i. 228 (1817); Syst. ii. 91 (1825) = **Borya**, Labill. (Liliac.).
lucens, Steud. Nom. ed. II. i. 192 = Borya nitida.
nitida, Spreng. Syst. ii. 91 = Borya nitida.
sphaerocephala, Spreng. l. c. = Borya nitida.

BAUMGARTIA, Moench, Meth. 650 (1794) = **Cocculus**, DC. (Menisp.).
scandens, Moench, l. c. = C. carolinus.

BAURSEA, Hort. ex Hoffmgg. Verz. Pfl. 42 (1824) = **Philodendron**, Schott (Aroid.).
glauca, Hoffmgg. l. c.—Hab.?
nervosa, Hort. Angl. ex Hoffmgg. l. c. = P. Karstenianum?

BAURSIA, Schott, Syn. Aroid. i. 73 (1856) = **Philodendron**, Schott (Aroid.).

BAUSCHIA, Seub. ex Warm. in Kjoeb. Vidensk. Meddel. (1872) 123 = **Aneilema**, R. Br. (Commel.).
bracteolata, Seub. ex Warm. l. c. 124 = A. ovato-oblongum.

BAUXIA, Neck. Elem. iii. 160 (1790) = **Marica**, Ker-Gawl. (Irid.).

BAVERA, Poir. in Dict. Sc. Nat. iv. Suppl. 52 (1816) = **Bauera**, Banks (Saxif.).

BAXTERA, Reichb. Consp. 131 (1828) = **Marsdenia**, R. Br. (Asclep.).
loniceroides, Steud. Nom. ed. II. i. 192 = M. loniceroides.

BAXTERIA, R. Br. ex Hook. Lond. Journ. Bot. ii. (1843) 494. t. 13–15. *JUNCACEAE*, Benth. & Hook. f. iii. 866.
australis, *R. Br. l. c.*—Austral.

BAZIASA, Steud. Nom. ed. II. i. 192 (1840) = **Sabazia**, Cass. (Compos.).
humilis, Steud. l. c. = S. humilis.
microglossa, Steud. l. c. = S. microglossa.
sarmentosa, Steud. l. c. = S. sarmentosa.
urticaefolia, Steud. l. c. = S. urticaefolia.

BDALLOPHYTON, Eichl. in Bot. Zeit. xxx. (1872) 709. t. 8 = **Cytinus**, Linn.
Andrieuxii, Eichl. l. c. 715. t. 8 = C. Andrieuxii.
ceratantherum, Eichl. l. c. = C. americanus.

BEATONIA, Herb. in Bot. Mag. sub t. 3779 (1839) = **Tigridia**, Ker-Gawl. (Irid.).
atrata, Herb. in Lindl. Bot. Reg. (1843) Misc. 72 = T. atrata.
bracteolata, Klatt, in Linnaea, xxxiv. (1865–66) 733.—Bolivia.
coelestina, Klatt, l. c. xxxi. (1861–62) 567.—Mexic.
curvata, Herb. in Lindl. Bot. Reg. (1843) Misc. 74 = T. curvata.
grandiflora, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 369 = T. lutea.
Houttei, Klatt, l. c. = T. Houttei.
lutea, Klatt, in Linnaea, xxxi. (1861–62) 566 = T. lutea.
purpurea, Herb. in Bot. Mag. sub t. 3799 = T. violacea.

BEATSONIA, Roxb. in Beats. St. Hel. App. 300 (1816) = **Frankenia**, Linn.
portulacifolia, Roxb. l. c. = F. portulacaefolia.

BEAUCARNEA, Lem. Illustr. Hortic. viii. (1861) Misc. 59 = **Nolina**, Michx. (Liliac.).
Bigelovii, Baker, in Journ. Bot. x. (1872) 326 = N. Bigelovii.
erumpens, Baker, l. c. = N. erumpens.
gracilis, Lem. Illustr. Hortic. viii. (1861) Misc. 61 = N. Hartwegiana.
Hartwegiana, Baker, in Journ. Bot. x. (1872) 327 = N. Hartwegiana.
Hookeri, Baker, l. c. = Dasylirion Hookeri.
humilis, Baker, in Journ. Linn. Soc. xviii. (1881) 237 = N. humilis.
Lindheimeriana, Baker, in Journ. Bot. x. (1872) 328 = N. Lindheimeriana.
longifolia, Baker, l. c. 324 = N. longifolia.
microcarpa, Baker, in Journ. Linn. Soc. xviii. (1881) 236 = N. microcarpa.
Palmeri, Baker, l. c. 235 = N. Palmeri.
parviflora, Baker, in Journ. Bot. x. (1872) 328 = N. parviflora.
recurvata, Lem. Illustr. Hortic. (1861) Misc. 59 = N. recurvata.
stricta, Lem. l. c. = N. recurvata.

BEAUCARNEA:—
texana, Baker, in Journ. Linn. Soc. xviii. (1881) 236 = Nolina texana.
Watsoni, Baker, l. c. = N. Watsoni.

BEAUFORTIA, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 418 (1812). *MYRTACEAE*, Benth. & Hook. f. i. 705.
SCHIZOPLEURA, Endl. Gen. 1228 (1840).
anisandra, *Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 13. t. 1.—Austral.
carinata, *A. Cunn. ex Loud. Hort. Brit.* 319.—Austral.
cyrtodonta, *Benth. Fl. Austral.* iii. 167.—Austral.
Dampieri, *A. Cunn. in Hook. Bot. Mag.* t. 3272.—Austral.
decussata, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.
elegans, *Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 16.—Austral.
empetrifolia, Schau. l. c. 17 = micrantha.
heterophylla, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 345.—Austral.
inops, Schau. in Nov. Act. Nat. Cur. xxi. (1844) 17 = Regelia inops.
interstans, *F. Muell. Fragm.* x. 30.—Austral.
macrostemon, *Lindl. Swan Riv. Bot.* 10. n. 31.—Austral.
micrantha, *Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 18.—Austral.
microphylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 24.—Austral.
orbifolia, *F. Muell. Fragm.* iii. 110.—Austral.
pinifolia, Cels, Cat. (1817) 8 = Calothamnus quadrifida.
puberula, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 345.—Austral.
purpurea, *Lindl. Swan Riv. Bot.* 10. n. 30.—Austral.
Schaueri, *Preiss, ex Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 14.—Austral.
sparsa, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.
splendens, Baxter, in Loud. Hort. Brit. Suppl. i. 582 = sparsa.
squarrosa, *Schau. in Nov. Act. Nat. Cur.* xxi. (1844) 11.—Austral.
velutina, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 345.—Austral.

BEAUHARNOISIA, Ruiz & Pav. in Ann. Mus. Par. xi. (1808) 71. t. 9 = **Tovomita**, Aubl. (Guttif.).
fructipendula, Ruiz & Pav. l. c. = T. fructipendula.

BEAUMARIA, Deless. ex Steud. Nom. ed. II. i. 192 (1840) = **Aristotelia**, L'Hérit. (Tiliac.).
Macqui, Deless. ex Steud. l. c. = A. Macqui.

BEAUMONTIA, Wall. Tent. Fl. Nep. 14. t. 7 (1824). *APOCYNACEAE*, Benth. & Hook. f. ii. 721.
grandiflora, *Wall. l. c.* 15. t. 7.—Ind. or.
Jerdoniana, *Wight, Ic.* iv. t. 1314.—Ind. or.
khasiana, *Hook. f. Fl. Brit. Ind.* iii. 661.—Reg. Himal.
longiflora, Hook. f. l. c. 660, in syn. = grandiflora.
longifolia, Lodd. ex Loud. Hort. Brit. 67; Hook. Bot. Mag. sub t. 3213 = grandiflora.
multiflora, Teijsm. & Binn. in Nederl. Kruidk. Arch. iii. (1855) 393 = khasiana.
Wallichii, Walp. Ann. iii. 39 = Wrightia tomentosa.

BEAUMULIX, Willd. ex Poir. in Dict. Sc. Nat. iv. Suppl. 54 (1816) = **Reaumuria**, Linn. (Tamariscin.).

BEAUPREA, Brongn. & Gris, in Bull. Soc. Bot. Fr. xviii. (1871) 243. *PROTEACEAE*, Benth. & Hook. f. iii. 175.
Balansae, *Brongn. & Gris, l. c.* 245.—N. Caled.
diversifolia, *Brongn. & Gris, l. c.* 244.—N. Caled.
gracilis, *Brongn. & Gris, l. c.* 243.—N. Caled.
Pancheri, *Brongn. & Gris, l. c.* 245.—N. Caled.
spathulaefolia, *Brongn. & Gris, l. c.* 244.—N. Caled.

BEAUTEMPSIA, Gaudich. Voy. Bonite, t. 56 (1844?) = **Capparis**, Linn.
avicenniaefolia, Gaudich. l. c.—Hab.?

BEBBIA, Greene, in Bull. Calif. Acad. i. (1885) 179. *COMPOSITAE*.
atripliciflora, *Greene, l. c.* 181.—Calif.
juncea, *Greene, l. c.* 179.—Calif.

BECCABUNGA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophul.).
Anagallis, Fourr. l. c. = V. Anagallis.
vulgaris, Fourr. l. c. = V. Beccabunga.

BECHIUM, DC. Prod. v. 70 (1836) = **Vernonia**, Schreb. (Compos.).
foliosum, Klatt, in Linnaea, xxxvii. (1871–73) 508.—Madag.
rubricaule, DC. Prod. v. 70 = V. Candollei.
scapiforme, DC. l. c. 71.—Madag.

BECHONNERIA, Hort. ex Carr. in Rev. Hortic. (1867) 320 = **Beschorneria**, Kunth (Amaryll.).

BECKEA, Pers. Syn. i. 234 (1805) = BAECKEA, Burm. f. = **Brunia**, Linn.

BECKEA, St. Hil. in Dict. Sc. Nat. iv. 208 (1804) = **Baeckea**, Linn. (Myrtac.).

BECIUM, Lindl. Bot. Reg. (1842) Misc. 42 = **Ocimum**, Tourn. (Labiat.).
bicolor, Lindl. l. c. = O. filamentosum.

BECKERA, Fresen. in Mus. Senckenb. ii. (1837) 132. t. 8. *GRAMINEAE*, Benth. & Hook. f. iii. 1117.
dioica, Heynh. Nom. ii. 63 = Panicum adoënse.
glabrescens, *Steud. Syn. Pl. Gram.* 117.—Abyssin.
gracilis, *Hochst. in Flora*, xxxviii. (1855) 199.—Abyssin.
mutica, *Hochst. l. c.* xxvii. (1844) 514.—Abyssin.
nubica, *Hochst. l. c.* 512.—Abyssin.
petiolaris, *Hochst. l. c.*—Abyssin.
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 358.—Abyssin.
polystachia, *Fresen. in Mus. Senckenb.* ii. (1837) 132.—Abyssin.
Schimperi, *Hochst. in Flora*, xxvii. (1844) 512.—Abyssin.
uniseta, Steud. Syn. Pl. Gram. 118 = Pennisetum unisetum.

BECKERIA, Bernh. Syst. Verz. Erf. 20 (1800) = **Melica**, Linn. (Gramin.).
ciliata, Bernh. l. c. 40 = M. ciliata.

BECKERIA, Heynh. Nom. ii. 63 (1846) = **Beckera**, Fresen. (Gramin.).

BECKEROPSIS, Fig. & De Not. in Mem. Acc. Torin. Ser. II. xiv. (1854) 365 = **Pennisetum**, Pers. (Gramin.).
nubica, Fig. & De Not. l. c. 366.—Nubia.
petiolaris, Fig. & De Not. l. c. 368.—Abyssin.

BECKIA, Rafin. Sylva Tellur. 103 (1838) = **Osbeckia**, Linn. (Melast.).

BECKMANNIA, Host, Gram. Austr. iii. 5. t. 6 (1805). *GRAMINEAE*, Benth. & Hook. f. iii. 1099.
BRUCKMANNIA, Nutt. Gen. Am. i. 48 (1818).
JOACHIMIA, Tenore, ex Roem. & Schult. Syst. ii. 695 (1817).
erucaeformis, *Host, l. c.*—Europ.; As. temp.; Am. bor.
erucoides, Beauv. Agrost. t. 19. f. 6 = erucaeformis.

BECLARDIA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 69 = **Cryptopus**, Lindl. (Orchid.).
brachystachya, A. Rich. l. c. 71 = Aeonia brachystachya.
elata, A. Rich. l. c. 69 = C. elatus.
macrostachya, A. Rich. l. c. 70 = Aeonia macrostachya.

BECQUERELA, Nees, in Linnaea, ix. (1834) 304 = seq.

BECQUERELIA, Brongn. in Duperr. Voy. Coq. Bot. Phaner. 161. t. 27 (1829). *CYPERACEAE*, Benth. & Hook. f. iii. 1069.
cymosa, *Brongn. l. c.*—Bras.
cymosa, Kunth, Enum. Pl. ii. 361 = Merkeliana.
discolor, *Kunth, l. c.* 362.—Bras.
gigantea, *Hochst. ex Steud. Syn. Pl. Cyp.* 180.—Guiana.
glomerata, Kunth, Enum. Pl. ii. 364 = seq.

BECQUERELIA :—
glomerulata, Brongn. in Duperr. Voy. Coq. Bot. Phan. 163 = Calyptrocarya Palmetto.
Merkeliana, *Nees, in Mart. Fl. Bras.* ii. 1. 191.—Bras.
muricata, *Nees, l. c.* 192.—Bras.
verruculosa, *Nees, l. c.* 191.—Bras.

BEDDOMEA, Hook. f. in Benth. & Hook. f. Gen. i. 336 (1862). *MELIACEAE*, Benth. & Hook. f. l. c.
indica, *Hook. f. l. c.*—Ind. or.
simplicifolia, *Bedd. Fl. Sylv.* i. t. 135.—Ind. or.

BEDFORDIA, DC. in Guillem. Arch. Bot. ii. 332 (1833). *COMPOSITAE*, Benth. & Hook. f. ii. 450.
linearis, *DC. l. c.* 333; *et Prod.* vi. 441.—Tasmania.
salicina, *DC. ll. cc.*—Austral.

BEDOUSIA, Dcnnst. Schluess. Hort. Malab. 31 (1818) = **Casearia**, Jacq. (Samyd.).
aromatica, Rafin. Sylva Tellur. 11 = C. tomentosa.
malabarica, Dcnnst. Schluess. Hort. Malab. 31 = C. tomentosa.

BEDUSIA, Rafin. Sylva Tellur. 11 (1838) = praec.

BEEHSA, Endl. Gen. 1354 (1841) = BEESHA, Kunth = **Bambusa**, Linn. (Gram.).

BEERA, Beauv. in Lestib. Essai Fam. Cyp. 43 (1819) = **Hypolytrum**, Rich. (Cyperac.).
caespitosa, Schrad. ex Kunth, Enum. Pl. ii. 270 = H. strictum.
laxa, Schrad. ex Kunth, l. c. = H. laxum.
silvatica, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 67 = H. fuscum.

BEESHA, Kunth, Syn. i. 253 (1822) = **Bambusa**, Linn. (Gram.).
baccifera, Kunth, l. c. = Melocanna bambusoides.
Fax, Schult. f. in Roem. & Schult. Syst. vii. 1336 = Melocanna humilis.
humilis, Kunth, Enum. Pl. i. 434 = Melocanna humilis.
Rheedii, Kunth, Rev. Gram. i. 141 = Melocanna bambusoides.
travancorica, Bedd. Forest. Man. Bot. 234 = Melocanna travancorica.

BEESHA, Munro, in Trans. Linn. Soc. xxvi. 144 (1868) = **Ochlandra**, Thw. (Gram.).
capitata, Munro, in Trans. Linn. Soc. xxvi. (1868) 145.—Madag.
elegantissima, Kurz, ex Munro, l. c. 146.—Java.
stridula, Munro, l. c. 145 = O. stridula.

BEETHOVENIA, Engl. in Linnaea, xxxiii. (1864–65) 677 = **Ceroxylon**, Humb. (Palm.).
cerifera, Engl. l. c. = C. andicola.

BEFARIA, Mutis, ex Linn. Mant. ii. 152 (1771). *ERICACEAE*, Benth. & Hook. f. ii. 599.
ACUNNA, Ruiz & Pav. Prod. 69. t. 12 (1794).
BEJARIA, Zea, in Anal. Cienc. Nat. iii. (1801) 151, in nota; Vent. Jard. Cels (1800), sub t. 51.
JURGENSENIA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) 1. 151.
aestuans, *Linn. Mant.* ii. 242.—N. Granat.
caxamarcensis, *H. B. & K. Nov. Gen. et Sp.* iii. 294.—Peruv.
cinnamomea, *Lindl. in Gard. Chron.* (1848) 175.—Peruv.
coarctata, Hook. Bot. Mag. t. 4433 = Lindeniana.
coarctata, *Humb. & Bonpl. Pl. Aequin.* ii. 125. t. 121.—Peruv.
cubensis, *Griseb. Cat. Pl. Cub.* 52.—Cuba.
densa, *Planch. ex Hérincq, in Rev. Hortic.* Sér. III. iv. (1850) 185.—Hab.?
denticulata, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 230.—Bolivia.
discolor, *Benth. Pl. Hartw.* 65.—Mexic.
drymifolia, *Linden, ex Hérincq, in Rev. Hortic.* Sér. III. iv. (1850) 185.—N. Granat.
floribunda, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) 543 = discolor.
glabra, Mart. & Gal. l. c. = laevis.
glauca, *Humb. & Bonpl. Pl. Aequin.* ii. 118. t. 117.—N. Granat.

BEFARIA :—
grandiflora, *Humb. & Bonpl. Pl. Aequin.* ii. 222. t. 119.—Ecuador.
guyanensis, *Klotzsch, in Schomb. Fauna et Fl. Guy.* 1088.—Guiana.
hispida, *Poepp. & Endl. Nov. Gen. et Sp.* i. 24. t. 39.—Chili.
laevis, *Benth. Pl. Hartw.* 65.—Mexic.
lanceolata, *Ruiz & Pav. Fl. Per.* iv. t. 413. f. b.—Peruv.
ledifolia, *Humb. & Bonpl. Pl. Aequin.* ii. 124. t. 120.—N. Granat.
Lindeniana, *Hérincq, in Rev. Hortic.* Sér. III. iv. (1850) 181.—Venezuela.
mexicana, *Benth. Pl. Hartw.* 15. n. 92.—Mexic.
Mathewsii, *Field. & Gardn. in Bot. Mag.* t. 4981.—Peruv.
microphylla, Paxt. & Lindl. Flow. Gard. i. 86 = densa.
myrtifolia, *Hérincq, in Rev. Hortic.* Sér. III. iv. (1850) 184.—N. Granat.
oblonga, *Ruiz & Pav. Fl. Per.* iv. t. 413; *Pers. Syn.* ii. 3.—Venezuela.
pallens, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 229.—Peruv.
paniculata, Michx. Fl. Am. Bor. i. 280 = racemosa.
parviflora, *Benth. Pl. Hartw.* 143.—Peruv.
phyllireaefolia, *Benth. l. c.* 225.—N. Granat.
racemosa, *Vent. Jard. Cels*, t. 51.—Am. bor.
resinosa, *Linn. f. Suppl.* 246.—N. Granat.
Schomburgkiana, *Klotzsch, in Schomb. Fauna et Fl. Guy.* 1088.—Guiana.
Sprucei, *Meissn. in Mart. Fl. Bras.* vii. 171.—Peruv.
subsessilis, *Benth. Pl. Hartw.* 143.—Peruv.
tricolor, *Linden, ex Hérincq, in Rev. Hortic.* Sér. III. iv. (1850) 185.—N. Granat.

BEGONIA, [Tourn.] Linn. Gen. ed. II. 516 (1742). *BEGONIACEAE*, Benth. & Hook. f. i. 841.
AUGUSTIA, BARYA, CASPARYA, CYATHOCNEMIS, DONALDIA, DORATOMETRA, EWALDIA, GAERDTIA, GIREOUDIA, GURLTIA, HAAGEA, HUSZIA, ISOPTERIX, ISOPTERYX, KNESEBECKIA, LAUCHEA, LEPSIA, MAGNUSIA, MITSCHERLICHIA, MOSCHKOWITZIA, NEPHROMISCHUS, PETERMANNIA, PILDERIA, PLATYCENTRUM, PRITZELIA, PUTZEYSIA, RACHIA, REICHENHEIMIA, REISSIA, ROSSMANNIA, SASSEA, SAUERIA, SCHEIDWEILERIA, STEINERIA, STIBADOTHECA, TITTELBACHIA, TRACHELANTHUS, TRENDELENBURGIA, WAGENERIA, et WEILBACHIA, Klotzsch, in Abh. Akad. Berl. (1854) 17–134. tt. 1–12.
CLADOMISCHUS, Klotzsch, ex DC. I. c. 278 (1864).
DIPLOCLINIUM, Lindl. Veg. Kingd. 319 (1847).
EUPETALUM, Lindl. Introd. Nat. Syst. ed. II. 440 (1836).
FALKEA, Koen. ex Steud. Nom. ed. II. i. 194, 625 (1840).
IRMA, Bout. ex A. DC. Prod. xv. 1. 406 (1864).
MITCHERLICHIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 124.
MEZIEREA, Gaudich. Voy. Bonite, t. 32 (1839).
PLATYCLINIUM, T. Moore, in Moore & Ayres, Gard. Mag. Bot. ii. (1850) 153.
SPHENDANTHERA, Hassk. in Versl. Kon. Akad. Wetensch. iv. (1855) 139.
STIRADOTHECA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 127.
TRACHELOCARPUS, C. Mucll. in Walp. Ann. iv. 909 (1857).
TRITOMISA, Rafin. Fl. Tellur. ii. 91 (1836).
WAGNERIA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 126.
acerifolia, *H. B. & K. Nov. Gen. et Sp.* vii. 186.—Ecuador.
acetosa, *Vell. Fl. Flum.* x. t. 50.—Bras.
acida, *A. DC. Prod.* xv. 1. 364.—Bras.
acida, Vell. Fl. Flum. x. t. 49 = praec.?
aconitifolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 127.—Bras.
aculeata, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 409 = maculata.
acuminata, *Dryand. in Trans. Linn. Soc.* i. (1791) 166. t. 14.—Jamaica.
acutifolia, Balb. ex A. DC. Prod. xv. 1. 295 = portoricensis.
acutifolia, Jacq. Coll. i. 122 = acuminata.
acutifolia, Herb. Vindob. ex A. DC. Prod. xv. 1. 378 = dentatiloba.

BEGONIA :—
acutiloba, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 14.—Mexic.
adenopoda, Lem. Jard. Fleur. ii. Misc. 17 = verticillata.
acquata, *A. Gray, Bot. U. St. Expl. Exped.* i. 658.—Ins. Luzon.
alaccida, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 637.—Burma.
albida-sctulosa, *Hassk. Cat. Hort. Bog. Alt.* 313.—Hab.?
albiflora, *Hort. ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 126, nomen.—Hab.?
albo-coccinea, *Hook. Bot. Mag.* t. 4172.—Ind. or.
alchemilloides, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 125.—Bras.
alnifolia, *A. DC. l. c.* 132.—N. Granat.
alto-peruviana, *A. DC. l. c.* 123.—Peruv.
amabilis, *Linden, Hort. Linden.* (1859) 1.—Assam.
amoena, *Wall. Cat.* n. 3682.—Reg. Himal.
ampla, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 574.—Afr. trop.
anaimalaiensis, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 48.—Ind. or.
anamalayana, Bedd. in Trans. Linn. Soc. xxv. (1866) 217 = praec.
andamensis, *Parish, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 650.—Ind. or.
anemonoides, *Azara, ex Steud. Nom.* ed. II. i. 193.—Hab.?
angularis, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 407.—Bras.
angulata, *Vell. Fl. Flum.* x. t. 52.—Bras.
angustifolia, *Blume, Enum. Pl. Jav.* 97.—Java.
angustiloba, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 126.—Mexic.
anisosepala, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 576.—Afr. trop.
annobonensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 127.—Afr. trop.
annulata, *C. Koch, in Berl. Gartenz.* i. (1837) 76.—Reg. Himal.
anodaefolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 126.—Mexic.
aptera, *Blume, Enum. Pl. Jav.* 97.—Ins. Celebes.
aptera, *Roxb. Hort. Beng.* 104; *Fl. Ind.* iii. 650.—Ins. Mascar.; Timor.; Ins. Philip.
arborescens, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 408.—Bras.
areolata, *Miq. Pl. Jungh.* 417.—Java.
argentea, Van Houtte, ex A. DC. Prod. xv. 1. 318 = maculata.
argyrocoelis, *Hérincq, in Rev. Hortic.* Sér. III. i. (1847) 111, 165.—Hab.?
argyrostigma, Fisch. ex Link & Otto, Ic. Sel. i. 23. t. 10 = maculata.
Arnottiana, A. DC. Prod. xv. 1. 322 = cordifolia.
asarifolia, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 8 = hydrocotilifolia.
ascotiensis, *J. B. Weber, in Rev. Hortic.* (1874) 17.—Hab.?
aspleniifolia, *Hook. f. ex A. DC. Prod.* xv. 1. 392.—Afr. occ.
atricha, *Miq. ex A. DC. l. c.* 321.—Sumatra.
attenuata, *A. DC. l. c.* 388.—Bras.
attenuata, Klotzsch, ex Regel, Ind. Sem. Hort. Petrop. (1866) 101 = hederacea.
aucubaefolia, Hort. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 122 = incarnata.
aurantiaca, Hort. ex Planch. in Fl. des Serres, Sér. I. v. (1849) 530 = cinnabarina.
auriculata, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 577.—Afr. trop.
auriformis, Hort. Berol. ex Klotzsch, in Abh. Akad. Berl. (1855) 38 = incana.
baccata, *Hook. Bot. Mag.* t. 5554.—Afr. trop.
bahiensis, *A. DC. in Ann. Sc. Nat.* Ser. IV. xi. (1859) 137.—Bras.
Balmisiana, *Ruiz, ex Klotzsch, in Abh. Akad. Berl.* (1855) 38.—Mexic.
barbata, *Wall. Cat.* n. 3679, A. & B. partim.—Ind. or.; Burma.
barbata, Wall. l. c. = rubro-venia.
Barkeri, *Knowl. & Westc. Flor. Cab.* iii. 179. t. 135.—Mexic.
Beddomei, *Hook. f. Bot. Mag.* t. 6767.—Assam.
Bernierii, *A. DC. in Ann. Sc. Nat.* Ser. IV. xi. (1859) 130.—Madag.

BEGONIA :—
besleriaefolia, *Schott, in Spreng. Syst.* iv. 408.—Bras.
bidentata, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 408. —Bras.
bipetala, *Lodd. ex Otto & Dietr. Allg. Gartenz.* ix. (1841) 58.—Ind. or.
biserrata, *Lindl. in Journ. Hort. Soc.* ii. (1847) 313.—Guatemala.
Boissieri, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 126.—Mexic.
Boiviniana, *A. DC. l. c.* 130.—Madag.
boliviensis, *A. DC. l. c.* 122.—Bolivia.
bombycina, Blume, Enum. Pl. Jav. i. 97 = isoptera.
borneensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 128.—Borneo.
Boucheana, *A. DC. Prod.* xv. i. 373.—Venezuela.
Bowringiana, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 120 = laciniata.
bracteata, *Jack, in Malay. Misc.* ii. (1822) vii. 13.—Sumatra.
bracteosa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 132.—Peruv.
Brandisiana, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 58; *et in Flora*, liv. (1871) 295.—Burma.
brasila, A. DC. in Mém. Soc. Phys. Genèv. vii. (1836) 295 = hirtella.
brasiliana, Schrank, ex Steud. Nom. ed. II. i. 193 = praec.
brasiliensis, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 122, nomen.—Bras.
brevicaulis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 134.—Reg. Himal.
Bridgesii, *A. DC. l. c.* 132.—Bolivia.
Brongniartiana, *Lem. Hort. Univ.* v. 355.—Hab.?
broussonetiaefolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 133.—Guatemala.
buddleiaefolia, *A. DC. l. c.* 141.—Peruv.
bulbifera, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 155 = discolor.
bulbifera, Lodd. ex Otto & Dietr. Allg. Gartenz. ix. (1841) 58 = bulbillifera.
bulbillifera, *Link & Otto, Ic. Pl. Rar.* 89. t. 45.—Mexic.
bulbillifera, Moric. ex A. DC. Prod. xv. i. 310 = gracilis.
caespitosa, *Jack, in Malay. Misc.* ii. (1822) vii. 8.—Sumatra.
caffra, Meissn. in Linnaea, xiv. (1840) 501 = Dregei.
canarana, *Miq. Anal. Bot. Ind.* iii. 18.—Ind. or.
candicans, A. DC. Prod. xv. i. 400, lapsu, *vide* Bignonia candicans.
capensis, Blanco, Fl. Filip. ed. I. 724 = rhombicarpa.
capensis, Linn. f. Suppl. 502 = diptera.
cardiocarpa, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 13.—Nicaragua.
carminata ×, *Hort. Fl. Mag.* t. 551 (err. typ. 451).
carnosa, *Teijsm. & Binn. Epim. Lugd. Bat.* (1863) 4. —Malaya.
caroliniaefolia, *Regel, Gartenfl.* i. (1852) 259. t. 25.—Mexic.
carpinifolia, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 20.—Costa Rica.
castaneaefolia, *Otto & Dietr. Allg. Gartenz.* iv. (1836) 356.—Bras.
castaneaefolia, Schott, in Spreng. Syst. iv. App. 407 = fruticosa.
Cathcartii, *Hook. f. & Thoms. Illustr. Pl. Himal.* t. 13. —Reg. Himal.
ciliata, Hort. ex Steud. Nom. ed. II. i. 193, 194 = insignis.
ciliata, H. B. & K. Nov. Gen. et Sp. vii. 136 = hirtella.
ciliata, Pav. ex A. DC. Prod. xv. i. 331 = subciliata.
cinnabarina, *Hook. Bot. Mag.* t. 4483.—Bolivia.
circumlobata, *Hance, in Journ. Bot.* xxi. (1883) 203. —China.
cladocarpa, *Baker, in Journ. Bot.* xx. (1882) 113.—Madag.
Clarkei, *Hook. f. Bot. Mag.* sub t. 5663 et t. 5675.—Bolivia.
clypeifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 576.—Afr. trop.
coccinea, *Hook. Bot. Mag.* t. 3990.—Bras.
coccinea, *Ruiz, ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 127; *A. DC. Prod.* xv. i. 273.—Peruv.
columnaris, *Benth. Pl. Hartw.* 131.—Peruv.

BEGONIA :—
columnaris, Ruiz, ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 127 = cordifolia.
complicata, *A. DC. Prod.* xv. i. 399.—Hab.?
concanensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 126.—Ind. or.
conchaefolia, *A. Dietr. Allg. Gartenz.* xix. (1851) 258. —Costa Rica.
concinna, Schott, in Spreng. Syst. iv. App. 408 = peltata?
confertiflora, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 134 = arborescens.
convolvulacea, *A. DC. in Mart. Fl. Bras.* iv. i. 367.—Bras.
corallina, *Carr. in Rev. Hortic.* (1875) 89.—Bras.
cordata, *Vell. Fl. Flum.* x. t. 34.—Bras.
cordifolia, *Thw. Enum. Pl. Zeyl.* 129.—Ind. or.
coriacea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 122.—Bolivia.
coriacea, Hassk. Cat. Hort. Bogor. Alt. 311 = peltata.
crassicaulis, *Lindl. in Bot. Reg.* (1842) *Misc.* 22.—Guatemala.
crenata, *Dryand. in Trans. Linn. Soc.* i. (1791) 162. t. 14. f. 3.—Ind. or.
crenata, Herb. Finlays. ex A. DC. Prod. xv. i. 401 = prolifera?
crenatiflora, *A. DC. Prod.* xv. i. 306.—Mexic.
crenatifolia, Hemsl. Biol. Centr. Am. Bot. i. 494 = crenatiflora.
crenulata, Schott, ex A. DC. in DC. Prod. xv. i. 358 = angularis.
crinita, *Oliver, in Hook. Bot. Mag.* t. 5897.—Peruv.
cruenta, *R. Grah. ex Steud. Nom.* ed. II. i. 193.—Bras.
cubensis, Hassk. Cat. Hort. Bogor. Alt. 322 = Wrightiana.
cubincola, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 124.—Cuba.
cucullata, Ruiz, ex A. DC. Prod. xv. i. 274 = columnaris.
cucullata, *Willd. Sp. Pl.* iv. 414.—Bras.
cucullifolia, Hassk. Cat. Hort. Bogor. Alt. 311 = semperflorens.
Cumingii, A. Gray, Bot. U. St. Expl. Exped. i. 658 = philippinensis.
Cummingiana, *A. DC. Prod.* xv. i. 320.—Ins. Fiji.
cuneata, Walp. Rep. ii. 214 = spathulata.
cyathophora, *Poepp. & Endl. Gen. et Sp.* i. 7. t. 14.—Peruv.
daedalea, Lem. Illustr. Hortic. vii. (1860) 54. t. 269 = strigillosa.
dasycarpa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 127.—Bras.
dasypoda, Meissn. ex A. DC. Prod. xv. i. 299 = hirtella.
Daveauiana, Godefr. in Rev. Hortic. (1880) 290 = Pellionia Daveauiana.
Davisii, *Veitch, in Hook. f. Bot. Mag.* t. 6252.—Peruv.
dealbata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 5.—Mexic.
decandra, *Pav. ex A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 122.—Mexic.
declinata, *Vell. Fl. Flum.* x. t. 44.—Bras.
delicatula, *Parish, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 652.—Burma.
dentata, Pav. ex A. DC. Prod. xv. i. 307 = angustiloba.
dentatiloba, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 140.—Bras.
denticulata, *H. B. & K. Nov. Gen. et Sp.* vii. 182.—Venezuela.
depauperata, Schott, in Spreng. Syst. iv. App. 408 = rhizocarpa.
derycxiana, Lem. Hort. Univ. (1843–44) Misc. 355, fide A. DC. Prod. xv. i. 343 = nelumbifolia.
Diadema, *Linden, in Illustr. Hortic.* xxix. (1882) t. 446.—Borneo.
dichotoma, *Jacq. Coll.* iii. 250.—Venezuela.
digitata, *Raddi, in Mem. Mod.* xvii. *Fis.* (1820) 406.—Bras.
Digwilliana, *Carr. in Rev. Hortic.* (1875) 420.—Hab.?
dimidiata, Vell. Fl. Flum. x. t. 46 = arborescens.
dioica, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 223 = scutata?
dipetala, R. Grah. in Bot. Mag. t. 2849 = malabarica.
diptera, *Dryand. in Trans. Linn. Soc.* i. (1791) 170.—Bras.

BEGONIA :—
discolor, [R. Br. in] Ait. Hort. Kew. ed. II. v. 184 = Evansiana.
disticha, Hort. Berol. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 122 = suaveolens.
disticha, Link, Enum. Hort. Berol. ii. 396 = stipulacea.
diversifolia, *Knowl. & Westc. Flor. Cab*, t. 14. i. (1837) 27.—Hab.?
diversifolia, R. Grah. in Edinb. N. Phil. Journ. (Jan.–Apr. 1830) 183 = gracilis.
domingensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 124.—Ins. S. Doming.
dominicalis, *A. DC. Prod.* xv. i. 366.—Ins. Dominica.
Donkelaariana, *Lem. Jard. Fleur.* i. (1851) *Misc.* 34. —Mexic.
Dregei, *Otto & Dietr. Allg. Gartenz.* iv. (1836) 357.—Bras.
dubia, *Vell. Fl. Flum.* x. t. 42.—Bras.
Dux, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 637.—Burma.
echinata, Royle, Illustr. Bot. Himal. 313. t. 80. fig. 1 = picta.
echinosepala, *Regel, in Act. Hort. Petrop.* i. (1871) 91.—Bras.
elaeagnifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 579. —Afr. trop.
elata, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 122.—Bras.
elatior, Hort. ex Steud. Nom. ed. II. i. 193 = Dregei.
elatostemmoides, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 576.—Afr. trop.
elegans, *H. B. & K. Nov. Gen. et Sp.* vii. 182.—N. Granat.
elliptica, H. B. & K. l. c. 138 = scandens.
elongata, Wall. Cat. n. 6291 = sinuata.
Epibaterium, *Mart. ex A. DC. in Mart. Fl. Bras.* iv. i. 363.—Bras.
epiphytica, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 580.—Afr. trop.
episcopalis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 644.—Reg. Himal.
erecta, *Vell. Fl. Flum.* x. t. 43.—Bras.
eriocaulis, Vis. Orto Padov. (1842) 135 = tomentosa.
eriocaulon, Neumann, in Rev. Hortic. Sér. III. i. (1847) 166 = tomentosa?
Ermani, Klotzsch, in Abh. Akad. Berl. (1855) 35 = uliginosa.
erminea, *L'Hérit. Stirp. Nov.* 97. t. 47.—Madag.
erosa, *Blume, Enum. Pl. Jav.* i. 96.—Java.
erosa, Wall. Cat. n. 3688, partim = picta, amoena.
erythrocarpa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 121.—Bras.
erythrophylla, *Neuman, in Rev. Hortic.* Sér. III. i. (1847) 166.—Hab.?
Evansiana, *Andr. Bot. Rep.* t. 627.—Malaya; China; Japon.
excelsa, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 571.—Afr. trop.
eximia, *Hort. ex Lem. Illustr. Hortic.* vii. (1860) t. 233.—Hab.?
fagifolia, *Fisch. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 356.—Bras.
fagopyroides, *Kunth & Bouché, Ind. Pl. Hort. Berol.* (1848).—Venezuela.
falcifolia, *Hook. f. Bot. Mag.* t. 5707.—Afr. trop.
falciloba, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 15.—Mexic.
fallax, *A. DC. Prod.* xv. i. 329.—Ind. or.
fasciculata, *Jack, in Malay. Misc.* ii. (1822) vii. 12. —Sumatra.
ferruginea, *Linn. f. Suppl.* 419.—N. Granat.
fibrosa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 652. —Burma.
filipes, *Benth. Bot. Voy. Sulph.* 101.—Panama.
fimbriata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 18.—Mexic.
fimbristipula, *Hance, in Journ. Bot.* xxi. (1883) 202.—China.
Finlaysoniana, Wall. Cat. n. 3684 = prolifera?
Fischeri, *Otto & Dietr. Allg. Gartenz.* iv. (1836) 354. —Bras.
Fischeri, Schrank, Pl. Rar. t. 59 = macroptera.
flaccidissima, *Kurz, in Journ. As. Soc. Beng.* xl. (1872) ii. 308.—Burma.
flexuosa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 142.—Panama.
floccifera, *Bedd. Ic. Pl. Ind. Or.* i. 23.—Ind. or.
floribunda, *Carr. in Rev. Hortic.* (1875) 420.—Hab.?

BEGONIA :—
foliosa, *H. B. & K. Nov. Gen. et Sp.* vii. 183. t. 642. —N. Granat.
foliosa, Poepp. ex A. DC. Prod. xv. 1. 376 = Poeppigiana.
Franconis, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 21.—Mexic.
frigida, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 143.—Hab.?
Froebelii, *A. DC. in Gard. Chron.* (1874) ii. 552.—Ecuador.
fruticosa, *A. DC. in Mart. Fl. Bras.* iv. 1. 377.—Bras.
fusca, *Liebm. ex Klotzsch, in Abh. Akad. Berl.* (1855) 39.—Bras.
fuchsioides, *Hook. Bot. Mag.* t. 4281.—Mexic.
furfuracea, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 571.—Afr. trop.
fusca, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 7.—Mexic.
Galeotti, Hort. Berol. ex Klotzsch, in Abh. Akad. Berl. (1855) 39 = lobata.
Galeottiana, *Lem. Jard. Fleur.* i. (1851) *Misc.* 14.—Bras.
Gardneri, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 138.—Bras.
Gaudichaudii, *Walp. Rep.* v. 769.—Peruv.
geminipara, *Hook. f. & Thoms. Illustr. Him. Pl.* t. 14.—Reg. Himal.
geniculata, Vell. Fl. Flum. x. t. 51 = convolvulacea.
geniculata, Jack, in Malay. Misc. ii. (1822) vii. 15 = isoptera.
geraniifolia, *Hook. Bot. Mag.* t. 3387.—Peruv.
geranioides, *Hook. f. l. c.* t. 5583.—Afr. austr.
gigantea, *Wall. Cat.* n. 3677.—Reg. Himal.
gigantea, Wall. l. c. n. 3677 B = silhetensis.
glabra, Aubl. Pl. Gui. ii. 916 = scandens.
glabra, *Ruiz, ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 128.—Peruv.
glandulifera, *Griseb. Fl. Brit. W. Ind.* 304.—Ind. occ.
glandulosa, *A. DC. Prod.* xv. 1. 339.—Am. centr.
glauca, *Ruiz & Pav. ex A. DC. l. c.* 330.—Peruv.
goëgoënsis, *N. E. Br. in Gard. Chron.* (1882) ii. 71.—Sumatra.
goniotis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 648.—Burma.
Goudotii, *A. DC. in Ann Sc. Nat.* Sér. IV. xi. (1859) 130.—Madag.
gracilis, *H. B. & K. Nov. Gen. et Sp.* vii. 184.—Mexico.
gracillima, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 120.—Peruv.
Grahamiana, Wight, Ic. t. 1811 = albo-coccinea.
grandiflora, Jacq. Coll. i. 128 = diptera.
grandiflora, Knowl. & West. Flor. Cab. i. (1837) 51 = octopetala.
grandis, Dryand. in Trans. Linn. Soc. i. (1791) 163 = Evansiana.
grandis, Otto, ex A. DC. in Mart. Fl. Bras. iv. 1. 369 = vitifolia.
Griffithii, *Hook. Bot. Mag.* t. 4984.—Reg. Himal.
grisea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 138.—Bras.
guaduensis, *H. B. & K. Nov. Gen. et Sp.* vii. 178.—N. Granat.
gunneraefolia, *Linden & André, Illust. Hortic.* xxii. (1875) 106. t. 212.—N. Granat.
guttata, *Wall. Cat.* n. 3671 A.—Malaya.
guttata, Wall. l. c. n. 3671 B = sinuata.
guyanensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 142.—Guiana.
haematotricha, Boiss. ex A. DC. Prod. xv. 1. 298 = humilis.
Hamiltoniana, Lehm. in E. Otto, Hamb. Allgem. Gartenz. vi. (1850) 456 = acuminata.
Hasskarli, *Zoll. & Mor. Syst. Verz. Zoll.* 31.—Java.
Hasskarliana, *Miq. ex A. DC. Prod.* xv. 1. 329.—Sumatra.
hastata, Vell. Fl. Flum x. t. 54 = angularis.
Hatacoa, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 223.—Nepal.
hederacea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 120.—N. Granat.
hederaefolia, *Linden, ex A. DC. Prod.* xv. 1. 401.—Mexic.
heracleifolia, *Cham. & Schlecht. in Linnaea*, v. (1830) 603.—Mexic.
herbacea, *Vell. Fl. Flum.* x. t. 53.—Bras.

BEGONIA :—
hernandiaefolia, Hort. ex Klotzsch, in Monatsb. Berl. Akad. (Maerz 1854) 125 = nelumbiifolia.
hernandiaefolia, Hook. Bot. Mag. t. 4676 = peltata.
hernandiaefolia, Seem. ex A. DC. Prod. xv. 1. 339 = glandulosa.
heterophylla, *Hort. Schoenb. ex Klotzsch, in Monatsb. Berl. Acad.* (Macrz 1854) 122.—Mexic.
heteropoda, *Baker, in Journ. Linn. Soc.* xxi. (1884) 347.—Madag.
Hilariana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 125.—Bras.
hirsuta, *Aubl. Pl. Gui.* ii. 913. t. 348.—Guiana.
hirsuta, Hort. ex A. DC. Prod. xv. 1. 298 = humilis.
hirsuta, Ruiz, ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 127 = cordifolia.
hirsutula, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 575.—Afr. trop.
hirtella, *Link, Enum. Hort. Berol.* ii. 296.—Bras.
hispida, *Schott, ex A. DC. Prod.* xv. 1. 363.—Bras.
holosericea, *Teijsm. & Binn. in Epim. Lugd. Bat.* (1863) 5.—Ins. Ternate.
Holtonis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 141.—N. Granat.
homonyma, Steud. Nom. ed. II. i. 194 = Dregei.
Hookeri, Sweet, Hort. Brit. ed. II. 437 = nitida.
Hookeriana, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 135.—Bras.
Horsfieldii, *Miq. ex A. DC. Prod.* xv. 1. 397.—Sumatra.
Hugelii, *Hort. Berol. ex A. DC. l. c.* 364.—Bras.
humilis, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 353.—Mexic.
humilis, Bonpl. ex Spreng. Syst. ii. 626 = hirsuta.
humilis, Dryand. in Trans. Linn. Soc. i. (1791) 166 = flexuosa.
humilis, Ker-Gawl. in Bot. Reg. t. 284 = diptera.
hydrocotylifolia, *Otto, ex Hook. Bot. Mag.* t. 3968.—Mexic.
hydrophila, Miq. Anal. Bot. Ind. iii. 18 = malabarica.
ignea, *Warsc. in A. DC. Prod.* xv. 1. 306.—Guatemala.
imperialis, *Lem. Illustr. Hortic.* vii. (1860) 53.—Mexic.
incana, *Lindl. in Bot. Reg.* (1841) *Misc.* 39.—Mexic.
incarnata, *Link & Otto, Ic. Pl. Sel.* 37. t. 19.—Mexic.
incarnata, Pav. ex A. DC. Prod. xv. 1. 273 = cordifolia.
incarnata, Seem. Bot. Voy. Herald, 128 = Seemanniana.
incisa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 129.—Ins. Philipp.
inciso-serrata, *A. DC. Prod.* xv. 1. 371.—Bras..
inflata, *C. B. Clarke, in Hook.f. Fl. Brit. Ind.* ii. 636.—Reg. Himal.
insignis, R. Grah. in Bot. Mag. t. 2900 = incarnata.
integerrima, *Spreng. Neue Entdeck.* ii. 174.—Bras.
integrifolia, *Dalz. in Hook. Kew Journ.* iii. (1851) 230.—Ind. or.
involucrata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 15.—Costa-Rica.
isoptera, *Dryand. in Sm. Ic.* t. 43; *et Trans. Linn. Soc.* i. (1791) 160.—Java.
jamaicensis, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 124 = acutifolia.
Jamesoniana, *A. DC. l. c.* 139.—Ecuador.
jatrophaefolia, Hort. ex Klotzsch, in Abh. Akad. Berl. (1855) 39 = heracleifolia.
Josephi, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 126.—Reg. Himal.
Junghuhniana, *Miq. Pl. Jungh.* 418.—Java.
Karwinskyana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 134.—Mexic.
khasiana, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 656.—Reg. Himal.
Kunthiana, *Walp. Ann.* ii. 650.—Bras.
laciniata, *Roxb. Hort. Beng.* 68; *Fl. Ind.* iii. 649.—Reg. Himal.; Burma; China.
laciniosa, *A. DC. Prod.* xv. 1. 340.—Am. centr.
laetevirens, *Van Houtte, ex A. DC. l. c.* 400.—Hab.?
lanceolata, *Vell. Fl. Flum.* x. t. 33.—Bras.
Langsdorffii, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 140 = bidentata.
lantaniaefolia, *A. DC. l. c.* 141.—N. Granat.
lanuginosa, *A. DC. l. c.* 131.—N. Granat.
laurina, Hort. ex A. DC. Prod. xv. 1. 292 = Ottonis.
Lazuli, Linden, in Belg. Hortic. viii. (1858) 166 = xanthina.
lepida, *Blume, Enum. Pl. Jav.* 98.—Java.

BEGONIA :—
lepidota, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 17 = manicata.
leprosa, *Hance, in Journ. Bot.* xxi. (1883) 202.—China.
Liebmannii, *A. DC. Prod.* xv. 1. 345.—Mexic.
Lindeniana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 123.—Cuba.
Lindleyana, *Walp. Rep.* ii. 209.—Guatemala.
Lindleyana, *Warc. ex Klotzsch. in Monatsb. Berl. Acad.* (Maerz 1854) 122.—Am. calid.
lineata, *N. E. Br. in Gard. Chron.* (1882) ii. 199.—Java.
lobata, Schott, in Spreng. Syst. iv. App. 408 = multinervia.
Lobbiana, A. DC. Prod. xv. 1. 355 = parviflora.
Lobbii, *A. DC. l. c.* 390.—Java.
lobulata, *A. DC. l. c.* 339.—Mexic.
loeellata, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 137.—Mexic.
longifolia, *Blume, Cat. Gew. Buitenz.* 102.—Java.
longipes, *Hook. Bot. Mag.* t. 3001.—Mexic.
longipila, Lem. Illustr. Hortic. vii. (1860) 307 = heracleifolia.
longirostris, *Benth. Pl Hartw.* 185.—Ecuador.
longovillosa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 130.—Ins. Philip.
loranthoides, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 580.—Afr. trop.
Lubbersii, *E. Morr. in Belg. Hortic.* xxxiii. (1883) 155. t. 13.—Bras.
lucida, Haw. Saxifr. Enum. 197 = humilis.
lucida, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 16 = Kunthiana.
lucida, Otto & Dietr. Allg. Gartenz. xvi. (1848) 162 = scandens.
lucida, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 208; *Contrib.* 57.—Paraguay.
ludicra, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 133.—Mexic.
luxurians, *Scheidw. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 131.—Bras.
Lyallii, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 144.—Madag.
Lyncheana, *Hook. f. Bot. Mag.* t. 6758.—N. Granat.
macra, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 121.—N. Granat.
macrophylla, *Dryand. in Trans. Linn. Soc.* i. (1791) 164.—Jamaica.
macrophylla, Lam. Encyc. i. 394 = diptera.
macroptera, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 122.—Bras.
macrotis, *Vis. in Act. Ist. Ven. Sc.* Ser. III. iv. (1858-59) 138.—Hab.?
maculata, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 406.—Bras.
magnifica, *Linden, in Belg. Hortic.* vi. (1856) 5.—N. Granat.
magnifica, Warsc. ex A. DC. Prod. xv. 1. 269 = ferruginea.
magnifolia, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 8.—Malaya.
malabarica, *Lam. Encyc.* i. 393.—Ind. or.
malabarica, Roxb. Hort. Beng. 68; Fl. Ind. iii. 648 = Roxburghii.
malabarica, Buch.-Ham. in Wall. Cat. n. 3676 D = megaptera.
malvacea, *Klotzsch, in Abh. Akad. Berl.* (1855) 33.—Bras.
manicata, *Cels, ex Vis. Orto Padov.* (1842) 135.—Mexic.
manillensis, *A. DC. in Ann. Sc. Nat.* Ser. IV. xi. (1859) 129.—Ins. Philipp.
Mannii, *Hook. f. Bot. Mag.* t. 5434.—Afr. trop.
maracayuensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 209; *Contrib.* 58.—Paraguay.
marmorata, Van Houtte, ex Planch. in Fl. des Serres, ix. (1853-54) 243. t. 952-53 = xanthina.
martabanica, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 136.—Burma.
Martiana, *Link & Otto, Ic. Pl. Select. Hort. Berol.* 49.—Mexic.
Martiana, Schlecht. in Linnaea, v. (1830) 604 = incarnata.
martinicensis, *A. DC. in Ann. Sc. Nat.* Ser. IV. xi. (1859) 123.—Ins. Martinic.
mascariensis, *Boj. Hort. Maurit.* 271.—Ins. Mascar.
Maurandiae, *A. DC. in Ann. Sc. Nat.* Ser. IV. xi. (1859) 119.—N. Granat.

BEGONIA :—
maxima, *Hort. Berol. ex Klotzsch, in Abh. Akad. Berl.* (1855) 39.—Mexic.
maynensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 126.—Peruv.
megaphylla, *A. DC. l. c.* 133.—Bras.
megaptera, *A. DC. l. c.* 134.—Reg. Himal.; Burma.
Meissneri, *Wall. Cat.* n. 6294.—Ind. or.
membranacea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 139.—Bras.
meridensis, *A. DC. l. c.* 123.—Venezuela.
metallica, L. Sm. in Floral Mag. (1876) t. 197 = incarnata.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 409.—Bras.
Meyeri, *Hook. Bot. Mag.* t. 4100.—Mexic.
Meyeri, Otto & Dietr. Allg. Gartenz. iv. (1836) 349 = tomentosa.
micrantha, Steud. Nom. ed. II. i. 194 = parviflora.
micranthera, *Griseb. in Goett. Abh.* xix. (1874) 148.—Reg. Argent.
microcarpa, *A. DC. Prod.* xv. i. 311.—Ecuador.
microphylla, *A. DC. l. c.* 375.—Reg. Granat.
microphylla, Willd. ex Klotzsch, in Abh. Akad. Berl. 1854 (1855) 62 = foliosa.
microptera, *Hook. f. Bot. Mag.* t. 4974.—Borneo.
miniata, Planch. & Linden, in Fl. des Serres, Sér. I. viii. (1852–53) 105 = fuchsioides.
minima, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 48 = crenata.
minor, Jacq. Coll. i. 126 = nitida.
modesta, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 20.—Mexic.
modestiflora, *Kurz, in Flora*, liv. (1871) 296.—Reg. Himal.; Burma.
mollis, *A. DC. Prod.* xv. i. 391.—Java.
monadelpha, *Ruiz & Pav. ex A. DC. l. c.* 286.—Peruv.
monophylla, *Pav. ex A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 121.—Mexic.?
monophylla, Pourr. ex A. DC. Prod. xv. i. 403 (est Orchidea).
monoptera, *Link & Otto, Ic. Pl. Rar.* t. 14.—Mexic.
Moritziana, Klotzsch, in Abh. Akad. Berl. (1854) 31 = tovarensis.
Moritziana, Kunth & Bouché, Ind. Sem. Hort. Berol. (1848) 16 = scandens.
moulmeinensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 643.—Burma.
multangula, *Blume, Enum. Pl. Jav.* i. 96.—Java.
multibulbillosa, *Mart. ex Steud. Nom.* ed. II. i. 194.—Bras.
multiflora, *Benth. Pl. Hartw.* 185.—N. Granat.
multinervia, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 18.—Costa Rica.
muricata, Blume, Cat. Gew. Buitenz. 103 = tuberosa.
muricata, *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 156.—Bras.
nana, *L'Hérit. Stirp. Nov.* 99. t. 48.—Madag.
natalensis, *Hook. Bot. Mag.* t. 4841.—Afr. austr.
neglecta, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 139.—Bras.
nelumbiifolia, *Cham. & Schlecht. in Linnaea*, v. (1830) 604.—Mexic.
nemophila, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 106, 108 = Cathcartii.
neocomensium, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 138.—Bras.
nervosa, [Desf.] Hort. Par. ex H. B. & K. Nov. Gen. et Sp. vii. 177 = cucullata.
nigrescens, *Van Houtte.—Cf. Hamb. Gartenz.* viii. (1852) 9.—Hab.?
nigritarum, Steud. Nom. ed. I. 104; ed. II. i. 194 = malabarica?
nigro-venia, Regel, Gartenfl. (1867) 163 = glandulosa.
nitida, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 352.—Jamaica.
nossibea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 130.—Ins. Nossibé.
Novo-Granatae, A. DC. Prod. xv. i. 517 = rosacea.
nivea, *Parish, ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 81.—Tenasserim.
nummulariaefolia, *Putzeys.—Cf. Bot. Zeit.* xi. (1853) 716—Hab.?
oaxacana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 127.—Mexic.
obliqua, L'Hérit. Stirp. Nov. ii. 46 = nitida.
obliqua, Jacq. Obs. ex Steud. Nom. ed. II. i. 194 = macrophylla.

BEGONIA :—
obliqua, Linn. Sp. Pl. 1056 = acuminata.
obliqua, Ruiz, ex A. DC. Prod. xv. i. 333 = cyathophora.
obliqua, Thunb. Fl. Jap. 231 = Evansiana.
obliqua, *Vell. Fl. Flum.* x. t. 48.—Bras.
octopetala. *L'Hérit. Stirp. Nov.* 101.—Peruv.
odorata, Willd. Enum. Hort. Berol. Suppl. 64 = diptera.
olbia, *Kerchove, in Rev. Hortic. Belge* (1883), 241—Bras.
oligophylla, *Blume, ex Miq. Fl. Ind. Bat.* i. 692.—Java.
opuliflora, *Putz. in Fl. des Serres*, Sér. I. x. (1854–55) 71. t. 995.—N. Granat.
opulifolia, Loud. Encyc. Pl. 1540 = pracc.?
orbiculata, *Jack, in Malay. Misc.* ii. (1812) VII. 9.—Sumatra.
oregana, *Hort. ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 126.—Bras.
Ottoniana ×, *A. DC. Prod.* xv. i. 399.
Ottonis, *Walp. Rep.* ii. 212.—N. Granat.
ovatifolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 132.—Reg. Himal.
oxyloba, *Welw. ex Hook. f. in Oliver, Fl. Trop. Afr.* ii. 573.—Afr. trop.
oxyphylla, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 140.—Bras.
oxysperma, *A. DC. l. c.* 122.—As. trop.
paleacea, *Kurz, in Flora*, liv. (1871) 297.—Burma.
paleata, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 137.—Bras.
palmaris, *A. DC. l. c.* 126.—Mexic.
palmata, D. Don, Prod. Fl. Nep. 223 = laciniata?
palmata, Pav. ex A. DC. Prod. xv. i. 307 = palmaris.
palustris, *Benth. Pl. Hartw.* 184. n. 1022.— N. Granat.
paniculata, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 209; *Contrib.* 59.—Paraguay.
pantherina, *Putz. ex Linden, Cat.* n. 17 (1862) 2.—Mexic.
papillosa, R. Grah. in Hook. Bot. Mag. t. 2846 = incarnata.
papillosa, *Lindl. Bot. Reg.* (1841) *Misc.* 39.—Hab.?
paraguayensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 207; *Contrib.* 56.—Paraguay.
Parishii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 651.—Burma.
parviflora, Liebm. ex A. DC. Prod. xv. i. 383 = Franconis.
parviflora, *Poepp. & Endl. Nov. Gen. et Sp.* i. 7. t. 12.—Chili.
parviflora, Steud. Nom. ed. II. i. 194 = parvifolia.
parviflora, Wall. Cat. n. 3674 = modestiflora.
parvifolia, R. Grah. in Bot. Mag. t. 3720 = Dregei.
parvifolia, *Klotzsch, in Abh. Akad. Berl.* (1855) 33.—Bras.
parvifolia, *Schott, in Spreng. Syst.* iv. *App.* 408.—Bras.
parvipeltata, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 139.—Bras.
parvuliflora, *A. DC. l. c.* 136.—Burma.
pastoënsis, *A. DC. l. c.* 121.—N. Granat.
patens, Griseb. ex A. DC. Prod. xv. i. 377 = arborescens.
patula, Haw. Pl. Succ. Suppl. 100 = macroptera.
pauciflora, Lindl. Bot. Reg. t. 471 = macroptera.
paulensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 125.—Bras.
Pavoniana, *A. DC. l. c.* 142.—Peruv.
Pearcei, *Hook. f. Bot. Mag.* t. 5545.—Bolivia.
pedata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 10.—Mexic.
pedunculosa, *Wall. Pl. As. Rar.* i. 82. t. 97.—Reg. Himal.
pedunculosa, Wall. Cat. n. 3672 B = khasiana.
peltata, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 133 = Hasskarlii.
peltata, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 138.—Bras.
peltata, Otto & Dietr. Allg. Gartenz. ix. (1841) 58 = incana.
peltifolia, Schott, in Spreng. Syst. iv. App. 408 = parvipeltata.
pentaphylla, Walp. Rep. ii. 209 = muricata.
peperomioides, *Hook. f. in Oliver, Fl. Trop Afr.* ii. 575.—Afr. trop.

BEGONIA :—
peponifolia, *Vis. Ind. Sem. Hort. Patav.* (1847).—*Cf. Linnaea*, xxiv. (1851) 180.—Jamaica.
perpusilla, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 143.—Ins. Nossibé.
peruviana, *A. DC. l. c.* 133.—Peruv.
petalodes, *Lindl. Bot. Reg.* t. 1757.—Bras.
philippinensis, *A. DC. in DC. Prod.* xv. i. 320.—Ins. Philipp.
phyllomaniaca, *Mart. in Ann. Sc. Nat.* Sér. III. xix. (1853) 366.—Bras.
physalifolia, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 19 = scandens.
picta, Hort. Henders. ex A. DC. Prod. xv. i. 350 = Griffithii.
picta, *Sm. Exot. Bot.* ii. 81. t. 101.—Reg. Himal.
Pilderia, *A. DC. Prod.* xv. i. 380.—Venezuela.
pilifera, *A. DC. l. c.* 337.—Am. centr.
pilosa, *Jack, in Malay. Misc.* ii. (1822) VII. 13.—Sumatra.
pinetorum, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 131.—Mexic.
plagiata, Linden, ex A. DC. Prod. xv. i. 312 = oaxacana.
plantanifolia, *Schott, in Spreng. Syst.* iv. *App.* 407.—Bras.
plebeja, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 8.—Nicaragua.
pleiopetala, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 121.—Bras.
Plumieri, *Kunth, in DC. Prod.* xv. i. 295.—Ins. S. Doming.
poculifera, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 574.—Afr. trop.
poecila, *C. Koch, in C. Koch & Fint. Wochenschr.* (1858) 338.—Hab.?
Poeppigiana, *A. DC. Prod.* xv. i. 376.—Bras.
Pohliana, Klotzsch, in Abh. Akad. Berl. (1855) 33 = macroptera.
polygonata. *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 12.—Mexic.
polygonifolia, *A. DC. in Mart. Fl. Bras.* iv. i. 362.—Bras.
polygonoides, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 571.—Afr. trop.
polypetala, *A. DC.—Cf. 'Garden'* (1878) 531.—Peruv.
populifolia, H. B. & K. Nov. Gen. et Sp. vii. 142 = Balmisiana.
populifolia, Klotzsch, in Abh. Akad. Berl. (1855) 30 = tovarensis.
populifolia, Schott, in Spreng. Syst. iv. App. 408 = scandens.
populnea, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 128.—Bras.
Porteriana, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 51 = humilis.
portoricensis, *A. DC. Prod.* xv. i. 295.—Porto Rico.
Prieurei, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 142.—Guiana.
Princeps, *Hort. Berol. ex Klotzsch, in Abh. Akad. Berl.* (1855) 39.—Bras.
prismatocarpa, *Hook. Bot. Mag.* t. 5307.—Afr. trop.
procridifolia, *Wall. Cat.* n. 6292.—Burma.
procumbens, *Vell. Fl. Flum.* x. t. 36.—Bras.
prolifera, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 135.—Burma; Malaya.
pruinata, *A. DC. Prod.* xv. i. 338.—Costa Rica.
pulchella, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 407.—Bras.
pulchra, Herb. Schreb. ex A. DC. Prod. xv. i. 293 = nitida.
punctata, Hort. ex Steud. Nom. ed. II. i. 194 = maculata.
punctata, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. Berol. i. 16 = heracleifolia.
Purdieana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 124.—Jamaica.
purpurea, Sw. Prod. Veg. Ind. Occ. 85 = nitida.
pusilla, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 120.—Bolivia.
pustulata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 6.—Mexic.
Putzeysiana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 139.—Venezuela.
quercifolia, *A. DC. l. c.* 129.—Ins. Philipp.
racemosa, *Jack, in Malay. Misc.* ii. (1822) VII. 14.—Sumatra.

BEGONIA:—
radiata, R. Grah. in Edinb. N. Phil. Journ. (July 1833) 182 = heracleifolia.
radicans, *Vell. Fl. Flum.* x t. 39.—Bras.
ramentacea, *Paxt. Mag. Bot.* x. 73.—Bras.
reniformis, Bedd. in Madr. Journ. Sc. xxii. (1861) 72 = anaimalaiensis.
reniformis, Dryand. in Trans. Linn. Soc. i. (1791) 161. t. 14. f. 1-2 = longipes.
reniformis, Hook. Bot. Mag. t. 3225 = vitifolia.
reniformis, Hort. Berol. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 24 = Dregei.
reniformis, Pav. ex A. DC. Prod. xv. i. 308 = Balmisiana.
reniformis, *Vell. Fl. Flum.* x. t. 40.—Bras.
repanda, *Blume, Enum. Pl. Jav.* i. 97.—Sumatra.
repens, Blume, l. c. 95 = mollis.
repens, Lam. Encyc. i. 394 = rotundifolia.
repens, Hemsl. Biol. Centr. Am. Bot. i. 497 = Liebmanni.
repens, Ruiz & Pav. ex A. DC. Prod. xv. i. 333 = Rossmanniae.
repens, Schott, ex A. DC. l. c. 365 = convolvulacea.
repens, *Vell. Fl. Flum.* x. t. 35.—Bras.
reptans, *Benth. Pl. Hartw.* 62.—Mexic.
reptans, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 5 = Liebmanni.
reticulata, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 134.—Bras.
Rex, *Putz. in Fl. des Serres*, ii. (1857) 141. t. 1255.—Reg. Himal.
rhizocarpa, *Fisch. ex A. DC. Prod.* xv. i. 388.—Bras.
rhizocaulis, *Hort. Berol. ex A. DC. l. c.* 340.—Am. centr.
rhombicarpa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 129.—Ins. Luzon.
Richardii ×, [*C. Moore*] *Cat. Bot. Gard. Sydney* (1857), 9.—Hab.?
Richardsiana, *T. Moore, in Gard. Chron.* (1871) 1065. —Afr. austr.
Richardsoniana, *Houll. in Rev. Hortic.* (1872) 333.—Hab.?
ricinifolia ×, A. Dietr. Allg. Gartenz. xv. (1847) 282.
Riedelii, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 137.—Bras.
rigida, *Regel, ex A. DC. Prod.* xv. i. 373.—Bras.
robusta, *Blume, Enum. Pl. Jav.* i. 96.—Java.
robusta, Zoll. ex A. DC. Prod. xv. i. 275 = multangula.
Rodwelli, *Hort.—Cf. Gard. Chron.* (1867) ii. 481.—Hab.?
Roezli, Regel, Gartenfl. (1876) 194; et in Act. Hort. Petrop. iv. (1876) 334 = Lynchiana.
rosacea, *Putz. in Belg. Hortic.* vii. (1857) 196; *et in Fl. des Serres*, Sér. II. ii. (1857) 25. t. 1194. —N. Granat.
rosaeflora, *Hook. f. Bot. Mag.* t. 5680.—Peruv.
rosea, *A. DC. Prod.* xv. i. 299.—Am. centr.
rosea, Pav. ex A. DC. l. c. 273 = cordifolia.
Rossmanniae, *A. DC. l. c.* 333.—Peruv.
rostrata, *Welw. ex Hook. f. in Oliver, Fl. Trop. Afr.* ii. 578.—Afr. trop.
rotata, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 11 = caroliniaefolia.
rotunda, *Vell. Fl. Flum.* x. t. 38.—Bras.
rotundifolia, *Lam. Encyc.* i. 394.—Ind. occ.
Roxburghii, *A. DC. Prod.* xv. i. 398.—Reg. Himal.; Burma.
Roylei, *Linden, in Belg. Hortic.* vii. (1857) 197.—Ind. or.
rubella, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 223.—Nepal.
rubella, Miq. Anal. Bot. Ind. iii. 18 = canarana.
rubella, Wall. Cat. n. 3687 = scutata.
rubicunda, Hort. Turic. ex A. DC. Prod. xv. i. 384 = Dregei.
rubra, *Blume, Enum. Pl. Jav.* i. 96.—Java.
rubricaulis, *Hook. Bot. Mag.* t. 4131.—Peruv.
rubro-nervia, *Hort. ex Klotzsch, in Abh. Akad. Berl.* (1855) 40.—Ind. or.
rubro-pilosa, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 125.—Bras.
rubro-setulosa, *A. DC. Prod.* xv. i. 390.—Hab.?
rubro-venia, *Hook. Bot. Mag.* t. 4689.—Reg. Himal.
rufa, *Thunb. in Flora*, iv. (1821) 330.—Bras.
rugosa, *Hort. Schoenb. ex Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 126.—Hab.?
rupestris, *Baxt. ex Loud. Hort. Brit. Suppl.* iii. 498. —Bras.

BEGONIA:—
rupicola, *Miq. Pl. Jungh.* 418.—Java.
rutilans, *Hort. Van-Houtte, ex A. DC. Prod.* xv. i. 373.—Hab.?
sagittata ×, Hort.—Cf. Gard. Chron. (1868) i. 1317.
salicifolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. —(1859) 128.—Bras.
sandalifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 649.—Burma.
Sandersii, *Hort. Kew. ex A. DC. Prod.* xv. i. 400, nomen.—Hab.?
sanguinea, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 409. —Bras.
sarcophylla, *Liebm. in Kjoeb. Vidensk. Meddel.* (1852) 12.—Mexic.
Sartorii, Liebm. l. c. 14 = praec.
Satrapis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 638. —Reg. Himal.
saxatilis, *Blume, Enum. Fl. Jav.* 95.—Ins. Philipp.
saxicola, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 144.—Bras.
Saxifraga, *A. DC. l. c.* 138.—Bras.
scabrida, *A. DC. Prod.* xv. i. 367.—Venezuela.
scandens, *Vell. Fl. Flum.* x. t. 41.—Bras.
scandens, Hort. Schoenb. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 126 = convolvulacea.
scandens, *Sw. Prod. Veg. Ind. Occ.* 86.—Am. austr.
scapigera, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 572.—Afr. trop.
schizolepis, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 17 = manicata.
Schlumbergeriana, *Lem. Illustr. Hortic.* (1858) *Misc.* 61.—Bras.
Schmidti, Hort. ex Wien. Illustr. Gart. Zeit. (1880) 250 = Schmidtiana.
Schmidtiana, *Regel, Gartenfl.* (1879) 321.—Bras.
Schottiana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 140.—Bras.
scutata, *Wall. Cat.* n. 3686. partim.—Reg. Himal.
scutata, Wall. l. c. A = Josephi.
scutellata, Liebm. in Kjoeb. Vidensk. Meddel. 1852 (1853) 9 = conchaefolia.
scutifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 572.—Afr. trop.
Scutulum, *Hook. f. l. c.* 575.—Afr. trop.
Sedeni ×, Hort.—Cf. Gard. Chron. (1870) 38.
Seemanniana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 133.—Am. centr.
Selloi, *Lodd. ex Steud. Nom.* ed. II. i. 194—Bras.
Sellowii, Klotzsch, in Abh. Akad. Berl. (1855) 28 = semperflorens.
semiovata, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 22.—Am. centr.
semperflorens, Hook. Bot. Mag. t. 2920 = Hookeri.
semperflorens, *Link & Otto, Ic. Pl. Rar.* t. 9.—Bras.
sericoneura, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 13.—Am. centr.
sericoneura, Seem. Bot. Voy. Herald, 128 = pilifera.
serotina, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 121.—Guayaquil.
sessilifolia, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 577. —Afr. trop.
setaria, Hort. Angl. ex A. DC. Prod. xv. i. 293 = semperflorens.
setifera, *A. DC. l. c.* 338.—Am. centr.
setosa, *Klotzsch, in Abh. Akad. Berl.* (1855) 31.—Bras.
setulosa, *Bertol. in Nov. Comm. Bon.* iv. (1840) 437.—Guatemala.
sikkimensis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 134.—Reg. Himal.
silhetensis, *C. B Clarke, in Hook. f. Fl. Brit. Ind.* ii. 636.—Reg. Himal.
sinensis, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 125 = Evansiana.
sinuata, R. Grah. in Edinb. N. Phil. Journ. (Jan. 1838) 192 = Dregei.
sinuata, *E. Mey. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 357.—Afr. austr.
sinuata, *Wall. Cat.* n. 3680.—Burma; Malaya.
smilacina, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 137.—Bras.
socotrana, *Hook. f. in Gard. Chron.* (1881) i. 8.—Ins. Socotr.
solananthera, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 128.—Bras.
sparsipila, *Baker, in Saund. Ref. Bot.* t. 340.—Am. centr.?
spathulata, Lodd. Bot. Cab. t. 107 = cucullata.

BEGONIA:—
speciosa, Hort. Van Houtte, ex A. DC. Prod. xv. i. 293 = nitida.
spilotophylla, *F. Muell. Not. Papuan Pl.* iv. 67.—N. Guinea.
splendens, Hort. Turic. ex A. DC. Prod. xv. i. 375 = Putzeysiana.
splendida, Hort. Rollison, ex A. DC. l. c. 275 = robusta.
Spruceana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 142.—Peruv.
squamulosa, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 579. —Afr. trop.
squarrosa, *Liebm. in Kjoeb. Vidensk. Meddel.* 1852 (1853) 7.—Mexic.
squarrosa, Seem. Bot. Voy. Herald, 128 = stigmosa.
Stella, *Regel, Gartenfl.* (1875) 88.—Hab.?
stenophylla, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 137.—Bras.
stictopoda, *Miq. ex A. DC. Prod.* xv. i. 391.—Sumatra.
stigmosa, *Lindl. Bot. Reg.* (1845) *Misc.* 32.—Mexic.
stipulacea, Willd. Sp. Pl. iv. 414 = angularis.
strigillosa, *A. Dietr. Allg. Gartenz.* xix. (1851) 330.—Am. centr.
strigulosa, *A. DC. Prod.* xv. i. 399.—Hab.?
suaveolens, Haw. in Lodd. Bot. Cab. t. 69 = diptera.
suaveolens, Klotzsch, in Abh. Akad. Berl. (1855) 26 = nitida.
subciliata, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 132.—Peruv.
subhumilis, *A. DC. l. c.* 124.—Guayaquil.
sublobata, *Jack, in Malay. Misc.* ii. (1822) n. vii. 10. —Sumatra.
subovata, Wall. Cat. n. 3683 = ovatifolia.
subpeltata, Regel, Ind. Sem. Hort. Petrop. (1866) 102 = incarnata.
subpeltata, *Wight, Ic.* t. 1812.—Ind. or.
subperfoliata, *Parish, ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 81.—Burma.
subrotunda, Wall. Cat. n. 6293 = sinuata.
subvillosa, *Klotzsch, in Abh. Akad. Berl.* (1855) 32.—Bras.
suffruticosa, *Meissn. in Linnaea*, xiv. (1840) 502.—Afr. austr.
suffruticosa, Pav. ex A. DC. Prod. xv. i. 311 = Boissieri.
sulcata, *Scheidw. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 130.—N. Granat.
surculigera, *Kurz, in Flora*, liv. (1871) 296.—Burma.
Sutherlandi, *Hook. f. Bot. Mag.* t. 5689.—Afr. austr.
sylvatica, *A. DC. Prod.* xv. i. 365.—Bras.
sylvestris, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 140.—Bras.
tanacetifolia, Hort. ex Steud. Nom. ed. II. i. 194 = heracleifolia.
Telfariae, Steud. l. c. sphalm. vide Bignonia Telfairiae.
tenella, D. Don, Prod. Fl. Nep. 223 = amoena.
tenera, *Dryand. in Trans. Linn. Soc.* i. (1791) 169. t. 16.—Zeylan.
tenuicaulis, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 120.—Bolivia.
tenuifolia, *Dryand. in Trans. Linn. Soc.* i. (1791) 162. t. 14.—Java.
tessaricarpa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 636.—Assam.
Teuscheri, *Linden, ex André, in Illustr. Hortic.* (1879) 137. t. 358.—Malaya.
Thomsonii, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 134.—Reg. Himal.
Thwaitesii, Hook. Bot. Mag. t. 4692 = tenera.
tomentosa, *Schott, in Spreng. Syst.* iv. *App.* 408.—Bras.
tovarensis, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 122.—Mexic.
trachyptera, *Benth. Pl. Hartw.* 184.—N. Granat.
trichocarpa, *Dalz. in Hook. Kew Journ.* iii. (1851) 230. —Ind. or.
trichopoda, *Miq. Fl. Ind. Bat.* i. i. 1098.—Sumatra.
tricuspidata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 637.—Burma.
triradiata, *C. B. Clarke, l. c.*—Burma.
tropaeolifolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 120.—N. Granat.
truncata, Vell. Fl. Flum. x. t. 47 = tomentosa vitifolia.
tuberosa, Herb. Wight, ex Wall. Cat. n. 3675 = malabarica.

BEGONIA:—
tuberosa, Heyne, ex A. DC. Prod. xv. 1. 391 = dipetala.
tuberosa, *Lam. Encyc.* i. 393.—Ins. Molucc.
tuberosa, Pav. ex A. DC. Prod. xv. 1. 311 = bulbillifera.
tuberosa, Ruiz, ex A. DC. l. c. 281 = geraniifolia.
uliginosa, *Klotzsch, in Abh. Akad. Berl.* (1855) 35.—Bras.
ulmifolia, *Willd. Sp. Pl.* iv. 418.—Venezuela.
umbellata, *H. B. & K. Nov. Gen. et Sp.* vii. 143.—N. Granat.
umbilicata, Hort. ex Planch. in Fl. des Serres, Sér. I. viii. (1852–53) sub t. 810 = hernandiaefolia.
umbrata, *Hort. ex A. DC. Prod.* xv. 1. 396.—Hab.?
uncinata, Hort. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 124 = Dregei.
undulata, *Schott, in Spreng. Syst.* iv. *App.* 408.—Bras.
urophylla, *Hook. Bot. Mag.* t. 4855.—Mexic.
Urticae, *Linn. f. Suppl.* 420.—N. Granat.
urticaefolia, Sm. Ic. Fasc. ii. 45 = tuberosa.
urticaefolia, Hort. ex Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 127 = B. Pilderia.
urticifolia, F. G. Dietr. Vollst. Gartenlex. ii. 177 = Urticae.
vaginans, *Vell. Fl. Flum.* x. t. 37.—Bras.
valdensium, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 138.—Bras.
variana, *A. DC. l. c.* 135.—Java.
Veitchii, *Hook. f. in Gard. Chron.* (1867) 734.—Peruv.
vellerea, *Klotzsch, in Monatsb. Berl. Acad.* (Maerz 1854) 122.—Bras.
Velloziana, Walp. Rep. ii. 216 = repens.
velutina, Hort. Vindob. ex Klotzsch, in Abh. Akad. Berl. (1855) 41 = lobata.
velutina, Parish, ex Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 81 = parvuliflora.
vernicosa, Hort. Berol. ex Klotzsch, in Abh. Akad. Berl. (1855) 40 = lobata.
Verschaffeltiana ×, Lem. Illustr. Hortic. ii. (1855) t. 68 = Verschaffeltii.
Verschaffeltii ×, *Regel, Gartenfl.* (1855) 248. t. 132.
verticillata, Vell. Fl. Flum. x. t. 45 = digitata.
verticillata, *Hook. Ic. Pl.* t. 811.—Burma.
villosa, Gardn. in Hook. Lond. Journ. i. (1842) 186 = macroptera.
villosa, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 122 = subvillosa.
villosa, Lindl. Bot. Reg. t. 1252 = hirtella.
violaefolia, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 134.—Mexic.
viridiflora, *A. DC. l. c.* 132.—Peruv.
vitifolia, Lindl. Bot. Reg. (1852) Misc. 21 = Lindleyana.
vitifolia, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 407.—Bras.
Wageneriana, *Hook. Bot. Mag.* t. 4988.—Bras.
Wallichiana, *Steud. Nom.* ed. II. i. 194; *Lehm. in Linnaea*, xxiv. (1851) 213 (sp. dubia).—Ind. or.
Walpersii, *Heynh. Nom.* ii. 63.—Venezuela.
Warscewiczii, Neuman, in Rev. Hortic. Sér. III. i. (1847) 166 = conchaefolia.
Weddeliana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 122.—Bolivia.
weltoniensis ×, *Hort.—Cf. J. B. Weber, in Rev. Hortic.* (1875) 105.
Wightiana, Wall. Cat. n. 3673 = albo-coccinea.
Wrightiana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 123.—Cuba.
xanthina, *Hook. Bot. Mag.* t. 4683 et t. 5107.—Reg. Himal.
zebrina, Hort. Angl. ex Lond. Encyc. Pl. Suppl. ii. 1506 = angularis.
Zollingeriana, *A. DC. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 135.—Java.

BEGONIELLA, Oliver, in Trans. Linn. Soc. xxviii. (1873) 513. *BEGONIACEAE.*
angustifolia, *Oliver, in Hook. Ic. Pl.* t. 1487.—N. Granat.
Kalbreyeri, *Oliver, l. c.* t. 1352.—N. Granat.
Whitei, *Oliver, in Trans. Linn. Soc.* xxviii. (1873) 513.—N. Granat.

BEHAIMIA, Griseb. Cat. Pl. Cub. 77 (1859). *LEGUMINOSAE*, Benth. & Hook. f. i. 1002.
cubensis, *Griseb. l. c.* 78.—Cuba.

BEHEN, Hill, Veg. Syst. iv. 41 (1762) = **Vernonia**, Schreb. (Compos.).
Jacea, Hill, Hort. Kew. 68 = Centaurea Jacea.
noveborascense, Hill, l. c. = V. noveboracensis.
praealtum, Hill, l. c. = V. noveboracensis et altissima.

BEHEN, Moench, Meth. 709 (1794) = **Silene**, Linn. (Caryoph.).
angustifolius, Schott, ex Steud. Nom. ed. II. i. 194 = S. Cucubalus.
glareosus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 343 = S. Cucubalus?
puberulus, Fourr. l. c. = S. Cucubalus.
trachiatus, Fourr. l. c. = S. Cucubalus.
vesicarius, Fourr. l. c. = S. Cucubalus.
vulgaris, Moench, Meth. 709 = S. Cucubalus.

BEHENANTHA, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 130 = **Silene**, Linn. (Caryoph.).
Cserei, Schur, l. c. 132 = S. Cucubalus.
glaucescens, Schur, l. c. = S. Cucubalus.
inflata, Schur, l. c. 131 = S. Cucubalus.
saponariaefolia, Schur, l. c. 133 = S. Cucubalus.

BEHNIA, F. Didrichs. in Kjoeb. Vidensk. Meddel. (1854) 182. *LILIACEAE*, Benth. & Hook. f. iii. 766.
DICTYOPSIS, Harv. in Bot. Mag. t. 5638 (1867).
HYLONOME, Baker, in Journ. Linn. Soc. xiv. (1875) 561.
reticulata, *F. Didrichs. l. c.* 183.—Afr. austr.

BEHRINIA, Sieber, ex Steud. Nom. ed. I. 105 (1821) = BERINIA, Brign. = **Crepis**, Linn. (Compos.).

BEHURIA, Cham. in Linnaea, ix. (1834) 376. *MELASTOMACEAE*, Benth. & Hook. f. i. 748.
insignis, *Cham. l. c.* 377.—Bras.

BEILIA, Eckl. Verz. 43 (1827) = **Watsonia**, Mill. (Irid.).
spicata, Eckl. l. c. = W. spicata.
triticea, Eckl. l. c. = W. plantaginea.

BEILSCHMIEDIA, Nees, in Wall. Pl. As. Rar. ii. 69 (1831). *LAURINEAE*, Benth. & Hook. f. iii. 152.
BIELSCHMEIDIA, Panch. & Sebert, Not. Bois Nouv. Caléd. 176 (1874).
HUFELANDIA, Nees, Prod. Laurin. ii. (1833); et Laurin. Expos. (1836) 187, 674.
NESODAPHNE, Hook. f. Fl. N. Zel. i. 217 (1853).
assamica, *Meissn. in DC. Prod.* xv. 1. 64.—Assam; Burma.
Baillonii, *Panch. & Sebert, Not. Bois Nouv. Caled.* 176.—N. Caled.
caloneura, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 47.—N. Guin.
chinensis, *Hance, in Journ. Bot.* xx. (1882) 79.—China.
fagifolia, *Nees, in Wall. Pl. As. Rar.* ii. 69.—Ind. or.
globularia, *Kurz, For. Fl. Brit. Burma*, ii. 294.—Burma.
javanica, *Miq. Pl. Jungh.* i. 176.—Java.
lanceolata, *Panch. & Sebert, Not. Bois Nouv. Caled.* 177.—Caled.
lancifolia, *Miq. Pl. Jungh.* 177.—Java.
macrophylla, *Meissn. in DC. Prod.* xv. 1. 63.—Burm.
Madang, *Blume, Mus. Bot. Lugd. Bat.* i. 332.—Java.
oppositifolia, *Benth. & Hook. f. Gen.* iii. 152.—Zeylan.
pendula, *Hemsl. Biol. Centr. Am. Bot.* iii. 70.—Mexic.
Roxburghiana, Dalz. & Gibs. Bomb. Fl. 222 = fagifolia.
Roxburghiana, *Nees, in Wall. Pl. As. Rar.* ii. 61.—Assam; Burma.
Thomaea, *Benth. & Hook. f. Gen.* iii. 152.—Ind. occ.
undulata, *Miq. Fl. Ind. Bat.* i. 1. 920.—Java.
Wightii, *Benth. & Hook. f. Gen.* iii. 152.—Ind. or.

BEJARIA, Zea, in Anal. Cienc. Nat. iii. (1801) 151, in nota; Vent. Jard. Cels, sub t. 51 = **Befaria**, Mutis (Ericac.).

BEJUCO, Loefl. Itin. 314 (1758) = **Hippocratea**, Linn.
pendulus, Loefl. l. c. = H. scandens.

BELAIRIA, A. Rich. Ess. Fl. Cub. 511. t. 40 (1845). *LEGUMINOSAE*, Benth. & Hook. f. i. 558.
mucronata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 179.—Cuba.
spinosa, *A. Rich. Ess. Fl. Cub.* 513.—Cuba.
ternata, *Wright, ex Griseb. Cat. Pl. Cub.* 81.—Cuba.

BELAMCANDA, Adans. Fam. ii. 60 (1763) (Belam-Canda). *IRIDIAE*, Benth. & Hook. f. iii. 697.
PARDANTHUS, Ker-Gawl. in Kon. & Sims, Ann. Bot. i. (1805) 246.
aristata, Moench, Meth. Suppl. 214 = Sparaxis grandiflora.
bulbifera, Moench, l. c. = Sparaxis bulbifera.
chinensis, [DC. in] Red. Lil. iii. t. 121 = punctata.
fenestrata, Moench, Meth. 529 = Tritonia crocata.
punctata, *Moench, l. c.*—China.
semiflexuosa, Moench, Meth. Suppl. 214 = Sparaxis grandiflora.
tripedalis, *Fisch. ex Roem. & Schult. Syst.* i. *Mant.* 306 (sp. dub.).
villosa, Moench, Meth. Suppl. 214 = Babiana villosa.

BELANGERA, Cambess. Cunon. Syn. Bras. Mer. 3 (1829). *SAXIFRAGEAE*, Benth. & Hook. f. i. 650.
LAMANONIA, Vell. Fl. Flum. v. t. 104 (1827).
POLYSTEMON, D. Don, in Edinb. N. Phil. Journ. ix. 95 (1830).
cuneata, *Cambess. l. c.*—Bras.
denticulata, *Moric. Pl. Nouv. Am.* 153. t. 90.—Bras.
glabra, *Cambess. Cunon. Syn. Bras. Mer.* (1829) 3.—Bras.
intermedia, *Mart. in Flora*, xx. (1837) II. *Beibl.* 95.—Bras.
Lamanonia, *Steud. Nom.* ed. II. i. 195.—Bras.
Riedeliana, Casar. ex Eichl. in Mart. Fl. Bras. xiv. II. 155 = speciosa.
speciosa, *Cambess. Cunon. Syn. Bras. Mer.* (1829) 4.—Bras.
tomentosa, *Cambess. l. c.*—Bras.

BELANTHERIA, Nees, in DC. Prod. xi. 96 (1847) = **Brillantaisia**, Beauv. (Gramin.).
Belvisiana, Nees, l. c. 97 = Brillantaisia owariensis.

BELENCITA, Karst. in C. Koch, Allg. Gartenz. (1857) 25. *CAPPARIDEAE.*
Hagenii, *Karst. l. c.* 24.—N. Granat.

BELENIA, Decne. in Jacquem. Voy. Bot. 113. t. 120 (1844) = **Physochlaina**, D. Don (Solanac.).
praealta, Decne. l. c. 114. t. 120 = P. praealta.

BELENIDIUM, Arn. ex DC. Prod. vii. 292, in syn. (1838) = **Hymenatherum**, Cass. (Compos.).
Candolleanum, Arn. ex DC. l. c. = H. Belenidium.

BELEROPONE, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 524 (1885), sphalm. = **Beloperone**, Nees (Acanth.).

BELHARNOSIA, Adans. Fam. ii. 432 (1763) = **Sanguinaria**, Linn. (Papav.).

BELIA, Steller, ex S. G. Gmel. Fl. Sibir. iv. 88 (1769) = **Claytonia**, Gron. (Portulac.).

BELILLA, Adans. Fam. ii. 159 (1763) = **Mussaenda**, Burm. (Rubiac.).

BELIS, Salisb. in Trans. Linn. Soc. viii. (1807) 315, nomen prius = **Cunninghamia**, R. Br. (1826) (Conifer.).
jaculifolia, Salisb. l. c. = C. sinensis.
lanceolata, Sweet, Hort. Brit. ed. II. 475 = C. sinensis.

BELLADONA, Adans. Fam. ii. 219 (1763) = **Atropa**, Linn. (Solanac.).

BELLADONNA, Sweet, Hort. Brit. ed. II. 506 (1830) = **Amaryllis**, Linn.
blanda, pallida, pudica, purpurascens, Sweet, l. c. = A. Belladonna.

BELLADONNA, [Tourn.] Rupp. Fl. Jcn. ed. Hall. 252 (1745) = **Atropa**, Linn. (Solanac.).
baccifera, Lam. Fl. Fr. ii. 255 = A. Belladonna.
peruviana, Hort. Par. ex Adans. Fam. ii. 219 = Nolana prostrata.
trichotoma, Scop. Fl. Carn. ed. II. i. 160 = A. Belladonna.

BELLARDIA, All. Fl. Pedem. i. 61 (1785) = **Bartsia**, Linn. (Scrophul.).
Trixago, All. l. c. = Bartsia Trixago.
viscosa, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 4 = Bartsia viscosa.

BELLARDIA, Colla, in Mem. Acc. Torin. xxxviii. (1835) 40. t. 34 = **Microseris**, D. Don (Compos.).
pusilla, Colla, l. c. 42. t. 34.—Chili.

BELLARDIA, Schreb. Gen. i. 790 (1789) = **Coccocipselum**, R. Br. (Rubiac.).
mollis, Willd. ex DC. Prod. iv. 397 = C. canescens.
repens, Willd. Sp. Pl. i. 626 = C. Tontanea.
Tontanea, Roem. & Schult. Syst. iii. 305 = C. Tontanea.

BELLENDENA, R. Br. in Trans. Linn. Soc. x. (1810) 166. *PROTEACEAE*, Benth. & Hook. f. iii. 175.
montana, *R. Br. l. c.* 48, 166.—Tasmania.

BELLENDENIA, Rafin. ex Endl. Gen. 169 (1837), in syn. = **Tritonia**, Ker-Gawl. (Irid.).

BELLENDENIA, Endl. Gen. 340 (1838) = **Bellendena**, R. Br. (Protcac.).

BELLEVALIA, Delile, ex Endl. Gen. 231 (1836) = **Althenia**, Petit (Naiad.).
australis, Delile, ex Parl. Fl. Ital. iii. 648 = A. filiformis.

BELLEVALIA, Lapeyr. in Journ. Phys. lxvii. (1808) 425 = **Hyacinthus**, Tourn. (Liliac.).
abortiva, Gren. in Gren. & Godr. Fl. Fr. iii. 217 = H. trifoliatus.
aleppica, Boiss. Diagn. Ser. II. iv. 111 = H. nervosus.
appendiculata, Lapeyr. Hist. Abr. Pl. Pyr. 186 = H. romanus.
atroviolacea, Regel, in Act. Hort. Petrop. viii. (1884) 654.—As. centr.
azurea, Boiss. Fl. Orient. v. 307 = H. ciliatus.
Battandieri, Freyn, in Flora, lxviii. (1885) 25.—Afr. bor.
Boissieri, Freyn, l. c. 95.—Graecia.
Bouviana, Heldr. ex Nym. Consp. 733.—Graecia.
calandriniana, Nym. Consp. 732 = Muscari comosum.
caucasica, Griseb. Spicil. Fl. Rumel. ii. 387 = Muscari caucasicum.
ciliata, T. Nees, Gen. 4. t. 8; Kunth, Enum. Pl. iv. 308 = H. ciliatus.
Clusiana, Griseb. Spicil. Fl. Rumel. ii. 387 = H. dubius.
comosa, Kunth, Enum. Pl. iv. 306 = Muscari comosum.
Cupaniana, Nym. Consp. 733 = Muscari Cupanianum.
curta, Nym. l. c. = Muscari Holzmanni.
cyanoleuca, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = H. romanus.
densiflora, Boiss. Diagn. Ser. I. vii. 109 = H. densiflorus.
dubia, Reichb. Fl. Germ. Excurs. 105 = H. dubius.
fallax, Pomel, Nouv. Mat. Fl. Atl. 254.—Algeria.
flexuosa, Boiss. Diagn. Ser. I. xiii. 36 = H. flexuosus.
Fontanesii, Nym. Consp. 733.—Creta.
glauca, Kunth, Enum. Pl. iv. 309 = H. ciliatus.
graeca, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876) 228 = Muscari graecum.
graminifolia, Nym. Consp. 733.—Ins. Graec.
Gussoniana, Griseb. Spicil. Fl. Rumel. ii. 387.—Sicil.
Hackeli, Freyn, in Oestr. Bot. Zeitschr. xxvii. (1877) 289.—Europ.
Haynei, Boiss. Fl. Orient. v. 308 = H. Haynei.
Heldreichii, Boiss. Diagn. Ser. II. iv. 111 = H. lineatus.
hispida, J. Gay, in Bull. Soc. Bot. Fr. iii. (1856) 240 = H. hispidus.
Holzmanni, Heldr. in Atti Congr. Bot. Firenze, 1874 (1876). 228 = Muscari Holzmanni.
leucophaea, Boiss. Fl. Orient. v. 305 = H. leucophaeus.
lineata, Kunth, Enum. Pl. iv. 309 = H. lineatus.

BELLEVALIA:—
macrobotrys, Boiss. Diagn. Ser. I. xiii. 35 = Hyacinthus macrobotrys.
maritima, Kunth, Enum. Pl. iv. 307 = Muscari maritimum.
mauritanica, Pomel, Nouv. Mat. Fl. Atl. 255.—Algeria.
micrantha, Boiss. Diagn. Ser. I. v. 63 = H. micranthus.
monophylla, J. Gay, ex Boiss. Fl. Orient. v. 294 = Muscari latifolium.
montana, Boiss. l. c. 304 = Muscari montanum.
muscarioides, Mast. in Journ. Linn. Soc. iii. (1859) 113 = Muscari latifolium.
nervosa, Boiss. Fl. Orient. v. 306 = H. nervosus.
nivalis, Boiss. & Kotschy, Diagn. Ser. II. iv. 110 = H. nivalis.
operculata, Lapeyr. in Journ. Phys. lxvii. (1808) 426, cum. tab. = H. romanus.
pallens, Vis. in Mem. Ist. Venet. xvi. (1871) 67 = H. dalmaticus.
paradoxa, Boiss. Fl. Orient. v. 308 = H. paradoxus.
persica, Boiss. l. c. = H. persicus.
pharmacusana, Nym. Consp. 733 = Muscari Sartorianum.
Pinardi, Boiss. Diagn. Ser. I. v. 62 = Muscari Pinardi.
pseudo-muscari, Boiss. & Buhse, Diagn. Ser. II. iv. 110 = H. paradoxus.
romana, Sweet, Hort. Brit. ed. I. 419; Reichb. Fl. Germ. Excurs. 105 = H. romanus.
Sartoriana, Nym. Consp. 733 = Muscari Sartorianum.
sessiliflora, Kunth, Enum. Pl. iv. 309 = H. sessiliflorus.
spicata, Boiss. Diagn. Ser. I. vii. 110 = H. spicatus.
syriaca, Herb. in Lindl. Bot. Reg. (1844) Misc. 88 = H. trifoliatus.
tenuiflora, Nym. Consp. 732 = Muscari caucasicum.
theraea, Nym. l. c. 733 = Muscari thereum.
trifoliata, Kunth, Enum. Pl. iv. 308 = H. trifoliatus.
turkestanica, Franch. in Ann. Sc. Nat. Sér. VI. xviii. (1884) 256.—Turkest.
variabilis, Freyn, in Flora, lxviii. (1885) 29.—Alger.
Webbiana, Parl. Nouv. Gen. e Spec. 19 = Hyacinthus dubius.
Weissii, Freyn, in Oestr. Bot. Zeitschr. xxviii. (1878) 87 = Muscari Weissii.

BELLIDASTRUM, Mich. ex Hall. Enum. Stirp. Helv. ii. 723 (1742) = **Aster**, Tourn. (Compos.).
Michelii, Plan. Ensayo Fl. Gall. 259 = Bellis sylvestris.
Michelii, Cass. in Dict. Sc. Nat. iv. 70 = A. Bellidiastrum.
montanum, Hoppe, ex DC. Prod. v. 226 = A. Bellidiastrum.
subalpinum, Schrank, Fl. Mon. iii. 251 = A. Bellidiastrum.

BELLIDIASTER, Dum. Fl. Belg. 66 (1827) = **Aster**, Tourn. (Compos.).
montanus, Dum. l. c. = A. Bellidiastrum.

BELLIDIASTRUM, [Vaill.] Less. Syn. Comp. 383 (1832) = **Osmites**, Linn. (Compos.).
osmitoides, Less. l. c. 384 = O. Bellidiastrum.

BELLIDIASTRUM, Cass. in Bull. Soc. Philom. (1816) 199 = BELLIDASTRUM, Mich. = **Aster**, Tourn. (Compos.).

BELLIDISTRUM, Reichb. Consp. 109 (1828) = praec.

BELLIDIOPSIS, Spach, Hist. Vég. Phan. x. 25 (1841) = **Osmites**, Linn. (Compos.)

BELLIDIUM, Bertol. Fl. Ital. ix. 316 (1853) = **Bellis**, Tourn. (Compos.).
pappulosum, Bertol. l. c. = Bellis sylvestris.
rotundifolium, Bertol. l. c. = Bellis rotundifolia.

BELLILLA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 82 = BELILLA, Rheede = **Mussaenda**, Burm. (Rubiac.).

BELLINIA, Roem. & Schult. Syst. iv. 56 (1819) = **Saracha**, Ruiz & Pav. (Solanac.).
biflora, Roem. & Schult. l. c. 689 = S. procumbens.
contorta, Roem. & Schult. l. c. = S. contorta.
dentata, Roem. & Schult. l. c. = S. dentata.
micrantha, Roem. & Schult. l. c. 690 = S. micrantha.

BELLINIA:—
peruviana, Steud. Nom. ed. I. 105 = *Saracha peruviana* (nomen delendum).
procumbens, Roem. & Schult. Syst. iv. 687 = S. procumbens.
pubescens, Roem. & Schult. l. c. 689 = S. pubescens.
punctata, Roem. & Schult. l. c. 688 = S. punctata.
umbellata, Roem. & Schult. l. c. = S. umbellata.

BELLIOPSIS, Pomel, Mat. Fl. Atl. 7 (1860) = **Bellium**, Linn. (Compos.).
cordifolia, Pomel, Nouv. Mat. Fl. Atl. 286.—Algeria.
rotundifolia, Pomel, Mat. Fl. Atl. 7 = Bellium rotundifolium.

BELLIS, [Tourn.] Linn. Gen. ed. I. 256 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 265.
ASTRANTHIUM, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 312.
KYBERIA, Neck. Elem. i. 42 (1790).
SEUBERTIA, H. Wats. in Hook. Lond. Journ. Bot. iii. (1840) 602.
aculeata, Labill. Nov. Holl. Pl. ii. 55. t. 206 = Brachycome Billardieri.
alpina, Hegetschw. Fl. Schw. 821 = perennis.
americana, Mirb. ex Steud. Nom. ed. II. i. 195 = integrifolia.
annua, *Linn. Sp. Pl.* 887.—Reg. Mediterr.
arabica, *Raeusch. Nom.* ed. III. 246, nomen.—Arabia.
armena, *Boiss. Fl. Orient.* iii. 174.—Armenia.
atlantica, *Boiss. & Reut. Pugill.* 54.—Hispan.
azorica, *Hochst. in Seub. Fl. Azor.* 31. t. 9. fig. 1-3.—Ins. Azores.
Bernardi, *Boiss. & Reut. Pugill.* 56.—Hispan.
caerulescens, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xxii. (1875) 60, nomen..—Marocco.
campestris, *Vell. Fl. Flum.* viii. t. 124.—Bras.
ciliaris, Labill. Nov. Holl. Pl. ii. 56. t. 209 = Brachycome ciliaris.
ciliata, *Rafin. New Fl. Am.* ii. 24.—Am. bor.
cordifolia, *Willk. Illustr. Fl. Hisp.* 47.—Hispan.
dentata, DC. Prod. v. 304 = annua.
droseraefolia, Gouan, Illustr. 69 = Bellium bellidioides.
dubia, *Spreng. Syst.* iii. 582.—Hab.?
geifolia, Banks & Soland. ex Hook. Fl. N. Zel. i. 125 = Lagenophora Forsteri.
glandulosa, Klatt, in Leopoldina, xix. (1883) 32 = Brachycome glandulosa.
graminea, Labill. Nov. Holl. Pl. ii. 54. t. 204 = Brachycome graminea.
hirta, Host, Fl. Austr. ii. 501 = sylvestris?
hortensis, Mill. Gard. Dict. ed. VIII. n. 3 = perennis.
hybrida, Tenore, Fl. Nap. v. 233. t. 194 = perennis.
integrifolia, DC. in Lam. Encyc. v. 7 = perennis.
integrifolia, *Michx. Fl. Bor. Am.* ii. 131.—Am. bor.
javanica, Blume, Bijdr. 917 = Myriactis Wightii.
longifolia, *Boiss. & Heldr. Diagn.* Ser. I. xi. 1.—Ins. Creta.
longifolia, Orph. ex Nym. Consp. 390 = sylvestris.
magellanica, DC. in Lam. Encyc. v. 7 = Lagenophora Commersonii.
mexicana, *A. Gray, Pl. Wright.* i. 93.—Mexic.
microcephala, Lange, in Kjoeb. Vidensk. Meddel. (1861) 66 = annua.
mossambicensis, Klatt, in Leopoldina, xix. (1883) 32 = Brachycome mossambicensis.
nutans, *Rafin. New Fl. Am.* ii. 23.—Am. bor.
pappulosa, Boiss. ex DC. Prod. vii. 276 = sylvestris.
parviflora, *Rafin. New Fl. Am.* ii. 23.—Am. bor.
pedunculata, *Vell. Fl. Flum.* viii. t. 125.—Bras.
perennis, *Linn. Sp. Pl.* 886.—Europ. occ.
pilosa, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 126 = Lagenophora lanata.
pilosa, Blume, Bijdr. 917 = Myriactis pilosa.
prostrata, *Pomel, Nouv. Mat. Fl. Atl.* 287.—Algeria.
racemosa, Steud. Nom. ed. I. 105, sphalm. = seq.
ramosa, Jacq. Enum. Pl. Carib. 28 = Eclipta punctata.
ramosa, Lam. Fl. Fr. ii. 122 = annua.
repens, Lam. l. c. = annua.
rotundifolia, *Boiss. & Reut. Pugill.* 55.—Algeria.
scandens, *Vell. Fl. Flum.* viii. t. 126.—Bras.
scaposa, Gilib. Fl. Lituan. i. 213 = perennis.
stipitata, Labill. Nov. Holl. Pl. ii. 55. t. 105 = Lagenophora Billardieri.
sylvestris, *Cyrill. Pl. Rar.* ii. 22. t. 4.—Reg. Mediterr.
velutina, *Pomel, Nouv. Mat. Fl. Atl.* 35.—Algeria.
xanthocomoides, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 118.—Mexic.

BELLIUM, Linn. Mant. 157 (1771). *COMPOSITAE*, Benth. & Hook. f. ii. 265.
BELLIOPSIS, Pomel, Mat. Fl. Atl. 7 (1860).
bellidioides, Desf. Fl. Atlant. ii. 279 = Bellis dentata.
bellidioides, *Linn. Mant.* 285.—Reg. Mediterr.
bellidioides, Urv. in Mém. Soc. Linn. Par. i. (1822) 368 = minutum.
bellioides, G. Don, in Sweet, Flow. Gard. ii. 175 = bellidioides.
cordifolium, Kunze, in Flora, xxix. (1846) 703 = Bellis rotundifolia.
crassifolium, *Moris, Pl. Sard. Elench.* i. 26.—Sardinia.
dentatum, Viviani, Fragm. 8. t. 10 = Bellis annua.
droseraefolium, Labill. ex Steud. Nom. ed. I. 105 = bellidioides.
giganteum, Cass. in Dict. Sc. Nat. xxxvii. 456 = rotundifolium.
minutum, *Linn. Mant.* 286.—Archip. Graec.
nivale, Req. in Ann. Sc. Nat. Sér. I. v. (1825) 383 = bellidioides.
pappulosum, Kunze, in Flora, xxix. (1846) 703 = Bellis sylvestris.
pumilum, Salisb. Prod. 202 = bellidioides.
rotundifolium, *DC. Prod.* v. 303.—Algeria.

BELLOA, Remy, in C. Gay, Fl. Chil. iii. 336. t. 38 (1849) = **Gnaphalium**, Linn. (Compos.).
chilensis, Remy, l. c. 337. t. 38. f. 2.—Chili.
subspicata, Wedd. Chlor. And. i. 159.—Bolivia.

BELLONIA, [Plum.] Linn. Gen. ed. I. 50 (1737). *GESNERACEAE*, Benth. & Hook. f. ii. 996.
BELONIA, Adans. Fam. ii. 158 (1763).
aspera, *Linn. Sp. Pl.* 172.—Ind. occ.
spinosa, *Sw. Prod. Veg. Ind. Occ.* 41.—Ins. S. Doming.

BELLOTA, C. Gay, Fl. Chil. v. 297. t. 59 (1849) = **Cryptocarya**, R. Br. (Laurin.).
Miersii, C. Gay, l. c. 298.—Chili.
nitida, Phil. in Linnaea, xxix. (1857–58) 39 = C. nitida.
pauciflora, Phil. in Anal. Univ. Chil. (1865) i. 649.—Chili.

BELLOTA, A. Rich. ex Phil. in Linnaea, xxix. (1857–58) 40 = **Boldu**, Nees (Laurin.).

BELLUCCIA, Adans. Fam. ii. 344 (1763) = **Ptelea**, Linn. (Rutac.).

BELLUCIA, Meissn. Gen. Comm. 46, in syn. = praec.

BELLUCIA, Neck. Elem. ii. 142 (1790). *MELASTOMACEAE*, Benth. & Hook. f. i. 768.
APATITIA, Desv. in Ham. Prod. Pl. Ind. Occ. 42 (1825).
AXINANTHERA, Karst. in Linnaea, xxx. (1859–60) 157.
ISCHYRANTHERA, Steud. ex Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 102.
WEEERA, J. F. Gmel. Syst. 820 (1790).
arborescens, Baill. Hist. Pl. vii. 34 = Loreya arborescens.
Aubletii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 102.—Guiana.
Aubletii, Seem. Bot. Voy. Herald, 122 = superba.
Axinanthera, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 142.—N. Granat.
brasiliensis, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 104.—Bras.
Cacatin, *Sagot, l. c.* Sér. VI. xv. (1883) 332.—Guiana.
grossularioides, Triana, in Trans. Linn. Soc. xxviii. (1871) 141 = Aubletii.
Hostmannii, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 103.—Guiana.
macrophylla, Triana, in Trans. Linn. Soc. xxviii. (1871) 142 = superba.
multiflora, Karst. in Linnaea, xxx. (1859–60) 158 = Aubletii.
nervosa, Rafin. Sylva Tellur. 93 = Blakea trinervia.
pentamera, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 105.—Peruv.
pluricaulis, Spruce, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 142 = Hostmannii.
riparia, Spruce, l. c. = Hostmannii.
superba, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 104.—Mexic.

BELLYNKXIA, Muell. Arg. in Flora, lviii. (1875) 465 = **Appunia**, Hook. f. (Rubiac.).
angulata, Muell. Arg. l. c. = A. angulata.

BELMONTIA, E. Mey. Comm. Pl. Afr. Austr. 183 (1837). *GENTIANEAE*, Benth. & Hook. f. ii. 804.
EXOCHAENIUM, Griseb. in DC. Prod. ix. 55 (1845).
PARASIA, Rafin. Fl. Tellur. iii. 78 (1836).
cordata, *E. Mey. l. c.*—Afr. austr.
gracilis, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 47.—Afr. trop.
grandis, *E. Mey. Comm. Pl. Afr. Austr.* 183.—Afr. austr.
Ohlendorffii, *Griseb. in DC. Prod.* ix. 54.—Afr. austr.
spathulata, *E. Mey. Comm. Pl. Afr. Austr.* 183.—Afr. austr.

BELOAKON, Rafin. Fl. Tellur. iii. 87 (1836) = **Phlomis**, Linn. (Labiat.).
luteum, Rafin. l. c. = P. fruticosa.
tomentosum, *Rafin. l. c.* = P. Samia.

BELOANTHERA, Hassk. in Flora, xxv. (1842) Beibl. ii. 55. '*GENUS DUBIUM.*'
oppositifolia, *Hassk. l. c.*—Java.

BELOËRE, Shuttlew. ex A. Gray, Pl. Wright. i. 21 (1852) = **Abutilon**, Gaertn. (Malvac.).
cistiflora, Shuttlew. ex A. Gray, l. c. = A. indicum.
crispa, Shuttlew. ex A. Gray, l. c. in nota = A. crispum.

BELONIA, Adans. Fam. ii. 158 (1763) = **Bellonia**, Plum. (Gesner.).

BELONITES, E. Mey. Comm. Pl. Afr. Austr. 187 (1837) = **Pachypodium**, Lindl. (Asclep.).
bispinosa, E. Mey. l. c. 188 = P. bispinosum.
succulenta, E. Mey. l. c. 187 = P. succulentum.

BELONOPHORA, Hook. f. Ic. Pl. t. 1127 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 109.
coffeoides, *Hook. f. l. c.*—Afr. trop.

BELOPERONE, Nees, in Wall. Pl. As. Rar. iii. 76, 102 (1832). *ACANTHACEAE*, Benth. & Hook. f. 1110.
BELOPERONIDES, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 162.
KUESTERA, Regel, Gartenfl. vi. (1857) 345.
SIMONISIA, Nees, in Mart. Fl. Bras. ix. 144. t. 23 (1847).
Amherstiae, Nees, l. c. 102 = Dianthera nodosa.
appendiculata, *Nees, in DC. Prod.* xi. 423.—Peruv.
atropurpurea, *Nees, in Mart. Fl. Bras.* ix. 138.—Bras.
aurantiaca, *Benth. & Hook. f. Gen.* ii. 1111.—Am. merid.?
auriculata, *Nees, in Mart. Fl. Bras.* ix. 138.—Bras.
bullata, *Nees, l. c.* 141.—Bras.
californica, *Benth. Bot. Voy. Sulph.* 38.—Am. bor. occ.
calycina, Nees, in Hook. Lond. Journ. Bot. iv. (1845) 637 = Dianthera calycina.
caracasana, *Fenzl, in Jacq. Eclog.* ii. 2, in obs.—Venezuela.
carthaginensis, Benth. & Hook. f. Gen. ii. 1111 = violacea.
ciliata, Hook. Bot. Mag. t. 5888 = Dianthera ciliata.
ciliata, *Nees, in DC. Prod.* xi. 422.—Peruv.
comosa, *Nees, l. c.* 416.—Mexic.
cordata, Mart. ex Nees, l. c. 361 (= *Sericographis cordata*).—Bras.
cristata, Mart. ex Nees, l. c. 419 = Dianthera nodosa.
denudata, *Nees, l. c.* 423.—Peruv.
dicliptcroides, *Nees, in Mart. Fl. Bras.* ix. 136.—Bras.
Eustachyana, *Benth. & Hook. f. Gen.* ii. 1111.—Ind. occ.
fimbriata, *Nees, in DC. Prod.* xi. 416.—Mexic.
formosa, *Fenzl, in Jacq. Eclog.* ii. *in Ind.* (*Justicia*, t. 102).—Venezuela.
fragilis, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 140.—Bras.
fulgida, Hassk. Cat. Hort. Bog. Alt. 150 = Clinacanthus Burmanni.
glandulosa, *Nees, in DC. Prod.* xi. 414.—Bras.

BELOPERONE:—
glomerata, *Nees, in Mart. Fl. Bras.* ix. 137.—Bras.
hirsuta, *Nees, l. c.* 143.—Bras.
involucrata, *Nees, l. c.*—Bras.
Lamarckiana, *Nees, in DC. Prod.* xi. 415.—Ind. occ.
lanceolata, *Mart. ex Nees, in Mart. Fl. Bras.* ix. 141.—Bras.
longipetiolata, *Nees, l. c.* 137.—Bras.
macrantha, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 516.—Mexic.
Mathewsiana, *Nees, in DC. Prod.* xi. 731.—Peruv.
Meyeniana, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 381.—Bras.
microstachya, *Nees, in Mart. Fl. Bras.* ix. 137.—Bras.
mollis, *Nees, l. c.* 142.—Bras.
monticola, *Nees, l. c.*—Bras.
nemorosa, Nees, in DC. Prod. xi. 415 = Adhatoda tetramera.
nodicaulis, *Nees, in Mart. Fl. Bras.* ix. 140.—Bras.
oblongata, *Lindl. Bot. Reg.* t. 1657.—Bras.
oblongata, Nees, in Wall. Pl. As. Rar. iii. 102 = plumbaginifolia.
obtusifolia, Nees, l. c. 339 = ovata.
plumbaginifolia, *Nees, in Wall. Pl. As. Rar.* iii. 102.—Bras.
portoricensis, *Nees, in DC. Prod.* xi. 414.—Porto Rico.
pulchella, Linden, Cat. n. 17 (1862) 2; ex Regel, Gartenfl. xiv. (1865) 289 = violacea.
Riedeliana, *Benth. & Hook. f. Gen.* ii. 1111.—Bras.
Rohrii, *Nees, in DC. Prod.* xi. 415.—Am. austr.
rosea, *Benth. & Hook. f. Gen.* ii. 1111.—Bras.
Schomburgkiana, *Nees, in Hook. Lond. Journ. Bot.* iv. (1845) 636.—Guiana.
scorpioides, *Nees, in DC. Prod.* xi. 422.—Bras.
Selloviana, *Nees, in Mart. Fl. Bras.* ix. 136.—Bras.
spathulata, *Nees, l. c.*—Bras.
sphaerosperma, *Benth. & Hook. f. Gen.* ii. 1111.—Ind. occ.
strobilacea, Mart. ex Nees, in DC. Prod. xi. 358 (= *Amphiscopia strobilacea*).—Bras.
surinamensis, *Miq. Stirp. Surin. Sel.* 130.—Guiana.
trifoliata, *Nees, in Mart. Fl. Bras.* ix. 141.—Bras.
villosa, Mart. ex Nees, in DC. Prod. xi. 417 = hirsuta.
violacea, *Planch. & Linden, Hort. Linden.* i. 5. t. 3.—Mexic.

BELOPERONIDES, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 162 = **Beloperone**, Nees (Acanth.).
macrantha, Oerst. l. c. = Beloperone macrantha.

BELOSPIS, Rafin. Fl. Tellur. iii. 90 (1836) = **Salvia**, Tourn. (Labiat.).
levigata, Rafin. l. c. 91 = S. involucrata.

BELOSTEMMA, Wall. ex Wight, Contrib. Bot. Ind. Or. 52 (1834) = **Tylophora**, R. Br. (Asclepiad.).
hirsutum, Wall. ex Wight, l. c. = T. Belostemma.

BELOSYNAPSIS, Hassk. in Flora, liv. (1871) 259 = **Cyanotis**, D. Don (Commelin.).
kewensis, Hassk. l. c. = C. kewensis.

BELOTIA, A. Rich. Ess. Fl. Cub. 207. t. 21 (1845). *TILIACEAE*, Benth. & Hook. f. i. 233.
ADENODISCUS, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) ii. 504.
Galleottii, *Turcz. l. c.*—Mexic.
greviaefolia, *A. Rich. Ess. Fl. Cub.* 209. t. 22.—Cuba.
insignis, *Baill. Adansonia*, x. (1872) 182.—Mexic.

BELOU, Adans. Fam. ii. 408 (1763) = **Aegle**, Correa (Rutac.).

BELOVIA, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1852) 462 = **Suaeda**, Forsk. (Chenopod.).
baccifera, Moq. in DC. Prod. xiii. ii. 168 = S. setigera.
paradoxa, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1852) 462 = S. setigera.

BELTOKON, Rafin. Fl. Tellur. iii. 86 (1836) = **Origanum**, Tourn. (Labiat.).
Tourneforti, Rafin. l. c. = O. Tournefortii.

BELUTTA, Rafin. Fl. Tellur. iii. 39 (1836) = **Allmania**, R. Br. (Amarant.).
peduncularis, Rafin. l. c. = A. nodiflora.
sessilis, Rafin. l. c. = A. albida, pyramidalis.

BELVALA, Adans. Fam. ii. 285 (1763) = **Struthiola**, Linn. (Thymel.).
spicata, Rafin. Fl. Tellur. iv. 105 = S. virgata?

BELVALIA, Delile, sec. Flora, xiii. (1830) ii. 455 = **Althenia**, Petit (Naiad.).
australis, Delile, l. c. = A. filiformis.

BELVEDERE, Clayt. ex Gronov. Fl. Virgin. 25 (1739) = **Galax**, Linn. (Diapens.).

BELVISIA, Desv. Journ. Bot. iv. (1814) 130 = **Napoleona**, Beauv. (Myrtac.).
caerulea, Desv. l. c. = N. imperialis.

BEMBECODIUM, Lindl. Veg. Kingd. 712 (1847) = BEMBYCODIUM, Kunze = **Athanasia**, Linn. (Compos.).

BEMBICIA, Oliver, in Hook. Ic. Pl. t. 1404. *SAMYDACEAE.*
axillaris, *Oliver, l. c.*—Malaya.

BEMBICIUM, Mart. ex Baker, in Mart. Fl. Bras. vi. ii. 341 (1876) = **Eupatorium**, Tourn. (Compos.).
gracile, Mart. ex Baker, l. c. 343 = E. kleinioides.

BEMBIX, Lour. Fl. Cochinch. 346 (1790). *MALPIGHIACEAE*?—Cf. Benth. & Hook. f. Gen. i. 251.
tectoria, *Lour. l. c.*—Cochineh.

BEMBYCODIUM, Kunze, in Linnaea, xvi. (1842) 316 = **Athanasia**, Linn. (Compos.).
Athanasiae, Kunze, l. c.—Afr. austr.

BEMSETIA, Rafin. Sylva Tellur. 12 (1838) = **Ixora**, Linn. (Rubiac.).
paniculata, Rafin. l. c. = I. stricta.

BENAUREA, Rafin. Fl. Tellur. ii. 77 (1836) = **Musschia**, Dum. (Campan.).
sempervirens, Rafin. l. c. 78 = M. aurea.

BENCOMIA, Webb & Berth. Phyt. Canar. ii. 10, t. 39 (1846). *ROSACEAE*, Benth. & Hook. f. i. 624.
caudata, *Webb & Berth. l. c.* 11.—Ins. Canar.
Moquiniana, *Webb & Berth. l. c.*—Ins. Canar.

BENEDICTA, Bernh. Syst. Verz. Erf. 108 (1800) = **Carbenia**, Adans. (Compos.).
officinalis, Bernh. l. c. 159 = C. benedicta.

BENINCASA, Savi, in Bibl. Ital. ix. 158 (1818); (Mem. sopra Piant. Cucurb. i. (1818) 6). *CUCURBITACEAE*, Benth. & Hook. f. i. 824.
cerifera, *Savi, ll. cc.*—As. et Afr. trop. (cult).
cylindrica, Hort. ex Ser. in DC. Prod. iii. 303 = cerifera.
hispida, Cogn. in DC. Monog. Phan. iii. 513 = cerifera.

BENITZIA, Karst. in Linnaea, xxviii. (1856) 420 = **Gymnosiphon**, Blume (Burman.).
Poeppigiana, Karst. l. c. 421.—Bras.
suaveolens, Karst. l. c. 420.—Venezuela.

BENJAMINA, Vell. Fl. Flum. 93 (1825); ii. t. 139 = **Dictyoloma**, DC. (Simarub.).
alata, Vell. ll. cc. = D. incanescens.

BENJAMINIA, Mart. ex Benj. in Mart. Fl. Bras. x. 255 (1847) = QUINQUELOBUS, Benj. (= **Utricularia**?).
glabra, Benj. in Linnaea, xx. (1847) 761 = Limnophila Griffithii.
minor, Benj. l. c. (= *Q. minor*).—As. trop.
splendens, Benj. l. c. (= *Q. splendens*).—As. trop.
utricularieaeformis, Mart. ex Benj. in Mart. Fl. Bras. x. 256 (= *Q. utricularioides*).—Bras.

BENKARA, Adans. Fam. ii. 85 (1763) = **Melastoma**, Burm.
galia, Rafin. Sylva Tellur. 98.—Hab.?

BENNETIA, DC. Prod. vi. 531, in syn. (1837) = BENNETTIA, S. F. Gray = **Saussurea**, DC. (Compos.).

BENNETIA, Rafin. in Ser. Bull. Bot. i. (1830) 220 = **Sporobolus**, R. Br. (Gramin.).
juncea, Rafin. l. c. = S. junceus.

BENNETTIA, R. Br. in Benn. Pl. Jav. Rar. 249. t. 50 (1838) = **Galearia**, Zoll. (Euphorb.).
affinis, R. Br. l. c. 251 = G. affinis.
aristifera, Muell. Arg. in DC. Prod. xv. ii. 1039 = G. aristifera.
filiformis, Muell. Arg. in Linnaea, xxxiv. (1865–66) 204.—Ind. or.
Finlaysoniana, R. Br. in Benn. Pl. Jav. Rar. 251 = G. Finlaysoni.
fulva, Muell. Arg. in Linnaea, xxxiv. (1865–66) 205 = G. fulva.
Jackiana, R. Br. in Benn. Pl. Jav. Rar. 251 = G. Jackiana.
javanica, R. Br. l. c. 249 = G. pedicellata.
Lindleyana, Muell. Arg. in Linnaea, xxxiv. (1865–66) 205.—Ind. or.
pedicellata, R. Br. in Benn. Pl. Jav. Rar. 251 = G. pedicellata.
phlebocarpa, R. Br. l. c. = G. phlebocarpa.
splendens, Muell. Arg. in Linnaea, xxxiv. (1865–66) 205 = G. splendens.
subulata, Muell. Arg. in DC. Prod. xv. ii. 1039.—Ins. Penang.
Wallichii, R. Br. in Benn. Pl. Jav. Rar. 251 = G. Wallichii.

BENNETTIA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 440 (1821) = **Saussurea**, DC. (Compos.).
alpina, S. F. Gray, l. c. = S. alpina.

BENNETTIA, Miq. Fl. Ind. Bat. i. ii. 105 (1859). *BIXINEAE*, Benth. & Hook. f. i. 128.
Horsfieldii, *Miq. l. c.*—Java.

BENTECA, Adans. Fam. ii. 166 (1763) = **Ambelania**, Aubl. (Apocyn.).
odorata, Rafin. Sylva Tellur. 13 = A. acida.

BENTEKA, Adans. Fam. ii. 525 (1763).—Cf. praec. = **Ambelania**, Aubl. (Apocyn.).
Rheedi, Roem. & Schult. Syst. iv. 706 = 'Malabaria.'

BENTHAMIA, Lindl. Bot. Reg. t. 1579 (1833) = **Cornus**, Tourn.
fragifera, Lindl. l. c. = C. capitata.
japonica, Sieb. & Zucc. Fl. Jap. i. 38. t. 16 = C. Kousa.

BENTHAMIA, Lindl. Nat. Syst. ed. I. 241 (1830) = **Amsinckia**, Lehm. (1831) (Boragin.).
lycopsoides, Lindl. ex A. DC. Prod. x. 118 = A. lycopsioides.

BENTHAMIA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 37. t. 7 = **Habenaria**, Willd. (Orchid.).
latifolia, A. Rich. l. c. 38 = H. chlorantha.
spiralis, A. Rich. l. c. 39 = Spiranthes africana.

BENTHAMANTHA, Alef. in Bonplandia, x. (1862) 264 = **Cracca**, Benth. (Legum.).
glabrescens, Alef. l. c. = C. glabrescens.
glandulifera, Alef. l. c. = C. glandulifera.
Grayi, Alef. l. c. = C. sericea.
mollis, Alef. l. c. = C. mollis.
ochroleuca, Alef. l. c. = C. ochroleuca.

BENTHAMIDIA, Spach, Hist. Vég. Phan. viii. 106 (1839) = **Cornus**, Tourn.
florida, Spach, l. c. 107 = C. florida.

BENTHAMIELLA, Speg. in Anal. Soc. Cient. Argent. xv. (1883) 109. *SOLANACEAE.*
patagonica, *Speg. l. c.* 110.—Reg. Argent.

BENTHECA, Neck. Elem. ii. 54 (1790) = BENTECA, Adans.? = **Ambelania**, Aubl. (Apocyn.).

BENTHEKA, Neck. ex A. DC. Prod. viii. 321, in syn. (1844) = **Willughbeia**, Roxb. (Apocyn.).

BENTINCKIA, Berry, ex Roxb. [Hort. Beng. 68 (1814) et] Fl. Ind. iii. 621 (1832). *PALMAE*, Benth. & Hook. f. iii. 916.
KEPPLERIA, Mart. ex Endl. Gen. 251 (1838).
ceramensis, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 8 = Cyrtostachys ceramica.
ceramica, H. Wendl. in Kerch. Palm. 235 = praec.
Condapanna, *Berry, ex Roxb.* [*Hort. Beng.* 68 *et*] *Fl. Ind.* iii. 621.—Ind. or.
nicobarica, *Becc. in Ann. Jard. Buitenz.* ii. (1885) 165.—Ins. Nicob.
Renda, Mart. Hist. Nat. Palm. iii. 316 = Cyrtostachys Renda.

BENZOIN, Hayne, Arzneigew. xi. t. 24 (1829) = **Styrax**, Linn.
officinale, Hayne, l. c. = S. Benzoin.

BENZOIN, Nees, in Wall. Pl. As. Rar. ii. 63 (1831) = **Lindera**, Thunb. (Laurin.).
aestivale, Nees, Syst. Laurin. 495 = L. Benzoin.
citriodorum, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 206, 78 (= *Aperula citriodora*).—Japon.
Diospyrus, Sweet, Hort. Brit. ed. III. 581 = L. melissaefolia.
geniculatum, G. Don, in Sweet, l. c. = L. geniculata.
glaucescens, Nees, ex Meissn. in DC. Prod. xv. i. 193 = Litsea glaucescens.
glaucum, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 205 = L. glauca.
melissaefolium, Nees, Syst. Laurin. 494 = L. melissaefolia.
Neesianum, Nees, in Wall. Pl. As. Rar. ii. 63 = L. Neesiana.
odoriferum, Nees, l. c. = L. Benzoin.
praecox, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 205 = L. praecox.
sericeum, Hook. f. & Thoms. ex Meissn. in DC. Prod. xv. i. 246 = L. heterophylla.
sericeum, Sieb. & Zucc. in Abh. Akad. Muench. iv. iii. (1846) 204 = L. sericea.
Thunbergii, Sieb. & Zucc. l. c. = L. umbellata.
trilobum, Sieb. & Zucc. l. c. 202 = L. triloba.

BENZOINA, Rafin. Sylva Tellur. 133 (1838) = **Styrax**, Linn.
vera, Rafin. l. c. = S. Benzoin.

BENZONIA, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 113 (1827). *RUBIACEAE*? Benth. & Hook. f. ii. 29.
corymbosa, *Schum. l. c.* 114.—Afr. trop.

BERARDIA, Brongn. in Ann. Sc. Nat. Sér. I. viii. (1826) 380. t. 37 = **Nebelia**, Neck. (Bruniac.).
affinis, Brongn. l. c. 381 = N. affinis.
affinis, Sond. in Harv. & Sond. Fl. Cap. ii. 320.—Afr. austr.
angulata, Sond. l. c.—Afr. austr.
aspera, Sond. l. c. 321.—Afr. austr.
Dregeana, Sond. l. c.—Afr. austr.
fragarioides, Schlecht. in Linnaea, vi. (1831) 190.—Afr. austr.
globosa, Sond. in Harv. & Sond. Fl. Cap. ii. 320.—Afr. austr.
laevis, E. Mey. ex Sond. l. c. 319.—Afr. austr.
microphylla, Sond. l. c. 320.—Afr. austr.
paleacea, Brongn. in Ann. Sc. Nat. Sér. I. viii. (1826) 381 = N. paleacea.
phylicoides, Brongn. l. c. = N. phylicoides.
sphaerocephala, Sond. in Harv. & Sond. Fl. Cap. ii. 319.—Afr. austr.

BERARDIA, Vill. Prosp. 27 (1779). *COMPOSITAE*, Benth. & Hook. f. ii. 474.
ARCTIO, Lam. Encyc. i. 235 (1783).
ARCTIUM, Lam. Fl. Fr. ii. 70 (1778).
VILARIA, Guett. Mém. Mineral. Dauph. i. 108. ii. t. 19 (1779).
subacaulis, *Vill. l. c.* 28.—Gallia.

BERBERIDOPSIS, Hook. f. Bot. Mag. t. 5848 (1862). *BERBERIDEAE*, Benth. & Hook. f. i. 964.
corallina, *Hook. f. l. c.*—Chili.

BERBERIS, [Tourn.] Linn. Gen. ed. I. 94 (1737).
BERBERIDEAE, Benth. & Hook. f. i. 43.
MAHONIA, Nutt. Gen. Am. i. 211 (1818).
ODOSTEMON, Rafin. in Am. Monthly Mag. (1817) 192.
abortiva, Renault, Fl. Dép. Orne, 179 = vulgaris.
acanthifolia, Wall. ex Walp. Rep. i. 103 = nepalensis.
acida, Gilib. Exercit. i. 284 = vulgaris.
actinacantha, *Mart. ex Schult. f. Syst.* vii. 12.—Chili.
acutifolia, *Prantl, Ind. Sem. Wirceburg.* (1873) 7.—Hab. ?
aethnensis, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 901 = vulgaris.
aetnensis, *Presl, Fl. Sic.* i. 28.—Cors.; Sardin.; Sicil.
affinis, G. Don, Gen. Syst. i. 115 = aristata.
africana, *Hebenstr. & Ludw. ex Schult. f. Syst.* vii. 11.—Afr. austr.
agapatensis, *Lechl. Berb. Am. Austr.* 44.—Peruv.
altaica, Pall. Fl. Ross. ii. 41. t. 67 = sibirica.
amurensis, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1857) 260.—Manchuria.
Andrieuxii, *Hook. & Arn. in Bot. Beech. Voy.* 318, adnot.—Mexic.
angulosa, *Wall. Cat.* n. 1475.—Ind. or.
angustifolia, *Benth. Pl. Hartw.* 34, 270.—Mexic.
angustifolia, Roxb. Hort. Beng. [87]; Fl. Ind. ii. 183 = B. Lycium.
apyrena, Hort. ex C. Koch, Dendrol. i. 395 = vulgaris.
Aquifolium, *Pursh, Fl. Am. Sept.* i. 219. t. 4.—Am. bor.
arborescens, Hort. ex C. Koch, Dendrol. i. 393 = vulgaris.
aristata, Sims, Bot. Mag. t. 2549 = umbellata.
aristata, *DC. Syst.* ii. 8.—Ind. or.
articulata, *Loisel. in Dict. Sc. Nat.* lxviii. 218.—Gallia.
asiatica, Griff. Ic. Pl. Ind. Or. iv. t. 648 = Wallichiana.
asiatica, *Roxb. ex DC. Syst.* ii. 13.—Reg. Himal.
asperma, Poit. & Turp. in Duham. ed. Nov. iv. 12 = vulgaris.
atrovirens, Wall. ex G. Don, Gen. Syst. i. 117 = Wallichiana.
atroviridis, Steud. Nom. ed. II. i. 196 = Wallichiana.
aurahuacensis, *Lem. in Fl. des Serres*, Sér. I. iv. (1848) 334.—N. Granat.
aurea, Tausch, in Flora, xvii. (1834) 495 = vulgaris.
Bealei, *Fortune, in Gard. Chron.* (1850) 212.—Japon.
bidentata, *Lechl. Berb. Am. Austr.* 11.—Chili.
Bigelovii, Schrad. in Linnaea, xii. (1838) 362 = vulgaris.
boliviana, *Lechl. Berb. Am. Austr.* 21.—Bolivia.
brachybotrys, Edgew. in Trans. Linn. Soc. xx. (1851) 29 = vulgaris.
brachybotrys, *C. Gay, Fl. Chil.* i. 81.—Chili.
brachypoda, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 308.—China.
brevifolia, Hort. ex C. Koch, Dendrol. i. 398 = canadensis.
Bussmul, C. Koch, ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 253 = aristata.
buxifolia, *Lam. Illustr.* t. 253. f. 3; *Poir. Encyc.* viii. 619.—Reg. Magell.
calliobotrys, Bienert, ex Aitch. in Journ. Linn. Soc. xix. (1882) 151 = vulgaris.
canadensis, Guimp. Abbild. Fremd. Holz. 79. t. 63 = Guimpeli.
canadensis, Hayne, ex Steud. Nom. ed. II. i. 196 = spathulata.
canadensis, Mill. Gard. Dict. ed. VIII. n. 2 = vulgaris.
canadensis, *Pursh, Fl. Am. Sept.* 219.—Am. bor.
caraganaefolia, Banks, ex DC. Syst. ii. 18 = Caragana Chamlagu.
carinata, *Lechl. Berb. Am. Austr.* 41.—Peruv.
caroliniana, Sweet, Hort. Brit. ed. III. 19 = canadensis.
cerasina, *Schrad. in Linnaea*, xii. (1838) 379.—Am. bor.
ceratophylla, G. Don, Gen. Syst. i. 115 = aristata.
chilensis, *Gill. in Hook. Bot. Misc.* iii. (1833) 135.—Chili.

BERBERIS:—
chinensis, Hort. ex C. Koch, Dendrol. i. 403 = Guimpeli.
chinensis, Poir. Encyc. viii. 617 = sinensis.
Chitria, D. Don, Prod. Fl. Nep. 204 = aristata.
Chococa, *Schlecht. in Bot. Zeit.* xii. (1854) 652.—Mexic.
ciliaris, *Lindl. in Journ. Hort. Soc.* v. (1850) 5, in adnot.—Bolivia.
coccinea, Hort. ex C. Koch, Dendrol. i. 406 = aristata.
coletioides, *Lechl. Berb. Am. Austr.* 38.—Chili.
oncinna, *Hook. f. Bot. Mag.* t. 4744.—Reg. Himal.
conferta, *H. B. & K. Nov. Gen. et Sp.* v. 69. t. 430.—Ecuador.
congestiflora, *C. Gay, Fl. Chil.* i. 75. t. 3.—Chili.
consanguinea, *Fortune, in Journ. Hort. Soc.* vii. (1852) 226.—China.
coriacea, *St. Hil. Fl. Bras. Mer.* i. 46.—Bras.
coriaria, Royle, ex Lindl. Bot. Reg. xxvii. (1841) t. 46 = aristata.
corymbosa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 135.—Ins. Juan. Fernand.
crataegina, DC. Syst. ii. 9 = vulgaris.
crenulata, Schrad. in Linnaea, xii. (1838) 362 = vulgaris.
cretica, Falk, Beitr. ii. 164 = sibirica.
cretica, Linn. Sp. Pl. 331 = vulgaris.
cretica, Thunb. Fl. Jap. 146 = sinensis.
cretica, Viv. Fl. Cors. 5 = aetnensis.
crispa, *C. Gay, Fl. Chil.* i. 86.—Chili.
cuneata, *DC. Syst.* ii. 16.—Am. austr.
cuneata, Hort. ex C. Koch, Dendrol. i. 415 = empetrifolia.
Darwinii, *Hook. Ic. Pl.* t. 672.—Chili.
dasystachya, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 308.—China.
daurica, *Lodd. ex Loud. Hort. Brit.* 139.—Davuria.
dealbata, *Lindl. Bot. Reg.* t. 1750.—Mexic.
declinata, *Schrad. in Linnaea*, xii. (1838) 368.—China ?
densa, Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 54 = rigidifolia.
densiflora, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 9 = vulgaris.
densiflora, *Rafin. Sylva Tellur.* 68.—Hab. ?
dentata, Tausch, in Flora, xxi. (1838) 729 = vulgaris.
diaphana, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 309.—China.
diffusa, *C. Gay, Fl. Chil.* i. 80.—Chili.
discolor, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 287.—Venezuela.
discolor, Willd. ex Steud. Nom. ed. II. i. 197 = rigidifolia.
diversifolia, *Steud. l. c.*—Reg. Argent.
dulcis, Hort. ex C. Koch, Dendrol. i. 395 = vulgaris.
dulcis, Sweet, Brit. Flow. Gard. Ser. I. ii. t. 100 = buxifolia.
dumetorum, Gouan, Fl. Monsp. 312 = vulgaris.
edulis, Hort. ex C. Koch, Dendrol. i. 395 = vulgaris.
Ehrenbergii, *Kunze, in Linnaea*, xx. (1847) 45.—Mexic.
elegans, Hort. ex C. Koch, Dendrol. i. 409 = aristata.
emarginata, Willd. Enum. Hort. Berol. 395 = vulgaris.
empetrifolia, *Lam. Illustr.* t. 253. f. 4.—Reg. Magellan.
fascicularis, Sims, Bot. Mag. t. 2396 = pinnata.
fasciculata, Schult. f. Syst. vii. 17 = pinnata.
Fendleri, *A. Gray, Pl. Fendl.* 5.—Am. bor. occ.
ferox, *C. Gay, Fl. Chil.* i. 80.—Chili.
ferruginea, *Lechl. Berb. Am. Austr.* 9.—Bolivia.
Fischeri, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
flexuosa, *Ruiz & Pav. Fl. Per.* iii. 52. t. 281. f. A.—Peruv.
floribunda, Wall. Cat. n. 1474; et G. Don, Gen. Syst. i. 115 = aristata.
floribunda, Hort.—Cf. Gard. Chron. (1869) 612 = B. Aquifolinm.
florida, *Phil. in Linnaea*, xxxiii. (1864-65) 5.—Chili.
Fortunei, *Lindl. in Journ. Hort. Soc.* i. (1846) 231, 300.—China.
fraxinifolia, *Hook. Ic. Pl.* t. 329.—Mexic.
Fremontii, *Torr. in Bot. Mex. Bound.* 30.—Am. bor. occ.
glauca, Benth. Pl. Hartw. 125. n. 710 = Jamesoni.
glauca, *DC. Syst.* ii. 10; *H. B. & K. Nov. Gen. et Sp.* v. 71.—N. Granat.

BERBERIS:—
glauca, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
glaucescens, *St. Hil. Fl. Bras. Mer.* i. 45.—Bras.
globosa, *Benth. Pl. Hartw.* 158.—N. Granat.
glomerata, *Hook. & Arn. Bot. Beech. Voy.* 5.—Chili.
glumacea, Spreng. Syst. ii. 120 = nervosa.
Gondotii, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 52.—N. Granat.
gracilis, Hort. ex Lindl. in Journ. Hort. Soc. v. (1850) 12 = aristata.
gracilis, *Hartw. ex Benth. Pl. Hartw.* 34. n. 271.—Mexic.
gracillima, C. Koch, in Miq. Ann. Mus. Bot. Lugd. Bat. i. 253 = aristata.
grandiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 287.—Ecuador.
gratissima, *Klotzsch, in Bot. Ergeb. Waldem. Reise*, 130.—Himal.
Grevilleana, *Gill. in Hook. Bot. Misc.* iii. (1833) 136.—Chili.
Grisebachii, *Lechl. Berb. Am. Austr.* 34.—Chili.
Guilache, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 54.—N. Granat.
Guimpeli, *C. Koch & Bouché, in App. Ind. Hort. Berol.* (1854) 13.—Am. bor.
Haenkeana, *Presl, ex Schult. f. Syst.* vii. 1616.—Chili.
Hartwegi, *Benth. Pl. Hartw.* 34. n. 272.—Mexic.
heterophylla, Hort. ex C. Koch, Dendrol. i. 395 = vulgaris.
heterophylla, *Juss. in Poir. Encyc.* viii. 622.—Reg. Magellan.
heteropoda, Schrenk, Enum. Pl. Nov. 102 = vulgaris.
hispanica, Boiss. & Reut. Pugill. Pl. Nov. (1852) 3 = vulgaris.
Hookeri, Hort. ex Lem. Illustr. Hortic. vi. (1859) t. 207 = Wallichiana.
horrida, *C. Gay, Fl. Chil.* i. 84.—Chili.
horrida, Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 42 = Wallichiana.
hypoleuca, Lindl. in Journ. Hort. Soc. ii. (1847) 246 = asiatica.
iberica, Sweet, Hort. Brit. ed. I. 13 = vulgaris.
ilicifolia, Bert. ex Steud. Nom. ed. II. i. 197 = glomerata, stipulacea.
ilicifolia, Booth, ex Steud. l. c. = vulgaris.
ilicifolia, *Forst. in Comm. Gott.* ix. (1789) 28.—Reg. Magellan.
ilicifolia, Hort. ex C. Koch, Dendrol. i. 402 = aetnensis.
ilicifolia, Roxb. in As. Res. vi. (1799) 357 = asiatica.
ilicifolia, Scheele, in Linnaea, xxi. (1848) 591 = trifoliolata.
ilicina, *Hemsl. Biol. Centr. Am. Bot.* i. 23.—Mexic.
inermis, *Pers. Syn.* i. 387.—Reg. Magell.
innominata, Kielm. Dec. Rar. Pl. Tueb. 18 = vulgaris.
insignis, *Hook. f. & Thoms. Fl. Ind.* i. 226.—Reg. Himal.
integerrima, Bunge, in Linnaea, xviii. (1844) 149 = vulgaris.
integerrima, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
intermedia, Hort. ex C. Koch, l. c. 404 = Guimpeli.
intermedia, Hort. Angl.—Cf. Gard. Chron. (1869) 612 = nepalensis.
irritabilis, Salisb. Prod. 213 = vulgaris.
Jacquini, Hort. ex C. Koch, Dendrol. i. 394 = vulgaris.
Jamesoni, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 286.—Ecuador.
Jamesonii, *Lindl. in Journ. Hort. Soc.* v. (1850) 8; *Lem. Illustr. Hortic.* vi. (1859) t. 201.—Ecuador.
japonica, *R. Br. in Tuckey, Congo Exped. App.* 22.—Japon.
japonica, Lindl. ex Lindl. & Paxt. Flow. Gard. i. (1850-51) 11 = Bealii.
kasgarica, *Rupr. Sert. Tiansch.* 38.—As. temp.
Klotzschiana, *Warcz. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 292, nomen.—Bolivia.
kunawurensis, Royle, Illustr. Bot. Himal. 64 = vulgaris.
Knightii, Hort. ex C. Koch, Dendrol. i. 413 = Bealei.
Lagenaria, Poir. Encyc. viii. 619 = ilicifolia.
lanceolata, *Benth. Pl. Hartw.* 34.—Mexic.
latifolia, *Ruiz & Pav. Fl. Per.* iii. 52. t. 282.—Peruv.
laurina, *Thunb. Pl. Bras.* i. 8.—Bras.

BERBERIS :—
laxa, *Rafin, Sylva Tellur.* 68.—Hab. ?
laxiflora, Schrad. in Linnaea, xii. (1838) 367 = vulgaris.
Leschenaultiana, Wall. Cat. n. 1479 = nepalensis.
Ligonii, Hort. ex C. Koch, Dendrol. ii. 411 = concinna.
linearifolia, *Phil. in Linnaea*, xxviii. (1856) 663.—Chili.
littoralis, *Phil. Fl. Atac.* 7.—Chili.
littoralis, *Rupr. Fl Cauc.* 49, in textu.—Reg. Caucas.
loxensis, *Benth. Pl. Hartw.* 125.—Peruv.
lucida, *Schrad. in Linnaea*, xii. (1838) 363.—Iberia.
lutea, *Ruiz & Pav. Fl. Per.* iii. 51. t. 280.—Ecuador.
Lycium, Hort. ex C. Koch, Dendrol. i. 393 = vulgaris.
Lycium, *Royle, Illustr. Bot. Himal.* 64.—Reg. Himal.
macracantha, Hort. ex C. Koch, Dendrol. i. 400 = canadensis.
macracantha, Schrad. in Linnaea, xii. (1838) 366 = vulgaris.
macrocarpa, Schrad. l. c. 370 = canadensis.
macrophylla, Hort. ex Lindl. in Journ. Hort. Soc. v. (1850) 4 = Wallichiana.
macrophylla, Hort. ex C. Koch, Dendrol. i. 408 = asiatica.
macrosepala, *Hook. f. & Thoms. Fl. Ind.* i. 228.—Reg. Himal.
macrotheca, Hort. ex C. Koch, Dendrol. i. 400 = canadensis.
maderensis, *Lowe, in Hook. Kew Journ.* viii. (1856) 289.—Ins. Madera.
magellanica, Hort. ex C. Koch, Dendrol. i. 414 = Bealei.
marginata, *C. Gay, Fl. Chil.* i. 88.—Chili.
marginata, Hort. ex C. Koch, Dendrol. i. 394 = vulgaris.
Maximowiczi, Regel, Gartenfl. (1872) 238 = vulgaris.
melanocarpa, *Schau. Jahrb. Ges. Vat. Cult.* xxviii. 180.—Am. bor.
Miccia, Buch.-Ham. in D. Don, Prod. Fl. Nep. 205 = nepalensis.
microcarpa, *Warcz. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 292.—Bolivia.
microphylla, F. G. Dietr. Vollst. Lex. Gaertn. ii. 184 = vulgaris.
microphylla, Forst. in Comm. Gott. ix. (1789) 29 = buxifolia.
microphylla, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
minor, Forst. ex DC. Syst. ii. 15 = buxifolia.
mitis, Schrad. in Linnaea, xii. (1838) 371 = vulgaris.
monosperma, Hort. ex C. Koch, Dendrol. i. 405 = sinensis.
monosperma, *Ruiz & Pav. Fl. Per.* iii. 52.—Peruv.
montana, *C. Gay, Fl. Chil.* i. 90.—Chili.
moranensis, *Schult. f. Syst.* vii. 17.—Mexic.
multiflora, *Benth. Pl. Hartw.* 124.—Peruv.
mutabilis, *Phil. in Anal. Univ. Chil.* (1872) 665.—Chili.
neglecta, *Hort. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 284.—Hab. ?
nepalensis, Hort. ex C. Koch, Dendrol. i. 393 = vulgaris.
nepalensis, *Spreng. Syst.* ii. 120.—Reg. Himal.
nervosa, *Pursh, Fl. Am. Sept.* 219. t. 5.—Am. bor. occ.
Neuberti ×, Hort. ex Lem. Illustr. Hortic. i. (1854) Misc. 111. t. c.
nigra, *Rafin. Sylva Tellur.* 67.—Am. bor.
nitens, Hort. ex C. Koch, Dendrol. i. 399 = canadensis.
nitens, Schrad. in Linnaea, xii. (1838) 382 = vulgaris.
nitida, Willd. ex Steud. Nom. ed. II. i. 197 = quinduensis.
Nummularia, Bunge, in Linnaea, xviii. (1844) 149 = vulgaris.
obovata, Schrad. l. c. xii. (1838) 385 = vulgaris.
orthobotrys, *Bienert, ex Aitch. in Journ. Linn. Soc.* xix. (1882) 150.—Persia.
pallida, *Hartw. ex Benth. Pl. Hartw.* 34.—Mexic.
paniculata, H. B. & K. ex DC. Syst. ii. 10 = glauca.
paniculata, *Juss. ex DC. l. c.* 12.—Peruv.
paniculata, *Hemsl. Biol. Centr. Am. Bot.* i. 24.—Costa-Rica.
paniculata, *Phil. in Anal. Univ. Chil.* (1872) 664.—Chili.

BERBERIS :—
parviflora, *Lindl. in Journ. Hort. Soc.* ii. (1847) 243.—Am. austr.
pauciflora, Salisb. Prod. 213 = vulgaris.
Pearcei, *Phil. in Linnaea*, xxxiii. (1864-65) 4.—Chili.
petiolaris, Wall. ex G. Don, Gen. Syst. i. 116 = aristata.
pichinchensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 286.—Ecuador.
pichinchensis, *Wedd. Chlor. And.* ii. 295 (= praec. ?).—Ecuador.
pinnata, Banks, ex DC. Syst. ii. 20 = B. Aquifolium.
pinnata, Buch, ex DC. l. c. 21 = nervosa.
pinnata, H. B. & K. Nov. Gen. et Sp. v. 434 = moranensis.
pinnata, *Lag. Elench. Hort. Madr.* (1803) 6.—Mexic.
pinnata, Roxb. Fl. Ind. ii. 184 = nepalensis.
pisifera, Rafin. Sylva Tellur. 68 = canadensis.
polymorpha, *Phil. in Linnaea*, xxviii. (1856) 664.—Chili.
polypetala, *Phil. in Anal. Univ. Chil.* (1872) 666.—Chili.
provincialis, Audib. ex Schrad. in Linnaea, xii. (1838) 381 = vulgaris.
psilopoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 285.—N. Granat.
purpurea, *Rafin. Sylva Tellur.* 67.—Hab. ?
quinduensis, *H. B. & K. Nov. Gen. et Sp.* v. 70. t. 432.—Ecuador.
racemosa, Stokes, Bot. Mat. Med. ii. 296 = vulgaris.
rariflora, *Lechl. Berb. Am. Austr.* 33.—Bolivia.
repens, Lindl. Bot. Reg. t. 1176 = B. Aquifolium.
reticulata, *Steud. Nom.* ed. II. i. 197.—Chili.
retinervis, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 52.—N. Granat.
revoluta, Sm. ex DC. Syst. ii. 16 = empetrifolia.
rigida, Willd. ex Steud. Nom. ed. II. i. 197 = conferta.
rigidifolia, *H. B. & K. Nov. Gen. et Sp.* v. 70. t. 431.—Am. austr.
Roemeriana, Scheele, in Linnaea, xxii. (1849) 352 = trifoliata.
rotundifolia, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 63.—Chili.
rotundifolia, Hort. ex Lindl. in Journ. Hort. Soc. v. (1850) 2 = buxifolia.
rubra, Poit. & Turp. in Duham. ed. Nov. iv. 12 = vulgaris.
ruscifolia, *Lam. Illustr.* t. 253. f. 2.—Reg. Argent.
sanguinea, Hort. ex C. Koch, Dendrol. i. 394 = vulgaris.
sanguinolenta, Hort. ex C. Koch, l. c. = vulgaris.
sanguinolenta, *Schrad. in Linnaea*, xii. (1838) 378.—Hab. ?
saxicola, *Lechl. Berb. Am. Austr.* 42.—Peruv.
Schiedeana, *Schlecht. in Bot. Zeit.* xii. (1854) 654.—Mexic.
serotina, *Lange, Ind. Sem. Haun.* (1872) 5, adnot.—Hab. ?
serratifolia, Hort. ex C. Koch, Dendrol. i. 406 = aristata.
serrato-dentata, *Lechl. Berb. Am. Austr.* 16.—Chili.
serrulata, Rafin. Sylva Tellur. 68 = canadensis.
sibirica, Hort. ex Schult. f. Syst. vii. 2 = vulgaris.
sibirica, *Pall. Fl. Ross.* ii. 42. t. 67.—Sibir. altaic.
sicula, Jan, Elench. ex Schult. f. Syst. vii. 3 = aetnensis.
Sieboldii, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 69.—Japon.
sinensis, *Desf. Tabl.* ed. I. (1804) 150.—China.
sinensis, DC. Syst. ii. 6 = Guimpeli.
spathulata, *Schrad. in Linnaea*, xii. (1838) 376.—Am. bor.
sphaerocarpa, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 376 = vulgaris.
spinosa, Comm. ex Decne. in Bot. Voy. Astrol. & Zél. 65 = buxifolia.
spinulosa, *St. Hil. Fl. Bras. Mer.* i. 45.—Bras.
stenophylla, *Hance, in Journ. Bot.* xx. (1882) 257.—China.
stenophylla ×, Hort.—Cf. Gard. Chron. (1864) 460; (1869) ii. 612; (1882) ii. 466; C. Koch, Dendrol. i. 412 = Darwinii.
stipulacea, *Steud. Nom.* ed. II. i. 197.—Chili.

BERBERIS :—
sulcata, *C. Koch & Bouché, in App. Ind. Sem. Hort. Berol.* (1854) 13.—Hab. ?
Swaseyi, *Buckley, in Young, Fl. Texas.* 152.—Texas.
tenuifolia, *Lindl. Bot. Reg.* (1838) *Misc.* 64.—Mexic.
Thunbergii, DC. Syst. ii. 9 = vulgaris.
tinctoria, Leschen. in Mém. Mus. Par. ix. (1822) 306 = aristata.
toluacensis, *Hort.—Cf. Gard. Chron.* (1868) 435.—Am. austr.
tolimensis, *Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 52.—N. Granat.
tomentosa, *Ruiz & Pav. Fl. Per.* iii. 52. t. 282 b.—Chili.
tortuosa, Domb. ex DC. Syst. ii. 11 = flexuosa.
tragacanthoides, *DC. Syst.* ii. 18.—Armenia.
tricuspidata, Sm. ex DC. l. c. 17 = heterophylla.
trifolia, *Schult. f. Syst.* vii. 1616.—Mexic.
trifoliata, Hartw. ex Lindl. Bot. Reg. xxvii. (1841) Misc. 68 = seq.
trifoliolata, *Moric. Pl. Nouv. Am.* 113. t. 69.—Texas.
trifurca, *Lindl. in Paxt. Flow. Gard.* iii. 57 (1852-53).—China.
trigona, *Kunze, in Poepp. & Endl. Nov. Gen. et Sp.* ii. 63. t. 187.—Chili.
truxillensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 285.—Venezuela.
Tschonoskyana, *Regel, in Act. Hort. Petrop.* ii. (1873) 421.—Japon.
turcomannica, Kar. ex Ledeb. Fl. Ross. i. 79 = vulgaris.
ulicina, *Hook. f. & Thoms. Fl. Ind.* i. 227.—Tibet. occ.
umbellata, Lindl. Bot. Reg. xxx. (1844) t. 44 = aristata.
umbellata, *Phil. in Anal. Univ. Chil.* (1872) 663.—Chili.
umbellata, *Wall. Cat.* n. 1475; *et ex G. Don, Gen. Syst.* i. 116.—Reg. Himal.
undulata, Hort. ex C. Koch, Dendrol. i. 408 = asiatica.
undulata, *Lindl. in Journ. Hort. Soc.* v. (1850) 7.—Peruv.
valdiviana, *Phil. in Linnaea*, xxviii. (1856) 609.—Chili.
verticillata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 288.—N. Granat.
violacea, Poit. & Turp. in Duham. ed. Nov. iv. 12; Hort. ex C. Koch, Dendrol. i. 394 = vulgaris.
virgata, Hort. ex Lindl. in Journ. Hort. Soc. v. (1850) 6 = parviflora.
virgata, Hort. ex C. Koch, Dendrol. i. 413 = actinacantha.
virgata, *Ruiz & Pav. Fl. Per.* iii. 51. t. 281. f. B.—Peruv.
vulgaris, *Linn. Sp. Pl.* 330.—Europ.; As. temp.
vulgaris, Vell. Fl. Flum. iii. t. 149 = glaucescens.
Wallichiana, *DC. Prod.* i. 107.—Reg. Himal.
Weddelli, *Lechl. Berb. Am. Austr.* 21.—Bolivia.
xanthoxylon, Hassk. Cat. Hort. Bog. Alt. 180 = Wallichiana ?

BERCHEMIA, Neck. Elem. ii. 122 (1790). *RHAMNEAE*, Benth. & Hook. f. i. 377.
OENOPLEA, Hedw. f. Gen. i. 151 (1806).
OENOPLIA, Schult. ex Roem. & Schult. Syst. v. 332 (1819).
affinis, *Hassk. Hort. Bog. Desc.* 117.—Malaya.
Burmanniana, *DC. Prod.* ii. 23.—Zeylan.
calophylla, G. Don, Gen. Syst. ii. 28 = Zizyphus calophylla.
congesta, S. Moore, in Journ. Bot. xiii. (1875) 226 = Michrorhamnus franguloides.
corollata, F. Muell. Census, 60 = ecorollata.
discolor, *Hemsl. in Oliver, Fl. Trop. Afr.* i. 381.—Afr. trop.
Edgeworthii, *Lawson, in Hook. f. Fl. Brit. Ind.* i. 638.—Reg. Himal.
ecorollata, *F. Muell. Fragm.* ix. 141.—Austral.
flavescens, *Wall. Cat.* n. 4255; *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 357.—Reg. Himal.
floribunda, *Wall. Cat.* n. 4256; *Brongn. l. c.*—Reg. Himal.
Fournieri, *Panch. & Sebert, Not. Bois. Nouv. Cald.* 236.—N. Caled.

BERCHEMIA:—
hamosa, Wall. Cat. n. 4253 = Sageretia hamosa.
laxa, Wall. l. c. n. 4257 = floribunda.
lineata, Benth. in Hook. Kew Journ. iv. (1852) 42 = racemosa.
lineata, *DC. Prod.* ii. 23.—China.
Loureiriana, *DC. l. c.*—China.
oppositifolia, Wall. Cat. n. 4254 = Sageretia oppositifolia.
parviflora, Wall. l. c. n. 4258; Wight, Ic. i. 19 = Sageretia oppositifolia.
Poiretiana, DC. Prod. ii. 23 = lineata.
polyphylla, *Wall. Cat.* n. 4259.—Burma.
pubiflora, *Miq. Fl. Ind. Bat.* i. 1. 645.—Malaya.
racemosa, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1843) 147.—Japon.
repanda, *Rafin. Sylva Tellur.* 33.—Hab.?
sessiliflora, *Benth. Fl. Hongk.* 68.—China.
trichantha, *Miq. Fl. Ind. Bat. Suppl.* i. 331.—Malaya.
undulata, *Rafin. Sylva Tellur.* 33.—Hab.?
volubilis, *DC. Prod.* ii. 23.—Am. bor.

BERCHTOLDIA, Presl, Rel. Haenk. 323. t. 43 (1830) = **Chaetium**, Nees (Gramin.).
bromoides, Presl, l. c. 324 = C. bromoides.
holciformis, Fourn. ex Benth. in Journ. Linn. Soc. xix. (1881) 46 = Oplismenus holciformis.
junciformis, Hemsl. Biol. Centr. Am. Bot. iii. 501, lapsu = Oplismenus holciformis.
oplismenoides, Fourn. ex Benth. in Journ. Linn. Soc. xix. (1881) 46 = Echinochloae sp.?

BERCKHEYA, Pers. Syn. ii. 480 (1807) = **Berkheya**, Ehrh. (Compos.).

BEREBERA, Baker, in Oliver, Fl. Trop. Afr. ii. 131 (1871), sphalm = BERREBERA, Hochst. = **Millottia**, Wight & Arn. (Legumin.).

BERENDTIA, A Gray, in Proc. Am. Acad. vii. (1868) 379. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 942.
Coulteri, *A. Gray, l. c.* 380.—Mexic.
Ghiesbrechtii, A. Gray, l. c. = rugosa.
rugosa, *A. Gray, l. c.*—Mexic.

BERENICE, Salisb. Gen. Pl. Fragm. 89 (1866) = **Allium**, Linn. (Liliac.).
Victorialis, Salisb. 490 = A. Victorialis.

BERENICE, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 156. *SAXIFRAGEAE*, Benth. & Hook. f. i. 646.
arguta, *Tul. l. c.*—Ins. Borbon.

BERGENA, Adans. Fam. ii. 345 (1763) = **Lecythis**, Linn. (Myrtac.).

BERGENIA, Moench, Meth. 664 (1794) = **Saxifraga**, Tourn.
biflora, Moench, l. c. = S. crassifolia.
cordifolia, Sternb. Rev. Sax. Suppl. ii. 2 = S. crassifolia.
purpurascens, Engl. in Verh. Zool. Bot. Ges. Wien, xix. (1869) 549 = S. purpurascens.

BERGENIA, Neck. Elem. ii. 108 (1790) = **Lythrum**, Linn.

BERGENIA, Rafin. Sylva Tellur. 102 (1838) = **Cuphea**, P. Br. (Lythrar.).
siphilitica, Rafin. l. c.—Hab.?

BERGERA, Koen. ex Linn. Mant. ii. 563 (1771) = **Murraya**, Linn. (Rutac.).
compressa, Blanco, Fl. Filip. ed. I. 360.—Ins. Philipp.
inodora, Blanco, l. c. 361.—Ins. Philipp.
integerrima, Buch.-Ham. ex Roxb. Hort. Beng. 32; Fl. Ind. ii. 376 = Micromelum pubescens.
intermedia, Zoll. & Moritzi, Syst. Verz. Zoll. 24.—Java.
Koenigii, Linn. Mant. 565 = Murraya Koenigii.
nitida, Thw. Enum. Pl. Zeyl. 46 = Clausena indica.
ternata, Blanco, Fl. Filip. ed. I. 360 = Glycosmis bilocularis.
villosa, Wall. Cat. n. 6372 = Micromelum pubescens.
Wallichii, A. DC. ex Oliver, in Journ. Linn. Soc. v. Suppl. ii. (1861) 35 = Clausena Wallichii.

BERGERETIA, Desv. Journ. Bot. iii. (1813) 161 = **Clypeola**, Linn. (Crucif.).
echinata, Desv. l. c. = C. lasiocarpa.

BERGERIA, Koen. ex Steud. Nom. ed. II. i. 849, in syn. (1840) = **Koenigia**, Linn. (Polygon.).

BERGERONIA, Micheli, in Mém. Soc. Phys. Genèv. xxviii. (1883) n. 7 (Contr. Fl. Parag. 38). *LEGUMINOSAE.*
sericea, *Micheli, ll. cc.* 39.—Paraguay.

BERGHAUSIA, Endl. Gen. Suppl. iii. 57 (1843) = **Garnotia**, Brongn. (Gramin.).
barbulata, Endl. ex Miq. Anal. Bot. Ind. 32.—Malaya.
courtallensis, Nees, ex Miq. l. c. = G. courtallensis.
elata, Arn. ex Miq. l. c.—Malaya.
Emodi, Endl. ex Miq. l. c.—Malaya.
mutica, Munro, in Proc. Am. Acad. iv. (1864) 362.—Hongkong.
pallens, Arn. ex Miq. Anal. Bot. Ind. 33.
patula, Munro, in Proc. Am. Acad. iv. (1864) 362 = G. patula.
tenella, Arn. ex Miq. Anal. Bot. Ind. 34.—Malaya.

BERGHESIA, Nees, in Linnaea, xx. (1847) 701. *RUBIACEAE*?, Benth. & Hook. f. ii. 29.
coccinea, *Nees, l. c.* 702.—Mexic.

BERGHIAS, Juss. in Mém. Mus. Par. vi. (1820) 399 = BERGKIAS, Sonn. = **Gardenia**, Ellis (Rubiac.).

BERGIA, Linn. Mant. ii. 152. n. 1309 (1771). *ELATINEAE*, Benth. & Hook. f. i. 163.
BERGIERIA, Neck. Elem. ii. 237 (1790).
LANCRETIA, Del. Fl. Egypte, 213. t. 25. f. 1 (1813).
MERIMEA, Cambess. in St. Hil. Fl. Bras. Mer. ii. 160. t. 107 (1829).
SPHONDYLOCOCCA, Willd. ex Roem. & Schult. Syst. vi. p. lxx. 799 (1820).
abyssinica, *A. Rich. Tent. Fl. Abyss.* i. 49.—Abyss.
aestivosa, *Steud. Nom.* ed. II. i. 198.—Ind. bor. occ.
ammannioides, *Roxb. Hort. Beng.* 34; *Fl. Ind.* ii. 456.—As. Afr. et Austral. trop.
anagalloides, *Walp. Rep.* ii. 286.—Afr. austr.
aquatica, Roxb. Pl. Corom. ii. 22. t. 142 = verticillata.
arenaroides, *Fenzl, in Denkschr. Baier. Bot. Ges.* iii. (1841) 183.—Bras.
capensis, *Linn. Mant.* 241.—Afr. austr.
decumbens, *Planch. ex Harv. Thes. Cap.* i. 15. t. 24.—Afr. austr.
erecta, *Guill. & Perr. Tent. Fl. Seneg.* 44.—Afr. trop.
glandulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 371.—Ins. Luzon.
glomerata, *Linn. f. Suppl.* 243.—Afr. austr.
odorata, *Edgew. in Journ. As. Soc. Beng.* vii. (1838) 765.—Afr. trop.; Persia; Ind. bor. occ.
oryzetorum, *Fenzl, ex Zoll. Syst. Verz. Ind. Archip.* 62.—Malaya.
pedicellaris, *F. Muell. Fragm.* ii. 145.—Austral.
pentandra, Cambess. ex Guill. & Perr. Tent. Fl. Seneg. 42. t. 12 = ammanoides.
peploides, *Guill. & Perr. l. c.* 44.—Afr. trop.
perennis, *F. Muell. Fragm.* ii. 146.—Austral.
polyantha, *Sond. in Linnaea*, xxiii. (1850) 16.—Afr. austr.
pusilla, *Benth. Fl. Austral.* i. 180.—Austral.
repens, *Blume, Bijdr.* 62.—Java.
serrata, *Blanco, Fl. Filip.* ed. I. 387.—Ins. Philipp.
sessiliflora, *Griseb. Cat. Pl. Cub.* 40.—Cuba.
suffruticosa, Fenzl, in Denkschr. Baier. Bot. Ges. iii. (1841) 183 = odorata.
texana, *Seub. ex Walp. Rep.* i. 285.—Texas.
trimera, Fisch. & Mey. in Linnaea, x. (1836) 74 = ammannioides.
tripetala, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 66.—Austral.
verticillata, *Willd. Sp. Pl.* ii. 770 = capensis.

BERGIERA, Neck. Elem. ii. 237 (1790) = praec.

BERGINIA, Harv. ex Benth. & Hook. f. Gen. ii. 1096 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.
virgata, *Harv. ex Benth. & Hook. f. l. c.* 1097.—Calif.

BERGKIAS, Sonner. Voy. 48. t. 17 (1776) = **Gardenia**, Ellis (Rubiac.).

BERGSMIA, Blume, Rumphia, iv. 23. t. 178 C. f. 2 (1848). *BIXINEAE*, Benth. & Hook. f. i. 129.
acuminata, *Miq. Fl. Ind. Bat. Suppl.* i. 389.—Sumatra.
javanica, *Blume, Rumphia*, iv. 23.—Java.
sumatrana, *Miq. Fl. Ind. Bat. Suppl.* i. 389.—Sumatra.

BERHARDIA, C. Muell. in Walp. Ann. v. 39 (1845), sphalm. = **Berardia**, Vill. (Compos.).

BERIESA, Steud. Nom. ed. II. i. 199 (1840) = SIEBERA, Presl = **Anredera**, Juss. (Chenop.).
baselloides, Steud. l. c. = A. spicata.

BERINGERIA, Neck. Elem. i. 312 (1790) = **Ballota**, Linn. (Labiat.).
acetabulosa, Neck. l. c. = Ballota acetabulosa.
africana, Neck. l. c. = Ballota africana.
cinerea, G. Don, in Sweet, Hort. Brit. ed. III. 545 = Ballota hirsuta.
crispa, G. Don, l. c. = Ballota africana.
frutescens, Reichb. Fl. Germ. Excurs. 325 = Ballota spinosa.
hirsuta, Neck. Elem. 312 = Ballota hirsuta.
hispanica, Neck. l. c. = Ballota hispanica.
mollissima, Heynh. Nom. ii. 66 = Ballota mollissima.
orientalis, G. Don, in Sweet, Hort. Brit. ed. III. 548 = Ballota orientalis (sp. dubia).
pseudodictamnus, Neck. Elem. i. 312; Link, Handb. 478 = Ballota pseudodictamnus.
rugosa, Heynh. Nom. ii. 66 = Ballota saxatilis.

BERINIA, Brignol. Pl. Forojul. 50 (1810) = **Crepis**, Vaill. (Compos.).
abyssinica, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 315 = C. abyssinica.
acuminata, Sch. Bip. l. c. = C. acuminata.
alpestris, Sch. Bip. l. c. = C. alpestris.
andryaloides, Brignol. Pl. Forojul. 50 = C. chondrilloides.
aurea, Sch. Bip. in Pollichia, xxii–xxiv. (1866) 318 = C. aurea.
auriculaefolia, Sch. Bip. l. c. 315 = C. auriculaefolia.
biennis, Sch. Bip. l. c. 318 = C. biennis.
chondrilloides, Sieb. ex Sch. Bip. l. c. 315 = C. chondrilloides.
chrysantha, Sch. Bip. l. c. 318 = C. chrysantha.
ciliata, Sch. Bip. l. c. 317 = C. ciliata.
crocea, Sch. Bip. l. c. = C. crocea.
foetida, Sch. Bip. l. c. 315 = C. foetida.
grandiflora, Sch. Bip. l. c. 317 = C. grandiflora.
heterantha, Sch. Bip. l. c. = Lactuca Dubyaea.
heterotricha, Sch. Bip. l. c. = C. heterotricha.
Jacquinii, Sch. Bip. l. c. 318 = C. Froelichii?
jubata, Sch. Bip. l. c. = C. jubata.
lacera, Sch. Bip. l. c. 315 = C. lacera.
nicaeënsis, Sch. Bip. l. c. 318 = C. nicaeënsis.
occidentalis, Sch. Bip. l. c. 315 = C. occidentalis.
oporinioides, Sch. Bip. l. c. = C. oporinioides.
pannonica, Sch. Bip. l. c. = C. rigida.
pinnatifida, Sch. Bip. l. c. = C. pinnatifida.
polytricha, Sch. Bip. l. c. 317 = C. polytricha.
runcinata, Sch. Bip. l. c. 315 = C. runcinata.
Sibthorpiana, Sch. Bip. l. c. = C. Sibthorpiana.
stricta, Sch. Bip. l. c. 317 = C. virens? vel tectorum.
tenuifolia, Sch. Bip. l. c. 316 = C. tenuifolia.

BERKHEYA, Ehrh. Beitr. iii. 137 (1788). *COMPOSITAE*, Benth. & Hook. f. ii. 460.
AGRIPHYLLUM, Juss. Gen. 190 (1789).
APULEIA, Gaertn. Fruct. ii. 439. t. 171 (1791).
ARELINA, Neck. Elem. i. 83 (1790).
ASPIDALIS, Gaertn. Fruct. ii. t. 171 (1791).
BASTERIA, Houtt. Nat. Hist. Pl. iv. 158 (1776).
BASTERA, J. F. Gmel. Syst. 1271 (1791).
BERCKHEYA, Schreb. Gen. ii. 577 (1791).
CROCODILOIDES, Adans. Fam. ii. 127 (1763).
CUSPIDIA, Gaertn. Fruct. ii. 454. t. 171 (1791).
EVOPIS, Cass. in Bull. Soc. Philom. (Févr. 1818) 32; et in Dict. Sc. Nat. xvi. 65 (1820).
GORTERIA, Linn. Sp. Pl. ed. II. 1282 (1763), pro parte; Lam. Illustr. t. 702 (1789?).
HETERORACHIS, Sch. Bip. in Flora, xxvii. (1844) 775.
ROHRIA, Vahl, in Skr. Naturh.-Selsk. Kjoeb. iii. (1793) 97.
STOBAEA, Thunb. Prod. Pl. Cap. 141 (1800).
ZARABELLIA, Neck. Elem. i. 10 (1790).

BERKHEYA :—
albida, Lichtst. ex DC. Prod. vii. 36 = Dicoma capensis.
annectens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 509.—Afr. austr.
bisulca, Pers. Syn. ii. 480 = Cullumia bisulca.
canescens, *DC. Prod.* vi. 507.—Afr. austr.
carduiformis, *DC. l. c.*—Afr. austr.
carlinoides, *Willd. Sp. Pl.* iii. 2275.—Afr. austr.
carthamoides, *Willd. l. c.* 2274.—Afr. austr.
cernua, R. Br. in Ait. Hort. Kew. ed. II. v. 139 = Didelta cernua.
ciliaris, Willd. Sp. Pl. iii. 2273 = Cullumia ciliaris.
coriacea, *Harv. in Harv. & Sond. Fl. Cap.* iii. 506.—Afr. austr.
corymbosa, *DC. Prod.* vi. 507.—Afr. austr.
cruciata, *Willd. Sp. Pl.* 2276.—Afr. austr.
cuneata, *Willd. l. c.* 2270.—Afr. austr.
cynaroides, *Willd. l. c.* 2275.—Afr. austr.
decurrens, Willd. l. c. 2276 = Stephanocoma berkheyoides.
Dregei, *Harv. in Harv. & Sond. Fl. Cap.* iii. 507.—Afr. austr.
Ecklonis, *Harv. l. c.* 504.—Afr. austr.
eryngiifolia, *Less. Syn. Comp.* 703.—Afr. austr.
fruticosa, *Ehrh. Beitr.* iii. 138.—Afr. austr.
gazanioides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 508.—Afr. austr.
gorterioides, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 429.—Afr. trop.
grandiflora, *Willd. Sp. Pl.* iii. 2271.—Afr. austr.
hirpicioides, *Sond. in Harv. & Sond. Fl. Cap.* iii. 509.—Afr. austr.
hispida, Willd. Sp. Pl. iii. 2274 = Cullumia hispida.
incana, Willd. l. c. 2269 = fruticosa.
Kraussii, Sch. Bip. in Flora, xxvii. (1844) 774 = cynaroides.
lanceolata, *Willd. Sp. Pl.* iii. 2270.—Afr. austr.
obovata, *Willd. l. c.* 2269.—Afr. austr.
palmata, *Willd. l. c.* 2271.—Afr. austr.
patula, Willd. l. c. 2272 = Cullumia patula.
pectinata, Willd. l. c. = Cullumia pectinata.
pinnata, *Less. Syn. Comp.* 75.—Afr. austr.
pungens, *Willd. Sp. Pl.* iii. 2275.—Afr. austr.
purpurea, *Benth. & Hook. f. ex Mast. in Gard. Chron.* (1872) 1262.—Afr. austr.
scolymoides, *DC. Prod.* vi. 504.—Afr. austr.
seminivea, *Harv. in Harv. & Sond. Fl. Cap.* iii. 507.—Afr. austr.
setifera, *DC. Prod.* vi. 507.—Afr. austr.
setosa, Willd. Sp. Pl. iii. 2273 = Cullumia setosa.
Spekeana, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 100.—Afr. trop.
spinosissima, *Willd. Sp. Pl.* iii. 2270.—Afr. austr.
squarrosa, Willd. l. c. 2272 = Cullumia bisulca.
stobaeoides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 505.—Afr. austr.
subulata, *Harv. l. c.* 507.—Afr. austr.
sulcata, Willd. Sp. Pl. 2271 = Cullumia sulcata.
umbellata, *DC. Prod.* vi. 507.—Afr. austr.
uniflora, *Willd. Sp. Pl.* iii. 2274.—Afr. austr.
Zeyheri, *Harv. & Sond. Fl. Cap.* iii. 496.—Afr. austr.

BERLANDIERA, DC. Prod. v. 517 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 350.
incisa, Torr. & Gray, Fl. N. Am. ii. 282 = lyrata.
longifolia, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 342 = texana.
lyrata, *Benth. Pl. Hartw.* 18.—Am. bor. occ.
pumila, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 342 = tomentosa.
subacaule, *Nutt. l. c.* 343.—Florida.
texana, *DC. in Deless. Ic. Sel.* iv. t. 26.—Am. bor.
tomentosa, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 343.—Am. bor.

BERLIERA, Buch.-Ham. ex Wall. Cat. sub n. 6225 (1832?) = **Myrioneuron**, R. Br. (Rubiac.).
nutans, Buch.-Ham. l. c. = M. nutans.

BERLINIA, Soland. ex Hook. f. & Benth. in Hook. Niger Fl. 326 (1849). *LEGUMINOSAE*, Benth. & Hook. f. i. 579.
acuminata, *Soland. ex Hook. f. & Benth. l. c.*—Afr. trop.
angolensis, *Welw. ex Benth. in Trans. Linn. Soc.* xxv. (1866) 310.—Afr. trop.
auriculata, *Benth. l. c.* 309.—Afr. trop.

BERLINIA :—
bracteosa, *Benth. in Trans. Linn. Soc.* xxv. (1866) 309.—Afr. trop.
Heudelotiana, *Baill. Adansonia*, vi. (1865) 185.—Afr. trop.
paniculata, *Benth. in Trans. Linn. Soc.* xxv. (1866) 311.—Afr. trop.
stipulacea, *Benth. l. c.* 310.—Afr. trop.

BERMUDIANA, [Tourn.] Linn. Syst. ed. I (1735) = **Sisyrinchium**, Linn. (Irid.).
graminea, Gaertn. Fruct. i. 32 = S. Bermudiana.
graminifolia, Medic. Malv. 77 = S. anceps.
guttata, Stokes, Bot. Mat. Med. i. 109 = Moraca chinensis.
iridifolia, Medic. Malv. 77 = S. Bermudiana.
striata, Moench, Meth. Suppl. 113 = S. striatum.

BERNARDIA, Endl. Gen. 469, in syn. (1838) = **Berardia**, Vill. (Compos.).

BERNARDIA, Houst. ex P. Br. Nat. Hist. Jamaic. 361 (1756). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 308.
ALEVIA, Baill. Etud. Gén. Euph. 508 (1858).
BIVONIA, Spreng. Neue Entdeck. ii. 116 (1821).
PASSAEA, Baill. Etud. Gén. Euph. 507. t. 18 (1858).
PHAEDRA, Klotzsch, in Endl. Gen. Suppl. iv. III. 88 (1847).
POLYBAEA, Klotzsch, l. c. (1847).
TRAGANTHUS, Klotzsch, in Wiegm. Archiv, vii. (1841) 188. t. 9.
TYRIA, Klotzsch, in Endl. Gen. Suppl. iv. III. 88 (1847).
axillaris, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 174.—Bras.
brevipes, *Muell. Arg. l. c.* 176.—Bras.
caperoniaefolia, *Muell. Arg. in DC. Prod.* xv. II. 920.—Reg. Argent.
capitellata, Muell. Arg. in Linnaea, xxxiv. (1865–66) 176 = eclastrinea.
carpinifolia, *Griseb. Fl. Brit. W. Ind.* 45.—Ind. occ.
celastrinea, *Muell. Arg. in DC. Prod.* xv. II. 921.—Bras.
confertifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 175.—Bras.
corensis, *Muell. Arg. l. c.* 173.—Ind. occ.
crassifolia, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 394.—Bras.
dichotoma, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 172.—Ind. occ.
fasciculata, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 153.—Am. bor. occ.
gambosa, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 391.—Bras.
Gardneri, *Muell. Arg. l. c.* 394.—Bras.
grandifolia, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 173.—Guiana.
hirsutissima, *Muell. Arg. in DC. Prod.* xv. II. 922.—Bras.
intermedia, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 160 = mexicana.
interrupta, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 171.—Mexic.
Jacquiniana, *Muell. Arg. l. c.* 173.—Venezuela.
leprosa, *Muell. Arg. l. c.* 172.—Ins. S. Doming.
Lorentzii, *Muell. Arg. in Journ. Bot.* xii. (1874) 229.—Reg. Argent.
megalophylla, Muell. Arg. in Linnaea, xxxiv. (1865–66) 173 = Adenophaedra megalophylla.
mexicana, *Muell. Arg. l. c.* 172.—Mexic.
microphylla, *Muell. Arg. l. c.*—Cuba.
multicaulis, *Muell. Arg. l. c.* 177.—Bras.
myricaefolia, *Benth. & Hook. f. Gen.* iii. 308.—Am. bor.
oligandra, Muell. Arg. in Linnaea, xxxiv. (1865–66) 173 = tamanduana.
pedunculoris, *Muell. Arg. l. c.* 176.—Bras.
pulchella, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 392.—Bras.
scabra, *Muell. Arg. l. c.* 396.—Bras.
Sellowii, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 177.—Bras.
sidoides, *Muell. Arg. l. c.*—Guiana.
spartioides, *Muell. Arg. l. c.*—Bras.
tamanduana, *Muell. Arg. in DC. Prod.* xv. II. 920.—Bras.
venosa, *Griseb. in Mem. Am. Acad.* N. S. viii. 1861) 159.—Ind. occ.

BERNARDINIA, Planch. in Linnaea, xxiii. (1850) 412. *CONNARACEAE*, Benth. &. Hook. f. i. 431.
fluminensis, *Planch. l. c.*—Bras.

BERNEUXIA, Decne. in Bull. Soc. Bot. Fr. xx. (1873) 159. *DIAPENSIACEAE*, Benth. & Hook. f. ii. 621.
thibetica, *Decne. l. c.*—Tibet.

BERNIERA, DC. Prod. vii. 18 (1838) = **Gerbera**, Gronov. (Compos.).
nepalensis, DC. l. c. = G. macrophylla.

BERNIERIA, Baill. in Bull. Soc. Linn. Par. i. (1884) 434. *LAURINEAE*.
madagascariensis, *Baill. l. c.*—Madag.

BERNOULLIA, Oliver, in Hook. Ic. Pl. t. 1169 (1873). *STERCULIACEAE*.
flammea, *Oliver, l. c.* t. 1169–70.—Guatemala.

BERREBERA, Hochst. in Flora, xxvii. (1844) 100; xxix. (1846) 597 = **Millettia**, Wight & Arn. (Legum.).
caffra, Hochst. ll. cc. 100 et 598 = M. caffra.
ferruginea, Hochst. ll. cc. 100 et 597 = M. ferruginea.

BERRIA, Roxb. Hort. Beng. 42 (1814); Pl. Corom. iii. 60. t. 264 (1819). *TILIACEAE*, Benth. & Hook. f. i. 232.
ESPERA, Willd. in Ges. Naturf. Fr. Berl. Neue Schr. iii. (1801) 449, nomen prius.
HEXAGONOTHECA, Turcz. in Bull. Soc. Nat. Mosc. (1846) II. 505.
Ammonilla, *Roxb. Hort. Beng.* 42; *Fl. Ind.* ii. 639.—Ceylan.; Ind. or.
mollis, Wall. Cat. n. 1186 = B. Ammonilla.
Vescoana, *Baill. Adansonia*, x. (1872) 240.—Ins. Tahiti.

BERRYA, DC. Prod. i. 517 (1824) = pracc.

BERRYA, Klein, ex Willd. Sp. Pl. ii. 840, in syn. (1800) = **Litsea**, Lam. (Laurin.).
chinensis, Klein, ex Willd. l. c. = L. sebifera.

BERSAMA, Fresen. in Mus. Senckenb. ii. (1837) 280. t. 17. *SAPINDACEAE*, Benth. & Hook. f. i. 412.
NATALIA, Hochst. in Flora, xxiv. (1841) 663.
RHAGANUS, E. Mey. Zwei Pfl. Docum. 216 (1843).
abyssinica, *Fresen. l. c.* 281.—Abyss.
integrifolia, *A. Rich. Tent. Fl. Abyss.* i. 107.—Abyss.
maxima, *Baker, in Oliver, Fl. Trop. Afr.* i. 434.—Afr. trop.
paullinioides, *Baker, l. c.* 435.—Afr. trop.
serrata, *A. Rich. Tent. Fl. Abyss.* i. 107.—Abyss.

BERTERA, Steud. Nom. ed. II. i. 199 (1840) = **Gladiolus**, Tourn. (Irid.).

BERTEROA, DC. in Mém. Mus. Par. vii. (1821) 232; et Syst. ii. 290 (1821) = **Alyssum**, Tourn. (Crucif.).
ascendens, Koch, in Linnaea, xv. (1841) 257 = A. mutabile.
graeca, Boiss. & Heldr. Diagn. Ser. II. i. 35 = A. mutabile.
incana, DC. Syst. ii. 291 = A. incanum.
mutabilis, DC. l. c. 292 = A. mutabile.
obliqua, DC. l. c. = A. mutabile.
orbiculata, DC. l. c. 293 = A. orbiculatum.
peruviana, DC. l. c.—Peruv.
Potanini, Maxim. in Bull. Acad. Pétersb. xxvi. (1880) 422.—Mongolia.
procumbens, Portenschl. Enum. Pl. Dalm. 15 = A. mutabile.
spathulata, C. A. Mey. in Ledeb. Fl. Alt. iii. 48 = A. spathulatum.
stricta, Boiss. Diagn. Ser. II. i. 35 = A. orbiculatum.
thessala, Heldr. ex Nym. Consp. 51 = A. orbiculatum.
viridis, Tausch, in Syll. Ratisb. ii. (1828) 245 = A. incanum.

BERTHELOTIA, DC. Prod. v. 375 (1836) = **Pluchea**, Cass. (Compos.).
lanceolata, DC. l. c. 376 = P. lanceolata.

BERTHIERA, Vent. Tabl. ii. 572 (1799) = **Bertiera**, Aubl. (Rubiac.).

BERTHOLETIA, Reichb. Nom. 91 (1841) = BERTHELOTIA, DC. = **Pluchea**, Cass. (Compos.).

BERTHOLETIA, Brongn. Enum. Gen. 123 (1843) = **Bertholletia**, Humb. & Bonpl. (Myrtac.).

BERTHOLLETIA, Humb. & Bonpl. Pl. Aequin. i. 122. t. 36 (1808). *MYRTACEAE*, Benth. & Hook. f. i. 723.
TONCA, Rich. Anal. Fruit. 84 (1811).
excelsa, Berg, in Mart. Fl. Bras. xiv. 1. 478. t. 60, 61 = nobilis.
excelsa, *Humb. & Bonpl. Pl. Aequin.* i. 122. t. 36.—Bras.
nobilis, *Miers, in Trans. Linn. Soc.* xxx. (1874) 197.—Guiana.

BERTIERA, Aubl. Pl. Gui. i. 180. t. 69 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 77.
POMATIUM, Gaertn. f. Fruct. iii. 252. t. 225 (1805).
ZALUZANIA, Comm. ex Gaertn. f. Fruct. iii. 74. t. 192 (1805).
aethiopica, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 83.—Afr. trop.
africana, *Rich. in Mém. Soc. Hist. Nat. Par.* v. (1830) 255.—Afr. trop.
angustifolia, *Benth. Bot. Voy. Sulph.* 103.—Ins. Cocos.
bistipulata, Boj. Hort. Maurit. 166 = B. Zaluzania.
borbonica, Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 254 = B. Zaluzania.
bracteolata, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 84.—Afr. trop.
breviflora, *Hiern, l. c.* 85.—Afr. trop.
chrysantha, *Hassk. Retzia*, i. 30.—Java.
coccinea, G. Don, Gen. Syst. iii. 506 = Mussaenda elegans.
ferruginea, Willd. ex Roem. & Schult. Syst. v. 227 = Psychotria rufescens.
gonzaleoides, *Griseb. Cat. Pl. Cub.* 124.—Cuba.
guianensis, *Aubl. Pl. Gui.* i. 180. t. 69.—Guiana.
lateriflora, *Blume, Bijdr.* 987.—Java.
laxa, *Benth. in Hook. Niger Fl.* 394.—Afr. trop.
macrocalyx, *Hassk. in Nederl. Kruidk. Arch.* iv. (1859) 7.—Malaya.
macrocarpa, *Benth. in Hook. Niger Fl.* 394.—Afr. trop.
montana, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 83.—Afr. trop.
mucronata, *Comm. ex Gaertn. f. Fruct.* iii. 74. t. 192.—Hab.?
palustris, *Rich. in Mém. Soc. Hist. Nat. Par.* v. (1830) 254.—Guiana.
Pomatium, *Benth. in Hook. Niger Fl.* 394.—Afr. trop.
rufa, Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 254 = B. Zaluzania.
subsessilis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 83.—Afr. trop.
Zaluzania, *Comm. ex Gaertn. f. Fruct.* iii. 74. t. 192.—Ins. Mascar.

BERTIERA, Blume, Bijdr. 987 (1826) = **Adenosacme**, Wall. (Rubiac.).
fasciculata, Blume, l. c. = A. longifolia.
javanica, Blume, l. c. = A. longifolia.
lateriflora, Blume, l. c.—Java.

BERTOLONIA, DC. in Ann. Mus. Par. xix. (1812) t. 5 (14) = **Leuchèria**, Lag. (Compos.).
purpurea, DC. l. c. = L. purpurea.

BERTOLONIA, Raddi, in Mem. Mod. xviii. Fis. 384 (1820). *MELASTOMACEAE*, Benth. & Hook. f. i. 755.
TRIBLEMMA, R. Br. ex DC. Prod. iii. 113 (1828).
acaulis, Cham. in Linnaea, ix. (1834) 383 = Eriocnema Hilariana.
acuminata, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 344.—Bras.
aenea, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 318 = marmorata.
guttata, Hook. f. Bot. Mag. t. 5524 = Gravesia guttata.
hirsuta, Benth. Bot. Voy. Sulph. 94 = Triolaena hirsuta.
Houtteana, *Van Houtte, in Fl. des Serres*, Sér. II. x. (1874) 109.—Hab.?
leuzeana, *DC. Prod.* iii. 113.—Bras.
longifolia, *Cham. in Linnaea*, ix. (1834) 384 = Salpinga longifolia.

BERTOLONIA:—
maculata, *DC. Prod.* iii. 114.—Bras.
margaritacea, Hort. Bull. ex Naud. in Fl. des Serres, xvi. (1865–67) 165 = Salpinga margaritacea.
marmorata, *Naud. in Ann. Sc. Nat.* Sér. III. xv. (1851) 318.—Bras.
Mirandaei ×, *Van Houtte, in Fl. des Serres*, xxi. (1875) 135.
nymphaeaefolia, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 384.—Bras.
ovata, *DC. Prod.* iii. 113.—Bras.
primulaeflora, Hort. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 80 = Monolena primulaeflora.
scabra, Rafin. ex J. C. Schau. in DC. Prod. xi. 584 = Lippia lanceolata.
scorpioides, Baill. Hist. Pl. vii. 11 = Triolena scorpioides.
superbissima, *Hort.—Cf. Gard. Chron.* (1873) 5.—Hab.?

BERTOLONIA, Rafin. in Am. Monthly Mag. (1818) 267 = **Lippia**, Linn. (Verben.).

BERTOLONIA, Moç. & Sessé, ex DC. Prod. ii. 589 (1825) = **Cereocarpus**, H. B. & K. (Rosac.).
guieroides, Moç. & Sessé, l. c. = C. fothergilloides.

BERTOLONIA, Spinola, Cat. Jard. St. Sebast. (1809) 24 = **Myoporum**, Banks.
glandulosa, Spinola, l. c. = M. viscosum.

BERTOLONIA, Spreng. Neue Entdeck. ii. 110. t. 1. f. 1 (1821) = **Chrysochlamys**, Poepp. (Guttif.).
paniculata, Spreng. l. c.—Bras.

BERTUCHIA, Dennst. Schluess. Hort. Malab. 30 (1818) = **Gardenia**, Ellis (Rubiac.).
speciosa, Dennst. l. c.—Ind. or.

BERTYA, Planch. in Hook. Lond. Journ. iv. (1845) 472. t. 16. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 264.
Cunninghamii, *Planch. l. c.* 473.—Austral.
dimerostigma, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 98.—Austral.
Findlayi, *F. Muell. Fragm.* viii. 141.—Austral.
gummifera, *Planch. in Hook. Lond. Journ. Bot.* iv. (1845) 473. t. 16.—Austral.
Mitchellii, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 63.—Austral.
oblongifolia, Muell. Arg. in Flora, xlvii. (1864) 471 = pomaderrioides.
oleaefolia, *Planch. in Hook. Lond. Journ. Bot.* iv. (1845) 473.—Austral.
oppositifolia, *F. Muell. & O'Shanesy, in Wing, South Sc. Record*, ii. (1882) 98.—Austral.
pedicellata, *F. Muell. Fragm.* iv. 143.—Austral.
pinifolia, *Planch. in Hook. Lond. Journ. Bot.* iv. (1845) 473.—Austral.
polymorpha, *Baill. Adansonia*, vi. (1865–66) 298.—N. Zel.
pomaderrioides, *F. Muell. Fragm.* iv. 34.—Austral.
psiloclada, Muell. Arg. in Flora, xlvii. (1864) 471 = Ricinocarpus psilocladus.
quadrisepala, *F. Muell. Fragm.* x. 52.—Austral.
rosmarinifolia, *Planch. in Hook. Lond. Journ. Bot.* iv. (1845) 473.—Austral.
rotundifolia, *F. Muell. Fragm.* iv. 34.—Austral.
tasmanica, Muell. Arg. in Linnaea, xxxiv. (1865–66) 63 = rosmarinifolia.

BERULA, Hoffm. ex Bess. Enum. Pl. Volh. 44 (1822) = **Sium**, Tourn. (Umbellif.).
angustifolia, Mert. & Koch, i Roehl. Deutschl. Fl. ii. 433 = S. erectum.
lancifolia, Bess. Enum. Pl. Volh. 44 = S. lancifolium.

BERYLLIS, Salisb. Gen. Pl. Fragm. 33 (1866) = **Ornithogalum**, Tourn. (Liliac.).
pyrenaica, Salisb. l. c. = O. pyrenaicum.
stachyoides, Salisb. l. c. = O. narbonense?

BERZELIA, Brongn. in Ann. Sc. Nat. Sér. I. viii (1826) 370. t. 35. *BRUNIACEAE*, Benth. & Hook. f. i. 671.
HETERODON, Meissn. Gen. 72 (1837).
RABENHORSTIA, Reichb. Nom. 159 (1841).
abrotanoides, *Brongn. l. c.* 371.—Afr. austr.

BERZELIA:—
alopecurioides, *Sond. in Harv. & Sond. Fl. Cap.* ii. 310.—Afr. austr.
arachnoidea, *Eckl. & Zeyh. Enum.* 138.—Afr. austr.
brevifolia, Eckl. & Zeyh. l. c. = abrotanoides.
callunoides, *Oliver, in Journ. Linn. Soc.* ix. (1867) 333.—Afr. austr.
commutata, *Sond. in Harv. & Sond. Fl. Cap.* ii. 310.—Afr. austr.
comosa, Eckl. & Zeyh. Enum. 137 = commutata.
cordifolia, *Schlecht. in Linnaea*, vi. (1831) 189.—Afr. austr.
ericoides, Eckl. & Zeyh. Enum. 137 = intermedia.
formosa, Eckl. & Zeyh. l. c. 138 = abrotanoides.
globosa, G. Don, Gen. Syst. ii. 46 (= *Berardia globosa*, Sond.).—Afr. austr.
intermedia, *Schlecht. in Linnaea*, vi. (1831) 188.—Afr. austr.
lanuginosa, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 372. t. 35.—Afr. austr.
rubra, Schlecht. in Linnaea, vi. (1831) 189 = squarrosa.
squarrosa, *Sond. in Harv. & Sond. Fl. Cap.* ii. 312.—Afr. austr.
superba, *Eckl. & Zeyh. Enum.* 138.—Afr. austr.
Wendlandiana, Eckl. & Zeyh. l. c. 137 = intermedia.

BERZELIA, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 292 = **Hermbstaedtia**, Reichb. (Amarant.).
glauca, Mart. l. c. = H. glauca.

BESCHORNERIA, Kunth, Enum. Pl. v. 844 (1850). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 738.
bracteata, *Jacobi, Index*, 12; *Baker, in Bot. Mag.* t. 6641.—Mexic.
Decosteriana, Baker, in Bot. Mag. t. 6768 = seq.
Dekosteriana, *C. Koch, Wochenschr.* vii. (1864) 187.—Mexic.
dubia, *Carr. in Rev. Hortic.* (1877) 155.—Texas.
floribunda, *Hort. ex C. Koch, Wochenschr.* v. (1862) 199.—Hab.?
Galleottii, *Jacobi, Index*, 12, nomen.—Mexic.?
multiflora, *Hort. ex C. Koch, Wochenschr.* iii. (1860) 63.—Hab.?
Parmentieri, *Jacobi, in Hamb. Gartenz.* xx. (1864) 503.—Hab.?
pumila, *Jacobi, Index*, 12, nomen.—Hab.?
Toneliana, Jacobi, l. c. = seq.
Tonelii, *Jacobi, in Hamb. Gartenz.* xx. (1864) 503, nomen.—Mexic.
Schlechtendalii, *Jacobi, Index*, 12, nomen.—Hab.?
tubiflora, *Kunth, Enum. Pl.* v. 844.—Mexic.
Verlindeniana, *Jacobi, Index*, 12, nomen.—Hab.?
yuccoides, Hort. ex Hook. Bot. Mag. t. 5203 = seq.
yuccoides, *C. Koch, Wochenschr.* ii. (1859) 337; iii. (186) 63; vi. (1864) 186.—Mexic.

BESENNA, A. Rich. Tent. Fl. Abyss. i. 253 (1847) = **Albizzia**, Durr. (Legumin.).
anthelmintica, A. Rich. l. c. = A. anthelmintica.

BESHA, D. Dietr. Syn. i. 139 (1839) = **Ochlandra**, Thw. (Gramin.).

BESLERIA, Plum. ex Linn. Gen. ed. I. 181 (1737). *GESNERACEAE*, Benth & Hook. f. ii. 1015.
CRANTZIA, Neck. Elem. i. 366 (1790).
CREMOSPERMA, Benth. Pl. Hartw. 234 (1846).
CYRTANTHEMUM, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 130.
DRUPINA, Linn. Pl. Surin. 11 (1775); Amoen. Acad. viii. 259 (1785).
ERIPHIA, P. Br. Hist. Jamaic. 270 (1756).
FRIMBROLINA, Rafin. Sylva Tellur. 71 (1838).
GASTERANTHOPSIS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 129.
GASTERANTHUS, Benth. Pl. Hartw. 233 (1846).
HEMATOPHYLA, Rafin. Sylva Tellur. 71 (1838).
PARABESLERIA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 126.
PSEUDOBESLERIA, Oerst. l. c. 128 (1861).
acutifolia, *Benth. Pl. Hartw.* 237.—N. Granat.
aggregata, *Hanst. in Mart. Fl. Bras.* viii. 400.—Bras.
ampla, *Vell. Fl. Flum.* 262; vi. t. 84.—Bras.
arborea, *Vell. ll. cc.* 261; vi. t. 82.—Bras.
ardens, *Decne. ex Linden, Cat.* n. 5 (1850) 1.—Venezuela.

BESLERIA :—
barbata, *Hanst. in Linnaea*, xxxiv. (1865–66) 327.—Peruv.
barbensis, *Hanst. l. c.* 319.—Costa Rica.
Berteriana, *DC. Prod.* vii. 538.—Jamaica.
bicolor, *H. B. & K. Nov. Gen. et Sp.* ii. 398.—N. Granat.
bicolor, Schott, in Oestr. Med. Jahrb. vi. II. 65 = Alloplectus bicolor.
biflora, Forst. f. Fl. Ins. Austr. 43. n. 236 = Cyrtandra biflora.
bivalvis, Linn. f. Suppl. 280 (= *Picria surinamensis*).—Guiana.
bonodora, Vell. Fl. Flum. 261; vi. t. 80 = Brunfelsia latifolia.
caerulea, Aubl. Pl. Gui. ii. 631 = violaeea.
calearata, *H. B. & K. Nov. Gen. et Sp.* ii. 399.—N. Granat.
campanulata, *Linden, Cat.* n. 5. (1850) 2.—Venezuela.
capitata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 2.—Peruv.
coccinea, Aubl. Pl. Gui. ii. 632 = Alloplectus coccineus.
columneoides, *Hanst. in Linnaea*, xxxiv. (1865–66) 322.—Costa Rica.
costarieensis, *Hanst. l. c.* 330.—Costa Rica.
crassifolia, *Schott, Neu. Bras. Pfl.* i. 7.—Bras.
cristata, Linn. Sp. Pl. 619 = Alloplectus cristatus.
cristata, Moç. & Sesse, ex DC. Prod. vii. 546, 540 = Alloplectus glaber.
cuneata, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 543.—Bras.
cymosa, Forst. f. Fl. Ins. Austr. 44. n. 237 = Cyrtandra cymosa.
Cyrtanthemum, *Hanst. in Linnaea*, xxxiv. (1865–66) 321.—Mexic.
deflexa, *Hanst. l. c.*—Mexic.
dichrus, Spreng. Syst. ii. 840 = Alloplectus dichrous.
divaricata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 2.—Peruv.
Drymonia, Steud. Nom. ed. II. i. 200 = Drymonia calearata.
elegans, *H. B. & K. Nov. Gen. et Sp.* ii. 397.—N. Granat.
fasciculata, *Wawra, in Oestr. Bot. Zeitschr.* xiii. (1863) 142.—Bras.
flavovirens, *Mart. in Nov. Act. Nat. Cur.* xi. (1823) 49.—Bras.
geminiflora, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 543.—Bras.
gibbosa, *Hanst. in Mart. Fl. Bras.* viii. 421.—Bras.
glabra, *Hanst. in Linnaea*, xxxiv. (1865–66) 325.—Mexic.
grandiflora, *H. B. & K. Nov. Gen. et Sp.* ii. 401.—N. Granat.
grandifolia, *Schott, Neu. Bras. Pfl.* i. 5. n. 5.—Bras.
guadalupensis, *DC. Prod.* vii. 538.—Guadalup.
hirsuta, *Hanst. in Linnaea*, xxxiv. (1865–66) 326.—Mexic.
hirtella, Schott, in Oestr. Med. Jahrb. vi. II. 66 = Alloplectus sparsiflorus.
hispida, H. B. & K. Nov. Gen. et Sp. ii. 39 = Alloplectus hispidus.
Imray, *Hook. Bot. Mag.* t. 6341.—Ind. occ.
incarnata, *Aubl. Pl. Gui.* ii. 635.—Guiana.
inodora, Vell. Fl. Flum. 261; vi. t. 81 = Brunfelsia paueiflora.
insignis, Mart. & Gal. in Bull. Acad. Brux. ix. (1842) II. 37 = Solenophora insignis.
labiosa, *Hanst. in Linnaea*, xxxiv. (1865–66) 324.—N. Granat.
laxiflora, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 361.—Guiana.
leucostoma, *Hanst. in Linnaea*, xxxiv. (1865–66) 326.—N. Granat.
lueida, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 2.—Peruv.
lutea, *Linn. Sp. Pl.* 619.—Ind. occ.
lutea, Spreng. Syst. ii. 841 = Berteriana.
luteo-virens, Mart. Nov. Gen. iii. 45 = flavovirens.
melittifolia, Linn. Sp. Pl. 619 = Episcia melittifolia.
mollis, *Hook. ex Steud. Nom.* ed. II. i. 200.—Am. austr.
mollissima, Spreng. ex Steud. Nom. ed. I. 107 = praec.?
mueronata, *Hanst. in Linnaea*, xxxiv. (1865–66) 331.—N. Granat.
oncogastra, *Hanst. l. c.* 335.—Ecuador.
Patrisii, *DC. Prod.* vii. 538.—Guiana.
pendula, *Hanst. in Linnaea*, xxxiv. (1865–66) 333.—N. Granat.

BESLERIA :—
pentaphylla, *Vell. Fl. Flum.* 262; vi. t. 83.—Bras.
Princeps, *Hanst. in Linnaea*, xxxiv. (1865–66) 317.—Costa Rica.
pulchella, Donn, Hort. Cantab. 140, ex Sims, Bot. Mag. t. 1146 = Tussaeia pulehella.
quitensis, *Hanst. in Linnaea*, xxxiv. (1865–66) 334.—Ecuador.
Rhytidophyllum, *Hanst. l. c.* 334.—N. Granat.
Riedeliana, *Hanst. in Mart. Fl. Bras.* viii. 399.—Bras.
sanguinea, Pers. Syn. ii. 165 = Alloplectus sanguineus.
scandens, Moç. & Sesse, ex D. Don, in Trans. Linn. Soc. xv. (1827) 349 = Maurandia erubescens?
Selloana, *Klotzsch & Hanst. in Mart. Fl. Bras.* viii. 398.—Bras.
serrulata, *Jacq. Hort. Schoenb.* iii. 21.—Ind. occ.
solanoides, *H. B. & K. Nov. Gen. et Sp.* ii. 398.—Am. austr.
spectabilis, H. B. & K. l. c. 400 = Drymonia speetabilis.
splendens, Hort. ex Steud. Nom. ed. II. i. 200 = Drymonia speetabilis.
surinamensis, *Miq. in Linnaea*, xxii. (1849) 471.—Guiana.
Symphytum, *Klotzsch & Hanst. ex Hanst. in Mart. Fl. Bras.* viii. 399.—Bras.
tetrangularis, *Ruiz, ex Hanst. in Linnaea*, xxxiv. (1865–66) 331.—Peruv.
triflora, *Hanst. l. c.* 329.—Costa Rica.
umbellata, Herb. Banks, ex Sims, Bot. Mag. t. 1146 = Episeia pulehella.
umbrosa, *Mart. Nov. Gen.* iii. 44. t. 218.—Bras.
violacea, *Aubl. Pl. Gui.* ii. 630. t. 254.—Guiana.
Wendlandiana, *Hanst. in Linnaea*, xxxiv. (1865–66) 318.—Costa Rica.

BESSERA, Sehult. f. in Linnaea, iv. (1829) 121. *LILIACEAE*, Benth. & Hook. f. iii. 801.
PHARIUM, Herb. in Bot. Reg. t. 1546 (1832).
elegans, *Schult. f. l. c.*—Mexic.
fistulosa, Pritz. Ie. Ind. 147 = elegans.
Herberti, G. Don, in Sweet, Hort. Brit. ed. III. 694 = elegans.
miniata, Lem. in Fl. des Serres, Sér. I. iv. (1848) t. 424 = elegans.
multiflora, Mart. & Gal. in Bull. Acad. Bras. ix. (1842) (reimpr. 14) = elegans.

BESSERA, Schult. Obs. Bot. 27 (1809) = **Pulmonaria**, Tourn. (Borag.).
azurea, Schult. l. e. = P. azurea.

BESSERA, Spreng. Pugill. ii. 90 (1815) = **Xylosma**, Forst. (Bixin.).
inermis, Spreng. l. c. = Flueggia microcarpa.
spinosa, Spreng. l. c. 91 = X. nitidum.

BESSERA, Vell. Fl. Flum. 147 (1825); iv. t. 2 = **Pisonia**, Linn. (Nyetagin.).
calycantha, Vell. ll. ee. = P. Olfersiana.

BESSIA, Rafin. Sylva Tellur. 13 (1838) = **Corypha**, Linn. (Palm.).
sanguinolenta, Rafin. l. c. = C. umbraculifera.

BESTRAM, Adans. Fam. ii. 354 (1763) = **Antidesma**, Linn. (Euphorb.).

BETA, [Tourn.] Linn. Syst. ed. I (1735). *CHENOPODIACEAE*, Benth. & Hook. f. i. 52.
alba, DC. Fl. Fr. iii. 383 = vulgaris.
altissima, Hort. ex Steud. Nom. ed. II. i. 200 = vulgaris.
bengalensis, Roxb. Hort. Beng. 21; Fl. Ind. ii. 59 = vulgaris.
Bourgaei, *Coss. Notes Crit.* 44.—Hispan.
campanulata, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 66.—Maroeco.
carnulosa, *Gren. in Gren. & Godr. Fl. Fr.* iii. 16.—Gallia.
chilensis, *Hort.—Cf. Gard. Chron.* (1870) I. 449.—Chili.
Cicla, Georgi, Beschr. Russ. Reich. iii. IV. 848 = trigyna.
Cicla, Linn. Syst. ed. XII. 195 = vulgaris.
crispa, Tratt. Arch. vii. 117 = vulgaris.
Cycla, Pall. Tabl. Taur. 48 = trigyna.
decumbens, Moench, Meth. 332 = vulgaris.

BETA :—
diffusa, *Coss. Notes Crit.* 178.—Hispan.
esculenta, Salisb. Prod. 152 = vulgaris.
foliosa, Ehrenb. ex Steud. Nom. ed. II. i. 200 = vulgaris.
hastata, Desf. Cat. Hort. Par. ed. III. (1829) 389 = procumbens.
hortensis, Mill. Gard. Dict. ed. VIII. n. 2 = vulgaris.
hybrida, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 124, nomen.—Rossia.
incarnata, Hort. ex Steud. Nom. ed. II. i. 200 = vulgaris.
intermedia, *Bunge, ex Boiss. Fl. Orient.* iv. 900.—Anatolia.
lomatogona, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1838) 360.—As. Min.; Syria.
longespicata, Moq. Chenop. Enum. 13 = lomatogona.
lutea, Hort. ex Steud. Nom. ed. II. i. 200 = vulgaris.
macrocarpa, Guss. Fl. Sieul. Prod. i. 302 = vulgaris.
macrorhiza, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 257.—Reg. Caucas.; Persia.
marina, Crantz, Inst. i. 120 = vulgaris.
maritima, Linn. Sp. Pl. ed. II. 322 = vulgaris.
nana, *Boiss. & Heldr. Diagn.* Ser. I. vii. 82.—Graeeia.
Noëana, Bunge, ex Boiss. Fl. Orient. iv. 899 = vulgaris.
orientalis, Roth, Nov. Pl. Sp. 181 = vulgaris.
patellaris, *Moq. in DC. Prod.* iii. II. 57.—Ins. Tenerif.
patula, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 315.—Ins. Madera.
procumbens, Lange, in Willk. & Lange, Prod. Fl. Hisp. i. 274 = diffusa.
procumbens, *Chr. Sm. in Hornem. Hort. Hafn. Suppl.* 31.—Ins. Canar.
pumila, Link, in Bueh, Besehr. Canar. Ins. 141 = Webbiana.
purpurea, *Hort. ex Steud. Nom.* ed. II. i. 200 = vulgaris.
Rapa, Dum. Fl. Belg. 21 = vulgaris.
rapacea, Hegetsehw. Fl. Sehw. 256 = vulgaris.
rosea, Hort. ex Steud. Nom. ed. II. i. 200 = vulgaris.
rubra, *Delile, Fl. Aeg. Illustr.* 57.—Aegypt.
rubra, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 8.—Malaya.
sativa, Bernh. Syst. Verz. Erf. 162 = vulgaris.
stricta, C. Koch, in Linnaea, xxii. (1849) 180 = vulgaris.
sulcata, Gasp. in Guss. Fl. Sieul. Syn. ii. 800 = vulgaris.
sylvestris, Hort. Panorm. ex Webb & Berth. Phyt. Canar. iii. 197 = procumbens.
triflora, Salisb. Prod. 152 = vulgaris.
trigyna, *Waldst. & Kit. Pl. Rar. Hung.* 34. t. 35.—Hungar.; As. Min.
vulgaris, *Linn. Sp. Pl.* 222.—Europ.; Afr. bor.; Oriens.
Webbiana, *Moq. Chenop. Enum.* 16.—Ins. Canar.

BETCKEA, DC. Mém. Valer. 18; et Prod. iv. 642 (1829) = **Plectritis**, DC. (Rubiae.).
caucasica, Boiss. Fl. Orient. iii. 94.—Reg. Caueas.
Gilliesii, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 366.—Reg. Argent.
heterophylla, Phil. in Linnaea, xxviii. (1856) 704.—Chili.
major, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 39 = P. samolifolia.
samolifolia, DC. Prod. iv. 642 = P. samolifolia.
samolifolia, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 366 = Gilliesii.

BETELA, Rafin. Sylva Tellur. 85 (1838) = **Piper**, Linn.
mastica, Rafin. l. e. = P. Betle.

BETENCOURTIA, St. Hil. Voy. i. 376 (1833) = **Galactia**, P. Br. (Legumin.).
rhynchosioides, St. Hil. l. e. = G. angustifolia.

BETHENCOURTIA, Choisy, in Bueh, Beschr. Canar. Ins. 148 (1825). = **Senecio**, Tourn. (Compos.).
palmensis, Choisy, l. e. in obs. = S. palmensis.

BETONICA, [Tourn.] Linn. Syst. ed. I (1735) = **Stachys**, Tourn. (Labiat.).
affinis, Wender. in Flora, ix. (1826) 353 = S. Betonica.
algeriensis, Noe, in Bull. Soc. Bot. Fr. ii. (1856) 582 = S. Betonica.
Alopecuros, Linn. Sp. Pl. 573 = S. Alopecuros.

BETONICA :—
alpestris, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 93 = Stachys densiflora.
alpigena, Schur, Enum. Pl. Transs. 540.—Hungaria.
alpina, Mill. Gard. Dict. ed. VIII. n. 3 = S. Alopecuros.
angustifolia, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 97 = S. Betonica.
annua, Linn. Sp. Pl. 573; Moench, Meth. 395 = S. annua.
brachystachya, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 98 = S. Betonica.
Brantii, Boiss. Fl. Orient. iv. 750 = S. Brantii.
capensis, Burm. f. Fl. Cap. Prod. 16.—Afr. austr.
Clementei, Perez-Lara, in Anal. Soc. Esp. Hist. Nat. xi. (1882) 400.—Hispan.
danica, Mill. Gard. Dict. ed. VIII. n. 2 = S. densiflora.
decumbens, Moench, Meth. 396 = S. recta.
densiflora, Schur, Enum. Pl. Transs. 890, Index = S. densiflora.
divulsa, Tenore, Syll. Fl. Neapol. App. 614 = S. Betonica.
drymophila, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 100 = S. Betonica.
elongata, Willd. ex Steud. Nom. ed. II. i. 200 = S. Betonica.
flava, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 120 = S. Alopecuros.
foliosa, Presl, Fl. Sicul. 37 = S. Betonica.
foliosa, Rupr. Sert. Tiansch. 66.—As. centr.
frutescens, Bontekoe, Jav. 146 = Justicia Betonica.
glabrata, C. Koch, in Linnaea, xxi. (1848) 684 = S. Betonica.
graeca, Boiss. & Sprun. Diagn. Ser. I. v. 27.—Graecia.
grandiflora, Thuill. Fl. Par. ed. II. 293 = S. longifolia.
grandiflora, Steph. ex Willd. Sp. Pl. iii. 96 = S. grandiflora.
grandifolia, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 101 = S. Betonica.
heraclea, Linn. Mant. i. 83 = S. integrifolia.
hirsuta, Linn. l. c. ii. 248 = S. densiflora.
hirsuta, C. A. Mey. Verz. Pfl. Cauc. 95 = S. longifolia.
hirta, Leyss. Fl. Halens. 109 = S. Betonica.
hirta, Linn. Cent. ii. 20; Gouan, Hort. Monsp. 276 = S. recta.
hylebium, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 101 = S. Betonica.
incana, Mill. Gard. Dict. ed. VIII. n. 5 = S. densiflora.
Jacquini, Gren. & Godr. Fl. Fr. ii. 694 = S. Alopecuros.
laevigata, D. Don, Prod. Fl. Nep. 110 = Nepeta spicata.
laxata, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 96 = S. Betonica.
legitima, Link, Handb. 476 = S. Betonica.
leiocalyx, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 95 = S. Betonica.
lutea, Lam. Fl. Fr. ii. 404 = S. Alopecuros.
macrantha, C. Koch, in Linnaea, xxi. (1848) 683 = S. grandiflora.
macrostachys, Wender. in Flora, ix. (1826) 353 = S. densiflora.
Monnieri, Gouan, Illustr. 36 = S. densiflora.
montana, Lej. Fl. Spa, ii. 24 = S. Betonica.
monticola, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 95 = S. Betonica.
nemorosa, Jord. & Fourr. l. c. 98 = S. Betonica.
nivea, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 266 = S. nivea.
nutans, Kit. ex Schult. Oestr. Fl. ed. II. ii. 146 = S. Betonica.
occitana, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 97 = S. Betonica.
officinalis, Linn. Sp. Pl. 573 = S. Betonica.
orientalis, Linn. l. c. = S. longifolia.
orientalis, Thuill. Fl. Par. 293 = S. grandiflora.
Orphanidea, Heldr. ex Boiss. Fl. Orient. iv. 750 = S. Alopecuros.
parvula, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 94 = S. Betonica.
polyclada, Jord. & Fourr. l. c. 102 = S. Betonica.
pradica, Zanted. ex Nym. Consp. 577 = S. densiflora.
pratensis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 96 = S. Betonica.
psilostachys, Jord. & Fourr. l. c. 97 = S. Betonica.
purpurea, Gilib. Fl. Lituan. i. 82 = S. Betonica.
pyrenaica, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 94 = S. Betonica.

BETONICA :—
recurva, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 102 = Stachys Betonica.
recurvidens, Peterm. Fl. Lips. Excurs. 444 = S. Betonica.
rigida, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 103 = S. Betonica.
rosea, Hort. ex Heynh. Nom. ii. 67 = S. grandiflora.
rubicunda, Wender. Ind. Sem. Hort. Marb. ex Benth. in DC. Prod. xii. 460 = S. densiflora.
rusticana, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 96 = S. Betonica.
sabauda, Jord. & Fourr. l. c. 93 = S. Betonica.
scardica, Griseb. Spicil. Fl. Rumel. ii. 136 = S. graeca.
serotina, Host, Fl. Austr. ii. 165 = S. Betonica.
stricta, [Soland. in] Ait. Hort. Kew. ed. I. ii. 299 = S. Betonica.
stricticaulis, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 99 = S. Betonica.
subcarnea, Jord. & Fourr. l. c. 98 = S. Betonica.
sylvulicola, Jord. & Fourr. l. c. 100 = S. Betonica.
valdepubens, Jord. & Fourr. l. c. 103 = S. Betonica.
validula, Jord. & Fourr. l. c. = S. Betonica.
virescens, Jord. & Fourr. l. c. 99 = S. Betonica.
virgultorum, Jord. & Fourr. l. c. 100 = S. Betonica.
vulgaris, Rota, Prosp. Piant. Pav. 272 = S. Betonica.

BETULA, [Tourn.] Linn. Syst. ed. I (1735). *CUPULIFERAE*, Benth. & Hook. f. iii. 404.
BETULASTER, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 198.
acuminata, Ehrh. Beitr. vi. 98 = alba.
acuminata, Wall. Pl. As. Rar. ii. 7. t. 109 = alnoides.
aetnensis, Rafin. Giorn. i. 17; J. & C. Presl, Delic. Prag. 144 = alba.
affinis, Endl. Gen. Suppl. iv. 20 = alnoides.
alba, *Linn. Sp. Pl.* 982.—Europ.; As. bor.; Am. bor.
Alnobetula, Ehrh. Beitr. ii. 72 = Alnus viridis.
alnoides, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 58.—Reg. Himal.
Alnus, Linn. Sp. Pl. 983 = Alnus glutinosa.
Alnus, Thunb. Fl. Jap. 76 = Alnus maritima.
Alnus-crispa, Steud. Nom. ed. II. i. 201 = Alnus crispa.
Alnus-incana, Du Roi, Obs. Bot. 31 = Alnus incana.
Alnus-rugosa, Du Roi, l. c. 32 = Alnus serrulata.
alpestris, *Fries, Summ. Veg. Scand.* i. 212.—Europ. bor.; Groenland.
alpina, Borckh. ex Reichb. Fl. Germ. Excurs. 174 = Alnus viridis.
ambigua, Hampe, ex Reichb. Ic. Fl. Germ. xii. 2 = alba.
angulata, *Hort. ex Steud. Nom.* ed. II. i. 201.—Am. bor.
antarctica, Forst. in Comm. Gott. ix. (1789) 45 = Fagus betuloides.
arguta, Schlecht. in Linnaea, vii. (1832) 139 = Alnus acuminata.
asplenifolia, Hort. ex Regel, in DC. Prod. xvi. II. 167 = alba.
aurata, Borkh. Forstb. i. 378 = alba.
aurea, Steud. Nom. ed. II. i. 201 = alba.
Bhojpattra, Wall. Pl. As. Rar. ii. 7 = utilis.
borealis, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 196 = pumila.
broccembergensis, Bechst. Forstb. 297 = alba.
carpatica, Waldst. & Kit. ex Willd. Sp. Pl. iv. 464 = alba.
canadensis, Hort. ex C. Koch, Dendrol. ii. II. 656 = alba.
carpinifolia, Ehrh. Beitr. vi. 99 = lenta.
carpinifolia, *Sieb. & Zucc. in Abh. Acad. Muench.* iv. *Abth.* III. (1846) 228.—Japon.
chinensis, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) I. 47.—China.
ciliata, Steud. Nom. ed. II. i. 201 = Populus ciliata.
cordata, Loisel. Notul. 139 = Alnus cordifolia.
cordifolia, Regel, in Nouv. Mém. Soc. Nat. Mosc. xiii. (1861) 86 = alba.
coriifolia, Tausch, ex Regel, in DC. Prod. xvi. II. 168 = alba.
corylifolia, *Regel & Maxim. in Bull. Soc. Nat. Mosc.* xxxviii. (1865) II. 417.—Japon.
costata, Trautv. in Maxim. Prim. Fl. Amur. 253 = ulmifolia.
crispa, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 339 = Alnus viridis.
cryptocarpa, Laestad. in Bot. Notiser, 1856 (1857) 109 = alba.

BETULA :—
cuspidata, Schrad. ex Regel, in DC. Prod. xvi. II. 164 = alba.
cycoviensis, Steud. Nom. ed. I. 107, sphalm. = oycoviensis.
cylindrostachya, Wall. Pl. As. Rar. ii. 7 = alnoides.
dalecarlica, Linn. f. Suppl. 416 = alba.
davurica, *Pall. Fl. Ross.* i. 60. t. 39. fig. A.—Sibir.; Am. bor.
dioica, *Pall. Reise*, iii. 321.—Sibir. transbaical.
divaricata, Ledeb. in Denkschr. Baier. Bot. Ges. iii. (1841) 59 = fruticosa.
emarginata, Ehrh. Beitr. iv. 158 = Alnus glutinosa.
Ermani, *Cham. in Linnaea*, vi. (1831) 537. t. 6. fig. D.—As. bor. or.
ethnensis, Rafin. ex Regel, in DC. Prod. xvi. II. 163 = alba.
exaltata, *S. Moore, in Journ. Linn. Soc.* xvii. (1880) 386. t. 16.—China.
excelsa, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 337 = papyracea ?
excelsa, Hook. Fl. Bor. Am. ii. 156 = lenta.
excelsa, Pursh, Fl. Am. Sept. ii. 621 = lutea.
Friesii, Larss. Fl. Verml. ed. II. ex Hartm. Scand. Fl. ed. XIII. 376 = alba.
fruticans, *Pall. Reise*, iii. t. kk = fruticosa.
fruticosa, Fries, in Flora, xxxiii. (1850) 152 = alpestris.
fruticosa, *Pall. Fl. Ross.* i. I. 62. t. 40. f. A. B. C.—Europ.; As. bor.; Sibir.; Manchur.
fruticosa, Willd. Sp. Pl. iv. 467 = humilis.
fusca, Pall. Reise, iii. 173 = fruticosa.
glabra, *Dum. Fl. Belg.* 11.—Belgia.
glandulosa, *Michx. Fl. Am. Bor.* ii. 180.—Am. bor.
glauca, Wender. in Bot. Zeit. iv. (1846) 291 = alba.
glutinosa, Lam. Encyc. i. 454 = Alnus glutinosa.
glutinosa, Wallr. Sched. Crit. 497 = alba.
Gmelini, Bunge, Verz. Suppl. Fl. Alt. 113 = fruticosa.
grandis, Schrad. in Ind. Sem. Hort. Gott. (1833) 2 = alba.
Grayii, Regel, in Bull. Soc. Nat. Mosc. xxxviii. (1865) II. 406 = pumila.
grossa, *Sieb. & Zucc. in Abh. Acad. Muench.* iv. *Abth.* III. (1846) 228.—Japon.
gummifera, Bertol. Fl. Ital. x. 229 = alba.
Hackelii, Opiz, ex Steud. Nom. cd. II. i. 201 = alba.
harcynica, Wender. ex Steud. l. c. = alba.
humilis, Fries, ex Regel, in DC. Prod. xvi. II. 172 = alpestris.
humilis, *Schrank, Bayer. Fl.* i. 420.—Europ. bor.; As. bor.; Am. bor.
hybrida, Blom, in Vet. Acad. Nya Handl. vii. (1786) 186. t. 6. f. 13 = alba.
hybrida, *Regel, in Nouv. Mém. Soc. Nat. Mosc.* xiii. (1861) 94.—Rossia.
incana, Roth, Tent. Fl. Germ. iii. 477 = Alnus incana.
intermedia, *Thomas, ex Reichb. Ic. Fl. Germ.* xii. t. 624. fig. 1283.—Europ.
Jacquemontii, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 189 = utilis.
japonica, Siebold, in Verh. Batav. Gen. xii. (1830) 25 = alba.
kamtschatica, Buek, in DC. Prod. Ind. iv. 52 = humilis.
laciniata, Blom, in Vet. Acad. Nya Handl. vii. (1786) 156. t. 6. f. 13 = alba.
laciniata, Ehrh. Beitr. iii. 21 = Alnus glutinosa.
laciniata, Thunb. Diss. Bet. 10 = alba.
lagopina, Hartm. ex Regel, in DC. Prod. xvi. II. 173 = alpestris.
lanulosa, Michx. Fl. Bor. Am. ii. 181 = nigra.
latifolia, Tausch, in Flora, xxi. (1838) 751 = alba.
lenta, Duroi, Harbk. Baumz. i. 92 = alba.
lenta, *Linn. Sp. Pl.* 983.—Am. bor.
lenta, Regel, in Nouv. Mém. Soc. Nat. Mosc. xiii. (1861) 125 = carpinifolia.
leptophylla, Regel, in DC. Prod. xvi. II. 181 = Alnus nepalensis.
leptostachya, Wall. Cat. n. 2799 = Alnus nepalensis.
Littelliana, Tuckerm. in Am. Journ. Sc. xlv. (1843) 30 = glandulosa.
lobulata, *Kit. in Linnaea*, xxxii. (1863) 351.—Europ.
longifolia, Bové, ex Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 208 = Alnus orientalis.
lutea, *Michx. Hist. Arb.* ii. 153. t. 5.—Am. bor.
Maackii, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 380 = davurica.
major, Gilib. Exercit. ii. 400 = alba.

BETULA:—
macrostachya, Schrad. ex Regel, in DC. Prod. xvi. 187; Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 7 = alba.
Maximowiczziana, Regel, in DC. Prod. xvi. 180 = Maximowiczii.
Maximowiczii, *Regel, in Bull. Soc. Nat. Mosc.* xxxviii. (1865) II. 418.—Japon.
Maximowiczii, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 139 = davurica.
Maximowitschii, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 435 = dahurica.
media, Lacstad. in Bot. Notiser, 1856 (1857) 107 = alba.
megalocarpa, Laestad. l. c. 106 = alba.
Michauxii, *Spach, in Ann. Sc. Nat.* Sér. II. xv. (1841) 195.—Am. bor.
microphylla, *Bunge, Verz. Suppl. Fl. Alt.* 112; *in Mém. Acad. Pétersb.* (1835) 506.—Sibir. altaic.
Middendorffii, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 84).—Manchuria.
Murithii, Gaudin, ex Regel, in DC. Prod. xvi. II. 168 = alba.
myrsinoides, Tausch, in Flora, xi. (1828) 753 = humilis.
nana, Hook. Fl. Bor. Am. ii. 156 = glandulosa.
nana, *Linn. Sp. Pl.* 983.—Reg. bor.
nana, Michx. Fl. Bor. Am. ii. 180 = Michauxii.
nigra, Duham. Arb. ed. Nov. iii. 203. t. 51 = alba.
nigra, Duroi, Harbk. Baumz. i. 93 = lenta.
nigra, *Linn. Sp. Pl.* 982.—Am. bor.
nigra, Murith, Guide Bot. Valais, 55 = intermedia.
nigricans, Wender. in Bot. Zeit. iv. (1846) 291 = alba.
nitida, D. Don, Prod. Fl. Nep. 58 = alnoides.
oblongata, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 338 = Alnus glutinosa.
occidentalis, *Hook. Fl. Bor. Am.* ii. 155.—Am. bor. occ.
odorata, Bechst. Forstb. 294 = alba.
ovalifolia, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 378 = humilis.
ovata, C. Koch, in Linnaea, xxii. (1849) 333 = alba.
ovata, Schrank, Baier. Fl. i. 419 = Alnus viridis.
oycoviensis, Bess. Prim. Fl. Galic. ii. 289 = alba.
oycowiensis, Laestad. ex Regel, in DC. Prod. xvi. II. 171 = intermedia.
oycowiensis, Reichb. Ic. Fl. Germ. xii. t. 622. f. 1281 = humilis.
palmaeformis, Laestad. in Bot. Notiser, 1856 (1857) 105 = alba.
palmata, Borkh. Handb. i. 500 = alba.
palustris, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 376 = humilis.
palustris, Salisb. Prod. 395 = Alnus glutinosa.
papyracea, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 337.—Am. bor.
papyrifera, Michx. Fl. Bor. Am. ii. 180 = papyracea.
pendula, Roth, Tent. Fl. Germ. i. 405 = alba.
persicifolia, Hort. ex C. Koch, Dendrol. ii. II. 641 = lutea.
pinnata, Landmark, in Vet. Acad. Nya Handl. Stockh. (1790) 131. f. 5 = Alnus incana.
pirifolia, Hort. ex C. Koch, Dendrol. ii. II. 646 = papyracea.
pontica, *Desf. ex Steud. Nom.* ed. II. i. 201.—As. Min.
pontica, Hort. ex Loud. Encyc. Pl. 720 = alba.
populifolia, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 336 = alba.
pubescens, Ehrh. Beitr. iv. 160; et vi. 98 = alba.
pumila, *Linn. Mant.* i. 124.—Am. bor.
pumila, J. Vahl, ex Lange, Consp. Fl. Groenl. i. 114 = alpestris?
quebeccensis, Burg. in Ges. Naturf. Fr. Berl. Schr. v. (1784) 196 = humilis.
reticulata, Laestad. ex Regel, in DC. Prod. xvi. II. 169 = alba.
reticulata, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 378 = humilis.
rhombifolia, Tausch, in Flora, xxxviii. (1838) 752 = alba.
rotundifolia, Regel & Tiling, Fl. Ajan. 118 = Middendorffii.
rotundifolia, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 194 = glandulosa.
rubra, Hort. ex Regel, in DC. Prod. xvi. II. 167 = alba.

BETULA:—
rubra, Michx. Hist. Arb. ii. 143. t. 3 = nigra.
rugosa, Ehrh. Beitr. iii. 21 = Alnus serrulata.
rustica, Laestad. in Bot. Notiser, 1856 (1857) 104 = alba.
Schmidtii, *Regel, in Bull. Soc. Nat. Mosc.* xxxviii. (1865) II. 412.—Manchuria.
serrulata, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 338 = Alnus serrulata.
sibirica, Lodd. ex Regel, in DC. Prod. xvi. II. 173 = humilis.
silvatica, Laestad. in Bot. Notiser, 1856 (1857) 108 = alba.
silvestris, Laestad. l. c. 107 = alba.
Socolowii, Jacq. f. ex Regel, in DC. Prod. xvi. II. 173 = humilis.
Sokolowii, Hort. Petrop. ex Regel, l. c. 163 = alba.
subaequalis, Laestad. in Bot. Notiser, 1856 (1857) 109 = alba.
subalpina, Laestad. l. c. 110 = alba.
tianschanica, *Rupr. Sert. Tiansch.* 72.—As. centr.
tiliaefolia, Laestad. in Bot. Notiser, 1856 (1857) 106 = alba.
tomentosa, Retz. & Abel, Beschr. Holzarten, i. 17. t. 15 = alba.
torfacea, Schleich. Cat. Pl. Helv. ed. IV. 9 = alba.
tortuosa, Ledeb. Fl. Ross. iii. 652 = alba.
transsilvanica, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 68.—Hungar.
triphylla, *Thunb. Diss. Betul.* 12.—Hab.?
tristis, *Wormsk. ex Link, Enum. Hort. Berol.* ii. 401.—Kamtschat.
turfosa, Weigel. ex Reichb. Fl. Germ. Excurs. 174 = fruticosa.
ulmifolia, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 228.—Japon.
urticaefolia, Curtois, ex Steud. Nom. ed. II. i. 201 = alba?
urticifolia, *Regel, in Nouv. Mém. Soc. Nat. Mosc.* xiii. (1861) 115.—Suecia (cult.).
utilis, *D. Don, Prod. Fl. Nep.* 58.—Reg. Himal.; Japon.
verrucosa, Ehrh. Beitr. v. 161; vi. 98 = alba.
virgata, Salisb. Prod. 395 = alba.
virgultosa, Fries, ex Regel, in Nouv. Mém. Soc. Nat. Mosc. xiii. (1861) 115 = urticifolia.
viridis, Chaix, in Vill. Hist. Pl. Dauph. iii. 789 = Alnus viridis.

BETULASTER, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 198 = **Betula**, Tourn. (Cupulif.).
acuminata, Spach, l. c. = Betula alnoides.
affinis, Spach, l. c. 199 = Betula alnoides.
cylindrostachya, Spach, l. c. 198 = Betula alnoides.
nitida, Spach, l. c. 200 = Betula alnoides.

BEURERIA, Ehret, Pl. & Pap. Rar. t. 13 (1755) = **Calycanthus**, Linn.

BEURRERIA, Jacq. Enum. Pl. Carib. 2 (1760) = **Bourreria**, P. Br. (Borag.).

BEVERNA, Adans. Fam. ii. 20 (1763) = **Iris**, Tourn.?

BEYERIA, Miq. in Ann. Sc. Nat. Sér. III. i. (1844) 350. t. 15. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 263.
BEYERIOPSIS, Muell. Arg. in Linnaea, xxxiv. (1865) 56.
CALYPTROSTIGMA, Klotzsch, in Lehm. Pl. Preiss. i. 175 (1844).
Backhousii, Hook. f. Fl. Tasm. i. 339 = opaca.
brevifolia, *Baill. Adansonia*, vi. (1865–66) 309.—Austral.
cinerea, *Baill. l. c.*—Austral.
cyanescens, *Benth. Fl. Austral.* vi. 66.—Austral.
cygnorum, *Baill. Adansonia*, vi. (1865–66) 309.—Austral.
Drummondi, *F. Muell. Census*, 18.—Austral.
lasiocarpa, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 59.—Austral.
latifolia, *Baill. Adansonia*, vi. (1865–66) 304.—Austral.
ledifolia, Sond. in Linnaea, xxviii. (1856) 565 = opaca.
lepidopetala, *F. Muell. Fragm.* i. 230.—Austral.
Leschenaultii, Baill. Adansonia, vi. (1865–66) 307 = opaca.

BEYERIA:—
loranthoides, *Baill. Etud. Gén. Euph.* 403.—Austral.
oblongifolia, Sond. in Linnaea, xxviii. (1856) 564 = viscosa.
opaca, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 16.—Austral.
similis, *Baill. Adansonia*, vi. (1865–66) 309.—Austral.
tristigma, *F. Muell. Fragm.* vi. 181.—Austral.
uncinata, *F. Muell. ex Baill. Adansonia*, vi. (1865–66) 306.—Austral.
viscosa, *Miq. in Ann. Sc. Nat.* Sér. III. i. (1844) 350. t. 15.—Austral.

BEYERIOPSIS, Muell. Arg. in Linnaea, xxxiv. (1865) 56 = **Beyeria**, Miq. (Euphorb.).
brevifolia, Muell. Arg. l. c. 58 = Beyeria brevifolia.
cinerea, Muell. Arg. l. c. 57 = Beyeria cinerea.
cyanescens, Muell. Arg. in DC. Prod. xv. II. 200 = Beyeria cyanescens.
cygnorum, Muell. Arg. in Linnaea, xxxiv. (1865) 56 = Beyeria cygnorum.
Drummondi, Muell. Arg. l. c. 58 = Beyeria Drummondi.
lasiocarpa, Muell. Arg. l. c. 59 = Beyeria lasiocarpa.
latifolia, Muell. Arg. l. c. 57 = Beyeria latifolia.
lepidopetala, Muell. Arg. l. c. = Beyeria lepidopetala.
similis, Muell. Arg. l. c. 58 = Beyeria similis.

BEYRICHIA, Cham. & Schlecht. in Linnaea, iii. (1828) 21. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 949.
ACHETARIA, Cham. & Schlecht. in Linnaea, ii. (1827) 566, nomen prius.
ajugoides, Benth. Scroph. Ind. 9, in nota = ocymoides.
floribunda, *Benth. in DC. Prod.* x. 379.—Bras.
ocymoides, *Cham. & Schlecht. in Linnaea*, iii. (1828) 21.—Bras.
scutellarioides, *Benth. Scroph. Ind.* 9, in nota.—Bras.
villosa, *Benth. l. c.*—Bras.

BEYTHEA, Endl. Gen. 1011 (1840) = **Elaeocarpus**, Linn. (Tiliac.).
bifida, Endl. l. c. = E. bifidus.

BEZANILLA, Remy, in C. Gay, Fl. Chil. iv. 109. t. 46 (1849) = **Psilocarphus**, Nutt. (Compos.).
chilensis, Remy, l. c. 110.—Chili.

BHESA, Buch.-Ham. ex Arn. in Edinb. Phil. Journ. xvi. (1834) 315 = **Kurrimia**, Wall. (Celastr.).
Moja, Buch.-Ham. l. c. = K. pulcherrima.
paniculata, Arn. l. c. = K. paniculata.

BIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 189 = **Tragia**, Linn. (Euphorb.).
alienata, F. Didr. in Kjoeb. Vidensk. Meddel. (1857) 131 = T. Sellowiana.
Lessertiana, Baill. Etud. Gén. Euph. 502 = T. Lessertiana.
Lhotzkyana, Klotzsch, in Wiegm. Archiv, vii. (1841) 190 = T. Sellowiana.
Sellowiana, Klotzsch, l. c. = T. Sellowiana.

BIANCAEA, Todaro, Nuov. Gen. 21 (1858) = **Caesalpinia**, Linn. (Legum.).
mimosoides, Todaro, Hort. Bot. Panorm. 3 = C. mimosoides.
Sappan, Todaro, l. c. = C. Sappan.
scandens, Todaro, Nuov. Gen. 22; et l. c. 4. t. 1 = C. sepiaria.

BIARUM, Schott, Melet. i. 17 (1832). *AROIDEAE*. Benth. & Hook. f. iii. 966.
CYLLENIUM, Schott, Gen. Aroid. t. 9 (1858).
HOMAID, Adans. Fam. ii. 470 (1763), nomen prius.
ISCHARUM, Blume, Rumphia, i. 144. t. 29 (1835).
LEPTOPETION, Schott, Gen. Aroid. t. 8 (1858).
STENURUS, Salisb. Gen. Pl. Fragm. t. 5 (1866).
abbreviatum, Schott, Prod. Aroid. 62 = tenuifolium.
alexandrinum, Boiss. Diagn. Ser. I. xiii. 6 = Olivieri.
Anguillarae, Schott, Prod. Aroid. 62 = tenuifolium.
angustatum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1880) 255.—Oriens; Syria.
arundanum, Boiss. & Reut. Pugill. 110 = tenuifolium.

BIARUM :—
Bovei, *Blume, Rumphia*, i. 114. t. 29.—Syria; As. Min.
carduchorum, *Engl. in DC. Monog. Phan.* ii. 575.—As. Min.; Syria.
constrictum, C. Koch, Ind. Sem. Hort. Berol. (1853) 12 = tenuifolium.
eximium, *Engl. in DC. Monog. Phan.* ii. 576.—Cilicia.
Fraasianum, *N. E. Br. in Journ. Linn. Soc.* xviii. (1880) 254.—Oriens.
gramineum, Schott, Melet. i. 17 = tenuifolium.
Haenseleri, *Willk. in Bot. Zeit.* v. (1847) 49. t. 2.—Hispan.
Homeid, *Blume, Rumphia*, i. 115.—Oriens.
Lehmanni, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 327 = Helicophyllum Lehmanni.
longifolium, *Pomel, Nouv. Mat. Fl. Atl.* 391.—Algeria.
macroglossum, *Pomel, l. c.* 390.—Algeria.
numidicum, *Parl. Fl. Ital.* ii. 243.—Numidia.
Olivieri, *Blume, Rumphia*, i. 115.—Aegypt.
Pyrami, *Engl. in DC. Monog. Phan.* ii. 576.—Cilicia; Syria.
rhopalospadix, C. Koch, Ind. Sem. Hort. Berol. (1853) 12 = Spruneri.
rupestre, *Pomel, Nouv. Mat. Fl. Atl.* 391.—Algeria.
Russellianum, *Schott, Prod. Aroid.* 63.—Syria.
Sewerzowi, *Regel, in Act. Hort. Petrop.* vi. (1879) 489.—Caucasus.
Spruneri, *Boiss. Diagn.* Ser. I. xiii. 5.—Graecia.
Spruneri, Schott, in Gen. Aroid. t. 7 = abbreviatum.
tenuifolium, Heldr. ex Nym. Consp. 755 = abbreviatum.
tenuifolium, *Schott, Melet.* i. 17.—Hispan.
Zeleborii, Schott, in Oestr. Bot. Wochenbl. (1857) 245 = tenuifolium.

BIASLIA, Vand. Fl. Lusit. et Bras. Spec. 4. t. 1. f. 2 (1788) = **Mayaca**, Aubl.

BIASOLETTIA, Koch, in Flora, xix. (1836) i. 163; Bertol. Fl. Ital. iii. 191 (1837) = **Chaerophyllum**, Linn. (Umbellif.).
congesta, Nym. Syll. Suppl. 28.—Graecia.
cretica, Nym. Consp. 303 = C. creticum.
divaricata, Nym. Syll. Suppl. 28.—Graecia.
macrocarpa, Nym. Consp. 302.—Graecia.
nodosa, Bertol. Fl. Ital. iii. 191 = C. nodosum.
parnassica, Nym. Syll. Suppl. 27.—Graecia.
pumila, Nym. Consp. 302.—Graecia.
tuberosa, Koch, in Flora, (1836) i. 163 = Carum cylindricum?

BIASOLETTIA, Pohl, ex Baker, in Mart. Fl. Bras. vi. ii. 311 = **Eupatorium**, Linn. (Compos.).
elliptica, Pohl, ex Baker, l. c. = E. trixoides.

BIASOLETTIA, Presl, Rel. Haenk. ii. 141 (1840?) = **Hernandia**, Linn.? (Laurin.).
nymphaeaefolia, Presl, l. c.—Ins. Marian.

BIASSOLETTIA, Endl. Gen. 1004 (1840) = BIASOLETTIA, Presl (= **Hernandia**, Linn.? Laurin.).

BIATHERIUM, Desv. Opusc. 72 (1831) = **Gymnopogon**, Beauv. (Gramin.).
foliosum, Desv. l. c. = G. foliosum.

BICCHIA, Parl. Fl. Ital. iii. 396 (1858) = **Habenaria**, Willd. (Orchid.).
albida, Parl. l. c. 397 = H. albida.

BICHEA, Stokes, Bot. Mat. Med. ii. 564 (1812) = **Cola**, Schott (Stercul.).
solitaria, Stokes, l. c. 565 = C. acuminata.

BICHENIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 236 = **Trichocline**, Cass. (Compos.).
aurea, D. Don, l. c. 237.—Chili.
auriculata, Wedd. Chlor. And. i. 26.—Bolivia.
Berteriana, Steud. Nom. ed. II. i. 202 = *aurea*.
Cineraria, D. Don, in Phil. Mag. (1832) 391 = T. Cineraria.
crenata, Remy, in C. Gay, Fl. Chil. 302.—Chili.
dealbata, D. Don, ex Hook. Comp. Bot. Mag. i. (1835) 104 = T. dealbata.

BICHENIA :—
plicata, D. Don, ex Wedd. Chlor. And. i. 26 = Trichocline plicata.
reptans, Wedd. l. c. 25.—Bolivia.
sinuata, D. Don, Phil. Mag. (1832) 391 = T. incana.

BICORNELLA, Lindl. Bot. Reg. sub t. 1701 (1735). *ORCHIDEAE*, Benth. & Hook. f. Gen. iii. 634.
gracilis, *Lindl. Gen. et Sp. Orch.* 334.—Madag.
longifolia, *Lindl. l. c.* 335.—Madag.
parviflora, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 500.—Madag.

BICORONA, A. DC. Prod. viii. 330 (1844) = **Melodinus**, Forst. (Apocyn.).
phyllireoides, A. DC. l. c. = M. phillyraeoides.

BICUCULLA, Borckh. in Roem. Arch. i. ii. 46 (1797) = **Adlumia**, Rafin. (1808) (Papav.).
fumarioides, Borckh. l. c. = A. cirrhosa.

BICUCULLARIA, Juss. ex Steud. Nom. ed. II. i. 202 (1840) = **Dicentra**, Bernh. (Papav.).
canadensis, Juss. ex Steud. l. c. 201 = D. Cucullaria.

BIDARIA, Decne. in DC. Prod. viii. 623 (1844) = **Gymnema**, R. Br. (Asclep.).
elegans, Dalz. & Gibs. Bomb. Fl. 151 = G. montanum.
elegans, Decne. in DC. Prod. viii. 623 = G. elegans.
erecta, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 59 = G. stenophyllum.
inodora, Decne. in DC. Prod. viii. 624 (Sp. dub.).—Ind. or.
leptophylla, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 60 = Marsdenia leptophylla.
pergularioides, Thw. Enum. Pl. Zeyl. 198 = G. pergularioides.
pubiflora, Miq. Fl. Ind. Bat. ii. 497.—Java.
syringaefolia, Decne. in DC. Prod. viii. 623 = Marsdenia syringaefolia.
tingens, Decne. l. c. = G. montanum.
trinervis, Decne. l. c. 624 = G. trinerve.

BIDENS, [Tourn.] Linn. Gen. ed. I. 248 (1737) = *COMPOSITAE*, Benth. & Hook. f. ii. 387.
CERATOCEPHALUS, Vaill. ex Cass. in Dict. Sc. Nat. vii. 432 (1817).
DELUCIA, DC. Prod. v. 633 (1836).
EDWARSIA, Neck. Elem. i. 87 (1790).
KERNERIA, Moench, Meth. 595 (1794).
PLURIDENS, Neck. Elem. i. 86 (1790).
Abadiae, *DC. Prod.* v. 601.—Peruv.
abortiva, Schum. & Thonn. Beskr. Guin. Pl. 381 = pilosa.
abyssinica, Sch. Bip. ex Walp. Rep. vi. 167 = pilosa, quadriseta.
acaulis, *Baker, in Mart. Fl. Bras.* vi. iii. 247.—Paraguay.
Acmella, Lam. Encyc. i. 415 = Spilanthes Acmella.
acmelloides, Berg. in Vet. Acad. Handl. xxix. ii. (1768) 245 = Spilanthes Acmella.
adhaerescens, *Vell. Fl. Flum.* 348; viii. t. 88.—Bras.
alausensis, *H. B. & K. Nov. Gen. et Sp.* iv. 235.—Ecuador.
alba, DC. Prod. v. 605 = pilosa.
andicola, *H. B. & K. Nov. Gen. et Sp.* iv. 237.—Ecuador.
angustifolia, Lam. Encyc. i. 416 = Spilanthes urens.
angustifolia, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 369.—Ins. Sandvic.
angustissima, *H. B. & K. Nov. Gen. et Sp.* iv. 233.—Mexic.
angustissima, Spreng. Syst. iii. 453 = andicola.
anthriscoides, *DC. Prod.* v. 600.—Mexic.
apiifolia, Linn. Syst. ed. X. 1203; Pl. Jam. Pugill. 22 = Chrysanthellum procumbens.
arborea, Roxb. in Beats. App. 301, 325 = Petrobium arboreum.
arguta, H. B. & K. Nov. Gen. et Sp. iv. 231 = heterophylla.
artemisiaefolia, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 49.—Peruv.
atriplicifolia, Linn. Cent. ii. 30; Amoen. Acad. iv. 329 = Isocarpha atriplicifolia.
atrosanguinea, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 269 = Dahlia Zimapani.

BIDENS :—
atrosanguinea, Ortg. ex Regel, Gartenfl. x. (1861) 406. t. 347 = Cosmos diversifolius.
australis, Spreng. Syst. iii. 458 = Coreopsis micrantha.
Bardanna, Wall. Cat. n. 3291 = Chrysanthellum indicum.
Beckii, *Torr. in Spreng. Neue Entdeck.* ii. 135.—Am. bor.
Berteriana, Spreng. Syst. iii. 454 = Cosmos caudatus.
Bigelovii, *A. Gray, in Torr. Bot. Mex. Bound.* 91.—Am. bor. occ.
bimucronata, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 184.—Ins. S. Doming.
bipinnata, Baill. Hist. Pl. viii. 50 = Cosmos bipinnatus.
bipinnata, Hook. & Arn. in Bot. Beech. Voy. i. 30 = chilensis.
bipinnata, *Linn. Sp. Pl.* 832.—Reg. trop.
Bonplandii, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos tenellus.
brachycarpa, *DC. Prod.* v. 600.—Mexic.
Buchingeri, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 142 = Schimperi.
bullata, *Linn. Sp. Pl.* 833.—Am. bor.; Europ.
californica, DC. Prod. v. 599 = pilosa.
calva, C. B. Clarke, Comp. Ind. 141 = Cosmos sulphureus.
canescens, *Bertol. Fl. Guatimal.* 31.—Guatemala.
cannabina, Lam. Fl. Fr. ii. 44 = tripartita.
caracasana, *DC. Prod.* v. 600.—Venezuela.
carnea, *Heer, Ind. Sem. Hort. Turic.* (1842).—Guiana.
carvifolia, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos carvifolius.
caucalidea, *DC. Prod.* v. 604.—Mexic.
caudata, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos caudatus.
cernua, *Linn. Sp. Pl.* 832.—Reg. bor. temp.
Cervantesii, Baill. Hist. Pl. viii. 305 = Dahlia coccinea.
chilensis, *DC. Prod.* v. 603.—Chili.
chinensis, Willd. Sp. Pl. iii. 1719 = pilosa.
chrysantha, *DC. Prod.* v. 605.—Ind. occ.
chrysanthemoides, *Michx. Fl. Bor. Am.* ii. 136.—Am. bor.
cicutaefolia, *Tausch, in Flora*, xix. (1836) 395.—Hab.?
ciliata, *Hoffmgg. ex Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 46.—Hab.?
coccinea, Baill. Hist. Pl. viii. 49 = Dahlia coccinea.
connata, *Muhl. in Willd. Sp. Pl.* iii. 1718.—Am. bor.
consolidaefolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 185.—Peruv.
cordifolia, *Sch. Bip. in Flora*, xxxix. (1856) 361.—Ins. Pacif.
coreopsidis, DC. Prod. v. 599 = rubifolia.
coronata, Fisch. ex Steud. Nom. ed. II. i. 202 = Coreopsis coronata.
Cosmanthus, *Griseb. in Goett. Abh.* xix. (1874) 185.—Reg. Argent.
costaricensis, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 94.—Costa Rica.
crithmifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 234.—Cuba.
crocata, Cav. Ic. i. 66. t. 99 = Verbesina crocata.
cynapifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 235.—Cuba.
daucifolia, *DC. Prod.* v. 601.—Mexic.
decolorata, *H. B. & K. Nov. Gen. et Sp.* iv. 233.—Mexic.
decomposita, Wall. Cat. n. 298 = pilosa.
decussata, Pav. ex Steud. Nom. ed. II. i. 202 = hispida.
delphinifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 234.—Mexic.
denudata, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 183 = Glossogyne tenuifolia.
dichotoma, *Desf. Tabl.* ed. I. (1804) 104.—Bras.
Dilleniana, *Hill, Veg. Syst.* iii. 123.—Hab.?
diversifolia, *Ortg. in Regel, Gartenfl.* (1861) 408.—Mexic.
diversifolia, Hort. ex DC. Prod. v. 602 = grandiflora.
diversifolia, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos diversifolius.
diversifolia, Willd. ex DC. Prod. v. 602 = alausensis.
dondiaefolia, *Less. in Linnaea*, v. (1830) 155.—Mexic.

BIDENS:—
elongata, *Tausch, in Flora*, xix. (1836) 395.—Hab.?
exaristata, *DC. Prod.* v. 600.—Mexic.
fastigiata, Michalet, in Mém. Soc. Emul. Doubs, Sér. II. v. (1854) 29 = tripartita.
ferulaefolia, *DC. Prod.* v. 603.—Mexic.
ferulaefolia, Hemsl. Biol. Centr. Am. Bot. ii. 202, partim = procera.
fervida, Lam. Encyc. i. 415 = Spilanthes Acmella.
fistulosa, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 250.—Bras.
fixa, Hook. f. Fl. Brit. Ind. iii. 307, sphalm. = fusca.
flagellaris, *Baker, in Mart. Fl. Bras.* vi. III. 248.—Bras.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iv. 238.—Ecuador.
foeniculifolia, DC. Prod. v. 603 = procera?
foliosa, *Willd. Enum. Hort. Berol. Suppl.* 56, nomen.—Hab.?
frondosa, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iii. 309 = tripartita.
frondosa, *Linn. Sp. Pl.* 832.—Am. bor.
frutescens, Mill. Gard. Dict. ed. VIII. n. 4 = Salmea grandiceps.
fruticosa, *DC. Prod.* v. 596.—Hab.?
fruticosa, Linn. Sp. Pl. 833 = Wulffia stenoglossa.
fruticulosa, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 271.—Peruv.
fusca, Lam. Encyc. i. 416 = Spilanthes Acmella.
Gardneri, *Baker, in Mart. Fl. Bras.* vi. III. 246.—Bras.
glaberrima, *DC. Prod.* v. 601.—Am. austr.
glycinaefolia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 247.—Bras.
gracilis, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 368.—Ins. Sandvic.
gracilis, Torr. in Ann. Lyc. N. York, ii. (1828) 215 = Tholesperma gracile.
grandiflora, *Balb. Cat. Hort. Taur.* (1812) 19.—Am. austr.
graveolens, *Mart. in Isis*, (1824) 590.—Bras.
hawaiensis, *A. Gray, in Proc. Am. Acad.* v. (1861) 128.—Ins. Hawai.
helianthoides, H. B. & K. Nov. Gen. et Sp. iv. 230 = chrysanthemoides.
heterophylla, *Orteg. Hort. Matr.* 99. t. 12.—Am. bor. occ.; Mexic.
heterosperma, *A. Gray, Pl. Wright.* ii. 90.—Arizona.
hirsuta, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 369 = pilosa.
hirsuta, Sw. Prod. Veg. Ind. Occ. 110 = Salmea hirsuta.
hirta, Jord. in Gren. & Godr. Fl. Fr. ii. 168 = bullata.
hirtella, *H. B. & K. Nov. Gen. et Sp.* iv. 232.—Mexic.
hispida, *H. B. & K. l. c.* 237.—Venezuela.
Humboldtii, Sch. Bip. in Seem. Bot. Voy. Herald, 307 = Cosmos parviflorus.
humilis, *H. B. & K. Nov. Gen. et Sp.* iv. 234.—Ecuador.
hybrida, Thuill. Fl. Par. ed. II. 422 = tripartita.
incisa, G. Don, in Sweet, Hort. Brit. ed. III. 360 = rubifolia.
insipida, Lam. Encyc. i. 416 = Spilanthes insipida.
intermedia, Opiz, ex Nym. Consp. 348 = tripartita.
Jardinii, *Sch. Bip. in Flora*, xxxix. (1856) 360.—Ins. Pacif.
Kotschyi, Sch. Bip. ex Walp. Rep. vi. 168 = bipinnata.
laciniata, *Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop.* 283, nomen.—Afr. trop.
lantanoides, *A. Gray, in Proc. Am. Acad.* v. (1861) 128.—Ins. Pacif.
Lemmoni, *A. Gray, Syn. Fl. N. Am.* i. II. 297.—Arizona.
leucantha, Poepp. ex DC. Prod. v. 604 = chilensis.
leucantha, Willd. Sp. Pl. iii. 1719 = pilosa.
leucorrhiza, *DC. Prod.* v. 605.—China.
Leyboldii, *Phil. in Anal. Univ. Chil.* ii. (1865) 338.—Chili.
Lindleyi, Sch. Bip. in Seem. Bot. Voy. Herald, 307 = Cosmos tennifolius.
linearifolia, Sch. Bip. l. c. = Cosmos lincarifolius.
linearifoba, *Oliver, in Trans. Linn. Soc.* xxix. (1873) 99. t. 60.—Afr. trop.
longifolia, DC. Prod. v. 597 = heterophylla.
luxurians, Willd. Enum. Hort. Berol. 847 = heterophylla.
macranthus, *Griseb. in Goett. Abh.* xix. (1874) 186.—Reg. Argent.

BIDENS:—
macrosperma, Fisch. Hort. Gorenk. (1808) 59; ed. II. (1812) 37 = parviflora.
marginata, Perr. ex DC. Prod. v. 631 = Chrysanthellum indicum.
megapotamica, *Spreng. Syst.* iii. 454.—Am. austr.
Messerschmidii, Turcz. ex Ledeb. Fl. Ross. ii. 518 = parviflora.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 271.—China.
micrantha, Gaudich. Voy. Freyc. Bot. 464, 85 = Coreopsis micrantha.
minima, Huds. Fl. Angl. 310 = cernua.
mollis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 49.—Ecuador.
monticola, Poepp. & Endl. l. c. 49. t. 255 = rubifolia.
multifida, Desf. Cat. Hort. Par. ed. III. 186 = parviflora.
mutica, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1834) 368 = Coreopsis macrocarpa.
myrrhidifolia, *Tausch, in Flora*, xix. (1836) 394.—Hab.?
nivea, Linn. Sp. Pl. 833 = Melanthera hastata.
nivea, Sw. ex Steud. Nom. ed. II. i. 203 = Melanthera deltoidea.
nodiflora, *Linn. Sp. Pl.* 832.—Ind. or.
Novae-Caledoniae, *Forst. ex Schlecht. in Linnaea*, x. (1835–36) 480 (Cf. Glossogyne pinnatifida).—N. Hebrid.
nudiflora, Steud. Nom. ed. II. i. 203 sphalm. = nodiflora.
ocymifolia, Lam. Encyc. i. 416; Illustr. t. 668. f. 3 = Spilanthes alba.
odorata, *Cav. Ic.* i. 9. t. 13.—Mexic.
odorata, Dum.-Cours. Bot. Cult. iv. 241; G. Don, in Loud. Hort. Brit. 236 = grandiflora.
oleracea, Cav. ex Steud. Nom. ed. II. i. 203 = Spilanthes Acmella.
ostruthioides, *Sch. Bip. in Seem. Bot. Voy. Herald*, 308.—Mexic.
oxyodonta, *DC. Prod.* v. 600.—Ins. S. Doming.
palcacea, *Vis. in Nuov. Saggi Acc. Padov.* v. (1840) 266.—Nubia.
paniculata, *Hook. & Arn. Bot. Beech. Voy.* 66.—Ins. Tahiti.
paradoxa, *D. Don, ex Hook. & Arn. in Hook. Journ. Bot.* iii. (1840) 319.—Chili.
parviflora, *Willd. Enum. Hort. Berol.* 840.—Sibiria; China.
patula, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 405 = rubifolia.
pauciflora, Poir. Encyc. Suppl. i. 630 = parviflora.
pectinata, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 528, nomen.—Bolivia.
peduncularis, *Gaudich. Voy. Freyc. Bot.* 464.—Ins. Sandvic.
perianthomega, Steud. Nom. ed. II. i. 203, sphalm. = Bignonia perianthomega.
petiolata, Nutt. in Journ. Acad. Philad. vii. (1834) 99 = connata.
pilosa, *Linn. Sp. Pl.* 832.—Ind. occ.; Am. austr.
pinnata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 8.—Malaya.
pinnatifida, Forst. ex DC. Prod. v. 593 = Coreopsis micrantha.
pinnatifida, Buch.-Ham. in Wall. Cat. n. 3190 = Glossogyne pinnatifida.
platycephala, Oerst. Ind. Sem. Haun. (1859) 27; et in Linnaea, xxx. (1859–60) 753 = tripartita.
polycephala, *Sch. Bip. in Flora*, xxxix. (1856) 360.—Ins. Pacif.
portoricensis, *Spreng. in DC. Prod.* v. 601.—Porto Rico.
procera, *D. Don, in Bot. Reg.* t. 684.—Mexic.; Arizona.
procumbens, *H. B. & K. Nov. Gen. et Sp.* iv. 232.—Mexic.
pulcherrima, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 528, nomen.—Bolivia.
pumila, Steud. Nom. ed. I. 108 = tripartita.
purpurea, DC. Prod. v. 604 = Cosmos purpureus.
quadrangularis, DC. l. c. 600 = pilosa.
quadriaristata, DC. l. c. 595 = chrysanthemoides.
quadriseta, *Hochst. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 393.—Afr. trop.
quinqueradiata, Zea, ex Steud. Nom. ed. I. 108 = grandiflora.
radiata, Thuill. Fl. Par. ed. II. 432 = tripartita.
Ramtilla, Wall. Cat. n. 3191 = Heliopsis platyglossa.
reflexa, *Link, Enum. Hort. Berol.* ii. 306.—Mexic.
repens, D. Don, Prod. Fl. Nep. 180 = tripartita.

BIDENS:—
reptans, G. Don, in Sweet, Hort. Brit. ed. III. 360 = tripartita.
reptans, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos diversifolius.
rhombifolia, *Sch. Bip. in Bonplandia*, iv. (1856) 54.—Peruv.
Riedelii, *Baker, in Mart. Fl. Bras.* vi. III. 246.—Bras.
rigida, Hort. Calc. ex C. B. Clarke, Comp. Ind. 141 = Glossogyne pinnatifida.
riparia, *H. B. & K. Nov. Gen. et Sp.* iv. 236.—Venezuela.
rosea, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos pilosus.
rubifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 237. t. 381.—Ecuador.
rugulosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) I. 184.—Peruv.
sambucifolia, *Cav. Ic.* iii. 15. t. 219.—Peruv.; Mexic.
sandwicensis, Less. in Linnaea, vi. (1831) 508 = peduncularis.
Sartorii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 307.—Mexic.
scabiosoides, *Ortg. in Regel, Gartenfl.* (1861) 408.—Mexic.
scabiosoides, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos scabiosoides.
scandens, Linn. Sp. Pl. 833 = Salmea grandiceps.
scandicina, *H. B. & K. Nov. Gen. et Sp.* iv. 235.—Ecuador.
Schimperi, *Sch. Bip. ex Walp. Rep.* vi. 168.—Abyss.
scorzoneraefolia, *Baker, in Mart. Fl. Bras.* vi. III. 247.—Bras.
Seemannii, *Sch. Bip. in Seem. Bot. Voy. Herald*, 307.—Mexic.
serrata, *Pav. ex DC. Prod.* v. 597.—Peruv.
serrulata, Desf. Cat. Hort. Par. ed. III. (1829) 186 = Coreopsis aurea.
serrulata, *Sch. Bip. in Flora*, xxxix. (1856) 361.—Ins. Pacif.
silvaticus, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 245 = rubifolia.
speciosa, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 126.—Bras.
squarrosa, *H. B. & K. Nov. Gen. et Sp.* iv. 238.—Venezuela.
squarrosa, Less. in Linnaea, v. (1830) 156 = rubifolia.
striata, Schott, ex Sweet, Brit. Flow. Gard. t. 237 = pilosa.
subalternans, DC. Prod. v. 600 = pilosa.
sulphurea, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos sulphureus.
sundiaca, Blume, Bijdr. 913 = pilosa.
tenella, *Linn. Pl. Rar. Afr.* 17; *Amoen. Acad.* vi. 96.—Afr. austr.
tenuifolia, Labill. Sert. Austr. Caled. 44. t. 45 = Glossogyne tenuifolia.
tenuifolia, *Tausch, in Flora*, xix. (1836) 395.—Hab.?
tenuisecta, *A. Gray, Pl. Fendl.* 86.—Am. bor. occ.
tereticaulis, DC. Prod. v. 598 = rubifolia.
tetragona, *DC. l. c.* 596.—Mexic.
tinctoria, Baill. Hist. Pl. viii. 305 = Coreopsis tinctoria.
tridentata, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 183 = Glossogyne tridentata.
trifida, Roxb. Fl. Ind. iii. 411 = tripartita.
trifoliata, Gueldenst. ex Ledeb. Fl. Ross. ii. 516 = tripartita.
tripartita, Bigel. Fl. Bost. ed. II. 294 = connata.
tripartita, Boj. Hort. Maurit. 183 = pilosa.
tripartita, *Linn. Sp. Pl.* 831.—Europ.; Sibir.
triplinervia, *H. B. & K. Nov. Gen. et Sp.* iv. 231.—Mexic.
tuberosa, *Sch. Bip. in Linnaea*, xxxiv. (1865–66) 528, nomen.—Bolivia.
valladolidensis, Sch. Bip. in Seem. Bot. Voy. Herald, 308 = Cosmos crithmifolius.
valparadisiaca, Colla, in Mem. Acc. Torin. xxviii. (1835) 12. t. 24 = chilensis.
variabilis, Baill. Hist. Pl. viii. 305 = Dahlia variabilis.
venosa, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 405 = graveolens.
verticillata, Baill. Hist. Pl. viii. 305 = Coreopsis verticillata.
verticillata, *Linn. Sp. Pl.* 833.—Mexic.
Wallichii, DC. Prod. v. 598 = bipinnata.
Warszewicziana, *Regel, in Flora*, xxxii. (1849) 183.—Guatemala.

BIDWILLIA, Herb. ex Lindl. Veg. Kingd. 205 (1847) nomen. (Liliac.). Benth. & Hook. f. Gen. iii. 763.

BIEBERSTEINIA, Steph. ex [Fisch.] Hort. Gorenk. ed. I. 88 (1808); et in Mém. Soc. Nat. Mosc. i. (1811) 89. t. 9. *GERANIACEAE*, Benth. & Hook. f. i. 271.
DAYENIA, Michx. ex Jaub. & Spach, in Ann. Sc. Nat. Sér. III. vi. (1846) 137.
ambigua, Jaub. & Spach, in Ann. Sc. Nat. Sér. III. vi. (1846) 141 = multifida.
Aucheri, Jaub. & Spach, Illustr. ii. 110 = multifida.
brachypetala, Jaub. & Spach, l. c. 113. t. 192 = multifida.
Emodii, *Jaub. & Spach, l. c.* 109.—Tibet. occ.
Iciosepala, *Jaub. & Spach, l. c.* 114. t. 193.—Persia.
multifida, *DC. Prod.* i. 708.—Syria; Persia; Soongaria.
odora, Royle, Illustr. Bot. Himal. i. 153. t. 30 = Emodii.
odora, *Steph. ex [Fisch.] Hort. Gorenk.* ed. I. (1808) 88; *et ex Mém. Soc. Nat. Mosc.* i. (1811) 89.—Persia.
Orphanidis, *Boiss. Diagn.* Ser. II. i. 113.—Graecia; Cilicia.
Rucheri, Walp. Ann. i. 153 = multifida.

BIELSCHMEIDIA, Panch. & Sebert, Not. Bois Nouv. Caléd. 176 (1874) = **Beilschmiedia**, Nees (Laurin.).

BIELZIA, Schur, Enum. Pl. Transs. 409 (1866) = **Centaurea**, Linn. (Compos.).
Schwarzenbergiana, Schur, l. c. = C. ruthenica.

BIENERIA, Reichb. f. in Bot. Zeit. xi. (1853) 3. t. 1 = **Chloraea**, Lindl. (Orchid.).
boliviana, Reichb. f. l. c.—Bolivia.

BIENERTIA, Bunge, in Act. Hort. Petrop. vi. (1879) II. 425; et in Boiss. Fl. Orient. iv. 945 (1879). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 68.
cycloptera, *Bunge, ll. cc.*—Persia.

BIFOLIUM, G. Gaertn. Mey. & Scherb. Fl. Wett. i. 175 (1799) = **Maianthemum**, Wiggers (Liliac.).
cordatum, G. Gaertn. Mey. & Scherb. l. c. 209 = M. Bifolium.

BIFORA, Hoffm. Umb. Gen. ed. II. 191 (1816). *UMBELLIFERAE*, Benth. & Hook. f. i. 927.
ATREMA, DC. Prod. iv. 250 (1830).
BIFORIS, Spreng. Anleit. ii. i. 635 (1817); et in Roem. & Schult. Syst. vi. 448 (1820).
CORION, Link & Hoffmgg. Fl. Port. ii. 457 (1820).
americana, *Benth. & Hook. f. ex S. Wats. Bibl. Index*, 415.—Am. bor.
benghalensis, Wall. [Cat. n. 588] ex DC. Prod. iv. 140 = Oenanthe benghalensis.
dicocca, *Hoffm. Umb. Gen.* ed. II. 192.—Europ.; Oriens.
flosculosa, Bieb. Fl. Taur. Cauc. iii. 234 = dicocca.
glauca, Wall. Cat. n. 587 = Oenanthe bengalensis.
Loureiri, *Kostel. Allg. Med.-Pharm. Fl.* iv. 1183.—Cochinch.
radians, *Bieb. Fl. Taur. Cauc. Suppl.* 233.—Europ. austr.; Oriens.
testiculata, DC. Prod. iv. 249 = dicocca.
testiculata, *Roth, Enum. Pl. Phan. Germ.* i. 888.—Reg. Mediterr.

BIFORIS, Spreng. Anleit. ii. i. 635 (1817); et in Roem. & Schult. Syst. vi. 448 (1820) = **Bifora**, Hoffm. (Umbellif.).

BIFRENARIA, Lindl. Gen. et Sp. Orchid. 152 (1833). *ORCHIDEAE*, Benth. & Hook. f. Gen. iii. 546.
ADIPE, Rafin. Fl. Tellur. ii. 101 (1836).
STENOCORYNE, Lindl. Bot. Reg. xxix. (1843) Misc. 68.
albiflora, *Rodrig. Orch. Nov.* ii. 212, nomen.—Bras.
atropurpurea, *Lindl. Gen. et Sp. Orch.* 152.—Bras.
aurantiaca, *Lindl. Bot. Reg.* (1836) t. 1875.—Ind. occ.
aurea, *Rodrig. Orch. Nov.* ii. 212.—Bras.
aureofulva, *Lindl. Bot. Reg.* (1843) *Misc.* 52.—Bras.
bella, Lem. Jard. Fleur. iii. (1852–53) t. 325 = Coelia bella.
bicornaria, *Reichb. f. in Hamb. Gartenz.* xix. (1863) 12.—Bras.

BIFRENARIA:—
chloroleuca, *Rodrig. Orch. Nov.* ii. 212, nomen.—Bras.
calcarata, *Rodrig. l. c.* 213.—Bras.
clavigera, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 296.—Bras.
fragrans, Rodrig. Orch. Nov. ii. 213 = inodora.
Hadwenii, Lindl. in Paxt. Fl. Gard. ii. 67 (1851–52) = Scuticaria Hadwenii.
Harrisoniae, *Reichb. f. in Bonplandia*, iii. (1855) 217.—Hab.?
inodora, *Lindl. Bot. Reg.* xxix. (1843) *Misc.* 48.—Bras.
leucopetala, *Rodrig. Orch. Nov.* ii. 212, nomen.—Bras.
leucorrhoda, *Reichb. f. in Otto, Hamb. Gartenz.* xv. (1859) 54.—Hab.?
longicornis, *Lindl. Bot. Reg.* (1838) *Misc.* 93.—Guiana.
melanopoda, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 105.—Bras.
mellicolor, *Reichb. f. in Gard. Chron.* (1878) i. 622.—Bras.?
parvula, *Reichb. f. in Walp. Ann.* vi. 547.—Bras.
racemosa, *Lindl. Bot. Reg.* (1843) *Misc.* 52.—Bras.
sabulosa, *Rodrig. Orch. Nov.* i. 111.—Bras.
vinosa, *Rodrig. l. c.* ii. 212, nomen.—Bras.
vitellina, *Lindl. Bot. Reg.* (1843) *Misc.* 52.—Bras.
tyrianthina, *Reichb. f. Xenia*, i. 61, 223.—Bras.
Wageneri, *Reichb. f. in Bonplandia*, ii. (1854) 17.—Venezuela.

BIGAMEA, Koen. ex Endl. Gen. 1183 (1840) = **Ancistrocladus**, Wall. (Dipterocarp.).

BIGELONIA, Rafin. in Journ. Phys. lxxxix. (1819) 289, sphalm. = BIGELOWIA, Rafin. = **Arenaria**, Rupp. (Caryoph.).
elongata, Rafin. l. c. = A. lanuginosa.
montana, Rafin. l. c. = A. montana.

BIGELOVIA, Spreng. Syst. i. 366, 404 (1827) = **Spermacoce**, Linn. (Rubiac.).
alata, Bartl. ex Steud. Nom. ed. II. i. 203 = S. latifolia, longifolia.
angustifolia, Span. in Linnaea, xv. (1841) 320 = S. stricta?
commutata, Spreng. Syst. i. 404 = Diodia maritima.
cornifolia, Heynh. Nom. i. 121 = S. capitata.
corymbosa, Bartl. ex DC. Prod. iv. 550 = S. corymbosa.
discolor, Bartl. ex DC. l. c. 545 (= *Borreria discolor*).
distans, Hook. & Arn. Bot. Beech. Voy. 295 = S. distans.
elata, Bartl. ex DC. Prod. iv. 544 = S. latifolia, longifolia.
eryngioides, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 360 = S. diffusa.
gracilis, Spreng. Syst. i. 404 = S. ocymoides.
gracilis, Miq. in Verh. Nederl. Inst. Reeks III. v. (1852) 13—Malaya.
Kleinii, Wight & Arn. Prod. 437 = S. stricta.
laevicaulis, Miq. Fl. Ind. Bat. ii. 335 = S. ocymoides.
lasiocarpa, Wight & Arn. Prod. 437 = S. stricta.
linearis, Spreng. Syst. i. 405 = Diodia hyssopifolia.
linifolia, Spreng. l. c. 404 = S. laevis.
mitracarpoides, Miq. Fl. Ind. Bat. ii. 334 = S. stricta.
myriantha, Miq. l. c. = S. stricta.
ocymoides, Miq. l. c. 335 = S. ocymoides.
parviflora, Sieber, ex DC. Prod. iv. 542 = S. ocymoides.
parviflora, Spreng. Syst. i. 405 = S. ocymoides.
psylloides, Bartl. ex DC. Prod. iv. 547 = S. tenella.
pumila, Span. in Linnaea, xv. (1841) 320 = S. pumila.
pusilla, Spreng. Syst. i. 405 = S. pusilla.
Roxburghiana, Wight & Arn. Prod. 437 = S. stricta.
Roxburghiana, Wight, 'Cat. n. 1360' = S. ocymoides.
setidens, Miq. Fl. Ind. Bat. ii. 336 = S. stricta.
sociata, Span. in Linnaea, xv. (1841) 320.—Ins. Timor.
striata, Span. l. c. 321 = S. stricta.
stricta, Blume, Bijdr. 945 = S. stricta.
suaveolens, Spreng. Syst. i. 404 = S. tenella.
tenera, Heynh. Nom. i. 122 = S. tenera.
tenuis, Heynh. l. c. = S. tenuis.
umbellata, Spreng. Syst. i. 405.—Bras.
vaginata, Spreng. l. c. = S. laevis.
verticillata, Spreng. l. c. 404 = S. verticillata.
Zollingeriana, Miq. Fl. Ind. Bat. ii. 336.—Malaya.

BIGELOVIA, Spreng. Neue Entdeck. ii. 150 (1821) = **Samyda**, Linn.
brasiliensis, Spreng. l. c. = S. Selloi.

BIGELOVIA, Sm. in Rees, Cyclop. xxxix. (1819) = **Forestiera**, Poir. (Oleac.).
acuminata, Sm. l. c. = F. acuminata.
cassinoides, Sm. l. c. = F. cassinoides.
ligustrina, Sm. l. c. = F. ligustrina.
porulosa, Sm. l. c. = F. porulosa.

BIGELOWIA, DC. ex Ging. in DC. Prod. i. 290 (1824) = **Noisettia**, H. B. & K. (Violar.).

BIGELOWIA, DC. Prod. iv. 540 (1830) = BIGELOVIA, Spreng. = **Spermacoce**, Linn. (Rubiac.).

BIGELOWIA, DC. Prod. v. 329 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 1232.
CHRYSOTHAMNUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 323.
ISOCOMA, Nutt. l. c. 320 (1841).
LINOSYRIS, Torr. & Gray, Fl. N. Am. ii. 232 (1841).
acradenia, *Greene, in Bull. Torrey Club*, x. (1883) 126.—Am. bor. occ.
albida, *M. E. Jones, ex A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 209.—Am. bor. occ.
arborescens, *A. Gray, l. c.* viii. (1873) 640.—Calif.
Bigelovii, *A. Gray, l. c.* xvii. (1881–82) 209.—N. Mexic.; Colorado.
Bolanderi, *A. Gray, l. c.* viii. (1873) 641.—Calif.
brachylepis, *A. Gray, in Bot. Calif.* i. 614.—Calif.
carnosa, Benth. & Hook. f. Gen. ii. 1232, 255 = Aster carnosus.
ceruminosa, *A. Gray, in Proc. Am. Acad.* viii. (1873) 643.—Calif.
Cooperi, *A. Gray, l. c.* 640.—Calif.
coronopifolia, *A. Gray, l. c.* 638.—Texas.
depressa, *A. Gray, l. c.* 643.—Am. bor. occ.
diffusa, *A. Gray, l. c.* 640.—Calif.
Douglasii, A. Gray, l. c. 644 = viscidiflora.
dracunculoides, *DC. Prod.* v. 329.—Am. bor.
Drummondii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 639.—Texas.
Engelmanni, *A. Gray, l. c.* xi. (1876) 75.—Am. bor. occ.
fuliginea, *A. Gray, l. c.* viii. (1873) 638.—Ecuador.
furfuracea, *Greene, in Bull. Calif. Acad.* i. n. 3 (1885) 87.—Calif.
graveolens, A. Gray, in Proc. Am. Acad. viii. (1873) 644 = dracunculoides.
Greenei, *A. Gray, l. c.* xi. (1876) 75.—Colorado.
Hartwegi, *A. Gray, ex Hemsl. Biol. Centr. Am. Bot.* ii. 115.—Mexic.
Howardi, *A. Gray, in Proc. Am. Acad.* viii. (1873) 641.—Colorado.
hypoleuca, *A. Gray, l. c.* 638.—Ecuador.
intricata, A. Gray, l. c. xvii. (1881–82) 208 = Aster carnosus.
juncea, *Greene, in Coult. Bot. Gazette*, vi. (1881) 184.—Arizona.
lanceolata, A. Gray, in Proc. Am. Acad. viii. (1873) 639 = viscidiflora.
leiosperma, *A. Gray, Syn. Fl. N. Am.* i. II. 139.—Am. bor. occ.
Mandonii, *Benth. & Hook. f. Gen.* ii. 1232, 255.—Bolivia?
Menziesii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 638.—Am. bor. occ.
missouriensis, DC. Prod. v. 329 = dracunculoides.
mohavensis, *Greene, ex A. Gray, Syn. Fl. N. Am.* i. II. 138.—Am. bor. occ.
nevadensis, *A. Gray, l. c.* 136.—Calif.
nudata, *DC. Coll. Mém.* ix. t. 5.—Am. bor. or.
oppositifolia, *A. Gray, in Proc. Am. Acad.* xv. (1880) 32.—Mexic.
paniculata, *A. Gray, l. c.* viii. (1873) 644.—Am. bor. occ.
Parishii, *Greene, in Bull. Torrey Club*, ix. (1882) 62.—Calif.
Parryi, *A. Gray, in Proc. Am. Acad.* viii. (1873) 642.—Colorado.
pluriflora, *A. Gray, l. c.* 639.—Colorado?
pulchella, *A. Gray, l. c.* 643.—Texas; N. Mexic.
ramulosa, *Benth. & Hook. f. Gen.* ii. 1232, 255.—Am. bor.
rupestris, Greene, in Coult. Bot. Gazette, vi. (1881) 183 = Aplopappus cuneatus.
spathulata, A. Gray, in Proc. Am. Acad. xi. (1876) 74 = Aplopappus cuneatus.
teretifolia, *A. Gray, l. c.* viii. (1873) 644.—Calif.; Arizona.
tridentata, Greene, in Bull. Torrey Club, x. (1883) 126 = veneta.

BIGELOWIA:—
uniligulata, DC. Prod. v. 329 = Solidago neglecta.
Vaseyi, *A. Gray, in Proc. Am. Acad.* xii. (1877) 58.—Colorado.
veneta, *A. Gray, l. c.* viii. (1873) 638.—Calif.; Mexic.
virgata, DC. Prod. v. 329 = nudata.
viscidiflora, *DC. l. c.* vii. 279.—Am. bor.
Wrightii, *A. Gray, in Proc. Am. Acad.* viii. (1873) 639.—Am. bor. occ.

BIGGINA, Rafin. Alsog. Am. 14 (1838) = **Salix**, Tourn.
angustata, Rafin. l. c. = S. cordata ?
argentea, Rafin. l. c. = S. argentea.
conifera, Rafin. l. c. = S. humilis.

BIGLANDULARIA, Karst. in Linnaea, xxviii. (1856) 416. *GENTIANEAE.*
azurea, *Karst. l. c.* 417.—N. Granat.

BIGLANDULARIA, Scem. in Gard. Chron. (1868) 738 = **Sinningia**, Nees (Gesneriac.).
conspicua, Scem. l. c.—Bras.

BIGNONIA, [Tourn.] Linn. Syst. ed. I (1735) *BIGNONIACEAE*, Benth. & Hook. f. ii. 1033.
Anisostichus, Bur. Monog. Bign. 43. t. 6 (1864).
Clematitaria, Bur. l. c. 44 (1864).
Cremastus, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 187.
Cupulissa, Rafin. Fl. Tellur. ii. 57 (1836).
Cydista, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 191.
Doxantha, Miers, l. c. 188 (1863).
Endoloma, Rafin. Sylva Tellur. 79 (1838).
Hieranthes, Rafin. l. c. (1838).
Odisca, Rafin. l. c. 80 (1838).
Panterpa, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 195.
Paragonia, Bur. in Bull. Soc. Bot. Fr. xix. (1872) 17.
Pentelesia, Rafin. Sylva Tellur. 146 (1838).
Petastoma, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 193.
Phaedranthus, Miers, l. c. 182 (1863).
Pleonotoma, Miers, l. c. 184 (1863).
Pongelia, Rafin. Sylva Tellur. 78 (1838).
Proterpia, Rafin. l. c. 80 (1838).
Pyrostegia, Presl, in Abh. Boehm. Ges. V. iii. (1845) 523 (Bot. Bemerk. 93).
Sererea, Rafin. Sylva Tellur. 107 (1838).
Stizophyllum, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 197.
Temnocydia, Mart. ex DC. Prod. ix. 143 (1845).
acutiflora, Mart. ex DC. Prod. ix. 181 = Lundia nitidula.
acutissima, *Cham. in Linnaea*, vii. (1832) 691.—Bras.
acutistipula, *Schlecht. l. c.* xxvi. (1853-55) 375.—Mexic.
adenophylla, Wall. Cat. n. 6502 = Heterophragma adenophyllum.
adhatodaefolia, *Jacq. Fragm.* 61.—Venezuela.
acquinoctialis, *Linn. Sp. Pl.* 623.—Am. trop.
aequinoctialis, Sieber, ex DC. Prod. ix. 146 = B. Unguiscati.
aesculifolia, H. B. & K. Nov. Gen. et Sp. iii. 140 = Tabebuia aesculifolia.
Aesculus, Desf. Cat. Hort. Par. ed. III. 397 = Tecoma Aesculus.
affinis, *G. Don, Gen. Syst.* iv. 217.—Bras.
africana, *Lam. Encyc.* i. 424.—Afr. trop.
Agnus-castus, Cham. in Linnaea, vii. (1832) 673 = Arrabidaea Agnus-castus.
alata, Pav. ex DC. Prod. ix. 224 = Tecoma Guarume.
alba, Aubl. Pl. Gui. ii. 653 = Spathodea bracteosa.
albida, Blume, in Verh. Bat. Gen. ix. (1825) 195 = Aeschynanthus purpurascens.
albida, Steud. Nom. ed. II. i. 204 = Aeschynanthus angustifolia.
albiflora, *Salzm. ex DC. Prod.* ix. 167.—Bras.
allamandaeflora, *Benth. ex Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 186, nomen.—Bras.
alliacea, *Lam. Encyc.* i. 421.—Guiana.
amara, Loud. Hort. Brit. 483; ex G. Don, Gen. Syst. iv. 222 = Stereospermum hypostictum.
amoena, Wall. Cat. n. 6512; et Pl. As. Rar. ii. 78. t. 183 = Stereospermum hypostictum.
anastomosans, *DC. Prod.* ix. 144.—Madag.

BIGNONIA:—
Andrieuxii, *DC. Prod.* ix. 156.—Mexic.
angrensis, *Vell. Fl. Flum.* 246; vi. t. 23.—Bras.
angustifolia, Blume, Cat. Hort. Buitenz. 82 = Aeschynanthus angustifolia.
antisyphilitica, Mart. Reise Bras. i. 283 = Cybistax antisyphilitica.
apurensis, *H. B. & K. Nov. Gen. et Sp.* iii. 138.—Venezuela.
aquatilis, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 780 = Tabebuia fluviatilis.
araliacea, Cham. in Linnaea, vii. (1832) 683 = Tecoma araliacea.
argyrea, *André, in Rev. Hortic.* (1865) 251.—Hab. ?
argyreo-violascens, *Hort. ex Lem. Illustr. Hortic.* (1866) t. 469.—Hab. ?
Arrabidae, *Steud. Nom.* ed. II. i. 204.—Bras.
Arthrerion, Mart. Herb. Fl. Bras. 287 = Distictis Arthrerion.
articulata, Desf. in Poir. Encyc. Suppl. i. 636 = Phyllarthron Noronhianum.
arvensis, Vell. Fl. Flum. 250; vi. t. 40 = Anemopaegma mirandum.
atrovirens, DC. Rev. Bign. ex Prod. ix. 171 = Tecoma atrovirens.
atrovirens, Roth, Nov. Sp. 284 = Dolichandrone falcata.
aurea, Manso, Enum. 40 = Tecoma aurea.
auriculigera, Mart. ex DC. Prod. ix. 194 = Pithecoctenium stipulacea.
australis, Ait. Hort. Kew. ed. II. iv. 34 = Tecoma australis.
Azedarachta, Kon. & Sims, Ann. Bot. i. (1805) 578 = Millingtonia hortensis.
Baclei, DC. Rev. Bign. ex Prod. ix. 171 = Pithecoctenium cynanchoides.
Balbisiana, *DC. Prod.* ix. 153.—Bras.
Banaibanai, Callery, ex Bur. in Baill. Adansonia, ii. (1861-62) 195 = Sterospermum Banaibanai.
barbata, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 782 = Tecoma barbata.
Bellas, *Sellow, ex Steud. Nom.* ed. II. i. 204.—Bras.
bibracteata, Bert. ex DC. Prod. ix. 240 = Bravaisia floribunda.
Bidwilliana, *Regel, Cat. Pl. Hort. Aksakov.* 19.—Hab. ?
bijuga, *Vahl, Symb. Bot.* ii. 68.—Madag.
binata, Thunb. Diss. Pl. Bras. iii. 35. n. 27 = Anemopaegma racemosum.
bipinnata, Salzm. ex DC. Prod. ix. 230 = Jacaranda nitida.
Blanchetii, *DC. l. c.* 164.—Bras.
Bojeri, *DC. l. c.* 165.—Madag.
botryoides, Cham. in Linnaea, vii. (1832) 100 = Pithecoctenium botryoides.
brachycalyx, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1085.—Guian.
brachypoda, *DC. Prod.* ix. 145.—Bras.
bracteata, Cham. in Linnaea, vii. (1832) 692 = Adenocalymna bracteatum.
bracteolata, *DC. Prod.* ix. 157.—Bras.
bracteosa, *DC. l. c.* 165.—Madag.
brasiliana, Lam. Encyc. i. 425 = Jacaranda brasiliana.
buccinatoria, *Mairet, ex DC. Prod.* ix. 195.—Mexic.
caerulea, Linn. Sp. Pl. 625 = Jacaranda caroliniana.
callistegioides, Cham. in Linnaea, vii. (1832) 712 = Cuspidaria callistegioides.
calycina, *DC. Prod.* ix. 160.—Bras.
campanulata, Cham. in Linnaea, vii. (1832) 711 = Cuspidaria campanulata.
candicans, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Arrabidaca candicans.
capensis, Thunb. Prod. Pl. Cap. 105 = Tecoma capensis.
capreolata, *Linn. Sp. Pl.* 624.—Am. bor.
carichanensis, *H. B. & K. Nov. Gen. et Sp.* iii. 137.—Venezuela.
Caroba, Vell. Fl. Flum. 250; vi. t. 43 = Jacaranda Caroba.
Carolinae, *Lindl. Bot. Reg.* (1842) sub t. 45.—Hab. ?
caryophyllea, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 293.—Porto Rico.
cassinoides, Lam. Encyc. i. 418 = Tabebuia cassinoides.
castaneaefolia, *DC. Prod.* ix. 145.—Peruv.
Catalpa, Linn. Sp. Pl. 622 = Catalpa bignonioides.
Catharinae, DC. Rev. Bign. ex Prod. ix. 172 = Pithecoctenium Catharinae.

BIGNONIA:—
caudata, DC. Prod. ix. 166 = Stereospermum caudatum.
caudata, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 382 = Stereospermum chelonoides.
cauliflora, Sieber, ex DC. Prod. ix. 241 = Colea floribunda.
Chamberlaynii, *Sims, Bot. Mag.* t. 2148.—Bras.
chelonoides, Linn. f. Suppl. 282 = Stereospermum chelonoides.
Cherere, Lindl. Bot. Reg. t. 1301 = buccinatoria.
Chica, *Humb. & Bonpl. Pl. Aequin.* i. 107. t. 31.—Venezuela.
chinensis, Lam. Encyc. i. 423 = Tecoma grandiflora.
chondrogona, *Miq. in Linnaea*, xviii. (1844) 611.—Guiana.
chrysantha, Jacq. Hort. Schoenb. ii. 45 = Tecoma chrysantha.
chrysoleuca, *H. B. & K. Nov. Gen. et Sp.* iii. 134.—Venezuela.
Cicutaria, Koen. ex Mart. in Denkschr. Akad. Muench. vi. (1820) 153 = Millingtonia hortensis.
ciliata, *Wawra, in Oestr. Bot. Zeitschr.* xii. (1862) 237.—Bras.
cinerea, DC. Rev. Bign. ex Prod. ix. 172 = Pithecoctenium cinereum.
cinnamomea, *DC. Prod.* ix. 164.—Bras.
citrifolia, *Vitm. Summa Veg.* iii. 511.—Hab. ?
Clematis, *H. B. & K. Nov. Gen. et Sp.* iii. 141.—Venezuela.
coccinea, Steud. Nom. ed. I. 109 = Tecoma radicans.
coccinea, *Vell. Fl. Flum.* 250; vi. t. 42.—Bras.
cognata, *Cham. in Linnaea*, vii. (1832) 703.—Bras.
Coito, Vell. Fl. Flum. 243; vi. t. 19 = Spathodea Coito.
Colais, Buch.-Ham. ex Wall. Cat. sub n. 6501 = Stereospermum chelonoides.
Colei, Boj. ex Hook. Bot. Mag. t. 2817 = Colea mauritiana.
colorata, Poir. Encyc. Suppl. i. 634 = Arrabidaea candicans.
comosa, Cham. in Linnaea, vii. (1832) 693 = Adenocalymna comosum.
comosa, *Roxb. Hort. Beng.* [95]; *Fl. Ind.* iii. 103.—Ins. Molucc.
compressa, Lam. Encyc. i. 424 = Colea decora.
conjugata, Vell. Fl. Flum. 243; vi. t. 18 = Arrabidaea conjugata.
conspicua, Rich. ex DC. Prod. ix. 221 = Tecoma conspicua.
convoluta, Vell. Fl. Flum. 248; vi. t. 31 = Cuspidaria puberula.
Copaia, Aubl. Pl. Gui. ii. 650. t. 262 = Jacaranda Copaia.
corallina, *Jacq. Fragm.* 37.—Venezuela.
corchoroides, Cham. in Linnaea, vii. (1832) 706 = Arrabidaea corchoroides.
cordata, Vell. Fl. Flum. 247; vi. t. 29 = Lundia cordata.
cordifolia, *DC. Prod.* ix. 162.—Bras.
coriacea, Sellow, ex Steud. Nom. ed. II. i. 204, 205 = Selloi.
corymbifera, *Vahl, Ecl. Am.* ii. 45. t. 17.—Am. austr.
craterophora, *DC. Prod.* ix. 147.—Bras.
crenata, *Lodd. ex Loud. Hort. Brit.* 242.—Ind. or.
crispa, Buch.-Ham. ex Roxb. Hort. Beng. 47; Fl. Ind. iii. 103 = Dolichandrone crispa.
crucifera, Bert. ex DC. Prod. ix. 213 = Tabebuia rufinervis.
crucigera, Linn. Sp. Pl. 624 = capreolata.
cujabana, *DC. Prod.* ix. 164.—Bras.
cuneifolia, *DC. l. c.* 157.—Bahia.
cuprea, *Cham. in Linnaea*, vii. (1832) 665.—Bras.
cupulata, *Splitg. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 6.—Bras.
curialis, Vell. Fl. Flum. 253; vi. t. 55 = Jacaranda curialis.
cuspidata, Boj. ex DC. Prod. ix. 243 = Periblema cuspidata.
Cymbalum, *Cham. in Linnaea*, vii. (1832) 716.—Bras.
cymosa, *Vell. Fl. Flum.* 249; vi. t. 36.—Bras.
dasyantha, DC. Rev. Bign. ex A. DC. Prod. ix. 172 (= *Pachyptera dasyantha*).—Bras.
decipiens, *Lodd. ex Loud. Hort. Brit.* 242.—Am. austr.
decomposita, *Miq. in Linnaea*, xviii. (1844) 260.—Guiana.
densiflora, Mart. ex DC. Prod. ix. 181 = Lundia densiflora.
dentata, *DC. l. c.* 162.—Bras.

BIGNONIA:—
dichotoma, *Jacq. Enum. Pl. Carib.* 25.—Am. austr.
dichotoma, *Vell. Fl. Flum.* 248; vi. t. 32.—Bras.
difficilis, *Cham. in Linnaea*, vii. (1832) 714.—Bras.
digitalis, Vell. Fl. Flum. 251; vi. t. 47 = Zeyheria montana.
digitata, E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 782 = Tecoma Meyeriana.
discolor, Boj. ex DC. Prod. ix. 165 = Bojeri.
discolor, *R. Br. in Salt, Voy. Abyss. App.* 64, nomen. —Abyss.
discolor, Rich. Sert. Astrol. p. xxix = moluccana.
diversifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 133.—Mexic.
dolichoides, *Cham. in Linnaea*, vii. (1832) 696.—Bras.
Dombeyana, DC. Rev. Bign. ex A. DC. Prod. ix. 172 = Delostoma nervosum?
Doniana, *DC. Prod.* ix. 145.—Bras.
dubia, *Linn. Pl. Surin.* n. 77; *Amoen. Acad.* viii. 258. —Guiana.
echinata, Aubl. Pl. Gui. ii. 648 = Pithecoctenium Aubletii.
echinata, Jacq. Enum. Pl. Carib. 25 = Pithecoctenium muricatum.
ehretioides, *Cham. in Linnaea*, vii. (1832) 704.—Bras.
elegans, *Cham. l. c.* 702.—Bras.
elegans, Vell. Fl. Flum. 247; vi. t. 27 = exserta.
elliptica, Cham. in Linnaea, vii. (1832) 686 = Tecoma elliptica.
elliptica, *Thunb. Diss. Pl. Bras.* iii. 34.—Bras.
elliptica, Vell. l. c. 250; vi. t. 44 = Jacaranda elliptica.
elongata, *Vahl. Ecl.* ii. 45. t. 16.—Am. austr.
erubescens, *DC. Prod.* ix. 157.—Bras.
euphorioides, Boj. Hort. Maurit. 219 = Stereospermum euphorioides.
exoleta, *Vell. Fl. Flum.* 248; vi. t. 30.—Bras.
exserta, *DC. Prod.* ix. 158.—Bras.
extensa, *Soland. ex Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 196, nomen.—Bras.
fagoides, *Cham. in Linnaea*, vii. (1832) 680.—Bras.
falcata, Koen. ex DC. Prod. ix. 206 = Dolichandrone falcata.
falcata, *Vell. Fl. Flum.* 245; vi. t. 17.—Bras.
fallax, *Cham. in Linnaea*, vii. (1832) 717.—Bras.
farinosa, Salzm. ex DC. Prod. ix. 185 = Arrabidaea virescens.
fasciculata, Vell. Fl. Flum. 247; vi. t. 25 = Arrabidaea fasciculata.
Ferdinandi, *Welw. Apont.* 584.—Afr. trop.
ferruginea, *Tweedie, ex Miers, in Proc. Roy. Hort. Soc.* iii. (1863) 190, nomen.—Reg. Argent.?
ficifolia, Steud. Nom. ed. II. i. 204, sphal. = filicifolia.
filicifolia, Anders. in Trans. Soc. Arts. xxv. (1807) 200 = Jacaranda filicifolia.
filiformis, A. Cunn. ex DC. Prod. ix. 209 = Spathodea filiformis.
fimbriata, Wall. Cat. n. 6500 = Stereospermum fimbriatum.
flava, *DC. Prod.* ix. 168.—Bras.
flavescens, Vell. Fl. Flum. 252; vi. t. 51 = Tecoma flavescens.
flavida, *DC. Prod.* ix. 168.—Bras.
floribunda, G. Don, Gen. Syst. iv. 216 = Doniana.
floribunda, *H. B. & K. Nov. Gen. et Sp.* iii. 134.—Mexic.
florida, Salisb. Prod. 106 = Tecoma radicans.
fluminensis, *Vell. Fl. Flum.* 246; vi. t. 24.—Bras.
fluviatilis, Aubl. Pl. Gui. ii. 655 = Tabebuia fluviatilis.
fluviatilis, H. B. & K. Nov. Gen. et Sp. ii. 139 = Tecoma barbata.
fluviatilis, G. F. W. Mey. Prim. Fl. Esseq. 212 = Tecoma Meyeriana.
Fockeana, *Miq. in Linnaea*, xviii. (1844) 609.—Guiana.
fraxinea, *Desf. Cat. Hort. Par.* ed. III. 398.—Hab.?
fraxinifolia, Spreng. Syst. ii. 833 = Spathodea fraxinifolia.
fraxinoides, *Perr. in Mém. Soc. Linn. Par.* iii. (1824) 102.—Java.
frutescens, Mill. Gard. Dict. ed. VIII. n. 3 = Tecoma stans.
fulva, Cav. Ic. iii. 58 = Tecoma fulva.
Gaudichaudi, DC. Rev. Bign. ex Prod. ix. 172 = Lundia pubescens.
Ghorta, Buch.-Ham. ex Wall. Cat. n. 6510 = Stereospermum Ghorta.

BIGNONIA:—
Giesbreghtii, *Heller, ex Peyr. in Linnaea*, xxx. (1859–60) 45.—Mexic.
gigantea, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 8.—Malaya.
glaberrima, *Cham. in Linnaea*, vii. (1832) 677.—Bras.
glabrata, *H. B. & K. Nov. Gen. et Sp.* iii. 135.—Am. austr.
glandulosa, *Mart. ex DC. Prod.* ix. 149.—Bras.
glandulosa, Schum. Beskr. Guin. Pl. 274 = Spathodea adenantha.
glauca, Decne. in Jacquem. Voy. Bot. 137. t. 142 = Tecoma undulata.
glauca, *Wall. Cat.* n. 6506.—Burma.
glutinosa, *DC. Prod.* ix. 162.—Bras.
gnaphalantha, *A. Rich. Fl. Cub. Fanerog.* ii. 105.—Cuba.
gracilis, Lodd. Bot. Cab. t. 1705 = B. Unguis-cati.
grandiflora, Spreng. Syst. ii. 830 = grandifolia.
grandiflora, Thunb. Fl. Jap. 253 = Tecoma grandiflora.
grandifolia, *Jacq. Hort. Schoenb.* iii. 19. t. 287.—Am. austr.
grandifolia, Vell. Fl. Flum. 247; vi. t. 28 = Adenocalymna grandifolium et subsessilifolium.
gratissima, Koen. ex Wall. Cat. sub n. 6499; et DC. Prod. ix. 211 = Stereospermum suaveolens.
Guarumé, Domb. ex A. DC. l. c. 225 = Tecoma Guarumé.
guayaquilensis, *DC. l. c.* 155.—Ecuador.
haemantha, Bert. ex Spreng. Syst. ii. 832 = Tabebuia haemantha.
hebantha, Mart. ex DC. Prod. ix. 181 = Lundia hebantha.
heptaphylla, Vell. Fl. Flum. 251; vi. t. 48 = Tecoma heptaphylla.
heterophylla, Willd. Sp. Pl. iii. 298 = B. Kerere.
heteropoda, DC. Rev. Bign. ex Prod. ix. 173 = Tecoma heteropoda.
heterotricha, DC. l. c. = Tabebuia heterotricha.
hexagona, DC. l. c. = Pithecoctenium hexagonum.
hibiscifolia, *Cham. in Linnaea*, vii. (1832) 705.—Bras.
hirsuta, Lam. Encyc. i. 222 = Tecoma hirsuta.
hirta, *Vell. Fl. Flum.* 249; vi. t. 35.—Bras.
hispida, *DC. Prod.* ix. 152.—Bras.
hondensis, *H. B. & K. Nov. Gen. et Sp.* iii. 135.—N. Granat.
Hostmanni, *E. Mey. in Nov. Act. Nat. Cur.* xii. (1825) ii. 779.—Guiana.
Hymenaea, *DC. Prod.* ix. 158.—Bras.
ignea, *Vell. Fl. Flum.* 244; iv. t. 15.—Bras.
ilicifolia, Pers. Syn. ii. 170 = Tabebuia ilicifolia.
inaequalis, *DC. Prod.* ix. 170.—Guiana.
Imperatoris-Maximiliana, *Wawra, Bot. Ergeb. Maxim. Bras.* 73.—Bras.
incarnata, *Aubl. Pl. Gui.* ii. 645. t. 261.—Guiana.
incisa, Hort. ex DC. Prod. ix. 224 = Tecoma stans.
indica, Linn. Sp. Pl. 625 = Oroxylum indicum.
indica, Lour. Fl. Cochinch. 460 = Pajanelia multijuga.
inflata, *Griseb. in Goett. Abh.* xxiv. (1879) 256.—Reg. Argent.
jasminifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 141.—Venezuela.
jasminoides, A. Cunn. ex DC. Prod. ix. 225 = Tecoma jasminoides.
jasminoides, *Thunb. Pl. Bras.* iii. 36. n. 28.—Bras.
Kerere, *Aubl. Pl. Gui.* ii. 644.—Guiana.
labiata, *Cham. in Linnaea*, vii. (1832) 701.—Bras.
lactiflora, Vahl, Symb. Bot. iii. 80. t. 66 = Distictis lactiflora.
laeta, *Wall. Cat.* n. 6505.—Burma.
lanata, Steud. Nom. ed. II. i. 205 = discolor.
lanceolata, *DC. Prod.* ix. 163.—Bras.
lanuginosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 491.—Mexic.
laserpitiifolia, *Mart. ex DC. Prod.* ix. 168.—Bras.
latifolia, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Callichlamys riparia.
latisiliqua, *Poir. Encyc. Suppl.* i. 632.—Guiana.
laurifolia, *Vahl. Ecl. Am.* ii. 44.—Guiana.
lexiflora, DC. Rev. Bign. 21; Prod. ix. 173 = Pithecoctenium laxiflorum.
lenta, *Mart. ex DC. Prod.* ix. 159.—Bras.
lepidophylla, *A. Rich. Fl. Cub. Fanerog.* ii. 104.—Ins. Pinet. (Ind. occ.).
lepidota, H. B. & K. Nov. Gen. et Sp. iii. 139 = Tecoma lepidota.

BIGNONIA:—
lepidota, *Seem. Bot. Voy. Herald*, 197.—Mexic.
leucantha, Vell. Fl. Flum. 251; vi. t. 49 = Sparattosperma lithontripticum.
leucopogona, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = seq.
Leucopogon, *Cham. in Linnaea*, vii. (1832) 707.—Bras.
leucoxylla, Vell. Fl. Flum. 252; vi. t. 54 = Tabebuia leucoxyla.
Leucoxylon, DC. Cat. Bign. ex Prod. ix. 173 = Tecoma Salzmanni.
Leucoxylon, Linn. Sp. Pl. 624 = Tecoma Leucoxylon.
Lindleyi, *DC. Prod.* ix. 147.—Reg. Argent.
linearis, Cav. Ic. iii. 35. t. 269 = Chilopsis saligna.
littoralis, *H. B. & K. Nov. Gen. et Sp.* iii. 139.—Mexic.
longa, Vell. Fl. Flum. 249; vi. t. 37 = Lundia longa.
longiflora, DC. Prod. ix. 206, sphalm. = Pajanelia Rheedii.
longiflora, Vell. Fl. Flum. 252; vi. t. 52 = Tecoma speciosa.
longiflora, *Cav. Ic.* vi. 58. t. 581.—Guayaquil.
longifolia, Willd. Sp. Pl. iii. 306 = Pajanelia Rheedii.
longisiliqua, Vell. Fl. Flum. 247; vi. t. 26 = Arrabidae.
longisiliqua, *Bert. ex Spreng. Syst.* ii. 830.—Venezuela.
longissima, Jacq. Enum. Pl. Carib. 25 = Catalpa longissima.
longissima, Lour. Fl. Cochinch. ii. 380 = Dolichandrone Rheedii.
lucida, *Lodd. ex Loud. Hort. Brit.* 242.—Am. austr.
lugubris, Salisb. Prod. 106 = Oroxylum indicum.
Lundii, DC. Rev. Bign. 21, ex Prod. ix. 173 = Pithecoctenium Lundii.
macrophylla, Cham. in Linnaea, vii. (1832) 689 = Adenocalymna macrophyllum.
macrostachya, *Wall. Cat.* n. 6504.—Burma.
magnifica, *Bull, in Gard. Chron.* (1879) ii. 73.—N. Granat.
magnoliaefolia, *H. B. & K. Nov. Gen. et Sp.* iii. 136.—Venezuela.
Mansoana, *DC. Prod.* ix. 157.—Bras.
maranhamensis, *DC. l. c.* 165.—Bras.
marginata, Cham. in Linnaea, vii. (1832) 695 = Adenocalymna marginatum.
marmoratum, *Hort. ex André, in Rev. Hortic.* (1865) 253.—Hab.?
Martini, *DC. Prod.* ix. 152.—Guiana.
Martiusiana, *DC. l. c.* 156.—Bras.
megapotamica, *Spreng. Syst.* iv. 237.—Bras.
meonantha, Link, Enum. ii. 130 = Tecoma australis.
meridionalis, Salisb. Prod. 106 = Tecoma Leucoxylon.
Meyeniana, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 366.—Peruv.
microcalyx, *G. F. W. Mey. Prim. Fl. Esseq.* 211.—Guiana.
microphylla, *Lam. Encyc.* i. 418.—Ins. S. Doming.
Mina, *Robin, Voy. Louis.* iii. 408.—Am. bor.
miranda, Cham. in Linnaea, vii. (1832) 687 = Anemopaegma mirandum.
mollis, *Vahl, Ecl. Am.* ii. 46.—Guiana.
mollissima, *H. B. & K. Nov. Gen. et Sp.* iii. 133.—Venezuela.
moluccana, *DC. Prod.* ix. 144.—Ins. Amboina.
moringaefolia, *DC. l. c.* 170.—Guiana.
multifida, *Lodd. ex Loud. Hort. Brit.* 242.—Ind. or.
multiflora, *Salzm. ex DC. Prod.* ix. 160.—Bras.
multijuga, Wall. Cat. n. 6503; Pl. Rar. As. i. 81. t. 95, 96 = Pajanelia Rheedii.
muricata, DC. Rev. Bign. 22, ex Prod. ix. 173 = Pithecoctenium muricatum.
muricata, Herb. Madr. ex Wall. Cat. n. 6511 = Stereospermum xylocarpum.
myriantha, Cham. ex Steud. Nom. ed. II. i. 205 = Tecoma myriantha.
nematocarpa, Boj. Hort. Maurit. 219 = Stereospermum nematocarpum.
nervosa, Domb. ex DC. Prod. ix. 198 = Delostoma nervosum.
nigricans, *Regel, Cat. Pl. Hort. Aksakov.* 19.—Hab.?
nitidissima, *DC. Prod.* ix. 160.—Venezuela.
nitidula, Mart. ex DC. l. c. 181 = Lundia nitidula.
noctiflora, Buch.-Ham. ex Wall. Cat. sub n. 6499 = Stereospermum suaveolens.
nodosa, *Manso, Enum. Cat. Bras.* 40.—Bras.
noterophila, *DC. Prod.* ix. 148.—Bras.

BIGNONIA :—
obliqua, *H. B. & K. Nov. Gen. et Sp.* iii. 135.—Venezuela.
obovata, Hook. & Arn. Bot. Becch. Voy. 439 = Stemmadenia pubescens.
obovata, Spreng. Syst. ii. 830 = Spathodea obovata.
obovata, Vell. Fl. Flum. 251; vi. t. 45 = Jacaranda subrhombea.
obtusifolia, Lam. Encyc. i. 418 = Spathodea obtusifolia *seu* Tabebuia uliginosa.
occidentalis, *Salzm. ex DC. Prod.* ix. 160.—Bras.
odorata, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 293.—Porto Rico.
ophthalmica, *Chisholm, ex Steud. Nom.* ed. II. i. 205.—Guiana.
orbiculata, Jacq. Enum. Pl. Carib. 25 = Anemopaegma orbiculatum.
ornata, *Linden, Cat.* n. 21 (1867) 1; *E. Morr. Belg. Hortic.* xvii. (1867) 102.—Bras.
oxyphylla, *DC. Prod.* ix. 169 (*Cf. Fl. Brit. Ind.* iv. 376).—Ind. or.?
pachyptera, DC. Rev. Bign. 21; ex Prod. ix. 174 (= *Pachyptera puberula*).—Bras.
Pajanelia, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 515 = Pajanella Rheedii.
pallida, *Lindl. Bot. Reg.* t. 965.—Ind. occ.
pandorana, Andr. Bot. Rep. t. 86 = Tecoma australis.
Pandorea, Vent. Jard. Malm. t. 43 = Tecoma australis.
paniculata, Linn. Sp. Pl. 623 = Amphilophum paniculatum.
pannosa, *DC. Prod.* ix. 148.—Peruv.
Parkeri, *DC. l. c.* 157.—Guiana.
parviflora, Rottl. ex Wall. Cat. sub n. 6501 = Stereospermum chelonoides.
parviflora, Salzm. ex DC. Prod. ix. 183 = Arrabidaea Salzmanni.
patellifera, *Schlecht. in Linnaea*, viii. (1833) 516.—Mexic.
Patrisiana, DC. Prod. ix. 174 = Tecoma Patrisiana.
pedunculata, *Vell. Fl. Flum.* 250; vi. t. 41.—Bras.
pentandra, Lour. Fl. Cochin. ii. 379 = Oroxylum indicum.
pentaphylla, Linn. Sp. Pl. ed. II. 870 = Tabebuia pentaphylla.
pentaphylla, Salzm. ex DC. Prod. ix. 219 = Tecoma Salzmanni.
perforata, *Cham. in Linnaea*, vii. (1832) 667.—Bras.
perianthomega, *Vell. Fl. Flum.* 248; vi. t. 34.—Bras.
peruviana, *Linn. Sp. Pl.* 625.—Peruv.
petiolaris, DC. Cat. Bign. 22; ex Prod. ix. 174 = Tecoma Leucoxylon.
phaseoloides, *Cham. in Linnaea*, vii. (1832) 698.—Bras.
physaloides, *Cham. l. c.* 668.—Bras.
picta, *H. B. & K. Nov. Gen. et Sp.* iii. 136.—Guiana.
picta, Lindl. Bot. Reg. (1842) t. 45 = Lindleyi.
pilosa, Dietr. ex Steud. Nom. ed. II. i. 205 = villosa.
pilulifera, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Guiana.
platyphylla, Cham. in Linnaea, vii. (1832) 679 = brachypoda.
Poeppigii, DC. Rev. Bign. ex Prod. ix. 174 = Anemopaegma Poeppigii.
populifolia, *DC. Prod.* ix. 159.—Bras.
Porteriana, Wall. Cat. n. 6509 = Stereospermum glandulosum.
Prieurei, *DC. Prod.* ix. 154.—Guiana.
procera, Willd. Sp. Pl. iii. 307 = Jacaranda Copaia.
pseudounguis, *Desf. Cat. Hort. Par.* ed. III. 397.—Ind. occ.
pterocarpa, Cham. in Linnaea, vii. (1832) 673 = Cuspidaria trifoliata.
pubescens, *Linn. Sp. Pl.* ed. II. 870.—Mexic.
pubescens, Pav. ex DC. Prod. ix. 148 = pannosa.
pulchella, *Cham. in Linnaea*, vii. (1832) 663.—Bras.
pulchra, *Cham. l. c.* 708.—Bras.
punctata, DC. Rev. Bign. ex Prod. ix. 160 = occidentalis.
punicea, *Mart. Herb. Fl. Bras.* 255.—Bras.
purpurea, *Lodd. ex Sweet, Hort. Brit.* 283, nomen; *Bot. Mag.* t. 5800.—Am. austr.
pyramidata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = Tabebuia pyramidata.
quadrangularis, *Lockart, ex Steud. Nom.* ed. II. i. 205.—Ins. Trinitat.
quadrilocularis, Roxb. Pl. Corom. ii. 24. t. 145 = Heterophragma Roxburghii.
quadripinnata, Blanco, Fl. Filip. ed. I. 499 = Oroxylum indicum.

BIGNONIA :—
quadrivalvis, *Jacq. Fragm.* 37.—Venezuela.
Quercus, Lam. Encyc. i. 417 = Catalpa longissima.
quinquefolia, Vell. Fl. Flum. 252; vi. t. 50 = Cybistax antisyphilitica.
racemosa, *Lam. Encyc.* i. 424.—Madag.
radiata, Linn. Sp. Pl. 624 = Argylia Feuillei.
radicans, Linn. l. c. = Tecoma radicans.
ramiflora, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 381 = Colea mauritiana.
Regnelliana, *Sond. in Linnaea*, xxii. (1849) 558.—Bras.
Rego, Vell. Fl. Flum. 249 = Arrabidaea Sego.
reticulata, Hb. Madr. ex Wall. Cat. n. 6511 = Stereospermum xylocarpum?
Reveili, *Hamon, in Bull. Soc. Hortic. Prat. Rhône*, (1851) 149. t. 3.—Hab.?
rhodosantha, *Zipp. ex Blume, Rumphia*, iv. 35.—N. Guin.
rigescens, Jacq. Hort. Schoenb. ii. 44. t. 210 = Distictis rigescens.
riparia, *H. B. & K. Nov. Gen. et Sp.* iii. 138.—N. Granat.
rivularis, Hort. Kew. (1830) ex DC. Prod. ix. 199 = Cybistax antisyphilitica.
rosea, DC. Rev. Bign. ex Prod. ix. 174 = Arrabidaea rosea.
rosea, Pav. ex DC. Prod. ix. 198 = Delostoma dentatum.
rotundata, *DC. l. c.* 163.—Bras.
rufescens, DC. Rev. Bign. ex Prod. ix. 174 = Pithecoctenium rufescens.
rufinervis, Hoffmgg. ex DC. Prod. ix. 213 = Callichlamys riparia.
rugosa, *Schlecht. in Linnaea*, xxvi. (1853–55) 656.—N. Granat.
rupestris, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 179.—Bras.
Sagraeana, *DC. Prod.* ix. 148.—Cuba.
salicifolia, Buch.-Ham. ex Wall. Cat. n. 6498 = Tecoma undulata.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 133.—Venezuela.
Salzmanni, DC. Rev. Bign. ex Prod. ix. 174 = Adenocalymna Salzmanni.
samydoides, *Cham. in Linnaea*, vii. (1832) 669.—Bras.
sambucifolia, *Hort. ex Carr. Rev. Hortic.* (1878) 50.—Hab.?
sarmentosa, Bertol. f. Fl. Guatim. 25 = aequinoctialis.
scandens, Vell. Fl. Flum. 246; vi. t. 22 = Anemopaegma racemosum.
Sceptrum, *Cham. in Linnaea*, vii. (1832) 710.—Bras.
Sego, Vell. Fl. Flum. vi. t. 39 = Arrabidaea Agnus-castus.
Selloi, *Spreng. Syst.* ii. 831.—Bras.
sempervirens, Linn. Sp. Pl. 623 = Gelsemium sempervirens.
serrata, Pav. ex G. Don, Gen. Syst. iv. 228 = castaneaefolia.
serratifolia, Vahl, Ecl. Am. ii. 46 = Tecoma serratifolia.
Serratula, Wall. ex DC. Prod. ix. 206 = Stereospermum serrulatum.
serrulata, Steud. Nom. ed. II. i. 205, sphalm. = praec.
Sieberi, DC. Rev. Bign. ex Prod. ix. 175 = Arrabidaea Sieberi.
simplex, *A. Rich. Fl. Cub. Fanerog.* ii. 104.—Cuba.
simplicifolia, Pav. ex DC. Prod. ix. 198 = Delostoma integrifolium.
Sinclairii, *Benth. Bot. Voy. Sulph.* 129 = laurifolia.
sorbifolia, Salisb. Prod. 106 = Tecoma stans.
sordida, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1158.—Guiana.
spathacea, Linn. f. Suppl. 283 = Dolichandrone Rheedii.
spathacea, Roxb. Pl. Corom. ii. 24. t. 144 = Dolichandrone falcata.
speciosa, *R. Grah. in Bot. Mag.* t. 3888.—Bras.
spectabilis, Salisb. Prod. 106 = Catalpa bignonioides.
spectabilis, Vahl, Symb. Bot. iii. 80 = aequinoctialis.
spicata, *Lockhart, ex Steud. Nom.* ed. II. i. 205.—Ins. Trinitat.
Splendor-sylvae, *Mart. ex DC. Prod.* ix. 154.—Bras.
squalus, Vell. Fl. Flum. 244; vi. t. 13 = Pithecoctenium squalum.
squamellulosa, DC. Rev. Bign. ex Prod. ix. 174 = Tecoma squamellulosa.
staminea, *Lam. Encyc.* i. 421.—Ins. S. Doming.
stans, Linn. Sp. Pl. ed. II. 871 = Tecoma stans.
stipulata, Roxb. Hort. Beng. 47; Fl. Ind. iii. 108 = Dolichandrone stipulata.

BIGNONIA :—
striata, DC. Rev. Bign. ex Prod. ix. 174 (= *Pachyptera striata*).—Bras.
suaveolens, Roxb. Hort. Beng. 47; Fl. Ind. iii. 104 = Stereospermum suaveolens.
suberosa, Roxb. ll. cc. 47, 111 = Millingtonia hortensis.
subincana, *Hemsl. Biol. Centr. Am. Bot.* ii. 491.—Panama.
subvernicosa, DC. Rev. Bign. ex Prod. ix. 220 = Tecoma subvernicosa.
Tababuya, Vell. Fl. Flum. 251; vi. t. 53 = Tabebuia uliginosa.
Tagada, Roxb. ex Wall. Cat. n. 6501 = Stereospernum chelonoides.
tecomoides, DC. Prod. ix. 166 = Tecoma mollis.
tenuiflora, *DC. l. c.*—Bolivia.
Telfairiae, *Boj. ex Hook. Bot. Mag.* t. 2976.—Madag.
tenuisiliqua, *Vahl, Ecl. Am.* ii. 43.—Am. austr.
ternata, G. Don, Gen. Syst. iv. 221 = maranhamensis.
ternata, *Vell. Fl. Flum.* 246; vi. t. 21.—Bras.
tetragona, Wall. ex DC. Prod. ix. 211 = Stereospermum tetragonum.
tetragonocaulos, *DC. Prod.* ix. 169.—Bras.
tetragonoloba, *Jacq. Fragm.* 36.—Venezuela.
tetraquetra, *Cham. in Linnaea*, vii. (1832) 675.—Bras.
thyrsoidea, *DC. Prod.* ix. 158.—Bras.
tiliaefolia, *H. B. & K. Nov. Gen. et Sp.* iii. 136.—Am. austr.
tinctoria, *Arruda, in Koster, Trav. Braz.* 497.—Bras.
tomentosa, Pav. ex DC. Prod. ix. 219 = Tecoma heteropoda.
tomentosa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = mollis.
tomentosa, Thunb. Fl. Jap. 252 = Paulownia imperialis.
triantha, DC. Rev. Bign. ex Prod. ix. 148 = exoleta.
trichoclada, *DC. l. c.* 158.—Bras.
triflora, Pav. ex DC. l. c. 148 (est Rubiacea).
triflora, *Vell. Fl. Flum.* 249; vi. t. 38.—Bras.
trifoliata, Vell. l. c. 245; vi. t. 16 = Adenocalymna longeracemosum.
triphylla, Vell. l. c. 244; vi. t. 14 = triternata.
triphylla, Linn. Sp. Pl. ed. II. 870 = Tabebuia triphylla.
triphylla, Willd. ex DC. Prod. ix. 151 = B. Chica.
tripinnata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 8.—Malaya.
triplinervia, *Mart. ex DC. Prod.* ix. 153.—Bras.
triternata, *Steud. Nom.* ed. II. i. 205.—Bras.
Tropaeolum, Jacquem. ex DC. Prod. ix. 223, in nota = Tecoma undulata.
tuberculata, Roxb. ex DC. Prod. ix. 177 = Oroxylum indicum.
tuberculosa, Vell. Fl. Flum. 251; vi. t. 46 = Zeyheria montana.
tulipifera, Schum. Beskr. Guin. Pl. 273 = Spathodea tulipifera.
Tweediana, *Lindl. Bot. Reg.* (1840) t. 45.—Bras.
uliginosa, Gomez, Obs. ii. 7. t. 2; Mem. Ac. Lisboa, iii. (1812) Mem. dos Corresp. 57 = Tabebuia uliginosa.
umbellulata, *DC. Prod.* ix. 148.—Bras.
umbrosa, *H. B. & K. Nov. Gen. et Sp.* iii. 138.—N. Granat.
uncata, Andr. Bot. Rep. t. 530 = Macfadyena uncinata.
uncata, Sims, Bot. Mag. t. 1511 = Spathodea uncata.
uncinata, G. F. W. Mey. Prim. Fl. Esseq. 210 = Macfadyena uncinata.
undulata, Sm. Exot. Bot. i. 35 = Tecoma undulata.
unguiculata, *Vell. Fl. Flum.* 248; vi. t. 33.—Bras.
unguiculata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = B. Unguis-cati.
Unguis, Linn. Syst. ed. X. xiii. n. 3 = B. Unguis-cati.
Unguis-cati, *Linn. Sp. Pl.* 623.—Am. trop.
Vargasiana, DC. Rev. Bign. ex Prod. ix. 175 = Anemopaegma Vargasianum.
variabilis, *Jacq. Hort. Schoenb.* ii. 45. t. 212.—Venezuela.
variabilis, Mart. Herb. Fl. Bras. 256 = varians.
variabilis, Sieber, ex E. Mey. in Nov. Act. Nat. Cur. xii. (1825) 779 = Hostmanni.
varians, *DC. Prod.* ix. 149.—Bras.
velutina, *DC. l. c.* 162.—Bras.
venusta, *Ker-Gawl. in Bot. Reg.* t. 249.—Bras.
verrucifera, *Schlecht. in Linnaea*, xxvi. (1853–55) 655. N. Granat.
verrucosa, *H. B. & K. Nov. Gen. et Sp.* iii. 137.—Venezuela.
villosa, Bert. ex Spreng. Syst. ii. 830 = Balbisiana.
villosa, *Vahl, Ecl. Am.* ii. 43.—Venezuela.

BIGNONIA:—
viminalis, H. B. & K. Nov. Gen. et Sp. iii. 132 = Tecoma viminalis.
violacea, *DC. Prod.* ix. 156.—Bolivia.
virginalis, Mart. ex DC. l. c. 181 = Lundia virginalis.
viridiflora, Lodd. Bot. Cab. xi. (1825) t. 1026 = Cybistax antisyphilitica.
Vitalba, Cham. in Linnaea, vii. (1832) 699 = Pithecoctenium Vitalbae.
xylocarpa, Roxb. Hort. Beng. 47; Fl. Ind. iii. 108 = Stereospermum xylocarpum.

BIKAI, Adans. Fam. ii. 67 (1763) = **Heliconia**, Linn. (Scitam.).

BIHANIA, Meissn. in DC. Prod. xv. i. 96 (1861) = **Eusideroxylon**, Teijsm. (Laurin.).
borneensis, Meissn. l. c.—Borneo.

BIKERA, Adans. Fam. ii. 130 (1763) = **Tetragonotheca**, Dill. (Compos.).

BIKKIA, Reinw. ex Blume, Bijdr. 1017, in syn. (1826); et in Syll. Ratisb. ii. (1828) 8. *RUBIACEAE*, Benth. & Hook. f. ii. 46.
BIKKIOPSIS, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 404.
CORMIGONUS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 83.
GRISIA, Brongn. in Bull. Soc. Bot. Fr. xii. (1865) 405.
TATEA, Seem. Fl. Vit. 125 (1866).
australis, DC. Prod. iv. 405 = grandiflora.
Forsteriana, *Brongn. in Bull. Soc. Bot. Fr.* xiii. (1866) 42.—Ins. Fiji.
Gaudichaudiana, *Brongn. l. c.* 43.—N. Guinea.
grandiflora, *Reinw. ex Blume, Bijdr.* 1017, *in syn.; et in Syll. Ratisb.* ii. (1828) 8.—Java; Ins. Amic.
Guilloviana, *Brongn. in Bull. Soc. Bot. Fr.* xiii. (1866) 42.—N. Guinea.
Hombroniana, *Brongn. l. c.* 43.—Ins. Pacif.
mariannensis, *Brongn. l. c.* 42.—Ins. Marian.

BIKKIOPSIS, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 404 = **Bikkia**, Reinw. (Rubiac.).
Pancheria, Brongn. l. c.—N. Caled.

BIKUKULLA, Adans. Fam. ii. 23 (1763) = **Diclytra**, Borckh. (Papav.).

BILABIUM, Miq. Fl. Ind. Bat. ii. 730 (1856). *DIDYMOCARPEAE*, Benth. & Hook. f. Gen. ii. 996.
limans, *Miq. l. c.*—Malaya.

BILABRELLA, Lindl. Bot. Reg. sub t. 1701 (1835) = **Habenaria**, Willd. (Orchid.).
falcicornis, Lindl. l. c. = H. tetrapetala.
falciformis, Bolus, in Journ. Linn. Soc. xix. (1882) 340 = praec.
salsicornis, Steud. Nom. ed. II. i. 205, sphalm. = praec.

BILAMISTA, Rafin. New Fl. Am. iv. 92 (1836) = **Gentiana**, Tourn.
grandiflora, Rafin. l. c. 93 = G. grandiflora?

BILDERDYKIA, Dum. Florul. Belg. 18 (1827) = **Polygonum**, Tourn.
Convulvulus, Dum. l. c. = P. Convulvulus.
dumetorum, Dum. l. c. = P. dumetorum.

BILITALIUM, Buch.-Ham. Journ. Mysore, 185 (1807), nomen.
farinosum, Buch.-Ham. l. c. (Quid?).—Ind. or.

BILLARDIERA, Moench, Meth. 369 (1794) = **Verbena**, Linn.
explanata, Moench, l. c. = V. Aubletia.

BILLARDIERA, Sm. Bot. Nov. Holl. i. t. 1 (1793). *PITTOSPOREAE*, Benth. & Hook. f. i. 132.
LABILLARDIERA, Roem. & Schult. Syst. v. p. xxvi. 330 (1819).
angustifolia, DC. Prod. i. 345 = scandens.
brachyantha, F. Muell. ex Klatt, in Linnaea, xxviii. (1856) 571 = scandens.
canariensis, Wendl. Beobacht. 43 = scandens.
coriacea, *Benth. Fl. Austral.* i. 124.—Austral.
cymosa, *F. Muell. in Trans. Vict. Inst.* i. (1855) 29.—Austral.

BILLARDIERA:—
cymosa, Klatt, in Linnaea, xxviii. (1856) 571 = cymosa.
daphnoides, *Knowles & Westc. Flor. Cab.* ii. (1838) 60.—Austral.?
elegans, *F. Muell. Pl. Vict.* i. 78.—Austral.
floribunda, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 1.—Austral.
fusiformis, Labill. Nov. Holl. Pl. t. 90 = Sollya heterophylla.
grandiflora, Putterl. Stirp. Nov. Decad. 48 = scandens.
Hambruchiana, Seem. & Schmidt, in Flora, xxvii. (1844) II. 496 = Sollya heterophylla.
latifolia, Putterl. Stirp. Nov. Decad. 47 = scandens.
Lehmanniana, *F. Muell. Pl. Vict.* i. 78.—Austral.
longiflora, *Labill. Nov. Holl. Pl.* i. 64. t. 89.—Tasmania.
macrantha, Hook. f. Fl. Tasm. i. 37 = longiflora.
melocarpa, *Hort. ex Steud. Nom.* ed. II. i. 206.—Austral.
mutabilis, *Salisb. Parad. Lond.* t. 48.—Austral.
ovalis, Lindl. Bot. Reg. t. 1719 = longiflora.
parviflora, DC. Prod. i. 346 = Marianthus tenuis.
pseudocymosa, Klatt, in Linnaea, xxviii. (1856) 571 = cymosa.
rosmarinifolia, *DC. Prod.* i. 345.—Austral.
scandens, *Sm. Exot. Bot.* i. t. 1.—Austral.
sericophora, F. Muell. in Linnaea, xxv. (1852) 371 = cymosa.
variifolia, *DC. Prod.* i. 346.—Austral.
versicolor, F. Muell. ex Klatt, in Linnaea, xxviii. (1856) 751 = cymosa.

BILLARDIERA, Vahl, Eclog. Am. i. 13. t. 10 (1796) = **Coussaria**, Aubl. (Rubiac.).
paniculata, Vahl, l. c. = C. Froelichia.

BILLBERGIA, Thunb. Dec. Pl. Bras. 30 (1823). *BROMELIACEAE*, Benth. & Hook. f. iii. 664.
CREMOBOTRYS, Beer, in Flora, xxxvii. (1854) 348.
EUCALLIAS, Rafin. Fl. Tellur. iv. 25 (1836).
HELICODEA, Lem. Illustr. Hortic. xi. (1864) t. 421.
JONGHEA, Lem. Jard. Fleur. ii. sub t. 180–181 (1852).
LIBONIA, Lem. Illustr. Hortic. ii. (1855) t. 48.
acaulis, Hort. ex Vis.—Cf. Otto & Dietr. Allgem. Gartenz. xvi. (1848) 29 = Cryptanthus zonatus.
amabilis, Beer, Bromel. 118 = vittata.
amoena, Lindl. Bot. Reg. sub t. 1068 = speciosa.
angustifolia, *C. Koch, Wochenschr.* ix. (1866) 181.—Hab.?
atro-rosea, *Drapiez, Encyc. et Fl. des Serr. Angl.* (1834) f. 1.—Bras.
Bakeri, *E. Morr. in Belg. Hortic.* xxx. (1880) 166. t. 8.—Bras.
Baraquiniana, Lem. Illustr. Hortic. (1864) 421 = decora.
bicolor, Lodd. Bot. Cab. t. 1819 = pyramidalis.
bicolor, Schult. f. Syst. vii. 1255 = Rhodostachys bicolor.
bifrons, Lindl. in Journ. Hort. Soc. viii. (1853) 54 = Pitcairnea bracteata?
bivittata, Hook. Bot. Mag. t. 5270 = Cryptanthus bivittatus.
Breauteana ×, André, in Rev. Hortic. (1884) 117.
Brongniarti, Regel, Gartenfl. (1875) 166; et in Act. Hort. Petrop. iii. II. (1875) 283 = Portea kermesina.
Chantini, *Carr. in Rev. Hortic.* (1878) 112.—Bras.
chlorocyanea, De Vriese, Hort. Acad. Lugd. Bat. Pl. Rar. (1854).—Cf. Linnaea, xxvi. (1853–55) 755 = speciosa.
chlorostica, Hort. Saunders; ex Gard. Chron. (1871) 1425 = Saundersii.
clavata, Lindl. Bot. Reg. sub t. 1068 = Aechmea bromeliaefolia.
coelestis, Hort. ex Morr. in Fl. des Serres, xxi. (1875) 1 = Aechmea coelestis.
Collevi ×, A. Van Geert, Cat. n. 81. p. 91; Belg. Hortic. xxxii. (1882) 334.
commixta, C. Koch, Wochenschr. iii. (1860) 146 = zebrina?
contracta, Mart. ex Schult. f. Syst. vii. II. 1263 = Aechmea contracta.
corymbosa, *Mart. ex Schult. f. l. c.* 1268.—Bras.
Croyiana, De Jongh, ex Lem. Jard. Fleur. iv. (1854) t. 413 = pyramidalis.
cruenta, *Hook. Bot. Mag.* lvi. t. 2892.—Bras.
decora, *Poepp. & Endl. Nov. Gen.* 42.—Peruv.
discolor, Beer, Bromel. 121 = speciosa.
distichostachya, Lem. Jard. Fleur. ii. (1852) Misc. 96 = Aechmea distichantha.

BILLBERGIA:—
elegans, Mart. ex Schult. f. Syst. vii. II. 1265 = speciosa.
Euphemiae, *E. Morr. Belg. Hortic.* xxii. (1872) 1. t. 1.—Bras.
excellens, *Miq. in Journ. Bot. Néerl.* i. (1861) 34.—Guiana.
fasciata, Lindl. Bot. Reg. xiii. t. 1130 = Aechmea fasciata.
fastuosa, Beer, Bromel. 110 = thyrsoidea.
filicaulis, *Griseb. in Goett. Nachr.* (1864) 13.—Venezuela.
Glazioviana, Regel, Gartenfl. (1885) 260 = Quesnelia strobilispica.
Glymiana, De Vriese, Jaarb. der Kon. Ned. Maatsch. Tuinb. (1853) 37 = Morelii.
Glymiana, Beer, Bromel. 119 = Morelii.
gracilis, Poepp. ex Beer, l. c. 254 = Aechmea tillandsioides.
granulosa, Brongn. in C. Koch, Wochenschr. iii. (1860) 145 = decora.
horrida, *Regel, Ind. Sem. Hort. Petrop.* (1856) 17; *Gartenfl.* (1857) 148.—Bras.
ianthina, Hort. ex E. Morr. Belg. Hortic. (1871) 2 = Leopoldi.
incarnata, *Schult. f. Syst.* vii. II. 1261.—Peruv.
iridifolia, *Lindl. Bot. Reg.* t. 1068.—Bras.
Joinvillei, Van Houtte, ex E. Morr. in Belg. Hortic. (1876) 161 = Rhodostachys pitcairniaefolia.
Joliboisi ×, A. Van Geert, Cat. n. 81. p. 31; Belg. Hortic. xxxii. (1882) 334.
lanuginosa, Hort. Hamb. ex C. Koch, Ind. Sem. Hort. Berol. (1856) App. 7 = Aechmea nudicaulis.
latifolia, Hort. Belg. ex Heynh. Nom. ii. 69 = Aechmeae sp.?
Legrelliana, Hort. ex Baker, in Saund. Ref. Bot. t. 285 = Ortgiesiae sp.
Leopoldi, *Linden, in Rev. Hortic.* (1869) 87; *E. Morr. in Belg. Hortic.* xxi. (1871) 1. t. 1.—N. Granat.; Bras.
Leopoldii, Hort. Belg. ex C. Koch, Ind. Sem. Hort. Berol. (1856) 5 = vittata.
Liboniana, *De Jongh. Journ. Hort. Prat.* (Mars. 1851); *et Lem. Jard. Fleur.* ii. (1852) t. 197.—Bras.
Lietzei, *E. Morr. in Belg. Hortic.* xxxi. (1881) 97. t. 5–7.—Bras.
Loddigesii, Steud. Nom. ed. II. i. 206 = pyramidalis.
longifolia, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1856) 5 = thyrsoidea.
lutea, Schult. f. Syst. vii. 1258 = Aechmea nudicaulis.
macrocalyx, *Hook. Bot. Mag.* t. 5114.—Bras.
marmorata, *Lem. Illustr. Hortic.* ii. (1855) t. 48.—Bras.
Mertensii, Miq. in Linnaea, xviii. (1844) 376 = Aechmea Mertensii.
Meyendorfii, Regel, Ind. Sem. Hort. Petrop. (1857) 27 = Karatas Carolinae.
miniata-rosea, Lem. Illustr. Hortic. ii. (1855) Misc. 94 = thyrsoidea.
mitis, *Mart. ex Schult. f. Syst.* vii. 1267.—Bras.
Moreliana, Brongn. ex Rev. Hortic. Sér. III. iii. (1849) 82 = Morelii.
Moreliana, Lem. Jard. Fleur. ii. (1852) t. 138 = vittata.
Morelii, *Brongn. in Portef. Hort.* ii. (1848) 97; *et E. Morr. Belg. Hortic.* x. (1860) 161.—Bras.
nudicaulis, Lindl. Bot. Reg. sub t. 1068 = Aechmea nudicaulis.
nuptialis, Hort. ex E. Morr. in Belg. Hortic. (1871) 2 = Leopoldi.
nutans, *H. Wendl. in Regel, Gartenfl.* (1869) 162.—Bras.
Oberthuri ×, A. Van Geert. Cat. n. 81. p. 92; Belg. Hortic. xxxii. (1882) 334.
oblonga, Mart. ex Schult. f. Syst. vii. 1256 (= Aechmeae sp.)—Bras.
odora, Miq. in Linnaea, xviii. (1844) 377 = Aechmea odora.
olens, Hook. f. Bot. Mag. t. 5502 = Karatas Carolinae.
pallescens, Baker, in Hook. f. l. c. t. 6342 = Bakeri.
pallescens, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1856) App. 5 = speciosa.
pallida, Beer, Bromel. 121 = speciosa.
pallidiflora, *Liebm. Ind. Sem. Hort. Hafn.* (1854) 26.—Nicaragua.
paniculata, Mart. ex Schult. f. Syst. vii. II. 1268 = Aechmea coelestis?
parviflora, Mart. ex Schult. f. l. c. 1270 = Aechmea parviflora.

BILLBERGIA:—
patentissima, Mart. ex Schult. f. Syst. vii. 11. 1270 = Aechmea patentissima.
Paxtonii, Beer, Bromel. 113 = thyrsoidea.
polystachya, Lindl. & Paxt. in Paxt. Flow. Gard. iii. (1852-53) 33. t. 80 = Aechmea distichantha.
Porteana, *Brongn. ex Beer, Bromel.* 115.—Bras.
pulcherrima, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1856) App. 6 = vittata.
punicca, *Beer, Bromel.* 112.—Hab.?
purpurea, *Beer, l. c.* 126 (= Pitcairniae sp. ?).—Am. trop.
purpureo-rosea, Hook. Bot. Mag. t. 3304 = Aechmea suaveolens.
pyramidalis, Forbes, Hort. Woburn. 64 = Aechmea nudicaulis.
pyramidalis, *Lindl. Bot. Reg.* sub t. 1068.—Bras.
pyramidata, Beer, Bromel. 123 = Aechmea nudicaulis.
Quesneliana, *A. Brongn. in Ann. Sc. Nat.* Sér. II. xv. (1841) 371.—Guiana.
Rancougnei ×, André, in Rev. Hortic. (1884) 524.
Reichardtii, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 115.—Bras.
rhedonensis ×, A. Van Geert, Cat. n. 81. p. 92; Belg. Hortic. xxxii. (1882) 334.
rhodocyanea, Lem. in Fl. des Serres, Sér. I. iii. (1847) 207 = Aechmea fasciata.
Rohaniana, De Vriese, in Linnaea, xxvi. (1853-55) 756 = vittata.
rosea, Beer, Bromel. 128 = Porteana.
roseo-marginata, Hort. ex C. Koch, Wochenschr. (1867) 141 = Quesnelia rufa.
rubra-marginata, Hort. ex Carr. in Rev. Hortic. (1880) 70 = Quesnelia rufa.
Sanderiana, *E. Morr. in Belg. Hortic.* xxxiv. (1884) 17.—Bras.
Saundersii, *Hort. Bull, ex Floral Mag.* N. S. t. 106.—Bras.
saxatilis, *Beer, Bromel.* 127.—Bras.
Skinneri, Hort. ex Carr. in Rev. Hortic. (1880) 70 = Quesnelia rufa.
speciosa, *Thunb. Dec. Pl. Bras.* iii. 30.—Bras.
sphacelata, Schult. f. Syst. vii. 1269 = Greigia sphacelata.
splendida, Lem. Jard. Fleur. ii. (1852) t. 180–181 = thyrsoidea.
tetrantha, Beer, Bromel. 127 = Tillandsia tetrantha.
thyrsoidea, *Mart. ex Schult. f. Syst.* vii. 11. 1260.—Bras.
tillandsioides, Mart. ex Schult. f. l. c. 1269 = Aechmea tillandsioides.
tinctoria, Mart. ex Schult. f. l. c. 1256 = Aechmea bromeliaefolia.
variegata, Schult. f. l. c. 1262 = speciosa.
variegata, *Morr. in Belg. Hortic.* xxxi. (1881) 73.—Bras.
violacea, *Beer, Bromel.* 116.—Guiana.
viridiflora, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 154.—Hab.?
vittata, *Brongn. ex Morel, Portef. Hortic.* ii. (1848) 353.—Bras.
Wetherelli, *Hook. Bot. Mag.* t. 4835.—Bras.
zebrina, *Lindl. Bot. Reg.* sub t. 1068.—Bras.
zonata, Linden, Cat. n. 6 (1851) 20 = vittata.

BILLIA, Peyr. in Bot. Zeit. xvi. (1858) 153; xvii. (1859) 221 = **Aesculus**, Linn. (Sapind.).
columbiana, Planch. & Linden, ex Triana & Planch. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 367.—N. Granat.
Hippocastanum, Peyr. in Bot. Zeit. xvi. (1858) 153 = Aesculus mexicana.

BILLIOTIA, G. Don, Gen. Syst. iii. 560 (1834) = **Billiottia**, DC. (Rubiac.).

BILLIOTIA, Reichb. Nom. 176 (1841) = BILLOTTIA, R. Br. = **Agonis**, Lindl. (Myrtac.).

BILLIOTTIA, Endl. Gen. 1229 (1840) = BILLOTTIA, R. Br. = **Agonis**, Lindl. (Myrtac.).

BILLIOTTIA, DC. Prod. iv. 618 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 80.
BILLIOTIA, G. Don, Gen. Syst. iii. 560 (1834).
MELANOPSIDIUM, Colla, Hort. Ripul. 88. t. 35 (1824).

BILLIOTTIA:—
VIVIANIA, Colla, in Mém. Soc. Linn. Par. iv. (1826) 25. t. 2.
polytrichoides, Steud. Nom. ed. II. ii. 114, sphalm. = seq.
psychotrioides, *DC. Prod.* iv. 618.—Bras.

BILLOTIA, G. Don, Gen. Syst. ii. 810, 827 (1832) = BILLOTTIA, R. Br. = **Agonis**, Lindl. (Myrtac.).

BILLOTIA, Sch. Bip. in Herb. & Winck. Jahrb. Pharmac. iv. (1841) in nota 155 = **Crepis**, Linn. (Compos.).
alpina, Sch. Bip. l. c. = C. alpina.

BILLOTTIA, R. Br. in Journ. Roy. Geogr. Soc. i. (1832) 19 = **Agonis**, DC. (Myrtac.).
elliptica, Swect, Hort. Brit. ed. III. 249 = Leptospermum ellipticum.
flexuosa, R. Br. in. Journ. Roy. Geogr. Soc. i. (1832) 19 = A. flexuosa.
hypericifolia, Hcynh. Nom. ii. 69 = A. hypericifolia.
linearifolia, R. Br. in Journ. Roy. Geogr. Soc. i. (1832) 19 = A. linearifolia.
marginata, R. Br. l. c. 19 = A. marginata.
ovatifolia, Sweet, Hort. Brit. ed. III. 249.—Austral.
saligna, Sweet, l. c.—Austral.
theaeformis, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1380 = A. hypericifolia.

BILLOTTIA, Colla, Hort. Ripul. 20. t. 23 (1824) = **Calothamnus**, Labill. (Myrtac.).
acerosa, Colla, l. c. = C. quadrifida.

BIMA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1 (Quid?).

BINARIA, Rafin. Sylva Tellur. 122 (1838) = **Bauhinia**, Linn. (Legumin.).
cumanensis, Rafin. l. c. = Bauhinia cumanensis.

BINDERA, Rafin. New Fl. Am. iv. 71 (1836) = **Aster**, Tourn. (Compos.).
ciliata, Rafin. l. c. = A. ciliatus.

BINECTARIA, Forsk. Fl. Aegypt. Arab. 82 (1775) = **Imbricaria**, Comm. (Sapot.)

BINNENDIJKIA, Kurz, in Tijdschr. Nederl. Ind. xxviii. (1865) 164 = **Leptonychia**, Turcz. (Sterculiac.).
trichostylis, Kurz, l. c. = L. glabra.

BIONDEA, Usteri, Ann. Bot. x. (Neue Ann. iv.) 90 (1794) = BLONDEA, Rich. = **Sloanea**, Linn. (Tiliac.).

BIONIA, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 130 = **Camptosema**, Hook. & Arn. (Legumin.).
acuminata, Benth. l. c. = C. acuminatum.
bella, Mart. in Walp. Rep. i. 764 = C. bellum.
coccinea, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 380 = C. coccineum.
coriacea, Benth. l. c. 130 = C. coriaceum.
marginata, Mart. ex Benth. l. c. = C. coriaceum.
nitens, Benth. l. c. = C. coccineum.
rigida, Benth. l. c. = C. coriaceum.

BIOPHYTUM, DC. Prod. i. 689 (1824). *GERANIACEAE*, Benth. & Hook. f. i. 276.
abyssinicum, *Steud. ex A Rich. Tent. Fl. Abyss.* i. 122.—Abyss.
adiantoides, *Wight, ex Edgew. & Hook. f. in Fl. Brit. Ind.* i. 437.—Ind. or.
Apodiscias, *Edgew. & Hook. f. l. c.*—Ind. or.
Candolleanum, Wight, Ill. i. 161. t. 62 = sensitivum.
castum, G. Don, Gen. Syst. i. 753 = Oxalis casta.
Cumingianum, Edgew. & Hook. f. in Fl. Brit. Ind. i. 437 = sensitivum.
Cumingii, Klotzsch, in Peters, Reise Mossamb. Bot. 85 = sensitivum.
dendroides, *DC. Prod.* i. 690.—N. Granat.
dormiens, G. Don, Gen. Syst. i. 753 = Oxalis dormiens.
fruticosum, *Blume, Bijdr.* 242.—Java.
intermedium, *Wight, Illustr.* i. 162.—Zeylan.
mimosoides, G. Don, Gen. Syst. i. 753 = Oxalis mimosoides.

BIOPHYTUM:—
nervifolium, Thw. Enum. Pl. Zeyl. 64 = sensitivum.
nudum, *Edgew. & Hook. f. Fl. Brit. Ind.* i. 438.—Zeylan.
Petersianum, Klotzsch, in Peters, Reise Mossamb. Bot. 81. t. 15 = sensitivum.
polyphyllum, *Munro, in Wight, Illustr.* i. 162.—Ind. or.
poterioides, Edgew. ex Hook. f. Fl. Brit. Ind. i. 437 = sensitivum.
proliferum, *Edgew. & Hook. f. l. c.* 438.—Zeylan.
Reinwardtii, *Klotzsch, in Peters, Reise Mossamb. Bot.* 85.—As. trop.
sensitivum, *DC. Prod.* i. 690.—Cosmopol. trop.
sesbanioides, Edgew. ex Hook. f. Fl. Brit. Ind. i. 437 = sensitivum.
sessile, Wall. Cat. n. 4344 = B. Apodiscias.
somnians, G. Don, Gen. Syst. i. 753 = Oxalis somnians.
Umbraculum, *Welw. Apont.* 590.—Afr. trop.
verticillatum, Wight, Illustr. i. 162 = intermedium.

BIOTA, D. Don, in Lamb. Pinet. ed. II. ii. 129 (1828); Endl. Conif. 46 (1847) = **Thuya**, Linn. (Conif.).
coraeana, Sieb. ex Gord. Pinet. Suppl. 17 = T. orientalis.
dumosa, Hort. ex Carr. Conif. ed. II. 96 = T. orientalis.
ericoides, Hort. ex Carr. l. c. 141 = T. occidentalis.
excelsa, Hort. ex Gord. Pinet. Suppl. 17 = T. orientalis.
falcata, Hort. ex Carr. Conif. ed. II. 96 = T. orientalis.
Fortunei, Hort. ex Carr. l. c. 94 = T. orientalis.
freneloides, Hort. Belg. ex Gord. Pinet. 35 = T. orientalis.
funiculata, Hort. ex Gord. l. c. ed. II. 52 = T. orientalis.
glauca, Hort. ex Carr. Conif. ed. II. 100 = T. orientalis.
gracilifolia, Knight, Syn. Conif. 16 = T. orientalis.
intermedia, Hort. ex Gord. Pinet. 35 = T. orientalis.
japonica, Sieb. ex Gord. l. c. 33 = T. orientalis.
macrocarpa, Hort. ex Gord. l. c. ed. II. 422 = T. orientalis.
meldensis, Laws. ex Gord. l. c. 37 = T. meldensis.
nepalensis, Endl. ex Gord. l. c. 35 = T. orientalis.
orientalis, Endl. Conif. 47 = T. orientalis.
pendula, Endl. l. c. 49 = T. orientalis.
prostrata, Hort. ex Gord. Pinet. ed. II. 401 = T. orientalis.
pyramidalis, Carr. Conif. ed. I. 96 = T. orientalis.
semperaurea, Hort.—Cf. Gard. Chron. (1874) 60 = T. orientalis.
stricta, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 205 = T. orientalis.
tatarica, Lindl. & Gord. l. c. = T. orientalis.
variegata, Hort. ex Gord. Pinet. 34 = T. orientalis.
Wareana, Hort. ex Gord. l. c. Suppl. 104 = T. orientalis.
Zuccarini, Siebold, ex Carr. Conif. ed. II. 94 = T. orientalis.

BIOTIA, Cass. in Dict. Sc. Nat. xxxiv. 308 (1825) = **Madia**, Molina (Compos.).
viscosa, Cass. l. c. 309 = M. viscosa.

BIOTIA, DC. Prod. v. 264 (1856) = **Aster**, Tourn. (Compos.).
commixta, DC. Prod. v. 265 = A. Herveyi.
corymbosa, DC. l. c. = A. corymbosus.
crenata, Steud. Nom. ed. II. i. 154, in syn. = Mairia crenata.
discolor, Maxim. Prim. Fl. Amur. 146 = A. scaber. *Thunb.*
glomerata, DC. Prod. v. 265 = A. macrophyllus.
japonica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 170.—Japon.
latifolia, DC. Prod. v. 264 = A. macrophyllus.
macrophylla, DC. l. c. 265 = A. macrophyllus.
Schreberi, DC. l. c. 264 = A. macrophyllus.

BIPINNULA, Comm. ex Juss. Gen. 65 (1789). *ORCHIDEAE*, Benth. & Hook. f. iii. 618.
bonariensis, *Spreng. Syst.* iii. 745.—Reg. Argent.
Commersonii, Lindl. in Quart. Journ. Roy. Inst. N. S. i. (1827) 52 = bonariensis.

BIPINNULA :—
Giberti, *Reichb. f. in Linnaea*, xli. (1877) 51.—Reg. Argent.
mystacina, *Lindl. in Hook. Journ. Bot.* i. (1834) 5.—Chili.
plumosa, *Lindl. in Quart. Journ. Roy. Inst.* N. S. i. (1827) 51.—Chili.

BIPONTINIA, Alef. in Pollichia, xxii–xxiv. (1866) 121 = **Psoralea**, Linn. (Legumin.).
bituminosa, Alef. l. c. 123 = P. bituminosa.
corylifolia, Alef. l. c. 124 = P. corylifolia.

BIPOREIA, Thou. Gen. Nov. Madag. 14. (1806) = **Samadera**, Gaertn. (Simarub.).

BIRAMIA, Neraud, in Freyc. Voy. Bot. 28 (1826), nomen = **Macleania**, Hook. (Vaccin.).
tuberosa, Auct. ex Walp. Ann. i. 478 = M. insignis.

BIRCHEA, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 66. t. 10 = **Luisia**, Gaudich. (Orchid.).
teretifolia, A. Rich. l. c. 67. t. 10 = L. tenuifolia.

BIRIS, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 257; et Phil. Bot. ii. 97 (1791) = **Iris**, Tourn.
sibirica, Medic. ll. cc. = I. sibirica.

BIRNBAUMIA, Kostel. Ind. Pl. Hort. Prag. (1844) 21 = **Anisacanthus**, Nees (Acanth.).
quadrifida, Kostel. l. c. = A. virgularia.

BIROLIA, Bell. in Mém. Acad. Turin, Ser. I. xviii. (1808) 403. t. 11 = **Elatine**, Linn.
palludosa, Bell. l. c. = E. triandra.

BIROLIA, Rafin. Sylva Tellur. 163 (1838) = **Clusia**, Linn. (Guttif.).
alba, Rafin. l. c. = C. alba.

BISCHOFIA, Blume, Bijdr. 1168 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 281.
Microëlus, Wight & Arn. in Edinb. N. Phil. Journ. xiv. (1833) 298.
Stylodiscus, Benn. Pl. Jav. Rar. 133. t. 29 (1838).
Cummingiana, Decne. in Jacquem. Voy. Bot. 153 = javanica.
javanica, *Blume, Bijdr.* 1168.—Ind. or.; Malaya; Ins. Pacif.
leptopoda, *Muell. Arg. in DC. Prod.* xv. II. 479.—Ins. Pacif.
oblongifolia, Decne. in Jacquem. Voy. Bot. 152. t. 154 = javanica.
Roperiana, Decne. l. c. 153 = javanica.
Toui, Decne. l. c. = javanica.
trifoliata, Hook. Ic. Pl. t. 844 = javanica.

BISCUTELLA, Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 91.
Dithyrea, Harv. in Hook. Lond. Journ. Bot. iv. (1845) 77. t. 5.
Jondraba, Medic. Pflanzeng. 27 (1792).
Thlaspidium, Tourn. ex Adans. Fam. ii. 422 (1763).
algeriensis, Jord. Diagn. i. 318 = didyma.
alpestris, Waldst. & Kit. Pl. Rar. Hung. iii. 253 = laevigata.
alpicola, Jord. Diagn. i. 292 = laevigata.
alsatica, Jord. l. c. 300 = laevigata.
ambigua, DC. Mon. Bisc. 9. t. 11. f. 1; Syst. ii. 415 = laevigata.
anchusaefolia, Marz. ex Reichb. Fl. Germ. Excurs. 660 = laevigata.
anchusaefolia, Tenore, ex Steud. Nom. ed. II. i. 206 = sempervirens.
angulata, Steud. l. c. = Megacarpaea angulata.
angustifolia, *Schkuhr, Handb.* ii. 238.—Europ.
apetala, Walt. Fl. Carol. 174 = Senebiera didyma.
apricorum, Jord. Diagn. i. 307 = laevigata.
apula, Gaertn. Fruct. ii. 279 = leiocarpa.
apula, Lam. Encyc. iii. 617 = ciliata.
apula, Linn. Mant. 254 = didyma.
apula, Webb, ex Willk. & Lange, Prod. Fl. Hisp. iii. 761 = scutulata.

BISCUTELLA :—
arvernensis, Jord. Diagn. i. 298 = laevigata.
auriculata, Herb. Un. ex Steud. Nom. ed. II. i. 207 = cichoriifolia.
auriculata, *Linn. Sp. Pl.* 652.—Reg. Mediterr.
austriaca, *Bonnet, in Bull. Soc. Dauph.* fasc. vi. 222.—Europ.
balearia, L'Hérit. ex DC. Syst. ii. 643 = Succowia balearia.
balearica, Jord. Diagn. i. 325, 328 = auriculata?
boetica, Boiss. & Reut. Diagn. Ser. II. i. 42 = laevigata.
Bourgaei, Jord. Diagn. i. 320 = lyrata.
brevicaulis, Jord. l. c. 303 = laevigata.
Burseri, Jord. l. c. 326 = auriculata.
californica, *Benth. & Hook. f. ex S. Wats. Bot. Calif.* i. 48.—Calif.
Candollii, Jord. Diagn. i. 326 = auriculata.
chilensis, Lag. ex DC. Mon. Bisc. 6 = Cremolobus chilensis.
Chouletti, Jord. Diagn. i. 319 = didyma.
cichoriifolia, *Loisel. Add. Fl. Gall.* 167.—Mont. Pyren.
ciliata, *DC. Mon. Bisc.* 6; *Syst.* ii. 410.—Ital.; Hispan.
collina, Jord. Diagn. i. 299 = laevigata.
Columnae, Tenore, Prod. Fl. Nap. 38 = didyma.
confusa, *Pomel, Nouv. Mat. Fl. Atl.* 231.—Afr. bor.
controversa, Bor. Fl. Centr. Fr. ed. III. 56 = laevigata.
coronaria, *Ehrenb. ex Willk. in Ind. Sem. Dorpat. Adn.* (1868).—Hab.?
coronopifolia, Willd. Sp. Pl. iii. 474 = ciliata.
coronopifolia, Linn. Mant. ii. 255; All. Fl. Pedem. i. 247 = laevigata.
cuneata, Lag. Cat. Hort. Madr. (1815) 20 = Cremolobus chilensis.
depressa, Thom. ex DC. Syst. ii. 415 = laevigata.
depressa, Willd. Enum. Hort. Berol. 673 = didyma.
didyma, *Linn. Sp. Pl.* 653.—Reg. Mediterr.; Persia.
didyma, Scop. Fl. Carn. ed. II. ii. 14 = laevigata.
didyma, Willd. Enum. Hort. Berol. 673 = ciliata.
dilatata, Vis. Stirp. Dalm. 14 = cichoriifolia.
divionensis, Jord. Diagn. i. 305 = laevigata.
dubia, Steud. Nom. ed. II. i. 207 = Cremolobus linearifolius.
erigerifolia, DC. Mon. Bisc. 4; Syst. ii. 408 = auriculata.
eriocarpa, DC. Mon. Bisc. 7. t. 9. f. 2; Syst. ii. 411 = didyma.
erucifolia, *Reichb. Ic. Fl. Germ.* vii. 5.—Europ.
flexuosa, Jord. Diagn. i. 300 = laevigata.
frutescens, *Coss. Notes Crit.* ii. 27.—Hispan.
glabra, [Clairv.] Man. Herb. 216 = laevigata.
glacialis, Jord. Diagn. i. 310 = laevigata.
glareosa, Jord. l. c. 304 = laevigata.
granitica, Bor. ex Perar. in Bull. Soc. Bot. Fr. xvi. (1869) 352 = laevigata.
Guilloni, Jord. Diagn. i. 302 = laevigata.
hispida, DC. Mon. Bisc. 4. t. i. f. 1; Syst. ii. 408 = cichoriifolia.
incana, *Tenore, Fl. Neap. Prod. App.* 5ta. (1826).—Italia.
intermedia, Gouan, Illustr. 42 = laevigata.
intricata, Jord. Diagn. i. 308 = laevigata.
laciniata, Willd. ex DC. Syst. ii. 417 = Megacarpaea laciniata.
laevigata, *Linn. Mant.* 225.—Reg. Medit.
laevigata, Sibth. & Sm. Fl. Graec. Prod. ii. 16 = didyma.
Lamarckii, Jord. Diagn. i. 323 = auriculata.
Lamottii, Jord. l. c. 302 = laevigata.
laxa, *Boiss. & Reut. Diagn.* Ser. II. i. 43.—Hispan.
laxiflora, Presl, ex Spreng. Syst. iv. Cur. Post. 239 = didyma.
leiocarpa, *DC. Mon. Bisc.* 8; *Syst.* ii. 413.—Oriens.
Lima, Reichb. Fl. Germ. Excurs. 661, sub n. 4206 = laevigata.
longa, Dulac, Fl. Hautes-Pyr. 190 = cichoriifolia.
longifolia, Vill. Prosp. 38 = laevigata.
lucida, Balb. ex DC. Mon. Bisc. 9. t. 7; Syst. ii. 414 = laevigata.
lusitanica, Jord. Diagn. i. 315 = laevigata.
lyrata, Linn. Mant. 354 = didyma.
macrocarpa, Hort. ex DC. Syst. ii. 409 = cichoriifolia.
macrocarpa, *Bieb. ex Tauscher, in Mém. Soc. Nat. Mosc.* iv. (1812) 232.—Transural.
major, Schkuhr, Handb. ii. 238. n. 1822 = auriculata.
marginata, Tenore, Prod. Fl. Nap. 38 = didyma.
maritima, Tenore, l. c. = didyma.
mauritanica, Jord. Diagn. i. 325 = auriculata.

BISCUTELLA :—
maxima, Pall. ex DC. Syst. ii. 417 = Megacarpaea laciniata.
mediterrannea, Jord. Diagn. i. 313 = laevigata?
Megacarpaea, Boiss. & Reut. Diagn. Ser. II. i. 44 = laevigata.
megalocarpa, Fisch. ex DC. Mon. Bisc. 5. t. 3 = Megacarpaea laciniata.
microcarpa, DC. l. c. 7; Syst. ii. 411 = didyma.
minor, Jord. Diagn. i. 297 = laevigata.
mollis, Loisel. Not. 168 = laevigata.
montana, Cav. Ic. ii. 59. t. 177 = laevigata.
neustriaca, Ed. Bontrel, in Act. Soc. Dauph. (1879) 198 = laevigata.
nicaeensis, Jord. Diagn. i. 314 = laevigata?
obcordata, Reichb. Ic. Fl. Germ. vii. 8. f. 615 = laevigata.
obovata, *Hort. Par. ex DC. Mon. Bisc.* 8; *Syst.* ii. 413.—Hab.?
orcelitana, Lag. ex DC. Syst. ii. 408 = auriculata.
oreites, Jord. Diagn. i. 294 = laevigata.
orientalis, [Desf.] ex DC. Mon. Bisc. 8 = leiocarpa.
patulipes, Jord. Diagn. i. 320 = didyma.
perennis, Spach, Hist. Vég. Phan. vi. 571 = ciliata, laevigata, sempervirens.
peruviana, Lam. Encyc. iii. 620 = Cremolobus peruvianus.
petraea, Jord. Diagn. i. 306 = laevigata.
picridifolia, Lapeyr. Hist. Abr. Pl. Pyr. 375 = cichoriifolia.
pinnatifida, Hort. ex Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 302 = laevigata?
pinnatifida, Jord. Diagn. i. 311 = praec.?
polyclada, Jord. l. c. 312 = laevigata.
pratensis, *Timb. in Bull. Soc. Hist. Nat. Toul.* iii. (1869) 114.—Gallia.
pumila, Balb. ex DC. Mon. Bisc. 6; et Syst. ii. 411 = didyma.
pyrenaica, *Huet, in Ann. Sc. Nat.* Sér. III. xix. (1853) 252.—Hispan.
radicata, *Coss. & Dur. in Bull. Soc. Bot. Fr.* xix. (1872) 224.—Afr. bor.; Hispan.
radicosa, Willk. & Lange, Prod. Fl. Hisp. iii. 778 = radicata.
raphanifolia, Poir. Voy. Barb. ii. 198 = didyma.
rosularis, *Boiss. & Reut. Diagn.* Ser. II. i. 45.—Hispan.
rupestris, *Ehrh. Beitr.* v. 177, nomen.—German.
saxatilis, Schleich. ex DC. Prod. i. 183 = laevigata.
saxatilis, Endr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 765 = pyrenaica.
scabra, Salisb. Prod. 268 = didyma.
sclerocarpa, *Revel, in Bull. Soc. Bot. Fr.* xxv. (1878) *Bibl.* 78.—Gallia.
scutulata, *Boiss. & Reut. Diagn.* Ser. II. i. 41.—Hispan.
secunda, Jord. Diagn. i. 310 = laevigata.
sempervirens, DC. Syst. ii. 416 = laxa.
sempervirens, *Linn. Mant.* 255.—Hispan.; Lusitan.
sinuata, Jord. Diagn. i. 301 = laevigata.
spathulata, DC. Prod. i. 183, in syn. = laevigata.
stenophylla, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 299 = laevigata.
stricta, Jord. Diagn. i. 312 = laevigata.
subspathulata, Lam. Encyc. iii. 620 = laevigata.
suffrutescens, *Coss. Notes Crit.* 27.—Hispan.
suffruticosa, DC. Mon. Bisc. 5. t. 5 = Cremolobns suffruticosus.
tergestina, Jord. Diagn. i. 295 = laevigata.
tenuicaulis, Jord. l. c. 310 = laevigata.
tomentosa, Lag. ex DC. Syst. ii. 416 = laevigata.
tumidula, Lag. ex DC. l. c. 411 = didyma.
varia, Dum. Fl. Belg. Prod. 118 = laevigata.
variabilis, Loisel. Fl. Gall. ii. 56 = laevigata.
variegata, Boiss. & Reut. Diagn. Ser. II. i. 44 = laevigata.
verna, Matth. ex Nym. Consp. 59 = laevigata.
virgata, Jord. Diagn. i. 309 = laevigata.
Wislizeni, *Benth. & Hook. f. ex S. Wats. Bot. Calif.* i. 432.—Am. bor. occ.

BISERRULA, Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 507.
Pelecinus, Tourn. ex Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 378.
leiocarpa. Hochst. ex A. Rich. Tent. Fl. Abyss. i. 194 = B. Pelecinus.
pelecina, Lam. Fl. Fr. ii. 634 = B. Pelecinus.
Pelecinus, *Linn. Sp. Pl.* 762.—Reg. Mediterr.

BISMALVA, Medic. Malv. 39 (1787) = **Malva**, Tourn.
Alcea, Medic. l. c. = M. Alcea.
fastigiata, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 350 = M. Alcea.
laciniata, Fourr. l. c. = M. moschata.
moschata, Medic. Malv. 39 = M. moschata.
Tournefortiana, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 195 = M. Tournefortiana.

BISMARCKIA, Hildebr. & H. Wendl. in Bot. Zeit. xxxix. (1881) 89, 90, 93. *PALMAE*, Benth. & Hook. f. Gen. iii. 882.
nobilis, *Hildebr. & H. Wendl. l. c.* 94.—Madag.

BISPHAERIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 1 (Gen. dub.).

BISTANIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 64 (1827) (Gen. dub.).

BISTELLA, Adans. Fam. ii. 226 (1763) (an Rubiacea?).

BISTELLA, Delile, in Caill. Voy. Méroé (Cent. Pl. Afr.) 97. t. 63 (1826) = **Vahlia**, Thunb. (Saxifr.).
geminiflora, Caill. & Delile, Voy. Méroé, 97. t. 2. f. 2 = V. viscosa.

BISTORTA, [Tourn.] Linn. Syst. ed. I. (1735); Gen. ed. I. 115 (1737) = **Polygonum**, Tourn.
americana, Rafin. Fl. Tellur. iii. 12 = P. viviparum?
major, S. F. Gray, Nat. Arr. Brit. Pl. ii. 267 = P. Bistorta.
officinalis, Rafin. Fl. Tellur. iii. 12 = B. Bistorta.
vivipara, S. F. Gray, Nat. Arr. Brit. Pl. ii. 268 = P. viviparum.

BISTROPOGON, Auct. ex Steud. Nom. ed. II. i. 207. (1840) = **Bysteropogon**, L'Hérit. (Labiat.).

BISWAREA, Cogn. in Comptes-rend. Soc. Bot. Belg. xxi. (1882) 16. *CUCURBITACEAE*.
WAREA, C. B. Clarke, in Journ. Linn. Soc. xv. (1876) 127.
tonglensis, *Cogn. l. c.*—Reg. Himal.

BIVINIA, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 78. *SAMYDACEAE*, Benth. & Hook. f. i. 800.
Jaubertia, *Tul. l. c.*—Madag.

BIVONAEA, DC. Syst. ii. 554 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 88.
lutea, *DC. Syst.* ii. 555.—Sicil.
praecox, *Bertol. Fl. Ital.* x. 520.—Sicil.
Saviana, *Caruel, Prod. Fl. Tosc.* 47.—Ital.

BIVONAEA, Moç. & Sesse, ex DC. Prod. iii. 372 (1828) = **Pentacaena**, Bartl. (Illecebr.).
multicaulis, Moç. & Sesse, ex DC. l. c. 373 = P. polycnemoides.

BIVONEA, Rafin. Fl. Ludov. 138 (1817) = **Jatropha**, Linn. (Euphorb.).
stimulosa, Rafin. l. c. = J. urens.

BIVONIA, Spreng. Neue Entdeck. ii. 116 (1821) = **Bernardia**, P. Br. (Euphorb.).
axillaris, Spreng. l. c. = B. axillaris.

BIWALDIA, Scop. Introd. 232 (1777) = **Garcinia**, Linn. (Guttif.).

BIXA, Linn. Coroll. Gen. 8 (1737). *BIXINEAE*, Benth. & Hook. f. i. 125.
URUCU, Adans. Fam. ii. 381 (1763).
acuminata, Boj. Hort. Maurit. 20 = Orellana.
americana, Poir. Encyc. vi. 229 = Orellana.
odorata, Ruiz & Pav. Fl. Per. v. t. 460, ex G. Don, Gen. Syst. i. 295 = Orellana?
Orellana, *Linn. Sp. Pl.* 512.—Am. austr.
orleana, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 8 = Orellana.
platycarpa, Ruiz & Pav. Fl. Per. v. t. 459, ex G. Don, Gen. Syst. i. 294 = Orellana.
purpurea, Sweet, Hort. Brit. ed. I. 33; G. Don, Gen. Syst. i. 295 = Orellana?
sphaerocarpa, *Triana, in Bull. Soc. Bot. Fr.* v. (1858) 369.—N. Granat.
tinctoria, Salisb. Prod. 369 = Orellana.
Urucurana, Willd. Enum. Hort. Berol. 565 = Orellana.

BIXAGREWIA, Kurz, in Journ. Bot. xiii. (1875) 325 = **Trichospermum**, Blume (Tiliac.).
nicobarica, Kurz, l. c.—Ins. Nicobar.

BLABEROPUS, A. DC. Prod. viii. 410 (1844) = **Alstonia**, R. Br. (Apocyn.).
lucidus, A. DC. l. c. 411 = Trachelospermum fragrans.
neriifolius, A. DC. l. c. = A. neriifolia.
Sebusi, Heurck & Muell. Arg. in Heurck, Obs. Bot. ii. 188 = A. neriifolia.
sericeus, A. DC. Prod. viii. 411 = A. sericea.
venenatus, A. DC. l. c. = A. venenata.
villosus, Miq. Fl. Ind. Bat. ii. 440 = A. villosa.

BLACHIA, Baill. Etud. Gén. Euph. 388. t. 19 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 301.
calycina, *Benth. in Journ. Linn. Soc.* xvii. (1880) 226.—Ind. or.
denudata, *Benth. l. c.*—Ind. or.
Pentzii, *Benth. l. c.*—China.
reflexa, *Benth. l. c.*—Ind. or.
umbellata, *Baill. Etud. Gén. Euph.* 387. t. 19. f. 18–20. —Ind. or.

BLACKBOURNEA, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 356 = seq.

BLACKBURNIA, Forst. Char. Gen. t. 6 (1776) = **Zanthoxylum**, Linn. (Rutac.).
monadelpha, Roxb. Hort. Beng. [83]; Fl. Ind. i. 415 = Dobera Roxburghii.
oppositifolia, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 191 = Dobera Roxburghii.
pinnata, Forst. Char. Gen. t. 6 = Z. Blackburnia.

BLACKIA, Schrank, ex DC. Prod. iii. 165 (1828) = **Myriaspora**, DC. (Melastom.).
egensis, Schrank, ex DC. l. c. = M. egensis.

BLACKSTONIA, Huds. Fl. Angl. ed. I. 146 (1762), nomen prius = **Chlora**, Ren. ex Adans. Fam. ii. 503 (1763).
perfoliata, Huds. l. c. = C. perfoliata.

BLACKSTONIA, A. Juss. in Orbign. Dict. vi. 432 (1849) = BLAKSTONIA, Scop. = **Moronobea**, Aubl. (Guttif.).

BLACKWELLIA, Gaertn. Fruct. ii. t. 117 (1791) (Blakwellia, p. 169) (Genus dubium).
antarctica, Gaertn. l. c.—Hab.?

BLACKWELLIA, Comm. ex Juss. Gen. 343 (1789) (Blakwellia) = **Homalium**, Jacq. (Samyd.).
acuminata, Wall. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 596 = H. nepalense.
africana, Hook. f. in Hook. Niger Fl. 361 = H. africanum.
austro-caledonica, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 107 = H. austro-caledonicum?
axillaris, Lam. Encyc. i. 428 = H. axillare.
brachybotrya, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 48 = H. brachybotrys.
brachystylis, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 59.—Madag.
caryophyllacea, Zoll. & Mor. Syst. Verz. Zoll. 33 = H. caryophyllaceum.
cerosifolia, Vent. Choix, 56. t. 56.—Madag.
ceylanica, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 452 = H. zeylanicum.
chinensis, Lindl. ex Steud. Nom. ed. II. i. 208 = H. fagifolium.
cochinchinensis, Blume, Mus. Bot. Lugd. Bat. ii. 27.—Cochinchina.
dasyantha, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 610 = H. Griffithianum.
decurrens, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 105.—N. Caled.
dentata, Harv. in Harv. & Sond. Fl. Cap. ii. 585.—Afr. austr.
Deplanchei, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 110.—N. Caled.
eriantha, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 62.—Madag.
fagifolia, Lindl. in Trans. Hort. Soc. Lond. vi. (1826) 269 = H. fagifolium.
foetida, Wall. Cat. n. 4899 = H. foetidum.

BLACKWELLIA:—
Jussieua, J. F. Gmel. Syst. 768 = Homalium paniculatum?
glauca, Vent. Choix, 55. t. 55 = H. paniculatum.
gracilis, Blume, Mus. Bot. Lugd. Bat. ii. 26 = H. axillare.
grandiflora, Lindl. ex Loud. Hort. Brit. 193 = H. fagifolium.
Guillaini, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 106.—N. Caled.
integrifolia, Lam. Encyc. i. 428 = H. paniculatum.
intermedia, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 109.—N. Caled.
kanaliensis, Vieill. l. c. 112.—N. Caled.
longiflora, Miq. Fl. Ind. Bat. i. 1. 715 = H. cordylanthus.
Loureiri, Benth. in Hook. Lond. Journ. Bot. i. (1842) 482 = H. fagifolium.
macrostachya, Turcz. in Bull. Soc. Nat. Mosc. (1863) 610 = H. longifolium.
micrantha, Boiv. ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 63 = H. micranthum.
Mathieuana, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 108.—N. Caled.
moluccana, Blume, Mus. Bot. Lugd. Bat. ii. 27.—Ins. Molucc.
monostachya, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 610.—Penang.
montana, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 110.—N. Caled.
napaulensis, DC. Prod. ii. 54 = seq.
nepalensis, Wall. Pl. As. Rar. t. 179 = H. nepalense.
padiflora, Lindl. Bot. Reg. t. 1308 = H. fagifolium.
padifolia, Steud. Nom. ed. II. i. 208 = H. fagifolium.
paniculata, Lam. Encyc. i. 428 = H. paniculatum.
planiflora, Boiv. ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 64.—Madag.
polystachya, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 107.—N. Caled.
propinqua, Wall. Cat. n. 4898 = H. propinquum.
pubiflora, Walp. Rep. i. 548 = H. fagifolium.
purpurea, Steud. Nom. ed. II. i. 208, sphalm. = H. propinquum.
rivularis, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 111.—N. Caled.
rubiginosa, Vieill. l. c. 108.—N. Caled.
rufescens, Presl, Bot. Bemerk. 42 = H. rufescens.
spiralis, Wall. in As. Res. xiii. (1820) 400 = H. tomentosum.
spiralis, Wall. Cat. n. 4897, partim = H. propinquum.
tetrandra, Wight, Ic. v. t. 1851 = H. zeylanicum.
Thouarsiana, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 60.—Madag.
tomentosa, Vent. Choix, 57. t. 57 = H. tomentosum.

BLADHIA, Thunb. Nov. Gen. 6 (1781), nomen prius = **Ardisia**, Sw. (1788) (Myrsin.).
crispa, Thunb. Fl. Jap. 97 = A. crenata, *Roxb.*
glabra, Thunb. in Trans. Linn. Soc. ii. (1794) 331 = A. japonica.
japonica, Blanco, Fl. Filip. ed. I. 126 = A. pyramidalis.
japonica, Thunb. Nov. Gen. 7 = A. japonica.
villosa, Thunb. Fl. Jap. 96 = A. pusilla.

BLAERIA, Linn. Coroll. Gen. 3 (1737). *ERICACEAE*, Benth. & Hook. f. ii. 592.
KOLBIA, Adans. Fam. ii. 164 (1763).
aggregata, Wendl. ex Steud. Nom. ed. II. i. 208; ii. 589 = Simocheilus multiflorus.
articulata, Linn. Mant. ii. 198 (= *Syndesmanthus articulatus*).—Afr. austr.
articulata, Willd. ex Steud. Nom. ed. II. i. 208 = Sympieza capitellata.
barbigera, *G. Don, Gen. Syst.* iii. 805.—Afr. austr.
bicolor, Klotzsch, in Linnaea, viii. (1833) 660 (err. typ. 606) = Simocheilus bicolor.
bracteata, Wendl. Collect. ii. 1. t. 37 = Sympieza capitellata.
bruniaefolia, G. Don, Gen. Syst. iii. 805 = Erica bruniaefolia.
caduca, *Thunb. Diss. Blaeria*, 10.—Afr. austr.
campanulata, *Benth. in DC. Prod.* vii. 698.—Afr. austr.
carnea, Klotzsch, in Linnaea, viii. (1833) 661 = Simocheilus carneus.
ciliaris, Linn. f. Suppl. 122 = Grisebachia ciliaris.

BLAERIA:—
ciliciiflora, G. Don, Gen. Syst. iii. 805 = Grisebachia velleriflora.
coccinea, *Klotzsch, in Linnaea*, viii. (1833) 657.—Afr. austr.
condensata, Hochst. in A. Rich. Tent. Fl. Abyss. ii. 13 = spicata.
depressa, Drège, ex Benth. in DC. Prod. vii. 705 = Sympieza Kunthii.
depressa, Lichst. ex Roem. & Schult. Syst. iii. 168 = Simocheilus depressus.
dumosa, Wendl. Collect. ii. 3. t. 38 = purpurea.
equisetifolia, G. Don, Gen. Syst. iii. 805 = purpurea.
eriantha, Willd. ex Steud. Nom. ed. II. i. 208 (= *Syndesmanthus articulatus*).—Afr. austr.
ericoides, *Linn. Sp. Pl.* 162.—Afr. austr.
fasciculata, Willd. Sp. Pl. i. 629 = Simocheilus glabellus.
fastigiata, *Benth. in DC. Prod.* vii. 697.—Afr. austr.
flexuosa, *Benth. l. c.* 698.—Afr. austr.
flosculosa, G. Don, Gen. Syst. iii. 805 (= *Syndesmanthus paucifolius*).—Afr. austr.
fuscescens, *Klotzsch, in Linnaea*, viii. (1833) 657.—Afr. austr.
glabella, Drège, ex Benth. in DC. Prod. vii. 698 = campanulata.
glabella, Willd. Sp. Pl. i. 631 = Simocheilus glabellus.
glabra, *Thunb. Diss. Blaeria*, 10.—Afr. austr.
gracilis, Bartl. in Linnaea, vii. (1832) 650 = Sympieza Kunthii.
hirsuta, Lichtst. in Roem. & Schult. Syst. iii. 170 (= *Syndesmanthus paucifolius*).—Afr. austr.
hirsuta, *Thunb. Diss. Blaeria*, 8.—Afr. austr.
incana, Bartl. in Linnaea, vii. (1832) 650 = Grisebachia incana.
jucunda, Reichb. ex Steud. Nom. ed. II. i. 208 = purpurea.
Kraussiana, *Klotzsch, in Walp. Rep.* ii. 728.—Afr. austr.
leucocephala, *Bory, Voy.* iii. 172.—Ins. Borbon.
multiflora, Klotzsch, in Linnaea, viii. (1833) 661 = Simocheilus multiflorus.
muscosa, Ait. Hort. Kew. i. 150 = Scyphogyne inconspicua.
nodiflora, G. Don, Gen. Syst. iii. 805 = Grisebachia plumosa.
nudiflora, *Thunb. Diss. Blaeria*, 6.—Afr. austr.
paniculata, *Thunb. l. c.* 10.—Afr. austr.
parviflora, Klotzsch, in Linnaea, viii. (1833) 665 (= *Codonanthemum parviflorum*).—Afr. austr.
pauciflora, G. Don, Gen. Syst. iii. 804, sphalm. = *paucifolia*.
paucifolia, Wendl. Collect. ii. 17. t. 43 (= *Syndesmanthus paucifolius*).—Afr. austr.
ptilota, E. Mey. in Drège, ex Benth. in DC. Prod. vii. 701 = Grisebachia hispida.
puberula, Klotzsch, in Linnaea, viii. (1833) 661 (= *Codonanthemum puberulum*).—Afr. austr.
purpurea, Berg. Desc. Pl. Cap. 34 = Simocheilus glabellus.
purpurea, *Linn. f. Suppl.* 122.—Afr. austr.
pusilla, *Klotzsch, in Linnaea*, viii. (1833) 659.—Afr. austr.
pusilla, *Linn. Mant.* 139 ('sp. dubia').—Afr. austr.
revoluta, Bartl. in Linnaea, vii. (1832) 650 = barbigera.
rubra, Hort. ex Stend. Nom. ed. II. i. 208 = ericoides.
scabra, Drège, ex Benth. in DC. Prod. vii. 698 = ericoides.
scabra, Willd. Sp. Pl. i. 629 (= *Syndesmanthus scaber*).—Afr. austr.
serrata, *Thunb. Diss. Blaeria*, 6.—Afr. austr.
sessiliflora, Wendl. ex Steud. Nom. ed. II. i. 208 = Simocheilus multiflorus.
spicata, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 13.—Abyss.
Thunbergii, G. Don, Gen. Syst. iii. 805 = Grisebachia ciliaris.
trigona, Wendl. ex Steud. Nom. ed. II. i. 208 = fuscescens.
turmalis, G. Don, Gen. Syst. iii. 805 = Erica turmalis.
xeranthemifolia, G. Don, l. c. = Eremia lanata.

BLAINVILLEA, Cass. in Dict. Sc. Nat. xxix. 493 (1823). *COMPOSITAE*, Benth. & Hook. f. ii. 369.
CALYPTOCARPUS, Less. Syn. Comp. 221 (1833).
EISENMANNIA, Sch. Bip. in Flora, xxv. (1842) II. 418.

BLAINVILLEA:—
GALOPHTHALMUM, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 7. t. 2.
OLIGOGYNE, DC. Prod. v. 629 (1837).
UCACEA, Cass. in Dict. Sc. Nat. xxvii. 9 (1823).
alba, Edgew. in Trans. Linn. Soc. xx. (1851) 70 = rhomboidea.
amazonica, *Benth. & Hook. f. Gen.* ii. 370.—Reg. Amazon.
bahiensis, *Baker, in Mart. Fl. Bras.* vi. III. 177.—Bras.
biaristata, *DC. Prod.* v. 492.—Bras.
Gayana, Benth. in Hook. Niger Fl. 434, in syn. = Aspilia Gayana.
Gayana, *Cass. in Dict. Sc. Nat.* xlvii. 90.—Afr. trop.
hispida, Edgew. in Trans. Linn. Soc. xx. (1851) 70 = rhomboidea.
lanceolata, *Baker, in Mart. Fl. Bras.* vi. III. 176.—Bras.
latifolia, DC. in Wight, Contrib. Bot. Ind. Or. 71 = rhomboidea.
polycephala, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 89 = rhomboidea.
Prieureana, Benth. in Hook. Niger Fl. 433, in syn. = Aspilia Prieureana.
Prieureana, *DC. Prod.* v. 492.—Afr. trop.
racemosa, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 89 = rhomboidea.
rhomboidea, *Cass. in Dict. Sc. Nat.* xxix. 493.—Cosmop. trop.; Ind. or.
synedrelloides, *Benth. & Hook. f. Gen.* ii. 370.—Hab. ?
tampicana, *Benth. & Hook. l. c.—Cf. Hemsl. Biol. Centr. Am. Bot.* ii. 169.—Mexic.
tenuicaulis, *Benth. & Hook. f. Gen.* ii. 370.—Ins. Galap.

BLAIRIA, Houst. ex Linn. Gen. ed. I. 334, in syn. (1737) = **Priva**, Adans. (Verben.).
cordifolia, Moench, Meth. 423 = P. hispida.
javanica, Gaertn. Fruct. i. t. 56 = Lippia javanica.
lappulacea, Steud. Nom. ed. II. i. 208 = P. echinata.
mexicana, Gaertn. Fruct. i. 265. t. 56. f. 1 = P. hispida.
nodiflora, Gaertn. l. c. 266. t. 56 = Lippia nodiflora.

BLAIRIA, Gled. in Mém. Acad. Berl. (1751) 125 = **Blaeria**, Linn. (Ericac.).

BLAKBURNIA, J. F. Gmel. Syst. 258 (1791) = BLACKBURNIA, Forst. = **Zanthoxylum**, Linn. (Rutac.).

BLAKEA, P. Br. Hist. Jamaic. 323 (1756). *MELASTOMACEAE*, Benth. & Hook. f. i. 770.
PYXIDANTHUS, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 150. t. 6.
VALDESIA, Ruiz & Pav. Prod. 67 (1794).
brevipes, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 144 = repens.
Cacatin, *D. Don, in Mem. Wern. Soc.* iv. (1823) 324.—Guiana.
caudata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 148.—N. Granat.
glabrescens, *Benth. Bot. Voy. Sulph.* 94.—Peruv.
gracilis, *Hemsl. Diag. Pl. Nov.* 13.—Costa Rica.
granatensis, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 143.—N. Granat.
grandiflora, *Hemsl. Diag. Pl. Nov.* 13.—Costa Rica.
hirsuta, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 148.—Bras.
laevigata, D. Don, in Mem. Wern. Soc. iv. (1823) 327 = Topobea laevigata.
latifolia, *D. Don, l. c.* 325.—Peruv.
laurifolia, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 143 = pulverulenta.
Lindeniana, *Triana in Trans. Linn. Soc.* xxviii. (1871) 149.—Venezuela.
macrophylla, D. Don, in Mem. Wern. Soc. iv. (1823) 326 = Bellucia superba.
mexicana, D. Don, l. c. 325 = repens.
multiflora, D. Don, l. c. 326 = Topobea multiflora.
ovalis, *D. Don, l. c.* 325.—Peruv.
ovalis, Griseb. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 148 = repens.
parasitica, D. Don, in Mem. Wern. Soc. iv. (1823) 327 = Topobea parasitica.
podagrica, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 149.—N. Granat.
pulverulenta, DC. Prod. iii. 196 = Conostegia subhirsuta.

BLAKEA:—
pulverulenta, *Vahl, Symb. Bot.* iii. 61.—Am. austr.
pyxidanthus, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 149.—N. Granat.
quadrangularis, *Triana, l. c.*—N. Granat.
quinquenervia, Aubl. Pl. Gui. i. 525. t. 210 = Bellucia Aubletii.
repens, *D. Don, in Mem. Wern. Soc.* iv. (1823) 325.—Peruv.
repens, Griseb. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 149 = ovalis.
rosea, *D. Don, in Mem. Wern. Soc.* iv. (1823) 325.—Peruv.
rostrata, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 148.—Peruv.
rotundifolia, *D. Don, in Mem. Wern. Soc.* iv. (1823) 326.—Peruv.
Schlimii, *Triana, in Trans. Linn. Soc.* xxvii. (1871) 149.—Venezuela.
sessiliflora, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 144.—Peruv.
sessiliflora, Poepp. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 148 = hirsuta.
subconnata, *O. Berg, ex Triana, l. c.*—Ecuador.
trinervia, *Linn. Sp. Pl.* ed. II. 635.—Jamaica.
trinervia, Pav. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 149 = Topobea laevigata.
trinervis, Pav. ex D. Don, in Mem. Wern. Soc. iv. (1823) 327 = Topobea laevigata.
triplinervia, Linn. Amoen. Acad. viii. 257; Pl. Surinam. n. 59 = Bellucia Aubletii.

BLAKSTONIA, Scop. Introd. 276 (1777) = **Moronobea**, Aubl. (Guttif.).

BLAKWELLIA, Comm. ex Juss. Gen. 343 (1789) (Blackwellia, ex J. F. Gmel. Syst. 768 (1791)) = **Homalium**, Jacq. (Samyd.).

BLAKWELLIA, Gaertn. Fruct. ii. 170 (1791) (Blackwellia, t. 117) (Genus dubium).

BLAKWELLIA, Scop. Introd. 326 (1777) = **Aralia**, Tourn.

BLANCHEA, Boiss. Fl. Orient iii. 212, in syn. (1875) = **Iphiona**, Cass. (Compos.).
iphionioides, Boiss. l. c.—Syria.

BLANCHETIA, DC. Prod. v. 75 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 226.
heterosticha, *DC. l. c.*—Bras.

BLANCKIA, Neck. Elem. i. 347 (1790) = **Conobea**, Aubl. (Scrophul.).

BLANCOA, Blume, Rumphia, ii. 128 (1836) = **Didymosperma**, H. Wendl. (Palm.).

BLANCOA, Blume, Rumphia, iii. 181 (1847) = **Harpullia**, Roxb. (Sapind.).
arborea, Blume, l. c. = H. cupanioides.

BLANCOA, Lindl. Swan Riv. App. 45 (1839). *HAEMODORACEAE*, Benth. & Hook. f. iii. 676.
canescens, *Lindl. l. c.*—Austral.

BLANDFORDIA, Andr. Bot. Rep. t. 343 (1804) = **Galax**, Linn. (Diapens.).
caudata, Steud. Nom. ed. II. i. 209, sphalm. = seq.
cordata, Andr. Bot. Rep. t. 343 = G. aphylla.

BLANDFORDIA, Sm. Exot. Bot. i. 5. t. 4 (1804). *LILIACEAE*, Benth. & Hook. f. iii. 774.
aurea, *Hook. f. Bot. Mag.* t. 5809.—Austral.
Backhousii, Lindl. Bot. Reg. (1845) sub t. 18 = grandiflora.
Cunninghamii, Lindl. l. c. = grandiflora.
Cunninghamii, Van Houtte, Fl. des Serres, t. 1829 = flammea.
flammea, *Lindl. in Journ. Hort. Soc.* v. (1850) 32.—Austral.
grandiflora, *R. Br. Prod.* 296.—Tasmania.
grandiflora, Hook. f. Fl. Tasm. ii. 49 = marginata.
grandiflora, Lindl. Bot. Reg. t. 924 = nobilis ?
intermedia, *Herb. in Lindl. Bot. Reg.* (1845) *Misc.* 64.—Tasmania.
marginata, *Herb. l. c.* (1842) *Misc.* 84.—Austral.

BLANDFORDIA :—
nobilis, *Sm. Exot. Bot.* i. 5. t. 4.—Austral.
Princeps, W. G. Smith, in Flor. Mag. N. S. (1875) t. 170 = flammea.
punicea, Sweet, Hort. Brit. ed. II. 517 = grandiflora.
speciosa, Hort. Angl. ex Link, Enum. Hort. Berol. i. 330 = grandiflora ?
uniflora, Willd. ex Kunth, Enum. Pl. iv. 478 = Milla biflora.

BLANDFORTIA, Poir. Encyc. Suppl. iv. 116 (1814) = **Blandfordia**, Sm. (Liliac.).

BLANDINA, Rafin. Fl. Tellur. iii. 88 (1836) = **Leucas**, Burm. (Labiat.).
biflora, Rafin. l. c. = L. biflora.
chinensis, Rafin. l. c. = L. chinensis.

BLANDOWIA, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 10. t. 4. f. 2 = **Apinagia**, Tul. (Podost.).
Preissii, Wedd. in DC. Prod. xvii. 85 = A. Preissii.
striata, Corda, in Sturm, Deutschl. Fl. Abth. ii. Heft v–vi. p. 107. t. 33 = A. Preissii.
striata, Lehm. ex Wedd. in DC. Prod. xvii. 71 = Dicraea Wallichii.
striata, Willd. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 100. t. 4 = A. Willdenowii.

BLANISIA, Pritz. Ind. Ic. 154 (1855), sphalm. = POLANISIA, Rafin. = **Cleome**, Linn. (Cappar.).

BLASTANIA, Kotschy & Peyr. Pl. Tinn. 15. t. 7 (1865–66). *CUCURBITACEAE.*
CTENOLEPIS, Hook. f. in Benth. & Hook. f. Gen. i. 832 (1867).
CTENOPSIS, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 12.
fimbristipula, *Kotschy & Peyr. l. c.*—Afr. trop.
Garcini, *Cogn. in DC. Monog. Phan.* iii. 629.—Ind. or.

BLASTEMANTHUS, Planch. in Hook. Lond. Journ. Bot. v. (1846) 644. *OCHNACEAE*, Benth. & Hook. f. i. 320.
gemmiflorus, *Planch. l. c.* 645.—Bras.
grandiflorus, *Spruce, ex Engl. in Mart. Fl. Bras.* xii. II. 355.—Bras.

BLASTOTROPHE, F. Didrichs. in Kjoeb. Vidensk. Meddel. (1854) 192 = **Alafia**, Thou. (Apocyn.).
scandens, F. Didrichs. l. c. 193.—Afr. trop.

BLASTUS, Lour. Fl. Cochinch. 526 (1790). *MELASTOMACEAE*, Benth. & Hook. f. i. 752.
cochinchinensis, *Lour. l. c.* 527.—Cochinch.
Hindsii, Hance, in Journ. Linn. Soc. xiii. (1873) 103 = Allomorphia pauciflora.
parviflorus, Triana, in Trans. Linn. Soc. xxviii. (1871) 74 = cochinchinensis.

BLATTARIA, [Tourn.] Rupp. Fl. Jen. ed. Hall. 247 (1745) = **Verbascum**, Linn. (Scrophul.).
alba, Medic. in Vorles. Churpf. Phys. Ges. iv. 1. (1789) 230 = V. Lychnitis.
Bastardi, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 125 = V. Bastardi.
virgata, Fourr. l. c. = V. virgatum.
vulgaris, Fourr. l. c. = V. Blattaria.

BLATTI, Adans. Fam. ii. 88 (1763) = **Sonneratia**, Linn. f. (Lythrar.).

BLAXIUM, Cass. in Dict. Sc. Nat. xxx. 328 (1824) = **Dimorphotheca**, Vaill. (Compos.).
decumbens, Cass. l. c. = D. fruticosa.

BLEASDALEA, F. Muell. Fragm. v. 90 (1865) = **Grevillea**, R. Br. (Proteac.).
cupanoides, F. Muell. l. c. = G. Bleasdalea.

BLECHUM, P. Br. Hist. Jamaic. 261 (1756). *ACANTHACEAE*, Benth. & Hook. f. ii. 1082.
ALVAREZIA, Pav. ex Nees, in DC. Prod. xi. 465 (1847).
angustifolium, *R. Br. Prod.* i. 478.—Ind. occ.
angustus, *Nees, in DC. Prod.* xi. 467.—N. Granat.
anisophyllum, Juss. in Ann. Mus. Par. ix. (1807) 270 = Phaylopsis longifolia.

BLECHUM :—
brasiliense, Mart. ex Nees, in DC. Prod. xi. 284 = Stenandrium mandioccanum.
brasiliense, *Lodd. Cat. ex G. Don, in Loud. Hort. Brit.* 245.—Bras.
Brownei, *Juss. in Ann. Mus. Par.* ix. (1807) 270.—Ind. occ.
Brownei, H. B. & K. Nov. Gen. et Sp. ii. 238 = Linnaei.
costaricense, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 168.—Costa Rica.
grandiflorum, *Oerst. l. c.*—Am. centr.
Haenkei, *Nees, in DC. Prod.* xi. 467.—Mexic.
hamatum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 220.—Afr. trop.
laxiflorum, *Juss. in Ann. Mus. Par.* ix. (1807) 270.—Jamaica.
Linnaei, *Nees, in DC. Prod.* xi. 465.—Mexic.
luzonium, Nees, l. c. 466 = Brownei.
mexicanum, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 169.—Mexic.
Sternbergii, Nees, in DC. Prod. xi. 468 = Tetramerium nervosum.
trinitense, *Nees, l. c.* 466.—Ins. Trinit.
Tweedii, *Nees, l. c.*—Mexic.

BLEEKERIA, Hassk. Retzia, i. 38 (1855) = **Ochrosia**, Juss. (Apocyn.).
calocarpa, C. Muell. in Walp. Ann. v. 492 = seq.
kalocarpa, Hassk. Retzia, i. 40 = O. elliptica.
salubris, Hassk. l. c. 41 = O. borbonica.

BLEEKERIA, Miq. Fl. Ind. Bat. i. II. 407 (1859) = **Alchornea**, Sw. (Euphorb.).
Zollingerii, Miq. l. c. = A. villosa.

BLEEKRODEA, Blume, Mus. Bot. Lugd. Bat. ii. 87. t. 28 (1852) [1856]. *URTICACEAE*, Benth. & Hook. f. iii. 358.
insignis, *Blume, l. c.* 88.—Borneo.
madagascariensis, *Blume, l. c.*—Madag.

BLENCOCOËS, Rafin. Fl. Tellur. iii. 75 (1836) = **Nierembergia**, Ruiz & Pav. (Solanac.).
filicaulis, Rafin. l. c. 76 = N. linearifolia.
longiflora, Rafin. l. c.—Hab. ?

BLENNODERMA, Spach, in Nouv. Ann. Mus. Par. iv. (1835) 406 = **Oenothera**, Linn. (Onagrar.).
Drummondii, Spach, l. c. 407 = Oe. Spachiana.

BLENNODIA, R. Br. in Sturt, Exped. ii. App. 67 (1849). *CRUCIFERAE*, Benth. & Hook. f. i. 82.
alpestris, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 100.—Austral.
blennodes, F. Muell. l. c. = lasiocarpa.
brevipes, *F. Muell. l. c.*—Austral.
canescens, *R. Br. in Sturt, Exped.* ii. *App.* 67.—Austral.
cardaminoides, *Benth. Fl. Austral.* i. 75.—Austral.
Cunninghamii, *Benth. l. c.* 76.—Austral.
curvipes, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 100.—Austral.
eremigera, *Benth. Fl. Austral.* i. 74.—Austral.
filifolia, *Benth. l. c.* 73.—Austral.
lasiocarpa, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 100.—Austral.
nasturtioides, *Benth. Fl. Austral.* i. 74.—Austral.
trisecta, *Benth. l. c.*—Austral.

BLENNOSPERMA, Less. Syn. Comp. 267 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 404.
APALUS, DC. Prod. v. 507 (1836).
CONIOTHELE, DC. Prod. v. 531 (1836).
UNXIA, Colla, in Mem. Acc. Torin. xxxviii. (1835) 37. t. 32.
californicum, *Torr. & Gray, Fl. N. Am.* ii. 272.—Calif.
chilense, *Less. Syn. Comp.* 267.—Chili.

BLENNOSPORA, A. Gray, in Hook. Kew Journ. iii. (1851) 172 = **Calocephalus**, R. Br. (Compos.).
Drummondii, A. Gray, l. c. 173 = C. Drummondii.

BLEPETALON, Rafin. Sylva Tellur. 30 (1838) = **Scutia**, Comm. (Rhamn.).
aculeatum, Rafin. l. c. = S. Commersoni.

BLEPHANTHERA, Rafin. Fl. Tellur. ii. 59 (1836) = **Bulbine**, Linn. (Liliac.).
depressa, Rafin. l. c. = Bulbine bulbosa.
Hookeri, Rafin. l. c. (= *Anthericum Hookeri*).—N. Zel.

BLEPHARACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836) = **Blepharis**, Juss. (Acanth.).
capensis, Nees, l. c. = Blepharis capensis.
carduifolius, Nees, ex Steud. Nom. ed. II. i. 209 = Blepharis carduifolia.
furcatus, Nees, ex Steud. l. c. = Blepharis furcata.
integrifolius, Nees, ex Steud. l. c. = Blepharis saturejaefolia.
procumbens, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 = Blepharis glomerata.
verticillata, Steud. Nom. ed. II. i. 209 = Cardanthera verticillata.

BLEPHARADEN, Dulac, Fl. Hautes-Pyr. 449 (1867) = **Swertia**, Linn. (Gentianac.).
perennis, Dulac, l. c. = S. perennis.

BLEPHARANDRA, Griseb. in Linnaea, xxii. (1849) 7. *MALPIGHIACEAE*, Benth. & Hook. f. i. 254.
hypoleuca, *Griseb. l. c.*—Guiana.

BLEPHARANTHEMUM, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. 20 (1841) = **Plagianthus**, Forst. (Malvac.).
sidoides, Klotzsch, l. c. = P. sidoides.

BLEPHARANTHES, Sm. Gramm. Bot. 188 (1821) = **Modecca**, Lam. (Passif.).

BLEPHARIGLOTIS, Rafin. Fl. Tellur. ii. 38 (1836) = **Habenaria**, Willd. (Orchid.).
albiflora, Rafin. l. c. = H. albida ?
bicolor, Rafin. l. c. 39.—Hab. ?
cristata, Rafin. l. c. = H. cristata.
flaviflora, Rafin. l. c.—Hab. ?
longicornis, Rafin. l. c. 38 = H. longicornis.

BLEPHARIPAPPUS, Hook. Fl. Bor. Am. i. 316 (1833). *COMPOSITAE*, Benth. & Hook. f. ii. 392.
PTILONELLA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 385.
glandulosus, Hook. l. c. = Layia glandulosa.
scaber, *Hook. l. c.*—Am. bor. occ.

BLEPHARIS, Juss. Gen. 103 (1789). *ACANTHACEAE*, Benth. & Hook. f. ii. 1089.
ACANTHODIUM, Delile, Fl. Egypte, 241. t. 33 (1813).
BLEPHARACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836).
ablephara, *Nees, in DC. Prod.* xi. 267.—Java.
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 150 = boerhaviaefolia.
acanthodioides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 212.—Afr. trop.
angusta, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 35.—Afr. austr.
arborea, *Spreng. Syst.* ii. 819.—Arabia.
asperrima, *Nees, in DC. Prod.* xi. 267.—Ind. or.
boerhaavifolia, Roth, Nov. Sp. 320 = seq.
boerhaviaefolia, *Pers. Syn.* ii. 180.—Ind. or.
calaminthaefolia, *Pers. l. c.*—Hab. ?
capensis, *Pers. l. c.*—Afr. austr.
carduifolia, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 35.—Afr. austr.
cuanzensis, *Welw. ex S. Moore, in Journ. Bot.* xviii. (1880) 230.—Afr. trop.
edulis, *Pers. Syn.* ii. 180.—Arab.; Persia; Ind. bor. occ.
furcata, *Pers. l. c.*; *T. Anders. in Journ. Linn. Soc.* vii. (1864) 35.—Afr. austr.
furcata, Eckl. ex Nees, in DC. Prod. xi. 276; et Zeyh. ex Nees, l. c. 277 = capensis.
glauca, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 36.—Afr. austr.
glomerata, *Poir. Encyc. Suppl.* i. 89.—Afr. austr.
glumacea, *S. Moore, in Journ. Bot.* xviii. (1880) 232.—Afr. trop.
grossa, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 36.—Afr. austr.
Guinzii, *T. Anders. l. c.* 34.—Afr. austr.
hirtinervia, *T. Anders. l. c.* 36.—Afr. austr.

BLEPHARIS:—
integrifolia, E. Mey. in Drège, Zwei Pfl. Docum. 168 = saturejaefolia.
involucrata, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 107.—Afr. trop.
linariaefolia, *Pers. Syn.* ii. 180.—Afr. trop.
maderaspatensis, Heyne, ex Roth, Nov. Sp. 320 = boerhaviaefolia.
molluginifolia, *Pers. Syn.* ii. 180.—Ind. or.
Noli-me-tangere, *S. Moore, in Journ. Bot.* xviii. (1880) 231.—Afr. trop.
pratensis, *S. Moore, l. c.* xv. (1877) 294.—Afr. trop.
procumbens, Heyne, ex Roth, Nov. Sp. 320 = boerhaviaefolia.
procumbens, Pers. Syn. ii. 180 = glomerata.
procurrens, Nees, in DC. Prod. xi. 267 = boerhaviaefolia.
pungens, *Klotzsch, in Peters, Reise Mossamb. Bot.* 211.—Afr. trop.
repens, Roth, Nov. Sp. 321 = molluginifolia.
rubiaefolia, Schum. in Dansk. Vidensk. Selsk. Afh. iv. (1828) 66 = boerhaviaefolia.
saturejaefolia, *Pers. Syn.* ii. 180.—Afr. austr.
scindica, Stocks, ex T. Anders. in Journ. Linn. Soc. vii. (1864) 36 = linariaefolia.
serrulata, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 24.—Afr. trop.
setosa, *Nees, in DC. Prod.* xi. 265.—Afr. austr.
sindica, Stocks, ex T. Anders. in Journ. Linn. Soc. ix. (1867) 500 = linariaefolia.
spathularis, *T. Anders. l. c.* vii. (1864) 35.—Afr. austr.
spiculifolia, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 85.—Ins. Socotr.
squarrosa, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 35.—Afr. austr.
Togodelia, *Solms, ex Schweinf. Beitr. Fl. Aethiop.* 108.—Abyss.
verticillata, Heyne, ex Wall. Cat. n. 2409 = Cardanthera verticillata.
Welwitschii, *S. Moore, in Journ. Bot.* xviii. (1880) 231.—Afr. trop.

BLEPHARISPERMUM, Benth. in DC. Prod. vii. 709 (1839) = BLEPHAROPHYLLUM, Klotzsch = **Scyphogyne**, Brongn. (Ericac.).

BLEPHARISPERMUM, Wight, ex DC. in Wight, Contrib. Bot. Ind. 11 (1834). *COMPOSITAE*, Benth. & Hook. f. ii. 295.
LEUCOBLEPHARIS, Arn. in Mag. Zool. & Bot. ii. (1838) 422.
petiolare, *DC. in Wight, l. c.* 12.—Ind. or.
spinulosum, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 335.—Afr. trop.
subsessile, *DC. in Wight, Contrib. Bot. Ind.* 12.—Ind. or.
zanguebaricum, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 336.—Afr. trop.

BLEPHARISTEMMA, Benth. in Wall. Cat. n. 6320 (1832). *RHIZOPHOREAE*, Benth. & Hook. f. i. 682.
corymbosum, *Benth. l. c.*—Ind. or.

BLEPHAROCALYX, Berg. in Linnaea, xxvii. (1854) 412 = **Myrtus**, Tourn.
acuminatissimus, Berg, l. c. 413; et in Mart. Fl. Bras. xiv. i. 421.—Reg. Argent.
acuminatus, Berg, ll. cc. 414, 426 (= Myrcia acuminata?).—Bras.
affinis, Berg, in Mart. Fl. Bras. xiv. i. 605.—Bras.
amarus. Berg, l. c. 422.—Reg. Argent.
angustifolius, Berg, l. c. 421.—Bras.
angustissimus, Berg, l. c.—Bras.
apiculatus, Berg, l. c. 425.—Bras.
brunneus, Berg, l. c. 427.—Bras.
canescens, Berg, l. c. 426.—Bras.
cisplatensis, Griseb. in Goett. Abh. xxiv. (1879) 126.—Reg. Argent.
cuspidatus, Berg, in Mart. Fl. Bras. xiv. i. 424.—Bras.
depauperatus, Berg, in Linnaea, xxvii. (1854) 415; et in Mart. l. c. 428.—Bras.
ellipticus, Berg, ll. cc. 415 et 429.—Bras.
lanceolatus, Berg, in Mart. Fl. Bras. xiv. i. 422.—Bras.
longipes, Berg, l. c. 423.—Bras.
parvifolius, Berg, l. c. 606.—Bras.
picrocarpus, Berg, l. c. 427.—Bras.
pilosus, Berg, l. c. 422.—N. Granat.
ramosissimus, Berg, l. c. 424.—Bras.

BLEPHAROCALYX:—
salicifolia, Berg, in Linnaea, xxvii. (1854) 413 = Myrtus salicifolia.
Serra, Berg, in Mart. Fl. Bras. xiv. i. 605.—Bras.
sessilifolius, Berg, l. c. 428.—Bras.
strictus, Berg, l. c. 423.—Bras.
Tweediei, Berg, in Linnaea, xxvii. (1854) 415; et in Mart. Fl. Bras. xiv. i. 429.—Bras.
villosus, Berg, in Mart. Fl. Bras. xiv. i. 425.—Bras.
Widgreni, Berg, l. c. 427.—Bras.

BLEPHAROCARYA, F. Muell. Fragm. ix. 15 (1878). *SAPINDACEAE*.
involucrigera, *F. Muell. l. c.* 16.—Austral.

BLEPHAROCHLAMYS, Presl, Epim. Bot. 245 (1849) = **Mystropetalon**, Harv. (Balanoph.).
capensis, Presl, l. c. 246 = M. Thomii.

BLEPHAROCHLOA, Endl. Gen. 1352 (1841) = **Leersia**, Sw. (Gramin.).
ciliata, Endl. l. c. = L. ciliata?

BLEPHARODON, Decne. in DC. Prod. viii. 603 (1844). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 756.
PTYCHOLEPIS, Griseb. in Lechl. Berb. Am. Austr. 58 (1857).
ampliflorus, *Fourn. in Mart. Fl. Bras.* vi. iv. 304.—Bras.
asparagoides, *Decne. in DC. Prod.* viii. 604.—Bras.
asterion, *Fourn. in Mart. Fl. Bras.* vi. iv. 307.—Bras.
bicolor, *Decne. in DC. Prod.* viii. 604.—Bras.
bicuspidatus, *Fourn. in Mart. Fl. Bras.* vi. iv. 306.—Bras.
bracteatus, *Fourn. l. c.* 307.—Bras.
Decaisnei, *Fourn. l. c.* 305.—Bras.
diffusum, *Decne. in DC. Prod.* viii. 605.—Bras.
grandiflorum, *Benth. Pl. Hartw.* 215.—N. Granat.
laurifolium, Decne. in DC. Prod. viii. 603 = Nephradenia laurifolia.
lineare, Decne. l. c. = Nephradenia linearis.
linearifolium, Benth. & Hook. f. Gen. ii. 756, sphalm. = laurifolium.
longipedicellatum, *Fourn. in Mart. Fl. Bras.* vi. iv. 306.—Bras.
mucronatum, *Decne. in DC. Prod.* viii. 603.—Mexic.
neriifolium, Decne. l. c. = Nephradenia neriifolia.
pallidum, *Decne. l. c.*—Bras.
salicinum, *Decne. l. c.* 604.—Bras.?
Spruceanum, *Fourn. in Mart. Fl. Bras.* vi. iv. 307.—Bras.
triplinerve, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) ii. 319.—Venezuela.

BLEPHAROLEPIS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 442 (1836) = **Polpoda**, Presl (Ficoid.).
Zeyheriana, Nees, l. c. = P. capensis.

BLEPHAROLEPIS, Nees, in Mart. Fl. Bras. ii. i. 91 (1843) = **Scirpus**, Linn. (Cyperac.).
leucoblepharis, Nees, l. c.—Bras.

BLEPHAROPHYLLUM, Klotzsch, in Linnaea, xii. (1838) 216 = **Scyphogyne**, Brongn. (Ericac.).
divaricatum, Klotzsch, l. c. 215 = S. divaricata.

BLEPHAROSTEMMA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 398 = **Asperula**, Linn. (Rubiac.).
taurinum, Fourr. l. c. = A. taurina.

BLEPHILIA, Rafin. in Am. Monthly Mag. (1819) 190; et Journ. Phys. lxxxix. (1819) 98. *LABIATAE*, Benth. & Hook. f. ii. 1198.
Becki, *Rafin. in Atl. Journ.* 146.—Am. bor.
brevifolia, Rafin. New Fl. Am. iv. 96 = ciliata.
brevipes, *Rafin. l. c.*—Am. bor.
ciliata, *Rafin. in Am. Monthly Mag.* (1819) 190.—Am. bor.
heterophyla, Rafin. New Fl. Am. iv. 96 = ciliata?
hirsuta, *Benth. Lab. Gen. et Sp.* 320.—Am. bor.
lanceolata, Rafin. New Fl. Am. iv. 96 = hirsuta?
nepetoides, Rafin. l. c. = hirsuta.
pratensis, Rafin. l. c. = ciliata.

BLEPHILOMA, Rafin. Fl. Tellur. iii. 95 (1836) = **Phlomis**, Tourn. (Labiat.).
amplifolia, Rafin. l. c.—Hab.?

BLEPHISTELMA, Rafin. Fl. Tellur. iv. 103 (1836) = **Passiflora**, Linn.
aurantia, Rafin. l. c. = P. aurantia.

BLETIA, Ruiz & Pav. Prod. 119. t. 26 (1794). *ORCHIDEAE*, Benth. & Hook. iii. 513.
BLETIANA, Rafin. in Am. Monthly Mag. (1818) 268.
BLETILLA, Reichb. f. in Fl. des Serres, viii. (1853) 246.
DILOMILIS, Rafin. Fl. Tellur. iv. 43 (1836).
GYAS, Salisb. in Trans. Hort. Soc. i. (1812) 299.
HECABE, Rafin. Fl. Tellur. iv. 44 (1836).
JIMENSIA, Rafin. l. c. 38 (1836).
THIEBAUTIA, Colla, Hort. Ripul. 139 (1824).
acaulis, Reichb. f. in Walp. Ann. vi. 435 = Brassavola acaulis.
acuminata, Reichb. f. l. c. 427 = Laelia acuminata.
acutipetala, Hook. Bot. Mag. t. 3217 = verecunda.
adenocarpa, *Reichb. f. in Bonplandia*, iv. (1856) 216.—Mexic.
albida, Reichb. f. in Walp. Ann. vi. 428 = Laelia albida.
amazonica, Reichb. l. c. 434 = Brassavola amazonica.
amboinensis, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 180 = Phaius amboinensis.
anceps, Reichb. f. in Walp. Ann. vi. 418 = Laelia anceps.
angustata, Gaudich. in Freyc. Voy. Bot. 421 = Spathoglottis plicata.
angustata, Reichb. f. in Walp. Ann. vi. 434 = Brassavola angustata.
anomala, *Rich. & Gal. in Ann. Sc. Nat.* Sér. III. i. (1845) 23.—Mexic.
aphylla, Nutt. Gen. Am. ii. 194 = Hexalectris aphylla.
attenuata, *Reichb. f. Xenia Orchid.* ii. 64.—Ind. occ.
autumnalis, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 19 = Laelia autumnalis.
bicallosa, D. Don, Prod. Fl. Nep. 30 = Liparis bituberculata?
Boothiana, Reichb. f. Xenia Orchid. ii. 51 = Laelia Boothiana.
bracteosa, *Griff. Notul.* iii. 326.—Ind. or.
campanulata, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 17.—Mexic.
capitata, [R. Br. in] Ait. Hort. Kew. ed. II. v. 206 = Elleanthus capitatus.
Casperiana, Reichb. f. in Walp. Ann. vi. 423 = Laelia purpurata.
catenulata, *Ruiz & Pav. Syst.* 229.—Peruv.
caulescens, Reichb. f. in Walp. Ann. vi. 431 = Laelia caulescens.
Ceboletta, Reichb. f. l. c. 435 = Brassavola Ceboletta.
ciliata, *Presl, Rel. Haenk.* 99.—Peruv.
cinnabarina, Reichb. f. in Walp. Ann. vi. 430 = Laelia cinnabarina.
cinnamomea, Reichb. f. l. c. 418 = Laelia cinnamomea.
coccinea, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 16.—Mexic.
cordata, Reichb. f. in Walp. Ann. vi. 436 = Brassavola nodosa.
crispa, Reichb. f. l. c. 423 = Laelia crispa.
crispina, Reichb. f. l. c. 419 = Schomburgkia crispa.
cucullata, Reichb. f. l. c. 433 = Brassavola cucullata.
Dabia, D. Don, Prod. Fl. Nep. 30 = Eulophia campestris?
densiflora, *Griff. Notul.* iii. 328.—Ind. or.
Digbyana, Reichb. f. in Walp. Ann. vi. 422 = Laelia Digbyana.
domingensis, Reichb. f. l. c. 432 = Broughtonia lilacina.
elegans, Reichb. f. l. c. 427 = Laelia elegans.
ensiformis, *Ruiz & Pav. Syst.* 230.—Peruv.
Eulophiae, Reichb. f. in Walp. Ann. vi. 439 = Tetramicra tenera.
euspatha, Reichb. f. l. c. 424 = Laelia euspatha.
flabellata, Reichb. f. l. c. 422 = Laelia xanthina.
flava, Reichb. f. l. c. 431 = Laelia flava.
flava, Wall. ex Lindl. Gen. et Sp. Orch. 127 = Phaius maculatus.
florida, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. v. 206.—Ind. occ.
fulgens, *Reichb. f. in Bonplandia*, iii. (1855) 221.—Mexic.
furfurea, Reichb. f. in Walp. Ann. vi. 428 = Laelia furfuracea.
gebina, Lindl. in Journ. Hort. Soc. ii. (1847) 307 = hyacinthina.
glauca, Reichb. f. in Walp. Ann. vi. 423 = Laelia glauca.

BLETIA:—
gloriosa, Reichb. f. in Walp. Ann. vi. 419 = Schomburgkia crispa.
gracilis, *Lodd. Bot. Cab.* t. 1977.—Mexic.
graminifolia, D. Don, Prod. Fl. Nep. 29 = Arundina bambusifolia.
grandiflora, La Llave & Lex. Nov. Veg. Desc. fasc. ii. (Orch. Opusc.) 17 = Laelia grandiflora.
grandis, Reichb. f. in Walp. Ann. vi. 424 = Laelia grandis.
guinecnsis, *G. Don, ex Sweet, Hort. Brit.* ed. I. 492, nomcn.—Guinea.
havannensis, *Lindl. Bot. Reg.* (1838) *Misc.* 28.—Cuba.
Humboldtii, Reichb. f. in Walp. Ann. vi. 430 = Schomburgkia Humboldtii.
humilis, Link & Otto, Ic. Pl. Rar. 53. t. 27 = Maxillaria gracilis.
hyacinthina, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. v. 206. —China; Japon.
irrorata, Reichb. f. in Walp. Ann. vi. 426 = Laelia irrorata.
jucunda, *Linden & Reichb. f. in Bonplandia*, iii. (1855) 221.—Mexic.
Landsbergii, *Reichb. f. in Nederl. Kruidk. Arch.* iv. (1859) 317.—Am. trop.
latifolia, *Gaudich. Bot. Freyc. Voy.* 421.—Ins. Molucc.
lilacina, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 23.—Mexic.
Lindeni, Reichb. f. in Walp. Ann. vi. 431 = Laelia Lindenii.
Lindleyana, Reichb. f. Xenia Orchid. ii. 65 = Laelia Lindleyana.
lineata, Reichb. f. in Walp. Ann. vi. 436 = Brassavola lineata.
lobata, Reichb. f. l. c. 424 = Laelia Boothiana.
longipes, Reichb. f. Xenia Orchid. ii. 59 = Laelia longipes.
Lundii, Reichb. f. & Warm. in Otia Bot. Hamb. 92 = Laelia Lundii.
Lyonsii, Reichb. f. in Walp. Ann. vi. 418 = Schomburgkia Lyonsia.
marginata, Reichb. f. l. c. 419 = Schomburgkia marginata.
Martiana, Reichb. f. l. c. 434 = Brassavola Martiana.
Masuca, D. Don, Prod. Fl. Nep. 30 = Calanthe Masuca.
montana, Reichb. f. in Walp. Ann. vi. 445 = Octadesmia serratifolia.
nipponica, *Franch. & Sav. Enum. Pl. Jap.* ii. 511.—Japon.
nodosa, Reichb. f. in Walp. Ann. vi. 437 = Brassavola nodosa.
obcordata, *Lindl. Gen. et Sp. Orch.* 123.—Ind. or.
Ortgiesiana, *Reichb. f. in Otto, Hamb. Gartenz.* xvi. (1860) 420.—Hab.?
pallida, Lodd. Bot. Cab. t. 629 = florida.
Parkinsonii, *Hook. Bot. Mag.* t. 3736.—Mexic.
parviflora, Reichb. f. in Walp. Ann. vi. 439 = Tetramicra parviflora.
parviflora, *Ruiz & Pav. Syst. Veg.* 231.—Peruv.
patula, *Hook. Bot. Mag.* t. 3518.—Ins. S. Doming.
Paulinae, F. Muell. Fragm. vi. 95 = Spathoglottis Paulinae.
peduncularis, Reichb. f. in Walp. Ann. vi. 426 = Laelia peduncularis.
Perrinii, Reichb. f. l. c. 421 = Laelia Perrinii.
platyphylla, Reichb. f. ex Griseb. Cat. Pl. Cub. 264 = sarcophylla.
praestans, Reichb. f. in Walp. Ann. vi. 425 = Laelia praestans.
pratensis, Reichb. f. l. c. 439 = Tetramicra rigida.
pulchella, Hort.—Cf. Gard. Chron. (1850) 168 = verecunda.
pumila, Reichb. f. in Walp. Ann. vi. 421 = Laelia pumila.
Pumilio, Reichb. f. l. c. 433 = Brassavola Pumilio.
punctata, *La Llave & Lex. Nov. Veg. Desc.* fasc. ii. (*Orch. Opusc.*) 15.—Mexic.
purpurata, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 23 = Arethusa rosea.
purpurata, Reichb. f. in Walp. Ann. vi. 423 = Laelia purpurata.
purpurea, DC. Huit. Not. Pl. Rar. Jard. Genèv. 23 = verecunda.
reflexa, *Lindl. Bot. Reg.* xxi. (1836) t. 1760.—Peruv.
repanda, *Ruiz & Pav. Syst. Veg.* 229.—Peruv.
retusa, Reichb. f. in Walp. Ann. vi. 435 = Brassavola retusa.

BLETIA:—
rhopalorrhachis, Reichb. f. l. c. 437 = Brassavola rhopalorrhachis.
rigida, Reichb. f. l. c. 438 = Tetramicra rigida.
Roezlii, *Reichb. f. in Linnaea*, xli. (1877) 7.—Mexic.
rosea, Reichb. f. in Walp. Ann. vi. 420 = Schomburgkia rosea.
rubescens, Reichb. f. l. c. 425 = Laelia rubescens.
rupestris, Reichb. f. l. c. 431 = Laelia rupestris.
sanguinea, Pocpp. & Endl. Nov. Gen. et Sp. i. 56. t. 95 = catenulata.
sarcophylla, *Reichb. f. in Flora*, xlviii. (1865) 278.—Ind. occ.
Schilleriana, Reichb. f. in Walp. Ann. vi. 424 = Laelia Schilleriana.
schoenina, Reichb. f. l. c. 439 = Tetramicra schoenina.
Schomburgkii, Reichb. f. l. c. = Tetramicra rigida.
secunda, *Lindl. Bot. Reg.* (1840) *Misc.* 57.—Mexic.
Shepherdii, Hook. Bot. Mag. t. 3319 = verecunda.
Sherrattiana, *Batem. in Hook. l. c.* t. 5646.—N. Granat.
speciosa, H. B. & K. Nov. Gen. et Sp. i. 342 = Laelia grandiflora.
splendens, *Linden, Illustr. Hortic.* xxviii. (1881) 64, nomen.—Bras.
stricta, *C. Presl, Rel. Haenk.* 98.—Ins. Luzon.
superbiens, Reichb. f. in Walp. Ann. vi. 418 = Laelia superbiens.
sylvatica, Boj. Hort. Maurit. 318 = Calanthe sylvatica.
Tankervilleae, R. Br. in Bot. Mag. t. 1924 = Phaius grandiflorus.
tenera, A. Rich. Fl. Cub. Fancrog. ii. 242 = Tetramicra tenera.
tibicinis, Reichb. f. in Walp. Ann. vi. 429 = Schomburgkia tibicina.
tuberculata, Reichb. f. l. c. 434 = Brassavola tuberculata.
tuberculosa, Spreng. Syst. iii. 744 = Phaius tuberculosus.
undulata, Reichb. f. in Walp. Ann. vi. 420 = Schomburgkia undulata.
uniflora, *Ruiz & Pav. Syst. Veg.* 230.—Peruv.
venosa, Reichb. f. in Walp. Ann. vi. 438 = Brassavola venosa.
verecunda, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. v. 206. —Florida; Ind. occ.
villosa, Spreng. Syst. iii. 743 = Phaius villosus.
violacea, Reichb. f. in Walp. Ann. vi. 426 = Laelia violacea.
Wagcnerii, *Reichb. f. in Bonplandia*, ii. (1854) 22.—Venczuela.
Wallisii, *Reichb. f. in Linnaea*, xli. (1877) 115.—N. Granat.
Wendlandi, Reichb. f. in Walp. Ann. vi. 431 = Laelia Wendlandi.
Woodfordii, Hook. Bot. Mag. t. 2719 = Phaius maculatus.
xanthina, Reichb. f. in Walp. Ann. vi. 425 = Laelia xanthina.

BLETIANA, Rafin. in Am. Monthly Mag. (1818) 268 = **Bletia**, Ruiz & Pav. (Orchid.).

BLETILLA, Reichb. f. in Fl. des Serres, Sér. I. viii. (1851–53) 246 = **Bletia**, Ruiz & Pav. (Orchid.).
florida, Reichb. f. l. c. = Bletia florida.
Gebinae, Reichb. f. l. c. = Bletia hyacinthina.

BLETTI, Steud. Nom. ed. II. i. 210 (1840) = BLATTI, Adans. = **Sonneratia**, Linn. f. (Lythrar.).

BLIGHIA, Kon. in Kon. & Sims, Ann. Bot. ii. (1806). 571. t. 16, 17. *SAPINDACEAE*. [Benth. & Hook. f. i. 399].
AKEA, Stokes, Bot. Mat. Med. ii. 353 (1812).
AKEESIA, Tussac, Fl. Ant. i. 66 (1808).
sapida, *Kon. in Kon. & Sims, Ann. Bot.* ii. (1806) 571. —Afr.; Ind. occ.
unijugata, Baker, in Oliver, Fl. Trop. Afr. i. 427.—Afr. trop.
zambesiaca, *Baker, l. c.*—Afr. trop.

BLINKWORTHIA, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 430. t. 4. *CONVOLVULACEAE*, Benth. & Hook. f. ii. 869.
lycioides, *Choisy, l. c.*—Burma.

BLITANTHUS, Reichb. Cat. Hort. Dresd. (1824); Handb. 238 (1837) = **Acroglochin**, Schrad. (Chenopod.).
nepalensis, Reichb. ll. cc. = A. chenopodioides.

BLITUM, [Tourn.] Linn. Gen. ed. I. 20 (1737) = **Chenopodium**, Linn.
acuminatum, Schur, Enum. Pl. Transs. 571 = C. rubrum.
antarcticum, Hook. f. Fl. Antarct. ii. 549 = C. antarcticum.
atriplicinum, F. Muell. in Trans. Vict. Inst. (1855) 133 = C. atriplicinum.
Bonus-Henricus, C. A. Mey. Fl. Alt. i. 11, nota = C. Bonus-Henricus.
botryoides, Drej. in Fl. Dan. t. 2894 = C. rubrum.
californicum, S. Wats. in Proc. Am. Acad. ix. (1874) 101 = C. californicum.
capitatum, Linn. Sp. Pl. 4 = C. capitatum.
carinatum, C. A. Mey. in Ledeb. Fl. Alt. i. 11 adnot. = C. carinatum.
chenopodioides, Bernh. ex Steud. Nom. ed. II. i. 210 = petiolare.
chenopodioides, Lam. Encyc. ii. 11 = C. capitatum.
chenopodioides, Nutt. Gen. Am. i. 4 = Monolepis chenopodioides.
crassifolium, Reichb. Fl. Germ. Excurs. 582 = C. rubrum.
cristatum, F. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 73 = C. cristatum.
gangeticum, Moench, Meth. 359 = Amaranthus gangeticus.
genuinum, Koch, ex Buek, Ind. DC. Prod. iii. 61 (est sectio generis).
glandulosum, Moq. in DC. Prod. xiii. II. 82 = C. carinatum.
glaucum, Koch, in Sturm, Deutschl. Fl. Heft 75 = C. glaucum.
graecizans, Moench, Meth. 359 = Amaranthus Blitum.
lividum, Moench, l. c. = Amaranthus lividus.
maritimum, Nutt. Gen. Am. ii. Add. post Ind. [12] C. rubrum.
maius, Scop. Fl. Carn. ed. II. ii. 237 = Amaranthus Blitum.
melancholicum, Moench, Meth. 359 = Amaranthus gangeticus.
Nuttallianum, Roem. & Schult. Mant. i. 65 = Monolepis chenopodioides.
oleraceum, Moench, Meth. 359 = Amaranthus oleraceus.
petiolare, Link, Enum. Hort. Berol. i. 8.—Tataria?
polymorphum, C. A. Mey. in Ledeb. Fl. Alt. i. 13 = C. rubrum.
Pumilio, C. A. Mey. l. c. 11, adnot. = C. Pumilio.
rubrum, C. A. Mey. l. c. = C. rubrum.
Sanguisorba, Pritz. Ic. Ind. 155, sphalm. = Poterium Sanguisorba.
tataricum, Mill. Gard. Dict. ed. VIII. n. 3 = C. capitatum.
tenue, Moq. in DC. Prod. xiii. II. 81 = Roubieva multifida.
terminale, Stokes, Bot. Comment. i. 11 = C. capitatum.
virgatum, Linn. Sp. Pl. 4 = C. capitatum.
viride, Moench, Meth. 359 = C. album.

BLOCHMANNIA, Reichb. Consp. 163 (1828) = **Triplaris**, Linn. (Polygon.).
peruviana, Pocpp. ex Meissn. in DC. Prod. xiv. 173 = T. Poeppigiana.
Weigeltiana, Reichb. Consp. 163 = T. surinamensis.

BLONDEA, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 110 = **Sloanea**, Linn. (Tiliac.).
latifolia, Rich. l. c.—Guiana.

BLONDIA, Neck. Elem. ii. 115 (1790) = **Tiarella**, Linn. (Saxifr.).
biternata, Rafin. Fl. Tellur. ii. 75 = Astilbe decandra.
trifoliata, Rafin. l. c. = T. trifoliata.

BLOOMERIA, Kellogg, in Proc. Calif. Acad. ii. 11 (1863). *LILIACEAE*, Benth. & Hook. f. Gen. iii. 801.
aurea, *Kellogg, l. c.*—Calif.
Clevelandi, *S. Wats. in Proc. Am. Acad.* xx. (1885) 376.—Calif.

BLUFFIA, Delile, Sem. Hort. Hamb. (1835) (Cf. Linnaea, x. (1836) 114); Nees, Fl. Afr. Austr. Gram. (1836) = **Panicum**, Linn. (Gramin.).
Eckloniana, Nees, in Lindl. Introd. Nat. Syst. ed. II. 447.—Afr. austr.

BLUMEA, G. Don, Gen. Syst. i. 578 (1831) = BLUMIA, Spreng. = **Saurauja**, Willd. (Ternstroem.).

BLUMEA, Reichb. Consp. 209 (1828) = **Neesia**, Blume (Malvac.).
altissima, Reichb. ex Steud. Nom. ed. II. i. 211 = N. altissima.

BLUMEA, DC. in Guill. Arch. Bot. ii. (1833) 514.
COMPOSITAE, Benth. & Hook. f. ii. 289.
DOELLIA, Sch. Bip. in Walp. Rep. ii. 953 (1843).
PLACUS, Lour. Fl. Cochinch. 496 (1790).
abyssinica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 392.—Abyss.
acutata, DC. in Nouv. Ann. Mus. Par. iii. (1834) 409 = laciniata.
alata, DC. Prod. v. 448 = Laggera alata.
amethystina, Hance, in Journ. Bot. vi. (1868) 173 = glomerata.
amplectens, *DC. in Wight, Contrib. Bot. Ind.* 13.—Ind. or.
amplectens, Thw. Enum. Pl. Zeyl. 163 = bifoliata.
anagallidifolia, DC. in Wight, Contrib. Bot. Ind. 13 = amplectens.
angustifolia, *Thw. Enum. Pl. Zeyl.* 164.—Zeylan.
appendiculata, *DC. Prod.* v. 447.—Java.
arenaria, DC. in Wight, Contrib. Bot. Ind. 13 = amplectens.
arfakiana, *Martelli, in Nuov. Giorn. Bot.* xv. (1883) 292.—N. Guin.
Arnottiana, Steud. Nom. ed. II. i. 210 = Inula Cappa.
aromatica, *DC. Prod.* v. 446.—Reg. Himal.; Burma.
aurita, DC. in Wight, Contrib. Bot. Ind. 16 = Laggera aurita.
axillaris, DC. Prod. v. 434 = lacera.
baccharoides, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 391 = Pluchea Dioscoridis.
balsamifera, *DC. Prod.* v. 447.—Ind. or.; Burma; Malaya.
barbata, *DC. in Wight, Contrib. Bot. Ind.* 14.—Ind. or.
Belangeriana, *DC. Prod.* v. 444.—Ind. or.
bifoliata, *DC. in Wight, Contrib. Bot. Ind.* 14.—Ind. or.; Burma; Malaya.
Bovei, Vatke, in Linnaea, xxxix. (1875) 485 = abyssinica.
bovina, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 173 = oxydonta.
cernua, DC. Prod. v. 436 = hieracifolia.
Chamissoniana, DC. l. c. 434 = Wightiana.
chinensis, *DC. l. c.* 444.—Ind. or.; China; Java.
chinensis, Hook. & Arn. Bot. Beech. Voy. 195 = Inula Cappa.
chinensis, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 294 = hieracifolia.
cichoriifolia, DC. ex Decne. in Nouv. Ann. Mus. Par. iii. (1834) 410 = laciniata.
cinerascens, DC. Prod. v. 439 = lacera.
Clarkei, Hook. f. Fl. Brit. Ind. iii. 267 = elongata.
Commersonii, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 38.—Ind. or.
crassifolia, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 394.—Abyss.
crepidifolia, DC. Prod. v. 437 = laciniata.
crinita, *Arn. in Nov. Act. Nat. Cur.* xviii. (1836) 348.—Zeylan.
crinita, Steetz, in Seem. Bot. Voy. Herald, 387 = hieracifolia.
cuneifolia, DC. Prod. v. 441 = lacera.
Cunninghami, *DC. l. c.* 435.—Austral.
densiflora, *DC. l. c.* 446.—Ind. or.; Malaya; Ins. Pacif.
diffusa, *R. Br. ex Benth. Fl. Austral.* iii. 525.—Austral.
diffusa, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 187.—Burma.
Dregeana, *DC. Prod.* v. 436.—Afr. austr.
dregeanoides, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 392 = lacera.
elongata, *DC. Prod.* v. 445.—Reg. Himal.; Burma; Penang.
eriantha, *DC. in Wight, Contrib. Bot. Ind.* 15.—Ind. or.
excisa, DC. Prod. v. 446 = densiflora.
fasciculata, DC. l. c. 442 = glomerata.
fistulosa, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 187 = glomerata.
flava, DC. Prod. v. 439 = Laggera flava.
flexuosa, *C. B. Clarke, Compos. Ind.* 86.—Ind. or.
fontinalis, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 173 = lacera.

BLUMEA:—
gariepina, *DC. Prod.* v. 448.—Afr. austr.
glandulosa, Benth. Fl. Hongk. 177 = laciniata.
glandulosa, DC. in Wight, Contrib. Bot. Ind. 14 = lacera.
glandulosa, Thw. Enum. Pl. Zeyl. 163 = membranacea.
glomerata, *DC. in Wight, Contrib. Bot. Ind.* 15.—Ind. or.; Malaya; China.
glutinosa, *DC. Prod.* v. 434.—Madag.
gracilis, DC. l. c. 443 = lacera.
grandiflora, Zipp. ex Span. in Linnaea, xv. (1841) 165 = Capparis spinosa.
grandis, DC. Prod. v. 447 = densiflora.
guineensis, DC. l. c. 449 = Laggera aurita.
Hamiltoni, DC. l. c. 439 = hieracifolia.
Heyneana, DC. in Wight, Contrib. Bot. Ind. 15 = lacera.
hieracifolia, *DC. l. c.* 15.—Ind. or.; Burma; Malaya.
hieracifolia, Hook. f. & Thoms. ex Fl. Brit. Ind. iii. 267 = elongata.
hispidula, DC. ex C. B. Clarke, in Fl. Brit. Ind. iii. 672 = membranacea.
holosericea, Dalz. & Gibs. Bomb. Fl. 126 = glomerata.
holosericea, DC. Prod. v. 442 = lacera.
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 269.—Reg. Himal.
hymenophylla, DC. Prod. v. 440 = membranacea.
integrifolia, *DC. Prod.* v. 433.—Austral.
Jacquemontii, *Hook. f. Fl. Brit. Ind.* iii. 265.—Ind. or.
javanica, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 239.—Java.
Kingii, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 265.—Ind. or.
lacera, *DC. in Wight, Contrib. Bot. Ind.* 14.—Ind. or.; Malaya; Ins. Mascar.
laciniata, *DC. Prod.* v. 436.—Ind. or.; Malaya; China.
lactucaefolia, DC. l. c. 435 = lacera.
lapsanoides, DC. in Wight, Contrib. Bot. Ind. 15 = virens.
leptoclada, DC. Prod. v. 443 = glomerata.
leptoclada, Dalz. & Gibs. Bomb. Fl. 125 = lacera.
Leschenaultiana, DC. Prod. v. 435 = Wightiana.
leucanthema, *DC. l. c.* 436 (sp. dubia).—Nepal.
longifolia, DC. l. c. 446 = myriocephala?
macrophylla, *DC. l. c.* 446.—Java.
macrostachya, DC. l. c. 442 = hieracifolia.
malabarica, *Hook. f. Fl. Brit. Ind.* iii. 267.—Ind. or.
Malcolmii, *Hook. f. l. c.* 266.—Ind. or.
manillensis, *DC. Prod.* v. 435.—Ins. Luzon.
membranacea, *DC. l. c.* 440.—Ind. or.; Malaya.
Metziana, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 266 = Belangeriana.
Milnei, Seem. in Bonplandia, ix. (1861) 257; Fl. Vit. 141 = densiflora,
muralis, DC. Prod. v. 440 = membranacea.
Musra, DC. l. c. 435 = lacera.
myriocephala, *DC. l. c.* 445.—Reg. Himal.; Burma.
myriocephala, Thw. Enum. Pl. Zeyl. 163 = spectabilis.
napifolia, DC. l. c. 440 = lacera.
natalensis, *Sch. Bip. in Walp. Rep.* ii. 971.—Afr. austr.
neilgherrensis, *Hook. f. Fl. Brit. Ind.* iii. 261.—Ind. or.
nitida, Wall. ex C. B. Clarke, Comp. Ind. 85 = chinensis.
nitida, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 244.—Java.
nodiflora, *Hook. f. Fl. Brit. Ind.* iii. 262.—Burma.
obovata, C. B. Clarke, Comp. Ind. 77 = Jacquemontii.
obovata, DC. Prod. v. 446.—Nepal.
oligocephala, DC. in Wight, Contrib. Bot. Ind. 13 = bifoliata.
oloptera, *DC. Prod.* v. 448.—Afr. trop.
oxydonta, *DC. in Wight, Contrib. Bot. Ind.* 15.—Ind. or.; Burma; Malaya; China.
parvifolia, DC. Prod. v. 437 = Wightiana.
paucifolia, DC. l. c. 440 = membranacea.
Perrottetiana, DC. l. c. 443 = lacera.
Petitiana, *A. Rich. Tent. Fl. Abyss.* i. 395.—Abyss.
phagnaloides, A. Rich. l. c. 393 = Phagnalon scoparium.
phyllostachya, DC. Prod. v. 438 = Wightiana.
procera, *DC. l. c.* 445.—Ind. or.; Burma.
pterodonta, DC. in Wight, Contrib. Bot. Ind. 15 = Laggera pterodonta.
pubiflora, DC. Prod. v. 434 = amplectens.

BLUMEA:—
purpurascens, A. Rich. Tent. Fl. Abyss. i. 395 = Laggera pterodonta.
purpurea, DC. Prod. v. 443 = glomerata.
racemosa, DC. l. c. 442 = glomerata.
riparia, DC. l. c. 444 = chinensis.
runcinata, DC. l. c. 438 = laciniata.
runcinata, Wall. Cat. n. 3087 = lacera.
salvifolia, *DC. Prod.* v. 448.—Ins. Borbon.
salviodora, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 246.—Java.
saxatilis, *Zoll. & Mor. l. c.* 243.—Java.
semivestita, DC. Prod. v. 445 = procera.
senecioides, DC. l. c. 439 = membranacea.
senecioides, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 38 = Pulicaria foliolosa.
senegalensis, DC. Prod. v. 449 = Conyza senegalensis.
sericans, *Hook. f. Fl. Brit. Ind.* iii. 262.—Burma.
sessiliflora, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 410.—Ins. Timor.
sessilifolia, DC. Prod. v. 447 = myriocephala.
sikkimensis, *Hook. f. Fl. Brit. Ind.* iii. 268.—Reg. Himal.
solidaginoides, *DC. Prod.* v. 443.—Afr. trop.
sonchifolia, DC. l. c. 438 = laciniata.
spectabilis, *DC. l. c.* 445.—Ind. or.
spinellosa, DC. l. c. 443 = hieracifolia?
squarrosa, C. B. Clarke, Comp. Ind. 87 = myriocephala.
subcapitata, *DC. Prod.* v. 439.—Reg. Himal.
subsimplex, DC. l. c. 441 = membranacea.
sylvatica, *DC. l. c.* 447.—Java.
tenella, DC. in Wight, Contrib. Bot. Ind. 13 = bifoliata?
tetraptera, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 313.—Ins. Philipp.
thyrsoidea, Sch. Bip. in Bot. Zeit. xxiv. (1866) 166 = lacera.
timorensis, *DC. ex Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 411.—Ins. Timor.
tomentosa, *A. Rich. Tent. Fl. Abyss.* i. 394.—Abyss.
trichophora, DC. Prod. v. 436 = Wightiana.
trigona, DC. l. c. 437 = lacera.
trisulca, DC. l. c. 440 = Conyza stricta.
vernonioides, DC. l. c. 447 = Laggera alata.
villosa, Sch. Bip. ex Hook. f. Fl. Brit. Ind. iii. 263 = lacera.
viminea, *DC. Prod.* v. 442.—Ins. Timor.
virens, *DC. in Wight, Contrib. Ind. Bot.* 14.—Ind. or.
viscosula, DC. Prod. v. 441 = membranacea.
Wallichii, *C. B. Clarke, Comp. Ind.* 87.—Assam.
Wightiana, *DC. in Wight, Contrib. Bot. Ind.* 14.—Geront. trop.
Zollingeriana, C. B. Clarke, Comp. Ind. 90 = balsamifera.

BLUMENBACHIA, Koel. Gram. Gall. et Germ. 28 (1802) = **Sorghum**, Moench (Gramin.).
haleppensis, Koel. l. c. 29 = S. halepense.

BLUMENBACHIA, Schrad. in Goett. Gel. Anz. (1825) 1705; Blumenb. (1827) 6. t. 1. *LOASEAE*, Benth. & Hook. f. i. 805.
CAJOPHORA, Endl. Gen. 931 (1839).
GRIPIDEA, Miers, in Trans. Linn. Soc. xxv. (1866) 227. t. 28.
RAPHISANTHE, Lilja, in Linnaea, xv. (1841) 263.
carduifolia, *Ball, in Journ. Linn. Soc.* xxii. (1885) 39.—Peruv.
cernua, *Griseb. in Goett. Abh.* xix. (1874) 152.—Reg. Argent.
chuquitensis, *Hook. f. Bot. Mag.* t. 6143.—Peruv.
contorta, Hook. f. l. c. t. 6134 = grandiflora.
coronata, *Haage & Schmidt, in Rev. Hortic.* (1874) 58.—Chili.
Domeykoana, *Phil. in Linnaea*, xxxiii. (1864–65) 75.—Chili.
Espigneera, *C. Gay, Fl. Chil.* ii. 434.—Chili.
grandiflora, *G. Don, Gen. Syst.* iii. 62.—Peruv.
Hieronymi, *Urb. in Eichl. Jahrb. Berl.* iii. (1884) 249.—Reg. Argent.
insignis, *Schrad. in Goett. Gel. Anz.* (1825) 1706.—Reg. Argent.
latifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 209.—Paraguay.
micrantha, *G. Don, Gen. Syst.* iii. 62.—Peruv.
multifida, Griseb. in Goett. Abh. xxiv. (1879) 139 = Hieronymi.

BLUMENBACHIA:—
multifida, *Hook. Bot. Mag.* t. 3599.—Reg. Argent.
palmata, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 208.—Paraguay.
parviflora, Gill. ex G. Don, Gen. Syst. iii. 62 = insignis.
Prietea, *C. Gay, Fl. Chil.* ii. 435.—Chili.
pterosperma, *G. Don, Gen. Syst.* iii. 62.—Peruv.
punicea, *G. Don, l. c.*—Peruv.
scandens, *Meyen, Reise*, i. 310.—Chili.
sepiaria, *G. Don, Gen. Syst.* iii. 62.—Peruv.
sylvestris, *Poepp. Fragm. Syn. Fl. Chil.* 25.—Chili.

BLUMEODENDRON, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 245 = **Mallotus**, Lour. (Euphorb.).
Muelleri, Kurz, l. c. = Paracroton pendulus.
Tokbrai, Kurz, l. c.—Ins. Andaman.

BLUMIA, Nees, ex Blume, in Verh. Bat. Gen. ix. (1823) 147 = **Talauma**, Juss. (Magnol.).
Candollii, Nees, l. c.; et in Flora, viii. (1825) 152 = T. Candollii.

BLUMIA, Spreng. Syst. iii. 126 (1826) = **Saurauja**, Willd. (Ternstr.).
japonica, Steud. Nom. ed. II. i. 212, sphalm. = seq.
javanica, Spreng. Syst. iii. 126 = S. Blumia.

BLUTAPARON, Rafin. New Fl. Am. iv. 45 (1836) **Philoxerus**, R. Br. (Amarant.).
brevifolium, Rafin. l. c. = P. vermicularis.
repens, Rafin. l. c. 46 = P. vermicularis.

BLYSMUS, Panz. ex Roem. & Schult. Syst. ii. Mant. 41 (1824) = **Scirpus**, Tourn. (Cyperac.).
Bonanni, Steud. Nom. ed. II. i. 212 = Schoenus Bonanni.
brevifolius, Decne. in Jacquem. Voy. Bot. 173. t. 173 = S. Caricis.
compressus, Panz. in Link, Hort. Berol. i. 278 = S. Caricis.
rufus, Link, l. c. = S. rufus.

BLYTTIA, Arn. in Mag. Zool. et Bot. ii. 420 (1838) = **Vincetoxicum**, Moench (Asclep.).
arabica, Arn. l. c. = V. fruticulosum.

BLYTTIA, Fries, Novit. Fl. Suec. Mant. ii. 2 (1839) = **Cinna**, Linn. (Gramin.).
suaveolens, Fries, l. c. = C. pendula.

BLYXA, Noronha, ex Thou. Gen. Nov. Madag. 4 (1806). *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 451.
DIPLOSIPHON, Decne. in Jacquem. Voy. Bot. 166. t. 167 (1844).
SAIVALA, Jones, in As. Res. iv. (1799) 275.
Aubertii, *Rich. in Mem. Inst. Fr.* (1811) 4.—Madag.
graminea, *Thou. ex Steud. Nom.* ed. II. i. 212.—Madag.
javanica, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 121.—Java.
octandra, *Planch. ex Thw. Enum. Pl. Zey.* 332.—Zeylan.
Roxburghii, *Rich. in Mém. Inst. Fr.* (1811) 77. t. 5.—As. trop.; Austral.
Saivala, *Steud. Nom.* ed. II. i. 212.—Ind. or.

BOADSCHIA, All. Fl. Pedem. i. 248 = BOHADSCHIA, Crantz = **Peltaria**, Linn. (Crucif.).

BOARIA, DC. Prod. viii. 299 (1844) = **Maytenus**, Molina (Celastr.).
Molinae, DC. l. c. = M. Boaria.

BOBAEA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 215 = **Bobea**, Gaudich. (Rubiac.).

BOBARTIA, Linn. Diss. Dass. 3 (1747); Fl. Zeyl. 17 (1747); Salisb. in Trans. Hort. Soc. i. (1812) 313. *IRIDEAE*, Benth. & Hook. f. iii. 698.
HECASTA, Soland. ex Schum. in Skr. Naturk-Selsk. iii. (1793) 10.
aphylla, *Ker-Gawl. Gen. Irid.* 30.—Afr. austr.
aurantiaca, Sweet, Hort. Brit. ed. III. 662 = Homeria aurantiaca.
filiformis, *Ker-Gawl. Gen. Irid.* 30.—Afr. austr.

BOBARTIA:—
gladiata, *Ker-Gawl. Gen. Irid.* 30; *Sweet, Hort. Brit.* ed. I. 395.—Afr. austr.
indica, *Linn. Sp. Pl.* 54.—Afr. austr.
juncea, Salisb. in Trans. Hort. Soc. i. (1812) 313 = indica.
Linnaei, Pritz. Ind. Ic. ii. 47 perperam = indica.
spathacea, Ker-Gawl. Gen. Irid. 30 = indica.
umbellata, *Ker-Gawl. l. c.* 31.—Afr. austr.

BOBEA, Gaudich. in Freyc. Voy. Bot. 473. t. 93 (1826). *RUBIACEAE*, Benth. & Hook. f. ii. 101.
BOBAEA, A. Rich. in Mém. Soc. Hist Nat. Par. v. (1830) 215.
BURNEYA, Cham. & Schlecht. in Linnaea, iv. (1829) 190.
brevipes, *A. Gray, in Proc. Am. Acad.* iv. (1860) 36.—Ins. Sandvic.
elatior, Gaudich. in Bot. Freyc. Voy. 473. t. 93 = Timonius Gaudichaudi.
glabra, Korth. in Nederl. Kruidk. Arch. ii. (1851) II. 211 = Timonius Jambosella.
hirsutiuscula, Teijsm & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 247 = Polyphragmon sericeum.
inacquisepala, *Miq. Fl. Ind. Bat. Suppl.* 545.—Ins. Banca.
mutabilis, Korth. in Nederl. Kruidk. Arch. ii. (1851) II. 212 = Polyphragmon mutabile.
putaminosa, F. Muell. Fragm. iv. 92 = Guettardella putaminosa.
sericantha, *Miq. Fl. Ind. Bat.* ii. 355.—Malaya.
Wallichiana, Korth. in Nederl. Kruidk. Arch. ii. (1851) II. 211 = Timonius Rumphii.

BOBU, Adans. Fam. ii. 88 (1763) = **Symplocos**, Linn. (Styrac.).

BOBUA, DC. Prod. iii. 23 (1828) = **Symplocos**, Linn. (Styrac.).
arborea, Miers, in Journ. Linn. Soc. xvii. (1879) 305 = S. arborea.
atroviridis, Miers, l. c. 304 = S. spicata.
baptica, Miers, l. c. 305 = S. baptica.
caerulescens, Miers, l. c. = S. caerulescens.
cerasifolia, Miers, l. c. 304 = S. cerasifolia.
glaucescens, Miers, l. c. 305 = S. glaucescens.
gracilis, Miers, l. c. = S. gracilis.
japonica, Miers, l. c. 306 = S. japonica.
lancifolia, Miers, l. c. = S. lancifolia.
laurina, DC. Prod. iii. 24 = S. spicata.
Lenormandiana, Miers, in Journ. Linn. Soc. xvii. (1879) 304 = S. Lenormandiana.
leptostachya, Miers, l. c. 306 = S. lancifolia.
lucida, Miers, l. c. 305 = S. lucida.
montana, Miers, l. c. = S. montana.
myrtacea, Miers, l. c. 306 = S. myrtacea.
neriifolia, Miers, l. c. = S. neriifolia.
nitida, Miers, l. c. 305 = S. nitida.
oligostachya, Miers, l. c. 304 = S. spicata.
prunifolia, Miers, l. c. 306 = S. caudata.
rotundifolia, Miers, l. c. 305 = S. rotundifolia.
stravadioides, Miers, l. c. 304 = S. stravadioides.
theophrastaefolia, Miers, l. c. 306 = S. theophrastaefolia.
Vieillardi, Miers, l. c. 305 = S. Vieillardi.

BOCA, Vell. Fl. Flum. 232; v. t. 113 (1825) = **Banara**, Aubl. (Samyd.).
serrata, Vell. ll. cc. = Banara Vellozii.

BOCAGEA, St. Hil. Fl. Bras. Mer. i. 41. t. 9 (1825). *ANONACEAE*, Benth. & Hook. f. i. 29.
OXANDRA, A. Rich. Ess. Fl. Cub. 45. t. 8 (1842).
SAGERAEA, Dalz. in Hook. Kew Journ. iii. (1851) 207.
acuminata, Steud. Nom. ed. II. i. 212 = Oropheа acuminata.
alba, *St. Hil. Fl. Bras. Mer.* i. 42.—Bras.
aromatica, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 36.—N. Granat.
cancscens, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) 70.—Am. austr.
celebica, *Blume, Fl. Jav. Anon.* 88.—Ins. Celebes.
coriacea, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 93.—Zeylan.
corymbosa, *Blume, Fl. Jav. Anon.* 85.—Java.

BOCAGEA:—
Dalzellii, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 92.—Ind. or.
elliptica, *Hook. f. & Thoms. l. c.*—Burma; Malaya.
enneandra, *Spreng. Syst.* iv. *Cur. Post.* 144.—Java.
espintana, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) 71.—Peruv.
Gaudichaudiana, *Baill. Adansonia*, viii. (1867–68) 183.—Cochinch.
heterantha, *Baill. l. c.* 173.—Madag.
hexandra, Blume, Fl. Jav. Anon. 83 = javensis.
javensis, *Spreng. Syst.* iv. *Cur. Post.* 144.—Java.
latifolia, *Blume, Fl. Jav. Anon.* 89.—Java.
leucodermis, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) 71.—Bras.
longipedunculata, *Mart. Fl. Bras.* xiii. I. 45.—Bras.
lutea, Baill. Hist. Pl. i. 215 = Alphonsea lutea.
multiflora, *Mart. Fl. Bras.* xiii. I. 45.—Bras.
nervosa, Pierre, Fl. For. Cochin. sub t. 16 = Kingstonia nervosa.
obliqua, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 93.—Zeylan.
philastreana, *Pierre, Fl. For. Cochin.* t. 16.—Cochinch.
pisocarpa, *Blume, Fl. Jav. Anon.* 90. t. 45.—Java.
polyandra, *C. Presl, Rel. Haenk.* ii. 77.—Ins. Luzon.
polycarpa, Steud. Nom. ed. II. i. 212 = Orophea polycarpa.
Thwaitesii, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 92.—Zeylan.
ventricosa, Baill. Hist. Pl. i. 215 = Alphonsea ventricosa.
verrucosa, Baill. l. c. sphalm. = praec.
viridis, *St. Hil. Fl. Bras. Mer.* i. 42. t. 9.—Bras.
Weddelliana, *Baill. Adansonia*, viii. (1867–68) 170.—Bras.

BOCCO, Steud. Nom. ed. I. 112 (1821) sphalm. = **Bocoa**, Aubl.

BOCCONIA, Plum. ex Linn. Gen. ed. I. 32 (1737). *PAPAVERACEAE*, Benth. & Hook. f. i. 53.
MACLEAYA, R. Br. App. Denh. & Clapp. Trav. 218, in adnot. (1826).
cordata, *Willd. Sp. Pl.* ii. 841.—China; Japon.
ferruginea, *Roezl, in Belg. Hortic.* xxiv. (1874) 39.—Peruv.
frutescens, *Linn. Sp. Pl.* 505.—Mexico; Peruv.
glauca, Salisb. Prod. 377 = frutescens.
integrifolia, *Humb. & Bonpl. Pl. Aequin.* i. 119. t. 35.—Mexico; Peruv.
japonica, Hort. ex André, in Rev. Hortic. (1866) 369 = cordata.
jedoënsis, Carr. l. c. 340 = cordata.
quercifolia, Moench, Meth. Suppl. 122 = frutescens.
sinuatifolia, Stokes, Bot. Mat. Med. iii. 8 = frutescens.
subtomentosa, L'Hérit. ex Stahl, Estud. Fl. Puertorico, Foll. ii. 34 = frutescens.

BOCKIA, Scop. Introd. 106 (1777) = **Mouriria**, Juss. (Melast.).

BOCOA, Aubl. Pl. Gui. Suppl. 38. t. 391 (1775) = **Inocarpus**, Forst. (Legum.).
edulis, Baill. Adansonia, ix. (1868–70) 237 = I. edulis.
guianensis, Steud. Nom. ed. I. 112 = seq.
prouacensis, Aubl. Pl. Gui. Suppl. 38.—Guiana.

BOCQUILLONIA, Baill. Adansonia, ii. 225 (1861–62). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 313.
brachypoda, *Baill. Adansonia*, xi. (1873) 127.—N. Caled.
brevipes, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 166.—N. Caled.
grandidens, *Baill. Adansonia*, xi. (1873) 128.—N. Caled.
sessiliflora, *Baill. l. c.* ii. (1861–62) 226.—N. Caled.
spicata, *Baill. l. c.* 227.—N. Caled.

BODWICHIA, Walp. Rep. i. 808 (1842), sphalm. = **Bowdichia**, H. B. & K. (Legum.).

BOEA, Comm. ex Lam. Encyc. i. 401 (1783). *GESNERACEAE*, Benth. & Hook. f. ii. 1023.
BAEA, Comm. ex Juss. Gen. 121 (1789).
CHLETERUS, Rafin. Prim. Somiol. 25 (1814).

BOEA :—
DORCOCERAS, Bunge, Enum. Pl. Chin. Bor. 54 (1832).
alata, Pers. Syn. i. 15 = Calceolaria petiolaris.
amplexicaulis, Parish, ex C. B. Clarke, Comm. et Cyrt. Beng. t. 84 = Phylloboea amplexicaulis.
borneensis, Scheff. ex H. O. Forbes, in Journ. Linn. Soc. xix. (1882) 298 = Didymocarpus Schefferi.
Cardwellia, F. Muell. ex C. B. Clarke, in DC. Monog. Phan. v. 145 = Commersonii.
cochinchinensis, *C. B. Clarke, l. c.* 143.—Cochinch.
Commersonii, *R. Br. in Benn. Pl. Jav.* 120.—China.
dictyoneura, *Hance, in Journ. Bot.* xxi. (1883) 169.—China.
flosculosa, C. B. Clarke, Comm. et Cyrt. Beng. t. 83 = multiflora.
Hancei, C. B. Clarke, in DC. Monog. Phan. v. 144 = dictyoneura.
herbacea, *C. B. Clarke, l. c.* 142.—Ins. Formosa.
hygrometrica, *R. Br. in Benn. Pl. Jav.* 120.—China.
hygroscopica, *F. Muell. Fragm.* iv. 146.—Austral.
magellanica, Lam. Illustr. i. 53. t. 15 = Commersonii.
Minahassae, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 415.—Ins. Celebes.
multiflora, *R. Br. in Benn. Pl. Jav.* 120.—Ind. or.
philippensis, *C. B. Clarke, in DC. Monog. Phan.* v. 146.—Ins. Philipp.
plantaginea, Pers. Syn. i. 15 = Calceolaria plantaginea.
praliniana, J. St. Hil. Expos. i. 279 = magellanica.
primuloides, Miq. in Ann. Mus. Lugd. Bat. iii. 190 = Oreocharis primuloides.
prolixa, *C. B. Clarke, in DC. Monog. Phan.* v. 143.—Burma.
punctata, Pers. Syn. i. 15 = Calceolaria punctata.
rufescens, *Franch. in Bull. Soc. Linn. Par.* i. (1885) 449.—China.
Swinhoii, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 231.—China.
Treubii, *H. O. Forbes, in Journ. Linn. Soc.* xix. (1882) 297.—Sumatra.
triandra, Pers. Syn. i. 15 = Calceolaria triandra.
Urvillei, *C. B. Clarke, in DC. Monog. Phan.* v. 147.—Ins. Waigiou.
violacea, Pers. Syn. i. 15 = Calceolaria violacea.
Wallichii, *R. Br. in Benn. Pl. Jav.* 120.—Burma.

BOEBERA, Willd. Sp. Pl. iii. 2125 (1803) = **Dyssodia**, Cav. (Compos.).
alternifolia, Moç. & Sessé, ex DC. Prod. v. 639 = Adenophyllum Porophyllum.
Cavanillesii, Spreng Syst. iii. 544 = Adenophyllum coccineum.
chrysanthemoides, Willd. Sp. Pl. iii. 2125.—D. chrysanthemoides.
fastigiata, H. B. & K. Nov. Gen. et Sp. iv. 198 = D. chrysanthemoides.
glandulosa, Pers. Syn. ii. 459 = D. chrysanthemoides, glandulosa.
incana, Lindl. Bot. Reg. t. 1602 = D. pubescens.
Porophyllum, H. B. & K. Nov. Gen. et Sp. iv. 198 = Adenophyllum Porophyllum.
pubescens, Spreng. Syst. iii. 544 = D. pubescens.
subintegerrima, Spreng. l. c. = D. subintegerrima.
tagetiflora, Spreng. l. c. = D. chrysanthemoides.

BOECKHIA, Kunth, Enum. Pl. iii. 448 (1841) = **Hypodiscus**, Nees (Restiac.).
laevigata, Kunth, l. c. 450 = H. albo-aristatus.
striata, Kunth, l. c. 449 = H. striatus.

BOEHMERIA, Jacq. Enum. Pl. Carib. 9 (1760).
URTICACEAE, Benth. & Hook. f. iii. 387.
GYMNOGYNE, F. Didr. Ind. Sem. Haun. (1850).—Cf. Linnaea, xxix. (1857–58) 737.
DURETIA, Gaudich. in Bot. Freyc. Voy. 500 (1826).
SPLITGERBERA, Miq. Comm. Phytogr. 133. t. 14 (1840).
acalyphoides, Blume, Mus. Bot. Lugd. Bat. ii. 225 = Pouzolzia acalyphoides.
acuminata, Hook. & Arn. ex Wedd. in DC. Prod. xvi. 1. 235 = Pipturus albidus.
acuminata, Pers. Syn. ii. 556 = Elatostema acuminatum.
acuminata, *Wedd. in DC. Prod.* xvi. 1. 209.—N. Caled.
albida, Hook. & Arn. Bot. Beech. Voy. 96 = Pipturus albidus.
alienata, Willd. Sp. Pl. iv. 341 = Pouzolzia indica.

BOEHMERIA :—
amplissima, Blume, Mus. Bot. Lugd. Bat. ii. 219 = stipularis.
ampullacea, H. B. & K. Nov. Gen. et Sp. ii. 35 = caudata.
angustata, Hassk. Cat. Hort. Bog. Alt. 79 = Debregeasia velutina.
angustifolia, H. B. & K. Nov. Gen. et Sp. ii. 34 = Phenax angustifolius.
arachnoidea, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 423 = Pouzolzia macrura.
arborea, Desf. Cat. Hort. Par. ed. III. 348 = Gesnouinia arborea.
arborescens, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 541 = caudata.
argentea, Guill. ex Wedd. in DC. Prod. xvi. 1. 235[19] = Pipturus argenteus.
argentea, *Linden, in Belg. Hortic.* viii. (1858) 166.—Mexic.
arguta, *Mart. ex Miq. in Mart. Fl. Bras.* iv. 1. 186.—Am. trop.
aspera, Blume, Mus. Bot. Lugd. Bat. ii. 206, in nota = Pouzolzia aspera.
aspera, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855–56) 349. t. 11.—N. Granat.
australis, *Endl. Prod. Fl. Norfolk.* 38.—Ins. Norfolk.
ballotaefolia, H. B. & K. Nov. Gen. et Sp. ii. 35 = Phenax ballotaefolius.
bifida, Blume, Mus. Bot. Lugd. Bat. ii. 222 = biloba, Wedd.
biloba, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 100.—Java.
biloba, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 199.—Japon.
Blumei, *Wedd. in DC. Prod.* xvi. 1. 204.—Ins. Philipp.
Boiviniana, Blume, Mus. Bot. Lugd. Bat. ii. 227 = Pouzolzia Boiviniana.
boliviana, Blume, l. c. 206, in nota = Pouzolzia Boiviniana.
brevirostris, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 201.—Peruv.
bullata, *H. B. & K. Nov. Gen. et Sp.* ii. 35.—Ecuador.
caerulea, Blume, Mus. Bot. Lugd. Bat. ii. 209 = Cyphophus coeruleus.
calophleba, *C. Moore & F. Muell. in F. Muell. Fragm.* viii. 11.—Austral.
candicans, Hassk. Pl. Jav. Rar. 203; Cat. Hort. Bog. Alt. 79 = nivea.
candidissima, Hassk. Cat. Hort. Bog. Alt. 79 = Leucosyke javanica.
Candolleana, Gaudich. in Freyc. Voy. Bot. 499 = Pipturus argenteus.
canescens, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 200 = platyphylla.
capensis, *Spreng. Syst.* iii. 844.—Afr. austr.
castaneaefolia, A. Cunn. ex Bureau, in DC. Prod. xvii. 250 = Pseudomorus Brunoniana.
caudata, Poir. in Bonpl. Jard. Malm. t. 15 = platyphylla.
caudata, *Sw. Prod. Veg. Ind. Occ.* 34.—Am. trop.
caudigera, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 199 = platyphylla.
celebica, *Blume, Mus. Bot. Lugd. Bat.* ii. 217.—Ins. Celebes.
celtidifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 34.—Ecuador.
cephalida, Pers. Syn. ii. 556 = Procris cephalida.
chinensis, Pers. l. c. 557 = Procris cantonensis.
cinerascens, Blume, Mus. Bot. Lugd. Bat. ii. 203 = Pouzolzia cinerascens.
cinerascens, Hassk. Cat. Hort. Bog. Alt. 79 = Pipturus argenteus.
clidemioides, Miq. Pl. Jungh. 34 = platyphylla.
cochinchinensis, Pers. Syn. ii. 557 = Procris cochinchinensis.
cochinchinensis, Spreng. Syst. iii. 844 = Pouzolzia indica.
comosa, Wedd. in DC. Prod. xvi. 1. 205 = sidaefolia.
compacta, *Blume, Mus. Bot. Lugd. Bat.* ii. 210.—Java.
conglobata, Blume, l. c. 204 = Pouzolzia conglobata.
crenata, Blume, l. c. 208 = Cyphophus macrocephalus?
Cummingiana, Blume, l. c. ii. 199 = heterophylla.
cuspidata, *Blume, l. c.* 216 (sp. dubia).—Nepal.
cuspidata, Wedd. in Arch. Mus. Hist. Nat. Par. viii. (1855–56) 345 = ramiflora.
cylindrica, Hb. ex Blume, Mus. Bot. Lugd. Bat. ii. 218 = elongata.

BOEHMERIA :—
cylindrica, *Sw. Prod. Veg. Ind. Occ.* 34.—Am. bor. et austr.
dasypoda, Miq. in Mart. Fl. Bras. iv. 1. 188 = cylindrica.
densiflora, *Hook. & Arn. Bot. Beech. Voy.* 271.—China.
depauperata, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 202 = malabarica.
dichotoma, Hassk. Cat. Hort. Bog. Alt. 79 = Debregeasia velutina.
didymogyne, *Wedd. in DC. Prod.* xvi. 1. 204.—Burma.
diffusa, Wedd. in Arch. Mus. Hist. Nat. Par. viii. (1855–56) 356 = sidaefolia.
discolor, Poepp. ex Blume, Mus. Bot. Lugd. Bat. ii. 206 = Pouzolzia discolor.
diversiflora, Miq. in Zoll. Syst. Verz. Ind. Archip. 101, 104 = Zollingeriana.
diversifolia, Miq. Pl. Jungh. 34 = Maoutia diversifolia.
diversifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 202 = Pavonii.
Drummondiana, Wedd. l. c. 201 = cylindrica.
elliptica, Wedd. l. c. 200 = platyphylla.
elongata, *Blume, Mus. Bot. Lugd. Bat.* ii. 218.—Hab.?
elongata, Hort. ex Steud. Nom. ed. II. i. 212 = platyphylla.
elongata, Fisch. ex Jacq. Eclog. t. 135 = Phenax vulgaris.
erythropoda, *Miq. in Zoll. Syst. Verz. Ind. Archip.* 101, 104.—Malaya.
excelsa, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 202.—Ins. Juan Fernand.
fagifolia, Pers. Syn. ii. 556 = Elatostema fagifolium.
fallax, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855–56) 346.—Peruv.
Fernandesiana, C. Gay, Fl. Chil. v. 360 = Urtica glomerulaeflora.
flagelliformis, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 310 = caudata.
florida, *Miq. in Mart. Fl. Bras.* iv. 1. 187.—Am. trop.
formicaria, Poepp. ex Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 201 = Pouzolzia formicaria.
frondosa, D. Don, Prod. Fl. Nep. 59 = Pouzolzia viminea.
frutescens, D. Don, l. c. = Maoutia Puya.
frutescens, Thunb. Fl. Jap. 367. n. 98 = Villebrunea frutescens.
glabra, Steud. Nom. ed. II. i. 192 = Neraudia melastomaefolia.
glochidiata, Hassk. Pl. Jav. Rar. 206 = Pouzolzia indica.
glomerulifera, Miq. in Zoll. Syst. Verz. Ind. Archip. 101, 104 = malabarica.
grandifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 199 = platyphylla.
greviaefolia, *Presl, Bot. Bemerk.* 110.—Hab.?
guatemalana, Blume, Mus. Bot. Lugd. Bat. ii. 206 = Pouzolzia guatemalana.
Hamiltoniana, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 199 = platyphylla.
Harveyi, Seem. Fl. Vit. t. 62 = Cyphophus heterophylla.
Helferii, *Blume, Mus. Bot. Lugd. Bat.* ii. 201.—Burma.
heteroidea, *Blume, l. c.* 216 (sp. dubia).—Ind. or.
heterophylla, Blume, l. c. 204 = Blumei.
heterophylla, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855–56) 351.—Ins. Philipp.
hirta, Roth, ex Blume, Mus. Bot. Lugd. Bat. ii. 231 = Pouzolzia Rothiana.
hirta, Sw. Prod. Veg. Ind. Occ. 34 = Phenax urticifolius.
hirta, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 202.—Peruv.
hispidula, *Blume, Mus. Bot. Lugd. Bat.* ii. 223.—Sumatra.
holosericea, *Blume, l. c.* 221.—Japon.
Huegeliana, *Blume, l. c.* 218 (sp. dubia).—Ind. or.
humilis, Miq. Pl. Jungh. 33 = platyphylla.
hypoleuca, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 264 = Debregeasia hypoleuca.
incana, Hassk. Cat. Hort. Bog. Alt. 79 = Pipturus velutinus.
interrupta, Guill. in Ann. Sc. Nat. Sér. II. vii. (1857) 182 = platyphylla.
interrupta, Willd. Sp. Pl. iv. 340 = Flenrya interrupta.
Jacquiniana, *Blume, Mus. Bot. Lugd. Bat.* ii. 219.—Hab.?
Janatsi-itsigo, Siebold, in Verh. Bat. Gen. xii. (1830) 70 = Debregeasia edulis.

BOEHMERIA :—
japonica, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iii. 131. —Japon.
javanica, Hassk. Cat. Hort. Bog. Alt. 79 = Fleurya interrupta.
laevis, Blume, Mus. Bot. Lugd. Bat. ii. 206 in nota = Pouzolzia virgata.
lateriflora, Muhl. ex Willd. Sp. Pl. iv. 342 = cylindrica.
latifolia, Blume, Mus. Bot. Lugd. Bat. ii. 207 = Cyphophus latifolius.
Lindeniana, *Blume, l. c.* 226.—N. Granat.
littoralis, Sw. Prod. Veg. Ind. Occ. 34 = cylindrica.
longispica, Steud. in Flora, xxxiii. (1850) 260 = japoniea.
lutescens, Blume, Mus. Bot. Lugd. Bat. ii. 209 = Cyphophus lutescens.
macrocephala, Blume, l. e. 208, in obs. = Cyphophus macroeephalus.
maerophylla, *Hornem. Hort. Hafn.* ii. 890.—Hab.?
maerophylla, *D. Don, Prod. Fl. Nep.* 60.—Reg. Himal.; Burma.
macrophylla, Sieb. & Zuec. in Abh. Akad. Mueneh. iv. III. (1846) 215 = japonica.
macrostachya, Wedd. in Ann. Se. Nat. Sér. IV. i. (1854) 199 = platyphylla.
macrothyrsa, Miq. in Zoll. Syst. Verz. Ind. Arehip. 100, 103 = Laportia decumana.
maculata, Pers. Syn. ii. 556 = Elatostema maculatum.
malabariea, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855-56) 355.—Ind. or.; Malaya.
massuriensis, Blume, Mus. Bot. Lugd. Bat. ii. 216 = platyphylla.
mauritiana, Wedd. in Ann. Se. Nat. Sér. IV. i. (1854) 200 = platyphylla.
melanocarpa, Blume, Mus. Bot. Lugd. Bat. ii. 210 = Cyphophus melanoearpus.
melastomaefolia, Hook. & Arn. Bot. Beeeh. Voy. 96 = Neraudia melastomaefolia.
melastomoidea, Griff. Notul. iv. 386 = Pouzolzia pentandra.
microcarpa, Wedd. in Ann. Se. Nat. Sér. IV. i. (1854) 201 = platyphylla.
microceltis, Miq. in Zoll. Syst. Verz. Ind. Arehip. 100, 104 = Pouzolzia viminea.
mollieoma, *Miq. in Zoll. l. c.*—Java.
mollis, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 203 = Cyphophus macrocephalus.
moluccana, Blume, Mus. Bot. Lugd. Bat. ii. 208 = Cyphophus macroeephalus.
moluccana, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 201 = Cyphophus macrocarpus.
montieola, *Blume, Mus. Bot. Lugd. Bat.* ii. 227.—Sumatra.
nana, D. Don, Prod. Fl. Nep. 60 = Pouzolzia indiea.
neglecta, *Blume, Mus. Bot. Lugd. Bat.* ii. 200.—Java.
nepalensis, Wedd. in Areh. Mus. Hist. Nat. Par. viii. (1855-56) 383 = Villebrunea fruteseens.
nervosa, Madden, in Journ. As. Soc. Beng. xvii. (1848) 1. 403 = rugulosa.
nivea, *Gaudich. in Bot. Freyc. Voy.* 499.—As. trop. (eult.).
nudiflora, *Willd. Sp. Pl.* iv. 343.—Venezuela.
obliqua, Poepp. ex Blume, Mus. Bot. Lugd. Bat. ii. 206, adnot. = Pouzolzia obliqua.
odontophylla, Miq. Pl. Jungh. 35 = Maoutia odontophylla.
ourantha, Miq. l. e. 33 = platyphylla.
ovalis, Miq. in Zoll. Syst. Verz. Ind. Archip. 100 = Pouzolzia viminea.
ovata, Blume, Mus. Bot. Lugd. Bat. ii. 206, adnot. = Pouzolzia ovata.
ovata, Steud. Nom. ed. II. i. 212 = Neraudia melastomaefolia.
palmata, *Hassk. Cat. Hort. Bog. Alt.* 79.—Hab.?
parvifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 203 = Droguetia diffusa.
pauciflora, Blume, Mus. Bot. Lugd. Bat. ii. 201 = Droguetia diffusa.
Pavonii, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 202.—Peruv.
penduliflora, Wedd. l. e. 199 = maerophylla.
peruviana, Blume, Mus. Lugd. Bat. ii. 224 = caudata.
petiolaris, H. B. & K. Nov. Gen. et Sp. ii. 36 = caudata.
phyllostachya, Miq. in Mart. Fl. Bras. iv. 1. 187 = cylindriea.
pilosiuscula, Hassk. Cat. Hort. Bog. Alt. 79 = platyphylla.

BOEHMERIA :—
platanifolia, *Franch. & Sav. Enum. Pl. Jap.* i. 440.—Japon.
platyphylla, *D. Don, Prod. Fl. Nep.* 60.—As. et Afr. trop.; Ins. Pacif.
polystachya, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 200.—Reg. Himal.; Burma.
procridioides, Blume, Mus. Bot. Lugd. Bat. ii. 204 = Pouzolzia proeridioides.
propinqua, Deene. Herb. Timor. 163 = Pipturus argenteus.
prostrata, Blume, Mus. Bot. Lugd. Bat. ii. 209 = Cyphophus prostratus.
Puya, Hook. in Hook. Kew Journ. iii. (1851) 316 = Maoutia Puya.
racemiflora, Forst. ex Blume, Rumphia, iv. 37 = Phytoerene hirsuta.
radicans, Nees & Mart. in Nov. Aet. Nat. Cur. xi. (1823) 35 = Pilea rhizobola.
ramiflora, Bedd. Forest. Man. Bot. t. 27. f. 2 = malabarica.
ramiflora, *Jacq. Enum. Pl. Carib.* 31.—Ind. oee.; Mexic.
repanda, Hassk. Cat. Hort. Bog. Alt. 79 = Pipturus repandus.
repens, Nees & Mart. in Nov. Aet. Acad. Nat. Cur. xi. (1823) 36 = Pilea hirtella.
repens, *Wedd. in DC. Prod.* xvi. 1. 201.—Cuba.
rhynehophylla, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 202.—Am. austr.
rigida, *Benth. in Hook. Niger Fl.* 519.—Afr. trop.
rotundifolia, D. Don, Prod. Fl. Nep. 60 = platyphylla.
rubeseens, *Hassk. Cat. Hort. Bog. Alt.* 79 = Villebrunea rubeseens.
rubescens, Jacq. Fragm. 8 = Gesnouinia arborea.
rugosa, Pers. Syn. ii. 556 = Phenax ballotaefolius.
rugosissima, Miq. Pl. Jungh. i. 32 = platyphylla.
rugosissima, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 200.—Java.
rugulosa, *Wedd. l. c.* 200.—Reg. Himal.
salicifolia, D. Don, Prod. Fl. Nep. 60 = Debregeasia hypoleuea.
sanguinea, Hassk. Cat. Hort. Bog. Alt. 79 = Pouzolzia viminea.
seabra, *Hassk. l. c.*—As. trop.?
scabrella, Gaudieh. Bot. Freyc. Voy. 499 = platyphylla.
scabrella, Hort. ex Blume, Mus. Bot. Lugd. Bat. ii. 215 = eaudata.
sericea, Blume, Mus. Bot. Lugd. Bat. ii. 208 = Cyphophus moluceanus.
sidaefolia, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 203.—Reg. Himal.; Burma; Java.
Sieboldiana, Blume, Mus. Bot. Lugd. Bat. ii. 220 = platyphylla.
spicata, Thunb. in Trans. Linn. Soe. ii. (1794) 330 = platyphylla.
spieiflora, *Blume, Mus. Bot. Lugd. Bat.* ii. 217 (sp. dubia).—Ind. or.
spieigera, *Spreng. Syst.* iii. 845.—Venezuela.
squamigera, Wedd. in Ann. Se. Nat. Sér. IV. i. (1854) 203 = Chamabainia euspidata.
stipularis, *Wedd. l. c.* 200.—Ins. Masear.
strigosa, Blume, Mus. Bot. Lugd. Bat. ii. 226 = Pavonii.
strigosula, Blume, l. c. 203 = Pouzolzia viminea.
subperforata, *Wedd. in Arch. Mus. Hist. Nat. Par.* viii. (1855-56) 383 (sp. dubia).—Ind. or.
sumatrana, Miq. Pl. Jungh. 32 = platyphylla.
sylvatica, Hassk. Cat. Hort. Bog. Alt. 79 = Villebrunca sylvatiea.
taitensis, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 200 = platyphylla.
tenacissima, Gaudieh. Bot. Freye. Voy. 500 = nivea.
ternifolia, D. Don, Prod. Fl. Nep. 59 = platyphylla.
tomentosa, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 200 = platyphylla.
travancorica, Bedd. Forest. Man. Bot. 225 = malabariea.
trinervata, Pers. Syn. ii. 556 = Gyrotaenia trinervata.
trinervis, Miq. Pl. Jungh. 36 = Pipturus repandus.
uliginosa, Mor. ex Wedd. in DC. Prod. xvi. 1. 235[37] = Phenax uliginosus.
ulmifolia, Wedd. in Ann. Sc. Nat. Sér. IV. i. (1854) 202 = fallax.
urantha, Blume, Mus. Bot. Lugd. Bat. ii. 221 = platyphylla.

BOEHMERIA :—
urticaefolia, Pers. Syn. ii. 556 = Phenax urticaefolius.
urticaefolia, Spreng. Syst. iii. 844 = Pilea rupipendia.
velutina, Deene. Herb. Timor. 163 = Pipturus velutinus.
vestita, Blume, Mus. Bot. Lugd. Bat. ii. 209 = Cyphophus vestitus.
vialis, Casar. ex Wedd. in DC. Prod. xvi. 1. 235[37] = Phenax Sonneratii.
viminea, Blume, Mus. Bot. Lugd. Bat. ii. 202 = Pouzolzia viminea.
virgata, Guillem. in Ann. Se. Nat. Sér. II. vii. (1837) 182 = platyphylla.
Waitziana, Blume, Mus. Bot. Lugd. Bat. ii. 203 = Pouzolzia viminea.
Wightiana, Blume, l. e. 217 = platyphylla.
Zollingeriana, *Wedd. in Ann. Sc. Nat.* Sér. IV. i. (1854) 201.—Java.

BOEICA, C. B. Clarke, Comm. et Cyrt. Beng. 118. tt. 85 ad 87 (1874). *GESNERACEAE*, Benth. & Hook. f. ii. 1023.
BAEICA, C. B. Clarke, in DC. Monog. Phan. v. 134 (1883).
filiformis, *C. B. Clarke, Comm. et Cyrt. Beng.* 118. t. 87.—Reg. Himal.
fulva, *C. B. Clarke, l. c.* t. 85.—Reg. Himal.
Griffithii, *C. B. Clarke, l. c.* t. 86.—Assam.
hirsuta, *C. B. Clarke, in DC. Monog. Phan.* v. 136.—Assam?
porosa, *C. B. Clarke, l. c.*—Burma.

BOELIA, Webb, Otia Hisp. 20. t. 15, 16 (1853) = **Genista**, Linn. (Legumin.).
sphaerocarpa, Webb, l. c. = G. sphaerocarpa.

BOENNINGHAUSENIA, Reichb. Consp. 197 (1828). *RUTACEAE*, Benth. & Hook. f. i. 287.
PODOSTAURUS, Jungh. in Nat. en Geneesk. Arch. Neêrl. Indie ii. (1845) 45.
albiflora, *Reichb. l. c.*—Reg. Himal.; Japon.
japonica, Sieb. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 21 = albiflora.

BOENNINGHAUSIA, Spreng. Syst. iii. 153, 245 (1826) = **Chaetocalyx**, DC. (Legumin.).
vincentina, Spreng. l. c. 245 = C. vineentinus.

BOERHAVIA, Linn. Fl. Zeyl. 4 (1747) = **Boerhaavia**, Vaill. (Nyetag.).

BOERHAAVIA, Vaill. ex Linn. Syst. ed. I. (1735); Linn. Gen. ed. I. 8 (1737). *NYCTAGINEAE*, Benth. & Hook. f. iii. 5.
adscendens, Willd. Sp. Pl. i. 19 = repens.
africana, *Lour. Fl. Cochinch.* i. 16.—Afr. trop.
aggregata, Pav. ex Choisy, in DC. Prod. xiii. II. 429 = Nyctaginia capitata.
angustifolia, *Linn. Syst.* ed. XII. 52 (sp. dubia).—Hab.?
angustifolia, Span. in Linnaea, xv. (1841) 342 = repens.
anisophylla, *Torr. in Bot. Mex. Bound.* 171.—Texas.
arborea, Roem. & Schult. Syst. i. 66 = Pisonia hirtella.
arborescens, Lag. & Rodr. in Anal. Cienc. Nat. iv. (1801) 257 = Pisonia hirtella.
ascendens, Schum. Beskr. Guin. Pl. 17 = repens.
bastica, Bueh.-Ham. ex Wall. Cat. sub n. 6769 = repanda.
bicolor, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 168, 169 = repens.
braeteosa, *S. Wats. in Proc. Am. Acad.* xx. (1885) 370.—Texas.
Burchellii, *Choisy, in DC. Prod.* xiii. II. 455.—Afr. austr.
caribaea, Jacq. Obs. Bot. iv. 5 = repens.
chaerophylloides, Willd. Sp. Pl. i. 22 = Asirephia chaerophylloides.
coccinea, Mill. Gard. Dict. ed. VIII. n. 4 = repens.
Commersoni, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 484.—Madag.
cortensis, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 168 = repens.
erispa, *Heyne, in Wall. Cat.* n. 6771.—Ind. or.
decumbens, Vahl, Enum. i. 284 = paniculata.

BOERHAAVIA:—
depressa, *C. Sm. in Tuckey, Congo*, 249, nomen.—Ins. Cap. Verd.
diandra, Aubl. Pl. Gui. i. 4 = paniculata?
diandra, Burm. f. Fl. Ind. 3. t. 1 = erecta.
diandra, Linn. Sp. Pl. 1194 = hirsuta, paniculata.
dichotoma, *Hochst. ex Walp. Ann.* iii. 298.—Abyss.
dichotoma, Schimp. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 142 = grandiflora.
dichotoma, Vahl, Enum. i. 290 = plumbaginea.
diffusa, Engelm. & A. Gray, in Boston, Journ. Nat. Hist. v. (1847) 259 = viscosa.
diffusa, Linn. Sp. Pl. 3 = repens.
diffusa, Sw. Obs. 10 = paniculata.
discolor, *H. B. & K. Nov. Gen. et Sp.* ii. 215.—Ecuador.
eglandulosa, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 169 = repens.
elegans, *Choisy, in DC. Prod.* xiii. II. 453.—Nubia; Arabia; Ind. bor. occ.
elongata, Salisb. Prod. 56 = erecta.
erecta, Burm. f. Fl. Ind. t. 3. f. 2 = glutinosa.
erecta, Ell. Sketch, i. 41 = repens.
erecta, *Linn. Sp. Pl.* 3.—Am. bor.; Ind. occ.
eriosolena, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 322.—Mexic.
excelsa, Willd. Phyt. 1. n. 5; Sp. Pl. i. 21 = Oxybaphus micranthus?
fruticosa, *Dalz. in Dalz. & Gibs. Fl. Bomb.* 213.—Ind. or.
fruticosa, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 167 = plumbaginea.
gibbosa, Pav. ex A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 323 = Senkenbergia annulata.
gibbosa, Pav. ex Choisy, in DC. Prod. xiii. II. 457 = Tinantia gypsophiloides.
glabrata, Blume, Bijdr. 733 = repens.
glandulosa, *Anderss. Om Oarn. Galapagos.* 64.—Ins. Galap.
glomerata, *Klotzsch, in Schomb. Fauna et Fl. Guy.* 1131, nomen.—Guiana.
glutinosa, Miers, ex Bert. in Merc. Chil. (1829) 596 = discolor.
glutinosa, Vahl, Enum. i. 287 = diffusa?
Grahamii, A. Gray, in Am. Journ. Sc. Ser. II. xv. (1853) 323 = scandens.
grandiflora, A. Rich. Tent. Fl. Abyss. ii. 209 = plumbaginea.
Helenae, Roem. & Schult. Syst. Mant. i. 73 = verticillata.
hirsuta, *Linn. Mant.* 1170.—Ind. occ.; Am. bor.
hirsuta, Willd. Phyt. 1 = repens.
insularis, Boj. Hort. Maurit. 264 = repens.
laxa, Pers. Syn. i. 36 = paniculata.
leiosolena, *Torr. Bot. Mex. Bound.* 172.—Texas.
libyca, *Pomel, Nouv. Mat. Fl. Atl.* 336.—Algeria.
linearifolia, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 322.—N. Mexic.
litoralis, *H. B. & K. Nov. Gen. et Sp.* ii. 216.—Peruv.
maroccana, Ball, in Journ. Bot. xiii. (1875) 176 = repens.
microphylla, Bory, ex Steud. Nom. ed. II. i. 213 (ordo ignot.).—Ins. Maurit.
minutiflora, Span. in Linnaea, xv. (1841) 342 = repens.
mutabilis, R. Br. Prod. 422 = repens.
nantocana, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 403 = viscosa.
nudicaulis, *Phil. in Linnaea*, xxix. (1857-58) 37.—Chili.
obtusifolia, Lam. Illustr. i. 10 = patula.
obtusifolia, Steud. Nom. ed. II. i. 213 = paniculata.
Palmeri, *S. Wats. in Proc. Am. Acad.* xviii. (1882-83) 142.—Texas.
paniculata, Lam. Illustr. i. 10 = hirsuta, erecta.
paniculata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 105.—Texas; Mexic.
patula, *Domb. ex Vahl, Enum.* i. 287.—Peruv.
pedunculosa, *A. Rich. Tent. Fl. Abyss.* ii. 210.—Abyss.
pentandra, Burch. Trav. i. 432 = Burchellii.
periplocaefolia, *Comm. ex Vahl, Enum.* i. 290.—Madag.
periplocifolia, Sieber, ex Choisy, in DC. Prod. xiii. II. 453 = repens.
plicata, *Boj. 2me Rapp. Soc. Hist. Nat. Maurit.* 43 (sp. dub.).—Maurit.?

BOERHAAVIA:—
plumbaginea, *Cav. Ic.* ii. 7. t. 112.—Hispan.; Afr. trop. et austr.; Syria.
polymorpha, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 105 = caribaea, hirsuta, repens.
procumbens, Banks, ex Roxb. Fl. Ind. i. 146 = repens.
pterocarpa, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 376.—Am. bor.
pubescens, R. Br. Prod. i. 422 = repens.
pulchella, *Griseb. in Goett. Abh.* xix. (1874) 88.—Reg. Argent.
pulverulenta, Dupuis, in Vahl, Enum. i. 291 = dichotoma.
purpurascens, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 321.—N. Mexic.
Reboudiana, *Pomel, Nouv. Mat. Fl. Atl.* 335.—Algeria.
repanda, Kotschy, ex Choisy, in DC. Prod. xiii. II. 454 = verticillata.
repanda, Roxb. Hort. Beng. 2; in Beatson, St. Helena Tracts, 302 = verticillata.
repanda, *Willd. Sp. Pl.* i. 22.—As. trop.; Austral.
repens, *Linn. Sp. Pl.* 3.—Cosmop. trop.
repens, Sieber.—Cf. Schweinf. Beitr. Fl. Aethiop. 168 = plumbaginea.
rhomboidea, Humb. in Spreng. Jahrb. Gew. i. III. 66 = Cryptocarpus globosus.
Rottleri, Steud. Nom. ed. II. i. 213 = crispa?
rubicunda, Steud. l. c. = elegans.
rugosa, Rottl. ex Wall. Cat. sub n. 6771 = crispa.
rupestris, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 169 = repens.
sarmentosa, Willd. Sp. Pl. i. 21 = scandens.
scabrida, Steud. & Hochst. ex Steud. Nom. ed. II. i. 213 = discolor.
scandens, Drège, ex Choisy, in DC. Prod. xiii. II. 455 = Burchellii.
scandens, Forsk. Fl. Aegypt. Arab. 3 = plumbaginea.
scandens, Gibs. Cat. Bomb. Pl. 167 = verticillata.
scandens, *Linn. Sp. Pl.* 3.—Am. bor.; Ind. occ.
Schomburgkiana, *Oliver, in Hook. Ic. Pl.* t. 1225.—Austral.
sessiliflora, Steud. Nom. ed. II. i. 213 = repens.
spicata, *Choisy, in DC. Prod.* xiii. II. 456.—Am. bor. occ.
stellata, *Boj. 2me Rapp. Soc. Hist. Nat. Maurit.* 43.—Ins. Maurit.?
stellata, Wight, Ic. t. 875 = verticillata.
suberosa, C. Sm. in Tuckey, Congo, 249 = repens.
sumata, Zipp. ex Span. in Linnaea, xv. (1841) 342 = repens.
surinamensis, Miq. l. c. xviii. (1844) 244 = repens.
tetrandra, *Forst. f. Prod.* 2. n. 5.—Ins. Societ.
tomentosa, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 168 = repens.
tuberosa, *Lam. Illustr.* i. 10.—Peruv.
umbellata, Wight, ex Hook. f. Fl. Brit. Ind. iv. 709 = repanda.
undulata, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 168 = repens.
verticillata, Boiss. Fl. Orient. iv. 1044 = plumbaginea.
verticillata, *Poir. Encyc.* v. 56.—Afr. trop.; Syria; Ind. or.
verticillata, Rottl. in Wall. Cat. n. 6772 = crispa.
virgata, *H. B. & K. Nov. Gen. et Sp.* ii. 215.—N. Mexic.
viscosa, Fresen. in Mus. Senckenb. i. (1834) 75 = repens.
viscosa, *Lag. & Rodr. in Anal. Cienc. Nat.* iv. (1801) 256. n. 12.—Peruv.
viscosa, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 167 = plumbaginea.
vulvariaefolia, Poir. Encyc. v. 55 = repens.
Wrightii, *A. Gray, in Am. Journ. Sc.* Ser. II. xv. (1853) 322.—N. Mexic.

BOGENHARDIA, Reichb. Nom. 200 (1841) = **Abutilon**, Tourn. (Malvac.).
crispa, Reichb. l. c. in nota = A. crispum.

BOHADSCHIA, Crantz, Stirp. Austr. ed. I. fasc. i. 2 (1762); ed. II. 5 (1769) = **Peltaria**, Linn. (Crucif.).
alliacea, Crantz, l. c. = P. alliacea.

BOHADSCHIA, Presl, Rel. Haenk. ii. 98. t. 68 (1831) = **Turnera**, Linn.
humifusa, Presl, l. c. = T. diffusa.
microphylla, Griseb. Cat. Pl. Cub. 114 = T. diffusa.

BOHADSCHIA, F. W. Schmidt, Samml. Phys. Aufs. i. 279 (1795) = **Hyoseris**, Linn. &c. (Compos.).
hirta, F. W. Schmidt, l. c. 280 = Leontodon hirtus.
lucida, F. W. Schmidt, l. c. 279 = H. lucida.

BOHERAVIA, Parodi, in Anal. Soc. Cient. Argent. v. 210 (1878); Contrib. 59 = **Boerhaavia**, Vaill. (Nyctag.).

BOISDUVALIA, Spach, Hist. Vég. Phan. iv. 383 (1835) = **Oenothera**, Linn. (Onagrar.).
andina, Phil. in Linnaea, xxxiii. (1864-65) 70 (= O. andina?).—Chili.
cleistogama, Curran, in Bull. Calif. Acad. (1884) 12.—Calif.
concinna, Spach, Hist. Vég. Phan. iv. 384 = O. concinna.
densiflora, S. Wats. Bot. Calif. i. 233 = O. densiflora.
Douglasii, Spach, Hist. Vég. Phan. iv. 385 = O. densiflora.
glabella, Walp. Rep. ii. 89 = O. glabella.
Tocornalii, C. Gay, Fl. Chil. ii. 344.—Chili.
Torreyi, S. Wats. Bot. Calif. 233 = O. Torreyi.
Volckmanni, Phil. in Linnaea, xxx. (1859-60) 187.—Chili.

BOISSIAEA, Lem. in Orb. Dict. ii. 638 (1842) = **Bossiaea**, Vent. (Legumin.).

BOISSIERA, Domb. ex DC. Syst. i. 512, in syn. (1818) = **Lardizabala**, Ruiz & Pav. (Berberid.).
trifoliata, Domb. ex DC. l. c. 513 = L. trifoliolata.
triternata, Domb. ex DC. l. c. 512 = L. biternata.

BOISSIERA, Haenseler, ex Willk. & Lange, Prod. Fl. Hisp. i. 218 (1861) = **Gagea**, Salisb. (Liliac.).
baetica, Haenseler, l. c. = G. polymorpha.

BOISSIERA, Hochst. & Steud. in Flora, xxi. (1838) I. Intell. 25. *GRAMINEAE*, Benth. & Hook. f. iii. 1175.
SCHNIZLEINIA, Steud. Nom. ed. II. i. 213, in syn. (1840).
WIESTIA, Boiss. Fl. Orient. v. 559 (1884).
bromoides, *Hochst. & Steud. l. c.; Syn. Pl. Gram.* 200.—Arab.; Syria; Persia; Affghan.
Danthoniae, A. Braun, Ind. Sem. Hort. Berol. (1857) 3 = Bromus macrostachyus.
Pumilio, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 9.—Persia.

BOJERIA, DC. Coll. Mém. ix. 18; et Prod. v. 94 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 331.
glabra, *Klatt, in Ann. Sc. Nat.* Sér. V. xviii. (1873) 364.—Afr. trop.
speciosa, *DC. Coll. Mém.* ix. 18. t. 3.—Madag.

BOJERIA, Rafin. Fl. Tellur. iv. 50 (1836) = **Amomum**, Linn. (Scitam.).
magnifica, Rafin. l. c. = A. magnificum.

BOLANDRA, A. Gray, in Proc. Am. Acad. vii. (1868) 341. *SAXIFRAGEAE*.
californica, *A. Gray, l. c.*—Calif.
oregana, *S. Wats. l. c.* xiv. (1879) 292.—Am. bor. occ.

BOLANOSA, A. Gray, Pl. Wright. i. 82 (1852). *COMPOSITAE*, Benth. & Hook. f. ii. 232.
Coulteri, *A. Gray, l. c.*—Mexic.

BOLAX, Comm. ex Juss. Gen. 226 (1789) = **Azorella**, Lam. (Umbell.).
angulatus, Presl, ex DC. Prod. iv. 80 = Mulinum angulatum.
aretioides, Spreng. Umb. Sp. 11 = A. aretioides.
caespitosus, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 62 = A. caespitosa.
caespitosus, Spreng. in Roem. & Schult. Syst. vi. 358 = A. aretioides.
cladorrhizus, Spreng. l. c. 362 = A. cladorrhiza.
complicatus, Spreng. Umb. Sp. 9, 10 = A. caespitosa.
crenatus, Spreng. Syst. i. 880 = A. crenata.
Echinus, Presl, ex DC. Prod. vi. 79 = Mulinum Echinus.

BOLAX :—
filamentosus, Spreng. in Roem. & Schult. Syst. vi. 359 = A. filamentosa.
floccipes, Sieber, ex Benth. Fl. Austral. iii. 380 = Astrotricha floccosa.
geraniifolius, Presl, ex DC. Prod. iv. 81 = Huanaca geraniifolia.
Gilliesii, Hook. Bot. Misc. i. (1830) 325 = A. Gilliesii.
Glebaria, Comm. ex Lam. Encyc. iii. 156 = A. caespitosa.
gummifer, Spreng. Syst. i. 879 = A. caespitosa.
integrifolius, Spreng. ex Steud. Nom. ed. II. i. 214 = A. filamentosa.
ledifolius, Sieber, ex DC. Prod. v. 74 = Astrotricha ledifolia.
lycopodioides, Spreng. in Roem. & Schult. Syst. vi. 361 = A. corymbosa.
multifidus, Spreng. Syst. i. 880 = A. multifida.
pedunculatus, Spreng. Umb. Sp. 10 = A. multifida.
pilifera, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 32.—Peruv.
prolifer, Spreng. in Roem. & Schult. Syst. vi. 361 = Mulinum proliferum.
spinosus, Spreng. l. c. 362 = Mulinum spinosum.
tridentatus, Spreng. Umb. Prod. 34 = Hydrocotyle tridentata.
trifurcatus, Spreng. ex Steud. Nom. ed. II. i. 214 = A. caespitosa.
trilobus, Spreng. Umb. Prod. 34 = Hydrocotyle triloba.

BOLBIDIUM, Lindl. Veg. Kingd. 181 (1847), nomen (Orchid.). In Herb. Lindl. includit species Dendrobii (D. subacaule & prasinum).

BOLBOPHYLLARIA, Reichb. f. in Bot. Zeit. x. (1852) 934 = **Bulbophyllum**, Thou. (Orchid.).
aristata, Reichb. f. Beitr. Orch. Centr. Am. 60 = Bulbophyllum aristatum.
biseta, Reichb. f. in Walp. Ann. vi. 242 = Bulbophyllum bisetum.
bracteolata, Reichb. f. in Bot. Zeit. x. (1852) 934 = Bulbophyllum bracteolatum.
clavata, S. Moore in Baker, Fl. Maurit. 346 = Bulbophyllum clavatum.
Oerstedii, Reichb. f. in Bonplandia, iii. (1855) 223 = Bulbophyllum Oerstedii.
pachyrhachis, Reichb. f. in Walp. Ann. vi. 241 = Bulbophyllum pachyrhachis.
sordida, Reichb. f. l. c. 242 = Bulbophyllum bracteolatum.

BOLBOPHYLLOPSIS, Reichb. f. in Bot. Zeit. x. (1852) 933 = **Cirrhopetalum**, Lindl. (Orchid.).
maculosa, Reichb. f. in Walp. Ann. vi. 243 = C. maculosum.
morphologorum, Reichb. f. in Bot. Zeit. x. (1852) 933 = C. maculosum.

BOLBOPHYLLUM, Spreng. Syst. iii. 681, 732 (1826) = **Bulbophyllum**, Thou. (Orchid.).

BOLBORCHIS, Zoll. & Moritzi, Syst. Verz. Zoll. 89 (1845–46) = **Coelogyne**, Lindl. (Orchid.).
crociformis, Zoll. & Moritzi, l. c. = C. javanica.

BOLBOSTYLIS, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 467 = **Eupatorium**, Tourn. (Compos.).

BOLDEA, Juss. in Ann. Mus. Par. xiv. (1809) 134 = **Peumus**, Molina (Monim.).
fragrans, C. Gay, Fl. Chil. v. 353; et Tul. in Arch. Mus. Par. viii. (1855) 412. t. 31 = P. Boldus.

BOLDOA, Endl. Gen. Suppl. i. 1378 (1841) = BOLDEA, Juss. = **Peumus**, Molina (Monim.).

BOLDOA, Cav. Cat. Hort. Madr. (1803) t. 7; Lag. Gen. et Sp. Nov. 10 (1816). *NYCTAGINEAE*, Benth. & Hook. f. iii. 8.
SALPIANTHUS, Humb. & Bonpl. Pl. Aequin. i. 154. t. 44 (1808).
lanceolata, *Lag. Gen. et Sp. Nov.* 10.—Mexic.
ovatifolia, Lag. l. c. = Cryptocarpus globosus.
paniculata, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 356 = Cryptocarpus globosus.
purpurascens, Cav. Hort. Madr. t. 7 = Cryptocarpus globosus.
repens, *Spreng. Syst.* i. 179.—Bras.

BOLDU, Feuill. ex Adans. Fam. ii. 446 (1763) = **Peumus**, Molina (Monim.).

BOLDU, Necs, Progr. 11 (1833); Syst. Laurin. 177 (1836) = **Cryptocarya**, R. Br. (Laurin.).
chilanum, Necs, Syst. Laurin. 178 = Peumus Boldus.
chilense, Poepp. ex Meissn. in DC. Prod. xv. 1. 67 = praec.
nitidum, Meissn. l. c. 506 = C. nitida.

BOLDUCIA, Neck. Elem. iii. 32 (1790) = **Dipteryx**, Schreb. (Legumin.).

BOLDUS, Schult. f. in Roem. & Schult. Syst. vii. p. x. 56 (1829) = **Peumus**, Molina (Monim.).
chilensis, *Schult. f. l. c.* 56.—P. Boldus.

BOLELIA, Rafin. Atl. Journ. 120 (1832) = **Downingia**, Torr. (Campan.).

BOLEUM, Desv. Journ. Bot. iii. (1814) 163. t. 26. *CRUCIFERAE*, Benth. & Hook. f. i. 98.
asperum, *Desv. l. c.* 163, 175.—Hispan.

BOLINA, Rafin. Sylva Tellur. 101 (1838) (Melastomac.).
conferta, Rafin. l. c. = Brachyotum confertum.
divaricata, Rafin. l. c. = Ptcrogastra divaricata.
excelsa, Rafin. l. c. = Centronia excelsa.

BOLIVARIA, Cham. & Schlecht. in Linnaea, i. (1826). 207. t. 4 = **Menodora**, Humb. & Bonpl. (Oleaceae).
chlorargantha, Remy, in Ann. Sc. Nat. Sér. III. viii. (1847) 225.—Bolivia.
decemfida, Gill. ex Hook. Journ. Bot. i. (1834) 284. —Chili.
Grisebachii, A. Scheele, in Linnaea, xxv. (1852) 254 = M. heterophylla.
integrifolia, Cham. & Schlecht. l. c. i. (1826) 208 = M. integrifolia.
mexicana, A. DC. Prod. viii. 315 = M. mexicana.
pinnatifida, Schlecht. in Linnaea, xxvi. (1853-55) 474 = M. pinnatifida.
robusta, Benth. in Hook. Lond. Journ. Bot. v. (1846) 190. t. 5 = M. robusta.
trifida, Cham. & Schlecht. in Linnaea, i. (1826) 209 = M. trifida.

BOLLAEA, Parl. in Bull. Soc. Bot. Fr. v. (1858) 509 = **Pancratium**, Linn. (Amaryll.).
canariensis, Parl. l. c. = P. canariense.

BOLLEA, Reichb. f. in Bot. Zeit. x. (1852) 667 = **Zygopetalum**, Hook. (Orchid.).
coelestis, Reichb. f. in Gard. Chron. (1876) I. 756 = Z. coeleste.
guianensis, Klotzsch, in Schomb. Fauna & Fl. Guy. 1206, nomen.—Guiana.
Lalindei, Reichb. f. in Gard. Chron. (1874) II. 33 = Z. Lalindei.
Lawrenceana, Reichb. f. l. c. (1878) II. 266 = Z. Lawrenceana.
pallens, Reichb. f. l. c. (1881) I. 462.—Hab.?
Patini, Reichb. f. l. c. (1874) II. 34 = Z. Patini.
pulvinaris, Reichb. f. in Linnaea, xli. (1877) 107 = Z. pulvinare.
violacea, Reichb. f. in Bot. Zeit. x. (1852) 668 = Z. violaceum.

BOLOPHYTA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 347 = **Parthenium**, Linn. (Compos.).
alpina, Nutt. l. c. 348 = P. alpinum.

BOLOSIA, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 211 (1870) = **Hispidella**, Barnad. (Compos.).
piloselloides, Pourr. ex Willk. l. c. = H. hispanica.

BOLTONIA, L'Hérit. Sert. Angl. 27 (1788). *COMPOSITAE*, Benth. & Hook. f. ii. 269.
ASTEROMOEA, Blume, Bijdr. 901 (1827).
DICHAETOPHORA, A. Gray, Pl. Fendl. 73 (1849).
HISUTSUA, DC. Prod. vi. 44 (1837).
MADEA, L'Hérit. ex DC. Prod. v. 301 (1836).
asteroides, Sims, Bot. Mag. t. 2554 = diffusa.
asteroides, *L'Hérit. Sert. Angl.* 27.—Am. bor.
campestris, *Hemsl. Biol. Centr. Am. Bot.* ii. 120.—Texas.
cantoniensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 398.—Japon.
diffusa, *Ell. Sketch*, ii. 400.—Am. bor.

BOLTONIA:—
glastifolia, L'Hérit. Sert. Angl. 27 = asteroides.
incisa, *Benth. Fl. Hongk.* 173 = Aster incisus.
indica, *Benth. l. c.* 174 = Aster indicus.
integrifolia, Benth. & Hook. f. Gen. ii. 269 = pekinensis.
japonica, *Benth. & Hook. f. l. c.*—Japon.
latisquama, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiii. (1862) 238.—Am. bor.
lautureana, Debeaux, in Act. Linn. Soc. Bord. xxxi. (1877) 215 = Aster lautureanus.
pekinensis, *Hance, in Journ. Bot.* v. (1867) 370.—China.
Seemannii, Benth. & Hook. f. Gen. ii. 269 = Achaetogeron Seemannii.

BOLUSIA, Benth. in Hook. Ic. Pl. xii. 57. t. 1163 (1873). *LEGUMINOSAE.*
capensis, *Benth. l. c.*—Afr. austr.

BOMAREA, Mirb. in Hist. Nat. Pl. ix. (1804) 71. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 736.
COLLANIA, Herb. Amaryll. 103. t. 8 (1837).
DANBYA, Salisb. Gen. Pl. Fragm. 57 (1866).
DODECASPERMA, Rafin. Fl. Tellur. iv. 35 (1836).
SPHAERINE, Herb. Amaryll. 106. t. 12, 13, 16, 18 (1837).
VANDESIA, Salisb. in Trans. Hort. Soc. i. (1812) 332.
WICHURAEA, M. Roem. Syn. Ensat. 277 (1847).
acutifolia, *Herb. Amaryll.* 112.—Mexic.
affinis, Kunth, Enum. Pl. v. 796 = edulis.
amoena, *M. Roem. Syn. Ensat.* 274.—Peruv.
anceps, Herb. Amaryll. 116 = fimbriata.
andimarcana, *Baker, in Journ. Bot.* xx. (1882) 201.—Peruv.
Andreana, *Baker, l. c.* 205.—N. Granat.
angulata, *Benth. Pl. Hartw.* 156.—Peruv.
angustifolia, *Benth. l. c.*—Peruv.
angustipetala, *Baker, in Journ. Bot.* xx. (1882) 202.—N. Granat.
aurantiaca, *Herb. Amaryll.* 399. t. 46. f. 1.—Peruv.
brachysepala, *Benth. Pl. Hartw.* 157.—Peruv.
bracteata, Herb. Amaryll. 112. t. 3. f. 3 = edulis.
Brauniana, Schenk, in Mart. Fl. Bras. iii. 1. 168 = edulis.
Bredemeyerana, Herb. Amaryll. 118 = multiflora.
brevis, *Baker, in Journ. Bot.* xx. (1882) 202.—Peruv.
Caldasiana, *Herb. Amaryll.* 119.—Guatemala; Peruv.
caraccensis, Herb. l. c. 111. t. 16. f. 2 = edulis.
Carderi, *Mast. in Gard. Chron.* (1876) I. 793, 795. —N. Granat.
chontalensis, Seem. in Gard. Chron. (1871) 479 = edulis.
coccinea, *Baker, in Journ. Bot.* xx. (1882) 202.—Peruv.
conferta, Benth. Pl. Hartw. 259 = patacocensis.
cordifolia, *Herb. Amaryll.* 113.—Peruv.
cornigera, *Herb. l. c.* 116. t. 17. f. 2.—Peruv.
cornuta, Herb. l. c. 114. t. 17. f. 4 = edulis.
crinita, *Herb. l. c.* 119. t. 15. f. 4.—Peruv.
crocea, *Herb. l. c.* 119.—Peruv.
cumbrensis, Herb. l. c. 115. t. 18. f. 2 = torta.
declinata, Klotzsch, ex Kunth, Enum. Pl. v. 802 dispar.
densiflora, Herb. Amaryll. 399. t. 46. f. 4 = tomentosa.
diffracta, *Baker, in Journ. Bot.* xx. (1882) 206.—N. Granat.
denticulata, *Herb. Amaryll.* 118.—Peruv.
dispar, *Herb. l. c.* 115. t. 17. f. 1.—Peruv.
dissitifolia, *Baker, in Journ. Bot.* xx. 1882) 203.—Ecuador.
distichophylla, *Baker, l. c.* 202.—Peruv.
edulis, *Herb. Amaryll.* 111.—Am. trop.
Fanningiana, Herb. l. c. 116. t. 16. f. 3 = multiflora.
fimbriata, *Herb. l. c.* 116.—Peruv.
floribunda, Herb. l. c. = fimbriata.
formosissima, *Herb. l. c.* 111. t. 14. f. 4.—Peruv.
frondea, *Mast. in Gard. Chron.* (1882) I. 668.—N. Granat.
furcata, Klotzsch, ex Kunth, Enum. Pl. v. 792 = edulis.
fuscata, *Klotzsch, in Schomb. Fauna & Fl. Guy.* 1120, nomen.—Guiana.
glaucescens, *Baker, in Journ. Bot.* xx. (1882) 201. Ecuador.
glomerata, *Herb. Amaryll.* 115. t. 15. f. 1.—Peruv.
goniocaulon, *Baker, in Journ. Bot.* xx. (1882) 204.—Ecuador.

BOMAREA:—
gloriosa, M. Roem. Syn. Ensat. 269 = edulis.
granatensis, M. Roem. l. c. 264 = B. Salsilla.
grandifolia, Herb. Amaryll. 113 = edulis.
Halliana, *Herb. Amaryll.* 117. t. 10. f. 3.—N. Granat.
Hartwegii, *Baker, in Journ. Bot.* xx. (1882) 203.—Ecuador.
hirsuta, *Herb. Amaryll.* 114.—N. Granat.
hirta, *Schenk, in Mart. Fl. Bras.* iii. 1. 169.—Bras.
hirtella, Herb. Amaryll. 112 = edulis.
hispida, *Baker, in Journ. Bot.* xx. (1882) 202.—N. Granat.
Hookeriana, *Herb. Amaryll.* 398. t. 46. f. 3.—Peruv.
involucrosa, *Baker, in Journ. Bot.* xx. (1882) 201.—Peruv.
Jacquesiana, *Kunth, Enum. Pl.* v. 800.—Bras.
janeirensis, M. Roem. Syn. Ensat. 267 = edulis.
Kalbreyeri, *Baker, in Journ. Bot.* xx. (1882) 204.—N. Granat.
lancifolia, *Baker, l. c.* 202.—Ecuador.
latifolia, Herb. Amaryll. 113 = edulis.
Lehmanni, *Baker, in Journ. Bot.* xxi. (1883) 373.—Am. austr.
linifolia, *Baker, l. c.* xx. (1882) 201.—N. Granat.
longipes, *Baker, l. c.* 204.—Ecuador.
lutea, *Herb. Amaryll.* 120. t. 5.—Peruv.
lyncina, *Herb. l. c.* 398. t. 46. f. 8.—Peruv.
Maakiana, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xx. (1852) 337.—Venezuela.
Macleanica, Herb. in Lindl. Bot. Reg. (1842) Misc. 66 = aurantiaca.
macrocarpa, Herb. Amaryll. 14 = edulis.
maranensis, *Herb. l. c.* 17.—Bras.
Martiana, Schenk, in Mart. Fl. Bras. iii. 1. 170 = edulis.
miniata, Kunth, Enum. Pl. v. 792 = edulis.
minima, *Baker, in Journ. Bot.* xx. (1882) 202.—N. Granat.
Moritziana, Klotzsch, ex Kunth, Enum. Pl. v. 797 = edulis.
multiflora, *Mirb. Hist. Nat. Pl.* ix. (1804) 72.—N. Granat.
multipes, *Benth. Pl. Hartw.* 156.—Peruv.
nervosa, *Baker, in Journ. Bot.* xx. (1882) 202.—Peruv.
nobilis, Herb. Amaryll. 422 = ovata.
obovata, Herb. l. c. 112. t. 14. f. 2 = edulis.
oculata, M. Roem. Syn. Ensat. 263 = B. Salsilla.
oligantha, *Baker, in Gard. Chron.* (1877) ii. 648.—Peruv.
ovata, *Mirb. Hist. Nat. Pl.* ix. (1804) 72.—Am. austr.
pachyphlebia, *Baker, in Journ. Bot.* xx. (1882) 204.—Ecuador.
pardina, *Herb. Amaryll.* 120. t. 14. f. 1.—N. Granat.
paltarumensis, Herb. l. c. 422 = fimbriata.
patacocensis, *Herb. l. c.* 120. t. 14. f. 3.—Ecuador; N. Granat.
pauciflora, Herb. l. c. 112 = edulis.
platypetala, Benth. Pl. Hartw. 156 = uncifolia.
podopetala, *Baker, in Journ. Bot.* xx. (1882) 202.—Ecuador.
polygonatoides, *Baker, l. c.*—Ecuador.
praecipua, Herb. Amaryll. 422 = B. Salsilla.
pudibunda, *Planch. & Linden, Cat.* n. 10. (1855) 4.—N. Granat.
pumila, *Griseb. in Lechl. Berb. Am. Austr.* 56, nomen.—Peruv.
punctata, *Herb. in Lindl. Bot. Reg.* (1842) *Misc.* 66.—Peruv.
purpurea, *Herb. Amaryll.* 118, 399.—Peruv.
rosea, *Herb. l. c.* 118.—Peruv.
Salsilla, *Mirb. in Hist. Nat. Pl.* ix. (1804) 71.—Chili.
Salsilla, Vell. Fl. Flum. iii. t. 20 = edulis.
salsilloides, M. Roem. Syn. Ensat. 264 = edulis.
secundifolia, *Baker, in Journ. Bot.* xx. (1882) 202.—Peruv.
setacea, Herb. Amaryll. 117 = tomentosa.
Shuttleworthii, *Mast. in Gard. Chron.* (1882) i. 76.—N. Granat.
simplex, Benth. Pl. Hartw. 156 = Hartwegii.
simplex, Herb. Amaryll. t. 15; et in Bot. Mag. t. 3863 = rosea.
spectabilis, Schenk, in Mart. Fl. Bras. iii. 1. 169 = edulis.
subfalcata, Herb. Amaryll. 422; Pritz. Ic. Ind. 158 = B. Salsilla.
subglobosa, Herb. in Lindl. Bot. Reg. (1842) Misc. 66 = fimbriata.

BOMAREA:—
superba, *Herb. Amaryll.* 117. t. 6. f. 1.—Peruv.
Tatiana, Herb. l. c. 423 = ovata.
tomentosa, *Herb. l. c.* 117.—Peruv.
torta, *Herb. l. c.* 115.—Ecuador.
Turneriana, Herb. in Lindl. Bot. Reg. (1842) Misc. 66 = formosissima.
uncifolia, *Herb. l. c.*—Peruv. vel Bolivia.
variabilis, Herb. l. c. = rosea.
villosa, M. Roem. Syn. Ensat. 268 = ovata.
vitellina, *Mast. in Gard. Chron.* (1882) i. 143.—N. Granat.
Williamsiae, *Mast. l. c.* (1882) ii. 553.—N. Granat.

BOMARIA, Kunth, Enum. Pl. v. 787 (1850) = **Bomarea**, Mirb. (Amaryll.).

BOMBAX, Linn. Sp. Pl. ed. I. 511 (1753). *MALVACEAE*, Benth. & Hook. f. i. 210.
ERIOTHECA, Schott & Endl. Meletem. 35 (1832).
SALMALIA, Schott & Endl. l. c. (1852).
aculeatum, *Vell. Fl. Flum.* 289; vii. t. 56.—Bras.
aesculifolium, H. B. & K. Nov. Gen. et Sp. v. 298 = Eriodendron aesculifolium.
axillare, Moç. & Sessé, ex DC. Prod. i. 479 = Eriodendron aesculifolium.
Barrigon, *Decne. in Fl. des Serres*, Sér. II. xiii. (1880) 44.—Panama.
buonopozense, *Beauv. Fl. Owar.* ii. 42. t. 83.—Afr. trop.
carolinoides, Donn, Cat. Hort. Cant. ed. IV. 156 = Pachira minor.
carolinum, *Vell. Fl. Flum.* 289; vii. t. 57.—Bras.
Ceiba, Burm. f. Fl. Ind. 145 = malabaricum.
Ceiba, *Linn. Sp. Pl.* 511.—Am. austr.
Congo, Burm. f. Fl. Ind. 145 = Cochlospermum Gossypium.
coriaceum, *Mart. Nov. Gen. et Sp.* i. 93, in obs.—Bras.
cumanense, *H. B. & K. Nov. Gen. et Sp.* v. 300.—Venezuela.
decaphyllum, Decne. in Fl. des Serres, Sér. II. xiii. (1880) 44 = endecaphyllum.
discolor, *H. B. & K. Nov. Gen. et Sp.* v. 299.—N. Granat.
ellipticum, *H. B. & K. l. c.*—Mexic.
emarginatum, *Decne. in Fl. des Serres*, Sér. II. xiii. (1880) 44.—Bras.
endecaphyllum, *Vell. Fl. Flum.* 288; vii. t. 50.—Bras.
Erianthus, Cav. Diss. v. 294 = Eriodendron leiantherum.
Fendleri, *Benth. in Journ. Linn. Soc.* vi. (1862) 108.—Panama.
festivum, Wall. Cat. n. 1841 = insigne.
glabrescens, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 211.—Bras.
globiferum, Raeusch. Nom. ed. III. 195 = globosum.
globosum, *Aubl. Pl. Gui.* ii. 701. t. 281.—Guiana.
Gossypium, Linn. Syst. ed. XII. 517 = Cochlospermum Gossypium.
grandiflorum, Cav. Diss. v. 295 = insigne.
grandiflorum, Sonner. Voy. Ind. ii. 235. t. 133 = Cochlospermum Gossypium.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 302 = Eriodendron anfractuosum.
heptaphyllum, Cav. Diss. v. 296; Roxb. Pl. Corom. iii. 43. t. 247 = malabaricum.
heptaphyllum, Linn. Sp. Pl. 960 = septenatum.
hexaphyllum, *Vell. Fl. Flum.* 288; vii. t. 52.—Bras.
hibiscifolium, Willd. ex Steud. Nom. ed. II. i. 215 = Cochlospermum hibiscoides.
Hilarianum, *Decne. in Fl. des Serres*, Sér. II. xiii. (1880) 44.—Bras.
humile, *Benth. in Journ. Linn. Soc.* vi. (1862) 108.—Bras.
insigne, *Wall. Pl. As. Rar.* i. 74. t. 79 et 80.—Ind. or.
jasminodorum, Baill. Hist. Pl. iv. 119 = Eriodendron jasminodorum.
malabaricum, *DC. Prod.* i. 479.—Ind. or.; Malaya; China.
mediterraneum, *Arruda, in Koster, Trav. Braz.* 489.—Bras.
mexicanum, *Hemsl. Diag. Pl. Nov.* 4.—Mexic.
mompoxense, *H. B. & K. Nov. Gen. et Sp.* v. 300.—Venezuela.
Munguba, *Mart. Nov. Gen. et Sp.* i. 93.—Bras.
occidentale, Spreng. Syst. iii. 124 = mompoxense; Eriodendron caribaeum.
octophyllum, *Vell. Fl. Flum.* 288; vii. t. 51.—Bras.
orientale, Spreng. Syst. iii. 124 = Eriodendron anfractuosum.

BOMBAX:—
orinoccense, H. B. & K. Nov. Gen. et Sp. v. 300 = Cochlospermum orinoccense.
parviflorum, *Mart. Nov. Gen. et Sp.* i. 91. t. 57.—Bras.
pentandrum, Jacq. Stirp. Am. 191 = Eriodendron caribaeum.
pentandrum, Linn. Sp. Pl. 511 = Eriodendron anfractuosum.
pentaphyllum, *Vell. Fl. Flum.* 289; vii. t. 55.—Bras.
plumosum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 8 = Eriodendron anfractuosum?
pubescens, *Mart. Nov. Gen. et Sp.* i. 91. t. 58.—Bras.
pyramidale, Cav. Diss. v. 294. t. 153 = Ochroma lagopus.
pyramidatum, Steud. Nom. ed. I. 114 = praec.
quinatum, Jacq. Stirp. Am. 192 = B. Ceiba.
religiosum, Linn. Sp. Pl. 512 = Cochlospermum Gossypium.
retusum, *Mart. Nov. Gen. et Sp.* i. 92. t. 59.—Bras.
rubrinervis, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 69.—Bras.
Samauma, Spreng. Syst. iv. Cur. Post. 259 = Eriodendron Samauma.
septenatum, *Jacq. Enum. Pl. Carib.* 26.—N. Granat.
serrata, *Desf. Cat. Hort. Par.* ed. III. 407.—Hab.?
sexdigitatum, *Vell. Fl. Flum.* 289; vii. t. 54.—Bras.
tomentosum, *A. Juss. in A. St. Hil. Fl. Bras. Mer.* i. 263.—Bras.
ventricosum, Arruda, in Koster, Trav. Braz. 489 = Chorisia crispiflora.
villosum, *Mill. Gard. Dict.* ed. VIII. n. 3.—Am. austr.
vitifolium, Willd. Enum. Hort. Berol. 720 = Cochlospermum vitifolium.

BOMBIX, Medic. Malv. 44 (1787) = **Hibiscus**, Linn. (Malvac.).
phoenicea, Medic. l. c. = H. phoeniceus.

BOMBYCELLA, Lindl. Veg. Kingd. 370 (1847) = **Hibiscus**, Linn. (Malvac.).
betulina, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 241 (= H. betulinus?).—Porto Rico.
bicolor, Welw. Apont. 589.—Afr. trop.
phoenicea, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 241 = H. phoeniceus.

BOMBYCIDENDRON, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 14 = **Hibiscus**, Linn. (Malvac.).
grewiaefolium, Zoll. & Mor. l. c. = H. grewiaefolius.

BOMBYCOSPERMUM, Presl, Rel. Haenk. ii. 137. t. 71 (1831) = **Ipomoea**, Linn. (Convolv.).
mexicanum, Presl, l. c. = I. bombycina.

BOMBYNIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1; ed. II. 64 (1827) (Quid?).

BOMBYX, Moench, Meth. 616 (1794) = **Hibiscus**, Linn. (Malvac.).
phoenicea, Moench, l. c. = H. phoeniceus.

BONA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 360; et Phil. Bot. i. 205 (1789) = **Vicia**, Linn. (Legumin.).
narbonensis, Medic. l. c. 360 = V. narbonensis.
speciosa, Medic. l. c. 361 = V. narbonensis.

BONAFIDIA, Neck. Elem. iii. 46 (1790) = **Amorpha**, Linn. (Legumin.).

BONAFOUSIA, A. DC. Prod. viii. 359 (1844) = **Tabernaemontana**, Linn. (Apocyn.).
attenuata, Miers, Apocyn. S. Am. 51.—Am. austr.
guyanensis, Miers, l. c. = T. guyanensis.
latiflora, Miers, l. c. 50 = T. flavicans.
obliqua, Miers, l. c. 49.—Am. austr.
oblongifolia, Miers, l. c. 50 = T. oblongifolia.
olivacea, Miers, l. c. 52 = T. olivacea.
Perrottetii, Miers, l. c. 51 = T. Perrottetii.
polyneura, Miers, l. c. 53.—Am. austr.
rariflora, Miers, l. c.—Am. austr.
rupicola, Miers, l. c. 52 = T. rupicola.
undulata, A. DC. Prod. viii. 359 = T. undulata.

BONAGA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 345; Phil. Bot. 206 (1789) = **Ononis**, Linn. (Legumin.).
antiquorum, Medic. l. c. 346 = O. antiquorum.
arvensis, Medic. Phil. Bot. 206 = O. arvensis.
Columnae, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 346 = O. Columnae.
hircina, Medic. l. c. = O. hircina.
minutissima, Medic. l. c. = O. minutissima.
mitis, Medic. l. c. = O. arvensis.
spinosa, Medic. l. c. = O. spinosa.

BONAMIA, A. Gray, in Proc. Am. Acad. v. (1862) 337 = **Breweria**, R. Br. (Convolv.).
aquatica, A. Gray, Man. Bot. N. U. St. ed. V. 376 = Brewcria aquatica.
humistrata, A. Gray, in Proc. Am. Acad. v. (1862) 337 = Brewcria humistrata.
Menziesii, A. Gray, l. c.—Ins. Sandvic.
Pickeringii, A. Gray, Man. Bot. N. U. St. ed. V. 376 = Brewcria Pickeringii.

BONAMIA, Thou. in Dict. Sc. Nat. v. 145 (1804); et Hist. Veg. Isl. Afr. 17. t. 8 (1805). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 877.
madagascariensis, *Poir. Encyc. Suppl.* i. 677.—Madag.

BONAMICA, Vell. Fl. Flum. 21 (1825); i. t. 50 = **Linociera**, Sw. (Olcac.).
filiformis, Vell. l. c.—Bras.

BONANIA, A. Rich. Fl. Cub. Fanerog. ii. 201. t. 68 (1853). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 335.
adenodon, *Benth. & Hook. f. Gen.* iii. 335.—Cuba.
cubana, *A. Rich. Fl. Cub. Fanerog.* ii. 201.—Cuba.
cubensis, A. Rich. l. c. t. 68 = cubana.
emarginata, *Wright, ex Griseb. in Goett. Nachr.* (1865) 78.—Cuba.
erythrosperma, *Benth. & Hook. f. Gen.* iii. 335.—Cuba.
myricifolia, *Benth. & Hook. f. l. c.*—Cuba.

BONANNIA, Guss. Fl. Sic. Syn. i. 355 (1842). *UMBELLIFERAE*, Benth. & Hook. f. i. 910.
resinifera, *Guss. l. c.*—Sicilia.
resinosa, Strobl, in Oestr. Bot. Zeitschr. xxxiv. (1884) 174 = resinifera.

BONANNIA, Presl, Fl. Sic. i. 99 (1826) = **Brassica**, Linn. (Crucif.).
dissecta, Presl, l. c. 100 = Brassica dissecta.
officinalis, Presl, l. c. 99 = Brassica alba.

BONANNIA, Rafin. Specchio, xv. 115 (1814) = **Blighia**, Kon. (Sapind.).
nitida, Rafin. l. c. 116 = Blighia sapida.

BONANOX, Rafin. Fl. Tellur. 77 (1836) = **Ipomoea**, Linn. (Convolv.).
indica, Rafin. l. c.—Ind. or.
muricata, Rafin. l. c. = I. Bona-nox.
orbiculata, Rafin. l. c. = I. orbiculata.
riparia, Rafin. l. c. = I. riparia.

BONAPARTEA, Haw. Syn. Pl. Succ. 68 (1812) = **Agave**, Linn. (Amaryll.).
filamentosa, Boucen. in Rev. Hortic. (1868) 65 = A. filifera.
flagelliformis, Henck. in Flora, iii. (1820) 45 = A. geminiflora.
gracilis, Sweet, Hort. Brit. ed. II. 597 = Dasylirion acrotichum.
Hystrix, Hort. ex Pasq. Cat. Orto Bot. Nap. 111 = A. striata.
juncea, Haw. Syn. Pl. Succ. 68 = A. geminiflora.
stricta, Hort. ex Vukot. in Rad. Jugos. Akad. Zagrab, xxvii. (1874) 213 = A. striata.

BONAPARTEA, Ruiz & Pav. Fl. Per. iii. 38 (1802) = **Tillandsia**, Linn. (Bromel.).
juncea, Ruiz & Pav. l. c. = T. setacea.
strobilantha, Ruiz & Pav. l. c. 39 = T. strobilantha.
vittata, Mart. ex Schult. f. Syst. vii. ii. 1198 = T. vittata.

BONAROTA, Adans. Fam. ii. 209 (1763) = **Paederota**, Linn. (Scrophul.).
chamaedrifolia, Scop. Fl. Carn. ed. II. i. 22 = P. Agcria.
rotundifolia, Caruel, in Parl. Fl. Ital. vi. 532 = P. Bonarota.

BONATEA, Willd. Sp. Pl. iv. 43 (1805) = **Habenaria**, Willd. (Orchid.).
Bilabrella, Lindl. Gen. et Sp. Orch. 328 = H. tetrapetala.
Boltoni, Harv. Thes. Cap. i. 55. t. 88 = H. Boltoni.
cassidea, Sond. in Linnaea, xix. (1847) 81 = H. cassidea.
cirrhata, Lindl. Gen. et Sp. Orch. 327 = H. cirrhata.
clavata, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 208 = H. clavata.
Darwinii, Weale, in Journ. Linn. Soc. x. (1869) 470 = H. cassidea.
densiflora, Sond. in Linnaea, xix. (1847) 80 = H. densiflora.
flexuosa, Lindl. Gen. et Sp. Orch. 328 = H. flexuosa.
foliosa, Lindl. l. c. 329 = H. foliosa.
gracilis, Lindl. l. c. = H. gracilis.
herbacea, Wall. ex Lindl. l. c. 307 = H. digitata.
incarnata, Lindl. l. c. 327 = Disa incarnata.
macilenta, Lindl. ex Benth. in Hook. Lond. Journ. Bot. ii. (1843) 673 = H. macilenta.
micrantha, Lindl. Gen. et Sp. Orch. 329 = H. arenaria.
panduana, Lindl. in Wall. Cat. n. 7063 = H. digitata.
pauciflora, Lindl. Gen. et Sp. Orch. 329 = H. pauciflora.
pratensis, Lindl. l. c. 328 = H. pratensis.
speciosa, Willd. Sp. Pl. iv. 43 = H. Bonatea.
tetrapetala, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 208 = H. tetrapetala.

BONAVERIA, Scop. Introd. 310 (1777) = **Securigera**, DC. (1805) (Legum.).
Securidaca, Desv. Journ. Bot. i. (1813) 120 = S. Coronilla.
Securigera, Endl. ex Heynh. Nom. ii. 73 = S. Coronilla.

BONDUC, Adans. Fam. ii. 318 (1763) = **Caesalpinia**, Linn. (Legum.).
canadense, Medic. Theod. 113 = C. Bonduc.
majus, Medic. l. c. 43 = C. Bonduc.
minus, Medic. l. c. 41 = C. Bonducella.

BONELLIA, Bert. ex Colla, Hort. Ripul. 21 (1826) = **Jacquinia**, Jacq. (Myrsin.).
Cavanillesii, Bert. l. c. = J. macrocarpa.

BONGARDIA, C. A. Mey. Verz. Pfl. Cauc. 174 (1831). *BERBERIDEAE*, Benth. & Hook. f. i. 43.
chrysogonum, Boiss. Fl. Orient. i. 99 = Rauwolfii.
Olivieri, C. A. Mey. Verz. Pfl. Cauc. 175 = Rauwolfii.
Rauwolfii, *C. A. Mey. l. c.* 174.—Graecia; Oriens; Afghan.

BONIFACIA, Manso, ex Steud. Nom. ed. II. i. 216 (1840) = **Augusta**, Pohl (Rubiac.).
riparia, Manso, ex Steud. l. c. = A. lanceolata.

BONJEANEA, Reichb. Fl. Germ. Excurs. 507 (1832) = **Dorycnium**, Linn. (Legum.).
cinerascens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 12 = D. hirsutum.
graeca, Griseb. Spicil. Fl. Rumel. i. 43 = D. latifolium.
hirsuta, Reichb. Ic. Fl. Germ. x. 32 = D. hirsutum.
hirta, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 11 = D. hirsutum.
italica, Jord. & Fourr. l. c. 12 = D. hirsutum.
Kotschyi, Boiss. & Reut. ex Boiss. Fl. Orient. ii. 161 = D. Kotschyi.
microphylla, Reichb. Fl. Germ. Excurs. 507, in nota = Indigofera gracilis.
prostrata, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 11 [emend. fasc. ii. 133] = D. hirsutum.
recta, Reichb. Fl. Germ. Excurs. 507 = D. rectum.
sericea, Jord. & Fourr. Ic. Fl. Eur. i. 24. t. 61 = D. hirsutum.
syriaca, Boiss. Diagn. Scr. I. ix. 31 = D. hirsutum.
venusta, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 12 = D. hirsutum.

BONNAYA, Link & Otto, Ic. Pl. Sel. 25. t. 11 (1820). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 956.
alata, *Spreng. Syst.* i. 41.—Ins. Molucc.

BONNAYA:—
alterniflora, *Wright, in Sauv. Fl. Cub.* 101.—Cuba.
bifolia, Spreng. Syst. iv. Cur. Post. 15 = Chirita bifolia.
brachiata, *Link & Otto, Ic. Pl. Sel.* ii. 25. t. 11.—Ind. or.; Malaya.
brachycarpa, Cham. & Schlecht. in Linnaea, ii. (1827) 568 = Ilysanthes capensis.
bracteata, Griff. Notul. iv. 107 = veronicaefolia.
ciliata, Spreng. Syst. i. 41 = reptans.
clausa, *F. Muell. ex Benth. Fl. Austral.* iv. 499.—Austral.
cordifolia, Spreng. Syst. i. 42 = Vandellia pedunculata.
cyanea, Griff. Notul. iv. 107 = veronicaefolia.
grandiflora, Spreng. Syst. i. 41 = veronicaefolia.
hispidula, G. Don, Gen. Syst. iv. 538 (Gen. dubium).—Ind. or.
humilis, Spreng. Syst. iv. Cur. Post. 15 = Chirita pumila.
hyssopioides, Benth. in Wall. Cat. n. 3866 = Ilysanthes hyssopioides.
integrifolia, Kostel. Allg. Med.-Pharm. Fl. iii. 893 = Vandellia erecta.
marginata, Spreng. Syst. i. 41 = veronicaefolia.
minima, G. Don, Gen. Syst. iv. 538 = oppositifolia.
minima, Wight, Ic. iii. t. 858 = Ilysanthes minima.
ocymifolia, G. Don, Gen. Syst. iv. 538 (= Vandelliae sp.).—Ind. or.
oppositifolia, *Spreng. Syst.* i. 41.—Ind. or.
origanifolia, Spreng. l. c. 42 = Vandellia diffusa.
parviflora, Benth. in Wall. Cat. n. 3867 = Ilysanthes parviflora.
pedunculatis, *Benth. in Wall. l. c.* n. 3865; *et Scroph. Ind.* 34.—Ind. or.
penicillata, *Nees, in DC. Prod.* xi. 161.—Japon.
personata, Hassk. in Flora, xlvii. (1864) 56 = brachiata.
procumbens, Benth. in Wall. Cat. n. 3859 = veronicaefolia.
pulegiifolia, Spreng. Syst. i. 41 = veronicaefolia.
pumila, Spreng. Syst. iv. Cur. Post. 15 = Chirita pumila.
punctata, G. Don, Gen. Syst. iv. 538 = Limnophila conferta.
pusilla, Benth. in Wall. Cat. n. 3861; Scroph. Ind. 33 = brachiata.
pusilla, Griff. Notul. iv. 107 = tenuifolia.
pusilla, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 121.—Afr. trop.
repens, G. Don, Gen. Syst. iv. 537 = seq.
reptans, *Spreng. Syst.* i. 41.—Ind. or.; Malaya; Ins. Maurit.
rigida, Benth. in Wall. Cat. n. 3860 = veronicaefolia.
Rothii, A. Dietr. Sp. Pl. i. 556 = veronicaefolia.
rotundifolia, Benth. Scroph. Ind. 34 = Ilysanthes rotundifolia.
Roxburghiana, A. Dietr. Sp. Pl. i. 559 = veronicaefolia.
ruellioides, Spreng. Syst. i. 41 = reptans.
serrata, A. Dietr. Sp. Pl. i. 556; Griff. Notul. iv. 108 = brachiata.
tenuifolia, *Spreng. Syst.* i. 42.—Ind. or.; China.
trichotoma, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 121.—Afr. trop.
Vahlii, G. Don, Gen. Syst. iv. 538 = Vandellia scabra.
verbenaefolia, Spreng. Syst. i. 42 = veronicaefolia.
veronicaefolia, *Spreng. l. c.* 41.—Ind. or.; Malaya; China.
Wightii, Benth. in Wall. Cat. n. 3864 = veronicaefolia.

BONNETIA, Mart. & Zucc. Nov. Gen. et Sp. i. 114. t. 100 (1824). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 187.
KIESERIA, Nees, in Neuwied, Reise nach Brasil. ii. 338 (1821).
anceps, *Mart. & Zucc. Nov. Gen. et Sp.* i. 115.—Bras.
bahiensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 246.—Bras.
coriacea, Spreng. Syst. iv. Cur. Post. 207 = Kielmeyera coriacea.
corymbosa, Spreng. l. c. = Kielmeyera corymbosa.
meridionalis, *Sw. Fl. Ind. Occ.* ii. 967.—Am. trop.
palustris, Vahl, Eclog. Am. ii. 42 = meridionalis.
paniculata, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1861) 63.—Peruv.
parviflora, *Spruce, ex Benth. l. c.*—Peruv.
petiolaris, Spreng. Syst. iv. Cur. Post. 207 = Kielmeyera petiolaris.
rosea, Spreng. l. c. = Kielmeyera rosea.

BONNETIA :—
sessilis, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 363.—Guiana.
stricta, *Mart. in Nov. Act. Nat. Cur.* xii. (1825) 87.—Bras.
trifoliata, Auct. ex Walp. Rep. i. 499 = Galipea officinalis.
variabilis, Spreng. Syst. iv. Cur. Post. 207 = Kielmeyera variabilis.
venulosa, *Mart. & Zucc. Nov. Gen. et Sp.* i. 115.—Bras.

BONNETIA, Neck. Elem. i. 368 (1790) = **Buchnera**, Linn. (Scrophular.).

BONNETIA, Schreb. Gen. 363 (1789) = **Mahurea**, Aubl. (Ternstroem.).
palustris, J. F. Gmel. Syst. 814 = M. palustris.
racemosa, Sw. Fl. Ind. Occ. ii. 965 = M. speciosa.

BONPLANDIA, Cav. in Anal. Hist. Nat. ii. (1800) 131. t. 20. *POLEMONIACEAE*, Benth. & Hook. f. ii. 824.
CALDASIA, Willd. Hort. Berol. t. 71 (1807).
geminiflora, *Cav. l. c.* 132.—Mexic.

BONPLANDIA, Willd. in Mém. Acad. Berl. (1802) 24 = **Galipea**, Aubl. (Rutac.).
Angostura, Rich. in Mém. Math.-Phys. Inst. Par. (1811) 82. t. 10 = G. Cusparia.
Candolleana, Spreng. Syst. i. 781 = G. Candolleana.
cuneifolia, Spreng. l. c. = G. cuneifolia.
Erythrochiton, Spreng. l. c. 780 = G. pentandra.
Fontanesiana, Spreng. l. c. = G. Fontanesiana.
heterophylla, Spreng. l. c. = G. heterophylla.
Lasiostemon, Spreng. l. c. = G. Lasiostemon.
pentagyna, Spreng. l. c. = G. pentagyna.
resinosa, Spreng. l. c. = G. resinosa.
trifoliata, Willd. in Mém. Acad. Berl. (1802) 35 = G. Cusparia.

BONTIA, Linn. in Loefl. Iter, 193 (1758) = **Avicennia**, Linn. (Verbenac.).
germinans, Linn. Sp. Pl. ed. II. 891 = A. officinalis.

BONTIA, Linn. Syst. ed. I (1735). *MYOPORINEAE*, Benth. & Hook. f. ii. 1126.
daphnifolia, Salisb. Prod. 109 = seq.
daphnoides, *Linn. Sp. Pl.* 638.—Ind. occ.
minor, Gaertn. f. Fruct. iii. 169. t. 212. f. 3 = daphnoides.

BONYUNIA, Schomb. Reise, iii. 1082, nomen; Prog. in Mart. Fl. Bras. vi. 1. 267. t. 72 (1868). *LOGANIACEAE*, Benth. & Hook. f. ii. 796.
antoniaefolia, *Prog. in Mart. Fl. Bras.* vi. 1. 288.—Bras.
superba, *Schomb. Reise*, iii. 1082.—Guiana.

BOOPHONE, Herb. App. 18 (1821) = **Buphane**, Herb. (Amaryll.).

BOOPIS, Juss. in Ann. Mus. Par. ii. (1803) 350. t. 58. *CALYCEREAE*, Benth. & Hook. f. ii. 161.
ACARPHA, Griseb. in Goett. Abh. vi. (1853-55) 125.
CHIONOPHILA, Miers, ex Lindl. Veg. Kingd. 701 (1847).
GAMOCARPHA, DC. Prod. v. 2 (1836).
NASTANTHUS, Miers, ex Lindl. Veg. Kingd. 701 (1847), nomen; et in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 185.
acaulis, *Phil. in Anal. Univ. Chil.* (1861) 49.—Chili.
alpina, *Poepp. ex Less. in Linnaea*, vi. (1831) 258.—Chili.
anthemoides, *Juss. in Ann. Mus. Par.* ii. (1803) 350.—Reg. Argent.
araucana, *Phil. in Anal. Univ. Chil.* (1872) 736.—Chili.
australis, *Decne. Voy. Pôle Sud Bot.* ii. 87. t. 18, B.—Reg. Magell.
balsamitaefolia, Juss. in Ann. Mus. Par. ii. (1803) 350 = Calycera balsamitaefolia.
bellidifolia, *Phil. in Anal. Univ. Chil.* (1872) 736.—Chili.
bupleuroides, *C. A. Muell. in Mart. Fl. Bras.* vi. IV. 355.—Bras.

BOOPIS :—
caespitosa, *Phil. Fl. Atac.* 26.—Chili.
compacta, *Phil. in Linnaea*, xxviii. (1856) 709.—Chili.
crassifolia, *A. Gray, in Proc. Am. Acad.* v. (1861) 321.—Patagon.
Diazi, *Phil. in Linnaea*, xxxiii. (1864-65) 106.—Chili.
Gayana, *Phil. l. c.* xxviii. (1856) 709.—Chili.
gracilis, Phil. l. c. 707 = leucanthema.
graminea, *Phil. l. c.*—Chili.
laciniata, *Ball, in Journ. Linn. Soc.* xxi. (1884) 220.—Patagon.
leucanthema, *Poepp. ex Less. in Linnaea*, vi. (1831) 259.—Chili.
multicaulis, *Phil. in Linnaea*, xxviii. (1856) 706.—Chili.
pozoaeformis, *Phil. in Anal. Univ. Chil.* (1872) 736.—Chili.
pusilla, *Phil. in Linnaea*, xxxiii. (1864-65) 105.—Chili.
rigidula, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 284.—Patag.
sanjuanina, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 31.—Reg. Argent.
scapigera, *Remy, in C. Gay, Fl. Chil.* iii. 250.—Chili.
spathulata, *Phil. in Linnaea*, xxviii. (1856) 708.—Chili.
squarrosa, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 286 = australis.
tribuloides, Spreng. Syst. iii. 674 = Acicarpha tribuloides.

BOOTHIA, Dougl. ex Benth. in Trans. Hort. Soc. Ser. II. i. (1835) 405 = **Platystemon**, Benth. (Papaverac.).
californica, Dougl. ex Loud. Encyc. Pl. Suppl. i. 1214 = P. californicus.

BOOTIA, Adans. Fam. ii. 320 (1763) = **Borbonia**, Linn. (Legumin.).

BOOTIA, Bigel. Fl. Boston. ed. II. 206 (1824) = **Potentilla**, Linn. (Rosac.).
sylvestris, Bigel. l. c. = P. arguta.

BOOTIA, Neck. Delic. Gallo-Belg. i. 193 (1768) = **Saponaria**, Linn. (Caryophyll.).
glutinosa, Neck. ex Reichb. Ic. Fl. Germ. v. 41 = S. glutinosa.
ocymoides, Neck. ex Reichb. l. c. = S. ocymoides.
Saponaria, Neck. in Act. Acad. Theod. Pal. ii. (1770) 484 = S. officinalis.
vulgaris, Neck. Delic. Gallo-Belg. i. 193 = S. officinalis.

BOOTROPHIS, Steud. Nom. ed. II. i. 22 (1840) = BOTROPHIS, Rafin. = **Cimicifuga**, Linn. (Ranunc.).

BOOTTIA, Ayres, ex Baker, Fl. Maurit. 49 (1877) = **Pleurostylia**, Wight & Arn. (Celastrac.).
obovata, Ayres, ex Baker, l. c. = P. leucocarpa.

BOOTTIA, Wall. Pl. As. Rar. i. 51. t. 65 (1829). *HYDROCHARIDEAE*, Benth. & Hook. f. iii. 453.
cordata, *Wall. l. c.*—Burma.

BOPUSIA, Presl, Bot. Bemerk. 91 (1844) = **Graderia**, Benth. (1846) (Scrophul.).
scabra, Presl, l. c. = G. scabra.

BOQUILA, Decne. in Comptes Rend. (1837) 394; et in Arch. Mus. Par. i. (1839) 188. t. 11 B. *BERBERIDEAE*, Benth. & Hook. f. i. 42.
discolor, *Decne. in Ann. Sc. Nat.* Sér. II. xii. (1839) 103.—Chili.
trifoliata, *Decne. l. c.*—Chili.

BORABORA, Steud. Syn. Pl. Cyp. 71 (1855) = **Mariscus**, Gaertn. (Cyperac.).
cyperoidea, Steud. l. c. = M. macrophyllus.

BORAEVA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 382 = **Boreava**, Jaub. & Spach (Crucif.).

BORAGO, Linn. Sp. Pl. i. 137 (1753). *BORAGINEAE*, Benth. & Hook. f. ii. 854.
BORRAGO, Tourn. ex Linn. Syst. ed. I (1735).
BUGLOSSITES, Moris, in Enum. Sem. Hort. Taur. (1845).—Cf. Ann. Sc. Nat. Sér. III. v. (1846) 365.
advena, Gilib. Fl. Lithuan. Suppl. 272 = officinalis.
africana, Linn. Sp. Pl. 138 = Trichodesma africanum.
arabica, Ehrenb. ex Boiss. Fl. Orient. iv. 281 = Trichodesma Ehrenbergii.
aspera, Gilib. Fl. Lithuan. i. 25 = officinalis.
constantinopolitana, *Hill, Veg. Syst.* vii. 45.—Oriens.
cordifolia, Moench, Meth. Suppl. 275 = orientalis.
crassifolia, Vent. Jard. Cels, 100. t. 100 = Caccinia glauca.
cretica, *Willd. Sp. Pl.* i. 778.—Graecia.
gruina, *C. Sm. in Tuckey, Congo*, 250, nomen.—Ins. Cap. Verd.
indica, Blanco, Fl. Filip. ed. I. 81 = Trichodesma zeylanicum.
indica, Linn. Sp. Pl. 137 = Trichodesma indicum.
inflata, Salisb. Prod. 116 = orientalis.
laxiflora, *Willd. Enum. Hort. Berol. Suppl.* 9.—Corsica.
longifolia, *Poir. Voy. Barb.* ii. 119.—Afr. bor.
macranthera, [Soland. in] Russell, Aleppo, ed. II. ii. 246 = Caccinia Russellii.
officinalis, *Linn. Sp. Pl.* 137.—Europ.; Afr. bor.; As. Min.
orientalis, *Linn. l. c.* 138.—As. Min.; Reg. Caucas.
spinulosa, Roxb. Hort. Beng. [13]; Fl. Ind. i. 459 = Trichodesma indicum.
verrucosa, Forsk. Fl. Aegypt. Arab. 41 = Trichodesma africanum.
zeylanica, Burm. f. Fl. Ind. 41 = Trichodesma zeylanicum.

BORASSUS, Linn. Mus. Cliff. 13 (1736). *PALMAE*, Benth. & Hook. f. iii. 939.
LONTARUS, Adans. Fam. ii. 25 (1763).
aethiopium, Mart. Hist. Nat. Palm. iii. 221 = flabellifer.
caudatum, Lour. Fl. Cochinch. 760 = Wallichia caudata.
dichotomus, *White, in J. Grah. Cat. Bomb. Pl.* (1839) 226.—Ind. or.
flabellifer, *Linn. Sp. Pl.* 1187.—Ind. or.
flabelliformis, Murr. Syst. ed. XIII. 827 = flabellifer.
Gomutus, Lour. Fl. Cochinch. ii. 618 = Arenga saccharifera.
Ihur, Giseke, Linn. Prael. 87 = Pholidocarpus Ihur.
madagascariensis. *Boj. Hort. Maurit.* 308.—Madag.
pinnatifrons, Jacq. Hort. Schoenb. ii. 65 = Chamaedorea gracilis.
secundiflorus, *Hort. ex H. Wendl. in Kerch. Palm.* 235, nomen.—Hab.?
Sonnerati, Giseke, Linn. Prael. 86 = Lodoicea sechellarum.
tunicata, Lour. Fl. Cochinch. 618 = Pholidocarpus tunicatus.

BORBONIA, Linn. Coroll. Gen. 12. n. 971 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 475.
BOOTIA, Adans. Fam. ii. 320 (1763).
alata, Willd. Hort. Berol. i. 48 = parviflora.
alpestris, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 461.—Afr. austr.
angustifolia, *Lam. Encyc.* i. 436.—Afr. austr.
axillaris, Lam. l. c. 438 = Amphithalea densa.
barbata, *Lam. l. c.* 436.—Afr. austr.
Candolleana, *Eckl. & Zeyh. Enum.* 163.—Afr. austr.
capitata, Poir. Encyc. Suppl. i. 680 = Priestleya capitata.
ciliata, DC. Prod. ii. 120 = Candolleana.
ciliata, *Willd. Sp. Pl.* iii. 923.—Afr. austr.
complicata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 462.—Afr. austr.
cordata, Andr. Bot. Rep. t. 31 = Rafnia triflora.
cordata, *Linn. Sp. Pl.* 707.—Afr. austr.
cordifolia, Lam. Encyc. i. 436 = cordata.
crenata, *Linn. Sp. Pl.* 708.—Afr. austr.
crenata, Willd. ex Steud. Nom. ed. II. i. 217 = parviflora.
decipiens, E. Mey. Comm. Pl. Afr. Austr. i. 15 = lanceolata.
ericifolia, Linn. Pl. Rar. Afr. 14; Amoen. Acad. vi. 92 = Amphithalea ericaefolia.
graminifolia, Lam. Encyc. i. 437 = Priestleya graminifolia.
hirsuta, *Poir. Encyc. Suppl.* i. 680 = Priestleya hirsuta.
laevigata, Linn. Mant. 100 = Priestleya Thunbergii.

BORBONIA:—
lanceolata, *Linn. Sp. Pl.* 707.—Afr. austr.
latifolia, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 462.—Afr. austr.
monosperma, *DC. Prod.* ii. 120.—Afr. austr.
myrtifolia, Poir. Encyc. Suppl. i. 680 = Priestleya myrtifolia.
obcordata, Spreng. Syst. iv. Cur. Post. 267 = Tephrosia obcordata.
ovata, Andr. Bot. Rep. t. 31 = Rafnia triflora.
parviflora, *Lam. Encyc.* i. 437.—Afr. austr.
perfoliata, Lam. l. c. = ciliata.
perfoliata, Thunb. Prod. Pl. Cap. 122 = Rafnia perfoliata.
perforata, *Eckl. & Zeyh. Enum.* 163.—Afr. austr.
perforata, Thunb. Prod. Pl. Cap. 122 = ciliata.
pungens, Mundt, ex Benth. in Hook. Lond. Journ. Bot. ii. (1843) 461 = trinervia.
ruscifolia, Sims, Bot. Mag. t. 2128 = parviflora.
sericea, Lam. Encyc. i. 438 = Priestleya sericea.
serrulata, Thunb. Diss. Borbonia, 6 = parviflora.
sphaerica, Lam. Encyc. i. 437 = Liparia sphaerica.
sphaerosperma, Spreng. Syst. iv. Cur. Post. 267 = Requienia sphaerosperma.
tecta, Poir. Encyc. Suppl. i. 680 = Priestleya tecta.
teres, Poir. l. c. = Priestleya teres.
tomentosa, Linn. Sp. Pl. 707 = Priestleya villosa.
triflora, Auct. ex Steud. Nom. ed. II. i. 217 = Rafnia triflora.
trinervia, E. Mey. Comm. Pl. Afr. Austr. i. 15 = lanceolata.
trinervia, Linn. Sp. Pl. 707 = Cliffortia ruscifolia.
trinervia, *Thunb. Prod. Pl. Cap.* 122.—Afr. austr.
umbellifera, *Poir, Encyc. Suppl.* i. 680.—Afr. austr.
undulata, *Thunb. Prod. Pl. Cap.* 122.—Afr. austr.
vestita, Poir. Encyc. Suppl. i. 680 = Priestleya vestita.
villosa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 28.—Afr. austr.
villosa, Thunb. Fl. Cap. 560 = Priestleya angustifolia.

BORBONIA, Plum. ex Adans. Fam. ii. 341 (1763) (Laurin.).
carolinensis, J. S. Presl, Rostl. ii. 57 = Persea carolinensis.
cupularis, Gaertn. f. Fruct. iii. 225. t. 222 (= *Mespilodaphne cupularis*).—Maurit.
exaltata, J. S. Presl, Rostl. ii. 57 = Nectandra exaltata.
foetens, J. S. Presl, l. c. = Ocotea foetens.
globosa, Gaertn. f. Fruct. iii. 224. t. 222 = Nectandra sanguinea.
indica, J. S. Presl, Rostl. ii. 57 = Persea indica.
martinicensis, J. S. Presl, l. c. (= *Oreodaphne Jacquiniana*).—Ind. occ.
membranacea, J. S. Presl, l. c. = Nectandra membranacea.
peruviana, Juss. ex Steud. Nom. ed. II. i. 217 (= *Mespilodaphne pretiosa*).—Peruv.

BORBOYA, Rafin. Fl. Tellur. iii. 52 (1836) = **Hyacinthus**, Tourn. (Liliac.).
spicata, Rafin. l. c. = H. spicatus.

BORCKHAUSENIA, Gaertn. Mey. & Scherb. Fl. Wett. iii. 4 (1801) = **Corydalis**, Vent. (1803) (Papav.).
capnoides, Fuss, Fl. Transs. 42 = C. capnoides.
cava, Gaertn. Mey. & Scherb. Fl. Wett. iii. 17 = C. bulbosa.
lutea, Gaertn. Mey. & Scherb. l. c. 19 = C. lutea.
ochroleuca, Fuss, Fl. Transs. 42 = C. ochroleuca.
solida, Gaertn. Mey. & Scherb. Fl. Wett. iii. 18 = C. Halleri.

BORCKHAUSENIA, Roth, Catalecta, fasc. ii. 56 (1800) = **Teedia**, Rudolphi (Scrophul.).
lucida, Roth, l. c. = T. lucida.

BORDEREA, Miégev. in Bull. Soc. Bot. Fr. xiii. (1866) 374 = **Dioscorea**, Linn.
pyrenaica, Miégev. l. c. = D. pyrenaica.

BOREA, Zipp. ex Mackl. in Bijdr. Nat. Wet. v. (1830) 142, nomen (absque descr.).

BOREA, Meissn. Gen. Comm. 221 (1836-42), sphalm. = BOVEA, Decne. = **Lindenbergia**, Lehm. (Scrophul.).

BOREAVA, Jaub. & Spach, Illustr. i. 3. t. 2 (1841).
CRUCIFERAE, Benth. & Hook. f. i. 95.
MARTINSIA, Godr. in Mém. Acad. Stanislas, Sér. III. [xvii.] 1853 (1854) 378.
aptera, *Boiss. & Heldr. Diagn.* Ser. I. viii. 49.—As. Min.
orientalis, *Jaub. & Spach, Illustr.* i. 3.—As. Min.; Syria.

BORELLIA, Neck. Elem. i. 275, 276 (1790) = **Cordia**, Linn. (Boragin.).
aspera, Rafin. Sylva Tellur. 41 = C. aspera.

BORETTA, Neck. Elem. i. 212 (1790) = **Daboëcia**, D. Don (Ericac.).

BORISSA, Rafin. ex Steud. Nom. ed. II. i. 217 (1840) = **Lysimachia**, Linn. (Primulac.).
Linum-stellatum, Rafin. ex Steud. l. c. = L. Linum-stellatum.

BORITH, Adans. Fam. ii. 262 (1763) = **Anabasis**, Linn. (Chenopod.).

BORKHAUSIA, Nutt. Gen. Am. ii. 125 (1818) = BARKHAUSIA, Moench = **Crepis**, Linn. (Compos.).

BORONELLA, Baill. Adansonia, x. (1872) 302.
RUTACEAE.
Pancheri, *Baill. l. c.* 303.—N. Caled.

BORONIA, Sm. Tracts, 288 (1798); et in Trans. Linn. Soc. viii. (1807) 285. t. 5-7. *RUTACEAE*, Benth. & Hook. f. i. 291.
CYANOTHAMNUS, Lindl. Swan River Bot. 18 (1839).
affinis, *R. Br. ex Benth. Fl. Austral.* i. 311.—Austral.
alata, *Sm. in Trans. Linn. Soc.* viii. (1807) 283.—Austral.
albiflora, *R. Br. ex Benth. Fl. Austral.* i. 317.—Austral.
algida, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 100.—Austral.
alulata, *Soland. ex Benth. Fl. Austral.* i. 313.—Austral.
anemonifolia, *A. Cunn. in Field, New South Wales*, 330.—Austral.
anemonifolia, Paxt. Mag. Bot. ix. 123 = Fraseri.
anethifolia, A. Cunn. ex Endl. in Enum. Pl. Hueg. 16 = anemonifolia.
arborescens, F. Muell. Fragm. i. 100 = Zieria Smithii.
artemisiaefolia, *F. Muell. l. c.* 66.—Austral.
artemisioides, F. Muell. in Hook. Kew Journ. ix. (1857) 196 = praec.
baeckeacea, *F. Muell. Fragm.* iv. 28.—Austral.
Barkeriana, *F. Muell. l. c.* xi. 96.—Austral.
bicolor, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 163 = tetrandra.
bipinnata, Lindl. in Mitch. Journ. Trop. Austr. 225 = anemonifolia.
Bowmani, *F. Muell. Fragm.* iv. 135.—Austral.
brachyphylla, F. Muell. l. c. i. 99 = oxyantha.
Busselliana, *F. Muell. l. c.* ix. 113.—Austral.
caerulescens, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 11.—Austral.
calophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 160.—Austral.
Candollii, G. Don, Gen. Syst. i. 793 = alata.
capitata, *Benth. Fl. Austral.* i. 323.—Austral.
chironiifolia, Bartl. in Lehm. Pl. Preiss. i. 167 = denticulata.
citriodora, Gunn, ex Hook. f. Fl. Tasm. i. 68 = pinnata.
clavellifolia, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 12 = inornata.
colorata, Lehm. Pl. Preiss. ii. 226 = viminea.
crassifolia, *Bartl. in Lehm. l. c.* i. 169.—Austral.
crassipes, *Bartl. l. c.* 168.—Austral.
crenulata, *Sm. in Trans. Linn. Soc.* viii. (1807) 284.—Austral.
cymosa, *Endl. in Enum. Pl. Hueg.* 16.—Austral.
defoliata, *F. Muell. Fragm.* ix. 113.—Austral.
denticulata, *Sm. in Trans. Linn. Soc.* viii. (1807) 284.—Austral.
dentigera, F. Muell. in Trans. Inst. Vict. i. (1855) 82 = anemonifolia.
dichotoma, Lindl. Bot. Reg. (1841) sub t. 47 = spathulata.

BORONIA:—
Drummondii, Planch. in Fl. des Serres, Sér. I. ix. (1853-54) 65. t. 881 = pulchella.
Edwardsii, *Benth. Fl. Austral.* i. 312.—Austral.
clatior, *Bartl. in Lehm. Pl. Preiss.* i. 170.—Austral.
criantha, *Lindl. in Mitch. Journ. Trop. Austr.* 298.—Austral.
cricifolia, *Benth. Fl. Austral.* i. 313.—Austral.
falcifolia, *A. Cunn. in Lindl. Bot. Reg.* (1841) sub t. 47.—Austral.
fasciculifolia, F. Muell. Fragm. i. 99 = thymifolia.
fastigiata, *Bartl. in Lehm. Pl. Preiss.* i. 168.—Austral.
filicifolia, *A. Cunn. ex Benth. Fl. Austral.* i. 311.—Austral.
filifolia, *F. Muell. Fragm.* i. 3.—Austral.
flexuosa, Bartl. in Lehm. Pl. Preiss. i. 166 = spathulata.
floribunda, Sieber, ex Spreng. Syst. iv. Cur. Post. 148 = pinnata.
Fraseri, *Hook. Bot. Mag.* t. 4052.—Austral.
gracilipes, *F. Muell. Fragm.* ii. 99.—Austral.
grandisepala, *F. Muell. l. c.* i. 66.—Austral.
granulata, F. Muell. l. c. 101 = Zieria granulata.
Gunnii, Hook. f. Fl. Tasm. i. 68. t. 10 = pinnata.
heterophylla, *F. Muell. Fragm.* ii. 98.—Austral.
hirsuta, F. Muell. l. c. i. 101 = Zieria pilosa.
humilis, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 160 = crassifolia.
hypericifolia, Regel, Gartenfl. (1857) 152 = fastigiata.
hyssopifolia, Sieber, in Spreng. Syst. iv. Cur. Post. 148 = polygalifolia.
inconspicua, *Benth. Fl. Austral.* i. 313.—Austral.
inornata, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 164.—Austral.
juncea, *Bartl. in Lehm. Pl. Preiss.* i. 166.—Austral.
laevigata, F. Muell. Fragm. i. 101 = Zieria laevigata.
lanceolata, *F. Muell. l. c.* 66.—Austral.
laniflora, Bartl. in Lehm. Pl. Preiss. i. 165 = juncea.
lanuginosa, *Endl. in Enum. Pl. Hueg.* 16.—Austral.
ledifolia, *J. Gay, in Mém. Mus. Par.* vii. (1821) 450, in nota.—Austral.
ledophylla, F. Muell. Fragm. i. 67 = ledifolia.
lepidota, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 596.—Austral.
leptophylla, Turcz. l. c. xxv. (1852) II. 164 = inornata.
Machardiana, *F. Muell. Fragm.* ix. 115.—Austral.
macra, Bartl. in Lehm. Pl. Preiss. i. 167 = spathulata.
megastigma, *Nees, ex Bartl. in Lehm. Pl. Preiss.* ii. 227.—Austral.
microphylla, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 148.—Austral.
minutiflora, F. Muell. Fragm. i. 100 = Zieria obcordata.
mollis, *A. Cunn. in Lindl. Bot. Reg.* (1841) sub t. 47.—Austral.
multicaulis, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 161 = crassifolia?
nana, Hook. Ic. Pl. t. 270 = polygalifolia.
nematophylla, *F. Muell. Fragm.* ii. 100.—Austral.
ovata, *Lindl. Bot. Reg.* (1841) sub t. 47.—Austral.
oxyantha, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 165.—Austral.
paleifolia, A. Cunn. in Enum. Pl. Hueg. 16 = falcifolia.
paradoxa, DC. Prod. i. 722 = ledifolia.
parviflora, *Sm. Tracts*, 295. t. 6.—Austral.
penicillata, *Benth. Fl. Austral.* i. 322.—Austral.
pilonema, Labill. Nov. Holl. Pl. i. 98. t. 126 = parviflora.
pilosa, *Labill. l. c.* 97. t. 124.—Austral.
pinnata, *Sm. Tracts*, 290.—Austral.
platyrrhachis, *F. Muell. Fragm.* vii. 37.—Austral.
polygalifolia, *Sm. Tracts*, 297. t. 7.—Austral.
psoraleoides, DC. Prod. i. 721 = elatior.
pteropoda, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 595.—Austral.
pubescens, Bartl. in Lehm. Pl. Preiss. ii. 227 = lanuginosa?
pulchella, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 162.—Austral.
ramosa, *Benth. Fl. Austral.* i. 320.—Austral.
rhomboidea, *Hook. Ic. Pl.* t. 722.—Austral.
Rinzii, *Hueg. ex Heynh. Nom.* ii. 73, nomen.—Austral.
rosmarinifolia, A. Cunn. ex Endl. in Enum. Pl. Hueg. 16 = ledifolia.
rubiginosa, A. Cunn. ex Endl. l. c. = ledifolia.
salicifolia, *A. Cunn. ex Heynh. Nom.* ii. 73, nomen.—Austral.
scabra, *Lindl. Swan River App.* 76.—Austral.
semifertilis, F. Muell. Fragm. ii. 98 = elatior.
serrulata, *Sm. Tracts*, 292. t. 5.—Austral.

BORONIA:—
spathulata, *Lindl. Swan River App.* 73.—Austral.
spinescens, *Benth. Fl. Austral.* i. 319.—Austral.
stricta, Bartl. in Lehm. Pl. Preiss. i. 169 = lanuginosa.
subcaerulea, F. Muell. Fragm. ii. 100 = ramosa.
subsessilis, *Benth. Fl. Austral.* i. 322.—Austral.
tenuifolia, Bartl. in Lehm. Pl. Preiss. i. 168 = viminea.
tenuis, *Benth. Fl. Austral.* i. 220.—Austral.
teretifolia, Lindl. Swan River App. 75 = cymosa.
ternata, *Endl. Nov. Stirp. Dec.* 6.—Austral.
tetrandra, Hook. Journ. Bot. ii. (1840) 419 = pinnata.
tetrandra, *Labill. Nov. Holl. Pl.* i. 98. t. 125.—Austral.
tetrandra, Lindl. & Paxt. Flow. Gard. i. (1850–51) 35. t. 8 = pulchella.
tetrathecoides, DC. Prod. i. 723 = polygalifolia.
thymifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 165.—Austral.
triphylla, Sieber, ex Spreng. Syst. iv. Cur. Post. 148 = ledifolia.
tristis, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 162 = megastigma.
variabilis, Hook. Journ. Bot. i. (1834) 255 = anemonifolia.
veronicea, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 11 = Zieria veronicea.
viminea, *Lindl. Swan River App.* 74.—Austral.

BORRAGINOIDES, Moench, Meth. 515 (1794) = **Trichodesma**, R. Br. (Boragin.).
aculeata, Moench, l. c. 516 = T. africanum.
sagittata, Moench, l. c. = T. indicum.

BORRAGO, Tourn. ex Linn. Syst. ed. I (1735) = **Borago**, Linn.

BORRERA, Spreng. Gen. i. 93 (1830) = BORRERIA, G. F. W. Mey. = **Spermacoce**, Linn. (Rubiac.).

BORRERIA, G. F. W. Mey. Prim. Fl. Esseq. 79. t. 1 (1818) = **Spermacoce**, Linn. (Rubiac.).
adscendens, Cham. & Schlecht. in Linnaea, iii. (1828) 340 = S. adscendens.
advena, Fisch. & Mey. Ind. Sem. Hort. Petrop. vi. 46 = Diodia verticillata?
affinis, DC. Prod. iv. 546 = S. diffusa.
alata, DC. l. c. 544 = S. alata.
alternans, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 283.—Porto Rico.
angustifolia, Cham. & Schlecht. in Linnaea, iii. (1828) 330.—Bras.
anthospermoides, DC. Prod. iv. 550.—Am. austr.
aralioides, Cham. & Schlecht. in Linnaea, iii. (1828) 339 = *umbellata*?
aretioides, Griseb. Cat. Pl. Cub. 286.—Cuba.
argentea, Cham. in Linnaea, ix. (1834) 215.—Bras.
arida, DC. Prod. iv. 549.—Porto-Rico.
asclepiadea, Cham. & Schlecht. in Linnaea, iii. (1828) 320 = S. Poaya.
aspera, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 130 = S. aspera.
asperula, DC. Prod. iv. 543 = Diodia rigida?
assurgens, Griseb. in Goett. Abh. xxiv. (1879) 156 = Diodia dasycephala.
aturensis, Cham. & Schlecht. in Linnaea, iii. (1828) 345 = S. aturensis.
Bartlingiana, DC. Prod. iv. 544 = S. latifolia.
basalis, Anderss. Om Galap.-Oarnes Vegetat. 191.—Ins. Galapag.
brachystemonoides, Cham. & Schlecht. in Linnaea, iii. (1828) 314.—Bras.
capitata, DC. Prod. iv. 545 = S. capitata.
capitellata, Cham. & Schlecht. in Linnaea, iii. (1828) 319 = S. laevis.
centranthoides, Cham. & Schlecht. l. c. 327.—Bras.
commutata, Spreng. Syst. i. 404 = S. verticillata.
coriacea, Scheele, in Linnaea, xvii. (1843) 339 = S. tenella.
cornifolia, DC. Prod. iv. 545 = S. capitata.
corymbosa, DC. l. c. 550 = S. corymbosa.
crassifolia, C. A. Mey. ex Heynh. Nom. ii. 74, nomen.—Am. austr.
cupularis, DC. Prod. iv. 543 = S. pusilla.
cymosa, Cham. & Schlecht. in Linnaea, iii. (1828) 323 = S. cymosa.
densiflora, DC. Prod. iv. 542 = S. longifolia, *L'Hérit.*
dichotoma, Cham. & Schlecht. in Linnaea, iii. (1828) 340 = Diodia glabra.
diffusa, DC. Prod. iv. 545 = S. diffusa.

BORRERIA:—
discolor, Benth. in Hook. Kew Journ. iv. (1852) 199 = Spermacoce semierecta.
discolor, DC. Prod. iv. 545.—Ins. Philipp.
dispersa, Hook. f. in Trans. Linn. Soc. xx. (1859) 217.—Ins. Galapag.
distans, Cham. & Schlecht. in Linnaea, iii. (1828) 340 = S. distans.
divaricata, Hook. f. in Trans. Linn. Soc. xx. (1859) 219.—Ins. Galapag.
domingensis, Griseb. Cat. Pl. Cub. 141 = S. ocymoides.
dubia, Cham. in Linnaea, ix. (1834) 216 = *valerianoides*.
elongata, DC. Prod. iv. 547 = S. capitata.
equisetoides, Cham. & Schlecht. in Linnaea, iii. (1828) 338.—Bras.
ericaefolia, Hook. f. in Trans. Linn. Soc. xx. (1859) 218.—Ins. Galapag.
ericoides, Cham. & Schlecht. in Linnaea, iii. (1828) 326 = S. corymbosa.
eritrichoides, Griseb. Cat. Pl. Cub. 142.—Cuba.
eryngioides, Cham. & Schlecht. in Linnaea, iii. (1828) 316 = S. diffusa.
eupatorioides, Cham. & Schlecht. l. c. 327.—Bras.
falcifolia, Hook. f. in Trans. Linn. Soc. xx. (1859) 219.—Ins. Galapag.
ferruginea, DC. Prod. iv. 547 = S. capitata.
ferruginea, Mart. & Gal. in Bull. Acad. Brux. Sér. I. xi. I. (1844) 128 = Mitracarpum villosum.
flavescens, DC. Prod. iv. 548 = S. Poaya.
Fockeana, Miq. in Linnaea, xviii. (1844) 299 = S. latifolia.
globularioides, Cham. & Schlecht. in Linnaea, iii. (1828) 312 = S. verticillata.
gracilis, Miq. ex Hook. f. Fl. Brit. Ind. iii. 200 = S. stricta.
gracilis, Scheele, in Linnaea, xvii. (1843) 340 = S. tenella.
gracillima, DC. Prod. iv. 543 = S. gracillima.
graminifolia, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 129 = S. verticillata.
gymnocephala, DC. Prod. iv. 549.—Bras.
Haenkeana, DC. l. c. 547 = S. tenella.
hebecarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 347 = S. stricta.
humifusa, Mart. in Flora, xxiv. (1841) Beibl. II. 68.—Bras.
Kleinii, Walp. Rep. ii. 464 = S. stricta.
kohautiana, Cham. & Schlecht. in Linnaea, iii. (1828) 311 = S. verticillata.
laevigata, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 130 = S. laevigata.
laevis, Griseb. in Goett. Abh. vii. (1857) 231 = S. laevis.
lasiocarpa, Walp. Rep. ii. 464 = S. stricta.
laxa, Cham. & Schlecht. in Linnaea, iii. (1828) 337.—Bras.
linearifolia, Hook. f. in Trans. Linn. Soc. xx. (1859) 217.—Ins. Galapag.
linifolia, DC. Prod. iv. 551 = S. laevis.
linoides, DC. l. c. 548 = S. tenella.
longiseta, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 128 = S. longiseta.
lutescens, DC. Prod. iv. 547 = S. tenella.
micrantha, Torr. & Gray, Fl. N. Am. ii. 28 = S. ocymoides.
microphylla, DC. Prod. iv. 546 = S. ocymoides.
minima, DC. l. c. 542 = S. verticillata.
Neesiana, DC. l. c. 542 = S. Neesiana.
neglecta, A. Rich. Tent. Fl. Abyss. i. 177.—Abyssin.
nervosa, DC. Prod. iv. 548 = S. Poaya.
oaxacana, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 130 = S. verticillata.
ocymoides, DC. Prod. iv. 544 = S. ocymoides.
organensis, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 111.—Bras.
ovalifolia, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 129 = S. ocymoides.
ovalis, Anderss. Om Galap.-Oarnes Vegetat. 192.—Ins. Galapag.
parviflora, G. F. W. Mey. Prim. Fl. Esseq. 83. t. 1 = S. ocymoides.
parvifolia, Hook. f. in Trans. Linn. Soc. xx. (1859) 218.—Ins. Galapag.
patula, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 128 = S. parviflora.
penicillata, Miq. in Verh. Nat. Wet. Haarl. vii. (1851) 176 = S. latifolia.

BORRERIA:—
perpusilla, Hook. f. in Trans. Linn. Soc. xx. (1859) 218.—Ins. Galapag.
Perrottetii, DC. Prod. iv. 548 = Spermacoce latifolia.
platyphylla, DC. l. c. 549 = S. Poaya.
Poaya, DC. l. c. = S. Poaya.
podocephala, DC. l. c. 542 = S. verticillata.
Pohliana, DC. l. c. 550 = *eupatorioides*.
pratensis, DC. l. c. 548 = S. Poaya.
procera, Cham. in Linnaea, ix. (1834) 593 = S. tenella.
prostrata, Miq. in Verh. Nat. Wet. Haarl. vii. (1851) 177 = S. ocymoides.
psyllioides, DC. Prod. iv. 548 = S. tenella.
pterophora, Presl, Bot. Bemerk. 86.—Bras.
pumila, DC. Prod. iv. 543 = S. pumila.
pusilla, DC. l. c. = S. stricta.
radiata, DC. l. c. 542 = S. radiata.
radicans, Griseb. Cat. Pl. Cub. 142 = Nodocarpaea radicans.
ramisparsa, DC. Prod. iv. 544 = S. ocymoides.
remotifolia, DC. l. c. = Mitracarpum villosum.
repens, DC. l. c. 542 = S. ocymoides.
Roxburghiana, Walp. Rep. ii. 264 = S. stricta.
rubrostipulata, DC. Prod. iv. 546 = S. tenella.
saponariaefolia, Cham. & Schlecht. in Linnaea, iii. (1828) 325.—Bras.
saturejaefolia, Cham. & Schlecht. l. c. 313 = S. tenella.
scabiosoides, Cham. & Schlecht. l. c. 318.—Bras.
scabrida, DC. Prod. iv. 548 = S. latifolia.
scandens, DC. l. c. 546 = S. scandens.
setosa, Mart. & Gal. in Bull. Acad. Brux. xi. I. (1844) 131 = S. setosa.
Sideritis, Cham. & Schlecht. in Linnaea, iii. (1828) 201 = S. latifolia.
simplex, Griseb. Fl. Brit. W. Ind. 350.—Jamaica.
sphaerica, DC. Prod. iv. 547 = S. laevis.
spinosa, Cham. & Schlecht. in Linnaea, iii. (1828) 340 = S. spinosa.
squamosa, Griseb. Cat. Pl. Cub. 142.—Cuba.
stricta, G. F. W. Mey. Prim. Fl. Esseq. 83. t. 1. f. 1–3 = S. verticillata.
strictissima, DC. Prod. iv. 547 = S. tenella.
strumpfioides, Griseb. Cat. Pl. Cub. 142.—Cuba.
suaveolens, G. F. W. Mey. Prim. Fl. Esseq. 81. t. 1 = S. tenella.
suberecta, Hook. f. in Trans. Linn. Soc. xx. (1859) 217.—Ins. Galapag.
subulata, DC. Prod. iv. 543 = Crusea subulata.
tampicana, DC. l. c. 544, pro parte = S. laevis.
tenella, Cham. & Schlecht. in Linnaea, iii. (1828) 317 = S. tenella.
tenera, DC. Prod. iv. 543 = S. tenera.
tenuis, DC. l. c. = S. tenuis.
tenuissima, Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr. iii. 234 = S. tenuissima.
tetracocca, Thw. Enum. Pl. Zeyl. 442 = Fergusonia zeylanica.
tetraptera, Miq. in Linnaea, xvii. (1843) 70 = S. latifolia.
thymifolia, Griseb. Fl. Brit. W. Ind. 350.—Ind. occ.
thymocephala, Griseb. Cat. Pl. Cub. 142 = S. verticillata.
trichantha, Miq. in Verh. Nat. Wet. Haarl. vii. (1851) 178.—Guiana.
umbellata, Spreng. Neue Entdeck. ii. 144.—Bras.
vaginata, Cham. & Schlecht. in Linnaea, iii. (1828) 340 = S. laevis.
valerianoides, Cham. & Schlecht. l. c. 335.—Bras.
verbenoides, Cham. & Schlecht. l. c. 331; iv. (1829) 550.—Bras.
verticillata, G. F. W. Mey. Prim. Fl. Esseq. 83 = S. verticillata.
virgata, Cham. & Schlecht. in Linnaea, iii. (1828) 324 = S. virgata.
Wydleriana, DC. Prod. iv. 545.—Porto Rico.

BORRICHIA, Adans. Fam. ii. 130 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 368.
ADELMANNIA, Reichb. Consp. 110 (1828).
DIOMEDEA, Cass. in Bull. Soc. Philom. (1815) 175.
HELICTA, Cass. l. c. (1818) 167.
TRIMETRA, Moç. ex DC. Prod. vii. 262 (1838).
arborescens, *DC. Prod.* v. 489.—Am. trop.
argentea, *DC. l. c.*—Am. trop.
demissa, *DC. l. c.*—N. Granat.
frutescens, *DC. l. c.*—Am. bor.
peruviana, *DC. l. c.*—Peruv.

BORSCZOWIA, Bunge, in Act. Hort. Petrop. v. (1877) 643. *CHENOPODIACEAE*, Benth. & Hook. f. iii. 68.
aralo-caspica, *Bunge, l. c.*—Reg. Casp.

BORSZCZOWIA, Bunge, in Act. Hort. Petrop. vi. (1879) 424 = praec.

BORYA, Labill. Nov. Holl. Pl. i. 81. t. 107 (1804). *LILIACEAE*, Benth. & Hook. f. iii. 797.
BAUMGARTENIA. Sprcng. Syst. ii. 91 (1825).
DAVIESIA, Poir. Encyc. Suppl. v. 480 (1817).
Cataractae, Endl. in Lehm. Pl. Preiss. ii. 43 = nitida.
gracilis, Endl. l. c. = nitida.
lucens, Poir. Encyc. viii. 615 = nitida.
nitens, Steud. Nom. ed. II. i. 192 = seq.
nitida, *Labill. Nov. Holl. Pl.* i. 81. t. 107.—Austral.
scirpoidea, Lindl. Swan River App. 57. t. 9 = nitida.
septentrionalis, *F. Muell. Fragm.* v. 41.—Austral.
sphaerocephala, R. Br. Prod. 286 = nitida.
sublanosa, F. Muell. ex Benth. Fl. Austral. vii. 71 = nitida.

BORYA, Willd. Sp. Pl. iv. 711 (1805) = **Forestiera**, Poir. (Oleac.).
acuminata, Willd. l. c. = F. acuminata.
cassinoides, Willd. l. c. = F. cassinoides.
distichophylla, Nutt. Gen. Am. ii. 232 = Buckleya distichophylla.
ligustrina, Willd. l. c. = F. ligustrina.
nitida, Willd. Enum. Hort. Berol. Suppl. [66] nomen = F. acuminata.
obovata, Desf. Cat. Hort. Par. ed. III. 88 = F. prinoides.
porulosa, Willd. Sp. Pl. iv. 711 = F. porulosa.
prinoides, Willd. Enum. Hort. Berol. Suppl. [66]; ex Spreng. Syst. iii. 900 = F. prinoides.
retusa, Willd. l. c. [66] = F. retusa.

BOSCA, Vell. Fl. Flum. 150. (1825); iv. t. 11 = **Funifera**, Leandr.? (Thymel.).
stupacea, Vell. ll. cc.—Bras.

BOSCHERIA, Carr. in Rev. Hortic. (1872) 199 = BOSSCHERIA, De Vriese & Teijsm. = **Ficus**, Tourn. (Urtic.).

BOSCHIA, Korth. Verh. Nat. Gesch. Bot. 257. t. 69 (1842). *MALVACEAE*, Benth. & Hook. f. i. 213.
HETEROPYXIS, Griff. Notul. iv. 524. t. 594 (1854).
acutifolia, *Mast. in Journ. Linn. Soc.* xiv. (1875) 503.—Borneo.
excelsa, *Korth. Verh. Nat. Gesch. Bot.* 258. t. 69.—Sumatra.
grandiflora, *Mast. in Journ. Linn. Soc.* xiv. (1875) 502.—Borneo.
Griffithii, *Mast. l. c.* 503.—Malaya.

BOSCHNIAKIA, C. A. Mey. (Bong. Veg. Sitka, 41) in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 157. *OROBANCHACEAE*, Benth. & Hook. f. ii. 985.
STELLARA, Fisch. & Reut. in DC. Prod. xi. 39 (1847).
glabra, *C. A. Mey. ex Bong. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 157.—Sibir.; Japon.; Am. bor.
himalaica, *Hook. f. & Thoms. in Fl. Brit. Ind.* iv. 327.—Reg. Himal.
Hookeri, *Walp. Rep.* iii. 479.—Am. bor. occ.
strobilacea, *A. Gray, in Pacif. Rail. Rep.* iv. 118.—Calif.

BOSCIA, Lam. Illustr. t. 395 (1804?). *CAPPARIDEAE*, Benth. & Hook. f. i. 108.
PODORIA, Pers. Syn. ii. 5 (1807).
angustifolia, Harv. in Harv. & Sond. Fl. Cap. i. Add. 19 = Niebuhria angustifolia.
angustifolia, *A. Rich. in Guill. & Perr. Tent. Fl. Seneg.* i. 26.—Afr. trop.
caffra, *Sond. in Linnaea*, xxiii. (1850) 8.—Afr. austr.
firma, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* xiv. [1884] 62, in nota.—Afr. trop.
integrifolia, Brunn. in Flora, xxiii. (1840) II. Beibl. 32 = angustifolia.
intermedia, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 27 = angustifolia.
microphylla, *Oliver, Fl. Trop. Afr.* i. 93.—Afr. austr.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 164.—Afr. trop.

BOSCIA:—
octandra, Fenzl, ex Radlk. in Sitzb. Math.-Phys. Acad. Muench. xiv. [1884] 62 = firma.
octandra, *Hochst. ex Radlk. l. c.*—Afr. trop.
reticulata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 28.—Abyss.
salicifolia, *Oliver, Fl. Trop. Afr.* i. 93.—Afr. trop.
senegalensis, Hochst. ex Walp. Ann. iii. 824 = angustifolia.
senegalensis, *Lam. Illustr.* t. 395.—Afr. trop.
urens, *Welw. in Oliver, Fl. Trop. Afr.* i. 93.—Afr. trop.

BOSCIA, Thunb. Prod. Pl. Cap. 32 (1794) = **Toddalia**, Juss. (Rutac.).
undulata, Thunb. l. c. = T. lanceolata.

BOSEA, Linn. Coroll. 5 (1737); [Bosia, Hort. Cliff. 84 (1737).] *AMARANTACEAE*, Benth. & Hook. f. iii. 26.
RODETIA, Moq. in DC. Prod. xiii. II. 323 (1849).
Amherstiana, *Hook. f. Fl. Brit. Ind.* iv. 716.—Reg. Himal.
cannabina, *Lour. Fl. Cochinch.* i. 176.—Cochinch.
cypria, *Boiss. ex Hook. f. Fl. Brit. Ind.* iv. 716, in nota.—Cyprus.
humilis, Salisb. Prod. 154 = B. Yervamora.
trinervia, Roxb. Hort. Beng. 86; Fl. Ind. ii. 87 = Celtis Wightii.
Yervamora, *Linn. Sp. Pl.* 225.—Ins. Canar.

BOSISTOA, F. Muell. ex Benth. Fl. Austral. i. 359 (1863). *RUTACEAE*, Benth. & Hook. f. i. 990.
euodiformis, *F. Muell. Fragm.* ix. 174.—Austral.
pentacocca, Baill. Hist. Pl. iv. 470 = sapindiformis.
sapindiformis, *F. Muell. ex Benth. Fl. Austral.* i. 359.—Austral.

BOSQUIEA, Thou. ex Baill. Adansonia, iii. (1862-63) 338. t. 10. *URTICACEAE*, Benth. & Hook. f. iii. 370.
Boiviniana, *Cord. ex Baill. l. c.* iii. (1862-63) 340.—Madag.
gymnandra, *Baker, Fl. Maurit.* 283.—Ins. Mascar.
Phoberos, *Baill. Adansonia*, viii. (1867) 72.—Afr. trop.
Thouarsiana, *Cord. ex Baill. l. c.* iii. (1862-63) 339.—Madag.

BOSSCHERIA, De Vriese & Teijsm. in Tijdschr. Nederl. Ind. xxiii. (1861) 213 = **Ficus**, Tourn. (Urtic.).
Minahassae, De Vriese & Teijsm. l. c. 214 = F. Minahassae.

BOSSIAEA, Vent. Jard. Cels, i. 7. t. 7 (1800). *LEGUMINOSAE*, Benth. & Hook. f. i. 473.
LALAGE, Lindl. Bot. Reg. t. 1722 (1834).
SCOTTEA, R. Br. in Ait. Hort. Kew. ed. II. iv. 269 (1812).
aculeata, *F. Muell. Fragm.* ii. 120.—Austral.
Aquifolium, *Benth. Fl. Austral.* ii. 157.—Austral.
Armitii, *F. Muell. Fragm.* ix. 44.—Austral.
biloba, *Benth. in Enum. Pl. Hueg.* 36.—Austral.
bracteosa, *F. Muell. ex Benth. Fl. Austral.* ii. 166.—Austral.
Brownei, *Benth. l. c.* 163.—Austral.
buxifolia, *A. Cunn. in Field, N. S. Wales*, 348.—Austral.
calycina, *Benth. Fl. Austral.* ii. 159.—Austral.
carinalis, *Benth. in Mitch. Journ. Trop. Austr.* 290.—Austral.
cinerea, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 266.—Austral.
coccinea, Bonpl. Jard. Malm. 128. t. 52 = cinerea.
concinna, *Benth. Fl. Austral.* ii. 161.—Austral.
cordifolia, Sweet, Fl. Austral. t. 20 = cinerea.
cordigera, *Benth. ex Hook. f. Fl. Tasm.* i. 95. t. 15.—Austral.
decumbens, F. Muell. Fragm. i. 9 = buxifolia.
dentata, *Benth. Fl. Austral.* ii. 156.—Austral.
disticha, *Lindl. Bot. Reg.* (1841) *Misc.* 38.—Austral.
distichoclada, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 39 = foliosa.
divaricata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 285.—Austral.
egena, *F. Muell. in Hook. Kew Journ.* viii. (1856) 43.—Austral.
Endlicheri, *Meissn. in Lehm. Pl. Preiss.* i. 83.—Austral.
ensata, Hook. f. Fl. Tasman. i. 94 = riparia.

BOSSIAEA:—
ensata, Meissn. in Lehm. Pl. Preiss. i. 81 = rufa.
ensata, *Sieber, ex DC. Prod.* ii. 117.—Austral.
eriocarpa, *Benth. in Enum. Pl. Hueg.* 36.—Austral.
foliosa, *A. Cunn. in Field, N. S. Wales*, 347.—Austral.
Gilbertii, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 286 = eriocarpa.
Hendersoni, *Hort. ex Regel, Gartenfl.* (1866) 322.—Austral.
heterophylla, *Vent. Jard. Cels*, t. 7.—Austral.
humilis, Meissn. in Lehm. Pl. Preiss. i. 85 = prostrata.
kiamensis, *Benth. Fl. Austral.* ii. 158.—Austral.
lalagoides, F. Muell. Fragm. iv. 12 = divaricata.
lanceolata, Sims, Bot. Mag. t. 1144 = heterophylla.
lenticularis, Lodd. Bot. Cab. t. 1238 = rhombifolia.
lenticularis, *Sieber, ex DC. Prod.* ii. 117.—Austral.
linnaeoides, G. Don, Gen. Syst. ii. 129 = prostrata.
linophylla, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 268.—Austral.
microphylla, *Sm. in Trans. Linn. Soc.* ix. (1808) 303.—Austral.
neo-angliea, *F. Muell. Fragm.* v. 106.—Austral.
nervosa, Meissn. in Bot. Zeit. xiii. (1855) 31 = eriocarpa.
Nummularia, Endl. Nov. Stirp. Dec. 27 = prostrata.
ornata, *Benth. Fl. Austral.* ii. 158.—Austral.
ovalifolia, Endl. Nov. Stirp. Dec. 26 = eriocarpa.
ovata, G. Don, Gen. Syst. ii. 128 = heterophylla.
ovata, Sm. in Trans. Linn. Soc. ix. (1808) 303 = prostrata.
oxyclada, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 284 = rufa.
paucifolia, Benth. ex Lindl. Bot. Reg. (1841) Misc. 53 = rufa.
peduncularis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 287.—Austral.
phylloclada, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 52.—Austral.
Preissii, *Meissn. in Lehm. Pl. Preiss.* i. 82.—Austral.
prostrata, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 268.—Austral.
pulchella, *Meissn. in Lehm. Pl. Preiss.* i. 84.—Austral.
rhombifolia, *Sieber, ex DC. Prod.* ii. 117.—Austral.
rigida, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 285 = Preissii.
riparia, *A. Cunn. ex Benth. Fl. Austral.* ii. 166.—Austral.
rosmarinifolia, *Lindl. in Mitch. Three Exped.* ii. 178.—Austral.
Rossii, F. Muell. Fragm. iii. 94, 168 = Templetonia sulcata.
rotundifolia, DC. Prod. ii. 117 = rhombifolia.
rufa, *R. Br. in Ait. Hort. Kew.* ed. II. iv. 266.—Austral.
rufa, Maund, Botanist, t. 81 = ensata.
rupicola, *A. Cunn. ex Benth. Fl. Austral.* ii. 162.—Austral.
scolopendria, *Sm. in Trans. Linn. Soc.* ix. (1808) 303.—Austral.
Scortechinii, *F. Muell. Census*, 140, nomen.—Austral.
spinescens, Meissn. in Lehm. Pl. Preiss. i. 82 = rufa.
stenophylla, *F. Muell. Fragm.* i. 9.—Austral.
strigillosa, *Benth. Fl. Austral.* ii. 157.—Austral.
sulcata, *Meissn. in Lehm. Pl. Preiss.* i. 81.—Austral.
tenuicaulis, R. Grah. in Edinb. N. Phil. Journ. xxix. (July 1840) 171 = cinerea.
virgata, Hook. Bot. Mag. t. 3986 = rufa.
Walkeri, *F. Muell. Fragm.* ii. 120.—Austral.
Webbii, *F. Muell. in Melb. Chem. et Drug.* (Dec. 1882).—Austral.

BOSSIENA, Pers. Syn. ii. 286 (1807) = **Bossiaea**, Vent. (Legum.).

BOSTRYCHANTHERA, Benth. in Benth. & Hook. f. Gen. ii. 1216 (1876). *LABIATAE*, Benth. & Hook. f. l. c.
deflexa, *Benth. l. c.*—China.

BOSWELLIA, Roxb. ex Colebr. in As. Res. ix. (1807) 379; Pl. Corom. iii. 4. t. 207 (1819). *BURSERACEAE*, Benth. & Hook. f. i. 322.
LIBANUS, Colebr. in As. Res. ix. 382, in nota (1807), t. 5. f. 1.
PLOESSLIA, Endl. Nov. Stirp. Dec. 39 (1839).

BOSWELLIA :—
Ameero, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 505.—Ins. Socotra.
balsamifera, Spreng. Syst. ii. 313 = glabra.
Bhaw-Dajiana, *Birdw. in Trans. Linn. Soc.* xxvii. (1870) 144. t. 31.—Afr. trop.
Carterii, *Birdw. l. c.* 143.—Arabia.
elongata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 505.—Ins. Socotra.
Frereana, *Birdw. in Trans. Linn. Soc.* xxvii. (1870) 146. t. 31.—Afr. trop.
glabra, Roxb. Hort. Beng. 90; Fl. Ind. ii. 384 = serrata.
hirsuta, Sm. in Rees, Cyclc. xxxix. n. 2 = Canarium hirsutum.
integra, Blanco, Fl. Filip. ed. II. 242 = Bursera javanica.
javanica, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) I. 612.—Java.
neglecta, *S. Moore, in Journ. Bot.* xv. (1877) 67. t. 185.—Afr. trop.
papyrifera, *Hochst. in Flora*, xxvi. (1843) 81.—Abyss.
sacra, Flueck. Lehrb. Pharmak. (1867) 31 = Carterii.
serrata, *Roxb. ex Colebr. in As. Res.* ix. (1807) 379. t. 5.—Ind. or.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 505.—Ins. Socotra.
thurifera, Roxb. ex Flem. in As. Res. xi. (1810) 158 = serrata.

BOTELUA, Lag. in Varied. Cienc. ii. IV. (1805) 134 = **Bouteloua**, Lag. (Gramin.).

BOTHERBE, Steud. ex Klatt, in Linnaea, xxxi. (1861–62) 562 = **Calydorea**, Herb. (Irid.).
bulbosa, Steud. ex Klatt, l. c. = C. speciosa.
campestris, Klatt, l. c. 563 = C. campestris.
gracilis, Klatt, l. c. 565 = C. nuda.
longispatha, Klatt, l. c. 564 = Sphenostigma longispatha.
luteola, Klatt, l. c. 563 = C. lutcola.

BOTHRIOCHILUS, Lem. Jard. Fleur. iii. (1852-53) t. 325; Illustr. Hortic. iii. (1856) Misc. 30, 31 = **Coelia**, Lindl. (Orchid.).
bellus, Lem. ll. cc. = C. bella.

BOTHRIOCLINE, Oliver, ex Benth. in Hook. Ic. Pl. xii. 30. t. 1133 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 226.
Schimperi, *Oliver & Hiern, ex Benth. l. c.*—Abyssin.

BOTHRIOSPERMUM, Bunge, Enum. Pl. Chin. Bor. 47 (1832). *BORAGINEAE*, Benth. & Hook. f. ii. 853.
asperugoides, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 150 = tenellum.
bicarinatum, Bunge, ex Steud. Nom. ed. II. i. 219 = seq.
bicarunculatum, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. (1835) 23 = chinense.
chinense, *Bunge, Enum. Pl. Chin. Bor.* 47.—China bor.
chinense, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 23 = Kusnetzowii.
Kusnetzowii, *Bunge, ex DC. Prod.* x. 116, cum cit. falsa.—Mongolia.
marifolium, A. DC. Prod. x. 116 = Heliotropium marifolium.
perenne, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 95 = tenellum.
secundum, *Maxim. Prim. Fl. Amur.* 202.—China.
tenellum, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. (1835) 24.—China bor.

BOTHRIOSPORA, Hook. f. in Hook. Ic. Pl. xi. 55. t. 1069 (1870). *RUBIACEAE*, Benth. & Hook. f. ii. 76.
corymbosa, *Hook. f. l. c.*—Am. austr.

BOTOR, Adans. Fam. ii. 326 (1763) = **Psophocarpus**, Neck. (Legumin.).

BOTRIA, Lour. Fl. Cochinch. 153 (1790) = **Vitis**, Tourn. (Ampel.).
africana, Lour. l. c. 154 = V. africana.

BOTROPHIS, Rafin. Med. Fl. i. 85 (1828) = **Cimicifuga**, Linn. (Ranunc.).
actaeoides, Rafin. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 = C. racemosa.
pumila, Rafin. Med. Fl. i. 86 = C. racemosa?
serpentaria, Rafin. l. c. 85 = C. racemosa.

BOTRYA, Juss. in Mém. Mus. Par. iii. (1817) 444 = BOTRIA, Lour. = **Vitis**, Tourn. (Ampelid.).

BOTRYADENIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 30 (1835) = **Myriactis**, Less. (Compos.).
Gmelini, Fisch. & Mey. l. c. = M. Gmelini.

BOTRYANTHE, Klotzsch, in Wiegm. Archiv, ii. (1841) 190. t. 9 = **Plukenetia**, Linn. (Euphorb.).
concolor, Klotzsch, l. c. 191.—Bras.
discolor, Klotzsch, l. c.—Bras.

BOTRYANTHUS, Kunth, Enum. Pl. iv. 310 (1843) = **Muscari**, Mill. (Liliac.).
albovirens, Tod. Nuovi Gen. 76 = M. commutatum.
alpestris, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
atlanticus, Nym. Consp. 734 = M. racemosum.
Aucheri, Boiss. Diagn. Ser. I. v. 63 = M. Aucheri.
autumnalis, Heynh. Nom. ii. 75 = M. parviflorum.
Boraeanus, Jord. & Fourr. Ic. Pl. Eur. ii. 22 = M. botryoides.
breviscapus, Tod. Hort. Panorm. t. 5.—Hab.?
candidus, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
commutatus, Kunth, Enum. Pl. iv. 311 = M. racemosum.
compactus, Jord. & Fourr. Ic. Pl. Eur. ii. 24 = M. botryoides.
conicus, Jord. & Fourr. l. c. = M. botryoides.
dilutus, Elwes, ex Baker, in Gard. Chron. (1878) I. 708 = M. dilutum.
festinus, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
Gussonei, Tod. Ind. Sem. Panorm. (1872) 38; Nuov. Giorn. Bot. v. (1873) 157 = M. Gussonii.
Heldreichii, Jord. & Fourr. Ic. Pl. Eur. ii. 24 = M. Heldreichii.
Kerneri, Marches.—Cf. Oestr. Bot. Zeitschr. xxxii. (1882) 339 = M. botryoides.
Lelievrii, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
micranthus, Hort. Berol. ex Baker, in Gard. Chron. (1878) I. 799 = M. micranthum.
Mordoanus, Nym. Consp. 734 = M. Mordoanum.
neglectus, Jord. & Fourr. Ic. Pl. Eur. ii. 24 = M. neglectum.
neglectus, Kunth, Enum. Pl. iv. 679 = M. neglectum.
odorus, Kunth, l. c. 311 = M. racemosum.
pallens, Kunth, Enum. Pl. iv. 312 = M. pallens.
paradoxus, Kunth, l. c. = Hyacinthus paradoxus.
parviflorus, Kunth, l. c. = M. parviflorum.
pulchellus, Jord. & Fourr. Ic. Pl. Eur. ii. 23 = M. botryoides.
racemosus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 160 = M. racemosum.
Sartorii, Tod. Hort. Panorm. t. 5 = M. racemosum.
Sauli, Jaub. & Spach, Illustr. iv. 44 = M. moschatum.
speciosus, Marches.—Cf. Oestr. Bot. Zeitschr. xxxii. (1882) 339 = M. neglectum.
stereophyllus, Herbich, Stirp. Rar. Bucov. 30 (1855) 182 = Hyacinthus leucophaeus.
Strangwaysii, Kunth, Enum. Pl. iv. 679 = M. racemosum.
Szovitsianus, Hort. Bonn. ex Baker, in Gard. Chron. (1878) I. 799 = M. Szovitsianum.
vulgaris, Kunth, Enum. Pl. iv. 311 = M. botryoides.

BOTRYCARPUM, A. Rich. Bot. Méd. ii. 487 (1823), in adnot. = **Ribes**, Linn. (Saxifrag.).
nigrum, A. Rich. l. c. 490 = R. nigrum.

BOTRYCERAS, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 396. *ANACARDIACEAE*, Benth. & Hook. f. i. 426.
DAPHNITIS, Spreng. Syst. i. 370 (1825).
LAUROPHILLUS, Roem. & Schult. Syst. iii. 30 (1818).
LAUROPHYLLUS, Thunb. Nov. Gen. vi. 104 (1792), nomen prius.
laurinum, *Willd. l. c.* 397.—Afr. austr.

BOTRYCOMUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 160 = **Muscari**, Mill. (Liliac.).
vulgaris, Fourr. l. c. = M. comosum.

BOTRYDIUM, Spach, Hist. Vég. Phan. v. 298 (1836) = **Chenopodium**, Linn.
aromaticum, Spach, l. c. 299 = C. Botrys.
Schraderi, Spach, l. c. = C. foetidum.

BOTRYMORUS, Miq. Fl. Ind. Bat. i. II. 282 (1859). *ARTOCARPEAE*.
paniculata, *Miq. l. c.* 283.—Malaya.

BOTRYODENDRUM, Endl. Prod. Fl. Norf. 62 (1833) = **Meryta**, Forst. (Araliac.).
angustifolium, Endl. l. c. = M. angustifolia.
capitatum, Endl. ex Heynh. Nom. ii. 75 = Oreopanax capitatum.
coriaceum, Panch. ex Baill. Adansonia, xii. (1878) 153 = M. coriacea.
latifolium, Endl. Prod. Fl. Norf. 62 = M. latifolia.
macrophyllum, Rich, ex A. Gray, Bot. U. St. Expl. Exped. i. 732. t. 97 = M. macrophylla.
Sinclairii, Hook. f. Fl. N. Zel. 97 = M. Sinclairii.
taitense, Guill. in Ann. Sc. Nat. Sér. II. vii. (1837) 349.—Ins. Tahiti.

BOTRYOLOTUS, Jaub. & Spach, Illustr. i. 124. t. 63 = **Trigonella**, Linn. (Legumin.).
cachemyrianus, Jaub. & Spach, l. c. 125 = T. Emodi.
persicus, Jaub. & Spach, l. c. 124. t. 63 = T. elliptica.

BOTRYOPANAX, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 5 (1863–64) = **Polyscias**, Forst. (Araliac.).
borbonica, Miq. l. c. = Ins. Borbon.

BOTRYOPSIS, Miers, in Ann. Nat. & Mag. Hist. Ser. II. vii. 43 (1851) = **Chondodendron**, Ruiz & Pav. (Menisperm.).
platyphylla, Miers, l. c. = C. platyphyllum.
Spruceana, Eichl. in Flora, xlvii. (1864) 394 = C. cretosum.

BOTRYOROPIS, Presl, Epim. Bot. 220 (1849) = **Barringtonia**, Forst. (Myrtac.).
luzonensis, Presl, l. c. = B. luzonensis.

BOTRYOSICYOS, Hochst. in Flora, xxvii. (1844) Bes. Beil. 3 = **Dioscorea**, Linn.
pentaphyllum, Hochst. l. c. = D. pentaphylla.

BOTRYPHILE, Salisb. Gen. Pl. Fragm. 25 (1866) = **Muscari**, Mill. (Liliac.).
botryoides, Salisb. l. c. = M. botryoides.
parviflorus, Salisb. l. c. = M. parviflorum.
racemosus, Salisb. l. c. = M. racemosum.

BOTRYS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 138 = **Teucrium**, Linn. (Labiat.).
chamaedryoides, Fourr. l. c. = T. Botrys.

BOTTIONEA, Colla, in Mem. Acc. Torin. xxxvii. (1834) 43. t. 1. *LILIACEAE*, Benth. & Hook. f. iii. 789.
ENDOCOMA, Rafin. Fl. Tellur. ii. 57 (1836).
TRICHOPETALUM, Lindl. Bot. Reg. t. 1535 (1832).
thysanthoides, *Colla, l. c.* 45. t. 1.—Chili.

BOUCEROSIA, Wight & Arn. Contrib. Bot. Ind. 34 (1834). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 782.
APTERANTHES, Mikan, in Nov. Act. Nat. Cur. xvii. (1835) 593. t. 41.
DESMIDOCHUS, Reichb. Consp. 130 (1828).
DESMIDORCHIS, Ehrenb. in Linnaea, iv. (1829) 94.
HUTCHINIA, Wight & Arn. Contrib. Bot. Ind. 34 (1834).
HUTSCHINIA, D. Dietr. Syn. i. 579 (1839).
Aaronis, *Hart, in Trans. Roy. Irish Acad.* xxviii. (1885) 436.—Palaestin.
acutangula, *Decne. in DC. Prod.* viii. 648.—Afr. trop.
adscendens, Wall. Cat. n. 8146, partim = Caralluma edulis.
Aucheriana, *Decne. in DC. Prod.* viii. 649.—Arabia; Persia; Ind. bor. occ.
campanulata, Wight, Ic. vi. t. 1287 = umbellata.
crenulata, *Wight & Arn. Contrib. Bot. Ind.* 34.—Ind. or.

BOUCEROSIA:—
cylindrica, *Brongn. in Bull. Soc. Bot. Fr.* vii. (1860) 900.—Abyss.
Decaisniana, *Lem. Herb. Gén. Amat.* Sér. II. iv. t. 21; *Decne. in DC. Prod.* viii. 648.—Afr. trop.
diffusa, *Wight, Ic.* vi. t. 1287.—Ind. or.
edulis, Edgew. in Journ. Linn. Soc. vi. (1862) 205 = Caralluma edulis.
Forskalii, *Decne. in DC. Prod.* viii. 648.—Arabia.
Gussoniana, *Hook. f. Bot. Mag.* t. 6137 in textu.—Algeria.
Hitchinia, Hook. f. Fl. Brit. Ind. iv. 78, lapsu = B. Hutchinia.
Hutchinia, *Decne. in DC. Prod.* viii. 649.—Ind. or.
incarnata, *N. E. Br. in Journ. Linn. Soc.* xvii. (1878) 166.—Afr. austr.
lasiantha, Wight, Ic. vi. t. 1286 = crenulata.
mammillaris, *N. E. Br. in Journ. Linn. Soc.* xvii. (1878) 165.—Afr. austr.
maroccana, *Hook. f. Bot. Mag.* t. 6137.—Marocco.
Munbyana, *Decne. ex Munby, Fl. Algér.* 25.—Algeria.
pauciflora, *Wight, in Madras Journ. Sc.* vi. (1837) t. ad p. 473.—Ind. or.
quadrangula, Decne. in DC. Prod. viii. 664, in syn. = Forskalii.
Russeliana, *Courb. ex Brongn. in Bull. Soc. Bot. Fr.* vii. (1860) 900.—Oriens.
sinaica, *Decne. in DC. Prod.* viii. 649.—Arabia.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 79.—Ins. Socotr.
Stocksiana, Boiss. Fl. Orient. iv. 63 = Caralluma edulis.
umbellata, *Wight & Arn. Contrib. Bot. Ind.* 34.—Ind. or.

BOUCHARDATIA, Baill. Adansonia, vii. (1867) 350 = **Melicope**, Forst. (Rutac.).
australis, Baill. l. c. 351. t. 10 = M. neurococca.
neurococca, Baill. l. c. ix. (1868) 110 = M. neurococca.

BOUCHEA, Cham. in Linnaea, vii. (1832) 252. *VERBENACEAE*, Benth. & Hook. f. ii. 1144.
Chascanum, E. Mey. Comm. Pl. Afr. Austr. 275 (1835).
Pleurostigma, Hochst. ex Schnizl. in Flora, (1842) I. Beib. 144.
Plexipus, Rafin. Fl. Tellur. ii. 104 (1836).
adenostachya, *Schau. in DC. Prod.* xi. 560.—Afr. trop.
agrestis, *Schau. l. c.* 558.—Bras.
cernua, *Schau. l. c.* 559.—Afr. austr.
copiapensis, C. Gay, Fl. Chil. v. 26. t. 55 = Priva laevis.
cuneifolia, *Schau. in DC. Prod.* xi. 559.—Afr. austr.
Ehrenbergii, *Cham. in Linnaea*, vii. (1832) 253.—Ins. S. Doming.
garepensis, *Schau. in DC. Prod.* xi. 560.—Afr. austr.
Hanningtonii, *Oliver, in Hook. Ic. Pl.* t. 1446.—Afr. trop.
hederacea, *Sond. in Linnaea*, xxiii. (1850) 36.—Afr. austr.
hyderabadensis, *Walp. Rep.* iv. 12.—Ind. or.
incrassata, *Lange, Ind. Sem. Haun.* (1870) 31.—Hab.?
lactevirens, *Schau. in DC. Prod.* xi. 557.—Bras.
latifolia, *Harv. Thes. Cap.* ii. 57.—Afr. austr.
linifolia, *A. Gray, in Am. Journ. Sc.* Ser. II. xvi. (1853) 98.—N. Mexic.
marrubiifolia, *Schau. in DC. Prod.* xi. 558.—Afr. trop.; Arab.; Scind.
pinnatifida, *Schau. l. c.* 560.—Afr. austr.
pseudogervaô, *Cham. in Linnaea*, vii. (1832) 252.—Bras.
pterygocarpa, *Schau. in DC. Prod.* xi. 558.—Nubia.
pubescens, Schau. l. c. 560 = pumila.
pumila, *Schau. l. c.*—Afr. austr.
sessilifolia, *Vatke, in Linnaea*, xliii. (1880-82) 529.—Afr. trop.
spathulata, *Torr. Bot. Mex. Bound.* 126.—Am. bor. occ.

BOUCHETIA, DC. ex Dun. in DC. Prod. xiii. i. 589 (1852). *SOLANACEAE*, Benth. & Hook. f. ii. 908.
Leucanthea, Scheele, in Linnaea, xxv. (1852) 258.
erecta, *DC. ex Dun. l. c.*—Texas; Mexic.
procumbens, *DC. ex Dun. l. c.*—Mexic.

BOUEA, Meissn. Gen. 75; Comm. 55 (1837). *ANACARDIACEAE*, Benth. & Hook. f. i. 420.
Cambessedea, Wight & Arn. Prod. i. 170 (1834).
angustifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 204.—Borneo.
Brandisiana, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 50 = burmanica.
burmanica, *Griff. Pl. Cantor*, 14.—Burma; Malaya.
diversifolia, Miq. Fl. Ind. Bat. Suppl. i. 522 = microphylla.
Gandaria, Blume, Mus. Bot. Lugd. Bat. i. 204 = macrophylla?
macrophylla, *Griff. Pl. Cantor*, 15; *Notul.* iv. 420.—Malaya.
microphylla, *Griff. ll. cc.* 15, 423.—Malaya.
myrsinoides, Blume, Mus. Bot. Lugd. Bat. i. 204 = microphylla?
oppositifolia, Meissn. Gen. 75; Walp. Rep. i. 556 = burmanica.

BOUGAINVILLAEA, Comm. ex Juss. Gen. 91 (1789) (Buginvillaea); Choisy, in DC. Prod. xiii. II. 437 (1849). *NYCTAGINEAE*, Benth. & Hook. f. iii. 7.
Josepha, Vell. Fl. Flum. 154 (1825); iv. t. 16.
bracteata, Pers. Syn. i. 418 = spectabilis.
brasiliensis, Lund, ex Choisy, in DC. Prod. xiii. II. 437 = virescens.
brasiliensis, Raeusch. Nom. ed. III. 112; [Nees, in] Princ. Neuw. Reise, i. 44 = spectabilis.
frondosa, *Griseb. in Goett. Abh.* xix. (1874) 89.—Reg. Argent.
glabra, *Choisy, in DC. Prod.* xiii. II. 437.—Bras.
infesta, *Griseb. in Goett. Abh.* xxiv. (1879) 40.—Reg. Argent.
patagonica, *Decne. in Orbign. Voy.* t. 8.—Patagon.
peruviana, *Humb. & Bonpl. Pl. Aequin.* i. 173. t. 49.—Peruv.
peruviana, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 39 = spectabilis.
pomacea, *Choisy, in DC. Prod.* xiii. II. 438.—Bras.
praecox, *Griseb. in Goett. Abh.* xxiv. (1879) 40.—Reg. Argent.
racemosa, Blanco, Fl. Filip. ed. I. 307 = Pisonia excelsa.
rubicunda, Schott, ex Rohrb. in Mart. Fl. Bras. xiv. II. 351 = glabra.
speciosa, Schnizl. Ic. Fam. Nat. t. 104 = spectabilis.
spectabilis, *Willd. Sp. Pl.* ii. 348.—Bras.
stipitata, *Griseb. in Goett. Abh.* xix. (1874) 88.—Reg. Argent.
virescens, *Choisy, in DC. Prod.* xiii. II. 437.—Bras.

BOUGAINVILLEA, Spach, Hist. Vég. Phan. x. 516 (1841) = **Bougainvillaea**, Juss. (Nyctag.).

BOUGUERIA, Decne. in Ann. Sc. Nat. Sér. II. v. (1836) 132. *PLANTAGINEAE*, Benth. & Hook. f. ii. 1224.
nubicola, *Decne. l. c.*—Peruv.; Boliv.

BOULARDIA, F. Schultz, Arch. Fl. Fr. et Allem. 103 (1847) = **Orobanche**, Tourn.
latisquama, F. Schult. l. c. 104.—Algeria.

BOUPHON, Lem. in Orb. Dict. ii. 686 (1842) = seq.

BOUPHONE, Lem. in Orb. Dict. ii. 778 (1842) = **Buphane**, Herb. (Amaryll.).

BOURGAEA, Coss. Notes Crit. 39 (1849) = **Cynara**, Vaill. (Compos.).
humilis, Coss. l. c. = C. humilis.
Tournefortii, Willk. in Linnaea, xxv. (1852) 39 = C. Tournefortii.

BOURGIA, Scop. Introd. 158 (1777). *BORAGIN.*?

BOURJOTIA, Pomel, Nouv. Mat. Fl. Atl. 89 (1874) = **Heliotropium**, Linn. (Boragin.).
Kralikii, Pomel, l. c. = H. Kralikii.
pterocarpa, Pomel, l. c. 90 = H. pterocarpum.

BOURRERIA, P. Br. Hist. Jamaic. 168 (1756). *BORAGINEAE*, Benth. & Hook. f. ii. 840.
Beurreria, Jacq. Enum. Pl. Carib. 2 (1760).
Crematomia, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. iii. (1869) 300.

BOURRERIA:—
Hymenesthes, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 26. t. 6.
Morelosia, La Llav. & Lex. Nov. Veg. Descr. i. 1 (1824).
Andrieuxii, *Hemsl. Biol. Centr. Am. Bot.* ii. 369.—Mexic.
aspera, G. Don, Gen. Syst. iv. 390 = Ehretia laevis.
baccata, *Rafin. Sylva Tellur.* 42.—Hab.?
calophylla, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 528.—Cuba.
cassinifolia, *Griseb. l. c.*—Cuba.
cestroides, *Spreng. Syst.* iv. *Cur. Post.* 66.—Am. austr.
clariuscula, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 204.—Am. austr.
dichotoma, Rottl. ex G. Don, Gen. Syst. iv. 390 = Ehretia laevis?
divaricata, *G. Don, l. c.* 387.—Cuba.
domingensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 528.—Ind. occ.
exsucca, *Jacq. Enum. Pl. Carib.* 14.—Am. austr.
formosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 369.—Mexic.
glabra, G. Don, Gen. Syst. iv. 390 = Cordia Myxa.
grandiflora, *Bertol. Fl. Guatim.* 10.—Guatemala.
havanensis, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 207.—Cuba.
Huanita, *Hemsl. Biol. Centr. Am. Bot.* ii. 370.—Mexic.
laevis, G. Don, Gen. Syst. iv. 390 = Ehretia laevis.
laxa, *G. Don, l. c.*—Ins. Mascar.
lincaris, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 210.—Ind. occ.
microphylla, *Griseb. Cat. Pl. Cub.* 210.—Cuba.
ovata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 203.—Ind. occ.
punctata, G. Don, Gen. Syst. iv. 390 = Ehretia laevis.
Radula, *G. Don, l. c.*—Ins. S. Doming.
recurva, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 203.—Ind. occ.
reticulata, *Griseb. Cat. Pl. Cub.* 210.—Cuba.
retusa, G. Don, Gen. Syst. iv. 390 = Ehretia retusa.
revoluta, *H. B. & K. Nov. Gen. et Sp.* iii. 67.—Mexic.
rigida, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. iii. (1869) 204.—Ind. occ.
setosa, G. Don, Gen. Syst. iv. 390 = Ehretia setosa.
spathulata, *Hemsl. Biol. Centr. Am. Bot.* ii. 370.—Mexic.
spinifex, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 528.—Cuba.
succulenta, *Jacq. Enum. Pl. Carib.* 14.—Ind. occ.
tomentosa, *G. Don, Gen. Syst.* iv. 390.—Ind. occ.
uniflora, G. Don, l. c. = Ehretia uniflora.
virgata, *G. Don, l. c.* 389.—Ind. occ.

BOUSSINGAULTIA, H. B. & K. Nov. Gen. et Sp. vii. 194. t. 645 (1825). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 77.
Tandonia, Moq. in DC. Prod. xiii. II. 226 (1849).
baselloides, *H. B. & K. Nov. Gen. et Sp.* vii. 196. t. 645.—Ecuador.
cordifolia, *Tenore, Ind. Sem. Neap.* (1852) *ex Ann. Sc. Nat.* Sér. III. xix. (1853) 355.—Hab.?
floribunda, *Moq. in DC. Prod.* xiii. II. 229.—N. Granat.
gracilis, *Miers, in Journ. Bot.* ii. (1864) 161. t. 18.—Bras.
Lachaumei, *Carr. in Rev. Hortic.* (1872) 279.—Cuba.
leptostachys, Moq. in DC. Prod. xiii. II. 229 = baselloides.
ramosa, *Hemsl. Biol. Centr. Am. Bot.* iii. 27.—Guatemala.

BOUTELOUA, Lag. in Varied. Cienc. ii. iv. (1805) 134 (Botelua). *GRAMINEAE*, Benth. & Hook. f. iii. 1168.
Actinochloa, Willd. ex Roem. & Schult. Syst. ii. 22, 417 (1818).
Aristidium, Endl. Gen. 94 (1836).
Atheropogon, Muhl. in Willd. Sp. Pl. iv. 937 (1805).
Chondrosium, Desv. Journ. Bot. iii. (1813) 68.
Corethrum, Vahl, in Skr. Naturk. Selsk. Kiøbenh. vi. (1810) 85.
Dinebra, DC. Cat. Hort. Mons. 104 (1813), pro parte.
Eutriana, Trin. Fund. Agrost. 161 (1820).

BOUTELOUA:—
HETEROSTECA, Desv. in Nov. Bull. Soc. Philom. ii. (1810) 188; Desv. Journ. Bot. iii. (1813) 68.
HETEROSTEGA, Kunth, in Mém. Mus. Par. ii. (1815) 73.
NESTLERA, Willd. ex Steud. Nom. ed. II. ii. 192 (1841).
POLYODON, H. B. & K. Nov. Gen. et Sp. i. 174. t. 55 (1815).
TRIAENA, H. B. & K. l. c. 178. t. 61 (1815).
TRIATHERA, Desv. in Nov. Bull. Soc. Philom. ii. (1810) 188.
TRIPLATHERA, Endl. Gen. 94 (1836).
aristidoides, *Griseb. Fl. Brit. W. Ind.* 537, in obs.—Mexic.
barbata, *Lag. Varied. Cienc.* ii. IV. (1805) 141; *Gen. et Sp. Nov.* 5.—Mexic.; Ins. Philipp.
brevifolia, Buckl. in Proc. As. Sc. Philad. '1862' (1863) 93 = eriopoda.
bromoides, *Lag. Gen. et Sp. Nov.* 5.—Mexic.
Burkei, *Scribn. ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 179.—Texas.
chrondrosioides, Benth. ex S. Wats. l. c. = ovata.
ciliata, *Griseb. in Goett. Abh.* xxiv. (1879) 302.—Reg. Argent.
curtipendula, Torr. in Emory, Notes Mil. Reconnois. 153 = racemosa.
disticha, *Benth. in Journ. Linn. Soc.* xix. (1882) 105.—Am. trop.
elatior, *Griseb. Fl. Brit. W. Ind.* 537.—Ind. occ.
eriopoda, *Torr. in Pacif. Rail. Rep.* iv. *Bot.* 155.—Am. bor.
foena, Bigelow, in Whipple Exped. Add. p. ii = hirsuta.
gracilis, Steud. Nom. ed. II. i. 219 = oligostachya.
gracilis, Vasey, in Wheeler, Exped. 287 = aristidoides.
Havardi, *Vasey, ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 179.—Texas.
hirsuta, *Lag. Varied. Cienc.* ii. IV. (1805) 141; *Gen. et Sp. Nov.* 5.—Mexic.
hirta, Hort. Madr. ex Lag. l. c. 141 = hirsuta.
Humboldtiana, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 532 = bromoides.
humilis, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 495.—Reg. Argent.
juncifolia, *Lag. Gen. et Sp. Nov.* 5.—Mexic.; Ind. occ.
juncifolia, Vasey, Cat. Gr. Un. St. 62 = bromoides.
litigosa, Lag. Gen. et Sp. Nov. 5 = juncifolia.
lophostachya, *Griseb. in Goett. Abh.* xxiv. (1879) 302.—Reg. Argent.
melicaeformis, Hornem. ex Steud. Nom. ed. I. 91, in syn. = racemosa.
melicoides, Steud. l. c. 115 = racemosa.
multiseta, *Griseb. in Goett. Abh.* xxiv. (1879) 303.—Bras.; Reg. Argent.
nana, *Griseb. l. c.*—Reg. Argent.
oligostachya, *Torr. ex A. Gray, Man.* ed. II. 553; *et in Pacif. Rail. Rep.* iv. *Bot.* 155.—Am. bor.
ovata, *Lag. Gen. et Sp. Nov.* 5.—Mexic.
pendula, Hort. Madr. ex Lag. in Varied. Cienc. ii. IV. (1805) 141 = racemosa.
pilosa, *Benth. ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 179.—Ins. Galap.
polystachya, *Torr. in Pacif. Rail. Rep.* v. 366. t. 10.—Am. bor.
porphyrantha, Wright, in Sauv. Fl. Cub. 192 = juncifolia.
prostrata, *Lag. in Varied. Cienc.* ii. IV. (1805) 141; *Gen. et Sp. Nov.* 5.—Mexic.
pumila, Buckl. in Proc. Ac. Sc. Philad. '1862' (1863) 93 = polystachya.
pusilla, *Vasey, in Bull. Torrey Bot. Club*, xi. (1884) 6.—N. Mexic.
racemosa, *Lag. in Varied. Cienc.* ii. IV. (1805) 141; *Gen. et Sp. Nov.* 5.—Mexic.
Rothrockii, *Vasey, Grasses U. St.* 33, nomen.—Am. bor.
scorpioides, *Lag. Gen. et Sp. Nov.* 5; *S. Wats. in Proc. Am. Acad.* xviii. (1883) 176.—Mexic.
simplex, *Lag. in Varied. Cienc.* ii. IV. (1805) 141.—Peruv.; Ins. Philipp.
tenuis, *Griseb. in Goett. Abh.* xxiv. (1879) 303.—Mexic.; Reg. Argent.
texana, *S. Wats. in Proc. Am. Acad.* xviii. (1883) 196.—Texas.
Triathera, *Benth. in Journ. Linn. Soc.* xix. (1882) 104.—Mexic.
trifida, *Thurb. ex S. Wats. in Proc. Am. Acad.* xviii. (1883) 177.—Mexic.

BOUTONIA, Boj. Hort. Maurit. 282 (1837) = **Mallotus**, Lour. (Euphorb.).
acuminata, Baill. Etud. Gén. Euphorb. 401 = M. Baillonianus.
mascarensis, Boj. Hort. Maurit. 282 = M. integrifolius.

BOUTONIA, DC. Rev. Bign. 18 (1838) = **Periblema**, DC. (Bignon.).
cuspidata, DC. l. c. = P. cuspidatum.

BOUTONIA, Hort. Erfurt. ex Steud. Nom. ed. II. i. 219 (1840) = **Goodenia**, Sm.
pomifera, Hort. Erfurt. ex Steud. l. c. = G. paniculata.

BOUVARDIA, Salisb. Parad. Lond. t. 88 (1805). *RUBIACEAE*, Benth. & Hook. f. ii. 36.
AEGINETIA, Cav. Ic. vi. 51. t. 572 (1801).
BOWVARDIA, Polak. in Linnaea, xli. (1877) 565.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 384.—Mexic.
bicolor, *Kunze, in Linnaea*, xx. (1847) 24.—Mexic.
candidissima ×, Hort.—Cf. Gard. Chron. (1873) 714.
Cavanillesii, *DC. Prod.* iv. 366.—Mexic.
chlorantha, *Bert. ex Roem. & Schult. Syst.* iii. *Mant.* 116.—Am. austr.
chrysantha, *Mart. Del. Sem. Hort. Monac.* (1848) 4.—Mexic.
coccinea, Link, Enum. Hort. Berol. i. 139 = triphylla.
coccinea, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 272.—Guiana.
cordifolia, *DC. Prod.* iv. 366.—Mexic.
corymbosa, *Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 46.—Nicarag.
crocata, *Van Houtte, Cat.* (1846).—Hab.?
Davisoni, *Hort.—Cf. Gard. Chron.* (1872) 7.—Hab.?
discolor, Hook. & Arn. Bot. Beech. Voy. 428 = Rondeletia dubia.
ferruginea, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 273.—Madag.
flava, *Decne. in Van Houtte, Fl. des Serres*, Sér. I. i. (1845) 90. t. 38.—Mexic.
glaberrima, *Engelm. in Wisliz. Tour N. Mexico*, 106.—Mexic.
glabra, *Polak. in Linnaea*, xli. (1877) 565.—Costa Rica.
havanensis, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 272.—Ind. occ.
hirsuta, *A. Rich. l. c.*—Ind. occ.
hirtella, *H. B. & K. Nov. Gen. et Sp.* iii. 384.—Mexic.
Houtteana, *Schlecht. in Fl. des Serres*, Sér. I. x. (1854-55) 149.—Mexic.
Humboldti, *Hort.—Cf. Gard. Chron.* (1873) 714, 716.—Hab.?
hypoleuca, *Benth. Pl. Hartw.* 288.—Mexic.
Jacquini, H. B. & K. Nov. Gen. et Sp. iii. 385 = triphylla.
jasminiflora, *Hort.—Cf. Gard. Chron.* (1872) 215.—Hab.?
laevis, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 236.—Mexic.
leiantha, *Benth. Pl. Hartw.* 85; *Decne. in Rev. Hortic.* Sér. III. v. (1851) 81.—Guatemala; Mexic.
linearis, *H. B. & K. Nov. Gen. et Sp.* iii. 383.—Mexic.
longiflora, H. B. & K. l. c. 386 = Houstonia longiflora.
microphylla, *Schlecht. in Linnaea*, xxvi. (1853-55) 112.—Mexic.
mollis, *Linden, Cat.* (1848).—*Cf. Schlecht. l. c.* 55.—Mexic.
multiflora, Schult. in Roem. & Schult. Syst. iii. 118 = Cavanillesii.
mutabilis, Hort. Berol. ex Walp. Ann. v. 127 = versicolor.
myrtifolia, *Schlecht. in Linnaea*, xxvi. (1853-55) 121.—Mexic.
obovata, Benth. Pl. Hartw. 16 = scabra.
obovata, *H. B. & K. Nov. Gen. et Sp.* iii. 385.—Mexic.
Oriana ×, *Pars. ex Van Houtte, Fl. des Serres*, xii. (1857) 159. t. 1265.
ovata, *A. Gray, Pl. Wright.* ii. 67.—N. Mexic.
suberula, Decne. in Fl. des Serres, Sér. I. i. (1845) 90, in descr. = flava.
quaterniflora, Steud. Nom. ed. II. i. 300, in syn. = seq.
quaternifolia, *DC. Prod.* iv. 365.—Mexic.
quinqueflora, *Dehnh. Revista Napol.* i. III. 167.—Mexic.

BOUVARDIA:—
racemosa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834).—Peruv.
rosea, *Schlecht. in Linnaea*, xxvi. (1853-55) 116.—Calif.
scabra, *Hook. & Arn. Bot. Beech. Voy.* 427.—Mexic.
scabrida, *Mart. & Gal. in Bull. Acad. Brux.* xi. (1844) 237.—Mexic.
scandens, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 272.—Hab.?
Schiediana, *Schlecht. in Linnaea*, xxvi. (1853-55) 123.—Mexic.
splendens, R. Grah. in Bot. Mag. t. 3781 = triphylla.
strigillosa, [Baxt. in] Loud. Hort. Brit. Suppl. iii. 502 = seq.?
strigosa, Benth. Pl. Hartw. 75 = Rondeletia strigosa.
tenuiflora, *Hort. Berol. ex Schlecht. in Linnaea*, xxvi. (1853-55) 97.—Mexic.
ternifolia, Schlecht. l. c. 98 = triphylla.
tolucana, *Hook. & Arn. Bot. Beech. Voy.* 427.—Mexic.
triflora, H. B. & K. Nov. Gen. et Sp. iii. 386. t. 288 = Houstonia triflora.
triphylla, *Salisb. Parad. Lond.* t. 88.—Mexic.
umbellata, *Hort.—Cf. Gard. Chron.* (1873) 714.—Hab.?
uniflora, *Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 272.—Am. trop.
versicolor, *Ker-Gawl. Bot. Reg.* t. 245.—Am. austr.
viminalis, *Schlecht. in Linnaea*, xxvi. (1853-55) 120.—Mexic.
viperalis, *Schlecht. l. c.* 114.—Mexic.
Vreelandi, Hort. ex Fl. et Pomol. (1872) 174 = Davisoni.
xylosteoides, *Hook. & Arn. Bot. Beech. Voy.* 428.—Mexic.

BOUZETIA, Montrous. in Mém. Acad. Lyon, x. (1860) 192. *RUTACEAE*?, Benth. & Hook. f. i. 989.
maritima, *Montrous. l. c.*—N. Caled.

BOVEA, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 253 = **Lindenbergia**, Lehm. (Scrophulac.).
sinaica, Decne. l. c. = L. sinaica.

BOWDICHIA, H. B. & K. Nov. Gen. et Sp. vi. 376 (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 557.
SEBIPIRA, Mart. Reise, 787 (1828).
brevipes, Benth. in Ann. Wien. Mus. ii. (1838) 89 = virgilioides.
densiflora, Benth. l. c. = virgilioides.
ferruginea, Walp. Rep. i. 808 = virgilioides.
floribunda, Benth. in Ann. Wien. Mus. ii. (1838) 89 = virgilioides.
major, Mart. Syst. Mat. Med. Bras. 53 = virgilioides.
nitida, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. i. 313.—Bras.
pubescens, Benth. in Ann. Wien. Mus. ii. (1838) 89 = virgilioides.
Sebipira, Steud. Nom. ed. II. i. 220 = virgilioides.
virgilioides, *H. B. & K. Nov. Gen. et Sp.* vi. 376.—Am. austr.

BOWENIA, Hook. Bot. Mag. t. 5398 (1863) et 6008. *CYCADACEAE*, Benth. & Hook. f. iii. 446.
spectabilis, *Hook. l. c.* 5398.—Austral.

BOWIEA, Harv. ex Hook. f. in Bot. Mag. t. 5619 (1867). *LILIACEAE*, Benth. & Hook. f. iii. 786.
volubilis, *Harv. l. c.*—Afr. austr.

BOWIEA, Haw. in Phil. Mag. lxiv. (1824) 299 = **Aloë**, Tourn. (Liliac.).
africana, Haw. l. c. = A. Bourea.
myriacantha, Haw. l. c.; et N. S. i. (1827) 122 = A. myriacantha.

BOWKERIA, Harv. Thes. Cap. i. 24. t. 37 (1859). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 937.
triphylla, *Harv. l. c.*—Afr. austr.

BOWLESIA, Ruiz & Pav. Prod. Fl. Per. 44. t. 34 (1794). *UMBELLIFERAE*, Benth. & Hook. f. i. 876.
DRUSA, DC. in Ann. Mus. Par. x. (1807) 466. t. 38.
ELSNERIA, Walp. Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 346. t. 8.

BOWLESIA:—
HOMALOCARPUS, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 348.
acutangula, *Benth. Pl. Hartw.* 186.—Peruv.
crenata, Desv. ex DC. Prod. iv. 70 = Dysopis Gayana.
dichotoma, *Poepp. ex DC. l. c.* 76.—Chili.
diversifolia, Meyen, ex Walp. Rep. v. 842 = lobata.
elata, *Clos, in C. Gay, Fl. Chil.* iii. 74.—Chili.
elegans, *Clos, l. c.* 74. t. 30.—Chili.
flexilis, *Meyen, Reise*, i. 348.—Chili.
geniculata, Spreng. Umbellif. 14. t. 5 = Angelica geniculata.
geraniifolia, *Schlecht. in Linnaea*, i. (1826) 382.—Chili.
incana, *Ruiz & Pav. Fl. Per.* iii. 28.—Peruv.
integerrima, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) 1. 170.—Chili.
lobata, *Ruiz & Pav. Prod.* 44. t. 34; *Fl. Per.* iii. 28.—Peruv.
multiradiata, *Colla, in Mem. Acc. Torino*, xxxvii. (1835) 81.—Chili.
nigripetala, *Clos, in C. Gay, Fl. Chil.* iii. 71.—Chili.
nodiflora, Presl, ex DC. Prod. iv. 75 = tenera.
oppositifolia, *Buch, in Abh. Acad. Berl.* 1816–17. (1819) 370.—Ins. Canar.
palmata, *Ruiz & Pav. Fl. Per.* iii. 28.—Peruv.
pulchella, *Wedd. Chlor. And.* ii. 188.—Peruv.; Boliv.
rotundifolia, *Phil. in Linnaea*, xxviii. (1856) 651.—Chili.
tenella, *Meyen, Reise*, i. 466.—Peruv.
tenera, *Spreng. Syst.* i. 880.—Bras.
tripartita, *Clos, in C. Gay, Fl. Chil.* iii. 76.—Chili.
tropaeolifolia, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 325.—Chili.
uncinata, *Colla, in Mem. Acc. Torino*, xxxvii. (1835) 81.—Chili.

BOWMANIA, Gardn. in Hook. Ic. Pl. 519 (1843); et in Hook. Lond. Journ. Bot. ii. (1843) 9 = **Trixis**, P. Br. (Compos.).
verbascifolia, Gardn. ll. cc. 519; 10 = T. Bowmanii.

BOWRINGIA, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 75. *LEGUMINOSAE*, Benth. & Hook. f. i. 553.
callicarpa, *Champ. ex Benth. l. c.*—Hongkong.

BOYKINIA, Nutt. in Journ. Acad. Philad. vii. (1834) 113. *SAXIFRAGEAE*, Benth. & Hook. f. i. 636.
THEROFON, Rafin. New Fl. Am. iv. 66 (1836).
aconitifolia, *Nutt. l. c.*—Am. bor.
humilis, Rafin. Aut. Bot. 9 ex S. Wats. Bibl. Ind. N. Am. Bot. i. 360 = Ammannia humilis.
major, *A. Gray, in S. Wats. Bot. Calif.* i. 196.—Calif.
occidentalis, *Torr. & Gray, Fl. N. Am.* i. 577.—Am. bor. occ.
ranunculifolia, A. Gray, in Am. Journ. Sc. Ser. I. xlii. (1842) 21 = Saxifraga ranunculifolia.
Richardsoni, *A. Gray, l. c.*—Am. bor. occ.
rotundifolia, *Parry, ex A. Gray, in Proc. Am. Acad.* xiii. (1878) 371.—Calif.
trispora, Nutt. ex Arn. in Hook. Journ. Bot. iii. (1840) 276 = Cayaponia Boykinii.

BOYMIA, A. Juss. in Mém. Mus. Par. xii. (1825) 507 (Mem. Rut. 124) = **Evodia**, Forst. (Rutac.).
glabrifolia, Champ. in Hook. Kew Journ. iii. (1851) 330 = E. meliaefolia.
martinicensis, G. Don, Gen. Syst. i. 805.—Ins. Martinic.
rutaecarpa, A. Juss. in Mém. Mus. Par. xii. (1825) 507 = E. rutaecarpa.

BRABEJARIA, Burm. f. Fl. Cap. Prod. 30, *err. typ.* 26 (1768) (Gen. dub.).
verticillata, Burm. f. l. c.—Afr. austr.

BRABEJUM, Linn. Gen. ed. I. 31 (1737). *PROTEACEAE*, Benth. & Hook. f. iii. 177.
BRABYLA, Linn. Mant. 137 (1767).
caliculatum, Blanco, Fl. Filip. ed. II. 39 = Loranthus Blancoanus.
concatenatum, Blanco, l. c. 40 = Alyxia laurina.
lucidum, Blanco, l. c. = Alyxia stellata.
pinnatum, Blanco, l. c. = Alyxia odorata.
stellatifolium, *Linn. Sp. Pl.* 121.—Afr. austr.
stellare, Knight, Prot. 98 = stellatifolium.

BRABEJUM:—
stellatum, Thunb. Prod. Pl. Cap. 31 = stellatifolium.
stellulifolium, Murr. Syst. Veg. ed. XIII. 764 = stellatifolium.

BRABYLA, Linn. Mant. 137 (1767) = **Brabejum**, Linn. (Proteac.).
capensis, Linn. l. c. = Brabejum stellatifolium.

BRACERA, Engelm. in Buchner, Rep. Pharm. lxviii. (1839) 369 = **Brayera**, Kunth (Rosac.).

BRACHANTHEMUM, DC. Prod. vi. 44 (1837) = **Chrysanthemum**, Tourn. (Compos.).
fruticulosum, DC. l. c. 45 = C. fruticulosum.

BRACHIATERA, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 = **Triodia**, R. Br. (Gramin.).
decumbens, Desv. l. c. = T. decumbens.

BRACHEILEMA, R. Br. in Salt, Abyss. App. 65 (1814) = **Vernonia**, Schreb. (Compos.).
paniculatum, R. Br. l. c. = V. amygdalina.

BRACHIARIA, Griseb. in Ledeb. Fl. Ross. iv. 469 (1853) = **Panicum**, Linn. (Gramin.).
cruciformis, Griseb. l. c. = P. eruciforme.
prostrata, Griseb. in Goett. Abh. vii. (1857) 263 = P. prostratum.

BRACHILOBOS et BRACHILOBUS, Desv. Journ. Bot. iii. 170 (1814) = seq.

BRACHIOLOBOS, All. Fl. Pedem. i. 278 (1785) = **Nasturtium**, R. Br. (Crucif.).
amphibius, All. l. c. = N. amphibium.
anceps, Link, Handb. ii. 311 = N. anceps.
armoracioides, Schur, Enum. Pl. Transs. 40 = N. armoracioides.
austriacus, Schur, l. c. 41 = N. austriacum.
barbaraefolius, Desv. Journ. Bot. iii. (1814) 170 = N. palustre.
brevistylus, Schur, Enum. Pl. Transs. 39 = N. brevistylon.
domingensis, Desv. Journ. Bot. iii. (1814) 183 = N. pyrenaicum.
hispidus, Desv. l. c. = N. palustre.
hybridus, Schur, Enum. Pl. Transs. 40 = N. austriacum.
lippizensis, Link, Handb. ii. 311 = N. lippizense.
palustris, [Clairv.] Man. Herb. 218; Schur, Enum. Pl. Transs. 39 = N. palustre.
pyrenaicus, All. Fl. Pedem. i. 278 = N. pyrenaicum.
riparius, Schur, Enum. Pl. Transs. 40 = N. amphibium.
sylvestris, All. Fl. Pedem. i. 278 = N. sylvestre.
tanacetifolius, Link, Handb. ii. 311 = N. tanacetifolium.

BRACHIOLOBUS, Bernh. Syst. Verz. Erf. 181 (1800) = praec.

BRACHIONIDIUM, Lindl. Fol. Orch. (1859). *ORCHIDEAE*, Benth. & Hook. f. iii. 491.
concolor, *Lindl. l. c.*—Bolivia.
parvifolium, *Lindl. l. c.*—N. Granat.
tuberculatum, *Lindl. l. c.*—N. Granat.

BRACHISTUS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. iii. (1849) 264. *SOLANACEAE*, Benth. & Hook. f. ii. 892.
FREGIRARDIA, Dun. Ind. Sem. Monsp. (1849). Cf. Ann. Sc. Nat. Sér. III. xii. (1849) 366.
SICKLERA, Sendtn. in Flora, xxix. (1846) 194 (err. typ. 178), nomen prius.
ciliatus, *Miers, l. c.* 263.—N. Granat.
dimorphus, *Miers, l. c.* 267.—N. Granat.
diversifolius, *Miers, l. c.* 268.—Mexic.
dumetorum, *Miers, l. c.* 265.—N. Granat.
hebephyllus, *Miers, l. c.* 266.—N. Granat.
Hookerianus, *Miers, l. c.* 268.—Ecuador.
lanceaefolius, *Miers, l. c.* 267.—Peruv.
Linnaeanus, Miers, l. c. 269 = Physalis arborescens.
macrophyllus, *Miers, l. c.* 263.—N. Granat.
mollis, *Miers, l. c.* 264.—Peruv.
Neesianus, Miers, l. c. 268 = Physalis arborescens.
oblongifolius, *Miers, l. c.* 266.—N. Granat.
rhomboideus, *Miers, l. c.* 264.—N. Granat.
riparius, *Miers, l. c.* 265.—N. Granat.
solanaceus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 424.—Nicaragua.
stramonifolius, *Miers, Illustr. S. Am. Pl.* ii. 7.—Mexic.

BRACHTIA, Reichb. f. in Linnaea, xxii. (1849) 853 *ORCHIDEAE*, Benth. & Hook. f. iii. 561.
ONCODIA, Lindl. Fol. Orch. (1853).
andina, *Reichb. f. in Bonplandia*, iv. (1856) 322.—N. Granat.
glumacea, *Reichb. f. in Linnaea*, xxii. (1849) 853.—N. Granat.
sulphurea, *Reichb. f. in Bonplandia*, ii. (1854) 14.—N. Granat.

BRACHYACHYRIS, Spreng. Syst. iii. 574 (1826) = BRACHYRIS, Nutt. = **Gutierrezia**, Lag. (Compos).
Euthamiae, Spreng. l. c. = G. Euthamiae.

BRACHYACTIS, Ledeb. Fl. Ross. ii. 495 (1846). *COMPOSITAE*, Benth. & Hook. f. ii. 279.
ciliata, Ledeb. l. c. = Aster angustus.
frondosa, A. Gray, in Proc. Am. Acad. viii. (1873) 647 = Aster frondosus.
gymnocephala, *Rupr. Sert. Tiansch.* 51.—As. centr.
iliensis, *Rupr. l. c.*—As. centr.
indica, C. B. Clarke, Compos. Ind. 49 = menthodora.
menthodora, *Benth. in Hook. Ic. Pl.* t. 1106.—Reg. Himal.
obovata, *Benth. l. c.* sub t. 1106.—Oriens.
pubescens, *Aitch. & Clarke, in Journ. Linn. Soc.* xviii. (1881) 68.—Ind. or.
robusta, *Benth. in Hook. Ic. Pl.* sub t. 1106.—Reg. Himal.
umbrosa, *Benth. l. c.*—Reg. Himal.; As. centr.
wangtuensis, Hook. f. Fl. Brit. Ind. iii. 253 = umbrosa.

BRACHYANDRA, Naud. in Ann. Sc. Nat. Sér. III. ii. 143 (1844) = **Pterolepis**, Miq. (Melastom.).
perpusilla, Naud. l. c. = P. perpusilla.
pusilla, Naud. l. c. xiii. (1844) 360 = P. perpusilla.

BRACHYANDRA, Phil. Fl. Atac. 34. t. 4 (1860). *COMPOSITAE*, Benth. & Hook. f. ii. 244.
macrogyne, *Phil. l. c.*—Chili.

BRACHYANTHES, Cham. ex Dunal, in DC. Prod. xiii. 1. 575 (1852) = **Petunia**, Juss. (Solanac.).
nierembergiensis, Cham. ex Dunal, l. c. = P. parviflora.

BRACHYBOTRYS, Maxim. ex Oliver, in Hook. Ic. Pl. xiii. 43. t. 1254 (1878). *BORAGINEAE*.
paridiformis, *Maxim. ex Oliver, l. c.*—China.

BRACHYCARPAEA, DC. in Mém. Mus. Par. vii. (1821) 247; et Syst. Veg. ii. 698 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 87.
laxa, *Sond. in Harv. & Sond. Fl. Cap.* i. 33.—Afr. austr.
linifolia, *Eckl. & Zeyh. Enum.* 8.—Afr. austr.
polygaloides, *Eckl. & Zeyh. l. c.* 7.—Afr. austr.
varians, *DC. Syst.* ii. 699.—Afr. austr.

BRACHYCENTRUM, Meissn. Gen. 114 (1837) = **Centronia**, D. Don (Melastom.).
excelsum, Meissn. Gen. Comm. 81 = C. excelsa.

BRACHYCHAETA, Torr. & Gray, Fl. N. Am. ii. 194 (1841). *COMPOSITAE*, Benth. & Hook. f. ii. 256.
cordata, *Torr. & Gray, l. c.* 195.—Am. bor.

BRACHYCHEILA, Harv. ex Eckl. & Zeyh. in Linnaea, xx. (1847) 192 = **Euclea**, Linn. (Ebenac.).
pubescens, Harv. l. c. = E. polyandra.

BRACHYCHITON, Schott & Endl. Meletem. 34 (1832) = **Sterculia**, Linn.
acerifolium, F. Muell. Fragm. i. 1 = S. accrifolia.
Bidwilli, Hook. Bot. Mag. t. 5133 = S. Bidwilli.
Delabechei, F. Muell. Pl. Vict. i. 157 = S. rupestris.
discolor, F. Muell. Fragm. i. 1 = S. discolor.
diversifolium, R. Br. in Benn. Pl. Jav. Rar. 234 = S. caudata.
Gregorii, F. Muell. in Hook. Kew Journ. ix. (1857) 199 = S. diversifolia.
incanum, R. Br. in Benn. Pl. Jav. Rar. 234 = S. incana.
luridum, C. Moore, ex F. Muell. Fragm. i. 1 = S. lurida.
paradoxum, Schott, Meletem. 34 = S. ramiflora.

BRACHYCHITON :—
platanoides, R. Br. in Benn. Pl. Jav. Rar. 234 = Sterculia trichosiphon.
populneum, R. Br. l. c. = S. diversifolia.
ramiflorum, R. Br. l. c. = S. ramiflora.

BRACHYCLADOS, D. Don, in Phil. Mag. (1832) 391. *COMPOSITAE*, Benth. & Hook. f. ii. 495.
lycioides, *D. Don, l. c.*—Chili.

BRACHYCODON, Prog. in Mart. Fl. Bras. vi. 1. 229. t. 42 (1865) = **Pagaea**, Griseb. (Gentian.).
pumilus, Prog. l. c. 230.—Bras.
ramosissimus, Prog. l. c.—Bras.

BRACHYCOME, Cass. in Dict. Sc. Nat. xxxvii. 491 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 264.
BRACHYSCOME, Cass. in Bull. Soc. Philom. (1816) 199.
BRACHYSTEPHIUM, Less. Syn. Comp. 388 (1832).
CTENOSPERMA, F. Muell. ex Pfeiff. Nom. i. 936 (1874).
PAQUERINA, Cass. in Dict. Sc. Nat. xxxvii. 492 (1825).
SILPHIOSPERMA, Steetz, in Lehm. Pl. Preiss. i. 433 (1844).
STEIROGLOSSA, DC. Prod. vi. 38 (1838).
aculeata, Less. Syn. Comp. 192 = Sieberi.
angustifolia, *A. Cunn. in DC. Prod.* v. (1834) 306.—Austral.
assamica, *C. B. Clarke, Compos. Ind.* 40.—Assam.
basaltica, *F. Muell. Fragm.* i. 50.—Austral.
bellidioides, Steetz, in Lehm. Pl. Preiss. i. 426 = pusilla.
Billardieri, *Benth. Fl. Austral.* iii. 518.—Austral.
calocarpa, *F. Muell. in Linnaea*, xxv. (1852) 399.—Austral.
capillacea, Walp. Rep. ii. 584 = iberidifolia.
cardiocarpa, *F. Muell. ex Benth. Fl. Austral.* iii. 517.—Austral.
cheilocarpa, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 172.—Austral.
chrysoglossa, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 44 = marginata.
ciliaris, *Less. Syn. Comp.* 192.—Austral.
collina, *Benth. Fl. Austral.* iii. 521.—Austral.
debilis, *Sond. in Linnaea*, xxv. (1852) 477.—Austral.
decipiens, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 114.—Austral.
discolor, *C. Stuart, ex Benth. Fl. Austral.* iii. 520.—Austral.
diversifolia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 31.—Austral.
Drummondii, Walp. Rep. ii. 584 = ciliaris.
exilis, *Sond. in Linnaea*, xxv. (1852) 473.—Austral.
glabra, Benth. in Enum. Pl. Hueg. 59 = multifida.
glandulosa, *Benth. Fl. Austral.* iii. 520.—Austral.
glauca, Walp. in Linnaea, xiv. (1840) 315 = stricta.
goniocarpa, *Sond. & F. Muell. in Linnaea*, xxv. (1852) 474.—Austral.
graminea, *F. Muell. Fragm.* i. 49.—Austral.
heterodonta, *DC. Prod.* v. 305.—Austral.
heterophylla, *Benth. in Enum. Pl. Hueg.* i. 50.—Austral.
iberidifolia, *Benth. l. c.* 59.—Austral.
lanuginosa, Steetz, in Lehm. Pl. Preiss. i. 427 = ciliaris.
latisquamea, *F. Muell. Fragm.* xi. 16.—Austral.
leptocarpa, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 43 = exilis.
leucanthemifolia, Benth. in Enum. Pl. Hueg. 60 = stricta.
linearifolia, *DC. Prod.* v. 306.—Austral.
linearifolia, Hook. f. Fl. Tasm. i. 185 = cardiocarpa.
marginata, *Benth. in Enum. Pl. Hueg.* 60.—Austral.
melanocarpa, *F. Muell. & Sond. in Linnaea*, xxv. (1852) 476.—Austral.
mexicana, Klatt, in Leopoldina, xix. (1883) 32 = Bellis mexicana.
microcarpa, *F. Muell. Fragm.* i. 50.—Austral.
mossambicensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 305.—Afr. trop.
Muelleri, *Sond. in Linnaea*, xxv. (1852) 475.—Austral.
multicaulis, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 43 = ciliaris.
multifida, *DC. Prod.* v. 306.—Austral.

BRACHYCOME :—
nivalis, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 43.—Austral.
oblongifolia, Benth. in Enum. Pl. Hueg. 60 = stricta.
odorata, *Hook. f. Handb. N. Zeal. Fl.* 138.—N. Zel.
pachyptera, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) 1. 175.—Austral.
parvula, *Hook. f. Fl. Tasm.* i. 185.—Austral.
pinnata, *Hook. f. Handb. N. Zeal. Fl.* 1381.—N. Zel.
ptychocarpa, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 43.—Austral.
pumilà, Walp. Rep. ii. 584 = Billardieri.
pusilla, *Steetz, in Lehm. Pl. Preiss.* i. 427.—Austral.
radicans, *Steetz, l. c.* 429.—Austral.
radicata, *Hook. f. Fl. N. Zel.* i. 127.—N. Zel.
scapiformis, *DC. Prod.* v. 306.—Austral.
scapigera, *DC. l. c.* viii. 277.—Austral.
segmentosa, *F. Muell. Fragm.* viii. 65; 144.—Ins. Dom. Howe.
Sieberi, *DC. Prod.* v. 306.—Austral.
simplicifolia, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 338.—N. Zel.
Sinclairii, *Hook. f. Handb. N. Zeal. Fl.* 137.—N. Zel.
squalida, Hook. f. in Hook. Lond. Journ. Bot. v. (1846) 115 = ciliaris.
stricta, *DC. Prod.* v. 305.—Austral.
strongylospermoides, Walp. in Linnaea, xiv. (1840) 305 = ciliaris.
Stuartii, *Benth. Fl. Austral.* iii. 513.—Austral.
sylvestris, Klatt, in Leopoldina, xix. (1883) 32 = Bellis sylvestris?
tenella, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) 1. 176 = collina.
tenera, Benth. in Enum. Pl. Hueg. 59 = multifida.
tenuiscapa, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 114 = scapiformis.
Thomsonii, *T. Kirk, in Trans. N. Z. Inst.* xvi. (1884) 372.—N. Zel.
trachycarpa, *F. Muell. in Linnaea*, xxv. (1852) 339.—Austral.
xanthocomoides, Less. in Linnaea, ix. (1834) 265 = Bellis xanthocomoides.
xeranthemoides, Less. ex Steud. Nom. ed. II. i. 220 = Xanthocephalum Alamani.

BRACHYCOME, Gaudich. Freyc. Voy. Bot. 467 (1826) = **Vittadinia**, Rich. (Compos.).
dentata, Gaudich. l. c. 468 = V. australis?
spathulata, Gaudich. l. c. = V. australis.
triloba, Gaudich. l. c. 467 = V. australis.

BRACHYCORYS, Schrad. Cat. Hort. Goetting. (1830).—Cf. Linnaea, vi. (1831) Litt. 72 = **Lindenbergia**, Linn. (Scrophular.).
parviflora, Schrad. l. c. = L. urticaefolia.

BRACHYCORYTHIS, Lindl. Gen. et Sp. Orch. 363 (1838). *ORCHIDEAE*, Benth. & Hook. f. iii. 632.
Bulbinella, *Reichb. f. in Flora*, l. (1867) 116 (sphalm. 161).—Afr. austr.
Gerardi, *Reichb. f. l. c.*—Afr. austr.
Kalbreyeri, *Reichb. f. l. c.* lxi. (1878) 77.—Afr. trop.
Mac-Owaniana, *Reichb. f. Otia Bot. Hamb.* 104.—Afr. austr.
pleistophylla, *Reichb. f. l. c.*—Afr. trop.
ovata, *Lindl. Gen. et Sp. Orch.* 363.—Afr. austr.
pubescens, *Harv. Thes. Cap.* i. 35.—Afr. austr.
Pumilio, *Reichb. f. in Flora*, lxv. (1882) 531.—Afr. trop.
Schweinfurthii, *Reichb. f. Otia Bot. Hamb.* 59.—Afr. trop.
tenuior, *Reichb. f. in Flora*, xlviii. (1865) 183.—Afr. trop.
Tysoni, *Bolus, in Journ. Linn. Soc.* xx. (1884) 485.—Afr. austr.
Welwitschii, *Reichb. f. in Flora*, l. (1867) 99.—Afr. trop.
Zeyheri, Reichb. f. l. c. 117 = Schizochilus Zeyheri.

BRACHYDEREA, Cass. in Dict. Sc. Nat. xlviii. 429 (1827) = **Crepis**, Linn. (Compos.).
alpestris, Sch. Bip. ex Nym. Consp. 455 = C. alpestris.
aurea, Sch. Bip. in Pollichia, xxii–iv. (1866) 318 = C. aurea.
biennis, Sch. Bip. l. c. = C. biennis.

BRACHYDEREA :—
chondrilloides, Sch. Bip. ex Nym. Consp. 456 = Crepis chondrilloides.
grandiflora, Sch. Bip. ex Nym. l. c. 455 = C. grandiflora.
heterantha, Sch. Bip. in Pollichia, xxii–iv. (1866) 318 = Lactuca Dubyaea.
Jacquinii, Sch. Bip. l. c. = C. Jacquinii.
jubata, Sch. Bip. l. c. = C. jubata.
nicaeënsis, Sch. Bip. l. c. = C. nicaeënsis.
rigida, Cass. in Dict. Sc. Nat. xlviii. 430 = C. rigida.
Rueppellii, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 283 = C. Rueppellii.
Schultzii, Sch. Bip. l. c. = C. Schultzii.
tenerrima, Sch. Bip. l. c. = C. carbonaria.
tenuifolia, Sch. Bip. in Bonplandia, x. (1862) 331 = C. tenuifolia.
xylorrhiza, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 284.—Afr. trop.

BRACHYELYTRUM, Beauv. Agrost. 39. t. 9. f. 2 (1812). *GRAMINEAE*, Benth. & Hook. f Gen. iii. 1144.
aristatum, Roem. & Schult. Syst. ii. 413 = erectum.
erectum, *Beauv. Agrost.* 39, 155.—Am. bor.

BRACHYGLOTTIS, Forst. Char. Gen. 91. t. 46 (1776). *COMPOSITAE*, Benth. & Hook. f. ii. 444.
Bidwillii, Hook. f. Fl. N. Zel. ii. 342 = Senecio Bidwillii.
Caledoniae, *DC. Prod.* v. 211.—N. Caled.
elaeagnifolia, Hook. f. Fl. N. Zel. t. 41 = Senecio elaeagnifolius.
Forsteri, *DC. Prod.* v. 211.—N. Hebrid.
Forsteri, Hook. f. Fl. N. Zel. ii. 342 = repanda.
perdicioides, Hook. f. l. c. = Senecio perdicioides.
Rani, A. Cunn. in Ann. Nat. Hist. ii. (1839) 132 = Olearia Cunninghami.
Rangiora, *J. Buch. in Trans. N. Z. Inst.* xiv. (1882) 357.—N. Zel.
repanda, *Forst. Char. Gen.* 92.—N. Zel.
rotundifolia, Forst. l. c. = Senecio rotundifolius.

BRACHYGYNE, Cass. in Dict. Sc. Nat. l. (1827) 493 = **Eriocephalus**, Linn. (Compos.).

BRACHYLAENA, R. Br. in Trans. Linn. Soc. xii. (1816) 115. *COMPOSITAE*, Benth. & Hook. f. ii. 288.
OLIGOCARPHA, Cass. in Bull. Soc. Philom. (1817) 151.
dentata, *DC. Prod.* v. 430.—Afr. austr.
discolor, *DC. l. c.*—Afr. austr.
elliptica, *Less. Syn. Comp.* 208.—Afr. austr.
grandifolia, *DC. Prod.* v. 430.—Afr. austr.
natalensis, Sch. Bip. ex Walp. Rep. ii. 972 = discolor.
nereifolia, *R. Br. in Trans. Linn. Soc.* xii. (1816) 115.—Afr. austr.
racemosa, *Less. Syn. Comp.* 208.—Afr. austr.
trinervia, *Sond. in Linnaea*, xxiii. (1850) 63.—Afr. austr.
uniflora, *Harv. in Harv. & Sond. Fl. Cap.* iii. 117.—Afr. austr.

BRACHYLEPIS, Wight & Arn. Contrib. Bot. Ind. 63 (1834). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 742.
nervosa, *Wight & Arn. l. c.*—Ind. or.

BRACHYLEPIS, C. A. Mey. in Ledeb. Fl. Alt. i. 370 (1829) = **Anabasis**, Linn. (Chenopod.).
elatior, C. A. Mey. in Bull. Sc. Acad. Pétersb. viii. (1840) 341 = A. phyllophora.
eriopoda, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 360 = A. phyllophora.
hispidula, Bunge, in Act. Hort. Petrop. vi. (1879) 444.—Turkestan.
humilis, Less. in Linnaea, ix. (1834) 198 = salsa.
intermedia, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 432 = A. phyllophora.
jaxartica, Bunge, in Act. Hort. Petrop. vi. (1879) 443.—Reg. Casp.
salsa, C. A. Mey. in Ledeb. Fl. Alt. i. 372.—Reg. Casp.
truncata, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. ii. (1844) 193 = A. truncata.

BRACHYLEPIS, Hook. & Arn. in Hook. Journ. Bot. (1834) 290 = **Melinia**, Decne. (Asclep.).
Candolleana, Hook. & Arn. l. c. 291 = M. Candolleana.

BRACHYLOBOS, Desv. Journ. Bot. iii. (1814) 170 = seq.

BRACHYLOBUS, Link, Handb. vii. 311 (1831) = BRACHIOLOBOS, All. = **Nasturtium**, R. Br. (Crucif.).

BRACHYLOBUS, Dulac, Fl. Hautes-Pyr. 279 (1867) = **Melilotus**, Tourn. (Legumin.).
officinalis, Dulac, l. c. = M. officinalis.

BRACHYLOMA, Hanst. in Linnaea, xxvi. (1853) 203 = **Isoloma**, Decne. (1848) (Gesneriac.).
elongatum, Hanst. l. c. xxix. (1857–58) 532.—Ecuador?
erianthum, Hanst. l. c. 530 = I. erianthum.
hirsutum, Hanst. in Linnaea, xxvi. (1853) 203; xxix. (1857–58) 526 = I. hirsutum.
hondense, Hanst. l. c. xxvi. (1853) 203, 213. t. f. 17 = I. hondense.
incurvum, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 105 = I. incurvum.
Karstenianum, Hanst. in Karst. Fl. Colomb. i. 101. t. 50.—N. Granat.
leucomallon, Hanst. in Linnaea, xxix. (1857–58) 528.—N. Granat.
longiflorum, Hanst. l. c. 530 = I. longipes.
longifolium, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 104 = I. longifolium.
longipes, Hanst. in Linnaea, xxix. (1857–58) 530 = I. longipes.
molle, Hanst. l. c. 528 = I. molle.
Moritzianum, Bouché & Hanst. l. c. 524.—N. Granat.
petiolare, Hanst. l. c. 534 = I. petiolare.
pictum, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 104 = I. pictum.
pilosum, Oerst. l. c. 103 = I. pilosum.
rhodomallon, Hanst. in Linnaea, xxix. (1857–58) 526.—Venezuela.
rhynchocarpum, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 103 = I. rhynchocarpum.
rubricaule, Hanst. in Linnaea, xxix. (1857–58) 526 = I. rubricaule.
strictum, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 103 = I. strictum.
Trianae, Hanst. in Linnaea, xxix. (1857–58) 532 = I. Trianaei.
tubiflora, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 105 = I. tubiflorum.
ventricosum, Hanst. in Linnaea, xxix. (1857–58) 528.—Hab.?
vestitum, Hanst. l. c. 530 = I. vestitum.

BRACHYLOMA, Sond. in Lehm. Pl. Preiss. i. 304 (1845). *EPACRIDEAE*, Benth. & Hook. f. ii. 613.
ciliatum, *Benth. Fl. Austral.* iv. 173.—Austral.
concolor, *F. Muell. Fragm.* vi. 39.—Austral.
daphnoides, *Benth. Fl. Austral.* iv. 173.—Austral.
depressum, *Benth. l. c.*—Austral.
ericoides, *Sond. in Linnaea*, xxvi. (1853–55) 247.—Austral.
Preissii, *Sond. in Lehm. Pl. Preiss.* i. 305.—Austral.
Scortechinii, *F. Muell. Fragm.* xi. 121.—Austral.

BRACHYMERIS, DC. Prod. vi. 76 (1837) = **Marasmodes**, DC. (Compos.).
scoparia, DC. l. c.—Afr. austr.

BRACHYNEMA, Benth. in Trans. Linn. Soc. xxii. (1859) 125. t. 22. *EBENACEAE*, Benth. & Hook. f. ii. 666.
ramiflorum, *Benth. l. c.* 126.—Bras.

BRACHYNEMA, Griff. Notul. iv. 176 (1854) = **Sphenodesma**, Jack (Verben.).
ferruginea, Griff. l. c. = S. microstylis.

BRACHYNEMA, F. Muell. Fragm. iii. 90 (1862) = **Abrophyllum**, Hook. f. (Saxifrag.).
ornans, F. Muell. l. c. = A. ornans.

BRACHYOGLOTIS, Lam. Encyc. i. 459 (1783) = **Brachyglottis**, Forst. (Compos.).

BRACHYOLOBOS, DC. Fl. Fr. v. 661 (1805) = BRACHIOLOBOS, All. = **Nasturtium**, R. Br. (Crucif.).

BRACHYOTUM, Triana, in Benth. & Hook. f. Gen. i. 743 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
ALIFANA, Rafin. Sylva Tellur. 101 (1838).
Benthamianum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 49.—Peruv.
campanulare, *Triana, l. c.* 48.—Peruv.
campylanthum, *Triana, l. c.* 49.—Peruv.
canescens, *Triana, l. c.* 48.—N. Granat.
cernuum, *Triana, l. c.*—N. Granat.
confertum, *Triana, l. c.* 49.—Peruv.
floribundum, *Triana, l. c.*—Peruv.
gracilescens, *Triana, l. c.*—Peruv.
hermannioides, *Triana, l. c.*—Bolivia.
Jamesonii, *Triana, l. c.*—Ecuador.
lediifolium, *Triana, l. c.* 48.—Peruv.
lutescens, *Triana, l. c.*—Peruv.
lycopodioides, *Triana, l. c.* 49.—Peruv.
Microdon, *Triana, l. c.*—Bolivia.
microphyllum, *Triana, l. c.*—Peruv.
Naudini, *Triana, l. c.* 48.—Ecuador.
Pentlandi, *Triana, l. c.* 49.—Bolivia.
quinquenerve, *Triana, l. c.* 48.—Peruv.
Radula, *Triana, l. c.*—Peruv.
rosmarinifolium, *Triana, l. c.* 49.—Peruv.
rostratum, *Triana, l. c.* 48.—Peruv.
sanguinolentum, *Triana, l. c.* 49.—Bolivia.
strigosum, *Triana, l. c.*—N. Granat.
trichocalyx, *Triana, l. c.* 48.—Ecuador.

BRACHYPAPPUS, Sch. Bip. in Flora, xxxviii. (1855) 119 = **Senecio**, Tourn. (Compos.).
candicans, Sch. Bip. l. c. 120 = S. candicans.
Smithii, Sch. Bip. l. c. = S. Smithii.

BRACHYPODIUM, Beauv. Agrost. 100 partim (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1201.
DISTICHEIA, Ehrh. Beitr. iv. 148 (1789).
HEMIBROMUS, Steud. Syn. Pl. Gram. 317 (1854).
TRACHYNIA, Link, Hort. Berol. i. 42 (1827).
abbreviatum, *Dum. Obs. Gram. Belg.* 99.—Belgium.
Allionii, Presl, Fl. Sic. p. xlv, in nota = ramosum.
asperum, Roem. & Schult. Syst. ii. 742 = distachyon.
Barrelieri, Roem. & Schult. l. c. 741 = pinnatum.
biflorum, Kunth, Enum. Pl. i. 543 (Ind.) = Agropyron biflorum.
biunciale, Roem. & Schult. Syst. ii. 746 = Festuca Poa.
Boissieri, Nym. Syll. 425 = ramosa.
brevisetum, Roem. & Schult. Syst. ii. 742 = Triticum brevisetum.
caespitosum, Presl, Fl. Sic. p. xlv, nomen = ramosum.
caespitosum, Roem. & Schult. Syst. ii. 737 = pinnatum.
cenisium, Beauv. Agrost. 155, nomen = Poa cenisia.
chinense, *S. L. Moore, in Journ. Bot.* xiii. (1875) 230.—China.
ciliatum, Beauv. Agrost. 101 = distachyum.
commutatum, Beauv. l. c. = Bromus racemosus.
contractum, Presl, Cyp. et Gram. Sic. 40 = pinnatum.
corniculatum, Lam. Fl. Fr. v. 608 = pinnatum.
dalmaticum, *Vis. Fl. Dalm. Suppl.* 23. t. 1.—Europ. austr.
distachyum, *Beauv. Agrost.* 101.—Reg. Mediterr.; Oriens.
distachyon, Roem. & Schult. Syst. ii. 741 = praec.
divaricatum, Presl, Cyp. et Gram. Sic. 40 = Festuca divaricata.
Festuca, *Beauv. Agrost.* 101.—Hab.?
festucoides, Beauv. l. c. 155, nomen = Festuca Poa.
festucoides, Link, Enum. Hort. Berol. i. 95 = Festuca tenella.
flexum, *Nees, Fl. Afr. Gram.* 456.—Afr. austr.
Fontanesianum, Nees, l. c. 457 = ramosum.
fragile, Beauv. Agrost. 155, nomen = Triticum fragile.
geniculatum, C. Koch, in Linnaea, xxi. (1848) 422 = distachyum.
genuense, Roem. & Schult. Syst. ii. 742 = Triticum genuense.
gracile, Beauv. Agrost. 101 = sylvaticum.
Halleri, Beauv. l. c. = Festuca Poa.
hispanicum, Reichb. Fl. Germ. Excurs. 140 = Festuca Poa.
Hostii, Link, Hort. Berol. i. 40 = ramosum.
intermedium, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 89.—Transsylv.
interruptum, Roem. & Schult. Syst. ii. 747 = Apera interrupta.
involutum, *Buese, in Miq. Pl. Jungh.* 347.—Malaya.
japonicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 286.—Japon.

BRACHYPODIUM:—
Kotschyi, *Boiss. Fl. Orient.* v. 659.—Cilicia.
latifolium, *Fourn. Mex. Pl. Enum. Gram.* 125; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 584.—Mexic.
littorale, Roem. & Schult. Syst. ii. 740 = ramosum.
loliaceum, Fries, Mant. iii. 15 = Glyceria fluitans.
loliaceum, Link, Hort. Berol. i. 42 = Festuca elatior.
loliaceum, Roem. & Schult. Syst. ii. 740 = Demazeria loliacea.
longearistatum, Boiss. Diagn. Sér. I. vii. 127 = Agropyron longearistatum.
longifolium, Beauv. Agrost. 101 = Bromus longifolius.
macropodum, *Hack. in Oestr. Bot. Zeitschr.* xxvii. (1877) 48.—Europ. austr.
macrostachyum, Bess. ex Roem. & Schult. Syst. Mant. iii. 651 = distachyum.
maritimum, Roem. & Schult. Syst. ii. 743 = Cutandia maritima.
megastachyum, Bess. ex Roem. & Schult. Syst. Mant. iii. 651 = distachyum.
mexicanum, *Link, Hort. Berol.* i. 41.—Mexic.
monostachyum, Nym. Consp. 843 = distachyum.
montanum, Nym. Syll. 426 = Festuca montana.
mucronatum, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* i. 111.—Hispan.
Nardus, Beauv. Agrost. 155, nomen = Festuca tenuiflora.
Nardus, Link, Enum. Hort. Berol. i. 95 = Festuca Poa.
nepalense, *Nees, ex Steud. Syn. Pl. Gram.* 316.—Ind. or.
nigricans, Roem. & Schult. Syst. ii. 743 = Triticum nigricans.
obtusifolium, Link, Enum. Hort. Berol. i. 95 = ramosum.
patens, Nym. Syll. 425 = Festuca loliacea.
pentastachyum, Nym. Consp. 843 = distachyum.
peruvianum, Kunth, Enum. Pl. 543 (Ind.) = Agropyron peruvianum.
phleoides, Beauv. Agrost. 155, nomen = Koeleria phleoides.
phoenicoides, Roem. & Schult. Syst. ii. 740 = ramosum.
pinnatum, *Beauv. Agrost.* 101.—Europ.; Reg. Mediterr.
Plukenetii, Beauv. l. c.; Link, Hort. Berol. i. 40 = ramosum.
Poa, Beauv. l. c.; Roem. & Schult. Syst. ii. 746 = Festuca Poa.
psilanthum, Link, Gram. Europ. Austr. Obs. Bot. 398 = Festuca tenuiflora.
ramosum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 118 = Glyceria maritima?
ramosum, *Roem. & Schult. Syst.* ii. 737.—Reg. Mediterr.
retusum, Beauv. Agrost. 101 = ramosum.
rigidulum, *Opiz & Bercht. Oek. Fl. Boehm.* i. 392.—Bohem.
rigidum, Link, Enum. Hort. Berol. i. 96 = Triticum asperum.
Rottboëlla, Beauv. Agrost. 155, nomen = ramosum.
rupestre, Roem. & Schult. Syst. ii. 736 = pinnatum.
sanctum, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 181.—Macedon.
scaberrimum, Wight & Arn. ex Steud. Nom. ed. II. i. 221 = Triticum scaberrimum.
Schrenkianum, *Ledeb. Fl. Ross.* iv. 346.—Sibiria.
setifolium, *Schur, Enum. Pl. Transs.* 800.—Hungar.
strigosum, Schult. in Roem. & Schult. Syst. Mant. ii. 404 = Triticum strigosum.
subtile, Dc Not. Prosp. Fl. Ligust. 49 = distachyum.
subulatum, *Fourn. Mex. Pl. Enum. Gram.* 125; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 584.—Mexic.
sylvaticum, *Beauv. Agrost.* 101.—Europ.; Oriens.
tataricum, Munro, ex Aitch. in Journ. Linn. Soc. (1880) 109 = Agropyron longearistatum.
tenellum, Beauv. Agrost. 101 = Festuca tenuiflora.
tenellum, Roem. & Schult. Syst. ii. 745 = Festuca montana?
Tenorianum, Schult. in Roem. & Schult. Syst. Mant. ii. 405 = Triticum Barrelieri.
tenue, Beauv. Agrost. 155, nomen = Lolium perenne.
tenue, Tineo, Cat. Hort. Panorm. (1827) 48 = Festuca incrassata.
tenuiculum, Roem. & Schult. Syst. ii. 744 = Festuca Poa.
tenuiflorum, Roem. & Schult. l. c. 745 = Festuca tenuiflora.
unilaterale, Beauv. Agrost. 155; Roem. & Schult. l. c. 747 = Triticum tenuiflorum.
unioloides, Link, Enum. Hort. Berol. i. 195 = Demazeria sicula.

BRACHYPTERUM, Benth. in Ann. Wien. Mus. ii. (1838) 101 = **Derris**, Lour. (Legumin.).
Benthamii, Thw. Enum. Pl. Zeyl. 93 = D. paniculata.
canarense, Dalz. & Gibs. Bomb. Fl. 76 = D. canarensis.
elegans, Thw. Enum. Pl. Zeyl. 93 = D. parviflora.
floribundum, Miq. Fl. Ind. Bat. i. 139 = D. floribunda.
microphyllum, Miq. Fl. Ind. Bat. Suppl. 296 = D. microphylla.
polyphyllum, Miq. Fl. Ind. Bat. i. 139 = D. polyphylla.
robustum, Dalz. & Gibs. Bomb. Fl. 77 = D. robusta.
scandens, Benth. in Ann. Wien. Mus. ii. (1838) 101 = D. scandens.
timorense, Benth. in Miq. Pl. Jungh. i. 255 = D. timorensis.

BRACHYPTERYS, A. Juss. in Deless. Ic. Sel. iii. 20 (1837). *MALPIGHIACEAE*, Benth. & Hook. f. i. 256.
australis, *A. Juss. l. c.*—Bras.
borealis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 356. t. 2. f. 16.—Am. centr.

BRACHYPUS, Ledeb. Fl. Ross. i. 133 (1842) = **Lunaria**, Tourn. (Crucif.).
asper, Ledeb. l. c. (= *Fibigia suffruticosa*).—Rossia.

BRACHYRAMPHUS, DC. Prod. vii. 176 (1838) = **Lactuca**, Tourn. (Compos.).
caribaeus, DC. l. c. 177 = L. intybacea.
goraeënsis, DC. l. c. = L. goraeënsis.
Heyneanus, Wight, Ic. t. 1146 = L. Heyneana.
intybaceus, DC. Prod. vii. 177 = L. intybacea.
lactucoides, T. Anders. in Journ. Linn. Soc. Bot. v. Suppl. i. (1860) 23 = Heterachaena massaviensis.
nudicaulis, Klatt, in Ann. Sc. Nat. Sér. V. xviii. (1873) 377 = Launaea nudicaulis.
obtusus, DC. in Deless. Ic. Sel. iv. t. 96 = Launaea nudicaulis.
ramosissimus, Benth. in Hook. Journ. Bot. i. (1834) 489 = L. denticulata.
sinicus, Miq. in Journ. Bot. Néerl. i. (1861) 105 = L. denticulata.
sonchifolius, DC. Prod. vii. 177 = L. remotiflora.
sonchifolius, Thw. Enum. Pl. Zeyl. 168 = L. Heyneana.
taraxacoides, DC. Prod. vii. 177 = L. Heyneana.

BRACHYRHYNCHOS, Less. Syn. Comp. 392 (1832) = **Senecio**, Tourn. (Compos.).
albicaulis, DC. Prod. vi. 438 = S. diversifolius, *Harv.*
cymbalarifolius, DC. l. c. = S. cymbalarifolius.
diversifolius, DC. l. c. 439 = S. diversifolius, *Harv.*
elongatus, Less. Syn. Comp. 393 = S. diversifolius, *Harv.*
eupatorioides, DC. Prod. iv. 439 (non seq.).—Afr. austr.
eupatorioides, Less. Syn. Comp. 393 = S. othonnaeflorus.
junceus, Less. l. c. = S. junceus.
trachycarpus, DC. Prod. vi. 438 = S. incertus.
tuberosus, DC. l. c. = S. incertus.

BRACHYRIS, Nutt. Gen. Am. ii. 163 (1818) = **Gutierrezia**, Lag. (Compos.).
californica, DC. Prod. v. 313 = G. californica.
divaricata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 313 = G. Euthamiae.
dracunculoides, DC. in Mém. Soc. Phys. Genèv. vii. (1836) II. 265. t. 1 = Amphiachyris dracunculoides.
Euthamiae, Nutt. Gen. Am. ii. 163 = G. Euthamiae.
floribunda, Phil. in Linnaea, xxxiii. (1864-65) 137.—Chili.
Gayana, Remy, in C. Gay, Fl. Chil. iv. 34.—Chili.
Iserni, Phil. in Anal. Univ. Chil. (1865) II. 335.—Chili.
Mandonii, Sch. Bip. in Linnaea, xxxiv. (1865-66) 534.—Bolivia.
microcephala, DC. Prod. v. 313 = G. Euthamiae.
microcephala, Hook. Ic. Pl. t. 147 = G. texana.
Neaeana, DC. Prod. v. 313.—Mexic.
ovatifolia, DC. l. c. = Brachychaeta cordata.
paniculata, DC. l. c. = G. paniculata.
ramosissima, Hook. Ic. Pl. t. 142 = Amphiachyris dracunculoides.
sectifolia, Ces. in Atti Soc. Acc. Sc. Nap. v. (1871) VII. 11.—Chili.
spathulata, Phil. in Anal. Univ. Chil. (1865) II. 336.—Chili.

BRACHYSCOME, Cass. in Bull. Soc. Philom. (1816) 199.—Cf. Dict. Sc. Nat. v. Suppl. 63 (1817) = **Brachycome**, Cass. (Compos.).
Billardieri, Cass. in Dict. Sc. Nat. v. Suppl. 63 = Brachycome heterodonta.

BRACHYSCYPHA, Baker, in Journ. Linn. Soc. xi. (1871) 393 = **Lachenalia**, Jacq. (Liliac.).
undulata, Baker, l. c. 394 = L. pusilla.

BRACHYSEMA, R. Br. in Ait. Hort. Kew. ed. II. iii. 10 (1811). *LEGUMINOSAE*, Benth. & Hook. f. i. 467.
BURGESIA, F. Muell. Fragm. i. 222 (1860 ?).
KALENICZENKIA, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) I. 252.
LEPTOSEMA, Benth. in Ann. Wien. Mus. ii. (1838) 84.
acuminatum, *Truff. in Rev. Hortic.* (1866) 413.—Hab. ?
aphyllum, *Hook. Bot. Mag.* t. 4481.—Austral.
bossiaeoides, *Benth. Fl. Austral.* ii. 12.—Austral.
bracteolosum, *F. Muell. Fragm.* iv. 10.—Austral.
Celsianum, Lem. Hortic. Univ. v. n. 2. 33, ex Walp. Rep. v. 423 = lanceolatum.
Chambersii, *F. Muell. ex Benth. Fl. Austral.* ii. 13.—Austral.
daviesoides, *Benth. l. c.*—Austral.
lanceolatum, *Meissn. in Lehm. Pl. Preiss.* i. 24.—Austral.
latifolium, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 10.—Austral.
macrocarpum, *Benth. Fl. Austral.* ii. 13.—Austral.
melanopetalum, F. Muell. Fragm. iv. 11 = undulatum.
oxyloboides, *Benth. Fl. Austral.* ii. 12.—Austral.
praemorsum, *Meissn. in Lehm. Pl. Preiss.* i. 25.—Austral.
subcordatum, *Benth. Fl. Austral.* ii. 11.—Austral.
tomentosum, *Benth. l. c.* 13.—Austral.
undulatum, *Ker-Gawl. in Bot. Reg.* t. 642.—Austral.
uniflorum, *R. Br. ex Benth. Fl. Austral.* ii. 12.—Austral.

BRACHYSIPHON, A. Juss. in Ann. Sc. Nat. Sér. III. vi. (1846) 24. t. 2 = **Sarcocolla**, Linn. (Penaeac.).
acutus, A. Juss. l. c. 25 = S. acuta.
ericaefolius, A. Juss. l. c. 24.—Afr. austr.
imbricatus, A. Juss. l. c. 25 = S. imbricata.
Mundii, Sond. in Linnaea, xxiii. (1850) 102.—Afr. austr.
rupestris, Sond. l. c. 101 = S. rupestris.
speciosus, Sond. l. c. 103 = S. speciosa.

BRACHYSPATHA, Schott, Syn. Aroid. 35 (1856) = **Amorphophallus**, Blume (Aroid.).
consimilis, Schott, Prod. Aroid. 127 = Hydrosme consimilis.
Konjac, C. Koch, in C. Koch & Fint. Wochenschr. (1858); Gart. Nachr. n. 4. p. 13 = A. Rivieri.
Muelleri, Schott, Syn. Aroid. 36 = A. Mulleri.
sylvatica, Schott, l. c. 35 = Synantherias sylvatica.
variabilis, Schott, l. c. = A. variabilis.
zeylanica, Schott, l. c. 36 = A. zeylanicus.

BRACHYSTACHYS, Klotzsch, in Hook. Lond. Journ. Bot. ii. (1843) 47 = **Croton**, Linn. (Euphorb.).
hirta, Klotzsch, l. c. = C. glandulosus.

BRACHYSTEGIA, Benth. in Benth. & Hook. f. Gen. i. 582 (1865). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
appendiculata, *Benth. in Trans. Linn. Soc.* xxv. (1866) 313. t. 42.—Afr. trop.
floribunda, *Benth. in Hook. Ic. Pl.* snb t. 1359.—Afr. trop.
globiflora, *Benth. l. c.*—Afr. trop.
longifolia, *Benth. l. c.* t. 1359.—Afr. trop.
spicaeformis, *Benth. in Trans. Linn. Soc.* xxv. (1866) 312.—Afr. trop.
tamarindoides, *Welw. ex Benth. l. c.*—Afr. trop.

BRACHYSTELMA, R. Br. in Bot. Mag. t. 2343 (1822). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 781.
ERIOPETALUM, Wight, Contrib. Bot. Ind. 35 (1834).
MICRASTER, Harv. Gen. Pl. Afr. Austr. ed. II. 242 (1868).
Arnottii, *Baker, in Saund. Ref. Bot.* t. 9.—Afr. austr.

BRACHYSTELMA:—
attenuatum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
Barberiae, *Harv. ex Hook. f. Bot. Mag.* t. 5607.—Afr. austr.
Beddomei, *Hook. f. Fl. Brit. Ind.* iv. 66.—Ind. or.
caudatum, N. E. Br. in Journ. Linn. Soc. xvii. (1880) 169 = crispum.
circinnatum, *E. Mey. Comm. Pl. Afr. Austr.* 196.—Afr. austr.
Comaru, *E. Mey. l. c.* 195.—Afr. austr.
crispum, *R. Grah. in Edinb. Phil. Journ.* (1830) 170.—Afr. austr.
ellipticum, *A. Rich. Tent. Fl. Abyss.* ii. 49.—Abyssin.
filiforme, *Harv. Thes. Cap.* i. 58. t. 93.—Afr. austr.
Gerrardi, *Harv. l. c.* ii. 61. t. 196.—Afr. austr.
glabrum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
hirsutum, *E. Mey. Comm. Pl. Afr. Austr.* 197.—Afr. austr.
laevigatum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
lineare, *A. Rich. Tent. Fl. Abyss.* ii. 49.—Abyssin.
macrorrhizum, *E. Mey. Comm. Pl. Afr. Austr.* 197.—Afr. austr.
maculatum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
micranthum, *E. Mey. Comm. Pl. Afr. Austr.* 196.—Afr. austr.
ovatum, *Oliver, in Saund. Ref. Bot.* t. 226.—Afr. austr.
parviflorum, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.
plocamoides, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 112.—Afr. trop.
spathulatum, *Lindl. Bot. Reg.* t. 1113.—Afr. austr.
tuberosum, *R. Br. in Bot. Mag.* t. 2343.—Afr. austr.
volubile, *Hook. f. Fl. Brit. Ind.* iv. 65.—Ind. or.

BRACHYSTEMMA, D. Don, Prod. Fl. Nep. 216 (1288) = **Arenaria**, Rupp. (Caryophyll.).
calycinum, D. Don, l. c. = A. nepalensis.

BRACHYSTEMUM, Michx. Fl. Bor. Am. ii. 5. t. 31, 32 (1803) = **Pycnanthemum**, Michx. (Labiat.).
lanceolatum, Willd. Enum. Hort. Berol. 623 = P. lanceolatum.
linifolium, Willd. l. c. = P. linifolium.
muticum, Michx. Fl. Bor. Am. ii. 6. t. 32 = P. muticum.
verticillatum, Michx. l. c. 6. t. 31 = P. lanceolatum.
virginicum, Michx. l. c. 6 = P. lanceolatum.

BRACHYSTEPHANUS, Nees, in DC. Prod. xi. 511 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1105.
Lyallii, *Nees, l. c.*—Madag.

BRACHYSTEPHIUM, Less. Syn. Comp. 388 (1832) = **Brachycome**, Cass. (Compos.).
leucanthemoides, Less. l. c. 389 = B. diversifolia.
scapigerum, DC. Prod. vi. 304 = B. scapigera.

BRACHYSTEPIS, Pritz. Ind. Ic. 162 (1855) = **Aeonia**, Lindl. (Orchid.).

BRACHYSTYLIS, E. Mey. ex DC. Prod. vi. 76 (1837) = **Marasmodes**, DC. (Compos.).
scoparia, E. Mey. ex DC. Prod. vi. 76 (= *Brachymeris scoparia*).—Afr. austr.

BRACHYSTYLUS, Dulac, Fl. Hautes-Pyr. 85 (1867) = **Koeleria**, Pers. (Gramin.).
cristatus, Dulac, l. c. = K. cristata.
setaceus, Dulac, l. c. = K. setacea.

BRACHYTOME, Hook. f. Ic. Pl. xi. 70. t. 1088 (1871). *RUBIACEAE*, Benth. & Hook. f. ii. 87.
Wallichii, *Hook. f. ll. cc.*—Reg. Himal.

BRACHYTROPIS, Reichb. Consp. 120 (1828) = **Polygala**, Tourn.
microphylla, Willk. in Oestr. Bot. Zeitschr. xxix. (1879) 288 = P. microphylla.

BRACKENRIDGEA, A. Gray, Bot. U. St. Expl. Exped. i. 361. t. 42 (1854). *OCHNACEAE*, Benth. & Hook. f. i. 318.
australiana, *F. Muell. Fragm.* v. 29.—Austral.
fascicularis, *Villar, in Blanco, Fl. Phil.* ed. III. *Nov. App.* 40.—Ins. Philipp.

BRACKENRIDGEA :—
Hookeri, *A. Gray, in Proc. Am. Acad.* iii. (1857) 51. —Ins. Fiji.
nitida, *A. Gray, Bot. U. St. Expl. Exped.* i. 362.—Ins. Fiji.
zanguebarica, *Oliver, in Hook. Ic. Pl.* t. 1096.—Afr. trop.

BRACONOTIA, Godr. Fl. Lorr. ed. I. iii. 191 (1844) [Ed. ii. (1861) Braconnotia.] = **Agropyron**, J. Gaertn. (Gramin.).
acuta, Godr. l. c. 192 = A. acutum.
elymoides, Godr. l. c. 193 = A. caninum.
glauca, Godr. l. c. 192 = A. glaucum.
juncea, Godr. l. c. = A. junceum.
officinarum, Godr. l. c. = A. repens.
pungens, Godr. l. c. = A. pungens.
rigida, Godr. l. c. = A. glaucum et rigidum.
stricta, Godr. l. c. = A. strictum.

BRACTEARIA, DC. ex Steud. Nom. ed. II. i. 222 (1840) = **Tibouchina**, Aubl. (Melast.).
Kappleri, Steud. ex Cogn. in Mart. Fl. Bras. xiv. III. 375 = T. aspera.

BRACTEOLARIA, Hochst. in Flora, xxiv. (1841) II. 658 = **Baphia**, Afzel. (Legum.).
polygalacea, Hook. f. in Hook. Niger Fl. 321 = Baphia polygalacea.
racemosa, Hochst. in Flora, xxiv. II. (1841) 658 = Baphia racemosa.

BRADBURIA, Torr. & Gray, Fl. N. Am. ii. 250 (1841). *COMPOSITAE*, Benth. & Hook. f. ii. 251.
hirtella, *Torr. & Gray, l. c.*—Texas.

BRADBURIA, Spreng. Syst. iii. 255 (1826) = scq.

BRADBURYA, Rafin. Fl. Ludov. 104 (1817) = **Wistaria**, Nutt. (Legumin.).
erecta, Rafin. l. c. 105 = W. speciosa.
scandens, Rafin. l. c. 104 = W. speciosa.

BRADDLEYA, Vell. Fl. Flum. 93 (1825); ii. t. 140 = **Amphirrhox**, Spreng. (Violar.).
legalis, Vell. l. c. 94; ii. t. 140 = A. longifolia.

BRADLAEIA, Neck. Elem. i. 169 (1790) = **Siler**, Scop. (Umbellif.).

BRADLEA, Adans. Fam. ii. 324 (1763) = **Apios**, Tourn. (Legumin.).

BRADLEIA, Banks, ex Gaertn. Fruct. ii. 127. t. 109 (1791) = **Glochidion**, Forst. (Euphorb.).
acuminata, Wall. Cat. sub n. 7855 = G. lanceolarium.
arborescens, Steud. Nom. ed. II. i. 222 = G. arborescens.
Blumei, Steud. l. c. = G. molle.
coccinea, Wall. Cat. n. 7868 = G. coccineum.
coriacea, Wall. l. c. 7872 = Chaetocarpus castanocarpus.
coronata, Wall. l. c. 7857 = G. leiostylum.
dioica, Gaertn. ex Vahl, fide Baill. Adansonia, i. (1860) 80 = Flueggia microcarpa.
Finlaysoniana, Wall. Cat. n. 7860 = G. superbum.
glauca, Labill. Sert. Austr. Caled. 76. t. 77 = G. glaucum.
glaucophylla, Hassk. Cat. Hort. Bog. Alt. 242 = G. glaucum.
Glochidion, Gaertn. Fruct. ii. 128. t. 109 = G. ramiflorum.
hirsuta, Roxb. Hort. Beng. 104; Fl. Ind. iii. 699 = G. hirsutum.
impuber, Roxb. ll. cc. 104, 698 = G. Lambertianum.
Kipareh, Steud. Nom. ed. II. i. 222 = G. glaucum.
laevigata, Wall. Cat. n. 7853.—Ind. or.
lanceolaria, Roxb. Hort. Beng. 69; Fl. Ind. iii. 697 = G. lanceolarium.
lanceolaria, Wall. Cat. n. 7855 = G. Hohenackeri et sphaerogynum.
littorea, Steud. Nom. ed. II. i. 222 = G. littorale.
lucida, Steud. l. c. = G. lucidum.
lutescens, Steud. l. c. = G. lutescens.
macrocarpa, Steud. l. c. = G. macrocarpum.
macrophylla, Labill. Sert. Austr. Caled. 77. t. 78 = G. macrophyllum.
mollis, Steud. Nom. ed. II. i. 222 = G. molle.

BRADLEIA :—
mollis, Wall. Cat. n. 7858 = Glochidion hirsutum.
moluccana, Steud. Nom. ed. II. i. 222 = G. moluccanum.
multiloculars, Spreng. Syst. iii. 19 = G. multiloculare.
Nephroia, Steud. Nom. ed. I. 116 = Cocculus Nephroia.
nitida, Roxb. Hort. Beng. 69; Fl. Ind. iii. 699 = G. zeylanicum.
nitida, Wall. Cat. n. 7867 = G. multiloculare.
obliqua, Spreng. Syst. iii. 19 = G. zeylanicum.
obliqua, Wight, ex Wall. Cat. n. 7947 = Phyllanthus Wightianus.
obtusa, Wall. l. c. 7869 = G. littorale.
ovata, Wall. l. c. 7052 = G. velutinum.
philippensis, Buch.-Ham. in Wall. l. c. 7871 = G. mutiloculare.
philippensis, Willd. Sp. Pl. iv. 592 = G. philippinense.
philippica, Cav. Ic. iv. 48. t. 371 = G. philippinense.
pinnata, Roxb. Hort. Beng. 69; Fl. Ind. iii. 700 = G. obscurum.
pubera, Roxb. ll. cc. 104; 698 = G. multiloculare.
rubra, Steud. Nom. ed. II. i. 222 = G. rubrum.
sinensis, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 128 = G. obovatum.
sinica, Gaertn. Fruct. ii. 127. t. 109. f. 1 = G. sinicum.
sinica, Hort. ex Mncll. Arg. in DC. Prod. xv. II. 414 = Phyllanthus nobilis.
timoriensis, Steud. Nom. ed. II. i. 222 = G. zeylanicum.
Wightiana, Wall. Cat. n. 7862 = G. ellipticum.
zeylanica, Gaertn. Fruct. ii. 128. t. 109 = G. zeylanicum.
zeylanica, Labill. Sert. Austr. Caled. 76. t. 76 = G. philippinense.
zeylanica, Wall. Cat. n. 7855 G. = G. lanceolarium.

BRAGANTIA, Lour. Fl. Cochinch. 528 (1790). *ARISTOLOCHIACEAE*, Benth. & Hook. f. iii. 122.
APAMA, Lam. Encyc. i. 91 (1783).
ASIPHONIA, Griff. in Trans. Linn. Soc. xix. (1845) 333. t. 37.
CERAMIUM, Blume, Bijdr. 1134 (1826).
CYCLODISCUS, Klotzsch, in Monatsb. Acad. Berl. (1859) 591.
MUNNICKIA, Reichb. Consp. 85 (1828).
STRAKAEA, Presl, Epim. Bot. 221 (1850).
TRIMERIZA, Lindl. Bot. Reg. sub t. 1543 (1832).
VANHALLIA, Schult. f. Syst. vii. 18 (1829).
corymbosa, *Griff. in Trans. Linn. Soc.* xix. (1845) 335. —Malaya.
hispida, Thw. ex Duch. in DC. Prod. xv. I. 430 = Wallichii.
khasiyana, Griff. in Trans. Linn. Soc. xix. (1845) 336 = tomentosa.
latifolia, Lindl. Bot. Reg. sub t. 1543 = Wallichii.
melastomaefolia, Duch. in DC. Prod. xv. I. 429 = corymbosa.
racemosa, *Lour. Fl. Cochinch.* 528.—China.
tomentosa, *Blume, Enum. Pl. Jav.* 82.—Ind. or.; Malaya.
Wallichii, *R. Br. in Wall. Cat.* n. 7415.—Ind. or.

BRAGANTIA, Vand. Fasc. Pl. Nov. (1771) 6 = **Gomphrena**, Linn. (Amarant.).
Vandelli, Steud. Nom. ed. II. i. 222 = G. officinalis.

BRAHEA, Mart. Hist. Nat. Palm. iii. 243, 319. t. 137, 162 (1830?). *PALMAE*, Benth. & Hook. f. iii. 926.
armata, S. Wats. in Proc. Am. Acad. xi. (1876) 146 = Erythea armata.
calcarata, Liebm. ex Linden, Cat. n. 87 (1871), sphalm. = seq.?
calcarea, *Liebm. in Mart. Hist. Nat. Palm.* iii. 319.—Mexic.
conduplicata, *Linden, Illustr. Hortic.* xxviii. (1881) 16. —Mexic.
dulcis, J. Cooper, in Smithson. Rep. (1860) 442 = Washingtonia filifera.
dulcis, *Mart. Hist. Nat. Palm.* iii. 244. t. 137. 162.—Peruv.
edulis, H. Wendl. ex S. Wats. in Proc. Am. Acad. xi. (1876) 120, 146 = Erythea edulis.
filamentosa, Hort. ex S. Wats. l. c. 147 = Washingtonia filifera.
minima, H. Wendl. in Kerch. Palm. 235 = Sabal Adansoni?
Roezli, *Linden, Illustr. Hortic.* xxviii. (1881) 38.— Arizona.
serrulata, *H. Wendl. in Kerch. Palm.* 235.—Georgia.

BRAMI, Adans. Fam. ii. 208 (1763) = **Herpestis**, Gaertn. f. (1805) (Scrophul.).

BRAMIA, Lam. Encyc. i. 459 (1783), nomen prius = **Herpestis**, Gaertn. f. (1805) (Scrophul.).
floribunda, F. Muell. Fragm. ix. 167 = H. floribunda.
indica, Lam. Encyc. i. 459 = H. Monniera.
lanigera, G. Don, Gen. Syst. iv. 546 = H. lanigera.
semiserrata, Mart. Amoen. Monac. t. 8 = H. gratioloides.

BRANCIONA, Salisb. Gen. Pl. Fragm. 36 (1866) = **Albuca**, Linn. (Liliac.).
setosa, Salisb. l. c. = A. setosa.

BRANDESIA, Mart. Nov. Gen. et Sp. ii. 25. t. 125 (1826) = **Telanthera**, R. Br. (Amarant.).
alba, Mart. ex Moq. in DC. Prod. xiii. II. 311 = Achyranthes alba.
echinocephala, Hook. f. in Trans. Linn. Soc. xx. (1851) 189 = T. echinocephala.
elongata, Mart. Beitr. Amarant. 105. n. 4 = T. elongata.
lanceolata, Benth. Pl. Hartw. 247 = T. lanceolata.
Martiana, Gill. ex Moq. in DC. Prod. xiii. II. 363 = T. philoxeroides.
mexicana, Schlecht. in Linnaea, vii. (1832) 392 = T. mexicana.
porrigens, Mart. Beitr. Amarant. 106 = T. porrigens.
puberula, Mart. Nov. Gen. et Sp. ii. 27 = puberula.
pubiflora, Benth. in Bot. Voy. Sulph. 157 = T. pubiflora.
pycnantha, Benth. l. c. = T. pycnantha.
rufa, Mart. Nov. Gen. ii. 26. t. 125 = T. rufa.
serpyllifolia, Mart. Beitr. Amarant. 106 = T. serpyllifolia.
villosa, Mart. Nov. Gen. et Sp. ii. 28. t. 128 = T. Martii.

BRANDISIA, Hook. f. & Thoms. in Journ. Linn. Soc. viii. (1865) 11. t. 4. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 938.
discolor, Hance, in Journ. Bot. xviii. (1880) 299 = Hancei.
discolor, *Hook. f. & Thoms. in Journ. Linn. Soc.* viii. (1865) 11.—Burma.
Hancei, *Hook. f. Fl. Brit. Ind.* iv. 257.—China.

BRANDONIA, Reichb. Consp. 127 (1828) = **Pinguicula**, Linn. (Lentibul.).

BRANDTIA, Kunth, Rev. Gram. ii. 511. t. 170 (1830) = **Arundinella**, Raddi (Gramin.).
holcoides, Kunth, l. c. = A. holcoides.

BRANICA, Endl. Ench. 339 (1841), sphalm. = BRAMIA, Lam. = **Herpestis**, Gaertn. f. (Scrophul.).

BRANICIA, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 476 = **Senecio**, Tourn. (Compos.).
insignis, Andrz. l. c. = S. racemosus.

BRANDZEIA, Baill. Adansonia, ix. (1869) 217. *LEGUMINOSAE*.
filicifolia, *Baill. l. c.*—Ins. Mascar.

BRASENIA, Schreb. Gen. 372 (1789). *NYMPHAEACEAE*, Benth. & Hook. f. i. 46.
HYDROPELTIS, Michx. Fl. Bor. Am. i. 323. t. 29 (1803).
IXODIA, Soland. ex DC. Syst. ii. 37 (1821).
RONDACHINE, Bosc; Encyc. Agric. vi. 1816) 180 (nomen gallicum, tamen citatur ab DC. Syst. et Steud.).
Hydropeltis, Muhl. Cat. 55; Rafin. Med. Fl. i. 90. t. 17 = Schreberi.
nymphoides, Baill. Hist. Pl. iii. 82 = Schreberi.
peltata, Pursh, Fl. Bor. Am. ii. 389 = Schreberi.
purpurea, Casp. in Jorn. Sc. Acad. Lisb. iv. (1873) 74 312 = Schreberi.
Schreberi, *J. F. Gmel. Syst.* 853.—Ind. or.: Japon; Austral.; Afr. trop.; Am. bor.

BRASILIASTRUM, Lam. Encyc. i. 462 (1783).
americanum, Lam. l. c. = Comocladia Brasiliastrum et Picramnia Antidesma.

BRASILIUM, J. F. Gmel. Syst. 417 (1791).
glabrum, J. F. Gmel. l. c. = Picramnia Antidesma.
hirsutum, J. F. Gmel. l. c. = Comocladia Brasiliastrum.

These are synonymous. The name to be kept up is

BRASSAIA, Endl. Nov. Stirp. Dec. i. 89 (1839). *ARALIACEAE*, Benth. & Hook. f. i. 943.
actinophora, *Endl. l. c.*—Austral.
actinophylla, *F. Muell. Fragm.* iv. 121.—Austral.
capitata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 732.—Reg. Himal.; Burma.
littorea, *Seem. in Journ. Bot.* ii. (1864) 244.—Amboina.
macrostachya, *Seem. l. c.*—N. Guinea.
palmata, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 106 = Trevesia sundaica.
sessilis, *Seem. in Journ. Bot.* ii. (1864) 244.—Sumatra.

BRASSAIOPSIS, Decne. & Planch. in Rev. Hortic. Sér. IV. iii. (1854) 106. *ARALIACEAE*, Benth. & Hook. f. i. 945.
ARALIOPSIS, Kurz, in Andam. Rep. 39 (1870).
aculeata, *Seem. in Journ. Bot.* ii. (1864) 293.—Reg. Himal.
alpina, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 736.—Reg. Himal.
confluens, Seem. in Journ. Bot. ii. (1864) 292 = B. Hainla et Trevesia palmata.
cyrtostyla, *Seem. l. c.* 293.—Sumatra.
floribunda, Seem. l. c. 262 = speciosa.
glomerulata, Regel, Gartenfl. (1863) 275. t. 411 = speciosa.
Griffithii. *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 736.—Reg. Himal.
Hainla, *Seem. in Journ. Bot.* ii. (1864) 291.—Reg. Himal.
hispida, *Seem. l. c.* 292.—Reg. Himal.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 737.—Reg. Himal.
mitis, *C. B. Clarke, l. c.* 736.—Reg. Himal.
palmata, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) II. 77.—Reg. Himal.; Burma.
ricinifolia, Seem. in Journ. Bot. ii. (1864) 291 = Acanthopanax ricinifolium.
simplicifolia, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 735.—Reg. Himal.
speciosa, *Decne. & Planch. in Rev. Hortic.* Sér. IV. iii. (1854) 106.—Reg. Himal.

BRASSAVOLA, Adans. Fam. ii. 127 (1763) = **Helenium**, Linn. (Compos.).

BRASSAVOLA, R. Br. in Ait. Hort. Kew. ed. II. v. 216 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 533.
LYSIMNIA, Rafin. Fl. Tellur. i. 43 (1836).
TULEXIS, Rafin. l. c. 42 (1836).
acaulis, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 152. f. 216.—Am. centr.
amazonica, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 3.—Bras.
angustata, *Lindl. Bot. Reg.* (1838) *Misc.* 41.—Guiana.
appendiculata, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 23 = cucullata.
Ceboletta, *Reichb. f. in Bonplandia*, iii. (1855) 221.—Hab.?
cordata, *Lindl. Bot. Reg.* t. 1913.—Bras.
cucullata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 216.—Ind. occ.
cuspidata, Hook. Bot. Mag. t. 3722 = cucullata.
Digbyana, Lindl. Bot. Reg. (1846) t. 53 = Laelia Digbyana.
elegans, Hook. Bot. Mag. t. 3098 = Tetramicra rigida.
filifolia, *Linden, Illustr. Hortic.* xxviii. (1881) 64.—N. Granat.
flagellaris, *Rodrig. Orch. Nov.* ii. 161.—Bras.
fragrans, Lem. in Jard. Fleur. iii. (1853) Misc. 78 = tuberculata.
fragans, *Rodrig. Orch. Nov.* i. 65.—Bras.
Gibbsiana, Hort. ex Nichols. Dict. Gard. i. 209 = tuberculata.
glauca, Lindl. Bot. Reg. (1839) Misc. 47 = Laelia glauca.
grandiflora, Lindl. l. c. 16 = nodosa.
lineata, *Hook. Bot. Mag.* t. 4734.—Am. centr.
Martiana, *Lindl. Bot. Reg.* (1836) sub t. 1914.—Bras.
Mathieuana, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxi. (1853) 290 = lineata.
nodosa, Hook. Bot. Mag. (1833) t. 3229 = subulifolia.

BRASSAVOLA:—
nodosa, *Lindl. Gen. et Sp. Orch.* 114.—Ind. occ.
odoratissima, Regel, Gartenfl. (1852) 325. t. 33 = cucullata.
Perrinii, *Lindl. Bot. Reg.* t. 1561.—Bras.
Pumilio, *Reichb. f. in Linnaea*, xviii. (1844) 402.—Mexic.
retusa, *Lindl. Bot. Reg.* (1847) sub t. 20.—Venezuela.
revoluta, *Rodrig. Orch. Nov.* ii. 161.—Bras.
rhopalorrhachis, *Reichb. f. in Bot. Zeit.* x. (1852) 855.—Guatemala.
Sloanei, Griseb. Fl. Brit. W. Ind. 620 = subulifolia.
Sloanei, Lindl. ex Hcynh. Nom. ii. 76, nomen (= nodosa?).—Jamaica.
suaveolens, Gal. ex Hemsl. in Gard. Chron. (1879) i. 235, 236 = Epidendri sp.?
subulifolia, *Lindl. Gen. et Sp. Orch.* 115.—Ins. Nevis.
surinamensis, Focke, in Tijdschr. Nat. Wetens. iv. (1851) 17 = angustata.
tuberculata, *Hook. Bot. Mag.* t. 2878.—Bras.
venosa, *Lindl. Bot. Reg.* (1840) *Misc.* 20. t. 39.—N. Granat.
vomeriformis, *Reichb. f. ex Griseb. Fl. Brit. W. Ind.* 621.—Ind. occ.

BRASSAVOLAEA, Poepp. & Endl. Nov. Gen. et Sp. ii. 3 (1838) = praec.

BRASSAVOLEA, Spreng. Syst. iii. 744 (1826) = praec.

BRASSENIA, Heynh. Nom. i. 130 (1840) = **Brasenia**, Schreb. (Nymphaeac.).

BRASSIA, R. Br. in Ait. Hort. Kew. ed. II. v. 215 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 564.
angusta, Lindl. Bot. Reg. (1844) Misc. 5 = Lawrenceana.
antherotes, *Reichb. f. in Gard. Chron.* (1879) II. 782.—Ecuador.
arachnoidea, *Rodrig. Orch. Nov.* i. 97.—Bras.
arcuigera, *Reichb. f. in Gard. Chron.* (1869) 388.—Peruv.
aristata, Lindl. Bot. Reg. (1844) Misc. 7 = verrucosa.
bidens, *Lindl. l. c.* 6.—Hab.?
brachiata, *Lindl. in Benth. Pl. Hartw.* 94.—Guatemala.
brachypus, *Reichb. f. in Gard. Chron.* (1875) I. 136.—Ecuador.
caudata, *Lindl. Bot. Reg.* t. 832.—Ind. occ.
chloroleuca, *Rodrig. Orch. Nov.* i. 97.—Bras.
chlorops, *Endr. & Reichb. f. in Gard. Chron.* (1873) 542.—Costa Rica.
cinnabarina, *Linden, ex Lindl. Fol. Orchid. Brassia*, 6.—N. Granat.
cinnamomea, Linden, ex Lindl. l. c. 5 = Keiliana.
Clowesii, Lindl. Bot. Reg. (1844) Misc. 7.—Bras.
cochleata, Knowl. & Westc. Flor. Cab. ii. (1838) 23. t. 53 = Lawrenceana.
coryandra, Morr. in Ann. Gand, iv. (1848) 295. t. 212 = verrucosa.
Cowani, Hort. ex Lindl. Fol. Orchid. Brassia, 4 = verrucosa.
cryptophthalma, *Reichb. f. in Gard. Chron.* (1879) II. 554.—Peruv.
elegantula, *Reichb. f. l. c.* (1885) II. 616.—Mexic.
euodes, *Reichb. f. l. c.* (1880) I. 680.—N. Granat.
farinifera, *Linden & Reichb. f. l. c.* (1870) 923.—Ecuador.
Gireoudiana, *Reichb. f. & Warsc. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 273.—Bras.
glumacea, *Lindl. Orch. Linden.* 17.—Venezuela.
guttata, Lindl. in Benth. Pl. Hartw. 94 = maculata.
Helenae, *Reichb. ex Linden, Illustr. Hortic.* xxviii. (1881) 64.—Ecuador.
hexodonta, *Reichb. f. in Bonplandia*, ii. (1854) 98.—Peruv.
imbricata, *Lindl. Fol. Orchid. Brassia*, 5.—N. Granat.
Josstiana, *Reichb. f. in Regel, Gartenfl.* iii. (1854) 309.—Bras.
Keiliana, *Reichb. f. ex Lindl. & Paxt. Flow. Gard.* iii. (1852–53) 114.—N. Granat.
Lanceana, *Lindl. Bot. Reg.* t. 1754.—Guiana.
Lawrenceana, *Lindl. l. c.* (1841) *Misc.* 2. t. 18.—Bras.
longicuspis, *Reichb. f. Otia Bot. Hamb.* 23.—Ecuador.

BRASSIA:—
longiloba, DC. in Mém. Soc. Phys. Genèv. xi. (1848) II. 427. t. 1 = verrucosa.
macrostachya, Lindl. Sert. Orch. t. 6 = Lanceana.
maculata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 215.—Jamaica.
neglecta, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 322.—Am. centr.
ocanensis, *Lindl. Fol. Orch. Brassia*, 5.—N. Granat.
odontoglossoides, Klotzsch & Karst. in Otto & Dietr. Allg. Gartenz. xv. (1847) 330 = verrucosa.
peruviana, *Poepp. & Endl. Nov. Gen. et Sp.* ii. t. 117.—Peruv.
pumila, *Lindl. Bot. Reg.* (1845) *Misc.* 60 = Lanceana.
signata, *Reichb. f. in Gard. Chron.* (1881) II. 6.—Hab.?
spathacea, *Linden & Reichb. f. in Illustr. Hortic.* xxviii. (1881) 64, nomen.—N. Granat.
thyrsodes, *Reichb. f. in Gard. Chron.* (1868) 842.—Peruv.
verrucosa, *Batem. Orch.* t. 22.—Mexic.
villosa, *Lindl. Fol. Orchid. Brassia*, 4.—Peruv.
Wageneri, *Reichb. f. in Bonplandia*, ii. (1854) 14.—N. Granat.
Warszewiczii, *Reichb. f. in Bot. Zeit.* x. (1852) 693.—Ecuador.
Wrayae, Skinner, ex Hook. Bot. Mag. t. 4003 = maculata.

BRASSICA, [Tourn.] Linn. Syst. ed. I (1735) *CRUCIFERAE*, Benth. & Hook. f. i. 84.
AGROSINAPIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 329.
BRASSICASTRUM, Link, Handb. ii. 318 (1831).
BRASSICELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 330.
BONANNIA, Presl, Fl. Sic. i. 99 (1826).
CORYNELOBUS, R. Roem. in Linnaea, xxv. (1852) 7.
ERUCASTRUM, Presl, Fl. Sic. i. 92 (1826).
GUENTHERA, Andrz. ex Bess. Enum. Pl. Volh. 83 (1822).
HETEROCRAMBE, Coss. & Dur. Ind. Sem. Hort. Burdig. (1867) 25.
HIRSCHFELDIA, Moench, Meth. 264 (1794).
MELANOSINAPIS, Schimp. & Spenn. in Spenn. Fl. Frib. iii. 944 (1829).
MICROPODIUM, Reichb. Consp. 185 (1828).
MUTARDA, Bernh. Syst. Verz. Erf. 184 (1800).
NAPUS, Schimp. & Spenn. in Spenn. Fl. Frib. iii. 939 (1829).
RAMPHOSPERMUM, Andrz. ex Reichb. Moessl. Handb. ed. II. i. p. lvii (1827).
RAPA, Tourn. ex Linn. Syst. ed. I (1735).
RHAMPHOSPERMUM, Andrz. ex Bess. Enum. Pl. Volh. 83 (1822).
SINAPI, Dulac, Fl. Hautes-Pyr. 194 (1867).
SINAPIDENDRON, Lowe, in Trans. Camb. Phil. Soc. iv. 36 (1831).
SINAPIS, Linn. Syst. ed. I (1735).
SINAPISTRUM, Spach, Hist. Vég. Phan. vi. 343 (1838).
STRANGALIS, Dulac, Fl. Hautes-Pyr. 195 (1867).
adpressa, *Boiss. Voy. Espagne*, ii. 38; *Vis. Fl. Dalm.* iii. 359.—Europ.; Reg. Mediterr.
agrestis, Steud. Nom. ed. I. 116, 117 = Conringia orientalis.
alba, *Boiss. Voy. Espagne*, ii. 39.—Europ.; Oriens; Ind. or. (cult.).
alba, Gilib. Fl. Lituan. ii. 63 = Conringia orientalis.
Allionii, Moretti, Ind. Sem. Hort. Taur. (1829) (fide DC. sched.) = oleracea.
alpina, Linn. Mant. 95 = Conringia alpina.
amplexicaulis, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 23.—Abyssin.
amplexicaulis, *Janka, in Termész. Füzetek*, vi. (1882) 177.—Europ.
amplexicaulis, *Pomel, Mat. Fl. Alt.* 15.—Algeria.
antarctica, Molina, Sagg. Chile, ed. II. 292 = magellanica.
anticaria, *Rouy & Janka, in Termész. Füzetek*, vi. (1882) 177.—Europ.
apula, *Janka, l. c.* 180.—Europ.
arborea, Steud. Nom. ed. I. 116, 117 = campestris.
arenosa, Jord. Diag. i. 181 = B. Cheiranthos.
armoracioides, *Czern. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 311.—Ucrania.
arvensis, Habl. Taur. 146 = campestris.

BRASSICA :—
arvensis, Linn. Mant. 95 = Moricandia arvensis.
asperifolia, Lam. Encyc. i. 746 = campestris.
atlantica, Coss. & Dur. ex Munby, Cat. Pl. Algér. 2 = Gravinae.
Aucheri, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 88 = Raphanus Aucheri.
austriaca, Jacq. Fl. Austr. iii. 45 = Conringia austriaca.
azurea, *Uspensky, in Bull. Soc. Nat. Mosc.* vii. (1834) 376, nomen.—Rossia.
balearica, Loisel. Fl. Gall. ii. 69 = Robertiana.
balearica, *Pers. Syn.* ii. 206.—Ins. Balear.
barbareaefolia, *Ball, in Journ. Linn. Soc.* xvi. (1878) 328.—Marocco.
Barrelieri, *Janka, in Termész. Füzetek*, vi. (1882) 179. —Europ.
Besseriana, *Andrz. ex Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) 1. 134.—Rossia; As. temp.
Blancoana, *Boiss. Diagn.* Ser. II. i. 29.—Hispan.
boetica, *Boiss. Voy. Espagne*, ii. 38. t. 9.—Hispan.
Boissieri, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 283. —Afr. bor.
botrytis, Mill. Gard. Dict. ed. VIII. n. 3 = oleracea.
Botteri, *Vis. Fl. Dalm.* iii. 135.—Dalmat.
brachyloma, *Boiss. & Reut. Diagn.* Ser. II. i. 30.— Marocco.
bracteata, *Janka, in Termész. Füzetek*, vi. (1882) 178. —Europ.
bractcolata, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 33.—Aegypt.
brevipes, Syme, Engl. Bot. ed. III. i. 140 = Diplotaxis muralis, viminea.
Briggsii, Varenne, in Journ. Bot. xix. (1881) 360 = B. Rapa.
bullata, Pasq. Cat. Ort. Bot. Nap. 17 = oleracea.
Bunias, *DC. Syst.* ii. 606.—Oriens.
campestris, *Linn. Sp. Pl.* 666.—Orb. Vet. (Cosmop. cult.).
campestris, Oed. in Fl. Dan. x. t. 550 = Turritis glabra.
capitata, Hort. ex Pers. Syn. ii. 206, in syn. = oleracea.
carinata, *A. Braun, in Flora*, xxiv. (1841) 267.— Abyss.
catholica, Brot. ex Willk. & Lange, Prod. iii. 867 = Diplotaxis catholica.
caulorapa, Pasq. Cat. Ort. Bot. Nap. 17 = oleracea.
cespitosa, *Pomel, Nouv. Mat. Fl. Atl.* 361.—Afr. bor.
cheiranthiflora, DC. Syst. ii. 601 = B. Cheiranthos.
Cheiranthos, *Vill. Prosp.* 40; *Hist. Pl. Dauph.* iii. 332. t. 36.—Europ. occ.; Afr. bor.
chinensis, Linn. Cent. 19; in Amoen. Acad. iv. 280 = campestris.
coryneloba, *Nym. Consp.* 47.—Hispan.
Cossoneana, *Boiss. & Reut. Diagn.* Ser. II. i. 31.— Hispan.
crassifolia, Forsk. Fl. Aegypt. Arab. 118 = Erucaria crassifolia.
cretica, Guss. Cat. Hort. Boccadif. (1821) 11 = rupestris.
cretica, *Lam. Encyc.* i. 747.—Graecia.
cretica, Moris, Stirp. Sard. Elench. i. 3 = insularis.
crispa, *Rafin. Specch.* ii. 69.—Sicilia
cyrenaica, *Spreng. Syst.* ii. 911.—Cyrenaic.
deflexa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 87.—Syria; Persia.
densiflora, Jord. Diag. i. 186 = B. Cheiranthos.
dimorpha, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 306.—Algeria.
dissecta, *Boiss. Voy. Espagne*, ii. 40.—Hispan.
elata, *Ball, in Journ. Bot.* xi. (1873) 298.—Marocco.
elongata, *Ehrh. Beitr.* vii. 159.—Europ. or.; Reg. Caucas.; Persia.
ensifera, *Dum. Fl. Belg. Prod.* 122.—Belgium.
erosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 311.—Afr. austr.
Eruca, Boiss. Voy. Bot. Espagne, ii. 41 = Eruca longirostris.
Eruca, Linn. Sp. Pl. 667 = Eruca sativa.
Erucastrum, Poll. Pl. Palat. ii. 249 = B. Cheiranthos.
Erucastrum, *Linn. Sp. Pl.* 667.—Europ.
erucoides, Boiss. Voy. Bot. Espagne, ii. 33 = Diplotaxis erucoides.
erucoides, Roxb. Hort. Beng. [48]; Fl. Ind. iii. 117 = Eruca sativa.
erysimoides, Pourr. in Mém. Acad. Toul. iii. (1788) 308 = B. Cheiranthos.
erysimoides, *Sieber, ex Spreng. Syst.* ii. 912.—Aegypt.

BRASSICA :—
fimbriata, Hort. ex Steud. Nom. ed. II. i. 223 = oleracea.
floridana, *Rafin. Atl. Journ.* 147.—Am. bor.
foetida, *Janka, in Termész. Füzetek*, vi. (1882) 177. —Europ.
foliosa, Willd. Enum. Hort. Berol. 688 = alba.
Forskahli, *Schult. Obs.* 131. n. 1012.—Oriens.
fragilis, Sieber, ex Spreng. Syst. ii. 912 = Diplotaxis acris.
frutescens, *Soland. ex DC. Syst.* ii. 624.—Ins. Madera.
fruticulosa, *Cyril. Pl. Rar.* ii. 7. t. 1.—Reg. Mediterr.
geniculata, Ball, in Journ. Linn. Soc. xvi. (1878) 327 = adpressa.
glaberrima, *Pomel, Nouv. Mat. Fl. Atl.* 364.— Algeria.
glareosa, Jord. Diagn. i. 183 = B. Cheiranthos.
gongylodes, Mill. Gard. Dict. ed. VIII. n. 8 = oleracea.
gracilis, *Pomel, Nouv. Mat. Fl. Atl.* 225.—Algeria.
grandiflora, Balansa, ex Coss. in Ann. Sc. Nat. Sér. IV. i. (1854) 221 = sabularia.
Gravinae, *Tenore, Prod. Fl. Nap.* 39. t. 62.—Ital.; Afr. bor.
Griffithii, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 171 = Diplotaxis Griffithii.
Havardi, *Pomel, Nouv. Mat. Fl. Atl.* 363.—Algeria.
heleniana, *Burch. ex Loud. Hort. Brit.* 264.—Ins. S. Helen.
heterophylla, Boiss. ex Nym. Syll. 197 = adpressa.
hirta, *Moench, Meth. Suppl.* 84.—Hab.?
hispida, *Boiss. Voy. Espagne*, ii. 40.—Hispan.; Afr. bor.
hispida, Tenore, Prod. Fl. Nap. 39 = Eruca sativa.
humilis, Boiss. Voy. Espagne, 35 = latisiliqua.
humilis, *DC. Syst.* ii. 598.—Gallia.
hybrida, *Tenore, Prod. Fl. Nap.* 59.—Italia.
incana, Maly, Enum. Stirp. Austr. 272 = Botteri.
incana, *Tenore, Prod. Fl. Nap.* 39.—Italia.
insularis, *Moris, Fl. Sard.* i. 168.—Sardinia.
intricata, *Janka, in Termész. Füzetek*, vi. (1882) 179. —Europ.
japonica, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. ii. 74 = auriculata?
juncea, *Coss. in Bull. Soc. Bot. Fr.* vi. (1859) 609.— As. temp. et trop. (cult.).
laciniata, Steud. Nom. ed. I. 116, 117 = oleracea.
laevigata, *Lag. in Varied. Cienc.* ii. iv. (1805) 40.— Hispan.
lanceolata, *Lange, in Fl. Dan.* t. 2658.—Dania.
lanceolata, *Sauv. Fl. Cub.* 2.—Cuba.
latisiliqua, *Boiss. & Reut. Diagn.* Ser. II. i. 30.— Hispan.
lativalvis, Boiss. Diagn. Ser. I. vi. 72 = Eruca lativalvis.
leptocarpa, *Boiss. Fl. Orient.* i. 392.—Persia.
leptopetala, *Sond. in Harv. & Sond. Fl. Cap.* i. 31.— Afr. austr.
leucantha, Boiss. ex Nym. Consp. 45 = nivea.
loncholoma, *Pomel, Nouv. Mat. Fl. Atl.* 360.— Algeria.
longirostra, *Boiss. Voy. Espagne*, ii. 40.—Hispan.
lutea, Gilib. Fl. Lituan. ii. 62 = Conringia austriaca.
lyrata, *Desf. Fl. Atlant.* ii. 96. t. 166.—Afr. bor.
macloviana, Urv. in Mém. Soc. Linn. Par. iv. (1826) 616 = Arabis macloviana.
macrocarpa, *Guss. in Presl, Fl. Sic.* i. 92.—Sicilia.
macrorhiza, S. F. Gray, Nat. Arr. Brit. Pl. ii. 682 = campestris.
magellanica, Gaudich. Freyc. Voy. Bot. i. 136 in nota = Arabis macloviana.
magellanica, *Pers. Syn.* ii. 207.—Reg. Magellan.
maritima, Tardent, Ess. Hist. Nat. Bessar. 79 = oleracea.
maurorum, *Dur. in Duch. Rev. Bot.* ii. (1847) 433. —Algeria.
mesopotamica, Bernh. ex Steud. Nom. ed. II. i. 223 = B. Sinapis.
mollis, *Vis. Fl. Dalm.* iii. 359.—Europ. austr.
monensis, *Huds. Fl. Angl.* ed. II. 291.—Europ. occ.
montana, DC. Fl. Fr. iv. 651 = B. Cheiranthos.
montana, Rafin. Specch. ii. 69 = rupestris.
Moricandia, Boiss. Voy. Espagne, ii. 341 = Moricandia arvensis.
moricandoides, Boiss. Elench. 10 = Moricandia Ramburei.
muralis, Huds. Fl. Angl. ed. II. 291 = Diplotaxis tenuifolia, muralis.
muscovita, Steud. Nom. ed. I. 117 = oleracea.

BRASSICA :—
Napella, Chaix, in Vill. Hist. Pl. Dauph. iii. 334 = campestris.
napobrassica, Mill. Gard. Dict. ed. VIII. n. 2 = campestris.
Napus, Linn. Sp. Pl. 666 = campestris.
nervosa, *Ball, in Journ. Bot.* xi. (1873) 299 (sp. dubia). —Marocco.
nigra, *Koch, in Roehl. Deutschl. Fl.* ed. III. iv. 713.— Orb. vet.; (Cosmop. cult.).
nivalis, *Boiss. & Heldr. in Boiss. Diagn.* Ser. II. i. 32. —Graecia.
nivea, *Boiss. & Sprun. l. c.* Ser. I. i. 72.—Graecia.
nudicaulis, *Pomel, Nouv. Mat. Fl. Atl.* 361.—Algeria.
obtusangula, *Reichb. in Moessl. Handb.* ed. II. ii. 1185. —Europ.
obtusangulata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = B. Erucastrum.
ochroleuca, Soyer-Willem. in Ann. Sc. Nat. ii. (1824) 116 = B. Cheiranthos.
odorata, *Schrank, ex Steud. Nom.* ed. I. 117.—Europ.?
oleifera, Moench, Meth. 253 = campestris.
oleracea, *Linn. Sp. Pl.* 667.—Europ. occ. (Cosmop. cult.).
oleracea, Lour. Fl. Cochinch. 396 = campestris.
oleronensis, *A. Savat, ex Foucaud, in Bull. Soc. Bot. Fr.* xxvi. (1879) 74.—Gallia.
orientalis, Linn. Sp. Pl. 666 = Conringia orientalis.
ovalis, *Viv. Fl. Libyc. Spec.* 35. t. 16. f. 3.—Libya.
oxyrrhina, *Coss. Notes Crit.* ii. 26.—Hispan.
oxyrrhyncha, Coss. ex Regel, Ind. Sem. Hort. Petrop. (1856) 34 = oxyrrhina.
palustris, Bivona, Fl. Forojuliensis Syllab. (1855) 18 = elongata.
papillaris, Boiss. Elench. 10 = sabularia.
pendula, *Boiss. Voy. Espagne*, ii. 35.—Hispan.
peregrina, Steud. Nom. ed. I. 117 = oleracea.
perfoliata, Crantz, Stirp. Austr. ed. II. 37; Lam. Fl. Fr. ii. 487 = Conringia orientalis.
persica, Boiss. & Hohen. Diagn. Ser. I. viii. 26 = elongata.
petrosa, Jord. Diagn. i. 185 = B. Cheiranthos.
pinnata, *Spreng. Syst.* ii. 911.—Afr. bor.
pinnatifida, *Desf. Fl. Atlant.* ii. 95. t. 165.—Hispan. Afr. bor.
Pollichii, Shuttl. in Mag. Zool. Bot. ii. (1838) 531 = B. Cheiranthos.
polymorpha, Murr. in Nov. Comm. Gott. vii. (1777) 35. t. 6 = juncea.
polymorpha, Syme, Engl. Bot. ed. III. i. 133 = campestris.
praecox, Waldst. & Kit. ex DC. Syst. ii. 593 = campestris.
Preslii, *Janka, in Termész. Füzetek*, vi. (1882) 182.— Europ.
Prolongi, *Boiss. Voy. Espagne*, ii. 38.—Hispan.
propera, Jord. Diagn. i. 183 = B. Cheiranthos.
pseudoerucastrum, *Brot. Fl. Lusit.* i. 581.—Lusitan.
psammophila, *Pomel, Nouv. Mat. Fl. Atl.* 363.— Algeria.
purpurascens, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 258.—Syria.
purpurea, Mill. Gard. Dict. ed. VIII. n. 6 = Moricandia arvensis.
pyrenaea, Jord. Diagn. i. 187 = B. Cheiranthos.
quadrivalvis, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 170.—Ind. or. (cult.).
racemiflora, Jord. Diagn. i. 184 = B. Cheiranthos.
Rapa, Linn. Sp. Pl. 666 = campestris.
rectangularis, *Viv. in Linnaea*, i. (1826) 502.—Corsica.
recurvata, Jord. Diagn. i. 182 = B. Cheiranthos.
repanda, *DC. Syst.* ii. 598.—Gallia.
retrocensis, *Ball, in Journ. Bot.* xi. (1873) 299.— Marocco.
Richerii, Lange, in Fl. Dan. xlv. t. 2658 = lanceolata, *Lange*.
Richerii, *Vill. Prosp.* 40.—Gallia.
Robertiana, *J. Gay, in Ann. Sc. Nat.* Sér. I. vii. (1826) 416.—Europ.
rostrata, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 500.—Ins. Socotr.
Rouyana, *Janka, in Termész. Füzetek*, vi. (1882) 181. —Europ.
rubra, Hort. ex Steud. Nom. ed. II. i. 224 = oleracea.
rupestris, *Rafin. Caratteri*, n. 191.—Sicil.
rupicola, *Pomel, Nouv. Mat. Fl. Atl.* 362.—Algeria.
sabellica, Hort. ex Pers. Syn. ii. 206, in syn. = oleracea.
sabularia, *Brot. Fl. Lusit.* i. 582.—Lusitan.; Afr. bor.

BRASSICA:—
sabularia, Willk. ex Willk. & Lange, Prod. Fl. Hisp. iii. 854 = oxyrrhina.
Saruna, Siebold, in Verh. Bat. Gen. xii. (1830) 52 = campestris?
Schimperi, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 86.—Arabia.
scoposa, *Janka, in Termész. Füzetek*, vi. (1882) 179.—Europ.
sempervirens, *Tenore, ex Schrank, Hort. Monac.* t. 10.—Europ. austr.
setigera, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* iii. 857.—Hispan.
sinaica, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 85 = Moricandia sinaica.
Sinapistrum, *Boiss. Voy. Espagne*, ii. 39.—Europ.
Sinapis, *Vis. Fl. Dalm.* iii. 136.—Europ. (Cosmop. cult.).
sisymbrioides, Fisch. ex DC. Prod. i. 217 = Tournefortii.
sphaerorhiza, S. F. Gray, Nat. Arr. Brit. Pl. ii. 683 = campestris.
spinescens, *Pomel, Nouv. Mat. Fl. Atl.* 364.—Algeria.
Stocksii, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 171 = Tournefortii.
stricta, Nestl. ex DC. Syst. ii. 593 = campestris.
strigosa, *DC. l. c.* 603.—Afr. austr.
subhastata, Willd. Sp. Pl. iii. 550 = Sisymbrium orientale.
subspinescens, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1839) II. 145 = Sisymbrium subspinescens.
subularia, Pers. Syn. ii. 207 (sphalm.) = sabularia.
suffruticosa, Desf. Fl. Atlant. ii. 94 = Moricandia arvensis.
sylvestris, Mill. Gard. Dict. ed. VIII. n. 4. = oleracea.
taraxacifolia, Boiss. Elench. 11 = Sisymbrium crassifolium.
tenuifolia, Baill. Hist. Pl. iii. 227 = Diplotaxis tenuifolia.
teretifolia, Desf. Fl. Atlant. ii. 94 = Moricandia teretifolia.
Tigridis, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 87 = deflexa.
timoriana, *F. Muell. Papuan Pl.* ii. 4.—Ins. Timor.
Tinei, *Tod. Ind. Sem. Panorm.* (1861) 32.—Hab.?
torulosa, Dur. in Duch. Rev. Bot. ii. (1847) 434 = Diplotaxis siifolia.
Tournefortii, *Gouan, Illustr.* 44. t. 20. f. A.—Reg. Mediterr.; Oriens; Ind. or.
trilocularis, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 169.—Reg. Himal.
trimestris, Boenn. Prod. Fl. Monast. 201 = campestris.
tuberosa, Salisb. Prod. 272 = campestris.
turgida, Pers. Syn. ii. 207 = Eruca sativa.
Turrita, Weig. Obs. 32 = Conringia orientalis.
Uechtritziana, *Janka, in Termész. Füzetek*, vi. (1882) 182.—Europ.
valentina, *DC. Syst.* ii. 603.—Hispan.
varia, *Dur. in Duch. Rev. Bot.* ii. (1847) 434.—Afr. bor.
vesicaria, Linn. Sp. Pl. 668 = Eruca vesicaria.
villosa, Bivon. ex Spreng. Syst. iv. Cur. Post. 243 = incana.
viminea, Boiss. Voy. Espagne, ii. 33 = Diplotaxis viminea.
violacea, Burm. f. Fl. Ind. 140 = campestris.
violacea, *Linn. Sp. Pl.* 667 (sp. dub.).—China.
virgata, Boiss. Voy. Espagne, ii. 35 = Diplotaxis virgata.
Washitana, Muhl. Cat. 63 = Streptanthus maculatus.
Willdenovii, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 88 = juncea.

BRASSICARIA, Pomel, Mat. Fl. Atl. 15 (1860) = **Brassica**, Tourn. (Crucifer.).
humilis, Lamotte, Prod. i. 87 = B. humilis.

BRASSICASTRUM, Link, Handb. ii. 318 (1831) = **Brassica**, Tourn. (Crucifer.).
elongatum, Link, l. c. = B. elongata.

BRASSICELLA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 330 = **Brassica**, Tourn. (Crucifer.).
arenosa, Fourr. l. c. = B. Cheiranthos.
Cheiranthos, Fourr. l. c. = B. Cheiranthos.

BRATHYDIUM, Spach, Hist. Vég. Phan. v. 442 (1836); et in Ann. Sc. Nat. Sér. II. v. (1836) 365 = **Hypericum**, Tourn.
canadense, Spach, l. c. 446 = H. pyramidatum?
Chamaerinum, Steud. Nom. ed. II. i. 224, sphalm. = seq.
Chamaenerium, Spach, Hist. Vég. Phan. v. 445 = H. sphaerocarpum.
grandiflorum, Spach, l. c. 443 = H. dolabriforme.
hyssopifolium, Spach, l. c. 445 = H. cistifolium.
sphaerocarpum, Spach, l. c. 444 = H. sphaerocarpum.

BRATHYS, Mutis, ex Linn. f. Suppl. 43 (1781) = **Hypericum**, Tourn.
acerosa, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 366 = H. laricifolium.
acicularis, Spach, l. c. = H. aciculare.
Billardieri, Spach, l. c. 367 = H. gramineum.
caespitosa, Blume, Mus. Bot. Lugd. Bat. ii. 20 = H. japonicum.
canadensis, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. canadense.
caracasana, Spach, l. c. 366 = H. caracasanum.
connata, Spach, l. c. = H. connatum.
debilis, Blume, Mus. Bot. Lugd. Bat. ii. 20 = H. japonicum.
denticulata, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. denticulatum.
Drummondii, Spach, l. c. = H. Drummondii.
Erythraea, Spach, l. c. 367 = H. angulosum.
euphorbioides, Spach, l. c. = H. mutilum?
Forsteri, Spach, l. c. = H. gramineum.
gentianoides, Spach, l. c. = H. Sarothra.
humifusa, Spach, l. c. = H. japonicum.
indecora, Spach, l. c. = H. thesiifolium.
japonica, Blume, Mus. Bot. Lugd. Bat. ii. 19 = H. japonicum.
juniperina, Linn. f. Suppl. 268 = H. Brathys.
lanceolata, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. angulosum.
laricifolia, Spach, l. c. 366 = H. laricifolium.
laxa, Blume, Mus. Bot. Lugd. Bat. ii. 19 = H. japonicum.
linoides, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. angulosum.
moranensis, Spach, l. c. = H. moranense.
Mutisiana, Spach, l. c. 366 = H. mexicanum.
nepalensis, Blume, Mus. Bot. Lugd. Bat. ii. 19 = H. japonicum.
oryzetum, Blume, l. c. 20 = H. japonicum.
paniculata, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 367 = H. paniculatum.
pauciflora, Spach, l. c. = H. pauciflorum.
prolifica, Payer, Organ. 8. t. 1 = H. densiflorum.
quinquenervia, Spach, l. c. = H. mutilum.
radicans, Blume, Mus. Bot. Lugd. Bat. ii. 20 = H. japonicum.
struthiolaefolia, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 366 = H. struthiaefolium.
tarquensis, Spach, l. c. 367 = H. thesiifolium.
thesiifolia, Spach, l. c. = H. thesiifolium.
thuyoides, Spach, l. c. 366 = H. thuyoides.
thymifolia, Spach, l. c. = H. Humboldtianum.
tomentosa, Spach, l. c. 367 = H. setosum.

BRAUNEA, Willd. Sp. Pl. iv. 797 (1805), *ex parte* = **Tiliacora**, Colebr. (Menisperm.).
menispermoides, Willd. l. c. = T. racemosa.

BRAUNERIA, Neck. Elem. i. 17 (1790) = **Echinacea**, Moench (Compos.).

BRAUNLOWIA, A. DC. Prod. xvii. 290 (1873) = **Brownlowia**, Roxb. (Tiliac.).

BRAVAISIA, DC. in Bibl. Univ. xvii. (1838) 124 (Rev. Bign. 16). *ACANTHACEAE*, Benth. & Hook. f. ii. 1084.
ONYCHACANTHUS, Nees, in DC. Prod. xi. 217 (1847).
floribunda, *DC. Prod.* ix. 240.—Mexic.

BRAVOA, Lex. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 6 (1824). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 737.
COETOCAPNIA, Link & Otto, Ic. Pl. Rar. 35 (1828).
ZETOCAPNIA, Link & Otto, l. c. i. t. 18 (sphalm.) (1828).

BRAVOA:—
Bulliana, *Baker, in Gard. Chron.* (1884) II. 328.—Am. austr.
Coetocapnia, M. Roem. Syn. Ensat. 245 = geminiflora.
geminiflora, *Lex. in La Llave & Lex. Nov. Veg. Desc.* fasc. i. 6.—Mexic.
sessiliflora, *Hemsl. Diag. Pl. Nov.* 54.—Mexic.

BRAXIPIS, Rafin. Sylva Tellur. 72 (1838) = **Sterculia**, Linn.
grandiflora, Rafin. l. c. = S. grandiflora.
nitida, Rafin. l. c. 73 = Cola acuminata.

BRAXIREON, Rafin. Fl. Tellur. iv. 23 (1836) = **Tapeinanthus**, Herb. (1837) (Amaryll.).
humile, Rafin. l. c. = T. humile.

BRAXYLIS, Rafin. Sylva Tellur. 51 (1838) = **Ilex**, Tourn.
obcordata, Rafin. l. c. = I. obcordata.

BRAYA, Sternb. & Hoppe, in Denkschr. Regensb. i. 1. (1815) 65. *CRUCIFERAE*, Benth. & Hook. f. i. 82.
APHRAGMUS, Andrz. ex DC. Prod. i. 209 (1824).
EUDEMA, Humb. & Bonpl. Pl. Aequin. ii. 133. t. 135 (1809), nomen prius.
OREAS, Cham. & Schlecht. in Linnaea, i. (1826) 29. t. 1.
OROBIUM, Reichb. Consp. 185 (1828).
PLATYPETALUM, R. Br. in Parry, Voy. App. 266 (1824).
aenea, Bunge, Del. Sem. Hort. Dorp. (1841) 8 = alpina.
alpina, *Sternb. & Hoppe, in Denkschr. Regensb.* i. 1. (1815) 66.—Alp. Europ.; Reg. arct.; Tibet.
arctica, Hook. in Parry, Voy. App. 387 = alpina.
calycina, *Wedd. Chlor. And.* t. 85.—Reg. Andin.
dubia, R. Br. ex Hook. f. in Trans. Linn. Soc. xxiii. (1861) 319 = alpina.
Eschscholtziana, *Benth. & Hook. f. ex S. Wats. Bibl. Ind. N. Am. Bot.* 51.—Am. arct.
falcata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 17 = Sisymbrium falcatum.
glabella, Richards. in Frankl. 1st Journ. 743 = alpina.
imbricatifolia, A. Gray, Bot. U. St. Expl. Exped. i. 58 = Sisymbrium imbricatifolium.
involucrata, *Ledeb. Fl. Ross.* i. 194.—Sibir. altaic.
Limosella, Bunge, Del. Sem. Hort. Dorp. (1841) 8 = Sisymbrium Limosella.
limoselloides, Bunge, ex Ledeb. Fl. Ross. i. 194 = Sisymbrium Limosella.
Meyeri, Bunge, ex Walp. Rep. ii. 755 (= Parryae sp.).—Sibir. altaic.
nana, *Bunge, Del. Sem. Hort. Dorp.* (1841) 8.—Sibir. baical.
Novae-Zelandiae, *Hook. f. Handb. New Zeal. Fl.* 13.—N. Zel.
oregonensis, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 199.—Am. bor. occ.
oxycarpa, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 169 = alpina.
pilosa, *Hook. Fl. Bor. Am.* i. 65. t. 17.—As. et Am. arct.
pinnatifida, Koch, Syn. Fl. Germ. ed. I. 50 = Sisymbrium pinnatifidum.
purpurascens, Bunge, ex Ledeb. Fl. Ross. i. 195 = alpina.
pusilla, *A. Gray, Bot. U. St. Expl. Exped.* i. 57.—Chili.
rosea, *Bunge, Del. Sem. Hort. Dorp.* (1841) 8.—As. arct.; Reg. altaic. et Himal.
Scharnhorsti, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 241.—Turkestan.
siliquosa, *Bunge, Del. Sem. Hort. Dorp.* (1839) 7.—*Cf. Linnaea*, xiv. 118.—Sibir. altaic.
supina, Koch, Syn. Fl. Germ. ed. I. 50 = Sisymbrium supinum.
Thomsoni, Hook. f. in Journ. Linn. Soc. v. (1861) 168 = alpina.
tibetica, *Hook. f. & Thoms. l. c.*—Tibet. occ.
Tilingii, *Regel & Tiling, in Nouv. Mém. Soc. Nat. Mosc.* xi. (1859) 61.—Ross. bor.
uniflora, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 168.—Tibet. occ.
versicolor, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 281.—Sibir. baical.

BRAYERA, Kunth, in Brayer, Not. Vermif. (1824). *ROSACEAE*, Benth. & Hook. f. i. 622.
BANKESIA, Bruce, Travels, v. 22, 23 (1790); Ed. II. vii. 181 (1805).
BRACERA, Engelm. in Buchner, Rep. Pharm. lxviii. (1839) 369.
HAGEA, Poir. in Dict. Sc. Nat. xii. 253 (1818).
HAGENIA, J. F. Gmel. Syst. 613 (1791).
anthelmintica, *Kunth, l. c.*—Abyssin.

BRAZORIA, Engelm. & Gray, in Bost. Journ. Nat. Hist. v. (Pl. Lindh.) 255 (1845). *LABIATAE*, Benth. & Hook. f. ii. 1204.
Roemeriana, *Scheele, in Linnaea*, xxii. (1849) 590.—Texas.
scutellarioides, *Engelm. & Gray, in Bost. Journ. Nat. Hist.* v. (1845) 257.—Texas.
truncata, *Engelm. & Gray, l. c.* 256.—Texas.

BREBISSONIA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 175 = **Fuchsia**, Linn. (Onagrar.).
microphylla, Spach, l. c. = F. microphylla.
thymifolia, Steud. Nom. ed. II. i. 225 = F. thymifolia.

BREDEMEYERA, Willd. in Ges. Naturf. Freunde Neue Schr. iii. (1801) 412. t. 6. *POLYGALEAE*, Benth. & Hook. f. i. 138.
CATOCOMA, Benth. in Hook. Journ. Bot. iv. (1842) 101.
HUALANIA, Phil. in Linnaea, xxxiii. (1864-65) 18.
acuminata, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 51 = myrtifolia.
acuminata, *Poepp. ex Klotzsch, in Peters, Reise Mossamb. Bot.* 117.—Afr. trop.
altissima, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 50. —Am. austr.
bracteata, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 187.—Guiana.
brevifolia, *Klotzsch, ex A. W. Benn. in Mart. Fl. Bras.* xiii. III. 49.—Bras.
cuneata, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 187.—Guiana.
densiflora, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 52.—Bras.
floribunda, *Willd. in Ges. Naturf. Freunde Neue Schr.* iii. (1801) 412.—Am. austr.
Hilaireana, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 52 = laurifolia.
Kunthiana, *Klotzsch, ex A. W. Benn. l. c.* 53.—Bras.
laurifolia, *Klotzsch, ex A. W. Benn. l. c.* 52.—Bras.
Lhotzkyana, Klotzsch, in Peters, Reise Mossamb. Bot. 117 = floribunda.
lucida, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 51.—Bras.
lucida, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 189.—Guiana.
Mansonii, Klotzsch, in Peters, Reise Mossamb. Bot. 117 = lucida.
Martiana, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 49. —Bras.
microphylla, *Hieron. in Roca, Exped. Rio-Negro*, ii. 182.—Patagonia.
Moritziana, *Klotzsch, ex Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 188.—Guiana.
myrtifolia, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 50.—Bras.
parviflora, *A. W. Benn. l. c.* 51.—Am. trop.
revoluta, *A. W. Benn. l. c.* 49.—Bras.
Sellowii, *Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 189.—Bras.
velutina, *A. W. Benn. in Mart. Fl. Bras.* xiii. III. 53. —Bras.

BREDIA, Blume, Mus. Bot. Lugd. Bat. i. 24. f. 4 (1849). *MELASTOMACEAE*, Benth. & Hook. f. i. 753.
hirsuta, *Blume, l. c.* 25.—Japon.
Oldhami, *Hook. f. Ic. Pl.* t. 1085.—Ins. Formosa.

BREEA, Less. Syn. Comp. 9 (1832) = **Cnicus**, Linn. (Compos.).
dioica, Less. l. c. 10 = C. arvensis.

BREHMIA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 25 = **Strychnos**, Linn. (Logan.).
spinosa, Harv. l. c. 26 = S. spinosa.

BREHMIA, Schrank, in Syll. Ratisb. i. (1824) 85 = **Pavonia**, Cav. (Malvac.).
arborescens, Schrank, l. c. = P. velutina.

BREHNIA, Baker, in Journ. Linn. Soc. xiv. (1875) 561 sphalm. = **Behnia**, Didrichs. (Liliac.).

BREMONTIERA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 93. *LEGUMINOSAE*, Benth. & Hook. f. i. 464.
Ammoxylon, *DC. l. c.*—Ins. Mascar.

BREONIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 290. *RUBIACEAE*, Benth. & Hook. f. ii. 32.
madagascaricnsis, *A. Rich. l. c.*—Madag.

BREPHOCTON, Rafin. Fl. Tellur. ii. 49 (1836) = **Rochonia**, DC. (Compos.).
cuneatum, Rafin. l. c. = R. cuneata.

BRETEUILLIA, Buchoz, ex DC. Prod. vi. 503, in syn. (1837) = **Didelta**, Less. (Compos.).

BREVIGLANDIUM, Dulac, Fl. Hautes-Pyr. 423 (1867) = **Hottonia**, Linn. (Primul.).
palustre, Dulac, l. c. = H. palustris.

BREVOORTIA, Wood, in Proc. Acad. Sc. Philad. (1867) 81. *LILIACEAE*, Benth. & Hook. f. iii. 800.
coccinea, S. Wats. in Proc. Am. Acad. xiv. (1879) 239 = Ida-Maia.
Ida-Maia, *Wood, in Proc. Acad. Sc. Philad.* (1867) 82.—Calif.

BREWERIA, R. Br. Prod. 487 (1810). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 876.
BONAMIA, A. Gray, in Proc. Am. Acad. v. (1862) 337.
CALYCOBOLUS, Willd. ex Roem. & Schult. Syst. v. 2 (1819).
CODONANTHUS, G. Don, Gen. Syst. iv. 166 (1856).
DETHARDINGIA, Nees, in Nov. Act. Nat. Cur. xi. (1823) 80.
DUFOUREA, H. B. & K. Nov. Gen. et Sp. iii. 113. t. 214 (1818).
PREVOSTEA, Choisy, in Ann. Sc. Nat. Sér. I. iv. (1825) 497.
REINWARDTIA, Spreng. Syst. i. 527 (1825).
SEDDERA, Hochst. in Flora, xxvii. (1844) Bes. Beil. 7. t. 5.
STYLISMA, Rafin. Neogenyt. 2 (1825).
TRICHANTHA, Karst. & Triana, in Linnaea, xxviii. (1856) 437.
WILBERFORCIA, Hook. f. ex Planch. in Hook. Ic. Pl. t. 796 (1848).
abscissa, *Choisy, in DC. Prod.* ix. 438.—Madag.
africana, *Benth. & Hook. f. Gen.* iii. 877.—Afr. trop.
aquatica, *A. Gray, Syn. Fl. N. Am.* ii. 1. 217.—Am. bor.
brevifolia, *Benth. Fl. Austral.* iv. 436.—Austral.
Burchellii, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 493.—Bras.
Choisyana, *Steud. Nom.* ed. II. i. 224.—Ind. or.
cordata, *Blume, Bijdr.* 722.—Ind. or.; Malaya.
elegans, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 494.—Burma.
evolvuloides, Choisy, l. c. = Choisyana.
evolvuloides, *R. Br. in Salt, Abyss. App.* 65.—Abyss.
fastigiata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 83.—Ins. Socotr.
ferruginea, *Benth. & Hook. f. Gen.* iii. 877.—Bras.
glomerata, *Balf. f. l. c.*—Ins. Socotr.
grandiflora, *A. Gray, in Proc. Am. Acad.* xv. (1880) 51. —Florida.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880-82) 523. —Afr. trop.
hispida, *Franch, in Révoil, Çomali*, 43.—Afr. trop.
humistrata, *A. Gray, Syn. Fl. N. Am.* ii. 1. 217.—Am. bor.
Langsdorffii, *Meissn. in Mart. Fl. Bras.* vii. 325.—Bras.
latifolia, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 224.—Afr. bor.; Arab.; Ind. bor. occ
linearis, *R. Br. Prod.* 488.—Austral.
linifolia, Spreng. Syst. i. 614 = Wahlenbergia linarioides.
madagascariensis, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 493.—Madag.

BREWERIA:—
malvacea, Klotzsch, in Peters, Reise Mossamb. Bot. 245 = Convolvulus malvaceus.
media, *R. Br. Prod.* 488.—Austral.
Menziesii, *Benth. & Hook. f. Gen.* iii. 877.—Ins. Sandvic.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* ii. 400.—Mexic.
minima, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 228.—Calif.
ovalifolia, *A. Gray, Syn. Fl. N. Am.* ii. 1. 217.—Am. bor. occ.
oxycarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 76 = evolvuloides.
pannosa, *R. Br. Prod.* 488.—Austral.
parviflora, Arn. ex Steud. Nom. ed. II. i. 224 = Convolvulus parviflorus.
pedunculata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 83.—Ins. Socotr.
Pickeringii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 217.—Am. bor.
rosea, *F. Muell. Fragm.* i. 233.—Austral.
Roxburghii, Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 493 = cordata.
scoparia, *Lindl. Fl. Med.* 400.—Hab.?
secunda, *Benth. in Hook. Niger Fl.* 470 (err. typ. 70). —Afr. trop.
sericea, *Griseb. in Goett. Abh.* xix. (1874) 229.—Reg. Argent.
somalensis, *Vatke, in Linnaea*, xliii. (1880-82) 523.— Afr. trop.
spectabilis, *Choisy, in Mém. Soc. Phys. Genèv.* vi. (1833) 493.—Madag.
valerianoides, *Villar, in Blanco, Fl. Phil.* ed. III. *Nov. App.* 143.—Ins. Philipp.
venulosa, *Meissn. in Mart. Fl. Bras.* vii. 326.—Bras.
virgata, *Vatke, in Linnaea*, xliii. (1880-82) 523.— Afr. austr.

BREWERINA, A. Gray, in Proc. Am. Acad. viii. (1872) 620 = **Arenaria**, Rupp. (Caryophyll.).
suffrutescens, A. Gray, l. c. = A. congesta.

BREWSTERA, M. Roem. Syn. i. Hesp. 132, 141 (1846) = **Ixonanthes**, Jack (Lineae).
crenata, M. Roem. l. c. 141 = I. icosandra.

BREXIA, Noronha, ex Thou. Gen. Nov. Madag. 20 (1806). *SAXIFRAGEAE*, Benth. & Hook. f. i. 645.
VENANA, Lam. Illustr. ii. 99. t. 131 (1800), nomen prius.
acanthifolia, Noronha, ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 159 = madagascariensis.
amplifolia, Noronha, l. c. = madagascariensis.
cauliflora, *Tul. l. c.* 161.—Madag.
chrysophylla, Sweet, Hort. Brit. ed. I. 492 = madagascariensis.
digyna, Noronha, ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 159 = madagascariensis.
heterophylla, Boj. Hort. Maurit. 52 = madagascariensis.
Lucanana, *Lubbers, Cat. Pl. Rar.* 6.—Hab.?
madagascariensis, *Thou. ex Ker-Gawl. in Bot. Reg.* t. 730.—Madag.
microcarpa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 160.—Ins. Seychell.
ovatifolia, Noronha, ex Tul. l. c. 159 = madagascariensis.
serrata, Presl, Rep. Bot. Syst. i. 190 = madagascariensis.
spinosa, Lindl. in Bot. Reg. t. 872 = madagascariensis.

BREYNIA, Forst. Char. Gen. 145. t. 73 (1776). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 276.
FOERSTERIA, Scop. Introd. 98 (1777).
MELANTHESA, Blume, Bijdr. 590 (1825).
MELANTHESIOPSIS, Muell. Arg. in Linnaea, xxxii. (1863) 74.
acuminata, *Muell. Arg. in DC. Prod.* xv. II. 442.— Ins. Philipp.
axillaris, Spreng. Pugill. ii. 92 = disticha.
cernua, *Muell. Arg. in DC. Prod.* xv. II. 439.— Austral.; Ins. Timor.
cinerascens, Baill. Adansonia, vi. (1866) 344 = oblongifolia.
discigera, *Muell. Arg. in DC. Prod.* xv. II. 440.— Malaya.
disticha, *Forst. Char. Gen.* 146. t. 73.—N. Caled.

BREYNIA:—
microphylla, *Muell. Arg. in DC. Prod.* xv. II. 442.—Java.
Muelleriana, Baill. Adansonia, vi. (1866) 344 = stipitata.
oblongifolia, *Muell. Arg. in DC. Prod.* xv. II. 440.—Austral.
paniculata, Spreng. Pugill. ii. 93 (gen. dub.).—Ind. or.
racemosa, *Muell. Arg. in DC. Prod.* xv. II. 441.—Java.
rhamnoides, *Muell. Arg. l. c.* 440.—As. trop.
rhynchocarpa, *Benth. Fl. Austral.* vi. 114.—Austral.
rubra, *Muell. Arg. in DC. Prod.* xv. II. 438.—Java.
stipitata, *Muell. Arg. l. c.* 442.—Austral.
virgata, *Muell. Arg. l. c.* 441.—Java.

BREYNIA, [Plum.] Linn. Gen. ed. I. 113 (1737) = **Capparis**, Tourn.
indica, Linn. Sp. Pl. 503 = C. Breyniana.

BREZIA, Moq. in DC. Prod. xiii. II. 167 (1849) = **Suaeda**, Forsk. (Chenop.).
heterophylla, Moq. l. c. = S. heterophylla.

BRICKELLIA, Ell. Sketch, ii. 290 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 247.
Bulbostyle, DC. Prod. v. 138 (1836).
Clavigera, DC. l. c. 127 (1836).
Coleosanthus, Cass. in Bull. Soc. Philom. (1817) 67.
Ismaria, Rafin. Sylva Tellur. 117 (1838).
Rosalesia, La Llav. & Lex. Nov. Veg. Desc. i. 9 (1824).
atractyloides, *A. Gray, in Proc. Am. Acad.* viii. (1870) 290.—Am. bor. occ.
baccharidea, *A. Gray, Pl. Wright.* i. 87.—Texas; Arizona.
betonicaefolia, *A. Gray, l. c.* ii. 72.—N. Mexic.; Arizona.
brachyphylla, *A. Gray, l. c.* i. 84.—Am. bor. occ.
californica, *A. Gray, Pl. Fendl.* 64.—Am. bor. occ.
Cavanillesii, *A. Gray, Pl. Wright.* i. 85.—Mexic.
cedrosensis, *Greene, in Bull. Torrey Club*, x. (1883) 86.—Ins. Cedros.
cordifolia, *Ell. Sketch*, ii. 290.—Am. bor.
corymbosa, *A. Gray, Pl. Wright.* i. 84.—Mexic.
Coulteri, *A. Gray, l. c.* 86.—Am. bor. occ.
Cumingii, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 325 = Chrysopsis oregana.
cylindracea, *A. Gray, ex Engelm. in Proc. Am. Acad.* i. (1846) 46.—Texas.
dentata, *Sch. Bip. in Seem. Bot. Voy. Herald*, 301.—Texas; Arizona.
diffusa, *A. Gray, Pl. Wright.* i. 86.—Am. trop.
Fendleri, *A. Gray, Pl. Fendl.* 63.—N. Mexic.
floribunda, *A. Gray, Pl. Wright.* ii. 73.—Arizona.
frutescens, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 207.—Calif.
Galeottii, *A. Gray, Pl. Wright.* i. 85.—Mexic.
grandiflora, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 287.—Am. bor. occ.
Greenei, *A. Gray, in Proc. Am. Acad.* xii. (1877) 58.—Calif.
Hartwegi, *A. Gray, Pl. Wright.* i. 85.—Mexic.
hastata, *Benth. Bot. Voy. Sulph.* 21.—Calif.
hebecarpa, *A. Gray, Pl. Wright.* i. 85.—Mexic.
hymenochlaena, *A. Gray, in Proc. Am. Acad.* xv. (1880) 29.—Mexic.
incana, *A. Gray, l. c.* vii. (1868) 350.—Calif.
laciniata, A. Gray, Pl. Wright. i. 87 = dentata.
lanata, *A. Gray. l. c.* 84.—Mexic.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 206.—Calif.
linearifolia, Klatt, in Abh. Naturf. Ges. Halle, xv. (1882) 325 = Bigelowia Douglasii.
linifolia, *Eaton, in Bot. King, Exp.* 137. t. 15. f. 1–6.—Am. bor. occ.
longifolia, *S. Wats. in Am. Natural.* vii. (1873) 301.—Am. bor. occ.
microphylla, *A. Gray, Pl. Wright.* i. 85.—Am. bor. occ.
mohavensis, *A. Gray, Syn. Fl. N. Am.* i. II. 104.—Calif.
multiflora, *Kellogg, in Proc. Calif. Acad.* vii. (1877) 49.—Calif.
Nevinii, *A. Gray, in Proc. Am. Acad.* xx. (1885) 297.—Calif.

BRICKELLIA:—
oblongifolia, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 288.—Am. bor. occ.
odontophylla, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 206.—Mexic.
oliganthes, *A. Gray, Pl. Wright.* i. 84.—Arizona; Mexic.
Palmeri, *A. Gray, in Proc. Am. Acad.* xv. (1880) 30.—Mexic.
Parryi, *A. Gray, l. c.* 31.—Mexic.
parvula, *A. Gray, Pl. Wright.* i. 84.—Texas.
paucidentata, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 326.—Ins. S. Cruc.
pendula, *A. Gray, Pl. Wright.* i. 85.—N. Mexic.
pinifolia, *A. Gray, l. c.* 84.—Bras.
Pringlei, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 206.—Arizona.
reniformis, A. Gray, Pl. Wright. i. 86 = Wrightii.
Riddellii, *A. Gray, l. c.* 83.—Texas.
Rosalesia, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 106.—Mexic.
Rusbyi, *A. Gray, Syn. Fl. N. Am.* i. II. 106.—N. Mexic.
scoparia, *A. Gray, Pl. Wright.* i. 84.—Mexic.
secundiflora, *A. Gray, l. c.* 85.—Mexic.
Seemanni, *A. Gray, in Proc. Am. Acad.* xv. (1880) 30.—N. Mexic.
simplex, *A. Gray, Pl. Wright.* ii. 73.—Arizona.
spinulosa, *A. Gray, l. c.* i. 84.—Mexic.
squamulosa, *A. Gray, in Proc. Am. Acad.* xv. (1880) 30.—Mexic.
tenera, A. Gray, Pl. Wright. ii. 72 = Wrightii.
thyrsiflora, *A. Gray, in Proc. Am. Acad.* xv. (1880) 30.—Mexic.
tomentella, *A. Gray, Pl. Wright.* i. 85.—Mexic.
veronicaefolia, *A. Gray, l. c.*—Mexic.
Wislizeni, *A. Gray, Pl. Fendl.* 64.—Arizona; Mexic.
Wrightii, Durand & Hilg. in Pacif. Rail. Rep. v. 8 = californica.
Wrightii, *A. Gray, Pl. Wright.* ii. 72.—Am. bor. occ.

BRICKELLIA, Rafin. in Med. Repos. New York, v. (1808) 353 = **Gilia**, Ruiz & Pav. (Polemon.).

BRICOUR, Adans. Fam. ii. 423 (1763) = **Myagrum**, Tourn. (Crucifer.).

BRIDELIA, Willd. Sp. Pl. iv. 978 (1805) (Briedelia). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 267.
Candelabria, Hochst. in Flora, xxvii. (1843) 79.
Cluytia, Roxb. ex Steud. Nom. ed. II. i. 225 (1840).
Pentameria, Klotzsch, ex Baill. Etud. Gén. Euph. 584 (1858).
Zarcoa, Llanos, in Bot. Zeit. xv. (1857) 423.
acuminata, Wall. Cat. n. 7885 = Phyllanthus bicolor.
alnifolia, *Griff. Notul.* iv. 481.—Burma.
amoena, Wall. ex Baill. Etud. Gén. Euph. 584 = retusa.
angolensis, *Muell. Arg. in Seem. Journ. Bot.* ii. (1864) 327.—Afr. trop.
atroviridis, *Muell. Arg. l. c.*—Afr. trop.
attenuata, *Wall. ex Voigt, Hort. Suburb. Calc.* 156.—Ind. or.
Berneriana, *Baill. Adansonia*, ii. (1861–62) 39.—Madag.
Berryana, Wall. Cat. n. 7876, 7960 = Phyllanthus indicus.
buxifolia, *Baill. Adansonia*, ii. (1861–62) 230.—N. Caled.
cathartica, *Bertol. f. in Mem. Acc. Sc. Bolog.* v. (1854) 476.—Afr. trop.
chartacea, Wall. Cat. n. 7881 = Cleistanthus chartaceus.
coccolobaefolia, *Baker, in Journ. Linn. Soc.* xxi. (1885) 441.—Madag.
collina, Hook. & Arn. Bot. Beech. Voy. 211 = Cleistanthus collinus.
crenulata, Roxb. Hort. Beng. 70; Fl. Ind. iii. 734 = retusa.
dasycalyx, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 241.—Burma.
diversifolia, *Hook. & Arn. Bot. Beech. Voy.* 211.—Malaya.
elegans, *Muell. Arg. in Seem. Journ. Bot.* ii. (1864) 327.—Afr. occ.

BRIDELIA:—
exaltata, *F. Muell. Fragm.* iii. 32.—Austral.
faginea, *F. Muell. ex Benth. Fl. Austral.* vi. 120.—Austral.
ferruginea, *Benth. in Hook. Niger Fl.* 511.—Afr. trop.
gambecola, *Baill. Adansonia*, i. (1860–61) 79.—Afr. trop.
glauca, *Blume, Bijdr.* 597.—Java.
glauca, Wall. Cat. n. 7875 = praec. ? (Glochidionis sp.).—Penang.
Hamiltoniana, *Wall. l. c.* 7882.—Ind. or.
heterantha, Wall. l. c. 7873 = Glochidion Wallichianum.
horrida, Dillw. Review Hort. Malab. 15 = Scleropyron Wallichianum.
insulana, *Hance, in Journ. Bot.* xv. (1877) 337.—China.
lancaefolia, Roxb. Hort. Beng. 70; Fl. Ind. iii. 737 = tomentosa.
lanceolata, *Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 45.—Malaya.
lancifolia, Buch.-Ham. in Wall. Cat. n. 7884 = tomentosa.
laurina, *Baill. Adansonia*, ii. (1861–62) 229.—N. Caled.
Leichardtii, Baill. Etud. Gén. Euph. 584 = faginea.
Loureirii, Hook. & Arn. Bot. Beech. Voy. 211 = tomentosa.
melanthesoides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 103.—Afr. trop.
micrantha, *Baill. Adansonia*, iii. (1862–63) 164.—Afr. trop.
montana, Wall. Cat. n. 7879 = retusa.
montana, *Willd. Sp. Pl.* iv. 978.—Ind. or.
Moonii, *Thw. Enum. Pl. Zeyl.* 279.—Zeylan.
multiflora, *Zipp. in Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 119.—Amboina.
oblongifolia, Hook. & Arn. Bot. Beech. Voy. 212 = Cleistanthus chartaceus.
ovata, *Decne. in Nouv. Ann. Mus. Par.* iii. (1835) 484.—Austral.; Burma; Malaya; Ins. Timor.
patula, Hook. & Arn. Bot. Beech. Voy. 212 = Cleistanthus patulus.
Pervilleana, *Baill. Adansonia*, ii. (1861–62) 38.—Madag.
polystachya, Baill. Etud. Gén. Euph. 584 = Cleistanthus polystachyus.
pubescens, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 241.—Reg. Himal.; Burma.
retusa, Baill. Etud. Gén. Euph. 584, nomen = Moonii.
retusa, A. Juss. Tent. Euphorb. 27. t. 7 = stipularis.
retusa, *Spreng. Syst.* iii. 48.—Ind. or.; Malaya.
rhamnoides, Griff. Notul. iv. 480 = tomentosa.
rhomboidalis, *Baill. Adansonia*, ii. (1861–62) 37.—Ins. Borbon.
rugosa, *Miq. Fl. Ind. Bat. Suppl.* 445.—Sumatra.
scandens, Willd. Sp. Pl. iv. 979 = stipularis.
scleroneura, *Muell. Arg. in Flora*, xlvii. (1864) 515.—Afr. trop.
sinica, J. Grah. Cat. Bomb. Pl. 179 = Cleistanthus malabaricus?
speciosa, *Muell. Arg. in Seem. Journ. Bot.* ii. (1864) 327.—Afr. trop.
spinosa, Hort. ex DC. in Mém. Soc. Phys. Genèv. vi. (1833) 563 = Damnacanthus indicus.
spinosa, Willd. Sp. Pl. iv. 979 = retusa.
stenocarpa, *Muell. Arg. in Flora*, xlvii. (1864) 515.—Afr. trop.
stipitata, *Baill. Adansonia*, ii. (1861–62) 229.—N. Caled.
stipularis, *Blume, Bijdr.* 597.—Ind. or.; Malaya; Afr. trop.
stipularis, Hook. & Arn. Bot. Beech. Voy. 211 = Cleistanthus stipularis.
tenuifolia, *Muell. Arg. in Journ. Bot.* ii. (1864) 328.—Afr. trop.
tomentosa, *Blume, Bijdr.* 597.—Ind. or.; Malaya; Austral.
Tulasneana, *Baill. Adansonia*, ii. (1861–62) 40.—Madag.
urticoides, Griff. Notul. iv. 481 = tomentosa?
Zollingeri, Miq. Fl. Ind. Bat. i. II. 364 = stipularis.

BRIDGESIA, Bert. ex Cambess. in Nouv. Ann. Mus. Par. iii. (1834) 234. t. 13. *SAPINDACEAE*. Benth. & Hook. f. i. 393.
Tripterocarpus, Meissn. Gen. 52 (1837).
incisifolia, *Bert. ex Cambess. l. c.* 235.—Chili.

BRIDGESIA, Hook. Bot. Misc. ii. (1831) 222. t. 92 = **Polyachyrus**, Lag. (Compos.).
echinopsoides, Hook. l. c. = P. sphaerocephalus.

BRIDGESIA, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 168. t. 102 = **Ercilla**, A. Juss. (Phytolacc.).
spicata, Hook. & Arn. l. c. 169 = E. volubilis.

BRIEDELIA, Willd. Sp. Pl. iv. 978 (1805) = **Bridelia**, 'Willd.' (Euphorb.).

BRIGHAMIA, A. Gray, in Proc. Am. Acad. vii. (1867) 185. *CAMPANULACEAE*, Benth. & Hook. f. ii. 545.
insignis, *A. Gray, l. c.* —Ins. Hawaii.

BRIGNOLIA, Bertol. in Desv. Journ. Bot. ii. (iv.) (1815) 76 = **Kundmannia**, Scop. (Umbellif.).
panacifolia, Heynh. Nom. i. 131, sphalm. = scq.
pastinacifolia, Bertol. in Desv. Journ. ii. (iv.) (1815) 76 = K. sicula.

BRIGNOLIA, DC. Prod. iv. 444 (1830) = **Isertia**, Schreb. (Rubiac.).
acuminata, DC. l. c.—Ins. Trinit.
pubigera, Benth. in Hook. Journ. Bot. iii. (1840) 219. —Guiana.

BRILLANTAISIA, Beauv. Fl. Owar. ii. 67. t. 100 (1805). *ACANTHACEAE*, Benth. & Hook. f. ii. 1076.
BELANTHERIA, Necs, in DC. Prod. xi. 96 (1847).
LEUCORHAPHIS, Nees, l. c. 97 (1847).
alata, *T. Anders. ex Oliver, in Trans. Linn. Soc.* xxix. (1875) 125.—Afr. trop.
hirsuta, *T. Anders. in Journ. Agri-Hort. Soc. Ind.* N. S. i. (1868) 267.—Afr. trop.
Lamium, *Benth. in Hook. Niger Fl.* 477.—Afr. trop.
owariensis, *Beauv. Fl. Owar.* ii. 68. t. 100. f. 2.—Afr. trop.
patula, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 21. —Afr. trop.
pubescens, *T. Anders. ex Oliver, in Trans. Linn. Soc.* xxix. (1875) 125.—Afr. trop.
Rutenbergiana, *Vatke, in Bremen Abh.* ix. (1885) 131. —Afr. trop.
Vogeliana, *Benth. in Hook. Niger Fl.* 477.—Afr. trop.

BRIMEURA, Salisb. Gen. Pl. Fragm. 26 (1866) = **Hyacinthus**, Tourn. (Liliac.).
amethystina, Salisb. l. c. = H. amethystinus.

BRIMYS, Scop. Introd. 252 (1777) err. typ. = **Drimys**, Forst. (Magnol.).

BRINDONIA, Thou. in Dict. Sc. Nat. v. 339 (1804) = **Garcinia**, Linn. (Guttif.).
celebica, Thou. l. c. 341 = G. cornea ?
cochinchinensis, Thou. l. c. = G. cochinchinensis.
indica, Thou. l. c. 340 = G. indica.

BRISEGNOA, Remy, in C. Gay, Fl. Chil. v. 291 (1849) = **Oxytheca**, Nutt. (Polygon.).
chilensis, Remy, l. c. 292 = O. dendroidcs.

BRISEIS, Salisb. Gen. Pl. Fragm. 92 (1866) = **Allium**, Tourn. (Liliac.).
triquetra, Salisb. l. c. 93 = A. triquetrum.

BRISSONIA, Neck. Elem. iii. 36 (1790); Desv. [Journ. i. (1814) 78]; in Ann. Sc. Nat. Sér. I. ix. (1826) 411 = **Indigofera**, Linn. et **Tephrosia**, Pers. (Legumin.).
coronillaefolia, Desv. Journ. i. (1814) 78 = T. coronillaefolia.
stipularis, Desv. l. c. = T. stipularis.
trapesicarpa, Desv. l. c. = seq. ?
trapezicarpa, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 411 = I. senegalensis.

BRITOA, Berg, in Linnaea, xxvii. (1854) 435 = **Campomanesia**, Ruiz & Pav. (Myrtac.).
acida, Berg, l. c. 436; et in Mart. Fl. Bras. xiv. t. 461. —Bras.
dichotoma, Berg, ll. cc. 436, 463.—Bras.
eriantha, Berg, ll. cc. = C. eriantha.
psidioides, Berg, ll. cc. 436, 462.—Bras.
rugosa, Berg, in Mart. Fl. Bras. xiv. t. 613.—Bras.
Sellowiana, Berg, in Linnaea, xxvii. (1854) 437; et in Mart. Fl. Bras. xiv. t. 463.—Bras.
sessiliflora, Berg, ll. cc. 436, 461.—Bras.
triflora, Berg, ll. cc. 436, 462 = C. triflora.

BRIZA, Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1194.
CALOTHECA et CHASCOLYTRUM, Desv. in Nov. Bull. Soc. Philom. ii. (1810) 190.
POA, Adans. Fam. ii. 34 (1763).
TREMULARIA, Heist. Syst. 12 (1748).
anceps, Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 44 = media.
aspera, Knapp, Gram. Brit. t. 61 = minor.
auriculata, Trev. ex Steud. Syn. Pl. Gram. 283 = triloba.
barbata, *Trin. in Mém. Acad. Pétersb.* Sér. VI. i. (1831) 363.—Bras.
Berteroniana, *Steud. Syn. Pl. Gram.* 283.—Chili.
bipinnata, Linn. Syst. Nat. x. 875 = Eragrostis cynosurioidcs.
canadensis, *Michx. Fl. Bor. Am.* i. 71.—Am. bor.
canadensis, Nutt. Gen. Am. i. 69 pro parte = Glyceria nervata.
capensis, Schrank, Hort. Monac. t. 43 = maxima.
capensis, Thunb. Prod. Fl. Cap. 21 = Eragrostis brizoidcs.
caroliniana, Lam. Encyc. i. 465; Illustr. i. 187. t. 45. f. 3 = Uniola paniculata.
caroliniana, Walt. Fl. Carol. 79 = Eragrostis major.
Clusii, Schult. in Roem. & Schult. Mant. ii. 294 = media.
cynosuroides, Scop. Delic. Insub. ii. 21 = Demazeria sicula.
deltoidea, *Burm. f. Fl. Cap. Prod.* 3.—Afr. austr.
disticha, Moench, Meth. 185 = Demazeria sicula.
dura, *Desv. Opusc.* 105.—Am. austr.
elatior, Sibth. & Sm. Fl. Graec. i. t. 75 = media.
elegans, *Doell, in Mart. Fl. Bras.* ii. III. 135.—Bras.
Eragrostis, Desf. Fl. Atlant. i. 78 = Eragrostis minor.
Eragrostis, Linn. Sp. Pl. 70 = Eragrostis major.
erecta, *Lam. Illustr. Gen.* i. 187.—Reg. Argent.
flava, *Desv. Opusc.* 105.—Am. austr.
geniculata, *Thunb. Prod. Pl. Cap.* 21.—Afr. austr.
grandis, Salisb. Prod. 21 = maxima.
humilis, Bieb. Fl. Taur. Cauc. i. 66 = spicata.
imbricata, Steud. in Flora, xii. (1829) 489 = Poa Uniolae.
Lamarkiana, Cham. & Schlecht. in Linnaea, vi. (1831) 39 = rotundata.
Lamarckiana, *Nees, Agrost. Bras.* 481.—Am. austr.
lutescens, [Foucault, in] Desv. Journ. iii. (1814) t. 24. f. 2 = media.
macrostachya, *Steud. Nom.* ed. II. i. 225.—Peruv.
major, Desv. ex Kunth, Enum. Pl. i. 371 = pulchella.
major, Presl, Cyp. et Gram. Sic. 42 = maxima.
maxima, *Linn. Sp. Pl.* 70.—Reg. Mediterr.; Afr. austr.
media, *Linn. l. c.*—Europ.; Oriens; As. bor. (Amphig. advcnt.).
megastachya, Hort. ex Steud. Nom. ed. II. i. 225 = Eragrostis major.
microstachya, *Steud. l. c.*—Peruv.
minima, Hort. ex Nichols. Dict. Gard. i. 212 = minor.
minor, *Linn. Sp. Pl.* 70.—Europ.; Oriens; As. bor. (Amphig. advent.).
monspessulana, Gouan, Hort. Monsp. 45 = maxima.
montana, *Brouss. Cat. Monsp.* (1804); *Roem. & Schult. Syst.* ii. 521.—Teneriffa ?
mucronata, Lam. Encyc. i. 465 = Lamarckiana.
multiflora, *Forst. ex Beauv. Agrost.* 155, nomen.—Oriens.
Neesii, *Doell, in Mart. Fl. Bras.* ii. III. 132.—Bras.
nigra, *Burch. ex Steud. Nom.* ed. II. i. 225.—Afr. austr.
oblonga, Moench, Meth. 185 = Eragrostis major.
pauciflora, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 86.—Europ.
pilosa, *Schur, Enum. Pl. Transs.* 766.—Hungaria.
Poa, Nees, ex Steud. Syn. Pl. Gram. 283 = Necsii.
pulchella, *Kunth, Rev. Gram.* i. 120.—Ins. Teneriff.
purpurascens, *Muhl. Descr. Uber.* 154.—Carolina.
reniformis, *Steud. Nom.* ed. II. i. 225.—Peruv.
rotundata, *Steud. Syn. Pl. Gram.* 284.—Mexic.
rubella, *Steud. l. c.* 282.—Afr. trop.
rubens, Poir. Suppl. i. 699 = maxima.
rubra, Lam. Illustr. i. 187; Pers. Syn. i. 93 = maxima.
rufa, Steud. Nom. ed. II. i. 225.—Peruv.
scabra, Nees, ex Steud. Syn. Pl. Gram. 276. n. 191 = Necsii.
Sellowii, *Nees, ex Steud. l. c.* 284. n. 26.—Bras.
serotina, Dum. Obs. Gram. Belg. 110 = media.
spicata, *Burm. f. Fl. Cap. Prod.* 3.—Afr. austr.

BRIZA :—
spicata, Lam. Encyc. i. 465 = Poa Michauxii.
spicata, *Sibth. & Sm. Fl. Graec.* i. t. 77.—Graecia; As. Min.
spicigera, *Steud. Nom.* ed. II. i. 225.—Peruv.; Chili.
stricta, *Steud. l. c.*—Chili.
subaristata, *Lam. Illustr.* i. 187.—Reg. Argent.
toluccensis, *H. B. & K. Nov. Gen. et Sp.* i. 148.—Mexic.
tremula, Lam. Fl. Fr. iii. 587 = media.
trichotoma, *Steud. in Flora*, xxix. (1846) 20.—Japon.
triloba, *Nees, Agrost. Bras.* 482.—Reg. Argent.
Uniolae, Nees, ex Steud. Syn. Pl. Gram. 283 = Eragrostis Uniolae.
violascens, Steud. l. c. = subaristata.
virens, Linn. Sp. Pl. ed. II. 103 = minor.
virens, Walt. Fl. Carol. 79 = Poa obtusa.
virens, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 362 = media.
viridis, Pall. ex Steud. Nom. ed. II. i. 226 = minor.

BRIZOPHILE, Salisb. Gen. Pl. Fragm. 34 (1866) = **Ornithogalum**, Tourn. (Liliac.).
nutans, Salisb. l. c. = O. nutans.

BRIZOPYRUM, Link, Hort. Berol. i. 159 (1827) = **Demazeria**, Dum. (Gramin.).
siculum, Link, l. c. = D. sicula.

BRIZOPYRUM, J. Presl, in Rel. Haenk. 280 (1830) = **Distichlis**, Rafin. (Gramin.).
acutiflorum, Nees, Fl. Afr. Austr. 371 = Eragrostis acutiflora.
alternans, Nees, l. c. 369 = Poa alternans.
americanum, Link, Hort. Berol. i. 160 = Distichlis maritima.
boreale, Presl, Rel. Haenk. 28 = Distichlis maritima.
bromoides, Presl, l. c. = Poa bromoides.
calycinum, Presl, l. c. 281 = Poa calycina.
capense, Nees, Fl. Afr. Austr. 371 = Poa Uniolae.
condensatum, Fourn. Mex. Pl. Enum. Gram. 121; ex Hemsl. Biol. Centr. Am. Bot. iii. 578 = Distichlis condensata.
cyperoides, Nees, Fl. Afr. Austr. 374 = Eragrostis cyperoides.
Douglasii, Hook. & Arn. Bot. Beech. Voy. 404 = Poa Douglasii.
obtusiflorum, Fourn. Mex. Pl. Enum. Gram. 120; ex Hemsl. Biol. Centr. Am. Bot. iii. 579 = Distichlis thalassica.
ovatum, Nees, ex Steud. Syn. Pl. Gram. 282.—Peruv.
pilosum, J. Presl, in Rel. Haenk. i. 280 = Distichlis thalassica.
prostratum, Benth. ex Fourn. Mex. Pl. Enum. Gram. 120; ex Hemsl. Biol. Centr. Am. Bot. iii. 578 = Distichlis prostrata.
scirpoideum, Steud. Syn. Pl. Gram. 282 = Schedonorus scirpoideus.
spicatum, Hook. & Arn. Bot. Beech. Voy. 403 = Distichlis maritima.
subspicatum, J. Presl, in Rel. Haenk. 281 = Poa subspicata.

BRIZULA, Hieron. in Bot. Zeit. xxx. (1872) 206 **Aphelia**, R. Br. (Centrolepid.).
Drummondii, Hieron. in Abh. Naturf. Ges. Halle, xii. (1873) 206 = A. Drummondii.
gracilis, Hieron. l. c. = A. gracilis.
Muelleri, Hieron. l. c. 207 = A. Brizula.
Pumilio, Hieron. l. c. = A. Pumilio.

BROCCHIA, Mauri, ex Tenore, Cat. Ort. Napoli. 80 (1845) = **Simmondsia**, Nutt. (Euphorb.).
dichotoma, Mauri, l. c. = S. californica.

BROCCHIA, Vis. Pl. Aeg. et Nub. 35 (1836) = **Cotula**, Linn. (Compos.).
cinerea, Vis. l. c. = C. cinercum.
Kotschyi, Schult. ex Schweinf. & Aschers. Enum. 283 = C. Kotschyi.

BROCCHINIA, Schult. f. Syst. vii. 1250 (1830). *BROMELIACEAE*, Benth. & Hook. f. Gen. iii. 665.
cordylinoides, *Baker, in Journ. Bot.* xx. (1882) 330.—Guiana.
paniculata, *Schult. f. Syst.* vii. 1250.—Bras.
Plumierii, *Griseb. Fl. Brit. W. Ind.* 593.—Ind. occ.
reducta, *Baker, in Journ. Bot.* xx. (1882) 331.—Guiana.

BROCHOSIPHON, Nees, in DC. Prod. xi. 492 (1847) = **Dicliptera**, Juss. (Acanthac.).
australis, Nees, l. c. = D. glabra.

BRODIAEA, Sm. in Trans. Linn. Soc. x. (1811) 2. t. 1.
LILIACEAE, Benth. & Hook. f. iii. 800.
CALLIPRORA, Lindl. Bot. Reg. t. 1590 (1833).
DICHELOSTEMMA, Kunth, Enum. Pl. iv. 469 (1843).
GARDINIA, Bertero, 'in Merc. Chil. i. (1829)'.—Cf. Férussac, Bull. xx. (1830) 111.
HESPEROSCORDUM, Lindl. Bot. Reg. sub t. 1293 (1829).
HOOKERA, Salisb. Parad. Lond. t. 98 (1808), nomen prius.
SCADUAKINTOS, Rafin. Fl. Tellur. iv. 23 (1836).
SEUBERTIA, Kunth, Enum. Pl. iv. 475 (1843).
THEMIS, Salisb. Gen. Pl. Fragm. 85 (1866).
TRITELEIA, Dougl. ex Lindl. Bot. Reg. sub t. 1293 (1829).
VEATCHIA, Kellogg, in Proc. Calif. Acad. ii. (1859?) 11.
alliacea, Miers, ex Baker, in Journ. Linn. Soc. xi. (1871) 375 = Leucocoryne alliacea.
Bridgesii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 237.—Calif.
californica, Lindl. ex Lem. in Fl. des Serres, Sér. I. v. (1849) 428 b = grandiflora.
capitata, *Benth. Pl. Hartw.* 339.—Am. bor. occ.
coccinea, *A. Gray, in Proc. Am. Acad.* vii. (1868) 389.—Calif.
congesta, *Sm. in Trans. Linn. Soc.* x. (1811) 3. t. 1.—Am. bor. occ.
crocea, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 238.—Calif.
Douglasii, *S. Wats. l. c.* 237.—Am. bor. occ.
filifolia, *S. Wats. l. c.* xvii. (1881–82) 381.—Calif.
gracilis, *S. Wats. l. c.* xiv. (1879) 238.—Calif.
grandiflora, Pursh, Fl. Am. Sept. i. 223 = lactea.
grandiflora, *Sm. in Trans. Linn. Soc.* x. (1811) 2.—Am. bor. occ.
grandiflora, Torr. in Stansb. Rep. 397 = Douglasii.
Howellii, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 301.—Am. bor.
ixioides, Sims, Bot. Mag. t. 2382 = Leucocoryne ixioides.
ixioides, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 238.—Am. bor. occ.
lactea, *S. Wats. l. c.*—Am. bor. occ.
laxa, *S. Wats. l. c.* 237.—Calif.
Lemmonae, *S. Wats. l. c.* xx. (1885) 376.—Arizona.
minor, *S. Wats. l. c.* xiv. (1879) 236.—Am. bor. occ.
multiflora, *Benth. Pl. Hartw.* 339.—Am. bor. occ.
parviflora, Torr. & Gray, in Pacif. Rail. Rep. ii. (1854) 125 = multiflora.
peduncularis, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 237.—Calif.
stellaris, *S. Wats. l. c.* xvii. (1881–82) 380.—Calif.
terrestris, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 6.—Calif.
Torreyi, Wood, in Proc. Acad. Sc. Philad. (1868) 172 = terrestris.
volubilis, Baker, in Journ. Linn. Soc. xi. (1871) 377 = Stropholirion californicum.

BROGNIARTIA, Walp. Rep. ii. 994 (1843) = **Brongniartia**, H. B. & K. (Legumin.).

BROMBYA, F. Muell. Fragm. v. 4 (1865). *RUTACEAE*, Benth. & Hook. f. i. 991.
platynema, *F. Muell. l. c.*—Austral.

BROMELIA, [Plum.] ex Linn. Syst. ed. I (1735).
BROMELIACEAE, Benth. & Hook. f. iii. 660.
AGALLOSTACHYS, Beer, Bromel. 35 (1857).
ANANAS, Gaertn. Fruct. i. 30. t. 11 (1788).
PINGUIN, [Dill.] Adans. Fam. ii. 67 (1763).
PSEDOMELIA, Neck. Elem. iii. 150 (1790).
Acanga, Linn. Syst. ed. XII. 232 = B. Pinguin et Karatas Plumieri.
Acanga, Schult. f. Syst. vii. II. 1281 = Araeococcus micranthus.
Acarna, Thunb. ex Schult. f. l. c. 1283, nomen.—Hab.?
acaulis, Stokes, Bot. Mat. Med. ii. 204 = Karatas Plumieri.
agavaefolia, Brongn. in Rev. Hortic. (1875) 247 = Karatas agavaefolia.
agavoides, *Carr. l. c.* (1881) 31.—Hab.?

BROMELIA :—
albo-bracteata, *Steud. in Lechl. Berb. Am. Austr.* 53, nomen.—Chili.
albo-rosea, Lem. Illustr. Hortic. ii. (1855) Misc. 64 = Aechmea suaveolens.
Ananas, Linn. Sp. Pl. 285 = Ananas sativus.
antiacantha, Bertol. in Virid. Bonon. Veg. (1824) 4; Colla, Hort. Ripul. App. ii. 342 = fastuosa.
aquilega, Salisb. Parad. Lond. t. 40 = Aechmea aquilega.
arvensis, Vell. Fl. Flum. 130; iii. t. 114 = Quesnelia rufa.
bicolor, Ruiz & Pav. Fl. Peruv. Chil. iii. 33 = Rhodostachys bicolor.
Binoti, Hort. Jacob-Makoy.—Cf. Gard. Chron. (1880) II. 151 = B. Pinguin.
bracteata, Schult. f. Syst. vii. II. 1280, ex parte = Aechmea laxiflora.
bracteata, Sw. Prod. Veg. Ind. Occ. 56 = Aechmea bracteata.
Caratas, Hill, Veg. Syst. vii. 19 = Karatas Plumieri.
carnea, Beer, Bromel. 31 = Rhodostachys andina.
Carolinae, Beer, l. c. 29; Ortg. ex Regel, Gartenfl. vi. (1857) 361 = Karatas Carolinae.
chrysantha, *Jacq. Hort. Schoenb.* i. 28.—Am. austr.
clandestina, Hort. ex Carr. in Rev. Hortic. (1880) 256 = Greigia sphacelata.
comata, *Beer, Bromel.* 34.—Bras.
Commeliniana, De Vriese, Del. Sem. Hort. Amst. 1844; ex Hoev. & De Vriese, Tijdschr. xii. (1845) 49 = fastuosa.
comosa, Hill, Veg. Syst. vii. 18 = Ananas sativus.
concentrica, Beer, Bromel. 29 = Karatas Laurentii?
crassa, Steud. in Lechl. Berb. Am. Austr. 53 = Greigia Landbecki.
cruenta, R. Grah. in Edinb. N. Phil. Journ. (Oct.–Dec. 1828) 174 = Billbergia cruenta.
denticulata, *C. Koch, Wochenschr.* ii. (1859) 151.—Bras.
discolor, *Lindl. Bot. Reg.* (1838) *Misc.* 48.—Am. austr.
edulis, Salisb. Prod. 247 = Ananas sativus.
elegans, *Hort. Hamb. ex Schult. f. Syst.* vii. II. 1283.—Hab.?
exsudans, Lodd. Bot. Cab. t. 801 = Aechmea exsudans.
fastuosa, *Lindl. Collect.* 1.—N. Granat.
Fernandae, *Morr. in Illustr. Hortic.* xviii. (1871) 114. t. 65.—Bras.
gigantea, *Hort. Vindob. ex Schult. f. Syst.* vii. II. 1283.—Hab.?
glabra, Schult. f. l. c. 1286 = Ananas sativus.
hemispherica, Lam. Encyc. i. 145 = Karatas humilis.
Hookeri, Sweet, Hort. Brit. ed. I. 425 = Aechmea nudicaulis.
humilis, Linn. Syst. ed. XII. 232; Jacq. Enum. Vindob. App. 306 = Karatas humilis.
ignea, Beer, Bromel. 35 = B. Pinguin.
incarnata, Ruiz & Pav. Fl. Per. iii. 32 = Billbergia incarnata.
inermis, Steud. Nom. ed. II. i. 226 = Ananas sativus.
iridifolia, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 16 = Billbergia iridifolia.
Itatiaiae, *Wawra, in Oestr. Bot. Zeitschr.* xxx. (1880) 114.—Bras.
Joinvillei, E. Morr. ex C. H. Morr. in Belg. Hortic. xxvi. (1876) 161. t. 10, 11 = Rhodostachys pitcairniaefolia.
Karatas, Linn. Sp. Pl. 285 = Karatas Plumieri.
laciniosa, *Mart. ex Schult. f. Syst.* vii. II. 1278.—Bras.
laevis, *Hort. Carlsruh. ex Schult. f. l. c.* 1283.—Hab.?
Landbecki, Lechl. ex Phil. in Linnaea, xxxiii. (1864–65) 246 = Greigia Landbecki.
lanuginosa, Beer, Bromel. 32 = Puya lanuginosa.
lasiantha, Willd. ex Schult. f. Syst. vii. II. 1233 = Puya lanata.
latifolia, *Willd. ex Schult f. l. c.* 1283.—Ins. Antigua.
Lindleyana, Lem. Jard. Fleur. iii. (1852–53) t. 223, in obs. = Araeococcus micranthus.
lingulata, Linn. Sp. Pl. 285 = Aechmea lingulata.
longifolia, Lindl. in Paxt. Flow. Gard. ii. t. 65 = Rhodostachys andina.
longifolia, *Rudge, Gui.* i. 31. t. 49.—Guiana.
longifolia, *Klotzsch, in Schomb. Reise*, iii. 903, nomen.—Guiana.
lucida, *Willd. Enum. Hort. Berol.* 345 = Ananas sativus.

BROMELIA :—
lutea, G. F. W. Mey. Prim. Fl. Esseq. 145 = Aechmea nudicaulis.
melanantha, Ker-Gawl. in Bot. Reg. t. 766 = Aechmea bromeliaefolia.
Mertensii, G. F. W. Mey. Prim. Fl. Esseq. 144 = Aechmea Mertensii.
mexicana, *Nois. ex Schult. f. Syst.* vii. II. 1283, nomen.—Mexic.
Moensi, *Linden, Cat.* (1873) n. I. 6.—Bras.
muricata, Arruda, Diss. Pl. Bras. 21 = Ananas bracteatus?
nudicaulis, Linn. Sp. Pl. 286 = Aechmea nudicaulis.
nudicaulis, Reichb. Ic. Exot. 185 = Billbergia pyramidalis.
pallida, Ker-Gawl. in Bot. Reg. t. 344 = Billbergia speciosa.
paniculata, J. F. Gmel. Syst. 529 = Aechmea paniculigera.
paniculigera, Reichb. Ic. Exot. t. 239, 240 = Aechmea exsudans.
paniculigera, Sw. Prod. Veg. Ind. Occ. 56 = Aechmea paniculigera.
pauciflora, *C. Koch, Wochenschr.* ix. (1866) 183.—Hab.?
Pinguin, *Linn. Sp. Pl.* 285.—Am. trop.
pitcairnifolia, C. Koch, Wochenschr. xi. (1868) 325 = Rhodostachys pitcairniaefolia.
pumila, *Hort. ex Schult. f. Syst.* vii. II. 1283.—Hab.?
pyramidalis, Sims, Bot. Mag. t. 1732 = Billbergia pyramidalis.
pyramidata, Beer, Bromel. 34 = Puya pyramidata.
Rohaniana, Walp. Ann. vi. 71 = Billbergia vittata.
rubra, Hort. ex Schult. f. Syst. vii. II. 1285 = Ananas sativus.
Sagenaria, Arruda, Diss. Pl. Bras. 14 = Ananas bracteatus.
scarlatina, Hort. ex Hérincq, Hortic. Franç. (1869) 246 = Distiacanthus scarlatinus.
scarlatina, *E. Morr. ex C. H. Morr. in Belg. Hortic.* xxxi. (1881) 164.—Bras.
Sceptrum, Hueg. Parad. Vind. fasc. viii. (sic) = fastuosa.
semiserrata, Willd. Enum. Hort. Berol. 345 = Ananas sativus.
sepiaria, *Hort. ex Schult. f. Syst.* vii. II. 1283.—Hab.?
Serra, *Griseb. in Goett. Abh.* xxiv. (1879) 328.—Reg. Argent.
sessiliflora, *Lodd. ex Loud. Hort. Brit.* 118.—Am. austr.
silvestris, Vell. Fl. Flum. 129; iii. t. 113 = Ananas sativus.
sphacelata, Ruiz & Pav. Fl. Per. iii. 32. t. 474 = Greigia sphacelata.
spicata, *Lam. Encyc.* i. 146.—Ins. Martinic.
subspinosa, Wendl. ex Schult. f. Syst. vii. II. 1286 = Ananas sativus.
surinamensis, *Miq. in Linnaea*, xviii. (1844) 378.—Guiana.
sylvestris, *Willd. ex Sims, Bot. Mag.* t. 2392.—Am. trop.
thyrsifolia, Willd. ex Schult. f. Syst. vii. II. 1282 = Aechmea spicata.
tinctoria, Mart. Reise Bras. ii. 554 = Aechmea bromeliaefolia.
tristis, *Beer, Bromel.* 30.—Bras.
variegata, Arruda, Diss. Pl. Bras. 7 = Billbergia speciosa.
violacea, Hort. ex Schult. f. Syst. vii. II. 1285 = Ananas sativus.
viridis, Hort. ex Schult. f. l. c. = Ananas sativus.
zebrina, Herb. in Bot. Mag. t. 2686 = Billbergia zebrina.

BROMFELDIA, Neck. Elem. ii. 347 (1790) = **Jatropha**?, Linn. (Euphorb.).

BROMHEADIA, Lindl. Bot. Reg. (1841) App. 90.
ORCHIDEAE, Benth. & Hook. f. iii. 540.
aporoides, *Reichb. f. Otia Bot. Hamb.* 44.—Burma.
Finlaysoniana, Reichb. f. in Walp. Ann. vi. 630, 882 = palustris.
palustris, *Lindl. Bot. Reg.* (1841) *Misc.* 89.—Malaya.

BROMIDIUM, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i.; (Pl. Meyen.) 154 (1843) = **Deyeuxia**, Clar. (Gramin.).

BROMIDIUM :—
anomalum, Doell, in Mart. Fl. Bras. ii. III. 103.—Bras.
brevifolium, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. 155 = D. brevifolia.
caespitosum, Nees & Meyen, l. c. = Agrostis Preslii.
hygrometricum, Nees & Meyen, l. c. 154.—Bras.
lobatum, Nees, in Hook. Lond. Journ. Bot. ii. (1843) 415 = D. quadriseta.
quadriseta, Nees, l. c. 416 = D. quadriseta.
rigescens, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. 155 = Agrostis rigescens.
spectabile, Nees & Meyen, l. c. 156.—Peruv.

BROMOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 187 = **Bromus**, Dill. (Gramin.).
aspera, Fourr. l. c. = Bromus asper.
erecta, Fourr. l. c. = Bromus erectus.

BROMUS, Dill. ex Linn. Syst. ed. I (1735). *GRAMINEAE*, Benth. & Hook. f. iii. 1200.
ANISANTHA, C. Koch, in Linnaea, xxi. (1848) 394.
BROMOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 187.
CERATOCHLOA, Beauv. Agrost. 75. t. 15. f. 7 (1812).
LASIOPOA, Ehrh. Beitr. iv. 147 (1789).
LIBERTIA, Lej. in Nov. Act. Nat. Cur. xii. (1825) 755. t. 65.
MICHELARIA, Dum. Obs. Gram. Belg. 77. t. 16 (1823).
SCHEDONORUS, Beauv. Agrost. 99. t. 19. f. 2 (1812), partim.
SERRAFALCUS, Parl. Pl. Nov. 75 (1842).
TRINIUSA, Steud. Syn. Pl. Gram. 328 (1854).
ZERNA, Panz. in Denkschr. Acad. Mucnch. 1813 (1814) 296, partim.
abortiflorus, St. Amans, Fl. Agen. 44 = tectorum.
adoënsis, *Hochst. ex Steud. Syn. Pl. Gram.* 326.—Abyss.
aegyptiacus, Tausch, in Flora, xx. (1837) 124 = brachystachys.
affinis, Dum. Obs. Gram. Belg. 118 = mollis ?
agrarius, Dum. l. c. = diffusus.
agrarius, Hornung, ex Steud. Nom. ed. II. i. 226 = racemosus.
agrestis, All. Fl. Pedem. ii. 249 = erectus.
albidus, *Bieb. Fl. Taur. Cauc.* iii. 79.—Reg. Cancas.
aleutensis, *Trin. ex Ledeb. Fl. Ross.* iv. 361.—Unalaska.
alopecuroides, Lag. Gen. et Sp. Nov. 4 = Koeleria phleoides.
alopecuroides, Moris, ex Steud. Nom. ed. II. i. 226, 228 = intermedius.
alopecuroides, Poir. Encyc. Suppl. i. 703 = seq.
Alopecuros, *Poir. Voy. Barb.* ii. 100; *Vahl, Symb. Bot.* ii. 22.—Reg. Mediterr.
Alopecurus, Pers. Syn. i. 95 = praec.
alpestris, *Steud. in Lechl. Berb. Am. Austr.* 52, nomen.—Chili.
altissimus, Gilib. Exercit. ii. 537 = arvensis.
altissimus, *Pursh, Fl. Am. Sept.* ii. 728.—Am. bor.
altissimus, [Weber, in] Wigg. Prim. Fl. Hols. 9 = asper.
ambigens, Jord. ex Nym. Consp. 821 = rigidus.
ambiguus, Cyrill. Pl. Rar. fasc. i. 10 = Festuca bromoides.
amplus, C. Koch, in Linnaea, xxi. (1848) 418 = sterilis.
anatolicus, Boiss. & Heldr. Diagn. Scr. I. xiii. 63 = patulus.
angustifolius, Schrank, Baier. Fl. i. 366 = erectus.
angustifolius, *Hornem. Hort. Hafn. Suppl.* 13.—Hab. ?
angustissimus, C. Koch, in Linnaea, xxi. (1848) 421 = Biebersteinii.
anomalus, *Rupr. in Bull. Acad. Brux.* ix. II. (1842) 236.—Mexic.
antarcticus, Hook. f. Fl. Antarct. 97 = Danthonia bromoides.
aquaticus, Schrank, Baier. Fl. i. 367 = Festuca gigantea.
arduennensis, *Dum. Obs. Gram. Belg.* 77.—Belgium.
arenaceus, Pourr. in Mém. Acad. Toul. iii. (1788) 308 = tectorum.
arenarius, *Labill. Nov. Holl. Pl.* i. 23. t. 28.—Austral.; N. Zel.
arenarius, Thomine Desm. in Mém. Soc. Linn. Calv. (1824) 40 = mollis.
aristatus, *Steud. Syn. Pl. Gram.* 320.—Persia.
armenus, *Boiss. Fl. Orient.* v. 642.—Armenia.
arrhenatheroides, *Baker, in Journ. Linn. Soc.* xx. (1884) 301.—Madag.
arundinaceus, Roth, Tent. Fl. Germ. ii. 141 = Festuca elatior.
arvensis, Knapp, Gram. Brit. t. 82 = racemosus.
arvensis, Lam. Fl. Fr. iii. 607 = erectus.
arvensis, *Linn. Sp. Pl.* 77.—Europ.; Oriens; As. bor.
arvensis, Oed. in Fl. Dan. t. 293 = secalinus.
asper, *Murr. Prod. Gotting.* 42.—Europ.; Oriens; As. bor.
asper, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. i. 73 = erectus.
asperipes, Jord. ex Nym. Consp. 821 = rigidus.
asperrimus, *Tenore, Fl. Nap.* t. 105.—Italia.
atticus, Steud. ex Nym. Consp. 823 = mollis.
auleticus, *Trin. ex Nees, in Mart. Fl. Bras.* ii. 468.—Reg. Argent.
auriculatus, Rasp. in Ann. Sc. Nat. Sér. I. v. (1825) 439 = arduennensis.
australis, R. Br. Prod. i. 178 = arenarius.
avenaceus, Lam. Illustr. i. 194 = tectorum.
avenaeformis, Steud. Syn. Pl. Gram. 326 = Trisetum flavescens.
avenoides, *Baker, in Journ. Linn. Soc.* xx. (1884) 302.—Madag.
badensis, C. C. Gmel. Fl. Bad. iv. 75. t. 5 = secalinus.
barbatus, Savi, Bot. Etrusc. ii. 61 = Festuca Alopecuros.
Baumgartenii, Steud. Syn. Pl. Gram. 320 = rigidus.
Berterianus, *Colla, in Mem. Acc. Torin.* xxxix. (1835) 25.—Chili.
bicuspis, Nees, ex Steud. Syn. Pl. Gram. 322.—Chili.
Biebersteinii, Roem. & Schult. Syst. ii. 638 = albidus.
bifidus, Thunb. Fl. Jap. 53 = Trisetum cernuum.
Billotii, Sch. Bip. in Flora, xxxii. (1849) 233 = secalinus.
bohemicus, F. W. Schmidt, ex Mert. & Koch, in Roehl. Deutschl. Fl. ed. III. i. 638 = Festuca Myurus.
Boraei, Jord. ex Nym. Consp. 821 = rigidus.
brachyantheca, *Doell, in Mart. Fl. Bras.* ii. III. 110.—Bras.
brachystachys, *Hornung, in Flora*, xvi. (1833) II. 417.—Oriens; Aegypt.
breviaristatus, *Buckl. in Proc. Acad. Sc. Philad.* '1862' (1863) 98.—Am. bor. occ.
brevis, *Steud. Syn. Pl. Gram.* 326.—Mendoza.
brevisetus, *Dum. Fl. Belg.* 155.—Belgium.
brizaeformis, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 30.—Reg. Caucas.; Persia.
brizaeformis, Willk. ex Willk. & Lange, Prod. Fl. Hisp. i. 101 = squarrosus.
brizioides, Lam. Illustr. i. 193. n. 1060 = Briza Lamarckiana.
brizoides, Willd. ex Steud. Nom. ed. II. i. 227 = Briza rotundata.
Brongniartii, *Kunth, Enum. Pl.* i. 421.—Bras.
Buxbaumii, Tenore, ex Steud. Nom. ed. II. i. 227 = Brachypodium distachynm.
caerulescens, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 9.—Hab. ?
caespitosus, Host, Gram. Austr. iv. t. 18 = Brachypodium pinnatum.
caespitosus, Steud. Nom. ed. I. 119 = Brachypodium ramosum.
Caldasii, H. B. & K. Nov. Gen. et Sp. i. 151 = Festuca Caldasii.
canadensis, Michx. Fl. Bor. Am. i. 65 = ciliatus.
canariensis, Zuccagni, Cent. n. 27; in Roem. Coll. 125 = lanceolatus.
canescens, Viv. Fl. Lyb. Spec. 5 = rubens.
canus, Willd. ex Steud. Nom. ed. II. i. 227 = Festuca mollis.
capensis, Steud. in Flora, xii. (1829) 491 = vestitus.
capillaris, Moench, Meth. 194 = Leptochloa domingensis.
capillaris, *Vahl, ex Hornem. Hort. Hafn.* i. 101.—Hab. ?
capillifolia, *Dufour, ex Roem. & Schult. Syst.* ii. 735.—Hispan.
cappadocicus, *Boiss. & Bal. in Bull. Soc. Bot. Fr.* iv. (1857) 306.—As. Min.
caprearum, Hack. in Oestr. Bot. Zeitschr. xxix. (1879) 209 = caprinus.
caprinus, *Kerner, ex Hack. l. c.*—Italia.
carinatus, *Hook. & Arn. Bot. Beech. Voy. Suppl.* 403.—Am. bor. occ.
catharticus, Vahl, Symb. Bot. ii. 22 = unioloides.
caucasicus, *Fisch. ex Steud. Nom.* ed. II. i. 227.—Reg. Caucas.
caudatus, *Brot. Phyt. Lusit.* t. 99.—Lusitan.
Cavanillesii, *Willk. ex Willk. & Lange, Prod. Fl. Hisp.* i. 101.—Hispan.
Cebadilla, *Steud. Syn. Pl. Gram.* 321.—Chili.
Chiapporianus, De Not. ex Nym. Consp. 824 = patulus.
chilensis, *Trin. in Linnaea*, x. (1836) 304.—Chili.
chloroticus, Bieb. ex Kunth, Enum. Pl. i. 420 = albidus.
Chrysopogon, Viv. Fl. Lyb. Spec. 4. t. 2 = scoparius.
ciliatus, Huds. Fl. Angl. ed. I. 40 = madritensis.
ciliatus, Lam. Fl. Fr. iii. 609 = Brachypodium distachyum.
ciliatus, *Linn. Sp. Pl.* 76.—Am. bor.
ciliatus, Muhl. Desc. Gram. 169 = Kalmii.
cinereus, S. G. Gmel. ex Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 66 = Trisetum Gmelini.
cincinnatus, Tenore, Lez. Bot. iv. I. 52; Fl. Nap. iii. 85 = Avena bromoides.
circinnatus, Kunth, Enum. Pl. i. 423 = praec.
coarctatus, Hornem. ex Kunth, l. c. 303 = Gaudinia geminiflora.
cognatus, *Steud. Syn. Pl. Gram.* 321.—Abyssin.
collinus, *Phil. in Linnaea*, xxix. (1857–58) 101.—Chili.
coloratus, Baumg. ex Nym. Consp. 824 = squarrosus.
coloratus, *Steud. Syn. Pl. Gram.* 429.—Reg. Magellan.
commutatus, Bieb. Fl. Taur. Cauc. iii. 77 = patulus.
commutatus, Guss. ex Steud. Syn. Pl. Gram. 325 = scabratus.
commutatus, Schrad. Fl. Germ. i. 353 = racemosus.
compactus, Hort. ex Steud. Nom. ed. II. i. 227 = mollis.
compressus, Lag. Elench. 4 = ciliatus.
condensatus, *Hack. in Oestr. Bot. Zeitschr.* xxix. (1879) 209.—Europ.
confertus, Bieb. Fl. Taur. Cauc. i. 71 = scoparius.
confertus, Bor. Fl. Centr. Fr. ed. II. 586 = molliformis.
confinis, *Nees, ex Steud. Syn. Pl. Gram.* 320.—Ind. or.
conformis, *Steud. l. c.* 323.—Japon.
contortus, Desf. Fl. Atlant. i. 95 = B. Alopecuros.
corniculatus, Lam. Fl. Fr. iii. 608 = Brachypodium pinnatum.
crinitus, *Boiss. & Hohen. Diagn.* Ser. I. xiii. 64.—Pers.; Afghan.; Turkest.
cristatus, Linn. Sp. Pl. 78 = Agropyron cristatum.
cristatus, Spreng. Nachr. i. Bot. Gart. Halle, 15 = Koeleria phleoides.
cultus, *Steud. Syn. Pl. Gram.* 323.—Europ.
cylindraceus, *Brot. Phyt. Lusit.* 98.—Lusitan.
Cyri, Trin. in C. A. Mey. Verz. Pfl. Cauc. 24 = patulus.
dactyloides, Roth, Nov. Sp. 60 = Koeleria phleoides.
Danthoniae, Trin. in C. A. Mey. Verz. Pfl. Cauc. 24 = macrostachys.
decipiens, Hort. Madr. ex Steud. Nom. ed. II. i. 227 = madritensis.
decolorans, Reichb. Fl. Germ. Excurs. 44 = Festuca elatior.
decumbens, Koel. Gram. 242 = decumbens.
denseciliatus, *Steud. Syn. Pl. Gram.* 422.—'Terra nova.'
depauperatus, *J. Presl, in Rel. Haenk.* i. 263.—Am. bor. occ.
dertonensis, All. Fl. Pedem. ii. 249 = Festuca bromoides.
diandrus, Curt. Fl. Lond. fasc. vi. t. 5 = madritensis.
diandrus, Roth, Bot. Abh. 44 = pilosus.
diffusus, *Dum. Obs. Gram. Belg.* 118.—Belgium.
digitatus, Poir. ex Beauv. Agrost. 156 = madritensis.
dilatatus, Lam. Encyc. i. 468 = madritensis.
dissitiflorus, *Baker, in Journ. Linn. Soc.* xx. (1884) 301.—Madag.
distachyos, Linn. Cent. ii. 8; Amoen. Acad. iv. 304 = Brachypodium distachyum.
distichus, Georgi, Bemerk. Reise Russ. Reich. i. 197 = Agropyron cristatum.
distichus, Moench, Meth. 192 = sterilis.
divaricatus, Rhode, ex Loisel. in Desv. Journ. Bot. II. (1809) 214 = macrostachys.

BROMUS:—
dortonensis, Kunth, Enum. Pl. i. 397, in syn. = Festuca bromoides.
dumetorum, Lam. Fl. Fr. iii. 605 = asper.
dumosus, Vill. Hist. Pl. Dauph. ii. 119 = Brachypodium sylvaticum.
Ehrharti, Gaud. in Roem. Collect. 19 = secalinus.
elatior, Koel. Gram. 214 = Festuca elatior.
elatior, Spreng. Syst. i. 359 = Festuca arundinacea.
elatus, Gueldenst. ex Ledeb. Fl. Ross. iv. 353 = Festuca elatior.
elongatus, Gaud. Agrost. i. 305 = racemosus.
erectus, *Huds. Fl. Angl.* ed. I. 39. — Europ.; As. Min.
erectus, Moris, Stirp. Sard. Elench. i. 50 = scoparius.
erectus, Ledeb. Fl. Ross. iv. 358 = inermis.
exaltatus, *Bernh. in Linnaea*, xv. (1841) *Litt.* 90.—Mexic.
fascicularis, Tenore, Fl. Nap. iv. 17, 161 = fasciculatus.
fasciculatus, *Presl, Cyp. et Gram. Sic.* 39.—Reg. Mediterr. or.
Ferronii, Mabelli, in Act. Soc. Linn. Bord. xxv. (1864) 613.—Cf. Bull. Soc. Bot. Fr. xiv. (1867) Bibl. 138 = mollis.
festucoides, *Link, in Schrad. Journ.* ii. (1799) 315.—Lusitan.
fibrosus, *Hack. in Oestr. Bot. Zeitschr.* xxix. (1879) 209.—Europ. or.
flabellatus, *Hack. ex Boiss. Fl. Orient.* v. 648.—Palaestin.
flavescens, *Tausch, in Flora*, xx. (1837) i. 124.—Sicilia.
Fonki, *Phil. in Linnaea*, xxix. (1857-58) 102.—Chili.
fragilis, Schrank, Baier. Fl. i. 370 = Brachypodium pinnatum.
fragilis, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 90.—Europ.
frigidus, *Ball, in Journ. Linn. Soc.* xxii. (1885) 63.—Peruv.
frigidus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 645.—Persia.
Gaudini, Roem. & Schult. Syst. ii. 632 = racemosus.
geniculatus, Linn. Mant. i. 33 = Festuca genieulata.
geniculatus, *Steud. Syn. Pl. Gram.* 318.—Oriens.
giganteus, Leers, Fl. Herborn. 38 = Festuca elatior.
giganteus, *Linn. Sp. Pl.* 77.—Europ.; Sibir.
giganteus, Vill. Hist. Pl. Dauph. ii. 118 = asper.
Gilliesii, *Nees, ex Steud. Syn. Pl. Gram.* 328.—Chili.
glaber, Reich. Fl. Franco-Moen. i. 19 = Festuca gigantea.
glaber, Scop. Fl. Carn. ed. II. i. 84 = Agropyron repens.
glaber, Willd. ex Steud. Nom. ed. II. i. 227 = pitensis.
glaberrimus, C. Koch, in Linnaea, xxi. (1848) 420 = erectus.
glaucus, Hort. ex Reichb. Fl. Germ. Excurs. 38 = Festuca ovina.
glaucus, Lapeyr. Hist. Abr. Pl. Pyr. t. 53 = erectus.
glaucus, Spreng. in Link, Hort. Berol. i. 170 = Festuca rubra.
glomeratus, Scop. Fl. Carn. ed. II. i. 76 = Dactylis glomerata.
glomeratus, Tausch, in Flora, xx. (1837) 124 = mollis.
gracilis, Leyss. Fl. Hal. 26. n. 116 = Brachypodium pinnatum.
gracilis, Weigel, Obs. 15 = Brachypodium sylvaticum.
gracillimus, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 527 = crinitus.
grandiflorus, *C. Koch, in Linnaea*, xxi. (1848) 419 (sp. dub.).—Pontus.
grandiflorus, Weigel, Obs. 9 = sterilis, tectorum.
grossus, DC. Fl. Fr. iii. 68 = secalinus.
grossus, Desf. in C. C. Gmel. Fl. Bad. iv. t. 2, 3 = velutinus.
Gussoni, Parl. Pl. Nov. 66 = rigidus.
gyganteus, Crantz, Inst. i. 390 = giganteus.
gynandrus, Roth, in Roem. & Ust. Mag. iv. 10tes St. (1790) 20 = madritensis.
Haenkeanus, Kunth, Enum. Pl. i. 416 = unioloides.
Hausknechtii, *Boiss. Fl. Orient.* v. 648.—Mesopotam.
hirsutissimus, Cyrill. ex Roem. & Schult. Syst. ii. 643 = asper.
hirsutus, Curt. Fl. Lond. fasc. ii. t. 8 = asper.
hirsutus, Schrank, in Denkschr. Bot. Ges. Regensb. (1818) 161 = squarrosus.
hirtus, Lichtst. in Roem. & Schult. Syst. ii. 654.—Afr. austr.

BROMUS:—
hispidus, Savi, Bot. Etrusc. ii. 62 = Koeleria hispida.
Hohenackeri, Hochst. ex Ledeb. Fl. Ross. iv. 416 = Ventenata macra.
holciformis, *Steud. & Hochst. ex Steud. Syn. Pl. Gram.* 324.—Chili.
Hookeri, Fourn. Mex. Pl. En. Gram. 127; ex Hemsl. Biol. Centr. Am. Bot. iii. 583 = ciliatus.
Hookerianus, Thurber, in Bot. U. St. Expl. Exped. ii. 493 = virens.
hordeaceus, C. C. Gmel. Fl. Bad. iv. 68 = secalinus.
hordeaceus, Linn. Sp. Pl. 77 = racemosus, mollis.
hordeiformis, Lam. Illustr. i. 195 = Festuca bromoides.
Hughi, Tod. ex Nym. Consp. 823 = intermedius.
humilis, Cav. Ic. vi. 65 = scoparius.
hyalinus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 90 = arvensis.
hybridus, *Schur, Enum. Pl. Transs.* 801.—Transsylv.
imperialis, Hort. ex Steud. Nom. ed. II. i. 227 = ciliatus.
incrassatus, Lam. Encyc. i. 468 = rigidus, Festuca geniculata.
indicus, Hort. ex Steud. Nom. ed. II. i. 227 = rigidus.
inermis, *Leyss. Fl. Hal.* 16; *Linn. Mant.* ii. 186.—Europ.; Reg. Caucas.; As. bor.
inermis, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 252 = albidus.
insignis, *Buese, in Miq. Pl. Jungh.* 347.—Java.
intermedius, *Guss. Fl. Sic. Prod.* i. 114.—Reg. Mediterr.
intermedius, *Zumag. Fl. Pedem.* i. 166.—Pedemont.
japonicus, *Thunb. Fl. Jap.* 51. t. 11.—Japon.
jubatus, Tenore, Fl. Nap. 10 = sterilis.
jubatus, *Vill. Hist. Pl. Dauph.* ii. 118.—Gallia.
Kalmii, *A. Gray, Man.* 600.—Am. bor.
Kochii, C. C. Gmel. Fl. Bad. iv. 76. t. 6 = patulus.
Krausei, *Regel, in Act. Hort. Petrop.* vii. (1880) 600.—As. centr.
lanatus, H. B. & K. Nov. Gen. et Sp. i. 150 = Festuca mollis.
lanceolatus, Roth, Catalect. i. 18 = macrostachys.
lanuginosus, Poir. Encyc. Suppl. i. 703 = macrostachys.
lateripronus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = tectorum.
latifolius, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 865.—Sibir.
laxiflorus, *Spreng. in Zeyh. ex Steud. Nom.* ed. II. i. 228.—Afr. austr.
laxus, Hornem. Hort. Hafn. ii. 954 = erectus.
laxus, Reichb. Ic. Fl. Germ. i. 32. t. 77. f. 1605 = inermis.
lenis, *J. Presl, in Rel. Haenk.* i. 262.—Peruv.
leptoclados, *Nees, Fl. Afr. Austr. Gram.* i. 453.—Afr. austr.
leptostachys, Ehrenb. ex Boiss. Fl. Orient. v. 654 = brachystachys.
leptostachys, [Pieri,] in Ionios Anthol. iii. (1834) 680 = squarrosus.
leptostachyus, *Steud. Syn. Pl. Gram.* 324.—Gallia.
ligusticus, All. Fl. Pedem. ii. 249 = rubens, scoparius, Festuca geniculata.
lithobius, *Trin. in Linnaea*, x. (1836) 303.—Chili.
littoralis, *Steud. Syn. Pl. Gram.* 325.—'Austr. littoralis.'
littoreus, Georgi, Beschr. Russ. Reich. Nachtr. 257 = inermis.
littoreus, Retz. Fl. Scand. Prod. ed. I. 19 = Festuca elatior.
lividus, H. B. & K. Nov. Gen. et Sp. i. 150 = Festuca livida.
loliaceus, Tenore, Fl. Nap. iii. 91 = Triticum Barrelieri.
longe-aristatus, Gilib. Exercit. ii. 536 = sterilis.
longiflorus, Spreng. Pugill. ii. 23 = tectorum.
longifolius, *Schousb. Vextr. Marokko*, 44.—Marocco.
longipilus, Kummer & Send. in Flora, xxxii. (1849) 757 = tectorum.
luzonensis, J. Presl, in Rel. Haenk. i. 262 = Triticum luzonense.
macranthes, Meyen, Reise, i. 311, nomen = setifolius.
macranthos, E. Desv. in C. Gay, Fl. Chil. vi. 443 = praec.
macrocladus, Boiss. Diagn. Ser. I. xiii. 64 = arvensis.
macrostachys, *Desf. Fl. Atlant.* i. 96. t. 19.—Reg. Mediterr.
macrostachyus, Guss. Fl. Sic. Prod. 115 = lanceolatus.
madritensis, Cav. ex Kunth, Enum. Pl. i. 419 = maximus.
madritensis, *Linn. Cent. Pl.* i. 5; *Amoen. Acad.* iv. 265.—Europ.; Afr. bor.; Oriens.
malacanthus, Trin. ex Ledeb. Fl. Ross. iv. 361 = ciliatus.

BROMUS:—
Mango, *E. Desv. in C. Gay, Fl. Chil.* vi. 440.—Chili.
marginatus, Steud. Syn. Pl. Gram. 322 = ciliatus.
massiliensis, *Forsk. Fl. Aegypt. Arab.* 203.—Massilia.
matritensis, 'Linn.' ex Roem. & Schult. Syst. ii. 651 = madritensis.
matritensis, Cav. Ic. vi. 67 = rigidus.
Matthewsii, *Steud. Syn. Pl. Gram.* 323.—Peruv.
maximus, Desf. Fl. Atlant. i. 95. t. 26 = rigidus.
maximus, Gilib. Exercit. ii. 535 = secalinus.
maximus, Roth, N. Beitr. i. 114 = pilosus.
megapotamicus, *Spreng. Syst.* iv. *Cur. Post.* 36.—Bras.
mexicanus, Hort. ex Steud. Nom. ed. II. i. 228 = Triticum mexicanum.
Micheli, Savi, Bot. Etrusc. i. 78 = festucoides, Avellinia Micheli.
modensis, *Steud. Syn. Pl. Gram.* 324.—Graecia.
molliformis, *Lloyd, Fl. Loire-Inf.* 315.—Gallia.
mollis, *Linn. Sp. Pl.* ed. II. 112.—Europ.; Afr. et As. bor.
mollis, Thunb. Prod. Pl. Cap. 22 = vestitus.
mollissimus, *Hornem. Hort. Hafn. Suppl.* 13.—Hab. ?
monostachyus, Steud. Nom. ed. II. i. 228 = Brachypodium distachyum.
montanus, Gaertn. Mey. & Scherb. Fl. Wett. ex Kunth, Enum. Pl. i. 418 = erectus.
montanus, Scop. Fl. Carn. ed. II. i. 82 = asper.
montanus, *Retz. Prod. Scand.* 124.—Europ. bor.
mucronatus, Willd. ex Steud. Nom. ed. II. i. 228 = unioloides.
multiflorus, Host, Gram. Austr. 1. t. 11 = patulus.
multiflorus, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 865 = macrostachys.
multiflorus, Roth, Tent. Fl. Germ. i. 47 = racemosus.
multiflorus, Sm. in Trans. Linn. Soc. iv. (1798) 283 = secalinus.
multiflorus, Weigel, Obs. 2. t. 1. f. 1 = arvensis.
multiflorus, Willd. Sp. Pl. i. 428 = commutatus.
multispicatus, Roem. & Schult. Syst. ii. 650 = madritensis.
Munroi, *Boiss. Fl. Orient.* v. 643.—Afghanist.
muralis, Huds. Fl. Angl. ed. II. i. 50 = madritensis.
murorum, Bernh. ex Roem. & Schult. Syst. ii. 651 = pilosus, tectorum.
mutabilis, *F. Schultz, in Flora*, xxxii. (1849) 234.—Europ.
nanus, Weigel, Obs. 8 = mollis.
neglectus, Nym. Syll. 419 = racemosus.
nemoralis, Huds. Fl. Angl. ed. II. i. 51 = asper.
nemorosus, Vill. Hist. Pl. Dauph. ii. 117 = asper.
nitidus, Clarke, Trav. ii. 260 = rigidus.
nitidus, Dum. Obs. Gram. Belg. 119 = secalinus.
Nočanus, Boiss. Diagn. Ser. II. iv. 141 = squarrosus.
nototropus, *Rupr. Sert. Tiansch.* 73.—As. centr.
odoratus, Gouan, Herbor. 11 = erectus.
oranensis, *Steud. Syn. Pl. Gram.* 319.—Afr. bor.
Orcuttianus, *Vasey, in Coult. Bot. Gaz.* x. (1885) 223.—Am. bor.
ovatus, Cav. Ic. vi. 67 = Trisetum ovatum.
ovatus, J. Gaertn. in Nov. Comm. Acad. Petrop. xiv. (1770) 537. t. 19. f. 1 = scoparius.
ovinus, Scop. Fl. Carn. ed. II. i. 77 = Festuca ovina.
oxyodon, Schrenk, Enum. Pl. Nov. ii. 1 = macrostachys.
oxyphloeus, Payne, Palest. Expl. Rep. (1874) 128 = macrostachys.
pallens, *Cav. Ic.* vi. 66. t. 591. f. 1.—Ins. Philipp.
pannonicus, *Kummer & Sendt. in Flora*, xxxii. (1849) 757.—Hungar.
paradoxus, Presl, Fl. Sicul. i. p. xlv, in nota = pentastachyus.
parviflorus, *Desf. Cat. Hort. Par.* ed. III. 387.—Hab. ?
patagonicus, *Phil. in Anal. Univ. Chil.* (1873) 577.—Chili.
patentissimus, Weigel, Obs. 12 = Festuca sylvatica.
patulus, *Mert. & Koch, in Roehl. Deutschl. Fl.* i. 685.—Europ.; Oriens; Afghanist.
pauciflorus, *Schum. in Fl. Dan.* t. 1383.—Island.
pauciflorus, Willd. ex Steud. Nom. ed. II. i. 228 = Festuca mollis.
pauper, J. F. Gmel. Syst. 191 = Brachypodium distachyum.
pauper, Schrank, Baier. Fl. i. 370 = Brachypodium sylvaticum.
pectinatus, *Thunb. Prod. Pl. Cap.* i. 22.—Afr. austr.

BROMUS:—
pendulinus, Desf. ex Steud. Nom. ed. I. 119, 120 = commutatus.
pendulinus, Hort. ex Schrad. N. Journ. iv. (1810) 70 = ciliatus.
pendulus, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 90 = patulus.
pentastachyus, Tinco, Pl. Rar. Sic. Pugill. i. 4 = Brachypodium distachyum.
perennis, Vill. Hist. Pl. Dauph. ii. 122 = erectus.
persicus, Boiss. Diagn. Ser. I. xiii. 66; Ledeb. Fl. Ross. iv. 357 = tomentosus.
Petitianus, *A. Rich. Tent. Fl. Abyss.* ii. 438.—Abyss.
phleoides, J. F. Gmel. Syst. 190 = Koeleria phleoides.
phoenicoides, Steud. Nom. ed. I. 120 = Brachypodium pinnatum.
Phoenix, *J. F. Gmel. Syst.* 190.—Hab. ?
phrygius, Boiss. Diagn. Ser. II. iv. 140 = patulus.
pictus, *Hook. f. Fl. Antarct.* 387.—Chili.
pilosus, *F. G. Dietr. Vollst. Lex. Gaertn.* i. 558.—Europ. austr.
pilosus, Willd. in Spreng. Syst. i. 358 = trichodes.
pinnatus, Linn. Sp. Pl. 78 = Brachypodium pinnatum.
pinnatus, Oed. in Fl. Dan. t. 164 = Brachypodium sylvaticum.
pitensis, *H. B. & K. Nov. Gen. et Sp.* i. 152.—Ecuador.
platystachys, Desf. Cat. Hort. Par. ed. III. 387 = arduennensis.
platystachyus, Lam. Illustr. i. 195 = Brachypodium distachyum.
Pluckenetii, All. Fl. Pedem. ii. 250 = Brachypodium ramosum.
poaeformis, *Forsk. Fl. Aegypt. Arab.* 23.—Aegypt.
poaeformis, Spreng. Nachr. i. Bot. Gart. Halle, 15; Fl. Hal. Mant. 34 = Diplachne fascicularis.
poaeformis, *Vahl, ex Hornem. Hort. Hafn.* i. 102.—Hab. ?
polymorphus, Hohen. Enum. Taluesch. 19 = patulus.
polymorphus, Hort. ex Steud. Nom. ed. II. i. 228 = arvensis.
polymorphus, Scop. Fl. Carn. ed. II. i. 79; Huds. Fl. Angl. ed. II. i. 48 = mollis, racemosus, secalinus.
polystachios, Forsk. Fl. Aegypt. Arab. 23 = Diplachne fusca.
polystachyus, DC. Fl. Fr. vi. 276 = madritensis.
polystachyus, Kunth, Enum. Pl. i. 447 = Brachypodium distachyum, Triticum ciliatum.
polystachyus, Steud. Nom. ed. II. i. 228 = arduennensis.
pratensis, Ehrh. Beitr. vi. 84, nomen = racemosus.
pratensis, Lam. Encyc. i. 468 = erectus.
pratensis, Spreng. Syst. i. 359 = Festuca elatior.
Preslii, *Kunth, Enum. Pl.* i. 416.—Peruv.
procerus, H. B. & K. Nov. Gen. et Sp. i. 150 = Festuca eminens.
procerus, Spreng. Syst. i. 357 = quitensis.
propendens, Jord. ex Nym. Consp. 821 = rigidus.
pseudoarvensis, Koel. Gram. 241 = erectus.
pseudoasper, Schur, Enum. Pl. Transs. 804 = asper.
pseudoinermis, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 90.—Transsylv.
pubescens, C. Koch, in Linnaea, xxi. (1848) 420 = variegatus.
pubescens, Muhl. in Willd. Enum. Hort. Berol. 120 = ciliatus.
pubescens, [Picri,] in Ionios Anthol. iii. (1834) 680 = squarrosus.
pulchellus, *Fig. & De Not. Agr. Fragm.* 16 (sp. dub.).—Arabia.
purgans, Linn. Sp. Pl. 76 = ciliatus.
purgans, Torr. Fl. N. York. ii. 468 = Kalmii.
purpurascens, Delile, Fl. Aeg. Illustr. 52 = rubens.
purpurascens, Turcz. ex Ledeb. Fl. Ross. iv. 357 = inermis.
quitensis, *Willd. ex Steud. Nom.* ed. II. i. 228.—Ecuador.
racemiferus, *Steud. Syn. Pl. Gram.* 323.—Japon.
racemosus, *Linn. Sp. Pl.* ed. II. 114.—Europ.
ramosus, Linn. Mant. i. 34 = Brachypodium ramosum.
ramosus, Huds. Fl. Angl. 40 = asper.
ramosus, Schousb. Vextr. Marok. 44 = Brachypodium pinnatum.
repens, Borb. ex Nym. Consp. 822 = vernalis.
Requieni, Loisel. Fl. Gall. i. 90 = intermedius.
retusus, Pers. Syn. i. 96 = Brachypodium ramosum.
Richardsoni, *Link, Hort. Berol.* ii. 281.—Am. bor.
rigens, Linn. Mant. 33 = rigidus.

BROMUS:—
rigens, Vandelli, in Roem. Script. Pl. Hisp. 49 = Koeleria phleoides.
rigidus, C. A. Mey. Verz. Pfl. Cauc. 23 = rigidus.
rigidus, Reichb. Ic. Fl. Germ. i. 31. n. 1586 = rubens.
rigidus, *Roth, in Roem. & Ust. Mag. Bot.* iv. (1790) 21.—Europ.; Reg. Mediterr. et Caucas.
riparius, *Rehm. Notiz. Veg. Gest. Schwarz. Meer.* 8.—Rossia.
Rochelianus, Opiz, Natural. n. 9 (1825) 119 = madritensis.
rotundatus, H. B. & K. Nov. Gen. et Sp. i. 152 = Briza rotundata.
rubens, Cav. Ic. i. t. 45 = scoparius.
rubens, Delile, Fl. Aeg. Illustr. 52 = fasciculatus.
rubens, Desv. ex Willk. & Lange, Prod. Fl. Hisp. i. 98 = madritensis.
rubens, Host. Gram. Austriac. i. t. 18 = rigidus.
rubens, *Linn. Cent.* i. 5.—Reg. Mediterr.
rupestris, Host, Gram. Austriac. iv. t. 17 = Brachypodium pinnatum.
sabulosus, *Guss. Fl. Sic. Prod.* i. 120.—Sicilia.
sardous, *Spreng. Syst.* iv. *Cur. Post.* 36.—Sardin.
scaberrimus, Tenore, Fl. Nap. iii. 89 = madritensis.
scabratus, *Link, in Linnaea*, xvii. (1843) 402.—Italia.
scabridus, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 231.—Afr. trop.
scabriflorus, Opiz, Natural. n. 9 (1825) 119 = tectorum.
Scheuchzeri, C. C. Gmel. Fl. Bad. iv. 79. t. 7 = Festuca gigantea.
Schraderi, Kunth, Enum. Pl. i. 416 = unioloides.
sclerophyllus, *Boiss. Diagn.* Ser. I. xiii. 65.—Lydia.
scoparius, Guss. Fl. Sic. Prod. 10 = alopecuroides.
scoparius, *Linn. Cent. Pl.* i. 6; *Amoen. Acad.* iv. 266.—Reg. Mediterr.; As. Min.; Persia.
secalinus, Bieb. Fl. Taur. Cauc. i. 71 = patulus.
secalinus, *Linn. Sp. Pl.* 76.—Europ.; Reg. Mediterr.; As. bor.
secalinus, Savi, Fl. Pis. i. 120 = arvensis.
secundus, *J. Presl, in Rel. Haenk.* i. 263.—Sinus Nootka.
segetalis, A. Braun, ex Nym. Consp. 822 = secalinus.
segetum, Cham. & Schlecht. in Linnaea, vi. (1831) 38 = ciliatus.
segetum, *H. B. & K. Nov. Gen. et Sp.* i. 152.—Am. austr.
sericeus, Tenore, Fl. Nap. Prod. 10 = alopecuroides.
serotinus, Beneken, in Bot. Zeit. iii. (1845) 724 = asper.
setaceus, Buckl. in Proc. Acad. Sc. Philad. '1863' (1863) 98 = sterilis.
setifolius, *J. Presl, in Rel. Haenk.* i. 261.—Chili; Mexic.
Sewerzowi, *Regel, in Act. Hort. Petrop.* vii. (1880) 601.—As. centr.
silvaticus, J. F. Gmel. Syst. 190 = sylvaticus.
simplex, Gaud. Agrost. Helv. i. 296 = racemosus.
simplex, *Pers. Syn.* i. 96.—Hab. ?
simplicissimus, Cesati, ex Nym. Consp. 823 = mollis.
sipyleus, *Boiss. Diagn.* Ser. I. xiii. 65.—As. Min.
sitchensis, *Bongard, Veget. Ins. Sitcha*, 173.—Ins. Sitcha.
speciosus, *Nees, Fl. Afr. Austr.* i. 454.—Afr. austr.
spicatus, Nees, in Mart. Fl. Bras. ii. 471 = Diplachne simplex.
spiculitenuata, Knapp, Brit. Gram. t. 81 = arvensis.
squarrosus, Brot. Fl. Lusit. i. 112 = macrostachys.
squarrosus, *Linn. Sp. Pl.* 76.—Europ. austr.; Oriens; As. bor.
squarrosus-muticus, C. A. Mey. Verz. Pfl. Cauc. 24 = briziformis.
stachyoides, *Link, Hort. Berol.* ii. 285.—Hab. ?
stamineus, E. Desv. in C. Gay, Fl. Chil. vi. 440 = unioloides.
stenophyllus, Link, Enum. Hort. Berol. i. 94 = erectus.
stenostachyus, *Boiss. Fl. Orient.* v. 643.—Afghanist.
sterilis, Guss. Fl. Sic. Prod. i. 118 = madritensis.
sterilis, Lang, ex Kumm. & Sendt. in Flora, xxxiii. (1849) 758 = tectorum.
sterilis, *Linn. Sp. Pl.* 77.—Europ.; Oriens; As. bor.
Steudelii, Frank, ex Steud. Nom. ed. II. i. 229 = ciliatus.
stipoides, Linn. Mant. ii. 557 = Festuca geniculata.
strictus, Brougn. in Duperr. Voy. 45 = Brongniartii.
strictus, *Kunth, Rev. Gram.* i. 134.—Ins. Borbon.
strictus, Scop. Fl. Carn. ed. II. i. 79 = Diplachne serotina.
strictus, Hall. f. in Roem. Arch. i. ii. 10 = Brachypodium pinnatum?

BROMUS:—
strigosus, Bieb. Fl. Taur. Cauc. iii. 81 = Triticum strigosum.
strigosus, Lam. Illustr. 191. n. 1063 = Festuca gigantea.
subalpinus, Rupr. in Bull. Acad. Brux. ix. (1842) 2, 236 = exaltatus.
submuticus, *Steud. Syn. Pl. Gram.* 321.—Missouri.
subulatus, Griseb. in Ledeb. Fl. Ross. iv. 358 = Melica acuminata.
Suksdorfii, *Vasey, in Coult. Bot. Gaz.* x. (1885) 223.—Am. bor.
sylvaticus, Pollich, Hist. Pl. Palat. i. 118 = Brachypodium sylvaticum.
sylvaticus, Vogler, Schediasma, 13 = asper.
syriacus, Boiss. & Blanche, Diagn. Ser. II. iv. 139 = erectus.
Tacna, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
subrectus, Fries, Mant. iii. 181 = erectus.
tectorius, Dulac, Fl. Hautes-Pyr. 96 = tectorum.
tectorum, *Linn. Sp. Pl.* 77.—Europ.; Oriens; As. bor.
temulentus, Bernh. Syst. Verz. Erf. 47 = Lolium temulentum.
Tenorianus, Roem. & Schult. Syst. Mant. ii. 360, obs. = sterilis.
tenuiflorus, Viv. Fl. Lyb. Spec. 5. t. 1. f. 1 = fasciculatus.
tenuis, *J. Presl, in Rel. Haenk.* i. 262.—Peruv.
tenuis, Tinco, Rar. Pl. Sic. Pugill. i. 3 = Festuca incrassata.
Thominii, Ard. ex Nym. Consp. 823 = mollis.
Tigridis, Boiss. & Noé, Diagn. Ser. II. iv. 141 = brachystachys.
tigurinus, Suter, Fl. Helv. i. 65 = Brachypodium pinnatum.
tomentellus, *Boiss. Diagn.* Ser. I. vii. 126.—Graecia; As. Min.; Persia.
tomentosus, Rohde, in Schrad. N. Journ. iii. (1809) 42 = macrostachys?
tomentosus, *Trin. in Mém. Acad. Pétersb.* (1813-14) t. 9.—Persia.
tortilis, *Presl, ex Steud. Nom.* ed. II. i. 229.—Europ. austr.
transsilvanicus, Schur, in Oestr. Bot. Zeitschr. x. (1860) 227 = fibrosus.
transylvanicus, Steud. Syn. Pl. Gram. 320 = pracc.
triaristatus, Loisel. Fl. Gall. i. 89 = arduennensis.
trichodes, *Kunth, Rev. Gram.* i. 135.—Ins. Mascar.
trichopodus, *A. Rich. Tent. Fl. Abyss.* ii. 437.—Abyss.
triflorus, Ehrh. Beitr. i. 90 = Festuca sylvatica.
triflorus, Linn. Sp. Pl. ed. II. 115 = Festuca gigantea.
triflorus, Pollich, Hist. Pl. Palat. i. 119 = Ventenata avenacea.
Trinianus, *Schult. Mant.* ii. 254.—Persia.
Trinii, *E. Desv. in C. Gay, Fl. Chil.* vi. 441.—Chili.
trisetoides, *Trin. ex Steud. Nom.* ed. II. i. 229, nomen.—Hab. ?
triticeus, *Raeusch. Nom.* ed. III. 27.—Helvet.
trivialis, Savi, Fl. Pis. i. 124 = Koeleria phleoides.
truncatus, Scop. Fl. Carn. ed. II. i. 85 = Agropyron junceum.
tunicatus, *Phil. in Linnaea*, xxxiii. (1864-65) 298.—Chili.
turgidus, Pers. Syn. i. 95 = macrostachys.
unilateralis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 90 = patulus.
unioloides, *H. B. & K. Nov. Gen. et Sp.* i. 151.—Am. calid.
valdivianus, *Phil. in Linnaea*, xxix. (1857-58) 101.—Chili.
variegatus, Fl. Danub. ex Boiss. Fl. Orient. v. 645 = fibrosus.
variegatus, *Bieb. Fl. Taur. Cauc.* iii. 79.—Reg. Caucas.; Persia.
varius, Brot. Fl. Lusit. i. 113 = rigidus.
velutinus, Schrad. Fl. Germ. i. 349. t. 6. f. 3 = secalinus.
Ventolana, Hort. ex Steud. Nom. ed. II. i. 229 = racemosus.
Ventolana, Schleich. ex Steud. l. c. = arvensis.
vernalis, *Panč. ex Hack. in Oestr. Bot. Zeitschr.* xxix. (1879) 205.—Europ.
versicolor, Pollich, Hist. Pl. Palat. i. 120 = arvensis.
verticillatus, *Cav. Ic.* vi. 66. t. 590.—Hispan.

BROMUS:—
vestitus, *Schrad. in Goett. Gel. Anz.* (1821) III. 2074. —Afr. austr.
villiferus, Steud. Syn. Pl. Gram. 326 = japonicus.
villosulus, *Steud. l. c.* 327.—Graecia.
villosus, Forsk. Fl. Aegypt. Arab. 23 = madritensis, rigidus.
villosus, C. C. Gmel. Fl. Bad. i. 229 = squarrosus.
villosus, Scop. Fl. Carn. ed. II. i. 83 = Agropyron repens.
virens, *Buckl. in Proc. Acad. Sc. Philad.* '1862' (1863) 98.—Am. bor. occ.
virgatus, *J. Presl, in Rel. Haenk.* i. 263.—Peruv.; Mexic.
vitiosus, Weigel, Obs. 4 = secalinus.
Willdenowii, Kunth, Rev. Gram. i. 134 = unioloides.
wolgensis, Willd. Enum. Hort. Berol. Suppl. 9; Jacq. Ecl. Gram. 20. t. 15 = squarrosus.

BROMUS, Scop. Introd. 74 (1777) = **Triticum**, Linn. (Gramin.).

BRONGNIARTIA, Blume, Bijdr. 435 (1825) = **Kibara**, Endl. (Monim.).
coriacea, Blume, l. c. = K. coriacea.

BRONGNIARTIA, H. B. & K. Nov. Gen. et Sp. vi. 465. t. 587 (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 495.
MEGASTEGIA, G. Don, Gen. Syst. ii. 468 (1832).
PERALTEA, H. B. & K. Nov. Gen. et Sp. vi. 469. t. 589 (1823).
Benthamiana, *Hemsl. Biol. Centr. Am. Bot.* i. 252. —Mexic.
foliolosa, *Benth. ex Hemsl. Diag. Pl. Nov.* 7.—Mexic.
galegoides, *Presl, Symb. Bot.* ii. 21. t. 67.—Am. trop.
glabrata, *Hook. & Arn. Bot. Beech. Voy.* 238.—Mexic.
gracilis, *Hemsl. Biol. Centr. Am. Bot.* i. 253.—Mexic.
intermedia, *Moric. in Mém. Soc. Phys. Genèv.* vii. (1836) 253.—Mexic.
magnibracteata, *Schlecht. in Linnaea*, xii. (1838) 338. —Mexic.
minutifolia, *S. Wats. in Proc. Am. Acad.* xx. (1885) 366.—Texas.
mollis, *H. B. & K. Nov. Gen. et Sp.* vi. 465. t. 587.—Mexic.
obliqua, *Schlecht. in Linnaea*, xii. (1838) 339.—Mexic.
oligosperma, *Baill. Adansonia*, ix. (1868–70) 240.—Mexic.
oxyphylla, *Hemsl. Biol. Centr. Am. Bot.* i. 254.—Mexic.
Parryi, *Hemsl. l. c.*—Mexic.
podalyrioides, *H. B. & K. Nov. Gen. et Sp.* vi. 465. t. 588.—Mexic.
retusa, *Benth. in Hemsl. Diag. Pl. Nov.* 8.—Mexic.
robinioides, *Kunth, Ind. Sem. Hort. Berol.* (1845) 12.—Chili.
sericea, *Schlecht. in Linnaea*, xii. (1838) 338.—Mexic.
stipitata, *Hemsl. Diag. Pl. Nov.* 8.—Mexic.
thermoides, *Spreng. Syst.* iv. *Cur. Post.* 287.—Mexic.
vicioides, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) 49.—Mexic.

BRONNIA, H. B. & K. Nov. Gen. et Sp. vi. 83. t. 528 (1823) = **Fouquieria**, H. B. & K. (Tamarisc.).
spinosa, H. B. & K. l. c. = F. spinosa.

BROOKEA, Benth. in Benth. & Hook. f. Gen. ii. 939 (1876). *SCROPHULARINEAE*, Benth. & Hook. f. l. c.
dasyantha, *Benth. in Hook. Ic. Pl.* xii. 84. t. 1197.—Borneo.
tomentosa, *Benth. l. c.*—Borneo.

BROSIMUM, Sw. Prod. Veg. Ind. Occ. 12 (1788). *URTICACEAE*, Benth. & Hook. f. iii. 369.
GALACTODENDRUM, Humb. Relat. Hist. ii. 108 (1819).
PIRATINERA, Aubl. Pl. Gui. ii. 888. t. 340 (1775), nomen prius.
Alicastrum, *Sw. l. c.*—Am. trop.

BROSIMUM:—
Aubletii, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 34. t. 148. f. a.—Am. trop.
costaricanum, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 334.—Costa Rica.
discolor, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 403.—Bras.
echinocarpum, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 34. t. 148.—Peruv.
Galactodendron, *D. Don, ex Sweet, Hort. Brit.* ed. II. 462.—Guiana.
Gaudichaudii, *Trécul, in Ann. Sc. Nat.* Sér. III. viii. (1847) 139.—Bras.
Humboldtii, Carr. in Rev. Hortic. (1874) 314. f. 42, 43 = B. Galactodendron.
microcarpum, Schott, in Spreng. Syst. iv. Cur. Post. 403 = B. Alicastrum.
refractum, Mart. ex Miq. in Mart. Fl. Bras. iv. 1. 108 = Gaudichaudii.
spurium, *Sw. Prod. Veg. Ind. Occ.* 12.—Jamaica.
turbinatum, *Spruce, ex Baill. Adansonia*, xi. (1875) 299.—Bras.

BROSSAEA, [Plum.] Linn. Gen. ed. I. 366 (1737) = **Gaultheria**, Kalm. (Ericac.).
anastomosans, Griseb. Fl. Brit. W. Ind. 142 = G. anastomosans.
coccinea, Linn. Sp. Pl. 1190 = G. cordifolia.

BROSSARDIA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 183. *CRUCIFERAE*, Benth. & Hook. f. i. 91.
papyracea, *Boiss. l. c.* 184.—Persia.
retusa, *Boiss. Fl. Orient.* i. 336.—Persia.

BROTERA, Cav. in Anal. Cienc. Nat. i. (1800) 33; Ic. v. 19. t. 433. (1799) = **Melhania**, Forsk. (Sterculiac.).
bracteosa, Guill. & Perr. Fl. Seneg. Tent. i. 86 = M. Denhami.
Leprieurii, Guill. & Perr. l. c. = Pentapetes Leprieurii.
ovata, Cav. Ic. v. 20 = M. abyssinica.

BROTERA, Spreng. in Schrad. Journ. Bot. iv. (1800) t. 5 = **Flaveria**, Juss. (Compos.).
Contrayerba, Spreng. l. c. = F. repanda.
Sprengelii, Cass. in Dict. Sc. Nat. xxxiv. 306 = F. repanda.
trinervata, Pers. Syn. ii. 498 = F. repanda.

BROTERA, Spreng. Nachr. i. Bot. Gart. Halle, (1801) 15; et in Trans. Linn. Soc. vi. (1802) 151. t. 12 = **Hyptis**, Jacq. (Labiat.).
persica, Spreng. ll. cc. = H. pectinata.

BROTERA, Vell. Fl. Flum. 322 (1825); vii. t. 163 (1827) = **Luhea**, Willd. (Tiliac.).
maritima, Vell. ll. cc. 323; t. 163 = L. speciosa.
mediterranea, Vell. ll. cc. 323; t. 164 = L. divaricata.

BROTERA, Willd. Sp. Pl. iii. 2399 (1803) = **Cardopatium**, Juss. (Compos.).
corymbosa, Willd. l. c. = C. corymbosum.

BROTEROA, DC. Prod. v. 635 (1836) = **Flaveria**, Juss. (Compos.).
trinervata, DC. l. c. 636 = F. repanda.

BROUGHTONIA, R. Br. in Ait. Hort. Kew. ed. II. v. 217 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 531.
alba, Spreng. Syst. iii. 735 = Maxillaria alba.
aurea, Lindl. Bot. Reg. (1840) Misc. 19 = Epidendrum aurantiacum.
chinensis, Lindl. in Hook. Lond. Journ. Bot. i. (1842) 492 = Epidendrum nonchinense.
coccinea, Hook. Bot. Mag. t. 3536 = sanguinea.
grandiflora, Spreng. Syst. iii. 735 = Maxillaria grandiflora.
lilacina, *Henfr. in Moore & Ayres, Gard. Mag. Bot.* iii. (1851) 201.—Ind. occ.?
maculata, Spreng. Syst. iii. 735 = Maxillaria maculata.
sanguinea, *R. Br. in Ait. Hort. Kew.* ed. II. v. 217.—Ind. occ.
tetragona, Spreng. Syst. iii. 734 = Phajus tetragonus.
violacea, Hort. ex Moore & Ayres, Gard. Mag. Bot. iii. (1851) 201 = lilacina.

BROUGHTONIA, Wall. ex Lindl. Gen. et Sp. Orch. 35 (1830) = **Otochilus**, Lindl. (Orchid.).
amoena, Wall. l. c. = O. fusca.
nitida, Herb. ex Sweet, Hort. Brit. ed. III. 641, nomen (= Coelogynes sp.?).—Hab.?
linearis, Wall. l. c. 42 = Coelogyne ovalis.
pendula, Wall. l. c. 35 = O. alba.
pilosa, Hook. ex Steud. Nom. ed. II. i. 230.—Ind. or.

BROUSSAISIA, Gaudich. Freyc. Voy. Bot. 479. t. 69. (1826). *SAXIFRAGEAE*, Benth. & Hook. f. i. 641.
arguta, *Gaudich. l. c.*—Ins. Sandvic.
pellucida, *Gaudich. Bot. Voy. Bonite*, t. 9.—Ins. Sandvic.

BROUSSONETIA, Orteg. Hort. Matr. 61. t. 7 (1798) = **Sophora**, Linn. (Legumin.).
secundiflora, Orteg. l. c. = S. secundiflora.
speciosa, Orteg. l. c. t. 7 = S. secundiflora.

BROUSSONETIA, L'Hérit. ex Vent. Tabl. Regn. Veget. iii. 547 (1799). *URTICACEAE*, Benth. & Hook. f. iii. 361.
PAPYRIUS, Lam. Illustr. t. 762 (1798).
STENOCHASMA, Miq. Pl. Jungh. i. 45 (1850).
Billardii, Carr. in Rev. Hortic. (1878) 375 = papyrifera.
brasiliensis, Mart. in Flora, xxiv. (1841) Beibl. 10 = Maclura brasiliensis.
cordata, *Blume, Mus. Bot. Lugd. Bat.* ii. 87.—China.
cordata, Hort. ex C. Koch, Dendrol. ii. II. 440 = papyrifera.
cucullata, Hort. ex Steud. Nom. ed. II. i. 230 = papyrifera.
dissecta, Hort. ex Bur. in DC. Prod. xvii. 225 = papyrifera.
elegans, Hort. ex C. Koch, Dendrol. ii. II. 440 = papyrifera.
integrifolia, *Buch.-Ham. in Trans. Linn. Soc.* xvii. (1837) 195.—Ind. or.
Kaempferi, *Siebold, in Verh. Bat. Gen.* xii. (1830) 28. —Japon.
Kazi, Siebold, ex Blume, Mus. Bot. Lugd. Bat. ii. 86 = papyrifera.
Kazinoki, *Siebold, in Verh. Bat. Gen.* xii. (1830) 28.—Japon.
luzonica, *Bur. in DC. Prod.* xvii. 224.—Ins. Philipp.
luzoniensis, Blanco, Fl. Filip. ed. II. 488 = Allaeanthus luzonicus.
maculata, Hort. ex Steud. Nom. ed. II. i. 230 = papyrifera.
monoica, *Hance, in Journ. Bot.* xx. (1882) 294.—China.
navicularis, Lodd. ex C. Koch, Dendrol. ii. II. 440 = papyrifera.
nana, Hort. ex Bur. in DC. Prod. xvii. 225 = papyrifera.
navicularis, Lodd. ex Bur. l. c. = papyrifera.
navifolia, Hort. ex Steud. Nom. ed. II. i. 230 = papyrifera.
papyrifera, *Vent. Tabl. Regn. Veget.* iii. 547.—Malaya; Ins. Pacific.
Plumerii, Spreng. Syst. iii. 901 = Chlorophora tinctoria.
Sieboldii, Blume, Mus. Bot. Lugd. Bat. ii. 86 = B. Kazinoki.
spathulata, Hort. ex Steud. Nom. ed. II. i. 230 = papyrifera.
tinctoria, Blanco, Fl. Filip. ed. II. 488 = Cudrania javanensis.
tinctoria, H. B. & K. Nov. Gen. et Sp. ii. 32 = Chlorophora tinctoria.
tinctoria, Spreng. Syst. ii. 901 = Maclura tinctoria.
tricolor, Hort. ex C. Koch, Dendrol. ii. II. 440 = papyrifera.
xanthoxylon, Mart. in Flora, xxiv. (1841) Beibl. 10 = Maclura xanthoxylon.

BROUVALEA, Adans. Fam. ii. 211 (1763) = seq.

BROVALLIA, Linn. Orb. Nat. (1764) = seq.

BROWALLIA, Linn. Gen. ed. I. 348 (1737). *SOLANACEAE*, Benth. & Hook. f. ii. 910.
abbreviata, *Benth. Bot. Voy. Sulph.* 144.—Peruv.
alienata, Linn. Syst. ed. X. 1118 = Ruellia paniculata.
americana, Linn. Sp. Pl. 631 = demissa.
cordata, *G. Don, Gen. Syst.* iv. 478.—Peruv.

BROWALLIA:—
demissa, *Linn. Syst.* ed. X. 1118.—Am. austr.
elata, Linn. l. c. = demissa.
elongata, H. B. & K. Nov. Gen. et Sp. ii. 372 = demissa.
grandiflora, *R. Grah. in Edin. N. Phil. Journ.* 'Jan.–Apl. 1830' [1829].—Peruv.
grandiflora, Lindl. Bot. Reg. t. 1384 = cordata.
humifusa, Forsk. Fl. Aegypt. Arab. 112 = Striga humifusa.
Jamesonii, *Benth. in DC. Prod.* x. 197.—Peruv.; N. Granat.
lactea, Hort. ex G. Don, in Sweet, Hort. Brit. ed. III. 508 = demissa.
Linnaeana, Spreng. Syst. ii. 808 = demissa.
nervosa, Miers, in Ann. & Mag. Nat. Hist. Ser. II. v. (1850) 206 = demissa.
pedunculariis, *Benth. Bot. Voy. Sulph.* 143.—Peruv.
pulchella, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 176 = viscosa.
pulchella, *Lehm. in Hamb. Gartenz.* x. (1854) 375.—Hab.?
speciosa, *Hook. Bot. Mag.* t. 4339.—N. Granat.
tenella, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. v. (1850) 205.—Bras.
texana, Torr. Bot. Mex. Bound. 156 = Nierembergia viscosa.
viscosa, *H. B. & K. Nov. Gen. et Sp.* ii. 373.—Am. austr.

BROWNEA, Jacq. Enum. Pl. Carib. 6 (1760) (Brownaea). *LEGUMINOSAE*, Benth. & Hook. f. i. 577.
Hermesias, Loefl. Iter, 278 (1758).
antioquensis, Linden, Cat. (1869) 3 = macrophylla.
Ariza, *Benth. Pl. Hartw.* 171.—N. Granat.
Ariza, Lindl. & Paxt. Flow. Gard. ii. 107. t. 59 = grandiceps.
Birschellii, *Hook. f. Bot. Mag.* t. 5998.—Venezuela.
capitella, *Jacq. Fragm.* 26. t. 18, 19.—Venezuela.
cauliflora, *Poepp. & Endl. Gen. et Sp. Pl. Chil.* iii. 32.—Chili.
coccinea, *Jacq. Enum. Pl. Carib.* 26.—Venezuela.
coccinea, Loefl. ex Griseb. Fl. Brit. W. Ind. 212 = B. Rosa-de-monte.
erecta, Hort. Belg. ex Planch. in Fl. des Serres, vi. (1850–51) 154 (= Theophrasta pinnata).
grandiceps, *Jacq. Collect.* iii. 287. t. 22.—Am. austr.
guianensis, *Klotzsch, in Schomb. Faun. & Fl. Guy.* 1209, nomen.—Guiana.
latifolia, *Jacq. Fragm.* 25. t. 17.—Venezuela.
leucantha, *Jacq. l. c.* 26.—Venezuela.
macrophylla, *Linden, Cat.* n. 18 (1863) 11; *Mast. in Gard. Chron.* (1873) 777. f. 149.—N. Granat.
negrensis, *Benth. in Mart. Fl. Bras.* xv. ii. 212.—Bras.
pauciflora, Willd. Sp. Pl. iii. 716 = Palovea guianensis.
Princeps, Linden, ex Otto, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 147 = B. Ariza.
racemosa, *Jacq. Fragm.* 25. t. 16.—Venezuela.
Rosa, Pers. Syn. ii. 237 = B. Rosa-de-monte.
Rosa-de-monte, *Berg. in Phil. Trans.* (1771) 171. t. 8, 9.—Am. austr.
rosea, Otto, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 147 = B. Rosa-de-monte.
speciosa, Reichb. ex DC. Prod. ii. 477 = B. Rosa-de-monte.

BROWNETERA, Rich. in Ann. Mus. Hist. Nat. Par. xvi. (1810) 299 = **Phyllocladus**, Rich. (Conifer.).
aspleniifolia, Rich. l. c. = Ph. rhomboidalis.

BROWNLEEA, Harv. ex Lindl. in Hook. Lond. Journ. Bot. i. (1842) 16. *ORCHIDEAE*, Benth. & Hook. f. iii. 631.
caerulea, *Harv. ex Lindl. l. c.*—Afr. austr.
macroceras, Sond. in Linnaea, xix. (1847) 106 = caerulea.
parviflora, *Harv. ex Lindl. in Hook. Lond. Journ. Bot.* i. (1842) 16.—Afr. austr.
recurvata, *Sond. in Linnaea*, xix. (1847) 107.—Afr. austr.

BROWNLOWIA, Roxb. Pl. Corom. iii. 61. t. 265 (1819). *TILIACEAE*, Benth. & Hook. f. i. 231.
Humea, Roxb. Fl. Ind. ii. 640 (1832).
argentata, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) ii. 67.—Ind. or.
elata, *Roxb. Pl. Corom.* iii. 61. t. 265.—Ind. or.; Burma.
lanceolata, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 57.—Ind. or.; Burma.
peltata, *Benth. l. c.* 56.—Borneo.

BROWNLOWIA:—
Riedelii, *Hemsl. Bot. Voy. Challenger*, i. iii. 128.—Ins. Molucc.

BRUCEA, J. S. Mill. Fasc. t. 25 (1780). *SIMARUBEAE*, Benth. & Hook. f. i. 311.
Gonus, Lour. Fl. Cochinch. ii. 658 (1790).
Nima, Buch.-Ham. ex A. Juss. in Mém. Mus. Par. xii. 516 (1825).
abyssinica, Spreng. Pugill. ii. 90, in obs. = ferruginea.
antidysenterica, Lam. in Mém. Acad. Par. (1784) 342; Eneye. i. 471 = ferruginea.
dubia, Steud. Nom. ed. II. i. 230 = Picrasma nepalensis.
ferruginea, *L'Hérit. Stirp. Nov.* 19. t. 10.—Abyss.
glabrata, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 447.—Ins. Timor.
gracilis, DC. Prod. ii. 88 = sumatrana.
guineensis, G. Don, Gen. Syst. i. 801 = paniculata.
macrophylla, *Oliver, Fl. Trop. Afr.* i. 310.—Afr. trop.
mollis, *Wall. Cat.* n. 8483.—Reg. Himal.
paniculata, *Lam. Encyc.* i. 472.—Afr. trop.
quercifolia, *Seem. Fl. Vit.* 33.—Ins. Fiji.
sumatrana, *Roxb. Hort. Beng.* 12; *Fl. Ind.* i. 449.—As. trop.; Austral.
sumatrensis, Spreng. Pugill. ii. 90 = sumatrana.
trichotoma, *Spreng. Syst.* i. 441.—Cochinch.
trifoliata, Spreng. l. c. iv. Cur. Post. 43 = Evodia drupacea.
virginica, *Hornem. ex Steud. Nom.* ed. I. 122, nomen.—Virginia.

BRUCHMANNIA, Nutt. Gen. Am. i. 48 (1818) = **Beckmannia**, Host (Gramin.).
erucaeformis, Nutt. l. c. = Beckmannia erucaeformis.

BRUCKENTHALIA, Reichb. Fl. Germ. Excurs. 414 (1831). *ERICACEAE*, Benth. & Hook. f. ii. 591.
spiculifolia, *Reichb. l. c.*—Europ. austr. or.; As. Min.

BRUEA, Gaudich. in Freyc. Voy. Bot. 511 (1826). *URTICACEAE*? (Gen. dubium) Benth. & Hook. f. iii. 351.
bengalensis, *Gaudich. l. c.* (= Laportea terminalis?).—Ind. or.

BRUECKEA, Klotzsch & Karst. in Karst. Ausw. Gew. Venez. 31. t. 10 (1848) = **Aegiphila**, Jacq. (Verbenac.).
grandifolia, Klotzsch & Karst. l. c. 32 = A. verrucosa.

BRUENNICHIA, Willd. Sp. Pl. ii. 751 (1800) = **Brunnichia**, Banks (Polygon.).

BRUGMANSIA, Blume, in Van Hall, Bijdr. Nat. Wet. ii. (1828) 422. *CYTINACEAE*, Benth. & Hook. f. iii. 119.
Mycetanthe, Reichb. Nom. 61 (1841).
Zippelia, Reichb. ex Endl. Gen. Suppl. ii. 6 (1842).
Lowi, Becc. in Atti Soc. Ital. Sc. Nat. xi. (1868) 198 = Zippelii.
Zippelii, *Blume, in Van Hall, Bijdr. Nat. Wet.* ii. (1828) 422.—Java.

BRUGMANSIA, Pers. Syn. i. 216 (1805) = **Datura**, Linn. (Solanac.).
arborea, Steud. Nom. ed. II. i. 230 = D. arborea.
aurantiaca, Hort. ex Walp. Rep. iii. 934 = Juanulloa parasitica.
bicolor, Pers. Syn. i. 216 = D. sanguinea.
candida, Pers. l. c. = D. arborea.
floribunda, Paxt. Mag. Bot. ix. 3 = Juanulloa aurantiaca.
Knightii, Hort. ex Dun. in DC. Prod. xiii. i. 690 = D. cornigera.
parviflora, Hort. ex Walp. Rep. iii. 934 = Juanulloa parasitica.
sanguinea, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. t. 272 = D. sanguinea.
suaveolens, Bercht. & Presl, Rostl. i. Solaneae. 45 = D. suaveolens.
Waymanni, Paxt. Mag. Bot. iv. 241 = D. fastuosa.

BRUGUIERA, Lam. Encyc. iv. 696 (1796). *RHIZOPHOREAE*, Benth. & Hook. f. i. 679.
Kanilia, Blume, Mus. Bot. Lugd. Bat. i. 140 (1849).
Paletuviera, Thou. ex DC. Prod. iii. 32 (1828).
Arnottiana, Wight, ex Arn. in Ann. Nat. Hist. i. (1838) 364 = Ceriops Candolleana.
australis, A. Cunn. ex Arn. l. c. 368 = Rheedii.
Candel, Steud. Nom. ed. II. i. 231 = Kandelia Rheedii.

BRUGUIERA:—
capensis, Blume, Mus. Bot. Lugd. Bat. i. 137 = gymnorrhiza.
caryophyllaeoides, *Blume, Enum. Pl. Jav.* i. 93.—Ind. or.; Malaya.
cylindrica, Blume, l. c. i. 91 = caryophyllaeoides, malabarica.
decangulata, Griff. Notul. iv. 669 = Rhizophora Mangle.
eriopetala, *Wight & Arn. in Ann. Nat. Hist.* i. (1838) 368.—As. trop.
gymnorrhiza, *Lam. Illustr.* t. 397; *Blume, Mus. Bot. Lugd. Bat.* i. 136.—Geront. trop.
malabarica, *Arn. in Ann. Nat. Hist.* i. (1838) 369.—Ind. or.
nemorosa, Blanco, Fl. Filip. ed. II. 275 = Carallia integerrima.
nubicola, Pritz. Ic. Ind. ii. 51, sphalm. = Bougueria nubicola.
obtusa, Steud. Nom. ed. II. i. 231 (= *Problastes cuneifolia*).—Java.
oxyphylla, *Miq. Fl. Ind. Bat. Suppl.* i. 324.—Sumatra.
parietosa, Griff. Notul. iv. 670; Ic. Pl. Asiat. t. 641 = eriopetala.
parviflora, *Wight & Arn. Prod.* 311.—Ind. or.; Malaya.
Rheedii, *Blume, Enum. Pl. Jav.* i. 92.—Geront. trop.
Rumphii, Blume, Mus. Bot. Lugd. Bat. i. 138 = Rheedii.
sexangula, Poir. Encyc. Suppl. iv. 262 = Rhizophora sexangula.
sexangularis, Spreng. Syst. ii. 602 = Rhizophora sexangula.
timoriensis, Wight & Arn. Prod. 311 = Ceriops Candolleana.
Wightii, Blume, Mus. Bot. Lugd. Bat. i. 138 = gymnorrhiza.
Zippelii, Blume, l. c. 139 = Rheedii.

BRUGUIERA, Thou. Gen. Nov. Madag. 21 (1806) = **Lumnitzera**, Willd. (Combret.).
littorea, Steud. Nom. ed. II. i. 231 = L. coccinea.
madagascariensis, DC. Prod. iii. 23 = L. racemosa.

BRUGUIERA, Rich. ex DC. Prod. iii. 173, in syn. (1828) = **Conostegia**, D. Don (Melastom.).

BRUINSMANIA, Miq. in Linnaea, xvii. (1843) 72 = **Isertia**, Schreb. (Rubiac.).
isertioides, Miq. l. c. 73.—Guiana.

BRUNELLA, Tourn. ex Linn. Syst. ed. I (1735) = **Prunella**, Linn. (Labiat.).

BRUNELLIA, Ruiz & Pav. Prod. Fl. Per. 71. t. 12 (1794). *SIMARUBEAE*, Benth. & Hook. f. i. 313.
aculeata, *Ruiz & Pav. Fl. Per.* iv. 71. t. 415; *Syst.* 127.—Peruv.
acutangula, *Humb. & Bonpl. Pl. Aequin.* i. 216. t. 62.—N. Granat.
comocladifolia, *Humb. & Bonpl. l. c.* i. 211. t. 59.—Peruv.
Funkiana, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 267.—N. Granat.
Goudotii, *Tul. l. c.* 270.—N. Granat.
inermis, *Ruiz & Pav. Fl. Per.* iv. 71. t. 414; *Syst.* 127.—Peruv.
ovalifolia, *Humb. & Bonpl. Pl. Aequin.* i. 216. t. 61.—Peruv.
propinqua, *H. B. & K. Nov. Gen. et Sp.* vii. 45.—Am. austr.
quadrilocularis, *Hook. & Arn. Bot. Beech. Voy.* 282.—Mexic.
racemifera, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 269.—N. Granat.
rugosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxxii. (1859) i. 274.—Bras.
sandwicensis, Gaudich. Freyc. Voy. Bot. 93; ex Hook. & Arn. Bot. Beech. Voy. 80 = Pelea sandwicensis.
tomentosa, *Humb. & Bonpl. Pl. Aequin.* 214. t. 60.—Am. austr.

BRUNFELSIA, Plum. ex Linn. Hort. Cliff. 5 (1737) (Brunsfelsia; Gen. ed. I. 332. *SOLANACEAE*, Benth. & Hook. f. ii. 911.
Franciscea, Pohl, Pl. Bras. Ic. i. 1. t. 1–7 (1827).
acuminata, Benth. in DC. Prod. x. 199 = ramosissima.
americana, *Linn. Sp. Pl.* 191.—Am. trop.

BRUNFELSIA :—
australis, Benth. in DC. Prod. x. 200 = Hopeana.
bahiensis, *A. DC. Prod.* x. 590.—Am. austr.
calycina, Benth. in DC. Prod. x. 199 = pauciflora.
capitata, *Benth. l. c.* 198.—Bras.
cestroides, *A. Rich. Fl. Cub. Fanerog.* ii. 151.—Cuba.
confertiflora, Benth. in DC. Prod. x. 199 = ramosissima.
cuneifolia, *J. A. Schmidt, in Mart. Fl. Bras.* viii. 259. —Bras.
eximia, Bosse, Handb. Blumeng. 524 = macrophylla.
falcata, *Regel, Cat. Pl. Hort. Aksakov.* 21.—Hab. ?
fallax, *Duchass. ex Griseb. in Goett. Abh.* vii. (1857) 242.—Ind. occ.
gracilis, *Regel, Cat. Pl. Hort. Aksakov.* 21.—Hab. ?
grandiflora, *D. Don, in N. Edin. Phil. Journ.* (Apr.–Oct. 1829) 86.—Peruv.
guianensis, *Benth. in DC. Prod.* x. 200.—Guiana.
Hopeana, *Benth. l. c.*—Bras.
hydrangeaeformis, *Benth. l. c.* 198.—Bras.
inodora, *Mart. Del. Sem. Hort. Monac.* (1847).—*Cf. Linnaea*, xxiv. (1851) 180.—Bras.
jamaicensis, *Griseb. Fl. Brit. W. Ind.* 432.—Ind. occ.
latifolia, *Benth. in DC. Prod.* x. 199.—Bras.
latifolia, Hort. ex Steud. Nom. ed. II. i. 231 = americana.
Lockharti, *Hort. Belg. ex Heynh. Nom.* ii. 79, nomen. —Ins. Trinitat. ?
longiflora, *Regel, Cat. Pl. Hort. Aksakov.* 21.—Hab. ?
longituba, *Lem. Jard. Fleur.* iv. (1854) *Misc.* 61.—Cuba.
macrophylla, *Benth. in DC. Prod.* x. 198.—Bras.
maritima, *Benth. l. c.* 200.—Bras.
montana, *Lodd. ex Loud. Hort. Brit.* 253, nomen.—Ind. occ.
multiflora, *Regel, Cat. Pl. Hort. Aksakov.* 21.—Hab. ?
mutabilis, *Poit. in Rev. Hortic.* Sér. II. i. (1842) 261. —Hab. ?
nitida, *Benth. in DC. Prod.* x. 201.—Am. trop.
obovata, *Benth. l. c.* 199.—Bras.
parviflora, A. Rich. Fl. Cub. Fanerog. ii. 327 = seq.
parvifolia, A. Rich. l. c. 151 = nitida.
pauciflora, *Benth. in DC. Prod.* x. 199.—Bras.
purpurea, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 523.—Ind. occ.
ramosissima, *Benth. in DC. Prod.* x. 199.—Bras.
Schomburgkiana, *Klotzsch, in Schomb. Fauna et Fl. Guy.* 1155, nomen.—Guiana.
Sieberi, *Regel, Cat. Pl. Hort. Aksakov.* 21.—Hab. ?
sinuata, *A. Rich. Fl. Cub. Fanerog.* ii. 151.—Am. trop.
spinosa, *Jacq. Enum. Pl. Carib.* 14.—Ind. occ.
Spruceana, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 258 = maritima.
terminalis, Salisb. Prod. 109 = americana.
undulata, *Sw. Prod. Veg. Ind. Occ.* 90.—Jamaica.
uniflora, *D. Don, in Edinb. N. Phil. Journ.* (Apr.–Oct. 1829) 85.—Hab. ?
vinciflora, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 523.—Ind. occ.
violacea, *Lodd. Bot. Cab.* t. 792.—Am. austr.

BRUNIA, Linn. Syst. ed. I (1375). *BRUNIACEAE*, Benth. & Hook. f. i. 671.
BAECKIA, Burm. f. Prod. Fl. Cap. 12 (1768).
abrotanifolia, F. G. Dietr. Vollst. Lex. Gaertn. ii. 320 = Berzelia abrotanoides.
abrotanoides, Linn. Sp. Pl. 199 = Berzelia abrotanoides.
alopecuroides, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 375.—Afr. austr.
alopecuroides, Thunb. Prod. Pl. Cap. 187 = Berzelia alopecuroides.
arachnoidea, Wendl. Coll. ii. 62 = Berzelia squarrosa.
brevifolia, D. Dietr. Syn. Pl. i. 848 = Berzelia abrotanoides.
candicans, *Hort. ex Steud. Nom.* ed. II. i. 231.—Afr. austr.
capitata, Desf. Tabl. Hort. Par. ed. II. 232 = Staavia capitella.
capitella, Thunb. Diss. Brun. 5 = Staavia capitella.
capitellata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 316 = staavioides.
ciliata, *Linn. Sp. Pl.* 199.—Afr. austr.
comosa, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 17 = Berzelia lanuginosa.
cordata, Walp. Rep. i. 544 = racemosa.
cordifolia, D. Dietr. Syn. Pl. i. 848 = Berzelia cordifolia.
coronata, Linn. Mant. ii. 210, in syn. = Staavia glutinosa.
cupressina, Linn. l. c. ii. 343 = Diosma cupressina.

BRUNIA :—
deusta, *Thunb. Prod. Pl. Cap.* 187.—Afr. austr.
deusta, Willd. in Denkschr. Acad. Muench. (1808) 127. t. 7. f. 2 = Nebelia phylicoides.
Dregeana, *Presl, Bot. Bemerk.* 39.—Afr. austr.
elegans, *Dum.-Cours. Bot. Cult.* ed. I. iii. 616.—Afr. austr.
ericoides, Wendl. Coll. ii. 57 = Berzelia squarrosa.
flagelliformis, *Hort. ex Steud. Nom.* ed. II. i. 231 (quid ?).—Hab. ?
formosa, D. Dietr. Syn. Pl. i. 848 = Berzelia abrotanoides.
formosa, *Dum.-Cours. Bot. Cult.* ed. I. iii. 616.—Afr. austr.
fragarioides, Willd. Sp. Pl. i. 1143 (= *Berardia globosa*). —Afr. austr.
glabrata, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1803) 18.—Afr. austr.
globosa, Eckl. & Zeyh. Enum. 139 = laevis.
globosa, Thunb. Diss. Brun. 4 (= *Berardia globosa*).—Afr. austr.
glutinosa, Berg. Desc. Pl. Cap. 57 = Staavia glutinosa.
hirsuta, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1803) 19.—Afr. austr.
imbricata, Sweet, Hort. Brit. ed. II. 115 = racemosa.
imbricata, *Wendl. f. ex Hoffm. Verz. Pfl. Nachtr.* i. 228, ii. & iii. 26, 71.—Afr. austr.
intermedia, D. Dietr. Syn. Pl. i. 848 = Berzelia intermedia.
laevis, *Thunb. Prod. Fl. Cap.* 187.—Afr. austr.
lancifolia, *Walp. Rep.* i. 544.—Afr. austr.
lanuginosa, Linn. Sp. Pl. 199 = Berzelia lanuginosa.
lanuginosa, Thunb. Prod. Pl. Cap. 41 = Berzelia squarrosa.
laurifolia, Sond. in Harv. & Sond. Fl. Cap. ii. 315 = lancifolia.
laxa, Thunb. Prod. Pl. Cap. 187 = Tittmannia laxa.
Levisanus, Linn. Sp. Pl. 199 = Leucadendron Levisanus.
macrocephala, *Willd. in Denkschr. Acad. Muench.* i. (1809) 152.—Afr. austr.
microcephala, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 320 (= *Berardia sphaerocephala*).—Afr. austr.
microphylla, Thunb. Prod. Pl. Cap. 187 (= *Berardia microphylla*).—Afr. austr.
nodiflora, *Linn. Sp. Pl.* 199.—Afr. austr.
paleacea, Berg. Desc. Pl. Cap. 56 = Nebelia paleacea.
paleacea, Wendl. Coll. t. 2 = Berzelia intermedia.
passerinoides, *Schlecht. in Linnaea*, vi. (1831) 190.—Afr. austr.
phylicoides, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 321 (= *Berardia Dregeana*).—Afr. austr.
phylicoides, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 18 = Nebelia phylicoides.
pinifolia, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 374.—Afr. austr.
plumosa, Lam. Encyc. i. 475 = Berzelia squarrosa.
pragarioides, DC. Prod. ii. 45, err. typ. = *fragarioides*.
Protea, Crantz, Inst. i. 357 = Berzelia abrotanoides.
racemosa, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 374.—Afr. austr.
racemosa, Sond. in Harv. & Sond. Fl. Cap. ii. 315 = lancifolia.
radiata, Berg. Desc. Pl. Cap. 58 = Staavia glutinosa.
radiata, Linn. Mant. ii. 209 = Staavia radiata.
rubra, Willd. in Denkschr. Acad. Muench. i. (1809) 131. t. 7 = Berzelia squarrosa.
sericea, Hort. ex Dum.-Cours. Bot. Cult. ed. II. vii. 329 = Berzelia superba.
speciosa, Hort. ex Dum.-Cours. l. c. vi. 279; vii. 329 = Berzelia superba.
squalida, *Sond. in Harv. & Sond. Fl. Cap.* ii. 315.—Afr. austr.
squarrosa, Sw. ex Harv. & Sond. l. c. 311 = Berzelia abrotanoides.
squarrosa, Thunb. Diss. Brun. 5 = Berzelia squarrosa.
staavioides, *Sond. in Harv. & Sond. Fl. Cap.* ii. 316. —Afr. austr.
superba, Donn, Ind. Hort. Cantab. 25 = Berzelia lanuginosa.
superba, Krauss, ex Harv. & Sond. Fl. Cap. ii. 314 = laevis.
tenuifolia, Willd. in Denkschr. Acad. Muench. i. (1809) 129 = Berzelia lanuginosa.
teres, *Oliver, in Journ. Linn. Soc.* ix. (1867) 333.—Afr. austr.
Thunbergiana, D. Dietr. Syn. Pl. i. 849 = Nebelia paleacea.
thyrsophora, *Walp. Rep.* i. 544.—Afr. austr.

BRUNIA :—
uniflora, Linn. Sp. Pl. 199 = Diosma cupressina.
verticillata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 322 = Staavia nuda.
verticillata, *Linn. f. Suppl.* 156.—Afr. austr.
verticillata, Thunb. Prod. Pl. Cap. 41 = virgata.
villosa, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 315.—Afr. austr.
virgata, *Brongn. in Ann. Sc. Nat.* Sér. I. viii. (1826) 376.—Afr. austr.

BRUNIERA, Franch. in Billotia, (1864) 25 = **Wolffia**, Horkel. (Lemnac.).
vivipara, Franch. l. c. = W. Michelii.

BRUNNERA, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 582 = **Anchusa**, Linn. (Boragin.).
myosotidiflora, Stev. l. c. = A. myosotidiflora.
sibirica, Stev. l. c.—Sibir.

BRUNNICHIA, Banks, ex Gaertn. Fruct. i. 213. t. 45 (1788). *POLYGONACEAE*, Benth. & Hook. f. iii. 103.
FALLOPIA, Adans. Fam. ii. 274, 277 (1763).
RAJANIA, Walt. Fl. Carol. 247 (1788).
africana, *Welw. in Trans. Linn. Soc.* xxvii. (1872) 61. —Afr. trop.
cirrhosa, *Gaertn. Fruct.* i. 214. t. 45.—Am. bor.

BRUNONIA, Sm. in Trans. Linn. Soc. x. (1811) 365. t. 28, 29. *GOODENOVIEAE*, Benth. & Hook. f. ii. 541.
australis, *Sm. l. c.* 367. t. 28.—Austral.
sericea, Sm. l. c. 367. t. 29 = australis.
simplex, Lindl. in Mitch. Trop. Austr. 82 = australis.

BRUNSFELSIA, Plum. ex Linn. Hort. Cliff. 5 (1737) = **Brunfelsia**, Linn. (Solanac.).

BRUNSVIA, Neck. Elem. iii. 337 (1790) = **Croton**, Linn. (Euphorb.).

BRUNSVIGIA, Heist. Beschr. Brunsv. Braunschweig (1755), 3, cum ic. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 728.
albiflora, *Eckl. Topogr. Verz.* 7, nomen.—Afr. austr.
Burchelliana, Herb. Amaryll. t. 22. f. 2 = radulosa.
ciliaris, [Ker-Gawl. in] Bot. Reg. sub t. 192, 193 = Buphane disticha.
Cooperi, *Baker, in Saund. Ref. Bot.* t. 330.—Afr. austr.
coranica, *Sweet, Hort. Brit.* ed. I. 404.—Am. austr.
disticha, Sweet, l. c. = Buphane disticha.
falcata, [*Ker-Gawl. in*] *Bot. Mag.* t. 1443.—Afr. austr.
fulva, Steud. Nom. ed. II. i. 231 = Stenomesson incarnatum.
gigantea, *Heist. Beschr. Brunsv. Braunschweig*, 3, *cum ic. fide Schult. f. Syst.* vii. 844 (nobis latet).—Afr. austr.
glauca, Salisb. Gen. Pl. Fragm. 117 = Josephinae.
grandiflora, *Lindl. Bot. Reg.* t. 1335.—Afr. austr.
Josephinae, [*Ker-Gawl. in*] *Bot. Reg.* t. 192, 193.—Afr. austr.
laticoma, Sweet, Hort. Brit. ed. I. 404 = Nerine lucida.
lucida, Herb. App. 16 = Nerine lucida.
magnifica, Linden & Rodigas, in Illustr. Hortic. xxxii. (1885) t. 552 = Crinum Forbesianum.
marginata, Ait. Hort. Kew. ed. II. ii. 230 = Nerine marginata.
minor, *Lindl. Bot. Reg.* t. 954.—Afr. austr.
multiflora, Ait. Hort. Kew. ed. II. ii. 230 = gigantea.
orientalis, Ait. ex Eckl. Topog. Verz. 7 = gigantea.
Radula, *Ait. Hort. Kew.* ed. II. ii. 230.—Afr. austr.
radulosa, *Herb. Amaryll.* 281. t. 22. f. 2.—Afr. austr.
rubricaulis, M. Roem. Syn. Ensat. 57 = gigantea.
Slateriana, *Benth. & Hook. f. Gen.* iii. 727.—Afr. austr.
striata, *Ait. Hort. Kew.* ed. II. ii. 231.—Afr. austr.
toxicaria, [Ker-Gawl. in] Bot. Reg. t. 567 = Buphane disticha.
vaginata, Schult. f. Syst. vii. 852 = Hessea vaginata.

BRUSCHIA, Bertol. in Mem. Acad. Sc. Bolog. viii. (1857) 238 (Misc. Bot. xviii. 16. t. 2) = **Nyctanthes**, Linn. (Oleac.).
macrocarpa, Bertol. ll. cc. 238; 17 = N. Arbor-tristis.

BRUXANELIA, Dennst. Schluess. Hort. Malab. 31 (1818). *RUBIACEAE* ? (Gen. dub.).
indica, *Dennst. l. c.*—Ind. or.

BRYA, Vell. Fl. Flum. 146 (1825); iv. t. 1 (1827) = **Hirtella**, Linn. (Rosac.).
racemosa, Vell. ll. cc. 147; iv. t. 1.—Bras.

BRYA, P. Br. Hist. Jamaic. 299. t. 31 (1756). *LEGUMINOSAE*, Benth. & Hook. f. i. 514.
Ebenus, *DC. Prod.* ii. 421.—Ind. occ.
leonensis, *Lodd. ex Loud. Hort. Brit.* 285, nomen.—Afr. trop.
nicaraguensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 13.—Nicaragua.
parviflora, *DC. Prod.* ii. 421.—Bras.

BRYANTEA, Rafin. Sylva Tellur. 165 (1838) = **Litsea**, Lam. (Laurin.).
dealbata, *Rafin. l. c.* = L. dealbata.

BRYANTHUS, S. G. Gmel. Fl. Sibir. iv. 133. t. 57. f. 3 (1769). *ERICACEAE*, Benth. & Hook. f. ii. 595.
aleuticus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 368.—As. bor. or.; Am. bor. occ.
Breweri, *A. Gray, l. c.* 367.—Calif.
empetriformis, *A. Gray, l. c.*—Am. bor. occ.
erectus ×, Lindl. & Paxt. Flow. Gard. i. (1850–51) 97.
glanduliflorus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 368.—Am. bor. occ.
Gmelini, *D. Don, in Edinb. N. Phil. Journ.* xvii. (July 1834) 160.—Sibir.; Kamtsch.
intermedius, A. Gray, in Proc. Am. Acad. vii. (1868) 367 = empetriformis.
Stelleri, D. Don, in Edinb. N. Phil. Journ. xvii. (July 1834) 160 = Cassiope Stelleriana.
taxifolius, *A. Gray, in Proc. Am. Acad.* vii. (1868) 368.—Reg. bor. frigid.

BRYANTIA, Webb, ex Gaudich. Bot. Voy. Bonite, t. 20 (1844?); Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 287. t. 15 = **Pandanus**, Linn.
butyrophora, Webb, l. c. = P. butyrophorus.
oblonga, Brongn. in Ann. Sc. Nat. Sér. VI. i. (1875) 288; in Comptes-Rendus, lxxx. (1875) 1196 = P. oblongus.
viscida, Brongn. ll. cc. 287; 1196 = P. viscidus.

BRYLKINIA, F. Schmidt, Reisen Amurl. 199. t. 8 (1868). *GRAMINEAE*, Benth. & Hook. f. iii. 1192.
caudata, *F. Schmidt, l. c.*—Japon; Ins. Sachal.

BRYOBIUM, Lindl. Introd. Nat. Syst. ed. II. 446 (1836) = **Eria**, Lindl. (Orchid.).
pubescens, Lindl. Bot. Reg. (1838) Misc. 79 = E. retusa.

BRYOCARPUM, Hook. f. & Thoms. in Hook. Kew Journ. ix. (1857) 200. t. 5. *PRIMULACEAE*, Benth. & Hook. f. ii. 634.
himalaicum, *Hook. f. & Thoms. l. c.*—Reg. Himal.

BRYOCLES, Salisb. in Trans. Hort. Soc. i. (1812) 335 = **Funkia**, Spreng. (Liliac.).
ventricosa, Salisb. l. c. = F. ovata.

BRYODES, Benth. in DC. Prod. x. 433 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 957.
micrantha, *Benth. l. c.*—Ins. Maurit.

BRYOMORPHA, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 172 = **Thylacospermum**, Fenzl (Caryophyll.).
rupifraga, Kar. & Kir. l. c. = T. rupifragum.

BRYOMORPHE, Harv. Thes. Cap. ii. 33. t. 151 (1863). *COMPOSITAE*, Benth. & Hook. f. ii. 324.
Zeyheri, *Harv. l. c.*—Afr. austr.

BRYONIA, Linn. Syst. ed. I (1735). *CUCURBITACEAE*, Benth. & Hook. f. i. 829.
abyssinica, Goualt, in Rev. Hortic. (1853) 61 = Maximowiczia Lindheimeri.
abyssinica, Lam. Encyc. i. 497 = Coccinia abyssinica.
acerifolia, D. Dietr. Syn. Pl. v. 367 = Coccinia indica.
acuta, Desf. Fl. Atlant. ii. 360 = dioica.
acutangula, Thunb. Prod. Pl. Cap. 13 = (Senecionis sp.).
affinis, Endl. Prod. Fl. Norfolk. 125 = Bryonopsis affinis.

BRYONIA:—
africana, E. Mey. Zwei Pfl. Docum. 169 = Kedrostis grossulariaefolia.
africana, Linn. Sp. Pl. 1013 = Kedrostis africana.
africana, Thunb. Prod. Pl. Cap. 13 = Kedrostis nana.
agrestis, Raeusch. Nom. Bot. ed. II. 283 = Blastania Garcini.
alba, Bull. Herb. t. 55 = dioica.
alba, *Linn. Sp. Pl.* 1012.—Europ.; Reg. Caucas.; Persia.
alceaefolia, Willd. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 223 = Coccinia indica.
althaeoides, Ser. in DC. Prod. iii. 306 = Melothria maderaspatana.
americana, Descourt. Fl. Antill. ii. 281. t. 136 = Melothria pervaga.
americana, Lam. Encyc. i. 498 = Cayaponia americana.
amplexicaulis, Lam. l. c. 496 = Melothria amplexicaulis.
angulata, Thunb. Prod. Pl. Cap. 13 = Melothria punctata.
arguta, *Zipp. ex Span. in Linnaea*, xv. (1841) 206.—Malaya.
aspera, Stev. in Cat. Hort. Gorenk. (1812) 32 = alba.
asperifolia, Presl, Bot. Bemerk. 72 = Cayaponia americana.
attenuata, Hook. & Arn. Bot. Beech. Voy. Suppl. 424 = Cayaponia attenuata.
barbata, Buch.-Ham. ex Cogn. in DC. Monog. Phan. iii. 530 = Coccinia indica.
biflora, Buch.-Ham. in Wall. Cat. n. 6707 = Melothria odorata.
Blumei, Ser. in DC. Prod. iii. 305 = Melothria Rumphiana.
bonariensis, Mill. Gard. Dict. ed. VIII. n. 6 = Cayaponia ficifolia.
Boykinii, Torr. & Gray, Fl. N. Am. i. 540 = Cayaponia Boykinii.
Cabocla, Vell. Fl. Flum. x. t. 88 = Cayaponia globosa.
callosa, Heyne, ex Wall. Cat. sub n. 6708 = Melothria maderaspatana.
callosa, Rottl. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 210; Wall. Cat. n. 6710 = Cucumis trigonus.
capillacea, Schum. & Thonn. Beskr. Guin. Pl. 430 = Melothria triangularis.
cheirophylla, Wall. Cat. n. 6715 A = Luffa aegyptiaca.
ciliata, Moench, Meth. 653 = Kedrostis africana.
cissoides, Wall. Cat. n. 6698 = Melothria perpusilla.
cochinchinensis, Lour. Fl. Cochinch. 595 = Gymnopetalum cochinchinense.
collosa, Rottl. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 210 = *sphalm. vide* callosa.
convolvulifolia, Schlecht. in Linnaea, xxvi. (1853–55) 640 = Melothria guadalupensis.
convolvuloides, A. Rich. Tent. Fl. Abyss. i. 288 = Coccinia adoënsis.
cordata, Thunb. Fl. Cap. 149 = Melothria cordata.
cordatifolia, Goday Torres, in Patriota, iii. n. 37 (1814) 71 = Cayaponia Tayuya.
cordifolia, Walp. Rep. ii. 199 = praec.
cordifolia, Linn. Sp. Pl. 1012 = Coccinia indica.
cordifolia, Russ. ex Steud. Nom. ed. II. i. 232 = Melothria maderaspatana.
cretica, *Linn. Sp. Pl.* 1013.—Graccia et Ins. Creta.
cucumeroides, Ser. in DC. Prod. iii. 308 = Trichosanthes cucumeroides.
deltoidea, Arn. Pugill. 19; in Nov. Act. Nat. Cur. xviii. (1836) 337 = Melothria zeylanica.
deltoidea, Schum. & Thonn. Beskr. Guin. Pl. 429 = Melothria deltoidea.
dentata, E. Mey. Zwei Pfl. Docum. 169 = Melothria punctata.
digitata, Thunb. Prod. Pl. Cap. 13 = Kedrostis digitata.
digyna, Pomel, Nouv. Mat. Fl. Atl. 155 = dioica.
dioica, Bieb. Fl. Taur. Cauc. iii. 625 = alba.
dioica, Boj. Hort. Maurit. 148 = Bryonopsis laciniosa.
dioica, *Jacq. Fl. Austr.* ii. 59. t. 199 (err. typ. disica).—Europ.; Reg. Mediterr.
dissecta, Thunb. Prod. Pl. Cap. 13 = Kedrostis africana.
divisa, *Willd. ex Steud. Nom.* ed. II. i. 232, nomen.—Hab.?
epigaea, Blume, Bijdr. 925 = Melothria Rumphiana.
epigaea, Rottl. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 223 = Corallocarpus epigaeus.

BRYONIA:—
ficifolia, Descourt. Fl. Antill. vi. 39. t. 391 = Cayaponia americana.
ficifolia, Lam. Encyc. i. 498 = Cayaponia ficifolia.
filicaulis, Wall. Cat. n. 6713 = Kedrostis rostrata.
filifolia, Spreng. Syst. iii. 16 = Cayaponia ficifolia.
filiformis, Roxb. Hort. Beng. 70; Fl. Ind. iii. 727 = Melothria heterophylla?
fimbristipula, Fenzl, ex Hook. f. in Oliver, Fl. Trop. Afr. ii. 558 = Blastania fimbristipula.
fluminensis, Vell. Fl. Flum. x. t. 87 = Cayaponia fluminensis.
foetidissima, Schum. & Thonn. Beskr. Guin. Pl. 428 = Kedrostis foetidissima.
Garcini, Willd. Sp. Pl. iv. 623 = Blastania Garcini.
geminata, Blume, Bijdr. 924 = Melothria indica.
glabra, Roxb. Hort. Beng. 104 = Corallocarpus epigaeus.
glandulosa, *Gill. in Hook. Bot. Misc.* iii. (1833) 324.—Chili.
glandulosa, Poepp. & Endl. Nov. Gen. et Sp. ii. 56. t. 175 = Cayaponia glandulosa.
gracilis, Wall. Cat. n. 6714 = Melothria maderaspatana.
grandis, Linn. Mant. i. 126 = Coccinia indica.
grandis, Wall. Cat. n. 6700 K = Gymnopetalum cochinchinense.
grandis, Wall. l. c. L = Momordica dioica.
grossularifolia, E. Mey. Zwei Pfl. Docum. 169 = Kedrostis grossulariaefolia.
guadalupensis, Spreng. Syst. iii. 15 = Melothria pervaga.
guineensis, G. Don, Gen. Syst. iii. 32 (sp. dub.).—Afr. trop.
hastata, Lour. Fl. Cochinch. ii. 594 = Melothria heterophylla.
hederaefolia, Jacq. Fragm. 73. n. 230. t. 113 = verrucosa.
heterophylla, Blume, Bijdr. 925 = Melothria Rumphiana.
heterophylla, Raeusch. Nom. ed. III. 282; Steud. Nom. ed. I. 123 = Melothria heterophylla.
heterophylla, Wall. Cat. n. 6704 = Melothria marginata?
hispida, Salisb. Prod. 158 = Melothria maderaspatana.
Hookeriana, Wight & Arn. Prod. 345 = Melothria perpusilla.
japonica, Thunb. Fl. Jap. 325 = Melothria japonica.
jatrophaefolia, A. Rich. Tent. Fl. Abyss. i. 289 = Coccinia jatrophaefolia.
Johnstoni, Cuzent, ex Seem. in Journ. Bot. ii. (1864) 47 = Melothria Grayana.
laciniosa, Linn. Sp. Pl. 1013 = Bryonopsis laciniosa.
laevis, Thunb. Prod. Pl. Cap. 13 = Melothria velutina.
lagenaria, E. Mey. Zwei Pfl. Docum. 169 = Coccinia sessilifolia.
latebrosa, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 384 = Cayaponia latebrosa.
leiosperma, Wight & Arn. Prod. 345 = Melothria leiosperma.
leucocarpa, Blume, Bijdr. 924 = Melothria odorata.
Lhotzkyana, Cogn. in Mart. Fl. Bras. vi. iv. 83 = Cayaponia Lhotzkyana.
lutea, Bast. ex Ser. in DC. Prod. iii. 307 = dioica.
macrophylla, *Kotschy, ex Boiss. Fl. Orient.* ii. 761.—Armenia.
macrophylla, *Ser. in DC. Prod.* iii. 308 (sp. dub.).—Hab.?
maderaspatana, Berg. Desc. Pl. Cap. 351 = Melothria punctata.
maderaspatana, Lam. Encyc. i. 496 = Melothria maderaspatana.
marginata, Blume, Bijdr. 924 = Melothria Rumphiana.
maysorensis, Wall. Cat. n. 6702 = Melothria mucronata.
maysorensis, Wight, Ic. iii. t. 758 = Melothria perpusilla.
Melothria, Crantz, Inst. i. 173 = Melothria pendula.
micrantha, Boiss. Diagn. Ser. I. x. 7 = syriaca.
micrantha, Hochst. ex Cogn. in DC. Monog. Phan. iii. 623 = Melothria maderaspatana.
micropoda, E. Mey. Zwei Pfl. Docum. 169 = Melothria maderaspatana.
Moimoi, Ser. in DC. Prod. iii. 305 = Coccinea Moghadd.
mucronata, Blume, Bijdr. 923 = Melothria mucronata.
multifida, E. Mey. Zwei Pfl. Docum. 169 = Kedrostis africana.
multiflora, *Boiss. & Heldr. Diagn.* Ser. I. x. 8.—Orient; Ind. bor. occ.

BRYONIA:—
mysorensis, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 623 = Melothria leiosperma.
mysorensis, Walp. Rep. ii. 198 = Melothria mucronata.
mysorensis, Wight & Arn. Prod. i. 345 = Melothria perpusilla.
nana, Lam. Encyc. i. 497 = Kedrostis nana.
nepaulensis, Ser. in DC. Prod. iii. 307 = Melothria heterophylla.
nigra, Gilib. Fl. Lituan. i. 60 = alba.
nitida, Link, Enum. Hort. Berol. ii. 404 = dioica.
obtusa, A. Rich. Tent. Fl. Abyss. i. 288 = Mukia scabrella.
obtusiloba, E. Mey. Zwei Pfl. Docum. 169 = Melothria obtusiloba.
odorata, Buch.-Ham. ex Wall. Cat. n. 6706 B, C = Melothria odorata.
odorata, Wall. Cat. n. 6706 D = Melothria Wallichii.
oxyphylla, Wall. Cat. n. 6698 = Melothria perpusilla.
palmata, Linn. Sp. Pl. 1012 = Bryonopsis laciniosa.
palmata, Wall. Cat. n. 6711 A, B, C = Coccinia indica.
palmata, Wall. l. c. D = Corallocarpus epigaeus.
Pancheri, Naud. in Ann. Sc. Nat. Sér. IV. ix. (1858) 398 = Bryonopsis affinis.
pectinata, E. Mey. Zwei Pfl. Docum. 169 = Trochomeria pectinata.
pedata, Hassk. in Flora, xxv. (1842) Beibl. 42 = Bryonopsis laciniosa.
pedunculosa, Ser. in DC. Prod. iii. 306 = Herpetospermum caudigerum
perpusilla, Blume, Bijdr. 926 = Melothria perpusilla.
Perrottetiana, Ser. in DC. Prod. iii. 304 = Kedrostis foetidissima.
pilosa, Vell. Fl. Flum. x. t. 86 = Cayaponia diffusa.
pilosa, Roxb. Hort. Beng. 104; Fl. Ind. iii. 726 = Kedrostis rostrata.
pinnatifida, Vell. Fl. Flum. x. t. 90 = Cayaponia diversifolia.
pinnatifida, Burch. Trav. i. 547 = Kedrostis africana.
pinnatiloba, M. Roem. Syn. fasc. ii. Pepon. 40 = Cayaponia diversifolia.
prostrata, D. Dietr. Syn. Pl. v. 363 = Kedrostis rostrata.
polymorpha, *Jensch. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 352, nomen.—Hab. ?
pubescens, *Poir. Encyc. Suppl.* i. 731 (sp. dub.).—Oriens.
punctata, Buch.-Ham. ex Wall. Cat. sub n. 6708 = Melothria maderaspatana.
punctata, Thunb. Prod. Pl. Cap. 13 = Melothria punctata.
quinquefolia, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 48 = Bryonopsis laciniosa.
quinqueloba, Thunb. Prod. Pl. Cap. 13 = Coccinia quinqueloba.
racemosa, *Mill. Gard. Dict.* ed. VIII. n. 4.—Ind. occ.
racemosa, Poepp. & Endl. Nov. Gen. et Sp. ii. 56 = Cayaponia Poeppigii.
racemosa, Sw. Prod. Veg. Ind. Occ. = Cayaponia racemosa.
reniformis, Roxb. ex Wight & Arn. Prod. 344 = Blastania Garcini.
repanda, Blume, Bijdr. 923 = Melothria punctata.
Rheedei, Blume, l. c. 925 = Melothria heterophylla.
rostrata, Rottl. in Ges. Naturf. Freunde Neue Schr. iv. (1803) 212 = Kedrostis rostrata.
Rottleri, Spreng. Syst. iii. 15 = Melothria maderaspatana.
ruderalis, Salisb. Prod. 158 = dioica.
ruderalis, *Zipp. ex Span. in Linnaea*, xv. (1841) 206, nomen.—Malaya.
sagittata, Blume, Bijdr. 925 = Melothria heterophylla.
scabra, Rottl. ex Wight & Arn. Prod. i. 345 = Melothria maderaspatana.
scabra, Thunb. Prod. Pl. Cap. 13 = Melothria punctata.
scabrata, Blume, Bijdr. 923 = Melothria perpusilla.
scabrella, Linn. f. Suppl. 424 = Melothria maderaspatana.
scabrella, Sieber, ex Cogn. in DC. Monog. Phan. iii. 786 = Cayaponia americana.
scabrella, Wall. Cat. n. 6708 = Melothria leiosperma.
scarlatina, Dum. Fl. Belg. 54 = dioica.
scrobiculata, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 287 = Melothria scrobiculata.
septemlobata, *Vell. Fl. Flum.* x. t. 85.—Bras.
sicula, Guss. Fl. Sic. Syn. ii. 621 = dioica.
sinuata, Wall. Cat. n. 6711 D = Corallocarpus epigaeus.
sinuosa, Wall. Cat. n. 6716 = Coccinia indica et Melothria heterophylla.

BRYONIA:—
Solena, Mirb. ex Steud. Nom. ed. II. i. 233 = Melothria heterophylla.
stipulacea, Willd. Sp. Pl. iv. 620 = Blastania Garcini.
subpinnata, Stend. Nom. ed. II. i. 232 = Cayaponia diversifolia.
syriaca, *Boiss. Diagn.* Ser. II. ii. 59.—Syria; Palaestin.
Tayuya, Mart. in Linnaea, xii. (1838) Litt. 86 = Cayaponia Martiana.
Tayuya, Vell. Fl. Flum. x. t. 89 = Cayaponia Tayuya.
Teedonda, Roxb. ex Willd. Sp. Pl. iv. 618 = Melothria heterophylla.
tenella, Roxb. Hort. Beng. 70; Fl. Ind. iii. 725 = Melothria indica.
tenuifolia, Gill. in Hook. Bot. Misc. iii. (1833) 323 = Abobra tenuifolia.
tenuifolia, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 56.—Chili.
tenuis, Klotzsch, in Peters, Reise Mossamb. Bot. 150 = Bryonopsis laciniosa.
ternata, Vell. Fl. Flum. x. t. 91 = Cayaponia ternata.
Thunbergiana, Dietr. ex Steud. Nom. ed. I. 123 = Melothria cordata.
Tinei, Huet, ex Cogn. in DC. Monog. Phan. iii. 471 = dioica.
tomentosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 8.—Malaya.
tricuspidata, Leschen. ex Steud. Nom. ed. II. i. 275 = Passiflora Leschenaultii.
triflora, Wall. Cat. n. 6707 = Melothria leucocarpa.
triloba, Lour. Fl. Cochinch. ii. 595 = Blastania Garcini.
triloba, Thunb. Prod. Pl. Cap. 13 = Kedrostis nana.
trilobata, Ser. in DC. Prod. iii. 307 = Kedrostis nana.
tubiflora, Wight & Arn. Prod. 347 = Gymnopetalum Wightii.
umbellata, Klein, ex Willd. Sp. Pl. iv. 618 = Melothria heterophylla.
umbellata, Wall. Cat. n. 6705 D = Trichosanthes cucumerina.
umbellata, Wall. l. c. K, L = Melothria amplexicaulis.
variegata, Mill. Gard. Dict. ed. VIII. n. 5 = Bryonopsis laciniosa.
verrucosa, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 385.—Ins. Canar.
vitifolia, *Hort. Belg. ex Heynh. Nom.* ii. 79, nomen.—Hab. ?
vittata, *Zipp. ex Span. in Linnaea*, xv. (1841) 206, nomen.—Malaya.
vulgaris, Gueldenst. ex Ledeb. Fl. Ross. ii. 1401 = alba.
Wallichiana, Ser. ex DC. Prod. iii. 309 = Melothria heterophylla.
Wightiana, Wall. Cat. n. 6703 = Melothria leiosperma.

BRYONOPSIS, Arn. in Hook. Journ. Bot. iii. (1841) 274. *CUCURBITACEAE*, [Benth. & Hook. f. i. 274].
affinis, *Cogn. in DC. Monog. Phan.* iii. 479.—Ins. Norfolk.
Bonnettii, Miq. Fl. Ind. Bat. i. 1. 657 = Cerasiocarpum zeylanicum.
courtallensis, Arn. in Hook. Journ. Bot. iii. (1841) 274 = laciniosa.
erythrocarpa, Naud. in Ann. Sc. Nat. Sér. IV. xviii. (1862) 194 = laciniosa.
laciniosa, *Naud. l. c.* Sér. V. vi. (1866) 30.—As. Afr. et Austral. trop.
leucocarpa, Miq. Fl. Ind. Bat. i. 1. 657 = Melothria odorata.
Pancheri, Naud. in Ann. Sc. Nat. Sér. V. vi. (1866) 30 = affinis.
pedata, Hassk. Cat. Pl. Hort. Bog. Alt. 189 = laciniosa.

BRYOPHTHALMUM, E. Mey. Preuss. Pflanzeng. 101 (1839) = **Moneses**, Salisb. (Ericac.).
uniflorum, E. Mey. l. c. = M. uniflora.

BRYOPHYLLUM, Salisb. Parad. Lond. t. 3 (1805). *CRASSULACEAE*, Benth. & Hook. f. i. 658.
PHYSOCALYCIUM, Vest, in Flora, iii. (1820) 409.
CRASSUVIA, Comm. ex Lam. Encyc. ii. 141 (1786).
calycinum, *Salisb. l. c.*—Mcxic.
cochleatum, Lem. Illustr. Hortic. vi. (1859) Misc. 100 = Kalanchoë cochleatum.
crenatum, *Baker, in Journ. Linn. Soc.* xx. (1884) 139. —Madag.
germinans, Blanco, Fl. Filip. ed. II. 220 = calycinum.
pinnatum, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 52 = calycinum.
proliferum. *Bowie, ex Hook. Bot. Mag.* t. 5147.—Afr. austr.

BRYOPHYLLUM:—
serratum, Blanco, Fl. Filip. ed. II. 220 = Kalanchoë laciniata.
triangulare, Blanco, l. c. 221 = Kalanchoë spathulata.
tubiflorum, *Harv. in Harv. & Sond. Fl. Cap.* ii. 380. —Afr. austr.

BUBALINA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 86 = **Burchellia**, R. Br. (Rubiac.).

BUBANIA, Girard, in Mém. Acad. Montp. (Sect. Sc.) i. (1848) 182 = **Limoniastrum**, Moench (Plumbagin.).
Feei, Girard, l. c. 185 = L. Guyonianum.
monopetala, Girard, l. c. 183 = L. articulatum.

BUBON, Linn. Gen. ed. I. 339 (1737) = **Seseli**, Linn. (Umbellif.).
aphyllus, Cham. & Schlecht. in Linnaea, i. (1826) 389 (= *Deverra tortuosa*).—Aegypt.
buchtormensis, Spreng. Pugill. ii. 55 = S. buchtormense.
capensis, Sond. in Harv. & Sond. Fl. Cap. iii. 561.—Afr. austr.
cuneifolius, Spreng, Syst. i. 900 = S. cuneifolium.
dichotomus, Link, Enum. Hort. Berol. i. (1821) 278 = S. dichotomum.
eriocephalus, Pall. ex Spreng. Syst. i. 900 = S. Lessingianum.
Galbanum, Hort. ex Steud. Nom. ed. II. i. 233 (= *Melanoselinum decipiens*).—Madera.
Galbanum, Linn. Sp. Pl. 253 = Peucedanum Galbanum.
garganicus, Tenore, Fl. Nap. i. 123 = Athamanta sicula.
glaucus, Spreng. Umb. Spec. 136 = S. petraeum.
globosus, Stev. ex Ledeb. Fl. Ross. ii. 274 = S. rigidum.
gummifer, Linn. Sp. Pl. 254 = S. gummiferum.
humilis, Salisb. Prod. Stirp. 165 = Athamanta macedonica.
hypoleucus, Meissn. in Hook. Lond. Journ. Bot. ii. (1840) 536.—Afr. austr.
laevigatus, Ait. Hort. Kew. ed. II. v. i. 352.—Afr. austr.
latifolius, Salisb. Prod. 165 = *laevigatus*.
Libanotis, Dum. Fl. Belg. 79 = S. Libanotis.
macedonicus, Linn. Sp. Pl. 253 = Athamanta macedonica.
montanus, Sond. in Harv. & Sond. Fl. Cap. iii. 560.—Afr. austr.
multiflorus, Moench, Meth. 95 = Athamanta macedonica.
persicus, Pall. ex Ledeb. Fl. Ross. ii. 329 = Zosima orientalis.
peucedanifolius, Spreng. Syst. i. 900 = S. rigidum.
pimpinellifolius, Eckl. & Zeyh. Enum. 353.—Afr. austr.
prolifer, Burm. f. Fl. Cap. Prod. 7 = Lichtensteinia pyrethrifolia.
pyrenaicus, Dum. Fl. Belg. 79 = S. vulgare.
rigidior, Linn. Sp. Pl. 254 = Ferula communis.
rigidus, Spreng. Pugill. ii. 53 = S. gummiferum.
rigidus, Spreng. Syst. i. 900 = S. rigidum.
siculus, Spreng. in Roem. & Schult. Syst. vi. 499 = S. Bocconi.
tenuifolius, Salisb. Prod. 165 = S. gummiferum.
tenuifolius, Sond. in Harv. & Sond. Fl. Cap. ii. 560.—Afr. austr.
tortuosus, Desf. Fl. Atlant. i. 257 (= *Deverra tortuosa*). —Afr. bor.

BUBONIUM, Hill, Veg. Syst. ii. 74 (1761); Hort. Kew. 13 (1769) (Compos.).
aquaticum, Hill, ll. cc. = Odontospermum aquaticum.
arborescens, Hill, Hort. Kew. 13 = Borrichia arborescens.
frutescens, Hill, Veg. Syst. ii. 74; Hort. Kew. 13 = Borrichia frutescens.
maritimum, Hill, Hort. Kew. 13 = Odontospermum maritimum.
spinosum, Hill, Veg. Syst. ii. 74 = Pallenis spinosa.

BUBROMA, Schreb. Gen. 513 (1791) = **Guazuma**, Plum. (Sterculiac.).
crinitum, Steud. Nom. ed. II. i. 233 = G. crinita.
grandiflorum, Willd. ex Steud. l. c. = G. grandiflora.
Guazuma, Willd. Sp. Pl. iii. 1423 = G. ulmifolia.
Invira, Willd. Enum. Hort. Berol. 806 = G. ulmifolia.
polybotryum, Willd. l. c. = G. polybotrya.
tomentosum, Spreng. Syst. iii. 332 = G. ulmifolia.

BUCAFER, Adans. Fam. ii. 469 (1763) = **Ruppia**, Linn. (Naiad.).

BUCANION, Stev. in Bull. Soc. Nat. Mosc. (1851) i. 568 = **Heliotropium**, Tourn. (Boragin.).
Szovitsii, Stev. l. c. = H. Szovitsii.

BUCCAFERREA, Bubani, in Nuovo Giorn. Bot. Ital. v. (1873) 316 = **Potamogeton**, Tourn. (Naiad.).
Hornemanni, Bubani, l. c. = P. Hornemanni.
obtusifolia, Bubani, l. c. 317 = P. obtusifolius.

BUCCAFERREA, Mich. ex Petagna, Inst. Bot. v. 1826 (1787) = **Ruppia**, Linn. (Naiad.).
cirrhosa, Petagna, l. c. = R. maritima.

BUCCO, Wendl. Coll. 13, tt. 2, 3, 13, 28, 77 (1808) = **Agathosma**, Willd. (Rutac.).
acuminata, Wendl. Coll. i. t. 28 = A. imbricata.
barbata, Roem. & Schult. Syst. v. 445 = Macrostylis lanceolata.
betulina, Roem. & Schult. l. c. 443 = Barosma betulina.
brevifolia, Roem. & Schult. l. c. 446 = A. erecta.
Cerefolium, Roem. & Schult. l. c. 439 = A. Cerefolium.
crenata, Roem. & Schult. l. c. 444 = Barosma crenata.
cuspidata, Wendl. Coll. iii. 17. t. 81 = A. cuspidata.
erecta, Wendl. l. c. i. 17. t. 3 = A. erecta.
hamata, Wendl. ex Bartl. & Wendl. f. Diosmeae, 114 = Barosma dioica.
hirta, Roem. & Schult. Syst. v. 446 = A. hirta.
hispida, Roem. & Schult. l. c. = A. hispida.
imbricata, Wendl. Coll. i. 33. t. 9 = A. imbricata.
Lamarckiana, Roem. & Schult. Syst. v. 447 = A. virgata.
linifolia, Lichtst. in Roem. & Schult. l. c. 448 = A. linifolia.
obtusa, Wendl. Coll. i. 45 = A. obtusa.
obtusata, Wendl. l. c. iii. 7. t. 76 = A. imbricata.
ovata, Roem. & Schult. Syst. v. 442 = Barosma ovata.
prolifera, Wendl. Coll. iii. t. 77 = A. prolifera.
pulchella, Roem. & Schult. Syst. v. 442 = Barosma pulchella.
rubra, Roem. & Schult. l. c. 441 = Diosma vulgaris.
scabra, Wendl. ex Bartl. & Wendl. f. Diosmeae, 156 = A. ciliata.
serpyllacea, Roem. & Schult. Syst. v. 447 = A. serpyllacea.
spicata, Roem. & Schult. l. c. 448 = A. spicata.
tetragona, Roem. & Schult. l. c. 444 = Acmadenia tetragona.
Ventenatiana, Roem. & Schult. l. c. 442 = A. Ventenatiana.
vestita, Roem. & Schult. l. c. 447 = A. imbricata.
villosa, Wendl. Coll. i. 14. t. 2 = A. villosa.

BUCCULINA, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 209 = **Holothrix**, Rich. (Orchid.).
aspera, Lindl. l. c. = H. aspera.

BUCEPHALANDRA, Schott, Gen. Aroid. t. 56 (1858). *AROIDEAE*, Benth. & Hook. f. iii. 984.
Mottleyana, *Schott, l. c.*—Borneo.

BUCEPHALON, [Plum.] Linn. Gen. ed. I. 375 (1737) = **Trophis**, Linn. (Urtic.).
racemosum, Linn. Sp. Pl. 1190 = T. americana.

BUCERAS, Hall. ex All. Fl. Pedem. i. 313 (1785) = **Trigonella**, Linn. (Legumin.).
corniculata, All. Fl. Pedem. i. 313 = T. corniculata.
elliptica, Moench, Meth. 143 = T. monspeliaca.
Foenum-graecum, All. Fl. Pedem. i. 313 = T. Foenum-graecum.
grandiflora, Moench, Meth. 143 = T. corniculata.
monspeliaca, All. Fl. Pedem. i. 313 = T. monspeliaca.
mutica, Moench, Meth. 142 = T. polycerata.
odoratissima, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 383 = T. Foenum-graecum.
platycarpos, Moench, Meth. Suppl. 51 = Medicago platycarpos.
polyceration, All. Fl. Pedem. i. 313 = M. polycerata.
spinosa, Moench, Meth. 143 = T. spinosa.
torosa, Moench, Meth. Suppl. 51 = T. striata.
tortuosa, Steud. Nom. ed. II. i. 233 = T. striata.

BUCERAS, P. Br. Hist. Jamaic. t. 23 (1756) = **Terminalia**, Linn. (Combretac.).
Bucida, Crantz, Inst. i. 133 = T. Catappa.

BUCEROSIA, Endl. Gen. 598 (1838) = **Boucerosia**, Wight & Arn. (Asclep.).

BUCETUM, Parnell, Grasses of Scotl. 104-109 (1842) = **Festuca**, Tourn. (Gramin.).
elatius, Parnell, l. c. 107 = F. elatior.
giganteum, Parnell, l. c. 108 = F. gigantea.
loliaceum, Parnell, l. c. 104 = F. loliacea.
pratense, Parnell, l. c. 105 = F. elatior.

BUCHANANIA, Sm. Exot. Bot. ii. tab. solum, 115 (1805) = **Colebrookea**, Sm. (Labiat.).
oppositifolia, Sm. l. c. = C. oppositifolia.

BUCHANANIA, Spreng. in Schrad. Journ. (1800) II. 234; Roxb. Pl. Corom. iii. 79. t. 282 (1819).
ANACARDIACEAE, Benth. & Hook. f. i. 421.
CAMBESSEDEA, Kunth, in Ann. Sc. Nat. ii. (1824) 366.
CONIOGETON, Blume, Bijdr. 1156 (1826).
LANZANA, Stokes, Bot. Mat. Med. ii. 543 (1812).
LAUNZAN, Buch.-Ham. in As. Res. v. (1799) 123.
LAUNZEA, Endl. Ench. 599 (1841).
LUNDIA, Puer. ex DC. Prod. ii. 64, in syn. (1825).
acuminata, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 472.—Burma; Malaya.
acuminata, Wall. Cat. n. 981 = lanceolata.
amboinensis, *Miq. in Ann. Mus. Lugd. Bat.* iv. 117. —Amboina.
angustifolia, Benth. Fl. Austral. i. 490 = Muelleri.
angustifolia, Roxb. Hort. Beng. 32; Fl. Ind. ii. 386 = B. Lanzan.
arborescens, Blume, Mus. Bot. Lugd. Bat. i. 183 = florida.
arborescens, F. Muell. Fragm. vii. 23 = Muelleri.
auriculata, Blume, l.c. 185 = Campnosperma auriculata.
bancana, Miq. Fl. Ind. Bat. Suppl. i. 523 = florida.
ferruginea, *Engl. in DC. Monog. Phan.* iv. 194.—Ind. or.
florida, A. Gray, in Bot. U. St. Expl. Exped. i. 366 = vitiensis.
florida, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 481.—As. trop.; Malaya.
glabra, *Wall. Cat.* n. 984.—Burma.
Halmaheirae, *Miq. in Ann. Mus. Lugd. Bat.* iv. 117. —Malaya.
insignis, *Blume, Mus. Bot. Lugd. Bat.* i. 184.—Borneo.
intermedia, Wight, Ic. i. t. 81 = florida.
lanceaefolia, Roxb. Hort. Beng. 32 = lancifolia.
lanceolata, *Wight, Ic.* i. t. 237.—Ind. or.
lancifolia, *Roxb. Fl. Ind.* ii. 386.—Ind. or.; Burma.
Lanzan, *Spreng. in Schrad. Journ.* (1800) II. 234.—Ind. or.
latifolia, *Roxb. Hort. Beng.* 32; *Fl. Ind.* ii. 385.—Ind. or.; Burma.
laxiflora, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 304.—Burma.
longifolia, Blume, Mus. Bot. Lugd. Bat. i. 184 = florida.
longifolia, *Span. in Linnaea*, xv. (1841) 188.—Ins. Timor.
lucida, Blume, Mus. Bot. Lugd. Bat. i. 184 = florida.
macrophylla, Blume, l. c. 185 = Campnosperma macrophylla.
mangoides, *F. Muell. Fragm.* vii. 23.—Ins. Pacif.
microphylla, *Engl. in DC. Monog. Phan.* iv. 185.—Ins. Philipp.
Muelleri, *Engl. l. c.* 190.—Austral.
nitida, *Engl. l. c.* 193.—Ins. Philipp.
obovata, *Engl. l. c.* 187.—Austral.
oxyphylla, *Miq. Fl. Ind. Bot. Suppl.* i. 522.—Sumatra.
oxyrhachis, Miq. l. c. 524 = Campnosperma oxyrhachis.
palembanica, Blume, Mus. Bot. Lugd. Bat. i. 186 = florida.
paniculata, Roxb. ex Steud. Nom. ed. II. i. 233 = Amoora cucullata.
petiolaris, Miq. Fl. Ind. Bat. i. II. 637 = florida.
platyneura, *Kurz, in Journ. As. Soc. Beng.* xlv. (1876) II. 125.—Ins. Nicobar.
polybotrya, Miq. Fl. Ind. Bat. i. II. 638 = florida.
racemiflora, Miq. Fl. Ind. Suppl. i. 523 = Campnosperma macrophylla.
reticulata, *Hance, in Journ. Bot.* xv. (1877) 332.—Cochinch.
scssilifolia, *Blume, Mus. Bot. Lugd. Bat.* i. 184.—Malaya.
siamensis, *Miq. in Ann. Mus. Lugd. Bat.* iv. 118.—Siam.
spicata, Roxb. ex Wall. Cat. n. 4895 = Amoora cucullata.
splendens, *Miq. Fl. Ind. Bat. Suppl.* i. 524.—Sumatra.
subobovata, Griff. Notul. iv. 413 = florida.

BUCHANANIA:—
vitiensis, *Engl. in DC. Monog. Phan.* iv. 186.—Ins. Fiji.
zeylanica, Blume, Mus. Bot. Lugd. Bat. i. 185 = Mangifera zeylanica.

BUCHAREA, Rafin. Fl. Tellur. iv. 84 (1836) = **Convolvulus**, Tourn.
atlantica, Rafin. l. c.—Afr. bor.
maderensis, Rafin. l. c.—Ins. Madera.

BUCHENAVIA, Eichl. in Flora, xlix. (1866) 164 = **Terminalia**, Linn. (Combret.).
capitata, Eichl. l. c. 165 = T. Hilariana.
macrophylla, Eichl. l. c. 166 = T. macrophylla.
megalophylla, Heurck & Muell. Arg. in Heurck, Obs. Bot. 211.—Am. trop.
ochroprumna, Eichl. in Flora, xlix. (1866) 165.—Am. trop.
oxycarpa, Eichl. l. c. 166 = T. oxycarpa.
punctata, Eichl. l. c. = T. punctata, *Spruce*.
reticulata, Eichl. l. c.—Am. trop.
suaveolens, Eichl. l. c. = T. suaveolens.
tomentosa, Eichl. l. c. = T. tomentosa, *Mart.*

BUCHENROEDERA, Eckl. & Zeyh. Enum. 194 (1836). *LEGUMINOSAE*, Benth. & Hook. f. i. 478.
COLOBOTUS, E. Mey. Comm. Pl. Afr. Austr. 156. (1836).
alpina, *Eckl. & Zeyh. Enum.* 195.—Afr. austr.
caerulescens, *Presl, Bot. Bemerk.* 47.—Afr. austr.
gracilis, *Eckl. & Zeyh. Enum.* 195.—Afr. austr.
holoscricea, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 581.—Afr. austr.
lanceolata, *Presl, Bot. Bemerk.* 47.—Afr. austr.
Meyeri, *Presl, l. c.*—Afr. austr.
multiflora, *Eckl. & Zeyh. Enum.* 195.—Afr. austr.
spicata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 93.—Afr. austr.
tenuifolia, *Eckl. & Zeyh. Enum.* 196.—Afr. austr.
teretifolia, Eckl. & Zeyh. l. e. = Aspalathus armata.
trichodes, *Presl, Bot. Bemerk.* 47.—Afr. austr.
umbellata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 93.—Afr. austr.
viminea, *Presl, Bot. Bemerk.* 47.—Afr. austr.

BUCHERIA, Heynh. Nom. ii. 80 (1846) = **Thryptomene**, Endl. (Myrtac.).
saxicola, Heynh. l. c. = T. saxicola.

BUCHIA, D. Dietr. Syn. Pl. iii. 370 (1843) = **Bouchea**, Cham. (Verbenac.).

BUCHIA, H. B. & K. Nov. Gen. et Sp. ii. 269. t. 132 (1817) = **Perama**, Aubl. (Rubiac.).
plantaginea, H. B. & K. l. c. 270. t. 132 = P. dichotoma.

BUCHINGERA, Boiss. & Hohen. Diagn. Ser. I. viii. 29 (1849). *CRUCIFERAE*, Benth. & Hook. f. i. 72.
axillaris, *Boiss. & Hohen. l. c.*—Persia.

BUCHINGERA, F. Schultz, in Jahrb. Pharm. (Mar. 1847).—Cf. Bot. Zeit. vi. (1848) 760 = **Cuscuta**, Linn. (Convolv.).
suaveolens, F. Schultz, l. c. = C. racemosa.

BUCHLOË, Engelm. in Trans. Acad. Sc St. Louis (1859), 432. t. 12, 14. *GRAMINEAE*, Benth. & Hook. f. iii. 1173.
BULBILIS, Rafin. in Journ. Phys. lxxxix. (1819) 262.
CALANTHERA, Nutt. ex Hook. Kew Journ. viii. (1856) 18.
SESLERIA, Nutt. Gen. Am. i. 64 (1818).
dactyloides, *Engelm. l. c.*—Texas; Mexic.

BUCHNERA, Linn. Hort. Cliff. 501 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 968.
BONNETIA, Neck. Elem. i. 368 (1790).
CHYTRA, Gaertn. f. Fruct. iii. 184. t. 214 (1805).
MICALIA, Rafin. Fl. Tellur. ii. 104 (1836).
PIRIPEA, Aubl. Pl. Gui. ii. 627. t. 253 (1775).
aethiopica, Linn. Mant. ii. 251 = Chaenostoma aethiopicum.
africana, Linn. Pl. Afr. Rar. 13; in Amoen. Acad. vi. 89 = Trixago Apula.
americana, *Linn. Sp. Pl.* 630.—Am. bor.
amethystina, Cham. & Schlecht. in Linnaea, ii. (1827) 588 = lobelioides.

BUCHNERA:—
angustifolia, D. Don, Prod. Fl. Nep. 91 = Striga euphrasioides.
angustifolia, *Rafin. New Fl. Am.* ii. 32.—Am. bor.
aquatica, Wight, ex Steud. Nom. ed. II. i. 234 = Striga lutea.
arguta, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 374 = ramosissima.
asiatica, Linn. Sp. Pl. 630; Vahl, Symb. Bot. iii. 81 = Striga densiflora.
aspera, Schum. & Thonn. Beskr. Guin. Pl. 280 = Striga aspera.
asperata, R. Br. Prod. 438 = linearis.
aurantica, Burch. Trav. i. 388 = Lyperia multifida.
aurea, Banks, ex Benth. in Hook. Comp. Bot. Mag. i. (1835) 372 = Polycerena aurea.
bifida, Buch.-Ham. in Wall. Cat. n. 3872 = Striga euphrasioides.
bilabiata, Thunb. Prod. Pl. Cap. 100 = Striga Thunbergii.
canadensis, Linn. Mant. 88 = Verbena Aubletia.
capensis, Linn. l. c. = Polycarena capensis.
capillaris, *Desv. in Ham. Prodr. Fl. Ind. Occ.* 46.—Guiana.
capitata, *Benth. in DC. Prod.* x. 495.—Madag.
capitata, *Burm. f. Fl. Cap. Prod.* 17.—Afr. austr.
cernua, Houtt. Handleid. ix. 542 = Bouchea cuneifolia.
cernua, Linn. Mant. ii. 251 = Bouchea cernua.
coccinea, Benth. Scroph. Ind. 40 = Striga lutea.
coccinea, Scop. Delic. Insub. ii. 10 = Achimenes coccinea.
cordifolia, Linn. f. Suppl. 287 = Streptium asperum.
cruciata, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 91.—As. trop.; Malaya.
cuneifolia, Thunb. Prod. Pl. Cap. 100 = Bouchea cuneifolia.
curviflora, R. Br. Prod. 438 = Striga curviflora.
densiflora, Benth. Scroph. Ind. 41 = Striga densiflora.
densiflora, Hook. & Arn. Bot. Beech. Voy. 203 = cruciata.
denticulata, Buch.-Ham. ex Wall. Cat. n. 3872 = Striga euphrasioides.
distichа, *H. B. & K. Nov. Gen. et Sp.* ii. 340.—N. Granat.
divaricata, Linn. ex Walp. Rep. iii. 308 = Zaluzianskya pusilla.
dura, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 366.—Afr. austr.
elegans, D. Dietr. Syn. Pl. iii. 525 = Striga elegans.
elongata, [Benth.] ex Griseb. Fl. Brit. W. Ind. 428 = longifolia.
elongata, *Sw. Prod. Veg. Ind. Occ.* 92.—Am. calid.
erinoides, *Jarosz, Pl. Nov. Cap.* 19.—Afr. austr.
euphrasioides, Vahl, Symb. Bot. iii. 81 = Striga euphrasioides.
exserta, *Fawc. in H. O. Forbes, Nat. Wand. East. Archip.* 512.—Ins. Timor.
foetida, Andr. Bot. Rep. t. 80 = Chaenostoma hispidum.
foetida, Jacq. Hort. Schoenb. iv. 23. t. 448 = Chaenostoma foetidum.
foetida, *Schrad. Journ.* 1800. II. (1801) 196.—Hab. ?
Forbesii, D. Dietr. Syn. Pl. iii. 526 = Striga Forbesii.
gesnerioides, Willd. Sp. Pl. iii. 338 = Striga orobanchoides.
glabrata, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 366.—Afr. austr.
gracilis, *R. Br. Prod.* 437.—Austral.
grandiflora, Linn. f. Suppl. 287 = Escobedia scabrifolia.
Hermonthica, Delile, Fl. Egypte, 245. t. 34 = Striga Hermonthica.
Heyneana, D. Dietr. Syn. Pl. iii. 525 = Striga densiflora.
hirsuta, Wall. Cat. n. 3869 = Striga euphrasioides.
hispida, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 91.—Ind. or.; Afr. trop.
humifusa, Vahl, Symb. Bot. iii. 81 = Striga humifusa.
hydrabadensis, Roth, Nov. Sp. Pl. 292 = Striga orobanchoides.
juncea, *Cham. & Schlecht. in Linnaea*, ii. (1827) 590.—Bras.
lavandulacea, Cham. & Schlecht. l. c. 589 = rosea.
leptostachya, *Benth. in DC. Prod.* x. 497.—Afr. trop.
levicaulis, *Rafin. New Fl. Am.* ii. 39.—Am. bor.
lilacina, Zipp. ex Span. in Linnaea, xv. (1841) 331 = ramosissima.
linearifolia, *Schum. & Thonn. Beskr. Guin. Pl.* 279.—Afr. trop.
linearis, *R. Br. Prod.* 437.—Austral.

BUCHNERA:—
lithospermifolia, H. B. & K. Nov. Gen. et Sp. ii. 341 = elongata.
lobelioides, *Cham. & Schlecht. in Linnaea*, ii. (1827) 589.—Bras.
longiflora, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 356 = Rhamphicarpa longiflora.
longifolia, H. B. & K. Nov. Gen. et Sp. Pl. ii. 340 = elongata.
longifolia, *Klotzsch, in Peters, Reise Mossamb. Bot.* 225.—Afr. trop.
macrantha, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 366.—Afr. trop.
macrocarpa, *A. Rich. Tent. Fl. Abyss.* ii. 128.—Abyssin.
macrocarpa, *H. B. & K. Nov. Gen. et Sp.* ii. 342.—N. Granat.
major, *Polak. in Linnaea*, xli. (1877) 588.—Costa Rica.
Masuria, Buch.-Ham. ex Benth. Scroph. Ind. 41 = Striga Masuria.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* ii. 457.—Mexic.
missurica, *Rafin. New Fl. Am.* ii. 32.—Am. bor.
montevidensis, Spreng. ex Walp. Rep. iv. 27 = Verbena venosa.
Moouria, Steud. Nom. ed. II. i. 234 = Striga Masurea.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 224.—Afr. trop.
multiflora, D. Dietr. Syn. Pl. iii. 525; F. Muell. Census, 98 = Striga multiflora.
nigrescens, Span. Ic. 69; et in Linnaea, xv. (1841) 331 = arguta.
obliqua, *Benth. in DC. Prod.* x. 498.—Mexic.
oppositifolia, Hort. ex Steud. Nom. ed. II. i. 234 = Chaenostoma hispidum.
orobanchoides, R. Br. in Salt, Abyss. App. 64 = Striga orobanchoides.
palustris, *Spreng. Syst.* ii. 805.—Guiana.
parviflora, R. Br. Prod. 438 = Striga parviflora.
pedunculata, Andr. Bot. Rep. t. 84 = Lyperia pedunculata.
Peripea, Steud. Nom. ed. I. 124 = palustris.
phoenicea, Wall. in Benth. Scroph. Ind. 41 = Striga lutea.
pilosa, Benth. Bot. Voy. Sulph. 144 = elongata.
pinnatifida, Linn. f. Suppl. 288 = Bouchea pinnatifida.
Piripea, Raeusch. Nom. ed. III. 179 = palustris.
pubescens, Benth. in Hook. Comp. Bot. Mag. i. (1835) 365 = ramosissima.
pusilla, *H. B. & K. Nov. Gen. et Sp.* ii. 340.—N. Granat.
ramosissima, *R. Br. Prod.* 438.—Austral.
rosea, *H. B. & K. Nov. Gen. et Sp.* ii. 342.—Venezuela.
rupestris, Sw. Fl. Ind. Occ. ii. 1063 = Rachicallis rupestris.
Schimperiana, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 128 = hispida.
stricta, Benth. in Hook. Comp. Bot. Mag. i. (1835) 367 = cruciata.
sumatrana, Miq. Fl. Ind. Bat. ii. 702 = cruciata.
tenella, *R. Br. Prod.* 437.—Austral.
ternifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 340.—Am. austr.
tetragona, *R. Br. Prod.* 437.—Austral.
tetrasticha, *Wall. Cat.* n. 3878; *et in Benth. Scroph. Ind.* 41.—Assam; Burma.
Thunbergii, D. Dietr. Syn. Pl. iii. 525 = Striga Thunbergii.
timorensis, *Fawc. in H. O. Forbes, Nat. Wand. East. Archip.* 512.—Ins. Timor.
tinctoria, *Bertol. Fl. Guatim.* 26.—Guatemala.
tomentosa, *Blume, Bijdr.* 741.—Java.
urticifolia, *R. Br. Prod.* 437.—Austral.
usuiensis, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 121.—Afr. trop.
verbenoides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 226.—Afr. trop.
virgata, *H. B. & K. Nov. Gen. et Sp.* ii. 342.—N. Granat.
viscosa, [Soland. in] Ait. Hort. Kew. ed. I. ii. 357 = Sphenandra viscosa.
Wallichii, Benth. in Wall. Cat. n. 3876 = Striga Masuria.

BUCHOLTZIA, Meissn. Gen. 318.; Comm. 230 (1841) = BUCHOLZIA, Mart. = **Telanthera**, P. Br. (Amaranth.).

BUCHOLZIA, Mart. Nov. Gen. et Sp. ii. 49. t. 147 (1826) = **Telanthera**, R. Br. (Amaranth.).
brachiata, Schrad. ex Seub. in Mart. Fl. Bras. v. I. 173 = T. ficoidea.
diffusa, Mart. Nov. Gen. et Sp. ii. tt. 149, 151 = T. polygonoides.
ficoidea, Mart. l. c. 52 = T. ficoidea.
filifolia, Hook. f. in Trans. Linn. Soc. xx. (1851) 192 = T. filifolia.
frutescens, Mart. Nov. Gen. et Sp. ii. 52, in obs. = T. frutescens.
glaucescens, Hook. f. in Trans. Linn. Soc. xx. (1851) 191 = T. glaucescens.
lupulina, Mart. Beitr. Amarant. 108 = T. lupulina.
maritima, Mart. Nov. Gen. et Sp. ii. 50. t. 147 = T. maritima.
nudicaulis, Hook. f. in Trans. Linn. Soc. xx. (1851) 191 = T. nudicaulis.
obovata, Mart. & Gal. in Bull. Acad. Brux. x. (1843) 348 = T. obovata.
philoxeroides, Mart. Beitr. Amarant. 107. n. 5 = T. philoxeroides.
polygonoides, Mart. Nov. Gen. et Sp. ii. 51 = T. polygonoides.
spinescens, Gill. ex Moq. in DC. Prod. xiii. II. 359 = Alternanthera Achyrantha.

BUCHOLZIA, Stadtm. ex Willem. in Ust. Ann. Bot. xviii. (1796) 54 = **Combretum**, vel **Terminalia**, Linn.

BUCHOZIA, L'Hérit. ex Juss. in Dict. Sc. Nat. v. (1804) 409; et in Mém. Mus. Par. vi. (1820) 395 = **Serissa**, Comm. (Rubiac.).
coprosmoides, L'Hérit. ex DC. Prod. iv. 575 = S. foetida.

BUCHOZIA, Vell. Fl. Flum. 33 (1825) = **Heteranthera**, Ruiz & Pav. (Ponteder.).
aquatica, Vell. l. c. 33; i. t. 80 = H. reniformis.

BUCIDA, Linn. Syst. ed. X. 1025 (1759) = **Terminalia**, Linn. (Combret.).
angustifolia, DC. Prod. ii. 10 = T. Buceras.
angustifolia, Spruce, ex Eichl. in Mart. Fl. Bras. xiv. II. 96 = T. Hilariana.
Buceras, Linn. Syst. ed. X. 1025 = T. Buceras.
Buceras, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 272 = T. Catappa.
Buceras, Vell. Fl. Flum. 172; iv. t. 87 = Laguncularia racemosa.
capitata, Vahl, Eclog. Am. i. 50. t. 8 = T. Hilariana.
Comintana, Blanco, Fl. Filip. ed. I. 856 = T. Chebula.
megapotamica, Spreng. Syst. iv. Cur. Post. 177.—Bras.
nitida, Hassk. in Flora, xxvii. (1844) 605.—Ins. Sandvic.

BUCKINGHAMIA, F. Muell. Fragm. vi. 248 (1868). *PROTEACEAE*, Benth. & Hook. f. iii. 182.
celsissima, *F. Muell. l. c.*—Austral.

BUCKLANDIA, R. Br. in Wall. Cat. n. 7414 (1832). *HAMAMELIDEAE*, Benth. & Hook. f. i. 668.
populifolia, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 86 = populnea.
populnea, *R. Br. in Wall. Cat.* n. 7414.—Reg. Himal.

BUCKLEYA, Torr. in Am. Journ. Sc. xlv. (1843) 170. *SANTALACEAE*, Benth. & Hook. f. iii. 226.
DARBYA, A. Gray, in Am. Journ. Sc. Ser. II. i. (1846) 388.
NESTRONIA, Rafin. New Fl. Am. iii. 12 (1836).
QUADRIALA, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1845) 194; (Fam. Nat. Fl. Jap. i. 86. t. 2).
distichophylla, *Torr. l. c.*—Am. bor.

BUCKNERA, Michx. Fl. Bor. Am. ii. 18 (1804) = **Buchnera**, Linn. (Scrophul.).

BUCQUETIA, DC. Prod. iii. 110 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 736.
glutinosa, *DC. l. c.*—N. Granat.
nigritella, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 23.—Ecuador.

BUDA, Adans. Fam. ii. 507 (1763); Dum. Fl. Belg. 110 (1827) = **Spergularia**, C. & J. Presl (1817) (Caryoph.).
marina, Dum. Fl. Belg. 110 = S. marina.
media, Dum. l. c. = S. marina.
rubra, Dum. l. c. = S. rubra.

BUDDLEIA, Houst. ex Linn. Gen. ed. I. 26 (1737). *LOGANIACEAE*, Benth. & Hook. f. ii. 793.
ADENOPLEA, Radlk. in Bremen Abh. viii. (1883) 406.
ROMANA, Vell. Fl. Flum. 54, pro parte (1825).
abbreviata, *H. B. & K. Nov. Gen. et Sp.* ii. 353.—Mexic.
acuminata, R. Br. in Salt, Abyss. App. 62 = polystachya.
acuminata, H. B. & K. l. c. 349 = Humboldtiana.
acuminata, *Poir. Encyc. Suppl.* i. 745.—Ind. or.
acuminatissima, Blume, Bijdr. 743 = asiatica.
alpina, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 25.—Costa Rica.
alternifolia, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 494.—China.
americana, H. B. & K. Nov. Gen. et Sp. ii. 351 = spicata.
americana, *Linn. Sp. Pl.* 112.—Am. austr.
angustata, Benth. in DC. Prod. x. 443 = vetula.
araucana, *Phil. in Anal. Univ. Chil.* (1873) 525.—Chili.
arborea, Meyen, Reise, i. 478 = pichinchensis.
aromatica, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 227.—Bolivia.
asiatica, *Lour. Fl. Cochinch.* 72.—As. trop.; Malaya.
auriculata, *Benth. in Hook. Comp. Bot. Mag.* ii. (1836) 60.—Afr. austr.
australis, Vell. Fl. Flum. 41; i. t. 104 = brasiliensis.
axillaris, Willd. in Roem. & Schult. Syst. iii. 97 = Adenoplusia Willdenowii.
barbata, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 12.—Mexic.
betonicaefolia, *Lam. Illustr.* i. 291.—Peruv.
braehiata, *Cham. & Schlecht. in Linnaea*, ii. (1827) 599.—Bras.
brasiliensis, *Jacq. f. Eclog.* t. 158.—Bras.
brevifolia, Willd. ex Roem. & Schult. Syst. iii. Mant. 97 = abbreviata.
bullata, *H. B. & K. Nov. Gen. et Sp.* ii. 348.—Ecuador.
callicarpoides, H. B. & K. l. c. 350 = americana.
calycina, *Benth. in DC. Prod.* x. 440.—Ecuador.
campestris, *Walp. Rep.* iii. 330.—Bras.
cana, Willd. ex Roem. & Schult. Syst. iii. Mant. 94 = verbascifolia.
capitata, Jacq. Ic. Rar. ii. 9; Coll. ii. 332 = globosa.
carnea, Carr. in Rev. Hortic. (1879) 90 = curviflora.
cestriflora, *Cham. in Linnaea*, viii. (1833) 20.—Bras.
Colvilei, *Hook. f. Illustr. Himal. Pl.* t. 18.—Reg. Himal.
comorensis, *Baker, in Journ. Linn. Soc.* xx. (1884) 206.—Ins. Comor.
connata, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 283 = brasiliensis.
connata, *Ruiz & Pav. Fl. Per.* i. 52. t. 18.—Peruv.
cordata, *H. B. & K. Nov. Gen. et Sp.* ii. 348. t. 185.—Mexic.
cordobensis, *Griseb. in Goett. Abh.* xix. (1874) 213.—Reg. Argent.
coriacea, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 226.—Bolivia.
crispa, Benth. in Wall. Cat. n. 6404; et Scroph. nd. 43 = paniculata.
crotonoides, *A. Gray, in Proc. Am. Acad.* v. (1861) 165.—Calif.
cuneata, *Cham. in Linnaea*, viii. (1833) 17.—Bras.
curviflora, *Hook. & Arn. Bot. Beech. Voy.* 267.—China.
curvifolia, *Hort. ex Carr. in Rev. Hortic.* (1866) 339.—Hab.?
decurrens, Cham. & Schlecht. in Linnaea, v. (1830) 105 = americana.
densiflora, *Blume, Bijdr.* 743.—Java.
dentata, *H. B. & K. Nov. Gen. et Sp.* ii. 352.—Venezuela.
diffusa, *Ruiz & Pav. Fl. Per.* i. 53. t. 80.—Peruv.
discolor, Roth, Nov. Pl. Sp. 83 = asiatica.
diversifolia, Vahl, Symb. Bot. iii. 15 = Nicodemia diversifolia.
elegans, *Cham. & Schlecht. in Linnaea*, ii. (1827) 594.—Bras.

BUDDLEIA:—
elliptica, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 24.—Mexic.
ferruginosa, Pohl, ex J. A. Schmidt, in Mart. Fl. Bras. 284 = elegans.
floccosa, *Kunth, in Ind. Sem. Hort. Berol.* (1844).—*Cf. Linnaea*, xviii. (1844) 501.—Mexic.
floribunda, *H. B. & K. Nov. Gen. et Sp.* ii. 352.—Mexic.
foliata, R. Br. ex Benth. in DC. Prod. x. 446 = polystachya.
fusca, *Baker, in Journ. Linn. Soc.* xx. (1884) 205.—Madag.
Gayana, *Benth. in DC. Prod.* x. 442.—Chili.
glaberrima, Hort. Amat. Cat. Hort. Taurin. 13 = Freylinia cestroides.
glabra, Hort. ex C. Koch, Dendrol. ii. II. 295 = curviflora.
glabrata, Spreng. Syst. i. 431 = Cephalanthus Sarandi.
globifera, Mirb. in Duham. Arb. ed. Nov. i. t. 25 = globosa.
globosa, *Hope, in Verh. Harlem.* xx. II. (1782) 417. t. 11.—Chili; Peruv.
glomerata, *Wendl. ex Benth. in DC. Prod.* x. 447, nomen.—Afr. austr.
gracilis, *Kunth, in Ind. Sem. Hort. Berol.* (1844).—*Cf. Linnaea*, xviii. (1844) 500.—Mexic.
grandiflora, *Cham. & Schlecht. in Linnaea*, ii. (1827) 596.—Bras.
guianensis, *Roxb. Hort. Beng.* [10], nomen.—Am. austr.
heterophylla, Lindl. Bot. Reg. t. 1259 = madagascariensis.
Humboldtiana, *Roem. & Schult. Syst.* iii. 93.—Texas; Mexic.
incana, *Ruiz & Pav. Fl. Per.* i. 52. t. 80.—Peruv.
incompta, *Linn. f. Suppl.* 123.—Afr. austr.
indica, Lam. Encyc. i. 513 = Nicodemia diversifolia.
insignis, *A. Van Geert, Cat.* n. 78 (1879–80), 8; *Carr. in Rev. Hortic.* (1878) 330.—Hab.?
integrifolia, Willd. ex Roem. & Schult. Syst. iii. Mant. 96 = abbreviata.
intermedia, *H. B. & K. Nov. Gen. et Sp.* ii. 350.—Mexic.
intermedia, *Lorenz, in Roca, Exped. Rio-Negro*, ii. 255, nota.—Patagon.
interrupta, Buch.-Ham. ex Wall. Cat. sub n. 6401 = asiatica.
interrupta, *H. B. & K. Nov. Gen. et Sp.* ii. 348.—Ecuador.
Jamesonii, *Benth. in DC. Prod.* x. 441.—Ecuador.
japonica, *Linden, Cat.* n. 19 (1865).—Japon.
lanata, *Benth. in DC. Prod.* x. 441.—Peruv.
lanceolata, *Benth. Pl. Hartw.* 48.—Am. centr.
lanceolata, Heyne, ex Wall. Cat. sub n. 6401 = asiatica.
Lindenii, *Benth. in DC. Prod.* x. 438.—N. Granat.
Lindleyana, *Fortune, in Lindl. Bot. Reg.* (1844) *Misc.* 25.—China.
longifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 349.—Peruv.
Lucae, *Niederl. in Roca, Exped. Rio-Negro*, ii. 254.—Patagon.
macrantha, Mart. ex J. A. Schmidt, in Mart. Fl. Bras. viii. 285 = grandiflora.
macrophylla, *Kunth, in Ind. Sem. Hort. Berol.* (1844).—*Cf. Linnaea*, xviii. (1844) 500.—Hab.?
macrostachya, *Benth. in Wall. Cat.* n. 6407; *et Scroph. Ind.* 42.—Reg. Himal.
madagascariensis, *Lam. Encyc.* i. 513.—Ins. Mascar.; Madag.
marrubiifolia, *Benth. in DC. Prod.* x. 441.—Texas; Mexic.
Martii, J. A. Schmidt, in Journ. Bot. vi. (1868) 228, 245 = macrostachya.
melliodora, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 10.—Mexic.
mendozinensis, *Gill. ex Benth. in DC. Prod.* x. 443.—Chili.
microphylla, *H. B. & K. Nov. Gen. et Sp.* ii. 353.—Mexic.
Missionis, *Benth. in Wall. Cat.* n. 6405; *et Scroph. Ind.* 43.—Ind. or.
mollis, *H. B. & K. Nov. Gen. et Sp.* ii. 349.—Ecuador.
Nappii, *Lorenz, in Roca, Exped. Rio-Negro*, ii. 254.—Patagon.
Neemda, Buch.-Ham. in Wall. Cat. n. 6401 = asiatica.
Neemda, Hort. ex Link, Enum. Hort. Berol. i. 125 = brasiliensis.

BUDDLEIA:—
nepalensis, Colla, in Mem. Acc. Torin. xxxv. (1835) 182 = paniculata.
nitida, *Benth. in DC. Prod.* x. 437.—Mexic.
oblonga, *Benth. l. c.* 442.—Bras.
obtusifolia, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 25 = microphylla.
occidentalis, *Linn. Sp. Pl.* ed. II. 162.—Jamaic.; Peruv.
occidentalis, Ruiz & Pav. Fl. Per. i. 53. t. 82. f. a = americana.
officinalis, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 496.—China.
otophylla, *Hassk. in Versl. en Med. Kon. Akad. Wetensch.* v. (1857) 97.—Malaya.
ovalifolia, *Kunth, in Ind. Sem. Hort. Berol.* (1844).—*Cf. Linnaea*, xviii. (1844) 500.—Mexic.
paniculata, *Wall. Cat.* n. 6403; *et in Roxb. Fl. Ind. ed. Carey*, i. 412.—Reg. Himal.; Burma.
parviflora, *H. B. & K. Nov. Gen. et Sp.* ii. 353.—Mexic.
parvifolia, Willd. ex Roem. & Schult. Syst. iii. Mant. 96 = microphylla.
perfoliata, *H. B. & K. Nov. Gen. et Sp.* ii. 346.—Mexic.
pichinchensis, *H. B. & K. l. c.*—Ecuador.
Poiretii, *Spreng. Syst.* i. 430.—Ind. or.
polycephala, *H. B. & K. Nov. Gen. et Sp.* ii. 346.—Peruv.
polystachya, *Fresen. in Flora*, xxi. (1838) 605.—Abyssin.
Pringlei, *A. Gray, in Proc. Am. Acad.* xix. (1883) 86.—Arizona.
propinqua, *Kunth, Ind. Sem. Hort. Berol.* (1844).—*Cf. Linnaea*, xviii. (1844) 501.—Mexic.
pseudoverticillata, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 24 = sessiliflora.
racemosa, *Torr. Bot. Mex. Bound.* 121.—Texas.
rondeletiaeflora, *Benth. in DC. Prod.* x. 445.—Ins. Comor.
rufa, *Fresen. in Flora*, xxi. (1838) 606.—Abyss.
rufescens, Willd. ex Roem. & Schult. Syst. iii. Mant. 97 = americana.
rugosa, *H. B. & K. Nov. Gen. et Sp.* ii. 347.—N. Granat.
salicifolia, Jacq. Hort. Schoenb. i. 12 = Chilianthus arboreus.
salicifolia, Vahl, Symb. Bot. iii. 14 = thyrsoidea.
salicina, Lam. Illustr. i. 291 = asiatica.
saligna, Willd. Enum. Hort. Berol. 159 = Chilianthus arboreus.
Saltiana, Steud. Nom. ed. II. i. 235 = polystachya.
salvifolia, *Lam. Encyc.* i. 513.—Afr. austr.
scordioides, *H. B. & K. Nov. Gen. et Sp.* ii. 345. t. 183.—Texas; Mexic.
serrulata, Roth, Nov. Pl. Sp. 82 = asiatica.
sessiliflora, *H. B. & K. Nov. Gen. et Sp.* ii. 345. t. 182.—Mexic.
sinuata, *Willd. ex Roem. & Schult. Syst. Mant.* iii. 97.—Madag.
spectabilis, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 11.—Mexic.
sphaerantha, Cham. & Schlecht. in Linnaea, v. (1830) 104 = perfoliata.
sphaerocephala, *Baker, in Journ. Linn. Soc.* xxi. (1885) 425.—Madag.
spicata, *Ruiz & Pav. Fl. Per.* i. 53. t. 81.—Peruv.
stachyoides, *Cham. & Schlecht. in Linnaea*, ii. (1827) 597.—Bras.
snaveolens, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1845) 11.—Chili.
subserrata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 92 = asiatica.
sundaica, *Blume, Bijdr.* 743.—Java.
tenuifolia, *Griseb. in Goett. Abh.* xix. (1874) 213.—Am. austr.
ternata, *Lour. Fl. Cochinch.* 72.—Cochinch.
thapsoides, Desf. Cat. Hort. Par. ed. III. 106 = brasiliensis.
thyrsoides, *Lam. Illustr.* i. 291.—Reg. Argent.
tubiflora, *Benth. in DC. Prod.* x. 443.—Bras.
tucumanensis, *Griseb. in Goett. Abh.* xix. (1874) 213.—Tucuman.
venusta, *Kunth, in Ind. Sem. Hort. Berol.* (1844).—*Cf. Linnaea*, xviii. (1844) 500.—Mexic.
Verbusci, Pers. Syn. i. 135 = americana.
verbascifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 351.—N. Granat.

BUDDLEIA :—
verticillata, *H. B & K. Nov. Gen. et Sp.* 346. t. 184.—Mexic.
vetula, *Cham. in Linnaea*, viii. (1833) 18.—Bras.
virgata, Blanco, Fl. Filip. ed. I. 57 = asiatica.
virgata, Linn. f. Suppl. 123 = Gomphostigma scoparioides.
volubilis, *Lam. Illustr.* i. 291.—Ins. Mascar.

BUECHNERIA, Roth, Nov. Pl. Sp. 292 (1821) = **Buchnera**, Linn. (Scrophul.).

BUECKIA, A. Rich. in Orb. Dict. ii. 759 (1842) = BUEKIA, Nees = **Elynanthus**, Linn. (Cyperac.).

BUEKIA, Giseke, Prael. Linn. 199, 204 (1792) = **Alpinia**, Linn. (Scitamin.).
malaccensis, Raeusch. Nom. ed. III. 1 = A. malaccensis.

BUEKIA, Nees, in Linnaea, ix. (1834) 300 = **Elynanthus**, Linn. (Cyperac.).
punctoria, Nees, l. c. = E. punctorius.

BUELOWIA, Schum. & Thonn. Beskr. Guin. Pl. 246 (1827) = **Smeathmannia**, Soland. (Passiflor.).
illustris, Schum. & Thonn. l. c. 247 = S. pubescens.

BUENA, Cav. in Anal. Hist. Nat. ii. (1800) 278 ; Ic. vi. 49 = GONZALAGUNIA, Ruiz & Pav. = **Gonzalea**, Pers. (Rubiac.).
panamensis, Cav. ll. cc. 279 ; 50 = G. panamensis.

BUENA, Pohl, Pl. Bras. Ic. i. 8 (1827) ; Wedd. in Journ. Linn. Soc. xi. (1869) 185 = **Cosmibuena**, Ruiz & Pav. et **Cascarilla**, Wedd. (Rubiac.).
acuminata, DC. Prod. iv. 356 = Cosmibuena acuminata.
acutifolia, Wedd. in Journ. Linn. Soc. xi. (1869) 186 = Cascarilla acutifolia.
bullata, Wedd. l. c. 187 = Cascarilla bullata.
calycina, Wedd. l. c. = Cascarilla calycina.
Carua, Wedd. l. c. = Cascarilla Carua.
citrifolia, Wedd. l. c. = Cascarilla citrifolia.
crassifolia, Wedd. l. c. = Cascarilla calyptrata.
Gaudichaudiana, Wedd. l. c. = Cascarilla Gaudichaudiana.
heterophylla, Wedd. l. c. 186 = Cascarilla heterophylla.
hexandra, Pohl, Pl. Bras. Ic. i. 10. t. 8 = Cascarilla hexandra.
Hookeriana, Wedd. in Journ. Linn. Soc. xi. (1869) 187 = Cascarilla Hookeriana.
Lambertiana, Wedd. l. c. = Cascarilla Lambertiana.
latifolia, Benth. Pl. Hartw. 191 = Cosmibuena latifolia.
macrocarpa, Benth. Bot. Voy. Sulph. 104. t. 38 = Cosmibuena macrocarpa.
macrocarpa, Wedd. in Journ. Linn. Soc. xi. (1869) 187 = Cascarilla macrocarpa.
magnifolia, Wedd. l. c. 186 = Cascarilla magnifolia.
muzonensis, Wedd. l. c. 187 = Cascarilla muzonensis.
nitida, Wedd. l. c. 186 = Cascarilla nitida.
obtusifolia, DC. Prod. iv. 356 = Cosmibuena obtusifolia.
Pavonii, Wedd. in Journ. Linn. Soc. xi. (1869) 186 = Cascarilla Pavonii.
Riedeliana, Wedd. l. c. = Cascarilla Riedeliana.
Riveroana, Wedd. l. c. = Cascarilla magnifolia.
Roraimae, Wedd. l. c. 187 = Cascarilla Roraimae.
Skinneri, Oerst. in Kjoeb. Vidensk. Meddel. (1852) 48 = Cosmibuena Skinneri.
stenocarpa, Wedd. in Journ. Linn. Soc. xi. (1869) 186 = Cascarilla stenocarpa.
triflora, Benth. in Hook. Journ. Bot. iii. (1841) 216 = Cosmibuena triflora.
undata, Wedd. in Journ. Linn. Soc. xi. (1869) 187 = Cascarilla undata.

BUERGERIA, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 53 (1867) = **Cladrastis**, Rafin. (Legumin.).
floribunda, Miq. l. c. = C. amurensis.

BUERGERIA, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1845) 186. t. 2 (Fl. Jap. Fam. i. 78. t. 2) = **Talauma**, Juss. (Magnol.).
obovata, Sieb. & Zucc. ll. cc. 187 = Magnolia obovata.
salicifolia, Sieb. & Zucc. l. c. = T. salicifolia.
stellata, Sieb. & Zucc. l. c. 186 = T. stellata.

BUETTNERA, J. F. Gmel. Syst. ii. 404 (1791) = seq.

BUETTNERIA, Loefl. Iter Hisp. 313 (1758) (Byttneria) ; Linn. Syst. ed. X. ii. 939 (1759). *STERCULIACEAE*, Benth. & Hook. f. i. 225.
BYTTNERIA, Loefl. Iter Hisp. 313 (1758).
CHAETAEA, Jacq. Enum. Pl. Carib. 2 (1760).
HETEROPHYLLUM, Boj. ex Hook. Bot. Misc. i. (1830) 291.
PENTACEROS, G. F. W. Mey. Prim. Fl. Esseq. 136 (1818).
TELFAIRIA, Newm. ex Hook. Bot. Misc. i. (1830) 291.
abutiloides, St. Hil. & Naud. in Ann. Sc. Nat. Sér. II. xviii. (1842) 31 = hirsuta.
aculeata, Domb. ex Lam. Encyc. i. 523 = cordata.
aculeata, Jacq. Stirp. Am. 76 = carthagenensis.
acuminata, *Bred. ex Roem. & Schult. Syst.* v. 470.—Venezuela.
affinis, Pohl, Pl. Bras. ii. 73 = melastomaefolia.
africana, *Mast. in Oliver, Fl. Trop. Afr.* i. 239.—Afr. trop.
andamanensis, *Kurz, in Journ. As. Soc.* Beng. ii. (1871) 47 ; *et Flora*, liv. (1871) 277.—Burma.
angulata, *Hassk. Cat. Pl. Hort. Bog.* 204.—Java.
aspera, *Colebr. in Roxb. Fl. Ind. ed. Carey*, ii. 383.—Reg. Himal. ; China.
australis, Sieber, ex Steud. Nom. ed. II. i. 241 = pannosa.
australis, St. Hil. Fl. Bras. Mer. i. 145 = tereticaulis.
biloba, *Baill. Adansonia*, x. (1872) 178.—Madag.
brasiliensis, *Spreng. Syst.* i. 790.—Bras.
brevipes, *Benth. Pl. Hartw.* 115.—Am. trop.
caledonica, *Turcz. in Bull. Soc. Nat. Mosc.* (1863) I. 568.—N. Caled.
carthagenensis, *Jacq. Stirp. Am. Piot.* 41.—N. Granat.
catalpaefolia, *Jacq. Hort. Schoenbr.* i. 21.—Am. trop.
catalpifolia, Wall. Cat. n. 1152 = crenulata ?
celtoides, *St. Hil. Fl. Bras. Mer.* i. 141.—Bras.
corchorifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 152.—Am. austr.
cordifolia, *Sagot, in Ann. Sc. Nat.* Sér. VI. xi. (1881) 155.—Guiana.
cordata, *Lam. Encyc.* i. 523.—Peruv.
corylifolia, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 470.—Am. austr.
crenulata, *Wall. Cat.* n. 1150.—Reg. Himal. ; Burma.
dasyphylla, J. Gay, in DC. Prod. i. 486 = Rulingia pannosa.
dentata, *Pohl, Pl. Bras.* ii. 79.—Bras.
discolor, *Benth. in Hook. Kew Journ.* iii. (1851) 165.—Bras.
divaricata, *Benth. in Hook. Journ. Bot.* iv. (1842) 124.—Guiana.
echinata, Wall. Cat. n. 1149 = crenulata ?
elliptica, *Mast. in Hook. f. Fl. Brit. Ind.* i. 377.—Malacca.
elliptica, Pohl, Pl. Bras. ii. 71 = melastomaefolia.
flaccida, *Span. Ic.* n. 18 ; *in Linnaea*, xv. (1841) 174.—Ins. Timor.
fulva, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 37.—Bras.
Gayana, *St. Hil. Fl. Bras. Mer.* i. 145.—Bras.
geminifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 153.—Ecuador.
genistella, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 333.—N. Granat.
glabrescens, *Benth. Bot. Voy. Sulph.* 71.—Am. trop.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 502.—Madag.
grandifolia, DC. Prod. i. 486 = aspera.
herbacea, *Roxb. Pl. Corom.* i. 28. t. 29.—Ind. or.
hermanniaefolia, J. Gay, in DC. Prod. i. 486 = Rulingia hermanniaefolia.
heterophylla, *Hook. Bot. Misc.* i. (1830) 291.—Madag.
hirsuta, *Ruiz & Pav. Fl. Per.* iii. 10.—Peruv.
hypoleuca, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 207 = Commersonia platyphylla.
inodora, J. Gay, ex DC. Prod. i. 486 = Rulingia pannosa.
Jackiana, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 386.—Ins. Penang.
jaculifolia, *Pohl, Pl. Bras.* ii. 77.—Bras.
javanica, Spreng. ex Miq. Illustr. Fl. Ind. Arch. 91 = angulata.
laevigata, Schott, in Spreng. Syst. Veg. iv. App. 404 = Gayana.
lanceolata, *Sesse, ex DC. Prod.* i. 487.—Mexic.

BUETTNERIA :—
lancifolia, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 158.—Borneo.
lancifolia, *St. Hil. & Naud. in Ann. Sc. Nat.* Sér. II. xviii. (1842) 32.—Bras.
lateralis, *Presl, Rel. Haenk.* ii. 145.—Mexic.
lobata, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 503.—Madag.
longicuspis, *Baill. l. c.* 502.—Madag.
longifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) II. 154.—Venezuela.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* v. 315.—N. Granat.
Maingayi, *Mast. in Hook. f. Fl. Brit. Ind.* i. 377.—Malacca.
melastomaefolia, *St. Hil. Fl. Bras. Mer.* i. 144.—Bras.
melastomoides, G. Don, Gen. Syst. i. 524 = praec.
microphylla, *Jacq. Hort. Vindob.* i. 10. t. 29 ; *Linn. Mant.* 209.—Ind. occ.
mollis, *H. B. & K. Nov. Gen. et Sp.* v. 314. t. 481 a et b.—N. Granat.
morifolia, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 332.—N. Granat.
nepalensis, Turcz. in Bull. Soc. Nat. Mosc. (1858) I. 207 = aspera.
obliqua, *Benth. in Hook. Journ. Bot.* iv. (1842) 124.—Guiana.
oblongata, *Pohl, Pl. Bras.* ii. 73.—Bras.
ovata, *Lam. Encyc.* i. 522.—Peruv.
pannosa, Bauer, ex Steetz, in Lehm. Pl. Preiss. ii. 350 = Keraudrenia Hookeriana.
pannosa, DC. Prod. i. 486 = Rulingia pannosa.
parviflora, *Benth. Pl. Hartw.* 114.—Ecuador.
peruviana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 207.—Peruv.
pilosa, *Roxb. Hort. Beng.* 18 ; *Fl. Ind.* i. 618.—Ind. or. ; Burma.
Pohliana, *Steud. Nom.* ed. II. i. 242.—Bras.
polyacantha, Wall. Cat. n. 1151 = crenulata ?
ramosissima, Pohl, Pl. Bras. ii. 75 = scabra.
Reinwardtii, *Korth. in Nederl. Kruidk. Arch.* (1848) i. 310.—Borneo.
reticaulis, Cav. Diss. 292, sphalm. = tereticaulis.
rhamnifolia, *Benth. in Hook. Kew Journ.* iii. (1851) 164.—Bras.
rivularis, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 333.—Bras.
rubricaulis, *Presl, Rel. Haenk.* ii. 145.—Mexic.
sagittaefolia, *St. Hil. Fl. Bras. Mer.* i. 142. t. 27.—Bras.
salicifolia, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 470.—Venezuela.
salicifolia, *Presl, Rel. Haenk.* ii. 144.—Mexic.
scabra, *Linn. Syst.* ed. X. 197.—Am. austr.
scabra, Pohl, Pl. Bras. ii. 81 = Pohliana.
scalpellata, Pohl, l. c. 78 = scabra.
scorpiura, *Wright, ex Griseb. Cat. Pl. Cub.* 29.—Cuba.
sidaefolia, St. Hil. Fl. Bras. Mer. i. 146 = catalpaefolia.
sulcata, *Ruiz & Pav. Fl. Per.* iii. 10.—Peruv.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 68.—Ins. Tahiti.
tereticaulis, *Lam. Encyc.* i. 523.—Peruv.
tiliaefolia, *Presl, Rel. Haenk.* ii. 144.—Mexic.
uncinata, *Mast. in Hook. f. Fl. Ind.* i. 377.—Malacca.
velutina, Wall. Cat. nn. 1148, 1925 = pilosa.
virgata, Pohl, Pl. Bras. ii. 80 = scabra.
vitifolia, *Baill. Adansonia*, x. (1872) 177.—Madag.
Voulily, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 502.—Madag.

BUFFONEA, Koch, Syn. Fl. Germ. ed. I. 108 (1836) = **Bufonia**, Linn. (Caryophyll.).

BUFONIA, Sauv. ex Linn. Diss. Dass. 2 (1747) ; Amoen. Acad. i. 386 (1749). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 151.
alsinoides, Hill, Veg. Syst. xi. 27 = annua ?
angustifolia, Cav. ex Willk. & Lange, Prod. Fl. Hisp. iii. 605 = tenuifolia.
annua, DC. Fl. Fr. iv. 768 = tenuifolia et macrosperma.
brachyphylla, *Boiss. & Heldr. Diagn.* Ser. I. viii. 94.—Graecia et Ins.
calyculata, *Boiss. & Bal. l. c.* Ser. II. v. 58.—Cilicia.
Duvaljouvii, *Batt. & Trab. in Bull. Soc. Bot. Fr.* xxvi. (1879) 56.—Afr. bor.
elata, Boiss. Diagn. Ser. I. i. 44 = macrocarpa.
enervis, *Boiss. l. c.*—Persia.

BUFONIA :—
hebecalyx, *Boiss. Diagn.* Ser. I. i. 45.—Persia.
Kotschyana, *Boiss. l. c.* Ser. II. i. 84.—Persia.
macrocarpa, *Ser. in DC. Prod.* i. 388. — Persia; Afghan.
macropetala, *Willk. in Flora*, xxxiv. (1851) 604; *et Ic. Pl. Hisp.* 112.—Hispan.
macrosperma, *J. Gay, in Mutel, Fl. Dauph.* ed. II. 88. —Europ.; As. Min.; Syria.
multiceps, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 259.—Arabia.
Oliveriana, *Ser. in DC. Prod.* i. 388.—Mesopot.; Persia: Afghan.
paniculata, Dubois, in Delarb. Fl. Auvergne, ed. II. 300 = tenuifolia.
parviflora, Griseb. Spicil. Fl. Rumel. i. 197 = tenuifolia.
perennis, *Pourr. in Mém. Acad. Toul.* iii. (1788) 309.—Gallia.
rotundifolia, Buch.-Ham. ex Steud. Nom. ed. II. i. 235 = Drymaria cordata.
spicata, Delarb. Fl. Auvergne, ed. II. 299 = macrosperma.
tenuifolia, *Linn. Sp. Pl.* 123. — Europ. austr.; Marocco; Reg. Caucas.
tenuifolia, Moç. & Sessé, ex DC. Prod. i. 390 = Hymenella moehringioides.
tenuifolia, Vill. Hist. Pl. Dauph. iii. 650 = macrosperma.
virgata, *Boiss. Fl. Orient.* i. 665.—As. Min.; Syria.
Willkommiana, Boiss. Diagn. Ser. II. i. 83 = perennis.

BUFORRESTIA, C. B. Clarke, in DC. Monog. Phan. iii. 233. t. 6, 7 (1881). *COMMELINACEAE*, Benth. & Hook. f. Gen. iii. 850.
Candolleana, *C. B. Clarke, l. c.* 233.—Guiana.
imperforata, *C. B. Clarke, l. c.* 234.—Afr. trop.
Mannii, *C. B. Clarke, l. c.* 233.—Afr. trop.
tenuis, *C. B. Clarke, l. c.* 234.—Afr. trop.

BUGENVILLEA, Endl. Gen. 312 (1837) = seq.

BUGINVILLAEA, Comm. ex Juss. Gen. 91 (1789) = **Bougainvillaea**, Juss. (Nyctag.).

BUGLOSSA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 351 (1821) = **Lycopsis**, Linn. (Boragin.).
arvensis, S. F. Gray, l. c. = L. arvensis.
sempervirens, S. F. Gray, l. c. 352 = Anchusa sempervirens.

BUGLOSSITES, Moris, in Enum. Sem. Hort. Taur. (1845).—Cf. Ann. Sc. Nat. Sér. III. v. (1846) 365 = **Borago**, Linn.
laxiflora, Moris, l. c. = Borago laxiflora.

BUGLOSSOIDES, [Riv.]. Moench, Meth. 418 (1794) = **Lithospermum**, Linn. (Borag.).
ramosissima, Moench, l. c. = L. tenuiflorum.

BUGLOSSUM, [Tourn.]. Adans. Fam. ii. 178 (1763); Gaertn. Fruct. i. 322. t. 67 (1788) = **Anchusa**, Linn. (Boragin.).
amoenum, Gaertn. l. c. = A. italica.
angustifolium, All. Fl. Pedem. i. 47 = A. undulata.
angustifolium, Moench, Meth. 418 = A. angustifolia.
aspermum, Gilib. Fl. Lituan. i. 21 = Anchusa officinalis.
Barrellieri, All. Fl. Pedem. i. 48 = A. Barrelieri.
caeruleum, Pers. Syn. i. 159 = A. italica.
echinatum, Tausch, in Flora, vii. (1824) 230 = A. strigosa.
elatum, Moench, Meth. 418 = A. italica.
italicum, Tausch, in Flora, vii. (1824) 230 = A. italica.
officinale, Lam. Fl. Fr. ii. 278 = A. officinalis.
paniculatum, Tausch, in Flora, vii. (1824) 228 = A. italica.
sempervirens, All. Fl. Pedem. i. 48 = A. sempervirens.
syriacum, Tausch, in Flora, vii. (1824) 231 = A. strigosa.
tinctorium, Lam. Fl. Fr. ii. 278 = Alkanna tinctoria.
vulgare, Tausch, in Flora, vii. (1824) 229 = A. italica.

BUGRANOPSIS, Pomel, Nouv. Mat. Fl. Atl. 171 (1874) = **Ononis**, Linn. (Legumin.).
alopecuroides, Pomel, l. c. 172 = O. alopecuroides.
cephalantha, Pomel, l. c. = O. cephalantha.

BUGRANOPSIS :—
crinita, Pomel, Nouv. Mat. Fl. Atl. 171 = O. crinita.
euphrasiaefolia, Pomel, l. c. 172 = O. euphrasiaefolia.
megalostachys, Pomel, l. c. 171 = O. megalostachys.
mitissima, Pomel, l. c. 172 = O. mitissima.
rosea, Pomel, l. c. 171 = O. rosea.
Salzmaniana, Pomel, l. c. 172 = O. Salzmanniana.
stricta, Pomel, l. c. = O. stricta.
Tournefortii, Pomel, l. c. = O. Tournefortii.
variegata, Pomel, l. c. = O. variegata.

BUGULA, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Ajuga**, Linn. (Labiat.).
alpina, All. Fl. Pedem. i. 45 = A. genevensis.
arvensis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 368 = A. Chamaepitys.
Chamaepithys, Scop. Fl. Carn. ed. II. i. 416 = A. Chamaepitys.
decumbens, Mill. Gard. Dict. ed. VIII. n. 2 = A. reptans.
densiflora, Benth. in DC. Prod. xii. 595, in syn. = A. reptans.
genevensis, Crantz, Inst. i. 518 = A. genevensis.
nana, Gilib. Fl. Lituan. i. 94 = A. pyramidalis.
obliqua, Moench, Meth. 381 = A. orientalis.
orientalis, Crantz, Inst. i. 518 = A. orientalis.
pyramidalis, Crantz, l. c. = A. pyramidalis.
reptans, Crantz, l. c. = A. reptans.
tomentosa, Gilib. Fl. Lituan. i. 94 = A. genevensis.
villosa, Steud. Nom. ed. II. i. 235 = A. pyramidalis.

BUHSIA, Bunge, Del. Sem. Hort. Dorpat. (1859) 41.—Cf. Linnaea, xxx. (1859–60) 752 = **Cleome**, Linn. (Capparid.).
coluteoides, Bunge, ll. cc. = C. coluteoides.

BUINALIS, Rafin. New Fl. Am. iv. 40 (1836) = **Paronychia**, Linn. (Illecebrac.).
floridana, Rafin. l. c. = P. Baldwinii.

BUJACIA, E. Mey. Comm. Pl. Afr. Austr. 127 (1835) = **Glycine**, Linn. (Legumin.).
anonychia, E. Mey. l. c. = G. javanica.
gampsonychia, E. Mey. l. c. = Teramnus labialis.

BULBEDULIS, Rafin. Fl. Tellur. ii. 26 (1836) = **Camassia**, Lindl. (Liliac.).

BULBILIS, Rafin. in Am. Monthly Mag. (1819) 190, nomen prius; et in Journ. Phys. lxxxix. (1819) 262 = **Buchloë**, Engelm. (Gramin.).
dactyloides, Rafin. in Am. Monthly Mag. (1819) 190 = Buchloë dactyloides.

BULBILLARIA, Zucc. in Abh. Akad. Mnench. iii. (1843) 229. t. 2 = **Gagea**, Salisb. (Liliac.).
gageoides, Zucc. l. c. = G. persica.

BULBINE, Linn. Hort. Cliff. 122 (1737); Gen. Pl. ed. I. 95 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 784.
BLEPHANTHERA, Rafin. Fl. Tellur. ii. 59 (1836).
NEMOPOGON, Rafin. l. c. 27 (1836).
abyssinica, A. Rich. Tent. Fl. Abyss. ii. 334 = asphodeloides.
acaulis, Linn. Sp. Pl. ed. II. 446 = alooides.
alooides, *Willd. Enum. Hort. Berol.* 372.—Afr. austr.
annua, *Willd. l. c.*—Afr. austr.
asiatica, Gaertn. Fruct. i. 42. t. 13 = Crinum asiaticum.
asphodeloides, *Spreng. Syst.* ii. 85.—Afr. austr.
australis, Spreng. Syst. ii. 86 = bulbosa.
bisulcata, *Haw. in Phil. Mag.* (Feb. 1827) 121.—Afr. austr.
brevifolia, Schult. f. Syst. vii. 450 = Anthericum brevifolium.
bulbosa, *Haw. Rev. Pl. Succ.* 33.—Austral.
canaliculata, Spreng. Syst. ii. 86 = Anthericum canaliculatum.
Cauda-felis, Schult. f. Syst. vii. 450 = Bulbinella caudata.
caudata, Spreng. Syst. ii. 86 = Bulbinella caudata.
caulescens, Linn. ex Steud. Nom. ed. I. 125; ed. II. i. 236 = frutescens.
ciliata, Link, Enum. Hort. Berol. i. 329 = Anthericum ciliatum.
crispa, Spreng. Syst. ii. 86 = Chlorophytum crispum.
densiflora, *Baker, in Journ. Linn. Soc.* xv. (1876) 347.—Afr. austr.

BULBINE :—
dubia, *Schult. f. Syst.* vii. 450.—Afr. austr.
falcata, Schult. f. l. c. 451 = Anthericum falcatum.
favosa, *Schult. f. l. c.* 444.—Afr. austr.
filifolia, *Baker, in Journ. Linn. Soc.* xv. (1876) 344. —Afr. austr.
floribunda, Sehrad. ex Benth. Fl. Austral. vii. 35 = semibarbata.
Fraseri, Kunth, Enum. Pl. iv. 565 = bulbosa.
frutescens, *Willd. Enum. Hort. Berol.* 372.—Afr. austr.
graminea, Haw. Revis. Pl. Succ. 33 = asphodeloides.
hispida, Sweet, Hort. Brit. ed. I. 416 = Anthericum hispidum.
Hookeri, Kunth, Enum. Pl. iv. 566 = bulbosa.
incurva, Spreng. Syst. ii. 86 = rostrata.
latifolia, *Spreng. l. c.*—Afr. anstr.
laxiflora, *Baker, in Journ. Linn. Soc.* xv. (1876) 347. —Afr. austr.
longiscapa, *Willd. Enum. Hort. Berol.* 372.—Afr. austr.
Mackenii, Hook. f. Bot. Mag. t. 5955 = Eriospermum Mackenii.
macrophylla, Salm-Dyck, Hort. Dyck. 333 = latifolia.
mesembryanthemoides, *Haw. in Phil. Mag.* (July 1825) 31.—Afr. anstr.
Mettinghii, Tenore, Syll. 563 = longiscapa.
minima, *Baker, in Journ. Linn. Soc.* xv. (1876) 344. —Afr. austr.
narcissifolia, *Salm-Dyck, Hort. Dyck.* 334.—Afr. austr.
nutans, *Spreng. Syst.* ii. 86.—Afr. austr.
nutans, Zeyh. ex Baker, in Journ. Linn. Soc. xv. (1876) 347 = Zeyheri.
pallida, *Baker, in Journ. Bot.* xiv. (1876) 184.—Afr. austr.
planifolia, Spreng. Syst. ii. 86 = Simethis bicolor.
praemorsa, *Spreng. l. c.*—Afr. anstr.
pugioniformis, *Link, Enum. Hort. Berol.* i. 389.—Afr. austr.
rostrata, *Willd. Enum. Hort. Berol.* 372.—Afr. austr.
scabra, Spreng. Syst. ii. 86 = Anthericum scabrum.
semibarbata, *Haw. Revis. Pl. Succ.* 33.—Austral.
semibarbata, Hook. Bot. Mag. t. 3129 = bulbosa.
suavis, Lindl. Bot. Reg. (1838) Misc. 45 = bulbosa.
subbarbata, Steud. Nom. ed. II. i. 74, sphalm. = bulbosa.
triquetra, Spreng. Syst. ii. 86 = Bulbinella triquetra.
uncinata, Moench, Meth. 641 = Crinum americanum.
undulata, *Spreng. Syst.* ii. 86 = Anthericum undulatum.
unifolia, Schult. f. ex Baker, in Journ. Linn. Soc. xv. (1876) 263 = Eriospermum abyssinicum.
urgineoides, *Baker, l. c.* 348.—Afr. austr.
Zeyheri, *Baker, l. c.* 347.—Afr. austr.

BULBINELLA, Kunth, Enum. Pl. iv. 569 (1843). *LILIACEAE*, Benth. & Hook. f. iii. 784.
CHRYSOBACTRON, Hook. f. Fl. Antarct. i. 72. t. 44 (1845).
brevifolia, Kunth, l. c. 573 = Anthericum brevifolium.
Burkei, *Benth. & Hook. f. Gen.* iii. 784.—Afr. austr.
capillaris, Kunth, Enum. Pl. iv. 572 = triquetra.
caudata, *Kunth, l. c.*—Afr. austr.
ciliolata, *Kunth, l. c.* 570.—Afr. austr.
filiformis, Kunth, l. c. 572 = triquetra.
gracilis, *Kunth, l. c.* 571.—Afr. austr.
latifolia, *Kunth, l. c.* 572.—Afr. anstr.
ornithogaloides, *Kunth, l. c.* 693.—Afr. austr.
peronata, *Kunth, l. c.* 570.—Afr. austr.
robusta, *Kunth, l. c.* 571.—Afr. austr.
setifolia, Kunth, l. c. 569 = triquetra.
squamea, Kunth, l. c. 573 = Anthericum hispidum.
triquetra, *Kunth, l. c.*—Afr. austr.

BULBISPERMA, Reinw. ex Blume, Cat. Gew. Buitenz. 59 [1823] = **Peliosanthes**, Andr. (Haemod.).
ovigera, Reinw. l. c. = P. javanica.

BULBOCAPNOS, Bernh. in Linnaea, viii. (1833) 469 = **Corydalis**, Vent. (Papaverac.).
bracteatus, Bernh. l. c. = C. bracteata.
cavus, Bernh. l. c. = C. cava.
digitatus, Bernh. l. c. = C. Halleri.
fabaceus, Bernh. l. c. = C. fabacea.
Halleri, Spach. Hist. Vég. Phan. vii. 77 = C. Halleri.
longiflorus, Bernh. in Linnaea, viii. (1833) 469 = C. longiflora.

BULBOCAPNOS:—
Marschallianus, Bernh. in Linnaea, viii. (1833) 469 = C. Marschalliana.
nobilis, Bernh. l. c. = C. nobilis.
tuberosus, Spach, Hist. Vég. Phan. vii. 78 = C. cava.

BULBOCASTANUM, Lag. Amen. Acad. Madr. 99 (1821) = **Conopodium**, Koch (Umbellif.).
majus, Lag. l. c. = C. denudatum.

BULBOCASTANUM, Schur, Enum. Pl. Transs. 249 (1866) = **Carum**, Rupp. (Umbellif.).
daucoides, Nym. Consp. 304 = C. daucoides.
ferulaceum, Nym. l. c. 303.—Europ.
incrassatum, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 88 = C. incrassatum.
Linnaei, Schur, Enum. Pl. Transs. 249 = C. Bulbocastanum.
mauritanicum, Willk. in Linnaea, xl. (1876) 84 = C. mauritanicum.

BULBOCODIUM, Linn. Coroll. Gen. 6 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 821.
ABANDIUM, Adans. Fam. ii. 54 (1763).
alpinum, Mill. Gard. Dict. ed. VIII. n. 1 = Lloydia alpina.
atticum, Nym. Syll. 379 = Merendera attica.
antumnale, Lapeyr. Hist. Abr. Pl. Pyr. 202 = Merendera Bulbocodium.
balearicum, Nym. Syll. 379 = Merendera filifolia.
Broteri, Welw. ex Baker, in Journ. Linn. Soc. xvii. (1880) 441 = Merendera Bulbocodium.
caucasicum, Endl. ex Heynh. Nom. ii. 81 = Merendera caucasica.
colchicoides, Nym. Syll. 379 = Merendera Bulbocodium.
dentatum, *Schur, in Verh. Siebenb. Ver. Naturw.* ii. (1851) 166.—Transsylv.
edentatum, Schur, in Oestr. Bot. Zeitschr. x. (1860) 355 = ruthenicum.
Eichleri, Regel, in Act. Hort. Petrop. v. (1877) 638 = Merendera Eichleri.
Fontanesii, *Heynh. Nom.* ii. 81.—Ins. Majorc.; Algeria.
hastulatum, Friwald. in Flora, xix. (1836) 434 = Merendera sobolifera.
lusitanicum, *Heynh. Nom.* ii. 81.—Lusitania.
montanum, Fisch. Ind. Pl. Hort. Gorenk. (1812) 10 = ruthenicum.
persicum, *Regel, Gartenfl.* xxx. (1881) 294. t. 1058.—Persia.
robustum, *Bunge, ex Regel, in Act. Hort. Petrop.* vi. (1879) 491.—Turkestan.
ruthenicum, *Bunge, in Del. Sem. Hort. Dorpat.* (1837) 11.—Transylv.; Reg. Caucas.
serotinum, Linn. Sp. Pl. 294 = Lloydia alpina.
soboliferum, Heynh. Nom. ii. 81 = Merendera sobolifera.
trigynum, Nym. Syll. 379 = Merendera sobolifera.
trigynum, Adam, in Weber & Mohr, Cat. i. 49; ex Schult. f. Syst. vii. 1524 = Merendera caucasica.
trigynum, Griseb. Spicil. Fl. Rumel. ii. 380 = Merendera sobolifera.
vernum, Bieb. Fl. Taur. Cauc. i. 261 = ruthenicum.
vernum, Desf. Fl. Atlant. i. 284 = Merendera filifolia.
vernum, Linn. Sp. Pl. 294 = ruthenicum.
versicolor, *Spreng. Syst.* ii. 40.—Rossia austr.

BULBOPHYLLARIA, S. Moore, in Baker, Fl. Maurit. 344 (1877), sphalm. = BOLBOPHYLLARIA, Reichb. f. = **Bulbophyllum**, Thou. (Orchid.).

BULBOPHYLLUM, Thou. Orch. Iles Afr. Tab. Esp. 3 et Ic. t. 93–97 (1822). *ORCHIDEAE*, Benth. & Hook. f. iii. 501.
ANISOPETALUM, Hook. Exot. Fl. t. 149 (1825).
BOLBOPHYLLARIA, Reichb. f. in Bot. Zeit. x. (1852) 934.
BOLBOPHYLLUM, Spreng. Syst. iii. 681 (1826).
COCHLIA, Blume, Bijdr. 320. t. 59 (1825).
DIDACTYLE, Lindl. Fol. Orch. (1852).
DIPHYES, Blume, Bijdr. 310. t. 66 (1825).
EPICRANTHES, Blume, l. c. 306. t. 9 (1825).
EPICRIANTHES, Blume, Fl. Jav. Praef. p. vii (1828).
GERSINIA, Neraud, in Gaudich. Bot. Freyc. Voy. 27 (1826).
IONE, Lindl. Fol. Orch. (1853).
LYRAEA, Lindl. Gen. et Sp. Orch. 46 (1830).
MACROLEPIS, A. Rich. Sert. Astrol. 25. t. 19 (1833).
MALACHADENIA, Lindl. Bot. Reg. (1839) Misc. 67.

BULBOPHYLLUM:—
ODONTOSTYLES, Breda, Orch. Kuhl et Hass. t. sine n. (1827).
OXYSEPALA, Wight, Ic. v. 17. t. 1736 (1852).
SARCOBODIUM, Beer, Prakt. Orch. 306 (1854).
SESTOCHILOS, Breda, Orch. Kuhl et Hass. t. sine n. (1827).
TAUROSTALIX, Reichb. f. in Bot. Zeit. x. (1852) 933.
TRIBRACHIA, Lindl. Bot. Reg. sub t. 832 (1824).
XIPHIZUSA, Reichb. f. in Bot. Zeit. x. (1852) 919.
abbreviatum, Reichb. f. in Gard. Chron. (1881) II. 70 = Cirrhopetalum abbreviatum.
acutiflorum, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 18 (Cirrhopetali sp.).—Ind. or.
adenopetalum, *Lindl. Bot. Reg.* (1842) *Misc.* 85.—Malaya.
affine, *Lindl. in Wall. Cat.* n. 1982.—Reg. Himal.
alcicorne, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 151.—Burma.
alopecurum, *Reichb. f. in Gard. Chron.* (1880) II. 70.—Burma.
amplebracteatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 307.—Malaya.
amplum, Reichb. f. in Walp. Ann. vi. 244 = Dendrobium amplum.
Andersonii, *Kurz, Rep. Andam.* 51, nomen.—Ins. Andam.
andongense, *Reichb. f. in Flora*, xlviii. (1865) 184.—Afr. occ.
angustifolium, *Lindl. Gen. et Sp. Orch.* 57.—Java.
antenniferum, *Reichb. f. in Walp. Ann.* vi. 250.—Ins. Philipp.
antenniferum, Reichb. f. l. c. 248 = geraënse.
apetalum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 127.—Afr. trop.
apiferum, *Batem. ex Steud. Nom.* ed. II. i. 241.—Afr. trop.
argyropus, *Reichb. f. in Linnaea*, xli. (1877) 42.—Ins. Norfolc.
aristatum, *Hemsl. Biol. Centr. Am. Bot.* iii. 213.—Am. centr.
atropurpureum, *Rodrig. Orch. Nov.* i. 41.—Bras.
aurantiacum, *F. Muell. Fragm.* iii. 39.—Austral.
aurantiacum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 219.—Afr. trop.
auratum, *Reichb. f. in Walp. Ann.* vi. 261.—Manilla.
auricomum, *Lindl. in Wall. Cat.* n. 1985.—Burma.
Baileyi, *F. Muell. Fragm.* ix. 5.—Austral.
balaeniceps, Reichb. f. in Hamb. Gartenz. xix. (1863) 208 = Napelli.
barbatum, *Rodrig. Orch. Nov.* ii. 119.—Bras.
barbigerum, *Lindl. Bot. Reg.* xxiii. (1837) t. 1942.—Afr. trop.
Baronii, *Ridl. in Journ. Linn. Soc.* xxi. (1885) 463.—Madag.
Beccarii, *Reichb. f. in Gard. Chron.* (1879) I. 41.—Borneo.
Berenicis, *Reichb. f. l. c.* xiv. (1880) II. 588.—Hab.?
Betchei, *F. Muell. in Wing, South. Sc. Record*, i. (1881) 173.—Ins. Samoa.
bicolor, *Lindl. Gen. et Sp. Orch.* 49.—China.
bifarium, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 219.—Afr. trop.
biflorum, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 397.—Java.
bisetum, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 186.—Reg. Himal.
Blepharistes, *Reichb. f. in Flora*, lv. (1872) 278 (Cirrhopetali sp.).—Burma.
Boukettae, *F. M. Bailey, in Proc. Roy. Soc. Queensl.* i. (1884) 89.—Austral.
bootanense, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 153 = Cirrhopetalum bootanense.
Bowringianum, *Reichb. f. in Gard. Chron.* (1881) I. 814.—Assam.
braccatum, *Reichb. f. in Linnaea*, xli. (1877) 117.—Ins. Philipp.
bracteolatum, *Lindl. Bot. Reg.* xxiv. (1838) 57.—Guiana.
Bufo, Reichb. f. in Walp. Ann. vi. 258 = Megaclinium Bufo.
caespitosum, *Thou. Orch. Afr.* t. 103.—Ins. Borbon.
calamarium, *Lindl. Bot. Reg.* (1843) *Misc.* 70.—Afr. trop.
capillipes, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 150.—Burma.
capitatum, *Lindl. Gen. et Sp. Orch.* 56.—Malaya.

BULBOPHYLLUM:—
Careyanum, *Spreng. Syst.* iii. 732.—Reg. Himal.
cariniflorum, Reichb. f. in Walp. Ann. vi. 253 = bisetum.
caudatum, *Lindl. in Wall. Cat.* n. 1983.—Reg. Himal.
cernuum, *Lindl. Gen. et Sp. Orch.* 48.—Java.
Cheiri, *Lindl. Bot. Reg.* (1844) *Misc.* 44.—Manilla.
chinense, *Reichb. f. in Walp. Ann.* vi. 260.—China.
chloroglossum, *Reichb. f. in Gard. Chron.* (1871) 1195.—Bras.
chloropterum, *Reichb. f. in Linnaea*, xxii. (1849) 835.—Bras.
ciliatum, *Lindl. Gen. et Sp. Orch.* 48.—Java.
cirrhopetaloides, Griff. Notul. iii. 290 = bisetum.
clandestinum, *Lindl. Bot. Reg.* (1841) *Misc.* 77.—Ind. or.; Malaya.
Clausseni, *Reichb. f. in Linnaea*, xix. (1847) 371.—Bras.
clavatum, *Thou. Orch. Afr.* t. 99.—Ins. Mascar.
clavigerum, *F. Muell. Census, Suppl.* i. 3.—Austral.
cochleatum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 125.—Afr. trop.
cocoinum, *Batem. ex Lindl. Bot. Reg.* xxiii. (1837) t. 1964.—Afr. trop.
colubrinum, Reichb. f. in Walp. Ann. vi. 257 = Megaclinium colubrinum.
comatum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 128.—Afr. trop.
Commersonii, *Thou. Orch. Afr.* t. 97.—Ins. Borbon.
compressum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 307.—Sumatra.
conchiferum, *Reichb. f. in Walp. Ann.* vi. 253.—Reg. Himal.
conicum, Thou. Orch. Afr. Tab. Esp. 3 = clavatum.
conitum, Thou. l. c. t. 100 = clavatum.
cornutum, Reichb. f. in Walp. Ann. vi. 247 = Cirrhopetalum cornutum.
coriscense, *Reichb. f. Otia Bot. Hamb.* 118.—Afr. trop.
crassifolium, *Thw. ex Trimen, in Journ. Bot.* xxiii. (1885) 244.—Zeylan.
croceum, *Lindl. Gen. et Sp. Orch.* 57.—Java.
Cumingii, *Reichb. f. in Walp. Ann.* vi. 261.—Ins. Philipp.
cupreum, Hook. Bot. Mag. t. 5316 = Careyanum.
cupreum, *Lindl. Bot. Reg.* (1838) *Misc.* 95.—Burma.
cuspidilingua, *Reichb. f. in Walp. Ann.* vi. 264.—Ins. Philipp.
cylindraceum, *Lindl. Gen. et Sp. Orch.* 53.—Reg. Himal.
Dayanum, *Reichb. f. in Gard. Chron.* (1865) 434.—Burma.
delitescens, Hance, in Journ. Bot. xiv. (1876) 44 = Cirrhopetalum delitescens.
densum, *Thou. Orch. Afr.* t. 108.—Ins. Mascar.; Madag.
distans, *Lindl. in Journ. Linn. Soc.* vi. (1862) 125.—Afr. trop.
Drallei, *Reichb. f. in Regel, Gartenfl.* (1885) 20.—Afr. trop.
elaidium, *Lindl. in Journ. Linn. Soc.* vi. (1862) 127.—Afr. trop.
elegans, *Gardn. ex Thw. Enum. Pl. Zeyl.* 298.—Zeylan.
Elisae, *F. Muell. Fragm.* vi. 120.—Austral.
Elliae, Reichb. f. in Walp. Ann. vi. 263 = Cirrhopetalum Wightii.
elongatum, *Hassk. Cat. Hort. Bog. Alt.* 39.—Malaya.
epiphytum, *Rodrig. Orch. Nov.* i. 40.—Bras.
erectum, *Thou. Orch. Afr.* t. 96.—Madag.
eublepharum, *Reichb. f. in Walp. Ann.* vi. 252.—Reg. Himal.
exaltatum, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 186.—Guiana.
exiguum, *F. Muell. Fragm.* ii. 72.—Austral.
falcatum, Reichb. f. in Walp. Ann. vi. 258 = Megaclinium falcatum.
falcipetalum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 128.—Afr. trop.
fimbriatum, Reichb. f. in Walp. Ann. vi. 260 = Cirrhopetalum fimbriatum.
flavescens, *Lindl. Gen. et Sp. Orch.* 54.—Java.
flavidum, *Lindl. Bot. Reg.* (1840) *Misc.* 83.—Afr. trop.
foenisecii, Par. ex Reichb. f. in Bot. Zeit. xxiii. (1865) 99 = auricomum.
fuscescens, Reichb. f. Walp. Ann. vi. 244 = Dendrobium fuscescens.
fusco-purpureum, *Wight, Ic.* v. t. 1651.—Ind. or.
fuscum, *Lindl. Bot. Reg.* (1839) *Misc.* 3.—Afr. trop.

BULBOPHYLLUM:—
Gabonis, *Linden & Reichb. f. in Flora*, xlviii. (1865) 185.—Afr. occ.
galeatum, *Lindl. Gen. et Sp. Orch.* 57.—Afr. trop.
geraënse, *Reichb. f. ex Rodrig. Orch. Nov.* ii. 125.—Bras.
gibbosum, *Lindl. Gen. et Sp. Orch.* 54.—Java.
gladiatum, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 185.—Bras.
gracile, *Lindl. Gen. et Sp. Orch.* 50.—Java.
gracile, Thou. Orch. Afr. t. 101 = Thouarsii.
gracile, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 152.—Burma.
grandiflorum, *Blume, Rumphia*, iv. 42.—N. Guinea.
grandiflorum, Griff. Notul. iii. 293 = reptans.
granulosum, *Rodrig. Orch. Nov.* i. 41.—Bras.
gravidum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 126.—Afr. trop.
Griffithianum, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 153 = Cirrhopetalum gamosepalum.
Griffithii, *Reichb. f. in Walp. Ann.* vi. 247.—Reg. Himal.
Henshallii, Hort. ex Gard. Chron. (1852) 422 = Lobbii.
herminiostachys, *Reichb. f. in Walp. Ann.* vi. 254.—Afr. trop.
Hildebrandtii, *Reichb. f. Otia Bot. Hamb.* 74; *et in Bot. Zeit.* xxxix. (1881) 449.—Madag.
hirsutum, *Lindl. Gen. et Sp. Orch.* 49.—Java.
hirtum, *Lindl. in Wall. Cat.* n. 1989.—Reg. Himal.
imbricatum, Griff. Notul. iii. 289 = cylindraceum.
imbricatum, *Lindl. Bot. Reg.* (1841) *Misc.* 37.—Afr. trop.
inaequale, *Lindl. Gen. et Sp. Orch.* 49.—Java.
incurvum, *Thou. Orch. Afr.* t. 95.—Ins. Mascar.; Madag.
iners, *Reichb. f. in Gard. Chron.* (1880) i. 776.—Assam?
inops, *Reichb. f. l. c.* (1880) ii. 620.—Hab.?
insectiferum, *Rodrig. Orch. Nov.* ii. 120.—Bras.
intertextum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 127.—Afr. trop.
javanicum, *Miq. Fl. Ind. Bat.* iii. 650.—Java.
kaitiense, Reichb. f. in Walp. Ann. vi. 262 = Cirrhopetalum nilgherense.
khasyanum, Griff. Notul. iii. 284 = cylindraceum.
khasyanum, Reichb. f. in Trans. Linn. Soc. xxx. (1874) 138 = repens.
lasianthum, *Lindl. in Gard. Chron.* (1855) 53.—Sumatra.
lasiochilum, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 153.—Burma.
laxiflorum, *Lindl. Gen. et Sp. Orch.* 57.—Java.
lemniscatum, *Par. ex Hook. f. Bot. Mag.* t. 5961.—Burma.
leopardinum, *Lindl. in Wall. Cat.* n. 1981.—Reg. Himal.
lichenastrum, *F. Muell. Fragm.* vii. 60.—Austral.
limbatum, *Lindl. Bot. Reg.* (1840) *Misc.* 74.—Singapora.
limbatum, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 152 (non Lindl.).—Burma.
Lindleyanum, *Griff. Notul.* iii. 287.—Burma.
Lobbii, *Lindl. Bot. Reg.* (1847) t. 29.—Bnrma; Malaya.
longiflorum, Thou. Orch. Afr. t. 98 = Cirrhopetalum Thouarsii.
longipes, *Reichb. f. in Walp. Ann.* vi. 253.—Burma.
Luckraftii, *F. Muell. in Wing, South. Sc. Record*, ii. (1882) 71.—Ins. Solomon.
Lundianum, *Reichb. f. & Warm. in Otia Bot. Hamb.* 94.—Bras.
lupulinum, *Lindl. in Journ. Linn. Soc.* vi. (1862) 126.—Afr. trop.
Macraei, Reichb. f. in Walp. Ann. vi. 263 = Cirrhopetalum Wightii.
macranthum, *Lindl. Bot. Reg.* (1844) t. 13.—Singapora.
macrocceras, *Rodrig. Orch. Nov.* ii. 118.—Bras.
maculatum, *Boxall, ex Naves, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 235.—Ins. Philipp.
Makoyanum, *Reichb. f. in Gard. Chron.* (1879) i. 234 in nota.—Bras.
mandibulare, *Reichb. f. l. c.* (1882) i. 366.—Borneo.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 218.—Afr. trop.
Mannii, Reichb. f. in Flora, lv. (1872) 275 (= Cirrhopetali sp.).—Reg. Himal.
maxillare, *Reichb. f. in Walp. Ann.* vi. 248.—Ins. Philipp.

BULBOPHYLLUM:—
maximum, Reichb. f. in Walp. Ann. vi. 259 = Megaclinium maximum.
Medusae, *Reichb. f. in Walp. Ann.* vi. 262.—Malaya.
megalanthum, *Griff. Notul.* iii. 286.—Malaya.
megalonyx, *Reichb. f. Otia Bot. Hamb.* 74; *et in Bot. Zeit.* xxxix. (1881) 449.—Ins. Comor.
melanorrhachis, Reichb. f. in Gard. Chron. (1875) ii. 162 = Megaclinium melanorrhachis.
Melleri, Reichb. f. in Flora, lxi. (1878) 78 = Megaclinium Melleri.
membranaceum, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 397.—Java.
mentosum, *Rodrig. Orch. Nov.* i. 42.—Bras.
merguense, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 154 (= Cirrhopetali sp.).—Burma.
meridense, *Reichb. f. in Linnaea*, xxii. (1849) 836.—Venezuela.
micranthum, *Rodrig. Orch. Nov.* i. 39.—Bras.
micropetalum, *Lindl. in Journ. Linn. Soc.* vi. (1861) 127.—Afr. trop.
micropetalum, *Rodrig. Orch. Nov.* ii. 121.—Bras.
microtepalum, *Reichb. f. in Walp. Ann.* vi. 253.—Burma.
minutum, *Thou. Orch. Afr.* t. 110.—Madag.
minutissimum, *F. Muell. Fragm.* xi. 53.—Austral.
moniliforme, *F. Muell. in Gard. Chron.* (1878) ii. 818.—Austral.
moniliforme, R. King, in Gard. Chron. (1879) i. 55, 149 = minutissimum.
moniliforme, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 151.—Burma.
monosepalum, *Rodrig. Orch. Nov.* ii. 121.—Bras.
monticolum, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 219.—Afr. trop.
moulmeinense, Reichb. f. in Walp. Ann. vi. 249 = Trias picta.
mucronatum, *Lindl. Gen. et Sp. Orch.* 50.—Java.
mucronifolium, *Reichb. f. & Warm. in Otia Bot. Hamb.* 94.—Bras.
multiflorum, *Ridley, in Journ. Linn. Soc.* xxi. (1885) 463.—Madag.
muscicolum, Reichb. f. in Flora, lv. (1872) 275 = Cirrhopetalum Wallichii.
mutabile, *Lindl. Gen. et Sp. Orch.* 48.—Java.
Napelli, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 185.—Bras.
nasutum, *Reichb. f. in Gard. Chron.* (1871) 148.—Burma.
neilgherrense, *Wight, Ic.* v. 1, t. 1680.—Ind. or.
nematopodum, *F. Muell. Fragm.* viii. 30.—Austral.
nutans, *Thou. Orch. Afr.* t. 107.—Ins. Mascar.; Madag.
nutans, *Reichb. f. in Walp. Ann.* vi. 260.—Ins. Philipp.
nutans, Thou. Orch. Afr. 63. t. 8. n. 2 = variegatum.
oblongum, Reichb. f. in Walp. Ann. vi. 249 = Trisa oblonga.
obtusum, *Lindl. Gen. et Sp. Orch.* 56.—Java.
occidentale, Spreng. Syst. iii. 732 = Epidendrum polybulbon.
occlusum, *Ridley, in Journ. Linn. Soc.* xxi. (1885) 464.—Madag.
occultum, *Thou. Orch. Iles Afr.* t. 93.—Ins. Mascar.
oculatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 309.—Malaya.
odoratissimum, *Lindl. in Wall. Cat.* n. 1987.—Reg. Himal.; Burma.
odoratum, *Lindl. Gen. et Sp. Orch.* 54.—Java.
Oerstedii, *Hemsl. Biol. Centr. Am. Bot.* iii. 213.—Nicaragua.
oligoglossum, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 297.—Burma.
oreonastes, *Reichb. f. Otia Bot. Hamb.* 118.—Afr. trop.
ovalifolium, *Lindl. Gen. et Sp. Orch.* 49.—Java.
oxypterum, Reichb. f. in Walp. Ann. vi. 258 = Megaclinium oxypterum.
pachyrrhachis, *Griseb. Fl. Brit. W. Ind.* 613.—Ind. occ.
Pahudi, *Reichb. f. in Walp. Ann.* vi. 264.—Malaya.
palcaceum, *Benth. & Hook. f. Gen.* iii. 503.—Reg. Himal.
Pangerangi, *Reichb. f. in Bonplandia*, v. (1857) 57.—Malaya.
parviflorum, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 152.—Burma.
parvulum, *Lindl. Gen. et Sp. Orch.* 49.—Java.

BULBOPHYLLUM:—
pavimentatum, *Lindl. in Journ Linn. Soc.* vi. (1861) 128.—Afr. trop.
pendulum, *Thou. Orch. Afr.* t. 104.—Ins. Mascar.
Penicillium, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 151.—Burma.
petiolare, *Thw. Enum. Pl. Zeyl.* 298.—Zeylan.
pictum, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 150 = Drymoda picta.
picturatum, Reichb. f. in Walp. Ann. vi. 262 = Cirrhopetalum picturatum.
pileatum, *Lindl. Bot. Reg.* (1844) *Misc.* 73.—Malaya.
Pipio, *Reichb. f. in Linnaea*, xli. (1877) 92.—Afr. trop.
polyrrhizum, *Lindl. Gen. et Sp. Orch.* 53.—Reg. Himal.
Prenticei, *F. Muell. in Wing, South. Sc. Record*, i. (1881) 173.—Austral.
prismaticum, *Thou. Orch. Afr.* t. 102.—Ins. Mascar.
psittacoglossum, *Reichb. f. in Bot. Zeit.* xxi. (1863) 236.—Burma.
Psychoon, *Reichb. f. in Gard. Chron.* (1878) ii. 170.—Assam.
Pumilio, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 153 = Cirrhopetalum Pumilio.
pumilum, *Lindl. Gen. et Sp. Orch.* 54.—Afr. trop.
punctatum, *Fitzgerald, in Journ. Bot.* xxi. (1883) 205.—Austral.
punctatum, *Rodrig. Orch. Nov.* i. 40.—Bras.
purpurascens, *F. M. Bailey, in Proc. Roy. Soc. Queensl.* i. (1884) 88.—Austral.
purpurascens, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 308.—Java.
purpureum, *Thw. Enum. Pl. Zeyl.* 298.—Zeylan.
pusillum, *Thou. Orch. Afr.* t. 102.—Ins. Maurit.
pygmaeum, *Lindl. Gen. et Sp. Orch.* 58.—N. Zeland.
quadrisetum, *Lindl. in Hook. Lond. Journ. Bot.* ii. (1843) 670.—Guiana.
radiatum, *Lindl. in Wall. Cat.* n. 1986.—Bnrma; China.
recurvum, *Lindl. Gen. et Sp. Orch.* 53.—Afr. trop.
refractum, Reichb. f. in Walp. Ann. vi. 259 = Cirrhopetalum refractum.
Regnelli, *Reichb. f. in Linnaea*, xxii. (1849) 835.—Bras.
Reinwardtii, *Reichb. f. in Walp. Ann.* vi. 246.—Malaya.
repens, *Griff. Notul.* iii. 293.—Reg. Himal.
reptans, *Lindl. in Wall. Cat.* n. 1988.—Reg. Himal.
reticulatum, *Batem. in Hook. f. Bot. Mag.* t. 5605.—Borneo.
retusiusculum, Reichb. f. in Gard. Chron. (1869) 1182 = Cirrhopetalum retusiusculum.
Rhizophorae, *Lindl. in Journ. Linn. Soc.* vi. (1861) 125.—Afr. trop.
rigens, *Reichb. f. in Hamb. Gartenz.* xxi. (1865) 296.—Burma.
rostriceps, *Reichb. f. Otia Bot. Hamb.* 55.—Ins. Fiji.
rotundatum, Reichb. f. in Walp. Ann. vi. 244 = Dendrobium rotundatum.
Roxburghii, Reichb. f. l. c. 263 = Cirrhopetalum Roxburghii.
rufinum, *Reichb. f. Xenia Orch.* iii. 45.—Burma.
rupicola, *Rodrig. Orch. Nov.* i. 39.—Bras.
rupincola, *Reichb. f. in Flora*, xlviii. (1865) 184.—Afr. trop.
salaccense, *Reichb. f. in Bonplandia*, v. (1857) 57.—Java.
saltatorium, *Lindl. Bot. Reg.* xxiii. (1837) t. 1970.—Afr. trop.
Sandersoni, Reichb. in Flora, lxi. (1878) 78 = Megaclinium Sandersoni.
sarcoscapum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 242.—Java.
scabratum, Reichb. f. in Walp. Ann. vi. 259 = Cirrhopetalum caespitosum.
Sceptrum, *Reichb. f. l. c.* 254.—Ins. Sunda.
Schillerianum, Reichb. f. in Hamb. Gartenz. xvi. (1866) 423 = Shepherdi.
Schmidtianum, *Reichb. f. l. c.* xxii. (1865) 357.—Ind. or.
sciadanthum, *F. Muell. in Wing, South. Sc. Record.* ii. (1882) 95.—Ins. Samoa.
serpens, Lindl. Gen. et Sp. Orch. 57 = nutans.
setigerum, *Lindl. Bot. Reg.* (1838) *Misc.* 21.—Guiana.
Seychellarum, *Reichb. f. in Linnaea*, xli. (1877) 93.—Ins. Mascar.
Shepherdi, *F. Muell. Fragm.* iii. 40.—Austral.

BULBOPHYLLUM :—
siamense, Reichb. f. in Gard. Chron. (1867) 572 = Lobbii.
sicyobulbon, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 152.—Burma.
Sillemianum, *Reichb. f. in Gard. Chron.* (1884) II. 166.—Burma.
simillinum, Par. & Reichb. f. Otia Bot. Hamb. 49 = Cirrhopetalum simillinum.
sordidum, Lindl. Bot. Reg. (1840) Misc. 89 = bracteolatum.
stenobulbon, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 153.—Burma.
striatum, *Reichb. f. in Walp. Ann.* vi. 257.—Reg. Himal.
suave, Griff. Notul. iii. 292 = hirtum.
sulcatum, *Lindl. Gen. et Sp. Orch.* 57.—Java.
taeniophyllum, Par. & Reichb. f. in Journ. Bot. xii. (1874) 198 (= Cirrhopetali sp).—Moulmein.
tahitense, *Nadeaud, Enum. Pl. Tahiti*, 36.—Ins. Pacif.
Taylori, *F. Muell. Fragm.* viii. 150.—Austral.
tenellum, *Lindl. Gen. et Sp. Orch.* 49.—Java.
tentaculigerum, Reichb. f. in Flora, lxi. (1878) 77 = Megaclinium tentaculigerum.
tenuicaule, *Lindl. in Journ. Linn. Soc.* vi. (1861) 126.—Afr. trop.
tenuifolium, *Lindl. Gen. et Sp. Orch.* 50.—Java.
tetragonum, *Lindl. l. c.* 53.—Afr. trop.
Thompsonii, *Ridley, in Journ. Linn. Soc.* xxi. (1885) 464.—Madag.
Thouarsii, *Steud. Nom.* ed. II. i. 214.—Ins. Maurit.
Thwaitesii, Par. & Reichb. f. in Journ. Bot. xii. (1874) 199 (= Cirrhopetali sp.).—Zeylan.
tigridum, *Hance, in Journ. Bot.* xxi. (1883) 232.—China.
tortuosum, *Lindl. Gen. et Sp. Orch.* 50.—Java.
tremulum, *Wight, Ic.* v. t. 1749.—Ind. or.
triadenium, *Reichb. f. in Walp. Ann.* vi. 257.—Java.
trigonopus, Reichb. f. in Gard. Chron. (1881) II. 71 = Cirrhopetalum abbreviatum.
tripetalum, *Lindl. in Ann. & Mag. Nat. Hist.* x. (1842) 185.—Bras.
tripudians, Par. & Reichb. f. in Trans. Linn. Soc. xxx. (1874) 154 = Cirrhopetalum refractum.
trisetosum, Griff. Notul. iii. 283 = clandestinum.
triste, *Reichb. f. in Walp. Ann.* vi. 253.—Burma.
tuberculatum, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 336.—N. Zel.
umbellatum, Lindl. in Wall. Cat. n. 1984 = Cirrhopetalum maculosum.
unguiculatum, *Reichb. f. in Linnaea*, xxii. (1849) 836.—Java.
uniflorum, *Griff. Notul.* iii. 293.—Reg. Himal.
uniflorum, *Hassk. Cat. Hort. Bogor. Alt.* 39.—Malaya.
vaginatum, Reichb. f. in Walp. Ann. vi. 261 = Cirrhopetalum vaginatum.
variegatum, *Thou. Orch. Afr.* t. 105, 106.—Ins. Borbon.
velutinum, Reichb. f. in Walp. Ann. vi. 258 = Megaclinium velutinum.
violaceum, *Lindl. Gen. et Sp. Orch.* 50.—Java.
vittatum, *Reichb. f. & Warm. in Otia Bot. Hamb.* 95.—Bras.
vittatum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 308.—Java.
Wallichii, Reichb. f. in Walp. Ann. vi. 259 = Cirrhopetalum Wallichii.
Weddelii, *Reichb. f. l. c.* 251.—Bras.
Wightii, Reichb. f. l. c. 262 = Cirrhopetalum grandiflorum.
xylophyllum, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 151.—Burma.

BULBOSPERMUM, Blume, Enum. Pl. Jav. 15 (1828) = **Peliosanthes**, Andr. (Haemodor.).
javanicum, Blume, l. c. = P. javanica.

BULBOSTYLIS, DC. Prod. v. 138 (1836) = **Brickellia**, Ell. (Compos.).
annua, Nutt. in Journ. Acad. Philad. Ser. II. i. (1847) 179 = Psathyrotes annua.
californica, Torr. & Gray, Fl. N. Am. ii. 79 = Brickellia californica
Cavanillesii, DC. Prod. v. 138 = Brickellia californica.
diffusa, DC. l. c. vii. 268 = Brickellia diffusa.
glabra, DC. l. c. v. 139 = Eupatorium Vauthierianum.
glabriuscula, C. Koch, in App. Sem. Hort. Berol. (1855) 13.—Am. trop.
hastata, Walp. Rep. Ind. p. 279 = Brickellia hastata.
hebecarpa, DC. Prod. v. 138 = Brickellia hebecarpa.

BULBOSTYLIS :—
lanata, DC. Prod. vii. 268 = Brickellia lanata.
microphylla, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 286 = Brickellia microphylla.
nepetaefolia, DC. Prod. v. 139 = Brickellia californica.
oliganthes, DC. l. c. = Brickellia oliganthes.
pauciflora, DC. l. c. = Eupatorium pauciflorum.
pedunculosa, DC. in Deless. Ic. Sel. iv. t. 17 = Eupatorium longipes.
pendula, DC. Prod. v. 138 = Brickellia pendula.
reticulata, DC. l. c. vii. 268 = Brickellia reticulata.
rigida, Hook. & Arn. Bot. Beech. Voy. 297 = Brickellia lanata.
scorodoniaefolia, Kunth, Ind. Sem. Hort. Berol. (1846) 12 = Brickellia californica ?
secundiflora, DC. Prod. v. 138 = Brickellia secundiflora.
spinaciaefolia, DC. l. c. 139 = Eupatorium spinaciaefolium.
subuligera, S. Schau. in Linnaea, xix. (1847) 718 = Barroetia subuligera.
triangularis, DC. Prod. vii. 268 = Carphephorus triangularis.
veronicaefolia, DC. l. c. v. 139 = Brickellia veronicaefolia.

BULBOSTYLIS, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 467 = **Eupatorium**, Tourn. (Compos.).
Cassiniana, Gardn. l. c. 471 = E. Vauthierianum.
deltoides, Buckl. in Proc. Acad. Sc. Philad. 1861 (1862) 456 = E. ageratifolium.
elegans, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 467 = E. elegans ?
glandulosa, Gardn. l. c. 469 = E. amygdalinum.
micrantha, Gardn. l. c. 449 = E. amygdalinum.
microcephala, Gardn. l. c. 468 = E. amygdalinum.
oblongifolia, Gardu. l. c. 469 = E. trixoides.
pumila, Gardn. l. c. 470 = E. amphidictyum.
ramosissima, Gardn. l. c. = E. ramosissimum.
scandens, Gardn. l. c. = E. Vitalbae.
tomentosa, Gardn. l. c. 468 = E. amygdalinum.

BULBOSTYLIS, Kunth, Enum. Pl. ii. 205 (1837) = **Fimbristylis**, Vahl (Cyperac.).
capillaris, Nees, in Mart. Fl. Bras. ii. 1. 84, in obs. = F. capillaris.
ciliata, Presl, ex Nees, l. c. 84, in obs.—Hab. ?
consanguinea, Nees, l. c.—Bras.
hirtella, Nees, l. c. 85.—Bras.
paradoxa, Nees, l. c. 82.—Bras.
tenuifolia, Nees, l. c. 83.—Bras.
trifida, Nees, l. c. 84, in obs. = F. capillaris.

BULBOSTYLIS, Stev. in Mém. Soc. Nat. Mosc. v. (1814) 355 = **Eleocharis**, R. Br. (Cyperac.).
capitatus, Stev. l. c. = E. capitata.
geniculatus, Stev. l. c. = E. geniculata.
maculosus, Stev. l. c. = E. maculosa.
ovatus, Stev. l. c. = E. ovata.
palustris, Stev. l. c. = E. palustris.
tuberculatus, Stev. l. c. = E. tuberculosa.

BULLIARDA, DC. in Bull. Soc. Philom. iii. (1801) n. 49. p. 1 = **Tillaea**, Mich. (Crassul.).
abyssinica, A. Rich. Tent. Fl. Abyss. i. 306 = T. aquatica.
alpina, Harv. in Harv. & Sond. Fl. Cap. ii. 330.—Afr. austr.
aquatica, DC. in Bull. Soc. Philom. (1801) n. 49. p. 2 = T. aquatica.
bonariensis, DC. l. c. = T. peduncularis.
brevifolia, Eckl. & Zeyh. Enum. 290.—Afr. austr.
capensis, DC. in Bull. Soc. Philom. (1801) n. 49. p. 2 = T. capensis.
decumbens, DC. l. c. = T. decumbens.
Dregei, Harv. in Harv. & Sond. Fl. Cap. ii. 330 = Crassula prostrata, *E. Mey.*
elatinoides, Eckl. & Zeyh. Enum. 290.—Afr. austr.
filiformis, Eckl. & Zeyh. l. c. = T. filiformis.
Linnaei, Spreng. Syst. i. 497 = T. aquatica.
magellanica, Comm. ex DC. in Bull. Soc. Philom. (1801) n. 49. p. 2 = T. moschata.
moschata, Urv. in Mém. Soc. Linn. Par. iv. (1826) 618 = T. moschata.
pedunculata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = T. Vaillantii.
perfoliata, DC. in Bull. Soc. Philom. (1801) n. 49. p. 2 = T. capensis.
prostrata, DC. l. c. = T. prostrata.

BULLIARDA :—
prostrata, Weinm. Fl. Petrop. 21 = T. aquatica.
recurva, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 472 [bis] = T. recurva.
Schkuhrii, Spreng. Syst. i. 498 = T. prostrata.
Sieberi, Spreng. Syst. iv. Cur. Post. 48 = T. verticillata.
trichotoma, Eckl. & Zeyh. Enum. 290 = T. decumbens.
Vaillantii, DC. in Bull. Soc. Philom. (1801) n. 49. p. 1 = T. Vaillantii.
Vaillantii, Schimp. ex Britten, in Oliver, Fl. Trop. Afr. ii. 388 = T. aquatica.

BULLIARDA, Neck. Elem. ii. 321 (1790) = **Unona**, Linn. f. et **Xylopia**, Linn. (Anonac.).

BULNESIA, C. Gay, Fl. Chil. i. 474. t. 15 (1846).
ZYGOPHYLLEAE, Benth. & Hook. f. i. 268.
GONOPTERA, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 150.
bonariensis, *Griseb. in Goett. Abh.* xix. (1874) 105.—Reg. Argent.
chilensis, *C. Gay, Fl. Chil.* i. 475.—Chili.
foliosa, *Griseb. in Goett. Abh.* xix. (1874) 106.—Reg. Argent.
macrocarpa, *Phil. Sert. Mend. Alt.* 9.—Chili.
Retamo, *Griseb. in Goett. Abh.* xix. (1874) 106.—Reg. Argent.
Sarmienti, *Lorentz, ex Griseb. l. c.* xxiv. (1879) 75.—Reg. Argent.
Schickendanzii, *Hieron. ex Griseb. l. c.*—Reg. Argent.

BULOWIA, Hook. Bot. Mag. sub t. 4364 (1848) = BUELOWIA, Schur = **Smeathmannia**, Soland. (Passifl.).

BULWERIA, F. Muell. Fragm. iv. 147 (1864) = **Diplanthera**, R. Br. (Bignon.).
nobilissima, F. Muell. l. c. = D. tetraphylla.

BUMALDA, Thunb. Nov. Gen. 62 (1783) = **Staphylea**, Linn. (Sapind.).
trifolia, Thunb. l. c. 63 = S. Bumalda.

BUMELIA, Sw. Prod. Veg. Ind. Occ. 49 (1788).
SAPOTACEAE, Benth. & Hook. f. ii. 660.
SCLEROCLADUS et SPONDOGONA, Rafin. Sylva Tellur. 35 (1838).
ambigua, Tenore, ex A. DC. in DC. Prod. v. 189 = lycioides.
angustifolia, *Nutt. Sylv. Am.* iii. 38. t. 93.—Am. bor.; Mexic.
arachnoidea, Rafin. New Fl. Am. iii. 28 = lanuginosa.
arborea, Buckl. in Proc. Ac. Sc. Philad. 1861 (1862) 461 = lanuginosa.
argentea, Roem. & Schult. in Ges. Naturf. Fr. Berl. Mag. iii. (1809) 57; Syst. iv. 499 = Heeria argentea.
Auzuba, *Roem. & Schult. Syst.* iv. 499.—Ins. S. Doming.
borbonica, *Lodd. ex Loud. Hort. Brit.* 69.—Ins. Borbon.
buxifolia, Roem. & Schult. Syst. iv. 802 = obtusifolia.
buxifolia, *Willd. ex H. B. & K. Nov. Gen. et Sp.* vii. t. 147.—Am. austr.
celastrina, *H. B. & K. l. c.* 212.—Mexic.
chrysophylloides, Wats. Dendrol. i. 10 = tenax.
crenulata, *Spreng. Syst.* i. 665.—Porto Rico.
Cruegerii, *Griseb. Fl. Brit. W. Ind.* 401.—Ind. occ.
cubensis, *Griseb. Cat. Pl. Cub.* 164.—Cuba.
cuneata, Sw. Fl. Ind. Occ. i. 496 = Sideroxylon obovatum, *Lam.*
cuneifolia, Rudge, Pl. Guian. i. 30, t. 47 = Chrysophyllum cuneifolium.
denticulata, *Rafin. New Fl. Am.* iii. 29.—Am. bor.
dulcifica, Schum. & Thonn. Beskr. Guin. Pl. 130 = Sideroxylon dulcificum.
Dunantii, *A. DC. in DC. Prod.* viii. 191.—Am. austr.
egensis, Poepp. ex A. DC. l. c. 182 = Sideroxylon rugosum.
excelsa, A. DC. l. c. 192 = obtusifolia.
ferox, *Cham. & Schlecht. in Linnaea*, vi. (1831) 392.—Mexic.
ferruginea, Nutt. Sylv. Am. 33 = lanuginosa.
foetidissima, Willd. Sp. Pl. i. 1086 = Sideroxylon foetidissimum.
glomerata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 518.—Ind. occ.
Hayesii, *Hemsl. Biol. Centr. Am. Bot.* ii. 297.—Guatemala.

BUMELIA :—
horrida, *Griseb. Cat. Pl. Cub.* 165.—Cuba.
laete-virens, *Hemsl. Biol. Centr. Am. Bot.* ii. 298.—Mexic.
lanuginosa, *Pers. Syn.* i. 237.—Am. bor.
lucida, Roem. & Schult. Syst. iv. 499 = Sideroxylon lucidum.
lycioides, *Willd. Enum. Hort. Berol.* 249.—Am. bor.
macrocarpa, Nutt. Sylva Am. iii. 37 = lanuginosa ?
Manglillo, Willd. Sp. Pl. i. 1087 = Myrsine Manglilla.
Mastichodendrum, Roem. & Schult. Syst. iv. 493 = Sideroxylon Mastichodendron.
micrantha, Willd. ex Roem. & Schult. l. c. 802 = Ardisia micrantha.
microphylla, *Griseb. Cat. Pl. Cub.* 165.—Cuba.
montana, Sw. Prod. Veg. Ind. Occ. 49 = Dipholis montana.
monticola, *Buckl. in Bull. Torrey Club*, x. (1883) 91.—Texas.
multiflora, Rocm. & Schult. Syst. iv. 498 (non Sapotacea).—Afr. austr.
myrsinifolia, A. DC. in DC. Prod. viii. 192 = Sideroxylon obovatum, *Lam.*
nervosa, Vahl, Eclog. Am. i. 28 = Lucuma Rivicoa.
nigra, Bert. ex Griseb. Fl. Brit. W. Ind. 400 = Dipholis salicifolia.
nigra, Sw. Prod. Veg. Ind. Occ. 49 = Dipholis nigra.
oblongifolia, Nutt. Gen. Am. i. 135 = lanuginosa.
obovata, *A. DC. in DC. Prod.* viii. 191.—Am. austr.
obtusifolia, *Roem. & Schult. Syst.* vi. 802.—Bras.
obtusifolia, Willd. ex Steud. Nom. ed. II. i. 237 = rotundifolia.
occidentalis, *Hemsl. Biol. Centr. Am. Bot.* ii. 298.—Mexic.
pallida, Sw. Prod. Veg. Ind. Occ. 49 = Sideroxylon Mastichodendron.
parvifolia, *A. DC. in DC. Prod.* viii. 190.—Ind. occ.
parvifolia, Chapm. Fl. S. U. St. 275 = angustifolia.
pauciflora, Engelm. ex A. Gray, Syn. Fl. ii. 11. 68 = lanuginosa.
pauciflora, Rocm. & Schult. Syst. iv. 493 = Sideroxylon pauciflorum.
pentagona, Sw. Prod. Veg. Ind. Occ. 50 = Dipholis salicifolia.
persimilis, *Hemsl. Biol. Centr. Am. Bot.* ii. 298.—Mexic.
pubescens, Tenore, Sem. Hort. Neap. (1827) = lycioides.
punctata, Roem. & Schult. Syst. iv. 498 = Myrsine Rapanea ?
reclinata, Torr. Bot. Mex. Bound. 109 = angustifolia.
reclinata, *Vent. Choix*, 22.—Am. bor.
retusa, Balbis, ex A. DC. Prod. viii. 194 = Ilicis sp.
retusa, *Sw. Prod. Veg. Ind. Occ.* 49.—Ind. occ.
rhamnoides, Casar. Stirp. Bras. Dec. 64 = obtusifolia.
rotundifolia, H. B. & K. Nov. Gen. et Sp. iii. 237 = obtusifolia.
rotundifolia, *Sw. Prod. Veg. Ind. Occ.* 50.—Ind. occ.
rufa, *Rafin. New Fl. Am.* iii. 29.—Am. bor.
salicifolia, Bert. ex A. DC. in Trans. Linn. Soc. xvii. (1837) 108 = Myrsine salicifolia.
salicifolia, Sw. Prod. Veg. Ind. Occ. 50 = Dipholis salicifolia.
sartorum, *Mart. Herb. Fl. Bras.* 233.—Bras.
serrata, Pursh, Fl. Am. Sept. i. 155 = Prunus caroliniana.
serrulata, *Rafin. New Fl. Am.* iii. 29.—Am. bor.
spiniflora, *A. DC. Prod.* viii. 191.—Mexic.
spinosa, *A. DC. l. c.*—Mexic.
spinosa, S. Wats. in Proc. Am. Acad. xviii. (1883) 112 = lanuginosa.
strigosa, *Spreng. Syst.* i. 655.—Hab. ?
subsessiflora, *Hemsl. Biol. Centr. Am. Bot.* ii. 299.—Mexic.
tenax, *Willd. Sp. Pl.* i. 1085.—Am. bor.
texana, *Buckl. in Bull. Torrey Club*, x. (1883) 90.—Texas.
tomentosa, A. DC. Prod. viii. 190 = lanuginosa.
tortuosa, *Wright, in Sauv. Fl. Cub.* 87.—Cuba.
undulata, *Rafin. New Fl. Am.* iii. 28.—Am. bor.

BUNBURIA, Harv. Gen. S. Afr. Pl. ed. I. 416 (1838) = **Vincetoxicum**, Moench (Asclep.).
elliptica, Harv. l. c. 417.—Afr. austr.

BUNBURYA, Meissn. ex Hochst. in Flora, xxvii. (1844) 553 = **Tricalysia**, A. Rich. (Rubiac.).
capensis, Meissn. ex Hochst. l. c.—Afr. austr.

BUNCHOSIA, Rich. ex Juss. in Ann. Mus. Par. xviii. (1811) 481. *MALPIGHIACEAE*, Benth. & Hook. f. i. 252.
Malacmaea, Griseb. in Linnaea, xiii. (1839) 248.
acutifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 336.—Mexic.
angustifolia, *A. Juss. l. c.* 331.—Peruv.
argentea, *DC. Prod.* i. 582.—Venezuela.
armeniaca, *DC. l. c.*—Peruv.
bilocellata, Hemsl. Biol. Centr. Am. Bot. i. 147 = seq.
biocellata, *Schlecht. in Linnaea*, x. (1836) 241.—Mexic.
Bonplandiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 332.—Am. trop.
bracteosa, Juss. in St. Hil. Fl. Bras. Mer. iii. 79 = Dicella bracteosa.
canescens, *DC. Prod.* i. 582.—Ind. occ.
cornifolia, *H. B. & K. Nov. Gen. et Sp.* v. 154.—N. Granat.
cumanensis, *G. Don, in Loud. Hort. Brit.* 170.—Venezuela.
deflexa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 310.—N. Granat.
discolor, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 266 = biocellata.
eglandulosa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 336.—Hab. ?
elliptica, *Tod. Ind. Sem. Hort. Panorm.* (1877) 38.—Hab. ?
emarginata, *Regel, Ind. Sem. Hort. Petrop.* (1856) 35.—Cuba.
fluminensis, *Griseb. in Mart. Fl. Bras.* xii. 1. 31.—Bras.
Gaudichaudiana, A. Juss. in Arch. Mus. Par. iii. (1843) 334 = fluminensis.
glandulifera, *H. B. & K. Nov. Gen. et Sp.* v. 154.—Venezuela.
glandulosa, *DC. Prod.* i. 581.—Am. trop.
glauca, *H. B. & K. Nov. Gen. et Sp.* v. 155.—N. Granat.
glauca, Seem. ex Hemsl. Biol. Centr. Am. Bot. i. 147 = cornifolia.
grandifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 341.—Ins. Martinic.
Hartwegiana, *Benth. Pl. Hartw.* 164.—N. Granat.
holosericea, Mart. ex A. Juss. in Arch. Mus. Par. iii. (1843) 344 = Dicella holosericea.
Hookeriana, *A. Juss. l. c.* 337.—Peruv.
hypoleuca, *Miq. Stirp. Surin. Sel.* 77.—Guiana.
lanceolata, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 266 = discolor.
laxiflora, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 168 = Tetrapteris Guilleminiana.
Lindeniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 335.—Am. trop.
Martiana, A. Juss. l. c. 334 = fluminensis.
media, *DC. Prod.* i. 581.—Ind. occ.
mollis, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 127.—Guiana.
mollis, Seem. ex Hemsl. Biol. Centr. Am. Bot. i. 148 = pilosa.
montana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 340.—Mexic.
multiflora, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 157.—Am. austr.
nitida, *Juss. in Ann. Mus. Par.* xviii. (1811) 481.—Mexic.; Peruv.; Ind. occ.
odorata, *Juss. l. c.*—N. Granat.
palustris, Poepp. ex A. Juss. in Arch. Mus. Par. iii. (1843) 311 = Burdachia prismatocarpa.
paniculata, DC. Prod. i. 582 = Triopteris ovata.
pilosa, *H. B. & K. Nov. Gen. et Sp.* v. 154.—N. Granat.
polystachya, *DC. Prod.* i. 581.—Ind. occ.
retusa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 311.—N. Granat.
rhombifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 582.—Venezuela.
sessilifolia, *DC. Prod.* i. 582.—Mexic.
sphaerocarpa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 313.—Bras.
squarrosa, Griseb. in Linnaea, xxii. (1849) 11 = Tetrapterys squarrosa.
strigosa, *Schlecht. in Linnaea*, x. (1836) 242.—Mexic.
Swartziana, *Griseb. Fl. Brit. W. Ind.* 115.—Jamaic.
tuberculata, *DC. Prod.* i. 581.—Venezuela.

BUNGEA, C. A. Mey. Verz. Pfl. Cauc. 108 (1831). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 976.
Sheareri, *S. Moore, in Journ. Bot.* xiii. (1875) 229.—China.
trifida, *C. A. Mey. Verz. Pfl. Cauc.* 108.—As. Min.; Persia.
turkestanica, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. No. III. 61.—Turkestan.

BUNIAS, [Tourn.] Linn. Syst. ed. I. (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 97.
Erucago, Tourn. ex Adans. Fam. ii. 423 (1763).
Laelia, Adans. Fam. ii. 423 (1763); Desv. Journ. Bot. iii. (1813) 160.
aegyptiaca, Linn. Syst. Nat. ed. XII. 231 = Ochthodium aegyptiacum.
altaica, Schang. ex DC. Syst. ii. 388 = Hutchinsia calycina.
americana, Rafin. Med. Fl. ii. 202 = Cakile maritima.
arvensis, *Jord. Cat. Gr. Jard. Dijon* (1848) 18.—Gallia.
aspera, Retz. Obs. ii. 21 = B. Erucago.
balearica, Linn. Syst. Nat. ed. XII. 446 = Succowia balearica.
Cakile, Linn. Sp. Pl. 670 = Cakile maritima.
cochlearioides, Bieb. Fl. Taur. Cauc. ii. 87 = Calepina Corvini.
cochlearioides, *Murr. in Comm. Gott.* viii. (1777) 42. t. 3.—Sibir. altaic.
cornuta, Linn. Sp. Pl. 669 = Pugionium cornutum.
cristata, Desv. Journ. Bot. iii. (1813) 163 = B. Erucago.
edentula, Bigel. Fl. Bost. ed. III. 267 = Cakile maritima.
Erucago, *Linn. Sp. Pl.* 670.—Europ. austr.; As. Min.; Syria.
glomerata, Lapeyr. Hist. Abr. Pl. Pyr. 362 = Senebiera Coronopus.
littoralis, Salisb. Prod. 273 = Cakile maritima.
macroptera, Reichb. Fl. Germ. Excurs. 654 = B. Erucago.
myagroides, Linn. Mant. i. 96 = Erucaria aleppica.
orientalis, *Linn. Sp. Pl.* 670.—Europ. or.; As. Min.
ovalis, Viv. Fl. Lib. Spec. t. 16 = Cakile cyrenaica.
paniculata, L'Hérit. ex DC. Syst. ii. 520 = Neslia paniculata.
paradoxa, Banks, ex DC. l. c. 677 = Chamira cornuta.
perennis, Moench, Meth. Suppl. 91 = orientalis.
perennis, Sibth. & Sm. Fl. Graec. Prod. ii. 2 = Rapistrum perenne.
prostrata, Desf. Fl. Atlant. ii. 76 = Muricaria prostrata.
pyramidata, *Fisch. Hort. Gorenk.* ed. II. (1812) 49, nomen.—Hab. ?
raphanifolia, Sibth. & Sm. Fl. Graec. t. 612 = Rapistrum orientale.
rostrata, L'Hérit. ex DC. Syst. ii. 422 = Euclidium syriacum.
sibirica, *Linn. Amoen. Acad.* viii. 8.—Sibir.
spinosa, Linn. Mant. 96 = Zilla myagroides.
syriaca, Gaertn. Fruct. ii. 290 = Euclidium syriacum.
tatarica, Willd. Sp. Pl. iii. 413 = Euclidium tataricum.
tcheliensis, *Debeaux, in Actes Linn. Soc. Bord.* xxxiii. (1879) 35.—China.
tenuifolia, Sibth. & Sm. Fl. Graec. Prod. ii. 3 (= *Didesmus tenuifolius*).—Graecia.
Tournefortii, Sibth. ex DC. Syst. ii. 659 (= *Didesmus tenuifolius*).—Graecia.
tricornis, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 82.—Hispan.
verrucosa, Moench, Meth. 277 = Rapistrum orientale.
virgata, Sibth. & Sm. Fl. Graec. t. 613 = Ochthodium aegyptiacum.
vulgaris, Andrz. ex Steud. Nom. ed. II. i. 237, 238 = B. Erucago.
Winterli, Schult. Obs. 121 = orientalis.

BUNION, St. Lag. in Ann. Soc. Lyon, vii. 55 (1880) = seq.

BUNIUM, Koch, Syn. Fl. Germ. ed. II. 315 (1843) = **Pimpinella**, Riv. (Umbellif.).
montanum, Koch, l. c.—Europ.

BUNIUM, Linn. Gen. ed. I. 72 (1737) = **Carum**, Linn. (Umbellif.).
acaule, Bieb. Fl. Taur. Cauc. i. 212 = Chamaesciadium acaule.

BUNIUM :—
alpinum, Waldst. & Kit. Pl. Rar. Hung. ii. 199. t. 182 = C. alpinum.
ammoides, Cesati, in Linnaea, xi. (1837) 323 = C. ammoides.
ammoides, Link, ex Spreng. Syst. i. 903 = Conopodium denudatum.
aphyllum, Jan, ex DC. Prod. iv. 117.—Sicilia.
arcuatum, Griseb. ex Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 267.—Europ.
aromaticum, Linn. Mant. 218 = C. copticum.
brevifolium, Lowe, in Trans. Camb. Phil. Soc. vi. (1838) Repr. 21.—Ins. Mader.
Bulbocastanum, Bert. ex DC. Prod. iv. 668 = Diposis Bulbocastanum.
Bulbocastanum, Huds. Fl. Angl. 122 = Conopodium denudatum.
Bulbocastanum, Linn. Sp. Pl. 243 = C. Bulbocastanum.
bulbosum, Dulac, Fl. Hautes-Pyr. 351 = C. Bulbocastannm.
butinioides, Boiss. & Reut. ex Boiss. Voy. Bot. App. 736.—Hispan.
capillifolium, Bertol. Fl. Ital. iii. 219.—Italia.
capillifolium, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 428 = Carum setaceum.
Carvi, Bieb. Fl. Taur. Cauc. i. 211 = C. Carvi.
carvifolium, DC. Prod. iv. 116 = C. flexuosum.
cassium, Boiss. Diagn. Ser. I. x. 24 = C. elegans.
cilicicum, Fenzl, in Tchihat. Asie Min. Bot. i. 411 = C. Pestalozzae.
collinum, Albert, in Bull. Soc. Sc. Draguign. (1884) repr. 23.—Europ.
copticum, Spreng. Prod. Umb. 28 = C. copticum.
corydalinum, DC. Prod. iv. 117.—Ins. Corsic.
creticum, Mill. Gard. Dict. ed. VIII. n. 2 = C. ferulaefolium.
creticum, Urv. in Bull. Soc. Linn. Par. i. (1822) 287 (= Conopodii sp.).—Graecia ; As. Min. ; Syria.
cynapioides, Bertol. Fl. Ital. iii. 217 = C. cylindricum ?
denudatum, DC. Fl. Fr. iv. 525 = Conopodium denudatum.
divaricatum, Bertol. Fl. Ital. iii. 222 (= Pimpinellae sp.).—Ital.
divaricatum, Cesati, in Linnaea, xi. (1837) 314 = C. ferulaefolium.
Falcaria, Bieb. Fl. Taur. Cauc. i. 211 = Falcaria Rivini.
ferulaceum, Sibth. & Sm. Fl. Graec. Prod. i. 186 = C. ferulaefolium.
ferulaefolium, Desf. in Ann. Mus. Par. xi. (1808) 275 ; Choix, 55. t. 43 = C. ferulaefolium.
flexuosum, Brot. Fl. Lusit. i. 437 = Conopodium capillifolium.
flexuosum, With. Bot. Arr. Brit. Pl. ed. II. 276 = Conopodium denudatum.
glaberrimum, DC. Prod. iv. 118.—Algeria.
glaucescens, DC. l. c. 117 (= *Scaligeria glaucescens*). —Armenia ; Persia.
glaucocarpum, Boiss. Diagn. Ser. I. x. 25 = C. Pestalozzae.
junceum, Margot & Reut. in Mém. Soc. Phys. Genèv. viii. (1839) 300 = C. creticum.
ligusticifolium, Cesati, ex Boiss. Fl. Orient. ii. 882 = C. multiflorum.
luteum, Hoffm. Gen. Umb. 108 = Muretia tanaicensis.
Macuca, Boiss. Elench. 44 ; Voy. Espagne, ii. 239. t. 66 = C. Macuca.
majus, Bieb. Fl. Taur. Cauc. i. 210 = C. ferulaefolium.
majus, Gouan, Illustr. 10 = Conopodium denudatum.
majus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 514 = C. Bulbocastanum.
minus, Gouan, Illustr. 10 = C. Bulbocastanum.
minus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 514 = Conopodium denudatum.
minutifolium, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 177 = Peucedanum vittijugum.
napiforme, Willd. ex Roem. & Schult. Syst. vi. 539 = C. creticum.
nepalense, Jacquem. ex Walp. Rep. ii. 422 = Chaerophyllum capnoides.
nivale, Boiss. Elench. 45 ; Voy. Espagne, ii. 240. t. 67 = C. nivale.
paucifolium, DC. Prod. iv. 117 = C. elegans.
Pestalozzae, Boiss. Diagn. Ser. I. x. 24 = C. Pestalozzae.

BUNIUM :—
petraeum, Loisel. Fl. Gall. ed. II. i. 195 = *corydalinum*.
petraeum, Tenore, Prod. Fl. Neap. p. lix = Carum alpinum.
peucedanoides, Bieb. Fl. Taur. Cauc i. 211 = Silaus carvifolius.
peucedanoides, DC. Prod. iv. 116.—Reg. Caucas.
pilosum, Willd. ex Roem. & Schult. Syst. vi. 500 = Conopodium denudatum.
pityranthon, Spreng. Syst. i. 901 = Pitnranthus denudatus.
pumilum, Sibth. & Sm. Fl. Graec. iii. 67. t. 274 (= *Biasolettia pumila*).—Graecia.
pyrenaeum, Loisel. Fl. Gall. 161. t. 5 = Conopodium denudatum.
rigens, Spreng. in Roem. & Schult. Syst. vi. 500.—Afr. austr.
rigidulum, Caruel, Prod. Fl. Tosc. 272 = C. rigidulum.
saxatile, Mill. Gard. Dict. ed. VIII. n. 3 = C. alpinum.
strictum, Griseb. Spicil. Fl. Rumel. i. 344 = C. strictum.
subcarneum, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 14 = Conopodium subcarneum.
tenuifolium, Salzm. ex DC. Prod. iv. 118 = Conopodium capillifolium.
tenuisectum, Griseb. ex Pantoč. in Oestr. Bot. Zeitschr. xxiii. (1873) 267.—Europ.
ternatum, Pers. Syn. i. 308 = Conopodium denudatum.
thalictrifolium, Boiss. Elench. 45.—Hispan.
verticillatum, Gren. & Godr. Fl. Fr. i. 729 = C. verticillatum.
virescens, DC. in Mém. Soc. Phys. Genèv. iv. (1828) 197 = Silaus carvifolius.

BUNNYA, F. Muell. Fragm. v. 36. t. 39 (1865) = **Cyanostegia**, Turcz. (Verbenac.).
cyanocalyx, F. Muell. l. c. = C. Bunnyana.

BUNOPHILA, Willd. ex Roem. & Schult. Syst. iii. Mant. 128, et Ind. (1827) (Ord. Nat. ?).
PUNOPHILA, Willd. l. c. 8 (1827).
lycioides, Willd. l. c. 128.—Am. austr.

BUONAPARTEA, G. Don, in Sweet, Hort. Brit. ed. III. 706 (1839) = BONAPARTEA, Ruiz & Pav. = **Tillandsia**, Linn. (Bromel.).

BUPHANE, Herb. in Bot. Mag. sub t. 2578 (1825) ; et Amaryll. 239. t. 22, 36. f. 7, 8 (1825) ; prius Boophane, ejusd. App. 18 (1821) ; serius Buphone in Bot. Mag. sub t. 2593 in nota (1825). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 730.
CROSSYNE, Salisb. Gen. Pl. Fragm. 116 (1866).
amaryllidea, M. Roem. Syn. Ensat. 61 = ciliaris.
angolensis, Baker, in Journ. Bot. xvi. (1878) 197 = Crinum angolense.
ciliaris, *Herb. App.* 18 (Boophane).—Afr. austr.
disticha, *Herb. in Bot. Mag.* sub t. 2578.—Afr. austr.
guttata, Herb. Amaryll. 240. t. 22. f. 1 = ciliaris.
intermedia, M. Roem. Syn. Ensat. 59 = disticha.
toxicaria, Herb. App. 18 (Boophane) = disticha.

BUPHTHALMUM, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 338.
ATHALMUM, Neck. Elem. i. 20 (1790).
BUSTIA, Adans. Fam. ii. 127 (1763).
MOLPADIA, Cass. Dict. Sc. Nat. xxiii. 565 (1822).
TELEKIA, Baumg. Fl. Transsylv. iii. 149 (1816).
angustifolium, Pursh, Fl. Am. Sept. ii. 564 = Marshallia angustifolia.
aquaticum, Linn. Sp. Pl. 903 = Odontospermum aquaticum.
arabicum, Delile, in Laborde, Voy. Arab. Pétr. 84 = Anvillea Garcini.
arborescens, Linn. Syst. ed. X. 1227 = Borrichia arborescens.
arvense, *Vell. Fl. Flum.* viii. t. 133.—Bras.
asperrimum, Spreng. Neue Entdeck. ii. 140 (= *Anomostephium buphthalmoides*).—Guadelup.
asperum, Hornem. Hort. Hafn. Suppl. 102 = Wedelia hispida.
asteriscoides, Baill. Hist. Pl. viii. 301 = Osmites asteriscoides.
asteroideum, Viv. Fl. Lib. Spec. 57. t. 25. f. 2 = Pallenis spinosa.
australe, Spreng. Fl. Hal. Mant. 52 = Wedelia aristata.

BUPHTHALMUM :—
aureum, Salzm. ex Steud. Nom. ed. II. i. 238 = Pallenis spinosa.
bipinnatum, Gilib. Fl. Lituan. i. 213 = Anthemis tinctoria.
bonariense, Pers. Syn. ii. 474 = Acicarpha spathulata ?
canum, L'Hérit. ex DC. Prod. v. 489 = Borrichia argentea.
capense, Linn. Syst. ed. X. 1227 = Oedera prolifera.
chrysanthemoides, Poir. Encyc. Suppl. i. 747 = Odontospermum imbricatum.
cordifolium, Waldst. & Kit. Pl. Rar. Hung. ii. 117 = speciosum.
coriaceum, Hort. ex Loud. Hort. Brit. 357 = Jasonia laevigata.
crassifolium, Rafin. Specch. 191 = Odontospermum maritimum.
dentatum, Baill. Hist. Pl. viii. 301 = Osmites dentata.
diffusum, Vahl, ex DC. Prod. v. 491 = Eclipta erecta.
durum, Linn. Sp. Pl. ed. II. 1275 = Osteospermum spinosum.
elegans, Brouss. ex Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 138 = Odontospermum imbricatum.
equinum, Vell. Fl. Flum. viii. t. 131.—Bras.
flexile, *Bertol. Fl. Ital.* ix. 413.—Ital.
flosculosum, Vent. Jard. Cels, 25 = Anvillea Garcini.
frutescens, Linn. Sp. Pl. 903 = Borrichia frutescens.
Garcini, Burm. f. Fl. Ind. t. 60 = Anvillea Garcini.
grandiflorum, Linn. Sp. Pl. 904 = salicifolium.
graveolens, Forsk. Fl. Aegypt. Arab. 151 = Odontospermum graveolens.
graveolens, Schimp. ex Boiss. Fl. Orient. iii. 180 (= *Asteriscus Schimperi*).—Arab.
Gussoni, *Pasquale, Notizie*, (1878) 2.—Sicil. ?
helianthoides, Forst. f. Prod. 57 = Wedelia aristata.
helianthoides, Linn. Sp. Pl. 904 = Heliopsis laevis.
heterophyllum, Willd. ex DC. Prod. v. 484 = Wedelia pedunculosa.
imbricatum, Cav. in Anal. Cienc. Nat. iv. (1801) 94 = Odontospermum imbricatum.
incanum, Mill. Gard. Dict. ed VIII. n. 10 = Borrichia frutescens.
inuloides, *Moris, Sard. App.* (1828).—Sardinia.
laevigatum, Brouss. ex Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 138 = Jasonia laevigata.
leucoifolium, *Burm. f. Fl. Cap. Prod.* 27.—Afr. austr.
lineare, Willd. Sp. Pl. iii. 2230 = Borrichia frutescens.
littorale, *Vell. Fl. Flum.* viii. t. 134.—Bras.
longipes, *Comm. ex Cass. in Bull. Soc. Philom.* (1822) 144.—Madag.
maritimum, Linn. Sp. Pl. 903 = Odontospermum maritimum.
mediterraneum, *Vell. Fl. Flum.* viii. t. 135.—Bras.
melissoides, Poir. Encyc. Suppl. i. 747 = Heliopsis scabra.
melitense, *Forsk. Fl. Aegypt. Arab.* 218, nomen.—Ins. Melita.
nutans, Vitm. Summa Veg. v. 123 = Bidens cernua.
odorum, Schousb. Vextr. Marokko, 199 = Odontospermum odorum.
oleraceum, *Lour. Fl. Cochinch.* ii. 506.—China.
Ortegae, *Spreng. Nachr. i. Bot. Gart. Halle*, 16, nomen.—Hab. ?
peruvianum, Lam. Encyc. i. 515 = Borrichia peruviana.
pratense, Vahl, Symb. Bot. i. 75 = Ceruana pratensis.
procumbens, Desf. ex Steud. Nom. ed. I. 127 = Wedelia carnosa.
ramosum, *Forsk. Fl. Aegypt. Arab.* 151.—Arab.
Ramtilla, Buch.-Ham. ex Wall. Cat. sub n. 3191 = Guizotia oleifera.
repens, Lam. Encyc. i. 515 = Wedelia carnosa.
sagittatum, Pursh, Fl. Am. Sept. ii. 564 = Balsamorhiza sagittata.
salicifolium, *Linn. Sp. Pl.* 904.—Europ. austr.
scabrum, Cav. Ic. ii. 53 = Wedelia hispida.
scandens, *Vell. Fl. Flum.* viii. t. 132.—Bras.
scandens, Schum. & Thonn. Beskr. Guin. Pl. 392 = Melanthera Brownei.
sericeum, Linn. f. Suppl. 379 = Odontospermum sericeum.
sessile, Mill. Gard. Dict. ed. VIII. n. 5 = Odontospermum maritimum.
silphioides, *Hornem. ex Steud. Nom.* ed. II. i. 239.—Hab. ?
speciosissimum, *Ard. Spec.* i. 26 ; *Linn. Mant.* i. 117.—Europ.
spcciosum, *Schreb. Dec.* 11. t. 6.—Oriens.
spinosum, Linn. Sp. Pl. 904 = Pallenis spinosa.

BUPHTHALMUM:—
stenophyllum, Link, in Buch, Beschr. Canar. Ins. 150 = Odontospermum sericeum.
strigosum, Spreng. Neue Entdeck. ii. 140 = Wedelia pedunculosa.
succisaefolium, Britting. in Flora, xxiv. (1841) i. 138 = salicifolium.
tenuifolium, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
trifolium, Pocpp. ex DC. Prod. v. 577 = Oyedaca buphthalmoides.
uniflorum, Forst. f. Prod. 541 = Wedelia Forsteriana.

BUPLEUROIDES, [Boerh.] Moench, Meth. 294 (1794) = **Phyllis**, Linn. (Rubiac.).
Nobla, Moench, l. c. = P. Nobla.

BUPLEURUM, [Tourn.] Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 886.
AGROSTANA, Hill, Veg. Syst. v. 32 (1772).
BUPRESTIS, Spreng. Pl. Umb. Prod. 27 (1813).
DIAPHYLLUM, Hoffm. Gen. Umb. 115 (1814).
DIATROPA, Dum. Fl. Belg. 76 (1827).
ISOPHYLLUM, Hoffm. Gen. Umb. 115 (1814).
ODONTEA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 391.
ODONTITES, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 258.
PERFOLIATA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 390.
TENORIA, Spreng. Pl. Umb. Prod. 32 (1813).
TRACHYPLEURUM, Reichb. Consp. 143 (1828).
acerosum, *E. Mey. in Drège, Zwei Pfl. Docum.* 170, nomen.—Afr. austr.
aciphyllum, *Webb & Berth. Phyt. Canar.* ii. 154. t. 77.—Ins. Canar.
acutifolium, Boiss. Elench. 47 = paniculatum.
aegyptiacum, Willd. ex Steud. Nom. ed. II. i. 239 = protractum.
aeneum, *Boiss. & Huet, Diagn.* Ser. II. ii. 85.—Armenia.
affine, *Sadl. Fl. Pesth.* 204.—Europ. austr.; Tauria.
aleppicum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 148.—Syria.
alpigenum, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 35 = falcatum.
alpinum, Nym. Consp. 311 = falcatum.
altaicum, Pall. ex Roem. & Schult. Syst. vi. 367, 368 = triradiatum.
amplexicaule, [Clairv.] Man. Herb. 87 = pyrenaicum.
angulosum, Cham. & Schlecht. in Linnaea, i. (1826) 383 = ranunculoides.
angulosum, *Linn. Sp. Pl.* 236.—Reg. arct.
angustifolium, *Ledeb. Fl. Ross.* ii. 265.—Sibir. altaic.
angustifolium, *Pall. Reise Statth. Russ. Reich.* i. 345.—Reg. Caucas.
apiculatum, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 147 = semidiaphanum.
apiculatum, *Friwald. in Flora*, xviii. (1835) 335.—Oriens.
arborescens, Jacq. Coll. ii. 343 = gibraltaricum.
arborescens, Thunb. Prod. Pl. Cap. 50 = Heteromorpha arborescens.
aristatum, *Bartl. in Reichb. Ic. Fl. Germ.* ii. 70. t. 178.—Europ.; Reg. Caucas.
armenum, Boiss. & Kotschy, ex Boiss. Fl. Orient. ii. 845 = kurdicum.
asperuloides, *Heldr. in Boiss. Diagn.* Ser. II. vi. 76.—Graecia.
Aucheri, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 148 = brevicaule.
aureum, *Fisch. in Hoffm. Gen. Umb.* 115.—Sibir.
australe, Jord. Pugill. 72 = Gerardi.
badium, Rochel & Lang, ex Saut. in Flora, ix. (1826) 177 = junceum.
Balansae, *Boiss. & Reut. Diagn.* Ser. II. ii. 83.—Afr. bor.
baldense, *Host, Syn.* 141.—Europ.; Oriens.
baldense, Turr. in Giorn. Ital. i. (1765) 120 = aristatum.
baldense, Waldst. & Kit. Ic. Pl. Hung. Rar. iii. 285. t. 257 = junceum.
Barcelloi, *Coss. ex Willk. in Linnaea*, xl. (1876) 83.—Afr. bor.
Baumgartenianum, Schur, Enum. Pl. Transs. 252 = petraeum.
bicalyculatum, Bell. Stirp. Nov. 4. t. 2 = stellatum.
bicaule, Helm. in Mém. Soc. Nat. Mosc. ii. (1809) 108. t. 8 = baldense.

BUPLEURUM:—
Bourgaei, *Boiss. & Reut. Diagn.* Ser. II. ii. 84.—Hispan.
brachiatum, *C. Koch, ex Boiss. Fl. Orient.* ii. 844.—As. Min.
Brasianum, *Timb. in Mém. Acad. Toulouse*, Sér. VIII. iv. (1882) 128.—Gallia.
brevicaule, *Schlecht. in Linnaea*, xvii. (1843) 124.—Syria.
Burserianum, Willd. Enum. Hort. Berol. Suppl. 14 = angulosum.
canalense, Wulf. ex Spreng. in Roem. & Schult. Syst. vi. 371 = ranunculoides.
canariense, Spreng. Syst. i. 882 = Astydamia canariensis.
Candollii, *Wall. Cat.* n. 552.—Reg. Himal.
canescens, *Schousb. Vextr. Marokkko*, 127.—Marocco.
capillare, *Boiss. & Heldr. Diagn.* Ser. II. ii. 82.—Graecia.
capitatum, Thunb. Prod. Pl. Cap. 50 = Hermas capitata.
cappadocicum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 146.—As. Min.; Syria.
caricifolium, Willd. Sp. Pl. i. 1373 = ranunculoides.
caricinum, Reichb. Ic. Crit. ix. f. 1109, 1110 = ranunculoides.
cernuum, Tenore, App. i. Cat. Hort. Neap.; DC. Prod. iv. 131 = baldense.
chinense, *Franch. in Nouv. Arch. Mus. Par.* Sér. II. vi. (1883) 18.—China.
chinense, DC. Prod. iv. 128 = falcatum.
chlorocarpum, *Timb. & Jeanb. in Mém. Acad. Toulouse*, Sér. VI. v. (1867) 471, nomen.—Gallia.
Choulettii, *Pomel, Nouv. Mat. Fl. Atl.* 140.—Algeria.
ciliatum, Thunb. Prod. Pl. Cap. 50 = Hermas ciliata.
citrinum, *Hochst. in Lorent, Wanderungen*, 337.—Armenia.
collinum, D. Dietr. Syn. Pl. ii. 951 = Heteromorpha arborescens.
coloratum, Schur, Enum. Pl. Transs. 254 = aureum.
Columnae, Guss. Fl. Sic. Prod. 70 = tenuissimum.
commutatum, *Boiss. & Bal. Diagn.* Ser. II. vi. 75.—Macedon.; As. Min.
compositum, *Spreng. in Ges. Naturf. Fr. Berl. Mag.* vi. (1812) 258.—Hab. ?
corbariense, *Timb. in Mém. Acad. Toulouse*, Sér. VIII. v. II. (1883) 139.—Mont. Pyren.
coriaceum, L'Hérit. Stirp. Nov. 139. t. 67 = gibraltaricum.
corsicum, *Coss. & Kral. Notes Crit.* 56.—Corsica.
croceum, *Fenzl, Pugill. Pl. Nov. Syr.* 16.—As. Min.; Syria.
cuspidatum, Bunge, in Mém. Acad. Pétersb. vii. (1854) 299; (Lehm. Rel. 123) = linearifolium.
dahuricum, *Fisch. & Mey. ex Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 720, 721.—Dahuria.
densiflorum, *Rupr. Sert. Tiansch.* 47.—As. centr.
denticulatum, Bellard, ex Spreng. in Roem. & Schult. Syst. vi. 396 = petraeum.
dianthifolium, *Guss. Suppl. Fl. Sic. Prod.* i. 71.—Sicilia.
dichotomum, *Boiss. Diagn.* Ser. I. x. 28.—Syria.
dichotomum, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 342, in textu = Gerardi.
difforme, *Linn. Sp. Pl.* 238.—Afr. austr.
dilatatum, Schur, Enum. Pl. Transs. 253 = falcatum.
distichophyllum, *Wight & Arn. Prod.* i. 370.—Ind. or.
divaricatum, Lam. Fl. Fr. iii. 410 = B. Odontites.
divaricatum, Steph. ex Ledeb. Fl. Ross. ii. 262 = gracile.
diversifolium, Rochel, Banat. 68. t. 28 = baldense.
dumosum, *Coss. & Bal. in Bull. Soc. Bot. Fr.* xx. (1873) 249; xxii. (1875) 59.—Marocco.
elatum, *Guss. Fl. Sic. Prod.* i. 316.—Sicil.
erubescens, *Boiss. Fl. Orient.* ii. 841.—Lycia.
exaltatum, Bieb. Beschr. Casp. 156 = baldense.
exaltatum, Schur, Enum. Pl. Transs. 253 = subfalcatum.
falcatum, Berg. Desc. Pl. Cap. 76, partim = Mundtii.
falcatum, Dalz. & Gibs. Bomb. Fl. 108 = mucronatum.
falcatum, *Linn. Sp. Pl.* 237.—Europ.; Oriens; As. bor.; Reg. Himal.
falcatum, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 721 = dahuricum.
filicaule, *Brot. Fl. Lusit.* i. 452.—Lusitan.

BUPLEURUM:—
flavicans, *Boiss. & Heldr. Diagn.* Ser. II. vi. 74.—Aetolia.
flavum, *Forsk. Fl. Aegypt. Arab.* 205 (sp. dub.) = glumaceum ?
flexuosum, *Ledeb. Fl. Ross.* ii. 267.—Dahuria.
flexuosum, Moench, Meth. 88 = falcatum.
flexuosum, Wall. Cat. n. 554 = tenue.
foliosum, *Salzm. ex DC. Prod.* iv. 133.—Marocco.
Fontanesii, Guss. Ind. Sem. Hort. Bocc. (1825) = B. Odontites.
frutescens, Hill, Veg. Syst. vi. 21 = fruticosum.
fruticans, Steud. Nom. ed. I. 128 = difforme.
fruticescens, Bourg. ex Nym. Consp. 310 = foliosum.
fruticescens, *Linn. Cent. Pl.* i. 9.—Reg. Mediterr. occ.
fruticosum, *Linn. Sp. Pl.* 238.—Reg. Mediterr.
Gerardi, Colm. ex Nym. Consp. 312 = junceum.
Gerardi, Gaertn. Mey. & Scherb. Fl. Wett. i. 403 = tenuissimum.
Gerardi, *Jacq. Fl. Austr.* iii. 31. t. 256.—Europ. austr.; Oriens.
Gerardi, Pollini, Viag. Lag. Gard. 15, 81 = B. Odontites.
Gerardi, Sibth. & Sm. Fl. Graec. t. 262 = trichopodum.
Gerhardi, Georgi, Beschr. Russ. Reich. iii. IV. 828 = Gerardi.
gibraltaricum, *Lam. Encyc.* i. 520.—Lusitan.; Marocco.
giganteum, Thunb. Prod. Pl. Cap. 50 = Hermas gigantea.
glaucum, *Robill. & Cast. ex DC. Fl. Fr. Suppl.* 515.—Reg. Mediterr.
glumaceum, *Sibth. & Sm. Fl. Graec. Prod.* i. 177.—Graecia; As. Min.
gracile, DC. Prod. iv. 128 = Marschallianum.
gracile, Urv. in Mém. Soc. Linn. Par. i. (1822) 28 = glumaceum.
gracillimum, Klotzsch, in Bot. Ergeb. Waldem. Reise, 148 = falcatum.
gramineum, Vill. Prosp. 23 = baldense.
graminifolium, Favre, in Bull. Soc. Murith. v. (1875) 85, 250 = stellatum.
graminifolium, Georgi, Beschr. Russ. Reich. iii. IV. 828 = exaltatum.
graminifolium, Lapeyr. Hist. Abr. Pl. Pyr. 139 = ranunculoides.
graminifolium, Salisb. Prod. 160 = B. Odontites.
graminifolium, Vahl, Symb. Bot. iii. 48 = petraeum.
Haussknechtii, *Boiss. Fl. Orient.* ii. 843.—Kurdistan.
Heldreichii, *Boiss. & Bal. Diagn.* Ser. II. vi. 75.—As. Min.
heterophyllum, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 675 = jucundum.
heterophyllum, Link, Enum. Hort. Berol. i. 262 = protractum.
heterophyllum, Rochel, Banat. t. 28 = baldense.
himalayense, Klotzsch, in Bot. Ergeb. Waldem. Reise, 146 = longicaule.
hispanicum, Pers. Syn. i. 305 = fruticescens.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 147 = falcatum.
humile, Vest, ex Reichb. Ic. Fl. Germ. ii. 71 = aristatum.
imaicolum, Kern. in Ber. Naturw. Ver. Innsbruck, i. (1870) 115 (Nov. Pl. Sp. Dec. ii. 19) = falcatum.
incurvum, Bell. App. Fl. Pedem. 77. t. 11 = petraeum.
intermedium, Steud. Nom. ed. I. 128 = protractum.
irregulare, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* ii. 850.—Syria.
Jacquinianum, Jord. Pugill. 71 = Gerardi.
jucundum, *Kurz, in Journ. Bot.* v. (1867) 240.—Himal. bor. occ.; Reg. Amur.
junceum, Bieb. Fl. Taur. Cauc. i. 204 = affine.
junceum, *Linn. Sp. Pl.* ed. II. 343.—Europ. austr.
junceum, Pall. Ind. Taur. ex Bieb. Beschr. Casp. 166 = exaltatum.
junceum, Pollich, Fl. Palat. i. 264 = tenuissimum.
Kargli, *Vis. Fl. Dalm.* iii. 35.—Dalmat.
Koechelii, *Fenzl, Pugill. Pl. Nov. Syr.* 17.—As. Min.
kokanicum, *Regel & Schmalh. ex Regel. Pl. Nov. Fedsch.* 29.—As. centr.
Kotschyanum, Boiss. Diagn. Ser. I. x. 29 = linearifolium.
kurdicum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 146.—As. Min.; Syria; Persia.
lanceolatum, *Wall. Cat.* n. 551.—Reg. Himal.
lancifolium, Hornem. Cat. Hort. Hafn. i. 267 = protractum.

BUPLEURUM:—
laricense, *Gaut. & Timb. in Mém. Acad. Toulouse*, Sér. VIII. iv. (1882) 130.—Gallia.
lateriflorum, *Coss. in Bull. Soc. Bot. Fr.* xxii. (1875) 59.—Marocco.
leucocladum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 144.—Assyria.
libanoticum, *Boiss. & Blanche, Diagn.* Ser. II. ii. 82.—Syria.
linearifolinm, *DC. Prod.* iv. 131.—Persia; Arab.; Turkest.
longepetiolatum, Weiss, in Verh. Zool.-Bot. Ges. Wien, xvi. (1866) 581 = Kargli.
longeradiatum, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 719.—Sibir. baical.
longicaule, *Wall. Cat.* n. 557.—Reg. Himal.
longifolium, Falk, Beitr. ii. 144 = aureum.
longifolium, *Linn. Sp. Pl.* 237.—Europ.
lophocarpum, *Boiss. & Bal. Diagn.* Ser. II. v. 97.—Cilicia.
Maddeni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 678.—Reg. Himal.
mareoticum, Delile, ex DC. Prod. iv. 129 = nodiflorum.
marginatum, Wall. Cat. n. 566 = falcatum.
Marschallianum, *C. A. Mey. Verz. Pfl. Cauc.* 123.—Graecia; As. Min.; Persia.
minimum, Clarke, Trav. ii. 644; Spreng. Neue Entdeck. iii. 163 = nodiflorum.
montanum, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iii. (1856) 706.—Algeria.
mucronatum, Brouss. ex Spreng. in Roem. & Schult. Syst. vi. 377 = fruticescens.
mucronatum, *Wight & Arn. Prod.* 370.—Ind. or.
multinerve, *DC. in Mém. Phys. Soc. Genèv.* iv. (1828) 500.—Sibir. altaic.
Mundtii, *Cham. & Schlecht. in Linnaea*, i. (1826) 384.—Afr. austr.
nanum, Poir. Encyc. Suppl. i. 750 = nodiflorum.
nebulosum, Haussk. ex Boiss. Fl. Orient. ii. 847 = cappadocicum.
neglectum, Cesati, in Linnaea, xi. (1837) 314 = baldense.
nervosum, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 97 = polyphyllum.
nervosum, Moon, Cat. 22 = mucronatum.
nervosum, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 464 = paniculatum.
nervosum, Trev. Symb. Phytol. i. 22 = multinerve.
nigrocarpa, Jacquem. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 676 = falcatum.
nodiflorum, *Sibth. & Sm. Fl. Graec.* t. 260.—Graecia; Syria; Aegypt.
Nordmannianum, Bal. ex Boiss. Fl. Orient. ii. 851 = polyphyllum.
Nordmannianum, Ledeb. Fl. Ross. ii. 265 = baldense.
nudum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 331.—Afr. austr.
obliquatum, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 466 = foliosum.
obliquum, Vahl, Symb. Bot. i. 24 = gibraltaricum.
oblongifolium, *Ball, in Journ. Linn. Soc.* xvi. (1878) 466.—Marocco.
obtusatum, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. ii. 42 = angulosum.
obtusifolium, Tenore, Prod. Fl. Nap. p. xviii = baldense.
octoradiatum, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1830) 106.—China bor.
Odontites, *Linn. Sp. Pl.* 237.—Europ.; Reg. Mediterr.
Odontites, Sm. Engl. Bot. t. 2468 (1812); Lapeyr. Hist. Abr. Pl. Pyr. 141 (1813) = aristatum.
odontitum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 67, 121 = B. Odontites.
oligactis, *Boiss. Diagn.* Ser. II. ii. 84.—Hispan.; Marocco.
olympicum, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 149 = baldense.
opacum, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 71 = aristatum.
oppositifolium, *Lapeyr. Hist. Abr. Pl. Pyr.* 141.—Mont. Pyren.
pachnospermum, *Panč. Fl. Princ. Serbiae* (1874) 329.—Banat.
panacifolium, Hornem. ex Steud. Nom. ed. II. i. 240 = heterophyllum.
paniculatum, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 74 = Bourgaei.

BUPLEURUM:—
paniculatum, *Brot. Fl. Lusit.* i. 455.—Lusit.; Marocco.
paniculatum, Willk. ex Willk. & Lange, Prod. Fl. Hisp. iii. 74 = rigidum.
papillosum, *DC. Prod.* iv. 127.—Mesopotam.
pauciradiatum, *Fenzl, in Tchihat. Asie Min. Bot.* i. 418.—Cilicia.
perfoliatum, Lam. Fl. Fr. iii. 405 = rotundifolium.
persicum, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 150 = baldense.
petiolare, Lapeyr. Hist. Abr. Pl. Pyr. 141 = falcatum.
petraeum, Geners. ex DC. Prod. iv. 131 = angulosum.
petraeum, Georgi, It. i. 204 = exaltatum.
petraeum, *Linn. Sp. Pl.* 236.—Europ.
petraeum, Wulf. in Jacq. Coll. ii. 209 = graminifolium.
petrogenes, Jord. ex Nym. Consp. 311 = falcatum.
petroselinoides, Spreng. Pl. Umb. Prod. 39 = Carum peregrinum.
plantagineum, *Desf. Fl. Atlant.* i. 233. t. 57.—Marocco.
plantaginifolium, *Wight, Ic.* i. t. 281.—Ind. or.
Pollichii, C. C. Gmel. Fl. Bad. i. 613 = tenuissimum.
polyphyllum, *Ledeb. in Mém. Acad. Pétersb.* v. (1815) 528.—Reg. Caucas.
procumbens, Desf. Fl. Atlant. i. 230. t. 56 = tenuissimum.
proliferum, Delile, Fl. Egypte, 205. t. 22 = nodiflorum.
prostratum, *Ledeb. ex Loud. Hort. Brit. Suppl.* ii. 615.—Sibir. altaic.
protractum, *Hoffmgg. & Link, Fl. Port.* ii. 387.—Reg. Mediterr. or.
provinciale, *Huet, ex Timb. in Mém. Acad. Toulouse*, Sér. VIII. iv. (1882) 131.—Gallia.
pruinosum, Cesati, ex Boiss. Fl. Orient. ii. 849 = baldense.
pulchellum, *Boiss. & Heldr. Diagn.* Ser. I. x. 27.—Pamphylia.
pyrenaeicum, Willd. Sp. Pl. i. 1371 = angulosum.
pyrenaeum, Gouan, Illustr. 8. t. 4 = angulosum.
quinqueden[ta]tum, Thunb. Prod. Pl. Cap. 50 = Hermas quinquedentata.
quinqueradiatum, Steud. Nom. ed. II. i. 240, sphalm. = praec.
ramosissimum, Wight & Arn. Prod. 370 = mucronatum.
ramosum, *Gaut. ex Timb. in Mém. Acad. Toulouse*, Sér. VIII. iv. (1882) 136.—Gallia.
ranunculiformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 67 = seq.
ranunculoides, *Linn. Sp. Pl.* 237.—Reg. bor. temp.
ranunculoides, Sibth. & Sm. Fl. Graec. Prod. i. 178 = baldense.
repens, Lapeyr. Hist. Abr. Pl. Pyr. 140 = ranunculoides.
rigidum, Frey. ex Reichb. Ic. Fl. Germ. xxi. 21 = baldense.
rigidum, Georgi, Beschr. Russ. Reichb. iii. IV. 828 = Gerardi.
rigidum, *Linn. Sp. Pl.* 238.—Gall. austr.
Rissoni, Rich. ex DC. Prod. iv. 128 = affine.
rotundifolium, *Linn. Sp. Pl.* 236.—Europ.; Reg. Caucas.; Persia.
rotundifolium, Schousb. Vext. Marokko, 126 = protractum.
rupestre, Edgew. in Trans. Linn. Soc. xx. (1851) 52 = longicaule.
rupestre, *Rafin. in Med. Repos. N. York*, v. (1808) 352, nomen.—Hab.?
sachalinense, F. Schmidt, Reisen Amurl. 135 = jucundum.
salicifolium, *Soland. ex Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) repr. 21.—Ins. Madera.
Savignonii, *De Not. in Mem. Acc. Tor.* ix. (1848) 208.—Italia.
saxatile, Pall. Reise, ii. 676 = baldense.
scabrum, Griseb. Spicil. Fl. Rumel. i. 348 = brevicaule.
Scheffleri, Hampe, ex Nym. Consp. 313 = filicaule.
Schimperi, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 151 = linearifolium.
scorzoneraefolium, Willd. Enum. Hort. Berol. Suppl. 30 = falcatum.
semicompositum, Hochst. & Steud. ex Steud. Nom. ed. II. i. 240 = aristatum.
semicompositum, *Linn. Amoen. Acad.* iii. 405.—Europ. austr.
semicompositum, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. i. 205 = junceum.
semicompositum, Sibth. & Sm. Fl. Graec. Prod. i. 176 = glaucum.
scmidiaphanum, *Boiss. Diagn.* Ser. II. vi. 73.—Graecia et Ins.

BUPLEURUM:—
semiperfoliatum, Griseb. Spicil. Fl. Rumel. i. 350 = croceum.
setaceum, *Fenzl, in Tchihat. Asie Min. Bot.* i. 418.—Bithynia; Himal. bor. occ.
sibiricum, Vest, ex Roem. & Schult. Syst. vi. 368 = angulosum.
Sibthorpianum, DC. ex Boiss. Fl. Orient. ii. 849 = linearifolium.
Sibthorpianum, Sibth. & Sm. Fl. Graec. Prod. i. 179 = baldense.
spinosum, *Gouan, Illustr.* 8.—Hispan.; Afr. bor. occ.
Sprunerianum, Hampe, in Flora, xxv. (1842) I. 63 = B. Odontites.
stellatum, Lapeyr. Hist. Abr. Pl. Pyr. 139 = angulosum.
stellatum, *Linn. Sp. Pl.* 236.—Europ. anstr.
striatum, Steph. ex Roem. & Schult. Syst. vi. 367 = multinerve.
subfalcatum, *Schur, Enum. Pl. Transs.* 253.—Transsylv.
subovatum, Link, ex Spreng. Sp. Umb. 19 = protractum.
subpinnatum, Ledeb. in Eichw. Pl. Nov. Casp. Cauc. 13. t. 11 = Carum segetum.
subrubeum, Dulac, Fl. Hautes-Pyr. 346 = ranunculoides.
subuniflorum, *Boiss. & Heldr. Diagn.* Ser. I. x. 28.—Pamphylia.
sulcatum, Steud. Nom. ed. II. i. 240 = Mundtii.
sulphureum, *Boiss. & Bal. Diagn.* Ser. II. vi. 74.—As. Min.
telonense, Gren. ex Nym. Consp. 311 = ranunculoides.
tenue, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 182.—Reg. Himal.
tenue, Salisb. Prod. 160 = tenuissimum.
tenuifolium, Pourr. ex Nym. Consp. 311 = baldense.
tenuissimum, *Linn. Sp. Pl.* 238.—Europ.; Afr. bor.
terminale, Salisb. Prod. 160 = fruticosum.
Thomsoni, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 675.—Reg. Himal.
tortuosum, *Schousb. ex Ball, in Journ. Linn. Soc.* xvi. (1878) 466.—Marocco.
transsilvanicum, Schur, Enum. Pl. Transs. 253 = baldense.
trichopodum, *Boiss. & Sprun. in Ann. Sc. Nat.* Sér. III. i. (1844) 145.—Graecia; As. Min.
trifidum, Tenore, Prod. Fl. Nap. 18 = junceum.
trifoliatum, Wendl. & Bartl. Beitr. ii. 13 = Heteromorpha trifoliata.
triradiatum, *Adams, ex Hoffm. Umb.* 115.—Sibir.
vapincense, Vill. Prosp. 23 = longifolium.
Vanheurckii, *Muell. Arg. in Heurck, Obs. Bot.* 207.—China.
veronense, Turra, Fl. Ital. Prod. 64 = B. Odontites.
verticale, Orteg. Fl. Esp. ex DC. Prod. iv. 133 = gibraltaricum.
villosum, Linn. Sp. Pl. 238 = Hermas villosa.
virgatum, Brot. ex Steud. Nom. ed. II. i. 240 = fruticescens.
virgatum, Cav. Desc. i. 121 = Gerardi.
virgatum, Wall. Cat. n. 555 = falcatum.
virgatum, Wight & Arn. Prod. 370 = mucronatum.
Wittmanni, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 342 = rotundifolium.

BUPRESTIS, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 258 = **Bupleurum**, Tourn. (Umbell.).
arborescens, Spreng. l. c. 259 = Heteromorpha arborescens.
coriacea, Spreng. l. c. = B. gibraltaricum.
difformis, Spreng. l. c. = B. difforme.
exaltata, Spreng. l. c. = B. baldense.
frutescens, Spreng. l. c. = B. fruticescens.
fruticosa, Spreng. l. c. = B. fruticosum.
Gerardi, Spreng. l. c. = B. Gerardi.
gigantea, Spreng. l. c. = Hermas gigantea.
juncea, Spreng. l. c. = B. junceum.
latifolia, Spreng. l. c. = Astydamia canariensis.

BURAEAVIA, Baill. Adansonia, xi. (1873) 83. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 280.
carunculata, *Baill. l. c.* 84.—N. Caled.
clnsiacea, *Baill. l. c.*—N. Caled.

BURASAIA, Thou. Gen. Madag. 18 (1806); et in Roem. Collect. 209 (1809); DC. Syst. Veg. i. 514 (1818). *MENISPERMACEAE*, Benth. & Hook. f i. 39, 960.

BURASAIA :—
congesta, *Decne. in Ann. Sc. Nat.* Sér. II. xii. (1839) 108.—Madag.
gracilis, *Decne. l. c.*—Madag.
madagascariensis, *DC. Syst.* i. 514.—Madag.
Vitis, *Mello, in Journ. Linn. Soc.* xiv. (1875) 4.—Bras.

BURBIDGEA, Hook. f. Bot. Mag. t. 6403 (1879). *SCITAMINEAE*, Benth. & Hook. f. iii. 647.
nitida, *Hook. f. l. c.*—Borneo.

BURCARDA, J. F. Gmel. Syst. 514 (1791) = scq.

BURCARDIA, Schreb. Gen. i. 530 (1789) = **Piriqueta**, Aubl. (Turnerac.).
aromatica, Rafin. Sylva Tellur. 106.—Hab. ?
grandiflora, Rafin. l. c.—Hab. ?
villosa, J. F. Gmel. Syst. ii. 514 = P. villosa.

BURCHARDIA, [Heist.] Duham. Arb. i. 111, t. 44 (1755) = **Callicarpa**, Linn. (Verbenac.).
americana, Duham. l. c. = C. americana.
Callicarpa, Crantz, Inst. ii. 360 = C. americana.

BURCHARDIA, R. Br. Prod. 272 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 822.
congesta, Lindl. Swan River App. 58 = umbellata.
multiflora, Lindl. l. c. = umbellata.
umbellata, *R. Br. Prod.* 273.—Austral.

BURCHELLIA, R. Br. in Bot. Reg. t. 466 (1820). *RUBIACEAE*, Benth. & Hook. f. ii. 85.
BUBALINA, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 86.
Bubalina, Sims, Bot. Mag. t. 2339 = capensis.
capensis, *R. Br. in Bot. Reg.* t. 466.—Afr. austr.
Kraussii, Hochst. in Flora, xxv. (1842) 237 = capensis.
major, Hort. ex Heynh. Nom. ii. 83 = capensis.
parviflora, Lindl. Bot. Reg. t. 891 = capensis.
speciosa, Hort. ex Heynh. Nom. ii. 83 = capensis.

BURDACHIA, Mart. ex A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 329. *MALPIGHIACEAE*, Benth. & Hook. f. i. 252.
macrocarpa, Benth. ex Griseb. in Mart. Fl. Bras. xii. 1. 23 = Glandonia macrocarpa.
prismatocarpa, *Mart. ex A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 330.—Bras.
pyramidata, Mart. ex Griseb. in Mart. Fl. Bras. xii. 1. 23 = prismatocarpa.
sphaerocarpa, *A. Juss. in Ann. Sc. Nat.* Sér. II. xiii. (1840) 330.—Bras.

BUREAVA, Baill. Adansonia, i. 71 (1860-61) = **Combretum**, Linn.
crotonoides, Baill. l. c. 72 = C. altum.

BURGESIA, F. Muell. Fragm. i. 222 (1859) = **Brachysema**, R. Br.
homaloclada, F. Muell. l. c. = Brachysema aphyllum.

BURGHARTIA, Scop. Introd. 229 (1777) = **Piriqueta**, Aubl. (Turnerac.).
villosa, J. F. Gmel. Syst. 514 = P. villosa.

BURGLARIA, Wendl. ex Steud. Nom. ed. I. 129 (1821) = **Ilex**, Tourn.
lucida, Wendl. l. c. = I. salicifolia.

BURGSDORFIA, Moench, Meth. 392 (1794) = **Sideritis**, Tourn. (Labiat.).
montana, Reichb. Fl. Germ. Excurs. 327 = S. montana.
rigida, Moench, Meth. 392 = S. romana.
romana, Hoffmgg. & Link, Fl. Port. i. 97 = S. romana.

BURKEA, Benth. in Hook. Ic. Pl. t. 593 (1843). *LEGUMINOSAE*, Benth. & Hook. f. i. 587.
africana, *Hook. l. c.* tt. 593, 594.—Afr. austr.
Caperangan, *Baill. Hist. Pl.* ii. 167.—Afr. trop.

BURKHARDIA, Benth. & Hook. f. Gen. i. 1007 (1867) = BURGHARTIA, Scop. = **Piriqueta**, Aubl. (Turnerac.).

BURLINGTONIA, Lindl. Bot. Reg. t. 1927 (1837) = **Rodriguezia**, Ruiz & Pav. (Orchid.).
Batemani, Linden, Illustr. Hortic. xxviii. (1881) 64 = R. Batemanni.
candida, Lindl. Bot. Reg. xxiii. (1837) sub t. 1927 = R. candida.
carnea, Lindl. ex Linden, Illustr. Hortic. xxviii. (1881) 80 = R. carnea.
decora, Lem. Jard. Fleur. ii. (1852) Misc. 96 = R. decora.
fragrans, Lindl. Bot. Reg. xxiii. (1837) sub t. 1927 = R. fragrans.
granadensis, Lindl. Orchid. Linden. 24 = R. granadensis.
maculata, Lindl. Bot. Reg. xxv. (1839) 44 = R. maculata.
obtusifolia, Lindl. Sert. Orch. (1840) sub t. 36 = R. obtusifolia.
pubescens, Lindl. & Paxt. Flow. Gard. i. (1851) 158 = R. pubescens.
refracta, Lindl. ex Linden, Illustr. Hortic. xxviii. (1881) 80 = R. refracta.
gida, Lindl. ex Lem. in Fl. des Serres, i. (1845) 11. t. 2 = R. rigida.
rubescens, Lindl. Bot. Reg. xxiii. (1837) sub t. 1927.—Peruv.
scintillulans, Lem. Jard. Fleur. iii. (1852-53) Misc. 66.—Bras.
venusta, Lindl. ex Lem. Illustr. Hortic. v. (1858) t. 188 = R. venusta.

BURMANNIA, Linn. Syst. ed. 1 (1735). *BURMANNIACEAE*, Benth. & Hook. f. iii. 457.
CYANANTHUS, Miers, in Wall. Cat. nn. 9007-8 (1849).
CYANOTIS, Miers, l. c. n. 9006 (1849).
CRYPTONEMA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) 1. 590.
GONIANTHES, Blume, Cat. Gew. Buitenz. (1823) 19.
MABURNIA, Thou. Gen. Nov. Madag. 4 (1806).
NEPHROCOELIUM, Turcz. in Bull. Soc. Nat. Mosc. (1853) 1. 287.
TETRAPTERA, Miers, in Lindl. Veg. Kingd. 172 (1847).
TRIPTERANTHUS, Wall. ex Miers, l. c. (1847).
TRIPTERELLA, Michx. Fl. Bor. Am. i. 19. t. 3 (1803).
VOGELIA, J. F. Gmel. Syst. 107 (1791).
alba, *Mart. Nov. Gen. et Sp.* i. 12.—Bras.
azurea, Griff. Notul. iii. 236 = coelestis.
bancana, *Miq. Fl. Ind. Bat. Suppl.* 617.—Ins. Banca.
bicolor, *Mart. Nov. Gen. et Sp.* i. 10. t. 5.—Bras.
biflora, *Linn. Sp. Pl.* 287.—Am. bor.
brachyphylla, *Willd. ex Schult. f. Syst.* vii. II. p. lxxv.—Venezuela.
brachystachya, *Miq. in Linnaea*, xix. (1847) 141.—Guiana.
capensis, *Mart. Nov. Gen. et Sp.* i. 12.—Afr. austr.
capitata, *Mart. l. c.*—Am. bor.
Championii, *Thw. Enum. Pl. Zeyl.* 325.—Zeylan.
coelestis, *D. Don, Prod. Fl. Nep.* 44.—As. trop.; Austral.
dasyantha, *Mart. Nov. Gen. et Sp.* i. 11.—Bras.
distachya, R. Br. Prod. 265 = disticha.
disticha, *Linn. Sp. Pl.* 287.—As. trop.; Austral.
flava, *Mart. Nov. Gen. et Sp.* i. 11. t. 5. f. 2.—Bras.
flavula, *Wright, in Sauv. Fl. Cub.* 165.—Cuba.
Geelvinkiana, *Becc. Malesia*, i. (1878) 244.—Malaya.
javanica, *Blume, Enum. Pl. Jav.* fasc. i. 28.—Java.
juncea, *Soland. ex R. Br. Prod.* 265.—Austral.
Kalbreyeri, *Oliver, in Hook. Ic. Pl.* t. 1357.—N. Granat.
longifolia, *Becc. Malesia*, i. (1878) 244.—Malaya.
lutescens, *Becc. l. c.* 246.—Malaya.
madagascariensis, *Baker, in Journ. Linn. Soc.* xx. (1884) 268.—Madag.
madagascariensis, *Mart. & Zucc. Nov. Gen. et Sp.* i. 12.—Madag.
paniculata, *Willd. ex Schult. f. Syst.* vii. II. p. lxxiv.—Madag.
pusilla, *Thw. Enum. Pl. Zeyl.* 325.—Zeylan.
quadriflora, *Willd. ex Schult. f. Syst.* vii. 72. p. lxxiv.—Guiana.
selebica, *Becc. Malesia*, i. (1878) 243.—Ins. Celebes
Sellowiana, *Seub. in Mart. Fl. Bras.* iii. 57. t. 7. f. 2.—Bras.
sphagnoides, *Becc. Malesia*, i. (1878) 246.—Malaya.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 617 = disticha.
tenella, *Benth. in Hook. Kew Journ.* vii. (1855) 12.—Bras.
tridentata, *Becc. Malesia*, i. (1878) 246.—Malaya.

BURMANNIA :—
triflora, Roxb. Hort. Beng. [87]; Fl. Ind. ii. 117 = coelestis.
tuberosa, *Becc. Malesia*, i. (1878) 245.—Malaya.
uniflora, Rottl. ex Spreng. Syst. iv. Cur. Post. 23 = coelestis.

BURMEISTERA, Karst. & Triana, in Linnaea, xxviii. (1856) 444. *CAMPANULACEAE*, Benth. & Hook. f. ii. 547.
acuminata, *Karst. l. c.* 445.—N. Granat.
glabrata, *Benth. & Hook. f. Gen.* ii. 547.—N. Granat.
ibaguensis, *Karst. in Linnaea*, xxviii. (1856) 444.—N. Granat.
lacerata, *Karst. l. c.* 445.—N. Granat.
Matthiaei, *Benth. & Hook. f. Gen.* ii. 547.—Peruv.
marginata, *Karst. in Linnaea*, xxviii. (1856) 445.—N. Granat.
pomifera, *Karst. l. c.*—N. Granat.
succulenta, *Karst. l. c.*—N. Granat.
tomentosula, *Karst. l. c.*—N. Granat.
virescens, *Benth. & Hook. f. Gen.* ii. 547.—Guatemala.

BURNATIA, Micheli, in DC. Monog. Phan. iii. 81 (1881). *ALISMACEAE*, Benth. & Hook. f. iii. 1007.
enneandra, *Micheli, l. c.*—Nubia.

BURNETTIA, Lindl. Gen. et Sp. Orch. 517 (1840). *ORCHIDEAE*, Benth. & Hook. f. iii. 612.
cuneata, *Lindl. l. c.* 518.—Tasmania.

BURNEYA, Cham. & Schlecht. in Linnaea, iv. (1829) 189 = **Timonius**, Rumph. et **Bobea**, Gaudich. (Rubiac.).
Forsteri, Cham. & Schlecht. l. c. = T. Forsteri.
Gaudichaudii, Cham. & Schlecht. l. c. 190 = B. Gandichaudi.

BURRIELIA, DC. Prod. v. 663 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 398.
anthemoides, *A. Gray, in Torr. Bot. Mex. Bound.* 97.—Am. bor.
chrysostoma, Torr. & Gray, Fl. N. Am. ii. 379 = Baeria chrysostoma.
Fremonti, Benth. Pl. Hartw. 317 = Baeria Fremontii.
gracilis, DC. Prod. v. 664 = Baeria gracilis.
hirsuta, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 381 = Baeria gracilis.
lanosa, A. Gray, in Pacif. Rail. Rep. iv. 107 = Eriophyllum lanosum.
leptalea, A. Gray, in Proc. Am. Acad. vi. (1864-65) 546 = Baeria leptalea.
longifolia, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 380 = Baeria gracilis.
maritima, A. Gray, in Proc. Am. Acad. vii. (1868) 358 = Baeria maritima.
microglossa, *DC. Prod.* v. 664.—Calif.
nivea, D. Eaton, in Bot. King, Exp. 174 = Eatonella nivea.
parviflora, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 381 = Baeria gracilis.
platycarpha, A. Gray, in Torr. Bot. Mex. Bound. 97 = Baeria platycarpha.
tenerrima, DC. Prod. v. 664 = Baeria gracilis ?

BURSA, [Weber, in] Wigg. Prim. Fl. Hols. 47 (1780) = **Capsella**, Medic. (1792) (Crucif.).
astoris, [Weber,] l. c. = C. Bursa-pastoris.

BURSA-PASTORIS, [Tourn.] Rupp. Fl. Jen. 77. ed. Hall. (1745) 87 = **Capsella**, Medic. (1792) (Crucif.).

BURSAIA, Steud. Nom. ed. II. i. 241 (1840) sphalm. = **Burasaia**, Thou. (Menisp.).

BURSARIA, Cav. Ic. iv. 30. t. 350 (1797). *PITTOSPOREAE*, Benth. & Hook. f. i. 132.
diosmoides, Putterl. Syn. Pittosp. 20 = Marianthus procumbens.
incana, Lindl. in Mitch. Journ. Trop. Austral. 224 = spinosa.
inermis, *Blanco, Fl. Filip.* ed. II. 124.—Ins. Philipp.
procumbens, Putterl. Syn. Pittosp. 20 = Marianthus procumbens.
spinosa, *Cav. Ic.* iv. 30. t. 350.—Austral.
Stuartiana, F. Muell. ex Klatt, in Linnaea, xxviii (1856) 586 = Marianthus procumbens.

BURSERA, ' Jacq. ' ex Linn. Sp. Pl. ed. II. 471 (1762).
BURSERACEAE, Benth. & Hook. f. i. 324.
BUSSERIA, Cramer, Disp. Syst. 144 (1803).
DAMMARA, Gaertn. Fruct. ii. 100. t. 103 (1791).
ELAPHRIUM, Jacq. Enum. Pl. Carib. 3 (1760); Stirp. Am. i. 105. t. 71 (1763), nomen prius.
ICICA, Aubl. Pl. Gui. i. 337. t. 130 ad 135 (1776).
MARIGNIA, Comm. ex Kunth, in Ann. Sc. Nat. Ser. I. ii. (1824) 351.
acuminata, Willd. Sp. Pl. iv. 1120 = Dacryodes hexandra.
acutifolia, Sieber, ex Steud. Nom. ed. II. i. 241 = Sorindeia madagascariensis.
Aloëxylon, *Engl. in DC. Monog. Phan.* iv. 52.—Mexic.
altissima, *Baill. Hist. Pl.* v. 297.—Guiana.
angustata, *Griseb. Cat. Pl. Cub.* 65.—Cuba.
bahiensis, Salzm. ex Engl. in DC. Monog. Phan. iv. 277, 537 = Tapiria guianensis.
balsamifera, Pers. Syn. i. 413 = Hedwigia balsamifera.
bicolor, *Engl. in DC. Monog. Phan.* iv. 53.—Mexic.
bipinnata, *Engl. l. c.* 49.—Mexic.
caudata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 614 = Protium Sagotianum.
cinerea, *Engl. in DC. Monog. Phan.* iv. 43.—Mexic.
cuneata, *Engl. l. c.* 56.—Mexic.
decandra, *Baill. Hist. Pl.* v. 262, 297.—Guiana.
Delpechiana, *Poiss. ex Engl. in DC. Monog. Phan.* iv. 53.—Mexic.
excelsa, *Engl. l. c.* 57.—Mexic.
fagaroides, *Engl. l. c.* 48.—Mexic.
Galeottiana, *Engl. l. c.* 47.—Mexic.
glabra, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 304.—N. Granat.
glauca, *Griseb. Cat. Pl. Cub.* 66.—Cuba.
gracilis, *Engl. in DC. Monog. Phan.* iv. 50.—Mexic.
grandifolia, *Engl. l. c.* 45.—Mexic.
graveolens, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 303.—Peruv.
guianensis, *Baill. Hist. Pl.* v. 297.—Guiana.
gummifera, *Linn. Sp. Pl.* ed. II. 471.—Am. trop.
heptaphylla, *Wright, in Sauv. Fl. Cub.* 20.—Cuba.
heterophylla, *Engl. in DC. Monog. Phan.* iv. 46.—Mexic.
Hindsiana, *Engl. l. c.* 58.—N. Granat.
Icicariba, *Baill. Hist. Pl.* v. 297.—Bras.
integerrima, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 303.—N. Granat.
javanica, *Baill. Hist. Pl.* v. 297.—Java.
jorullensis, *Engl. in DC. Monog. Phan.* iv. 57.—Mexic.
Karsteniana, *Engl. l. c.* 41.—Venezuela.
Karwinskii, *Engl. l. c.*—Mexic.
Kerberi, *Engl. l. c.*—Mexic.
lancifolia, *Engl. l. c.* 42.—Mexic.
lanuginosa, *Engl. l. c.* 58.—Mexic.
leptophloeos, *Mart. in Flora*, xi. (1828) ii. 393.—Bras.
macrostachya, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 613.—Guiana.
Martiana, *Engl. in Mart. Fl. Bras.* xii. ii. 253.—Bras.
mexicana, *Engl. in DC. Monog. Phan.* iv. 51.—Mexic.
microphylla, *A. Gray, in Proc. Am. Acad.* v. (1861) 155.—Mexic.
multijuga, *Engl. in DC. Monog. Phan.* iv. 42.—Mexic.
obovata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 614.—Mexic.
obtusifolia, Lam. Encyc. ii. 768 = Protium obtusifolium.
orinocensis, *Engl. in Mart. Fl. Bras.* xii. ii. 254.—Bras.
ovalifolia, *Engl. in DC. Monog. Phan.* iv. 40.—Mexic.
paniculata, Lam. Encyc. ii. 768 = Canarium mauritianum.
pannosa, *Engl. in DC. Monog. Phan.* iv. 54.—Mexic.
penicillata, *Engl. l. c.* 52.—Mexic.
Schiedeana, *Engl. l. c.* 57.—Mexic.
Schlechtendalii, *Engl. l. c.* 41.—Mexic.
Sellowii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 613.—Bras.
serrata, Wall. ex Coleb. in Trans. Linn. Soc. xv. (1827 361. t. 4 = Protium serratum.

BURSERA:—
sessiliflora, *Engl. in DC. Monog. Phan.* iv. 55.—Mexic.
simplicifolia, *DC. Prod.* ii. 78.—Jamaic.
submoniliformis, *Engl. in DC. Monog. Phan.* iv. 55.—Mexic.
Tacahamaca, Baill. Hist. Pl. v. 297 = Protium heptaphyllum.
Tatamaco, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xiv. (1872) 303.—N. Granat.
tomentosa, *Triana & Planch. l. c.* 304.—N. Granat.

BURSERIA, Loefl. It. Hisp. 194 (1758) = **Verbena**, Linn.

BURSERIA, Jacq. Stirp. Am. 94. t. 65 (1763) = **Bursera**, Linn.

BURSHIA, Rafin. in Med. Repos. N. York, v. (1808) 357 = **Myriophyllum**, Linn. (Haloragˌ).
humilis, Rafin. l. c. et vi. (1808) 422 = M. ambiguum.

BURSINOPETALUM, Wight, Ic. t. 956 (1847) = **Mastixia**, Blume (Cornac.).
arboreum, Wight, l. c. = M. arborea.
tetrandrum, Wight, ex Thw. Enum. Pl. Zeyl. 42 = M. tetrandra.

BURTONIA, R. Br. in Ait. Hort. Kew. ed. II. iii. 12 (1811). *LEGUMINOSAE*, Benth. & Hook. f. i. 468.
WEIHEA, Reichb. Consp. 212 [b] (1828).
conferta, *DC. Prod.* ii. 106.—Austral.
diosmaefolia, Benth. Enum. Pl. Hueg. 82 = Latrobea diosmifolia.
foliolosa, *Benth. Fl. Austral.* ii. 50.—Austral.
gompholobioides, *F. Muell. Fragm.* x. 34.—Austral.
Hendersonii, *Benth. Fl. Austral.* ii. 51.—Austral.
minor, DC. Prod. ii. 106 = Gompholobium minus.
polyzyga, *Benth. Fl. Austral.* ii. 51.—Austral.
pulchella, Meissn. in Lehm. Pl. Preiss. i. 41 = scabra.
scabra, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 12.—Austral.
sessilifolia, DC. Prod. ii. 106 = scabra.
solanacea, *Hort. ex Steud. Nom.* ed. II. i. 241.—Austral.
subalpina, F. Muell. in Trans. Phil. Soc. Victoria, i. (1855) 39 = Pultenaea rosea.
subulata, *Benth. Fl. Austral.* ii. 50.—Austral.
tenella, Meissn. in Lehm. Pl. Preiss. i. 42 = Latrobea tenella.
villosa, *Meissn. l. c.* 41.—Austral.

BURTONIA, Salisb. Parad. Lond. t. 73 (1807) = **Hibbertia**, Andr. (Dillen.).
grossulariaefolia, Salisb. l. c. = H. grossulariaefolia.

BUSBECKEA, Endl. Prod. Fl. Ins. Norf. 64 (1833) = **Capparis**, Tourn.
arborea, F. Muell. Fragm. i. 163 = C. nobilis.
canescens, F. Muell. l. c. v. 104 = C. canescens.
corymbiflora, F. Muell. l. c. vii. 163 = C. lucida.
humistrata, F. Muell. l. c. v. 156 = C. humistrata.
lorantifolia, F. Muell. l. c. i. 164 = C. loranthifolia.
Mitchellii, F. Muell. l. c.; et Vict. Pl. i. 53 = C. Mitchellii.
nobilis, Endl. Prod. Fl. Ins. Norf. 64 = C. ornans.
ornans, F. Muell. Fragm. v. 104 = C. ornans.
Shanesiana, F. Muell. l. c. x. 94 = C. Shanesiana.
Thozetiana, F. Muell. l. c. v. 104 = C. Thozetiana.
umbonata, F. Muell. l. c. i. 164; v. 104 = C. umbonata.

BUSBECKEA, Mart. Cat. Hort. Monac. (1829) 69 = **Salpichroa**, Miers (Solanac.).
radicans, Mart. l. c. = S. rhomboidea.

BUSBEQUIA, Salisb. Gen. Pl. Fragm. 25 (1866) = **Hyacinthus**, Tourn. (Liliac.).
ciliaris, Salisb. l. c. = H. ciliatus.

BUSEA, Miq. Fl. Ind. Bat. ii. 732 (1856) = **Cyrtandromoea**, Zoll. (Gesnerac.).
acuminata, Miq. l. c. 733 = C. Miqueliana.
angustifolia, Miq. l. c. 734 = C. angustifolia.
asperifolia, Miq. l. c. 733 = Dichrotrichum asperifolium.

BUSEA:—
decurrens, Miq. Fl. Ind. Bat. ii. 733 = C. decurrens.
subsessilis, Miq. l. c. = C. acuminata.

BUSIPHO, Salisb. Gen. Pl. Fragm. 76 (1866) = **Aloë**, Tourn. (Liliac.).
ferox, Salisb. l. c. = A. ferox.

BUSSERIA, Cramer, Disp. Syst. 144 (1803) = **Bursera**, Jacq.

BUSTAMENTA, Alam. ex DC. Prod. v. 166 (1836) = **Eupatorium**, Tourn. (Compos.).
cordata, Alam. ex DC. l. c. = E. petiolare.
ovata, Mairet, ex DC. l. c. 168 = E. Bustamenta.

BUSTELINA, Fourn. in Mart. Fl. Bras. vi. iv. 288 (1885). *ASCLEPIADEAE*.
Warmingii, *Fourn. l. c.*—Bras.

BUSTIA, Adans. Fam. ii. 127 (1763) = **Buphthalmum**, Linn. (Compos.).

BUSTILLOSIA, Clos, in C. Gay, Fl. Chil. iii. 106. t. 32 (1847) = **Asteriscium**, Cham. (Umbellif.).
chilensis, Clos, l. c. 108 = A. chilense.

BUTEA, Koen. ex Roxb. Pl. Corom. i. 22. t. 21 (1795). *LEGUMINOSAE*, Benth. & Hook. f. i. 533.
MEGALOTROPIS, Griff. Notul. iv. 441 (1854).
MEIZOTROPIS, Voigt, Hort. Suburb. Calc. 239 (1845).
PLASO, Adans. Fam. ii. 325 (1763).
acuminata, Wall. Cat. n. 5443 = Spatholobus acuminatus.
Braamiana, *DC. Prod.* ii. 415.—China.
frondosa, *Roxb. Pl. Corom.* i. 21.—Ind. or.; Burma.
frondosa, Wall. Cat. n. 5569, partim = superba.
gyrocarpa, R. Grah. in Wall. l. c. n. 5442 = Spatholobus gyrocarpus.
Loureirii, *Spreng. Syst.* iii. 186.—Cochinch.
minor, *Buch.-Ham. ex Wall. Cat.* n. 5439.—Reg. Himal.
parviflora, Roxb. Hort. Beng. 53 = Spatholobus Roxburghii.
peltata, Pers. Syn. ii. 279 = Rudolphia peltata.
sericophylla, Wall. Cat. n. 5441 = Spatholobus Roxburghii.
superba, *Roxb. Pl. Corom.* i. t. 22.—Ind. or.
volubilis, Pers. Syn. ii. 279 = Rudolphia volubilis.

BUTERAEA, Nees, in Wall. Pl. As. Rar. iii. 75, 196 (1832) = **Strobilanthes**, Blume (Acanthac.).
rhamnifolia, Nees, in DC. Prod. xi. 725 = S. rhamnifolius.
ulmifolia, Nees, in Wall. Pl. As. Rar. iii. 84 = S. rufescens.

BUTINIA, Boiss. Elench. 53 (1838) = **Conopodium**, Koch (Umbellif.).
bunioides, Boiss. l. c. 54 = Conopodium butinioides.
capnoides, Decne. in Jacquem. Voy. Bot. 70 = Chaerophyllum capnoides.
cretica, Boiss. & Heldr. Diagn. Ser. I. x. 50 = Carum creticum.
glaucescens, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 61 (= *Elaeosticta glaucescens*).—Persia.
libanotica, Boiss. l. c. 62 (= *Elaeosticta meifolia*).—Syria.
macrocarpa, Boiss. & Sprun. in Ann. Sc. Nat. Sér. III. ii. (1844) 62 = Carum macrocarpum.
nodosa, Boiss. l. c. 61 (= *Elaeosticta nodosa*).—Persia.
parnassica, Heldr. ex Nym. Consp. 302 (= *Biasolettia parnassica*).—Graecia.
stylosa, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 63 (= *Freyera stylosa*).—Graecia.

BUTNERIA, Duham. Arb. i. 113. t. 45 (1755) = **Calycanthus**, Linn.

BUTOMISSA, Salisb. Gen. Pl. Fragm. 91 (1866) = **Allium**, Tourn. (Liliac.).
tatarica, Salisb. l. c. = A. tataricum.

BUTOMOPSIS, Kunth, Enum. Pl. iii. 164 (1841). *ALISMACEAE*, Benth. & Hook. f. iii. 1008.
TENAGOCHARIS, Hochst. in Flora, xxiv. (1841) 369.
cordofana, Kunth, ex Walp. Ann. i. 769 = lanceolata.

BUTOMOPSIS:—
lanceolata, *Kunth, Enum. Pl.* iii. 165.—Austral.
latifolia, Kunth, l. c. = lanceolata.

BUTOMUS, Linn. Syst. ed. I (1735). *ALISMACEAE*, Benth. & Hook. f. iii. 1008.
Caesalpini, Neck. Delic. Gallo-Belg. i. 189 = umbellatus.
floridus, Gaertn. Fruct. i. 74. t. 19. f. 3 = umbellatus.
junceus, Turez. in Bull. Soc. Nat. Mosc. (1837) n. VII. 157; (1838) 101, nomen; et xxvii. (1854) II. 60 = umbellatus.
lanceolatus, Roxb. Hort. Beng. 89; Fl. Ind. ii. 315 = Butomopsis lanceolata.
latifolius, D. Don, Prod. Fl. Nep. 22 = Butomopsis lanceolata.
umbellatus, *Linn. Sp. Pl.* 372.—Europ.; As. temp.
vulgaris, Gueldenst. Iter, ii. 22 = umbellatus.

BUTONICA, Lam. Encyc. i. 515 (1783); Juss. Gen. 326 (1789) = **Barringtonia**, Forst. (Myrtac.).
acutangula, Lam. l. c. 591 = Barringt. acutangula.
alata, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 70 = Barringt. conoidea.
alba, Miers, l. c. 65 = Barringt. racemosa.
apiculata, Miers, l. c. 78.—Madag.
caffra, Miers, l. c. 77 = Barringt. racemosa.
calyptrata, Miers, l. c. 76 = Barringt. speciosa?
ceylanica, Miers, l. c. 77 = Barringt. ceylanica.
edulis, Miers, l. c. 76 = Barringt. edulis.
inclyta, Miers, l. c. 71 = Barringt. racemosa.
intermedia, Miers, l. c. 73 = Barringt. intermedia.
procera, Miers, l. c. 74.—Ins. Pacif.
racemosa, Juss. Gen. 326; Miers, l. c. 65 = Barringt. racemosa.
rosata, Miers, l. c. 72.—Ins. Philipp.
rubra, Miers, l. c. 70 = Barringt. racemosa.
Rumphiana, Miers, l. c. 68 = Barringt. Butonica.
samoënsis, Miers, l. c. 75 = Barringt. samoënsis.
speciosa, Lam. Encyc. i. 521 = Barringt. speciosa.
terrestris, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 69 = Barringt. elongata.

BUTTNERIA, Duham. Arb. ed. II. 217. t. 47 (1801) = **Calycanthus**, Linn.

BUTTNERIA, Schreb. Gen. i. 145 (1789) = **Buettneria**, Loefl.

BUTTONIA, MacKen, ex Benth. in Hook. Ic. Pl. xi. 63. t. 1080 (1871). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 970.
natalensis, *MacKen, ex Benth. in Hook. l. c.*—Afr. austr.
natalis, MacKen, ex Benth. in Hook. l. c. = natalensis.

BUTYROSPERMUM, Kotschy, in Sitzb. Acad. Wien, Math. Nat. l. Abth. I. (1865) 357. *SAPOTACEAE*, Benth. & Hook. f. ii. 661.
MICADANIA, R. Br. in Denh. et Clapp. Voy. App. 239 (1826).
Kirkii, *Baker, in Oliver, Fl. Trop. Afr.* iii. 504.—Afr. trop.
niloticum, Kotschy, in Sitzb. Acad. Wien, Math. Nat. l. Abth. I. (1865) 358. t. 1. = seq.
Parkii, *Kotschy, l. c.* 359. t. 2.—Afr. trop.

BUXUS, Linn. Syst. ed. I (1735). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 266.
CRANTZIA, Sw. Prod. Veg. Ind. Occ. 38 (1788).
TRICERA, Schreb. Gen. 630 (1791); Sw. Fl. Ind. Occ. 331. t. 7 (1797).
acuminata, *Muell. Arg. in DC. Prod.* xvi. I. 15.—Cuba.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 2 = sempervirens.
aquartiana, Rich. ex Baill. Monog. Bux. 69 = citrifolia.
arborescens, Mill. Gard. Dict. ed. VIII. n. 1; Lam. Fl. Fr. ii. 203 = sempervirens.
argentea, Hort. ex Steud. Nom. ed. II. i. 242 = sempervirens.
aurea, Hort. ex Steud. l. c. = sempervirens.
australis, *A. Cunn. ex Steud. l. c.*—Austral.
balearica, *Lam. Encyc.* i. 511.—Ins. Balear.
californica, Hort. ex Baill. Monog. Bux. 66 = Simmondsia californica?

BUXUS:—
caucasica, Hort. ex C. Koch, Dendrol. ii. II. 476 = sempervirens.
chinensis, Link, Enum. Hort. Berol. ii. 386 = Simmondsia californica.
citrifolia, *Spreng. Syst.* iii. 847.—Am. austr.
cordifolia, Spreng. Syst. iii. 847 = Tricera cordifolia.
coriacea, Spreng. Syst. iv. Cur. Post. 314 = Sarcococca pruniformis.
crispa, Hort. ex C. Koch, Dendrol. ii. II. 476 = sempervirens.
cruciata, Rich. ex Baill. Monog. Bux. 67 = Vahlii.
cubana, *Baill. l. c.* 71.—Cuba.
cucullata, Hort. ex C. Koch, Dendrol. ii. II. 476 = sempervirens.
dioica, Forsk. Fl. Aegypt. Arab. 159 = Myrsine bottensis.
elegantissima, Hort. ex C. Koch, Dendrol. ii. II. 477 = sempervirens.
Fortunei, *Carr. in Rev. Hortic.* (1870–71) 519.—China.
glomerata, *Muell. Arg. in DC. Prod.* xvi. I. 17.—Cuba.
gonoclada, *Muell. Arg. l. c.* 16.—Cuba.
haleppica, Hort. ex C. Koch, Dendrol. ii. II. 477 = longifolia.
Handsworthii, Hort. ex C. Koch, l. c. 476 = sempervirens.
Harlandi, *Hance, in Journ. Linn. Soc.* xiii. (1873) 123.—China.
Hildebrandtii, *Baill. Adansonia*, xi. (1873–76) 268.—Afr. trop.
japonica, *Muell. Arg. in DC. Prod.* xvi. I. 20.—Japon.
lacvigata, *Spreng. Syst.* iii. 847.—Jamaic.
longifolia, *Boiss. Diagn.* Ser. I. xii. 107.—Syria.
madagascarica, *Baill. Monog. Bux.* 65.—Madag.
marginata, Hort. ex Steud. Nom. ed. II. i. 242 = sempervirens.
microphylla, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1846) 142 = japonica.
mucronata, Hort. ex Baill. Monog. Bux. 62 = sempervirens.
myrtifolia, Lam. Encyc. i. 511 = sempervirens.
obcordata-variegata, Fortune, in Gard. Chron. (1861) 735 = microphylla.
ovalifolia, Siebold, ex C. Koch, Dendrol. ii. II. 479 = microphylla.
pulchella, *Baill. Monog. Bux.* 68.—Jamaic.
Purdieana, *Baill. l. c.* 70 (1859).—Jamaic.
retusa, *Muell. Arg. in DC. Prod.* xvi. I. 16.—Cuba.
rosmarinifolia, Hort. ex Baill. Monog. Bux. 62 = sempervirens.
rotundifolia, Hort. ex C. Koch, Dendrol. ii. II. 479 = microphylla.
salicifolia, Hort. ex C. Koch, l. c. 476 = sempervirens.
saligna, D. Don, Prod. Fl. Nep. 63 = Sarcococca pruniformis.
sempervirens, *Linn. Sp. Pl.* 983.—Europ.; Oriens; As. temp.
sempervirens, Thunb. Fl. Jap. 77 = japonica.
stenophylla, *Hance, in Journ. Bot.* vi. (1868) 331.—China.
subcolumnaris, *Muell. Arg. in DC. Prod.* xvi. I. 14.—Ind. occ.
suffruticosa, Mill. Gard. Dict. ed. VIII. n. 3 = sempervirens.
tenuifolia, Hort. ex Baill. Monog. Bux. 61 = sempervirens.
Vahlii, *Baill. Monog. Bux.* 67.—Ind. occ.
variegata, Hort. ex Steud. Nom. ed. II. i. 242 = sempervirens.
virens, Thunb. Fl. Jap. 77 = microphylla.
Wallichiana, Baill. Monog. Bux. 63 = sempervirens.
Wrightii, *Muell. Arg. in DC. Prod.* xvi. I. 17.—Cuba.

BYBLIS, Salisb. Parad. Lond. t. 95 (1808). *DROSERACEAE*, Benth. & Hook. f. i. 664.
caerulea, R. Br. ex Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 306 = linifolia.
filifolia, Planch. l. c. 305 = linifolia.
gigantea, *Lindl. Swan River App.* 92.—Austral.
Lindleyana, Planch. in Ann. Sc. Nat. Sér. III. ix. (1848) 307 = gigantea.
linifiora, *Salisb. Parad. Lond.* t. 95.—Austral.

BYGNONIA, Barcena, in La Naturaleza, ii. (1873) 173 = **Bignonia**, Linn.

BYRONIA, Endl. in Ann. Wien. Mus. i. (1836) 184. *ILICINEAE*, Benth. & Hook. f. i. 357.
POLYSTIGMA, Meissn. Gen. 252; Comm. 161 (1839).
arnhemensis, *F. Muell. Fragm.* ii. 119.—Austral.
Hookeri, Steud. Nom. ed. II. i. 242 = seq.
sandwicensis, *Endl. in Ann. Wien. Mus.* i. (1836) 184.—Ins. Sandvic.
taitensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 297. t. 25.—Ins. Tahiti.

BYRSA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. i. (Gen. ignot.).

BYRSANTHES, Presl, Prod. Monog. Lobel. 41 (1836) = **Siphocampylus**, Pohl (Campanul.).
Halliana, Presl, l. c. 42 = S. Hallianus.
Humboldtiana, Presl, l. c. = S. niveus.
nivea, A. DC. in DC. Prod. vii. 408 = S. niveus.

BYRSANTHUS, Guillem. in Deless. Ic. Sel. iii. 30. t. 52 (1837). *SAMYDACEAE*, Benth. & Hook. f. i. 800.
ANETIA, Endl. Gen. 923 (1839).
Brownii, *Guillem. l. c.*—Afr. trop.
epigynus, *Mast. in Oliver, Fl. Trop. Afr.* ii. 499.—Afr. trop.

BYRSOCARPUS, Schum. & Thonn. Beskr. Guin. Pl. 226 (1827). *CONNARACEAE*, Benth. & Hook. f. i. 431.
Boivinianus, Baill. Adansonia, vii. (1866) 231 = Rourea Boiviniana.
coccineus, *Schum. & Thonn. Beskr. Guin. Pl.* 226.—Afr. trop.
maximus, *Baker, in Oliver, Fl. Trop. Afr.* i. 453.—Afr. trop.
orientalis, *Baill. Adansonia*, vii. (1866) 230.—Afr. trop.
ovatifolius, *Baker, in Oliver, Fl. Trop. Afr.* i. 452.—Afr. trop.
parviflorus, Baill. Adansonia, vii. (1866) 229 = coccineus.
parvifolius, Planch. in Linnaea, xxiii. (1850) 412 = coccineus.
Pervilleanus, Baill. Adansonia, vii. (1866) 232 = Rourea Pervilleana.
puniceus, Schum. & Thonn. Beskr. Guin. Pl. 227 = coccineus.

BYRSONIMA, Rich. ex Juss. in Ann. Mus. Par. xviii. (1811) 481. *MALPIGHIACEAE*, Benth. & Hook. f. i. 251.
adenophylla, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 314.—N. Granat.
Aerugo, *Sagot, l. c.* Sér. VI. xii. (1881) 178.—Guiana.
altissima, *DC. Prod.* i. 579.—Guiana.
amazonica, *Griseb. in Mart. Fl. Bras.* xii. I. 14.—Bras.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* v. 153. t. 449.—Venezuela.
arbutifolia, *Griseb. in Mart. Fl. Bras.* xii. I. 14.—Bras.
arthropoda, A. Juss. in Arch. Mus. Par. iii. (1843) 307 = laxiflora.
basiloba, *A. Juss. l. c.* 279.—Bras.
Berteroana, A. Juss. l. c. 293 = spicata.
bicorniculata, *A. Juss. l. c.* 284.—Bras.
biflora, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 167.—Cuba.
Blanchetiana, Miq. in Linnaea, xxii. (1849) 799 = bicorniculata.
brachystachya, *DC. Prod.* i. 581.—Bras.
brachystachya, Poepp. ex Griseb. in Mart. Fl. Bras. xii. I. 10 = crispa.
bracteolaris, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 123.—Guiana.
brasiliensis, Griseb. in Linnaea, xiii. (1839) 255 = sericea.
bumeliaefolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 79.—Bras.
ceranthera, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 121.—Guiana.
chrysophylla, *H. B. & K. Nov. Gen. et Sp.* v. 150.—Venezuela.
chrysophylla, A. Juss. in St. Hil. Fl. Bras. Mer. iii. 81 = sericea.
cinerea, *DC. Prod.* i. 580.—Am. trop.
Clausseniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 281.—Bras.

BYRSONIMA:—
coccolobaefolia, *H. B. & K. Nov. Gen. et Sp.* v. 148.—Am. austr.
coleostachya, *Griseb. in Linnaea*, xxii. (1849) 3.—Guiana.
concinna, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 122.—Guiana.
coniophylla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 310.—Bras.
coriacea, DC. Prod. i. 580 = spicata.
correaefolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 78.—Bras.
cotinifolia, Benth. Pl. Hartw. 277 = oaxacana.
cotinifolia, *H. B. & K. Nov. Gen. et Sp.* v. 152. t. 447.—Mexic.
crassifolia, *H. B. & K. Nov. Gen. et Sp. Pl.* v. 149.—Guiana.
crassifolia, Lunan, ex Griseb. Fl. Brit. W. Ind. 114 = cinerea.
crassifolia, Steud. Nom. ed. II. i. 242 = verbascifolia.
crispa, *A. Juss. in Arch. Mus. Par.* iii. (1843) 306.—Bras.
cujabensis, *A. Juss. ex Mart. Herb. Fl. Bras.* 344. n. 657, nomen.—Bras.
cubensis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 292.—Cuba.
Cumingiana, A. Juss. l. c. 282 = crassifolia.
cuprea, *Griseb. in Mart. Fl. Bras.* xii. 1. 19.—Bras.
cydoniaefolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 77.—Bras.
cydoniaefolia, Mart. in Flora, xxiv. (1841) Beibl. ii. 104 = Clausseniana.
dealbata, *Griseb. in Linnaea*, xiii. (1839) 253.—Bras.
densa, *DC. Prod.* i. 580.—Guiana.
diphylla, *DC. l. c.* 581.—N. Granat.
dispar, *Griseb. in Mart. Fl. Bras.* xii. 1. 10.—Bras.
eglandulosa, A. Juss. in Arch. Mus. Par. iii. (1843) 273 = sericea.
elegans, *DC. Prod.* i. 580.—Guiana.
eriopoda, *DC. l. c.* 579.—Am. austr.
fastigiata, Mart. in Flora, xxiv. (1841) II. Beibl. 61 = laevigata.
Fendleri, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 582.—Panama.
ferruginea, H. B. & K. Nov. Gen. et Sp. v. 151 = crassifolia.
frondosa, *Mart. ex A. Juss. in Arch. Mus. Par.* iii. (1843) 303.—Guiana.
Gardnerana, *A. Juss. l. c.* 296.—Bras.
guadalupensis, *G. Don, Gen. Syst.* i. 637.—Ind. occ.
Guilleminiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 276.—Bras.
gymnocalycina, *A. Juss. l. c.* 304.—Guiana.
Hostmanni, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 120.—Guiana.
hypargyrea, *Planch. ex Linden, Cat.* n. 8 (1853) 2.—N. Granat.
hypoleuca, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) I. 391.—N. Granat.
intermedia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 82.—Bras.
inundata, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 122.—Bras.
japurensis, *A. Juss. in Arch. Mus. Par.* iii. (1843) 305.—Bras.
Karwinskiana, *A. Juss. l. c.* 288.—Mexic.
laevigata, *DC. Prod.* i. 580.—Guiana.
lanceolata, DC. l. c. = crassifolia.
lancifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 275.—Bras.
laurifolia, *H. B. & K. Nov. Gen. et Sp.* v. 147.—Venezuela.
laxiflora, *Griseb. in Linnaea*, xiii. (1839) 256.—Bras.
leucophlebia, *Griseb. in Mart. Fl. Bras.* xii. 1. 15.—Bras.
ligustrifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 82.—Bras.
ligustrifolia, Mart. Herb. Fl. Bras. n. 1176 = intermedia.
linearifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 279.—Bras.
longibracteata, A. Juss. ex Mart. in Flora, xxiv. (1841) II. Beibl. 61 = stipulacea.
lucida. *DC. Prod.* i. 580.—Am. bor.; Ind. occ.
Martiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 286.—Bras.
microphylla, *A. Juss. l. c.* 298.—Bras.
mollis, *DC. Prod.* i. 581.—Guiana.
montana, H. B. & K. Nov. Gen. et Sp. v. 149 = crassifolia.

BYRSONIMA:—
monticola, A. Juss. & Mart. ex Griseb. in Mart. Fl. Bras. xii. 1. 17 = coccolobaefolia.
Moritziana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 391.—Venezuela.
Mourèila, Loud. Hort. Brit. 182 = crassifolia.
myricifolia, *Griseb. in Kjoeb. Vidensk. Meddel.* (1875) 125.—Bras.
nervosa, *DC. Prod.* i. 579.—Bras.
nitida, *D. Don, Gen. Syst.* i. 636.—Peruv.
nitidifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 296.—Bras.
nitidissima, *H. B. & K. Nov. Gen. et Sp.* v. 152.—Venezuela.
oaxacana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 283.—Mexic.
oblongifolia, *A. Juss. l. c.* 300.—Bras.
obversa, *Miq. Stirp. Surin. Sel.* 76.—Guiana.
Orbignyana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 284.—Bolivia.
oxyphylla, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 84.—Bras.
pachyphylla, Griseb. in Linnaea, xiii. (1839) 250 = verbascifolia.
pachyphylla, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 77.—Bras.
pallida, *DC. Prod.* i. 581.—Guiana.
panamensis, *Beurling, in Vet. Akad. Handl. Stockh.* (1854) 117.—Panama.
Paulista, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 85.—Bras.
perseaefolia, *Griseb. in Linnaea*, xiii. (1839) 257.—Bras.
peruviana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 289.—Peruv.
phlomoides, *G. Don, Gen. Syst.* i. 636.—Bras.
piauhensis, A. Juss. in Arch. Mus. Par. iii. (1843) 274 = sericea.
pinetorum, *Griseb. Cat. Pl. Cub.* 42.—Cuba.
Poeppigiana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 308.—Bras.
propinqua, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 120.—Guiana.
psilandra, *Griseb. in Linnaea*, xiii. (1839) 251.—Bras.
pulchra, *DC. Prod.* i. 580.—Mexic.
punctulata, *A. Juss. in Arch. Mus. Par.* iii. (1843) 309.—Bras.
reticulata, DC. Prod. i. 581 = Heteropteris platyptera.
reticulata, *Klotzsch & Karst. ex Griseb. in Linnaea*, xxii. (1849) 6.—N. Granat.
rhombifolia, *A. Juss. in Arch. Mus. Par.* iii. (1843) 290.—Bras.
rhopalaefolia, H. B. & K. Nov. Gen. et Sp. v. 148 = crassifolia.
rigida, *A. Juss. in Arch. Mus. Par.* iii. (1843) 298.—Bras.
rotunda, *Griseb. in Mart. Fl. Bras.* xii. 1. 20.—Bras.
rufescens, *Bertol. Fl. Guatim.* 18.—Guatemala.
rugosa, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 118.—Guiana.
Salzmanniana, *A. Juss. in Arch. Mus. Par.* iii. (1843) 277.—Bras.
Schomburgkiana, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 123.—Guiana.
sericea, *DC. Prod.* i. 580.—Bras.
sessilifolia, *Benth. in Hook. Lond. Journ. Bot.* vii. (1848) 124.—Guiana.
sparsibilis, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) I. 391.—Bras.
spicata, *Rich. ex Juss. in Ann. Mus. Par.* xviii. (1811) 481; *DC. Prod.* i. 580.—Am. trop.
spicata, Poepp. ex Spreng. Syst. ii. 383 = Heteropteris platyptera.
stigmatophora, *Schlecht. in Linnaea*, x. (1836) 241.—Mexic.
stipulacea, *A. Juss. in Arch. Mus. Par.* iii. (1843) 287.—Bras.
trinitensis, *A. Juss. l. c.* 296.—Ind. occ.
triopterifolia, *A. Juss. l. c.* 298.—Am. austr.
umbellata, *Mart. ex A. Juss. l. c.* 301.—Bras.
vacciniaefolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* i. 84.—Bras.
variabilis, *A. Juss. l. c.* 78.—Bras.
verbascifolia, *Rich. ex Juss. in Ann. Mus. Par.* xviii. (1811) 481; *DC. Prod.* i. 579.—Am. trop.
viminifolia, *A. Juss. in St. Hil. Fl. Bras. Mer.* iii. 83.—Bras.
volubilis, DC. Prod. i 581 = Hiraea Simsiana.

BYRSOPHYLLUM, Hook. f. in Benth. & Hook. f. Gen. ii. 83 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
ellipticum, *Hook. f. l. c.*—Zeylan.
tetrandrum, *Hook. f. l. c.*; *Bedd. Forest. Man.* 134/2.—Ind. or.

BYSTROPOGON, L'Hérit. Sert. Angl. 19. t. 22, 23 (1788). *LABIATAE*, Benth. & Hook. f. ii. 1184.
ASTEMON, Regel, Ind. Sem. Hort. Petrop. (1860) 30.
MINTOSTACHYS, Spach, Hist. Vég. Phan. ix. 164 (1840).
bipinnatus, Roth, Nov. Pl. Sp. 225 = Lavandula Burmanni.
canariensis, *L'Hérit. Sert. Angl.* 20.—Ins. Canar.
canus, *Benth. Lab. Gen. et Sp.* 326.—Peruv.
coarctatus, *Thonn. & Schum. in Vidensk. Selsk. Skr.* iv. (1828) 34 = Hyptis pectinata.
confertus, Willd. ex Steud. Nom. ed. II. i. 243 = Mentha mollis.
dentatus, *Poir. Encyc. Suppl.* i. 755.—Peruv.
erianthus, Benth. Lab. Gen. et Sp. 727 = Cuminia eriantha.
glabrescens, *Benth. Pl. Hartw.* 145.—Peruv.
graveolens, Blume, Bijdr. 824 = Hyptis suaveolens.
maderensis, *Webb & Berth. Phyt. Canar.* iii. 66.—Ins. Madera.
marifolius, Juss. ex Steud. Nom. ed. II. i. 243, 521 = Micromeria marifolia.
Meridiani, *C. Bolle, in Bonplandia*, viii. (1860) 281.—Ins. Canar.
mollis, *H. B. & K. Nov. Gen. et Sp.* ii. 317.—N. Granat.
mollis, Salisb. Prod. 80 = canariensis.
odoratissimus, *C. Bolle, in Bonplandia*, viii. (1860) 281.—Ins. Canar.
origanifolius, *L'Hérit. Sert. Angl.* 20.—Ins. Canar.
pectinatus, L'Hérit. l. c. 19 = Hyptis pectinata.
piperitus, *Lowe, in Hook. Kew Journ.* viii. (1856) 299.—Ins. Madera.
plumosus, *L'Hérit. Sert. Angl.* 20. t. 22.—Ins. Canar.
punctatus, *L'Hérit. l. c.* 20. t. 23.—Ins. Madera.
reticulatus, Willd. ex Steud. Nom. ed. II. i. 243 = Mentha mollis.
serrulatus, *Webb & Berth. Phyt. Canar.* iii. 65. t. 151.—Ins. Canar.
sidaefolius, L'Hérit. Sert. Angl. 19 = Hyptis polyantha.
Smithii, Webb & Berth. Phyt. Canar. iii. 281 = punctata.
spicata, *Benth. Lab. Gen. et Sp.* 326.—Peruv.
suaveolens, L'Hérit. Sert. Angl. 19 = Hyptis suaveolens.
tomentosus, *Benth. Lab. Gen. et Sp.* 326.—Peruv.

BYTHOPHYTON, Hook. f. Fl. Brit. Ind. iv. 286 (1884). *SCROPHULARINEAE.*
indicum, *Hook. f. l. c.*—Reg. Himal.

BYTNERIA, Jacq. Hort. Vindob. i. 10 (1770) = seq.

BYTTNERIA, Loefl. Iter Hisp. 313 (1758) = **Buettneria**, Locfl. (Sterculiac.).

BYTTNERIA, Steud. Nom. ed. II. i. 243 (1840) = **Calycanthus**, Linn.

C

CAAPEBA, Plum. ex Adans. Fam. ii. 357 (1763) = **Cissampelos**, Linn. (Menisp.).

CABALLERIA, Ruiz & Pav. Prod. Fl. Per. et Chil. 141. t. 30 (1794) = **Myrsine**, Linn.
coriacea, G. F. W. Mey. Prim. Fl. Esseq. 118 = M. floribunda.
dentata, Ruiz & Pav. Syst. Veg. 281 = M. dentata.
dependens, Ruiz & Pav. l. c. = M. dependens.
ferruginea, Ruiz & Pav. l. c. 250 = M. ferruginea.
latifolia, Ruiz & Pav. l. c. 279 = M. latifolia.
myrtifolia, Ruiz & Pav. ex DC. Prod. viii. 102 = M. dependens.
oblonga, Ruiz & Pav. Syst. Veg. 280 = M. Manglilla.
pellucida, Ruiz & Pav. l. c. = M. pellucida.
venosissima, Ruiz & Pav. l. c. 282 = M. venosissima.

CABANISIA, Klotzsch, ex Schlecht. in Abh. Naturf. Ges. Halle, vi. (1862) 176 = **Eichhornia**, Kunth (Ponted.).
caracasana, Schlecht. l. c. = E. paradoxa.
Martiana, *Schlecht. l. c.* = E. paniculata.

CABOMBA, Aubl. Pl. Gui. i. 321. t. 124 (1775). *NYMPHAEACEAE*, Benth. & Hook. f. i. 46.
NECTRIS, Schreb. Gen. i. 237 (1789).
VILLARSIA, Neck. Elem. ii. 110 (1790).
aquatica, *Aubl. l. c.*—Mexic.
aquatica, DC. Syst. ii. 36 = caroliniana.
Aubletii, Michx. Fl. Bor. Am. i. 206 = caroliniana.
australis, *Speg. in Anal. Soc. Cient. Argent.* x. (1880) 219.—Reg. Argent.
caroliniana, *A. Gray, in Ann. Lyc. N. York*, iv. (1848) 46.—Am. bor.
furcata, *Schult. f. Syst.* vii. 1379.—Bras.
peltata, F. Muell. Native Pl. Vict. 15 = Brasenia peltata.
piauhiensis, *Gardn. in Hook. Ic. Pl.* vii. t. 641.—Bras.
pinnata, Schult. f. Syst. vii. 1379 = caroliniana.
Warmingii, *Casp. in Mart. Fl. Bras.* iv. II. 142.—Bras.

CABRALEA, A. Juss. Mém. Mus. Par. xix. (1830) 229. t. 16. *MELIACEAE*, Benth. & Hook. f. i. 333.
affinis, A. Juss. l. c. 230 = polytricha.
Burchellii, *C. DC. in Mart. Fl. Bras.* xi. 1. 171.—Bras.
Cangerana, *Saldanha, in Ann. Sc. Nat.* Sér. V. xix. (1874) 210.—Bras.
Clausseniana, *C. DC. in Mart. Fl. Bras.* xi. 1. 173.—Bras.
corcovadensis, *C. DC. l. c.* 177.—Bras.
coriacea, *C. DC. l. c.* 181.—Bras.
Eichleriana, *C. DC. l. c.* 174.—Bras.
estrellensis, *C. DC. l. c.* 171.—Bras.
Gaudichaudii, *C. DC. l. c.* 172.—Bras.
glaberrima, *A. Juss. in Mém. Mus. Par.* xix. (1830) 270.—Bras.
Glaziovii, *C. DC. in Mart. Fl. Bras.* xi. 1. 172.—Bras.
inaequilatera, *Casar. Decad. Nov. Stirp. Bras.* ii. 22.—Bras.
Jussiaeana, *C. DC. in Mart. Fl. Bras.* xi. 1. 171.—Bras.
laevis, *C. DC. l. c.* 176. t. 53.—Bras.
lagoënsis, *C. DC. l. c.* 178.—Bras.
Lundii, *C. DC. l. c.* 180.—Bras.
macrophylla, *C. DC. l. c.* 174.—Bras.
montana, *C. DC. l. c.*—Bras.
multijuga, *C. DC. l. c.* 175.—Bras.
oblongiflora, *C. DC. l. c.* 180.—Bras.
oligotricha, A. Juss. in Mém. Mus. Par. xix. (1830) 230 = polytricha.
pallescens, *C. DC. in Mart. Fl. Bras.* xi. 1. 175. t. 52.—Bras.
pedunculata, *C. DC. l. c.* 172.—Bras.
pilosa, *C. DC. l. c.* 177.—Bras.
Poeppigii, *C. DC. in DC. Monog. Phan.* i. 471.—Am. trop.
polytricha, *A. Juss. in Mém. Mus. Par.* xix. (1830) 230.—Bras.
Richardiana, *C. DC. in Mart. Fl. Bras.* xi. 1. 176.—Bras. (vel Cuba).
Riedelii, *C. DC. l. c.* 180.—Bras.
rubiginosa, *C. DC. l. c.* 179.—Bras.
Selloi, *C. DC. l. c.* 176.—Bras.
silvatica, *C. DC. l. c.* 173.—Bras.
sulcata, *C. DC. l. c.* 171.—Bras
tomentosa, Casar. Decad. Nov. Stirp. Bras. ii. 22 = polytricha.
Warmingiana, *C. DC. in Mart. Fl. Bras.* xi. 1. 178.—Bras.

CABRERA, Lag. Gen. et Sp. Nov. 5 (1816) = **Paspalum**, Linn. (Gramin.).
chrysoblepharis, Lag. l. c. = P. exasperatum.

CACABUS, Bernh. in Linnaea, xiii. (1839) 360. *SOLANACEAE*, Benth. & Hook. f. ii. 896.
DICTYOCALYX, Hook. f. in Trans. Linn. Soc. xx. (1847) 202
STREPTOSTIGMA, Regel, Gartenfl. (1853) 322. t. 68.
THINOGETON, Benth. Bot. Voy. Sulph. 142 (1844).

CACABUS:—
inflatus, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 255.—Peruv.
maritimus, *Benth. & Hook. f. Gen.* ii. 897.—N. Granat.
mexicanus, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 127.—Mexic.
nolanoides, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. iv. (1849) 255.—Peruv.
prostratus, *Bernh. in Linnaea*, xiii. (1839) 360.—Peruv.

CACALIA, Linn. Sp. Pl. 834 (1753) = **Senecio**, Tourn. (Compos.).
abyssinica, R. Br. in Salt, Abyss. App. 65 = S. abyssinicus.
abyssinica, Walp. Ann. ii. 908 = S. tuberosus.
acaulis, Linn. f. Suppl. 364 = S. acaulis, *Sch. Bip.*
acerifolia, Siebold, ex Maxim. in Bull. Acad. Pétersb. xix. (1874) 487, nomen.—Japon.
acetosellaefolia, Jacq. ex Steud. Nom. ed. II. ii. 243 = S. articulatus.
aconitifolia, Bunge, Enum. Pl. Chin. Bor. 37 = S. aconitifolius.
aconitifolia, Miq. ex Maxim. in Bull. Acad. Pétersb. xix. (1874) 487 = S. adenostyloides.
acuminata, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) 1. 215 (= *Mulgedium hastatum*).
adenostyloides, Franch. & Sav. ex Maxim. in Bull. Acad. Pétersb. xix. (1874) 486 = S. adenostyloides.
albida, Cass. ex Schur, Enum. Pl. Transs. 301 = Adenostyles alba.
albifrons, Linn. f. Suppl. 353 = Adenostyles alba.
Alliariae, Gouan, Illustr. 65 = Adenostyles alba.
alliariaefolia, Lam. Encyc. i. 532 = Adenostyles viridis.
alliariaefolia, Poir. Voy. Barb. ii. 235 = Petasites fragrans.
alpina, Linn. Sp. Pl. 836 = Adenostyles viridis.
alpina, Mill. Gard. Dict. ed. VIII. n. 1 = Adenostyles alba.
amplifolia, DC. Prod. vi. 328 = S. amplifolius.
angulata, Vahl, Symb. Bot. iii. 92 = Emilia angulata.
angulata, Vell. Fl. Flum. 336; viii. t. 55 = Mikania cordifolia.
angulosa, Wall. Cat. n. 3152 = Gynura angulosa.
angustifolia, H. B. & K. Nov. Gen. et Sp. iv. 159 = S. Humboldtianus.
angustifolia, Wall. Cat. n. 3163 = Emilia prenanthoidea.
Anteuphorbium, Linn. Sp. Pl. 834 = S. Anteuphorbium.
appendiculata, Linn. f. Suppl. 352 = S. populifolius.
arborea, H. B. & K. Nov. Gen. et Sp. iv. 163 (= *Psacalium arboreum*).—Am. austr.
arbuscula, Thunb. Prod. Pl. Cap. 142 = Othonna arbuscula.
arenaria, H. B. & K. Nov. Gen. et Sp. iv. 165 = S. sabulosus.
articularis, Salisb. Prod. 187 = seq.
articulata, Linn. f. Suppl. 364 = S. articulatus.
asarifolia, Steud. Nom. ed. II. i. 243 = Faujasia reticulata.
asclepiadea, Linn. f. Suppl. 352 = S. asclepiadeus.
atriplicifolia, Linn. Sp. Pl. 835 = S. atriplicifolius.
aurantiaca, Blume, Bijdr. 908 = Gynura aurantiaca.
aurantiaca, Wall. Cat. n. 3146 = Gynura nepalensis.
auriculata, DC. Prod. vi. 329 = S. dahuricus.
baccharidifolia, Poepp. ex Phil. f. Cat. Pl. Vasc. Chil. 156 = S. baccharidifolius.
Berlandieri, DC. Prod. vi. 328 = S. desertorum.
bicolor, Roxb. Hort. Beng. 61; Fl. Ind. iii. 412 = Gynura bicolor.
bipinnata, Linn. f. Suppl. 353 = S. bipinnatus.
bulbosa, Lour. Fl. Cochinch. 485 = Gynura ovalis.
cancellata, Rudolphi, in Ledeb. Diss. Pl. Doming. (1805) = S. cancellatus.
candicans, Poepp. ex DC. Prod. vi. 325 = Culcitium Poeppigii.
candicans, Vahl, Symb. Bot. iii. 91 = S. candicans.
canescens, Poepp. ex DC. Prod. vi. 414 = S. Neaei.
canescens, Willd. Enum. Hort. Berol. Suppl. 427 = S. Haworthii.
Calliariae, Gouan, Illustr. 65 = Adenostyles alpina.
carnosa, Ait. Hort. Kew. iii. 156 = S. elegans.
caucasica, Hoffm. Hort. Mosq. (1808) n. 598 = S. platyphyllus.
cervariaefolia, DC. Prod. vi. 328 = S. cervariaefolius.

CACALIA;—
chiapensis, A. Gray, in Proc. Am. Acad. xix. (1883) 53 = Senecio chiapensis.
cinerarioides, H. B. & K. Nov. Gen. et Sp. iv. 166 = S. Bonplandianus.
cirsiifolia, Hook. & Arn. Bot. Beech. Voy. 436 = S. cirsiifolius.
cissampelina, DC. Prod. vi. 331 = S. cissampelinus.
citrina, Boj. ex DC. l. c. 302 = Emilia citrina.
Clematites, Poir. Encyc. Suppl. ii. 4 = Faujasia flexuosa.
coccinea, Sims, Bot. Mag. t. 564 = Emilia flaminea.
cordata, Vell. Fl. Flum. 336; viii. t. 53 = Mikania cordifolia.
Cor-Jesu, Vell. l. c. 340; viii. t. 71 = Mikania officinalis.
cordifolia, H. B. & K. Nov. Gen. et Sp. iv. 168. t. 360 = S. cardiophyllus.
cordifolia, Linn. f. Suppl. 351 = Mikania cordifolia.
cordiformis, Lam. Fl. Fr. ii. 75 = Adenostyles alpina.
crassissima, Wall. Cat. n. 3155 = Notonia crassissima.
crenata, Vell. Fl. Flum. 335; viii. t. 51.—Bras.
cuneifolia, Linn. Mant. i. 110 = S. cuneifolius.
Cusimbua, D. Don, Prod. Fl. Nep. 179 = Gynura angulosa.
cuspidata, Bertol. Fl. Guatimal. 32 = S. guatimalensis.
cuspidata, Klatt, in Ann. Sc. Nat. Sér. V. xviii. (1873) 374.—Ins. Borbon.
cylindrica, Lam. Encyc. i. 529 = Othonna cylindrica.
cylindriflora, Wall. Cat. n. 3150 = Gynura sarmentosa.
decomposita, A. Gray, Pl. Wright. ii. 99 = S. Grayanus.
decurrens, Vell. Fl. Flum. 340; viii. t. 72 = Baccharis genistelloides.
delphiniifolia, Sieb. & Zucc. in Abh. Acad. Muench. iv. III. (1846) 190 = S. Zuccarinii.
denticulata, Hook. & Arn. Bot. Beech. Voy. 29 = S. denticulatus.
dichroa, Boug. Nov. Comp. Bras. 40. t. 7 = S. dichrous.
discolor, Griseb. Cat. Pl. Cub. 157 = S. discolor.
disticha, Heyne, ex DC. Prod. v. 384 = Conyza Wallichii.
diversifolia, Torr. & Gray, Fl. N. Am. ii. 435 = S. floridanus.
dulcis, Salisb. Prod. 187 = S. ficoides.
echinata, Linn. f. Suppl. 353 = S. echinatus.
elata, Spreng. ex DC. Prod. vi. 314 = S. Ligularia.
emarginata, Vell. Fl. Flum. 336; viii. t. 52.—Bras.
farfaraefolia, Sieb. & Zucc. in Abh. Acad. Muench. iv. III. (1846) 190 = S. farfaraefolius.
ficoides, Linn. Sp. Pl. 834 = S. ficoides.
fimbrillifera, Cass. in Dict. Sc. Nat. xlviii. 460 = S. deltoideus.
Finlaysoniana, Wall. Cat. n. 3162 = Gynura sarmentosa.
flexuosa, Wall. Cat. n. 3160 = Faujasia flexuosa.
flexuosa, Willd. ex DC. Prod. vii. 69 = Trixis divaricata.
floridana, A. Gray, in Proc. Am. Acad. xix. (1883) 52 = S. floridanus.
foetens, Wall. Cat. n. 3156 = Gynura nepalensis.
frutescens, Vell. Fl. Flum. 338; viii. t. 64.—Bras.
gigantea, Nees, Del. Sem. Hort. Vrat. (1841) 3 = S. atriplicifolius.
glabra, Ledeb. Ind. Sem. Hort. Dorpat. (1820) 16 = S. sagittatus.
glabra, Heyne, in Wall. Cat. n. 3145 = Emilia sonchifolia.
glabra, Mill. Gard. Dict. ed. VIII. n. 2 = Adenostyles alpina.
glubra, Vill. Prosp. 30 = Adenostyles viridis.
glabrata, H. B. & K. Nov. Gen. et Sp. iv. 161 (= *Psacalium glabratum*).—Am. trop.
glandulosa, Salisb. Prod. 187 = Porophyllum ruderale.
glauca, Salisb. l. c. = S. succulentus, *Sch. Bip.*
graciliflora, Wall. Cat. n. 3149 = S. graciliflorus.
gracilis, Boj. ex DC. Prod. vi. 302 = Emilia gracilis.
grandiflora, Wall. Cat. n. 3147 = Notonia grandiflora.
hastata, Linn. Sp. Pl. 835 = S. sagittatus.
Haworthii, Sweet, in Loud. Hort. Brit. 336 = S. Haworthii.
heterogama, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. 1852) 107 = S. heterogamus.
heterophylla, Poepp. ex DC. Prod. vi. 414 = S. pilquensis.

CACALIA :—
hieracioides, Willd. Sp. Pl. iii. 1731 = Gynura auriculata.
hirsuta, Vill. Prosp. 31 = Adenostyles alba.
humifusa, Boj. ex DC. Prod. vi. 302 = Emilia humifusa.
humilis, H. B. & K. Nov. Gen. et Sp. iv. 166 = Senecio humilis.
hybrida, Steud. Nom. ed. II. i. 244 = Adenostyles alpina, etc.
iatrophoides, H. B. & K. Nov. Gen. et Sp. iv. 169 = S. jatrophoides.
ilicifolia, H. B. & K. l. c. 163 = S. scrobicarius.
incana, Bosc, ex DC. Prod. vi. 330 = S. Boscianius.
incana, Linn. Sp. Pl. ed. II. 1169 = Gynura nitida.
indica, Lam. Encyc. i. 530.—Ind. or.
involucrata, H. B. & K. Nov. Gen. et Sp. iv. 166 = S. involucratus.
Kleinia, J. Grah. Cat. Bomb. Pl. 98 = Notonia balsamica.
Kleinia, Herb. Madr. ex C. B. Clarke, Comp. Ind. 176 = Notonia grandiflora.
Kleinia, Linn. Sp. Pl. 834 = S. Kleinia.
Kleiniana, Hill, Veg. Syst. iii. 144 = praec.
kleinioides, H. B. & K. Nov. Gen. et Sp. iv. 164 (= *Psacalium kleinioides*).—Am. trop.
laciniata, Jacq. Ic. Rar. 1. t. 68 = S. articulatus.
laciniata, Wall. Cat. n. 3153 = Gynura lycopersicifolia.
laeta, Vell. Fl. Flum. 341; viii. t. 76.—Bras.
lanata, H. B. & K. Nov. Gen. et Sp. 162 = S. lanatus.
lanceolata, Nutt. Gen. Am. ii. 138 = S. Boscianus.
laurifolia, Linn. f. Suppl. 351 = Mikania laurifolia.
ledifolia, H. B. & K. Nov. Gen. et Sp. iv. 162 = S. ledifolius.
leucopappa, DC. Prod. vi. 330 = S. leucopappus.
leucophylla, Willd. Sp. Pl. iii. 1736 = Adenostyles candidissima.
Linaria, Cav. Ic. iii. 29. t. 257 = Porophyllum Linaria.
linearis, Labill. Nov. Holl. Pl. ii. 36. t. 178 = Bedfordia linearis.
linifolia, Heyne, ex DC. Prod. v. 25 = Vernonia cinerea.
linifolia, Linn. Mant. 109 = Porophyllum linifolium.
lobata, Moris, in Flora, xv. (1832) Lit. ii. 79.—Chili.
longifolia, Haw. Rev. Pl. Succ. 193 = S. pugioniformis.
lutea, Mill. Gard. Dict. ed. VIII. n. 10.—Ins. S. Helena.
Lycopersicum, Heyne, ex DC. Prod. vi. 301 = Gynura lycopersicifolia.
macrophylla, Bieb. Fl. Taur. Cauc. ii. 286 = S. platyphyllus.
macrophylla, Rafin. in Med. Repos. N. York, v. (1808) 354, nomen.—Hab. ?
maculata, Buch.-Ham. ex Steud. Nom. ed. II. i. 244 = Gynura sinuata.
Mentrasto, Vell. Fl. Flum. 339; viii. t. 69 = Ageratum conyzoides.
mikaniaefolia, DC. Prod. vi. 328 = S. mikaniaefolia.
mucronata, Heyne, in Wall. Cat. n. 3161 = Emilia flammea.
multiflora, Auct. ex Walp. Rep. vi. 266 = S. multiflorus.
napeaefolia, DC. Prod. vi. 328 = S. napeaefolius.
napellifolia, S. Schau. in Linnaea, xix. (1847) 732 = S. napellifolius.
Nardosmia, A. Gray, in Proc. Am. Acad. vii. (1868) 361 = Cacaliopsis Nardosmia.
nepalensis, Spreng. ex DC. Prod. vi. 300 = Gynura nepalensis.
nipponica, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 181.—Japon.
nitida, H. B. & K. Nov. Gen. et Sp. iv. 162 = S. nitidus.
nitida, Vell. Fl. Flum. 337; viii. t. 59.—Bras.
nivea, Kunze, ex DC. Prod. vi. 325 = Culcitium Poeppigii.
obsoleta, Vell. Fl. Flum. 337; viii. t. 57 = Mikania biformis.
odora, Forsk. Fl. Aegypt. Arab. 146 = S. odorus.
odorata, Desf. Cat. Hort. Par. ed. III. 165, 400 = S. odoratus.
oligocephala, Scheele, in Linnaea, xviii. (1844) 480.—Hab. ?
oppositifolia, Poir. Encyc. Suppl. ii. 7 = Porophyllum oppositifolium.
ovalifolia, Salisb. Prod. 186 = S. Anteuphorbium.

CACALIA :—
ovalis, Ker-Gawl. in Bot. Reg. t. 101 = Gynura pseudochina.
ovata, Walt. Fl. Carol. 196 = Senecio Walteri ?
pachyphylla, Sch. Bip. ex Secm. Bot. Voy. Herald, 310 = S. sclerophyllus.
paniculata, Rafin. Ann. Nat. (1820) 14 = S. Nuttallii.
papillaris, Linn. Sp. Pl. 834 = S. papillaris.
parasitica, Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot. ii. 244 = S. parasiticus.
patens, H. B. & K. Nov. Gen. et Sp. iv. 164 = S. patens.
pedunculata, Vell. Fl. Flum. 339; viii. t. 68.—Bras.
peltata, H. B. & K. Nov. Gen. et Sp. iv. 170. t. 361 = S. peltiferus.
pendula, Forsk. Fl. Aegypt. Arab. 145 = S. pendulus.
penduliflora, A. Gray, in Proc. Am. Acad. xix. (1883) 53 = S. subpeltatus, *Sch. Bip.*
penicillata, Cass. in Dict. Sc. Nat. xlviii. 460 = S. penicillatus.
Petasites, Lam. Encyc. i. 531 = Adenostyles alba.
peucedanifolia, Jacq. Coll. Suppl. 149 = S. paniculatus.
pilosa, Vell. Fl. Flum. 338; viii. t. 61 = Mikania cordifolia.
pinnata, Willd. Enum. Hort. Berol. 850 = S. Othonnae.
pinnatifida, Berg. Desc. Pl. Cap. 230 = S. pinnatifidus.
pinnatifida, Linn. Mant. ii. 281 = S. bipinnatus.
pinnatifida, Lour. Fl. Cochinch. ii. 486 = Gynura pinnatifida.
pinnatifida, Wight, ex DC. Prod. vi. 301 = Gynura lycopersicifolia.
Porophyllum, Linn. Sp. Pl. 834 = Porophyllum ruderale.
praestans, Baker, in Mart. Fl. Bras. vi. III. 384, sphalm. = Trixis mollissima.
prenanthoides, H. B. & K. Nov. Gen. et Sp. iv. 167 = S. albiflorus.
prenanthoides, Sieber, ex DC. Prod. vi. 302 = Emilia sonchifolia.
procumbens, Lour. Fl. Cochinch. ii. 485 = Gynura sarmentosa.
Pteranthus, Rafin. Ann. Nat. (1820) 14 = S. Nuttallii.
pubescens, Rafin. Fl. Ludov. 61 = Eupatorium ludovicianum.
pugioniformis, Salzm. ex Link, Enum. Hort. Berol. ii. 510 = S. pugioniformis.
pulchella, H. B. & K. Nov. Gen. et Sp. iv. 160 = S. pulchellus.
pumila, Boj. ex DC. Prod. vi. 302 = Emilia pumila.
punctulata, Vell. Fl. Flum. 339; viii. t. 66.—Bras.
pungens, H. B. & K. Nov. Gen. et Sp. iv. 160 = S. pungens.
purpurascens, Wall. Cat. n. 3157 = Gynura purpurascens.
quinqueloba, Thunb. Prod. Pl. Cap. 142 = S. quinquelobus.
racemosa, DC. Prod. vi. 330 = S. exsertus.
racemosa, Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot. ii. 241 = S. helodes.
radicans, Linn. f. Suppl. 354 = S. radicans.
radulaefolia, H. B. & K. Nov. Gen. et Sp. iv. 169 = S. radulaefolius.
ramosa, Spreng. ex DC. Prod. vi. 368 = S. triligulatus.
reclinata, Roxb. Hort. Beng. 101; Fl. Ind. iii. 412.—Ins. Molucc.
reclinata, Wall. Cat. n. 3151 = Gynura sarmentosa.
regia, Baker, in Mart. Fl. Bras. vi. III. 384, sphalm. = Trixis divaricata.
reniformis, Lam. Fl. Fr. ii. 75 = Adenostyles viridis.
reniformis, Muhl. in Willd. Sp. Pl. iii. 1735 = S. Muehlenbergii.
repens, Linn. Mant. 110 = S. succulentus.
reticulata, Vahl, Symb. Bot. iii. 91 = Faujasia reticulata.
reticulata, Vell. Fl. Flum. 337; viii. t. 60.—Bras.
rhombifolia, Willd. Enum. Hort. Berol. 850 = S. platyphyllus.
rigida, Thunb. Prod. Pl. Cap. 142 = S. spinescens.
Ros-marinus, Vell. Fl. Flum. 339; viii. t. 67.—Bras.
rotundifolia, Willd. Phyt. 11. t. 6. f. 1 = Vernonia cinerea.
ruderalis, Sw. Prod. Veg. Ind. Occ. 110 = Porophyllum ruderale.

CACALIA :—
runcinata, H. B. & K. Nov. Gen. et Sp. iv. 168 = Senecio roseus.
runcinata, Lam. Encyc. i. 529 = S. articulatus.
sagittaria, Heyne, in Wall. Cat. n. 3159 = Gynura pseudochina.
sagittata, Vahl, Symb. Bot. iii. 91 = Emilia flammea.
sagittifolia, Mertens, in Linnaea, v. (1830) 63, nomen.—Ins. Koragin.
salicina, Labill. Nov. Holl. Pl. ii. 37. t. 179 = Bedfordia salicina.
sarmentosa, Blume, Bijdr. 907 = Gynura sarmentosa.
sarmentosa, Lesch. ex Hook. f. Fl. Brit. Ind. iii. 352 = S. candicans.
Sarracenia, Blanco, Fl. Filip. ed. I. 618 = Gynura angulosa.
sarracenica, Linn. Sp. Pl. ed. II. 1169 = S. Cacaliaster.
scabra, Vahl, Symb. Bot. iii. 92 = Trixis brasilicnsis.
scandens, Ait. Hort. Kew. iii. 157.—Afr. austr.
scandens, Thunb. Prod. Pl. Cap. 142 = S. deltoidcus.
scandens, Zeyh. ex DC. Prod. vi. 404 = S. volubilis.
Schaffneri, A. Gray, in Proc. Am. Acad. xix. (1883) 53.—Mexic.
sempervirens, Spreng. ex DC. Prod. vi. 442 = Notonia grandiflora.
sempervirens, Vahl, Symb. Bot. iii. 92 = S. sempervivus.
semperviva, Forsk. Fl. Acgypt. Arab. 146 = S. sempervivus.
senecioides, H. B. & K. Nov. Gen. et Sp. iv. 163 = S. assuayensis.
septemnata, Vell. Fl. Flum. 338; viii. t. 63.—Bras.
sessilifolia, Hook. & Arn. Bot. Bcech. Voy. 436 = S. sessilifolius.
sessilis, Vell. Fl. Flum. 341; viii. t. 73 = Baccharis articulata.
sinuata, Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc. i. 25 = S. albo-lutescens.
sonchifolia, Forsk. Fl. Aegypt. Arab. p. cxix = Emilia angulata.
sonchifolia, Hort. ex Steud. Nom. ed. II. i. 244 = Emilia sagittata.
sonchifolia, Linn. Sp. Pl. 835 = Emilia sonchifolia.
sonchifolia, Wall. Cat. n. 3144 D = Lactuca remotiflora.
spathulata, Moç. & Sesse, ex D. Don, in Trans. Linn. Soc. xvi. (1830) 250 = Chaptalia spathulata.
suaveolens, Linn. Sp. Pl. 835 = S. suaveolens.
suffruticosa, Linn. Mant. i. 109 = Porophyllum linifolium.
tabularis, A. Gray, in Proc. Am. Acad. xix. (1883) 52 = S. tabularis.
tenuissima, Boj. in Rapp. Soc. Maur. vi. 27 = Emilia gracilis.
teres, Wall. Cat. n. 3164 = Emilia prenanthoidea.
teretifolia, H. B. & K. Nov. Gen. et Sp. iv. 159 = S. teretifolius.
terminalis, Salisb. Prod. 187 = S. Kleinia.
ternata, Vell. Fl. Flum. 336; viii. t. 56 = Mikania apiifolia.
thyrsoidea, Spreng. ex DC. Prod. vi. 368 = S. alatus.
toluccana, DC. Prod. vi. 328 = S. callosus.
tomentosa, Eckl. ex DC. l. c. 403 = S. scaposus.
tomentosa, Haw. Misc. 189 = S. Haworthii.
tomentosa, Jacq. Fl. Austr. iii. 20 = Adenostyles alba.
tomentosa, Linn. f. Suppl. 353 = S. oliganthus.
tomentosa, Vell. Fl. Flum. 341; viii. t. 75.—Bras.
tomentosa, Vill. Prosp. 31 = Adenostyles candidissima.
triangularis, Vell. Fl. Flum. 338; viii. t. 62 = Mikania cordifolia.
trilobata, Vell. l. c. 336; viii. t. 54 = Mikania glomerata.
trinervia, Vell. l. c. 338; viii. t. 65.—Bras.
triphylla, Vell. l. c. 341; viii. t. 74.—Bras.
tuberosa, Bosc, ex DC. Prod. vii. 89 = Krigia Dandelion.
tuberosa, Delile, in Ferr. & Gall. Voy. Abyss. iii. 142 = S. tuberosus.
tuberosa, Nutt. Gen. Am. ii. 138 = S. Nuttallii ?
tussilaginoides, H. B. & K. Nov. Gen. et Sp. iv. 168 = S. farfarus.
uniflora, Schum. & Thonn. Beskr. Guin. Pl. 382 = Gynura cernua.
vaccinoides, H. B. & K. Nov. Gen. et Sp. iv. 162 (= *Psacalium vaccinoides*).—Am. trop.

CACALIA :—
verbascifolia, Sibth. & Sm. Fl. Graec. ix. 35 = Senecio thapsoides.
villosa, Jacq. Coll. ii. 347 = S. purpureus.
villosa, Vell. Fl. Flum. 337 ; viii. t. 58.—Bras.
volubilis, Blume, Bijdr. 903 = S. araneosus.
volubilis, Buch.-Ham. ex DC. Prod. vi. 370 = S. Buimalia.
Wightiana, Wall. ex DC. Prod. vi. 370 = S. scandens.

CACALIOPSIS, A. Gray, in Proc. Am. Acad. xix. (1883) 50. *COMPOSITAE.*
Nardosmia, *A. Gray, l. c.*—Am. bor. occ.

CACAO, Tourn. ex Mill. Gard. Dict. ed. VI. (1752) = **Theobroma**, Linn. (Sterculiac.).
bicolor, Poir. Encyc. Suppl. ii. 7 = T. bicolor.
guianensis, Aubl. Pl. Gui. ii. 683 = T. Cacao.
minor, Gaertn. Fruct. ii. 190 = T. Cacao.
sativa, Aubl. Pl. Gui. ii. 689 = T. Cacao.
sylvestris, Aubl. l. c. 687 = Genipa Merianac.
Theobroma, Tussac, Fl. Med. Ant. t. 13 = T. Cacao.

CACARA, Rumph. ex Thou. in Dict. Sc. Nat. vi. 35 (1805) = **Pachyrhizus**, Rich. (Legumin.).

CACATALI, Adans. Fam. ii. 213 (1763) = **Pedalium**, Royen (Pedalin.).

CACCINIA, Savi, Cose Bot. 1. t. 1 (1832). *BORAGINEAE*, Benth. & Hook. f. ii. 846.
ANISANTHERA, Rafin. Fl. Tellur. iii. 80 (1836).
Actinobole, *Bunge, Helioc.* 8.—Persia.
Celsii, Boiss. Diagn. Ser. I. xi. 133 = glauca.
dubia, *Bunge, Lehm. Rel.* (1847) 242.—As. centr.
glauca, *Savi, Cose Bot.* 1. t. 1. f. 1–6 (1832).—Persia ; Beluchist.
Kotschyi, *Boiss. Diagn.* Ser. I. vii. 31.—Persia.
Rauwolfii, A. DC. in DC. Prod. x. 167 = Russellii.
Rauwolfii, *C. Koch, in Linnaea*, xvii. (1843) 303.—As. Min. ; Persia.
Russellii, *Boiss. Diagn.* Ser. I. xi. 134.—Syria.
strigosa, *Boiss. l. c.*—Persia.

CACHRIS, D. Dietr. Syn. Pl. i. 586 (1839) = **Cachrys**, Tourn. (Umbellif.).

CACHRYDIUM, Link, Handb. i. 339 (1829) = **Hippomarathrum**, Link (Umbellif.).
Libanotis, Link, Handb. i. 339 = H. Libanotis.

CACHRYS Linn. Syst. ed. I (1735) ; Gen. ed. I. 75 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 904.
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 333 = Diplolophium africanum.
acaulis, DC. Prod. iv. 238 = Prangos Szovitzii.
alata, Hoffm. Umb. i. 176 ; Bieb. Fl. Taur. Cauc. i. 217 = Prangos ferulacea.
alpina, *Bieb. l. c.*—Tauria.
ambigua, Salzm. ex DC. Prod. iv. 241 = Magydaris panacina.
amplexifolia, Pritz. Ind. Ic. 179 = scq.
amplifolia, Ledeb. in Eichw. Casp. 6. t. 9 = Hippomarathrum crispum.
athamanthoides, Bieb. Fl. Taur. Cauc. iii. 217 = Rumia athamanthoides.
callosperma, Pall. Reise, iii. 663 = odontalgica.
cheilanthifolia, *Boiss. Fl. Orient.* ii. 936.—Persia.
cretica, Lam. Encyc. i. 259 = Lecokia cretica.
crispa, Pers. Syn. i. 311 = Hippomarathrum crispum.
crispata, *Pomel, Nouv. Mat. Fl. Atl.* 151.—Algeria.
cristata, DC. Prod. iv. 238 = Hippomarathrum cristatum.
cylindracea, Guss. ex DC. l. c. 239 = Prangos cylindracea.
decursiva, Hornem. Hort. Hafn. ii. 960. Add. = Siler caucasicum.
dichotoma, Spreng. Pl. Umb. Prod. 21 = Capnophyllum dichotomum.
didyma, *Regel, in Act. Hort. Petrop.* v. (1877) 601.—Turkestan.
echinophora, Guss. Fl. Sic. Syn. i. 347 = Hippomarathrum cristatum.
eriantha, *DC. Prod.* iv. 238.—Persia.
goniocarpa, *Boiss. Diagn.* Ser. I. x. 53.—Syria.
Herderi, *Regel, in Act. Hort. Petrop.* v. (1877) 601.—Turkest.
hirta, *Bertol. Disquisit. de Pl. Nov.* 6.—Hab. ?
humilis, *Schousb. Vextr. Marokko*, 133.—Marocco.
hungarica, Mill. Gard. Dict. ed. VIII. n. 5 = Magydaris panacina.
involucrata, Pall. ex Willd. fide Roem. & Schult. Syst. vi. 447 = Polylophium orientale.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 602.—Turkestan.
laevigata, Brouss. ex Ball, in Journ. Linn. Soc. xvi. (1878) 463 = Hippomarathrum cristatum.
laevigata, *Lam. Encyc.* i. 256.—Lusitan.
lanata, Hort. ex Steud. Nom. ed. II. i. 245 = Magydaris panacina, tomentosa.
latifolia, Bieb. Fl. Taur. Cauc. i. 219 = Siler caucasicum.
Libanotis, Georgi, Beschr. Russ. Reich. iii. IV. 845 = odontalgica.
Libanotis, Gouan, Illustr. 12 = laevigata.
Libanotis, Guss. Fl. Sic. Prod. i. 358 = Prangos ferulacea.
Libanotis, Linn. Sp. Pl. 346 = Hippomarathrum Libanotis.
Libanotis, Salzm. ex Ball, in Journ. Linn. Soc. xvi. (1878) 463 = Hippomarathrum cristatum.
Linearia, Mill. Gard. Dict. ed. VIII. n. 4 = Hippomarathrum Libanotis.
littoralis, Spreng. Umbell. Prod. iv. 20 = Daucus bessarabicus.
longiloba, DC. Prod. iv. 237 = Hippomarathrum crispum.
macrocarpa, *Ledeb. Fl. Alt.* i. 364.—Sibir. altaic.
maritima, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 259 = Crithmum maritimum.
microcarpa, Bieb. Fl. Taur. Cauc. i. 218 = Hippomarathrum crispum.
Morisoni, All. Auctuar. Fl. Pedem. 23 = laevigata.
nudicaulis, Godet, ex DC. Prod. iv. 237 = Hippomarathrum crispum.
odontalgica, *Pall. Reise*, iii. 720.—Reg. Caucas. ; Sibir.
panacifolia, Vahl, Symb. Bot. i. 25 = Magydaris panacina.
panax-siculum, Ucria, in Roem. Arch. i. 68 = Magydaris tomentosa.
papillaris, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 75.—Mesopotam.
pastinacea, Lam. Encyc. i. 260 = Magydaris panacina.
peregrina, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 259 = Capnophyllum dichotomum.
peucedanoides, *Desf. Fl. Atlant.* i. 250.—Afr. bor.
prangoides, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 76.—Persia.
pterochlaena, DC. Prod. iv. 237 = Hippomarathrum pterochlaenum.
pungens, Jan, ex Guss. Fl. Sic. Prod. i. App. 7 = Hippomarathrum cristatum.
seseloides, Bieb. Fl. Taur. Cauc. iii. 217 = Rumia seseloides.
sibirica, Steph. ex Fisch. in Cat. Sem. Hort. Petrop. (1823) = Phloidicarpos dahuricus.
sibirica, Turcz. ex Ledeb. Fl. Ross. ii. 331 = Phloidicarpos villosus.
sicula, Gueldenst. ex Ledeb. Fl. Ross. ii. 357 = Hippomarathrum crispum.
sicula, Guss. Fl. Sic. Prod. i. 359 = Hippomarathrum cristatum.
sicula, Linn. Sp. Pl. ed. II. 355 = Hippomarathrum pterochlaenum.
sicula, Urv. in Mém. Soc. Linn. Par. i. (1822) 287 = Hippomarathrum Bocconi.
sphaerocarpa, Tenore, Prod. Fl. Nap. xix. ; Syn. 120 = Hippomarathrum Libanotis.
taurica, Willd. Sp. Pl. i. 1410 = Rumia taurica.
tomentosa, Desf. Fl. Atlant. i. 249 = Magydaris tomentosa.
tortuosa, Webb & Berth. Phyt. Canar. iii. 281 = Ferula tomentosa.
trifida, Mill. Gard. Dict. ed. VIII. n. 1 = laevigata.
triquetra, Spreng. in Ges. Naturf. Fr. Berl. Mag. vi. (1812) 259 = Prangos triquetra.
vaginata, Ledeb. Fl. Alt. i. 366 = Hippomarathrum vaginatum.

CACHYRIS, Zumag. Fl. Pedem. i. 39, 425 (1849) = praec.

CACONAPEA, Cham. in Linnaea, viii. (1833) 28 = **Herpestis**, Gaertn. (Scrophularin.).
gratioloides, Cham. l. c. 29 = H. gratioloides.

CACONOBEA, Walp. Rep. iii. 281 (1845) = praec.

CACOSMANTHUS ,Miq. Fl. Ind. Bat. ii. 1040 (1856) = KAKOSMANTHUS, Hassk. = **Payena**, Linn. (Sapotac.).

CACOSMIA, H. B. & K. Nov. Gen. et Sp. iv. 289. t. 404 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 396.
CLAIRVILLEA, DC. Prod. v. 636 (1836).
XANTHOLEPIS, Willd. ex Less. in Linnaea, iv. (1829) 338.
quinquenervia, Steud. Nom. ed. II. i. 377, in syn. = seq.
rugosa, *H. B. & K. Nov. Gen. et Sp.* iv. 289.—Peruv.

CACOTANIS, Rafin. New Fl. Am. ii. 43 (1836) = **Eclipta**, Linn. ? (Compos.).
ciliata, Rafin. l. c. 44.—Am. bor.

CACOUCIA, Aubl. Pl. Gui. i. 450. t. 179 (1775). *COMBRETACEAE*, Benth. & Hook. f. i. 688.
HAMBERGERA, Scop. Introd. 106 (1777).
SCHOUSBOEA, Willd. Sp. Pl. ii. 578 (1799).
bractcata, *Laws. in Oliver, Fl. Trop. Afr.* ii. 434.—Afr. trop.
chinensis, *A. Juss. ex DC. Prod.* iii. 22.—China.
coccinea, *Aubl. Pl. Gui.* i. 450. t. 179.—Guiana.
cordifolia, *Walp. Rep.* ii. 68.—Afr. trop.
exannulata, *O. Hoffm. in Linnaea*, xliii. (1880–82) 132.—Afr. trop.
lucida, Hassk. in Flora, xxvii. (1844) II. 607 = Combretum trifoliatum.
paniculata, *Laws. in Oliver, Fl. Trop. Afr.* ii. 434.—Afr. trop.
platyptera, *Welw. in Trans. Linn. Soc.* xxvii. (1869) 24.—Afr. trop.
trifoliata, DC. Prod. iii. 22 = Combretum trifoliatum.
velutina. *S. Moore, in Journ. Bot.* xviii. (1880) 2.—Afr. trop.
villosa, *Laws. in Oliver, Fl. Trop. Afr.* ii. 433.—Afr. trop.

CACTUS, Linn. Syst. ed. I (1735) = *CACTEAE*, Benth. & Hook. f. Gen. i. 1008.
abnormis, Willd. Enum. Hort. Berol. Suppl. 31 = Cereus peruvianus.
Ackermanni, Haw. in Bot. Reg. t. 1331 = Phyllocactus Ackermanni.
aculeatissimus, Zeyh. ex Steud. Nom. ed. I. 131 (quid ?).
alatus, Hook. Bot. Mag. t. 2820 = Rhipsalis pachyptera.
alatus, Sw. Prod. Veg. Ind. Occ. 77 = Rhipsalis Swartziana.
alatus, Willd. Enum. Hort. Berol. Suppl. 35 = Phyllocactus phyllanthoides.
albisetosus, Haw. ex Spreng. Syst. ii. 496 = Cereus albisetosus.
albisetus, Steud. Nom. ed. II. i. 245 = praec.
ambiguus, Bonpl. Jard. Malm. t. 36 = Cereus ambiguus.
ambiguus, Gill. ex Loud. Hort. Brit. 194 = Mammillaria ambigua.
americanus, Vitm. Summa Pl. iii. 209 (quid ?).
anizogonus, Hort. Angl. ex Foerst. Handb. Cact. ed. II. i. 764 = Cereus triangularis.
arboreus, Vell. Fl. Flum. 207 ; v. t. 28 = Opuntia arborea.
aurantiaca, Gill. ex Loud. Hort. Brit. Suppl. ii. 652 = Opuntia aurantiaca.
aurantiiformis, Thiery, ex Steud. Nom. ed. I. 131 ; ed. II. i. 245 (quid ?).
aureus, Meyen, Reise. i. 447 = Echinocactus aureus.
bicolor, Berland. ex Engelm. Cact. Mex. Bound. 21 = Echinocactus uncinatus.
Bleo, H. B. & K. Nov. Gen. et Sp. vi. 69 = Pereskia Bleo.
Bleo, Torr. in Ann. Lyc. N. York, ii. (1828) 202 = Opuntia arborescens.
bolivianus, Lem. Cact. 88.—Bolivia.
Bonplandii, H. B. & K. Nov. Gen. et Sp. vi. 69 = Opuntia Tuna.
bradypus, Lehm. Ind. Sem. Hort. Hamb. (1826) 17 = Cereus senilis.
brasiliensis, Willd. Enum. Hort. Berol. Suppl. 33 = Opuntia brasiliensis.
bulbispinus, Lem. Cact. 88.—Hab. !
californicus, Torr. & Gray, Fl. N. Am. i. 555 = Opuntia serpentina ?

CACTUS :—
campechianus, Thiery, ex Steud. Nom. ed. I. 131 ; ed. II. i. 243 (quid ?).
canescens, Moç. & Sesse, ex DC. Prod. iii. 459 = Mammillaria lanifera.
caripensis, H. B. & K. Nov. Gen. et Sp. vi. 66 = Cereus caripensis.
cassythoides. Moç. & Sesse, ex DC. Prod. iii. 476 = Rhipsalis Cassytha.
caudata, Gill. ex Loud. Hort. Brit. 194 = Mammillaria caudata.
chiloënsis, Colla, ex Steud. Nom. ed. II. i. 245 = Cereus chiloënsis.
chinensis, Roxb. Hort. Beng. [37] ; Fl. Ind. ii. 476 = Opuntia Dillenii.
chlorocarpus, H. B. & K. Nov. Gen. et Sp. vi. 67 = Cereus chlorocarpus.
clavatus, Lem. Cact. 88 = Opuntia clavata, *Engelm.*
coccinellifer, Steud. Nom. ed. I. 131, vide cochenillifer.
coccineus, Gill. ex Loud. Hort. Brit. 194 = Mammillaria coccinea.
cochenillifer, Linn. Sp. Pl. 468 = Nopalea coccinellifera.
cochenillifer, Moç. & Sesse, ex DC. Prod. iii. 474 = Opuntia Hernandesii.
columnaris, Moç. & Sesse, ex DC. Prod. iii. 459 = Mammillaria acanthophlegma.
communis, Steud. Nom. ed. II. i. 245 = Melocactus communis.
compressus, Salisb. Prod. 348 = Opuntia vulgaris.
coquimbanus, Molina, Saggio Chile, 170 = Cereus eburneus.
cornigerus, Moç. & Sesse, ex DC. Prod. iii. 461 = Echinocactus cornigerus.
coronarius, Cav. Desc. 490 = Cereus cylindricus.
coronatus, Lam. Encyc. i. 537 = Melocactus communis.
coronatus, Willd. Enum. Hort. Berol. Suppl. 30 = Mammillaria coronaria.
corrugatus, Gill. ex Loud. Hort. Brit. 194 = Echinocactus corrugatus.
crispatus, Moç. & Sesse, ex DC. Prod. iii. 462 = Echinocactus arrigens.
cruciformis, Vell. Fl. Flum. 207 ; v. t. 29 = Cereus cruciformis.
curassavicus, Linn. Sp. Pl. 469 = Opuntia curassavica.
curvispinus, Bert. in 'Merc. Chil. 1829.'—Cf. Bull. Férussac, xx. (1830) 108.—Chili.
cylindricus, James, Cat. 182 = Opuntia arborescens.
cylindricus, Lam. Encyc. i. 539 = Opuntia cylindrica.
cylindricus, Orteg. Hort. Matr. 128. t. 16 = Mammillaria coronaria.
cylindricus, Vell. Fl. Flum. 207 ; v. t. 31 = Rhipsalis cylindrica.
decumanus, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia decumana.
depressus, DC. Cat. Hort. Monsp. 84 = Mammillaria discolor.
depressus, Haw. Syn. Pl. Succ. 173 = Echinocactus depressus.
Dillenii, Ker-Gawl. in Bot. Reg. t. 255 = Opuntia Dillenii.
divaricatus, Lam. Encyc. i. 540 = Cereus divaricatus.
eburneus, Link, Enum. Hort. Berol. ii. 22 = Cereus eburneus.
eburneus, Salm. ex Spreng. Syst. ii. 494 = Cereus eburneus.
echinocarpus, Lem. Cact. 88.—Hab. ?
elatior, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia elatior.
elegans, Link, Enum. Hort. Berol. ii. 25 = Phyllocactus phyllanthoides.
elongatus, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia decumana ?
Emoryi, Lem. Cact. 88 = Cereus Emoryi.
ensiformis, *Biden*, *in Gard. Chron.* xx. (1883) 53.—Hab. ?
Epidendrum, Linn. Amoen. Acad. viii. 257 = Rhipsalis undulata ?
erinaceus, Haw. Suppl. 74 = Cereus erinaceus.
euphorbioides, Spreng. Syst. ii. 496 = Cereus euphorbioides.
fascicularis, Meyen, Reise, i. 447 = Echinocactus fascicularis.
fasciculatus, Willd. Enum. Hort. Berol. Suppl. 33 = Rhipsalis fasciculata.
ferox, Nutt. Gen. Am. i. 296 = Opuntia missouriensis.
ferox, Willd. Enum. Hort. Berol. Suppl. 35 = Opuntia ferox.

CACTUS :—
Ficus, Thunb. Fl. Jap. 198 – seq.
Ficus-indica, Linn. Sp. Pl. 468 = Opuntia Ficus-indica, decumana.
fimbriatus, Desc. Fl. Antill. vi. 160 = Cereus Swartzii.
fimbriatus, Lam. Encyc. i. 539 = Cereus fimbriatus.
fimbriatus, Moç. & Sesse, ex DC. Prod. iii. 475 = Pereskia lychnidiflora.
flagelliformis, Linn. Sp. Pl. 467 = Cereus flagelliformis.
flavescens, DC. Cat. Hort. Monsp. 83 = Mammillaria flavescens.
flavispinus, Colla, Hort. Rip. 24 = Cereus flavispinus.
foliosus, Gill. ex Loud. Hort. Brit. 194 = Echinocactus foliosus.
foliosus, Willd. Enum. Hort. Berol. Suppl. 35 = Opuntia foliosa.
fragilis, Nutt. Gen. Am. i. 296 = Opuntia fragilis.
frutescens, Moç. & Sesse, ex Pfeiff. Enum. Diag. Cact. 178 = Pereskia rotundifolia.
fulvispinosus, Spreng. Syst. ii. 497 = Cereus Royeni.
funalis, Spreng. l. c. 479 = Rhipsalis funalis.
garipensis, Kunth, Syn. Pl. Aeq. iii. 370 = Cereus caripensis.
gibbosus, Haw. in Bot. Reg. t. 137 = Echinocactus gibbosus.
glomeratus, Lam. Encyc. i. 257 = Mammillaria glomerata.
grandiflorus, Linn. Sp. Pl. 467 = Cereus grandiflorus.
grandifolius, Link, Enum. Hort. Berol. ii. 25 = Pereskia grandifolia.
Haworthii, Spreng. Syst. ii. 495 = Cereus Haworthii.
Helicteres, Moç. & Sesse, ex DC. Prod. iii. 460 = Mammillaria Helicteres.
heptagonus, Linn. Sp. Pl. 466 = Cereus heptagonus.
heptagonus, Vell. Fl. Flum. 205 ; v. t. 19.—Bras.
heterogonus, DC. Prod. iii. 470, sphalm. = praec.
hexagonus, Lam. Encyc. i. 538 = Melocactus monoclonos.
hexagonus, Linn. Sp. Pl. 466 = Cereus hexagonus.
hexagonus, Vell. Fl. Flum. 205 ; v. t. 18.—Bras.
Histrix, Salm-Dyck, Obs. Bot. (1822) = Cereus Hystrix.
horridus, H. B. & K. Nov. Gen. et Sp. vi. 70 = Pereskia horrida.
horridus, Salisb. Prod. 348 = Opuntia Tuna.
Humboldtii, H. B. & K. Nov. Gen. et Sp. vi. 66 = Cereus Humboldtii.
humifusus, Rafin. Ann. Nat. (1820) 15 = Opuntia vulgaris.
humilis, Desf. ex Steud. Nom. ed. II. i. 246 = Opuntia vulgaris.
humilis, Haw. Misc. 187 = Opuntia Tuna.
humilis, Lag. Gen. et Sp. Nov. 16 = Opuntia foliosa.
hybridus, Hort. Angl. ex P. C. Van Geel, Sert. Bot. iv. (1832) t. 118.—Hab. ?
hyptiacanthus, Lem. ex Steud. Nom. ed. II. i. 246 = Echinocactus hyptiacanthus.
icosagonus, H. B. & K. Nov. Gen. et Sp. vi. 67 = Cereus icosagonus.
imbricatus, Lem. Cact. 88.—Hab. ?
indicus, Roxb. Hort. Beng. [37] ; Fl. Ind. ii. [475] = Opuntia Dillenii.
intortus, Mill. Gard. Dict. ed. VIII. n. 2 = Echinocactus intortus.
italicus, Tenore, ex Steud. Nom. ed. II. ii. 246 (quid ?).
Jamacaru, Kostel. Allg. Med.-Pharm. Fl. iv. 1393 = Cereus Jamacaru.
Kageneckii, Gmel. ex Salm-Dyck, fide DC. Prod. iii. 463 = Cereus multangularis.
laetus, H. B. & K. Nov. Gen. et Sp. vi. 68 = Cereus laetus.
Lamarkii, Colla, Hort. Ripul. App. iii. 15. t. 7 = Melocactus communis.
lanatus, H. B. & K. Nov. Gen. et Sp. vi. 68 = Cereus lanatus.
lanceolatus, Haw. Misc. 188 = Opuntia lanceolata.
Langsdorfii, Lehm. Ind. Sem. Hort. Hamb. (1826) = Echinocactus Langsdorfii.
lanuginosus, Linn. Sp. Pl. 467 = Cereus repandus.
latispinus, Haw. in Phil. Mag. lxiii. (1824) 41 = Echinocactus latispinus.
Lecchii, Colla, Hort. Ripul. App. iv. 6. t. 2 = Cereus Lecchii.
Lemarii, Monvill. ex 'Lem. Cact. Gen. 105' = Echinocactus Lemarii.
Linkii, Lehm. Ind. Sem. Hort. Hamb. (1827) 16 = Cereus Linkii.
lucidus, Salisb. Prod. 349 = Pereskia aculeata.
luteus, Thiery, ex Steud. Nom. ed. I. 132 ; ed. II. i. 246 (quid ?).

CACTUS :—
lyratus, Vell. Fl. Flum. 205 ; v. t. 21 = Rhipsalis salicornoides.
macracanthus, Salm-Dyck, Obs. (1820) 1 = Melocactus macracanthus.
magnimamma, Salm-Dyck, ex Steud. Nom. ed. II. i. 246 = Mammillaria magnimamma.
Mallisoni, Hort. ex Loud. Encyc. Pl. Suppl. i. 1202 = Cereus Smithii.
mammillaris, Linn. Sp. Pl. 466 = Mammillaria simplex.
mammillaris, Nutt. Gen. Am. i. 295 = Mammillaria missourensis.
melocactoides, Hoffmgg. Verz. Pfl. Nachtr. iii. 24 = Melocactus melocactoides.
Melocactus, Linn. Sp. Pl. 466 = Melocactus communis.
mensarum, Thiery, ex Steud. Nom. ed. I. 132 ; ed. II. i. 246 (quid ?).
meonacanthus, Lem. ex Steud. Nom. ed. II. i. 246 = Echinocactus meonacanthus.
micranthus, H. B. & K. Nov. Gen. et Sp. vi. 65 = Rhipsalis micrantha.
microdasys, Lehm. Ind. Sem. Hort. Hamb. (1827) 16 = Opuntia microdasys.
microthele, Spreng. Syst. ii. 494 = Mammillaria parvimamma.
monacanthos, Willd. Enum. Hort. Berol. Suppl. 33 = Opuntia monacantha.
moniliformis, Lam. Encyc. i. 541 = Cereus moniliformis.
moniliformis, Linn. Sp. Pl. 468 = Opuntia moniliformis.
multangularis, Moç. & Sesse, ex DC. Prod. iii. 462 = Echinocactus melocactiformis.
multangularis, Voigt, ex Steud. Nom. ed. I. 132 = Echinocactus recurvus.
multangularis, Willd. Enum. Hort. Berol. Suppl. 33 = Cereus multangularis.
nanus, H. B. & K. Nov. Gen. et Sp. vi. 68 = Cereus nanus.
nanus, Hort. ex Steud. Nom. ed. II. ii. 246 = Opuntia vulgaris.
Napoleonis, Hort. ex Loud. Encyc. Pl. Suppl. i. 1202 = Cereus Napoleonis.
neglectus, Dehnhardt, Revist. Napol. i. III. 166 (= Pereskiae sp.).
niger, Spreng. Syst. ii. 495 = Cereus niger.
nigricans, Haw. Misc. 187 = Opuntia nigricans.
nobilis, Haw. Syn. Pl. Succ. 174 = Cereus reductus.
nobilis, Lam. Encyc. i. 537 = Echinocactus intortus.
nobilis, Linn. Mant. 243 = Echinocactus recurvus.
nudus, Moç. & Sesse, ex DC. Prod. iii. 460 = Mammillaria nuda.
obvallatus, Moç. & Sesse, ex DC. l. c. 462 = Echinocactus obvallatus.
octogonus, Page, ex Steud. Nom. ed. II. i. 246 = Cereus hexagonus.
Opuntia, Blanco, Fl. Filip. ed. I. 414 = Nopalea coccinellifera.
Opuntia, Guss. Fl. Sicul. Prod. 559 = Opuntia Ficus-indica.
Opuntia, Linn. Sp. Pl. 468 = Opuntia vulgaris.
opuntiaeflorus, Moç. & Sesse, ex Pfeiff. Enum. Diag. Cact. 178 = Pereskia opuntiaeflora.
Ottonis, Lehm. Ind. Sem. Hort. Hamb. (1827) 16 = Echinocactus Ottonis.
ovatus, Gill. ex Loud. Hort. Brit. 195 = Cereus ovatus.
ovoides, Lem. Cact. 88.—Hab. ?
oxypetalus, Moç. & Sesse, ex DC. Prod. iii. 470 = Phyllocactus latifrons.
paniculatus, Lam. Encyc. i. 540 = Cereus paniculatus.
paradoxus, Hornem. Hort. Hafn. ii. 443 – Opuntia brasiliensis.
parasiticus, Lam. Encyc. i. 541 = Rhipsalis fasciculata.
parasiticus, Linn. Syst. ed. X. 1054 = Rhipsalis undulata.
Parryi, Lem. Cact. 88—Hab. ?
parvifolius, Ehrh. ex F. G. Dietr. Vollst. Lex. Gaertn. ii. 416.—Hab. ?
parvispinus, Haw. Suppl. 73 = Echinocactus parvispinus.
pendulinus, Sieber, ex DC. Prod. iii. 476 = Rhipsalis Cassyta.
pendulus, Sw. Prod. Veg. Ind. Occ. 77 = Rhipsalis Cassutha.
pentagonus, Linn. Sp. Pl. 467 = Cereus pentagonus.
pentagonus, Vell. Fl. Flum. 205 ; v. t. 22.—Bras.
Pentlaudi, Lem. Cact. 88.—Hab. ?

CACTUS :—
Pereskia, Linn. Sp. Pl. 469 = Pereskia aculcata.
peruvianus, DC. Pl. Grasscs, t. 58 = Cereus haxagonus.
peruvianus, Linn. Sp. Pl. 467 = Cereus peruvianus.
peruvianus, Willd. Enum. Hort. Berol. Suppl. 32 = Cereus eburneus.
phyllanthoides, DC. Hort. Monsp. 84 = Phyllocactus phyllanthoidcs.
Phyllanthus, Linn. Sp. Pl. 469 = Phyllocactus Phyllanthus.
Phyllanthus, Vell. Fl. Flum. 208 ; v. t. 33.—Bras.
Pitajaya, Blanco, Fl. Filip. ed. II. 289 = Cereus triangularis.
Pitajaya, Jacq. Enum. Pl. Carib. 23 = Cereus variabilis.
placentiformis, Lchm. Ind. Sem. Hort. Hamb. (1826) 17 = Melocactus placcntiformis.
polyanthos, Sims, Bot. Mag. t. 2691 = Opuntia polyantha.
polygonus, Lam. Encyc. i. 539 = Cereus polygonus.
polymorphus, Gill. ex Loud. Hort. Brit. 195 = Cereus polymorphus.
portulacaefolius, Linn. Sp. Pl. 469 = Percskia portulacaefolia.
prismaticus, Desf. Tabl. Hort. Par. 165 = Cereus triqueter.
prismaticus, Willd. Enum. Hort. Berol. Suppl. 32 = Cercus pcntagonus, prismaticus.
prolifer, Willd. ex Pfeiff. Enum. Diagn. Cact. 9 = Mammillaria parvimamma.
proliferus, Mill. Gard. Dict. ed. VIII. n. 6 = Mammillaria prolifera.
proteiformis, Desf. Tabl. Hort. Par. ed. II. 192 = Cereus proteiformis.
pruinosus, Monvill. ex Stcud. Nom. ed. II. i. 246 = Echinocactus pruinosus.
pseudococcinillifer, Bertol. ' Exc. 11 virid. (1824) 4 '; ex DC. Prod. iii. 473 = Opuntia nigricans.
pseudomammillaris, Desf. Tabl. Hort. Par. ed. II. 191 = Mammillaria discolor.
Pseudotuna, Hort. ex Stcud. Nom. ed. II. i. 246 = Opuntia Pseudotuna.
pubescens, Lem. Cact. 87.—Hab. ?
pusillus, DC. Cat. Hort. Monsp. 184 = Mammillaria pusilla.
pusillus, Haw. Misc. 188 = Opuntia fragilis.
pyramidalis, Salm-Dyck, Obs. (1820) 4 = Melocactus pyramidalis.
quadrangularis, Haw. Syn. Pl. Succ. 181 = Cereus caripensis.
quadratus, Gill. ex Loud. Hort. Brit. 194 = Mammillaria quadrata.
recurvus, Mill. Gard. Dict. ed. VIII. n. 3 = Echinocactus recurvus.
reductus, Link, Enum. Hort. Berol. ii. 21 = Echinocactus gibbosus.
regalis, Spreng. Syst. ii. 496 = Cereus regalis.
repandus, Linn. Sp. Pl. 467 = Cereus gracilis.
reptans, Salm-Dyck, ex DC. Prod. iii. 468 = Cereus pentagonus.
reptans, Willd. Enum. Hort. Berol. Suppl. 33 = Cereus radicans.
reticulata, Desc. Fl. Antill. i. t. 68.—Ind. occ.
Rosa, Vell. Fl. Flum. 206 ; v. t. 27 = Pereskia grandifolia.
rotundifolius, Moç. & Sesse, ex DC. iii. 475 = Pereskia rotundifolia.
Roycni, Linn. Sp. Pl. 467 = Cereus Roycni.
Royeni, Willd. Enum. Hort. Berol. Suppl. 32 = Cereus crenulatus.
salicornoides, Steud. Nom. cd. II. i. 246 = Rhipsalis salicornoides.
Salmianus, Lem. Cact. 87.—Hab. ?
Scopa, Link, Enum. Hort. Bcrol. ii. 21 = Cereus Scopa.
senilis, Haw. in Phil. Mag. lxiii. (1824) 41 = Cereus senilis.
sepium, H. B. & K. Nov. Gen. et Sp. vi. 66 = Cereus sepium.
sericeus, Gill. ex Loud. Hort. Brit. 196 = Opuntia sericea.
serpens, H. B. & K. Nov. Gen. et Sp. vi. 68 = Cereus serpens.
serpentinus, Lag. & Rodr. in Anal Cienc. Nat. iv. (1801) 261 = Cereus serpentinus.
serpentinus, Pers. Syn. ii. 23 = Cereus flagelliformis.
solitarius, Gill. ex Loud. Hort. Brit. 194 = Mammillaria solitaria.
speciosissimus, Desf. in Mém. Mus. Par. iii. (1817) 193. t. 9 = Cereus speciosissimus.

CACTUS :—
speciosus, Cav. in Anal. Cienc. Nat. vi. (1803) 339 = Cereus speciosissimus.
speciosus, Desf. Tabl. Hort. Par. 191 = Phyllocactus phyllanthoidcs.
speciosus, Gill. ex Swect, Hort. Brit. ed. II. 235 = Mammillaria speciosa.
speciosus, Hook. Bot. Mag. t. 2820 = Rhipsalis Swartziana.
Spinii, Colla, Antol. vi. 501 = Mammillaria discolor.
spinosissimus, Martyn, Cat. Hort. Cant. 88 = Opuntia spinosissima.
spinosus, Gill. ex Loud. Hort. Brit. 194 = Mammillaria spinosa.
stellaris, Salm-Dyck, Hort. Dyck. ed. II. 7 = Mammillaria pusilla.
stellatus, Willd. Enum. Hort. Berol. Suppl. 30 = Mammillaria pusilla.
stramineus, Spreng. Syst. ii. 494 = Mammillaria flavescens.
strictus, Haw. Misc. 188 = Opuntia stricta.
strictus, Willd. Enum. Hort. Bcrol. Suppl. 32 = Cereus strictus.
subinermis, Link, ex Steud. Nom. ed. II. i. 246 = Opuntia coccinellifera.
subquadriflorus, Moç. & Sesse, ex DC. Prod. iii. 471 = Opuntia rosca.
subrepandus, Spreng. Syst. ii. 495 = Cercus subrepandus.
sulphureus, Gill. ex Loud. Hort. Brit. 196 = Opuntia sulphurea.
sylvestris, Thiery, ex Steud. Nom. ed. I. 132 ; cd. II. i. 246 (quid ?).
tenuis, Schott, ex DC. Prod. iii. 469 = Cereus tenuis.
teres, Vell. Fl. Flum. 207 ; v. t. 30 = Rhipsalis fasciculata.
tetragonus, Linn. Sp. Pl. 466 = Cereus tetragonus.
tomentosus, Link, Enum. Hort. Berol. ii. 24 = Opuntia tomentosa.
torquatus, Hort. Lugd. ex Walp. Rep. ii. 342 = Rhipsalis rhombea.
triacanthos, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia triacantha.
triangularis, Linn. Sp. Pl. 468 = Cereus compressus, trigonus.
triangularis, Vell. Fl. Flum. 206 ; v. t. 24 = Cereus triangularis.
trichotomus, Tenore, ex Steud. Nom. ed. II. i. 246 (quid ?).
triqueter, Haw. Misc. 189 = Cereus triqueter.
triqueter, Vell. Fl. Flum. 206 ; v. t. 25 = Cereus Arrabidae.
triqueter, Willd. Enum. Hort. Berol. Suppl. 33 = Cereus trigonus.
truncatus, Link, Enum. Hort. Berol. ii. 24 = Epiphyllum Altensteinii ?
tuberculatus, Willd. Enum. Hort. Berol. Suppl. 34 = Opuntia tuberculata.
Tuna, Linn. Sp. Pl. 468 = Opuntia Tuna.
Tuna-elatior, Roxb. Hort. Beng. [37] nomen.—Am. austr.
Tuna-major, Roxb. l. c.—Am. austr.
tunicatus, Lchm. Ind Sem. Hort. Hamb. (1827) 17 = Opuntia tunicata.
undulosus, Kostel. Allg. Med.-Pharm. Flora, iv. 1393 = Cereus variabilis.
Urumbeba, Vell. Fl. Flum. 207 ; v. t. 32 = Opuntia triacantha.
Urumbella, Steud. Nom. ed. II. i. 246 = praec.
verticillatus, Brot. ex Heynh. Nom. ii. 103, nomen (quid ?).
viviparus, Nutt. Gen. Am. i. 295 = Mammillaria vivipara.
zinniaeflorus, Moç. & Sesse, ex DC. Prod. iii. 475 = Pereskia zinniaeflora.

CACUCIA, J. F. Gmel. Syst. 693 (1791) = **Cacoucia**, Aubl. (Combret.).

CACUVALLUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 399 = **Mucuna**, Adans. (Legum.).
altissimus, Medic. l. c. 400 = M. urens.

CADABA, Forsk. Fl. Aegypt. Arab. 67 (1775). *CAPPARIDEAE*, Benth. & Hook. f. i. 108.
Desmocarpus, Wall. Cat. n. 6278 (1832).
Macromerum, Burch. Trav. i. 388 (1822) ; ii. 615.
Schepperia, Neck. Elem. iii. 67 (1790).
Stroemeria, Roxb. Fl. Ind. ii. 78 (1832).
Stroemia, Vahl, Symb. Bot. i. 19 (1790).

CADABA :—
capparoides, *DC. Prod.* i. 244.—Ins. Molucc.
dubia, DC. l. c. = farinosa.
farinosa, *Forsk. Fl. Aegypt. Arab.* 68.—Ind. bor. occ. ; Arab. ; Afr. trop.
glandulosa, *Forsk. l. c.*—Arabia.
glandulosa, Miq. in Verh. Nederl. Inst. III. v. (1852) 2 = Miqueliana.
heterotricha, *Stocks, in Hook. Ic. Pl.* t. 839.—Sindh.
indica, *Lam. Encyc.* i. 544.—Ind. or.
juncea, *Harv. Gen. S. Afr. Pl.* ed. II. 13.—Afr. austr.
Kirkii, *Oliver, Fl. Trop. Afr.* i. 90.—Afr. trop.
longifolia, *DC. Prod.* i. 244.—Abyss.
madagascariensis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 464.—Madag.
Miqueliana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 321.—Malaya.
monopetala, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1212 = glandulosa.
natalensis, *Sond. in Linnaea*, xxiii. (1850) 8.—Natal.
rotundifolia, *Forsk. Fl. Aegypt. Arab.* 68.—Arab. ; Abyss.
somalensis, *Franch. in Revoil, Çomali*, 12.—Afr. trop.
trifoliata, *Wight & Arn. Prod.* 24.—Ind. or.
trinervia, DC. Prod. i. 244 = Cleome coluteoides.
triphylla, Wight, in Hook. Bot. Misc. iii. (1833) 296 = trifoliata.
virgata, *Boj. in Ann. Sc. Nat.* Sér. II. xx. (1843) 58.—Madag.

CADALVENA, Fenzl, in Sitz. Akad. Wien, Math. Nat. li. Abth. ii. (1865) 139 = **Kaempferia**, Linn. (Scitamin.).
spectabilis, Fenzl, l. c. 140.—Afr. trop.

CADAMBA, Sonner. Voy. ii. 228. t. 128 (1782) = **Guettarda**, Linn. (Rubiac.).
jasminiflora, Sonner. l. c. ii. t. 128 ; iii. 253 = G. speciosa.

CADELARI, Adans. Fam. ii. 268 (1763) = **Pupalia**, Juss. (Amarantac.).
lappacea, Medic. Malv. 92 = P. atropurpurea.

CADELARIA, Rafin. Fl. Tellur. iii. 39 (1836) = **Achyranthes**, Linn. (Amarantac.).
indica, Rafin. l. c. = A. aspera.
punctata, Rafin. l. c. = A. aspera ?
sicula, Rafin. l. c. = A. aspera.

CADELIUM, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 352 ; Phil. Bot. i. 206 (1789) = **Phaseolus**, Tourn. (Legumin.).
nigrum, Medic. l. c. 352 = P. Max.
viride, Medic. l. c. = P. Max.

CADELLIA, F. Muell. Fragm. ii. 25 (1860). *SIMARUBEAE*, Benth. & Hook. f. i. 313.
Guilfoylia, F. Muell. Fragm. viii. 33 (1873).
monostylis, *Benth. Fl. Austral.* i. 375.—Austral.
pentastylis, *F. Muell. Fragm.* ii. 26. t. 12.—Austral.

CADETIA, Gaudich. in Freyc. Voy. Bot. 422. t. 33 (1826) = **Dendrobium**, Sw. (Orchid.).
angustifolia, Blume, Rumphia, iv. 39 = D. angustifolium ?
biloba, Blume, Mus. Bot. Lugd. Bat. i. (1849) 30 = D. bilobum.
recurvata, Blume, l. c.—N. Guinea.
similis, Blume, Rumphia, iv. 29, in obs.—N. Guinea.
umbellata, Gaudich. in Freyc. Voy. Bot. 422. t. 33 = D. umbellatum.

CADIA, Forsk. Fl. Aegypt. Arab. 90 (1775). *LEGUMINOSAE*, Benth. & Hook. f. i. 560.
Panciatica, Picciv. Hort. Panciat. 9 (1783).
Spaendoncea, Desf. Dec. Phil. vii. 259 ; et in Usteri, N. Ann. xiii. (1796) 15.
anomala, *Vatke, in Linnaea*, xliii. (1880–82) 337.—Madag.
arabica, Raeusch. Nom. ed. III. 117 = purpurea.
Commersoniana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 370.—Madag.
Ellisiana, *Baker, in Journ. Linn. Soc.* xx. (1884) 135.—Madag.
pedicellata, *Baker, l. c.* 338.—Madag.
pubescens, *Boj. ex Baker, in Oliver, Fl. Trop. Afr.* ii. 256.—Madag.
purpurea, *Forsk. Fl. Aegypt. Arab.* 90.—Arab. felix
varia, L'Herit. Diss. in Mag. Enc. v. 29 = purpurea.

CADISCUS, E. Mey. ex DC. Prod. vii. 254 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 408.
aquaticus, *E. Mey. l. c.* 255.—Afr. austr.

CADSURA, Spreng. Syst. ii. 642 (1827) = **Kadsura**, Juss. (Magnoliac.).

CAELESTINA, Cass. in Dict. Sc. Nat. vi. Suppl. 8. t. 93 (1817) = **Ageratum**, Linn. (Compos.).
ageratoides, H. B. & K. Nov. Gen. et Sp. iv. 151 = A. corymbosum.
albida, DC. Prod. v. 107 = A. albidum.
caerulea, Cass. in Dict. Sc. Nat. vi. Suppl. 8 = A. corymbosum.
cordata, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 345 = Eupatorium macrophyllum.
corymbosa, Benth. Bot. Voy. Sulph. 111 = A. salicifolium.
corymbosa, DC. Prod. v. 108 = A. corymbosum.
Hartwegi, Walp. Rep. ii. 545 = Eupatorium Hartwegi.
hastata, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 363 = Eupatorium betonicaeforme.
isocarphoides, DC. Prod. v. 107 = A. isocarphoides.
latifolia, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 71 = A. latifolium.
Lessingiana, Klotzsch, ex Walp. Rep. ii. 545 = A. corymbosum.
linearifolia, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 191 = Alomia polyphylla.
longifolia, Pohl, ex Baker, l. c. 361 = Eupatorium organense.
maritima, Torr. & Gray, Fl. N. Am. ii. 64 = A. littorale.
micrantha, Spreng. Syst. iii. 446 = A. coelestinum.
microcarpa, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 72 = A. microcarpum.
paleacea, J. Gay, ex DC. Prod. v. 107 = A. paleacenm.
parvifolia, DC. l. c. 108.—Am. austr.
petiolata, Hook. & Arn. Bot. Beech. Voy. 433 = A. petiolatum.
Pohlii, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 190 = Alomia Pohlii.
repanda, Pohl, ex Baker, l. c. 309 = Eupatorium viridiflorum.
repens, Sch. Bip. in Schomb. Fauna et Flora Guy. 1138.—Guiana.
scabriuscula, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 72 = A. scabriusculum.
suffruticosa, Sweet, Hort. Brit. ed. I. 229 = A. corymbosum.
tomentosa, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 71 = A. tomentosum.
urticaefolia, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 362 = Eupatorium betonicaeforme.
viscosa, Pohl, ex Baker, l. c. = Eupatorium apiculatum.

CAELIA, G. Don, in Sweet, Hort. Brit. ed. III. 637 (1839) = **Coelia**, Lindl.

CAELOCLINE, Auct. ex Steud. Nom. ed. II. i. 247 (1840) = COELOCLINE, A. DC. = **Xylopia**, Linn. (Anonac.).

CAELODEPAS, Benth. & Hook. f. Gen. iii. 313 (1880) = **Coelodepas**, Hassk. (Euphorb.).

CAELOGLOSSUM, Steud. Nom. ed. II. i. 247 (1840) = COELOGLOSSUM, Lindl. = **Habenaria**, Willd. (Orchid.).

CAELOGYNE, Wall. ex Steud. Nom. ed. II. i. 247 (1840) = **Coelogyne**, Lindl. (Orchid.).

CAELOSPERMUM, Blume, Bijdr. 994 (1826), vide **Coelospermum**, Blume.

CAENOTUS, Rafin. Fl. Tellur. ii. 50 (1836) = **Erigeron**, Linn. (Compos.).
canadensis, Rafin. l. c. = E. canadensis.
longifolius, Rafin. l. c. = E. longifolius?
pusillus, Rafin. l. c. = E. canadensis.

CAESALPINA, Plum. ex Linn. Gen. ed. I. 363 (1753) = seq.

CAESALPINIA, Linn. Sp. Pl. 380 (1753). *LEGUMINOSAE*, Benth. & Hook. f. i. 565.
BALSAMOCARPON, Clos, in C. Gay, Fl. Chil. ii. 226. t. 20 (1846).
BIANCAEA, Todaro, Nuov. Gen. 21 (1858).
BONDUC, Adans. Fam. ii. 318 (1763).
CAMPECIA, Adans. l. c. (1763).
CINCLIDOCARPUS, Zoll. in Nat. en Geneesk. Arch. Neêrl. Indie, iii. (1846) 81.
CLADOTRICHIUM, Vog. in Linnaea, xi. (1837) 401.
COULTERIA, H. B. & K. Nov. Gen. et Sp. vi. 328. t. 568 (1823).
ERYTHROSTEMON, Klotzsch, in Link, Klotzsch & Otto, Ic. Pl. i. 97. t. 39 (1841).
GUILANDIA, P. Br. Hist. Jamaic. 228 (1756).
GUILANDINA, Linn. Gen. ed. I. 124 (1737).
LIBIDIBIA, Schlecht. in Linnaea, v. (1830) 192.
POMARIA, Cav. Ic. v. 1. t. 402 (1799).
TARA, Molina, Saggio Chile, ed. II. 153 (1810).
TICANTO, Adans. Fam. ii. 319 (1763).
acanthobotrya, *Miq. Fl. Ind. Bat. Suppl.* i. 292.—Sumatra.
acinaciformis, *Mart. ex Steud. Nom.* ed. II. i. 247.—Bras.
affinis, *Hemsl. Diag. Pl. Nov.* 8.—Guatemala.
angulicaulis, *Clos, in C. Gay, Fl. Chil.* ii. 223.—Chili.
angustifolia, Salisb. Prod. 326 = C. Sappan.
arborea, Zoll. ex Miq. Fl. Ind. Bat. i. 112 = Peltophorum ferrugineum.
Arista, Mill. Gard. Dict. ed. VIII. n. 2, sphalm. = Crista.
armata, R. Grah. in Wall. Cat. n. 5840 = mimosoides.
axillaris, DC. Prod. ii. 481 = C. sepiaria? et Mezoneurum cucullatum.
bahamensis, Lam. Encyc. i. 461 = Crista.
Barbon, *Bert. in 'Merc. Chil.* (1829)'; *et in Bull. Férussac*, xx. (1830) 109.—Chili.
Berterii, Spreng. ex DC. Prod. ii. 482 = bijuga.
bijuga, *Sw. Obs.* 166.—Ind. occ.
bijuga, Wall. Cat. n. 5833 = C. Nuga.
Bonduc, *Roxb. Hort. Beng.* 32; *Fl. Ind.* ii. 362.—Cosmop. trop.
Bonduc, Wight & Arn. Prod. 280 = C. Bonducella.
Bonducella, *Fleming, in As. Res.* xi. (1810) 159.—Cosmop. trop,
bracteosa, *Tul. in Arch. Mus. Par.* iv. (1844) 141.—Bras.
brasiliensis, *Linn. Sp. Pl.* 380.—Am. trop.
Brea, Gill. ex Steud. Nom. ed. II. i. 247 = praecox.
brevifolia, *Baill. Hist. Pl.* ii. 168.—Chili.
Cacalaco, *Humb. & Bonpl. Pl. Aeq.* ii. 173. t. 137.—Mexic.
calycina, *Benth. in Mart. Fl. Bras.* xv. II. 71.—Am. trop.
cassioides, *Willd. Enum. Hort. Berol.* 444.—Am. austr.
Cavanillesii, Spreng. Syst. v. Ind. 123 = bijuga.
chinensis, Roxb. Hort. Beng. 32; Fl. Ind. ii. 361 = C. Nuga.
Cinclidocarpus, *Miq. Fl. Ind. Bat.* i. 110.—Ind. or.; Malaya.
coluteifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 111.—Reg. Argent.
compressa, *D. Dietr. Syn. Pl.* ii. 1494.—Mexic.; Peruv.
coriacea, Poir. in Rev. Hortic. Sér. II. ii. (1843–44) 269 = coriaria.
coriaria, *Willd. Sp. Pl.* ii. 532.—Am. austr.
corymbosa, *Benth. Pl. Hartw.* 117.—Guayaqnil.
coulterioides, *Griseb. in Goett. Abh.* xxiv. (1879) 113.—Reg. Argent.
Courboniana, *Baill. Adansonia*, x. (1871) 103.—Mare rubrum.
Crista, *Linn. Sp. Pl.* 380.—Ind. occ.
Crista, Thunb. Fl. Jap. 179 = sepiaria.
cucullata, Roxb. Hort. Beng. 32; Fl. Ind. ii. 358 = Mezoneurum cucullatum.
dasyrachis, Miq. Fl. Ind. Bat. Suppl. i. 292 = Peltophorum dasyrachis.
desertorum, Mart. ex G. Don, Gen. Syst. ii. 431 = bracteosa.
digyna, *Rottl. in Ges. Naturf. Freunde Neue Schr.* iv. (1803) 200.—Ind. or.; Malaya.
dubia, Spreng. Syst. ii. 343 = Peltophornm Vogelianum.
Ebano, *Karst. Fl. Columb.* ii. 57. t. 129.—N. Granat.
echinata, *Lam. Encyc.* i. 461.—Bras.
elata, Sw. Obs. 166 = Poinciana elata.

CAESALPINIA:—
enneaphylla, Roxb. Hort. Beng. 32; Fl. Ind. ii. 363 = Mezoneurum enneaphyllum.
eriostachys, *Benth. Bot. Voy. Sulph.* 88.—Am. centr.
exilifolia, *Griseb. in Goett. Abh.* xix. (1874) 128.—Reg. Argent.
exostemma, *Moç. & Sesse, ex DC. Prod.* ii. 483.—Nicaragua.
ferox, Hassk. Ind. Sem. Hort. Amst. (1841); Pl. Jav. Rar. 400 = sepiaria.
ferox, Hohen. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 261 = Wagatea spicata.
ferrea, *Mart. Reise Bras.* ii. 611.—Bras.
ferruginea, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 462 = Peltophorum ferrugineum.
fimbriata, *Tul. in Arch. Mus. Par.* iv. (1844) 145.—Bolivia.
Finlaysoniana, R. Grah. in Wall. Cat. n. 5971 = Peltophorum dasyrachis.
Fischeri, *Regel & Koern. in Gartenfl.* (1858) 251.—Ind. or.?
flavicans, R. Grah. in Wall. Cat. n. 5825 = digyna.
floribunda, *Tul. in Arch. Mus. Par.* iv. (1844) 140.—Bras.
furfuracea, Wall. Cat. n. 5835 = Mezoneurum glabrnm.
Gardneriana, *Benth. in Mart. Fl. Bras.* xv. II. 68.—Bras.
Gilliesii, *Wall. ex Hook. Bot. Misc.* i. (1830) 129.—Am. austr.
glabrata, *H. B. & K. Nov. Gen. et Sp.* vi. 326.—Peruv.
glandulosa, Bert. ex DC. Prod. ii. 482 = bijuga.
Glenieii, *Thw. Enum. Pl. Zeyl.* 414 = Peltophorum ferrugineum.
gracilis, *Benth. ex Hemsl. Diag. Pl. Nov.* 9.—Mexic.
gracilis, Miq. Fl. Ind. Bat. i. 110 = digyna.
grandis, *Heyne, in Wall. Cat.* n. 5830.—Ind. or.
Heyneana, Steud. Nom. ed. II. i. 247 = Wagatea spicata.
Hildebrandtii, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 371.—Afr. trop.
hispida, *D. Dietr. Syn. Pl.* ii. 1495.—Mexic.
horrida, Herb. Madr. ex Wall. Cat. n. 5836 = mimosoides.
horrida, A. Rich. Ess. Fl. Cub. 411 = sepiaria.
horrida, *Spreng. Syst.* iv. *Cur. Post.* 169.—N. Granat.
hymenocarpa, Wall. Cat. n. 5832 = Mezoneurum pubescens.
ignota, Blanco, Fl. Filip. ed. I. 336 = Mezoneurum pubescens.
inermis, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 367 = Peltophorum ferrugineum.
insignis, *Steud. Nom.* ed. I. 133; *D. Dietr. Syn. Pl.* ii. 1494.—Reg. Amazon.
japonica, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1845?) 117 = sepiaria.
kavaiensis, *H. Mann, in Proc. Am. Acad.* vii. (1867) 164.—Ins. Hawaii.
Kilaroe, Span. in Linnaea, xv. (1841) 200 = Peltophorum ferrugineum.
lacerans, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 367 = Pterolobium indicum.
languida, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 72.—Ind. or.
latifolia, Salisb. Prod. 326 = vesicaria.
laxa, *Benth. Pl. Hartw.* 60.—Mexic.
laxiflora, *Tul. in Arch. Mus. Par.* iv. (1844) 143.—Bras.
lebbekkioides, DC. Prod. ii. 483 = Albizzia Millettii.
ligulata, Heyne, ex DC. Prod. ii. 484 = Pterolobium lacerans.
Macklottii, Zipp. ex Miq. Fl. Ind. Bat. i. 112 = Peltophorum ferrugineum.
macrantha, *Delile, Ind. Sem. Hort. Monsp.* (1838) 3.—Chili.
marginata, *Tul. in Arch. Mus. Par.* iv. (1844) 147.—Bras.
melanocarpa, *Griseb. in Goett. Abh.* xix. (1874) 128.—Reg. Argent.
Melanosticta, Spreng. Syst. iv. Cnr. Post. 169 = Melanosticta Burchellii.
mexicana, *A. Gray, in Proc. Am. Acad.* v. (1861) 157.—Mexic.
microphylla, Buch.-Ham. in Wall. Cat. n. 5826 = C. Cinclidocarpa.
microphylla, *Mart. Reise Bras.* ii. 611.—Bras.
Millettii, *Hook. & Arn. Bot. Beech. Voy.* 182.—China.
mimosifolia, *Griseb. in Goett. Abh.* xix. (1874) 128.—Reg. Argent.

CAESALPINIA :—
mimosoides, Heyne, in Wall. Cat. n. 5837 = Wagatea spicata.
mimosoides, *Lam. Encyc.* i. 452.—Ind. or.; Burma.
Minax, *Hance, in Journ. Bot.* xxii. (1884) 365.—China.
mollis, *Spreng. Syst.* iv. *Cur. Post.* 169.—Am. trop.
monosperma, Buch.-Ham. ex Wall. Cat. sub n. 5828 = Mezoneurum cucullatum.
monosperma, *Tul. in Arch. Mus. Par.* iv. (1844) 148.—Ind. occ.
mucronata, *Willd. Enum. Hort. Berol.* 444.—Bras.
nitida, *Hassk. Cat. Hort. Bog. Alt.* 285.—Malaya.
Nuga, *Ait. Hort. Kew.* ed. II. iii. 32.—As. trop.; Austral.; Polynes.
obliqua, Vog. in Linnaea, xi. (1837) 407 = echinata.
oleosperma, Roxb. Hort. Beng. 32; Fl. Ind. ii. 357 = digyna.
Paipae, *Ruiz & Pav. Fl. Per.* iv. t. 375.—Peruv.
paniculata, Desf. Tabl. Hort. Par. ed. II. 210 = C. Nuga.
Parahyba, Allem. in Trab. Soc. Vell. 56 = Schizolobium excelsum.
parvifolia, Steud. Nom. ed. II. i. 247 = C. Cinclidocarpa.
pauciflora, *Benth. & Hook. f. ex Chapm. Fl. S. U. St. Suppl.* 18.—Am. bor.; Ind. occ.
pectinata, Cav. Desc. 467 = tinctoria.
pellucida, *Vog. in Linnaea*, x. (1836) 601.—Ins. S. Doming.
peltophoroides, *Benth. in Mart. Fl. Bras.* xv. II. 72.—Bras.
pilosa, *Benth. l. c.* 73.—Bras.
pluviosa, *DC. Prod.* ii. 483.—Bras.
polyphylla, *Roxb. Hort. Beng.* 32, nomen.—Pegu.
porcina, *Mart. ex Steud. Nom.* ed. II. i. 247.—Bras.
praecox, *Ruiz & Pav. Fl. Per.* iv. t. 376.—Chili.
procera, *Poepp. ex Spreng. Syst.* ii. 343.—Cuba.
pubescens, Buch.-Ham. ex Wall. Cat. sub n. 5834 = sepiaria.
pulcherrima, *Sw. Obs.* 166.—Cosmop. trop.
Pumilio, *Griseb. in Goett. Abh.* xxiv. (1879) 113.—Reg. Argent.
punctata, *Willd. Enum. Hort. Berol.* 445.—Bras.
pyramidalis, *Tul. in Arch. Mus. Par.* iv. (1844) 139.—Bras.
regia, D. Dietr. Syn. Pl. ii. 1495 = Poinciana regia.
resupinata, Roxb. Hort. Beng. 32; Fl. Ind. ii. 362 = mimosoides.
Roxburghii, D. Dietr. Syn. Pl. ii. 1494 = Peltophorum ferrugineum.
rubicunda, *Benth. in Mart. Fl. Bras.* xv. II. 73.—Bras.
Sapang, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 12 = C. Sappan.
Sappan, *Linn. Sp. Pl.* 381.—As. trop.
scandens, *Heyne, ex Roth, Nov. Pl. Sp.* 209.—Ind. or.
scandens, Koen. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 255 = Nuga.
Selloi, G. Don, Gen. Syst. ii. 431 = Peltophorum Vogelianum.
sepiaria, *Roxb. Hort. Beng.* 32; *Fl. Ind.* ii. 360.—As. trop.
sepiaria, Wall. Cat. n. 5834 H. = Mezoneurum enneaphyllum.
Simora, Buch.-Ham. ex Roxb. Hort. Beng. 32; Fl. Ind. ii. 359 = mimosoides.
smilacea, Hort. ex Steud. Nom. ed. II. i. 248 = Cocculus carolinus.
spicata, Dalz. in Hook. Kew Journ. iii. (1851) 89 = Wagatea spicata.
stipularis, *Benth. in Mart. Fl. Bras.* xv. II. 72.—Bras.
suberosa, *Lodd. ex Loud. Hort. Brit.* 167.—Ind. or.
sumatrana, Roxb. Hort. Beng. 32; Fl. Ind. ii. 366 = Mezoneurum sumatranum.
sumatrana, Wall. Cat. n. 5831 B, C = Nuga.
Tara, *Ruiz & Pav. Fl. Per.* iv. t. 374.—Peruv.
thomaea, Spreng. Syst. ii. 343 = coriaria.
tinctoria, *Domb. ex DC. Prod.* ii. 481.—N. Granat.
torquata, Blanco, Fl. Filip. ed. I. 336 = Mezoneurum glabrum.
tortuosa, *Roxb. Hort. Beng.* 32; *Fl. Ind.* ii. 365.—Burma; Malaya; Sumatra.
tortuosa, Wall. Cat. n. 5827. D = C. Cinclidocarpus.
trichocarpa, *Griseb. in Goett. Abh.* xxiv. (1879) 112.—Reg. Argent.

CAESALPINIA :—
ulosperma, *Hort. ex Steud. Nom.* ed. II. i. 248.—Hab.?
ventricosa, Griseb. Fl. Brit. W. Ind. 205 = bijuga.
vernalis, *Champ. ex Benth. in Hook. Kew Journ.* iv. (1852) 77.—China.
vesicaria, Lam. Encyc. i. 461 = bijuga.
vesicaria, *Linn. Sp. Pl.* 381.—Bras.
vesicaria, Vell. Fl. Flum. 172; iv. t. 89 = echinata.
wynaadensis, Heyne, ex Wall. Cat. sub n. 5830 = grandis.

CAESAREA, Cambess. in Mém. Mus. Par. xviii. (1829) 373. t. 18 = **Viviania**, Cav. (Geraniac.).
albiflora, Cambess. l. c. 374.—Bras.
grandiflora, St. Hil. ex Steud. Nom. ed. II. i. 248.—Bras.
hypoleuca, Mart. ex Steud. l. c.—Bras.
lanceolata, Klotzsch, in Linnaea, x. (1836) 437.—Bras.
montevidensis, Klotzsch, l. c. 436.—Reg. Argent.
ovata, Klotzsch, l. c.—Bras.
petiolata, Klotzsch, l. c. 437.—Bras.
rubriflora, Cambess. in Mém. Mus. Par. xviii. (1829) 374.—Bras.

CAESIA, R. Br. Prod. 277 (1810). *LILIACEAE*, Benth. & Hook. f. iii. 792.
acanthoclada, F. Muell. Fragm. i. 215 = Corynotheca acanthoclada.
africana, *Baker, in Trans. Linn. Soc.* xxix. (1875) 160.—Afr. trop.
alpina, Hook. f. Fl. Tasm. ii. 373 = Chlorophytum alpinum.
asparagoides, *Eckl. ex Steud. Nom.* ed. II. i. 248.—Afr. austr.
chlorantha, F. Muell. Fragm. i. 63 = vittata.
coarctata, Spreng. Syst. ii. 88 = Eccremis coarctata.
comosa, Spreng. l. c. = Chlorophytum comosum.
corymbosa, R. Br. Prod. 277 = Chamaescilla corymbosa.
cyanea, A. Cunn. ex Baker, in Journ. Linn. Soc. xv. (1876) 358 = vittata.
dichotoma, F. Muell. Fragm. i. 215 = Corynotheca dichotoma.
Dregeana, Kunth, Enum. Pl. iv. 611 = Anthericum Dregeanum.
Eckloniana, Schult. f. Syst. vii. 1691 = Anthericum scilliflorum.
Hclleri, *Hueg. ex Heynh. Nom.* ii. 103, nomen.—Austral.
hirsuta, Lindl. Swan Riv. App. 57. n. 273 = Agrostocrinum stypandroides.
lateriflora, R. Br. Prod. 277 = Corynotheca lateriflora.
micrantha, Lindl. Swan Riv. App. 57 = parviflora.
occidentalis, R. Br. Prod. 277 = parviflora.
paradoxa, Endl. in Lehm. Pl. Preiss. ii. 34 = Chamaescilla spiralis?
parviflora, *R. Br. Prod.* 277.—Austral.
physodes, Spreng. Syst. ii. 88 = Urginea physodes.
pusilla, Spreng. l. c. = Urginea pusilla.
rigidifolia, *F. Muell. Fragm.* x. 48.—Austral.
scabra, Baker, in Journ. Linn. Soc. xv. (1876) 359 = Agrostocrinum stypandroides.
setifera, *Baker, l. c.*—Austral.
spiralis, Endl. in Lehm. Pl. Preiss. ii. 34 = Chamaescilla spiralis.
Thunbergii, Schult. f. Syst. vii. 1692 = Anthericum contortum.
versicolor, Lindl. Swan Riv. App. 57 = Chamaescilla corymbosa.
vittata, *R. Br. Prod.* 277.—Austral.

CAESIA, Vell. Fl. Flum. 107 (1825); iii. t. 23 (1827) = **Cormonema**, Reissek (Rhamneae).
spinosa, Vell. l. c. = C. spinosum.

CAESULIA, Roxb. Pl. Corom. i. 64. t. 93 (1759). *COMPOSITAE*, Benth. & Hook. f. ii. 318.
axillaris, *Roxb. l. c.*—Ind. or.
lancifolia, *Sweet, Hort. Brit.* ed. II. 288 (quid?).—Ind. or.
radicans, Willd. Sp. Pl. iii. 1797 = Cryphiospermum repens.
scandens, Poepp. ex DC. Prod. v. 564 = Wulffia scandens.

CAETA, Steud. Nom. ed. II. i. 248 (1840) = **Torenia**, Linn. (Scrophular.).

CAETOCAPNIA, Endl. Gen. 180 (1837) = COETOCAPNIA, Link & Otto = **Bravoa**, Lex.

CAFFEA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 12, an **Coffea**, Linn.

CAHOTA, Karst. in Linnaea, xxviii. (1856) 448 = **Clusia**, Linn. (Guttifer.).
carachensis, Karst. l. c.—Venezuela.

CAIDBEJA, Forsk. Fl. Aegypt. Arab. 82 (1775) = **Forskohlea**, Linn. (Urticac.).
adhaerens, Forsk. l. c. = F. tenacissima.

CAILLIEA, Guillem. & Perr. Tent. Fl. Seneg. 239 (1833) = **Dichrostachys**, DC. (Legumin.).
dichrostachys, Guillem. & Perr. l. c. 240 = D. nutans.
macrostachya, Steud. in Flora, xxvi. (1843) 759 = Pentaclethra filamentosa.

CAINA, Panch. ex Baill. in Bull. Soc. Linn. Par. i. (1880) 264 = **Couthovia**, A. Gray (Loganiac.).
erratica, Panch. ex Baill. l. c. = C. corynocarpa.

CAINITO, Plum. ex Adans. Fam. ii. 166 (1763); Tuss. Fl. Antill. iii. 41. t. 9 (1824) = **Chrysophyllum**, Linn. (Sapotac.).
pomiferum, Tuss. Fl. Antill. iii. 41. t. 9 = Chrysophyllum Cainito.

CAIOPHORA, Presl, Rel. Haenk. ii. 41. t. 56 (1831) = **Blumenbachia**, Schrad. (Loas.).
absinthiifolia, Presl, l. c. 42 = B. coronata.
carduifolia, Presl, l. c. = B. carduifolia.
cirsiifolia, Presl, l. c. t. 56.—Peruv.
contorta, Presl, l. c. 42 = Loasa compressa.
coronata, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 327 = B. coronata.
Herberti ×, G. Don, ex Lonr. Encyc. Pl. Suppl. ii. 1438.—Hab.?
lateritia, Benth. in Maund. & Hensl. Botanist, 111. t. 119.—Chili.
Pentlandii, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1438.—Peruv.
scandens, Mey. & Klotzsch, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 339 = B. scandens.

CAJAN, Adans. Fam. ii. 326 (1763) = **Cajanus**, DC. (Legumin.).
inodorum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 363 = Cajanus indicus.
scarabaeoides, Thou. in Dict. Sc. Nat. vi. 167 = Atylosia scarabaeoides.

CAJANUM, Rafin. Sylva Tellnr. 25 (1838) = seq.
Thora, Rafin. l. c. = Cajanus indicus.

CAJANUS, DC. Cat. Hort. Monsp. 85 (1813). *LEGUMINOSAE*, Benth. & Hook. f. i. 541.
albicans, R. Grah. in Wall. Cat. n. 5582 = Atylosia albicans.
argenteus, Spreng. Syst. iii. 248 = Argyrolobium Linnaeanum.
bicolor, DC. Cat. Hort. Monsp. 85 = indicus.
candicans, Wall. Cat. n. 5576 = Rhynchosia aevensis.
cinereus, *F. Muell. Census*, 41.—Austral.
confertiflorus, *F. Muell. Pl. Fitzalan*, 9.—Austral.
flavus, DC. Cat. Hort. Monsp. 85 = indicus.
glandulosus, Dalz. & Gibs. Bomb. Fl. 73 = Atylosia mollis.
goënsis, Dalz. in Hook. Kew Journ. ii. (1850) 264 = Atylosia barbata.
grandifolins, *F. Muell. Pl. Fitzalan*, 9.—Austral.
helvolus, Spreng. Syst. iii. 248 = Glycine helvola.
indicus, *Spreng. l. c.*—Ind. or.
kulnensis, Dalz. in Hook. Kew Journ. ii. (1850) 264 = Atylosia kulnensis.
lineatus, R. Grah. in Wall. Cat. n. 5578 = Atylosia lineata.
luteus, Bello. in Anal. Soc. Esp. Hist. Nat. x. (1881) 260 = indicus?
marmoratus, *F. Muell. Census*, 41.—Austral.
megalanthus, Spreng. ex DC. Prod. ii. 218 = Astragalus megalanthus.
niveus, R. Grah. in Wall. Cat. n. 5581 = Atylosia nivea.
quinquepetalus, Blanco, Fl. Filip. ed. II. 417 = Desmodium Cephalotes.

CAJANUS :—
reticulatus, *F. Muell. Census*, 41.—Austral.
scarabaeoides, Thou. ex R. Grah. in Wall. Cat. n. 5580 = Atylosia scarabaeoides.
striatus, Boj. Hort. Maurit. 109 = indicus.
suaveolens, R. Grah. in Wall. Cat. n. 5579 = Rhynchosia suaveolens.
volubilis, Blanco, Fl. Filip. ed. II. 417 = Atylosia mollis.
Wightianus, R. Grah. in Wall. Cat. n. 5583 = Atylosia albicans.
wolgaricus, Spreng. ex Steud. Nom. ed. II. i. 248 = Calophaca wolgarica.

CAJU-PUTI, Adans. Fam. ii. 84 (1763) = **Melaleuca**, Linn. (Myrtac.).

CAKILE, Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 99.
aegyptia, Spreng. Syst. ii. 852 = Didesmus aegyptiacus.
aegyptiaca, Willd. Sp. Pl. iii. 417 = maritima.
aequalis, L'Hérit. ex DC. Syst. ii. 430 = maritima.
americana, Nutt. Gen. Am. ii. 62 = maritima.
bipinnata, Spreng. Syst. ii. 852 = Didesmus bipinnatus.
clavata, Spreng. l. c. = Rapistrum clavatum.
crenata, Jord. Diagn. i. 346 = maritima.
cubensis, H. B. & K. Nov. Gen. et Sp. v. 75 = maritima.
cyrenaica, *Spreng. Syst.* ii. 852.—Cyrenia.
domingensis, Tuss. Fl. Ant. i. 119 = maritima.
edentula, Hook. Fl. Bor. Am. i. 59 = maritima.
hispanica, L'Hérit. ex DC. Syst. ii. 619 = Brassica incana.
hispanica, Jord. Diagn. i. 345 = maritima.
latifolia, Poir. Encyc. Suppl. ii. 88 = maritima.
littoralis, Jord. Diagn. i. 345 = maritima.
maritima, *Scop. Fl. Carn.* ed. II. ii. 35.—Europ.; Afr. bor.; Am. bor.
monosperma, *Lange, Pl. Nov. Hisp.* fasc. i. 5. t. 7.—Hispan.
myagroides, Poir. Encyc. Suppl. ii. 88 = Erucaria aleppica.
perennis, L'Hérit. ex DC. Syst. ii. 431 = Rapistrum perenne.
perfoliata, L'Hérit. ex DC. l. c. 575 = Conringia orientalis.
pinnatifida, Stokes, Bot. Mat. Med. iii. 484 = maritima.
rugosa, L'Hérit. ex DC. Syst. ii. 432 = Rapistrum rugosum.
Serapionis, Gaertn. Fruct. ii. 287 = maritima.
sinuatifolia, Stokes, Bot. Mat. Med. iii. 485 = maritima.
striata, L'Hérit. ex DC. Syst. ii. 659 = Didesmus tennifolius.

CALACANTHA, T. Anders. ex Benth. & Hook. f. Gen. ii. 1088 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.
Dalzellianus, *T. Anders. ex Benth. & Hook. f. l. c.*—Ind. or.
grandiflorus, Radlk. in Sitzb. Math.-Phys. Acad. Muench. xiii. '1884' (1883) 279 = praec.

CALACINUM, Rafin. Fl. Tellur. ii. 33 (1836) = **Muehlenbeckia**, Meissn. (Polygon.).
adpressum, Rafin. l. c. 34 = M. adpressa.
australis, Rafin. l. c. = M. australis.

CALADENIA, R. Br. Prod. 323 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 612.
LEPTOCERAS, Lindl. Gen. et Sp. Orch. 415 (1840).
alata, R. Br. l. c. 324 = carnea.
alba, R. Br. l. c. 323 = carnea.
angustata, Hook. f. Fl. Tasm. ii. 30. t. 125 = carnea.
angustata, Lindl. Gen. et Sp. Orch. 420 = testacea.
antarctica, Reichb. f. Beitr. Syst. Pfl. 67 = Lyperanthus antarcticus.
aphylla, *Benth. Fl. Austral.* vi. 387.—Austral.
arenaria, *Fitzg. Austral. Orch.* i. VII. t. 1.—Austral.
Barbarossa, *Reichb. f. Beitr. Syst. Pfl.* 64.—Austral.
barbata, Lindl. Gen. et Sp. Orch. 418 = deformis.
Behrii, Schlecht. in Linnaea, xx. (1847) 569 = Patersoni.
bifolia, *Hook. f. Fl. Nov. Zel.* i. 247.—N. Zel.
Brunonis, Reichb. Beitr. Syst. Pfl. 67 = Glossodia Brunonis.
caerulea, *R. Br. Prod.* 324.—Austral.
Cairnsiana, *F. Muell. Fragm.* vii. 31.—Austral.
carnea, *R. Br. Prod.* 324.—Austral.

CALADENIA :—
clavigera, *A. Cunn. ex Lindl. Gen. et Sp. Orch.* 422.—Austral.
concolor, *Fitzg. Austral. Orch.* i. VII. t. 1.—Austral.
congesta, *R. Br. Prod.* 324.—Austral.
cornuta, Reichb. f. Beitr. Syst. Pfl. 67 = Chiloglottis cornuta.
cucullata, *Fitzg. Austral. Orch.* i. II. t. 4.—Austral.
cuneata, Reichb. f. Beitr. Syst. Pfl. 67 = Burnettia cuneata.
deformis, *R. Br. Prod.* 324.—Austral.
denticulata, Lindl. Swan Riv. App. 52 = filamentosa.
dilatata, R. Br. Prod. 325 = Patersoni.
dimorpha, *Fitzg. Austral. Orch.* i. I. t. 3.—Austral.
diphylla, Reichb. f. Beitr. Syst. Pfl. 67 = Chiloglottis diphylla.
discoidea, *Lindl. Swan Riv. App.* 52.—Austral.
Drummondii, *Benth. Fl. Austral.* vi. 383.—Austral.
elliptica, Reichb. f. Beitr. Syst. Pfl. 67 = Lyperanthus ellipticus.
elongata, Lindl. Swan Riv. App. 52 = latifolia.
emarginata, Reichb. f. Beitr. Syst. Pfl. 67 = Glossodia emarginata.
filamentosa, Lindl. Gen. et Sp. Orch. 421 = Patersoni.
filamentosa, *R. Br. Prod.* 324.—Austral.
filifera, Lindl. Swan Riv. App. 52 = filamentosa.
fimbriata, *Reichb. f. Beitr. Syst. Pfl.* 65.—Austral.
flava, *R. Br. Prod.* 324.—Austral.
gemmata, *Lindl. Swan Riv. App.* 52.—Austral.
Gigas, *Reichb. f. in Linnaea*, xli. (1877) 56.—N. Caled.
gracilis, R. Br. Prod. 324 = testacea.
Gunnii, Reichb. f. Beitr. Syst. Pfl. 67 = Chiloglottis Gunnii.
hirta, *Lindl. Swan Riv. App.* 52.—Austral.
ixioides, *Lindl. l. c.*—Austral.
javanica, *Benn. ex Ridl. in H. O. Forbes, Nat. Wand. East. Archip.* 518.—Timor.
latifolia, *R. Br. Prod.* 324.—Austral.
leptochila, *Fitzg. in Gard. Chron.* (1882) I. 462.—Austral.
lobata, *Fitzg. l. c.* 461.—Austral.
longicauda, Lindl. Swan Riv. App. 52. t. 8 = Patersoni.
Lyallii, *Hook. f. Fl. Nov. Zel.* i. 247.—N. Zel.
macrophylla, R. Br. Prod. 325 = Menziesii.
macrostylis, *Fitzg. in Gard. Chron.* (1882) I. 462.—Austral.
major, Reichb. f. Beitr. Syst. Pfl. 67 = Glossodia major.
marginata, Lindl. Swan Riv. App. 51 = latifolia.
Menziesii, *R. Br. Prod.* 325.—Austral.
minor, *Hook. f. Fl. Nov. Zel.* i. 247.—N. Zel.
minor, Reichb. f. Beitr. Syst. Pfl. 67 = Glossodia minor.
mollis, Endl. in Lehm. Pl. Preiss. ii. 8 = hirta.
mollis, Lindl. Swan Riv. App. 51 = latifolia.
multiclavia, *Reichb. f. Beitr. Syst. Pfl.* 64.—Austral.
nana, Endl. in Lehm. Pl. Preiss. ii. 7 = reptans.
nigricans, Reichb. f. Beitr. Syst. Pfl. 67 = Lyperanthus nigricans.
Nortoni, *F. Muell. Census*, 113.—Austral.
oblonga, Reichb. f. Beitr. Syst. Pfl. 67 = Cyrtostylis oblonga.
ochreata, Lindl. Swan Riv. App. 51 = latifolia.
pallida, Lindl. Gen. et Sp. Orch. 421 = Patersoni.
paniculata, *Fitzg. in Gard. Chron.* (1882) I. 461.—Austral.
Patersoni, *R. Br. Prod.* 324.—Austral.
pellita, Endl. in Lehm. Pl. Preiss. ii. 8 = gemmata.
plicata, *Fitzg. in Gard. Chron.* (1882) I. 461.—Austral.
Preissii, Endl. in Lehm. Pl. Preiss. ii. 7 = reptans.
pulcherrima, F. Muell. Fragm. v. 93 = Patersoni.
reniformis, Reichb. f. Beitr. Syst. Pfl. 67 = Cyrtostylis reniformis.
reptans, *Lindl. Swan Riv. App.* 52.—Austral.
reticulata, *Fitzg. in Gard. Chron.* (1882) I. 462.—Austral.
Roei, *Benth. Fl. Austral.* vi. 383.—Austral.
rotundifolia, Reichb. f. Beitr. Syst. Pfl. 67 = Cyrtostylis rotundifolia.
saccharata, *Reichb. f. l. c.* 63.—Austral.
sericea, *Lindl. Swan Riv. App.* 52.—Austral.
serrata, *Reichb. f. Beitr. Syst. Pfl.* 67.—Austral.
suaveolens, *Reichb. f. l. c.*—Austral.
sulphurea, A. Cunn. in Field, N. S. Wales, 361 = suaveolens.

CALADENIA :—
tentaculata, Schlecht. in Linnaea, xx. (1847) 571 = Patersoni.
tenuis, *Fitzg. in Gard. Chron.* (1882) I. 462.—Austral.
tessellata, *Fitzg. Austral. Orch.* i. II. t. 3.—Austral.
testacea, *R. Br. Prod.* 324.—Austral.
unguiculata, Lindl. Swan Riv. App. 51 = deformis.
unita, *Fitzg. in Gard. Chron.* (1882) I. 461.—Austral.
variegata, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 248.—N. Zel.

CALADIUM, Vent. in Roem. Arch. ii. (1800) 347. *AROIDEAE*, Benth. & Hook. f. iii. 976.
AROSMA, Rafin. Fl. Tellur. iii. 66 (1836).
CYRTOSPADIX, C. Koch, Ind. Sem. Hort. Berol. (1853).
acre, R. Br. Prod. 336 = Colocasia antiquorum.
aculeatum, *G. F. W. Mey. Prim. Fl. Esseq.* 274.—Guiana.
acuminatum, F. G. Dietr. Vollst. Lexik. Gaertn. i. 616 = Xanthosoma belophyllum.
albo-punctatissimum, J. Makoy, ex Karst. in C. Koch & Fint. Wochenschr. (1858) 331 = bicolor?
amabile, *Versch. Cat.* n. 70 (1862) 2.—Bras.
amoenum, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
appendiculatum, Hort. ex Engl. l. c. 476 = Xanthosoma atrovirens.
Appunianum, Hort. ex Engl. l. c. 466 = picturatum.
arborescens, *Vent. Jard. Cels*, sub t. 30; *Willd. Sp. Pl.* iv. 490.—Ind. or.
arboreum, *H. B. & K. Nov. Gen. et Sp.* i. 80.—N. Granat.
argyrites, Lem. in Illustr. Hortic. (1858) t. 185. f. 3. Misc. 57 = Humboldtii.
argyrospilum, Lem. l. c. 59 = bicolor.
argyrostigma, *Schott, Melet.* i. 18, nomen.—Am. trop.
asperulum, Schott, in Oestr. Bot. Wochenbl. (1853) 313 = smaragdinum.
auritum, Vent. Jard. Cels, sub t. 30; Willd. Sp. Pl. iv. 491 = Syngonium auritum.
Baraquini, Hérincq, in Fl. des Serres, Sér. II. iii. (1858) 163 = bicolor.
Barilletii, *Hort.—Cf. Gard. Chron.* (1867) 461.—Hab.?
Bauersii, *Reichb. Hort. Bot.* t. 160.—Bras.
Belleyenei, Hort. Chantin, ex Lem. in Illustr. Hortic. (1860) t. 252 = picturatum.
belophyllum, Willd. Enum. Hort. Berol. 987 = Xanthosoma belophyllum.
bicolor, *Vent. Jard. Cels*, t. 30.—Am. austr.
Bongniartii, C. Koch & Fint. Wochenschr. (1858) 287, sphalm. = bicolor.
Bowei, Steud. Nom. ed. II. i. 249 = Biarum Bovei.
brasiliense, Desf. Cat. Hort. Par. ed. III. 7, 386 = Xanthosoma hastifolium.
Brongniartii, Lem. in Illustr. Hortic. (1858) Misc. 58 = bicolor.
caracasanum, Engl. in DC. Monog. Phan. ii. 666, Index = Xanthosoma belophyllum.
Chantinii, Lem. in Illustr. Hortic. (1858) Misc. 58 = bicolor.
Colocasia, Schott, ex Wight, Ic. t. 787 = Alocasia cucullata.
colocasioides, Brongn. in Nouv. Ann. Mus. Par. iii. 156 = Colocasia antiquorum.
concolor, *C. Koch, in Berl. Allg. Gartenz.* (1857) 135.—Venezuela.
Connaertii, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
cordatum, Hort. Versch. ex Engl. l. c. 455 = Schomburgkii.
crassipes, Hort. ex Engl. l. c. 369 = Philodendron cannaefolium.
cucullatum, Pers. Syn. ii. 575 = Alocasia cucullata.
cupreum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1854) 6 = Alocasia cuprea.
Curwadlii, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
Devosianum, Lem. in Illustr. Hortic. (1862) t. 322. f. 1 = bicolor.
digitatum, *Blanco, Fl. Filip.* ed. II. 459.—Ins. Philipp.
discolor, Hort. ex Engl. in DC. Monog. Phan. ii. 462 = bicolor.
Duchartrei, Hort. ex Engl. l. c. = bicolor.

CALADIUM:—
Eckhartii, Lem. ex Engl. in DC. Monog. Phan. ii. 463 = bicolor.
edule, G. F. W. Mey. Prim. Fl. Esseq. 272 = Xanthosoma edule.
Engelii, *Karst. in C. Koch & Fint. Wochenschr.* (1858) 330.—Venezuela.
Enkeanum, C. Koch, Wochenschr. (1861) 267 = bicolor.
erythropus, Mart. ex Engl. in Mart. Fl. Bras. iii. II. 172 = Adelonema erythropus.
esculentum, Vent. Jard. Cels, sub t. 30; et in Roem. Arch. ii. III. 351 = Colocasia antiquorum.
firmulum, Schott, in Oestr. Bot. Wochenbl. (1854) 417 = bicolor.
fragrantissimum, Hook. Bot. Mag. t. 3314 = Philodendron fragrantissimum.
Gaerdtii, C. Koch & Fint. Wochenschr. (1858) 288 = bicolor.
giganteum, Blume, Cat. Gew. Buitenz. 103 = Colocasia indica.
glycyrrhizum, Fraser, in Hook. Bot. Misc. i. (1830) 259 = Alocasia macrorrhiza.
grandifolium, Sims, Bot. Mag. t. 2643 = Philodendron Simsii.
grandifolium, Willd. Sp. Pl. iv. 490 = Philodendron grandifolium.
griseo-argenteum, Hort. ex Engl. in DC. Monog. Phan. ii. 463 = bicolor.
guttiferum, Poepp. ex Kunth, Enum. Pl. iii. 52 = Philodendron guttiferum.
Haageanum, C. Koch, Wochenschr. (1862) 136 = bicolor.
haematostigma, Kunth, Enum. Pl. iii. 42 = bicolor.
hastatum, Lem. in Illustr. Hortic. (1858) Misc. 61 = picturatum.
hastatum, Willd. ex Engl. in DC. Monog. Phan. ii. 279 = Urospatha Poeppigiana.
hastifolium, Steud. Nom. ed. II. i. 249 = Xanthosoma hastifolium.
helleborifolium, Vent. Jard. Cels, sub t. 30; et in Roem. Arch. ii. III. 351 = Xanthosoma helleborifolium.
Hendersoni, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
heterophyllum, *Presl*, *Reliq. Haenk.* ii. 148.—Ins. Luzon.
Houbyanum, Hort. ex Engl. in DC. Monog. Phan. ii. 464 = bicolor.
Houlletii, Lem. in Illustr. Hortic. (1858) 59 = bicolor.
Humboldtii, *Schott, in Oestr. Bot. Wochenbl.* (1854) 417.—Bras.
Imbe, Hort. ex Kunth, Enum. Pl. iii. 50 = Philodendron Imbe.
indica, Hort. ex C. Koch, Berl. Allg. Gartenz. (1857) 136 = Alocasia indica.
Ketteleri, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
Kochii, Hort. Versch. ex C. Koch, Wochenschr. v. (1862) 135 = bicolor.
Kramerianum, Hort. ex Engl. in DC. Monog. Phan. ii. 459 = bicolor.
lacerum, Willd. Sp. Pl. iv. 491 = Philodendron lacerum.
Laucheanum, C. Koch, Wochenschr. (1861) 266 = bicolor.
Lemaireanum, Barraq. in Illustr. Hortic. (1862) t. 311 = picturatum.
Leopoldii, Hort. Versch. ex Engl. in DC. Monog. Phan. ii. 460 = bicolor.
Lindeni, Hort. ex Engl. l. c. 462 = bicolor.
liniferum, *Nees, in Neuwied, Reise nach Brasil.* ii. 338.—Bras.
lividum, Lodd. Bot. Cab. t. 1590 = Staurostigma concinnum.
Lowii, Lem. in Illustr. Hortic. (1863) t. 360 = Alocasia Lowii.
luridum, Kunth, Enum. Pl. iii. 50 = Staurostigma concinnum.
macrophyllum, Lem. in Illustr. Hortic. (1862) t. 316, 322 = bicolor.
macropus, Hort. Petrop. ex Kunth, Enum. Pl. iii. 47 = Philodendron cannaefolium.
macrorrhizon, R. Br. Prod. 336 = Alocasia macrorrhiza.
macrotites, *Schott, in Oestr. Bot. Wochenbl.* (1855) 20.—Bras.
maculatum, Lodd. Bot. Cab. t. 608 = Dieffenbachia Seguine.
Mafaffa, Hort. ex Engl. in DC. Monog. Phan. ii. 471 = Xanthosoma Mafaffa.
marantaefolium, Zipp. ex Kunth, Enum. Pl. iii. 55 = Aglaonema marantifolium.
marginatum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1854) 6 = bicolor.
marmoratum, *L. Math. ex C. Koch, l. c.*—Ecuador.
Martersteigianum, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
metallicum, Hamburg. Gart. Blumenzeit. (1853) 517 = Alocasia cuprea.
metallicum, Hort. ex Engl. l. c. 502 = Alocasia indica.
mirabile, Lem. Illustr. Hortic. (1863) t. 354 = bicolor.
Mooreanum, Hort. ex Engl. in DC. Monog. Phan. ii. 462 = bicolor.
myriostigma, C. Koch, Wochenschr. v. (1862) 135 = Humboldtii.
Neumanni, Lem. in Illustr. Hortic. (1858) 58 = bicolor.
nymphaeaefolium, Vent. Jard. Cels, sub t. 30; Willd. Sp. Pl. iv. 488 = Colocasia antiquorum.
odoratissimum, Hort. ex C. Koch, Berl. Allg. Gartenz. (1857) 20 = Alocasia odora.
odoratum, Lodd. Bot. Cab. t. 416 = Alocasia macrorrhiza.
odorum, Lindl. Bot. Reg. t. 641 = Alocasia macrorrhiza.
Ottonis, Hort. ex Engl. in DC. Monog. Phan. ii. 464 = bicolor.
ovatum, Vent. Jard. Cels, sub t. 30; et in Roem. Arch. ii. III. 351 = Lagenandra toxicaria.
pallidinervium, Hort. ex Engl. in DC. Monog. Phan. ii. 463 = bicolor.
pallidum, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1853) 13 = smaragdinum.
pedatum, Hook. Exot. Fl. t. 206 = Philodendron laciniatum.
pellucidum, DC. in Pl. Jard. Genèv. (1826) 38 = bicolor.
pendulinum, *Manso, Enum.* 21.—Bras.
pendulum, *Blume, Cat. Gew. Buitenz.* 103.—Malaya.
Perrieri, Lem. in Illustr. Hortic. (1858) 59 = bicolor.
petiolatum, Hook. Bot. Mag. t. 3428 = Anchomanes difformis.
pictum, DC. 4me Note, Pl. Jard. Genèv. 38 = bicolor.
pictum, Lodd. Bot. Cab. t. 608 = Dieffenbachia picta.
picturatum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1854) 6.—Am. austr.
pinnatifidum, Willd. Sp. Pl. iv. 487 = Philodendron pinnatifidum.
platynervium, Hort. ex Kunth, Enum. Pl. iii. 47 = Philodendron crassinervium.
platynevron, *Salm-Dyck, Hort. Dyck.* (1834) 334.—Bras.
plumbeum, C. Koch, Berl. Allg. Gartenz. (1857) 136 = Alocasia cuprea.
poecile, Schott, Melet. i. 18 = bicolor.
porphyroneuron, C. Koch, Wochenschr. v. (1862) 135 = picturatum.
Princeps, Hort. ex Kunth, Enum. Pl. iii. 55 = Aglaonema simplex.
pumilum, D. Don, Prod. Fl. Nep. 21 = Gonatanthus sarmentosus.
punctatissimum, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
punctatum, *Desf. Tabl. Hort. Par.* ed. II. 6, nomen.—Hab.?
Purdieanum, Schott, in Oestr. Bot. Zeitschr. (1859) 38 = bicolor.
pusillum, C. Koch, in Fl. des Serres, Sér. II. iii. (1858) 110 = bicolor.
regale, Lem. in Illustr. Hortic. (1862) t. 316 = bicolor.
Reichenbachianum, Stange, ex Engl. in Mart. Fl. Bras. iii. II. 182 = bicolor.
reticulatum, *Desf. Tabl. Hort. Par.* ed. II. 6, nomen.—Hab.?
Rougieri, [*Versch.*].—*Cf. Gard. Chron.* (1864) 414.—Bras.
rubellum, C. Koch & Fint. Wochenschr. (1858) 288 = bicolor.
rubricaule, Hort. ex Lem. in Fl. des Serres, Sér. II. iii. (1858) 109 = bicolor.
rubrovenium, Hort. ex Engl. in DC. Monog. Phan. ii. 460 = bicolor.
rugosum, Desf. Cat. Hort. Par. ed. III. 386 = Alocasia cucullata.
sagittaefolium, Sieber, ex Engl. in DC. Monog. Phan. ii. 464 = bicolor.
sagittaefolium, Vent. Jard. Cels, sub t. 30; et in Roem. Arch. ii. III. 351 = Xanthosoma sagittifolium.
sarmentosum, Hort. ex Kunth, Enum. Pl. iii. 36 = Gonatanthus sarmentosus.
scandens, Willd. Sp. Pl. iv. 489 = Culcasia scandens.
Schmitzii, Lem. Illustr. Hortic. (1861) t. 297. f. 3 = Schomburgkii.
Schoelleri, Lem. l. c. f. 2 = Schomburgkii.
Schomburgkii, *Schott, in Oestr. Bot. Wochenschr* (1858) 122.—Guiana.
seguinum, Vent. Jard. Cels, sub t. 30; Willd. Sp. Pl. iv. 490 = Dieffenbachia Seguine.
Sieboldii, Hort. ex Engl. in DC. Monog. Phan. ii. 461 = bicolor.
simplex, Blume, Cat. Gew. Buitenz. 103 = Aglaonema simplex.
Simsii, Hook. Bot. Mag. t. 3345 = Philodendron Simsii.
smaragdinum, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1853) 13.—Am. trop.
sororium, *Schott, in Oestr. Bot. Zeitschr.* ix. (1859) 38.—Bras.
splendens, C. Koch & Fint. Wochenschr. (1858) 288 = bicolor.
splendidum, *Versch. Cat.* n. 70 (1862) 4.—Bras.
Spruceanum, Schott, in Oestr. Bot. Zeitschr. ix. (1859) 38 = bicolor.
Stangeanum, C. Koch, Wochenschr. (1861) 136 = bicolor.
steudneriaefolium, *Engl. Bot. Jahrb.* vi. (1885) 284.—N. Granat.
striatipes, *Schott, Syn. Aroid.* 51.—Bras.
subrotundum, Lem. in Illustr. Hortic. (1858) Misc. 60 = Schomburgkii.
surinamense, Miq. Hort. Amstelod. (1853) = bicolor.
Thelemanni, *Versch. Cat.* n. 72 (1863) 2.—Hab.?
thripedestum, Lem. in Illustr. Hortic. v. (1858) Misc. 59 = marmoratum.
trifoliatum, Desf. Cat. Hort. Par. ed. III. 386 = seq.
tripartitum, Willd. Sp. Pl. iv. 491 = Philodendron tripartitum.
Troubetzkoyi, Hérincq, in Fl. des Serres, Sér. II. iii. (1858) 164 = picturatum.
undulatum, Steud. Nom. ed. II. ii. 219 = Peltandra undulata.
utile, Hort. ex Engl. in DC. Monog. Phan. ii. 469 = Xanthosoma sagittifolium.
variegatum, Desf. Cat. Hort. Par. ed. III. 7, 385 = Xanthosoma helleborifolium.
Veitchii, Henders. ex Hook. f. Bot. Mag. t. 5376 = Alocasia Lowii.
Veitchii, Lindl. in Gard. Chron. (1859) 740 = Alocasia cuprea.
Vellozianum, Schott, in Bonplandia, vii. (1859) 163 = bicolor.
Verschaffeltii, Lem. in Illustr. Hortic. (1858) 59. t. 185 = bicolor.
violaceum, Desf. Cat. Hort. Par. ed. III. 7 = seq.?
violaceum, Hort. ex Engl. in DC. Monog. Phan. ii. 492 = Colocasia antiquorum.
virginicum, Hook. Exot. Fl. t. 182 = Peltandra undulata.
viviparum, Nees, in Schott, Melet. i. 18; Lodd. Bot. Cab. t. 281 = Remusatia vivipara.
Wagneri, Hort. ex Engl. in DC. Monog. Phan. ii. 464 = bicolor.
Wightii, Lem. Illustr. Hortic. (1861) 294 = bicolor.
xanthorrhizon, Boos, Cat. 348 = Xanthosoma Jacquini.
xanthorrhizum, Willd. Sp. Pl. iv. 490 = Xanthosoma sagittifolium.
zamiaefolium, Lodd. Bot. Cab. t. 1408 = Zamioculcas Loddigesii.

CALAENA, Schlecht. in Linnæa, ii. (1827) 528, sphalm. = **Caleana**, R. Br. (Orchid.).

CALAIS, DC. Prod. vii. 85 (1839) = **Microseris**, D. Don (Compos.).
aphantocarpha, A. Gray, in Proc. Am. Acad. vi. (1864–65) 552 = M. aphantocarpha.
Bigelovii, A. Gray, Pacif. Rail. Rep. iv. 113 = M. Bigelovii.
Bolanderi, A. Gray, in Proc. Am. Acad. vii. (1868) 365 = M. Bolanderi.
cyclocarpha, A. Gray, in Pacif. Rail. Rep. iv. 113 = M. Douglasii.

CALAIS :—
Douglasii, DC. Prod. vii. 85 = Microseris Douglasii.
eriocarpha, A. Gray, in Proc. Am. Acad. vi. (1864–65) 552 = M. Douglasii.
glauca, A. Gray, l. c. vii. (1868) 364 = M. procera.
gracililoba, Kellogg, in Proc. Calif. Acad. v. (1873) 48 = M. major.
laciniata, A. Gray, in Pacif. Rail. Rep. iv. 113 = M. laciniata.
Lindleyi, DC. Prod. vii. 85 = M. Lindleyi.
linearifolia, DC. l. c. = M. linearifolia.
macrochaeta, A. Gray, Pl. Fendl. 112 = M. macrochaeta.
major, A. Gray, in Pacif. Rail. Rep. iv. 113 = M. major.
nutans, A. Gray, l. c. = M. nutans.
Parryi, A. Gray, l. c. 112 = M. Parryi.
platycarpha, A. Gray, l. c. 113 = M. platycarpha.
sylvatica, A. Gray, l. c. 114 = M. sylvatica.
tenella, A. Gray, l. c. = M. aphantocarpha.

CALAMAGROSTIS, Adans. Fam. ii. 31 (1763). *GRAMINEAE*, Benth. & Hook. f. iii. 1150.
AMAGRIS, Rafin. Princ. Somiol. 27 (1814); et in Am. Monthly Mag. (1819) 190.
ATHERNOTUS, Dulac, Fl. Hautes-Pyr. 74 (1867).
LECHLERA, Steud. Syn. Pl. Gram. 101, in nota (1855).
PTEROPODIUM, Willd. ex Steud. Nom. ed. II. ii. 414 (1841).
abietina, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 155 = Deyeuxia acutiflora.
acrathera, *Peterm. in Flora*, xxvii. (1844) 225.—Saxonia.
acutiflora, DC. Fl. Fr. v. 255 = Deyeuxia acutiflora.
aemula, Steud. Nom. ed. II. i. 249; Syn. Pl. Gram. 192 = Deyeuxia Forsteri.
agapatea, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
agrostoides, *Boiss. Fl. Orient.* v. 522.—Reg. Caucas.
agrostoides, Spreng. Syst. i. 252 = Deyeuxia canadensis.
airoides, Steud. Nom. ed. II. i. 249 = Arundo airoides.
alba, *Steud. l. c.*—Peruv.
albicans, Buckl. in Proc. Acad. Sc. Philad. '1862' (1863) 92 = Deyeuxia aleutica.
aleutica, Bong. Veg. Ins. Sitcha, 171 = Deyeuxia aleutica.
alpina, Host, Gram. iv. t. 51 = tenella.
altissima, Host, Fl. Austr. i. 101 = Arundo mauritanica.
ambigua, *Hartm. Skand. Fl.* ed. V.—Scandin.
andina, *Nutt. in Journ. Acad. Sc. Philad.* N. S. i. (1847) 187.—Calif.
anomala, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
anthoxanthoides, *Regel, in Act. Hort. Petrop.* vii. (1880) 640.—Turkestan.
Antoniana, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
arenaria, Roth, Tent. Fl. Germ. i. 34 = Ammophila arundinacea.
argentea, DC. Fl. Fr. iii. 25 = Stipa Calamagrostis.
arundinacea, Roth, Tent. Fl. Germ. i. 33 = Deyeuxia sylvatica.
arundinacea, Wibel, Prim. Fl. Werth. 122 = Deyeuxia acutiflora.
Arundo, J. F. Gmel. Syst. 171 = Stipa Calamagrostis.
Arundo, Roth, Tent. Fl. Germ. i. 33 = Aira caespitosa.
australis, *Buese, in Miq. Pl. Jungh.* 343.—Malaya.
azorica, Steud. Syn. Pl. Gram. 189 = Agrostis pallida.
Balansae, *Boiss. Fl. Orient.* v. 522.—Pontus.
baltica, Trin. Gram. Unifl. 227; Hartm. Skand. Fl. ed. II. 20 = Ammophila baltica.
Beyrichiana, *Nees, ex Doell, in Mart. Fl. Bras.* ii. III. 53.—Bras.
bicolor, J. F. Gmel. Syst. i. 172 = Ampelodesma bicolor.
Billardieri, Stend. Nom. ed. II. i. 249 = Deyeuxia Billardieri.
Bolanderi, Thurb. in S. Wats. Bot. Calif. ii. 280 = Deyeuxia Bolanderi.
borealis, Laest. Bidr. Vaextl. Torn. Lappm. Ups. 44 = Deyeuxia neglecta.
brachytricha, *Steud. Syn. Pl. Gram.* 189.—Japon.
brevifolia, Steud. Nom. ed. II. i. 249 = Deyeuxia brevifolia.
brevipilis, Beck, Bot. N. & Mid. St. 401 = Ammophila brevipilis.

CALAMAGROSTIS:—
Breweri, Thurb. ex Boland. in Trans. Calif. Agr. Soc. (1864–65) 136 = Deyeuxia Breweri.
bromoides, Nutt. Gen. Am. i. 44 = Stipa Aristella.
caespitosa, Steud. Syn. Pl. Gram. 189 = Agrostis pallida.
canadensis, Beauv. Agrost. 157, nomen = Deyeuxia canadensis.
canadensis, Nutt. Gen. Am. i. 46 = Deyeuxia Nuttalliana.
canescens, Beauv. ex Steud. Nom. ed. II. i. 249, 250 = Epigeios.
capillaris, *Nees, ex Steud. Nom.* ed. II. i. 219; *Syn. Pl. Gram.* 188.—Bras.
caucasica, Trin. in C. A. Mey. Verz. Pfl. Cauc. 15 = Deyeuxia acutiflora.
ehalybaea, *Fries, Summa Pl.* 241.—Norveg.
Chamissonis, Steud. Nom. ed. II. i. 250 = Deyeuxia Chamissonis.
chilensis, *Phil. in Linnaea*, xxix. (1857–58) 89.—Chili.
chrysantha, Steud. Nom. ed. II. i. 250 = Deyeuxia chrysantha.
ciliata, *Steud. Syn. Pl. Gram.* 193.—Nepal.
cinnoides, Spreng. Syst. i. 252 = Deyeuxia Nuttalliana.
Clarionis, Loisel. Fl. Gall. 719 = Deyeuxia Halleriana.
coarctata, Steud. Nom. ed. II. i. 250 = Deyeuxia coarctata.
coarctata, Torr. Fl. N. York, ii. 444 = Deyeuxia Nuttalliana.
colorata, Nutt. Gen. Am. i. 46 = Phalaris intermedia.
colorata, Sibth. Fl. Oxon. 37 = Phalaris arundinacea.
columbiensis, Nutt. ex A. Gray, in Proc. Acad. Sc. Philad. '1862' (1863) 334 = Deyeuxia Langsdorfii.
compressa, *Doell, in Mart. Fl. Bras.* ii. III. 56.—Bras.
confinis, *Nutt. Gen. Am.* i. 47.—Am. bor.
conspicua, Bergeret, Fl. Pyr. i. 60 = Stipa Calamagrostis.
conspicua, J. F. Gmel. Syst. 172 = Arundo conspicua.
crassiglumis, Thurb. ex S. Wats. in Bot. Calif. ii. 281 = Deyeuxia crassiglumis.
contracta, Dum. Obs. Gram. Belg. 126 = stricta?
cuspidata, *Spreng. Syst.* i. 253.—Mexic.
densiflora, C. Muell. in Walp. Ann. vi. 986 = phragmitoides.
densiflora, Steud. Nom. ed. II. i. 250 = Deyeuxia densiflora.
deschampsioides, Trin. Ic. Gram. iii. t. 354 = Deyeuxia deschampsioides.
dioica, *Steud. Syn. Pl. Gram.* 193.—Cochinch.
donaciformis, Loisel. Fl. Gall. i. 53 = Arundo Pliniana.
dubia, *Bunge, Lehm. Rel.* (1847) 348.—As. centr.
effusa, Steud. Nom. ed. II. i. 250 = Deyeuxia effusa.
elata, Blytt, Norges Fl. i. 92 = phragmitoides.
eminens, *Steud. Nom.* ed. II. i. 250 = Deyeuxia eminens.
emodensis, *Griseb. in Goett. Nachr.* (1868) 80.—Reg. Himal.
epigeios, Huds. Fl. Angl. 43; ed. II. 54 = lanceolata.
epigeios, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1842) 529 = laxa.
epigeios, *Roth, Tent. Fl. Germ.* i. 34.—Europ.; As. temp.
epigeios, Steud. Nom. ed. II. i. 250, sphalm. = Saccharum cylindricum.
eriantha, Steud. Syn. Pl. Gram. 190 = Deyeuxia eriantha.
Falklandiae, *Steud. l. c.* 192.—Ins. Falkland.
farcta, *J. F. Gmel. Syst.* 172.—Guiana.
filiformis, *Griseb. in Goett. Nachr.* (1868) 79.—Ind. or.
flexuosa, Rupr. Beitr. Pfl. Russl. iv. 34 = phragmitoides.
Forsteri, Steud. Nom. ed. II. i. 250 = Deyeuxia Forsteri.
Funckii, *Steud. Syn. Pl. Gram.* 192.—Venezuela.
fuscata, Steud. Nom. ed. II. i. 250 = Deyeuxia fuscata.
Gaudiniana, Reichb. Fl. Germ. Exc. 27 = lanceolata.
georgica, C. Koch, in Linnaea, xxi. (1848) 387 = epigeios.
glauca, Hohen. Enum. Talusch. 14 = laxa.
glauca, Reichb. Fl. Germ. Excurs. 27 = epigeios.
glauca, Steud. Nom. ed. II. i. 250 = Deyeuxia Nuttalliana.
glauca, Steud. Syn. Pl. Gram. 187 = littorea.
glaucescens, C. Muell. in Walp. Ann. vi. 986 = phragmitoides.
glomerata, Boiss. & Buhse, in Mém. Soc. Nat. Mosc. xii. (1860) 229 = epigeios.

CALAMAGROSTIS:—
graciliseens, *Blytt, Norges Fl.* i. 88.—Norveg.
gracilis, Seenus, Reise Istr. 73 = Stipa Aristella.
gracilis, Schumach. ex Nym. Consp. 800 = lanceolata.
groenlandica, *Kunth, Rev. Gram.* i. 75.—Groenl.; Labrad.
hakonensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 599.—Japon.
Halleriana, DC. Fl. Fr. Suppl. 256 = Deyeuxia Halleriana.
Halleriana, Fries, Summa Veg. 240 = gracilescens.
Hartmaniana, Fries, l. c. 241 = ambigua.
hirtigluma, *Steud. Syn. Pl. Gram.* 188.—Am. bor.
holciformis, *Jaub. & Spach, Illustr.* iv. 61.—Tibet.
Hookeriana, *Steud. Syn. Pl. Gram.* 192.—Zeylan.
Howelli, Vasey, in Coult. Bot. Gaz. vi. (1881) 271 = Deyeuxia Howellii.
Huebneriana, Reichb. Fl. Germ. Excurs. 27 = epigeios.
Humboldtiana, *Steud. Nom.* ed. II. i. 250.—N. Granat.
hyperborea, *Lange, in Fl. Dan.* t. 2942.—Groenl.
inexpansa, A. Gray, ex Torr. Fl. N. York, ii. 445. t. 152 = Deyeuxia confinis.
intermedia, Lindeb. ex Nym. Consp. 799 = Hartmaniana.
intermedia, Steud. Nom. ed. II. i. 250 = Deyeuxia intermedia.
Jamesoni, *Steud. Syn. Pl. Gram.* 191.—Ecu dor.
javana, *Steud. l. c.* 193.—Java.
junciformis, Steud. Nom. ed. II. i. 250 = Deyeuxia junciformis.
Karka, J. F. Gmel. Syst. 172 = Arundo Karka.
laguroides, *Regel, in Act. Hort. Petrop.* vii. (1880) 641.—Turkestan.
Lagurus, Koel. Desc. Gram. 112 = Imperata arundinacea.
lanceolata, *Roth, Tent. Fl. Germ.* i. 34.—Europ.; As. temp.
lanceolata, Stokes, in With. Arr. Brit. Pl. ed. III. ii. 122 = epigeios.
lanceolata, Trin. ex Steud. Nom. ed. II. i. 250 = purpurea.
Langsdorffii, Auct. ex Nym. Consp. 799 = phragmitoides.
Langsdorffii, Trin. Gram. Unifl. 225. t. 4. f. 10 = Deyeuxia Langsdorffii.
lapponica, Hartm. Scand. Fl. ed. I. 46; Trin. Gram. Unifl. 226 = Deyeuxia lapponica.
lapponica, Hook. Brit. Fl. ed. IV. 34 = Deyeuxia neglecta.
latifolia, Blytt, Norges Fl. i. 90 = phragmitoides.
laxa, *Hegetschw. Fl. Schw.* 65.—Helvet.
laxa, Host, Gram. iv. 25 = littorea.
laxior, C. Muell. in Walp. Ann. vi. 1115 = Deyenxia Hartmanniana.
Leersii, Koel. Desc. Gram. 107 = Deschampsia juncea.
Leersii, Trin. ex Steud. Nom. ed. II. i. 850, ii. 643 = Stipa Calamagrostis.
lenkoranensis, *Steud. Syn. Pl. Gram.* 188.—Reg. Caucas.
leucantha, *Delile, ex Steud. Nom.* ed. II. i. 250 = Stipa Calamagrostis.
lithuanica, Bess. in Roem. & Schult. Add. ad Mant. ii. 602 = phragmitoides.
littorea, *DC. Fl. Fr.* vi. 255.—Europ.; As. temp.
longearistata, *Fisch. ex Ledeb. Fl. Ross.* iv. 433, nomen. —Hab.?
longifolia, Hook. Fl. Bor. Am. ii. 241 = Ammophila longifolia.
macloviana, *Steud. Syn. Pl. Gram.* 192.—Ins. Falkl.
maritima, *Kunth, Rev. Gram.* i. 75.—Scania.
mauritanica, Spreng. Syst. i. 252 = Arundo mauritanica.
megapotamica, Steud. Syn. Pl. Gram. 192 = Andropogon laguroides.
mexicana, Kunth, Syn. iv. 473 = Muehlenbergia calamagrostidea.
mexicana, Nutt. Gen. Am. i. 46 (excl. syn.) = Deyeuxia canadensis.
Michauxii, Trin. ex Steud. Nom. ed. II. i. 250 = Deyeuxia canadensis.
montana, *DC. Fl. Fr.* vi. 254.—Germ.; Austr.; Gall.
montana, Host, Gram. t. 146 = Deyeuxia sylvatica.
montevidensis, *Nees, in Mart. Fl. Bras.* ii. 401.—Reg. Argent.
Munroana, *Boiss. Fl. Orient.* v. 523.—Afghan.
Munroi, *Boiss. l. c.* 526.—Afghan.
mutica, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.

CALAMAGROSTIS:—
Necsii, *Steud. Nom.* ed. II. i. 250; *Syn. Pl. Gram.* 193. —Ind. or.
neglecta, *Gaertn. Mey. & Scherb. Fl. Wett.* i. 94 = Deyeuxia neglecta.
nepalensis, *Nees, ex Steud. Syn. Pl. Gram.* 193.—Nepal.
nigricans, Mérat, Fl. Paris, ed. II. ii. 33 = Phragmites communis.
nipponica, *Franch. & Sav. Enum. Pl. Jap.* ii. 599.—Japon.
nutans, Saut. ex Reichb. Fl. Germ. Exc. 26 = Deyeuxia Halleriana.
nutkaënsis, Steud. Syn. Pl. Gram. 190 = Deyeuxia nutkaënsis.
Nuttalliana, Steud. Nom. ed. II. i. 251 = Deyeuxia Nuttalliana.
obscura, *Downar, in Bull. Soc. Nat. Mosc.* xxxv. (1862) 1. 605.—Podol.
obtusata, *Trin. Gram. Unifl.* 225. t. 4. f. 11.—Sibir.
obtusata, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104; xxix. (1856) 1. 26 = Deyeuxia deschampsioides.
odessana, *Bess. ex Andrz. Rys Botan.* ii. (1830) 26.—Tauria.
olympica, *Boiss. Diagn.* Ser. I. v. 70.—Reg. Caucas.; Persia.
Onaei, *Franch. & Sav. Enum. Pl. Jap.* ii. 598.—Japon.
oregonensis, Buckl. in Proc. Acad. Sc. Philad. '1862' (1863) 92 = Deyeuxia Langsdorffii.
Orizabae, Steud. Syn. Pl. Gram. 193 = Deyeuxia Orizabae.
ovata, Steud. Nom. ed. II. i. 251 = Deyeuxia ovata.
pallens, Steud. l. c. = Deyeuxia pallens.
pallida, C. Muell. in Walp. Ann. vi. 986 = phragmitoides.
pallidiflora, C. Muell. l. c. = phragmitoides.
parviflora, Rupr. in Beitr. Pfl. Russ. Reich. iv. 36 = Deyeuxia sylvatica.
patula, *Steud. Syn. Pl. Gram.* 422.—Chili.
persica, Boiss. Diagn. Ser. I. vii. 120 = littorea.
phalaroides, *Regel, in Act. Hort. Petrop.* vii. (1880) 641, in textu.—Turkestan.
phleoides, *Steud. Nom.* ed. II. i. 251.—Am. austr.?
phragmitoides, *Hartm. Scand. Fl.* ed. IV. 25.—Reg. bor.
Pickeringii, *A. Gray, Man. Bot. U. St.* ed. I. 547 = Deyeuxia Pickeringii.
piscatoria, *Steud. Syn. Pl. Gram.* 193.—Cochinch.
planifolia, Trin. ex Steud. Nom. ed. II. i. 251 = Deyeuxia planifolia.
Plinii, J. F. Gmel. Syst. 172 = Arundo mauritanica.
plumosa, Spreng. Syst. i. 253 = Stipa papposa.
poacoides, *Steud. Syn. Pl. Gram.* 423.—Reg. Magell.
poaeoides, Trin. ex Steud. Nom. ed. II. i. 251 = Poa subuniflora.
Porteri, A. Gray, in Proc. Am. Acad. vi. (1862) 79 = Deyeuxia Porteri.
Pseudophragmites, Aschers. Fl. Brandenb. 821 = littorea.
Pseudophragmites, Blytt, Norges Fl. i. 89 = phragmitoides.
Pseudophragmites, Link, ex Reichb. Fl. Germ. Exc. 26 = Deyeuxia Halleriana.
Pseudophragmitis, Kocl. Desc. Gram. 106 = littorea.
pulchella, Blytt, Norges Fl. i. 93 = phragmitoides.
pulchella, *Griseb. in Goett. Nachr.* (1868) 78.—Ind. or.
pulchella, Saut. ex Reichb. Fl. Germ. Excurs. 26 = Deyeuxia Halleriana.
pumila, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 176. —Ins. Galap.
purpurascens, R. Br. in Richards. App. Frankl. Journ. 3 = Deyeuxia sylvatica.
purpurea, *Trin. Gram. Unifl.* 219.—Sibir. baical.
pyramidalis, Host, Gram. iv. t. 49 = Deyeuxia sylvatica.
quadriseta, Spreng. Syst. i. 253 = Deyeuxia quadriseta.
quitensis, H. B. & K. Nov. Gen. et Sp. i. 133 = Muehlenbergia calamagrostidea.
ramosa, Host, Gram. iv. t. 44 = lanceolata.
rariflora, C. Muell. in Walp. Ann. vi. 986 = Deyeuxia Halleriana.
rariflora, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 156 = Deyeuxia sylvatica.
recta, Trin. ex Steud. Nom. ed. II. i. 251 = Deyeuxia recta.
retrofracta, Link, ex Steud. l. c. = Deyeuxia Forsteri.
rigens, Fries, Summa Veg. 240 = epigeios.

CALAMAGROSTIS:—
rigida, Trin. ex Steud. Nom. ed. II. i. 251 = Deyeuxia rigida.
robusta, C. Muell. in Walp. Ann. vi. 985 = Deyeuxia Halleriana.
robusta, *Franch. & Sav. Enum. Pl. Jap.* ii. 600.—Japon.
Roylcana, *Steud. Nom.* ed. II. i. 251.—Ind. or.
Roylei, Steud. Syn. Pl. Gram. 193 = pracc.
rubella, Boiss. Fl. Orient. v. 525 = littorea.
rubescens, Buckl. in Proc. Acad. Sc. Philad. '1862' (1863) 92 = Deyeuxia rubescens.
rubicunda, Blytt, Norges Fl. i. 91 = phragmitoides.
rubra, Bosc, ex Kunth, Enum. Pl. 478 = Erianthus saccharoides.
rudis, Steud. Syn. Pl. Gram. 192 = Deyeuxia scabra.
rupestris, *Trin. ex Steud. Nom.* ed. II. i. 251, *nomen; Doell. in Mart. Fl. Bras.* ii. III. 53.—Bras.
sachalinensis, *F. Schmidt, Reisen Amurl.* 202.—Terr. Amur.
scabra, *Presl, Rel. Haenk.* i. 234.—Sinus Nootka.
scabrescens, *Griseb. in Goett. Nachr.* (1868) 79.—Ind. or.
Schiedeana, Steud. Syn. Pl. Gram. 193 = Deyeuxia Schiedeana.
Schimperiana, *Hochst. in Flora*, xxxviii. (1855) 202. —Afr. trop.
Schleicheri, Bess. Prim. Fl. Galic. i. 69 = lanceolata.
Schwabii, Spreng. Syst. i. 253 = Gastridium australe.
sciuroides, *Franch. & Sav. Enum. Pl. Jap.* ii. 600.—Japon.
sedenensis, Loisel. Fl. Gall. ed. II. i. 54 = varia.
sesquitriflora, Steud. Syn. Pl. Gram. 190 = Deyeuxia tolucensis.
simplex, *Boiss. Fl. Orient.* v. 523.—Reg. Caucas.
sitchensis, *Trin. ex Ledeb. Fl. Ross.* iv. 433.—Ins. Sitcha.
speciosa, Host, Gram. iv. t. 45 = Stipa Calamagrostis.
spicigera, Steud. Nom. ed. II. i. 251 = Deyeuxia spicigera.
splendens, *Steud. l. c.*—Ins. Catharin.
stricta, Koel. Desc. Gram. 105 = Deyeuxia neglecta?
stricta, *Trin. Gram. Unifl.* 226.—Europ.; Am. bor.
strigosa, Hartm. Skand. Fl. ed. I. 46 = Deyeuxia strigosa.
subalpina, Schur, Enum. Pl. Transs. 740 = Deyeuxia sylvatica.
subspicata, Desv. Opusc. 60 = Trisetum subspicatum?
subulata, Dum. Agr. Belg. t. 9. f. 36 = montana.
subulata, Steud. Nom. ed. II. i. 251 = varia.
sylvatica, *Bess. Prim. Fl. Galic.* i. 70.—Europ.
sylvatica, DC. Fl. Fr. v. 253 = Deyeuxia arundinacea.
sylvatica, Host, Gram. iv. t. 48 = varia.
sylvatico-stricta, Kuehlew. ex Ledeb. Fl. Ross. iv. 429 = Deyeuxia neglecta.
tenella, *Host, Gram.* iv. t. 50.—Eur. med.; As. bor.
tenuifolia, H. B. & K. Nov. Gen. et Sp. i. 134 = Muehlenbergia tenuifolia.
thrysoidea, C. Koch, in Linnaea, xxi. (1848) 388 = epigeios.
tianschanica, *Rupr. Sert. Tiansch.* 34.—Turkestan.
toluccensis, Trin. ex Steud. Nom. ed. II. i. 250 = Deyeuxia tolucensis.
transsylvanica, Schur, in Oestr. Bot. Zeitschr. ix. (1859) 154 = laxa.
Trinii, Rupr. in Beitr. Pfl. Russ. Reich. iv. 36 = Deyeuxia acutiflora.
varia, *Beauv. Agrost.* 157.—Europ.; As. bor.
varia, Boland. ex Bot. Calif. ii. 280 = Deyeuxia Bolanderi.
varia, Gren. & Godr. Fl. Fr. iii. 477 = montana.
varia, Host, Gram. iv. t. 47 = Deyeuxia Halleriana.
variegata, With. Bot. Arr. Brit. Pl. ed. III. ii. 124 = Phalaris arundinacea.
variflora, C. Muell. in Walp. Ann. vi. 1115 = Deyeuxia Halleriana.
velutina, *Steud. Nom.* ed. II. i. 251.—Chili.
villosa, J. F. Gmel. Syst. 172 = Deyeuxia sylvatica.
villosa, Mutel, Fl. Fr. iv. 41 = Deyeuxia Halleriana.
vilnensis, Bess. ex Roem. & Schult. Add. ad Mant. ii. 602 = phragmitoides.
vincalis, *Baumg. ex Steud. Syn. Pl. Gram.* 188.—Transsylv.
viridescens, Steud. l. c. 192 = seq.
viridiflavescens, Steud. Nom. ed. II. i. 251 = Deyeuxia viridiflavescens.
Willdenowii, Steud. Syn. Pl. Gram. i. 192 = Deyeuxia Forsteri.

CALAMINA, Beauv. Agrost. 128. t. 23. f. 1. (1812) = **Apluda**, Linn. (Gramin.).
gigantea, Beauv. l. c. 129 = Apluda mutica.
gigantea, Roem. & Schult. Syst. ii. 810 = Anthistiria gigantea.
humilis, Presl, Rel. Haenk. i. 344 = Apluda mutica.
imberbis, Beauv. Agrost. 129 = Anthistiria imberbis.
mutica, Beauv. l. c. = Apluda mutica.
Sehima, Beauv. l. c. 157 = Ischaemum laxum.
Themeda, Beauv. l. c. = Anthistiria imberbis.

CALAMINTHA, Adans. Fam. ii. 192 (1763) = **Nepeta**, Linn. (Labiat.).

CALAMINTHA, [Tourn.] Lam. Fl. Fr. ii. 393 (1778). *LABIATAE*, Benth. & Hook. f. ii. 1190.
Acinos, Moench, Meth. 407 (1794).
Clinopodium, Tourn. ex Linn. Gen. ed. I. 170 (1737), nomen prius.
Faucibarba, Dulac, Fl. Hautes-Pyr. 402 (1867).
abyssinica, *A. Rich. Tent. Fl. Abyss.* ii. 191.—Abyssin.
acinoides, Nym. Consp. 589 = suaveolens.
Acinos, [*Clairv.*] *Man. Herb.* 197 (*err. typ.* 917).—Europ.; Reg. Caucas.
acuminata, Frivald. in Flora, xviii. (1835) 332 = suaveolens.
adscendens, Willk. & Lange, Prod. Fl. Hispan. ii. 413 = officinalis.
aegyptiaca, Lam. ex Schnr, Enum. Pl. Transs. 531 = C. Clinopodium.
aetnensis, Strobl, in Oestr. Bot. Zeitschr. xxiv. (1874) 29 = alpina.
alba, Lam. ex Steud. Nom. ed. II. i. 251 = Melissa alpina.
alba, Reichb. Fl. Germ. Exc. 328 = Micromeria rupestris.
alpina, *Lam. Fl. Fr.* ii. 394.—Europ. austr.; Reg. Mediterr.
annua, *Schrenck, in Bull. Sc. Acad. Pétersb.* x. (1842) 353.—Rossia.
arundana, *Benth. in DC. Prod.* xii. 233.—Hispan.
arvensis, Lam. Fl. Fr. ii. 394 = C. Acinos.
ascendens, Jord. Obs. Pl. Crit. iv. (1846) 8 = officinalis.
athonica, Reichb. Fl. Germ. Exc. 329 = C. Nepeta.
atlantica, Ball, in Journ. Bot. xiii. (1875) 175 = C. Clinopodium.
atlantica, Coss. in Bull. Soc. Bot. Fr. ix. (1862) 176 = grandiflora.
baetica, Boiss. & Reut. Pugill. Pl. Nov. (1852) 92 = officinalis.
betulifolia, *Boiss. & Bal. Diagn.* Ser. II. iv. 14.—Cilicia.
Biebersteinii, C. Koch, ex Ledeb. Fl. Ross. iii. 355 = umbrosa.
byzantina, C. Koch, in Linnaea, xxi. (1848) 672 = C. Nepeta.
caerulescens, *Wedd. Chlor. And.* ii. 149. t. 62.—N. Granat.
cana, Auct. ex Heynh. Nom. ii. 104 = graveolens.
candidissima, *Munby, ex Benth. in DC. Prod.* xii. 226. —Algeria.
canescens, Presl, Fl. Sicul. 37 = C. Nepeta.
canescens, Torr. & Gray, ex Benth. in DC. Prod. xii. 229 = Conradina canescens.
caroliniana, *Sweet, Hort. Brit.* 409.—Am. bor.
chinensis, *Benth. in DC. Prod.* xii. 233.—China.
Clinopodium, *Benth. l. c.*—Reg. bor. temp.
coccinea, *Benth. l. c.* 229.—Am. bor.
confinis, *Hance, in Journ. Bot.* vi. (1868) 331.—China.
corsica, *Benth. in DC. Prod.* xii. 231.—Ins. Corsica.
cretica, *Benth. l. c.* 227.—Ins. Creta.
cretica, Lam. Fl. Fr. ii. 395 = Micromeria marifolia.
croatica, Host, Fl. Austr. ii. 132 = Micromeria Piperella.
cryptantha, *Vatke, in Linnaea*, xxxvii. (1871-73) 328. —Abyss.
debilis, *Benth. in DC. Prod.* xii. 232.—Sibiria.
dentata, *Chapm. Fl. S. U. St.* 318.—Florida.
depauperata, *Bunge, Lab. Pers.* 38.—Persia.
dilatata, Schrad. Ind. Sem. Hort. Goetting. (1835) = C. Nepeta.
Einseleana, *F. Schultz, Arch. Fl. Fr. et Allem.* 178.—Europ.
exigua, Nym. Consp. 589 = graveolens.
Fenzlii, Vis. Revis. Pl. Min. Cogn. (1855) 7 = subnuda.
florida, *Boiss. Diagn.* Ser. I. xii. 59.—Cilicia.
foliosa, *Opiz, Natural.* iii. (1823) [20].—Europ.

CALAMINTHA:—
fruticosa, Steud. Nom. ed. II. i. 251 = Micromeria marifolia.
glabella, *Benth. in DC. Prod.* xii. 230.—Am. bor.
glabella, A. Gray, Man. ed. II. 307 = Nuttallii.
glabrescens, Boiss. Fl. Orient. iv. 580 = staminea.
glandulosa, *Benth. Lab. Gen. et Sp.* 387.—Corsica.
glomerulosa, Boiss. & Heldr. ex Benth. in DC. Prod. xii. 234 = origanifolia.
gracilis, *Benth. l. c.* 232.—Java.
granatensis, Boiss. & Reut. Pugill. Pl. Nov. 94 = alpina.
grandiflora, *Moench, Meth.* 408.—Europ.
grandiflora, Pursh, Fl. Am. Sept. ii. 414 = caroliniana.
graveolens, *Benth. in DC. Prod.* xii. 231.—Reg. Mediterr.; As. Min.
graveolens, Coss. ex Willk. & Lange, Prod. Fl. Hispan. ii. 415 = umbrosa.
Gussonei, Tod. ex Nym. Consp. 588 = C. Nepeta.
Haussknechti, *Bunge, Lab. Pers.* 36.—Persia.
hederacea, Scop. Fl. Carn. ed. II. i. 423 = Nepeta Glechoma.
Herba-barona, Heynh. Nom. i. 144 = Thymus Herba-barona.
heterophylla, Heynh. l. c. = C. Acinos.
heterotricha, *Boiss. & Reut. Pugill. Pl. Nov.* 93.—Hispan.
hispidula, *Boiss. & Reut. l. c.*—Hispan.
ilicifolia, A. Gray, in Proc. Am. Acad. viii. (1873) 368 = Acanthomentha ilicifolia.
incana, *Boiss. ex Benth. in DC. Prod.* xii. 226.—Graecia et Ins.; Syria.
Jahniana, *Simk. in Termész. Füzetek*, ix. (1885) 25.—Europ.
Langei, Nym. Consp. 589 = suaveolens.
longicaulis, *Benth. in DC. Prod.* xii. 234.—Reg. Himal.
macrostema, *Benth. l. c.* 229.—Mexic.
marifolia, Steud. Nom. ed. II. i. 251 = Micromeria marifolia.
maritima, *Benth. in DC. Prod.* xii. 231.—Thracia.
menthaefolia, Host, Fl. Austr. ii. 129 = officinalis.
meridionalis, Nym. Consp. 589 = alpina.
mimuloides, *Benth. Pl. Hartw.* 331.—Calif.
mollis, Jord. ex Lamotte, Prod. ii. 599 = C. Nepeta.
moluccana, *Miq. Fl. Ind. Bat.* ii. 968.—Ins. Molucc.; Timor.
montana, Lam. Fl. Fr. ii. 396 = officinalis.
multicaulis, *Maxim. in Bull. Acad. Pétersb.* xx. (1875) 466.—Japon.
nebrodensis, Kern. & Strobl, in Oestr. Bot. Zeitschr. xxiv. (1874) 171 = alpina.
neglecta, *Opiz, Natural.* iii. (1823) [20].—Europ.
nepalensis, Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 53 = umbrosa.
Nepeta, *Savi, Fl. Pis.* (1798) 197.—Europ.; Afr. bor.; Oriens.
Nepeta, Willk. in Flora, xxxv. (1852) 277 = officinalis.
nepetoides, Jord. Obs. Pl. Crit. iv. (1846) 16 = C. Nepeta.
nervosa, *Pomel, Nouv. Mat. Fl. Atl.* 307.—Afr. bor.
nivea, *Boiss. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 253.—As. Min.
Nuttallii, *Benth. in DC. Prod.* xii. 230.—Am. bor.
obliqua, Host, Fl. Austr. ii. 131 = C. Nepeta.
officinalis, *Moench, Meth.* 409.—Europ.; Oriens.
origanifolia, *Boiss. Diagn.* Ser. I. xii. 50.—As. Min.; Syria.
origanifolia, Host, Fl. Austr. ii. 130 = Micromeria Pulegium.
origanifolia, Vis. Fl. Dalm. ii. 199 = Micromeria origanifolia.
Palmeri, *A. Gray, in Proc. Am. Acad.* xi. (1876) 100.—Calif.
pamphylica, *Boiss. & Heldr. Diagn.* Ser. I. xii. 52.—Pamphylia.
paradoxa, *Vatke, in Linnaea*, xxxvii. (1871–73) 327.—Abyssin.
parviflora, Lam. Fl. Fr. ii. 396 = C. Nepeta.
patavina, Heldr. ex Nym. Consp. 588 = suaveolens.
patavina, *Host, Fl. Austr.* ii. 133.—Oriens; Europ. austr. or.
Piperella, Reichb. Fl. Germ. Excurs. 328 = Thymus Piperella.
Piperella, Schloss. & Vukot. Syll. Fl. Croat. 80 = Micromeria croatica.
piperelloides, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 95.—Lycia.
Pulegium, Reichb. f. Ic. Fl. Germ. xviii. 45. f. 1278 = Micromeria Pulegium.
purpurascens, *Benth. in DC. Prod.* xii. 231.—Hispan.

CALAMINTHA:—
repens, Benth. l. c. 233 = umbrosa.
rotundifolia, Benth. l. c. 232 = umbrosa.
rotundifolia, Ces. Pass. & Gib. Comp. Fl. Ital. 303 = alpina.
rotundifolia, Host, Fl. Austr. ii. 131 = C. Nepeta.
rotundifolia, Losc. & Pard. Ser. Inconf. 82 = Zizyphora acinoides.
rotundifolia, Reichb. Ic. Fl. Germ. xviii. 43 = patavina.
rupestris, Host, Fl. Austr. ii. 131 = Micromeria rupestris.
simensis, *Benth. in DC. Prod.* xii. 230.—Abyssin.
spicigera, C. Koch, in Linnaea, xxi. (1848) 671 = Micromeria serpyllifolia.
Spruneri, Boiss. Diagn. Ser. I. xii. 53 = C. Nepeta.
staminea, *Boiss. l. c.* 51.—Kurdistan.
stenostoma, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 95.—Lycia.
stricta, Reichb. Ic. Fl. Germ. xviii. 44 = officinalis.
suaveolens, *Boiss. Diagn.* Ser. I. xii. 54.—Graecia.
subnuda, Host, Fl. Austr. ii. 130 = Micromeria Pulegium.
subnuda, Reichb. Fl. Germ. Excurs. 329 = thymifolia.
sylvatica, Bromf. in Engl. Bot. Suppl. t. 2897 = officinalis.
thymifolia, *Host, Fl. Austr.* ii. 132.—Carniola.
thymifolia, Reichb. Fl. Germ. Excurs. 328 = Micromeria rupestris.
trichotoma, Moench, Meth. 409 = C. Nepeta.
umbrosa, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vi. 6; ix. *Suppl.* 8.—Oriens; Ind. or.
umbrosa, Reichb. Fl. Germ. Excurs. 329 = officinalis.
ussuriensis, *Regel & Maack, in Regel, Tent. Fl. Ussur.* 116.—Reg. Amur.
vulgaris, [Clairv.] Man. Herb. 197 (err. typ. 917) = C. Nepeta.

CALAMOCHLOA, Fourn. in Bull. Soc. Bot. Fr. xxiv. (1877) 178. *GRAMINEAE*, Benth. & Hook. f. iii. 1178.
filifolia, *Fourn. l. c.*—Mexic.

CALAMOCHLOË, Reichb. Consp. 52 (1828) = **Arundinella**, Raddi (Gramin.).

CALAMOSAGUS, Griff. in Calc. Journ. Nat. Hist. v. (1845) 22 = **Korthalsia**, Blume (Palm.).
harinaefolius, Griff. Palms Brit. E. Ind. 29. t. 184 = K. wallichiaefolia.
laciniosus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 23 = K. laciniosa,
ochriger, Griff. Palms Brit. E. Ind. 31. t. 216 = K. polystachys.
polystachys, H. Wendl. in Kerch. Palm. 235, sphalm. = K. polystachys.
scaphiger, Griff. Palms Brit. E. Ind. 30. t. 148 a = K. scaphigera.
wallichiaefolius, Griff. in Calc. Journ. Nat. Hist. v. (1845) 24 = K. wallichiaefolia.

CALAMPELIS, D. Don, in Edinb. N. Phil. Journ. vii. (1829) 89 = **Eccremocarpus**, Ruiz & Pav. (Bignoniac.).
scaber, D. Don, l. c. = E. scaber.

CALAMUS, Auct. ex Linn. Gen. ed. I. 104 (1737) = **Acorus**, Linn. (Aroid.).
aromaticus, Gueldenst. ex Ledeb. Fl. Ross. iv. 13 = Acorus Calamus.

CALAMUS, Linn. Sp. Pl. 325 (1753). *PALMAE*, Benth. & Hook. f. iii. 931.
ROTANG, Adans. Fam. ii. 24 (1763).
ROTANGA, Boerh. ex Crantz, Inst. i. 127 (1766).
acanthopis, Griff. Palms Brit. E. Ind. t. 216 = Daemonorops malaccensis.
acanthospathus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 39.—Reg. Himal.
accedens, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 28 = Daemonorops Draco.
adspersus, Blume, Rumphia, iii. 40 = Daemonorops adspersus.
albus, *Pers. Syn.* i. 383.—Amboina.
amarus, Lour. Fl. Cochinch. 210 = tenuis?
amboinensis, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 20 = Daemonorops calapparius.
americanus, Hort. ex Steud. Nom. ed. II. i. 252 = Desmoncus americanus.

CALAMUS:—
amplectens, *Becc. Malesia*, ii. (1884) 78.—Borneo.
anceps, Blume, Rumphia, iii. 65. t. 149 = melanoloma.
andamanicus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 211.—Ins. Andaman.
angustifolius, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 89 = Daemonorops angustifolius.
arborescens, *Griff. l. c.* 33.—Burma.
asperrimus, *Blume, in Schult. f. Syst.* vii. 1327.—Java.
assamicus, *Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Assam.
aureus, Reinw. ex Mart. Hist. Nat. Palm. iii. 341 = ornatus.
australis, *Mart. l. c.* 342.—Austral.
barbatus, Macklot, in Bijdr. Nat. Wetensch. v. (1830) 178, nomen; Blume, Rumphia, iii. 42 = Daemonorops barbatus.
Blancoi, *Kunth, Enum. Pl.* iii. 595.—Ins. Philipp.
borneensis, Miq. Anal. Bot. Ind. 4 = javensis.
brevifrons, Mart. Hist. Nat. Palm. iii. 338 = Blancoi.
buroënsis, *Mart. l. c.* 336.—Ins. Buro.
caesius, *Blume, Rumphia*, iii. 57.—Malaya.
calapparius, Mart. Hist. Nat. Palm. ed. prior, 209 (iii. 331) = Daemonorops calapparius.
calicarpus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 92 = Daemonorops calicarpus.
calolepis, Miq. Pl. Jungh. 159 = melanoloma.
caryotoides, *A. Cunn. ex Mart. Hist. Nat. Palm.* iii. 338.—Austral.
castaneus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 28.—Malaya.
Cawa, *Blume, Rumphia*, iii. 31.—Amboina.
ciliaris, *Blume, in Schult. f. Syst.* vii. 1330.—Malaya.
cochleatus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 29 = Daemonorops cochleatus.
collinus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 31.—Ind. or.
concinnus, *Mart. Hist. Nat. Palm.* iii. 332. t. 116.—Burma.
crinitus, Miq. Anal. Bot. Ind. 6 = Daemonorops crinitus.
cuspidatus, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 434. t. 41 = Eremospatha cuspidata.
dealbatus, Hort. ex H. Wendl. in Kerch. Palm. 236 = Acanthophoenix rubra.
deerratus, *Mann & H. Wendl. in Trans. Linn. Soc.* xxiv. (1864) 429.—Afr. trop.
delicatulus, *Thw. Enum. Pl. Zeyl.* 330.—Zeylan.
depressiusculus, *Teijsm. & Binn. Cat. Hort. Bog.* 74.—Sumatra.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 594.—Malaya.
dioicus, *Lour. Fl. Cochinch.* 211.—Cochinch.
discolor, Mart. Hist. Nat. Palm. iii. 341 = siphonospathus?
Draco, Willd. Sp. Pl. ii. 203 = Daemonorops Draco.
elegans, *Hort. ex H. Wendl. in Kerch. Palm.* 236.—Hab. ?
elongatus, Miq. Anal. Bot. Ind. 6 = Daemonorops elongatus.
epetiolaris, *Mart. Hist. Nat. Palm.* iii. 336.—Java.
equestris, Blume, in Schult. f. Syst. vii. 1330 = javensis.
equestris, *Willd. Sp. Pl.* ii. 204.—Ins. Molucc.
erectus, *Roxb. Hort. Beng.* 72; *Fl. Ind.* iii. 774.—Reg. Himal.
exilis, *Griff. ex Mart. Hist. Nat. Palm.* iii. 333.—Malaya.
extensus, Mart. Hist. Nat. Palm. iii. 200, partim, t. 116. f. 4. = viminalis.
extensus, Roxb. Hort. Beng. 73; Fl. Ind. iii. 777 = palustris?
farinosus, *Linden, in Illustr. Hortic.* xix. (1872) t. 109.—Sumatra.
fasciculatus, Roxb. Hort. Beng. 72; Fl. Ind. iii. 779 = viminalis.
Fernandezii, H. Wendl. in Kerch. Palm. 236 = Oxleyanus.
fissus, Miq. Anal. Bot. Ind. 6 = Daemonorops fissus.
Flagellum, Griff. ex Mart. Hist. Nat. Palm. iii. 333 = polygamus.
floribundus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 56.—Reg. Himal.
Gaudichaudii, H. Wendl. in Kerch. Palm. 236 = Daemonorops Gaudichaudii.
geminiflorus, Griff. ex Mart. Hist. Nat. Palm. iii. 338 = Plectocomia geminiflora.
geniculatus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 67 = Daemonorops geniculatus.

CALAMUS:—
glaucescens, Blume, Rumphia, iii. 65 = caesius.
glaucescens, D. Dietr. Syn. Pl. ii. 1064 = Ceratolobus glaucescens.
gracilipes, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 28 = Daemonorops longipes, *Miq.*
gracilis, Blanco, Fl. Filip. ed. I. 267 = Blancoi.
gracilis, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 781.—Reg. Himal.
gracilis, Thwaites, Enum. Pl. Zeyl. 330 = pachystemonus.
graminosus, *Blume, Rumphia*, iii. 31.—Amboina.
grandiflorus, Steud. Nom. ed. II. i. 252, sphalm. = Ancistrophyllum secundiflorum.
grandis, Griff. in Calc. Journ. Nat. Hist. v. (1845) 84 = Daemonorops grandis.
Griffithianus, *Mart. Hist. Nat. Palm.* iii. 332.—Malaya.
Guruba, *Buch.-Ham. ex Kunth, Enum. Pl.* iii. 210.—Reg. Himal.; Burma.
Haenkeanus, Mart. Hist. Nat. Palm. iii. 337 = mollis.
Helferianus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 213.—Burma.
Heliotropium, Buch.-Ham. ex Kunth, Enum. Pl. iii. 210 = tenuis.
heteracanthus, *Macklot, in Bijdr. Nat. Wetensch.* v. (1830) 178.—N. Guinea.
heteroideus, Blume, Rumphia, iii. 46 = Reinwardtii.
hirsutus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 28 = Daemonorops Hystrix.
Hookeri, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 434 = Eremospatha Hookeri.
horrens, *Blume, Rumphia*, iii. 43.—Java.
hostilis, Wall. ex Voigt, Hort. Suburb. Calc. 639; Miq. Fl. Ind. Bat. iii. 113, nomen = arborescens.
Huegelianus, *Mart. Hist. Nat. Palm.* iii. 338.—Malabar.
humilis, Roxb. Hort. Beng. 72; Fl. Ind. iii. 773 = latifolius?
hygrophilus, Griff. Palms Brit. Ind. t. 213 C = Daemonorops hygrophilus.
hypoleucus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 208.—Burma.
Hystrix, Griff. in Calc. Journ. Nat. Hist. v. (1845) 70 = Daemonorops Hystrix.
inermis, T. Anders. in Journ. Linn. Soc. xi. (1871) 11 = latifolius.
insignis, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 59.—Malaya.
intermedius, Griff. l. c. 86 = Daemonorops intermedius.
javensis, *Blume, Rumphia*, iii. 62.—Malaya.
Jenkinsianus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 81 = Daemonorops Jenkinsianus.
Jenkinsianus, Griff. Palms Brit. Ind. 40. t. 186 A. f. 3 (non p. 89) = polygamus.
Korthalsii, Miq. Anal. Bot. Ind. i. 6 = Daemonorops Korthalsii.
laevigatus, *Mart. Hist. Nat. Palm.* iii. 339.—Malaya.
laevis, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 430. t. 38 (Ancistrophylli sp.).—Afr. trop.
latifolius, Kurz, in Journ. As. Soc. Beng. xliii. (1874) II. 20. t. 31. A. = palustris.
latifolius, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 775.—Burma.
latispinus, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1868) n. v. 29.—Sumatra.
leptopus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 73 = Daemonorops leptopus.
leptospadix, *Griff. l. c.* 49.—Reg. Himal.
Lewisianus, Griff. l. c. 87 = Daemonorops Lewisianus.
Lindeni, *Rodigas, in Illustr. Hortic.* xxx. (1883) t. 499.—Malaya.
litoralis, Blume, Rumphia, iii. 43 = viminalis.
longipes, Griff. in Calc. Journ. Nat. Hist. v. (1845) 68 = Daemonorops longipes, *Mart.*
longisetus, *Griff. l. c.* 36.—Burma.
macracanthus, T. Anders. in Journ. Linn. Soc. xi. (1871) 10 = latifolius.
macrocarpus, Griff. ex Mart. Hist. Nat. Palm. iii. 333 = erectus.
macrocarpus, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 435 = Eremospatha macrocarpa.
macropterus, Miq. in Verh. Kon. Acad. Wetensch. xi. (1868) n. v. 19.—Ins. Celebes.
Manan, *Miq. Fl. Ind. Bat. Suppl.* 595.—Sumatra.

CALAMUS:—
manicatus, Teijsm. & Binn. ex Miq. Fl. Ind. Bat. iii. 135 = Daemonorops crinitus.
manillensis, *H. Wendl. in Kerch. Palm.* 237.—Ins. Philipp.
Mannii, H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 436 = Oncocalamus Mannii.
Margaritae, *Hance, in Journ. Bot.* xii. (1874) 266.—China.
marginatus, *Mart. Hist. Nat. Palm.* iii. 342.—Borneo.
maritimus, *Blume, Rumphia*, iii. 31, in adnot.—Java.
Mastersianus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 76 = C. Guruba.
maximus, Blanco, Fl. Filip. ed. I. 266 = manillensis?
maximus, Reinw. ex Schult. f. Syst. vii. 1333 = Plectocomia elongata.
melanacanthus, *Mart. Hist. Nat. Palm.* iii. 333. t. 116. f. 13.—Burma.
melanochaetes, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 28 = Daemonorops melanochaetes.
melanolepis, H. Wendl. in Kerch. Palm. 237 = Huegelianus.
melanoloma, *Mart. Hist. Nat. Palm.* iii. 209, 339. t. 116. f. 3.—Java.
Metzianus, *Schlecht. in Linnaea*, xxvi. (1853-55) 727.—Ind. or.
Meyenianus, *Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 425.—Ins. Philipp.
micracanthus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 62.—Malacca.
micranthus, *Blume, Rumphia*, iii. 53.—Malaya.
mirabilis, Mart. ex Kunth, Enum. Pl. iii. 213 = Daemonorops mirabilis.
mishmeensis, Griff. in Calc. Journ. Nat. Hist. v. (1845) 55 = floribundus.
mollis, *Blanco, Fl. Filip.* ed. I. 264.—Ins. Philipp.
monoecus, Roxb. Hort. Beng. 73; Fl. Ind. iii. 783 = C. Rotang.
montanus, T. Anders. in Journ. Linn. Soc. xi. (1871) 9 = acanthospathus.
monticolus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 90 = Daemonorops monticolus.
Muelleri, *H. Wendl. & Drude, in Linnaea*, xxxix. (1875) 193.—Austral.
niger, Willd. Sp. Pl. ii. 203 = Daemonorops niger.
nitidus, *Mart. Hist. Nat. Palm.* iii. 334.—Burma; Ind. or.
nivalis, Thw. ex Trimen, in Journ. Bot. xxiii. (1885) 268, sphalm. = rivalis.
nutantiflorus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 79 = Daemonorops Jenkinsianus.
oblongus, *Reinw. in Schult. f. Syst.* vii. 1323.—Java.
obstruens, F. Muell. Fragm. v. 48 = australis.
ochreatus, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1868) n. v. 29.—Sumatra.
opacus, *Blume, Rumphia*, iii. 50.—Sumatra.
opacus, Mann & H. Wendl. in Trans. Linn. Soc. xxiv. (1864) 431. t. 41 = Laccosperma opaca.
ornatus, *Blume, ex Schult. f. Syst.* vii. 1326.—Malaya.
ovatus, Reinw. ex Kunth, Enum. Pl. iii. 205 = ornatus.
ovoideus, *Thw. ex Trimen, in Journ. Bot.* xxiii. (1885) 269.—Zeylan.
Oxleyanus, *Teijsm. & Binn. Cat. Hort. Bog.* 75, nomen.—Malaya.
pachystemonus, *Thw. Enum. Pl. Zeyl.* 431.—Zeylan.
palembanicus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 29 = Daemonorops palembanicus.
pallens, Blume, Rumphia, iii. 51 = Reinwardtii.
palustris, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 60.—Burma.
paniculatus, *Roxb. Hort. Beng.* [105], nomen.—Penang.
paradoxus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 213. t. 29, 30.—Burma.
penicillatus, Roxb. Fl. Ind. iii. 781 = javensis?
periacanthus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 28 = Daemonorops periacanthus.
petiolaris, Griff. in Calc. Journ. Nat. Hist. v. (1845) 93 = Daemonorops petiolaris.
petraeus, *Lour. Fl. Cochinch.* 209.—Cochinch.
philippensis, *Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Ins. Philipp.

CALAMUS:—
pisicarpus, *Blume, Rumphia*, iii. 39.—Ins. Molucc.
platyacanthos, Mart. ex Kunth, Enum. Pl. iii. 205 = oblongus.
platyspathus, *Mart. ex Kunth, l. c.* 209.—Burma.
plicatus, *Blume, Rumphia*, iii. 67.—Ins. Celebes.
polygamus, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 780.—Reg. Himal.
Pseudo-Rotang, Mart. ex Kunth, Enum. Pl. iii. 207 = viminalis.
pulcher, *Miq. Anal. Bot. Ind.* i. 3.—Borneo.
quinquenervius, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 777.—Reg. Himal.
radiatus, *Thw. Enum. Pl. Zeyl.* 431.—Zeylan.
radicalis, *H. Wendl. & Drude, in Linnaea*, xxxix. (1875) 195.—Austral.
ramosissimus, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 78.—Malaya.
Reinwardtii, *Mart. Hist. Nat. Palm.* iii. 208. t. 112.—Malaya.
Rheedei, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 73, nota.—Malabar.
rhomboideus, *Blume, in Schult. f. Syst.* vii. 133.—Malaya.
Riedelianus, *Miq. in Verh. Kon. Akad. Wetensch.* xi. (1868) n. v. 18.—Ins. Celebes.
rivalis, *Thw. ex Trimen, in Journ. Bot.* xxiii. (1885) 386, 268.—Zeylan.
Rotang, *Linn. Sp. Pl.* 325.—Ind. or.
Roxburghii, Griff. in Calc. Journ. Nat. Hist. v. (1845) 43 = C. Rotang.
Royleanus, Griff. l. c. 40 = tenuis.
ruber, Reinw. ex Kunth, Enum. Pl. iii. 208 = Daemonorops Draco.
rudentum, *Lour. Fl. Cochinch.* 209.—Cochinch.
rudentum, Mart. Hist. Nat. Palm. iii. 340 = rivalis.
rudentum, Roxb. Hort. Beng. 72; Fl. Ind. iii. 776 = albus.
Rumphii, Blume, Rumphia, iii. 38 = Daemonorops Rumphii.
Salakka, Willd. ex Steud. Nom. ed. II. i. 252 = Zalacca edulis.
schistocanthos, *Blume, Rumphia*, iii. 49.—Sumatra.
schizospathus, Griff. in Calc. Journ. Nat. Hist. v. (1845) 32 = erectus.
Scipionum, Lam. Encyc. vi. 304 = C. Rotang.
Scipionum, *Lour. Fl. Cochinch.* 210.—Malaya.
secundiflorus, Beauv. Fl. Owar. i. 15. t. 9, 10 = Ancistrophyllum secundiflorum.
siphonospathus, *Mart. Hist. Nat. Palm.* iii. 342.—Ins. Philipp.
spectabilis, *Blume, Rumphia*, iii. 55.—Sumatra.
stoloniferus, Teijsm. & Binn. Cat. Hort. Bog. 75 = horrens.
strictus, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 28 = Daemonorops longipes.
subangulatus, *Miq. Fl. Ind. Bat. Suppl.* 594.—Sumatra.
symphysipus, *Mart. Hist. Nat. Palm.* iii. 336.—Ins. Celebes.
tenuis, *Roxb. Hort. Beng.* 73; *Fl. Ind.* iii. 780.—Reg. Himal.; Burma; Cochinch.
tetradactylus, *Hance, in Journ. Bot.* xiii. (1875) 289.—China.
tetrastichus, Blume, Rumphia, iii. 62 = javensis.
thysanolepis, *Hance, in Journ. Bot.* xii. (1874) 265.—China.
tigrinus, Kurz, in Journ. As. Soc. Beng. xliii. (1874) II. 211. t. 25 = longisetus.
trichrous, Miq. in Verh. Kon. Akad. Wetensch. xi. (1868) n. v. 28 = Daemonorops trichrous.
unifarius, *H. Wendl. in Bot. Zeit.* xvii. (1859) 158.—Ins. Nicobar.; Java.
usitatus, Blanco, Fl. Filip. ed. I. 265 = Daemonorops Gaudichaudii.
usitatus, Mart. Hist. Nat. Palm. iii. 340 = Blancoi.
Verschaffelti, Hort. ex H. Wendl. in Kerch. Palm. 238 = Acanthophoenix crinita.
verticillaris, Griff. in Calc. Journ. Nat. Hist. v. (1845) 63 = Daemonorops verticillaris.
verus, Lour. Fl. Cochinch. 210 = pisicarpus.
viminalis, *Reinw. ex Mart. Nat. Hist. Palm.* iii. 335. t. U. 111.—Celebes; Java?
viminalis, *Willd. Sp. Pl.* ii. 203.—Ind. or.; Malaya.
Walkeri, *Hance, in Journ. Bot.* xii. (1874) 266.—China.
Wightii, Griff. Palms Brit. Ind. 102. t. 216 G = Huegelianus.

CALAMUS:—
Zalacca, Gaertn. Fruct. ii. 267. t. 139 = Zalacca edulis.
Zalacca, Roxb. Hort. Beng. 72; Fl. Ind. iii. 773 = Zalacca Wallichiana.

CALANCHOË, Pers. Syn. i. 445 (1805) = **Kalanchoë**, Adans. (Crassulac.).

CALANDARIUM, Juss. ex Steud. Nom. ed. II. i. 252 (1840) = seq.

CALANDRINIA, H. B. & K. Nov. Gen. et Sp. vi. 77. t. 526 (1823). *PORTULACEAE*, Benth. & Hook. f. i. 158.
BAITARIA, Ruiz & Pav. Prod. 63. t. 36 (1794).
CISTANTHE, Spach, Hist. Vég. Phan. v. 229 (1833).
COSMIA, Domb. ex Juss. Gen. 312 (1789).
DIAZIA, Phil. Fl. Atac. 22. t. 1 E (1830).
GEUNSIA, Moç. & Sesse, ex DC. Prod. iii. 358 (1828).
PHACOSPERMA, Haw. in Phil. Mag. i. (1827) 124.
RHODOPSIS, Lilja, Fl. Sverig. Suppl. i. 42 (1840).
TEGNERIA, Lilja, Fl. Sverig. odl. Vext. 76 (1839).
acaulis, *H. B. & K. Nov. Gen. et Sp.* vi. 78.—Ecuador.
acuta, Meyen, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 340 = biflora.
adscendens, Hort. ex Phil. f. Pl. Vasc. Chil. 85 = pilosiuscula.
affinis, *Gill. ex Arn. in Cheek, Edinb. Journ.* iii. (1831) 355.—Chili.
alba, *DC. Prod.* iii. 359.—Peruv.
amoena, *Vis. in Nuov. Sagg. Padov.* vi. (1847) 94.—Hab.?
andicola, *Gill. in Hook. Bot. Misc.* iii. (1833) 332.—Chili.
Andrewsii, *Sweet, Hort. Brit.* ed. II. 219.—Ind. occ.
Andrewski, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 865 = Talinum patens.
angulata, Hort. ex Fisch. & Mey. Ind. Sem. Hort. Petrop. xi. 53 = procumbens.
arenaria, *Cham. in Linnaea*, vi. (1831) 563.—Chili.
aurea, *Barn. in C. Gay, Fl. Chil.* ii. 510.—Chili.
axilliflora, *Barn. l. c.* 486.—Chili.
balonensis, *Lindl. in Mitch. Journ. Trop. Austral.* 148.—Austral.
Berteroana, *Phil. in Linnaea*, xxviii. (1856) 643.—Chili.
biflora, *Meyen, Reise*, i. 349.—Chili.
brevipedata, *F. Muell. Fragm.* x. 69.—Austral.
Breweri, *S. Wats. in Proc. Am. Acad.* xi. (1876) 124.—Am. bor. occ.
Buridgii, *Hort. ex Regel, Gartenfl.* (1855) 4.—Am. austr.
cachinalensis, *Phil. Fl. Atac.* 20.—Chili.
caespitosa, *Gill. ex Arn. in Cheek, Edinb. Journ.* iii. (1831) 356.—Chili.
calycina, *Phil. in Linnaea*, xxxiii. (1864–65) 75.—Chili.
calycotricha, *Phil. l. c.* xxx. (1859–60) 188.—Chili.
calyptrata, *Hook. f. in Ic. Pl.* t. 296.—Austral.
capitata, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 334.—Chili.
caulescens, *H. B. & K. Nov. Gen. et Sp.* vi. 78. t. 526.—Ecuador; Peruv.
Chamissoi, *Barn. in C. Gay, Fl. Chil.* ii. 497.—Chili.
chromantha, *Griseb. in Goett. Abh.* xxiv. (1879) 30.—Reg. Argent.
ciliata, *DC. Prod.* iii. 359.—Peruv.
cistiflora, *Gill. ex Arn. in Cheek, Edinb. Journ.* iii. (1831) 355.—Chili.
colchaguensis, *Barn. in C. Gay, Fl. Chil.* ii. 483.—Chili.
compacta, *Barn. l. c.* 481.—Chili.
composita, *Benth. Fl. Austral.* i. 175.—Austral.
compressa, *Schrad. ex DC. Prod.* iii. 359.—Chili.
conferta, *Gill. ex Arn. in Cheek, Edinb. Journ.* iii. (1831) 356.—Chili.
coquimbensis, *Barn. in C. Gay, Fl. Chil.* ii. 483.—Chili.
corrigioloides, *F. Muell. ex Benth. Fl. Austral.* i. 175.—Austral.
corymbosa, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 341.—Chili.
Cotyledon, *S. Wats. in Proc. Am. Acad.* xx. (1885) 355.—Calif.
Cumingii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 334.—Chili.

CALANDRINIA:—
curvifolia, Hort. ex Heynh. Nom. ii. 105 = procumbens.
demissa, *Phil. in Linnaea*, xxviii. (1856) 644.—Chili.
densiflora, *Barn. in C. Gay, Fl. Chil.* ii. 503.—Chili.
densiflora, *Phil. in Linnaea*, xxviii. (1856) 642.—Chili.
denticulata, *Gill. in Hook. Bot. Misc.* iii. (1833) 333.—Chili.
dianthoides, *Phil. in Anal. Univ. Chil.* (1872) 717.—Chili.
diffusa, *Gill. ex Arn. in Cheek, Edinb. Journ.* iii. (1831) 355.—Chili.
discolor, Lindl. Bot. Reg. (1839) t. 4 = Lindleyana.
discolor, *Schrad. in Linnaea*, viii. (1833) Lit. 22.—Chili.
elegans, Hort. ex Vilm. Fl. Pl. Terre, ed. I. 149 = discolor.
elegans, Spach, Hist. Vég. Phan. v. 232 = Menziesii.
erythrocoma, *Phil. in Linnaea*, xxxiii. (1864–65) 76.—Chili.
fasciculata, *Phil. l. c.* xxviii. (1856) 645.—Chili.
Fenzlii, *Barn. in C. Gay, Fl. Chil.* ii. 493.—Chili.
ferruginea, *Barn. l. c.* 513.—Chili.
floribunda, *Phil. in Linnaea*, xxxiii. (1864–65) 78.—Chili.
frigida, *Barn. in C. Gay, Fl. Chil.* ii. 495.—Chili.
fruticulosa, Kunze, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 340 = trifida.
Gaudichaudii, *Barn. in C. Gay, Fl. Chil.* ii. 490.—Chili.
Gayana, Barn. l. c. 503 = biflora.
Gilliesii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 333.—Chili.
glandulosa, Hort. ex Steud. Nom. ed. II. i. 253 = procumbens.
glauca, Schrad. ex DC. Prod. iii. 359 = grandiflora.
glomerata, *Phil. in Linnaea*, xxviii. (1856) 644.—Chili.
gracilis, *Benth. Fl. Austral.* i. 173.—Austral.
graminifolia, *Phil. in Linnaea*, xxviii. (1856) 690.—Chili.
grandiflora, *Lindl. Bot. Reg.* t. 1194.—Chili.
granulifera, *Benth. Fl. Austral.* i. 176.—Austral.
hirtella, *Phil. in Linnaea*, xxviii. (1856) 691.—Chili.
Landbeckii, *Phil. l. c.* xxxiii. (1864–65) 77.—Chili.
Leana, *Porter, in Coult. Bot. Bull.* i. (1876) 49.—Calif.
Lehmanni, *Endl. in Lehm. Pl. Preiss.* ii. 235.—Austral.
leucocephala, *Phil. Fl. Atac.* 21.—Chili.
leucotricha, *Phil. in Anal. Univ. Chil.* (1861) 91.—Chili.
Lindleyana, Hort. ex Vilm. Fl. Pl. Terre, ed. II. 162 = discolor.
Lindleyana, *Walp. Rep.* ii. 235.—Hab.?
linearifolia, *DC. Prod.* iii. 359.—Mexic.?
lingulata, *DC. l. c.* 358.—Peruv.
lingulata, Hort. ex Steud. Nom. ed. II. i. 25? = procumbens.
liniflora, *Fenzl, in Enum. Pl. Hueg.* 52.—Austral.
litoralis, *Phil. Fl. Atac.* 20.—Chili.
Llavea, Loud. Hort. Brit. Suppl. i. 616, sphalm. = Cuphea Llavea.
Lockharti, *Sweet, Hort. Brit.* ed. II. 219.—Ins. Trinit.
longiscapa, *Barn. in C. Gay, Fl. Chil.* ii. 492.—Chili.
macilenta, *Barn. l. c.* 488.—Chili.
maritima, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 197.—Calif.
megarhiza, *Hemsl. Diag. Pl. Nov.* 23.—Guatemala.
Menziesii, *Torr. & Gray, Fl. N. Am.* i. 197.—Am. bor. occ.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 341.—Chili.
micrantha, *Schlecht. in Linnaea*, xiii. (1839) Lit. 97.—Mexic.
minima, *Bert. ex Steud. Nom.* ed. II. i. 253.—Chili.
minutissima, *Barn. in C. Gay, Fl. Chil.* ii. 504.—Chili.
modesta, *Phil. Fl. Atac.* 20.—Chili.
monandra, DC. Prod. iii. 359 = Monocosmia corrigioloides.
mucronulata, *Meyen, Reise*, i. 314.—Chili.
multicaulis, *Phil. in Linnaea*, xxviii. (1856) 641.—Chili.
nevadensis, *A. Gray, in Proc. Am. Acad.* viii. (1873) 623.—Am. bor. occ.
nitida, *DC. Prod.* iii. 359.—Chili.

CALANDRINIA:—
oblongifolia, *Barn. in C. Gay, Fl. Chil.* ii. 482.—Chili.
occulta, *Phil. Fl. Atac.* 20.—Chili.
oppositifolia, *S. Wats. in Proc. Am. Acad.* xx. (1885) 355.—Calif.
paniculata, *DC. Prod.* iii. 358.—Peruv.
parviflora, Hort. ex Steud. Nom. ed. II. i. 253 = procumbens.
Phacosperma, *DC. Prod.* iii. 359.—Peruv.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 144.—Chili.
picta, Gill. ex Arn. in Cheek, Edinb. Journ. iii. (1831) 356 = arenaria.
pilosiuscula, *DC. Prod.* iii. 359.—Chili.
pleiopetala, *F. Muell. Fragm.* x. 70.—Austral.
Poeppigiana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 340.—Chili.
pogonophora, *F. Muell. Fragm.* x. 69.—Austral.
polyandra, *Benth. Fl. Austral.* i. 172.—Austral.
polycarpoides, *Phil. in Linnaea*, xxviii. (1856) 644.—Chili.
polyclados, *Phil. l. c.* xxxiii. (1864–65) 77.—Chili.
polypetala, *Fenzl, in Enum. Pl. Hueg.* 51.—Austral.
potentilloides, *Barn. in C. Gay, Fl. Chil.* ii. 512.—Chili.
procumbens, *Moris, in Mem. Acc. Tor.* xxxvii. (1834) 100.—Chili.
prostrata, *Phil. in Linnaea*, xxxiii. (1864–65) 76.—Chili.
ptychosperma, *F. Muell. Fragm.* iv. 137.—Austral.
pulchella, Lilja, in Linnaea, xvii. (1843) 109 = Menziesii.
pumila, *F. Muell. Fragm.* x. 68.—Austral.
pusilla, *Barn. in C. Gay, Fl. Chil.* ii. 485.—Chili.
pusilla, *Lindl. in Mitch. Trop. Austral.* 360.—Austral.
pygmaea, *A. Gray, in Proc. Am. Acad.* viii. (1873) 623.—Am. bor. occ.
pygmaea, *F. Muell. Fragm.* i. 175.—Austral.
quadrivalvis, *F. Muell. l. c.* 176.—Austral.
racemosa, Steud. Nom. ed. II. i. 253 = arenaria.
ramosissima, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 334.—Chili.
rupestris, *Barn. in C. Gay, Fl. Chil.* ii. 485.—Chili.
salsoloides, *Barn. l. c.* 502.—Chili.
Saxifraga, *Barn. l. c.* 506.—Chili.
sericea, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 334.—Chili.
sericea, Poepp. ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 340 = Poeppigiana.
setosa, *Phil. in Linnaea*, xxviii. (1856) 690.—Chili.
speciosa, Lehm. in Linnaea, vi. (1831) Litt. 74 = discolor.
speciosa, Lindl. Bot. Reg. t. 1598 = Menziesii.
spectabilis, *Otto & Dietr. Allg. Gartenz.* i. (1833) 161.—Chili.
spergularina, *F. Muell. Fragm.* i. 175.—Austral.
spicata, *Phil. Fl. Atac.* 21.—Chili.
splendens, *Barn. in C. Gay, Fl. Chil.* ii. 499.—Chili.
strophiolata, *F. Muell. Fragm.* xi. 82.—Austral.
tenella, Hook. Bot. Beech. Voy. 24 = pilosiuscula.
tenuifolia, *Phil. in Linnaea*, xxviii. (1856) 691.—Chili.
tricolor, *Phil. l. c.* 642.—Chili.
trifida, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 333.—Chili.
tuberosa, *Benth. Pl. Hartw.* 9.—Mexic.
umbellata, *DC. Prod.* iii. 358.—Chili.
umbellata, Gill. ex Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 334 = Gilliesii.
uniflora, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 41.—Austral.
venulosa, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 336 = arenaria.
venusta, Steud. Nom. ed. II. i. 253 = arenaria.
vinulosa, Walp. Rep. iii. 236 = arenaria.
virgata, *Phil. in Linnaea*, xxviii. (1856) 642.—Chili.
volubilis, *Benth. Fl. Austral.* i. 174.—Austral.

CALANTHE, R. Br. in Bot. Reg. sub t. 573 (1821). *ORCHIDEAE*, Benth. & Hook. f. iii. 520.
AMBLYGLOTTIS, Blume, Bijdr. 369. t. 64 (1825).
CENTROSIS, Thou. Orch. Iles Afr. t. 35, 36 (1822).
GHIESBREGHTIA, A. Rich. in Ann. Sc. Nat. Sér. III. iii. (1845) 28.
PREPTANTHE, Reichb. f. in Fl. des Serres, Sér. I. viii. (1853) 245.
STYLOGLOSSUM, Breda, Orch. Kuhl et Hass. (1827).
ZEDUBA, Buch.-Ham. ex Meissn. Gen. Comm. 280 (1842).

CALANTHE:—
abbreviata, *Lindl. Gen. et Sp. Orch.* 251.—Java.
alismaefolia, *Lindl. Fol. Orchid. Calanth.* 8.—Reg. Himal.
alpina, *Hook. f. ex Lindl. l. c.* 4.—Reg. Himal.
alta, *Reichb. f. Otia Bot. Hamb.* 53.—Ins. Fiji.
anchorifera, *Reichb. f. in Gard. Chron.* (1883) II. 166.—Polynesia.
angraeciflora, *Reichb. f. in Linnaea*, xli. (1877) 75.—N. Caled.
angusta, *Lindl. Fol. Orchid. Calanth.* 7.—Reg. Himal.
angustifolia, *Lindl. Gen. et Sp. Orch.* 250.—Malaya.
aristulifera, *Reichb. f. in Bot. Zeit.* xxxvi. (1878) 74.—Japon.
australis, Ait. ex Loud. Hort. Brit. Suppl. ii. 615 = Spiranthes australis.
Barberiana ×, Reichb. f. in Gard. Chron. (1881) I. 136.
bella ×, Reichb. f. l. c. 234.
bicolor, Lindl. Scrt. Orch. sub t. 9 = striata.
biloba, *Lindl. Fol. Orchid. Calanth.* 3.—Reg. Himal.
bracteosa, *Reichb. f. in Flora*, lxv. (1882) 533.—Ins. Samoa.
brevicornu, *Lindl. in Wall. Cat.* n. 7338.—Reg. Himal.
catilligera, *Reichb. f. in Bonplandia*, v. (1857) 37.—Java.
Ceciliae, *Hort. Low.—Cf. Gard. Chron.* (1883) I. 432.—Malaya.
chloroleuca, *Lindl. Fol. Orchid. Calanth.* 10.—Reg. Himal.
citrina, *Scheidw. in L'Hortic. Belge*, v. (1838) 185. t. 104.—Hab.?
clavata, *Lindl. in Wall. Cat.* n. 7343.—Reg. Himal.; Burma.
colorans, *Reichb. f. in Gard. Chron.* (1885) II. 360.—Hab.?
comosa, Reichb. f. in Linnaea, xix. (1847) 374 = veratrifolia.
conspicua, *Lindl. Fol. Orchid. Calanth.* 4.—Malaya.
corymbosa, *Lindl. in Journ. Linn. Soc.* vi. (1862) 129.—Afr. trop.
curculigoides, *Lindl. in Wall. Cat.* n. 7340.—Malaya.
Curtisii, *Reichb. f. in Gard. Chron.* (1884) II. 262.—Ins. Sondaic.
densiflora, *Lindl. in Wall. Cat.* n. 7344.—Reg. Himal.
dipteryx, *Reichb. f. in Gard. Chron.* (1884) II. 394.—Ins. Sondaic.
discolor *Lindl. Scrt. Orch.* sub t. 9.—Japon.
Dominii ×, Lindl. in Gard. Chron. (1858) 4.
emarginata, *Lindl. Gen. et Sp. Orch.* 249.—Java.
flava, *Morr. in L'Hortic. Belge*, ii. (1834) 238. t. 46.—Java.
Foerstermanni, *Reichb. f. in Gard. Chron.* (1883) I. 814.—Burma.
fulgens, Lindl. Fol. Orchid. Calanth. 10 = C. Masuca.
furcata, *Batem. ex Lindl. Bot. Reg.* (1838) *Misc.* 28.—Ins. Luzon.
galeata, Lindl. Fol. Orchid. Calanth. 5 = chloroleuca.
Gebina, Lodd. Cat. ex Lindl. l. c. 11 = Bletia hyacinthina.
gracilis, *Lindl. in Wall. Cat.* n. 7341.—Reg. Himal.; China.
gracillima, *Lindl. Fol. Orchid. Calanth.* 8.—Ins. Carolin.
granatensis, *Reichb. f. in Bonplandia*, iv. (1856) 322. N. Granat.
grandiflora, *Nadeaud, Enum. Pl. Tahiti*, 37.—Ins. Pacif.
Griffithii, *Lindl. in Paxt. Fl. Gard.* iii. (1852–53) 37.—Reg. Himal.
herbacea, *Lindl. Fol. Orchid. Calanth.* 10.—Reg. Himal.
hololeuca, *Reichb. f. in Seem. Fl. Vit.* 298.—Ins. Fiji.
japonica, *Blume, in Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 205.—Japon.
Jusnerii, *Boxall, ex Naves, in Blanco, Fl. Filip.* ed. III. *Nov. App.* 247.—Ins. Philipp.
labrosa, *Reichb. f. in Gard. Chron.* (1883) I. 44.—Burma.
Langei, *F. Muell. in Wing, South Sc. Record*, N.S. i. (1885).—N. Caled.
lentiginosa ×, Reichb. f. in Gard. Chron. (1883) I. 44.
lyroglossa, *Reichb. f. Otia Bot. Hamb.* 53.—Afr. trop.
Masuca, *Lindl. in Wall. Cat.* n. 7337.—Ind. or.; Malaya.
mexicana, *Reichb. f. in Linnaea*, xviii. (1844) 406.—Mexic.

CALANTHE:—
natalensis, *Reichb. f. in Bonplandia*, iv. (1856) 322.—Afr. austr.
nivalis, *Boxall, ex Naves, in Blanco, Fl. Filip.* ed. III. *Nov. App.* 247.—Ins. Philipp.
occidentalis, Lindl. Fol. Orchid. Calanth. 3 = tricarinata.
odora, *Griff. Notul.* iii. 365.—Reg. Himal.
parviflora, *Lindl. in Paxt. Fl. Gard.* iii. (1852–53) 37, sub t. 61.—Java.
Perrottetii, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 68 = veratrifolia.
Petri, *Reichb. f. in Gard. Chron.* (1880) II. 326.—Polynesia.
phajoides, Reichb. f. in Bonplandia, v. (1857) 37 = angustifolia.
pilosa, *Lindl. Fol. Orchid. Calanth.* 11.—Sumatra.
plantaginea, Griff. Notul. iii. 368 = Griffithii.
plantaginea, *Lindl. in Wall. Cat.* n. 7346.—Reg. Himal.
pleiochroma, *Reichb. f. in Gard. Chron.* (1871) 938.—Japon.
porphyrea ×, Reichb. f. l. c. (1884) I. 76.
proboscidea, *Reichb. f. l. c.* 476.—Ins. Sondaic.
pubcrula, *Lindl. in Wall. Cat.* n. 7342.—Reg. Himal.
pulchra, *Lindl. Gen. et Sp. Orch.* 250.—Java.
purpurea, *Lindl. l. c.* 249.—Zeylan.
reflexa, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 68.—Japon.
Regnieri, *Reichb. f. in Gard. Chron.* (1883) I. 274.—Cochinch.
rosea, *Benth. in Journ. Linn. Soc.* xviii. (1880) 309.—Burma.
Sandhurstiana ×, Gosse, ex Reichb. f. in Gard. Chron. (1881) I. 395.
Sedeni ×, Reichb. f. l. c. (1878) I. 168.
Sieboldi, *Decne. ex Regel, Ind. Sem. Hort. Petrop.* (1868) 80.—Japon.
speciosa, *Lindl. Gen. et Sp. Orch.* 250.—Java.
speciosa, *Vieill. in Ann. Sc. Nat.* Sér. IV. xvi. (1861) 44.—N. Caled.
Stevensiana, *Régnier, ex Reichb. f. in Gard. Chron.* (1883) I. 274, in nota.—Java?
striata, *R. Br. in Bot. Reg.* 573, in nota.—China; Japon.
sulphurea, *Hort. Hueg. ex Heynh. Nom.* ii. 105, nomen.—Ind. or.
sumatrana, *Blanco, ex Boerl. in Hasselt & Boerl. Bijdr. Midd. Sum.* 33.—Sumatra.
sylvatica, *Lindl. Gen. et Sp. Orch.* 250.—Madag.
sylvestris, Lindl. ex Steud. Nom. ed. II. i. 253 = sylvatica.
tahitensis, *Nadeaud, Enum. Pl. Tahiti*, 37.—Ins. Tahiti.
Textori, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 204.—Japon.
triantherifera, *Nadeaud, Enum. Pl. Tahiti*, 38.—Ins. Tahiti.
tricarinata, *Lindl. in Wall. Cat.* n. 7339.—Reg. Himal.
Turneri, *Reichb. f. in Gard. Chron.* (1883) I. 274.—Java?
uncata, *Lindl. Fol. Orchid. Calanth.* 6.—Reg. Himal.
vaginata, *Lindl. l. c.* 7.—Assam.
varicgata, *Scheidw. in L'Hortic. Belge*, v. (1838) 185. t. 105.—Hab.?
Veitchii ×, Hort.—Cf. Gard. Chron. (1859) 1016.
Ventilabrum, *Reichb. f. in Seem. Fl. Vit.* 298.—Ins. Fiji.
veratrifolia, *R. Br. in Bot. Reg.* sub t. 573.—Ind. or.; Malaya; Austral.
veratrifolia, Hook. Bot. Beech. Voy. 71 = gracillima.
versicolor, Lindl. Scrt. Orch. 42 = C. Masuca.
vestita, *Lindl. in Wall. Cat.* n. 7345.—Burma; Malaya.
viridi-fusca, Hook. Bot. Mag. t. 4669 = Tainia viridifusca.
Wightii, Reichb. f. in Walp. Ann. vi. 932 = C. Masuca.
Zollingeri, *Miq. Fl. Ind. Bat.* iii. 711.—Java.
Zollingeri, *Reichb. f. in Bonplandia*, v. (1857) 38.—Java.

CALANTHERA, Kunth (?), ex Hook. Kew Journ. viii. (1856) 18 = **Buchloë**, Engelm. (Gramin.).
dactyloides, Kunth, l. c. = Buchloë dactyloides.

CALANTHUS, Oerst. in Vidensk. Selsk. Scr. V. v. (1861) 121; (Gesn. Centr. Am. 47. t. 4) = **Alloplectus**, Mart. (Gesnerac.).
multiflorus, Oerst. l. c. = A. multiflorus.

CALANTICA, Jaub. ex Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 74. *SAMYDACEAE*, Benth. & Hook. f. i. 799.
cerasifolia, *Tul. l. c.* 76.—Madag.
grandiflora, *Jaub. ex Tul. l. c.* 75.—Madag.
Jalberti, Baill. Hist. Pl. iv. 277 = Bivinia Jaubertia.

CALASIAS, Rafin. Fl. Tellur. iv. 64 (1836) (nom. prius) = **Anisotes**, Nees (Acanthac.).
bracteata, Rafin. l. c. = A. trisulcus.

CALATHEA, G. F. W. Mey. Prim. Fl. Esseq. 6 (1818). *SCITAMINEAE*, Benth. & Hook. f. iii. 653.
ENDOCODON, Rafin. Fl. Tellur. iv. 49 (1836).
GOEPPERTIA, Nees, in Linnaea, vi. (1831) 337.
MONOSTICHIE, Koern. in Regel, Gartenfl. (1858) 88.
PSYDARANTA?, Neck. Elem. iii. 145 (1790).
ZELMIRA, Rafin. Fl. Tellur. iv. 50 (1836).
Ackermanni, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 133.—Bras.
aemula, *Koern. l. c.* 131.—Bras.
albicans, *A. Brongn. ex Gris, in Ann. Sc. Nat.* Sér. IV. xii. (1859) 193.—Hab.?
affinis, *Fenzl, ex Regel, Gartenfl.* (1879) 294.—Bras.?
Allouga, Steud. Nom. ed. II. i. 253 = C. Allouia.
Allouia, *Lindl. Bot. Reg.* sub t. 1210.—Guiana.
altissima, *Horan. Prod. Scitam.* 13; *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 141.—Am. trop.
angustifolia, *Koern. in Regel, Gartenfl.* (1858) 87.—Am. trop.
applicata, *J. Makoy, ex E. Morr. in Belg. Hortic.* xxiv. (1874) 228.—Bras.
argyraea, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 142.—Bras.
Arrabida, *Koern. l. c.* 144.—Bras.
arrecta, *Lindl. & André, in Illustr. Hortic.* (1871) 174. t. 77.—Ecuador.
Backemiana, *E. Morr. in Belg. Hortic.* xxv. (1875) 271. f. 4.—Bras.
bambusacea, Poepp. & Endl. Nov. Gen. et Sp. ii. 23 = Ischnosiphon bambusacea.
bella, *Regel, Gartenfl.* (1879) 297.—Hab.?
bellula, *Linden, Cat.* n. 89 (1872) 2.—Reg. Amazon.
bicolor, Steud. Nom. ed. II. i. 253 = Maranta bicolor.
blanda, *Steud. l. c.*—Bras.
brasiliensis, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 118.—Bras.
brevipes, *Koern. l. c.* 140.—Bras.
Cachibou, Lindl. ex Horan. Prod. Scitam. 12 = lutea.
capitata, *Lindl. Bot. Reg.* t. 1210.—Peruv.
Cassupito, *G. F. W. Mey. Prod. Fl. Esseq.* 10.—Venezuela.
chrysantha, *Horan. Prod. Scitam.* 13.—Bras.
chimboracensis, *Linden, Illustr. Hortic.* (1870) t. 6.—Ecuador.
chlorosticta, Hort. Vind. ex Regel, Gartenfl. (1879) 294 = angustifolia.
chrysoleuca, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 142.—Peruv.
cinerea, *Regel, Gartenfl.* (1876) 2.—Reg. Amazon.
colorata, *Benth. & Hook. f. Gen.* iii. 654.—Bras.
comosa, *Lindl. Bot. Reg.* sub t. 1210.—Ind. occ.
composita, Klotzsch, in Schomb. Brit. Guian. iii. 918 = Ischnosiphon Parkeri.
conferta, *Benth. in Benth. & Hook. f. Gen.* iii. 653.—Afr. trop.
crocata, *E. Morr. & Joriss. in Belg. Hortic.* xxv. (1875) 141. t. 8.—Bras.
densa, *Regel, Ind. Sem. Hort. Petrop.* (1866) 83.—Bras.
dicephala, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 127.—Peruv.
discolor, *G. F. W. Mey. Prim. Fl. Esseq.* 7.—Guiana.
dubia, Lindl. Bot. Reg. sub t. 1210 = Ischnosiphon spicatus.
eximia, *Koern. ex Regel, in Gartenfl.* (1858) 87.—Am. trop.
exscapa, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) I. 122.—Peruv.
fasciata, *Regel & Koern. in Gartenfl.* (1858) 348.—Bras.
fasciculata, Presl, Reliq. Haenk. i. 108. t. 16 = Phrynium fasciculatum.
flavescens, *Lindl. Bot. Reg.* t. 932.—Bras.

CALATHEA :—
Glaziovi, *Benth. in Benth. & Hook. f. Gen.* iii. 654.—Bras.
grandifolia, *Lindl. Bot. Reg.* t. 1210.—Bras.
hieroglyphica, *Linden & André, Illustr. Hortic.* (1873) 63. t. 123.—N. Granat.
Jagoriana, *Regel, Gartenfl.* (1879) 297.—Bras.
juncea, Spreng. Syst. i. 8 = Ischnosiphon Arouma.
Kappleriana, *Koern. ex Horan. Prod. Scitam.* 12.—Bras.
Koernickeana, Horan. l. c. = brasiliensis.
Koernickiana, *Regel, Gartenfl.* (1874) 33.—Bras.
Kummeriana, *E. Morr. in Belg. Hortic.* xxv. (1875) 270. t. 15–17. f. 1.—Bras.
lanceolata, Koern. in Bull. Soc. Nat. Mosc. xxxv. (1862) i. 134 = longifolia.
lateralis, *Lindl. Bot. Reg.* sub t. 1210.—Peruv.
latifolia, *Klotzsch, in Schomb. Faun. et Fl. Guian.* 918.—Guiana.
laxa, Poepp. & Endl. Nov. Gen. et Sp. ii. 22. t. 130 = Ischnosiphon plurispicatus.
Legrelliana, *Regel, Gartenfl.* (1879) 301.—Ecuador.
leopardina, *Regel, l. c.* (1877) 35. t. 893.—Bras.
leucocephala, D. Dietr. Syn. Pl. i. 7 = Ischnosiphon leucophaeus.
leuconeura, Hort. ex Nichols. Dict. Gard. i. 237 = Maranta leuconeura.
leucophaea, Poepp. & Endl. Nov. Gen. et Sp. ii. 21. t. 129 = Ischnosiphon leucophaeus.
leucostachys, *Hook. f. Bot. Mag.* t. 6205.—Costa Rica.
Lietzei, *E. Morr. in Belg. Hortic.* xxv. (1875) 273. f. 8.—Bras.
Lindeni, Wallis & André, Illustr. Hortic. (1871) 211. t. 82 = seq.
Lindeniana, *Wallis, in Belg. Hortic.* xvi. (1866) 200.—Peruv.
littoralis, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 144.—Bras.
longibracteata, *Lindl. Bot. Reg.* t. 1020.—Bras.
longifolia, *Klotzsch, ex Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 134.—Bras.
lutea, *G. F. W. Mey. Prod. Fl. Esseq.* 10.—Venezuela.
macilenta, *Lindl. Bot. Reg.* sub t. 1210.—Bras.
macrostachya, *Griseb. in Goett. Abh.* xxiv. (1879) 335.—Reg. Argent.
macrostachya, Klotzsch, in Schomb. Faun. et Fl. Brit. Guian. 918 = Ischnosiphon obliquus.
Makoyana, *E. Morr. in Belg. Hortic.* xxii. (1872) 321. t. 24 & 25.—Bras.
Mandioccae, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 138.—Bras.
Mannii, *Benth. in Benth. & Hook. f. Gen.* iii. 653.—Afr. trop.
Mansonis, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 119.—Bras.
marantina, C. Koch, Allg. Gartenz. (1857) 163 = C. Casupito.
medio-picta, *Makoy, ex E. Morr. in Belg. Hortic.* xxiv. (1874) 228.—Bras.
metallica, *Koern. ex Regel, Gartenfl.* (1858) 88.—Hab. ?
micans, *Koern. l. c.* 87.—Peruv.
microcephala, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 125.—Peruv.
mirabilis, *Makoy, ex E. Morr. in Belg. Hortic.* xxiv. (1874) 228.—Bras.
modesta, *A. Brongn. ex Gris, in Ann. Sc. Nat.* Sér. IV. xii. (1859) 193.—Hab. ?
monophylla, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 144.—Bras.
Myrosma, *Koern. in Regel, Gartenfl.* (1858) 87.—Guiana.
nigro-costata, *Linden & André, Illustr. Hortic.* xx. (1873) t. 144.—N. Granat.
nobilis, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 144.—Hab. ?
oblonga, *Koern. l. c.* 138.—Bras.
olivaris, Hort. ex Nichols. Dict. Gard. i. 238 = Makoyana.
Oppenheimiana, *E. Morr. in Belg. Hortic.* xxv. (1875) 271. f. 3.—Bras.
orbiculata, *Lodd. Bot. Cab.* t. 1879.—Ind. occ.
ornata, *Koern. in Regel, Gartenfl.* (1858) 87.—Am. trop.
ovata, *Lindl. Bot. Reg.* sub t. 1210.—Bras.
pachystachya, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 142.—Peruv.
pacifica, *Linden & André, Illustr. Hortic.* xix. (1872) t. 101.—Peruv.

CALATHEA :—
pardina, Planch. & Linden, in Fl. des Serres, Sér. II. i. (1856) 53 = villosa.
Parkeri, Poepp. & Endl. Nov. Gen. et Sp. ii. 22 = Ischnosiphon Parkeri.
Pavonii, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 116.—Peruv.
pavonina, *C. Koch & Linden, Belg. Hortic.* xv. (1865) 99.—Bras.
peruviana, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 128.—Peruv.
picturata, *C. Koch & Linden, ex C. Koch, Wochenschr.* vi. (1863) 346.—Bras.
polyphylla, Poepp. & Endl. Nov. Gen. et Sp. ii. 23. t. 131 = Ischnosiphon polyphyllus.
Princeps, *Regel, Gartenfl.* (1879) 302.—Reg. Amazon.
propinqua, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 122.—Peruv.
pulchella, *Koern. in Regel, Gartenfl.* (1858) 87.—Bras.
pumila, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 138.—Bras.
Riedeliana, Paull, ex Regel, in Act. Hort. Petrop. iii. (1875) ii. 284 = Koernickiana.
rosea-picta, *Regel, Ind. Sem. Hort. Petrop.* (1869) 12.—Am. trop.
Rossii, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 144.—Bras.
rotundifolia, *Koern. l. c.* 125.—Hab. ?
rotundifolia, Poepp. & Endl. Nov. Gen. et Sp. ii. 23 = Ischnosiphon rotundifolius.
rufibarba, *Fenzl, in Regel, Gartenfl.* (1879) 294.—Bras. ?
Sellowii, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 120.—Bras.
smaragdina, *Linden & André, Illustr. Hortic.* (1870) 78.—Ecuador ?
spicata, *Steud. Nom.* ed. II. i. 253.—Guiana; Bras.
strobilifera, *Miq. in Linnaea*, xxii. (1849) 79.—Surinam.
taeniosa, *G. Joriss. in Belg. Hortic.* xxvi. (1876) 83. t. 5.—Bras.
trifasciata, *Koern. in Regel, Gartenfl.* (1858) 86.—Guiana.
tuberosa, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 124.—Bras.
tubispatha, *Hook. Bot. Mag.* t. 5542.—Bolivia.
umbrosa, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 137.—Bras.
undulata, *Regel, Illustr. Hortic.* xix. (1872) t. 98.—Ecuador.
Vandenheckei, *Regel, Ind. Sem. Hort. Petrop.* (1869) 14.—Hab. ?
varians, *C. Koch & Math. in App. Cat. Sem. Hort. Berol.* (1855) 12.—Am. trop.
variegata, *Koern. in Regel, Gartenfl.* (1858) 88.—Am. trop.
Veitchiana, *Veitch, ex Hook. f. Bot. Mag.* t. 5535.—Bolivia ?
velutina, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 127.—Peruv.
vestita, *Baker, in Saund. Ref. Bot.* v. t. 311.—Bras.
villosa, *Lindl. Bot. Reg.* (1834) *Misc.* 61.—Guiana.
violacea, *Lindl. l. c.* sub t. 932; t. 962.—Bras.
virginalis, *Linden, ex Regel, Gartenfl.* (1879) 299.—Bras.
vittata, *Koern. l. c.* (1858) 88.—Hab. ?
Wallisi, *Regel, Ind. Sem. Hort. Petrop.* (1869) 14.—Reg. Amazon.
Warscewiczii, *Koern. in Regel, Gartenfl.* (1858) 87.—Am. trop.
Widgreni, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 117.—Bras.
Wiotiana, *Makoy, ex E. Morr. in Belg. Hortic.* xxiv. (1874) 228.—Bras.
zebrina, *Lindl. Bot. Reg.* sub t. 1210.—Bras.
zingiberina, *Koern. in Bull. Soc. Nat. Mosc.* xxxv. (1862) i. 122.—Bras.

CALATHIANA, Delarb. Fl. Auv. ed. II. i. 28 (1800) = **Gentiana**, Tourn.
nivalis, Delarb. l. c. = G. nivalis.

CALATHINUS, Rafin. Fl. Tellur. iv. 21 (1836) = **Narcissus**, Tourn. (Amaryllid.).
cernuus, Rafin. l. c. = N. calathinus.
multiflorus, Rafin. l. c. = N. Tazetta ?
serotinus, Rafin. l. c. = N. serotinus.

CALATHODES, Hook. f. & Thoms. Fl. Ind. i. 40 (1855). *RANUNCULACEAE*, Benth. & Hook. f. i. 7.
palmata, *Hook. f. & Thoms. l. c.* 41.—Reg. Himal.

CALATHOSTELMA, Fourn. in Mart. Fl. Bras. vi. iv. 219 (1885). *ASCLEPIADEAE.*
ditassoides, *Fourn. l. c.*—Bras.

CALBOA, Cav. Ic. v. 51. t. 476 (1799) = **Ipomoea**, Linn. (Convolvul.).
globosa, Lindl. in Journ. Hort. Soc. v. (1850) 82 = I. Llaveana.
vitifolia, Cav. Ic. v. 51. t. 476 = I. peduncularis.

CALCALIA, Krock. Fl. Siles. ii. ii. 381 = CACALIA, Linn. = **Adenostyles**, Cass. (Compos.).
alpina, Krock. l. c. = A. glabra.

CALCARUNIA, Rafin. Med. Fl. ii. 106 (1830) = **Pontederia**, Linn.
hastata, Rafin. l. c. = P. cordata.

CALCATRIPPA, Heist. Syst. 8 (1748) = **Delphinium**, Linn. (Ranunculac.).

CALCEARIA, Blume, Bijdr. 417. t. 33 (1825) = **Corysanthes**, R. Br. (Orchid.).
fornicata, Blume, l. c. = Cor. fornicata.
picta, Blume, l. c. = Cor. picta.

CALCEOLARIA, Heist. Syst. 5 (1748) = **Cypripedium**, Linn. (Orchid.).

CALCEOLARIA, Linn. Mant. ii. 143 (1771). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 929.
FAGELIA, Schwencke, in Verh. Bat. Gen. Roterd. i. (1774) 474. t. 13.
JOVELLANA, Ruiz & Pav. Fl. Per. i. 12 (1798).
adscendens, *Lindl. Bot. Reg.* t. 1215.—Chili.
alba, *Ruiz & Pav. Fl. Per.* i. 19. t. 27.—Chili.
alternifolia, *Cav. Ic.* v. 28. t. 445. f. 2.—Peruv.
ambigua, *Phil. in Linnaea*, xxix. (1857–58) 32.—Chili.
amplexicaulis, H. B. & K. Nov. Gen. et Sp. ii. 384.—Ecuador.
andina, *Benth. in DC. Prod.* x. 219.—Chili.
angustiflora, Hook. Bot. Mag. t. 3094 = verticillata.
angustiflora, *Ruiz & Pav. Fl. Per.* i. 17, 28. t. 28.—Ecuador.
angustifolia, Sweet, Hort. Brit. ed. II. 396 = integrifolia.
anomala, Pers. Syn. i. 16 = crenatiflora.
aquatica, *A. Br. & Bouché, Ind. Sem. Hort. Berol.* (1852) 2.—Hab. ?
arachnoidea, *R. Grah. in Edinb. N. Phil. Journ.* (Sept. 1828) 371; *et Bot. Mag.* t. 2874.—Chili.
araucana, *Phil. in Linnaea*, xxix. (1857–58) 33.—Chili.
argentea, *H. B. & K. Nov. Gen. et Sp.* ii. 387. t. 170.—Peruv.
Atkinsiana, Sweet, Brit. Flow. Gard. Ser. II. t. 168.
bartsiaefolia, *Wedd. Chlor. And.* ii. 134.—Bolivia.
bellidifolia, *Gill. ex Benth. in DC. Prod.* x. 207.—Chili.
bellidifolia, Salisb. Prod. 89 = Fothergillii.
Benthami, *Phil. in Anal. Univ. Chil.* (1873) 533.—Chili.
Bergii, *Hieron. in Bol. Acad. Nac. Cordova*, iii. (1879) 363.—Reg. Argent.
Berterii, *Colla, in Mem. Acc. Tor.* xxxix. (1836) 138.—Chili.
bicolor, *Ruiz & Pav. Fl. Per.* i. 16. t. 25.—Peruv.
bicrenata, *Ruiz & Pav. l. c.* 15. t. 23.—Peruv.
bifida, Willd. ex Link, in Spreng. Jahrb. i. iii. (1820) 53 = graminifolia.
biflora, Lam. Ill. i. 52; Encyc. i. 556 = plantaginea.
biflora, Ruiz & Pav. Fl. Per. i. 16. t. 20 = ovata.
bipinnatifida, *Phil. Fl. Atac.* 46.—Chili.
brachiata, Willd. ex Link, in Spreng. Jahrb. i. iii. (1820) 52 = flexuosa.
brevistyla, *Clos, in C. Gay, Fl. Chil.* v. 175.—Chili.
Bridgesii, *Kunze, in Linnaea*, xx. (1847) 35.—Chili.
calycina, *Benth. in DC. Prod.* x. 211.—Peruv.
cana, *Cav. Ic.* v. 27. t. 443. f. 2.—Chili.
candicans, Willd. ex Link, in Spreng. Jahrb. i. iii. (1820) 53 = argentea.
canescens, Humb. ex Link, l. c. 54 = hyssopifolia.

CALCEOLARIA :—
canescens, *Willd. ex Roem. & Schult. Syst. Mant.* i. 165.—Am. austr.
Cavanillesii, Pers. ex Steud. Nom. ed. I. 135 = Fothergillii.
Cavanillesii, *Phil. in Anal. Univ. Chil.* (1873) 532. —Chili.
cerasifolia, *Benth. in DC. Prod.* x. 218.—Peruv.
chelidonioides, *H. B. & K. Nov. Gen. et Sp.* ii. 378.— Ecuador.
chiloënsis, Lindl. Bot. Reg. t. 1476 = dentata.
chimboracensis, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 52 = saxatilis.
collina, *Phil. in Linnaea*, xxix. (1857-58) 30.—Chili.
concinna, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 51 = chelidonioides.
connata, Hook. ex R. Grah. in Edinb. N. Phil. Journ. (Sept. 1828) 372 = petiolaris.
corymbosa, *Ruiz & Pav. Fl. Per.* i. 14. t. 20.—Chili.
crassifolia, *Phil. in Anal. Univ. Chil.* (1873) 532.— Chili.
crenata, Bot. Reg. t. 790 = adscendens.
crenata, *Lam. Encyc.* i. 556.—Ecuador.
crenatiflora, *Cav. Ic.* v. 28. t. 446.—Ins. Chiloe.
cuneiformis, *Ruiz & Pav. Fl. Per.* i. 19. t. 27.— Ecuador.
Cunninghamii, *Vatke, in Linnaea*, xl. (1876) 222.— Chili.
Darwinii, *Benth. in DC. Prod.* x. 207.—Reg. Magellan.
deflexa, *Ruiz & Pav. Fl. Per.* i. 18. t. 30.—Ecuador.
deflexa, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 52 = padifolia.
densifolia, *Phil. in Linnaea*, xxxiii. (1864-65) 216.— Chili.
dentata, *Ruiz & Pav. Fl. Per.* i. 18. t. 29.—Chili.
denticulata, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 53 = stricta.
dichotoma, Lam. Illustr. i. 32 = ovata.
diffusa, Lindl. Bot. Reg. t. 1374 = bicolor.
dilatata, *Benth. in DC. Prod.* x. 211.—Ecuador.
divaricata, *H. B. & K. Nov. Gen. et Sp.* ii. 381.— Peruv.
elatior, *Griseb. in Goett. Abh.* xxiv. (1879) 237.—Reg. Argent.
elliptica, *Wedd. Chlor. And.* ii. 137.—Peruv. ; Boliv.
ericoides, *Juss. ex Vahl, Enum.* i. 190.—Ecuador.
erioclada, *Benth. in DC. Prod.* x. 215.—Ecuador.
excelsa, *Hort. ex Court. in Mag. Hortic.* i. (1833) 64. —Hab. ?
exoniensis ×, Hort. Angl. ex Steud. Nom. ed. II. i. 254.
extensa, *Benth. in DC. Prod.* x. 214.—Peruv.
fasciculata, *Willd. ex Link, in Spreng. Jahrb.* i. III. (1820) 53.—Peruv.
ferruginea, *Cav. Ic.* v. 27. t. 445. f. 1.—Chili.
ferruginea, Colla, in Mém. Acc. Tor. xxxviii. (1835) 137. t. 46 = integrifolia.
ferruginosa, Kunze, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 397 = integrifolia.
filicaulis, *Clos, in C. Gay, Fl. Chil.* v. 181.—Chili.
flexuosa, *Ruiz & Pav. Fl. Per.* i. 17, 26.—Peruv.
floribunda, *H. B. & K. Nov. Gen. et Sp.* ii. 385.— Ecuador.
floribunda, Hook. Bot. Mag. t. 4154 = crenata.
floribunda, Lindl. Bot. Reg. t. 1214 = petiolaris.
foliosa, *Griseb. in Goett. Abh.* xix. (1874) 210.—Reg. Argent.
foliosa, Meyen, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 395 = cuneiformis.
foliosa, *Phil. in Anal. Univ. Chil.* (1873) 531.—Chili.
Fothergillii, [*Soland. in*] *Ait. Hort. Kew.* ed. I. i. 30. t. 1.—Ins. Falkl.
fuchsiaefolia, *Hort.—Cf. Gard. Chron.* (1881) I. 269. —Hab. ?
Gellandana, Lindl. ex Steud. Nom. ed. II. i. 254 = scq.
Gellaniana, Penny, in Loud. Hort. Brit. Suppl. i. 583 = purpurea.
gemelliflora, *Cav. Ic.* v. 30. t. 450. f. 1.—Peruv.
glabrata, *Phil. in Linnaea*, xxix. (1857-58) 31.— Chili.
glacialis, *Wedd. Chlor. And.* ii. 143.—Bolivia.
glandulosa, Benth. in DC. Prod. x. 210 = pseudoglandulosa.
glandulosa, *Poepp. ex Benth. l. c.*—Chili.
glauca, *Ruiz & Pav. Fl. Per.* i. 17. t. 27.—Peruv.
glutinosa, *Heer & Regel, Ind. Sem. Hort. Turic.* (1848) 4.—*Cf. Linnaea*, xxiv. (1851) 196.—Guatemala.

CALCEOLARIA :—
gnidiifolia, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 54 = lavandulaefolia.
gossypina, *Benth. in DC. Prod.* x. 223.—N. Granat.
gracilis, *H. B. & K. Nov. Gen. et Sp.* ii. 379.— Ecuador.
gracilis, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 52 = ovata.
graminifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 386.— Ecuador.
guttata, *Willmore, ex Knowles & Westc. Flor. Cab.* i. (1837) 21.—Hab. ?
Hartwegii, *Benth. Pl. Hartw.* 147.—Peruv.
helianthemoides, *H. B. & K. Nov. Gen. et Sp.* ii. 387 (err. typ. helianthoides).—Peruv.
Henrici, *Hook. f. Bot. Mag.* t. 5772.—Ecuador.
Herbertiana, Lindl. Bot. Reg. t 1313 = racemosa.
heterophylla, *Ruiz & Pav. Fl. Per.* i. 16. t. 21.— Peruv.
heterophylla, Willd. Enum. Hort. Berol. 29 = scabiosaefolia.
hians, Penny, ex G. Don, Gen. Syst. iv. 603 = corymbosa ?
hispida, *Benth. in DC. Prod.* x. 212.—Peruv.
Hopeana, Morrison, ex G. Don, Gen. Syst. iv. 606 = corymbosa ?
Hoppeana, Morrison, ex Loud. Hort. Brit. Suppl. i. 583 = plantaginea.
hypericina, *Poepp. ex Benth. in DC. Prod.* x. 222.— Chili.
hypoleuca, *Benth. l. c.*—Ecuador.
hypoleuca, *Meyen, Reise*, i. 224 ; *Walp. & Schau. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 396.— Chili.
hyssopifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 386.— Ecuador.
inflexa, Ruiz & Pav. Fl. Per. i. 16. t. 25 = serrata.
integrifolia, Linn. f. Suppl. 86 = ovata.
integrifolia, *Murr. Syst.* ed. XIII. 61.—Chili.
involuta, *Ruiz & Pav. Fl. Per.* i. 15. t. 23.—Ecuador.
Kellyana, *Hort.—Cf. Gard. Chron.* (1883) II. 12.— Hab. ?
knypersliensis, Hort. ex Steud. Nom. ed. II. i. 254 = crenatiflora.
lamiifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 383.— Ecuador.
lanata, *H. B. & K. l. c.*—Ecuador.
lanceolata, *Cav. Ic.* v. 27. t. 144. f. 2.—Patagon.
lancifolia, Link, in Spreng. Jahrb. i. III. (1820) 5 = lamiifolia.
Landbeckii, *Phil. in Linnaea*, xxxiii. (1864-65) 215.— Chili.
lanigera, *Phil. in Anal. Univ. Chil.* (1884) reimpr. 9. —Chili.
latifolia, *Benth. in DC. Prod.* x. 212.—Chili.
lavandulaefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 386. —Ecuador.
laxa, *Benth. in DC. Prod.* x. 223.—Peruv.
linearis, *Ruiz & Pav. Fl. Per.* i. 19.—Peruv.
linifolia, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 54 = hyssopifolia.
lobata, *Cav. Ic.* v. 26. t. 443. f. 1.—Peruv.
Lorentzii, *Griseb. in Goett. Abh.* xix. (1874) 210.
Martineauae ×, Low, ex Sweet, Brit. Fl. Gard. Ser. II. t. 162.
Matthewsii, *Benth. in DC. Prod.* x. 207.—Peruv.
melissaefolia, *Benth. l. c.* 214.—Peruv.
mendocina, *Phil. in Anal. Univ. Chil.* (1862) II. 404. —Chili.
mexicana, *Benth. Pl. Hartw.* 47.—Mexic.
Meyeniana, *Phil. in Linnaea*, xxix. (1857-58) 32.— Chili.
mimuloides, *Clos, in C. Gay, Fl. Chil.* v. 174.— Chili.
mirabilis, Knowl. & Westc. Flor. Cab. i. (1837) 25 = crenatiflora.
mollissima, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 396 = integrifolia.
monanthos, Poir. Encyc. Suppl. ii. 24 = uniflora.
montana, *Cav. Ic.* v. 27. t. 444. f. 1.—Chili.
Morisii, *Walp. Rep.* iii. 157.—Chili.
Morrisoni, G. Don, in Sweet, Hort. Brit. ed. III. 521 = plantaginea.
multiflora, *Cav. Ic.* v. 29. t. 449.—Peruv.
Nahuelbutae, *Phil. in Anal. Univ. Chil.* (1873) 531. —Chili.
nana, *Sm. Ic. Ined.* i. 1. t. 1.—Reg. Magellan.
Neeana, Spreng. Syst. i. 44 = Fothergillii.

CALCEOLARIA :—
nitida, Colla, in Mem. Acc. Tor. xxxix. (1836) 138 = adscendens.
nivalis, *H. B. & K. Nov. Gen. et Sp.* ii. 381.—Chili.
nivalis, Steud. & Hochst. ex Benth. in DC. Prod. x. 219 = adscendens.
nudicaulis, *Benth. l. c.* 208.—Chili.
nudicaulis, Meyen, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 397 = plantaginea.
obovata, Stokes, Bot. Comment. i. 119 = Fothergillii.
oblonga, *Ruiz & Pav. Fl. Per.* i. 15. t. 24.—Peruv.
obtusifolia, Kunze, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 397 = plantaginea.
oligantha, *Phil. in Linnaea*, xxxiii. (1864-65) 216.— Chili.
ovata, *Sm. Ic. Ined.* i. 3. t. 3.—Peruv.
padifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 380.—Peruv.
Palpe, Steud. & Hochst. ex Benth. in DC. Prod. x. 219 = thyrsiflora.
panicea, Roem. & Schult. Syst. i. 181, sphalm. = punicea.
paniculata, Herb. ex G. Don, Gen. Syst. iv. 606 = petiolaris.
paniculata, Willd. ex Link, in Spreng. Jahrb. i. III. (1820) 53 = floribunda.
pannosa, *Phil. in Linnaea*, xxxiii. (1864-65) 219.— Chili.
paposana, *Phil. Fl. Atac.* 46.—Chili.
paralensis, Spreng. Syst. i. 45 = seq.
paralia, *Cav. Ic.* v. 29. t. 447.—Chili.
paralia, Hook. Exot. Fl. t. 75 = corymbosa.
pardanthera ×, W. Baxt. ex Loud. Hort. Brit. Suppl. ii. 616.
parviflora, *Gill. ex Benth. in DC. Prod.* x. 207.—Chili.
parvifolia, *Wedd. Chlor. And.* ii. 134.—Boliv.
Pavonii, *Benth. in DC. Prod.* x. 211.—Peruv.
pendula, Sweet, Brit. Flow. Gard. Ser. II. 155 = crenatiflora.
perfoliata, *Linn. f. Suppl.* 86.—N. Granat.
perfoliata, Ruiz & Pav. Fl. Per. i. 15. t. 21. f. a = Pavonii.
petioalaris, *Cav. Ic.* v. 30, 451.—Ecuador.
petiolaris, Vahl, Enum. i. 186 = pracc.
picta, *Phil. in Linnaea*, xxix. (1857-58) 34.—Chili.
pinifolia, *Cav. Ic.* v. 26. t. 442. f. 2.—Chili.
pinnata, *Linn. Mant.* 171.—Peruv.
pinnata, Ruiz & Pav. Fl. Per. i. 14. t. 19 = scabiosaefolia.
pisacomensis, *Meyen, Reise*, i. 469.—Peruv.
plantaginea, *Sm. Ic. Ined.* i. 2. t. 2.—Reg. Magellan.
plectranthifolia, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 394.—Peruv.
polifolia, *Hook. Bot. Mag.* t. 2897.—Chili.
polyantha, Penny, ex Loud. Hort. Brit. Suppl. i. 583 = petioalaris.
polyrrhiza, *Cav. Ic.* v. 25. t. 441.—Patagon.
pratensis, *Phil. in Linnaea*, xxix. (1857-58) 33.—Chili.
pseudoglandulosa, *Clos, in C. Gay, Fl. Chil.* v. 175.— Chili.
pulchella, Salisb. Prod. 89 = pinnata.
pulverulenta, *Ruiz & Pav. Fl. Per.* i. 19.—Peruv.
punctata, *Vahl, Enum.* i. 177.—Peruv.
puncticulata, *Phil. in Linnaea*, xxix. (1857-58) 36.— Chili.
punicea, *Ruiz & Pav. Fl. Per.* i. 15. t. 24.—Peruv.
purpurea, *R. Grah. in Bot. Mag.* t. 2775.—Peruv.
quadriradiata, *Phil. in Linnaea*, xxix. (1857-58) 31.— Chili.
racemosa, *Cav. Ic.* v. 29. t. 448.—Patagon.
refugens, Sweet, Brit. Flow. Gard. Ser. II. t. 227 = arachnoidea.
repens, *Hook. f. Fl. Nov. Zel.* i. 187.—N. Zel.
robusta, A. Dietr. in Otto & Dietr. Allg. Gartenz. i. (1833) 73 = integrifolia.
rosmarinifolia, *Lam. Encyc.* i. 556.—Ecuador.
rotundifolia, *H. B. & K. Nov. Gen. et Sp.* ii. 382.—Peruv.
rugosa, Hook. Exot. Fl. t. 99 = adscendens.
rugosa, Ruiz & Pav. Fl. Per. i. 19. t. 28 ; Lodd. Bot. Cab. t. 942 = integrifolia.
salicifolia, Colla, in Mem. Acc. Tor. xxxix. (1836) 139 = sessilis ?
salicifolia, *Ruiz & Pav. Fl. Per.* i. 18. t. 19.—Peruv.
salviaefolia, Pers. Syn. i. 17 = integrifolia.
saxatilis, *H. B. & K. Nov. Gen. et Sp.* ii. 383.— Ecuador.
scabiosaefolia, Nees, ex G. Don, Gen. Syst. iv. 606 = rugosa.
scabiosaefolia, Roem. & Schult. Syst. i. 187 = chelidonioides.

CALCEOLARIA :—
scabiosaefolia, *Sims, Bot. Mag.* t. 2405.—Peruv.
scabra, *Ruiz & Pav. Fl. Chil.* i. 19. t. 29.—Peruv.
scapiflora, *Benth. in DC. Prod.* x. 206.—Peruv.
Segethi, *Phil. in Linnaea*, xxix. (1857–58) 29.—Chili.
serrata, *Lam. Encyc.* i. 555.—Ecuador.
sessilis, *Ruiz & Pav. Fl. Per.* i. 18.—Chili.
sibthorpioides, *H. B. & K. Nov. Gen. et Sp.* ii. 382.—Peruv.
Sinclairii, *Hook. Ic. Pl.* t. 561.—N. Zel.
sparsiflora, *Kunze, in Linnaea*, xxiv. (1851) 228.—Boliv.
spectabilis, Kunze, ex Walp. & Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 396 = dentata.
Standishi ×, Paxt. Mag. ix. t. 75.
stricta, *H. B. & K. Nov. Gen. et Sp.* ii. 380.—N. Granat.
tenella, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 76. t. 287.—Chili.
tenera, *Phil. in Linnaea*, xxix. (1857–58) 34.—Chili.
tenuis, *Benth. in DC. Prod.* x. 205.—Chili.
terniflora, *Cav. Ic.* v. 30. t. 450. f. 2.—Peruv.
tetragona, *Benth. in DC. Prod.* x. 218.—Peruv.
tetraphylla, *Phil. in Linnaea*, xxix. (1857–58) 35.—Chili.
teucrioides, *Griseb. in Goett. Abh.* xix. (1874) 212.—Am. austr.
thyrsiflora, *R. Grah. in Edinb. N. Phil. Journ.* (Sept. 1828) 372 (err. typ. 272).—Chili.
tinctoria, Gill. ex G. Don, Gen. Syst. iv. 606 = arachnoidea.
tomentosa, *Ruiz & Pav. Fl. Per.* i. 15. t. 22.—Peruv.
tracheliifolia, Mart. & Gal. in Bull. Acad. Brux. xii. (1845) reimpr. 2 = mexicana.
triandra, *Vahl, Enum.* i. 181.—Peruv.
trifida, *Ruiz & Pav. Fl. Per.* i. 17. t. 26.—Peruv.
trilobata, *Hemsl. Biol. Centr. Am. Bot.* ii. 439.—Guatemala.
tripartita, *Ruiz & Pav. Fl. Per.* i. 14. t. 20.—Peruv.
umbellata, *Wedd. Chlor. And.* ii. 142.—Boliv.
undulata, *Benth. in DC. Prod.* x. 212.—Chili.
uniflora, Lam. Tabl. Meth. i. 52 = nana.
uniflora, *Ruiz & Pav. Fl. Per.* i. 16. t. 20.—Peruv.
utricularioides, *Hook. ex Benth. in DC. Prod.* x. 213.—Peruv.
valdiviana, *Phil. in Linnaea*, xxix. (1857–58) 35.—Chili.
verbascifolia, *Phil. l. c.* 30.—Chili.
verbascifolia, *Bert. in 'Merc. Chil.* 1829' *in Bull. Férussac*, xx. (1830) 109, nomen.—Chili.
vernicosa, *Phil. in Linnaea*, xxix. (1857–58) 29.—Chili.
verticillata, *Ruiz & Pav. Fl. Per.* i. 17.—Ecuador.
violacea, *Cav. Ic.* v. 31. t. 452.—Chili.
virgata, *Ruiz & Pav. Fl. Per.* i. 20. t. 31.—Peruv.
viscosa, *Ruiz & Pav. l. c.* 18. t. 30.—Peruv.
viscosissima, *Lindl. Bot. Reg.* t. 1611.—Chili.
Volckmanni, *Phil. in Linnaea*, xxxiii. (1864–65) 218.—Chili.
Wheeleri, Sweet, Brit. Flow. Gard. Ser. II. t. 130 = corymbosa.
Youngii ×, Lindl. Bot. Reg. t. 1448.

CALCEOLUS, [Tourn.] Adans. Fam. ii. 70 (1763) = **Cypripedium**, Linn. (Orchid.).
alternifolius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 62, 124 = C. Calceolus.
Marianus, Crantz, Stirp. Austr. ed. II. fasc. vi. 454 = C. Calceolus.

CALCITRAPA, Hall. Enum. Stirp. Helv. ii. 689 (1742) = **Centaurea**, Linn. (Compos.).
Adami, Schur, Enum. Pl. Transs. 409 = Cent. solstitialis.
aegyptiaca, Sweet, Hort. Brit. ed. I. 218, nomen = Cent. aegyptiaca.
altissima, Lam. Fl. Fr. ii. 31 = Microlonchus salmanticus.
aspera, Cass. in Dict. Sc. Nat. viii. 252 = Cent. aspera.
Balsamita, Sweet, Hort. Brit. ed. I. 218 = Cent. Balsamita.
benedicta, Sweet, l. c. = Carbenia benedicta.
brevispina, Moench, Meth. 563 = Microlonchus salmanticus.
Calcitrapa, Hill, Hort. Kew. 62 = Cent. Calcitrapa.
calcitrapoides, Hill. l. c. = Cent. calcitrapoides.
centauroides, Sweet, Hort. Brit. ed. I. 218 = Cent. centaurioides.
cichoracea, Lam. Fl. Fr. ii. 32 = Cent. Seridis.

CALCITRAPA :—
collina, Lam. Fl. Fr. ii. 33 = Centaurea collina.
conferta, Moench, Meth. 564 = Cent. melitensis.
cucullata, Moench, l. c. 563 = Cent. Crocodylium.
elongata, Moench, l. c. = Volutarella muricata.
eriophora, Moench, l. c. 564 = Cent. eriophora.
ferox, Moench, Meth. Suppl. 229 = Cent. ferox.
fusca, Moench, Meth. 564 = Cent. sicula.
Galactites, Lam. Fl. Fr. ii. 30 = Galactites tomentosa.
heterophylla, Moench, Meth. Suppl. 230 = Cent. aspera.
Hippophaestrum, Gaertn. Mey. & Scherb. Fl. Wett. iii. 178 = Cent. Calcitrapa.
hybrida, Sweet, Hort. Brit. ed. I. 218 = Cent. Calcitrapa.
Hypophaestum, Gaertn. Fruct. ii. t. 163 = Cent. Calcitrapa.
iberica, Trev. ex Spreng. Syst. iii. 406 = Cent. iberica.
lanceolata, Lam. Fl. Fr. ii. 34 = Cent. calcitrapoides.
lanuginosa, Steud. Nom. ed. II. i. 845 = Kentrophyllum lanatum.
lanuginosa, Lam. Fl. Fr. ii. 35 = Carbenia benedicta.
longifolia, Willk. & Lange, Prod. Fl. Hisp. ii. 141, sphalm. = Cent. sonchifolia.
lutea, Delarb. Fl. Auv. ed. II. 199 = Cent. solstitialis.
myacantha, Cass. in Dict. Sc. Nat. viii. 251 = Cent. Calcitrapa.
napifolia, Moench, Meth. 564 = Cent. napifolia.
nudicaulis, Lam. Fl. Fr. ii. 30 = Serratula nudicaulis.
parviflora, Lam. l. c. 32 = Cent. aspera.
patibilcensis, H. B. & K. Nov. Gen. et Sp. iv. 23 = Cent. americana.
phoenicea, Sweet, Hort. Brit. ed. I. 218 = Cent. calcitrapoides.
pubescens, Sweet, l. c. = Cent. incana.
reflexa, Sweet, l. c. = Cent. reflexa.
rigida, Sweet, l. c. = Cent. rigida.
rupestris, Sweet, l. c. = Cent. rupestris.
sessiliflora, Lam. Fl. Fr. ii. 35 = Cent. melitensis.
solstitialis, Lam. l. c. 34 = Cent. solstitialis.
sonchifolia, Lam. l. c. 32; Moench, Meth. 564 = Cent. sonchifolia.
sordida, Sweet, Hort. Brit. ed. I. 218 = Cent. sordida.
stellaris, Hill, Herb. Brit. i. 76 = Cent. Calcitrapa.
stellata, Lam. Fl. Fr. ii. 34 = Cent. Calcitrapa.
subspinosa, Moench, Meth. Suppl. 228 = Cent. diluta.
Verutum, Moench, Meth. 563 = Cent. Verutum.
vulgaris, Bernh. Syst. Verz. Erf. 158 = Cent. Calcitrapa.

CALCOA, Salisb. Gen. Pl. Fragm. 67 (1866) = **Geitonoplesium**, A. Cunn. (Liliac.).

CALDASIA, Lag. Amen. Nat. ii. 98 (1821) = **Oreomyrrhis**, Endl. (Umbellif.).
andicola, Lag. ex DC. Coll. Mém. v. t. 2 = O. andicola.
argentea, Hook. f. in Hook. Ic. Pl. iii. t. 300 = O. andicola.
brachycarpa, Hook. f. l. c. nota = O. andicola.
chaerophyllaea, DC. Prod. iv. 229 = O. andicola.
chaerophylloides, Lag. Amen. Nat. ii. 99 = O. andicola.
daucoides, Hook. f. Fl. Antarct. ii. t. 101 = O. andicola.
eriopoda, DC. Coll. Mem. v. 60 t. 2; et Prod. iv. 229 = O. andicola.
eriopus, DC. Coll. Mém. v. 75 = O. andicola.
lasiopetala, Lag. ex DC. Prod. iv. 229 = O. andicola.

CALDASIA, Mutis, ex Caldas, in Seman. Nuov. Gran. II. 26 (1810) = **Helosis**, Rich. (Balanoph.).
cayennensis, Mutis, ex Steud. Nom. ed. II. i. 255 = H. guyanensis.
fungiformis, Mutis, ex Eichl. in DC. Prod. xvii. 136 = H. guyanensis.

CALDASIA, Willd. Hort. Berol. t. 71 (1807) = **Bonplandia**, Cav. (Acanthac.).
heterophylla, Willd. l. c. = B. geminiflora.

CALDCLUVIA, D. Don, in Edinb. N. Phil. Journ. ix. (1830) 92. *SAXIFRAGEAE*, Benth. & Hook. f. i. 653.
DIETERICA, Ser. in DC. Prod. iv. 8 (1830).
paniculata, *D. Don, l. c.*—Chili.

CALDENBACHIA, Pohl, ex Nees, in DC. Prod. xi. 281 (1847) = **Stenandrium**, Nees (Acanthac.).
elegans, Pohl, ex Nees, l. c. 283 = S. Pohlii.
macrogranum, Nees, l. c. 282 = S. dulce.

CALDESIA, Parl. Fl. Ital. iii. 598 (1858) = **Alisma**, Linn.
parnassifolia, Parl. l. c. 599 = A. parnassifolium.

CALEA, Linn. Sp. Pl. ed. II. 1179 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 390.
ALLOCARPUS, H. B. & K. Nov. Gen. et Sp. iv. 291, t. 405 (1820).
ALLOISPERMUM, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 139.
AMPHICALEA, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 411.
CALEACTE, R. Br. in Trans. Linn. Soc. xii. (1818) 109.
CALEBRACHYS, Cass. in Dict. Sc. Nat. lv. 277 (1828).
CALYDERMOS, Lag. Gen. et Sp. Nov. 24 (1816).
LEMMATIUM, DC. Prod. v. 669 (1836).
LEONTOPHTHALMUM, Willd. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 140.
MEYERIA, DC. Prod. v. 670 (1836).
MOCINNA, Lag. Gen. et Sp. Nov. 31 (1816).
OTEIZA, La Llave, Reg. Trim. Mex. (1832) 41.
STENOPHYLLUM, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 391 (1873).
TETRACHYRON, Schlecht. in Linnaea, xix. (1847) 744.
acaulis, *Baker, in Mart. Fl. Bras.* vi. III. 266.—Bras.
aculeata, Labill. Nov. Holl. Pl. ii. 41. t. 185 = Cassinia aculeata.
albida, *A. Gray, in Proc. Am. Acad.* xv. (1880) 38.—Mexic.
Amellus, Linn. Sp. Pl. ed. II. 1179 = Salmea scandens.
angustifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 417 = multiplinervia.
angustifolia, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 256.—Bras.
aspera, Jacq. Collect. ii. 290 = Melananthera deltoidea.
axillaris, *DC. Prod.* v. 673.—Mexic.
Berteriana, *DC. l. c.* 672.—N. Granat.
brachiata, *DC. l. c.* 673.—Panama.
cacosmoides, *Less. in Linnaea*, v. (1830) 157.—Mexic.
Candolleana, *Baker, in Mart. Fl. Bras.* vi. III. 256.—Bras.
caroliniana, *Walt. Fl. Carol.* 200.—Am. bor.
Clausseniana, *Baker, in Mart. Fl. Bras.* vi. III. 265.—Bras.
clematidea, *Baker, l. c.* 262.—Bras.
collina, *Klatt, in Leopoldina*, xx. (1884) 95.—N. Granat.
cordata, *Lour. Fl. Cochinch.* ii. 488.—China.
cordifolia, Sw. Fl. Ind. Occ. iii. 1326 = jamaicensis.
coriacea, *DC. Prod.* v. 675.—Peruv.
crenata, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 256 = Candolleana.
cuneifolia, *DC. Prod.* v. 674.—Bras.
cymosa, *Less. in Linnaea*, v. (1830) 158.—Bras.
densiflora, *Klatt, in Leopoldina*, xx. (1884) 96.—Peruv.
discolor, *A. Gray, in Proc. Am. Acad.* xv. (1880) 38.—Mexic.
divaricata, *Benth. in Hook. Journ. Bot.* ii. (1840) 44.—Guiana.
divergens, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 262.—Bras.
elegans, *DC. Prod.* v. 674.—Mexic.
elongata, *Baker, in Mart. Fl. Bras.* vi. III. 255.—Bras.
eupatorioides, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 417 = serrata.
ferruginea, *Baker, in Mart. Fl. Bras.* vi. III. 261.—Bras.
fruticosa, Benth. & Hook. f. Gen. ii. 390 = rotundifolia.
Gardneriana, *Baker, in Mart. Fl. Bras.* vi. III. 255.—Bras.
gentianoides, DC. Prod. v. 672 = Geissopappus gentianoides.
glabra, *DC. l. c.* 674.—Bras.
Grayii, *Klatt, in Leopoldina*, xx. (1884) 96.—Peruv.
hispida, *Baker, in Mart. Fl. Bras.* vi. III. 261.—Bras.
hymenolepis, *Baker, l. c.* 258.—Bras.
hypericifolia, *Baker, l. c.*—Bras.
integrifolia, *Hemsl. Biol. Centr. Am. Bot.* ii. 205.—Mexic.
jamaicensis, *Linn. Sp. Pl.* ed. II. 1179.—Jamaica.
lantanoides, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 416.—Bras.
lemmatioides, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 252.—Bras.

CALEA :—
Leontophthalmum, *Less. Syn. Comp.* 241.—Peruv.
leptophylla, Forst. f. Prod. 55 = Cassinia leptophylla.
lobata, Sw. Prod. Veg. Ind. Occ. 113 = Neurolaena lobata.
longifolia, *Baker, in Mart. Fl. Bras.* vi. III. 260.—Bras.
longifolia, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 418 = multiplinervia.
manicata, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 206.—Mexic.
Martiana, *Baker, in Mart. Fl. Bras.* vi. III. 256.—Bras.
martinicensis, *Hort. ex Steud. Nom.* ed. II. i. 255.—Ins. Martinic.
melissaefolia, *Baker, in Mart. Fl. Bras.* vi. III. 257.—Bras.
microphylla, *Baker, l. c.* 259.—Bras.
montana, *Klatt, in Leopoldina*, xx. (1884) 96.—Peruv.
multiplinervia, *Less. in Linnaea*, v. (1830) 159.—Bras.
myrtifolia, *Benth. & Hook. f. Gen.* ii. 391; *Baker, in Mart. Fl. Bras.* vi. III. 259.—Bras.
nitida, *Less. in Linnaea*, v. (1830) 158.—Bras.
oligocephala, *DC. Prod.* v. 675.—Bras.
oppositifolia, Linn. Sp. Pl. ed. II. 1179 = Isocarpha oppositifolia.
oxylepis, *Baker, in Mart. Fl. Bras.* vi. III. 354.—Bras.
parvifolia, *Baker, l. c.* 259.—Bras.
paucidentata, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 391 = myrtifolia.
pedunculares, *H. B. & K. Nov. Gen. et Sp.* iv. 295. t. 408.—Mexic.
pedunculosa, DC. Prod. v. 673 = uniflora.
perforata, *Klatt, in Leopoldina*, xx. (1884) 95.—N. Granat.
phyllolepis, *Baker, in Mart. Fl. Bras.* vi. III. 260.—Bras.
pilosa, *Baker, l. c.* 257.—Bras.
pinifolia, Forst. f. Prod. 55. n. 288 = Ozothamnus pinifolius.
pinnatifida, *Banks, ex Steud. Nom.* ed. I. 136.—Bras.
platylepis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 267.—Bras.
Pohliana, *Sch. Bip. ex Baker, l. c.* 266.—Bras.
prunifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 294. t. 406.—N. Granat.
ramosissima, *Baker, in Mart. Fl. Bras.* vi. III. 257.—Bras.
reticulata, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 416.—Bras.
rotundifolia, *Benth. & Hook. f. Gen.* ii. 390; *Baker, in Mart. Fl. Bras.* vi. III. 253.—Bras.
rugosa, *Hemsl. Biol. Centr. Am. Bot.* ii. 206.—Mexic.
sabazioides, *Hemsl. l. c.*—Mexic.
salmiaefolia, *Hemsl. l. c.*—Mexic.
scaberrima, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 255 = elongata.
scabrifolia, *Benth. & Hook. f. Gen.* ii. 391.—Mexic.
scandens, *Griseb. in Goett. Abh.* xxiv. (1879) 199.—Reg. Argent.
scoparia, Linn. Syst. Nat. ed. XII. 234 = Baccharis scoparia.
senecioides, *Baker, in Mart. Fl. Bras.* vi. III. 258.—Bras.
Senecionis, Benth. & Hook. f. Gen. ii. 391 = praec.
serrata, F. G. Dietr. Vollst. Lex. Gaertn. i. 642 = cordifolia.
serrata, *Less. in Linnaea*, v. (1830) 158.—Bras.
sessiliflora, *Less. l. c.*—Mexic.
sessiliflora, Stokes, Bot. Mat. Med. iv. 172 = Pacourina edulis.
solidaginea, *H. B. & K. Nov. Gen. et Sp.* iv. 295.—Venezuela; Ind. occ.
spectabilis, Labill. Nov. Holl. Pl. i. 42. t. 186 = Apalochlamis spectabilis.
stenophylla, *Baker, in Mart. Fl. Bras.* vi. III. 255.—Bras.
subrotunda, Gardn. in Hook. Lond. Journ. Bot. vii. (1848) 415 = lantanoides.
Suriani, Cass. in Dict. Sc. Nat. vi. Suppl. 32 = Neuroleana integrifolia.
terrifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 294.—N. Granat.
teucriifolia, *Baker, in Mart. Fl. Bras.* vi. III. 259.—Bras.
tomentosa, *Gardn. in Hook. Lond. Journ. Bot.* vii. (1848) 418.—Bras.

CALEA :—
tomentosa, *A. Gray, in Proc. Am. Acad.* xv. (1880) 38.—Mexic.
trifida, Moench, Meth. 581 = jamaicensis.
uniflora, *Less. in Linnaea*, v. (1830) 159.—Bras.
urticaefolia, *DC. Prod.* v. 674.—Mexic.
verbenaefolia, *DC. l. c.* 673.—Am. trop.
villosa, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. III. 262.—Bras.
Zacatechichi, *Schlecht. in Linnaea*, ix. (1834) 589.—Mexic.

CALEA, Sw. Prod. Veg. Ind. Occ. 113 (1788) = **Neurolaena**, R. Br. (Compos.).
lobata, Sw. l. c. = N. lobata.

CALEACTE, R. Br. in Trans. Linn. Soc. xii. (1818) 109 = **Calea**, Linn. (Compos.).
pinnatifida, R. Br. l. c. = Calea pinnatifida.
rotundifolia, Less. in Linnaea, v. (1830) 158 = Calea rotundifolia.
urticifolia, R. Br. in Trans. Linn. Soc. xii. (1818) 109 = Calea urticaefolia.

CALEANA, R. Br. Prod. 329 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 610.
CALEYA, R. Br. in Ait. Hort. Kew. ed. II. v. 204 (1813).
major, *R. Br. Prod.* 329.—Austral.
minor, *R. Br. l. c.*—Austral.
nigrita, *Lindl. Swan Riv. App.* 54.—Austral.

CALEBRACHYS, Cass. in Dict. Sc. Nat. lv. 277 (1828) = **Calea**, Linn. (Compos.).
pedunculares, Cass. l. c. = Calea pedunculares.

CALECTASIA, R. Br. Prod. i. 263 (1810); et in Flind. Voy. ii. 609. t. 9 (1814). *JUNCACEAE*, Benth. & Hook. f. iii. 866.
HUTTIA, Preiss, ex Hook. Bot. Mag. t. 3834 (1840).
cyanea, *R. Br. Prod.* 264.—Austral.
grandiflora, Preiss, in Lehm. Pl. Preiss. ii. 53 = cyanea.
intermedia, Sond. in Linnaea, xxviii. (1856) 222 = cyanea.

CALENDULA, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 454.
CALTHA, Tourn. ex Adans. Fam. ii. 126 (1763).
aegyptiaca, *Desf. Tabl. Hort. Par.* ed. I. 100.—Reg. Mediterr. or.; Mesopot.
algarbiensis, Boiss. Diagn. Ser. II. vi. 106 = suffruticosa.
algeriensis, *Boiss. & Reut. in Boiss. l. c.* 109.—Algeria.
americana, *Mill. Gard. Dict.* ed. VIII. n. 10.—Mexic.
amplexicaulis, Thunb. Prod. Pl. Cap. 164 = Tripteris amplexicaulis.
amplexifolia, Reichb. in Flora, xiii. (1830) 382 = arvensis.
anemoneflora, *Brcit. ex DC. Prod.* vi. 454, nomen.—Hab.?
arborescens, Jacq. Collect. iii. 167 = Tripteris arborescens.
arragonica, *Fisch. ex Steud. Nom.* ed. II. i. 256.—Hispan.
arvensis, Bieb. Fl. Taur. Cauc. iii. 596 = persica.
arvensis, Boiss. Voy. Bot. Espagne, ii. 338 = aegyptiaca.
arvensis, *Linn. Sp. Pl.* ed. II. 1303.—Europ.; Oriens.
aspera, Thunb. Prod. Pl. Cap. 164 = Osteospermum asperum.
Asterias, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 31 = stellata.
aurantiaca, Kotschy, ex Boiss. Diagn. Ser. I. vi. 96 = sinuata.
aurea, *Hort. ex Steud. Nom.* ed. II. i. 256.—Afr. austr.
Balansae, *Boiss. & Reut. Diagn.* Ser. II. vi. 107.—Hispan.
bicolor, Rafin. Caratt. 82. n. 199 = arvensis.
brachyglossa, Rupr. in Ind. Sem. Hort. Petrop. (1852); ex Bull. Phys.-Math. Acad. Pétersb. xiv. (1856) 230 = arvensis.
byzantina, DC. Prod. vi. 453 = arvensis.
cariensis, *Boiss. Diagn.* Ser. I. x. 82.—As. Min.
ceratosperma, Viv. Fl. Lib. Spec. 59. t. 20 = palestina.
chrysanthemifolia, Vent. Jard. Malm. 56 = Dimorphotheca chrysanthemifolia.
cornuta, Poir. Encyc. vii. 277 = stellata.

CALENDULA :—
Cristagalli, Viv. Fl. Lib. Spec. 59. t. 26. f. 2 = sicula, stellata.
cuneata, Thunb. Prod. Pl. Cap. 164 = Dimorphotheca cuneata.
Dalgesirum, Hort. ex DC. Prod. vi. 452 = marginata.
decumbens, *Mill. Gard. Dict.* ed. VIII. n. 9.—Mexic.
decurrens, Thunb. Prod. Pl. Cap. 163 = Dimorphotheca pluvialis.
dentata, Hort. ex Steud. Nom. ed. I. 136 = denticulata.
denticulata, *Schousb. in Willd. Enum. Hort. Berol.* 936.—Maurit.
diffusa, Salisb. Prod. 209 = Dimorphotheca fruticosa.
echinata, DC. Prod. vi. 453 = arvensis.
eriocarpa, *DC. l. c.*—Hab.?
flaccida, Vent. Jard. Malm. 20 = Dimorphotheca aurantiaca.
Forskohleana, Hort. ex Steud. Nom. ed. II. i. 256 = stellata.
fruticosa, Hort. ex DC. Prod. vi. 452 = suffruticosa.
fruticosa, Linn. Pl. Afr. Rar. 25 (Amoen. Acad. v. 25) = Dimorphotheca fruticosa.
fulgida, Rafin. Caratt. 83. n. 200 = suffruticosa.
glabrata, Thunb. Prod. Pl. Cap. 163 = Osteospermum glabratum.
glauca, *Salisb. Prod.* 209.—Hab.?
gracilis, DC. Prod. vi. 453 = persica.
graminifolia, Linn. Sp. Pl. 922 = Dimorphotheca graminifolia.
grandiflora, *Hort. ex Steud. Nom.* ed. II. i. 256.—Afr. austr.
hispanica, Hort. ex DC. Prod. vi. 452 = suffruticosa.
hispida, Thunb. Prod. Pl. Cap. 164 = Aster hispidus.
hybrida, Linn. Sp. Pl. 921 = Dimorphotheca hybrida.
hybrida, Sweet, Brit. Flow. Gard. i. t. 39 = Dimorphotheca pluvialis.
incana, Willd. Sp. Pl. iii. 2341 = suffruticosa.
incanescens, *Schrad. ex Steud. Nom.* ed. II. i. 256.—Hab.?
linifolia, *Wendl. ex DC. Prod.* vi. 454, nomen.—Hab.?
lusitanica, Boiss. Diagn. Ser. I. x. 83 = suffruticosa.
maderensis, *DC. Prod.* vi. 454.—Ins. Madera.
magellanica, Willd. Sp. Pl. iii. 2344 = Lagenophora Commersonii.
malacitana, Boiss. & Reut. Pugill. (1852) 61 = aegyptiaca.
malvaecarpa, *Pomel, Nouv. Mat. Fl. Atl.* 33.—Algeria.
marginata, Willd. Enum. Hort. Berol. 935 = suffruticosa.
maritima, *Guss. Ind. Sem. Hort. Bocc.* (1825).—Sicilia.
maritima, Lowe, ex DC. Prod. vi. 454 = maderensis.
maroccana, *Ball, in Journ. Bot.* xi. (1873) 365.—Marocco.
micrantha, Boiss. & Noé, Diagn. Ser. II. vi. 111 = aegyptiaca.
micrantha, Tineo & Guss. Fl. Sic. Syn. ii. 874 = arvensis.
microcephala, Kral. ex Reichb. Ic. Fl. Germ. xv. 99. t. 891 = aegyptiaca.
microcephala, Lange, ex Ficalho, in Bol. Soc. Broter. i. (1883) 51 = suffruticosa?
microphyla, *Lange, ex Ficalho, l. c.*—Hispan.
Monardi, *Boiss. & Reut. Diagn.* Ser. II. vi. 108.—Hispan.
monstrosa, Burm. f. Fl. Cap. Prod. 28 = Tripteris scariosa.
muricata, Thunb. Prod. Pl. Cap. 164 = Tripteris arborescens.
Nočana, Boiss. Fl. Orient. iii. 416 = suffruticosa.
nudicaulis, Linn. Sp. Pl. 922 = Dimorphotheca nudicaulis.
officinalis, Hohen. in Bull. Soc. Nat. Mosc. (1833) 256 = persica.
officinalis, *Linn. Sp. Pl.* 921.—Europ. austr.
oppositifolia, Ait. Hort. Kew. iii. 272 = Dimorphotheca oppositifolia.
palestina, *Boiss. Diagn.* Ser. I. x. 83.—Palestin.; Mesopotam.
parviflora, DC. partim, ex Boiss. Fl. Orient. iii. 419 = aegyptiaca.
parviflora, Rafin. Caratt. 83. n. 201 = stellata.
parviflora, Thunb. Prod. Pl. Cap. 163 = Oligocarpus calendulaceus.

CALENDULA:—
persica, *C. A. Mey. Verz. Pfl. Cauc.* 72.—Reg. Caucas.; Persia; Afghan.
pinnata, Thunb. Prod. Pl. Cap. 164 = Chrysanthemum lidbeckioides.
platycarpa, Coss. in Bull. Soc. Bot. Fr. iii. (1856) 655 = stellata.
pluvialis, Linn. Sp. Pl. 921 = Dimorphotheca pluvialis.
pluvialis, Thunb. Fl. Cap. 704 = Dimorphotheca clandestina.
prolifera, Hort. ex Steud. Nom. ed. II. i. 256 = officinalis.
pumila, Forst. f. Prod. 57 = Lagenophora Commersonii.
pusilla, Thou. Fl. Trist. Acun. 40 = Lagenophora Commersonii.
repanda, Boiss. & Noé, Diagn. Ser. II. vi. 110 = palestina.
rigida, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 271 = Tripteris arborescens.
rosmarinifolia, Houtt. Handl. xi. 84 = Osteospermum polygaloides.
sancta, Linn. Sp. Pl. ed. II. 1304 = aegyptiaca?
scabra, Berg. Desc. Pl. Cap. 310 = Dimorphotheca pluvialis.
sicula, DC. Prod. vi. 452 (pl. aegypt.) = aegyptiaca.
sicula, Cyrill. ex Balb. Hort. Taur. (1800) = stellata.
sicula, Poir. Encyc. vii. 277 = praec.?
sinuata, *Boiss. & Gaill. Diagn.* Ser. II. vi. 109.—Syria; Persia.
spatulata, Hoffmgg. Verz. Pfl. Nachtr. iii. 24 = stellata.
stellata, *Cav. Ic.* i. 3. t. 5.—Reg. Mediterr.
stellata, Lowe, ex DC. Prod. vi. 454 = maderensis.
stipitata, Spreng. Syst. iii. 624 = Lagenophora Billardieri.
subinermis, *Pomel, Nouv. Mat. Fl. Atl.* 34.—Algeria.
sublanata, Reichb. f. Ic. Fl. Germ. xv. 99. t. 891 = arvensis.
suffruticosa, *Vahl, Symb. Bot.* ii. 94.—Reg. Mediterr. occ.
thapsiaecarpa, *Pomel, Nouv. Mat. Fl. Atl.* 34.—Algeria.
tomentosa, Desf. Fl. Atlant. ii. 305. t. 245 = suffruticosa.
tomentosa, Thunb. Prod. Pl. Cap. 163 = Tripteris tomentosa.
Tragus, Ait. Hort. Kew. iii. 271; Jacq. Hort. Schoenb. ii. 14. t. 153 = Dimorphotheca Tragus.
Tragus, Curt. Bot. Mag. t. 408 = Dimorphotheca aurantiaca.
tripterocarpa, *Rupr. in Ind. Sem. Hort. Petrop.* (1853); *ex Bull. Phys.-Math. Acad. Pétersb.* xiv. (1856) 231.—Mesopot.
undulata, J. Gay, ex Gaud. Fl. Helv. v. 417 = arvensis.
undulata, Rafin. Caratt. 81. n. 198 = stellata.
versicolor, Salisb. Prod. 209 = Dimorphotheca hybrida.
viscosa, Andr. Bot. Rep. t. 412 = Dimorphotheca cuneata.

CALEPINA, Adans. Fam. ii. 423 (1763); DC. Syst. Veg. ii. 648. *CRUCIFERAE*, Benth. & Hook. f. i. 96.
cochlearioides, Dum. Fl. Belg. 121 = Bunias cochlearioides.
Corvini, *Desv. Journ. Bot.* iii. (1814) 158.—Europ. med.; Reg. Caucas.; Persia.
prostrata, Spreng. Syst. ii. 850 = Muricaria prostrata.

CALESIA, Rafin. Princ. Somiol. 29 (1814) = seq.

CALESIAM, Adans. Fam. ii. 446 (1763) = **Odina**, Roxb. (Anacard.).

CALETIA, Baill. Etud. Gén. Euph. 553. t. 26 (1858) = **Micrantheum**, Desf. (Euphorbiac.).
divaricatissima, Muell. Arg. in Linnaea, xxxii. (1863) 79 = Pseudanthus divaricatissimus.
hexandra, Muell. Arg. in DC. Prod. xv. II. 194 = M. hexandrum.
linearis, Muell. Arg. in Linnaea, xxxii. (1863) 79 = Pseudanthus orientalis.
micrantheoides, Baill. Etud. Gén. Euph. 554. t. 26. f. 1 = M. hexandrum.
orientalis, Baill. Adansonia, vi. (1865-66) 327 = Pseudanthus orientalis.

CALETIA:—
ovalifolia, Muell. Arg. in Linnaea, xxxiv. (1865-66) 55 = Pseudanthus ovalifolius.
Wilhelmi, F. Muell. ex Muell. Arg. in DC. Prod. xv. II. 194 = Pseudanthus ovalifolius.

CALEYA, R. Br. in Ait. Hort. Kew. ed. II. v. 204 (1813) = **Caleana**, R. Br. (Orchid.).
major, R. Br. l. c. = Caleana major.
minor, Sweet, Hort. Brit. ed. I. 385 = Caleana minor.
Sullivani, F. Muell. in Melb. Chemist (1882) 44.—Austral.

CALIBRACHOA, Cerv. in La Llave & Lex. Nov. Veg. Desc. fasc. ii. (1825) 3 = **Petunia**, Juss. (Solanac.).
mexicana, Lex. l. c. = P. parviflora.
procumbens, Cerv. l. c. = P. viscidula?

CALICERA, Cav. Ic. iv. 34. t. 358 (1797) = **Calycera**, Cav.

CALICOREMA, Hook. f. in Benth. & Hook. f. Gen. iii. 34 (1880). *AMARANTACEAE*, Benth. & Hook. f. l. c.
capitata, *Hook. f. l. c.* 35.—Afr. austr.

CALICOTOME, Link, in Schrad. Neues Journ. ii. (1808) II. 50 = **Calycotome**, Link (Legumin.).

CALIGULA, Klotzsch, in Linnaea, xxiv. (1851) 28 = **Agapetes**, G. Don (Vacciniac.).
odontocera, Klotzsch, l. c. = A. variegata.
pulcherrima, Klotzsch, l. c. = A. variegata.

CALIMERIS, Nees, Gen. et Sp. Ast. 225 (1833) = **Aster**, Tourn. (Compos.).
Alberti, Regel, Gartenfl. (1884) 130.—Turkestan.
altaica, Nees, ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. viii. 52 = A. altaicus.
alyssoides, DC. Prod. v. 259 = A. alyssoides.
amplexifolia, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 182 = A. trinervius.
biennis, Ledeb. Fl. Ross. ii. 483 = Heteropappus hispidus.
biennis, Nees, Gen. et Sp. Ast. 231 = A. altaicus.
canescens, Nees, l. c. 229 = A. altaicus.
cassiarabica, G. Don, in Sweet, Hort. Brit. ed. III. 343 = Aster cassiarabicus.
chinensis, Sch. Bip. in Flora, xxxv. (1852) 48 = A. altaicus.
ciliata, A. Gray, in Mem. Am. Acad. N. S. vi. (1858-59) 394 = A. altaicus.
ciliosa, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 61 = A. altaicus.
cordata, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 126, nomen.—Malaya.
coronata, Sch. Bip. in Flora, xxxv. (1852) 48 = A. altaicus.
distortus, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) II. 152 = A. altaicus.
exilis, Lindl. ex DC. Prod. v. 259 = A. altaicus.
fastigiata, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 126 = A. fastigiatus.
flexuosa, Royle, ex Lindl. in DC. Prod. v. 258 = A. Thomsoni.
hispida, Lindl. l. c. 259 = A. altaicus.
hispida, Nees, Gen. et Sp. Ast. 227 = Heteropappus hispidus.
incisa, DC. Prod. v. 258 = A. incisus.
indica, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 125 = Boltonia indica.
integrifolia, Turcz. in DC. Prod. v. 259.—Reg. Amur.
japonica, Sch. Bip. in Zoll. Syst. Verz. Ind. Archip. 126 = A. fastigiatus.
microcephala, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 169.—Japon.
platycephala, Nees, Gen. et Sp. Ast. 226 = A. incisus.
rosea, DC. Prod. v. 259 = A. roseus.
tatarica, Lindl. ex DC. Prod. v. 259 = Heteropappus hispidus.

CALINEA, Aubl. Pl. Gui. i. 556. t. 221 (1775) = **Doliocarpus**, Boland. (Dilleniac.).
scandens, Aubl. l. c. = D. Calinea.

CALINUX, Rafin. in Med. Repos. New York, v. (1808) 352 = **Pyrularia**, Michx. (Santalac.).

CALIPHRURIA, Herb. in Bot. Reg. (1844) Misc. 87. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 731. Hartwegiana, *Herb. l. c.*—N. Granat.
subedentata, Baker. in Bot. Mag. t. 6289 = Eucharis subedentata.

CALIPOGON, Rafin. Atl. Journ. 148 (1832) = **Calopogon**, R. Br. (Orchid.).

CALISAYA, Hort. ex Pav. Nuev. Quinol. sub t. 5. p. 6 (1862); Howard, Quinol. E. Ind. 86 (1869) = **Cinchona**, Linn. (Rubiac.).
anglica ×, Hort. ex Howard, l. c. 87.
javanica, Hort. ex Howard, l. c. 86 = Cinch. Calisaya.

CALISPERMUM, Lour. Fl. Cochinch. 157 (1790) = **Embelia**, Juss. (Myrsin.).
scandens, Lam. l. c. = E. Ribes.

CALISTA, Ritg. in Marb. Ges. Schrift. ii. 128 (1831) = CALLISTA, Lour. = **Dendrobium**, Sw. (Orchid.).

CALISTACHYA, Rafin. in Med. Repos. New York, v. (1808) 352 = **Veronica**, Tourn. (Scrophular.).
alba, Rafin. l. c. = V. virginica.

CALIUS, Blanco, Fl. Filip. ed. I. 698 (1837) = **Streblus**, Lour. (Urticac.).
lactescens, Blanco, l. c. = S. asper.

CALIXNOS, Rafin. Fl. Tellur. iv. 82 (1836) (quid?).
trinervius, Rafin. l. c. = Convolvulus trinervis; 'non Convolvulacea,' sec. Choisy.

CALLA, Linn. Syst. ed. I (1735). *AROIDEAE*, Benth. & Hook. f. iii. 989.
CALLARIA, Rafin. in Am. Monthly Mag. (1818) 267.
HOUTTINIA, Neck. Elem. iii. 291 (1790).
PROVENZALIA, Adans. Fam. ii. 469 (1763).
aethiopica, Gaertn. Fruct. ii. 20. t. 84 = palustris.
aethiopica, Linn. Sp. Pl. 968 = Zantedeschia aethiopica.
ambigua, Salisb. Prod. 262 = Zantedeschia aethiopica.
angustifolia, Jack, in Malay. Misc. i. (1820) n. 1. p. 24 = Chamaecladon angustifolium.
aromatica, Roxb. Hort. Beng. 65; Fl. Ind. iii. 513 = Homalomena aromatica.
Badian, Blanco, Fl. Filip. ed. I. 658 = Alocasia indica.
calyptrata, Roxb. Hort. Beng. 65; Fl. Ind. iii. 514 = Schismatoglottis calyptrata.
cordifolia, Stokes, Bot. Mat. Med. iv. 326 = palustris.
Dracontium, G. F. W. Mey. Prim. Fl. Esseq. 197 = Monstera pertusa.
Gaby, *Blanco, Fl. Filip.* ed. I. 659.—Ins. Philipp.
humilis, Jack, in Malay. Misc. i. (1820) n. 1. p. 22 = Chamaecladon humile.
maxima, *Blanco, Fl. Filip.* ed. I. 658.—Ins. Philipp.
montana, Blume, Cat. Gew. Buitenz. 62 = Anadendrum montanum *vel* Rhaphidophora montana.
moschata, Moench, Meth. 346 = Richardia elliptica.
nitida, Jack, in Malay. Misc. i. (1820) n. 1. p. 24 = Aglaonema nitidum.
oblongifolia, Roxb. Hort. Beng. 65; Fl. Ind. iii. 516 = Aglaonema marantifolium.
occulta, Lodd. Bot. Cab. t. xii. = Homalomena aromatica.
occulta, Lour. Fl. Cochinch. ii. 532 = Zantedeschia occulta.
oculata, Lindl. in Gard. Chron. (1859) 788 = Richardia melanoleuca.
orientalis, Linn. Sp. Pl. ed. II. 1373 = Biarum Bovei.
ovatofolia, Gilib. Exercit. ii. 452 = palustris.
palustris, *Linn. Sp. Pl.* 968.—Europ.; As. bor.; Am. bor.
pertusa, Kunth, Syn. i. 129 = Monstera pertusa.
picta, Roxb. Hort. Beng. 65; Fl. Ind. iii. 516 = Aglaonema pictum.
polyphylla, *Blanco, Fl. Filip.* ed. I. 659.—Ins. Philipp.
radicans, Ruiz & Pav. ex Engl. in DC. Monog. Phan. ii. 377 = Philodendron deltoideum.
rubescens, Roxb. Hort. Beng. 65; Fl. Ind. iii. 515 = Homalonema rubescens.
sagittifolia, Michx. Fl. Bor. Am. ii. 187 = Peltandra alba.
sylvestris, Blume, Cat. Gew. Buitenz. 62 = Rhaphidophora angustifolia.
virginica, Michx. Fl. Bor. Am. ii. 187 = Peltandra virginica.
virosa, Roxb. Hort. Beng. 65; Fl. Ind. iii. 517 = Colocasia virosa.

CALLAEOCARPUS, Miq. Pl. Jungh. 13 (1851-55) = **Castanopsis**, Spach (Cupulif.).
rhamnifolia, Miq. Fl. Ind. Bat. Suppl. 353 = Cast. rhamnifolia.
sumatrana, Miq. Pl. Jungh. 14 = Cast. sumatrana.

CALLAEOLEPIUM, Karst. Fl. Colomb. ii. 123. t. 165 (1869?) = **Fimbristemma**, Turcz. (Asclepiad.).
Warscewiczii, Karst. l. c. = F. Warscewiczii.

CALLANTHUS, Reichb. Consp. 44 (1828) = **Watsonia**, Mill. (Irid.).

CALLARIA, Rafin. in Am. Monthly Mag. (1818) 267 = **Calla**, Linn. (Aroid.).

CALLERYA, Endl. Gen. Suppl. iii. 104 (1843) = **Milletia**, Wight & Arn. (Legumin.).
tomentosa, Endl. l. c. = M. nitida.

CALLIACHYRIS, Torr. & Gray, in Journ. Bost. Soc. Nat. Hist. v. (1844) 110 = **Layia**, Hook. (Compos.).
Fremontii, Torr. & Gray, l. c. = L. Fremontii.

CALLIANASSA, Webb & Berth. Phyt. Canar. iii. 143 (1836-50) = **Isoplexis**, Lindl. (Scrophularin.).
canariensis, Webb & Berth. l. c. 144 = I. canariensis.
Isabelliana, Webb & Berth. l. c. 145.—Ins. Canar.

CALLIANDRA, Benth. in Hook. Journ. Bot. ii. (1840) 138. *LEGUMINOSAE*, Benth. & Hook. f. i. 596.
ANNESLIA, Salisb. Parad. Lond. t. 64 (1807), nomen prius.
CLELIA, Casar. Nov. Stirp. Dec. 83 (1842).
CODONANDRA, Karst. Fl. Colomb. ii. 43. t. 122 (1862?).
abbreviata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 108.—Bras.
aculeata, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 541.—Ecuador.
aeschynomenoides, *Benth. in Mart. Fl. Bras.* xv. II. 415.—Am. trop.
alba, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = portoricensis.
alternans, *Benth. in Trans. Linn. Soc.* xxx. (1875) 548.—Madag.
amazonica, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 94.—Bras.
angelica, *Benth. l. c.* 100.—Mexic.
angustifolia, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 539.—Am. trop.
asplenioides, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = mertensioides.
axillaris, *Benth. in Mart. Fl. Bras.* xv. II. 415.—Bras.
bella, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 110.—Am. trop.
bicolor, *Benth. in Hook. Journ. Bot.* ii. (1840) 139.—Bras.
Blancheti, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 102.—Bras.
bombycina, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 538.—Am. trop.
bracteosa, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 104.—Bras.
brevicaulis, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 61.—Paraguay.
brevipes, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.—Bras.
californica, *Benth. Bot. Voy. Sulph.* 14. t. 11.—Calif.
Callistemon, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) 527.—Mexic.
calothyrsus, *Meissn. in Linnaea*, xxi. (1848) 251.—Guiana.
calycina, *Benth. in Mart. Fl. Bras.* xv. II. 419.—Am. trop.
canescens, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 96.—Mexic.
capillata, *Benth. l. c.* 98.—Mexic.
caracasana, *Benth. in Trans. Linn. Soc.* xxx. (1875) 543.—Am. trop.
carbonaria, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 95.—N. Granat.
Chamaedrys, Engelm. ex A. Gray, Pl. Fendl. 39 = eriophylla.
chilensis, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 103.—Chili.

CALLIANDRA:—
clavellina, Karst. Fl. Colomb. i. 159. t. 79 = Purdiaei.
Codonandra, *Benth. in Trans. Linn. Soc.* xxx. (1875) 547.—Am. trop.
collctioides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 180.—Cuba.
comosa, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 104.—Am. trop.
conferta, *Benth. in A. Gray, Pl. Wright.* i. 63.—Am. bor. occ.
coriacea, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 95 = cmarginata.
corocensis, Karst. Fl. Colomb. ii. 41 = purpurea.
Coulteri, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 352.—Am. bor. occ.
crassipes, *Benth. in Trans. Linn. Soc.* xxx. (1875) 555.—Am. trop.
Crugerii, *Griseb. Fl. Brit. W. Ind.* 224.—Ind. occ.
Cumingii, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.—N. Granat.
cylindrocarpa, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 96 = Harrisii.
cynometroides, Bedd. Fl. Sylv. t. 317 = Inga cynometroides.
dasyantha, Walp. Rep. i. 926, sphalm. = dysantha.
depauperata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 546.—Am. trop.
diademata, Lem. Jard. Fleur. iii. (1852-53) t. 305 = bicolor.
dysantha, *Benth. in Hook. Journ. Bot.* ii. (1840) 138.—Bras.
emarginata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 95.—Mexic.
eriophylla, *Benth. l. c.* 105.—N. Mexic.
expansa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 549.—Am. trop.
falcata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 97.—Am. cent.
fasciculata, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.—Bras.
filipes, *Benth. l. c.* 139.—Am. trop.
foliolosa, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 110.—Am. trop.
foliosa, *Benth. l. c.*—Bras.
formosa, *Benth. l. c.* 98.—Am. trop.
Gardneri, *Benth. l. c.* 108.—Bras.
geminata, Benth. in Trans. Linn. Soc. xxx. (1875) 548 Pithecolobium geminatum.
glomerulata, *Karst. Fl. Colomb.* ii. 5. t. 103.—N. Granat.
glyphoxylon, *Spruce, ex Benth. in Trans. Linn. Soc.* xxx. (1875) 539.—Ecuador.
gracilis, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 180 = formosa.
gracilis, *Klotzsch, ex Baker, in Saund. Refug. Bot.* t. 294.—Am. trop.
grandiflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 139.—Am. trop.
Griffithii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 537.—Reg. Himal.
Guildingii, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 96.—Bras.
haematocephala, *Hassk. Retzia*, 216.—Hab.?
haematomma, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 103.—Am. trop.
Harrisii, *Benth. l. c.* 95.—Mexic.
herbacea, Engelm. ex A. Gray, Pl. Fendl. 39 = humilis.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880-82) 110.—Ins. Mascar.
hirsuta, *Benth. in Trans. Linn. Soc.* xxx. (1875) 554.—Mexic.
hirtiflora, *Benth. l. c.* 550.—Am. trop.
Hookeriana, *Schomb. in Linnaea*, xx. (1847) 754 = rigida.
Houstoni, *Benth. in Hook. Journ. Bot.* ii. (1840) 139.—Mexic.
humilis, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 103.—N. Mexic.
hymenaeoides, *Benth. in Trans. Linn. Soc.* xxx. (1875) 537.—Am. trop.
Hystrix, Griseb. Cat. Pl. Cub. 83 = Pithecolobium Hystrix.
Kunthii, Benth. in Hook. Journ. Bot. ii. (1840) 139 = grandiflora.
Lambertiana, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 100.—Mexic.

CALLIANDRA:—
lanata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 553.—Am. trop.
lasiopus, Griseb. Fl. Brit. W. Ind. 711 = Pithecolobium cauliflorum.
latifolium, Griseb. l. c. 225 = Pithecolobium latifolium.
laxa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 551.—Am. trop.
leptopoda, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 101.—Bras.
linearis, *Benth. l. c.* v. (1846) 103.—Bras.
longipes, *Benth. in Trans. Linn. Soc.* xxx. (1875) 538.—Bras.
longipinna, *Benth. l. c.* 546.—Am. trop.
macrocephala, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.—Bras.
Magdalenae, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 102.—Am. trop.
malacophylla, *Benth. l. c.* iii. (1844) 100.—Mexic.
marginata, Benth. in Trans. Linn. Soc. xxx. (1875) 586 = Pithecolobium marginatum.
marginata, Griseb. Fl. Brit. W. Ind. 710 = formosa.
mertensioides, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 106.—Am. trop.
metrosideriflora, *Benth. l. c.* i. (1842) 527.—Mexic.
microphylla, *Benth. in Hook. Journ. Bot.* ii. (1840) 139.—Bras.
mollissima, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 97.—Bras.
myriophylla, *Benth. l. c.* 111.—Bras.
obovata, Benth. l. c. 101 = Pithecolobium lusorium.
obtusifolia, Karst. Fl. Colomb. ii. 41. t. 121 = purpurea.
Pacara, *Griseb. in Goett. Abh.* xxiv. (1879) 123.—Reg. Argent.
pallens, Benth. in Hook. Lond. Journ. Bot. v. (1846) 102 = Pithecolobium albicans.
parviflora, *Benth. l. c.* iii. (1844) 112.—Bras.
Patrisii, *Sagot, in Ann. Sc. Nat.* Sér. VI. xiii. (1882) 324.—Guiana.
pauciflora, *Griseb. Cat. Pl. Cub.* 284.—Cuba.
Peckoltii, *Benth. in Trans. Linn. Soc.* xxx. (1875) 555.—Bras.
pedicellata, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 102.—Ins. S. Doming.
portoricensis, *Benth. l. c.* 99.—Am. bor.; Ind. occ.
prehensilis, Wright, in Sauv. Fl. Cub. 35 = Pithecolobium prehensile.
prostrata, *Benth. in Trans. Linn. Soc.* xxx. (1875) 554.—Peruv.
Purdiaei, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 104.—N. Granat.
purpurea, *Benth. l. c.* iii. (1844) 104.—Am. trop.
quadrangularis, Walp. Rep. i. 927 = tetragona.
reticulata, *A. Gray, Pl. Wright.* ii. 53.—N. Mexic.
revoluta, Griseb. Cat. Pl. Cub. 83 = Pithecolobium obovale.
rigida, *Benth. in Hook. Lond. Journ. Bot.* v. (1846) 103.—Guiana.
Rutenbergiana, *Vatke, in Bremen Abh.* vii. (1882) 250.—Madag.
Saman, Griseb. Fl. Brit. W. Ind. 225 = Pithecolobium Saman.
Sancti-Pauli, Hassk. Retzia, i. 214 = Tweedii.
Schottii, *Torr. ex S. Wats. in Proc. Am. Acad.* xx. (1885) 364.—Arizona.
scutellifera, *Benth. in Hook. Journ. Bot.* ii. (1840) 139.—Am. trop.
Seemanni, *Benth. in Seem. Bot. Voy. Herald.* 116. t. 22.—Venezuela.
sessilis, *Benth. in Hook. Journ. Bot.* ii. (1840) 141.—Bras.
Spruceana, Benth. in Trans. Linn. Soc. xxx. (1875) 598 = Pithecolobium parvifolium.
squarrosa, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 104.—Am. trop.
stenophylla, *Wawra, in Oestr. Bot. Zeitschr.* xxix. (1879) 215.—Bras.
stipulacea, *Benth. in Hook. Journ. Bot.* ii. (1840) 137.—Guiana.
subnervosa, *Benth. in Trans. Linn. Soc.* xxx. (1875) 540.—Am. trop.
subspicata, *Benth. l. c.* 556.—Am. trop.
surinamensis, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 105.—Guiana.
taxifolia, *Benth. l. c.* 104.—N. Granat.

CALLIANDRA:—
tenuiflora, *Benth. in Trans. Linn. Soc.* xxx. (1875) 547.—Am. trop.
tergemina, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 96.—Am. trop.
tetragona, *Benth. in Hook. Journ. Bot.* ii. (1840) 139.—Am. trop.
tetraphylla, *Benth. in Trans. Linn. Soc.* xxx. (1875) 544.—Mexic.
Thouarsiana, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 365.—Madag.
trinervia, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 94.—Bras.
tubulosa, Benth. l. c. 101 = Pithecolobium Saman.
turbinata, *Benth. l. c.* 109.—Bras.
Tweedii, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.—Bras.
umbellifera, *Benth. l. c.* 141.—Bras.
umbrosa, *Benth. in Benth. & Hook. f. Gen.* i. 597.—Ind. or.; Penang.
ungulata, Benth. in Hook. Lond. Journ. Bot. i. (1842) 527 = portoricensis.
vespertina, Benth. l. c. = portoricensis.
virgata, *Benth. in Hook. Journ. Bot.* ii. (1840) 140.—Bras.
viridiflora, *Benth. in Hook. Lond. Journ. Bot.* iii. (1844) 98.—Bras.
viscidula, *Benth. l. c.* 109.—Bras.
Wendlandi, *Benth. in Trans. Linn. Soc.* xxx. (1875) 556.—Guatemala.
xalapensis, Benth. in Hook. Lond. Journ. Bot. iii. (1844) 105 = laxa.

CALLIANIRA, Miq. Syst. Pip. 344 (1843) = **Piper**, Linn.
melastomoides, Miq. l. c. 345 = P. melastomoides.

CALLIANTHEMUM, C. A. Mey. in Ledeb. Fl. Alt. ii. 336 (1830). *RANUNCULACEAE*, Benth. & Hook. f. i. 5.
anemonoides, Endl. ex Heynh. Nom. ii. 106 = rutaefolium.
bipinnatum, Dulac, Fl. Hautes-Pyr. 216 = rutaefolium.
cachemirianum, *Cambess. in Jacquem. Voy. Bot.* 5.—Reg. Himal.
coriandrifolium, Reichb. Fl. Germ. Excurs. 727 = rutaefolium.
Endlicheri, Walp. Rep. i. 33 = Oxygraphis polypetala.
pimpinelloides, Royle, Illustr. Bot. Himal. 45 (Hook. f. & Thoms. Fl. Ind. 26) = cachemirianum.
rutaefolium, *C. A. Mey. in Ledeb. Fl. Alt.* ii. 336.—Sibir.
rutaefolium, Reichb. Fl. Germ. Excurs. 727 = Ranunculus anemonoides.
semiverticillatum, *Phil. in Linnaea*, xxxiii. (1864–65) 3.—Chili.

CALLIAS, Cass. in Dict. Sc. Nat. xxiv. 327 (1822) = KALLIAS, Cass. = **Heliopsis**, Pers. (Compos.).

CALLICARPA, Linn. in Act. Soc. Ups. (1741) 80. *VERBENACEAE*, Benth. & Hook. f. ii. 1150.
AGANON, Rafin. Sylva Tellur. 161 (1838).
AMICTONIS, Rafin. l. c. (1838).
BURCHARDIA, [Heist.] Duham. Arb. i. 111. t. 44 (1755).
ILLA, Adans. Fam. ii. 446 (1763).
JOHNSONIA, T. Dale, ex Mill. Gard. Dict. ed. VI. App. 75 (1752).
PORPHYRA, Lour. Fl. Cochinch. 69 (1790).
SPONDYLOCOCCA, Mitch. in Act. Phys.-Med. Acad. Nat. Cur. viii. (1748) 218.
TOMEA, Linn. Diss. Dass. 5 (1747) (Amoen. Acad. i. 389).
aculeolata, *Schau. in DC. Prod.* xi. 642.—Ins. S. Doming.
acuminata, *H. B. & K. Nov. Gen. et Sp.* ii. 252.—N. Granat.
acuminata, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 394.—Ind. or.
acuminatissima, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 409 = Geunsia farinosa.
acutidens, *Schau. in DC. Prod.* xi. 645.—Cochinch.
adenanthera, R. Br. Prod. 513 = cana.
albida, *Blume, Bijdr.* 818.—Java.
americana, Blanco, Fl. Filip. ed I. 517 = Blancoi.
americana, *Linn. Sp. Pl.* 111.—Am. bor.
americana, Lour. Fl. Cochinch. 70 = cana.

CALLICARPA:—
ampla, *Schau. in DC. Prod.* xi. 642.—Jamaica.
angusta, *Schau. l. c.*—Ins. Philipp.
arborea, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 567 = lanata.
arborea, *Roxb. Hort. Beng.* [10]; *Fl. Ind.* i. 390.—As. trop.
arborea, Wall. Cat. n. 1826, partim = vestita.
attenuata, Wall. l. c. n. 1835 = longifolia.
bicolor, Juss. in Ann. Mus. Par. vii. (1806) 77 = cana.
bicolor, F. Vill. in Blanco, Fl. Philipp. ed. III. Nov. App. 158 = Blancoi.
Blancoi, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 315.—Ins. Philipp.
Blumei, *Zoll. & Moritzi, Syst. Verz. Zoll.* 53.—Malaya.
Bonplandiana, Schult. Mant. iii. 50 = acuminata.
breviceps, Hance, in Journ. Linn. Soc. xiii. (1873) 116 = brevipes.
brevipes, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 233.—China.
cana, Dalz. & Gibs. Bomb. Fl. 200 = lanata.
cana, Gamble, Darjeeling List, 60; Ind. Timbers, 283 = macrophylla.
cana, *Linn. Mant.* ii. 198.—As. trop.; Austral.
cana, Wall. Cat. n. 1834 = pedunculata et longifolia.
cinerea, *A. Rich. Fl. Cub. Fanerog.* ii. 145.—Cuba.
cordifolia, Ruiz & Pav. Fl. Per. i. 50. t. 57. f. a = Aegiphila Mutisii.
Cumingiana, Schau. in DC. Prod. xi. 644 = Geunsia Cumingiana.
cuspidata, Roxb. Hort. Beng. 83; Fl. Ind. i. 394 = pedunculata.
dentata, Roth, Nov. Pl. Sp. 81; Wall. Cat. n. 6319 = pedunculata.
dentata, Roxb. ex Wall. l. c. p. 87. n. 1834 = cana.
dentata, Wall. l. c. n. 6319, partim = longifolia.
dichotoma, Raeusch. Nom. ed. III. 37 = purpurea.
discolor, Willd. ex Steud. Nom. ed. II. i. 257 = Aegiphila arborescens.
erioclona, Schau. in DC. Prod. xi. 643 = cana.
farinosa, Roxb. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 567 = lanata.
ferruginea, *Sw. Prod. Veg. Ind. Occ.* 31.—Jamaica.
formosana, *Rolfe, in Journ. Bot.* xx. (1882) 358.—Ins. Formosa.
fulva, *A. Rich. Fl. Cub. Fanerog.* ii. 145.—Ins. Cuba.
globiflora, *Ruiz & Pav. Fl. Per.* i. 49. t. 57. f. b.—Am. austr.
gracilis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 154.—Japon.
hexandra, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 410 = Geunsia farinosa?
Heynii, Roth, Nov. Pl. Sp. 82 = cana.
Horsfieldii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 217.—Java.
incana, Roxb. Hort. Beng. [10]; Fl. Ind. i. 393 = macrophylla.
integerrima, *Champ. ex Benth. in Hook. Kew Journ.* v. (1853) 135.—China.
integrifolia, Jacq. Enum. Pl. Carib. 12 = Aegiphila arborescens.
japonica, *Thunb. Fl. Jap.* 60.—Japon.
lanata, Gamble, Darjeeling List, 60 = vestita.
lanata, Lam. ex Steud. Nom. ed. I. 137 = Premna lanata (sp. ignot.).
lanata, *Linn. Mant.* ii. 331.—Ind. or.
lanata, Schau. in DC. Prod. xi. 644 = pedunculata.
lanceolaria, Roxb. Hort. Beng. [10]; Fl. Ind. i. 395 = longifolia.
latifolia, *Zipp. ex Span. in Linnaea*, xv. (1841) 330, nomen.—Ins. Timor.
lobata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 566.—Reg. Himal.
longifolia, *Lam. Encyc.* i. 563.—Malaya; Austral.
macrocarpa, Raeusch. Nom. ed. III. 37 = cana?
macrophylla, *Vahl, Symb. Bot.* iii. 13. t. 53.—Ind. or.
magna, *Schau. in DC. Prod.* xi. 641.—Ins. Philipp.
micrantha, *Vidal, Phan. Cuming. Philipp.* 187.—Ins. Philipp.
Mimurazaki, Hassk. Cat. Hort. Bog. Alt. 136 = japonica.
mollis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 155.—Japon.
mollis, Willd. ex Schau. in DC. Prod. xi. 644 = acuminata, *H. B. & K.*

CALLICARPA:—
nudiflora, Hook. & Arn. Bot. Beech. Voy. 206 = Reevesii.
oblongifolia, Hassk. Cat. Hort. Bog. Alt. 136 = longifolia.
paniculata, Lam. Encyc. i. 563 = Chilianthus arboreus.
parvifolia, *Hook. & Arn. Bot. Beech. Voy.* 305.—Mexic.
pedunculata, *R. Br. Prod.* 513.—Malaya; Austral.
pentandra, Roxb. Hort. Beng. 83; Fl. Ind. i. 395 = Geunsia farinosa.
psilocalyx, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 569.—Reg. Himal.
purpurea, *Juss. in Ann. Mus. Par.* vii. (1806) 67.—China.
Reevesii, *Wall. Cat.* n. 1830.—China.
reticulata, Poepp. ex Walp. Rep. iv. 72 = Petitia Poeppigii.
reticulata, *Sw. Prod. Veg. Ind. Occ.* 31.—Jamaica.
Rheedii, Kostel. Allg. Med.-Pharm. Fl. iii. 829 = cana.
rhynchophylla, *Miq. Fl. Ind. Bat.* ii. 888.—Malaya.
Roxburghiana, Schult. Mant. iii. 54 = longifolia.
Roxburghii, Wall. Cat. n. 1833 = macrophylla.
rubella, *Lindl. Bot. Reg.* t. 883.—Reg. Himal.; China.
Schlimii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) I. 217.—N. Granat.
serrata, Moench, Meth. 468 = americana.
sessilifolia, Wall. Cat. n. 1837 = rubella.
sinensis, Hort. ex Steud. Nom. ed. II. i. 257 = cana.
subintegerrima, H. B. & K. Nov. Gen. et Sp. ii. 252 = acuminata, *H. B. & K.*
subpubescens, *Hook. & Arn. Bot. Beech. Voy.* 305.—Mexic.
sumatrana, Miq. Fl. Ind. Bat. ii. 888 = cana.
tectonaefolia, Wall. Cat. n. 1827 = arborea.
tenuiflora, Champ. ex Benth. in Hook. Kew Journ. v. (1853) 135 = rubella.
tiliaefolia, Teijsm. & Binn. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 569 = pedunculata.
tomentosa, Koen. ex Vahl, Symb. Bot. iii. 13 = macrophylla.
tomentosa, Lam. Encyc. i. 562 = cana.
tomentosa, Murr. Syst. Veg. ed. XIII. 153 = lanata.
tomentosa, *Willd. Enum. Hort. Berol.* 158; *Hook. & Arn. Bot. Beech. Voy.* 305.—China.
Tomex, Poir. Encyc. Suppl. ii. 32 = lanata.
triloba, *Lour. Fl. Cochinch.* 70.—Cochinch.
umbellata, *Lour. l. c.*—Cochinch.
velutina, *Presl, Bot. Bemerk.* 98.—Bras.
vestita, *Wall. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 567.—Reg. Himal.
viburnifolia, Salisb. Prod. 53 = americana.
villosa, Roxb. Hort. Beng. [10] = arborea.
villosa, *Vahl, Symb. Bot.* iii. 14.—Ind. or.?
Wallichiana, Walp. Rep. iv. 125 = lanata.
Zollingeriana, Schau. in DC. Prod. xi. 640 = mollis.

CALLICEPHALUS, C. A. Mey. Verz. Pfl. Cauc. 66 (1831) = **Centaurea**, Linn. (Compos.).
nitens, C. A. Mey. l. c. = Cent. nitens.

CALLICHLAMYS, Miq. in Linnaea, xviii. (1844) 254. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1039.
riparia, *Miq. l. c.*—Panama.
rubiginosa, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 201 = riparia.
rufinervis, Miers, l. c. 201 = riparia.
splendida, Miers, l. c. = riparia.

CALLICHLOË, Pfeiff. Nom. i. 536 (1871) = seq.

CALLICHLOEA, Spreng. ex Steud. Nom. ed. II. i. 257 (1840) = **Elionurus**, Humb. & Bonpl. (Gramin.).
elegans, Spreng. l. c. = E. elegans.

CALLICHROA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 31 (1835) = **Layia**, Hook. & Arn. (Compos.).
angustifolia, Walp. Rep. ii. 632 = L. platyglossa.
angustifolia, Walp. l. c. iii. 396 = Castilleja angustifolia.
Bradburii, Walp. l. c. = Castilleja Bradburii.
Douglasii, Torr. & Gray, Fl. N. Am. ii. 396 = L. Douglasii.
hirsuta, Walp. Rep. ii. 632 = L. platyglossa.
platyglossa, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 31 = L. platyglossa.

CALLICOCCA, Schreb. Gen. i. 126 (1789) = **Cephaëlis**, Sw. (Rubiac.).
alba, J. F. Gmel. Syst. 371 = Ceph. alba.
elata, J. F. Gmel. l. c. 372 = Ceph. elata.
glabra, J. F. Gmel. l. c. = Ceph. glabra.
guianensis, J. F. Gmel. l. c. = Ceph. involucrata.
Ipecacuanha, Brot. in Roem. Arch. iii. III. 455 = Psychotria Ipecacuanha.
muscosa, Steud. Nom. ed. II. i. 257 = Ceph. muscosa.
purpurea, J. F. Gmel. Syst. 371 = Ceph. purpurea.
tomentosa, J. F. Gmel. l. c. = Psychotria tomentosa.

CALLICOMA, Andr. Bot. Rep. t. 566 (1809). *SAXIFRAGEAE*, Benth. & Hook. f. i. 649.
CALYCOMIS, R. Br. in Flind. Voy. App. 549 (1814).
STUTZERIA, F. Muell. Fragm. v. 31 (1865).
Billardieri, *D. Don, in Edinb. N. Phil. Journ.* (Apr.–June 1830) 94.—Austral.
ferruginea, D. Don, l. c. = serratifolia.
serratifolia, *Andr. Bot. Rep.* t. 566.—Austral.
Stutzeri, *F. Muell. Fragm.* v. 34.—Austral.
ternata, *Montr. in Mém. Acad. Lyon*, x. (1860) 211.—N. Caled.

CALLICORE, Link, Handb. i. 193 (1829) = **Amaryllis**, Linn.
crocata, Link, l. c. = Hippeastrum rutilum.
reticulata, Link, l. c. = Hippeastrum reticulatum.
rosea, Link, l. c. = A. Belladonna.
rutila, Link, l. c. 194 = Hippeastrum rutilum.

CALLICORNIA, Burm. f. Prod. Fl. Cap. 24 (1768) = **Leyssera**, Linn. (Compos.).
gnaphaloides, Burm. f. l. c. = L. gnaphaloides.
tenuifolia, Burm. f. l. c.—Afr. austr.
triflora, Burm. f. l. c.—Afr. austr.

CALLICYSTHUS, Endl. Prod. Fl. Norf. 90 (1833) = **Vigna**, Savi (Legumin.).
volubilis, Endl. l. c.—Ins. Norfolk.

CALLIDRYNOS, Neraud, in Freyc. Voy. Bot. 29 (1829), nomen (Sapind.).

CALLIGLOSSA, Hook. & Arn. Bot. Beech. Voy. 356 (1840) = **Layia**, Hook. & Arn. (Compos.).
Douglasii, Hook. & Arn. l. c. = L. Calliglossa.

CALLIGONUM, Linn. Gen. ed. I. 345 (1737). *POLYGONACEAE*, Benth. & Hook. f. iii. 95.
CALLIPHYSA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 24 (1835).
GIBSONIA, Stocks, in Proc. Linn. Soc. i. (1848) 367.
PALLASIA, Linn. f. Suppl. 37 (1781).
PTEROCOCCUS, Pall. Reise, i. App. 738. t. 5 (1776); Reise, ii. App. 43 (1777).
TRAXILISA, Rafin. Sylva Tellur. 161 (1838).
acanthopterum, *Borszcz. Aralo-Casp. Callig.* 34.—Reg. Aral. Casp.
anfractuosum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 487 (*Lehm. Rel.* 311).—As. centr.
aralense, *Borszcz. Aralo-Casp. Callig.* 31.—Reg. Aral. Casp.
asperum, Lour. Fl. Cochinch. 342 = Tetracera sarmentosa.
bucocladum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 485 (*Lehm. Rel.* 309).—As. centr.
Bungei, *Boiss. Fl. Orient.* iv. 999.—Persia.
Calliphysa, *Bunge, Del. Sem. Dorpat.* (1839) 8.—Reg. Casp.; Turkest.
canescens, Pursh, Fl. Am. Sept. ii. 370 = Atriplex canescens.
Caput-Medusae, *Schrenk, Enum. Pl. Nov.* i. 9.—Soongaria.
colubrinum, *Borszcz. Aralo-Casp. Callig.* 38.—Reg. Aral. Casp.
comosum, *L'Hérit. in Trans. Linn. Soc.* i. (1791) 180.—Aegypt.; Arab.; Persia.
crinitum, *Boiss. Diagn.* Ser. II. iv. 77.—Beluchist.
crispum, *Bunge, Del. Sem. Dorp.* (1839) 8.—Hab.?
cristatum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 485 (*Lehm. Rel.* 309).—As. centr.
densum, *Borszcz. Aralo-Casp. Callig.* 36.—Reg. Aral. Casp.
denticulatum, *Bunge, ex Boiss. Fl. Orient.* iv. 999.—Persia.
erinaceum, *Borszcz. Aralo-Casp. Callig.* 37.—Reg. Aral. Casp.

CALLIGONUM:—
eriopodum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 485, 486 (*Lehm. Rel.* 309, 310).—Turkest.
flavidum, *Bunge, Del. Sem. Dorp.* (1839) 8.—Hab.?
gobicum, Bunge, ex Meissn. in DC. Prod. xiv. 29 = mongolicum.
horridum, *Borszcz. Aralo-Casp. Callig.* 32.—Reg. Aral. Casp.
leucocladum, *Bunge, in Mém. Sav. Etr. Pétersb.* (1851) 485 (*Lehm. Rel.* 309).—As. centr.
macrocarpum, *Borszcz. Aralo-Casp. Callig.* 35.—Reg. Aral. Casp.
medusecephalum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69.—C. Caput-Medusae.
microcarpum, *Borszcz. l. c.* 41.—Reg. Aral. Casp.
mongolicum, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 204.—Mongolia.
Murex, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 485, 486 (*Lehm. Rel.* 309, 310).—As. centr.
Pallasia, *L'Hérit. in Trans. Linn. Soc.* i. (1791) 180.—Reg. Casp.; Turkest.
Panderi, *Fisch. ex Sweet, Hort. Brit.* ed. II. 440, nomen.—Reg. Casp.
paniculatum, *Borszcz. Aralo-Casp. Callig.* 42.—Reg. Aral. Casp.
persicum, *Boiss. Fl. Orient.* iv. 999.—Persia.
platyacanthum, *Borszcz. Aralo-Casp. Callig.* 37.—Reg. Aral. Casp.
polygonoides, *Linn. Sp. Pl.* 530.—Armenia; Persia; Ind. bor. occ.
polygonoides, Pall. Reise, iii. 530 = C. Pallasia.
Rotula, *Boszcz. Aralo-Casp. Callig.* 35.—Reg. Aral. Casp.
rubicundum, *Bunge, Del. Sem. Dorp.* (1839) 8.—Hab.?
songaricum, *Endl. Gen. Suppl.* iv. II. 50.—Soongaria.
stenopterum, *Bunge, ex Boiss. Fl. Orient.* iv. 999.—Persia; Afghan.
tetrapterum, *Jaub. & Spach, Illustr.* v. 80. t. 471.—Mesopotam.
tripterum, Bunge, ex Boiss. Fl. Orient. iv. 999 = Bungei.

CALLIGONUM, Lour. Fl. Cochinch. 342 (1790) = **Tetracera**, Linn. (Dilleniac.).
asperum, Lour. l. c. = T. sarmentosa.

CALLILEPIS, DC. Prod. v. 671 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 339.
glabra, *DC. l. c.*—Afr. austr.
hispida, *DC. l. c.*—Afr. austr.
Laureola, *DC. l. c.*—Afr. austr.
leptophylla, *Harv. in Harv. & Sond. Fl. Cap.* iii. 136.—Afr. austr.
salicifolia, *Oliver, in Hook. Ic. Pl.* t. 1482.—Afr. austr.

CALLIOPEA, D. Don, in Edinb. N. Phil. Journ. (1828–29) 309 = **Crepis**, Linn. (Compos.).
aurea, D. Don, l. c. = Crepis aurea.

CALLIOPSIS, Reichb. Ic. et Desc. Pl. t. 70 (1824) = **Coreopsis**, Linn. (Compos.).
Atkinsoniana, Hook. Fl. Bor. Am. i. 310 = Cor. Atkinsoniana.
atropurpurea, Hort. ex Steud. Nom. ed. II. i. 258 = Cor. tinctoria.
basalis, A. Dietr. in Otto & Dietr. Allgem. Gartenz. iii. (1835) 330.—Am. bor.
bicolor, Reichb. Ic. et Desc. Pl. t. 70 = Cor. tinctoria.
cardaminefolia, DC. Prod. v. 568 = Cor. cardaminefolia.
Drummondi, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 315 = Cor. Drummondii.
fimbriata, Lange, Sem. Hort. Haun. Adn. (1866) 5.—Hab.?
nudata, Spreng. Syst. iii. 611 = Cor. nudata.
palmata, Spreng. l. c. = Cor. palmata.
rosea, Spreng. l. c. = Cor. rosea.
tinctoria, DC. Prod. v. 568 = Cor. tinctoria.

CALLIPELTIS, Stev. in Mém. Soc. Nat. Mosc. vii. (1829) 275 (Obs. Pl. Ross. 69). *RUBIACEAE*, Benth. & Hook. f. ii. 148.
aperta, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 110.—Aegypt.; Persia.
Cucullaria, *Stev. in Mém. Soc. Nat. Mosc.* vii. (1829) 275.—Orient; Reg. Mediterr.
microstegia, *Boiss. Fl. Orient.* iii. 84.—Persia.

CALLIPHYSA, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 24 (1835) = **Calligonum**, Linn. (Polygonac.).
juncea, Fisch. & Mey. l. c. = Calligonum Calliphysa.
polygonea, C. A. Mey. ex Borszcz. Aralo-Casp. Callig. 43 = Calligonum Calliphysa.

CALLIPRENA, Salisb. Gen. Pl. Fragm. 89 (1855) = **Allium**, Tourn. (Liliac.).
cernua, Salisb. l. c. = A. cernuum.

CALLIPRORA, Lindl. Bot. Reg. t. 1590 (1833) = **Brodiaea**, Sm. (Liliac.).
aurantea, Kellogg, in Proc. Calif. Acad. ii. (1863) 20 = B. ixioides.
flava, Hort. ex Steud. Nom. ed. II. i. 258 = B. ixioides.
lutea, Lindl. Bot. Reg. t. 1590 = B. ixioides.

CALLIPSYCHE, Herb. in Bot. Reg. (1842) Misc. 49. *AMARYLLIDEAE*, Benth. & Hook. f. iii. 731.
aurantiaca, *Baker, in Saund. Ref. Bot.* t. 167.—Ecuador.
auratum, Hort. ex Gard. Chron. (1871) 6 = pracc.
cucrosioides, *Herb. in Bot. Reg.* (1842) *Misc.* 50.—Ecuador.
mexicana, M. Roem. Syn. Ensat. 155 = eucrosioides.
mirabilis, *Baker, in Saund. Ref. Bot.* t. 168.—Peruv.

CALLIRHOË, Nutt. in Journ. Acad. Philad. ii. (1821) 181. *MALVACEAE*, Benth. & Hook. f. i. 201.
MONOLIX, Rafin. Cat. 14 (1824).
NUTTALLIA, Bart. Fl. N. Am. ii. 74. t. 62 (1822).
alcaeoides, *A. Gray, Pl. Fendl.* 18.—Am. bor.
digitata, *Nutt. in Journ. Acad. Philad.* ii. (1821) 181.—Am. bor. occ.
involucrata, *A. Gray, Pl. Fendl.* 16.—Am. bor. occ.
linearriloba, *A. Gray, in Proc. Am. Acad.* xix. (1883) 74.—Am. bor.
macrorrhiza, *A. Gray, Pl. Fendl.* 18 = alcaeoides.
palmata, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 449 = involucrata.
Papaver, *A. Gray, Pl. Fendl.* 17.—Am. bor. occ.
pedata, *A. Gray, l. c.*—Am. bor. occ.
spicata, Regel, Gartenfl. (1872) 291. t. 737 = Sidalcea malvaeflora.
triangulata, *A. Gray, Pl. Fendl.* 16.—Am. bor.
verticillata, Grocnl. in Rev. Hortic. (1862) 171 = involucrata.

CALLIRRHOË, A. Gray, Pl. Fendl. 16 (1849) = praec.

CALLISACE, Fisch. in Hoffm. Gen. Umb. ed. II. 170 = **Angelica**, Linn. (Umbellif.).
cantabrigiensis, Hoffm. ex Steud. Nom. ed. I. 762. 138 = Peucedanum palustre.
dahurica, Fisch. in Hoffm. Gen. Umb. ed. II. 170 = A. dahurica.
ruthenica, Fisch. l. c. = Peucedanum ruthenicum?
Schiefereckii, Hoffm. l. c. = Peucedanum palustre.

CALLISEMA, Steud. Nom. ed. II. i. 258 (1840) = seq.

CALLISEMAEA, Benth. in Ann. Wien. Mus. ii. (1838) 105 = **Platypodium**, Vog. (Legumin.).
elegans, Walp. Rep. i. 802 = P. elegans.
grandiflora, Benth. in Ann. Wien. Mus. ii. (1838) 105 = P. grandiflorum.
pubescens, Benth. l. c. = P. elegans.
sericea, Benth. l. c. = P. elegans.
viridis, Walp. Rep. i. 802 = P. viride.

CALLISIA, Linn. in Loefl. It. Hisp. 305 (1758). *COMMELINACEAE*, Benth. & Hook. f. iii. 854.
HAPALANTHUS, Jacq. Stirp. Am. 11. t. 11 (1763).
ciliata, H. B. & K. Nov. Gen. et Sp. i. 261 = Tradescantia elongata.
ciliata, Pers. Syn. i. 56 = repens.
delicatula, Kunth, Enum. Pl. iv. 63 = umbellulata.
diffusa, Pohl, ex Seub. in Mart. Fl. Bras. iii. 1. 251 = Tradescantia effusa.
glandulosa, Pohl, ex Seub. l. c. 255 = Dithyrocarpus glandulosus.
insignis, *C. B. Clarke, in Hemsl. Diagn. Pl. Mex.* 56.—Mexic.

CALLISIA:—
Martensiana, *C. B. Clarke, in DC. Monog. Phan.* iii. 312.—Mexic.
meiandra, Wright, in Sauv. Fl. Cub. 159 = Tradescantia cordifolia.
monandra, Schult. f. Syst. vii. 1179 = umbellulata.
orientalis, Buch.-Ham. ex Wall. Cat. n. 5222 = Aneilema Hamiltonianum.
repens, *Linn. Sp. Pl.* ed. II. 62.—Am. austr.
rosea, Pohl, ex Seub. in Mart. Fl. Bras. iii. 1. 252 = Tradescantia Gaudichaudiana.
umbellulata, *Lam. Illustr.* i. 130. t. 35. f. 2.—Ins. Martinic.

CALLISTA, Lour. Fl. Cochinch. 519 (1790) = **Dendrobium**, Sw. (Orchid.).
amabilis, Lour. l. c.—Cochinch.

CALLISTA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 154 = **Erica**, Tourn.
acuminata, G. Don, Gen. Syst. iii. 813 = E. Massoni.
armata, G. Don, l. c. = E. Sparmanni.
Bandonia, G. Don, l. c. = E. Irbyana.
Bedfordiana, G. Don, l. c. 814 = E. Massoni.
blanda, G. Don, l. c. 815 = E. blanda.
Broadleyana, G. Don, l. c. 813 = E. Broadleyana.
buccinaeflora, G. Don, l. c. = E. Massoni.
calostoma, G. Don, l. c. 815 = E. Massoni.
Celsiana, G. Don, l. c. 813 = E. Thunbergii.
Cliffortiana, G. Don, l. c. 815 = E. tenuiflora.
comosa, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. comosa.
Comptoniana, G. Don, Gen. Syst. iii. 813 = E. curvifolia.
Coventryana, G. Don, l. c. 815 = E. pavettaeflora.
daphniflora, G. Don, l. c. = E. daphniflora.
daphnoides, G. Don, l. c. = E. praestans.
denticulata, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. denticulata.
fastigiata, G. Don, Gen. Syst. iii. 814 = E. fastigiata.
ferruginea, G. Don, l. c. 813 = E. squarrosa.
fragrans, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. fragrans.
Humeana, G. Don, Gen. Syst. iii. 813 = Erica fastigiata.
hyacinthoides, G. Don, l. c. 814 = E. praestans.
inflata, G. Don, l. c. = E. inflata.
infundibuliformis, G. Don, l. c. 813 = E. pavettaeflora.
Juliana, G. Don, l. c. = E. Walkeriana.
Lawsoniana, G. Don, l. c. = E. Lawsoniana.
magnifica, G. Don, l. c. 813 = E. Massoni.
metulaeflora, G. Don, l. c. = E. Massoni.
moschata, G. Don, l. c. 815 = E. Muscari.
mundula, G. Don, l. c. = E. fastigiata.
Muscari, G. Don, l. c. 814 = E. Muscari.
nidiflora, G. Don, l. c. 815 = E. Muscari.
Parmentieri, G. Don, l. c. = E. praestans.
pavettaeflora, G. Don, l. c. = E. pavettaeflora.
pellucida, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. Walkeriana.
pellucida, G. Don, Gen. Syst. iii. 814 = E. venusta.
praegnans, G. Don, l. c. 813 = E. ventricosa.
praestans, G. Don, l. c. 815 = E. praestans.
primuloides, G. Don, l. c. 816 = E. fastigiata.
quadrangularis, G. Don, l. c. = E. Massoni.
rigida, G. Don, l. c. 815 = E. Massoni.
Russeliana, G. Don, l. c. = E. Russeliana.
Sainsburiana, G. Don, l. c. 813 = E. inflata.
stellifera, G. Don, l. c. 815 = E. Massoni.
struthiolaeflora, G. Don, l. c. = E. teretiuscula.
Swainsoniana, G. Don, l. c. 816 = E. Massoni.
Templeana, G. Don, l. c. 814 = E. Massoni.
tenuiflora, G. Don, l. c. 812 = E. tenuiflora.
tetragona, G. Don, l. c. = E. tetragona.
thymifolia, Steud. Nom. ed. II. i. 258 = E. thymifolia.
trossula, G. Don, Gen. Syst. iii. 815 = E. Muscari.
undulata, G. Don, l. c. 813 = E. Massoni.
varia, G. Don, l. c. = E. varia.
ventricosa, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = E. ventricosa.
venusta, G. Don, Gen. Syst. iii. 815 = E. venusta.
verecunda, G. Don, l. c. = E. Meuroni.
Walkeria, G. Don, l. c. 814 = E. Walkeriana.

CALLISTACHYA, Rafin. in Med. Repos. New York v. (1808) 60 = **Veronica**, Tourn. (Scrophular.).
sibirica, Rafin. l. c. = V. sibirica.
virginica, Rafin. l. c. = V. virginica.

CALLISTACHYA, Sm. in Trans. Linn. Soc. ix. (1808) 267 = CALLISTACHYS, Vent.

CALLISTACHYS, Heuffel, in Flora, xxvii. (1844) 528 = **Carex**, Linn. (Cyperac.).
pyrenaica, Heuffel, l. c. = C. pyrenaica.

CALLISTACHYS, Vent. Jard. Malm. 115. t. 115 (1803) = **Oxylobium**, Andr. (Legumin.).
carinata, Meissn. in Lehm. Pl. Preiss. i. 27 = O. carinatum.
cuneata, Heynh. Nom. i. 148 = Isotropis stricta.
cuneifolia, Sm. in Trans. Linn. Soc. ix. (1808) 266 = Isotropis striata.
elliptica, Vent. Jard. Malm. 115 = O. ellipticum.
lanceolata, Vent. l. c. = O. Callistachys.
linariifolia, Hort. Mackay, ex Sweet, Hort. Brit. ed. I. 474, nomen = O. lineare.
linariaefolia, G. Don, Gen. Syst. ii. 117 = O. lineare.
linearis, Benth. in Enum. Pl. Hueg. 27 = O. lineare.
longifolia, Paxt. Mag. viii. 31 = O. Callistachys.
ovata, Sims, Bot. Mag. t. 1925 = O. Callistachys.
oxylobioides, Meissn. in Lehm. Pl. Preiss. i. 27 = O. capitatum.
parviflora, Benth. in Enum. Pl. Hueg. 27 = O. lineare.
purpurea, Van Houtte, ex Heynh. Nom. ii. 107 = O. ellipticum.
retusa, Lodd. Bot. Cab. t. 1983 = O. Callistachys.
sparsa, A. Cunn. ex Benth. in Ann. Wien. Mus. ii. (1838) 69 = O. Pultenaea.
spathulata, Meissn. in Lehm. Pl. Preiss. ii. 208 = O. spathulatum.
tetragona, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) 1. 249 = O. retusum.

CALLISTEMMA, Boiss. Fl. Orient. iii. 146 (1875) = **Scabiosa**, Tourn. (Dipsac.).
brachiata, Boiss. l. c. = S. brachiata.
Sibthorpianum, Nym. Consp. 341 = S. brachiata.

CALLISTEMMA, Cass. in Dict. Sc. Nat. vi. Suppl. 45 (1817), nomen prius = **Callistephus**, Cass. (Compos.).
capitatum, Reichb. Hort. Bot. i. t. 84, sphalm. = *Callistemon capitatus*.
hortense, Cass. in Dict. Sc. Nat. vi. Suppl. 45; et Bull. Soc. Philom. (1817) 32 = Callistephus chinensis.
indicum, G. Don, in Loud. Hort. Brit. 348 = Boltonia indica.

CALLISTEMON, R. Br. in App. Flind. Voy. ii. 547 (1814). *MYRTACEAE*, Benth. & Hook. f. i. 704.
acerosus, Miq. in Nederl. Kruidk. Arch. iv. (1859) 141 = brachyandrus.
acerosus, Tausch, in Flora, xix. (1836) 411 = pinifolius.
amoenus, *Lem. in Illustr. Hortic.* vii. t. 247.—Hab.?
arborescens, F. Muell. in Linnaea, xxv. (1852) 388 = brachyandrus.
brachyandrus, *Lindl. in Journ. Hort. Soc.* iv. (1849) 112.—Austral.
capitatus, Reichb. Ic. Exot. i. 59. t. 84 = Kunzea capitata.
coccineus, *F. Muell. Fragm.* i. 13.—Austral.
Cunninghami, *Hort. Berol. ex C. Koch, Wochenschr.* vi. (1863) 294.—Austral.
flavescens, *Regel, Gartenfl.* (1861) 51.—Austral.
glaucus, Sweet, Fl. Austral. sub t. 29 = speciosus.
Hainesii, F. Muell. l. c. iii. 153 = Kunzea Baxteri.
hybridus, *DC. Prod.* iii. 224.—Austral.
intermedius, *Hort. Berol. ex Steud. Nom.* ed. II. i. 259.—Austral.
juniperinus, *Heynh. Nom.* i. 148.—Austral.
lanceolatus, *Sweet, Fl. Austral.* sub t. 29; *DC. Prod.* iii. 223.—Austral.
leptostachyus, Sweet, l. c. = salignus.
linearis, *DC. Prod.* iii. 223.—Austral.
linearifolius, DC. l. c. = rigidus.
lophanthus, Sweet, Fl. Austral. t. 29 = salignus.
macrostachyus, Lindl. Bot. Reg. (1838) t. 7 = Kunzea Baxteri.
marginatus, Sweet, Hort. Brit. ed. I. 155; DC. Prod. iii. 224 = lanceolatus.
macrostachyus, Lindl. Bot. Reg. (1838) t. 7 = Kunzea Baxteri.

CALLISTEMON:—
nervosus, Lindl. in Mitch. Journ. Trop. Austral. 235 = Melaleuca Leucadendron.
pallidus, DC. Prod. iii. 223 = salignus.
paludosus, F. Muell. Fragm. i. 14 = salignus.
Pancheri, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 183; *in Ann. Sc. Nat.* Sér. V. ii. (1864) 138.—N. Caled.
pendulus, *Regel, Gartenfl.* (1859) 288.—Austral.
phoeniceus, *Lindl. Swan River App.* 10.—Austral.
pinifolius, *Sweet, Hort. Brit.* ed. I. 155; *DC. Prod.* iii. 223.—Austral.
pithyoides, Miq. in Nederl. Kruidk. Arch. iv. (1859) 142 (sp. dub.).—Austral.
pungens, *G. Don, ex Loud. Hort. Brit.* 197.—Austral.
rigidus, *R. Br. in Bot. Reg.* t. 393.—Austral.
rugulosus, DC. Prod. iii. 223 = rigidus.
rugulosus, Miq. in Nederl. Kruidk. Arch. iv. (1859) 141 = coccineus.
salignus, *Sweet, Fl. Austral.* sub t. 29; *DC. Prod.* iii. 223.—Austral.
scaber, Lodd. Bot. Cab. t. 1288 = lanceolatus.
semperflorens, *Lodd. l. c.* t. 523.—Austral.
Sieberi, DC. Prod. iii. 223 = salignus.
speciosus, *DC. l. c.* 224.—Austral.
suberosus, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xi. (1864) 183.—N. Caled.
teretifolius, *F. Muell. in Linnaea*, xxv. (1852) 387.—Austral.
viminalis, G. Don, ex Loud. Hort. Brit. 197 = rigidus.
viridiflorus, Sweet, Hort. Brit. ed. I. 155 = salignus.

CALLISTEPHANA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 367 = **Coronilla**, Linn. (Legumin.).
varia, Fourr. l. c. = C. varia.

CALLISTEPHUS, Cass. in Dict. Sc. Nat. xxxvii. 491(1825). *COMPOSITAE*, Benth. & Hook. f. ii. 270.
CALLISTEMMA, Cass. l. c. vi. Suppl. 45 (1817), nomen prius.
biennis, Lindl. ex DC. Prod. v. 275 = Heteropappus hispidus.
chinensis, Nees, Gen. et Sp. Ast. 222 = hortensis.
concolor, Dalz. in Hook. Kew Journ. ii. (1850) 344 = Pulicaria angustifolia.
Heyneanus, DC. Prod. v. 275 = Pulicaria Wightiana.
hortensis, *Cass. in Dict. Sc. Nat.* xxxvii. 491.—Sibiria; China.
indicus, DC. Prod. v. 275 = Boltonia indica.
Wightianus, DC. l. c. = Pulicaria Wightiana.

CALLISTHENE, Mart. Nov. Gen. et Sp. i. 123. t. 75, 76 (1824). *VOCHYSIACEAE*, Benth. & Hook. f. i. 976.
Blanchetii, *Warm. in Mart. Fl. Bras.* xiii. II. 26.—Am. trop.
erythroclada, *Warm. l. c.* 28.—Am. trop.
fasciculata, *Mart. Nov. Gen. et Sp.* i. 126.—Bras.
major, *Mart. l. c.* 124. t. 75.—Bras.
microphylla, *Warm. in Mart. Fl. Bras.* xiii. II. 28.—Am. trop.
minor, *Mart. Nov. Gen. et Sp.* ii. 126. t. 76.—Bras.
mollissima, *Warm. in Mart. Fl. Bras.* xiii. II. 26.—Am. trop.

CALLISTHENIA, Spreng. Gen. i. 8 (1830) = praec.

CALLISTROMA, Fenzl, in Flora, xxvi. (1843) 457; Endl. Gen. Suppl. iii. 82 (1843) = **Oliveria**, Vent. (Umbell.).
erubescens, Fenzl, l. c. = O. orientalis.

CALLITHAMNA, Herb. Amaryll. 225 (1837) = **Stenomesson**, Herb. (Amaryllid.).
angustifolium, Herb. in Bot. Mag. t. 3866. f. b = S. viridiflorum.
Hartwegianum, Van Houtte, ex Heynh. Nom. ii. 108, nomen = S. aurantiacum?
spathulatum, Herb. Amaryll. 225.—'Truxillo.'
viridiflorum, Herb. l. c. = S. viridiflorum.

CALLITRICHE, Linn. Syst. ed. VI. 82 (1748). *HALORAGEAE*, Benth. & Hook. f. i. 676.
STELLARIA, Hall. Enum. Stirp. Helv. i. 198 (1742).
STELLINA, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 318.
aestivalis, Thuil. Fl. Par. 2 = stagnalis.

CALLITRICHE:—
alpina, Schur, in Oestr. Bot. Zeitschr. (1858) 403 = verna.
amblyocarpa, Scheidw. ex Hegelm. Monog. Callit. 61 = truncata.
androgyna, Linn. Cent. Pl. Rar. i. 31 = verna.
angustifolia, Gilib. Exercit. ii. 421 = autumnalis.
angustifolia, Hoppe, Taschenb. (1792) 160 = verna.
antaretica, *Engelm. ex Hegelm. in Verh. Bot. Ver. Brandenb.* ix. (1867) 20.—Reg. Antaret.
aquatica, Huds. Fl. Angl. ed. II. 439 = verna ?
Asagraei, *Hegelm. Monog. Callit.* 54. t. 3. f. 9.—Am. bor.
attenuata, *Delastre, Fl. Dép. Vienne*, 144.—Gallia.
autumnalis, A. Gray, ex Hegelm. Monog. Callit. 55 = Asagraei.
autumnalis, Kuetz. in Linnaea, vii. (1832) 186 = hamulata.
autumnalis, Bertol. ex Steud. Nom. ed. II. i. 259 = Berteroniana.
autumnalis, Brot. ex Steud. l. c. = latifolia.
autumnalis, *Linn. Fl. Suec.* ed. II. n. 4.—Europ.; Am. bor.
autumnalis, Michx. Fl. Bor. Am. i. 2 = Peplis diandra.
autumnalis, Pollich, Fl. Palat. 3 = verna.
Berteroniana, *Steud. Nom.* ed. II. i. 259.—Chili.
Bolanderi, *Hegelm. in Verh. Bot. Ver. Brandenb.* x. (1868) 116.—Am. bor. occ.
brachycarpa, *Hegelm. l. c.* 115.—Austral.
brevifolia, *Pursh, Fl. Am. Sept.* i. 3.—Am. bor.
brutia, Petagna, Inst. Bot. ii. 10 = hamulata.
capillaris, *Parl. Fl. Ital.* iv. 423.—Italia.
cespitosa, Schultz, Prod. Fl. Stargard. 2 = verna.
confervoides, Thuil. ex Link, in Spreng. Jahrb. Gew. iii. 30 = autumnalis.
connata, Rafin. in Med. Rep. New York, Ser. III. ii. (1811) 408 = hamulata.
cophocarpa, Sendtn. ex Hegelm. Monog. Callit. 55 = verna.
cruciata, Lebel, ex Hegelm. in Verh. Bot. Ver. Brandenb. ix. (1867) 38 = truncata.
cruciata, *Rafin. in Med. Rep. New York*, Ser. III. ii. (1811) 408.—Am. bor.
cuneifolia, A. Br. ex Hegelm. in Verh. Bot. Ver. Brandenb. ix. (1867) 21 = verna.
cyclocarpa, *Hegelm. l. c.* x. (1868) 116.—Austral.
decussata, Link, in Spreng. Jahrb. Gew. iii. 30 = autumnalis.
decussata, Schur, Enum. Pl. Transs. 217 = hamulata.
deflexa, *A. Br. in Hegelm. Monog. Callit.* 58. t. 3.—Am. bor. et austr.
dioica, *Petagn. Inst. Bot.* ii. 10.—Italia.
dioica, Schur, in Verh. Siebenb. Ver. Naturw. (1859) 98 = stagnalis.
Drummondi, *Hegelm. Monog. Callit.* 60.—Austral.
dubia, *Hoffm. Obs. Bot.* n. 1; *ex Roth, Tent. Fl. Germ.* i. 389 = verna.
euautumnalis, Syme, Engl. Bot. ed. III. viii. 122 = autumnalis.
euverna, Syme, l. c. 119 = verna.
foliosa, Rafin. in Med. Repos. New York, V. (1808) 354 = autumnalis.
fontana, Scop. Fl. Carn. ed. II. ii. 251 = verna.
genuina, Ducommun, Taschenb. 243 = hamulata.
glomerata, Thuill. ex Link, in Spreng. Jahrb. Gew. iii. 30 = autumnalis.
graminea, Link, l. c. = truncata.
hamulata, *Kuetz. ex Koch, Syn. Fl. Germ.* ed. II. 271.—Europ.
hermaphrodita, Linn. Cent. Pl. Rar. i. 31 = verna.
heterophylla, Pursh, Fl. Am. Sept. i. 3 = verna.
heteropoda, *Engelm. ex Hegelm. in Verh. Bot. Ver. Brandenb.* ix. (1867) 40.—Am. austr.
intermedia, Hoffm. Fl. Germ. (1791) 2 = hamulata.
intermedia, *Hoppe, Taschenb.* (1792) 159 = verna.
japonica, *Engelm. ex Hegelm. in Verh. Bot. Ver. Brandenb.* x. (1868) 113.—Japon.
Kuetzingi, Rupr. Fl. Ingric. 378 = stagnalis.
latifolia, Gilib. Exercit. ii. 421 = hamulata *vel* verna.
latifolia, *Link, ex Steud. Nom.* ed. II. i. 259.—Lusitan.
linearis, Pursh, Fl. Am. Sept. i. 3 = autumnalis.
macropteryx, *Hegelm. Monog. Callit.* 59.—Austral.
marginata, *Torr. in Pacif. Rail. Rep.* iv. 135.—Calif.; Chili.
Mandonis, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 136.—Ecuador.
microcarpa, *Engelm. ex Hegelm. in Verh. Bot. Ver. Brandenb.* x. (1868) 113.—Cuba.
minima, Hoppe, Taschenb. (1792) 157 = verna.

CALLITRICHE:—
minor, Bellynck, Fl. Namur (1855), 232 = stagnalis ?
Muelleri, *Sond. in Linnaea*, xxviii. (1856) 229.—Austral.
Nuttallii, *Torr. in Pacif. Rail. Rep.* iv. 135.—Am. bor.
obtusangula, *Le Gall, ex Hegelm. Monog. Callit.* 54.—Europ.
occidentalis, *Hegelm. l. c.* 57. t. iv. 5.—Cuba.
ovata, *Rafin. in Med. Rep. New York*, Ser. III. ii. (1811) 408.—Am. bor.
pallens, Bieb. Fl. Taur. Cauc. iii. 6 = hamulata.
pallens, Goldb. in Mém. Soc. Nat. Mosc. v. (1817) 118 = verna.
palustris, Linn. Sp. Pl. 969 = autumnalis.
pedunculata, DC. Fl. Fr. iv. 415 = hamulata.
pedunculata, A. Gray, Man. ex Hegelm. Monog. Callit. 58 = deflexa.
pendunculata, F. Muell. ex Hegelm. l. c. 61 = truncata.
pedunculosa, *Nutt. in Trans. Am. Phil. Soc.* N. S. v. (1837) 140.—Am. bor.
peploides, *Nutt. l. c.* 141.—Am. bor.
platycarpa, Kuetz. in Linnaea, vii (1832) 174 = stagnalis.
plena, *Rafin. in Med. Rep. New York*, Ser. III. ii. (1811) 408.—Am. bor.
polymorpha, *Loennroth, Obs. Crit. Pl. Suec.* 19 (1854).—Scania.
reflexa, Lange, in Cutanda, Fl. Matrit. 292 = hamulata.
sepulta, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 298.—Am. bor. occ.
sessilis, DC. Fl. Fr. iv. 414 = autumnalis, verna.
Sonderi, *Hegelm. in Verh. Bot. Verh. Brandenb.* ix. (1867) 18.—Austral.
spuria, *Rafin. in Med. Repos. New York*, v. (1808) 354.—Am. bor.
stagnalis, *Scop. Fl. Carn.* ed. II. ii. 251.—Europ.; Ind. or.
Stellaria, Rafin. in Med. Rep. New York, Ser. III. ii. (1811) 408 = autumnalis.
stenocarpa, *Hegelm. in Verh. Bot. Ver. Brandenb.* x. (1868) 114.—Calif.
stellata, Hoppe, Taschenb. (1792) 158 = verna.
tenuifolia, Fries, Novit. Fl. Suec. 280 = hamulata.
tenuifolia, Gennari, ex Parl. Fl. Ital. iv. 424 = capillaris.
tenuifolia, Thuil. ex Pers. Syn. i. 6 = verna.
terrestris, DC. Prod. iii. 71 = deflexa.
terrestris, *Muhl. ex Rafin. l. c.* Ser. III. ii. (1811) 408.—Am. bor.
terrestris, Rafin. in Med. Repos. New York, v. (1808) 358 = brevifolia.
transsilvanica, Schur, Enum. Pl. Transs. 216 = stagnalis.
truncata, *Guss. Pl. Rar.* 4.—Italia.
turfosa, *Bert. in Am. Journ. Sc.* xix. (1831) 308, nomen; *Hegelm. Monog. Callit.* 59.—Chili.
turfosa, *Schur, Enum. Pl. Transs.* 216.—Transsylv.
umbonata, *Hegelm. in Verh. Bot. Ver. Brandenb.* ix. (1867) 19.—Tasmania.
verna, Hook. f. Fl. Antarct. i. 11 = antarctica.
verna, *Linn. Fl. Suec.* ed. II. ii. n. 3.—Europ.; Am. bor.
verna, Linn. l. c. partim = polymorpha.
verna, Savi, Fl. Pis. i. 4 = stagnalis.
vernalis, Kuetz. in Linnaea, vii. (1832) 175 = verna.
virens, Goldb. in Mém. Soc. Nat. Mosc. v. (1817) 119 = autumnalis.
Wightiana, Wall. Cat. n. 7008 = stagnalis.

CALLITRIS, Vent. Decad. (1808) 10. *CONIFERAE*, Benth. & Hook. f. iii. 424.
Frenela, Mirb. in Mém. Mus. Par. xiii. (1826) 30.
Fresnelia, Steud. Nom. ed. II. i. 648 (1840).
Leichhardtia, Sheph. Cat. Pl. Cult. Sydn. (1851) 15.
Pachylepis, Brongn. in Ann. Sc. Nat. Sér. I. xxx. (1833) 189.
Octoclinis, E. Muell. in Trans. Phil. Inst. Vict. ii. (1858) 21.
Parolinia, Endl. Gen. Suppl. i. 1372 (1841).
Widdringtonia, Endl. l. c. ii. 25 (1842).
Actinostrobus, F. Muell. Rep. Burdek. Exp. 19 = Actinostrobus pyramidalis.
acuminata, *F. Muell. Census*, 109.—Austral.
arborea, *Schrad. ex E. Mey. Zwei Pfl. Docum.* 73, nomen.—Afr. austr.
arenosa, Sweet, Hort. Brit. ed. II. 473, nomen = rhomboidea.
articulata, Hort. ex Gord. Pinet. ed. II. 117 = rhomboidea.
australis, Sweet, Hort. Brit. ed. II. 474 = rhomboidea.
calcarata, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 74.—Austral.
capensis, Schrad. ex Gord. Pinet. 333 = cupressoides.

CALLITRIS:—
columellaris, F. Muell. Fragm. v. 198 = robusta.
conglobata, *Sieber, ex Heynh. Nom.* i. 148.—Austral.
cupressiformis, Vent. Decad. 10 = rhomboidea.
cupressoides, *Schrad. ex E. Mey. Zwei Pfl. Docum.* 126.—Afr. austr.
Drummondii, *Benth. & Hook. f. ex F. Muell. Census*, 109.—Austral.
elegans, *Sieber, ex Heynh. Nom.* i. 149.—Austral.
Fothergillii, Loud. Encycl. Trees, 1072 = australis.
fruticosa, R. Br. ex Rich. Conif. 49 = calcarata.
glauca, R. Br. ex Mirb. in Mém. Mus. Par. xiii. (1825) 74 = robusta.
Gunnii, Hook. f. in Hook. Lond. Journ. iv. (1845) 147 = australis.
Hugelii, *Knight, Syn. Conif.* 14.—Austral.
Macleayana, *F. Muell. in Rep. Burdek. Exped.* 17; *Extra-Trop. Pl.* Am. ed. (1884) 65.—Austral.
macrostachya, Hort. ex Steud. Nom. ed. II. i. 260 = quadrivalvis.
montana, *Sieber, ex Heynh. Nom.* i. 149.—Austral.
Muelleri, *Benth. & Hook. f. ex F. Muell. Census*, 109.—Austral.
oblonga, *Rich. Conif.* 49. t. 18. f. 2.—Austral.
Parlatorei, *F. Muell. Fragm.* v. 186.—Austral.
Preissii, Miq. in Lehm. Pl. Preiss. i. 643 = robusta.
propinqua, R. Br. ex Endl. Syn. Conif. 38 = robusta.
pyramidalis, Sweet, Hort. Brit. ed. II. 473 = calcarata.
quadrivalvis, *Vent. Decad.* 10.—Afr. bor.
rhomboidea, *R. Br. in Rich. Conif.* 47. t. 18.—Austral.
robusta, *R. Br. ex Mirb. in Mém. Mus. Par.* xiii. (1825) 74.—Austral.
Roei, *Benth. & Hook. f. ex F. Muell. Census*, 109.—Austral.
stricta, Schrad. ex Carr. Conif. ed. I. 66 = cupressoides.
triquetra, Hort. ex Loud. Encyc. Trees, 1072 = quadrivalvis ?
tuberculata, R. Br. ex Endl. Syn. Conif. 38 = robusta.
Ventenatii, R. Br. ex Mirb. in Mém. Mus. Par. xiii. (1825) 74 = rhomboidea.
verrucosa, R. Br. ex Mirb. l. c. = robusta.

CALLIXENE, Comm. ex Juss. Gen. 41 (1789) = **Luzuriaga**, Ruiz & Pav. (Liliac.).
magellanica, Raeusch. Nom. ed. III. 98 = L. marginata.
marginata, Lam. Illustr. t. 248 = L. marginata.
melantha, Colenso, in Trans. N. Z. Inst. xvii. (1885) 250.—N. Zel.
parviflora, Hook. f. in Hook. Ic. Pl. t. 632 = L. parviflora.
polyphylla, Hook. l. c. t. 674 = L. erecta.

CALLOPISMA, Mart. Nov. Gen. et Sp. ii. 107. t. 183, 184 (1827) = **Deianira**, Cham. & Schlecht. (Gentian.).
amplexifolium, Mart. l. c. 109 = D. nervosa.
cordifolium, Lhotsky, ex Griseb. in DC. Prod. ix. 42 = D. erubescens.
perfoliatum, Mart. Nov. Gen. et Sp. ii. 108 = D. erubescens.

CALLOSMIA, Presl, Bot. Bemerk. 103 (1844) = **Anneslea**, Wall. (Ternstroem.).
fragrans, Presl, l. c. = A. fragrans.

CALLOSTYLIS, Blume, Bijdr. 340. t. 74 (1825) = **Eria**, Lindl. (Orchid.).
rigida, Blume, l. c. = E. pulchella.

CALLOTROPIS, G. Don, Gen. Syst. ii. 228 (1832) = **Galega**, Linn. (Legumin.).
tricolor, G. Don, l. c. = G. officinalis.

CALLUNA, Salisb. in Trans. Linn. Soc. vi. (1802) 317. *ERICACEAE*. Benth. & Hook. f. ii. 589.
atlantica, Seem. in Journ. Bot. iv. (1866) 306 = vulgaris.
ciliaris, Schur, Enum. Pl. Transs. 447 = vulgaris.
Erica, DC. Fl. Fr. iii. 680 = vulgaris.
genuina, Ducommun, Taschenb. 491 = vulgaris.
sagittaefolia, S. F. Gray, Nat. Arr. ii. 399 = vulgaris.
vulgaris, *Salisb. in Trans. Linn. Soc.* vi. (1802) 317.—Europ.; Am. bor.

CALOBOTA, Eckl. & Zeyh. Enum. 192 (1836) = **Lebeckia**, Thunb. (Legumin.).
cytisoides, Eckl. & Zeyh. l. c. 191 = L. cytisoides.
decipiens, Benth. in Ann. Wien. Mus. ii. (1838) 142 = L. decipiens.
decutiens, Benth. l. c. = L. decutiens.
flexuosa, Benth. l. c. = L. flexuosa.
multiflora, Benth. l. c. = L. multiflora.
pulchella, Eckl. & Zeyh. Enum. 191 = L. cytisoides.

CALOBOTRYA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 21 = **Ribes**, Linn. (Saxifr.).
sanguinea, Spach, l. c. = R. sanguineum.

CALOCEDRUS, Kurz, in Journ. Bot. xi. (1873) 196. t. 133 = **Libocedrus**, Endl. (Conifer.).
macrolepis, Kurz, l. c. = L. macrolepis.

CALOCEPHALUS, R. Br. in Trans. Linn. Soc. xii. (1817) 106. *COMPOSITAE*, Benth. & Hook. f. ii. 520.
ACHRYSUM, A. Gray, in Hook. Kew Journ. iv. (1852) 228.
BLENNOSPORA, A. Gray, l. c. iii. (1851) 172.
LEUCOPHYTA, R. Br. in Trans. Linn. Soc. xii. (1817) 106.
PACHYSURUS, Steetz, in Lehm. Pl. Preiss. i. 441 (1845).
aeruoides, *Benth. Fl. Austral.* iii. 576.—Austral.
angianthoides, *Benth. l. c.* 575.—Austral.
Brownii, *F. Muell. Rep. Babb. Exp.* 13.—Austral.
chrysanthus, Schlecht. in Linnaea, xx. (1847) 592 = Craspedia chrysantha.
citreus, *Less. Syn. Comp.* 271.—Austral.
Drummondii, *Benth. Fl. Austral.* iii. 574.—Austral.
Francisii, *Benth. l. c.* 576.—Austral.
gnaphalioides, Hook. in Mitch. Journ. Trop. Austral. 378 = Helipterum moschatum.
lacteus, *Less. Syn. Comp.* 271.—Austral.
multiflorus, *Benth. Fl. Austral.* iii. 576.—Austral.
platycephalus, *Benth. l. c.*—Austral.
Sonderi, *F. Muell. 2nd Gen. Rep.* (1854); *et Rep. Babb. Exp.* 13.—Austral.

CALOCHILUS, R. Br. Prod. 320 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 614.
campestris, *R. Br. l. c.*—Austral.
herbaceus, Lindl. Gen. et Sp. Orch. 459 = campestris.
paludosus, *R. Br. Prod.* 320.—Austral.
Robertsoni, *Benth. Fl. Austral.* vi. 315.—Austral.

CALOCHLAMYS, Presl, Bot. Bemerk. 148 (1844) = **Congea**, Roxb. (Verbenac.).
capitata, Presl, l. c. 149 = Congea tomentosa.

CALOCHORTUS, Pursh, Fl. Am. Sept. i. 240 (1814). *LILIACEAE*, Benth. & Hook. f. iii. 820.
CYCLOBOTHRA, G. Don, in Sweet, Brit. Flow. Gard. t. 273 (1829).
albus, *Dougl. ex Benth. in Maund & Hensl. Botanist*, t. 98.—Calif.
apiculatus, *Baker, in Journ. Linn. Soc.* xiv. (1875) 305.—Am. bor. occ.
aureus, *S. Wats. in Am. Natural.* vii. (1873) 303.—Calif.
Barnardi, *Dougl. ex Steud. Nom.* ed. II. i. 260.—Am. bor. occ.
Benthami, *Baker, in Journ. Linn. Soc.* xiv. (1875) 304.—Calif.
Bonplandianus, *Schult. f. Syst.* vii. 1532.—Mexic.
caeruleus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 263.—Calif.
Catalinae, *S. Wats. l. c.* 268.—Calif.
citrinus, *Baker, in Bot. Mag.* t. 6200.—Calif.
clavatus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 265.—Calif.
Douglasianus, *Schult. f. in Van Hall, Bijdr.* iv. 127.—Am. bor. occ.
elegans, Baker, in Journ. Linn. Soc. xiv. (1875) 305 = lilacinus, nudus et Tolmiei.
elegans, Hook. f. Bot. Mag. t. 5976 = Maweanus.
elegans, *Pursh, Fl. Am. Sept.* i. 240.—Am. bor. occ.
eurycarpus, S. Wats. in Bot. King, Exp. 348 = nitidus.
flavus, *Schult. f. Syst.* vii. 1535.—Mexic.
flexuosus, *S. Wats. in Am. Natural* vii. (1873) 303.—Am. bor. occ.
fuscus, *Schult. f. Syst.* vii. 1534.—Mexic.

CALOCHORTUS:—
Ghiesbreghtianus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 268.—Mexic.
glaucus, Regel, Gartenfl. (1875) 260; et in Act. Hort. Petrop. iii. II. (1875) 285 = caeruleus.
Greenei, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 264.—Am. bor. occ.
Gunnisoni, *S. Wats. in Bot. King, Exp.* 348.—Am. bor. occ.
Hartwegii, *Benth. Pl. Hartw.* 26.—Mexic.
Kennedyi, *Porter, in Coult. Bot. Gaz.* ii. (1877) 79.—Calif.
Leichtlinii, Hook. f. Bot. Mag. t. 5862 = Nuttallii.
lilacinus, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 5.—Calif.
longebarbatus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 381.—Am. bor. occ.
luteus, *Dougl. ex Kunth, Enum. Pl.* iv. 233.—Calif.
luteus, Nutt. in Journ. Acad. Phil. vii. (1834) 53 = Nuttallii.
Lyallii, Baker, in Journ. Linn. Soc. xiv. (1875) 305 = elegans.
macrocarpus, *Dougl. in Trans. Hort. Soc.* vii. (1830) 276. t. 8.—Am. bor. occ.
Maweanus, *Leichtl. ex Baker, in Journ. Linn. Soc.* xiv. (1875) 305.—Am. bor. occ.
monophyllus, *Lem. in Fl. des Serres*, Sér. I. v. (1849) 430 b.—Calif.
nitidus, *Dougl. in Trans. Hort. Soc.* vii. (1830) 277. t. 9.—Calif.
nitidus, Torr. Pacif. Rail. Rep. iv. 146 = Benthami.
nudus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 263.—Calif.
Nuttallii, *Torr. & Gray, in Pacif. Rail. Rep.* ii. 124.—Am. bor. occ.
Palmeri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 266.—Calif.
pallidus, Schult. f. Syst. vii. 1533 = flavus.
parviflorus, Nutt. ex Baker, in Journ. Linn. Soc. xiv. (1875) 306 = Nuttallii.
pulchellus, *Dougl. ex Benth. in Trans. Hort. Soc.* Ser. II. i. (1835) 412.—Calif.
purpureus, Baker, in Journ. Linn. Soc. xiv. (1875) 308, pro parte = Bonplandianus.
pusillus, *Dougl. ex Steud. Nom.* ed. II. i. 260.—Am. bor. occ.
spatulatus, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 267.—Mexic.
splendens, *Dougl. ex Benth. in Trans. Hort. Soc.* Ser. II. i. (1835) 411. t. 15.—Calif.
Tolmiei, *Hook. & Arn. Bot. Beech. Voy.* 398.—Calif.
umbellatus, Wood, in Proc. Nat. Sc. Philad. (1868) 168 = lilacinus.
uniflorus, *Hook. & Arn. Bot. Beech. Voy. Suppl.* 398.—Calif.
uniflorus, Hook. f. Bot. Mag. t. 5804 = lilacinus.
venustus, *Dougl. ex Benth. in Trans. Hort. Soc.* Ser. II. i. (1835) 412 t. 15. f. 3.—Calif.
vestitus, *Benth. ex Steud. Nom.* ed. II. i. 260.—Calif.
Weedii, Wood, in Proc. Nat. Sc. Philad. (1868) 169 = luteus.

CALOCOCCUS, Kurz, ex Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind. xxvi. (1864) 48 = **Phyllanthus**, Linn. (Euphorbiac.).
sundaicus, Kurz, l. c. = P. sundaicus.

CALODENDRUM, Thunb. Nov. Gen. ii. 41 (1782). *RUTACEAE*, Benth. & Hook. f. i. 288.
PALLASIA, Houtt. Handleid. ii. 382 (1775).
capensis, *Thunb. l. c.* 43; *et Prod. Pl. Cap.* 44.—Afr. austr.

CALODIUM, Lour. Fl. Cochinch. 247 (1790) = **Cassytha**, Linn. (Laurin.).
cochinchinensis, Lour. l. c. = Cas. filiformis.

CALODONTA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 448 = **Tolpis**, Adans. (Compos.).
azorica, Nutt. l. c. = T. virgata.

CALODRACON, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 132, 137 = **Cordyline**, Comm. (Liliac.).
heliconiaefolia, Planch. l. c. = Cord. heliconiaefolia.
Jacquini, Planch. l. c. = Cord. Jacquinii.
nobilis, Planch. l. c. vii. (1851–52) 105. t. 682, 683.—Japon.
Sieberi, Planch. l. c. vi. (1850–51) 132 = Cord. Sieberi.

CALODRACON:—
Sieboldii, Planch. in Fl. des Serres, Sér. I. vi. 137 = Dracaena Sieboldii.
terminalis, Planch. l. c. = Cord. terminalis.
venusta, Planch. l. c. 132 = Dracaena venusta.

CALODRYUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 401. t. 51 = **Quivisia**, Comm. (Meliac.).
tubiflorum, Desv. l. c. = Q. lanceolata.

CALOGYNE, R. Br. Prod. 579 (1810). *GOODENOVIEAE*, Benth. & Hook. f. ii. 539.
DISTYLIS, Gaudich. in Freyc. Voy. Bot. 460. t. 80 (1826).
Berardiana, *F. Muell. Fragm.* vi. 7.—Austral.
chinensis, *Benth. in Journ. Linn. Soc.* v. (1861) 78.—China.
distylis, F. Muell. Fragm. vi. 6 = Berardiana.
heteroptera, *F. Muell. Fragm.* x. 43.—Austral.
pilosa, *R. Br. Prod.* 579.—Austral.
purpurea, *F. Muell. Fragm.* viii. 57.—Austral.

CALOMECON, Spach, Hist. Vég. Phan. vii. 7 (1839) = **Papaver**, Tourn.
bracteatum, Spach, l. c. 9 = P. bracteatum.
orientale, Spach, l. c. = P. orientale.

CALOMERIA, Vent. Jard. Malm. 73. t. 73 (1804) = **Humea**, Sm. (Compos.).
amaranthoides, Vent. l. c. = H. elegans.

CALOMORPHE, Kunze, ex Walp. Rep. i. 680 (1840) = **Lennea**, Klotzsch (Legumin.).
Ehrenbergiana, Kunze, ex Walp. l. c. = L. robinioides.

CALOMYRTUS, Blume, Mus. Bot. Lugd. Bat. i. 76, in obs. (1849) = **Myrtus**, Tourn.
excelsa, Blume, l. c. = M. excelsa.
ovalifolia, Blume, l. c. = M. ovalifolia.

CALONNEA, Buchoz, Icon. (1786) t. 126 = **Gaillardia**, Fouger. (Compos.).
pulcherrima, Buchoz, l. c. = G. pulchella.

CALONYCTION, Choisy, in Mém. Soc. Phys. Genèv. vi. 441 (1833) (Conv. Or. 59) = **Ipomoea**, Linn. (Convolvul.).
acanthocarpum, Choisy, in DC. Prod. ix. 346 = I. acanthocarpa.
asperum, Choisy, Conv. Or. 60 = I. grandiflora.
Bona-nox, Boj. Hort. Maurit. 228 = I. Bona-nox.
capillatum, Miq. Fl. Ind. Bat. ii. 598 = I. capillata.
chenopodifolium, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 269 = I. chenopodifolia.
clavatum, G. Don, Gen. Syst. iv. 264.—Ecuador.
comorensis, Boj. Hort. Maurit. 228 = I. grandiflora.
comespermum, Boj. l. c. = I. grandiflora.
diversifolium, Hassk. Pl. Jav. Rar. 523 = I. trichosperma.
dubium, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 268 = I. dubia.
Galeottii, Martens, ex Mart. & Gal. l. c.—Mexic.
grandiflorum, Choisy, Conv. Or. 60 = I. grandiflora.
Jacquinii, G. Don, Gen. Syst. iv. 264 = I. grandiflora.
longiflorum, Hassk. Pl. Jav. Rar. 523 = I. grandiflora.
macrantholeucum, Colla, N. Sp. Calon. 15 = I. Bona-nox.
macranthum, Van Houtte, ex Poit. in Rev. Hortic. Sér. II. iv. (1845–46) 409.—Hab. ?
megalocarpum, A. Rich. Fl. Cub. Fanerog. ii. 129 = I. Bona-nox.
mollissimum, Zoll. Syst. Verz. Ind. Archip. 128, 131.—Java.
muricatum, G. Don, Gen. Syst. iv. 264 = I. Bona-nox.
muticum, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 390 = I. grandiflora.
noctolucum, G. Don, in Sweet, Hort. Brit. ed. III. 482 = I. latiflora.
proximum, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 268 = I. proximum.
pseudomuricatum, G. Don, Gen. Syst. iv. 264 = I. grandiflora.
pteripes, G. Don, l. c.—Ecuador.
pterodes, Choisy, in DC. Prod. ix. 346 = praec.
pulcherrimum, Parodi, Contrib. 12.—Paraguay.
Rheedi, Colla, N. Sp. Calon. 15 = I. Bona-nox.
Roxburghii, G. Don, Gen. Syst. iv. 263 = I. Bona-nox.
sanguineum, Hort.—Cf. Gard. Chron. (1865) 934 = Lettsomia Kurzii.
sanguinolentum, Guillon, in Rev. Hortic. (1880) 20.—Hab. ?

CALONYCTION :—
speciosum, Choisy, Conv. Or. 59 = I. Bona-nox.
trichospermum, Choisy, l. c. 60 = I. trichosperma.
Tuba, Colla, Nov. Sp. Calon. 15 = I. grandiflora ?
venustum, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 269 = I. venustum.

CALOPAPPUS, Meyen, Reisc, i. 315 (1834) = **Nassauvia**, Juss. (Compos.).
acanthifolius, Remy, in C. Gay, Fl. Chil. iii. 298.—Chili.
acerosus, Meyen, Reise, i. 315 = N. acerosa.

CALOPETALON, J. Drum. ex Harv. in Hook. Kew Journ. vii. (1855) 52 = **Marianthus**, Hueg. (Pittospor.).
ringens, J. Drum. l. c. 53 = M. ringens.

CALOPHACA, Fisch. Hort. Gorenk. ed. II. 67 (1812); et in DC. Prod. ii. 270 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 505.
CHESNEYA, Lindl. in Endl. Gen. 1275 (1840).
Hovenii, *Schrenk, Enum. Pl. Nov.* 74.—Soongaria.
juncea, Steud. Nom. ed. II. i. 260, sphalm. = Caligonum Calliphysa.
soongorica, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 401 = Hovenii.
wolgarica, *Fisch. Hort. Gorenk.* ed. II. (1812) 68.—Ross. austr.

CALOPHANES, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. t. 181 (1833). *ACANTHACEAE*, Benth. & Hook. f. ii. 1077.
CHAETACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836).
DYSCHORISTE, Necs, in Wall. Pl. As. Rar. iii. 75 (1832), nomen prius.
LINOSTYLES, Fenzl, in Linnaea, xxiii. (1850) 94.
adscendens, *Hochst. ex Nees, in DC. Prod.* xi. 111.—Abyss.
amoenus, *Nees, Mart. Fl. Bras.* ix. 27.—Bras.
angustus, *A. Gray, Syn. Fl. N. Am.* ii. 1. 456.—Florida.
angustifolius, *Hemsl. Biol. Centr. Am. Bot.* ii. 502.—Mexic.
biflorus, Shuttlew. ex Nees, in DC. Prod. xi. 107 = oblongifolius.
bilabiatus, *Seem. Bot. Voy. Herald*, 324. t. 65.—Mexic.
Buchenavii, *Vatke, in Bremen Abh.* ix. (1885) 131.—Madag.
Burkei, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 24.—Afr. austr.
capitatus, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 121.—Mexic.
Clarkei, *Vatke, in Bremen Abh.* ix. (1885) 132.—Ins. Mascar.
ciliatus, *Nees, in DC. Prod.* xi. 110.—Peruv.
costatus, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Afr. austr.
crinitus, *Nees, in Mart. Fl. Bras.* ix. 26.—Bras.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 160.—Cuba.
Dalzellii, *T. Anders. ex Bedd. Ic. Ind. Or.* i. 60.—Ind. or.
decumbens, *A. Gray, Syn. Fl. N. Am.* ii. 1. 325.—Arizon.; Mexic.
depressus, T. Anders. in Thw. Enum. Pl. Zeyl. 225 = Nagchana.
gracilis, *Nees, in DC. Prod.* xi. 111.—Madag.
Heudelotianus, *Nees, l. c.* 112.—Afr. trop.
Hildebrandtii, *S. Moore, in Journ. Bot.* xviii. (1880) 8, 197.—Afr. trop.
hirsutissimus, *Nees, in DC. Prod.* xi. 109.—Mexic.
hirsutus, *Oerst. ex Hiern, in Kjoeb. Vidensk. Meddel.* (1877-78) 71.—Mexic.
humistratus, *Shuttlew. ex Nees, in DC. Prod.* xi. 108.—Am. bor.
hygrophyloides, *Nees, in Mart. Fl. Bras.* ix. 26.—Bras.
hyssopifolius, *Nees, in DC. Prod.* xi. 111.—Abyss.
Jasminum-mexicanum, *Nees, l. c.* 110.—Mexic.
lavandulaceus, *Nees, in Mart. Fl. Bras.* ix. 27.—Bras.
linearis, *A. Gray, Syn. Fl. N. Am.* ii. 1. 324.—Texas; Mexic.
littoralis, *T. Anders. in Thw. Enum. Pl. Zeyl.* 225.—Ind. or.
madagascariensis, *Nees, in DC. Prod.* xi. 112.—Madag.
Maranhonis, *Nees, in Mart. Fl. Bras.* ix. 25.—Bras.

CALOPHANES:—
mauritianus, T. Anders. in Journ. Linn. Soc. vii. (1864) 23, nomen = Hypoëstes serpens.
microphyllus, *Nees, in DC. Prod.* xi. 113.—Mexic.
multicaulis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Abyss.
Nagchana, *Nees, in DC. Prod.* xi. 109.—Ind. or.
natalensis, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Afr. austr.
oblongifolius, *D. Don, in Sweet, Flow. Gard.* t. 181.—Am. bor.
oblongifolius, Torr. Bot. Mex. Bound. 122 = decumbens.
ovatus, Benth. Pl. Hartw. 89 = linearis.
Perrottetii, *Nees, in DC. Prod.* xi. 111.—Afr. trop.
Persoonii, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Afr. austr.
Pulegium, *Nees, in Mart. Fl. Bras.* ix. 25.—Bras.
quadrangularis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1854) 120.—Mexic.
quitensis, *Nees, in DC. Prod.* xi. 110.—Ecuador.
radicans, *T. Anders. in Journ. Linn. Soc.* vii. (1864) 23.—Afr. trop.
repens, *Nees, in DC. Prod.* xi. 109.—Peruv.
rigidus, Dalz. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 411 = Dalzellii.
Schiedeanus, *Nees, in DC. Prod.* xi. 111.—Mexic.
Serpyllum, *Nees, in Mart. Fl. Bras.* ix. 26.—Bras.
setosus, *Nees, in DC. Prod.* xi. 112.—Afr. austr.
siphonanthus, *Nees, l. c.*—Madag.
thunbergiaeflora, *S. Moore, in Journ. Bot.* xviii. (1880) 8, 197.—Afr. trop.
trichocalyx, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 126.—Afr. trop.
Tweedianus, *Nees, in DC. Prod.* xi. 108.—Reg. Argent.
vagans, *Wight, Ic.* xi. t. 1526.—Ind. or.
verticillaris, *T. Anders. ex Oliver, in Trans. Linn. Soc.* xxix. (1874) 126, in nota.—Afr. trop.

CALOPHTHALMUM, Reichb. Nom. 82 (1841) = GALOPHTHALMUM, Nees & Mart.

CALOPHYLICA, Presl, Bot. Bemerk. 39 (1844) = **Phylica**, Linn. (Rhamn.).
gnidioides, Presl, l. c. = P. gnidioides.
juniperifolia, Presl, l. c. = P. gnidioides.

CALOPHYLLOIDES, Smeathm. ex DC. Prod. iii. 272 (1828) = **Eugenia**, Mich. (Myrtac.).
lucida, Smeathm. l. c. = E. Calophylloides.

CALOPHYLLUM, Linn. Gen. ed. I. 154 (1737). *GUTTIFERAE*, Benth. & Hook. f. i. 175.
APOTERIUM, Blume, Bijdr. 218 (1825).
AUGIA, Lour. Fl. Cochinch. 337 (1790).
acuminatum, Lam. Encyc. i. 553 = spectabile.
acuminatum, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 80 = Rheedia acuminata.
Akara, Burm. f. Fl. Ind. 121 = Tetracera laevis.
amoenum, Wall. Cat. n. 4849 = retusum.
amplexicaule, *Choisy, ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 281.—Ins. Philipp.
angustifolium, *Roxb. Hort. Beng.* 41; *Fl. Ind.* ii. 608.—Penang.
apetalum, *Blanco, Fl. Filip.* ed. II. 429.—Ins. Philipp.
apetalum, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 79 = C. Inophyllum ?
Augia, *Steud. Nom.* ed. II. i. 260.—China.
bancanum, Miq. Fl. Ind. Bat. Suppl. i. 499 = pulcherrimum.
Bingator, Roxb. Hort. Beng. 41; Fl. Ind. ii. 607 = C. Inophyllum.
Blancoi, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 272.—Ins. Philipp.
Blumei, *Wight, Illustr.* i. 128.—Java.
bracteatum, *Thw. Enum. Pl. Zeyl.* 51.—Zeylan.
brasiliense, *Camb. in St. Hil. Fl. Bras. Mer.* i. 321. t. 67.—Bras.
Burmanni, Wight, Illustr. i. 129 = C. Calaba, *Linn.*
Calaba, *Jacq. Am.* 269. t. 165.—Ins. Carib.
Calaba, *Linn. Sp. Pl.* 514.—Zeylan.
calaboides, *G. Don, Gen. Syst.* i. 622.—Ind. or.
calcdonicum, *Vieill. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 291.—N. Caled.
canum, *Hook. f. Fl. Brit. Ind.* i. 271.—Malacca.
chinense, T. Anders. in Hook. f. Fl. Brit. Ind. i. 270 = Ochrocarpus longifolius.
cordato-oblongum, *Thw. Enum. Pl. Zeyl.* 407.—Zeylan.

CALOPHYLLUM:—
Cumingii, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 259.—Ins. Philipp.
cuncifolium, *Thw. Enum. Pl. Zeyl.* 51, 407.—Zeylan.
Cupi, H. B. & K. Nov. Gen. et Sp. v. 203 = Rheedia Cupi.
Cussi, G. Don, Gen. Syst. i. 623 = praec.
cymosum, Miq. Fl. Ind. Bat. Suppl. i. 497 = spectabile.
dasypodium, *Miq. Fl. Ind. Bat.* i. II. 511.—Java.
decipiens, Thw. Enum. Pl. Zeyl. 51 = Thwaitesii.
decipiens, Wight, Illustr. 128; Ic. i. 106 = Wightianum.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* i. 497.—Sumatra.
dongnaiense, *Pierre, Fl. For. Cochinch.* t. 108.—Cochinch.
dryobalanoides, *Pierre, l. c.* t. 106.—Cochinch.
dubium, Steud. Nom. ed. II. i. 261 = Ochrocarpus longifolius.
edule, Seem. Bot. Voy. Herald, 89 = Rheedia edulis.
elatum, Bedd. Fl. Sylv. i. t. 2 = tomentosum.
excelsum, Zoll. & Morr. in Nat. en Geneesk. Arch. Neerl. Indie, ii. (1845) 582 = Ochrocarpus ovalifolius.
floribundum, *Hook. f. Fl. Brit. Ind.* i. 272.—Malacca.
gracile, Miq. Fl. Ind. Bat. Suppl. i. 498 = pulcherrimum.
Griffithii, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 273.—Malacca.
Hasskarlii, *Teijsm. & Binn. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 273.—Java.
hexapetalum, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 163.—Borneo.
hirtellum, *Miq. Pl. Jungh.* 291.—Java.
Inophyllum, Lam. Encyc. i. 552 = C. Tacamahaca.
Inophyllum, *Linn. Sp. Pl.* 513.—Geront. trop.
Inophyllum, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = C. Calaba, *Jacq.*
javanicum, *Miq. Pl. Jungh.* 292.—Java.
lanceolarium, Roxb. Hort. Beng. 41; Fl. Ind. ii. 608 = C. Tacamahaca.
lanceolatum, Blume, Bijdr. 217 = C. Tacamahaca.
lanceolatum, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 395.—Java.
lanigerum, *Miq. Fl. Ind. Bat. Suppl.* i. 498.—Ins. Banca.
longifolium, Wall. Cat. n. 4851 = Ochrocarpus longifolius.
longifolium, *Willd. Ges. Naturf. Fr. Berl. Mag.* v. (1811) 80.—N. Granat.
Lowei, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 271.—Borneo.
lucidum, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 370.—Guiana.
macrocarpum, *Hook. f. Fl. Brit. Ind.* i. 273.—Malacca.
macrophyllum, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 405.—Malaya.
Madruño, H. B. & K. Nov. Gen. et Sp. v. 202 = Rheedia Madruno.
marginatum, *Wall. Cat.* n. 4845 (sp. ?).—Ins. Penang.
Mariae, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xv. (1861) 251.—Am. anstr.
membranaceum, *Gardn. & Champ. in Hook. Kew Journ.* i. (1849) 309.—China.
mesuaefolium, Wall. Cat. n. 4850 = pulcherrimum.
microphyllum, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 272.—Malaya.
microphyllum, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 282 = parvifolium.
microphyllum, *Scheff. in Tijdschr. Nederl. Ind.* xxxii. (1871) 406.—Ins. Gebeh.
montanum, *Vieill. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 292.—N. Caled.
Moonii, Wight, Illustr. i. 129 = spectabile.
Nagassarium, Burm. f. Fl. Ind. 121 = Mesua ferrea.
obovale, *Miq. Fl. Ind. Bat. Suppl.* i. 498.—Sumatra.
odoratissimum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 12 = spectabile.
ovatifolium, Noronha, l. c. 13 = C. Inophyllum.
pachyphyllum, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 255.—Bras.
parvifolium, *Boj. ex Baker, Fl. Maurit.* 16.—Ins. Maurit.
parvifolium, *Choisy, in Mém. Soc. Hist. Nat. Par.* i. (1823) 229.—Ins. Molucc.
pisiferum, Planch. & Triana, in Ann. Sc. Nat. Ser. IV. xv. (1861) 294 = retusum.
plicipes, Miq. Fl. Ind. Bat. Suppl. i. 499 = pulcherrimum.

CALOPHYLLUM:—
polyanthum, *Wall. Cat.* n. 4844; *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 278.—Ind. or.
pseudotacamacha, *Planch. & Triana, l. c.* 270.—Ind. or.
pulcherrimum, *Wall. Cat.* n. 4848; *Planch. & Triana, l. c.* 274.—Malaya.
retusum, *Wall. Cat.* n. 4846; *Planch. & Triana, l. c.* 265.—Malaya.
rigidum, *Miq. Fl. Ind. Bat. Suppl.* i. 497.—Sumatra.
saigonense, *Pierre, Fl. For. Cochinch.* t. 105.—Cochinch.
Soualatti, Burm. f. Fl. Ind. 121 = spectabile.
spectabile, Hook. & Arn. Bot. Beech. Voy. 174 = membranaceum.
spectabile, *Willd. in Ges. Naturf. Fr. Berl. Mag.* v. (1811) 80.—Malaya; Polynes.
spurium, Boj. Hort. Maurit. 52 = parviflorum.
spurium, Choisy, in Mém. Soc. Hist. Nat. Par. i. (1823) 229 = C. Inophyllum.
Suriga, *Buch.-Ham. ex Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 608.—Ind. or.
Suringa, Buch.-Ham. ex Steud. Nom. ed. II. i. 261 = praec.
Tacamahaca, *Willd. in Ges. Naturf. Fr. Berl. Mag.* v. (1811) 79.—Borbon.; Madag.
tetrapetalum, Roxb. Hort. Beng. [93]; Fl. Ind. ii. 608 = spectabile.
tetrapetalum, Wall. Cat. n. 4843 = Wallichianum.
tetrapterum, *Miq. Pl. Jungh.* 291.—Sumatra.
Teysmanni, *Miq. Fl. Ind. Bat. Suppl.* i. 499.—Sumatra.
Teysmannii, Zoll. ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xv. (1861) 275 = lanceolatum.
thuriferum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 16.—Peruv.
Thorelii, *Pierre, Fl. For. Cochinch.* t. 103.—Cochinch.
Thwaitesii, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 260.—Zeylan.
tomentosum, *Wight, Illustr.* i. 128.—Ind. or.
trapezifolium, *Thw. Enum. Pl. Zeyl.* 51.—Zeylan.
venulosum, *Zoll. Syst. Verz. Ind. Archip.* 149, 150.—Java.
Walkeri, *Wight, Illustr.* i. 128.—Ind. or.
Wallichiana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 277.—Malaya.
Wightianum, *Wall. Cat.* n. 4847; *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 256.—Ind. or.

CALOPHYSA, DC. Prod. iii. 166 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 766.
ciliata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 140.—Am. trop.
dentata, *Triana, l. c.*—Peruv.
dispar, *Triana, l. c.* 141.—Peruv.
flexuosa, *Triana, l. c.*—Venezuela.
heterophylla, *Triana, l. c.* 140.—Peruv.
pilosa, *Triana, l. c.*—Am. trop.
setosa, *Triana, in Journ. Bot.* v. (1867) 209.—Nicaragua.
testiculata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 140.—N. Granat.
tococoidea, *DC. Prod.* iii. 166.—Bras.
vesiculosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 140.—Mexic.

CALOPLECTUS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 119 = **Alloplectus**, Mart. (Melastom.).
macrophyllus, Oerst. l. c. = A. macrophyllus.

CALOPOGON, R. Br. in Ait. Hort. Kew. ed. II. v. 204 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 615.
CATHEA, Salisb. in Trans. Hort. Soc. i. (1812) 300, nomen prius.
Lessonii, Brongn. Bot. Voy. Coquille, 188. t. 37 = Pogonia tetraphylla.
multiflorus, *Lindl. Gen. et Sp. Orch.* 425.—Am. bor.
pallidus, *Chapman, Fl. S. U. St.* 457.—Am. bor.
parviflorus, *Lindl. Gen. et Sp. Orch.* 424.—Am. bor.
parviflorus, *Rafin. Atl. Journ.* 148 (= praec.?).—Am. bor.
pulchellus, *R. Br. in Ait. Hort. Kew.* ed. II. v. 204.—Am. bor.

CALOPOGONIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 423. *LEGUMINOSAE*, Benth. & Hook. f. i. 534.
CYANOSTREMMA, Benth. ex Hook. & Arn. Bot. Beech. Voy. 415 (1841).

CALOPOGONIUM:—
STENOLOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 125.
brachycarpum, Benth. ex Hemsl. in Biol. Centr. Am. Bot. i. 300 = mucunoides.
caeruleum, *Hemsl. l. c.* 301.—Mexic.
galactoides, *Benth. ex Hemsl. l. c.*—Mexic.
mucunoides, *Desv. in Ann. Sc. Nat.* Sér. I. ix. (1826) 423.—Guiana.

CALOPSIS, Beauv. ex Juss. in Dict. Sc. Nat. xlv. 272 (1827); et ex Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 44. t. 3 = **Leptocarpus**, R. Br. (Restiac.).
chilensis, Steud. Syn. Pl. Cyp. 258 = L. chilensis.
festucacea, Kunth, Enum. Pl. iii. 425 = L. incurvatus.
hirtella, Kunth, l. c. 426 = L. peronatus.
incurvata, Kunth, l. c. 427 = L. incurvatus.
neglecta, Hochst. in Flora, xxviii. (1845) 338 = L. neglectus.
oxylepis, Kunth, Enum. Pl. iii. 427 = L. oxylepis.
paniculata, Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 44. t. 3. f. 2 = L. paniculatus.
peronata, Kunth, Enum. Pl. iii. 426 = Restio ocreatus et L. peronatus.
ramiflora, Kunth, l. c. 423 = L. paniculatus.
triticea, Kunth, l. c. 424 = Restio triticeus.

CALOPTILIUM, Lag. Amen. Nat. i. 34 (1811) = **Nassauvia**, Juss. (Compos.).
Lagascae, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 36 = N. Lagascae.

CALOPYXIS, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 86 = **Combretum**, Linn.
alata, Tul. l. c. 89.—Madag.
eriantha, Tul. l. c. 90.—Madag.
oxygonia, Tul. l. c. 88.—Madag.
sphaeroides, Tul. l. c. 86.—Madag.
velutina, Tul. l. c. 87.—Madag.

CALORCHIS, Rodrig. Orch. Nov. i. 195 (1877). *ORCHIDEAE*.
phaenoleuca, *Rodrig. l. c.* 196.—Bras.

CALORHABDOS, Benth. Scroph. Ind. 44 (1835). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 963.
Brunoniana, *Benth. l. c.*—Reg. Himal.
cauloptera, *Hance, in Journ. Bot.* xv. (1877) 298.—China.

CALOROPHUS, Labill. Nov. Holl. Pl. ii. 78. t. 228 (1806) = **Hypolaena**, R. Br. (Restiac.).
asper, Nees, in Lehm. Pl. Preiss. ii. 67 = Restio Loxocarya.
crispatus, Nees, l. c. = Restio crispatus.
densus, Nees, l. c. = H. densa.
dentatus, Nees, l. c.—Austral.
elongatus, Labill. Nov. Holl. Pl. ii. 78. t. 228 = H. elongata.
elongatus, F. Muell. Fragm. viii. 86 = H. elongata.
elongatus, Nees, in Lehm. Pl. Preiss. ii. 68 = H. gracillima.
flexuosus, Nees, in Lehm. Pl. Preiss. ii. 68 = H. flexuosa.
minor, Hook. f. Fl. N. Zel. i. 267 = H. elongata.
Sieberianus, Steud. Syn. Pl. Cyp. 265 = Restio fastigiatus.

CALOSACME, Wall. Cat. nn. 800–806 (1829) = **Chirita**, Buch.-Ham. (Gesnerac.).
acuminata, Wall. l. c. n. 802 = Ch. acuminata et Didymocarpus acuminata.
amplectens, Wall. l. c. n. 806 = Ch. bifolia.
dimidiata, Wall. l. c. n. 803 = Ch. dimidiata.
flava, Wall. l. c. n. 801 = Ch. pumila.
grandiflora, Wall. l. c. n. 800 = Ch. urticaefolia.
macrophylla, Wall. l. c. n. 805 = Ch. macrophylla.
polycarpa, Wall. l. c. n. 804 = Lysionotus serrata.

CALOSANTHES, Blume, Bijdr. 760 (1826) = **Oroxylum**, Vent. (Bignoniac.).
indica, Blume, l. c. = O. indicum.

CALOSANTHOS, Reichb. in Moessl. Handb. ed. II. i. 50 (1827) = KALOSANTHES, Haw. = **Rochea**, DC. (Crassul.).

CALOSCIADIUM, Endl. Gen. Suppl. iv. III. 7 (1850) = **Aciphylla**, Forst. (Umbellif.).
antipodum, Endl. ex Walp. Ann. ii. 702.—N. Zel.
latifolium, Endl. ex Walp. l. c.—N. Zel.

CALOSCILLA, Jord. & Fourr. Ic. Fl. Eur. ii. 14. t. 245 (1869) = **Scilla**, Linn. (Liliac.).
Clusii, Jord. & Fourr. l. c. = S. peruviana.
elegans, Jord. & Fourr. l. c. 16. t. 250 = S. peruviana.
flaveola, Jord. & Fourr. l. c. 17. t. 254 = S. peruviana.
grandiflora, Jord. & Fonrr. l. c. 16. t. 252 = S. peruviana.
hipponensis, Jord. & Fourr. l. c. 17. t. 255 = S. peruviana.
Hughii, Jord. & Fourr. l. c. 15. t. 246 = S. peruviana.
livida, Jord. & Fourr. l. c. = S. peruviana.
pallidiflora, Jord. & Fourr. l. c. t. 248 = S. peruviana.
subalbida, Jord. & Fourr. l. c. 16. t. 253 = S. peruviana.
subcarnea, Jord. & Fourr. l. c. t. 250 = S. peruviana.
venusta, Jord. & Fourr. l. c. 15. t. 247 = S. peruviana.

CALOSCORDUM, Herb. in Lindl. Bot. Reg. (1844) Misc. 66 = **Nothoscordum**, Kunth (Liliac.).
neriniflorum, Herb. l. c. 67 = N. neriniflorum.

CALOSERIS, Benth. Pl. Hartw. 88 (1841) = **Onoseris**, DC. (Compos.).
rupestris, Benth. l. c. = O. isotypus.

CALOSMON, J. S. Presl, Rostl. ii. 71 (1823) = **Lindera**, Thunb. (Laurin.).
aestivale, Kostel. Allg. Med.-Pharm. Fl. ii. 478 = L. Benzoin.
Benzoin, J. S. Presl, Rostl. ii. 71 = L. Benzoin.
Neesianum, Kostel. Allg. Med.-Pharm. Fl. ii. 478 = L. Neesiana.

CALOSPHACE, Rafin. Fl. Tellur. iii. 91 = **Salvia**, Tourn. (Labiat.).

CALOSTELMA, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. sub t. 184 (1833) = **Liatris**, Schreb. (Compos.).
elegans, D. Don, l. c. = L. elegans.

CALOSTEMMA, R. Br. Prod. 297 (1810). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 735.
album, *R. Br. l. c.* 298.—Austral.
aureum, Link, Enum. Hort. Berol. i. 312 = luteum.
candidum, Lindl. in Mitch. Three Exped. i. 54 = luteum.
carneum, Lindl. l. c. ii. 39 = purpureum.
Cunninghami, *Ait. ex Loud. Hort. Brit. Suppl.* ii. 616.—Austral.
luteum, *Sims, Bot. Mag.* t. 2101.—Austral.
purpureum, *R. Br. Prod.* 298.—Austral.

CALOSTEPHANE, Benth. in Hook. Ic. Pl. t. 1111 (1872). *COMPOSITAE*, Benth. & Hook. f. ii. 335.
divaricata, *Benth. l. c.*—Afr. trop.

CALOSTIGMA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 343. t. 12. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 752.
Burchellii, *Fourn. in Mart. Fl. Bras.* vi. IV. 286.—Bras.
glabrum, *Decne. in DC. Prod.* viii. 581.—Bras.
Glaziovii, *Fourn. in Mart. Fl. Bras.* vi. IV. 286.—Bras.
Guilleminianum, *Decne. in DC. Prod.* viii. 581.—Bras.
insigne, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 343.—Bras.

CALOSTIGMA, Schott, Melet. Bot. 19 (1832) = **Philodendron**, Schott (Aroid.).
Imbe, Schott, l. c. = P. Imbe.

CALOSTIMA, Rafin. Fl. Tellur. iii. 47 (1836) = **Urera**, Gaudich. (Urtic.).
aculeata, Rafin. l. c. 48 = U. baccifera.

CALOSTROPHUS, F. Muell. Fragm. viii. 86 (1873) = **Hypolaena**, R. Br. (Restiac.).
elongatus, F. Muell. l. c. = H. longissima.
elongatus, Mast. in DC. Monog. Phan. i. 378 = H. lateriflora.

CALOSTROPHUS :—
exsulcus, F. Muell. Census, 124 = H. exsulca.
fastigiatus, F. Muell. in Papers Roy. Soc. Tasm. (1878) 117 = H. fastigiata.
gracillimus, F. Muell. Fragm. viii. 88 = H. gracillima.
lateriflorus, F. Muell. l. c. 87 = H. lateriflora.
Sieberianus, Mast. in DC. Monog. Phan. i. 378 = H. lateriflora.

CALOTHAMNUS, Labill. Nov. Holl. Pl. ii. 25. t. 164 (1806). *MYRTACEAE*, Benth. & Hook. f. i. 706.
BAUDINIA, Leschen. ex DC. Prod. iii. 211 (1828).
BILLOTTIA, Colla, Hort. Ripul. 20. t. 23 (1824).
affinis, Turcz. in Bull. Phys.-Math. Acad. Pétersb. x. (1852) 346 = microcarpus ?
asper, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 25.—Austral.
Baxteri, Hort. Angl. ex Klotzsch, in Otto & Dietr. Allg. Gartenz. iv. (1836) 115 = Kunzea Baxteri.
blepharanthcrus, *F. Muell. Fragm.* iii. 112.—Austral.
blepharospermus, *F. Muell. l. c.* 111.—Austral.
chrysanthereus, *F. Muell. l. c.* 112.—Austral.
clavatus, Mackay, ex Lodd. Bot. Cab. t. 1447 = quadrifidus.
eriocarpus, Lindl. Swan River App. 9 = sanguineus.
gibbosus, *Benth. Fl. Austral.* iii. 175.—Austral.
Gilesii, *F. Muell. Fragm.* x. 31.—Austral.
gracilis, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.
homalophyllus, *F. Muell. Fragm.* iii. 111.—Austral.
Huegelii, Schau. in Nov. Act. Nat. Cur. xxi. (1845) 30 = lateralis.
Knightii, Hort. ex Schau. l. c. 23 = robusta.
laevigatus, Schau. l. c. 26 = quadrifidus.
lateralis, *Lindl. Swan River App.* 9.—Austral.
laxus, Kunze, in Linnaea, xx. (1847) 58 = Preissii.
Lehmanni, *Schau. in Nov. Act. Nat. Cur.* xxi. (1845) 27.—Austral.
longifolius, Lehm. Del. Sem. Hort. Hamb. (1842) 7 = lateralis.
longissimus, *F. Muell. Fragm.* iii. 112.—Austral.
microcarpus, *F. Muell. l. c.* 113.—Austral.
nodosus, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 168 = lateralis.
Oldfieldii, *F. Muell. Fragm.* iii. 113.—Austral.
pachystachyus, *Benth. Fl. Austral.* iii. 173.—Austral.
pinifolius, *F. Muell. Fragm.* iii. 153.—Austral.
planifolius, *Lehm. Del. Sem. Hort. Hamb.* (1842) 7.—Austral.
plumosus, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 168 = Lehmanni.
Preissii, *Schau. in Nov. Act. Nat. Cur.* xxi. (1845) 27.—Austral.
purpureus, Endl. in Enum. Pl. Hueg. 48 = quadrifidus.
quadrifidus, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.
robustus, Schau. in Nov. Act. Nat. Cur. xxi. (1845) 22 = villosus.
rupestris, *Schau. l. c.*—Austral.
sanguineus, *Labill. Nov. Holl. Pl.* ii. 25. t. 164.—Austral.
Schaueri, *Lehm. Del. Sem. Hort. Hamb.* (1842) 7.—Austral.
schoenophyllus, Schau. in Nov. Act. Nat. Cur. xxi. (1845) 29 = Schaueri.
spathulatus, Hort. ex Steud. Nom. ed. II. i. 261 = Kunzea Baxteri.
suberosus, Schau. in Lehm. Pl. Preiss. i. 156 = Melaleuca exarata.
torulosus, *Schau. in Nov. Act. Nat. Cur.* xxi. (1845) 21.—Austral.
villosus, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 418.—Austral.

CALOTHECA, Spreng. Anleit. ii. 1. 167 (1817); Syst. i. 347 (1825) = **Aeluropus**, Trin. (Gramin.).
arabica, Spreng. Syst. i. 348 = A. arabicus.
cuspidata, Spreng. l. c. = Poa rufescens.
litoralis, Spreng. l. c. 347 = A. littoralis.
massariensis, Hochst. & Steud. ex Steud. Nom. ed. II. i. 261 = A. littoralis.
mucronata, Spreng. Syst. i. 348 = Poa mucronulata.
niliaca, Spreng. l. c. = A. niliacus.
poaeformis, Spreng. l. c. = Briza Lamarckiana.
pungens, Spreng. l. c. = Sesleria echinata.
purpurascens, Spreng. l. c. = Briza purpurascens.
repens, Spreng. l. c. 347 = A. villosus.

CALOTHECA, Desv. in Nov. Bull. Soc. Philom. ii. (1810) 190 = **Briza**, Linn. (Gramin.).
barbata, Nees, Agrost. Bras. 477 = B. barbata.
brizoidea, Beauv. Agrost. 86. t. 17. f. 6 = B. erecta.
brizoidea, Desv. Journ. Bot. i. (1813) 71 = B. Lamarckiana.
bromoidea, Lejeune, Messager des Sciences (1823); et in Nov. Act. Nat. Cur. xii. (1825) 757 = Bromus arduennensis.
dilatata, Link, Hort. Berol. i. 157 = B. erecta.
elegans, Beauv. Agrost. 157. t. 17 = B. elegans.
macrostachya, Presl, Rel. Haenk. i. 268 = B. macrostachya.
microstachya, Presl, l. c. = B. microstachya.
montevidensis, Spreng. ex Steud. Nom. ed. II. i. 261, 225 = B. subaristata.
reniformis, Presl, Rel. Haenk. i. 268 = B. reniformis.
rotundata, Roem. & Schult. Syst. ii. 632 = B. rotundata.
sabulosa, Steud. in Flora, xii. (1829) II. 488 = Sporobolus virginicus.
stricta, Hook. & Arn. Bot. Beech. Voy. 50 = B. stricta.
triloba, Kunth, Enum. Pl. i. 374 = B. triloba.

CALOTHYRSUS, Spach, in Ann. Sc. Nat. Sér. II. ii. (1834) 62 = **Aesculus**, Linn. (Sapindac.).
californica, Spach, l. c. = A. californica.

CALOTIS, R. Br. in Bot. Reg. t. 504 (1820). *COMPOSITAE*, Benth. & Hook. f. ii. 267.
CHEIROLOMA, F. Muell. in Linnaea, xxv. (1852) 401.
GONIOPOGON, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 173.
HUENEFELDIA, Walp. in Linnaea, xiv. (1840) 307.
anthemoides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 44.—Austral.
breviseta, *Benth. in Enum. Pl. Hueg.* 60.—Austral.
cuneifolia, *R. Br. in Bot. Reg.* t. 504.—Austral.
cymbacantha, *F. Muell. in Linnaea*, xxv. (1852) 400.—Austral.
dentex, *R. Br. in Bot. Reg.* t. 504.—Austral.
dilatata, A. Cunn. in DC. Prod. v. 302 = cuneifolia.
erinacea, *Steetz, in Lehm. Pl. Preiss.* i. 424.—Austral.
glandulosa, *F. Muell. in Trans. Vict. Inst.* (1855) 129.—Austral.
hispidula, *F. Muell. l. c.* 130.—Austral.
Kempei, *F. Muell. in Trans. Roy. Soc. S. Austral.* iv. (1881) 112.—Austral.
lappulacea, *Benth. in Enum. Pl. Hueg.* 60.—Austral.
microcephala, *Benth. Fl. Austral.* iii. 504.—Austral.
microphylla, *Benth. in Enum. Pl. Hueg.* 60.—Austral.
Muelleri, Sond. in Linnaea, xxv. (1852) 470 = scabiosifolia.
palmata, *A. Gray, in Proc. Am. Acad.* v. (1861) 121.—Austral.
plumulifera, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 57.—Austral.
polyseta, Sond. in Linnaea, xxv. (1852) 470 = lappulacea.
porphyroglossa, *F. Muell. ex Benth. Fl. Austral.* iii. 505.—Austral.
pterosperma, *R. Br. ex Benth. l. c.*—Austral.
scabiosifolia, *F. Muell. in Linnaea*, xxv. (1852) 471.—Austral.
scapigera, *Hook. in Mitch. Journ. Trop. Austral.* 75.—Austral.
tropica, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 58 = breviseta.

CALOTROPIS, R. Br. in Mem. Wern. Soc. i. (1809) 39. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 754.
Acia, *Buch.-Ham. in Trans. Linn. Soc.* xiv. (1825) 247.—Ind. or.
gigantea, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 78.—Ind. or.
Hamiltoni, Wight, Contrib. Ind. Bot. 53 = procera.
herbacea, Wight, l. c. 54 = C. Acia.
heterophylla, Wall. Cat. n. 8217 = procera.
procera, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. ii. 78.—Peruv.; Afr. trop.; Ind. or.
Sussuela, G. Don, Gen. Syst. iv. 147 = Hoya imperialis ?
Wallichii, Wight, Contrib. Ind. Bot. 53 = procera.

CALPANDRIA, Blume, Bijdr. 178 (1825) = **Camellia**, Linn. (Ternstroem.).
lanceolata, Blume, l. c. = Cam. lanceolata.
Quiscosaura, Korth. in Verh. Nat. Gesch. Bot. 149 = Cam. Quinosaura.

CALPICARPUM, G. Don, Gen. Syst. iv. 100 (1837) = **Kopsia**, Blume (Apocyn.).
albiflorum, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 402.—Malaya.
Lamarkii, G. Don, Gen. Syst iv. 100 = K. fruticosa.
Roxburghii, G. Don, l. c. = K. fruticosa.

CALPIDIA, Thou. Hist. Vég. Isles Afr. Austr. (1805) 37. t. 10 = **Pisonia**, Linn. (Nyctagin.).
costata, Boj. Hort. Maurit. 265 = P. costata.
lanceolata, Thou. Hist. Vég. Isles Afr. Austr. 37. t. 10 = P. Calpidia.
macrophylla, Boj. Hort. Maurit. 265 = P. macrophylla.
ovalifolia, Boj. l. c. = P. Calpidia.

CALPIGYNE, Blume, Mus. Bot. Lugd. Bat. ii. 193 (1856). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 323.
frutescens, *Blume, l. c.*—Borneo.

CALPURNIA, E. Mey. Comm. Pl. Afr. Austr. 2 (1835). *LEGUMINOSAE*, Benth. & Hook. f. i. 554.
aurea, *Baker, in Oliver, Fl. Trop. Afr.* ii. 252.—Afr. trop.
aurea, Benth. in Ann. Wien. Mus. ii. (1840) 90 = lasiogyne.
capensis, Steud. Nom. ed. II. i. 262 = Virgilia capensis.
floribunda, *Harv. in Harv. & Sond. Fl. Cap.* ii. 267.—Afr. austr.
intrusa, *E. Mey. Comm. Pl. Afr. Austr.* 2.—Afr. austr.
lasiogyne, *E. Mey. l. c.* 3.—Afr. austr.
robinioides, *E. Mey. l. c.*—Afr. austr.
sericea, *Harv. in Harv. & Sond. Fl. Cap.* ii. 267.—Afr. austr.
sylvatica, *E. Mey. Comm. Pl. Afr. Austr.* 2.—Afr. austr.
villosa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 268.—Afr. austr.

CALSIAMA, Rafin. Sylva Tellur. 12 (1838) = **Odina**, Roxb. (Anacard.).
malabarica, Rafin. l. c. = O. Wodier.

CALTHA, Tourn. ex Adans. Fam. ii. 126 (1763) = **Calendula**, Linn. (Compos.).
arvensis, Moench, Meth. 585 = Calend. arvensis.
dentata, Moench, Meth. Suppl. 238 = Calend. stellata.
graveolens, Gilib. Fl. Lituan. i. 212 = Calend. arvensis.
officinalis, Moench, Meth. 585 = Calend. officinalis.
officinalis, Scop. Fl. Carn. ed. II. ii. 147 = Calend. arvensis.
stellata, Moench, Meth. Suppl. 238 = Calend. stellata.

CALTHA, [Rupp.] Linn. Syst. ed. I (1735). *RANUNCULACEAE*, Benth. & Hook. f. i. 6.
PSYCHROPHILA, Bercht. & Presl, Rostl. i. Ranunc. 80 (1823).
PSYCROPHILA, Rafin. Atl. Journ. 144 (1832).
THACLA, Spach, Hist. Vég. Phan. vii. 295 (1839).
alba, Jacquem. Voy. Bot. 6 = palustris.
alpestris, Schott, Nym. & Kotschy, Analect. Bot. 33 = palustris.
alpina, Schur, Enum. Pl. Transs. 25 = palustris.
andicola, *C. Gay, Fl. Chil.* i. 49. t. 2.—Chili.
appendiculata, Lechl. ex C. Muell. in Walp. Ann. vii. 26 = limbata.
appendiculata, *Pers. Syn.* ii. 107.—Reg. Magell.
arctica, R. Br. in Parry, 1st Voy. App. 265 = palustris.
asarifolia, DC. Syst. i. 309 = palustris.
baicalensis, Demid. ex Steud. Nom. ed. II. i. 262 = natans.
biflora, DC. Syst. i. 310 = palustris.
Bisma, Buch.-Ham. in Brewst. Edinb. Journ. Sc. i. (1824) 251 = Aconitum palmatum.
camtschatica, Spreng. Syst. ii. 660 = Oxygraphis glacialis.
Codua, Buch.-Ham. in Brewst. Edinb. Journ. Sc. i. (1824) 251 = Aconitum ferox.

CALTHA:—
cornuta, Schott, Nym. & Kotschy, Analect. Bot. 31 = palustris.
coronata, *Schur, Enum. Pl. Transs.* 25.—Transsylvan.
dentata, Muhl. Cat. ed. II. 56 = palustris.
De-Ranco, *Steud. in Flora*, xxxix. (1856) 407.—Chili.
dioneaefolia, *Hook. f. in Hook. Lond. Journ. Bot.* ii. (1843) 306.—Reg. Magell.
emodorum, Spreng. Syst. iv. Cur. Post. 220 = palustris.
ficarioides, Pursh, Fl. Am. Sept. ii. 289 = palustris.
flabellifolia, Pursh, l. c. 390. t. 17 = palustris.
glabra, Gilib. Fl. Lituan. ii. 279 = palustris.
glacialis, Spreng. Syst. ii. 660 = Oxygraphis glacialis.
Govaniana, Wall. Cat. n. 4710 = palustris.
grosse-serrata, Pantoč. in Verh. Ver. Naturw. Heilk. Presb. ii. (1871–72) 86 = palustris.
Guerangerii, Bor. in Billot, Annot. 11 = palustris.
himalensis, D. Don, Prod. Fl. Nep. 195 = palnstris.
holophylla, *Leyb. in Anal. Univ. Chil.* (1859) 678.—Chili.
integerrima, Pursh, Fl. Am. Sept. ii. 390 = palustris.
intermedia, Schott, Nym. & Kotschy, Analect. Bot. 32 = palustris.
introloba, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 98.—Austral.
kamtschatica, Spreng. Syst. ii. 666 = *camtschatica*.
laeta, Schott, Nym. & Kotschy, Analect. Bot. 32 = palustris.
lasopetala, Steud. Nom. ed. II. i. 139, sphalm. = leptosepala.
latifolia, Schott, Nym. & Kotschy, Analect. Bot. 32 = palustris.
leptosepala, *DC. Syst.* i. 310.—Am. bor. occ.
leptostachya, G. Don, Gen. Syst. i. 44 = praec.
limbata, *Schlecht. in Linnaea*, xxviii. (1854) 556.—Chili.
major, Mill. Gard. Dict. ed. VIII. n. 1 = palustris.
minor, Mill. l. c. n. 2 = palustris.
multicapsularis, Sond. ex DC. Prod. i. 44 = sagittata.
natans, *Pall. Reise*, iii. 248.—Reg. bor.
Nirbisia, Buch.-Ham. in Brewst. Edinb. Journ. Sc. i. (1824) 251 = Aconitum ferox.
Novae-Zelandiae, *Hook. f. Fl. Nov. Zeal.* i. 12. t. 6.—N. Zel.
orthorhyncha, *Rupr. Fl. Caucas.* 28.—Reg. Caucas.
pallidiflora, Martr. in Bull. Soc. Bot. Fr. (1862) 130 = palustris.
palustris, *Linn. Sp. Pl.* 558.—Reg. bor. temp. et arct.
paniculata, Wall. Cat. n. 4711 = palustris.
paradoxa, T. F. Forst. in Trans. Linn. Soc. viii. (1807) 324 = appendiculata.
parnassifolia, Rafin. in Med. Rep. New York v. (1808) 361 = palustris.
polypetala, *Hochst. in Lorent, Wanderungen*, 339.—As. Min.; Persia.
Populago, Goüan, Hort. Monsp. 268 = palustris.
pumila, Schur, Enum. Pl. Transs. 26 = palustris.
radicans, *T. F. Forst. in Trans. Linn. Soc.* viii. (1807) 324. t. 17.—Scotia.
ranunculoides, Schur, Enum. Pl. Transs. 26 = palustris.
riparia, G. Don, Gen. Syst. i. 44 = palustris.
sagittata, *Cav. Ic.* v. 8, 456. t. 414.—Reg. Magell.
scaposa, *Hook. f. & Thoms. Fl. Ind.* i. 40.—Reg. Himal.
vulgaris, Pall. ex Steud. Nom. ed. II. i. 262 = palustris.
vulgaris, Schott, Nym. & Kotschy, Analect. Bot. 33 = palustris.

CALTHOIDES, B. Juss. ex DC. Prod. vi. 473 (1837) = **Othonna**, Linn. (Compos.).

CALUCECHINUS, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 19 (1853) = **Fagus**, Tourn. (Cupulif.).
antarctica, Hombr. & Jacquinot, l. c. = F. antarctica.
Montagnei, Hombr. & Jacquinot, l. c. = F. antarctica.

CALUSIA, Bert. ex Steud. Nom. ed. II. i. 262 (1840) = **Myrospermum**, Jacq. (Legumin.).
emarginata, Bert. ex Steud. l. c. ii. 175 = M. frutescens.

CALUSPARASSUS, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 20 (1853) = **Fagus**, Tourn. (Cupulif.).

CALUSPARASSUS:—
betuloides, Hombr. & Jacquinot, ex Decne. Bot. Voy. Astrol. et Zél. 20 = F. betuloides.
Forsteri, Hombr. & Jacquinot, l. c. = F. betuloides.
Pumilio, Hombr. & Jacquinot, l. c. 21 = F. betuloides.

CALVARIA, Comm. ex Gaertn. f. Fruct. Suppl. 116. t. 200, 201 (1805) (Gen. dub.).
cerebellina, Comm. ex Gaertn. f. l. c.—Hab.?
globosa, Gaertn. f. l. c. 117.—Hab.?
hexangularis, Gaertn. f. l. c.—Hab.?
major, Gaertn. f. l. c. 116 = *cerebellina*?

CALVELIA, Moq. in DC. Prod. xiii. II. 167 (1849) = **Suaeda**, Forsk. (Chenopod.).
pterantha, Moq. l. c. 168.—Soongar.

CALVOA, Hook. f. in Benth. & Hook. f. Gen. i. 755 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
crassinoda, *Hook. f. ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 78.—Afr. trop.
hirsuta, *Hook. f. ex Triana, l. c.*—Afr. trop.
sinuata, *Hook. f. ex Triana, l. c.*—Afr. trop.

CALYCADENIA, DC. Prod. v. 695 (1836) = **Hemizonia**, DC. (Compos.).
Cephalotes, DC. l. c. = H. multiglandulosa.
Fremontii, A. Gray, in Torr. Bot. Mex. Bound. 101, in nota = H. multiglandulosa.
Harrisii, Lem. Illustr. Hortic. (1855) Misc. 9 = Odontadenia speciosa.
mollis, A. Gray, in Proc. Am. Acad. vii. (1868) 360 = H. mollis.
multiglandulosa, DC. Prod. v. 695 = H. multiglandulosa.
pauciflora, A. Gray, in Torr. Bot. Mex. Bound. 101, in nota = H. pauciflora.
plumosa, Kellogg, in Proc. Calif. Acad. v. (1873) 49 = H. plumosa.
tenella, Torr. & Gray, Fl. N. Am. ii. 402 = H. tenella.
truncata, DC. Prod. v. 695 = H. truncata.
villosa, DC. l. c. = H. Douglasii.

CALYCAMPE, Berg, in Linnaea, xxvii. (1854) 129 = **Myrcia**, DC. (Myrtac.).
angustifolia, Berg, l. c. 131 = M. hebepetala.
latifolia, Berg, l. c. 130.—Guiana.

CALYCANDRA, Lepr. ex A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 30 et 232. t. 9 (1832) = **Cordyla**, Lour. (Legumin.).
pinnata, A. Rich. l. c. 31. t. 9 = Cord. africana.

CALYCANTHEMUM, Klotzsch, in Peters, Reise Mossamb. Bot. 243, t. 40 (1861) = **Ipomoea**, Linn. (Convolvul.).
leucanthemum, Klotzsch, l. c. 244.—Afr. trop.

CALYCANTHUS, Linn. Syst. ed. X. 1066 (1759). *CALYCANTHACEAE*, Benth. & Hook. f. i. 16.
BASTERIA, Mill. ex Adans. Fam. ii. 294 (1763).
BEURERIA, Ehret, Pl. et Pap. Rar. t. 13 (1755).
BUTNERIA, Duham. Arb. i. 113. t. 45 (1755).
POMPADOURA, Buchoz, ex DC. Prod. iii. 2 (1828).
acuminatus, Hort. ex C. Koch, Hort. Dendrol. 183 = floridus.
asplenifolius, Hort. ex C. Koch, Dendrol. i. 365 = floridus.
bullatus, Hort. ex C. Koch, l. c. = floridus.
ferax, Michx. Fl. Bor. Am. i. 305 = laevigatus.
fertilis, Andr. Rep. viii. t. 539 = laevigatus.
fertilis, Walt. Fl. Carol. 151 = glaucus.
floridus, *Linn. Syst.* ed. X. 1066.—Am. bor.
glaucus, *Willd. Enum. Hort. Berol.* 559.—Am. bor.
inodorus, Ell. Sketch, i. 576 = laevigatus.
laevigatus, *Willd. Hort. Berol.* t. 80; *Enum. Hort. Berol.* 559.—Am. bor.
macrophyllus, Hort. ex C. Koch, Dendrol. i. 364 = occidentalis.
mollis, *Rafin. Alsog. Am.* 7.—Am. bor.
nanus, Loisel. in Duham. ed. Nov. i. 219 = laevigatus.
oblongifolius, *Loud. Hort. Brit.* 214.—Am. bor.
occidentalis, *Hook. & Arn. Bot. Beech. Voy.* 340. t. 84.—Calif.
pensylvanicus, *Lodd. in Loud. Hort. Brit.* 214.—Am. bor.

CALYCANTHUS:—
philadelphicus, *Nois. ex Steud. Nom.* ed. II. i. 262.—Am. bor.
praecox, Linn. Sp. Pl. ed. II. 718 = Chimonanthus fragrans.
prunifolius, *Sweet, Hort. Brit.* ed. II. 195, nomen.—Carolina.
reticulatus, *Rafin. Alsog. Am.* 7.—Am. bor.
sterilis, Walt. Fl. Carol. 151 = floridus.
suaveolens, Salisb. Prod. 379 = Chimonanthus fragrans.
tomentosus, *Rafin. Alsog. Am.* 7.—Am. bor.
verrucosus, *Rafin. l. c.* 8.—Am. bor.
versicolor, *Regel, Cat. Pl. Hort. Aksakov.* 25.—Am. bor.

CALYCERA, Cav. Ic. iv. 34. t. 358 (1797) (Calicera). *CALYCEREAE*, Benth. & Hook. f. ii. 162.
ANOMOCARPUS, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 351; Contrib. Bot. ii. 28.
DISCOPHYTUM, Miers, ex Lindl. Veg. Kingd. 701 (1847).
GYMNOCAULUS, Phil. in Linnaea, xxviii. (1856) 705.
LEUCOCERA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 582.
balsamitaefolia, *Rich. in Mem. Mus. Par.* vi. (1820) 77.—Chili.
Calcitrapa, *Griseb. in Goett. Abh.* xix. (1874) 163.—Reg. Argent.
Cavanillesii, Rich. in Mem. Mus. Par. vi. (1820) 77 = herbacea.
eryngioides, *Remy, in C. Gay, Fl. Chil.* iii. 254.—Chili.
herbacea, Cav. Ic. iv. 34. t. 358.—Chili.
intermedia, *Phil. in Anal. Univ. Chil.* (1870) ii. 174.—Chili.
involucrata, *Phil. Sert. Mend. Alt.* 16.—Chili.
nudicaulis, Phil. ex Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 399 = viridiflora.
pulvinata, *Remy, in Ann. Sc. Nat.* Sér. III. vi. (1846) 352.—Bolivia.
sessiliflora, *Phil. in Linnaea*, xxviii. (1856) 706.—Chili.
sinuata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 398.—Chili.
spinulosa, *Gill. ex Miers, l. c.* 399.—Chili.
squarrosa, *Miers, l. c.* 398.—Chili.
tenuifolia, *Phil. ex Miers, l. c.* 356.—Chili.
ventosa, *Meyen, Reise*, i. 356.—Chili.
viridiflora, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 399.—Chili.

CALYCIUM, Ell. Sketch, ii. 339 (1824) = **Heterotheca**, Cass. (Compos.).
divaricatum, Ell. l. c. 339, 338 = H. Lamarckii.
scabrum, Ell. l. c. 339 = H. Lamarckii.

CALYCOBOLUS, Willd. ex Roem. & Schult. Syst. v. 2 (1819) = **Breweria**, R. Br. (Convolvul.).
emarginatus, Willd. l. c. 4.—N. Granat.
pulchellus, Willd. l. c.—N. Granat.

CALYCOCARPUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 48 (1838). *MENISPERMACEAE*, Benth. & Hook. f. i. 35.
Lyonii, *Nutt. ex A. Gray, Gen. Am. Bor.* i. 76.—Am. bor.

CALYCOCORSUS, F. W. Schmidt, Samml. Phys. Aufs. i. 271 (1795) = **Chondrilla**, Linn. (Compos.).
hieracioides, F. W. Schmidt, l. c. = Ch. Peltidium.

CALYCODAPHNE, Boj. Hort. Maurit. 273 (1837) = **Ocotea**, Aubl. (Laurin.).
cupularis, Boj. l. c. = O. floribunda.
dysoxylon, Boj. l. c. = O. floribunda.
floribunda, Boj. l. c. = O. floribunda.

CALYCODON, Nutt. in Journ. Acad. Phil. Ser. II. i. 186 (1847) = **Muehlenbergia**, Schreb. (Gramin.).
montanum, Nutt. l. c. = M. gracilis.

CALYCOGONIUM, DC. Prod. iii. 168 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 761.
CALYCOPTERIS, Rich. ex DC. Prod. iii. 168 (1828).
angulatum, *Griseb. Cat. Pl. Cub.* 93.—Cuba.
clidemioides, *Griseb. l. c.* 94.—Cuba.
cubense, Griseb. l. c. 93 = stellatum.

CALYCOGONIUM :—
cubense, A. Rich. Ess. Fl. Cub. 555 = Pachyanthus Poiretii.
echinatum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 184.—Cuba.
ellipticum, *Sauvalle, Fl. Cub.* 50.—Cuba.
glabratum, *DC. Prod.* iii. 168.—Jamaic.
Griscbachii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 97.—Cuba.
hetcrophyllum, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 85.—Cuba.
hetcrotrichum, *Triana, in Trans. Linn Soc.* xxviii. (1871) 96.—Cuba.
hypoglaucum, *Wright, ex Griseb. Cat. Pl. Cub.* 94.—Cuba.
involucratum, Griscb. in Mem. Am. Acad. N. S. viii. (1861) 184 = Ossaca involucrata.
lanceolatum, Griseb. Cat. Pl. Cub. 93 = glabratum.
Lima, Griscb. l. c. 95 = Oxymeris asperifolia.
Lindenianum, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 84.—Cuba.
microphyllum, Sauvalle, Fl. Cub. 50 = glabratum.
muricatum, Griseb. Cat. Pl. Cub. 95 = Octopleura muricata.
pauciflorum, Triana, in Trans. Linn. Soc. xxviii. (1871) 96 = echinatum.
plicatum, *Griseb. Cat. Pl. Cub.* 94.—Cuba.
productum, Sauvalle, Fl. Cub. 49 = heterophyllum.
rhamnoideum, Griseb. Cat. Pl. Cub. 93 = glabratum.
rhamnoideum, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 85.—Jamaic.
rosmarinifolium, *Griseb. Cat. Pl. Cub.* 93.—Cuba.
stellatum, *DC. Prod.* iii. 168.—Ins. S. Doming.
stellatum, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 183 = Grisebachii.
verrucosum, *Griseb. Cat. Pl. Cub.* 95.—Cuba.
Wrightianum, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 184 = glabratum.

CALYCOLPUS, Berg, in Linnaea, xxvii. (1854) 378. *MYRTACEAE*, Benth. & Hook. f. i. 713.
calophyllus, *Berg, in Mart. Fl. Bras.* xiv. I. 412.—Bras.
glaber, *Berg, l. c.* 411.—Peruv.
Goetheanus, *Berg, l. c.* 412.—Bras.
Kegclianus, *Berg, in Linnaea*, xxvii. (1854) 381.—Guiana.
ovalifolius, *Berg, l. c.* 379.—Ind. occ.
parviflorus, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 181.—Guiana.
revolutus, *Berg, in Linnaea*, xxvii. (1854) 383.—Guiana.
Schomburgkianus, *Berg, l. c.* 380.—Guiana.
Warscewiczianus, *Berg, l. c.* 382.—N. Granat.

CALYCOMELIA, Kostcl. Allg. Med.-Pharm. Fl. iii. 1003 (1834) = **Fraxinus**, Linn. (Oleac.).
acuminata, Kostel. l. c. 1004 = F. americana.
alba, Kostcl. Ind. Hort. Prag. 26 = F. alba.
americana, Kostcl. Allg. Med.-Pharm. Fl. iii. 1004 = F. americana.
caroliniana, Kostcl. Ind. Hort. Prag. 26 = F. platycarpa.
elliptica, Kostel. l. c. = F. elliptica.
epiptera, Kostel. l. c. = F. epiptera.
expansa, Kostel. l. c. = F. pubescens.
juglandifolia, Kostel. Allg. Med.-Pharm. Fl. iii. 1004 = F. americana.
lancea, Kostel. Ind. Hort. Prag. 26 = F. lancea.
nigra, Kostel. l. c. = F. nigra.
ovata, Kostel. l. c. = F. ovata.
pallida, Kostel. l. c. = F. pallida.
platycarpa, Kostel. l. c. = F. platycarpa.
pubescens, Kostel. l. c. = F. pubescens.
quadrangulata, Kostel. Allg. Med.-Pharm. Fl. iii. 1004 = F. quadrangulata.
Richardi, Kostel. Ind. Hort. Prag. 26 = F. Richardi.
tomentosa, Kostel. Allg. Med.-Pharm. Fl. iii. 1004 = F. pubescens.
viridis, Kostel. Ind. Hort. Prag. 26 = F. viridis.

CALYCOMIS, R. Br. in Flind. Voy. App. 549 (1814) = **Callicoma**, Andr. (Saxifrag.).

CALYCOMIS, D. Don, in Edinb. N. Phil. Journ. ix. (1830) 93 = **Acrophyllum**, Benth. (Saxifr.).
verticillatum, D. Don. l. c. = A. venosum.

CALYCOMORPHUM, Presl, Symb. Bot. i. 50 (1830) = **Trifolium**, Tourn. (Legumin.).
globosum, Presl, l. c. = T. globosum.
subterraneum, Presl, l. c. = T. subterrancum.

CALYCOPEPLUS, Planch. in Bull. Soc. Bot. Fr. viii. (1861) 30. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 261.
cphedroides, *Planch. l. c.* 31.—Austral.
marginatus, *Benth. Fl. Austral.* vi. 53.—Austral.
paucifolius, Baill. Adansonia, vi. (1865–66) 319 = ephedroides.

CALYCOPHYLLUM, DC. Prod. iv. 367 (excl. sp. 2 & 3) (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 38.
EUKYLISTA, Benth. in Hook. Kew Journ. v. (1853) 230.
candidissimum, *DC. Prod.* iv. 367.—Am. austr.
coccineum, DC. l. c. = Warscewiczia coccinea.
grandiflorum, Meyen, Reise, ii. 234 = Mussaenda grandiflora.
multiflorum, *Griseb. in Goett. Abh.* xxiv. (1879) 155.—Reg. Argent.
Stanleyanum, *Schomb. in Hook. Lond. Journ. Bot.* iii. (1844) 622.—Guiana.
tubulosum, DC. Prod. iv. 367 (Pogonopi sp.).—Bras.
tubulosum, Secm. Bot. Voy. Herald, 135 = Pogonopus exsertus.

CALYCOPHYSUM, Karst. & Triana, in Linnaea, xxviii. (1856) 427. *CUCURBITACEAE*, Benth. & Hook. f. i. 828.
gracile, *Cogn. in Bull. Acad. Belg.* Sér. II. xlix. (1880) 191.—N. Granat.
pedunculatum, *Karst. & Triana, in Linnaea*, xxviii. (1856) 428.—N. Granat.

CALYCOPLECTUS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 120 = **Alloplectus**, Mart. (Gesner.).
concolor, Oerst. l. c. = A. dichrous.
dichrous, Oerst. l. c. = A. dichrous.
Pinelianus, Oerst. l. c. = A. Pinelianus.

CALYCOPTERIS, Lam. Illustr. t. 357 (1794?). *COMBRETACEAE*, Benth. & Hook. f. i. 686.
GETONIA, Roxb. Pl. Corom. i. 61. t. 87 (1795).
floribunda, *Lam. Tabl. Encyc.* ii. 485.—Ind. or.; Malaya.
nutans, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 59 = floribunda.

CALYCOPTERIS, Rich. ex DC. Prod. iii. 168 (1828) = **Calycogonium**, DC. (Melastom.).

CALYCORECTES, Berg, in Linnaea, xxvii. (1854) 317. *MYRTACEAE*, Benth. & Hook. f. i. 720.
SCHIZOCALYX, Berg, l. c. 319 (1854).
cubensis, *Griseb. Cat. Pl. Cub.* 90.—Cuba.
grandifolius, *Berg, in Linnaea*, xxvii. (1854) 317.—Guiana.
Langsdorffi, *Berg, in Mart. Fl. Bras.* xiv. I. 597.—Bras.
latifolius, *Berg, in Linnaea*, xxx. (1859–60) 701.—Guiana.
Martianus, *Berg, in Mart. Fl. Bras.* xiv. I. 596.—Bras.
mexicanus, *Berg, in Linnaea*, xxvii. (1854) 318.—Mexic.
protractus, *Griseb. Cat. Pl. Cub.* 284.—Cuba.
Ricdelianus, *Berg, in Mart. Fl. Bras.* xiv. I. 596.—Bras.
Schottianus, *Berg, l. c.* 357.—Bras.
Sellowianus, *Berg, l. c.* 356.—Bras.

CALYCOSERIS, A. Gray, Pl. Wright. ii. 104. t. 14 (1853). *COMPOSITAE*, Benth. & Hook. f. ii. 523.
Parryi, *A. Gray, in Torr. Bot. Mex. Bound.* 106.—Californ.
Wrightii, *A. Gray, Pl. Wright.* ii. 104. t. 14.—N. Mexic.

CALYCOSORUS, Endl. Gen. 499, in syn. (1838) = CALYCOCORSUS, F. W. Schmidt = **Chondrilla**, Linn. (Compos.).

CALYCOSIA, A. Gray, in Proc. Am. Acad. iv. (1860) 47. *RUBIACEAE*, Benth. & Hook. f. ii. 122.
Hunteri, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1884) 364.—Ins. Fiji.

CALYCOSIA :—
Milnei, *A. Gray, in Proc. Am. Acad.* iv. (1860) 307.—Ins. Fiji.
petiolata, *A. Gray, l. c.* 48.—Ins. Fiji.
pubiflora, *A. Gray, l. c.* 306.—Ins. Fiji.
sessilis, *A. Gray, l. c.* 48.—Ins. Samoa.

CALYCOSTEGIA, Lem. in Orb. Dict. iii. 86 (1849) = **Calystegia**, R. Br. (Convolv.).

CALYCOSTEMMA, Hanst. in Linnaea, xxix. (1857–58) 506 = **Isoloma**, Decne. (Gesnerac.).
Lindenii, Hanst. l. c. 516.—N. Granat.

CALYCOTHRIX, Meissn. Gen. 107 (1837) = **Calythrix**, Labill. (Myrtac.).

CALYCOTOME, E. Mey. Comm. Pl. Afr. austr. 113 (1835) = **Dichilus**, DC. (Legumin.).
pusilla, E. Mey. l. c. 114 = D. gracilis.

CALYCOTOME, Link, in Schrad. Neues Journ. ii. (1808) II. 50 (Calicotome). *LEGUMINOSAE*, Benth. & Hook. f. i. 481.
cretica, Presl, ex Nym. Consp. 158 = villosa.
infesta, *Guss. Fl. Sic. Syn.* ii. 247.—Dalmatia.
intermedia, Boiss. ex Reichb. f. Ic. Fl. Germ. xxii. 33. t. 2094 = villosa.
spinosa, *Link, Enum. Hort. Berol.* ii. 225.—Lusitan.
villosa, *Link, in Schrad. Neues Journ.* ii. (1808) II. 51.—As. Min.

CALYCOTOMON, Hoffmgg. Verz. Pfl. 166 (1824) = **Calicotome**, Link (Legumin.).

CALYCOTOMUS, Rich. ex DC. Prod. iii. 173 (1828) = **Conostegia**, D. Don (Melastom.).

CALYCOTROPIS, Turcz. in Bull. Soc. Nat. Mosc. xxxv. (1862) II. 327. *PARONYCHIEAE* vel *ELATINEAE*? Benth. & Hook. f. Gen. i. 978.
minuartioides, *Turcz. l. c.* 328.—Mexic.

CALYDERMOS, Lag. Gen. et Sp. Nov. 24 (1816) = **Calea**, Linn. (Compos.).
atriplicifolius, Spreng. Syst. iii. 457 = Isocarpha atriplicifolia.
jamaicensis, Spreng. Syst. iii. 457 = Eupatorium Dalea.
longifolius, Lag. Gen. et Sp. Nov. 24 = Calea pedunculatis.
pedunculatis, DC. Prod. v. 669 = Calea pedunculatis.
repens, Spreng. Neue Entdeck. i. 278 = Diospermum involucratum.
rugosus, DC. Prod. v. 670 = Calea rugosa.
salmeaefolius, DC. l. c. = Calea salmiaefolia.
scaber, Lag. Gen. et Sp. Nov. 24 = Calea pedunculatis.
spilanthoides, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 514.—Cuba.

CALYDERMOS, Ruiz & Pav. Fl. Per. ii. 43 (1799) = **Nicandra**, Adans. (Solanac.).
erosus, Ruiz & Pav. l. c. = N. physalodes.

CALYDOREA, Herb. in Lindl. Bot. Reg. (1843) Misc. 85. *IRIDEAE*, Benth. & Hook. f. iii. 695.
BOTHERBE, Steud. ex Klatt, in Linnaea, xxxi. (1860–61) 562.
ROTERBE, Klatt, in Mart. Fl. Bras. iii. I. 543. t. 71 (1871).
azurea, *Klatt, in Abh. Naturf. Ges. Halle.* xii. (1882) 387.—Hab.?
campestris, *Baker, in Journ. Bot.* xiv. (1876) 188.—Am. trop.
cipuroides, *Klatt, in Abh. Naturf. Ges. Halle.* xii. (1882) 387.—Hab.?
furcata, *Baker, in Journ. Bot.* xiv. (1876) 189.—Reg. Argent.
Gardneri, *Baker, l. c.*—Bras.
longispatha, Baker, l. c. = Sphenostigma longispatha.
luteola, *Baker, l. c.*—Bras.
nuda, *Baker, l. c.*—Reg. Argent.
pallens, *Griseb. in Goett. Abh.* xxiv. (1879) 324.—Reg. Argent.
punctata, Baker, in Journ. Bot. xiv. (1876) 189 = Nemastylis punctata.
speciosa, *Herb. in Bot. Reg.* (1843) *Misc.* 85.—Chili.
texana, *Baker, in Journ. Bot.* xiv. (1876) 189.—Texas.

CALYGOGONIUM, G. Don, Gen. Syst. ii. 731 (1832) = **Calycogonium**, DC. (Melastom.).

CALYGONIUM, D. Dietr. Syn. ii. 1376 (1840) = praec.

CALYLOPHIS, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 272 = **Oenothera**, Linn. (Onagrar.).
Berlandieri, Spach, l. c. 273 = Oe. serrulata ?
Drummondiana, Spach, l. c. = Oe. serrulata.
Drumondii, Spach, ex Steud. Nom. ed. II. i. 262 = praec.
Nuttallii, Spach, Hist. Vég. Phan. iv. 350 = Oe. serrulata.

CALYLOPHUS, Spach, Hist. Vég. Phan. iv. 351 = praec.

CALYMENIA, Pers. Syn. i. 36 (1805) = **Oxybaphus**, Vahl (Nyctagin.).
aggregata, Pers. Syn. i. 37 = O. aggregatus.
albida, Nutt. Gen. Am. i. 26 = O. aggregatus.
angustifolia, Nutt. in Fras. Cat. (1813); Gen. Am. i. 26 = O. angustifolius.
Cervantesii, Heynh. Nom. ii. 110 = O. Cervantesii.
chilensis, Heynh. l. c. = O. chilensis.
corymbosa, Pers. Syn. i. 37 = O. glabrifolius.
decumbens, Nutt. Gen. Am. i. 26 = O. decumbens.
expansa, Pers. Syn. i. 36 = O. expansus.
hirsuta, Nutt. Gen. Am. i. 26 = O. hirsutus.
linearis, Nutt. l. c. = O. angustifolius.
nyctaginea, Nutt. l. c. = O. nyctagineus.
ovata, Pers. Syn. i. 36 = O. ovatus.
paniculata, Heynh. Nom. ii. 110 = O. glabrifolius.
pilosa, Nutt. Gen. Am. i. 26 = O. ovatus.
prostrata, Pers. Syn. i. 36 = O. prostratus.
viscosa, Pers. l. c. = O. viscosus.

CALYMMANDRA, Torr. & Gray, Fl. N. Am. ii. 262 (1841) = **Evax**, Gaertn. (Compos.).
candida, Torr. & Gray, l. c. = E. candida.

CALYMNANDRA, Lindl. Veg. Kingd. 710 (1847) = praec.

CALYNUX, Rafin. in Journ. Phys. lxxxix. 257 (1819) = **Pyrularia**, Rich. (Santalac.).

CALYPLECTUS, Ruiz & Pav. Prod. 73. t. 13 (1794) = **Lafoensia**, Vand. (Lythrar.).
acuminatus, Ruiz & Pav. Fl. Per. iv. t. 423 = L. acuminata.
adenophyllus, Schott, ex Steud. Nom. ed. II. i. 263 = L. replicata.
dependens, Ruiz, ex Koehne, in Engl. Bot. Jahrb. iii. (1882) 155 = L. acuminata.
punicaefolius, Bert. ex Koehne, l. c. 154 = L. punicaefolia.
speciosus, H. B. & K. Nov. Gen. et Sp. vi. 183. t. 548 = L. speciosa.

CALYPSO, Salisb. Parad. Lond. t. 89 (1806).
ORCHIDEAE, Benth. & Hook. f. iii. 496.
CYTHEREA, Salisb. in Trans. Hort. Soc. i. (1812) 301.
NORNA, Wahlenb. Fl. Suec. 561 (1826).
ORCHIDIUM, Sw. in Svensk Bot. t. 518 (1819).
americana, R. Br. in Ait. Hort. Kew. ed. II. v. 208 = borealis.
borealis, *Salisb. Parad. Lond.* t. 89.—Reg. bor. temp.
bulbosa, Reichb. f. Ic. Fl. Germ. xiii. 158. t. 137 = borealis ?

CALYPSO, Thou. Hist. Vég. Afr. Austr. i. 33. t. 6 (1805) = **Salacia**, Linn. (Celastr.).
africana, G. Don, Gen. Syst. i. 629 = S. africana.
campestris, Cambess. in St. Hil. Fl. Bras. Mer. ii. 111. t. 104 = S. campestris.
debilis, G. Don, Gen. Syst. i. 629 = S. debilis.
erecta, G. Don, l. c. = S. erecta.
madagascariensis, Cambess. in St. Hil. Fl. Bras. Mer. ii. 111. t. 104 = S. madagascariensis.
pyriformis, G. Don, Gen. Syst. i. 629 = S. pyriformis.
salacioides, Thou. ex Cambess. in St. Hil. Fl. Bras. Mer. ii. 111. t. 104 = S. Calypso.
senegalensis, Cambess. l. c. = S. senegalensis.
sylvestris, Mart. in Flora, xx. (1837) II. Beibl. 96 = S. sylvestris.

CALYPSODIUM, Link, Handb. i. 252 (1829) = **Calypso**, Salisb. (Orchid.).
boreale, Link, l. c. = Calypso borealis.

CALYPSOGYNE, Neraud, in Freyc. Voy. Bot. 25, 26 (1826), nomen.

CALYPTERIOPETALON, Hassk. in Flora, xl. (1857) 531 = **Croton**, Linn. (Euphorbiac.).
brasiliensis, Hassk. l. c. = Croton urticaefolius.

CALYPTERIS, Zipp. ex Mackl. in Alg. Konst. en Letter-bode, i. (1829) 298 (Ord. Nat. ?).
miniata, *Zipp. ex Mackl. l. c.* nomen.—Hab. ?

CALYPTHRANTES, Raeusch. Nom. ed. III. 144 (1797) = **Calyptranthes**, Sw. (Myrtac.).

CALYPTOCARPUS, Less. Syn. Comp. 221 (1832) = **Blainvillea**, Cass. (Compos.).
bahiensis, Sch. Bip. in Bot. Zeit. xxiv. (1866) 165 = B. bahiensis.
Burchellii, Sch. Bip. l. c. = B. rhomboidea.
megapotamica, Sch. Bip. l. c. = B. biaristata.
vialis, Less. Syn. Comp. 221 = B. tampicana.
Wendlandii, Sch. Bip. in Bot. Zeit. xxiv. (1866) 165. —Costa Rica.

CALYPTRANTHES, Sw. Prod. Veg. Ind. Occ. 79 (1788). *MYRTACEAE*, Benth. & Hook. f. i. 717.
CHYTRALIA, Adans. Fam. ii. 80 (1763).
MITRANTHES, Berg, in Linnaea, xxvii. (1854) 316.
SUZYGIUM, P. Br. ex Adans. Fam. ii. 244 (1763).
affinis, *Berg, in Mart. Fl. Bras.* xiv. I. 43.—Bras.
anceps, *Berg, l. c.* 40.—Bras.
aromatica, Blume, Bijdr. 1092 (= *Caryophyllus antisepticus*).
aromatica, *St. Hil. Pl. Us. Bras.* fasc. iii. t. 14.—Bras.
axillaris, *Berg, in Mart. Fl. Bras.* xiv. I. 53.—Bras.
bimarginata, *Berg, l. c.* 543.—Bras.
bipennis, *Berg, in Linnaea*, xxxi. (1861-62) 248.—Peruv.
Blanchetiana, *Berg, in Mart. Fl. Bras.* xiv. I. 40.—Bras.
brasiliensis, *Spreng. Syst.* ii. 499.—Bras.
bullata, *DC. Prod.* iii. 258.—Honduras.
calyptrata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 181.—Cuba.
capitellata, Buch.-Ham. ex Wall. Cat. n. 3560 B = Eugenia Jambolana.
caryophyllata, Pers. Syn. 32 = Eugenia caryophyllaea.
caryophyllifolia, Willd. in Usteri, Ann. xvii. (1796) 22 = Eugenia Jambolana.
caudata, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 102.—Bras.; N. Granat.
cephalantha, *Berg, in Mart. Fl. Bras.* xiv. I. 46.—Bras.
Chytraculia, *Sw. Prod. Veg. Ind. Occ.* 79.—Jamaica.
clusiaefolia, *Berg, in Mart. Fl. Bras.* xiv. I. 39.—Bras.
concinna, *DC. Prod.* iii. 258.—Bras.
cordata, *Berg, in Mart. Fl. Bras.* xiv. I. 48.—Bras.
cordifolia, Moon, Cat. Pl. Ceyl. 39 = Eugenia Neesiana *vel* cordifolia.
corymbosa, Blume, Bijdr. 1091 (= *Syzygium corymbosum*).
costaricensis, *Berg, in Linnaea*, xxvii. (1854) 20.—Costa Rica.
costata, Buch.-Ham. in Wall. Cat. n. 3556 = Eugenia operculata.
cubensis, *Berg, in Linnaea*, xxix. (1857-58) 215.—Ind. occ.
Cumini, Pers. Syn. ii. 32 = Eugenia Jambolana.
cuminodora, Stokes, Bot. Mat. Med. iii. 67 = Eugenia caryophyllifolia.
cuneata, Buch.-Ham. ex Wall. Cat. n. 3557 = Eugenia operculata.
cuprea, *Berg, in Linnaea*, xxx. (1859-60) 653.—Cuba.
cuspidata, *Mart. ex DC. Prod.* iii. 258.—Bras.
Danca, Buch.-Ham. ex Wall. Cat. n. 3560 = Eugenia Heyneana.
decandra, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 181.—Cuba.
densa, *DC. Prod.* iii. 257.—Bras.
densiflora, *Poepp. ex Berg, in Linnaea*, xxvii. (1854) 30.—Peruv.

CALYPTRANTHES:—
dichotoma, *Casar. Decad. Nov. Stirp. Bras.* v. 47.—Bras.
enneantha, *Wright, in Sauv. Fl. Cub.* 40.—Cuba.
eriopoda, *DC. Prod.* iii. 257.—Bras.
eugenioides, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 370.—Bras.
eugenioides, *Seem. Fl. Vit.* 81.—Ins. Fiji.
fasciculata, *Berg, in Linnaea*, xxvii. (1854) 31.—Guiana.
fastigiata, *Berg, in Mart. Fl. Bras.* xiv. I. 50.—Bras.
fastigiata, Blume, Bijdr. 1090 (= *Caryophyllus fastigiatus*).
floribunda, Blume, l. c. 1091 = Eugenia floribunda.
Forsteri, *Berg, in Linnaea*, xxvii. (1854) 23.—Ins. S. Lucia.
fragrans, Ruiz, ex Berg, l. c. 20 = paniculata.
glomerata, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 371. —Bras.
gracilipes, *Wright, in Sauv. Fl. Cub.* 39.—Cuba.
grandiflora, *Berg, in Mart. Fl. Bras.* xiv. I. 542.—Bras.
grandifolia, *Berg, l. c.* 48.—Bras.
grandis, Buch.-Ham. ex Wall. Cat. n. 3554 = Eugenia operculata.
guineensis, *Willd. Sp. Pl.* ii. 974.—Afr. trop.
Jambolana, Moon, Cat. Pl. Ceyl. 39 = Eugenia sylvestris.
Jambolana, Willd. in Usteri, Ann. xvii. (1796) 23 = Eugenia Jambolana.
jambolifera, Stokes, Bot. Mat. Med. iii. 68 = Eugenia Jambolana.
Karwinskyana, *Berg, in Linnaea*, xxix. (1857-58) 214. —Mexic.
lanceolata, *Berg, in Mart. Fl. Bras.* xiv. I. 51.—Bras.
Langsdorffi, *Berg, l. c.* 539.—Bras.
lateriflora, DC. Prod. iii. 258 (= *Rubachia lateriflora*). —Bras.
latifolia, *Poir. Encyc. Suppl.* ii. 43.—Ins. Maurit.
laxiflora, *Blume, Bijdr.* 1090.—Java.
Lindeniana, *Berg, in Linnaea*, xxix. (1857-58) 213.—Mexic.
longifolia, *Berg, in Mart. Fl. Bras.* xiv. I. 46.—Bras.
loranthifolia, *DC. Prod.* iii. 258.—Bras.
lucida, *Mart. ex DC. l. c.*—Bras.
macrophylla, *Berg, in Mart. Fl. Bras.* xiv. I. 45.—Bras.
Makal, Raeusch. Nom. ed. III. 144 (= *Syzygium Makul*).
Makul, Blanco, Fl. Filip. ed. I. 419 = Eugenia operculata.
malabarica, Dennst. Schluess. Hort. Malab. 31 = Eugenia zeylanica.
mangiferifolia, Hance, ex Walp. Ann. ii. 629 = Eugenia operculata.
Martiusiana, *DC. Prod.* iii. 257.—Bras.
maschalantha, *Berg, in Mart. Fl. Bras.* xiv. I. 53.—Bras.
melanoclada, *Berg, l. c.* 50.—Bras.
micrantha, *Wright, ex Griseb. Cat. Pl. Cub.* 85.—Cuba.
multiflora, *Poepp. ex Berg, in Mart. Fl. Bras.* xiv. I. 42.—Bras.
musciflora, *Berg, l. c.* 54.—Bras.
mutabilis, *Berg, l. c.* 540.—Bras.
myrtoides, *Seem. Fl. Vit.* 81.—Ins. Fiji.
nigricans, *DC. Prod.* iii. 258.—Bras.
nitida, Raeusch. Nom. ed. III. 144 = *Syzygium lucidum*.
Nummularia, *Berg, in Linnaea*, xxx. (1859-60) 652. —Ins. S. Doming.
obscura, *DC. Prod.* iii. 257.—Bras.
obscura, Mart. in Flora, xx. (1837) Beibl. ii. 87 = tuberculata.
obtusa, *Benth. in Hook. Journ. Bot.* ii. (1840) 319.—Guiana.
obtusa, Miq. in Linnaea, xxii. (1849) 172 = Marlierea suffruticosa.
obtusifolia, Buch.-Ham. ex Wall. Cat. n. 3600 B = Eugenia inophylla.
obversa, *Berg, in Mart. Fl. Bras.* xiv. I. 51.—Bras.
oleina, Wight, Illustr. ii. 15 = Eugenia myrtifolia.
Ottonis, *Wright, in Sauv. Pl. Cub.* 40.—Cuba.
ovalifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 371. —Bras.
ovata, *Berg, in Mart. Fl. Bras.* xiv. I. 50.—Bras.
pallens, *Griseb. in Goett. Abh.* vii. (1857) 215.—Ind. occ.

CALYPTRANTHES:—
paniculata, Raeusch. Nom. ed. III. 144 = Eugenia paniculatum.
paniculata, *Ruiz & Pav. Fl. Per.* iv. t. 424; *Syst.* 131.—Peruv.
paniculata, Spreng. Neuc Entdeck. ii. 170 = brasiliensis.
pauciflora, *Berg, in Mart. Fl. Bras.* xiv. 1. 543.—Bras.
pendula, Berg, in Linnaea, xxvii. (1854) 21 = C. Chytraculia.
platyphylla, *Berg, in Mart. Fl. Bras.* xiv. 1. 539.—Bras.
Poeppigiana, *Berg, l. c.* 45.—Bras.
Pohliana, *Berg, l. c.* 42.—Bras.
pollicina, Willem. in Usteri, Ann. xviii. (1796) 39 = Eugenia obovata.
polyantha, *Berg, in Mart. Fl. Bras.* xiv. 1. 541.—Bras.
pteropoda, *Berg, l. c.* 47.—Bras.
pulchella, *DC. Prod.* iii. 257.—Bras.
punctata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 181.—Cuba.
pyrifolia, Blume, Bijdr. 1090 = Eugenia pyrifolia.
racemosa, Blume, l. c. 1089 (= *Syzygium racemosum*).
ramiflora, Blanco, Fl. Filip. ed. I. 420 = Eugenia bracteata.
Regeliana, *Berg, in Mart. Fl. Bras.* xiv. 1. 542.—Bras.
rigida, Macf. ex Griseb. Fl. Brit. W. Ind. 233 = C. Syzygium.
rigida, *Sw. Prod. Veg. Ind. Occ.* 80.—Jamaic.; Cuba.
rigida, Tussac, Fl. Antill. iii. t. 26 = Tussaceana.
rostrata, Blume, Bijdr. 1092 (= *Syzygium rostratum*).
rostrata, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 181.—Cuba.
rotundata, *Griseb. l. c.*—Cuba.
rufa, *Berg, in Mart. Fl. Bras.* xiv. 1. 541.—Bras.
Ruiziana, *Berg, in Linnaea*, xxvii. (1854) 22.—Peruv.
Schiediana, *Berg, l. c.* 28.—Mexic.
Schlechtendaliana, *Berg, l. c.* 29.—Mexic.
scoparia, *Berg, l. c.* xxxi. (1861–62) 247.—Bras.
Seemanni, *Seem. Fl. Vit.* 81.—Ins. Fiji.
sericea, *Griseb. Fl. Brit. W. Ind.* 233.—Ind. occ.
speciosa, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 187.—Guiana.
Spruceana, *Berg, in Mart. Fl. Bras.* xiv. 1. 45.—Bras.
strigipes, *Berg, l. c.* 540.—Bras.
Syzygium, *Sw. Fl. Ind. Occ.* ii. 919.—Ind. occ.
Tatna, Buch.-Ham. ex Wall. Cat. n. 3555 = Eugenia operculata.
tenuis, Buch.-Ham. ex Wall. l. c. sub n. 3570 = Eugenia tenuis.
tetraptera, *Berg, in Mart. Fl. Bras.* xiv. 1. 53.—Bras.
Thomasiana, *Berg, in Linnaea*, xxvii. (1854) 26.—Ind. occ.
tuberculata, *Berg, in Mart. Fl. Bras.* xiv. 1. 52.—Bras.
Tussaccana, *Berg, in Linnaea*, xxvii. (1854) 25.—Ind. occ.
uniflora, *Spreng. in Flora*, xx. (1837) *Beibl.* ii. 87.—Bras.
variabilis, *Berg, in Mart. Fl. Bras.* xiv. 1. 49.—Bras.
Widgreniana, *Berg, l. c.* 39.—Bras.
Wilsonii, *Griseb. Fl. Brit. W. Ind.* 233.—Ind. occ.
Zuzygium, Blanco, Fl. Filip. ed. II. 293 = Eugenia operculata.
Zuzygium, Sw. Prod. Veg. Ind. Occ. 79 = C. Syzygium.

CALYPTRANTHUS, Thou. Obs. Pl. Isles Afr. 26 (1811) = **Capparis**, Tourn.

CALYPTRANTHUS, Blume, Bijdr. 1089 (1826) = **Eugenia**, Mich. (Myrtac.).

CALYPTRANTHUS, Juss. in Dict. Sc. Nat. vi. 274 (1817) = **Calyptranthes**, Sw. (Myrtac.).

CALYPTRARIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 132 = **Centronia**, D. Don (Melastom.).
africana, DC. Prod. ix. 469, in syn. = Varronia abyssinica.
brachycera, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 134 = Cent. brachycera.
eximia, Naud. l. c. 133 = Cent. eximia.
haemantha, Planch. & Linden, in Fl. des Serres, Sér. 1. ix. (1853–54) t. 924 = Cent. haemantha.
insignis, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1851) 133 = Cent. insignis.
Mutisii, Naud. l. c. 132 = Cent. Mutisii.

CALYPTRELLA, Naud, in Ann. Sc. Nat. Sér. III. xviii. (1852) 115. *MELASTOMACEAE*, Benth. & Hook. f. i. 750.
cucullata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 72.—Peruv.
Galeotti, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 115.—Mexic.
gracilis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 72.—Peruv.
tristis, *Triana, l. c.*—Peruv.

CALYPTRIDIUM, Nutt. ex Torr. & Gray, Fl. N. Am. i. 198 (1838). *PORTULACEAE*, Benth. & Hook. f. i. 159.
monandrum, *Nutt. l. c.*—Calif.
quadripetalum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 356.—Calif.
roseum, *S. Wats. in Bot. King, Exp.* 44. t. 6.—Calif.

CALYPTRION, Ging. in DC. Prod. i. 288 (1824) = **Corynostylis**, Mart. (Violac.).
Aubletii, Ging. l. c. 289 = Cor. Hybanthus.
Berterii, Ging. l. c. = Cor. Berterii.
citrifolium, Ging. l. c. = Cor. Loeflingii.
diandrum, Ging. l. c. = Cor. diandra.
frangulaefolium, Ging. l. c.—Peruv.
nitidum, Benth. in Hook. Journ. Bot. iv. (1842) 106 = Cor. Hybanthus.
orinocense, H. B. & K. Nov. Gen. et Sp. v. 384.—Venezuela.
pyrifolium, Sweet, ex Steud. Nom. ed. II. i. 263 = Anchietea pyrifolia.

CALYPTRIOPETALUM, Hassk. ex Mucll. Arg. in DC. Prod. xv. II. 513 (1866) = **Croton**, Linn. (Euphorb.).

CALYPTROCALYX, Blume, Rumphia, ii. 103. t. 102 D, 118, 161 (1836). *PALMAE*, Benth. & Hook. f. Gen. iii. 902.
LACCOSPADIX, H. Wendl. & Drude, in Linnaea, xxxix. (1875) 205.
australasicus, *Scheff. in Ann. Jard. Bot. Buitenz.* i. (1876) 132, 140.—Austral.
spicatus, *Blume, Rumphia*, ii. 103.—Amboina.

CALYPTROCARPUS, Reichb. Nom. 82 (1841) = CALYPTOCARPUS, Less. = **Blainvillea**, Cass. (Compos.).

CALYPTROCARYA, Nees, in Linnaea, ix. (1834) 304. *CYPERACEAE*, Benth. & Hook. f. Gen. iii. 1069.
angustifolia, *Nees, l. c.*—Guiana.
bicolor, Nees, l. c. = Poeppigiana.
brevicaulis, *Nees, l. c.* 365.—Guiana.
fragifera, *Nees, l. c.* 304.—Am. trop.
longifolia, Boeck. in Linnaea, xxxviii. (1874) 431 = C. Palmetto.
longifolia, *Kunth, Enum. Pl.* ii. 365.—Guiana.
Martii, *Nees, in Mart. Fl. Bras.* ii. 1. 195.—Bras.
Palmetto, *Nees, l. c.*—Am. trop.
Poeppigiana, *Kunth, Enum. Pl.* ii. 364.—N. Granat.
Schottmuelleriana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 509.—Bras.

CALYPTROCORYNE, Schott, in Oestr. Bot. Wochenschr. (1857) 262 = **Theriophonum**, Blume (Aroid.).
cochinchinensis, Schott, Prod. Aroid. 105 = T. Wightii.
minuta, Schott, l. c. = T. Wightii.
Wightii, Schott, in Oestr. Bot. Wochenschr. (1857) 262 = T. Wightii.

CALYPTROGYNE, H. Wendl. in Bot. Zeit. xvii. (1859) 72. *PALMAE*, Benth. & Hook. f. iii. 914.
CALYPTRONOMA, Griseb. Fl. Brit. W. Ind. 518 (1864).
Ghiesbrechtiana, *H. Wendl. l. c.*—Mexic.
Ghiesbreghti, H. Wendl. in Kerch. Palm. 238 = Ghiesbrechtiana.
glauca, *H. Wendl. l. c.*—Nicaragua.
sarapiquensis, *H. Wendl. l. c.*—Costa Rica.
spicigera, *H. Wendl. in Bot. Zeit.* xvii. (1859) 72.—Guatemala.

CALYPTROLEPIS, Steud. Syn. Pl. Cyp. 151 (1855) = **Rynchospora**, Vahl (Cyperac.).
junciformis, Steud. l. c. = R. oligantha.

CALYPTROMYRCIA, Berg, in Linnaea, xxvii. (1854) 34 = **Myrcia**, DC. (Myrtac.).
cordata, Berg, in Mart. Fl. Bras. xiv. 1. 56 = M. cordata.
costata, Berg, l. c. = M. costata.
cymosa, Berg, l. c. 58.—Bras.
elegans, Berg, l. c. = M. elegans.
eugenioides, Berg, l. c. 56 = M. eugenioides.
paniculata, Berg, l. c. 57.—Bras.
puberula, Berg, l. c. = M. puberula.
Spixiana, Berg, l. c. 59 = M. Spixiana.
venosa, Berg, l. c. 57.—Bras.

CALYPTRONOMA, Griseb. Fl. Brit. W. Ind. 518 (1864) = **Calyptrogyne**, H. Wendl. (Palm.).
dulcis, H. Wendl. in Kerch. Palm. 238 (= *Geonoma dulcis*).
intermedia, H. Wendl. l. c. nomen.—Hab.?
robusta, Traill, in Journ. Bot. xiv. (1876) 330.—Bras.
Schwartzii, H. Wendl. in Kerch. Palm. 238 = seq.
Swartzii, Griseb. Fl. Brit. W. Ind. 518 (= *Geonoma Swartzii*).

CALYPTROON, Miq. Fl. Ind. Bat. Suppl. 471 (1860) = **Baccaurea**, Lour. (Euphorbiac.).
sumatranum, Miq. l. c. 472 = B. sumatrana.

CALYPTROPSIDIUM, Berg, in Linnaea, xxvii. (1854) 349 = **Psidium**, Linn. (Myrtac.).
Friedrichsthalianum, Berg, l. c. 350.—Guatemala.

CALYPTROSPATHA, Klotzsch, ex Baill. Etud. Gen. Euph. 440 (1858); et in Peters, Reise Mossamb. Bot. t. 18 (1861) = **Acalypha**, Linn. (Euphorb.).
pubiflora, Klotzsch, l. c. 97. t. 18 = A. pubiflora.

CALYPTROSPERMUM, A. Dietr. Sp. Pl. i. 226 (1831) = **Menodora**, Humb. & Bonpl. (Oleac.).
integrifolium, A. Dietr. l. c. 227 = M. integrifolia.
trifidum, A. Dietr. l. c. = M. trifida.

CALYPTROSTEGIA, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 72 pro parte = **Pimelea**, Banks (Thymel.).
angustifolia, C. A. Mey. l. c. 74 = P. angustifolia.
argentea, C. A. Mey. l. c. = P. argentea.
brevifolia, C. A. Mey. l. c. = P. brevifolia.
cernua, C. A. Mey. l. c. = P. spathulata.
cluytioides, Walp. Ann. iii. 324 = P. serpyllifolia.
Cornucopiae, Endl. Gen. Suppl. iv. II. 60 = P. Cornucopiae.
curviflora, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. curviflora.
Drummondii, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 178 = P. suaveolens.
flava, Endl. Gen. Suppl. iv. II. 60 = P. flava.
glauca, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. glauca.
graciliflora, Endl. Gen. Suppl. iv. II. 61 = P. sylvestris.
gracilis, Endl. l. c. = P. curviflora.
hypericina, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. ligustrina.
intermedia, C. A. Mey. l. c. = P. glauca.
lanata, Endl. Gen. Suppl. iv. II. 61 = P. hispida.
latifolia, Endl. l. c. = P. latifolia.
Lehmanniana, Endl. l. c. = P. Lehmanniana.
ligustrina, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. ligustrina.
linifolia, C. A. Mey. l. c. = P. linifolia.
linoides, Endl. Gen. Suppl. iv. II. 61 = P. linifolia.
longiflora, Endl. l. c. = P. longiflora.
macrocephala, Walp. Ann. iii. 325 = P. suaveolens.
Menkeana, Endl. Gen. Suppl. iv. II. 61 = P. suaveolens.
microcephala, Endl. l. c. = P. microcephala.
myriantha, Endl. l. c. = P. argentea.
nana, Endl. l. c. = P. imbricata.
nervosa, Walp. Ann. iii. 324 = P. nervosa.
phylicoides, Walp. l. c. = P. phylicoides.
punicea, Endl. Gen. Suppl. iv. II. 60 = P. punicea.
Schlechtendaliana, Walp. Ann. iii. 324 = P. glauca.
spathulata, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. spathulata.
spicata, Endl. Gen. Suppl. iv. II. 61 = P. spicata.

CALYPTROSTEGIA :—
suaveolens, Endl. Gen. Suppl. iv. II. 61 = Pimelea suaveolens.
sulphurea, Walp. Ann. iii. 325 = P. sulphurea.
sylvestris, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iv. (1845) 74 = P. sylvestris.
tinctoria, Endl. Gen. Suppl. iv. II. 61 = P. suaveolens.
trichostachya, Walp. Ann. iii. 325 = P. trichostachya.
villifera, Walp. l. c. 324 = P. villifera.
villosa, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 178 = P. longiflora.

CALYPTROSTIGMA, Klotzsch, in Lehm. Pl. Preiss. i. 175 (1844) = **Beyeria**, Miq. (Euphorbiac.).
ledifolium, Klotzsch, l. c. 176 = B. Drummondii.
oblongifolium, Klotzsch, l. c. = B. viscosa.
viscosum, Klotzsch, l. c. = B. viscosa.

CALYPTROSTIGMA, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 46) (1847) = **Diervilla**, Tourn. (Caprifol.).
Middendorffianum, Trautv. & Mey. l. c. = D. Middendorffiana.

CALYPTROSTYLIS, Nees, in Linnaea, ix. (1834) 295 = **Rynchospora**, Vahl (Cyperac.).
articulata, Nees, in Hook. Journ. Bot. ii. (1840) 394, in obs. = R. aurea.
asperula, Nees, in Mart. Fl. Bras. ii. I. 139 = R. aurea.
capitata, Nees, l. c. 140 = R. minarum.
divergens, Nees, l. c. 139 = R. aurea.
fascicularis, Nees, l. c. 139 = R. aurea.
florida, Nees, l. c. 138 = R. aurea.
Gaudichaudii, Nees, in Linnaea, ix. (1834) 295 = R. aurea.
gigantea, Nees, in Mart. Fl. Bras. ii. I. 137 = R. gigantea.
longirostris, Nees, in Hook. Journ. Bot. ii. (1840) 394 = R. amazonica.
paniculata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 252 = R. Schiedeana.
Rudgei, Nees, in Linnaea, ix. (1834) 295 = R. gigantea.
Schiedeana, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 252 = R. Schiedeana.
terminalis, Nees, ex Steud. Syn. Pl. Cyp. 143 = R. minarum.
zeylanica, Nees, in Hook. Journ. Bot. ii. (1840) 394, in obs. = R. triflora.

CALYSACCION, Wight, Illustr. i. 130 (1840) = **Ochrocarpus**, Thou. (Guttifer.).
chinense, Walp. in Bot. Zeit. ix. (1851) 368 = O. longifolius.
Horstii, Teijsm. & Binn. Cat. Hort. Bog. 205.—Borneo.
longifolium, Wight, Illustr. i. 130 = O. longifolius.
obovale, Miq. Fl. Ind. Bat. Suppl. i. 500 = O. ovalifolius.
ovalifolium, Choisy, Guttif. Ind. 45 = O. ovalifolia.
siamense, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 7, 209 = O. siamensis.
tinctorium, Seem. Fl. Vit. 13.—Ins. Fiji.

CALYSERICOS, Eckl. & Zeyh. ex Meissn. in DC. Prod. xiv. 573 (1856–57) = **Cryptadenia**, Meissn. (Thymelaeac.).
argentea, Eckl. & Zeyl. l. c. 582 = Gnidia Cephalotes.
canaliculata, Eckl. & Zeyh. l. c. 573 = Cr. grandiflora.
grandiflora, Eckl. & Zeyh. l. c. = Cr. grandiflora.
linarifolia, Eckl. & Zeyh. l. c. 583 = Gnidia linoides.
parviflora, Eckl. & Zeyh. l. c. 574 = Cr. breviflora.
subulata, Eckl. & Zeyh. l. c. = Cr. breviflora.
tabularis, Eckl. & Zeyh. l. c. = Cr. breviflora.
typica, Eckl. & Zeyl. l. c. 581 = Gnidia pubescens.
uniflora, Eckl. & Zeyh. l. c. 573 = Gnidia uniflora.

CALYSPHYRUM, Bunge, in Mém. Sav. Etr. Acad. Pétersb. ii. (1835) 107; Enum. Pl. Chin. Bor. 33 = **Diervilla**, Tourn. (Caprifol.).
floridum, Bunge, l. c. = D. florida.
pauciflorum, Bunge, in Walp. Rep. ii. 450 = D. florida.
roseum, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. xiii. (1855) 220 = D. rosea.

CALYSTEGIA, R. Br. Prod. 483 (1810). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 874.
MILHANIA, Rafin. Fl. Tellur. iv. 71 (1836).
VOLVULUS, Medic. in Staatsw. Vorles. Churpf. Phys. Oek. Ges. i. (1791) 202.

CALYSTEGIA :—
acetosifolia, Turcz. Fl. Baic. Dahur. ii. II. 289 = hederacea.
affinis, *Endl. Prod. Fl. Norf.* 51.—Ins. Norfolk.
asarifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 344 = C. Soldanella.
barbara, *Pomel, Nouv. Mat. Fl. Atl.* 83.—Afr. bor.
Basniniana, Turcz. ex Ledeb. Fl. Ross. iii. 95 = pellita.
Catesbeiana, *Pursh, Fl. Am. Sept. Suppl.* ii. 729.—Am. bor.
chinensis, *Baill. in Bull. Soc. Linn. Par.* i. (1881) 284.—China.
dahurica, Choisy, in DC. Prod. ix. 433 = pellita.
dahurica, *Fisch. ex Planch. in Fl. des Serres*, x. (1854–55) 245. t. 1075.—Sibir.
grandiflora, *G. Don, Gen. Syst.* iv. 297.—Ind. or.
Hantelmanni, *Phil. in Linnaea*, xxx. (1859–60) 196.—Chili.
hederacea, *Wall. Cat.* n. 1328.—Ind. or.; China; Afghan.
inflata, G. Don, in Sweet, Hort. Brit. ed. III. 486 = sepium.
japonica, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 130, 132.—Japon.
Keriana, Sweet, Hort. Brit. ed. I. 287 = Hewittia bicolor.
Krauseana, *Phil. in Linnaea*, xxxiii. (1864–65) 184.—Chili.
lucana, G. Don, Gen. Syst. iv. 296 = sylvestris.
marginata, *R. Br. Prod.* 484.—Australas.
Maximiliana, *Nees, in Neuwied, Reise Nord. Am.* ii. 443.—Am. austr.?
mucronata, Spreng. ex Choisy, in DC. Prod. ix. 430 = Ipomoea martinicensis.
mutabilis, Rafin. Fl. Ludov. 46 = sepium?
obtusa, *Pomel, Nouv. Mat. Fl. Atl.* 295.—Afr. bor.
ochroleuca, *Boj. Hort. Maurit.* 231.—Afr. trop.
palmato-partita, Benth. & Hook. f. Gen. ii. 874 = seq.
palmato-pinnata, Meissn. in Mart. Fl. Bras. vii. 317 = Ipomoea palmato-pinnata.
paradoxa, Pursh, Fl. Am. Sept. Suppl. ii. 729 = Convolvulus hirsutus.
pellita, *G. Don, Gen. Syst.* iv. 296.—Sibir. baical.
physoides, *Pomel, Nouv. Mat. Fl. Atl.* 294.—Afr. bor.
pubescens, Lindl. in Journ. Hort. Soc. i. (1846) 70 = hederacea.
reniformis, R. Br. Prod. 483 = C. Soldanella.
riparia, *Rafin. New Fl. Am.* ii. 29.—Am. bor.
rosea, *Phil. in Linnaea*, xxix. (1857–58) 15.—Chili.
rugosa, *Ledeb. Ind. Hort. Dorp.* (1821) 4.—Hab.?
sagittata, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 356.—Am. bor.
Scammonia, Pritz. Ind. Ic. ii. 58 = Convolvulus Scammonia.
sepium, *R. Br. Prod.* 483.—Cosmop. temp.
setifera, Meissn. in Mart. Fl. Bras. vii. 316 = Ipomoea setifera.
silvatica, *Choisy, in DC. Prod.* ix. 483.—Europ. austr. or.; Reg. Caucas.; Afr. bor.
Soldanella, *R. Br. Prod.* 483.—Cosmop. temp.
spithamaea, *Pursh, Fl. Am. Sept.* i. 143.—Am. bor.
stans, Steud. Nom. ed. II. i. 264 = spithamaea?
subacaulis, *Hook. & Arn. Bot. Beech. Voy.* 363.—Calif.
subvolubilis, G. Don, Gen. Syst. iv. 296 = pellita.
sylvestris, *Roem. & Schult. Syst.* iv. 183.—Oriens.
tomentosa, Pursh, Fl. Am. Sept. i. 143 = spithamaea.
tuguriorum, *R. Br. ex Hook. Fl. Nov. Zel.* i. 183. t. 47.—N. Zeland.; Chili.
valdiviana, *Miq. in Lechl. Berb. Am. Austr.* 54.—Chili.
villosa, *Kellogg, in Proc. Calif. Acad.* v. (1873) 17.—Calif.
villosa, *Rafin. Fl. Ludov.* 45.—Am. bor.

CALYTHRIX, Labill. Nov. Holl. Pl. ii. 8. t. 146 (1806) (Calytrix). *MYRTACEAE*, Benth. & Hook. f. i. 699.
CALYCOTHRIX, Meissn. Gen. 107 (1837).
TRICHOCALYX, Schau. Myrt. Xeroc. 86, adnot. (1840).
achaeta, *F. Muell. in Trans. Phil. Vict. Inst.* iii. (1859) 43.—Austral.
angulata, *Lindl. Swan River App.* 6.—Austral.
arborescens, *F. Muell. in Trans. Phil. Vict. Inst.* iii. (1859) 42.—Austral.
asperula, *Schau. in Lehm. Pl. Preiss.* i. 104.—Austral.
aurea, *Lindl. Swan River App.* 5. t. 3.—Austral.
Baueri, Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 261 = tetragona.

CALYTHRIX :—
Behriana, Schlecht. in Linnaea, xx. (1847) 650 = tetragona.
Billardieri, Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 245 = tetragona.
Birdii, *F. Muell. Fragm.* x. 26.—Austral.
brachychaeta, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 43.—Austral.
brachyphylla, *A. Cunn. ex Steud. Nom.* ed. II. i. 264.—Austral.
brachyphylla, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 161.—Austral.
brevifolia, *Meissn. in Journ. Linn. Soc.* i. (1857) 46.—Austral.
breviseta, *Lindl. Swan River App.* 5.—Austral.
Brownii, Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 260 = tetragona.
brunioides, A. Cunn. in Bot. Mag. sub t. 3323 = tetragona.
Candolleana, Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 260 = C. decandra.
ciliata, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 161 = empetroides.
conanthera, F. Muell. Fragm. i. 146 = decandra.
conferta, *A. Cunn. in Bot. Mag.* sub t. 3323.—Austral.
Creswelli, *F. Muell. Fragm.* x. 27.—Austral.
cupressifolia, A. Rich. Sert. Astrol. 41. t. 16 = microphylla.
cupressoides, A. Rich. l. c. 43 = microphylla.
curtophylla, A. Cunn. in Bot. Mag. sub t. 3323 = flavescens.
curtophylla, Schau. in Nov. Act. Nat. Cur. xix. Suppl. ii. (1841) 242 = Leschenaultii.
cuspidata, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 162 = breviseta.
decandra, *DC. Prod.* iii. 208.—Austral.
depressa, *Turcz. in Bull. Soc. Nat. Mosc.* xx. (1847) I. 162.—Austral.
diversifolia, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 328.—Austral.
Drummondii, Meissn. in Journ. Linn. Soc. i. (1857) 47 = flavescens.
empetroides, *Schau. in Nov. Act. Nat. Cur.* xix. Suppl. ii. (1841) 254.—Austral.
ericoides, A. Cunn. in Field, N. S. Wales, 350 = tetragona.
exstipulata, DC. Prod. iii. 208 = microphylla.
flavescens, *A. Cunn. in Bot. Mag.* sub t. 3323.—Austral.
Fraseri, *A. Cunn. l. c.*—Austral.
glabra, R. Br. in Bot. Reg. t. 409 = tetragona.
glutinosa, *Lindl. Swan River App.* 5.—Austral.
gracilis, *Benth. Fl. Austral.* iii. 45.—Austral.
granulosa, *Benth. l. c.* 49.—Austral.
laricina, *R. Br. ex Benth. l. c.* 52.—Austral.
lasiantha, Meissn. in Journ. Linn. Soc. i. (1857) 46 = strigosa.
lasiostachya, F. Muell. Fragm. i. 224 = sapphirina.
leptophylla, *Benth. Fl. Austral.* iii. 50.—Austral.
Leschenaultii, *Schau. in Lehm. Pl. Preiss.* i. 104.—Austral.
leucantha, Miq. in Nederl. Kruidk. Arch. iv. (1859) 117 = tetragona.
longiflora, *F. Muell. Fragm.* i. 12.—Austral.
luteola, Schau. in Lehm. Pl. Preiss. i. 106 = flavescens.
megaphylla, *F. Muell. Fragm.* i. 146.—Austral.
microphylla, *A. Cunn. in Bot. Mag.* sub t. 3323.—Austral.
monticola, Miq. in Nederl. Kruidk. Arch. iv. (1859) 118 = tetragona.
Muelleri, Miq. l. c. 119 = tetragona.
muricata, *F. Muell. Fragm.* i. 224.—Austral.
Oldfieldii, *Benth. Fl. Austral.* iii. 46.—Austral.
plumulosa, *F. Muell. Fragm.* x. 27.—Austral.
propinqua, A. Cunn. ex Walp. Rep. ii. 158 = tetragona.
puberula, *Meissn. in Journ. Linn. Soc.* i. (1857) 48.—Austral.
pubescens, Sweet, ex Loud. Hort. Brit. 199 = tetragona.
pulchella, *Turcz. in Bull. Phys.-Math. Acad. Pétersb.* x. (1852) 328.—Austral.
rosea, Meissn. in Journ. Linn. Soc. i. (1847) 46 = tenuifolia.
rosea, Miq. in Nederl. Kruidk. Arch. iv. (1859) 117 = tetragona.
sapphirina, *Lindl. Swan River App.* 5.—Austral.
scabra, DC. Prod. iii. 208 = tetragona.

CALYTHRIX:—
Schlechtendahlii, Miq. in Nederl. Kruidk. Arch. iv. (1859) 116 = tetragona.
simplex, *Lindl. Swan River App.* 5.—Austral.
sparsa, A. Cunn. ex Walp. Rep. ii. 160 = strigosa.
squarrosa, Miq. in Nederl. Kruidk. Arch. iv. (1859) 118 = tetragona.
strigosa, *A. Cunn. in Bot. Mag.* sub t. 3323.—Austral.
Sullivani, *F. Muell. Fragm.* ix. 1.—Austral.
tenella, Meissn. in Journ. Linn. Soc. i. (1857) 47 = flaveseens.
tenuifolia, *Meissn. l. c.* 46.—Austral.
tenuiramea, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 20.—Austral.
tetragona, *Labill. Nov. Holl. Pl.* ii. 8. t. 146.—Austral.
tetragonophylla, Meissn. in Journ. Linn. Soc. i. (1857) 47 = flaveseens.
tetraptera, DC. Prod. iii. 208 = tetragona.
variabilis, *Lindl. Swan River App.* 5.—Austral.
virgata, A. Cunn. in Bot. Mag. sub t. 3323 = tetragona.

CALYTRIPLEX, Ruiz & Pav. Prod. 96 (1794) = **Herpestis**, Gaertn. (Scrophular.).
obovata, Ruiz & Pav. Syst. Veg. 165 = H. Monniera.

CALYXHYMENIA, Orteg. Hort. Matr. Dec. 5. t. 1 (1798) = **Oxybaphus**, Vahl (Nyetagin.).
aggregata, Orteg. l. c. 81. t. 11 = O. aggregatus.
Cervantesii, Desf. Cat. Hort. Par. ed. III. 390 = O. Cervantesii.
chilensis, Desf. ex Steud. Nom. ed. II. i. 264 = O. chilensis.
expansa, Ruiz & Pav. Fl. Per. i. 45 = O. expansus.
glabrifolia, Orteg. Hort. Matr. 5. t. 1 = O. glabrifolius.
ovata, Ruiz & Pav. Fl. Per. i. 45 = O. ovatus.
paniculata, Desf. Cat. Hort. Par. ed. III. 390 = O. ovatus.
pilosa, Choisy, in DC. Prod. xiii. II. 433 = O. angustifolius.
prostrata, Ruiz & Pav. Fl. Per. i. 46 = O. prostratus.
viscosa, Orteg. Hort. Matr. Dec. i. 6, in obs. = O. viseosus.

CAMACUM, 'Adans.' ex Steud. Nom. ed. II. ii. 174, in syn. (1841) = **Myristica**, Linn.

CAMAION, Rafin. Sylva Tellur. 75 (1838) = **Helicteres**, Linn. (Stereuliae.).
hirsuta, Rafin. l. e. = H. hirsuta.
undulata, Rafin. l. e. = Sterculia lanceolata.

CAMARA, Plum. ex Linn. Syst. ed. I (1735); Adans. Fam. ii. 199 (1763) = **Lantana**, Linn. (Verbenae.).
tiliaefolia, Benth. in Hook. Journ. Bot. ii. (1840) 53 = L. tiliaefolia.
vulgaris, Benth. Bot. Voy. Sulph. 154 = L. Camara.

CAMAREA, St. Hil. in Bull. Soc. Philom. (1823) 133.
MALPIGHIACEAE, Benth. & Hook. f. i. 261.
affinis, *St. Hil. Pl. Rem.* 57.—Bras.
axillaris, *St. Hil. in Bull. Soc. Philom.* (1823) 133.—Bras.
discolor, *Griseb. in Mart. Fl. Bras.* xii. 1. 105.—Bras.
ericoides, *St. Hil. in Bull. Soc. Philom.* (1823) 133.—Bras.
hirsuta, *St. Hil. l. c.*—Bras.
juncea, *Griseb. in Mart. Fl. Bras.* xii. 1. 105.—Bras.
linearifolia, *St. Hil. in Bull. Soc. Philom.* (1823) 133.—Bras.
phylicoides, Mart. ex Walp. Rep. v. 351 = axillaris.
pulchella, *Griseb. in Mart. Fl. Bras.* xii. 1. 105.—Bras.
sericea, *St. Hil. in Bull. Soc. Philom.* (1823) 133.—Bras.
triphylla, *A. Juss. in Arch. Mus. Par.* iii. (1843) 601.—Bras.

CAMARIDIUM, Lindl. Bot. Reg. sub t. 844 (1824).
ORCHIDEAE, Benth. & Hook. f. Gen. iii. 556.
AMARIDIUM, Hort. ex Lubbers, Cat. Pl. Rar. San Donato, 15 (1880).
alatum, *Lindl. in Benth. Pl. Hartw.* 153.—Ecuador.
Arbuscula, *Lindl. l. c.*—Ecuador.
brevifolium, Lindl. l. c. 154.—N. Granat.

CAMARIDIUM:—
graminifolium, *Reichb. f. in Linnaea*, xxii. (1849) 857.—Peruv.
inflexum, *Lindl. in Ann. Nat. Hist.* v. (1840) 116.—Ins. Dominica.
laneifolium, *Reichb. f. in Linnaea*, xxii. (1849) 857.—Peruv.
longibraeteatum, *Lindl. in Benth. Pl. Hartw.* 154.—Peruv.
luteo-rubrum, *Lindl. Orch. Linden.* 22.—Venezuela.
lutescens, *Reichb. f. in Walp. Ann.* vi. 541.—Bras.
ochroleucum, Lindl. Bot. Reg. x. (1824) 844 = Maxillaria Camardii.
pendulum, *Rodrig. Orch. Nov.* i. 123.—Bras.
purpuratum, *Lindl. Orch. Linden.* 22.—Venezuela.
purpureum, Spreng. Syst. iii. 735 = Cymbidium vestitum.
robustum, *Rodrig. Orch. Nov.* ii. 210.—Bras.
vestitum, Lindl. in Ann. & Mag. Nat. Hist. Ser. III. i. (1858) 333 = Ornithidium vestitum.

CAMARILLA, Salisb. Gen. Pl. Fragm. 91 (1866) = **Allium**, Tourn. (Liliae.).
obliqua, Salisb. l. c. = A. obliquum.

CAMAROTIS, Lindl. Gen. et Sp. Orch. 219 (1833) = **Sarcochilus**, R. Br. (Orchid.).
apiculata, Reichb. f. in Bonplandia, v. (1857) 39.—Malaya.
cochinchinensis, Reichb. f. in Gard. Chron. (1876) II. 740.—Cochineh.
obtusa, Lindl. Bot. Reg. xxx. (1844) Misc. 71 = S. obtusus.
pallida, Lindl. in Journ. Linn. Soc. iii. (1859) 37 = S. pallidus.
philippinensis, Lindl. l. e. = S. philippinensis.
purpurea, Lindl. in Wall. Cat. n. 7329 = S. purpureus.
rostrata, Reichb. f. in Walp. Ann. vi. 881 = S. purpureus.

CAMASSIA, Lindl. Bot. Reg. t. 1486 (1832). *LILIACEAE*, Benth. & Hook. f. iii. 815.
BULBEDULIS, Rafin. Fl. Tellur. ii. 26 (1836).
CYANOTRIS, Rafin. in Am. Monthly Mag. (1811) 356; (1819) 192, nomen prius.
LEMOTRIS, Rafin. Fl. Tellur. ii. 26 (1836).
QUAMASIA, Rafin. Cat. 15 (1824); QUAMASSIA, Rafin. Fl. Tellur. ii. 25 (1836).
SITOCODIUM, Salisb. Gen. Pl. Fragm. 27 (1866).
alba, Hort.—Cf. Gard. Chron. (1875) I. 662 = Leichtlinii.
esculenta, *Lindl. Bot. Reg.* t. 1486.—Am. bor. oec.
Fraseri, *Torr. in Rep. Pacif. Rail.* iv. 91.—Am. bor. oec.
Leichtlinii, *S. Wats. in Proc. Am. Acad.* xx. (1885) 376.—Calif.

CAMAX, Sehreb. Gen. i. 144 (1789) = **Ropourea**, Aubl. (Verbenae.?).
fraxinea, Willd. Sp. Pl. i. 1117 = R. guianensis.
guianensis, J. F. Gmel. Syst. 403 = R. guianensis.

CAMBANIA, Comm. ex M. Roem. Syn. Hesper. 83, 102 (1846) = **Dysoxylum**, Blume (Meliac.).
Fraseriana, M. Roem. l. c. 102 = D. Fraseranum.

CAMBDERIA, Steud. Nom. ed. II. i. 264 (1840) = CAMPDERIA, Rich. = **Vellozia**, Vand. (Amaryll.).

CAMBEA, Endl. Gen. 1234, in Syn. (1840) sphalm. = CUMBIA, Buch.-Ham. = **Careya**, Roxb. (Myrtae.).

CAMBESSEDEA, Wight & Arn. Prod. i. 170 (1834) = **Bouea**, Meissn. (Anacard.).
oppositifolia, Wight & Arn. l. e. = B. burmanica.

CAMBESSEDEA, Kunth, in Ann. Sc. Nat. Sér. I. ii. 336 (1824) = **Buchanania**, Roxb. (Anacard.).
axillaris, Kunth, l. c. = B. angustifolia.

CAMBESSEDESIA, DC. Prod. iii. 110 (1828).
MELASTOMACEAE, Benth. & Hook. f. i. 737.
ACIPETALUM, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) I. 577, pro parte.
adamantium, *DC. Prod.* iii. 111.—Bras.
adamantium, Miq. in Linnaea, xxii. (1849) 537 = ilicifolia.
arenaria, *Cogn. in Mart. Fl. Bras.* xiv. III. 15.—Bras.

CAMBESSEDESIA:—
balsamifera, DC. Prod. iii. 110 = Microlicia balsamifera.
bidentata, DC. l. c. 111 = Hilariana.
corymbosa, *DC. l. c.* 110.—Bras.
corymbosa, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 61 = rugosa.
crenulata, DC. Prod. iii. 111 = Mierolicia crenulata.
Eichleri, *Cogn. in Mart. Fl. Bras.* xiv. III. 11.—Bras.
Espora, *DC. Prod.* iii. 111.—Bras.
Espora, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 62 = ilicifolia.
Hilariana, *DC. Prod.* iii. 111.—Bras.
ilicifolia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 24.—Bras.
intermedia, Mart. Herb. Fl. Bras. n. 940 = rugosa.
latevenosa, *DC. Prod.* iii. 110.—Bras.
membranaeea, *Gardn. in Field. Sert. Pl.* t. 64.—Bras.
paraguayensis, *Hook. f. Bot. Mag.* t. 6604.—Paraguay.
purpurata, *DC. Prod.* iii. 110.—Bras.
Regnelliana, *Cogn. in Mart. Fl. Bras.* xiv. III. 19.—Bras.
rugosa, *Cogn. l. c.* 14.—Bras.
setaeea, *Cogn. l. c.* 23.—Bras.
sincorensis, DC. Prod, iii. 111 = Mierolicia sincorensis.
terminalis, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 61 = Hilariana,
Weddelii, *Naud. l. c.*—Bras.

CAMBOGIA, Linn. Pl. Mart. Burser. (1745); Diss. Dass. 18 (1747) (Amoen. Acad. i. 171, 402) = **Garcinia**, Linn. (Guttifer.).
Binucao, Blanco, Fl. Filip. ed. I. 434 = G. Cambogia.
crassifolia, Blaneo, l. c. ed. II. 304 = G. Cowa.
Gutta, Linn. Gen. ed. V. Add. Fin. = G. Cambogia.
solitaria, Stokes, Bot. Mat. Med. iii. 175 = G. Cambogia.
venulosa, Blaneo, Fl. Filip. ed. I. 435 = G. venulosa.

CAMDENIA, Seop. Introd. 190 (1777) = **Evolvulus**, Linn. (Convolvul.).

CAMDERIA, Dum. Anal. Fam. 80 (1829) = **Lachnanthes**, Ell. (Haemod.).

CAMELINA, Crantz, Stirp. Austr. ed. I. fasc. i. 18 (1762); ed. II. 18 (1769). *CRUCIFERAE*, Benth. & Hook. f. i. 83.
CHAMAELINUM, Host, Fl. Austr. ii. 224 (1831).
LINOSTROPHUM, Schrank, Fl. Prim. Salisb. 163 (1792).
SINISTROPHORUM, Sehrank, ex Endl. Gen. 882 (1840).
albiflora, Kotsehy, ex Boiss. Fl. Orient. i. 312 = sylvestris.
ambigua, Bess. ex Steud. Nom. ed. II. i. 264 = sativa.
anomala, *Boiss. & Haussk. ex Boiss. Fl. Orient.* i. 313.—Syria.
aquatica, Brot. Fl. Lusit. i. 560 = Nasturtium amphibium.
armeniaca, Desv. Journ. Bot. iii. (1814) 182 = sylvestris.
austriaca, Pers. Syn. ii. 191 = Nasturtium austriaeum.
barbaraefolia, DC. Syst. ii. 517 (= *Tetrapoma barbaraefolium*.).
Caisir, Wall. Cat. n. 4802 = Coehlearia flava.
Camisir, Buch.-Ham. ex Steud. Nom. ed. II. i. 264, sphalm. = prace.
campestris, Schimp. & Spenn. in Spenn. Fl. Friburg. iii. 959 = sylvestris.
dentata, Pers. Syn. ii. 191 = sativa.
edentula, Desv. Journ. Bot. iii. (1814) 179 = Alyssum edentulum.
foetida, *Fries, Nov. Mant.* iii. 70.—Europ.
grandiflora, *Boiss. Diagn.* Ser. I. v. 82.—As. Min.
hirsuta, Bernh. Syst. Verz. Erf. 190 = sativa!
hispida, *Boiss. in Ann. Sc. Nat.* Ser. II. xvii. (1842) 176.—Armenia; Persia.
lasiocarpa, *Boiss. & Bal. ex Boiss. Fl. Orient.* i. 312.—Syria.
laxa, *C. A. Mey. Verz. Pfl. Cauc.* 193.—As. Min.; Persia.
linicola, *Schimp. & Spenn. in Spenn. Fl. Friburg.* iii 958.—Europ.

CAMELINA:—
macrocarpa, Wierzb. ex Reichb. Ic. Fl. Germ. ii. 10 = sativa.
microcarpa, Andrz. ex DC. Syst. ii. 517 = sylvestris.
myagrodes, Moretti, in Giorn. Fis. Pav. (1820) 279 = Cochlearia saxatilis.
pinnatifida, Hornem. Hort. Hafn. ii. 598 = foctida.
sagittata, Moench, Meth. 265 = sativa.
sativa, Boiss. Voy. Bot. Espagne, 49 = sylvestris.
sativa, *Crantz, Stirp. Austr.* ed. I. fasc. i. 14; ed. II. 18.—Europ.; As. temp.
saxatilis, Pers. Syn. ii. 191 = Cochlearia saxatilis.
sylvestris, *Wallr. Sched. Crit.* 347.—Europ.; As. Min.

CAMELLIA, Linn. Syst. ed. I (1735). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 187.
Calpandria, Blume, Bijdr. 178 (1825).
Desmitus, Rafin. Sylva Tellur. 139 (1838).
Drupifera, Rafin. l. c. 140 (1838).
Kemelia, Rafin. l. c. 138, 139 (1838).
Salceda, Blanco, Fl. Filip. ed. II. 374 (1854).
Sasanqua, Nees, in Flora, iv. (1834) Lit. 144.
Thea, Linn. Gen. ed. I. 154 (1737).
Theaphylla, Rafin. in Loud. Gard. Mag. vii. (1832) 246.
Tsia, Adans. Fam. ii. 450 (1763).
Tsubaki, Adans. l. c. 399 (1763).
assimilis, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 309.—Hongkong.
axillaris, Roxb. ex Ker-Gawl. in Bot. Reg. t. 349; et Sims, Bot. Mag. t. 2047 = Gordonia anomala.
axillaris, Wall. Cat. n. 158 = Adinandra acuminata.
Bohea, Lindl. in Loud. Encycl. Pl. 592; Griff. Notul. iv. 553 = C. Thea.
caudata, Griff. Notul. iv. 559 = drupifera.
caudata, *Wall. Cat.* n. 978.—Reg. Himal.
Chamgota, Buch.-Ham. ex Wall. in As. Res. xiii. (1820) 429 = C. Sasanqua.
drupifera, *Lour. Fl. Cochinch.* ii. 411.—Reg. Himal.; Burma; China.
Edithae, *Hance, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 221.—China.
euryoides, *Lindl. Bot. Reg.* t. 983.—China.
florida, Salisb. Prod. 370 = japonica.
fraterna, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 218.—China.
grandiflora, *Forsk. Fl. Aegypt. Arab.* 126.—Arab.
Grijsii, *Hance, in Journ. Bot.* xvii. (1879) 9.—China.
hongkongensis, *Seem. in Trans. Linn. Soc.* xxii. (1859) 342. t. 60.—China.
integrifolia, Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 150 = Actinodaphne chinensis.
japonica, Champ. in Hook. Kew Journ. iii. (1851) 309 = hongkongensis.
japonica, *Linn. Sp. Pl.* 698.—Japon.
japonica, Wall. Cat. n. 3667 = Adinandra integerrima.
Kaempferia, *Reboul, in Atti 3ª Reun. Sc. Ital.* (1841) 494.—Hab.?
Keina, Buch.-Ham. in D. Don, Prod. Fl. Nep. 224 = drupifera.
Kissi, Wall. in As. Res. xiii. (1820) 429 = drupifera.
lanceolata, *Seem. in Trans. Linn. Soc.* xxii. (1859) 345.—Malaya.
lutescens, *Dyer, in Hook. f. Fl. Brit. Ind.* i. 293.—Reg. Himal.
maliflora, Lindl. Bot. Reg. t. 1078 = rosaeflora.
Mastersia, Griff. Notul. iv. 559 = drupifera.
mutabilis, Paxt. Mag. Bot. ii. t. 122 = japonica.
oleifera, Abel, Journ. in China, 174 = C. Sasanqua.
oleifera, Wall. Cat. n. 976 = drupifera.
oleosa, Lour. Fl. Cochinch. 339 = C. Sasanqua.
planipetala, Lem. Illustr. Hortic. (1865) t. 426 = japonica.
Quinosaura, *Seem. in Trans. Linn. Soc.* xxii. (1859) 345.—Java.
reticulata, *Lindl. Bot. Reg.* t. 1078.—China.
rosaeflora, *Hook. Bot. Mag.* t. 5044.—China.
salicifolia, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 309.—Hongkong.
Sasanqua, Blanco, Fl. Filip. ed. I. 530 = drupifera.
Sasanqua, *Thunb. Fl. Jap.* 273. t. 30.—Japon.; China.
Scottiana, Choisy, Mém. Ternstr. 90 = Adinandra dumosa.
Scottiana, Wall. Cat. n. 3668 = C. Thea.
simplicifolia, Griff. Notul. iv. 560. t. 604. f. 2 = drupifera.

CAMELLIA:—
spectabilis, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 310 = reticulata.
symplocifolia, Griff. Itin. Notes, 40. n. 652 = drupifera.
Thea, *Link, Enum. Hort. Berol.* ii. 73.—China.
theifera, Griff. Notul. iv. 558. t. 601. f. 1, 3 = C. Thea.
theiformis, Hance, in Ann. Sc. Nat. Sér. IV. xv. (1861) 221 = euryoides.
Tsubakki, Crantz, Inst. ii. 172 = japonica.
viridis, Link, Enum. Hort. Berol. ii. 73 = C. Thea.

CAMERARIA, Dill. ex Moench, Meth. 520 (1794) = **Montia**, Linn. (Portulac.).
fontana, Moench, l. c. = M. fontana.

CAMERARIA, [Plum.] Linn. Syst. ed. I (1735). *APOCYNACEAE*, Benth. & Hook. f. ii. 701.
angustifolia, *Linn. Sp. Pl.* 210.—Am. austr.
dubia, Sims, Bot. Mag. t. 1646 = Wrightia dubia.
guianensis, Aubl. Pl. Gui. i. 262 = Malouetia Tamaquarina.
latifolia, *Linn. Sp. Pl.* 210.—Ind. occ.
lutea, Lam. Encyc. i. 573 = Malouetia Tamaquarina.
obesa, Spreng. Syst. i. 641 = Adenium obesum.
retusa, *Griseb. Fl. Brit. W. Ind.* 410.—Ind. occ.
Tamaquarina, Aubl. Pl. Gui. i. 256 = Malouetia Tamaquarina.
zeylanica, Retz. Obs. iv. 24 = Hunteria corymbosa.
zeylanica, Wight, ex Wall. Cat. n. 4459 = Strophanthus Wightianus.

CAMERIDIUM, Reichb. f. in Linnaea, xxii. (1849) 857, sphalm. = **Camaridium**, Lindl. (Orchid.).

CAMFOROSMA, Spreng. Syst. i. 453 (1825) = **Camphorosma**, Linn. (Chenop.).

CAMIRIUM, Gaertn. Fruct. ii. 194. t. 125 (1791) = **Aleurites**, Forst. (Euphorbiac.).
cordifolium, Gaertn. l. c. 195 = A. moluccana.
oleosum, Reinw. ex Blume, Cat. Gew. Buitenz. 104 = A. moluccana.

CAMISSONIA, Link, in Jahrb. Gew. i. (1818) 1. 186 = **Oenothera**, Linn. (Onagrar.).
flava, Link, l. c. = O. dentata.

CAMMARUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 327 = **Aconitum**, Tourn. (Ranuncul.).
paniculatum, Fourr. l. c. = A. paniculatum.

CAMOCLADIA, Linn. Syst. ed. X. 861 (1756) = **Comocladia**, P. Br. (Anacard.).

CAMOENSIA, Welw. ex Benth. & Hook. f. Gen. i. 557 (1865). *LEGUMINOSAE*, Benth. & Hook. f. l. c.
brevicalyx, *Benth. in Trans. Linn. Soc.* xxv. (1866) 302.—Afr. trop.
maxima, *Welw. ex Benth. l. c.* 302. t. 36.—Afr. trop.

CAMOMILLA, Gilib. Exercit. i. 178 (1792) = **Matricaria**, Tourn. (Compos.).
deflexa, Gilib. l. c. 179 = M. suaveolens.
inodora, Gilib. l. c. = M. inodora.
patens, Gilib. l. c. 178 = M. Chamomilla.

CAMONEA, Rafin. Fl. Tellur. iv. 81 (1836) = **Ipomoea**, Linn. (Convolvul.).
bifida, Rafin. l. c. = I. cymosa.

CAMPANEA, Decne. in Rev. Hortic. (1849) 241. t. 13. *GESNERACEAE*, Benth. & Hook. f. ii. 1003.
grandiflora, *Decne. l. c.*—N. Granat.
Humboldtii, *Hanst. in Linnaea*, xxvi. (1853) 214. t. 1. f. 24.—Costa Rica.
Oerstedii, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 108.—Costa Rica.

CAMPANULA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 46 (1737). *CAMPANULACEAE*, Benth. & Hook. f. ii. 561.
Apenula, Neck. Elem. i. 234 (1790).
Ballela, Rafin. Fl. Tellur. iv. 70 (1836).
Čeněkia, Opiz, Seznam, 27 (1852).
Decaprisma, Rafin. l. c. ii. 78 (1836).
Depierrea, Anon. ex Schlecht. in Linnaea, xvi. (1842) 374.

CAMPANULA:—
Drymocodon, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 111.
Erinia, Noulet, Fl. Sous-Pyr. 407.
Hecale, Rafin. Fl. Tellur. ii. 79 (1836).
Heterocodon, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1843) 255.
Lacara, Rafin. Fl. Tellur. iv. 70 (1836).
Loreia, Rafin. l. c. iii. 82 (1836).
Marianthemum, Schrank, in Denkschr. Regensb. Ges. ii. (1822) 34.
Medium, [Tourn.] Fisch. ex A. DC. Monog. Camp. 213 (1830).
Nenningia, Opiz, Seznam, 68 (1852).
Pentropis, Rafin. Fl. Tellur. ii. 80 (1836).
Quinquelocularia, C. Koch, in Linnaea, xxiii. (1850) 630.
Rapunculus, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 111.
Roncelia, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 289 (1870).
Roucelia, Dum. Comm. Bot. 14 (1822).
Sykoraea, Opiz, Seznam, 94 (1852).
Syncodon, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 111.
Talanelis, Rafin. Fl. Tellur. iv. 70 (1836).
Tracheliopsis, Opiz, Seznam, 98 (1852).
Weitenwebera, Opiz, l. c. 104 (1852).
abietina, *Griseb. & Schenk, in Wiegm. Archiv*, xv. (1852) 1. 333.—Europ.
acuminata, Michx. Fl. Bor. Am. i. 108 = americana.
acutangula, *Leresche & Levier, in Journ. Bot.* xvii. (1879) 198.—Hispan.
acutiloba, *Vatke, in Linnaea*, xxxviii. (1874) 709.—Kurdistan.
Adami, Bieb. Fl. Taur. Cauc. i. 153 = bellidifolia.
Adami, Willd. ex Steud. Nom. ed. II. i. 265 = lasiocarpa.
adhaerens, Ehrenb. ex Boiss. Fl. Orient. iii. 943 = retrorsa.
adpressa, Linn. f. Suppl. 140 = Wahlenbergia adpressa.
adscendens, *Vest. ex Roem. & Schult. Syst.* v. 90.—Sibir.
adsurgens, *Leresche & Levier, in Journ. Bot.* xvii. (1879) 199.—Hispan.
affinis, Fisch. ex Rupr. in Bull. Acad. Pétersb. xi. (1867) 205 = Aucheri.
affinis, Reichb. ex Nym. Consp. 480 = Portenschlagiana.
affinis, *Roem. & Schult. Syst.* v. 140.—Hispan.
afganica, *Pomel, Nouv. Mat. Fl. Atl.* 257.—Afr. bor.
afra, Cav. in Anal. Cienc. Nat. iii. (1801) 21 = dichotoma.
aggregata, Willd. Enum. Hort. Berol. Suppl. 10 = glomerata.
agrestis, Wall. Cat. n. 1292 = Wahlenbergia gracilis.
Aizoon, *Boiss. & Sprun. Diagn.* Ser. I. iv. 34.—Graecia.
ajugaefolia, *Sestini, ex Spreng. Syst.* i. 732 (sp. dub.).—As. Min.
alata, *Desf. Fl. Atlant.* i. 178. t. 50.—Afr. bor.
Alberti, *Trautv. in Act. Hort. Petrop.* vi. (1879) 83.—Turkestan.
albiflora, C. Koch, in Linnaea, xxiii. (1850) 634 = sarmatica.
alburnica, Brigant. Pempt. i. 3. t. 2 = Phyteuma amplexicaule.
algida, Fisch. ex A. DC. Monog. Camp. 338 = lasiocarpa.
alliariaefolia, Reichb. Pl. Crit. Cent. iii. 345 = Symphyandra pendula.
alliariaefolia, *Willd. Sp. Pl.* i. 910.—Reg. Caucas.; As. Min.
Allionii, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa.
Allionii, *Vill. Fl. Delph.* 18; *Prosp.* 22.—Europ.
allophylla, Rafin. ex A. DC. Monog. Camp. 282 = rotundifolia.
alopecuroides, Willd. ex Steud. Nom. ed. II. i. 265 = Phyteuma Sibthorpianum.
alpestris, All. Fl. Pedem. 113. t. 6. f. 3 = Allionii.
Alphonsii, *Wall. Cat.* n. 1296.—Ind. or.
alpigena, C. Koch, in Linnaea, xxii. (1849) 638 = Aucheri.
alpina, *Jacq. Enum. Vind.* 36; 210.—Europ.
Alpini, Linn. Sp. Pl. ed. II. 1669 = Adenophora lilifolia.
Alpini, Pichl. ex Vis. Fl. Dalm. Suppl. 74 = Phyteuma amplexicaule.
alsinoides, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 24.—Reg. Himal.
altaica, A. DC. Monog. Camp. 229. t. 10 = pilosa.

CAMPANULA :—
altaica, Ledeb. Ind. Sem. Hort. Dorpat. (1824) 2 = Steveni.
altiflora, Pers. Syn. i. 192 = Prismatocarpus altiflorus.
americana, Hort. ex Steud. Nom. ed. II. i. 266 = Phyteuma canescens.
americana, *Linn. Sp. Pl.* 164.—Am. bor.
americana, Mill. Gard. Dict. ed. VIII. n. 13 = planiflora.
amplexicaulis, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma amplexicaule.
amplexicaulis, Michx. Fl. Am. Bor. i. 108 = Specularia perfoliata.
amygdalifolia, Salisb. Prod. 126 = persicaefolia.
anagalloides, Royle, Illustr. Bot. Himal. 253 = Cephalostigma hirsutum.
anchusiflora, *Sibth. & Sm. Fl. Graec. Prod.* i. 141.—Archip. Graec.
andina, Rupr. in Bull. Acad. Pétersb. xi. (1867) 215 = gumbetica.
Andrewsii, A. DC. Monog. Camp. 220 = tomentosa.
androsacea, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia androsacea.
angulata, Rafin. Fl. Ludov. 55 = Specularia perfoliata.
angustifolia, Lam. Tabl. Encyc. ii. 53 = linifolia.
anticensis, Kunze, ex A. DC. in DC. Prod. vii. 491 = Specularia perfoliata.
antirrhina, Schleich. Cat. Pl. Helv. ed. III. 10 = caespitosa, rotundifolia.
aparinoides, *Pursh, Fl. Am. Sept.* i. 159.—Am. bor.
appendiculata, A. DC. Prod. vii. 462 = tomentosa.
arborescens, P. Br. (err. R. Br.) ex Steud. Nom. ed. II. i. 266 = Hamelia ventricosa.
arcuata, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 138 = rotundifolia.
ardonensis, *Rupr. in Bull. Acad. Pétersb.* xi. (1867) 212.—Reg. Caucas.
argaea, *Boiss. & Bal. Diagn.* Ser. II. vi. 119.—Cappadoc.
argentea, *Lam. Encyc.* i. 584.—Armenia.
argunensis, Rupr. in Bull. Acad. Pétersb. xi. (1867) 209 = Aucheri.
argyrotricha, *Wall. Cat.* n. 7138.—Reg. Himal.
arida, H. B. & K. Nov. Gen. et Sp. iii. 301 = Wahlenbergia linarioides.
aristata, *Wall. Cat.* n. 1291; *et in Roxb. Fl. Ind. ed. Carey*, ii. 98.—Reg. Himal.
armena, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 236 = Symphyandra armena.
arvatica, Lag. in Varied. Cienc. ii. IV. (1805) 40 = Wahlenbergia hederacea.
arvensis, Pers. Syn. i. 193 = Specularia Speculum.
aspera, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma asperum.
aspera, Donn, ex A. DC. Monog. Camp. 290 = aparinoides.
aspera, *Moench, Meth. Suppl.* 188.—Hab. ?
asperrima, Zuccagni, in Roem. Collect. 128 = ajugaefolia.
asteroides, Lam. Tabl. Encyc. ii. 55 = americana.
athoa, Boiss. & Heldr. Diagn. Ser. II. iii. 110 = C. Trachelium.
attenuata, Ledeb. ex Spreng. Syst. i. 729 = persicifolia.
attica, Boiss. & Heldr. Diagn. Ser. I. xi. 67 = drabaefolia.
Aucheri, *A. DC. in DC. Prod.* vii. 460.—Reg. Caucas.; As. Min.
aurea, Linn. f. Suppl. 141 = Musschia aurea.
axillaris, *Boiss. & Bal. Diagn.* Ser. II. iii. 109.—Cilicia.
azurea, Soland. & Banks, ex Sims, Bot. Mag. t. 551 = rhomboidea.
baicalensis, Pall. ex A. DC. Prod. vii. 479 = silenifolia.
Balansae, *Boiss. & Haussk. Fl. Orient.* iii. 931.—Cappadoc.
baldensis, Balb. Cat. Hort. Taur. (1813) 20 = ramosissima.
Banksiana, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia Banksiana.
barbata, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa.
barbata, *Linn. Syst.* ed. X. 927.—Europ.
barbata, Spreng. Fl. Halens. 77 = glomerata.
Barrelieri, *Marnock, Floric. Mag.* i. (1836) 161. t. 12.—Italia.
Barrelieri, Presl, Symb. Bot. 30. t. 19 = fragilis.
Baumgartenii, Becker, Fl. Francof. i. 264 = rotundifolia.

CAMPANULA :—
Bayerniana, *Rupr. in Bull. Acad. Pétersb.* xi. (1867) 214.—Reg. Caucas.
Bellardi, All. Fl. Pedem. i. 108 = caespitosa.
bellidifolia, *Adam, in Web. & Mohr, Beitr.* i. (1805) 47.—Reg. Caucas.
bellidifolia, Frivald. ex Nym. Consp. 476 = Orphanidea.
bellidifolia, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 36 = patula.
Benthamii, Wall. Cat. n. 1290 = canescens.
Bergiana, D. Dietr. Syn. Pl. i. 757 = Prismatocarpus subulatus.
betonicaefolia, Biehl. Pl. Nov. Hort. Spreng. 14 = sarmatica.
betonicaefolia, Gilib. Fl. Lituan. i. 53 = glomerata.
betonicaefolia, *Sibth. & Sm. Fl. Graec. Prod.* i. 141.—As. Min.
betulaefolia, *C. Koch, in Linnaea*, xxiii. (1850) 635.—Armenia.
bicaulis, Lapeyr. Hist. Abr. Pl. Pyr. 107. t. 7 = speciosa.
bicolor, Boiss. Fl. Orient. iii. 931 = camptoclada.
Biebersteiniana, C. A. Mey. Verz. Pfl. Cauc. 83, pro parte = Aucheri.
Biebersteiniana, Roem. & Schult. Syst. v. 147 = tridentata.
Bielzii, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = rotundifolia.
biflora, Ruiz & Pav. Fl. Per. ii. 55 = Specularia biflora.
Billardieri, A. DC. Monog. Camp. 303. t. 13 = C. Cymbalaria.
biserrata, C. Koch, in Linnaea, xix. (1847) 29 = lactiflora.
bithynica, A. DC. in DC. Prod. vii. 460 = tridentata.
Bocconi, J. F. Gmel. Syst. 348 = rhomboidea.
Bocconi, Vill. Hist. Pl. Dauph. ii. 502 = pusilla.
Boissieri, Sprun. ex Boiss. Fl. Orient. iii. 915 = radicosa.
Boissieri, Vatke, in Linnaea, xxxviii. (1874) 703 = Phyteuma asperum.
Bolosii, *Vayr. in Anal. Soc. Esp. Hist. Nat.* viii. (1879) 451.—Hispan.
bononiensis, *Linn. Sp. Pl.* 165.—Europ. austr.; Reg. Caucas.; Sibir.
brachiata, Salzm. ex A. DC. Prod. vii. 462 = dichotoma.
brachiata, Seidl, ex Opiz, Boeheims Gew. 168; Natural. ii. (1823) [6] = patula.
brachysepala, Vukot. in Linnaea, xxvi. (1853–55) 325 = cenisia.
bracteata, Thunb. in Hoffm. Phytogr. Blaett. i. (1803) 20 = Roella bracteata.
brevibracteata, D. Dietr. Syn. Pl. i. 757 = Microcodon glomeratum.
Broussonetiana, Roem. & Schult. Syst. v. 104 = Loeflingii.
Brunonis, Wall. Cat. n. 1285 = latifolia.
Buchii, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia pilosa.
caespitosa, *Scop. Fl. Carn.* ed. II. i. 143. t. 4.—Europ.
calamenthifolia, Aucher, ex A. DC. Prod. vii. 462 = tomentosa.
calamenthifolia, *Lam. Encyc.* i. 585.—Ins. Naxos.
calaminthifolia, Griseb. Spicil. Fl. Rumel. i. 286 = orphanidea.
calycantha, Turcz. ex Herd. in Act. Hort. Petrop. i. (1873) 288, 296 = pilosa, silenifolia.
calycina, Boeber, ex Roem. & Schult. Syst. v. 104 = C. Rapunculus.
camptoclada, *Boiss. Diagn.* Ser. I. xi. 63.—Syria.
camtschatica, *Pall. ex Roem. & Schult. Syst.* v. 152.—Kamtschat.
cana, *Wall. Cat.* n. 1284; *et in Roxb. Fl. Ind. ed. Carey*, ii. 101.—Reg. Himal.
canaria, *Schrank, ex Steud. Nom.* ed. II. i. 266.—Ins. Canar.
canariensis, Linn. Sp. Pl. 168 = Canarina campanulata.
candida, *A. DC. Monog. Camp.* 234.—Kurdistan.
Candolleana, D. Dietr. Syn. Pl. i. 756 = Prismatocarpus Candolleanus.
canescens, Roth, Enum. Pl. Phan. Germ. i. 716 = Phyteuma canescens.
canescens, *Wall. Cat.* n. 1289.—Ind. or.; Afghan.
capensis, Linn. Sp. Pl. 169 = Wahlenbergia capensis.
caperonoides, Klotzsch, in Bot. Ergeb. Waldem. Reise, 73 = sylvatica.

CAMPANULA :—
capillacea, Thunb. Prod. Pl. Cap. 38 = Wahlenbergia capillacea.
capillaris, Lodd. Bot. Cab. t. 1406 = Wahlenbergia gracilis.
capitata, Schur, in Verh. Siebenb. Ver. Naturw. iii. (1852) 88 = C. Cervicaria.
capitata, Sims, Bot. Mag. t. 811 = lingulata.
cappadocica, Boiss. Diagn. Ser. I. xi. 73 = Phyteuma cappadocica ?
cariensis, A. DC. in DC. Prod. vii. 459 = betonicaefolia.
carnica, *Schiede, ex Mert. & Koch, Deutschl. Fl.* ii. 158.—Carniola.
carnosa, Wall. Cat. n. 1282; et in Roxb. Fl. Ind. ed. Carey, ii. 102 = Peracarpa carnosa.
carpatica, *Jacq. Hort. Vind.* i. 22. t. 57.—Hungar.
cashmiriana, *Royle, Illustr. Bot. Himal.* i. 254. t. 62.—Reg. Himal.
caucasica, Bieb. Casp. 140; App. n. 14 = sibirica ?
caudata, *Vis. Fl. Dalm.* ii. 136. t. 33.—Dalmat.
Cavolini, Tenore, Fl. Nap. t. 213 = fragilis.
celebica, D. Dietr. Syn. Pl. i. 758 = Campanumoca celebica.
Celsii, A. DC. Monog. Camp. 217 = tomentosa.
celtidifolia, Boiss. Diagn. Ser. II. iii. 111 = lactiflora.
cenisia, *Linn. Sp. Pl.* ed. II. Snppl. 1669.—Europ.
cephalantha, Fisch. ex A. DC. Monog. Camp. 254 = glomerata.
Cephalaria, *Vukot. in Linnaea*, xxvi. (1853–55) 333.—Hungar.
Cephalotes, Fisch. ex A. DC. Monog. Camp. 254 = glomerata.
cernua, Hort. ex Hornem. Hort. Hafn. 201 = sibirica.
cernua, Thunb. Prod. Pl. Cap. 39 = Wahlenbergia cernua.
cerviana, Pall. Reise, iii. 687 = C. Cervicaria.
Cervicaria, *Linn. Sp. Pl.* 167.—Europ.
cervicarioides, Roem. & Schult. Syst. v. 128 = glomerata.
Cervicina, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia Cervicina.
chilensis, Bert. ex Steud. Nom. ed. II. i. 266 = Wahlenbergia hirsuta, reflexa.
chilensis, Molina, Sagg. Chile, ed. II. 281 = Wahlenbergia linarioides.
chondrophylla, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia chondrophylla.
cichoracea, Sibth. & Sm. Fl. Graec. t. 209 = lingulata.
cichoriformis, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma cichoriforme.
ciliaris, Salisb. Prod. 127 = Roella ciliata.
ciliata, Patr. ex A. DC. Prod. vii. 479 = silenifolia.
ciliata, *Stev. in Mém. Soc. Nat. Mosc.* iii. (1812) 256.—Reg. Caucas.
ciliata, Thunb. Fl. Cap. ii. 24 = Lightfootia tenella.
cilicia, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma amplexicaule.
cinerea, Linn. f. Suppl. 139 = Lightfootia grisea.
circaeoides, *F. Schmidt, Reisen Amurl.* 154.—Japon.
cissophylla, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. Dec.* 1, 2.—Kurdistan.
clisophylla, *Moretti, ex Reichb. Ic. Fl. Germ.* xix. t. 1611.—Hab. ?
clivosa, Banks, ex A. DC. Monog. Camp. 161 = Wahlenbergia clivosa.
clivosa, Herb. Roxb. ex A. DC. Prod. vii. 486 = Wahlenbergia linifolia.
coa, D. Dietr. Syn. Pl. i. 758 = Specularia pentagonia.
coarctata, Gilib. Fl. Lituan. i. 48 = C. Rapunculus.
coccinea, *Sieber, ex A. DC. Prod.* vii. 486 (quid ?).—Ins. Maurit.
cochlearifolia, Lam. Encyc. i. 578 = caespitosa.
cochlearifolia, Vahl, Symb. Bot. 18 = fragilis.
collina, *Bieb. Fl. Taur. Cauc.* i. 152.—Reg. Caucas.
coloradoënse, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 460 = Specularia Lindheimeri.
colorata, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 101.—Ind. or.; Afghan.
commutata, Roem. & Schult. Syst. v. 143 = sarmatica.
compacta, *Boiss. & Heldr. Diagn.* Ser. I. xi. 71.—Pisidia.
conferta, *A. DC. Prod.* vii. 468.—Cappadoc.
confertifolia, Kent. ex Nym. Consp. 479 = rotundifolia.
congesta, Roem. & Schult. Syst. v. 125 = glomerata.
conglomerata, Gueldenst. ex Ledeb. Fl. Ross. ii. 880 = glomerata.

CAMPANULA:—
consanguinea, *Schott, Nym. & Kotschy, Analect. Bot.* 8.—Europ.
contracta, Mutis, ex Nym. Consp. 478 = rapunculoides.
controversa, Boiss. Diagn. Ser. II. iii. 115 = Phyteuma rigidum.
cordata, *Peterm. Fl. Lips. Excurs.* 187.—Saxonia.
cordata, Tausch, in Flora, xxv. (1842) i. 288 = Adenophora cordata.
cordata, Vis. Stirp. Dalm. 5 = Specularia Speculum.
cordifolia, C. Koch, in Linnaea, xix. (1847) 29 = rapunculoides.
cordifolia, *Vukot. in Linnaea*, xxvi. (1853) 328.—Europ.
coriacea, Boiss. & Kotschy, ex Fl. Orient. iii. 910 = C. Radula.
coronata, Ker-Gawl. in Bot. Reg. t. 149 = Adenophora marsupiiflora.
coronopifolia, Fisch. ex Roem. & Schult. Syst. v. 157 = Adenophora coronopifolia.
corymbifera, Poir. Encyc. Suppl. ii. 64 = seq.
corymbosa, Desf. in Ann. Mus. Par. xi. (1808) 139 = pelviformis.
corymbosa, Tenore, Prod. Fl. Nap. 15 = versicolor.
Costae, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 294 = patula.
crassifolia, Nees, in Syll. Ratisb. i. (1824) 6 = fragilis.
crassipes, *Heuff. in Oestr. Bot. Zeitschr.* viii. (1858) 27.—Europ.
crenata, Link, Enum. Hort. Berol. i. 214 = rapunculoides.
cretica, *D. Dietr. Syn. Pl.* i. 758.—Ins. Creta.
crispa, Banks, ex A. DC. Prod. vii. 444 = Prismatocarpus crispus.
crispa, *Lam. Encyc.* i. 581.—Armenia.
Cymbalaria, *Sibth. & Sm. Fl. Graec. Prod.* i. 139.—As. Min.; Syria.
dalmatica, D. Dietr. Syn. i. 752 = Wahlenbergia dalmatica.
dalmatica, Tausch, in Flora, x. (1827) 246 = C. Cervicaria.
damascena, *Labill. Ic. Pl. Syr. Dec.* v. 7. t. 5.—Syria.
dasyantha, Bieb. Fl. Taur. Cauc. iii. 147 = pilosa.
dasycarpa, Kit. ex Schult. Oestr. Fl. ed. II. 404 = persicifolia.
dasycarpa, Schur, Enum. Pl. Transs. 440 = carpatica.
davurica, *Siev. in Pall. N. Nord. Beitr.* vii. 218.—Sibir.
debilis, D. Dietr. Syn. Pl. i. 752 = Wahlenbergia debilis.
decipiens, Roem. & Schult. Syst. v. 142 = dichotoma.
declinata, Moench, Meth. Suppl. 187 = americana.
Decloëtiana, Ortm. in Weitenw. Beitr. iii. (1838) 220 = rotundifolia.
decumbens, A. DC. Monog. Camp. 334. t. 11 = Loeflingii.
decurrens, Linn. Sp. Pl. 164 = patula, persicaefolia.
decurrens, Thore, Ess. Chlor. 64 = C. Rapunculus.
decurrens, Zuccagni, in Roem. Collect. 128 = peregrina.
dehiscens, Roxb. Hort. Beng. 85; Fl. Ind. i. 505 = Wahlenbergia gracilis.
delicatula, *Boiss. Diagn.* Ser. I. xi. 67.—As. Min.
delicatula, Jord. ex Nym. Consp. 479 = caespitosa.
denticulata, Boiss. & Huet, Diagn. Ser. II. iii. 107 = betulaefolia.
denticulata, Burch. Trav. i. 538 = Lightfootia capillaris.
denticulata, Spreng. Syst. i. 735 = Adenophora tricuspidata.
denudata, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia denudata.
desertorum, *Weinm. in Bull. Soc. Nat. Mosc.* (1837) n. VII. 58.—Ruthenia.
dichotoma, *Linn. Cent. Pl.* ii. 10; (*Amoen. Acad.* iv. 306).—Reg. Mediterr. occ.
diffusa, Banks, ex A. DC. Prod. vii. 436 = Wahlenbergia nudicaulis.
diffusa, D. Dietr. Syn. i. 752, 756 = Microcodon lineare.
diffusa, Vahl, Symb. Bot. i. 48 = fragilis.
dilecta, Schott, Nym. & Kotschy, Analect. Bot. 8 = rotundifolia.
dimorphantha, *Schweinf. Beitr. Fl. Aethiop.* 140.—Aegypt.
dissoluta, Schott & Kotschy, ex Boiss. Fl. Orient. iii. 943 = retrorsa.

CAMPANULA:—
divaricata, *Michx. Fl. Bor. Am.* i. 109.—Am. bor
divergens, Willd. Enum. Hort. Berol. i. 212 = sibirica.
diversifolia, D. Dietr. Syn. Pl. i. 752 = Wahlenbergia diversifolia.
diversifolia, Dum. Fl. Belg. 58 = rotundifolia.
Draba, *Burm. f. Fl. Cap. Prod.* 5.—Afr. austr.
drabaefolia, Frivald. ex Boiss. Fl. Orient. iii. 932 = scutellata.
drabaefolia, *Sibth. & Sm. Fl. Graec.* iii. 11. t. 215.—Graecia.
drabaefolia, Steud. Nom. ed. II. i. 267 = fastigiata.
dracunculifolia, Boiss. & Kotschy, Fl. Orient. iii. 952 = Phyteuma rigidum.
dubia, A. DC. Monog. Camp. 286 = rotundifolia.
dulcis, *Decne. in Ann. Sc. Nat.* Sér. II. ii. (1834) 258.—Arab.
dumetorum, Boiss. Fl. Orient. iii. 922 = rapunculoides.
Dunantii, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia Dunantii.
Duriaei, Boiss. Voy. Bot. Espagne, 401 = Loeflingii.
echiifolia, Gilib. Fl. Lituan. i. 56 = C. Cervicaria.
Ecklonii, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia Ecklonii.
edulis, *Forsk. Fl. Aegypt. Arab.* 106.—Arab.
elatina, Hill, Veg. Syst. viii. 13 = C. Elatines.
Elatines, Bout. ex Willk. & Lange, Prod. Fl. Hisp. ii. 295 = specularioides.
Elatines, *Linn. Syst.* ed. X. 927.—Europ. austr.
Elatines, Petag. Inst. Bot. ii. 367 = garganica.
Elatines, Pollini, Fl. Veron. i. 278 = elatinoides.
elatinoides, *Moretti, in Brugnat. Giorn. Fis.* II. v. (1822) 110.—Italia.
elatior, Link & Hoffmgg. Fl. Port. ii. 11. t. 80 = C. Rapunculus.
elegans, *Roem. & Schult. Syst.* v. 105.—Sibir.
elliptica, Kit. in Schult. Oestr. Fl. ed. II. 410 = glomerata.
elongata, Banks, ex A. DC. Prod. vii. 436 = Wahlenbergia nudicaulis.
elongata, Portensch. ex A. DC. Monog. Camp. 277 = rapunculoides.
elongata, Willd. Enum. Hort. Berol. Suppl. 10 = Wahlenbergia capensis.
ensifolia, Lam. Encyc. i. 582 = Heterochaenia ensifolia.
ephesia, *Boiss. Fl. Orient.* iii. 898.—Lydia.
eriantha, Hampe, in Flora, xxv. (1842) i. 76 = tomentosa.
ericoides, Lam. Tabl. Encyc. ii. 63 = Prismatocarpus paniculatus.
erinoides, Cav. in Anal. Cienc. Nat. iii. (1800) 20 = Loeflingii.
erinoides, *Linn. Mant.* i. 44.—Hisp.; Afr. bor.
erinoides, Muhl. Cat. n. 22 = aparinoides.
Erinus, *Linn. Sp. Pl.* 169.—Reg. Mediterr.
eriocarpa, Bieb. Fl. Taur. Cauc. i. 149 = latifolia.
erysimoides, Vest, ex Roem. & Schult. Syst. v. 102 = Adenophora Gmelini.
esculenta, *A. Rich. Tent. Fl. Abyss.* ii. 4.—Abyssin.
esculenta, Salisb. Prod. 126 = C. Rapunculus.
euclasta, *Boiss. Diagn.* Ser. I. xi. 70.—Syria.
evolvulacea, Royle, Illustr. Bot. Himal. i. 254 = cachemiriana.
excisa, *Schleich. ex Murith, Guide Val.* 33, 35.—Helvet.
exilis, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia exilis.
expansa, *Frivald. in Flora*, xix. (1836) ii. 434.—Thessalia; Macedon.
expansa, Rudolph, in Mém. Acad. Pétersb. iv. (1809) 340. t. 2. f. 1 = Wahlenbergia homallanthina.
Exul, *Schott, Nym. & Kotschy, Analect. Bot.* 9.—Austria.
falcata, Roem. & Schult. Syst. v. 154 = Specularia falcata.
fallax, Rupr. in Bull. Acad. Pétersb. xi. (1867) 205, in textu = Aucheri.
farinosa, Andrz. ex Bess. Enum. Pl. Volh. 10 = glomerata.
fasciculata, Linn. f. Suppl. 139 = Lightfootia fasciculata.
fastigiata, *Dufour, ex A. DC. Monog. Camp.* 340. t. 12.—Hispan.; Afr. bor.; Reg. Caucas.
fastigiata, S. G. Gmel. Iter, i. 153. t. 33 = C. Rapunculus.
Fedtschenkiana, *Trautv. in Act. Hort. Petrop.* vi. (1879) 77.—As. temp.

CAMPANULA:—
fernandeziana, D. Dietr. Syn. Pl. i. 744 = Wahlenbergia fernandeziana.
ficarioides, Timb. in Mém. Acad. Toulouse, Sér. V. vi. (1862) 33 = linifolia.
filicaulis, *Dur. Expl. Algér.* t. 62.—Algeria.
filiflora, Kellogg, in Proc. Calif. Acad. ii. (1863) 5 = prenanthoides.
filiformis, Gilib. Fl. Lituan. i. 53 = rotundifolia.
filiformis, Moretti, in Brugnat. Giorn. Fis. II. ix. (1826) 155 = Morettiana.
filiformis, Ruiz & Pav. Fl. Per. ii. 55 = Wahlenbergia linarioides.
firma, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma Aucheri.
Firmiana, Vand. Fasc. Pl. 9 = barbata.
Fischeri, Roem. & Schult. Syst. v. 116 = Adenophora communis.
Fischeriana, Spreng. Syst. iv. Cur. Post. 77 = Adenophora coronopifolia.
flaccida, D. Dietr. Syn. Pl. i. 752 = Wahlenbergia flaccida.
flaccidula, *Vatke, in Linnaea*, xxxviii. (1874) 712.—Mesopot.
flagellaris, H. B. & K. Nov. Gen. et Sp. iii. t. 265 = Specularia perfoliata.
flexuosa, Michx. Fl. Bor. Am. i. 109 = divaricata.
flexuosa, Waldst. & Kit. Pl. Rar. Hung. ii. 145 = Waldsteiniana.
floribunda, Viv. Fl. Lib. Spec. 67 = isophylla.
florida, Salisb. Prod. 127 = C. Medium.
floridana, *S. Wats. ex A. Gray, Syn. Fl. N. Am.* ii. i. 13.—Florida.
foliosa, *Tenore, Fl. Nap.* i. 71. t. 18.—Italia.
Fontanesiana, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma lanceolatum.
Foudrasi, Jord. Pugill. Pl. Nov. (1852) 124 = caespitosa.
fragilis, *Cyrill. Pl. Fasc.* i. 32. t. 11. f. 2.—Italia.
frigida, Pall. ex A. DC. Prod. vii. 461 = pilosa.
Frivaldskyi, Steud. Nom. ed. II. i. 567 = expansa.
fruticosa, Banks, ex A. DC. Prod. vii. 444 = Prismatocarpus fruticosus.
fruticosa, Hill, Syst. Veg. viii. t. 8 = Prismatocarpus paniculatus.
fruticosa, Linn. Sp. Pl. 168 = Lightfootia subulata.
fulgens, *Wall. Cat.* n. 1283; *et in Roxb. Fl. Ind. ed. Carey*, ii. 99.—Ind. or.
garganica, *Tenore, Cat. Sem. Hort. Nap.* (1827) 3; *ex Fl. Nap.* iii. 203.—Italia; Graecia.
Gautieri, Jeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul. iii. (1875-76) 621 = Scheuchzeri.
gentianoides, Lam. Encyc. i. 181 = Platycodon grandiflorum.
ghilanensis, Pall. ex Roem. & Schult. Syst. v. 154 = Specularia ghilanensis.
ghilanensis, *Schlecht. ex Boiss. Fl. Orient.* iii. 943.—Persia.
Gieseckiana, Vest, ex Roem. & Schult. Syst. v. 89 = uniflora.
gilanica, Rupr. ex Boiss. Fl. Orient. iii. 905 = Ruprechtii.
Gisleri ×, *Bruegg, in Jahresb. Naturf. Ges. Graub.* II. xxiii-xxiv. (1880) 104.—Europ.
glabrata, Banks, ex A. DC. Prod. vii. 435 = Wahlenbergia undulata.
glabricarpa, Schleich. Cat. Pl. Helv. ed. IV. 10 = rapunculoides.
glacialis, *Shuttl. in Mag. Zool. & Bot.* ii. (1838) 188.—Helvetia.
glandulosa, Banks, ex A. DC. Monog. Camp. 326 = C. Rapunculus.
glareosa, Huet, ex Nym. Consp. 480 = caespitosa.
glauca, *Thunb. Fl. Jap.* 88.—Japon.
glaucophylla, Schloss. & Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 111 = glomerata.
glomerata, *Linn. Sp. Pl.* 166.—Europ.; As. temp.
Gmelini, Roem. & Schult. Syst. v. 103 = Adenophora marsupiiflora.
Gmelini, Spreng. Fl. Hal. Mant. 36 = Adenophora Gmelini.
gracilis, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma lobelioides.
gracilis, Bert. ex Colla, in Mem. Acc. Tor. xxxviii. (1835) 119 = Wahlenbergia Berteroi.
gracilis, Forst. f. Prod. 15 = Wahlenbergia gracilis.
gracilis, Jord. Pugill. Pl. Nov. (1852) 122 = caespitosa.
graminifolia, Host, Fl. Austr. i. 268 = Wahlenbergia tenuifolia.

CAMPANULA:—
graminifolia, Linn. Sp. Pl. 166 = Wahlenbergia graminifolia.
graminifolia, Waldst. & Kit. Pl. Rar. Hung. ii. 166 = Wahlenbergia Kitaibelii.
graminifolia, Willerm. ex A. DC. Monog. Camp. 254 = glomerata.
grammosepala, Vukot. in Linnaea, xxvi. (1853–55) 323 = spp. plurimi.
grandiflora, Jacq. Hort. Vindob. iii. 4 = Platycodon grandiflorus.
grandiflora, Lam. Fl. Fr. iii. 334 = C. Medium.
grandiflora, Pourr. ex Nym. Consp. 476 = speciosa.
grandis, Fisch. & Mey. Ind. Sem. Hort. Petrop. v. 34 = latiloba.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 22.—Afghan.
Grossekii, *Heuff. in Flora*, xvi. (1833) i. 353.—Hungar.
gumbetica, *Boiss. Fl. Orient.* iii. 914.—Reg. Caucas.
gummifera, Willd. ex Roem. & Schult. Syst. v. 144 = sarmatica.
Gundelia, C. Koch, in Linnaea, xxiii. (1850) 634 = sibirica.
hagiclia, *Boiss. Fl. Orient.* iii. 899.—Ins. Rhodos.
hastaefolia, Salisb. Prod. 127 = Canarina campanulata.
Hauryi, Schott, Nym. & Kotschy, Analect. Bot. 10 = rotundifolia.
Hausmanni ×, Reichb. f. in Flora, lx. (1877) 31.
haylodgensis, *Hort.—Cf. Gard. Chron.* (1885) ii. 147.—Hab. ?
hederacea, Linn. Sp. Pl. 169 = Wahlenbergia hederacea.
hederaefolia, Salisb. Prod. 127 = Wahlenbergia hederacea.
hemschinica, *C. Koch, in Linnaea*, xxiii. (1850) 644.—As. Min.
Hendersoni, *Hort.—Cf. Gard. Chron.* (1882) ii. 502.—Hab. ?
Herminii, Bory & Chaub. ex Boiss. Fl. Orient. iii. 937 = Spruneri.
Herminii, *Hoffmgg. & Link, Fl. Port.* ii. 9. t. 79.—Lusitan.
hesperidifolia, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma lobelioides.
heterodoxa, Bong. Veg. Sitcha, 144 = rotundifolia.
heterodoxa, Vest, in Roem. & Schult. Syst. v. 98 = linifolia.
heterophylla, Baumg. Fl. Transs. Suppl. iii. 342 = Wanneri.
heterophylla, S. F. Gray, Nat. Arr. Brit. Pl. ii. 408 = rotundifolia.
heterophylla, Linn. Sp. Pl. ed. I. errata, et 169 (sine nom. triv.).—Archip. Graec.
hierosolymitana, *Boiss. Diagn.* Ser. I. xi. 62.—Palaestin.
Hilsenbergi, Boj. ex A. DC. Prod. vii. 429 = Wahlenbergia Hilsenbergii.
himalayensis, Klotzsch, in Bot. Ergeb. Waldem. Reise, 73 = colorata.
hirsuta, Guss. Syn. Fl. Sic. i. 250, in syn. = Specularia Speculum.
hirsuta, *Pantoč. in Verh. Ver. Natur- u. Heilk. Presb.* ii. (1871–72) 54.—Europ.
hirsutissima, Guss. ex A. DC. Monog. Camp. 316 = peregrina.
hirta, Roem. & Schult. Syst. v. 153 = Specularia Speculum.
hispanica, *Willk. in Willk. & Lange, Prod. Fl. Hisp.* ii. 291.—Hispan.
hispida, Eckl. ex A. DC. Prod. vii. 421 = Microcodon glomeratum.
hispida, Fisch. ex Hornem. Hort. Hafn. 199 = lactiflora.
hispida, Lejeune, Fl. Spa, Suppl. ii. 298 = persicifolia.
hispida, Portensch. ex A. DC. Monog. Camp. 304 = Morettiana.
hispida, Spreng. ex Steud. Nom. ed. II. i. 268 = ciliata.
hispidissima, Hochst. in Lorent, Wanderungen, 331 = propinqua.
hispidula, Eckl. ex A. DC. Monog. Camp. 127 = Microcodon glomeratum.
hispidula, Link, Enum. Hort. Berol. i. 215 = Microcodon sparsiflorum.
hispidula, Linn. f. Suppl. 142 = Microcodon depressum.
Hochstetteri, Schott, Nym. & Kotschy, Analect. Bot. 12 = caespitosa.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 74 = colorata.

CAMPANULA:—
Hohenackeri, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. Suppl. 9 = sibirica.
homallanthina, Ledeb. in Mém. Acad. Pétersb. iv. (1811) 340 = Wahlenbergia homallanthina.
Hornungiana, *Schur, Enum. Pl. Transs.* 442.—Transsylv.
hortensis, Meerb. Pl. Sel. 10 = rapunculoides.
Hostii, Baumg. Enum. Stirp. Transs. iii. 342 = rotundifolia.
humilis, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia humilis.
humilis, Schur, Enum. Pl. Transs. 438 = persicifolia.
humillima, *A. DC. Prod.* vii. 474.—Persia.
hungarica, Pers. Syn. i. 191 = lingulata.
hybrida, Bout. ex Willk. & Lange, Prod. Fl. Hisp. ii. 295 = Loeflingii.
hybrida, Linn. Sp. Pl. 168 = Specularia hybrida.
hybrida, Rodigas.—Cf. Illustr. Hortic. xxx. (1883) 118. t. 499 = Van-Houttei.
hygrophila, Rupr. in Bull. Acad. Pétersb. xi. (1867) 209 = Aucheri.
hypopolia, *Trautv. in Act. Hort. Petrop.* iv. (1876) 389.—As. temp.
hyrcania, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 69.—Persia.
illinoënsis, Fresen. Ind. Sem. Hort. Franc. (1836) = americana.
imbricata, Hayne, ex A. DC. Monog. Camp. 284 = caespitosa.
imeretina, *Rupr. in Bull. Acad. Pétersb.* xi. (1867) 217.—Reg. Caucas.
incanescens, *Boiss. Diagn.* Ser. I. vii. 16.—Persia.
inconcessa, Schott, Nym. & Kotschy, Analect. Bot. 10 = Scheuchzeri.
inconspicua, Banks, ex A. DC. Monog. Camp. 157 = Wahlenbergia lobelioides.
incurva, Aucher, ex A. DC. Prod. vii. 464 = Leutweinii.
indica, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia indica.
infundibuliformis, Sims, Bot. Mag. t. 2632 = rapunculoides.
Infundibulum, Reichb. Ic. Pl. Crit. i. t. 75. f. 158 = silenifolia.
Infundibulum, Vest, ex Roem. & Schult. Syst. v. 106 = Steveni.
integerrima, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 155 = sylvatica.
intermedia, Engelm. ex Nutt. in Trans. Am. Phil. Soc. N. S. viii. (1843) 256 = biflora.
intermedia, Roem. & Schult. Syst. v. 110 = Adenophora communis.
interrupta, Pers. Syn. i. 192 = Prismatocarpus interruptus.
Intybus, Schott & Kotschy, ex Tchihat. Asie Min. Bot. ii. 402 = trachyphylla.
involucrata, *Aucher, ex A. DC. Prod.* vii. 467.—As. Min.; Persia.
isophylla, *Moretti, in Brugnat. Giorn. Fis.* II. vii. (1824) 44, 98 (*App. Schomb. Fl. Ital.* 22).—Italia.
Jacobaea, *C. Sm. ex Hook. Niger Fl.* 148. t. 12.—Ins. Cap. Verd.
Jacquini, A. DC. Monog. Camp. 251 = Trachelium Jacquini.
japonica, Vatke, in Linnaea, xxxviii. (1874) 705 = Phyteuma japonicum.
jasionefolia, A. DC. in DC. Prod. vii. 463 = stricta.
Jaubertiana, *Timb. in Bull. Soc. Bot. Fr.* xv. (1868) p. xlviii.—Gallia.
javanica, D. Dietr. Syn. Pl. i. 759 = Campanumoea javanica.
juncea, D. Dietr. l. c. 757 = Lightfootia juncea.
juncea, *Hill, Veg. Syst.* viii. 14.—German.
juncea, Hort. ex A. DC. Monog. Camp. 157 = Wahlenbergia lobelioides.
juncea, *Wettstein, in Stapf, in Denkschr. Acad. Wien*, l. (1885) 117.—Lycia.
juncifolia, Hort. ex A. DC. Monog. Camp. 157 = Wahlenbergia lobelioides.
jurjurensis, *Pomel, Nouv. Mat. Fl. Atl.* 257.—Afr. bor.
kamtschatica, Pall. ex Ledeb. Fl. Ross. ii. 889 = camtschatica.
kashmiriana, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 24 = cashmiriana.
khasiana, *Hook. f. & Thoms. l. c.* 25.—Reg. Himal.
Kitaibeliana, *Roem. & Schult. Syst.* v. 90.—Banatus.
kolenatiana, *C. A. Mey. ex Rupr. in Bull. Acad. Pétersb.* xi. (1867) 216.—Reg. Caucas.
Kotschyana, *A. DC. Prod.* vii. 481.—Cilicia.

CAMPANULA:—
Krebsii, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia Zeyheri.
Kremeri, Boiss. & Reut. Pugill. Pl. Nov. 75 = dichotoma.
kryophila, Rupr. in Bull. Acad. Pétersb. xi. (1867) 211 = ardonensis.
laciniata, Andr. Bot. Rep. t. 385 = tomentosa.
laciniata, *Linn. Sp. Pl.* 165.—Archip. Graec.
lactiflora, *Bieb. Fl. Taur. Cauc.* i. 153.—Reg. Caucas.
Lamarckii, D. Dietr. Syn. Pl. i. 755 = Adenophora Lamarkii.
Lambertiana, A. DC. Monog. Camp. 327 = C. Rapunculus.
lamiifolia, Bieb. Fl. Taur. Canc. i. 154 = alliariaefolia.
lanata, *Frivald. in Flora*, xix. (1836) 434.—Thracia.
lanceolata, Boiss. & Reut. ex Willk. & Lange, Prod. Fl. Hisp. ii. 291 = hispanica.
lanceolata, *Lapeyr. Hist. Abr. Pl. Pyr.* 105.—Mont. Pyren.
lanceolata, Pall. ex A. DC. Monog. Camp. 321 = Steveni.
lanceolata, J. & C. Presl, Fl. Čech. 49 = persicifolia.
lanceolata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = rotundifolia.
lancifolia, Roxb. Fl. Ind. ed. Carey, ii. 96 = Campanumoea celebica.
lancifolia, Schur, Enum. Pl. Transs. 445 = rotundifolia.
Langsdorffiana, Fisch. ex A. DC. Monog. Camp. 279 = linifolia.
Langsdorffiana, A. Gray, in Am. Journ. Sc. Ser. II. xxxiv. (1862) 254 = Parryi.
laniflora, *Hornem. ex Steud. Nom.* ed. II. i. 268.—Hab. ?
lanuginosa, Hort. ex Steud. l. c. = divergens.
lanuginosa, Lam. Encyc. i. 584 = peregrina.
lanuginosa, Willd. Enum. Hort. Berol. 213 = rupestris.
Larreinii, Bert. ex Colla, in Mem. Acc. Tor. xxxviii. (1835) 118. t. 35 = Wahlenbergia fernandeziana.
lasiocarpa, *Cham. in Linnaea*, iv. (1829) 39.—Am. bor. occ.
latifolia, *Linn. Sp. Pl.* 165.—Europ.; Oriens; As. temp.
latiloba, *A. DC. Prod.* vii. 478.—Bithynia.
lavandulaefolia, Reinw. ex Blume, Bijdr. 726 = Wahlenbergia gracilis.
Ledebouriana, *Trautv. in Act. Hort. Petrop.* ii. (1873) 477.—Armenia.
Ledebourii, Boiss. Fl. Orient. iii. 905 = praec.
Lehmanniana, *Bunge, in Lehm. Reliq. Bot.* 211.—Turkestan.
leptocarpa, Engelm. ex A. Gray, in Proc. Am. Acad. xi. (1876) 82 = Specularia leptocarpa.
leptopetala, Ehrenb. ex Boiss. Fl. Orient. iii. 952 = Phyteuma rigidum.
leucanthemifolia, Pourr. in Mém. Acad. Toul. iii. (1788) 309 = caespitosa.
leucoclada, *Boiss. Diagn.* Ser. II. iii. 109.—Afghan.
leucosiphon, *Boiss. & Heldr. Diagn.* Ser. I. xi. 64.—As. Min.
Leutweinii, *Heldr. in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 379.—Graecia.
libanotica, *A. DC. Prod.* vii. 463.—Syria.
ligularis, *Lam. Encyc.* i. 585.—Europ.
ligulata, Pers. Syn. i. 191 = lingulata.
ligulata, Steud. Nom. ed. II. i. 268 = C. Cervicaria.
liliflora, Roth, Enum. Pl. Phan. Germ. i. 715 = Adenophora communis.
lilifolia, Lam. Encyc. i. 581 = Adenophora Lamarkii.
lilifolia, Linn. Sp. Pl. 165 = Adenophora communis.
limonia, Murr. Syst. Veg. ed. XIII. 175 = seq.
limonifolia, Linn. Sp. Pl. ed. II. 239 = Phyteuma limonifolium.
linarioides, Lam. Encyc. i. 580 = Wahlenbergia linarioides.
linearifolia, D. Dietr. Syn. Pl. i. 756 = Prismatocarpus campanuloides.
linearis, Linn. f. Suppl. 140 = Microcodon lineare.
lingulata, Reichb. Ic. Pl. Crit. vi. 12 = C. Cervicaria.
lingulata, *Waldst. & Kit. Pl. Rar. Hung.* i. 65. t. 64.—Europ. austr. or.
linifolia, *Scop. Ann.* ii. *Hist. Nat.* 47.—Europ.
linifolia, Schrank, Prim. Fl. Sal. 70 = rotundifolia.
liniphylla, Pall. ex A. DC. Monog. Camp. 363 = Adenophora marsupiifolia.
linnaeifolia, *A. Gray, in Proc. Am. Acad.* vii. (1868) 366.—Calif.
linoides, Presl, ex A. DC. Monog. Camp. 158 = Wahlenbergia linarioides.

CAMPANULA :—
littoralis, Labill. Nov. Holl. Pl. i. 49. t. 70 = Wahlenbergia gracilis.
lobata, *Schloss. & Vukot. in Rad Jugos. Akad. Zagreb.* xxxi. (1875) 105.—Europ.
lobelioides, Linn. f. Suppl. 140 = Wahlenbergia lobelioides.
lobelioides, Spreng. Syst. i. 736 = Wahlenbergia nutabunda.
lobelioides, Vatke, in Linnaea, xxxviii. (1874) 704 = Phyteuma lobelioides.
Loeflingii, *Brot. Fl. Lusit.* i. 287; *Phyt. Lusit.* i. 41. t. 18.—Lusitan.; Marocco.
longebracteata, D. Dietr. Syn. Pl. i. 757 = Treichelia longebracteata.
longifolia, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa.
longipes, Coss. ex Nym. Consp. 482 = Loeflingii.
longirostris, Banks, ex A. DC. Monog. Camp. 168 = Prismatocarpus crispus.
Loreyi, A. Blanco, ex Nym. Consp. 282 = Loeflingii.
Loreyi, *Pollini, Elem. Bot.* ii. 149 (Lorei).—Europ. austr.
Lostritti, *Tenore, Syll.* 96.—Italia.
lourica, *Boiss. Diagn.* Ser. I. xi. 70.—Persia.
ludoviciana, Torr. ex A. Gray, Syn. Fl. N. Am. ii. 1. 11 = Specularia biflora.
lunariaefolia, Reichb. Ic. Fl. Germ. vi. 750 = rapunculoides.
lunariaefolia, Willd. ex Roem. & Schult. Syst. v. 92 = Phyteuma campanuloides.
lusitanica, Linn. in Loefl. It. Hisp. 111, 126 = Loeflingii.
Lychnitis, Hornem. Hort. Hafn. i. 199 = bononiensis.
lycia, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma lycium.
lyraefolia, Salisb. Prod. 127 = Michauxia campanuloides.
lyrata, Bourg. ex Nym. Consp. 476 = hagielia.
lyrata, *Lam. Encyc.* i. 588.—As. Min.; Reg. Caucas.
macedonica, Boiss. Fl. Orient. iii. 932 = scutellata.
macedonica, Boiss. & Orph. ex Boiss. l. c. 925 = multiflora.
macrantha, Fisch. ex Hornem. Hort. Hafn. Suppl. 23 = latifolia.
macrocalyx, Schur, Enum. Pl. Transs. 441 = pulla.
macrocephala, Willd. ex Steud. Nom. ed. II. i. 268 = Wahlenbergia tenuifolia.
macrochlamys, *Boiss. & Huet, Diagn.* Ser. II. iii. 111.—Armenia.
macrophylla, Sims, Bot. Mag. t. 912 = alliariaefolia.
macrorhiza, Vukot. in Linnaea, xxvi. (1853–55) 329 = spp. plurim.
macrorrhiza, *J. Gay, ex A. DC. Monog. Camp.* 302.—Gallia.
macrostachya, Panz. ex Roem. & Schult. Syst. v. 121 = rapunculoides.
macrostachya, Willd. Enum. Hort. Berol. i. 213 = multiflora.
macrostyla, *Boiss. & Heldr. Diagn.* Ser. I. xi. 65.—Pisidia.
madagascariensis, D. Dietr. Syn. Pl. i. 752 = Wahlenbergia oppositifolia.
Malyi, Schott, Nym. & Kotschy, Analect. Bot. 11 = rotundifolia.
marginata, Thunb. Fl. Jap. 89 = Wahlenbergia gracilis.
maroccana, *Ball, in Journ. Linn. Soc.* xvi. (1878) 554.—Marocco.
Marschalliana, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma campanuloides.
marsupiiflora, Fisch. ex Roem. & Schult. Syst. v. 116 = Adenophora marsupiiflora.
Massonii, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia Massonii.
Mathoneti, Jord. ex Gren. & Godr. Fl. Fr. ii. 418 = caespitosa.
matritensis, A. DC. Monog. Camp. 332 = Loeflingii.
mauritanica, *Pomel, Nouv. Mat. Fl. Atl.* 3.—Afr. bor.
Medium, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa.
Medium, *Linn. Sp. Pl.* 167.—Europ. austr.
Meyeriana, Rupr. in Bull. Acad. Pétersb. xi. (1867) 207 = Aucheri.
michauxioides, *Boiss. Diagn.* Ser. I. iv. 35.—Caria.
micrantha, Schur, Enum. Pl. Transs. 438 = bononiensis.
Microcodon, D. Dietr. Syn. Pl. i. 757 = Microcodon glomeratum.
microphylla, Cav. in Anal. Cienc. iii. (1800) 19 = mollis.
microphylla, Kit. ex Schult. Oestr. Fl. ed. II. 400 = Kitaibeliana.
Minae, *P. G. Stroebl, in Flora*, lxvi. (1883) 549.—Sicil.

CAMPANULA :—
minor, [Honck.] Verz. aller Gew. Teutschl. 617 = caespitosa.
minor, Lem. Fl. Fr. iii. 339 = rotundifolia.
minuta, Agardh, ex Roem. & Schult. Syst. v. 87 = Jasione foliosa.
minuta, Savi, Cent. ii. Fl. Etrusc. 54 = rotundifolia.
minutissima, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 47.—Transs.
mixta, *Hegetschw. Fl. Schw.* 231.—Helvet.
modesta, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 24.—Reg. Himal.
modesta, Schott, Nym. & Kotschy, Analect. Bot. 13 = caespitosa.
mollis, *Linn. Sp. Pl.* ed. II. 237.—Reg. Mediterr. occ.
monadelpha, Pall. ex A. DC. Monog. Camp. 363 = Adenophora coronopifolia.
monantha, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = Scheuchzeri.
monanthos, Pantoč. in Verh. Ver. Natur- u. Heilk. Presb. ii. (1871–72) 54 = praec.?
monocephala, *Trautv. in Act. Hort. Petrop.* vi. (1879) 64.—Turkestan.
montana, *Delarb. Fl. Auvergne*, ed. II. 40.—Gallia.
montevidensis, Spreng. Syst. i. 738 = Specularia biflora?
Moorcroftiana, Wall. Cat. n. 1288 = colorata.
Morettiana, *Reichb. Ic. Fl. Germ.* iv. 18.—Dalmat.
morifolia, Salisb. Prod. 126 = rapunculoides.
multicaulis, Boiss. Diagn. Ser. I. vii. 19 (= *Podanthum persicum*).—Persia.
multiflora, Hort. ex A. DC. Monog. Camp. 271 = bononiensis.
multiflora, *Waldst. & Kit. Pl. Rar. Hung.* iii. 292.—Hungar.
multiflora, Willd. ex Roem. & Schult. Syst. v. 130 = polyantha.
muralis, Portenschl. ex A. DC. Monog. Camp. 303 = Portenschlagiana.
musarum, Heldr. ex Nym. Consp. 477 = rupicola.
myosotidifolia, Boiss. in Tchihat. Asie Min. Bot. ii. 392 = argentea.
muscosa, [*Pieri,*] *in Ionios Anthol.* v. 204.—Ins. Corcyra.
myrtifolia, Boiss. & Heldr. Diagn. Ser. I. xi. 69 = Trachelium myrtifolium.
nana, Lam. Encyc. i. 585 = Allionii.
napuligera, Schur, Enum. Pl. Transs. 444 = rotundifolia.
neglecta, Bess. Cat. Hort. Cremen. (1816) 28 = rapunculoides.
neglecta, Downar, in Bull. Soc. Nat. Mosc. xxxiv. (1861) 1. 182 = praec.?
neglecta, Roem. & Schult. Syst. v. 104 = patula.
nemorosa, A. DC. Monog. Camp. 274 = rapunculoides.
nervosa, Royle, Illustr. Bot. Himal. 253 = colorata.
nicaeensis, Risso, ex A. DC. Monog. Camp. 302 = macrorrhiza.
nicaeensis, Roem. & Schult. Syst. v. 126 = glomerata.
nitida, [Soland. in] Ait. Hort. Kew. i. 346 = planiflora.
nobili-macrantha, *Regel, Ind. Sem. Hort. Petrop.* (1869) 14.—Hab.?
nobilis, Lindl. in Journ. Hort. Soc. i. (1846) 232 = punctata.
notata, Schott, Nym. & Kotschy, Analect. Bot. 13 = caespitosa.
nudicaulis, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia nudicaulis.
numidica, *Dur. Expl. Algér.* t. 62.—Afr. bor.
nutabunda, Guss. in Tenore, Append. v. ad Fl. Nap. Prod. 8 = Wahlenbergia nutabunda.
nutans, Lam. Fl. Fr. ii. 336 = rapunculoides.
nutans, Sieber, ex A. DC. Monog. Camp. 366 = Symphyandra cretica.
nutans, Vahl, ex Hornem. Hort. Hafn. 201 = sibirica.
Nuttallii, *Penny, ex Loud. Hort. Brit. Suppl.* i. 583.—Am. bor.
obliqua, Jacq. Hort. Schoenb. iii. 47 = americana.
obliquifolia, Tenore, Fl. Nap. Prod. 15 = bononiensis.
odontosepala, *Boiss. Diagn.* Ser. I. xi. 66.—Persia.
oenipontana, Moretti, ex A. DC. Monog. Camp. 269 = rapunculoides.
oligosantha, Schur, Enum. Pl. Transs. 436 = C. Trachelium.
Olivieri, A. DC. Monog. Camp. 233. t. 9 = calamenthifolia.
olympica, *Boiss. Diagn.* Ser. I. iv. 34.—As. Min.
orbelica, *Panč. Elem.* 48.—Europ.

CAMPANULA :—
oreadum, *Boiss. & Heldr. Diagn.* Ser. II. iii. 107.—Thessalia.
oreophila, Schur, Enum. Pl. Transs. 441 = carpatica.
orientalis, Steud. Nom. ed. II. i. 269 = peregrina?
Orphanidea, *Boiss. Fl. Orient.* iii. 897.—Macedon.
ossetica, Bieb. Fl. Taur. Cauc. iii. 145 = Symphyandra ossetica.
Ottoniana, Roem. & Schult. Syst. v. 115 = Lightfootia oxyococcoides.
ovata, Spreng. Syst. iv. Cur. Post. 78 = Wahlenbergia ovata.
oxyphylla, Vukot. in Linnaea, xxvi. (1853–55) 331 = spp. plurim.
paenina, *Reut. ex Tissière*, Bot. Gr. St. Bernard, 63.—Helvet.
Pallasiana, Roem. & Schult. Syst. v. 138 = pilosa.
pallida, Wall. in As. Research. xiii. (1820) 375 = colorata.
paniculata, Linn. f. Suppl. 139 = Wahlenbergia paniculata.
paniculata, Pohl, ex A. DC. Monog. Camp. 244 = sibirica.
paniculata, Sm. ex A. DC. l. c. 154 = Wahlenbergia Banksiana.
paniculata, *Turra, ex Sacc. in Atti Ist. Venet.* Ser. III. ix. (1863–64) 486.—Italia.
paniculata, Wall. Cat. n. 1295 A, B = Cephalostigma paniculatum.
pannonica, Bernh. ex Schrank, in Denkschr. Regensb. Ges. ii. (1822) 34 = sibirica.
parnassica, *Boiss. & Sprun. Diagn.* Ser. I. vii. 17.—Graecia.
Parryi, *A. Gray, in Syn. Fl. N. Am.* ii. 1. 395.—Am. bor. occ.
parviflora, Lam. Encyc. i. 588 = sibirica.
parviflora, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = C. Erinus.
parviflora, Salisb. Prod. 126 = Wahlenbergia lobelioides.
parvula, Jord. Pugill. Pl. Nov. (1852) 123 = caespitosa.
patens, Gueldenst. ex Ledeb. Fl. Ross. ii. 887 = patula.
patula, *Linn. Sp. Pl.* 163.—Europ.
patula, Sibth. & Sm. Fl. Graec. Prod. i. 137 = Spruneriana.
pauciflora, Desf. Choix Pl. 36. t. 26 = Spruneriana.
pauciflora, Lam. ex Steud. Nom. ed. II. i. 269 = *planiflora*, Lam.
peduncularis, Wall. Cat. n. 7139 = Wahlenbergia peduncularis.
pedunculata, J. F. Gmel. Syst. 352 = Wahlenbergia paniculata.
peirescifolia, Spreng. Syst. i. 735 = Adenophora latifolia.
pelviformis, *Lam. Encyc.* i. 586.—Ins. Creta.
pendula, Bieb. Fl. Taur. Cauc. i. 154 = Symphyandra pendula.
pennica, Reut. ex Payot, in Bull. Soc. Bot. Fr. xxix. (1882).—Cf. Bibl. 43 = seq.
pennina, Reut. in Comptes-Rend. Soc. Hallér. (1854–56) 115 = rotundifolia.
Pentagonia, Linn. Sp. Pl. 169 = Specularia Pentagonia.
pentagonophylla, Vukot. in Linnaea, xxvi. (1853–55) 325 = Wahlenbergia hederacea.
peregrina, Hoffmgg. & Link, Fl. Port. ii. 15. t. 83 = primulaefolia.
peregrina, *Linn. Mant.* ii. 204.—As. Min.; Syria.
Pereskia, Fisch. ex A. DC. Monog. Camp. 357 = Adenophora latifolia.
pereskiaefolia, Fisch. ex Roem. & Schult. Syst. v. 116 = Adenophora latifolia.
perfoliata, Linn. Sp. Pl. 169 = Specularia perfoliata.
periplocifolia, Lam. Encyc. i. 580 = Adenophora periplocaefolia.
perneglecta, *Schott, Nym. & Kotschy, Analect. Bot.* 11.—Austria.
Perpentiae, Moretti, ex A. DC. Monog. Camp. 305 = Rainerii.
perpusilla, *A. DC. Prod.* vii. 474.—Persia.
persepolitana, *Kotschy, ex Boiss. Diagn.* Ser. I. vii. 19.—Persia.
persica, A. DC. Prod. vii. 483 (= *Podanthum persicum*).—Persia.
persicifolia, *Linn. Sp. Pl.* 164.—Europ.; As. bor.
Pestalozzae, Boiss. Diagn. Ser. I. xi. 62 = propinqua.
petiolata, A. DC. Monog. Camp. 278 = rotundifolia.
petraea, All. Fl. Pedem. i. 112 = glomerata.

CAMPANULA:—
petraea, Habl. Taur. 157 = bononiensis.
petraea, Kotschy, ex A. DC. Prod. vii. 467 = involucrata.
petraea, *Linn. Syst.* ed. X. 927.—Italia.
petraea, Zanted. ex Moretti, in Syll. Ratisb. i. (1824) 154 = elatinoides.
petrophila, *Rupr. in Bull. Acad. Pétersb.* xi. (1867) 212.—Reg. Caucas.
phrygia, *Jaub. & Spach, Illustr. Pl. Or.* iii. 42.—Macedon.; As. Min.
phyctidocalyx, *Boiss. & Noé, Diagn.* Ser. II. iii. 114.—Armenia.
phyteumoides, Sibth. ex Zuccagni, in Roem. Collect. 128 = Phyteuma limoniifolium.
Pichleri, Vis. in Mem. Ist. Venet. xvi. (1871) 106 = Phyteuma amplexieaule.
pilosa, Less. ex Ledeb. Fl. Ross. ii. 894 = Adenophora tricuspidata.
pilosa, *Pall. ex Roem. & Schult. Syst.* v. 148.—Sibir.; Am. bor. occ.
pinifolia, Vukot. in Rad Jugos. Akad. Zagreb. lvii. (1881) 92 = patula.
planiflora, Engelm. in Coult. Bot. Gaz. vii. (1882) 5 = Parryi.
planiflora, Lam. Eneyc. i. 580 = pyramidalis?
planiflora, Willd. Enum. Hort. Berol. 210 = versicolor.
plicata, Pers. Syn. i. 193 = Prismatocarpus crispus.
plicatula, Dum. Fl. Belg. 58 = C. Trachelium.
podanthoides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 938.—Cataonia.
podoearpa, *Boiss. Diagn.* Ser. I. xi. 68.—Pamphylia.
polyantha, *Roem. & Schult. Syst.* v. 130.—Sibiria.
polymorpha, Banks & Soland. ex A. DC. Monog. Camp. 142 = Wahlenbergia gracilis.
pomponiifolia, Fisch. ex A. DC. l. c. 363 = Adenophora Gmelini.
porosa, Thunb. Prod. Pl. Cap. 39 = Salmolus Valerandi.
Portenschlagiana, *Roem. & Schult. Syst.* v. 93.—Dalmatia.
portensis, Linn. in Loefl. It. Hisp. 111, 127 = C. Erinus.
Pourretii, *Jeanb. & Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* vi. (1883–84) 153.—Gallia.
pratensis, Lapyl. ex A. DC. Monog. Camp. 287 = rotundifolia.
pratensis, Reut. ex Nym. Consp. 478 = C. Trachelium.
precatoria, Timb. in Mém. Acad. Sc. Toul. Sér. VII. v. (1873) 271 = lanceolata.
Preissii, De Vriese, in Lehm. Pl. Preiss. ii. 241 = Wahlenbergia gracilis.
prenanthoides, *Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 93.—Calif.
primulaefolia, *Brot. Fl. Lusit.* i. 288.—Lusitan.
Prismatocarpus, L'Hérit. Sertum, t. 3 = Prismatocarpus nitidus.
procumbens, Linn. f. Suppl. 141 = Wahlenbergia procumbens.
propinqua, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1835) 32.—As. Min.; Persia.
pruinosa, [*Pieri*] *in Ionios Anthol.* v. 204.—Ins. Corcyra.
prostrata, Dulac, Fl. Hautes-Pyr. 457 = Scheuchzeri.
pseudo-carpatica, Schur, Enum. Pl. Transs. 441 = reniformis.
pseudolaneeolata, *Pantoč. in Magyar Növent. Lapok*, vi. (1882) 162.—Hungar.
pseudo-pulla, Schur, Enum. Pl. Transs. 441 = pulla.
pseudo-valdensis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = rotundifolia.
psilostachya, *Boiss. & Kotschy, Diagn.* Ser. II. iii. 113.—Cilicia.
ptarmicaefolia, *Lam. Encyc.* i. 570.—As. Min.
pubescens, Reichb. ex Nym. Consp. 479 = rotundifolia.
pubescens, Schmidt, Fl. Boëm. Cent. ii. 67 = caespitosa.
pubiflora, Rupr. in Bull. Acad. Pétersb. xi. 207 = Aucheri.
pulchella, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma pulchellum.
pulchella, Jord. Pugill. Nov. Pl. (1852) 126 = caespitosa.
pulchella, Salisb. Prod. 127 = Specularia Speculum.
pulcherrima, *Schrank & Zeyh. ex Steud. Nom.* ed. II. i. 269.—Hab.?

CAMPANULA:—
pulla, Baumg. ex Stend. Nom. ed. II. i. 268, 269 = linifolia.
pulla, *Gueldenst. It.* ii. 24; *ex Ledeb. Fl. Ross.* ii. 891.—Reg. Caucas.
pulla, *Linn. Sp. Pl.* 163.—Europ.
pulla, Parol. ex Pollini, Fl. Ver. i. 268 = Morettiana.
pumila, Curt. Bot. Mag. t. 512 = caespitosa.
pumila, Schmidt, Fl. Boem. ii. 71 = persicifolia.
Pumilio, Portenschl. ex Roem. & Schult. Syst. v. 136 = Wahlenbergia Pumilio.
punctata, *Lam. Encyc.* i. 586.—Sibir.; Japon.
punduana, D. Dietr. Syn. Pl. i. 757 = Campanumoea parviflora.
purpurea, Spreng. Syst. iv. Cur. Post. 78 = Codonopsis purpurea.
pusilla, Haenke, in Jacq. Collect. ii. 79 = caespitosa.
pygmaea, D. Dietr. Sp. Pl. i. 757 = Microcodon glomeratum.
pygmaea, Lam. ex DC. Fl. Fr. iii. 705 = Borago laxiflora.
pyramidalis, Gilib. Fl. Lituan. i. 52 = bononiensis.
pyramidalis, *Linn. Sp. Pl.* 164.—Europ.
pyramidata, Gilib. Hist. Pl. Eur. ii. 209 = praec.
pyramidiflora, Reichb. Cat. Sem. Hort. Dresd. (1822) = rapuneuloides.
pyrenaica, *A. DC. Monog. Camp.* 324.—Mont. Pyren.
pyrenaica, 'Hecht. Willd.' ex Steud. Nom. ed. II. i. 269, 270 = rhomboidea.
quadrifida, R. Br. Prod. 561 = Wahlenbergia gracilis.
Quartiniana, *A. Rich. Tent. Fl. Abyss.* ii. 5.—Abyss.
Rabelaisiana, Roem. & Schult. Syst. v. 158 = Adenophora Gmelini.
racemosa, S. G. Gmel. It. i. 9 = sibirica.
racemosa, Opiz, ex Steud. Nom. ed. II. i. 269 = rapuneuloides.
racemosa, Vukot. in Linnaea, xxvi. (1851) 332 = spp. plurim.
Raddeana, *Trautv. in Bull. Acad. Pétersb.* x. (1866) 395.—Reg. Caucas.
radicosa, *Bory & Chaub. Fl. Pélop.* 14. t. 7.—Graecia.
Radula, *Fisch. ex Tchihat. Asie Min. Bot.* ii. 395.—Kurdistan.
Rainerii, *Perpent. in Bibl. Ital.* v. (1817) 134.—Italia.
ramosissima, Griseb. Spicil. Fl. Rumel. ii. 282 = phrygia.
ramosissima, Host, Fl. Austr. i. 264 = Loreyi.
ramosissima, *Sibth. & Sm. Fl. Graec. Prod.* i. 137.—Graecia.
ramosissima, Sprun. ex Nym. Consp. 481 = Spruneriana.
ramosissima, Willd. ex Steud. Nom. ed. II. i. 269 = Loeflingii.
ramulosa, Wall. Cat. n. 1286; in Roxb. Fl. Ind. ed. Carey, ii. 100 = colorata.
ranunculoides, Uspensky, in Bull. Soc. Nat. Mosc. vii. (1834) 369 = rapunculoides.
rapuncula, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = C. Rapunculus.
rapunculoides, *Linn. Sp. Pl.* 165.—Europ.; As. temp.
rapunculoides, Pall. ex Steud. Nom. ed. II. i. 270 = bononiensis.
Rapunculus, *Linn. Sp. Pl.* 164.—Europ.; Oriens; Afr. bor.; As. bor.
Rapunculus, O. F. Muell. Fl. Dan. t. 855 = rotundifolia.
Ravcyi, *Boiss. Diagn.* Ser. I. iv. 32.—Caria.
Reboudiana, Gren. & Godr. Fl. Fr. ii. 415 = rotnndifolia.
Reboudiana, *Pomel, Nouv. Mat. Fl. Atl.* 2.—Algeria.
recta, Dulac, Fl. Hautes-Pyr. 458 = linifolia.
Redowskiana, Cham. & Schlecht. in Linnaea, iv. (1829) 41 = Wahlenbergia hemallanthina.
Redowskyi, Fisch. ex Herd. in Act. Hort. Petrop. i. (1873) 288 = pilosa.
redux, Schott, Nym. & Kotschy, Analect. Bot. 9 = Scheuchzeri.
reflexa, Hausm. ex Nym. Consp. 480 = carnica.
reflexa, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = caespitosa.
remotiflora, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 180.—Japon.
reniformis, *Schur, Enum. Pl. Transs.* 441.—Transsylv.
repanda, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma limonifolium.
repens, Hort. ex Steud. Nom. ed. II. i. 270 = Wahlenbergia procumbens.
repens, *Lour. Fl. Cochinch.* 139.—Cochinch.
retrorsa, *Labill. Ic. Pl. Syr. Dec.* v. 5.—Syria; Cilicia.

CAMPANULA:—
Reverchoni, *A. Gray, Syn. Fl. N. Am.* ii. 1. 396.—Texas.
Reuteriana, *Boiss. & Bal. Diagn.* Ser. II. iii. 108.—As. Min.; Persia.
Reyneri, D. Dietr. Syn. Pl. i. 749 = Rainerii.
rhodensis, A. DC. Monog. Camp. 297 = drabaefolia.
Rhodii, Loisel. Fl. Gall. ed. II. i. 140 = linifolia.
rhomboidalis, Auct. ex Steud. Nom. ed. II. i. 270 = lanceolata.
rhomboidalis, Gorter, Fl. Ingr. 34 = rapuneuloides.
rhomboidalis, *Linn. Sp. Pl.* 165.—Europ.
rhomboidalis, Schur, Enum. Pl. Transs. 445 = arenata.
rhomboidea, Endr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 291 = lanceolata.
rhomboidea, *Murr. Syst.* ed. XIII. 173.—Europ.
rigescens, Pall. ex Roem. & Schult. Syst. v. 102 = silenifolia.
rigida, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma rigidum.
rigida, Gilib. Fl. Lituan. i. 51 = rapuneuloides.
rigidipila, *Steud. & Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 3.—Abyss.
rimarum, *Boiss. Fl. Orient.* iii. 931.—Lycia.
riparia, Boj. ex A. DC. Prod. vii. 432 = Wahlenbergia emirnensis.
riparia, Leprieur & Perr. ex A. DC. l. c. 435 = Wahlenbergia riparia.
Rochelii, Schur, in Oestr. Bot. Zeitschr. (1861) 45 = Steveni.
roëlloides, Juss. ex A. DC. Monog. Camp. 165 = Prismatocarpus paniculatus.
Roezli, Regel, Gartenfl. (1872) 239; et in Act. Hort. Petrop. ii. (1873) 310 = prenanthoides.
Rohdii, Loisel. Fl. Gall. ed. II. i. 140. t. 24 = linifolia.
Rosani, Tenore, Fl. Nap. iii. 205. t. 120 = versicolor.
rosea, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 12.—Malaya.
rosulata, Vukot. in Linnaea, xxvi. (1853–55) 326 = cenisia.
rotunda, Gilib. Fl. Lituan. i. 46 = rotundifolia.
rotundifolia, All. ex Steud. Nom. ed. II. i. 270 = rhomboidea.
rotundifolia, Boiss. Voy. Espagne, 399 = macrorrhiza.
rotundifolia, Ledeb. Fl. Alt. i. 238 = linifolia.
rotundifolia, *Linn. Sp. Pl.* 163.—Reg. bor. temp.
rotundifolia, Pall. ex A. DC. in DC. Prod. vii. 484 = adscendens.
rubioides, Banks, ex A. DC. Monog. Camp. 137 = Microcodon lineare.
rubra, Hort. Jonrn. & Fl. des Jard. (1832) 120. t. 120 = rhomboidalis.
ruderalis, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 174.—Afghan.
rumeliana, *Vatke, in Linnaea*, xxxviii. (1874) 705.—Rumel.
rupestris, Adams, ex Ledeb. Fl. Ross. ii. 876 = bellidifolia.
rupestris, Bieb. Fl. Taur. Cauc. i. 154 = Bieberstciniana.
rupestris, Host, Fl. Austr. i. 263 = Waldsteiniana.
rupestris, Risso, ex A. DC. Monog. Camp. 302 = macrorrhiza.
rupestris, Sibth. & Sm. Fl. Graec. Prod. i. 142 = tomentosa.
rupestris, Sprun. ex Nym. Consp. 477 = rupicola.
rupicola, *Boiss. & Sprun. Diagn.* Ser. I. vii. 17.—Graecia.
Ruprechtii, *Boiss. Fl. Orient.* iii. 905.—Persia.
ruscinonensis, Timb. in Mém. Acad. Sc. Toul. Ser. VII. v. (1873) 275 = linifolia.
Russeliana, Roem. & Schult. Syst. v. 142 = strigosa.
ruthenica, Bieb. Fl. Taur. Cauc. i. 151 = bononiensis.
sabatia, *De Not. Prosp. Fl. Ligust.* 52.—Italia.
sajanensis, [Fisch.] Hort. Gorenk. ed. I. (1808) 52 = Adenophora marsupiiflora.
sajanensis, Spreng. ex Steud. Nom. ed. II. i. 270 = lasiocarpa.
salicifolia, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma canescens.
salicifolia, Boiss. & Kotschy, ex Boiss. Fl. Orient. iii. 949 = Phyteuma amplexicaule.
salicifolia, Juss. ex Ledeb. Fl. Ross. ii. 894 = Adenophora coronopifolia.
sanensis, Fisch. Cat. Hort. Gorenk. (1808), ex A. DC. in Prod. vii. 461 = pilosa.
sarmatica, *Ker-Gawl. in Bot. Reg.* t. 237.—Reg. Caucas.

CAMPANULA :—
sarmentosa, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 4. —Abyssin.
sarmentosa, Fenzl & Vukot. in Rad Jugos. Akad. Zagreb. xxxi. (1875) 105 = lobata.
Sartorii, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iii. 919. —Archip. Graec.
saxatilis, Georgi, Beschr. Russ. Reich. iii. iv. 776 (sp. dub.).—Rossia.
saxatilis, *Linn. Sp. Pl.* 167.—Ins. Creta.
saxicola, R. Br. Prod. 561 = Wahlenbergia saxicola.
Saxifraga, *Bieb. Fl. Taur. Cauc.* i. 155.—Reg. Caucas.
scabrella, *Engelm. in Coult. Bot. Gazette*, vi. (1881) 237.—Am. bor. occ.
scabrida, *Hochst. in Lorent, Wanderungen*, 331.—Mesopotam.
scandens, *Pall. ex Roem. & Schult. Syst.* v. 120.—Sibir.
Scheuchzeri, A. Gray, Syn. Fl. N. Am. ii. 1. 12 = Parryi.
Scheuchzeri, Lodd. Bot. Cab. t. 485 = rotundifolia.
Scheuchzeri, *Vill. Prosp.* 22.—Mont. Pyren.
Schimperi, Vatke, in Linnaea, xxxviii. (1874) 712 = rigidipila.
Schleicheri, Suter, Fl. Helv. i. 124 = linifolia.
Schotti, *Hort.—Cf. Gard. Chron.* (1885) ii. 147.—Hab. ?
sclerotricha, *Boiss. Diagn.* Ser. I. xi. 66.—Persia.
Scouleri, *Hook. ex A. DC. Monog. Camp.* 312.—Am. bor. occ.
scutellata, *Griseb. Spicil. Fl. Rumel.* ii. 282.—Macedon.
secunda, Schmidt, Fl. Boëm. ii. 74 = rapunculoides.
secundiflora, Vis. & Panč. in Mem. Ist. Venet. x. (1861) 442 = rapunculoides.
seminuda, Vest, in Roem. & Schult. Syst. v. 91 = Steveni.
serpyllifolia, *Vis. Fl. Dalm.* ii. 136. t. 15.—Dalmatia.
serratifolia, Vukot. in Linnaea, xxvi. (1853-55) 335 = C. Trachelium, latifolium.
sessiliflora, *Boiss. Fl. Orient.* iii. 947.—Armenia.
sessiliflora, D. Dietr. Syn. Pl. i. 756 = Prismatocarpus campanuloides.
sessiliflora, C. Koch, in Linnaea, xix. (1847) 30 = latiloba.
sessiliflora, Linn. f. Suppl. 139 = Lightfootia longifolia, subulata.
sessiliflora, Vukot. in Linnaea, xxvi. (1853-55) 327 = Rainerii.
sessilis, D. Dietr. Syn. Pl. i. 756 = Prismatocarpus sessilis.
setacea, D. Dietr. l. c. 754 = Wahlenbergia capillacea.
setosa, Fisch. ex DC. Monog. Camp. 269 = rapunculoides.
setosa, *Wendl. Beobacht.* 42.—Hab. ?
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) ii. 188.—As. temp.
sibirica, *Linn. Sp. Pl.* 167.—Europ.; As. temp.
Sibthorpiana, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma Sibthorpianum.
sidoniensis, *Boiss. & Blanche, Diagn.* Ser. II. iii. 114. —Syria.
Sieberi, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia gracilis.
silenifolia, *Fisch. ex A. DC. Monog. Camp.* 320.—As. bor.
silenifolia, Host, Fl. Austr. i. 268 = Wahlenbergia Pumilio.
simplex, Lam. ex DC. Fl. Fr. iii. 730 = bononiensis.
simplex, Stev. in Mém. Soc. Nat. Mosc. iii. (1812) 255 = Steveni.
Sinai, Boiss. Diagn. Ser. I. xi. 76 = Phyteuma lanceolatum.
sinensis, D. Dietr. Syn. Pl. i. 755 = Adenophora sinensis.
singarensis, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. Dec.* i. 2.—Mesopotam.
sparsiflora, D. Dietr. Syn. Pl. i. 757 = Microcodon sparsiflorum.
spathulaefolia, Sprun. ex Nym. Consp. 483 = ramosissima.
spathulata, Ehrenb. ex Boiss. Fl. Orient. iii. 918 = trichopoda.
spathulata, Sibth. & Sm. Fl. Graec. Prod. i. 137 = Spruneriana.
spathulata, Waldst. & Kit. Pl. Rar. Hung. ii. 286. t. 258 = sibirica.
speciosa, Gilib. Fl. Lituan. i. 50 = persicifolia.

CAMPANULA :—
speciosa, Hornem. Hort. Hafn. ii. 957, Add. = glomerata.
speciosa, *Pourr. in Mém. Acad. Sc. Toul.* iii. (1788) 309.—Mont. Pyren.
speciosa, Willd. ex Spreng. Syst. i. 729 = elegans.
specularioides, *Coss. Notes Crit.* 41.—Hispan.
Speculum, Linn. Sp. Pl. 168 = Specularia Speculum.
sphaerothrix, Griseb. Spicil. Fl. Rumel. ii. 280 = expansa.
spicata, *Linn. Sp. Pl.* 166.—Europ.
spicata, Geners. Elench. Scepus. n. 192; ex Wahlenb. Fl. Carp. 62 = C. Cervicaria.
spinulosa, D. Dietr. Syn. Pl. i. 754 = Wahlenbergia spinulosa.
spreta, Roem. & Schult. Syst. v. 123 = Adenophora communis.
Spruneri, Boiss. Fl. Orient. iii. 937 = seq.
Spruneriana, *Hampe, in Flora*, xxv. (1842) i. 76.—Graecia et Ins.
spuria, Pall. ex Roem. & Schult. Syst. v. 154 = Specularia hybrida.
St.-Helenae, D. Dietr. Syn. Pl. i. 755 = Wahlenbergia linifolia.
stellarioides, D. Dietr. l. c. 753 = Wahlenbergia stellarioides.
stellaris, *Boiss. Diagn.* Ser. I. xi. 63.—Syria; Palestin.
stellata, *Thunb. in Hoffm. Phyt. Blaett.* i. (1803) 20.—Afr. austr.
Stelleri, Steph. ex Herd. in Act. Hort. Petrop. i. (1873) 299 = lasiocarpa.
stenanthina, Ledeb. in Mém. Acad. Pétersb. v. (1814) 525 = Adenophora marsupiiflora.
stenocarpa, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 61).—As. temp.
stenocodon, Boiss. & Reut. Diagn. Ser. II. iii. 112 = rotundifolia.
stenophylla, Boiss. & Heldr. l. c. Ser. I. xi. 77 = Phyteuma linifolium.
stenosiphon, Boiss. & Heldr. l. c. vii. 18 = glomerata.
stenosiphon, Reichb. f. ex Nym. Consp. 479 = rotundifolia.
Steveni, *Bieb. Fl. Taur. Cauc.* iii. 138.—Reg. Caucas.; Persia.
Steveniana, Roem. & Schult. Syst. v. 91 = praec.
stolonifera, *Mign. in Bull. Soc. Bot. Fr.* xii. (1865) 342.—Hispan.
stricta, Labill. Ic. Pl. Syr. Dec. v. 6 = libanotica.
stricta, *Linn. Sp. Pl.* ed. II. 238.—As. Min.
stricta, Sibth. & Sm. Fl. Graec. Prod. i. 140 = lyrata.
stricta, Wall. in As. Research. xiii. (1820) 374 = sylvatica.
strigillosa, *Boiss. in Ann. Sc. Nat.* Sér. IV. ii. (1854) 251.—Cappadocia.
strigosa, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 246.—Cilicia; Syria.
strigosa, *Vahl, Symb. Bot.* iii. 34.—As. Min.
stylidioides, Boiss. Diagn. Ser. I. xi. 77 = Phyteuma lobelioides.
stylosa, Bess. Cat. Hort. Crem. (1816) = Adenophora communis.
stylosa, Lam. Encyc. i. 580 = Adenophora stylosa.
styriaca, Schott, Nym. & Kotschy, Analect. Bot. 9 = Scheuchzeri.
suanetica, *Rupr. in Bull. Acad. Pétersb.* xi. (1867) 215. —Reg. Caucas.
suaveolens, Gilib. Fl. Lituan. i. 50 = Adenophora communis.
suaveolens, *Wender. ex Steud. Nom.* ed. II. i. 270.—Hab. ?
subalpina, *Delponte & Gras, Ind. Sem. Hort. Taurin.* (1873) 35.—Hab. ?
subpyrenaica, *Timb. in Mém. Acad. Toul.* Sér. IV. v. (1855) 387.—Hispan.
subramulosa, Jord. in Gren. & Godr. Fl. Fr. ii. 418 = caespitosa.
subulata, Beauv. ex A. DC. Prod. vii. 478 = americana.
subulata, Spreng. Syst. i. 809 = Lightfootia subulata.
subulata, Thunb. Prod. Pl. Cap. 38 = Prismatocarpus subulatus.
subuniflora, Lam. Tabl. Encyc. ii. 53 = liniflora.
sulphurea, *Boiss. Diagn.* Ser. I. xi. 64.—Syria.
swellendamensis, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia swellendamensis.
sylvatica, *Wall. Cat.* n. 1293; *et in Roxb. Fl. Ind.* ii. 97.—Reg. Himal.

CAMPANULA :—
syngenesiflora, Pall. ex A. DC. Monog. Camp. 363 = Adenophora marsupiiflora.
syriaca, Ehrenb. ex Boiss. Fl. Orient. iii. 928 = stellaris.
syriaca, Willd. ex Roem. & Schult. Syst. v. 133 = Specularia falcata.
syspirensis, *C. Koch, in Linnaea*, xxiii. (1850) 639.—Armenia.
tauricola, Boiss. & Bal. Diagn. Ser. II. iii. 116 = Phyteuma rigidum.
telephioides, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. Dec.* i. 1.—Cataonia.
tenella, Jord. Pugill. Pl. Nov. (1852) 125 = caespitosa.
tenella, Lange, ex Nym. Consp. 479 = hispanica.
tenella, Linn. f. Suppl. 141 = Lightfootia tenella, Thunbergiana.
tenerrima, D. Dietr. Syn. Pl. i. 757 = Prismatocarpus tenerrimus.
Tenorii, Moretti, Prosp. Geogr. Bot. Ital. Schouw, 19 = versicolor.
tenuiflora, Schur, Enum. Pl. Transs. 438 = bononiensis.
tenuiflora, Tenore, Fl. Nap. iii. 207. t. 210 = linguulata.
tenuifolia, Hoffm. Fl. Germ. i. 100 = rotundifolia.
tenuifolia, Waldst. & Kit. Pl. Rar. Hung. ii. 168 = Wahlenbergia tenuifolia.
tetraphylla, Thunb. Pl. Jap. 87 = Adenophora verticillata.
teucrioides, *Boiss. Diagn.* Ser. I. iv. 33.—Lydia.
Teymonsii, *Hort.—Cf. Gard. Chron.* (1885) ii. 147, 182.—Hab. ?
Thaliana, Wallr. Scheď. Crit. 86 = bononiensis.
thalictriflora, Spreng. Syst. iv. Cur. Post. 77 = Codonopsis thalictrifolia.
Thomasii, Tenore, ex A. DC. Monog. Camp. 309 = versicolor.
Thomsonii, Hook. f. in Journ. Linn. Soc. ii. (1858) 25 = Phyteuma Thomsoni.
Thunbergii, Roem. & Schult. Syst. v. 135 = Lightfootia ciliata.
thyrsoidea, Lapeyr. Hist. Abr. Pl. Pyr. 107 = speciosa.
thyrsoides, *Linn. Sp. Pl.* 167.—Europ.
tomentosa, *Lam. Encyc.* i. 584.—As. Min.
tomentosa, *Vent. Hort. Cels*, 18.—Graecia et Ins.
Tommasiniana, Koch, ex F. Schultz, Arch. Fl. Fr. et Allem. 229; Reut. in Cat. Genev. (1865) 4 = Waldsteiniana.
trachelioides, Bieb. Fl. Taur. Cauc. i. 150 = rapunculoides.
trachelioides, *Munby, in Bull. Soc. Bot. Fr.* ii. (1855) 285.—Afr. bor.
Trachelium, Brot. Fl. Lusit. i. 287 = primulaefolia.
Trachelium, Bull. Herb. Fr. t. 319 = rapunculoides.
Trachelium, *Linn. Sp. Pl.* 166.—Europ.; As. temp.; Afr. bor.
Trachelium, Thunb. Fl. Jap. 88 = punctata.
trachyphylla, *Schott & Kotschy, ex Boiss. Fl. Orient.* iii. 926.—Cilicia.
transsilvanica, *Schur, ex Andrae, in Bot. Zeit.* xiii. (1855) 328.—Transsylv.
trichocalycina, Tenore, Fl. Nap. i. 67. t. 16 = Phyteuma amplexicaule.
trichopoda, *Boiss. Diagn.* Ser. I. xi. 68.—Syria.
tricuspidata, Fisch. ex Roem. & Schult. Syst. v. 158 = Adenophora denticulata.
tridens, Rupr. in Bull. Acad. Pétersb. xi. (1867) 204 = tridentata.
tridentata, *Schreb. Ic.* 3. t. 2; *Linn. Mant.* i. 44.—As. Min.
trigona, Ehrenb. ex Boiss. Fl. Orient. iii. 959 = Specularia Speculum.
trilocularis, Turra, Prod. Fl. Ital. 64 = Allionii.
triphylla, Thunb. Fl. Jap. 87 = Adenophora triphylla.
truncata, D. Dietr. Syn. Pl. i. 757 = Campanumoea celebica.
tubiflora, Tausch, ex Ledeb. Fl. Ross. ii. 880 = glomerata.
tubulosa, *Lam. Encyc.* i. 588.—Ins. Creta.
turbinata, Schott, Nym. & Kotschy, Analect. Bot. 14 = carpatica.
tyrolensis, Schott, Nym. & Kotschy, l. c. 12 = caespitosa.
ucranica, Bess. Cat. Hort. Cremen. (1816) 28 = rapunculoides.
umbellifera, Vukot. in Linnaea, xxvi. (1853-55) 333 = pyramidalis.

CAMPANULA:—
umbrosa, F. G. Dietr. Vollst. Lex. Gaertn. i. 675 = Adenophora communis.
undulata, Linn. f. Suppl. 142 = Wahlenbergia undulata.
undulata, Moench, Meth. Suppl. 189 = sibirica.
unidentata, Linn. f. Suppl. 139 = Lightfootia unidentata.
uniflora, Georgi, Beschr. Russ. Reich. iii. iv. 772 = silenifolia.
uniflora, Gortcr, Fl. Ingr. 33 = rotundifolia.
uniflora, Honck. Syn. Pl. Germ. ii. 251 = cenisia.
uniflora, *Linn. Sp. Pl.* 163.—Reg. Arctic.
uniflora, Schult. Oestr. Fl. ed. II. 400 = caespitosa.
uniflora, Vill. Prosp. 22 = linifolia.
urticaefolia, All. Fl. Pedem. i. 110 = latifolia.
urticaefolia, Gilib. Fl. Lithuan. i. 52 = bononiensis.
urticaefolia, Turra, Prod. Fl. Ital. 64 = rapunculoides.
urticifolia, Schmidt, Fl. Boëm. ii. 73 = C. Trachelium.
valdensis, All. Fl. Pedem. i. 109 = linifolia.
Van-Houttei, *Carr. in Rev. Hortic.* (1878) 420.—Hab.?
Vandesi, *G. Don, ex Loud. Hort. Brit.* 77.—Hab.?
variifolia, Salisb. Prod. 126 = rotundifolia.
Vayredae, *Leresche & Levier, in Journ. Bot.* xvii. (1879) 199.—Hispan.
velata, *Pomel, Nouv. Mat. Fl. Atl.* 2.—Algeria.
velebica, *Borb.—Cf. in Oestr. Bot. Zeitschr.* xxxiii. (1883) 201.—Austria.
velutina, Desf. Fl. Atlant. i. 180. t. 51 = mollis.
venosa, Willd. Enum. Hort. Berol. Suppl. 10 = rhomboidalis.
venusta, Schur, Enum. Pl. Transs. 442 = caespitosa.
verbenacfolia, *Sieber, in Ann. Fl. Pom.* (1838) 375.—Hab.?
veronicifolia, *Hance, in Journ. Bot.* ix. (1871) 133.—China.
verruculosa, Hoffmgg. & Link, Fl. Port. ii. 12. t. 81 = C. Rapunculus.
versicolor, *Sibth. & Sm. Fl. Graec. Prod.* i. 138.—Italia; Graccia.
verticillata, Guss. Pl. Rar. 92 = versicolor.
verticillata, *Hill, Veg. Syst.* viii. 10.—Am. bor.
verticillata, Pall. Reise, iii. 719 = Adenophora verticillata.
vesula, All. Fl. Pedem. i. 108 = persicifolia.
Vidalii, *H. C. Wats. in Hook. Ic. Pl.* t. 684.—Ins. Azores.
vincaeflora, Vent. Jard. Malm. 12 = Wahlenbergia gracilis.
Violae, Pers. Syn. i. 192 = seq.
violaefolia, *Lam. Encyc.* i. 587.—Sibiria.
virgata, A. DC. Prod. vii. 483 = C. Rapunculus.
virgata, Labill. Ic. Pl. Syr. Dec. ii. 11. t. 6 = Phyteuma limonifolium.
virgata, Rafin. ex Roem. & Schult. Syst. v. 100 = divaricata, linifolia.
viridis, Spreng. Syst. iv. Cur. Post. 78 = Codonopsis viridis.
Vitinghofiana, Roem. & Schult. Syst. v. 102 = Steveni, persicifolia.
Vlachovae, Orphan. ex Boiss. Fl. Orient. iii. 927 = glomerata.
volubilis, Willd. ex Roem. & Schult. Syst. v. 98 = lactiflora.
vulgaris, *Gueldenst. It.* ii. 10, *ex Ledeb. Fl. Ross.* ii. 891.—Reg. Caucas.
Waldsteiniana, *Roem. & Schult. Syst.* v. 99.—Croatia.
Wanneri, Rochel, Pl. Banat. Rar. 41. t. 5. f. 12 = Symphyandra Wanneri.
Welandii, Heuff. in Oestr. Bot. Wochenschr. (1857) 118 = expansa.
Willdenowiana, Roem. & Schult. Syst. v. 107 = versicolor.
Willdenowii, Boiss. Diagn. Ser. II. iii. 117 = Phyteuma lobelioides.
Zeyheri, D. Dietr. Syn. Pl. i. 753 = Wahlenbergia Zeyheri.
Zoysii, *Wulf. in Jacq. Collect.* ii. 122.—Europ. austr.

CAMPANULOIDES, Hort. Kew. ex A. DC. Monog. Camp. 107 (1830) = **Lightfootia**, L'Hérit. (Campan.).
subulata, Hort. Kew. ex A. DC. l. c. = L. subulata.

CAMPANULOPSIS, Zoll. & Mor. in Nat. en Genecsk. Arch. Neérl. Indie, i. (1844) 484, nomen. *CAMPANULACEAE*?
cyanea, *Zoll. & Mor. l. c.*—Malaya.

CAMPANUMOEA, Blume, Bijdr. 726 (1825). *CAMPANULACEAE*, Benth. & Hook. f. ii. 557.
CYCLOCODON, Griff. Notul. iv. 277 (1854).
celebica, *Blume, l. c.* 727.—Ind. or.; Malaya; China.
cordata, Maxim. in Bull. Acad. Pétersb. xii. (1868) 68 = javanica.
inflata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 436.—Reg. Himal.
japonica, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 67.—Japon.
japonica, Siebold, ex Morr. Belg. Hortic. (1863) 337 = lanceolata?
javanica, *Blume, Bijdr.* 726.—Reg. Himal.; Java.
lanceolata, Sieb. & Zucc. Fl. Japon. i. 174. t. 91. = Codonopsis lanceolata.
leucocarpa, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 436 = celebica.
parviflora, *Benth. & Hook. f. Gen.* ii. 558.—Reg. Himal.
pilosula, *Franch. Pl. Davidian.* i. (1884) 192.—Mantchur.
truncata, Endl. Gen. Plant. i. 515; C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 436 = celebica.

CAMPBELLIA, Wight, Ic. iv. 5. t. 1424, 1425 (1850) = **Christisonia**, Gardn. (Orobanch.).
albida, Benth. & Hook. f. Gen. ii. 967 = Christisonia albida.
aurantiaca?, Wight, Ic. iv. t. 1424 = Chr. neilgherrica.
cytinoides, Wight, l. c. 1425 = Chr. neilgherrica.
Lawii, Benth. & Hook. f. Gen. ii. 967 = Chr. Lawii.
subacaulis, Benth. & Hook. f. l. c. = Chr. subacaulis.

CAMPDERIA, Benth. Bot. Voy. Sulph. 159. t. 52. (1844). *POLYGONACEAE*, Benth. & Hook. f. iii. 102.
floribunda, *Benth. l. c.* 160.—Guatemala.
gracilis, *Meissn. in Mart. Fl. Bras.* v. 1. 23.—Bras.
Lindeniana, *Benth. & Hook. f. Gen.* iii. 103.—Mexic.
mexicana, Meissn. in DC. Prod. xiv. 171 = floribunda.
nematostachya, *Griseb. in Bonplandia*, vi. (1858) 4.—Panama.

CAMPDERIA, Lag. Amen. Nat. ii. 99 (1821) = **Kundmannia**, Scop. (Umbellif.).
sicula, Lag. l. c. = K. sicula.

CAMPDERIA, A. Rich. in Bull. Soc. Philom. (1822) 79 = **Barbacenia**, Vand. (Amaryllid.).

CAMPE, Dulac, Fl. Hautes-Pyr. 199 (1867) = **Barbarea**, R. Br. (Crucifer.).
praecox, Dulac, l. c. = B. praecox.
vulgaris, Dulac, l. c = B. vulgaris.

CAMPECARPUS, H. Wendl. ex Benth. & Hook. f. Gen. iii. 893 (1883) = **Cyphophoenix**, H. Wendl. (Palm.).
fulcitus, H. Wendl. l. c. = Cyph. fulcita.

CAMPECIA, Adans. Fam. ii. 567 (1763) = **Caesalpinia**, Linn. (Legumin.).

CAMPELEPIS, Falc. in Trans. Linn. Soc. xix. (1845) 101 = **Periploca**, Linn. (Asclepiad.).
viminea, Falc. l. c. = P. aphylla.

CAMPELIA, Kunth, Enum. Pl. i. 286, 287 (1833) = CAMPELLA, Link.

CAMPELIA, Rich. Anal. Fruit. 46 (1808). *COMMELINACEAE*, Benth. & Hook. f. iii. 855.
GONATANDRA, Schlecht. in Linnaea, xxiv. (1851) 659.
SARCOPERIS, Rafin. Fl. Tellur. ii. 16 (1836).
ZANONIA, [Plum.] Cram. Diss. Syst. 75 (1803).
bibracteata, Wied-Neuwied, Beitr. Bras. 15. t. 2. f. 4 = C. Zanonia.
Boucheana, Schult. f. Syst. vii. 1180 = C. Zanonia.
fastigiata, Schlecht. in Linnaea, xxv. (1852) 188 = C. Zanonia.
Fendleri, Hassk. in Flora, xlviii. (1865) 102 = C. Zanonia.
glabrata, Hassk. in Miq. Pl. Jungh. 154 = Forrestia glabrata.
glabrata, Kunth, Enum. Pl. iv. 109 = C. Zanonia.
Hoffmanni, Hassk. in Flora, xlviii. (1865) 105 = C. Zanonia.
marginata, Blume, Enum. Pl. Jav. i. 7 = Forrestia marginata.
marginata, Wall. Cat. n. 8977 = Forrestia Hookeri.

CAMPELIA:—
mexicana, Mart. ex Kunth, Enum. Pl. iv. 109 = C. Zanonia.
mollissima, Blume, Enum. Pl. Jav. i. 7 = Forrestia mollis.
Pseudo-Zanonia, Kunth, Enum. Pl. iv. 108 = C. Zanonia.
scandens, *Hassk. in Flora*, xlviii. (1865) 103.—Peruv.
Zanonia, *H. B. & K. Nov. Gen. et Sp.* i. 264.—Am. austr.

CAMPELLA, Link, Hort. Berol. i. 122 (1827) = **Deschampsia**, Beauv. (Gramin.).
caespitosa, Link, l. c. 122 = D. caespitosa.
media, Link, l. c. 123 = D. juncea.

CAMPHORA, [Bauh.] Linn. Syst. ed. I (1735); Nees, Syst. Laurin. 87 = **Cinnamomum**, Tourn. (Laurin.).
angustifolia, Rafin. Sylva Tellur. 136.—Hab.?
chinensis, Nees, in Otto, Gartenz. (1833) 91; Syst. Laurin. 92 = Cinn. Parthenoxylon.
glandulifera, Nees, in Wall. Pl. As. Rar. ii. 72 = Cinn. glanduliferum.
glauca, G. Don, in Sweet, Hort. Brit. ed. III. 582 = Litsea glauca.
inodora, Blume, ex Miq. Fl. Ind. Bat. i. 904.—Arch. Sunda.
inuncta, Nees, in Wall. Pl. As. Rar. iii. 32 = Cinn. inunctum.
japonica, Rafin. Sylva Tellur. 136 = Cinn. pedunculatum?
officinalis, Steud. Nom. ed. II. i. 271 = Cinn. Camphora.
officinarum, Nees, ex Wall. Pl. As. Rar. ii. 72 = Cinn. Camphora.
Parthenoxylon, Nees, l. c. = Cinn. Parthenoxylon.
porrecta, Voigt, Hort. Suburb. Calc. 308 = Cinn. Parthenoxylon.
Pseudo-Sassafras, Miq. in Zoll. Syst. Verz. Ind. Archip. 113, 114 = Cinn. Parthenoxylon.
sumatrana, Miq. Pl. Jungh. 174.—Sumatra.
vera, Rafin. Sylva Tellur. 136 = Cinn. Camphora.

CAMPHORATA, Tourn. ex Crantz, Inst. i. 113 (1766) = **Camphorosma**, Linn. (Chenopod.).
acuta, Crantz, l. c. 114 = Halimocnemis juniperina.
glabra, Crantz, l. c. = Camphorosma glabrum.
hirsuta, Moench, Meth. 700 = Camphorosma monspeliacum.
monspeliensium, Crantz, Inst. i. 114 = Camphorosma monspeliacum.

CAMPHORINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 1 = **Cinnamomum**, Tourn.? (Laurin.).

CAMPHOROMOEA, Nees, Laurin. Expos. 16 (1833) = **Ocotea**, Aubl. (Laurin.).
divaricata, Nees, Syst. Laurin. 467.—Bras.
laxa, Nees, l. c. 468.—Bras.
litsaeifolia, Meissn. in DC. Prod. xv. 1. 145.—Bras.
ovalifolia, Meissn. l. c. 144.—Bras.
rhamnoides, Meissn. l. c. 145.—Bras.
subtriplinervia, Nees, Syst. Laurin. 466.—Bras.
surinamensis, Meissn. in DC. Prod. xv. 1. 145.—Guinea.
tenuiflora, Meissn. l. c. 144.—Bras.
venulosa, Nees, Syst. Laurin. 469.—Bras.
zizyphoides, Meissn. in DC. Prod. xv. 1. 145.—Bras.

CAMPHOROMYRTUS, Schau. in Linnaea, xvii. (1843) 240 = **Baeckia**, Linn. (Myrtac.).
Behrii, Schlecht. l. c. xx. (1847) 651 = B. Behrii.
Brownii, Schau. l. c. xvii. (1843) 240 = B. camphorata.
crenulata, F. Muell. in Trans. Vict. Inst. i. (1855) 123 = B. virgata.
pluriflora, F. Muell. l. c. = B. virgata.

CAMPHOROSMA, Linn. Diss. Dass. 8 (1747) (Amoen. Acad. i. 392). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 56.
CAMPHORATA, Tourn. ex Crantz, Inst. i. 113 (1766).
acutum, Falk, Beitr. ii. 120 = Nanophyton caspicum.
acutum, Linn. Sp. Pl. 122 = Halimocnemis juniperina.
acutum, Pollich, Fl. Palat. iii. 317 = Kochia arenaria.
annua, *Pall. Illustr. Pl.* 67. t. 58.—Rossia.
glabrum, Auct. ex DC. Prod. iii. 368 = Siphonychia americana.
glabrum, *Linn. Sp. Pl.* 122.—Helvetia.
hirsutum, Stokes, Bot. Mat. Med. i. 232 = monspeliacum.
monandrum, *Bunge, ex Boiss. Fl. Orient.* iv. 922.—Afghan.

CAMPHOROSMA:—
monspeliacum, *Linn. Sp. Pl.* 122.—Reg. Mediterr.
monspeliacum, Pall. Reise, ii. 471 = ruthenicum.
monspeliacum, Pollich, Fl. Palat. i. 165 = Kochia arenaria.
monspeliense, Murr. Syst. Veg. ed. XIII. 139 = monspeliacum.
ovatum, Biasol. ex Moq. in DC. Prod. xiii. II. 126 = monspeliacum.
ovatum, *Waldst. & Kit. Pl. Rar. Hung.* i. 64. t. 63.—Europ. austr. or.
ovatum, Bong. & Mey. Suppl. ii. Fl. Alt. 68. n. 274 = annuum.
paleaceum, Linn. f. Suppl. 128 = Monetia paleacea.
perennis, *Pall. Illustr. Pl.* iii. 66. t. 57.—Reg. Caucas.; Persia.
polygamum, *Bunge, ex Boiss. Fl. Orient.* iv. 921.—Afghan.
Pteranthus, Linn. Mant. i. 41 = Pteranthus echinatus.
ruthenicum, *Bieb. Fl. Taur. Cauc.* iii. *Suppl.* 112.—Europ. austr. or.; As. bor.
sabulosa, *Kit. ex Moq. in DC. Prod.* xiii. II. 126.—Hungar.
songoricum, *Bunge, in Act. Hort. Petrop.* vi. (1879) 415.—Sibir.
vingintimilleis, *Tineo, Cat. Hort. Panorm.* 277.—Sicil.

CAMPHUSIA, De Vriese, in Nederl. Kruidk. Arch. ii. (1850) 148 = **Scaevola**, Linn. (Goodeniac.).
glabra, De Vriese, l. c. 149.—Malaya.

CAMPHYLEIA, Spreng. Gen. ii. 486 (1831) = **Striga**, Lour. (Scrophulac.).

CAMPIA, 'Domb.' ex Endl. Ench. 91, in syn. (1841) = CAPIA, Juss. = **Lapageria**, Ruiz & Pav. (Liliac.).

CAMPILOSTACHYS, A. Juss. in Orb. Dict. xii. 38 (1849) = **Campylostachys**, Kunth (Verbenac.).

CAMPNOSPERMA, Thw. in Hook. Kew Journ. vi. (1854) 65. t. 1. *ANACARDIACEAE*, Benth. & Hook. f. i. 425.
auriculata, *Hook. f. Fl. Brit. Ind.* ii. 41.—Ind. or.
Griffithii, *March. Rev. Anac.* 174.—Penang.
gummifera, *March. l. c.* 173.—Bras.
macrophylla, *Hook. f. Fl. Brit. Ind.* ii. 41.—Malacca.
Micranteia, *March. Rev. Anac.* 173.—Madag.
oxyrhachis, *Engl. in A. DC. Monog. Phan.* iv. 319.—Sumatra.
seychellarum, *March. Rev. Anac.* 173.—Ins. Mascar.
zeylanica, *Thw. in Hook. Kew Journ.* vi. (1854) 65.—Zeylan.

CAMPOMANESIA, Ruiz & Pav. Prod. 72. t. 13 (1797). *MYRTACEAE*, Benth. & Hook. f. i. 712.
ABBEVILLEA, ACRANDRA, BRITOA, LACERDAEA, Berg, in Linnaea, xxvii. (1854) 425–437.
adamantium, *Blume, Mus. Bot. Lugd. Bat.* i. 73.—Bras.
affinis, *Berg, in Mart. Fl. Bras.* xiv. I. 446.—Bras.
aprica, *Berg, l. c.* 459.—Bras.
arenaria, *Berg, l. c.* 448.—Reg. Argent.
aromatica, *Griseb. Fl. Brit. W. Ind.* 242.—Ind. occ.
aurea, *Berg, in Mart. Fl. Bras.* xiv. I. 454.—Bras.
australis, *Berg, l. c.* 445.—Bras.
caerulea, *Berg, l. c.* 455.—Bras.
caerulescens, *Berg, l. c.* 612.—Bras.
Cambessedeana, *Berg, l. c.* 457.—Bras.
campestris, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
cerasoides, *A. Gray, Bot. U. St. Expl. Exped.* i. 549.—Bras.
ciliata, *Berg, in Mart. Fl. Bras.* xiv. I. 453.—Bras.
coaetanea, *Berg, l. c.* 444.—Am. trop.
cornifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 150. t. 147.—N. Granat.
corymbosa, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
crassifolia, *Benth. Bot. Voy. Sulph.* 97. t. 37.—N. Granat.
crenata, *Berg, in Mart. Fl. Bras.* xiv. I. 456.—Bras.
cuneata, *Berg, l. c.* 446.—Bras.
cyanea, *Berg, l. c.* 454.—Bras.
dentata, *Berg, l. c.* 457.—Bras.
desertorum, *Berg, l. c.* 450.—Bras.
dimorpha, *Berg, l. c.* 609.—Bras.
discolor, *Berg, l. c.* 439.—Bras.

CAMPOMANESIA:—
eriantha, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
eugenioides, *Blume, l. c.*—Bras.
fruticosa, *Berg, in Mart. Fl. Bras.* xiv. I. 458.—Bras.
fusca, *Berg, l. c.* 447.—Bras.
Gardneriana, *Berg, l. c.* 455.—Bras.
glabra, Benth. in Hook. Journ. Bot. ii. (1843) 319 = Calycolpus glaber.
glabra, *Berg, in Mart. Fl. Bras.* xiv. I. 450.—Bras.
Goetheana, Hemsl. Biol. Centr. Am. Bot. i. 407 = Calycolpus glaber.
grandiflora, *Sagot, in Ann. Sc. Nat.* Sér. VI. xx. (1885) 182.—Guiana.
Guaviroba, *Benth. & Hook. f. Gen.* i. 712.—Bras.
guazumaefolia, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
heterophylla, *Berg, in Mart. Fl. Bras.* xiv. I. 440.—Bras.
hirsuta, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 353.—Bras.
Houlletii, *Berg, in Mart. Fl. Bras.* xiv. I. 449.—Bras.
lanceolata, *Berg, l. c.* 612.—Bras.
Langsdorffii, *Berg, l. c.* 610.—Bras.
laurifolia, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 353.—Bras.
lineatifolia, *Ruiz & Pav. Fl. Per.* iv. t. 422; *Syst.* 128.—Peruv.
malifolia, *Berg, in Mart. Fl. Bras.* xiv. I. 452.—Bras.
Martiana, *Berg, l. c.* 610.—Bras.
mediterranea, *Berg, l. c.* 458.—Bras.
microcarpa, *Berg, l. c.* 455.—Bras.
multiflora, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
obscura, *Berg, in Mart. Fl. Bras.* xiv. I. 451.—Bras.
obversa, *Berg, l. c.* 445.—Bras.
ovalifolia, *Berg, l. c.* 452.—Bras.
Pohliana, *Berg, l. c.* 441.—Bras.
Poiteaui, *Berg, in Linnaea*, xxvii. (1854) 432.—Bras.
pubescens, *Berg, in Mart. Fl. Bras.* xiv. I. 443.—Bras.
racemosa, *Berg, l. c.* 458.—Bras.
repanda, *Berg, l. c.* 456.—Bras.
reticulata, *Berg, l. c.* 439.—Bras.
rhombea, *Berg, l. c.* 453.—Bras.
rhytidophylla, *Berg, l. c.* 608.—Bras.
Riedeliana, *Berg, l. c.* 610.—Bras.
rugosa, *Berg, l. c.* 439.—Bras.
salviaefolia, *Berg, l. c.* 442.—Bras.
suaveolens, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
suffruticosa, *Berg, in Mart. Fl. Bras.* xiv. I. 448.—Bras.
synchrona, *Berg, l. c.* 444.—Bras.
tenuifolia, *Berg, l. c.* 452.—Bras.
tomentosa, *H. B. & K. Nov. Gen. et Sp.* vi. 151. t. 147.—N. Granat.
transalpina, *Berg, in Mart. Fl. Bras.* xiv. I. 459.—Bras.
triflora, *Baill. Hist. Pl.* vi. 342.—Bras.
vaccinioides, *Berg, in Mart. Fl. Bras.* xiv. I. 450.—Bras.
velutina, *Blume, Mus. Bot. Lugd. Bat.* i. 72.—Bras.
virescens, *Berg, in Mart. Fl. Bras.* xiv. I. 611.—Bras.
Widgreniana, *Berg, l. c.* 447.—Bras.
xanthocarpa, *Berg, l. c.* 451.—Bras.

CAMPSANTHUS, Steud. Nom. ed. II. i. 272 (1840), sphalm. = COMPSANTHUS, Spreng. = **Tricyrtis**, Wall. (Liliac.).

CAMPSIANDRA, Benth. in Hook. Journ. Bot. ii. (1840) 93. *LEGUMINOSAE*, Benth. & Hook. f. i. 563.
angustifolia, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 55.—Am. trop.
comosa, *Benth. in Hook. Journ. Bot.* ii. (1840) 93.—Guiana.
laurifolia, *Benth. l. c.* 94.—Bras.
rosea, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 62.—Bras.

CAMPSIDIUM, Seem. in Bonplandia, x. (1862) 147; in Ann. & Mag. Nat. Hist. Ser. III. x. (1862) 32 = **Tecoma**, Juss. (Bignon.).
chilense, Reiss. & Seem. in Bonplandia, x. (1862) 147.—Chili.
filicifolium, A. Van Geert. Cat. n. 71 (1874); Fl. et Pom. (1874) 280.—Ins. Fiji.

CAMPSIS, Lour. Fl. Cochinch. 377 (1790) = **Tecoma** Juss. (Bignon.).
adrepens, Lour. l. c. = Incarvillea sinensis.
amboinensis, Seem. in Journ. Bot. v. (1867) 374 = T. amboinensis.
Curtisii, Seem. l. c. 371 = T. radicans?
dendrophila, Seem. l. c. 373 = T. dendrophila.
Fortunei, Seem. l. c.—China.
Haussmannii, F. Muell. Fragm. iv. 148 = Haussmannia jucunda.
radicans, Seem. in Journ. Bot. v. (1867) 372 = T. radicans.

CAMPTEDERIA, Steud. Nom. ed. II. i. 272 (1840), sphalm. = CAMPDERIA, A. Rich.

CAMPTOCARPUS, Decne. in DC. Prod. viii. (1844) 493. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 744.
Bojerianus, *Decne. l. c.* 494.—Madag.
crassifolius, *Decne. l. c.*—Madag.
linearis, *Decne. l. c.*—Madag.
mauritianus, *Decne. l. c.* 493.—Ins. Borbon.

CAMPTOCARPUS, C. Koch, in Linnaea, xvii. (1842) 304 = **Alkanna**, Tausch (Boragin.).
orientalis, C. Koch, l. c. = A. orientalis.

CAMPTOLOMA, Benth. in DC. Prod. x. 430 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 960.
rotundifolia, *Benth. l. c.* 431.—Afr. austr.
villosa, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 84.—Ins. Socotra.

CAMPTOPUS, Hook. f. Bot. Mag. t. 5755 (1869) = **Cephaëlis**, Sw. (Rubiac.).
Mannii, Hook. f. l. c. = Cephaëlis Mannii.

CAMPTOSEMA, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 200. *LEGUMINOSAE*, Benth. & Hook. f. i. 536.
BIONIA, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 130.
acuminatum, *Benth. in Mart. Fl. Bras.* xv. I. 155.—Am. austr.
bellum, *Benth. l. c.* 156.—Am. austr.
coccineum, *Benth. l. c.* 154.—Am. austr.
coriaceum, *Benth. l. c.* 155.—Am. austr.
erythrinoides, *Benth. l. c.* 156.—Am. austr.
grandiflorum, *Benth. in Linnaea*, xxii. (1849) 516.—Bras.
pedicellatum, *Benth. in Hook. Journ. Bot.* ii. (1840) 60.—Guiana.
pinnatum, *Benth. in Mart. Fl. Bras.* xv. I. 325.—Bras.
rubicundum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 201.—Reg. Argent.
tomentosum, *Benth. in Mart. Fl. Bras.* xv. I. 155.—Am. austr.

CAMPTOSTEMON, Mast. in Hook. Ic. Pl. xii. 18. t. 1119 (1872). *MALVACEAE*.
Schultzii, *Mast. l. c.*—Austral.

CAMPTOTHECA, Decne. in Bull. Soc. Bot. Fr. xx. (1873) 157. *CORNACEAE*.
acuminata, *Decne. l. c.*—Tibet.

CAMPULEIA, Thou. Gen. Nov. Madag. 7 (1806) = **Striga**, Lour. (Scrophularin.).
coccinea, Hook. Exot. Fl. 3. t. 203 = S. lutea.
coerulea, Boj. ex Benth. in DC. Prod. x. 501 = S. orobanchioides.
rosea, Boj. ex Benth. l. c. = S. orobanchioides.

CAMPULOA, Desv. Journ. Bot. i. (1813) 69 = **Ctenium**, Panz. (Gramin.).
brachystachya, Nees, ex Doell, in Mart. Fl. Bras. ii. III. 73 = Ct. brachystachyum.
chapadensis, Trin. ex Doell, l. c. 74 = Ct. chapadense.
elegans, J. Gay, ex Kunth, Rev. Gram. i. 93 = Ct. elegans.
gracilis, Desv. Journ. Bot. i. (1813) 69 = Ct. carolinianum.
hirsuta, Desv. l. c. (= *Campulosus falcatus*).—Afr. austr.
monostachya, Roem. & Schult. Syst. ii. 516 = Ct. carolinianum.

CAMPULOCLINIUM, DC. Prod. v. 136 (1836) = **Eupatorium**, Tourn. (Compos.).
alternifolium, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 438 = Ageratum alternifolium.
arenarium, Gardn. l. c. v. (1846) 467.—Bras.
ascendens, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 363 = E. betonicaeforme.
corymbosum, Sch. Bip. ex Baker, l. c. 356 = E. Burchellii.
crenatum, Schlecht. in Mart. Herb. Fl. Bras. n. 668 = E. sordescens.
decumbens, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 344 = E. decumbens.
frutescens, Bartl. Ind. Sem. Hort. Goett. (1843).—Hab.?
hirsutum, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 438 = E. hirsutum.
kleinioides, DC. Prod. v. 137 = E. kleinioides.
macrocephalum, DC. l. c. = E. macrocephalum.
palustre, DC. in Deless. Ic. Sel. iv. t. 16 = E. palustre.
Pohlianum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 358 = E. macrocephalum.
polyphyllum, Sch. Bip. ex Baker, l. c. 354 = E. megacephalum.
siegesbeckioides, Sch. Bip. ex Baker, l. c. 359 = E. hirsutum.
strigosum, Sch. Bip. ex Baker, l. c. 358 = E. macrocephalum.
subpaniculatum, DC. Prod. v. 137 = E. sordescens.
surinamense, Miq. in Linnaea, xvii. (1843) 69 = E. Vitalbae.
turbinatum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 357 = E. chlorolepis.
urticaefolium, DC. Prod. v. 137 = E. urticaefolium.
viridiflorum, Bartl. ex Baker, in Mart. Fl. Bras. vi. II. 309 = E. viridiflorum.

CAMPULOSUS, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 = **Ctenium**, Panz. (1814) (Gramin.).
aciculatus, Trin. Sp. Gram. iii. t. 301 = Ct. elegans.
aromaticus, Trin. ex Steud. Nom. ed. II. i. 272 = Ct. carolinianum.
brachystachyus, Nees, Agrost. Bras. 417 = Ct. brachystachyum.
chapadensis, Trin. Sp. Gram. iii. 26. t. 303 = Ct. chapadense.
cirrhosus, Nees, Agrost. Bras. 416 = Ct. cirrhosum.
concinnus, Steud. Nom. ed. II. i. 272 = Ct. concinnum.
dolichostachys, Trin. ex Steud. l. c.—Hab.?
falcatus, Beauv. Agrost. 64.—Afr. austr.
gracilior, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 = Ct. carolinianum.
gracilis, Bertol. in Mem. Acad. Sc. Bolog. ii. (1850) 602.—Am. bor.
hirsutus, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 = *falcatus*.
macrostachys, Steud. Nom. ed. II. i. 29 = Ct. carolinianum.
monostachyos, Beauv. Agrost. 64. t. 13. f. 1 = Ct. carolinianum.
planifolius, Presl, Reliq. Haenk. i. 287 = Ct. planifolium.
tricorys, Trin. ex Steud. Nom. ed. II. i. 272.—Hab.?

CAMPYDORUM, Salisb. Gen. Pl. Fragm. 64 (1866) = **Polygonatum**, Adans. (Liliac.).
verticillatum, Salisb. l. c. = P. verticillatum.

CAMPYLANDRA, Baker, in Journ. Linn. Soc. xiv. (1875) 582. t. 20 = **Tupistra**, Ker-Gawl. (Liliac.).
aurantiaca, Baker, l. c. 582 = T. aurantiaca.
dasystachys, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 256 = Dasystachys campanulata.

CAMPYLANTHERA, Hook. Ic. Pl. t. 82 (1837) = **Pronaya**, Hueg. (Pittospor.).
ericoides, Lindl. in Mitch. Three Exped. ii. 277 = Marianthus procumbens.
Fraseri, Hook. Ic. Pl. t. 82 = P. elegans.

CAMPYLANTHERA, Schott, Melet. 35 (1832) = **Eriodendron**, DC. (Malvac.).
Samauma, Schott, l. c. = E. Samauma.

CAMPYLANTHUS, Roth, Nov. Pl. Sp. 4 (1821). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 961.
Benthamii, *Webb, in Hook. Niger Fl.* 163.—Ins. Cap. Verd.

CAMPYLANTHUS:—
glabra, *Benth. in DC. Prod.* x. 508.—Ins. Cap. Verd.
junceus, *Edgew. in Journ. As. Soc. Beng.* xvi. (1847) 1217.—Ind. or.
ramosissimus, *Wight, Ic.* iv. t. 1416.—Ind. or.; Beluchist.
salsoloides, *Roth, Nov. Pl. Sp.* 4.—Ins. Teneriff.
salsoloides, Stocks, ex Boiss. Fl. Orient. iv. 433 = ramosissimus.
salsoloides, Webb, Phyt. Canar. iii. 126 = Benthamii.
spinosus, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 84.—Ins. Socotra.

CAMPYLEIA, Spreng. Syst. iv. Cur. Post. 222 (1831) = CAMPULEIA, Thou. = **Striga**, Lour. (Scrophular.).

CAMPYLIA, Lindl. ex Sweet, Geran. sub t. 43 (1820) = **Pelargonium**, L'Hérit. (Geraniac.).
blattaria, Sweet, Geran. t. 88 = P. ovale.
cana, Colv. ex Sweet, l. c. t. 114 = P. ovale.
capillaris, Eckl. & Zeyh. Enum. 74 = P. capillare.
carinata, Lindl. ex Sweet, Geran. sub t. 43 = P. carinatum.
cartilaginea, Eckl. & Zeyh. Enum. 74 = P. cartilagineum.
coronopifolia, Sweet, Hort. Brit. ed. I. 75 = P. coronopifolium.
elata, Eckl. & Zeyh. Enum. 74 = P. elatum.
elegans, Colv. ex Sweet, Geran. t. 222 = P. elegans.
eriostemon, Eckl. & Zeyh. Enum. 73 = P. ovale.
holosericea, Sweet, Geran. t. 75 = P. ovale.
laciniata, Sweet, l. c. t. [401] = P. laciniatum.
Oenotherae, Sweet, Hort. Brit. ed. I. 75 = P. Oenotherae.
staticephylla, Eckl. & Zeyh. Enum. 74 = P. staticaefolium.
tricolor, Eckl. & Zeyh. l. c. = P. violaceum.
variegata, Sweet, Geran. t. 266 = P. marginatum.
verbasciflora, Sweet, l. c. ii. t. 157 = P. ovale.
veronicaefolia, Eckl. & Zeyh. Enum. 73 = P. veronicaefolium.

CAMPYLOBOTRYS, Lem. in Fl. des Serres, Sér. I. iii. (1847) sub t. 260 = **Hoffmannia**, Sw. (Rubiac.).
argyroneura, Linden, in Belg. Hortic. viii. (1858) 166.—Mexic.
discolor, Hook. Bot. Mag. t. 4530 = seq.?
discolor, Lem. in Fl. des Serres, Sér. I. iii. (1847) sub t. 260 = H. discolor.
Ghiesbreghtii, Lem. Illustr. Hortic. viii. (1861) t. 279 = H. Ghiesbreghtii.
pyrophylla, Linden, Cat. n. 16 (1861) 2.—Mexic.
regalis, Linden, l. c. n. 15 (1860) = H. regalis.

CAMPYLOCARYUM, DC. ex A. DC. in Prod. x. 97, in syn. (1846) = **Alkanna**, Tausch (Boragin.).
smyrnaeum, Boiss. ex A. DC. Prod. x. 98 = A. Pinardi.
syriacum, Boiss. Fl. Orient. iv. 226 = A. syriaca.
syriacum, Boiss. & Hohen. ex A. DC. Prod. x. 101 = A. strigosa.
verrucosum, Boiss. ex A. DC. l. c. 102 = A. incana.

CAMPYLOCENTRON, Benth. in Journ. Linn. Soc. xviii. (1881) 337. *ORCHIDEAE*, Benth. & Hook. f. iii. 585.
TODAROA, A. Rich. in Ann. Sc. Nat. Sér. III. iii. (1845) 28.
Schiedei, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* iii. 292.—Mexic.

CAMPYLOCERA, Nutt. in Trans. Am. Phil. Soc. Ser. II. viii. (1843) 257 = **Specularia**, Heist. (Campanul.).
leptocarpa, Nutt. l. c. = S. leptocarpa.

CAMPYLOCHINIUM, Endl. Gen. 369 (1838) = CAMPULOCHINIUM, DC. = **Eupatorium**, Tourn. (Compos.).

CAMPYLONEMA, Poir. Illustr. 598 (1823); Schult. f. Syst. vii. 96, 1507 (1830) = **Campynema**, Labill. (Amaryllid.).
lineare, Schult. f. l. c. = Campynema lineare.

CAMPYLOPTERA, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1841) 194 = **Aethionema**, R. Br. (Crucif.).
heterocarpa, Baill. Hist. Pl. iii. 291 = Ae. campylopterum.
syriaca, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1841) 194 = Ae. campylopterum.

CAMPYLOPUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 360 = **Hypericum**, Tourn.
cerastioides, Spach, Hist. Vég. Phan. v. 412 = H. rhodopeum.

CAMPYLOSIPHON, Benth. in Hook. Ic. Pl. t. 1384 (1882). *BURMANNIACEAE*, Benth. & Hook. f. iii. 458.
purpurascens, *Benth. l. c.*—Am. austr.

CAMPYLOSIPHON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 135 = **Siphocampylus**, Pohl (Campanul.).
lycioideus, St. Lag. l. c. = S. lycioides.

CAMPYLOSPORUS, Spach, in Ann. Sc. Nat. Sér. II. v. (1836) 363 = **Hypericum**, Tourn.
angustifolius, Spach, Hist. Vég. Phan. v. 425 = H. lanceolatum.
madagascariensis, Spach, l. c. 426 = H. madagascariense.
reticulatus, Spach, l. c. 424 = H. lanceolatum.

CAMPYLOSTACHYS, Kunth, in Abh. Akad. Berl. (1831) 206. *VERBENACEAE*, Benth. & Hook. f. ii. 1137.
CAMPILOSTACHYS, A. Juss. in Orb. Dict. xii. 38 (1849).
abbreviata, *E. Mey. Comm. Pl. Afr. Austr.* i. 279.—Afr. austr.
cernua, *Kunth, in Abh. Akad. Berl.* (1831) 201.—Afr. austr.
cernua, Zeyh. ex DC. Prod. xii. 607 = Stilbe pinastra.
phylicoides, Sond. in Linnaea, xix. (1847) 202 = Stilbe phylicoides.

CAMPYLOSTACHYS, E. Mey. in Drège, Zwei Pfl. Docum. 83 (1843) = **Fimbristylis**, Vahl (Cyperac.).
abbreviata, E. Mey. l. c. = F. ferruginea.

CAMPYLOSTEMON, E. Mey. in Drège, Zwei Pfl. Docum. 170 (1843), nomen = **Justicia**, Linn. (Acanthac.).
campanulatus, E. Mey. l. c. (= *Leptostachya Campylostemon*).—Afr. austr.
paniculatus, Nees, in DC. Prod. ix. 378 = praec.

CAMPYLOSTEMON, Welw. in Benth. & Hook. f. Gen. i. 998 (1862). *CELASTRINEAE*, Benth. & Hook. f. l. c.
angolense, *Welw. ex Oliver, in Journ. Linn. Soc.* x. (1869) 44.—Afr. trop.

CAMPYLOTHECA, Cass. in Dict. Sc. Nat. li. 476 (1827) = **Coreopsis**, Linn. (Compos.).
angustifolia, Walp. Rep. ii. 618 = Bidens angustifolia.
australis, Less. in Linnaea, vi. (1831) 509 = Cor. micrantha.
gracilis, Walp. Rep. ii. 618 = Bidens gracilis.
grandiflora, DC. Prod. v. 593.—Ins. Sandvic.
micrantha, Cass. in Dict. Sc. Nat. lix. 321 = Cor. micrantha.
mutica, Walp. Rep. ii. 618 = Bidens mutica.

CAMPYLOTROPIS, Bunge, Pl. Mongholico-Chin. 6 (1835) = **Lespedeza**, Michx. (Legumin.).
chinensis, Bunge, l. c. 7 = L. macrocarpa.
cytisoides, Benth. in Pl. Jungh. i. 230.—Java.
macrostyla, Lindl. ex Miq. Fl. Ind. Bat. i. 230 = L. criocarpa.
virgata, Miq. l. c. 230 = L. virgata.

CAMPYLUS, Lour. Fl. Cochinch. 113 (1790) (Genus dub.).
sinensis, *Lour. l. c.*—China.

CAMPYNEMA, Labill. Nov. Holl. Pl. i. 93. t. 121 (1804). *AMARYLLIDEAE*, Benth. & Hook. f. Gen. iii. 716.
CAMPYLONEMA, Poir. Illustr. 598 (1823); Schult. f. Syst. vii. 96, 1507 (1830).
lineare, *Labill. l. c.*—Ins. Tasman.
pygmaeum, *F. Muell. ex Benth. Fl. Austral.* vi. 415.—Ins. Tasman.; Austral.

CAMUNIUM, Adans. Fam. ii. 166 (1763) = **Aglaia**, Lour. (Meliac.).
bengalense, Buch.-Ham. ex Wall. Cat. n. 1279 = A. edulis.
chinense, Roxb. Hort. Beng. 18; Fl. Ind. i. 636 = A. odorata.

CAMUTIA, Bonat. ex Steud. Nom. ed. II. i. 273 (1840) = **Melampodium**, Linn. (Compos.).
perfoliata, Bonat. ex Steud. l. c. = M. perfoliatum.

CANABIS, Roth, Tent. Fl. Germ. i. 422 (1788), sphalm. = **Cannabis**, Tourn. (Urtic.).

CANAHIA, Steud. Nom. ed. I. 146 (1821) = **Kanahia**, R. Br. (Asclepiad.).
laniflora, Steud. l. c. = K. laniflora.

CANALA, Pohl, Pl. Bras. Ic. ii. 62. t. 142 (1831) = **Spigelia**, Linn. (Loganiac.).
heliotropioides, Pohl, l. c. 63. t. 142 = S. Olfersiana.
macrophylla, Pohl, l. c. 64 = S. macrophylla.
rubiaefolia, Pohl, l. c. 65 = S. Schlechtendaliana.

CANALIA, F. W. Schmidt, Neue u. Selt. Pfl. 30 (1793) = **Gnidia**, Linn. (Thymel.).
daphnoides, F. W. Schmidt, l. c. = G. pinifolia.

CANANGA, Aubl. Pl. Gui. i. 607. t. 244 (1775) = **Guatteria**, Ruiz & Pav. (1794) (Anonac.).
Candolleana, Warm. in Kjoeb. Vidensk. Meddel. (1873) 145 = G. Candolleana.
Klotzschiana, Warm. l. c. = G. Klotzschiana.
lancea, Poit. ex DC. Syst. i. 506 = G. virgata.
Ouregon, Aubl. Pl. Gui. i. 608 = G. Ouregon.
pseudolancea, Poit. ex DC. Syst. i. 507 = G. laurifolia.
Schomburgkiana, Baill. Hist. Pl. i. 204 = G. Schomburgkiana.
Sellowiana, Warm. in Kjoeb. Vidensk. Meddel. (1873) 144 = G. Sellowiana.
villosissima, Warm. l. c. = G. villosissima.

CANANGA, Rumph. ex Hook. f. & Thoms. Fl. Ind. i. 129 (1855). *ANONACEAE*, Benth. & Hook. f. i. 24.
monosperma, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 57.—Malacca.
odorata, *Hook. f. & Thoms. Fl. Ind.* i. 130.—Burma; Malaya.
virgata, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 57.—Malaya.

CANARIA, Linn. Mant. ii. 148 (1767) = **Canarina**, Linn. (Campanul.).
campanulata, Linn. l. c. = Canarina campanulata.

CANARINA, Linn. Mant. 588 (1767). *CAMPANULACEAE*, Benth. & Hook. f. ii. 558.
CANARIA, Linn. Mant. ii. 148 (1767).
PERNETYA, Scop. Introd. 150 (1777).
Campanula, Lam. Encyc. i. 598 = seq.
campanulata, *Linn. Mant.* ii. 225.—Ins. Canar.
laevigata, G. Don, ex Loud. Hort. Brit. 139 = campanulata.
moluccana, *Roxb. Hort. Beng.* 87, nomen.—Ins. Molucc.
Zanguebar, *Lour. Fl. Cochinch.* 195.—Afr. trop.

CANARION, St. Lag. in Ann. Soc. Bot. Lyon, vii. 121 (1880) = **Canarium**, Rumph. (Burserac.).
pimelum, St. Lag. l. c. = Canarium Pimela.

CANARIOPSIS, Miq. Fl. Ind. Bat. i. II. 651 (1859) = **Canarium**, Rumph. (Burserac.).
altissima, Miq. l. c. = Canarium altissimum.
angustifolia, Miq. l. c. 653 = Canarium angustifolium.
aspera, Miq. l. c. 653 = Canarium asperum.
decumana, Miq. l. c. = Canarium decumanum.
denticulata, Miq. l. c. 654 = Canarium denticulatum.
glabra, Miq. l. c. 652 = Canarium balsamiferum.
hirsuta, Miq. l. c. 653 = Canarium hirsutum.
hispida, Miq. l. c. = Canarium hispidum.
paucijuga, Miq. l. c.—Ins. Molucc.
villosa, Miq. l. c. 652 = Canarinm villosum.

CANARIUM, [Rumph.] Linn. Herb. Amboin. 9 (1754) (Amoen. Acad. iv. 121); Mant. 127 (1767). *BURSERACEAE*, Benth. & Hook. f. i. 324.
CANARIOPSIS, Miq. Fl. Ind. Bat. i. II. 651 (1859).
CENARIUM, Linn. Amoen. Acad. iv. 121, in nota (1759).
COLOPHONIA, Comm. ex Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 352.
MEHENBETHENE, Besl. ex Gaertn. Fruct. ii. 98 (1788); Walp. Ann. ii. 291 (1851).

CANARIUM:—
PACHYLOBUS, G. Don, Gen. Syst. ii. 89 (1832).
PIMELA, Lour. Fl. Cochinch. 407 (1790).
SCUTINANTHE, Thwaites, in Hook. Kew Journ. viii. (1856) 267. t. 8.
acutum, *Engl. in DC. Monog. Phan.* iv. 113.—Borneo.
album, Blanco, Fl. Filip. ed. I. 793 = luzonicum.
album, *Raeusch. Nom.* ed. III. 287; *DC. Prod.* ii. 80.—Cochinch.
altissimum, *Blume, Bijdr.* 1163.—Java.
altissimum, Korth. ex Blume, Mus. Bot. Lugd. Bat. i. 211 = Santiria laevigata.
angustifolium, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 117.—N. Guin.
asperum, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 215.—N. Guin.
australianum, *F. Muell. Fragm.* iii. 15.—Austral.
Balansae, *Engl. in DC. Monog. Phan.* iv. 117.—N. Caled.
balsamiferum, Moon, Cat. 68 = zeylanicum.
balsamiferum, *Willd. Sp. Pl.* iv. 760.—Ins. Molucc.
Beccarii, *Engl. in DC. Monog. Phan.* iv. 107.—Borneo.
bengalense, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 136.—Ind. or.
Bennettii, *Engl. in DC. Monog. Phan.* iv. 119.—Burma; Malacca.
Boivini, *Engl. l. c.* 110.—Madag.
brunneum, *Bedd. Fl. Sylv.* t. 127.—Zeylan.
coccineo-bracteatum, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 296.—Ins. Andaman.
Colophania, Baker, Fl. Maurit. 44 = mauritianum.
commune, Blanco, Fl. Filip. ed. I. 791 = Cumingii.
commune, *Linn. Mant.* i. 127.—Ins. Molucc.
commune, Wight & Arn. Prod. 175 = zeylanicum.
Cumingii, *Engl. in DC. Monog. Phan.* iv. 132.—Ins. Philipp.
decumanum, *Gaertn. Fruct.* ii. 99. t. 102.—Ins. Molucc.
denticulatum, *Blume, Bijdr.* 1162.—Java.
dichotomum, *Miq. Fl. Ind. Bat.* i. II. 648.—Sumatra.
divergens, *Engl. in DC. Monog. Phan.* iv. 143.—Borneo.
edule, *Hook. f. in Hook. Niger Fl.* 285.—Afr. trop.
euphyllum *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 295.—Ins. Andaman.
eupteron, *Miq. Fl. Ind. Bat.* i. II. 648.—Sumatra.
fissistipulum, *Miq. l. c. Suppl.* 525.—Sumatra.
fragile, *Engl. in DC. Monog. Phan.* iv. 138.—Borneo.
fuscum, *Engl. l. c.* 116.—Java.
giganteum, *Engl. l. c.* 106.—Borneo.
gilvescens, Miq. Fl. Ind. Bat. i. II. 650 = Santiria rostrata.
glaucum, *Blume, Mus. Bot. Lugd. Bat.* i. 219.—Sumatra.
gracile, *Engl. in DC. Monog. Phan.* iv. 140.—Ins. Philipp.
grandiflorum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 533.—Malacca.
Harami, *Boj. Hort. Maurit.* 83.—Madag.
Harveyi, *Seem. Fl. Vit.* 35.—Ins. Tonga.
hirsutum, *Willd. Sp. Pl.* iv. 760.—Ins. Molucc.
hirtellum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 534.—Penang.
hispidum, *Blume, Bijdr.* 1163.—Java.
Horami, Engl. in DC. Monog. Phan. iv. 150 = Harami.
incurvatum, *Engl. l. c.* 138.—Borneo.
Kadondon, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 535.—Malacca.
Kipella, *Miq. Fl. Ind. Bat.* i. II. 646.—Java.
Kitenga, *Miq. l. c.* 647.—Java.
Korthalsii, Miq. l. c. 645 = Santiria tomentosa.
laevigatum, Miq. l. c. 648 = Santiria laevigata.
laxiflorum, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 477 = microcarpum.
laxiflorum, *Zipp. ex Blume, Mus. Bot. Lugd. Bat.* i. 221.—Ins. Molucc.
laxum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 535.—Malacca.
legitimum, *Miq. Fl. Ind. Bat.* i. II. 647.—Ins. Amboina.
littorale, *Blume, Bijdr.* 1164.—Java.
longiflorum, *Zipp. ex Miq. Fl. Ind. Bat.* i. II. 647.—Ins. Molucc.
luxurians, *Engl. in DC. Monog. Phan.* iv. 146.—Ins. Philipp.
luzonicnm, *Miq. Fl. Ind. Bat.* i. II. 651.—Ins. Philipp.

CANARIUM:—
macrophyllum, *Oliver, Fl. Trop. Afr.* i. 327.—Afr. trop.
madagascariense, *Engl. in DC. Monog. Phan.* iv. 111.—Madag.
Mahassan, *Miq. Fl. Ind. Bat. Suppl.* 526.—Sumatra.
mauritianum, *Blume, Mus. Bot. Lugd. Bat.* i. 217.—Ins. Maurit.
Mehenbethene, *Gaertn. Fruct.* ii. 98. t. 102.—Ins. Molucc.
microcarpum, *Willd. Sp. Pl.* iv. 760.—Ins. Molucc.
minutiflorum, *Engl. in DC. Monog. Phan.* iv. 123.—Ins. Philipp.
molle, *Engl. l. c.* 109.—Ins. Borbon.
moluccanum, Blume, Mus. Bot. Lugd. Bat. i. 216 = C. Mehenbethene.
montanum, Korth. ex Blume, l. c. 212 = Santiria montana.
Motleyanum, *Engl. in DC. Monog. Phan.* iv. 133.—Borneo.
multiflorum, *Engl. l. c.* 128.—Madag.
multipinnatum, *Llanos, Fragm.* 107.—Ins. Philipp.
nigrum, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 138.—Ins. Amboina.
nitidum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 533.—Malaya.
oblongifolium, Miq. Fl. Ind. Bat. i. II. 645 = Santiria oblongifolia.
odontophyllum, *Miq. l. c. Suppl.* 525.—Sumatra.
oleiferum, *Baill. Adansonia*, x. (1871–72) 341.—N. Caled.
ovatum, *Engl. in DC. Monog. Phan.* iv. 110.—Ins. Philipp.
paniculatum, Benth. ex Engl. l. c. 124 = mauritianum.
parvifolium, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 536.—Malacca.
patentinervium, *Miq. Fl. Ind. Bat. Suppl.* 526.—Sumatra.
pilosum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 533.—Malacca.
Pimela, Blanco, Fl. Filip. ed. II. 545 = villosum.
Pimela, Blume, Bijdr. 1162 = C. Kipella.
Pimela, *Kon. in Kon. & Sims, Ann. Bot.* i. (1805) 361.—China.
Pimela, Zoll. ex Engl. in DC. Monog. Phan. iv. 127 = Zollingeri.
pruinosum, *Engl. l. c.* 106.—Borneo.
purpurascens, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 532.—Malacca.
Riedelianum, *Engl. in DC. Monog. Phan.* iv. 130.—Ins. Celebes.
rigidum, *Zipp. ex Miq. Fl. Ind. Bat.* i. II. 648.—N. Guinea.
rostratum, *Zipp. ex Blume, Mus. Bot. Lugd. Bat.* i. 221.—Ins. Molucc.
rostriferum, *Miq. Fl. Ind. Bat. Suppl.* 526 — Sumatra.
rubiginosum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 535.—Malacca.
rubiginosum, Miq. Fl. Ind. Bat. i. II. 651 = Santiria rubiginosa.
rufescens, Miq. l. c. 650 = Santiria rufescens.
rufum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 533.—Malacca.
rugosum, Miq. Fl. Ind. Bat. i. II. 649 = Santiria rugosa.
Sajiga, Buch.-Ham. ex Wight & Arn. Prod. 175 = Turpinia pomifera.
samoënse, *Engl. in DC. Monog. Phan.* iv. 134.—Ins. Samoa.
Schweinfurtii, *Engl. l. c.* 145.—Afr. trop.
secundum, *A. W. Benn. in Hook. f. Fl. Brit. Ind.* i. 532.—Malaya.
sericuspe, *Miq. Fl. Ind. Bat.* i. II. 649.—Sumatra.
serrulatum, *Miq. l. c.* 646.—Java.
simplicifolium, *Engl. in DC. Monog. Phan.* iv. 146.—Ins. Amboina.
spectabile, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 116.—Sumatra.
strictum, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 138.—Ind. or.
subrepandum, *Miq. Fl. Ind. Bat. Suppl.* 525.—Sumatra.
subtruncatum, *Engl. in DC. Monog. Phan.* iv. 109.—Java.
sylvestre, *Gaertn. Fruct.* ii. 99. t. 102.—Ins. Amboina.
tomentosum, *Blume, Mus. Bot. Lugd. Bat.* i. 219.—Malaya.
triandrum, *Engl. in DC. Monog. Phan.* iv. 145.—Malaya.

CANARIUM:—
trifoliolatum, *Engl. in DC. Monog. Phan.* iv. 124.—N. Caled.
villosum, *Benth. & Hook. f. ex Naves, in Blanco, Fl. Philipp.* ed. III. *App.* 40.—Ins. Philipp.
virgatum, Miq. Fl. Ind. Bat. i. II. 650 = Santiria virgata.
vitiense, *A. Gray, Bot. U. St. Expl. Exped.* i. 373.—Ins. Fiji.
Vriesseanum, *Engl. in DC. Monog. Phan.* iv. 142.—Ins. Celebes.
zephyrinum, March. in Baill. Adansonia, viii. (1867-68) 53 = communc?
zeylanicum, *Blume, Mus. Bot. Lugd. Bat.* i. 218.—Zeylan.
Zollingeri, *Engl. in DC. Monog. Phan.* iv. 127.—Java.

CANAVALI, Adans. Fam. ii. 325 (1763); Thou. in Desv. Journ. Bot. i. (1813) 77 = **Canavalia**, DC. (Legumin.).

CANAVALIA, DC. Mém. Lég. ix. 375; Prod. ii. 403 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 537.
CANAVALI, Adans. Fam. ii. 325 (1763); Thou. in Desv. Journ. Bot. i. (1813) 77.
CLEMENTEA, Cav. in Anal. Cienc. Nat. vii. (1804) 63. t. 47.
MALOCCHIA, Savi, in Nuov. Giorn. Pisa, viii. (1824) 113.
WENDEROTHIA, Schlecht. in Linnaea, xii. (1838) 330.
altissima, *Macfad. Fl. Jamaic.* i. 292.—Jamaica.
Baueriana, Endl. Prod. Fl. Norf. 91 = obtusifolia.
bonariensis, *Lindl. Bot. Reg.* t. 1199.—Bras.
Bouquete, *Montr. in Mém. Acad. Lyon*, x. (1860) 197. —N. Caled.
brasiliensis, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 135 = ensiformis.
cathartica, *Thou. in Desv. Journ. Bot.* i. (1813) 81.—Madag.
cryptodon, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 96 = bonariensis.
cubensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 178.—Cuba.
dolichoides, Knrz, in Journ. As. Soc. Beng. xliii. (1874) II. 185 = Vigna dolichoides.
emarginata, G. Don, Gen. Syst. ii. 362 = obtusifolia.
ensiformis, *DC. Prod.* ii. 404.—Cosmop. trop.
Finlaysoniana, *R. Grah. in Wall. Cat.* n. 5535.—Siam.
galeata, *Gaudich. Freyc. Voy. Bot.* 486.—Ins. Sandvic.
Gaudichaudii, Endl. in Ann. Wien. Mus. i. (1836) 186 = galeata.
gladiata, DC. Prod. ii. 404 = ensiformis.
grandiflora, *Benth. in Ann. Wien. Mus.* ii. (1838) 135. —Bras.
grandis, Kurz, in Journ. As. Soc. Beng. xliii. (1874) II. 185 = Phaseolus velutinus.
incurva, DC. Prod. ii. 404 = ensiformis.
incurva, Thou. in Desv. Journ. Bot. i. (1813) 80 = ensiformis.
lenta, *Benth. in Ann. Wien. Mus.* ii. (1838) 135.—Bras.
lineata, DC. Prod. ii. 404 = obtusifolia.
Loureirii, G. Don, Gen. Syst. ii. 363 = ensiformis.
lucens, Kurz, in Journ. As. Soc. Beng. xliii. (1874) II. 185 = Vigna lucens.
machacroides, Steud. Nom. ed. II. i. 273 = cathartica.
maritima, Thou. in Desv. Journ. Bot. i. (1813) 80 = obtusifolia.
maxima, *Thou. l. c.*—Ind. or.
miniata, DC. Prod. ii. 404 = obtusifolia.
mollis, Wall. Cat. n. 5533 = ensiformis.
moneta, *Welw. Apont.* 588. n. 62.—Afr. trop.
monodon, E. Mey. Comm. Pl. Afr. Austr. 149 = bonariensis.
multiflora, Hook. & Arn. Bot. Beech. Voy. 416 = villosa.
obcordata, Voigt, Hort. Suburb. Calc. 235 = obtusifolia.
obtusifolia, *DC. Prod.* ii. 404.—Cosmop. trop.
paranensis, Benth. in Ann. Wien. Mus. ii. (1838) 135 = parviflora.
paranensis, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 200 = bonariensis.
parviflora, *Benth. in Mart. Fl. Bras.* xv. i. 177.—Am. trop.

CANAVALIA:—
picta, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 135.—Bras.
polystachya, Schweinf. Rel. Kotsch. 25. t. 20 = ensiformis.
pubescens, Hook. & Arn. Bot. Beech. Voy. 81 = galeata.
rosea, DC. Prod. ii. 404 = obtusifolia.
rostrata, Benth. in Ann. Wien. Mus. ii. (1838) 135 = villosa.
rutilans, DC. Prod. ii. 404 = obtusifolia.
scricea, *A. Gray, Bot. U. St. Expl. Exped.* i. 440.—Ins. Fiji.
Stocksii, Dalz. & Gibs. Bomb. Fl. 69 = ensiformis.
turgida, R. Grah. in Wall. Cat. n. 5534 = ensiformis.
villosa, *Benth. in Ann. Wien. Mus.* ii. (1838) 135.—Mexic.
virosa, *Naves, ex Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 64.—Ins. Philipp.
virosa, Wight & Arn. Prod. 253 = ensiformis.

CANBYA, Parry, ex A. Gray, in Proc. Am. Acad. xii. (1877) 51. *PAPAVERACEAE.*
candida, *Parry, ex A. Gray, l. c.*—Calif.

CANCELLARIA, Sch. Bip. ex Oliver, Fl. Trop. Afr. iii. 321 (1877) = **Adelostigma**, Steetz (Compos.).
senegalensis, Sch. Bip. ex Oliver, l. c. = A. senegalensis.

CANCRINIA, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 124. *COMPOSITAE*, Benth. & Hook. f. ii. 424.
chrysocephala, *Kar. & Kir. l. c.* 125.—As. centr.

CANDARUM, Schott, Melet. i. 17 (1832) = **Amorphophallus**, Blume (Aroid.).
Hookeri, Schott, l. c. = A. campanulatus.
Roxburghii, Schott, l. c. = A. campanulatus.
Rumphii, Schott, l. c. = A. campanulatus.

CANDELABRIA, Hochst. in Flora, xxvi. (1843) 79 = **Bridelia**, Willd. (Euphorbiac.).
micrantha, Hochst. l. c. = B. micrantha.
polystachya, Planch. in Ann. Sc. Nat. Sér. IV. ii. (1854) 264 = Cleistanthus polystachyus.

CANDIDEA, Tenore, in Atti Acc. Nap. iv. (1839) 104. t. 1, 2 = **Vernonia**, Schreb. (Compos.).
senegalensis, Tenore, l. c. = V. Tenoreana.

CANDJERA, Decne. in Ann. Sc. Nat. Sér. II. xix. (1843) 37, sphalm. = **Cansjera**, Juss. (Olacin.).

CANDOLLEA, Baumg. Cat. Hort. Bollw. (1810) = **Menziesia**, Sm. (Ericac.).
azaleoides, Baumg. l. c. = M. globularis.

CANDOLLEA, Labill. in Ann. Mus. Par. vi. (1805) 453. t. 63 = **Stylidium**, Sw.
adnata, F. Mucll. Census, 86 = S. adnata.
adpressa, F. Muell. l. c. = S. adpressa.
alsinoides, F. Muell. l. c. = S. alsinoides.
amoena, F. Muell. l. c. = S. amoena.
Armeria, Labill. in Ann. Mus. Par. vi. (1805) 455 = S. graminifolium.
articulata, F. Muell. Census, 86 = S. articulata.
assimilis, F. Muell. l. c. = S. assimilis.
Barleei, F. Mucll. l. c. = S. Barleei.
brachyphylla, F. Mucll. l. c. = S. brachyphylla.
breviscapa, F. Muell. l. c. = S. breviscapa.
Brunoniana, F. Muell. l. c. = S. Brunoniana.
bulbifera, F. Muell. l. c. = S. bulbifera.
caespitosa, F. Muell. l. c. = S. caespitosa.
calcarata, F. Mucll. l. c. = S. calcarata.
canaliculata, Steud. Nom. ed. II. i. 273 = S. canaliculata.
capillaris, F. Mucll. Census, 86 = S. capillaris.
carnosa, F. Muell. l. c. 85 = S. carnosa.
ciliata, F. Muell. l. c. 86 = S. ciliata.
corymbosa, F. Muell. l. c. = S. corymbosa.
crassifolia, F. Muell. l. c. = S. crassifolia.
crossocephala, F. Muell. l. c. 85 = S. crossocephala.
debilis, F. Muell. l. c. 86 = S. debilis.
despecta, F. Muell. l. c. = S. despecta.
dichotoma, F. Muell. l. c. = S. dichotoma.
diffusa, F. Muell. l. c. = S. diffusa.
disperma, F. Muell. l. c. = S. disperma.

CANDOLLEA:—
diuroides, F. Muell. Census, 86 = Stylidium diuroides.
diversifolia, F. Muell. l. c. = S. diversifolia.
eglandulosa, F. Muell. l. c. = S. eglandulosa.
elongata, F. Muell. l. c. 85 = S. elongata.
emarginata, F. Muell. l. c. 86 = S. emarginata.
eriorrhiza, F. Muell. l. c. = S. eriorrhiza.
falcata, F. Muell. l. c. = S. falcata.
fasciculata, F. Muell. l. c. = S. fasciculata.
filifera, F. Muell. l. c. = S. filifera.
fissiloba, F. Muell. l. c. = S. fissiloba.
Floodii, F. Muell. l. c. = S. Floodii.
floribunda, F. Muell. l. c. = S. floribunda.
glandulosa, F. Muell. l. c. = S. glandulosa.
glauca, Labill. in Ann. Mus. Par. vi. (1805) 454 = S. glaucum.
graminifolia, F. Muell. Census, 85 = S. graminifolia.
guttata, F. Muell. l. c. = S. guttata.
hirsuta, F. Muell. l. c. = S. hirsuta.
imbricata, F. Muell. l. c. 86 = S. imbricata.
juncea, F. Muell. l. c. 85 = S. juncea.
laricifolia, F. Muell. l. c. 86 = S. laricifolia.
lepida, F. Mucll. l. c. = S. lepida.
leptophylla, F. Muell. l. c. = S. leptophylla.
leptorrhiza, F. Muell. l. c. = S. leptorrhiza.
limbata, F. Muell. l. c. = S. limbata.
linearis, F. Muell. l. c. 85 = S. linearis.
lineata, F. Muell. l. c. 86 = S. lineata.
lobuliflora, F. Muell. l. c. = S. lobuliflora.
longituba, F. Muell. l. c. = S. longituba.
lutea, F. Muell. l. c. = S. lutea.
musicola, F. Muell. l. c. = S. musicola.
pachyrrhiza, F. Muell. l. c. = S. pachyrrhiza.
pedunculata, F. Muell. l. c. = S. pedunculata.
perpusilla, F. Muell. l. c. = S. perpusilla.
petiolaris, F. Muell. l. c. = S. petiolaris.
pilosa, Labill. in Ann. Mus. Par. vi. (1805) 453 = S. pilosum.
Preissii, F. Muell. Census, 86 = S. Preissii.
pubigera, F. Muell. l. c. = S. pubigera.
pulchella, F. Muell. l. c. = S. pulchella.
pychnostachya, F. Muell. l. c. = S. pychnostachya.
pygmaea, F. Muell. l. c. = S. pygmaea.
reduplicata, F. Muell. l. c. 85 = S. reduplicata.
repens, F. Muell. l. c. = S. repens.
rhynchocarpa, F. Muell. l. c. 86 = S. rhynchocarpa.
rotundifolia, F. Muell. l. c. = S. rotundifolia.
rupestris, F. Muell. l. c. = S. rupestris.
scabrida, F. Muell. 85 = S. scabrida.
scandens, F. Muell. l. c. 86 = S. scandens.
schizantha, F. Muell. l. c. = S. schizantha.
serrulata, Labill. in Ann. Mus. Par. vi. (1805) 454 = S. graminifolium.
setacea, Labill. l. c. 455 = S. spinulosum?
sobolifera, F. Muell. Census, 86 = S. sobolifera.
spathulata, F. Muell. l. c. = S. spathulata.
spinulosa, F. Mucll. l. c. 85 = S. spinulosa.
squamellosa, F. Muell. l. c. 86 = S. squamellosa.
streptocarpa, F. Muell. l. c. = S. streptocarpa.
striata, F. Muell. l. c. = S. striata.
tenerrima, F. Mucll. l. c. = S. tenerrima.
trichopoda, F. Mucll. l. c. = S. trichopoda.
uliginosa, F. Muell. l. c. = S. uliginosa.
umbellata, Labill. in Ann. Mus. Par. vi. (1805) 456 = S. graminifolium.
uniflora, F. Muell. Census, 86 = S. uniflora.
utricularioides, F. Muell. l. c. = S. utricularioides.
verticillata, F. Muell. l. c. = S. verticillata.
violacea, F. Muell. l. c. = S. violacea.

CANDOLLEA, Labill. Nov. Holl. Pl. ii. 33 t. 176 (1806). *DILLENIACEAE*, Benth. & Hook. f. i. 14.
assimilis, Steud. in Lehm. Pl. Preiss. i. 273 = pedunculata.
calycina, Steud. l. c. 274 = tetrandra.
cuneiformis, *Labill. Nov. Holl. Pl.* ii. 34. t. 176.—Austral.
Cunninghami, Benth. in Maund. Bot. ii. t. 85 = Hibbertia Cunninghamii.
cygnorum, *Steud. in Lehm. Pl. Preiss.* i. 275.—Austral.
desmophylla, *Benth. Fl. Austral.* i. 43.—Austral.
exasperata, *Steud. in Lehm. Pl. Preiss.* i. 276.—Austral.
fasciculata, *R. Br. ex DC. Syst.* i. 424.—Austral.
glaberrima, *Steud. in Lehm. Pl. Preiss.* i. 274.—Austral.

CANDOLLEA:—
glomerosa, *Benth. Fl. Austral.* i. 43.—Austral.
helianthemoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 8.—Austral.
Huegelii, *Benth. in Enum. Pl. Hueg.* 4.—Austral.
kochioides, Turcz. in Bull. Soc. Nat. Mosc. xxii. (1849) II. 7 = fasciculata.
latifolia, Steud. in Lehm. Pl. Preiss. i. 273 = tetrandra.
pachyrrhiza, *Benth. Fl. Austral.* i. 44.—Austral.
parviflora, Steud. in Lehm. Pl. Preiss. i. 276 = pedunculata.
pedunculata, *R. Br. ex DC. Syst.* i. 424.—Austral.
Preissiana, *Steud. in Lehm. Pl. Preiss.* i. 274.—Austral.
racemosa, Endl. in Enum. Pl. Hueg. 5 = pedunculata.
rupestris, Steud. in Lehm. Pl. Preiss. i. 275 = glaberrima.
striata, Steud. in Lehm. Pl. Preiss. i. 274 = Huegelii.
subvaginata, Steud. l. c. 275 = glaberrima.
teretifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxii. (1849) II. 6.—Austral.
tetrandra, *Lindl. Bot. Reg.* (1842) *Misc.* 40.—Austral.
tridentata, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 140 = pedunculata.
uncinata, *Benth. Fl. Austral.* i. 46.—Austral.
vaginata, *Benth. l. c.* 45.—Austral.

CANDOLLEA, Steud. Nom. ed. II. i. 273 (1840) = DECANDOLIA, Bast. = **Agrostis**, Linn. (Gramin.).

CANELLA, P. Br. Hist. Jamaic. 275 t. 27. f. 3 (1756); Sw. in Trans. Linn. Soc. i. (1791) 96. t. 8. *CANELLACEAE*, Benth. & Hook. f. i. 121.
WINTERANIA, Linn. Hort. Cliff. 488 (1737); et Syst. ed. X. (1759) 1045 (Wintirania).
alba, *Murr. Syst.* ed. XIV. 443.—Ind. occ.
alba, Nees, Pl. Officin. t. 418 = Cinnamodendron corticosum.
axillaris, Nees & Mart. in Neuwied, Reise nach Brasil. ii. 336 = Cinnamodendron axillare.
laurifolia, Lodd. Cat. ex Sweet, Hort. Brit. 65 = alba.
obtusifolia, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 348.—Venezuela.
Winterana, Gaertn. Fruct. i. t. 77 = alba.

CANEPHORA, Juss. Gen. 208 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 74.
axillaris, Willd. Sp. Pl. i. 976 = madagascariensis.
capitata, Willd. l. c. = Burchellia capensis.
madagascariensis, *J. F. Gmel. Syst.* 372.—Madag.

CANICIDIA, Vell. Fl. Flum. 184 (1825); iv. t. 139 (1827) = **Connarus**, Linn.
rostrata, Vell. ll. cc. = Connarus cymosa.

CANIDIA, Salisb. Gen. Pl. Fragm. 92 (1866) = **Allium**, Tourn. (Liliac.).
magica, Salisb. l. c. = A. magicum.

CANIRAM, Thou. ex Steud. Nom. ed. II. i. 274 (1840) = **Strychnos**, Linn. (Loganiac.).

CANISTRUM, E. Morr. Belg. Hortic. (1873) 257. t. 15 = **Aechmea**, Ruiz & Pav. (Bromeliac.).
aurantiacum, E. Morr. l. c. = A. aurantiaca.
eburneum, E. Morr. l. c. xxix. (1879) 168. t. 13, 14.—Bras.
fuscum, E. Morr. l. c. xxxiii. (1883) 195.—Bras.
purpureum, E. Morr. l. c.—Bras.
roseum, E. Morr. l. c. xxix. (1879) 301.—Bras.
viride, E. Morr. ex Ch. Morr. l. c. xxiv. (1874) 376. t. 16 = A. viridis.

CANKRIENIA, De Vriese, in Jaarb. Nederl. Maatsch. Tuinb. (1850) = **Primula**, Tourn.
chrysantha, De Vriese, l. c. 30 = P. imperialis.
farinosa, Zoll. Syst. Verz. Ind. Archip. 59.—Java.

CANNA, Linn. Syst. ed. I (1735); Gen. ed. I. 1 (1737). *SCITAMINEAE*, Benth. & Hook. f. iii. 654.
ACHIRIDA, Horan. Prod. Scitam. 18. t. 2 (1862).
CANNACORUS, Tourn. ex Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 378.
DISTEMON, Bouché, in Linnaea, xviii. (1844) 494.
EURYSTYLUS, Bouché, l. c. (1844) 485.

CANNA:—
KATUBALA, Adans. Fam. ii. 67 (1763).
XYPHOSTYLIS, Rafin. Fl. Tellur. iv. 52 (1836).
Achiras, *Gill. in Bot. Reg.* t. 1358.—Chili.
Altensteinii, *Bouché, in Linnaea*, xi. (1837) 326.—Am. austr.
angustifolia, *Linn. Sp. Pl.* 1.—Bras.
angustifolia, Walt. Fl. Carol. 59 = flaccida.
aurantiaca, Rosc. Scit. t. 21 = lutea.
aurantiaca ×, *Tineo, ex Tod. in Ann. Sc. Nat.* Sér. IV. xi. (1859) 377.—Hab.?
aurea, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Hab.?
aureo-vittata, Lodd. Bot. Cab. t. 449 = limbata.
barbadica, Hort. ex Bouché, in Linnaea, viii. (1833) 145 = occidentalis.
bidentata, *Bertol. in Mem. Acc. Sc. Bolog.* x. (1859) 33.—Afr. trop.
bifida, Herb. ex Roem. & Schult. Syst. i. Add. I. 69 = speciosa.
brasiliensis, Hort. Vind. ex Bouché, in Linnaea, xii. (1838) 148 = Jacquini.
brasiliensis, *Rosc. ex Spreng. Syst.* iv. *Cur. Post.* 5.—Bras.
Buekii, Weinm. in Syll. Ratisb. i. (1824) 119; Flora, vii. (1824) 10 = pedunculata.
caripensis, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Hab.?
carnea, *Rosc. Scit.* t. 15.—Bras.
chinensis, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 170 = orientalis.
cinnabarina, *Bouché, in Linnaea*, xviii. (1844) 490.—Mexic.
coccinea, Link, Enum. Hort. Berol. i. 1 = patens.
coccinea, *Mill. Gard. Dict.* ed. VIII. n. 3.—Ind. or.
commutata, Bouché, in Linnaea, viii. (1833) 147 = lutea.
compacta, Hort. ex Bouché, l. c. 145 = occidentalis.
compacta, *Rosc. Scit.* t. 22.—Hab.?
concinna, *Bouché, in Linnaea*, xviii. (1844) 491.—Am. trop.
crocea, *Lag. ex Reichb. Hort. Bot.* t. 148.—Hab.?
crocea, Roem. & Schult. Syst. i. 11 = patens.
curviflora, *Hort. ex Horan. Prod. Scitam.* 18.—Am. centr.
densifolia, *Bouché, in Linnaea*, xviii. (1844) 489.—China.
denudata, *Rosc. Scit.* t. 23.—Bras.
discolor, *Lindl. Bot. Reg.* t. 1231.—Ins. Trinitat.
edulis, *Ker-Gawl. in Bot. Reg.* ix. t. 775.—Am. trop.
Ehrenbergii, *Bouché, in Linnaea*, viii. (1833) 150.—Ins. S. Thomas.
elegans, Rafin. Fl. Ludov. 143 = flaccida?
ellipticifolia, Stokes, Bot. Mat. Med. i. 2 = indica.
esculenta, Lodd. Cat. ex Loud. Hort. Brit. 1 = edulis.
excelsa, *Lodd. Bot. Cab.* n. 743*.—Bras.
exigua, *Bouché, in Linnaea*, xviii. (1844) 490.—Nepal.
eximia, *Hort. ex Horan. Prod. Scitam.* 18.—Bras.
Fintelmanni, *Bouché, in Linnaea*, xviii. (1844) 487.—Hab.?
flaccida, *Rosc. in Trans. Linn. Soc.* viii. (1807) 339.—Am. bor.
flava, Michx. ex Lam. in Journ. Nat. Hist. Par. i. (1792) 416 = flaccida.
flavescens, *Link, Handb.* i. 226.—Ind. or.
floribunda, *Bouché, in Linnaea*, xviii. (1844) 489.—Hab.?
formosa, *Bouché, l. c.* 491.—Bras.
fulgida, *Bouché, l. c.* 490.—Ins. S. Barthol.
gemella, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 22 = gigantea.
gigantea, *Desf. Tabl. Hort. Par.* ed. I. 32.—Bras.
glauca, *Linn. Sp. Pl.* 1.—Ind. occ.; Mexic.
glauca, Walt. Fl. Carol. 59 = flaccida.
heliconiaefolia, *Hort. Berol. ex Bouché, in Linnaea*, viii. (1833) 164.—Mexic.
humilis, *Bouché, l. c.* 153.—China.
indica, Curt. Bot. Mag. t. 454 = patens.
indica, [DC. in] Red. Liliac. t. 201 = speciosa.
indica, *Linn. Sp. Pl.* 1.—Cosmop. trop.
indica, Rosc. in Trans. Linn. Soc. viii. (1807) 338 = orientalis.
indica, Ruiz & Pav. Fl. Per. (1820) i. 1 = edulis.
iridiflora, *Ruiz & Pav. l. c.*—Peruv.
iridiflora, Willd. Enum. Hort. Berol. Suppl. 1 = gigantea.

CANNA:—
Jacquini, *Bouché, in Linnaea*, xii. (1838) 148.—Bras.
juncea, *Retz. Obs.* fasc. i. 9.—China.
lacta, *Bouché, in Linnaea*, xviii. (1844) 489.—Hab.?
lagunensis, Lindl. Bot. Reg. t. 1311 = lutea.
Lamberti, *Lindl. l. c.* t. 470.—Ind. occ.; Guiana.
lanceolata, *Lodd. ex Loud. Hort. Brit.* 1.—Bras.
lanuginosa, *Rosc. Scit.* t. 16.—Am. trop.
latifolia, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. bor.
latifolia, Rosc. Scit. t. 4 = gigantea.
leptochila, *Bouché, in Linnaea*, viii. (1833) 152.—Hab.?
leucocarpa, *Bouché, l. c.* xviii. (1844) 493.—Am. austr.
liliiflora, *Warscew. ex Planch. in Fl. des Serres*, Sér. I. x. (1854–55) 211.—Am. centr.
limbata, *Rosc. Scit.* t. 9.—Bras.
Linkii, *Bouché, in Linnaea*, viii. (1833) 166.—Bras.
liturata, Link, ex A. Dietr. Sp. Pl. i. 12 = glauca.
longifolia, *Bouché, in Linnaea*, xviii. (1844) 486.—Mexic.
lutea, *Mill. Gard. Dict.* ed. VIII. n. 4.—Am. trop.
macrocarpa, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Am. trop.
macrophylla, *Hort. ex Horan. l. c.*—Am. trop.
maculata, *Link, Handb.* i. 227.—Am. trop.
maxima, Lodd. ex Rosc. Scit. t. 2 = Lamberti.
mexicana, A. Dietr. Sp. Pl. i. 11 = glauca.
miniata, *Bouché, in Linnaea*, viii. (1833) 165.—Bras.
montana, *Blume, Enum. Pl. Jav.* 35.—Java.
Moritziana, *Bouché, in Linnaea*, xviii. (1844) 492.—Venezuela.
napalensis, Wall. Cat. n. 6622 = indica.
neglecta, Hort. ex Steud. Nom. ed. II. i. 274 = lanuginosa.
neglecta, Weinm. in Flora, iii. (1820) 607 = gigantea.
nepalensis, Hort. Par. ex D. Dietr. Syn. Pl. i. 4 = indica.
nudata, *Lodd. Cat. ex Steud. Nom.* ed. II. i. 274.—Hab.?
occidentalis, *Rosc. in Bot. Reg.* t. 772.—Ind. occ.
orientalis, Bouché, in Linnaea, viii. (1833) 152 = pulchra.
orientalis, *Rosc. Scit.* t. 12.—Ind. or.
Ottonis, *Bouché, ex Heynh. Nom.* ii. 116, nomen.—Venezuela.
ovata, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Hab.?
pallida, Rosc. Scit. t. 19 = lutea.
paniculata, *Ruiz & Pav. Fl. Per.* i. 1.—Peruv.
patens, Hook. Exot. Fl. t. 228 = limbata.
patens, *Rosc. in Trans. Linn. Soc.* viii. (1807) 338.—Hab.?
pedicellata, *Presl, Reliq. Haenk.* 106.—Chili.
pedunculata, *Sims, Bot. Mag.* t. 2323.—Am. trop.
pentaphylla, D. Dietr. Syn. Pl. i. 5 = platyphylla.
petiolata, *Wall. ex Horan. Prod. Scitam.* 17, nomen.—Hab.?
platyphylla, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 22.—Bras.
Poeppigii, *Bouché, in Linnaea*, xii. (1838) 143.—Bras.
polyclada, *Wawra, in Oestr. Bot. Zeitschr.* xiii. (1863) 7.—Bras.
polymorpha, *Lodd. ex Loud. Hort. Brit.* 1; *Bouché, in Linnaea*, viii. (1833) 151.—Am. austr.
portoricensis, *Bouché, l. c.* xii. (1838) 147.—Porto Rico.
pruinosa, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 79.—Bras.
pulchra, *Bouché, ex Horan. Prod. Scitam.* 16.—Ind. or.
pulchra, *Hassk. Cat. Bogor. Alt.* 52; *et Pl. Jav. Rar.* i. 140.—Java.
recurvata, *Bouché, in Linnaea*, xviii. (1844) 488.—Hab.?
reflexa, Nees, ex D. Dietr. Syn. Pl. i. 4 = pedunculata.
Reevesii, *Lindl. Bot. Reg.* t. 2004.—China.
Roscoeana, *Hort. Berol. ex Bouché, in Linnaea*, viii. (1833) 146.—Am. trop.
rotundifolia ×, E. André, in Rev. Hortic. (1862) 371.
rubra, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 169 = coccinea.
rubricaulis, Link, Enum. Hort. Berol. i. 1 = edulis.
sanguinea, Hort. Angl. ex Bouché, in Linnaea, viii. (1833) 154 = speciosa.
sanguinea, Warsz. ex Otto & Dietr. Allg. Gartenz. xix. (1851) 290 = Warszewiczii.
saturate-rubra, *Bouché, ex C. Koch, Allg. Gartenz.* (1858) 386.—Hab.?

CANNA :—
Schlechtendaliana, *Bouché, in Linnaea*, xviii. (1844) 487.—Hab.?
Schubertii, *Hort. ex Horan. Prod. Scitam.* 18.—Venezuela.
Sellowi, *Hort. Berol. ex Bouché, in Linnaea*, viii. (1833) 162.—Bras.
speciosa, Herb. ex Sims, Bot. Mag. t. 2317 = coccinea.
speciosa, Rosc. Scit. t. 17 = indica.
spectabilis, *Bouché, in Linnaea*, xviii. (1844) 487.—Hab.?
stolonifera, [*Hort. Berol. ex*] *A. Dietr. Syn. Pl.* i. 11.—Reg. Argent.
stricta, *Bouché, in Linnaea*, xii. (1838) 144.—Hab.?
sulphurea, Hort. ex Bouché, l. c. viii. (1833) 150 = lutea.
surinamensis, *Bouché, in Linnaea*, xviii. (1844) 491.—Guiana.
sylvestris, *Rosc. Scit.* t. 10.—Am. austr.
tenuiflora, *Bouché, in Linnaea*, viii. (1833) 150.—Ind. occ.
thyrsiflora, Hegetschw. Comm. Bot. t. 1. f. 8 = patens.
Tinci, *Tod. Cat. Hort. Panorm.* (1858) 25.—Hab.?
tubiflora, *Regel, Cat. Pl. Hort. Aksakov.* 27.—Hab.?
variabilis, Willd. in Ges. Naturf. Fr. Berl. Mag. ii. (1808) 169 = carnea.
variegata, *Bouché, in Linnaea*, xviii. (1844) 487.—Hab.?
ventricosa, *Bouché, l. c.* 488.—Bras.
violacea, *Bouché, l. c.* xii. (1838) 146.—Hab.?
vitellina, *Hort. ex Horan. Prod. Scitam.* 18, nomen.—Hab.?
Warszewiczii, *Hort. ex Otto & Dietr. Allg. Gartenz.* xix. (1851) 289.—Am. centr.
xalapensis, Hort. Berol. ex Bouché, in Linnaea, viii. (1833) 163 = patens.

CANNABINA, Tourn. ex Medic. Phil. Bot. i. 53 (1789) = **Datisca**, Gaertn.
laevis, Moench, Meth. 363 = D. Cannabina.

CANNABIS, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 304 (1737). *URTICACEAE*, Benth. & Hook. f. iii. 357.
chinensis, Delile, Ind. Sem. Hort. Monsp. (1849) = sativa.
erratica, Siev. in Pall. N. Nord. Beitr. vii. 174 = sativa.
foetens, Gilib. Exercit. ii. 450 = sativa.
indica, Lam. Encyc. i. 695 = sativa.
Lupulus, Scop. Fl. Carn. ed. II. ii. 263 = sativa.
macrosperma, Stokes, Bot. Mat. Med. iv. 539 = sativa.
sativa, *Linn. Sp. Pl.* 1027.—As. centr.; Himal. bor. occ.

CANNACORUS, Tourn. ex Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 378 = **Canna**, Linn. (Scitamin.).
indicus, Medic. l. c. 379 = Canna indica.
ovatus, Moench, Meth. 526 = Canna indica.

CANNELLA, Schott, ex Meissn. in DC. Prod. xv. 1. 103, in syn. (1864) = **Ocotea**, Aubl. (Laurin.).
Sassafras, Schott, l. c. = O. indecora.

CANNOMOIS, Beauv. ex Desv. in Ann. Sc. Nat. Sér. I. xiii. (1828) 43. t. 3. f. 1. *RESTIACEAE*, Benth. & Hook. f. iii. 1036.
CUCULLIFERA et MESANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 451 (1836).
Cephalotes, *Beauv. l. c.*—Afr. austr.
robustus, Hochst. in Flora, xxviii. (1845) 340 = C. Cephalotes.
scirpoides, *Hochst. l. c.; Mast. in Journ. Linn. Soc.* x. (1869) 236.—Afr. austr.
simplex, *Kunth, Enum. Pl.* iii. 448.—Afr. austr.
strictus, Hochst. in Flora, xxviii. (1845) 340 = simplex?
virgata, Hochst. l. c. = C. Cephalotes.

CANONANTHUS, G. Don, Gen. Syst. iii. 718 (1834) = **Siphocampylus**, Pohl (Campanulac.).
campanulatus, G. Don, l. c. 719 = S. campanulatus.

CANOPHOLIS, G. Don, Gen. Syst. iv. 634 (1837), sphalm. = **Conopholis**, Wallr. (Orobanch.).

CANOPUS, Presl, Epim. Bot. 248 (1849) = **Exocarpus**, Labill. (Santalac.).
luzonensis, Presl, l. c. = E. latifolius.

CANSCHI, Adans. Fam. ii. 443 (1763) = **Trewia**, Linn. (Euphorb.).

CANOTHUS, Rafin. in Med. Repos. New York, v. (1808) 360 = **Ceanothus**, Linn. (Rhamn.).

CANOTIA, Torr. in Pacif. Rail. Rep. iv. 68 (1856). *RUTACEAE*, Benth. & Hook. f. i. 164, 616.
holacantha, *Torr. l. c.*—Calif.

CANSCORA, Lam. Encyc. i. 601 (1783). *GENTIANEAE*, Benth. & Hook. f. ii. 811.
COMBAMBA, Blanco, Fl. Filip. 510 (1837).
FLEMINGIA, Roxb. ex Steud. Nom. ed. II. i. 643 (1840).
HETEROCANSCORA, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 431.
ORTHOSTEMON, R. Br. Prod. 451 (1810).
PHYLLOCYCLUS, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 235.
PLADERA, Soland. ex Roxb. Fl. Ind. ed. Carey, i. 400 (1832).
POOTIA, Dennst. Schluess. Hort. Malab. 37 (1818).
alata, Wall. Cat. n. 4363 = perfoliata.
andrographioides, *Griff. ex C. B. Clarke, in Journ. Linn. Soc.* xiv. (1875) 431.—Reg. Himal.; Burma.
concanensis, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 104.—Ind. or.
decurrens, *Dalz. in Hook. Kew Journ.* ii. (1850) 136.—Ind. or.
decussata, *Schult. Mant.* iii. 229.—Ind. or.; Afr. trop.
diffusa, *R. Br. Prod.* 451.—Geront. trop.
diffusa, Wall. Cat. n. 4361 K = andrographioides.
divaricata, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 103 = decurrens.
foliosa, D. Don, in G. Don, Gen. Syst. iv. 199 = diffusa?
grandiflora, Wight, Ic. iv. t. 1326 = perfoliata.
Helferiana, *Wall. ex C. B. Clarke, in Journ. Linn. Soc.* xiv. (1875) 432.—Burma.
justicioides, *Griff. ex Voigt, Hort. Suburb. Calc.* 520.—Ind. or.
lancifolia, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 103 = pauciflora.
Lauri, C. B. Clarke, in Journ. Linn. Soc. xiv. (1875) 430 = pauciflora.
Lawii, Wight, Ic. iv. t. 1327 = diffusa.
macrocalyx, Miq. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 104 = perfoliata.
macrocalyx, *Miq. Fl. Ind. Bat.* ii. 558.—Java.
Parishii, *Hook. f. Bot. Mag.* t. 5429.—Burma.
pauciflora, *Dalz. in Hook. Kew Journ.* ii. (1850) 136.—Ind. or.
perfoliata, *Lam. Encyc.* i. 601.—Ind. or.
perfoliata, Wall. Cat. n. 4365 = Wallichii.
pusilla, Roem. & Schult. Syst. iii. 301 = Hoppea dichotoma.
Roxburghii, *Arn. ex Miq. Anal. Bot. Ind.* iii. 11.—Ind. or.
Schultesii, *Wall. Cat.* n. 4368.—Burma.
sessiliflora, *Schult. Mant.* iii. 230.—Ind. or.
tenella, Wall. Cat. n. 4362; Wight, Ic. t. 1327 = diffusa.
ventricosa, J. F. Gmel. Syst. 262 = perfoliata.
Wallichii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 105.—Ind. or.

CANSENIA, Rafin. Sylva Tellur. 122 (1838) = **Bauhinia**, Linn. (Legumin.).
angulosa, Rafin. l. c. = B. angulosa.
tomentosa, Rafin. l. c. = B. tomentosa.

CANSJERA, Juss. Gen. 448 (1789). *OLACINEAE*, Benth. & Hook. f. i. 349.
OCTOPLIS, Rafin. Fl. Tellur. iv. 105 (1836).
grossularioides, *Blanco, Fl. Filip.* ed. I. 73.—Ins. Philipp.
lanceolata, Benth. in Hook. Lond. Journ. Bot. i. (1842) 491 = Rheedii.
leptostachya, *Benth. l. c.* ii. (1843) 231.—Austral.
madagascariensis, Spreng. Syst. i. 453 = Potameia Thouarsii.
malabarica, Lam. Encyc. iii. 433 = Rheedii.
manillana, *Blume, Mus. Bot. Lugd. Bat.* i. 246.—Ins. Philipp.

CANSJERA :—
martabanica, Wall. Cat. n. 7266 = Rheedii.
monostachya, M. Roem. Syn. Hesper. 16 = Rheedii.
parvifolia, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 298.—Burma.
pentandra, Blanco, Fl. Filip. ed. I. 73 = Antidesma Cumingii.
pentandra, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 190 = Antidesma Ghesaembilla.
polystachya, M. Roem. Syn. Hesper. 144 = Rheedii.
Rheedii, Blanco, Fl. Filip. ed. I. 73 = Antidesma Ghaesembilla.
Rheedii, *J. F. Gmel. Syst.* i. 280.—As. et Austral. trop.
Rheedii, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 190 = Antidesma Cumingiana.
scandens, Roxb. Pl. Corom. ii. [2]. t. 103 = Rheedii.
timorensis, *Decne. Bot. Vénus*, 12. t. 4.—Ins. Timor.
zizyphifolia, Griff. in. Calc. Journ. Nat. Hist. iv. (1844) 236 = Rheedii.

CANTALEA, Rafin. Sylva Tellur. 54 (1838) = **Lycium**, Linn. (Solanac.).

CANTHAROSPERMUM, Wight & Arn. Prod. 255 (1834) = **Atylosia**, Wight & Arn. (Legumin.).
albicans, Wight & Arn. l. c. 256 = A. albicans.
calycinum, Miq. Anal. Bot. Ind. i. 8.—Borneo.
distans, Royle, ex Baker, in Hook. f. Fl. Brit. Ind. ii. 216 = A. platycarpa.
nervosum, Royle, ex Baker, l. c. = A. scarabaeoides.
pauciflorum, Wight & Arn. Prod. 255 = A. scarabaeoides.
scarabeoideum, Baill. in Bull. Soc. Linn. Par. i. (1888) 384 = A. scarabaeoides.

CANTHIOPSIS, Seem. Fl. Vit. 166. t. 46 (1865–73) = **Randia**, Linn. (Rubiac.).
odorata, Seem. l. c.—Ins. Fiji.

CANTHIUM, Lam. Encyc. i. 602 (1783) = **Plectronia**, Linn. (Rubiac.).
acuminatum, D. Dietr. Syn. Pl. i. 779 = P. acuminata.
acutiflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 136.—Afr. trop.
Afzelianum, Hiern, l. c. 142.—Afr. trop.
angulosum, Wall. Cat. n. 8285 = Randia longiflora.
angustifolium, Roxb. Hort. Beng. 85; Fl. Ind. i. 533 = P. angustifolia.
anomocarpum, DC. Prod. iv. 475.—Guiana.
anonaefolium, Webb, in Hook. Kew Journ. ii. (1850) 370.—Ins. Cap. Verd.
approximatum, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 234.—Borneo.
arboreum, Vidal, Phan. Cuming. Philipp. 181.—Ins. Philipp.
attenuatum, R. Br. ex Benth. Fl. Austral. iii. 421.—Austral.
Balansae, Baill. Adansonia, xii. (1878) 184.—N. Caled.
barbatum, Seem. Fl. Vit. 132.—Ins. Societ.
Barteri, Hiern, in Oliver, Fl. Trop. Afr. iii. 143.—Afr. trop.
Beecheyi, Steud. Nom. ed. II. i. 275.—Ins. Societ.
Benthamianum, Baill. Adansonia, xii. (1878) 190 = Vangueria canthoides.
bibracteatum, Hiern, in Oliver, Fl. Trop. Afr. iii. 145 = P. bibracteatum.
Billardieri, D. Dietr. Syn. Pl. i. 779 = Coprosma Billardieri.
Breoni, Baill. Adansonia, xii. (1878) 221.—Madag.
buxifolium, Benth. Fl. Austral iii. 422 = P. buxifolia.
campanulatum, Thw. Enum. Pl. Zeyl. 152 = P. campanulatum.
cardiocarpum, Baill. Adansonia, xii. (1878) 185.—N. Caled.
caudatiflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 137.—Afr. trop.
celastroides, Baill. Adansonia, xii. (1878) 190 = P. celastroides.
chinense, Pers. Syn. i. 200 = Randia dumetorum.
confertum, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 235.—Borneo.
congense, Hiern, in Oliver, Fl. Trop. Afr. iii. 141.—Afr. trop.
coprosmoides, F. Muell. Fragm. ii. 134 = [illegible] foliosum.

CANTHIUM :—
Cornelia, Cham. & Schlecht. in Linnaea, iv. (1829) 14.—Afr. trop.
cornelianum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = praec.
coronatum, Lam. Encyc. i. 602 = Randia dumetorum.
corymbosum, D. Dietr. Syn. Pl. i. 779 = P. ventosa, Celastrus Plectronia.
corymbosum, Pers. Syn. i. 200 = Webera corymbosa.
crassum, Hiern, in Oliver, Fl. Trop. Afr. iii. 145.—Afr. trop.
cymosum, Pers. Syn. i. 200 = P. didyma.
Deplanchei, Baill. Adansonia, xii. (1878) 183, 186 = Cyclophyllum Deplanchei.
didymum, Gaertn. f. Fruct. iii. 94. t. 196 = P. didyma.
discolor, Baill. Adansonia, xii. (1878) 191 = Vangueria lasiantha.
discolor, Benth. in Hook. Niger Fl. 409.—Afr. trop.
dubium, Lindl. Bot. Reg. t. 1026 = Diplospora viridiflora.
edule, Baill. Adansonia, xii. (1878) 191 = Vangueria edulis.
ellipticum, Hochst. ex Delile, in Ferr. & Galin. Voy. Abyss. iii. 136 = Feretia apodanthera.
erythroxyloides, Baill. Adansonia, xii. (1878) 220.—Madag.
evonymoides, Baill. l. c. 189 = Vangueria euonymoides.
fasciculatum, Blume, Bijdr. 967.—Java.
ficiforme, Hook. f. Fl. Brit. Ind. iii. 133.—Ind. or.
flavidum, Seem. Fl. Vit. 132.—Ins. Fiji.
floribundum, Wall. ex Voigt, Hort. Suburb. Calc. 389, nomen.—Hab.?
foetidum, Hiern, in Oliver, Fl. Trop. Afr. iii. 142.—Afr. trop.
fraternum, Miq. Fl. Ind. Bat. ii. 253.—Malaya.
furcatum, Miq. l. c. 252.—Malaya.
Gaertneri, Steud. Nom. ed. II. i. 275 = Damnacanthus indicus.
glabriflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 140.—Afr. trop.
glabrum, Blume, Bijdr. 967 = P. glabra.
glaucum, Hiern, in Oliver, Fl. Trop. Afr. iii. 134.—Afr. trop.
glomeratum, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 565 (sphalm. *glomerulatum*) = seq.
glomerulatum, Miq. Fl. Ind. Bat. Suppl. 545 = *confertum*.
gracile, Hiern, in Oliver, Fl. Trop. Afr. iii. 139.—Afr. trop.
gracilipes, Kurz, in Jouru. As. Soc. Beng. xli. (1872) II. 314 = P. gracilipes.
Guenzii, Sond. in Linnaea, xxiii. (1850) 54.—Natal.
Gynochthodes, Baill. Adansonia, xii. (1878) 199.—Ins. Philipp.
hebecladum, DC. Prod. iv. 474.—Ins. Luzon.
Henriettae, Baill. Adansonia, xii. (1878) 186.—N. Caled.
Heudelotii, Hiern, in Oliver, Fl. Trop. Afr. iii. 139.—Afr. trop.
hispidum, Benth. in Hook. Niger Fl. i. 409.—Afr. trop.
horizontale, Benth. l. c. 408 = *Afzelianum*.
horizontale, Hiern, in Oliver, Fl. Trop. Afr. iii. 137.—Afr. trop.
horridum, Benth. Fl. Hongk. 159 = P. parvifolia.
horridum, Blume, Bijdr. 966 = P. horrida.
indicum, D. Dietr. Syn. Pl. i. 779 = Damnacanthus indicus.
infaustum, Baill. Adansonia, xii. (1878) 191 = Vangueria infausta.
Korthalsianum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 252.—Sumatra.
laeve, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 32.—Ins. Banca.
lagoënse, Baill. Adansonia, xii. (1878) 190 = Vangueria euonymoides?
lamprophyllum, F. Muell. Fragm. ii. 133 = *lucidum*.
lanceolatum, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 342 = P. didyma.
lanciflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 146.—Afr. trop.
lasianthoides, Miq. Fl. Ind. Bat. ii. 253.—Java.
latifolium, F. Muell. ex Benth. Fl. Austral. iii. 421.—Austral.
Leschenaultii, Wight & Arn. Prod. 426 = P. angustifolia.
leucophloeum, Baill. Adansonia, xii. (1878) 191 = Fadogia leucophloea.

CANTHIUM :—
lividum, Hiern, in Oliver, Fl. Trop. Afr. iii. 144.—Afr. trop.
lucidulum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 254 = Plectronia didyma.
lucidum, R. Br. in Salt, Abyss. App. lxii.—Abyssin.
lucidum, Hook. & Arn. Bot. Beech. Voy. 65 = Beecheyi.
lucidum, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 132 = P. umbellata?
lycioides, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 188.—Ins. Philipp.
macrocalyx, Baill. Adansonia, xii. (1878) 191 = Vangueria macrocalyx.
macrocarpum, Boiv. ex Baill. l. c. 226 = *pallens*.
macrocarpum, Thw. Enum. Pl. Zeyl. 152 = P. macrocarpa.
Mannii, Hiern, in Oliver, Fl. Trop. Afr. iii. 143.—Afr. trop.
microphyllum, F. Muell. Fragm. ii. 134 = *vaccinifolium*.
mite, Bartl. in DC. Prod. iv. 474.—Ins. Luzon.
moluccanum, Roxb. Hort. Beng. 85; Fl. Ind. i. 536.—Ins. Molucc.
mombazense, Baill. Adansonia, xii. (1878) 188.—Afr. trop.
monoflorum, Blanco, Fl. Filip. ed. I. 166 = *pedunculare*.
montanum, Thw. Enum. Pl. Zeyl. 152.—Zeylan.
multiflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 144.—Afr. trop.
Mundtianum, Cham. & Schlecht. in Linnaea, iv. (1829) 131 = P. Mundtiana.
neglectum, Hiern, in Oliver, Fl. Trop. Afr. iii. 135.—Afr. trop.
neilgherrense, Wight, Ic. t. 1064 *bis* = P. neilgherrensis.
nervosum, Hiern, in Oliver, Fl. Trop. Afr. iii. 143.—Afr. trop.
nitens, DC. Prod. iv. 474.—Ind. or.
nitens, Hiern, in Oliver, Fl. Trop. Afr. iii. 135.—Afr. trop.
obovatum, Klotzsch, in Eckl. & Zeyh. Enum. 361.—Afr. austr.
odoratum, Seem. Fl. Vit. 132.—Ins. Fiji.
oleifolium, Hook. in Mitch. Journ. Trop. Austr. 397.—Austral.
oligocarpum, Hiern, in Oliver, Fl. Trop. Afr. iii. 138.—Afr. trop.
pallens, Baill. Adansonia, xii. (1878) 226.—Ins. Nossibe.
Pancheri, Baill. l. c. 186.—N. Caled.
parviflorum, Bartl. ex DC. Prod. iv. 474 = *hebecladum*.
parviflorum, Lam. Encyc. i. 602 = P. parviflora.
parviflorum, Schlecht. ex Hook. f. Fl. Brit. Ind. iii. 111 = Randia malabarica.
parvifolium, Baill. Adansonia, xii. (1878) 191 = Vangueria parvifolia.
parvifolium, Roxb. Hort. Beng. 15; Fl. Ind. i. 534 = P. parvifolia.
pauciflorum, Baill. Adansonia, xii. (1878) 189 = P. pauciflora.
pauciflorum, Blanco, Fl. Filip. ed. I. 165 = P. horrida.
pedunculare, Cav. Ic. v. 21. t. 436.—Ins. Philipp.
Pervilleanum, Baill. Adansonia, xii. (1878) 197.—Madag.
phyllanthoideum, Baill. l. c. 220.—Afr. trop.
polyanthum, Miq. Fl. Ind. Bat. Suppl. 545.—Sumatra.
polycarpum, Schweinf. ex Oliver, Fl. Trop. Afr. iii. 139.—Afr. trop.
puberulum, Thw. ex Hook. f. Fl. Brit. Ind. iii. 134.—Zeylan.
pyrifolium, Klotzsch, ex Eckl. & Zeyh. Enum. 361.—Afr. austr.
quadrifidum, Labill. Nov. Holl. Pl. i. 69 = Coprosma Billardieri.
recurvum, Wall. Cat. n. 8284 = Randia malabarica.
Rheedii, DC. Prod. iv. 474 = P. Rheedii.
rubens, Hiern, in Oliver, Fl. Trop. Afr. iii. 142.—Afr. trop.
ruminatum, Baill. Adansonia, xii. (1878) 200.—Afr. trop.
sagittatum, Baill. l. c. 184.—N. Caled.
scandens, Blume, Bijdr. 966.—Java.
Schimperianum, A. Rich. Tent. Fl. Abyss. i. 350 = P. Schimperi.
senegalense, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1834) 188 = P. hirsuta.
sessilifolium, A. Gray, in Proc. Am. Acad. iv. (1860) 39.—Ins. Fiji.

CANTHIUM :—
setiflorum, Hiern, in Oliver, Fl. Trop. Afr. iii. 134.—Afr. trop.
setosum, Hiern, l. c. 141.—Afr. trop.
spirostylum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 251.—Sumatra.
subcordatum, DC. Prod. iv. 473.—Afr. trop.
sumatranum, Miq. Fl. Ind. Bat. ii. 254.—Sumatra.
tetraphyllum, Baill. Adansonia, xii. (1878) 192 = Vangueria tetraphylla.
Thonningii, Benth. in Hook. Niger Fl. 410.—Afr. trop.
Thunbergianum, Cham. & Schlecht. in Linnaea, iv. (1829) 130 = Wendlandia thyrsoidea.
thyrsoideum, Roem. & Schult. Syst. vi. 207 = Wendlandia Notoniana.
tomentosum, D. Dietr. Syn. Pl. i. 779 (= *Gynopachis tomentosa*).—Malaya.
travancoricum, Bedd. Ic. Pl. Ind. Or. t. 239 = P. travancorica.
triacanthum, Webb, in Hook. Kew Journ. ii. (1850) 371.—Ins. Cap. Verd.
umbellatum, Wight, Ic. t. 1034; Spicil. Neilgh. i. 81. t. 97 = P. umbellata.
umbelligerum, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 252.—Borneo.
umbellulatum, Korth. in Nederl. Kruidk. Arch. ii. II. (1851) 235.—Borneo.
undulatum, Champ. ex Benth. in Hook. Kew Journ. iv. (1852) 198 = P. didyma.
vacciniifolium, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 47.—Austral.
vangueriioides, Hiern, in Oliver, Fl. Trop. Afr. iii. 146.—Afr. trop.
Vatkeanum, Hiern, l. c. 136.—Afr. trop.
venosum, Hiern, l. c. 144.—Afr. trop.
venulosum, Boiv. ex Baill. Adansonia, xii. (1878) 221.—Ins. Comor.
Vieillardi, Baill. l. c. 185.—N. Caled.
Villarii, Vidal, Phan. Cuming. Philipp. 182.—Ins. Philipp.
violaceum, Zoll. & Moritzi, Syst. Verz. Zoll. 64.—Java.
viridiflorum, D. Dietr. Syn. Pl. i. 780 = Diplospora viridiflora.
zanzibaricum, Klotzsch, in Peters, Reise Mossamb. Bot. 291 = P. zanzibarica.
zizyphinum, Wall. Cat. n. 8288 A, C = P. horrida.

CANTHOPSIS, Miq. Fl. Ind. Bat. ii. 256 (1856). *RUBIACEAE*, Benth. & Hook. f. ii. 113.
pubiflora, *Miq. l. c.* 257.—Ins. Timor.

CANTUA, Juss. Gen. 136 (1789). *POLEMONIACEAE*, Benth. & Hook. f. ii. 824.
PERIPHRAGMOS, Ruiz & Pav. Prod. 26. t. 24 (1794).
aggregata, Pursh, Fl. Am. Sept. i. 147 = Gilia aggregata.
bicolor, *Lem. in Fl. des Serres*, Sér. I. iii. (1847) 242, verso.—Bolivia.
breviflora, Juss. in Ann. Mus. Par. iii. (1804) 119 = Gilia laciniata.
brevifolia, Steud. Nom. ed. II. i. 275 = praec.
buxifolia, *Lam. Encyc.* i. 603.—Peruv.
caerulea, Poir. Encyc. Suppl. ii. 80 = Loeselia caerulea.
coccinea, Poir. l. c. = Loeselia coccinea.
cordata, *Juss. in Ann. Mus. Par.* iii. (1804) 119.—Peruv.
coronopifolia, Willd. Sp. Pl. i. 879 = Gilia coronopifolia.
cuneifolia, *Juss. ex Roem. & Schult. Syst.* iv. 369.—Hab.?
dependens, Pers. Syn. i. 187 = buxifolia.
elegans, Poir. Encyc. Suppl. ii. 80 = Gilia coronopifolia.
fasciculata, Humb. ex Roem. & Schult. Syst. iv. 369 = Fouquiera spinosa.
flexuosa, Pers. Syn. i. 187 = pyrifolia.
floridana, Nutt. ex A. Gray, Syn. Fl. N. Am. ii. 1. 145 = Gilia coronopifolia.
foetida, Pers. Syn. i. 187 = Vestia lycioides.
glandulosa, Poir. Encyc. Suppl. ii. 80 = Loeselia glandulosa.
glomeriflora, Juss. in Ann. Mus. Par. iii. (1804) 119 = Collomia Cavanillesiana.
glutinosa, *Presl, Epim. Bot.* 213.—Chili.
Hoitzia, Willd. Sp. Pl. i. 878 = Loeselia coccinea.
laciniata, Poir. Encyc. Suppl. ii. 81 = Gilia laciniata.
ligustrifolia, Juss. in Ann. Mus. Par. iii. (1804) 118 = Vestia lycioides.

CANTUA:—
longiflora, Torr. in Ann. Lyc. N. York, iii. (1828) 221 = Collomia longiflora.
loxensis, Willd. ex Roem. & Schult. Syst. iv. 369 = pyrifolia.
megapotamica, *Spreng. Syst.* iv. *Cur. Post.* 64 (quid?). —Bras.
ovata, *Cav. Ic.* iv. 43.—Peruv.
parviflora, Pursh, Fl. Am. Sept. ii. 730 = Gilia parviflora.
peruviana, J. F. Gmel. Syst. 347 = pyrifolia.
pinnatifida, Lam. Illustr. i. 473 = Gilia coronopifolia.
pungens, Torr. in Ann. Lyc. N. York. ii. (1828) 221 = Gilia pungens.
pyrifolia, *Juss. in Ann. Mus. Par.* iii. (1804) 117.—Peruv.
quercifolia, *Juss. l. c.* 118.—Peruv.
theaefolia, *D. Don, in Edinb. Phil. Journ.* (Apl.–Oct. 1822) 289.—Peruv.
thyrsoidea, Juss. in Ann. Mus. Par. iii. (1804) 119 = Gilia coronopifolia.
tomentosa, Cav. Ic. iv. 43 = buxifolia.
tuberosa, Roem. & Schult. Syst. iv. 793 = Ipomoea muricata.
uniflora, Pers. Syn. i. 187 = ovata.

CANTUFFA, J. F. Gmel. Syst. 677 (1791) = **Pterolobium**, R. Br. (Legumin.).
exosa, J. F. Gmel. l. c. = P. lacerans.

CAOUTCHOUA, J. F. Gmel. Syst. 1007 (1791) = **Hevea**, Aubl. (Euphorbiac.).
elastica, J. F. Gmel. l. c. = H. guianensis.

CAPANEA, Deene. in Van Houtte, Fl. des Serres, t. 499 (1850) = **Campanea**, Deene. (Gesnerac.).

CAPANEMIA, Rodrig. Orch. Nov. i. 137 (1877). *ORCHIDEAE.*
carinata, *Rodrig. l. c.* ii. 243.—Bras.
micromera, *Rodrig. l. c.* i. 138.—Bras.
Therasiae, *Rodrig. l. c.* ii. 244.—Bras.
uliginosa, *Rodrig. l. c.* i. 137.—Bras.

CAPASSA, Klotzsch, in Peters Reise Mossamb. Bot. 27. t. 5 (1861) = **Lonchocarpus**, Humb. & Bonpl. (Legumin.).
violacea, Klotzsch, l. c. 28 = L. violaceus.

CAPELLENIA, Hassk. Cat. Hort. Bog. Alt. 178 (1844) = CAPELLIA, Blume = **Wormia**, Rottb. (Dilleniac.).

CAPELLENIA, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1866) 239 = **Endospermum**, Benth. (Euphorb.).
moluccana, Teijsm. & Binn. l. c.—Ins. Molucc.

CAPELLIA, Blume, Bijdr. 5 (1825) = **Wormia**, Rottb. (Dilleniac.).
biflora, A. Gray, Bot. U. St. Expl. Exped. i. 15. t. 1 = W. biflora.
membranifolia, A. Gray, l. c. 17 = W. membranifolia.
multiflora, Blume, Bijdr. 5.—Java.
pauciflora, Zoll. & Moritzi, Syst. Verz. Zoll. 35.—Java.

CAPERONIA, St. Hil. Pl. Rem. Bras. 244 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 304.
ANDROPHORANTHUS, Karst. in C. Koch & Fint. Wochenschr. (1859) 5.
CAVANILLA, Vell. Fl. Flum. 226 (1825).
LEPIDOCOCCA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 588.
METERANA, Rafin. Sylva Tellur. 65 (1838).
acalyphifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 58. —Reg. Argent.
aculeolata, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 152.—Bras.
angustissima, *Klotzsch, in Hook. Lond. Journ. Bot.* ii. (1843) 50.—Guiana.
bahiensis, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 325.—Bras.
buettneriacea, *Mucll. Arg. l. c.* 320.—Bras.
castaneaefolia, Miq. in Linnaea, xxi. (1848) 478 = corchoroides.
castaneaefolia, *St. Hil. Pl. Rem. Bras.* 245.—Java.
corchoroides, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 153.—Guiana.
cordata, *St. Hil. Pl. Rem. Bras.* 245.—Bras.

CAPERONIA:—
Gardneri, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 321. —Bras.
heteropetala, *F. Didr. in Kjoeb. Vidensk. Meddel.* (1857) 148.—Bras.
heteropetaloides, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 152.—Bras.
Langsdorffii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 319.—Bras.
linearifolia, *St. Hil. Pl. Rem. Bras.* 246.—Bras.
multicostata, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 323.—Bras.
nervosa, A. Rich. Fl. Cub. Fanerog. ii. 213 = castaneaefolia.
paludosa, *Klotzsch, in Hook. Lond. Journ. Bot.* ii. (1843) 51.—Mexic.
palustris, *St. Hil. Pl. Rem. Bras.* 245.—Ind. occ.; Mexic.
palustris, H. B. & K. Nov. Gen. et Sp. ii. 71 = castaneaefolia.
panamensis, Klotzsch, in Seem. Bot. Voy. Herald, 103 = paludosa.
Regnellii, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 321.—Bras.
Rutenbergii, *Muell. Arg. in Bremen Abh.* vii. (1880) 25.—Madag.
senegalensis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 153.—Afr. trop.
serrata, Baill. in Adansonia, i. (1860–61) 66 = senegalensis.
serrata, *Presl, Epim. Bot.* 213.—Afr. trop.
stenophylla, *Muell. Arg. in Mart. Fl. Bras.* xi. II. 326. —Bras.
Velloziana, *Muell. Arg. l. c.* 323.—Bras.

CAPIA, Domb. ex Juss. in Dict. Sc. Nat. vi. 500 (1805) = **Lapageria**, Ruiz & Pav. (Liliac.).

CAPIRONA, Spruce, in Journ. Linn. Soc. iii. (1859) 200. *RUBIACEAE*, Benth. & Hook. f. ii. 39.
Boiviniana, *Baill. in Bull. Soc. Linn. Par.* i. (1880) 270.—Madag.
decorticans, *Spruce, in Journ. Linn. Soc.* iii. (1859) 200.—Peruv.

CAPITELLARIA, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 103 = **Sagraea**, DC. (Melastom.).
Benthami, Naud. l. c. = S. capitata.

CAPNITES, Dum. Fl. Belg. 117 (1827) (err. typ. 177) = **Corydalis**, Vent. (Papaverac.).
alpestris, Rupr. Fl. Caucas. 59 = Cor. pauciflora.
angustifolia, Rupr. l. c. 57 = Cor. angustifolia.
araratica, Rupr. l. c. 61.—Armenia.
Bayerniana, Rupr. l. c. 59. t. 4 = Cor. pallida.
caucasica, Rupr. l. c. 56 = Cor. caucasica.
cava, Dum. Fl. Belg. 117 (err. typ. 177) = Cor. tuberosa, bulbosa.
conorhiza, 'Ledeb.' ex Rupr. Fl. Caucas. 61 = Cor. conorhiza.
digitata, Dum. Fl. Belg. 117 (err. typ. 177) = Cor. solida.
Emanueli, Rupr. Fl. Caucas. 58 = Cor. pauciflora.
fabacea, Dum. Fl. Belg. 117 (err. typ. 177) = Cor. fabacea.
macrosepala, Rupr. Fl. Caucas. 61.—Reg. Caucas.
Marschalliana, Rupr. l. c. 55 = Cor. Marschalliana.
ochroleuca, Rupr. l. c. 61 = Cor. lutea.
pallidiflora, Rupr. l. c. 58 = Cor. pallida.
roseo-purpurea, Rupr. l. c. 288.—Reg. Caucas.
solida, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 329 = Cor. solida.

CAPNITIS, E. Mey. Comm. Pl. Afr. Austr. 81 (1835) = **Lotononis**, Eckl. & Zeyh. (Legumin.).
clandestina, E. Mey. l. c. = L. clandestina.
porrecta, E. Mey. l. c. = L. porrecta.

CAPNOCYSTIS, Juss. in Ann. Mus. Par. xviii. (1811) 473 = CYSTICAPNOS, Boerh. = **Corydalis**, Vent. (Papaverac.).

CAPNOGONIUM, Bernh. Ind. Sem. Hort. Erfurt. (1844) ex Endl. Gen. Suppl. ii. 107 = **Corydalis**, Vent. (Papaverac.).
nobile, Bernh. l. c. = Cor. nobilis.

CAPNOIDES, Tourn. ex Adans. Fam. ii. 431 (1763) = **Corydalis**, Vent. (Papaverac.).
albida, Bernh. ex Steud. Nom. ed. II. i. 276 = Cor. lutea.
cava, Moench, Meth. 52 = Cor. tuberosa.
glauca, Moench, l. c. = Cor. glauca.
lutea, Gaertn. Fruct. ii. 163 = Cor. lutea.
minor, Schum. ex DC. Syst. ii. 119 = Cor. fabacea.
nobilis, Moench, Meth. 52 = Cor. nobilis.
rigidula, Bernh. ex Steud. Nom. ed. II. i. 276 = Cor. ochroleuca.
scandens, Moench, Meth. Suppl. 215 = Adlumia cirrhosa.
sempervirens, Borckh. in Roem. Arch. i. II. 44 = Cor. glauca.
solida, Moench, Meth. 52 = Cor. tuberosa.

CAPNOPHYLLUM, Gaertn. Fruct. ii. 32. t. 85 (1791). *UMBELLIFERAE*, Benth. & Hook. f. i. 908.
ACTINOCLADUS, E. Mey. Ind. Sem. Hort. Regiom. (1846).
KRUBERA, Hoffm. Gen. Umb. 103 (1814).
SCLEROSCIADIUM, Koch, in DC. Coll. Mém. v. 43 (1829); Prod. iv. 140 (1836).
ULOSPERMUM, Link, Enum. Hort. Berol. i. 267 (1822).
africanum, *Gaertn. Fruct.* ii. 32. t. 85.—Afr. austr.
dichotomum, *Lag. Gen. et Sp. Nov.* 13.—Reg. Mediterr.
Jacquini, *DC. Prod.* iv. 187.—Afr. austr.
peregrinum, Lange, in Willk. & Lange, Prod. Fl. Hisp. iii. 33 = dichotomum.
simplex, Lag. Gen. et Sp. Nov. 13 = africanum.

CAPNORCHIS, Borck. in Roem. Arch. i. II. 46 (1797) = **Dicentra**, Bernh. (Papaverac.).
chrysantha, Planch. in Fl. des Serres, Sér. I. viii. (1852–53) 193. t. 820 = D. chrysantha.
Cucullaria, Planch. l. c. 193 = D. Cucullaria.
eximia, Planch. l. c. = D. eximia.
formosa, Planch. l. c. = D. formosa.
spectabilis, Borck. in Roem. Arch. i. II. 46 = D. spectabilis.

CAPNOREA, Rafin. Fl. Tellur. iii. 74 (1836) = **Nicotiana**, Linn. (Solanac.).
nana, Rafin. l. c. 75 = N. nana.

CAPPARIS, [Tourn.] Linn. Syst. ed. I (1735). *CAPPARIDEAE*, Benth. & Hook. f. i. 108.
BEAUTEMPSIA, Gaudich. Bot. Voy. Bonite, t. 56 (1844?).
BUSBECKIA, Endl. Prod. Fl. Ins. Norf. 64 (1833).
CALYPTRANTHUS, Thou. Obs. Pl. Isles Afr. 26 (1811).
COLICODENDRON, Mart. Herb. Fl. Bras. 201 (1837).
DESTRUGUEZIA, Gaudich. Bot. Voy. Bonite, t. 57 (1844?).
HOMBAK, Adans. Fam. ii. 408 (1763).
INTUTIS, MARSESINA, OCTANEMA, OLIGLORON, OLOFUTON, PLEUTERON, TRICLANTHERA, Rafin. Sylva Tellur. 108–112 (1838).
LINDACKERA, Sieber, ex Endl. Gen. 893, in syn. (1841).
PETERSIA, Klotzsch, in Peters, Reise Mossamb. Bot. 168. t. 30 (1861).
QUADRELLA, Meissn. Gen. 15 (1867).
SODADA, Forsk. Fl. Aegypt. Arab. 81 (1775).
UTERVERIA, Bertol. Pl. Nov. Hort. Bonon. ii. 8 (1839).
aculeata, Steud. Nom. ed. I. 147 = spinosa.
acuminata, Lindl. Bot. Reg. t. 1320 = acutifolia.
acuminata, Roxb. Hort. Beng. 41; Fl. Ind. ii. 566 = zeylanica.
acuminata, *Willd. Sp. Pl.* ii. 1131.—Ind. or.
acuminata, Willd. ex Wall. Cat. n. 6990 C. horrida.
acutifolia, *Sweet, Hort. Brit.* ed. II. 585.—China.
aegyptia, Lam. Encyc. i. 605 = spinosa.
aegyptia, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 274 = Fontanesii.
Afzelii, DC. Prod. i. 246 = erythrocarpos.
Aguba, Banks, ex DC. Prod. i. 248 = Roxburghii.
albitrunca, *Burch. Trav.* i. 343.—Afr. austr.
ambigua, *Kurz, For. Fl. Brit. Burma*, i. 65.—Ins. Andam.
amplissima, *Lam. Encyc.* i. 607.—Ind. occ.; Mexic.
amygdalifolia, *Jacq. Enum. Pl. Carib.* 24.—Ind. occ.
amygdalina, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 155 = Grisebachii.

CAPPARIS:—
amygdalina, Lam. Encyc. i. 608 = Capparis Breynia.
angulata, *Ruiz & Pav. Fl. Per.* t. 431 *ex DC. Prod.* i. 253.—Ecuador.
angustifolia, *H. B. & K. Nov. Gen. et Sp.* v. 96. t. 438.—Mexic.
antanossarum, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 462.—Madag.
apetala, Roth, Sp. Pl. Nov. 238 = Niebuhria linearis.
aphylla, *Roth, l. c.*—Afr. bor.; Arab.; Ind. or.
arborescens, Mill. Gard. Dict. ed. VIII. n. 3 = pulcherrima.
Arrabidae, *Steud. Nom.* ed. II. i. 276.—Bras.
artensis, *Montr. in Mém. Acad. Lyon*, x. (1860) 177.—N. Caled.
asperifolia, *Presl, Rel. Haenk.* ii. 86.—Mexic.
assamica, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 177.—Assam.
aurantioides, *Presl, Rel. Haenk.* ii. 86.—Ins. Luzon.
avicennifolia, *H. B. & K. Nov. Gen. et Sp.* v. 94.—Am. austr.
Baducca, Blanco, Fl. Filip. ed. I. 438 = spinosa.
Baducca, Linn. Sp. Pl. 504 = zeylanica ?
barcellonensis, H. B. & K. Nov. Gen. et Sp. v. 92 = Breynia.
Billardieri, *DC. Prod.* i. 247.—Ins. Molucc.
bisperma, Roxb. Hort. Beng. [93]; Fl. Ind. ii. 568 = grandis.
bisperma, Wight, ex Hook. f. Fl. Brit. Ind. i. 176 = Roxburghii.
brachyscias, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 323.—Java.
brasiliana, *DC. Prod.* i. 249.—Bras.
Brassii, *DC. l. c.* 248.—Am. austr.
brevipes, Benth. Bot. Voy. Sulph. 65 = verrucosa.
brevis, Spreng. Fl. Hal. Mant. 43 = eustachiana.
brevisiliqua, *Moç. & Sessé, ex DC. Prod.* i. 251.—Mexic.
brevispina, DC. Prod. i. 246 = zeylanica.
brevispina, Wight & Arn. Prod. 24 = divaricata.
Breynia, H. B. & K. Nov. Gen. et Sp. v. 97 = odoratissima.
Breynia, *Linn. Syst.* ed. X. 1071; *Jacq. Am. Select.* 161. t. 103.—Am. austr.
caerulea, Heyne, ex Wall. Cat. n. 6986 = diversifolia.
callophylla, *Blume, Bijdr.* 53.—Java.
callosa, Blume, l. c. = micracantha.
canescens, *Banks, ex DC. Prod.* i. 246.—Austral.
canescens, G. Don, Gen. Syst. i. 281 = Doniana.
cantoniensis, *Lour. Fl. Cochinch.* 330.—China.
capensis, Thunb. Prod. Pl. Cap. 92 = citrifolia.
Carandas, *Burm. f. Fl. Ind.* 118.—Ind. or.
cardiophylla, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 325.—Venezuela.
cartilaginea, Decne. in Ann. Sc. Nat. Sér. II. iii. (1835) 273 = galeata.
celebica, *Miq. Illustr. Fl. Arch. Ind.* 26.—Ins. Celebes.
cerasifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 71.—Archip. Sulu.
chinensis, G. Don, Gen. Syst. i. 278 = acutifolia.
chrysomela, *Boj. in Ann. Sc. Nat.* Sér. II. xx. (1843) 59.—Madag.
citrifolia, *Lam. Encyc.* i. 606.—Afr. austr.
clutiaefolia, *Burch. ex DC. Prod.* i. 248.—Afr. austr.
coccolobifolia, *Mart. ex Eichl. in Mart. Fl. Bras.* xiii. I. 284.—Am. trop.
coluteoides, Boiss. Diagn. Ser. I. xiii. 83 = Cleome coluteoides.
commutata, Spreng. Neue Entdeck. iii. 57 = frondosa.
comosa, *Jacq. Enum. Pl. Carib.* 23.—Ins. S. Doming.
conferta, Mill. Gard. Dict. ed. VIII. n. 8 = laurifolia.
conspicua, Wall. Cat. n. 6991 = micracantha.
cordata, Ruiz & Pav. Fl. Per. t. 432; ex DC. Prod. i. 251 = crotonoides.
cordifolia, Lam. Encyc. i. 609 = spinosa.
coriacea, *Burch. ex DC. Prod.* i. 248.—Afr. austr.
corymbifera, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 62.—Afr. austr.
corymbosa, *Lam. Encyc.* i. 605.—Afr. trop.
corymbosa, Roxb. Hort. Beng. [93]; Fl. Ind. ii. 569 = Roxburghii.
crassifolia, *A. Gray, Bot. U. St. Expl. Exped.* i. 71.—Archip. Sulu.
crassifolia, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 227.—Burma.
crotonoides, *H. B. & K. Nov. Gen. et Sp.* v. 95. t. 437.—N. Granat.
cuneata, DC. Prod. i. 249 = frondosa.
Cynophallophora, Linn. Sp. Pl. ed. I. 504 = jamaicensis.

CAPPARIS:—
Cynophallophora, *Linn. Sp. Pl.* ed. II. 721; *Jacq. Am. Select.* t. 98.—Am. austr.
Dahi, *Forsk. Fl. Aegypt. Arab.* 212 (Zygophylli sp. ?).—Arab.
dasypetala, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 322.—Java.
dealbata, *DC. Prod.* i. 246.—Ins. Timor.
decidua, Edgew. in Journ. Linn. Soc. vi. (1862) 184 = aphylla.
declinata, Vell. Fl. Flum. 230; v. t. 111 = C. Cynophallophora.
detonsa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 80.—N. Granat.
didymobotrys, *Ruiz & Pav. Fl. Per.* t. 423; *ex DC. Prod.* i. 253.—Ecuador.
disperma, Walp. Rep. i. 199 = grandis.
disticha, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 70.—Burma.
divaricata, Lam. Encyc. i. 606.; Klein, ex Wall. Cat. n. 6988 A = diversifolia.
diversifolia, *Wight & Arn. Prod.* 27.—Ind. or.
domingensis, *Spreng. ex DC. Prod.* i. 253.—Ins. S. Doming.
Doniana, *D. Dietr. Syn. Pl.* iii. 227.—Afr. trop.
Eichleriana, *Urban, in Linnaea*, xliii. (1880–82) 265.—Bras.
elaeagnifolia, *Jacq. Enum. Pl. Carib.* 23.—Ind. occ.
elegans, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 24.—Bras
elliptica, Span. in Linnaea, xv. (1841) 166 = Roxburghii ?
emarginata, *Presl, Reliq. Haenk.* ii. 85.—Ins. Luzon.
emarginata, A. Rich. Ess. Fl. Cub. 78 = jamaicensis.
emarginata, Zipp. ex Span. in Linnaea, xv. (1841) 166 = sepiaria.
erythrocarpos, *Isert, in Ges. Naturf. Fr. Berl. Schrift.* ix. (1789) 334. t. 9.—Guiana.
erythrodasys, Miq. Pl. Jungh. 397 = horrida.
eustachiana, *Jacq. Enum. Pl. Carib.* 23.—Ins. S. Eustach.
exstipulata, J. F. Gmel. Syst. 807 = inermis ?
falcata, Lour. Fl. Cochinch. 330 = Crataeva falcata.
fascicularis, DC. Prod. i. 248 = corymbosa.
ferruginea, *Linn. Syst.* ed. X. 1071.—Ind. occ.
ferruginea, Sieber, ex Eichl. in Mart. Fl. Bras. xiii. I. 271 = C. Breynia.
ferruginea, Willd. ex Eichl. l. c. = odoratissima.
Finlaysoniana, *Wall. Cat.* n. 6992 B.—Malaya.
Finlaysoniana, Wall. l. c. A = micracantha.
flavicans, *Wall. l. c.* n. 7003.—Burma.
flexicaulis, *Hance, in Journ. Bot.* vi. (1878) 225.—China.
flexuosa, *Blume, ex Hassk. Pl. Jav. Rar.* 179.—Java.
flexuosa, Linn. Sp. Pl. ed. II. 722 = C. Cynophallophora.
flexuosa, Vell. Fl. Flum. 230; v. t. 108 = Arrabidae.
floribunda, Lepr. ex Walp. Rep. i. 197 = tomentosa.
floribunda, *Wight, Illustr. Ind. Bot.* i. 35. t. 14.—Ind. or.; Ins. Philipp.
foetida, *Blume, Bijdr.* 52.—Java.
Fontanesii, *DC. Prod.* i. 245.—Ins. Maurit.
formosa, Dalz. in Hook. Kew Journ. ii. (1850) 40 = Heyneana.
formosa, Wall. Cat. n. 6982 A = horrida.
Forsteniana, *Miq. Illustr. Fl. Arch. Ind.* 32.—Malaya.
frondosa, *Jacq. Enum. Pl. Carib.* 24; *Linn. Sp. Pl.* ed. II. *App.* 1674.—Am. austr.
fruticosa, *Mill. Gard. Dict.* ed. VIII. n. 7.—Am. austr.
furfuracea, *Ruiz & Pav. ex DC. Prod.* i. 252.—Mexic.
galeata, *Fresen. in Mus. Senckenb.* ii. (1837) 111.—Aegypt.
Gaudichaudiana, *Eichl. in Mart. Fl. Bras.* xiii. I. 273.—Peruv.
glandulosa, *Desf. Cat. Hort. Par.* ed. III. 405.—Hab. ?
glauca, *Wall. Cat.* n. 7005.—Burma.
globifera, *Delile, in Rochet, Sec. Voy. Choa*, 340.—Afr. trop.
Gomeziana, Steud. Nom. ed. II. i. 277 = pumila.
Grandidiera, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 463.—Madag.
grandiflora, *Wall. Cat.* n. 6984.—Ind. or.
grandis, *Linn. f. Suppl.* 263.—Ind. or.; Burma.
grandis, Russ. ex Wall. Cat. n. 7001 = Russeliana.
Grisebachii, *Eichl. in Mart. Fl. Bras.* xiii. I. 275.—Cuba.
guayaquilensis, H. B. & K. Nov. Gen. et Sp. v. 89 = C. Cynophallophora.
Guenezii, *Sond. in Harv. & Sond. Fl. Cap.* i. 62.—Afr. austr.

CAPPARIS:—
Hasseltiana, *Miq. Illustr. Fl. Arch. Ind.* 24. t. 13.—Java.
hastata, Jacq. Enum. Pl. Carib. 23 = C. Cynophallophora.
hastigera, Hance, in Journ. Bot. vi. (1868) 296 = Swinhoii.
herbacea, Willd. Enum. Hort. Berol. 560 = spinosa.
heteracantha, *DC. Prod.* i. 246.—Mesopotam.
heteroclita, Roxb. Hort. Beng. 41; Fl. Ind. ii. 570 = Maerua arenaria.
heterogenea, Roxb. ex Steud. Nom. ed. II. i. 277 = Maerua oblongifolia.
heterophylla, *Ruiz & Pav. ex DC. Prod.* i. 250.—Ecuador.
Heyneana, *Wall. Cat.* n. 6985.—Ind. or.
horrida, Banks, ex Wight & Arn. Prod. 25 = stylosa.
horrida, *Linn. f. Suppl.* 264.—As. trop.; Malaya.
Humbloti, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 463.—Madag.
humistrata, *F. Muell. Fragm.* v. 156.—Austral.
hypericoides, *Hochst. in Flora*, xxvii. (1844) i. 290.—Afr. occ.
hypoleuca, *Presl, Rel. Haenk.* ii. 87.—Peruv.
incana, *H. B. & K. Nov. Gen. et Sp.* v. 94.—Am. austr.
incanescens, DC. Prod. i. 247 = sepiaria.
inermis, *Forsk. Fl. Aegypt. Arab.* 100.—Arab.
intermedia, H. B. & K. Nov. Gen. et Sp. v. 98 = odoratissima.
isthmensis, *Eichl. in Mart. Fl. Bras.* xiii. I. 269.—Am. trop.
Jacobinae, *Moric. ex Eichl. l. c.* 277.—Bras.
jamaicensis, *Jacq. Enum. Pl. Carib.* 23.—Jamaica.
Karwinskiana, *Schlecht. in Linnaea*, x. (1836) 237.—Mexic.
Kirkii, *Oliver, Fl. Trop. Afr.* i. 98.—Afr. trop.
Korthalsiana, *Miq. Illustr. Fl. Arch. Ind.* 31.—Borneo.
laeta, H. B. & K. Nov. Gen. et Sp. v. 88 = verrucosa.
laetevirens, Mart. in Flora (1839) I. Beibl. 23 = C. Cynophallophora.
laevigata, Mart. l. c. 21 = C. Cynophallophora.
lanceolaris, *Lesch. ex Steud. Nom.* ed. II. i. 277.—Java.
lanceolata, Ruiz & Pav. Fl. Per. t. 423; ex DC. Prod. i. 249 = C. Cynophallophora.
lasiantha, *R. Br. ex DC. Prod.* i. 247.—Austral.
lasiopodia, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 322.—Ins. Luzon.
laurifolia, *Mill. Gard. Dict.* ed. VIII. errat.—N. Granat.
laurina, *H. B. & K. Nov. Gen. et Sp.* v. 93.—Am. austr.
leucophylla, DC. Prod. i. 246 = spinosa.
linearifolia, Hook. f. in Hook. Niger Fl. 217 = Thonningii.
linearifolia, *Linden, ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 324.—Venezuela.
linearis, Blanco, Fl. Filip. ed. I. 438 = viminea.
linearis, *Jacq. Enum. Pl. Carib.* 24.—Am. austr.
lineata, *Pers. Syn.* ii. 60.—Bras.
linifolia, Roxb. Hort. Beng. 41 = Niebuhria linearis.
Lobbiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 323.—Singapore.
Loddigesii, *Steud. Nom.* ed. II. i. 277.—Hab. ?
longifolia, *Sw. Prod. Veg. Ind. Occ.* 81.—Ind. occ.
longispina, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 176.—Burma.
loranthifolia, *Lindl. in Mitch. Trop. Austral.* 220.—Austral.
lucida, *Banks, ex DC. Prod.* i. 254.—Austral.
luzonensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 324.—Ins. Luzon.
macrocarpa, Ruiz & Pav. ex DC. Prod. i. 250 = petiolaris.
macrophylla, *H. B. & K. Nov. Gen. et Sp.* v. 91.—N. Granat.
macrosperma, *Delile, in Rochet, Sec. Voy. Choa*, 344.—Afr. trop.
magna, Lour. Fl. Cochinch. 330 = Crataeva magna.
malabaria, Wight, ex Steud. Nom. ed. II. i. 277 = Heyneana.
malabarica, Wall. Cat. n. 7000 = floribunda.
mariana, Jacq. Hort. Schoenbr. i. 57. t. 109 = spinosa.
maxima, Roth, Sp. Pl. Nov. 237 = grandis.
membranacea, *Gardn. & Champ. in Hook. Kew Journ.* i. (1849) 241.—China.
membranifolia, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 70.—Burma.
micracantha, *DC. Prod.* i. 247.—Burma; Malaya.

CAPPARIS:—
micracantha, Teijsm. ex Miq. Illustr. Fl. Arch. Ind. 33 = Forsteniana.
micrantha, *A. Rich. Tent. Fl. Abyss.* i. 31.—Abyss.
Mitchellii, *Lindl. in Mitch. Three Exped.* i. 315.—Austral.
mithridatica, *Forsk. Fl. Aegypt. Arab.* 99.—Arabia.
mollis, *H. B. & K. Nov. Gen. et Sp.* v. 88.—Peruv.
molliuscula, Wall. Cat. n. 7006 (= Cansjerae sp.).—Ind. or.
Moonii, *Wight, Illustr. Ind. Bot.* i. 35.—Zeylan.
Morisonia, Sw. Obs. 272 = Morisonia amcricana.
mucronifolia, Boiss. Diagn. Ser. I. i. 3 = spinosa.
multiflora, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 178.—Reg. Himal.; Burma.
Murrayana, J. Grah. in Cat. Pl. Bomb. 9 = spinosa.
Murrayi, Stew. ex Dalz. in Journ. Linn. Soc. xiii. (1873) 74 = spinosa?
nectaria, *Vell. Fl. Flum.* 230; v. t. 107.—Bras.
nemorosa, *Blanco, Fl. Filip.* ed. I. 438.—Ins. Philipp.
nemorosa, *Jacq. Enum. Pl. Carib.* 24.—N. Granat.
nepaulensis, DC. Prod. i. 246; Wall. Cat. n. 6979 = spinosa.
neriifolia, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* xiv. (1884) 180.—Hab.?
nigricans, *Span. in Linndea*, xv. (1841) 165.—Ins. Timor.
nitida, *Ruiz & Pav. ex DC. Prod.* i. 252.—Peruv.
nobilis, *F. Muell. ex Benth. Fl. Austral.* i. 95.—Austral.
noctiflora, Spreng. ex Eichl. in Mart. Fl. Bras. xiii. 1. 285 = nitida.
nummularia, *DC. Prod.* i. 246.—Austral.
oblongifolia, *Forsk. Fl. Aegypt. Arab.* 99.—Arabia.
obovata, Buch.-Ham. ex DC. Prod. i. 248 = grandis.
obovata, Royle, Illustr. Bot. Himal. 73 = spinosa.
obovatifolia, H. B. & K. Nov. Gen. et Sp. v. 92 = tenuisiliqua.
obtusa, Rafin. Sylva Tellur. 109 = C. Cynophallophora.
octandra, Jacq. Am. Sclect. 160. t. 100 = ferruginea.
odorata, *Blanco, Fl. Filip.* ed. I. 439.—Ins. Philipp.
odoratissima, *Jacq. Hort. Schoenb.* t. 110.—Venezuela.
olacifolia, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 178.—Reg. Himal.
olcoides, *Burch. ex DC. Prod.* i. 248.—Afr. austr.
oligandra, Griff. Notul. iv. 577 = floribunda.
orbiculata, *Wall. ex Hook. f. & Thoms. in Fl. Brit. Ind.* i. 176.—Burma.
ornans, *F. Muell. ex Benth. Fl. Austral.* i. 95.—Austral.
ovalifolia, Ruiz & Pav. Fl. Per. t. 432, ined. ex DC. Prod. i. 253 = avicennifolia.
ovalis, Risso, Fl. Nice, 44 = spinosa.
ovata, Bieb. Fl. Taur. Cauc. ii. 1 = spinosa.
ovata, Desf. Fl. Atlant. i. 404 = Fontanesii.
oxyphylla, *Miq. Pl. Jungh.* 397.—Java.
oxyphylla, Wall. Cat. n. 6997 = tenera.
oxysepala, *Wright, ex Radlk. in Sitzb. Math.-Phys. Acad. Muench.* xiv. (1884) 172.—Nicarag.
Pachaca, *H. B. & K. Nov. Gen. et Sp.* v. 93.—Venezuela.
panduriformis, Lam. Encyc. i. 609 = Thylachium panduriforme.
panduriformis, Thou. Obs. Afr. Austr. 26 = Thylachium heterophyllum.
paradoxa, Jacq. Hort. Schoenb. i. 58 = Steriphoma paradoxum.
parviflora, Boiss. Diagn. Ser. I. i. 4 = spinosa.
parviflora, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 176.—Ind. or.
pauciflora, H. B. & K. Nov. Gen. et Sp. v. 89 = C. Cynophallophora.
pauciflora, Presl, Rel. Haenk. ii. 86 = incana.
pedunculuris, *C. & J. Presl, Delic. Prag.* i. 20.—Sicilia.
pedunculosa, *Wall. Cat.* n. 6999.—Ind. or.
peltata, G. Don, Gen. Syst. i. 285.—Ins. Trinitat.
pendula, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 76.—N. Granat.
persicaefolia, A. Rich. Tent. Fl. Abyss. i. 31 = tomentosa.
petiolaris, Banks, ex DC. Prod. i. 249 = frondosa.
petiolaris, *H. B. & K. Nov. Gen. et Sp.* v. 91.—Guayaquil.
platyacantha, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 323.—Java.
pluvialis, Mart. in Flora, xxii. (1839) 1. Beibl. 22 = C. Cynophallophora.

CAPPARIS:—
polyantha, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 76.—N. Granat.
polymorpha, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 227.—Burma.
polymorpha, Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 24 = tomentosa.
populifolia, Willd. ex Eichl. in Mart. Fl. Bras. xiii. 1. 274 = crotonoides.
prninosa, *Griseb. in Goett. Abh.* xxiv. (1879) 18.—Reg. Argent.
puberula, DC. Prod. i. 248 = tomentosa.
pubiflora, *DC. l. c.* 246.—Malaya.
pulcherrima, *Jacq. Enum. Pl. Carib.* 24.—N. Granat.
pumila, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 260.—China; Ind. or.
punctata, *Burch. Trav.* i. 492.—Afr. austr.
punctata, Wall. Cat. n. 7002 = pumila.
pyracantha, *Boj. in Ann. Sc. Nat.* Sér. II. xx. (1843) 59.—Madag.
pyrifolia, *Lam. Encyc.* i. 606.—Ind. or.
pyrifolia, Wight & Arn. Prod. 25 = grandiflora.
quadriflora, DC. Prod. i. 247 = horrida.
quiniflora, *DC. l. c.*—Austral.
quinqueflora, Steud. Nom. ed. II. i. = quiniflora.
racemifera, DC. Prod. i. 248 = grandis.
racemosa, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
racemulosa, *DC. Prod.* i. 248.—Afr. austr.
radiatiflora, Ruiz & Pav. Fl. Per. v. t. 433 = Crataeva radiatiflora.
reflexa, *Thonn. & Schum. in Dan. Vid. Sel. Afh.* iv. (1828) 11.—Guinea.
reticulata, Klein, ex Wall. Cat. n. 6986 = diversifolia.
retusa, *Griseb. in Goett. Abh.* xxiv. (1879) 18.—Am. austr.
retusella, Thw. Enum. Pl. Zeyl. 16, 400 = sepiaria.
Rheedii, DC. Prod. i. 246 = zeylanica.
Richardi, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 463.—Madag.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 69.—Ins. Fiji.
rosea, *Oliver, Fl. Trop. Afr.* i. 99.—Afr. trop.
Rothii, *Oliver, l. c.* 97.—Afr. trop.
rotundifolia, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 185 = zeylanica.
Roxburghii, *DC. Prod.* i. 247.—Ind. or.
roydsiaefolia, *Kurz, in Journ. As. Soc. Beng.* xxxix. (1870) II. 62.—Burma.
rufescens, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 321.—Ins. Sumbawa.
rugulosa, *Turcz. l. c.* 326.—Venezuela.
rupestris, Sibth. & Sm. Fl. Graec. t. 487 = spinosa.
Russeliana, *Wall. Cat.* n. 7001.—Ind. or.
sabiaefolia, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 179.—Reg. Himal.
salaccensis, *Blume, Bijdr.* 54.—Java.
salicifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 17.—Reg. Argent.
saligna, Vahl, Symb. Bot. iii. 66 = C. Cynophallophora.
sandwicensis, Walp. Rep. v. 54 = seq.
sandwichiana, DC. Prod. i. 245 = spinosa.
sarmentosa, *A. Cunn. ex Benth. Fl. Austral.* i. 95.—Austral.
sativa, Pers. Syn. ii. 59 = spinosa.
saxatilis, *H. B. & K. ex DC. Prod.* i. 250.—Venezuela.
scabrida, *H. B. & K. Nov. Gen. et Sp.* v. 95.—Am. austr.
scandens, Vell. Fl. Flum. 230; v. t. 109 = lincata.
Schimperi, *Steud. Nom.* ed. II. i. 278.—Arab.
sciaphila, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 206 = pumila.
securidacea, *Triana & Planch. l. c.* Sér. IV. xvii. (1862) 77.—N. Granat.
sepiaria, *Linn. Syst.* ed. X. 1071.—Ind. or.; Malaya.
sepiaria, Wall. Cat. n. 6993 G = pedunculosa.
sessilis, *Banks, ex DC. Prod.* i. 249.—N. Granat.
Shanesiana, *F. Muell. Fragm.* x. 94.—Austral.
siamensis, *Kurz, For. Fl. Brit. Burma*, i. 63.—Burma.
sicula, Duham. Arbr. i. 159 = spinosa.
sidaefolia, Ruiz & Pav. ex DC. Prod. i. 252 = crotonoides.
sikkimensis, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 181.—Reg. Himal.
siliquosa, Linn. Syst. ed. X. 1071 = jamaicensis.
Sinclairii, *Benth. Bot. Voy. Sulph.* 65. t. 27.—Am. austr.

CAPPARIS:—
singalensis, Korth. ex Miq. Illustr. Fl. Arch. Ind. 28 = pubiflora.
Sodada, R. Br. in Denh. Clapp. & Oudn. Trav. 225 = aphylla.
solanoides, Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 59 = chrysomeia.
speciosa, *Griseb. in Goett. Abh.* xxiv. (1879) 18.—Reg. Argent.
speciosa, Moric. ex Eichl. in Mart. Fl. Bras. xiii. 1. 272 = C. Yco.
spinosa, *Linn. Sp. Pl.* 503.—Reg. Mediterr.; Ind. or.; Oriens.
Sprucei, *Eichl. in Mart. Fl. Bras.* xiii. 1. 281.—Peruv.
stylosa, DC. Prod. i. 246 = divaricata.
subacuta, *Miq. Fl. Ind. Bat.* i. 1. 101.—Borneo.
subbiloba, H. B. & K. Nov. Gen. et Sp. v. 90 = C. Cynophallophora.
subcordata, *Span. in Linnaea*, xv. (1841) 166.—Ins. Timor.
subspinosa, *Roxb. Hort. Beng.* [93]; *Fl. Ind.* ii. 568.—Ins. Molucc.
Swinhoii, *Hance, in Journ. Bot.* vi. (1868) 296.—China.
tarapotensis, *Eichl. in Mart. Fl. Bras.* xiii. 1. 284.—Peruv.
tenera, *Dalz. in Hook. Kew Journ.* ii. (1850) 41.—Ind. or.
tenuiflora, Fl. Brit. Ind. i. 178, sphalm. = terniflora.
tenuifolia, J. F. Gmel. Syst. 1519, sphalm. = seq.
tenuisiliqua, *Jacq. Enum. Pl. Carib.* 24.—N. Granat.
terniflora, DC. Prod. i. 247 = horrida.
tetrasperma, Thw. Enum. Pl. Zeyl. 15 = tenera.
Thonningii, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 236.—Afr. trop.
Thozetiana, *F. Muell. Fragm.* v. 104.—Austral.
tomentosa, *Lam. Encyc.* i. 606.—Afr. trop.
torulosa, Griseb. ex Hemsl. Biol. Centr. Am. Bot. i. 44 = odoratissima.
torulosa, Sw. Prod. Veg. Ind. Oce. 81 = jamaicensis.
trapeziflora, *Span. in Linnaea*, xv. (1841) 165.—Ins. Timor.
triflora, Mill. Gard. Dict. ed. VIII. n. 10 = frondosa.
trifoliata, Roxb. Hort. Beng. 41; Fl. Ind. ii. 571 = Crataeva religiosa.
trifoliata, Spreng. ex DC. Prod. i. 243 = Crataeva tapioides.
trinervia, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 175.—Burma.
triphylla, *Thunb. Prod. Pl. Cap.* 92.—Afr. austr.
Tweediana, *Eichl. in Mart. Fl. Bras.* xiii. 1. 273.—Reg. Argent.
tylophylla, *Spreng. Syst.* iv. *Cur. Post.* 204.—Java.
uberiflora, *F. Muell. Fragm.* ix. 172.—Austral.
umbellata, *R. Br. in DC. Prod.* i. 247.—Austral.
umbellata, Willd. ex Eichl. in Mart. Fl. Bras. xiii. 1. 273 = angustifolia.
umbonata, *Lindl. in Mitch. Journ. Trop. Austral.* 257.—Austral.
uncinata, Edgew. in Journ. As. Soc. Beng. xvi. (1847) II. 1213 = spinosa.
uncinata, Lodd. ex Eichl. in Mart. Fl. Bras. xiii. 1. 270 = jamaicensis.
uncinata, Wall. Cat. n. 6998 = tenera.
undulata, Lodd. ex Loud. Hort. Brit. 215 = Loddigesii.
undulata, *Zeyh. ex Eckl. & Zeyh. Enum.* 14.—Afr. austr.
variabilis, Wall. Cat. n. 7004 (non Capparid.).—Ind. or.
Vellosiana, Mart. in Flora, xxii. (1839) 1. Beibl. 23 = C. Cynophallophora.
verrucosa, *Jacq. Enum. Pl. Carib.* 23.—Ind. occ.; N. Granat.
verrucosa, Wight, ex Wall. Cat. n. 6980 = stylosa.
versicolor, *Griff. Notul.* iv. 577.—Burma.
verticillaris, *Turcz. in Bull. Soc. Nat. Mosc.* (1863) i. 552.—Bras.
vimiuea, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 179.—Reg. Himal.; Burma.
viridiflora, H. B. & K. Nov. Gen. et Sp. v. 92 = tenuisiliqua.
Volkameria, *DC. Prod.* i. 247.—Afr. austr.
Wallichiana, Wight & Arn. Prod. 25 = zeylanica.
Wightiana, Wall. Cat. n. 6987 = zeylanica.
Yco, *Mart. Reise*, ii. 599; *Mart. & Eichl. in Mart. Fl. Bras.* xiii. 1. 272.—Bras.

CAPPARIS:—
Zeyheri, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 324.—Afr. austr.
zeylanica, *Linn. Sp. Pl.* ed. II. 720.—Ind. or.
zeylanica, Roxb. Hort. Beng. 47; Fl. Ind. ii. 567 = horrida.
zeylanica, Wight & Arn. Prod. 25 = acuminata.
Zippeliana, *Miq. Illustr. Fl. Arch. Ind.* 25. t. 14.—N. Guinea.

CAPRAEA, Opiz, Seznam, 25 (1852) = **Salix**, Tourn.
acuminata, Opiz, l. c. = S. acuminata.
aurita, Opiz, l. c. = S. aurita.
cinerea, Opiz, l. c. = S. cinerea.
grandifolia, Opiz, l. c. = S. grandifolia.
vulgaris, Opiz, l. c. = S. Caprea.

CAPRARIA, [Tourn.] Linn. Gen. ed. I. 28, 75 (1737). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 959.
POGOSTOMA, Schrad. Ind. Sem. Hort. Goett. (1831). XUAREZIA, Ruiz & Pav. Prod. 24. t. 4 (1794).
aegyptiaca, Steud. & Hochst. ex Endl. Nov. Dec. 23 = Anticharis arabica.
arabica, Steud. & Hochst. ex Steud. Nom. ed. II. i. 278 = Anticharis arabica.
biflora, *Linn. Sp. Pl.* 628.—Am. trop.
calycina, A. Gray, in Proc. Am. Acad. vi. (1862) 49 = Myoporum debile.
crustacea, Linn. Mant. 87 = Vandellia crustacea.
cuneata, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 45 = saxifragaefolia.
diffusa, Roxb. Hort. Beng. 47; Fl. Ind. iii. 93 = Ebermaiera thysoidea.
dissecta, Delile, Fl. Egypte, 239. t. 32. f. 2 = Sutera glandulosa.
durantifolia, Linn. Syst. ed. X. 1116 = Stemodia durantaefolia.
gratioloides, Linn. l. c. 1117 = Ilysanthes gratioloides.
gratissima, Roxb. Hort. Beng. 47; Fl. Ind. iii. 92 = Limnophila Roxburghii.
hirsuta, H. B. & K. Nov. Gen. et Sp. ii. 355 = saxifragaefolia.
humifusa, Buch.-Ham. ex Wall. Cat. n. 3883 = Centranthera humifusa.
humilis, [Soland. in] Ait. Hort. Kew. ed. I. ii. 354 = Stemodia parviflora.
integerrima, *Miq. in Linnaea*, xxii. (1849) 476.—Guiana.
integrifolia, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 20.—Mexic.
lanceolata, Hort. ex Steud. Nom. ed. II. i. 278 = salicifolia.
lanceolata, Linn. f. Suppl. 284 = Freylinia lanceolata.
lanceolata, *Vahl, Ecl. Am.* ii. 47.—N. Granat.
lucida, [Soland. in] Ait. Hort. Kew. ed. I. ii. 353 = Teedia lucida.
mexicana, Moric. ex Benth. in DC. Prod. x. 429 = biflora.
Monnieria, Roxb. Hort. Beng. 47 = Herpestis Monnieria.
multifida, Michx. Fl. Bor. Am. ii. 22. t. 35 = Conobea multifida.
multiflora, Steud. Nom. ed I. 148 = Conobea multifida.
peruviana, Benth. in DC. Prod. x. 430 = biflora.
pusilla, Torr. in Ann. Lyc. N. York, i. (1824) 36 = Mimulus floribundus.
rigida, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 301 = Centranthera hispida.
rigida, *Thunb. Prod. Pl. Cap.* 103.—Afr. austr.
salicifolia, Link & Otto, Ic. Pl. Rar. Hort. Berol. 11. t. 4 = Freylinia cestroides.
salicifolia, Salisb. Prod. 94 = Freylinia lanceolata.
saxifragaefolia, *Cham. & Schlecht. in Linnaea*, v. (1830) 105.—Mexic.
semiserrata, *Vahl, Ecl. Am.* ii. 47.—Am. mer.
undulata, *Linn. f. Suppl.* 284.—Afr. austr.
uniflora, *Burm. f. Fl. Ind.* 133.—Ind. or.

CAPRIFICUS, Gasp. Nov. Gen. Fic. 6 (1844) = **Ficus**, Tourn. (Urticac.).
gigantea, Gasp. Ric. Caprif. 80 = F. fulva.
insectifera, Gasp. Nov. Gen. Fic. 6 = F. Carica.
leucocarpa, Gasp. Ric. Caprif. 80. t. 2, 3 = F. albescens.
oblongata, Gasp. l. c. = F. neapolitana.
pedunculata, Gasp. l. c. = F. pedunculata, *Miq.*
rugosa, Gasp. l. c. = F. rugosa, *Miq.*
sphaerocarpa, Gasp. l. c. = F. globosa, *Miq.*

CAPRIFOLIUM, Tourn. ex Linn. Syst. ed. I (1735) = **Lonicera**, Linn. (Caprifol.).
alpigenum, Gaertn. Fruct. i. 136 = L. alpigena.
alpinum, Lam. Fl. Fr. iii. 367 = L. alpigena.
arboreum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 9.—Malaya.
atropurpureum, Hort. ex C. Koch, Dendrol. ii. II. 8 = L. Caprifolium.
balearicum, Dum.-Cours. Bot. Cult. ed. II. iv. 358 = L. implexa.
bracteosum, Michx. Fl. Bor. Am. i. 105 = L. glauca.
caeruleum, Lam. Fl. Fr. iii. 366 = L. caerulea.
chinense, Wats. ex Loud. Hort. Brit. 79 = L. japonica.
ciliosum, Pursh, Fl. Am. Sept. i. 160 = L. ciliosa.
cristatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 9.—Malaya.
dentatum, Rafin. Atl. Journ. 151.—Am. bor.
dioicum, Roem. & Schult. Syst. v. 260 = L. glauca.
distinctum, Moench, Meth. 501 = L. Periclymenum.
Douglasii, Lindl. in Trans. Hort. Soc. Lond. vii. (1830) 244 = L. hirsuta.
dumetorum, Lam. Fl. Fr. iii. 367 = L. Xylosteum.
etruscum, Roem. & Schult. Syst. v. 261 = L. etrusca.
flavum, Ell. Sketch, i. 271 = L. flava.
flexuosum, Hort. ex Steud. Nom. ed. II. i. 278 = L. japonica.
Fraseri, Pursh, Fl. Am. Sept. i. 160 = L. flava.
germanicum, Delarb. Fl. Auv. ed. II. 128 = L. Caprifolium.
germanicum, Quer, Fl. Esp. t. 79 = L. etrusca.
germanicum, Roehl. Deutschl. Fl. i. 175 = L. Periclymenum.
glaucum, Moench, Meth. 502 = L. glauca.
gratum, Pursh, Fl. Am. Sept. i. 161 = L. grata.
hirsutum, Denson, ex Loud. Hort. Brit. Suppl. iii. 509 = L. hirsuta.
hispidulum, Dougl. ex Lindl. Bot. Reg. t. 1761 = L. hispidula.
hortense, Lam. Fl. Fr. iii. 365 = L. Caprifolium.
implexum, Roem. & Schult. Syst. v. 261 = L. implexa.
italicum, Medic. Pfl. Anat. 97 = L. Caprifolium.
japonicum, D. Don, Prod. Fl. Nep. 140 = L. macrantha.
japonicum, Roem. & Schult. ex Steud. Nom. ed. II. i. 278 = L. longiflora.
javanicum, Blume, Bijdr. 653 = L. javanica.
longiflorum, Sabine, in Bot. Reg. t. 1232 = L. longiflora.
Loureiri, Blume, Bijdr. 653 = L. confusa.
macranthum, D. Don, Prod. Fl. Nep. 140 = L. macrantha.
nepalense, G. Don, ex Loud. Hort. Brit. 79 = L. macrantha.
oblongifolium, Sweet, Hort. Brit. ed. II. 259 = L. oblongifolia.
occidentale, Lindl. Bot. Reg. t. 1457 = L. ciliosa.
ovatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 9 = Clerodendron serratum.
pallidum, Schur, Enum. Pl. Transs. 243 = L. Caprifolium.
paniculatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 9.—Malaya.
parviflorum, Pursh, Fl. Am. Sept. i. 161 = L. glauca.
perfoliatum, Roehl. Deutschl. Fl. i. 175 = L. Caprifolium.
Periclymenum, Delarb. Fl. Auv. ed. II. 129 = L. Periclymenum.
pilosum, H. B. & K. Nov. Gen. et Sp. iii. 427. t. 298 = L. pilosa.
pubescens, Goldie, in Edinb. Phil. Journ. (Apr. 1822) 323 = L. hirsuta.
pyrenaicum, Lam. Fl. Fr. iii. 366 = L. pyrenaica.
quercifolium, Meigen, Syst. Verz. 35 = L. Periclymenum.
reclinatum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 9.—Malaya.
roseum, Lam. Fl. Fr. iii. 368 = L. japonica.
rotundifolium, Moench, Meth. 501 = L. Caprifolium.
rubrum, Rafin. Fl. Ludov. 78.—Am. bor.
rupestris, Rafin. New Fl. Am. iii. 17.—Am. bor.
scarlatinum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 9.—Malaya.
sempervirens, Moench, Meth. Suppl. 194 = L. sempervirens.
sericeum, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 9.—Malaya.
sylvaticum, Lam. Fl. Fr. iii. 365 = L. Caprifolium.
viticifolium, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 9.—Malaya.
vulgare, Medic. Pfl. Anat. 387 = L. Caprifolium.
Xylosteum, Gaertn. Fruct. i. 135 = L. Xylosteum.

CAPRIOLA, Adans. Fam. ii. 31 (1763) = **Cynodon**, Pers. (Gramin.).

CAPROXYLON, Tussac, Fl. Antill. iv. t. 30 (1827) = **Hedwigia**, Sw. (Burserac.).
Hedwigii, Tussac, l. c. = H. balsamifera.

CAPSELLA, Medic. Pflanzeng. i. 85 (1792). *CRUCIFERAE*, Benth. & Hook. f. i. 86.
BURSA, [Web. in] Wigg. Prim. Fl. Holsat. 47 (1780).
BURSA-PASTORIS, [Tourn.] Rupp. Fl. Jen. ed. Hall. 87 (1745), nomen prius.
HYMENOLOBUS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 117 (1838).
MARSYPOCARPUS, Neck. Elem. iii. 91 (1790).
MICROLEPIDIUM, F. Muell. in Linnaea, xxv. (1852) 371.
acutifolia, Rafin. New Fl. Am. ii. 27 = C. Bursa-pastoris?
agrestis, Jord. Diagn. i. 339 = C. Bursa-pastoris.
agrestis-rubella, Paill. in Mém. Soc. Emul. Doubs, Sér. IV. vi. (1872) 80 = C. Bursa-pastoris.
amblodes, Rafin. New Fl. Am. ii. 28 = C. Bursa-pastoris.
Andraeana, *F. Muell. in Wing, South. Sc. Record* (March, 1885).—Austral.
antipoda, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 34.—Austral
antipoda, F. Muell. Pl. Vict. i. 44 = australis.
apetala, Opiz, in Flora, v. (1822) 269 = C. Bursa-pastoris.
australis, *Hook. f. ex Benth. Fl. Austral.* i. 81.—Austral.
Berengeriana, Hort. Petrop. ex Bull. Phys.-Math. Acad. Pétersb. xiv. (1856) 234 = procumbens.
bifida, *Rafin. New Fl. Am.* ii. 28 = C. Bursa-pastoris?
blennodina, F. Muell. Pl. Vict. i. 42 = Blennodia alpestris.
Bursa, Rafin. New Fl. Am. ii. 28 = C. Bursa-pastoris.
Bursa-pastoris, *Medic. Pflanzeng.* 85.—Reg. temp.
ceratocarpa, Moench, Meth. 271 = Thlaspi ceratocarpon.
cochlearina, *F. Muell. Pl. Vict.* i. 51.—Austral.
cornigera, Medic. Pflanzeng. 86 = Thlaspi ceratocarpon.
dentata, Rafin. New Fl. Am. ii. 28 = C. Bursa-pastoris?
divaricata, *Walp. Rep.* i. 175.—Am. bor. occ.
Drummondii, *F. Muell. Fragm.* xi. 26.—Austral.
elliptica, C. A. Mey. in Ledeb. Fl. Alt. iii. 199 = procumbens.
erecta, Walp. Rep. i. 175 = divaricata.
furcata, Rafin. New Fl. Am. ii. 28 = C. Bursa-pastoris?
gracilis, Gren. in Mém. Soc. Emul. Doubs, Sér. III. ii. (1858) 403 = C. Bursa-pastoris.
grandiflora, *Boiss. Diagn.* Ser. I. i. 76.—Archip. Graec.
humistrata, *F. Muell. Fragm.* xi. 25.—Austral.
hybrida, *Albert, in Bull. Soc. Sc. Draguignan* (1884) [reimpr. 9].—Gallia.
integrifolia, Rafin. New Fl. Am. ii. 27 = C. Bursa-pastoris.
mexicana, *Hemsl. Diagn. Pl. Nov.* 19.—Mexic.
pastoris, Rupr. Fl. Caucas. 128 = C. Bursa-pastoris.
pauciflora, Koch, in Sturm, Deutschl. Fl. Heft x. t. 66 = Hutchinsia pauciflora.
pastoralis, Dulac, Fl. Hautes-Pyr. 189 = C. Bursa-pastoris.
pilosula, *F. Muell. Vict. Pl.* i. 44.—Austral.
polymorpha, Cav. Desc. 411 = C. Bursa-pastoris.
poimenobalantion, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 71 = C. Bursa-pastoris.
praecox, Jord. Diagn. i. 342 = C. Bursa-pastoris.
procumbens, *Fries, Novit. Fl. Suec. Mant.* i. 14.—Reg. temp.
pubens, *Benth. & Hook. f. ex S. Wats. Bibliog. Ind. N. Am.* 52.—Am. bor. occ.
puberula, *Rupr. Fl. Caucas.* 129.—Reg. Caucas.
rubella, *Reut. in Comptes-Rend. Soc. Hallér.* (1853-54) 18.—Europ.
rubello-agrestis, Paill. in Mém. Soc. Emul. Doubs, Sér. IV. vi. (1872) 80 = C. Bursa-pastoris.
rubescens, Personnat, in Bull. Soc. Bot. Fr. vii. (1860) 511 = rubella.
ruderalis, Jord. Diagn. i. 340 = C. Bursa-pastoris.
sabulosa, Jord. l. c. 341 = C. Bursa-pastoris?
Schaffneri, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 322.—Mexic.
spinosa, Moench, Meth. Suppl. 89 = Lepidium spinosum.

CAPSELLA:—
stenocarpa, *Timb. in Bull. Soc. Hist. Nat. Toulouse*, iii. (1869) 115.—Gallia.
tasmanica, *F. Muell. Fragm.* xi. 26.—Ins. Tasman.
Thomsoni, *Hook. f. in Journ. Linn. Soc.* v. (1861) 172.—Tibet. occ.
triangularis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69 = C. Bursa-pastoris.
virgata, Jord. Diagn. i. 339 = C. Bursa-pastoris.

CAPSICUM, [Tourn.] Linn. Syst. ed. I (1735). *SOLANACEAE*, Benth. & Hook. f. ii. 892.
abyssinicum, *A. Rich. Tent. Fl. Abyss.* ii. 96.—Abyssin.
aggregatum, *Willd. ex Roem. & Schult. Syst.* iv. 809.—Am. bor.
angulosum, Mill. Gard. Dict. ed. VIII. n. 4 = annuum?
angustifolium, *Dun. in DC. Prod.* xiii. 1. 420.—Ind. or.
annuum, *Linn. Sp. Pl.* 188.—Reg. trop.
anomalum, *Franch. & Sav. Enum. Pl. Jap.* ii. 452.—Japon.
Axi, Vell. Fl. Flum. 61; ii. t. 6 = annuum.
baccatum, Hort. Genev. ex Dun. in DC. Prod. xiii. 1. 420 = angustifolium.
baccatum, Buch.-Ham. ex Wall. Cat. n. 2644 = minimum.
baccatum, H. B. & K. Nov. Gen. et Sp. iii. 39 = cumanense.
baccatum, *Linn. Mant.* i. 46.—Reg. trop.
baccatum, Rodsch. Obs. 38 = globiferum.
baccatum, Vell. Fl. Flum. 60; ii. t. 3 = frutescens.
Bauhini, *Dun. in DC. Prod.* xiii. 1. 428.—Hab.?
bicolor, Jacq. Fragm. 66. t. 99. f. 1 = nigrum.
caerulescens, *Bess. Cat. Hort. Crem.* 27.—Hab.?
campylopodium, *Sendt. in Mart. Fl. Bras.* x. 144.—Bras.
cerasiflorum, *Link, Enum. Hort. Berol.* i. 190.—Hab.?
cerasiforme, Hort. ex Dun. in DC. Prod. xiii. 1. 420 = baccatum.
cerasiforme, Mill. Gard. Dict. ed. VIII. n. 5 = annuum?
cerasiforme, Willd. Enum. Hort. Berol. i. 242 = grossum.
ceratocarpum, *Fingerh. Monog.* 22.—Hab.?
cereolum, Bertol. Hort. Bonon. Pl. Nov. i. 6. t. 2 = frutescens.
chamaecerasus, Nees, in Trans. Linn. Soc. xvii. (1837) 65 = grossum.
chinense, *Jacq. Hort. Vindob.* iii. 38. t. 67.—Hab.?
chlorocladum, *Dun. in DC. Prod.* xiii. 1. 415.—Mexic.
ciliare, Willd. Enum. Hort. Berol. 243 = microcarpum.
Comarum, Vell. Fl. Flum. 60; ii. t. 2 = frutescens.
conicum, Lam. Illustr. n. 2390 = conoideum.
conicum, *G. F. W. Mey. Prim. Fl. Esseq.* 112.—Guiana.
conicum, Vell. Fl. Flum. ii. t. 9 = baccatum.
conoides, Roem. & Schult. Syst. iv. 562 = frutescens.
conoideum, Mill. Gard. Dict. ed. VIII. n. 1 = annuum?
cordiforme, Mill. l. c. n. 2 = annuum?
crispum, *Dun. in DC. Prod.* xiii. 1. 415.—Ins. Maurit.
cumanense, *Fingerh. Monog.* 17.—Venezuela.
curvipes, *Dun. in DC. Prod.* xiii. 1. 423.—Guiana.
cydoniforme, Hort. ex Roem. & Schult. Syst. iv. 561 = tetragonum.
dichotomum, *Vell. Fl. Flum.* 61; ii. t. 9.—Bras.
dulce, *Hort. ex Dun. in DC. Prod.* xiii. 1. 428.—Reg. trop.
fastigiatum, Blume, Bijdr. 705 = minimum.
flexuosum, *Sendt. in Mart. Fl. Bras.* x. 143.—Bras.
frutescens, *Linn. Sp. Pl.* 189.—Reg. trop.
frutescens, Rodsch. Obs. 38 = conicum.
glandulosum, *Dun. in DC. Prod.* xiii. 1. 417.—Bras.
globiferum, *G. F. W. Mey. Prim. Fl. Esseq.* 113.—Guiana.
globosum, Bess. Cat. Hort. Crem. 27 = grossum.
gracilipes, *Dun. in DC. Prod.* xiii. 418.—Bras.
grossum, *Linn. Mant.* i. 47; *Roxb. Hort. Beng.* 17; *Fl. Ind.* i. 574.—Reg. trop.
grossum, Willd. Enum. Hort. Berol. 241 = annuum.
Hamiltonii, G. Don, Gen. Syst. iv. 447 = nigrum.
havanense, *H. B. & K. Nov. Gen. et Sp.* iii. 49.—Am. austr.
hispidum, *Dun. in DC. Prod.* xiii. 1. 419.—Texas.
Hornemanni, *Dun. l. c.* 429.—Hab.?

CAPSICUM:—
inaequale, *Vell. Fl. Flum.* 61; ii. t. 5.—Bras.
laurifolium, *Dun. in DC. Prod.* xiii. 1. 418.—Ind. occ.; Bras.
leucocarpon, *Mill. ex Fingerh. Monog.* 32.—Hab.?
longum, Bouton, ex Dun. in DC. Prod. xiii. 1. 414 = conoides.
longum, DC. Cat. Hort. Monsp. (1813) 86 = annuum.
luteum, *Lam. Illustr.* n. 2392.—Ind. or.
Maximowiczii, *Regel & Rach, Ind. Sem. Hort. Petrop.* (1858) 40.—Hab.?
micranthum, *Link, Enum. Hort. Berol.* i. 190.—Bras.
microcarpum, *Cav. Desc.* 371; *DC. Cat. Hort. Monsp.* 86.—Ind. occ.
microphyllum, Dun. in DC. Prod. xiii. 1. 421 = baccatum.
Milleri, Roem. & Schult. Syst. iv. 563 = annuum?
minimum, *Blanco, Fl. Filip.* ed. I. 133.—Ins. Philipp.
minimum, Mill. Gard. Dict. ed. VIII. n. 10 = baccatum, microcarpum.
minimum, *Roxb. Hort. Beng.* 17; *Fl. Ind.* i. 574.—Reg. trop.
mirabile, *Mart. Fl. Bras.* x. 144.—Bras.
Naranca, Hort. Matr. ex Dun. in DC. Prod. xiii. 1. 414 = nigrum.
nigrum, *Willd. Enum. Hort. Berol.* 242.—Hab.?
odoratum, Steud. Nom. ed. II. i. 279 = frutescens.
odoriferum, Vell. Fl. Flum. 61; ii. t. 8 = frutescens.
oliveaeforme, Mill. Gard. Dict. ed. VIII. n. 6 = annuum?
ovatum, *DC. Cat. Hort. Monsp.* (1813) 86.—Hab.?
oxycarpum, *Dun. in DC. Prod.* xiii. 1. 426.—Bras.
parvifolium, *Sendt. in Mart. Fl. Bras.* x. 145.—Bras.
pendulum, *Willd. Enum. Hort. Berol.* i. 242.—Cuba.
pomiferum, Mart. ex Steud. Nom. ed. II. i. 279 = grossum.
pubescens, Dun. in DC. Prod. xiii. 1. 415 = chlorocladum.
pubescens, *Ruiz & Pav. Fl. Per.* ii. 30.—Peruv.
pulchellum, Salisb. Prod. 134 = baccatum.
purpureum, Vahl, ex Hornem. Hort. Hafn. i. 224 = grossum.
pyramidale, Mill. Gard. Dict. ed. VIII. n. 7.—Aegypt.
quitense, Willd. ex Roem. & Schult. Syst. iv. 809 = violaceum.
Rabenii, *Sendt. in Mart. Fl. Bras.* x. 145.—Bras.
salicifolium, *Dun. in DC. Prod.* xiii. 1. 418.—Bras.
Schottianum, *Sendt. in Mart. Fl. Bras.* x. 143.—Bras.
silvestre, Vell. Fl. Flum. 60; ii. t. 1 = annuum.
sinense, Murr. Syst. ed. XIV. 226 = chinense.
sphaerium, Willd. Enum. Hort. Berol. i. 241 = annuum?
strictum, *Fingerh. Monog.* 21.—Am. austr.
testiculatum, *Vis. ex Dun. in DC. Prod.* xiii. 1. 424.—Hab.?
tetragonum, Mill. Gard. Dict. ed. VIII. n. 3 = annuum?
tomatiforme, Fingerh. ex Steud. Nom. ed. II. i. 279 = dulce.
torulosum, Hort. Matr. ex Hornem. Hort. Hafn. Suppl. 27 = pyramidale.
torulosum, *Vell. Fl. Flum.* 60; ii. t. 4.—Bras.
Tournefortii, Bess. Cat. Hort. Crem. (1811) 27 = annuum.
toxicarium, Poepp. ex Fingerh. Monog. 32 = frutescens.
umbilicatum, Vell. Fl. Flum. 61; ii. t. 7 = annuum.
ustulatum, *Paxt. Mag. Bot.* v. 197.—Chili.
villosum, *Sendt. in Mart. Fl. Bras.* x. 144.—Bras.
violaceum, Desf. Tabl. Hort. Par. ed. I. 70; Desv. in Ham. Prod. 25 = nigrum.
violaceum, *H. B. & K. Nov. Gen. et Sp.* iii. 49.—Ecuador.

CAPURA, Blanco, Fl. Filip. ed. I. 264 (err. typ. 644). *SAPINDACEAE*, Benth. & Hook. f. i. 405.
OTOLEPIS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 572.
OTOPHORA, Blume, Rumphia, iii. 142 (1850).
nigrescens, Vidal, Sinops. t. 34. f. E = pinnata.
pinnata, *Blanco, Fl. Filip.* ed. I. 264.—Ins. Philipp.

CAPURA, Linn. Mant. ii. 149 (1771) = **Wikstroemia**, Endl. (Thymelac.).
purpurata, Linn. l. c. 225 = W. indica.

CAQUEPIRIA, J. F. Gmel. Syst. 651 (1791) = **Gardenia**, Ellis (Rubiac.).
Bergkia, J. F. Gmel. l. c. = G. Thunbergia.

CARACALLA, Tod. Ind. Sem. Hort. Panorm. (1861) 32; et in Lem. Illustr. Hortic. ix. (1862) Misc. 13 = **Phaseolus**, Tourn. (Legumin.).
pulcherrima, Tod. ll. cc. = P. Caracalla.

CARACHERA, Juss. in Dict. Sc. Nat. vii. 11 (1817) = **Charachera**, Forsk. (Inc. sed.).

CARADESIA, Rafin. New Fl. Am. iv. 80 (1836) = **Eupatorium**, Tourn. (Compos.).
pauciflora, Rafin. l. c. = E. pauciflorum, *Rafin.*

CARAGANA, Lam. Encyc. i. 615 (1783). *LEGUMINOSAE*, Benth. & Hook. f. i. 505.
acaulis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 44.—Afghanist.
Altagana, Poir. Encyc. Suppl. ii. 89 = microphylla.
ambigua, *Stocks, in Hook. Kew Journ.* iv. (1852) 145.—Beluchist.
arborescens, *Lam. Encyc.* i. 615.—Sibir.
arenaria, *Donn, ex Sims, Bot. Mag.* t. 1886.—Sibir.
argentea, Lam. Encyc. i. 616 = Halimodendron argenteum.
brevispina, *Benth. in Royle, Illustr. Bot. Himal.* 198.—Reg. Himal.; Afghan.
Bungei, *Ledeb. Fl. Alt.* iii. 264.—Sibir. alt.
Chamlagu, *Lam. Encyc.* i. 616.—China.
chinensis, Turcz. ex Maxim. Prim. Fl. Amur. 470 = C. Chamlagu.
conferta, *Benth. ex Baker, in Hook. f. Fl. Brit. Ind.* ii. 116.—Tibet. occ.
crassicaulis, *Benth. ex Baker, l. c.* 117.—Reg. Himal.
cuneata, *Baker, l. c.*—Reg. Himal.
cuneata, Moench, Meth. 135 = frutescens.
digitata, Lam. Encyc. i. 616 = frutescens.
ferox, Lam. l. c. 315 = spinosa.
flava, Poir. Encyc. Suppl. ii. 90 = Robinia flava.
frutescens, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 365.—Reg. Caucas.; Soongaria.
frutex, C. Koch, Dendrol. i. 48 = seq.?
fruticosa, Bess. Cat. Hort. Crem. 116 = arborescens.
Gerardiana, *Benth. in Royle, Illustr. Bot. Himal.* 198.—Reg. Himal.
glomerata, Hort. ex C. Koch, Dendrol. i. 48 = microphylla.
gracilis, *Hort. ex C. Koch, l. c.* 52.—Hab.?
grandiflora, *DC. Prod.* ii. 268.—As. Min.; Reg. Casp.
Halodendrum, Hoffmgg. Verz. Pfl. 46 = Halimodendron argenteum.
inermis, Moench, Meth. 135 = arborescens.
jubata, *Poir. Encyc. Suppl.* ii. 89.—Sibir.
microphylla, *Lam. Encyc.* i. 615.—Sibir. alt.
mollis, Bess. Enum. Pl. Volh. 29 = frutescens.
mongolica, *Fisch. ex Steud. Nom.* ed. II. i. 279.—Mongolia.
mongolica, Hort. ex C. Koch, Dendrol. i. 47 = arborescens.
Moorcroftiana, Benth. in Royle, Illustr. Bot. Himal. 198 = Sophora Moorcroftiana.
nubigena, Bunge, Monog. Astrag. 240 = crassicaulis.
parvifolia, *Hoffmgg. Preisv. Pfl.* viii. *Anh.* 39; *ex Walp. Rep.* i. 681.—Hab.?
polyacantha, *Benth. in Royle, Illustr. Bot. Himal.* 198.—Reg. Himal.
pygmaea, *DC. Prod.* ii. 268.—Reg. Himal.; As. bor.
quadrifolia, Medic. Pfl.-Anat. 62 = frutescens.
Redowskii, *DC. Mém. Lég.* 94. t. 11. f. 45.—Sibir.
rosea, Turcz. ex Maxim. in Prim. Fl. Amur. 470 = frutescens.
sibirica, *Medic. in Vorles. Churpf. Phys. Ges.* ii. (1787) 365.—Sibir.
sophoraefolia, Hort. ex C. Koch, Dendrol. i. 47 = arborescens.
sophoraefolia, *Tausch, in Flora*, xxi. (1838) i. *Beibl.* 77.—Hab.?
spinosa, *DC. Prod.* ii. 269.—Sibir.
spinosa, Rich. ex DC. l. c. 314 = Pictetia squamata.
spinosissima, Benth. in Royle, Illustr. Bot. Himal. 198 = Gerardiana.
tenuis, *Hoffmgg. Preisv. Pfl.* viii. *Anh.* 40; *ex Walp. Rep.* i. 681.—Hab.?
tortumensis, *Boiss. ex Tchihat. Asie Min. Bot.* i. 49. nomen.—As. Min.

CARAGANA :—
tragacanthoides, *Poir. Encyc. Suppl.* ii. 90.—Sibir.
triflora, Lindl. Bot. Reg. (1845) Misc. 41 = brevispina ?
ulicina, *Stocks, in Hook. Kew Journ.* iv. (1852) 145.—Beluchist.
versicolor, Benth. in Royle, Illustr. Bot. Himal. 198. t. 34 = pygmaea

CARAGUATA, [Plum.] Lindl. Bot. Reg. sub t. 1068 (1827). *BROMELIACEAE*, Benth. & Hook. f. iii. 668.
DEVILLEA, Bert. ex Schult. f. Syst. vii. p. lxvii. 1229 (1830).
MASSANGIA, E. Morr. in Belg. Hortic. (1877) 59, 199. t. 8.
Andreana, *E. Morr. in Rev. Hortic.* (1884) 247.—N. Granat.
angustifolia, *Baker, in Gard. Chron.* (1884) ii. 616.—N. Granat.
Augustae, *Benth. & Hook. f. Gen.* iii. 668.—Guiana.
Berteroniana, *Schult. f. Syst.* vii. ii. 1229.—Porto Rico.
cardinalis, E. André, in Illustr. Hortic. xxvii. (1880) t. 374; Rev. Hortic. (1883) 12 = lingulata.
coriostachya, *Griseb. in Goett. Nachr.* (1864) 21.—Venezuela.
fulgens, *Griseb. l. c.*—Venezuela.
Fuerstenbergiana, *Kirch. & Wittm. in Berl. Gartenzeit.* (1883) 299.—Ecuador.
latifolia, Beer, Bromel. 78 = lingulata.
lingulata, *Lindl. Bot. Reg.* sub t. 1068.—Am. trop.
macrostachya, *Bello, in Anal. Soc. Esp. Hist. Nat.* xii. (1883) 122.—Porto Rico.
musaica, *E. André, in Illustr. Hortic.* xxiv. (1877) t. 268.—N. Granat.
nitida, *Griseb. in Goett. Nachr.* (1864) 21.—Venezuela.
Osyana, *E. Morr. in Belg. Hortic.* xxxv. (1885) 254.—Ecuador.
Peacockii, *E. Morr. l. c.* 82.—Reg. Andin.?
sanguinea, *E. André, in Rev. Hortic.* (1883) 468.—N. Granat.
serrata, Hort. Petrop. ex Antoine, Bromel. 46 = Karatas Scheremetiewii.
serrata, Schult. f. Syst. vii. ii. 1231 = Tillandsia serrata.
sphaerocephala, *Gaudich. Bot. Voy. Bonite*, t. 61.—Hab.?
splendens, Hort. ex Planch. in Fl. des Serres, (1856) 31. t. 1091 = lingulata.
Van-Volxemi, *E. André, in Illustr. Hortic.* xxv. (1878) t. 326.—N. Granat.
Zahnii, *Hook. f. Bot. Mag.* t. 6059.—Am. centr.

CARAIPA, Aubl. Pl. Gui. i. 561 (1775), ex parte. *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 188.
africana, *Oliver, Fl. Trop. Afr.* i. 172.—Afr. trop.
angustifolia, *Aubl. Pl. Gui.* i. 561.—Guiana.
calophylla, *Benth. in Journ. Linn. Soc.* v. (1860) 61.—Bras.
calophylla, Spreng. ex Choisy, in Mém. Soc. Phys. Genèv. xiv. (1855) 166 = glabrata.
costata, *Spruce, ex Benth. in Journ. Linn. Soc.* v. (1860) 61.—Am. austr.
densiflora, *Mart. Nov. Gen. et Sp.* i. 105.—Bras.
fasciculata, *Cambess. in Mém. Mus. Par.* xvi. (1828) 416.—Guiana.
glabrata, *Mart. Nov. Gen. et Sp.* i. 105.—Bras.
grandiflora, *Mart. l. c.*—Bras.
latifolia, *Aubl. Pl. Gui.* i. 561.—Guiana.
laurifolia, *Spruce, ex Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 166.—Bras.
laxiflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 364.—Guiana.
leiantha, *Benth. l. c.*—Guiana.
longifolia, *Aubl. Pl. Gui.* i. 561.—Guiana.
myrciaefolia, Spruce, ex Benth. in Journ. Linn. Soc. v. (1860) 61 = fasciculata.
paniculata, *Mart. Nov. Gen. et Sp.* i. 104.—Bras.
parvifolia, *Aubl. Pl. Gui.* i. 561.—Guiana.
racemosa, *Cambess. in Mém. Mus. Par.* xvi. (1828) 415. t. 3.—Bras.
Richardiana, *Cambess. l. c.* 414.—Guiana.
Sellowii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 578.—Bras.
suaveolens, *Planch. ex Benth. in Journ. Linn. Soc.* v. (1860) 61.—Guiana.
surinamensis, *Miq. Stirp. Surin. Sel.* 75.—Guiana.

CARAIPA :—
tereticaulis, *Tul. in Ann. Sc. Nat.* Sér. III. viii. (1847) 341.—Guiana.
variabilis, *Cambess. in Mém. Mus. Par.* xvi. (1828) 416.—Guiana.

CARAJAEA, Wedd. in DC. Prod. xvii. 84 (1873) = **Castelnavia**, Tul. (Podostemac.).
orthocarpa, Wedd. l. c. 85 = Castelnavia orthocarpa.

CARALLIA, Roxb. ex R. Br. in Flind. Voy. ii. 549 (1814). *RHIZOPHOREAE*, Benth. & Hook. f. i. 680.
BARRALDEIA, Thou. Gen. Nov. Madag. 24 (1809).
BARAULTIA, Spreng. Syst. ii. 313 (1825).
CATALIUM, Buch.-Ham. ex Wall. Cat. sub n. 4880 (1831?).
DEMIDOFIA, Dennst. Schluess. Hort. Malab. 31 (1818).
DIATOMA, Lour. Fl. Cochinch. 296 (1790), nomen prius.
KARE-KANDEL, Adans. Fam. ii. 88 (1763).
PETALOTOMA, DC. Prod. iii. 294 (1828).
SYMMETRIA, Blume, Bijdr. 1130 (1826).
Baraldeia, *Arn. in Ann. Nat. Hist.* i. (1838) 371.—Madag.
calycina, *Benth. in Journ. Linn. Soc.* iii. (1859) 75.—Zeylan.
celebica, *Blume, Mus. Bot. Lugd. Bat.* i. 131.—Ins. Celebes.
cerisopsifolia, Miq. Anal. Bot. Ind. iii. 8 = integerrima.
ceylanica, Arn. in Ann. Nat. Hist. i. (1838) 371 = integerrima.
confinis, Blume, Mus. Bot. Lugd. Bat. i. 129 = lanceaefolia.
corymbosa, Arn. in Ann. Nat. Hist. i. (1838) 371 = integerrima.
cuspidata, *Blume, Mus. Bot. Lugd. Bat.* i. 129.—Borneo.
densiflora, *Griff. Notul.* iv. 668.—Burma.
integerrima, *DC. Prod.* iii. 33.—As. et Austral. trop.
lanceaefolia, *Roxb. Hort. Beng.* [37]; *Fl. Ind.* ii. 481.—Burma; Malaya.
lanceolaria, Wall. Cat. n. 4881 = lanceaefolia.
lucida, Roxb. ex Kurz, For. Fl. Brit. Burm. i. 451 = integerrima.
lucida, Roxb. Hort. Beng. [92]; Fl. Ind. ii. 481 = lanceaefolia.
madagascariensis, *Tul. in Ann. Sc. Nat.* Sér. IV. iv. (1856) 117. t. 6.—Madag.
multiflora, *Blume, Mus. Bot. Lugd. Bat.* i. 131.—Borneo.
multiflora, Miq. Fl. Ind. Bat. i. 1. 594 = floribunda.
obcordata, Wight, ex Walp. Rep. ii. 71 = integerrima.
octopetala, F. Muell. Pl. Austr. Trop. Occid. ex Benth. in Journ. Linn. Soc. iii. (1859) 74 = integerrima.
sinensis, Arn. in Ann. Nat. Hist. i. (1838) 371 = integerrima.
symmetria, Blume, Mus. Bot. Lugd. Bat. i. 130 = integerrima.
timorensis, Blume, l. c. 128 = integerrima.
Tulasnei, *Baill. Adansonia*, iii. (1862-63) 20.—Ins. Marian.
zeylanica, Arn. in Wight, Illustr. Ind. Bot. t. 90 = integerrima.

CARALLUMA, R. Br. in Mem. Wern. Soc. i. (1809) 25. *ASCLEPIADEAE*, Benth. & Hook. f. ii. 782.
adscendens, *R. Br. l. c.*—Ind. or.
attenuata, *Wight, Ic.* iv. 15. t. 1268.—Ind. or.
crenulata, Wall. Pl. As. Rar. i. 6. t. 7 = Boucerosia crenulata.
edulis, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 76.—Ind. or.
fimbriata, *Wall. Pl. As. Rar.* i. 7. t. 8.—Ind. or.; Burma.
subulata, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 267.—Arabia.
umbellata, Haw. Syn. Pl. Succ. 47 = Boucerosia umbellata.

CARAMANICA, Tineo, Pl. Rar. Sic. 3 (1847?) = **Taraxacum**, Linn. (Compos.).
taraxacoides, Tineo, l. c. 4 = T. minimum.

CARAMBOLA, Adans. Fam. ii. 508 (1763) = **Averrhoa**, Linn. (Oxalid.).

CARANDA, Gaertn. Fruct. ii. 17. t. 83. f. 5 (1791) (Gen. dub. fructu deformato).
pedunculata, Gaertn. l. c.—Zeylan.

CARANDAS, Rumph. ex Adans. Fam. ii. 171 (1763) = **Carissa**, Linn. (Apocyn.).

CARANGA, Vahl, Enum. i. 100 (1805) = **Curanga**, Juss. (Scrophular.).
amara, Vahl, l. c. = Curanga amara.

CARAPA, Aubl. Pl. Gui. ii. Suppl. 32. t. 387 (1775). *MELIACEAE*, Benth. & Hook. f. i. 338.
PERSOONIA, Willd. Sp. Pl. ii. 331 (1800).
RACAPA, M. Roem. Syn. Hesper. 123 (1846).
TOULOUCOUNA, M. Roem. l. c. (1846).
XYLOCARPUS, Koen. ex A. Juss. in Mém. Mus. Par. xix. (1830) 243.
ZELEA, Hort. ex Tenore, in Atti 3ª Reun. Sc. Ital. (1841) 504.
ZURLOA, Tenore, l. c. 503 (1841).
cauliflora, Poepp. & Endl. Nov. Gen. et Sp. iii. 38 = Gaurea Poeppigii.
guianensis, *Aubl. Pl. Gui. Suppl.* 32. t. 387.—Guiana; Afr. trop.
guineensis, Sweet, Hort. Brit. ed. I. 72; Juss. in Mém. Mus. Par. xix. (1830) 242 = procera.
guyanensis, Oliver, Fl. Trop. Afr. i. 336 = procera.
indica, A. Juss. in Dict. Sc. Nat. vii. 31 = moluccensis.
latifolia, Willd. ex C. DC. in DC. Monog. Phan. i. 718 = guianensis.
moluccensis, *Lam. Encyc.* i. 621.—Geront. trop.
nicaraguensis, *C. DC. in DC. Monog. Phan.* i. 717.—Am. centr.
obovata, Blume, Bijdr. i. 179 = moluccensis.
procera, *DC. Prod.* i. 626.—As. et Afr. trop.
Rumphii, Kostel. Allg. Med.-Pharm. Fl. v. 1988 = moluccensis?
sericea, Poepp. & Endl. Nov. Gen. et Sp. iii. 39. t. 245 = Guarea trunciflora.
surinamensis, *Miq. Stirp. Surin. Sel.* 75. t. 19.—Guiana.
Touloucouna, Guill. & Perr. Fl. Seneg. Tent. i. 128 = procera.
trijuga, Willd. ex C. DC. in DC. Monog. Phan. i. 564 = Guarea humilis.

CARAPICHEA, Aubl. Pl. Gui. 167. t. 64 (1775) = **Cephaëlis**, Sw. (Rubiac.).
Aubletii, DC. Prod. iv. 536 = Ceph. involucrata.
guianensis, Aubl. Pl. Gui. i. 168 = Ceph. involucrata.
Kappleri, Miq. Stirp. Surin. Sel. 181.—Guiana.
Patrisii, DC. Prod. iv. 536 = Ceph. Patrisii.

CARARA, Medic. Pflanzeng. 34 (1792) = **Senebiera**, DC. (1799) (Crucifer.).
Coronopus, Medic. l. c. 35 = S. Coronopus.

CARAXERON, Vaill. ex Rafin. Fl. Tellur. iii. 38 (1836) = **Philoxerus**, R. Br. (Amarantac.).
brasiliense, Rafin. l. c. = Telanthera dentata.
conicus, Rafin. l. c. = P. conicus.
diffusus, Rafin. l. c. = Gomphrena diffusa.
vermicularis, Rafin. l. c. = P. vermicularis.

CARBENIA, Adans. Fam. ii. 116 (1763) (Carbeni). *COMPOSITAE*, Benth. & Hook. f. ii. 482.
BENEDICTA, Bernh. Syst. Verz. Erf. 108 (1800).
benedicta, *Adans. l. c.* 116.—Reg. Mediterr. et Caucas.

CARCIA, Raeusch. Nom. ed. III. 275 (1797) = **Garcia**, Rohr. (Euphorb.).

CARDA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid?).

CARDAMINDUM, Tourn. ex Adans. Fam. ii. 388 (1763) = **Tropaeolum**, Linn. (Geraniac.).
majus, Moench, Meth. Suppl. 20 = T. majus.
minus, Moench, l. c. 21 = T. minus.

CARDAMINE, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 176 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 69.
DENTARIA, [Tourn.] Linn. Syst. ed. I (1735).
HETEROCARPUS, Phil. in Bot. Zeit. xiv. (1856) 641.
KARDANOGLYPHOS, Schlecht. in Linnaea, xxviii. (1856) 472.
PTERONEURON, DC. Prod. i. 150 (1824).

CARDAMINE:—
acaulis, *Berg, in Bot. Zeit.* xiv. (1856) 874.—German.
acris, *Griseb. Spicil. Fl. Rumel.* i. 253.—Macedon.; Armen.
affinis, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 137.—Chili.
africana, *Linn. Sp. Pl.* 655.—Ind. or.; Afr.; Japon.
allevia, Comm. ex DC. Syst. ii. 252 = africana.
alpina, *Willd. Sp. Pl.* iii. 481.—Europ.
amara, Bieb. Fl. Taur. Cauc. ii. 109 = uliginosa.
amara, Lam. Encyc. ii. 185 = pratensis.
amara, *Linn. Sp. Pl.* 656.—Europ.; As. bor.
amplexicaulis, *Haensel. in Bot. Zeit.* iv. (1846) 312.—Hispan.
angulata, *Hook. Bot. Misc.* i. (1830) 343. t. 69.—Am. bor. occ.
angulata, Regel, in Bull. Soc. Nat. Mosc. xxxiv. (1861) ii. 172 = hirsuta.
angulata, Torr. in Pacif. Rail. Rep. iv. 65, 77 = paucisecta.
angustifolia, *Rafin. Fl. Ludov.* 84.—Louisiana.
anteniquana, Burch. ex DC. Syst. ii. 252 = africana.
antiscorbutica, Banks, ex Grisch. in Goett. Abh. vi. (1854) 115 = glacialis.
apetala, Gilib. Fl. Lituan. ii. 67 = impatiens.
apetala, Moench, Meth. 259 = hirsuta.
appendiculata, *Franch. & Sav. Enum. Pl. Jap.* ii. 281.—Japon.
arenosa, Roth, Man. Bot. 926 = Arabis arenosa.
armoracioides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 293.—Venezuela.
articulata, Pursh, Fl. Am. Sept. ii. 430 = Parrya nudicaulis.
asarifolia, *Linn. Sp. Pl.* 654.—Europ.
auriculata, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 319.—Calif.
axillaris, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 290.—Bolivia.
ballidifolia, Hill, Herb. Brit. ii. 275 = bellidifolia.
belgaumensis, Dalz. ex Hook. in Hook. Kew Journ. iv. (1852) 294 = subumbellata.
bellidifolia, All. Fl. Pedem. i. 260. t. 18. f. 3 = alpina.
bellidifolia, *Linn. Sp. Pl.* 654.—Reg. bor. et arct.
Bergeriana, Andrz. ex DC. Syst. ii. 266 = glauca
Berro, *Steud. Nom.* ed. II. i. 280.—Ins. Juan. Fern.
bicolor, Opiz, ex J. & C. Presl, Fl. Čech. 136 = amara.
Bielzii, Schau. in Verh. Siebenb. Ver. Naturw. iv. (1853) 64 = amara.
Bocconi, Vis. Prod. Fl. Cors. App. 4 = thalictroides.
bonariensis, Juss. ex Pers. Syn. ii. 195 = flaccida.
borbonica, Boj. Hort. Maurit. 11 = hirsuta.
borbonica, Pers. Syn. ii. 195 = africana.
borealis, Andrz. ex DC. Syst. ii. 256 = prorepens.
borealis, Laest. ex Nym. Consp. 37 = amara.
Boryi, Boiss. Elench. 9 = Sisymbrium pinnatifidum.
brachycarpa, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 83.—Japon.
brachycarpa, Opiz, Natural. xi. (1826) 411 = impatiens.
bracteata, *S. Moore, in Journ. Bot.* xvi. (1878) 130.—Japon.
brasiliensis, *D. Dietr. Syn. Pl.* iii. 698.—Bras.
Breweri, *S. Wats. in Proc. Am. Acad.* x. (1875) 339.—Am. bor. occ.
buchtormensis, Willd. ex DC. Syst. ii. 258 = pratensis.
bulbifera, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 101.—Europ.
Burchellii, Spreng. Syst. ii. 886 = africana.
calabrica, *Archang. in Nuov. Giorn. Bot.* x. (1878) 163.—Italia.
calbucana, *Phil. in Anal. Univ. Chil.* (1872) 668.—Chili.
caldcirarum, *Guthn. in Seub. Fl. Azor.* 43.—Ins. Azores.
carnosa, *Waldst. & Kit. Pl. Rar. Hung.* ii. 137. t. 129.—Thessalia.
caucasica, Willd. ex Ledeb. Fl. Ross. i. 126 = uliginosa.
chelidonia, S. G. Gmel. It. iv. 108 = impatiens.
chelidonia, Lam. Encyc. ii. 183 = latifolia.
chelidonia, *Linn. Sp. Pl.* 655.—Italia; Hungar.
chelidonia, Pall. Reise, iii. 34 = macrophylla.
chelidonioides, *S. Moore, in Journ. Bot.* xvi. (1878) 130.—Japon.
chenopodifolia, *Pers. Syn.* ii. 195.—Am. austr.
chilensis, *DC. Syst.* ii. 254.—Chili.
cireaeoides, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 144.—Reg. Himal.

CARDAMINE:—
Clematitis, *Shuttl. ex S. Wats. Bibliog. Ind. N. Am. Bot.* i. 53.—Am. bor.
cognata, *Steud. in Flora*, xxxix. (1856) 409.—Chili.
colchaguensis, *Barn. in C. Gay, Fl. Chil.* i. 115.—Chili.
cordata, *Barn. l. c.* 109.—Chili.
cordifolia, *A. Gray, Pl. Fendl.* 8.—N. Mexic.
corsica, Jord. ex Nym. Consp. 38 = graeca.
corsica, *Sieber, ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 293.—Ins. Corsica.
corydaloides, Guss. ex Spreng. Syst. ii. 888 = glauca.
corymbosa, Hook. f. in Ic. Pl. t. 686 = hirsuta.
crassifolia, Opiz, Natural. xi. (1826) 412 = amara.
cretica, Nym. Consp. 38 = graeca.
croatica, Schott, Nym. & Kotschy, Analect. Bot. 46 = carnosa.
cuneata, *Greene, in Bull. Calif. Acad.* n. 3 (1885) 74.—Calif.
Cupani, Jord. ex Nym. Consp. 38 = graeca.
curvisiliqua, *Shuttl. ex Chapm. Fl. S. U. St. Suppl.* 205.—Florida.
dalmatica, Nym. Consp. 38 = maritima.
dasycarpa, Bieb. Fl. Taur. Cauc. iii. 437 = C. impatiens.
dasyloba, Bernh. ex Steud. Nom. ed. II. i. 280 = macrophylla.
debilis, Banks, ex DC. Syst. ii. 265 = hirsuta.
debilis, D. Don, Prod. Fl. Nep. 201 = hirsuta.
decumbens, *Barn. in C. Gay, Fl. Chil.* i. 109.—Chili.
decurrens, *Zoll. & Moritzi, Syst. Verz. Zoll.* 35.—Java.
demissa, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 60.—N. Granat.
dentariaefolia, Royle, ex Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 145 = macrophylla.
dentata, Schult. Obs. 126 = pratensis.
depressa, *Hook. f. Fl. Antarct.* 6.—N. Zel.
dictyosperma, *Hook. Journ. Bot.* i. (1834) 246.—Austral.
digitata, Richards. in Frankl. Narr. 1st Journ. 743 = pratensis?
diphylla, *Wood, Bot. & Fl.* 37.—Am. bor.
divaricata, Hook. f. Fl. N. Zel. i. 13 = stylosa.
diversifolia, Sternb. & Hoppe, in Denkschr. Bot. Ges. Regensb. i. (1815) 156. t. 2 = Arabis Halleri.
drymeja, Schur, Enum. Pl. Transs. 47 = hirsuta.
duraniensis, *Revel, ex Des Moul. in Actes Linn. Soc. Bord.* xx. (1855) 466.—Gallia.
elegantula, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 146.—Reg. Himal.
Engelmanniana, Ind. Sem. Hort. Berol. (1840); ex Heynh. Nom. ii. 118 = Arabis ludoviciana.
enneaphylla, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 101.—Europ.
Eschscholtziana, Andrz. ex Ledeb. Fl. Ross. i. 128 = purpurea.
eustylis, *F. Muell. in Trans. Vict. Inst.* i. (1854) 114.—Austral.
faëroënsis, Hornem. in Fl. Dan. t. 1392 = Arabis petraea.
fagetina, Schur, Enum. Pl. Transs. 47 = umbrosa.
fastigiata, *Hook. f. Handb. New Zeal. Fl.* 13.—N. Zel.
flaccida, Bert. ex C. Muell. in Walp. Ann. vii. 100 = macrorrhiza.
flaccida, *Cham. & Schlecht. in Linnaea*, i. (1826) 21.—Chili.
flavescens, *Phil. in Anal. Univ. Chil.* (1872) 667.—Chili.
flexuosa, With. Bot. Arr. Brit. Pl. ed. III. iii. 578 = sylvatica.
fluminensis, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 185. t. 3.—Bras.
foliosa, Wall. Cat. n. 4779 = macrophylla.
fontana, Lam. Fl. Fr. ii. 499 = Nasturtium officinale.
fontinalis, Schur, Enum. Pl. Transs. 48 = pratensis.
fossicola, Godet, Suppl. Fl. Jura, 13 = pratensis.
Gambellii, *S. Wats. in Proc. Am. Acad.* xi. (1876) 147.—Calif.
gelida, *Schott, in Oestr. Bot. Wochenbl.* (1855) 145.—Tyrol.
geraniifolia, *DC. Syst.* ii. 268.—Reg. Magellan.
gilanensis, Willd. ex DC. l. c. 262 = impatiens.
glacialis, *DC. l. c.* 264.—Reg. Magellan.
glandulosa, *Blanco, Fl. Filip.* ed. I. 521.—Ins. Philipp.
glauca, *Spreng. ex DC. Syst.* ii. 266.—Calabria.
glaucescens, Reichb. Ic. Pl. Crit. v. Index, 4 (Quid?).—Europ.

CARDAMINE:—
Gmelini, Tausch, in Flora, xix. (1836) 402 = macrophylla.
gongylodes, *Phil. in Linnaea*, xxviii. (1856) 664.—Chili.
graeca, Janka, ex Nym. Consp. 38 = longirostris.
graeca, *Linn. Sp. Pl.* 655.—Reg. Mediterr. or.
grandiflora, Hallier, in Bot. Zeit. xxiv. (1866) 209 = pratensis.
grandiflora, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 238 = tenuifolia, *Turcz.*
grandis, Schur, Enum. Pl. Transs. 49 = amara.
granulata, *Phil. in Linnaea*, xxxiii. (1864–65) 8.—Chili.
granulosa, All. Auct. Fl. Pedem. 16 = pratensis.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 146.—Reg. Himal.
Hamiltonii, G. Don, Gen. Syst. i. 167 = hirsuta.
hamulosa, Bertol. Mant. Fl. Alp. Apuan. 43 = resedifolia.
hastata, Willd. ex Ledeb. Fl. Ross. i. 125 = uliginosa.
hastulata, Sm. Engl. Bot. t. 469 = Arabis petraea.
hastulata, Steud. Nom. ed. II. i. 280 = resedifolia.
hederacea, DC. Syst. ii. 264 = thalictroides.
herbivaga, Jord. Diagn. i. 129 = pratensis.
heterophylla, Bory, in Ann. Sc. Gén. Phys. iii. (1820) 6 = Sisymbrium pinnatifidum.
heterophylla, Hook. Comp. Bot. Mag. i. (1835) 273 = hirsuta?
heterophylla, Host, Syn. Pl. Austr. 366 = resedifolia.
heterophylla, *Lapeyr. Hist. Abr. Pl. Pyr.* 377.—Mont. Pyren.
heterophylla, *Wood, Bot. & Flor.* 38.—Am. bor.
Heyneana, Welw. ex Reichb. Fl. Germ. Exeurs. 676 = pratensis.
Hilariana, *Walp. Rep.* i. 137.—Bras.
hirsuta, *Linn. Sp. Pl.* 655.—Reg. temp. et subtrop.
hirsuta, Oed. in Fl. Dan. t. 148 = amara.
hirsuta, Pall. ex DC. Syst. ii. 256 = prorepens.
Huetii, Boiss. Diagn. Ser. II. v. 18 = Sisymbrium Huetii.
humilis, *Kit. in Linnaea*, xxxii. (1863) 496.—Hungar.
ibaguense, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 60.—N. Granat.
impatiens, *Linn. Sp. Pl.* 655.—Europ.; As. bor.; Reg. Himal.
impatiens, O. F. Muell. in Fl. Dan. t. 735 = hirsuta.
indica, *Burm. f. Fl. Ind.* 140.—Java.
integrifolia, Gilib. Fl. Lituan. ii. 68 = pratensis.
intermedia, Hook. Ic. Pl. t. 258 = tenuifolia?
intermedia, *Steud. in Flora*, xxxix. (1856) 410.—Chili.
Jamesoni, *Hook. Lond. Journ. Bot.* vi. (1847) 293.—Ecuador.
javanica, *Miq. Illustr. Fl. Archip.* t. 10.—Java.
Keckii ×, A. Kern. in Zeitschr. Ferdinand. Tirol. III. xv. (1870) 280.—Europ.
kopaonikensis, Panč. ex Nym. Consp. 37 = glauca.
laciniata, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 34.—Austral.
laciniata, Steud. Nom. ed. II. i. 281 = Thelypodium laciniatum.
laciniata, *Wood, Bot. & Flor.* 38.—Am. bor.
Lamontii, *Hance, in Journ. Bot.* xiv. (1876) 363.—China.
latesiliqua, *Cheesem. in Trans. N. Z. Inst.* xv. (1883) 298.—N. Zel.
latifolia, Lej. Fl. Spa, ii. 348 = pratensis.
latifolia, *Vahl, Symb. Bot.* ii. 77.—Europ. austr.
laxa, *Benth. Pl. Hartw.* 158.—N. Granat.
Lechleriana, *Steud. in Flora*, xxxix. (1856) 409.—Chili.
legionensis, *Reut. Ind. Sem. Hort. Genev.* (1861) 4.—Hispan.
lenensis, Andrz. in Ledeb. Fl. Alt. iii. 33 = bellidifolia.
Libertiana, Lej. Rev. Fl. Spa, 135 = amara.
lilacina, Hook. Comp. Bot. Mag. i. (1835) 273 = tenuifolia.
litoralis, *Phil. in Anal. Univ. Chil.* (1865) ii. 314.—Chili.
lobata, Moench, Meth. 260 = P. graeca.
longirostris, *Janka, 'Adat. Magyar.* 164.'—*Cf. Fuss, Bot. Jahresb.* (1876) 1061.—Banat.
lucorum, Boiss. & Huet, ex Boiss. Fl. Orient. i. 161 = pectinata.
ludoviciana, Hook. Journ. Bot. i. (1834) 191 = Arabis ludoviciana.

CARDAMINE:—
Lunaria, Linn. Sp. Pl. 656 = Ricotia Lunaria.
lyrata, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 573.—China.
macrophylla, Schur, Enum. Pl. Transs. 49 = amara.
macrophylla, *Willd. Sp. Pl.* iii. 484.—Reg. Himal.; Sibir.; Japon.
macrorrhiza, *Bert. ex Steud. Nom.* ed. II. i. 281.—Chili.
magellanica, *Phil. in Anal. Univ. Chil.* (1872) 666.—Chili.
marginata, *Phil. l. c.* (1865) ii. 324.—Chili.
maritima, *Portenschl. Enum. Pl. Dalm.* 15.—Dalmat.
Matthioli, Moretti, ex Comol. Fl. Com. v. 157 = pratensis.
maxima, *Wood, Bot. & Flor.* 38.—Am. bor.
melananthera, Steud. Nom. ed. II. i. 280, 281 = seq.
melanthera, Stokes, Bot. Mat. Med. iii. 445 = amara.
Menziesii, DC. Syst. ii. 267 = Sisymbrium canescens.
micrantha, Spenn. Fl. Frib. iii. 922 = hirsuta.
microphylla, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 111.—Sibir. or.
microphylla, J. & C. Presl, Delic. Prag. 15 = thalictroides.
minima, *Steud. in Flora*, xxxix. (1856) 410.—Peruv.
minuta, Willd. ex DC. Prod. i. 150 = microphylla.
monticola, *Timb. in Bull. Soc. Hist. Nat. Toulouse*, iii. (1869) 112.—Gallia.
multicaulis, Hoppe, ex Schur, Enum. Pl. Transs. 47 = hirsuta.
multifida, Pursh, Fl. Am. Sept. ii. 440 = Sisymbrium canescens.
multifida, *Wood, Bot. & Flor.* 38.—Am. bor.
muscosa, Vahl, ex DC. Syst. ii. 260 = hirsuta.
nana, *Barn. in C. Gay, Fl. Chil.* i. 108.—Chili.
nana, *C. Muell. in Walp. Ann.* vii. 106.—Peruv.
nasturtiana, Thuill. Fl. Par. ed. II. i. 330 = amara.
Nasturtii, Spreng. Syst. iv. Cur. Post. 241 = nasturtiodes, *D. Don.*
nasturtiifolia, Steud. Nom. ed. II. i. 281 = nasturtioides, *Bert.*; C. Berro.
nasturtioides, *Bert. in 'Merc. Chil. (1829)'*; *Bull. Férussac*, xx. (1830) 109; *Barn. in C. Gay, Fl. Chil.* i. 113.—Chili.
nasturtioides, Camb. in St. Hil. Fl. Bras. Mer. ii. 89 = Hilariana.
nasturtioides, *D. Don, Prod. Fl. Nep.* 201.—Nepal.
nasturtioides, Schur, Enum. Pl. Transs. 48 = pratensis.
Nasturtium, Wallr. Beitr. Fl. Hercyn. 235 = amara?
nemophila, *Phil. in Linnaea*, xxx. (1859-60) 186.—Chili.
nemorosa, *Lej. Fl. Spa*, ii. 62.—Belgium.
nevadensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 295.—Hispan.
nilagirica, Schlecht. ex Hook. f. & T. Anders. in Fl. Brit. Ind. i. 138 = subumbellata.
nipponica, *Franch. & Sav. Enum. Pl. Jap.* i. 281.—Japon.
nivalis, *Gill. ex Hook. Bot. Misc.* iii. (1833) 136.—Chili.
nivalis, Pall. Reise, ii. 740 = Macropodium nivale.
nivalis, Schur, Enum. Pl. Transs. 46 = resedifolia.
nivea, Hook. Comp. Bot. Mag. i. (1835) 273 = dictyosperma.
nudicaulis, Linn. Sp. Pl. 654 = Parrya macrocarpa.
nudicaulis, Pall. ex DC. Syst. ii. 454 = Macropodium nivale.
nudicaulis, S. Wats. Bibl. Ind. N. Am. Bot. 54 = Parrya nudicaulis.
obliqua, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 19.—Abyssin.
occulta, *Hornem. Hort. Hafn. Suppl.* 71.—China.
oligosperma, A. Gray, in Proc. Am. Acad. viii. (1873) 376 = Breweri.
oligosperma, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 85.—Am. bor. occ.
olympica, Boiss. Diagn. Ser. I. viii. 19 = uliginosa.
Opičii, C. & J. Presl, Fl. Čech. 136 = amara.
orophila, *Timb. in Bull. Soc. Hist. Nat. Toulouse*, iii. (1869) 112.—Gallia.
ovata, *Benth. Pl. Hartw.* 158.—N. Granat.
oxycarpa, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 56 = pectinata.
oxyphylla, Andrz. ex Ledeb. Fl. Ross. i. 128 = pratensis?
paludosa, Knaf, in Flora, xxix. (1846) 293 = pratensis.
palustris, Peterm. in Rabenh. Bot. Centralbl. i. (1846) 47 = pratensis.

CARDAMINE:—
paradoxa, *Hance, in Journ. Bot.* vi. (1868) 111.—China.
parviflora, Linn. Syst. ed. X. 1131 = hirsuta.
parviflora, Suter, Fl. Helv. ii. 59 = tetrandra.
paucijuga, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 295 = hirsuta.
paucisecta, *Benth. Pl. Hartw.* 297.—Calif.
pectinata, Pall. ex DC. Syst. ii. 264 = impatiens.
pedata, *Regel & Tiling, in Nouv. Mém. Soc. Nat. Mosc.* xi. (1859) 47.—Sibir.
pensylvanica, Muhl. Cat. 60, ex Willd. Sp. Pl. iii. 486 = hirsuta.
pentaphylla, [R. Br. in] Ait. Hort. Kew. ed. II. iv. 101 = pratensis.
pentaphylla, *Phil. in Linnaea*, xxxiii. (1864-65) 6.—Chili.
petiolaris, DC. Syst. ii. 264 = graeca.
petraea, Linn. Sp. Pl. 654 = Arabis petraea.
petraea, Townson, Trav. Hung. 490, 348 = Arabis arenosa.
picta, Hook. Lond. Journ. Bot. vi. (1847) 292. t. 12 = Porphyrocodon pictum.
pilosa, Willd. ex DC. Syst. ii. 256 = prorepens.
pinnata, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 101.—Europ.
Plumieri, Vill. Prosp. 38 = thalictroides.
polyphylla, D. Don, Prod. Fl. Nep. 201 = macrophylla.
praecox, Pall. ex Ledeb. Fl. Ross. i. 127 = hirsuta.
pratensis, Hook. f. Fl. Tasm. i. 19 = tenuifolia.
pratensis, *Linn. Sp. Pl.* 656.—Reg. temp.
praticola, Jord. Diagn. i. 128 = pratensis.
propinqua, Carm. in Trans. Linn. Soc. xii. (1819) 507 = hirsuta.
prorepens, *Fisch. ex DC. Syst.* ii. 256.—Sibir.
pseudopratensis, Schur, ex Nichols. in Journ. Bot. xviii. (1880) 200 = pratensis.
pubescens, Stev. ex DC. Syst. ii. 256 = prorepens.
punicea, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 295.—Venezuela.
purpurea, *Cham. & Schlecht. in Linnaea*, i. (1826) 20.—Am. bor. occ.
purpurea, Torr. & Gray, Fl. N. Am. i. 667 = paucisecta.
pusilla, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 18.—Abyssin.
pusilla, *Phil. in Linnaea*, xxviii. (1856) 665.—Chili.
pusilla, Schur, Enum. Pl. Transs. 47 = hirsuta.
radicata, *Hook. f. in Ic. Pl.* t. 882.—Tasman.
ramosissima, *Steud. in Flora*, xxxix. (1856) 409.—Chili.
raphanifolia, Pourr. in Mém. Acad. Toul. iii. (1788) 310 = latifolia.
reflexa, *Rafin. Fl. Ludov.* 84.—Am. bor.
Regeliana, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 73 = hirsuta.
reniformis, *Phil. in Anal. Univ. Chil.* (1865) ii. 313.—Chili.
repanda, Sm. ex DC. Syst. ii. 251; Prod. i. 150 = flaccida.
resedifolia, *Linn. Sp. Pl.* 656.—Europ.
rhomboidea, *DC. Syst.* ii. 246.—Am. bor.
rhomboidea, Durand, Fl. Utah, 159 = cordifolia.
rigida, *Bieb. Fl. Taur. Cauc.* iii. 439.—Reg. Caucas.
rivularis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 61 = pratensis.
rostrata, *Griseb. in Goett. Abh.* vi. (1854) 115.—Chili.
rotundifolia, Bigel. Fl. Bost. ed. II. 252 = bellidifolia.
rotundifolia, Torr. & Gray, Fl. N. Am. i. 83, pro parte = rhomboidea.
rubifolia, Sm. ex DC. Syst. ii. 252; Prod. i. 150 = flaccida.
runcinata, Pourr. in Mém. Acad. Toul. iii. (1788) 310 = latifolia, Sisymbrium pinnatifidum.
sarmentosa, *Forst. f. Prod. App.* 92, nomen.—Ins. Pacif.
saxatilis, Salisb. Prod. 269 = impatiens.
scaposa, *Franch. Pl. Davidian.* i. (1884) 33.—Mongol.
scaturiginosa, Wahlenb. ex Steud. Nom. ed. II. i. 281 = pratensis.
Schaffneri, Hook. f. in Hemsl. Diagn. Pl. Nov. 2 = Gambelii.
scutata, *Thunb. in Trans. Linn. Soc.* ii. (1794) 339.—Japon.
senanensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 280.—Japon.

CARDAMINE:—
serbica, Panč. ex Aschers. & Kan. Cat. Corm. Serb. 76 = carnosa.
setigera, *Tausch, in Flora*, xix. (1836) 404.—Hab.?
silvatica, Auct. vide *sylvatica*.
simensis, *Hochst. ex Oliver, Fl. Trop. Afr.* i. 61.—Abyssin.
Solisi, *Phil. in Anal. Univ. Chil.* (1856) 410.—Chili.
spathulata, *Michx. Fl. Bor. Am.* ii. 29.—Carolina.
stellata, Hook. f. Fl. Antarct. 7. t. 4 A = depressa.
stolonifera, Scop. Fl. Carn. ed. II. ii. 22 = Arabis stolonifera.
stolonifera, Tausch, Cat. 5 = pratensis.
strictula, *Steud. in Flora*, xxxix. (1856) 410.—Reg. Magell.
stylosa, *DC. Syst.* ii. 248.—Austral.; N. Zel.
sublyrata, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 73 = Nasturtium sublyratum.
subumbellata, *Hook. ex Hook. f. & T. Anders. in Fl. Brit. Ind.* i. 138.—Ind. or.
sylvatica, Bess. Primit. Fl. Galic. ii. 76 = pratensis.
sylvatica, Link, in Hoffm. Phyt. Blaett. i. (1803) 50 = hirsuta.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* i. 36.—Japon.
tenella, Clark, Trav. ii. 117 = hirsuta.
tenera, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 56 = pectinata.
tenera, *S. G. Gmel. ex C. A. Mey. Verz. Pfl. Cauc.* 179.—Reg. Caucas.; Persia.
tenuifolia, *Hook. Journ. Bot.* i. (1834) 247.—Austral.
tenuifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xv. (1842) 238.—Sibir.
tenuirostris, *Hook. & Arn. Bot. Beech. Voy.* 6.—Chili.
teres, Michx. Fl. Bor. Am. ii. 29 = Nasturtium tanacetifolium.
ternata, Bory, ex DC. Syst. ii. 252 = africana.
tetrandra, *Hegetschw. Fl. Schw.* 641.—Helvet.
thalictroides, *All. Fl. Pedem.* i. 261.—Europ. austr.
tolimensis, *Planch. & Linden, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 59.—N. Granat.
trichocarpa, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 18 = graeca.
trifolia, *Linn. Sp. Pl.* 654.—Europ. austr.
trifolia, Pall. Reise, iii. 316 = amara.
trifolia, Thunb. Fl. Jap. 260 = scutata.
trifolia, Wahlenb. Fl. Lapp. 179 = amara.
trifoliolata, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 145.—Reg. Himal.
triphylla, Pall. Reise, ii. 35 = amara.
tuberosa, Bert. ex Steud. in Flora, xxxix. (1856) 409 = cognata.
tuberosa, *DC. Syst.* ii. 254.—Chili.
udicola, Jord. Diagn. i. 130 = pratensis.
uliginosa, *Bieb. Fl. Taur. Cauc.* iii. 438.—Ross. austr.; Oriens.
umbrosa, *Andrz. in DC. Syst.* ii. 260.—Volhynia.
umbrosa, Lej. Fl. Spa, ii. 63 = amara.
unalaschcensis, Andrz. ex Ledeb. Fl. Ross. i. 127 = hirsuta.
undulata, *Laramb. & Timb. in Bull. Soc. Bot. Fr.* xiv. (1867) 62.—Gallia.
uniflora, Michx. Fl. Bor. Am. ii. 29 = Leavenworthia Michauxii.
valdiviana, *Phil. in Anal. Univ. Chil.* (1865) II. 314.—Chili.
variabilis, *Phil. in Linnaea*, xxxiii. (1864-65) 5.—Chili.
violacea, *Wall. Cat.* n. 4782.—Reg. Himal.
virginica, Linn. Sp. Pl. 656 = Arabis ludoviciana.
virginica, Michx. Fl. Bor. Am. ii. 29 = hirsuta.
viscosa, S. G. Gmel. ex DC. Syst. ii. 216 = Arabis albida.
Volckmanni, *Phil. in Linnaea*, xxxiii. (1864-65) 7.—Chili.
vulgaris, Phil. l. c. xxviii. (1856) 665 = pratensis.
Wiedmanniana, *Boiss. Fl. Orient.* i. 162.—Anatolia.
Wightiana, Wall. Cat. n. 4780 = africana.
Willdenowii, Tausch, in Flora, xix. (1836) 403 = macrophylla.
yezoënsis, *Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 277.—Japon.
Zollingeri, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854 II. 294.—Java?

CARDAMINUM, Moench, Meth. 262 (1794) = **Nasturtium**, R. Br. (Crucifer.).
Nasturtium, Moench, l. c. = N. officinale.

CARDAMOMUM, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2; Salisb. in Trans. Hort. Soc. i. (1812) 282 = **Elettaria**, Maton (Scitamin.).
malabaricum, Pritz. Ic. Ind. 207 = E. Cardamomum.
medium, Schult. Mant. i. 24 = Amomum costatum.
officinale, Salisb. in Trans. Hort. Soc. i. (1812) 282 = E. Cardamomum.

CARDAMON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 338 = **Lepidium**, Linn. (Crucif.).
sativum, Fourr. l. c. = L. sativum.

CARDANTHERA, Buch.-Ham. ex Nees, in DC. Prod. xi. 67 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1074.
ADENOSMA, Nees, in Wall. Pl. As. Rar. iii. 75 (1832).
SYNNEMA, Benth. in DC. Prod. x. 538 (1846), nomen prius.
africana, *Benth. & Hook. f. Gen.* ii. 1075.—Afr. trop.
avana, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 405.—Burma.
balsamica, *Benth. l. c.* 404.—Ind. or.
glutinosa, Buch.-Ham. ex Nees, in DC. Prod. xi. 68 = triflora.
Griffithii, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 404.—Assam; Burma.
justicioides, *S. Moore, in Journ. Bot.* xviii. (1880) 6.—Afr. trop.
longifolia, Buch.-Ham. ex Nees, in DC. Prod. xi. 90 = Hygrophila phlomoides.
pinnatifida, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 405.—Ind. or.
Thwaitesii, *Benth. & Hook. f. Gen.* ii. 1075.—Zeylan.
triflora, *Buch.-Ham. ex Nees, in DC. Prod.* xi. 68.—Ind. or.: Malaya.
uliginosa, *Buch.-Ham. l. c.* 69.—Ind. or.
verticillata, *Benth. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 405.—Ind. or.

CARDARIA, Desv. Journ. Bot. iii. (1813) 163 = **Lepidium**, Linn. (Crucif.).
amplexicaulis, Spach, Hist. Vég. Phan. vi. 544 = L. amplexicaule.
brachypetala, Opiz, Seznam, 26 = L. Draba.
cochlearia, Spach, Hist. Vég. Phan. vi. 542 = L. Draba.
crassifolia, Spach, l. c. 545 = L. crassifolium.
Draba, Desv. Journ. Bot. iii. (1813) 163 = L. Draba.
latifolia, Spach, Hist. Vég. Phan. vi. 546 = L. latifolium.

CARDERINA, Cass. in Dict. Sc. Nat. xxxv. 272 (1825) = **Senecio**, Tourn. (Compos.).
reclinata, Cass. l. c. xlviii. 447 = S. reclinatus.

CARDIA, Dulac, Fl. Hautes-Pyr. 387 (1867) = **Veronica**, Tourn. (Scrophularin.).
agrestis, Dulac, l. c. 388 = V. agrestis.
alpina, Dulac, l. c. 391 = V. alpina.
amplexicaulis, Dulac, l. c. 390 = V. Anagallis.
aphylla, Dulac, l. c. = V. aphylla.
arvensis, Dulac, l. c. 388 = V. arvensis.
Beccabunga, Dulac, l. c. 390 = V. Beccabunga.
ciliata, Dulac, l. c. = V. Chamaedrys.
didyma, Dulac, l. c. 388 = V. didyma.
elliptica, Dulac, l. c. 390 = V. anagalloides.
filiformis, Dulac, l. c. 389 = V. Tournefortii.
fruticulosa, Dulac, l. c. 391 = V. fruticulosa.
latifolia, Dulac, l. c. 390 = V. latifolia.
montana, Dulac, l. c. 389 = V. montana.
multiflora, Dulac, l. c. 391 = V. serpyllifolia.
Nummularia, Dulac, l. c. 392 = V. Nummularia.
obtusifolia, Dulac, l. c. 391 = V. bellidioides.
officinalis, Dulac, l. c. 390 = V. officinalis.
orbicularis, Dulac, l. c. 388 = V. acinifolia.
plana, Dulac, l. c. 391 = V. Teucrium.
praecox, Dulac, l. c. 389 = V. praecox.
prostrata, Dulac, l. c. 391 = V. prostrata.
quadriloba, Dulac, l. c. 389 = V. hederaefolia.
scutellata, Dulac, l. c. = V. scutellata.
serrata, Dulac, l. c. 391 = V. Ponae.
spicata, Dulac, l. c. 392 = V. spicata.
verna, Dulac, l. c. 388 = V. verna.

CARDIACA, [Tourn.] Linn. Syst. ed I (1735) = **Leonurus**, Linn. (Labiat.).
arvensis, Lam. Fl. Fr. ii. 383 = Stachys arvensis.

CARDIACA :—
crispa, Moench, Meth. 401 = Leonurus Cardiaca.
glabra, Gilib. Fl. Lituan. i. 85 = L. tataricus.
Marrubiastrum, Medic. Beobacht. (1783) 127 = L. Marrubiastrum.
quinquelobata, Gilib. Fl. Lituan. i. 84 = L. Marrubiastrum.
Stachys, Medic. Beobacht. (1783) 126 = L. Cardiaca.
sylvatica, Lam. Fl. Fr. ii. 384 = Lamium Galeobdolon.
trilobata, Lam. l. c. 383 = L. Cardiaca.
vulgaris, Moench, Meth. 401 = L. Cardiaca.

CARDIACANTHUS, Schau. in Linnaea, xx. (1847) 714; et in DC. Prod. xi. 331 (1847) = **Jacobinia**, Moric. (Acanthac.).
Neesianus, Schau. l. c. 715 = J. Neesiana.

CARDIANDRA, Sieb. & Zucc. Fl. Jap. 119. t. 65, 66 (1835). *SAXIFRAGEAE*, Benth. & Hook. f. i. 643.
alternifolia, *Sieb. & Zucc. l. c.* 121.—Japon.

CARDIANTHERA, Hance, in Journ. Linn. Soc. xiii. (1873) 114, sphalm. = **Cardanthera**, Buch.-Ham.

CARDINALIS, Rupp. Fl. Jen. ed. Hall. 248 (1745) = **Lobelia**, Linn.

CARDIOCARPUS, Reinw. in Syll. Ratisb. ii. (1828) 14 = **Soulamea**, Lam. (Simarub.).
amarus, Reinw. l. c. = S. amara.

CARDIOCHLAMYS, Oliver, in Hook. Ic. Pl. t. 1403 (1883). *CONVOLVULACEAE*.
madagascariensis, *Oliver, l. c.*—Madag.

CARDIOCRINUM, Lindl. Veg. Kingd. 205 (1847) = **Lilium**, Tourn.

CARDIOGYNE, Bur. in DC. Prod. xvii. 232 (1873) = **Plecospermum**, Tréc. (Urticac.).
africana, Bur. l. c. 233.—Afr. trop.

CARDIOLEPIS, Rafin. Neogen. 1 (1825) = **Rhamnus**, Tourn.
nigra, Rafin. l. c. 1 = R. lanceolata.
obtusa, Rafin. Sylva Tellur. 28 = R. Purshiana.
rubra, Rafin. Neogen. 1 = R. lanceolata.
spinosa, Rafin. l. c.—Am. bor.

CARDIOLEPIS, Wallr. Sched. Crit. 340 (1822) = **Lepidium**, Linn. (Crucif.).
dentata, Wallr. l. c. = L. Draba.

CARDIOLOCHIA, Rafin. ex Reichb. Consp. 85 (1828) = **Aristolochia**, Tourn.

CARDIOLOPHUS, Griff. in Madr. Journ. Sc. iv. (1836) 376 = **Herpestis**, Gaertn. (Scrophular.).
decussata, Griff. l. c. = H. Hamiltoniana.

CARDIONEMA, DC. Prod. iii. 372 (1828) = **Pentacaena**, Bartl. (Illecebrac.).
multicaule, DC. l. c. = P. polycnemoides.

CARDIOPETALUM, Schlecht. in Linnaea, ix. (1834) 328 = **Duguetia**, St. Hil. (Anonac.).
calophyllum, Schlecht. l. c.—Bras.

CARDIOPHORA, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 216 = **Soulamea**, Lam. (Simarub.).
Hindsii, Benth. l. c. = S. amara.

CARDIOPTERIS, Wall. Cat. n. 8033 (1836?).
OLACINEAE, Benth. & Hook. f. i. 355.
PERIPTERYGIUM, Hassk. in Hoev. & De Vriese, Tijdschr. x. (1843) 142.
STOJA, Buch.-Ham. ex Lindl. Introd. Nat. Syst. Bot. ed. II. 82 (1836).
hamulosa, Griff. Notul. iv. 542; Ic. Pl. Asiat. t. 598 = lobata.
javanica, Blume, Rumphia, iii. 206 = lobata.
lobata, *Wall. Cat.* n. 8033.—Ind. or.
moluccana, Blume, Rumphia, iii. 207 = lobata.
quinqueloba, Hassk. Retzia, i. 64 = lobata.
Rumphii, Baill. Adansonia, x. (1871–72) 280 = lobata.
subhamata, Wall. ex Baill. in DC. Prod. xvii. 26 = lobata.

CARDIOSPERMUM, Linn. Syst. ed. I (1735). *SAPINDACEAE*, Benth. & Hook. f. i. 393.
acuminatum, Miq. in Linnaea, xvii. (1843) 579 = microcarpum.
altissimum, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 38.—Bras.
anomalum, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 351.—Bras.
barbicaule, *Baker, in Oliver, Fl. Trop. Afr.* i. 418.—Afr. trop.
canescens, *Wall. Pl. As. Rar.* i. 14.—As. et Afr. trop.
clematideum, *A. Rich. Tent. Fl. Abyss.* i. 100.—Abyss.
coluteoides, *H. B. & K. Nov. Gen. et Sp.* v. 100.—Am. austr.
Corindum, Linn. Sp. Pl. 526 = C. Halicacabum.
corycodes, *Kunze, in Linnaea*, xviii. (1844) 359.—Cuba.
Duarteanum, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 349.—Bras.
elegans, *H. B. & K. Nov. Gen. et Sp.* v. 99.—Peruv.
ferrugineum, *A. Rich. Ess. Fl. Cub.* 276.—Cuba.
glabrum, Schum. & Thonn. Beskr. Guin. Pl. 197 = C. Halicacabum.
grandiflorum, *Sw. Prod. Veg. Ind. Occ.* 64.—Jamaica.
Halicacabum, *Linn. Sp. Pl.* 366.—Reg. trop.
hirsutum, *Willd. Sp. Pl.* iii. 467.—Afr. occ.
hispidum, *H. B. & K. Nov. Gen. et Sp.* v. 100.—Am. austr.
inflatum, Salisb. Prod. 279 = C. Halicacabum.
inflatum, *Vell. Fl. Flum.* 158; iv. t. 22.—Bras.
integerrimum, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 260.—Bras.
Leuarthianum, G. Don, Gen. Syst. i. 657, sphalm. = Duarteanum.
loxense, *H. B. & K. Nov. Gen. et Sp.* v. 104.—Peruv.
macrolophum, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 261.—Venezuela.
macrophyllum, *H. B. & K. Nov. Gen. et Sp.* v. 100.—Venezuela.
microcarpum, H. B. & K. l. c. 104 = C. Halicacabum.
microspermum, E. Mey. in Drège, Zwei Pfl. Docum. 170 = C. Halicacabum.
molle, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 100 = clematideum.
molle, *H. B. & K. Nov. Gen. et Sp.* v. 104.—Mexic.
moniliferum, *Sw. ex Steud. Nom.* ed. II. i. 150.—Ind. occ.
oblongum, A. Rich. Tent. Fl. Abyss. i. 101 = canescens.
ovatum, Wall. Cat. n. 8031 = canescens.
parviflorum, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 351.—Bras.
pilosum, *Vell. Fl. Flum.* 159; iv. t. 26.—Bras.
pilosum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 395.—Bras.
procumbens, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 262.—Bras.
pubescens, Griff. Journ. 108 = canescens.
pubescens, Lag. Gen. et Sp. Nov. 14 = molle.
Schmiedelia, Dalz. & Gibs. Bomb. Fl. 34 = Schmiedelia villosa.
strictum, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 262.—Bras.
tortuosum, *Benth. Bot. Voy. Sulph.* 9.—Calif.
triphyllum, *Vell. Fl. Flum.* 159; iv. t. 25.—Bras.
truncatum, A. Rich. Tent. Fl. Abyss. i. 101 = C. Halicacabum.
velutinum, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 158.—Bras.
vesicarum, Humb. Rel. Hist. i. 39 = coluteoides.
villosum, Dalz. & Gibs. Bomb. Fl. 34 = Schmiedelia villosa.
villosum, *Macfad. Fl. Jamaic.* i. 154.—Ind. occ.
villosum, Mill. ex DC. Prod. i. 602 = C. Halicacabum.

CARDIOSTEGIA, Presl, Epim. Bot. 249 (1849) = **Melhania**, Forsk. (Sterculiac.).
Kotschyi, Presl, l. c. 249 = M. Denhami.

CARDIOSTIGMA, Baker, in Journ. Linn. Soc. xvi. (1878) 102 = **Sphenostigma**, Baker (Irid.).
longispatha, Baker, l. c. = S. longispatha.
nuda, Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 388 = Calydorea nuda.

CARDIOTHECA, Ehrenb. ex Steud. Nom. ed. II. i. 282 (1840) = **Anarrhinum**, Desf. (Scrophularin.).
pubescens, Ehrenb. ex Steud. l. c. = A. pubescens.
virgata, Ehrenb. ex Boiss. Fl. Orient. iv. 362 = A. orientale.

CARDOPATIUM, Juss. in Ann. Mns. Par. vi. (1805) 324 in adn. *COMPOSITAE*, Benth. & Hook. f. ii. 463.
BROTERA, Willd. Sp. Pl. iii. 2399 (1803).
amethystinum, *Spach, in Ann. Sc. Nat.* Sér. III. v. (1846) 242.—Afr. bor.
apulum, Spach, l. c. 240 = corymbosum.
Billardieri, Spach, l. c. 247 = corymbosnm.
Boryi, *Spach, l. c.* 245.—As. Min.
corymbosum, *Pers. Syn.* ii. 500.—Reg. Mediterr. or.; As. Min.
Fontanesii, *Spach, in Ann. Sc. Nat.* Sér. III. v. (1846) 244.—Reg. Tunetan.
orientale, Jaub. & Spach, Illnstr. t. 426 = corymbosum.
orientale, *Spach, in Ann. Sc. Nat.* Sér. III. v. (1846) 237.—Graecia.
Vrionis, Heldr. in Boiss. Fl. Orient. iii. 443 = corymbosum.

CARDUNCELLUS, Adans. Fam. ii. 116 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 483.
LAMOTTEA, Pomel, Mat. Fl. Atl. 3 (1860 ?).
ONOBROMA, Gaertn. Fruct. ii. 380. t. 160 (1791).
acaulis, Presl, Fl. Sic. i. p. xxx = pinnatus.
araneosus, *Boiss. & Reut. Pl. Nov. Hisp. Diagn.* 18.—Hispan.
arborescens, Sweet, Hort. Brit. ed. I. 216 = Carthamus arborescens.
atlanticus, *Coss. & Dur. ex Coss. in Ann. Sc. Nat.* Sér. IV. i. (1854) 240.—Algeria.
atractyloides, *Coss. & Dur. ex Pomel, Nouv. Mat. Fl. Atl.* 278.—Algeria.
caeruleus, *Less. Syn. Comp.* 6.—Reg. Mediterr.
calvus, *Boiss. & Reut. Pugill.* 64.—Afr. bor.
dianius, Webb, Iter Hisp. 33 = caeruleus.
eriocephalus, *Boiss. Diagn.* Ser. I. x. 100.—Afr. bor.
hispanicus, Boiss. ex DC. Prod. vii. 304 = caernleus.
ilicifolius, *Pomel, Nouv. Mat. Fl. Atl.* 277.—Afr. bor.
lucens, *Ball, in Journ. Bot.* xi. (1873) 370.—Marocco.
mitissimus, *DC. Fl. Fr.* iv. 73.—Gallia.
monspeliensis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = seq.
monspeliensinm, *All. Fl. Pedem.* i. 154.—Gall. austr.
pectinatus, *DC. Prod.* vi. 615.—Afr. bor.
pinnatns, *DC. l. c.* 614.—Oriens.
plumosus, *Pomel, Nouv. Mat. Fl. Atl.* 277.—Afr. bor.
rhaponticoides, *Coss. & Dur. ex Pomel, Nouv. Mat. Fl. Atl.* 278.—Algeria.
tingitanus, DC. ex Duby, Bot. Gall. i. 281 = caerulens.
vulgaris, Loud. Hort. Brit. 332 = monspeliensium.

CARDUUS, [Tonrn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 467.
ALFREDIA, Cass. in Bull. Soc. Philom. (1815) 175.
CLAVENA, DC. Prod. vi. 633 (1837).
ONOPYXUS, Bubani, in Nnov. Giorn. Bot. Ital. v. (1873) 319.
POLYACANTHA, Hill, Herb. Brit. i. 69 (17 9).
PTERNIX, Hill, Veg. Syst. iv. 16 (1762).
abyssinicns, *Sch. Bip. in Linnaea*, xix. (1847) 332.—Abyssin.
acanthifolius, Lam. Encyc. i. 703 = Cnicns oleraceus.
Acanthium, Baill. Hist. Pl. viii. 289 = Onopordon Acanthium.
acanthocephalus, *C. A. Mey. Verz. Pfl. Cauc.* 71.—Reg. Cancas.
acanthoides, Gren. & Godr. Fl. Fr. ii. 231, sec. Nym. Consp. 411 = nntans.
acanthoides, Hornem. in Fl. Dan. t. 1341 = Cuicns palustris.
acanthoides, Huds. Fl. Angl. ed. II. ii. 351 = pycnocephalus.
acanthoides, *Linn. Sp. Pl.* 821.—Enrop.; Reg. Cancas.
acanthoides, Pall. ex Bieb. Fl. Taur. Cauc. ii. 268 = hamulosns.
acanthoides, Urv. in Mém. Soc. Linn. Par. i. (1822) 361 = nutans.
acantholepis, *Benth. in Benth. & Hook. f. Gen.* ii. 468. —As.centr.
Acarna, Linn. Sp. Pl. 820 = Cnicus Acarna.
acaulis, Linn. l. c. 1199 = Cnicns acaulis.
acaulis, Thunb. Fl. Jap. 306 = Cuicus japonicns.
acicularis, *Bertol. in Ann. Sc. Nat.* Sér. I. i. (1829) 274.—Italia.
acuminatus, Gaud. ex Reichb. Ic. Fl. Germ. xv. 90. t. 872 = acanthoides.
adpressns, *C. A. Mey. Verz. Pfl. Cauc.* 71.—Reg. Caucas.

CARDUUS:—
afer, Jacq. Hort. Schoenb. ii. 10 = Cnicus afer.
affinis, *Guss. Pl. Rar.* 334. t. 57.—Neapol.
aggregatus, Schleich. Cat. Pl. Helv. ed. IV. 11 = nutans ?
agrestis, Presl, Fl. Sicul. p. xxx, in nota = corymbosns.
Aitonii, Raensch. Nom. ed. III. 229 (Qnid ?).
alatus, D. Don, Prod. Fl. Nep. 167 = Jurinea alata.
albidus, Bieb. Fl. Tanr. Cauc. ii. 269 = pycnocephalns.
albidus, Lam. Fl. Fr. ii. 18 = praec. ?
algeriensis, Munby, in Bull. Soc. Bot. Fr. ii. (1855) 285. —Algeria.
alpestris, Waldst. & Kit. Pl. Rar. Hnng. iii. 296. t. 267 = arctioides.
altaicus, Patrin, ex S. G. Gmel. Fl. Sib. t. 39. f. 2 = Serratula glauca.
altissimus, Gilib. Fl. Litnan. iii. 186 = Cnicns palustris.
altissimus, Linn. Sp. Pl. 824 = Cnicus altissimus.
ambiguus, Pers. Syn. ii. 389 = Cnicus heterophyllns.
ammophilus, *Hoffmgg. & Link, Fl. Port.* ii. 187.—Lusitan.
amplexicaulis, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 11.—Malaya.
Amsteinii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii.-xxiv. (1880) 106.—Helvet.
anglicus, Lam. Encyc. i. 705 = Cnicus pratensis.
angustifolins, *Lam. l. c.* 703.—Hab. ?
antarcticus, Auct. ex Stend. Nom. ed. I. 152 = autareticus.
appeninus, Jan, ex DC. Prod. vi. 622 = nntans.
arabicus, Jacq. Coll. i. 56 = pycnocephalus.
arachnoideus, Bieb. ex Steud. Nom. ed. II. i. 283 = Cnicus arachnoideus.
araneosus, Steud. Nom. ed. I. 151 = Cnicns araneosus.
arctioides, Vill. Hist. Pl. Danph. ii. 22 = Personata.
arctioides, *Willd. Sp. Pl.* iii. 1656.—Enrop.
arenarius, DC. Fl. Fr. vi. 457 = Candollei.
arenarius, *Desf. Fl. Atlant.* ii. 99.—Afr. bor.
Argemone, Pourr. in Lam. Encyc. i. 700 = carlinaefolins.
argentatus, *Linn. Mant.* ii. 280.—Reg. Mediterr. or.; Mesopot.
argentatus, Willd. ex Stend. Nom. ed. II. i. 283 = pycnocephalns.
argiroa, Nym. Consp. 413 = argyroa.
argutus, Sweet, Hort. Brit. ed. I. 213 = Serratula tinctoria.
argyracanthus, Wall. Cat. n. 2903 = Cnicns Wallichii.
argyroa. *Bivona. Stirp. Rar. Sicil.* i. [7].—Sicilia.
argyroa, Kunze, in Flora, xxix. (1846) 760 = malacitanns.
armatus, Boiss. & Heldr. Diagn. Ser. I. vi. 104 = cronius.
armatus, Stend. Nom. ed. I. 151 = Cnicus armatus.
armenus, Boiss. Fl. Orient. iii. 516 = untans.
arvensis, Robs. Brit. Fl. 163 = Cnicus arvensis.
Aschersonianus ×, *Ruhm. in Eichl. Jahrb. Berl.* i. (1881) 239.—German.
astracanicus, *Spreng. Fl. Hal. Mant.* 49.—Astracan.
atlanticus, *Pomel, Nouv. Mat. Fl. Atl.* 22.—Afr. bor.
atriplicifolius, Trevir. Hort. Wratisl. 1820 = Serratula atriplicifolia.
auriculatus, Wall. Cat. n. 2899 = Saussurea hypolenca.
auroficus, Chaix, ex Vill. Fl. Delph. 90 sphalm. = seq.
aurosicns, *Vill. Hist. Pl. Dauph.* iii. 7.—Gallia.
australis, Jord. Gr. Rec. Bot. Gren. (1849) 14 = vivariensis.
australis, Linn. f. Suppl. 348 = pycnocephalus.
autareticus, Vill. Hist. Pl. Dauph. iii. 12 = Cnicns heterophyllus.
axillaris, Gaud. Fl. Helv. ii. 169 = acanthoides.
baeocephalus, *Webb & Berth. Phyt. Canar.* ii. 373.—Ins. Canar.
baeticus, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 63.—Hispan.
Balansae, *Boiss. Diagn.* Ser. II. iii. 44.—Algeria.
Ballii, *Hook. f. in Journ. Bot.* xi. (1873) 368.—Marocco.
Barrelieri, *Bertol. ex Nym. Consp.* 411.—Liguria.
benedictus, Auct. ex Stend. Nom. ed. I. 151 = Carbenia benedicta.
bicolor, Visiani, Fl. Dalm. ii. 48. t. 49. fig. 2 = argyroa.
bihariensis ×, Simk. in Termész. Füzetek, v. (1881) 51. —Transsylv.

CARDUUS:—
Boujarti, Pill. & Mitterp. Iter Poseq. i. 143. t. 13 = Cuicus arachnoideus.
Boujarti, Savi, Fl. Pis. ii. 243 = Cnicus ferox.
Bourgaeanus, Boiss. & Reut. Pugill. Pl. Nov. (1852) 62 = Reuterianus.
Bourgeaenus, *Sch. Bip. ex Boiss. Diagn.* Ser. II. iii. 44. —Ins. Canar.
brachycephalus, Schur, in Verh. Siebenb. Ver. Naturw. x. (1859) 68 = collinns.
bulbosus, Lam. Encyc. i. 705 = Cnicns tuberosns.
caeruleus, *Brot. Fl. Lusit.* i. 342.—Lusitan.
caesius, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 498, nomen.—Rossia.
Caflischii ×, Bruegg. in Jahresb. Naturf. Ges. Granb. II. xxiii.-xxiv. (1880) 106.—Helvet.
callosus, *Steud. Nom.* ed. II. i. 283.—Hab. ?
camporum, *Boiss. Diagn.* Ser. II. iii. 45.—Rumel.
canaliculatus, Dulac, Fl. Hautes-Pyr. 522 = carlinaefolius.
candicans, Waldst. & Kit. Pl. Rar. Hung. i. t. 83 = colliuus.
Candollei, *Moretti, Pl. Ital. Dec.* ii. 10.—Gallia; Ital.
canus, Linn. Mant. 105 = Cnicus canns.
Canduelis, Gren. in Billotia, i. (1864) 14 = arctioides.
carlinaefolius, *Lam. Encyc.* i. 700.—Europ.
carlinaefolius, Nym. Consp. 412 = defloratus.
carlinaefolius, Tenore, Prod. Fl. Neap. 48 = chrysacanthus.
carlinoides, Dur. ex DC. Prod. vi. 625 = confertus.
carlinoides, *Gouan, Illustr.* 62. t. 23.—Mont. Pyren.
carlinoides, Steud. Nom. ed. II. i. 283 = Cnicus carliuoides.
carlinoides, Tenore, Prod. Fl. Neap. 48 = chrysacanthns.
carniolicus, Pers. Syn. ii. 388 = Cnicus rufescens.
carolinianus, Walt. Fl. Carol. 195 = Cnicus mnticns.
Carolorum ×, Howie & Jenner, in Trans. Bot. Soc. Edinb. ix. (1868) 260.—Scotia.
carpetanus, Boiss. & Rent. Diagn. Pl. Hisp. 19. n. 35 = Gayanus.
carthamoides, Stend. Nom. ed. I. 151 = Leuzea carthamoides.
Casabonae, Linn. Sp. Pl. 823 = Cnicns Casabonae.
caucasicus, Adam, in Weber & Mohr, Beitr. i. 64 = Cnicus horridus.
caulescens, Pers. ex Steud. Nom. ed. I. 151 = Cuicns acaulis.
centauroides, Bnch.-Ham. ex DC. Prod. vi. 563 = Volutarella divaricata.
centauroides, D. Don, Prod. Fl. Nep. 167 = Serratnla centauroides.
centauroides, Hoppe, ex DC. Prod. vi. 628 = arctioides.
centauroides, Linn. Sp. Pl. ed. II. 1148 (= *Cnicus cynaroides*).
cephalanthus, *Viv. Fl. Cors. Sp. Nov.* 14.—Sardin.
cerinthifolius, Vill. Prosp. 30 = Serratula nndicaulis.
cerinthoides, Willd. Sp. Pl. iii. 1660 = Serratula nudicaulis.
cernnns, *Bertol. Fl. Guatimal.* 31.—Gnatemal.
cernnns, *Steud. Nom.* ed. I. 151.—Sibir.
Chailleti, Godr. Fl. Lorr. ii. 42 = Cnicus palustris.
chamaecephalns, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 434.—Abyssin.
chinensis, DC. Prod. vi. 629 = Cnicns chinensis.
chius, *Jacq. Hort. Vindob.* iii. 7.—Ins. Chios.
chrysacanthus, *Tenore, Sem. Hort. Neap.* (1825) 12.—Hispan.; Italia.
cichoraceus, Cyr. Pl. Rar. Neap. Fasc. ii. 17. t. 7 = Serratula cichoracea.
ciliatus, Murr. in Comm. Gott. (1784) 35. t. 5 = Cnicns ciliatus.
ciliatus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 193 = Gayanus.
ciliatus, Vill. Préc. Voy. 45 = Cnicus spathulatus.
cinerascens, *Hort. Huber, in Rev. Hortic.* (1872) 60 —Am. bor. ?
cinereus, Bieb. Fl. Taur. Cauc. ii. 270 = pycnocephalus.
cirsiformis, *Vukot. in Rad Jugos. Akad. Zagreb.* xxxix. (1877) 206.—Europ.
cirsioides, Vill. Prosp. 30 = defloratus.
Cirsium, Mill. Gard. Dict. ed. VIII. n. 5 = Cnicus dissectus.
clavulatus, *Link, in Buch, Beschr. Canar. Ins.* 147.—Ins. Canar.
collinus, *Waldst. & Kit. Pl. Rar. Hung.* iii. 257. t. 232.—Europ. austr. or.

CARDUUS :—

comosus, Lam. Encyc. i. 703 = Cnicus spinosissimus.
confertus, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 194 = baeticus.
confertus, Moris, Elench. Pl. Sard. ii. 5 = myriacanthus.
congestus, Guss. in Tenore, Syll. 412 = cephalanthus.
costariensis, *Timb. & Thévén. in Mém. Acad. Toulouse*, Ser. VII. vi. (1874) 347.—Europ.
coronata, D. Don, Prod. Fl. Nep. 167 = Serratula coronata.
corticus, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 373 = clavulatus.
corymbosus, *Tenore, Prod. Fl. Neap.* 48.—Italia.
Cosmelii, Adam, ex DC. Prod. vi. 637, 638 = Cnicus lappaceus.
crassifolius, Willd. Enum. Hort. Berol. 838 = defloratus.
creticus, Lam. Encyc. i. 701 = Cnicus pungens.
crispus, Gouan, Fl. Monsp. 327 = tenuiflorus.
crispus, Guir. ex Nym. Consp. 413 = Kernerianus.
crispus, Huds. Fl. Angl. ed. II. ii. 350 = acanthoides.
crispus, *Linn. Sp. Pl.* 821.—Europ.; As. bor.
cronius, *Boiss. & Heldr. Diagn.* Ser. I. vi. 105.—Graecia; As. Min.
cyanoides, Linn. Sp. Pl. 822 = Jurinea cyanoides.
cyanoides, Pall. Reise, iii. 534 = Jurinea polyclonos.
Cynara, Lam. ex Steud. Nom. ed. II. i. 283 = Cnicus cynaroides.
cynaroides, Lam. Encyc. i. 702 = Cnicus cynaroides.
Decandollei, Auct. ex Steud. Nom. ed. II. i. 283 = Candollei.
defloratus, *Linn. Syst.* ed. X. 1200.—Europ.
demidecurrens ×, Franch. Fl. Cher-et-Loire, 317.—Gallia.
dentatus, Pers. Syn. ii. 389 = Kentrophyllum dentatum.
desertorum, Fisch. ex Steud. Nom. ed. II. i. 283 (cf. *Cirsium desertorum*).—Sibir.
diacantha, Labill. Ic. Pl. Syr. dec. i. 7 = Cnicus diacanthus.
discolor, Hook. Fl. Bor. Am. i. 302 = Cnicus Hookerianus et undulatus.
discolor, Nutt. Gen. Am. ii. 130 = Cnicus altissimus.
dissectus, Linn. Sp. Pl. 822 = Cnicus dissectus.
dissectus, Vill. Hist. Pl. Dauph. iii. 15 = Cnicus pratensis.
divaricatus, Gilib. Fl. Lituan. i. 167 = Cnicus lanceolatus.
diversifolius, *Wall. Beitr. Fl. Hercyn.* 267.—Europ.
dubius ×, Balb. Cat. Hort. Taur. (1813) = crispus.
Durisei, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 64.—Algeria.
echinatus, Desf. Fl. Atlant. ii. 247 = Cnicus echinatus.
echinocephalus, Pers. ex Steud. Nom. ed. I. 152 = Cnicus echinocephalus.
elegans, *Pall. ex Georgi, Beschr. Russ. Reich.* iii. iv. 1224.—Rossia.
elegans, Salisb. Prod. 185 = Cnicus Casabonae.
elegans, Stephan, ex Ledeb. Fl. Ross. ii. 673 = Cousinia wolgensis.
elegans, Willd. ex Steud. Nom. ed. II. i. 283 = Cnicus igniarius.
elongatus, Wall. Cat. n. 2900 = Tricholepis elongata.
eriophorus, Linn. Sp. Pl. 823 = Cnicus eriophorus.
eriophorus, Pall. ex Bieb. Fl. Taur. Cauc. ii. 276 = Cnicus fimbriatus.
eriophorus, Thunb. Fl. Jap. 306 = Cnicus japonicus.
Erisithales, Jacq. Enum. Vindob. 146, 279 = Cnicus ochroleucus.
Erisithales, Vill. Hist. Pl. Dauph. ii. 20 = Cnicus rivularis.
erucagineus, Lam. Encyc. i. 704 = Cnicus ochroleucus!
esdraelonicus, Boiss. Diagn. Ser. I. x. 90 = argentatus.
fasciculiflorus, *Viv. App. Alt. Fl. Cors.* 5. t. 1.—Corsic.; Sardin.
fastigiatus, Moris, Elench. Pl. Sard. ii. 5 = argyroa.
ferox, Vill. Prosp. 30; Lam. Encyc. i. 703 = Cnicus ferox.
fimbriatus, Steud. Nom. ed. II. i. 284 = Cnicus fimbriatus.
firmus, Steud. l. c. ed. I. 152 = Cnicus firmus.
flavescens, *C. Koch, in Linnaea*, xxiv. (1851) 394.—Pontus.
flavescens, Linn. Sp. Pl. 825 = Serratula flavescens.
foliosus, Hook. Fl. Bor. Am. i. 303 = Cnicus foliosus.
Forsteri ×, Bab. Man. Brit. Bot. ed. I. 173.—Anglia.
fruticosus, Desf. Tabl. Hort. Par. ed. I. 91 = Cnicus fruticosus.

CARDUUS :—

fruticosus, Poir. Encyc. Suppl. ii. 186 = Carlina xeranthemoides.
Galactites, Bory & Chaub. Fl. Pélop. 58 = Galactites tomentosa.
Galactites, Lam. Encyc. i. 701 = pycnocephalus.
Gayanus, *Dur. ex DC. Prod.* vi. 625.—Hispan.
getulus, *Pomel, Nouv. Mat. Fl. Atl.* 273.—Afr. bor.
giganteus, Desf. Fl. Atlant. ii. 95. t. 221 = Cnicus giganteus.
Gigas, Ucria, in Roem. Arch. i. 70 = Cnicus giganteus.
glaber, Buch.-Ham. ex Wall. Cat. n. 2907 = Tricholepis radicans.
glaber, Nutt. Gen. Am. ii. 130 = Cnicus muticus!
glaber, Steud. Nom. ed. I. 152 = Cn. spinosissimus!
glaucophyllus, Desf. ex DC. Prod. vi. 622 = defloratus.
glaucus, Baumg. Transs. iii. 54 = defloratus.
glaucus, Cav. Ic. iii. 18. t. 236 = Serratula nudicaulis.
glaucus, *Ledeb. Hort. Dorp. Suppl.* i. 2.—Rossia.
glomeratus, Lam. Fl. Fr. ii. 20 = Cnicus pratensis.
glutinosus, Beck, Bot. N. & Mid. St. 178 = Cnicus muticus.
glycacanthus, Sibth. & Sm. Fl. Graec. Prod. ii. 150 = Jurinea glycacantha.
Gmelini, Steud. Nom. ed. I. 152 = Cnicus acaulis.
gnaphalodes, Cyr. Pl. Rar. Neap. Fasc. i. 27 = Cnicus gnaphalodes.
Gouani, Steud. Nom. ed. I. 152 = Cnicus tuberosus.
granatensis, Willk. in Linnaea, xxx. (1859–60) 113 = platypus.
Grenieri, Sch. Bip. ex Nym. Consp. 412 = C. Personata.
haemorrhoidalis, Auct. ex DC. Prod. vi. 643 = Cnicus arvensis.
hamulosus, *Ehrh. Beitr.* vii. 166.—Europ. austr. or.; Reg. Caucas.
hamulosus, Tenore, Prod. Fl. Neap. 46 = nutans.
hastatus, Lam. Encyc. i. 704 = Cnicus heterophyllus.
helenifolius, Salisb. Prod. 185 = Cnicus heterophyllus.
helenioides, Linn. Sp. Pl. 826 = Cnicus heterophyllus.
heteromallus, D. Don, Prod. Fl. Nep. 166 = Saussurea candicans.
heterophyllus, Linn. Sp. Pl. 824 = Cnicus heterophyllus.
Hippolyti, Bory & Chaub. Exp. Mor. 241. t. 29 = Cnicus pungens.
hispanicus, Bory, ex DC. Prod. vii. 304 = carlinoides.
hispanicus, Lam. Encyc. i. 701 = Cnicus hispanicus.
horridulus, Pers. Syn. ii. 389 = Cnicus horridulus.
horridus, Adam, in Weber & Mohr, Beitr. i. 66 = Cnicus munitus.
hybridus, Steud. Nom. ed. I. 152 = Cnicus ochroleucus.
hymenolepis, *Trautv. in Act. Hort. Petrop.* iii. ii. (1875) 272.—Daghestan.
hyssopifolius, Schur, Enum. Pl. Transs. 395. Index = Serratula heterophylla.
Hystrix, *C. A. Mey. Verz. Pfl. Caucas.* 71.—Persia.
igniarius, Pall. ex DC. Prod. vi. 637 = Cnicus igniarius.
illyricus, Baill. Hist. Pl. viii. 289 = Onopordon illyricum.
inclinans, Stokes, in With. Bot. Arr. Brit. Pl. ed. II. ii. 870 = crispus.
inclinatus, Dulac, Fl. Hautes-Pyr. 502 = medius.
indicus, Klein, ex Wall. in DC. Prod. vi. 564 = Tricholepis angustifolia.
inermis, Pers. Syn. ii. 388 = Cnicus centauroides.
intermedius, Heller, ex Steud. Nom. ed. I. 152 = Cnicus intermedius.
intricatus, Reichb. f. Ic. Fl. Germ. xv. 51. t. 856 = Candollei.
intricatus, *Timb. in Bull. Soc. Bot. Fr.* xv. (1868) p. lxxxvi.—Mont. Pyren.
involucratus, Wall. Cat. n. 2902 = Cnicus involucratus.
italicus, Savi, Bot. Etrus. iii. 144 = Cnicus italicus.
Jacquini, Steud. Nom. ed. I. 152 = Cnicus pyrenaicus.
japonicus, Franch. Pl. David. i. 175 = Cnicus japonicus.
Kabenoe, *Pomel, Nouv. Mat. Fl. Atl.* 273.—Afr. bor.
Killiasii ×, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxiii.–xxiv. (1880) 106.—Europ.
Kosmelii, Adam, in Weber & Mohr, Beitr. i. 66 = Cnicus lappaceus.
laciniatus, Lam. Encyc. i. 706 = Cnicus palustris.
laciniatus, Ledeb. Fl. Ross. ii. 722 = adpressus.
lactifolius, Stokes, Bot. Mat. Med. iv. 141 = Silybum Marianum.
lanatus, *Brot. Fl. Lusit.* 342.—Lusitan.

CARDUUS :—

lanatus, Roxb. ex Poir. Encyc. vi. 505 = Cnicus arvensis.
lanceatus, Hill, Veg. Syst. iv. 14 = Cnicus lanceolatus.
lanceolatus, Linn. Sp. Pl. 821 = Cnicus lanceolatus.
lanceolatus, Lour. Fl. Cochinch. ii. 483 = chinensis.
laniflorus, Steud. Nom. ed. I. 153 = Cnicus laniflorus.
lanuginosus, *Willd. Sp. Pl.* iii. 1646.—As. Min.
lappaceus, Bieb. Casp. 198 = Cnicus lappaceus.
lappoides, Less. in Linnaea, v. (1830) 128 = Cnicus lappoides.
latifolius, Banks, ex Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 28 = *Cirsium latifolium*.—Ins. Madera.
laxiflorus, *Tausch, in Flora*, xii. (1829) Erg. 42.—Austria.
leptacanthus, *Fresen. in Mus. Senckenb.* iii. (1845) 71.—Abyssin.
leptocladus, *Dur. in Duch. Rev. Bot.* i. (1845–46) 362.—Algeria.
leptophyllus, Gaud. Fl. Helv. v. 171 = defloratus.
leucanthemus, Schur, Enum. Pl. Transs. 417 = C. Personata.
leucanthus, Cav. Ic. ii. 61. t. 166 = Serratula flavescens.
leucocephalus, Pers. Syn. ii. 387 = Cnicus leucocephalus.
leucographus, Linn. Sp. Pl. 820 = Tyrimnus leucographus.
leucophyllus, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 194.—Mongolia.
linearis, Thunb. Fl. Jap. 306 = Cnicus linearis.
litigiosus, Moris, Elench. Pl. Sard. i. 26 = pycnocephalus.
litigiosus, Nocca & Balb. Fl. Ticin. ii. 99. t. 11 = Candollei.
litoralis ×, *Borb. in Magyar Növent. Lapok.* ix. (1885) 56.—Europ.
Lomonossowii, *Trautv. in Act. Hort. Petrop.* i. (1871–72) 183.—Mongolia.
lucidus, Moench, Meth. 564 = pycnocephalus.
lucidus, Wall. Cat. n. 2908 = nutans.
lycopifolius, Vill. Prosp. 30 = Serratula heterophylla.
macranthus, Haussk. ex Boiss. Fl. Orient. iii. 436 = Cousinia Hablitzii.
macrocephalus, Coss. in Bourg. ex Willk. & Lange, Prod. Fl. Hisp. ii. 188 = chrysacanthus.
macrocephalus, Desf. Fl. Atlant. ii. 245 = nutans.
macrocephalus, Wall. Cat. n. 2904 = Jurinea macrocephala.
macrolepis, Peterm. in Flora, xxvii. (1844) 492 = nutans!
macropterus, Fisch. ex Spreng. Pugill. i. 64 = Jurinea polyclonos.
malacitanus, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 62.—Hispan.
Mariae, Crantz, Inst. i. 141 = Silybum Marianum.
Marianus, Linn. Sp. Pl. 823 = Silybum Marianum.
marmoratus, *Boiss. & Heldr. Diagn.* Ser. II. vi. 93.—Peloponnes.
Martrinii, Timb. in Marc. Fl. Tarn. 377 = nigrescens.
medius, *Gouan, Illustr.* 62. t. 24.—Mont. Pyren.
medius, Steud. Nom. ed. I. 158 = Cnicus tuberosus.
meonanthus, *Hoffmgg. & Link, Fl. Port.* ii. 198.—Hispan.
mexicanus, Moric. ex DC. Prod. vi. 637 = Cnicus mexicanus.
microcephalus, Gaud. Fl. Helv. v. 168 = tenuiflorus.
microcephalus, *Tenore, Syll. App.* iv. *ex Fl. Nap.* v. 206.—Italia.
mitis, Steud. Nom. ed. I. 153 = Cnicus mitis.
mollis, Bieb. Fl. Taur. Cauc. ii. 270 = Jurinea arachnoidea.
mollis, Gouan, Illustr. 63 = Jurinea humilis.
mollis, Linn. Cent. Pl. Rar. ii. 30 = Jurinea mollis.
mollis, Pollich, Fl. Palat. ii. 421 = Jurinea cyanoides.
mollis, Vill. Hist. Pl. Dauph. ii. 17 = Cnicus acaulis.
monocephalus, *Schur, ex Verh. Siebenb. Ver. Naturw.* iii. (1852) 87.—Europ.
monspessulanus, Linn. Sp. Pl. 822 = Cnicus monspessulanus.
montanus, Pers. Syn. ii. 388 = Cnicus montanus.
montanus, Salis. ex DC. Prod. vi. 622 = Candollei.
montanus, Pollini, Viag. Lag. Gard. 115 = defloratus.
Morisii, Balb. in Moris, Elench. Pl. Sard. 26 = fasciculiflorus.
Moritzii ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii.–xxiv. (1880) 106.—Europ.

CARDUUS:—
moschatus, Guss. Ind. Sem. Hort. Boccadif. (1825) = Jurinea mollis.
multiflorus, Gaud. Fl. Helv. v. 166 = acanthoides.
multiflorus, Panč. ex Nym. Consp. 414 = Pancicii.
multijugus, C. Koch, in Linnaea, xxiv. (1851) 392 = adpressus.
muticus, Pers. Syn. ii. 386 = Cnicus muticns.
myriacanthus, Boiss. Voy. Bot. Espagne, ii. 361 = Reuterianus.
myriacanthus, *Salzm. in DC. Prod.* vi. 624.—Reg. Mediterr. occ.
Naegeli ×, Bruegg. in Jahresb. Nat. Ges. Graub. II. xxiii–xxiv. (1880) 107.—Europ.
neglectus, Steud. Nom. ed. I. 153 = Cnicus uliginosus.
neglectus, Tenore, Ind. Sem. Hort. Neap. (1830) 14 = acicularis.
nepalensis, Spreng. ex DC. Prod. vi. 542 = Jurinea macrocephala.
nervosus, *C. Koch, in Linnaea*, xvii. (1843) 42.—Armenia.
nidulans, *Rupr. Sert. Tiansch.* 55.—As. centr.
nigrescens, Mutel, ex Nym. Consp. 412 = vivariensis.
nigrescens, *Vill. Prosp.* 30.—Europ.
nitidus, Waldst. & Kit. Pl. Rar. Hung. i. 52 = Serratula heterophylla.
nitidus, Wall. Cat. n. 2908 = Tricholepis angustifolia.
niveus, *Benth. in Benth. & Hook. f. Gen.* ii. 468.—As. centr.
nudicaulis, Pers. Syn. ii. 386 = Serratula nudicaulis.
numidicus, *Coss. & Dur. Explor. Pl. Algér.* t. 49.—Algeria.
nutans, Boiss. ex Nym. Consp. 411 = platypus.
nntans, *Linn. Sp. Pl.* 821.—Europ. ; As. temp.
obvallatus, *Bieb. ex Willd. Enum. Hort. Berol.* 842.—Reg. Caucas.
obvallatus, Steud. Nom. ed. I. 153 = Cnicus obvallatus.
obvallatus, Wall. Cat. n. 2095 = Saussurea obvallata.
occidentalis, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 418 = Cnicus occidentalis.
ochroleucus, Pers. Syn. ii. 389 = Cnicus ochroleucus.
odoratus, Muhl. Cat. Darl. Fl. Cestr. 85 = Cnicus pumilus.
oleraceus, Vill. Hist. Pl. Dauph. iii. 21 = Cnicus oleraceus.
olympicus, *Boiss. Diagn.* Ser. II. iii. 43.—As. Min.
onopordoides, *Fisch. ex Bieb. Fl. Taur. Cauc.* iii. 552.—Reg. Caucas.
Onopyxos, *Hill, Veg. Syst.* iv. 13.—German.
orgyalis, Steud. Nom. ed. I. 153 = Cnicus orgyalis.
orientalis, Adam, in Weber & Mohr, Beitr. i. 63 = Cousinia carduiformis.
orientalis, Pers. Syn. ii. 387 = Galactites tomentosa.
orthocephalus ×, Wallr. Beitr. Fl. Hercyn. 268 ; et in Linnaea, xiv. (1840) 86 = acanthoides.
osseticus, Adam, in Weber & Mohr, Beitr. i. 65 = Cnicus fimbriatus.
palatinus ×, Sch. Bip. ex Nym. Consp. 413 = acanthoides.
palustris, Linn. Sp. Pl. 822 = Cnicus palustris.
palustris, Savi, Bot. Etrusc. iii. 141 = Cnicus polyanthemus.
Pancicii, *Sch. Bip. ex Nym. Consp.* 414.—Serbia.
paniculatus, *Ait. Hort. Kew.* ed. I. iii. 143.—Europ.
paniculatus, Dulac, Fl. Hautes-Pyr. 522 = carlinoides.
paniculatus, Vahl, Symb. Bot. i. 68 = Cnicus flavispinus.
pannonicus, Linn. f. Suppl. 348 = Cnicus pannonicus.
pannonicus, Schleich. ex DC. Prod. vi. 629 = tenuifolius.
pannosus, *Trautv. in Act. Hort. Petrop.* iii. II. (1875) 271.—Armenia.
parviflorus, Buch, Beschr. Ins. Canar. 147 = pycnocephalus.
parviflorus, Linn. Mant. 279 = Cnicus parviflorus.
pauciflorus, Pers. Syn. ii. 388 = Cnicus pauciflorus.
pectinatus, Linn. Mant. 279 = defloratus.
pedemontanus, Pers. Syn. ii. 389 = Cnicus tuberosus.
peregrinus, Retz. Obs. i. 27 = pycnocephalus.
Personata, *Jacq. Fl. Austr.* iv. 25. t. 348.—Europ.
personatus, Gaertn. Fruct. ii. 378 = praec.
petrophilus, *Timb. in Bull. Soc. Bot. Fr.* xv. (1868) p. lxxxvii.—Mont. Pyren.
pinnatifidus, Cav. Ic. i. 58. t. 83 = Serratula pinnatifida.
Pitcheri, Steud. Nom. ed. II. i. 284 = Cnicus Pitcheri.
platylepis, Sauter, in Flora, xiii. (1830) 410 = nutans.
platypus, *Lange, Ind. Sem. Hort. Haun.* (1857) 26.—Hispan.
podacanthus, DC. Fl. Fr. iii. 80 = aurosicus.

CARDUUS:—
poliochrus, *Trautv. in Act. Hort. Petrop.* iii. II. (1875) 272.—Reg. Caucas.
polyacanthos, Curt. Fl. Lond. fasc. vi. t. 54 = acanthoides.
polyacanthus, Lam. Fl. Fr. ii. 20 = Cnicus Casabonae.
polyanthemos, *Hegel, ex Heer, Fl. Schw.* 801.—Helvet.
polyanthemos, Schur, Enum. Pl. Transs. 417 = collinus.
polyanthemus, Boeber, in Pall. N. Nord. Beitr. vi. 261 ; Bieb. Casp. App. 210 (= *Cirsium elodes*).—Reg. Caucas.
polyanthemus, Linn. Mant. i. 109 = Cnicus polyanthemus.
polyanthemus ×, Schleich. Spicil. 15 = nutans.
polyanthos, Boiss. Fl. Orient. iii. 548, 988 = Cnicus pungens.
polyanthos, Schreb. Spicil. 15 = acanthoides.
polyclonos, Willd. Sp. Pl. iii. 1655 = Jurinea polyclonos.
polymorphus, Lapeyr. in Mém. Acad. Toulouse, i. (1782) 217. t. 19, 20 = Cnicus heterophyllus.
Pontederae, Steud. Nom. ed. I. 153 = Cnicus rivularis.
Poolii ×, Bruegg. in Jahresb. Nat. Ges. Graub. II. xxiii–xxiv. (1880) 107.—Europ.
pratensis, Huds. Fl. Angl. ed. II. 353 ; Lam. Encyc. i. 700 = Cnicus pratensis.
pseudosyriacus, *Lojac. in Natural. Sicil.* iv. (1885) 109.—Sicil.
ptarmicaefolius, Mill. Gard. Dict. ed. VIII. n. 1 = Cnicus stellatus.
pteracanthus, Dur. in Duch. Rev. Bot. i. (1845–46) 362 = Reuterianus.
pterocaulos, Steph. ex DC. Prod. vi. 553 = Cousinia arachnoidea.
pulcher, *Clark, Trav.* i. 739, nomen.—Tauria.
pumilus, *D. Don, Prod. Fl. Nep.* 167.—Nepal.
pumilus, Hook. Fl. Bor. Am. i. 302 = Cnicus Drummondii.
pumilus, Nutt. Gen. Am. ii. 130 = Cnicus pumilus.
pumilus, Vill. Hist. Pl. Dauph. ii. 17 = Cnicus heterophyllus.
punctatus, Herb. Madr. ex Wall. Cat. n. 2907 = Tricholepis radicans.
pungens, Pers. ex DC. Prod. vi. 643 = Cnicus pungens.
purpureus, *Vill. Fl. Delphin.* 90.—Gallia.
pycnocephalus, *Linn. Sp. Pl.* ed. II. 1151; *Jacq. Hort. Vind.* i. 17. t. 44.—Europ. ; Reg. Mediterr. ; Oriens.
pycnocephalus, Spreng. Syst. iii. 385 = cephalanthus.
pycnocephalus-alatus, Stev. ex DC. Prod. vi. 553 = Cousinia arachnoidea.
pygmaeus, Jacq. Enum. Vindob. 147, 282 = Saussurea pygmaea.
pyrenaicus, Gouan, Illustr. 63 = Cnicus pyrenaicus.
pyrenaicus, *Jacq. Obs. Bot.* iv. 11. t. 95.—Mont. Pyren.
pyrochros, Less. in Linnaea, v. (1830) 130 = Cnicus conspicuus.
quinquefolius, D. Don, Prod. Fl. Nep. 167 = Serratula quinquefolia.
radiatus, Waldst. & Kit. Pl. Rar. Hung. i. 9 = Serratula radiata.
radicans, Roxb. Hort. Beng. 60 ; Fl. Ind. iii. 408 = Tricholepis radicans.
ramosus, Roxb. ll. cc. 101, 407 = Volutarella divaricata.
ramosissimus, *Panč. Elench. Crna-Gora*, 51.—Monteneg.
recurvatus, Jord. ex Bor. Fl. Cent. Fr. ed. III. ii. 359, in obs. = nigrescens.
recurvatus, Willk. in Linnaea, xxx. (1859–60) 113 = granatensis.
remotifolius, Hook. Fl. Bor. Am. i. 302 = Cnicus remotifolius.
repandus, Pers. Syn. ii. 386 = Cnicus repandus.
Reuterianus, *Boiss. Diagn.* Ser. II. iii. 44.—Hispan.
rhaeticus, *Dalla Torre, Atl. Alpenfl.* 138.—Europ.
rheginus, Steud. Nom. ed. I. 153 = Cnicus strictus.
rigens, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 144 = Cnicus rigens.
rigens ×, *Godr. Fl. Lorr.* ii. 40.—Gallia.
rivularis, Jacq. Fl. Austr. i. 57 = Cnicus rivularis.
Rosenii, Vill. Fl. Delphin. 91 = Cnicus acaulis.
ruderalis, Tausch, ex Nym. Consp. 412 = acanthoides.
rufescens, Pers. Syn. ii. 389 = Cnicus rufescens.
salinus, Bieb. ex Steud. Nom. ed. II. i. 353 = Centaurea serratuloides.
salisburgensis, Pers. Syn. ii. 388 = Cnicus rivularis.
samniticus, Tenore, Sem. Hort. Neap. (1825) 12 = Cnicus italicus.
Sanctae-Balmae, Loisel. Nouv. Not. 34 = Candollei.
sardous, DC. Prod. vi. 626 = pycnocephalus.
scaber, Poir. Voy. Barb. ii. 231 = Cnicus giganteus.

CARDUUS:—
Schimperi, *Sch. Bip. in Linnaea*, xix. (1847) 334.—Abyssin.
Schulzeanus ×, Ruhm. in Eichl. Jahrb. i. (1881) 240.—Europ.
Scolymus, Baill. Hist. Pl. viii. 7, 289 = Cynara Cardunculus.
Scordium, Stokes, Bot. Mat. Med. iv. 142 = Serratula Scordium.
segetum, Franch. Pl. Davidian. i. 178 = Cnicus segetum.
seminudus, *Bieb. Fl. Taur. Cauc.* ii. 271.—Reg. Caucas. et Casp.
semipectinatus, Lam. Encyc. i. 705 = Cnicus semipectinatus.
semipinnatus, Desf. ex Steud. Nom. ed. I. 153 = Cnicus pratensis.
serratuloides, Jacq. Fl. Austr. 127 = Cnicus pannonicus.
serratuloides, Linn. Sp. Pl. 825 = Cnicus serratuloides.
serratuloides, Neck. Delic. Gallo-Belg. 338 = Cnicus arvensis.
serrulatus, Steud. Nom. ed. I. 154 ; ed. II. i. 285 = Cnicus serrulatus.
setosus, Bab. in Gard. Chron. (1844) 718 = Cnicus arvensis ?
simplicifolius, *Sang. in Atti Acc. Pont. Lincei*, Ser. I. xviii. (1865) t. 2.—Italia.
sinuatus, Gilib. Fl. Lituan. i. 187 = acanthoides.
Spachianus, *Dur. in Duch. Rev. Bot.* i. (1845–46) 361.—Afr. bor.
spinigerus, Jord. Obs. Pl. Crit. iii. (1846) 215 = hamulosus.
spinosissimus, Gueldenst. ex Ledeb. Fl. Ross. ii. 743 = Cnicus acaulis.
spinosissimus, C. A. Mey. in Beitr. Pfl. Russ. Reich. iv. 43 = Cnicus esculentus.
spinosissimus, Steud. Nom. ed. II. i. 285 = Cnicus echinocephalus.
spinosissimus, Vill. Hist. Pl. Dauph. iii. 11 = Cnicus spinosissimus.
spinosissimus, Walt. Fl. Carol. 194 = Cnicus horridulus.
spinulosus, Bertol. Amoen. Ital. 41 = defloratus.
spinosus, Gueldenst. ex Ledeb. Fl. Ross. ii. 743 = Cnicus acaulis.
spurius, Linn. Hort. Ups. 249 = Cnicus eriophorus.
squarrosus, *DC. ex Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimpr. 18.—Madera.
Stangii, Buek, ex Nym. Consp. 411 = nutans.
stellatus, Linn. Sp. Pl. 823 = Cnicus stellatus.
stenolepis, *Benth. in Benth. & Hook. f. Gen.* ii. 468.—As. centr.
strictus, Steud. Nom. ed. I. 154 = Cnicus strictus.
strigosus, Bieb. Tabl. Casp. 194 = Cnicus strigosus.
suaveolens, *Benth. in Benth. & Hook. f. Gen.* ii. 468.—As. centr.
subcoriaceus, Less. in Linnaea, v. (1830) 130 = Cnicus subcoriaceus.
subdecurrens, *Bertol. Fl. Ital.* viii. 623.—Liguria.
sumanus, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 96 = defloratus.
syncephalus, Reichb. f. Ic. Fl. Germ. xv. 87. t. 864 = pycnocephalus.
syriacus, Linn. Sp. Pl. 823 = Notabasis syriaca.
tataricus, Lam. Encyc. i. 703 = Cnicus rigens.
tataricus, Linn. Sp. Pl. 825 = Cnicus oleraceus ?
tauricus, Boeber, ex Georgi, Beschr. Russ. Reich. iii. IV. 1224 = Cnicus echinocephalus.
taygeteus, Boiss. & Heldr. Diagn. Ser. II. iii. 42 = nutans.
tectus, Wall. Cat. n. 2906 = Saussurea obvallata.
tenuiflorus, Curt. Fl. Lond. fasc. vi. t. 55 = pycnocephalus.
tenuifolius, Gaud. Fl. Helv. v. 174 = defloratus.
thessalus, Boiss. & Heldr. Diagn. Ser. II. iii. 46 = acanthoides.
Thoermeri, Weinm. in Bull. Soc. Nat. Mosc. (1837) n. VII. 69 = nutans.
Thomsoni, *Hook. f. Fl. Brit. Ind.* iii. 361.—Tibet. occ.
Timbali, Martr. Fl. Tarn, 377 = nigrescens.
tinctorius, Falk, Beitr. ii. 237 = Carthamus tinctorius.
tinctorius, Scop. Fl. Carn. ed. II. ii. 132 = Serratula tinctoria.
tmoleus, *Boiss. Diagn.* Ser. I. iv. 21.—As. Min.
tomentosus, Gilib. Fl. Lituan. i. 188 = Cnicus eriophorus.
transalpinus, Suter, Fl. Helv. 172 = defloratus.
tricephaloides, Lam. Encyc. i. 704 = Cnicus rivularis.
trichocephalus, Wall. Cat. n. 2901 = Tricholepis furcata.
trilobatus, Buch.-Ham. ex DC. Prod. vi. 639 = Cnicus involucratus.

CARDUUS:—
tuberosus, Georgi, It. i. 228 (= *Cirsium setigerum*).
tuberosus, Jacq. Fl. Austr. 280 = Cnicus canus.
tuberosus, Linn. Sp. Pl. 824 = Cnicus pratensis.
tuberosus, Vill. Hist. Pl. Dauph. iii. 15 = Cnicus tuberosus.
uliginosus, Hort. ex Steud. Nom. ed. I. 154 = Cnicus Willdenowii.
uliginosus, Steud. l. c. = Cnicus uliginosus.
uncinatus, *Bieb. Fl. Taur. Cauc.* iii. 553.—Reg. Caucas.; Tauria.
undulatus, Nutt. Gen. Am. ii. 130 = Cnicus undulatus.
uniflorus, Turcz. ex Ledeb. Fl. Ross. ii. 758 = Serratula glauca.
valentinus, *Boiss. & Reut. Diagn.* Scr. II. iii. 45.—Hispan.
Verdii, *Carr. in Rev. Hortic.* (1870–71) 520.—Hispan.
versicolor, Salisb. Prod. 185 = Silybum Marianum.
virginianus, Bosc, ex DC. Prod. vi. 651 = Cnicus Nuttallii.
virginianus, Linn. Sp. Pl. 824 = Cnicus virginianus.
virginianus, Walt. Fl. Carol. 195 = Cnicus repandus.
viridis, *A. Kern. Sched. Fl. Exsicc.* i. 74.—Europ.
vivariensis, *Jord. Obs.* iii. 212.—Gallia.
vulgaris, Savi, Fl. Pis. ii. 241 = Cnicus italicus.
Winklerianus ×, Dalla Torre, in Lotos, xxvii. 1877 (1878) 45.—Europ.
wolgensis, Bieb. ex Willd. Enum. Hort. Berol. 839 = Cousinia wolgensis.
xanthacanthus, *Freyn, in Flora*, lxiv. (1881) 211.—Lusitan.
xeranthemoides, Sweet, Hort. Brit. ed. I. 213 = Serratula xeranthemoides.

CARDWELLIA, F. Muell. Fragm. v. 23 (1865). *PROTEACEAE*, Benth. & Hook. f. iii. 182.
sublimis, *F. Muell. l. c.* 24, 38, 73.—Austral.

CARELIA, Adans. Fam. ii. 123 (1763) = **Ageratum**, Linn. (Compos.).

CARELIA, Less. Syn. Comp. 156 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 241.
cistifolia, *Less. l. c.*—Bras.

CAREX, [Dill.] Linn. Syst. ed. I (1735). *CYPERACEAE*, Benth. & Hook. f. iii. 1073.
CALLISTACHYS et CRYPTOGLOCHIN, Heuff. in Flora, xxvii. (1844) 528.
CARICELLA, Ehrh. Beitr. iv. 146 (1789).
DISTINAX, Rafin. ex Steud. Nom. ed. II. i. 521 (1840).
DORNERA, Heuff. ex Schur, Enum. Pl. Transs. 697 (1866).
DRYMEIA, Ehrh. Beitr. iv. 148 (1789).
GENERSICHIA, Heuff. in Flora, xxvii. (1844) 528.
HELEONASTES, Ehrh. Beitr. iv. 147 (1789).
LEPTOSTACHYS, Ehrh. l. c. (1789).
LEUCOGLOCHIN, Heuff. in Flora, xxvii. (1844) 528.
LIMONAETES, Ehrh. Beitr. iv. 148 (1789).
MAUKSCHIA, Heuff. in Flora, xxvii. (1844) 527.
MONDO, Adans. Fam. ii. 496 (1763).
NEILREICHIA, Kotule, in Spraw. Kom. Fis. Krakow, xvii. (1883) 136.
PHYLLOSTACHYS, Torr. ex Steud. Nom. ed. II. i. 297, in syn. (1840); ii. 328.
PHYSIGLOCHIS, Neck. Elem. iii. 245 (1790).
POLYGLOCHIN, Ehrh. Beitr. iv. 146 (1789).
PSEUDOCAREX, Miq. in Ann. Mus. Lugd. Bat. ii. 146 (1866).
PSYLLOPHORA, Ehrh. Beitr. iv. 146 (1789).
PTACOSEIA, Ehrh. l. c. 147 (1789).
SCHELHAMMERIA, Moench, Meth. Suppl. 119 (1802).
SCURIA, Rafin. in Journ. Phys. lxxxix. (1819) 106.
TRASUS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 53 (1821).
TRIODUS, Rafin. in Journ. Phys. lxxxix. (1819) 106.
TRIPLIMA, Rafin. in Am. Monthly Mag. (1819) 195.
ULVA, Hall. Enum. Stirp. Helv. i. 242 (1742).
UVA, Steud. Nom. ed. II. i. 285 (1840).
VIGNANTHA, Schur, Enum. Pl. Transs. 705 (1866).
VIGNEA, Beauv. in Lestib. Essai Fam. Cyp. 22 (1819).
abbreviata, Presc. ex Boott, in Ann. Nat. Hist. xvii. (1846) 57 = Torreyi.
abjiciens, *Steud. Syn. Pl. Cyp.* 209.—Missouri.
acaulis, *Urv. in Mém. Soc. Linn. Par.* iv. (1826) 599.—Ins. Falkl.

CAREX:—
acicularis, *Boott, in Hook. Fl. N. Zel.* 280. t. 63.—Austral.; N. Zel.
acicularis, *Turcz. ex Steud. Nom.* ed. II. i. 285.—Sibir.; Reg. Caucas.
acroandra, *Schur, Enum. Pl. Transs.* 699.—Transsylv.
acrolepis, Ledeb. in Denkschr. Baier. Bot. Ges. iii. (1841) 56 = Gmelini.
acrolepis, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 271.—Costa-Rica.
acuminata, Reichb. Ic. Fl. Germ. viii. 28. t. 267 = nutans.
acuminata, Willd. Sp. Pl. iv. 300 = glauca.
acuta, All. Fl. Pedem. ii. 247 = riparia.
acuta, Curt. Fl. Lond. t. 281 = paludosa.
acuta, Darling. Florul. Cestric. 98 = virginiana.
acuta, Dewey, ex Hook. Fl. Bor. Am. ii. 218 = angustata.
acuta, *Linn. Sp. Pl.* 978.—Reg. bor. temp.; Austral.
acuta, Muhl. Cat. 84; ed. (1818) 85, nomen = commutata.
acuta, Suter, Fl. Helv. ii. 261 = glauca.
acutaeformis, Brot. Fl. Lusit. i. 66 = echinata.
acutata, *Boott, in Proc. Linn. Soc.* i. (1846) 287.—Am. austr.
acutiformis, Ehrh. Beitr. iv. 43 = paludosa.
acutissima, Déségl. in Loisel. Fl. Gall. ed. I. 628 = pyrenaica.
Aederi, Desv. Fl. Anj. 74, sphalm. = Oederi.
adusta, *Boott, in Hook. Fl. Bor. Am.* ii. 215.—Am. bor.
adusta, Carey, ex Boott, Illustr. Carex, iii. 118 = foenea.
aematorhyncha, Desv. in C. Gay, Fl. Chil. vi. 224 = filiformis.
aemulans, *Liebm. & Drej. in Flora*, xxii. (1839) 142 = stricta.
aequabilis, *Boott, Illustr. Carex*, i. 66. t. 178.—Ins. S. Helena.
aequata, Nees, ex Boott, l. c. ii. 83 = myosurus.
aestivalis, *M. A. Curt. ex A. Gray, in Am. Journ. Sc.* xlii. (1842) 28.—Am. bor.
aethiopica, *Schkuhr, Riedgr.* i. 107. t. Z. f. 83.—Afr. bor.
aethostachya, Schkuhr, l. c. ii. 53 = atrata.
affinis, R. Br. in Richards. in Frankl. Narr. 1st Journ. 750 = obtusata.
agastachys, Linn. f. Suppl. 414 = pendula.
Aitchisoni, Boeck. in Flora, lxiii. (1880) 456 = fissirostris.
alascana, *Boeck. in Engl. Bot. Jahrb.* vii. (1885) 277.—Am. bor. occ.
alata, Torr. in Ann. Lyc. N. York, iii. (1836) 396 = straminea.
alba, Bast. Ess. Fl. Maine-et-Loire, 338 = Bastardiana.
alba, Dewey, in Am. Journ. Sc. vii. (1824) 266 = eburnea.
alba, *Scop. Fl. Carn.* ed. II. ii. 216.—Europ.; Am. bor.
albata, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iii. 193, nomen.—Japon.
albescens, F. Nyland. Spicil. Pl. Fenn. ii. n. 90 = gracilis.
albicans, *Willd. ex Spreng. Syst.* iii. 818.—Am. bor.
albo-atra, Willd. ex Kunth, Enum. Pl. ii. 432, in obs. = fusca.
albo-lutescens, Schwein. in Ann. Lyc. N. York, i. (1828) 66 = straminea.
algida, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 104; xxviii. (1855) 1. 341.—Sibir. baical.
alligata, *Boott, Illustr. Carex*, iv. 129.—Ins. Sandvic.
allomacros, *Steud. Syn. Pl. Cyp.* 189.—Chili.
alopecoidea, *Tuckerm. Enum. Caric.* 18.—Am. bor.
alopecuroides, *D. Don, in Trans. Linn. Soc.* xiv. (1825) 332; *Prod. Fl. Nep.* 43.—Reg. Himal.
Alopecurus, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 141 = Eriophorum latifolium.
alpestris, All. Fl. Pedem. ii. 270 = Halleriana.
alpestris, Dewey, in Am. Journ. Sc. vii. (1824) 268 = varia.
alpestris, Gaud. Etrennes Fl. 173 = sempervirens.
alpestris, Lam. Encyc. iii. 389 = caespitosa.
alpigena, Kern. ex Dalla Torre, Atl. Alpenfl. 224 = ferruginea.
alpina, Honck. Syn. Pl. Germ. i. 374 = vulgaris.
alpina, Hoppe, ex Kunth, Enum. Pl. ii. 466 = ferruginea.
alpina, Schrank, Baier. Fl. i. 298 = sempervirens.
alpina, Sw. ex Wahlenb. Vet. Akad. Nya Handl. Stockh. (1803) 160 = Vahlii.

CAREX:—
alsophila, *F. Muell. Fragm.* viii. 257.—Austral.
alta, *Boott, in Proc. Linn. Soc.* i. (1849) 254; *et in Trans. Linn. Soc.* xx. (1846) 130.—Java.
alveata, *Boott, Illustr. Carex*, i. 21. t. 56.—Texas.
ambigua, *Link, in Schrad. Journ.* ii. (1799) 308.—Lusitan.; Marocc.
ambigua, Moench, Meth. 324 = acuta.
ambleocarpa, Willd. Sp. Pl. iv. 307 = glauca.
amblyolepis, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 99).—Sibir.
ambusta, *Boott, Illustr. Carex*, i. 64. t. 172.—Alaska.
amgunensis, *F. Schmidt, Reisen Amurl.* 69.—Reg. Amur.
amicta, *Boott, Illustr. Carex*, iv. 131.—N. Granat.
ammophila, Willd. Sp. Pl. iv. 226 = divisa.
amoena, *Boott, Illustr. Carex*, iii. 106.—Ind. or.
amphibola, *Steud. Syn. Pl. Cyp.* 234.—Am. bor.
amphilogos, C. Koch, in Linnaea, xxi. (1848) 615 = incurva.
amphora, *Franch. & Sav. Enum. Pl. Jap.* ii. 566.—Japon.
amplifolia, *Boott, in Hook. Fl. Bor. Am.* ii. 228.—Am. bor.
ampullacea, *Good. in Trans. Linn. Soc.* ii. (1794) 207. Reg. bor. temp.
ampullacea, Wulf. ex Wahlenb. in Vet. Akad. Nya Handl. Storkh. (1803) 155 = Michellii.
anceps, *Muhl. ex Schkuhr, Riedgr.* ii. 66. t. F. ff. fig. 128.—Am. bor.
anceps, Willd. ex Kunth, Enum. Pl. ii. 457 = grisea.
Andersoni, Boott, in Hook. f. Fl. Antarct. 364 = decidua.
Andoiniana, *De Not. Sem. Hort. Genov.* (1847); *ex Steud. Syn. Pl. Cyp.* 233.—Europ.
androgyna, Balb. Flencho, 97 = bicolor.
Angarae, *Steud. Syn. Pl. Cyp.* 190.—Sibir.
anguillata, Drej. in Fl. Dan. t. 2846; Revis. Caric. 36 = aquatilis.
angustata, Boott, in Hook. Fl. Bor. Am. ii. 218 = stricta.
angustifolia, Linn. ex Steud. Nom. ed. I. 156 = filiformis.
angustifolia, Seguier, ex Boott, Illustr. Carex, iv. 199 = Schreberi.
angustifolia, Sm. Engl. Fl. iv. 127 = vulgaris.
anisostachys, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 266.—Mexic.
anomala, Boott, ex Perry, Jap. Exp. ii. 327 = gibba.
anomala, *Janka, in Linnaea*, xxx. (1859–60) 609.—Europ. austr.
anomala, Pall. ex Kunth, Enum. Pl. ii. 479 = Halleriana.
anomala, *Steud. Syn. Pl. Cyp.* 230.—Abyssin.
anthericoides, *Presl, Rel. Haenk.* iii. 204.—Am. bor. occ.
anthoxanthera, *Presl, l. c.* 203.—Alaska.
antucensis, Kunze, ex Kunth, Enum. Pl. ii. 412 = vulgaris.
aperta, Boott, in Hook. Fl. Bor. Am. ii. 218 = stricta.
aphanandra, *Franch. & Sav. Enum. Pl. Jap.* ii. 564.—Japon.
aphanolepis, *Franch. & Sav. l. c.* 580.—Japon.
aphylla, *Kunth, Enum. Pl.* ii. 421.—Chili.
aporandra, Kunze, ex Boott, Illustr. Carex, ii. 92 = Shortiana.
appressa, R. Br. Prod. 242 = paniculata.
approximata, All. Fl. Pedem. ii. 267 = cricetorum.
approximata, Hoppe, ex DC. Fl. Fr. Suppl. 290 = lagopina.
approximata, Willd. ex Kunth, Enum. Pl. ii. 411 = caespitosa.
aprica, *Turcz. ex Bess. in Flora*, xvii. (1834) 1. *Beibl.* 27.—Sibir. baical.
aquatilis, Tenore, Fl. Nap. iv. 133 = microcarpa.
aquatilis, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 165.—Reg. bor.
Archeri, Boott, in Hook. f. Fl. Tasm. ii. 98. t. 150 = acicularis.
arcta, Boott, Illustr. Carex, iv. 155 = canescens.
arctata, *Boott, in Hook. Fl. Bor. Am.* ii. 227.—Am. bor.
arctica, Deinb. in Fl. Dan. t. 101; Fries, Summ. Scand. i. 222 = stenophylla.
arctica, Dewey, in Am. Journ. Sc. xxvii. (1835) 239 = Parryana.

CAREX:—
arctophila, Nyland. ex Fries, Summ. Scand. 561 = aquatilis.
Ardoiniana, De Not. in Ann. Sc. Nat. Sér. III. ix. (1848) 324 = olbiensis.
arenaria, Dubois, ex Steud. Nom. ed. II. i. 286 = praecox.
arenaria, Lapeyr. ex Kunth, Enum. Pl. ii. 389 = paniculata.
arenaria, Leers, Fl. Herborn. 195. t. 14. f. 2 = disticha.
arenaria, *Linn. Sp. Pl.* 973.—Reg. bor. temp.
arenaria, Mohr, ex Boott, Illustr. Carex, iv. 210 = incurva.
arenicola, *F. Schmidt, Reisen Amurl.* 195.—Ins. Sachalin.
argentea, J. F. Gmel. Syst. 143 = alba.
argentea, Vill. Fl. Hist. Pl. Dauph. ii. 206 = humilis.
argunensis, *Turcz. ex Ledeb. Fl. Ross.* iv. 267.—Dahuria.
argyrantha, Tuckerm. ex Boott, Illustr. Carex, iii. 119 = adusta.
argyroglochin, Hornem. in Fl. Dan. x. 7. t. 1710 = leporina.
argyrolepis, Maxim. in Franch. & Sav. Enum. Pl. Jap. i. 126 = albata.
arida, Schleich. ex Kunth, Enum. Pl. ii. 464 = sempervirens.
arida, Schwein. & Torr. in Ann. Lyc. N. York, i. (1824) 312. t. 24 = muskingumensis.
aristata, R. Br. in Richards. in Frankl. Narr. 1st Journ. 751 = trichocarpa.
aristata, Clairv. ex Steud. Nom. ed. II. i. 286 = Uncinia microglochin.
aristata, Dewey, in Am. Journ. Sc. vii. (1824) 277. t. A. f. 1 = Davisii.
aristata, Honck. Syn. Pl. Germ. 361 = praecox.
aristata, Siegert, ex Wimm. Fl. Schles. ed. III. (1857) 72 = hirta.
aristata, Urv. in Mém. Soc. Linn. Par. iv. (1826) 599 = trifida.
armena, Boiss. Fl. Orient. v. 426 = fulva.
Arnottiana, *Boott, in Ann. & Mag. Nat. Hist.* xvii. (1846) 56.—Ind. or.; Malaya.
aspera, Willd. Prod. Fl. Berl. 32 = glauca.
asperifolia, *Steud. Syn. Pl. Cyp.* 195.—Ins. Borbon.
asperula, *Nees, in Wight, Contrib.* 124.—Reg. Himal.
asperula, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104; xxviii. (1855) 1. 347 = Meyeriana.
assimilis, *Steud. Syn. Pl. Cyp.* 195.—Ins. Maurit.
assiniboinensis, *Boott, in Coult. Bot. Gaz.* ix. (1884) 91.—Am. bor.
astracanica, Willd. ex Kunth, Enum. Pl. ii. 385 = muricata.
asturica, *Boiss. Pugill. Pl. Nov.* 117.—Hispan.
aterrima, *Hoppe, in Denkschr. Bot. Ges. Regensb.* (1815) t. 3. f. o, p, q.—Alp. Europ.
aterrima, Koch, Syn. Fl. Germ. ed. II. 874 = atrata.
atherodes, Frank, ex Kunth, Enum. Pl. ii. 498 = stenolepis.
atherodes, Spreng. Syst. iii. 828 = aristata.
athrostachya, *Olney, in Proc. Am. Acad.* viii. (1868) 393.—Am. bor. occ.
atrata, Banks, ex Boott, in Proc. Linn. Soc. i. (1849) 257 = Banksii.
atrata, *Linn. Sp. Pl.* 976.—Reg. bor. et arct.
atrofusca, Schkuhr, Riedgr. 106 = ustulata.
atrofusca, Sieber, ex Kunth, Enum. Pl. ii. 435 = fuliginosa.
atrofusca, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 67 = atrata.
atrofusca, Willd. ex Steud. Nom. ed. II. i. 286 = pichinchensis.
atropicta, *Steud. Syn. Pl. Cyp.* 204.—Reg. Magell.
atropurpurea, *Boeck. in Linnaea*, xxxix. (1875) 150.—Am. austr.
atropurpurea, Fisch. ex Trevir. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 538 = atrata.
atrorufa, Willd. ex Kunth, Enum. Pl. ii. 392 = pichinchensis.
attenuata, R. Br. in Richards. in Frankl. Narr. 1st Journ. 750 = rupestris.
anrea, *Nutt. Gen. Am.* ii. 205.—Am. bor.
aureolensis, *Steud. Syn. Pl. Cyp.* 223.—Am. bor.
austriaca, Schkuhr, Riedgr. ii. 10. t. 2 qq. f. 157 = divisa.

CAREX:—
axillaris, *Good. in Trans. Linn. Soc.* ii. (1794) 151. t. 19. f. 1.—Europ.
axillaris, Link, Enum. Hort. Berol. ii. 329 = diluta.
axillaris, Linn. Sp. Pl. ed. II. 1382 = remota.
axillaris, Schrank, ex Steud. Nom. ed. I. 155 = loliacea.
azorica, *J. Gay, in Ann. Sc. Nat.* Sér. II. xi. (1839) 185.—Ins. Azoric.
baazasana, *Steud. Syn. Pl. Cyp.* 236.—Texas.
baccans, *Nees, in Wight, Contrib.* 122.—Ind. or.
Backiana, Dewey, in Am. Journ. Sc. xxix. (1836) 250 = obtusata.
Backii, *Boott, in Hook. Fl. Bor. Am.* ii. 210.—Am. bor.
badia, Pers. Syn. ii. 541 = canescens.
baetica, Auersw. ex Willk. in Bot. Zeit. vi. (1848) 414 = distans.
Balbisii, Schkuhr, ex Spreng. Pugill. ii. 86 = extensa, punctata.
baldensis, *Linn. Cent. Pl.* ii. 32.—Europ.
baldensis, Vill. Hist. Pl. Dauph. ii. 196 = foetida.
Baldwinia, Dewey, in Am. Journ. Sc. xxvi. (1834) 107. t. T. f. 61 = Elliottii.
Baldwiniana, Steud. Nom. ed. II. i. 286 = praec.
Baltzellii, *Chapm. List of Florida Pl.* (1845).—Am. bor.
banata, Sm. in Rees, Cyclop. xxxix. = incurva.
banatica, *Heuff. in Verh. Zool.-Bot. Ges. Wien*, viii. (1858) 222.—Banat.
Banksii, *Boott, in Trans. Linn. Soc.* xx. (1851) 119; *ex Hook. f. Fl. Antarct.* 465. t. 142.—Reg. Magell.
Barbarae, Dewey, ex Torr. in Bot. Mex. Bound. 231 = Prescottiana.
barbata, Boott, Illustr. Carex, i. 68 = Gunniana.
Baroni, *Baker, in Journ. Linn. Soc.* xxi. (1885) 451.—Madag.
Barratti, Torr. ex Schwein. in Ann. Lyc. N. York, i. (1824) 361 = littoralis.
basiantha, *Steud. Syn. Pl. Cyp.* 232.—Am. bor.
basilaris, *Ford. Obs. Pl. Crit.* iii. (1846) 246.—Europ. austr. occ.; Marocco.
Bastardiana, DC. Fl. Fr. vi. 293 = pilulifera.
Beckeri, *C. A. Mey. ex Claus, in Beitr. Pfl. Russ. Reich.* viii. 55.—Rossia.
Beechyana, Boott, Illustr. Carex, i. 67 = Hookeri.
Bellardi, All. Fl. Pedem. ii. 264 = Elyna spicata.
Bella-Villa, *Dewey, in Am. Journ. Sc.* Ser. II. xli. (1866) 229.—Am. bor.
bengalensis, *Roxb. Hort. Beng.* 103; *Fl. Ind.* iii. 572.—Ind. or.
Benthamiana, Boott, in Royle, Illustr. Bot. Himal. (Pref.) ex Boott, Illustr. Carex, 8 = notha.
beringiana, *Cham. ex Steud. Syn. Pl. Cyp.* 229.—Alaska.
bermudiana, *Hemsl. in Journ. Bot.* xxi. (1883) 260.—Ins. Bermuda.
Berteroana, Desv. in C. Gay, Fl. Chil. vi. 198 = aphylla?
Berteroana, Hohen. ex Boott, Illustr. Carex, iv. 149 = setifolia.
Berteroniana, *Steud. in Flora*, xxv. (1842) 604.—Ins. Juan. Fernand.
Bertolonii, Schkuhr, Riedgr. ii. 5. t. D. f. 18 = divisa.
Beyrichiana, *Boeck. in Linnaea*, xli. (1877) 239.—Am. bor.
Bichenoviana, *Boott, in Hook. f. Fl. Tasm.* ii. 101.—Tasmania.
bicolor, *All. Fl. Pedem.* ii. 267.—Europ.; Am. bor.
bicolor, Nyland. ex Nym. Consp. 776 = salina.
bicostata, *Olney, Car. Bor. Am.* 9.—Am. bor.
bifida, *Boott, in Proc. Am. Acad.* vii. (1868) 394.—Am. bor. occ.
bifida, *Roth, ex Steud. Nom.* ed. I. 155.—Hab.?
bifurca, Moench, Meth. 326 = riparia.
bifurca, Schrank, Baier. Fl. i. 304 = ampullacea.
Bigelowii, Torr. & Schwein. ex Boott, Illustr. Carex, iv. 167 = vulgaris.
biligularis, DC. Fl. Fr. v. 296 = laevigata.
bina, Schkuhr, Riedgr. i. 36 = distachya.
binata, *Poir. Encyc. Suppl.* iii. 257.—Afr. bor.
binervis, Dewey, in Am. Journ. Sc. xxx. (1836) 61 = laevigata.
binervis, *Sm. in Trans. Linn. Soc.* v. (1800) 268.—Europ.
binervis, Wahlenb. ex Kunth, Enum. Pl. ii. 451 = fulva.
binervis, Willd. ex Kunth, l. c. 447 = extensa.

CAREX:—
bipartita, All. Fl. Pedem. ii. 265. t. 89. f. 3 = Kobresia caricina.
bipartita, F. G. Dietr. Vollst. Lex. Gaertn. Nachtr. ii. 16 = bifida, *Roth*.
bispicata, Hook. & Arn. in Bot. Beech. Voy. 118. t. 28 = longerostrata.
blanda, Dewey, in Am. Journ. Sc. x. (1826) 45 = laxiflora.
blepharophora, A. Gray, in Ann. Lyc. N. York, iii. (1835) 235 = flexilis.
Blyttii, F. Nyland. Spicil. Pl. Fenn. ii. 35 = tenella.
Boenninghauseniana, Kunth, Enum. Pl. ii. 404 = axillaris.
Boenninghausiana, Weihe, in Flora, ix. (1826) 759 = axillaris.
bohemica, Schreb. Beschr. Graes. ii. 52 = cyperoides.
Boissieri, *Steud. Syn. Pl. Cyp.* 215.—Afr. bor.
Bolanderi, *Olney, in Proc. Am. Acad.* vii. (1868) 393.—Calif.
bolina, Lang, in Linnaea, xxiv. (1851) 551 = vulgaris.
boliviensis, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 32.—Bolivia.
Bolliana, *Boeck. in Flora*, lxi. (1878) 40.—Am. bor.
bonariensis, *Desf. in Poir. Encyc. Suppl.* iii. 250.—Reg. Argent.
bonariensis, Schlecht. in Linnaea, x. (1836) 116 = involucrata.
Bongardi, *Boott, in Proc. Linn. Soc.* i. (1846) 260.—Ins. Loo-choo.
Bongardiana, *C. A. Mey. in Middend. Reise* (*Fl. Ochot.* 101).—Sibir. or.
Bonplandii, *Kunth, Enum. Pl.* ii. 380.—Am. bor. occ. et austr.
Boottiana, Benth. ex Boott, in Bost. Journ. Nat. Hist. v. (1845) 112 = picta.
Boottiana, Hook. & Arn. Bot. Beech. Voy. 273 = Bongardi.
borbonica, *Lam. Encyc.* iii. 387.—Ins. Mascar.
borealis, Lang, in Flora, xxvi. (1843) 142 = aquatilis.
borotalicola, *Regel, in Act. Hort. Petrop.* vii. (1880) 566.—As. centr.
Boryana, *Schkuhr, Riedgr.* ii. 43.—Ins. Mascar.
Boscii, *Willd. ex Spreng. Syst.* iii. 812.—Am. bor.
brachycalama, *Griseb. in Goett. Abh.* xxiv. (1879) 315.—Reg. Argent.
brachycephala, Poepp. ex Boott, Illustr. Carex, i. 67 = indecora.
brachylepis, Turcz. ex Steud. Nom. ed. II. i. 286; et ex Boott, l. c. iii. 113 = alpina.
brachyphylla, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104; xxviii. (1855) 1. 346 = ericetorum.
brachyrhyncha, *Gsaller, in Oestr. Bot. Zeitschr.* xx. (1870) 199, 292.—Europ.
brachystachys, Schrank, Baier. Fl. i. 294 = tenuis.
bracteata, Giesecke, ex Boott, Illustr. Carex, iv. 160 = salina.
bracteata, Suter, Fl. Helv. ii. 250 = mucronata.
bracteosa, *Kunze, in Schkuhr, Riedgr. Suppl.* 12.—Chili.
bracteosa, Schwein. in Ann. Lyc. N. York, i. (1824) 65 = vulpinoidea.
brasiliensis, St. Hil. Voy. Bras. i. 369 = procera.
brazatana, Boeck. in Flora, xxxix. (1856) 321 = baazasana.
Breutelii, Schnizl. ex Lange, Consp. Fl. Groenl. i. 136 = canescens.
brevicaulis, *Thou. ex Steud. Nom.* ed. II. i. 287.—Ins. Trist. d'Acunna.
brevicollis, *DC. Fl. Fr.* vi. 295.—Europ. austr.; As. Min.
breviculmis, *R. Br. Prod.* 242.—Reg. Himal.; Japon.; Australas.
breviculmis, Thw. Enum. Pl. Zeyl. 356 = Thwaitesii.
brevifolia, Host, Gram. iv. 50. t. 89 = ferruginea.
breviglumis, *Le Gall, Fl. Morb.* 658.—Gallia.
brevipes, *S. Wats. in Bot. Calif.* ii. 246.—Calif.
brevirostrata, Poir. Encyc. Suppl. iii. 273 = obesa.
brevirostris, *Blytt, in Bot. Notiser*, (1857) 133.—Europ.
brevirostris, Cederstr. in Vet. Akad. Oefv. Stockh. xiv. (1857) 199 = incurva.
Breweri, *Boott, Illustr. Carex*, iv. 142. t. 455.—Am. bor. occ.
brizoides, Geners. Elench. n. 857 = leporina.
brizoides, Huds. Fl. Angl. 349; ed. II. 406 = canescens.

CAREX:—
brizoides, Lam. Encyc. iii. 382 = praecox.
brizoides, *Linn. Cent. Pl.* i. 31.—Europ.
brizoides, Pall. ex Boott, Illustr. Carex, iv. 199 = Schreberi.
Brizopyrum, Kunze, in Schkuhr, Riedgr. Suppl. i. 169. t. 43 = alta.
bromoides, Dubois, ex Boott, Illustr. Carex, iv. 199 = Schreberi.
bromoides, *Schkuhr, Riedgr.* ii. 8.—Am. bor.
Brongniartii, *Kunth, Enum. Pl.* ii. 380.—Chili.
Brownei, Steud. Syn. Pl. Cyp. 209 = longifolia.
Brownii, Tuckerm. Enum. Car. 21 = lacistoma.
brunnea, *Thunb. Fl. Jap.* 38.—Reg. bor. et austr.
brunnescens, Poir. Encyc. Suppl. iii. 286 = Gebhardi.
Buchanani, *Berggr. in Journ. Bot.* xviii. (1880) 104.—N. Zel.
Buckleyi, Dewey, in Am. Journ. Sc. xlviii. (1845) 143 = canescens.
Buekii, Wimm. Fl. Schles. ed. III. (1857) 81 = stricta.
bufonia, *Gueldenst. ex Ledeb. Fl. Ross.* iv. 320.—Ross. austr.
bulbosa, *Boeck. in Flora*, xxxviii. (1855) 597.—Am. bor.
bullata, Ell. Sketch, ii. 556 = striata.
bullata, Dewey, in Am. Journ. Sc. ix. (1825) 71 = Tuckermani.
bullata, *Rudolph, ex Georgi, Besch. Russ. Reich.* iii. v. 1286.—Tauria.
bullata, *Schkuhr, Riedgr.* ii. 85.—Am. bor.
Bungeana, *Debeaux, in Act. Linn. Soc. Bord.* xxxiii. (1879) 68.—China.
Burchelliana, *Boeck. in Linnaea*, xli. (1877) 234.—Afr. austr.
Buxbaumii, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 163 = fusca.
caduca, *Boott, Illustr. Carex*, iv. 157.—Am. austr.
caesia, Griseb. Spicil. Fl. Rumel. ii. 412 = tomentosa.
caespititia, *Nees, in Wight, Contrib.* 127.—Reg. Himal.
caespitosa, Good. in Trans. Linn. Soc. ii. (1794) 195. t. 21 = vulgaris.
caespitosa, Hook. Fl. Scot. 286 = saxatilis.
caespitosa, Huds. Fl. Angl. ed. II. 412 = stricta.
caespitosa, Ledeb. Fl. Ross. iv. 310 = elcusinoides.
caespitosa, *Linn. Sp. Pl.* 978.—Reg. bor. et austr.
caespitosa, Soland. ex Boott, Illustr. Carex, i. 63 = decidua.
caespitosa, Tenore, Viaggio in Calab. 129 = frigida.
caespitosa, Thunb. Fl. Jap. 39 = Thunbergii ?
Caflischii ×, Brucgg. in Jahresb. Nat. Ges. Graub. II. xxiii–xxiv. (1880) 119.—Europ.
campestris, Hort. ex Boott, Illustr. Carex, i. 161 = obesa.
campestris, Host, Gram. iv. t. 38 = supina.
Camposii, *Boiss. & Reut. Pugill. Pl. Nov.* 117.—Hispan.
Campyloxys, *Steud. Syn. Pl. Cyp.* 213.—Reg. Magell.
camtschatcensis, Kunth, Enum. Pl. ii. 477 = longerostrata.
canadensis, *Dewey, in Am. Journ. Sc.* Ser. II. xli. (1866) 229.—Am. bor.
canescens, Host, Gram. i. t. 57 = paradoxa.
canescens, Huds. Fl. Angl. ed. II. 405 = divulsa.
canescens, Leers, Fl. Herborn. 712. t. 14. f. 3 = muricata.
canescens, *Linn. Sp. Pl.* 974.—Reg. temp. bor. et austr.
canescens, Poll. Palat. Hist. Pl. ii. 571 = elongata.
capensis, *Schkuhr, Riedgr.* ii. 39.—Afr. austr.
capensis, Thunb. Prod. Pl. Cap. 14 = Schoenoxyphium Thunbergii.
capillacea, *Boott, Illustr. Carex*, i. 44. t. 110.—Reg. Himal.; Austral.
capillaris, Leers, Fl. Herborn. t. 15. f. 2 = sylvatica.
capillaris, *Linn. Sp. Pl.* 977.—Reg. bor. temp. et arct.
capillaris, Willd. ex Kunth, Enum. Pl. ii. 464 = firma.
capillata, Schkuhr, ex Kunth, l. c. 430 = hirsuta.
capillipes, *Drej. in Fl. Dan.* t. 2844; *Revis. Caric.* 50.—Ins. Island.
capitata, All. Fl. Pedem. ii. 264 = Davalliana.
capitata, *Linn. Syst.* ed. X. 1261.—Europ.; Am. bor.
capitata, F. Muell. Fragm. viii. 251 = Cephalotes.
capitata, Oed. in Fl. Dan. t. 372 = Eleocharidis sp.
capitata, Suter, ex Boott, Illustr. Carex, iv. 143 = dioica.
capitellata, *Boiss. & Bal. ex Boiss. Fl. Orient.* v. 399.—As. Min.

CAREX:—
cardiolepis, *Nees, in Wight, Contrib.* 127.—Reg. Himal.
Careyana, *Torr. ex Dewey, in Am. Journ. Sc.* Ser. I. xxx. (1836) 60.—Am. bor.
Carlantoniana, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 293 = enervis.
carltoniana, *Dewey, in Am. Journ. Sc.* Ser. I. xxviii. (1835) 238.—Am. bor.
caroliniana, *Buckley, l. c.* xlv. (1843) 173.—Am. bor.
caryophyllaea, J. F. Gmel. Syst. 142 = seq.
caryophyllata, Kunth, Enum. Pl. ii. 437 = emarginata.
caryophyllea, Latour. Chlor. 27 = praecox.
Castanea, Ell. Sketch, ii. 546 = Elliottii.
Castanea, Mielich. in Flora, xxxii. (1849) 665 = capillaris.
Casteriana, Heer, ex Boott, Illustr. Carex, iv. 143 = dioica.
cataractae, R. Br. Prod. 242 = flava.
caucasica, *Stev. in Mém. Soc. Nat. Mosc.* iv. (1813) 68.—Reg. Caucas.
celsa, *Boott, Illustr. Carex*, iii. 108.—Reg. Himal.
cenisia, Balb. in Roem. Arch. iii. 137 = bicolor.
cephaloidea, *Dewey, ex Boott, Illustr. Carex*, iii. 123.—Am. bor.
cephalophora, *Muhl. ex Schkuhr, Riedgr.* ii. 4.—Am. bor.
Cephalotes, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 110.—Austral.
cerina, Dum. Fl. Belg. 147; Boott, Illustr. Carex, iv. 170 = vulgaris.
cernua, *Boott, l. c.* 171.—Reg. Himal.; Japon.
cernua, *Phil. in Linnaea*, xxix. (1857–8) 83.—Chili.
cernua, *Raeusch. Nom.* ed. III. 270.—'Helvet.'
ceylanica, *Boeck. in Linnaea*, xl. (1876) 341.—Ind. or.
Chaberti, F. Schult. in Flora, liv. (1871) 21 = divulsa.
chaetophylla, Steud. Syn. Pl. Cyp. 187 = divisa.
chaetorhiza, *Franch. & Sav. Enum. Pl. Jap.* ii. 552.—Japon.
chalaros, *Steud. Syn. Pl. Cyp.* 231.—Am. bor.
Chamissoi, *Boeck. in Linnaea*, xli. (1877) 145.—Hab.?
Chapmani, Torr. ex Boott, Illustr. Carex, i. 16 = Baltzellii.
Chapmanni, Sartwell, ex Dewey, in Am. Journ. Sc. Ser. II. xix. (1855) 254 = tenax.
Chapmanni, *Steud. Syn. Pl. Cyp.* 222.—Florida.
Cheesemanii, *Petrie, in Trans. N. Z. Inst.* xv. (1883) 358.—N. Zel.
cherokeensis, *Schwein. & Torr. in Ann. Lyc. N. York*, i. (1824) 369. t. 25. f. 1.—Am. bor.
chilensis, Brongn. in Duperr. Voy. Bot. 156 = riparia.
chillanensis, *Phil. in Anal. Univ. Chil.* II. (1862) 384.—Chili.
chinensis, *Retz. Obs.* iii. 42.—China.
chlorantha, *R. Br. Prod.* 242.—Austral.
chlorocarpa, Wimm. in Flora, xxxiii. (1849) 620 = vulgaris.
chlorocarpos, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 265 = Liebmanni.
chlorocystis, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 520.—Hongkong.
chlorolepis, *Steud. Syn. Pl. Cyp.* 204.—Chili.
chlorostachys, D. Don, in Till. Mag. (Dec. 1823) 455 = japonica.
chlorostachys, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 68 = sempervirens.
chordalis, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 269.—Mexic.
chordorrhiza, *Linn. f. Suppl.* 414.—Europ.; Am. bor.
Christyana, Boott, in Bost. Journ. Nat. Hist. v. (1847) 114 = cherokeensis.
chrysolepis, *Franch. & Sav. Enum. Pl. Jap.* ii. 571.—Japon.
ciliata, Willd. in Mém. Acad. Berol. 1794 (1797) 47. t. 3. f. 2 = cricetorum.
cilicica, *Boiss. Diagn.* Ser. II. iv. 124.—Cilicia.
cinerea, Poll. Hist. Pl. Palat. ii. 571 = canescens.
cinnamomea, Boott, in Proc. Linn. Soc. i. (1846) 257; Trans. Linn. Soc. xx. (1851) 136 = nivalis.
cinnamomea, *Cheesm. in Trans. N. Z. Inst.* xiv. (1882) 301.—N. Zel.
cinnamomea, *Olney, in Proc. Am. Acad.* vii. (1868) 396.—Calif.
circinnata, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 209. t. 6.—Unalaschka.
cirrhosa, *Bergg. in Minneskr. Fisiog. Sallks. Lund* (1877) n. VIII. 29.—N. Zel.

CAREX:—
cirrhulosa, *Nees, in Hook. Kew Journ.* vi. (1854) 29.—Ins. Philipp.
cladostachya, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 149 = polystachya.
clandestina, Good. in Trans. Linn. Soc. ii. (1794) 167 = humilis.
clavaeformis, *Hoppe, in Sturm, Deutschl. Fl. Heft* 61 (1833).—Carinthia.
clavata, *Thunb. Prod. Pl. Cap.* 14.—Afr. austr.
coacta, Boott, in Proc. Linn. Soc. i. (1846) 285; et Trans. Linn. Soc. xx. (1846) 133 = curaica.
coarctata, *Kit. in Linnaea*, xxxii. (1863) 319.—Hungar.
cognata, *Kunth, Enum. Pl.* ii. 502.—Afr. austr.
colchica, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 303 = arenaria.
Colensoi, *Boott, in Hook. f. Fl. N. Zeal.* 281. t. 63.—N. Zel.
collata, Boott, in Hook. Lond. Journ. Bot. iii. (1844) 417 = virgata.
collecta, *Dewey, in Am. Journ. Sc.* xi. (1826) 314. tab. N. fig. 44.—Am. bor.
collina, Willd. Sp. Pl. iv. 260 = montana.
Collinsii, Nutt. Gen. Am. ii. 205 = subulata.
columbiana, Dewey, ex Torr. in Ann. Lyc. N. York, iii. (1836) 406 = Mertensii.
comans, *Bergg. in Minneskr. Fisiog. Sallk. Lund* (1877) n. VIII. p. 28.—N. Zel.
Commersoniana, Sieber, ex Kunth, Enum. Pl. ii. 391 = brunnea.
commixta, *Steud. Syn. Pl. Cyp.* 207.—Java.
commutata, *J. Gay, in Ann. Sc. Nat.* Sér. II. xi. (1839) 198.—Am. bor.
commutata, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) 1. 205.—Soongaria.
comosa, Boott, in Trans. Linn. Soc. xx. (1846) 117 = C. Pseudo-cyperus.
compacta, *R. Br. in App. Ross, Voy.* 163.—Am. arct.
compacta, Krock. ex Hoppe, Caric. 37 = vulgaris.
compacta, Lam. Fl. Fr. ii. 172 = vulpina.
compacta, Poir. Encyc. Suppl. iii. 282 = Uncinia compacta.
compacta, *Wimm. in Flora*, xxxiii. (1850) 618.—Europ.
complanata, Torr. in Ann. Lyc. N. York, iii. (1836) 408 = triceps.
composita, *Boott, Illustr. Carex*, i. 3.—Reg. Himal.
compressa, Gaud. Etiennes Fl. 111 = stricta.
compressa, *Hose, in Usteri, Ann. Bot.* xxi. (1797) 33.—Europ.
compressa, Kit. ex Willd. Sp. Pl. iv. 276 = brachystachys.
compressa, Vahl, ex Kunth, Enum. Pl. ii. 378 = bonariensis.
concinna, *R. Br. in Richards. Frankl. Narr. 1st Journ.* 751—Am. arct.
concolor, *R. Br. in Parry, Voy. App.* 283.—Am. arct.
concolor, Nees, in Wight, Contrib. 125 = speciosa.
condensata, *Nees, in Wight, Contrib.* 123.—Reg. Himal.
conferta, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 512.—Abyssin.
confertiflora, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 418.—Japon.
congesta, C. A. Mey. ex Boott, Illustr. Carex, i. 17 = Hoodii.
conglobata, All. Fl. Pedem. ii. 268 = montana.
conglobata, Kit. in Willd. Sp. Pl. iv. 281 = obesa.
conica, *Boott, ex Perry, Jap. Exped.* ii. 325.—Japon.
conjugata, Willd. ex Kunth, Enum. Pl. ii. 391 = Commersoniana.
conjuncta, *Boott, Illustr. Carex*, iii. 122.—Am. bor.
conoidea, Muhl. Desc. Gram. Uber. 248 = laxiflora.
conoidea, *Schkuhr, Riedgr.* ii. 67.—Am. bor.
consanguinea, Kunth, Enum. Pl. ii. 374 = divisa.
consocialis, *Steud. Syn. Pl. Cyp.* 222.—Japon.
contigua, Hoppe, in Sturm, Deutschl. Fl. Heft 61 (1833) = muricata.
contracta, *F. Muell. Fragm.* viii. 258.—Austral.
convexa, *Kit. in Linnaea*, xxxii. (1863) 317.—Hungaria.
Cooleyi, Dewey, in Ann. Journ. Sc. Ser. I. xlviii. (1845) 144 = hystericina.
coriophora, Fisch. & Mey. ex Kunth, Enum. Pl. ii. 463 = ustulata.
corrabensis, *Phil. in Linnaea*, xxxiii. (1864–65) 273.—Chili.

CAREX:—
corsica, Degl. ex Loisel. Fl. Gall. ii. 307 = microcarpa.
corsicana, Link, Enum. Hort. Berol. i. 358 = punctata.
Cortesii, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 268.—Mexic.
corynophora, Peterm. in Flora, xxvii. (1844) 335 = acuta.
costata, *J. & C. Presl, Fl. Čech.* 191.—Bohemia.
costata, Schwein. in Ann. Lyc. N. York, i. (1824) 67 = virescens.
costata, Turcz. ex Ledeb. Fl. Ross. iv. 305 = obesa.
Coulteri, *Boott, ex Hemsl. Biol. Centr. Am. Bot.* iii. 473.—Mexic.
courtallensis, *Nees, ex Boott, Illustr. Carex*, i. 52.—Ind. or.
crassa, Ehrh. Beitr. iv. 43 = riparia.
crassipes, *Boeck. in Linnaea*, xl. (1876) 329.—Ind. or.
Crawei, *Dewey, ex Torr. Fl. N. York*, ii. 408.—Am. bor.
crinalis, *Boott, in Proc. Linn. Soc.* i. (1846) 256.—N. Granat.
crinigera, *Boott, Illustr. Carex*, ii. 102.—Ins. Borbon.
crinita, *Lam. Encyc.* iii. 393.—Am. bor.
cristata, [Clairv.] Man. Herb. 291 = ferruginea.
cristata, Schwein. in Ann. Lyc. N. York, i. (1824) 66 = tribuloides.
cruciata, *Nees, ex Wight, Contrib.* 123.—Ind. or.
cruciata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 149 = ramosa.
cruenta, *Nees, in Wight, Contrib.* 128.—Reg. Himal.
Crus-corvi, *Shuttl. ex Kunze, in Schkuhr, Riedgr. Suppl.* 128. t. 32.—Am. bor.
cryptandra, Franch. & Sav. Enum. Pl. Jap. ii. 148 = macroglossa.
cryptocarpa, *Cheesem. in Trans. N. Z. Inst.* xvi. (1884) 412.—N. Zel.
cryptocarpa, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 226 = halophila.
cryptostachys, *Brongn. in Duperr. Voy. Bot.* 152.—N. Guinea.
Csetzii, Nym. Consp. 773 = Czetzii.
Cumingiana, *Steud. Syn. Pl. Cyp.* 206.—Ins. Philipp.
Cunninghamii, Boott, Illustr. Carex, iv. 171. t. 579 = acuta.
curaica, Franch. & Sav. Enum. Pl. Jap. ii. 252 = chaetorhiza.
curaica, Kunth, Enum. Pl. ii. 375 = vulpinaris.
curta, Good. in Trans. Linn. Soc. ii. (1794) 145 = canescens.
curvata, *Boott, Illustr. Carex*, i. 2.—Reg. Himal.
curvata, Knaf, in Flora, xxx. (1847) 184 = Schreberi.
curvicollis, *Franch. & Sav. Enum. Pl. Jap.* ii. 579.—Japon.
curvirostra, Hartm. Skand. Fl. ed. II. 221 = vaginata.
curvirostris, Kunze, in Schkuhr, Riedgr. Suppl. 79. t. 20 = baccans.
curvula, *All. Fl. Pedem.* ii. 264. t. 92. f. 3.—Europ.
curvula, Lam. Encyc. iii. 380 = Schreberi.
curvula, Stev. in Mém. Soc. Nat. Mosc. vi. (1823) 67 = Elyna schoenoides.
curvula, Willd. ex Kunth, Enum. Pl. ii. 269 = Davalliana.
cuspidata, Bertol. in Mem. Soc. Med. Emul. Gen. ii. (1803) 145 = divisa.
cuspidata, Degl. in Loisel. Fl Gall. 294 = echinata.
cuspidata, Host, Gram. i. 71. t. 97 = glauca.
cuspidata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 164 = salina.
Custoriana, Heer, ex Hegetschw. Fl. Schw. 906 = dioica.
cylindrica, Mielich. ex Kunth, Enum. Pl. ii. 434 = atrata.
cylindrica, Schwein. in Ann. Lyc. N. York, i. (1824) 71 = bullata.
cylindrica, Tuckerm. ex Boott, in Hook. Lond. Journ. Bot. v. (1846) 74 = Tuckermani.
cyperoides, Dewey, in Am. Journ. Sc. Ser. II. iii. (1847) 171 = sychnocephala.
cyperoides, *Murr. in Linn. Syst.* ed. XIII. 703.—Europ.; As. temp.
Czetzii, *Janka, in Oestr. Bot. Zeitschr.* viii. (1858) 93.—Europ.
dacica, Heuff. in Flora, xviii. (1835) 247 = caespitosa.
Daltonii, *Boott, Illustr. Carex*, i. 5.—Ind. or.

CAREX:—
Darlingtonii, Schwein. ex M. A. Curt. in Am. Journ. Sc. Ser. II. vii. (1849) 410 = aestivalis.
Darwinii, *Boott, in Proc. Linn. Soc.* i. (1845) 261.—Reg. Magell.
dasycarpa, *Muhl. Desc. Gram. Uber.* 236.—Am. bor.
Davalliana, Dewey, ex Steud. Nom. ed. II. i. 288 = pyrenaica.
Davalliana, *Sm. in Trans. Linn. Soc.* v. (1800) 266.—Reg. bor. temp.
Davalliana, Willd. ex Kunth, Enum. Pl. ii. 368 = dioica.
Davidi, *Franch. Pl. Davidian.* i. (1884) 319.—China.
Davisii, Dewey, in Am. Journ. Sc. x. (1826) 279 = varia.
Davisii, *Schwein. & Torr. in Ann. Lyc. N. York*, i. (1824) 326.—Am. bor.
debilis, Boiss. & Bal. ex Boiss. Fl. Orient. v. 410 = transylvanica.
debilis, Forst. f. Prod. 92 = Forsteri.
debilis, *Michx. Fl. Bor. Am.* ii. 172.—Am. bor.
debilis, Soland. ex Boott, Illustr. Carex, i. 65 = Solandri.
decidua, *Boott, in Proc. Linn. Soc.* i. (1845) 255.—Calif.; Reg. Magell.
decipiens, J. Gay, in Ann. Sc. Nat. Sér. I. xxvi. (1832) 209 = macrostyla.
decipiens, Turcz. in Bull. Soc. Nat. Mosc. (1838) 102; xxviii. (1855) i. 323 = obtusata.
declinata, *Boott, Illustr. Carex*, iv. 171. t. 580.—Austral.
decolorans, Wimm. in Flora, xxxiii. (1850) 622 = vulgaris.
decomposita, *Muhl. Descr. Gram. Uber.* 264.—Am. bor.
decora, *Boott, Illustr. Carex*, i. 5. t. 15.—Reg. Himal.
decumbens, Ehrh. Beitr. vi. 82 = pilulifera.
deflexa, *Hornem. in Fl. Dan.* t. 3051.—Am. bor.; Groenl.
Deinbolliana, J. Gay, in Ann. Sc. Nat. Sér. II. xi. (1839) 183 = stenophylla.
Dematranea, *Lagger, in Flora*, xxxviii. (1855) 207.—Europ.
demissa, *Hornem. in Fl. Dan.* t. 1342.—Europ.
densiflora, *Presl, Rel. Haenk.* iii. 204.—Ins. Marian.
denudata, Lapeyr. ex Steud. Nom. ed. I. 160 = pyrenaica.
depauperata, *Good. in Trans. Linn. Soc.* ii. (1794) 181.—Europ.; As. bor.
depauperata, Hornem. Plantelaere, ii. 829 = vaginata.
depressa, Link, in Schrad. Journ. ii. (1799) 309 = Halleriana.
depressa, Willd. ex Kunth, Enum. Pl. ii. 479 = alpestris.
desponsa, *Boott, Illustr. Carex*, ii. 82.—Reg. Himal.
Despreauxii, *Steud. Syn. Pl. Cyp.* 237.—'Terra nova.'
devia, *Cheesem. in Trans. N. Z. Inst.* xv. (1883) 301.—N. Zel.
Deweyana, *Schwein. in Ann. Lyc. N. York*, i. (1824) 65, 310.—Am. bor.
Dianae, *Steud. Syn. Pl. Cyp.* 238.—Ins. S. Helena.
diandra, Roth, Tent. Fl. Germ. ii. ii. 437 = paniculata.
diandra, Schrank, in Act. Acad. Mogunt. (1782) 49 (err. typ. 57) = teretiuscula.
diaphana, *Boott, Illustr. Carex*, iv. 156.—Austral.
Dickinsii, *Franch. & Sav. Enum. Pl. Jap.* ii. 581.—Japon.
diclina, *Phil. in Linnaea*, xxxiii. (1864–65) 271.—Chili.
Dietrichiae, Boeck. in Flora, lviii. (1875) 122 = fissilis?
digitalis, Schwein. & Torr. in Ann. Lyc. N. York, i. (1824) 324. t. 27 = gracillima.
digitalis, *Willd. Sp. Pl.* iv. 298.—Am. bor.
digitata, *Linn. Sp. Pl.* 975.—Europ.; As. temp.
dilatata, Hohen. ex Boiss. Fl. Orient. v. 425, sphalm. = seq.
diluta, *Bieb. Fl. Taur. Cauc.* ii. 388.—Europ. austr.; As. temp.
dimorpha, Brot. Fl. Lusit. i. 64 = Linkii.
dimorpholepis, Steud. Syn. Pl. Cyp. 214 = pruinosa.
dimorphotheca, *Stschégl. in Bull. Soc. Nat. Mosc.* (1854) 206.—Sibir. altaic.
dioica, Carey, ex Boott, Illustr. Carex, iv. 143 = chordorrhiza.

CAREX:—
dioica, Host, ex Steud. Nom. ed. II. i. 288 = Davalliana.
dioica, Lam. ex Steud. l. c. = Elyna spicata.
dioica, *Linn. Sp. Pl.* 972.—Reg. bor.
dioicotrigona, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = Davalliana.
dioiscostrongyla, St. Lag. l. c. = dioica.
dipsacea, *Bergg. in Minneskr. Fisiog. Sallks. Lund* (1877) n. VIII. p. 28.—N. Zel.
discoidea, Boott, in Mem. Am. Acad. N. S. vi. (1858–59) 419 = breviculmis.
discolor, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 519.—Japon.
discolor, F. Nyland. Spicil. Pl. Fenn. iii. n. 12 = salina.
discolor, Reinw. ex De Vriese, Pl. Ind. Bat. Reinw. 143, partim = virgata.
dispalatha, *Boott, in Perry, Jap. Exp.* ii. 325.—Japon.
disperma, Dewey, in Am. Journ. Sc. viii. (1820) 266. t. A. f. 3 = tenella.
disperma, Herb. Un. It. ex Kunth, Enum. Pl. ii. 480 = Steudelii.
dissimilis, *J. F. Gmel. Syst.* 144.—Helvet.
dissita, *Soland. ex Hook. f. Fl. N. Zel.* 284.—N. Zel.
distachya, Desf. Fl. Atlant. ii. 336 = Linkii.
distachya, Willd. Sp. Pl. iv. 220 = atrata.
distans, Bert. ex Steud. Nom. ed. II. i. 288 = indecora.
distans, Host, Gram. i. 57. t. 77 = fulva.
distans, Lightf. Fl. Scot. ii. 561 = binervis.
distans, *Linn. Syst.* ed. X. 1263.—Europ.; As. occ.; Am. bor.
distans, Willd. ex Kunth, Enum. Pl. ii. 488 = nutans.
distenta, Kunze, ex Kunth, l. c. 449 = indecora.
disticha, A. Gray, Man. N. U. St. ed. V. 574 = Sartwellii.
disticha, *Huds. Fl. Angl.* ed. I. 347.—Europ.
divergens, Thuill. Fl. Par. ii. 481 = elongata.
diversicolor, *Crantz, Inst.* i. 405.—Italia?
diversiflora, Host, Gram. i. 53. t. 70 = Halleriana.
divisa, *Huds. Fl. Angl.* ed. I. 348; ed. II. 405.—Reg. bor. temp.
divisa, Oed. in Fl. Dan. t. 371 = flava.
divulsa, Gaud. Fl. Helv. vi. 47 = muricata.
divulsa, *Good. in Trans. Linn. Soc.* ii. (1794) 160.—Reg. bor. temp.
Doenitzii, *Boeck. in Flora*, lxv. (1882) 61.—Japon.
Doniana, Spreng. Syst. iii. 825 = japonica.
Douglasii, *Boott, in Hook. Fl. Bor. Am.* ii. 213.—Am. bor. occ.
Dregeana, *Kunth, Enum. Pl.* ii. 511.—Afr. austr.
Drejeri, Lange, in Flora, xxv. (1842) 548 = caespitosa.
Drejeriana, *Lange, in Fl. Dan.* t. 2975.—Groenl.
Drummondiana, Dewey, in Am. Journ. Sc. xxix. (1836) 251. t. Y. f. 82 = rupestris.
Drymeia, Linn. f. Suppl. 414 = sylvatica.
drymophila, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 104.—Sibir.
dubia, Gaud. Etrennes Fl. 178 = atrata.
dubia, *Hose, in Usteri, Ann. Bot.* xxi. (1797) 34.—Europ.
dubia, Sibth. in Sibth. & Sm. Fl. Graec. Prod. i. 53 = Sesleria alba.
dubitata, Dewey, ex Boott, Illustr. Carex, iv. 167 = vulgaris.
Dufourei, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 140 = rupestris.
dura, Boott, in Proc. Linn. Soc. i. (1845) 255 = pichinchensis.
Durieui, *Steud. Nom.* ed. II. i. 289.—Hispan.
duriuscula, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 214. t. 8 = stenophylla.
Duvaliana, *Franch. & Sav. Enum. Pl. Jap.* ii. 568.—Japon.
ebracteata, *Phil. in Linnaea*, xxxiii. (1864-65) 271.—Chili.
ebracteata, *Trautv. in Act. Hort. Petrop.* v. (1877) 125.—Sibir.
eburnea, *Boott, in Hook. Fl. Bor. Am.* ii. 226.—Am. bor.
echinata, Lam. Fl. Fr. ii. 177 = flava.
echinata, *Murr. Prod. Stirp. Gotting.* 76.—Reg. bor. temp.; Austral.
echinata, Roth, Tent. Fl. Germ. i. 395 = stellulata.
echinochloe, *Kunze, in Schkuhr, Riedgr. Suppl.* 47. t. 12.—Abyssin.
Ecklonei, *Nees, in Linnaea*, x. (1836) 203.—Afr. austr.

CAREX:—
efflata, Lacstad. ex Ledeb. Fl. Ross. iv. 291 = panicea.
eggytera, Steud. Syn. Pl. Cyp. 220 = breviculmis.
Ehrenbergiana, *Boeck. in Linnaea*, xli. (1877) 175.—Mexic.
Ehrhartiana, Hoppe, ex Boott, Illustr. Carex, iv. 145 = terctiuscula.
elata, All. Fl. Pedem. ii. 272 = stricta.
elata, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1831) 11.—Ins. Madera.
elata, Turner, ex Boott, Illustr. Carex, i. 66 = praelata.
elatior, *Boeck. in Bremen Abh.* vii. (1880) 41.—Madag.
elegans, Willd. Prod. Fl. Berl. 34 = limosa.
eleopsammodes, Steud. Syn. Pl. Cyp. 213 = Darwinii.
eleusinoides, *Turcz. ex Kunth, Enum. Pl.* ii. 407.—Sibir. baical.
Elliottii, *Torr. in Ann. Lyc. N. York*, i. (1824) 357.—Am. bor.
elongata, Leers, Fl. Herborn. 197. t. 14. f. 7 = canescens.
elongata, *Linn. Sp. Pl.* 974.—Europ.; Sibir.
elytroides, Frics, in Bot. Notiser, (1846) 106 = saxatilis.
emarcida, Suter, Fl. Helvet. ii. 263 = sylvatica.
emarginata, Willd. Sp. Pl. iv. 262 = montana.
eminens, *Nees, in Wight, Contrib.* 122.—Reg. Himal.
emirnensis, *Baker, in Journ. Bot.* xxi. (1883) 129.—Madag.
Emmonsii, Dewey, in Ann. Lyc. N. York, iii. (1836) 411 = varia.
emodorum, Sprcng. Syst. iii. 818 = alopecuroides.
Emoryi, *Dewey, ex Torr. Bot. Mex. Bound.* 230.—Am. bor. occ.
enervis, Dulac, Fl. Hautes-Pyr. 53 = brizoides.
enervis, *C. A. Mey. in Ledeb. Fl. Alt.* iv. 209.—Sibir. altaic.
ensifolia, Turcz. ex Ledeb. Fl. Ross. iv. 309 = saxatilis.
epigejos, Frics, Summa, 223 = salina.
erecta, DC. Fl. Fr. iii. 120 = sempervirens.
erecta, Gilib. Exercit. ii. 549 = caespitosa.
ericetorum, *Poll. Hist. Pl. Palat.* ii. 580.—Europ.; As. temp.
erinacea, Cav. Ic. v. 40 = Uncinia erinacea.
erinacea, Mubl. ex Steud. Nom. ed. II. i. 288 = hystericina.
eriophora, *Fisch. ex Steud. l. c.* 289.—Dahuria.
erythraea, *Schrad. in Linnaea*, xi. (1837) 87.—Reg. Himal.
erythrorrhiza, *Boeck. in Linnaea*, xxxix. (1875) 103.—Abyss.
erythrorrhiza, Steud. Syn. Pl. Cyp. 194 = Koestlini.
erythrostachys, Hoppe, in Sturm, Deutschl. Fl. Heft 69 (1835) = glauca.
erythrostachys, Kral. ex Nym. Consp. 774 = serrulata.
Esenbeckiana, *Boeck. in Linnaea*, xl. (1876) 372.—Afr. austr.
Esenbeckii, *Kunth, Enum. Pl.* ii. 522.—Reg. Himal.
Essenbeckii, Boott, Illustr. Carex, i. 52 = linearis.
evoluta, *Hartm. in Vet. Akad. Handl. Stockh.* (1818) 156.—Europ.
exaltata, *Kit. in Linnaea*, xxxii. (1863) 322.—Hungaria.
exaltata, Peterm. in Flora, xxvii. (1844) 340 = riparia.
excelsa, Poepp. ex Kunth, Enum. Pl. ii. 502 = C. Pseudocyperus.
excisa, *Boott, ex Perry, Jap. Exp.* ii. 326.—Japon.
excurrens, Cham. ex Steud. Syn. Pl. Cyp. 228 = podocarpa.
excurva, *Boott, Illustr. Carex*, i. 57.—Reg. Himal.
exigua, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 514.—Zeylan.
exilis, *Dewey, in Am. Journ. Sc.* xiv. (1828) 351.—Am. bor.
extensa, *Good. in Trans. Linn. Soc.* ii. (1794) 175. t. 21. f. 7.—Reg. temp. bor.; Am. austr.
extensa, C. A. Mey. Verz. Pfl. Cauc. 31 = diluta.
Fabri, *Hance, in Journ. Linn. Soc.* xiii. (1873) 90.—China.
falcata, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104; xxviii. (1855) t. 338 = saxatilis.
fallax, *Steud. Syn. Pl. Cyp.* 189.—Java.
familiaris, *Steud. l. c.* 226.—Texas.
fascicularis, Sievers, ex Trevir. in Bull. Soc. Nat. Mosc. xxxvi. (1863) t. 543 = caespitosa.

CAREX:—
fascicularis, Soland. ex Hook. f. Fl. N. Zel. i. 283 = C. Pseudocyperus.
fasciculata, Link, in Schkuhr, Riedgr. i. 119 = echinata.
fasciculata, *Willd. Sp. Pl.* iv. 248.—Lusitan.
Fauriae, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 89.—Japon.
fecunda, *Steud. Syn. Pl. Cyp.* 194.—Am. trop.
Fedia, Nees, ex Wight, Contrib. 129 = Wallichiana.
feminea, Steud. ex Boott, Illustr. Carex, iv. 138 = acutata.
Fendleriana, *Boeck. in Linnaea*, xxxix. (1875) 135.—Am. bor.
ferax, Racusch. Nom. ed. III. 270 = crecta.
ferruginea, Schkuhr, Riedgr. i. 92 = sempervirens.
ferruginea, *Scop. Fl. Carn.* ed. II. ii. 225.—Europ.
ferruginea, Stev. in Mém. Soc. Mat. Mosc. iv. (1813) 68 = sempervirens.
festiva, *Dewey, in Am. Journ. Sc.* xxix. (1835) 246.—Reg. bor.; Ins. Sandvic.
festucacea, Schkuhr, Riedgr. ii. 23 = straminea.
fibrillosa, *Franch. & Sav. Enum. Pl. Jap.* ii. 564.—Japon.
filicina, *Nees, in Wight, Contrib.* 123.—Ind. or.; Malaya.
filiculmis, *Franch. & Sav. Enum. Pl. Jap.* ii. 563.—Japon.
filifolia, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 125 = Duricui.
filifolia, Nutt. Gen. Am. ii. 204 = Uncinia breviseta.
filiformis, Leers, Fl. Herborn. t. 16. f. 5 = praecox.
filiformis, *Linn. Sp. Pl.* 976.—Europ.; Am. bor. et austr.; As. bor.
filiformis, Thuill. Fl. Par. 440 = tomentosa.
filiformis, Vahl, in Fl. Dan. t. 1048 = pilulifera.
filipendula, Drej. l. c. t. 2370 = halophila.
filipes, *Franch. & Sav. Enum. Pl. Jap.* ii. 576.—Japon.
fimbriata, *Schkuhr, Riedgr.* ii. 61.—Europ.
finitima, *Boott, Illustr. Carex*, i. 44.—Reg. Himal.
firma, *Host, Syn.* 509.—Europ.
Fischeriana, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 286 = chordorrhiza.
fissilis, *Boott, Illustr. Carex*, ii. 86. t. 245.—Austral.; N. Caled.
fissirostris, *Ball, in Journ. Bot.* xiii. (1875) 206.—Marocco; Afghanist.
flacca, Carey, ex Boott, Illustr. Carex, i. 69 = Barrattii.
flacca, Schreb. Spicil. 669 = glauca.
flaccida, Sw. ex Kunth, Enum. Pl. ii. 150 = cladostachya.
flaccidula, *Steud. Syn. Pl. Cyp.* 199.—Am. bor.
flacciformis, Hoffmgg. ex Kunth, Enum. Pl. ii. 493 = laevigata.
flaccosperma, *Dewey, in Am. Journ. Sc.* Ser. II. ii. (1846) 245.—Am. bor.
flagellifera, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 342.—N. Zel.
Flagellum, J. F. Gmel. Syst. 138 = Scleria Flagellum.
flava, Boiss. Voy. Espagne, ii. 631 = Oederi.
flava, *Linn. Sp. Pl.* 975.—Reg. bor. temp.; Tasman.
flavescens, Host, Gram. iv. 53. t. 96 = fulva.
flavicans, F. Nyland. Spicil. Pl. Fenn. iii. 15 = salina.
flavocuspis, *Franch. & Sav. Enum. Pl. Jap.* ii. 574.—Japon.
flavo-fulva, Beurl. in Bot. Notiser (1853) 37 = flava.
flectens, *Boott, Illustr. Carex*, iv. 171.—Japon.
flexilis, D. Don, in Till. Mag. (Dec. 1823) 455 = gracilis.
flexilis, Rudge, in Trans. Linn. Soc. vii. (1804) 98. t. 10 = castanea.
flexuosa, Muhl. ex Schkuhr, Riedgr. ii. 74 = debilis.
floresiana, Hochst. ex Seub. Fl. Azor. 22. t. 3. f. 1 = Vulcani.
floribunda, *Boeck. in Linnaea*, xl. (1876) 335.—Reg. Himal.
floridana, *Schwein. in Ann. Lyc. N. York*, i. (1824) 66.—Florida.
fluviatilis, *Boott, Illustr. Carex*, iv. 172.—Burma.
foeminea, Steud. Syn. Pl. Cyp. 203 = acutata.
foenea, *Muhl. ex Torr. in Ann. Lyc. N. York*, iii. (1836) 315.—Am. bor.
foenea, Willd. Enum. Hort. Berol. ii. 957 = straminea.
foetida, *All. Fl. Pedem.* ii. 265.—Europ.
foliosa, All. l. c. 270 = flava, caespitosa.
foliosa, *D. Don, Prod. Fl. Nep.* 42.—Reg. Himal.
foliosissima, *F. Schmidt, Reisen Amurl.* 195.—Reg. Amur.

CAREX:—
folliculacea, *Crantz, Inst.* i. 405.—Hab. ?
folliculata, Ell. Sketch, ii. 545 = turgescens.
folliculata, *Linn. Sp. Pl.* 978.—Am. bor.
folliculata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 152 = intumescens.
Fonki, *Phil. in Linnaea*, xxix. (1857–58) 83.—Chili.
Fontanesiana, DC. Fl. Fr. iii. 102 = pyrenaica.
Fontanesii, Poir. Encyc. Suppl. iii. 257 = divisa.
Forficula, *Franch. & Sav. Enum. Pl. Jap.* ii. 557.—Japon.
formosa, *Dewey, in Am. Journ. Sc.* viii. (1824) 98. t. B. fig. 6.—Am. bor.
Forsteri, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 154 = C. Pseudo-cyperus.
fragilis, *Boott, Illustr. Carex*, i. 7.—Reg. Himal.
Frankii, Kunth, Enum. Pl. ii. 498 = stenolepis.
Frankii, *Steud. Nom.* ed. II. i. 290.—Am. bor.
Franklinii, *Boott, in Hook. Fl. Bor. Am.* ii. 217.—Am. bor.
Fraseri, *Andr. Bot. Rep.* t. 639.—Am. bor.
Fraseriana, Ker-Gawl. in Bot. Mag. t. 1391 = Fraseri.
fraterna, *Rupr. Symb. Pl. Ross.* 208.—Rossia.
Fridrichsthaliana, Steud. Syn. Pl. Cyp. 211 = caespitosa.
frigida, *All. Fl. Pedem.* ii. 270.—Reg. bor.
frigida, Vill. Hist. Pl. Dauph. ii. 215 = ferruginea.
frigida, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 154 = fuliginosa.
frigida, Willd. ex Kunth, Enum. Pl. ii. 464, 465 = firma, fimbriata.
frisica, H. Koch, in Flora, xxix. (1846) 273 = trinervis.
fuircnioides, *Gaudich. Freyc. Voy. Bot.* 412.—Ins. Marian.
fuliginosa, Host, Gram. iv. 52. t. 93 = fimbriata.
fuliginosa, *Schkuhr, Riedgr.* i. 91.—Lappon.
fulva, Boiss. Voy. Bot. Espagne, ii. 631 = Hornschuchiana.
fulva, Gaertn. Mey. & Scherb. Fl. Wett. iii. 330 = distans.
fulva, *Good. in Trans. Linn. Soc.* ii. (1794) 177. t. 20. f. 6.—Europ.; Am. bor.
fulva, Hornem. in Fl. Dan. t. 1768 = binervis?
fulva, Host, Gram. iv. 53. t. 95 = varia.
fulva, Muhl. Desc. Gram. Uber. 246 = Elliotii.
fulva, Thuill. Fl. Par. 483 = teretiuscula, paradoxa.
fulvicoma, Dewey, in Am. Journ. Sc. xxix. (1836) 249 = chordorrhiza.
funiformis, [Clairv.] Man. Herb. 287 = chordorrhiza.
furcata, Ell. Sketch, ii. 552 = comosa.
furcata, *Lapeyr. Hist. Abr. Pl. Pyr.* 568.—Mont. Pyren.
furculata, Peterm. in Flora, xxvii. (1844) 329 = muricata.
furva, Webb, It. Hisp. 5 = lagopina.
fusca, All. Fl. Pedem. ii. 269 = caespitosa.
fusca, Schkuhr, Riedgr. i. 64 = pulla.
fusca, Soland. ex Boott, Illustr. Carex, iv. 217 = pumila.
fuscescens, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 517.—Japon.
fuscescens, Willd. ex Kunth, Enum. Pl. ii. 373 = divisa.
fuscoatia, *Boeck. in Linnaea*, xl. (1876) 378.—Peruv.
fuscula, Urv. in Mém. Soc. Linn. Par. iv. (1826) 599 = indecora.
fusiformis, Chapm. ex Boott, Illustr. Carex, i. 37 = laxiflora.
fusiformis, *Nees, in Wight, Contrib.* 128.—Reg. Himal.
gaimardioides, *Desv. in C. Gay, Fl. Chil.* vi. 202.—Chili.
Galeottiana, C. A. Mey. in Bull. Acad. Brux. ix. (1842) ii. 248 = Jamesoni.
Gasparrinii, *Parl. Fl. Ital.* iv. 149.—Italia.
Gaudichaudiana, Kunth, Enum. Pl. ii. 417 = vulgaris.
Gaudiniana, *Guthnick, in Flora*, xv. (1832) 241.—Helvet.
Gayana, *Desv. in C. Gay, Fl. Chil.* vi. 205.—Chili.
Gayiana, Steud. Nom. ed. II. i. 290 = asturica.
Gebhardi, Willd. Sp. Pl. iv. 240; Schkuhr, Riedgr. ii. 22 = elongata.
Gebhardii, Hoppe, Caric. t. 30 = canescens.
Gebleri, Prescott, ex Boott, in Proc. Linn. Soc. i. (1846) 285 = songarica.

CAREX:—
gemella, *Hochst. ex Steud. Syn. Pl. Cyp.* 214.—Ind. or.
geminata, Schkuhr, Riedgr. i. 65 = ternaria.
geniculata, Host, Gram. iv. 52. t. 94 = frigida.
geniculata, Schleich. ex Kunth, Enum. Pl. ii. 466 = ferruginea.
genuensis, DC. Cat. Hort. Monsp. 87 = glauca.
georgiana, *Dewey, ex Steud. Syn. Pl. Cyp.* 225.—Am. bor.
germana, *Boott, in Hook. f. Fl. Antarct.* 466.—Chili.
Gersneri, Suter, Fl. Helv. ii. 248 = humilis.
Geyerii, *Boott, in Trans. Linn. Soc.* xx. (1846) 118.—Am. bor. occ.
gibba, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 148.—Japon.
Gibsoni, Bab. in Jard. Ann. Nat. Hist. xi. (1843) 168. t. 5 = vulgaris.
gigantea, *Rudge, in Trans. Linn. Soc.* vii. (1804) 99. t. 10. f. 2.—Am. bor.
Gilliesii, *Phil. in Anal. Univ. Chil.* (1873) 556.—Chili.
Giraudiana, *Steud. Syn. Pl. Cyp.* 209.—Ins. Borbon.
glabra, Boott, Illustr. Carex, ii. 93 = venusta.
glandulosa, Ell. ex Boott, l. c. 89 = glaucescens.
glareosa, *Schkuhr, ex Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 146.—Scandinav.
glauca, Bosc, ex Boott, Illustr. Carex, ii. 89 = glaucescens.
glauca, *Scop. Fl. Carn.* ed. II. ii. 223.—Europ.; Afr. bor.; Am. bor.
glaucescens, *Ell. Sketch*, ii. 553.—Am. bor.
glaucina, *Boeck. in Linnaea*, xl. (1876) 353.—Ind. or.
glaucodea, *Tuckerm. ex Olney, in Proc. Am. Acad.* vii. (1868) 395.—Am. bor.
Glazioviana, *Boeck. in Flora*, lxv. (1882) 59.—Bras.
Glehni, *F. Schmidt, Reisen Amurl.* 194.—Reg. Amur.
globosa, *Boott, in Proc. Linn. Soc.* i. (1846) 259.—Calif.
globularis, Leyss. ex Kunth, Enum. Pl. ii. 439 = praecox.
globularis, *Linn. Sp. Pl.* 976.—Scandinav.
globularis, Oed. in Fl. Dan. t. 443 = panicea.
globularis, Roth, Tent. Fl. Germ. i. 397 = obesa.
globularis, Suter, Fl. Helv. ii. 249 = ericetorum.
globularis, Vahl, in Skr. Naturh. Selsk. ii. 1. (1792) 45 = rotundata.
globularis, Vill. Hist. Pl. Dauph. ii. 211 = canescens.
globularis, Willd. in Ges. Naturf. Fr. Neue Schr. i. (1794) 45. t. 2. f. 1 = montana.
glomerata, Gilib. Exercit. ii. 545 = vulpina.
glomerata, Host, Gram. i. t. 44 = stenophylla.
glomerata, Schkuhr, Riedgr. i. 79. t. 1. f. 41 = obesa.
glomerata, *Thunb. Prod. Pl. Cap.* 14.—Afr. austr.; Am. bor. et austr.
Gmelini, *Hook. & Arn. in Bot. Beech. Voy.* 118. t. 27.—Kamtschatka.
Godefrini, Willem. Phyt. iii. 1114 = laevigata.
Goodenowii, J. Gay, in Ann. Sc. Nat. Sér. II. xi. (1839) 191 = vulgaris.
Goyeni, *Petrie, in Trans. N. Z. Inst.* xiv. (1882) 363.—N. Zel.
gracilescens, Hartm. ex Anderss. Cyp. Scand. 16 = riparia.
gracilescens, *Steud. Syn. Pl. Cyp.* 226.—Am. bor.
gracilipes, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 151.—Japon.
gracilis, *R. Br. Prod.* 242.—Austral.
gracilis, Curt. Fl. Lond. fasc. iv. t. 60 = acuta.
gracilis, Ehrh. Beitr. vi. 83, nomen; Schkuhr, Riedgr. i. 48 = loliacea.
gracilis, A. Gray, in Am. Journ. Sc. Ser. II. iv. (1847) 19 = tenella.
gracilis, Honck. Syn. Pl. Germ. i. 368 = mucronata.
gracilis, Moench, Meth. 323 = montana.
gracilis, Wimm. in Jahresb. Schles. Ges. (1849) 79; et in Flora, xxxiii. (1850) 620 = stricta.
gracillima, Hoppe, ex Nym. Consp. 775 = panicea.
gracillima, *Schwein. in Ann. Lyc. N. York*, i. (1824) 66.—Am. bor.
gracillima, Steud. & Hochst. ex Steud. Nom. ed. II. i. 290 = ferruginea.
Graeffeana, *Boeck. in Flora*, lviii. (1875) 123.—Ins. Fiji.
Grahami, Boott, in Proc. Linn. Soc. i. (1843) 180; et Trans. Linn. Soc. xix. (1845) 215 = saxatilis.
granularioides, Schwein. in Ann. Lyc. N. York, i. (1824) 69 = conoidea.

CAREX:—
granularis, *Muhl. ex Schkuhr, Riedgr.* ii. 67.—Am. bor.
Grasmanniana, Rabenh. Fl. Lusat.; ex Steud. Syn. Pl. Cyp. 218 = tomentosa.
Grayana, Dewey, in Am. Journ. Sc. xxv. (1834) 141 = livida.
Grayii, *Carey, in A. Gray, Man. Bot. U. St.* ed. I. 563.—Am. bor.
Greeniana, Dewey, in Am. Journ. Sc. xxx. (1836) 61 = fulva.
Greenii, *Boeck. in Flora*, xli. (1858) 649.—Am. bor.
Griffithii, *Boott, in Proc. Linn. Soc.* i. (1846) 286.—Afghan.
Grioletti, Roem. in Schkuhr, Riedgr. ii. 76 = grisea.
grisea, *Viv. Ann. Bot.* i. 186.—Reg. Caucas.; Persia.
grisea, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 154.—Am. bor.
Grosseckii, Heuff. ex Boott, Illustr. Carex, iv. 148 = pyrenaica.
Grypos, Schkuhr, Riedgr. ii. 18 = echinata.
guestphalica, Boenn. ex Koch, Syn. Fl. Germ. ed. II. 867 = muricata.
Gunniana, *Boott, in Proc. Linn. Soc.* i. (1846) 258.—Austral.
Guthnickiana, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 289 = decipiens.
gynandra, *Schwein. in Ann. Lyc. N. York*, i. (1824) 70.—Am. bor.
gynandra, Sieber, ex Kunth, Enum. Pl. ii. 478 = Linkii.
gynobasis, Vill. Hist. Pl. Dauph. ii. 206 = Halleriana.
gynocrates, Wormsk. in Drej. Rev. Caric. 16 = chordorrhiza.
gynodynama, *Olney, in Proc. Am. Acad.* vii. (1868) 394.—Calif.
Gynomane, Bertol. Pl. Rar. Ital. Dec. ii. 43 = Linkii.
Haastiana, *Boeck. in Flora*, lxi. (1878) 168.—N. Zel.
haematolepis, Drej. in Fl. Dan. t. 2370 = salina.
haematostoma, *Nees, in Wight, Contrib.* 125.—Nepal.; Reg. Himal.
Haenkeana, Presl, Rel. Haenk. iii. 203 = C. Pseudocyperus.
Hailstoni, Gibs. in Phytol. i. (1841) 870 = axillaris.
hakonensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 550.—Japon.
Haleana, *Olney, Car. Bor. Am.* 6.—Am. bor.
Halei, Dewey, in Am. Journ. Sc. Ser. II. ii. (1846) 248 = C. Crus-corvi.
Halei, *Carey, in Chapm. Fl. S. U. St.* 543.—Am. bor.
Halleri, Vest, ex Steud. Nom. ed. II. i. 290.—Cf. Suteri.
Halleriana, *Asso, ex Honck. Syn.* i. 372.—Europ.; Reg. Casp.; Marocco.
Halliana, *L. Bailey, in Coult. Bot. Gaz.* ix. (1884) 117.—Am. bor. occ.
Hallii, *Olney, in Hayden, Rep.* (1871) 496.—Am. bor. occ.
halmaturina, Boeck. in Linnaea, xxxix. (1875) 100 = paniculata, tereticaulis.
halophila, Heuff. in Linnaea, xxxi. (1861–62) 695 = fulva.
halophila, *F. Nyland. Spicil. Pl. Fenn.* ii. n. 50.—Lappon.
Halseyana, Dewey, in Am. Journ. Sc. xi. (1826) 313 = polymorpha.
hamata, Forst. ex Steud. Nom. ed. II. i. 290 = Forsteri.
hamata, Sw. Prod. Veg. Ind. Occ. 18 = Uncinia jamaicensis.
Hancockiana, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) 1. 66.—China.
Harlandi, *Boott, Illustr. Carex*, ii. 87. t. 255.—Hongkong.
Hartii, *Dewey, in Am. Journ. Sc.* Ser. II. xli. (1866) 226.—Am. bor.
Hartwegii, *Boott, in Benth. Pl. Hartw.* 96.—N. Granat.
Haydeniana, *Olney, in Bot. King, Exp.* 366.—Am. bor. occ.
Haydenii, Dewey, in Am. Journ. Sc. Ser. II. xviii. (1854) 103 = aperta.
hebecarpa, Hook. & Arn. Bot. Beech. Voy. 50 = Hookeri.
hebecarpa, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 223. t. 12.—Reg. Himal.
hebetata, *Boott, Illustr. Carex*, iv. 172.—Am. austr.
Heleonastes, *Ehrh. ex Linn. f. Suppl.* 414.—Europ.
Heleonastes, Liljebl. in Vet. Akad. Nya Handl. Stockh. (1793) 73 = glareosa.

CAREX:—
Helferi, *Boeck. in Linnaea*, xl. (1876) 365.—Burma.
helodes, Link, in Schrad. Journ. ii. (1799) 309 = laevigata.
helvola, *Blytt, in Fries, Bot. Notiser* (1849), 58.—Norveg.
helvetica, Honck. Syn. Pl. Germ. i. 375 = frigida.
helvetica, Schleich. ex Kunth, Enum. Pl. ii. 448 = punctata.
Hepburnii, Boott, in Hook. Fl. Bor. Am. ii. 209. t. 207 = nardina.
heptastachys, *Boeck. in Linnaea*, xxxix. (1875) 114.—Am. trop.
hermaphrodita, J. F. Gmel. Syst. 138 = Elyna spicata.
hermaphrodita, Jacq. Coll. iv. 174 = Mariscus Jacquini.
heterolepis, *Bunge, Enum. Pl. Chin. Bor.* 69.—China.
heteroneura, *S. Wats. in Bot. Calif.* ii. 240.—Calif.
heterophylla, *Krock. ex Steud. Nom.* ed. II. i. 290.—Silesia.
heterosperma, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 151 = anceps.
heterostachya, Bunge, Enum. Pl. Chin. Bor. 69 = nutans.
heterostachya, Torr. in Am. Journ. Sc. Ser. II. ii. (1846) 248 = Crawei.
hians, *Steud. Syn. Pl. Cyp.* 236.—Hab. ?
Hilairei, *Boott, Illustr. Carex*, iv. 146.—Bras.
Hildebrandtiana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 516.—Madag.
Hilgendorfiana, *Boeck. l. c.* 518.—Japon.
hirsuta, Suter, Fl. Helv. ed. II. ii. 324 = Halleri.
hirsuta, Willd. Sp. Pl. iv. 252 = triceps.
hirta, *Linn. Sp. Pl.* 975.—Europ.; As. temp.
hirta, Oed. in Fl. Dan. t. 379 = filiformis.
hirtaeformis, Pers. Syn. ii. 547 = hirta.
hirtella, *Drej. Symb.* 21.—Afghan.; Reg. Himal.
hirtella, J. F. Gmel. Syst. 138 = Scleria hirtella.
hirtissima, *S. Wats. in Bot. Calif.* ii. 247.—Calif.
hispida, *Gaud. Agrost. Helv.* ii. 136.—Helvet.
hispida, Link, Hort. Berol. ii. 331 = lasiochlaena.
hispida, Willd. ex Schkuhr, Riedgr. i. 63 = echinata.
hispidula, Gaud. Agrost. Helv. 136 = fimbriata.
Hitchcockiana, *Dewey, in Am. Journ. Sc.* x. (1826) 274.—Am. bor.
Hochstetteri, Frank, ex Steud. Nom. ed. II. i. 291 = polytrichoides.
Hochstetteriana, *J. Gay, ex Seub. Fl. Azor.* 22. t. 2. f. 2.—Ins. Azoric.
holostoma, *Drej. in Fl. Dan.* t. 2428.—Groenl.
homalocarpa, Peterm. in Flora, xxvii. (1844) 333 = vulgaris.
homoiolepis, *Franch. & Sav. Enum. Pl. Jap.* ii. 567.—Japon.
Hoodii, *Boott, in Hook. Fl. Bor. Am.* ii. 211.—Am. arct.
Hookerana, *Dewey, in Am. Journ. Sc.* xxix. (1836) 248.—Am. arct.
Hookeri, *Kunth, Enum. Pl.* ii. 490.—Chili.
Hoppneri, Boott, in Hook. Fl. Bor. Am. ii. 219 = salina.
hordeiformis, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 152 = hordeistichos.
hordeistichos, *Vill. Prosp.* 18.—Europ.; As. temp.
Hornschuchiana, *Hoppe, in Flora*, vii. (1824) 599.—Europ.
Horsfieldii, *Boott, in Proc. Linn. Soc.* i. (1845) 257.—Java.
Hosteana, DC. Cat. Hort. Monsp. 88 = fulva.
Hostii, Hoppe, ex Kunth, Enum. Pl. ii. 372 = divisa.
Hostii, Schkuhr, Riedgr. Nachtr. 7 = stenophylla.
Houghtonii, *Torr. in Ann. Lyc. N. York*, iii. (1836) 413.—Am. bor. occ.
Huetiana, *Boiss. Diagn.* Ser. II. iv. 123.—Reg. Caucas.
huitensis, Steud. Syn. Pl. Cyp. 223 = indecora.
Humboldtiana, *Steud. l. c.* 208.—N. Granat.
humilis, *Leyss. Fl. Hal.* 175.—Europ.; Reg. Caucas.
humilis, Willd. ex Kunth, Enum. Pl. ii. 411 = caespitosa.
hyalina, Boott, in Bost. Journ. Nat. Hist. v. (1847) 112 = straminea.
hyalinolepis, *Steud. Syn. Pl. Cyp.* 235.—Am. bor.
hybrida, Belev. ex Lam. Encyc. iii. 382 = divisa.
hybrida, Schkuhr, ex Kunth, Enum. Pl. ii. 533 = Kobresia caricina.
hydrophila, Dum. Fl. Belg. 146 = echinata.
hymelayensis, *Steud. Syn. Pl. Cyp.* 204.—Reg. Himal.
hymenina, *Nees, ex Boott, Illustr. Carex*, i. 67.—Chili.

CAREX:—
hymenocarpa, Drej. in Fl. Dan. t. 2849 = ampullacea.
hymenolepis, Nees, in Wight, Contrib. 126 = setigera.
hypandra, *F. Muell. Fragm.* viii. 259.—Austral.
hyperborea, Drej. Revis. Caric. 43 = salina.
hypoxanthos, Steud. Syn. Pl. Cyp. 193 = Brongniartii.
hypsophila, *Miq. Fl. Ind. Bat.* iii. 354.—Malaya.
hystericina, *Muhl. ex Schkuhr, Riedgr.* ii. 69.—Am. bor.
Hystrix, A. Gray, ex Boott, Illustr. Carex, i. 26 = Cruscorvi.
Idzuroei, *Franch. & Sav. Enum. Pl. Jap.* ii. 583.—Japon.
ignota, Dewey, in Am. Journ. Sc. Ser. II. viii. (1849) 348 = laxiflora.
illegitima, *Cesati, in Frieder. Reise*, 271.—Dalmat.; Graecia.
illinoënsis, *Dewey, in Am. Journ. Sc.* Ser. II. iv. (1847) 245.—Am. bor.
illustranda, Steud. Syn. Pl. Cyp. 190 = bracteosa.
Ilseana ×, Ruhm. in Eichl. Jahrb. i. (1881) 258.—Europ.
impunctata, *Boott, Illustr. Carex*, iii. 107.—Java.
inaequalis, Raeusch. Nom. ed. III. 270 = caespitosa?
inanis, *Kunth, Enum. Pl.* ii. 522.—Reg. Himal.
incerta, Steud. Syn. Pl. Cyp. 210 = saxatilis.
incisa, *Boott, in Perry, Jap. Exp.* ii. 327.—Japon.
incisodentata, Lechl. ex Boott, Illustr. Carex, i. 26 = festiva.
incisodentata, *Steud. Syn. Pl. Cyp.* 189.—Reg. Magellan.
inclusa, Turcz. ex Boott, Illustr. Carex, iv. 177 = alba.
inconspicua, Steud. Syn. Pl. Cyp. 221 = indecora.
incrassata, Schlecht. in Linnaea, x. (1836) 119 = riparia.
incrassata, Soland. ex Boott, Illustr. Carex, iv. 138 = trifida.
incurva, *Lightf. Fl. Scot.* ii. 544. t. 24.—Reg. bor. temp. et arct.
incurva, Sibth. & Sm. Fl. Graec. Prod. i. 227 = divisa.
indecora, *Kunth, Enum. Pl.* ii. 448.—Ins. Falk.
indica, Koen. ex Willd. Sp. Pl. iv. 315 = Scleria tessellata.
indica, *Linn. Mant.* ii. 574.—Ind. or.; Malaya.
indica, F. Muell. ex Benth. Fl. Austral. vii. 441 = fissilis.
indica, Schkuhr, Riedgr. i. 37 = Uncinia spartea.
indica, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 149 = Wahlenbergiana.
inflata, Huds. Fl. Angl. ed. I. 354 = vesicaria.
inflata, Suter, Fl. Helvet. ii. 265 = ampullacea.
infuscata, Nees, in Wight, Contrib. 125 = alpina.
innocua, *Wallr. Beitr. Fl. Hercyn.* 299.—Europ.
inops, Kunze, in Schkuhr, Riedgr. Suppl. 131 = tenella.
inornata, Turcz. ex Ledeb. Fl. Ross. iv. 303, nomen = ericetorum.
insignis, *Boott, Illustr. Carex*, i. 5.—Reg. Himal.
insularis, *Carmich. in Trans. Linn. Soc.* xii. (1818) 508.—Ins. Trist. d'Acunha.
intermedia, Good. in Trans. Linn. Soc. ii. (1794) 154 = disticha.
intermedia, Miégev. in Bull. Soc. Bot. Fr. x. (1863) 83 = vulgaris.
intermedia, Retz. Fl. Scand. 178 = muricata.
intermedia, Suter, Fl. Helvet. ii. 261, 262 = paludosa.
intermedia, Urv. in Mém. Soc. Linn. Par. i. (1822) 378 = arenaria.
interrupta, *Boeck. in Linnaea*, xl. (1876) 432.—Am. bor.
intricata, Tineo, in Guss. Fl. Sic. Syn. ii. 574 = vulgaris.
intumescens, *Rudge, in Trans. Linn. Soc.* vii. (1804) 97. f. 3.—Am. bor.
inundata, Willd. ex Kunth, Enum. Pl. ii. 492 = glaucescens.
inversa, *R. Br. Prod.* 242.—Austral.; N. Zeland.
involucrata, *Boeck. in Flora*, xxxviii. (1855) 594.—Texas.
involucrata, *Boott, Illustr. Carex*, ii. 76.—Am. austr.
iridifolia, Kunth, Enum. Pl. ii. 492 = aethiopica.
irregularis, Schwein. in Ann. Lyc. N. York, i. (1824) 66 = flava.
irrigua, *Sm. in Hoppe, Caric.* t. 92.—Reg. bor. temp. et arct.
ischnantha, Steud. Syn. Pl. Cyp. 204 = brunnea.
ischnostachya, *Steud. l. c.* 222.—Japon.
Jackiana, *Boott, in Proc. Linn. Soc.* i. (1846) 260.—Java.
jamaicensis, Poir. Encyc. iii. 246 = Uncinia jamaicensis.
Jamesii, Schwein. in Ann. Lyc. N. York, i. (1824) 67 = Steudelii.

CAREX:—
Jamesii, *Torr. in Ann. Lyc. N. York*, iii. (1836) 398.—Am. bor.
Jamesoni, *Boott, in Proc. Linn. Soc.* i. (1845) 258.—Ecuador.
japonica, Fisch. ex Boott, l. c. 259 = breviculmis.
japonica, *Thunb. Fl. Jap.* 38.—Japon.
juncea, Scop. Fl. Carn. ed. II. ii. 227 = fulva.
juncea, *Willd. Enum. Hort. Berol. Suppl.* 63.—Am. bor.
juncella, T. M. Fries, in Bot. Notiser, (1857) 207 = vulgaris.
juncifolia, All. Fl. Pedem. ii. 264 = incurva.
juncifolia, J. F. Gmel. Syst. 142 = mucronata.
juncifolia, Host, Syn. 504 = stenophylla.
juncoides, *J. & C. Presl, Fl. Čech.* 190.—Bohemia.
jungenda, Steud. Syn. Pl. Cyp. 210 = saxatilis.
kaloides, *Petrie, in Trans. N. Z. Inst.* xiii. (1881) 332.—N. Zel.
Kanitzii, *Floriano, in Magyar Növent. Lapok*, ix. (1885) 131.—Europ.
kattegatensis, Fries, Ind. Sem. Hort. Upsal. (1857).—Cf. Hartm. Skand. Fl. ed. XIII. 466 = salina.
Keiskei, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 148.—Japon.
Kelloggii, *S. Wats. Bot. Calif.* ii. 240.—Calif.
Kelvingtoniana, *Steud. Syn. Pl. Cyp.* 215.—Am. bor.
Kerneri, *Kohts, in Oestr. Bot. Zeitschr.* xx. (1870) 164.—Europ.
kiotensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 556.—Japon.
Kirilowii, *Turcz. in Bull. Soc. Nat. Mosc.* xxviii. (1855) i. 340.—As. centr.
Kneiskernii ×, Dewey, in Am. Journ. Sc. xxxii. (1846) 247.—Am. bor.
Kochiana, DC. Cat. Hort. Monsp. 8 = paludosa.
Kochiana, Schueb. Fl. Wuerttb. 601 = evoluta.
Koestlini, *Hochst. ex Steud. Syn. Pl. Cyp.* 193.—Abyssin.
Kotschyana, Boiss. & Hohen. Diagn. Ser. I. xiii. 38 = vulgaris.
Krameri, *Franch. & Sav. Enum. Pl. Jap.* i. 551.—Japon.
Krullii, *Boeck. in Flora*, lxv. (1882) 59.—N. Zel.
Kunzei, *Olney, ex A. Gray, in Proc. Am. Acad.* viii. (1872) 406.—Am. bor.
Lachenalii, Schkuhr, Riedgr. i. 51 = lagopina.
lachnosperma, Wall. Cat. n. 3379 = gracilis.
laciniata, *Boott, Illustr. Carex*, iv. 175.—Calif.
lacistoma, *R. Br. Prod.* 243.—Austral.
lacustris, Balb. ex Kunth, Enum. Pl. ii. 533 = Elyna caricina.
lacustris, *Schkuhr, Riedgr.* ii. 84. t. 600. f. 152.—Europ.
lacustris, Willd. Sp. Pl. iv. 306 = riparia.
laeta, *Boott, Illustr. Carex*, i. 69.—Reg. Himal.
laevicaulis, Hochst. in Seub. Fl. Azor. 21 = punctata.
laevi-conica, *Dewey, in Am. Journ. Sc.* xxiv. (1857) 47.—Am. bor.
laevigata, Boiss. Voy. Bot. Espagne, ii. 631 = Camposii.
laevigata, *Sm. in Trans. Linn. Soc.* v. (1799) 272.—Europ.
laevigata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 153 = aethiopica.
laevirostris, Anderss. Cyp. Scand. 17 = rhynchophysa.
laevis, J. F. Gmel. Syst. 138 = Scleria laevis.
laevis, Hoppe, Taschenb. (1800) 243 = dioica.
laevis, *Kit. ex Willd. Sp. Pl.* iv. 292.—Enrop. austr. or.
Laggeri, Wimm. in Flora, xxxvii. (1854) 161 = repens.
lagopina, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 145.—Reg. bor.
lagopodioides, Schkuhr, Riedgr. ii. 20 = tribuloides.
Lagopus, Muhl. Desc. Gram. Uber. 265 = Fraseri.
Lamarckii, [W. Wood] in Rees, Cycl. vi. n. 103 = obesa.
Lambertiana, *Boott, ex Hook. f. Fl. N. Zel.* 284.—N. Zel.
lanceata, Dewey, in Am. Journ. Sc. xxix. (1836) 249 = salina.
lanceolata, *Boott, in Perry, Jap. Exp.* ii. 326.—Japon.
lanescens, *Fellm. in Bull. Soc. Nat. Mosc.* viii. (1835) 283.—Lappon.
Langii, *Steud. Syn. Pl. Cyp.* 227.—Norveg.
Langsdorffii, Boott, in Proc. Linn. Soc. i. (1845) 259 = leucochlora.
lanosa, *Gueldenst. It.* i. 107; *ex Ledeb. Fl. Ross.* iv. 320.—Ross. austr.
lanuginosa, *Michx. Fl. Bor. Am.* ii. 175.—Am. bor.

CAREX:—
lapponica, Lang, in Linnaea, xxiv. (1851) 539 = canescens.
lasiocarpa, Ehrh. in Hann. Mag. 9^tes^ Stück (1784), 132 = filiformis.
lasiochlaena, *Kunth, Enum. Pl.* ii. 485.—Sardinia.
latifolia, *Boiss. & Bal. in Boiss. Fl. Orient.* v. 421.—Pontus.
latifolia, J. F. Gmel. Syst. 138 = Scleria latifolia.
latifolia, Moench, Meth. 324 = plantaginea.
latifolia, Soland. ex Boott, Illustr. Carex, i. 52 = C. Pseudo-cyperus.
laxa, Dewey, in Am. Journ. Sc. xxvi. (1834) 376 = limosa.
laxa, Sw. ex Knnth, Enum. Pl. ii. 513 = scabrella.
laxa, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 156.—Lappon.
laxiculmis, *Schwein. in Ann. Lyc. N. York*, i. (1824) 70.—Am. bor.
laxiflora, Ell. Sketch, ii. 549 = grisea.
laxiflora, *Lam. Encyc.* iii. 378.—Am. bor.
laxula, Tineo, ex Boott, Illustr. Carex, iv. 202 = sylvatica.
lazica, Boiss. & Bal. in Boiss. Fl. Orient. v. 424 = sempervirens.
Leavenworthii, Dewey, in Am. Journ. Sc. Ser. II. ii. (1846) 246 = cephalophora.
Lechleri, *Phil. in Linnaea*, xxix. (1857–58) 83.—Reg. Magellan.
Lechleri, *Steud. Syn. Pl. Cyp.* 234.—Chili.
Ledebouriana, C. A. Mey. ex Trevir. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 540 = capillaris.
Ledebourii, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 222 = natans.
Leersii, F. Schult. in Flora, liii. (1870) 455, & liv. (1871) 25 = muricata.
Leersii, Willd. Prod. Fl. Berl. 28 = stellulata.
Lehmanniana, Boott, ex Walp. Ann. i. 910 = Lemanniana.
Lehmannii, *Drej. Symb.* 13. t. 2.—Ind. or.
leiocarpa, J. Gay, in Ann. Sc. Nat. Sér. II. vi. (1836) 351 = asturica.
leiocarpa, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 208. t. 5.—Unalask.
leiorhyncha, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 217. t. 9.—Dahuria.
Lemanniana, *Boott, in Proc. Linn. Soc.* i. (1846) 256.—Am. austr.
Lemmoni, *W. Boott, in Coult. Bot. Gaz.* ix. (1884) 93.—Am. bor. occ.
lenta, D. Don, ex Spreng. Syst. iii. 841, sphalm. = gracilis.
lenticularis, Dewey, in Am. Journ. Sc. vii. (1824) 263; ix. (1825) 263 = magellanica.
lenticularis, D. Don, in Trans. Linn. Soc. xiv. (1825) 331 = phacota.
lenticularis, *Michx. Fl. Bor. Am.* ii. 172.—Am. bor.
leonura, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 161 = crinita, hystericina.
lepida, *Boott, Illustr. Carex*, iv. 211.—Peruv.
lepidocarpa, Tausch, in Flora, xvii. (1834) 179 = flava.
leporina, *Linn. Sp. Pl.* 973.—Reg. bor. temp.
leporina, Michx. Fl. Bor. Am. ii. 170 = scoparia.
leporina, Oed. in Fl. Dan. t. 294 = lagopina.
leporina, Pall. ex Kunth, Enum. Pl. ii. 393 = divisa.
leporina, Schkuhr, ex Ledeb. Fl. Ross. iv. 279 = C. Heleonastes.
leptalea, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 139 = polytrichoides.
leptophylla, *Heuff. in Linnaea*, xxxi. (1861–62) 728. t. 5.—Europ.
leptostachya, Boiss. Fl. Orient. v. 418 = seq.
leptostachys, Ehrh. ex Linn. f. Suppl. 414 = strigosa.
Lerchenfeldiana, Schur, Enum. Pl. Transs. 711 = hordeistichos.
Lessoniana, *Steud. Nom.* ed. II. i. 292.—N. Zel.
leucantha, *Arn. ex Boott, in Proc. Linn. Soc.* i. (1845) 257.—Ind. or.
leucantha, *Schur, Enum. Pl. Transs.* 714.—Europ.
leucocarpa, *Boeck. in Flora*, xxxix. (1856) 227.—Am. bor.
leucocarpa, *Phil. in Linnaea*, xxix. (1857–58) 82.—Chili.
leucochlora, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 142 = breviculmis.
leucoglochin, Ehrh. ex Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 139, 140 = obtusata.
leucoglochin, Linn. f. Suppl. 413 = pauciflora.

CAREX:—
leucolepis, Turcz. ex Steud. Nom. ed. II. i. 292 = tenuiflora.
leucorhizia, Dulac, Fl. Hautes-Pyr. 452 = Davalliana.
Lhotzkiana, Nees, ex Steud. Nom. ed. II. i. 292 = Fintelmannia restioides.
Liddoni, *Boott, in Hook. Fl. Bor. Am.* ii. 214.—Am. bor.
Liddoni, Carey, in A. Gray, Man. Bot. U. St. ed. I. 545 = pallida.
Liebmanni, *Walp. Ann.* iii. 705.—Mexic.
ligata, *Boott, ex Benth. Fl. Hongk.* 402.—China.
ligerica, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 360 = arenaria.
ligerina, Guep. ex Bor. in Mém. Soc. Acad. Maine-et-Loire, vi. (1859) 171 = praec.
ligulata, *Nees, in Wight, Contrib.* 127.—Ind. or.
limosa, Engelm. ex Steud. Syn. Pl. Cyp. 209 = abjiciens.
limosa, *Linn. Sp. Pl.* 977.—Europ.; Am. bor.
limosa, Scop. Fl. Carn. ed. II. ii. 222 = ustulata.
limula, Fries, Summa Veg. 229 = vulgaris.
Lindleyana, *Nees, in Wight, Contrib.* 121.—Ind. or.
linearis, *Boott, Illustr. Carex*, i. 51.—Reg. Himal.
linearis, Clairv. ex Boott, l. c. iv. 204 = tenuis.
lineola, *Fries, in Bot. Notiser* (1844), 53.—Lappon.
Linkii, *Willd. ex Schkuhr, Riedgr.* ii. 39.—Reg. Mediterr.
Linnaeana, Host, Gram. iii. 77 = dioica.
Linnaei, Dégl. in Loisel. Fl. Gall. 627 = dioica.
liparocarpus, Gaud. Etrennes Fl. 153 = obesa.
lipsiensis, Peterm. Fl. Lips. Excurs. 58 = flava.
lithophila, *Turcz. in Bull. Soc. Nat. Mosc.* (1838) 104; xxviii. (1855) I. 328.—Sibir.
lithosperma, Forst. f. Prod. 65 = Scleria margaritifera.
lithosperma, Linn. Syst. ed. XII. 706 = Scleria Flagellum.
lithosperma, Walt. Fl. Carol. 227 = Scleria reticularis.
littoralis, Krock. ex Steud. Nom. ed. II. i. 292 = stricta.
littoralis, *Petrie, in Trans. N. Z. Inst.* xv. (1883) 358.—N. Zel.
littoralis, *Schwein. in Ann. Lyc. N. York*, i. (1824) 70.—Am. bor.
littorea, Labill. Nov. Holl. Pl. ii. 69. t. 219 = pumila.
livida, *Willd. Sp. Pl.* iv. 285.—Reg. bor. temp.
lobata, Bell. App. Fl. Pedem. 42 = curvula.
lobata, Lam. Encyc. iii. 379; Schkuhr, Riedgr. i. 28 = divisa.
lobata, Schleich. ex Steud. Nom. ed. II. i. 293 = microstyla.
lobata, Vill. Hist. Pl. Dauph. ii. 197 = foetida.
lobata, Wahlenb. ex Steud. Nom. ed. II. i. 292 = curvula.
lobata, Willd. ex Kunth, Enum. Pl. ii. 533 = Elyna caricina.
lobolepis, *F. Muell. Fragm.* viii. 258.—Austral.
lobulirostris, *Drej. Symb.* 27. t. 14.—Zeylan.
loliacea, *Linn. Sp. Pl.* 974.—Reg. bor. et arct.
loliacea, Schkuhr, Riedgr. i. 22 = tenella.
loliacea, Schreb. ex Steud. Nom. ed. II. i. 292 = divulsa.
loliacea, Steph. ex Boott, Illustr. Carex, iv. 187 = elongata.
loliacea, Thuill. ex Boott, Illustr. Carex, iv. 192 = muricata.
lonchicarpa, Willd. ex Kunth, Enum. Pl. ii. 500 = folliculata.
longearistata, Bivona, in Sicil. Sponte Prov. Desc. iv. 8. t. 2 = echinata.
longebrachiata, *Boeck. in Linnaea*, xli. (1877) 282.—Austral.
longebracteata, *Steud. Syn. Pl. Cyp.* 205.—Java.
longerostrata, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 220.—Kamtschatka.
longibracteata, Dulac, Fl. Hautes-Pyr. 57 = C. Pseudocyperus.
longibracteata, Schleich. Cat. Pl. Helv. ed. IV. 11 = intermedia.
longicaulis, *Boeck. in Flora*, lxv. (1882) 62.—Mexic.
longicruris, *Nees, in Wight, Contrib.* 124.—Ind. or.
longiculmis, *Petrie, in Trans. N. Z. Inst.* xiv. (1882) 363.—N. Zel.
longifolia, *R. Br. Prod.* 242.—Austral.
longifolia, Burch. ex Hemsl. Bot. Challeng. Exped. i. II. 87 = aequabilis.

CAREX:—
longifolia, Host, Gram. iv. t. 85 = praecox.
longifolia, Thuill. Fl. Par. i. 490 = ampullacea.
longipes, *D. Don, in Trans. Linn. Soc.* xiv. (1825) 329.—Ind. or.
longirostris, Krock. ex Kunth, Enum. Pl. ii. 493 = laevigata.
longirostris, *Torr. ex Schwein. in Ann. Lyc. N. York*, i. (1824) 71.—Am. bor.
longiseta, *Brot. Fl. Lusit.* i. 63.—Lusitan.
Lorentziana, *Griseb. in Goett. Abh.* xix. (1874) 271.—Reg. Argent.
Loscosii, *Lange, in Kjoeb. Vidensk. Meddel.* (1877-78) 223.—Hispan.
lucida, *Boott, in Journ. Linn. Soc.* xx. (1884) 377.—N. Zel.
lucida, Dicks. ex Steud. Nom. ed. II. i. 292 = Elyna spicata.
lucorum, Sartw. ex Boott, Illustr. Carex, ii. 99 = nigromarginata.
lucorum, *Willd. Enum. Hort. Berol. Suppl.* 63.—Am. bor.
ludibunda, *J. Gay, in Ann. Sc. Nat.* Sér. II. x. (1838) 357.—Gallia.
lupuliformis, Sartw. ex Dewey, in Am. Journ. Sc. Ser. II. ix. (1850) 29 = lupulina.
lupulina, *Muhl. ex Schkuhr, Riedgr.* ii. 54.—Am. bor.
lurida, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 153 = lupulina.
lusitanica, Schkuhr, Riedgr. ii. 23 = paniculata.
lutensis, *Kunth, Enum. Pl.* ii. 487.—Afr. austr.
luteola, Nees, ex Boott, Illustr. Carex, iv. 181 = bracteosa.
luzulaefolia, *S. Wats. in Bot. Calif.* ii. 250.—Calif.
luzulina, *Olney, in Proc. Am. Acad.* vii. (1868) 395.—Calif.
Lyallii, *Boott, Illustr. Carex*, iv. 150.—Am. bor. occ.
Lyngbyei, Hornem. in Fl. Dan. t. 1888 = maritima.
Lyoni, *Boott, in Hook. Fl. Bor. Am.* ii. 209.—Am. bor.
Maackii, *Maxim. Prim. Fl. Amur.* 308.—Reg. Amur.
macella, *Kunth, Enum. Pl.* ii. 482.—Reg. Argent.
macilenta, Fries, Summa Veg. 224 = gracilis.
macloviana, Urv. in Mém. Soc. Linn. Par. v. (1826) 599 = leporina.
Macounii, *Dewey, in Am. Journ. Sc.* Ser. II. xli. (1866) 228.—Am. bor.
macra, Steud. Syn. Pl. Cyp. 212 = vulgaris.
macrocarpa, *Phil. in Linnaea*, xxix. (1857-58) 86.—Chili.
macrocephala, *Willd. ex Kunth, Enum. Pl.* ii. 428.—Sibir.; Japon.
macrochaeta, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 224. t. 13 = podocarpa.
macroglossa, *Franch. & Sav. Enum. Pl. Jap.* ii. 576.—Japon.
macrogyna, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104 = ferruginea.
macrokolea, *Steud. Syn. Pl. Cyp.* 223.—Am. bor.
macrolepis, *DC. Cat. Hort. Monsp.* 89.—Italia; Graecia.
macrolepis, D. Don, in Till. Mag. (Dec. 1823) 455 = longipes.
macrophylla, *Hochst. ex Steud. Syn. Pl. Cyp.* 207.—Ind. or.
macrorhyncha, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 521 = parva.
macrosolen, *Steud. Syn. Pl. Cyp.* 210.—Reg. Magell.
macrostachyos, *Link, in Schrad. Journ.* ii. (1799) 311.—Lusitan.
macrostachys, Bertol. Amoen. Ital. 421 = ferruginea.
macrostyla, Lapeyr. Hist. Abr. Pl. Pyr. 562 = decipiens.
macrothrysa, *Miq. Fl. Ind. Bat.* iii. 351.—Malaya.
maculata, *Boott, in Trans. Linn. Soc.* xx. (1851) 128.—Zeylan; As. or.; Austral.
maculata, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 268 = spilocarpa.
maculata, Turcz. ex Ledeb. Fl. Ross. iv. 295 = ustulata.
maculata, Turcz. ex Steud. Nom. ed. II. i. 292 = coriophora.
madagascariensis, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 517.—Madag.
magellanica, Banks, ex Boott, in Proc. Linn. Soc. i. (1884) 257 = Banksii.
magellanica, *Lam. Encyc.* iii. 385.—Reg. bor. et austr.

CAREX:—
Mairii, *Coss. & Germ. Obs. Pl. Crit.* 18.—Mont. Pyren.
malaena, Steud. Syn. Pl. Cyp. 211 = vulgaris.
manca, *Boott, ex Benth. Fl. Hongk.* 402.—Hongkong.
manostachys, Spreng. Syst. iii. 824, sphalm. = macrostachys.
Marchandiana, Lapeyr. ex Spreng. l. c. 806 = pyrenaica.
marcida, *Boott, in Hook. Fl. Bor. Am.* ii. 212.—Am. bor. occ.
marcida, *Raeusch. Nom.* ed. III. 270 (quid?).—Helvet.
margaritifera, Steud. Nom. ed. II. i. 292 = Scleria Flagellum.
marginata, Gort. Fl. Belg. 247 = divisa, intermedia.
marginata, Muhl. ex Schkuhr, Riedgr. ii. 49 = pensylvanica.
marina, Dewey, in Am. Journ. Sc. xxix. (1836) 247 = C. Heleonastes.
maritima, *O. F. Muell. in Fl. Dan.* t. 703.—Reg. bor. et arct.
Marssoni, Auersw. in Bot. Zeit. x. (1852) 409 = flava.
Maubertiana, *Boott, Illustr. Carex*, i. 45.—Cochinch.
mauritanica, *Boiss. & Reut. Pugill. Pl. Nov.* 116.—Afr. bor.
mauritanica, *Steud. Syn. Pl. Cyp.* 204.—Ins. Maurit.
maxima, Scop. Fl. Carn. ed. II. ii. 229 = pendula.
Maximowiczii, *Boeck. in Linnaea*, xli. (1877) 237.—Reg. Amur.
Maximowiczii, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 180 = picta.
Maximowiczii, *F. Schmidt, Reisen Amurl.* 71.—As. temp.
Meadii, Dewey, in Am. Journ. Sc. xliii. (1842) 90 = tetanica.
media, R. Br. in Richards. in Frankl. Narr. 1st Journ. 750 = alpina.
meiogyna, Nees, in Wight, Contrib. 123 = filicina.
melaena, Wimm. in Flora, xxxiii. (1850) 619 = vulgaris.
melanantha, *C. A. Mey. in Ledeb. Fl. Ross.* iv. 216.—As. centr.
melanocarpa, *Cham. in Middend. Sibir. Reise*, i. II. 21.—Sibiria.
melanocephala, Turcz. ex Kunth, Enum. Pl. ii. 437 = alpina.
melanochloros, Thuill. Fl. Par. i. 488 = stricta.
melanocystis, Desv. in C. Gay, Fl. Chil. vi. 203 = incurva.
melanosperma, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 270.—Mexic.
melanostachya, Willd. Sp. Pl. iv. 299 = nutans.
melanostylis, *Desv. in. C. Gay, Fl. Chil.* vi. t. 73.—Chili.
melastoma, Fisch. ex Boott, Illustr. Carex, ii. 100 = Tolmiei.
membranacea, Hook. in Parry, 2 Voy. App. 406 = vesicaria.
membranacea, Hoppe, Ic. t. 71 = ericetorum.
mendocinensis, *Olney, in S. Wats. Bot. Calif.* ii. 249.—Calif.
Menziesiana, Sm. ex Boott, Illustr. Carex, i. 27 = macrocephala.
mercarensis, *Hochst. ex Steud. Syn. Pl. Cyp.* 194.—Ind. or.
Mertensii, *Presc. ex Bong. Vég. Sitcha*, 51.—Alaska.
mertensis, Weihe, ex Kunth, Enum. Pl. ii. 383 = vulpina.
Metteniana, Lehm. ex Schur, Enum. Pl. Transs. 697 = dioica.
mexicana, Presl, Rel. Haenk. i. 204 = polystachya.
Meyeni, Nees, in Nov. Act. Nat. Cur. xix. (1843) Suppl. i. 123 = brunnea.
Meyeriana, *Kunth, Enum. Pl.* ii. 438.—Sibir. alt.
micans, *Boott, in Mem. Am. Acad.* N. S. vi. (1858-59) 419 = maculata.
Michauxiana, *Boeck. in Linnaea*, xli. (1877) 336.—Am. bor.
Michauxii, Dewey, in Am. Journ. Sc. x. (1826) 273 = subulata.
Michauxii, Schwein. in Ann. Lyc. N. York, i. (1824) 64 = scirpoidea.
Micheliana, Sm. in Trans. Linn. Soc. v. (1800) 279 = glauca.
Michelii, *Host, Syn. Pl. Austr.* 507.—Europ. austr.; As. temp.

CAREX:—
microcarpa, *Bertol. ex Moris, Stirp. Sard. Elench.* 48. —Sardinia.
microcephala, C. A. Mey. in Ledeb. Fl. Alt. iv. 205 = obtusata.
microdontos, *Torr. in Ann. Lyc. N. York*, iii. (1836) 423.—Texas.
microglochin, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 140.—Reg. bor. et arct.
microlepis, *Boeck. in Linnaea*, xli. (1877) 269.—Hab.?
micropoda, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 210.—Unalaschk.
microsperma, *Steud. Syn. Pl. Cyp.* 231.—Texas.
microsperma, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 144 = vulpinoidea.
microstachya, *Ehrh. in Hannov. Mag.* 9^tes^ Stück (1784), 132.—Europ. bor.
microstachya, Michx. Fl. Bor. Am. ii. 169 = polytrichoides.
microstyla, *J. Gay, in Gaud. Fl. Helv.* vi. 37.—Alp. Helv.
Middendorffii, *F. Schmidt, Reisen Amurl.* 70.—Reg. Amur.
Mielichhoferi, Schkuhr, Riedgr. ii. 66 = ferruginea.
Mielichhoferi, Sm. Engl. Bot. t. 2293 = vaginata.
Mielichhoferi, Willd. ex Kunth, Enum. Pl. ii. 465 = brachystachys, fimbriata.
miliacea, Muhl. ex Schkuhr, Riedgr. ii. 77 = prasina.
miliacea, Schrank, Baier. Fl. i. 300 = distans.
miliaris, *Michx. Fl. Bor. Am.* ii. 174.—Am. bor.
minima, *Boullu, in Ann. Soc. Bot. Lyon*, v. (1878) 88. —Gallia.
mirabilis, Dewey, in Am. Journ. Sc. xxx. (1836) 63 = straminea.
mirabilis, Host, Gram. iv. t. 78 = Elyna caricina.
mirabilis, Schleich. Cat. (1821) ex Gaud. Fl. Helv. vi. 63 = curvula.
mirata, *Dewey, in Am. Journ. Sc.* Ser. II. xxxix. (1865) 71.—Am. bor.
misandra, *R. Br. in Parry, Voy. App.* 283.—Am. arct.
miser, Buckl. in Am. Journ. Sc. xlv. (1843) 173 = juncea.
misera, *Phil. Fl. Atac.* 53.—Chili.
Mitchelliana, *M. A. Curt. in Am. Journ. Sc.* xliv. (1843) 84.—Am. bor.
mitis, J. F. Gmel. Syst. 138 = Scleria mitis.
mixta, J. F. Gmel. l. c. 146 = stricta, paludosa.
mixta, *Miégev. in Bull. Soc. Bot. Fr.* xii. (1865) 343. —Gallia.
modesta, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 304 = disticha.
Moenchiana, Wender. in Flora, vi. (1823) 718 = acuta.
moesta, *Kunth, Enum. Pl.* ii. 482.—Reg. Argent.
mollicula, *Boott, Illustr. Carex*, iv. 192.—Ins. Loochoo.
mollis, Gilib. Exercit. ii. 546 = leporina.
mollis, Host, Gram. iv. 46. t. 82 = praecox.
monadelpha, Boott, in Perry, Jap. Exped. ii. 323 = tristachya.
Moniezi, *Timb. in Bull. Soc. Bot. Fr.* iv. (1857) 164.—Gallia.
monile, *Tuckerm. ex Boott, in Hook. Lond. Journ. Bot.* v. (1846) 69.—Am. bor.
monilifera, Thuill. Fl. Par. i. 490 = depanperata.
Moniziana, *Lowe, ex Boott, Illustr. Carex*, iv. 174.—Ins. Mader.
monostachya, *A. Rich. Tent. Fl. Abyss.* ii. 512.—Abyssin.
montana, Leers, Fl. Herborn. 716. t. 16. f. 6 = ericetorum.
montana, Linn. Sp. Pl. 975 = pilulifera.
montana, *Linn. Fl. Suec.* ed. II. 328.—Europ.; Reg. Caucas.; Japon.
montana, Poll. ex Lightf. Fl. Scot. ii. 551 = praecox.
montana, Scop. Fl. Carn. ed. II. ii. 219 = tomentosa.
monticola, Dewey, ex Torr. Bot. Mex. Bound. 229 = triquetra.
Moorcroftii, *Falc. in Proc. Linn. Soc.* i. (1846) 288.—Reg. Himal.
Morrowii, *Boott, in Perry, Jap. Exp.* ii. 326.—Japon.
Motoskei, *Miq. in Ann. Mus. Bot. Lugd. Bat.* ii. 148. —Japon.
mucronata, *All. Fl. Pedem.* ii. 268.—Europ. med.; Reg. Caucas.
Muehlenbergii, Brongn. Voy. Duperr. Bot. 151 = Brongniartii.
Muehlenbergii, *Schkuhr, Riedgr.* ii. 12.—Am. bor.

CAREX:—
Muelleri, Steud. Syn. Pl. Cyp. 204 = Linkii.
Muhlenbergii, Kunth, ex Boott, Illustr. Carex, iii. 125 = vulpinoidea.
multicaulis, *L. Bailey, in Coult. Bot. Gaz.* ix. (1884) 118.—Calif.
multiceps, Gaud. Etrennes Fl. 78 = elongata.
multiculmis, Ehrh. Beitr. vi. 83 = elongata.
multiflora, Muhl. ex Schkuhr, Riedgr. ii. 14 = vulpinoidea.
multiformis, Thuill. Fl. Par. i. 479 = disticha.
multinervis, *Boeck. in Flora*, xxxviii. (1855) 596.—Texas.
multinervis, Krock. ex Steud. Nom. ed. II. i. 293 = binervis.
multispicata, *Kunze, in Schkuhr, Riedgr. Suppl.* 42. t. 10.—Chili.
munda, *Boott, Illustr. Carex*, i. 7.—Reg. Himal.
muricata, Desf. Fl. Atlant. ii. 335 = divulsa.
muricata, Huds. Fl. Angl. 406 = echinata,
muricata, Jungh. in Linnaea, vi. (1831) 29 = glomerata.
muricata, Lecrs, Fl. Herborn. t. 13. f. 8 = stellulata.
muricata, *Linn. Sp. Pl.* 974.—Reg. bor. temp.
muricata, Schlecht. in Linnaea, vi. (1831) 29 = Brongniartii.
musei, *Steud. Syn. Pl. Cyp.* 205.—Ins. Borbon.
muskingumensis, *Schwein. in Ann. Lyc. N. York*, i. (1824) 66.—Am. bor.
mutabilis, Willd. Prod. Fl. Berl. 37 = pendula.
mutans, Boott, ex C. B. Clarke, in Journ. Linn. Soc. xx. (1883) 383 = Hemicarex Hookeri.
mutica, R. Br. in Richards. in Frankl. Narr. 1st Journ. 751 = aurea.
myosuroides, *Lowe, in Trans. Camb. Phil. Soc.* iv. (1833) 10.—Ins. Azoric.
myosuroides, Vill. Prosp. 17 = Elyna spicata.
myosurus, *Nees, in Wight, Contrib.* 122.—Ind. or.
nana, *Boott, in Mem. Am. Acad.* N. S. vi. (1858-59) 418.—Japon.
nana, *Cham. ex Steud. Syn. Pl. Cyp.* 228.—Alaska.
nana, Lam. Encyc. iii. 389 = ferruginea.
nardifolia, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 165 = aquatilis.
nardina, *Fries, Nov. Mant.* ii. 55.—Europ. et Am. bor. et arct.
Naufragii, Hochst. & Steud. Nom. ed. II. i. 293; Syn. Pl. Cyp. 206 = illegitima.
Naumanniana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 518.—Japon.
nebrascensis, Dewey, in Am. Journ. Sc. Ser. II. xviii. (1854) 102 = Jamesii.
Neesiana, Arn. ex Boott, in Proc. Linn. Soc. i. (1845) 260 = lobulirostris.
Neesiana, *Endl. Prod. Fl. Norf.* 24.—Ins. Norfolk.
neglecta, Degl. in Loisel. Fl. Gall. ii. 298 = distans.
neglecta, Peterm. in Flora, xxvii. (1844) 331 = vulgaris.
neglecta, Tuckerm. Enum. 19; et in Am. Journ. Sc. xlv. (1843) 40 = rosea.
nematosperma, Willd. ex Kunth, Enum. Pl. ii. 456 = anceps.
nemorensis, J. F. Gmel. Syst. 143 = pilosa.
nemorosa, J. F. Gmel. l. c. 1520 = pilosa.
nemorosa, Hort. Berol. ex Kunth, Enum. Pl. ii. 388 = leiorhyncha.
nemorosa, Lumnitz. ex Honck. Syn. i. 355 = muricata.
nemorosa, Rebent. ex Boott, Illustr. Carex, iii. 122 = vulpina.
nemorosa, Schrank, Baier. Fl. i. 291 = alba.
nemostachys, *Steud. in Flora*, xxix. (1846) 23.—Japon.
nepalensis, Spreng. Syst. iii. 811 = longipes.
nervata, *Franch. & Sav. Enum. Pl. Jap.* ii. 566.—Japon.
nervina, *L. Bailey, in Coult. Bot. Gaz.* x. (1885) 203. —Calif.
nervosa, Desf. Fl. Atlant. ii. 337 = extensa.
nervosa, Willd. ex Kunth, Enum. Pl. ii. 449 = distans.
neurocarpa, *Maxim. Prim. Fl. Amur.* 306.—Reg. Amur.
neurochlamys, F. Muell. Fragm. viii. 258 = maculata.
nevadensis, Boiss. & Reut. Pugill. Pl. Nov. 118 = Oederi.
nexa, *Boott, ex Benth. Fl. Hongk.* 402.—China.
nigella, *Boott, ex Hook. Fl. Bor. Am.* ii. 225.—Am. bor.
nigra, All. Fl. Pedem. ii. 267 = atrata.
nigra, Reich. Fl. Moeno-Franc. ii. 96 = acuta.
nigra, Schkuhr, Riedgr. ii. 38 = aterrima.

CAREX:—
nigra, Schwein. ex Torr. in Ann. Lyc. N. York, i. (1824) 336 = vulgaris.
nigra, *Siev. in Pall. N. Nord. Beitr.* vii. 198.—Sibir. baical.
nigricans, Dewey, in Am. Journ. Sc. xxix. (1836) 249 = dioica.
nigricans, Hombr. ex Boott, Illustr. Carex, iv. 156 = Darwinii.
nigricans, *Kit. in Linnaea*, xxxii. (1863) 320.—Europ.
nigricans, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 210. t. 7.—Unalaschk.
nigritella, Drej. in Fl. Dan. t. 2369 = stylosa.
nigrolutea, Gaud. Agrost. Helv. ii. 196 = glauca.
nigro-marginata, *Schwein. in Ann. Lyc. N. York*, i. (1824) 68.—Am. bor.
nikoënsis, *Franch. & Sav. Enum. Pl. Jap.* ii. 558.—Japon.
nilagirica, *Hochst. ex Steud. Syn. Pl. Cyp.* 207.—Ind. or.
nipponica, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 89.—Japon.
nitens, *Phil. in Anal. Univ. Chil.* (1873) 557.—Chili.
nitida, Hoppe, Taschenb. (1800) 198 = mucronata.
nitida, *Host, Gram.* i. 71.—Europ.
nitida, Willd. ex Kunth, Enum. Pl. ii. 411 = caespitosa.
nivalis, *Boott, in Proc. Linn. Soc.* i. (1845) 256.—Reg. Himal.
nivalis, Cham. ex Steud. Syn. Pl. Cyp. 184 = micropoda.
nivicola, G. Don, ex Boott, Illustr. Carex, iv. 149 = vaginata.
nobilis, *Boott, l. c.* i. 4.—Reg. Himal.
nodiflora, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 516.—Ins. Philipp.
norvegica, *Willd. ex Schkuhr, Riedgr.* i. 50.—Europ. et Am. bor.
notha, *Kunth, Enum. Pl.* ii. 421.—Reg. Himal.
notoleia, Nees, in Wight, Contrib. 121 = foliosa.
Novae-Angliae, *Schwein. in Ann. Lyc. N. York*, i. (1824) 67.—Am. bor.
Novae-Selandiae, *Boeck. in Flora*, lxi. (1878) 169.—N. Zel.
nubigena, *D. Don, in Trans. Linn. Soc.* xiv. (1825) 326.—Ind. or.
nuda, Lam. Fl. Fr. ii. 172 = leporina.
nudata, *S. Wats. Bot. Calif.* ii. 241.—Calif.
nuptialis, *Boott, Illustr. Carex*, iv. 175.—Ins. Sandvic.
nutans, *Host, Gram.* i. 61. t. 83.—Europ.; Japon.
nutans, Raeusch. Nom. ed. III. 270 = erecta.
Nuttallii, *Dewey, in Am. Journ. Sc.* xliii. (1842) 92.—Am. bor. occ.
Oakesiana, Torr. in Ann. Lyc. N. York, iii. (1836) 421 = oligosperma.
obesa, *All. Fl. Pedem.* ii. 270.—Europ.; Tauria.
obesa, Schleich. ex Kunth, Enum. Pl. ii. 411 = caespitosa.
obliqua, Turcz. in Bull. Soc. Nat. Mosc. (1838) 104 = pediformis.
oblita, *Steud. Syn. Pl. Cyp.* 231.—Am. bor.
oblonga, *J. F. Gmel. Syst.* 146.—German.
obovata, Goldb. ex Ledeb. Fl. Ross. iv. 290 = pediformis.
obscura, *Nees, in Wight, Contrib.* 126.—Reg. Himal.
obtusa, *Kit. in Linnaea*, xxxii. (1863) 319.—Hungar.
obtusangula, Ehrh. Beitr. vi. 82 = ampullacea.
obtusangula, Hort. Berol. ex Kunth, Enum. Pl. ii. 459 = glauca.
obtusangula, Salzm. ex Boott, Illustr. Carex, iv. 206 = echinata.
obtusangula, Salzm. ex Nym. Consp. 774 = hispida.
obtusata, *Liljeb. in Vet. Akad. Nya Handl. Stockh.* (1793) 69. t. 4.—Reg. bor. et arct.
ochroleuca, *Cham. ex Steud. Syn. Pl. Cyp.* 216.—Alaska.
odontolepis, *Phil. in Linnaea*, xxix. (1857-58) 82.—Chili.
Oederi, *Retz. ex Ehrh. Beitr.* vi. (1791) 83.—Reg. bor. temp.
Oederi, Willk. in Flora, xxxv. (1852) 521 = nevadensis.
oedipostyla, *Duval-Jouve, in Mém. Acad. Montp. (Sect. Sc.)* vii. (1869) 442.—Gallia.
oedocarpa, Anderss. Cyp. Scand. 26 = Oederi.
Oerstedii, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 272.—Costa Rica.
ohiotica, *Hochst. & Steud. ex Steud. Nom.* ed. II. i. 293.—Am. bor.

CAREX:—
Ohmuelleriana, Lang, in Flora, xxvi. (1843) 240 = arenaria.
Okesiana, Dewey, in Am. Journ. Sc. xiv. (1828) 351 = oligosperma?
olbiensis, *Jord. Obs. Pl. Crit.* iii. (1846) 241.—Europ.
Olgae, *Regel, Pl. Nov. Fedsch.* 83.—As. centr.
oligandra, F. Muell. ex Boott, Illustr. Carex, iv. 136 = Buxbaumii.
oligantha, *Boott, l. c.* 174. t. 589.—Am. austr.
oligantha, *Steud. Syn. Pl. Cyp.* 203.—Reg. Caucas.
oligocarpa, Hornem. ex Boott, Illustr. Carex, iv. 161 = obesa.
oligocarpa, Muhl. Descr. Gram. Uber. 242; ex Boott, l. c. i. 40 = digitalis.
oligocarpa, *Schkuhr, Riedgr.* ii. 58.—Am. bor.
oligogyna, Less. ex Kunth, Enum. Pl. ii. 442 = globularis.
oligosperma, *Michx. Fl. Bor. Am.* ii. 174.—Am. bor.
oligostachya, *Nees, in Hook. Kew Journ.* vi. (1854) 29.—Ins. Philipp.
olivacea, *Boott, in Proc. Linn. Soc.* i. (1845) 286.—Reg. Himal.
olivacea, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 267.—Mexic.
Olivcri, *Boeck. in Flora*, lxiii. (1880) 455.—Afghanist.
Oliveriana, J. Gay, ex Boiss. Fl. Orient. v. 408 = illegitima.
Olneyi ×, Boott, Illustr. Carex, i. 15.—Am. bor.
olyssiponensis, Steud. Syn. Pl. Cyp. 204 = Linkii.
omiana, *Franch. & Sav. Enum. Pl. Jap.* ii. 554.—Japon.
Onoei, *Franch. & Sav. l. c.* i. 551.—Japon.
ontakensis, *Franch. & Sav. l. c.* 550.—Japon.
orbicularis, Boott, in Proc. Linn. Soc. i. (1845) 254 = vulgaris.
oreades, C. A. Mey. in Bull. Acad. Brux. ix. II. (1842) 248 = festiva.
oregonensis, *Olney, ex A. Gray, in Proc. Am. Acad.* viii. (1872) 407.—Am. bor. occ.
orcophila, *C. A. Mey. Verz. Pfl. Cauc.* 29.—Reg. Caucas.; As. Min.
Orizabae, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 263 = festiva.
ornithopoda, *Willd. Sp. Pl.* iv. 255.—Europ.; As. bor.
ornithopodioides, Hausm. in Flora, xxxvi. (1853) 225 = ornithopoda.
ornithorhyncha, Boott, Illustr. Carex, i. 26 = C. Cruscorvi.
Orsiniana, Tenore, Fl. Nap. v. 244 = muricata.
orthostachys, C. A. Mey. in Ledeb. Fl. Alt. iv. 231 = aristata.
orthostachys, Trev. ex Nym. Consp. 769 = hirta.
ovalis, Good. in Trans. Linn. Soc. ii. (1794) 148 = leporina.
ovalis, Willd. ex Kunth, Enum. Pl. ii. 397 = scoparia.
ovata, *Burm. f. Fl. Ind.* 194 (err. typ. 294).—Java.
ovata, Honck. Syn. i. 341 = Halleri.
ovata, C. A. Mey. in Ledeb. Fl. Alt. iv. 207 = curaica.
ovata, Rudge, in Trans. Linn. Soc. vii. (1804) 96. t. 9. f. 1 = atrata.
oxylepis, *Torr. & Hook. in Ann. Lyc. N. York*, iii. (1836) 409.—Texas.
pachygyna, *Franch. & Sav. Enum. Pl. Jap.* ii. 560.—Japon.
pachystachya, *Cham. ex Steud. Syn. Pl. Cyp.* 197.—Alaska.
pachystylis, J. Gay, in Ann. Sc. Nat. Sér. II. x. (1838) 301 = stenophylla.
pacifica, Drej. Revis. Car. 292 = caespitosa.
Pairaei, F. Schultz, in Flora, li. (1868) 303 = divulsa.
paleacea, Wahlenb. in Vet. Acad. Nya Handl. Stockh. (1803) 164 = maritima.
paleata, *Boott, in Trans. Linn. Soc.* xx. (1846) 127.—Ins. Juan Fernandez.
pallescens, *Linn. Sp. Pl.* 977.—Reg. bor. temp.
pallida, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 215 = siccata.
pallida, Salisb. Prod. 29 = pallescens.
pallidior, Degl. in Loisel. Fl. Gall. ii. 299 = punctata.
paludosa, *Good. in Trans. Linn. Soc.* ii. (1794) 202.—Reg. bor. temp.
palustris, Suter, Fl. Helvet. ii. 261 = paludosa.
panicea, Carey, in A. Gray, Man. Bot. U. St. ed. I. 521 = Meadii.
panicea, Bunge, ex Boott, in Proc. Linn. Soc. i. (1845) 288 = tenuissima?

CAREX:—
panicea, *Linn. Sp. Pl.* 977.—Reg. bor. temp.
paniculata, *Linn. Cent. Pl.* i. 31 (*Amoen. Acad.* iv. 294).—Reg. temp. bor. et austr.
paniculata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 143 = paradoxa.
paniculata, Willd. ex Kunth, Enum. Pl. ii. 390 = teretiuscula.
panormitana, Guss. Fl. Sic. Syn. ii. 575 = acuta.
papillosa, *Nees, in Hook. Journ. Bot.* ii. (1840) 398.—Guiana.
papulosa, *Boott, in Mem. Am. Acad.* N. S. vi. (1858-59) 418.—Japon.
paradoxa, Benth. Cat. Pl. Pyr. 67 = divisa.
paradoxa, J. F. Gmel. Syst. 145 = seq.?
paradoxa, *Willd. in Schrift. Naturf. Fr. Berl.* (1794) 30. t. 1.—Europ.; As. bor.
parallela, Laest. in Vet. Akad. Nya Handl. Stockh. (1822) 338 = dioica.
parciflora, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 418.—Japon.
Parkeri, *Petrie, in Trans. N. Z. Inst.* xiii. (1881) 332.—N. Zel.
Parryana, *Dewey, in Am. Journ. Sc.* xxvii. (1835) 239.—Am. bor.
parva, *Nees, in Wight, Contrib.* 120.—Reg. Himal.
parvibracteata, *Nees, l. c.* 125.—Reg. Himal.
parviflora, Gaud. Etrennes Fl. 84 = lagopina.
parviflora, Host, Gram. i. 2, 221 = atrata.
parviflora, C. A. Mey. Verz. Pfl. Cauc. 30 = alpina.
patula, Host, Gram. i. 2, 221. t. 64 = flava.
patula, Huds. Fl. Angl. ed. II. ii. 402 = pauciflora.
patula, Schkuhr, Riedgr. i. 115. t. B. bb. f. 16 = laevigata.
patula, Scop. Fl. Carn. ed. II. ii. 226 = sylvatica.
pauciflora, Balb. ex Boott, Illustr. Carex, iv. 174 = microglochin.
pauciflora, *Lightf. Fl. Scot.* ii. 543. t. 6. f. 2.—Reg. bor. temp.
pauciflora, Usteri, ex Schkuhr, Riedgr. i. 83 = firma.
paupercula, Michx. Fl. Am. Bor. ii. 172 = magellanica.
pedata, All. Fl. Pedem. ii. 168 = ornithopoda.
pedata, *Linn. Sp. Pl.* ed. II. 1384.—Europ. arct.
pediformis, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 219. t. 10.—As. bor.
pedunculariis, Wall. ex Boott, Illustr. Carex, i. 53 = speciosa.
pedunculata, *Muhl. ex Schkuhr, Riedgr.* ii. 39.—Am. bor.
pedunculata, Willd. ex Roxb. in Beats. St. Helena Tracts, 302 = praealta.
pelia, Lang, in Linnaea, xxiv. (1851) 575 = panicea.
pellita, Muhl. ex Schkuhr, Riedgr. ii. 84 = lanuginosa.
pellucida, Turcz. ex Boott, Illustr. Carex, iv. 196 = pediformis.
pendula, Geners. Elench. n. 871 = capillaris.
pendula, *Huds. Fl. Angl.* ed. I. 352; ed. II. 411.—Europ.; As. bor.
pendula, Moench, Meth. 324 = vesicaria.
pennsylvanica, Torr. in Am. Lyc. N. York, iii. (1836) 410 = varia.
pensylvanica, *Lam. Encyc.* iii. 388.—Am. bor.
pentacarpa, *Boeck. in Flora*, lviii. (1875) 265.—Java.
pentastachys, *Fisch. ex Steud. Nom.* ed. II. i. 294.—Kamtschatka.
peregrina, *Link, Hort. Berol.* i. 334.—Afr. trop.
perennis, Krock. ex Steud. Nom. ed. II. i. 294 = marginata.
Perraudieriana, *J. Gay, in Bull. Soc. Bot. Fr.* iii. (1856) 685, nomen.—Ins. Teneriff.
Personii, Schkuhr, ex Kunth, Enum. Pl. ii. 509 = Wahlenbergiana.
Persoonii, Lang, in Flora, xxv. (1842) 748, 767 = canescens.
Persoonii, Sieber, ex Koch, Syn. Fl. Germ. ed. II. 871 = elongata, Gebhardi.
peruviana, Domb. ex Boott, Illustr. Carex, iv. 138 = acutata.
peruviana, Presl, Rel. Haenk. i. 205 = extensa.
petasata, Dewey, in Am. Journ. Sc. xxix. (1836) 246 = leporina.
Petersii, C. A. Mey. ex F. Schmidt, Reisen Amurl. 194 = falcata.
Petitiana, *A. Rich. Tent. Fl. Abyss.* ii. 513.—Abyssin.
petraea, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 139 = rupestris.

CAREX:—
petricosa, *Dewey, in Am. Journ. Sc.* xxix. (1836) 246.—Am. bor.
Petriei, *Cheesem. in Trans. N. Z. Inst.* xvi. (1884) 413.—N. Zel.
phacota, *Spreng. Syst.* iii. 826.—Ind. or.; Malaya.
phaecostachya, Sm. Engl. Bot. Suppl. t. 2731 = vaginata.
phalaroides, *Kunth, Enum. Pl.* ii. 482.—Bras.
pharensis, Vis. Fl. Dalm. iii. 346 = illegitima.
phleoides, Cav. Ic. v. 40 = Uncinia phleoides.
phyllomanica, *S. Wats. Bot. Calif.* ii. 233.—Calif.
phyllophora, Kunth, Enum. Pl. ii. 424 = pauciflora.
phyllostachys, *C. A. Mey. Verz. Pfl. Cauc.* 30.—Reg. Cauc.; Persia.
physema, *Dewey, in Am. Journ. Sc.* Ser. II. xxix. (1860) 347.—Am. bor.
physocarpa, Nees, ex Boott, in Proc. Linn. Soc. i. (1846) 287 = acutata.
physocarpa, *Presl, Rel. Haenk.* iii. 202.—Am. arct.
physodes, *Bieb. in Mém. Soc. Nat. Mosc.* ii. (1809) 104. t. 7.—Reg. Caucas.; As. bor.
physorhyncha, *Steud. Syn. Pl. Cyp.* 719.—Mexic.
pichinchensis, *H. B. & K. Nov. Gen. et Sp.* i. 233.—Ecuador.
picta, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 418.—Japon.
picta, *Steud. Syn. Pl. Cyp.* 184.—Am. bor.
Pierotii, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 148.—Japon.
pilosa, *Scop. Fl. Carn.* ed. II. ii. 226.—Italia.
pilosiuscula, *Boeck. in Flora*, lxv. (1882) 61.—Am. bor.
pilosiuscula ×, *Gobi, in Just, Jahresb.* iv. (1876) 1078.—Rossia.
pilulifera, Geners. ex Boott, Illustr. Carex, iv. 216 = praecox.
pilulifera, *Linn. Sp. Pl.* 976.—Europ.; Japon.
pilulifera, *Willd. ex Kunth, Enum. Pl.* ii. 439.—Reg. bor. et arct.?
pinetorum, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 263 = festiva.
pinetorum, Willd. ex Kunth, Enum. Pl. ii. 381 = Muehlenbergii.
piniaria, Bosc, ex Boott, Illustr. Carex, iii. 124 = Muehlenbergii.
pinnata, Moench, ex Reichb. Fl. Germ. Excurs. 59 = elongata.
piptolepis, Desv. in C. Gay, Fl. Chil. vi. 201 = setifolia.
pisiformis, *Boott, in Perry, Jap. Exped.* ii. 324.—Japon.
Pissisi, *Phil. in Linnaea*, xxxiii. (1864–65) 272.—Chili.
planata, *Franch. & Sav. Enum. Pl. Jap.* ii. 555.—Japon.
planifolia, *Kohts, in Oestr. Bot. Zeitschr.* xx. (1870) 141.—Europ.
planostachys, Kunze, in Schkuhr, Riedgr. Suppl. 138. t. 35 = Hallcriana.
plantaginea, *Lam. Encyc.* iii. 392.—Am. bor.
plantaginea, Muhl. Descr. Gram. Uber. 235 = laxiflora.
platycarpa, Hochst. ex Steud. Syn. Pl. Cyp. 214 = phacota.
platyphylla, *Carey, in Am. Journ. Sc.* II. iv. (1847) 23.—Am. bor.
platyrhyncha, *Franch. & Sav. Enum. Pl. Jap.* ii. 582.—Japon.
platystachys, Tenore, Fl. Nap. iv. 133 = macrolepis.
plumbea, Willd. Sp. Pl. iv. 308 = riparia.
poaeformis, J. F. Gmel. Syst. 138 = Scleria poaeformis.
pocilliformis, *Boott, Illustr. Carex*, iv. 175.—Japon.
podocarpa, *R. Br. in Richards. in Frankl. Narr. 1st Journ.* 751.—Am. arct.
podogyna, *Franch. & Sav. Enum. Pl. Jap.* ii. 557.—Japon.
podostachys, *Steud. Syn. Pl. Cyp.* 232.—Am. bor.
Poireti, J. F. Gmel. Syst. 140 = Fuirena pubescens.
polyandra, Schkuhr, Riedgr. i. 59 = caespitosa.
polyantha, *Franch. & Sav. Enum. Pl. Jap.* ii. 556.—Japon.
polyantha, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 110 = acuta.
polycephala, *Boott, Illustr. Carex*, i. 4.—Reg. Himal.
polygama, Schkuhr, Riedgr. i. 84 = Buxbaumii.

CAREX:—
polymorpha, *Muhl. ex Boott, Illustr. Carex*, i. 21. t. 57.—Am. bor.
polymorpha, Schwein. ex Boott, l. c. iii. 125 = vulpinoidea.
polymorpha, Torr. in Ann. Lyc. N. York, iii. (1836) 413 = striata.
polyphylla, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 859.—As. centr.
polyrrhiza, Wallr. Sched. Crit. 492 = praecox.
polystachya, Lesson & A. Rich. Voy. Astrol. 118 = Lessoniana.
polystachya, A. Rich. Voy. Astrol. 111. t. 21 = ternaria.
polystachya, *Sw. ex Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 149.—Am. trop.
polysticha, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 157.—Bras.
polytrichoides, *Muhl. ex Schkuhr, Riedgr.* ii. 34.—Am. bor.
Porteri, *Olney, Car. Bor. Am.* 12.—Am. bor.
potamostauros, *Steud. in Lechl. Berb. Am. Austr.* 52.—Chili.
potosina, *Hemsl. Biol. Centr. Am. Bot.* iii. 474.—Mexic.
praealta, *Boott, Illustr. Carex*, i. 66.—Ins. S. Helen.
praeceps, Borck. ex Reichb. Fl. Germ. Excurs. 64 = praecox.
praecox, *Jacq. Fl. Austr.* v. 23.—Reg. bor. temp.
praecox, Jundz. Fl. Lithuan. n. 889 = brizoides.
praecox, Poll. ex Steud. Nom. ed. II. i. 294 = emarginata.
praecox, Schreb. Spicil. 63 = Schreberi.
praegracilis, *W. Boott, in Coult. Bot. Gaz.* ix. (1884) 87.—Am. bor.
Praetutiana, Parl. Fl. Ital. ii. 182 = glauca.
prairea, Dewey, ex Woods, Bot. 750 = teretiuscula.
prasina, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 161 = miliacea.
pratensis, Drej. Revis. Caric. 24 = adusta.
pratensis, *Hose, in Usteri, Ann. Bot.* xxi. (1797) 35.—Europ.
pratensis, *Phil. in Linnaea*, xxix. (1857–58) 81.—Chili.
Preissii, *Nees, in Lehm. Pl. Preiss.* ii. 94.—Austral.
Prescottiana, *Boott, in Proc. Linn. Soc.* i. (1845) 286.—Calif.; Ins. Hawaii.
Preslii, *Steud. Syn. Pl. Cyp.* 242.—Sinus Nutka.
Princeps, Kunth, Enum. Pl. ii. 439 = praecox.
procera, *Kunth, l. c.* 491.—Bras.
prolifera, *Kit. in Linnaea*, xxxii. (1863) 320.—Hungar.
prolixa, Fries, Nov. Mant. iii. 228 = acuta.
prolixa, Schur, Enum. Pl. Transs. 707 = stricta.
propinqua, Nees & Meyen [Reise, i. 116], ex Kunth, Enum. Pl. ii. 396 = festiva.
prostrata, All. Fl. Pedem. ii. 267 = humilis.
prostrata, J. F. Gmel. Syst. 142 = pilosa.
protracta, *Steud. Syn. Pl. Cyp.* 234.—Am. bor.
provincialis, Degl. in Loisel. Fl. Gall. ii. 307 = echinata.
pruinosa, *Boott, in Proc. Linn. Soc.* i. (1845) 255.—Java.
psammogaea, Steud. Syn. Pl. Cyp. 187 = incurva.
psammophila, Schur, Enum. Pl. Transs. 700 = brizoides.
pseudo-arenaria, Pers. Syn. ii. 536 = disticha.
pseudo-arenaria, Reichb. Ic. Fl. Germ. viii. 8. t. 208 = arenaria.
pseudo-brizoides, *Chavaud, in Bull. Soc. Pharm. Bord.* xiii. (1873) 156.—Gallia.
pseudo-Buxbaumii, *Winkler, in Oestr. Bot. Zeitschr.* xviii. (1868) 72.—Europ.
pseudo-conica, *Franch. & Sav. Enum. Pl. Jap.* ii. 570.—Japon.
pseudo-curaica, *F. Schmidt, Reisen Amurl.* 67.—Reg. Amur.
Pseudo-cyperus, R. Br. Prod. 243 = fascicularis.
Pseudo-cyperus, *Linn. Sp. Pl.* 978.—Reg. temp. bor. et austr.
pseudo-flava, *Schur, Enum. Pl. Transs.* 712.—Transsylv.
pseudo-loliacea, *F. Schmidt, Reisen Amurl.* 193.—Ins. Sachalin.
pseudo-nutans, *Bor. ex Martin, Cat. Romor.* 309.—Gallia.
pseudo-paradoxa, S. Gibson, in Phytol. i. (1844) 778 = teretiuscula.
pseudo-praecox, Schur, Enum. Pl. Transs. 716 = praecox.

CAREX:—
psilostachya, Kit. ex Willd. Sp. Pl. iv. 289 = sylvatica.
psilostachys, *Steud. Syn. Pl. Cyp.* 215.—Chili.
psychrophila, *Nees, in Wight, Contrib.* 127.—Reg. Himal.
psyllophora, Hartm. ex Reichb. Fl. Germ. Excurs. 61 = pauciflora.
psyllophora, Linn. f. Suppl. 413 = pulicaris.
pteroloma, Kunze, ex Steud. Syn. Pl. Cyp. 242 = gibba.
pterygocarpa, *Steud. l. c.* 222.—Chili.
ptychocarpa, Link, in Schrad. Journ. ii. (1799) 309 = alpestris.
ptychocarpa, *Steud. Syn. Pl. Cyp.* 234.—Am. bor.
puberula, Boott, in Perry, Jap. Exp. ii. 324 = breviculmis.
pubescens, Gilib. Exercit. ii. 547 = montana.
pubescens, *Muhl. ex Schkuhr, Riedgr.* ii. 60. t. Eee. fig. 126.—Am. bor.
pubescens, Poir. Voy. Barb. ii. 254 = Fuirena pubescens.
pulchella, *Bergg. in Minneskr. Fisiog. Sallks. Lund* (1877) n. VIII. 29.—N. Zel.
pulchella, Nyland. ex Trevir, in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 537 = gracilis.
pulchra, *Boott, Illustr. Carex*, i. 4.—Reg. Himal.
pulicarioides, Ramond, ex DC. Fl. Fr. iii. 101 = pyrenaica.
pulicaris, All. ex Boott, Illustr. Carex, iv. 174 = microglochin.
pulicaris, *Linn. Sp. Pl.* 972.—Europ.; As. bor.
pulicaris, Linn. f. ex Boott, Illustr. Carex, iv. 196 = pauciflora.
pulla, Good. in Trans. Linn. Soc. iii. (1797) 78. t. 14 = saxatilis.
pulla, Soland, ex Boott, Illustr. Carex, i. 64 = lucida.
pullulans, Dulac, Fl. Hautes-Pyr. 57 = Oederi.
pumila, Honck. Syn. Pl. Germ i. 373 = montana ?
pumila, *Thunb. Fl. Jap.* 39.—China; Japon; Reg. austr. temp.
punctata, *Gaud. Agrost. Helv.* ii. 152.—Europ.; Afr. bor.
punctata, Nees, ex Wight, Contrib. 127 = notha.
punctulata, Leers, Fl. Herborn. 713. t. 14. f. 4 = paniculata.
punctulata, Lesson, ex A. Rich. Voy. Astrol. Bot. 119. t. 21 = C. Pseudo-cyperus.
pungens, *Boeck. in Linnaea*, xxxix. (1875) 46.—Chili.
Purdiei, Boott, Illustr. Carex, i. 26 = Bonplandii.
purpurea, *Boott, l. c.* iv. 157. t. 514.—Hab. ?
purpureovaginata, *Boeck. in Kjoeb. Vidensk. Meddel.* (1879–80) 30.—Bras.
pusilla, Pers. Syn. ii. 536 = bicolor.
pycnostachya, *Desv. in C. Gay, Fl. Chil.* vi. 204.—Chili.
pycnostachya, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 522 = C. Curaica.
pygmaea, *Boeck. in Linnaea*, xl. (1876) 379.—N. Granat.
pyrenaica, F. Muell. Fragm. viii. 251 = acicularis.
pyrenaica, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 139.—Europ.; Reg. Caucas.; Am. bor.; N. Zel.
pyriformis, F. Schultz, in Pollichia, xv. (1855) 122 = aurea.
pyriformis, Tausch, in Schultz, Arch. Fl. ii. 285 = flava.
quadrangulata, *Colenso, in Trans. N. Z. Inst.* xvii. (1885) 254.—N. Zel.
quaternaria, Spreng. Syst. iii. 809 = loliacea.
quinata, Raeusch. Nom. ed. III. 270 = caespitosa ?
radicalis, *Boott, Illustr. Carex*, i. 56.—Reg. Himal.
Raeana, *Boott, in Richardson, Arctic Exp.* ii. 344.—Am. bor.
Rafflesiana, *Boott, in Proc. Linn. Soc.* i. (1845) 285.—Java.
Ramondiana, DC. Fl. Fr. iii. 101 = pyrenaica.
ramosa, Eckl. ex Kunth, Enum. Pl. ii. 531 = Schoenoxiphium Sickmannianum.
ramosa, *Schkuhr, Riedgr.* ii. 40.—Ind. or.; China.
Rasulii, *Boott, in Hook. f. Fl. N. Zeal.* 288.—N. Zel.
raphidocarpa, *Nees, ex Wight, Contrib.* 122.—Ind. or.
rara, *Boott, in Trans. Linn. Soc.* xx. (1846) 139.—Zeylan.
rariflora, *Sm. Engl. Fl.* iv. 100.—Reg. bor. et arct.
Raynoldsii, *Dewey, in Am. Journ. Sc.* Ser. II. xxxii. (1861) 39.—Am. bor.
reclinata, Facch. in Ambr. Fir. Mer. i. (1854) 333, 828 = ornithopoda.
recta, Boott, in Hook. Fl. Bor. Am. ii. 220 = salina.

CAREX:—
recurva, Huds. Fl. Angl. ed. II. 413 = glauca.
recurva, Muhl. Desc. Gram. Uber. n. 55 = cherokensis.
recurva, Schkuhr, Riedgr. i. 120 = C. Pseudo-cyperus.
recurva, Willd. ex Kunth, Enum. Pl. ii. 413 = acuta.
recurvirostra, Hall. f. ex Steud. Nom. ed. I. 156 = Davalliana.
recurvirostra, *Kunze, in Schkuhr, Riedgr. Suppl.* 79. t. 20.—Java.
Redowskiana, C. A. Mey. in Mém. Sav. Etr. Pétersb. i. (1831) 207. t. 4 = chordorrhiza.
reducta, Drej. Revis. Caric. 36 = salina.
reflexa, Dietr. ex Kunth, Enum. Pl. ii. 384 = glomerata.
reflexa, Gaud. Etrennes Fl. 53 = Davalliana.
reflexa, Hoppe, Caric. 65 = praecox.
refracta, Roth, Tent. Fl. Germ. ii. II. 451 = praecox.
refracta, *Schkuhr, Riedgr.* ii. 62.—Mons Cenis.
Reinii, *Franch. & Sav. Enum. Pl. Jap.* ii. 559.—Japon.
remota, *Linn. Cent. Pl.* i. 31.—Reg. bor. temp. et arct.
remota, Richards. in Frankl. Narr. 1st Journ. 750 = Deweyana.
remota, Thunb. Fl. Jap. 37 = gibba.
remotiuscula, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 147 = cristata.
Renschiana, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 515.—Madag.
Remyi, Boeck. in Flora, lviii. (1875) 264 = brunnea.
repens, *Bell. App. Fl. Pedem.* 42.—Italia.
repens, Schleich. ex Kunth, Enum. Pl. ii. 376 = disticha.
reticulosa, *Peterm. in Flora*, xxvii. (1844) 333.—Saxonia.
retrocurva, Dewey, in Am. Journ. Sc. Scr. II. xlii. (1866) 243 = laxiculmis.
retroflexa, Muhl. ex Schkuhr, Riedgr. ii. 11 = rosea.
retrorsa, Nees, in Linnaea, x. (1835–36) 204 = cognata.
retrorsa, *Schwein. in Ann. Lyc. N. York*, i. (1824) 71.—Am. bor.
retusa, Degl. in Loisel. Fl. Gall. 307. t. 30 = echinata.
Retzii, Nees, in Wight, Contrib. 128 = chinensis.
Reuteriana, Boiss. & Reut. in Fl. Orient. v. 411 = Huetiana.
Reuteriana, *Boiss. Pugill. Pl. Nov.* 116.—Hispan.
reversa, Gilib. Exercit. ii. 549 = C. Pseudo-cyperus.
reversa, Spreng. Syst. iii. 827 = retrorsa.
rhaphidocarpa, Kunth, Enum. Pl. ii. 512 = raphidocarpa.
rhizantha, J. F. Gmel. Syst. 144 = alpestris.
rhizodes, Blytt, ex Boott, Illustr. Carex, iv. 196 = pediformis.
rhizomatosa, Steud. Syn. Pl. Cyp. 206 = Cumingiana.
rhynchocarpa, Heuff. in Flora, xvi. (1833) 365 = brevicollis.
rhynchophysa, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop.* ix. *Suppl.* 9.—Europ. et As. bor.
rhynchophysa, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 264 = physorhyncha.
rhynchosporoides, *Steud. Syn. Pl. Cyp.* 195.—Ins. Borbon.
Richardi, Michx. Fl. Bor. Am. ii. 170 = tribuloides.
Richardi, Thuill. Fl. Par. i. 482 = canescens.
Richardsonii, *R. Br. in Richards. in Frankl. Narr. 1st Journ.* 751, 35.—Am. arct.
rigens, *L. Bailey, in Coult. Bot. Gaz.* ix. (1884) 117.—Am. bor.
rigens, Boott, in Mem. Am. Acad. N. S. vi. (1858–59) 419 = Brownii.
rigens, Thuill. Fl. Par. i. 488 = paludosa, stricta.
rigida, Clairv. ex Steud. Nom. ed. II. i. 295 = caespitosa.
rigida, Good. in Trans. Linn. Soc. ii. (1794) 198. t. 22 = vulgaris.
rigida, Schrank, Baier. Fl. i. 290 = firma.
rigida, Tuckerm. Enum. Caric. 11 = stylosa.
rigidifolia, Hochst. ex Seub. Fl. Azor. 21 = punctata.
Ringgoldiana, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 419.—Japon.
riparia, *Curt. Fl. Lond. Fasc.* iv. t. 60.—Reg. bor. et austr.
riparia, Geners. Elench. n. 874 = paludosa.
riparia, Moench, ex Steud. Nom. ed. II. i. 295 = acuta.
riparia, Muhl. Desc. Gram. Uber. 259 = lacustris.
riparia, Poir. Encyc. Suppl. iii. 282 = Uncinia riparia.

CAREX:—
rivalis, Willk. & Lange, Prod. Fl. Hisp. i. 130 = divisa.
rivularis, Koch, ex Kunth, Enum. Pl. ii. 487 = paludosa.
rivularis, Schkuhr, Riedgr. i. 30 = divisa.
robusta, F. Nyland, Spicil. Fenn. sub n. 33 = rhynchophysa.
Rochebruni, *Franch. & Sav. Enum. Pl. Jap.* ii. 555.—Japon.
Roemeriana, *Scheele, in Linnaea*, xxii. (1849) 346.—Texas.
Romanzowiana, *Cham. ex Steud. Syn. Pl. Cyp.* 216.—Alaska.
rosea, *Schkuhr, Riedgr.* ii. 15.—Am. bor.
Rossii, *Boott, in Hook. Fl. Bor. Am.* ii. 222.—Am. bor. occ.
rostellata, Gay, ex Boott, Illustr. Carex, i. 8 = notha.
rostrata, Hoppe, ex Schkuhr, Riedgr. i. 99 = Michelii.
rostrata, Michx. Fl. Bor. Am. ii. 173 = Michauxiana.
rostrata, Muhl. ex Schkuhr, l. c. ii. 54 = tentaculata.
rostrata, Stokes, in With. Bot. Arr. Brit. Pl. ed. II. ii. 1059 = ampullacea.
Rotae, De Not. Ind. Sem. Hort. Gen. (1848) 24 = paniculata.
rotundata, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 153.—Lappon.
Royleana, Nees, in Wight, Contrib. 127 = breviculmis.
rubella, *Boott, Illustr. Carex*, iv. 176.—Zeylan.
rubescens, *Boeck. in Flora*, lxv. (1882) 60.—Japon.
rudis, *Wimm. in Flora*, xxxiii. (1850) 619.—Silesia.
rufa, Lam. Fl. Fr. ii. 184 = paludosa.
rufa, Schrank, Baier. Fl. i. 305 = paludosa.
rufidula, *Steud. Syn. Pl. Cyp.* 220.—Am. bor.
rufina, *Drej. Revis. Crit. Car.* 28.—Norveg.
Rugeliana, Kunze, in Linnaea, xx. (1847) 8 = juncea.
Ruiziana, *Boeck. in Linnaea*, xxxix. (1875) 30.—Peruv.
rupestris, *All. Fl. Pedem.* ii. 264. t. 92. f. 1.—Europ.
rupestris, Turcz. ex Ledeb. Fl. Ross. iv. 267 = argunensis.
Rutenbergiana, *Boeck. in Bremen Abh.* vii. (1880) 40.—Abyssin.
sabuletorum, Bieb. ex Boott, Illustr. Carex, iv. 178 = arenaria.
sabulosa, *Turcz. ex Kunth, Enum. Pl.* ii. 432.—Sibir. baical.
sabynensis, Less. ex Kunth, l. c. 440 = praecox.
sachalinensis, *F. Schmidt, Reisen Amurl.* 194.—Ins. Sachalin.
Sachapata, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
sagittifera, Lowe, in Camb. Phil. Soc. Trans. vi. (1838) 531 = macrostyla.
salina, Boott, in Hook. Fl. Bor. Am. 219 = cryptocarpa.
salina, Bongard, ex Ledeb. Fl. Ross. iv. 281 = canescens.
salina, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1808) 165.—Norveg.
salinoides, Beurl. in Bot. Notiser, (1853) 35 = salina.
Salisiana ×, Bruegg. in Jahresb. Nat. Ges. Graub. II. xxiii–xxiv. (1880) 120.—Europ.
Sampsoni, *Hance, in Journ. Bot.* vi. (1868) 89.—China.
sandwicensis, *Boeck. in Flora*, lviii. (1875) 265.—Ins. Pacif.
sanguinea, *Boott, in Trans. Linn. Soc.* xx. (1846) 137.—Afghanist.
sarda, Link, Hort. Berol. ii. 328 = Linkii.
Sartwelliana, Dewey, in Am. Journ. Sc. xliii. (1842) 90 = disticha.
Sartwelliana, *Olney, in Proc. Am. Acad.* vii. (1868) 396.—Calif.
Sartwellii, Dewey, ex J. Carey, in A. Gray, Man. Bot. U. St. ed. I. 539 = Sartwelliana.
saskatschawana, *Boeck. in Linnaea*, xli. (1877) 159.—Am. bor.
satsumensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 558.—Japon.
saxatilis, Huds. Fl. Angl. 408 = praecox.
saxatilis, *Linn. Sp. Pl.* 976.—Europ.
saxatilis, Matusch. ex Kunth, Enum. Pl. ii. 450 = fulva.
saxatilis, Oed. in Fl. Dan. t. 159 = vulgaris.
saxatilis, Scop. Fl. Carn. ed. II. ii. 221 = atrata.
saxatilis, Suter, ex Boott, Illustr. Carex, iv. 218 = sempervirens.

CAREX:—
scaberrima, *Scheele, in Linnaea*, xxii. (1849) 345.—Texas.
scabra, Hoppe, Taschenb. (1800) 242 = Davalliana.
scabrata, *Schwein. in Ann. Lyc. N. York*, i. (1824) 69.—Am. bor.
scabrella, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 149.—Ind. occ.
scabrifolia, *Steud. Syn. Pl. Cyp.* 237.—Java.
scabrior, Sartwell, ex Boott, Illustr. Carex, iii. 125 = vulpinoidea.
scariosa, Lam. Encyc. iii. 388 = humilis.
Schaffneri, *Boeck. in Flora*, lxi. (1878) 39.—Mexic.
Schaffneri, W. Boott, ex S. Wats. in Proc. Am. Acad. xviii. (1882–83) 172 = potosina.
schedonautos, *Steud. Syn. Pl. Cyp.* 189.—Reg. Magell.
Scheuchzeri, Gaud. ex Boott, Illustr. Carex, iv. 204 = tenuis.
Scheuchzeri, J. F. Gmel. Syst. 144 = paludosa?
Schiedeana, *Kunze, in Schkuhr, Riedgr. Suppl.* 119. t. 30.—Mexic.
Schimperiana, *Boeck. in Linnaea*, xl. (1876) 373.—Abyssin.
Schkuhrii, Willd. Sp. Pl. iv. 264 = obesa.
Schmidtii, *Boeck. in Flora*, lxv. (1882) 63.—Ins. Sachalin.
Schmidtii, *Meinshaus. Nachr. Wilni-Geb.* 224.—Manchuria.
schoenoides, DC. Fl. Fr. iii. 104 = intermedia.
schoenoides, Desf. Fl. Atlant. ii. 336 = divisa.
schoenoides, *Schrank, Baier. Fl.* i. 274.—Europ.
schoenoides, *Thuill. Fl. Par.* 480.—Gallia.
Schottii, Boiss. Diagn. Ser. II. iv. 123 = sempervirens.
Schottii, *Dewey, ex Torr. Bot. Mex. Bound.* 231.—Calif.
Schraderi, Schkuhr, Riedgr. ii. 81 = laevigata.
Schreberi, Desv. Fl. Auv. 73 (Scherberi) = arenaria.
Schreberi, *Schrank, Baier. Fl.* i. 278.—Europ.; As. bor.
Schreberi, Tenore, ex Boott, Illustr. Carex, iv. 186 = divisa.
Schreberi, Willd. in Mém. Acad. Berl. 1794 (1797) 38 = praecox.
Schrenkiana, C. A. Mey. ex Ledeb. Fl. Ross. iv. 317 = vesicaria.
Schummelii, *Siegert, in Verh. Bot. Brandenb.* (1860) 112.—Europ.
Schweinitzii, *Dewey, in Am. Journ. Sc.* ix. (1825) 68.—Am. bor.
scirpina, Tuckerm. ex Boott, Illustr. Carex, iv. 150 = scirpoidea.
scirpoidea, *Michx. Fl. Bor. Am.* ii. 171.—Am. bor.
scirpoides, Schkuhr, Riedgr. ii. 19 = echinata.
scitula, *Boott, Illustr. Carex*, iv. 177.—Reg. Himal.
scoparia, *Schkuhr, Riedgr.* ii. 20.—Am. bor.
scoparia, Torr. in Ann. Lyc. N. York, iii. (1836) 394 = arida.
Scopoliana, Willd. Sp. Pl. iv. 292 = ferruginea.
Scopolii, Gaud. Agrost. Helv. ii. 168 = ferruginea.
scotica, Spreng. Syst. iii. 819 = vaginata.
Scouleri, Torr. in Ann. Lyc. N. York, iii. (1836) 398 = cryptocarpa.
secale, Ledeb. Fl. Ross. iv. 300 = hordeistichos.
secalina, Lapeyr. Hist. Abr. Pl. Pyr. ii. 576 = riparia.
secalina, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 151 = hordeistichos.
secta, *Boott, in Hook. f. Fl. N. Zel.* i. 281.—N. Zel.
seditiosa, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
Sellowiana, *Schlecht. in Linnaea*, x. (1836) 117.—Bras.
semicylindrica, *Kit. l. c.* xxxii. (1863) 319.—Hungar.
sempervirens, Ell. ex Kunth, Enum. Pl. ii. 460 = commutata?
sempervirens, Schwein. in Ann. Lyc. N. York, i. (1824) 70 = glaucescens.
sempervirens, *Vill. Hist. Pl. Dauph.* ii. 214.—Europ.
Sendtneriana ×, Bruegg. in Jahresb. Nat. Ges. Graub. II. xxiii–xxiv. (1880) 119.—Europ.
senta, *Boott, Illustr. Carex*, iv. 174, in nota.—Calif.
serotina, Tenore, Fl. Nap. iv. 133 = muricata.
serratodens, *S. Wats. Bot. Calif.* ii. 245.—Calif.
serrulata, Bivona, ex Spreng. Syst. iii. 827 = glauca.
serrulata, Mutel, Fl. Fr. 395 = echinata.
setacea, Dewey, in Am. Journ. Sc. ix. (1825) 61. t. B. f. 5 = vulpinoidea.
setacea, Frank, ex Kunth, Enum. Pl. ii. 382 = stipata.

CAREX:—
setacea, Froel. ex Kunth, Enum. Pl. ii. 409 = mucronata.
setariaeformis, Turcz. ex Boott, Illustr. Carex, iv. 189 = leiorhyncha.
seticulmis, *Boeck. in Kjoeb. Vidensk. Meddel.* (1869) 156.—Bras.
setifolia, Godr. in Mém. Soc. Emul. Doubs, Sér. II. v. (1854) 14 = divisa.
setifolia, *Kunze, Syn. Pl. Am. Austr.* 26.—Chili.
setigera, *D. Don, Prod. Fl. Nep.* 43.—Reg. Himal.
setosa, *Boott, Illustr. Carex*, iii. 108.—Reg. Himal.
setosa, Franch. & Sav. Enum. Pl. Jap. 142 = homoiolepis.
sexspicata, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 342.—N. Zel.
Shortiana, *Dewey & Torr. in Am. Journ. Sc.* xxx. (1836) 60.—Calif.
Shortii, Steud. Syn. Pl. Cyp. 202 = stenolepis.
Shortii, Torr. in Ann. Lyc. N. York, iii. (1836) 407 = Shortiana.
sibirica, Willd. ex Kunth, Enum. Pl. ii. 406 = loliacea.
sicaeformis, *Boott, in Bost. Journ. Nat. Hist.* v. (1845) 113.—Am. bor.
siccata, *Dewey, in Am. Journ. Sc.* x. (1826) 27.—Am. bor.; As. bor. or.
sicula, Presl, ex Boott, Illustr. Carex, iv. 185 = distans.
sicula, *Tineo, Pl. Rar. Sicil.* 30.—Sicilia.
sicyocarpa, Lebel, Obs. Pl. Manche (1848) 18 = praecox.
siderosticha, *Hance, in Journ. Linn. Soc.* xiii. (1873) 89.—China.
Sieberi, Nees, ex Kunth, Enum. Pl. ii. 118 = Mariscus flavus.
Sieberiana, Opiz, Natural. xi. (1826) 413 = Davalliana.
Sieboldi, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 150.—Japon.
Siegertiana, Uechtritz, ex Garcke, Fl. Deutschl. ed. VIII. 435 = hirta.
sikokiana, *Franch. & Sav. Enum. Pl. Jap.* ii. 573.—Japon.
silicea, *Olney, in Proc. Am. Acad.* vii. (1878) 393.—Am. bor.
simensis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 514 = aethiopica.
similis, Urv. in Mém. Soc. Linn. Par. iv. (1826) 599 = curta.
simplicissima, F. Muell. Fragm. ix. 191 = capillacea.
simpliciuscula, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 141 (Quid?).—Anglia.
Sinai, *Boott, in Trans. Linn. Soc.* xx. (1851) 146.—Arab.
sinaica, Nees, ex Steud. Syn. Pl. Cyp. 223 = distans.
sinensis, J. F. Gmel. Syst. 144 = chinensis.
sitchensis, *Presc. ex Bong. Veg. Sitcha*, 51.—Alaska; Am. bor. occ.
Smithii, Tausch, in Flora, iv. (1821) 559 = vaginata.
socia, Boott, in Proc. Linn. Soc. i. (1845) 254 = Arnottiana.
sociata, *Boott, in Mem. Am. Acad.* N. S. vi. (1858–59) 420.—China.
Solandri, *Boott, in Hook. f. Fl. N. Zel.* 284.—N. Zel.
Soleirolii, DC. & Duby, Bot. Gall. ii. 498 = echinata.
songorica, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 525.—Afghanist.; Soongar.
sordida, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 33.—Kamtschatka.
sororia, *Kunth, Enum. Pl.* ii. 379.—Reg. Argent.
sororia, Meinsh. Fl. Ingr. 404 = aperta.
Spachiana, *Boott, Illustr. Carex*, i. 43.—China.
spadicea, Gilib. Exercit. ii. 546 = arenaria.
spadicea, J. F. Gmel. Syst. 144 = firma.
spadicea, Host, Gram. iv. 51. t. 91; DC. Fl. Fr. iii. 120 = ferruginea.
spadicea, Roth, Tent. Fl. Germ. ii. 11. 461 = paludosa.
spadicea, Schkuhr, Riedgr. i. 90 = frigida.
spaniocarpa, Steud. Syn. Pl. Cyp. 225 = obesa.
sparganioides, Le Conte, ex Kunth, Enum. Pl. ii. 382 = stipata.
sparganioides, *Muhl. ex Schkuhr, Riedgr.* ii. 14.—Am. bor.
sparsiflora, Steud. Nom. ed. II. i. 296 = vaginata.
spartea, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 149 = Uncinia spartea.
spatiosa, *Boott, Illustr. Carex*, ii. 86.—Cochinch.

CAREX:—
speciosa, *Kunth, Enum. Pl.* ii. 504.—Ind. or.
spectabilis, Dewey, in Am. Journ. Sc. xxix. (1836) 248. t. X. f. 76 = podocarpa.
speirostachya, Sw. ex Sm. Engl. Fl. iv. 98 = fulva.
sphaerica, *Lapeyr. Hist. Abr. Pl. Pyr.* 570.—Mont. Pyren.
sphaerocarpa, Ehrh. Beitr. vi. 83 = tomentosa.
sphaerocarpa, Willd. Sp. Pl. iv. 265 = obesa.
sphaerogyne, *Baker, in Journ. Bot.* xxi. (1883) 129.—Madag.
sphaerostachya, *Dewey, in Am. Journ. Sc.* xlix. (1845) 44. t. Ee. f. 110.—Am. bor.
spicata, Brot. ex Kunth, Enum. Pl. ii. 373 = divisa.
spicata, Huds. Fl. Angl. ed. I. 353 = muricata.
spicata, Lam. Encyc. iii. 381; Pollich, Palat. ii. 562 = disticha.
spicata, Schkuhr, Riedgr. i. 11 = obtusata.
spicata, Thuill. Fl. Par. 480 = vulpina.
spicigera, *Nees, in Wight, Contrib.* 121.—Zeylan.
spiculata, *Boott, in Trans. Linn. Soc.* xx. (1851) 139.—Reg. Himal.
spiculosa, F. Nyland. Spicil. Pl. Fenn. ii. n. 49 = salina.
spilocarpa, *Steud. Syn. Pl. Cyp.* 195.—Mexic.
spinirostris, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 335.—N. Zel.
splendens, Thuill. ex Pers. Syn. iii. 536 = divisa.
splendida, Willd. Prod. Fl. Berl. 33 = filiformis.
Sprengelii, *Boeck. in Linnaea*, xl. (1876) 371.—Afr. austr.
Sprengelii, Dewey, ex Spreng. Syst. iii. 837 = longirostris.
spreta, Steud. Syn. Pl. Cyp. 212 = caespitosa.
squarrosa, *Linn. Sp. Pl.* 973.—Am. bor.
stans, Drej. Revis. Caric. 40 = aquatilis.
Steenstrupii, *Liebm. in Forhand. Scand. Naturf.* (1840) 321.—Islandia.
stellulata, Bieb. Fl. Taur. Cauc. ii. 383 = muricata.
stellulata, Good. in Trans. Linn. Soc. ii. (1794) 144 = echinata.
stenantha, *Franch. & Sav. Enum. Pl. Jap.* ii. 573.—Japon.
stenocarpa, Turcz. ex Steud. Nom. ed. II. i. 296 = frigida.
stenolepis, *Less. ex Steud. l. c.*—Norveg.
stenolepis, *Torr. in Am. Journ. Sc.* xxx. (1836) 59.—Am. bor.
stenophylla, Thien. ex Boott, Illustr. Carex, iv. 210 = incurva.
stenophylla, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 142.—Reg. bor. temp.
stenostachys, *Franch. & Sav. Enum. Pl. Jap.* ii. 569.—Japon.
sterilis, *Schkuhr, Riedgr.* ii. 3.—Am. bor.
Steudelii, Kunth, Enum. Pl. ii. 480 = Jamesii.
Steudneri, *Boeck. in Linnaea*, xl. (1876) 364.—Abyssin.
stictocarpa, Sm. Engl. Bot. t. 2772 = glauca.
stipata, *Muhl. ex Schkuhr, Riedgr.* ii. 12.—Am. bor.
stipitata, Boeck. in Linnaea, xxxix. (1875) 85 = stipata.
stolonifera, Ehrh. Beitr. vi. 83 = praecox.
stolonifera, Hoppe, Ic. n. 41 = vulgaris.
straminea, *Willd. ex Schkuhr, Riedgr.* i. 49.—Am. bor.
striata, R. Br. Prod. 243 = lacistoma.
striata, Gilib. Exercit. ii. 550 = riparia.
striata, *Michx. Fl. Bor. Am.* ii. 174.—Am. bor.
striata, Torr. ex Boott, Illustr. Carex, i. 22 = polymorpha.
striatula, Michx. Fl. Bor. Am. ii. 173 = laxiflora.
stricta, *Good. in Trans. Linn. Soc.* ii. (1794) 196. t. 21. f. 9.—Europ.; Am. bor.
stricta, Lam. Encyc. iii. 387 = angustata.
strictifolia, Opiz, ex Nym. Consp. 777 = acuta.
strictior, Dewey, ex Wood, Cl. Book. 755 = stricta.
strigosa, All. Fl. Pedem. ii. 270 = tenuis.
strigosa, *Huds. Fl. Angl.* 411.—Europ.; Reg. Caucas.
strigosa, Suter, ex Boott, Illustr. Carex, iv. 208 = firma.
strigosa, Willd. ex Kunth, Enum. Pl. ii. 470 = sylvatica.
stygia, Fries, Mant. iii. 141 = rariflora.
styloflexa, Buckl. in Am. Journ. Sc. xlv. (1843) 174 = laxiflora.
stylosa, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 222. t. 12.—Unalaschka.
subanceps, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 520.—Japon.

CAREX:—
subbiflora, *Steud. Syn. Pl. Cyp.* 234.—Am. bor.
subcompressa, *Steud. l. c.* 221.—Am. bor.
subdola, *Boott, in Proc. Linn. Soc.* i. (1845) 255.—N. Zel.
suberea, Boott, Illustr. Carex, iv. 165 = Pierotii.
subfusca, *S. Wats. in Bot. Calif.* ii. 234.—Calif.
subglobosa, Miclich. in Flora, xxii. (1839) 256 = Oederi.
sublivida, *Norrl. in Notis. Saellsk. Fauna et Fl. Fenn.* II. *Haeft* x. 51.—Europ. bor.
subramosa, Willd. ex Kunth, Enum. Pl. ii. 385 = echinata.
subrotunda, *Serres, in Bull. Soc. Bot. Fr.* iv. (1857) 440.—Gallia.
subsalsa, Nym. Syll. Suppl. 69 = fulva.
subspathacea, Wormsk. in Fl. Dan. ix. 4. t. 1530 = salina.
subulata, J. F. Gmel. Syst. 138 = Scleria subulata.
subulata, *Michx. Fl. Bor. Am.* ii. 173.—Am. bor.
subulata, Schum. ex Steud. Nom. ed. II. i. 296 = Buxbaumii.
subulata, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 152 = Boryana.
subuniflora, *Steud. Syn. Pl. Cyp.* 234.—Am. bor.
subvillosa, Bieb. Fl. Taur. Cauc. ii. 386 = tomentosa.
sudetica, Opiz, Natural. xi. (1826) 413 = atrata.
sudetica, Presl, ex Kunth, Enum. Pl. ii. 451 = vaginata.
suilla, J. Fellm. in Bull. Soc. Nat. Mosc. viii. (1835) 285 = pulla ?
sulcata, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 80.—Transsylv.
Sullivantii ×, Boott, in Am. Journ. Sc. xlii. (1842) 29.—Am. bor.
supina, *Wahlenb. in Akad. Vet. Nya Handl. Stockh.* (1803) 158.—Reg. bor. temp.
Suteri, *Kunth, Enum. Pl.* ii. 523.—Europ.
sychnocephala, *Carey, in Am. Journ. Sc.* Ser. II. x. (1847) 24.—Am. bor.
sylvatica, Dewey, l. c. Ser. I. x. (1826) 40 = arctata.
sylvatica, *Huds. Fl. Angl.* ed. I. 353.—Europ.; As. temp.
sylvatica, Maxim. ex Boeck. in Linnaea, xli. (1877) 238 = Maximowiczii.
tabina, *Steud. in Lechl. Berb. Am. Austr.* 56.—Am. austr.
tabularia, M. A. Curt. ex Boott, Illustr. Carex, i. 54 = aestivalis.
tartarea, *Ridl. in Journ. Bot.* xxi. (1883) 262.—Sumatra.
tauricola, Boiss. Fl. Orient. v. 415 = atrata.
teinogyna, *Boott, Illustr. Carex*, i. 60.—Reg. Himal.
tenebrosa, *Boott, l. c.* 88. t. 256.—China.
tenax, *Bergg. in Minneskr. Fisiog. Sallks. Lund* (1877), n. VIII. p. 27.—N. Zel.
tenax, *Chapm. ex Dewey, in Am. Journ. Sc.* Ser. II. xix. (1855) 254.—Am. bor.
tenax, *Reut. in Comptes-Rend. Soc. Hallér.* (1854-56) 130.—Europ.
tenella, Ehrh. Beitr. vi. 83 = canescens.
tenella, Froel. ex Kunth, Enum. Pl. ii. 445 = supina.
tenella, Poir. Encyc. Suppl. iii. 282 = Uncinia tenella.
tenella, *Schkuhr, Riedgr.* i. 23.—Am. bor.
tenella, Thuill. Fl. Par. i. 429 = praecox.
tenera, Dewey, in Am. Journ. Sc. viii. (1824) 97 = straminea.
Tenorii, Kunth, Enum. Pl. ii. 437 = macrolepis.
tentaculata, *Muhl. ex Willd. Sp. Pl.* iv. 266.—Am. bor.
tenuiflora, Hartm. ex Kunth, Enum. Pl. ii. 405 = loliacea.
tenuiflora, *Wahlenb. in Vet. Acad. Nya Handl. Stockh.* (1803) 147.—Europ.; Am. bor.
tenuifolia, Poir. Voy. Barb. ii. 254 = Scirpus pubescens.
tenuirostris, *Olney, in Am. Natural.* viii. (1874) 214.—Am. bor.
tenuis, Gmel. ex Steud. Nom. ed. II. i. 296 = Scleria tenuis.
tenuis, *Host, Gram. Austr.* iv. 51. t. 92.—Europ. austr.
tenuis, Nees, in Wight, Contrib. 128 = hirtella.
tenuis, Rudge, in Trans. Linn. Soc. vii. (1804) 97. t. 9. f. 2 = debilis.
tenuis, Schleich. ex Steud. Nom. ed. II. i. 296 = leporina.
tenuispica, Boeck. in Flora, xxxix. (1856) 225 = tristicha, *Boeck.*

CAREX:—
tenuispica, *Steud. Syn. Pl. Cyp.* 319.—Ins. Borbon.
tenuissima, *Boott, in Proc. Linn. Soc.* i. (1845) 288.—Japon.
teres, *Boott, Illustr. Carex*, i. 62.—Reg. Himal.
tercticaulis, *F. Muell. Fragm.* viii. 256.—Austral.
teretiuscula, *Good. in Trans. Linn. Soc.* ii. (1794) 163.—Europ.; Am. bor.
teretiuscula, Willd. ex Kunth, Enum. Pl. ii. 389 = paniculata.
teretiuscula, Willk. in Bot. Zeit. vi. (1848) 414 = chordorrhiza.
tergestina, Hoppe, ex Boott, Illustr. Carex, iv. 193 = muricata.
ternaria, *Forst. f. Prod.* 92.—N. Zel.
testacea, *Soland. ex Boott, in Hook. f. Fl. N. Zel.* i. 282.—N. Zel.
tetanica, Reichb. Fl. Germ. Excurs. 63 = vaginata.
tetanica, *Schkuhr, Riedgr.* ii. 68.—Groenl.; Am. bor.
tetanica, Schwein. & Torr. in Ann. Lyc. N. York, i. (1824) 347 = conoidea.
tetrastachya, Schcele, in Linnaea, xxii. (1849) 347 = straminea.
tetrastachya, *Traunst. in Flora*, xxxiii. (1850) 366.—Europ.
Textori, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 151 = incisa.
thecata, Boott, in Trans. Linn. Soc. xx. (1851) 143 = Preissii.
thermarum, *Phil. in Anal. Univ. Chil.* xxi. (1862) II. 385.—Chili.
Thomsonii, *Boott, Illustr. Carex*, i. 1.—Reg. Himal.
Thouarsii, *Carmich. in Trans. Linn. Soc.* xii. (1818) 508.—Ins. Tristan. D'Acunha.
Thunbergii, *Steud. in Flora*, xxix. (1846) 23.—Japon.
Thurberi, *Dewey, ex Torr. Bot. Mex. Bound.* 232.—Am. bor. occ.
thuringiaca, Willd. Sp. Pl. iv. 250 = glauca.
Thwaitesii, *Hance, in Journ. Bot.* v. (1867) 235.—Zeylan.
thyrsiflora, *Boott, Illustr. Carex*, i. 12. t. 34.—Ind. or.
tokioënsis, *Boeck. in Flora*, lxv. (1882) 63.—Japon.
Tolmiei, *Boott, in Hook. Fl. Bor. Am.* ii. 224.—Am. bor. occ.
tomentosa, Hornem. in Fl. Dan. t. 1404 = globularis.
tomentosa, Lightf. Fl. Scot. 553 = filiformis.
tomentosa, *Linn. Mant.* 123.—Europ.; Reg. Caucas.
tomentosa, C. A. Mey. Verz. Pfl. Canc. 192 = grisea.
torfacea, J. F. Gmel. Syst. 145 = ampullacea.
Torreyana, Dewey, in Am. Journ. Sc. x. (1826) 47 = Davisii.
Torreyi, *Tuckerm. Enum. Caric.* 21.—Am. bor.
torta, *Boott, ex Tuckerm. l. c.* 11.—Am. bor.
Touranginiana, Bor. Fl. Cent. Fr. ed. II. 532 = acuta.
trachyantha, Dorner, ex Heuff. in Verh. Zool. Bot. Ges. Wien, viii. (1858) 219 = praecox.
trachycarpos, Link, in Schrad. Journ. ii. (1799) 310 = glauca.
trachycystis, *Griseb. in Goett. Abh.* xxiv. (1879) 314.—Reg. Argent.
traiziscana, *F. Schmidt, Reisen Amurl.* 192.—Ins. Sachalin.
transversa, *Boott, in Hook. Fl. Bor. Am.* ii. 215. t. 216.—Japon.
transversa, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 148 = rigens.
transsilvanica, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 80 = Halleriana ?
triandra, *Gromow, ex Trautv. in Act. Hort. Petrop.* ix. (1884) 311, nomen.—Rossia.
triangularis, *Boeck. in Flora*, xxxix. (1856) 226.—Texas.
tribuloides, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 145 = lagopodioides.
tricephala, *Boeck. in Flora*, lviii. (1875) 263.—Malaya.
triceps, *Michx. Fl. Bor. Am.* ii. 170.—Am. bor.
triceps, *Schrank, Baier. Fl.* i 287.—Europ.
trichocarpa, *Muhl. ex Schkuhr, Riedgr.* ii. 47, 81.—Am. bor.
trichocarpa, Schur, Enum. Pl. Transs. 709 = sempervirens.
trichodes, *Steud. in Linnaea*, xxxix. (1875) 29.—Peruv.

CAREX:—
trichostachya, Fisch. ex Boott, Illustr. Carex, iv. 198 = podocarpa.
trichostyles, *Franch. & Sav. Enum. Pl. Jap.* ii. 581. —Japon.
tricostata, Fries, Novit. Mant. iii. 152 = vulgaris.
trifida, *Cav. Ic.* v. 41. t. 465.—Reg. Magell.
triflora, Schkuhr, Riedgr. i. 94, ii. 66 = depauperata.
triflora, Vahl, ex Kunth, Enum. Pl. ii. 460 = vaginata.
trigona, All. Fl. Pedem. ii. 269 = fulva.
trinervis, *Degl. in Loisel. Fl. Gall.* ii. 731.—Gallia.
trinervis, Nees, in Wight, Contrib. 120 = Esenbeckii.
tripartita, All. Fl. Pedem. ii. 265 = curvula.
tripartita, Lam. Encyc. iii. 339 = divisa.
tripartita, Suter, Fl. Helv. ii. 241 = lagopina.
triquetra, *Boott, in Trans. Linn. Soc.* xx. (1851) 126. —Calif.
trisperma, *Dewey, in Am. Journ. Sc.* ix. (1825) 63.—Am. bor.
trisperma, A. Gray, ex Boott, Illustr. Carex, ii. 81 = rosea.
tristachya, *Thunb. Fl. Jap.* 38.—Japon.
tristicha, *Boeck. in Flora*, xli. (1858) 651.—Am. bor.
tristicha, *Spruce, ex Boott, Illustr. Carex*, iv. 153.—Ecuador.
tristis, Bieb. Fl. Taur. Cauc. iii. 615 = sempervirens.
triticea, Wahlenb. ex Steud. Nom. ed. II. i. 299 = clavata.
truncata, *Boeck. in Flora*, xli. (1858) 649.—Am. bor.
truncata, *Fenzl, in Friederichsthal, Reise Griechenl.* 278.—Graecia.
truncatula, *Kit. in Linnaea*, xxxii. (1863) 318.—Hungar.
tuberculata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 270.—Mexic.
tuberculata, Nees, ex Boott, Illustr. Carex, ii. 76 = bonariensis.
tuberosa, Blanco, Fl. Filip. ed. I. 35 = Eleocharis tuberosa.
tuberosa, Degl. in Loisel. Fl. Gall. ii. 629 = Linkii.
tubulata, Schum. ex Boott, Illustr. Carex, iv. 136 = Buxbaumii.
Tuckermanni, *Boott, in Hook. Lond. Journ. Bot.* v. (1846) 73.—Am. bor.
tumida, Beilschmid, in Hornsch. Archiv Skand. ii. 214 = vulgaris.
tumida, *Boott, Illustr. Carex*, i. 66.—Reg. Himal.
tumidicarpa, *Anderss. in Bot. Notiser* (1849); *et Flora*, xxxiii. (1850) 297.—Scandinav.
turbinata, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 265.—Mexic.
Turczaninofii, Steud. ex Boott, Illustr. Carex, iv. 161 = obesa.
turfacea, Auct. vide *torfacea*.
turfosa, Fries, Bot. Notiser (1846), 104, 107 = caespitosa.
turfosa, Fries, Summa, 228 = vulgaris.
turgescens, Dewey, in Am. Journ. Sc. Ser. II. iii. (1847) 356 = Halei.
turgescens, *Torr. in Ann. Lyc. N. York*, iii. (1836) 419. —Am. bor.
turkestanica, *Regel, in Act. Hort. Petrop.* vii. (1880) 570.—Turkestan.
Tweediana, *Nees, in Hook. Journ. Bot.* ii. (1840) 398. —Reg. Argent.
typhina, Michx. Fl. Bor. Am. ii. 169 = squarrosa.
typhinoides, Schwein. in Ann. Lyc. N. York, i. (1824) 66 = squarrosa.
typhoides, Bory, Voy. iii. 24 = Boryana.
uda, *Maxim. Prim. Fl. Amur.* 303.—Reg. Amur.
uetliaca, Suter, Fl. Helv. i. 251 = flava.
uliginosa, Linn. Sp. Pl. 973 = Scirpus Caricis.
umbellata, *Schkuhr, Riedgr.* ii. 75.—Am. bor.
umbrosa, Host, Gram. i. 52. t. 69 = praecox.
uncifolia, *Cheesem. in Trans. N. Z. Inst.* xvi. (1884) 412.—N. Zel.
uncinata, Linn. f. Suppl. 413 = Uncinia australis.
uncinata, Schkuhr, ex Steud. Nom. ed. II. i. 297 = Uncinia jamaicensis.
uncinioides, *Boott, Illustr. Carex*, i. 8.—Reg. Himal.
undulata, Kunze, in Schkuhr, Riedgr. Suppl. 23. n. 8. t. 4. f. 2 = pallescens.
undulata, Schur, Enum. Pl. Transs. 714 = leucantha.
Urbani, *Boeck. in Engl. Bot. Jahrb.* vii. (1885) 280.—Alaska.
ursina, *Dewey, in Am. Journ. Sc.* xxvii. (1835) 240.—Am. arct.

CAREX:—
Urvillei, Brongn. in Duperr. Voy. Bot. 157 = pumila.
ustulata, *Wahlenb. in Vet. Akad. Nya Handl. Stockh.* (1803) 156.—Europ.
utriculata, *Boott, in Hook. Fl. Bor. Am.* ii. 221.—Am. bor.
vacillans, *Soland. ex Boott, in Hook. f. Fl. N. Zel.* i. 285.—N. Zel.
vacillans, Steud. Syn. Pl. Cyp. 235 = riparia.
vaginata, *Tausch, in Flora*, iv. (1821) 557.—Europ.
Vahlii, *Schkuhr, Riedgr.* i. 87.—Reg. bor. et arct.
valdiviana, *Phil. in Linnaea*, xxix. (1857–58) 84.—Chili.
valesiaca, Suter, Fl. Helv. 259 = tenuis.
valesiaca, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 155 = ambigua.
valida, *Nees, in Wight, Contrib.* 123.—China; Zeylan.
vallicola, *Dewey, in Am. Journ. Sc.* Ser. II. xxxii. (1861) 40.—Calif.
Vanheurckii, *Muell. Arg. in Heurck, Obs. Bot.* 30.—Kamtschatka.
varia, Host, Gram. i. 59. t. 80 = sempervirens.
varia, *Muhl. ex Schkuhr, Riedgr.* ii. 48.—Am. bor.
variegata, Lam. Encyc. iii. 389 = sempervirens.
variegata, *Scheele, in Linnaea*, xxiii. (1850) 565.—Am. bor.
Vaseyi, Dewey, in Am. Journ. Sc. Ser. II. xxix. (1860) 347 = monile.
ventricosa, Curt. Fl. Lond. fasc. vi. t. 68 = depauperata.
venusta, *Dewey, in Am. Journ. Sc.* xxvi. (1834) 107.—Am. bor.
verna, Chaix & Vill. Hist. Pl. Dauph. i. 312, ii. 204 = praecox.
verna, Lam. Fl. Fr. ii. 185, partim = glauca.
verna, Schkuhr, Riedgr. i. 89 = obesa.
verna, Torr. ex Kunth, Enum. Pl. ii. 439 = praecox.
verrucosa, Muhl. Desc. Gram. Uber. 261 = glaucescens.
verrucosa, Schwein. ex Torr. in Ann. Lyc. N. York, iii. (1836) 400 = commutata.
versicolor, *Dum. Fl. Belg.* 148.—Belgium.
verticillata, *Boott, Illustr. Carex*, i. 67.—Am. bor.
verticillata, *Zoll. & Moritzi, Syst. Verz. Zoll.* 98.—Java.
vesicaria, Bunge, ex Boott, in Proc. Linn. Soc. i. (1845) 261 = songorica.
vesicaria, Dewey, ex Steud. Nom. ed. II. i. 297 = bullata.
vesicaria, Leers, Fl. Herborn. t. 16. f. 2. 1 = riparia.
vesicaria, Lightf. Fl. Scot. ii. 566 = ampullacea.
vesicaria, *Linn. Sp. Pl.* 979.—Reg. bor. temp.
vesicaria, Thunb. Prod. Pl. Cap. 14 = clavata.
vesiciformis, *Peterm. in Flora*, xxvii. (1844) 339.—Saxonia.
vesiculosa, *Boott, Illustr. Carex*, iii. 107.—Ind. or.
vestita, *Willd. Sp. Pl.* iv. 263.—Am. bor.
vexilis, Wormsk. ex Boott, Illustr. Carex, iv. 191 = Michelii.
vicinalis, *Boott, l. c.* 133.—Ind. or.
victorialis, Nees, in Wight, Contrib. 126 = notha.
Villarsii, *Schkuhr, Riedgr.* ii. 5.—Gallia.
villosa, *Boott, in Perry, Jap. Exped.* ii. 327.—Japon.
villosa, Franch. & Sav. Enum. Pl. Jap. ii. 142 = Davaliana.
villosa, Stokes, Bot. Mat. Med. iv. 346 = hirta.
virens, Lam. Encyc. iii. 384 = muricata.
virens, Thuill. Fl. Par. i. 489 = acuta.
virescens, *Muhl. ex Schkuhr, Riedgr.* ii. 45.—Am. bor.
virescens, Willd. ex Kunth, Enum. Pl. ii. 430 = triceps.
virgata, *Miq. Fl. Ind. Bat.* iii. 351.—Java.
virgata, Soland. ex Hook. f. Fl. N. Zel. i. 282 = paniculata.
virginiana, Woods, in Rees, Cycl. vi. n. 100 = acuta.
virginica, Steud. Syn. Pl. Cyp. 217 = praec.
viridis, Honck. Syn. Pl. Germ. i. 372 = flava.
viridis, *Petrie, in Trans. N. Z. Inst.* xiii. (1881) 332. —N. Zel.
viridis, *Schlecht. & Cham. in Linnaea*, vi. (1831) 30. —Mexic.
viridis, *Spenn. Fl. Friburg.* i. 48.—Europ.
viridula, *Franch. & Sav. Enum. Pl. Jap.* ii. 579.—Japon.
viridula, *Michx. Fl. Bor. Am.* ii. 170.—Am. bor.
viridula, Schwein. & Torr. in Ann. Lyc. N. York, i. (1824) 320 = triceps.

CAREX:—
vitilis, Fries, Novit. Mant. iii. 137 = canescens.
Vleckii, Schwein. in Ann. Lyc. N. York, i. (1824) 69 = oligocarpa.
Vulcani, *Hochst. ex Seub. Fl. Azor.* 22. t. 3. f. 2.—Ins. Azoric.
vulgaris, *Fries, Nov. Mant.* iii. 153.—Reg. temp. bor. et austr.
vulpina, Carey, ex Dewey, in Am. Journ. Sc. Ser. II. viii. (1849) 348 = conjuncta.
vulpina, Hohen. in Bull. Soc. Nat. Mosc. (1833) 259 = muricata.
vulpina, *Linn. Sp. Pl.* 973.—Europ.; As. temp.
vulpina, Nees, in Hook. Journ. Bot. ii. (1840) 398 = bonariensis.
vulpinaeformis, Tuckerm. Enum. 9 = vulpinoidea.
vulpinaris, Nees, in Wight, Contrib. 121.—Reg. Himal.; As. bor.
vulpinoidea, *Michx. Fl. Bor. Am.* ii. 169.—Reg. Cauc.; Am. bor.
vulpinoidea, Torr. in Ann. Lyc. N. York, i. (1824) 390 = stipata.
Wahlenbergiana, *Boott, Illustr. Carex*, ii. 101. t. 301. —Ins. Maurit.
wahuensis, *C. A. Mey. in Mém. Sav. Etr. Pétersb.* i. (1831) 218. t. 10.—Ins. Oahu; Japon.
Wakatipu, *Petrie, in Trans. N. Z. Inst.* xiv. (1882) 363.—N. Zel.
Walkeri, *Arn. ex Boott, in Proc. Linn. Soc.* i. (1845) 257.—Zeylan.
Wallichiana, *Presc. in Wall. Cat.* n. 3380.—Reg. Himal.
Wallichiana, Spreng. Syst. iii. 812 = foliosa.
Washingtoniana, Dewey, in Am. Journ. Sc. x. (1826) 272 = vulgaris.
Watsoni, Boott, Illustr. Carex, iv. 203 = Vulcani.
Watsoni, *Olney, in Bot. King, Exp.* 370.—Am. bor. occ.
Watsoniana, *Steud. Syn. Pl. Cyp.* 215.—Am. bor.
Wazmanni, *Schrank, Baier. Fl.* i. 297.—Europ.
Weiheana, Boeng. ex Kunth, Enum. Pl. ii. 439 = praecox?
Welwitschi, *Boiss. ex Steud. Syn. Pl. Cyp.* 230.—Lusitan.
Whitneyi, *Olney, in Proc. Am. Acad.* vii. (1868) 394. —Calif.
Wichurai, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 519. —China.
Wightiana, *Nees, in Wight, Contrib.* 122.—Ind. or.
Wilkesii, *Olney, in Bot. U. St. Expl. Exped.* ii. 477. t. 17.—Hab.?
Willdenowii, *Schkuhr, Riedgr.* ii. 33.—Am. bor.
wiluica, *Meinsh. ex Maack, Vilyuisk. Okruge Yakutsk. Obl.* II. 308.—Sibir.
Wimmeri, *Steud. Syn. Pl. Cyp.* 215.—Silesia.
Winkleri, *Lange, in Kjoeb. Vidensk. Meddel.* (1877–78) 224.—Hispan.
Witheringii, S. F. Gray, Nat. Arr. Brit. Pl. ii. 48 = arenaria.
Wormskioldiana, Hornem. in Fl. Dan. t. 1528 = scirpoidea.
Wormskioldii, Drej. Revis. Caric. 18 = scirpoidea.
Woodii, Dewey, in Am. Journ. Sc. Ser. II. ii. (1846) 249 = tetanica.
Wrightii, *Dewey, ex Torr. Bot. Mex. Bound.* 232.—Am. bor. occ.
xalapensis, Kunth, Enum. Pl. ii. 380 = Brongniartii.
xanthophysa, Wahlenb. in Vet. Akad. Nya Handl. Stockh. (1803) 152 = folliculata.
xerocarpa, *S. H. Wright, in Am. Journ. Sc.* Ser. II. xlii. (1866) 334, in nota.—Am. bor.
yodoënsis, *Boeck. in Engl. Bot. Jahrb.* v. (1884) 515. —Japon.
Zollingeri, Kunze, ex Steud. Syn. Pl. Cyp. 221 = nemostachys.
zonata, F. Nyland. Spicil. Pl. Fenn. ii. 19 = vulgaris?

CAREYA, Roxb. Hort. Beng. 52 (1814); Fl. Ind. ii. 636 (1832). *MYRTACEAE*, Benth. & Hook. f. i. 721.
CUMBEA, Wight & Arn. Prod. i. 334 (1834).
CUMBIA, Buch.-Ham. Mysore, iii. 187 (1807).
arborea, *Roxb. ll. cc.*—Ind. or.
australis, F. Muell. Fragm. v. 183 = Barringtonia Careya.
herbacea, *Roxb. Hort. Beng.* 52; *Fl. Ind.* ii. 638.—Ind. or.
macrostachya, Jack, in Malay Misc. i. n. v. 47 = Barringtonia macrostachya.

CAREYA :—
orbiculata, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 98 = arborea.
pendula, Griff. Notul. iv. 661 = Barringtonia macrostachya.
sphaerica, *Roxb. Hort. Beng.* 52 ; *Fl. Ind.* ii. 636.—Ind. or.
valida, *Kurz, in Journ. As. Soc. Beng.* xlvi. (1877) II. 72.—Ins. Andam.

CARGILLA, Adans. Fam. ii. 130 (1763) = **Chrysogonum**, Linn. (Compos.).

CARGILLIA, R. Br. Prod. 526 (1810) = **Diospyros**, Lam. (Ebenac.).
arborea, A. Cunn. ex Hiern, in Trans. Camb. Phil. Soc. xii. (1873) 239 = D. pentamera.
australis, R. Br. Prod. 526 = D. Cargillia.
laxa, R. Br. l. c. = D. maritima.
mabacea, F. Mucll. Fragm. v. 162 = D. mabacea.
maritima, Hassk. Cat. Hort. Bog. Alt. 159 = D. maritima.
megalocarpa, F. Muell. Fragm. v. 163 = D. maritima.
pentamera, F. Muell. l. c. iv. 82 = D. pentamera.

CARICA, Linn. Gen. ed. I. 309 (1737). *PASSIFLOREAE*, Benth. & Hook. f. i. 815.
MOCINNA, Cerv. ex La Llave, in La Naturaleza, vii. (1885) Apénd. 70.
PAPAYA, Tourn. ex Linn. Syst. ed. I (1735).
VASCONCELLEA, A. St. Hil. Mém. Resed. ii. 13 (1837).
Boissieri, *Hemsl. Biol. Centr.-Am. Bot.* i. 481.—Mexic.
candamarcensis, *Hook. f. Bot. Mag.* t. 6198.—Ecuador.
candicans, *A. Gray, Bot. U. St. Expl. Exped.* i. 640.—Peruv.
cauliflora, *Jacq. Hort. Schoenb.* iii. 33.—Venezuela.
citriformis, *Jacq. f. ex Spreng. Syst.* iii. 905.—Afr. trop.
cundinamarcensis, J. Linden, Cat. n. 87 (1871) = candamarcensis.
digitata, Spreng. Syst. iii. 906 = spinosa.
dodecaphylla, *Vell. Fl. Flum.* x. t. 132.—Bras.
erythrocarpa, *Linden & André, Illustr. Hortic.* xviii. (1871) t. 51.—Ecuador.
glandulosa, *Pav. ex A. DC. in DC. Prod.* xv. 1. 417.—Peruv.
gossypifolia, *Griseb. in Goett. Abh.* xxiv. (1879) 137.—Reg. Argent.
heptaphylla, Vell. Fl. Flum. x. t. 133 = spinosa.
hermaphrodita, Blanco, Fl. Filip. ed. I. 205 = C. Papaya.
heterophylla, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 60.—N. Granat.
Mamaya, *Vell. Fl. Flum.* x. t. 131.—Bras.
microcarpa, *Jacq. Hort. Schoenb.* iii. 32.—Am. austr.
monoica, *Desf. in Ann. Mus. Par.* i. (1802) 273. t. 18.—Peruv.
nana, *Benth. Pl. Hartw.* 288.—Mexic.
Papaya, *Linn. Sp. Pl.* 1036.—Am. trop.
peltata, *Hook. & Arn. Bot. Beech. Voy.* 425. t. 98.—Nicaragua.
Posopora, *Linn. Sp. Pl.* 1036.—Peruv.
pubescens, *Lenné & C. Koch, in Ind. Sem. Hort. Bot. Berol.* (1854) 12.—Guatemala.
pyriformis, Willd. Sp. Pl. iv. 815 = C. Posopora.
scandens, *A. Gray, Bot. U. St. Expl. Exped.* i. 640.—Peruv.
spinosa, *Aubl. Pl. Gui.* ii. 908.—Guian. ; Bras.

CARICELLA, Ehrh. Beitr. iv. 146 (1789) = **Carex**, Dill.

CARICTERIA, Scop. Introd. 104 (1777) = **Corchorus**, Tourn. (Tiliac.).

CARIGOLA, Rafin. Fl. Tellur. ii. 10 (1836) = **Monochoria**, Presl (Pontederac.).
hastata, Rafin. l. c. = M. hastaefolia.

CARIMA, Rafin. Fl. Tellur. iv. 62 (1836) = **Adhatoda**, Nees (Acanthac.).
sulcata, Rafin. l. c. = A. sulcata.

CARINIANA, Casar. Nov. Stirp. Bras. 35 (1842) = **Couratari**, Aubl. (Myrtac.).
brasiliensis, Casar. l. c. = Couratari legalis.

CARINIANA :—
domestica, Miers, in Trans. Linn. Soc. xxx. (1874) 286 = Couratari domestica.
excelsa, Casar. Nov. Stirp. Bras. 46 = Couratari estrellensis.
exigua, Miers, in Trans. Linn. Soc. xxx. (1874) 289.—Ind. occ.
pyriformis, Miers, l. c. 290.—Bolivia.
rubra, Miers, l. c. 288 = Couratari rubra.
uahupensis, Miers, l. c. = Couratari uaupensis.

CARIONIA, Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 311. t. 15. *MELASTOMACEAE*, Benth. & Hook. f. i. 759.
elegans, *Naud. l. c.* 312.—Ins. Philipp.
triplinervia, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 310.—Ins. Philipp.

CARISSA, Linn. Mant. i. 7 (1767). *APOCYNACEAE*, Benth. & Hook. f. ii. 695.
ANTURA, Forsk. Fl. Aegypt. Arab. 63 (1775).
ARDUINA, Mill. Ic. t. 300 (1759) ; Linn. Mant. 7 (1767).
abyssinica, *R. Br. in Salt, Abyss. App.* p. lxiv, nomen.—Abyss.
acuminata, *A. DC. Prod.* viii. 335.—Afr. austr.
africana, *A. DC. l. c.* 332.—Afr. trop.
Arduina, *Lam. Encyc.* i. 555.—Afr. austr.
axillaris, *Roxb. Hort. Beng.* 84 ; *Fl. Ind.* i. 691.—Ins. Molucc.
bispinosa, Desf. Tabl. ed. I. 78 = C. Arduina.
Brownii, F. Muell. Fragm. iv. 45 = lanceolata, ovata, et scabra.
Candolleana, *Jaub. & Spach, Illustr.* v. 107. t. 497.—Abyssin.
Carandas, *Linn. Mant.* 52.—Ind. or. ; Malaya.
Carandas, Lodd. Bot. Cab. t. 663 = paucinervia.
Carandas, Lour. Fl. Cochinch. 155 = africana.
congesta, Wight, Ic. iv. t. 1289 = C. Carandas.
coriacea, *Wall. Cat.* n. 1680.—Ind. or.
cornifolia, *Jaub. & Spach, Illustr.* v. 108. i. 498.—Abyssin.
cryptophlebia, *Baker, in Journ. Linn. Soc.* xx. (1884) 204.—Madag.
Dalzellii, Bedd. Forest. Fl. Anal. Gen. 156 ; Man. Bot. 157 = macrophylla.
densiflora, *Baker, in Journ. Linn. Soc.* xx. (1884) 204.—Madag.
diffusa, Roxb. Hort. Beng. 19 ; Fl. Ind. i. 689 = spinarum.
diffusa, Wall. Cat. n. 1678, partim = paucinervia.
dulcis, Schum. & Thonn. Beskr. Guin. Pl. 146 = edulis.
edulis, *Vahl, Symb. Bot.* i. 22.—Aegypt.
erythrocarpa, *A. DC. Prod.* viii. 335.—Afr. austr.
ferox, *A. DC. l. c.*—Afr. austr.
grandiflora, *A. DC. l. c.*—Afr. austr.
grandis, *Bert. ex A. DC. l. c.* 336.—Ins. Tahiti.
haematocarpa, *A. DC. l. c.*—Afr. austr.
hirsuta, Roth, Nov. Pl. Sp. 128 = spinarum.
inermis, Vahl, Symb. Bot. iii. 43 = macrophylla.
lanceolata, *R. Br. Prod.* 468.—Austral.
lanceolata, Dalz. in Dalz. & Gibs. Bomb. Fl. 143 = macrophylla.
laxiflora, *Benth. Fl. Austral.* iv. 305.—Austral.
macrocarpa, *A. DC. Prod.* viii. 336.—Afr. austr.
macrophylla, *Wall. Cat.* n. 1679.—Ind. or.
madagascariensis, *Thou. Obs. Pl. Iles Afr. Austr.* 80.—Madag.
mitis, Heynh. ex A. DC. Prod. viii. 332 = macrophylla.
mitis, Vahl, Symb. Bot. iii. 44 = Holarrhena mitis.
myrtoides, *Desf. Cat. Hort. Par.* ed. III. 398.—Hab. ?
oblongifolia, *Hochst. in Flora*, xxvii. (1844) ii. 827.—Afr. austr.
ovata, *R. Br. Prod.* 468.—Austral.
paucinervia, *A. DC. Prod.* viii. 333.—Ind. or.
pubescens, *A. DC. l. c.* 334.—Afr. trop.
Richardiana, Jaub. & Spach, Illustr. v. 106 t. 496 = edulis.
salicina, Lam. Encyc. i. 554 = spinarum ?
scabra, *R. Br. Prod.* 468.—Austral.
Schimperi, *A. DC. Prod.* viii. 675.—Abyssin.
sechellensis, *Baker, Fl. Maurit.* 222.—Ins. Seychell.
spinarum, *Linn. Mant.* t. 559.—Ins. Molucc.
spinarum, Lodd. ex A. DC. Prod. viii. 336 = Dammacanthus indicus.
suavissima, *Bedd. ex Hook. f. Fl. Brit. Ind.* iii. 632.—Ind. or.

CARISSA :—
tomentosa, *A. Rich. Tent. Fl. Abyss.* ii. 30.—Abyssin.
villosa, Roxb. Hort. Beng. 19 ; Fl. Ind. i. 690 = spinarum.
Xylopicron, *Thou. Obs. Pl. Iles Afr. Austr.* 80. t. 2.—Ins. Mascar.

CARLEA, Presl, Epim. Bot. 216 (1849) = **Symplocos**, Linn. (Styrac.).
oblongifolia, Presl, l. c. 217 = S. oblongifolia.

CARLEMANNIA, Benth. in Hook. Kew Journ. v. (1853) 308. *RUBIACEAE*, Benth. & Hook. f. ii. 63.
congesta, *Hook. f. Fl. Brit. Ind.* iii. 85.—Reg. Himal.
Griffithii, *Benth. in Hook. Kew Journ.* v. (1853) 308.—Reg. Himal.
tetragona, *Hook. f. Fl. Brit. Ind.* iii. 85.—Reg. Himal.

CARLINA, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 465.
ATHAMUS, Neck. Elem. i. 85 (1790).
CARLOWIZIA, Moench, Meth. Suppl. 225 (1802).
CHROMATOLEPIS, Dulac, Fl. Hautes-Pyr. 526 (1867).
MITINA, Adans. Fam. ii. 116 (1763).
acanthifolia, *All. Fl. Pedem.* i. 156. t. 51.—Reg. Mediterr.
Acarna, Bieb. Fl. Taur. Cauc. ii. 282 = Cnicus Acarna.
acaulis, Lam. Encyc. i. 623 = acanthifolia.
acaulis, *Linn. Sp. Pl.* 828.—Europ.
acaulis, Stokes, Bot. Mat. Med. iv. 156 = Cnicus acaulis.
aculeata, Burm. f. Fl. Cap. Prod. 23 (= *Stobaea rigida*).—Afr. austr.
aggregata, Willd. Sp. Pl. iii. 1694 = acaulis.
alpina, Jacq. Enum. Stirp. Vindob. 274 = acaulis.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 19.—Algeria.
atractyloides, Linn. Sp. Pl. 1161 (= *Stobaea atractyloides*).—Afr. austr.
Biebersteinii, Bernh. ex Hornem. Hort. Hafn. Suppl. 94 = vulgaris.
bracteata, J. & C. Presl, Delic. Prag. 103 = sicula.
caulescens, Gilib. Fl. Lituan. iii. 195 = vulgaris.
caulescens, Lam. Fl. Fr. ii. 7 = acaulis.
Chamaeleon, Vill. Hist. Pl. Dauph. iii. 30 = acaulis.
Chardousse, Vill. Prosp. 27 = acanthifolia.
complanata, Schur, Enum. Pl. Transs. 413 = vulgaris.
corymbosa, *Linn. Sp. Pl.* 828.—Reg. Mediterr.
corymbosa, Scop. ex Steud. Nom. ed. II. i. 299 = vulgaris.
Curetum, Heldr. in Boiss. Fl. Orient. iii. 449 = corymbosa.
Cynara, Pourr. ex DC. Fl. Fr. iv. 123 = acanthifolia.
discolor, Soleir. ex Nym. Consp. 400 = macrocephala.
Echinus, Bieb. Fl. Taur. Cauc. ii. 283.—Reg. Cauc. (= *Cirsium scleranthum*).
elatior, Wallr. in Linnaea, xiv. (1840) 636 ; et Beitr. Fl. Hercyn. 266 = acaulis.
foliosa, Boiss. & Haussk. in Boiss. Fl. Orient. iii. 450 = oligocephala.
Fontanesii, DC. Prod. vi. 548 = Atractylis macrocephala.
frigida, *Boiss. & Heldr. Diagn.* Ser. I. vi. 109.—Graecia.
gorterioides, Lam. Encyc. i. 625 (= Berkheyae sp.).—Afr. austr.
graeca, Heldr. & Sart. ex Boiss. Fl. Orient. iii. 449 = corymbosa.
grandiflora, Moench, Meth. 558 = acaulis.
gummifera, Less. Syn. Comp. 12 = Atractylis gummifera.
hispanica, Lam. Encyc. i. 624 = corymbosa.
intermedia, Schur, Enum. Pl. Transs. 413 = vulgaris.
involucrata, *Poir. Voy. Barb.* ii. 234.—Reg. Mediterr.
lanata, Habl. Taur. 160 = vulgaris.
lanata, *Linn. Sp. Pl.* 828.—Reg. Mediterr.
lanata, Pall. ex DC. Prod. vi. 546 = vulgaris.
libanotica, Boiss. Diagn. Ser. I. x. 95 = corymbosa.
Lobeliana, Tausch, in Flora, xii. (1829) Erg. 41 = corymbosa.
longifolia, Reichb. Ic. Fl. Germ. viii. 25 = vulgaris.
longifolia, Viv. Fl. Lib. Spec. 68 = corymbosa.
lyrata, *Thunb. Prod. Pl. Cap.* 141.—Afr. austr.
macrocephala, Less. Syn. Comp. 12 = Atractylis macrocephala.

CARLINA:—
macrocephala, *Moris, Elench. Sard.* ii. 5.—Reg. Mediterr.
macrophylla, DC. Prod. vi. 548 = Atractylis macrophylla.
nebrodensis, *Guss. in DC. Prod.* vi. 546.—Sicilia.
nebrodensis, Koch, ex Nym. Consp. 401 = vulgaris.
oligocephala, *Boiss. & Kotschy, Diagn.* Ser. II. iii. 51.—As. Min.; Syria.
onopordifolia, Bess. ex DC. Prod. vi. 545 = acanthifolia.
orophila, Lamotte, Prod. ii. 443 = vulgaris.
patula, Burm. f. Fl. Cap. Prod. 23 (= *Stobaeae sp.*).—Afr. austr.
Pola, Hacq. Pl. Alp. Carn. 29. t. 4. f. 3 = lanata.
pyrenaica, Linn. Sp. Pl. 829 = Carduus carlinoides.
racemosa, Gilib. Chlor. Grodn. 34 = vulgaris.
racemosa, Gouan, Illustr. 64 = corymbosa.
racemosa, *Linn. Sp. Pl.* 829.—Reg. Mediterr. occ.
radiata, Viv. Fl. Lib. Spec. 68 = corymbosa.
Reboudiana, *Pomel, Nouv. Mat. Fl. Atl.* 273.—Algeria.
Rothii, Heldr. & Sart. ex Boiss. Fl. Orient. iii. 449 = corymbosa.
salicifolia, *Cav. in Anal. Cienc. Nat.* iv. (1801) 81; *Less. in Linnaea*, vi. (1831) 88.—Ins. Madera.
scabra, Burm. f. Fl. Cap. Prod. 23 = Tripteris arborescens.
sicula, *Tenore, Cat. Hort. Neap. App.* (1819).—Sicilia.
simplex, Waldst. & Kit. Pl. Rar. Hung. ii. 164. t. 152 = acaulis.
subacaulis, DC. Fl. Fr. iv. 122 = acaulis.
sulphurea, Desf. Fl. Atlant. ii. 251 = racemosa.
tragacanthifolia, *Klatt, in Leopoldina*, xx. (1884) 94.—Ins. Rhodus.
transsilvanica, Schur, in Verh. Siebenb. Ver. Nat. iv. (1853) 34 = acanthifolia.
tristis, Salisb. Prod. 185 = vulgaris.
Utzka, Hacq. Pl. Alp. Carn. 7. t. 1 = acanthifolia.
vulgaris, *Linn. Sp. Pl.* 828.—Europ.; As. bor.
xeranthemoides, *Linn. f. Suppl.* 349.—Ins. Tenerif.

CARLOTEA, Arruda, ex Koster, Trav. 493 (1816) (Gen. ignot.).
formosissimum, Arruda, l. c.—Pernambuc.
speciosa, Arruda, l. c.—Pernambuc.

CARLOWIZIA, Moench, Meth. Suppl. 225 (1802) = **Carlina**, Linn. (Compos.).
corymbosa, Cass. in Dict. Sc. Nat. xxv. 53 = Carlina salicifolia.
salicifolia, Moench, Meth. Suppl. 226 = Carlina salicifolia.

CARLOWRIGHTIA, A. Gray, in Proc. Am. Acad. xiii. (1878) 364. *ACANTHACEAE.*
arizonica, *A. Gray, l. c.*—Arizona.
linearifolia, *A. Gray, l. c.*—Texas.

CARLUDOVICA, Ruiz & Pav. Prod. 146. t. 31 (1794). *CYCLANTHACEAE*, Benth. & Hook. f. iii. 953.
EVODIANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1875) 194.
LUDOVIA, Pers. Syn. ii. 576 (1807).
SALMIA, Willd. in Ges. Naturf. Fr. Berl. Mag. v. (1811) 399.
SARCINANTHUS, Oerst. in Kjoeb. Vidensk. Meddel. (1875) 196.
acuminata, Poepp. & Endl. Nov. Gen. et Sp. ii. 37 = divergens.
acuminata, *Ruiz & Pav. Syst.* 293.—Am. austr.
angustifolia, Ruiz & Pav. l. c. 292 = Cyclanthus Plumerii.
atrovirens, *H. Wendl. Ind. Palm.* (1854) 67.—N. Granat.
brachypus, *Drude, in Mart. Fl. Bras.* iii. II. 237.—Bras.
chelidonura, *Drude, l. c.* 238.—Bras.
disticha, *Poit. in Rev. Hortic.* Sér. III. i. (1847) 86.—Guiana.
divergens, *Drude, in Mart. Fl. Bras.* iii. II. 241.—Bras.
Drudei, *Mast. in Gard. Chron.* (1877) II. 714.—Bras.
ensiformis, *Hook. f. Bot. Mag.* t. 6418.—Costa Rica.
funifera, *Kunth, Enum. Pl.* iii. 106.—Guiana.
Gardneri, *Hook. Journ. Bot.* ii. (1840) 29. t. 3, 4.—Bras.
heterophylla, *Mart. ex Drude, in Mart. Fl. Bras.* iii. II. 242.—Bras.

CARLUDOVICA:—
humilis, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 37.—Peruv.
imperialis, *Linden & André, Illustr. Hortic.* xxi. (1874) t. 165.—Ecuador.
incisa, *H. Wendl. Ind. Palm.* (1854) 67.—Am. centr.
insignis, *Duchass. ex Griseb. Fl. Brit. W. Ind.* 513.—Ind. occ.
Kegeliana, *Lem. Illustr. Hortic.* ii. (1855) *Misc.* 23.—Guiana.
latifolia, *Ruiz & Pav. Syst.* 292.—Ind. occ.
latifrons, *Drude, in Mart. Fl. Bras.* iii. II. 237.—Bras.
longicrura, *Drude, l. c.* 240.—Bras.
macropoda, *Klotzsch, in Linnaea*, xx. (1847) 469.—N. Granat.
microphylla, *Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 197.—Costa Rica.
Moritziana, *Klotzsch, in Linnaea*, xx. (1847) 468.—N. Granat.
Oerstedii, *Hemsl. Biol. Centr. Am. Bot.* iii. 416.—Costa Rica.
palmata, Griseb. in Goett. Abh. vii. (1857) 259 = insignis.
palmata, *Ruiz & Pav. Syst.* 291.—Am. austr.
plicata, *Klotzsch, in Linnaea*, xx. (1847) 468.—N. Granat.
plicata, Linden, ex H. Wendl. Ind. Palm. (1854) 67 = atrovirens.
Plumerii, *Kunth, Enum. Pl.* iii. 106.—Ins. Martinic.
Plumieri, Griseb. Fl. Brit. W. Ind. 513 = gracilis.
purpurascens, J. Linden, Illustr. Hortic. xxviii. (1881) 64.—N. Granat.
rupestris, *Klotzsch, in Linnaea*, xx. (1847) 469.—N. Granat.
sarmentosa, *Sagot, ex Drude, in Mart. Fl. Bras.* iii. II. 239.—Guiana.
Sartoni, *Teijsm. & Binn. Cat. Hort. Bog.* 381, nomen.—Hab.?
schizophylla, *Drude, in Mart. Fl. Bras.* iii. II. 235.—Bras.
speciosa, Linden.—Cf. Gard. Chron. (1877) II. 753 = Drudei.
subacaulis, *Kunth, Enum. Pl.* iii. 107.—Guiana.
tetragona, *H. B. & K. Nov. Gen. et Sp.* i. 79.—N. Granat.
tetragonopus, *Mart. ex Drude, in Mart. Fl. Bras.* iii. II. 238.—Bras.
Trailiana, *Drude, l. c.* 241.—Bras.
trigona, *Ruiz & Pav. Syst.* 293.—Am. austr.
utilis, *Benth. & Hook. f. ex Hemsl. Biol. Am. Centr. Bot.* iii. 416.—Costa Rica.
Wallisi, *Regel, Gartenfl.* (1879) 325.—N. Granat.

CARMELITA, C. Gay, ex DC. Prod. vii. 14 (1838) = **Chaetanthera**, Ruiz & Pav. (Compos.).
formosa, C. Gay, l. c. 15 = Chaet. villosa.

CARMENTA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 (Quid?).

CARMICHAELA, Reichb. Nom. 147 (1841) = seq.

CARMICHAELIA, R. Br. in Bot. Reg. xi. t. 912 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 502.
australis, *R. Br. l. c.*—N. Zel.
compacta, *Petrie, in Trans. N. Z. Inst.* xvii. (1885) 272.—N. Zel.
corrugata, *Colenso, l. c.* xv. (1883) 320.—N. Zel.
crassicaulis, *Hook. f. Handb. New Zeal. Fl.* 48.—N. Zel.
Cunninghamii, *Raoul, Choix*, 29.—N. Zel.
Enysii, *T. Kirk, in Trans. N. Z. Inst.* xvi. (1884) 379.—N. Zel.
Exsul, *F. Muell. Fragm.* vii. 126.—Ins. Dom. Howe.
flagelliformis, *Colenso, in Hook. f. Fl. N. Zel.* i. 51.—N. Zel.
gracilis, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 336.—N. Zel.
grandiflora, *Hook. f. Handb. New Zeal. Fl.* 49.—N. Zel.
juncea, *Colenso, ex Hook. f. Fl. N. Zel.* i. 51.—N. Zel.
Kirkii, *Hook. f. Ic. Pl.* t. 1332.—N. Zel.
Munroi, *Hook. f. Handb. New Zeal. Fl.* 49.—N. Zel.
nana, *Colenso, ex Hook. f. l. c.*—N. Zel.
odorata, *Colenso, ex Hook. f. Fl. N. Zel.* i. 50.—N. Zel.
pilosa, *Colenso, ex Hook. f. l. c.*—N. Zel.
stricta, *Lehm. in Ind. Hort. Bot. Hamb.* (1852) 8.—*Cf. Hamb. Gartenz.* viii. (1852) 372.—Hab.?

CARMICHAELIA:—
uniflora, *J. Buch. in Trans. N. Z. Inst.* xvi. (1884) 394.—N. Zel.
uniflora, *T. Kirk, in Gard. Chron.* (1884) I. 512; *Trans. N. Z. Inst.* xvi. (1884) 379.—N. Zel.
Williamsi, *T. Kirk, in Trans. N. Z. Inst.* xii. (1880) 394.—N. Zel.

CARMINATIA, Moç. ex DC. Prod. vii. 267 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 243.
tenuiflora, *DC. l. c.; et in Deless. Ic. Sel.* iv. t. 98.—Mexic.
tenuifolia, Steud. Nom. ed. II. i. 299, sphalm. = praec.

CARMONA, Cav. Ic. v. 22. t. 438 (1799) = **Ehretia**, Linn. (Borag.).
heterophylla, Cav. l. c. 23 = E. buxifolia.
lycioides, G. Don, Gen. Syst. iv. 391 = E. lycioides.
microphylla, G. Don, l. c. = E. buxifolia.
viminea, G. Don, l. c. = Rhabdia lycioides.

CARMONEA, Pers. Syn. i. 290 (1805) = praec.

CARMOREA, Steud. Nom. ed. II. i. 299 (1840) sphalm. = praec.

CARNARVONIA, F. Muell. Fragm. vi. 80. t. 55, 56 (1867). *PROTEACEAE*, Benth. & Hook. f. iii. 182.
araliaefolia, *F. Muell. l. c.* 81.—Austral.

CAROLI-GMELINA, Gaertn. Mey. & Scherb. Fl. Wett. ii. 419 (1800) = **Nasturtium**, R. Br. (Crucif.).
lancifolia, Gaertn. Mey. & Scherb. l. c. 468 = N. amphibium.
palustris, Gaertn. Mey. & Scherb. l. c. 470 = N. amphibium.
sylvestris, Gaertn. Mey. & Scherb. l. c. 470 = N. sylvestre.

CAROLINEA, Linn. f. Suppl. 51 (1781) = **Pachira**, Aubl. (Malvac.).
affinis, Mart. Nov. Gen. et Sp. i. 85 = P. affinis.
alba, Lodd. Bot. Cab. t. 752 = P. alba.
arenaria, G. Don, Gen. Syst. i. 510 = P. arenaria.
campestris, Mart. Nov. Gen. et Sp. i. 86 = P. campestris.
emarginata, D. Dietr. Syn. Pl. iv. 861 = P. emarginata.
fastuosa, Sesse, ex DC. Prod. i. 478 = P. fastuosa.
grandiflora, Spach, Hist. Veg. Phan. iii. 426; xiv. 206 = P. grandiflora.
insignis, Sw. Prod. Veg. Ind. Occ. 101 = P. insignis.
longiflora, Mart. Nov. Gen. et Sp. i. 86 = P. longiflora.
macrantha, G. Don, Gen. Syst. i. 510 = P. macrantha.
macrocarpa, Cham. & Schlecht. in Linnaea, vi. (1831) 423 = P. macrocarpa.
marginata, G. Don, Gen. Syst. i. 511 = P. marginata.
minor, Sims, Bot. Mag. t. 1412 = P. minor.
pompalis, Moç. & Sesse, ex DC. Prod. i. 478 = P. minor.
Princeps, Linn. f. Suppl. 314 = P. aquatica.
robusta, Lodd. ex Steud. Nom. ed. II. i. 299.—Am. austr.
tomentosa, Mart. Nov. Gen. et Sp. i. 84 = P. tomentosa.
villosa, Mart. ex Steud. Nom. ed. II. i. 300 = P. tomentosa.

CAROMBA, Steud. Nom. ed. II. i. 300 (1840) sphalm. = **Cabomba**, Aubl. (Nymph.).

CAROPYXIS, Benth. & Hook. f. Gen. i. 688 (1865) . sphalm. = CALOPYXIS, Tul. = **Combretum**, Linn.

CAROSELINUM, Griseb. Spicil. Fl. Rumel. i. 374 (1843) = **Peucedanum**, Linn. (Umbellif.).
Chabraei, Griseb. l. c. = P. carvifolium.
distans, Griseb. l. c. = Johrenia graeca.

CAROTA, Rupr. Sert. Tiansch. 49: in Mém. Acad. Pétersb. Sér. VII. xiv. (1869) = **Daucus**, Tourn. (Umbellif.).
sativa, Rupr. Fl. Ingric. 468 = D. Carota.
sylvestris, Rupr. Sert. Tiansch. 49 = D. Carota.

CAROXYLON, Thunb. Nov. Gen. ii. 37 (1782) = **Salsola**, Linn. (Chenopod.).
acutifolium, Moq. in DC. Prod. xiii. II. 173 = S. Auricula.

CAROXYLON :—
articulatum, Moq. in DC. Prod. xiii. II. 175 = Haloxylon articulatum.
Auricula, Moq. l. c. 172 = S. Auricula.
Bottae, Moq. in DC. l. c. 178.—Afr. austr.
brevifolium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 121 = S. aphylla.
divaricatum, Moq. in DC. Prod. xiii. II. 177 = S. longifolia.
foetidum, Moq. l. c. 178 = S. foetida.
glaucum, Moq. l. c. 173 = S. glauca.
Griffithii, Moq. l. c. 175 = Haloxylon Griffithii.
hispidulum, Bunge, Lehm. Reliq. Bot. 293 = S. hispidula.
imbricatum, Moq. in DC. Prod. xiii. II. 177 = S. gemmascens.
indicum, Wight, Ic. t. 1794 = Haloxylon recurvum.
lancifolium, Boiss. Diagn. Ser. I. xii. 98 = S. lancifolia.
nodulosum, Moq. in DC. Prod. xiii. II. 177 = S. gemmascens.
recurvum, Moq. l. c. 175 = Haloxylon recurvum.
salicornicum, Moq. l. c. 174 = Haloxylon salicornicum.
Salsola, Thunb. Diss. Med. Afric. 8 = S. aphylla.
subaphyllum, Moq. in DC. Prod. xiii. II. 173 = S. subaphylla.
tamariscifolium, Moq. l. c. 174 = S. genistoides.
tamariscinum, Moq. l. c. = S. tamariscina.
tetragonum, Moq. l. c. 176 = S. tetragona.
tuberculatum, Moq. l. c. 178 = S. tuberculata.
verrucosum, Moq. l. c. = S. tuberculata.
Zeyheri, Moq. l. c. 176.—Afr. austr.

CARPACOCE, Sond. in Harv. & Sond. Fl. Cap. iii. 32 (1864-65). *RUBIACEAE*, Benth. & Hook. f. ii. 141.
LAGOTIS, E. Mey. in Drège, Zwei Pfl. Docum. 197 (1843).
scabra, *Sond. l. c.* 33.—Afr. austr.
Spermacoce, *Sond. l. c.*—Afr. austr.

CARPENTARIA, Becc. in Ann. Jard. Buitenz. ii. (1885) 128. *PALMAE*.
acuminata, *Becc. l. c.*—Austral.

CARPENTERIA, Torr. in Smithson. Contrib. vi. 12. t. 7 (1854). *SAXIFRAGEAE*, Benth. & Hook. f. i. 643.
californica, *Torr. l. c.*—Calif.

CARPENTIERA, Steud. Nom. ed. II. i. 344 in syn. (1840) sphalm. = **Charpentiera**, Gaudich. (Amarant.).

CARPESIUM, Linn. in Act. Soc. Upsal. (1741) 80. *COMPOSITAE*, Benth. & Hook. f. ii. 336.
CONYZOIDES, Tourn. ex DC. Prod. vi. 281 (1837).
abrotanoides. *Linn. Sp. Pl.* 860.—Europ.; As. temp.
cernuum, *Linn. l. c.* 859.—Europ.; As. temp.
ciliatum, Wall. Cat. n. 3214 = cernuum.
divaricatum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 187.—Japon.
glossophyllum, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 475.—Japon.
macrocephalum, *Franch. & Sav. Enum. Pl. Jap.* ii. 408.—Japon.
nepalense, Less. in Linnaea, vi. (1831) 234 = cernuum.
pedunculosum, Wall. Cat. n. 3200 = cernuum.
pubescens, Wall. l. c. n. 3199 = cernuum.
racemosum, Wall. l. c. n. 3201 = abrotanoides.
rosulatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 179.—Japon.
scandens, Schult. f. ex Miq. l. c. 172 = Rhynchospermum verticillatum.
Thunbergianum, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1845) 187.—Japon.
trachelifolium, *Less. in Linnaea*, vi. (1831) 233.—Reg. Himal.
triste, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 479.—Japon.
Wulfenianum, Schreb. ex DC. Prod. vi. 282 = abrotanoides.

CARPEZIUM, Gouan, Fl. Monsp. 465 (1765) = praec.

CARPHA, Banks & Soland. ex R. Br. Prod. 230 (1810), pro parte. *CYPERACEAE*, Benth. & Hook. f. iii. 1061.
alpina, *R. Br. Prod.* 230.—Austral.; N. Zel.; Patag.
andina, Phil. in Anal. Univ. Chil. (1873) 555 = paniculata.
arundinacea, Brongn. in Duperr. Voy. Bot. 169. t. 30 = Schoenus arundinaceus.
Aubertii, Nees, in Linnaea, ix. (1834) 300 = Cyclocampe Aubertei.
avenacea, R. Br. Prod. 230 = Cyathochaete avenacea.
capensis, Martens, ex Steud. Nom. ed. II. i. 300 = Ecklonea capensis.
capitellata, *Boeck. in Linnaea*, xxxviii. (1874) 266.—Afr. austr.
clandestina, R. Br. Prod. 231 = Cyathochacte clandestina.
deusta, R. Br. l. c. 230 = Mesomolaena deusta.
diandra, R. Br. l. c. 231 = Cyathochacte diandra.
elongata, Boeck. l. c. xxxviii. (1874) 273 = Asterochaeta elongata.
glomerata, Nees, l. c. vii. (1832) 529 = Asterochaete glomcrata.
hexandra, Nees, l. c. ix. (1834) 300; x. (1835-36) 193 = Macrochactium Dregei.
junciformis, Boeck. in Linnaea, xxxviii. (1874) 267 = Cladium undulatum.
nivicola, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 111 = alpina.
paniculata, *Phil. in Linnaea*, xxix. (1857-58) 80.—Chili.
schoenoides, Banks & Soland. ex Hook. f. Fl. Antarct. 362. t. 148 = alpina.
Schweinfurthiana, *Boeck. in Flora*, lxii. (1879) 569.—Afr. trop.
Urvilleana, Gaudich. ex Nees, in Linnaea, ix. (1834) 300 = Asterochaete arundinacea.

CARPHALEA, Juss. Gen. 198 (1789). *RUBIACEAE*, Benth. & Hook. f. ii. 52.
angulata, *Baill. in Bull. Soc. Linn. Par.* i. (1878) 188.—Madag.
corymbosa, *Willd. Sp. Pl.* i. 583.—Madag.
Kirondron, *Baill. in Bull. Soc. Linn. Par.* i. (1878) 188.—Madag.
madagascariensis, Lam. Illustr. i. 258. t. 59 = corymbosa.
Pervilleana, *Baill. in Bull. Soc. Linn. Par.* i. (1878) 188.—Madag.
pubiflora, Moç. & Sesse, ex DC. Prod. iv. 365 = Bouvardia quaternifolia.

CARPHEPHORUS, Cass. in Bull. Soc. Philom. (1816) 198. *COMPOSITAE*, Benth. & Hook. f. ii. 249.
atriplicifolius, *A. Gray, in Proc. Am. Acad.* v. (1861) 159.—Calif.
baicalensis, DC. Prod. v. 132 = Saussurea pycnocephala.
bellidifolius, *Torr. & Gray, Fl. N. Am.* ii. 66.—Am. bor.
coridifolius, DC. Prod. vii. 267 = Brickellia pinifolia.
corymbosus, *Torr. & Gray, Fl. N. Am.* ii. 67.—Am. bor.
junceus, *Benth. Bot. Voy. Sulph.* 21.—Calif.
junceus, Durand, in Pacif. Rail. Rep. v. 8 = Linosyris squamata.
Pseudo-liatris, *Cass. in Bull. Soc. Philom.* (1816) 149.—Am. bor.
revolutifolius, *DC. Prod.* v. 133.—Mexic.
tomentosus, *Torr. & Gray, Fl. N. Am.* ii. 66.—Am. bor.
triangularis, *A. Gray, Pl. Wright.* i. 86.—Mexic.

CARPHOBOLUS, Schott, in Spreng. Syst. iv. Cur. Post. 409 (1827) = **Piptocarpha**, R. Br. (Compos.).
asterotrichius, Sch. Bip. in Pollichia, xx.-xxi. (1863) 426 = P. Vauthieriana.
axillaris, Sch. Bip. l. c. 417 = P. axillaris.
Blanchetianus, Sch. Bip. l. c. 423 = P. Lundiana.
cinereus, Sch. Bip. l. c. 419 = P. cinerea.
cuneifolius, Sch. Bip. l. c. 413 = P. oblonga.
latifolius, Sch. Bip. l. c. 426 = P. opaca.
Lechleri, Sch. Bip. l. c. 428 = P. Lechleri.
lepidotus, Sch. Bip. l. c. 414 = P. oblonga.
leprosus, Sch. Bip. l. c. 419 = P. leprosa.
lucidus, Sch. Bip. l. c. 413 = P. lucida.

CARPHOBOLUS :—
Lundianus, Sch. Bip. in Pollichia, xx.-xxi. (1863) 421 = Piptocarpha Lundiana.
macropodus, Sch. Bip. l. c. 418 = P. macropoda.
neurophyllus, Sch. Bip. l. c. 414 = P. pyrifolia.
notatus, Sch. Bip. l. c. 420 = P. notata.
oblongus, Sch. Bip. l. c. 416 = P. oblonga.
opacus, Sch. Bip. l. c. 427 = P. polycephala.
oxyphyllus, Sch. Bip. l. c. 415 = P. oxyphylla.
pellucidus, Sch. Bip. l. c. 420 = P. pellucida.
Poeppigianus, Sch. Bip. l. c. 422 = P. Poeppigiana.
pyrifolius, Sch. Bip. l. c. 415 = P. pyrifolia.
ramiflorus, Sch. Bip. l. c. 429 = P. ramiflora.
Regnellii, Sch. Bip. l. c. 417 = P. axillaris.
Riedelii, Sch. Bip. l. c. 425 = P. Riedelii.
rotundifolius, Sch. Bip. l. c. 424 = P. rotundifolia.
Sellovii, Sch. Bip. l. c. 421 = P. Sellowii.
semiserrulatus, Sch. Bip. l. c. 416 = P. axillaris.
sessiliflorus, Schott, in Spreng. Syst. iv. Cur. Post. 409 = P. lucida.
tereticaulis, Sch. Bip. in Pollichia, xx.-xxi. (1863) 422 = P. tereticaulis.
umbellulatus, Schott, in Spreng. Syst. iv. Cur. Post. 409 = P. umbellulata.

CARPHOCHAETE, A. Gray, Pl. Fendl. 65; in Mem. Am. Acad. N. S. iv. (1849). *COMPOSITAE*, Benth. & Hook. f. ii. 247.
Bigelovii, *A. Gray, Pl. Wright.* i. 89.—N. Mexic.
Grahami, *A. Gray, l. c.*—Mexic.
Wislizeni, *A. Gray, Pl. Fendl.* 65.—Mexic.

CARPHOLOMA, D. Don, in Mem. Wern. Soc. v. (1826) 555 = **Lachnospermum**, Willd. (Compos.).
rigidum, D. Don, l. c. = L. ericaefolium.

CARPHOPAPPUS, Sch. Bip. in Walp. Rep. ii. 954, 973 (1843) = **Iphiona**, Cass. (Compos.).
baccharidifolius, Sch. Bip. l. c. 974 = I. baccharidifolia.

CARPHOSTEPHIUM, Cass. in Dict. Sc. Nat. xliv. 62 (1826) = **Tridax**, Linn. (Compos.).
trifidum, Cass. l. c. = T. trifida.

CARPIDOPTERIX, Karst. Fl. Columb. ii. 45. t. 123 (1862?) = **Thouinia**, Poit. (Sapindac.).
macroptera, Karst. l. c. = T. macroptera.

CARPINUS, Linn. Gen. ed. I. 292 (1737). *CUPULIFERAE*, Benth. & Hook. f. iii. 405.
CARPINUM, Rafin. in Am. Monthly Mag. (1818) 268.
DISTEGOCARPUS, Sieb. & Zucc. Fam. Nat. Fl. Jap. ii. 102. t. 3 (1846).
americana, *Michx. Fl. Bor. Am.* ii. 201.—Am. bor.
Betulus, *Linn. Sp. Pl.* 998.—Europ.; Reg. Cauc.: Persia.
caroliniana, Walt. Fl. Carol. 236 = americana.
Carpinizza, Kit. in Host, Fl. Austr. ii. 626 = C. Betulus.
compressus, Gilib. Exercit. ii. 399 = C. Betulus.
cordata, *Blume, Mus. Bot. Lugd. Bat.* i. 309.—Japon.
duinensis, Scop. Fl. Carn. ed. II. ii. 243. t. 60 = orientalis.
edentulus, *Kit. in Rochel, Pl. Banat. Rar.* nomen.—Banat.
crosa, *Blume, Mus. Bot. Lugd. Bat.* i. 308.—Japon.
faginea, *Lindl. in Wall. Pl. As. Rar.* ii. 5.—Reg. Himal.
intermedia, Wierzb. ex Reichb. Ic. Fl. Germ. xii. 4. t. 633 = C. Betulus.
italica, Scop. ex Steud. Nom. ed. II. i. 300 = Ostrya carpinifolia.
japonica, *Blume, Mus. Bot. Lugd. Bat.* i. 308.—Japon.
laxiflora, *Blume, l. c.* 309.—Japon.
minor, *Pall. Reise*, ii. 95.—Tauria.
nervata, Dulac, Fl. Hautes-Pyr. 141 = C. Betulus.
nigra, Moench, Verz. Ausl. Baeume, 19 = orientalis.
orientalis, *Mill. Gard. Dict.* ed. VIII. n. 3.—Europ. austr.
Ostrya, Linn. Sp. Pl. 998 = Ostrya virginica.
ostryoides, Rafin. in Med. Repos. N. York, Ser. III. ii. (1811) 333 americana?
quercifolia, Desf. ex Steud. Nom. ed. II. i. 300 = C. Betulus.

CARPINUS:—
sepium, Lam. Fl. Fr. ii. 212 = C. Betulus.
triflora, Mocnch, Meth. 694 = Ostrya virginica.
Tschonoskii, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 534.—Japon.
Turczaninovii, *Hance, in Journ. Linn. Soc.* x. (1869) 203.—China.
ulmifolia, Salisb. Prod. 392 = C. Betulus.
ulmoides, S. F. Gray, Nat. Arr. Brit. Pl. ii. 245 = C. Betulus.
viminea, *Wall. ex Lindl. in Wall. Pl. As. Rar.* ii. 4. t. 106.—Reg. Himal.
virginiana, Michx. Hist. Arb. For. iii. 57. t. 8 = americana.
virginiana, Mill. Gard. Dict. ed. VIII. n. 4; Lam. Encyc. i. 708 = Ostrya virginiana.
vulgaris, Mill. l. c. n. 1 = C. Betulus.
yedoënsis, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 535.—Japon.

CARPIPHEA, Rafin. Sylva Tellur. 39 (1838) = **Cordia**, Tourn. (Boragin.).
dentata, Rafin. l. c. = Cordia dentata.

CARPOCALYMNA, Zipp. in Alg. Konst. en Letterbode, i. (1829) 297 = **Epithema**, Blume (Gesnerac.).
monophylla, Zipp. ex Span. in Linnaea, xv. (1841) 331 = E. difformis.

CARPOCERAS, Link, Handb. ii. 289 (1831) = **Thlaspi**, Tourn. (Crucif.).
cappadocicum, Boiss. Diagn. Ser. II. vi. 19.—Cappadoc.
cilicicum, Boiss. & Kurz, ex Boiss. Fl. Orient. i. 332.—Cilicia.
Griffithianum, Boiss. Diagn. Ser. II. i. 40 = T. alpestre.
hastulatum, Boiss. l. c. Ser. I. viii. 38 = T. hastulatum
oxyceras, Boiss. l. c. 37 = T. oxyceras.
sibiricum, Link, Handb. ii. 289 = T. ceratocarpon.
stenocarpum, Boiss. Diagn. Ser. I. viii. 38 = T. stenocarpum.

CARPOCERAS, A. Rich. Elem. Bot. (1846) 706 = **Martynia**, Linn. (Pedalin.).
longiflora, A. Rich. l. c. = M. diandra.

CARPODETES, Herb. App. 41 (1821) = **Stenomesson**, Herb. (Amaryllid.).
recurvata, Herb. l. c. = S. recurvatum.

CARPODETUS, Forst. Char. Gen. 33. t. 17 (1776). *SAXIFRAGEAE*, Benth. & Hook. f. i. 646.
dentatus, Poir. Encyc. Suppl. ii. 120 = serratus.
Forsteri, Roem. & Schult. Syst. v. 604 = serratus.
serratus, *Forst. Char. Gen.* 34. t. 17 A.—N. Zel.

CARPODINUS, R. Br. ex Sabine, in Trans. Hort. Soc. v. (1823) 455. *APOCYNACEAE*, Benth. & Hook. f ii. 693.
acida, *Sabine, l. c.* 456.—Afr. trop.
dulcis, *Sabine, l. c.* 455.—Afr. trop.

CARPODIPTERA, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 163. *TILIACEAE*, Benth. & Hook. f. i. 232.
africana, *Mast. in Oliver, Fl. Trop. Afr.* i. 241.—Afr. trop.
Boivini, *Baill. Adansonia*, x. (1872) 180.—Ins. Comor.
cubensis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 164.—Cuba.
Schomburgkii, *Baill. Adansonia*, x. (1872) 181.—Guiana.

CARPODON, Spreng. Syst. ii. 634 (1825) = seq.

CARPODONTOS, Labill. Voy. ii. 16. t. 18 (1798) = **Eucryphia**, Cav. (Rosac.).
lucida, Labill. l. c. = E. Billardieri.

CARPOLOBIA, G. Don, Gen. Syst. i. 370 (1831), partim. *POLYGALEAE*, Benth. & Hook. f. i. 139, 553.
Afzeliana, *Oliver, Fl. Trop. Afr.* i. 136.—Afr. trop.
alba, *G. Don, Gen. Syst.* i. 370.—Afr. trop.
dubia, G. Don, l. c. = Baphia polygalacea.
lutea, *G. Don, l. c.*—Afr. trop.
versicolor, G. Don, l. c. = Baphia nitida.

CARPOLIZA, Steud. Nom. ed. II. i. 300 (1840) = seq.

CARPOLYZA, Salisb. Parad. Lond. t. 63 (1807). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 721.
HESSEA, Berg, ex Schlecht. in Linnaea, i. (1826) 252, non Herb.
spiralis, *Salisb. Parad. Lond.* t. 63.—Afr. austr.
stellaris, Heynh. Nom. ii. 122 = Hessea stellaris, *Herb.*

CARPONEMA, Eckl. & Zeyh. Enum. 8 (1835). *CRUCIFERAE*, Benth. & Hook. f. i. 102.
aggregata, Eckl. & Zeyh. l. c. = Heliophila gracilis.
filiformis, *Eckl. & Zeyh. l. c.*—Afr. austr.

CARPOPHILLUS, Neck. Elem. ii. 84 (1790) = **Pereskia**, Plum.? (Cacteae).

CARPOPHORA, Klotzsch, in Bot. Ergeb. Waldem. Reise, 139. t. 32 (1862) = **Silene**, Linn. (Caryophyll.).
Hoffmeisterii, Klotzsch, l. c. = S. Griffithii.

CARPOPHYLLUM, Miq. Fl. Ind. Bat. Suppl. i. 401 (1860) = **Sterculia**, Linn.
macropodum, Miq. l. c.—Sumatra.

CARPOPHYLLUM, Neck. Elem. iii. 402, 411 (1790) = CARPOPHILLUS, Neck.

CARPOPODIUM, Eckl. & Zeyh. Enum. i. 13 (1835) = **Heliophila**, Linn. (Crucif.).
carnosum, Eckl. & Zeyh. l. c. = H. carnosa.
cleomoides, Eckl. & Zeyh. l. c. = H. glauca.
Thunbergii, Eckl. & Zeyh. l. c. = H. Thunbergii.

CARPOPOGON, Roxb. Hort. Beng. 54 (1814); Fl. Ind. iii. 283 (1832) = **Mucuna**, Adans. (Legumin.).
anguineum, Roxb. ex Baker, in Hook. f. Fl. Brit. Ind. ii. 185 = M. monosperma.
atropurpureum, Roxb. Hort. Beng. 54; Fl. Ind. iii. 287 = M. atropurpurea.
bracteatum, Roxb. Hort. Beng. 54 = M. bracteata.
capitatum, Roxb. l. c. [98]; Fl. Ind. iii. 284 = M. capitata.
giganteum, Roxb. ll. cc. 54, 287 = M. gigantea.
imbricatum, Roxb. Hort. Beng. 54 = M. imbricata.
monospermum, Roxb. l. c.; Fl. Ind. iii. 283 = M. monosperma.
niveum, Roxb. ll. cc. 54, 285 = M. nivea.
pruriens, Roxb. ll. cc. 54, 283 = M. pruriens.
volubilis, Roxb. ex Steud. Nom. ed. II. i. 300 = M. volubilis.

CARPOTHALIS, E. Mey. in Drège, Zwei Pfl. Docum. 170 (1843) = **Kraussia**, Harv. (Rubiac.).
lanceolata, E. Mey. l. c. = K. lanceolata.

CARPOTRICHE, Reichb. Nom. 189 (1841) = scq.

CARPOTROCHE, Endl. Gen. 918 (1839). *BIXINEAE*, Benth. & Hook. f. i. 125.
brasiliensis, *Endl. l. c.*—Bras.
denticulata, *Benth. in Hook. Journ. Bot.* iv. (1842) 115.—Guiana.
grandiflora, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiii. 1. 437.—Bras.
laxiflora, *Benth. in Hook. Journ. Bot.* iv. (1842) 114.—Guiana.
longifolia, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 82.—Bras.; Peruv.
paludosa, *Benth. in Hook. Journ. Bot.* iv. (1842) 114.—Guiana.

CARPOXIS, Rafin. New Fl. Am. iii. 89 (1836) = **Forestiera**, Poir. (Oleac.).
inermis, Rafin. l. c. 92.—Am. bor.
nitida, Rafin. l. c. 91 = F. acuminata.
prinoides, Rafin. l. c. 92 = F. prinoides.
retusa, Rafin. l. c. = F. retusa.
spinosa, Rafin. l. c.—Am. bor.

CARPOXYLON, H. Wendl. & Drude, in Kerch. Palm. 238 (1878), nomen. *PALMAE*.
macrospermum, *H. Wendl. & Drude, l. c.*—Hab.?

CARPUNYA, Presl, Epim. Bot. 228 (1849) = **Piper**, Linn.
Lessertiana, Presl, l. c. 229 = P. Carpunya.
oblongata, Presl, l. c. = P. oblongatum.
peruviana, Presl, l. c. = P. Carpunya.

CARPUPICA, Rafin. Sylva Tellur. 85 (1838) = **Piper**, Linn.
odorata, Rafin. l. c. = P. Carpunya.

CARRADORIA, A. DC. Prod. xii. 610 (1848) = **Globularia**, Linn. (Selagin.).
incanescens, A. DC. l. c. = G. incanescens.

CARREGNOA, Boiss. Voy. Bot. Espagne, ii. 605 (1842) = **Tapeinanthus**, Herb. (Amaryllid.).
dubia, Perez-Lara, in Anal. Soc. Esp. Hist. Nat. xi. (1882) 399 = Narcissus serotinus?
humilis, J. Gay, in Ann. Sc. Nat. Sér. IV. x. (1858) 98 = T. humilis.
lutea, Boiss. Voy. Bot. Espagne, ii. 605 = T. humilis.

CARRIA, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 6 = **Gordonia**, Ellis (Ternstroem.).
speciosa, Gardn. l. c. 7 = G. speciosa.

CARRICHTERA, Adans. Fam. ii. 421 (1763); DC. Syst. Veg. ii. 641. *CRUCIFERAE*, Benth. & Hook. f. i. 86.
Vella, *DC. l. c.* 642.—Hispan.

CARROA, Presl, Symb. Bot. ii. 25 (1858) = **Dalea**, Linn. (Legumin.).
diffusa, Presl, l. c. = D. diffusa.
grandulosa, Presl, l. c. = D. diffusa.

CARRONIA, F. Muell. Fragm. ix. 171 (1875). *MENISPERMACEAE*.
multisepalea, *F. Muell. l. c.*—Austral.

CARRUTHERSIA, Seem. Fl. Vit. 155. t. 30 (1865–73). *APOCYNACEAE*, Benth. & Hook. f. ii. 718.
pilosa, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 132.—Ins. Philipp.
scandens, *Seem. Fl. Vit.* 156. t. 30.—Ins. Fiji.

CARTALINIA, Szov. ex Kunth, Enum. Pl. v. 119 (1850) = **Paris**, Rupp. (Liliac.).

CARTERETIA, A. Rich. Sert. Astrol. 10. t. 4 (1834) = **Doritis**, Lindl. (Orchid.).
paniculata, A. Rich. l. c. = D. paniculata.

CARTESIA, Cass. in Bull. Soc. Philom. (1816) 198 = **Stokesia**, L'Hérit. (Compos.).
centauroides, Cass. l. c. = S. cyanea.

CARTHAMUS, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 483.
DURANDOA, Pomel, Mat. Fl. Atl. 2 (1860).
HERACANTHA, [Hoffmgg. &] Link, Fl. Port. ii. 205 (1820).
HETERACANTHA, Steud. Nom. ed. II. i. 755 (1840).
HOHENWARTHA, Vest, in Flora, iii. (1820) 1.
KENTROPHYLLUM, Neck. Elem. i. 86 (1790).
ONOBROMA, DC. Prod. vi. 613 (1837).
PHONUS, Hill, Veg. Syst. iv. 5 (1762).
acaulis, Presl, ex Steud. Nom. ed. II. i. 300 = Carduncellus pectinatus.
africanus, Lam. Encyc. i. 640 = Berckheya grandiflora.
albus, *Desf. Tabl.* ed. I. 92.—Oriens.
araneosus, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 363 = Carduncellus araneosus.
arborescens, *Linn. Sp. Pl.* 831.—Hispan.
armenus, Willd. Enum. Hort. Berol. 845 = flavescens.
baeticus, Nym. Consp. 419 = lanatus.
brasiliensis, *Vell. Fl. Flum.* 342; viii. t. 77.—Bras.
caeruleus, Linn. Sp. Pl. 830 = Carduncellus caeruleus.
cancellatus, Lam. Fl. Fr. ii. 10 = Atractylis cancellata.
canescens, Lam. l. c. = Cnicus Acarna.
canescens, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 262.—Aleppo.
Carduncellus, Linn. Sp. Pl. 831 = Carduncellus monspeliensium.
carolinianus, Walt. Fl. Carol. 194 = Elephantopus carolinianus?

CARTHAMUS:—
corymbosus, Linn. Sp. Pl. ed. II. 1164 = Cardopatium corymbosum.
creticus, Linn. l. c. 1163 = lanatus.
creticus, Nym. Consp. 419 = glaucus.
cyaneus, Banks, ex Steud. Nom. ed. II. i. 300 = Stokesia cyanea.
cynaroides, Bieb. Tabl. Casp. 195 = Cousinia cynaroides.
dentatus, *Vahl, Symb. Bot.* i. 69.—As. Min.
dianius, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 363 = Carduncellus caeruleus.
elatus, *Nym. Consp.* 419.—Italia.
ferox, Lam. Fl. Fr. ii. 11 = Cnicus ferox.
flavescens, *Willd. Sp. Pl.* iii. 1706.—As. Min.; Syria.
fluminensis, *Vell. Fl. Flum.* 342; viii. t. 78.—Bras.
glaber, *Burm. f. Fl. Cap. Prod.* 23.—Afr. austr.
glaucus, *Bieb. Fl. Taur. Cauc.* ii. 284.—Aegypt.; As. Min.; Persia.
gummiferus, Lam. Encyc. i. 639 = Atractylis gummifera.
helenioides, *Desf. Fl. Atlant.* ii. 258. t. 230.—Algeria.
hircinus, Lag. ex Spreng. Syst. iii. 391 = arborescens.
hispanicus, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 363 = Carduncellus caeruleus.
humilis, Lam. Encyc. i. 638 = Carduncellus pinnatus.
integrifolius, Hort. ex Webb & Berth. Phyt. Canar. ii. 344 = Carlina salicifolia.
involucratus, Lam. Fl. Fr. ii. 12 = Cnicus spinosissimus.
laevis, Hill, Hort. Kew. 57 = Stokesia cyanea.
lanatus, *Linn. Sp. Pl.* 830.—Europ.; Oriens.
leucocaulos, *Sibth. & Sm. Fl. Graec.* t. 842.—Archip. Graec.
linearifolius, *Molina, Sagg. Chile*, ed. II. 294.—Chili.
maculatus, Lam. Encyc. i. 638 = Silybum Marianum.
magellanicus, *Lam. l. c.* 640.—Reg. Magellan.
mareoticus, *Delile, Fl. Egypte*, 276.—Aegypt.
mitissimus, Linn. Sp. Pl. 831 = Carduncellus mitissimus.
multifidus, Desf. Fl. Atlant. ii. 256 = Carduncellus caeruleus.
nitidus, *Boiss. Fl. Orient.* iii. 708.—Syria.
Oxyacantha, *Bieb. Fl. Taur. Cauc.* ii. 283.—Reg. Caucas.; As. occ.
pectinatus, Desf. Fl. Atlant. ii. 257 = Carduncellus pectinatus.
persicus, Desf. ex Willd. Sp. Pl. iii. 1707 = flavescens.
persicus, Willd. ex Boiss. Fl. Orient. iii. 708 = leucocaulos.
pinnatus, Cutand. Fl. Madrit. 412 = Carduncellus monspeliensium.
pinnatus, Desf. Fl. Atlant. ii. 258 = Carduncellus pinnatus.
polyacantha, *Bieb. Tabl. Casp.* 58.—Transcauc.
pterocaulos, C. A. Mey. Verz. Pfl. Cauc. 67 = Cousinia Hystrix.
rigidus, *Burm. f. Fl. Cap. Prod.* 23.—Afr. austr.
rigidus, Willd. Enum. Hort. Berol. 845 = arborescens.
salicifolius, Linn. f. Suppl. 350 = Carlina salicifolia.
tauricus, Bieb. Fl. Taur. Cauc. ii. 285 = lanatus.
Teneriffae, *J. F. Gmel. ex Steud. Nom.* ed. I. 164.—Ins. Teneriffa.
tinctorius, *Linn. Sp. Pl.* 830.—Geront. cult.
tingitanus, Linn. l. c. ed. II. 1163 = Carduncellus caeruleus.
tomentosus, J. F. Gmel. Syst. 1521. Index (quid?).
turbinatus, *Nym. Consp.* 419.—Sicilia.

CARTODIUM, Soland. ex R. Br. in Trans. Linn. Soc. xii. (1817) 106 = **Craspedia**, Forst. f. (Compos.).

CARTONEMA, R. Br. Prod. 271 (1810). *COMMELINACEAE*, Benth. & Hook. f. iii. 852.
brachyantherum, *Benth. Fl. Austral.* vii. 92.—Austral.
parviflorum, *Hassk. in Flora*, lii. (1869) 365.—Austral.
philydroides, F. Muell. Fragm. i. 62 = spicatum.
spicatum, *R. Br. Prod.* 271.—Austral.
spicatum, F. Muell. Fragm. i. 62 = parviflorum.
tenue, *Caruel, in Nuov. Giorn. Bot.* xi. (1879) 216.—Austral.
trigonospermum, *C. B. Clarke, in DC. Monog. Phan.* iii. 264.—Austral.

CARTREMA, Rafin. Sylva Tellur. 184 (1838) = **Osmanthus**, Lour. (Oleac.).
americana, Rafin. l. c. = O. americanus.

CARUELIA, Parl. Nuov. Gen. e Spec. 21 (1854) = **Ornithogalum**, Tourn. (Liliac.).
arabica, Parl. l. c. 22 = O. arabicum.

CARUM, Rup. ex Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 890.
ANISACTIS, Dulac, Fl. Hautes-Pyr. 347 (1867).
ATAENIA, Hook. & Arn. Bot. Beech. Voy. 349 (1840).
BULBOCASTANUM, Schur, Enum. Pl. Transs. 249 (1866).
BUNIUM, Linn. Gen. ed. I. 72 (1737).
CHAMAREA, Eckl. & Zeyh. Enum. iii. 346 (1837).
EDOSMIA, Nutt. in Torr. & Gray, Fl. N. Am. i. 612 (1840).
ELWENDIA, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 140.
FRORIEPIA, C. Koch, in Linnaea, xvi. (1842) 362.
GEOCARYUM, Coss. & Dur. Pl. Nouv. et Crit. Esp. 112 (1851).
HLADNIKIA, Reichb. Ic. Pl. Crit. ix. 9 (1831); et Fl. Germ. Excurs. 476 (1832).
HUETIA, Boiss. Diagn. Ser. II. ii. 103 (1856).
PETROSELINUM, Hoffm. Umb. i. 78 (1814).
PHYMATIS, E. Mey. in Drège, Zwei Pfl. Docum. 211 (1843).
PTYCHOTIS, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 124.
RIDOLFIA, Moris, Ind. Sem. Hort. Taur. (1841); Fl. Sard. ii. 212 (1842).
SELINOPSIS, Coss. & Dur. ex Munby, Cat. Pl. Alg. ed. II. 13 (1859), nomen.
SYMPODIUM, C. Koch, in Linnnea, xvi. (1842) 356.
TRACHYSPERMUM, Link, Enum. Hort. Berol. i. 267 (1821).
TRACHYSSIADIUM, Eckl. & Zeyh. Enum. 341 (1837).
WYDLERIA, DC. Coll. Mém. v. 36 (1829).
ZIZIA, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 128.
Ajowan, Benth. & Hook. f. Gen. i. 891 = copticum.
allioides, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 294.—Turkest.
alpinum, *Benth. & Hook. f. Gen.* i. 891.—Europ.; Persia.
ammoides, *Benth. & Hook. f. l. c.* 892.—Reg. Mediterr.
anethifolium, *Benth. & Hook. f. l. c.* 891.—Reg. Himal.
Anisum, Baill. Hist. Pl. vii. 119, 178 = Pimpinella Anisum.
apiculatum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 358.—As. centr.
armenum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 137.—Reg. Caucas.
aromaticum, Salisb. Prod. 168 = C. Carvi.
atrosanguineum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 359.—As. centr.
avromanum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 888.—Persia.
aureum, Benth. & Hook. f. Gen. i. 891 = Thaspium aureum.
Bourgaei, *Boiss. Fl. Orient.* ii. 885.—As. Min.
brachycarpum, Boiss. Diagn. Sér. I. x. 23 = ferulaefolium.
Bulbocastanum, *Koch, in Nov. Act. Nat. Cur.* xii. 1. (1824) 121.—Europ.; As. bor.
Bunius, *Linn. Syst. Nat.* ed. XII. 733.—Europ.
bupleuroides, *Schrenk, ex Fisch. & Mey. in Bull. Phys.-Math. Acad. Pétersb.* iii (1845) 305.—Soongar.
buriaticum, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 713.—Sibir. baical.
calcicolum, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 514.—Ins. Socotra.
capense, *Sond. in Harv. & Sond. Fl. Cap.* ii. 538.—Afr. austr.
Capusi, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 293.—Turkestan.
Carvi, *Linn. Sp. Pl.* 263.—Europ.; Oriens; As. bor.
carvifolium, *Benth. & Hook. f. Gen.* i. 892.—Afr. austr.
caucasicum, *Boiss. Fl. Orient.* ii. 880.—Reg. Caucas.
chaerophylloides, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 585.—Turkest.
copticum, *Benth. & Hook. f. Gen.* i. 891.—Europ.; Afr. bor.; As. bor.
cordatum, Benth. & Hook. f. l. c. 891, 913 = Thaspium trifoliatum.

CARUM:—
cornigerum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii 887.—Kurdistan.
creticum, *Benth. & Hook. f. Gen.* i. 891.—Creta.
cylindricum, *Boiss. & Haussk. Diagn.* Ser. I. x. 23.—Persia; Beluchist.
daucoides, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 139.—Macedon.; Lydia.
decussatum, Gilib. Fl. Lituan. ii. 37 = C. Carvi.
dichotomum, *Benth. & Hook. f. Gen.* i. 891.—Reg. Mediterr. occ.
didymum, *Benth. & Hook. f. l. c.*—Afr. austr.
dissectum, Baill. Hist. Pl. vii. 179 = Pimpinella magna.
divaricatum, Koch, Syn. Fl. Germ. 315 = ferulaefolium.
divergens, *Boiss. & Huet, Diagn.* Ser. II. ii. 80.—Armenia.
diversifolium, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 681.—Reg. Himal.
elegans, *Fenzl, Pugill. Pl. Nov. Syr.* 16.—Oriens.
Elvendia, *Boiss. Fl. Orient.* ii. 888.—Persia.
Falcaria, *Lange, in Willk. & Lange, Prod. Fl. Hisp.* iii. 92.—Hispan.
falcarioides, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 96 = elegans.
Falconeri, *C. B. Clarke, in Hook. Fl. Brit. Ind.* ii. 683.—Reg. Himal.
ferulaefolium, *Boiss. Diagn.* Ser. I. x. 22.—Reg. Mediterr. or.
flexuosum, Fries, Summa Veg. Scand. 24, 180 = Conopodium denudatum.
flexuosum, *Nym. Consp.* 307.—Italia.
Gairdneri, *A. Gray, in Proc. Am. Acad.* vii. (1868) 344.—Am. bor. occ.
gracile, *Lindl. in Royle, Illustr. Bot. Himal.* 232.—Ind. or.
graecum, *Boiss. & Heldr. Diagn.* Ser. I. vi. 58.—Graecia; Serbia.
Hallii, *S. Wats. Bibliog. Ind. N. Am. Bot.* 416.—Calif.
Heldreichii, *Boiss. Diagn.* Ser. II. ii. 78.—Graecia.
heterophyllum, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 586.—As. temp.
hispanicum, *Mill. Gard. Dict.* ed. VIII. n. 2.—Hispan.
hispidum, *Benth. & Hook. f. Gen.* i. 891.—Afr. austr.
Huetii, Boiss. Diagn. Ser. II. ii. 81 = Bourgaei.
humile, Boiss. & Bal. ex Boiss. Fl. Orient. ii. 881 = caucasicum.
incrassatum, *Boiss. Voy. Bot. Espagne*, ii. 239, 731.—Hispan.
indicum, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxix. (1866) ii. 72.—Soongaria.
inodorum, *Siev. in Pall. N. Nord. Beitr.* vii. 229.—Rossia.
intermedium, Benth. & Hook. f. Gen. i. 892 = inundatum.
inundatum, *Lespin. in Act. Soc. Bord.* xiv. (1845) 270.—Europ.
involucratum, Baill. Hist. Pl. vii. 179 = Roxburghianum.
Kelloggii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 344.—Calif.
khasianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 682.—Reg. Himal.
latifolium, *Benth. & Hook. f. Gen.* i. 892.—Europ.
leucocoleon, *Boiss. & Huet, Diagn.* Ser. II. ii. 80.—Armenia.
lomatocarum, *Boiss. Fl. Orient.* ii. 879.—Reg. Caucas.
lutescens, Turcz. ex Walp. Rep. v. 866 (= *Cnidium multicaule*).—Sibir.
macrocarpum, *Benth. & Hook. f. Gen.* i. 891.—Graecia.
Macuca, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 34.—Hispan.
magnum, Baill. Hist. Pl. vii. 178 = Pimpinella magna.
mauritanicum, *Boiss. & Reut. Pugill.* (1852) 49.—Afr. bor. occ.
meifolium, Boiss. Fl. Orient. ii. 880 = armenum.
Meum, Stokes, Bot. Mat. Med. ii. 138 = Meum athamanticum.
microcarpum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 137.—Lydia.
multiflorum, *Boiss. Fl. Orient.* ii. 882.—Graecia; As. Min.
neurophyllum, *Franch. & Sav. Enum. Pl. Jap.* i. 180.—Japon.
nigrum, Baill. Hist. Pl. vii. 178 = Pimpinella Saxifraga.

CARUM :—
nivale, *Lange, in Kjoeb. Vidensk. Meddel.* (1865) 34. —Hispan.
Noëanum, Boiss. Diagn. Ser. II. ii. 77 = elegans.
nothum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 681.—Zeylan.?
officinale, S. F. Gray, Nat. Arr. Brit. Pl. ii. 515 = C. Carvi.
oreganum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 368.—Am. bor. occ.
Panatjan, Baill. Hist. Pl. vii. 178 = Pimpinella Panatjan.
Pansil, *Griseb. in Goett. Abh* xxiv. (1879) 146.—Reg. Argent.
paucifolium, DC. Prod. iv. 117 = elegans.
peregrinum, *Linn. Diss. Dem. Pl. in Amoen. Acad.* iii. 406.—Peruv.?
persicum, *Boiss. in Ann. Sc. Nat.* Sér. III. i. (1844) 138.—Persia.
Pestalozzae, *Boiss. Fl. Orient.* ii. 886.—Persia; Beluchist.
Petroselinum, *Benth. & Hook. f. Gen.* i. 891.—Geront. cult.
pimpinelloides, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 514.—Ins. Socotra.
Podagraria, Roth, Enum. Pl. Phan. Germ. i. 946 = Aegopodium Podagraria.
polyphyllum, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 881.—Syria.
purpurascens, Boiss. in Tchihat. Asie Min. Bot. i. 411 = elegans.
rectangulum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 884.—Persia.
Ridolfia, *Benth. & Hook. f. Gen.* i. 891.—Reg. Mediterr.; Oriens.
rigidulum, *Koch, in DC. Prod.* iv. 115.—Italia.
rigidum, Steud. Nom. ed. II. i. 301 = rigidulum.
Roxburghianum, *Benth. & Hook. f. Gen.* i. 891.—Ind. or.; Malaya.
rupestre, *Boiss. & Heldr. Diagn.* Ser. II. ii. 79.—Graecia.
Saxifraga, Baill. Hist. Pl. vii. 178 = Pimpinella Saxifraga.
Saxifraga, Benth. & Hook. f. Gen. i. 892 = C. Bunius.
segetum, *Benth. & Hook. f. l. c.*—Europ.
setaceum, *Schrenk, Enum. Pl. Nov.* 61.—Soongaria.
Sewerzowi, *Regel, in Act. Hort. Petrop.* v. (1877) 587, 588.—Turkestan.
simplex, Steph. ex Willd. Sp. Pl. i. 1470 = Seseli annuum.
Sisarum, Baill. Hist. Pl. vii. 179 = Sium Sisarum.
stictocarpum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 681.—Ind. or.
strictum, *Boiss. Fl. Orient.* ii. 887.—Macedon.
sulcatum, Steud. Nom. ed. I. 164, 49 = Selinum Carvifolia.
sylvestre, *Baill. Hist. Pl.* vii. 179.—Ind. or.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* ii. 371.—Japon.
tenuifolium, *Benth. & Hook. f. Gen.* i. 891.—Hispan.; Marocco.
Thorei, Benth. & Hook. f. l. c. 888, in obs. = inundatum.
trichophyllum, *Schrenk, Enum. Pl. Nov.* 61.—Soongaria.
verticillatum, *Koch, in Nov. Act. Nat. Cur.* xii. 1. (1824) 122.—Europ.

CARUMBIUM, Kurz, For. Fl. Brit. Burma, ii. 411 (1877) = **Sapium**, P. Br. (Euphorb.).
baccatum, Kurz, l. c. 412 = S. baccatum.
insigne, Kurz, l. c.—Burma.
sebiferum, Kurz, l. c. = S. sebiferum.

CARUMBIUM, Reinw. Elench. Sem. Hort. Leid. ex Isis (1823) 319; et in Syll. Ratisb. ii. (1828) 6 = **Homalanthus**, A. Juss. (Euphorb.).
acuminatum, Muell. Arg. in DC. Prod. xv. II. 1144. —Ins. Pacif.
amboinicum, Miq. Fl. Ind. Bat. i. II. 413.—Malaya.
fastuosum, Muell. Arg. in DC. Prod. xv. II. 1144 = H. fastuosus.
giganteum, Muell. Arg. ex Miq. Fl. Ind. Bat. i. II. 687 = H. giganteus.
Moerenhoutianum, Muell. Arg. in DC. Prod. xv. II. 1146.—Ins. Pacif.
nutans, Muell. Arg. l. c. = H. nutans.
pallidum, Muell. Arg. in Linnaea, xxxii. (1863) 85 = H. Leschenaultianus.
pedicellatum, Miq. Fl. Ind. Bat. i. II. 414 = H. pedicellatus.

CARUMBIUM :—
platyneuron, Muell. Arg. in DC. Prod. xv. II. 1145 = Homalanthus Leschenaultianus.
polyandrum, Hook. f. Handb. New Zeal. Fl. 248.—N. Zel.
populifolium, Reinw. in Blume, Cat. Hort. Buitenz. 105 = H. Leschenaultianus.
populneum, Muell. Arg. in DC. Prod. xv. II. 1144 = praec.
Sieberi, Muell. Arg. in Linnaea, xxxii. (1863) 85 = H. Leschenaultianus.
stillingiaefolium, Baill. Adansonia, vi. (1865–66) 325 = H. stillingiaefolius.

CARUNCULARIA, Haw. Syn. Pl. Succ. 333 (1812) = **Stapelia**, Linn. (Asclepiad.).
aperta, Sweet, Hort. Brit. ed. II. 359 = S. aperta.
Jacquini, Sweet, l. c. = S. pedunculata.
Massoni, Sweet, l. c. = S. pedunculata.
pedunculata, Haw. Syn. Pl. Succ. 334 = S. pedunculata.
penduliflora, Sweet, Hort. Brit. ed. II. 359 = S. penduliflora.
serrata, Decne. in DC. Prod. viii. 658, sphalm. = seq.
serrulata, G. Don, Gen. Syst. iv. 122 = Piaranthus serrulatus.
Simsii, Sweet, Hort. Brit. ed. II. 358 = S. pedunculata.

CARVI, [Tourn.] Bernh. Syst. Verz. Erf. 114 (1800) = **Selinum**, Tourn. (Umbellif.).
sulcatum, Bernh. l. c. 169 = S. Carvifolia.

CARVIFOLIA, C. Bauh. ex Vill. Hist. Pl. Dauph. ii. 629 (1787) = **Selinum**, Tourn. (Umbellif.).

CARYA, Nutt, Gen. Am. ii. 220 (1818). *JUGLANDEAE*, Benth. & Hook. f. iii. 398.
HICORIUS, Rafin. Fl. Ludov. 109 (1817).
SCORIA, Rafin. in Med. Repos. N. York, v. (1808) 352, nomen prius.
alba, *Nutt. l. c.* 221.—Am. bor.
amara, *Nutt. l. c.* 222.—Am. bor.
angustifolia, Sweet, Hort. Brit. ed. I. 97 = olivaeformis.
aquatica, *Nutt. Gen. Am.* ii. 222.—Am. bor.
compressa, G. Don, in Loud. Hort. Brit. 384 = alba.
cordiformis, C. Koch, Dendrol. i. 597 = sulcata.
glabra, Sweet, Hort. Brit. ed. I. 97 = porcina.
illinoënsis, C. Koch, Dendrol. i. 593 = olivaeformis.
integrifolia, *Spreng. Syst.* iii. 849.—Louisian.
laciniosa, Loud. Hort. Brit. 384, nomen = sulcata.
mexicana, *Engelm. ex Hemsl. Biol. Centr. Am. Bot.* iii. 162.—Mexic.
microcarpa, *Nutt. Gen. Am.* ii. 221.—Am. bor.
myristicaeformis, *Nutt. l. c.* 222.—Am. bor.
obcordata, Sweet, Hort. Brit. ed. I. 97 = porcina.
olivaeformis, *Nutt. Gen. Am.* ii. 221.—Am. bor.
ovata, C. Koch, Dendrol. i. 598 = alba.
porcina, *Nutt. Gen. Am.* ii. 222.—Am. bor.
pubescens, Sweet, Hort. Brit. ed. I. 97 = sulcata.
sulcata, *Nutt. Gen. Am.* ii. 221.—Am. bor.
tetraptera, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 80 = olivaeformis.
texana, *C. DC. in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 34.—Texas.
tomentosa, *Nutt. Gen. Am.* ii. 221.—Am. bor.

CARYOCAR, Linn. Mant. ii. 247. n. 1314 (1771). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 180.
ACANTACARYX, Arruda, ex Koster, Trav. Braz. 491 (1816).
BAROLLAEA, Neck. Elem. ii. 322 (1790).
PEKEA, Aubl. Pl. Gui. 594 (1775).
RHIZOBOLUS, Gaertn. ex Schreb. Gen. 369 (1789); et Fruct. ii. 93. t. 98 (1791).
SAOUARI, Aubl. Pl. Gui. 599 (1775).
amygdaliferum, *Cav. Ic.* iv. 37. tt. 361, 362.—N. Granat.
amygdaliforme, *Ruiz & Pav. Fl. Per.* v. t. 570.—Peruv.
barbinerve, *Miq. in Linnaea*, xxii. (1849) 802.—Bras.
brasiliense, *St. Hil. Fl. Bras. Mer.* i. 322.—Bras.
butyrosum, *Willd. Sp. Pl.* ii. 1243.—Guiana.
edule, *Casar. Decad. Nov. Stirp. Bras.* viii. 67.—Bras.
glabrum, *Pers. Syn.* ii. 84.—Guiana.
nuciferum, *Linn. Mant.* ii. 247.—Guiana.
tomentosum, *Willd. Sp. Pl.* ii. 1244.—Guiana.
tuberculosum, Baill. Hist. Pl. iv. 251 = tomentosum.
villosum, *Pers. Syn.* ii. 84.—Guiana.

CARYOCHLOA, Spreng. Syst. iv. Cur. Post. 22 (1827) = **Oryzopsis**, Michx. (Gramin.).
chilensis, Spreng. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 123 (= *Stipa Caryochloa*).
montevidensis, Spreng. Syst. iv. Cur. Post. 30 (= *Stipa setifolia*).—Reg. Argent.
refracta, Spreng. ex Trin. in Mém. Acad. Pétersb. Sér. VI. iii. II. (1835) 124 (= *Stipa Caryochloa*).

CARYOCHLOA, Trin. Gram. Panic. Diss. ii. 54 (1826) = **Luziola**, Juss. (Gramin.).
bahiensis, Steud. Syn. Pl. Gram. 5.—Bras.
brasiliensis, Trin. ex Nees, Agrost. Bras. 299.—Bras.
caudata, Trin. ex Steud. Nom. ed. II. i. 301.—Bras.

CARYOCOCCA, Willd. ex Roem. & Schult. Syst. iii. Mant. 131 (1827) = **Gonzalea**, Pers. (Rubiac.).
dependens, Willd. ex Roem. & Schult. l. c. = G. pendula.
multiflora, Willd. ex Roem. & Schult. l. c. 132 = G. pendula.

CARYODAPHNE, Blume, ex Nees, Syst. Laurin. 225 (1836) = **Cryptocarya**, R. Br. (Laurin.).
australis, A. Br. Ind. Sem. Hort. Berol. (1851) App. 13 = Crypt. australis.
Browniana, Nees, Syst. Laurin. 230 = Crypt. triplinervis.
costata, Miq. in Zoll. Syst. Verz. Ind. Archip. 113, 115 = Crypt. costata.
densiflora, Blume, ex Nees, Syst. Laurin. 228 = Crypt. densiflora.
laevigata, Blume, l. c. 227 = Crypt. laevigata.

CARYODENDRON, Karst. Fl. Colomb. i. 91. t. 45 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 314.
CENTRODISCUS, Muell. Arg. in Mart. Fl. Bras. xi. II. 325. t. 102 (1874).
orinocense, *Karst. l. c.*—N. Granat.

CARYOLOBIS, Gaertn. Fruct. i. 215. t. 45 (1788) = **Doona**, Thw. (Dipteroc.).
indica, Gaertn. l. c. = D. cordifolia?

CARYOLOBIUM, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 266 = **Astragalus**, Tourn. (Legumin.).
macrocarpos, Stev. l. c. = A. macrocarpus.

CARYOLOPHA, Fisch. & Trautv. Ind. Sem. Hort. Petrop. iii. 31 (1837) = **Anchusa**, Linn. (Boragin.).
sempervirens, Fisch. & Trautv. l. c. = A. sempervirens.

CARYOPHYLLATA, Tourn. ex Scop. Fl. Carn. ed. II. i. 364 (1772) = **Geum**, Tourn. (Rosac.).
alba, Moench, Meth. 660 = G. album.
aquatica, Lam. Fl. Fr. iii. 123 = G. rivale.
kamtchatica, Lam. Encyc. i. 400 = G. anemonoides.
montana, Scop. Fl. Carn. ed. II. i. 365 = G. montanum.
nutans, Lam. Encyc. i. 399 = G. nutans.
nutans, Moench, Meth. 661 = G. rivale.
obliqua, Moench, l. c. Suppl. 280 = G. obliquum.
officinalis, Moench, Meth. 660 = G. urbanum.
potentilloides, Lam. Encyc. i. 400 = Coluria geoides.
prolifera, Gilib. Fl. Lituan. 416.—Europ.
reptans, Lam. Fl. Fr. iii. 122 = G. reptans.
rivalis, Scop. Fl. Carn. ed. II. i. 365 = G. rivale.
urbana, Scop. l. c. 364 = G. urbanum.
virginiana, Lam. Encyc. i. 399 = G. virginianum.
vulgaris, Lam. Fl. Fr. iii. 122 = G. urbanum.

CARYOPHYLLEA, Opiz, Seznam, 27 (1852) = **Aira**, Linn. (Gramin.).
airoides, Opiz, l. c. = A. caryophyllea.
praecox, Opiz, l. c. = A. praecox.

CARYOPHYLLUS, Linn. Syst. ed. I (1735) = **Eugenia**, Linn. (Myrtac.).
acutangulus, Stokes, Bot. Mat. Med. iii. 75 = Barringtonia acutangula.
adenolepis, Miq. Fl. Ind. Bat. i. 466.—Java.
antisepticus, Blume, ex DC. Prod. iii. 262.—Java.
aromaticus, Linn. Sp. Pl. 515 = E. caryophyllata.
baladensis, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 185.—N. Caled.
corticosus, Stokes, Bot. Mat. Med. iii. 75 = E. Jambolana.

CARYOPHYLLUS :—
cotinifolius, Mill. Gard. Dict. ed. VIII. n. 4 = Myrcia coriacea.
elegans, Brongn. & Gris, in Bull. Soc. Bot. Fr. xii. (1865) 184.—N. Caled.
ellipticus, Labill. Sert. Austr. Caled. 64. t. 63.— N. Caled.
fastigiatus, Blume, ex DC. Prod. iii. 262.—Java.
floribundus, Blume, ex DC. l. c.—Java.
fruticosus, Mill. Gard. Dict. ed. VIII. n. 3 = E. biflora.
Jambos, Stokes, Bot. Mat. Med. iii. 73 = E. Jambos.
hortensis, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 11 = E. caryophyllata ?
malaccensis, Stokes, Bot. Mat. Med. iii. 72 = E. malaccensis.
Pimenta, Mill. Gard. Dict. ed. VIII. n. 2 = Pimenta officinalis.
pterocarpus, Vieill. ex Panch. & Sebert, Not. Bois. Nouv. Caléd. 256.—N. Caled.
racemosus, Mill. Gard. Dict. ed. VIII. n. 5 = E. acris.
racemosus, Stokes, Bot. Mat. Med. iii. 76 = Barringtonia racemosa.

CARYOPHYLLUS, Tourn. ex Moench, Meth. 58 (1794) = **Dianthus**, Linn. (Caryophyll.).
aridus, Moench, Meth. 59 = D. prolifer.
armerius, Moench, l. c. = D. Armeria.
atrorubens, Moench, Meth. Suppl. 23 = D. atrorubens.
barbatus, Moench, Meth. 59 = D. barbatus.
Carthusianorum, Moench, l. c. = D. Carthusianorum.
deltoides, Moench, l. c. = D. deltoides.
glaucus, Moench, l. c. = D. deltoides.
plumarius, Moench, l. c. = D. plumarius.
superbus, Moench, l. c. = D. superbus.

CARYOPTERIS, Bunge, Pl. Mongholico-Chin. 27 (1835). *VERBENACEAE*, Benth. & Hook. f. ii. 1157.
BARBULA, Lour. Fl. Cochinch. 366 (1790), nomen prius.
MASTACANTHUS, Endl. Gen. 638 (1839).
divaricata, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 390.—Japon.
grata, *Benth. & Hook. f. Gen.* ii. 1158.—Reg. Himal.
incana, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 97.—Japon.
Mastacanthus, *Schau. in DC. Prod.* xi. 625.—China.
mongholica, *Bunge, Fl. Mongholico-Chin.* 28.—China bor.
nepetaefolia, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 390.—China.
ovata, *Miq. in Journ. Bot. Néerl.* i. (1861) 114.—China.
paniculata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 597.—Reg. Himal.
tangutica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 525.—China.
terniflora, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 40.—China.
Wallichiana, *Schau. in DC. Prod.* xi. 625.—Reg. Himal.

CARYOSPERMUM, Blume, Mus. Bot. Lugd. Bat. i. 175 (1850). *CELASTRINEAE*, Benth. & Hook. f. i. 367.
arborescens, *F. Muell. Fragm.* viii. 279.—Austral.
moluccanum, *Blume, Mus. Bot. Lugd. Bat.* i. 176.—Ins. Molucc.
serrulatum, *Miq. Fl. Ind. Bat.* i. II. 592.—Java.

CARYOTA, Linn. Mus. Cliff. 12 ; et Gen. ed. I. 355 (1737). *PALMAE*, Benth. & Hook. f. iii. 918.
Alberti, F. Muell. ex H. Wendl. in Linnaea, xxxix. (1875) 221 = Rumphiana.
Cumingii, *Lodd. ex Mart. Hist. Nat. Palm.* iii. 315.—Ins. Philipp.
elegans, *Hort. ex H. Wendl. in Kerch. Palm.* 238, nomen.—Hab. ?
excelsa, *Hort. ex H. Wendl. l. c.* nomen.—Hab. ?
furfuracea, Blume, in Mart. Hist. Nat. Palm. iii. 195 = mitis.
Griffithii, Becc. in Nuov. Giorn. Bot. iii. (1871) 15 = mitis.
horrida, Jacq. Fragm. 20 = Bactris caryotaefolia.

CARYOTA :—
humilis, Reinw. ex Kunth, Enum. Pl. iii. 193 = Didymosperma porphyrocarpa.
javanica, Zipp. ex Miq. Fl. Ind. Bat. iii. 41 = mitis.
majestica, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Ins. Philipp.
maxima, *Blume, in Mart. Hist. Nat. Palm.* iii. 195.—Java.
mitis, *Lour. Cochinch.* 697.—Malaya ; Cochinch.
nana, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Ins. Philipp.
nana, Wall. Cat. n. 8595 = mitis.
Nó, *Becc. in Nuov. Giorn. Bot.* iii. (1871) 12.—Borneo.
obtusa, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) 480.—Reg. Himal.
obtusidentata, Griff. Palms Brit. Ind. t. 236 A, B = praec.
ochlandra, Hance, in Journ. Bot. xvii. (1879) 174 = obtusa ?
onusta, Blanco, Fl. Filip. ed. I. 741 = Arenga saccharifera.
Palindan, Blanco, l. c. ed. I. 714; ed. II. 513 = Orania regalis.
Princeps, *Voigt, in Syll. Ratisb.* ii. (1828) 51.—Hab. ?
propinqua, Blume, in Mart. Hist. Nat. Palm. iii. 195 = mitis.
Rumphiana, *Mart. l. c.*—Malaya ; Austral.
sobolifera, Wall. in Mart. l. c. 194. t. 107. f. 2 = mitis.
speciosa, *Linden, Illustr. Hortic.* xxviii. (1881) 16.—Ins. Philipp.
tremula, Blanco, Fl. Filip. ed. I. 744 = Didymosperma tremula.
urens, Blanco, l. c. 740 = Cumingii.
urens, Jacq. Fragm. 20. t. 12. f. 1 = mitis.
urens, *Linn. Sp. Pl.* 1189.—As. trop.

CARYOTAXUS, Zucc. ex Endl. Syn. Conif. 240 (1847) = **Torreya**, Arn. (Conif.).
grandis, Henk. & Hochst. Nadelhoelz. 367 = T. grandis.
Myristica, Henk. & Hochst. l. c. 368 = T. californica.
nucifera, Zucc. ex Endl. Syn. Conif. 241 = T. nucifera.
taxifolia, Henk. & Hochst. Nadelhoelz. 367 = T. taxifolia.

CASALEA, St. Hil. Fl. Bras. Mer. i. 8. t. 1 (1825) = **Ranunculus**, Tourn.
ascendens, St. Hil. l. c. 11 = R. cordifolius.
biternata, St. Hil. l. c. 9, in obs. = R. biternatus.
breviscapa, St. Hil. l. c. = R. breviscapus.
chilensis, St. Hil. l. c. = R. chilensis.
ficarifolia, St. Hil. l. c. 10 = R. bonariensis.
flagelliformis, St. Hil. l. c. 11 = R. flagelliformis.
hederacea, Mart. ex Eichl. in Mart. Fl. Bras. xiii. i. 156 = R. sessiliflorus.
phyteumaefolia, St. Hil. Bras. Mer. i. 9 = R. bonariensis.
sessiliflora, St. Hil. l. c. = R. platensis.

CASARETTOA, Walp. Rep. iv. 91 (1844) = **Vitex**, Linn. (Verbenac.).
diversifolia, Walp. l. c. 92 = V. triflora.
mollissima, Walp. l. c. 91 = V. polygama.

CASASIA, A. Rich. Fl. Cub. Fanerog. ii. 9. t. 49 (1853). *RUBIACEAE*, Benth. & Hook. f. ii. 84.
calophylla, *A. Rich. l. c.*—Cuba.

CASCABELA, Rafin. Sylva Tellur. 162 (1838) = **Cerbera**, Linn. (Apocyn.).
peruviana, Rafin. l. c. = Cerb. peruviana.

CASCARILLA, Adans. Fam. ii. 355 (1763) = **Croton**, Linn. (Euphorbiac.).
aromatica, Rafin. Sylva Tellur. 62 = Crot. aromaticus.
discolor, Rafin. l. c. = Crot. discolor.
linearis, Rafin. l. c. = Crot. linearis.
officinalis, Rafin. l. c. = Crot. Cascarilla.

CASCARILLA, Ruiz, ex Steud. Nom. ed. I. 165 (1821) = **Cinchona**, Linn. (Rubiac.).
pallida, Steud. l. c. = Cinch. cordifolia.

CASCARILLA, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 10, 77. *RUBIACEAE*, Benth. & Hook. f. ii. 32.

CASCARILLA :—
BUENA, Pohl, ex Wedd. in Journ. Linn. Soc. xi. (1871) 185.
LADENBERGIA, Klotzsch, in Hayne, Arzn. Gew. xiv. in not. ad t. 15 (1846).
acutifolia, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 11.—Pernv.
bullata, *Wedd. l. c.* 11.—Peruv.
caduciflora, *Wedd. l. c.* 10.—Peruv.
calycina, *Wedd. l. c.* 12.—N. Granat.
calyptrata, *Wedd. l. c.* 13.—Peruv.
Carua, *Wedd. l. c.* 12.—Peruv.
citrifolia, *Wedd. l. c.* 11.—Venezuela.
Gaudichaudiana, *Wedd. l. c.* 12.—Bras.
gavanensis, *Schlecht. in Linnaea*, xxvi. (1853) 730.—Peruv.
grandiflora, *Linden, in Belg. Hortic.* vii. (1857) 197.—N. Granat.
heterophylla, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 10.—N. Granat.
hexandra, *Wedd. l. c.* 12.—Bras.
Hookeriana, *Wedd. Hist. Nat. Quinq.* 91.—N. Granat.
Lambertiana, *Wedd. l. c.* 85.—Bras.
macrocarpa, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 13.—N. Granat. ; Peruv.
magnifolia, *Wedd. l. c.* 10.—Bolivia.
muzonensis, *Wedd. l. c.* 11.—N. Granat.
nitida, *Wedd. l. c.* 11.—Venezuela.
oblongifolia, Wedd. l. c. 12 = magnifolia.
officinalis, *Ruiz, Quinol.* 56.—Peruv.
Pavonii, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 11.—Peruv.
Riedeliana, *Wedd. l. c.* 12.—Bras.
Riveroana, *Wedd. l. c.* 11.—Peruv.
Roraimae, *Wedd. l. c.* 12.—Guiana.
rostrata, *Wedd. l. c.* 10.—Boliv. ; Pernv.
stenocarpa, *Wedd. Hist. Nat. Quinq.* 81.—Peruv.
undata, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 12.—Venezuela.

CASCARONIA, Griseb. in Goett. Abh. xxiv. (1879) 100. *LEGUMINOSAE*.
astragalina, *Griseb. l. c.*—Reg. Argent.

CASCOELYTRUM, Beauv. Agrost. 85 (1812) = CHASCOLYTRUM, Desv. = **Briza**, Linn. (Gramin.).

CASEARIA, Jacq. Enum. Pl. Carib. 4 (1760). *SAMYDACEAE*, Benth. & Hook. f. i. 796.
ANAVINGA, Lam. Encyc. i. 147 (1783).
ANTIGONA, Vell. Fl. Flum. iv. t. 145 (1827).
ATHENAEA, Schreb. Gen. i. 259 (1789).
BEDUSIA, Rafin. Sylva Tellur. 11 (1838).
CHAETOCRATER, Ruiz & Pav. Prod. Fl. Per. 61. t. 36 (1794).
CHETOCRATER, Rafin. Sylva Tellur. 149 (1838).
CLASTA, Comm. ex Vent. Choix, sub t. 47 (1803).
CORIZOSPERMUM, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 255 (1850).
CRATERIA, Pers. Syn. i. 485 (1805).
GUIDONIA, Griseb. Fl. Brit. W. Ind. 24 (1864).
IROUCANA, Anbl. Pl. Gui. i. 328. t. 127 (1775).
LANGLEIA, Scop. Introd. 231 (1777).
LINDLEYA, Kunth, Malv. 10 (1822).
MELISTAURUM, Forst. Char. Gen. 143. t. 72 (1776).
MOELLERIA, Scop. Introd. 335 (1777).
PIPAREA, Aubl. Pl. Gui. ii. App. 30. t. 386 (1775).
PITUMBA, Aubl. l. c. 29. t. 385 (1775).
THIODIA, Griseb. Fl. Brit. W. Ind. 22 (1864).
VALENTINIA, Sw. Prod. Veg. Ind. Occ. 63 (1788).
VARECA, Gaertn. Fruct. i. 290. t. 60 (1788).
ZUELANIA, A. Rich. Ess. Fl. Cub. 88. t. 12 (1845).
aculeata, Jacq. Enum. Pl. Carib. 21 = spinosa.
acuminata, *DC. Prod.* ii. 50.—Guiana.
acuminata, Wall. Cat. n. 7198 (non Casearia).—Ind. or.
acuminatissima, *A. Gray, Bot. U. St. Expl. Exped.* i. 80.—Ins. Fiji.
adamantium, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 230.—Bras.
adstringens, *Mart. Mat. Med. Veg. Bras.* 51.—Bras.
affinis, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 529 = sylvestris.
alba, *A. Rich. Ess. Fl. Cub.* 388.—Cuba.
albicans, *Wall. Cat.* n. 7197.—Malaya.
amplissima, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 327.—Madag.
Anavinga, Dalz. & Gibs. Bomb. Fl. 11 = tomentosa.

CASEARIA:—
Anavinga, Pers. Syn. i. 485 = ovata.
angustata, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 409.—Java.
aquifolia, *Sauv. Fl. Cub.* 6.—Cuba.
arguta, H. B. & K. Nov. Gen. et Sp. v. 363 = parvifolia.
astyla, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 608.—Ind. or.
avellana, Miq. ex Eichl. in Mart. Fl. Bras. xiii. 1. 463 = spinosa.
Barteri, *Mast. in Oliver, Fl. Trop. Afr.* ii. 494.—Afr. trop.
Benthamiana, *Miq. in Ann. Sc. Nat.* Sér. III. i. (1844) 38.—Bras.
Berteriana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 461.—Porto Rico.
Blanchetiana, *Miq. in Linnaea*, xxii. (1849) 801.—Bras.
brasiliensis, *Eichl. in Mart. Fl. Bras.* xiii. 1. 477.—Bras.
brevipes, *Benth. in Hook. Journ. Bot.* iv. (1842) 110.—Guiana; Ind. occ.
Cambessedesii, *Eichl. in Mart. Fl. Bras.* xiii. 1. 475.—Bras.
Canziala, Buch.-Ham. in Wall. Cat. n. 7192 = tomentosa.
capitata, Spreng. Syst. ii. 355 = sylvestris.
capitellata, *Blume, Mus. Bot. Lugd. Bat.* i. 254.—Borneo.
carpinifolia, Benth. in Hook. Journ. Bot. iv. (1842) 112 = parvifolia.
celastroides, *Klotzsch, in Schomb. Faun. et Fl. Guy.* 1167, nomen.—Guiana.
celtidifolia, *H. B. & K. Nov. Gen. et Sp.* v. 362.—Venezuela.
Championi, Thw. Enum. Pl. Zeyl. 19 = esculenta.
cinerea, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 462 = grewiaefolia.
Clausseniana, *Miq. in Linnaea*, xix. (1847) 441.—Bras.
clutiaefolia, *Blume, Mus. Bot. Lugd. Bat.* i. 255.—N. Guinea.
combaymensis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 362.—N. Granat.
Commersoniana, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 235.—Bras.
comocladifolia, *Vent. Choix*, t. 44.—Ins. S. Doming.
congestiflora, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 608.—Bras.
contermina, *Miq. Fl. Ind. Bat.* i. 1. 707.—Java.
coriacea, *Thw. Enum. Pl. Zeyl.* 20.—Zeylan.
coriacea, *Vent. Choix*, t. 45.—Java.
coriacea, Wall. Cat. n. 7196 = Chaetocarpus castanocarpus.
corymbosa, *H. B. & K. Nov. Gen. et Sp.* v. 366.—N. Granat.
cuspidata, *Blume, Mus. Bot. Lugd. Bat.* i. 255.—Sumatra.
Dallachii, F. Muell. Fragm. v. 107 = tomentosa.
decandra, Jacq. Enum. Pl. Carib. 21 = parvifolia.
densiflora, *Benth. in Hook. Journ. Bot.* iv. (1842) 113.—Guiana.
dentata, *Moç. & Sesse, in DC. Prod.* ii. 51.—Mexic.
disticha, A. Gray, Bot. U. St. Expl. Exped. i. 81 = C. Melistaurum.
dubia, *Moç. & Sesse, in DC. Prod.* ii. 51.—Mexic.
elliptica, *Klotzsch, in Schomb. Faun. & Fl. Guy.* 1167, nomen.—Guiana.
elliptica, Willd. Sp. Pl. ii. 628 = tomentosa.
emarginata, *Griseb. Cat. Pl. Cub.* 10.—Cuba.
eriophora, *Wright, ex Griseb. l. c.* 11.—Cuba.
esculenta, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 422.—As. trop.; Austral.
fallax, *Miq. in Linnaea*, xix. (1847) 128.—Guiana.
fasciculata, Boj. Hort. Maurit. 71 = fragilis.
flavovirens, *Blume, Mus. Bot. Lugd. Bat.* i. 254.—Java.
Fockeana, *Miq. in Ann. Sc. Nat.* Sér. III. i. (1844) 39.—Guiana.
fragilis, *Vent. Choix*, 47, verso.—Ins. Mascar.
fuliginosa, *Blanco, Fl. Filip.* ed. II. 262.—Ins. Philipp.
glabra, Kurz, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 591 = glomerata.
glabra, Roxb. Hort. Beng. 33; Fl. Ind. ii. 421 = tomentosa.
glabrata, *Blume, Mus. Bot. Lugd. Bat.* i. 253.—Java.

CASEARIA:—
gladiiformis, *Mast. in Oliver, Fl. Trop. Afr.* ii. 493.—Afr. trop.
glomerata, *Roxb. Hort. Beng.* 33; *Fl. Ind.* ii. 419.—Ind. or.
gonocarpa, *Miq. Fl. Ind. Bat.* i. 1. 1093.—Sumatra.
grandiflora, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 232.—Bras.
grandifolia, *Miq. Fl. Ind. Bat.* i. 1. 712.—Sumatra.
graveolens, *Dalz. in Hook. Kew Journ.* iv. (1852) 107.—Ind. or.
grewiaefolia, *Vent. Choix*, 48.—Malaya.
Guidonia, Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 89 = brevipes.
guineensis, *G. Don, Gen. Syst.* ii. 53.—Afr. trop.
Hamiltonii, Wall. Cat. n. 7195 = graveolens.
hexagona, *Decne. in Ann. Nouv. Mus. Par.* iii. (1834) 429.—Ins. Timor.
hirsuta, *Sw. Fl. Ind. Occ.* ii. 755.—Ind. occ.
hirta, *Sw. l. c.* 756.—Ind. occ.
Hostmanniana, Steud. ex Griseb. Fl. Brit. W. Ind. 23 = stipularis.
hypoleuca, *Mart. in Flora*, xx. (1837) *Beibl.* ii. 128.—Bras.
icosandra, *Planch. & Triana, ex Hemsl. Biol. Centr. Am. Bot.* i. 469.—Panama.
ilicifolia, *Vent. Choix*, t. 44.—Ins. S. Doming.
impunctata, *Hook. & Arn. Bot. Beech. Voy.* 61.—Ins. Societat.
inaequilatera, *Cambess. in A. St. Hil. Fl. Bras. Mer.* ii. 237.—Bras.
integrifolia, Vahl, ex DC. Prod. ii. 49 = sylvestris.
javitensis, *H. B. & K. Nov. Gen. et Sp.* v. 366.—Am. austr.
Kurzii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 594.—Burma.
laevigata, Dalz. in Hook. Kew Journ. iv. (1852) 108 = esculenta.
lanceolata, Miq. in Linnaea, xviii. (1844) 753 = stipularis.
lasiophylla, *Eichl. in Mart. Fl. Bras.* xiii. 1. 468.—Bras.
lasiosperma, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 113.—N. Granat.
laurifolia, *Benth. in Hook. Journ. Bot.* iv. (1842) 113.—Guiana.
laurina, *Blume, Mus. Bot. Lugd. Bat.* i. 253.—Malaya.
leucolepis, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) 1. 463.—Malaya.
Lingua, Cambess. in St. Hil. Fl. Bras. Mer. ii. 236 = sylvestris.
Lobbiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 463.—Burma; Malaya.
lucida, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 325.—Madag.
lucida, Wall. Cat. n. 7199 (Euphorbiac.).—Ind. or.
macrocarpa, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 593.—Penang.
macrogyna, Turcz. in Bull. Soc. Nat. Mosc. (1858) 1. 463 = graveolens.
macrophylla, *Vahl, Eclog. Am.* ii. 32.—Guiana.
Mannii, *Mast. in Oliver, Fl. Trop. Afr.* ii. 494.—Afr. trop.
mariquitensis, *H. B. & K. Nov. Gen. et Sp.* v. 363.—N. Granat.
Mathewsii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 608.—Peruv.
Maximiliana, *Eichl. in Mart. Fl. Bras.* xiii. 1. 487.—Bras.
meliodora, *Eichl. l. c.* 469.—Bras.
Melistaurum, *Spreng. Syst.* ii. 354.—Ins. Pacif.
membranacea, *Hance, in Journ. Bot.* vi. (1868) 113.—China.
micrantha, G. Don, Gen. Syst. ii. 52 = parviflora.
microdon, *Miq. Fl. Ind. Bat. Suppl.* 333.—Sumatra.
microphylla, *Eichl. in Mart. Fl. Bras.* xiii. 1. 474.—Bras.
mollis, H. B. & K. Nov. Gen. et Sp. v. 362. t. 480 = hirsuta.
moluccana, *Blume, Mus. Bot. Lugd. Bat.* i. 255.—Amboina.
montana, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 335 = inaequilatera.
multiflora, *Spreng. Pugill.* ii. 61.—Ind. or.
myriantha, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 609.—Panama.
negrensis, *Eichl. in Mart. Fl. Bras.* xiii. 1. 466.—Bras.
nigrescens, *Tul. in Ann. Sc. Nat.* Sér. V. ix. (1868) 326.—Madag.

CASEARIA:—
nitida, *Jacq. Enum. Pl. Carib.* 21.—N. Granat.
nitida, Sieber, ex Griseb. Fl. Brit. W. Ind. 23 = parvifolia.
obliqua, Spreng. Syst. ii. 355 = inaequilatera.
oblongifolia, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 234.—Bras.
obovalis, Poepp. ex Griseb. in Goett. Abh. ix. (1860) 27 = obovata.
obovata, *Poepp. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 472.—Peruv.
obovata, *Schlecht. in Linnaea*, xiii. (1839) 434.—Mexic.
odorata, Macfad. Fl. Jamaic. i. 216 = ramiflora.
odorata, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 408.—Java.
oligantha, *Eichl. in Mart. Fl. Bras.* xiii. 1. 476.—Bras.
ovata, Roxb. Hort. Beng. [90]; Fl. Ind. ii. 420 = tomentosa.
ovata, Wall. Cat. n. 7192 E = glomerata.
ovata, *Willd. Sp. Pl.* ii. 629.—Ind. or.
paniculata, Gardn. ex Thw. Enum. Pl. Zeyl. 20 = Osmelia zeylanica.
parviflora, J. F. Gmel. Syst. 700 = parvifolia.
parviflora, Willd. Sp. Pl. ii. 627 = sylvestris.
parvifolia, *Willd. l. c.* 628.—Ind. occ.
pauciflora, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 235.—Bras.
petiolaris, *Poepp. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 471.—Peruv.
petraea, *Benth. in Hook. Journ. Bot.* iv. (1842) 111.—Guiana.
Pitumba, DC. Prod. ii. 50, in syn. = macrophylla.
Poeppigii, *Eichl. in Mart. Fl. Bras.* xiii. 1. 475.—Bras.
Pohliana, *Eichl. l. c.* 474.—Bras.
praecox, *Griseb. Cat. Pl. Cub.* 10.—Cuba.
prismatocarpa, *Mast. in Oliver, Fl. Trop. Afr.* ii. 494.—Afr. trop.
propinqua, *Blume, Mus. Bot. Lugd. Bat.* i. 253.—Java.
prunifolia, *H. B. & K. Nov. Gen. et Sp.* v. 362.—Am. austr.
prunifolia, Tul. in Ann. Sc. Nat. Sér. III. vii. (1847) 363 = spinosa.
pubiflora, Benth. Bot. Voy. Sulph. 66 = parvifolia.
punctata, Spreng. Neue Entdeck. ii. 154 = parvifolia.
punctata, *Spreng. Syst.* ii. 355.—Am. trop.
quinduensis, *Tul. in Ann. Sc. Nat.* Sér. III. vii. (1847) 360.—Ecuador.
ramiflora, Seem. Bot. Voy. Herald, 98 = hirsuta.
ramiflora, *Vahl, Symb. Bot.* ii. 50.—Am. trop.
ramosissima, *Griseb. Cat. Pl. Cub.* 10.—Cuba.
resinifera, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiii. 1. 466.—Bras.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 82.—Ins. Fiji.
rubescens, *Dalz. in Hook. Kew Journ.* iv. (1852) 108.—Ind. or.
rufescens, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 231.—Bras.
rufidula, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 107.—N. Granat.
rufinervis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 462.—Cuba.
rugulosa, *Blume, Mus. Bot. Lugd. Bat.* i. 255.—Malaya.
rupestris, *Eichl. in Mart. Fl. Bras.* xiii. 1. 468.—Bras.
salacioides, *Blume, Mus. Bot. Lugd. Bat.* i. 252.—N. Guinea.
salicifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 460.—Mexic.
Samyda, *DC. Prod.* ii. 51.—Porto Rico.
Selloana, *Eichl. in Mart. Fl. Bras.* xiii. 1. 483.—Bras.
serrata, Macfad. Fl. Jamaic. i. 215 = serrulata.
serrulata, Sieber, ex Griseb. Fl. Brit. W. Ind. 23 = parvifolia.
serrulata, *Sw. Fl. Ind. Occ.* ii. 754.—Ind. occ.
sessiliflora, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 231.—Bras.
singularis, *Eichl. in Mart. Fl. Bras.* xiii. 1. 473.—Guiana.
spinescens, *Griseb. Cat. Pl. Cub.* 10.—Cuba.
spinosa, Willd. Sp. Pl. ii. 626 = ramiflora.
Spruceana, *Benth. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 486.—Bras.

CASEARIA:—
squarrosa, *Klotzsch, in Schomb. Faun. et Fl. Guy.* 1167, nomen.—Guiana.
stipitata, *Mast. in Oliver, Fl. Trop. Afr.* ii. 493.—Afr. trop.
stipularis, Cambess. in St. Hil. Fl. Bras. Mer. ii. 169 = brasiliensis.
stipularis, *Vent. Choix*, 46. t. 46.—Am. austr.
subciliaris, *Klotzsch, in Schomb. Faun. et Fl. Guy.* 984, nomen.—Guiana.
subcuneata, Miq. Fl. Ind. Bat. i. 706 = grewiaefolia.
subopaca, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 110.—N. Granat.
subrhombea, *Hance, in Journ. Bot.* xxiii. (1885) 323.—China.
sylvestris, *Sw. Fl. Ind. Occ.* ii. 752.—Ind. occ.
tinifolia, *Vent. Choix*, 47. t. 47.—Hab. ?
tomentosa, *Roxb. Hort. Beng.* [90]; *Fl. Ind.* ii. 421.—As. trop.; Austral.
truncata, *Blume, Mus. Bot. Lugd. Bat.* i. 252.—Malaya.
tuberculata, *Blume, l. c.* 254.—Java.
tuberculata, *Blume, l. c.* 255.—Sumatra.
ulmifolia, Cambess. in St. Hil. Fl. Bras. Mer. ii. 169 = Cambessedesii.
ulmifolia, Vahl, ex Vent. Choix, 46, verso = serrulata.
uniflora, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) 428.—Ins. Timor.
Vareca, *Roxb. Hort. Beng.* 33; *Fl. Ind.* ii. 418.—Reg. Himal.; Burma.
variabilis, Blume, Mus. Bot. Lugd. Bat. i. 252 = grewiaefolia.
varians, Thw. Enum. Pl. Zeyl. 19 = esculenta.
velutina, *Blume, Mus. Bot. Lugd. Bat.* i. 253.—Sumatra.
viridiflora, *DC. Prod.* ii. 51.—Ind. or.
viridiflora, Lam. Encyc. vi. 493 = leucolepis.
wynadensis, *Bedd. Ic. Pl. Ind. Or.* i. 34.—Ind. or.
zeylanica, Thw. Enum. Pl. Zeyl. 19 = esculenta.
zizyphoides, *H. B. & K. Nov. Gen. et Sp.* v. 362.—Am. austr.

CASEOLA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 (Quid ?).

CASIMIRA, Scop. Introd. 233 (1777) = **Melicocca**, Linn. (Sapind.).

CASIMIROA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 2 (1825). *RUTACEAE*, Benth. & Hook. f. i. 302.
edulis, *La Llave, l. c.*—Mexic.
Sapota, *Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 187.—Nicaragua.
tomentosa, Domb. ex Miers, in Journ. Linn. Soc. xvii. (1880) 79 = Cervantesia tomentosa.

CASINGA, Griseb. in Goett. Abh. ix. (1861) 27, 29 = **Laetia**, Linn. (Olacin.).
procera, Griseb. Fl. Brit. W. Ind. 710 = L. procera.
suaveolens, Griseb. in Goett. Abh. ix. (1861) 27 = L. suaveolens.

CASIOSTEGA, Rupr. ex Galeotti, in Bull. Acad. Brux. ix. II. (1842) 232, nomen = **Opizia**, Presl (Gramin.).
anomala, Rupr. l. c. = O. stolonifera.
dactyloides, Fourn. in Bull. Soc. Bot. Belg. xv. (1876) 470 = Buchloë dactyloides.
Hookeri, Rupr. ex Fourn. l. c. 471 = Buchloë dactyloides.
humilis, Rupr. [ex Munro] in Benth. Pl. Hartw. 347 = = Buchloë dactyloides.

CASPAREA, H. B. & K. Nov. Gen. et Sp. vi. 317 (1823) = **Bauhinia**, Linn. (Legumin.).
acuminata, Heynh. Nom. ii. 123 = B. acuminata.
aurita, Griseb. Fl. Brit. W. Ind. 213 = B. aurita.
castrata, Hassk. Cat. Hort. Bogor. 288 = B. castrata.
divaricata, H. B. & K. Nov. Gen. et Sp. vi. 317 = B. divaricata.
furcata, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 429 = B. furcata.
latifolia, H. B. & K. Nov. Gen. et Sp. vi. 317 = B. latifolia.
lunaria, H. B. & K. l. c. = Bauhinia lunaria.
Pes-caprae, H. B. & K. l. c. 318 = B. Pes-caprae.
porrecta, H. B. & K. l. c. 317 = B. porrecta.
subrotundifolia, H. B. & K. l. c. = B. subrotundifolia.

CASPARYA, Klotzsch, in Monatsb. Berl. Acad. (Maerz 1854) 127 = **Begonia**, Linn.
antioquensis, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 116.—N. Granat.
brevipetala, A. DC. l. c. 118.—Venezuela.
coccinea, Klotzsch, in Monatsb. Berl. Acad. (1854) 127 = B. columnaris.
columnaris, Klotzsch, l. c. = B. columnaris.
cordifolia, A. DC. in DC. Prod. xv. I. 273 = *hirta*.
crassicaulis, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 119.—Java.
elegans, Klotzsch, in Monatsb. Berl. Acad. (1854) 127 = B. elegans.
erosa, A. DC. in DC. Prod. xv. I. 276 = B. erosa.
ferruginea, A. DC. l. c. 269 = B. ferruginea.
fuchsiaeflora, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 116.—Ecuador.
grewiaefolia, A. DC. l. c. 117.—N. Granat.
hirta, Klotzsch, in Monatsb. Acad. Berl. (1854) 127.—Peruv.
longirostris, A. DC. in DC. Prod. xv. I. 272 = B. longirostris.
montana, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 118.—Venezuela.
multiangula, A. DC. in DC. Prod. xv. I. 275 = B. multiangula.
oligocarpa, A. DC. in Ann. Nat. Sér. IV. xi. (1859) 118 = B. Roxburghii.
polycarpa, A. DC. l. c. = B. Roxburghii.
robusta, A. DC. in DC. Prod. xv. I. 275 = B. robusta.
silhetensis, A. DC. l. c. 277 = B. silhetensis.
Teysmanniana, Miq. ex A. DC. l. c. 276.—Sumatra.
trachyptera, A. DC. l. c. 274 = B. trachyptera.
Trianaei, A. DC. in Ann. Sc. Nat. Sér. IV. xi. (1859) 117.—N. Granat.
trispathulata, A. DC. l. c.—N. Granat.
trisulcata, A. DC. l. c. 119.—Java.
umbellata, A. DC. in DC. Prod. xv. I. 270 = B. umbellata.
Urticae, A. DC. l. c. 274 = B. Urticae.

CASPIA, Pison. ex Scop. Introd. 276 (1777) = **Vismia**, Vell. (Hypericin.).

CASSANDRA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 158. *ERICACEAE*, Benth. & Hook. f. ii. 584.
DIPLARIA, Rafin. ex DC. Prod. vii. 610 (1839).
angustifolia, D. Don, l. c. = calyculata.
calyculata, *D. Don, l. c.* 158.—Europ.; As. et Am. bor.
racemosa, Spach, Hist. Vég. Phan. ix. 478 = Leucothoë racemosa.

CASSEBEERIA, Dennst. Schluess. Hort. Malab. 35 (1818) = **Sonerila**, Roxb. (Melast.).
maculata, Dennst. l. c. = S. Wallichii.

CASSELIA, Dum. Comm. Bot. 21 (1822) = **Mertensia**, Roth (Boragin.).
bracteata, Dum. l. c. 25 = M. sibirica.
davurica, Dum. l. c. 22 = M. dahurica.
denticulata, Dum. l. c. 23 = M. sibirica.
diversifolia, Regel, Ind. Sem. Hort. Petrop. (1857) 46.—Hab. ?
gracilis, Dum. Comm. Bot. 23 = M. dahurica.
lanceolata, Dum. l. c. 24 = M. lanceolata.
maritima, Dum. l. c. 25 = M. maritima.
paniculata, Dum. l. c. 22 = M. paniculata.
parviflora, Dum. l. c. 25 = M. maritima.
sibirica, Dum. l. c. 23 = M. sibirica.
simplicissima, Dum. l. c. 24 = M. simplicissima.
villosula, Dum. l. c. 23 = M. villosula.
virginica, Dum. l. c. 24 = M. pulmonarioides.

CASSELIA, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 73. t. 6. *VERBENACEAE*, Benth. & Hook. f. ii. 1148.
chamaedryfolia, *Cham. in Linnaea*, vii. (1832) 365.—Bras.
integrifolia, *Nees & Mart. in Nov. Act. Acad. Nat. Cur.* xi. (1823) 76.—Bras.
Mansoi, *Schau. in Mart. Fl. Bras.* ix. 175.—Bras.
serrata, *Nees & Mart. in Nov. Act. Acad. Nat. Cur.* xi. (1823) 75.—Bras.
veronicaefolia, *Cham. in Linnaea*, vii. (1832) 364.—Bras.

CASSIA, Tourn. ex Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 571.
ADIPERA, DIALANTHERA, DIALLOBUS, DISTEREPTA, DIPLOTAX, DITREMEXA, EMELISTA, HEPTEIRECA, HERPETICA, ISANDRINA, NICTITELLA, OCTELISIA, OPHIOCAULON, PANISIA, PEIRANISIA, SCOLODIA, TAGERA, XAMACRISTA, Rafin. Sylva Tellur. 127–129 (1838).
BACTYRILOBIUM, Willd. Enum. Hort. Berol. 439 (1809).
CASSIANA, Rafin. in Am. Monthly Mag. (1818) 266.
CATHARTOCARPUS, Pers. Syn. i. 459 (1805).
CHAMAECRISTA, Moench, Meth. 272 (1794).
CHAMAEFISTULA, G. Don, Gen. Syst. ii. 451 (1832).
GRIMALDIA, Schrank, in Denkschr. Akad. Muench. (1808) 103. t. 3.
MACLEAYA, Montrouz. in Mém. Acad. Lyon, x. (1860) 198.
SENNA, Tourn. ex Mill. Gard. Dict. ed. VIII. (1768).
abbreviata, *Oliver, Fl. Trop. Afr.* ii. 271.—Afr. trop.
Absus, Aubl. Pl. Gui. i. 381 = hispidula.
Absus, *Linn. Sp. Pl.* 376.—Cosmop. trop.
acanthoclada, *Griseb. in Goett. Abh.* xxiv. (1879) 116.—Reg. Argent.
acapulcensis, H. B. & K. Nov. Gen. et Sp. vi. 353 = biflora.
acclinus, F. Muell. Fragm. iv. 13 = glauca.
aciphylla, Benth. ex A. Gray, Bot. U. St. Expl. Exped. i. 465 = australis, *Sims*.
acisperma, *Schrank, in Colla, Hort. Ripul. App.* ii. 343.—Hab. ?
acosmifolia, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 133.—Bras.
aculeata, *Pohl, ex Benth. l. c.* 128.—Bras.
acuminata, Moench, Meth. 273 = marylandica.
acuminata, Willd. Sp. Pl. ii. 517 = C. Apoucouita.
acurucnsis, *Benth. in Mart. Fl. Bras.* xv. II. 122.—Bras.
acuta, Meyen, Reise, i. 376 = Cumingii.
acutifolia, *Delile, Fl. Egypte*, 219. t. 27.—Aegypt.
acutisepala, *Benth. in Mart. Fl. Bras.* xv. II. 97.—Am. trop.
adenantha, Zoll. & Moritz. Syst. Verz. Zoll. 2 = divaricata.
adenensis, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 553.—Arab.
adenopoda, Miq. in Linnaea, xix. (1847) 436 = C. Apoucouita.
adiantifolia, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 130.—Bras.
advena, Willd. ex Vog. Syn. Cass. 18 = bicapsularis.
aegyptiaca, Willd. Enum. Hort. Berol. 442 = C. Sophera.
Aeschynomene, DC. in Collad. Hist. Cass. 127. t. 17 = mimosoides.
aethiopica, *Guibourt, Hist. Drogues*, ed. III. iii. 219.—Aegypt.
affinis, *Benth. in Mart. Fl. Bras.* xv. II. 98.—Am. trop.
Afzeliana, Vog. in Linnaea, xv. (1841) 70 = podocarpa.
alata, *Linn. Sp. Pl.* 378.—Cosmop. trop.
albida, Orteg. Hort. Matr. Dec. 92 = tomentosa.
Alcaparilla, H. B. & K. Nov. Gen. et Sp. vi. 355 = bicapsularis.
Alcaparra, *Phil. in Linnaea*, xxxiii. (1864–65) 61.—Chili.
amoena, Buch.-Ham. in Wall. Cat. n. 5321 = mimosoides.
ampliflora, Steud. in Flora, xxvi. (1843) ii. 760 = multijuga.
anceps, *Benth. in Mart. Fl. Bras.* xv. II. 166.—Am. trop.
andina, Phil. in Linnaea, xxviii. (1856) 685 = Arnottiana.
Andrieuxii, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 548.—Mexic.
andromedea, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 136.—Am. trop.
angulata, *Vog. Syn. Cass.* 16.—Bras.
angustifolia, *Vahl, Symb. Bot.* i. 29.—Afr. trop.
angustisiliqua, Lam. Encyc. i. 649 = biflora.
angustisiliqua, Torr. & Gray, Fl. N. Am. i. 396 = bahamensis.

CASSIA :—

angustissima, Lam. Encyc. i. 650 = mimosoides.
annulata, *Willd. ex Klotzsch, in Schomb. Faun. & Fl. Gui.* 1207, nomen.—Guiana.
aphylla, *Cav. Ic.* vi. 41. t. 561.—Reg. Argent.
apiculata, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 304 = crotalarioides.
Apoucouita, *Aubl. Pl. Gui.* i. 379. t. 146.—Guiana.
appendiculata, *Vog. Syn. Cass.* 24.—Bras.
applanata, Anderss. Galap.-Oarnes Vegetat. 254 = picta.
aprica, Vell. Fl. Flum. 169 ; iv. t. 76 = cathartica.
arachoides, Burch. Trav. i. 341 = obovata.
arayatensis, Llanos, Fragm. 71 = timoriensis.
arayatensis, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 71 = alata.
arborea, Macfad. Fl. Jamaic. i. 343 = siamea.
arborescens, Mill. Gard. Dict. ed. VIII. n. 15 = emarginata.
arborescens, Vahl, Symb. Bot. iii. 56 = glauca.
arenaria, H. B. & K. Nov. Gen. et Sp. vi. 370 = flexuosa.
arequipensis, *Meyen, in Vog. Syn. Cass.* 43.—Peruv.
Arereh, *Delile, Cent. Pl. Afr.* 29.—Aegypt.
argentea, *H. B. & K. Nov. Gen. et Sp.* 358.—Mexic.
aristata, *Benth. in Mart. Fl. Bras.* xv. II. 170.—Bras.
aristata, Willd. ex Walp. Rep. i. 836 = patellaria.
armata, *S. Wats. in Proc. Am. Acad.* xi. (1876) 136.—Calif.
Arnottiana, *Gill. in Hook. Bot. Misc.* iii. (1833) 211.—Chili.
Arnottiana, Wight, ex Benth. in Trans. Linn. Soc. xxvii. (1871) 577 = nigricans.
arowana, *Schomb. Faun. & Fl. Gui.* 1206.—Guian.
artemisioides, *Gaudich. ex DC. Prod.* ii. 495.—Austral.
Aschrek, *Forsk. Fl. Aegypt. Arab.* 86.—Arab.
aspera, Muhl. Cat. ex Ell. Sketch, 474 = nictitans.
astroites, Willd. ex Cham. & Schlecht. in Linnaea, ii. (1827) 597 = villosa.
atomaria, *Linn. Mant.* 68.—Am. austr.
atropurpurea, Benth. in Trans. Linn. Soc. xxvii. (1871) 533, sphalm. = seq.
atroviridis, Span. in Linnaea, xv. (1841) 201 = C. Sophera.
aurantia, *Ruiz & Pav. ex G. Don, Syst.* ii. 442.—Peruv.
aurata, Roxb. Hort. Beng. 32, nomen = laevigata.
auricoma, R. Grah. in Wall. Cat. n. 5322 = mimosoides.
auriculata, *Linn. Sp. Pl.* 379.—Ind. or.
aurita, *Collad. Hist. Cass.* 131.—Bras.
aurivilla, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 152.—Am. trop.
australis, *Reinw. ex Steud. Nom.* ed. II. i. 303, nomen. —Java.
australis, *Sims, Bot. Mag.* t. 2676.—Austral.
australis, Vell. Fl. Flum. 166 ; iv. t. 65 = appendiculata.
aversiflora, Hook. Bot. Mag. t. 2138 = biflora.
bacillaris, *Linn. f. Suppl.* 231.—Am. trop.
bacillaris, Willd. ex Steud. Nom. ed. II. i. 303 = sclerocarpa.
Bacillus, Gaertn. Fruct. ii. 313 = javanica.
bahamensis, *Mill. Gard. Dict.* ed. VIII. n. 9.—Am. trop.
barbata, *Mart. & Nees, in Nov. Act. Nat. Cur.* xii. (1824) 32.—Bras.
Barclayana, Sweet, Fl. Austral. t. 32 = C. Sophera.
Barrenfieldii, Colla, Hort. Ripul. App. ii. 343 = australis.
basifolia, *Vog. Syn. Cass.* 56.—Bras.
batramensis, Steud. Nom. ed. II. i. 303, sphalm. = bahamensis.
bauhiniaefolia, Kunth, Mimos. 123. t. 37 = rotundifolia.
bauhinioides, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 180.—Texas ; N. Mexic.
Berlandieri, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 520.—Mexic.
Berryana, Voigt, Hort. Suburb. Calc. 249 = montana.
Berteriana, Balb. ex DC. Prod. ii. 496 = biflora.
Berterii, Colla, Hort. Ripul. 30. t. 24 = bicapsularis.
bicapsularis, *Linn. Sp. Pl.* 376.—Am. trop.
bifida, Zoll. & Mor. in Nat. en Genesk. Arch. Neerl. Ind. iii. (1846) 68, 80 = divaricata.
biflora, Griseb. Cat. Pl. Cub. 79 = robiniaefolia.
biflora, *Linn. Sp. Pl.* 378.—Am. trop.

CASSIA :—

biflora, Mill. Gard. Dict. ed. VIII. n. 14 (= Aeschynomenes sp.).—Ind. occ. ?
bifoliolata, DC. in Collad. Hist. Cass. 120. t. 9 = rotundifolia.
biglandulosa, Bertol. in Bologn. Opusc. Sc. iii. (1819) 409 = patellaria.
bijuga, Vog. Syn. Cass. 17 = speciosa.
bimarginata, *Griseb. in Bonplandia*, vi. (1858) 5.—Panama.
birostris, *Domb. ex Vog. Syn. Cass.* 43.—Chili.
Blancheti, *Benth. in Hook. Journ. Bot.* ii. (1840) 78. —Bolivia.
bonariensis, Hort. ex Colla, Hort. Ripul. 88 = corymbosa.
Bonplandiana, DC. Prod. ii. 490 = C. Fistula.
borbonioides, Vog. Syn. Cass. 53 = Langsdorffii.
borneensis, *Miq. Anal. Bot. Ind.* i. 9.—Borneo.
Botteriana, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 541.—Mexic.
brachypoda, *Benth. in Mart. Fl. Bras.* xv. II. 172.—Am. trop.
brachystachya, *Benth. in Hook. Journ. Bot.* ii. (1840) 78.—Bras.
brachystachya, Moç. & Sesse, ex DC. Prod. ii. 500 = pauciflora.
bracteata, Linn. f. Suppl. 232 = alata.
bracteolata, *Vog. Syn. Cass.* 52.—Bras.
bracteosa, Welw. ex Oliver, Fl. Trop. Afr. ii. 277 = didymobotrya.
brasiliana, Lam. Encyc. i. 649 = grandis.
brasiliana, Saldanha, Descr. Rio Madeiras Jan. 43 = ferruginea.
brevicalyx, *Benth. in Mart. Fl. Bras.* xv. II. 134.—Bras.
brevifolia, *Lam. Encyc.* i. 651.—Madag.
brevipes, *DC. in Collad. Hist. Cass.* 119. t. 9.—Panama.
Brewsterii, *F. Muell. 4th Ann. Rep.* 17.—Austral.
Brongniartii, *Gaudich. Bot. Voy. Bonite*, t. 10.—Hab. ?
Browniana, Kunth, Mimos. 135. t. 41 = polyantha.
Buchanani, Kostel. Allg. Med.-Pharm. Fl. iv. 1333 = obovata.
Burchellii, *Benth. in Mart. Fl. Bras.* xv. II. 165.—Bras.
Burmanni, DC. Prod. ii. 502 = capensis ?
Burmanni, Wight, in Madras Journ. Sc. vi. (1837) t. 5 = obovata.
caespitosa, *Benth. in Mart. Fl. Bras.* xv. II. 136.—Bras.
calliantha, G. F. W. Mey. Prim. Fl. Esseq. 169 = multijuga.
calycioides, *DC. in Collad. Hist. Cass.* 125. t. 20 B.—Am. trop.
camporum, Benth. in Hook. Journ. Bot. ii. (1840) 79 = pauciflora.
cana, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1824) 34.—Bras.
cana, Schrank, ex Steud. Nom. ed. II. i. 304 = tomentosa.
cana, Wender. in Linnaea, xii. (1838) 22 = holosericea.
canaliculata, R. Br. in Sturt, Centr. Austral. ii. 611 = eremophila.
Canca, Cav. Desc. 132 = C. Sophera.
candenatensis, Dennst. Schluess. Hort. Malab. 32 = C. Tora.
Candolleana, Vog. Syn. Cass. 42 = bicapsularis.
canescens, H. B. & K. Nov. Gen. et Sp. vi. 357 = emarginata.
capensis, *Thunb. Prod. Pl. Cap.* 79.—Afr. austr.
caracasana, Jacq. Hort. Schoenb. iii. 11 = hirsuta.
cardiosperma, *F. Muell. Fragm.* x. 50.—Austral.
caroliniana, Walt. Fl. Carol. 135 = occidentalis.
carthaginensis, Willd. ex Steud. Nom. ed. II. 304 = bacillaris.
Castiglionia, Collad. Hist. Cass. 121 = C. Chamaecrista.
cathartica, *Mart. Reise, Bras.* i. 548.—Bras.
centijuga, Wawra, in Flora, xlvii. (1864) 248 = verrucosa.
cernua, Balb. Cat. Hort. Taur. (1813) 22 = sulcata.
Chamaecrista, Boj. Hort. Maur. 122 = glandulosa.
Chamaecrista, *Linn. Sp. Pl.* 379.—Am. bor. et austr.
Chamaecrista, Mill. Gard. Dict. ed. VIII. n. 17 = procumbens.
Chamaecrista, Mill. ex Benth. in Trans. Linn. Soc. xxvii. (1871) 576 = virgata.
chamaecristoides, Collad. Hist. Cass. 134 = procumbens, *Linn. Herb.*
Chatelainiana, *Gaudich. in Freyc. Voy. Bot.* 485. t. 111.—Austral.

CASSIA :—

chinensis, Jacq. Coll. i. 64 = C. Sophera.
chinensis, Lam. Encyc. i. 644 = bicapsularis.
choriophylla, *Vog. Syn. Cass.* 56.—Bras.
chrysocarpa, *Desv. Journ. Bot.* iii. (1814) 72.—Bras.
chrysogyne, Miq. in Linnaea, xviii. (1844) 579 = chrysocarpa.
chrysoloma, De Not. Ind. Sem. Hort. Gen. (1840) = bicapsularis.
chrysophylla, A. Rich. Ess. Fl. Cub. 500 = emarginata.
chrysotricha, Collad. Hist. Cass. 99. t. 13 = chrysocarpa.
chrysotricha, Mart. Herb. Fl. Bras. 103 = angulata.
ciliaris, Collad. Hist. Cass. 98 = C. Tagera.
ciliata, Hoffmgg. Verz. Pfl. Nachtr. i. 208 = sericea.
ciliata, Rafin. Fl. Ludov. 100 = occidentalis.
ciliolata, *Benth. in Mart. Fl. Bras.* xv. II. 146.—Bras.
cinerascens, *Vog. Syn. Cass.* 60.—Bras.
cinerea, *Cham. & Schlecht. in Linnaea*, v. (1830) 599. —Mexic.
circinnata, *Benth. in Mitch. Journ. Trop. Austral.* 384.—Austral.
Clausseni, *Benth. in Hook. Journ. Bot.* ii. (1840) 79. —Bras.
coccinea, Wall. Cat. n. 5315 = C. Absus.
Collae, G. Don, Gen. Syst. ii. 442 = bicapsularis.
coluteoides, Collad. Hist. Cass. 102. t. 12 = bicapsularis.
comosa, *Vog. Syn. Cass.* 65.—Afr. austr.
concinna, *Benth. Fl. Austral.* ii. 291.—Austral.
conferta, *Benth. in Mart. Fl. Bras.* xv. II. 142.—Bras.
conjugata, *Ruiz & Pav. ex Benth. in Trans. Linn. Soc.* xxvii. (1871) 540.—Am. trop.
conspicua, Vog. Syn. Cass. 47 = Sieberiana.
contorta, Vog. l. c. 17 = C. Tora.
coquimbensis, *Vog. l. c.* 27.—Chili.
cordistipula, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 164.—Bras.
coriacea, *Bong. ex Benth. in Hook. Journ. Bot.* ii. (1840) 78.—Bras.
corifolia, *Benth. in Mart. Fl. Bras.* xv. II. 120.—Am. trop.
corneliana, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 79.—Afr. trop.
coromandeliana, Jacq. Fragm. 67. t. 100 = C. Sophera.
coronilloides, A. Cunn. ex Benth. in Mitch. Journ. Trop. Austral. 384 = australis.
corymbosa, *Lam. Encyc.* i. 644.—Am. calid.
corymbosa, Orteg. Hort. Matr. Dec. 124 = laevigata.
cotinifolia, *G. Don, Gen. Syst.* ii. 445.—Bras.
Covesii, *A. Gray, in Proc. Am. Acad.* vii. (1868) 399. —Calif.
crassifolia, Orteg. Hort. Matr. Dec. 122 = corymbosa.
crassiramea, *Benth. in Hook. Ic. Pl.* t. 1063.—Bolivia.
crassisepala, Benth. in Linnaea, xxii. (1849) 527 = bicapsularis.
crenulata, *Benth. in Hook. Journ. Bot.* ii. (1840) 80. —Bras.
Crista, Jacq. Coll. i. 82 = biflora.
crotalarioides, *Kunth, Mimos.* 132. t. 40.—Am. austr.
Cruikshanksii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 210.—Chili.
cubensis, Hoffmgg. Verz. Pfl. Nachtr. i. 209 = corymbosa.
cultrifolia, *H. B. & K. Nov. Gen. et Sp.* vi. 363.—Venezuela.
Cumingii, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 211.—Chili.
cuneata, *DC. in Collad. Hist. Cass.* 121.—Am. austr.
cuneata, Griseb. Cat. Pl. Cub. 80 = lineata.
cuneifolia, Vog. Syn. Cass. 49 = viscosa.
curvifolia, *Vog. l. c.* 55.—Bras.
cuspidata, Willd. Enum. Hort. Berol. 443 = pistaciaefolia.
cytisoides, *DC. in Collad. Hist. Cass.* 116. t. 14.—Bras.
dalbergiaefolia, *Benth. in Mart. Fl. Bras.* xv. II. 149. —Am. trop.
debilis, *Vog. Syn. Cass.* 53.—Bras.
decipiens, Desv. Journ. Bot. iii. (1814) 72 = angustifolia.
decipiens, Griseb. Cat. Pl. Cub. 79 = stenophylla.
decrescens, *Benth. in Hook. Journ. Bot.* ii. (1840) 80. —Bras.
decumbens, *Benth. l. c.* 79.—Am. trop.
delagoënsis, *Harv. in Fl. Cap.* ii. 272.—Afr. austr.
densiflora, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 304.—Mexic.

CASSIA :—
densifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 80.—Bras.
dentata, *Vog. Syn. Cass.* 49.—Bras.
Deplanchei, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 554.—N. Caled.
desertorum, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 133.—Bras.
desolata, *F. Muell. in Linnaea*, xxv. (1852) 389.—Austral.
Desvauxii, *Collad. Hist. Cass.* 131.—Am. trop.
didymobotrya, *Fresen. in Flora*, xxii. (1839) 53.—Abyssin.
diffusa, DC. in Mém. Soc. Phys. Genèv. ii. (1824) 130; Prod. ii. 504 = nictitans.
dimidiata, Klein, ex Wall. Cat. n. 5328 = Kleinii.
dimidiata, Roxb. Hort. Beng. 32, nomen = mimosoides.
diphylla, *Linn. Sp. Pl.* 376.—Am. trop.
disadena, Steud. in Flora, xxvi. (1843) II. 760 = glandulosa.
discolor, Desv. Journ. Bot. iii. (1814) 73 = glauca.
dispar, *Willd. Enum. Hort. Berol.* 441.—Am. austr.
disperma, Vell. Fl. Flum. 167; iv. t. 69 = Peltophorum Vogelianum.
distichoclada, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 168.—Am. trop.
divaricata, *Nees, in Syll. Ratisb.* i. (1824) 94.—As. trop.; Malaya.
Dombeyana, Vog. Syn. Cass. 28 = aurantia.
domingensis, *Spreng. Neue Entdeck.* iii. 55.—Ind. occ.
dormiens, Vell. Fl. Flum. 167; iv. t. 67 = bicapsularis.
drepanophylla, *Benth. in Mart. Fl. Bras.* xv. II. 170.—Bras.
dumetorum, Bert. ex DC. Prod. ii. 499 = reticulata.
dysophylla, *Benth. in Mart. Fl. Bras.* xv. II. 117.—Am. trop.
echinocarpa, *Benth. l. c.* 136.—Am. trop.
edulis, *Posada, in Baill. Adansonia*, x. (1872) 187.—N. Granat.
eglandulosa, *Dum.-Cours. Bot. Cult.* ii. VI. 34.—Hab. ?
Ehrenbergii, Bisch. in Bot. Zeit. ii. (1844) 51 = angustifolia.
elegans, H. B. & K. Nov. Gen. et Sp. vi. 342 = laevigata.
elegans, *Voigt, in Syll. Ratisb.* ii. (1828) 55.—Hab. ?
elliptica, H. B. & K. Nov. Gen. et Sp. vi. 356 = emarginata.
elongata, Lemaire-Lis. in Journ. Pharm. vii. (1821) 345 = angustifolia.
emarginata, Bert. ex Steud. Nom. ed. II. i. 304, 305 = atomaria.
emarginata, Clos, in C. Gay, Fl. Chil. ii. 237 = bicapsularis.
emarginata, *Linn. Sp. Pl.* 376.—Am. trop.
emarginata, Mill. Gard. Dict. ed. VIII. n. 13 = pilosa.
enneaphylla, Koen. ex Wight & Arn. Prod. 289 = glauca.
ensiformis, Vell. Fl. Flum. 170; iv. t. 79 = C. Apoucouita.
erecta, Willd. ex Steud. Nom. ed. II. i. 303 = glandulosa.
eremophila, *A. Cunn. ex Vog. Syn. Cass.* 47.—Austral.
ericifolia, *Benth. in Mart. Fl. Bras.* xv. II. 145.—Am. trop.
esculenta, Roxb. Hort. Beng. [31], nomen = C. Sophera.
exaltata, *Reinw. in Blume, Cat. Gew. Buitenz.* 68; *Span. in Linnaea*, xv. (1841) 201.—Ins. Timor.
excelsa, H. B. & K. Nov. Gen. et Sp. vi. 339 = C. Fistula.
excelsa, *Schrad. in Goett. Gel. Anz.* i. (1821) 717.—Bras.
exigua, Roxb. Hort. Beng. [31], nomen = C. Absus.
exsudans, *Benth. in Hook. Journ. Bot.* ii. (1840) 80.—Bras.
fabaginifolia, H. B. & K. Nov. Gen. et Sp. vi. 363 = rotundifolia.
fagifolia, Bertol. Fl. Guatim. 14 = oxyphylla.
fagonioides, Vog. Syn. Cass. 50 = hispidula.
falcata, Dum.-Cours. Bot. Cult. ed. II. vi. 35 = corymbosa.
falcata, Linn. Sp. Pl. 377 = occidentalis.
falcinella, *Oliver, Fl. Trop. Afr.* ii. 281.—Afr. trop.
fasciculata, Michx. Fl. Bor. Am. i. 262 = C. Chamaecrista.

CASSIA :—
fastigiata, Necs, in Flora, iv. (1821) 303 = excelsa.
fastigiata, Vahl, Symb. Bot. iii. 57 = glauca.
fastuosa, *Willd. ex Vog. in Linnaea*, xi. (1837) 654.—Am. trop.
felipensis, H. B. & K. Nov. Gen. et Sp. vi. 368 = C. Chamaecrista.
ferruginea, *Schrad. ex DC. Prod.* ii. 489.—Am. trop.
Fieldii, Colla, Hort. Ripul. App. iv. 23. t. 11 = australis.
filicifolia, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 148.—Bras.
filipendula, Boj. Hort. Maurit. 122 = mimosoides.
filipes, Benth. in Hook. Journ. Bot. ii. (1840) 81 = rotundifolia.
fimbriata, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 9.—Malaya.
Fistula, Herbb. ex Oliver, Fl. Trop. Afr. 271 = C. Arerch.
Fistula, *Linn. Sp. Pl.* 377.—As. trop.
fistuloides, Collad. Hist. Cass. 87. t. 1 = C. Fistula.
flaccida, Clos, in C. Gay, Fl. Chil. ii. 238 = coquimbensis.
flavicoma, Benth. Pl. Hartw. 171 = tristicula.
flavicoma, *H. B. & K. Nov. Gen. et Sp.* vi. 366.—Venezuela.
flexuosa, A. DC. Not. 7^mo^ Pl. Rar. Jard. Genèv. 34. t. 4 = bicapsularis.
flexuosa, Burm. f. Fl. Cap. Prod. 12 = capensis.
flexuosa, *Linn. Sp. Pl.* 379.—Bras.
flexuosa, Mill. Gard. Dict. ed. VIII. n. 16 = serpens.
floribunda, Cav. Desc. 132 = laevigata.
floribunda, Collad. Hist. Cass. 88. t. 11 = corymbosa.
floribunda, Griseb. ex Benth. in Trans. Linn. Soc. xxvii. (1871) 549 = racemosa.
florida, Vahl, Symb. Bot. iii. 57 = siamea.
fluminensis, Vell. Fl. Flum. 168; iv. t. 72 = Dimorphandra exaltata.
Fockeana, Miq. in Linnaea, xviii. (1844) 579 = bacillaris.
foetida, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 443 = stipulacea.
foetida, Salisb. Prod. 325 = C. Tora.
foetida, Willem. in Usteri, Ann. Bot. xviii. (1796) 32 = occidentalis.
foetidissima, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 441 = mexicana.
foliolosa, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 544.—Guatemala.
foliolosa, Steud. Nom. ed. II. i. 305 = seq.
foliosa, G. Don, Gen. Syst. ii. 746 = tenella ?
foliosa, *Phil. in Anal. Univ. Chil.* (1862) II. 379.—Chili.
Fraseri, A. Cunn. ex Vog. Syn. Cass. 48 = australis.
fraxinifolia, H. B. & K. Nov. Gen. et Sp. vi. 349 = pistaciaefolia.
frondosa, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 210 = bicapsularis.
frondosa, [Soland. in] Ait. Hort. Kew. ed. I. ii. 53 = biflora.
frutescens, Mill. Gard. Dict. ed. VIII. n. 2 = C. Sophera.
fruticescens, Bert. ex Spreng. Syst. ii. 335 = viminea.
fruticosa, Mill. Gard. Dict. ed. VIII. n. 10 = bacillaris.
fulgens, Macfad. Fl. Jamaic. i. 342 = biflora.
fulgens, Wall. Cat. n. 5310 = multijuga.
fuscescens, *Benth. in Mart. Fl. Bras.* xv. II. 153.—Am. trop.
galegaefolia, Linn. Syst. Nat. ed. X. 1017 = biflora.
Galeottiana, *Martens, in Bull. Acad. Brux.* x. II. (1843) 305.—Mexic.
gallinaria, Collad. Hist. Cass. 96 = C. Tora.
Gardneri, *Benth. in Mart. Fl. Bras.* xv. II. 120.—Bras.
Gaudichaudii, *Hook. & Arn. Bot. Beech. Voy.* 81.—Ins. Sandvic.
gemina, Vell. Fl. Flum. 169; iv. t. 74 = C. Chamaecrista.
geminata, *Benth. in Mart. Fl. Bras.* xv. II. 135.—Bras.
geminata, Vahl, ex DC. Prod. ii. 505 = mimosoides.
geminiflora, *Collad. Hist. Cass.* t. 3.—Mexic.
geminiflora, Moç. & Sesse, in Collad. Hist. Cass. 103. t. 3 = biflora.
geminiflora, Schrank, Hort. Monac. t. 26 = occidentalis.
geniculata, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 440 = villosa.

CASSIA :—
gigantea, Bert. ex DC. Prod. ii. 492 = siamea.
glabra, Collad. Hist. Cass. 131 = cytisoides.
glandulifera, Reinw. ex Steud. Nom. ed. II. i. 305 = seq.
glanduliflora, Reinw. in Blume, Cat. Gew. Buitenz. 68 = bicapsularis.
glandulosa, Hook. Bot. Mag. t. 3435 = virgata.
glandulosa, *Linn. Syst. Nat.* ed. X. 1017.—Am. trop.
glauca, *Lam. Encyc.* i. 647.—As. trop.; Austral.; Polynes.
glaucescens, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 539.—Chili.
glaucescens, *Hoffmgg. Verz. Pfl. Nachtr.* i. 209.—Cuba.
glutinosa, *DC. Prod.* ii. 495.—Austral.
goënsis, *Dalz. in Hook. Kew Journ.* iv. (1852) 112.—Ind. or.
gonioides, *A. Cunn. ex Benth. in Hook. Ic. Pl.* xi. 48. t. 1061.—Austral.
gonoclada, *Benth. in Mart. Fl. Bras.* xv. II. 169.—Bras.
gonoclada, Benth. in Trans. Linn. Soc. xxvii. (1871) 572 = choriophylla.
goratensis, *Fresen. in Flora*, xxii. (1839) 53.—Abyssin.
gracilis, *Kunth, Mimos.* 120. t. 36.—Am. trop.
gracillima, *Benth. in Mart. Fl. Bras.* xv. II. 143.—Bras.
gracillima, Welw. Apont. 590 = mimosoides.
grammica, *Spreng. Neue Entdeck.* iii. 55.—Am. trop.
grandiflora, Desf. Cat. Hort. Par. ed. III. 302 = laevigata.
grandiflora, Pers. Syn. i. 457 = bicapsularis.
grandis, *Linn. f. Suppl.* 230.—Panama.
grandistipula, Vog. Syn. Cass. 58 = rotundata.
Grantii, *Oliver, Fl. Trop. Afr.* ii. 279.—Afr. trop.
graveolens, Colla, Hort. Ripul. App. ii. 343 = atomaria.
Greggii, *A. Gray, Pl. Wright.* i. 59.—Mexic.
grisea, A. Rich. Ess. Fl. Cub. 493 = emarginata.
guineensis, G. Don, Gen. Syst. ii. 449 = mimosoides.
Hartwegi, Benth. Pl. Hartw. 117 = oxyphylla.
hecatophylla, DC. in Collad. Hist. Cass. 124. t. 18 = mimosoides.
hedysaroides, *Vog. Syn. Cass.* 52.—Bras.
heptanthera, *F. Muell. Fragm.* x. 8.—Austral.
Herbertiana, Lindl. Bot. Reg. t. 1422 = laevigata.
herpetica, Jacq. Obs. Bot. ii. 24. t. 45. f. 2 = alata.
heteroloba, Lindl. in Mitch. Three Exped. Austral. ed. II. ii. (1839) 122 = eremophila.
heterophylla, Walp. Rep. i. 838, sphalm. = mimosoides.
Hibbertiana, Steud. Nom. ed. II. i. 305, sphalm. = Herbertiana.
Hilariana, *Benth. in Mart. Fl. Bras.* xv. II. 108.—Am. trop.
Hildebrandtii, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 80.—Afr. trop.
hirsuta, *Linn. Sp. Pl.* 378.—Am. trop.
hirsuta, Vell. Fl. Flum. 170; iv. t. 80 = cathartica.
hirta, *Willd. Enum. Hort. Berol. Suppl.* 23.—Hab. ?
hispida, Collad. Hist. Cass. 118 = hispidula.
hispidula, *Vahl, Eclog. Am.* iii. 10.—Mexic.
Hoffmanseggii, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 104.—Bras.
holosericea, *Fresen. in Flora*, xxii. (1839) I. 54.—Abyssin.
homophylla, *Hoffmgg. Verz. Pfl. Nachtr.* i. 209.—Hab. ?
Hookeriana, *Gill. in Hook. Bot. Misc.* iii. (1833) 210.—Chili.
Hornemanni, DC. Prod. ii. 507 = C. Chamaecrista.
Horsfieldii, Miq. Fl. Ind. Bat. i. 99 = glauca.
Houstoniana, Collad. Hist. Cass. 132 = *biflora*, Mill.
Humboldtiana, DC. Prod. ii. 489 = spectabilis.
humilis, Collad. Hist. Cass. 96 = C. Tora.
humilis, Willd. ex Steud. Nom. ed. II. i. 305 = C. Tagera.
hybrida, Tenore. Cat. Hort. Neap. 80 = laevigata.
hymeneaefolia, *Benth. in Mart. Fl. Bras.* xv. II. 130.—Am. trop.
hypnotica, Vell. Fl. Flum. 169; iv. t. 75 = C. Chamaecrista.
hypoleuca, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 117.—Bras.
inaequilatera, *Balb. ex DC. Prod.* ii. 490.—N. Granat.
incana, *Vog. Syn. Cass.* 52.—Bras.

CASSIA :—
incarnata, *Pav. ex Benth. in Trans. Linn. Soc.* xxvii. (1871) 545.—Peruv.
incurvata, *Benth. in Mart. Fl. Bras.* xv. II. 137.—Am. trop.
indecora, H. B. & K. Nov. Gen. et Sp. vi. 313 = bicapsularis.
indica, Poir. Encyc. Suppl. ii. 126 = C. Sophera.
inflata, Spreng. Syst. ii. 336 = bicapsularis.
Isidorea, *Benth. in Mart. Fl. Bras.* xv. II. 154.—Bras.
italica, Lam. ex Steud. Nom. ed. I. 166, 167 = C. Senna.
itambana, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 151.—Am. trop.
Jacobinea, *Benth. in Mart. l. c.* 132.—Am. trop.
javanica, Aubl. Pl. Gui. i. 382 = grandis.
javanica, *Linn. Sp. Pl.* 379.—Malaya.
javanica, Sieber, ex Benth. in Trans. Linn. Soc. xxvii. (1871) 517 = Sieberiana.
javanica, Vell. Fl. Flum. 168; iv. t. 73 = ferruginea.
Johannae, *Vatke, in Oestr. Bot Zeitschr.* xxx. (1880) 77.—Afr. trop.
Ketschta, *Hasselq. It. Palaest.* 469.—Palaest.
Kirkii, *Oliver, Fl. Trop. Afr.* ii. 281.—Afr. trop.
kituiensis, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 81.—Afr. trop.
Kleinii, *Wight & Arn. Prod.* 293.—Ind. or.; Malaya.
Kotschyana, *Oliver, Fl. Trop. Afr.* ii. 271.—Afr. trop.
Kunthiana, Cham. & Schlecht. in Linnaea, v. (1830) 598 = C. Tagera.
lactea, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 78.—Madag.
laeta, *H. B. & K. Nov. Gen. et Sp.* vi. 340.—N. Granat.
laevigata, *Willd. Enum. Hort. Berol.* 441.—Cosmop. trop.
lamprosperma, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 146.—Am. trop.
lanceolata, Collad. Hist. Cass. 93. t. 15 = acutifolia.
lanceolata, Forsk. Fl. Aegypt. Arab. 85 = C. Sophera.
lanceolata, Pers. Syn. i. 456 = uniflora.
lanceolata, Wight & Arn. Prod. 288 = angustifolia.
Langsdorffii, *Kunth, ex Vog. Syn. Cass.* 55.—Bras.
latifolia, *G. F. W. Mey. Prim. Fl. Esseq.* 166.—Guiana.
latistipula, *Benth. in Mart. Fl. Bras.* xv. II. 156.—Bras.
latopetiolata, *Domb. ex Vog. Syn. Cass.* 70.—Peruv.
lavradioides, *Benth. in Mart. Fl. Bras.* xv. II. 137.—Am. trop.
laxiflora, *Benth. Fl. Austral.* ii. 283.—Austral.
leiandra, *Benth. in Mart. Fl. Bras.* xv. II. 94.—Bras.
leiantha, Benth. in Hook. Journ. Bot. ii. (1840) 78 = hispidula.
leiophylla, *Vog. Syn. Cass.* 25.—Bras.
lenitiva, Bisch. in Bot. Zeit. viii. (1850) 885 = acutifolia.
lentiscifolia, *Benth. in Mart. Fl. Bras.* xv. II. 144.—Am. trop.
leptocarpa, *Benth. in Linnaea*, xxii. (1849) 528.—N. Mexic.
leptoclada, *Benth. Fl. Austral.* ii. 290.—Austral.
leptophylla, *Vog. Syn. Cass.* 13.—Bras.
Leschenaultiana, DC. in Mém. Soc. Phys. Genèv. ii. (1824) 132 = mimosoides.
Leschenaultii, Wall. Cat. n. 5325 = mimosoides.
leucoxylon, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1207, nomen.—Guiana.
Liebmannii, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 549.—Mexic.
ligustrina, Forsk. ex Benth. l. c. 533 = C. Sophera.
ligustrina, *Linn. Sp. Pl.* 378.—Am. trop.
ligustrina, Mill. Gard. Dict. ed. VIII. n. 12 = obovata.
liguştrinoides, Schrank, in Denkschr. Akad. Muench. vi. (1816–17) 179 = angustifolia.
limensis, Lam. Encyc. i. 643 = bicapsularis.
Lindheimeriana, *Schlecht. in Linnaea*, xxi. (1848) 457.—Texas.
Lindleyana, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 341 = multijuga.
linearifolia, *G. Don, Gen. Syst.* ii. 446.—Bras.
linearis, Michx. Fl. Bor. Am. i. 261 = occidentalis.
lineata, *Sw. Prod. Veg. Ind. Occ.* 66.—Am. trop.

CASSIA :—
lomatopoda, *Benth. in Mart. Fl. Bras.* xv. II. 146.—Bras.
longeglandulosa, *Benth. l. c.* 119.—Am. trop.
longicuspis, *Benth. l. c.* 150.—Am. trop.
longifolia, Steud. Nom. ed. II. i. 304, in syn. = angustifolia.
longiracemosa, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 80.—Afr. trop.
longisiliqua, Blanco, Fl. Filip. ed. I. 338 = hirsuta.
longisiliqua, Linn. f. Suppl. 230 = occidentalis.
Lorentzii, *Niederl. in Roca, Exped. Rio Negro*, ii. 210.—Patagon.
lotoides, H. B. & K. Nov. Gen. et Sp. vi. 361 = hispidula.
lucens, Vog. Syn. Cass. 46 = racemosa.
lucida, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 18.—Hab.?
Lundii, *Benth. in Mart. Fl. Bras.* xv. II. 143.—Am. trop.
lutescens, G. Don, Gen. Syst. ii. 442 = tomentosa.
machaeriifolia, *Benth. in Mart. Fl. Bras.* xv. II. 151.—Am. trop.
macradenia, *Collad. Hist. Cass.* 132.—Bras.
macranthera, *DC. in Collad. l. c.* 99. t. 8.—Bras.
macrocarpa, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 43.—Paraguay.
macrophylla, *Kunth, Mimos.* 126. t. 38.—N. Granat.
macrophylla, Ruiz & Pav. ex Benth. in Trans. Linn. Soc. xxvii. (1871) 520 = Ruiziana.
madagascariensis, *Boj. Hort. Maurit.* 120.—Madag.
madagascariensis, Poir. ex Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 424 = Swartzia madagascariensis.
magnifica, Mart. Herb. Fl. Bras. 106 = multijuga.
magnifolia, *F. Muell. Fragm.* i. 166.—Austral.
malacophylla, *Vog. Syn. Cass.* 55.—Bras.
Mandoni, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 540.—Peruv.
Mannii, *Oliver, Fl. Trop. Afr.* ii. 272.—Afr. trop.
marginata, *Roxb. Hort. Beng.* [31]; *Fl. Ind.* ii. 338.—Ind. or.
marginata, Willd. Enum. Hort. Berol. 443 = polyphylla.
Marimari, Aubl. Pl. Gui. 382 = biflora.
maritima, Willd. ex Vog. Syn. Cass. 23 = pilifera.
Martiana, *Benth. in Mart. Fl. Bras.* xv. II. 127.—Am. trop.
marylandica, *Linn. Sp. Pl.* 378.—Am. bor.
medellinensis, *Posado, in Baill. Adansonia*, x. (1872) 187.—N. Granat.
medica, Forsk. Fl. Aegypt. Arab. p. cxi = angustifolia.
medica, Vell. Fl. Flum. 166; iv. t. 62 = quinquangulata.
medicinalis, Bisch. in Bot. Zeit. viii. (1850) 887 = angustifolia.
megalantha, *Decne. Herb. Tim. Descr.* 136.—Ins. Timor.
melanocarpa, Bert. ex DC. Prod. ii. 491 = viminea.
mexicana, *Jacq. Hort. Schoenbr.* ii. t. 203.—Am. trop.
micans, Nees, in Flora, iv. (1821) 303 = speciosa.
micrantha, Guill. & Perr. Fl. Seneg. Tent. i. 262 = nigricans.
microphylla, Willd. Sp. Pl. ii. 529 = mimosoides.
Milleri, Collad. Hist. Cass. 132 = pilosa.
Mimosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 9.—Malaya.
mimosoides, DC. Prod. ii. 503 = Kleinii.
mimosoides, *Linn. Sp. Pl.* 379.—Cosmop. trop.
misera, *Phil. Fl. Atac.* 17.—Chili.
mollis, Vahl, Symb. Bot. iii. 57 = grandis.
mollissima, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 440 = atomaria.
monaden, Vell. Fl. Flum. 166; iv. t. 63 = splendida.
monantha, DC. Prod. ii. 506 = sericea.
monophylla, Vell. Fl. Flum. 166; iv. t. 61 = rotundifolia.
montana, *Heyne, in Roth, Nov. Sp. Pl.* 214.—Ind. or.
montana, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 71 = timoriensis.
montana, *Vell. Fl. Flum.* 169; iv. t. 77.—Bras.
monticola, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 134.—Am. trop.
moschata, Benth. in Hook. Journ. Bot. ii. (1840) 75 = leiandra.
moschata, *H. B. & K. Nov. Gen. et Sp.* vi. 358.—Am. austr.

CASSIA :—
mucronata, *Spreng. Syst.* ii. 341.—Bras.
mucronifera, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 116.—Bras.
multiflora, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 307.—Am. trop.
multiflora, Scheele, in Linnaea, xvii. (1843) 337 = excelsa.
multiflora, Vog. Syn. Cass. 39 = macranthera.
multiglandulosa, Jacq. Ic. Rar. i. t. 72 = tomentosa.
multijuja, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 108.—Guiana.
multinervia, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 166.—Am. trop.
multiseta, *Benth. l. c.* 141.—Am. trop.
multisiliqua, J. F. Gmel. Syst. 1521 (quid?).
Mutisiana, *Kunth, Mimos.* 142. t. 43.—Am. trop.
myriophylla, Wall. Cat. n. 5326 = mimosoides.
nana, *Benth. in Mart. Fl. Bras.* xv. II. 110.—Am. trop.
Neesiana, *Mart. ex Benth. l. c.* 148.—Bras.
neglecta, *Vog. in Linnaea*, x. (1836) 394.—Bras.
nemophila, Walp. Rep. i. 827, sphalm. = eremophila.
nemorosa, H. B. & K. Nov. Gen. et Sp. vi. 358 = biflora.
nervosa, Vog. Syn. Cass. 39 = speciosa.
nicaraguensis, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 552.—Am. trop.
nictitans, *Linn. Sp. Pl.* 380.—Am. bor. et austr.
nictitans, Wydl. ex Steud. Nom. ed. II. i. 303, 306 = mimosoides.
nigricans, *Vahl, Symb. Bot.* i. 30.—Geront. trop.
nitida, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = viminea.
nodosa, *Buch.-Ham. ex Roxb. Hort. Beng.* [31]; *Fl. Ind.* ii. 336.—As. trop.
notabilis, *F. Muell. Fragm.* iii. 28.—Austral.
nummularia, Vahl, ex DC. Prod. ii. 501 = rotundifolia.
nummulariaefolia, *Benth. in Mart. Fl. Bras.* xv. II. 143.—Bras.
nutans, Collad. Hist. Cass. 113. t. 4 = atomaria.
obcordata, Sw. ex Wickstr. in Vet. Acad. Handl. (1825) 429 = lineata.
obliqua, *Ruiz & Pav. ex G. Don, Gen. Syst.* ii. 451.—Peruv.
obliquifolia, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1822) 40 = occidentalis.
oblongifolia, *Vog. Syn. Cass.* 23.—Bras.
obovata, *Collad. Hist. Cass.* 92. t. 15.—Geront. trop.
obtecta, *Benth. in Mart. Fl. Bras.* xv. II. 136.—Am. trop.
obtusa, Clos, in C. Gay, Fl. Chil. ii. 235 = bicapsularis.
obtusa, Roxb. Hort. Beng. [31]; Wight, Ic. t. 757 = obovata.
obtusata, Hayne, Arzneig. ix. t. 43 = obovata.
obtusata, Hochst. & Steud. Nom. ed. II. i. 306 = holosericea.
obtusifolia, Linn. Sp. Pl. 377 = C. Tora.
occidentalis, Hort. ex Steud. Nom. ed. II. i. 306 = homophylla.
occidentalis, *Linn. Sp. Pl.* 377.—Cosmop. trop.
occidentalis, Naves, in Blanco, Fl. Philipp. ed. III. t. 73 = C. Sophera.
ochnacea, *Vog. Syn. Cass.* 51.—Bras.
odorata, *Morris, Fl. Conspic.* t. 57.—Austral.
olesiphylla, *Vog. Syn. Cass.* 68.—Bras.
oligoclada, *F. Muell. Fragm.* iii. 49.—Austral.
oligophylla, *F. Muell. l. c.*—Austral.
oligosperma, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 147.—Bras.
opaca, R. Grah. ex Vog. Syn. Cass. 42 = stipulacea.
orbiculata, *Benth. in Hook. Journ. Bot.* ii. (1840) 79.—Bras.
orinocensis, *Spruce, ex Benth. in Mart. Fl. Bras.* xv. II. 150.—Bras.
orientalis, Pers. Syn. i. 457 = acutifolia.
ornata, Bernh. ex Benth. in Trans. Linn. Soc. xxvii. (1871) 522 = latifolia.
ornata, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1104, nomen.—Guiana.
ornithopodioides, Steud. Nom. ed. I. 166 = seq.
ornithopoides, Lam. Encyc. i. 644 = sericea.
Otterbeynii, G. F. W. Mey. Prim. Fl. Esseq. 169 = glandulosa.
ovalifolia, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 305 = bicapsularis.

CASSIA :—
ovata, *Mérat & Lens, ex Geiger, Pharm. Bot.* ed. II. ii. 1125.—Arab.
oxyadena, DC. Prod. ii. 495 = biflora.
oxyphylla, *Kunth, Mimos.* 129. t. 39.—N. Granat.
pachycalyx, Vog. Syn. Cass. 52 = cathartica.
pachyphylla, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 166.—Bras.
pallida, Vahl, Eclog. Am. iii. 12 = biflora.
palmata, Wall. Cat. n. 5306 = timoriensis.
paniculata, *Benth. in Mart. Fl. Bras.* xv. II. 153.—Am. trop.
paposana, *Phil. Fl. Atac.* 17.—Chili.
papulosa, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 81.—Bras.
paradyction, *Vog. Syn. Cass.* 45.—Bras.
paraguensis, *Micheli, in Mém. Soc. Phys. Genèv.* xxviii. (1883) n. VII. 44.—Paraguay.
Parahyba, Vell. Fl. Flum. 168 ; iv. t. 71 = Schizolobium excelsum.
paramariboënsis, *Miq. in Bot. Zeit.* i. (1843) 461.—Guian.
paratyensis, *Vell. Fl. Flum.* 168 ; iv. t. 70.—Bras.
Parkeriana, DC. Prod. ii. 504 = riparia.
parvifolia, Wender. in Syll. Ratisb. i. (1824) 65 = glandulosa.
parvistipula, *Benth. in Mart. Fl. Bras.* xv. II. 170.—Am. trop.
pascuorum, *Mart. ex Benth. l. c.* 165.—Am. trop.
patellaria, *DC. in Collad. Hist. Cass.* 125. t. 16.—Am. trop.
patula, [Soland. in] Ait. Hort. Kew. ed. I. ii. 51 = C. Sophera.
pauciflora, *H. B. & K. Nov. Gen. et Sp.* vi. 360.—Am. trop.
Pavoniana, G. Don, Gen. Syst. ii. 447 = glandulosa.
Pearcii, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 552.—Bolivia.
pedicellaris, *DC. Prod.* ii. 504.—Ins. S. Doming.
pendula, Willd. Enum. Hort. Berol. 440 = bicapsularis.
pentagonia, *Mill. Gard. Dict.* ed. VIII. n. 18.—Am. trop.
pentandra, Raddi, in Mem. Mod. xviii. Fis. (1820) 399 = rotundifolia.
peralteana, *H. B. & K. Nov. Gen. et Sp.* vi. 356.—Mexic.
Persoonii, Collad. Hist. Cass. 119 = uniflora.
peruviana, Vog. Syn. Cass. 39 = obliqua.
Petersiana, Bolle, in Peters, Reise Mossamb. Bot. 13 = delagoënsis.
phyllodinea, *R. Br. App. Sturt, Centr. Austral.* 15.—Austral.
phyllostachya, *Benth. in Mart. Fl. Bras.* xv. II. 150.—Am. trop.
picta, *G. Don, Gen. Syst.* ii. 444.—Ecuador ; Ins. Galap.
pilidens, Steud. Nom. ed. II. i. 307 = nictitans.
pilifera, *Vog. Syn. Cass.* 23.—Bras.
pilosa, *Linn. Syst. Nat.* ed. X. 1017.—Am. trop.
pinnata, Hort. ex Voigt, in Syll. Ratisb. ii. (1828) 55 = elegans.
pistaciaefolia, *H. B. & K. Nov. Gen. et Sp.* vi. 349.—N. Granat.
planisiliqua, Burm. f. Fl. Ind. 96 = Albizzia Lebbek.
planisiliqua, Linn. Sp. Pl. 377 = occidentalis.
planisiliqua, *Lam. Encyc.* i. 645.—Ins. Guadelup.
platypoda, R. Br. in Sturt, Centr. Austr. ii. 78 = eremophila.
pleurocarpa, *F. Muell. Fragm.* i. 223.—Austral.
Plumieri, *DC. Prod.* ii. 506.—Ins. Guadelup.
plumosa, Vog. Syn. Cass. 64 = mimosoides.
podocarpa, *Guill. & Perr. Fl. Seneg. Tent.* 259.—Afr. trop.
Pohliana, *Benth. in Hook. Journ. Bot.* ii. (1840) 80.—Bras.
polyadena, *DC. Pl. Rar. Jard. Genèv.* 2e *Rapp.* 12.—Ind. occ.
polyantha, *Moç. & Sesse, ex Collad. Hist. Cass.* 112. t. 2.—Mexic.
polyphylla, *Jacq. Coll.* iv. 104 ; *Ic. Rar.* t. 460.—Ind. occ.
polystachya, *Benth. in Hook. Journ. Bot.* ii. (1840) 77.—Guiana.
porturegalensis, Bancroft, ex Macf. Fl. Jamaic. i. 340 = obovata.
Potentilla, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 168.—Am. trop.
praetexta, *Vog. Syn. Cass.* 67.—Am. trop.
procumbens, Forsk. Fl. Aegypt. Arab. p. cxi = nigricans.
procumbens, Linn. Sp. Pl. ed. I. 380 = nictitans.

CASSIA :—
procumbens, *Linn. Sp. Pl.* ed. II. 543, pro parte, et Herb.—Am. bor.
procumbens, Mill. Gard. Dict. ed. VIII. n. 20 = hispidula.
procumbens, Thunb. Fl. Jap. 179 = mimosoides.
prominens, D. Dietr. Syn. Pl. ii. 1479 = macranthera.
propinqua, Cham. & Schlecht. in Linnaea, v. (1830) 599 = tristicula.
propinqua, H. B. & K. Nov. Gen. et Sp. vi. 369 = glandulosa.
prostrata, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 441 = serpens.
prostrata, Koen. ex Roxb. Hort. Beng. 32 = pumila.
pruinosa, *F. Muell. Fragm.* iii. 48.—Austral.
pseudo-nictitans, Hance, in Journ. Linn. Soc. xi. (1871) 79 = nictitans.
psammophila, Benth. in Trans. Linn. Soc. xxvii. (1871) 571 = cordistipula.
psilocarpa, Welw. Apont. 587. n. 40 = Siebcriana.
puberula, H. B. & K. Nov. Gen. et Sp. vi. 341 = bacillaris.
pubescens, R. Br. in Salt, Abyss. App. 64 = holosericea.
pubescens, Ehrenb. & Hempr. ex Benth. in Proc. Linn. Soc. ii. (1855) 283 = holosericea.
pubescens, *Jacq. Fragm.* 46. t. 57.—Bras.
pubescens, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 442 = tomentosa.
pubigera, Lag. Gen. et Sp. Nov. 14 = hirsuta.
pudibunda, *Mart. ex Benth. in Mart. Fl. Bras*, xv. II. 102.—Bras.
pulchella, *Boj. Hort. Maurit.* 122.—Ins. Maurit.
pulchella, Salisb. Prod. 326 = C. Chamaecrista.
pulcherrima, *Dehn. Rivista Nap.* i. III. 174.—Guiana.
pulchra, H. B. & K. Nov. Gen. et Sp. vi. 362 = Desvauxii.
pumila, *Lam. Encyc.* i. 651.—Cosmop. trop.
pumila, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 307 = leiophylla.
pumila, F. Muell. Fragm. iii. 47 = concinna.
Pumilio, *A. Gray, Pl. Lindh.* 180.—Am. bor. occ.
punctata, Scheele, in Linnaea, xvii. (1843) 336 = sylvestris.
punctata, *Vog. Syn. Cass.* 51.—Bras.
punctulata, Hook. & Arn. Bot. Beech. Voy. 420 = pauciflora.
purgans, *Steud. Nom.* ed. II. i. 307.—Chili.
purpurea, Roxb. Hort. Beng. [31] = C. Sophera.
pygmaea, DC. Prod. ii. 502 = procumbens, *Linn. Herb.*
quadrangularis, Zoll. & Mor. Syst. Verz. Zoll. 2 = laevigata.
quadriflora, Vog. Syn. Cass. 47 = glutinosa.
Quebracho, Steud. '1833' ex Nom. ed. II. i. 307, 304 = bicapsularis.
quinquangulata, *Rich. in Act. Soc. Hist. Nat. Par.* (1792) 108.—Am. trop.
racemosa, *Mill. Gard. Dict.* ed. VIII. n. 19.—Bras.
ramiflora, Vog. Syn. Cass. 48 = C. Apoucouita.
ramosa, Vog. l. c. 55 = uniflora.
ramosissima, H. B. & K. Nov. Gen. et Sp. vi. 366 = glandulosa.
reflexa, Salisb. Prod. 326 = marylandica.
Reinwardtii, *Hassk. in Hoev. & De Vriese, Tijdschr.* v. (1838-39) 269.—Java.
reniformis, *G. Don, Gen. Syst.* ii. 440.—Bras.
renigera, *Wall. Cat.* n. 5307.—Burma.
repens, *Vog. Syn. Cass.* 60.—Bras.
reticulata, *Willd. Enum. Hort. Berol.* 443.—Am. austr.
retusa, *Soland. ex Vog. in Linnaea*, xv. (1841) 72.—Austral.
revoluta, F. Muell. in Trans. Vict. Inst. (1855) 120 = australis.
rhombifolia, Roxb. Hort. Beng. [31] ; Fl. Ind. ii. 334 = C. Fistula.
Richardiana, Kunth, Mimos. 139. t. 142 = multijuga.
Riedelii, *Benth. in Mart. Fl. Bras.* xv. II. 122.—Am. trop.
rigidifolia, *Benth. l. c.* 142.—Bras.
riparia, *H. B. & K. Nov. Gen. et Sp.* vi. 369.—Am. trop.
robiniaefolia, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 545.—Am. trop.
robinioides, Willd. Enum. Hort. Berol. 443 = C. Sophera.
Roemeriana, *Scheele, in Linnaea*, xvi. (1842) 457.—Texas.

CASSIA :—
Roraimae, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 574.—Guiana.
rostrata, Voigt, in Syll. Ratisb. ii. (1828) 55 = biflora.
rotundata, *Vog. Syn. Cass.* 58.—Bras.
rotundifolia, *Pers. Syn.* i. 456.—Mexic.
Roxburghiana, R. Grah. in Wall. Cat. n. 5323 = mimosoides.
Roxburghii, DC. Prod. ii. 489 = marginata.
Roxburghii, Steud. Nom. ed. II. i. 307, sphalm. = *Roxburghiana*.
rufa, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 306.—Mexic.
rugosa, *G. Don, Gen. Syst.* ii. 440.—Bras.
Ruiziana, *Vog. Syn. Cass.* 40.—Am. trop.
rupicola, Benth. in Trans. Linn. Soc. xxvii. (1871) 574 = olesiphylla.
rusa, Mart. & Gal. in Bull. Acad. Brux. x. II. (1848) 306 = tristicula.
ruscifolia, Jacq. Ic. Rar. i. t. 71 = C. Sophera.
Sabak, *Delile, Cent. Pl. Afr.* 28.—Aegypt.
Sagera, Lam. Encyc. i. 643 = C. Tora.
Sagotiana, *Benth. in Mart. Fl. Bras.* xv. II. 93.—Guiana.
Saltiana, Steud. Nom. ed. II. i. 307 = holosericea.
sapindifolia, *Vog. Syn. Cass.* 34.—Bras.
savannensis, Miq. in Bot. Zeit. i. (1843) 461 = uniflora.
scabra, *Pohl, ex Benth. in Mart. Fl. Bras.* xv. II. 135.—Am. trop.
scabrella, *Hoffmgg. Verz. Pfl. Nachtr.* ii. 81.—Bras.
scandens, *Ruiz & Pav. ex G. Don, Gen. Syst.* ii. 440.—Ecuador.
Schimperi, Steud. Nom. ed. II. i. 307 = holosericea.
schinifolia, A. DC. 7e Not. Pl. Rar. Hort. Genèv. 35 = C. Sophera.
Schlimii, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 571.—Am. trop.
Schomburgkii, *Klotzsch, ex Schomb. Faun. & Fl. Gui.* 1207, nomen.—Guiana.
Schultesii, Colla, Hort. Ripul. App. ii. 343 = australis.
sclerocarpa, *Vog. Syn. Cass.* 15.—Bras.
secunda, *Benth. in Mart. Fl. Bras.* xv. II. 148.—Bras.
Selloi, G. Don, Gen. Syst. ii. 442 = multijuga.
Selloi, Spreng. ex Vog. Syn. Cass. 39 = speciosa
semifalcata, Vell. Fl. Flum. 167 ; iv. t. 68 = multijuga.
semperflorens, DC. Cat. Hort. Monsp. 90 = biflora.
Senna, Linn. Sp. Pl. 377 = obovata, acutifolia.
sennoides, Jacq. Coll. i. 74 = bicapsularis.
sensitiva, Jacq. l. c. ii. 367 = sericea.
sensitiva, Roxb. Hort. Beng. 32 = mimosoides.
septemtrionalis, *Viv. Elench. Pl. Hort. Dinegro*, 14.—Hab. ?
septentrionalis, Zuccagni, in Roem. Collect. 141 = laevigata.
sericea, *Sw. Fl. Ind. Occ.* 724.—Am. trop.
serpens, Griseb. Cat. Pl. Cub. 80 = strigillosa.
serpens, *Linn. Syst. Nat.* ed. X. 1018.—Am. trop.
sessilis, Benth. in Mart. Fl. Bras. xv. II. 135, in syn. = andromedea.
setigera, DC. Prod. ii. 499 = montana.
setosa, *Vog. Syn. Cass.* 51.—Bras.
siamea, *Lam. Encyc.* i. 648.—Ind. or. ; Malaya.
Sieberi, Presl, in Oken, Isis, xxi. (1828) 272 = glauca.
Sieberiana, *DC. Prod.* ii. 489.—Afr. trop.
silvestris, *Vell. Fl. Flum.* 169 ; iv. t. 78.—Bras.
simplicifolia, Desv. Journ. Bot. iii. (1814) 72 (= *Delaria ovalifolia*).—Bras.
sinensis, J. F. Gmel. Syst. 669 = chinensis.
singueana, Delile, in Caill. Voy. Meroë, Bot. iv. 27 (Cent. Pl. Afr. 28) = goratensis.
Skinneri, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 542.—Guatemal.
smaragdina, Macfad. Fl. Jamaic. i. 347 = nictitans.
Sophera, *Linn. Sp. Pl.* 379.—Geront. trop.
Sophera, Wall. Cat. n. 5317 = occidentalis.
sopheroides, Collad. Hist. Cass. 133 = C. Sophera.
Sophora, Auct. = C. Sophera.
sophoroides, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 152.—Am. trop.
speciosa, H. B. & K. Nov. Gen. et Sp. vi. 358 = spectabilis.
speciosa, Roxb. Hort. Beng. [31], nomen = glauca.
speciosa, *Schrad. in Gott. Gel. Anz.* i. (1821) 718.—Bras.
spectabilis, *DC. Cat. Hort. Monsp.* 90.—Am. trop.

CASSIA:—
spinescens, *Hoffmgg. ex Vog. Syn. Cass.* 27.—Bras.
splendida, *Vog. l. c.* 17.—Bras.
Spruceana, *Benth. in Mart. Fl. Bras.* xv. II. 92.—Bras.
staminea, Vog. Syn. Cass. 14 = ferruginea.
stenocarpa, *Vog. l. c.* 68.—Bras.
stenophylla, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 535.—Cuba.
stipulacea, [*Soland. in*] *Ait. Hort. Kew.* ed. II. ii. 52.—Chili.
stipulata, G. Don, Gen. Syst. ii. 448 = flavicoma.
striata, Vog. Syn. Cass. 39 = splendida.
stricta, Schrank, Hort. Monac. t. 34 = C. Chamaecrista.
stricta, *Steud. Nom.* ed. II. i. 308.—Afr. austr.
strictifolia, *Benth. in Mart. Fl. Bras.* xv. II. 145.—Bras.
strigillosa, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 581.—Ind. occ.
strobilacea, H. B. & K. Nov. Gen. et Sp. vi. 347 = reticulata.
Sturtii, *R. Br. in Sturt, Centr. Austral. App.* 14.—Austral.
subtriflora, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 176.—Bras.
subulata, *Griseb. in Goett. Abh.* xix. (1874) 130.—Am. austr.
succedana, Bell. ex DC. Prod. ii. 498 = marylandica.
suffruticosa, Koen. ex Roth, Nov. Sp. Pl. 213 = glauca.
sulcata, Blanco, Fl. Filip. ed. II. 236 = hirsuta.
sulcata, *DC. Cat. Hort. Monsp.* 90.—Am. austr.
sulphurea, DC. ex Collad. Hist. Cass. 84 = glauca.
sumatrana, Roxb. Hort. Beng. [31], nomen = siamea.
Sunsub, Forsk. Fl. Aegypt. Arab. 86 = C. Tora.
supplex, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 163.—Bras.
surattensis, Burm. f. Fl. Ind. 97 = glauca.
surinamensis, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 18.—Guiana.
Swainsoni, *Benth. in Mart. Fl. Bras.* xv. II. 167.—Bras.
Swartzii, Wikst. in Vet. Akad. Handl. (1825) 430 = virgata.
sylvestris, Auct. = silvestris.
Tagera, Lam. Encyc. i. 643 = C. Tora.
Tagera, *Linn. Sp. Pl.* 376.—Am. trop.
Tala, Desv. Journ. Bot. iii. (1814) 23. t. 73 = C. Tora.
Tarantan, H. B. & K. Nov. Gen. et Sp. vi. 348 = reticulata.
tarda, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 9.—Malaya.
tecta, *Vog. Syn. Cass.* 56.—Bras.
Telfairiana, Wall. Cat. n. 5324 = mimosoides.
tenella, *H. B. & K. Nov. Gen. et Sp.* vi. 365.—Venezuela.
tenella, Roxb. Hort. Beng. 32, nomen = mimosoides.
tenuifolia, *Vog. Syn. Cass.* 16.—Bras.
tenuisepala, *Benth. in Mart. Fl. Bras.* xv. II. 164.—Am. trop.
tenuissima, Linn. Sp. Pl. 378 = biflora.
tephrosiaefolia, *Benth. in Mart. Fl. Bras.* xv. II. 153.—Am. trop.
teretifolia, *A. Cunn. ex Lindl. in Ann. Sc. Nat.* Sér. II. xv. (1841) 60.—Austral.
teretifolia, Lindl. in Mitch. Three Exped. i. 289 = artemisioides.
teretiuscula, F. Muell. in Linnaea, xxv. (1852) 389 = artemisioides.
tetrafoliata, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 428 = Langsdorffii.
tetraphylla, Collad. Hist. Cass. 130 = hispidula.
tetraphylla, Desv. Journ. Bot. iii. (1814) 72 = Desvauxii.
tettensis, *Bolle, in Peters, Reise Mossamb. Bot.* 14.—Afr. trop.
texana, *Buckl. in Proc. Acad. Sc. Philad.* '1861' (1862) 452 = procumbens, *Linn. Herb.*
Thonningii, DC. Prod. ii. 500 = C. Absus.
Thunbergiana, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 54 = mimosoides.
timorensis, *DC. Prod.* ii. 499.—As. trop.; Austral.
tomentosa, Ehrenb. & Hempr. ex Benth. in Proc. Linn. Soc. ii. (1855) 283 = holosericea.
tomentosa, *Linn. f. Suppl.* 231.—Am. trop.
tomentosa, Wall. Cat. n. 5304 = hirsuta.
Tora, *Linn. Sp. Pl.* 376.—Cosmop. trop.
toroides, Roxb. Hort. Beng. [31], nomen = C. Tora.
torosa, Cav. Desc. 131 = C. Sophera.
torulosa, Poir. Encyc. Suppl. ii. 127 = C. Sophera.
trachycarpa, *Vog. Syn. Cass.* 52.—Bras.

CASSIA:—
trachypus, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 122.—Bras.
tragacanthoides, *Mart. ex Benth. l. c.* 166.—Am. trop.
trichopoda, *Benth. l. c.* 163.—Am. trop.
triflora, Jacq. Hort. Schoenb. iv. 40 = C. Chamaecrista.
triflora, *Vahl, Eclog. Am.* iii. 11.—Ins. S. Cruc.
trinitatis, Benth. in Hook. Journ. Bot. ii. (1840) 77 = racemosa.
trinitatis, Reichb. ex DC. Prod. ii. 489 = spectabilis.
tristicula, *H. B. & K. Nov. Gen. et Sp.* vi. 367.—Am. austr.
tropica, Vell. Fl. Flum. 166; iv. t. 64 = laevigata.
tuberculata, *Collad. Hist. Cass.* 133.—Bras.
Tula, Desv. ex Steud. Nom. ed. II. i. 308 = C. Tora.
umbellata, Reichb. Ic. Exot. t. 206 = australis.
undulata, *Benth. in Hook. Journ. Bot.* ii. (1840) 76.—Am. trop.
uniflora, Mill. Gard. Dict. ed. VIII. n. 5 = sericea.
uniflora, *Spreng. Neue Entdeck.* i. 291.—Am. trop.
Urmenetae, *Phil. in Anal. Univ. Chil.* (1872) 708.—Chili.
ursina, *Mart. ex Benth. in Mart. Fl. Bras.* xv. II. 151.—Am. trop.
Vauthieri, *Benth. l. c.* 138.—Am. trop.
velutina, *Vog. Syn. Cass.* 24.—Bras.
venenifera, Rodsch. in G. F. W. Mey. Prim. Fl. Esseq. 167 = hirsuta.
venosa, Castigl. ex Zuccagni, in Roem. Collect. 140 = C. Chamaecrista.
venosa, Desv. Journ. Bot. iii. (1814) 72 = cytisoides.
venulosa, *Benth. in Mart. Fl. Bras.* xv. II. 170.—Bras.
venusta, *F. Muell. Fragm.* i. 165.—Austral.
venustula, H. B. & K. Nov. Gen. et Sp. vi. 358 = biflora.
vernicosa, Clos, in C. Gay, Fl. Chil. ii. 244 = laevigata.
verrucosa, *Vog. Syn. Cass.* 38.—Bras.
versicolor, Benth. Pl. Hartw. 128 = incarnata.
versicolor, *Meyen, in Vog. Syn. Cass.* 29; *et Pl. Meyen*, 39.—Peruv.
vestita, *Vog. Syn. Cass.* 60.—Bras.
viciaefolia, *Benth. in Trans. Linn. Soc.* xxvii. (1871) 544.—N. Granat.
villosa, *Mill. Gard. Dict.* ed. VIII. n. 4.—Mexic.
viminea, *Linn. Syst. Nat.* ed. X. 1016.—Am. trop.
virgata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = chrysocarpa.
virgata, Sieber, ex Steud. Nom. ed. II. i. 308 = glandulosa.
virgata, *Sw. Prod. Veg. Ind. Occ.* 66.—Ind. occ.
viscida, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie*, iii. (1846) 80.—Malaya.
viscosa, *H. B. & K. Nov. Gen. et Sp.* vi. 360.—N. Granat.
viscosa, Schum. & Thonn. Beskr. Guin. Pl. 205 = C. Absus.
viscosopilosa, Steud. in Flora, xxvi. (1843) II. 760 = viscosa.
Vogeliana, *Schlecht. in Linnaea*, xii. (1838) 342.—Mexic.
vulgaris, *Roxb. ex Steud. Nom.* ed. II. i. 308.—Ind. or.
Wallichiana, DC. in Mém. Soc. Phys. Genèv. ii. (1824) 132 = mimosoides.
Wightiana, R. Grah. in Wall. Cat. n. 5329 = tomentosa.
Wislizeni, *A. Gray, Pl. Wright.* i. 60.—N. Mexic.
Wrightii, *A. Gray, l. c.* ii. 50.—Mexic.
xanthocoma, *Miq. Anal. Bot. Ind.* i. 10.—Borneo.
xiphoidea, Bertol. Fl. Guatim. 14 = biflora.
zambesica, *Oliver, Fl. Trop. Afr.* ii. 280.—Afr. trop.
zanzibarensis, *Vatke, in Oestr. Bot. Zeitschr.* xxx. (1880) 77.—Afr. trop.
zygophylla, Benth. in Mitch. Journ. Trop. Austral. 288 = eremophila.

CASSIANA, Rafin. in Am. Monthly Mag. (1818) 266 = **Cassia**, Linn. (Legumin.).

CASSIDA, Tourn. ex Adans. Fam. ii. 188 (1763) = **Scutelleria**, Herm. (Labiat.).
alpina, Moench, Meth. Suppl. 144 = S. alpina.
altissima, Moench, l. c. 145 = S. altissima.
galericulata, Scop. Fl. Carn. ed. II. i. 430 = S. galericulata.
hastifolia, Scop. l. c. = S. hastifolia.
lateriflora, Moench, Meth. Suppl. 145 = S. lateriflora.
major, Gilib. Fl. Lituan. i. 90 = S. galericulata.
minor, Gilib. l. c. 91 = S. minor.
racemosa, Moench, Meth. 414 = S. peregrina.

CASSIDOCARPUS, Presl, ex DC. Prod. iv. 82 (1830) = **Asteriscium**, Cham. & Schlecht. (Umbellif.).
chilensis, Presl, l. c. = A. chilense.

CASSIERA, Raeusch. Nom. ed. III. 42 (1797) sphalm. = **Cansjera**, Juss. (Olacin.).

CASSINE, Linn. Gen. ed. I. 338 (1737). *CELASTRINEAE*, Benth. & Hook. f. i. 363.
MAUROCENIA, Linn. Gen. ed. I. 85 (1737).
aethiopica, Eckl. & Zeyh. Enum. 128 = capensis.
aethiopica, *Thunb. Fl. Cap.* ii. 227.—Afr. austr.
affinis, *Sond. in Harv. & Sond. Fl. Cap.* i. 465.—Afr. austr.
albanensis, *Sond. l. c.* 467.—Afr. austr.
articulata, E. Mey. ex Sond. l. c. 457 = Celastrus procumbens.
Barba, Murr. Syst. Veg. ed. XIII. 243, sphalm. = seq.
barbara, *Linn. Mant.* ii. 220.—Afr. austr.
capensis, *Linn. l. c.*—Afr. austr.
caroliniana, Lam. Encyc. i. 652 = Ilex Cassine.
caroliniana, Walt. Fl. Carol. 242 = Ilex ambigua.
Colpoon, Eckl. & Zeyh. Enum. 128 = affinis.
Colpoon, Thunb. Prod. Pl. Cap. 52 = capensis.
concava, Lam. Encyc. i. 652 = Celastrus lucidus.
Congonha, St. Hil. in Guill. Arch. Bot. i. (1832) 31 = Villaresia mucronata.
corymbosa, Mill. Gard. Dict. ed. VIII. n. 1 = Ilex Cassine.
crocea, Presl, Bot. Bemerk. 37 = Elaeodendron capense.
discolor, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 378 = Microtropis discolor.
domingensis, *Spreng. Syst.* i. 939.—Ins. S. Doming.
excelsa, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 376 = Ilex excelsa.
Gongonha, Mart. Reise Bras. i. 285 = Ilex Gongonha.
humilis, Hort. ex Steud. Nom. ed. II. i. 308 = Celastrus lucidus.
ilicifolia, Hochst. ex Sond. in Harv. & Sond. Fl. Cap. i. 474 = Cassinopsis capensis.
integrifolia, Salisb. Prod. 173 = C. Maurocenia.
Kraussiana, Bernh. ex Harv. & Sond. Fl. Cap. i. 466 = capensis.
Kraussiana, *Hochst. in Flora*, xxvii. (1844) I. 305.—Afr. austr.
laevigata, Lam. Encyc. i. 652 = Pterocelastrus rostratus.
latifolia, *Eckl. & Zeyh. Enum.* 129.—Afr. austr.
magellanica, Lam. Illustr. ii. 92. n. 2690 = Maytenus magellanious.
Maurocenia, *Linn. Sp. Pl.* ed. II. 385.—Afr. austr.
mucronata, Turcz. in Bull. Soc. Nat. Mosc. (1858) I. 455 = Cassinopsis capensis.
oleifolia, Pers. Syn. i. 327 = Celastrus oleoides.
oleoides, Lam. Encyc. i. 652 = Celastrus oleoides.
oppositifolia, *Mill. Gard. Dict.* ed. VIII. n. 3.—Hab.?
parvifolia, E. Mey. ex Harv. & Sond. Fl. Cap. i. 468 = Elaeodendron croceum.
parvifolia, *Sond. l. c.* 466.—Afr. austr.
Peragua, Linn. Mant. ii. 220 = Ilex Cassine.
ramulosa, Rafin. Fl. Ludov. 110 = Ilex Cassine.
scandens, *Eckl. & Zeyh. Enum.* 128.—Afr. austr.
serratifolia, Salisb. Prod. 173 = capensis.
sphaerocarpa, Cels, ex Steud. Nom. ed. II. i. 308 = capensis.
xylocarpa, Vent. Choix, 23 = Elaeodendron xylocarpum.

CASSINIA, R. Br. in Trans. Linn. Soc. xii. (1817) 126. *COMPOSITAE*, Benth. & Hook. f. ii. 313.
ACHROMOLAENA, Cass. in Dict. Sc. Nat. lvi. 222 (1828).
APALOCHLAMYS, Cass. l. c. 223 (1828).
CHROMOCHITON, Cass. l. c. 220 (1828).
RHYNEA, DC. in Deless. Ic. Sel. iv. t. 52 (1837).
aculeata, *R. Br. in Trans. Linn. Soc.* xii. (1817) 127.—Austral.
aculeata, A. Cunn. ex DC. Prod. v. 156 = Cunninghami.
adunca, F. Muell. in Linnaea, xxv. (1852) 496 = aculeata.
affinis, R. Br. in Trans. Linn. Soc. xii. (1817) 127 = aculeata.
arcuata, *R. Br. l. c.*—Austral.
argophylla, DC. Prod. vi. 155 = Helichrysum ferrugineum.
aurea, *R. Br. in Trans. Linn. Soc.* xii. (1818) 127.—Austral.

CASSINIA :—
aurea, R. Br. in Ait. Hort. Kew. ed. II. v. (1813) 185 = Angianthus tomentosus.
berberifolia, A. Cunn. ex DC. Prod. vi. 155 = denticulata.
cinerea, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 132 = leptophylla.
compacta, *F. Muell. Fragm.* i. 18.—Austral.
cuneifolia, A. Cunn. ex DC. Prod. vi. 155 = Helichrysum Backhousii.
Cunninghami, DC. Prod. vi. 156 = Helichrysum Cunninghamii.
cuprea, F. Muell. Fragm. iii. 139 = Helipterum polycephalum.
denticulata, *R. Br. in Trans. Linn. Soc.* xii. (1817) 127.—Austral.
fulvida, *Hook. f. Handb. New Zeal. Fl.* 145.—N. Zel.
glossophylla, *Cass. in Dict. Sc. Nat.* lvi. 219.—Austral.
hygrophila, A. Cunn. in DC. Prod. vi. 156 = quinquefaria.
laevis, *R. Br. in Trans. Linn. Soc.* xii. (1818) 127.—Austral.
laevis, Endl. ex DC. l. c. = longifolia.
ledifolia, A. Cunn. ex DC. Prod. vi. 155 = Helichrysum ledifolium.
leptocephala, *F. Muell. Fragm.* iii. 138.—Austral.
leptophylla, *R. Br. in Trans. Linn. Soc.* xii. (1817) 126.—N. Zel.
longifolia, *R. Br. l. c.* 127.—Austral.
obovata, A. Cunn. in DC. Prod. vi. 155 = Helichrysum obcordatum.
paniculata, Behr & F. Muell. in Linnaea, xxv. (1852) 496 = arcuata.
patens, A. Cunn. ex DC. Prod. vi. 156 = Helichrysum Cunninghami.
pholidota, F. Muell. Fragm. ii. 131 = Helichrysum pholidotum.
quinquefaria, *R. Br. in Trans. Linn. Soc.* xii. (1817) 127.—Austral.
quinquefaria, Sond. in Linnaea, xxv. (1852) 496 = Humea punctulata.
retorta, *A. Cunn. ex DC. Prod.* vi. 154.—N. Zel.
rosmarinifolia, A. Cunn. ex DC. l. c. 156 = laevis.
spathulata, Endl. ex Steud. Nom. ed. II. i. 309 = spectabilis.
spectabilis, *R. Br. in Trans. Linn. Soc.* xii. (1817) 128.—Austral.
subtropica, *F. Muell. Fragm.* i. 17.—Austral.
tenuifolia, *Benth. Fl. Austral.* iii. 585.—Austral.
Theodori, *F. Muell. Fragm.* v. 148.—Austral.
uncata, A. Cunn. in DC. Prod. vi. 156 = aculeata.
Vauvilliersii, *Hook. f. Fl. N. Zel.* 133.—N. Zel.

CASSINIOLA, F. Muell. Fragm. iii. 139 (1863) = **Helipterum**, DC. (Compos.).
cuprea, F. Muell. l. c. = H. polycephalum.

CASSINOPSIS, Sond. in Harv. & Sond. Fl. Cap. i. 473 (1860). *OLACINEAE*, Benth. & Hook. f. i. 354.
capensis, *Harv. & Sond. l. c.* 474.—Afr. austr.
ciliata, *Baker, in Journ. Linn. Soc.* xx. (1884) 118.—Madag.
madagascariensis, *Baill. Adansonia*, xi. (1873-76) 180.—Madag.
tinifolia, *Harv. Thes. Cap.* ii. 44.—Afr. austr.

CASSIOPE, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 157. *ERICACEAE*, Benth. & Hook. f. ii. 584.
ericoides, *D. Don. l. c.* 158.—Sibiria.
fastigiata, *D. Don, l. c.* 156.—Reg. Himal.
hypnoides, *D. Don, l. c.* 158.—Reg. arct. et subarct.
lycopodioides, *D. Don, l. c.* 158.—Sibiria; Am. subarct. occ.
Mertensiana, *G. Don, Gen. Syst.* iii. 829.—Am. bor. occ.
oxycoccoides, *A. Gray, in Proc. Am. Acad.* xx. (1885) 300.—Kamtschat.
Redowskii, *G. Don, Gen. Syst.* iii. 829.—Sibir. or.; Ross.
selaginoides, *Hook. f. & Thoms. in Hook. Kew Journ.* vii. (1855) 126. t. 4.—Reg. Himal.
Stelleriana, *DC. Prod.* vii. 611.—Alaska.
tetragona, *D. Don, in Edinb. N. Phil. Journ.* xvii. (1834) 158.—Reg. arct. et subarct.

CASSIPOUREA, Aubl. Pl. Gui. i. 528. t. 211 (1775). *RHIZOPHOREAE*, Benth. & Hook. f. i. 682.
LEGNOTIS, Sw. Prod. Veg. Ind. Occ. 84 (1788).
TITA, Scop. Introd. 219 (1777).
africana, Benth. in Hook. Niger Fl. 341 = Weihea africana.
alba, *Griseb. in Goett. Abh.* vii. (1857) 223.—Ind. occ.
congensis, *R. Br. in Tuckey, Voy. Congo*, 18.—Afr. trop.
elliptica, *Poir. Encyc. Suppl.* ii. 131.—Jamaica.
guianensis, *Aubl. Pl. Gui.* i. 529. t. 211.—Am. austr.
gummiflua, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 123 = Dactylopetalum sessiliflorum.
lanceolata, *Tul. l. c.* 121.—Ins. Mascar.
leptoclada, *Tul. l. c.* 125.—Ins. Mascar.
macrophylla, *DC. Prod.* iii. 34.—Bras.
madagascariensis, *DC. l. c.*—Madag.
microcarpa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 162.—Ins. Mascar.
microphylla, *Tul. l. c.* vi. (1856) 124.—Madag.
myriocarpa, *Tul. l. c.* viii. (1857) 162.—Madag.
ovata, *Tul. l. c.* vi. (1856) 121.—Ins. Comor.
phaeotricha, *Tul. l. c.* 122.—Madag.
quadrilocularis, Spruce, ex Benth. in Mart. Fl. Bras. xii. II. 429 = guianensis.
serrata, *Benth. in Hook. Journ. Bot.* ii. (1840) 223.—Guiana.
Spruceana, *Benth. in Mart. Fl. Bras.* xii. II. 429.—Am. trop.

CASSITHA, Hill, Veg. Syst. ix. 17 (1765) = **Cassytha**, Linn. (Laurin.).

CASSUMBIUM, Benth. & Hook. f. Gen. i. 404 (1862) = CUSSAMBIUM, Rumph. = **Schleichera**, Willd. (Sapind.).

CASSUMUNAR, Colla, Nov. Scit. Gen. 9 (1830) = **Zingiber**, Adans. (Scitamin.).
Roxburghii, Colla, l. c. 10 = Z. Cassumunar.

CASSUPA, Humb. & Bonpl. Pl. Aequin. i. 43. t. 12 (1808). *RUBIACEAE*, Benth. & Hook. f. ii. 65.
laevis, *Triana, in Ann. Sc. Nat.* Sér. IV. ix. (1858) 44.—N. Granat.
verrucosa, *Humb. & Bonpl. Pl. Aequin.* i. 43. t. 12.—Mexic.

CASSUTHA, [J. Bauh.] Des Moul. Etud. Org. Cusc. 66 (1853) = **Cuscuta**, Linn. (Convolvul).
americana, Des Moul. l. c. 70 = Cusc. americana.
arabica, Des Moul. l. c. 72 = Cusc. arabica.
chrysocoma, Des Moul. l. c. 71 = Cusc. australis.
suaveolens, Des Moul. l. c. 66 = Cusc. racemosa.

CASSUVIUM, Lam. Encyc. i. 22 (1783) = **Anacardium**, Rottb.
pomiferum, Lam. l. c. = A. occidentale.
reniforme, Blanco, Fl. Filip. ed. I. 322 = A. occidentale.
solitarium, Stokes, Bot. Mat. Med. ii. 475 = A. occidentale.

CASSYTA, Linn. Gen. ed. VI. 291 (1764) = **Cassytha**, Linn. (Laurin.).

CASSYTHA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 345 (1821) = **Cuscuta**, Tourn. (Convolvul.).
major, S. F. Gray, l. c. 346 = Cusc. europaea.

CASSYTHA, Linn. Sp. Pl. ed. I. 35 (1753). *LAURINEAE*, Benth. & Hook. f. iii. 164.
CALODIUM, Lour. Fl. Cochinch. 247 (1790).
OZARTHRIS, Rafin. Fl. Tellur. iv. 92 (1836).
ROMBUT, Rumph. ex Adans. Fam. ii. 284 (1763).
RUMPUTRIS, Rafin. Fl. Tellur. iv. 92 (1836).
VOLUTELLA, Forsk. Fl. Aegypt. Arab. 84 (1775).
americana, Nees, Syst. Laurin. 644 = filiformis.
aphylla, Raeusch. Nom. ed. III. 116 = filiformis.
brasiliensis, Mart. ex Nees, Syst. Laurin. 648 = filiformis.
capensis, *Meissn. in DC. Prod.* xv. I. 254.—Afr. austr.
capillaris, *Meissn. l. c.* 252.—Zeylan.; Borneo.
Casuarinae, Nees, in Lehm. Pl. Preiss. i. 619 = glabella.
ceratopoda, Meissn. in DC. Prod. xv. I. 257 = pomiformis.
ciliolata, *Nees, Syst. Laurin.* 646.—Afr. austr.
corniculata, *Burm. f. Fl. Ind.* 93.—Ind. or.
coronata, *Nees, in Lehm. Pl. Preiss.* i. 620.—Austral.
digitata, Nees, l. c. = racemosa.

CASSYTHA :—
dispar, Schlecht. in Linnaea, xx. (1847) 578 = glabella.
dissitiflora, *Meissn. in Kjoeb. Vidensk. Meddel.* (1870) 145.—Bras.
filiformis, *Linn. Sp. Pl.* 35.—Cosmop. trop.
filiformis, Thunb. Prod. Pl. Cap. 78 = glabella.
flava, *Nees, in Lehm. Pl. Preiss.* i. 620.—Austral.
glabella, *R. Br. Prod.* 404.—Austral.
glabella, Sieber, ex Meissn. in DC. Prod. xv. I. 256 = paniculata.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 199 = filiformis.
melantha, *R. Br. Prod.* 404.—Austral.
micrantha, *Meissn. in DC. Prod.* xv. I. 256.—Austral.
microcephala, Meissn. l. c. 253 = glabella.
Muelleri, Meissn. l. c. 257 = racemosa.
multiflora, Nees, in Lehm. Pl. Preiss. i. 621 = pomiformis.
nodiflora, *Meissn. in DC. Prod.* xv. I. 252.—Austral.
paniculata, *R. Br. Prod.* 404.—Austral.
phaeolasia, *F. Muell. Fragm.* v. 167.—Austral.
piligera, Schlecht. in Linnaea, xxi. (1848) 446 = pubescens.
pomiformis, *Nees, in Lehm. Pl. Preiss.* i. 620.—Austral.
pubescens, *R. Br. Prod.* 404.—Austral.
racemosa, *Nees, in Lehm. Pl. Preiss.* i. 621.—Austral.
remotiflora, F. Muell. ex Meissn. in DC. Prod. xv. I. 256 = paniculata.
robusta, Meissn. l. c. 255 = melantha.
rubiginosa, *E. Mey. in Drège, Zwei Pfl. Docum.* 171, nomen.—Afr. austr.
rugulosa, Meissn. in DC. Prod. xv. I. 255 = pubescens.
subcapitata, Meissn. l. c. 253 = pomiformis, racemosa.
tasmanica, Meissn. l. c. 252 = pubescens.
triflora, E. Mey. in Drège, Zwei Pfl. Docum. 171 = capensis.
umbellata, Meissn. in DC. Prod. xv. I. 257 = racemosa?
zeylanica, Gaertn. Fruct. i. 134 = filiformis?

CASSYTHA, Mill. Gard. Dict. ed. VIII (1768) = **Rhipsalis**, Gaertn. (Cacteae).
baccifera, Mill. ex DC. Prod. iii. 476 = seq.
filiformis, Mill. Gard. Dict. ed. VIII = R. Cassytha.
polysperma, Ait. ex Gaertn. Fruct. i. 137 = R. Cassytha.

CASTALIA, Salisb. Parad. Lond. i. t. 14 (1805); et in Kon. & Sims, Ann. Bot. ii. (1806) 71 = **Nymphaea**, Linn.
ampla, Salisb. ll. cc. t. 14, 73 = N. ampla.
caerulea, Tratt. in Flora, v. (1822) 604 = N. stellata.
edulis, Salisb. Parad. Lond. i. sub t. 14; et in Kon. & Sims, Ann. Bot. ii. (1806) 73 = N. Lotus.
Lotus, Tratt. in Flora, v. (1822) 598 = N. Lotus.
magnifica, Salisb. Parad. Lond. i. t. 14 = N. Lotus.
mystica, Salisb. l. c.; et in Kon. & Sims, Ann. Bot. ii. (1806) 73 = N. Lotus.
pubescens, Blume, Bijdr. 48 = N. Lotus.
pudica, Salisb. in Kon. & Sims, Ann. Bot. ii. (1806) 72 = N. odorata.
pygmaea, Salisb. Parad. Lond. (1806) i. 2. t. 68 = N. tetragona.
rubra, Tratt. in Flora, v. (1822) 595 = N. Lotus.
Rudgeana, Tratt. l. c. 599 = N. tetragona.
sacra, Salisb. Parad. Lond. i. sub t. 14 = N. Lotus.
scutifolia, Salisb. l. c.; et in Kon. & Sims, Ann. Bot. ii. (1806) 72 = N. stellata.
speciosa, Salisb. in Kon. & Sims, Ann. Bot. ii. (1806) 72 = N. alba.
stellaris, Salisb. Parad. Lond. i. sub t. 14 = N. stellata.
stellata, Blume, Bijdr. 49 = N. stellata.
versicolor, Tratt. in Flora, v. (1822) 600 = N. stellata.

CASTALIS, Cass. in Dict. Sc. Nat. xxx. 332 (1824) = **Dimorphotheca**, Moench (Compos.).
Ventenati, Cass. l. c. = D. aurantiaca.

CASTANEA, Tourn. ex Linn. Syst. ed. I. (1735). *CUPULIFERAE*, Benth. & Hook. f. iii. 409.
CASTANOPHORUM, Neck. Elem. iii. 257 (1790) (Casanophorum).
acuminatissima, Blume, Mus. Bot. Lugd. Bat. i. 283 = Quercus Junghuhnii.
alnifolia, Nutt. Gen. Am. ii. 217 = pumila.
americana, G. Don, in Sweet, Hort. Brit. ed. III. 614 = sativa.
americana, Rafin. Fl. Ludov. 134; New Sylva. 82 = sativa.
argentea, Blume, Bijdr. 525 = Castanopsis argentea.

CASTANEA :—
argyrophylla, Buek, in DC. Prod. Index, iv. 67 = Castanopsis argentea.
brevicuspis, Miq. Fl. Ind. Bat. i. 1. 866 = Castanopsis javanica.
Bungeana, Blume, Mus. Bot. Lugd. Bat. i. 284 = sativa.
buruana, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 377 = Castanopsis buruana.
castanicarpa, Oerst. l. c. = Castanopsis castanicarpa.
Chincapin, Hort. ex C. Koch, Dendrol. ii. II. 24 = pumila.
chinensis, Hassk. Cat. Hort. Bog. Alt. 73 = sativa.
chinensis, Spreng. Syst. iii. 856 = Castanopsis tribuloides.
chrysophylla, Dougl. ex Hook. Fl. Bor. Am. ii. 159 = Castanopsis chrysophylla.
concinna, Champ. ex Benth. in Hook. Kew Journ. vi. (1854) 115 = Castanopsis concinna.
cooperta, *Oerst. in Vidensk. Selsk. Skr.* V. ix. (1873) 379.—Ins. Philipp.
costata, Blume, Mus. Bot. Lugd. Bat. i. 284 = Castanopsis javanica.
crenata, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 324 = sativa.
divaricata, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 379 = Castanopsis argentea.
diversifolia, *Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 198.—Ind. or.
echidnocarpa, Hook. f. et Thoms. ex A. DC. in Prod. xvi. II. 112 = Castanopsis tribuloides.
Fagus, Scop. Fl. Carn. ed. II. ii. 242 = Fagus sylvatica.
Falconeri, Hance, in Journ. Bot. xiii. (1875) 367 = Castanopsis armata.
furfurella, Miq. Fl. Ind. Bat. Suppl. 352 = Castanopsis Hystrix.
glomerata, Blume, Mus. Bot. Lugd. Bat. i. 283 = Castanopsis sumatrana.
Hystrix, Hook. f. & Thoms. ex Miq. in Ann. Mus. Bot. Lugd. Bat. i. 119 = Castanopsis Hystrix.
indica, Roxb. Hort. Beng. 68; Fl. Ind. iii. 643 = Castanopsis indica.
inermis, Lindl. in Wall. Pl. As. Rar. ii. 6 = Castanopsis sumatrana.
japonica, Blume, Mus. Bot. Lugd. Bat. i. 284 = sativa.
javanica, Blume, Bijdr. 525 = Castanopsis javanica.
lanceaefolia, Kurz, For. Fl. Brit. Burma, ii. 482 = Quercus lanceaefolia.
latifolia, *Blume, Bijdr.* 526.—Java.
martabanica, Wall. Cat. n. 2764; Pl. As. Rar. ii. t. 107 = Castanopsis argentea.
microcarpa, Lindl. ex Wall. Cat. n. 3735 = Castanopsis tribuloides.
mollissima, *Blume, Mus. Bot. Lugd. Bat.* i. 286.—China.
montana, Blume, Bijdr. 526 = Castanopsis javanica.
nana, Muhl. Cat. 86 = pumila.
pumila, Hassk. Cat. Hort. Bog. Alt. 73 = sativa.
pumila, *Mill. Gard. Dict.* ed. VIII. n. 2.—Am. bor.
pumila, Siebold, ex Blume, Mus. Bot. Lugd. Bat. i. 286 = Castanopsis tribuloides.
purpurella, Miq. Fl. Ind. Bat. Suppl. 352 = Castanopsis Hystrix.
regia, Hance, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 230 = Quercus fissa.
rhamnifolia, Kurz, For. Fl. Brit. Burma, ii. 481 = Castanopsis rhamnifolia.
rhamnifolia, *Oerst. in Vidensk. Selsk. Skr.* V. ix. (1873) 378.—Am. centr.
Roxburghii, Lindl. in Wall. Pl. As. Rar. ii. 6 = Castanopsis castanicarpa.
sativa, *Mill. Gard. Dict.* ed. VIII. n. 1.—Europ.; Japon.; Am. bor.
sempervirens, Kellogg, in Proc. Calif. Acad. i. 71, ed. II. (1873) 75 = Castanopsis chrysophylla.
sessi[li]folia, Blume, Mus. Bot. Lugd. Bat. i. 284 = Quercus Junghuhnii.
Sloanea, Mill. Gard. Dict. ed. VIII. n. 3 = Sloanea dentata.
spectabilis, Miq. Fl. Ind. Bat. i. 1. 866 = Castanopsis sumatrana.
sphaerocarpa, Lindl. in Wall. Pl. As. Rar. ii. 5 = Castanopsis armata.
stricta, Sieb. & Zucc. in Abh. Akad. Muench. iv. III. (1846) 225 = sativa.
sumatrana, Oerst. in Vidensk. Selsk. Skr. V. ix. (1873) 378 = Castanopsis sumatrana.

CASTANEA :—
tribuloides, Lindl. in Wall. Pl. As. Rar. ii. 6 = Castanopsis tribuloides.
tribuloides, Wall. Cat. n. 2765 B. = Quercus lanceaefolia.
Tungurrut, Blume, Bijdr. 525 = Castanopsis Tungurrut.
vesca, Gaertn. Fruct. i. 181. t. 37 = sativa.
vulgaris, Lam. Encyc. i. 708 = sativa.

CASTANELLA, Spruce, ex Benth. & Hook. f. Gen. i. 394 (1862). *SAPINDACEAE*, Benth. & Hook. f. l. c.
granatensis, Planch. & Linden, in Ann. Sc. Nat. Sér. IV. xviii. (1862) 365 = Paullinia granatensis.
paullinioides, *Spruce, ex Radlk. Monog. Serjan.* 75.—Bras.
riparia, *Spruce, ex Radlk. l. c.*—Peruv.

CASTANOCARPUS, Sweet, Hort. Brit. ed. II. 589 (1830) = **Castanospermum**, A. Cunn. (Legumin.).
australis, Sweet, l. c. = Castanosp. australe.

CASTANOLA, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 503 = **Connarus**, Linn.
trinervis, Llanos, l. c. = Conn. monocarpus.

CASTANOPHORUM, Neck. Elem. iii. 257 (1790) = **Castanea**, Tourn. (Cupulif.).

CASTANOPSIS, Spach, Hist. Vég. Phan. xi. 185 (1842). *CUPULIFERAE*, Benth. & Hook. f. iii. 409.
CALLAEOCARPUS, Miq. Pl. Jungh. i. 13 (1851).
acuminatissima, A. DC. ex Hance, in Journ. Bot. i. (1863) 182 = Quercus lanceaefolia.
argentea, *A. DC. in Journ. Bot.* i. (1863) 182.—Burma; Malaya.
armata, *Spach, Hist. Vég. Phan.* xi. 185.—Reg. Himal.; Burma.
brevicuspis, A. DC. ex Hance, in Journ. Bot. i. (1863) 183 = javanica.
buruana, *Miq. Ann. Mus. Lugd. Bat.* i. 120.—Ins. Molucc.
castanicarpa, *Spach, Hist. Vég. Phan.* xi. 185.—Reg. Himal.
caudata, *Franch. Pl. Davidian.* i. 277.—Mongol.
chinensis, Hance, in Journ. Linn. Soc. x. (1869) 199 = tribuloides.
chrysophylla, *A. DC. in Journ. Bot.* i. (1863) 182.—Am. bor. occ.
concinna, *A. DC. l. c.*—Hongkong.
costata, A. DC. l. c. = javanica.
discocarpa, Hance, l. c. xvi. (1878) 201 = Quercus discocarpa.
echidnocarpa, A. DC. l. c. i. (1863) 182 = tribuloides.
echinocarpa, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 119, sphalm. = praec.
Fabri, *Hance, in Journ. Bot.* xxii. (1884) 230.—China.
Falconeri, Hance, l. c. xiii. (1875) 368 = armata.
ferox, Spach, Hist. Vég. Phan. xi. 185 = tribuloides.
Fordii, *Hance, in Journ. Bot.* xxii. (1884) 230.—China.
Hystrix, *A. DC. l. c.* i. (1863) 182.—Reg. Himal.; Malaya.
indica, *A. DC. l. c.*—Reg. Himal.
inermis, Benth. & Hook. f. Gen. iii. 409 = sumatrana.
javanica, *A. DC. in Journ. Bot.* i. (1863) 182.—Malaya.
jucunda, *Hance, l. c.* xxii. (1884) 230.—China.
Lamontii, *Hance, l. c.* xiii. (1875) 369.—China.
lanceaefolia, Kurz, For. Fl. Brit. Burma, ii. 482 = Quercus lanceaefolia.
mitifica, Hance, in Journ. Bot. xvi. (1878) 200 = sumatrana.
Pierrei, *Hance, l. c.* xiii. (1875) 369.—Cambodia.
rhamnifolia, *A. DC. Prod.* xvi. II. 113.—Burma; Sumatra.
Schefferiana, *Hance, in Journ. Bot.* xvi. (1878) 200.—Sumatra.
sessilifolia, A. DC. l. c. i. (1863) 182 = Quercus Junghuhnii.
spectabilis, A. DC. l. c. = javanica.
sumatrana, *A. DC. l. c.*—Burma; Malaya.
tibetana, *Hance, l. c.* xiii. (1875) 367.—Tibet; China.
tribuloides, *A. DC. l. c.* i. (1863) 182.—Reg. Himal.; Burma.
trisperma, Scheff. in Tijdschr. Nederl. Ind. xxxi. (1870) 362; in Flora, liii. (1870) 252 = javanica.
Tungurrut, *A. DC. in Journ. Bot.* i. (1863) 182.—Malaya.

CASTANOSPERMUM, A. Cunn. in Hook. Bot. Misc. i. (1830) 241. t. 51. *LEGUMINOSAE*, Benth. & Hook. f. i. 556.
CASTANOCARPUS, Sweet, Hort. Brit. ed. II. 589 (1830).
VIELLARDIA, Montr. in Mém. Acad. Lyon, x. (1860) x. 196.
australe, *A. Cunn. & Fraser, in Hook. l. c.*—Austral.

CASTANOSPORA, F. Muell. Fragm. ix. 92 (1875). *SAPINDACEAE*.
Alphandi, *F. Muell. l. c.*—Austral.

CASTELA, Turp. in Ann. Mus. Par. vii. (1806) 78. t. 5. *SIMARUBEAE*, Benth. & Hook. f. i. 310.
alaternifolia, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 569.—Chili.
coccinea, *Griseb. in Goett. Abh.* xix. (1874) 107.—Reg. Argent.
depressa, *Turp. in Ann. Mus. Par.* vii. (1806) 79. t. 5 A.—Ins. S. Doming.
erecta, *Turp. l. c.* 80. t. 5 B.—Ind. occ.
Galapagei, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 229.—Ins. Galap.
laevigata, *Zipp. ex Span. in Linnaea*, xv. (1841) 186, nomen.—Hab.?
lychnophoroides, *Liebm. in Kjoeb. Vidensk. Meddel.* (1853) 111.—Mexic.
Nicholsoni, *Hook. Bot. Misc.* i. (1830) 271. t. 56.—Ind. occ.
retusa, *Liebm. in Kjoeb. Vidensk. Meddel.* (1853) 110.—Mexic.
tortuosa, *Liebm. l. c.* 108.—Mexic.
Tweediei, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 569.—Reg. Argent.

CASTELIA, Cav. in Anal. Cienc. Nat. iii. (1801) 134 = **Priva**, Adans. (Verbenac.).
cuneato-ovata, Cav. l. c. 135 = laevis.

CASTELIA, Liebm. in Kjoeb. Vidensk. Meddel. (1853) 108 = **Castela**, Turp. (Simarub.).

CASTELLIA, Tin. Pl. Rar. Sicul. 17 (1817) = **Festuca**, Linn. (Gramin.).
tuberculata, Tin. l. c. 18 = F. tuberculata.

CASTELNAVIA, Tul. & Wedd. in Ann. Sc. Nat. Sér. III. xi. (1849) 108. *PODOSTEMACEAE*, Benth. & Hook. f. iii. 114.
CARAJAEA, Wedd. in DC. Prod. xvii. 84 (1873).
fimbriata, *Tul. & Wedd. l. c.*—Bras.
fluitans, *Tul. & Wedd. l. c.* 109.—Bras.
fruticulosa, *Tul. & Wedd. l. c.* 110.—Bras.
monandra, *Tul. & Wedd. l. c.* 109.—Bras.
multipartita, *Tul. & Wedd. l. c.*—Bras.
orthocarpa, *Tul. & Wedd. l. c.* 110.—Bras.
Princeps, *Tul. & Wedd. l. c.* 108.—Bras.
pusillina, *Tul. & Wedd. l. c.* 109.—Bras.
serpens, *Tul. & Wedd. l. c.*—Bras.

CASTIGLIONIA, Ruiz & Pav. Prod. 139. t. 37 (1794) = **Jatropha**, Linn. (Euphorbiac.).
lobata, Ruiz & Pav. l. c.; Syst. Veg. 277 = J. Curcas.

CASTILLA, Cerv. in Supl. Gaz. Lit. Mexico (Jul. 2, 1794), 7 = **Castilloa**, 'Cerv.' (Urticac.).

CASTILLEJA, Mutis, ex Linn. f. Suppl. 47 (1781). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 973.
EUCHROMA, Nutt. Gen. Am. ii. 54 (1818).
acuminata, Spreng. Syst. ii. 775 = pallida.
acuminata, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 321 = pallida.
affinis, Benth. Pl. Hartw. 329 = stenantha.
affinis, *Hook. & Arn. Bot. Beech. Voy.* 154.—Calif.
ambigua, Hook. & Arn. l. c. = Orthocarpus australis.
angustifolia, G. Don, Gen. Syst. iv. 616 = parviflora.
angustifolia, A. Gray, in Torr. Bot. Mex. Bound. 118 = integra.
angustifolia, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 29.—Mexic.
anthemidifolia, Benth. in DC. Prod. x. 528 = tenuiflora.
arvensis, *Cham. & Schlecht. in Linnaea*, v. (1830) 103.—Mexic.
Bradburii, G. Don, Gen. Syst. iv. 616 = parviflora.
breviflora, *Benth. in DC. Prod.* x. 534.—Venezuela.
breviflora, *A. Gray, in Am. Journ. Sc.* Ser. II. xxxiv. (1862) 250.—Am. bor. occ.

CASTILLEJA:—
candens, Durand & Hilg. in Pacif. Rail. Rep. v. 12 = linariaefolia.
canescens, *Benth. in DC. Prod.* x. 533.—Mexic.
cinerea, *A. Gray, in Proc. Am. Acad.* xix. (1883) 93.—Calif.
coccinea, Dougl. in Lindl. Bot. Reg. t. 1136 = parviflora.
coccinea, *Spreng. Syst.* ii. 775.—Am. bor.
communis, *Benth. in DC. Prod.* x. 529.—Am. trop.
coronopifolia, *Vent. Choix*, 59.—N. Granat.
desertorum, Geyer, in Hook. Kew Journ. v. (1853) 258 = parviflora.
divaricata, *Benth. in DC. Prod.* x. 534.—Venczuela.
Douglasii, Benth. l. c. 530 = parviflora.
fissifolia, *Linn. f. Suppl.* 293.—N. Granat.
flava, *S. Wats. in Bot. King, Exped.* 230.—Am. bor. occ.
foliolosa, *Hook. & Arn. Bot. Beech. Voy.* 154.—Calif.
fulgens, Nutt. ex A. Gray, in Torr. Bot. Mex. Bound. 119 = linariaefolia?
gracilis, *Benth. in DC. Prod.* x. 528.—Mexic.
grandiflora, Spreng. Syst. ii. 755 = sessiliflora.
hirsuta, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 29.—Mexic.
hispida, Benth. in Hook. Fl. Bor. Am. ii. 105 = parviflora.
hyssopifolia, Willd. ex G. Don, Gen. Syst. iv. 615 = lithospermoides.
indivisa, *Engelm. in Engelm. & A. Gray, Pl. Lindh.* i. 47.—Texas.
integra, *A. Gray, in Torr. Bot. Mex. Bound.* 119.—N. Mexic.
integrifolia, H. B. & K. Nov. Gen. et Sp. ii. 330 = canescens.
integrifolia, *Linn. f. Suppl.* 293.—Am. trop.
irasuensis, *Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 27.—Costa Rica.
laciniata, *Hook. & Arn. Bot. Beech. Voy.* 60.—Chili.
lanata, *A. Gray, in Torr. Bot. Mex. Bound.* 118.—N. Mexic.
latifolia, *Hook. & Arn. Bot. Beech. Voy.* 154.—Calif.
laxa, *A. Gray, in Torr. Bot. Mex. Bound.* 119.—N. Mexic.
Lemmoni, *A. Gray, Syn. Fl. N. Am.* ii. 1. 297.—Calif.
linariaefolia, *Benth. in DC. Prod.* x. 532.—Mexic.
Lindheimeri, *A. Gray, Syn. Fl. N. Am.* ii. 1. 298.—Texas.
linoides, *A. Gray, l. c.* 299.—Am. bor. occ.
lithospermoides, *H. B. & K. Nov. Gen. et Sp.* ii. 331. t. 164.—Mexic.
longibracteata, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 28.—Mexic.
longiflora, *Kunze, in Linnaea*, xvi. (1842) 312.—Mexic.
macrocarpa, Benth. Scroph. Ind. 13 = latifolia.
miniata, *Dougl. in Hook. Fl. Bor. Am.* ii. 106.—Am. bor. occ.
minor, *A. Gray, Bot. Calif.* i. 573.—Am. bor. occ.
moranensis, *H. B. & K. Nov. Gen. et Sp.* 329.—Mexic.
nubigena, *H. B. & K. l. c.* 330.—Ecuador.
oblongifolia, *A. Gray, Syn. Fl. N. Am.* ii. 1. 296.—Calif.
obovata, Benth. in DC. Prod. x. 528 = hirsuta.
occidentalis, Torr. in Ann. Lyc. N. York, ii. (1828) 230 = pallida.
Orisabae, Benth. in DC. Prod. x. 533 = pectinata.
pallida, *Kunth, Syn. Pl. Aequin.* ii. 100.—Sibir.; Am. bor.
pallida, Spreng. Syst. ii. 774 = septentrionalis, sibirica.
parviflora, *Bong. in Mém. Acad. Pétersb.* Sér. VI. v. II. (1840) 157.—Am. bor. occ.
pectinata, *Mart. & Gall. in Bull. Acad. Brux.* xii. II. (1845) 27.—Mexic.
plagiotoma, *A. Gray, in Proc. Am. Acad.* xix. (1883) 93.—Calif.
pumila, Wedd. Chlor. And. t. 61 = fissifolia.
purpurea, *G. Don, Gen. Syst.* iv. 615.—Am. bor. occ.
Schaffneri, *Hemsl. Biol. Centr. Am. Bot.* ii. 462.—Mexic.
scorzoneraefolia, *H. B. & K. Nov. Gen. et Sp.* ii. 331.—Mexic.
septentrionalis, *Lindl. Bot. Reg.* t. 925.—Am. bor.
serrata, *Hort. ex Loud. Hort. Brit. Suppl.* iii. 511.—Hab.?
sessiliflora, *Pursh, Fl. Am. Sept.* ii. 738.—Am. bor. occ.
sibirica, Lindl. Bot. Reg. t. 925 = pallida.

CASTILLEJA:—
speciosa, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 30 = scorzoneraefolia.
stenantha, *A. Gray, Syn. Fl. N. Am.* ii. 1. 295.—Am. bor. occ.
stricta, *Benth. in DC. Prod.* x. 534.—Ecuador.
tenuiflora, *Benth. Pl. Hartw.* 22.—Mexic.
tenuifolia, *Mart. & Gal. Bull. Acad. Brux.* xii. II. (1845) 30.—Mexic.
toluccensis, Cham. & Schlecht. in Linnaea, ii. (1827) 579 = parviflora?
toluccensis, *H. B. & K. Nov. Gen. et Sp.* ii. 329.—Mexic.
tomentosa, A. Gray, in Torr. Bot. Mex. Bound. 118 = integra.
trifida, Willd. ex Walp. Rep. iii. 394 = nubigena.
viscidula, *A. Gray, Syn. Fl. N. Am.* ii. 1. 297.—Am. bor. occ.

CASTILLOA, Cervant. in Supl. Gaz. Lit. Mexic. (Jul. 2, 1794) 7 (Castilla). *URTICACEAE*, Benth. & Hook. f. iii. 372.
costaricana, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 319.—Costa Rica.
elastica, *Cerv. in Supl. Gaz. Lit. Mexico* (Jul. 2, 1794), 7.—Mexic.
Markhamiana, *Collins, Rep. Caoutch. Comm.* 12. t. 3.—Am. trop.

CASTRA, Vell. Fl. Flum. 242 (1825); viii. 79–81 = **Trixis**, P. Br. (Compos.).
illustris, Vell. Fl. Flum. 343; viii. t. 81 = T. brasiliensis.
nobilis, Vell. l. c. 344; viii. t. 81 = T. glaberrima.
praestans, Vell. l. c. 343; viii. t. 80 = T. mollissima.
regia, Vell. l. c. 343; viii. t. 79 = T. divaricata.

CASTRATELLA, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 139. *MELASTOMACEAE*, Benth. & Hook. f. i. 736.
piloselloides, *Naud. l. c.*—N. Granat.

CASTREA, A. St. Hil. Morphol. 451 (1840) = **Phoradendron**, Nutt. (Loranthac.).
falcata, St. Hil. l. c.—Bras.?

CASTRONIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 = **Helicia**, Lour. (Proteac.).

CASUARINA, Linn. Amoen. Acad. iv. 143, nomen (1759); Adans. Fam. ii. 481 (1763); Forst. Char. Gen. 103. t. 52 (1776). *CASUARINEAE*, Benth. & Hook. f. iii. 402.
acuaria, *F. Muell. in Journ. Bot.* v. (1867) 212.—Austral.
acutivalvis, *F. Muell. Fragm.* x. 61.—Austral.
africana, Lour. Fl. Cochinch. ii. 549 = equisetifolia.
angulata, *Poiss. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 49.—N. Caled.
Baxteriana, Miq. Rev. Crit. Casuar. 37. t. 3 C = distyla.
bicuspidata, *Benth. Fl. Austral.* vi. 202.—Austral.
Brunoniana, *Miq. Rev. Crit. Casuar.* (1848) 21. t. 1. C.—Java.
Chamaecyparis, *Poiss. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 52.—N. Caled.
collina, *Poiss. ex Panch. & Sebert, Not. Bois Nouv. Calléd.* 174.—N. Caled.
corniculata, *F. Muell. Fragm.* x. 62.—Austral.
cristata, Miq. Rev. Crit. Casuar. 70. t. 10. A = stricta.
Cunninghamiana, *Miq. l. c.* 56. t. 6 A.—Austral.
Decaisneana, *F. Muell. Fragm.* i. 61.—Austral.
decussata, *Benth. Fl. Austral.* vi. 200.—Austral.
Deplancheana, *Miq. in DC. Prod.* xvi. II. 342.—N. Caled.
distyla, *Vent. Jard. Cels*, 62. t. 62.—Austral.
Drummondiana, *Miq. Rev. Crit. Casuar.* 26. t. 1 D.—Austral.
dumosa, A. Cunn. ex Miq. in DC. Prod. xvi. II. 338 = distyla.
echinata, R. Br. ex Benth. Fl. Austral. vi. 202 = thuyoides.
equisetifolia, Blanco, Fl. Filip. ed. I. 661 = sumatrana.
equisetifolia, *Linn. Amoen. Acad.* iv. 143; *Forst. Char. Gen.* 103. t. 52.—Malaya; Ins. Pacif.

CASUARINA:—
excelsa, Dehnhardt, ex Miq. Rev. Crit. Casuar. 23. t. 1 F = equisetifolia.
excelsa, Salisb. Prod. 2 = stricta.
Fraseriana, *Miq. Rev. Crit. Casuar.* 59. t. 6. D.—Austral.
glauca, *Sieber, in Spreng. Syst.* iii. 803.—Austral.
Gunnii, Hook. f. ex Miq. in Neder. Kruidk. Arch. iv. (1859) 100 = stricta.
hexagona, *Dehnhardt, in Revista Napolitana*, i. III. 176.—Austral.
Huegeliana, *Miq. in Lehm. Pl. Preiss.* i. 640.—Austral.
humilis, Otto & Dietr. Allg. Gartenz. ix. (1841) 163 = Fraseriana.
incana, A. Cunn. ex Benth. Fl. Austral. vi. 197 = equisetifolia.
indica, Pers. Syn. ii. 531 = equisetifolia.
inophloia, *F. Muell. & Bailey, in Melb. Chemist* (Apr. 1882).—Austral.
Junghuhniana, *Miq. Pl. Jungh.* 7.—Malaya.
lateriflora, Poir. Encyc. Suppl. ii. 658 = equisetifolia.
Lehmanniana, Miq. in Lehm. Pl. Preiss. i. 639 = humilis.
lepidophloia, *F. Muell. Fragm.* x. 115.—Austral.
leptoclada, Miq. Rev. Crit. Casuar. 41. t. 4. C = suberosa.
leucodon, *Poiss. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 52.—N. Caled.
littoralis, Salisb. Prod. 2 = equisetifolia.
lucida, *Dehnhardt, in Revista Napolitana*, i. III. 175.—Austral.
lugubris, Salisb. Prod. 2 = torulosa.
macrocarpa, A. Cunn. ex Miq. in DC. Prod. xvi. II. 335 = stricta.
Mertensiana, Rupr. ex Miq. l. c. 339 = equisetifolia.
microstachya, *Miq. in Lehm. Pl. Preiss.* i. 642.—Austral.
Miquelii, Hook. f. ex Miq. in DC. Prod. xvi. II. 337 = suberosa.
moesta, F. Muell. ex Miq. in Neder. Kruidk. Arch. iv. (1859) 98 = suberosa.
montana, *Leschen. ex Miq. in Zoll. Syst. Verz. Ind. Archip.* 86.—Malaya.
Muelleriana, Miq. in Neder. Kruidk. Arch. iv. (1859) 99 = distyla.
muricata, Herb. Paris. ex Miq. in DC. Prod. xvi. II. 335 = montana.
muricata, Roxb. Hort. Beng. 66; Fl. Ind. iii. 519 = equisetifolia.
nana, A. Cunn. ex Benth. Fl. Austral. vi. 200 = humilis.
nana, Schlecht. in Linnaea, xx. (1847) 574 = Fraseriana.
nana, *Sieber, ex Spreng. Syst.* iii. 804.—Austral.
nodiflora, *Forst. f. Prod. Fl. Ins. Austr.* 64.—Ins. N. Hebrid.
obesa, Miq. in Lehm. Pl. Preiss. i. 643 = Huegeliana.
obtusa, Hort. ex Miq. in DC. Prod. xvi. II. 334 = glauca.
oxyclada, Miq. l. c. 344 = acuaria.
paludosa, Sieber, ex Spreng. Syst. iii. 803 = distyla.
Pauper, *F. Muell. ex Miq. in Neder. Kruidk. Arch.* iv. (1859) 100.—Austral.
Preissiana, Miq. in Lehm. Pl. Preiss. i. 640 = humilis.
pumila, Otto & Dietr. in Allg. Gartenz. ix. (1841) 162 = distyla.
quadridentata, Desf. Tabl. Cat. Hort. Par. ed. II. 246 = quadrivalvis.
quadrivalvis, Labill. Nov. Holl. Pl. ii. 67. t. 218 = stricta.
ramuliflora, Otto & Dietr. Allg. Gartenz. ix. (1841) 163 = humilis.
repens, Hort. ex Hoffmgg. Verz. Pfl. Nachtr. 83 = equisetifolia.
rigida, Miq. in DC. Prod. xvi. II. 337 = distyla.
rigida, Miq. Rev. Crit. Casuar. 61. t. 7. D = suberosa.
rugosa, *Roxb. Hort. Beng.* 66, nomen.—Ind. or.?
Rumphiana, *Miq. in Flora*, xlviii. (1865) 23, 38.—Amboina.
selaginoides, Miq. in Lehm. Pl. Preiss. i. 643 = humilis.
sparsa, Tausch, in Flora, xxii. (1839) 480 = equisetifolia?
stricta, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 320.—Austral.
stricta, Miq. in DC. Prod. xvi. II. 336 = distyla.

CASUARINA:—
suberosa, *Otto & Dietr. Allg. Gartenz.* (1841) 155.—Austral.
sumatrana, *Jungh. in Hoev. & De Vriese, Tijdschr.* xi. (1844) 115.—Sumatra.
tenuissima, Sieber, ex Spreng. Syst. iii. 804 = torulosa.
tephrosperma, Lehm. Hort. Hamb. ex Miq. Rev. Crit. Casuar. 31 = humilis.
thuyoides, *Miq. in Lehm. Pl. Preiss.* i. 641.—Austral.
tortuosa, Henry, in Nov. Act. Nat. Cur. xx. 1. (1847) 217. t. 19 = stricta?
torulosa, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 320.—Austral.
torulosa, Miq. Rev. Crit. Casuar. 75. t. 11 B = glauca.
torulosa, Miq. in Lehm. Pl. Preiss. i. 639 = Fraseriana.
trichodon, *Miq. l. c.* 641.—Austral.
truncata, Willd. Enum. Hort. Berol. Suppl. 63 = equisetifolia?
verticillata, Lam. Encyc. ii. 502 = nodiflora.
Webbiana, *Miq. in Flora*, xlviii. (1865) 33.—Borneo; N. Caled.?

CATABROSA, Beauv. Agrost. 97. t. 19. f. 8 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1186.
agrostidea, Presl, Fl. Sicul. p. xliv = Airopsis agrostidea.
algida, Fries, Nov. Fl. Suec. Mant. iii. 174 = Phippsia algida.
altaica, Boiss. Fl. Orient. v. 579 = Colpodium altaicum.
antarctica, *Hook. f. Fl. Antarct.* 102.—N. Zel.
aquatica, *Beauv. Agrost.* 97. t. 19. f. 8.—Reg. bor. temp.
Balansae, *Boiss. in Bull. Soc. Bot. Fr.* xxi. (1874) 15.—As. Min.
Capusii, *Franch. in Ann. Sc. Nat.* Sér. VI. xviii. (1884) 272.—Turkest.
concinna, T. Fries, in Ofver. Vet. Akad. xxvi. (1869) 140 = Phippsia algida?
distans, Link, ex Heynh. Nom. ii. 126 = Glyceria distans.
elodes, Auct. = helodes.
Engelmanni, Link, ex Heynh. l. c. = Glyceriae sp.?—Hab.?
fibrosa, *Boiss. Fl. Orient.* v. 577.—Armenia.
frigida, *Phil. Fl. Atac.* 55.—Chili.
glaucescens, Phil. in Anal. Univ. Chil. (1873) 569 = = Glyceria glaucescens.
helodes, Roem. & Schult. Syst. ii. 696 = Poa nemoralis.
humilis, *Trin. Fund. Agrost.* 136.—Persia; Reg. Casp.
latifolia, Fries, Summ. Veg. Scand. i. 245 = Arctagrostis latifolia.
magellanica, Hook. f. Fl. Antarct. 387 = Glyceria magellanica.
micrantha, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 426 = Eragrostis interrupta.
minuta, Trin. Fund. Agrost. 136 = Aira minuta.
ochroleuca, Dum. Obs. Gram. Belg. 108 = aquatica.
parviflora, *Boiss. Fl. Orient.* v. 578.—Persia.
pontica, *Balansa, in Bull. Soc. Bot. Fr.* xxi. (1874) 15.—As. Min.
pumila, Roem. & Schult. Syst. ii. 696 = Poa annua.
scabra, *Kunth, Rev. Gram.* i. 119.—Hab.?
stricta, *Link, ex Steud. Nom.* ed. II. i. 310.—Hab.?
tenuifolia, *J. Presl, in Rel. Haenk.* i. 256.—Chili.
vacillans, T. Fries, in Ofver. Vet. Akad. Stockh. xxvi. (1869) 142 = Phippsia algida.
variegata, *Boiss. Diagn.* Ser. i. v. 71.—As. Min.
versicolor, Boiss. Fl. Orient. v. 579 = Colpodium Steveni.
verticillata, *Beauv. Agrost.* 157, nomen.—Hab.?
vilfoidea, Anders. ex Malm. in Ofver. Vet. Akad. Stockh. xix. (1862) 254 = Glyceria vilfoidea.
violacea, Boiss. Diagn. Ser. I. vii. 126 = Colpodium violaceum.
viridula, Roem. & Schult. Syst. ii. 697 = Aira viridula.

CATABROSIA, Roem. & Schult. Syst. ii. 696 (1817) = **Catabrosa**, Beauv. (Gramin.).

CATACHAENIA, Griseb. in Goett. Abh. ix. (1861) 54 = **Miconia**, Ruiz & Pav. (Melastom.).
alternifolia, Griseb. l. c. 55 = M. Grisebachii.
parviflora, Griseb. Cat. Pl. Cub. 102 = M. Grisebachii.
rufa, Griseb. l. c. = M. rufa.

CATACHAETUM, Hoffmgg. Verz. Orch. 22 (1842); ex Linnaea, xvi. (1842) Litt. 231 = **Catasetum**, Rich. (Orchid.).

CATACLINE, Edgew. in Journ. As. Soc. Bengal, xvi. (1847) 1214 = **Tephrosia**, Pers. (Legumin.).
sericea, Edgew. l. c. = T. sericea.

CATACOMA, Walp. Rep. i. 243 (1842) = CATOCOMA, Benth. = **Bredemeyera**, Willd. (Polygal.).

CATAGYNA, Beauv. in Lestib. Essai Fam. Cyp. 26 (1819) = **Eriospora**, Hochst. (Cyperac.).

CATAKIDOZAMIA, T. Hill, in Gard. Chron. (1865) 1107 = **Macrozamia**, Miq. (Cycad.).
Hopei, T. Hill, l. c. = M. Perowskiana.
Macleayi, Hort. ex Miq. in Verh. Akad. Wetensch. II. iii. (1869) 56 = M. Perowskiana.

CATALIUM, Buch.-Ham. ex Wall. Cat. sub n. 4880 (1831?) = **Carallia**, Roxb. (Rhizoph.).
nitidum, Buch.-Ham. l. c. = Carallia lanceaefolia.

CATALPA, Scop. Introd. 170 (1777). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1041.
CATALPIUM, Rafin. Princ. Somiol. 27 (1814); Fl. Ludov. 139 (1817).
CUMBALU, Adans. Fam. ii. 199 (1763).
bignonioides, *Walt. Fl. Carol.* 64.—Am. bor.
Bungei, *C. A. Mey. in Bull. Sc. Acad. Pétersb.* ii. (1837) 51.—China.
cassinoides, *Spreng. Syst.* i. 70.—Bras.
communis, Dum.-Cours. Bot. Cult. ed. II. iii. 242 = bignonioides.
cordifolia, Jaume, in Duham. Arb. ed. Nov. ii. 13. t. 5 = bignonioides.
hirsuta, *Spreng. Syst.* i. 70.—Bras.
Kaempferi, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 142.—Japon.
longisiliqua, Cham. in Linnaea, vii. (1832) 720 = longissima.
longissima, *Sims, Bot. Mag.* sub t. 1094.—Ind. occ.
microphylla, *Spreng. Syst.* i. 70.—Ins. S. Doming.
ovata, G. Don, Gen. Syst. iv. 230 = Kaempferi.
Pottsii, *Seem. in Otto & Dietr. Allg. Gartenz.* xix. (1851) 321.—Mexic.
punctata, *Griseb. Cat. Pl. Cub.* 192.—Cuba.
purpurea, *Griseb. l. c.*—Cuba.
speciosa, *Warder, ex Engelm. in Coult. Bot. Gaz.* v. (1880) 1.—Am. bor.
syringaefolia, Bunge, Enum. Pl. Chin. Bor. 45 = Bungei.
syringifolia, Sims, Bot. Mag. t. 1094 = bignonioides.
ternifolia, Cav. Desc. 26 = bignonioides.

CATALPIUM, Rafin. Princ. Somiol. 27 (1814); Fl. Ludov. 139 (1817) = **Catalpa**, Scop. (Bignoniac.).
amena, Rafin. Fl. Ludov. 139 = Catalpa bignonioides.

CATAMIXIS, Thoms. in Journ. Linn. Soc. ix. (1865) 342. t. 4. *COMPOSITAE*, Benth. & Hook. f. ii. 494.
baccharoides, *Thoms. l. c.* 343. t. 4.—Himal. bor. occ.

CATANANCHE, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 505.
PIPTOCEPHALUM, Sch. Bip. in Bonplandia (1860) 369.
arenaria, *Coss. & Dur. in Bull. Soc. Bot. Fr.* ii. (1855) 253.—Algeria.
caerulea, Georgi, Beschr. Russ. Reich. iii. IV. 1217 = Centaurea pulchella.
caerulea, *Linn. Sp. Pl.* 812.—Reg. Mediterr. occ.
caespitosa, *Desf. Fl. Atlant.* ii. 238.—Afr. bor.
carpholepis, Nym. Consp. 472 = caerulea.
graeca, Bory & Chaub. Nouv. Fl. Pélop. 55 = Hymenonema laconicum.
graeca, Linn. Sp. Pl. 813 = Hymenonema Tournefortii.
lutea, *Linn. l. c.* 812.—Reg. Mediterr.
montana, *Coss. & Dur. in Bull Soc. Bot. Fr.* iii. (1856) *Bibl.* 743.—Afr. bor.
propinqua, *Pomel, Nouv. Mat. Fl. Atl.* 19.—Algeria.
Webbii, Sch. Bip. in Bonplandia, viii. (1860) 369 = montana.

CATANGA, Steud. Nom. ed. II. i. 311 (1840), sphalm. = **Cananga**, Aubl.

CATAPODIUM, Link, Hort. Berol. i. 380, 44 (1827) = **Festuca**, Linn. (Gramin.).
aquaticum, Willk. in Willk. & Lange, Prod. Fl. Hisp. i. 77 = Catabrosa aquatica.
durum, Link, in Linnaea, xvii. (1843) 399 = Sclerochloa dura.
fusiforme, Nees, ex Benth. in Journ. Linn. Soc. xix. (1881) 128 = Tripogon bromoides.
Halleri, Reichb. Fl. Germ. Excurs. 140 = F. Poa.
loliaceum, Link, Hort. Berol. i. 45 = Desmazeria loliacea.
nepalensis, Link, ex Steud. Nom. ed. II. i. 311 (quid?).
pungens, Boiss. Fl. Orient. v. 635.—Afghan.
Salzmanni, Boiss. l. c. 634.—Reg. Mediterr.
siculum, Link, Hort. Berol. ii. 194 = Desmazeria sicula.
tuberculosum, Moris, in Atti 2da Riun. Sc. Ital. (1841) 481 = F. tuberculosa.

CATAPPA, Gaertn. Fruct. ii. 206. t. 127 (1791) = **Terminalia**, Linn. (Combretac.).
Benzoin, Gaertn. l. c. 206 = T. angustifolia.
brasiliensis, Raddi, in Mem. Mod. xviii. Fis. (1820) 405 = T. januarensis.
guyanensis, Gaertn. f. Fruct. iii. 207 = T. mauritiana.
guyanensis, Gaertn. f. l. c. t. 217 = T. Tanibouca.
mauritiana, Gaertn. f. l. c. 207 = T. mauritiana.

CATAPUNTIA, Muell. Arg. in DC. Prod. xv. II. 1020 (1866), sphalm. = seq.

CATAPUTIA, Ludw. Gen. 81 (1760) = **Ricinus**, Tourn. (Euphorbiac.).
major, Ludw. Ectyp. t. 7 = R. communis.
minor, Ludw. Gen. 81 = R. communis.

CATARIA, Adans. Fam. ii. 192 (1763) = **Nepeta**, Linn. (Labiat.).
canescens, Moench, Meth. 388 = N. orientalis.
Nepetella, Moench, l. c. = N. Nepetella.
nuda, Moench, l. c. = N. nuda.
paniculata, Moench, Meth. Suppl. 134 = N. nuda.
tomentosa, Gilib. Fl. Lituan. i. 78 = N. Cataria.
violacea, Moench, Meth. 388 = N. nuda.
vulgaris, Moench, l. c. 387 = N. Cataria.

CATAS, Domb. ex Lam. Encyc. ii. 354 (1786) = **Embothrium**, Forst. (Proteac.).
grandiflora, Juss. ex Lam. Encyc. ii. 354 = E. grandiflorum.
parviflora, Juss. l. c. 355 = E. coccineum.

CATASETUM, Rich. ex Kunth, Syn. Pl. Aequin. i. 330 (1822). *ORCHIDEAE*, Benth. & Hook. f. iii. 551.
CATACHAETUM, Hoffmgg. Verz. Orch. 22 (1842). Cf. Linnaea, xvi. (1842) Litt. 231.
CLOWESIA, Lindl. Bot. Reg. xxix. (1843) Misc. 25. t. 39.
CUCULINA, Rafin. Fl. Tellur. iv. 49 (1836).
MONACHANTHUS, Lindl. Bot. Reg. sub t. 1538 (1832).
MYANTHUS, Lindl. Bot. Reg. sub t. 1538 (1832).
abruptum, *Hook. Bot. Mag.* t. 3929.—Bras.
adnatum, Steud. Nom. ed. II. i. 311, sphalm. = atratum.
albovirens, *Rodrig. Orch. Nov.* i. 129.—Bras.
atratum, *Lindl. Bot. Reg.* (1838) *Misc.* 61.—Bras.
Baraquinianum, *Lem. Illustr. Hortic.* ix. (1862) *Misc.* 81.—Bras.
barbatum, *Lindl. Bot. Reg.* (1844) *Misc.* 38.—Bras.
bicolor, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 337.—N. Granat.
calceolatum, *Lem. Jard. Fleur.* i. (1851) *Misc.* 45.—Guatemala.
callosum, *Lindl. Bot. Reg.* (1840) *Misc.* 77.—Venezuela.
carunculatum, *Reichb. f. & Warsc. in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 274.—Peruv.
cassideum, *Linden & Reichb. f. Xen. Orch.* ii. 171.—Bras.
cernuum, *Reichb. f. in Walp. Ann.* vi. 570.—Bras.
Christyanum, *Reichb. f. in Gard. Chron.* (1882) i. 588.—Hab.?
ciliatum, *Rodrig. Orch. Nov.* i. 130.—Bras.
Claveringi, Lindl. ex Van Geel, Sert. Bot. ii. (1829) t. 3 = macrocarpum.
cornutum, *Lindl. Bot. Reg.* (1840) *Misc.* 77.—Guiana.

CATASETUM:—
craniomorphum, *Hoffmgg. ex Heynh. Nom.* ii. 126, nomen.—Hab.?
cristatum, *Lindl. Bot. Reg.* t. 966.—Bras.
deltoideum, *Lindl. l. c.* xxvii. (1840) *Misc.* 157.—Guiana.
dilectum, *Reichb. f. Beitr. Orch. Centr. Am.* 73.—Costa Rica.
discolor, *Lindl. Bot. Reg.* (1844) *Misc.* 34.—Bras.
expansum, *Reichb. f. Otia Bot. Hamb.* 9.—Ecuador.
fimbriatum, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 124.—Hab.?
fimbriatum, Reichb. f. in Saund. Refug. Bot. t. 83 = discolor.
floribundum, Hook. Exot. Fl. ii. 151 = macrocarpum.
fuliginosum, *Lindl. Bot. Reg.* (1841) *Misc.* 78.—Hab.?
glaucoglossum, *Reichb. f. in Gard. Chron.* (1885) II. 552.—Mexic.
globiflorum, *Hook. Bot. Mag.* t. 3942.—Bras.
Gnomus, *Linden & Reichb. f. Xen. Orch.* ii. 171.—Bras.
heteranthum, Rodrig. Orch. Nov. i. 127 = C. Gnomus.
Hookeri, *Lindl. Bot. Reg.* sub t. 840.—Bras.
inapertum, Steud. Nom. ed. II. i. 311, sphalm. = purum.
incurvum, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxii. (1854) 178.—Peruv.
integerrimum, Hook. Bot. Mag. t. 3823 = maculatum.
labiatum, *Rodrig. Orch. Nov.* ii. 218.—Bras.
laminatum, *Lindl. in Ann. Nat. Hist.* iv. (1840) 384.—Mexic.
lanciferum, *Lindl. Bot. Reg.* xxvii. (1841) 5. f. 5.—Bras.
Landsbergii, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 156.—Venezuela.
lituratum, *Hoffmgg. Verz. Orch.* 22; *ex Linnaea*, xvi. (1842) *Litt.* 231.—Bras.
longifolium, *Lindl. Bot. Reg.* (1839) *Misc.* 94.—Guiana.
luridum, *Lindl. Gen. et Sp. Orch.* 156.—Bras.
macrocarpum, *Rich. ex Kunth, Syn. Pl. Aequin.* i. 330.—Guiana.
macroglossum, *Reichb. f. in Gard. Chron.* (1877) II. 552.—Ecuador.
maculatum, *Kunth, Syn. Pl. Acquin.* i. 331.—N. Granat.
medium, *Reichb. f. in Gard. Chron.* (1885) II. 6.—Hab.?
mentosum, *Lem. Jard. Fleur.* iii. (1852–53) *Misc.* 65.—Bras.
micranthum, *Rodrig. Orch. Nov.* ii. 219.—Bras.
Milleri, *Lodd. ex Lindl. Bot. Reg.* (1838) *Misc.* 80.—Bras.
Naso, *Lindl. Bot. Reg.* (1843) *Misc.* 71.—Venezuela.
ochraceum, *Lindl. l. c.* (1844) *Misc.* 44.—N. Granat.
Oerstedii, *Reichb. f. in Bonplandia*, iii. (1855) 218.—Nicaragua.
ollare, *Linden, Illustr. Hortic.* xxviii. (1881) *Misc.* 80.—Bras.
pallidum, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 217.—Bras.
Phasma, *Reichb. f. in Gard. Chron.* (1877) II. 488.—Bras.
pileatum, *Reichb. f. l. c.* (1882) I. 492.—Venezuela.
planiceps, *Lindl. Bot. Reg.* (1843) t. 9.—Guiana.
poriferum, *Lindl. l. c.* (1838) *Misc.* 89.—Guiana.
proboscideum, *Lindl. l. c.* (1839) *Misc.* 86.—Bras.
purpurascens, *Hoffmgg. ex Heynh. Nom.* ii. 126, nomen.—Bras.
purum, *Nees & Sinning, Pl. Hort. Bonn. Nutrit.* i. 1.—Bras.
recurvatum, *Link, Klotzsch & Otto, Ic. Pl.* ii. 105. t. 42.—Guiana.
Regnellii, *Rodrig. Orch. Nov.* ii. 219.—Bras.
rivularium, *Rodrig. l. c.* i. 130.—Bras.
roseo-album, Lindl. Bot. Reg. (1840) Misc. 65 = discolor.
roscum, *Reichb. f. in Gard. Chron.* (1872) 1003.—Bras.
roseum, *Rodrig. Orch. Nov.* i. 128.—Bras.
Russellianum, *Hook. Bot. Mag.* t. 3777.—Guatemala.
saccatum, *Lindl. Bot. Reg.* (1840) *Misc.* 76.—Peruv.
sanguineum, *Lindl. & Paxt. Flow. Gard.* ii. (1851–52) 168.—Am. centr.
Scurra, *Reichb. f. in Gard. Chron.* (1872) 1003.—Guiana.
semiapertum, Hook. Exot. Fl. iii. 213 = purum.
serratum, Lindl. Bot. Reg. (1847) t. 24 = viridi-flavum.
spinosum, Lindl. l. c. (1840) Misc. 65 = barbatum?

CATASETUM:—
squalidum, *Hoffmgg. Verz. Orch.* 23; *ex Linnaea*, xvi. (1842) *Litt.* 231.—Bras.
tabulare, *Lindl. Bot. Reg.* xxx. (1840) *Misc.* 40.—N. Granat.
tigrinum, *Reichb. f. in Gard. Chron.* (1881) I. 40.—Hab.?
thylaciochilum, *Lem. Illustr. Hortic.* iii. (1856) *Misc.* 90.—Mexic.
tricolor, *Reichb. f. in Hamb. Gartenz.* xiii. (1857) 313.—Guatemal.
tridentatum, Hook. Exot. Fl. ii. 90, 91 = macrocarpum.
trifidum, *Hook. Bot. Mag.* t. 3262.—Ind. occ.
trimerochilum, Lem. Illustr. Hortic. ix. (1862) t. 374 = Mormodes lineatum.
triodon, *Reichb. f. in Otto, Hamb. Gartenz.* (1857) 313.—Guatemal.
triste, Reichb. f. in Bonplandia, iii. (1855) 218 = Hookeri.
Trulla, *Lindl. Bot. Reg.* (1840) *Misc.* 75.—Am. trop.
turbinatum, *Hoffmgg. ex Heynh. Nom.* ii. 126, nomen.—Bras.
umbrosum, *Rodrig. Orch. Nov.* i. 129.—Bras.
variabile, *Rodrig. l. c.* ii. 217.—Bras.
violascens, *Reichb. f. & Warsc. in Bonplandia*, ii. (1854) 97.—N. Granat.
viride, *Lindl. Bot. Reg.* (1841) *Misc.* 79.—Bras.
viridi-flavum, *Hook. Bot. Mag.* t. 4017.—Am. centr.
Wailesii, Hook. l. c. t. 3937 = maculatum.
Warczewitzii, *Lindl. & Paxt. Flow. Gard.* i. (1850–51) 45. f. 29.—Panama.

CATATEROPHORA, Steud. Nom. ed. II. i. 311 (1840), sphalm. = seq.

CATATHEROPHORA, Steud. in Flora, xii. (1829) 465 = **Pennisetum**, Pers. (Gramin.).
hordeiformis, Steud. l. c. = P. macrourum.

CATENARIA, Benth. in Miq. Pl. Jungh. 217, 220 (1852) = **Desmodium**, Desv. (Legumin.).
laburnifolia, Benth. l. c. = D. laburnifolium.

CAEPHA, Leschen. ex Reichb. Handb. 221 (1837) = **Platysace**, Bunge (Umbell.).

CATESBAEA, Linn. Gen. ed. I. 336 (1737). *RUBIACEAE*, Benth. & Hook. f. ii. 78.
campanulata, *Sagra, ex DC. Prod.* iv. 401.—Cuba.
elliptica, Spreng. ex. DC. l. c. = Exostemma Vavassorii.
erecta, *Moç. & Sesse, ex DC. l. c.*—Mexic.
Grayii, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 503.—Cuba.
holacantha, *Griseb. Cat. Pl. Cub.* 122.—Cuba.
inermis, Spreng. Syst. i. 416 = Rondeletia buxifolia.
latifolia, *Lindl. Bot. Reg.* t. 858.—Ind. occ.
Lindeniana, *Baxt. ex Loud. Hort. Brit. Suppl.* iii. 511, nomen.—Hab.?
longiflora, Sw. Prod. Veg. Ind. Occ. 30 = spinosa.
longispina, *A. Rich. Fl. Cub. Fanerog.* ii. 12.—Cuba.
macrantha, A. Rich. l. c. = spinosa?
macrantha, *Wright, in Sauv. Fl. Cub.* 59.—Cuba.
paralybensis, *Vell. Fl. Flum.* 54; i. t. 145.—Bras.
parviflora, Griseb. Cat. Pl. Cub. 122 = Scolosanthus parviflorus.
parviflora, Lam. Tabl. i. 288. n. 1481 = Scolosanthus versicolor.
parviflora, Spreng. ex Steud. Nom. ed. II. i. 311 = Rondeletia buxifolia.
parviflora, *Sw. Prod. Veg. Ind. Occ.* 30.—Jamaic.
parvifolia, *DC. Prod.* iv. 401.—Ins. S. Doming.
phyllacantha, Griseb. Cat. Pl. Cub. 122 = Phyllacantha Grisebachiana.
spinosa, *Linn. Sp. Pl.* 109.—Ind. occ.
strigosa, Willd. ex Steud. Nom. ed. II. i. 311 = Randia tetrandra.
triacantha, Spreng. Neue Entdeck. iii. 47 = Scolosanthus triacanthus.
Vavassorii, Spreng. Syst. i. 416 = Exostemma Vavassorii.

CATEVALA, Medic. Theodora, 67 (1786) = **Aloë**, Linn. (Liliac.).
arachnoidea, Medic. l. c. 68 = Haworthia arachnoides.
arborescens, Medic. l. c. 67 = A. arborescens.
atroviridis, Medic. l. c. 69 = Haworthia atrovirens.
humilis, Medic. l. c. = A. humilis.
retusa, Medic. l. c. 68 = Haworthia retusa.

CATHA, Forsk. Fl. Aegypt. Arab. 63 (1775). *CELASTRINEAE*, Benth. & Hook. f. i. 361.
METHYSCOPHYLLUM, Eckl. & Zeyh. Enum. 152 (1836).
TRIGONOTHECA, Hochst. in Flora, xxiv. (1841) 662.
edulis, *Forsk. l. c.*—Afr. trop.
Forskalei, A. Rich. Tent. Fl. Abyss. i. 134 = edulis.
inermis, J. F. Gmel. Syst. 411 = edulis.
spinosa, Forsk. Fl. Aegypt. Arab. 64 = Celastrus parviflorus.

CATHA, G. Don, Gen. Syst. ii. 11 (1832) = **Celastrus**, Linn.
acuminata, Presl, Bot. Bemerk. 33 = Cel. acuminatus.
alaternifolia, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 98.—Ins. Comor.
articulata, G. Don, Gen. Syst. ii. 10 = Cel. articulatus.
Benthamii, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 310 = Cel. Championi.
buxifolia, G. Don, Gen. Syst. ii. 10 = Cel. buxifolius.
campestris, Presl, Bot. Bemerk. 34 = Cel. pyracanthus.
cassinoides, Webb & Berth. Phyt. Canar. ii. 142 = Cel. cassinoides.
crenata, A. Gray, Bot. U. St. Expl. Exped. i. 288 in textu = Cel. crenatus.
Cunninghamii, Hook. in Mitch. Journ. Trop. Austral. 387 = Cel. Cunninghamii.
cymosa, G. Don, Gen. Syst. ii. 10 = Cel. buxifolius.
decolor, Webb, Fragm. Fl. Aethiop. 60 = Gymnosporia montana.
Dryandri, Lowe, Man. Fl. Madeir. i. 107.—Ins. Madera.
emarginata, G. Don, Gen. Syst. ii. 9 = Gymnosporia emarginata.
europaea, Boiss. Voy. Espagne, ii. 725 = Gymnosporia montana.
excisa, G. Don, Gen. Syst. ii. 9 = Cel. excisa.
fasciculata, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 98.—Malaya.
flexuosa, G. Don, Gen. Syst. ii. 10 = Cel. flexuosus.
fruticosa, Thw. Enum. Pl. Zeyl. 72 = Gymnosporia emarginata.
Grossulariae, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 98.—Madag.
heterophylla, Presl, Bot. Bemerk. 33 = Cel. heterophyllus.
Heyneana, Walp. Rep. i. 532 = Gymnosporia Heyneana.
integer, G. Don, Gen. Syst. ii. 9 = Cel. pyracanthus.
integrifolia, G. Don, l. c. = Cel. integrifolius.
lanceolata, Presl, Bot. Bemerk. 33 = Cel. linearis?
leptopus, Tul. in Ann. Sc. Nat. Sér. IV. viii. (1857) 100.—Madag.
linearis, G. Don, Gen. Syst. ii. 9 = Cel. linearis.
monosperma, Benth. in Hook. Lond. Journ. Bot. i. 483 = Cel. monospermus.
montana, G. Don, Gen. Syst. ii. 10 = Gymnosporia montana.
multiflora, G. Don, l. c. = Cel. multiflorus.
ovata, Walp. Rep. i. 532 = Gymnosporia ovata.
paniculata, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1842) 275 = Cel. paniculatus.
parviflora, G. Don, Gen. Syst. ii. 10 = Cel. parviflorus.
parvifolius, Presl, Bot. Bemerk. 34 = Cel. parvifolius.
patens, Presl, l. c. 33 = Cel. patens.
Pyracantha, G. Don, Gen. Syst. ii. 10 = Cel. pyracanthus.
rigida, G. Don, l. c. 9 = Cel. rigidus.
Rothiana, Walp. Rep. i. 532 = Gymnosporia Rothiana.
rotundifolia, G. Don, Gen. Syst. ii. 10 = Dovyalis celastroides.
rufa, G. Don. l. c. = Gymnosporia rufa.
rupestris, Presl, Bot. Bemerk. 33 = Cel. acuminatus.
senegalensis, G. Don, Gen. Syst. ii. 10 = Gymnosporia montana.
senegalensis, Webb, Fragm. Fl. Aethiop. 61 = Cel. senegalensis.
spathulata, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 275.—Ind. or.
spathyphylla, Presl, Bot. Bemerk. 33 = Cel. buxifolius.
trigyna, Presl, Bot. Bemerk. 33 = Gymnosporia trigyna.
venenata, Presl, l. c. 34 = Cel. buxifolius.
vitiensis, A. Gray, Bot. U. St. Expl. Exped. i. 287. t. 23.—Ins. Fiji.
Wallichii, G. Don, Gen. Syst. ii. 10 = Gymnosporia Wallichiana.
zeylanica, G. Don, l. c. = Scutia Commersonii.

CATHANTHES, Rich. in Mém. Mus. Par. i. (1815) 365 = **Tetroncium**, Willd. (Naiad.).

CATHARANTHUS, G. Don, Gen. Syst. iv. 95 (1836) = **Vinca**, Linn. (Apocynac.).
pusillus, G. Don, l. c. = V. parviflora.
roseus, G. Don, l. c. = V. rosea.

CATHARTOCARPUS, Pers. Syn. i. 459 (1805) = **Cassia**, Linn. (Legumin.).
Atteleh, G. Don, Gen. Syst. ii. 453 = Cassia Arereh.
bacillaris, Pers. Syn. i. 459 = Cassia bacillaris.
Bacillus, Lindl. Bot. Reg. t. 881 = Cassia bacillaris.
brasilianus, Jacq. Fragm. 59. t. 85. f. 3 = Cassia grandis.
Brewsteri, F. Muell. Fragm. i. 110 = Cassia Brewsteri.
conspicuus, G. Don, in Loud. Hort. Brit. 167 = Cassia Sieberiana.
excelsus, G. Don. Gen. Syst. ii. 453 = Cassia Fistula.
ferrugineus, G. Don, l. c. = Cassia ferruginea.
Fistula, Pers. Syn. i. 459 = Cassia Fistula.
fistuloides, G. Don, in Loud. Hort. Brit. 167 = Cassia Fistula.
grandis, Pers. Syn. i. 459 = Cassia grandis.
Humboldtianus, Loud. Hort. Brit. 167 = Cassia spectabilis.
javanicus, Pers. Syn. i. 459 = Cassia javanica.
laetus, G. Don, Gen. Syst. ii. 454 = Cassia laeta.
marginatus, G. Don, l. c. 453 = Cassia marginata.
moschatus, G. Don, l. c. = Cassia moschata.
nodosus, Steud. Nom. ed. II. i. 311 = Cassia nodosa.
rhombifolius, G. Don, Gen. Syst. ii. 453 = Cassia Fistula.
Roxburghii, Loud. Hort. Brit. 167 = Cassia marginata.
Sieberianus, G. Don, Gen. Syst. ii. 453 = Cassia Sieberiana.
speciosus, G. Don, l. c. = Cassia spectabilis.
Trinitatis, G. Don, in Loud. Hort. Brit. 167 = Cassia spectabilis.

CATHARTOLINUM, Reichb. Handb. 306 (1837) = **Linum**, Linn.
agreste, Reichb. l. c. 307 = L. angustifolium.
alternum, Reichb. l. c. = L. strictum.
bicolor, Reichb. l. c. = L. bicolor.
corymbulosum, Reichb. l. c. = L. strictum.
gallicum, Reichb. l. c. = L. gallicum.
pratense, Reichb. l. c. = L. catharticum.
salsoloides, Reichb. l. c. = L. salsoloides.
sardoum, Reichb. l. c. = L. Muelleri.
spicatum, Reichb. Ic. Fl. Germ. xvi. 62 = L. strictum.
strictum, Reichb. l. c. xvi. t. 327 = L. strictum.
suffruticosum, Reichb. Handb. 307 = L. suffruticosum.
tenuifolium, Reichb. l. c. = L. tenuifolium.
virginianum, Reichb. l. c. = L. virginianum.

CATHASTRUM, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 448. *CELASTRINEAE*, Benth. & Hook. f. i. 362.
capense, *Turcz. l. c.*—Afr. austr.

CATHCARTIA, Hook. f. in Bot. Mag. t. 4596 (1851). *PAPAVERACEAE*, Benth. & Hook. f. i. 52.
integrifolia, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 310.—China.
villosa, *Hook. f. in Bot. Mag.* t. 4596.—Reg. Himal.

CATHEA, Salisb. in Trans. Hort. Soc. i. (1812) 300 = **Calopogon**, R. Br. (Orchid.).
pulchella, Salisb. l. c. = Calop. pulchellus.

CATHEDRA, Miers, in Ann. Nat. Hist. Ser. II. vii. (1851) 452, 457. *OLACINEAE*, Benth. & Hook. f. i. 348.
Diplocrater, Benth. in Hook. Kew Journ. iii. (1851) 367.
acuminata, *Miers, l. c.* Ser. III. iv. (1859) 361.—Bras.
crassifolia, *Miers, l. c.*—Bras.
Gardneriana, *Miers, l. c.* Ser. II. vii. (1851) 459.—Bras.
rubricaulis, *Miers, l. c.* 458.—Bras.

CATHESTECUM, J. Presl, in Rel. Haenk. i. 294. t. 42 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1122.
erectum, *Vasey & Hack. in Bull. Torrey Bot. Club.* xi. (1884) 37. t. 45.—Mexic.
prostratum, *J. Presl, in Rel. Haenk.* i. 295. t. 42.—Mexic.

CATHETOSTEMMA, Blume, Rumphia, iv. 30 (1848) = **Hoya**, R. Br. (Asclepiad.).
laurifolium, Blume, l. c. = H. laurifolia.

CATHETUS, Lour. Fl. Cochinch. 607 (1790) = **Phyllanthus**, Linn. (Euphorbiac.).
fasciculata, Lour. l. c. 608 = P. cochinchensis.

CATHISSA, Salisb. Gen. Pl. Fragm. 34 (1866) = **Ornithogalum**, Tourn. (Liliac.).
concinna, Salisb. l. c. = O. unifolium.
unifolia, Salisb. l. c. = O. unifolium.

CATHORMION, Hassk. Retzia, i. 231 (1855) = **Pithecolobium**, Mart. (Legumin.).
Junghuhnianum, Hassk. l. c. 232 = P. Junghuhnianum.
moniliferum, Hassk. l. c. 231 = P. moniliferum.

CATIMBIUM, Juss. Gen. 62 (1789) = **Alpinia**, Linn. (Scitamin.).
erectum, Juss. ex Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 342 = A. calcarata.
exaltatum, Mirb. ex Steud. Nom. ed. II. i. 311 = Renealma exaltata.
nutans, Juss. ex Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 342 = A. nutans.

CATINGA, Aubl. Pl. Gui. i. 511. t. 203 (1775) = **Eugenia**, Linn. (Myrtac.).
aromatica, Aubl. l. c. f. 2 = E. aromatica.
moschata, Aubl. l. c. f. 1 = E. Catinga.

CATOBLASTUS, H. Wendl. in Bonplandia, viii. (1860) 104, 106. *PALMAE*, Benth. & Hook. f. iii. 901.
maynensis, *Drude, in Mart. Fl. Bras.* iii. II. 544.—Peruv.
praemorsus, *H. Wendl. in Bonplandia*, viii. (1860) 104.—N. Granat.
pubescens, *H. Wendl. l. c.*—Bras.

CATOCOMA, Benth. in Hook. Journ. Bot. iv. (1842) 101 = **Bredemeyera**, Willd. (Polygal.).
altissima, Poepp. & Endl. Nov. Gen. et Sp. iii. 65. t. 273 = B. altissima.
brevifolia, Benth. in Hook. Journ. Bot. iv. (1842) 102 = B. brevifolia.
emarginata, Spruce, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 50 = B. altissima.
floribunda, Benth. in Hook. Journ. Bot. iv. (1842) 102 = B. floribunda.
Kunthiana, Benth. l. c. 103 = B. Kunthiana.
laurifolia, Benth. l. c. = B. laurifolia.
lucida, Benth. l. c. 101 = B. lucida.
Mansonii, Klotzsch, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 51 = B. lucida.
mollis, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) v. 133 = B. altissima.
myrtifolia, Spruce, ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 51 = B. myrtifolia.
parviflora, Spruce, ex A. W. Benn. l. c. 51 = B. parviflora.

CATOCORYNE, Hook. f. in Benth. & Hook. f. Gen. i. 765 (1867). *MELASTOMACEAE*, Benth. & Hook. f. l. c.
linneoides, *Hook. f. l. c.*—Peruv.

CATODIACRUM, Dulac, Fl. Hautes-Pyr. 369 (1867) = **Orobanche**, Tourn.
cruentum, Dulac, l. c. 371 = O. cruenta.
foetidum, Dulac, l. c. = O. foetida.
incurvum, Dulac, l. c. 370 = O. caryophyllacea.
luteum, Dulac, l. c. = O. Hederae.
pallidiflorum, Dulac, l. c. = O. pallidiflora.
rubidum, Dulac, l. c. = O. Teucrii.
sparsiflorum, Dulac, l. c. = O. Epithymum.
variegatum, Dulac, l. c. = O. variegata.

CATONIA, P. Br. Hist. Jamaic. 148 (1756); Juss. Gen. 441 (1789). *INC. SED.*
Brownei, *Roem. & Schult. Syst.* iii. 480.—Jamaica.
marginata, Jaume St. Hil. Expos. ii. 347 = Brownei.

CATONIA, Moench, Meth. 535 (1794) = **Crepis**, Linn. (Compos.).
blattarioides, Cass. in Dict. Sc. Nat. v. 274; xxvi. 9 = Crepis blattarioides.
cordifolia, Moench, Meth. 536 = Hieracium amplexicaule.
crocea, Reichb. ex Steud. Nom. ed. II. i. 311 = Crepis Columnae.
sagittata, Moench, Meth. 536 = Crepis blattarioides.

CATONIA, Rafin. Fl. Tellur. ii. 36 (1836) (Quid?).
lantanoides, Rafin. l. c.—Hab.?

CATONIA, Vahl, in Skrivt. Nat. Hist. Selsk. Kiobenh. vi. (1810) 98 = **Erycibe**, Roxb. (Convolvulac.).
elliptica, Vahl, l. c. 101 = E. paniculata.
glauca, Vahl, l. c. 98 = E. paniculata.

CATONIA, Vell. Fl. Flum. 155 (1825); t. iv. t. 18 ('Oct. Mon.' Quid?).
rustica, Vell. l. c.—Bras.

CATOPHERIA, Benth. in Benth. & Hook. f. Gen. ii. 1173 (1876). *LABIATAE*, Benth. & Hook. f. l. c.
capitata, *Benth. l. c.*—Mexic.
chiapensis, *A. Gray, ex Benth. in Hook. Ic. Pl.* sub t. 1215.—Mexic.
spicata, *Benth. l. c.* t. 1215.—N. Granat.

CATOPHRACTES, D. Don, in Proc. Linn. Soc. i. 4 (1839); Trans. Linn. Soc. xviii. (1841) 306. t. 22. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1048.
Alexandri, *D. Don, ll. cc.* 4; 307.—Afr. austr.
Kolbeana, *Harv. Gen. S. Afr. Pl.* ed. II. 276.—Afr. austr.
Welwitschii, *Seem. in Journ. Bot.* iii. (1865) 331.—Afr. trop.

CATOPHYLLUM, Pohl, ex Baker, in Mart. Fl. Bras. vi. II. 222 (1876) = **Mikania**, Willd. (Compos.).
deltoideum, Pohl, ex Baker, l. c. = M. officinalis.
tropaeolifolium, Pohl, ex Baker, l. c. = M. officinalis.

CATOPODIUM, Link, Hort. Berol. i. 44 (1827), sphalm. = **Catapodium**, Link (Gramin.).

CATOPSIS, Griseb. in Goett. Nachr. 1864 (1865) 21. *BROMELIACEAE*, Benth. & Hook. f. iii. 670.
Pogospermum, Brongn. in Ann. Sc. Nat. Sér. V. i. (1864) 237.
Tussacia, Klotzsch, ex Beer, Bromel. 99 (1856).
fulgens, *Griseb. l. c.* 21.—Am. trop.
nitida, *Griseb. Fl. Brit. W. Ind.* 599.—Ind. occ.
nutans, *Griseb. l. c.*—Ind. occ.

CATOSPERMA, Benth. in Hook. Ic. Pl. t. 1028 (1868). *GOODENOVIEAE*, Benth. & Hook. f. ii. 539.
Careyi, *F. Muell. in Melb. Chemist* (Apr. 1884).—Austral.
Muelleri, *Benth. in Hook. Ic. Pl.* t. 1028.—Austral.

CATOSTEMMA, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 365. *MALVACEAE*, Benth. & Hook. f. i. 724.
fragrans, *Benth. l. c.*—Guiana.

CATTLEYA, Lindl. Collect. Bot. tt. 33, 37 (1824). *ORCHIDEAE*, Benth. & Hook. f. iii. 531.
Maelenia, Dum. in Mém. Acad. Brux. ix. (1835) 13. t. 10.
Acklandiae, Planch. in Fl. des Serres, Sér. I. vii. (1851–52) 83 = seq.
Aclandiae, *Lindl. Bot. Reg.* xxvi. (1840) t. 48.—Bras.
acuminata, Beer, Prakt. Orch. 208 = Laelia acuminata.
albida, Beer, l. c. = Laelia albida.
alutacea, *Rodrig. Orch. Nov.* ii. 157.—Bras.
amethystina, Morr. in Ann. Gand. iv. (1848) 217. t. 201 = intermedia.
amethystoglossa, Linden & Reichb. f. ex Warner, Sel. Orch. Ser. I. t. 2 = guttata.
anceps, Beer, Prakt. Orch. 208 = Laelia anceps.
Arembergii, Scheidw. in Otto & Dietr. Allg. Gartenz. xi. (1843) 109 = Loddigesii.
Aucklandiae, Heynh. Nom. ii. 126, sphalm. = Aclandiae.

CATTLEYA :—
Auclandii, Beer, Prakt. Orch. 208, sphalm. = Aclandiae.
aurantiaca, P. N. Don, in Florist's Journ. (1840) 185 = Epidendrum aurantiacum.
aurea, *Linden*, *Illustr. Hortic.* xxx. (1883) t. 493.—N. Granat.
autumnalis, Beer, Prakt. Orch. 208 = Laelia autumnalis.
bicolor, *Lindl. Bot. Reg.* (1836) sub t. 1919.—Am. austr.
Bluntei, *Hort. ex Gard. Chron.* (1876) I. 735.—Hab.
bogotensis, Linden, in Belg. Hortic. xv. (1865) 102 = labiata.
Bowringiana, *Veitch*, *ex Gard. Chron.* (1885) II. 683. —Honduras.
Brabantiae ×, Veitch.—Cf. Gard. Chron. (1864) 413; et Floral Mag. vi. (1867) t. 360.
Brymeriana ×, Reichb. f. in Gard. Chron. (1883) II. 492.
Brysiana, Lem. in Jard. Fleur. iii. (1852–53) t. 275 = Laelia purpurata.
bulbosa, Lindl. Bot. Reg. (1847) t. 42 = Walkeriana.
calummata ×, André, in Rev. Hortic. (1883) 564.
candida, Williams, in Gard. Chron. (1851) 453 = Loddigesii.
Carrierei, Houllet, in Rev. Hortic. (1876) 250 = labiata.
cernua, Beer, Prakt. Orch. 209 = Sophronitis cernua.
Chamberlainiana ×, Reichb. f. in Gard. Chron. (1881) II. 427.
chlorantha, Beer, Prakt. Orch. 209 = Sobralia macrophylla.
chocoënsis, Linden, in Illustr. Hortic. xvii. (1870) 37 = labiata.
cinnabarina, Beer, Prakt. Orch. 209 = Laelia cinnabarina.
citrina, *Lindl. Coll. Bot.* sub t. 37.—Mexic.
coccinea, Lindl. Bot. Reg. sub t. 1919 = Sophronitis grandiflora.
crispa, Beer, Prakt. Orch. 209 = Schomburgkia crispa.
crispa, Lindl. Bot. Reg. t. 1172 = Laelia crispa.
Dawsoni, Warner, Sel. Orch. Pl. i. t. 16 = labiata.
Deckeri, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiii. (1855) 81 = Skinneri.
decora, Beer, Prakt. Orch. 209 = Sobralia decora.
devoniensis ×, Hort. ex Floral Mag. N. S. (1881) sub t. 437 = Laelia elegans?
dolosa, *Reichb. f. Xenia Orchid.* ii. 224, *in synon.*— Bras.
domingensis, Lindl. Gen. et Sp. Orch. 118 = Broughtonia lilacina.
Dominiana ×, Lindl. in Gard. Chron. (1859) 948.
Dormaniana ×, Reichb. f. in Gard. Chron. (1882) I. 216.
Dowiana, *Batem. l. c.* (1866) 922.—Costa Rica.
elatior, Lindl. Gen. et Sp. Orch. 117 = guttata.
Eldorado, Linden, in Fl. des Serres, Sér. II. viii. (1869–70) 13. t. 1826 = labiata.
elegans, Morr. in Ann. Gand. iv. (1848) 93. t. 185 = Laelia elegans.
elegantissima, *Linden*, *Illustr. Hortic.* xxviii. (1881) 80.—Venezuela.
elongata, *Rodrig. Orch. Nov.* i. 72.—Bras.
eximia, *Rodrig. l. c.* 70.—Bras.
exoniensis ×, Reichb. f. in Gard. Chron. (1867) 1144.
fausta ×, Reichb. f. l. c. (1873) 289.
felix ×, Reichb. f. l. c. (1876) II. 68.
flava, Beer, Prakt. Orch. 210 = Laelia flava.
flavida, Klotzsch, in Otto & Dietr. Allg. Gartenz. xxiv. (1856) 73 = luteola.
Forbesii, *Lindl. Coll. Bot.* sub t. 37.—Bras.
fragrans, *Rodrig. Orch. Nov.* i. 72.—Bras.
furfuracea, Beer, Prakt. Orch. 210 = Laelia furfuracea.
fulva, *Beer*, *l. c.* 210.—Hab.?
Galeottiana, Beer, l. c. = Epidendrum Galeottianum.
Gardneriana, Reichb. f. in Gard. Chron. (1870) 1373, in nota = Walkeriana.
Gaskelliana, Hort. Sander, l. c. (1883) I. 243, 310 = labiata.
gigantea, *Hort. ex P. N. Don*, *in Florist's Journ.* (1840) 187.—Hab.?
Gigas, Linden & André, Illustr. Hortic. xx. (1873) 70 = labiata.
gloriosa, Carr. in Rev. Hortic. (1885) 333 = labiata.
Grahami, Lindl. Gen. et Sp. Orch. 116 = Laelia majalis.
grandiflora, Beer, Prakt. Orch. 209 = Sophronitis grandiflora.
granulosa, *Lindl. Bot. Reg.* (1842) t. 1.—Am. austr.
guatemalensis, *T. Moore*, *in Flor. Mag.* (1861) t. 61.— Guatemala.

CATTLEYA :—
guttata, *Lindl. Bot. Reg.* t. 1406.—Bras.
Hardyana, Hort. Sander.—Cf. Gard. Chron. (1883) I. 243 = labiata.
Harrisoniae, Paxt. Mag. Bot. iv. (1838) 247 = Loddigesii.
Harrisoniana, Batem. ex Lindl. in Bot. Reg. sub t. 1919 = Loddigesii.
Harrisonii, Hort. ex Duchartre, in Journ. Soc. Hort. Fr. iii. (1857) 725. t. 7 = praec.
hybrida ×, Veitch, in Gard. Chron. (1863) 602.
ianthina, Hort. ex Gard. Chron. (1853) 247, nomen = intermedia?
imperialis, J. O'Brien, in Gard. Chron. (1883) II. 404 = labiata.
intermedia, *R. Grah. in Bot. Mag.* t. 2851.—Bras.
intricata, *Reichb. f. in Gard. Chron.* (1884) II. 7.— Hab.?
iricolor, *Reichb. f. l. c.* (1874) II. 162.—Hab.?
Isabella, *Reichb. f. in C. Koch & Fint. Wochenschr.* (1859) 336.—Hab.?
isopetala, *Beer*, *Prakt. Orch.* 211.—Hab.?
Karwinskii, Mart. Ausw. Merkw. Pfl. 14. t. 10 = citrina.
labiata, *Lindl. Coll. Bot.* t. 33.—Am. trop.
laelioides, Lem. Jard. Fleur. iii. (1852–53) Misc. 42, 50 = Skinneri.
Lawrenceana, *Reichb. f. in Gard. Chron.* (1885) I. 338.—Guiana.
Lawrenceana, Warsc. ex Reichb. f. in Gard. Chron. (1883) I. 243 = Dowiana.
Leeana, Hort. Sander.—Cf. Gard. Chron. (1883) I. 43 = labiata.
Lemoniana, Lindl. Bot. Reg. (1846) t. 35 = labiata.
Leopoldii, Hort. Versch. ex Illustr. Hortic. i. (1854) Misc. 68; ii. (1855) t. 69 = guttata.
Liliastrum, Beer, Prakt. Orch. 212 = Sobralia Liliastrum.
Lindleyana, *Reichb. f. in C. Koch*, *Allg. Gartenz.* (1857) 118.—Bras.
lobata, Lindl. Gard. Chron. (1848) 403 = Laelia Boothiana.
Loddigesii, *Lindl. Coll. Bot.* t. 37.—Bras.
Lucieniana ×, Reichb. f. in Gard. Chron. (1885) II. 456.
Luddemanniana, Reichb. f. Xenia Orch. i. 29 = labiata.
luteola, *Lindl. in Gard. Chron.* (1853) 774.—Bras.
macrantha, Beer, Prakt. Orch. 212 = Sobralia macrantha.
majalis, Beer, l. c. = Laelia majalis.
Manglesii ×, Reichb. f. in Gard. Chron. (1880) II. 556.
Mardellii ×, Seden, ex Reichb. f. l. c. (1879) I. 234.
marginata, Beer, Prakt. Orch. 212 = Schomburgkia marginata.
marginata, Paxt. Mag. Bot. x. (1843) 265 = Laelia pumila.
maritima, Lindl. Bot. Reg. (1836) sub t. 1919 = intermedia.
Marstersoniae ×, Seden, ex Reichb. f. in Gard. Chron. (1878) II. 556.
Massangeana, Reichb. f. in Gard. Chron. (1883) I. 242 = labiata.
maxima, *Lindl. Gen et Sp. Orch.* 116.—Peruv.
Mendelli, *Hort.*—*Cf. Gard. Chron.* (1883) II. 404.— Hab.?
Meyeri, Regel, in Ind. Sem. Hort. Petrop. (1855) 17. Cf. Linnaea, xxviii. (1856) 369 = luteola.
Mitchelli ×, Reichb. f. in Gard. Chron. (1876) II. 386.
Morganae, Warner, Orch. Album, i. (1882) t. 6 = labiata.
Mossiae, Parker, in Hook. Bot. Mag. t. 3669 = labiata.
Nalderiana, Reichb. f. in Gard. Chron. (1885) I. 756 = labiata.
nobilior, Reichb. f. in Illustr. Hortic. xxx. (1883) t. 485 = Walkeriana.
odoratissima, P. N. Don, in Florist's Journ. (1840) 185 = Sobralia violacea?
ovata, Lindl. Bot. Reg. sub t. 1919 = intermedia.
pallida, Lindl. & Paxt. Flow. Gard. ii. (1851–52) 51 = labiata.
paludosa, Beer, Prakt. Orch. 215 = Sobralia paludosa.
papeiansiana, Morr. in Ann. Gand. i. (1845) 57. t. 5 = Loddigesii.
peduncularis, Beer, Prakt. Orch. 213 = Laelia peduncularis.
Peetersii, André, in Rev. Hortic. (1885) 271 = labiata.
Percivaliana, Hort.—Cf. Gard. Chron. (1883) II. 404 = labiata.

CATTLEYA :—
Perrinii, Endl. in Harting. Parad. Viudob. t. 20 = labiata.
Perrinii, Lindl. Bot. Reg. (1838) t. 2 = Laelia Perrinii.
picturata ×, Reichb. f. in Gard. Chron. (1877) II. 584.
Pinellii, Hort. ex Lindl. in Fl. des Serres, Sér. I. vii. (1851–52) 43 = Laelia pumila.
piperita, *Hort.*—*Cf. Gard. Chron.* (1850) 168, nomen. —Hab.?
porphyroglossum, *Linden & Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 98.—Bras.
porphyrophlebia ×, Reichb. f. in Gard. Chron. (1885) II. 552.
Princeps, *Rodrig. Orch. Nov.* i. 68.—Bras.
pterocarpa, Beer, Prakt. Orch. 209 = Sophronitis pterocarpa.
pumila, Hook. Bot. Mag. t. 3656 = Laelia pumila.
purpurata, Beer, Prakt. Orch. 213 = Laelia purpurata.
purpurina, *Rodrig. Orch. Nov.* ii. 158.—Bras.
quadricolor, Lindl. in Paxt. Flow. Gard. i. 6 = labiata.
Quindois, *Hort.*—*Cf. Gard. Chron.* (1853) 247, nomen. —Hab.?
quinquecolor ×, Dombr. in Floral Mag. t. 511.
reflexa, Parment. ex Walp. Ann. vi. 423 = Laelia crispa.
regalis, Hort.—Cf. Gard. Chron. (1883) II. 404 = labiata?
Regnellii, Warner, Sel. Orch. Ser. II. t. 22 = Schilleriana.
Reineckiana, Reichb. f. in Bonplandia, iv. (1856) 237 = labiata.
resplendens ×, *Reichb. f. in Gard. Chron.* (1885) I. 692.
Roezlii, Reichb. f.—Cf. Gard. Chron. (1882) II. 457 = labiata.
Rollissonii, T. Moore, in Floral Mag. i. (1861) t. 8 = labiata.
rubescens, Beer, Prakt. Orch. 214 = Laelia rubescens.
Ruckeri, *Linden*, *in Belg. Hortic.* xv. (1865) 102.—N. Granat.
Rueckerii, Beer, Prakt. Orch. 214 = Sobralia Rueckeri.
Sanderiana, Hort. Low.—Cf. Gard. Chron. (1882) II. 151 = labiata.
Schilleriana, *Reichb. f. in C. Koch*, *Allg. Gartenz.* (1857) 335.—Bras.
Schofieldiana, *Reichb. f. in Gard. Chron.* (1882) II. 808.—Hab.?
Schroederiana, *Reichb. f. l. c.* (1883) II. 102.—Hab.?
Scita ×, Reichb. f. l. c. (1885) II. 489.
sessilis, Beer, Prakt. Orch. 214 = Sobralia sessilis.
Skinneri, *Batem. Orch. Mex. & Guat.* t. 13.—Guatemala.
speciosissima, Hort.—Cf. Gard. Chron. (1868) 404, in nota = labiata.
spectabilis, Hort.—Cf. Florist (1850) 91 = marginata.
sphenophora, Morr. Ann. Gand. iv. (1848) 17. t. 175 = guttata.
sulfurina, Lem. Illustr. Hortic. i. (1854) Misc. 2 = citrina.
superba, Schomb. ex Lindl. Sert. Orch. t. 22 = violacea.
superbiens, Beer, Prakt. Orch. 214 = Laelia superbiens.
tibicinis, Beer, l. c. 215 = Schomburgkia tibicinis.
tichotoma, Beer, l. c. = Sobralia dichotoma.
tigrina, *A. Rich. ex Beer*, *l. c.* 215.—Hab.?
Trianaei, Linden & Reichb. f. in Bot. Zeit. xviii. (1860) 74 = labiata.
Trianaei, *Linden & Reichb. f. in C. Koch*, *Wochenschr.* iii. (1860) 67.—N. Granat.
trichopiliochila, *Rodrig. Orch. Nov.* i. 70.—Bras.
trilabiata, *Rodrig. l. c.* 69.—Bras.
triophthalma ×, Reichb. f. in Gard. Chron. (1883) II. 526.
tuberculata, *Hort. ex Reichb. f. in Walp. Ann.* xi. 820. —Hab.?
undulata, Beer, Prakt. Orch. 215 = Schomburgkia undulata.
Veitchi, Hort. ex Pfitz, Verg. Morph. Orch. 124 = seq.?
Veitchiana ×. Cf. Gard. Chron. (1874) I. 385, 566 = Laelia Veitchiana.
velutina, *Reichb. f. in Gard. Chron.* (1870) 140, 1373. —Hab.?
vestalis, Hoffmgg. Verz.—Cf. Bot. Zeit. i. (1843) 831 = Forbesii.
violacea, Beer, Prakt. Orch. 209 = Sophronitis violacea.
violacea, Beer, l. c. 215 = Sobralia violacea.
virginalis, Linden & André, Illustr. Hortic. xxiii. (1876) t. 257 = labiata.

CATTLEYA:—
Wageneri, Reichb. f. in Bonplandia (1854) 21 = labiata.
Walkeriana, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 662.—Bras.?
Wallisii, Linden, in Gard. Chron. (1882) i. 557 = labiata.
Warneri, T. Moore, ex Warner, Orch. Sel. Ser. I. t. 8 = labiata.
Warscewiczii, Reichb. f. in Bonplandia (1854) 112 = labiata.
Whitei, *Hort. Low.—Cf. Gard. Chron.* ii. (1882) 151, 586.—Hab.?
Wilsoniana, *Reichb. f. in Gard. Chron.* (1877) ii. 72.—Am. austr.

CATTLEYOPSIS, Lem. Jard. Fleur. iv. (1854) Misc. 59. *ORCHIDEAE.*
delicatula, *Lem. l. c.*—Cuba.

CATU-ADAMBOE, Adans. Fam. ii. 85 (1763) = **Lagerstroemia**, Linn. (Lythrar.) (L. hirsuta).

CATU-NAREGAM, Adans. Fam. ii. 85 (1763) = **Randia**, Houst. (Rubiac.) (R. virosa).

CATURUS, Linn. Mant. i. 19 (1767) = **Acalypha**, Linn. (Euphorbiac.).
ramiflorus, Linn. l. c. 127 = Boehmeria ramiflora.
sessilis, Thou. ex Baill. Adansonia, i. (1860–61) 267 = A. integrifolia.
spiciflorus, Jacq. f. ex Wedd. in DC. Prod. xvi. i. 214 = Boehmeria spiciflora.
spiciflorus, Linn. Mant. i. 127 = A. hispida.
pedunculatus, Thou. ex Baill. in Adansonia, i. (1860–61) 267 = A. arborea.

CATURUS, Lour. Fl. Cochinch. 612 (1790) = **Malaisia**, Blanco (Urtic.).
scandens, Lour. l. c. = M. tortuosa.
acuminatus, *Cunninghamii*, *Deplanchei*, *fagifolius*, *oblongatus*, *pelagicus*, *torulosus*, *virescens*, Seem. Fl. Vit. 254 = M. tortuosa.

CATYONA, Lindl. Veg. Kingd. 715 (1847), sphalm. = GATYONA, Cass. = **Crepis**, Linn. (Compos.).

CAUCALIS, Linn. Syst. ed. I (1735); Gen. ed. I. 71 (1737). *UMBELLIFERAE*, Benth. & Hook. f. i. 928.
Agrocharis, Hochst. in Flora, xxvii. (1844) 19.
Daucalis, Pomel, Nouv. Mat. Fl. Atl. 148 (1874).
Glochidotheca, Fenzl, in Russegg. Reise, i. ii. 970, obs. (1843).
Lappularia, Pomel, Nouv. Mat. Fl. Atl. 149 (1874).
Lisaea, Boiss. in Ann. Sc. Nat. Sér. III. ii. 54 (1844).
Torilis, Adans. Fam. ii. 99 (1763).
Turgenia, Hoffm. Gen. Umb. 59 (1814).
Turgeniopsis, Boiss. in Ann. Sc. Nat. Sér. III. ii. 53 (1844).
abyssinica, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 331 = Daucus abyssinicus.
aequicolorum, All. Fl. Pedem. ii. 33 = Anthriscus vulgaris.
africana, Crantz, Cl. Umb. Emend. 109 = Capnophyllum africanum.
africana, Thunb. Prod. Pl. Cap. 49 = capensis.
angustifolia, *Forsk. Fl. Aegypt. Arab. Suppl.* 206.—Arabia.
Anthriscus, *Huds. Fl. Angl.* 99.—Europ.
arvensis, *Huds. l. c.* 98.—Europ.; As. temp.
aspera, Lam. Fl. Fr. iii. 424 = C. Anthriscus.
aspersa, Savi, Fl. Pis. i. 290 = arvensis.
bifrons, *Coss. & Dur. ex Ball, in Journ. Linn. Soc.* xvi. (1878) 478.—Algeria.
caerulescens, *Boiss. Elench.* 93. p. 53.—Hispan.; Marocco.
canadensis, Crantz, Cl. Umb. Emend. 109 = Sanicula canadensis.
capensis, *Lam. Encyc.* i. 651.—Afr. austr.
capitata, Stokes, Bot. Mat. Med. ii. 64 = Sanicula europaea.
carnosa, Roth, Tent. Fl. Germ. i. 119 = Daucus Carota.
Carota, Huds. Fl. Angl. ed. II. 114 = Daucus Carota.
coniifolia, Wall. ex DC. Prod. iv. 220 = C. Anthriscus.

CAUCALIS:—
cretica, Salzm. ex Boiss. Fl. Orient. ii. 1071 = Daucus maritimus.
daucoides, *Linn. Sp. Pl.* 241.—Europ.; As. temp.
Daucus, Crantz, Stirp. Austr. ed. I. fasc. iii. 125; ed. II. 227 = Daucus Carota.
divaricata, Heller, Fl. Wirceb. ed. I. 169 = arvensis.
elata, D. Don, Prod. Fl. Nep. 183 = C. Anthriscus.
elongata, Link, ex Steud. Nom. ed. I. 169 = leptophylla.
erythrotricha, *Boiss. & Haussk. ex Boiss. Fl. Orient.* ii. 1085.—Cilicia.
fallax, *Boiss. & Balansa, ex Boiss. l. c.* 1086.—Syria; Persia.
Gaillardoti, *Boiss. l. c.* 1085.—Syria.
Gerardi, *Schwaegr. ex Steud. Nom.* ed. I. 169.—Europ.
Gingidium, Crantz, Cl. Umb. Emend. 113 = Daucus Gingidium.
glabra, Forsk. Fl. Aegypt. Arab. 206 = Daucus littoralis.
glochidiata, Poir. Encyc. Suppl. ii. 137 = Daucus brachiatus.
grandiflora, Linn. Sp. Pl. 240.—Daucus grandiflorus.
helvetica, Jacq. Hort. Vindob. iii. 12. t. 16 = arvensis.
heterophylla, *Guss. Fl. Sic. Prod.* i. 326.—Reg. Mediterr.
hispanica, Crantz, Cl. Umb. Emend. 113 = Daucus mauritanicus.
hispanica, Lam. Encyc. i. 658 = Daucus Durieui.
hispida, Desf. Cat. Hort. Par. ed. III. 205 = Laserpitium hispidum.
humilis, Jacq. Hort. Vindob. ii. 92. t. 195 = leptophylla.
humilis, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 277 = nodosa.
infesta, Curt. Fl. Lond. ii. fasc. vi. t. 23 = arvensis.
infesta, *Vest, Ench.* 495; *ex DC. Prod.* iv. 219.—Europ.
japonica, *Houtt. Handleid.* viii. 42.—Japon.
lappulacea, Poepp. ex DC. Prod. iv. 219 = nodosa.
latifolia, Lam. Fl. Fr. iii. 426 = daucoides.
latifolia, *Linn. Syst. Nat.* ed. XII. ii. 205.—Europ.; As. temp.
leptophylla, Crantz, Stirp. Austr. ed. I. fasc. iii. 121; ed. II. 225 = daucoides.
leptophylla, *Linn. Sp. Pl.* 242.—Europ.; Reg. Mediterr.; Ind. or.
leptophylla, Viv. Fl. Lib. Spec. 16 = nodosa.
linearifolia, *Requien, ex DC. Prod.* iv. 219.—Europ. austr.
littoralis, Bieb. Fl. Taur. Cauc. i. 208 = Daucus bessarabicus.
marilandica, Crantz, Cl. Umb. Emend. 110 = Sanicula marilandica.
maritima, Gouan, Hort. Monsp. 135 = Daucus maritimus.
mauritanica, *Linn. Sp. Pl.* 241.—Afr. bor.
melanantha, *Benth. & Hook. f. Gen.* i. 929.—Afr. trop.
microcarpa, *Hook. & Arn. Bot. Beech. Voy.* 348.—Calif.
muricata, Crantz, Cl. Umb. Emend. 114 = daucoides.
nodiflora, Lam. Fl. Fr. iii. 424 = nodosa.
nodosa, *Scop. Fl. Carn.* ed. II. i. 192.—Europ.
orientalis, Linn. Sp. Pl. 241 = Daucus pulcherrimus.
orientalis, Lour. Fl. Cochinch. 177 = C. Anthriscus.
parviflora, Lam. Encyc. i. 657 = leptophylla.
platycarpos, Linn. Sp. Pl. 241 = daucoides.
pulcherrima, Willd. Enum. Horti Berol. 303 = Daucus pulcherrimus.
pumila, *Forsk. Fl. Aegypt. Arab.* 206.—Arab.
pumila, Gouan, Fl. Monsp. 285 = Daucus pumilus.
pumila, Lam. Fl. Fr. iii. 425 = leptophylla.
pumila, Linn. Syst. ed. X. 955 = arvensis.
pumila, Willd. Sp. Pl. i. 1385 = Daucus maritimus.
purpurea, Tenore, Fl. Nap. t. 131 = infesta.
Royeni, Crantz, Cl. Umb. Emend. 109 = daucoides.
Sanicula, Crantz, Stirp. Austr. ed. I. fasc. iii. 125; ed. II. 228 = Sanicula europaea.
scandicina, Roth, Tent. Fl. Germ. i. 121 = C. Anthriscus.
Scandix, Scop. Fl. Carn. ed. II. i. 191 = C. Anthriscus.
segetalis, Steud. Nom. ed. II. i. 312 = arvensis.
segetum, Thuill. Fl. Par. ed. II. 136 = arvensis.
silifolia, Roehl. ex Steud. Nom. ed. II. i. 312 = Heracleum austriacum.

CAUCALIS:—
Stocksiana, *Boiss. Diagn.* Ser. II. vi. 89.—Persia; Beluchist.
strigosa, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 248.—Syria; Mesopotam.
tenella, *Delile, Fl. Egypt*, 202. t. 21. f. 3.—Syria; Persia; Aegypt.
tenerrima, Scheele, in Linnaea, xxv. (1852) 256 = nodosa.
tenuifolia, Salisb. Prod. 161 = daucoides.
trichosperma, *Delile, in Steud. Nom.* ed. II. i. 312.—Syria.
tuberculata, Poir. Encyc. Suppl. ii. 137 = Pimpinella peregrina.
virgata, Poir. Voy. Barb. ii. 133 = Daucus laserpitioides.
xanthotricha, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 353 = leptophylla.

CAUCANTHUS, Forsk. Fl. Aegypt. Arab. 91 (1775). *MALPIGHIACEAE*?
arabicus, *Lam. Encyc.* i. 658.—Arabia.
Forskahlei, Raeusch. Nom. ed. III. 132 = arabicus.
platanifolia, Rafin. Sylva Tellur. 72 = Sterculia platanifolia.

CAULANTHUS, S. Wats. Bot. King, Exped. 19, 27 (1871). *CRUCIFERAE.*
amplexicaulis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 364.—Calif.
Coulteri, *S. Wats. in Bot. King, Exped.* 27.—Calif.
crassicaulis, *S. Wats. l. c.*—Calif.
glaucus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 364.—Calif.
hastatus, *S. Wats. in Bot. King, Exped.* 28. t. 3.—Am. bor. occ.
inflatus, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 364.—Calif.
pilosus, *S. Wats. in Bot. King, Exped.* 27.—Calif.
procerus, *S. Wats. l. c.*—Calif.

CAULARTHRON, Rafin. Fl. Tellur. ii. 40 (1836) = **Epidendrum**, Linn. (Orchid.).
bicornutum, Rafin. l. c. 41 = Diacrium bicornutum.
umbellatum, Rafin. l. c. = E. umbellatum.

CAULINIA, DC. Fl. Fr. iii. 156 (1805) = **Posidonia** Koen. (Naiad.).
antarctica, R. Br. Prod. 339 = Cymodocea antarctica.
australiana, F. Muell. Fragm. vi. 198.—Austral.
oceanica, R. Br. Prod. 339 = Posidonia australis.
oceanica, DC. Bot. Gall. iii. 156 = Posidonia Caulini.
ovalis, R. Br. l. c. 339 = Halophila ovata.
prorepens, F. Muell. Fragm. vii. 225.—Austral.
serrulata, R. Br. Prod. 339 = Cymodocea serrulata.
spinulosa, R. Br. l. c. = Halophila spinulosa.

CAULINIA, Moench, Meth. Suppl. 47 (1802) = **Kennedya**, Vent. (Legumin.).
Beckxiana, F. Muell. Fragm. xi. 98 = K. Beckxiana.
rubicunda, Moench, Meth. Suppl. 47 = K. rubicunda.

CAULINIA, Willd. in Mém. Acad. Berl. (1798) 87 = **Najas**, Linn.
alagnensis, Pollini, Fl. Veron. iii. 49 = N. alagnensis.
alternifolia, Willd. ex Cham. in Linnaea, iv. (1829) 501 = N. indica.
composita, Buch.-Ham. in Wall. Cat. n. 5184.—Ind. or.
flexilis, Willd. in Mém. Acad. Berl. (1798) 89. t. i. f. 1 = N. flexilis.
fragilis, Willd. l. c. 87 = N. fragilis.
guadalupensis, Spreng. Syst. i. 20 = N. tenuifolia.
indica, Willd. in Mém. Acad. Berl. (1798) 89 = N. indica.
intermedia, Balb. & Nocca, Fl. Tic. 11. 163 = N. alagnensis.
microphylla, Balb. & Nocca, l. c. = N. graminea.
minor, Coss. & Germ. Fl. Paris, 575 = N. muricata.
pectinata, Parl. Fl. Ital. iii. 665.—Italia.
tenella, Nees, in Neuwied, Reise Bras. ii. 345 = N. tenera.

CAULLINIA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Hippuris**, Linn. (Halorag.).
hippuroides, Rafin. l. c. = H. vulgaris.

CAULOBRYON, Klotzsch, ex C. DC. in DC. Prod. xvi. i. 240 (1869) = **Piper**, Linn.

CAULOPHYLLUM, Michx. Fl. Bor. Am. i. 204. t. 21 (1803). *BERBERIDEAE*, Benth. & Hook. f. i. 43.
PHTHEIROTHECA, Maxim. ex Regel, in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 223.
gracile, Dougl. ex Hook. Fl. Bor. Am. i. 30 = Vancouveria hexandra.
robustum, *Maxim. Prim. Fl. Amur.* 33.—Manchuria.
thalictroides, *Michx. Fl. Bor. Am.* i. 205.—Am. bor.
thalictroides, Regel, in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 223 = robustum.

CAULOPSIS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 332 = **Arabis**, L nn. (Crucif.).
alpina, Fourr. l. c. = A. brassicaeformis.

CAULOTRETUS, Rich. ex Spreng. Syst. iv. Cur. Post. 406 (1827) = **Bauhinia**, Linn. (Legumin.).
cordifolius, Rich. ex DC. Prod. ii. 517 = B. Richardiana.
eriophorus, Tenore, in Rendic. Soc. Borb. Ncap. i. (1842) 410 = B. eriophora.
smilacinus, Schott, ex Spreng. Syst. iv. Cur. Post. 406 = B. smilacina.

CAULOTULIS, Rafin. Fl. Tellur. iv. 76 (1836) = **Ipomoea**, Linn. (Convolv.).

CAUSONIA, Rafin. Sylva Tellur. 87 (1838) = seq.

CAUSONIS, Rafin. Med. Fl. ii. 122 (1830) = **Vitis**, Linn. (Ampelid.).
japonica, Rafin. l. c. = V. japonica.
trifolia, Rafin. l. c. = V. trifolia.

CAUSTIS, R. Br. Prod. 229 (1810). *CYPERACEAE* Benth. & Hook. f. iii. 1067.
EUROSTORHIZA, Steud. Syn. Pl. Cyp. 265 (1855).
dioica, *R. Br. l. c.*—Austral.
flexuosa, *R. Br. l. c.*—Austral.
hexandra, Nees, in Lehm. Pl. Preiss. ii. 88 = dioica.
pentandra, *R. Br. Prod.* 239.—Austral.
recurvata, *Spreng. Syst.* iv. *Cur. Post.* 26.—Austral.
restiacea, *F. Muell. ex Benth. Fl. Austral.* vii. 421.—Austral.
Sieberi, Kunth, Enum. Pl. ii. 307 = Gahnia Sieberi.

CAUTLEA, Royle, Illustr. Bot. Himal. 361 (1839) = **Roscoea**, Sm. (Scitam.).
lutea, Royle, l. c. = R. lutea.

CAVALAM, Adans. Fam. ii. 357 (1763) = **Sterculia**, Linn.

CAVALLIUM, Schott, Melet. 33 (1832) = **Sterculia**, Linn.
comosum, Schott & Endl. l. c. = S. comosa.
urens, Schott & Endl. l. c. = S. urens.

CAVANALIA, Griseb. Cat. Pl. Cub. 76 (1864), sphalm. = **Canavalia**.

CAVANILLA, J. F. Gmel. Syst. 1037 (1791) = **Dombeya**, Cav. (Sterculiac.).
acerifolia, J. F. Gmel. l. c. = Pterospermum acerifolium.
acutangula, J. F. Gmel. l. c. = D. acutangula.
angulata, J. F. Gmel. l. c. = D. acutangula.
decanthera, J. F. Gmel. l. c. = Trochetia decanthera.
ferruginea, J. F. Gmel. l. c. 1038 = D. ferruginea.
ovata, J. F. Gmel. l. c. = D. ovata.
palmata, J. F. Gmel. l. c. 1037 = D. acutangula.
phoenicea, J. F. Gmel. l. c. 1038 = Pentapetes phoenicea.
punctata, J. F. Gmel. l. c. 1037 = D. punctata.
tiliaefolia, J. F. Gmel. l. c. = D. acutangula.
tomentosa, J. F. Gmel. l. c. = D. tomentosa.
umbellata, J. F. Gmel. l. c. = D. umbellata.

CAVANILLA, Salisb. Prod. 385 (1796) = **Stuartia**, Linn. (Ternstroem.).
florida, Salisb. l. c. = S. Malachodendron.

CAVANILLA, Thunb. Nov. Gen. 105 (1795) = **Pyrenacantha**, Hook. (Olacin.).
scandens, Thunb. l. c. = P. scandens.

CAVANILLA, Vell. Fl. Flum. 226 (1825); v. t. 102 (1827) = **Caperonia**, St. Hil. (Euphorbiac.).
spinosa, Vell. il. cc. = Caper. Velloziana.

CAVANILLEA, Desr. in Lam. Encyc. iii. 663 (1789) = **Diospyros**, Linn. (Ebenac.).
philippinensis, Desr. l. c. = D. Embryopteris.
Mabolo, Hiern, Ebenac. 260 = D. discolor.

CAVANILLEA, Medic. Malv. 19 (1787) = **Anoda**, Cav. (Malvac.).
hastata, Medic. l. c. = A. hastata.

CAVANILLESIA, Ruiz & Pav. Prod. 97. t. 20 (1794). *MALVACEAE*, Benth. & Hook. f. i. 211.
POURRETIA, Willd. Sp. Pl. iii. 844 (1801).
cordata, *Spreng. Syst.* iii. 125.—Peruv.
platanifolia, *H. B. & K. Nov. Gen. et Sp.* v. 306.—Panama.
tuberculata, Kostel. Allg. Med. Pharm. Fl. v. 1872 = platanifolia?
umbellata, *Ruiz & Pav. Syst. Veg.* 166.—Peruv.

CAVARIA, Steud. Nom. ed. I. 169 (1821), sphalm.? = **Tovaria**, Ruiz & Pav. (Capparid.).

CAVENDISHIA, Lindl. Bot. Reg. sub t. 1791 (1836). *VACCINIACEAE*, Benth. & Hook. f. ii. 570.
POLYBAEA, Klotzsch, in Linnaea, xxiv. (1851) 30.
PROELESIA, Klotzsch, l. c. 32.
SOCRATESIA, Klotzsch, l. c. 22.
acuminata, *Benth. ex Hemsl. Biol. Centr. Am. Bot.* ii. 272.—Mexic.
complectens, *Hemsl. l. c.*—Costa Rica.
crassifolia, *Hemsl. l. c.* 273.—Mexic.
Endresii, *Hemsl. l. c.*—Costa Rica.
latifolia, *Hemsl. l. c.*—Mexic.
laurifolia, *Benth. & Hook. f. Gen.* ii. 570.—Guatemala.
melastomoides, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 273.—Costa Rica.
nobilis, *Lindl. Bot. Reg.* sub t. 1791.—Am. austr.
pubescens, *Hemsl. Biol. Centr. Am. Bot.* ii. 273.—Panama.
Quercma, *Benth. & Hook. f. Gen.* ii. 570.—N. Granat.
tarapotana, *Benth. & Hook. f. l. c.*—Bras.
veraguensis, *Hemsl. Biol. Centr. Am. Bot.* ii. 273.—Panama.
Warscewiczii, *Hemsl. l. c.* 274.—Guatemala.

CAVINIUM, Thou. Gen. Nov. Madag. 11 (1806) = **Vaccinium**, Linn.
madagascariense, Thou. ex Poir. in Dict. Sc. Nat. vii. 310 = *Agapetes madagascariensis*.

CAVOLINIA, Rafin. in Am. Monthly Mag. (1818) 175 = CAULINIA, Willd. = **Najas**, Linn.

CAYAPONIA, Silva Manso, Enum. Subst. Brazil. 31 (1836). *CUCURBITACEAE*.
ALLAGOSPERMA, M. Roem. Syn. Pepon. 15 (1846).
ANTAGONIA, Griseb. Pl. Lorentz. 144 (1874).
ALTERNASEMINA, Silva Manso, Cathart. Bras. (1836).
CIONANDRA, Griseb. Fl. Brit. W. Ind. 286 (1860).
DERMOPHYLLA, Silva Manso, Enum. Subst. Brazil. 30 (1836).
DRUPARIA, Silva Manso, l. c. 35 (1836).
PERIANTHOPODUS, Silva Manso, l. c. 28 (1836)
TRIANOSPERMA, Mart. Syst. Mat. Med. Brazil. 79 (1843).
alata, *Cogn. in DC. Monog. Phan.* iii. 746.—Yucatan.
amazonica, *Cogn. l. c.* 797.—Bras.
americana, *Cogn. l. c.* 785.—Am. cent.
Andreana, *Cogn. in Bull. Acad. Belg.* Sér. II. xlix. (1880) 195.—Ecuador.
angustiloba, *Cogn. in DC. Monog. Phan.* iii. 771.—Bras.
attenuata, *Cogn. l. c.* 769.—Mexic.
Bonplandii, *Cogn. l. c.* 796.—Bras.
Boykinii, *Cogn. l. c.* 766.—Am. bor.
Buraeavi, *Cogn. l. c.* 790.—N. Granat.
Cabocla, Mart. Syst. Mat. Med. Veg. Bras. 81 = globosa.
calycina, *Cogn. in Mart. Fl. Bras.* vi. iv. 78.—Bras.
citrullifolia, *Cogn. ex Griseb. in Goett. Abh.* xxiv. (1879) 135.—Reg. Argent.
coccinea, André, ex Cogn. in Bull. Acad. Belg. Sér. II. xlix. (1880) 196 = micrantha.
cordifolia, *Cogn. in Mart. Fl. Bras.* vi. iv. 77. t. 21.—Bras.
coriacea, *Cogn. l. c.* 79. t. 22.—Bras.

CAYAPONIA:—
Crugeri, *Cogn. in DC. Monog. Phan.* iii. 778.—Ins. Trinit.
diffusa, *Silva Manso, Enum. Subst. Brazil.* 32.—Bras.
diversifolia, *Cogn. in DC. Monog. Phan.* iii. 787.—Bras.
elliptica, *Silva Manso, Enum. Subst. Brazil.* 32.—Bras.
Espelina, *Cogn. in DC. Monog. Phan.* iii. 792.—Bras.
excisa, *Cogn. l. c.* 747.—Cuba.
ficifolia, *Cogn. l. c.* 782.—Bras.
floribunda, *Cogn. l. c.* 779.—Bras.
fluminensis, *Cogn. in Mart. Fl. Bras.* vi. iv. 76.—Bras.
glandulosa, *Cogn. in DC. Monog. Phan.* iii. 755.—Peruv.
Glaziovii, *Cogn. in Mart. Fl. Bras.* vi. iv. 74.—Bras.
globosa, *Silva Manso, Enum. Subst. Brazil.* 32.—Bras.
gracillima, *Cogn. in DC. Monog. Phan.* iii. 784.—Bras.
granatensis, *Cogn. l. c.* 794.—N. Granat.
grandiflora, *Cogn. l. c.* 779.—Yucatan.
heterophylla, *Cogn. l. c.* 758.—N. Granat.
hirsuta, *Cogn. in Mart. Fl. Bras.* vi. iv. 76.—Bras.
latebrosa, *Cogn. in DC. Monog. Phan.* iii. 776.—Ins. Canar.
latifolia, *Cogn. l. c.* 752.—Paraguay.
Lhotzkyana, *Cogn. l. c.* 765.—Bras.
longifolia, *Cogn. l. c.* 796.—Bras.
Martiana, *Cogn. l. c.* 777.—Bras.
Maximowiczii, *Cogn. l. c.* 745.—Mexic.
micrantha, *Cogn. in Bull. Acad. Belg.* Sér. II. xlix. (1880) 196.—N. Granat.
Ottoniana, *Cogn. in DC. Monog. Phan.* iii. 755.—Venezuela.
ovata, *Cogn. l. c.* 754.—N. Granat.
palmata, *Cogn. in Bull. Soc. Bot. Belg.* xvii. (1878) 295.—Bras.
pedata, *Cogn. in Mart. Fl. Bras.* vi. iv. 80. t. 23.—Bras.
pentaphylla, *Cogn. in DC. Monog. Phan.* iii. 760.—Bras.
peruviana, *Cogn. l. c.* 795.—Peruv.
petiolulata, *Cogn. in Bull. Soc. Bot. Belg.* xvii. (1878) 295.—Bras.
piauhiensis, *Cogn. in DC. Monog. Phan.* iii. 770.—Bras.
pilosa, Cogn. in Mart. Fl. Bras. vi. iv. 75 = diffusa.
podantha, *Cogn. in DC. Monog. Phan.* iii. 753.—Paraguay.
Poeppigii, *Cogn. l. c.* 756.—Peruv.
racemosa, *Cogn. l. c.* 768.—Porto Rico.
rigida, *Cogn. l. c.* 790.—Guiana.
Ruizii, *Cogn. l. c.* 794.—Peruv.
Sandia, *Cogn. ex Griseb. in Goett. Abh.* xxiv. (1879) 135.—Reg. Argent.
setulosa, *Cogn. in DC. Monog. Phan.* iii. 767.—Bras.
simplicifolia, *Cogn. l. c.* 774.—N. Granat.
Tayuya, *Cogn. l. c.* 772.—Bras.
ternata, *Cogn. in Mart. Fl. Bras.* vi. iv. 79.—Bras.
Tibiricae, *Cogn. in DC. Monog. Phan.* iii. 784.—Bras.
tomentosa, *Cogn. l. c.* 747.—Peruv.
triangularis, *Cogn. l. c.* 774.—Guiana.
trifoliolata, *Cogn. l. c.* 789.—Bras.
trilobata, *Cogn. l. c.* 780.—Bras.
tubulosa, *Cogn. in Mart. Fl. Bras.* vi. iv. 78.—Bras.
villosissima, *Cogn. l. c.* 81. t. 24.—Bras.
Weddellii, *Cogn. in DC. Monog. Phan.* iii. 793.—Bras.

CAYLUSEA, A. St. Hil. 2me Mém. Resed. 29 (1837). *RESEDACEAE*, Benth. & Hook. f. i. 111.
STYLEXIA, Rafin. Fl. Tellur. iv. 121 (1836).
SYNTROPHE, G. Ehrenb. ex Muell. Arg. in Neue Denks. Schw. Ges. Zuerich, xvi. (1858) 227.
abyssinica, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* vii. (1840) 43.—Abyss.
canescens, *A. St. Hil. 2me Mém. Resed.* 29.—Afr. bor.; Arabia.
Moquiniana, *Webb, Fragm. Fl. Aethiop.* 28.—Afr. bor.

CAYRATIA, Juss. ex Guill. in Dict. Class. Hist. Nat. iv. 346 (1823) = **Vitis**, Linn. (Ampelid.) (V. pedata).

CEANOTHUS, Linn. in Act. Soc. Sc. Upsal. 77 (1741). *RHAMNEAE*, Benth. & Hook. f. i. 378.
FORRESTIA, Rafin. in Med. Repos. N. York, II. iii. (1806) 422.
africanus, Linn. Sp. Pl. 196 = Noltea africana.
Alamani, DC. Prod. ii. 31 = Colubrina Alamani.
americanus, *Linn. Sp. Pl.* 195.—Am. bor.
americanus, Mill. Ic. t. 86 = Milleri.
arborescens, Mill. Gard. Dict. ed. VIII. n. 3 = Colubrina ferruginosa?
asiaticus, Linn. Sp. Pl. 196 = Colubrina asiatica.
atropurpureus, Rafin. New Fl. Am. iii. 56 = Nolteae sp.?
axillaris, Carr. in Rev. Hortic. (1876) 87 = azureus?
azureus, *Desf. Tabl.* ed. II. 232.—Mexic.
azureus, Kellogg, in Proc. Calif. Acad. i. (1855) 55 = hirsutus.
Baumannianus, Spach, Hist. Vég. Phan. ii. 460 = ovatus.
Bertini, Carr. in Rev. Hortic. (1872) 440 = azureus?
bicolor, Willd. in Schult. f. Syst. vii. 65 = hirsutus.
Burmannianus, Steud. Nom. ed. II. i. 313, sphalm. = Baumannianus.
buxifolius, *Willd. ex Schult. f. Syst.* vii. 62.—Mexic.
caeruleus, *Lag. Gen. et Sp. Nov.* 11.—Mexic.
californicus, Kellogg, in Proc. Calif. Acad. i. (1855) 55 = integerrimus.
capensis, DC. Prod. ii. 30 = Scutia Commersonii.
capsularis, Forst. f. Prod. 18 = Colubrina asiatica.
celtidifolius, Cham. & Schlecht. in Linnaea, v. (1830) 602 = Colubrina celtidifolia.
Chloroxylon, *Nees, Syst. Laurin.* 660.—Jamaica.
circumscissus, Gaertn. Fruct. ii. 110. t. 106 = Scutia Commersonii.
collinus, Dougl. ex Knight & West. Fl. Cab. i. t. 13 = sorediatus?
colubrinus, Lam. Tabl. Encyc. ii. 90 = Colubrina ferruginosa.
cordulatus, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 124. f. 39.—Calif.
crassifolius, *Torr. in Pacif. Rail. Rep.* iv. 75.—Calif.
cubensis, *Lam. Tabl. Encyc.* ii. 90.—Ind. occ.
cuneatus, A. Gray, in Ives, Rep. Color. Riv. 9 = Greggii.
cuneatus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 267.—Calif.
decumbens, Hort. ex Steud. Nom. ed. II. i. 313 = americanus.
decumbens, *S. Wats. in Proc. Am. Acad.* x. (1875) 335.—Calif.
Delilianus, *Spach, Hist. Vég. Phan.* ii. 459.—Hab.?
dentatus, *Torr. & Gray, Fl. N. Am.* i. 268.—Calif.
depressus, *Benth. Pl. Hartw.* 8.—Mexic.
Dillenianus, Hort. ex C. Koch, Dendrol. i. 620 = americanus.
discolor, Vent. Jard. Malm. 58 = Pomaderris elliptica.
divaricatus, Boland. Cat. Pl. Calif. 8 = cordulatus.
divaricatus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 266.—Calif.
diversifolius, Kellogg, in Proc. Calif. Acad. i. (1855) 57 = decumbens.
elegans, Lem. Illustr. Hortic. vii. (1860) t. 268 = thyrsiflorus.
ellipticus, *Rafin. New Fl. Am.* iii. 56.—Am. bor.
elongatus, Salisb. Prod. 140 = Noltea africana.
Fendleri, *A. Gray, Pl. Fendl.* 29.—Am. bor. occ.
ferreus, DC. Prod. ii. 30 = Scutia ferrea.
ferrugineus, Wendl. ex Steud. Nom. ed. II. i. 313 = Pomaderris lanigera.
floribundus, *Hook. Bot. Mag.* t. 4806.—Calif.
Fontanesianus, Spach, Hist. Vég. Phan. ii. 460 = ovatus.
glaber, *Spach, l. c.* 459.—Am. bor.
glandulosus, Rafin. New Fl. Am. iii. 57 = ovatus.
glandulosus, *Schlecht. in Linnaea*, xv. (1841) 474.—Mexic.
globulosus, Labill. Nov. Holl. Pl. i. 61. t. 85 = Spyridium globulosum.
glomeratus, Rafin. New Fl. Am. iii. 55 = americanus.
grandis, Dougl. ex Hook. Fl. Bor. Am. i. 125 = velutinus.
granulosus, *Ruiz & Pav. Fl. Per.* iii. 5. t. 228.—Peruv.
Greggii, *A. Gray, Pl. Wright.* ii. 28.—Arizona.
Guadalupae, Steud. Nom. ed. II. i. 313 = laevigatus.
guineensis, DC. Prod. ii. 30 = Chailletia toxicaria?
Hartwegii, *Hook. ex Heynh. Nom.* ii. 128, nomen.—Mexic.
herbaceus, Rafin. in Med. Repos. N. York, v. (1808) 360 = americanus.

CEANOTHUS:—
hirsutus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 266.—Calif.
hybridus, Hort. ex C. Koch, Dendrol. i. 620 = americanus.
hypericoides, L'Hérit. ex DC. Prod. ii. 32 = microphyllus.
incanus, Torr. & Gray, Fl. N. Am. i. 265.—Calif.
infestus, H. B. & K. Nov. Gen. et Sp. vii. 61 = Adolphia infesta.
integerrimus, *Hook. & Arn. Bot. Beech. Voy.* 329.—Am. bor. occ.
intermedius, Hook. Fl. Bor. Am. i. 124 = ovatus.
intermedius, Pursh, Fl. Am. Sept. i. 167 = americanus.
laevigatus, *DC. Prod.* ii. 30.—Ind. occ.
laevigatus, Hook. Fl. Bor. Am. i. 125 = velutinus.
lancifolius, Moench, Meth. 651 = Noltea africana.
laniger, Andr. Bot. Rep. t. 569 = Pomaderris lanigera.
latifolius, *Rafin. New Fl. Am.* iii. 55.—Am. bor.
Leschenaultii, *DC. Prod.* ii. 31.—Zeylan.
levigatus, Rafin. New Fl. Am. iii. 55 = americanus.
Lobbianus, Hook. Bot. Mag. t. 4810 = dentatus.
macrocarpus, *Cav. Ic.* iii. 38. t. 270.—Mexic.
macrocarpus, Hort. ex Steud. Nom. ed. II. i. 313 = americanus.
macrocarpus, Moç. & Sesse, ex DC. Prod. ii. 32, 33 = Mocinianus.
macrocarpus, Nutt. in Torr. & Gray, Fl. N. Am. i. 267 = cuneatus.
macrophyllus, Desf. Tabl. ed. II. 232 = americanus.
megacarpus, Nutt. Sylv. Am. ii. 46 = cuneatus.
microphyllus, *Michx. Fl. Bor. Am.* i. 154.—Am. bor.
Milleri, *Tausch, in Flora*, xxi. (1838) 1. *Beibl.* 79.—Am. bor.
Mocinianus, *DC. Prod.* ii. 32.—Mexic.
mystacinus, DC. l. c. 31 = Helinus scandens.
napalensis, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 375 = Rhamnus nipalensis.
Neumanni, *Tausch, in Flora*, xxi. (1838) 738.—Am. bor.
nevadensis, Kellogg, in Proc. Calif. Acad. ii. (1863) 152 = integerrimus.
nitidus, Torr. in Pacif. Rail. Rep. iv. 75 = sorediatus.
officinalis, Rafin. Med. Fl. ii. 205 = americanus.
oliganthus, Nutt. in Torr. & Gray, Fl. N. Am. i. 266 = hirsutus.
oreganus, Nutt. l. c. 265 = sanguineus.
ovalifolius, *Wender. in Schr. Naturf. Ges. Marb.* ii. (1830) 247.—Am. bor.
ovalis, Bigel. Fl. Bost. ed. II. 92 = ovatus.
ovatus, *Desf. Arb.* ii. 381.—Am. bor.
ovatus, Hort. ex Lindl. Bot. Reg. (1840) t. 20 = seq.
pallidus, *Lindl. l. c.*—Hab.?
paniculatus, Roth, Nov. Pl. Sp. 154 = Celastrus paniculata.
papillosus, *Torr. & Gray, Fl. N. Am.* i. 268.—Calif.
pauciflorus, *Moç. & Sesse, ex DC. Prod.* ii. 33.—Mexic.
perennis, Pursh, Fl. Am. Sept. i. 167 = americanus.
Pitcheri, Pickering, ex Torr. & Gray, Fl. N. Am. i. 264 = americanus.
procumbens, Hort. ex C. Koch, Dendrol. i. 619 = americanus.
prostratus, *Benth. Pl. Hartw.* 302.—Calif.
pubescens, *Ruiz & Pav. Fl. Per.* iii. 6. t. 228.—Peruv.
pubiflorus, DC. Prod. ii. 30 = Zizyphus pubiflorus.
pulchellus, Delile, ex Spach, Hist. Vég. Phan. ii. 459 = Delilianus.
reclinatus, Bosc, ex Steud. Nom. ed. II. i. 313 = americanus.
reclinatus, *L'Hérit. Sert. Angl.* 6.—Ind. occ.
rigidus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 268.—Calif.
sanguineus, *Pursh, Fl. Am. Sept.* i. 167.—Am. bor. occ.
sanguineus, Nutt. Gen. Am. i. 153 = americanus.
Sarcomphalus, DC. Prod. ii. 30 = Sarcomphalus laurinus.
scandens, D. Dietr. Syn. Pl. i. 812 = Noltea africana.
serpyllifolius, *Nutt. Gen. Am.* i. 154.—Florida.
sorediatus, *Hook. & Arn. Bot. Beech. Voy.* 328.—Calif.
spathulatus, Labill. Nov. Holl. Pl. i. 60. t. 84 = Trymalium Billardieri.
sphaerocarpus, DC. Prod. ii. 30 = Rhamnus sphaerospermus.
spinosus, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 267.—Calif.

CEANOTHUS:—
tardiflorus, Hornem. Hort. Hafn. i. 230 = americanus.
thyrsiflorus, *Eschw. in Mém. Acad. Pétersb.* Sér. VI. x. (1826) 285.—Calif.
triflorus, Steud. Nom. ed. II. i. 313 = Colubrina triflora.
trinervus, Moench, Meth. 651 = americanus.
triqueter, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 376 = Rhamnus triqueter.
Veitchianus, *Hook. Bot. Mag.* t. 5127.—Calif.
velutinus, *Dougl. in Hook. Fl. Bor. Am.* i. 125. t. 45.—Am. bor. occ.
verrucosus, Nutt. in Torr. & Gray, Fl. N. Am. i. 267 = cuneatus.
virgatus, Rafin. New Fl. Am. iii. 56 = americanus.
Wendlandianus, Roem. & Schult. Syst. v. 299 = Pomaderris ferruginea.
Wightianus, Wall. Cat. n. 4264 = Rhamnus Wightii.
zeylanicus, Heyne, in Roth, Nov. Pl. Sp. 153 = Scutia Commersonii.

CEARIA, Dum. Comm. Bot. 65 in adnot. (1822) = **Eurycles**, Salisb. (Amaryllid.).
amboinensis, Dum. l. c. = E. sylvestris.

CEBATHA, Forsk. Fl. Aegypt. Arab. 171 (1775) = **Cocculus**, DC. (Menisperm.) (C. Cebatha).

CECICODAPHNE, Nees, in Wall. Pl. As. Rar. ii. 70 (1831) = **Cinnamomum**, Blume (Laurin.).
glaucescens, Nees, l. c. = Cinnam. Cecicodaphne.

CECROPIA, Linn. in Loefl. It. 272 (1758). *URTICACEAE*, Benth. & Hook. f. iii. 378.
AMBAIBA, Adans. Fam. ii. 377 (1763).
acutifolia, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 81.—Peruv.
adenopus, *Mart. ex Miq. in Mart. Fl. Bras.* iv. 1. 147.—Am. trop.
albicans, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 82.—Peruv.
angustifolia, *Tréc. l. c.* 83.—Peruv.
argentea, *Visiani, Ort. Bot. Padov.* 135.—Hab.?
bicolor, *Klotzsch, in Linnaea*, xx. (1847) 531.—Peruv.
carbonaria, *Mart. ex Miq. in Mart. Fl. Bras.* iv. 1. 144.—Am. trop.
cinerea, *Miq. l. c.* 142.—Am. trop.
commutata, *Schott, ex Miq. l. c.* 148.—Am. trop.
concolor, Hort. Schoenbr. ex Miq. l. c. = Schiedeana.
concolor, *Willd. Sp. Pl.* iv. 652.—Bras.
cyrtostachya, *Miq. in Mart. Fl. Bras.* iv. 1. 145.—Am. trop.
dentata, *Klotzsch, in Linnaea*, xx. (1847) 533.—Peruv.
digitata, *Tenore, ex Miq. in Mart. Fl. Bras.* iv. 1. 149.—Bras.
digitata, Klotzsch, in Linnaea, xx. (1847) 534 = angustifolia.
flagellifera, Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 81 = bicolor.
frigida, *Dufresne, in Illustr. Hortic.* xx. (1873) 70.—Hab.?
Goudotiana, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 83.—N. Granat.
hololeuca, *Miq. in Mart. Fl. Bras.* iv. 1. 148.—Am. trop.
Humboldtiana, *Klotzsch, in Linnaea*, xx. (1847) 530.—Am. trop.
insignis, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 318.—Nicaragua.
Klotzschiana, *Miq. in Mart. Fl. Bras.* iv. 1. 151.—Am. trop.
latiloba, *Miq. l. c.* 147.—Am. trop.
leucocoma, *Miq. l. c.* 142.—Am. trop.
leucophaea, *Poepp. ex Miq. l. c.* 151.—Am. trop.
lyratiloba, *Miq. l. c.* 144.—Am. trop.
membranacea, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 83.—Ind. occ.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* iii. 151.—Mexic.
nivea, *Poepp. ex Klotzsch, in Linnaea*, xx. (1847) 532.—Peruv.
obtusa, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 79.—Bras.
obtusifolia, *Bertol. Fl. Guatimal.* 39.—Guatemala.
pachystachya, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 80.—Bras.
palmata, *Willd. Sp. Pl.* iv. 652.—Bras.

CECROPIA :—
panamensis, *Hemsl. Biol. Centr. Am. Bot.* iii. 151.—Panama.
peltata, *Linn. Pugill. Pl. Jamaic.* 28 (*Amoen. Acad.* v. 410).—Am. trop.
peltata, G. F. W. Mey. Prim. Fl. Esseq. 280 = surinamensis.
peltata, Ruiz, ex Klotzsch, in Linnaea, xx. (1847) 532 = Ruiziana.
peltata, Schreb. ex Miq. in Mart. Fl. Bras. iv. 1. 150 = Schreberiana.
peltata, Vell. Fl. Flum. x. t. 101 = adenopus.
peltata, Willd. ex Klotzsch, in Linnaea, xx. (1847) 530 = Humboldtiana.
pinnatiloba, *Klotzsch, in Linnaea*, xx. (1847) 533.—Peruv.
polystachya, Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 80 = pinnatiloba.
propinqua, *Miq. in Mart. Fl. Bras.* iv. 1. 149.—Am. trop.
Ruiziana, *Klotzsch, in Linnaea*, xx. (1847) 532.—Peruv.
scabra, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 95.—Bras.
scabra, Ruiz, ex Klotzsch, in Linnaea, xx. (1847) 531 = Klotzschiana.
Schiedeana, *Klotzsch, l. c.*—Mexic.
Schreberiana, *Miq. in Mart. Fl. Bras.* iv. 1. 150.—Am. trop.
sciadophylla, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 93.—Bras.
strigosa, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 82.—Peruv.
surinamensis, *Miq. in Mart. Fl. Bras.* iv. 1. 143.—Am. trop.
tubulosa, *Ruiz, ex Klotzsch, in Linnaea*, xx. (1847) 534.—Peruv.

CEDRELA, P. Br. Hist. Jamaic. 158 (1756). *MELIACEAE*, Benth. & Hook. f. i. 339.
CEDRUS, Mill. Gard. Dict. ed. VII (1759).
CUVERACA, Jones, in As. Res. iv. (1795) 281.
JOHNSONIA, Adans. Fam. ii. 343 (1763).
PTEROSIPHON, Turcz. in Bull. Soc. Nat. Mosc. (1863) i. 589.
TOONA, M. Roem. Syn. Hesper. 131 (1846).
adenophylla, Mart. ex C. DC. in DC. Monog. Phan. i. 738 = paraguaiensis.
alternifolia, *Steud. Nom.* ed. I. 170.—Honduras.
angustifolia, *Moç. & Sesse, ex DC. Prod.* i. 624.—Mexic.
australis, F. Muell. Fragm. i. 4 = C. Toona.
bogotensis, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 377.—N. Granat.
brasiliensis, Juss. in St. Hil. Fl. Bras. Mer. ii. 86 = fissilis.
Cedro, Loefl. It. 183 = odorata.
Dugesii, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 190.—Mexic.
febrifuga, Forsten, Diss. Cedrel. 16 = C. Toona.
fissilis, *Vell. Fl. Flum.* 75; ii. t. 68.—Bras.
glabra, C. DC. in DC. Monog. Phan. i. 742 = C. Toona.
Glaziovii, *C. DC. in Mart. Fl. Bras.* xi. 1. 224. t. 65. f. 2.—Am. trop.
guianensis, *A. Juss. in Mém. Mus. Par.* xix. (1830) 255.—Guiana.
hexandra, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 425 = C. Toona.
inodora, *Hassk. Hort. Bogor.* i. 131.—Java.
Kotschyi, *Schweinf. Reliq. Kotsch.* 36. t. 35.—Afr. trop.
longifolia, Wall. Cat. n. 1273 = C. Toona.
Mahagoni, Linn. Syst. ed. X. 940 = Swietenia Mahagoni.
mexicana, M. Roem. Syn. Hesper. 137 = Swietenia humilis.
microcarpa, C. DC. in DC. Monog. Phan. i. 745 = C. Toona.
montana, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) 415.—Venezuela.
multijuga, Kurz, in Journ. As. Soc. Beng. xli. (1872) II. 297 = C. Toona.
odorata, Blanco, Fl. Filip. ed. I. 184 = C. Toona.
odorata, Cham. & Schlecht. in Linnaea, vi. (1831) 422 = Swietenia humilis.
odorata, Griseb. in Mem. Am. Acad. N. S. viii. (1860) 169 = Glaziovii.
odorata, *Linn. Syst.* ed. X. 940.—Am. austr. (? Guiana)
W. Indies (Jamaica, Guadeloupe)

CEDRELA :—
odorata, Ruiz & Pav. Fl. Per. iii. 9 = angustifolia.
odorata, Vell. Fl. Flum. 74; ii. t. 67 = Velloziana.
paraguaiensis, *Mart. in Flora*, xx. (1837) *Beibl.* 93.—Paraguay.
Rosmarinus, Lour. Fl. Cochinch. 160 = Baeckia cochinchinensis.
serrata, Royle, Illustr. Bot. Himal. 144. t. 25 = C. Toona.
serrulata, *Miq. Fl. Ind. Bat. Suppl.* i. 508.—Sumatra.
sinensis, *Juss. in Mém. Mus. Par.* xix. (1830) 255.—China.
Teysmanni, *Hassk. Hort. Bogor.* i. 131.—Java.
Toona, *Roxb. ex Rottl. in Ges. Naturf. Fr. Neue Schr.* ii. (1803) 198.—Malaya; Austral.
Velloziana, *M. Roem. Syn. Hesper.* 137.—Am. trop.
velutina, DC. Prod. i. 625 = Chickrassia tabularis.
villosa, Roxb. Hort. Beng. 18 = Chickrassia tabularis.

CEDRELLA, Scop. Introd. 324 (1777) = **Cedrela**, R. Br.

CEDRONELLA, Riv. ex Rupp. Fl. Jen. ed. Hall. 225 (1745); Moench, Meth. 411 (1794). *LABIATAE*, Benth. & Hook. f. ii. 1200.
breviflora, *A. Gray, in Proc. Am. Acad.* xx. (1885) 309.—Arizona.
cana, *Hook. Bot. Mag.* t. 4618.—Mexic.
canariensis, *Webb & Berth. Phyt. Canar.* iii. 87.—Ins. Canar.
cordata, *Benth. Lab. Gen. et Sp.* 502.—Am. bor.
hastifolia, Regel, Ind. Sem. Hort. Petrop. (1868) 85 = cana.
japonica, *Hassk. in Verh. Natuurk. Ver. Ned. Ind.* i. (1856) 36.—Japon.
mexicana, *Benth. Lab. Gen. et Sp.* 502.—Mexic.
micrantha, *A. Gray, in Proc. Am. Acad.* viii. (1870) 369.—Am. bor. occ.
pallida, *Lindl. Bot. Reg.* (1846) t. 29.—Am. bor. occ.
triphylla, *Moench, Meth.* 411.—Ins. Canar.

CEDROTA, Schreb. Gen. i. 259 (1789) = **Ocotea**, Aubl. (Laurin.).
guianensis, Blanco, Fl. Filip. ed. II. 213 = Pisonia umbellifera.
guianensis, Raeusch. Nom. ed. III. 111 = O. commutata.
longifolia, Willd. Sp. Pl. ii. 338 = O. commutata.

CEDRUS, Mill. Gard. Dict. ed. VII (1759) = **Cedrela**, Linn. (Meliac.).
alternifolia, Mill. l. c. ed. VIII. n. 3 = Cedrela alternifolia.
Mahogani, Mill. l. c. n. 2 = Swietenia Mahagoni.
odorata, Mill. l. c. n. 1 = Cedrela odorata.

CEDRUS, [Tourn.] Mill. Gard. Dict. ed. III (1737); Loud. Arb. Brit. iv. 2402 (1838). *CONIFERAE*, Benth. & Hook. f. iii. 439.
africana, Gord. ex Knight, Syn. Conif. 42 = Libani.
argentea, Hort. ex Carr. Conif. ed. I. 286 = Libani.
atlantica, Manetti, Cat. Hort. Madoet. Suppl. 8 = Libani.
Deodara, Loud. Arbor. Brit. iv. 2428 = Libani.
elegans, Knight, Syn. Conif. 42 = Libani.
indica, Chambr. Trait. Prat. Arbr. Résin. Conif. 341 = Libani.
Libani, *Barrel. Ic.* 499.—As. Min.; Syria; Affghan.; Reg. Himal.; Alger.
patula, C. Koch, Dendrol. ii. II. 268 = Libani.

CEIBA, Medic. Malv. 15 (1787); Gaertn. Fruct. ii. 244. t. 133 (1791) = **Eriodendron**, DC. (Malvac.).
Casearia, Medic. l. c. 16 = E. orientale.
pentandra, Gaertn. Fruct. ii. 244. t. 133 = E. anfractuosum.

CELAENA, Wedd. Chlor. And. i. 230 (1855) = **Oligandra**, Less. (Compos.).
pachymorpha, Wedd. l. c. = O. pachymorpha.

CELASINE, Pritz. Ic. Ind. 284, sphalm. (1855) = **Gelasine**, Herb. (Irid.).

CELASTRUS, Linn. Gen. ed. I. 59 (1737). *CELASTRINEAE*, Benth. & Hook. f. i. 364 [365].
CATHA, G. Don, Gen. Syst. ii. 11 (1832).
GUEVINIA, Hort. Par. ex Decne. in Rev. Hortic. Sér. II. iv. (1845–46) 425.
ORIXA, Thunb. Nov. Gen. 56 (1783); Fl. Jap. 3 (1784).
SEMARILLA, Rafin. Sylva Tellur. 146 (1838).
acuminatus, *Linn. f. Suppl.* 154.—Afr. austr.
acuminatus, *Rafin. New. Fl. Am.* iii. 14.—Am. bor.
acuminatus, Wall. Cat. n. 4342 = Chailletia gelonioides.
adenophylla, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 85 = Ilex crenata.
alatus, Thunb. Fl. Jap. 98 = Evonymus Thunbergianus.
alnifolius, D. Don, Prod. Fl. Nep. 190 = paniculatus.
alpestris, *Blume, Bijdr.* 1145.—Java.
andongensis, *Oliver, Fl. Trop. Afr.* i. 361.—Afr. trop.
angularis, *Sond. in Harv. & Sond. Fl. Cap.* i. 460.—Afr. austr.
angulata, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 455.—China.
aphyllus, *Schlecht. in Linnaea*, xv. (1841) 458.—Mexic.
aquifolius, *Regel, Ind. Sem. Hort. Petrop.* (1856) 36.—Hab. ?
arbutifolius, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 133.—Abyss.
articulatus, *Thunb. Fl. Jap.* 97.—Japon.
Atkaio, *A. Rich. Tent. Fl. Abyss.* i. 132.—Abyssin.
attenuatus, Wall. Cat. n. 4319 = Gymnosporia neglecta.
auriculatus, Vitm. Summa, ii. 31, in syn. = articulatus.
australis, *Harv. & F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 41.—Austral.
barbarus, Linn. Mant. ii. 211, in obs. = Cassine barbara.
bilocularis, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 31.—Austral.
bivalvis, Jack, in Malay Misc. i. (1820) n. v. 19 = Microtropis bivalvis.
Boaria, Baill. Hist. Pl. vi. 26 = Maytenus chilensis.
bullatus, Hort. ex Steud. Nom. ed. II. i. 314 = pyracanthus.
bullatus, Linn. Sp. Pl. 196 = scandens.
buxifolia, Drège, ex Walp. Ann. i. 189 = Catha patens.
buxifolius, *Linn. Sp. Pl.* 197.—Afr. austr.
buxifolius, Roxb. Hort. Beng. 18, nomen = Gymnosporia Wallichiana.
campestris, Eckl. & Zeyh. Enum. 119 = pyracanthus.
cantonensis, *Hance, in Journ. Bot.* xxiii. (1885) 323.—China.
capensis, Thunb. ex Steud. Nom. ed. II. i. 314 = excisus.
capitatus, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 458.—Afr. austr.
cassinifolius, Salisb. Prod. 141 = lucidus.
cassinoides, L'Hérit. Sert. Angl. 6. t. 10 = Gymnosporia cassinoides.
cernuus, *Thunb. Prod. Pl. Cap.* 42.—Afr. austr.
Championi, *Benth. in Hook. Kew Journ.* iii. (1851) 334.—Ins. Hongkong.
ciliidens, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 85 = flagellaris.
collinus, Eckl. & Zeyh. Enum. 119 = undatus.
Colpoon, Hort. ex Steud. Nom. ed. II. i. 315 = pyracanthus.
concavus, Lam. Tabl. Encyc. ii. 93 = lucidus.
confertus, *Ruiz & Pav. Fl. Per.* iii. 7.—Peruv.
cordatus, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* 454.—Afr. austr.
coriaceus, Guill. & Perr. Fl. Seneg. Tent. i. 142 = Gymnosporia montana, senegalensis.
corniculatus, *L'Hérit. Sert. Angl.* 6.—Afr. austr.
crassifolius, Salisb. Prod. 141 = cassinoides.
crenatus, *Forst. f. Prod.* 19.—Ins. Marchion.
crenatus, Roth, Nov. Pl. Sp. 156 = Gymnosporia montana.
crenulatus, *Wall. Cat.* n. 4323.—Siam.
crispulus, *Regel, Gartenfl.* (1860) 407.—China.
crispus, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 23 = Euclea ovata.
Cunninghamii, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 30.—Austral.
cymatodes, Spreng. Syst. i. 775 = undatus.

CELASTRUS :—

cymosus, Eckl. & Zeyh. Enum. 121 = heterophyllus.
cymosus, Soland. in Bot. Mag. t. 2070 = buxifolius.
decolor, Delile, Voy. Méroë, 100. t. 64. f. 6 = Gymnosporia montana, senegalensis.
dependens, Wall. Cat. 4302; et in Roxb. Fl. Ind. ed. Carey, i. 389 = paniculatus.
diffusus, G. Don, Gen. Syst. ii. 6 = Ventilago leiocarpa.
dilatatus, *Thunb. in Trans. Linn. Soc.* ii. (1794) 332. —Japon.
dispermus, *F. Muell. in Trans. Phil. Inst. Vict.* iii. (1859) 31.—Austral.
dubius, *Spreng. Syst.* i. 774.—Ind. occ.
dumetorum, Eckl. & Zeyh. Enum. 119 = undatus.
edulis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 131 = serratus.
edulis, Vahl, Symb. Bot. i. 21 = Catha edulis.
ellipticus, *Thunb. in Hoffm. Phytogr. Blaetter*, i. (1803) 22.—Afr. austr.
emarginatus, Ruiz & Pav. Fl. Per. iii. 6 = retusus.
emarginatus, Willd. Sp. Pl. i. 1128 = Gymnosporia emarginata.
empleurifolius, Eckl. & Zeyh. Enum. 121 = buxifolius.
enonymifolius, *Burm. f. Fl. Cap. Prod.* 6.—Afr. austr.
enonymoides, *Welw. ex Oliver, Fl. Trop. Afr.* i. 362.—Afr. trop.
europaeus, Boiss. Elench. 29 = Gymnosporia montana.
excisus, *Thunb. Fl. Cap.* ii. 119.—Afr. austr.
filiformis, Linn. f. Suppl. 153 (= *Mystroxylon filiforme*).—Afr. austr.
Finlaysonianus, *Wall. Cat.* n. 4324.—Siam; China.
flagellaris, *Rupr. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1857) 357.—Reg. Amur.
flexuosus, *Thunb. Fl. Cap.* ii. 124.—Afr. austr.
floribundus, *Span. in Linnaea*, xv. (1841) 186.—Hab.?
Fournieri, *Panch. & Sebert, Not. Bois Nouv. Caléd.* 234.—N. Caled.
glaucus, R. Br. in Salt, Voy. Abyss. App. 64 = Gymnosporia montana.
glaucus, Vahl, Symb. Bot. ii. 42 = Elaeodendron glaucum.
glomeratus, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 458 = heterophyllus.
glomeratus, E. Mey. ex Walp. Rep. v. 401 = patens.
goniocaulis, Eckl. & Zeyh. Enum. 120 = buxifolius.
gracilipes, *Welw. ex Oliver, Fl. Trop. Afr.* i. 361.—Afr. trop.
Haenkea, Spreng. Syst. iv. Cur. Post. 88 = Schoepfia flexuosa.
Hamelii, Spreng. Syst. i. 774 = Rhamnus racemosus?
heterophyllus, *Eckl. & Zeyh. Enum.* 120.—Afr. austr.
heterophyllus, E. Mey. ex Sond. in Harv. & Sond. Fl. Cap. i. 467 = Cassine latifolia.
heterophyllus, *Savi, in Mem. Torino*, xxxviii. (1835) 173.—Hab.?
Heyneana, Roth, in Roem. & Schult. Syst. v. 421 = Gymnosporia Heyneana.
Hindsii, *Benth. in Hook. Kew Journ.* iii. (1851) 334 = monospermus.
hispanicus, Hort. ex DC. Prod. ii. 8 = multiflorus.
huillensis, *Welw. ex Oliver, Fl. Trop. Afr.* i. 364.—Afr. trop.
humilis, Eckl. & Zeyh. Enum. 120 = buxifolius.
ilicifolius, Schrad. in Goett. Gel. Anz. i. (1821) 716 = Maytenus truncatus.
ilicinus, *Burch. ex Loud. Hort. Brit.* 83, nomen.—Afr. austr.
ilicinus, Krauss. ex Sond. in Harv. & Sond. Fl. Cap. i. 457 = procumbens.
inflexus, Zeyh. ex Sond. l. c. 455 = polyacanthus.
integer, Thunb. Fl. Cap. ii. 123 = pyracanthus.
integrifolius, *Linn. f. Suppl.* 153.—Afr. austr.
Jackianus, Steud. Nom. ed. II. i. 314 = lucidus.
japonicus, *C. Koch, Dendrol.* i. 625.—Japon.
Jodina, Steud. Nom. ed. II. i. 314 = Iodina rhombifolia.
kiusiana, *Franch. & Sav. Enum. Pl. Jap.* ii. 314.—Japon.
lanceolatus, *E. Mey. ex Sond. in Harv. & Sond. Fl. Cap.* i. 456.—Afr. austr.
lanceolatus, *Spreng. Syst.* i. 775.—Ins. Mascar.
lancifolius, *Thonn. in Schum. & Thonn. Beskr. Guin. Pl.* 132.—Afr. trop.
laurifolius, *A. Rich. Tent. Fl. Abyss.* i. 130.—Abyssin.

CELASTRUS :—

laurinus, *Thunb. Prod. Pl. Cap.* 42.—Afr. austr.
leptopus, Bernh. ex Walp. Rep. v. 401 = buxifolius.
linearis, Eckl. & Zeyh. Enum. 121 = polyacanthus.
linearis, *Linn. f. Suppl.* 153.—Afr. austr.
lineatus, D. Dietr. Syn. i. 815 = linearis.
lucidus, *Linn. Mant.* i. 49.—Afr. austr.
lucidus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 400.—Ind. or.
luteolus, *Delile, in Ann. Sc. Nat.* Sér. II. xx. (1843) 90.—Afr. trop.
lycioides, *Brouss. ex Willd. in Roem. & Schult. Syst.* v. 427.—Ins. Canar.
macrocarpus, *Ruiz & Pav. Fl. Per.* iii. 8. t. 230. f. 16.—Peruv.
macrophyllus, Welw. ex Oliver, Fl. Trop. Afr. i. 364 = lancifolius.
magellanicus, DC. Prod. ii. 8 = Maytenus magellanicus.
mauritianus, *Stadtm. ex Willem. in Ust. Ann. Bot.* xviii. (1796) 22.—Ins. Maurit.
Maytenus, Willd. Sp. Pl. i. 1127 = Maytenus Boaria.
Metzianus, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) 1. 448 = paniculatus.
mexicanus, *Moç. & Sesse, ex DC. Prod.* ii. 8.—Mexic.
micrantha, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 625.—Ins. Molucc.
microcarpus, *D. Don, Prod. Fl. Nep.* 191.—Reg. Himal.
microphyllus, Eckl. & Zeyh. Enum. 119 = acuminatus.
microphyllus, *Linn. f. Suppl.* 154.—Afr. austr.
mollis, *Decne. in Rev. Hortic.* Sér. II. iv. (1845–46) 425.—Hab.?
monospermus, *Roxb. Hort. Beng.* 18; *Fl. Ind.* i. 625.—Ind. or.; China.
montanus, Roth, Nov. Pl. Sp. 154 = Gymnosporia montana.
mossambicensis, *Klotzsch. in Peters, Reise Mossamb. Bot.* i. 112.—Afr. trop.
mucronatus, Eckl. & Zeyh. Enum. 119 = acuminatus.
Muelleri, *Benth. Fl. Austral.* i. 399.—Austral.
multiflorus, Eckl. & Zeyh. Enum. 121 = buxifolius.
multiflorus, *Lam. Encyc.* i. 661.—Afr. austr.
multiflorus, Roxb. Hort. Beng. 18; Fl. Ind. 622 = paniculatus.
myrtifolius, *Linn. Sp. Pl.* 196.—Jamaica.
neglecta, Wall. Cat. n. 4341 = Gymnosporia neglecta.
nemorosus, *Eckl. & Zeyh. Enum.* 120.—Afr. austr.
nepalensis, Steud. Nom. ed. II. i. 315 = Pittosporum floribundum.
nossibaeus, *O. Hoffm. Sert. Madag.* 12 (= Alsodeia).—Madag.
nutans, Roxb. Hort. Beng. 18; Fl. Ind. i. 623 = paniculatus.
obovatus, Hochst. ex Oliver, Fl. Trop. Afr. i. 362 = Gymnosporia montana.
obscurus, A. Rich. Tent. Fl. Abyss. i. 131 = serratus.
obtusatus, Presl, Bot. Bemerk. 34 = Simmondsia californica.
obtusifolius, Roxb. Hort. Beng. 86; Fl. Ind. i. 625 = Gymnosporia trigyna.
obtusus, Thunb. in Hoffm. Phytogr. Blaetter, i. (1803) 22 = pyracanthus.
octogonus, L'Hérit. Sert. Angl. 6 = Maytenus octogonus.
octogonus, Hort. ex Steud. Nom. ed. II. i. 315 = pyracanthus.
oleifolius, Pers. Syn. i. 327 = oleoides.
oleoides, *Lam. Tabl. Encyc.* ii. 93.—Afr. austr.
oppositus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 398 = Pleurostylia Wightii.
orbicularis, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 423 = Maytenus uliginosus.
orbiculatus, Lam. Tabl. Encyc. ii. 94 = articulatus.
Orixa, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1846) 150.—Japon.
ovalifolius, Steud. Nom. ed. II. i. 315 = Gymnosporia ovata.
ovatus, *Hill, Veg. Syst.* xiii. 62; *Hort. Kew.* 437. t. 16.—Ins. Bahama.
ovatus, Wall. Cat. n. 4308 = Gymnosporia ovata.
oxyphyllus, Wall. Cat. n. 4312 = Gymnosporia acuminata.
pallidus, Wall. Cat. n. 4307 = Gymnosporia montana.
paniculatus, *Willd. Sp. Pl.* i. 1125.—As. trop.; Malaya.
parvifolius, *A. Rich. Ess. Fl. Cub.* 349.—Cuba.
parvifolius, Eckl. & Zeyh. Enum. 121 = heterophyllus.
parviflorus, *Vahl, Symb. Bot.* i. 21.—Arabia.
patens, *Eckl. & Zeyh. l. c.* 120.—Afr. austr.

CELASTRUS :—

pauciflorus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 400.—Ind. or.
pedunculatus, *Sond. in Harv. & Sond. Fl. Cap.* i. 454.—Afr. austr.
pentagynus, *Zipp. ex Span. in Linnaea*, xv. (1841) 186.—Malaya?
peruvianus, F. G. Dietr. Vollst. Lexik. Gaertn. ii. 111 = retusus.
phyllacanthus, L'Hérit. Sert. Angl. 6 = Gymnosporia montana.
Plectronia, DC. Prod. ii. 9 = acuminatus.
polyacanthus, *Sond. in Harv. & Sond. Fl. Cap.* i. 455.—Afr. austr.
polyanthemos, *Eckl. & Zeyh. Enum.* 121.—Afr. austr.
polybotrys, *Turcz. in Bull. Soc. Nat. Mosc.* xxxi. (1858) 1. 449.—Ins. Philipp.
populifolius, *Lam. Tabl. Encyc.* ii. 94 = acuminatus.
procumbens, *Linn. f. Suppl.* 153.—Afr. austr.
pterocarpus, DC. Prod. ii. 5 = Pterocelastrus tricuspidatus.
pubescens, *Wall. Cat.* n. 4303.—Ind. or.
punctatus, Thunb. Fl. Jap. 97 = articulatus.
punctulatus, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. II. (1846) 150.—Japon.
pyracanthifolius, Salisb. Prod. 141 = pyracanthus.
pyracanthus, *Linn. Sp. Pl.* 197.—Afr. austr.
pyrifolia, Herb. Franq. ex Oliver, Fl. Trop. Afr. i. 362 = serratus.
pyrius, Willem. in Ust. Ann. Bot. xviii. (1796) 21 = Gymnosporia trigyna.
quadrangulatus, *Schrad. in Goett. Gel. Anz.* i. (1821) 716.—Bras.
racemosus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 599.—Japon.
racemosus, Wall. Cat. n. 4320 = Parastemon urophyllum.
racemulosus, *Hassk. Hort. Bogor.* i. 155.—Java.
refractus, E. Mey. ex Sond. & Harv. Fl. Cap. i. 467 = Cassine scandens.
repandus, *Blume, Bijdr.* 1145.—Java.
retusus, *Poir. Encyc. Suppl.* ii. 146.—Peruv.
rhamnoides, *Poir. l. c.* 145.—Afr. austr.
rhombifolius, *Eckl. & Zeyh. Enum.* 120.—Afr. austr.
rhombifolius, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 170 = Iodina rhombifolia.
Richii, *A. Gray, Bot. U. St. Expl. Exped.* i. 289.—Ins. Fiji.
rigidus, Eckl. & Zeyh. Enum. 121 = capitatus.
rigidus, *Thunb. Fl. Cap.* ii. 124.—Afr. austr.
rigidus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 396 = Gymnosporia Wallichiana.
robustus, Roxb. Hort. Beng. 18; Fl. Ind. i. 626 = Kurrimia pulcherrima.
rostratus, Thunb. Prod. Pl. Cap. 42 = Pterocelastrus rostratus.
Rothianus, Roem. & Schult. Syst. v. 423 = paniculatus.
Rothianus, Wight & Arn. Prod. i. 159 = Gymnosporia Rothiana.
rotundifolius, Thunb. Prod. Pl. Cap. 42 = Dovyalis cclastroides.
Royleanus, Wall. Cat. n. 4317 = Gymnosporia Royleana.
ruber, Gerrard, ex Oliver, Fl. Trop. Afr. i. 363 = mossambicensis.
ruber, *Harv. in Harv. & Sond. Fl. Cap.* ii. 592.—Afr. austr.
rufus, Wall. Cat. n. 4309 = Gymnosporia rufa.
rupestris, Eckl. & Zeyh. Enum. 119 = acuminatus.
saxatilis, *Burch. Trav.* ii. 264.—Afr. austr.
scandens, *Linn. Sp. Pl.* 196.—Am. bor.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 132 = serratus.
scmiarillatus, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) 1. 599.—Java.
senegalensis, Lam. Encyc. i. 661 = Gymnosporia montana.
sepiarius, *Dennst. Schluess. Hort. Malab.* 31.—Ind. or.
serratus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 131.—Abyssin.
serrulatus, R. Br. in Salt, Abyss. App. 64 = serratus.
serrulatus, Roth, Nov. Pl. Sp. 155 = Gymnosporia Heyneana.
Sieberi, Bernh. ex Harv. & Sond. Fl. Cap. i. 473 = Ilex capensis.
sinuato-dentatus, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 130 = luteolus.
spathephyllus, Eckl. & Zeyh. Enum. 121 = buxifolius.

CELASTRUS:—
spicatus, Vell. Fl. Flum. 92; ii. t. 138 = Gouania corylifolia.
spinosus, Royle, Illustr. Bot. Himal. 167 = Gymnosporia Royleana.
stenophyllus, Eckl. & Zeyh. Enum. 122 = linearis.
striatus, Thunb. Fl. Jap. 98 = Enonymus Thunbergianus.
stylosus, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 401.—Reg. Himal.
subspicatus, *Hook. Ic. Pl.* t. 482.—Hab.?
Tatarinowii, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 357 = articulatus.
tenuispinus, *Sond. in Harv. & Sond. Fl. Cap.* i. 456.—Afr. austr.
tetragonus, Thunb. Prod. Pl. Cap. 42 = Cassine scandens.
tricuspidatus, Lam. Tabl. Encyc. ii. 93 = Pterocelastrus tricuspidatus.
trigynus, Lam. l. c. 94 = Gymnosporia trigyna.
Tsaad, *Ferr. & Gal. ex Walp. Ann.* iii. 841.—Abyssin.
umbellatus, Vell. Fl. Flum. 92; ii. t. 137 = Reissekia cordifolia.
uncinatus, Ruiz & Pav. Fl. Per. iii. 7 = Maytenus chilensis.
undatus, *Thunb. Prod. Pl. Cap.* 42.—Afr. austr.
undulatus, Lam. Encyc. i. 662 = Pittosporum Senacia.
venenatus, Eckl. & Zeyh. Enum. 121 = buxifolius.
venulosus, *Wall. Cat.* n. 4321.—Reg. Himal.
verrucosus, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 453.—Afr. austr.
verticillatus, Roxb. Hort. Beng. 18; Fl. Ind. i. 624 = Pittosporum floribundum.
verticillatus, Ruiz & Pav. Fl. Per. iii. 6 = Maytenus verticillatus.
Wallichianus, Spreng. Syst. v. Index, 150 = Gymnosporia Wallichiana.
Wallichianus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 400 = Gymnosporia Heyneana.
Wightianus, Wall. Cat. n. 4322 = Pleurostylia Wightii.
Zeyheri, *Sond. in Harv. & Sond. Fl. Cap.* i. 456.—Afr. austr.
zeylanicus, Roth, ex Roem. & Schult. Syst. v. 427 = Scutia Commersonii.

CELEBNIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid?).

CELMISIA, Cass. in Dict. Sc. Nat. vii. 356 (1817) = **Alciope**, DC. (Compos.).
rotundifolia, Cass. l. c. 357 = A. tabularis.

CELMISIA, Cass. in Dict. Sc. Nat. xxxvii. 259 (1825). *COMPOSITAE*, Benth. & Hook. f. ii. 278.
asteliaefolia, Hook. f. Fl. Antarct. 35 = longifolia.
bellidioides, *Hook. f. Handb. New Zeal. Fl.* 135.—N. Zel.
cordatifolia, *J. Buch. in Trans. N. Z. Inst.* xi. (1879) 427. t. 18.—N. Zel.
coriacea, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 119.—N. Zel.
Dallii, *J. Buch. in Trans. N. Z. Inst.* xiv. (1882) 355. t. 35.—N. Zel.
densiflora, *Hook. f. Handb. New Zeal. Fl.* 130.—N. Zel.
discolor, *Hook. f. Fl. N. Zel.* i. 123.—N. Zel.
glandulosa, *Hook. f. l. c.* 124.—N. Zel.
gracilenta, *Hook. f. Fl. Antarct.* 35.—N. Zel.
graminifolia, *Hook. f. l. c.*—N. Zel.
Haastii, *Hook. f. Handb. New Zeal. Fl.* 131.—N. Zel.
Hectori, *Hook. f. l. c.* 135.—N. Zel.
hieracifolia, *Hook. f. Fl. N. Zel.* 124. t. 34.—N. Zel.
holosericea, *Hook. f. Fl. Antarct.* 36.—N. Zel.
incana, *Hook. f. Fl. N. Zel.* 123. t. 34.—N. Zel.
laricifolia, *Hook. f. l. c.* 331.—N. Zel.
lateralis, *J. Buch. in Trans. N. Z. Inst.* iv. (1872) 226.—N. Zel.
Lindsayi, *Hook. f. Handb. New Zeal. Fl.* 131.—N. Zel.
linearis, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 337.—N. Zel.
longifolia, *Cass. in Dict. Sc. Nat.* xxxvii. 259.—Austral.
Lyallii, *Hook. f. Handb. N. Zeal. Fl.* 133.—N. Zel.
Mackaui, *Raoul, Choix Pl. Nouv. Zél.* 19. t. 14.—N. Zel.
Munroi, *Hook. f. Handb. New Zeal. Fl.* 133.—N. Zel.
petiolata, *Hook. f. l. c.* 135.—N. Zel.
ramulosa, *Hook. f. l. c.* 733.—N. Zel.
rupestris, *Cheesem. in Trans. N. Z. Inst.* xvi. (1884) 409.—N. Zel.
Leehleri, *Sch. Bip. in Bonplandia*, iv. (1856) 54, nomen.—Peruv.
sessiliflora, *Hook. f. Handb. New Zeal. Fl.* 135.—N. Zel.
Sinclairii, *Hook. f. l. c.* 132.—N. Zel.
spathulata, A. Cunn. in DC. Prod. v. 209 = Trichocline scapigera.
spectabilis, *Hook. f. Fl. Antarct.* 35.—N. Zel.
Traversii, *Hook. f. Handb. New Zeal. Fl.* 134.—N. Zel.
verbascifolia, *Hook. f. Fl. N. Zel.* 121.—N. Zel.
vernicosa, *Hook. f. Fl. Antarct.* 34. t. 26, 27.—Ins. Auckland.
viscosa, *Hook. f. Handb. New Zeal. Fl.* 133.—N. Zel.
Walkeri, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 549.—N. Zel.

CELOSIA, Linn. Gen. ed. I. 34 (1737). *AMARANTACEAE*, Benth. & Hook. f. iii. 24.
AMARANTHUS, Adans. Fam. ii. 269 (1763).
GONUFAS, Rafin. Sylva Tellur. 124 (1838).
KOKERA, Adans. Fam. ii. 269 (1763).
LAGREZIA, Moq. in DC. Prod. xiii. ii. 252 (1849).
LESTIBUDESIA, Thou. Hist. Vég. Iles Afr. 53. t. 16 (1806).
LOPHOXERA, Rafin. Fl. Tellur. iii. 42 (1836).
SUKANA, Adans. Fam. ii. 269 (1763).
acroporoides, *Hochst. ex Benth. & Hook. f. Gen.* iii. 25.—Abyss.
acroprosoides, Hochst. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 140 = trigyna.
adoënsis, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 211 = trigyna.
albida, Willd. Sp. Pl. i. 1197 = Allmania albida.
allmanoides, *Moq. in DC. Prod.* xiii. ii. 243.—Burma.
altissima, Salzm. ex Moq. l. c. 381 = Telanthera ramosissima.
amaranthoides, Medic. Malv. 92 = trigyna.
amphibia, Salzm. ex Moq. in DC. Prod. xiii. ii. 363 = Telanthera philoxeroides.
anthelminthica, *Aschers. in Schweinf. Beitr. Fl. Aethiop.* 176.—Afr. trop.
arborescens, Spreng. Syst. i. 815 = Deeringia altissima.
arenaria, Salzm. ex Seub. in Mart. Fl. Bras. v. i. 170 = Telanthera maritima.
argentea, *Linn. Sp. Pl.* 205.—Cosmop. trop.
aspera, Roth, Nov. Pl. Sp. 173 = Allmania nodiflora.
aurea, T. Moore, Flor. Mag. i. (1861) t. 47 = cristata.
axillaris, Mart. Hort. Erlang. 70 = Alternanthera spinosa.
baccata, Retz. Obs. fasc. v. 23 = Deeringia celosioides.
bicolor, Blanco, Fl. Filip. ed. I. 191 = philippica.
Boivinii, *Hook. f. in Benth. & Hook. f. Gen.* iii. 25.—Madag.
brasiliensis, *Moq. in DC. Prod.* xiii. ii. 239.—Bras.
castrensis, Linn. Sp. Pl. ed. II. 297 = cristata.
caudata, *Vahl, Symb. Bot.* i. 21.—Arabia.
cernua, *Andr. Bot. Rep.* x. t. 635.—Ind. or.
cernua, Mairet, ex Moq. in DC. Prod. xiii. ii. 351 = Alternanthera canescens.
cernua, Roxb. Hort. Beng. 19; Fl. Ind. i. 680 = cristata.
coccinea, Linn. Sp. Pl. ed. II. 297 = cristata.
comosa, Retz. Obs. fasc. vi. 26 = cristata.
corymbifera, *Didr. Ind. Sem. Hort. Havn.* (1849) 13.—Hab.?
corymbosa, Retz. Obs. fasc. i. 13 = Polycarpaea corymbosa.
cristata, *Linn. Sp. Pl.* 205.—Cosmop. trop.
cymosa, *Seub. in Mart. Fl. Bras.* v. i. 245.—Bras.
decumbens, Hort. ex. Moq. in DC. Prod. xiii. ii. 410 = Gomphrena decumbens.
dichotoma, Heyne, in Roth, Nov. Pl. Sp. 172 = Allmania nodiflora.
diffusa, Hoffmgg. ex Seub. in Mart. Fl. Bras. v. i. 179 = Telanthera ramosissima.
echinata, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 531 = Alternanthera Achyrantha.
elongata, Spreng. in Schrad. Journ. (1800) ii. 196 = Telanthera portigens.
eriantha, Vahl, ex Moq. in DC. Prod. xiii. ii. 386 = Gomphrena eriantha.
exstipulata, *Hornem. ex Schrank, in Syll. Ratisb.* i. (1824) 202.—Nepal.
floribunda, *A. Gray, in Proc. Am. Acad.* v. (1861) 167.—Calif.
glauca, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 198 = patula.
glauca, Wendl. Beobacht. 43; Hort. Herrenh. 4. t. 2 = Hermbstaedtia glauca.
gnaphalioides, Linn. f. Suppl. 161 = Gomphrena gnaphalioides.
grandifolia, *Moq. in DC. Prod.* xiii. ii. 238.—Bras.
humifusa, Willd. ex Roem. & Schult. Syst. v. 531 = Telanthera caracasana.
Huttoni, *Mast. in Gard. Chron.* (1872) 215.—Java.
intermedia, Hochst. ex Oliver, in Trans. Linn. Soc. xxix. (1875) 140 = trigyna.
interrupta, Seub. in Mart. Fl. Bras. v. i. 165 = Froelichia interrupta.
japonica, *Mart. Hort. Erlang.* 70.—Japon.?
lactea, Hort. ex Moq. in DC. Prod. xiii. ii. 300 = Aerva sanguinolenta.
lanata, Blanco, Fl. Filip. ed. II. 134 = Aerva lanata.
lanata, Linn. Sp. Pl. 205 = Aerva tomentosa.
lanata, Poir. ex Steud. Nom. ed. II. i. 315 = Gomphrena gnaphalioides.
lappacea, Medic. Beobacht. (1783) 147 = Pupalia atropurpurea.
latifolia, *Steud. Nom.* ed. II. i. 315.—Java.
laxa, *Schur & Thonn. Beskr. Guin. Pl.* 141.—Afr. trop.
leptostachya, *Benth. in Hook. Niger Fl.* 491.—Afr. trop.
linearis, Sweet, ex Hook. f. Fl. Brit. Ind. iv. 714 = argentea.
longifolia, *Mart. Nov. Gen. et Sp.* ii. 63. t. 157, 158. f. 2.—Bras.
madagascariensis, *Poir. Encyc.* v. 39.—Madag.
major, *Griseb. in Goett. Abh.* xix. (1874) 79.—Reg. Argent.
margaritacea, Linn. Sp. Pl. ed. II. 297 = argentea.
marylandica, Retz. Obs. fasc. iii. 27 = argentea.
maritima, Salzm. ex Moq. in DC. Prod. xiii. ii. 341 = Philoxerus vermicularis.
melanocarpos, Poir. Encyc. Suppl. iv. 318 = trigyna.
missionis, Wall. Cat. n. 6915 = Banalia thyrsiflora.
Monsonia, Retz. Obs. fasc. ii. 13 = Aerva Monsonia.
Moquini, *Guill. ex DC. Prod.* xiii. ii. 239.—Mexic.
mucronata, Herb. ex Moq. in DC. l. c. 248 = Allmania albida.
muricata, Spreng. Syst. iv. Cur. Post. 102 = Cladostachys muricata.
nana, Blanco, Fl. Filip. ed. I. 192 = Ammania baccifera.
nitida, Vahl, Symb. Bot. iii. 44 = paniculata.
nodiflora, Linn. Sp. Pl. 205 = Allmania nodiflora.
nodiflora, Salzm. ex Moq. in DC. Prod. xiii. ii. 364 = Telanthera polygonoides.
nodiflora, Wall. Cat. n. 6890 = Allmania albida.
nudiflora, Pav. ex Moq. in DC. Prod. xiii. ii. 341 = Philoxerus vermicularis.
odorata, *Burch. ex Moq. l. c.* 244.—Afr. austr.
pallida, Hort. Vindob. ex Moq. l. c. 248 = Allmania albida.
pallida, Salisb. Prod. 145 = argentea.
Palmeri, *S. Wats. in Proc. Am. Acad.* xviii. (1882 83) 143.—Calif.
paniculata, *Linn. Sp. Pl.* 206.—Florida; Ind. occ.
parviflora, Vahl, ex Moq. in DC. Prod. xiii. ii. 346 = Iresine gracilis.
patula, *Willd. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 198.—Ind. or.
peruviana, Van Spand. ex Moq. in DC. Prod. xiii. ii. 365 = Telanthera frutescens.
peruviana, Zuccagni, in Roem. Collect. 133 = Telanthera porrigens.
philippica, *Steud. Nom.* ed. II. i. 316.—Ins. Philipp.
phytolaccaefolia, Juss. ex Moq. in DC. Prod. xiii. ii. 253 = madagascariensis.
piloselloides, Poit. ex Moq. l. c. 337 = Guilleminea lanuginosa.
polygonoides, *Retz. Obs. fasc.* ii. 12.—Ind. or.
polygonoides, Wall. Cat. n. 6913 = pulchella.
polysperma, Roxb. Hort. Beng. 84; Fl. Ind. i. 681 = Deeringia polysperma.
populifolia, *Moq. in DC. Prod.* xiii. ii. 239.—Abyss.
procumbens, Hort. ex Moq. l. c. 410 = Gomphrena decumbens.

CELOSIA :—
procumbens, Jacq. Misc. ii. 344 = Froelichia interrupta.
prostrata, Hort. ex Moq. in DC. Prod. xiii. II. 410 = Gomphrena decumbens.
pulchella, *Moq. l. c.* 238.—Ind. or.
purpurea, St. Hil. ex Steud. Nom. ed. II. i. 315, 316 = cristata.
pyramidalis, Burm. f. Fl. Ind. 62 = cristata, Allmania albida.
recurva, Burch. Trav. ii. 226 = Hermbstaedtia elegans.
sanguinea, *Hoffmgg. Verz. Pfl.* ii. 85.—Ind. or.
sinuata, *Mart. ex Steud. Nom.* ed. I. 171.—Hab.?
sparsa, Forsyth, ex Moq. in DC. Prod. xiii. II. 250 = Chamissoa altissima.
spathulata, *Boj. Hort. Maurit.* 267.—Madag.
spathulata, Saint-Elme, ex Moq. in DC. Prod. xiii. II. 253 = madagascariensis.
spicata, André, ex Moq. l. c. 143 = Anthochlamys polygaloides.
spicata, *Spreng. Syst.* i. 815.—Madag.
spinescens, Russ. ex Wall. Cat. sub n. 6932 = Achyranthes aquatica.
splendens, *Schum. & Thonn. Beskr. Guin. Pl.* 140.—Afr. trop.
stolonifera, Leprieur, ex Moq. in DC. Prod. xiii. II. 365 = Telanthera maritima.
stricta, Fisch. ex Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. ix. 65 = brasiliensis.
stricta, *Hornem. Hort. Hafn. Suppl.* 28.—Bras.
tetragyna, Roxb. ex Moq. in DC. Prod. xiii. II. 237 = Deeringia tetragyna.
texana, *Scheele, in Linnaea*, xxii. (1849) 148.—Texas.
thyrsiflora, Wall. Cat. n. 6914 = Banalia thyrsiflora.
tomentosa, Willd. in Roem. & Schult. Syst. v. 531 = Chamissoa macrocarpa.
trigyna, *Linn. Mant.* 212.—Afr. trop.
trigyna, Wight, ex Steud. Nom. ed. II. i. 316 = Banalia thyrsiflora.
trigyna, Willd. ex Wall. Cat. n. 6915 = polygonoides.
triloba, *E. Mey. ex Meissn. in Hook. Lond. Journ. Bot.* ii. (1843) 448.—Afr. austr.
tumida, *Hort. Hall. ex Moq. in DC. Prod.* xiii. II. 245.—Hab.?
virgata, *Jacq. Coll.* ii. 279.—Am. trop.

CELSA, Vell. Fl. Flum. 173 (1825); iv. t. 93. *ZYGOPHYLLEAE.*
frutescens, *Vell. ll. cc.* 174.—Bras.

CELSIA, Heist. Syst. 5, 20 (1748) = **Ornithogalum**, Tourn. (Liliac.).

CELSIA, Linn. Syst. ed. I (1735). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 928.
DITOXIA, Rafin. ex Desv. Journ. Bot. iv. (1814) 270.
JANTHE, Griseb. Spicil. Fl. Rumel. ii. 40 (1844).
NEFFLEA, Spach, Hist. Vég. Phan. ix. 266 (1840).
THAPSANDRA, Griseb. Spicil. Fl. Rumel. ii. 40 (1844).
acaulis, *Bory & Chaub. Expéd. Mor. Bot.* 177. t. 18.—Graecia.
affinis, *A. Rich. Tent. Fl. Abyss.* ii. 110.—Abyssin.
agrimonifolia, C. Koch, in Linnaea, xxii. (1849) 732 = heterophylla.
alpina, *Boiss. & Kotschy, Diagn.* Ser. II. iii. 150.—Syria.
arbuscula, *A. Rich. Tent. Fl. Abyss.* ii. 110.—Abyssin.
arcturis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Arcturus.
Arcturus, *Jacq. Hort. Vindob.* ii. 53. t. 117.—Creta.
armeniaca, Boiss. Fl. Orient. iv. 361 = aurea.
Aucheri, *Boiss. Diagn.* Ser. I. xii. 29.—Persia.
aurea, *C. Koch, in Linnaea*, xxii. (1849) 731.—Armenia.
australis, Hort. Bonn. ex Flora, xvi. (1833) 125 = cretica.
Bernadesii, *G. Don, Gen. Syst.* iv. 499.—Hispan.; Marocco.
betonicaefolia, *Desf. Fl. Atlant.* ii. 58.—Afr. bor.
Boissieri, *Heldr. & Sart. ex Boiss. Fl. Orient.* iv. 353.—Graecia.
brachysepala, *Fisch. & Trautv. in Ind. Sem. Hort. Petrop.* iv. 33.—As. Min.
bugulifolia, *Jaub. & Spach, Illustr.* v. 7. t. 407.—Rumelia; As. Min.
caduca, Moench, Meth. 447 = orientalis.

CELSIA :—
Cavanillesii, Kunze, in Flora, xxix. (1846) 698 = cretica.
chrysurus, Fenzl, ex Boiss. Fl. Orient. iv. 359 = heterophylla.
cilicia, *Boiss. & Heldr. Diagn.* Ser. I. xii. 30.—Cilicia.
collina, Benth. in DC. Prod. x. 247 = Suwarowiana.
coromandelina, *Vahl, Symb. Bot.* iii. 79.—Ind. or.; Afghanist.
coronopifolia, *Boiss. & Bal. Diagn.* Ser. II. vi. 126.—Phrygia.
cretica, *Linn. f. Suppl.* 281.—Reg. Mediterr.
cyllenea, *Boiss. & Heldr. Diagn.* Ser. II. iii. 151.—Graecia.
Daenzeri, *Fauché & Chaub. in Expéd. Mor. Bot.* 342.—Graecia.
densiflora, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 208.—Afr. trop.
dentata, Moench, Meth. Suppl. 156 = coromandelina.
Doniana, *Walp. Rep.* iii. 150.—Hab.?
elata, *Fisch. ex Steud. Nom.* ed. I. 172.—Hab.?
erecta, '*Fisch. Hort. Gorenk.*' *ex Steud. Nom.* ed. II. i. 316.—Hab.?
floccosa, *Benth. in DC. Prod.* x. 245.—Abyssin.
glabrata, *Roth, ex Steud. Nom.* ed. II. i. 316.—Hab.?
glandulosa, *Bouché, in Verh. Ges. Naturf. Fr. Berl.* i. (1829) 395.—As. Min.
grandiflora, *Hort. ex Steud. Nom.* ed. II. i. 316.—Asia.
hetcrophylla, *Desf. ex Pers. Syn.* ii. 161.—Persia; As. Min.; Ind. or.
horizontalis, Moench, Meth. 448 = C. Arcturus.
incana, Benth. in DC. Prod. x. 247 = lanceolata.
incarnata ×, Reichb. ex Heynh. Nom. ii. 129, nomen.
interrupta, *Fresen. in Flora*, xvi. (1838) 605.—Abyssin.
intricata, *Benth. in DC. Prod.* x. 247.—Persia.
laciniata, Coss. ex Ball, in Journ. Linn. Soc. xvi. (1878) 585 = cretica.
laciniata, Poir. Encyc. Suppl. ii. 147 = Bernadesii.
lanata, Spreng. Syst. ii. 810, err. typ. = sublanata.
lanceolata, *Vent. Jard. Cels*, t. 27.—Mesopot.; Persia.
lepturus, Schott & Kotschy, in Oestr. Bot. Wochenbl. (1854) 186 = brachysepala.
linearis, Jacq. Coll. ii. 270 = Alonsoa linearis.
Luciliae, *Boiss. Diagn.* Ser. I. iv. 65.—As. Min.
lycia, *Boiss. Fl. Orient.* iv. 357.—Lycia.
lyrata, G. Don, Gen. Syst. iv. 499 = cretica.
maroccana, *Ball, in Journ. Bot.* xiii. (1875) 172.—Marocco.
orientalis, *Linn. Sp. Pl.* 621.—Graecia; As. Min.; Persia.
parviflora, *Decne. in Ann. Sc. Nat.* Sér. II. v. (1836) 254.—Arabia.
parviflora, G. Don, Gen. Syst. iv. 499 = Doniana.
pedunculosa, *Hochst. & Steud. ex Benth. in DC. Prod.* x. 245.—Abyssin.
persica, *C. A. Mey. Verz. Pfl. Cauc.* 111.—Reg. Cauc.; Persia.
pinetorum, *Boiss. Diagn.* Ser. I. xii. 30.—Cilicia.
pinnatifida, Boiss. & Reut. Diagn. Pl. Nov. Hisp. 22 = Bernadesii.
pontica, *Boiss. in Tchihat. Asie Min. Bot.* ii. 11.—As. Min.
purpurea, *Janka, in Természt. Füzetek*, iv. (1880) 296.—Hungar.
pyroliformis, *Boiss. & Heldr. Diagn.* Ser. I. xii. 28.—Lycaonia.
ramosissima, *Benth. in DC. Prod.* x. 244.—Marocco.
scrophulariaefolia, *Hochst. in A. Rich. Tent. Fl. Abyss.* ii. 112.—Abyssin.
sinuata, Cav. in Anal. Cienc. Nat. iii. (1801) 68 = cretica.
sinuata, *Colla, Hort. Ripul. App.* ii. 344; iii. 30.—Hab.?
speciosa, *Fenzl, in Friedric. Reise*, 280.—Graecia.
striatiflora, *Hort. ex Steud. Nom.* ed. II. i. 316.—Hab.?
sublanata, Jacq. Fragm. 79. t. 126 = C. Arcturus.
Suwarowiana, *C. Koch, in Linnaea*, xvii. (1844) 284.—Armenia; Persia.
tomentosa, *Hochst. ex Benth. in DC. Prod.* x. 245.—Abyssin.
tomentosa, Zucc. Pl. Nov. Fasc. ii. 22 = Verbascum graecum.
trapaefolia, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 88.—Lycia.

CELSIA :—
urticaefolia, Sims, Bot. Mag. t. 417 = Alonsoa incisifolia.
valerianaefolia, *A. Rich. Tent. Fl. Abyss.* ii. 112.—Abyssin.
viscosa, Roth, Catalect. ii. 69 = coromandeliana.
viscosa, Wight, ex Benth. in DC. Prod. x. 229 = Verbascum virgatum.

CELTIS, Tourn. ex Linn. Gen. ed. I. 337 (1737). *URTICACEAE*, Benth. & Hook. f. iii. 354.
MERTENSIA, H. B. & K. Nov. Gen. et Sp. ii. 30 (1817).
MOMISIA, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. v. 122 (1819).
SAUROBROMA, Rafin. Sylva Tellur. 32 (1838).
SOLENOSTIGMA, Endl. Prod. Fl. Norf. 41 (1833).
Acata, Buch.-Ham. in Trans. Linn. Soc. xvii. (1837) 211 = australis.
aculeata, *Sw. Prod. Veg. Ind. Occ.* 53.—Ind. occ.; Am. trop.
africana, *Burm. f. Fl. Cap. Prod.* 31.—Afr. austr.
alba, *Rafin. Fl. Ludov.* 25.—Am. bor.
albicans, Willd. ex Steud. Nom. ed. II. i. 316 = Trema micrantha.
alnifolia, *Miq. in Mart. Fl. Bras.* iv. I. 176.—Am. trop.
amblyphylla, *F. Muell. Fragm.* ix. 76.—Austral.
amboinensis, Buch.-Ham. ex Wall. Cat. sub n. 3693 = Trema politoria.
amboinensis, Willd. Sp. Pl. iv. 997 = Trema amboinensis.
americana, Hort. ex Planch. in DC. Prod. xvii. 176 = mississipicnsis.
americana, *Mill. Gard. Dict.* ed. VIII. n. 4.—Ind. occ.
amplifolia, *Noronha, in Verh. Batav. Gen.* v. (1790) cd. I. *Art.* IV. 11.—Malaya.
anfractuosa, *Liebm. in Vidensk. Selsk. Skr.* V. ii. (1851) 338.—Mexic.
angustifolia, Lindl. in Wall. Cat. n. 3691 = Trema angustifolia.
angustifolia, *Planch. in DC. Prod.* xvii. 186.—Sumatra.
appendiculata, E. Mey. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 341 = Chactachme aristata.
arcata, Buch.-Ham. ex Wall. Cat. sub n. 3695 = australis.
aristata, E. Mey. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 341 = Chaetachme aristata.
aspera, Brongn. in Duperr. Bot. Voy. Coq. 213. t. 48 = Trema aspera.
aspera, Lodd. ex G. Don, in Loud. Hort. Brit. 413 = australis.
asperula, *Miq. in Mart. Fl. Bras.* iv. I. 176.—Am. trop.
Audibertiana, Spach, in Ann. Sc. Nat. Sér. II. xvi. (1841) 41 = occidentalis.
australis, *Linn. Sp. Pl.* 1043.—Europ.; As. temp.; Ind. or.
australis, A. Rich. Tent. Fl. Abyss. 257 = Kraussiana.
azcurrensis, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 152.—Reg. Argent.
Balansae, *Planch. in DC. Prod.* xvii. 183.—N. Caled.
Berlandieri, *Klotzsch, in Linnaea*, xx. (1847) 541.—Mexic.
biflora, Ruiz, ex Miq. in Mart. Fl. Bras. iv. I. 182 = dichotoma.
boliviensis, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 311.—Bolivia.
Bonplandiana, *Planch. in DC. Prod.* xvii. 190.—Am. austr.
brasiliensis, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 310.—Bras.
brevifolia, *Miq. in Mart. Fl. Bras.* iv. I. 180.—Am. trop.
brevinervis, Blume, Mus. Bot. Lugd. Bat. ii. 67 = paniculata.
brevipes, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 297.—Arizona.
Bungeana, *Blume, Mus. Bot. Lugd. Bat.* ii. 71.—China.
Burmannii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 296 = Kraussiana.
canescens, H. B. & K. Nov. Gen. et Sp. ii. 28 = Trema micrantha.
canina, *Rafin. in Am. Monthly Mag.* ii. (1817) 43.—Am. bor.
caucasica, Hohen. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 285 = australis.
caucasica, Willd. Sp. Pl. iv. 994 = australis.
caudata, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 241 = Prunus pogonostyla.

CELTIS :—
caudata, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 294.—Mexic.
caudata, Wall. ex Planch. in DC. Prod. xvii. 199 = Trema amboinensis.
Chichilea, Ruiz & Pav. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 335 = Trema micrantha.
Chicope, Miq. in Mart. Fl. Bras. iv. 1. 181 = C. Tala.
chinensis, Bunge, Enum. Pl. Chin. Bor. 61 = Bungeana.
cinerea, Torr. Bot. Mex. Bound. t. 50 = C. Tala.
cinnamomea, *Lindl. in Wall. Cat.* n. 3696.—Ind. or.; Malaya.
Clausseniana, *Miq. in Mart. Fl. Bras.* iv. 1. 178.—Am. trop.
Commersonii, A. Brongn. in Duperr. Voy. Coq. Bot. 215 = Trema orientalis.
conferta, *Planch. in DC. Prod.* xvii. 183.—Madag.
cordata, Pers. Syn. i. 292 = occidentalis.
cordifolia, L'Hérit. ex Duham. Arb. ed. Nov. ii. 37 = occidentalis.
crassifolia, Lam. Encyc. iv. 138 = occidentalis.
crenata, Miq. in Mart. Fl. Bras. iv. 1. 181 = C. Tala.
crenata, *Planch. in DC. Prod.* xvii. 191.—Bolivia.
Curiandiuba, Gomez, ex Planch. l. c. 204 = Trema micrantha.
Davidiana, *Carr. in Rev. Hortic.* (1868) 300.—China.
dichotoma, *Ruiz, ex Miq. in Mart. Fl. Bras.* iv. 1. 182.—Peruv.
diffusa, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 314.—Bras.
discolor, A. Brongn. in Duperr. Voy. Coq. Bot. i. 215. t. 47. f. B = Trema discolor.
disticha, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 96 (*Contrib.* 44).—Paraguay.
Djungiel, *Planch. in DC. Prod.* xvii. 185.—Java.
Douglasii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 293 = reticulata.
dumosa, Casar. ex Planch. in DC. Prod. xvii. 188 = ferruginea.
dysodoxylon, Thw. Enum. Pl. Zeyl. 267 = cinnamomea.
Ehrenbergiana, Liebm. in Dansk. Vidensk. Selsk. Skrift. V. ii. (1851) 339 = aculeata.
elongata, Wall. Cat. n. 3692 C = Villebrunea integrifolia.
epiphylladena, Orteg. Hort. Matr. 79 = aculeata.
eriantha, *E. Mey. ex Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 296.—Afr. austr.
eriocarpa, Decne. in Jacquem. Voy. Bot. 150 = australis.
excelsa, Salisb. Prod. 175 = australis.
ferruginea, *Planch. in DC. Prod.* xvii. 188.—Bras.
ferruginea, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 424 = Zizyphus exserta.
flagellaris, Casar. ex Planch. in DC. Prod. xvii. 189 = brasiliensis.
flexuosa, *Miq. in Mart. Fl. Bras.* iv. 1. (1852-62) 181. —Am. trop.
flexuosa, Planch. in DC. Prod. xvii. 192 = bolivicnsis?
floridana, Rafin. New Fl. Am. iii. 37 = occidentalis?
fuscata, Rafin. l. c. 33 = mississipiensis?
Gardneri, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 311.—Bras.
glabra, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. *Art.* IV. 12.—Malaya.
glabra. Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 298 = tetrandra.
glabrata, Spreng. Syst. v. 150 = aculeata.
glabrata, Stev. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 285 = australis.
glomerata, Hochst. in Flora, xxviii. (1845) 87 (= *Sponia guineensis*).—Afr. trop.
glycycarpa, *Mart. ex Miq. in Mart. Fl. Bras.* iv. 1. 174.—Am. trop.
Goudotii, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 312.—N. Granat.
grandidentata, Tenore, Ind. Sem. Hort. Neap. (1833) 15 = mississipiensis?
guineensis, *Schum. & Thonn. Beskr. Guin. Pl.* 160.—Afr. trop.
hamata, *Blume, Mus. Bot. Lugd. Bat.* ii. 72.—Malaya.
Hamiltonii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 301 = tetrandra.
Harperi, *Horne, ex Baker, in Journ. Linn. Soc.* xx. (1883) 371.—Ins. Fiji.
Hasseltii, *Planch. in DC. Prod.* xvii. 185.—Java.
heterophyla, Rafin. New Fl. Am. iii. 37 = occidentalis?
Hilariana, *Planch. in DC. Prod.* xvii. 189.—Bras.
hypoleuca, *Planch. l. c.* 183.—N. Caled.
ingens, F. Muell. Fragm. iv. 88 = paniculata.

CELTIS :—
integrifolia, Lam. Encyc. iv. 140 = mississipiensis.
jamaicensis, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 290.—Jamaica.
japonica, *Planch. in DC. Prod.* xvii. 172.—Japon.
Kotschyana, Stev. in Bull. Soc. Nat. Mosc. xxx. (1857) 1. 392 = australis.
Kraussiana, *Bernh. in Flora*, xxviii. (1845) 87.—Afr. trop. et austr.
laeta, Salisb. Prod. 175 = orientalis.
laevigata, Spreng. Syst. i. 932 = aculeata.
laevigata, Willd. Enum. Hort. Berol. Suppl. 67 = mississipiensis.
Lamarckiana, Roem. & Schult. Syst. vi. 311 = Trema Lamarckiana.
lanceolata, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 152 (*Contrib.* 45).—Paraguay.
lancifolia, *Miq. in Mart. Fl. Bras.* iv. 1. 175.—Bras.
lancifolia, *Planch. in DC. Prod.* xvii. 192.—Bras.
latifolia, *Planch. l. c.* 186.—N. Guinea.
laurifolia, *Planch. l. c.* 185.—Java.
Lima, Blanco, Fl. Filip. ed. I. 197 = Trema Blancoi.
Lima, Lam. Encyc. iv. 40 = Trema Lamarckiana.
Lima, Sw. Prod. Veg. Ind. Occ. 53 = Trema Lima.
Lindheimeri, *Engelm. ex C. Koch, Dendrol.* ii. II. 434. —Am. bor.
littoralis, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 337 = caudata.
longifolia, Rafin. Atl. Journ. 177 = mississipiensis?
lutea, Pers. Syn. i. 292 = australis.
macrophylla, H. B. & K. Nov. Gen. et Sp. ii. 24 = Trema macrophylla.
madagascariensis, Boj. Hort. Maurit. 296 = Trema orientalis.
maritima, *Rafin. in Am. Monthly Mag.* ii. (1817) 44. —Am. bor.
mauritiana, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 307.—Ins. Maurit.
membranacea, *Miq. in Mart. Fl. Bras.* iv. 1. 175.—Bras.
micrantha, Sw. Prod. Veg. Ind. Occ. 53 = Trema micrantha.
microcarpa, Salzm. ex Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 333 = Trema micrantha.
microphylla, Zipp. ex Blume, Mus. Bot. Lugd. Bat. ii. 59 = Trema timorensis.
mississippiensis, *Bosc, Encyc. Agric.* vii. 577.—Am. bor.
mollis, Humb. & Bonpl. ex Willd. Sp. Pl. iv. 996 = Trema micrantha.
mollis, Wall. Cat. n. 7203 = tetrandra.
monoica, *Hemsl. Biol. Centr. Am. Bot.* iii. 139.—Mexic.
montana, Jungh. ex Planch. in DC. Prod. xvii. 193 = Parasponia parviflora.
morifolia, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 311.—Bras.
morifolia, Rafin. New Fl. Am. iii. 34 = occidentalis.
Mukii, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. III. (1846) 223.—Japon.
napalensis, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 298 = tetrandra.
obliqua, Moench, Meth. 344 = occidentalis.
occidentalis, *Linn. Sp. Pl.* 1044.—Am. bor.
opaca, C. Moore, ex Benth. Fl. Austral. vi. 157 = paniculata.
Opegrapha, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 294 = Kraussiana.
orientalis, Hort. ex Steud. Nom. ed. II. i. 316 = occidentalis.
orientalis, Lam. Encyc. iv. 138 = Trema Commersonii.
orientalis, Linn. Sp. Pl. 1044 = Trema orientalis.
orientalis, Mill. Gard. Dict. ed. VIII. n. 3 = australis.
orientalis, Thunb. Fl. Jap. 114 = japonica.
orthacanthos, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 309 = aculeata.
pacifica, *Planch. l. c.* 308.—Ins. Marquesas.
pallida, Torr. in Bot. Mex. Bound. 203 = C. Tala.
paniculata, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 305.—Austral.; Ins. Pacific.
parvifolia, Rafin. New Fl. Am. iii. 36 = occidentalis.
parvifolia, *A. Rich. Fl. Cub. Fanerog.* ii. 219.—Cuba.
patula, Rafin. New Fl. Am. iii. 37 = reticulata?
Pavonii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 313 = dichotoma.
philippinensis, *Blanco, Fl. Filip.* ed. I. 197.—Ins. Philipp.; Austral.

CELTIS :—
politoria, Wall. Cat. n. 3693 = Trema politoria.
procera, Salisb. Prod. 175 = occidentalis.
pubescens, *H. B. & K. Nov. Gen. et Sp.* ii. 32.—N. Granat.
pumila, Pursh, Fl. Am. Sept. i. 200 = occidentalis.
reticulata, Hook. f. & Thoms. ex Planch. in DC. Prod. xvii. 209 = Gironniera reticulata.
reticulata, *Torr. in Ann. Lyc. N. York*, i. (1824) 247.—Am. bor. occ.
reticulosa, Miq. Pl. Jungh. 69 = cinnamomea.
rhamnoides, DC. ex Planch. in DC. Prod. xvii. 189 = brasiliensis.
rhamnoides, Willd. Sp. Pl. iv. 998 = aculeata.
rigescens, *Planch. in DC. Prod.* xvii. 182.—Sumatra.
rigida, Blume, Bijdr. 486 = Trema rigida.
rigida, *Parodi, in Anal. Soc. Cient. Argent.* v. (1878) 153 (*Contrib.* 45).—Paraguay.
riparia, H. B. & K. Nov. Gen. et Sp. ii. 28 = Trema riparia.
riparia, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 337 = caudata.
rostrata, Zipp. ex Span. in Linnaea, xv. (1841) 343 (= *Sponia rostrata*).—Hab.?
Roxburghii, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 302 = tetrandra.
rufescens, Banks, ex Planch. l. c. 334 = Trema micrantha.
rugosa, Willd. Sp. Pl. iv. 996 = Trema rugosa.
salicifolia, Rafin. New Fl. Am. iii. 34 = mississipiensis?
scabra, Casar. ex Planch. in DC. Prod. xvii. 204 = Trema carinata.
Schiedeana, Schlecht. in Linnaea, vii. (1832) 140 = Trema micrantha.
Selloviana, *Miq. in Mart. Fl. Bras.* iv. 1. 179.—Bras.
serotina, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 301 = tetrandra.
Shortii, *Planch. l. c.* 292.—Am. bor.
sinensis, Planch. l. c. 286 = Bungeana.
sinensis, *Pers. Syn.* i. 292.—China.
sinensis, Willd. Enum. Hort. Berol. Suppl. 67 = japonica.
spinosa, Ruiz, ex Miq. in Mart. Fl. Bras. iv. 1. 182 = dichotoma.
spinosa, *Spreng. Syst.* i. 932 (sp. dub.).—Bras.
spinosissima, *Miq. in Mart. Fl. Bras.* iv. 1. 176.—Bras.
strychnoides, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 306 = philippinensis.
subdentata, E. Mey. ex Planch. l. c. 341 = Chaetachme aristata.
sumatrana, *Planch. in DC. Prod.* xvii. 181.—Sumatra.
Swartzii, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 304.—Ins. S. Doming.
Tala, *Gill. ex Planch. l. c.* 310.—Am. austr.
tenuifolia, Nutt. Gen. Am. i. 202 = pumila?
tenuifolia, Rafin. New Fl. Am. iii. 36 = occidentalis?
tetrandra, *Roxb. Hort. Beng.* 21; *Fl. Ind.* ii. 63.—Ind. or.; Malaya.
tetrandra, Wall. Cat. n. 3695 F = Villebrunea integrifolia.
texana, Scheele, in Linnaea, xxii. (1849) 146 = Berlandieri.
timorensis, *Span. l. c.* xv. (1841) 343.—Ins. Timor.
tomentosa, Roxb. Hort. Beng. 86; Fl. Ind. ii. 66 = Trema amboinensis.
Tournefortii, Lam. Encyc. iv. 138 = australis.
triflora, *Ruiz, ex Miq. in Mart. Fl. Bras.* iv. 1. 181.—Peruv.
trinervia, Lam. Encyc. iv. 140 (= *Sponia trinervia*).—Ins. S. Doming.
trinervia, Roxb. Hort. Beng. 86; Fl. Ind. ii. 65 = tetrandra.
urticifolia, Rafin. New Fl. Am. iii. 32 = occidentalis.
velutina, *Planch. in Ann. Sc. Nat.* Ser. III. x. (1848) 313.—Peruv.
vesiculosa, Hochst. ex Planch. l. c. 295 = Kraussiana.
virgata, Roxb. in Wall. Cat. n. 3694 = Trema timorensis.
Waitzii, Blume, Mus. Bot. Lugd. Bat. ii. 71 = cinnamomea.
Wallichii, Steud. Nom. ed. II. i. 316 = tetrandra.
Wightii, *Planch. in Ann. Sc. Nat.* Sér. III. x. (1848) 307.—Ind. or.; Malaya; Austral.
Willdenowiana, Roem. & Schult. Syst. vi. 306 = japonica.
Zippelii, *Planch. in DC. Prod.* xvii. 185.—N. Guinea.
zizyphoides, *Spreng. Syst.* i. 932.—N. Granat.

CEMBRA, Opiz, Seznam, 27 (1852) = **Pinus**, Tourn. (Conif.).
montana, Opiz, l. c. = P. Cembra.

CENARIUM, Linn. Amoen. Acad. iv. 121, in nota (1759) = **Canarium**, Rumph. (Burser.).

CENARRHENES, Labill. Nov. Holl. Pl. i. 36. t. 50 (1804). *PROTEACEAE*, Benth. & Hook. f. iii. 176.
nitida, *Labill. l. c.*—Austral.
paniculata, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* xii. (1865) 41.—N. Caled.
spathulaefolia, *Brongn. & Gris, l. c.*—N. Caled.
spathulata, Panch. & Sebert, Not. Bois Nouv. Caléd. 180 = spathulaefolia.

CENCHRUS, Linn. Coroll. Gen. 20 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1105.
ECHISACHYS, Neck. Elem. iii. 228 (1790).
RARAM, Adans. Fam. ii. 35 (1763).
aegyptius, Beauv. Agrost. 157 = Eleusine aegyptiaca.
agrimonoides, *Trin. Diss. Gram. Panic.* 72.—Ins. Societ.
alopecuroides, *J. & C. Presl, Rel. Haenk.* i. 317.—Am. austr.
alopecuroides, Thunb. Prod. Pl. Cap. 24 = Pennisetum Linnaei.
annularis, *Anderss. in Peters, Reise Mossamb. Bot.* 553.—Afr. trop.
anomoplexis, Labill. Sert. Austr. Caled. 14. t. 19 = calyculatus.
asperifolius, Desf. Fl. Atlant. ii. 388 = Pennisetum asperifolium.
australis, *R. Br. Prod.* i. 196.—Austral.
barbatus, *Schum. in Schum. & Thonn. Beskr. Guin. Pl.* 43.—Afr. trop.
biflorus, *Roxb. Hort. Beng.* [81]; *Fl. Ind.* i. 233.—Ind. or.
brevisetus, *Fourn. Mex. Pl. Enum. Gram.* 50; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 506.—Mexic.
Brownei, Roem. & Schult. Syst. ii. 258 = inflexus.
bulbifer, Hochst. ex Boiss. Fl. Orient. v. 448 = montanus.
bulbosus, Fresen. ex Steud. Nom. ed. II. i. 317 = pennisetiformis.
calyculatus, *Cav. Ic.* v. 39. t. 463.—Polynes.
capitatus, Linn. Sp. Pl. 1049 = Echinaria capitata.
carolinianus, Walt. Fl. Carol. 79 = tribuloides.
catharticus, *Delile, Cat. Hort. Monsp.* (1838).—*Cf. Linnaea*, xiii. (1839) *Litt.* 103.—As. et Afr. trop.
Cavanillesii, *Tausch, in Flora*, xx. (1837) 97.—Bras.
ciliaris, *Fig. & De Not. in Mem. Acc. Torin.* (1854) 383.—Nubia?
ciliaris, Linn. Mant. 302 = Pennisetum cenchroides.
dactylolepis, Steud. Syn. Pl. Gram. 109 = viridis.
digynus, Ehrenb. ex Boiss. Fl. Orient. v. 449 = Pennisetum cenchroides.
distichophyllus, *Griseb. Cat. Pl. Cub.* 234.—Cuba.
echinatus, Cav. Ic. v. 39. t. 462 = Cavanillesii.
echinatus, *Linn. Sp. Pl.* 1050.—Cosmop. trop.
echinatus, A. Rich. Tent. Fl. Abyss. ii. 389 = catharticus.
echinatus, Steud. ex Doell, in Mart. Fl. Bras. ii. II. 309 = viridis.
echinatus, Torr. Fl. N. Am. i. 68 = tribuloides.
echinoides, Wight, ex Steud. Nom. ed. II. i. 317 = Pennisetum cenchroides.
Elliotii, *Kunth, Rev. Gram.* i. 51.—Am. bor.
elymoides, *F. Muell. Fragm.* viii. 107.—Austral.
frutescens, *Linn. Sp. Pl.* 1050.—'Am.'
frutescens, *Sieber, ex Steud. Nom.* ed. II. i. 317, nomen.—Armenia.
fusiformis, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 170 = agrimonoides.
geniculatus, Thunb. Prod. Pl. Cap. 24 = Pennisetum Thunbergii.
gracilis, *Beauv. Agrost.* 157, nomen.—Hab.?
granularis, *Anderss. Enum. Pl. Ins. Galap.* 140.—Ins. Galapag.
granularis, Linn. Mant. ii. App. 575 = Manisuris granularis.
heterochaetus, *Steud. Syn. Pl. Gram.* 110.—Hab.?
hexaflorus, Blanco, Fl. Filip. ed. I. 36 = Pennisetum nigricans.
hirsutus, Spreng. Neue Entdeck. iii. 15 = Scleria hirtella.
hordeiformis, Rottl. ex Steud. Syn. Pl. Gram. 102 = Pennisetum Alopecuros.

CENCHRUS:—
hordeiformis, Thunb. Prod. Pl. Cap. 24 = Pennisetum compressum.
Hystrix, *Fig. & De Not. in Mem. Acc. Torin.* (1854) 382.—Nubia.
incertus, *M. A. Curt. in Bost. Journ. Nat. Hist.* i. (1837) 135.—Am. bor.
inflexus, Poir. Encyc. vi. 50 = Echinolaena hirta.
inflexus, *R. Br. Prod.* i. 195.—Austral.
laevigatus, Trin. Fund. Agrost. 172 = Anthephora elegans.
laniflorus, *Steud. Syn. Pl. Gram.* 110.—Ins. Tahiti.
lappaceus, Linn. Sp. Pl. ed. II. 1488 = Centotheca lappacea.
lappaceus, *Tausch, in Flora*, xx. (1837) 97.—Hab.?
Lechleri, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
limensis, *Meyen, Reise*, ii. 71.—Peruv.
linearis, Lam. Fl. Fr. iii. 631 = Tragus racemosus.
longifolius, *Hochst. ex Steud. Syn. Pl. Gram.* 109.—Nubia.
macrocarpus, Ledeb. ex Steud. Nom. ed. II. i. 317 = echinatus.
macrostachyus, *Hochst. ex Steud. Syn. Pl. Gram.* 109.—Abyssin.
marginalis, Rudge, Pl. Guian. 19. t. 25 = Echinolaena hirta.
mitis, *Anderss. in Peters, Reise Mossamb. Bot.* 553.—Afr. trop.
montanus, *Nees, in Royle, Illustr. Bot. Himal.* 416.—As. et Afr. trop.
mucronatus, Pers. ex Steud. Nom. ed. I. 172 = muricatus.
multiflorus, *J. & C. Presl, Rel. Haenk.* i. 318.—Mexic.
muricatus, Linn. Mant. 302 = Trachys mucronata.
muricatus, *Phil. Sert. Mend. Alt.* 44.—Chili.
myosuroides, *H. B. & K. Nov. Gen. et Sp.* i. 115. t. 35.—Am. trop.
niloticus, Fig. & De Not. in Mem. Acc. Torin. (1854) 380 = catharticus.
orientalis, Willd. ex Kunth, Enum. Pl. i. 162 = Pennisetum orientale.
ovatus, *Lam. ex Poir. Encyc.* vi. 51.—Afr. austr.
pallidus, *Fourn. Mex. Pl. Enum. Gram.* 50; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 509.—Mexic.
parviflorus, Poir. Encyc. vi. 52 = Panicum Ventenatii.
pauciflorus, Benth. Bot. Voy. Sulph. 56 = tribuloides.
pennisetiformis, *Hochst. & Steud. ex Steud. Nom.* ed. II. i. 317.—Arabia.
pilosus, *H. B. & K. Nov. Gen. et Sp.* i. 116. t. 36.—Am. austr.
platyacanthus, *Anderss. Enum. Pl. Ins. Galap.* 139.—Ins. Galapag.
pubescens, *Steud. Nom.* ed. II. i. 317.—Afr. austr.
pungens, *H. B. & K. Nov. Gen. et Sp.* i. 115.—Mexic.
purpurascens, Thunb. in Trans. Linn. Soc. ii. (1794) 329 = Gynothrix japonica.
pycnostachyus, *Steud. Syn. Pl. Gram.* 109.—Afr. trop.
racemosus, Linn. Sp. Pl. 1049 = Tragus racemosus.
ramosissimus, Poir. Encyc. vi. 51 = Pennisetum dichotomum.
rigidifolius, *Fig. & De Not. in Mem. Acc. Torin.* (1854) 386.—Nubia.
rigidus, *Willd. ex Doell, in Mart. Fl. Bras.* ii. II. 310, in obs.—Bras.
roseus, *Fourn. Mex. Pl. Enum. Gram.* 50; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 507.—Mexic.
rufescens, Desf. Fl. Atlant. ii. 388 = Pennisetum cenchroides.
Schimperi, Hochst. & Steud. ex Steud. Nom. ed. II. i. 317 = montanus.
setigerus, Steud. l. c. = Pennisetum Vahlii.
setigerus, Vahl, Enum. ii. 395 = agrimonoides.
setosus, Sw. Prod. Veg. Ind. Occ. 26 = Pennisetum setosum.
spicatus, Cav. Desc. 304, 614 = Pennisetum typhoideum.
spinifex, Cav. Ic. v. 38. t. 461 = tribuloides.
strictus, Chapm. in Coult. Bot. Gaz. iii. (1878) 20 = incertus.
taitensis, *Steud. Syn. Pl. Gram.* 419.—Ins. Tahiti.
tomentosus, *Poir. Encyc.* vi. 51.—Afr. austr.
tribuloides, *Linn. Sp. Pl.* 1050.—Am. bor.
tripsaceus, Herb. Linn. ex Munro, in Journ. Linn. Soc. vi. (1862) 55 = Trachys mucronata.
tripsacoides, R. Br. in Salt, Abyss. App. 62 = montanus.

CENCHRUS:—
uniflorus, Ehrenb. ex Boiss. Fl. Orient. v. 448 = montanus.
vaginatus, *Steud. Syn. Pl. Gram.* 110.—Hab.?
villosus, Spreng. Syst. i. 301 = Anthephora elegans.
viridis, *Spreng. l. c.*—Am. trop.

ČENĚKIA, Opiz, Seznam, 36 (1852) = **Campanula**, Tourn.
rapunculoides, Opiz, l. c. = Camp. rapunculoides.
simplex, Opiz, l. c. = Camp. bononiensis.

CENIA, Comm. ex Juss. Gen. 183 (1789). *COMPOSITAE*, Benth. & Hook. f. ii. 429.
LANCISIA, Gaertn. Fruct. ii. 422 (1791).
anthemoidea, DC. Prod. vi. 83 = turbinata.
barbata, *Benth. & Hook. f. Gen.* ii. 430.—Afr. austr.
debilis, DC. Prod. vi. 83 = turbinata.
discoidea, Less. Syn. Comp. 261 = Cotula nudicaulis.
discolor, DC. Prod. vi. 82 = sericea.
flosculosa, DC. l. c. 83 = microglossa.
geminata, *Kunze, Del. Sem. Hort. Lips.* (1846) 1.—Afr. austr.
hispida, *Benth. & Hook. f. Gen.* ii. 430.—Afr. austr.
microcephala, *E. Mey. ex DC. Prod.* vi. 79.—Afr. austr.
microglossa, *DC. l. c.* 83.—Afr. austr.
nudicaulis, E. Mey. ex Heynh. Nom. ii. 130 = Cotula nudicaulis.
pectinata, *DC. Prod.* vi. 83.—Afr. austr.
pruinosa, DC. l. c. 82 = turbinata.
sericea, *DC. l. c.*—Afr. austr.
subheterocarpa, Less. Syn. Comp. 261 = turbinata.
Thunbergii, *Benth. & Hook. f. Gen.* ii. 430.—Afr. austr.
turbinata, *Pers. Syn.* ii. 465.—Afr. austr.
villosa, Heynh. Nom. ii. 130 = Cotula villosa.

CENOCLINE, C. Koch, in Bot. Zeit. i. (1843) 41 = **Cotula**, Tourn. (Compos.).
acutiloba, C. Koch, l. c. = Matricaria acutiloba.
albida, C. Koch, l. c. = Matricaria albida.
aurea, C. Koch, l. c. = Matricaria aurea.
chamaemelifolia, C. Koch, l. c. = Cotula chamaemelifolia.
cinerea, C. Koch, l. c. = Cotula cinerea.
globifera, C. Koch, l. c. = Matricaria globifera.
grandiflora, C. Koch, l. c. = Matricaria grandiflora.
pauciflora, C. Koch, l. c. = Matricaria discoidea.

CENOLOPHIUM, Koch, Gen. Umbell. 103 (1824) = **Selinum**, Linn. (Umbellif.).
divaricatum, Bess. in Flora, xv. (1832) II. Beibl. 27 = *Fischeri*.
Fischeri, Koch, Gen. Umbell. 103.—Rossia; Reg. Caucas.; As. bor.
lapponicum, Nyl. ex Nyl. & Sael. Herb. Mus. Fenn. 33 = *Fischeri*.

CENOLOPHON, Blume, Enum. Pl. Jav. 60 (1830) = **Alpinia**, Linn. (Scitamin.).
rubrum, Blume, l. c. = A. rubra.
vitellinum, Horan. Prod. Scitam. 36 = Amomum vitellinum.

CENOSTIGMA, Tul. in Ann. Sc. Nat. Sér. II. xx. 140. t. 3 (1843). *LEGUMINOSAE*, Benth. & Hook. f. i. 564.
angustifolium, Tul. l. c. 141 = Gardnerianum.
Gardnerianum, *Tul. l. c.*—Bras.
macrophyllum, *Tul. l. c.*—Bras.

CENTAUREA, Linn. Gen. ed. I. 263 (1737) (Centauria). *COMPOSITAE*, Benth. & Hook. f. ii. 477.
ACROCENTRON, Cass. in Dict. Sc. Nat. xliv. 37 (1826).
ACROLOPHUS, Cass. l. c. l. 253 (1827).
ACROPTILON, Cass. l. c. 464 (1827).
AEGIALOPHILA, Boiss. & Heldr. Diagn. Ser. I. x. 105 (1849).
AETHEOPAPPUS, Cass. in Dict. Sc. Nat. l. 250 (1827).
ALOPHIUM, Cass. l. c. liv. 493 (1828).
AMBLYOPOGON, Fisch. & Mey. ex DC. Prod. vi. 561 (1837).
ANTAUREA, Neck. Elem. i. 70 (1790).
BIELZIA, Schur, Enum. Pl. Transs. 409 (1866).

CENTAUREA :—

Calcitrapa, Hall. Enum. Stirp. Helv. ii. 689 (1742).
Callicephalus, C. A. Mey. Verz. Pfl. Cauc. 66 (1831).
Centaurium, Tourn. ex Linn. Syst. ed. I (1735).
Cestrinus, Cass. Bull. Soc. Philom. 33 (1817); et in Dict. Sc. Nat. viii. 24 (1817).
Chartolepis, Cass. Dict. Sc. Nat. xliv. 36 (1826).
Cheirolepis, Boiss. Diagn. Ser. I. x. 106 (1849).
Cheirolophus, Cass. in Dict. Sc. Nat. l. 250 (1827).
Chryseis, Cass. l. c. ix. 154 (1817).
Colymbada, Hill, Veg. Syst. iv. 31 (1762).
Crepula, Hill. l. c. 9 (1762).
Crocodilium, [Vaill.] Juss. Gen. 173 (1789).
Cyanus, [Tourn.] Linn. Syst. ed. I (1735); Juss. Gen. 174 (1789).
Eriopha, Hill, Hort. Kew. 69 (1769).
Fornicium, Cass. in Bull. Soc. Philom. 93 (1819); et in Dict. Sc. Nat. xvii. 249 (1820).
Halocharis, Bieb. ex DC. Prod. vi. 665 (1837).
Heraclea, Hill, Veg. Syst. iv. 37 (1762).
Heterolophus, Cass. in Dict. Sc. Nat. l. 250 (1827).
Hippophaestum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 443 (1821).
Hookia, Neck. Elem. i. 70 (1790).
Hyalea, Jaub. & Spach, Illustr. iii. 19. t. 214, 217, 292 (1847).
Hymenocentron, Cass. in Dict. Sc. Nat. xliv. 37 (1826).
Hymenocephalus, Jaub. & Spach, Illustr. iii. 12. t. 209 (1847).
Jacea, [Tourn.] Linn. Syst. ed. I (1735); Juss. Gen. 173 (1789).
Lepteranthus, Neck. Elem. i. 73 (1790).
Leucantha, S. F. Gray, Nat. Arr. Brit. Pl. ii. 443 (1821).
Lopholoma, Cass. in Dict. Sc. Nat. xliv. 37 (1826).
Malacocephalus, Tausch, in Flora, xi. (1828) ii. 481.
Mantisalca, Cass. in Bull. Soc. Philom. (1818) 142.
Melanoloma, Cass. in Dict. Sc. Nat. xxix. 472 (1823).
Menomphalus, Pomel, Nouv. Mat. Fl. Atl. 32 (1874).
Mesocentron, Cass. in Dict. Sc. Nat. xliv. 38 (1826).
Microlonchus, Cass. l. c. (1826); l. 247 (1827).
Microlophus, Cass. l. c. xliv. 37 (1826).
Odontolophus, Cass. l. c. l. 252 (1827).
Pachycentron, Pomel, Nouv. Mat. Fl. Atl. 32 (1874).
Pectinastrum, Cass. in Dict. Sc. Nat. xliv. 38 (1826).
Phaeopappus, Boiss. Diagn. Ser. I. vi. 122 (1845).
Phalolepis, Cass. in Dict. Sc. Nat. l. 248 (1827).
Phillostizus, Cass. l. c. xxxix. 498 (1826).
Phrygia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 441 (1821).
Piptoceras, Cass. in Dict. Sc. Nat. l. 469 (1827).
Platylophus, Cass. l. c. xliv. 36 (1826).
Plectocephalus, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. t. 51 (1838).
Psora, Hill, Veg. Syst. iv. 30 (1762).
Polyacantha, S. F. Gray, Nat. Arr. Brit. Pl. ii. 443 (1821).
Pterolophus, Cass. in Dict. Sc. Nat. xliv. 34 (1826).
Ptosimopappus, Boiss. Voy. Bot. Espagne, 739 (1839–45).
Psephellus, Cass. in Dict. Sc. Nat. xliii. 488 (1826).
Podia, Neck. Elem. i. 72 (1790).
Pycnocomus, Hill, Syst. Veg. iv. 38 (1762).
Rhapontica, Hill. l. c. 47 (1762).
Rhaponticum, [Vaill.] Adans. Fam. ii. 117 (1763); Juss. Gen. 174 (1789).
Seridia, Juss. Gen. 173 (1789).
Setachna, Dulac, Fl. Hautes-Pyr. 518 (1867).
Sagmen, Hill, Veg. Syst. iv. 35 (1762).
Spilacron, Cass. in Dict. Sc. Nat. l. 238 (1827).
Solstitiaria, Hill, Veg. Syst. iv. 21 (1762).
Sphaerocephala, Hill, l. c. 48 (1762).
Stenolophus, Cass. in Dict. Sc. Nat. l. 499 (1827).

CENTAUREA :—

Stizolophus, Cass. in Dict. Sc. Nat. xliv. 35 (1826), nomen; li. 49 (1827).
Staebe, Hill, Veg. Syst. iv. 33 (1762).
Tetramorphaea, DC. in Guill. Arch. Bot. ii. (1833) 331.
Tomanthea, DC. Prod. vi. 564 (1837).
Triplocentron, Cass. in Dict. Sc. Nat. xliv. 38 (1826).
Veltis, Adans. Fam. ii. 116 (1763).
Verutina, Cass. in Dict. Sc. Nat. xliv. 38 (1826).
Xanthopsis, C. Koch, in Linnaea, xxiv. (1851) 422.

abrotanifolia, Lam. Encyc. i. 670 = paniculata.
abyssinica, *Sch. Bip. ex Boiss. Diagn.* Ser. II. iii. 62. —Abyssin.
acanthodes, *Spreng. Syst.* iii. 403.—Hab.?
acaulis, Cav. ex Willk. & Lange, Prod. Fl. Hisp. ii. 150 = Cavanillesiana.
acaulis, Forsk. Fl. Aegypt. Arab. 52 = glomerata.
acaulis, Haensel. ex Willk. & Lange, Prod. Fl. Hisp. ii. 151 = Haenseleri.
acaulis, Lag. l. c. = Graelsii.
acaulis, *Linn. Sp. Pl.* 914.—Afr. bor.
achaia, *Boiss. & Heldr. Diagn.* Ser. II. iii. 79.—Graecia.
acicularis, *Sibth. & Sm. Fl. Graec.* t. 911.—Archip. Graec.
acmophylla, Boiss. Diagn. Ser. II. iii. 68 = axillaris.
acrolepis, *Hoffmgg. ex Heynh. Nom.* ii. 131, nomen.—Hab.?
acromaura, *Hoffmgg. ex Heynh. l. c.* nomen.—Hab.?
acuta, Lam. Fl. Fr. ii. 49 = Crupina vulgaris.
acuta, Vahl, ex Bieb. Casp. 82 = C. Behen.
acutangula, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 68. —Hispan.
acutifolia, Jord. Pugill. 105 = pectinata.
acutifolia, Schur, Enum. Pl. Transs. 899, Index = plumosa.
acutiloba, DC. Prod. vi. 581 = cristata.
Adami, Willd. Sp. Pl. iii. 2310 = solstitialis.
adonidifolia, Reichb. in Moessl. Handb. ed. II. ii. 1545 = rupestris.
adpressa, Ledeb. ex Steud. Nom. ed. II. i. 316 = stereophylla.
adulterina, Moretti, ex DC. Prod. vi. 596 = C. Calcitrapa.
aegialophila, Boiss. & Heldr. ex Boiss. Fl. Orient. iii. 704 = cretica.
aegyptiaca, *Linn. Mant.* i. 118.—Aegypt.; Arab.
aegyptiaca, Sibth. & Sm. Fl. Graec. t. 907 = iberica.
aeolica, Guss. ex DC. Prod. vi. 584 = aplolepis.
affinis, Frivald. in Flora, xix. (1836) 435 = dissecta.
africana, *Lam. Encyc.* i. 664.—Afr. bor.
aggregata, *Fisch. & Mey. in DC. Prod.* vi. 585.—As. Min.; Persia.
Aichingeriana, Welw. ex Reichb. Fl. Germ. Excurs. 848 = variegata.
ainetensis, Boiss. Diagn. Ser. I. x. 117 = eryngioides.
alata, Lam. Encyc. i. 665 = C. Behen.
alba, Bieb. Fl. Taur. Cauc. ii. 353 = sterilis.
alba, Herb. Calc. ex C. B. Clarke, Comp. Ind. 239 = Tricholepis elongata.
alba, *Linn. Sp. Pl.* 914.—Eur. austr.; Reg. Caucas.
alba, Suter, Fl. Helv. ii. 204 = amara.
albida, C. Koch, in Linnaea, xxiv. (1851) 433 = aggregata.
albiflora, C. Koch, l. c. 430 = carduiformis.
alexandrina, *Delile, Fl. Egypte*, 280. t. 49. f. 3.—Aegypt.
algeriensis, *Dur. & Coss. Notes Pl. Crit.* 136.—Algeria.
Alophium, *DC. Prod.* vi. 601.—Hispan.
alpestris, Hegetschw. Fl. Schw. 854 = C. Scabiosa.
alpina, *Linn. Sp. Pl.* 910.—Europ.
altaica, Fisch. ex Spreng. Pugill. i. 59 = Leuzea salina.
amaena, Boiss. & Bal. Diagn. Ser. II. vi. 112 = amoena.
amara, Habl. Taur. 155 = trinervia.
amara, *Linn. Sp. Pl.* ed. II. 1292.—Europ. austr.; As. Min.
Amberboi, Mill. Gard. Dict. ed. VIII. n. 9 = moschata.
ambigua, *Guss. Ind. Sem. Hort. Boccad.* (1826) 3; *ex DC. Prod.* vi. 582.—Italia.
ambigua, Thom. ex DC. l. c. 573 = nervosa.
amblensis, *Graells, in Mem. Acad. Cienc. Madr.* ii. (1859) 462.—Hispan.
amblyolepis, *Ledeb. Fl. Ross.* ii. 703.—Reg. Caucas.

CENTAUREA :—

ambracea, Sehkuhr, Handb. t. 261 = moschata.
americana, *Nutt. in Journ. Acad. Philad.* (1821) 117. —Am. bor.
americana, Spreng. Syst. iii. 407 = melitensis.
ammocyanus, *Boiss. Diagn.* Ser. I. x. 109.—Arab.; Palaest.
ammophila, Bess. ex DC. Prod. vi. 586 = diffusa.
amoena, *Boiss. & Bal. Diagn.* Ser. II. vi. 112.—Cappadoc.
Amoi, Campo, ex Campo & Amo, in Revista, v. (1855) 56 = Boissieri.
amourensis, *Pomel, Nouv. Mat. Fl. Atl.* 30.—Algeria.
amplexicaulis, S. G. Gmel. It. i. 136. t. 24 = Jurinea polyclonos.
amplifolia, *Boiss. & Heldr. Diagn.* Ser. II. iii. 68.—Graecia.
anatolica, Griseb. Spicil. Fl. Rumel. ii. 234 = depressa.
angulosa, *Pomel, Nouv. Mat. Fl. Atl.* 26.—Algeria.
angustifolia, Mill. Gard. Dict. ed. VIII. n. 7 = montana.
angustifolia, Schrank, Baier. Fl. ii. 376 = amara.
antennata, *Dufour, in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 158.—Hispan.
antiochia, *Boiss. Diagn.* Ser. I. x. 115.—Syria.
aphrodisea, *Boiss. l. c.* iv. 17.—Caria.
apiculata, Ledeb. Ind. Hort. Dorp. Suppl. (1824) 3 = C. Scabiosa.
aplolepis, *Moretti, in Brugnat. Giorn. Fis.* II. ix. (1826) 154.—Italia.
appendicigera, C. Koch, in Linnaea, xxiv. (1851) 425 = Balansae.
Apula, Lam. Encyc. i. 674 = melitensis.
arachnoidea, Viv. Ann. Bot. i. ii. 185 = rupestris.
arancosa, *Boiss. Diagn.* Ser. I. x. 121.—Syria.
arborea, *Webb & Berth. Phyt. Canar.* ii. 356.—Ins. Canar.
arenaria, *Bieb. Fl. Taur. Cauc.* ii. 347.—Europ. or.; Reg. Caucas.
arenaria, Reichb. Pl. Crit. v. 26. t. 443. f. 634 = Reichenbachii.
arenaria, Urv. in Mém. Soc. Linn. Par. i. (1822) 371 = polyclada.
argentea, Frivald. ex Nym. Consp. 429 = rupestris.
argentea, *Linn. Sp. Pl.* 912.—Ins. Creta et Istria.
argophylla, Steud. Nom. ed. I. 173 = argyrophylla.
arguta, *Nees, Hor. Phys. Berol.* 116. t. 25.—Ins. Canar.
argyrophylla, *Willd. Enum. Hort. Berol. Suppl.* 61.—Hab.?
arifolia, *Boiss. Diagn.* Ser. I. x. 112.—Syria.
aristata, *Benth. & Hook. f. Gen.* ii. 480.—Cappadoc.
aristata, Hoffmgg. & Link, Fl. Port. ii. 226 = caerulescens.
Armena, *Boiss. Fl. Orient.* iii. 676.—As. Min.
armoraeifolia, *Sibth. & Sm. Fl. Graec. Prod.* ii. 205.—Graecia.
arrectispina, Bertol. Fl. Ital. ix. 487 = C. Calcitrapa.
aspera, *Linn. Sp. Pl.* 916.—Europ. austr.
astrachanica, Spreng. Syst. iii. 397 = stereophylla.
athoa, DC. Prod. vi. 588 = rupestris.
atlantica, *Pomel, Nouv. Mat. Fl. Atl.* 283.—Algeria.
atrata, Willd. Sp. Pl. iii. 2290 = axillaris.
atropurpurea, Griseb. Spicil. Fl. Rumel. ii. 237 = C. Scabiosa.
atropurpurea, Olivier, Voy. i. 313 = Olivieriana.
atropurpurea, *Waldst. & Kit. Pl. Rar. Hung.* ii. 121. —Hungar.
atropurpurea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 65, 122 = C. Centaurium.
attica, Nym. Syll. 33 = graeca.
Aucheriana, *DC. Prod.* vi. 576.—Assyr.
augustana, Reichb. f. Ic. Fl. Germ. xv. 37. t. 785 = maculosa.
aurea, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 265.—Europ. austr.
aurea, Hort. ex Steud. Nom. ed. II. i. 318 = nicaeensis.
auriculata, Balb. ex Pers. Syn. ii. 486 = C. Seridis.
auriculata, Spreng. Pugill. i. 59 = stereophylla.
australis, *Benth. & Hook. f. Gen.* ii. 479.—Austral.
australis, Pané. Elem. 44 = sublanata.
australis, Poepp. ex DC. Prod. vi. 575 = chilensis.
austriaca, C. A. Mey. Verz. Pfl. Cauc. 64 = trichocephala.
austriaca, Reichb. Ic. Fl. Germ. xv. 37. t. 785, 3–4 = phrygia.

CENTAUREA :—
austriaca, Rochel, Pl. Banat. Rar. 76. t. 36. f. 5 = nervosa.
austriaca, *Willd. Sp. Pl.* iii. 2283.—Europ.; Reg. Caucas.; As. bor.
axillaris, *Willd. l. c.* 2290.—Europ.; Oriens.
babylonica, Bieb. Casp. 82 = C. Behen.
babylonica, *Linn. Mant.* ii. 460.—As. Min.; Syria.
badensis, Tratt. Arch. Gew. i. 28 = C. Scabiosa.
Balansae, *Boiss. & Reut. Diagn.* Ser. II. iii. 82.—Afr. bor.
baldensis, Pers. Syn. ii. 481 = alpina.
balearica, *Rodrig. in Bull. Soc. Bot. Fr.* xvi. (1869) 237.—Ins. Minorca.
Balsamita, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. t. 355 = Phaeopappus spectabilis.
Balsamita, *Lam. Encyc.* i. 667.—As. Min.; Persia.
banatica, Kern. ex Nym. Consp. 427 = arenaria.
barbara, *Pomel, Nouv. Mat. Fl. Atl.* 27.—Algeria.
Barrelieri, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 301 = Serratula pinnatifida.
Barrelieri, *Spreng. Syst.* iii. 403.—Hungar.
Behen, DC. Prod. vi. 567 = Szovitsiana.
Behen, Lam. Encyc. i. 665 = cerinthifolia.
Behen, *Linn. Sp. Pl.* 914.—As. Min.; Syria; Persia.
Belangeriana, Stapf, in Denkschr. Acad. Wien, l. (1885) 66 = phyllocephala.
bella, *Trautv. in Bull. Acad. St. Pétersb.* x. (1866) 394.—Reg. Caucas.
benedicta, Linn. Sp. Pl. ed. II. 1296 = Carbenia benedicta.
Berini, Sieber, ex Steud. Nom. ed. II. i. 319 = nervosa.
Besseriana, *DC. Prod.* vi. 585.—Europ. or.
Besseriana, Janka, ex Nym. Consp. 427 = tenuiflora.
bicolor, *C. Koch, in Linnaea*, xxiv. (1851) 428.—Reg. Caucas.
Biebersteinii, DC. Prod. vi. 583 = maculosa.
Biebersteinii, Schur, Verh. Siebenb. Ver. Naturf. (1859) 75 = Reichenbachii.
biflora, *Kalen. in Bull. Soc. Nat. Mosc.* xviii. (1845) i. 237.—Astrakhan.
bimorpha, Viv. Fl. Lib. Spec. 58. t. 24 = sonchifolia.
Boissieri, *DC. Prod.* vii. 303.—Hispan.
Boissieri, Walp. Rep. vi. 298 = graeca.
bombycina, *Boiss. in DC. Prod.* vii. 302.—Hispan.
borealis, Salisb. Prod. 208 = C. Scabiosa.
borysthenica, *Gruner, in Bull. Soc. Nat. Mosc.* xli. (1868) ii. 426.—Tauria.
Bourgaei, *Boiss. Fl. Orient.* iii. 637.—Lycia.
Bourgaei, Schult. ex Nym. Consp. 426 = Langei.
Brachtii, Reichb. f. Ic. Fl. Germ. xv. 35. t. 786 = maculosa.
brachyptera, DC. Prod. vi. 567 = C. Behen.
bracteata, Balb. ex Steud. Nom. ed. II. i. 319 = nigrescens.
bracteata, *Benth. & Hook. f. Gen.* ii. 481.—Syria.
bracteata, Scop. Delic. Insub. ii. 17 = amara.
brevicaulis, Boiss. Diagn. Ser. I. x. 120 = pallescens.
brevipappa, Boiss. & Reut. l. c. Ser. II. iii. 71 = nigrescens.
Brosseana ×, *Bonnet, in Bull. Soc. Bot. Fr.* xxvii. (1880) p. xi.—Gallia.
Bubanii, Timb. in Mém. Acad. Sc. Toul. Sér. VIII. i. ii. (1879) 187 = caerulescens.
budziackea, *Tardent, Ess. Hist. Nat. Bessar.* 61, nomen.—Bessarab.
bulbosa, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 111.—Chili.
busambarensis, *Guss. Fl. Sic. Syn.* ii. 873.—Sicilia.
cachinalensis, *Phil. Fl. Atac.* 34.—Chili.
cadmea, *Boiss. Diagn.* Ser. I. iv. 16.—Caria.
caerulescens, *Willd. Sp. Pl.* iii. 2319.—Europ. austr.
caespitosa, Cyrill. Pl. Neap. Nov. Fasc. i. 24 = sphaerocephala.
calalepis, *Boiss. in Tchihat. Asie Min. Bot.* t. 31.—As. Min.
calcaria, Jord. Obs. Pl. Crit. vii. (1849) 32 = C. Scabiosa.
Calcitrapa, Bieb. Fl. Taur. Cauc. iii. 594 = iberica.
Calcitrapa, *Linn. Sp. Pl.* 917.—Europ.; Afr. bor.; As. temp.
calcitrapoides, DC. Prod. vi. 597 = iberica.
calcitrapoides, Gouan, Fl. Monsp. 461 = Pouzini.
calcitrapoides, *Linn. Cent.* i. 29; *Amoen. Acad.* iv. 291.—Reg. Caucas.; As. Min.
calcitrapoides, Thuill. Fl. Par. 446 = C. Calcitrapa.
calendulacea, Lam. Encyc. i. 668 = Zoegea leptaurea.

CENTAUREA :—
calliacantha, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 32 = patula.
calocephala, Willd. Enum. Hort. Berol. 928 = atropurpurea.
calolepis, Boiss. Diagn. Ser. I. vi. 129 = diffusa.
calophylla, Steud. Nom. ed. I. 173 = atropurpurea.
calva, *Ledeb. ex DC. Prod.* vi. 583.—Mont. Altai.
calva, Reut. ex Nym. Consp. 429 = alba.
cana, C. A. Mey. ex Steud. Nom. ed. II. i. 320 = Ledebourii.
cana, Sibth. & Sm. Fl. Graec. Prod. ii. 198 = axillaris.
canadensis, *Hort. Huber, ex Regel, Ind. Sem. Hort. Petrop.* (1866) 90.—Am. bor.
canariensis, *Willd. Enum. Hort. Berol.* 928.—Ins. Teneriff.
cancellata, Sieber, ex Spreng. Syst. iv. 406 = aegyptiaca.
candidissima, Lam. Encyc. i. 669 = C. Cineraria.
Candolleana, *Boiss. Diagn.* Ser. I. vi. 127.—Persia.
Candollei, *C. Koch, in Linnaea*, xxiv. (1851) 424.—Reg. Caucas.
capillata, *Linn. Sp. Pl.* 910 (sp. dub.).—Sibir.; Hispan.
cappadocica, *DC. Prod.* vi. 595.—As. Min.
caprina, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 394 = ovina.
carduifolia, Salisb. Prod. 208 = C. Calcitrapa.
carduiformis, DC. Prod. vi. 590.—As. Min.; Persia.
carduncella, DC. l. c. 591 = raphanina.
Cardunculus, *Boiss. Diagn.* Ser. I. vi. 135.—Assyria.
Cardunculus, Pall. Reise, i. 500 = Serratula glauca.
Carduus, Forsk. Fl. Aegypt. Arab. 152 = Atractylis flava.
Carelli, *Ucria, in Roem. Arch.* i. 70.—Sicilia.
cariensis, *Boiss. Diagn.* Ser. I. iv. 19.—As. Min.
carneola, *Boiss. l. c.* Ser. II. iii. 83.—Cilicia.
carniolica, Host, Fl. Austr. ii. 517 = montana.
caroliniana, Walt. Fl. Carol. 214 = americana?
carpathica, Geners. Elem. n. 824 = montana.
carpatica, *Floriano, in Magyar Növént. Lapok*, ix. (1885) 128.—Hungar.
carpetana, Boiss. & Reut. Pugill. Pl. Nov. (1852) 65 = nigra.
carratracensis, *Langé, in Kjoeb. Vidensk. Meddel.* (1881) 94.—Hispan.
carthamoides, *Benth. & Hook. f. Gen.* ii. 479.—Sibir.
cassia, *Boiss. Diagn.* Ser. I. x. 108.—Syria.
Castellana, *Boiss. l. c.* vi. 129.—Hispan.
cataonica, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 669.—Cataonia.
caucasica, Bieb. ex DC. Prod. vi. 579 = montana.
Cavanillesiana, *Graells, in Mem. Acad. Cienc. Madr.* ii. (1859) 464.—Hispan.
Ceccariniana, *Boiss. & Heldr. Diagn.* Ser. II. iii. 81.—Macedon.
Centaurium, Host, ex Nym. Consp. 417 = Serratula coronata.
Centaurium, *Linn. Sp. Pl.* 910.—Italia.
centauroides, Bieb. Fl. Taur. Cauc. ii. 359 = salonitana.
centauroides, *Linn. Sp. Pl.* 918.—Europ.
cephalariaefolia, *Willk. in Flora*, xxxiv. (1851) 762.—Hispan.
ceratophylla, Moris, Elench. i. 55 = filiformis.
ceratophylla, Tenore, Cat. Hort. Neap. (1819) 72 = rupestris.
cerinthifolia, Sibth. & Sm. Fl. Graec. Prod. ii. 197.—Graecia.
chamaerhaponticum, *Ball, in Journ. Linn. Soc.* xvi. (1878) 526.—Marocco.
Chaubardi, Reichb. f. Ic. Fl. Germ. xv. 34. t. 781 = maculosa.
cheiracantha, *Fenzl, ex Boiss. Fl. Orient.* iii. 693.—As. Min.; Syria.
cheiranthifolia, Willd. Phytog. 12 = axillaris.
chilensis, *Bert. ex Bull. Férussac* (1830), 109, nomen; *et ex Hook. & Arn. in Bot. Beech. Voy.* i. 33.—Chili.
chlorantha, Adams, ex Ledeb. Fl. Ross. ii. 699 = montana.
Choulettiana, *Pomel, Nouv. Mat. Fl. Atl.* 31.—Algeria.
chrysanthemifolia, *Tausch, in Flora*, xii. (1829) i. *Erg.* 43.—Hab.?
chrysoleuca, *Boiss. Diagn.* Ser. I. iv. 21, in textu; *Fl. Orient.* iii. 678.—As. Min.
chrysolepis, Vis. in Mem. Ist. Venet. ix. (1860) 172 = orientalis.

CENTAUREA :—
chrysolopha, Boiss. & Kotschy, Diagn. Ser. II. iii. 75 = aggregata.
chthonocephala, Boiss. Fl. Orient. iii. 677 = ustulata.
cichoracea, Linn. Syst. ed. V. 1231 = Serratula cichoracea.
cichoracea, Stev. in Mém. Soc. Nat. Mosc. iv. (1813) 66 = Serratula caucasica.
cicutaefolia, Hornem. Hort. Hafn. ii. 849 = collina.
ciliata, Frivald. in Flora, xviii. (1835) 334 = pallida.
cilicica, *Boiss. & Bal. Diagn.* Ser. II. v. 113.—Cilicia.
Cineraria, Jacq. ex Nym. Consp. 425 = cinerea.
Cineraria, *Linn. Sp. Pl.* 912.—Europ.; Afr. bor.
Cineraria, Pall. Tabl. Taur. 58 = declinata.
Cineraria, Urv. in Mém. Soc. Linn. i. (1822) 371 = rutifolia.
cinerascens, *Bubani, in Nuov. Giorn. Bot. Ital.* v. (1873) 319.—Hispan.
cinerea, Griseb. Spicil. Fl. Rumel. ii. 236 = dissecta.
cinerea, *Lam. Encyc.* i. 669.—Italia.
cirrata, *Reichb. Fl. Germ. Excurs.* 214.—Europ. austr.
Clementei, *Boiss. in DC. Prod.* vii. 303.—Hispan.
coarctata, Spreng. Syst. iii. 407 = glomerata.
cochinchinensis, *DC. Prod.* vi. 601.—Cochinch.
coelolepis, *Fisch. & Mey. ex Heynh. Nom.* ii. 130, nomen.—Rossia.
collina, Asso, Syn. 127 = ornata.
collina, *Linn. Sp. Pl.* 918.—Reg. Mediterr.
collina, Schur, Enum. Pl. Transs. 408 = pseudo-collina.
collina, Scop. ex Nym. Consp. 429 = rupestris.
collina, Sibth. & Sm. Fl. Graec. t. 914 = salonitana.
comata, Jord. Pugill. 106 = pectinata.
commutata, Timb. ex Nym. Consp. 421 = nigra.
Comperiana, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) ii. 393 = diffusa.
concinna, *Trautv. in Act. Hort. Petrop.* ii. (1873) 553.—Armenia.
concinna, Willd. ex Steud. Nom. ed. II. i. 319 = pulcherrima.
congesta, Willd. ex DC. Prod. vi. 593 = melitensis.
conglomerata, *C. A. Mey. in Beitr. Pfl. Russl.* v. 44.—Rossia.
conifera, *Linn. Sp. Pl.* 915.—Reg. Mediterr.
conocephala, *C. Bolle, in Bonplandia*, viii. (1860) 134.—Ins. Canar.
consanguinea, *DC. Prod.* vi. 585.—As. Min.
consimilis, Bor. Fl. Cent. Fr. ed. III. ii. 351 = C. Jacea.
contracta, Viv. Fl. Lib. Spec. 58. t. 24. f. 1, 2 = Delilei.
Cordonis, Boiss. & Orph. ex Boiss. Fl. Orient. iii. 663 = Ceccariniana.
coriacea, Waldst. & Kit. Pl. Rar. Hung. ii. 214 = C. Scabiosa.
corinthiaca, Boiss. & Heldr. Diagn. Ser. II. vi. 113 = achaia.
Conuti, Reichb. Hort. t. 146 = alpina.
coronopifolia, *Lam. Encyc.* i. 667.—Iberia.
corymbosa, Pourr. in Mém. Acad. Sc. Toul. iii. (1788) 310 = maculosa.
Cossoniana, *Ball, in Journ. Bot.* xi. (1873) 369.—Marocco.
Costae, *Willk. in Linnaea*, xxx. (1859-60) 115.—Hispan.
crassifolia, *Bertol. in Ann. Stor. Nat.* ii. (1829) 359.—Ins. Melita.
cretica, *Nym. Syll.* 34.—Ins. Creta.
cretica, Spreng. Syst. iii. 408 = Carthamus creticus.
crinita, *Willd. ex Spreng. l. c.* 400.—Hab.?
cristata, *Bartl. in Wendl. Beitr.* ii. 119.—Austria.
crithmifolia, *Vis. Fl. Dalmat.* ii. 40.—Dalmatia.
crocodylioides, Boiss. Diagn. Ser. I. x. 118 = C. Crocodylium.
Crocodylium, *Linn. Sp. Pl.* 919.—Syria.
cruenta, *Willd. Enum. Hort. Berol.* 929.—Hab.?
Crupina, Bory & Chaub. Nouv. Fl. Pélop. 59 = Crupina pseudocrupina.
Crupina, Desf. Fl. Atlant. ii. 292 = Crupina intermedia.
Crupina, Linn. Sp. Pl. 909 = Crupina vulgaris.
Crupina, Sibth. & Sm. Fl. Graec. t. 900 = Crupina Crupinastrum.
Crupinastrum, Moris, Ind. Sem. Hort. Taur. (1852) 12; et in Bot. Zeit. i. (1843) 168 = Crupina Crupinastrum.
crupinoides, Desf. Fl. Atlant. ii. 293 = Volutarella bicolor.
Csatoi, *Borbas, in Ertek. Koré Magyar Tudom. Akad.* ix. (1879) 29.—Hungar.
cucullata, *Spreng. ex Steud. Nom.* ed. I. 174.—Hab.?

CENTAUREA :—
cuneifolia, Orphan. ex Boiss. Fl. Orient. iii. 645 = sublanata.
cuneifolia, *Sibth. & Sm. Fl. Graec.* t. 901.—Macedon.
cuspidata, Aucher, ex DC. Prod. vi. 577 = Aucheriana.
cuspidata, *Vis. in Flora*, xii. (1829) 1. *Erg.* 22.—Dalmatia.
cyaneum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Cyanus.
cyanifolia, *C. Koch, in Linnaea*, xxiv. (1851) 436.—Reg. Caucas.
cyanifolia, Poir. Encycl. Suppl. ii. 158 = solstitialis.
cyanoides, *Bergg. & Wahlenb. in Isis*, V. xxi. (1828) 971.—Syria.
cyanoides, DC. Prod. vi. 578 = axillaris.
cyanoides, *C. Sm. in Buch, Beschr. Canar. Ins.* 147.—Ins. Canar.
Cyanus, *Linn. Sp. Pl.* 911.—Europ.; Reg. Caucas.
cylindrocarpa, Reichb. f. Ic. Fl. Germ. xv 34. t. 781 = cuneifolia et sublanata.
cynaroides, C. Sm. ex Buch, in Berl. Acad. Abh. 1816–17 (1819) 369 = Serratula canariensis.
cynaroides, *Link, in Buch, Beschr. Canar. Ins.* 147.—Ins. Canar.
cyrtolepis, *Ledeb. Fl. Ross.* ii. 709.—As. Min.
damascena, *Boiss. Diagn.* Ser. I. x. 112.—Syria.
dealbata, Moris, ex Nym. Consp. 425 = C. Cineraria.
dealbata, *Willd. Sp. Pl.* iii. 2295.—As. Min.; Reg. Caucas.; Persia.
Debeauxii, Godr. & Gren. Fl. Fr. ii. 243 = nigra.
decipiens, Thuill. Fl. Par. ed. II. 445 = C. Jacea.
declinata, Aucher, ex DC. Prod. vi. 579 = lanigera.
declinata, Bieb. Fl. Taur. Cauc. iii. 590 = leucophylla.
decumbens, Boiss. ex DC. Prod. vii. 303 = Boissieri.
decumbens, DC. Prod. vi. 576 = leucophylla.
decumbens, Dubois, Fl. Orl. ex Pers. Syn. 485 = C. Jacea.
decurrens, *DC. Prod.* vi. 596.—Hab.?
deflexa, *Tatarinow, in Beitr. Kenntn. Russ. Reich.* xx. 252.—Soongaria.
deinacantha, *Boiss. & Haussk. in Fl. Orient.* iii. 665.—Kurdistan.
Delilei, *Godr. in Mém. Acad. Montp.* (*Sect. Médic.*) i. (1853) 435.—Gallia?
denticulata, Lejeune, Rev. Fl. Spa, 183 = nigrescens.
depressa, *Bieb. Fl. Taur. Cauc.* ii. 346.—Reg. Caucas.; Iber.
dertosensis, Costa, Fl. Catal. (1864) 141 = podospermifolia.
derventana, Janka, in Oestr. Bot. Zeitschr. xxi. (1871) 287 = Kerneriana.
derventana, *Vis. & Panč. in Mem. Ist. Venet.* xii. (1864) 472.—Oriens.
desmicephala, Fenzl, Ind. Hort. Vind. (1855) = aggregata.
deusta, Tenore, Prod. Fl. Nap. p. li = alba.
Devauxii, Bor. ex Nym. Consp. 431 = C. Calcitrapa.
dianthoides, *Remy, in C. Gay, Fl. Chil.* iv. 312.—Chili.
dichroa, *Boiss. & Heldr. Diagn.* Ser. I. x. 111.—Syria.
dichroantha, Kern. in Oestr. Bot. Zeitschr. xxiv. (1874) 104 = sordida.
diffusa, *Lam. Encyc.* i. 675.—Europ. austr. or.; As. Min.
diluta, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 261.—Afr. bor.
dimorpha, *Viv. Fl. Lib. Spec.* 58. t. 24. f. 3.—Afr. bor.
diomedea, *Gasp. Descr. Trem.* 19, 23.—Ital.; Ins. Trem.
discoidea, Balb. ex DC. Prod. vi. 573 = flosculosa.
dissecta, *Hill, Veg. Syst.* iv. 33.—Germ.
dissecta, *Tenore, Prod. Fl. Nap.* p. li—Italia; Graecia.
divaricata, Wall. Cat. n. 2894 = Volutarella divaricata.
divergens, Lange, in Kjoeb. Vidensk. Meddel. (1861) 87 = Langei.
divergens, *Vis. Fl. Dalmat.* ii. 37. t. 12 b.—Europ. austr.
diversifolia, Amo, ex Campo & Amo, in Revista, v. (1855) 56 = Amoi.
diversifolia, Lag. ex Boiss. Voy. Bot. Espagne, ii. 354 = aspera.
Dornacina, Amo, Descr. Pl. Nuev. ex Willk. & Lange, Prod. Fl. Hisp. ii. 173 = Serratula nudicaulis.

CENTAUREA :—
doryphora, Fenzl, ex Boiss. Fl. Orient. iii. 665 = Urvillei.
drabaefolia, Auct. = seq.
drabifolia, *Sibth. & Sm. Prod.* ii. 202; *Fl. Graec.* t. 910.—Bithynia.
dracunculifolia, *Dufour, in Ann. Sc. Nat.* xxiii. Sér. I. (1831) 157.—Hispan.
druentica ×, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 342, 348.—Gallia.
dubia, S. G. Gmel. ex Steud. Nom. ed. I. 174 = Xeranthemum annuum.
dubia, Krock. Fl. Sil. Suppl. ii. 331 = humilis?
dubia, Suter, Fl. Helv. ii. 202 = nigrescens.
Duboisii, Bor. Fl. Centr. Fr. ed. III. ii. 350 = C. Jacea.
Dumortieri, *Thys, in Bull. Soc. Bot. Belg.* viii. (1869) 464.—Belg.
dumulosa, *Boiss. Diagn.* Ser. I. x. 111.—Syria.
dutosensis, Costa, ex Nym. Consp. 429 = podospermifolia.
ebenoides, *Heldr. ex S. Moore, in Journ. Bot.* xvi. (1878) 133.—Graecia.
elata, *Poir. Encyc. Suppl.* ii. 150.—Mauritan.
elbrusensis, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 131.—Persia.
elegans, All. Fl. Pedem. i. 163 = Galactites tomentosa.
elegans, Salisb. Prod. 207 = C. Cineraria.
elongata, Desf. ex DC. Prod. vi. 604 (Quid?).
elongata, Schousb. Vextr. Marokko, 199 = diluta.
elongata, *Webb & Berth. Phyt. Canar.* ii. 360.—Ins. Canar.
emigrantis, *Bubani, in Nuov. Giorn. Bot. Ital.* v. (1873) 318.—Hispan.
Endresii, Hochst. & Steud. ex Steud. Nom. ed. II. i. 320 = nigra.
eriocephala, Boiss. & Reut. Diagn. Ser. II. iii. 86 = dimorpha.
eriophora, Forsk. Fl. Aegypt. Arab. 74. n. 465 = aegyptiaca.
eriophora, *Linn. Sp. Pl.* 916.—Gall.; Lusit.
eriophylla, Boiss. & Bal. ex Boiss. Fl. Orient. iii. 678 = chrysoleuca.
eriophylla, *Spreng. Syst.* iii. 404.—Hab.?
erucifolia, Linn. Sp. Pl. 909 = Volutarella erucifolia.
eryngioides, *Lam. Encyc.* i. 675.—Arab.; Syria.
eryngioides, Sibth. & Sm. Fl. Graec. t. 913 = achaia.
Escobilla, Coss. ex Nym. Consp. 426 = castellana.
euphratica, Boiss. Diagn. Ser. I. vi. 125 = polypodiifolia.
exaltata, Adam, in Weber & Mohr, Beitr. i. 74 = coronopifolia.
exarata, *Boiss. ex Coss. Notes Pl. Crit.* 116.—Afr. bor.
exscapa, Urv. in Mém. Soc. Linn. Par. i. (1822) 373 = acicularis.
Fabrei ×, *Bonnet, ex Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 349.—Gallia.
Fenzlii, *Reichardt, in Verh. Zool. Bot. Ges. Wien*, xiii. (1863) 1039.—As. temp.
Ferdinandi, Gren. Cat. Sem. Hort. Bot. Grat. (1847) Collect. 20 = nervosa.
ferox, *Desf. Fl. Atlant.* ii. 297. t. 242.—Numidia.
filiformis, Bieb. ex DC. Prod. vi. 566 = pulchella.
filiformis, *Viv. Fl. Cors. App.* 6.—Corsic.; Sardin.
Fischeri, Willd. Enum. Hort. Berol. Suppl. 61 = axillaris.
fistulosa, *Wender. ex Steud. Nom.* ed. II. i. 320.—Hab.?
flavicans, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 149 = C. Jacea.
floccosa, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 110.—Chili.
floccosa, Schloss. ex Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 149 = montana.
floccosa, Schloss. & Vukot. Syll. Croat. 46 = axillaris.
flosculosa, Balb. ex Nym. Consp. 422 = nervosa.
flosculosa, *Willd. Sp. Pl.* iii. 2285.—Europ. austr.
foliosa, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iii. 672.—Syria.
Fontanesii, *Spach, ex Dur. in Duch. Rev. Bot.* ii. (1847) 429.—Algeria.
Forardi, *Hornem. Hort. Hafn. Suppl. Add.* 159.—Italia.
Fraasii, Noë & Bal. ex Boiss. Fl. Orient. iii. 681 = monacantha.
Fraasii, Sch. Bip. ex Nym. Consp. 429 = seq.
Fraatii, Sch. Bip. ex Tchihat. Asie Min. Pot. ii. 323 = Spruneri.

CENTAUREA :—
fragilis, *Dur. in Duch. Rev. Bot.* ii. (1847) 429.—Afr. sept.
fraylensis, *Sch. Bip. ex Nym. Consp.* 420.—Hispan.
Friderici, *Vis. Fl. Dalmat.* ii. 40. t. 12 C.—Dalmat.
fruticosa, Linn. Syst. ed. X. 1229 = Stachelina fruticosa.
fuliginosa, Dollin. ex Nym. Consp. 424 = C. Scabiosa.
fulva, Huet, in Ann. Sc. Nat. Sér. III. xix. (1853) 254 = nigra.
Funkii, Boiss. & Reut. Diagn. Ser. II. iii. 83 = macrorrhiza.
Funkii, Sch. Bip. ex Willk. in Linnaea, xxv. (1852) 37 = Boissieri.
furfuracea, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 363.—Algeria.
fuscata, Desf. Fl. Atlant. ii. 302. t. 244 = nicaeënsis.
fuscata, Jord. Pugill. 105 = pectinata.
fuscescens, *Pomel, Nouv. Mat. Fl. Atl.* 28.—Algeria.
Galactites, Linn. Sp. Pl. 919 = Galactites tomentosa.
garganica, Sch. Bip. ex Nym. Consp. 425 = subtilis.
Gaudini, Boiss. & Reut. Diagn. Ser. II. iii. 70 = amara.
Gayana, *Remy, in C. Gay, Fl. Chil.* iv. 310.—Chili.
geluensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 674.—Persia.
geocephala, Boiss. Fl. Orient. iii. 677 = ustulata.
Gerberi, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) II. 391, in textu.—Ross. austr.
Germaniciae, Haussk. ex Boiss. Fl. Orient. iii. 682 = C. Behen.
gigantea, *Sch. Bip. ex Boiss. l. c.* 672.—Persia.
glaberrima, *Tausch, in Flora*, x. (1827) II. 249.—Dalmatia.
glandulifera, *Vukot.—Cf. Oestr. Bot. Zeitschr.* xxx. (1880) 163.—Europ.
glastifolia, *Linn. Sp. Pl.* 915.—As. Min.; Reg. Cauc.; Sibir.
glauca, Desf. ex DC. Prod. vi. 587 = calocephala.
glauca, Puschk. ex Willd. Sp. Pl. iii. 2276 = moschata.
glaucescens, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. i. (1854) 31.—Galatia.
Glehni, *Trautv. in Act. Hort. Petrop.* iv. (1876) 382.—Rossia.
globosa, Hort. ex DC. Prod. vi. 569 = alba.
glomerata, *Vahl, Symb. Bot.* ii. 94.—Aegypt.
glomerata, Webb & Berth. Phyt. Canar. ii. 360 = melitensis.
Gmelini, *Steud. Nom.* ed. II. i. 320.—Sibir.
goniocaula, *Boiss. Fl. Orient.* iii. 647.—Lydia.
gracilior, Reut. ex Nym. Consp. 421 = amara.
gracilis, Bor. ex Nym. l. c. = C. Jacea.
gracca, *Boiss. & Sprun. Diagn.* Ser. I. vi. 128.—Graecia.
graeca, Griseb. Spicil. Fl. Rumel. ii. 242 = Ceccariniana.
Graëlsii, *Nym. Consp.* 429.—Hispan.
Grafiana, DC. Prod. vi. 587 = sordida.
graminifolia, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 159 = montana.
granatensis, *Boiss. ex DC. Prod.* vii. 303.—Hispan.
grandiflora, Moç. & Sesse, ex DC. Prod. vi. 575 = americana.
grandiflora, Pall. Reise, iii. 237 = monanthos.
grinensis, Reut. Ind. Sem. Hort. Genev. (1857) 4.—Cf. Linnaea, xxix. (1857–58) 722 = C. Scabiosa.
Grisebachii, Nym. Consp. 427 = graeca.
gudrunensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 673.—Kurdist.
Guicciardii, *Boiss. l. c.* 661.—Bacotia.
gymnocarpa, *Moris & De Not. Fl. Insul. Caprar.* 76. t. 2.—Ins. Caprar.
gymnoclada, *Benth. & Hook. f. Gen.* ii. 482.—Mesopot.
Haenseleri, *Boiss. Voy. Bot. Espagne*, ii. 349.—Hispan.
Hanrii, Jord. Obs. Pl. Crit. v. (1847) 70 = coerulescens, limbata.
Haussknechtii, *Boiss. Fl. Orient.* iii. 670.—Syria.
Haynaldi, Borb. ex Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 149 = C. Jacea.
helenioides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 683.—Pontus?
hellenica, *Boiss. & Sprun. Diagn.* Ser. I. vi. 131.—Graecia.
hemiptera ×, *Borb. in Oestr. Bot. Zeitschr.* xxviii. (1878) 392.—Europ. austr.

CENTAUREA:—
Herbichii, *Janka, in Flora*, xli. (1858) 441.—Transsylv.
Hermonis, Boiss. Diagn. Ser. I. x. 122 = iberica.
heterocarpa, *Boiss. & Gaill. ex Boiss. Fl. Orient.* iii. 680.—Syria.
heterophylla, Sprun. ex Boiss. l. c. 663 = Spruneri.
heterophylla, Willd. Sp. Pl. iii. 2314 = aspera.
Heerffelii, Reichb. f. Ic. Fl. Germ. xv. 27. t. 801 = Kotschyana.
hierapolitana, *Boiss. Diagn.* Ser. I. iv. 15.—Caria.
hispida, DC. Prod. vi. 589 = reflexa.
Hladnickiana, *Fisch. ex Heynh. Nom.* ii. 130, nomen.—Europ. austr.
Hochstetteri, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 438.—Afr. trop.
Hoefftiana, *C. A. Mey. in Mém. Acad. Pétersb.* Sér. VI. *Sc. Nat.* vii. (1855) 7.—Reg. Casp.
Hohenackeri, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 395 = ovina.
Hohenackeriana, Trautv. in Act. Hort. Petrop. viii. (1883) 493 = praec.
hololeuca, *Boiss. Fl. Orient.* iii. 694.—Syria.
horrida, *Badaro, in Brugnat. Giorn. Fis.* (1824) 363.—Liguria; Sardin.
horrida, Tenore, Ind. Sem. Hort. Neap. (1829) 15; Syll. 449 = C. Calcitrapa.
Huetii, *Boiss. Diagn.* Ser. II. iii. 69.—Armenia.
humifusa, Gueldenst. ex Ledeb. Fl. Ross. ii. 696 = decumbens.
humilis, Schrank, Baier. Fl. ii. 376 = pullata?
hyalina, L'Hérit. ex DC. Prod. vi. 566 = pulchella.
hyalolepis, Boiss. Diagn. Ser. I. vi. 133 = pallescens.
hybrida, All. Fl. Pedem. i. 161 = corymbosa.
hybrida, Chaix, in Vill. Hist. Pl. Dauph. iii. 54 = C. Calcitrapa.
hymenolepis, *Trautv. in Act. Hort. Petrop.* iii. (1874) 272.—Reg. Caucas.
hypericifolia, Brouss. ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 355 = canariensis.
hypoleuca, *DC. Prod.* vi. 575.—As. Min.
hyssopifolia, Georgi, Beschr. Russ. Reich. iii. IV. 1263 = trichocephala.
hyssopifolia, *Vahl, Symb. Bot.* i. 75.—Hispan.
Hystrix, Boiss. Voy. Bot. Espagne, ii. 351 = polyacantha.
iberica, *Trevir. ex Spreng. Syst.* iii. 406.—Iber.; Armen.; Caucas.
idaea, *Boiss. & Heldr. Diagn.* Ser. I. x. 119.—Creta.
incana, Burm. f. Fl. Cap. Prod. 28 = Arctotis argentea.
incana, *Desf. Fl. Atlant.* ii. 301.—Marocco.
incana, Lag. Gen. et Sp. Nov. 32 = Lagascae.
incana, Losc. & Pardo, Ser. Inconf. 59 = tenuifolia.
incana, *Tenore, Prod. Fl. Nap.* p. li.—Italia.
incompta, *Vis. Fl. Dalmat.* 38. t. 13^{b}. fig. 1.—Dalmat.
indica, Less. in Linnaea, vi. (1831) 84 = Goniocaulon glabrum.
indurata, Janka, in Flora, xli. (1858) 444 = nigra.
infestans, *Dur. in Duch. Rev. Bot.* ii. (1847) 430.—Afr. bor.
intacta, *Ledeb. Fl. Ross.* ii. 705.—Reg. Caucas.
integrifolia, *C. A. Mey. Verz. Pfl. Cauc.* 64.—Reg. Caucas.
integrifolia, Tausch, in Flora, xi. (1828) II. 485 = nigra.
integrifolia, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 153 = C. Scabiosa.
integrisquama, Vukot. l. c. 152 = C. Scabiosa.
intermedia, Cariot, Etud. Fl. ed. V. 317 = axillaris.
intermedia, *Mutel, in Duch. Rev. Bot.* i. (1845-46) 400.—Algeria.
interrupta, *Hoffmgg. & Link, Fl. Port.* ii. 227.—Lusitan.
intricata, *Boiss. Diagn.* Ser. I. vi. 130.—Persia.
intybacea, *Lam. Encyc.* i. 671.—Gallia; Hispan.
inuloides, Fisch. ex Nym. Consp. 418 = trinervia.
inusta, *Hoffmgg. ex Heynh. Nom.* ii. 131, nomen.—Hab.?
involucrata, *Desf. Fl. Atlant.* ii. 295.—Alger.
iserniana, Graells, ex Willk. & Lange, Prod. Fl. Hisp. ii. 162 = Janerii.
Isernii, *Willk. in Willk. & Lange, l. c.* 155.—Hispan.
Isnardi, Linn. Sp. Pl. 916 = aspera.
ispahanica, *Boiss. Diagn.* Ser. I. vi. 133.—Persia.
italia, Steud. Nom. ed. II. i. 320 = seq.
italica, Pers. Syn. ii. 484 = C. Scabiosa.

CENTAUREA:—
ixodes, *Pomel, Nouv. Mat. Fl. Atl.* 284.—Algeria.
Jacea, *Linn. Sp. Pl.* 914.—Europ.; As. med.
jacobaeaefolia, *Lam. Encyc.* i. 675.—Hab.?
jacobaeaefolia, Spreng. Syst. iii. 404 = argyrophylla.
Jacobi, Dufour, in Bull. Soc. Bot. Fr. vii. (1860) 348 = sonchifolia.
Janerii, *Graells, in Mem. Acad. Cienc. Madr.* ii. (1859) 466 (*Ramill.* 8).—Hispan.
Jankeana, *Simk. in Termész. Füzetek*, i. (1817) 168.—Europ. austr.
Jordaniana, *Gren. & Godr. Fl. Fr.* ii. 245.—Gallia.
jurineaefolia, Boiss. Diagn. Ser. II. iii. 73 = rutifolia.
juvenalis ×, *Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.)* i. (1853) 433.—Gallia.
karduchorum, *Boiss. Fl. Orient.* iii. 633.—Armenia.
karlowensis, Frivald. ex Hampe, in Flora, xx. (1837) I. 228 = napulifera.
Karschiana, Auersw. ex Nym. Consp. 426 = cristata.
Karstiana, Griseb. Spicil. Fl. Rumel. ii. 240 = pallida.
Kartschiana, *Scop. Carn.* ed. II. ii. 140. t. 55.—Carniol.
Kerneriana, *Janka, in Oestr. Bot. Zeitschr.* xxii. (1872) 178.—Thracia.
kestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 325.—Turkest.
kilaea, *Boiss. Fl. Orient.* iii. 643.—Byzant.
Kochii, F. Schultz, ex Nym. Consp. 421 = nigrescens.
Kolczygini, *Kalen. in Bull. Soc. Nat. Mosc.* xviii. (1845) I. 236.—Reg. Caucas.
Kotschyana, Heuff. in Koch, Syn. Fl. Germ. 437 = C. Scabiosa.
Kralikii, Boiss. Diagn. Ser. II. iii. 84 = dimorpha.
kurdica, *Reichardt, in Verh. Zool.-Bot. Ges. Wien*, xiii. (1863) 1042.—Kurdist.
laciniata, Guss. in DC. Prod. vi. 582 = ambigua.
laconica, *Boiss. Fl. Orient.* iii. 660.—Laconia.
lactucaefolia, *Boiss. l. c.* 622.—Ins. Rhodus.
Lagascae, *Nym. Syll.* 33.—Hispan.
Lagascana, Graells, in Mem. Acad. Cienc. Madr. ii. (1859) 465 (Ind. Pl. Nov. 7) = Graëlsii.
lanata, DC. Prod. vi. 610 = Carthamus lanatus.
lanata, Roxb. Hort. Beng. 62; Fl. Ind. iii. 444 = C. Cyanus.
lancifolia, *Sieber, ex Spreng. Syst.* iii. 406.—Creta.
Langeana, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 157 = Langei.
Langei, *Nym. Suppl. Syll.* 6.—Hispan.
lanigera, *DC. Prod.* vi. 579.—Cappadoc.
lappacea, Tenore, Ind. Sem. Hort. Neap. (1829) 15 = solstitialis.
lasiorhiza, *DC. Prod.* vi. 600.—Persia.
latifolia, Pers. Syn. ii. 487 = centauroides.
latisquama, DC. Prod. vi. 589 = salonitana.
laxa, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 640.—Mesopot.
Ledebourii, Steud. Nom. ed. II. i. 320 = cana.
leptocephala, *Boiss. Diagn.* Ser. I. x. 110.—Syria.
leptoloma, Panč. ex Nym. Consp. 426 = maculosa.
leptophylla, *C. Koch, in Linnaea*, xxiv. (1851) 419.—Lazistan.
leucantha, Pourr. in Mém. Acad. Toul. iii. (1788) 310 = intybacea.
leucantha, Web. & Mohr, Beitr. i. 74. n. 49 = montana.
leucocephala, Schur, Enum. Pl. Transs. 403 = phrygia.
leucolepis, DC. Prod. vi. 568 = alba.
leucolepis, Ledeb. ex Nym. Consp. 420 = sterilis.
leucolepis, Tenore, ex Nym. l. c. = splendens.
leucophaea, *Jord. Obs. Pl. Crit.* v. (1847) 64.—Gallia.
leucophylla, *Bieb. Fl. Taur. Cauc.* iii. 591.—Reg. Caucas.
leuzeoides, *Walp. Ann.* i. 447.—Media.
ligerina ×, *Franch. in Bull. Soc. Bot. Fr.* xxvii. (1880) p. xxii.—Gallia.
limbata, *Hoffmgg. & Link, Fl. Port.* ii. 221.—Lusitan.
linarifolia, Steud. Nom. ed. II. i. 321 = linifolia.
linearifolia, Lam. Encyc. i. 667 = linifolia.
lingulata, Lag. Gen. et Sp. Nov. 32 = variegata.
linifolia, Lam. Encyc. i. 667 = hyssopifolia.
linifolia, *Linn. Mant.* i. 117; *Vahl, Symb. Bot.* i. 75.—Hispan.; Italia.
Linnaeana, Walp. Ann. i. 450 = Fontanesii.
Lippii, Linn. Sp. Pl. 910 = Volutarella Lippii.
livoniana, Weinm. ex Ledeb. Hort. Dorp. 38 = C. Jacea.
longepedunculata, *Sch. Bip. ex Boiss. Fl. Orient.* iii. 668.—Persia.

CENTAUREA:—
lugdunensis, Jord. Obs. Pl. Crit. v. (1847) 49 = montana.
Luschaniana, *Heimerl, ex Stapf, in Denkschr. Acad. Wien*, l. (1885) 113.—Lycia.
lusitanica, Boiss. & Reut. Diagn. Ser. II. iii. 85 = sphaerocephala.
lutea, *Gueldenst. It.* ii. 43, *ex Ledeb. Fl. Ross.* ii. 712.—Ross. austr.
lycaonica, *Boiss. & Heldr. Diagn.* Scr. II. v. 111.—As. Min.
lycia, *Boiss. Diagn.* Ser. I. x. 109.—Lycia.
lycopifolia, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iii. 693.—Syria.
lydia, *Boiss. Diagn.* Ser. I. iv. 20.—Bithynia.
lyraefolia, Salisb. Prod. 208 = salmantica.
lyrata, Bell. App. Fl. Pedem. 40 = rhapontica.
lyrata, *Pers. Syn.* ii. 486.—Uruguay.
lyrata, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 270.—Syria.
lyrophylla, *Griseb. Spicil. Fl. Rumel.* ii. 238.—Europ. austr.
macedonica, *Boiss. Diagn.* Ser. I. vi. 130.—Graecia.
macracantha, Heldr. ex Boiss. Fl. Orient. iii. 690 = iberica.
macroacantha, Guss. Fl. Sic. Syn. ii. 518 = calcitrapoides.
macrocarpa, *Boiss. Fl. Orient.* iii. 677.—Persia.
macrocephala, *Puschk. ex Willd. Sp. Pl.* iii. 2299.—Armen.
macrolopha, *Fenzl, ex Boiss. Fl. Orient.* iii. 595.—Oriens.
macrorrhiza, *Willk. in Linnaea*, xxv. (1852) 38.—Hispan.
maculata, Andrz. Rys Botan. i. (1823) 12 = maculosa.
maculosa, Eichw. Skizze, 146 = Biebersteinii.
maculosa, Guss. Pl. Rar. 359 = cinerea.
maculosa, *Lam. Encyc.* i. 669.—Gallia; Hispan.; Italia.
maculosa, Noé, ex Nym. Consp. 425 = cuneifolia.
malacitana, Boiss. Diagn. Ser. II. iii. 85 = sonchifolia.
margaritacea, Tenore, Syll. Add. iii. A. 628 = splendens.
marginata, Tenore, Prod. Fl. Nap. p. lii = nicaeënsis.
mariana, Nym. Suppl. Syll. 6 = macrorrhiza.
mariolensis, *Rouy, in Bull. Soc. Bot. Fr.* xxix. (1882) 110.—Gallia.
maritima, Dufour, in Ann. Sc. Nat. Sér. I. xxiii. (1831) 165 = sonchifolia.
maroccana, *Ball, in Journ. Bot.* xi. (1873) 370.—Marocco.
Marschalliana, *Spreng. Syst.* iii. 398.—Podol.; Caucas.
Massoniana, *Lowe, in Hook. Kew Journ.* viii. (1856) 297.—Ins. Madera.
mathiolifolia, Boiss. Diagn. Ser. I. iv. 19 = axillaris.
maxima, Forsk. Fl. Aegypt. Arab. 152 = verbascifolia.
melitensis, *Linn. Sp. Pl.* 917.—Europ.; Am. austr.
membranacea, Jacquem. Voy. Bot. 97 = Saussurea denticulata.
membranacea, Lam. Encyc. i. 666 = monanthos.
menteyerica, Vill. Hist. Pl. Dauph. i. 365; iii. 48 = C. Scabiosa.
Meryonis, DC. Prod. vi. 598 = iberica.
mexicana, DC. l. c. 575 = americana.
micracantha, *Dufour, in Ann. Sc. Nat.* Sér. I. xxiii. (1831) 164.—Hispan.
micrantha, Hoffmgg. & Link, Fl. Port. ii. 220 = castellana.
micranthos, S. G. Gmel. It. i. t. 23. f. 1 = maculosa.
microcephala, *Benth. & Hook. f. Gen.* ii. 480.—Cappadoc.
microlepis, *Boiss. Fl. Orient.* iii. 651.—Lycia.
microlonchoides, *Boiss. Diagn.* Ser. I. vi. 127.—Persia.
microptilon, Godr. & Gren. Fl. Fr. ii. 242 = nigra.
Mierghii, Jord. in Cat. Grain. Grenob. (1850) 15; et Pugill. 110 = paniculata.
Minoa, Heldr. ex Boiss. Fl. Orient. iii. 662 = Guicciardii.
minor, Willd. ex Spreng. Syst. iii. 403 = polymorpha.
mixta, DC. Prod. vi. 594 = hellenica.
mollis, Bess. [Fl. Galic. ii. 207 ?] = axillaris.
mollis, Schleich. Cat. Pl. Helv. ed. IV. 12 = C. Jacea.
mollis, Waldst. & Kit. Pl. Rar. Hung. iii. 243 t. 219 = montana.
monacantha, *Boiss. Fl. Orient.* iii. 681.—Thracia.

CENTAUREA :—
monacantha, *Clark, Trav.* ii. 354.—Cyprus.
monantha, *Georgi, Bemerk. Reise Russ. Reich.* i. 231. —Rossia.
monocephala, *Remy, in C. Gay, Fl. Chil.* iv. 311.—Chili.
montana, Auct. ex Willk. & Lange, Prod. Fl. Hisp. ii. 159 = variegata, axillaris.
montana, Burm. f. Prod. Fl. Cap. 28 = sempervirens.
montana, Costa, Fl. Catal. 139 = semidecurrens, lugdunensis.
montana, *Linn. Sp. Pl.* 911.—Europ.
monticola, *Boiss. ex DC. Prod.* vii. 302.—Hispan.
Moritziana, Hegetschw. Fl. Schw. 851 = phrygia.
Moschata, *Linn. Sp. Pl.* 909.—Oricns.
Moschus, Habl. in S. G. Gmel. It. iv. 176 = Jurinea depressa.
Mouelli, *Blanchet, in Bull. Soc. Bot. Fr.* xxviii. (1880) 21.—Gallia.
mucronata, Forsk. Fl. Aegypt. Arab. 151 = pumila.
mucronifera, *DC. Prod.* vi. 569.—Cappadoc.
multifida, *Pers. Syn.* ii. 487.—Afr. bor.
multiflora, Pers. l. c. 486 = sicula.
Mureti, Jord. Pugill. 108 = coerulescens.
muricata, Linn. Sp. Pl. 918 = Volutarella muricata.
muricata, Spreng. Syst. iii. 402 = Volutarella Lippii.
musarum, *Boiss. & Orph. Diagn.* Ser. II. v. 112.—Graecia.
mutabilis, *St. Amans, in Mém. Mus. Par.* i. (1815) 477.—Mont. Pyren.
myacantha, DC. Fl. Fr. iv. 101 = C. Calcitrapa.
myconia, Boiss. & Sart. Diagn. Ser. II. vi. 113 = Urvillei.
myriocephala, Clark, Trav. i. 23 = diffusa.
myriocephala, *Sch. Bip. ex Boiss. Fl. Orient.* iii. 682. —Syria.
myriostigma, *Lem. in Illustr. Hortic.* (1856) *Misc.* 88. —Hab.?
myriostoma, *Vis. & Panč. in Mem. Ist. Venet.* xii. (1864) 470.—Orient.
nana, Chesn. ex Steud. Nom. ed. II. i. 320, 318 = amara.
nana, *Desf. Fl. Atlant.* ii. 296. t. 241.—Mont. Atl.
nana, Sieber, ex Boiss. Fl. Orient. iii. 675 = raphanina.
napifolia, *Linn. Sp. Pl.* 916.—Italia; Graecia; Afr. bor.
napulifera, *Rochel, in 'Act. Hungar. Evhony.'* (1834) t. 3.—Europ.
neapolitana, Boiss. Diagn. Ser. II. iii. 72 = nigrescens.
neglecta, Bess. Enum. Pl. Volh. 35 = centauroides.
neglecta, Martr. ex Nym. Consp. 431 = sonchifolia.
neglecta, Schott, ex Spreng. Syst. iii. 400 = orientalis.
nemophila, Jord. ex Nym. Consp. 421 = C. Jacea.
nemoralis, Jord. Pugill. Pl. Nov. 104 = nigra.
nervosa, Griseb. Spicil. Fl. Rumel. ii. 233 = Kerneriana.
nervosa, Reichb. ex Steud. Nom. ed. II. i. 321 = phrygia.
nervosa, *Willd. Enum. Hort. Berol.* 925.—Europ. austr.
nevadensis, *Boiss. & Reut. Diagn.* Ser. II. iii. 71.—Hispan.
nicaeensis, *All. Fl. Pedem.* i. 162. t. 74. f. 1.—Gall.; Ital.
nidulans, *Pomel, Nouv. Mat. Fl. Atl.* 280.—Algeria.
Niederi, *Heldr. in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 380.—Graecia.
nigra, Frivald. ex Boiss. Fl. Orient. iii. 630 = Kerneriana.
nigra, Lam. Encyc. i. 666 = pratensis.
nigra, *Linn. Sp. Pl.* 911.—Europ.
nigrescens, DC. Prod. vi. 571 = nigra.
nigrescens, Gren. & Godr. Fl. Fr. ii. 241 = C. Jacea.
nigrescens, *Willd. Sp. Pl.* iii. 2288.—Europ. med. austr.
nigricans, DC. ex Steud. Nom. ed. II. i. 321 = Endressii.
nigricans, Link, ex Spreng. Syst. iii. 401 = nigra.
Nimrodis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 664.—Mesopot.
nitens, *Bieb. ex Willd. Sp. Pl.* iii. 2305.—Tauria; Reg. Cauc.
nitida, Lam. Fl. Fr. ii. 49 = alba.
nitida, Naldi, ex DC. Prod. vi. 601 = crassifolia.
Noëana, Boiss. Diagn. Sér. II. iii. 84 = iberica.
Novelii ×, *Franch. in Bull. Soc. Bot. Fr.* xxvii. (1880) p. xxi.—Gallia.
nudicaulis, Linn. Syst. ed. X. 1232 = Serratula nudicaulis.

CENTAUREA :—
Nuttallii, Spreng. Syst. iv. Cur. Post. 298 = americana.
obscura, Jord. in F. Schultz, Arch. Fl. Fr. et Allem. 320 = nigra.
ochrolepis, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 149 = axillaris.
ochroleuca, Pushk. ex Willd. Sp. Pl. iii. 2289 = axillaris.
ochrolopha, Costa, Fl. Catal. 140 = leucophaea.
odorata, *Burm. f. Fl. Cap. Prod.* 28.—Afr. austr.
odorata, *C. Koch, in Linnaea*, xxiv. (1851) 417.—Reg. Caucas.
Olivieri, *Pomel, Nouv. Mat. Fl. Atl.* 282.—Algeria.
Olivieriana, *DC. Prod.* vi. 590.—Ins. Naxos.
olympica, *Boiss. Diagn.* Ser. II. iii. 77.—Bithynia.
omphalodes, *Coss. & Dur. in Bull. Soc. Bot. Fr.* iv. (1857) 397, nomen.—Algeria.
onopordifolia, *Boiss. Diagn.* Ser. I. x. 114.—Syria.
orientalis, Baumg. ex Schur, Enum. Pl. Transs. 408 = pseudo-collina.
orientalis, *Linn. Sp. Pl.* 918.—Taur.; Sibir.
ornata, *Willd. Sp. Pl.* iii. 2320.—Hispan.
Orphanidea, *Heldr. & Sart. ex Boiss. Diagn.* Ser. II. iii. 73.—Graecia.
orthoacantha, *Tausch, in Flora*, xii. (1829) i. *Erg.* 44. —Hab.?
ossica, *C. Koch, in Linnaea*, xvii. (1843) 40.—Reg. Caucas.
ovina, Hort. ex Steud. Nom. ed. II. i. 321 = pterolopha.
ovina, Pall. ex Steud. l. c. = diffusa.
ovina, *Pall. ex Willd. Sp. Pl.* iii. 2292.—Taur.; Sibir.
pallescens, Bové, ex DC. Prod. vi. 597 = alexandrina.
pallescens, *Delile, Fl. Egypte*, 278. t. 49.—Aegypt.
pallida, *Frivald. in Flora*, xviii. (1835) 333.—Europ.
pallida, Wall. Cat. n. 2983 = Serratula pallida.
pamphylica, *Boiss. & Heldr. Diagn.* Ser. I. x. 120.—Pamphylia.
paniculata, Auct. Hisp. ex Willk. & Lange, Prod. Fl. Hisp. ii. 157 = castellana.
paniculata, Bieb. Fl. Taur. Cauc. ii. 346 = maculosa.
paniculata, Lag. ex DC. Prod. vi. 583 = caerulescens.
paniculata, *Linn. Sp. Pl.* 912.—Europ. austr.
paniculata, Pett. ex Nym. Consp. 426, 427 = cristata, divergens.
paniculata, Sibth. & Sm. Fl. Graec. Prod. ii. 199 = graeca.
paniculata, Vill. Hist. Pl. Dauph. iii. 53 = leucophaea.
pannosa, *DC. Prod.* vi. 582.—Mons Athos.
Parlatorii, Heldr. in Flora, xxvii. (1844) i. 68 = seq.?
Parlatoris, Heldr. in Ann. Acc. Asp. Nat. i. (1843) 287 = dissecta.
Parolinii, *DC. Prod.* vi. 592.—Mons Ida; As. Min.
parviflora, Bieb. Fl. Taur. Cauc. iii. 594 = diffusa.
parviflora, *Desf. Fl. Atlant.* ii. 301.—Pers.; Afr. bor.
parviflora, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 450 = squarrosa.
parviflora, Lam. Fl. Fr. ii. 32 = aspera.
parviflora, Sm. Fl. Graec. x. 9. t. 912 = diffusa.
patibilcensis, DC. Prod. vi. 593 = melitensis.
patula, *DC. l. c.* 583.—Persia; Cappadoc.
pauciflora, C. Koch, in Linnaea, xxiv. (1851) 433 = aggregata.
pauciloba, *Trautv. in Act. Hort. Petrop.* ii. (1873) 474.—Daghestania.
pectinata, Auct. Ross. ex Boiss. Fl. Orient. iii. 632 = trichocephala.
pectinata, *Linn. Sp. Pl.* ed. II. 1287.—Gall. austr.
pectinata, Siev. in Pall. N. Nord. Beitr. vii. 323 = calva.
Pelia, *DC. Prod.* vi. 586.—Graecia.
penicillata, Delile, Fl. Aeg. Illustr. 74 = trichacantha.
peregrina, *Linn. Sp. Pl.* 918.—Europ. austr.
pergamacea, *DC. Prod.* vi. 569.—Cappadoc.
Perrottettii, *DC. l. c.* 598.—Afr. trop.
persica, *Boiss. Diagn.* Ser. I. vi. 132.—Persia.
Pestalotii, De Not. ex Cesati, in Bibl. Ital. xci. (1838) 348 = alba.
Pestalozzae, *Boiss. Diagn.* Ser. I. x. 118.—Lycia.
petrocaula, *Trautv. in Act. Hort. Petrop.* ii. (1873) 552.—Armenia.
Petteri, Reichb. f. Ic. Fl. Germ. xv. 36. t. 783 = divergens.
peucedanifolia, *Boiss. & Orph. in Boiss. Fl. Orient.* iii. 647.—Macedon.
phaeolepis, *Coss. Notes Pl. Crit.* 138.—Mauritania.
phaocephala, *Wallr. Beitr. Fl. Hercyn.* 270.—Europ.
Pharaonis, *Pomel, Nouv. Mat. Fl. Atl.* 281.—Algeria.

CENTAUREA :—
phlomoides, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 673.—Persia.
phrygia, Jacq. Hort. Vindob. 167 = austriaca.
phrygia, Lapeyr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 164 = nigra.
phrygia, *Linn. Sp. Pl.* 910.—Europ.
phrygia, Lour. Fl. Cochinch. ii. 681 = cochinchinensis.
phyllocephala, *Boiss. Diagn.* Ser. I. vi. 134.—Afghan.
Pichleri, *Boiss. Fl. Orient.* iii. 638.—Bithynia.
Pieris, *Pall. Tabl. Taur.* 58; *Willd. Sp. Pl.* iii. 2302. —Reg. Casp.
Pinardi, *Boiss. Diagn.* Ser. I. iv. 17.—Caria.
pindicola, *Griseb. Reise Rumel.* ii. 164.—Maced.
pinnatifida, Schur, Sert. Transs. 35 = axillaris.
pinnatispina, *Turra, Farset. Nov. Gen.* 13.—Sicilia.
pityocephala, Brot. Fl. Lusit. i. 369 = Leuzea conifera.
platylepis, Peterm. in Flora, xxvii. (1844) 475 = C. Jacea.
plumosa, Hort. ex Vilm. Fl. Pl. Terre, ed. III. 245 = gymnocarpa.
plumosa, *Kern. in Oestr. Bot. Zeitschr.* xxii. (1872) 44.—Europ. austr.
podospermifolia, *Losc. & Pard. Ser. Inconf.* 58.—Hispan.
polyacantha, Spreng. Syst. iii. 406 = sonchifolia.
polyacantha, *Willd. Sp. Pl.* iii. 2312.—Hispan.
polycephala, Jord. Obs. Pl. Crit. v. (1847) 67 = paniculata.
polyclada, *DC. Prod.* vi. 584.—Graecia.
polymorpha, *Lag. Gen. et Sp. Nov.* 32.—Hispan.
polyphylla, Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb. ii. (1837) 311 = ossica.
polyphylla, *Pomel, Nouv. Mat. Fl. Atl.* 281.—Algeria.
polypodifolia, *Boiss. Diagn.* Ser. I. vi. 126.—Persia.
Postii, *Boiss. Fl. Orient.* iii. 688.—Syria.
Pottii, [Spreng.] Bot. Gart. Halle, 23 = sicula, sulphurea.
Pourretiana, *Timb. & Theven. in Mém. Acad. Toul.* Sér. VII. vi. (1874) 645.—Gallia.
Pouzini, DC. Cat. Hort. Monsp. 91 = C. Calcitrapa.
pracalta, *Boiss. & Bal. Diagn.* Ser. II. v. 112.—Cilicia.
praecox, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 438.—Afr. trop.
praetermissa, Martr. in Soc. Sc. Tarn et Gar. (1852); Pl. Crit. Tarn, 382 = aspera.
praetoria, Reichb. f. Ic. Fl. Germ. xv. 34. t. 856 = leucophaea.
pratensis, *Hornem. Hort. Hafn. Suppl.* 104.—Hungaria.
pratensis, Salisb. Prod. 208 = C. Jacea.
pratensis, Thuill. Fl. Paris. ed. II. 444 = nigrescens.
Preauxiana, Webb & Berth. Phyt. Canar. ii. 350 = Atractylis Preauxiana.
princeps, Boiss. & Heldr. Diagn. Ser. II. vi. 111 = alba.
procera, Heynh. Nom. ii. 131 lapsu = Cephalaria procera.
procumbens, *Balb. Misc. Alt.* 31. t. 1.—Pedemont.
procumbens, Habl. in S. G. Gmel. It. iv. 167 = decalbata.
procumbens, Jord. Obs. v. 57 = Jordaniana.
procurrens, DC. Prod. vi. 595 = sinaica.
procurrens, *Sieber, ex Spreng. Syst.* iii. 407.—Palaestin.; Arab.
prolifera, Delile, ex Godr. in Mém. Acad. Montp. (Sect. Médic.) i. (1853) 435 = Delilei.
prolifera, Vent. Jard. Cels, 16 = glomerata.
Prolongi, *Boiss. ex DC. Prod.* vii. 303.—Hispan.
prostrata, *Coss. Notes Pl. Crit.* 115.—Hispan.
prostrata, Huet, ex Nym. Consp. 425 = cinerea.
provincialis, Boiss. ex Lindem. in Bull. Soc. Nat. Mosc. lv. (1880) ii. 181 = jurineaefolia?
pseudo-behen, *Boiss. Diagn.* Ser. I. vi. 126.—Cappadoc.
pseudo-collina, *Schur, Enum. Pl. Transs.* 408.—Transsylv.
pseudo-crupina, *Mutel, in Duch. Rev. Bot.* i. (1845-46) 400.—Morea.
pseudo-philostigus, Godr. in Mém. Acad. Stanislas, Sér. III. [xvii] 1853 (1854) 405 = dimorpha.
pseudo-phrygia, C. A. Mey. in Bull. Phys.-Math. Acad. Petersb. vi. (1848) 134 = phrygia.
pseudo-scabiosa, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 131 = C. Scabiosa.
pseudo-uniflora, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 85 = nervosa.
psilacantha, *Boiss. & Heldr. Diagn.* Ser. II. iii. 82.—Graecia.

CENTAUREA :—

pterocaula, *Trautv. in Act. Hort. Petrop.* ii. (1873) 552.—Armenia.
pterocaulos, *Pomel, Nouv. Mat. Fl. Atl.* 29.—Algeria.
pterocephala, Hort. ex DC. Prod. vi. 597 = trichacantha.
pterodonta, *Pomel, Nouv. Mat. Fl. Atl.* 28.—Algeria.
pterolopha, *DC. Prod.* vi. 572.—Hab.?
pubescens, Willd. Sp. Pl. iii. 2322 = incana.
pubigera, Pers. Syn. ii. 488 = Volutarella muricata.
pulchella, *Ledeb. Ic. Pl. Alt.* 1. t. 93.—Persia; Caucas.
pulcherrima, Wight, ex DC. Prod. vi. 579 = pulchra.
pulcherrima, *Willd. Sp. Pl.* iii. 2298.—Armenia; Caucas.
pulchra, *DC. Prod.* vi. 578.—Ind. or.
pullata, *Linn. Sp. Pl.* 911.—Europ. austr.
pumila, *Linn. Cent.* i. 30; *Amoen. Acad.* iv. 292.—Aegypt.; Graecia.
pumila, Sprun. ex Nym. Consp. 430 = hellenica.
pumila, Urv. in Mem. Soc. Linn. Par. i. (1822) 374 = raphanina.
Pumilio, Hill, Veg. Syst. iv. 26 = pumila?
punctata, Dufour, in Ann. Sc. Nat. Sér. I. xxiii. (1831) 159 = hyssopifolia.
punctata, Vis. in Flora, xix. (1829) 23 = glaberrima.
pungens, *Pomel, Nouv. Mat. Fl. Atl.* 29.—Algeria.
pusilla, *Thuill. ex Pers. Syn.* ii. 482.—Aegypt.
pygmaea, *Benth. & Hook. f. Gen.* ii. 482.—Syria.
pygmaea, Chesn. ex DC. Prod. vi. 571 = amara.
pygmaea, Hoffm. Hort. Mosc. (1808) n. 783 = depressa.
pyracantha, *Boiss. Fl. Orient.* iii. 685.—Tauria.
pyrenaica, Spreng. Fl. Hal. Mant. 53 = variegata.
pyrrhoblephara, *Boiss. Diagn.* Ser. II. iii. 66.—Armenia.
Rabenhorstiana, Sch. Bip. ex Nym. Consp. 426 = cristata.
Raddeana, *Trautv. in Act. Hort. Petrop.* ii. (1873) 475.—Armenia.
radiata, Linn. Syst. ed. XII. 656 = Serratula xeranthemoides.
ragusina, Clem. ex Boiss. Voy. Bot. Espagne, ii. 346 = Clementei.
ragusina, *Linn. Sp. Pl.* 912.—Dalmat.; Creta.
ragusina, Sm. Fl. Graec. x. 2. t. 903 = argentea.
ramosa, *Vukot.—Cf. Oestr. Bot. Zeitschr.* xxx. (1880) 163.—Europ.
ramosissima, Tausch, in Flora, xii. (1829) Erg. 45 = nitens.
raphanifolia, Salzm. ex DC. Prod. vi. 591 = diluta.
raphanina, *Sibth. & Sm. Fl. Graec.* t. 917.—Creta.
raphanina, Sieber, ex Steud. Nom. ed. II. i. 322 = nana.
raphanina, Sprun. ex Nym. Consp. 430 = hellenica.
raphanina, Urv. in Mem. Soc. Linn. Par. i. (1822) 372 = Urvillei.
recta, *Krock. Fl. Sil. Suppl.* ii. 333.—Europ.
recurvata, C. Koch, in Linnaea, xxiv. (1851) 428 = carduiformis.
reflexa, *Lam. Encyc.* i. 675.—Armenia; Caucas.
regia, *Boiss. Diagn.* Ser. I. vi. 135.—Assyria.
Reichenbachii, *DC. Prod.* vi. 583.—Europ.?
reichenbachioides, Schur, Enum. Pl. Transs. 409 = Reichenbachii.
repens, *Linn. Sp. Pl.* ed. II. 1233.—Oriens.
resupinata, *Coss. Notes Pl. Crit.* 114.—Hispan.
reticulata, Bieb. ex Willd. Sp. Pl. iii. 2287 = coronopifolia.
Reuteri, Reichb. f. Ic. Fl. Germ. xv. 33. t. 780 = paniculata.
Reuteriana, *Boiss. Diagn.* Ser. I. iv. 18.—Graecia.
rhaetica, Moritz. in Denkschr. Allg. Schw. Ges. Natur. iii. (Pfl. Graub. 81. t. 3) = phrygia.
rhapontica, Wulf. in Jacq. Coll. i. 223 = rhaponticum.
rhaponticoides, *Benth. & Hook. f. Gen.* ii. 479.—Hispan.
Rhaponticum, *Linn. Sp. Pl.* 915.—Europ.
rhenana, Bor. ex Nym. Consp. 426 = maculosa.
rhizantha, C. A. Mey. Verz. Pfl. Cauc. 64 = sessilis.
rhizantha, Tchihat. Asie Min. Bot. ii. 320 = armena.
rhizocalathium, *C. Koch, in Linnaea*, xxiv. (1851) 417.—Armenia.
rhizocephala, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 438.—Afr. trop.
rhizocephala, *Trautv. in Act. Hort. Petrop.* ii. (1873) 476.—Armenia.
rhutenica, Baumg. Enum. Stirp. Transs. iii. 70 = Serratula coronata.
rigens, *Lag. Gen. et Sp. Nov.* 32.—Hispan.

CENTAUREA :—

rigescens, *Hornem. Hort. Hafn.* ii. 852.—Hab.?
rigida, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 263.—Syria.
rigida, Willd. Enum. Hort. Berol. Suppl. 61 = rigescens.
rigidifolia, Bess. ex DC. Prod. vi. 587 = orientalis.
rigidula, Jord. Obs. Pl. Crit. v. (1847) 69 = paniculata.
riparia, *A. Cunn. ex DC. Prod.* vi. 602.—N. Cambria.
rivularis, Brot. Fl. Lusit. i. 367 = nigra.
romana, *Linn. Sp. Pl.* 916.—Campania.
Ropalon, *Pomel, Nouv. Mat. Fl. Atl.* 25.—Algeria.
rosea, Steud. Nom. ed. I. 175 = trinervia.
rubescens, *Bess. ex DC. Prod.* vi. 587.—Tauria.
rufescens, Jord. Pugill. 107 = pectinata.
rufescens, Noul. ex Nym. Consp. 422, partim = nigra.
rumelica, *Boiss. Diagn.* Ser. II. iii. 78.—Rumelia.
rupestris, Kit. ex Steud. Nom. ed. II. i. 322 = dissecta.
rupestris, *Linn. Sp. Pl.* ed. II. 1298.—Austr.; Ital.
rupicola, Bad. ex Nym. Consp. 426 = aplolepis.
rupicola, *Pomel, Nouv. Mat. Fl. Atl.* 31.—Algeria.
ruscinonensis, Boiss. Diagn. Ser. II. iii. 70 = C. Jacea?
ruthenica, Baumg. ex Nym. Consp. 417 = *rhutenica*.
ruthenica, *Lam. Encyc.* i. 663.—Transsylv.; Caucas.; Sibir.
rutifolia, Boiss. Fl. Orient. iii. 642 = pannosa.
rutifolia, *Sibth. & Sm. Prod. Fl. Graec.* ii. 205. t, 916.—Byzant.
sabulosa, Ledeb. ex Spreng. Syst. iii. 405 = diffusa.
Sadleriana, *Janka, in Termész. Füzetek*, ii. (1878) 142.—Europ.
Saharae, *Pomel, Nouv. Mat. Fl. Atl.* 30.—Afr. bor.
salicifolia, Bess. ex Nym. Consp. 422 = austriaca.
salicifolia, Bieb. ex Willd. Sp. Pl. iii. 2283 = nigra.
salina, *Benth. & Hook. f. Gen.* ii. 479.—Sibir.
Salisiana ×, *Bruegg. in Jahresb. Naturf. Ges. Graub.* II. xxiii–xxiv. (1880) 110.—Europ.
salmantica, *Linn. Sp. Pl.* 918.—Europ.; Afr. bor.
salonitana, *Vis. in Flora*, xii. (1829) 1. *Erg.* 23.—Europ. or.
sanguinea, *Spreng. Pugill.* i. 60.—Hab.?
saxatilis, Bertol. ex Nym. Consp. 429 = rupestris.
saxatilis, *C. Koch, in Linnaea*, xxiv. (1851) 419.—Creta.
saxicola, *Lag. Gen. et Sp. Nov.* 32.—Hispan.
Scabiosa, Asso, Syn. 126 = cephalariaefolia.
Scabiosa, *Linn. Sp. Pl.* 918.—Europ.
Scabiosa, Sadl. Fl. Pesth. ed. II. 409 = Sadleriana.
scabiosaeformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Scabiosa.
scabiosifolia, St. Lag. l. c. viii. (1881) 203 = praec.
scabra, Presl, Fl. Sicul. p. xxx = sicula.
schemnitzensis, Hort. ex Steud. Nom. ed. II. i. 322 = nigrescens.
Schimperi, *DC. Prod.* vi. 596.—Arab.
schizolepis, *Trautv. in Act. Hort. Petrop.* ii. (1873) 553.—Armenia.
Schmidtiana, *Tausch, in Flora*. xii. (1829) 1. *Erg.* 44.—Hab.?
Schousboei, *Lange, in Kjoeb. Vidensk. Meddel.* (1861) 85 (*Pugill.* 135).—Hispan.
Schouwii, DC. Prod. vi. 593 = sicula.
Schwarzenbergiana, Schur, in Verh. Siebenb. Ver. Naturw. vi. (1855) 4 = ruthenica.
sciaphila, Vukot. in Rad Jugos. Akad. Zagreb. lviii. (1881) 150 = nigrescens.
sclerolepis, *Boiss. Diagn.* Ser. I. vi. 136.—Mesopot.
scoparia, *Sieber, ex DC. Prod.* vi. 596.—Palaest.; Aegypt.
Scopolii, Vest, in Flora, xii. (1829) 1. Erg. 31 = cristata.
scopulorum, *Boiss. & Heldr. Diagn.* Ser. I. x. 115.—Graecia.
scorpiurifolia, Dufour, in Ann. Sc. Nat. xxiii. (1831) 163 = aspera.
segetalis, Salisb. Prod. 207 = C. Cyanus.
semidecurrens, Jord. Obs. Pl. Crit. v. (1847) 52 = axillaris.
sempervirens, *Linn. Sp. Pl.* 913.—Lusit.; Ital.
senegalensis, *DC. Prod.* vi. 598.—Afr. trop.
Seridis, *Linn. Sp. Pl.* 915.—Europ. austr.
Seridis, Loisel. Fl. Gall. ed. II. ii. 212 = aspera.
serotina, Bor. Fl. Cent. Fr. ed. III. ii. 350 = nigrescens.
serrata, Kitt. Taschenb. ed. II. 563 = montana.
serratuloides, *Benth. & Hook. f. Gen.* ii. 479.—As. Min.

CENTAUREA :—

serratuloides, Georgi, Bemerk. Reise Russ. Reich. i. 231 = salina.
serratuloides, Krock. Fl. Sil. ii. 11. 484 (delenda).—Silesia.
sessiliflora, Lam. Fl. Fr. ii. 35 = melitensis.
sessiliflora, Linn. Sp. Pl. 912 = nigra.
sessilifolia, Hill, Veg. Syst. iv. 32 = praec.?
sessilis, *Willd. Sp. Pl.* iii. 2300.—Armenia.
setifera, Lag. Gen. et Sp. Nov. 32 = nitens.
setosa, *Cav. in Anal. Ciencias Nat.* iv. (1801) 96.—Marocco.
seusana, Benth. Cat. Pl. Pyr. 68 = montana.
seusana, Chaix, in Vill. Hist. Pl. Dauph. iii. 52 = variegata, axillaris.
sibirica, Bieb. Fl. Taur. Cauc. ii. 348 = Marschalliana.
sibirica, *Linn. Sp. Pl.* 913.—Sibiria.
sibirica, Willd. ex Ledeb. Fl. Ross. ii. 695 = declinata.
sicula, Lam. ex Willk. & Lange, Prod. Fl. Hisp. ii. 145 = nicaeënsis.
sicula, Leyss. ex DC. Prod. vi. 594 = solstitialis.
sicula, *Linn. Sp. Pl.* 918.—Sicilia.
simplex, *Cav. in Anal. Ciencias Nat.* iv. (1801) 95.—Afr. bor.
simplex, Murr. ex Steud. Nom. ed. II. i. 322 = C. Jacea.
simplicicaulis, *Boiss. & Huet, Diagn.* Ser. II. iii. 67.—Armenia.
sinaica, *DC. Prod.* vi. 592.—Arab.
singarensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 654.—Assyria.
Smyrneae, Boiss. Diagn. Ser. I. iv. 20, in textu = Urvillei.
solstitialis, Asso, Syn. 127 = melitensis.
solstitialis, *Linn. Sp. Pl.* 917.—Europ.
somalensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 438.—Afr. trop.
sonchifolia, Boiss. Voy. Bot. Espagne, ii. 352 = malacitana.
sonchifolia, *Linn. Sp. Pl.* 915.—Europ. austr.
sonchifolia, Welw. ex Nym. Consp. 432 = sphaerocephala.
sordida, Graf, ex DC. Prod. vi. 587 = Grafiana.
sordida, Salisb. Prod. 207 = sempervirens.
sordida, *Willd. Sp. Pl.* iii. 2323.—Austria.
Spachii, *Sch. Bip. ex Willk. & Lange, Prod. Fl. Hisp.* ii. 154.—Hispan.
Sparmanni, *DC. Prod.* vi. 598.—Africa.
spathulata, Tenore, Prod. Fl. Nap. p. li = nigra?
spathulata, Zeraph. Fl. Melit. i. 11 = crassifolia.
speciosa, *Boiss. Diagn.* Ser. I. x. 116.—Palaestina.
speciosa, Mor. ex Nym. Consp. 428 = filiformis.
sphaerocephala, J. Gay, ex Dur. in Duch. Rev. Bot. ii. (1846–47) 429 = Fontanesii.
sphaerocephala, *Linn. Sp. Pl.* 916.—Europ. med.
sphaerocephala, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1878) 531 = fragilis.
spicata, *Boiss. Diagn.* Ser. I. x. 113.—Syria.
Spina-badia, Bubani, ex Timb. in Mém. Acad. Toulouse, Sér. VIII. i. 11. (1879) 187 = coerulescens.
spinosa, Griseb. Spicil. Fl. Rumel. ii. 241 = diffusa.
spinosa, *Linn. Sp. Pl.* 912.—Creta; Oriens.
spinulosa, Rochel, Pl. Ban. Rar. 76. t. 36 = C. Scabiosa.
splendens, *Linn. Sp. Pl.* 914.—Hispan.; Ital.
splendens, Tenore, Syll. Add. iii. 628 = alba.
Spruneri, *Boiss. & Heldr. Diagn.* Ser. I. vi. 132.—Graecia.
Spruneri, Boiss. & Orph. ex Boiss. Fl. Orient. iii. 662 = Guicciardii.
spuria ×, *Kern. in Oestr. Bot. Zeitschr.* xxii. (1872) 51.—Austria.
squarrosa, Roth, Catalecta, fasc. ii. 118 = virgata.
stenolepis, Kern. in Oestr. Bot. Zeitschr. xxii. (1872) 45 = austriaca.
stenophylla, Dufour, in Ann. Sc. Nat. Sér. I. xxiii. (1831) 162 = aspera.
stenophylla, *Walp. Ann.* i. 448.—Persia.
stereophylla, *Bess. Enum. Pl. Volh.* 35.—Podol.
stereophylla, Griseb. Spicil. Fl. Rumel. ii. 237 = C. Scabiosa.
sterilis, *Stev. in Bull. Soc. Nat. Mosc.* xxix. (1856) 11. 390.—Tauria.
Steveni, *Bieb. Fl. Taur. Cauc.* ii. 356.—Reg. Caucas.
Steveni, Kotschy, ex Boiss. Fl. Orient. iii. 693 = cheiracantha.
Stoebe, Ledeb. Fl. Alt. iv. 49 = calva.
Stoebe, *Linn. Sp. Pl.* 913.—Austria; Gallia.

CENTAUREA:—
Stoebe, Tausch, in Flora, xii. (1829) i. Erg. 42 = diffusa.
Stoebe, Tenore, Fl. Nap. ii. 264 = subtilis.
stolensis, Panč. ex Nym. Consp. 428 = orientalis.
straminea, Willd. Sp. Pl. iii. 2312 = glomerata.
strepens, Hoffmgg. & Link, Fl. Port. ii. 236 = splendens?
stricta, Waldst. & Kit. Pl. Rar. Hung. ii. 194 = axillaris.
strictissima, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 130 = Serratula coriacea.
strobilacea, Scop. Delic. Insub. i. 38 = alba.
suaveolens, Willd. Sp. Pl. iii. 2279 = moschata.
subacaulis, Ledeb. Fl. Ross. ii. 709 = Urvillei.
subalbida, Jord. Pugill. 109 = maculosa.
subalpina, Schur, Enum. Pl. Transs. 407 = atropurpurea.
subciliaris, *Boiss. & Heldr. ex Boiss. Fl. Orient.* iii. 627.—Cephalon.
sublanata, *Boiss. l. c.* 645.—Graecia.
sublanuginosa, Schur, Enum. Pl. Transs. 899, Index = nervosa.
subspinosa, Steud. Nom. ed. II. i. 322 = diluta.
subtilis, *Bertol. Fl. Ital.* ix. 451.—Ital.
sulphurea, Lag. Gen. et Sp. Nov. 32 = dissecta, orientalis.
sulphurea, *Willd. Enum. Hort. Berol.* 930.—Sicilia.
sumensis, *Kalen. in Bull. Acad. Soc. Nat. Mosc.* xviii. (1845) i. 238.—Reg. Caucas.
supina, Jord. Pugill. 108 = pectinata.
sylvatica, Pourr. in Mém. Acad. Toul. iii. (1788) 310 = C. Scabiosa.
Szovitsiana, *Boiss. Fl. Orient.* iii. 683.—Armenia; Anatol.
tagana, *Brot. Fl. Lusit.* i. 369.—Lusit.; Afr. bor.
taraxacifolia, Boiss. in Tchihat. Asie Min. Bot. ii. 321 = rhizocalathium.
taraxacifolia, D. Don, Prod. Fl. Nep. 167 = Saussurea criostemon.
tatarica, Linn. f. Suppl. 383 = orientalis.
tatarica, Pall. Ind. Taur. 58 = centauroides.
tauromenitana, *Guss. Fl. Sicil. Syn.* ii. 512.—Sicil.
Tauscheri, Kern. in Oestr. Bot. Zeitschr. xxii. (1872) 119 = arenaria.
Tchihatcheffii, *Fisch. & Mey. in Ann. Sc. Nat.* Sér. IV. (1854) i. 31.—Lycaonia.
Teidis, Steud. Nom. ed. II. i. 323 = arguta.
tenella, Hort. ex Spreng. Syst. iii. 401 = salmantica.
Tenoreana, Willk. ex Nym. Consp. 420 = incana.
tenuiflora, *DC. Prod.* vi. 584.—Tauria.
tenuiflora, Fisch. Hort. Gorenk. ed. II. (1812) 36 = salmantica.
tenuifolia, *Dufour, in Ann. Sc. Gén. Phys.* ii. (1820) 303.—Hispan.
tenuifolia, Hort. ex Steud. Nom. ed. II. i. 323 = salmantica.
tenuifolia, Schleich. ex Steud. l. c. = C. Scabiosa.
tenuifolia, Salisb. Prod. 207 = Crupina vulgaris.
tenuiramea, *Delile, ex Herd. in Ind. Sem. Hort. Petrop.* (1860) 37.—Hab.?
tenuisecta, Jord. Pugill. 110 = maculosa.
Teydis, C. Sm. ex Buch, in Berl. Acad. Abh. 1816–17 (1819) 369 = arguta.
theiantha, [Spreng.] Bot. Gart. Halle, 24 = orientalis.
Thirkei, *Sch. Bip. in Linnaea*, xix. (1847) 314.—Phrygia.
Thomasiana, *Dalla Torre, Atl. Alpenfl.* 141.—Europ.
thrinciaefolia, *DC. Prod.* vi. 590.—Cappadoc.
Timbali, Martr. Pl. Crit. Taur. 382 = amara.
tinctoria, [Clairv.] Man. Herb. 253 = Serratula tinctoria.
tingitana, Linn. Sp. Pl. ed. II. 1300 = Carduncellus caeruleus.
toletana, *Boiss. & Reut. Diagn. Pl. Nov. Hisp.* 18.—Hispan.
tomentosa, Salisb. Prod. 207 = ragusina.
Torreana, Tenore, Ind. Sem. Hort. Neap. (1829) 15 = C. Calcitrapa.
tougourensis, *Boiss. & Reut. Diagn.* Ser. II. iii. 76.—Algeria.
Tournefortii, *Walp. Ann.* i. 448.—Armenia.
transalpina, Schleich. Cat. Pl. Helv. ed. III. 11 = nigrescens.
trapezuntina, Boiss. Fl. Orient. 630 = nigrescens.
trichacantha, Willd. ex Spreng. Syst. iii. 404 = C. Calcitrapa.
trichocephala, *Bieb. ex Willd. Sp. Pl.* iii. 2286.—Tauria.

CENTAUREA:—
trifurcata, *Pomel, Nouv. Mat. Fl. Atl.* 27.—Afr. bor.
trinervia, *Stev. ex Willd. Sp. Pl.* iii. 2301.—Tauria; Cauc.
triniaefolia, Heuff. in Oestr. Bot. Zeitschr. viii. (1858) 27 = maculosa.
Triumfetti, All. Fl. Pedem. i. 158 = variegata.
tuberosa, Vis. Fl. Dalmat. ii. 33. t. 12. f. 2 = napulifera.
tubulosa, Chab. ex Nym. Consp. 421 = nigrescens.
turkestanica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 325.—Turkestan.
Tweedieii, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 110.—Bonaria.
ucrainica, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 497, nomen.—Ross. austr.
uliginosa, *Brot. Fl. Lusit.* i. 368.—Lusitan.
umbrosa, *Huet, ex Reut. Ind. Sem. Hort. Genev.* (1856) 4.—*Cf. Linnaea*, xxix. (1857–58) 719.—Sicilia.
uniflora, *Linn. Mant.* i. 118.—Europ. austr.
uniflora, Pall. Reise, iii. 252 = Serratula glauca.
Urvillei, *DC. Prod.* vi. 592.—Ins. Samos.
ustulata, Bess. Enum. Pl. Volh. 108 = Bieberstеinii.
ustulata, *DC. Prod.* vi. 576.—Persia.
vallesiaca, Jord. Pugill. 111 = paniculata.
vallesiaca, *Reichb. f. Ic. Fl. Germ.* xv. 33. t. 783.—Europ.
variabilis, Bartl. ex Steud. Nom. ed. II. i. 323 = rupestris.
varians, *A. Rich. Tent. Fl. Abyss.* i. 453.—Abyss.
variegata, *Lam. Encyc.* i. 668.—Europ. austr.
variifolia, Loisel. Not. 130 = C. Scabiosa.
venetum, J. F. Gmel. Syst. 1267, sphalm. = C. Verutum.
verbascifolia, *Vahl, Symb. Bot.* i. 75.—Arab.
Verutum, *Linn. Cent.* i. 30; *Amoen. Acad.* iv. 292.—As. Min.
veseritensis, *Boiss. Diagn.* Ser. II. iii. 76.—Algeria.
Villarsii, *Mutel, Fl. Dauph.* ii. 254.—Gallia.
viminea, Less. in Linnaea, vi. (1831) 83 = pulchella.
viretorum, Jord. ex Nym. Consp. 421 = C. Jacea.
virgata, Bess. Enum. Pl. Volh. 35 = Besseriana.
virgata, Cav. Ic. iii. 16 = caerulescens.
virgata, *Lam. Encyc.* i. 670.—Armenia.
virgata, Portenschl. ex Nym. Consp. 426 = glaberrima.
vochinensis, Bernh. ex Reichb. Ic. Fl. Germ. xv. 15 = nigrescens.
vulgaris, Pers. Syn. ii. 488 = Crupina vulgaris.
vulnerariaefolia, *Pomel, Nouv. Mat. Fl. Atl.* 26.—Afr. bor.
Wallichiana, *Less. in Linnaea* vi. (1831) 84.—Ind. or.
Webbiana, *Sch. Bip. in Webb & Berth. Phyt. Canar.* ii. 353.—Ins. Canar.
Weldeniana, Reichb. Ic. Fl. Germ. xv. 12. t. 753 = amara.
Wiedemanniana, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1835) 32.—Anatolia.
Willkommii, *Sch. Bip. ex Willk. in Linnaea*, xxv. (1852) 36.—Hispan.
wolgensis, DC. Prod. vi. 581 = arenaria.
xanthina, Boiss. & Heldr. in Boiss. Fl. Orient. iii. 650 = Pelia.
xanthina, Spreng. Syst. iii. 407 = fuscata.
xanthocephalus, *Benth. & Hook. f. Gen.* ii. 480.—Armen.
Zanonii, Sebast. & Maur. Fl. Rom. 298 = romana.
Zuccariniana, *DC. Prod.* vi. 574.—Graecia.

CENTAURELLA, Delarb. Fl. Auv. ed. II. 28 (1800) = **Erythraea**, Borckh. (Gentian.).
dichotoma, Delarb. l. c. 29 = E. Centaurium.

CENTAURELLA, Michx. Fl. Bor. Am. i. 97. t. 12 (1803) = **Bartonia**, Muhl. (Gentian.).
aestivalis, Pursh, Fl. Am. Sept. i. 100 = B. verna.
autumnalis, Pursh, l. c. = B. tenella.
Moseri, Steud. & Hochst. in Griseb. Gen. et Sp. Gentian. 308 = B. tenella.
paniculata, *Michx. Fl. Bor. Am.* i. 98. t. 12. f. 1 = B. tenella.
verna, Michx. l. c. 98 = B. verna.
vernalis, Pursh, Fl. Am. Sept. i. 99 = B. verna.

CENTAUREUM, Rupp. Fl. Jen. ed. Hall. 22 (1745) = **Erythraea**, Borckh. (Gentian.).

CENTAURIA, Linn. Gen. ed. I. 263 (1737) = **Centaurea**, Linn. (Compos.).

CENTAURIDIUM, Torr. & Gray, Fl. N. Am. ii. 246 (1841) = **Xanthisma**, DC. (Compos.).
Drummondii, Torr. & Gray. l. c. = X. texanum.

CENTAURIUM, Borckh. in Roem. Arch. i. (1796) 29 = **Canscora**, Lam. (Gentian.).
malabaricum, Borckh. l. c. = C. sessiliflora.

CENTAURIUM, Gilib. Fl. Lituan. i. 35 (1781); Moench, Meth. 449 (1794) = **Erythraea**, Borkh. (Gentian.).
Erythraea, Rafn, Danm. og Holst. Fl. ii. 75 = E. Centaurium.
inapertum, Rafn, l. c. 77.—Europ.
minus, Moench, Meth. 449 = E. Centaurium.
umbellatum, Gilib. Fl. Lituan. i. 35 = E. Centaurium.
vulgare, Rafn, Danm. og Holst. Fl. ii. 73 = E. Centaurium.

CENTAURIUM, Pers. Syn. i. 137 (1805) = CENTAURELLA, Michx. = **Bartonia**, Muhl. (Gentian.).
autumnale, Pers. l. c. = B. tenella.
vernum, Pers. l. c. = B. verna.

CENTAURIUM, Tourn. ex Linn. Syst. ed. I (1735) = **Centaurea**, Linn. (Compos.).
africanum, Cass. in Dict. Sc. Nat. lviii. 10 = Centaurea africana.
alpina, Pomel, Nouv. Mat. Fl. Atl. 32.—Algeria.
moschatum, Cass. in Dict. Sc. Nat. vii. 378 = Centaurea moschata.
officinale, Cass. l. c. = Centaurea Centaurium.
ruthenicum, Cass. l. c. lviii. 10 = Centaurea rhuteniea.
suaveolens, Cass. l. c. vii. 379 = Centaurea moschata.
taganum, Pomel, Nouv. Mat. Fl. Atl. 32.—Algeria.
tatarica, Pomel, l. c.—Algeria.

CENTAUROPSIS, Boj. ex DC. Prod. v. 93 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 231.
frutieosa, *Boj. l. c.*—Madag.
lanuginosa, *Boj. l. c.*—Madag.
Rutenbergiana, *Vatke, in Bremen Abh.* ix. (1885) 119.—Madag.

CENTELLA, Linn. Pl. Rar. Afr. 28 (1760); Amoen. Acad. vi. 112 (1764) = **Hydrocotyle**, Linn. (Umbellif.).
asiatica, Urban, in Mart. Fl. Bras. xi. i. 287 = H. asiatica.
cuneifolia, F. Muell. in Hook. Kew Journ. vii. (1855) 379. t. 12 = Azorella cuneifolia.
glabra, DC. Prod. iv. 69, sphalm. = seq.
glabrata, Linn. Pl. Rar. Afr. 28; Amoen. Acad. vi. 112 = H. Centella.
renifolia, Urban, in Mart. Fl. Bras. xi. i. 286, in obs. = Micropleura renifolia.
villosa, Linn. Pl. Rar. Afr. 28; Amoen. Acad. vi. 112 = H. villosa.

CENTEMA, Hook. f. in Benth. & Hook. f. Gen. iii. 31 (1880). *AMARANTACEAE*, Benth. & Hook. f. l. c.
angolensis, *Hook. f. l. c.*—Afr. trop.
Kirkii, *Hook. f. l. c.*—Afr. trop.

CENTHRISCUS, Spreng. ex Steud. Nom. ed. I. 176, 55 (1821) = **Anthriscus**, Bernh. (Umbellif.).
fumarioides, Spreng. l. c. = A. fumarioides.
nemorosa, Spreng. l. c. = A. nemorosa.

CENTINODIA, Reichb. Handb. 236 (1837) = **Polygonum**, Tourn.
agrestina, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 146 = P. aviculare.
arenaria, Fourr. l. c. = P. arenarium.
avicularis, Fourr. l. c. = P. aviculare.
Bellardi, Fourr. l. c. = P. Bellardi.
humifusa, Fourr. l. c. = P. aviculare.
maritima, Fourr. l. c. = P. maritimum.
microsperma, Fourr. l. c. = P. aviculare.
rurivaga, Fourr. l. c. = P. aviculare.

CENTIPEDA, Lour. Fl. Cochinch. 492 (1790). *COMPOSITAE*, Benth. & Hook. f. ii. 430.
MYRIOGYNE, Less. in Linnaea, vi. (1831) 219.
capensis, Less. Syn. Comp. 201 = Dicnrocephala latifolia.
Cunninghami, *A. Br. & Aschers. Ind. Sem. Hort. Berol.* 1867 *App.* 6; *F. Muell. Fragm.* viii. 143.—Austral.
minima, A. Br. & Aschers. l. c. = orbicularis.

CENTIPEDA :—
minuta, C. B. Clarke, Compos. Ind. 151 = orbicularis.
orbicularis, C. B. Clarke, l. c. = Sphaeromorphaea Russeliana.
orbicularis, *Lour. Fl. Cochinch.* ii. 493.—As. et Austral. trop.; Ins. Pacif.
racemosa, *F. Muell. Census*, 84.—Austral.
thespidioides, *F. Muell. Fragm.* viii. 143.—Austral.

CENTOPODIUM, Burch. Trav. i. 340 (1824) = **Emex**, Neck. (Polygon.).
spinosum, Burch. l. c. = E. spinosa.

CENTOTHECA, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 189 (Centosteca). *GRAMINEAE*, Benth. & Hook. f. iii. 1190.
lappacea, *Desv. l. c.*—Geront. trop.
latifolia, Trin. Fund. Agrost. 141 = lappacea.
parviflora, *Anderss. in Peters, Reise Mossamb. Bot.* 559.—Afr. trop.

CENTRACHAENA, Less. Syn. Comp. 255 (1833) = seq.

CENTRACHENA, Schott, Cat. Ind. Sem. Hort. Vindob. (1823); ex Reichb. Ic. Pl. Crit. iv. 47 (1827) = **Chrysanthemum**, Tourn. (Compos.).
viscida, Schott, l. c.; ex Reichb. l. c. = Chrys. viscosum.

CENTRADENIA, G. Don, Gen. Syst. ii. 755 (1832). *MELASTOMACEAE*, Benth. & Hook. f. i. 739.
Doncklaeria, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1350 (1855).
Plagiophyllum, Schlecht. in Linnaea, xiii. (1839) 429.
divaricata, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xix. (1851) 354.—Am. centr.
floribunda, *Planch. in Fl. des Serres*, Sér. I. v. (1849) t. 453.—Mexic.
grandifolia, *Endl. ex Walp. Rep.* ii. 119.—Mexic.
inaequilateralis, *G. Don, Gen. Syst.* ii. 765.—Mexic.
Oerstediana, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xix. (1851) 354.—Am. centr.
ovata, *Klotzsch, l. c.*—Am. centr.
rosea, Lindl. Bot. Reg. (1843) t. 20 = inaequilateralis.

CENTRANDRA, Karst. in Linnaea, xxviii. (1856) 440; = **Julocroton**, Mart. (Euphorbiac.).
hondensis, Karst. l. c. = J. hondensis.

CENTRANTHERA, Scheidw. in Otto & Dietr. Allg. Gartenz. x. (1842) 293 = **Pleurothallis**, R. Br. (Orchid.).
punctata, Scheidw. l. c. = P. Centranthera.

CENTRANTHERA, R. Br. Prod. 438 (1810). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 969.
Gumteolis, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 87 (1825).
Purshia, Dennst. Schluess. Hort. Malab. 35 (1818).
Razumovia, Spreng. Syst. ii. 682 (1825).
Brunoniana, Thw. Enum. Pl. Zeyl. 220 = procumbens.
Brunoniana, *Wall. Cat.* n. 3882; *Benth. Scroph. Ind.* 50.—Reg. Himal.; Burma.
grandiflora, *Benth. in Wall. Cat.* n. 3880; *Scroph. Ind.* 50.—Reg. Himal.
hispida, Benth. Fl. Hongk. 254 = Brunoniana.
hispida, *R. Br. Prod.* 438.—Austral.; Afghan.
humifusa, *Wall. Cat.* n. 3883.—Ind. or.
nepalensis, D. Don, Prod. Fl. Nep. 88 = hispida.
procumbens, *Benth. in DC. Prod.* x. 525.—Ind. or.
prostrata, Boj. Hort. Maurit. 246 = Radamaea prostrata.

CENTRANTHUS, DC. Fl. Fr. iv. 238 (1805) *VALERIANEAE*, Benth. & Hook. f. ii. 155.
Hybidium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 399.
Kentranthus, Neck. Elem. i. 122 (1790), nomen prius.
angustifolius, Bieb. Fl. Taur. Cauc. i. 67 = longiflorus.
angustifolius, Boiss. Voy. Bot. Espagne, 291 = nevadensis.
angustifolius, *DC. Fl. Fr.* iv. 239.—Europ. austr.
Calcitrapa, *Dufr. Hist. Valér.* 39; *DC. l. c.* v. 492.—Europ.
Clausonis, *Pomel, Nouv. Mat. Fl. Atl.* 73.—Afr. bor.
dasycarpus, Kunze, Del. Sem. Hort. Lips. (1847); et Linnaea, xxiv. (1851) 176 = macrosiphon.
elatus, Boiss. & Heldr. Diagn. Ser. I. x. 73 = longiflorus.
junceus, *Boiss. & Heldr. l. c.*—Graecia.

CENTRANTHUS:—
latifolius, Dufr. Hist. Valér. 38 = ruber.
Lecoqii, Jord. Pugill. Pl. Nov. 76 = angustifolius.
longiflorus, *Stev. Obs. Pl. Ross.* 76.—As. Min.; Syria; Reg. Caucas.
macrosiphon, *Boiss. Diagn.* Ser. I. iii. 57.—Hispan.
marinus, S. F. Gray, Nat. Arr. Brit. Pl. ii. 479 = ruber.
maritimus, DC. Prod. iv. 632 = ruber.
nervosus, *Moris, Elench. Sard.* ii. 4.—Corsic.; Sardin.
nevadensis, *Boiss. Diagn.* Ser. II. ii. 120.—Hispan.
orbiculatus, Dufr. Hist. Valér. 39 = C. Calcitrapa.
orbiculatus, *Roem. & Schult. Syst.* i. 35.—Cyprus.
pinnatifidus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Calcitrapa.
ruber, *DC. Fl. Fr.* iv. 239.—Europ.; Syria.
Sibthorpii, *Heldr. & Sart. Diagn.* Ser. II. ii. 119.—Graecia.

CENTRAPALUS, Cass. in Dict. Sc. Nat. vii. 382 (1817) = **Vernonia**, Schreb. (Compos.).
galamensis, Cass. l. c. 383 = V. pauciflora.

CENTRATHERUM, Cass. in Bull. Soc. Philom. (1817) 31. *COMPOSITAE*, Benth. & Hook. f. ii. 225.
Ampherephis, H. B. & K. Nov. Gen. et Sp. iv. 31. t. 314, 315 (1820).
Amphibecis, Schrank, in Syll. Ratisb. i. (1824) 86.
Crantzia, Vell. Fl. Flum. Ic. viii. t. 153 (1827).
Decaneurum, DC. in Guill. Arch. Bot. ii. (1833) 516.
Phyllocephalum, Blume, Bijdr. 888 (1826).
Rolfinkia, Zenk. Pl. Ind. 13. t. 14 (1837).
Spixia, Schrank, Pl. Rar. Hort. Monac. t. 80 (1819).
Wightia, Spreng. ex DC. Prod. v. 67 (1836).
aristatum, Cass. in Dict. Sc. Nat. lvii. 342 = punctatum.
brachylepis, *Sch. Bip. ex Baker, in Mart. Fl. Bras.* vi. II. 12.—Bras.
brevispinum, Cass. in Dict. Sc. Nat. lvii. 346 = intermedium.
chinense, Less. in Linnaea, iv. (1829) 320 = Vernonia chinensis.
courtallense, *Benth. & Hook. f. Gen.* ii. 225.—Ind. or.
courtallense, C. B. Clarke, Comp. Ind. 3 = Ritchiei.
Hookeri, *C. B. Clarke, l. c.*—Ind. or.
intermedium, *Less. in Linnaea*, iv. (1829) 320.—Bras.
longispinum, Cass. in Dict. Sc. Nat. lvii. 346 = punctatum.
Metzianum, *Hook. f. Fl. Brit. Ind.* iii. 228.—Ind. or.
molle, *Benth. & Hook. f. Gen.* ii. 225.—Ind. or.
muticum, *Less. in Linnaea*, iv. (1829) 320.—Mexic.
phyllolaenum, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 228.—Ind. or.
pulchellum, *Steud. Nom.* ed. II. i. 324.—N. Zel.
punctatum, *Cass. in Dict. Sc. Nat.* vii. 384.—Venezuela.
reticulatum, *Benth. & Hook. f. Gen.* ii. 225.—Ind. or.
Ritchiei, *Hook. f. Fl. Brit. Ind.* iii. 228.—Ind. or.
scariosum, C. B. Clarke, Comp. Ind. 4 = Vernonia scariosa.
tenue, *C. B. Clarke, l. c.*—Ind. or.

CENTROCARPHA, D. Don, in Sweet, Hort. Brit. ed. II. 310 (1830); et in Brit. Fl. Gard. Ser. II. iv. t. 87 (1831) = **Rudbeckia**, Linn. (Compos.).
acutifolia, Sweet, Hort. Brit. ed. II. 310, nomen = R. Newmanni.
aristata, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. sub t. 87 = R. triloba.
chrysantha, Steud. Nom. ed. II. i. 321 = R. Newmanni.
chrysomela, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. sub t. 87 = R. fulgida.
discolor, D. Don, l. c. = R. fulgida?
fulgida, D. Don, l. c. = R. fulgida.
gracilis, D. Don, in Sweet, Hort. Brit. ed. II. 310; et l. c. = R. hirta.
grandiflora, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. t. 87 = R. grandiflora.
hirta, D. Don, in Sweet, Hort. Brit. ed. II. 310; et l. c. = R. hirta.
moschata, J. Cree, ex Steud. Nom. ed. II. i. 321 = R. moschata.
Radula, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. sub t. 87 = Helianthus Radula.
spathulata, D. Don, l. c. = R. spathulata.
subtomentosa, Sweet, Hort. Brit. ed. I. 310, nomen = R. subtomentosa.
triloba, D. Don, in Sweet, Brit. Fl. Gard. Ser. II. iv. sub t. 87 = R. subtomentosa.
triloba, Loud. Hort. Brit. Suppl. i. 584 = R. triloba.

CENTROCHILUS, J. C. Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (Pl. Meyen.) 435. t. 113 (1843) = **Habenaria**, Willd. (Orchid.).
gracilis, Schau. l. c. = H. tipuloides.

CENTROCLINIUM, D. Don, in Trans. Linn. Soc. xvi. (1830) 254 = **Onoseris**, DC. (Compos.).
adpressum, Hook. Bot. Mag. t. 3115 = O. adpressa.
albicans, D. Don, in Trans. Linn. Soc. xvi. (1830) 254 = O. integrifolia.
altissimum, Poepp. & Endl. Nov. Gen. et Sp. iii. 52. t. 259.—Peruv.
hyssopifolium, D. Dietr. Syn. Pl. iv. 1565 = O. hyssopifolia.
reflexum, Hook. Bot. Mag. t. 3115 = O. reflexa.
salicifolium, D. Dietr. Syn. Pl. iv. 1565 = O. salicifolia.

CENTRODISCUS, Muell. Arg. in Mart. Fl. Bras. xi. II. 325. t. 102 (1874) = **Caryodendron**, Karst. (Euphorb.).
grandifolius, Muell. Arg. l. c. 327.—Bras.

CENTROGLOSSA, Rodrig. Orch. Nov. ii. 234 (1881). *ORCHIDEAE*.
macroceras, *Rodrig. l. c.* 235.—Bras.
tripollinica, *Rodrig. l. c.*—Bras.

CENTROLEPIS, Labill. Nov. Holl. Pl. i. 7. t. 1 (1804). *CENTROLEPIDEAE*, Benth. & Hook. f. iii. 1026.
Alepyrum, R. Br. Prod. 253 (1810).
Centrosepis, R. Hedw. Gen. 51 (1806).
Devauxia, R. Br. Prod. 253 (1810) ed. Necs, 108 (1827).
aemula, Rudge, in Trans. Linn. Soc. x. (1811) 284 = strigosa.
alepyroides, *Walp. Ann.* i. 897.—Austral.
aristata, *Roem. & Schult. Syst.* i. 44.—Austral.
Banksii, *Roem. & Schult. l. c.*—Austral.
brevifolia, Walp. Ann. i. 896 = Drummondii.
cambodiana, *Hance, in Journ. Bot.* xiv. (1876) 14.—Cambodia.
cuspidigera, Rudge, in Trans. Linn. Soc. x. (1811) 283 = fascicularis.
Drummondii, *Walp. Ann.* i. 896.—Austral.
exserta, *Roem. & Schult. Syst.* i. 44.—Austral.
fascicularis, *Labill. Nov. Holl. Pl.* i. 7. t. 1.—Austral.
glabra, *Hieron. in Abh. Naturf. Ges. Halle*, xii. (1873) 209.—Austral.
humillima, *F. Muell. ex Benth. Fl. Austral.* vii. 203.—Austral.
longifolia, *Steud. Nom.* ed. II. i. 324; *et Kunth, Enum. Pl.* iii. 489.—Austral.
monogyna, *Benth. Fl. Austral.* vii. 205.—Austral.
muscoides, *Hieron. in Abh. Naturf. Ges. Halle*, xii. (1873) 209.—Tasmania.
mutica, *Hieron. l. c.* 211.—Austral.
Patersoni, Roem. & Schult. Syst. i. 43 = strigosa.
pilosa, *Hieron. in Abh. Naturf. Ges. Halle*, xii. (1873) 216.—Austral.
polygona, Hieron. Centrolep. 96 = polygyna.
polygyna, *Hieron. in Abh. Naturf. Ges. Halle*, xii. (1873) 210.—Austral.
pulchra, Hieron. l. c. 213 = Drummondi.
pulvinata, *Roem. & Schult. Syst.* i. 43.—Tasmania.
pusilla, *Roem. & Schult. l. c.* 44.—Austral.
strigosa, *Roem. & Schult. l. c.* 43.—Austral.
tenuior, Roem. & Schult. l. c. = strigosa.
Urvillei, Hieron. in Abh. Naturf. Ges. Halle, xii. (1873) 214 = Drummondii.

CENTROLOBIUM, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 95. *LEGUMINOSAE*, Benth. & Hook. f. i. 546.
minus, Presl, Bot. Bemerk. 62 = robustum.
paraënse, *Tul. in Arch. Mus. Par.* iv. (1844) 87.—Am. austr.
robustum, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 95.—Bras.
tomentosum, *Guill. ex Benth. in Hook. Journ. Bot.* ii. (1840) 66.—Bras.

CENTRONIA, Blume, Bijdr. 776 (1826) = **Aeginetia**, Linn. (Orobanchac.).
mirabilis, Blume, l. c. = A. Centronia.

CENTRONIA, D. Don, in Mem. Wern. Soc. iv. (1823) 314. *MELASTOMACEAE*, Benth. & Hook. f. i. 750.
BRACHYCENTRUM, Meissn. Gen. 114 (1837).
CALYPTRARIA, Naud. in Ann. Sc Nat. Sér. III. xviii. (1852) 132.
STEPHANOGASTRA, Karst. & Triana, in Linnaea, xxviii. (1856) 425.
brachycera, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 72.—N. Granat.
crassiramus, *Triana, l. c.*—Guiana.
excelsa, *Triana, l. c.*—Peruv.; Ecuador.
eximia, *Triana, l. c.*—Ecuador.
haemantha, *Triana, l. c.*—N. Granat.
insignis, *Triana, l. c.*—Ecuador.
laurifolia, *D. Don, in Mem. Wern. Soc.* iv. (1823) 315.—Peruv.
Mutisii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 72.—N. Granat.
phlomoides, *Triana, l. c.*—Costa Rica.
reticulata, *Triana, l. c.* 71.—Peruv.

CENTRONOTA, DC. in Meissn. Gen. 303 [213] (1840); et Prod. ix. 286 (1843) = **Aeginetia**, Linn. (Orobanchac.).
mirabilis, DC. Prod. ix. 186 = Ae. Centronia.

CENTROPAPPUS, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 124 = **Senecio**, Tourn. (Compos.).
Brunonis, Hook. f. l. c. = S. Centropappus.

CENTROPETALUM, Lindl. Sert. Orch. sub t. 21 (1839). *ORCHIDEAE*, Benth. & Hook. f. iii. 570.
FERNANDEZIA, Ruiz & Pav. Prod. Fl. Per. 123. t. 27 (1794).
NASONIA, Lindl. in Benth. Pl. Hartw. 150 (1844).
distichum, *Lindl. l. c.*—Peruv.
Myrtillus, *Pfitzer, Verg. Morph. Orch.* 16.—Peruv.
Warszewiczii, *Reichb. f. in Bot. Zcit.* x. (1852) 706.—Peruv.

CENTROPHORUM, Trin. Fund. Agrost. 106. t. 5 (1820) = **Chrysopogon**, Trin. (Gramin.).
chinense, Trin. l. c. = Chrys. aciculatus.

CENTROPHYLLUM, Dum. Anal. Fam. 32 (1829) = KENTROPHYLLUM, Neck. = **Carthamus**, Linn. (Compos.).

CENTROPHYTA, Reichb. Nom. 148 (1841) = KENTROPHYTA, Nutt. = **Astragalus**, Tourn. (Legum.).

CENTROPODIUM, Lindl. Introd. Nat. Syst. ed. II. 442 (1836), sphalm. = CENTOPODIUM, Burch. = **Emex**, Neck. (Polygon.).

CENTROPOGON, Presl, Prod. Monog. Lobel. 48 (1836). *CAMPANULACEAE*, Benth. & Hook. f. ii. 547.
affine, *Mart. & Gal. in Bull. Acad. Brux.* ix. II. (1842) 40.—Mexic.
amplifolius, *Vatke, in Linnaea*, xxxviii. (1874) 716.—Peruv.
Andropogon, *DC. Prod.* vii. 345.—Ecuador.
argentinus, *Griseb. in Goett. Abh.* xxiv. (1879) 219.—Reg. Argent.
Bertcrianus, *DC. Prod.* vii. 345.—Ins. Guadalup.
beslerioides, *DC. l. c.* 346.—N. Granat.
Bonplandianus, *Presl, Prod. Monog. Lobel.* 48.—Am. austr.
calycinus, *Benth. Pl. Hartw.* 212.—N. Granat.
Chamissonianus, *Kanitz, in Mart. Fl. Bras.* vi. IV. 133.—Bras.
coccineus, *Regel, Gartenfl.* (1858) 374, in textu.—Bras.
cordatus, Mart. & Gal. in Bull. Acad. Brux. ix. II. (1842) 40 = cordifolius.
cordifolius, *Benth. Pl. Hartw.* 77.—Mexico.
costaricanus, *Planch. & Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 156.—Costa Rica.
cuspidatus, *DC. Prod.* vii. 346.—Peruv.
densiflorus, *Benth. Pl. Hartw.* 138.—Peruv.
discolor, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1847) 13.—Venezuela.
edulis, Presl, Prod. Monog. Lobel. 48 = surinamensis.
exasperatus, *Presl, l. c.*—Peruv.
fasciculatus, *Regel, Cat. Pl. Hort. Aksakov.* 29.—Hab.?

CENTROPOGON:—
fastuosus, *Scheidw. in Otto & Dietr. Allg. Gartenz.* ix. (1841) 396.—Mexico.
floccosus, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 16.—N. Granat.
glandulosus, *Decne. in Rev. Hortic.* Sér. III. ii. (1848) 421.—N. Granat.
grandis, *Presl, Prod. Monog. Lobel.* 48.—N. Granat.
granulosus, *Presl, l. c.* 49.—Peruv.
hirtus, *Presl, l. c.* 48.—Ecuador.
laevigatus, *DC. Prod.* vii. 344.—Guiana.
longipes, *Regel, Gartenfl.* iii. (1854) 3. t. 75.—Am. centr.
nutans, *Planch. & Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 156.—Costa Rica.
oblongus, *Benth. Pl. Hartw.* 214.—N. Granat.
prostratus, *Benth. l. c.* 212.—Peruv.
reflexus, *Presl, Prod. Monog. Lobel.* 49.—Peruv.
scandens, *Planch. & Oerst. in Kjoeb. Vidensk. Meddel.* (1857) 157.—Costa Rica.
solanifolius, *Benth. Pl. Hartw.* 139.—Peruv.
speciosus, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 16.—Venezuela.
surinamensis, *Presl, Prod. Monog. Lobel.* 48.—Am. austr.
tovarensis, *Planch. & Linden, in Fl. des Serres*, Sér. I. viii. (1852–53) 145.—Venezuela.
virescens, Planch. & Oerst. in Kjoeb. Vidensk. Meddel. (1857) 157 = Burmeistera virescens.
Warscewiczii, *Van Houtte, ex Regel, Gartenfl.* (1858) 374.—Hab.?
Warscewiczii, *Vatke, in Linnaea*, xxxviii. (1874) 716.—Costa Rica.

CENTROPSIS, Endl. Gen. Suppl. ii. 33 (1842) = KENTROPSIS, Moq. = **Sclerolaena**, R. Br. (Chenop.).

CENTROSEMA, Benth. in Ann. Wien. Mus. ii. (1838) 117. *LEGUMINOSAE*, Benth. & Hook. f. i. 527.
CRUMINIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 423.
PILANTHUS, Poit. ex Endl. Gen. 1289 (1841).
PLATYSEMA, Hoffmgg. ex Benth. in Ann. Wien. Mus. ii. (1838) 117.
STEGANOTROPIS, Lehm. Ind. Sem. Hort. Hamb. (1826).
VEXILLARIA, Hoffmgg. Verz. Pfl. 119 (1824); Benth. in Ann. Wien. Mus. ii. (1838) 117.
acutifolium, Benth. l. c. 118 = brasilianum.
angustifolium, *Benth. l. c.*—Venezuela.
arenarium, *Benth. l. c.* 119.—Bras.
bifidum, *Benth. l. c.* 118.—Bras.
biflorum, Mart. ex Benth. l. c. 120 = virginianum.
brachypodum, Benth. in Ann. Nat. Hist. iii. (1839) 435 = brasilianum.
bracteosum, *Benth. in Ann. Wien. Mus.* ii. 119.—Bras.
brasilianum, *Benth. l. c.* 118.—Bras.
conjugatum, *Steud. Nom.* ed. II. i. 325.—Am. austr.
coriaceum, *Benth. in Ann. Wien. Mus.* ii. (1838) 118.—Bras.
cuspidatum, *Hassk. in Flora*, li. (1868) 27.—Ind. or.
dasyanthum, *Benth. in Ann. Nat. Hist.* iii. (1839) 436.—Bras.
decumbens, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 120 = virginianum.
dubium, Hemsl. Biol. Centr. Am. Bot. i. 294 = hastatum.
fasciculatum, *Benth. in Ann. Wien. Mus.* ii. (1838) 120.—Bras.
ferrugineum, *A. Rich. Ess. Fl. Cub.* 410.—Cuba.
glabrum, Benth. in Ann. Wien. Mus. ii. (1838) 119 = coriaceum.
grandiflorum, *Benth. l. c.*—Bras.
grandiflorum, Walp. Rep. v. 529 = Clitoria grandiflora.
hastatum, *Benth. in Ann. Wien. Mus.* ii. (1838) 120.—Bras.
heptaphyllum, *Moric. Pl. Nouv. Rar. Am.* 96. t. 61.—Calif.
insulanum, Steud. Nom. ed. II. i. 325 = brasilianum.
intermedium, A. Rich. Ess. Fl. Cub. 410 = pubescens.
longifolium, Benth. in Ann. Wien. Mus. ii. (1838) 118 = angustifolium.
macrocarpum, *Benth. in Ann. Nat. Hist.* iii. (1839) 436.—Guiana.
micranthum, Steud. Nom. ed. II. i. 386 in syn. sphalm. = virginianum.

CENTROSEMA:—
molle, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 119 = pubescens.
oblongum, Benth. l. c. 118 = coriaceum.
pascuorum, *Mart. ex Benth. l. c.* 120.—Bras.
perspicuum, *Klotzsch, in Schomb. Faun. & Fl. Gui.* 1203, nomen.—Guiana.
platycarpum, *Benth. in Mart. Fl. Bras.* xv. I. 126.—Peruv.
Plumieri, *Benth. in Ann. Wien. Mus.* ii. (1838) 118.—Am. austr.
pubescens, *Benth. l. c.* 119.—Bras.
rigidulum, Benth. in Hook. Journ. Bot. ii. (1846) 58 = arenarium.
rotundifolium, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 119.—Bras.
Salzmanni, Benth. in Ann. Nat. Hist. iii. (1839) 436 = pubescens.
scabriusculum, Hassk. Pl. Jav. Rar. 377 = virginianum.
sinuatum, *Benth. in Ann. Wien. Mus.* ii. (1838) 120.—Bras.
sordidum, *Klotzsch, in Schomb. Faun. & Fl. Guy.* 1203, nomen.—Guiana.
venosum, *Mart. ex Benth. in Mart. Fl. Bras.* xv. I. 133.—Bras.
verticillatum, *Benth. in Hook. Journ. Bot.* ii. (1848) 58.—Guiana.
vestitum, *Klotzsch, in Schomb. Faun. & Fl. Guy.* 1203, nomen.—Guiana.
vetulum, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 119.—Bras.
vexillatum, *Benth. in Ann. Nat. Hist.* iii. (1839) 435.—Guiana.
virginianum, *Benth. in Ann. Wien. Mus.* ii. (1838) 120.—Am. trop.
virginianum, Griseb. in Goett. Abh. vii. (1857) 204 = pubescens.

CENTROSEPIS, R. Hedw. Gen. 51 (1806), sphalm. = **Centrolepis**, Labill.

CENTROSIA, A. Rich. in Mém. Soc. Hist. Nat. Par. iv. (1828) 39 = **Calanthe**, R. Br. (Orchid.).
Auberti, A. Rich. l. c. 40 = Cal. sylvatica.

CENTROSIS, Thou. Orch. Iles Afr. t. 35, 36 (1822) = **Calanthe**, R. Br. (Orchid.).
corymbosa, Thou. l. c. 1er Tabl. = Cal. sylvatica?
plantaginea, Thou. l. c. 1er Tabl. = Cal. sylvatica?
sylvatica, Thou. l. c. tt. 35, 36 = Cal. sylvatica.

CENTROSOLENIA, Benth. in Hook. Lond. Journ. Bot. v. (1846) 362 = **Episcia**, Mart. (Gesnerac.).
aenea, Linden & André, in Illustr. Hortic. xxii. (1875) t. 222.—N. Granat.
bracteata, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 322.—N. Granat.
bractescens, Hook. Bot. Mag. t. 4675 = E. bractescens.
bullata, Lem. Illustr. Hortic. xvi. (1869) t. 607 = E. tessellata.
glabra, Benth. in Hook. Bot. Mag. t. 4552 = E. glabra.
glabra, Hook. Ic. Pl. t. 873 = E. Hookeri.
hirsuta, Benth. in Hook. Lond. Journ. Bot. v. (1846) 362 = E. hirsuta.
hirtiflora, Sprucc, ex Hanst. in Linnaea, xxxiv. (1865–66) 450 = E. hirtiflora.
picta, Hook. Bot. Mag. t. 4611 = E. picta.

CENTROSPERMUM, H. B. & K. Nov. Gen. et Sp. iv. 270. t. 397 (1820) = **Acanthospermum**, Schrank (Compos.).
humile, Less. Syn. Comp. 217 = A. humile.
xanthioides, H. B. & K. Nov. Gen. et Sp. iv. 270 = A. brasilum.

CENTROSPERMUM, Spreng. Nov. Prov. 9 (1818); et Syst. iii. 588 (1826) = **Chrysanthemum**, Linn. (Compos.).
Chrysanthemum, Spreng. l. c. = C. viscosum.

CENTROSTACHYS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 497 (1824) = **Achyranthes**, Linn. (Amarant.).
aquatica, Wall. Cat. n. 6932 = A. aquatica.
diandra, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 504 = A. diandra.
flabelligera, Fenzl, ex Moq. in DC. Prod. xiii. II. 321 = A. aquatica.

CENTROSTEGIA, A. Gray, ex Benth. in DC. Prod. xiv. 27 (1856); et in Pacif. Rail. Rep. vii. 19 (1857).
POLYGONACEAE, Benth. & Hook. f. iii. 83.
leptoceras, *A. Gray, in Proc. Am. Acad.* viii. (1870) 192.—Calif.
Thurberi, *A. Gray, ex Benth. in DC. Prod.* xiv. 27; *et in Pacif. Rail. Rep.* vii. 20.—Calif.

CENTROSTEMMA, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 271. t. 12 = **Hoya**, R. Br. (Asclepiad.).
coriaceum, Meissn. Gen. Comm. 177 = H. coriacea.
Cyrtoceras, Meissn. l. c. = H. multiflora.
elegans, Blume, Mus. Bot. Lugd. Bat. i. 46 (= *Cyrtoceras elegans*).—Borneo.
laurifolium, Blume, l. c. (= *Cyrtoceras laurifolium*).—Java.
Lindleyanum, Decne. in DC. Prod. viii. 634 = H. multiflora.
micranthum, Blume, Mus. Bot. Lugd. Bat. i. 46 (= *Cyrtoceras micranthum*).—Ins. Celebes.
multiflorum, Decne. in Ann. Sc. Nat. Sér. II. ix. (1838) 272 = H. multiflora.

CENTROSTYLIS, Baill. Etud. Gén. Euph. 469 (1858) = **Adenochlaena**, Baill. (Euphorb.).
zeylanica, Baill. l. c. 470. t. 2. f. 28–29 = A. zeylanica.

CENTUNCULUS, Adans. Fam. ii. 256 (1763) = **Cerastium**, Linn. (Caryophyll.).
alpinus, Scop. Fl. Carn. ed. II. i. 321 = Cerast. trigynum.
angustifolius, Scop. l. c. 322 = Cerast. strictum.
arvensis, Scop. l. c. 321 = Cerast. arvense.
rigidus, Scop. l. c. 322 = Cerast. strictum.
semidecandrus, Scop. l. c. 331 = Cerast. semidecandrum.

CENTUNCULUS, Dill. ex Linn. Syst. ed. I (1735).
PRIMULACEAE, Benth. & Hook. f. ii. 637.
ANAGALLIDASTRUM, Mich. ex Adans. Fam. ii. 230 (1763).
MICROPYXIS, Duby, in DC. Prod. viii. 71 (1844).
erectus, *Phil. Fl. Atac.* 35.—Chili.
lanceolatus, Michx. Fl. Bor. Am. i. 94 = minimus.
minimus, *Linn. Sp. Pl.* 116.—Europ.; Am. bor. et austr.
pentandrus, *R. Br. Prod.* 427.—Am. et Austral. trop.
simplex, Hornem. in Fl. Dan. ix. t. 1567 = minimus.
tenellus, Duby, in DC. Prod. viii. 72 = pentandrus.

CEODES, Forst. Char. Gen. 141. t. 71 (1776) = **Pisonia**, Linn. (Nyctagin.).
umbellifera, Forst. l. c. 142 = P. umbellifera.

CEPA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 103 (1737) = **Allium**, Tourn. (Liliac.).
alba, Renault, Fl. Dep. Orne, 33 = A. Cepa?
angulosa, Bernh. Syst. Verz. Erf. 202 = A. angulosum.
ascalonica, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 11 = A. ascalonicum.
carinata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 185 = A. carinatum.
cernua, Moench, Meth. Suppl. 80 = A. cernuum.
esculenta, S. F. Gray, Nat. Arr. Brit. Pl. ii. 183 = A. Cepa.
fistulosa, S. F. Gray, l. c. 184 = A. fistulosum.
flava, Moench, Meth. 243 = A. flavum.
flexuosa, Moench, Meth. Suppl. 80 = A. roseum.
Moly, Moench, Meth. 244 = A. Moly.
obliqua, Moench, l. c. 243 = A. obliquum.
oleracea, Bernh. Syst. Verz. Erf. 202 = A. oleraceum.
pallens, Renault, Fl. Dép. Orne, 33 = A. Cepa?
paniculata, Moench, Meth. 243 = A. paniculatum.
prolifera, Moench, l. c. 244 = A. Cepa.
rubra, Renault, Fl. Dép. Orne, 33 = A. Cepa?
Schoenoprasa, Moench, Meth. 244 = A. Schoenoprasum.
senescens, Moench, l. c. = A. senescens.
tenuifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 114 = A. Schoenoprasum.
ursina, Bernh. Syst. Verz. Erf. 201 = A. ursinum.
ventricosa, Moench, Meth. 244 = A. fistulosum.
Victorialis, Moench, l. c. 243 = A. Victorialis.
vulgaris, Renault, Fl. Dép. Orne, 33 = A. Cepa.

CEPAEA, Caesalp. ex Fourr. in Ann. Sc. Linn. Lyon, N. S. xvi. (1868) 384 = **Sedum**, Tourn. (Crassulac.).
Caesalpini, Fourr. l. c. = S. Cepaea.

CEPALARIA, Rafin. Fl. Tellur. iv. 95 (1836) = **Cephalaria**, Schrad. (Dipsac.).

CEPHAËLIS Sw. Prod. Veg. Ind. Occ. 45 (1788).
RUBIACEAE, Benth. & Hook. f. ii. 127.
CALLICOCCA, Schreb. Gen. i. 126 (1789).
CARAPICHEA, Aubl. Pl. Gui. 167. t. 64 (1775).
CHESNEA, Scop. Introd. 119 (1777).
EURHOTIA, Neck. Elem. i. 207 (1790).
EVEA, Aubl. Pl. Gui. 103. t. 39 (1775).
IPECACUANHA, Arruda, Diss. 44 (1810), ex St. Hil. Pl. Bras. t. 6 (1824).
NETTLERA, Rafin. Sylva Tellur. 147 (1838).
TAPOGOMEA, Aubl. Pl. Gui. i. 357. t. 60 ad 63 (1775).
alba, *Willd. Sp. Pl.* i. 978.—Guiana.
ambigua, DC. Prod. iv. 534 = Psychotria Salzmanniana.
attenuata, *Miq. in Linnaea*, xix. (1847) 444.—Bras.
axillaris, *Sw. Prod. Veg. Ind. Occ.* 45.—Ind. occ.
Beeri, *Teijsm. & Binn. Cat. Hort. Bog.* 384, nomen.—Hab.?
Beskeana, Schlecht. in Linnaea, xxviii. (1856) 533 = Psychotria Beskeana,
bidentata, Benth. in Hook. Niger Fl. 421 = peduncularis.
bidentata, Thunb. ex Roem. & Schult. Syst. v. 214 = Psychotria bidentata.
bracteocardia, DC. Prod. iv. 534 = Psychotria bracteocardia.
brasiliensis, *Cham. in Linnaea*, ix. (1834) 237.—Bras.
bubalina, Pers. Syn. i. 202 = Burchellia capensis.
calycina, Lindl. Collect. t. 21 = Psychotria Lindleyana.
Carapichea, Raeusch. ex Steud. Nom. ed. II. i. 326 = involucrata.
Chamissonis, Steud. l. c. = Salzmannia nitida.
clausa, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 225.—Afr. trop.
colorata, *Hoffmgg. ex Roem. & Schult. Syst.* v. 213.—Bras.
congensis, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 226.—Afr. trop.
coriacea, D. Dietr. Syn. Pl. i. 774 = Psychotria xanthophylla.
coriacea, G. Don, Gen. Syst. iii. 606 = peduncularis.
cornuta, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 224.—Afr. trop.
costaricensis, *Schlecht. in Linnaea*, xxviii. (1856) 546.—Costa Rica.
crocea, *G. F. W. Mey. Prim. Fl. Esseq.* 107.—Guiana.
cuneata, *Korth. in Nederl. Kruidk. Arch.* ii. (1851) 248.—Malaya.
cyanocarpa, Moç. & Sesse, ex DC. Prod. iv. 533 = tomentosa.
cymosa, Spreng. Syst. i. 749 = Psychotria maypurensis.
dichotoma, *Rudge, Pl. Gui.* i. 29. t. 44.—Guiana.
dichotoma, Willd. ex Roem. & Schult. Syst. v. 214 = Psychotria Hoffmannseggiana.
discolor, *Polak. in Linnaea*, xli. (1877) 572.—Costa Rica.
diversifolia, Blume, Bijdr. 1004 = Geophila reniformis.
elata, *Sw. Prod. Veg. Ind. Occ.* 45.—Jamaica.
emetica, Pers. Syn. i. 203 = Psychotria Ipecacuanha.
Evea, *DC. Prod.* iv. 535.—Guiana.
ferruginea, G. Don, Gen. Syst. iii. 605 = Sabicea ferruginea.
fragrans, *Hook. & Arn. Bot. Beech. Voy.* 64. t. 13.—Ins. Societ.
furcata, Bartl. ex DC. Prod. iv. 513 = Psychotria furcata.
fuscescens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 224.—Afr. trop.
gemmiflora, *Willd. ex Roem. & Schult. Syst.* v. 213.—Hab.?
glabra, *Willd. Sp. Pl.* i. 979.—Guiana.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 178.—Malacca.
herbacea, Kurz, in Journ. As. Soc. Beng. (1877) II. 140 = Geophila reniformis.
hirsuta, *Mart. & Gal. in Bull. Acad. Brux.* xi. 1. (1844) 135.—Mexic.
hirta, *Miq. in Linnaea*, xvii. (1843) 71.—Guiana.
Hoffmannseggiana, Roem. & Schult. Syst. v. 214 = Psychotria Hoffmannseggiana.
Humboldtiana, Cham. & Schlecht. in Linnaea, iv. (1829) 136 = Psychotria Humboldtiana.
involucrata, *Willd. Sp. Pl.* i. 979.—Guiana.
Ipecachuanha, Rich. in Bull. Fac. Med. iv. (1818) 92 = Psychotria Ipecacuanha.
justiciaefolia, *Rudge, Pl. Gui.* i. 28. t. 43.—Guiana.
latifolia, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 226.—Afr. trop.
ligularis, *Rich. ex DC. Prod.* iv. 533.—Guiana.

CEPHAËLIS:—
Mannii, *Benth. & Hook. f. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 225.—Afr. trop.
maranhaënsis, *G. Don, Gen. Syst.* iii. 605.—Bras.
microcephala, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 214 = Psychotria microcephala.
microcephala, Miq. in Linnaea, xviii. (1844) 748 = Psychotria Hoffmannseggiana.
muscosa, *Sw. Prod. Veg. Ind. Occ.* 46.—Ins. Martinic.
nuda, *Cham. & Schlecht. in Linnaea*, iv. (1829) 135.—Bras.
oblonga, *DC. Prod.* iv. 535.—Guiana.
ornata, *Spruce, ex Benth. & Hook. f. Gen.* ii. 128, nomen.—Bras.
Patabaea, Steud. Nom. ed. I. 177, 596 (= *Patabea coccinea*).—Bras.
Patrisii, *D. Dietr. Syn. Pl.* i. 773.—Guiana.
peduncularis, *Salisb. Parad. Lond.* t. 99.—Afr. trop.
pedunculata, G. Don, Gen. Syst. iii. 605 = peduncularis.
peruviana, Spreng. Syst. i. 749 = Psychotria capitata.
polycephala, *Schlecht. in Linnaea*, xxviii. (1856) 532.—Peruv.
prunifolia, H. B. & K. Nov. Gen. et Sp. iii. 377 = Psychotria microcephala.
psychotriaefolia, *Seem. Bot. Voy. Herald.* 138.—Panama.
pubescens, Hoffmgg. ex Roem. & Schult. Syst. v. 213 = Psychotria bractcocardia.
punicea, Vahl, Eclog. Am. i. 19 = elata.
purpurea, *Willd. Sp. Pl.* i. 978.—Guiana.
reniformis, H. B. & K. Nov. Gen. et Sp. iii. 377 = Geophila reniformis.
rosea, Benth. in Hook. Journ. Bot. iii. (1841) 224 = Psychotria rosea.
rubescens, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 225.—Afr. trop.
rubra, Hoffmgg. ex Roem. & Schult. Syst. v. 214 = Psychotria rubra.
ruelliaefolia, Cham. & Schlecht. in Linnaea, iv. (1829) 134 = Psychotria ruelliaefolia.
salicifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 376.—N. Granat.
sessiliflora, Willd. Sp. Pl. i. 979 (= *Patabea coccinea*).—Bras.
spathacea, *Hiern, in Oliver, Fl. Trop. Afr.* iii. 225.—Afr. trop.
speciosa, Spreng. Syst. i. 749 = Psychotria speciosa.
stipulacea, *Blume, Bijdr.* 1005.—Java.
suaveolens, *Schweinf. ex Hiern, in Oliver, Fl. Trop. Afr.* iii. 224.—Afr. trop.
subconnata, *Sagot, ex Benth. & Hook. f. Gen.* ii. 128.—Guiana.
subtomentosa, Spreng. Syst. i. 749 = Psychotria subtomentosa.
Swartzii, *DC. Prod.* iv. 534.—Ind. occ.
tenuiflora, D. Dietr. Syn. Pl. i. 774 (= *Patabea tenuiflora*).—Guiana.
tenuis, D. Dietr. l. c. = Psychotria tenerior.
tetranda, Willd. Sp. Pl. i. 979 = C. Evea.
tomentosa, Vahl, Eclog. Am. i. 19 = Psychotria tomentosa.
triplocephala, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 282.—Porto Rico.
turbinata, *Benth. & Hook. f. Gen.* ii. 128, nomen.—Bras.
violacea, Sw. Prod. Veg. Ind. Occ. 45 = Swartzii.
violacea, *Willd. Sp. Pl.* i. 977.—Ind. occ.
violaefolia, H. B. & K. Nov. Gen. et Sp. iii. 379 = Geophila violaefolia.

CEPHALANDRA, Schrad. ex Eckl. & Zeyh. Enum. 280 (1836) = **Coccinia**, Wight & Arn. (Cucurb.).
decipiens, Hook. f. in Oliver, Fl. Trop. Afr. ii. 552 = Cocc. decipiens.
diversifolia, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 18. t. 7 = Cocc. Hartmanniana.
grandis, Kurz, in Journ. As. Soc. Beng. xlvi. (1877) II. 103 = Cocc. indica.
indica, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 16 = Cocc. indica.
McKennii, Naud. l. c. 17 = Cocc. Mac-Kennii.
palmata, Sond. in Harv. & Sond. Fl. Cap. ii. 493 = Cocc. palmata.
pubescens, Sond. l. c. = Cocc. adoënsis.
quinqueloba, Schrad. ex Eckl. & Zeyh. Enum. 280 = Cocc. quinqueloba.
Schimperi, Naud. in Ann. Sc. Nat. Sér. V. v. (1866) 16 = Cocc. indica.

CEPHALANDRA :—
senensis, Klotzsch, in Peters, Reise Mossamb. Bot. 151 = Coccinia senensis.
sessilifolia, Sond. in Harv. & Sond. Fl. Cap. ii. 493 = Cocc. sessilifolia.
Wightiana, Hook. f. Fl. Brit. Ind. ii. Index 757, sphalm. = Cocc. indica.

CEPHALANOPHLOS, Neck. Elem. i. 68 (1790) = **Cnicus**, Tourn. (Compos.).

CEPHALANTHERA, Rich. in Mém. Mus. Par. iv. (1818) 51. *ORCHIDEAE*, Benth. & Hook. f. iii. 619.
Dorycheile, Reichb. Nom. 56 (1841).
acuminata, Ledeb. Fl. Ross. iv. 78 = pallens.
acuminata, Lindl. in Wall. Cat. n. 7405 = ensifolia.
chloidophylla, *Reichb. f. in Linnaea*, xxv. (1852) 228. —Japon.
comosa, Tineo, in Guss. Fl. Sic. Syn. ii. 877 = rubra.
cucullata, *Boiss. & Heldr. Diagn.* Ser. I. xiii. 12.—Graecia; As. Min.; Persia.
ensifolia, *Rich. in Mém. Mus. Par.* iv. (1818) 60.—Europ.; Afr. bor.; As. temp.
epipactoides, Fisch. & Mey. in Ann. Sc. Nat. Sér. IV. i. (1854) 30 = cucullata.
erecta, *Blume, Orch. Archip. Ind.* 188.—Japon.
falcata, *Blume, l. c.* 187.—Japon.
grandiflora, S. F. Gray, Nat. Arr. Brit. Pl. ii. 210 = pallens.
japonica, A. Gray, in Perry, Exp. Jap. ii. 319 = falcata.
lancifolia, *Dum. Fl. Belg.* 134; *Tod. Orch. Sic.* 123.—Sicilia.
lonchophylla, Reichb. f. Ic. Fl. Germ. xiii. et xiv. 136 = pallens.
longibractcata, *Blume, Orch. Archip. Ind.* 188.—Japon.
Maravignae, Tineo, in Guss. Fl. Sic. Syn. ii. 877 = ensifolia.
ochroleuca, Reichb. Fl. Germ. Excurs. 140[20] = pallens.
oregana, *Reichb. f. in Linnaea*, xli. (1877) 53.—Am. bor. occ.
pallens, Hohen. in Bull. Soc. Nat. Mosc. (1838) 257 = ensifolia.
pallens, *Rich. in Mém. Mus. Par.* iv. (1818) 60.—Europ.; Reg. Caucas.
palustris, Rich. ex Pritz. Ic. Ind. ii. 71, sphalm. = Epipactis palustris.
platycheila, Reichb. f. in Bot. Zeit. iii. (1845) 335 = falcata.
Royleana, Regel, in Act. Hort. Petrop. vi. (1879) 490 = Epipactis Royleana.
rubra, *Rich. in Mém. Mus. Par.* iv. (1818) 60.—Europ.; As. Min.; Persia.
Thomsoni, *Reichb. f. in Linnaea*, xli. (1877) 54.—Reg. Himal.
xiphophylla, Reichb. f. Ic. Fl. Germ. xiii. et xiv. 135 = ensifolia.

CEPHALANTHUS, Linn. Gen. ed. I. 61 (1737). *RUBIACEAE*, Benth. & Hook. f. ii. 30.
Acrodryon, Axolus, Eresimus, Gilipus, Silamnus, Rafin. Sylva Tellur. 60–61 (1838).
acuminatus, *Rafin. New Fl. Am.* iii. 25.—Am. bor.
africanus, Reichb. ex DC. Prod. iv. 345 = Mitragyna africana.
angustifolius, *Lour. Fl. Cochinch.* i. 67.—China.
aralioides, Zoll. & Moritzi, Syst. Verz. Zoll. 61 = naucleoides.
chinensis, Lam. Encyc. i. 678 = Nauclea purpurea.
montanus, *Lour. Fl. Cochinch.* i. 67.—China.
natalensis, *Oliver, in Hook. Ic. Pl.* t. 1331.—Natal.
naucleoides, *DC. Prod.* iv. 539.—Reg. Himal.; Malaya; China.
obtusifolia, *Rafin. New Fl. Am.* iii. 25.—Am. bor.
occidentalis, *Linn. Sp. Pl.* 95.—Am. bor.; Mexic.; Ind. occ.
occidentalis, Lour. Fl. Cochinch. i. 67 = Nauclea orientalis.
oppositifolius, Moench, Meth. 487 = occidentalis.
orientalis, Blume, Cat. Gew. Buitenz. 38 = Nauclea purpurea.
orientalis, Linn. Sp. Pl. 95 = Anthocephalus morindaefolius.
orientalis, *Roem. & Schult. Syst.* iii. 105.—China.
pilulifer, Lam. Encyc. i. 679 = Mitragyna parvifolia.
piluliflorus, *Willd. ex Roem. & Schult. Syst.* iii. 525. —Hab.?
procumbens, *Lour. Fl. Cochinch.* i. 7.—Cochinch.
pubescens, *Rafin. New Fl. Am.* iii. 5.—Am. bor.

CEPHALANTHUS:—
salicifolius, Humb. & Bonpl. Pl. Aequin. ii. 63. t. 98 = occidentalis.
Sarandi, *Cham. & Schlecht. in Linnaea*, ii. (1827) 610. —Bras.
spathelliferus, *Baker, in Journ. Bot.* xx. (1882) 137.—Madag.
stellatus, *Lour. Fl. Cochinch.* i. 68.—China.

CEPHALARIA, Schrad. Cat. Sem. Hort. Gotting. (1814) ex Roem. & Schult. Syst. iii. 1 (1818). *DIPSACEAE*, Benth. & Hook. f. ii. 159.
Cerionanthus, Schott, ex Roem. & Schult. l. c. 43 (1818).
Lepicephalus, Lag. Gen. et Sp. Nov. 7 (1816).
Leucopsora, Rafin. Fl. Tellur. iv. 95 (1836).
Picnocomum, DC. Prod. iv. 647 (1830) in syn.
Pterocephalus, Adans. Fam. ii. 152 (1763).
Pycnocomon, Wallr. Sched. Crit. 46 (1822) in syn.
Succisa, Wallr. l. c. (1822).
acaulis, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 368.—Abyssin.
albescens, Roem. & Schult. Syst. iii. 48 = leucantha.
alpina, *Schrad. Cat. Sem. Hort. Gotting.* (1814); *ex Roem. & Schult. Syst.* iii. 43.—Europ.
ambrosioides, *Roem. & Schult. l. c.* 45.—Graecia.
amplexicaulis, *Steud. Nom.* ed. II. i. 327.—Graecia; Oriens?
appendiculata, Schrad. Cat. Sem. Hort. Gotting. (1814); ex Roem. & Schult. Syst. iii. 42 = Dipsacus pilosus.
aristata, *C. Koch, in Linnaea*, xxiv. (1851) 445.—As. Min.
attenuata, *Roem. & Schult. Syst.* iii. 44.—Afr. austr.
balearica, *Coss. ex Willk. in Linnaea*, xl. (1876) 37.—Hispan.
bidens, Roem. & Schult. Syst. iii. 45 = Scabiosa bidens.
boetica, Boiss. Voy. Bot. Espagne, 738 = leucantha.
Boissieri, Reut. in Boiss. Diagn. Ser. II. ii. 122 = syriaca.
cachemirica, Decne. in Jacquem. Voy. Bot. 86. t. 94 = Dipsacus inermis.
calva, Boiss. & Bal. Diagn. Ser. II. vi. 95 = aristata.
centauroides, Roem. & Schult. Syst. iii. 49 = corniculata.
centauroides, Schur, Enum. Pl. Transs. 294 = laevigata.
cilicica, *Boiss. & Kotschy, Diagn.* Ser. II. v. 106.—Cilicia.
coriacea, Steud. Nom. ed. II. i. 327 = corniculata.
corniculata, *Roem. & Schult. Syst.* iii. 49.—Bannatus.
cretacea, Roem. & Schult. l. c. 51 = corniculata.
decurrens, Roem. & Schult. l. c. 50 = attenuata.
diandra, Roem. & Schult. l. c. 52 = Scabiosa papposa.
dichaetophora, *Boiss. Diagn.* Ser. I. vi. 71.—Persia.
dipsacoides, *Boiss. & Bal. Diagn.* Ser. II. v. 107.—Cilicia; Syria.
dipsacoides, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 434 = Dipsacus azureus.
elata, Schrad. in Cat. Sem. Hort. Gotting. (1814); ex Roem. & Schult. Syst. iii. 51 = tatarica.
Fussiana, Heuff. ex Schur, Enum. Pl. Transs. 293 = radiata.
graeca, *Roem. & Schult. Syst.* iii. 43.—Graecia.
hirsuta, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 55. —Persia.
humilis, Roem. & Schult. Syst. iii. 50 = attenuata.
involucrata, *Roem. & Schult. l. c.* 53.—Creta; Cyprus.
joppensis, *Coult. ex DC. Prod.* iv. 648.—Italia; Cilicia; Syria.
juncea, *Boiss. Diagn.* Ser. I. vi. 72.—Persia.
Kotschyi, Boiss. & Hohen. l. c. x. 76 = procera.
laevigata, Panč. ex Nym. Consp. 346 = alpina.
laevigata, Schrad. Cat. Sem. Hort. Gotting. (1814) = corniculata.
lavandulacea, *Sond. in Harv. & Sond. Fl. Cap.* iii. 41. —Afr. austr.
leucantha, Baumg. ex Nym. Consp. 346 = radiata.
leucantha, Ledeb. Fl. Ross. i. 449 = corniculata.
leucantha, *Schrad. in Cat. Sem. Hort. Gotting.* (1814); *ex Roem. & Schult. Syst.* iii. 47.—Hispan.; Gall.
leucanthema, Roem. & Schult. l. c. 48 = leucantha.
linearifolia, Lange, in Kjoeb. Vidensk. Meddel. (1877–78) 226 = leucantha.
longifolia, E. Mey. Zwei Pfl. Docum. 171 = attenuata.
macrophylla, Griseb. Spicil. Fl. Rumel. ii. 174 = ambrosioides.

CEPHALARIA:—
mauritanica, *Pomel, Nouv. Mat. Fl. Atl.* 66.—Afr. bor.
microcephala, *Boiss. Diagn.* Ser. II. ii. 123.—Persia.
neglecta, *Verlot, Ind. Sem. Hort. Gratianop.* (1852) 10; *et Linnaea*, xxvi. (1853–55) 764.—Hab.?
papposa, Roem. & Schult. Syst. iii. 52 = syriaca.
pilosa, *Boiss. & Huet, Diagn.* Ser. II. ii. 122.—Armenia.
pilosa, Gren. & Godr. Fl. Fr. ii. 69 = Dipsacus pilosus.
plumosa, Roem. & Schult. Syst. iii. 53 = Scabiosa lusitanica?
procera, *Fisch. & Avé-Lall. Ind. Sem. Hort. Petrop.* vii. 46.—As. Min.
procera, Panč. ex Nym. Consp. 346 = alpina.
radiata, *Griseb. & Schenk, in Wiegm. Archiv*, xviii (1852) i. 351.—Transsylv.
rigida, Eckl. & Zeyh. Enum. 371 = rigida.
rigida, *Roem. & Schult. Syst.* iii. 44.—Afr. austr.
rupestris, Griseb. ex Nym. Consp. 346 = alpina.
scabra, Eckl. & Zeyh. Enum. 371 = rigida.
scabra, *Roem. & Schult. Syst.* iii. 44.—Afr. austr.
setosa, *Boiss. & Hohen. Diagn.* Ser. I. ii. 107.—Syria.
setulifera, Boiss. & Heldr. in Boiss. Fl. Orient. iii. 124 = graeca.
speciosa, *Boiss. & Kotschy, in Boiss. l. c.* 121.—Armenia.
stellipilis, *Boiss. Diagn.* Ser. I. x. 76.—Syria.
syriaca, *Schrad. Cat. Sem. Hort. Gotting.* (1814); *ex Roem. & Schult. Syst.* iii. 45.—Reg. Mediterr.; Oriens.
tatarica, *Schrad. l. c.*; *ex Roem. & Schult. l. c.* 51. —Rossia; As. bor.; Oriens.
Tchihatchewi, *Boiss. in Tchihat. Asie Min. Bot.* ii. 220. t. 26.—Armenia.
transylvanica, *Schrad. Cat. Sem. Hort. Gotting.* (1814); *ex Roem. & Schult. Syst.* iii. 45.—Europ. austr.; As. Min.; Reg. Caucas.
uralensis, Roem. & Schult. Syst. iii. 50 = corniculata.
ustulata, E. Mey. Zwei Pfl. Docum. 171 = attenuata.
ustulata, *Roem. & Schult. Syst.* iii. 43.—Afr. austr.
Vaillantii, Schott, ex Roem. & Schult. l. c. 46 = syriaca.
virginea, Janka, in Oestr. Bot. Zeitschr. xxii. (1872) 178 = graeca.
Willichii, Roem. & Schult. Syst. iii. 53 = Scabiosa plumosa.

CEPHAELIS, Vahl, Eclog. i. 19 (1796), sphalm. = **Cephaëlis**, Sw. (Rubiac.).

CEPHALIDIUM, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) = **Anthocephalus**, A. Rich. (Rubiac.).

CEPHALINA, Thonn. in Schum. & Thonn. Beskr. Guin. Pl. 105 (1827) = **Sarcocephalus**, Afzel. (Rubiac.).
esculenta, Schum. & Thonn. l. c. = S. esculentus.
scandens, DC. Prod. iv. 368, in syn. = S. esculentus.

CEPHALIPTERUM, A. Gray, in Hook. Kew Journ. iv. (1852) 271. *COMPOSITAE*, Benth. & Hook. f. ii. 321.
Drummondii, *A. Gray, l. c.* 272.—Austral.

CEPHALOCARPUS, Nees, in Mart. Fl. Bras. ii. 1. 162. t. 18 (1842). *CYPERACEAE*, Benth. & Hook. f. iii. 1068.
Dracaenula, *Nees, l. c.*—Bras.

CEPHALOCEREUS, Pfeiff. in Otto & Dietr. Allg. Gartenz. vi. (1838) 142 = **Cereus**, Haw. (Cacteae).
senilis, Pfeiff. l. c. = C. senilis.

CEPHALOCHLOA, Coss. & Dur. in Ann. Sc. Nat. Sér. IV. i. (1854) 229 = **Ammochloa**, Boiss. (Gramin.).
echinata, Coss. & Dur. l. c. = A. pungens.

CEPHALOCROTON, Hochst. in Flora, xxiv. (1841) 370. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 307.
albicans, *Muell. Arg. in DC. Prod.* xv. ii. 760.—Malaya.
cordofanus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 155.—Afr. trop.
discolor, *Muell. Arg. in DC. Prod.* xv. ii. 761.—Malaya.

CEPHALOCROTON:—
indicus, Bedd. Fl. Sylv. ii. t. 261 = Adenochlaena indica.
leucocephalus, *Baill. Adansonia*, v. (1864–65) 148.—Madag.
mollis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 99. t. 17.—Afr. trop.
orientalis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 120.—Ins. Timor.
socotranus, *Balf. in Proc. Roy. Soc. Edinb.* xii. (1884) 95.—Ins. Socotra.
zeylanicus, Baill. Adansonia, v. (1864–65) 148 = Adenochlaena zeylanica.

CEPHALODES, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 159 = **Cephalaria**, Schrad. (Dipsac.).

CEPHALOMA, Neck. Elem. i. 314 (1790) = **Dracocephalum**, Linn. (Labiat.).

CEPHALOMAPPA, Baill. Adansonia, xi. 130 (1874). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 323.
Beccariana, *Baill. l. c.* 131.—Borneo.

CEPHALONOPLOS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 95 = **Cnicus**, Tourn. (Compos.).
arvensis, Fourr. l. c. = Cnicus arvensis.

CEPHALOPAPPUS, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 5. t. 1. *COMPOSITAE*, Benth. & Hook. f. ii. 504.
sonchifolius, *Nees & Mart. l. c.* 6.—Bras.

CEPHALOPHORA, Cav. Ic. vi. 79. t. 599 (1801). *COMPOSITAE*, Benth. & Hook. f. ii. 413.
ACTINEA, A. Juss. in Ann. Mus. Par. ii. (1803) 425. t. 61.
ACTINELLA, Pers. Syn. ii. 469 (1807).
GRAEMIA, Hook. Exot. Fl. t. 189 (1827).
GRAHAMIA, Spreng. Syst. iv. Cur. Post. 297, 299 (1827).
PTILEPIDA, Rafin. in Am. Monthly Mag. (1818) 268.
acaulis, DC. Prod. v. 663 = Actinella acaulis.
anthemoides, Less. in Linnaea, vi. (1831) 518 = Actinella odorata.
aromatica, *Schrad. Ind. Sem. Hort. Gotting.* (1830); *Less. Syn. Comp.* 240.—Chili.
Berteroana, *Phil. in Linnaea*, xxix. (1857–58) 9.—Chili.
collina, *Phil. l. c.* 8.—Chili.
decurrens, Less. l. c. vi. (1831) 517 = Helenium puberulum.
Doniana, *Hook. & Arn. in Hook. Journ. Bot.* iii. (1841) 324.—Reg. Argent.
elongata, D. Don, ex DC. Prod. vii. 293, pro parte = Gaillardia megapotamica.
elongata, D. Don, ex Hook. Journ. Bot. iii. (1841) 324, pro parte = Doniana.
fasciculata, Pohl, ex J. A. Schmidt, in Mart. Fl. Bras. viii. 89 = Hyptis caprariaefolia.
foliosa, *Phil. in Linnaea*, xxix. (1857–58) 7.—Chili.
glauca, *Cav. Ic.* vi. 80. t. 599.—Chili.
grandiflora, *Walp. Ann.* i. 416.—Am. bor. occ.
heterophylla, *Less. Syn. Comp.* 240.—Reg. Argent.
insularis, *Phil. in Anal. Univ. Chil.* (1873) 506.—Chili.
integrifolia, Steud. Nom. ed. II. i. 328 = Helenium integrifolium.
litoralis, *Phil. Fl. Atac.* 34.—Chili.
plantaginea, *DC. Prod.* v. 662.—Chili.
punctata, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 328.—Chili.
radiata, Less. in Linnaea, vi. (1831) 516 = heterophylla.
rigida, *Phil. in Linnaea*, xxxiii. (1864–65) 170.—Chili.
scabiosoides, D. Don, ex DC. Prod. vii. 293 = Gaillardia megapotamica.
scaposa, DC. Prod. v. 663 = Actinella scaposa.
setigera, *Phil. in Anal. Univ. Chil.* (1873) 505.—Chili.
suffruticosa, D. Don, ex Hook. Journ. Bot. iii. (1841) 324 = Doniana.
tenera, Cass. Opusc. iii. 93 = aromatica.
Urmenetae, *Phil. in Anal. Univ. Chil.* (1873) 505.—Chili.

CEPHALOPHORUS, Lem. Cact. Aliq. Nov. Desc. p. xii (1838) = **Cereus**, Haw. (Cacteae).
Columna-Trajani, Lem. l. c. = Cereus Columna-Trajani.
senilis, Lem. l. c. = Cereus senilis.

CEPHALOPHYTON, Hook. f. ex Baker, in Journ. Linn. Soc. xx. (1883) 250, nomen = **Thonningia**, Vahl, (Balanoph.).
Parkeri, Hook. f. ex Baker, l. c.—Madag.

CEPHALORRHYNCHUS, Boiss. Diagn. Ser. I. iv. 28 (1844) = **Lactuca**, Tourn. (Compos.).
Aucheri, Boiss. l. c. vii. 11 = L. microcephala.
Candolleanus, Boiss. Fl. Orient. iii. 820 = L. hispida.
glandulosus, Boiss. Diagn. Ser. I. iv. 28.—As. Min.
hispidus, Boiss. Fl. Orient. iii. 821 = L. microcephala.
Kotschyi, Boiss. & Reut. ex Boiss. l. c. 816 = L. seticuspis.

CEPHALOSCHOENUS, Nees, in Linnaea, ix. (1834) 295 = **Rynchospora**, Vahl (Cyperac.).
Armeria, Nees, l. c. 296 = R. globosa.
articulatus, Nees, in Edinb. N. Phil. Journ. xvii. (1834) 266 = R. aurea.
divergens, Nees, in Linnaea, ix. (1834) 296 = R. aurea.
globosus, Nees, l. c. = R. globosa.
longisetis, Nees, l. c. = R. longisetis.
marginatus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 250 = R. globosa.
oligocephalus, Hochst. in Flora, xxviii. (1845) 759 = R. cyperoides.
parvus, Nees, in Linnaea, ix. (1834) 296 = R. Wallichiana.
Pohlianus, Nees, in Mart. Fl. Bras. ii. 1. 131 = R. globosa.
polycephalus, Nees, in Linnaea, ix. (1834) 296 = R. cyperoides.
pseudomariscus, Nees, l. c. = R. gigantea.
tenuirostris, Nees, l. c. = R. cyperoides, tenuirostris.
zeylanicus, Nees, in Edinb. N. Phil. Journ. xvii. (1834) 265 = R. triflora.

CEPHALOSCIRPUS, Kurz, in Journ. As. Soc. Beng. xxxviii. (1869) II. 83 = **Mapania**, Aubl. (Cyperac.).
macrocephalus, Kurz, l. c. 84.—Ins. Molucc.

CEPHALOSERIS, Poepp. ex Reichb. Consp. 212* (1828) = **Polyachyrus**, Lag. (Compos.).
Poeppigii, Kunze, ex Steud. Nom. ed. II. i. 328 = P. Poeppigii.

CEPHALOSORUS, A. Gray, in Hook. Kew Journ. iii. (1851) 152 = **Angianthus**, Wendl. (Compos.).
brevipapposus, F. Muell. Fragm. iii. 159 = A. phyllocephalus.
gnephosoides, F. Muell. l. c. ii. 158 = Gnephosis cyathopappa.
gymnocephalus, A. Gray, in Hook. Kew Journ. iii. (1851) 153 = Gnephosis macrocephala.
leptocladus, F. Muell. Fragm. iii. 158 = Gnephosis leptoclada.
microcephalus, F. Muell. l. c. = A. microcephalus.
phyllocephalus, A. Gray, in Hook. Kew Journ. iii. (1851) 152 = A. phyllocephalus.

CEPHALOSTACHYUM, Munro, in Trans. Linn. Soc. xxvi. (1868) 138. *GRAMINEAE*, Benth. & Hook. f. iii. 1213.
capitatum, *Munro, l. c.* 139.—Reg. Himal.
Chapelieri, *Munro, l. c.* 140.—Madag.
flavescens, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 252.—Burma.
Griffithii, Kurz, For. Fl. Brit. Burma, ii. 566 = Teinostachyum Griffithii.
latifolium, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 140.—Reg. Himal.
pallidum, *Munro, l. c.* 139.—Reg. Himal.
pergracile, *Munro, l. c.* 141.—Burma.
schizostachyoides, *Kurz, For. Fl. Brit. Burma*, ii. 565.—Ins. Andaman.
virgatum, *Kurz, l. c.*—Burma.

CEPHALOSTEMON, Rob. Schomb. Rap. & Saxo-Frid. 9 (1845). *RAPATEACEAE*, Benth. & Hook. f. iii. 858.
affinis, *Koern. in Linnaea*, xxxvii. (1871–73) 447.—Venezuela.

CEPHALOSTEMON:—
gracilis, *Rob. Schomb. Rap. & Saxo-Frid.* 9.—Bras.
Riedelianus, *Koern. in Linnaea*, xxxvii. (1871–73) 445.—Bras.
squarrosus, *Koern. l. c.* 450.—Venezuela.

CEPHALOSTIGMA, A. DC. Monog. Camp. 117 (1830). *CAMPANULACEAE*, Benth. & Hook. f. ii. 555.
anagalloides, Royle, Illustr. Bot. Himal. 253 = hirsutum.
bahiense, A. DC. Prod. vii. 421 = Perrottetii.
erectum, Vatke, in Linnaea, xxxviii. (1874) 699 = Schimperi.
flexuosum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 9.—Ind. or.
hirsutum, *Edgew. in Trans. Linn. Soc.* xx. (1851) 81.—Reg. Himal.
hirsutum, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 9 = Schimperi.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 429.—Reg. Himal.
paniculatum, *A. DC. Monog. Camp.* 117.—Burma.
paniculatum, Wall. Cat. n. 1295 C = hirsutum.
Perrottetii, *A. DC. Monog. Camp.* 118.—Afr. trop.
Perrottetii, Hook. f. in Journ. Linn. Soc. vii. (1864) 204 = ramosissimum.
Prieurei, *A. DC. Monog. Camp.* 118.—Afr. trop.
ramosissimum, *Hemsl. in Oliver, Fl. Trop. Afr.* iii. 472.—Afr. trop.
Schimperi, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 2.—Ind. or.; Afr. trop.
spathulatum, Thw. Enum. Pl. Zeyl. 422 = Campanula canescens.

CEPHALOTAXUS, Sieb. & Zucc. ex Endl. Gen. Suppl. II. 27 (1842). *CONIFERAE*, Benth. & Hook. f. iii. 430.
Buergeri, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 169.—Japon.
drupacea, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* ii. 108.—Japon.
Fortuni, Hook. Bot. Mag. t. 4499 = drupacea.
Harringtonia, C. Koch, Dendrol. ii. II. 102 = pedunculata.
pedunculata, *Sieb. & Zucc. Fl. Jap. Fam. Nat.* ii. 108.—Japon.
sumatrana, *Miq. Fl. Ind. Bat.* ii. 1076.—Sumatra.
tardiva, Siebold, ex Endl. Syn. Conif. 239 = Taxus brevifolia.
umbraculifera, Siebold, ex Endl. l. c. = Taxus brevifolia.

CEPHALOTOMANDRA, Karst. & Triana, in Linnaea, xxviii. (1856) 429. *NYCTAGINEAE*, Benth. & Hook. f. iii. 10.
fragrans, *Karst. & Triana, l. c.* 430.—N. Granat.

CEPHALOTROPHIS, Blume, Mus. Bot. Lugd. Bat. ii. 75. t. 27. 1852 [1856] = **Malaisia**, Blanco (Urticac.).
javanica, Blume, l. c. 76 = M. tortuosa.
puberula, Miq. Fl. Ind. Bat. Suppl. 416 = M. puberula.

CEPHALOTUS, Labill. Nov. Holl. Pl. ii. 6. t. 145 (1806). *SAXIFRAGEAE*, Benth. & Hook. f. i. 655.
follicularis, *Labill. l. c.* 7.—Austral.

CEPHALOXIS, Desv. Journ. Bot. i. (1808) t. 11 = seq.

CEPHALOXYS, Desv. Journ. Bot. i. (1808) 321 = **Juncus**, Tourn.
flabellata, Desv. l. c. 324 = J. repens.
graminifolia, Nees, ex Kunth, Enum. Pl. iii. 343 = J. cyperoides.

CERADIA, Lindl. Bot. Reg. (1845) Misc. 12 = **Othonna**, Linn. (Compos.).
furcata, Lindl. l. c. (= *Doria Ceradia*).—Afr. austr.

CERAIA, Lour. Fl. Cochinch. ii. 518 (1790) = **Dendrobium**, Sw. (Orchid.).
simplicissima, Lour. l. c. = D. Ceraia.

CERAMANTHE, Dum. Not. Scroph. 7 (1834) = **Scrophularia**, Linn.
vernalis, Dum. l. c. = S. vernalis.

CERAMANTHUS, Hassk. Cat. Hort. Bogor. Alt. 240 (1844) = **Phyllanthus**, Linn. (Euphorbiac.).
gracilis, Hassk. l. c. = P. gracilis.

CERAMIA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 153 = **Erica**, Linn.
auricularis, G. Don, Gen. Syst. iii. 806 = E. urceolaris.
blenna, G. Don, l. c. 807 = E. Vernix.
calathiflora, G. Don, l. c. = E. bicolor.
cordata, G. Don, l. c. = E. cordata.
helianthemifolia, G. Don, l. c. 806 = E. marifolia.
humifusa, G. Don, l. c. 807 = E. humifusa.
latifolia, G. Don, l. c. = E. latifolia.
marifolia, G. Don, l. c. 806 = E. marifolia.
obliqua, G. Don, l. c. 807 = E. obliqua.
oxycoccifolia, G. Don, l. c. = E. oxycoccifolia.
planifolia, G. Don, l. c. 806 = E. planifolia.
serpyllifolia, G. Don, l. c. 807 = E. hispidula.
teucriifolia, G. Don, l. c. = E. teucriifolia.
thymifolia, G. Don, l. c. = E. thymifolia.
urceolaris, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 153 = E. urceolaris.

CERAMICALYX, Blume, Mus. Bot. Lugd. Bat. i. 50 (1849) = **Osbeckia**, Linn. (Melastomac.).
calvus, Blume, l. c. = O. rostrata?
pulchellus, Blume, l. c. = O. rostrata.
stellatus, Blume, l. c. = O. crinita.
ternifolius, Blume, l. c. = O. rostrata.

CERAMIOCEPHALUM, Sch. Bip. in Bull. Soc. Bot. Fr. ix. (1862) 284 = **Crepis**, Linn. (Compos.).
patulus, Sch. Bip. l. c. 285 = Crep. patula.

CERAMIUM, Blume, Bijdr. 1134 (1826) = **Bragantia**, Lour. (Aristolochiac.).
tomentosum, Blume, l. c. 1135 = B. tomentosa.

CERAMOCARPIUM, Nees, ex Meissn. in DC. Prod. xv. i. 111 (1864) = **Ocotea**, Aubl. (Laurin.).
cubense, Nees, l. c. 31 = Phoebe cubensis.
lanatum, Nees, l. c. 102 (= *Mespilodaphne lanata*).—Bras.
marginatum, Nees, l. c. 106 (= *Mespilodaphne marginata*).—Madag.
Maximilianum, Nees, l. c. 108 (= *Mespilodaphne prolifera*).—Bras.
obtusatum, Nees, l. c. 105 (= *Mespilodaphne heteromorpha*).—Ins. Maurit.
polyanthes, Nees, l. c. 133 = O. polyantha.
psychotrioides, Nees, l. c. 124 = O. psychotrioides.
semicompletum, Nees, l. c. 133 (= *Oreodaphne semicompleta*).—Bras.
umbrosum, Nees, l. c. 129 = O. umbrosa, *H. B. & K.*

CERAMOPHORA, Nees, ex Meissn. in DC. Prod. xv. i. 105, 111 (1864) = **Ocotea**, Aubl. (Laurin.).
cymosa, Nees, l. c. 105 (= *Mespilodaphne cymosa*).—Madag.

CERANTHE, Opiz, Seznam, 27 (1852) = **Cerinthe**, Tourn. (Borag.).

CERANTHERA, Beauv. Fl. Ow. et Ben. ii. 10 (1805) = **Alsodeia**, Thon. (Violar.).
dentata, Beauv. l. c. 11. t. 65 = A. guianensis.
ilicifolia, Welw. in Trans. Linn. Soc. xxvii. (1869) 94. t. 2 = A. ilicifolia.
subintegrifolia, Beauv. Fl. Ow. et Ben. ii. 11. t. 66 = A. owariensis.

CERANTHERA, Ell. Sketch, ii. 93 (1824). *LABIATAE*, Benth. & Hook. f. ii. 1191.
DICERANDRA, Benth. in Bot. Reg. sub t. 1300 (1830).
densiflora, *A. Gray, Syn. Fl.* ii. ii. 365.—Florida.
lincarifolia, *Ell. Sketch*, ii. 94.—Am. bor.

CERANTHERA, Endl. Gen. Suppl. ii. 22, in syn. (1842) = CERATANTHERA, Hornem. = **Colebrookia**, Sm. (Labiat.).

CERANTHERA, Rafin. in Am. Monthly Mag. (1818) 176 = **Solanum**, Tourn.
heterandra Rafin. l. c. = S. rostratum.

CERANTHUS, Schreb. Gen. i. 14 (1789) = **Linociera**, Sw. (Oleac.).
corniculatus, Schreb. ex Eichl. in Mart. Fl. Bras. vi. i. 307 = L. compacta.
Schreberi, J. F. Gmel. Syst. 26 = L. compacta.

CERASEIDOS, Sieb. & Zucc. Abh. Akad. Muench. iii. (1843) 743. t. 5 = **Prunus**, Linn. (Rosac.).
apetala, Sieb. & Zucc. l. c. 744.—Japon.

CERASIOCARPUM, Hook. f. in Benth. & Hook. f. Gen. i. 832 (1867). *CUCURBITACEAE*, Benth. & Hook. f. l. c.
Bennettii, Cogn. in DC. Monog. Phan. iii. 729 = zeylanicum.
Maingayi, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 629 = Melothria Rumphiana.
penangense, C. B. Clarke, l. c. = Melothria Rumphiana.
zeylanicum, *Hook. f. in Benth. & Hook. f. Gen.* i. 832.—Zeylan.; Java.

CERASITES, Steud. Nom. ed. II. i. 528 (1840) = CERASTITES, S. F. Gray = **Papaver**, Tourn.

CERASOPHORA, Neck. Elem. ii. 71 (1790) = **Prunus**, Linn. (Rosac.).

CERASTITES, S. F. Gray, Nat. Arr. Brit. Pl. ii. 703 (1821) = **Papaver**, Tourn.
cambrica, S. F. Gray, l. c. 704 = Meconopsis cambrica.
dubia, S. F. Gray, l. c. = P. dubium.
hybrida, S. F. Gray, l. c. = P. hybridum.
laciniata, S. F. Gray, l. c. = P. dubium.
macrocephala, S. F. Gray, l. c. 705 = P. Argemone.

CERASTIUM, [Dill.] Linn. Syst. ed. I (1735). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 148.
CENTUNCULUS, Adans. Fam. ii. 256 (1763).
DICHODON, Bartl. ex Reichb. Nom. 205 (1841).
DUFOUREA, Gren. in Act. Soc. Linn. Bord. ix. (1837) 25.
ESMARCHIA, Reichb. Fl. Germ. Excurs. 793 (1832).
LEUCODONIUM, Opiz, Seznam, 59 (1852).
MOENCHIA, Ehrh. Beitr. ii. 177 (1788).
MYOSOTIS, Tourn. ex Moench, Meth. 224 (1794).
PENTAPLE, Reichb. Ic. Fl. Germ. v. 37. t. 227 (1841).
PREVOTIA, Adans. Fam. ii. 256 (1763).
QUATERNELLA, Ehrh. Beitr. iv. 149 (1789).
acutifolium, Schur, Enum. Pl. Transs. 122 = ovatum.
adscendens, *Wender. ex Steud. Nom.* ed. II. i. 328.—Hab.?
africanum, *Oliver, Fl. Trop. Afr.* i. 141.—Afr. trop.
aggregatum, Dur. ex Brignoli, in Flora, xxiii. (1840) 123 = siculum.
album, Presl, Fl. Sic. 167 = tomentosum.
alpestre, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 151 = triviale.
alpigenum, Schur, l. c. 149 = triviale.
alpinum, All. Fl. Pedem. ii. 116 = arvense.
alpinum, Bieb. Fl. Taur. Cauc. i. 361 = ovatum.
alpinum, Bunge, in Ledeb. Fl. Alt. ii. 180 = glomeratum.
alpinum, *Linn. Sp. Pl.* 438.—Reg. bor. frigid.
alsinifolium, Tausch, in Syll. Ratisb. ii. (1828) 243 = ovatum, arvense.
alsinoides, Pers. Syn. i. 521 = glutinosum, *Fries*.
ambiguum, Fisch. ex Ser. in DC. Prod. i. 419 = arvense.
amplexicaule, Sims, Bot. Mag. t. 1789 = dahuricum.
andinum, *Benth. Pl. Hartw.* 162.—N. Granat.
andinum, *Phil. in Anal. Univ. Chil.* (1862) II. 318.—Chili.
androsaceum, Ser. in DC. Prod. i. 416 = illyricum.
angustifolium, Vitm. Summa Pl. iii. 137 = arvense.
anomalum, *Waldst. & Kit. ex Willd. Sp. Pl.* ii. 812; *et Pl. Rar. Hung.* i. 21.—Europ.; Oricns.
apetalum, Dum. Comm. Bot. 47 = glomeratum.
apricum, Schlecht. in Linnaea, xii. (1838) 208 = nutans.
apuanum, *Parl. in Nuov. Giorn. Bot.* vii. (1875) 69.—Italia.
aquaticum, Linn. Sp. Pl. 439 = Stellaria aquatica.
arabicum, *Fisch. & Mey. ex Heynh. Nom.* ii. 133, nomen.—Arab.
arabidis, *E. Mey. ex Fenzl, in Ann. Wien. Mus.* i. (1836) 340.—Afr. austr.

CERASTIUM:—
araraticum, *Rupr. Fl. Cauc.* 234, in textu.—Reg. Caucas.
arcticum, *Lange, in Fl. Dan.* t. 2963.—Groenland.
arenarioides, Crantz, Inst. ii. 402 = Arenaria spathulata.
arenarium, Tenore, Fl. Nap. iv. 232 = semidecandrum.
arenosum, Kit. in Linnaea, xxxii. (1863) 518 = glomeratum?
argaeum, *Boiss. & Bal. Diagn.* Ser. II. vi. 38.—Cappadoc.
argenteum, Bieb. Fl. Taur. Cauc. i. 361 = grandiflorum.
armeniacum, *Gren. Monog. Cerast.* 19. t. 1.—Armenia.
arvense, Cham. & Schlecht. in Linnaea, v. (1830) 233 = orithales.
arvense, *Linn. Sp. Pl.* 438.—Reg. temp. bor. et austr.
arvensiforme, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 296.—Bolivia.
atlanticum, *Dur. in Duch. Rev. Bot.* ii. (1846–47) 437.—Hispan.
atratum, Lapeyr. Hist. Abr. Pl. Pyr. 265 = alpinum.
atrovirens, Bab. in Mag. Zool. & Bot. ii. (1838) 317. t. 9 = tetrandrum.
atticum, Boiss. & Heldr. Diagn. Ser. II. i. 93 = brachypetalum.
azoricum, *Hochst. ex Seub. Fl. Azor.* 45.—Ins. Azoric.
banaticum, Kit. in Rochel, Banat. Rar., in ind. = grandiflorum.
barbulatum, Wahlenb. Fl. Carp. 137 = brachypetalum.
beeringianum, Cham. & Schlecht. in Linnaea, i. (1826) 62 = alpinum.
Biebersteinii, *DC. in Mém. Soc. Phys. Genèv.* i. (1823) 436.—Tauria.
biflorum, Crantz, Inst. ii. 402 = Stellaria biflora.
biflorum, *Kit. in Linnaea*, xxxii. (1863) 525.—Hungar.
blepharostemon, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 403 = longifolium.
Boissieri, *Gren. in Mém. Soc. Emul. Doubs*, i. (1841) 67.—Hispan.
bombycinum, Schur, Enum. Pl. Transs. 123 = alpinum.
brachycarpum, E. Mey. Zwei Pfl. Docum. 171 = Dregeanum.
brachycarpum, Schur, Enum. Pl. Transs. 124 = arvense.
brachypetalum, *Desp. in Pers. Syn.* i. 520.—Europ.; Reg. Mediterr.
brachypterum, Steud. Nom. ed. II. i. 328, sphalm. = praec.?
bracteatum, Rafin. Préc. Découv. 36 = oblongifolium?
breviflorum, Gilib. Fl. Lituan. ii. 158 = semidecandrum.
bulgaricum, *Uechtritz, in Oestr. Bot. Zeitschr.* xxvi. (1876) 221.—Bulgaria.
caeruleum, *Boiss. Diagn.* Ser. I. i. 53.—As. Min.
caespitosum, Gilib. Fl. Lituan. ii. 159 = glomeratum.
caespitosum, Kit. ex Reichb. Fl. Germ. Excurs. 799 = arvense.
caespitosum, Malmgr. in Vet. Akad. Oefvers. xix. (1862) 242 = alpinum.
caespitosum, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 152.—N. Granat.
calycinum, Fenzl, Verbreit. Alsin. tab. ad p. 56 = Arenaria calycina.
campanulatum, *Viviani, in Ann. Bot.* i. II. 171. t. 1.—Italia; Hung.; Graecia.
camphoratifolium, Stend. Nom. ed. I. 178, 179 = arvense.
candicans, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 295.—Ecuador.
canescens, Hornem. ex Ser. in DC. Prod. i. 416 = brachypetalum.
capense, *Sond. in Harv. & Sond. Fl. Cap.* i. 131.—Afr. austr.
cardiopetalum, C. Gay, Fl. Chil. i. 274 = Commersonianum.
carinthiacum, Vest, in Hoppe, Neues Taschenb. (1808) 229 = ovatum.
carnosulum, Turcz. ex Ledeb. Fl. Ross. i. 413 = arvense.
caucasicum, Fisch. ex Ser. in DC. Prod. i. 414 = nemorale.
chilense, *Bartl. in Presl, Rel. Haenk.* ii. 17.—Chili.

CERASTIUM :—
chloraefolium, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iv. 34.—As. Min.
ciliatum, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 616 = glomeratum.
ciliatum, Waldst. & Kit. Pl. Rar. Hung. iii. 250 = arvense.
collinum, Hort. Dorpat. ex Flora, xvi. (1833) 122 = purpurascens.
collinum, Salisb. Prod. 300 = arvense.
Colsmanni, *Lehm. ex Spreng. Syst.* ii. 418.—Reg. Magell.
Columnae, Tenore, Prod. Fl. Nap. p. xxvii = tomentosum.
Commersonianum, *Ser. in DC. Prod.* i. 417.—Reg. Argent.
Commersonii, Steud. Nom. ed. II. i. 329 = praec.
connatum, Beck, N. et M. St. Bot. 55 = viscosum.
connatum, S. G. Gmel. It. ii. 196 ; ex Ledeb. Fl. Ross. i. 401 = davuricum.
consanguineum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 296.—Peruv.
constantinopolitanum, Nym. Consp. 109 = glomeratum.
cordatum, Crantz, Inst. ii. 400 = Drymaria cordata.
cordifolium, Roxb. Hort. Beng. 35 ; Fl. Ind. ii. 458 = Drymaria cordata.
corniculatum, Crantz, Inst. ii. 403 = dichotomum.
coronense, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 154 = ovatum.
corsicum, Soleir. ex Nym. Consp. 108 = arvense.
crassipes, *Bartl. in Presl, Rel. Haenk.* ii. 18.—Peruv.
cuspidatum, *Hemsl. Diag. Pl. Nov.* 21.—Mexic.
davuricum, *Fisch. ex Spreng. Pugill.* ii. 65.—As. occ. et bor.
dasyphyllum, *Kit. in Linnaea*, xxxii. (1863) 521.—Hungar.
decalvans, Schloss, ex Nym. Consp. 107 = tomentosum.
deflexum, Ser. in DC. Prod. i. 417 = Stellaria aquatica.
densiflorum, *Guss. Suppl. Fl. Sic. Prod.* 136.—Sicilia.
Diazi, *Phil. in Anal. Univ. Chil.* (1862) II. 391 ; *et in Linnaea*, xxxiii. (1864-65) 21.—Chili.
dichotomum, Crantz, Inst. ii. 401 = Stellaria dichotoma.
dichotomum, *Linn. Sp. Pl.* 438.—Reg. Mediterr. ; As. Min.
dichotomum, Schangin, in Pall. N. Nord. Beitr. vi. 93 = davuricum.
dicrotrichum, *Fenzl, ex Rohrb. in Mart. Fl. Bras.* xiv. II. 281.—Bras.
diffusum, Pers. Syn. i. 520 = tetrandrum.
dioicum, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 120.—Hispan.
divaricatum, *Herbich, in Flora*, vii. (1824) 184.—Neapol.
divaricatum, Kit. in Rochel, Pl. Banat. Rar. in ind. = Kitaibelii.
Dregeanum, *Fenzl, in Ann. Wien. Mus.* i. (1836) 341.—Afr. austr.
echinulatum, *Coss. & Dur. ex Debeaux, in Act. Linn. Soc. Bord.* xxiii. (1860) 180, nomen.—China.
elatum, Huet, ex Nym. Consp. 107 = grandiflorum.
elatum, Tenore, Fl. Nap. iv. 236 = tomentosum.
elbrusense, *Boiss. Fl. Orient.* i. 729.—Persia.
elegans, Fisch. ex Ser. in DC. Prod. i. 400 = trigynum.
elongatum, Bieb. Fl. Taur. Cauc. iii. 316 = nemorale.
elongatum, Pursh, Fl. Am. Sept. i. 321 = arvense.
Endressianum, Prol. ex Willk. & Lange, Prod. Fl. Hisp. iii. 637 = pyrenaicum.
eriophorum, Kit. ex Rochel, Pl. Banat. Rar. in ind. = alpinum.
erectum, Coss. & Germ. Fl. Env. Paris (1845) 39 = quaternellum.
falcatum, *Bunge, Verz. Supp. Fl. Alt.* 37.—Sibir. altaic.
fallax, Guss. Suppl. Fl. Sic. Prod. 139 = semidecandrum.
fasciculatum, *Bartl. in Presl, Rel. Haenk.* ii. 16.—Mexic.
filifolium, *Vest, in Flora*, iii. (1820) 353.—Styria.
filiforme, Schleich. ex Gaud. Fl. Helv. iii. 252, in syn. = latifolium.
filiforme, Ser. in DC. Prod. i. 420 = filifolium.
fimbriatum, Ledeb. in Mém. Acad. Pétersb. v. (1815) 540 = Stellaria radians.
Fischeri, Gren. in Mém. Soc. Emnl. Doubs, i. (1841) 3 = chloraefolium.

CERASTIUM :—
Fischerianum, Ser. in DC. Prod. i. 419 = alpinum.
flaccidum, *Andrz. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 155, nomen.—Rossia.
flexuosum, Hegetschw. in Suter, Fl. Helvet. ed. II. App. 454 = latifolium.
floccosum, *Benth. Pl. Hartw.* 162.—N. Granat.
fontanum, Baumg. Enum. Stirp. Transs. i. 425 = triviale.
fragillimum, *Boiss. Diagn.* Ser. I. i. 54.—As. Min. ; Syria.
frigidum, Bieb. Fl. Taur. Cauc. i. 362 = purpurascens.
fulvum, Rafin. Préc. Découv. 36 = glomeratum.
furcatum, Cham. & Schlecht. in Linnaea, i. (1826) 61 = alpinum.
Gayanum, *Boiss. Diagn.* Ser. II. i. 92.—Hispan.
gibraltaricum, Boiss. Elench. 24 = Boissieri.
glabellum, Turcz. ex Ledeb. Fl. Ross. i. 413 = arvense.
glaberrimum, Lapeyr. Hist. Abr. Pl. Pyr. 265 = alpinum.
glabratum, Hartm. Handb. Skand. Fl. ed. I. 180 = alpinum.
glaciale, Gaud. ex Ser. in DC. Prod. i. 419 = latifolium.
glandulosum, *Schur, in Oestr. Bot. Zeitschr.* xix. (1869) 306.—Europ.
glaucum, Gren. in Mém. Soc. Emul. Doubs, i. (1841) 47 = manticum, quaternellum.
glomeratum, *Thuill. Fl. Par.* ed. II. 226.—Reg. temp. et subtrop.
glutinosum, *Fries, Novit.* ed. II. 132.—Europ. ; As. bor.
glutinosum, Hegetschw. Fl. Schweiz, 433 = latifolium.
glutinosum, *H. B. & K. Nov. Gen. et Sp.* vi. 29.—N. Granat.
glutinosum, Nutt. Gen. Am. i. 291 = nutans.
gnaphalodes, Fenzl, in Russegg. Reis. i. 929 = tomentosum.
gracile, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 304 = Riaei.
gracile, Wallr. Beitr. Fl. Hercyn. 201 = brachypetalum.
gramineum, Crantz, Inst. ii. 401 = Stellaria graminea.
grandiflorum, D. Don, Prod. Fl. Nep. 216 = glomeratum ?
grandiflorum, Gilib. Fl. Lituan. ii. 159 = arvense.
grandiflorum, *Waldst. & Kit. Pl. Rar. Hung.* ii. 168.—Europ. or. ; Reg. Caucas.
grandiflorum, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 636 = Boissieri.
Grenieri, F. Schultz, in Arch. Fl. Fr. et Allem. 24 = glutinosum.
Haussknechtii, *Boiss. & Haussk. in Oestr. Bot. Zeitschr.* xxvi. (1876) 411.—Mesopot.
heterophyllum, Schur, Enum. Pl. Transs. 121 = ovatum.
heterophyllum, Viviani, Fl. Lib. Spec. 67 ; et Fl. Cors. Sp. Nov. 17 = illyricum.
hirsutum, Crantz, Inst. ii. 400 = Holosteum hirsutum.
hirsutum, Fisch. ex Ser. in DC. Prod. i. 419 = alpinum.
hirsutum, Muhl. Cat. 46 = glomeratum.
hirsutum, Tenore, Prod. Fl. Nap. p. xxvii = arvense.
hirtellum, *Pomel, Nouv. Mat. Fl. Atl.* 206 —Afr. bor.
Hochstetterianum, *Fenzl, Verbreit. Alsin.* tab. ad p. 56, nomen.—Hispan. ; Gall.
Holostea, Hornem. Hort. Hafn. 434 = nemorale.
holosteiforme, Schur, Enum. Pl. Transs. 119 = semidecandrum.
holosteoides, Fries, Novit. ed. II. 126 = glomeratum.
holosteum, Crantz, Inst. ii. 401 = Stellaria Holostea.
holosteum, Fisch. ex Ser. in DC. Prod. i. 415 = dahuricum.
humifusum, *Camb. in St. Hil. Fl. Bras. Mer.* ii. 166.—Bras.
hybridum, Schur, Enum. Pl. Transs. 125 = tomentosum.
illyricum, *Ard. Animad.* ii. 26.—Graecia ; As. Min.
imbricatum, *H. B. & K. Nov. Gen. et Sp.* vi. 28.—Am. austr.
incanum, Hoffm. Hort. Mosq. 1808. n. 805 ; ex Bieb. Fl. Taur. Cauc. iii. 320 = grandiflorum.
incanum, Ledeb. in Mém. Acad. Pétersb. v. (1815) 540 = arvense.
indicum, *Wight & Arn. Prod.* 43.—Ind. or.
inflatum, *Link, ex Sweet, Hort. Brit.* ed. II. 57 ; Boiss. Fl. Orient. i 721.—As. Min. ; Persia.
insubricum, Moretti, ex Reichb. Fl. Germ. Excurs. 799 = arvense.

CERASTIUM :—
juvenale, *Godr. in Mém. Acad. Stanislas*, Sér. III. [xvii.] 1853 (1854) 383.—Gallia ?
Kablikianum, Wolfner, in Lotos, iv. (1854) 153 = arvense.
Kasbek, *Parrot, Reise Cauc.* ii. t. 3. f. 3.—Reg. Caucas.
Kitaibelii, *Steud. Nom.* ed. II. i. 329.—Bannatus.
Kotschyi, *Boiss. Fl. Orient.* i. 715.—Syria.
Lamottei, Le Grand, ex Nym. Consp. 110 = Riaei.
Lamyi, F. Schultz, in Arch. Fl. Fr. et Allem. 24 = litigiosum.
lanatum, Lam. Encyc. i. 680 = alpinum.
laniferum, Schur, Enum. Pl. Transs. 123 = alpinum.
lanigerum, Clem. in Atti Terza Riun. Scienz. Ital. (1841) 520 = tomentosum.
lanuginosum, Willd. ex Reichb. Fl. Germ. Excurs. 797 = alpinum.
lapponicum, Crantz, Inst. ii. 402 = trigynum.
laricifolium, Vill. Hist. Pl. Dauph. iii. 644 = arvense.
latifolium, Edmonst. in Phytolog. i. (1843) 498 = alpinum.
latifolium, Hartm. Skand. Fl. ed. XI. 239 = arcticum.
latifolium, Lapeyr. Hist. Abr. Pl. Pyr. 265 = pyrenaicum.
latifolium, Linn. Sp. Pl. 439.—Europ. ; Reg. Caucas.
laxiferum, *Schur, Enum. Pl. Transs.* 125.—Transsylvan.
laxum, *Boiss. & Heldr. Diagn.* Ser. I. vi. 25.—Graecia.
Ledebourianum, Ser. in DC. Prod. i. 420 = pilosum.
Lensii, F. Schultz, in Arch. Fl. Fr. et Allem. 24 = litigiosum.
Lerchenfeldianum, Schur, Enum. Pl. Transs. 122 = ovatum.
ligusticum, Viv. Cat. Hort. Dinegro, 15 = campanulatum.
lineare, All. Fl. Pedem. ii. 365 = tomentosum.
lithospermifolium, Bunge, in Ledeb. Fl. Alt. ii. 179 = falcatum.
lithospermifolium, *Fisch. in Mém. Soc. Nat. Mosc.* iii. (1812) 81.—Sibir.
litigiosum, *De Lens, in Loisel. Fl. Gall.* i. 323.—Gallia.
Loefflingii, Crantz, Inst. ii. 403 = pentandrum.
longepedunculatum, Muhl. Cat. 47 = nutans.
longifolium, S. G. Gmel. ex Ledeb. Fl. Ross. i. 414 = grandiflorum.
longifolium, Juss. ex Poir. Encyc. Suppl. ii. 164 = Commersonianum.
longifolium, Tenore, Prod. Fl. Nap. p. xxvii = tomentosum.
longifolium, *Willd. Sp. Pl.* ii. 814.—Armenia.
longirostre, Wichura, in Jahresb. Schles. Ges. (1854) 75 = triviale.
lucorum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 150 = triviale.
luridum, Guss. Fl. Sic. Syn. i. 510 = brachypetalum.
macilentum, Aspegr. Blek. Fl. 34 = semidecandrum.
macranthum, *Boiss. Diagn.* Ser. I. v. 85.—As. Min.
macranthum, *Link, ex Steud. Nom.* ed. II. i. 329, nomen.—Hab. ?
macranthum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 14 = latifolium.
macrocarpum, *Boiss. & Haussk. in Boiss. Fl. Orient.* i. 716.—Mesopotam.
macrocarpum, Schur, Enum. Pl. Transs. 120 = triviale.
macrocarpum, Stev. ex Ledeb. Fl. Ross. i. 407 = purpurascens.
magellanicum, *Phil. in Anal. Univ. Chil.* (1872) 678.—Chili.
malachiforme, Schur, in Verh. Naturf. Ver. Bruenn. xv. II. (1877) 149 = triviale.
manticum, *Linn. Cent. Pl.* ii. 18.—Europ. austr.
matrense, Kit. ex Spreng. Pugill. i. 33 = arvense.
mauritanicum, *Pomel, Nouv. Mat. Fl. Atl.* 206.—Afr. bor.
mauritianum, Bouton, ex Baker, Fl. Maurit. 14 = glomeratum.
maximum, Gilib. Fl. Lituan. ii. 158 = Stellaria aquatica.
maximum, *Linn. Sp. Pl.* 439.—Sibir. ; Am. arct.
medium, Crantz, Inst. ii. 401 = Spergularia media.
melanandrum, *Maxim. in Bull. Acad. Pétersb.* xxvi. (1880) 429.—China.
membranaceum, Jacqnem. ex Hook. f. Fl. Brit. Ind. i. 228 = glomeratum ?
Meratii, *Jullien, in Mém. Soc. Acad. Maine-et-Loire*, xii. (1862) 67.—Gallia.

CERASTIUM:—
Meyerianum, *Rupr. Fl. Cauc.* 221, in textu.—Reg. Caucas.
mieranthum, *C. A. Mey. ex Heynh. Nom.* ii. 133, nomen. —Rossia.
mierocarpum, *Kit. ex Reichb. Fl Germ. Excurs.* 799. —Croatia.
microphyllum, Schur, Enum. Pl. Transs. 123 = alpinum.
microspermum, *C. A. Mey. Verz. Pfl. Cauc.* 222.— Reg. Caucas.
minus, Schur, in Oestr. Bot. Zeitschr. xxi. (1871) 46 = triviale.
minutulum, Desmoul. ex Steud. Nom. ed. II. i. 329 = glomeratum.
moesiacum, Frivald. in Flora, xix. (1836) 435 = tomentosum.
molle, *Bartl. in Presl, Rel. Haenk.* ii. 17.—Mexic.
molle, Vill. Hist. Pl. Dauph. iii. 644 = arvense, tomentosum.
mollissimum, *Poir. Encyc. Suppl.* ii. 164.—Peruv.
montanum, *C. Gay, Fl. Chil.* i. 275.—Chili.
montioides, *C. Gay, l. c.* 279.—Chili.
mosanum, *Cloet, ex Steud. Nom.* ed. II. i. 329.— Belgium.
mucronatum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 294.—Peruv.; Boliv.
multiflorum, *C. A. Mey. Verz. Pfl. Cauc.* 222.—Reg. Caucas.
murale, Desp. in DC. Fl. Fr. vi. 609 = triviale.
murale, Schur, Enum. Pl. Transs. 119 = glutinosum.
murale, Salisb. Prod. 300 = Holosteum umbellatum.
mutabile, Gren. in Mém. Soc. Emul. Doubs, i. (1841) 68 = arvense.
mutabile, Spenn. Fl. Friburg. iii. 847 = triviale, glomeratum, brachypetalum, semidecandrum.
nemorale, *Bieb. Fl. Taur. Cauc.* iii. 317.—Reg. Caucas.
nemorum, Crantz, Inst. ii. 401 = Stellaria nemorum.
nepalense, Wall. Cat. n. 628 = glomeratum?
nervosum, *C. Gay, Fl. Chil.* i. 277.—Chili.
nigrescens, Edmondst. ex Syme, Engl. Bot. ed. III. ii. 87 = alpinum.
nitens, Stev. ex Ser. in DC. Prod. i. 415 = nemorale.
nivale, D. Don, ex Nym. Consp. 110 = trigynum.
nutans, *Rafin. Préc. Découv.* 36.—Am. bor.
oblongifolium, Anders. Cat. Pl. Nevad. 118 = nutans.
oblongifolium, Torr. Pacif. Rail. Rep. iv. 70 = pilosum.
oblongifolium, *Torr. Fl. U. St.* 460.—Am. bor.
obscurum, Chaub. in St. Amans, Fl. Agen. 180 = glutinosum, *Fries*.
obtusatum, *Kit. in Linnaea*, xxxii. (1863) 523.— Hungar.
obtusifolium, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 393.—Sibir. altaic.
obtusifolium, Lam. Fl. Fr. iii. 58 = glomeratum?
octandrum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 45 = glomeratum.
Opizii, Seidl, in Opiz, Seznam, 28 = glutinosum.
orithales, *Schlecht. in Linnaea*, xii. (1838) 209.— Mexic.
orophyllum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. (1864) 295.—Peruv.; Bolivia.
ovale, Bess. Prim. Fl. Galic. i. 294 = glutinosum.
ovale, Pers. Syn. i. 521 = glomeratum?
ovatum, *Hoppe, ex Willd. Enum. Hort. Berol.* 493.— Europ.; Reg. Caucas.
Pallasii, Vest, in Flora, iii. (1820) 356 = arvense.
pallens, F. Schultz, Arch. Fl. 124 = litigiosum.
pallidum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 146 = semidecandrum.
palustre, Moris, in Mem. Accad. Torino, xxxviii. (1835) p. xxviii = campanulatum.
pauciflorum, Bunge, in Ledeb. Fl. Alt. ii. 176 = lithospermifolium.
pauciflorum, Stev. ex Ser. in DC. Prod. i. 414 = pilosum.
pauciflorum, Turcz. ex Steud. Nom. ed. II. i. 330 = alpinum.
pedunculare, *Bory & Chaub. Nouv. Fl. Pélop.* 29. t. 13.—Graecia; As. Min.
pedunculatum, Gaud. ex Ser. in DC. Prod. i. 419 = latifolium.
pedunculatum, Willd. ex Steud. Nom. ed. II. i. 330 = nutans.
pellucidum, Chaub. in St. Amans, Fl. Agen. 181 = semidecandrum.
pensylvanicum, Hook. Fl. Bor. Am. i. 104 = oblongifolium.
pensylvanicum, Hornem. Hort. Hafn. i. 435 = arvense.

CERASTIUM:—
pentandrum, Bieb. Fl. Taur. Cauc. i. 359 = glutinosum.
pentandrum, Krock. Fl. Siles. ii. 1. 81 = semidecandrum.
pentandrum, *Linn. Sp. Pl.* 438.—Europ.
pentandrum, Moris, Fl. Sard. i. 265 = pumilum.
pentandrum, Tenore, Syll. 220 = densiflorum.
peplidifolium, Kotschy, ex Boiss. Fl. Orient. i. 725 = fragillimum.
perfoliatum, Falk, Beitr. ii. 182 = davuricum.
perfoliatum, *Linn. Sp. Pl.* 437.—Reg. Mediterr.; As. Min.
persicum, *Boiss. Diagn.* Ser. I. i. 54.—Persia.
petiolare, Hance, in Hook. Kew Journ. i. (1849) 143 = Stellaria aquatica.
petraeum, F. Schultz, Arch. Fl. 124 = glutinosum.
petricola, *Panč. Elem. Fl. Bulg.* 20.—Bulgaria.
petrosum, Schur, Enum. Pl. Transs. 122 = ovatum.
pilosissimum, *Kit. in Linnaea*, xxxii. (1863) 519.— Hungar.
pilosum, Fisch. ex Ledeb. Fl. Ross. i. 409 = glomeratum.
pilosum, Hornem. Hort. Hafn. ii. 965 = grandiflorum?
pilosum, *Ledeb. in Mém. Acad. Pétersb.* v. (1815) 539. —Sibir.; Am. bor. occ.
pilosum, Sibth. & Sm. Fl. Graec. Prod. i. 316 = illyricum.
pilosum, Tenore, Prod. Fl. Nap. p. xxvii = brachypetalum.
Pohlianum, Fenzl, ex Rohrb. in Mart. Fl. Bras. xiv. II. 282 = dicrotrichum.
polymorphum, *Rupr. Fl. Cauc.* 237.—Reg. Caucas.
polyphyllum, Stev. ex Ledeb. Fl. Ross. i. 413 = arvense.
praecox, Tenore, Prod. Fl. Nap. p. xxvii = campanulatum.
procumbens, *Forsk. Fl. Aegypt. Arab.* 211.—Afr. bor.
prostratum, Jan, Elench. 7, ex Bertol. Fl. Ital. iv. 756 = campanulatum.
pseudo-sylvaticum, Schur, in Oestr. Bot. Zeitschr. xxi. (1871) 46 = triviale.
pseudo-viscosum, Schur, l. c. 99 = glomeratum.
pubescens, Goldie, in Edinb. Phil. Journ. vi. (1822) 327 = oblongifolium.
pumilum, Bourg. ex Willk. & Lange, Prod. Fl. Hisp. iii. 634 = Gayanum.
pumilum, *Curt. Fl. Lond. fasc.* vi. t. 30.—Europ.
pumilum, Griseb. Spicil. Fl. Rumel. i. 208 = glutinosum.
pumilum, Rafin. in Am. Monthly Mag. (1818) 266 = glomeratum.
purpurascens, *Adams, in Weber & Mohr, Beitr.* i. 60.— As. Min.; Reg. Caucas.
purpurascens, Fenzl, Verbreit. Alsin. tab. ad p. 56 = Arenaria purpurascens.
pusillum, Ser. in DC. Prod. i. 418 = glomeratum.
pyrenaicum, *J. Gay, in Ann. Sc. Nat.* Sér. I. xxvi. (1832) 231.—Mont. Pyren.
quaternellum, *Fenzl, Verbreit. Alsin.* tab. ad p. 56.— Europ.; Am. bor.
raceemosum, *Bartl. in Presl, Rel. Haenk.* ii. 18.—Chili.
radians, Crantz, Inst. ii 401 = Stellaria radians.
ramigerum, *Bartl. in Presl, Rel. Haenk.* ii. 16.— Mexic.
ramosissimum, Boiss. Elench. 23 = Riaei.
rectum, *Frivald. in Flora*, xix. (1836) 435.—Rumelia.
refractum, All. Fl. Pedem. ii. 117 = trigynum.
repens, Bertol. Fl. Ital. iv. 757 = sylvaticum.
repens, Bieb. Fl. Taur. Cauc. i. 360 = Biebersteinii.
repens, Boiss. Voy. Bot. Espagne, 105 = Boissieri.
repens, Linn. Sp. Pl. 439 = tomentosum.
revolutum, *Meinsh. in Baer & Helm. Beitr. Kenntn. Russ. Reich.* xxvi. 146.—Ins. Koraginsk.
Riaei, *Des Moul. ex J. Gay, in Ann. Sc. Nat.* Sér. II. vi. (1836) 348.—Europ.; As. Min.
rigidum, Ledeb. in Mém. Acad. Pétersb. v. (1815) 538 = alpinum.
rigidum, Vitm. Summa Veg. iii. 137 = arvense.
Ripartianum, *F. Schultz, in Flora*, xlv. (1862) 458.— Mexic.
rivulare, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 166.— Bras.
Roeseri, Boiss. & Heldr. Diagn. Ser. II. i. 93 = brachypetalum.
rotundatum, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 146 = semidecandrum.
rotundifolium, Fisch. Cat. Hort. Gorenk. (1812) 58 = glomeratum.
rotundifolium, Schur, Enum. Pl. Transs. 123 = alpinum.
ruderale, *Bieb. Fl. Taur. Cauc.* i. 357.—Ross.; Reg. Caucas.

CERASTIUM:—
rupestre, Fisch. ex Ser. in DC. Prod. i. 417 = trigynum.
salatavicum, *Rupr. Fl. Cauc.* 229.—Reg. Caucas.
samnianum, Ser. in DC. Prod. i. 421 = grandiflorum.
saxatile, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 616 = alpinum.
saxigenum, Schur, in Oestr. Bot. Zeitschr. xxi. (1871) 100 = glutinosum.
scandens, *Lejeune, Fl. Spa*, i. 211.—Belgium.
scaposum, *Boiss. & Heldr. Diagn.* Ser. I. viii. 104.— Ins. Creta.
Scarani, Tenore, Prod. Fl. Nap. p. xxvii = arvense.
Selloi, *Schlecht. ex Rohrb. in Mart. Fl. Bras.* xiv. II. 283.—Bras.
semidecandrum, Cham. & Schlecht. in Linnaea, v. (1830) 233 = vulcanicum.
semidecandrum, Chaub. Expéd. Sc. Morée, 129 = brachypetalum.
semidecandrum, *Linn. Sp. Pl.* 438.—Europ.; As. Min.
semidecandrum, Sibth. & Sm. Fl. Graec. Prod. i. 315 = glutinosum.
sericeum, *S. Wats. in Proc. Am. Acad.* xx. (1885) 354. —Arizona.
sericeum, *Pourr. in Mém. Acad. Toul.* iii. (1788) 311. —Europ.
serpyllifolium, Bieb. ex Ser. in DC. Prod. i. 417 = glomeratum.
serpyllifolium, Willd. Enum. Hort. Berol. Suppl. 26 = arvense.
sibiricum, Stev. ex. Ledeb. Fl. Ross. i. 407 = alpinum.
sibiricum, Turcz. ex Ledeb. l. c. 410 = glomeratum.
siculum, *Guss. Suppl. Fl. Sic. Prod.* 137.—Sicilia.
simense, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 45 = glomeratum.
Soleirolii, Ser. ex Duby, Bot. Gall. i. 87 = arvense.
soratense, *Rohrb. in Linnaea*, xxxvii. (1871–73) 291. —Bolivia.
spathulatum, *Pers. Syn.* i. 520.—Jamaica.
speciosum, Sprun. ex Boiss. Fl. Orient. i. 727 = grandiflorum.
Sprengelii, *Ser. in DC. Prod.* i. 421.—Hab.?
squalidum, Ram. in Mém. Acad. Paris, vi. (1826) 158 = alpinum.
Stellaria, *Tenore, ex Steud. Nom.* ed. II. i. 330.— Neapol.
stellarioides, Hartm. Handb. Skand. Fl. ed. I. 181 = trigynum.
stellarioides, Moç. ex Ser. in DC. Prod. i. 415 = pilosum.
stenopetalum, *Fenzl, ex Gren. & Godr. Fl.* i. 272.—Ins. Corsica.
strictum, Linn. Sp. Pl. 439 = arvense.
strigosum, Fries, Fl. Halland. 78 = brachypetalum.
subacaule, Hegetschw. in Suter, Fl. Helvet. ed. II. App. 455 = latifolium.
subspicatum, *Wedd. in Ann. Sc. Nat.* Sér. V. i. 1864 295.—Peruv.
succulentum, Crantz, Inst. ii. 400 = Arenaria peploides.
suffruticosum, *Lam. Encyc.* i. 681.—Gallop.; As. Min.
suffruticosum, Linn. Sp. Pl. 439 = arvense.
sylvaticum, Opiz, Seznam, 28 = triviale.
sylvaticum, Stev. ex Ledeb. Fl. Ross. i. 410 = glomeratum.
sylvaticum, *Waldst. & Kit. Pl. Rar. Hung.* i. 100.— Austria.
szalaberense, *Kit. in Linnaea*, xxxii. (1863) 524.— Hungar.
Szowitsii, *Boiss. Fl. Orient.* i. 717.—Armenia.
tauricum, Spreng. ex Ser. in DC. Prod. i. 415 = brachypetalum.
Tauschianum, *Wolfner, in Lotos*, iv. (1854) 57.— Europ.
tenellum, Gaud. ex Ser. in DC. Prod. i. 416 = glomeratum.
Tenoreanum, Ser. l. c. 421 = brachypetalum.
tenue, Viviani, Fl. Cors. Sp. Nov. 7 = quaternellum.
tenuifolium, Pursh, Fl. Am. Sept. i. 321 = arvense.
tenuifolium, Spreng. in Hornem. Hort. Hafn. Suppl. 138 = Sprengelii.
tenuifolium, Visiani, ex Reichb. Fl. Germ. Excurs. 799 = grandiflorum.
tetragonum, Baumg. Enum. Stirp. Transs. i. 424 = Stellaria crassifolia.
tetrandrum, *Curt. Fl. Lond. fasc.* vi. t. 31.—Europ.
Thomasii, *Tenore, Fl. Neap. Prod. App.* iv. 21.— Europ.
Thomsonii, *Hook. f. Fl. Brit. Ind.* i. 228.—Reg. Himal.
tmoleum, Boiss. Diagn. Ser. I. v. 86 = fragillimum.

CERASTIUM:—
tomentosum, Boj. Hort. Maurit. 24 = glomeratum.
tomentosum, Georgi, Beschr. Russ. Reich. iv. 988 = Biebersteinii.
tomentosum, Haensel. ex Willk. & Lange, Prod. Fl. Hisp. iii. 636 = Boissieri.
tomentosum, Huds. Fl. Angl. ed. I. 176 = alpinum.
tomentosum, *Linn. Sp. Pl.* 440.—Europ.; As. Min.
Tournefortii, *Gren. in Mém. Soc. Emul. Doubs*, i. (1841) 11. t. 7.—Armenia.
transsilvanicum, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 14.—Transsylv.
trigynum, *Vill. Prosp.* 48; *Hist. Pl. Dauph.* iii. 645.—Reg. bor. temp. et arct.
triviale, *Link, Enum. Hort. Berol.* i. 433.—Europ.; As.; Afr. bor.
uliginosum, Hegetschw. Fl. Schweiz, 437 = triviale?
umbellatum, Crantz, Inst. ii. 401 = Holosteum umbellatum.
umbellatum, *C. Koch, in Linnaea*, xv. (1841) 708.—Armenia.
umbrosum, Kit. in Linnaea, xxxii. (1863) 515 = triviale?
uniflorum, Thom. ex Reichb. Fl. Germ. Excurs. 797 = latifolium.
vagans, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimp. 26.—Ins. Madera.
valachicum, *Kit. in Linnaea*, xxxii. (1863) 519.—Hungar.
varians, Coss. & Germ. Obs. Pl. Crit. 26 = glutinosum *Fries*, semidecandrum.
velutinum, Rafin. in Med. Repos. N. York, v. (1808) 359 = vulgatum?
Vidali, *Phil. in Anal. Univ. Chil.* (1872) 677.—Chili.
villosum, Baumg. Enum. Stirp. Transs. i. 424 = alpinum.
villosum, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) i. 322 = glomeratum.
viscarium, Schur, Enum. Pl. Transs. 118, in syn. = glutinosum.
viscidum, Link, Enum. Hort. Berol. i. 433 = semidecandrum.
viscosissimum, Schur, Enum. Pl. Transs. 123 = alpinum.
viscosum, Linn. Sp. Pl. 437 = glomeratum.
viscosum, Pollich, Hist. Pl. Palat. i. 448 = brachypetalum.
vulcanicum, *Schlecht. in Linnaea*, xii. (1838) 208.—Mexic.
vulgare, Hartm. Handb. Skand. Fl. ed. I. 182 = triviale.
vulgatum, Linn. Fl. Suec. ii. 158 = glomeratum.
Wahlenbergii, Rupr. Fl. Caucas. 237 = latifolium.
Willdenowii, *H. B. & K. Nov. Gen. et Sp.* vi. 29.—Ecuador.

CERASUS, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 141 (1737) = **Prunus**, Linn. (Rosac.).
acida, Borck. in Roem. Arch. i. ii. 38 = P. Cerasus.
acuminata, Wall. Pl. As. Rar. ii. 78 = P. acuminata.
adenophylla, M. Roem. Syn. Rosifl. 81 = P. undulata.
affinis, Sweet, Hort. Brit. ed. III. 209.—Hab.?
americana, Hook. & Arn. in Comp. Bot. Mag. i. (1835) 24 = P. americana.
aspera, Loisel. in Duham. Arb. ed. nov. v. 33 = P. aspera.
asplenifolia, Hort. ex C. Koch, Dendrol. i. 106 = P. Avium.
austera, Borck. in Roem. Arch. i. ii. 38 = P. Cerasus.
avicularis, Dulac, Fl. Hautes-Pyr. 301 = P. Avium.
Avium, Moench, Meth. 672 = P. Avium.
beroliensis, Nym. Consp. 213 = P. Cerasus.
bigarella, Dum. Fl. Belg. 91 = P. Cerasus.
borealis, Michx. Fl. Bor. Am. i. 286 = P. pensylvanica.
brachypetala, Boiss. Diagn. Ser. I. vi. 52 = *incisa*.
brasiliensis, Cham. & Schlecht. in Linnaea, ii. (1827) 542 = P. brasiliensis.
Bungei, Walp. Rep. ii. 9 = P. Cerasus.
campestris, Gueldenst. ex Ledeb. Fl. Ross. ii. 7 = P. Chamaecerasus.
canadensis, Loisel. in Duham. Arb. ed. nov. v. 3 = P. americana.
canadensis, Mill. Gard. Dict. ed. VIII. n. 5 = P. Chamaecerasus.
Capollin, Ser. in DC. Prod. ii. 539 = P. Capollin.
capricida, Wall. Cat. n. 718 = P. undulata.
Caproniana, DC. Fl. Fr. iv. 482 = P. Cerasus.

CERASUS:—
Capuli, Ser. in DC. Prod. ii. 541 = Prunus Capollin.
caroliniana, Michx. Fl. Bor. Am. i. 285 = P. caroliniana.
cerasifera, Eaton & Wright, N. Am. Bot. 189 = P. maritima.
Chamaecerasus, Loisel. in Duham. Arb. ed. nov. v. 29 = P. Chamaecerasus.
Chicasa, Ser. in DC. Prod. ii. 538 = P. Chicasa.
chinensis, G. Don, Gen. Syst. ii. 514 = P. japonica.
collina, Lej. & Court. Comp. Fl. Belg. ii. 130 = P. Cerasus.
cornuta, Wall. Cat. n. 716 = P. Padus.
corymbosa, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = P. Mahaleb.
cucullata, Hort. ex C. Koch, Dendrol. i. 111 = P. Cerasus.
decumana, Mordant, ex DC. Prod. ii. 536 = P. Avium.
demissa, Nutt. in Torr. & Gray, Fl. N. Am. i. 411 = P. demissa.
densiflora, Spach, Hist. Vég. Phan. i. 415 = P. virginiana.
depressa, Ser. in DC. Prod. ii. 538 = P. pumila.
diffusa, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 647.—Persia.
domestica, Cat. Perck, ex Wesmael, in Bull. Congr. Bot. Brux. (1864) 255 = P. Cerasus.
Duerinckii, Mart. Bull. Acad. Belg. viii. i. (1841) 68 = P. virginiana.
dulcis, Borck. ex Steud. Nom. ed. II. i. 331 = P. Avium.
dulcis, Gaertn. Mey. & Scherb. Fl. Wett. ii. 181 = P. Cerasus.
duracina, DC. Fl. Fr. iv. 483 = P. Avium.
ebroliensis, Lamotte, Prod. i. 238 = P. Cerasus.
effusa, Host, Fl. Austriac. ii. 6 = P. Cerasus.
elliptica, Loisel. in Duham. Arb. ed. nov. v. 4 = P. elliptica.
emarginata, Dongl. ex Hook. Fl. Bor. Am. i. 169 = P. emarginata.
erecta, Presl, Epim. Bot. 194 = P. emarginata.
ferruginea, DC. Prod. ii. 540 = P. ferruginea.
fimbriata, Spach, Hist. Vég. Phan. i. 416 = P. virginiana.
floribunda, Hort. ex C. Koch, Dendrol. i. 119 = P. pumila.
Fontanesiana, Spach, Hist. Vég. Phan. i. 410 = P. graeca.
frutĭcosa, Pall. Fl. Ross. t. 8^{b}. = P. Chamaecerasus.
glandulosa, Kellogg, in Proc. Calif. Acad. i. 1855 (1873) 59 = P. emarginata.
glandulosa, Loisel. in Duham. Arb. ed. nov. v. 33 = P. japonica.
glauca, Moench, Meth. 672 = P. pumila.
glaucifolia, Wall. Cat. n. 717 = P. nepaulensis.
graeca, Hort. ex C. Koch, Dendrol. i. 705, Index = P. graeca.
Griffithii, Boiss. Fl. Orient. ii. 648.—Afghanist.
Heaumiana, M. Roem. Syn. Rosifl. 69 = P. Cerasus.
heterophylla, Hort. ex C. Koch, Dendrol. i. 106 = P. Avium.
hirsuta, Spach, Hist. Vég. Phan. i. 417 = P. virginiana.
Hixa, C. Sm. ex Spach, Hist. Nat. Veg. i. 419 = P. Hixa.
hortensis, Mill. Gard. Dict. ed. VIII. n. 3 = P. Cerasus.
humilis, Host, Fl. Austr. ii. 7 = P. Chamaecerasus.
humilis, Moris, Elench. Sard. 17 = P. prostrata.
hyemalis, Ser. in DC. Prod. ii. 538 = P. americana.
ilicifolius, Nutt. ex Hook. & Arn. Bot. Beech. Voy. 340. t. 83 = P. ilicifolia.
incana, Boiss. Fl. Orient. ii. 647 = P. incana.
incana, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 126 = P. prostrata.
incisa, Boiss. Fl. Orient. ii. 648.—Persia.
incisa, Loisel. in Duham. Arb. ed. nov. v. 33 = P. incisa.
integerrima, Wall. Cat. n. 722 = P. acuminata.
integrifolia, Presl, Epim. Bot. 194.—Peruv.
intermedia, Host, Fl. Austriac. ii. 7 = P. Avium.
intermedia, Loisel. in Duham. Arb. ed. nov. v. 30 = P. Chamaecerasus.
japonica, Ker-Gawl. ex Steud. Nom. ed. II. i. 331 = P. Kerii.
japonica, Loisel. in Duham. Arb. ed. nov. v. 33 = P. japonica.
javanica, Teijsm. & Binn. in Nederl. Kruidk. Arch. iii. (1855) 412 = P. javanica.
Jenkinsii, Hook. f. & Thoms. ex Fl. Brit. Ind. ii. 317 = P. Jenkinsii.
Juliana, DC. Fl. Fr. iv. 482 = P. Avium.

CERASUS:—
Juliana, Hort. ex Steud. Nom. ed. II. i. 331 = Prunus Cerasus.
Launesiana, Carr. in Rev. Hortic. (1872) 198.—Japon.
Laurocerasus, Loisel. in Duham. Arb. ed. nov. v. 6 = P. Laurocerasus.
Lindleyana, Wall. Cat. n. 1008 = P. nepaulensis.
littoralis, Eaton & Wright, N. Am. Bot. 189 = P. maritima.
lusitanica, Loisel. in Duham. Arb. ed. nov. v. 5 = P. lusitanica.
macedonica, M. Roem. Syn. Rosifl. 64 = P. Chamaecerasus.
macrophylla, Sweet, Hort. Brit. ed. I. 485 = P. Avium.
maeda, Hort. ex Lavall. Arbor. Segrez. 119 = P. paniculata.
Mahaleb, Mill. Gard. Dict. ed. VIII. n. 4 = P. Mahaleb.
Marasca, Host, Fl. Austriac. ii. 6 = P. Cerasus.
martabanica, Wall. Cat. n. 4902 = P. martabanica.
micrantha, Spach, Hist. Vég. Phan. i. 414 = P. virginiana.
microcarpa, Boiss. Fl. Orient. ii. 646 = P. microcarpa.
minutiflora, A. Gray, Pl. Wright. i. 68 = P. minutiflora.
mollis, Dougl. in Hook. Fl. Bor. Am. i. 169 = P. emarginata.
multicarpa, Hort. ex Rev. Hortic. (1875) 409 = P. Cerasus.
napaulensis, Ser. in DC. Prod. ii. 540 = P. nepaulensis.
nicotianaefolia, Hort. ex DC. Prod. ii. 536 = P. Cerasus.
nigra, Loisel. in Duham. Arb. ed. nov. v. 32 = P. americana.
nigra, Mill. Gard. Dict. ed. VIII. n. 2 = P. Avium.
nigricans, Borck. in Roem. Arch. i. ii. 38 = P. Avium.
obovata, Beck, N. et M. St. Bot. 37 = P. virginiana.
occidentalis, Loisel. in Duham. Arb. ed. nov. v. 4 = P. occidentalis.
opaca, Benth. Pl. Hartw. 172 = P. opaca.
orientalis, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 128 = P. microcarpa.
oxypyrena, Spach, Hist. Vég. Phan. i. 413 = P. Padus.
Padus, Delarb. Fl. Auv. ed. II. 323; DC. Fl. Fr. iv. 480 = P. Padus.
pallida, M. Roem. Syn. Rosifl. 69 = P. Avium.
paniculata, Loisel. in Duham. Arb. ed. nov. v. 7 = P. paniculata.
Pattoniana, Carr. in Rev. Hortic. (1872) 135 = P. mollis.
pectinata, Spach, in Ann. Sc. Nat. Sér. II. xix. (1843) 127 = P. pectinata.
pendula, Sweet, Hort. Brit. ed. I. 485 = P. fruticosa,
pensylvanica, Loisel. in Duham. Arb. ed. nov. v. 9 = P. pensylvanica.
persicifolia, Loisel. l. c. = P. pensylvanica.
Phoshia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 239 = P. Puddum.
polygyna, DC. ex Ser. in DC. Prod. ii. 537 = P. Cerasus.
prostrata, Hort. ex C. Koch, Dendrol. i. 119 = P. pumila.
prostrata, Ser. in DC. Prod. ii. 538 = P. prostrata.
Pseudocerasus, G. Don, in Loud. Hort. Brit. 200 = P. Puddum.
pubescens, Ser. in DC. Prod. ii. 538 = P. maritima.
Puddum, Wall. Pl. As. Rar. ii. 37. t. 143 = P. Puddum.
pumila, Borck. in Roem. Arch. i. ii. (1797) 38 = P. Chamaecerasns.
pumila, Michx. Fl. Bor. Am. ii. 286 = P. pumila.
punctata, Hook. f. & Thoms. ex Fl. Brit. Ind. ii. 317 = P. punctata.
pygmaea, Loisel. in Duham. Arb. ed. nov. v. 32 = P. maritima.
racemosa, Gilib. Exercit. i. 349 = P. Padus.
reflexa, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 342 = P. reflexa.
Rhexii, Hort. Gall. ex Van Houtte, Fl. des Serres, Sér. II. vii. (1867–68) 159 = P. Cerasus.
rosea, Hort. ex Steud. Nom. ed. II. i. 331 = P. Cerasus.
rubicunda, Bechst. Forstb. 160, 355 = P. Avium.
rubra, Gilib. Fl. Lituan. ii. 229 = P. Cerasus.
rufa, Wall. Cat. n. 721 = P. rufa.
salicifolia, Hort. ex C. Koch, Dendrol. i. 106 = P. Avium.
salicifolia, Ser. in DC. Prod. ii. 540 = P. salicifolia.

CERASUS :—
salicina, [G. Don, in] Loud. Hort. Brit. 480 = Prunus salicina.
samydoides, Walp. Rep. iv. Index, 311 = P. samydoides.
semperflorens, Borckh. in Roem. Arch. i. II. 38 = P. Cerasus.
serotina, Hook. Fl. Bor. Am. 1. 169 = P. virginiana.
serotina, Loisel. in Duham. Arb. ed. nov. v. 3 = P. serotina.
serrulatus, [G. Don, in] Loud. Hort. Brit. 480 = P. paniculata.
sibirica, Hort. ex C. Koch, Dendrol. i. 114 = P. fruticosa.
Sieboldii, Carr. in Rev. Hortic. (1866) 279. ; 370 = P. paniculata.
sinensis, Spach, Hist. Vég. Phan. i. 425 = P. sinensis.
sphaerocarpa, Loisel. in Duham. Arb. ed. nov. v. 4 = P. sphaerocarpa.
Susquehannae, Sweet, Hort. Brit. ed. I. 485 = P. pumila.
sylvestris, S. G. Gmel. It. i. 102 = P. Chamaecerasus.
tomentosa, Wall. Cat. n. 715 = P. tomentosa.
tortuosa, Boiss. & Haussk. ex Boiss. Fl. Orient. ii. 647. —As. Min. ; Persia.
tridentina, M. Roem. Syn. Rosifl. 76 = P. Cerasus.
triflora, Wall. ex Steud. Nom. ed. II. i. 331 = P. triflora.
umbellata, Torr. & Gray, Fl. N. Am. i. 409 = Prunus umbellata.
undulata, Ser. in DC. Prod. ii. 540 = P. undulata.
varia, Borck. in Roem. Arch. i. II. 38 = P. Avium.
virginiana, Michx. Fl. Bor. Am. i. 285 = P. serotina.
virginica, Michx. ex Steud. Nom. ed. II. i. 331 = P. virginiana.
vulgaris, Mill. Gard. Dict. ed. VIII. n. 1 = P. Cerasus.
Wallichii, M. Roem. Syn. Rosifl. 81 = P. acuminata.
Wattererii, Hort. ex Lavall. Arbor. Segrez. 119 = P. Puddum.

CERATANDRA, Eckl. ex Bauer, Illustr. Gen. Orch. t. 16 (1837). *ORCHIDEAE*, Benth. & Hook. f. iii. 634.
affinis, *Sond. in Linnaea*, xix. (1847) 108.—Afr. austr.
auriculata, Lindl. Gen. et Sp. Orch. 364 = chloroleuca.
bicolor, *Sond. ex Drège, in Linnaea*, xx. (1847) 220, nomen.—Afr. austr.
chloroleuca, *Eckl. ex Bauer, Illustr. Gen. Orch.* t. 16. —Afr. austr.
globosa, *Lindl. Gen. et Sp. Orch.* 364.—Afr. austr.
grandiflora, *Lindl. l. c.*—Afr. austr.
Harveyana, *Lindl. l. c.* 365.—Afr. austr.
Harveyana, Sond. in Linnaea, xix. (1847) 108 = bicolor.
ochroleuca, G. Don, in Sweet, Hort. Brit. ed. III. 651 = chlorolcuca.
parviflora, *Lindl. Gen. et Sp. Orch.* 364.—Afr. austr.

CERATANTHERA, Hornem. Hort. Hafn. i. 2, 5 (1813) ; Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 335, 341 = **Globba**, Linn. (Scitamin.).
amomoides, Hornem. l. c. i. 5 = G. marantina.
pendula, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 335 = G. pendula.

CERATELLA, Hook. f. Fl. Antarct. i. 25. t. 18 (1845) = **Abrotanella**, Cass. (Compos.).
rosulata, Hook. f. l. c. = A. rosulata.

CERATIA, Adans. Fam. ii. 319 (1763) = **Ceratonia**, Linn. (Legumin.).

CERATIOLA, Michx. Fl. Bor. Am. ii. 221 (1803). *EMPETRACEAE*, Benth. & Hook. f. iii. 415.
ericoides, *Michx. l. c.* 222.—Am. bor.

CERATIOSICYOS, Nees, in Eckl. & Zeyh. Enum. 281 (1836). *PASSIFLOREAE*, Benth. & Hook. f. i. 814.
Ecklonii, *Nees, l. c.*—Afr. austr.

CERATOSYCIOS, Walp. Rep. ii. 222 (1843) = pracc.

CERATIUM, Blume, Bijdr. 341. t. 46 (1825) = **Eria**, Lindl. (Orchid.).
compressum, Blume, l. c. 342 = E. compressa.

CERATOCALYX, Coss. in Ann. Sc. Nat. Sér. III. ix. (1848) 145. t. 10 = **Orobanche**, Linn.
fimbriata, Lange, in Kjoeb. Vidensk. Meddel. (1863) 52.—Hispan.
macrolepis, Coss. in Ann. Sc. Nat. Sér. III. ix. (1848) 146. t. 10 = O. macrolepis.
Reuteri, Lange, in Kjoeb. Vidensk. Meddel. (1863) 52, in obs. = O. gamosepala.

CERATOCAPNOS, Dur. in Parlat. Giorn. Bot. Ital. i. I. (1844) 336 = **Corydalis**, Vent. (Papaverac.).
heterocarpa, Dur. l. c. = C. heterocarpa.
palaestina, Boiss. Diagn. Ser. I. viii. 12.—Syria.
umbrosa, Walp. Ann. ii. 30 = Cor. heterocarpa.

CERATOCARPUS, Buxb. ex Linn. Diss. Dass. 28 (1747) ; Amoen. Acad. i. 412 (1749). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 55.
arenarius, *Linn. Sp. Pl.* 969.—Sibir.
maritimus, Pall. Ind. Taur. ex Bieb. Fl. Taur. Cauc. ii. 397 = Atriplex pedunculata.
salinus, Pall. Reise Statth. Russ. Reich. i. 268, 270 = Atriplex pedunculata.

CERATOCARPUS, Dur. (lapsu) in Parlat. Giorn. Bot. Ital. i. I. (1844) 336 = CERATOCAPNOS, Dur. = **Corydalis**, Vent.

CERATOCARYUM, Nees, in Lindl. Nat. Syst. ed. II. 451 (1836). *RESTIACEAE*, Benth. & Hook. f. iii. 1037.
argenteum, *Nees, l. c.* ; *Kunth, Enum. Pl.* iii. 483.—Afr. austr.
fistulosum, *Mart. in Journ. Linn. Soc.* x. (1869) 274. —Afr. austr.
speciosum, Nees, ex Mast. l. c. 273 = argenteum.

CERATOCAULOS, Reichb. Handb. 201 (1837) = **Datura**, Linn. (Solanac.).
daturoides, Spach, Hist. Vég. Phan. ix. 69 = D. Ceratocaula.

CERATOCEPHALUS, Moench, Meth. 218 (1794) (Ceratocephala) = **Ranunculus**, Linn.
falcatus, Pers. Syn. i. 341 = R. falcatus.
furfurascens, Pomel, Nouv. Mat. Fl. Atl. 248.—Afr. bor.
incurvus, Stev. in Bull. Soc. Nat. Mosc. xxi. II. (1848) 269. t. 7 = R. falcatus.
leiocarpus, Stev. l. c. = R. falcatus.
orthoceras, DC. Syst. i. 231 = R. falcatus.
platyceras, Stev. in Bull. Soc. Nat. Mosc. xxi. II. (1848) 269 = R. falcatus.
reflexus, Stev. l. c. 268 = R. falcatus.
spicatus, Moench, Meth. 218 = R. falcatns.
syriacus, Stev. in Bull. Soc. Nat. Mosc. xxi. II. (1848) 268 = R. falcatus.
testiculatus, Roth, Enum. Pl. Phan. Germ. i. I. 1014 = R. falcatus.

CERATOCEPHALUS, Vaill. ex Cass. in Dict. Sc. Nat. vii. 432 (1817) = **Bidens**, Linn. (Compos.).
pilosus, Rich. ex Cass. l. c. = B. pilosa.

CERATOCHILUS, Blume, Bijdr. 358. t. 25 (1825) = **Saccolabium**, Blume (Orchid.).
biglandulosus, Blume, l. c. 359 = S. distichum ?
micranthus, Lindl. Gen. et Sp. Orch. 232 = S. micranthum.
orchideus, Lindl. l. c.—Ind. or.

CERATOCHILUS, Lindl. in Lodd. Bot. Cab. t. 1414 (1828) = **Stanhopea**, Frost (Orchid.).
grandiflorus, Lodd. l. c. = S. eburnea.
insignis, Lindl. ex Steud. Nom. ed. II. i. 331 = S. insignis.
oculatus, Lodd. Bot. Cab. t. 1764 = S. oculata.

CERATOCHLOA, Beauv. Agrost. 75. t. 15. f. 7 (1812) = **Bromus**, Dill. (Gramin.).
australis, Spreng. ex Steud. Nom. ed. II. i. 332 ; 229 = B. unioloides.
Boryana, Roem. & Schult. Syst. ii. 597 = B. strictus.
breviaristata, Hook. Fl. Bor. Am. ii. 253 = B. breviaristatus.
brevis, Nees, ex Steud. Syn. Pl. Gram. 326 = B. brevis.
festucoides, Beauv. Agrost. 75 = B. unioloides.

CERATOCHLOA :—
grandiflora, Hook. Fl. Bor. Am. ii. 253 = Bromus virens.
Haenkeana, J. Presl, in Rel. Haenk. i. 285 = B. unioloides.
marginata, Nees, ex Steud. Syn. Pl. Gram. 322 = B. ciliatus.
pendula, Schrad. Ind. Sem. Hort. Gotting. (1830) ; Linnaea, vi. (1831) Litt. 72 = B. unioloides.
secunda, J. Presl, in Rel. Haenk. i. 285 = B. Preslii.
unioloides, DC. Cat. Hort. Monsp. 92 = B. unioloides.

CERATOCNEMUM, Coss. & Balansa, in Bull. Soc. Bot. Fr. xx. (1873) 239. *CRUCIFERAE*.
rapistroides, *Coss. & Balansa, l. c.*—Marocco.

CERATOCOCCA, Willd. ex Roem. & Schult. Syst. vi. p. lxx (1820) = **Microtea**, Sw. (Phytolaccac.).
maypurensis, Humb. & Bonpl. ex Roem. & Schult. l. c. 800 = M. maypurensis.

CERATOCOCCUS, Meissn. Gen. Comm. 369 (1843) = **Plukenetia**, Linn. (Euphorbiac.).

CERATOGONON, Meissn. in Wall. Pl. As. Rar. iii. 63 (1832) = **Oxygonum**, Burch. (Polygonac.).
atriplicifolium, Meissn. l. c.—Afr. trop.
cordofanum, Meissn. in DC. Prod. xiv. 39 = *atriplicifolium*.
sinuatum, Hochst. & Steud. ex A. Rich. Tent. Fl. Abyss. ii. 231 = *atriplicifolium*.

CERATOGYNE, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 68. *COMPOSITAE*, Benth. & Hook. f. ii. 431.
DIOTOSPERMA, A. Gray, in Hook. Kew Journ. iv. (1852) 275.
obionoides, *Turcz. l. c.* 69.—Austral.

CERATOGYNUM, Wight, Ic. v. II. 26. t. 1900 (1852) = **Sauropus**, Blume (Euphorbiac.).
concolor, Pritz. Ic. Ind. 245, sphalm. = Ceratolobus concolor.
rhamnoides, Wight, Ic. v. II. 26. t. 1900 = S. quadrangularis.

CERATOLACIS, Wedd. in DC. Prod. xvii. 66 (1873) *PODOSTEMACEAE*, Benth. & Hook. f. iii. 118.
erythrolichen, *Wedd. l. c.*—Bras.

CERATOLEPIS, Cass. in Bull. Soc. Philom. (1819) 111 = **Pamphalea**, DC. (Compos.).

CERATOLOBUS, Blume, in Schult. f. Syst. vii. p. lxxx (1830). *PALMAE*, Benth. & Hook. f. iii. 983.
concolor, *Blume, ex Rumphia*, ii. 165. t. 130. f. 1.—Sumatra.
glaucescens, *Blume, in Schult. f. Syst.* vii. 1334.—Java.
glaucescens, Blume, Rumphia, ii. t. 137 = Korthalsiae sp.
plicatus, Zipp. ex Blume, l. c. 171 = Korthalsia Zippelii.
Zippelii, Blume, l. c. = Korthalsia Zippelii.

CERATONIA, Linn. Gen. ed. I. 35 (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 574.
CERATIA, Adans. Fam. ii. 319 (1763).
chilensis, Molina, Sagg. Chil. 172 = Prosopis juliflora.
coriacea, Salisb. Prod. 323 = C. Siliqua.
inermis, Stokes, Bot. Mat. Med. i. 487 = C. Siliqua.
Siliqua, *Linn. Sp. Pl.* 1026.—Europ. austr. ; Orient.

CERATONYCHIA, Edgew. in. Journ. As. Soc. Beng. xvi. (1847) 1215 = **Cometes**, Linn. (Amaranthac.).
Nidus, Edgew. l. c. = Com. surattensis.

CERATOPETALUM, Sm. Bot. N. Holl. 9. t. 3 (1793). *SAXIFRAGEAE*, Benth. & Hook. f. i. 651.
apetalum, *D. Don, in Edinb. N. Phil. Journ.* ix. (1830) 94.—Austral.
gummiferum, *Sm. Bot. N. Holl.* 9. t. 3.—Austral.
monopetalum, Caley, ex D. Don, in Edinb. N. Phil. Journ. ix. (1830) 94 = apetalum.
montanum, D. Don, l. c. = apetalum.
ovatum, Caley, ex D. Don, l. c. 95 = Schizomeria ovata.

CERATOPHORUS, Hassk. Retzia, i. 100 (1856) (= Keratophorus) = **Payena**, Linn. (Sapotac.).
Leerii, Hassk. l. c. 101 = P. Leerii.
longepetiolatus, Teijsm. & Binn. Cat. Hort. Bog. 161 = P. longepetiolata.
Wightii, Hassk. Retzia, i. 101 = P. lucida ?
Wightii, Maing. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 547 = P. malaccensis.

CERATOPHORUS, Sond. in Linnaea, xxiii. (1850) 120 = **Gelonium**, Roxb. (Euphorbiac.).
africanus, Sond. l. c. 121 = G. africannm.

CERATOPHYLLUM, Linn. Syst. ed. I (1735). *CERATOPHYLLEAE*, Benth. & Hook. f. iii. 415.
DICHOTOPHYLLUM, Dill. ex Moench, Meth. 345 (1794).
apiculatum, Cham. in Linnaea, iv. (1829) 504 = demersum.
aquaticum, H. C. Wats. Topogr. Bot. i. 165 = demersum ?
asperum, Lam. Fl. Fr. ii. 196 = demersum.
australe, *Griseb. in Goett. Abh.* xxiv. (1879) 14.—Reg. Argent.
chilense, *Leyb. in Anal. Univ. Chil.* (1859) 682.—Chili.
cornutum, Rich. in Ann. Mus. Par. xvi. (1810) 299.—demersum.
cristatum, *Guill. & Perr. Fl. Seneg. Tent.* 295.—Afr. trop.
demersum, Klein, ex Cham. in Linnaea, iv. (1829) 505 = tuberculatum.
demersum, *Linn. Sp. Pl.* 992.—Reg. temp. et trop.
demersum, Sieber, ex Cham. in Linnaea, iv. (1829) 504 = muricatum.
echinatum, *A. Gray, Man. Bot. U. St.* 401.—Am. bor.
gibbum, Laforet, ex Nym. Consp. 251 = demersum.
granulosum, Schur, Enum. Pl. Transs. 218 = submersum.
Haynaldianum, *Borb. in Magy. Növént. Lapok*, viii. (1884) 20.—Hungar.
indicum, Willd. ex Cham. in Linnaea, iv. (1829) 504 = tuberculatum.
inerme, Dum. Fl. Belg. 165 = submersum.
laeve, Lam. Fl. Fr. ii. 197 = submersum.
missionis, *Wall. Cat.* sub n. 7007.—Ind. or.
muricatum, *Cham. in Linnaea*, iv. (1829) 504.—As. et Afr. trop.
muticum, Cham. l. c. 505 = submersum.
oxyacanthum, Cham. l. c. 504 = tuberculatum.
oxyacanthum, Schur, Enum. Pl. Transs. 218 = muricatum.
pentacanthum, *Haynald, in Magy. Növént. Lapok*, v. (1881) 109.—Hungar.
platyacanthum, Cham. in Linnaea, iv. (1829) 504 = muricatum.
polyacanthum, Schur, Enum. Pl. Transs. 218 = muricatum.
submersum, *Linn. Sp. Pl.* ed. II. 1409.—Europ. ; As. trop.
triacanthum, Schur, Enum. Pl. Transs. 218 = muricatum.
tricorne, Dum. Fl. Belg. 165 = demersum.
tricuspidatum, Dum. l. c. = demersum.
tuberculatum, *Cham. in Linnaea*, iv. (1829) 504.—Ind. or.
unicorne, Dum. Fl. Belg. 165 = demersum.
verrucosum, S. F. Gray, Nat. Arr. Brit. Pl. ii. 554 = submersum.
verruculosum, Rich. in Ann. Mus. Par. xvi. (1810) 299 = submersum.
verticillatum, Roxb. Hort. Beng. 68 ; Fl. Ind. iii. 624 = demersum.
vulgare, Schleid. in Linnaea, xi. (1837) 540 = muricatum, tuberculatum, submersum.

CERATOPSIS, Lindl. Gen. et Sp. Orch. 383 (1840) = **Epipogum**, S. G. Gmel. (Orchid.).
rosea, Lindl. l. c. 384 = E. roseum.

CERATOPYXIS, Hook. f. Ic. Pl. t. 1125 (1872). *RUBIACEAE*, Benth. & Hook. f. ii. 105.
verbenacea, *Hook. f. l. c.*—Cuba.

CERATOSANTHES, Burm. ex Adans. Fam. ii. 139 (1763) ; Juss. Gen. 396 (1789). *CUCURBITACEAE*, Benth. & Hook. f. i. 833.

CERATOSANTHES :—
corniculata, Cogn. in DC. Monog. Phan. iii. 724 = tuberosa.
gracilis, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) ; *Diagn. Cucurb.* fasc. ii. 33.—N. Granat.
Hilariana, *Cogn. l. c.* 31.—Bras.
latiloba, *Cogn. l. c.* 33.—Venezuela.
multiloba, *Cogn. l. c.* 29.—Bras.
tomentosa, *Cogn. l. c.*—Bras.
trifoliata, *Cogn. l. c.* 34.—Bras.
tuberosa, *J. F. Gmel. Syst.* 102.—Ind. occ.
Warmingii, *Cogn. in Mém. Cour. Acad. Belg.* 8vo. xxviii. (1878) ; *Diagn. Cucurb.* fasc. ii. 30.—Bras.

CERATOSTANTHUS, Schur, in Verh. Sieb. Ver. Naturf. (1853) 46 ; et Enum. Pl. Transs. 30 (1866) = **Delphinium**, Tourn. (Ranunculac.).
Ajacis, Schur, Enum. Pl. Transs. 30 = D. Ajacis.
Consolida, Schur, in Verh. Sieb. Ver. Naturf. (1853) 46 = D. Consolida.
divaricatus, Schur, l. c. = D. divaricatum.
paniculatus, Schur, l. c. = D. paniculatum.

CERATOSCHOENUS, Nees, in Linnaea, ix. (1834) 296 = **Rynchospora**, Vahl (Cyperac.).
capitatus, Chapman, Fl. S. U. St. 529 = R. cyperoides.
corniculatus, Nees, in Linnaea, ix. (1834) 296 = R. laxa.
longirostris, Torr. in Ann. Lyc. N. York, iii. (1836) 369 = R. laxa.
macrophyllus, Tuckerm. in Am. Journ. Sc. Ser. II. vi. (1848) 232 = R. laxa.
macrostachys, Torr. in Ann. Lyc. N. York, iii. (1836) 369 = R. macrostachya.

CERATOSEPALUM, Oerst. Rech. Fl. Am. Centr. 18. t. 17 (1863) = **Passiflora**, Linn.
micranthum, Oerst. l. c. t. 17 = P. ceratosepala.
parviflorum, Mast. in Mart. Fl. Bras. xiii. 1. 555 = P. ceratosepala.

CERATOSPERMUM, Pers. Syn. ii. 551 (1807) = **Eurotia**, Adans. (Chenop.).
papposum, Pers. l. c. 552 = E. ceratoides.

CERATOSTACHYS, Blume, Bijdr. 644 (1825) = **Nyssa**, Linn. (Cornac.).
arborea, Blume, l. c. = N. sessiliflora.

CERATOSTEMA, Juss. Gen. Pl. 163 (1789). *VACCINIACEAE*, Benth. & Hook. f. ii. 570.
SIPHONANDRA, Klotzsch, in Linnaea, xxiv. (1851) 24.
SIPHONOSTEMA, Griseb. in Lechl. Berb. Am. Austr. 58 (1857).
angulatum, Griff. Notul. iv. 302 = Agapetes angulata.
biflorum, Poepp. & Endl. Nov. Gen. et Sp. i. 6. t. 10 = Eurygania biflora.
buxifolium, *Gardn. & Field, Sert. Pl.* t. 7.—Bras.
calycinum, *Benth. & Hook. f. Gen.* ii. 570.—Ecuador.
cordifolium, *Dun. in DC. Prod.* vii. 553.—Peruv.
costatum, *Benth. & Hook. f. Gen.* ii. 570.—Peruv.
ellipticum, *Benth. & Hook. f. l. c.*—Am. austr.
emarginatum, *Ruiz & Pav. Fl. Per.* iv. t. 384.—Peruv.
grandiflorum, *Ruiz & Pav. l. c.* 383.—Peruv.
hirsutum, *Ruiz & Pav. l. c.*—Peruv.
lanceolatum, *Benth. Pl. Hartw.* 142.—Peruv.
longiflorum, *Benth. & Hook. f. Gen.* ii. 570.—Ecuador.
longiflorum, Lindl. ex Lem. in Fl. des Serres, Sér. I. iv. (1848) 346[b] = grandiflorum.
loranthiflorum, *Benth. Pl. Hartw.* 142.—Peruv.
miniatum, Griff. Notul. iv. 302 = Agapetes miniata.
myrtifolium, *Griseb. ex Benth. & Hook. f. Gen.* ii. 570, nomen.—Am. austr.
nanum, Griff. Notul. iv. 303 = Agapetes nana.
oblongifolium, *Dun. in DC. Prod.* vii. 553.—Peruv.
odoratissimum, Willd. ex Klotzsch, in Linnaea, xxiv. (1851) 31 = Cavendishia Qnerema.
parvifolium, Benth. Pl. Hartw. 220 = Enrygania parvifolia.
peruvianum, *Pers. Syn.* i. 480.—Peruv.
pubiflorum, *Wedd. Chlor. And.* ii. 181.—N. Granat.
rigidum, *Benth. Pl. Hartw.* 220.—N. Granat.
Salapa, *Benth. l. c.* 141.—Peruv.
speciosum, *André, Illustr. Hortic.* xvii. (1870) t. 9.—Ecuador.

CERATOSTEMA :—
vaccinaceum, Roxb. Hort. Beng. 33 ; Fl. Ind. ii. 412 = Vaccinium serratum.
variegatum, D. Don, ex Steud. Nom. ed. II. i. 332 = Agapetes variegata.
variegatum, Griff. Notul. iv. 302 = Agapetes grandiflora.
variegatum, Roxb. Hort. Beng. 33 ; Fl. Ind. ii. 413 = Agapetes macrantha.

CERATOSTIGMA, Bunge, in Mém. Sav. Etr. Acad. Pétersb. ii. (1835) 129 (Enum. Pl. Chin. Bor. 55). *PLUMBAGINEAE*, Benth. & Hook. f. ii. 628.
VALORADIA, Hochst. in Flora, xxv. (1842) 239.
abyssinicum, *Aschers. in Schweinf. Beitr. Fl. Aethiop.* 288.—Afr. trop.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 481.—Reg. Himal.
plumbaginioides, *Bunge, Enum. Pl. Chin. Bor.* 55.—China.

CERATOSTYLIS, Blume, Bijdr. 304. t. 56 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 517.
ACORIDIUM, Nees & Meyen, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 131.
anceps, *Blume, Bijdr.* 305.—Java.
braccata, *Reichb. in Bonplandia*, v. (1857) 53.—Java.
capitata, *Zoll. & Moritzi, in Nat. en Geneesk. Arch. Neêrl. Indie*, i. (1844) 404.—Java.
Cepula, *Reichb. f. in Bonplandia*, v. (1857) 53.—Java.
Gigas, *Reichb. f. l. c.*—Java.
gracilis, *Blume, Bijdr.* 306.—Java.
graminea, *Blume, l. c.* 305.—Java.
latifolia, *Blume, l. c.*—Java.
retisquama, *Reichb. f. in Bonplandia*, v. (1857) 53.—Ins. Philipp.
senilis, *Reichb. in Otia Bot. Hamb.* 54.—Ins. Philipp.
simplex, *Blume, Bijdr.* 305.—Java.
subulata, *Blume, l. c.* 306.—Java.
teres, *Reichb. f. in Bonplandia*, ii. (1854) 89.—Reg. Himal.

CERATOTHECA, Endl. in Linnaea, vii. (1832) 5. t. 1, 2. *PEDALINEAE*, Benth. & Hook. f. ii. 1059.
SPORLEDERA, Bernh. in Linnaea, xvi. (1842) 41.
Kraussiana, *Bernh. in Linnaea*, xvi. (1842) 32.—Afr. austr.
melanosperma, *Hochst. ex Bernh. l. c.*—Afr. trop.
sesamoides, *Endl. l. c.* vii. (1832) 5. t. 1, 2.—Afr. trop.
triloba, *E. Mey. ex Bernh. l. c.* xvi. (1842) 29.—Afr. austr.

CERATOZAMIA, Brongn. in Ann. Sc. Nat. Sér. III. v. (1846) 7. t. 1, partim. *CYCADACEAE*, Benth. & Hook. f. iii. 446.
DIPSACOZAMIA, Lehm. in Lindl. Veg. Kingd. 225 (1847).
angustifolia, *Linden, in Illustr. Hortic.* xxviii. (1881) 32, nomen.—Mexic.
boliviana, *Brongn. in Ann. Sc. Nat.* Sér. III. v. (1846) 9.—Bolivia.
brevifrons, Miq. in Tijdschr. Wis- en Nat. Wetens. i. (1848) 41 = mexicana.
fusco-viridis, D. Moore, in Proc. Roy. Dubl. Soc. (1878) 113 = mexicana.
Ghiesbrechti, Hort. ex Regel, in Act. Hort. Petrop. iv. (1876) 299 = Miqueliana.
intermedia, Miq. in Tijdschr. Wis- en Nat. Wetens. i. (1848) 40 = mexicana.
Karsteniana, Hort. Kew. ex Dyer, in Hemsl. Biol. Centr. Am. Bot. iii. 192 = latifolia.
Katzeriana, *Regel, Gartenfl.* (1876) 261.—Hab. ?
Kuesteriana, *Regel, in Bull. Soc. Nat. Mosc.* xxx. (1857) 1. 187.—Mexic.
latifolia, *Hort. Belg. ex Miq. in Tijdschr. Wis- en Nat. Wetens.* i. (1848) 206.—Mexic.
longifolia, Miq. l. c. 40 = mexicana.
mexicana, *Brongn. in Ann. Sc. Nat.* Sér. III. v. (1846) 8.—Mexic.
Miqneliana, *H. Wendl. Ind. Palm.* 68.—Mexic.
muricata, *Miq. in Linden, Illustr. Hortic.* xxviii. (1881) 32, nomen.—Mexic.
robusta, Miq. in Tijdschr. Wis- en Nat. Wetens. i. (1848) 42 = mexicana.

CERAUNIA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 (1827) (Quid?).

CERBERA, Linn. Gen. ed. I. 62 (1737). *APOCYNACEAE*, Benth. & Hook. f. ii. 699.
CASCABELA, Rafin. Sylva Tellur. 162 (1838).
ELCANA, Blanco, Fl. Filip. ed. II. 584 (1845).
MANGHAS, Burm. Thes. Zeyl. 150 (1737).
NEIOSPERMA, Rafin. Sylva Tellur. 162 (1838).
ODOLLAM, Adans. Fam. ii. 171 (1763).
ODOLLAMIA, Rafin. Sylva Tellur. 162 (1838).
TANGHINIA, Thou. Gen. Nov. Madag. 10 (1806).
Ahouai, Linn. Sp. Pl. 208 = Thevetia Ahouai.
alliodora, Willd. ex Roem. & Schult. Syst. iv. 798 = Thevetia cuneifolia.
batjanica, *Teijsm. & Binn. Cat. Hort. Bog.* 385, nomen. —Malaya.
borbonica, Spreng. Syst. i. 642 = Ochrosia borbonica.
chinensis, Spreng. l. c. = Dissolaena verticillata.
cuneifolia, H. B. & K. Nov. Gen. et Sp. iii. 224 = Thevetia cuneifolia.
dichotoma, Lodd. Bot. Cab. t. 1516 = Tabernaemontana dichotoma.
Forsteri, Seem. Fl. Vit. 157 = C. Odollam.
fruticosa, Ker-Gawl. in Bot. Reg. t. 391 = Kopsia fruticosa.
fruticosa, *Roxb. Hort. Beng.* 19; *Fl. Ind.* i. 691.—Burma.
glomerata, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 158 = Pseudochrosia glomerata.
lactaria, Buch.-Ham. ex D. Dietr. Syn. Pl. i. 623 = C. Odollam.
laurifolia, Lodd. Bot. Cab. t. 989 = C. Odollam.
linearifolia, Stokes, Bot. Mat. Med. i. 490 = Thevetia nereifolia.
Linnaei, *Montr. in Mém. Acad. Lyon*, x. (1860) 233. —N. Caled.
maculata, Willd. Sp. Pl. i. 1223 = Ochrosia borbonica.
Manghas, Linn. Sp. Pl. 208 = C. Odollam, et Tabernaemontana dichotoma.
museuliformis, Lam. Encyc. i. 62 = Banksia musculiformis.
nereifolia, Zipp. in Bijdr. Nat. Wetensch. v. (1830) 178 = Podocarpus Rumphia.
nitida, H. B. & K. Nov. Gen. et Sp. iii. 225 = Thevetia nitida.
obovata, *Roem. & Schult. Syst.* iv. 798.—Hab.?
obtusifolia, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 203.—N. Caled.
Odollam, *Gaertn. Fruct.* ii. 193.—Ind. or.; Malaya.
oppositifolia, *Lam. Encyc.* i. 62.—Ins. Molucc.
ovata, Cav. Ic. iii. 35. t. 270 = Thevetia ovata.
parviflora, Forst. f. Prod. 19 = Ochrosia elliptica.
parviflora, Hook. & Arn. Bot. Beech. Voy. 90 = Ochrosia sandwicensis.
parviflora, Wall. Cat. n. 1584 = Ochrosia borbonica.
peruviana, Pers. Syn. i. 267 = Thevetia nereifolia.
platyspermos, Gaertn. Fruct. ii. 193. t. 124. f. 2 = Ochrosia borbonica.
quaternifolia, *Roxb. Hort. Beng.* 19, nomen.—Sumatra.
salutaris, Blume, Bijdr. 1033 = oppositifolia.
Tanghin, *Hook. Bot. Mag.* t. 2968.—Madag.
Tanquin, Steud. Nom. ed. II. i. 332 = praec.
Thevetia, Linn. Sp. Pl. 209 = Thevetia nereifolia.
thevetioides, H. B. & K. Nov. Gen. et Sp. iii. 223 = Thevetia Yccotli.
triphylla, Rudge, Pl. Gui. 31. t. 48 = Couma guianensis.
tubiflora, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 156 = Neuburgia tubiflora.
undulata, Andr. Bot. Rep. t. 130 = Ochrosia borbonica.
venenifera, Steud. Nom. ed. II. i. 332 = C. Tanghin.

CERBERA, Lour. Fl. Cochin. 136 (1790) = **Scaevola**, Linn. (Goodeniac.).
ovata, Sieber, ex DC. Prod. vii. 507 = S. Lobelia.
salutaris, Lour. Fl. Cochinch. 136 = S. Koenigii?

CERBERIOPSIS, Vieill. ex Panch. & Sebert, Not. Bois Nouv. Caléd. 187 (1874). *APOCYNACEAE*.
candelabra, *Vieill. l. c.*—N. Caled.

CERCESTIS, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 414. *AROIDEAE*, Benth. & Hook. f. iii. 980.
Afzelii, *Schott, l. c.*—Afr. trop.

CERCIDIPHYLLUM, Sieb. & Zucc. in Abh. Math. Acad. Muench. iv. III. (1846) 238. *MAGNOLIACEAE*?
japonicum, *Sieb. & Zucc. l. c.*—Japon.

CERCIDIUM, Tul. in Arch. Mus. Par. iv. (1844) 133. *LEGUMINOSAE*, Benth. & Hook. f. i. 570.
HOOPESIA, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 453, partim.
RHETINOPHLOEUM, Karst. Fl. Columb. ii. 25. t. 113 (1862?).
andicolum, *Griseb. in Goett. Abh.* xxiv. (1879) 114.—Reg. Argent.
floridum, *Benth. ex A. Gray, Pl. Wright.* i. 58.—Mexic.
floridum, Torr. in Pacif. Rail. Rep. v. 360. t. 3 = Parkinsonia Torreyana.
spinosum, *Tul. in Arch. Mus. Par.* iv. (1844) 134.—Bras.
texanum, *A. Gray, Pl. Wright.* i. 58.—Mexic.

CERCIS, Linn. Sp. Pl. 374, ed. I (1753). *LEGUMINOSAE*, Benth. & Hook. f. i. 576.
SILIQUASTRUM, Tourn. ex Adans. Fam. ii. 317 (1763).
ealifornica, Torr. ex Benth. Pl. Hartw. 361, nomen = occidentalis.
canadensis, *Linn. Sp. Pl.* 374.—Am. bor.
chinensis, *Bunge, in Mém. Sav. Etr. Pétersb.* ii. (1835) 95.—China.
florida, Salisb. Prod. 327 = C. Siliquastrum.
Griffithii, *Boiss. Fl. Orient.* ii. 633.—Afghan.
japonica, Siebold, ex Planch. in Fl. des Serres, Sér. I. viii. (1852–53) 269 = chinensis.
occidentalis, *Torr. ex A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 177.—Texas.
reniformis, Engelm. ex A. Gray, l. c. = occidentalis.
siliquosa, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Siliquastrum.
Siliquastrum, *Linn. Sp. Pl.* 374.—Europ.; Oriens.

CERCOCARPUS, H. B. & K. Nov. Gen. et Sp. vi. 232. t. 559 (1823). *ROSACEAE*, Benth. & Hook. f. i. 618.
betulaefolius, Nutt. ex Hook. Ic. Pl. t. 322 = parvifolius.
betuloides, Nutt. in Torr. & Gray, Fl. N. Am. i. 427 = parvifolius.
breviflorus, *A. Gray, Pl. Wright.* ii. 54.—N. Mexic.
fothergilloides, *H. B. & K. Nov. Gen. et Sp.* vi. 233.—Mexic.; Am. bor.
fothergilloides, Torr. in Ann. Lyc. N. York, ii. (1828) 198 = parvifolius.
intricatus, *S. Wats. in Proc. Am. Acad.* x. (1875) 346. —Am. bor. occ.
ledifolius, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 427. —Am. bor. occ.
montanus, *Rafin. Atl. Journ.* 146.—Am. bor.
parvifolius, *Nutt. ex Hook. & Arn. Bot. Beech. Voy.* 337.—Calif.

CERCOCOMA, Wall. Cat. n. 1623 (1832) = **Strophanthus**, DC. (Apocynac.).
macrantha, Teijsm. & Binn. Cat. Hort. Bog. 126.—Hab.?
singaporiana, Wall. Cat. n. 1623 = S. brevicaudatus.

CERCOCOMA, Miq. Fl. Ind. Bat. ii. 445 (1856) = **Rhynchodia**, Benth. (Apocynac.).
Wallichii, Miq. l. c. = R. Wallichii.

CERCODEA, Soland. ex Lam. Encyc. i. 682 (1783) = seq.

CERCODIA, Murr. in Comm. Gotting. iii. (1780) 3. t. 1 = **Haloragis**, Forst.
alternifolia, A. Cunn. in Ann. Nat. Hist. iii. (1839) 29 = H. alata.
erecta, Murr. in Comm. Gotting. iii. (1780) 3. t. 1 = H. alata.
incana, A. Cunn. in Ann. Nat. Hist. iii. (1839) 30 = H. tetragyna.
racemosa, DC. Prod. iii. 67 = H. racemosa.

CERCOPHORA, Miers, in Trans. Linn. Soc. xxx. (1874) 301. *MYRTACEAE*.
anomala, *Miers, l. c.* 302.—Reg. Amazon.

CERCOSTYLOS, Less. Syn. Comp. 239 (1832) = **Gaillardia**, Fouger. (Compos.).
brasiliensis, Less. l. c. 240 = G. megapotamica.
scabiosoides, Arn. ex DC. Prod. vii. 293 = G. megapotamica.
sesamoides, Steud. Nom. ed. II. i. 333, sphalm. = prace.

CERDANA, Ruiz & Pav. Prod. 37. t. 6 (1794) = **Cordia**, Linn. (Boragin.).
alliodora, Ruiz & Pav. Fl. Per. ii. 47 = Cordia Cerdana.
eujabensis, Manso, ex DC. Prod. ix. 473 = Cordia cujabensis.

CERDIA, Moç. & Sesse, ex DC. Prod. iii. 377 (1828). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 153.
congestiflora, *Hemsl. Diagn. Pl. Nov.* 23.—Mexic.
glauca, *Hemsl. l. c.* 22.—Mexic.
purpurascens, *Moç. & Sesse, ex DC. Prod.* iii. 377.—Mexic.
virescens, *Moç. & Sesse, l. c.*—Mexic.

CEREFOLIUM, Riv. ex Rupp. Fl. Jen. ed. Hall. 283 (1745) = **Anthriscus**, Hoffm. (Umbellif.).
nitidum, Čelak. in Arch. Naturw. Landesd. Boehm. iii. (1875) 586 = A. nitida.
sativum, Bess. Prim. Fl. Galic. i. 219 = A. Cerefolium.
sylvestre, Bess. l. c. 218 = A. Cerefolium.
trichospermum, Bess. Enum. Pl. Volh. 44 = A. Cerefolium.

CEREOPSIS, Blanco, Fl. Filip. ed. II. Suppl. 591 (1845), sphalm. = **Coreopsis**, Linn. (Compos.).

CERESIA, Pers. Syn. i. 85 (1805) = **Paspalum**, Linn. (Gramin.).
aristata, Willd. ex Steud. Nom. ed. II. i. 333 = Panicum Thrasya.
elegans, Pers. Syn. i. 85 = Pasp. membranaceum.
fluitans, Ell. Sketch, i. 109 = Pasp. repens.
gracilis, Schlecht. in Linnaea, xxvi. (1853–55) 134 = Pasp. gracile.
membranacea, Beauv. Agrost. 9. t. 5. f. 4 = Pasp. membranaceum.
Wageneriana, Schlecht. in Linnaea, xxvi. (1853–55) 133 = Pasp. Wagenerianum.

CEREUS, Mill. Gard. Dict. ed. VIII (1768); Haw. Syn. Pl. Succ. 178 (1812). *CACTEAE*, Benth. & Hook. f. i. 849.
CEPHALOCEREUS, Pfeiff. in Otto & Dietr. Allg. Gartenz. vi. (1838) 142.
CEPHALOPHORUS, Lem. Cact. Aliq. Nov. Desc. p. xii. (1838).
CLEISTOCACTUS, Lem. Illust. Hortic. viii. (1861) Misc. 35.
ECHINOCEREUS, Engelm. in Wisliz. Tour North Mexico, 91 (1848).
ECHINONYCTANTHUS, Lem. Cact. Gen. Nov. et Sp. 10 (1839).
ECHINOPSIS, Zucc. in Abh. Akad. Muench, ii. (1837) 675.
PILOCEREUS, Lem. Cact. Gen. Nov. et Sp. 6 (1839).
abnormis, Sweet, Hort. Brit. ed. I. 171 = peruvianus.
acifer, *Otto, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 189.—Hab.?
acinaciformis, Steud. Nom. ed. II. i. 333 = cinerascens.
aeiniformis, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 101 = cinerascens.
Ackermanni, Hort. Berol. ex Pfeiff. l. c. 123 = Phyllocactus Ackermanni.
acromelas, Hort. Berol. Index Cact. 1833, ex Pfeiff. Enum. Diagn. Cact. 84 = crenulatus.
aculeatus, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 433, nomen.—Hab.?
acutangulus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 107.—Mexic.
adustus, *Engelm. in A. Gray, Pl. Fendl.* 50.—N. Mexic.
Aethiops, Haw. in Phil. Mag. (1830) 109 = caerulescens.
affinis, Hort. ex Pfeiff. Enum. Diagn. Cact. 99, 106 = variabilis, virens.
alacriportanus, *Hort. Monac. ex Pfeiff. Enum. Diagn. Cact.* 87.—Bras.
alatus, DC. Prod. iii. 470 = Rhipsalis pachyptera.
albisetosus, *Haw. Suppl. Pl. Succ.* 77.—Ins. S. Doming.
albispinus, *Salm-Dyck, Obs. Bot.* (1822) 5.—Am. austr.
ambiguus, *DC. Prod.* iii. 467.—Hab.?
amblygonus, *G. Don, in Sweet, Hort. Brit.* ed. III. 284, nomen.—Am. austr.
amoenus, *Hemsl. Biol. Centr. Am. Bot.* i. 540.—Mexic.

CEREUS:—
anisacanthus, *DC. in Mém. Mus. Par.* xvii. (1828) 116.—Mexic.
anizogonus, Hort. Angl. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 52 = triangularis.
Antoini, Hort. Vindob. ex Pfeiff. Enum. Diagn. Cact. 114 = nycticalus.
aquicaulensis, Hort. ex Pfeiff. l. c. 90 = geometrizans.
arequipensis, Meyen, Reise, ii. 41 = peruvianus.
armatus, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 81.—Ins. S. Thom.
Arrabidae, *Steud. Nom.* ed. II. i. 333.—Bras.
articulatus, *Pfeiff. Enum. Diagn. Cact.* 103.—Chili.
assurgens, *Wright, ex Griseb. Cat. Pl. Cub.* 116.—Cuba.
atacamensis, *Phil. Fl. Atac.* 23.—Chili.
auratus, *Labour. in Rev. Hortic.* Sér. IV. iv. (1855) 27.—Mexic.
aureus, *Salm-Dyck, ex DC. Prod.* iii. 465.—Hab.?
azureus, *Parm. ex Pfeiff. Enum. Diagn. Cact.* 86.—Bras.
bacciferus, Hemsl. Biol. Centr. Am. Bot. i. 548 = Rhipsalis Cassutha.
barbatus, Wendl. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 29 = floccosus.
Baumannii, *Lem. in Hortic. Univ.* (1844) 126, 315; *ex Lem. Jard. Fleur.* i. (1851) t. 48.—Peruv.?
baxaniensis, *Karw. ex Pfeiff. Enum. Diagn. Cact.* 109.—Mexic.
Beneckei, *Ehrenb. in Bot. Zeit.* ii. (1844) 835.—Mexic.
Berlandieri, *Engelm. Cact. Mex. Bound.* 38. t. 58.—Texas.
Bertini, *E. Cels, ex Hérincq, L'Horticult. Franç.* (1863) 251.—Chili.
biformis, Hort. ex Pfeiff. Enum. Diagn. Cact. 114 = radicans.
biformis, Lindl. Bot. Reg. (1843) Misc. 51 = Phyllocactus biformis.
bifrons, Haw. Suppl. Pl. Succ. 76 = coccineus.
Bigelovii, Engelm. in Pacif. Rail. Rep. iv. t. 4. f. 8 = mojavensis.
Blanckii, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 134.—Mexic.
Boeckmanni, *Otto, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 217.—Mexic.
bonariensis, *Sweet, Hort. Brit.* ed. III. 283.—Reg. Argent.
bonariensis, Hort. Berol. ex Foerst. Handb. Cact. ed. I. 388 = alacriportanus.
Bonplandii, *Parm. ex Pfeiff. Enum. Diagn. Cact.* 108.—Bras.
brachiatus, *Gal. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 195.—Mexic.
bradypus, Steud. Nom. ed. II. i. 333 = senilis.
Brandii, Hort. Angl. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 49 = variabilis.
brevispinulus, Salm-Dyck, Hort. Dyck. 339 = nycticalus.
Bridgesii, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 208.—Bolivia.
caerulescens, *Salm-Dyck, Hort. Dyck.* 335.—Bras.
caesius, *Salm-Dyck, ex Pfeiff. Enum. Diagn. Cact.* 89.—Hab.?
caespitosus, *Engelm. & A. Gray, in Bost. Journ. Nat. Hist.* v. (1845) 247.—Texas.
californicus, Torr. & Gray, Fl. N. Am. i. 555 = Opuntia serpentina?
callicoche, Gal. ex Scheidw. in Bull. Acad. Brux. vi. 1. (1839) 88 = Echinocactus myriostigma.
calvescens, DC. in Mém. Mus. Par. xvii. (1828) 116 = peruvianus.
candelabrius, Hort. ex Pfeiff. Enum. Diagn. Cact. 109 = tenellus.
candelaris, Meyen, Reise, i. 447 = peruvianus.
candicans, *Gill. ex Salm-Dyck, Hort. Dyck.* 335.—Chili.
caripensis, *DC. Prod.* iii. 467.—Mexic.
caudatus, Gill. ex Sweet, Hort. Brit. ed. III. 285 = Mammillaria caudata.
Cavendishii, *Monv. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 210.—Hab.?
chalibaeus, *Hort. Berol. ex Salm-Dyck, l. c.* 201.—Hab.?
chalybaeus, Otto, ex Walp. Rep. ii. 340 = polychaetus.
chiloënsis, *DC. Prod.* iii. 465.—Ins. Chiloe.
chloranthus, *Engelm. Cact. Mex. Bound.* 29.—Texas.

CEREUS:—
chlorocarpus, *DC. Prod.* iii. 466.—Ecuador.
chrysomallus, *Hemsl. Biol. Centr. Am. Bot.* i. 541.—Mexic.
cinerascens, *DC. in Mém. Mus. Par.* xvii. (1828) 11.—Mexic.
cirrhiferus, *Labour. Cact.* 311.—Mexic.
clavarioides, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 173 = Opuntia clavarioides.
clavatus, *Otto & Dietr. Allg. Gartenz.* vi. (1838) 28.—Venezuela.
claviformis, *Regel & Klein, Ind. Sem. Hort. Petrop.* (1860) 46.—Mexic.
coccineus, DC. Prod. iii. 469 = setaceus.
coccineus, Engelm. in A. Gray, Pl. Fendl. 51 = phoeniceus.
coccineus, *Salm-Dyck, ex Pfeiff. & Otto, Cact.* i. t. 15.—Bras.
cognatus, Hort. ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
colicochus, Booth, Cat. ex Foerst. Handb. Cact. ed. I. 336 = Echinocactus myriostigma.
colubrinus, *Otto, ex Foerst. l. c.* 409; *ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 50, 215.—Hab.?
Columna, Walp. Rep. v. 818 = seq.
Columna-Trajani, *Karw. ex Pfeiff. Enum. Diagn. Cact.* 76.—Mexic.
columnaris, *Lodd. ex Voigt, Hort. Suburb. Calc.* 61, nomen.—Hab.?
Colvillii ×, *Sweet, Hort. Brit.* ed. II. 236, nomen, et p. 590.
cometes, *Scheidw. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 339.—Mexic.
compressus, *Mill. Gard. Dict.* ed. VIII. n. 10.—Mexic.
concolor, Bigel. in Whipple, Exped. Add. p. ii = caespitosus.
conformis, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 203.—Mexic.
conicus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 97.—Mexic.
conoideus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 36.—Am. bor. occ.
coquimbanus, Hort. ex DC. Prod. iii. 465 = chiloensis.
Coryne, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 205.—Hab.?
cossyrensis, *Tin. ex Todaro, in Ind. Sem. Hort. Panorm.* (1857) 39.—Hab.?
crenatus, Lindl. Bot. Reg. (1844) t. 31 = Phyllocactus crenatus.
crenoides, *Engelm. Cact. Mex. Bound.* 31. t. 42.—Mexic.
crenulatus, *Salm-Dyck, Obs. Bot.* (1822) 6.—Am. austr.
Crimsonii, Pritz. Ic. Ind. 246, sphalm. = Smithii.
crispatus, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 130 = Rhipsalis crispata.
cruciformis, *Steud. Nom.* ed. II. i. 333.—Bras.
cubensis, Zuccar. ex Pfeiff. Enum. Diagn. Cact. 94 = eriophorus.
cupulatus, Hort. ex Pfeiff. l. c. 97 = marginatus.
Curtisi, *Otto, in Otto & Dietr. Allg. Gartenz.* i. (1833) 365.—N. Granat.
curvispinus, Pfeiff. Enum. Diagn. Cact. 89 = peruvianus.
cylindricus, Haw. Syn. Pl. Succ. 183 = Opuntia cylindrica.
dasyacanthus, *Engelm. in A. Gray, Pl. Fendl.* 50.—Texas.
decagonus, Hort. ex Pfeiff. Enum. Diagn. Cact. 85 = albispinus.
Decandollii, *Pfeiff. ex Foerst. Handb. Cact.* ed. I. 389.—Hab.?
decorus, *Lodd. ex Voigt, Hort. Suburb. Calc.* 62, nomen.—Hab.?
deficiens, *Otto & Dietr. Allg. Gartenz.* vi. (1838) 28.—Venezuela.
deflexispinus, Monv. Cat. (1846), ex Labour. Cact. 321 = dasyacanthus.
denudatus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 73.—Bras.
Deppei, Hort. Berol. ex Pfeiff. l. c. 101 = cinerascens.
dichroacanthus, *Mart. ex Pfeiff. l. c.* 76.—Mexic.
divaricatus, *DC. Prod.* iii. 466.—Ins. S. Doming.
divergens, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 95.—Ins. S. Doming.
Donkelaarii, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 355.—Bras.
dubius, *Engelm. Cact. Mex. Bound.* 36.—Texas.

CEREUS:—
Duledevanti, *Lem. Illustr. Hortic.* v. (1858) *Misc.* 10.—Hab.?
Dumortieri, *Scheidw. in Bull. Acad. Brux.* vi. 1. (1839) 88.—Reg. Argent.
Dyckii, *Mart. ex Pfeiff. Enum. Diagn. Cact.* 87.—Mexic.
eburneus, *Salm-Dyck, Obs. Bot.* (1822) 6.—Ins. Curaçao; Chili.
Ehrenbergii, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 282.—Mexic.
elegans, Hort. ex Pfeiff. Enum. Diagn. Cact. 138 = Rhipsalis Mittleri.
elegans, *Lem. Illustr. Hortic.* v. (1858) *Misc.* 10.—Peruv.
Emoryi, *Engelm. in Am. Journ. Sc.* Ser. II. xiv. (1852) 338.—Calif.
Engelmanni, *Parry, ex Engelm. l. c.*—Calif.
enneacanthus, *Engelm. in A. Gray, Pl. Fendl.* 50.—Texas.
erectus, *Karw. ex Pfeiff. Enum. Diagn. Cact.* 95.—Mexic.
ericomus, Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 47 = virens.
ericomus, *Reichb. f. in Terscheck. Suppl. Cact. Verz.* 3.—Bras.
erinaceus, *Haw. ex Steud. Nom.* ed. II. i. 334.—Ind. occ.
eriophorus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 94.—Cuba.
euphorbioides, *Haw. Suppl. Pl. Succ.* 75.—Am. austr.
exerens, Link, ex Pfeiff. Enum. Diagn. Cact. 99 = virens.
extensus, *Salm-Dyck, ex DC. Prod.* iii. 469.—Am. austr.
Eyriesii, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 72.—Reg. Argent.
farinosus, Haage, ex Salm-Dyck, in Otto & Dietr. Allg. Gartenz. xiii. (1845) 355 = Beneckei.
Fendleri, *Engelm. in A. Gray, Pl. Fendl.* 51.—N. Mexic.
Ferchecki, *Parm. ex Pfeiff. in Hort. Belg.* v. (1838) 66.—Reg. Argent.
fernambucensis, Lem. sphalm. vide pernambucensis.
ferox, *Haw. in Phil. Mag.* (1830) 109.—Bras.
fimbriatus, DC. Prod. iii. 464 = grandispinus.
fimbriatus, Hort. ex Pfeiff. Enum. Diagn. Cact. 95 = divaricatus.
flagelliformis, *Mill. Gard. Dict.* ed. VIII. n. 12.—Am. austr.
flagriformis, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 111.—Mexic.
flavescens, *Otto, ex Pfeiff. l. c.* 79.—Hab.?
flavicomus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 202.—Hab.?
flavispinus, *Haw. ex Steud. Nom.* ed. II. i. 334.—Am. austr.
flavispinus, *Salm-Dyck, Obs. Bot.* (1822) 5.—Am. calid.
floccosus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 81.—Ins. S. Thomas.
fluminensis, *Miq. in Bull. Sc. Phys. Nat. Néerl.* (1838) 48.—Bras.
Forbesii, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 398; *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 206.—Hab.?
formosus, *Hort. ex Foerst. Handb. Cact.* ed. I. 404.—Reg. Argent.
formosus, Monv. Cat. (1834) = Schrankii.
foveolatus, *Haage, ex Pfeiff. Enum. Diagn. Cact.* 77.—Hab.?
fulgidus, *Hook. f. Bot. Mag.* t. 5856.—Hab.?
fulvibarbis, Otto & Dietr. Allg. Gartenz. vi. (1838) 28 = chiloënsis.
fulvispinosus, Haw. Syn. Pl. Succ. 183 = Royeni.
fulvispinus, Salm-Dyck, Cact. Hort. Dyck. ed. II. 46, sphalm. = praec.
Funkii ×, *Hort. Monac. ex Foerst. Handb. Cact.* ed. I. 411.
Garambello, Haage, ex Foerst. l. c. 433 = geometrizans.
geminisetus, *Reichb. in Terscheck. Suppl. Cact. Verz.* 3.—Hab.?
gemmatus, Otto, ex Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 314 = turbinatus.
gemmatus, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 96.—Mexic.
geometrizans, *Mart. ex Pfeiff. l. c.* 90.—Mexic.

CEREUS:—
gibbosus, *Sweet, Hort. Brit.* ed. I. 171.—Jamaica.
giganteus, *Engelm. in Emory, Notes Mil. Reconnois.* 158.—Texas.
gilvus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 355.—Hab.?
Gladiator, Otto & Dietr. l. c. vi. (1838) 34 = geometrizans.
gladiatus, Lem. Cact. Aliq. Nov. Desc. 28 = candicans.
Gladiger, Cels, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 47 = resupinatus.
glaucescens, *Tweedie, ex Sweet, Hort. Brit.* ed. III. 284, nomen.—Reg. Argent.
glaucus, Hort. ex Pfeiff. Enum. Diagn. Cact. 89 = caesius, variabilis.
glaucus, *Salm-Dyck, Hort. Dyck.* 335.—Bras.
gloriosus, Hort. ex Pfeiff. Enum. Diagn. Cact. 80 = Royeni.
gonacanthus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 33. t. 5. f. 2. 3.—N. Mexic.
gracilis, *Haw. in Phil. Mag.* (1827) 125.—Hab.?
gracilis, *Mill. Gard. Dict.* ed. VIII. n. 8.—Ind. occ.
gracilis, Salm-Dyck, ex DC. Prod. iii. 468 = humilis.
grandiflorus, *Mill. Gard. Dict.* ed. VIII. n. 11.—Ind. occ.
grandis, *Haw. Suppl. Pl. Succ.* 76.—Bras.
grandispinus, *Haw. in Phil. Mag.* (1830) 109.—Ind. occ.
Greggii, *Engelm. in Wisliz. Rep. N. Mexico*, 102.—Texas.
griseus, Haw. Syn. Pl. Succ. 182 = cburneus.
Haageanus, *Salm-Dyck, ex Foerst. Handb. Cact.* ed. I. 381, nomen.—Hab.?
hamatus, [Scheidw.] Hort. Belg. ex Pfeiff. in Otto & Dietr. Allg. Gartenz. v. (1837) 371 = rostratus.
Hansii ×, *Baumann, in Rev. Hortic.* Sér. II. i. (1842) 267.
Haworthii, *DC. Prod.* iii. 465.—Ind. occ.
heptagonus, *Mill. Gard. Dict.* ed. VIII. n. 6; *Haw. Syn. Pl. Succ.* 178.—Am. austr.
hermentianus, *Monv. ex Lem. in Illustr. Hortic.* vi. (1859) *Misc.* 90.—Hab.?
heteracanthus, *Tweedie, ex Sweet, Hort. Brit.* ed. III. 284.—Reg. Argent.
hexaedrus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 34. t. 5. f. 1.—N. Mexic.
hexagonus, *Mill. Gard. Dict.* ed. VIII. n. 1; *Haw. Syn. Pl. Succ.* 179.—Am. austr.
hexangularis, Hort, ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
Histrix, Salm-Dyck, ex DC. Prod. iii. 464.—Cf. Hystrix.
Hoffmannseggii, Hort. ex Foerst. Handb. Cact. ed I. 292 = Echinocactus exsculptus.
Hookeri, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 125.—Bras.
horizontalis, *Gill. ex Sweet, Hort. Brit.* ed. III. 285.—Chili.
horribarbis, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 205.—Hab.?
horridus, *Otto, ex Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 370.—Venezuela.
Humboldtii, *DC. Prod.* iii. 467.—Ecuador.
humilis, *DC. l. c.* 468.—Am. austr.
hybridus, Otto, in Otto & Dietr. Allg. Gartenz. i. (1833) 72 = speciosissimus.
hypogaeus, *Weber, ex Regel, Gartenfl.* (1882) 165. t. 1085.—Hab.?
Hystrix, *Sweet, Hort. Brit.* ed. I. 171.—Ind. occ.
icosigonus, *DC. Prod.* iii. 467.—Ecuador.
imbricatus, Haw. Rev. Pl. Succ. 70 = Opuntia imbricata.
imbricatus, Hort. ex Pfeiff. Enum. Diagn. Cact. 93 = = subrepandus.
incrustans, Steud. Nom. ed. II. i. 334 = seq.
incrustatus, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 97 = marginatus.
incurvispinus, Hort. Darmst. ex Otto & Dietr. Allg. Gartenz. iii. (1835) 244 = leucanthus.
inermis, *Otto, ex Pfeiff. Enum. Diagn. Cact.* 116.—Venezuela.
inermis, Scheidw. in Bull. Acad. Brux. vi. (1839) 88 = Echinocactus myriostigma.
insularis, *Hemsl. Bot. Voy. Challenger*, i. 11. 16.—Mons S. Mich.

CEREUS:—
intricatus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 194.—Hab.?
inversus, *Hort. Berol. ex Salm-Dyck, l. c.* 219, nomen.—Hab.?
irradians, *Lem. Illust. Hortic.* xi. (1864) *Misc.* 74.—Cuba.
Jamacaru, Hort. Vindob. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 31 = variabilis.
Jamacaru, *DC. Prod.* iii. 467.—Bras.
janthothele, Monv. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 41 = Rhipsalis cereiformis.
jasmineus, Hort. Darmst. ex Pfeiff. Enum. Diagn. Cact. 72 = turbinatus.
Jenkinsoni ×, Sweet, Hort. Brit. ed. II. 237, nomen.
jubatus, Salm-Dyck, ex Foerst. Handb. Cact. ed. I. 357 = cometes.
Kageneckii, Gmel. ex Pfeiff. Enum. Diagn. Cact. 77 = multangularis.
Karstenii, Hort. Angl. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 207 = lepidotus.
Karstenii, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 218.—N. Granat.
Knightii, Parment. ex Pfeiff. Enum. Diagn. Cact. 139 = Rhipsalis Knightii.
Kunthianus, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 217.—Hab.?
laetevirens, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 99 = lividus.
laetevirens, Salm-Dyck, Hort. Dyck. 336 = variabilis.
lactus, *DC. Prod.* iii. 466.—Ecuador.
laetus, Salm-Dyck, Hort. Dyck. 336 = formosus.
laevigatus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 204.—Hab.?
lagenaeformis, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 164.—Peruv.
lamprochlorus, *Lem. Cact. Aliq. Nov. Desc.* 30.—Hab.?
lanatus, *DC. Prod.* iii. 464.—Ecuador.
Lanceanus, *Hort. ex G. Don, in Sweet, Hort. Brit.* ed. III. 285, nomen.—Hab.?
Landbecki, *Phil. ex Regel, Gartenfl.* (1875) 162. t. 832.—Reg. Argent.
lanuginosus, Haw. Syn. Pl. Succ. 182 = repandus.
lanuginosus, *Mill. Gard. Dict.* ed. VIII. n. 3.—Am. trop.
lateribarbatus, Hort. ex Lem. in Rev. Hortic. (1862) 427 = C. Columna-Trajani.
latifrons, Zucc. in Denkschr. Akad. Mucneh. ii. (1837) 735 = Phyllocactus latifrons.
Lecchii, *Pfeiff. Enum. Diagn. Cact.* 78.—Hab.?
Leeanus, *Hook. Bot. Mag.* t. 4417.—Mexic.
Lehmanni, Hort. ex Foerst. Handb. Cact. ed. I. 245 = Mammillaria Lehmanni.
leiocarpus, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 276.—Porto Rico.
Lemairii, *Hook. Bot. Mag.* t. 4814.—Hab.?
lepidotus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 207.—Venezuela.
leptacanthus, DC. ex Pfeiff. Enum. Diagn. Cact. 101 = pentalophus.
leptophis, *DC. in Mém. Mus. Par.* xvii. (1828) 117.—Mexic.
leucanthus, *Pfeiff. Enum. Diagn. Cact.* 71.—Chili.
limensis, Lodd. ex Foerst. Handb. Cact. ed. I. 375 = flavescens.
limensis, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 353.—Bolivia.
Linkii, Pfeiff. Enum. Diagn. Cact. 48 = Echinocactus Linkii.
Linnaei, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 165.—Mexic.
lividus, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* (1835) 380.—Am. austr.
longifolius, *Karw. ex Sweet, Hort. Brit.* ed. III. 286, nomen.—Mexic.
longipedunculatus, *Hort. Berol. ex Foerst. Handb. Cact.* ed. I. 433, nomen.—Hab.?
longisetus, *Engelm. Cact. Mex. Bound.* 32. t. 45.—Texas.
longispinus, *Salm-Dyck, ex Otto & Dietr. Allg. Gartenz.* xiii. (1845) 354.—Bolivia.
lumbricoides, *Lem. Cact. Gen. Nov. et Sp.* 60.—Bras.
lutescens, *Salm-Dyck, ex Pfeiff. Enum. Diagn. Cact.* 84.—Hab.?
MacDonaldiae, *Hook. Bot. Mag.* t. 4707.—Honduras.

CEREUS:—
macrogonus, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 203.—Hab.?
Maelenii, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 379.—Mexic.
magnus, *Haw. in Phil. Mag.* (1830) 109.—Ins. S. Doming.
Mallisoni, Hort. ex Pfeiff. Enum. Diagn. Cact. 111 = Smithii.
marginatus, *DC. in Mém. Mus. Par.* xvii. (1828) 116.—Mexic.
marginatus, Salm-Dyck, Hort. Dyck. 340 = Hookeri.
Mariculi, Hort. ex Pfeiff. Enum. Diagn. Cact. 115 = humilis.
Martianus, *Zucc. ex Pfeiff. l. c.* 110.—Mexic.
Mendory, Hort. ex Pfeiff. l. c. 85 = caeruleseens.
meonacanthus, Link & Otto, ex Steud. Nom. ed. II. i. 334 = Echinocactus meonacanthus.
mexicanus, *Lem. ex Foerst. Handb. Cact.* ed. I. 430.—Mexic.
micracanthus, *DC. in Mém. Mus. Par.* xvii. (1828) 115.—Mexic.
militaris, *Audot, in Rev. Hortic.* Sér. II. iv. (1845–46) 307.—Mexic.
Mirbelii, Hort. Belg. ex Pfeiff. Enum. Diagn. Cact. 97 = marginatus.
mohavensis, S. Wats. Bibliog. Ind. Fl. Am. Bor. 398 = scq.
mojavensis, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 33.—Am. bor. occ.
mollis, Hort. ex Pfeiff. Enum. Diagn. Cact. 83 = strictus.
mouiliformis, *DC. in Mém. Mus. Par.* xvii. (1828) 60.—Ins. S. Doming.
monoclonus, DC. Prod. iii. 464 = peruvianus.
monstrosus, Steud. Nom. ed. II. i. 334 = peruvianus.
montevidensis, Hort. ex Pfeiff. Enum. Diagn. Cact. 65 = Echinocactus exsculptus.
Montezumae, Hort. ex Pfeiff. l. c. 91 = candicans.
Moritzianus, *Otto, ex Pfeiff. l. c.* 84.—Bras.
multangularis, *Haw. Suppl. Pl. Succ.* 75.—Am. austr.
multiplex, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 70.—Bras.
Myosurus, Salm-Dyck, in DC. Prod. iii. 469 = Rhipsalis Myosurus.
myriocaulon, Mart. ex Pfeiff. Enum. Diagn. Cact. 115 = humilis.
myriophyllus, Gill. ex Otto, in Otto & Dietr. Allg. Gartenz. i. (1833) 365 = strigosus.
nanus, *DC. Prod.* iii. 470.—Ecuador.
Napoleonis, *R. Grah. in Bot. Mag.* t. 3458.—Ind. occ.
niger, *Salm-Dyck, Obs. Bot.* (1822) 4.—Am. austr.
nigricans, Hort. ex Pfeiff. Enum. Diagn. Cact. 83 = strictus.
nigricans, *Lem. Cact. Gen. Nov. et Sp.* 57.—Hab.?
nigripilis, *Phil. Fl. Atac.* 23.—Chili.
nitens, *Salm-Dyck, ex Otto & Dietr. Allg. Gartenz.* xiii. (1845) 354.—Bolivia.
nitidus, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 211.—Venezuela.
nobilis, Haw. Syn. Pl. Succ. 179 = Haworthii.
nycticalus, *Link, in Verh. Preuss. Ver. Gartenb.* x. (1834) 373. t. 4.—Mexic.
nyriocaulon, Sweet, Hort. Brit. ed. III. 285, sphalm. = *myriocaulon*.
obtusus, Haw. Rev. Pl. Succ. 70 = variabilis.
obtusus, Hort. ex Pfeiff. Enum. Diagn. Cact. 114 = nycticalus.
Ocamponis, *Salm-Dyck, Cact. Hort. Dyck.* ed. II. 220.—Mexic.; N. Granat.
ochracanthus, Hort. ex Pfeiff. Enum. Diagn. Cact. 78 = multangularis.
octogonus, Hort. ex Otto, in Otto & Dietr. Allg. Gartenz. i. (1833) 365 = albispinus, Curtisii.
Olfersii, *Salm-Dyck, Hort. Dyck.* 335.—Bras.
olivaceus, *Lem. in Rev. Hortic.* Sér. IV. viii. (1859) 643.—Ind. occ.
Ottonis, Pfeiff. Enum. Diagn. Cact. 47, sphalm. = Echinocactus Ottonis.
ovatus, *G. Don, in Loud. Hort. Brit.* 195; *Pfeiff. Enum. Diagn. Cact.* 102.—Chili.
oxygonus, *Otto, in Otto & Dietr. Allg. Gartenz.* i (1833) 365.—Bras.
oxygonus, Salm-Dyck, ex Foerst. Handb. Cact. ed. I. 393 = euphorbioides.
oxypetalus, DC. Prod. iii. 470 = Phyllocactus latifrons.

CEREUS:—
oxypetalus, Hort. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 37, nomen = Phyllocactus marginatus.
paniculatus, *DC. Prod.* iii. 466.—Ins. S. Doming.
panoplaeatus, Cels, ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 44 = pycnacanthus.
paradoxus, Hort. ex Steud. Nom. ed. II. i. 335 = Opuntia brasiliensis.
parasiticus, Haw. ex Steud. l. c. = Rhipsalis fasciculata.
parvisetus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 79.—Bras.
paucispinus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 34, 36; *et in Proc. Am. Acad.* iii. (1857) 285.—Texas.
Paxtonianus, *Monv. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 211.—Hab.?
pectinatus, Engelm. in A. Gray, Pl. Fendl. 50 = caespitosus.
pectiniferus, Labour. Monog. Cact. 320 = caespitosus.
pellucidus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 108.—Cuba.
pentagonus, *Haw. Syn. Pl. Succ.* 180.—Am. austr.
pentagonus, Hort. ex Foerst. Handb. Cact. ed. I. 389 = peruvianus.
pentalophus, *DC. in Mém. Mus. Par.* xvii. (1828) 117.—Mexic.
pentapterus, *Otto, ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 220.—N. Granat.
Pepinianus, *Lem. ex Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 354.—Bolivia.
pernambucensis, Lem. Nov. Cact. Gen. et Sp. 58 (err. typ. *fernambucensis*) = variabilis.
Perrotteti, Hort. ex Pfeiff. Enum. Diagn. Cact. 98 = lividus.
peruvianus, Hort. ex Foerst. Handb. Cact. ed. I. 392 = eburneus.
peruvianus, *Mill. Gard. Dict.* ed. VIII. n. 4; *Haw. Syn. Pl. Succ.* 179.—Am. austr.
Pfeifferi, *Parm. ex Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 370.—Reg. Argent.
Philippi, *Regel & Schmidt, Gartenfl.* (1882) 98.—Chili.
phoenicœus, *Engelm. & Bigel. in Pacif. Rail. Rep.* iv. 34. t. 4.—Am. bor. occ.
phyllanthoides, DC. Prod. iii. 469 = Phyllocactus phyllanthoides.
Phllyanthus, DC. l. c. = Phyllocactus Phyllanthus.
Pitajaya, DC. l. c. 466 = variabilis.
platycarpus, Zucc. ex Foerst. Handb. Cact. ed. I. 452 = Rhipsalis platycarpa.
platygonus, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 199.—Hab.?
polyacanthus, *Engelm. in A. Gray, Pl. Fendl.* 51.—Texas.
polyachaetus, *Reichb. in Tersch. Suppl. Cact. Verz.* 4.—Hab.?
polygonatus, Hort. ex Pfeiff. Enum. Diagn. Cact. 91 = eburneus.
polygonus, *DC. Prod.* iii. 466.—Ins. S. Doming.
polylophus, *DC. in Mém. Mus. Par.* xvii. (1828) 115.—Mexic.
polymorphus, *G. Don, in Loud. Hort. Brit.* 195.—Chili.
polymorphus, Hort. Monac. ex Foerst. Handb. Cact. ed. I. 472 = articulatus.
polyptychus, *Lem. Cact. Gen. Nov. et Sp.* 56.—Hab.?
Pottsii, Salm-Dyck, Cact. Hort. Dyck. ed. II. 208 = Greggii.
Princeps, *Hort. Wuerzb. ex Pfeiff. Enum. Diagn. Cact.* 108.—Am. bor. occ.
Pringlei, *S. Wats. in Proc. Am. Acad.* xx. (1885) 368.—Am. bor. occ.
prismaticus, Haw. ex Steud. Nom. ed. II. i. 335 = pentagonus.
prismaticus, *Salm-Dyck, in DC. Prod.* iii. 469.—Am. austr.
prismatiformis, Hort. ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
procumbens, *Engelm. in A. Gray, Pl. Fendl.* 50.—Mexic.
procumbens, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 203.—Texas.
propinquus, Salm-Dyck, ex Otto, in Otto & Dietr. Allg. Gartenz. i. (1833) 366 = pentalophus.
proteiformis, *Steud. Nom.* ed. II. i. 335.—Hab.?
pruinosus, Salm-Dyck, Cact. Hort. Dyck. ed. II. 47 = Echinocactus pruinosus.

CEREUS:—
pterandrus, Hemsl. Biol. Centr. Am. Bot. i. 544, sphalm. = seq.
pteranthus, Link, ex Dietr. in Otto & Dietr. Allg. Gartenz. ii. (1834) 209 = nycticalus.
pterocaulis, Hort. ex Pfeiff. Enum. Diagn. Cact. 140 = Rhipsalis paradoxa.
pterogonus, *Lem. Cact. Gen. Nov. et Sp.* 59.—N. Granat.
pugioniferus, Lem. Cact. Aliq. Nov. Desc. 30 = geometrizans, Gladiator.
pulchellus, *Pfeiff. Enum. Diagn. Cact.* 74.—Mexic.
pycnacanthus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 355.—Bolivia.
quadrangularis, Haw. Syn. Pl. Succ. 181 = caripensis.
quadrangularis, Hort. ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
quadrangulispinus, *Lem. ex Ehrenb. in Linnaea*, xix. (1847) 363.—Mexic.
quadricostatus, *Bello, in Anal. Soc. Esp. Hist. Nat.* x. (1881) 276.—Porto Rico.
Quintero, Hort. Gotting. ex Pfeiff. Enum. Diagn. Cact. 86 = chilensis,
Quisco, *C. Gay, Fl. Chil.* iii. 19.—Chili.
radicans, *DC. Prod.* iii. 468.—Am. austr.
ramosus, Karw. in Pfeiff. Enum. Diagn. Cact. 108 = baxaniensis.
ramulosus, Salm-Dyck, Hort. Dyck. 340 = Rhipsalis ramulosa.
recurvus, Link & Otto, ex Steud. Nom. ed. II. i. 335 = Echinocactus recurvus.
reductus, DC. Prod. iii. 463 = Echinocactus gibbosus.
reflexus, Hort. ex Steud. Nom. ed. II. i. 335 = virens.
regalis, *Haw. ex Spreng. Syst.* ii. 496.—Am. austr.
Reichenbachianus, Labour. Monog. Cact. 318 = caespitosus.
repandus, Haw. Syn. Pl. Succ. 183 = gracilis.
repandus, *Mill. Gard. Dict.* ed. VIII. n. 5.—Ind. occ.
reptans, Haw. ex Steud. Nom. ed. II. i. 335 = pentagonus.
reptans, Salm-Dyck, ex DC. Prod. iii. 468 = radicans.
resupinatus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* viii. (1840) 10.—Hab.?
retroflexus, Hort. ex Pfeiff. in Otto & Dietr. l. c. iii. (1835) 380 = virens.
rhombeus, Salm-Dyck, Hort. Dyck. 341 = Rhipsalis rhombea.
rigidispinus, *Lem. Hort. Univ.* (1840) 223. t. 1.—Mexic.
rigidispinus, *Muehlenp. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 12.—Mexic.
rigidus, Lem. ex Salm-Dyck, Cat. Hort. Dyck. ed. I. 34; ed. II. 52 = humilis.
Roemeri, *Engelm. in A. Gray, Pl. Fendl.* 50.—Texas.
Roemeri, *Muehlenp. in Otto & Dietr. Allg. Gartenz.* xvi. (1848) 19.—Hab.?
Roetteri, *Engelm. Cact. Mex. Bound.* 33. t. 41. f. 3–5.—Am. bor. occ.
roridus, Hort. Berol. ex Pfeiff. Enum. Diagn. Cact. 54. = Echinocactus pruinosus.
rosaceus, Hort. ex Pfeiff. l. c. 114 = nycticalus.
rostratus, *Lem. Cact. Aliq. Nov. Desc.* 29.—Ins. Antigua.
Royeni, Haw. Syn. Pl. Succ. 182 = lanuginosus.
Royeni, Hook. Bot. Mag. t. 3125 = Curtisi.
Royeni, *Mill. Gard. Dict.* ed. VIII. n. 7.—Am. austr.
rufispinus, *Engelm. in A. Gray, Pl. Fendl.* 50.—N. Mexic.
Russelianus, *Forb. Hort. Tour Germ.* 154, nomen.—Guiana.
Russellianus, Gardn. ex Hook. Bot. Mag. t. 3717 = Epiphyllum Russellianum.
Russellianus, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 201.—Laguayra.
Salm-Dyckianus, *Hemsl. Biol. Centr. Am. Bot.* i. 545.—Mexic.
Salm-Dyckianus, Hort. Vindob. ex Salm-Dyck, Cact. Hort. Dyck. ed. I. 31; ed. II. 49 = variabilis.
scandens, *Salm-Dyck, l. c.* ed. II. 219.—Guiana.
Scheerii, *Salm-Dyck, l. c.* 190.—Mexic.
Schelhasii, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iii. (1835) 314.—Hab.?
Schomburgkii, *Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 219.—Guiana?

CEREUS:—
Schottii, *Engelm. in Proc. Am. Acad.* iii. (1857) 288.—Am. bor. occ.
Schrankii, *Zucc. ex Pfeiff. Enum. Diagn. Cact.* 122.—Mexic.
Scopa, Salm-Dyck, ex DC. Prod. iii. 464 = Echinocactus Scopa.
Siedelii, *Lehm. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 200.—Hab.?
senilis, *Salm-Dyck, in DC. Prod.* iii. 464.—Am. austr.
sepium, *DC. l. c.* 467.—Ecuador.
sericeus, Hort. ex Pfeiff. Enum. Diagn. Cact. 173 = Opuntia clavarioides.
serpens, *DC. Prod.* iii. 470.—Ecuador.
serpentinus, *DC. l. c.* 467.—Mexic.
serruliflorus, *Haw. in Phil. Mag.* (1830) 109.—Ind. occ.
setaceus, *Salm-Dyck, in DC. Prod.* iii. 469.—Bras.
setiger, *Haw. in Phil. Mag.* (1830) 109.—Bras.
similis, *Meinsh. in C. Koch & Fint. Wochenschr.* i. (1858) 29.—Mexic.
Smithianus ×, Sweet, Hort. Brit. ed. II. 237, nomen = seq.?
Smithii ×, *Hort. Angl. ex Pfeiff. Enum. Diagn. Cact.* 111.
Spachianus, *Lem. Hort. Univ.* i. 225.—Mexic.
speciosissimus, *DC. Prod.* iii. 468.—Mexic.
speciosus, Sweet, Hort. Brit. ed. I. 172 = variabilis?
spinibarbis, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 86.—Chili.
spinosissimus, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 165.—Peruv.
spinulosus, *DC. in Mém. Mus. Par.* xvii. (1828) 117.—Mexic.
splendens, *Hort. Angl. ex Salm-Dyck, Cact. Hort. Dyck.* ed. II. 214.—Hab.?
splendidus, *Paxt. ex Steud. Nom.* ed. II. i. 336 (= Epiphylli sp.?).—Mexic.
squamulosus, Salm-Dyck, in DC. Prod. iii. 469 = Rhipsalis Mittleri.
stellatus, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iv. (1836) 258.—Mexic.
stramineus, *Engelm. Cact. Mex. Bound.* 35. t. 46.—Texas.
strictus, *DC. Prod.* iii. 465.—Am. austr.
strigosus, *Salm-Dyck, Hort. Dyck.* 334.—Chili.
subinermis, *Hemsl. Biol. Centr. Am. Bot.* i. 546.—Mexic.
sublanatus, *Salm-Dyck, Hort. Dyck.* 337.—Hab.?
subrepandus, *Haw. Suppl. Pl. Succ.* 78.—Ind. occ.
subrepandus, Hort. ex Pfeiff. Enum. Diagn. Cact. 86, 94 = chilensis, eriophorus.
subsquamatus, Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 380 = extensus.
subtortuosus, Hort. ex Foerst. Handb. Cact. ed. I. 409 = colubrinus.
subuliferus, *Salm-Dyck, in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 354.—Mexic.
superbus, *Ehrenb. in Bot. Zeit.* iv. (1846) 324.—Mexic.
Swartzii, *Griseb. Fl. Brit. W. Ind.* 301.—Ind. occ.
syringacanthus, *Pfeiff. Enum. Diagn. Cact.* 103.—Chili.
tenellus, *Salm-Dyck, ex Pfeiff. l. c.* 109.—Bras.
tenuis, DC. Prod. iii. 469 = Rhipsalis Myosurus.
tenuis, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* viii. (1840) 407.—Am. austr.
tenuispinus, Haw. in Phil. Mag. (1827) 125 = Rhipsalis Myosurus.
tenuissimus, *G. Don, in Sweet, Hort. Brit.* ed. III. 285, nomen.—Hab.?
tephracanthus, *Link & Otto, ex Steud. Nom.* ed. II. i. 336.—Am. austr.
Terscheckii, *Parm. ex Pfeiff. in Otto & Dietr. Allg. Gartenz.* v. (1837) 370.—Reg. Argent.
tetracanthus, *Labour. in Rev. Hortic.* Sér. IV. iv. (1855) 25.—Bolivia.
tetragonus, *Mill. Gard. Dict.* ed. VIII. n. 2; *Haw. Syn. Pl. Succ.* 180.—Am. austr.
thalassinus, *Otto & Dietr. Allg. Gartenz.* vi. (1838) 34.—Venezuela.
Thurberi, *Engelm. in Am. Journ. Sc.* Ser. II. xvii. (1854) 234.—N. Mexic.
tilophorus, Pfeiff. in Otto & Dietr. Allg. Gartenz. iii. (1835) 380 = virens.
Tinei, *Todaro, Ind. Sem. Hort. Panorm.* (1857) 39.—*Cf. in Linnaea*, xxix. (1857–58) 739.—Bras.?

CEREUS:—
Tonclianus, *Lem. Illustr. Hortic.* ii. (1855) *Misc.* 63.—Mexic.
tortuosus, *Forbes, ex Otto & Dietr. Allg. Gartenz.* vi. (1838) 35.—Reg. Argent.
triangularis, Haw. Syn. Pl. Succ. 180 = compressus.
triangularis, *Mill. Gard. Dict.* ed. VIII. n. 9.—Hab. ?
trichacanthus, Hort. Berol. ex Salm-Dyck, Cact. Hort. Dyck. ed. II. 46 = lutescens.
triglochidiatus, *Engelm. in A. Gray, Pl. Fendl.* 51.—N. Mexic.
trigonus, *Haw. Syn. Pl. Succ.* 181.—Ind. occ.
trigonus, Hort. ex Pfeiff. Enum. Diagn. Cact. 106 = variabilis.
trinitatensis, *Lem. & Herm. in Rev. Hortic.* Sér. IV. viii. (1859) 642.—Ind. occ.
triptcris, *Salm-Dyck, in DC. Prod.* iii. 468.—Am. austr.
triquetcr, *Haw. Syn. Pl. Succ.* 181.—Am. austr.
truncatus, Sweet, Hort. Brit. ed. I. 172 = Epiphyllum Altenstcinii.
tubcrosus, *Pfeiff. Enum. Diagn. Cact.* 102.—Mexic.
tubcrosus, *Poselg. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 135.—Texas.
tubiflorus, Endl. in Harting. Parad. Vindob. t. 71 (= *Echinopsis Zuccarini*).—Hab. ?
tubiflorus, Pfeiff. Enum. Diagn. Cact. 71 = Echinocactus tubiflorus.
tunicatus, Pfeiff. l. c. 173 in syn. = Opuntia tunicata.
turbinatus, *Pfeiff. in Otto & Dietr. Allg. Gartenz.* iii. (1835) 314.—Hab. ?
Tweedici, *Hook. Bot. Mag.* t. 4498.—Reg. Argent.
undatus, Haw. in Phil. Mag. (1830) 110 = Napoleonis.
undatus, *Hort. Berol. ex Pfeiff. Enum. Diagn. Cact.* 94.—Hab. ?
undiflorus, *Wright, in Sauv. Fl. Cub.* 59.—Cuba.
undulatus, Hort. Berol. ex D. Dietr. Syn. Pl. iii. 104 = undatus.
undulatus, Hort. Dresd. ex Pfeiff. Enum. Diagn. Cact. 107 = acutangulus.
undulosus, DC. Prod. iii. 467 = variabilis.
urcacanthus, *C. F. Foerst. in Hamb. Gartenz.* xvii. (1861) 166.—Peruv.
validus, *Haw. in Phil. Mag.* x. (1831) 414.—Am. austr.
variabilis, Engelm. Cact. Mex. Bound. 40 = Princeps.
variabilis, *Pfeiff. Enum. Diagn. Cact.* 105.—Am. austr.
Vasmerii, Young, Fl. Texas, 276 = Princeps.
violaceus, *Lem. Cact. Gen. Nov. et Sp.* 57.—Hab. ?
virens, *DC. in Mém. Mus. Par.* xvii. (1828) 116.—Mexico.
viridiflorus, *Engelm. in A. Gray, Pl. Fendl.* 50.—Texas.
xanthochactus, *Reichb. in Tersch. Suppl. Cact. Verz.* 4.—Hab. ?

CERINTHE, [Tourn.] Linn. Syst. ed. I. (1735). *BORAGINEAE*, Benth. & Hook. f. ii. 864.
CERANTHE, Opiz, Seznam, 27 (1852).
acuta, Moench, Meth. 520 = major.
alpina, *Kit. ex Schult. Oestr. Fl.* ed. II. i. 353.—Europ.; As. Min.
alpina, Vis. Fl. Dalmat. Suppl. (1872) 94; fide Nym. Consp. 513 = longiflora.
aperta, [Clairv.] Man. Herb. 53 = major.
aspera, Roth, Catalect. i. 33 = major.
atropurpurea, Vis. ex A. DC. in DC. Prod. viii. 4 = retorta.
auriculata, Tenore, Ind. Scm. Hort. Neap. (1830) 10 = minor.
cilicica, Kotschy, ex Boiss. Pl. Or. Nov. dec. ii. 7 = minor.
cleiostoma, Boiss. & Sprun. Diagn. Scr. I. iv. 44 = minor.
connivens, [Clairv.] Man. Herb. 54 = minor.
echioides, Linn. Sp. Pl. 137 = Onosma echioides.
glabra, Mill. Gard. Dict. ed. VIII. n. 2; DC. Fl. Fr. iii. 619 = alpina.
glabra, Scop. ex Nym. Consp. 513 = minor.
glauca, Moench, Meth. 520 = major.
gymnandra, Gasp. in Rendic. Acc. Sc. Nap. i. (1863) 72 = major.
hirsuta, *Wettstein, ex Stapf, in Denkschr. Acad. Wien*, I. (1885) 85.—Lycia.
longiflora, *Viv. App. Alt. ad Fl. Cors.* (1830) 3.—Italia.

CERINTHE:—
lurida, Salisb. Prod. 116 = minor.
macrophylla, Boiss. ex Boiss. Pl. Or. Nov. dec. ii. 7. = minor.
maculata, All. Fl. Pedem. i. 178 = alpina.
maculata, Linn. Sp. Pl. 137 = minor.
major, *Linn. l. c.* 136.—Europ.; As. Min.
minor, *Linn. l. c.* 137.—Europ.; As. Min.; Persia.
orientalis, Linn. Cent. i. 7 = Onosma orientalis.
pseudo-aspera, Schur, Enum. Pl. Transs. 470 = minor.
purpurea, Vis. in Flora, xii. (1829) Erg. 8 = retorta.
quinquemaculata, Wahlenb. Fl. Carpat. 51 = alpina.
retorta, *Sibth. & Sm. Fl. Graec.* ii. 60. t. 171.—Graecia.
Smithiae, Kern. ex Nym. Consp. 513 = longiflora.
strigosa, Reichb. Fl. Germ. Excurs. 340 = major.
suevica, Martens, Reise Vened. i. 44; ex Reichb. Ic. Crit. v. 40. t. 658 = alpina.
tenuiflora, *Bertol. Fl. Ital.* ii. 325 = longiflora.
versicolor, Hall. ex Steud. Nom. ed. II. i. i. 336 = major.

CERINTHOPSIS, Kotschy, ex Benth. & Hook. f. Gen. ii. 849 (1876) = **Solenanthus**, Ledeb. (Boragin.).
kurdica, Kotschy, ex Boiss. Fl. Orient. iv. 271 = S. cerinthioides.

CERIONANTHUS, Schott, ex Roem. & Schult. Syst. iii. 43 (1818) = **Cephalaria**, Schrad. (Dipsac.).
alpinus, Schott, ex Roem. & Schult. l. c. = Ceph. alpina.
corniculatus, Schott, ex Roem. & Schult. l. c. 49 = Ceph. corniculata.
cretaceus, Schott, ex Roem. & Schult. l. c. 51 = Ceph. corniculata.
laevigatus, Schott, ex Roem. & Schult. l. c. 49 = Ceph. corniculata.
leucanthus, Schott, ex Roem. & Schult. l. c. 47 = Ceph. leucantha.
syriacus, Schott, ex Roem. & Schult. l. c. 46 = Ceph. syriaca.
tataricus, Schott, ex Roem. & Schult. l. c. 51 = Ceph. tatarica.
Vaillantii, Schott, ex Roem. & Schult. l. c. 46 = Ceph. syriaca.

CERIOPS, Arn. in Ann. Nat. Hist. i. (1838) 363. *RHIZOPHOREAE*, Benth. & Hook. f. i. 679.
Boviniana, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 112.—Madag.
Candolleana, *Arn. in Ann. Nat. Hist.* i. (1838) 364.—Reg. trop.
Candolleana, Naves, in Blanco, Fl. Filip. ed. III. t. 415 = Roxburghiana.
Forsteniana, *Blume, Mus. Bot. Lugd. Bat.* i. 143.—Ins. Molucc.
globulifera, Bor. ex Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 112 = Boviniana.
lucida, *Miq. Fl. Ind. Bat. Suppl.* i. 325.—Malaya.
mossambicensis, Klotzsch, in Peters, Reise Mossamb. Bot. i. 71 = Candolleana.
pauciflora, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 218.—Polynes.
Roxburghiana, *Arn. in Ann. Nat. Hist.* i. (1838) 364.—Reg. trop.
Zippeliana, *Blume, Mus. Bot. Lugd. Bat.* i. 143.—N. Guinea.

CERISCUS, Gaertn. Fruct. i. t. 28 (1788) = **Randia**, Linn. (Rubiac.).
malabaricus, Gaertn. l. c. = R. dumetorum.

CERISCUS, Nees, in Flora, viii. (1825) 116 = **Tarenna**, Gaertn. (Rubiac.).
fragrans, Nees, l. c. (= *Webera fragrans*).—Ind. or.; Malaya.

CERIUM, Lour. Fl. Cochinch. 135. *INC. SED.*
spicatum, *Lour. l. c.* 136.—China.

CEROCARPUS, Hassk. in Flora, xxv. (1842) Beibl. ii. 36 = **Eugenia**, Linn. (Myrtac.).
aqueus, Hassk. l. c. = E. aquea.
bifaria, Colebr. in Wall. Cat. n. 3606 = E. Wallichii.

CEROCHILUS, Lindl. in Gard. Chron. (1854) 87 = **Hetaeria**, Blume (Orchid.).
rubens, Lindl. l. c. = H. rubens.

CEROPEGIA, Linn. Gen. ed. I. 65 (1737). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 779.
APEGIA, Neck. Elem. i. 251 (1790).
NIOTA, Adans. Fam. ii. 172 (1763).
SYSTEPHA, Burch. Trav. i. 546 (1822).
abyssinica, *Decne. in DC. Prod.* viii. 644.—Abyss.
acuminata, Dalz. & Gibs. Bomb. Fl. 117 = tuberosa.
acuminata, *Roxb. Pl. Corom.* i. 12. t. 8.—Ind. or.
affinis, *Vatke, in Linnaea*, xl. (1876) 218.—Abyss.
africana, *R. Br. in Bot. Reg.* t. 626.—Afr. austr.
albiflora, *Hook. f. Fl. Brit. Ind.* iv. 75.—Ind. or.
ampliata, *E. Mey. Comm. Pl. Afr. Austr.* 194.—Afr. austr.
angustifolia, Dalz. in Hook. Kew Journ. ii. (1850) 259 = attenuata.
angustifolia, Vahl, ex Decne. in DC. Prod. viii. 503 = Secamone saligna.
angustifolia, *Wight, Contrib.* 31.—Reg. Himal.
aphylla, Haw. Syn. Pl. Succ. 13 = dichotoma.
arcta, *Nimmo, in J. Grah. Cat. Bomb. Pl.* 118, nomen.—Ind. or.
aristolochioides, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 263.—Afr. trop.
Arnottiana, *Wight, Contrib.* 32.—Ind. or.; Burma.
attenuata, *Hook. Ic. Pl.* t. 867.—Ind. or.; Burma.
australis, *Lodd. Cat. ex Loud. Hort. Brit.* 95, nomen.—Austral.
Barklyi, *Hook. f. Bot. Mag.* t. 6315.—Afr. austr.
Beddomei, *Hook. f. Fl. Brit. Ind.* iv. 75.—Ind. or.
biflora, *Linn. Sp. Pl.* 211.—Zeylan.
Bowkeri, *Harv. in Proc. Dubl. Zool. & Bot. Assoc.* i. (1859) 254.—Afr. austr.
brevicollis, *Hook. f. Fl. Brit. Ind.* iv. 74.—Ind. or.
brevitubulata, Bedd. Ic. Pl. Ind. Or. i. 38 = Brachystelma Beddomei.
bulbosa, *Roxb. Pl. Corom.* i. 11. t. 7.—Ind. or.
campanulata, *G. Don, Gen. Syst.* iv. 112.—Afr. trop.
candelabriformis, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Candelabrum.
Candelabrum, Heyne, ex Decne. in DC. Prod. viii. 643 = bulbosa.
Candelabrum, *Linn. Sp. Pl.* 211.—Ind. or.
Candelabrum, Lour. Fl. Cochinch. 114 = Loureiri.
Candelabrum, Roxb. Hort. Beng. 21; Fl. Ind. ii. 27 = tuberosa.
Candelabrum, Thw. Enum. Pl. Zeyl. 199 = intermedia.
carnosa, *E. Mey. Comm. Pl. Afr. Austr.* 193.—Afr. austr.
ciliata, *Wight, Ic.* t. 1262.—Ind. or.
convolvuloides, *A. Rich. Tent. Fl. Abyss.* ii. 47.—Abyssin.
cordata, *Lour. Fl. Cochinch.* 114.—Cochinch.
Cumingiana, *Decne. in DC. Prod.* viii. 643.—Ins. Philipp.
curviflora, *Hassk. in Flora*, xl. (1857) 102.—Java.
Decaisneana, *Wight, Ic.* t. 1259.—Ind. or.
Decaisneana, Miq. ex Hook. f. Fl. Brit. Ind. iv. 42 = Tylophora tenuis.
dichotoma, *Haw. Syn. Pl. Succ.* 13.—Ind. or.
edulis, Hort. ex Decne. in DC. Prod. viii. 643 = bulbosa.
elegans, *Wall. in Bot. Mag.* t. 3015.—Ind. or.
Elliottii, *Hook. f. Fl. Brit. Ind.* iv. 70.—Ind. or.
ensifolia, *Bedd. in Madr. Journ. Sc.* Scr. III. i. (1864) 52.—Ind. or.
esculenta, Edgew. in Journ. Linn. Soc. vi. (1862) 204 = bulbosa.
fimbriata, *E. Mey. Comm. Pl. Afr. Austr.* 194.—Afr. austr.
fimbriifera, *Bedd. in Madr. Journ. Sc.* Scr. III. i. (1864) 53.—Ind. or.
fusca, *C. Bolle, in Bonplandia*, ix. (1861) 51.—Ins. Canar.
Gardneri, *Hook. Bot. Mag.* t. 5306; *Thw. Enum. Pl. Zeyl.* 199.—Zeylan.
gracilis, *Bedd. in Madr. Journ. Sc.* Scr. III. i. (1864) 53.—Ind. or.
hirsuta, Hochst. ex Decne. in DC. Prod. viii. 644 = abyssinica.
hirsuta, *Wight & Arn. Contrib.* 30.—Ind. or.
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 73.—Reg. Himal.
Horsfieldiana, *Miq. Fl. Ind. Bat.* ii. 528.—Java.
infundibuliformis, *E. Mey. Comm. Pl. Afr. Austr.* 194.—Afr. austr.
intermedia, *Wight, Ic.* t. 1263.—Ind. or.

CEROPEGIA:—
Jacquemontiana, Decne. in DC. Prod. viii. 641 = hirsuta ?
juncea, *Roxb. Pl. Corom.* i. 12.—Ind. or.
lanceolata, Wight & Arn. Contrib. 31 = longifolia.
Lawii, *Hook. f. Fl. Brit. Ind.* iv. 67.—Ind. or.
linearis, *E. Mey. Comm. Pl. Afr. Austr.* 194.—Afr. austr.
longiflora, Poir. Encyc. ii. 177 = bulbosa.
longifolia, Hook. f. & Tboms. in Fl. Brit. Ind. iv. 72 = angustifolia.
longifolia, *Wall. Pl. As. Rar.* i. 56. t. 73.—Reg. Himal.
Loureirii, *G. Don, Gen. Syst.* iv. 112.—Cochinch.
lucida, *Wall. Pl. As. Rar.* ii. 33. t. 139.—Reg. Himal.; Burma.
Lushii, R. Grab. in Bot. Mag. t. 3300 = bulbosa.
macrantha, *Wight, Contrib.* 31.—Reg. Himal.
maculata, *Bedd. in Madr. Journ. Sc.* Ser. III. i. (1864) 52.—Ind. or.
madagascariensis, *Decne. in DC. Prod.* viii. 642.—Madag.
Metziana, *Miq. Anal. Bot. Ind.* iii. 11.—Ind. or.
Meyeri, *Decne. in DC. Prod.* viii. 645.—Afr. austr.
micans, *Nimmo, in J. Grah. Cat. Bomb. Pl.* 118, nomen.—Ind. or.
mucronata, Roth, Nov. Pl. Sp. 179 = tuberosa.
multiflora, *Baker, in Saund. Refug. Bot.* t. 10.—Afr. austr.
Munroi, Wight, Ic. t. 1254 = fimbriifera ?
mysorensis, *Wight, l. c.* t. 846.—Ind. or.
nilotica, *Kotschy, in Sitzb. Acad. Wien. Math. Nat.* li. (1865) 356.—Afr. trop.
nitida, *Poir. Encyc. Suppl.* ii. 178.—Ins. S. Doming.
obtusa, *Lour. Cochinch.* 114.—Cochinch.
obtusiloba, *Fawc. in Forbes, Nat. Wand. in East. Archip.* 511.—Ins. Timor.
oculata, *Hook. Bot. Mag.* t. 4093.—Ind. or.
odorata, *Nimmo, in J. Grah. Cat. Bomb. Pl.* 118.—Ind. or.
ophiocephala, Dalz. in Hook. Kew Journ. ii. (1850) 259 = hirsuta.
palustris, Pursb, Fl. Am. Sept. i. 184 = Vincetoxicum palustre.
peltata, *Regel, Cat. Pl. Hort. Aksakov.* 31.—Hab. ?
parasitica, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 12.—Malaya.
pubescens, *E. Mey. Comm. Pl. Afr. Austr.* 193.—Afr. austr.
pubescens, *Wall. Pl. As. Rar.* ii. 81. t. 187.—Reg. Himal.
pusilla, *Wight & Arn. in Wight, Contrib.* 31.—Ind. or.
ringens, *A. Rich. Tent. Fl. Abyss.* ii. 48.—Abyssin.
sagittata, Linn. Mant. 215 = Microloma sagittatum.
Sandersoni, *Decne. ex Hook. Bot. Mag.* t. 5792.—Natal.
sinuata, *Decne. ex A. Rich. Tent. Fl. Abyss.* ii. 47.—Abyssin.
sinuata, Poir. Encyc. Suppl. ii. 179 = Microloma lineare.
Sororia, *Harv. in Hook. Bot. Mag.* t. 5578.—Afr. austr.
sphenanantha, Wigbt & Arn. in Wigbt, Contrib. 31 = elegans.
sphenanthera, Decne. in DC. Prod. viii. 643 = praec.
spiralis, Hook. f. & Thoms. in Fl. Brit. Ind. iv. 67 = attenuata.
spiralis, *Wight, Ic.* t. 1267.—Ind. or.
squamulata, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 263. t. 9.—Arabia.
stapeliaeformis, *Haw. in Phil. Mag.* (1827) 121.—Afr. austr.
stellata, *Regel, Cat. Pl. Hort. Aksakov.* 31.—Hab. ?
Stephanotis, Poir. ex Decne. in DC. Prod. viii. 621 = Stephanotis Isaura.
Stepbanotis, *Roem. & Schult. Syst.* vi. 4.—Madag.
Steudneri, *Vatke, in Linnaea*, xl. (1876) 217.—Abyssin.
Stocksii, *Hook. f. Fl. Brit. Ind.* iv. 74.—Ind. or.
tenuiflora, Willd. Sp. Pl. i. 1276 = Microloma lineare.
tenuifolia, Linn. Mant. ii. 346 = Microloma lineare.
Tbwaitesii, *Hook. Bot. Mag.* t. 4758.—Zeylan.
torulosa, *E. Mey. Comm. Pl. Afr. Austr.* 194.—Afr. austr.
trichantha, *Hemsl. in Journ. Bot.* xxiii. (1885) 286.—China.
tuberosa, *Roxb. Pl. Corom.* i. 12. t. 9.—Ind. or.
tuberosa, Wall. Cat. n. 8145 A = intermedia.
variegata, *Decne. in Ann. Sc. Nat.* Sér. II. ix. (1838) 262.—Arabia.

CEROPEGIA:—
Vignaldiana, *A. Rich. Tent. Fl. Abyss.* ii. 48.—Abyss.
vincaefolia, Hook. Bot. Mag. t. 3740 = hirsuta.
Walkeriae, *Wight, Ic.* t. 1266.—Zeylan.
Wallichii, *Wight, Contrib.* 32.—Reg. Himal.
Wightii, *R. Grah. in Bot. Mag.* t. 3267.—Ind. or.

CEROPHORA, Rafin. Alsog. Am. 11 (1838) = **Myrica**, Tourn.
angustifolia, Rafin. l. c. = M. cerifera.
inodora, Rafin. l. c. = M. cerifera.
lanceolata, Rafin. l. c. = M. cerifera.
spicans, Rafin. l. c. 12 = M. cerifera.

CEROPHYLLUM, Spacb, Hist. Vég. Phan. vi. 152 (1838) = **Ribes**, Linn. (Saxifrag.).
Douglasii, Spach, l. c. 153 = R. cereum.
inebrians, Spach, l. c. 154 = R. cereum.

CEROXYLON, Humb. & Bonpl. Pl. Aequin. l. t. 1, 1 b (1807). *PALMAE*, Benth. & Hook. f. iii. 905.
BEETHOVENIA, Engelm. in Linnaea, xxxiii. (1864–65) 677.
KLOPSTOCKIA, Karst. in Linnaea, xxviii. (1856) 251.
alpinum, Steud. Nom. ed. II. i. 337 = seq.
andicolum, *Humb. & Bonpl. Pl. Aequin.* l. t. 1.—N. Granat.
australe, Mart. Hist. Nat. Palm. iii. 314.—Juania australis.
coarctatum, *H. Wendl. in Kerch. Palm.* 238.—N. Granat.
ferrugineum, *Regel, Gartenfl.* (1879) 163. t. 977.—N. Granat.
interruptum, *H. Wendl. in Bonplandia*, viii. (1860) 70.—N. Granat.
Klopstockia, *Mart. Hist. Nat. Palm.* iii. 314.—Venezuela.
niveum, Hort. Belg. ex H. Wendl. in Kercb. Palm. 239 = Diplothemium caudescens.
parvifrons, *H. Wendl. l. c.*—N. Granat.
pityrophyllum, Mart. ex H. Wendl. l. c. = Cocos pityrophylla.
quindiuense, *H. Wendl. in Bonplandia*, viii. (1860) 70.—N. Granat.
utile, *H. Wendl. l. c.*—N. Granat.
Vogelianum, *H. Wendl. in Kerch. Palm.* 239.—N. Granat.

CERQUEIRIA, Berg, in Linnaea, xxvii. (1854) 5 = **Myrcia**, DC. (Myrtac.).
Sellowiana, Berg, l. c. 6.—Bras.

CERQUIERIA, Bentb. & Hook. f. Gen. i. 716 (1865) sphalm. = praec.

CERRARIA, Tausch, in Flora, xvii. (1834) i. 347, sphalm. = CERVARIA, Minuart = **Peucedanum**, Linn. (Umbellif.).

CERRIS, Rafin. Alsog. Am. 29 (1838) = **Quercus**, Linn.
australis, Rafin. l. c. = Q. Cerris.
austriaca, Rafin. l. c. = Q. Cerris.
crinita, Rafin. l. c. = Q. Cerris.
dulcis, Rafin. l. c.—Am. bor.
macrocarpa, Rafin. l. c. = Q. macrocarpa.
microphyla, Rafin. l. c. = Q. micropbylla.
oliveformis, Rafin. l. c. = Q. macrocarpa.
paliphleos, Rafin. l. c.—Am. bor.
salicifolia, Rafin. l. c. = Q. salicifolia.

CERUANA, Forsk. Fl. Aegypt. Arab. 153 (1775). *COMPOSITAE*, Benth. & Hook. f. ii. 261.
fruticosa, Less. Syn. Comp. 202 = pratensis.
montana, Aucher, ex DC. Prod. vii. 287 = Chrysophtbalmum sternutatorium.
pratensis, *Forsk. Fl. Aegypt. Arab.* p. lxxiv.—Aegypt.
rotundifolia, Cass. in Dict. Sc. Nat. xli. 123 = pratensis.
Schimperi, Boiss. Diagn. Ser. I. xi. 4 = Asteriscus Schimperi.
senegalensis, *DC. Prod.* v. 488.—Afr. trop.

CERUCHIS, Gaertn. ex Schreb. Gen. 543 (1791) = **Spilanthes**, Jacq. (Compos.).

CERVANTESIA, Ruiz & Pav. Prod. 39. t. 7 (1794). *SANTALACEAE*, Benth. & Hook. f. iii. 222.
bicolor, *Cav. Ic.* 49. t. 475.—Peruv.
Kunthiana, *Baill. Adansonia*, ii. (1861–62) 373. t. 11.—Peruv.
tomentosa, A. DC. Prod. xiv. 692 = bicolor.
tomentosa, H. B. & K. Nov. Gen. et Sp. vii. 139 = Kunthiana.
tomentosa, *Ruiz & Pav. Prod. Fl. Per.* iii. 20. t. 7.—Peruv.

CERVARIA, Minuart, ex Linn. Diss. Dass. 31 (1747) (Amoen. Acad. i. 142); Gaertn. Fruct. i. 90. t. 21 = **Peucedanum**, Linn. (Umbellif.).
alsatica, Gaud. Fl. Helv. ii. 327 = P. alsaticum.
angustifolia, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 395 = P. angustifolium ?
glauca, Gaud. Fl. Helv. ii. 326 = P. Cervaria.
laevis, Gaud. l. c. 327, in obs. = P. Cervaria ?
latifolia, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 395 = P. latifolium.
nigra, Bernh. Syst. Verz. Erf. 171 = P. Cervaria.
Oreoselinum, Gaud. Fl. Helv. ii. 324 = P. Oreoselinum.
rigida, Moench, Metb. 95 = P. Cervaria.
Rivini, Gaertn. Fruct. 91. t. 21 = P. Cervaria.

CERVIA, Rodrig. ex Lag. Gen. et Sp. Nov. 7 (1816). *CONVOLVULACEAE* ? (Cf. Benth. & Hook. f. ii. 868).
saturejaefolia, *Rodrig. ex Lag. l. c.* —Hispan.

CERVICINA, Delile, Fl. Egypte, 150 (1813) = **Wahlenbergia**, Schrad. (Campanulac.).
campanuloides, Delile, l. c. t. 5. f. 2 = W. Cervicina.
tomentosa, Perr. ex Steud. Nom. ed. II. i. 337; ii. 740. = Vahlia tomentosa.

CERVISPINA, [Dill.] Moench, Meth. 686 (1794) = **Rhamnus**, Tourn.
cathartica, Mocnch, l. c. = R. cathartica.
infectoria, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 356 = R. infectoria.
saxatilis, Fourr. l. c. = R. saxatilis.
Villarsii, Fourr. l. c. = R. infectoria.

CESATIA, Endl. in Ann. Wien. Mus. ii. (1838) 200 = **Trachymene**, Rudge (Umbellif.).
ornata, Endl. l. c. = T. eriocarpa.
ovata, Steud. Nom. ed. II. i. 337, sphalm. = praec.

CESPEDESIA, Goudot, in Ann. Sc. Nat. Sér. III. ii. (1844) 368. *OCHNACEAE*, Bentb. & Hook. f. i. 320.
Bonplandi, *Goudot, l. c.* 370.—N. Granat.
macrophylla, *Seem. Bot. Voy. Herald*, 97.—Panama.
spathulata, *Planch. in Hook. Lond. Journ. Bot.* v. (1846) 647.—Peruv.

CESTICHIS, Thou. Orch. Afr. t. 90 (1822) = **Liparis**, Rich. (Orchid.).

CESTRINUS, Cass. in Dict. Sc. Nat. viii. 24 (1817) = **Centaurea**, Linn. (Compos.).
carthamoides, Cass. l. c. = Cent. Cbamaerhaponticum.

CESTRON, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = seq.

CESTRUM, Linn. Hort. Cliff. 491 (1737); Coroll. Gen. 3 (1737). *SOLANACEAE*, Benth. & Hook. f. ii. 904.
HABROTHAMNUS, Endl. Gen. 667 (1836).
LOMERIA, Rafin. Sylva Tellur. 56 (1838).
MEYENIA, Schlecbt. in Linnaea, viii. (1833) 251.
PARQUI, Adans. Fam. ii. 219 (1763).
WADEA, Rafin. Sylva Tellur. 56 (1838).
acuminatissimum, *Dun. in DC. Prod.* xiii. i. 627.—Guiana.
acuminatum, *Sweet, Hort. Brit.* ed. II. 388.—Mexic.
affine, *H. B. & K. Nov. Gen. et Sp.* iii. 60.—Ecuador.
affine, Pers. Syn. i. 230 = auriculatum.
alaternoides, *Desf. Tabl.* ed. I. 70.—Ind. occ.
albo-punctatum, *Dun. in DC. Prod.* xiii. i. 635.—Peruv.
album, *Ferrero, ex Dun. l. c.* 605.—Cuba.
Amelanchier, *Dun. l. c.* 662.—Hab. ?
amictum, *Schlecht. in Linnaea*, vii. (1832) 64.—Bras.
amygdalifolium, *Dun. in DC. Prod.* xiii. i. 653.—Mexic.

CESTRUM:—
anagyridium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = seq.
Anagyris, *Dun. in DC. Prod.* xiii. 1. 608.—Mexic.
angustifolium, *Lodd. Bot. Cab.* t. 618.—Ind. occ.
aronioideum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Amelanchier.
arvense, *Vell. Fl. Flum.* 101; iii. t. 7.—Bras.
atrovirens, Dun. in DC. Prod. xiii. 1. 648 = Sessea atrovirens.
aurantiacum, *Lindl. Bot. Reg.* (1844) *Misc.* 71.—Guatimal.
aurantiacum, Mey. ex Steud. Nom. ed. II. i. 337 = Freylinia cestroides.
auriculatum, *L'Hérit. Stirp. Nov.* 71. t. 35.—Am. austr.
auriculatum, Ruiz & Pav. Fl. Per. ii. 28. t. 155 f. a = hediundinum.
axillare, Vell. Fl. Flum. 101; iii. t. 6 = laevigatum.
Bella-Sombra, *Dun. in DC. Prod.* xiii. 1. 632.—Ins. Canar.
Benthami, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 151.—Mexic.
Billbergianum, *Beurling, in Vet. Acad. Handl.* (1854) 140.—Panama.
bogotense, Willd. ex Roem. & Schult. Syst. iv. 807 = mariquitense.
bonariense, *Dun. in DC. Prod.* xiii. 1. 666.—Reg. Argent.
braeteatum, *Link & Otto, Ic. Pl. Rar.* i. 11. t. 6.—Bras.
bupleurifolium, *Dun. in DC. Prod.* xiii. 1. 628.—Bras.
buxifolium, *H. B. & K. Nov. Gen. et Sp.* iii. 57.—N. Granat.
calycinum, *H. B. & K. l. c.* 58.—Peruv.
calycinum, Willd. ex Schlecht. in Linnaea, vii. (1832) 64 = viridiflorum.
campanulatum, Lam. Encyc. i. 688 = Acnistus aggregatus.
campestre, *Griseb. in Goett. Abh.* xxiv. (1879) 244.—Reg. Argent.
cancellatum, *Dun. in DC. Prod.* xiii. 1. 657.—Peruv.
candidum, *Seem. & Schmidt, in Flora*, xxvii. (1844) 496.—Mexic.
cauliflorum, Jacq. Hort. Schoenb. iii. 41. t. 325 = Acnistus arborescens.
cauliflorum, Sieber, ex Bercht. & Presl, Rostl. i. Solanae. 52 = Acnistus Plumieri.
chlamidatum, *Dun. in DC. Prod.* xiii. 1. 645.—Bras.
chloranthum, *Dun. l. c.* 636.—Ins. Trinitat.
citrifolium, *Retz. in Hoffm. Phyt. Blaetter*, i. (1803) 36.—Hab.?
Claussenі, *Dun. in DC. Prod.* xiii. 1. 637.—Bras.
collinum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 160.—Bras.
confertiflorum, *Schlecht. in Linnaea*, xix. (1847) 263.—Mexic.
confertum, Mill. Gard. Dict. ed. VIII. n. 5 = vespertinum.
confertum, *Ruiz & Pav. Fl. Per.* ii. 29. t. 153.—Peruv.
conglomeratum, *Ruiz & Pav. l. c.* 29. t. 156.—Peruv.
corcovadense, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 160.—Bras.
cordatum, *Schott, ex Sendt. in Mart. Fl. Bras.* x. 224.—Bras.
coriaceum, *Miers, in Hook. Lond. Journ. Bot.* v (1846) 161.—Bras.
corymbosum, *Schlecht. in Linnaea*, vii. (1832) 57.—Bras.
cuspidatum, *Sendt. in Mart. Fl. Bras.* x. 223.—Bras.
cyaneum, Schlecht. ex Dun. in DC. Prod. xiii. 1. 673 = Iochroma Schlechtendalianum.
daphnoides, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 524.—Ind. occ.
depauperatum, *Dun. in DC. Prod.* xiii. 1. 660.—Ins. Guadalup.
discolor, *Dun. l. c.* 609.—Peruv.
diurnum, Domb. ex Dun. in DC. l. c. 667 = leptanthum.
diurnum, H. B. & K. Nov. Gen. et Sp. iii. 61 = Forsythi.
diurnum, *Linn. Sp. Pl.* 191.—Ind. occ
dubium, *Steud. Nom.* ed. II. i. 337.—Hab.?
dumetorum, *Schlecht. in Linnaea*, vii. (1832) 61.—Bras.

CESTRUM:—
Ehrenbergi, *Schlecht. ex Dun. in DC. Prod.* xiii. 1. 613.—Mexic.
elegans, *Schlecht. in Linnaea*, xix. (1847) 261.—Mexic.
elongatum, Hort. ex Steud. Nom. ed. II. i. 337 = candidum.
Endlicheri, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 151.—Peruv.
eriochiton, *Sendt. in Mart. Fl. Bras.* x. 217.—Bras.
euanthes, *Schlecht. in Linnaea*, vii. (1832) 60.—Bras.
exstipulatum, *Ledeb. in Schrad. N. Journ. Bot.* iv. (1810) 65.—Am. austr.
fasciculare, Otto, ex Dun. in DC. Prod. xiii. 1. 673 = Toxicophlaea cestroides.
fasciculatum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 151.—Mexic.
fastigiatum, *Jacq. Hort. Schoenb.* iii. 44. t. 330.—Ind. occ.
fastigiatum, Jan, Cat. Hort. Parm. 189, 10 = diurnum.
Faucheri, *Dun. in DC. Prod.* xiii. 1. 640.—Hab.?
ferrugineum, *A. Rich. Fl. Cub. Fanerog.* ii. 126.—Cuba.
floribundum, Willd. ex Roem. & Schult. Syst. iv. 807 = tenuiflorum.
foetidissimum, Jacq. Hort. Schoenb. iii. 43. t. 329 = nocturnum.
foetidum, *Medic. in Act. Acad. Theod. Palat.* iv. *Phys.* (1780) 192.—Hab.?
foetidum, Salisb. Prod. 111 = auriculatum.
Forsythi, *Dun. in DC. Prod.* xiii. 1. 604.—Ind. occ.
frutescens, *Aubl. Pl. Gui.* i. 233.—Guiana.
Gardneri, *Sendt. in Mart. Fl. Bras.* x. 208.—Bras.
glaucescens, *Sendt. l. c.* 222.—Bras.
glomeratum, *Schott, ex Sendt. l. c.* 212.—Bras.
graciliflorum, *Dun. in DC. Prod.* xii. 1. 669.—Mexic.
granadense, Willd. ex Roem. & Schult. Syst. iv. 807 = tomentosum.
grandiflorum, Hort. ex Steud. Nom. ed. II. i. 337 = laurifolium.
grandistipulum, *Sendt. in Mart. Fl. Bras.* x. 214.—Bras.
Hartwegi, *Dun. in DC. Prod.* xiii. 1. 599.—Mexic.
Hediunda, Lam. Encyc. i. 687 = auriculatum.
Hediunda, Matr. ex Dun. in DC. Prod xiii. 1. 647 = hediundinum.
hediundinum, *Dun. l. c.*—Am. austr.
heterofractum, Buek, in DC. Prod. Index, iii. 93, sphalm. = retrofractum.
hirsutum, *Jacq. Hort. Schoenb.* iii. 41. t. 324.—Hab.?
hirtellum, *Schlecht. in Linnaea*, vii. (1832) 62.—Mexic.
hirtum, Sieber, ex Sendt. in Mart. Fl. Bras. x. 210 = Poeppigii.
hirtum, *Sw. Prod. Veg. Ind. Occ.* 49.—Jamaic.
intermedium, *Sendt. in Mart. Fl. Bras.* x. 221.—Bras.
involucratum, *Dun. in DC. Prod.* xiii. 1. 610.—Mexic.
jamaicense, Lam. Encyc. i. 687 = C. Parqui, vespertinum.
jasminiflorum, *Moric. ex Dun. in DC. Prod.* xiii. 1. 660.—Hab.?
Kohauti, Bercht. & Presl, Rostl. i. Solaneae. 52 = cauliflorum.
laevigatum, *Schlecht. in Linnaea*, vii. (1832) 59.—Bras.
Lamberti, *Dun. in DC. Prod.* xiii. 1. 624.—Hab.?
lanatum, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) (reimpr. 18).—Mexic.
lanceolatum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 159.—Bras.
lanceolatum, *Schott, ex Sendt. in Mart. Fl. Bras.* x. 214.—Bras.
lancifolium, *Schlecht. in Linnaea*, xix. (1847) 267.—Mexic.
lanuginosum, *Ruiz & Pav. Fl. Per.* ii. 30. t. 157.—Peruv.
lasianthum, *Dun. in DC. Prod.* xiii. 1. 649.—Peruv.
latifolium, *Lam. Illustr.* ii. 5.—Ins. Trinitat.
latifolium, Sieber, ex Dun. in DC. Prod. xiii. 1. 636 = chloranthum.
laurifolium, *L'Hérit. Stirp. Nov.* 69. t. 34.—Am. austr.
laxiflorum, *Dun. in DC. Prod.* xiii. 1. 655.—Venezuela.
laxum, *Benth. Pl. Hartw.* 33. n. 207.—Mexic.
leptanthum, *Dun. in DC. Prod.* xiii. 1. 666.—Peruv.
leucocarpum, *Dun. l. c.* 630.—Am. austr.

CESTRUM:—
Lindeni, *Dun. in DC. Prod.* xiii. 1. 611.—Venezuela.
longiflorum, *Ruiz & Pav. Fl. Per.* ii. 28. t. 154.—Peruv.
longifolium, Poepp. ex Dun. in DC. Prod. xiii. 1. 657 = unibracteatum.
Lorentzianum, *Griseb. in Goett. Abh.* xix. (1874) 217.—Am. austr.
lucidum, Pav. ex Dun. in DC. Prod. xiii. 1. 665 = servatum.
Lundianum, *Dun. l. c.* 658.—Bras.
luridum, *Dun. l. c.* 659.—Bras.
lycioides, Lichtst. ex Roem. & Schult. Syst. iv. 558 = Lycium cinereum.
lycioides, *Sendt. in Mart. Fl. Bras.* x. 208.—Bras.
macrophyllum, Hort. Par. ex Dun. in DC. Prod. xiii. 1. 626 = laurifolium.
macrophyllum, Salzm. ex Dun. l. c. = Salzmanni.
macrophyllum, *Vent. Choix* 18.—Ind. occ.
mariquitense, *H. B. & K. Nov. Gen. et Sp.* iii. 57.—N. Granat.
Martii, *Sendt. in Mart. Fl. Bras.* x. 221.—Bras.
Mathewsii, *Dun. in DC. Prod.* xiii. 1. 637.—Peruv.
megalophyllum, Dun. l. c. 638 = macrophyllum.
melanochloranthum, *Dun. l. c.* 622.—N. Granat.
microphyllum, *Linden, ex Dun. l. c.* 623.—N. Granat.
Miersianum, *Wedd. Chlor. And.* ii. 97.—N. Granat.
Miersii, *Dun. in DC. Prod.* xiii. 1. 671.—Bras.
montanum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 161.—Am. austr.
Moquinianum, *Dun. in DC. Prod.* xiii. 1. 646.—Cuba.
Moritzi, *Dun. l. c.* 619.—N. Granat.
mucronatum, *Miers, in Hook. Lond. Journ. Bot.* v. (1846) 160.—Bras.
multiflorum, *Roem. & Schult. Syst.* iv. 553.—Am. austr.
multiflorum, Schlecht. in Linnaea, vii. (1832) 59 = Schlechtendalii.
multiflorum, Schott, ex Sendt. in Mart. Fl. Bras. x. 216 = laevigatum.
multinervium, Dun. in DC. Prod. xiii. 1. 611 = C. Parqui?
Mutisii, *Willd. ex Roem. & Schult. Syst.* iv. 807.—Am. austr.
nemanthum, *Dun. in DC. Prod.* xiii. 1. 637.—Bras.
nervosum, Mill. Gard. Dict. ed. VIII. n. 3 = Tabernaemontana amygdalaefolia.
nitidum, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) (reimpr. 19).—Mexic.
nocturnum, Lam. Encyc. i. 687 = leucocarpum.
nocturnum, *Linn. Sp. Pl.* 191.—Am. austr.
oblongifolium, *Schlecht. in Linnaea*, xix. (1847) 266.—Mexic.
obovatum, *Sendt. in Mart. Fl. Bras.* x. 209.—Bras.
odontospermum, *Jacq. Hort. Schoenb.* iii. 44. t. 331.—Am. austr.
oliganthum, *Dun. in DC. Prod.* xiii. 1. 634.—Guiana.
oppositifolium, Lam. Illustr. ii. t. 112. f. 2. n. 2279 = Acokanthera Lamarkii.
organense, Miers, in Hook. Lond. Journ. Bot. v. (1846) 158 = amictum.
ovatum, *Willd. ex Roem. & Schult. Syst.* iv. 807 = tenuiflorum.
pallidum, *Lam. Encyc.* i. 688.—Jamaic.
paniculatum, *H. B. & K. Nov. Gen. et Sp.* iii. 62.—Venezuela.
paniculatum, Schlecht. in Linnaea, xix. (1847) 262 = sylvaticum.
Parqui, Benth. Pl. Hartw. 24 = terminale.
Parqui, *L'Herit. Stirp. Nov.* 73.—Am. trop.
parviflorum, *Dun. in DC. Prod.* xiii. 1. 624.—Peruv.
parvifolium, *Willd. ex Roem. & Schult. Syst.* iv. 808.—Hab.?
pauciflorum, *Nees & Mart. in Nov. Act. Acad. Caes. Leop.* xi. (1823) 79.—Bras.
pauciflorum, Willd. ex Roem. & Schult. Syst. iv. 808 = Willdenowii.
pedicellatum, *Sendt. in Mart. Fl. Bras.* x. 224.—Bras.
pedunculare, *Pav. ex Dun. in DC. Prod.* xiii. 1. 618.—Mexic.
pendulinum, Hort. Monsp. ex Dun. in DC. Prod. xiii. 1. 630 = undulatum.
pendulinum, *Jacq. Hort. Schoenb.* iii. 42. t. 327.—N. Granat.
persicaefolium, *Dun. in DC. Prod.* xiii. 1. 668.—Mexic.
peruvianum, Hort. Roth. ex Dun. l. c. 662 = C. Amelanchier.

CESTRUM:—
peruvianum, Willd. ex Roem. & Schult. Syst. iv. 807 = affine.
petiolare, H. B. & K. Nov. Gen. et Sp. iii. 58 = Sessea petiolaris.
Plukenetii, *Tausch, in Flora*, xix. (1836) 387.—Hab.?
Poeppigii, *Schlecht. in Linnaea*, xxvii. (1854) 510.—Ins. Martinic.
Poeppigii, *Sendt. in Mart. Fl. Bras.* x. 210.—Bras.
polyanthum, *Sendt. in Mart. l. c.* vi. 216.—Bras.
polyanthum, Steud. Nom. ed. II. i. 337 = Schlechtendalii.
porphyreum, *Dun. in DC. Prod.* xii. 1. 602.—Hab.?
porphyroneuron, *Schlecht. in Linnaea*, xxv. (1852) 199.—Hab.?
potaliaefolium, *Dun. in DC. Prod.* xiii. 1. 638.—N. Granat.
Prieurei, *Dun. l. c.* 635.—Guiana.
propinquum, *Mart. & Gal. in Bull. Acad. Brux.* xii. 1. (1845) 147.—Mexic.
Pseudo-Quina, *Mart. in Flora*, xxi. (1838) *Beibl.* 21.—Bras.
psychotrifolium, *Schlecht. in Linnaea*, xix. (1847) 265.—Mexic.
pubens, *Griseb. in Goett. Abh.* xix. (1874) 217.—Reg. Argent.
pubescens, Lichst. ex Roem. & Schult. Syst. iv. 558 = Acokanthera pubescens.
racemosum, *Ruiz & Pav. Fl. Per.* ii. 29. t. 154.—Peruv.
racemosum, Tausch, in Flora, xix. (1836) 387 = dubium.
ramosissimum, *Nocca, ex Dun. in DC. Prod.* xiii. 1. 673, nomen.—Hab.?
reflexum, *Sendt. in Mart. Fl. Bras.* x. 218.—Bras.
Regelii, *Planch. in Fl. des Serres*, Sér. I. ix. (1853-54) 229. t. 946.—Guatimal.
Requieni, *Dun. in DC. Prod.* xiii. 1. 661.—Hab.?
reticulatum, Willd. ex Roem. & Schult. Syst. iv. 808 = conglomeratum.
retrofractum, *Dun. in DC. Prod.* xiii. 1. 655.—Bras.
roseum, Hort. ex Dun. l. c. 602 = porphyreum.
roseum, *H. B. & K. Nov. Gen. et Sp.* iii. 59. t. 197.—Mexic.
rugulosum, Schott, ex Sendt. in Mart. Fl. Bras. x. 215 = lanceolatum.
salicifolium, Hort. Monsp. ex Dun. in DC. Prod. xiii. 1. 616 = foetidissimum.
salicifolium, *Jacq. Hort. Schoenb.* iii. 421.—Venezuela.
salicifolium, Kunth, ex Spreng. Syst. i. 673 = C. Parqui.
Salzmanni, *Dun. in DC. Prod.* xiii. 1. 626.—Bras.
scandens, Thib. ex Dun. l. c. 631 = nocturnum.
scandens, *Vahl, Eclog. Am.* i. 24.—Am. austr.
Schlechtendalii, *G. Don, Gen. Syst.* iv. 482.—Bras.
Schottii, Pohl, ex Sendt. in Mart. Fl. Bras. x. 218 = amictum.
Schottii, *Sendt. l. c.* 213.—Bras.
Sellowianum, *Sendt. l. c.* 219.—Reg. Argent.
semivestitum, *Dun. in DC. Prod.* xiii. 1. 651.—Mexic.
Sendtnerianum, *Mart. ex Sendt. in Mart. Fl. Bras.* x. 215.—Bras.
servatum, *Dun. in DC. Prod.* xiii. 1. 665.—Peruv.
sessiliflorum, *Schott, ex Sendt. in Mart. Fl. Bras.* x. 213.—Bras.
spermacocifolium, Willd. ex Roem. & Schult. Syst. iv. 808 = Bouvardia triflora.
spicatum, *Mill. Gard. Dict.* ed. VIII. n. 4.—N. Granat.
spurium, Hort. Genev. ex Dun. in DC. Prod. xiii. 1. 617 = C. Parqui.
stipulatum, Vell. Fl. Flum. 101; iii. t. 5 = bracteatum.
strictum, *Schott, ex Sendt. in Mart. Fl. Bras.* x. 208.—Bras.
strigillatum, *Ruiz & Pav. Fl. Per.* ii. 29. t. 156.—Peruv.
suberosum, *Jacq. Hort. Schoenb.* iv. 26. t. 452.—Hab.?
subpulverulentum, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 65.—Bras.
subsessile, *Vell. Fl. Flum.* 102; iii. t. 8.—Bras.
subtriflorum, *Dun. in DC. Prod.* xiii. 1. 623.—Ins. Trinitat.
subuniflorum, *Dun. l. c.* 654.—Bras.
sylvaticum, *Dun. l. c.* 603.—Mexic.
tenuiflorum, *H. B. & K. Nov. Gen. et Sp.* iii. 61.—Venezuela.
terminale, *Dun. in DC. Prod.* xiii. 1. 608.—Mexic.

CESTRUM:—
thrysoideum, *H. B. & K. Nov. Gen. et Sp.* iii. 63.—Mexic.
tinctorium, *Jacq. Hort. Schoenb.* iii. 45. t. 332.—Am. austr.
Tinus, *Dun. in DC. Prod.* xiii. 1. 614.—Bras.
tomentosum, *Linn. f. Suppl.* 150.—Am. austr.
tomentosum, Moç. & Sessé, ex Dun. in DC. Prod. xiii. 1. 601 = Benthami.
toxicarium, J. F. Gmel. Syst. 389, 1524 = Antiaris toxicaria.
tubulosum, *Sendt. in Mart. Fl. Bras.* x. 207.—Bras.
umbellatum, E. Mey. ex Meissn. in DC. Prod. xiv. 528 = Peddiea Dregei.
umbellatum, *Pauq. Diss. Bellad.* 16.—Hab.?
undulatum, *Ruiz & Pav. Fl. Per.* ii. 28. t. 155.—Peruv.
unibracteatum, *Dun. in DC. Prod.* xiii. 1. 656.—Peruv.
velutinum, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877-78) 63.—Am. trop.
venatorium, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 79 = Acokanthera venenata.
venenatum, Burm. f. Fl. Cap. Prod. 5 = Acokanthera venenata.
venenatum, Lam. Encyc. i. 688 = laurifolium.
venenatum, Mill. Gard. Dict. ed. VIII. n. 6 = pallidum.
venosum, Willd. ex Roem. & Schult. Syst. iv. 807 = Sessea petiolaris.
vespertinum, Bertero, ex Dun. in DC. Prod. xiii. 1. 660 = depauperatum.
vespertinum, Hort. Val. ex Roem. & Schult. Syst. iv. 366 = Vestia lycioides.
vespertinum, *Linn. Mant.* ii. 206.—Ind. occ.
vespertinum, Poepp. ex Sendt. in Mart. Fl. Bras. x. 218 = Poeppigii.
vestioides, *Schlecht. in Linnaea*, vii. (1832) 65.—Bras.
vestitum, Hook. Ic. Pl. t. 381 = Sessea vestita.
viminale, *Sendt. in Mart. Fl. Bras.* x. 220.—Bras.
virgatum, Ruiz & Pav. Fl. Per. ii. 27 = C. Parqui.
viride, *Moric. ex Dun. in DC. Prod.* xiii. 1. 606.—Mexic.
viridiflorum, *Hook. Bot. Mag.* t. 4022.—Bras.
Warscewiczii, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xix. (1851) 362.—Am. austr.
Willdenowii, *Steud. Nom.* ed. II. i. 338.—Hab.?

CETRA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 (Quid?).

CEVALLIA, Lag. in Varied. Cienc. ii. IV. (1805) 35. *LOASEAE*, Benth. & Hook. f. i. 803.
PETALANTHERA, Nutt. in Journ. Acad. Philad. vii. (1834) 107.
sinuata, *Lag. l. c.* 36.—N. Mexic.

CEYTOSIS, Munro, in Journ. Linn. Soc. vi. (1862) 54, err. typ. = **Crypsis**, Ait. (Gramin.).

CHABOISSAEA, Fourn. Gram. Mex. 112; ex Benth. & Hook. f. Gen. iii. 1096 (1883). Gen. dub. *GRAMIN.* Benth. & Hook. f. l. c.
ligulata, *Fourn. l. c.*; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 588.—Mexic.

CHABRAEA, Adans. Fam. ii. 237 (1763) = **Peplis**, Linn. (Lythrar.).

CHABRAEA, DC. in Ann. Mus. Par. xix. (1812) 65. t. 5 = **Leuceria**, Lag. (Compos.).
abbreviata, Bert. 'Merc. Chil. (1829) 601.'—Cf. Bull. Férussac, xx. (1830) 109 = L. abbreviata.
Barrasiana, Remy, in C. Gay, Fl. Chil. iii. 399.—Chili.
Berteroniana, Steud. Nom. ed. II. i. 338.—Chili.
candidissima, DC. Prod. vii. 59 = L. candidissima.
canescens, Phil. in Linnaea, xxviii. (1856) 716.—Chili.
cinerea, DC. Prod. vii. 60 = L. cinerea.
concinna, Phil. in Linnaea, xxxiii. (1864-65) 123.—Chili.
coquimbana, Phil. in Anal. Univ. Chil. (1861) 50.—Chili.
daucifolia, Wedd. Chlor. And. i. 35.—Peruv.
elongata, Bert. 'Merc. Chil. (1829) 601.'—Cf. Bull. Férussac, xx. (1830) 109 = L. senecioides.
fragrans, Phil. in Linnaea, xxix. (1857-58) 3.—Chili.

CHABRAEA:—
Gayana, Remy, in C. Gay, Fl. Chil. iii. 400.—Chili.
glabra, DC. in Deless. Ic. Select. iv. t. 89.—Chili.
glabriuscula, Phil. in Linnaea, xxviii. (1856) 715.—Chili.
glacialis, DC. in Deless. Ic. Select. iv. t. 91 = L. pulchella.
glandulosa, DC. Prod. vii. 59 = L. glandulosa, *D. Don.*
integrifolia, Phil. in Linnaea, xxviii. (1856) 716.—Chili.
laciniata, Wedd. Chlor. And. i. 34. t. 10 = L. laciniata.
Landbeckii, Phil. in Linnaea, xxxiii. (1864-65) 122.—Chili.
lithospermifolia, DC. Prod. vii. 60.—Chili.
lyrata, Spreng. Syst. iii. 506 = Gerbera Anandria.
modesta, Phil. Fl. Atac. 28.—Chili.
multifida, DC. Prod. vii. 60.—Patagonia.
nutans, Bland. C. Gay, Fl. Chil. viii. 397.—Chili.
nutans, Remy, in Ann. Sc. Nat. Sér. III. xii. (1849) 182.—Chili.
oligocephala, Phil. in Anal. Univ. Chil. (1872) 745 = L. oligocephala.
Poeppigii, Phil. in Linnaea, xxix. (1857-58) 3.—Chili.
polyclados, Remy, in C. Gay, Fl. Chil. iii. 399.—Chili.
prenanthoides, Bert. in 'Merc. Chil. 1829.'—Cf. Bull. Férussac, xx. (1830) 109.—Chili.
pulchella, Phil. in Anal. Univ. Chil. (1872) 744 = L. pulchella.
purpurea, DC. in Ann. Mus. Par. xix. (1812) 65, 71 = L. purpurea.
rosea, DC. in Deless. Ic. Select. iv. t. 90 = L. runcinata.
runcinata, Hook. Bot. Mag. t. 4116 = L. runcinata.
salina, Remy, in C. Gay, Fl. Chil. iii. 393. t. 41.—Chili.
scrobiculata, DC. Prod. vii. 59 = L. scrobiculata.
suaveolens, DC. l. c. = L. gossypina.
tenerifolia, Phil. in Linnaea, xxviii. (1856) 715.—Chili.
tenuior, Bert. in 'Merc. Chil. 1829.'—Cf. Bull. Férussac, xx. (1830) 109 = *Berteroniana.*
tenuisecta, Sch. Bip. in Bonplandia, iv. (1856) 55, nomen.—Chili.
thermarum, Phil. in Anal. Univ. Chil. (1862) 380.—Chili.
tomentosa, DC. Prod. vii. 60 = L. Bridgesii.
viscida, Bert. in 'Merc. Chil. (1829) 601.'—Cf. Bull. Férussac, xx. (1830) 109.—Chili.
viscosa, Bert. ex DC. Prod. vii. 59 = L. runcinata.

CHADARA, Forsk. Fl. Aegypt. Arab. 105 (1775) = **Grewia**, Linn. (Tiliac.).
arborea, Forsk. l. c. = G. excelsa.
tenax, Forsk. l. c. = G. populifolia.
velutina, Forsk. l. c. = G. velutina.

CHADRA, T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 11, err. typ. = praec.

CHADSIA, Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 104. *LEGUMINOSAE*, Benth. & Hook. f. i. 497.
andravinensis, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 391.—Madag.
coluteifolia, *Baill. l. c.* 392.—Madag.
flammea, *Boj. in Ann. Sc. Nat.* Sér. II. xx. (1843) 106.—Madag.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1883) 391.—Madag.
granitica, *Baill. l. c.* 392.—Madag.
Lantziana, *Baill. l. c.* 391.—Madag.
salicina, *Baill. l. c.* 392.—Madag.
versicolor, *Boj. in Ann. Sc. Nat.* Sér. II. xx. (1843) 106.—Madag.

CHAELANTHUS, Poir. in Dict. Sc. Nat. viii. 50 (1817) sphalm. = **Chaetanthus**, R. Br. (Restiac.).

CHAELOTHILUS, Neck. Elem. ii. 14 (1790) = **Gentiana**, Tourn.

CHAENACTIS, DC. Prod. v. 659 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 401.
ACARPHAEA, Harv. & A. Gray, Pl. Fendl. 98 (1849).
ACICARPHAEA, Walp. Ann. ii. 877 (1852).
MACROCARPHUS, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 376.
achilleaefolia, Hook. & Arn. Bot. Beech. Voy. 354 = Douglasii.

CHAENACTIS:—
artemisiaefolia, *A. Gray, in Proc. Am. Acad.* x. (1875) 74.—Calif.
attenuata, *A. Gray, l. c.* 73.—Arizona.
brachypappa, *A. Gray, l. c.* viii. (1872) 390.—Am. bor. occ.
carphoclinia, *A. Gray, in Bot. Mex. Bound.* 94.—Am. bor. occ.
Cusickii, *A. Gray, Syn. Fl. N. Am.* ii. II. 452.—Am. bor. occ.
denudata, Nutt. in Journ. Acad. Sc. Philad. Ser. II. i. (1847) 177 = glabriuscula.
Douglasii, *Hook. & Arn. Bot. Beech. Voy.* 354.—Am. bor. occ.
filifolia, Harv. in A. Gray, Pl. Fendl. 98 = tenuifolia.
Fremonti, *A. Gray, in Proc. Am. Acad.* xix. (1883) 30.—Calif.
glabriuscula, *DC. Prod.* v. 659.—Calif.
heterocarpha, *A. Gray, Pl. Fendl.* 98.—Calif.
lanosa, *DC. Prod.* v. 659.—Calif.
macrantha, *Eaton, in Bot. King, Exp.* 171. t. 18.—Am. bor. occ.
nevadensis, *A. Gray, in Bot. Calif.* i. 391.—Calif.
Nevii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 30.—Am. bor. occ.
Parishii, *A. Gray, l. c.* xx. (1885) 299.—Calif.
santolinoides, *Greene, in Bull. Torrey Club*, ix. (1882) 17.—Calif.
stevioides, *Hook. & Arn. Bot. Beech. Voy.* 353.—Am. bor. occ.
suffrutescens, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 100.—Calif.
tanacetifolia, A. Gray, l. c. vi. (1864–65) 545 = heterocarpha.
tenuifolia, *Nutt. in Trans. Am. Phil. Soc.* N. S. vii. (1841) 375.—Calif.
thysanocarpha, *A. Gray, in Proc. Am. Acad.* xix. (1883) 30.—Calif.
xantiana, *A. Gray, l. c.* vi. (1865) 545.—Calif.

CHAENANTHE, Lindl. Bot. Reg. (1838) Misc. 38 = **Diadenium**, Poepp. & Endl. (Orchid.).
Barkeri, Lindl. l. c. = D. Barkeri.

CHAENANTHERA, Rich. ex DC. Prod. iii. 196 (1828) = **Charianthus**, D. Don (Melastomac.).
mucronata, Rich. l. c. 197 = Charianthus glaberrimus.

CHAENARRHINUM, Reichb. Consp. 123 (1828) = **Linaria**, Tonrn. (Scrophul.).
crassifolium, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 579 = L. origanifolia.
exile, Lange, in Kjoeb. Vidensk. Meddel. (1877–78) 232 = L. exilis.
flexuosum, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 580 = L. Langei.
macrocalyx, Pomel, Nouv. Mat. Fl. Atl. 98 = L. macrocalyx.
macropodum, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 579 = L. macropoda.
minus, Lange, l. c. 577 = L. minor.
origanifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 127 = L. origanifolia.
praetermissum, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 578 = L. minor.
rubriflorum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 127 = L. rubrifolia.
rubrifolium, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 578 = L. rubrifolia.
serpyllifolium, Lange, l. c. = L. serpyllifolium.
tenellum, Lange, l. c. 581 = Anarrhinum tenellum.
thymiflorum, Losc. ex Lange, in Kjoeb. Vidensk. Meddel. (1877–78) 232 = L. exilis.
villosum, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 580 = L. villosa.

CHAENESTHES, Micrs, in Hook. Lond. Journ. Bot. iv. (1845) 336 = **Iochroma**, Benth. (Solanac.).
cornifolia, Micrs, l. c. 338 = I. cornifolia.
fuchsioides, Micrs, l. c. 337 = I. fuchsioides.
gesnerioides, Micrs, l. c. 338 = I. gesnerioides.
lanceolata, Miers, l. c. = I. lanceolata.
longipes, Dun. in DC. Prod. xiii. 1. 489 = I. longipes.
loxensis, Micrs, in Hook. Lond. Journ. Bot. iv. (1848) 338 = I. loxensis.
umbrosa, Miers, l. c. 337 = I. umbrosa.

CHAENOCARPUS, Juss. in Dict. Sc. Nat. viii. 50 (1817) = CHENOCARPUS, Neck. = **Spermacoce**, Linn. (Rubiac.).

CHAENOCEPHALUS, Griseb. Fl. Brit. W. Ind. 374 (1861). *COMPOSITAE*, Benth. & Hook. f. ii. 382.
heterophyllus, *Griseb. in Goett. Abh.* xxiv. (1879) 196.—Reg. Argent.
macrophyllus, *Griseb. l. c.*—Reg. Argent.
petrobioides, *Griseb. Fl. Brit. W. Ind.* 374.—Ind. occ.
Suncho, *Griseb. in Goett. Abh.* xxiv. (1879) 196.—Reg. Argent.

CHAENOLOBIUM, Miq. Fl. Ind. Bat. Suppl. i. 302 (1860) = **Ormosia**, Jack (Legumin.).
decemjugum, Miq. l. c. = O. microsperma.
septemjugum, Miq. l. c. = O. microsperma.

CHAENOMELES, Lindl. Veg. Kingd. 560 (1847) = CHOENOMELES, Lindl. = **Pyrus**, Linn. (Rosac.).

CHAENOPHORA, Rich. ex Crueger, in Linnaca, xx. (1847) 112, err. typ. = seq.

CHAENOPLEURA, Rich. ex DC. Prod. iii. 197 (1828) = **Miconia**, Ruiz & Pav. (Melastomac.).
densiflora, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 349 = M. pusilliflora.
ferruginea, H. Crueger, in Linnaea, xx. (1847) 112 = M. hypoleuca.
fulva, Griseb. in Goett. Abh. ix. (1861) 55 = M. fulva.
hypoleuca, Benth. in Hook. Journ. Bot. ii. (1840) 315 = M. hypoleuca.
lanceolata, Gardn. in Hook. Lond. Jonrn. Bot. ii. (1843) 349 = M. pnsilliflora.
longifolia, Griseb. in Goett. Abh. ix. (1861) 55 = M. longifolia.
parviflora, Gardn. in Hook. Lond. Journ. Bot. ii (1843) 348 = M. pusilliflora.
quadrangularis, Macf. Fl. Jamaic. ii. 96 = M. quadrangularis.
stelligera, Macf. l. c. 97.—Jamaic.
stenobotrys, DC. Prod. iii. 197 = M. quadrangularis.

CHAENORRHINUM, Lange, in Willk. & Lange, Prod. Fl. Hisp. ii. 577 (1870), err. typ. = CHAENARRHINUM, Reichb. = **Linaria**, Tourn. (Scrophul.).

CHAENOSTOMA, Benth. in Hook. Comp. Bot. Mag. i. (1835) 374. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 945.
PALMSTRUCKIA, Retz. f. Obs. Bot. Pugill. 15 (1810).
SUTERA, Roth, Bot. Bemerk. 172 (1878).
aethiopicum, *Benth. l. c.* 375.—Afr. austr.
affine, *Bernh. in Flora*, xxvii. (1844) 834.—Afr. austr.
calycinum, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 374.—Afr. austr.
campanulatum, *Benth. l. c.*—Afr. austr.
cordatum, *Benth. l. c.* 377.—Afr. austr.
cuneatum, *Benth. l. c.* 376.—Afr. austr.
denudatum, *Benth. l. c.* 375.—Afr. austr.
fastigiatum, *Benth. l. c.* 376.—Afr. austr.
floribundum, *Benth. l. c.*—Afr. austr.
foetidum, *Benth. l. c.* 377.—Afr. austr.
glabratum, *Benth. l. c.* 375.—Afr. austr.
halimifolium, *Benth. l. c.*—Afr. austr.
hispidum, *Benth. l. c.* 376.—Afr. austr.
integrifolium, *Benth. l. c.*—Afr. austr.
Kraussianum, Bernh. in Flora, xxvii. (1844) 835 = Lyperia Kraussiana.
laxiflorum, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 374.—Afr. austr.
linifolium, *Benth. l. c.* 375.—Afr. austr.
marifolium, *Benth. l. c.* 376.—Afr. austr.
natalensis, *Bernh. in Flora*, xxvii. (1844) 835.—Afr. austr.
pauciflorum, *Benth. in Hook. Comp. Bot. Mag.* i. (1835) 374.—Afr. austr.
pedunculosum, *Benth. l. c.* 377.—Afr. austr.
polyanthum, *Benth. l. c.* 375.—Afr. anstr.
procumbens, *Benth. l. c.* 374.—Afr. austr.
pumilum, *Benth. l. c.* 375.—Afr. austr.
racemosum, *Benth. l. c.* 377.—Afr. austr.
revolutum, *Benth. l. c.* 375.—Afr. austr.
rotundifolium, *Benth. l. c.* 374.—Afr. austr.
subspicatum, *Benth. l. c.* 376.—Afr. austr.
villosum, [G. Don, in] Sweet, Hort. Brit. ed. III. 520 = Polycarena capensis.

CHAERADOPLECTRON, Benth. & Hook. f. Gen. iii. 626 (1883), err. typ. = CHOERADOPLECTRON, Schau. = **Habenaria**, Willd. (Orchid.).

CHAEROPHYLLUM, Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 898.
BALANSAEA, Boiss. & Reut. Pugill. Pl. Nov. 49 (1852).
BIASOLETTIA, Bertol. Fl. Ital. iii. 191 (1837).
FIEBERA, Opiz, Seznam, 44 (1852).
FREYERA, Reichb. Handb. 291 (1837).
GRAMMOSCIADIUM, DC. Prod. iv. 232 (1830).
PHYSOCAULIS, Tausch, in Flora, xvii. (1834) 342.
RHYNCHOSTYLIS, Tansch, l. c. 343 (1834).
acuminatum, *Lindl. in Royle, Illustr. Bot. Himal.* 232.—Reg. Himal.
affine, Steud. in A. Rich. Tent. Fl. Abyss. i. 332 = Anthriscus sylvestris.
alpestre, Jord. Pugill. 75 = Villarsii.
alpinum, Vill. Prosp. 26 = Anthriscus sylvestris.
angelicaefolium, C. A. Mey. Verz. Pfl. Cauc. 130 = Meycri.
angelicaefolium, *Bieb. Fl. Taur. Cauc.* iii. 240.—Oriens; Iberia.
angelicaefolium, DC. Prod. iv. 227 = byzantinum.
angulatum, Kit. ex Spreng. in Comm. Gott. Rec. ii. (1813) Umbell. Animad. 7. t. 1. f. 6 = Anthriscus sylvestris.
Anthriscus, Lam. Encyc. i. 685 = Anthriscus vulgaris.
arborescens, Linn. Sp. Pl. 259 = Aralia spinosa?
aristatum, Thunb. Fl. Jap. 119 = Osmorhiza Claytoni.
aromaticum, *Linn. Sp. Pl.* 259.—Europ.
articulatum, Bosc, ex DC. Prod. iv. 225 = procumbens.
Astrantiae, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 906.—As. Min.
aurenm, *Linn. Sp. Pl.* ed. II. 370.—Europ.; As. Min.
aureum, Linn. Mant. ii. 356 = temulum.
australe, Crantz, Cl. Umb. Emcnd. 76 = Scandix australis.
Biebersteinii, *Lag. ex Sweet, Hort. Brit.* ed. I. 190.—Reg. Caucas.
bifidum, Willd. ex DC. Prod. iv. 225 = procumbens.
Boscii, Steud. Nom. ed. II. i. 339 = procnmbens.
brachycarpum, Bieb. ex Hoffm. Gen. Umb. i. 212 = millefolium.
bulbosum, *Linn. Sp. Pl.* 258.—Europ.; As. Min.
byzantinum, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 65.—Byzant.; As. Min.
cachemiricum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 691.—Reg. Himal.
cadonense, Schult. ex Stend. Nom. ed. II. i. 339 = Anthriscus sylvestris.
calabricum, *Guss. ex DC. Prod.* iv. 227.—Calabria.
californicum, Torr. in Pacif. Rail. Rep. iv. 93 = Podosciadium californicum.
canadense, Pers. Syn. i. 320 = Cryptotaenia canadensis.
capense, Thunb. Prod. Pl. Cap. 51 = Annesorhiza capensis.
capnoides, *Benth. & Hook. f. Gen.* i. 898.—Reg. Himal.
Cerefolium, Crantz, Stirp. Austr. ed. I. fasc. iii. 70 = Anthriscus Cerefolium.
chilense, Poir. Encyc. v. 145 = Scandix chilensis.
Cicutaria, Vill. Prosp. 26 = hirsntum.
ciliatum, *Kit. ex Kanitz, Reliq. Kitaib.* 103.—Croatia.
Claytoni, Pers. Syn. i. 320 = Osmorrhiza brevistylis.
coloratnm, *Linn. Mant.* 57.—Dalmat.; Banat.
creticum, *Boiss. & Heldr. Diagn.* Ser. I. x. 51.—Ins. Creta.
crinitum, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 63.—Mesopot.; Persia.
Cyminum, Fisch. ex DC. Prod. iv. 230 = Conopodium Cyminum.
dasycarpum, Nutt. ex Torr & Gray, Fl. N. Am. i. 638 = Tainturicir.
daucifolium, Desf. Cat. Hort. Par. ed. III. 200, 405 = Anthriscns alpestris.
daucophyllum, Nutt. ex Torr. & Gray, Fl. N. Am. i. 638 = Tainturieri.
divaricatum, Poir. Encyc. Suppl. iv. 34 = Anthriscus fumarioides.
dulce, Fisch. ex Steud. Nom. ed. II. i. 339 = Osmorrhiza Claytoni.
elegans, *Gaud. Fl. Helv.* ii. 364.—Helvet.
Friedrichsstahlii. Cesati. in Bibl. Ital. lxxv. 388 = Anthriscus tenerrima.
geniculatum, Gilib. Fl. Lituan. ii. 29 = temulum.

CHAEROPHYLLUM:—
glaberrimum, Pers. Syn. i. 320 (= *Bunium glaberrimum*).—Algeria.
gracile, Bess. Hort. Cremen. (1822) = Conopodium Cyminum.
gracillimum, Klotzsch, in Bot. Ergeb. Waldem. Reise, 149 = Vicatia coniifolia.
grandiflorum, Crantz, Cl. Umb. Emend. 77; Lam. Encyc. i. 686 = Scandix grandiflora.
Heldreichii, *Orph. in Boiss. Diagn.* Ser. II. ii. 104.—Graecia.
hirsutum, *Linn. Sp. Pl.* 258.—Europ.; As. Min.
hirsutum, Pall. Tabl. Taur. 49 = temulum.
hirsutum, Vill. ex Koch, Syn. Fl. Germ. ed. I. 317 = Villarsii.
hispidum, Thunb. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 64 = Caucalis Anthriscus.
Hladnikianum, Reichb. ex Heynh. Nom. ii. 135 = Anthriscus fumarioides.
humile, *Bieb. Fl. Taur. Cauc.* iii. 240.—Reg. Caucas.
hybridum, Tenore, Fl. Neap. Prod. p. lxvi = aureum.
infestum, Salisb. Prod. 167 = Anthriscus sylvestris.
Kotschyi, Fenzl, ex Boiss. Fl. Orient. ii. 913 = Anthriscus Kotschyi.
lactescens, Rochel, ex Steud. Nom. ed. II. i. 339 = Anthriscus sylvestris.
laevigatum, Vis. Fl. Dalm. iii. 65 = bulbosum.
lasiolaenum, *Boiss. & Bal. ex Boiss. Fl. Orient.* ii. 903.—As. Min.
leucolaenum, *Boiss. l. c.* 905.—Armenia.
libanoticum, *Boiss. & Kotschy, Diagn.* Ser. II. vi. 89.—Syria.
lucidum, Desf. Cat. Hort. Par. ed. III. 200 = Anthriscus nemorosa.
macropodum, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 64.—Persia.
macrospermum, *Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc.* (1838) 327.—Reg. Caucas.
maculatum, Willd. Enum. Hort. Berol. Suppl. 15 = aureum.
magellense, *Tenore, Prod. Fl. Neap. App.* iv. 15.—Neapol.
Meyeri, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 103.—Persia.
millefolium, *DC. Prod.* iv. 226.—Reg. Caucas.
millefolium, Klotzsch, in Bot. Ergeb. Waldem. Reise, 149 = Vicatia millefolia.
minimum, *Vand. Fasc. Pl.* 11.—Italia.
monogonum, Kit. ex Link, Enum. Hort. Berol. i. 281 = aureum.
neglectum, *Zinger, in Bull. Soc. Nat. Mosc.* lvi. (1881) i. 313.—Rossia.
nemorosum, Hoffm. Gen. Umb. 45 = Anthriscus nemorosa.
nemorosum, Lag. ex DC. Prod. iv. 224 = Anthriscus Cerefolium.
nitens, Turcz. ex Bernh. in Linnaea, xii. (1838) Litt. 76 = Anthriscus nitens.
nitidum, Wahlenb. Fl. Carp. 85 = Anthriscus alpestris.
nodosum, *Crantz, Cl. Umb. Emend.* 76; *Lam. Encyc.* i. 685.—Europ.; As. Min.; Afr. bor.
odoratum, Crantz, l. c. 75; Lam. l. c. 683; Fl. Fr. iii. 439 = Myrrhis odorata.
orientale, Willd. ex Boiss. Fl. Orient. ii. 908 = byzantinum.
orthostylum, *Trautv. in Act. Hort. Petrop.* iii. 11. (1875) 273.—Reg. Caucas.
pallescens, Presl, ex DC. Prod. iv. 227 = hirsutum.
palustre, Lam. Encyc. i. 683 = hirsutum.
Pecten-Veneris, Crantz, Stirp. Austr. ed. I. fasc. ii. 66; ed. II. 189 = Scandix Pecten-Veneris.
pinnatifidum, Poir. Encyc. v. 144 = Scandix pinnatifida.
polonicum, *Jastrzebowski, ex Rostaf. Fl. Polon. Prod.* 112.—Europ.
Prescottii, DC. Prod. iv. 225 = bulbosum.
procumbens, *Crantz, Cl. Umbell. Emend.* 77.—Am. bor.
reflexum, Aitch. in Journ. Linn. Soc. xiii. (1882) 164 = villosum.
reflexum, *Lindl. in Royle, Illustr. Bot. Himal.* 233.—Reg. Himal.
rigidum, Huet, ex Nym. Consp. 300 = Villarsii.
roseum, *Bieb. Fl. Taur. Cauc.* i. 234.—Reg. Caucas.
rostratum, Lam. Encyc. i. 685 = Scandix Pecten-Veneris.
sativum, Bieb. Fl. Taur. Cauc. i. 232 = Caucalis trichosperma.
sativum, Lam. Fl. Fr. iii. 438 = Anthriscus Cerefolium.

CHAEROPHYLLUM:—
scabrum, Thunb. Fl. Jap. 119 = Caucalis Anthriscus.
siculum, Guss. Fl. Sic. Prod. i. 352 = Anthriscus sicula.
sphallerocarpus, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 367.—Soongaria.
sylvestre, Bieb. Fl. Taur. Cauc. i. 231 = maculatum.
sylvestre, Linn. Sp. Pl. 258 = Anthriscus sylvestris.
syriacum, *Hempr. & Ehrenb. ex Boiss. Fl. Orient.* ii. 904.—Syria.
Tainturieri, *Hook. & Arn. in Hook. Comp. Bot. Mag.* i. (1835) 47.—Texas.
temulentum, Linn. Fl. Suec. 94 = temulum.
temuloides, *Boiss. in Ann. Sc. Nat.* Sér. III. ii. (1844) 64.—Persia.
temulum, *Linn. Sp. Pl.* 258.—Europ.; Afr. bor.; Reg. Caucas.
tenuifolium, *Poir. Encyc. Suppl.* iv. 342.—Marocco.
tenuifolium, Stev. in Hoffm. Gen. Umbell. ed. II. 212 = millefolium.
tomentosum, *Poir. ex Steud. Nom.* ed. II. i. 339.—Hab. ?
torquatum, DC. Fl. Fr. Suppl. 505 = Anthriscus alpestris.
trapezuntinum, Boiss. Fl. Orient. ii. 903 = temuloides.
trichospermum, Lam. Encyc. i. 685 = Caucalis trichosperma.
tuberculosum, Poir. Encyc. Suppl. iv. 342 = Anthriscus fumarioides.
tuberosum, C. A. Mey. ex Regel, in Act. Hort. Petrop. v. (1877) 587 = Carum cylindricum.
tuberosum, *Royle, Illustr. Bot. Himal.* 231.—Reg. Himal.
tumidum, Gilib. Fl. Lituan. ii. 28 = Anthriscus sylvestris.
umbrosum, Jord. Obs. Pl. Crit. vii. (1849) 30 = hirsutum.
verticillatum, Pers. Syn. i. 320 = bulbosum, Anthriscus vulgaris.
Villarsii, *Koch, Syn. Fl. Germ.* ed. I. 317.—Europ.
villosum, *Wall. Cat.* n. 558.—Reg. Himal.
villosum, Wall. l. c. partim = reflexum.

CHAETACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836) = **Calophanes**, D. Don (Acanthac.).
Burchellii, Nees, in DC. Prod. xi. 462 = Cal. Burkei.
costatus, Nees, l. c. = Calophanes costatus.
glandulosus, Nees, l. c. = Calophanes Persoonii.
Persoonii, Nees, in Linnaea, xv. (1841) 357 = Calophanes Persoonii.
repandus, A. Gray, in Proc. Am. Acad. v. (1861–62) 349.—Ins. Fiji.
setigera, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 = Calophanes Persoonii.

CHAETACHLAENA, D. Don, in Trans. Linn. Soc. xvi. (1830) 256 = **Onoseris**, DC. (Compos.).
acerifolia, D. Dietr. Syn. Pl. iv. 1565 = O. acerifolia.
annua, D. Dietr. l. c. = O. annua.
Cumingii, D. Dietr. l. c. = O. Cumingii.
odorata, D. Don, in Trans. Linn. Soc. xvi. (1830) 256 = O. odorata.

CHAETACHME, Planch. in Ann. Sc. Nat. Sér. III. x. (1848) 340. *URTICACEAE*, Benth. & Hook. f. iii. 356.
aristata, *Planch. l. c.* 341.—Afr. austr.
madagascariensis, *Baker, in Journ. Linn. Soc.* xxi. (1885) 443.—Madag.
Meyeri, Harv. Thes. Cap. i. 16 = aristata.
nitida, *Planch. & Harv. in Harv. Thes. Cap.* 16.—Afr. austr.

CHAETADELPHA, A. Gray, ex S. Wats. in Am. Natural. vii. (1873) 301; et in Proc. Am. Acad. ix. (1874) 218. *COMPOSITAE*, Benth. & Hook. f. ii. 1238.
Wheeleri, *A. Gray, ll. cc.*—Arizona.

CHAETAEA, Jacq. Enum. Pl. Carib. 2 (1760) = **Buettneria**, Loefl. (Stercul.).
aculeata, Jacq. l. c. 17 = B. carthaginensis.

CHAETAGASTRA, Crueg. in Linnaea, xx. (1849) 101, sphalm. = CHAETOGASTRA, DC. = **Tibouchina**, Aubl. (Melast.).

CHAETANTERA, Less. Syn. Comp. 111 (1832) = **Chaetanthera**, Ruiz & Pav. (Compos.).

CHAETANTHERA, Nutt. in Journ. Acad. Philad. vii. (1834) 111 = **Chaetopappa**, DC. (Compos.).
asteroides, Nutt. l. c. = Chaetopappa asteroides.

CHAETANTHERA, Ruiz & Pav. Prod. Fl. Per. 106. t. 23 (1794). *COMPOSITAE*, Benth. & Hook. f. ii. 496.
ALDUNATEA, Remy, in C. Gay, Fl. Chil. iii. 320. t. 38 (1849).
CARMELITA, C. Gay, in DC. Prod. vii. 14 (1838).
CHERINA, Cass. in Dict. Sc. Nat. viii. 437 (1817).
CHONDROCHILUS, Phil. in Linnaea, xxviii. (1856) 711.
EGANIA, Remy, in C. Gay, Fl. Chil. iii. 324 (1849).
ELACHIA, DC. Prod. vii. 256 (1838).
EUTHRIXIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 257.
MINYTHODES, Phil. ex Benth. & Hook. f. Gen. ii. 496 (1873).
ORIASTRUM, Poepp. & Endl. Nov. Gen. et Sp. iii. 50. t. 257 (1847).
PROSELIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 234.
TYLLOMA, D. Don, l. c. 238 (1830).
argentea, D. Don, in Phil. Mag. (1832) 392 = chilensis.
argentea, *Phil. in Anal. Univ. Chil.* (1872) 739.—Chili.
Berteriana, *Less. Syn. Comp.* 111.—Chili.
Berteroana, Hook. & Arn. in Comp. Bot. Mag. i. (1835) 104 = praec.
Caput-Tringae, Less. ex DC. Prod. vii. 32 = limbata.
chilensis, *DC. in Ann. Mus. Par.* xix. (1812) 70.—Chili.
chilensis, Hook. & Arn. in Bot. Beech. Voy. 29 = incana.
ciliaris, Bert. ex Steud. Nom. ed. II. i. 340 = spathulata.
ciliata, *Ruiz & Pav. Syst. Veg. Per. et Chil.* 190.—Chili.
crenata, *Remy, in C. Gay, Fl. Chil.* iii. 302.—Chili.
crispa, *Phil. in Anal. Univ. Chil.* (1872) 739.—Chili.
dealbata, *Hook. & Arn. in Comp. Bot. Mag.* i. (1835) 104.—Chili.
debilis, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 287.—Chili.
elegans, *Phil. in Linnaea*, xxviii. (1856) 712.—Chili.
eryngioides, D. Don, in Phil. Mag. (1832) 391 = tenuifolia.
flabellata, D. Don, l. c. = multicaulis.
glandulosa, *Remy, in C. Gay, Fl. Chil.* iii. 311. t. 35. f. 1.—Chili.
grandiflora, *Steud. Nom.* ed. II. i. 340.—Chili.
humilis, *Phil. in Anal. Univ. Chil.* (1872) 740.—Chili.
incana, *Poepp. ex Less. in Linnaea*, v. (1830) 284.—Chili.
Kunthiana, Less. Syn. Comp. 114 = tenella.
limbata, *Less. l. c.* 116.—Chili.
linearifolia, Poepp. ex Steud. Nom. ed. II. i. 340 = linearis.
linearis, *Poepp. ex Less. Syn. Comp.* 112.—Chili.
linifolia, *Bert. ex Less. l. c.*—Chili.
magellanica, Spreng. Syst. iii. 503 = Perezia lactucoides.
microphylla, Hook. & Arn. in Comp. Bot. Mag. i. (1835) 104 = linifolia.
moenchioides, *Less. Syn. Comp.* 113.—Chili.
multicaulis, *DC. Prod.* vi. 31.—Chili.
multiflora, Humb. & Bonpl. Pl. Aequin. ii. 168 = Perezia multiflora.
peruviana, *A. Gray, in Proc. Am. Acad.* v. (1861) 144.—Peruv.
pinnatifida, Humb. & Bonpl. Pl. Aequin. ii. 170 = Perezia pinnatifida.
prostrata, D. Don, in Phil. Mag. (1832) 391 = multicaulis.
pungens, Humb. & Bonpl. Pl. Aequin. 146. t. 127 = Perezia pungens.
pusilla, *Hook. & Arn. in Comp. Bot. Mag.* i. (1835) 106.—Chili.
ramosissima, Hook. & Arn. l. c. 104 = moenchioides, tenella.
recurvata, Spreng. Syst. iii. 503 = Perezia recurvata.
scariosa, D. Don, in Phil. Mag. (1832) 391 = incana.
sericea, Lag. ex DC. Prod. vii. 30 = chilensis.
serrata, Ruiz & Pav. Syst. Veg. Per. et Chil. 191 = chilensis.
spathulata, *Poepp. ex Less. in Linnaea*, v. (1830) 285.—Chili.
spinulosa, Cass. Opusc. ii. 103 = tenuifolia.
tenella, *Less. Syn. Comp.* 114; *DC. in Deless. Ic. Sel.* iv. t. 82.—Chili.
tenuifolia, *D. Don, in Phil. Mag.* (1832) 391.—Chili.
valdiviana, *Phil. in Linnaea*, xxviii. (1856) 712.—Chili.
villosa, *D. Don, in Phil. Mag.* (1832) 391.—Chili.
Volkmanni, *Phil. in Anal. Univ. Chil.* (1872) 740.—Chili.

CHAETANTHUS, R. Br. Prod. 251 (1810). *RESTIACEAE*, Benth. & Hook. f. iii. 1034.
PRIONOSEPALUM, Steud. Syn. Pl. Cyp. 266 (1855).
leptocarpoides, *R. Br. l. c.*—Austral.

CHAETARIA, Beauv. Agrost. 30. t. 8 (1812) = **Aristida**, Linn. (Gramin.).
Adscensionis, Roem. & Schult. Syst. ii. 390 = A. Adscensionis.
affinis, Roem. & Schult. Syst. ii. Mant. 210 = A. purpurascens.
Antillarum, Roem. & Schult. Syst. ii. 395 = A. Antillarum.
Ascensionis, Beauv. Agrost. 30 = A. Adscensionis.
bipartita, Nees, Fl. Afr. Austr. i. 187 = A. bipartita.
bromoides, Roem. & Schult. Syst. ii. 396 = A. setifolia.
caerulescens, Beauv. Agrost. 30 = A. coerulescens.
calicina, Beauv. l. c. = A. calycina.
canariensis, Beauv. l. c. = A. coerulescens.
capensis, Beauv. l. c. = A. capensis.
capillacea, Beauv. l. c. = A. capillacea.
capillaris, Nees, Agrost. Bras. 388 = A. capillacea.
coarctata, Roem. & Schult. Syst. ii. 396 = A. setifolia.
congesta, Nees, Fl. Afr. Austr. 189 = A. congesta.
curvata, Nees, l. c. 186 = A. coerulescens.
cyanantha, Nees, ex Steud. Syn. Pl. Gram. 141 = A. cyanantha.
depressa, Beauv. Agrost. 30 = A. Adscensionis.
divaricata, Beauv. l. c. = A. divaricata.
divaricata, Nees, Agrost. Bras. 387 = A. longifolia.
elatior, Beauv. l. c. = A. coerulescens.
elliptica, Nees, Agrost. Bras. 389 = A. elliptica.
fasciculata, Roem. & Schult. Syst. ii. Mant. 578 = A. fasciculata.
festucoides, Beauv. Agrost. 30 = A. Adscensionis.
Forskolii, Nees, Fl. Afr. Austr. 188 = A. paniculata.
furcata, Beauv. Agrost. 30 (= *Eutriana juncifolia*).—Ind. occ.
gibbosa, Nees, Agrost. Bras. 383 = A. gibbosa.
gigantea, Beauv. Agrost. 30 = A. Adscensionis.
glauca, Nees, in Linnaea, xix. (1847) 688 = A. longiramea.
gossypina, Beauv. Agrost. 30; Roem. & Schult. Syst. ii. 391 = A. gossypina.
Histrix, Beauv. l. c. = A. Hystrix.
humilis, Roem. & Schult. Syst. ii. 396 = A. humilis.
Hystrix, Roem. & Schult. l. c. 390 = A. Hystrix.
interrupta, Beauv. Agrost. 30 = A. interrupta.
laevis, Nees, Agrost. Bras. ii. 384 = A. laevis.
Lamarckii, Roem. & Schult. Syst. ii. 393 = A. stipiformis.
luzoniensis, Beauv. Agrost. 30 = A. luzoniensis.
mauritiana, Nees, Fl. Afr. Austr. 188 = A. Adscensionis.
megapotamica, Roem. & Schult. Syst. ii. Mant. 578 = A. megapotamica.
murina, Nees, ex Trin. & Rupr. in Mém. Acad. Pétersb. Sér. VI. vii. (1849) 160 = A. Royleana.
nana, Nees, ex Steud. Nom. ed. II. i. 340 = A. nana.
olygantha, Beauv. Agrost. 30 = A. oligantha.
pallens, Beauv. l. c. = A. oligantha.
purpurascens, Beauv. l. c. = A. coerulescens.
racemosa, Beauv. l. c. = A. racemosa.
ramosa, Roem. & Schult. Syst. ii. 397 = A. ramosa.
recurvata, Roem. & Schult. l. c. = A. recurvata.
Royleana, Nees, ex Steud. Syn. Pl. Gram. 143 = A. Royleana.
setacea, Beauv. Agrost. 30 = A. setacea.
setifolia, Roem. & Schult. Syst. ii. 396 = A. setifolia.
Sieberiana, Roem. & Schult. Syst. ii. Mant. 578 = A. Hystrix.
spadicea, Roem. & Schult. Syst. ii. 397 = A. spadicea.
spadicea, Nees, Agrost. Bras. 385 = A. tincta.
squarrosa, Roem. & Schult. Syst. ii. Mant. 577 = A. squarrosa.
stipaeformis, Beauv. Agrost. 30 = Avena stipaeformis.
stricta, Beauv. l. c. = A. stricta.
teneriffa, Beauv. l. c. = Panicum Teneriffae.
torta, Nees, Agrost. Bras. 386 = A. tincta.
trichodes, Nees, in Hook. Kew Journ. ii. (1850) 101 = A. Cumingiana.
tuberculosa, Roem. & Schult. Syst. ii. Mant. 211 = A. tuberculosa.
vestita, Beauv. Agrost. 30 = A. vestita.

CHAETHYMENIA, Hook. & Arn. Bot. Beech. Voy. 298. t. 62 (1841) = **Jaumea**, Pers. (Compos.).
pedunculatis, Hook. & Arn. l. c. = J. pedunculatis.

CHAETIUM, Nees, Agrost. Bras. 269 (1829). *GRAMINEAE*, Benth. & Hook. f. iii. 1104.
BERCHTOLDIA, Presl, Rel. Haenk. 323. t. 43 (1830).
bromoides, *Benth. in Journ. Linn. Soc.* xix. (1881) 46.—Mexic.
festucoides, *Nees, Agrost. Bras.* 270.—Bras.

CHAETOBROMUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836) = **Danthonia**, DC. (Gramin.).
Dregeanus, Nees, Fl. Afr. Austr. 343 = D. Dregeana.
fascicularis, Nees, l. c. 341 = D. fascicularis.
interceptus, Nees, l. c. 342 = D. intercepta.
involucratus, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 = D. involucrata.

CHAETOCALYX, DC. Mém. Lég. 262 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 513.
BONNINGHAUSIA, Spreng. Syst. iii. 245 (1826).
PLANARIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 416.
RHADINOCARPUS, Vog. in Linnaea, xii. (1838) 108.
acutiflora, *Benth. in Mart. Fl. Bras.* xv. 1. 75.—Am. austr.
brasiliensis, *Benth. l. c.*—Am. austr.
hebecarpa, *Benth. l. c.* 76.—Am. austr.
latifolia, *Benth. l. c.* 75.—Am. austr.
latisiliqua, *Benth. Bot. Voy. Sulph.* 81. t. 30.—Panama.
longiflorus, *Benth. ex A. Gray, Bot. U. St. Expl. Exped.* i. 423.—Bras.
parviflora, *Benth. in Mart. Fl. Bras.* xv. 1. 74.—Am. austr.
polyphylla, *Benth. l. c.* 76.—Am. austr.
pubescens, *DC. Prod.* ii. 243.—Ins. S. Doming.
Schottii, *Torr. in Bot. Mex. Bound.* 56.—N. Mexic.
vincentina, *DC. Prod.* ii. 243.—Ins. S. Vincent.
Wislizeni, *A. Gray, Pl. Wright.* i. 51.—Am. bor. occ.

CHAETOCAPNIA, Sweet, Hort. Brit. ed. III. 685 (1839) = COETOCAPNIA, Link & Otto = **Bravoa**, Lex. (Amaryllid.).
geminiflora, Sweet, l. c. = B. geminiflora.

CHAETOCARPUS, Schreb. Gen. 75 (1789) = **Pouteria**, Aubl. (Sapotac.).
Ponteria, J. F. Gmel. Syst. 248 = P. guianensis.

CHAETOCARPUS, Thw. in Hook. Kew Journ. vi. (1854) 300. t. 10. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 323.
REGNALDIA, Baill. Adansonia, i. (1860–61) 187.
Blanchetii, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 507.—Bras.
castanocarpus, *Thw. Enum. Pl. Zeyl.* 275.—Ind. or.; Malaya.
coriaceus, *Thw. l. c.*—Zeylan.
myrsinites, *Baill. Adansonia*, xi. (1873) 95.—Bras.
Pohlii, *Muell. Arg. in Mart. Fl. Bras.* xi. 11. 508.—Bras.
pungens, Thw. in Hook. Kew Journ. vi. (1854) 301 = castanocarpus, coriaceus.

CHAETOCEPHALA, Barb. Rodr. Orch. Nov. ii. 37 (1881) = **Pleurothallis**, R. Br. (Orchid.).
lonchophylla, Barb. Rodr. l. c. 39.—Bras.
punctata, Barb. Rodr. l. c. 38.—Bras.

CHAETOCHILUS, Vahl, Enum. i. 101 (1805) = **Schwenkia**, Linn. (Solanac.).
lateriflorus, Vahl, l. c. = S. brasiliensis.

CHAETOCLADUS, Senilis [Nelson], Pinaceae, 161 (1866) = **Ephedra**, Linn. (Gnetac.).
altissima, Senilis [Nelson], l. c. 162 = E. altissima.
distachys, Senilis [Nelson], l. c. = E. vulgaris.
fragilis, Carr. Conif. ed. II. 860, Index = E. fragilis.
monostachys, Senilis [Nelson], Pinac. 162 = E. vulgaris.

CHAETOCRATER, Ruiz & Pav. Prod. Fl. Per. 61. t. 36 (1799) = **Casearia**, Jacq. (Samydac.).
capitatus, Ruiz & Pav. Fl. Per. iv. t. 396; Syst. 108 = Casearia sylvestris.
fasciculatus, Ruiz & Pav. ll. cc. 396, 107 = Casearia punctata.

CHAETOCYPERUS, Nees, in Linnaea, ix. (1834) 289 = **Eleocharis**, R. Br. (Cyperac.).
acicularis, Nees, in Mart. Fl. Bras. ii. 1. 94 = E. acicularis.
albibracteatus, Nees & Meyen, in Act. Nov. Nat. Cur. xix. Suppl. i. (1843) 95 = E. albibracteata.
arenicola, Steud. Syn. Pl. Cyp. 73 = E. Chaetaria.
Baldwinii, Torr. in Ann. Lyc. N. York, iii. (1836) 295 = E. Baldwinii.
bonariensis, Nees, in Mart. Fl. Bras. ii. 1. 96 = E. bonariensis.
capillaceus, Nees, l. c. 93 = E. capillacea.
costulatus, Nees & Meyen, in Act. Nov. Nat. Cur. xix. Suppl. i. (1843) 96 = E. acicularis.
depauperatus, Steud. Syn. Pl. Cyp. 73 = E. Chaetaria.
emarginatus, Nees, in Mart. Fl. Bras. ii. 1. 96.—Bras.
Jamesoni, Steud. Syn. Pl. Cyp. 74.—Ecuador.
Limnocharis, Nees, in Wight, Contrib. 96 = E. Chaetaria.
membranaceus, Buckl. in Proc. Acad. Sc. Philad. '1862' (1863) 10 = Scirpus nanus.
niveus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 242 = E. Chaetaria.
obtusatus, Nees, in Mart. Fl. Bras. ii. 1. 94 = E. Chaetaria.
polymorphus, Nees, in Linnaea, ix. (1834) 289 = E. acicularis, ochreata, Scirpus cernuus, nanus.
punctatus, Nees, in Mart. Fl. Bras. ii. 1. 93.—Bras.
pusillus, Steud. Syn. Pl. Cyp. 74 = Scirpus nanus.
pygmaeus, Torr. ex A. Gray, Man. Bot. N. U. St. 524 = Scirpus nanus.
radicans, Steud. Syn. Pl. Cyp. 74 = E. acicularis.
reptans, Boeck. in Flora, xl. (1857) 34 = E. minima?
rugulosus, Nees, in Bonplandia, iii. (1855) 86 = E. Chaetaria.
setaceus, Nees, in Mart. Fl. Bras. ii. 1. 94 = E. Chaetaria.
stoloniferus, Nees, in Linnaea, xix. (1847) 695 = E. montana.
subarticulatus, Nees, in Mart. Fl. Bras. ii. 1. 96.—Bras.
tenuiculus, Nees, l. c. = E. Rothiana.
trichoides, Steud. Syn. Pl. Cyp. 73 = E. acicularis.
urceolatus, Liebm. in Vidensk. Selsk. Skr. V. ii. (1851) 243 = E. minima.
viviparus, Nees, in Mart. Fl. Bras. ii. 1. 93 = E. tenuissima.

CHAETODISCUS, Steud. Syn. Pl. Cyp. 261 (1855) = **Eriocaulon**, Linn.
Gilberti, Steud. l. c.—Austral.

CHAETOGASTRA, DC. Prod. iii. 131 (1828); Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 127 = **Tibouchina**, Aubl. (Melast.).
alpestris, Mart. Nov. Gen. et Sp. iii. 101 = Pterolepis alpestris.
asperior, Cham. in Linnaea, ix. (1834) 435 = T. asperior.
bicolor, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 128 (= *Pleroma bicolor*).—Boliv.
Bonplandiana, Naud. l. c. 137 = Brachyotum lediifolium.
Callichaeta, Benth. in Hook. Journ. Bot. ii. (1840) 291 = Pterolepis glomerata.
campanularis, Benth. Pl. Hartw. 131 = Brachyotum Benthamianum.
campanularis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 130 = Brachyotum campanulare.
canescens, DC. Prod. iii. 134 = Brachyotum canescens.
cardinalis, DC. l. c. = T. cardinalis.
cataphracta, Cham. in Linnaea, ix. (1834) 451 = Pterolepis cataphracta.
cernua, DC. Prod. iii. 135 = Brachyotum cernuum.
Chamaecistus, Griseb. Fl. Brit. W. Ind. 267 (= *Pleroma Chamaecistus*).—Ind. occ.
ehironioides, Griseb. l. c. (= *Pleroma strigosum, ehironioides*).—Ind. occ.
ciliaris, DC. Prod. iii. 132 (= *Pleroma ciliare*).—N. Granat.
cistoides, Griseb. Fl. Brit. W. Ind. 267 (= *Pleroma cistoides*).—Ins. S. Vincent.
clinopodifolia, DC. Prod. iii. 133 = T. clinopodiifolia.
conferta, DC. l. c. 135 = Brachyotum confertum.
cubensis, Griseb. Cat. Pl. Cub. 103 = Chaetolepis saturacioides.

CHAETOGASTRA :—
debilis, Cham. in Linnaea, ix. (1834) 499 = Tibouchina debilis.
debilis, Miq. in Linnaea, xxii. (1849) 540 = T. ursina.
depauperata, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 129 = T. longifolia.
divaricata, DC. Prod. iii. 132 = Pterogastra divaricata.
echinata, DC. Prod. iii. 135 (= *Pleroma echinatum*).—Peruv.
ferruginea, Hook. & Arn. Bot. Beech. Voy. 423 = T. longifolia.
floribunda, Griseb. in Lechl. Berb. Am. Austr. 59 = Brachyotum floribundum.
fraterna, DC. Prod. iii. 133 = T. gracilis.
Geitneriana, Schlecht. in Linnaea, xxvii. (1854) 523.—N. Granat.
glomerata, Benth. in Hook. Journ. Bot. ii. (1840) 290 = Pterolepis glomerata.
Goudotii, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 131 = Brachyotum strigosum.
gracilis, DC. Prod. iii. 133 = T. gracilis.
havanensis, DC. l. c. 132 = T. longifolia.
hermannioides, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 183 = Brachyotum hermannioides.
hieracioides, DC. Prod. iii. 133 = T. hieracioides.
hirsuta, DC. l. c. = T. gracilis.
hispida, DC. l. c. 132 = T. longifolia.
hypericoides, DC. l. c. = Desmoscelis villosa.
ladanoides, Mart. Herb. Fl. Bras. n. 500 = Pterolepis trichotoma.
lanceolata, DC. Prod. iii. 131 = T. longifolia.
lasiophylla, Benth. in Hook. Journ. Bot. ii. (1840) 291 = T. lasiophylla.
lepidota, DC. Prod. iii. 134 = T. lepidota.
Lhotskyana, Steud. in Flora, xxvii. (1844) 720 = T. Lhotzkyana.
Lindeniana, Planch. in Fl. des Serres, Sér. I. (1854–55) 113. t. 1011–12 = T. grossa.
longifolia, DC. Prod. iii. 132 = T. longifolia.
lutescens, Griseb. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 48 = Brachyotum lutescens.
lutescens, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 134 = Brachyotum Naudini.
lychnitoides, DC. Prod. iii. 132 = Desmoscelis villosa.
Martiana, Benth. in Hook. Journ. Bot. ii. (1840) 290 = T. herbacea.
microdon, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 134 = Brachyotum microdon.
microphylla, Naud. l. c. 136 = Brachyotum microphyllum.
mollis, DC. Prod. iii. 134 (= *Pleroma exappendiculatum*).—Peruv.
Moritziana, Klotzsch, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 46 (= *Pleroma Naudinianum*).—N. Granat.
multiflora, Benth. Pl. Hartw. 181 = Monochaetum Hartwegianum.
muricata, DC. Prod. iii. 134 = T. grossa.
Naudiniana, Decne. in Rev. Hortic. Sér. III. i. (1847) 86 (= *Pleroma Naudinianum*).—Mexic.
nigritella, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 137 = Bucquetia nigritella.
origanoides, Griseb. Cat. Pl. Cub. 103 = Chaetolepis cubensis.
panicularis, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 129 (= *Pleroma paniculare*).—Bolivia.
Pentlandii, Naud. l. c. 133 = Brachyotum Pentlandi.
quinquenervis, Naud. l. c. 130 = Brachyotum quinquenerve.
repanda, Mart. Nov. Gen. et Sp. iii. 100. t. 246 = Pterolepis repanda.
reticulata, DC. Prod. iii. 131 = T. reticulata.
rhynchanthera, Benth. Pl. Hartw. 180 = Monochaetum Bonplandii.
rosea, Regel, Cat. Pl. Hort. Aksakov. 31.—Hab. ?
rosmarinifolia, Benth. Pl. Hartw. 131 = Miconia ledifolia.
rosmarinifolia, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 131 = Brachyotum rosmarinifolium.
rostrata, Naud. l. c. 135 = Brachyotum rostratum.
rufipilis, Schlecht. in Linnaea, xiii. (1839) 430 (= *Pleroma rufipile*).—Mexic.
sanguinolenta, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 131 = Brachyotum sanguinolentum.
sarmentosa, DC. Prod. iii. 134 (= *Pleroma laxum*).—Peruv.
saturejoides, Griseb. Cat. Pl. Cub. 103 = Chaetolepis saturaeioides.

CHAETOGASTRA :—
scabriuscula, Schlecht. in Linnaea, xiii. (1839) 431 (= *Pleroma scabriusculum*).—Mexic.
Schiedeana, Cham. & Schlecht. in Linnaea, v. (1830) 565 (= *Pleroma rufipile*).—Mexic.
sherardioides, DC. Prod. iii. 133 = Pterolepis glomerata.
speciosa, DC. l. c. 131 = Meriania speciosa.
stachyoides, DC. l. c. 132 = Desmoscelis villosa.
stricta, Benth. Pl. Hartw. 180 = Brachyotum cernuum.
stricta, DC. Prod. iii. 134 = Brachyotum strigosum.
strigillosa, DC. l. c. 133 = T. gracilis.
strigosa, DC. l. c. iii. 134 (= *Pleroma strigosum*).—Am. austr.
striphnocalyx, Mart. Nov. Gen. et Sp. iii. 101. t. 246 = Pterolepis striphnocalyx.
sulphurea, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 135 = Brachyotum ledifolium.
tortuosa, DC. Prod. iii. 132 (= *Pleroma mexicanum*).—Mexic.
Weddelliana, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 129 (= *Pleroma paniculare*).—Bolivia.

CHAETOLEPIS, Miq. Comm. Phyt. ii. 72 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 744.
HAPLODESMIUM, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 154. t. 5.
TRIMERANTHUS, Karst. in Linnaea, xxx. (1859–60) 159.
alpestris, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Venezuela.
alpina, *Naud. in Ann. Sc. Nat.* Sér. III. xiv. (1850) 140.—Bras.
anisandra, *Naud. l. c.*—Bras.
cubensis, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Cuba.
ericoides, Triana, l. c. 51. t. 3. f. 35 *g* = Marcetia ericoides.
heterophylla, *Triana, l. c.* 51.—Uruguay.
Lindeniana, *Triana, l. c.*—N. Granat.
loricarella, *Triana, l. c.*—N. Granat.
microphylla, *Miq. Comm. Phyt.* 72.—N. Granat.
saturaeioides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 51.—Cuba.
thymifolia, *Triana, l. c.* 50.—N. Granat.

CHAETONYCHIA, Sweet, Hort. Brit. ed. III. 263 (1839) = **Paronychia**, Juss.
cymosa, Sweet, l. c. = P. cymosa.

CHAETOPAPPA, DC. Prod. v. 301 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 268.
CHAETANTHERA, Nutt. in Journ. Acad. Sc. Philad. vii. (1834) 111.
CHAETOPHORA, Nutt. ex DC. Prod. v. 301, in syn. (1836).
DIPLOSTELMA, Rafin. ex Torr. & Gray, Fl. N. Am. ii. 187 (1840).
DISTASIS, DC. Prod. v. 279 (1836).
asteroides, *DC. l. c.* 301.—Am. bor.
modesta, *A. Gray, in Proc. Am. Acad.* xvi. (1881) 82.—Mexic.
Parryi, *A. Gray, l. c.*—Texas.

CHAETOPHORA, Nutt. ex DC. Prod. v. 301, in syn. (1836) = **Chaetopappa**, DC. (Compos.).
asteroides, Nutt. ex DC. l. c. = Chaetopappa asteroides.

CHAETOPTELEA, Liebm. in Kjoeb. Vidensk. Meddel. (1850) 76 = **Ulmus**, Linn. (Urticac.).
mexicana, Liebm. l. c. 77 = U. mexicana.

CHAETOSCIADIUM, Boiss. Fl. Orient. ii. 1078 (1872) = TORILIS, DC. = **Caucalis**, Linn. (Umbellif.).
trichospermum, Boiss. l. c. = Cauc. trichosperma.

CHAETOSPORA, H. B. & K. Nov. Gen. et Sp. i. 229 (1815) = **Rynchospora**, Vahl (Cyperac.).
aurea, Cham. & Schlecht. in Linnaea, vi. (1831) 28 = R. aristata.
aurea, H. B. & K. Nov. Gen. et Sp. i. 231 = R. aurea.
capitata, H. B. & K. l. c. 229 = R. capitata.
ferruginea, Cham. & Schlecht. in Linnaea, vi. (1831) 28, partim = R. eximia.
ferruginea, H. B. & K. Nov. Gen. et Sp. ii. 85 = R. glauca.
pterocarpa, H. B. & K. Nov. Gen. et Sp. i. 230 = R. pterocarpa.

CHAETOSPORA :—
triceps, Cham. & Schlecht. in Linnaea, vi. (1831) 29 = Rynchospora triceps.
vicozensis, Schrad. ex Roem. & Schult. Syst. ii. Mant. 46 = R. glauca.

CHAETOSPORA, R. Br. Prod. 232 (1810) = **Schoenus**, Linn. (Cyperac.).
albescens, Franch. & Sav. Enum. Pl. Jap. ii. 548 = S. apogon.
alpina, F. Muell. Fragm. ix. 39 = Carpha alpina.
anceps, R. Br. Prod. 233 = Mesomelaena anceps.
antarctica, Hook. f. Fl. Antarct. ii. 361. t. 147.—Chili.
arundinacea, A. Dietr. Sp. Pl. ii. 31 = Asterochaete arundinacea.
aurata, Nees, in Ann. Nat. Hist. Ser. I. vi. (1841) 49 = S. curvifolius.
avenacea, F. Muell. Fragm. ix. 40 = Cyathochaeta avenacea.
axillaris, R. Br. Prod. 233 = S. axillaris.
brevisetis, R. Br. l. c. 232 = S. brevisetis.
brevisetis, F. Muell. Fragm. ix. 37 = S. capitatus.
Burmanni, Schrad. Anal. Cap. Cyp. 32. t. 3. f. 4.—Afr. austr.
calostachys, R. Br. Prod. 233 = S. calostachyus.
capillacea, Hook. f. Fl. Tasm. ii. 81. t. 141 A = Elynanthus capillaceus.
capillacea, Nees, in Linnaea, ix. (1834) 299; x. (1836) 192 = S. capillaceus.
capillaris, F. Muell. Fragm. ix. 34 = Elynanthus capillaceus.
Capitellum, A. Dietr. Sp. Pl. ii. 27 = Ficinia setiformis.
ciliata, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 128 = Rynchospora elatior.
circinalis, Schrad. in Goett. Gel. Anz. (1821) III. 2070 = S. circinalis.
clandestina, F. Muell. Fragm. ix. 40 = Cyathochaeta clandestina.
compressa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 71; Nees, in Lehm. Pl. Preiss. ii. 85 = Scirpus Caricis.
concava, Nees, in Ann. & Mag. Nat. Hist. Ser. I. vi. (1841) 47 = Lepidospermum concavum.
concinna, Hook. f. Fl. N. Zeal. i. 274. t. 62 B = S. concinnus.
cruenta, Nees, in Lehm. Pl. Preiss. ii. 85 = S. cruentus.
curvifolia, R. Br. Prod. 232 = S. curvifolius.
cuspidata, Nees, in Linnaea, vii. (1832) 529 = Elynanthus cuspidatus.
cygnea, Nees, in Ann. Nat. Hist. Ser. I. vi. (1841) 49 = S. brevisetis.
dactyloides, A. Dietr. Sp. Pl. ii. 32 = Asterochaete glomerata.
deformis, R. Br. Prod. 232 = S. deformis.
deusta, F. Muell. Fragm. ix. 39 = Mesomelaena deusta.
diandra, F. Muell. l. c. = Cyathochaeta diandra.
diodon, Schrad. ex Nees, in Mart. Fl. Bras. ii. 1. 148 = Rynchospora lunata.
dispar, A. Dietr. Sp. Pl. ii. 32 = Ficinia secunda.
distachya, Nees, in Linnaea, x. (1836) 192.—Afr. austr.
distans, F. Muell. Fragm. ix. 35 = S. calostachyus.
effusa, Schrad. ex Kunth, Enum. Pl. ii. 286 = Pleurostachys sparsiflora.
elongata, Nees, in Lehm. Pl. Preiss. ii. 275 = Mesomelaena anceps.
ferruginea, Reichb. Fl. Germ. Excurs. 74 = S. ferrugineus.
fimbriolata, Nees, in Linnaea, x. (1836) 191.—Afr. austr.
fimbristyloides, F. Muell. Fragm. ix. 34 = Cladium undulatum.
flexuosa, Schrad. Anal. Cap. Cyp. 33. t. 3. f. 3 = S. flexuosus.
globosa, H. B. & K. Nov. Gen. et Sp. i. 135 = Rynchospora globosa.
hexandra, Boeck. in Flora, lxi. (1878) 37.—Afr. austr.
imberbis, R. Br. Prod. 233 = S. apogon.
japonica, Franch. & Sav. Enum. Pl. Jap. ii. 548 = S. apogon.
laeta, Kunze, ex Steud. Nom. ed. II. i. 341 = Fimbristylis aspera.
lanata, R. Br. Prod. 232 = S. lanatus.
laxa, Hook. f. Fl. Antarct. ii. 361. t. 146 = Carpha paniculata.
lucens, Poir. in Dict. Sc. Nat. viii. 58 = S. nitens.
madagascariensis, Steud. Syn. Pl. Cyp. 161.—Madag.

CHAETOSPORA :—
Mauritii, Steud. Syn. Pl. Cyp. 161 = Rynchospora glauca.
microstachya, Nees, in Lehm. Pl. Preiss. ii. 84 = Schoenus Drummondii.
microstachys, Schrad. Anal. Fl. Cap. i. 33, in obs. = S. circinalis.
nana, Nees, in Lehm. Pl. Preiss. ii. 85 = S. nanus.
natans, F. Muell. Fragm. ix. 38 = S. natans.
Neesii, Boeck. in Linnaea, xxxviii. (1874) 297 = Tricostularia Neesii.
nigricans, Kunth, Enum. Pl. ii. 323 = S. nigricans.
nitens, R. Br. Prod. 233 = S. nitens.
oligostachya, Boeck. in Flora, xlii. (1859) 101 = S. nigricans.
oligostachya, F. Muell. Fragm. ix. 38 = S. bifidus.
paludosa, R. Br. Prod. 233 = Tricostularia paludosa.
paniculata, Steud. Syn. Pl. Cyp. 161.—Ins. Borbon.
pedicellata, R. Br. Prod. 232 = S. pedicellatus.
pterosperma, H. B. & K. Nov. Gen. et Sp. i. 230 = Haplostylis barbata.
punctoria, A. Dietr. Sp. Pl. ii. 28 = Bucckia punctoria.
rhynchospermoides, Phil. f. Pl. Vasc. Chil. 306 = seq.
rhynchosporoides, Steud. Syn. Pl. Cyp. 162.—Chili.
robusta, Kunth, Enum. Pl. ii. 325.—Afr. austr.
rufa, S. F. Gray, Nat. Arr. Brit. Pl. ii. 71 = Scirpus rufus.
sphaerocephala, R. Br. Prod. 233 = Mesomelaena sphaerocephala.
spicata, Boeck. in Linnaea, xxxviii. (1874) 297 = Tricostularia compressa.
striata, A. Dietr. Sp. Pl. ii. 27 = Ficinia setiformis.
stygia, R. Br. Prod. 233 = Mesomelaena stygia.
tenax, Hook. f. Fl. N. Zel. i. 273 = S. brevifolius.
Tendo, Hook. f. l. c. = S. Tendo.
tenella, Rupr. Sert. Tiansch. 34.—Turkestan.
tenera, A. Dietr. Sp. Pl. ii. 32 = S. tener.
tenuissima, Hook. f. Fl. Tasm. ii. 81. t. 14 = Lepidospora tenuissima.
tenuissima, Steud. Syn. Pl. Cyp. 319, 162 = S. apogon.
tetragona, R. Br. Prod. 233 = Mesomelaena tetragona.
turbinata, R. Br. l. c. 232 = S. turbinatus.
umbellulifera, Boeck. in Flora, lxv. (1882) 28 = S. apogon.
villosa, Nees, ex Steud. Nom. ed. II. i. 341 = S. villosus.

CHAETOSTACHYS, Benth. in Wall. Pl. As. Rar. ii. 19 (1831) = **Lavandula**, Tourn. (Labiat.).
multifida, Benth. l. c. = L. Burmanni.

CHAETOSTEMMA, Reichb. Consp. 176 (1828) = seq.

CHAETOSTOMA, DC. Prod. iii. 112 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 737.
acuminatum, *Cogn. in Mart. Fl. Bras.* xiv. III. 35.—Bras.
armatum, Cogn. l. c. 31 = diosmoides.
castratum, *Cogn. l. c.* 36.—Bras.
diosmoides, *Mart. Nov. Gen. et Sp.* iii. 130. t. 264.—Bras.
ericoides, DC. Prod. iii. 112 = Marcetia ericoides.
fastigiatum, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 191.—Bras.
Gardneri, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 25. t. 1. f. 8. c.—Bras.
Glaziovii, *Cogn. in Mart. Fl. Bras.* xiv. III. 30.—Bras.
inermc, *Naud. in Ann. Sc. Nat.* Sér. III. iii. (1845) 191.—Bras.
longiflorum, *Cogn. in Mart. Fl. Bras.* xiv. III. 32.—Bras.
microlicioides, Cham. in Linnaea, ix. (1834) 382 = Microlicia Chamissois.
oxyantherum, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 25, pro parte.—Bras.
oxyantherum, Triana, l. c. pro parte = acuminatum, castratum.
pnngens, *DC. Prod.* iii. 112.—Bras.
Riedelianum, *Cogn. in Mart. Fl. Bras.* xiv. III. 33.—Bras.
tetrastichum, DC. Prod. iii. 112 = Marcetia ericoides, tetrasticha.

CHAETOSUS, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 226. *APOCYNACEAE*, Benth. & Hook. f. ii. 695.
CHOETOSUS, Benth. in Journ. Linn. Soc. i. (1857) 87.
volubilis, *Benth. l. c.*—N. Guinea.

CHAETOTHYLAX, Nees, in Mart. Fl. Bras. ix. 153. t. 26 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1107.
CHAETOTHYLOPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 163.
HEINZELIA, Nees, in Mart. Fl. Bras. ix. 153 (1847).
lythroides, *Benth. & Hook. f. Gen.* ii. 1108.—Am. austr.
micranthus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 515.—Nicaragua.
phyllostachyus, *Nees, in DC. Prod.* xi. 313.—Mexic.
tocantinus, *Nees, in Mart. Fl. Bras.* ix. 153.—Bras.
umbrosus, *Nees, in DC. Prod.* xi. 313.—N. Granat.

CHAETOTHYLOPSIS, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 163 = **Chaetothylax**, Nees (Acanthac.).
micrantha, Oerst. l. c. = Chactothylax micranthus.

CHAETOTROPIS, Kunth, Rev. Gram. i. 72, 271. t. 47 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1151.
andina, *Ball, in Journ. Linn. Soc.* xxii. (1885) 58.—Peruv.
chilensis, *Kunth, Rev. Gram.* i. 72, 271. t. 47.—Chili.
latifolia, Phil. in Linnaea, xxx. (1859–60) 205 = chilensis.

CHAETOTROPSIS, D. Dietr. Syn. Pl. i. 134 (1839) = praec.

CHAETURUS, Link, in Schrad. Journ. ii. (1799) 313. *GRAMINEAE*, Benth. & Hook. f. iii. 1149.
divaricatus, DC. Cat. Hort. Monsp. 93 = fasciculatus.
fasciculatus, *Link, in Schrad. Journ.* ii. (1799) 313.—Hispan.
prostratus, *Hack. & Lange, in Kjoeb. Vidensk. Meddel.* (1877–78) 223.—Lusitan.
spicatus, Schrad. Ind. Sem. Hort. Gotting. ex Steud. Nom. ed. II. i. 342 = fasciculatus.

CHAILLETIA, DC. in Ann. Mus. Par. xvii. (1811) 153. *CHAILLETIACEAE*, Benth. & Hook. f. i. 341.
DICHAPETALUM, Thou. Gen. Madag. 23 (1806).
LEUCOSIA, Thou. l. c. (1806).
MESTOTES, Soland. ex Steud. Nom. ed. II. i. 342 (1842).
MOACURRA, Roxb. Hort. Beng. 21 (1814); Fl. Ind. ii. 69 (1832).
PATRISIA, Rohr, ex Steud. Nom. ed. II. i. 342 (1840).
PLAPPERTIA, Reichb. Consp. 146 (1828).
QUILESIA, Blanco, Fl. Filip. ed. I. 176 (1837).
SYMPHYLLANTHUS, Vahl, in Skriv. Nat. Hist. Selsk. vi. 86 (1810).
WAHLENBERGIA, R. Br. in Wall. Cat. p. 250, sub n. 4342 (1831).
affinis, Planch. ex Benth. in Hook. Niger Fl. 276 = toxicaria.
Bangii, *F. Didr. in Kjoeb. Vidensk. Meddel.* (1854) 195.—Afr. trop.
Benthamiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 610.—Ins. Philipp.
Benthamii, *F. Didr. in Kjoeb. Vidensk. Meddel.* (1854) 196.—Afr. trop.
Bojeri, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 85.—Malacca.
Brunoniana, Wall. Cat. n. 4038 = Helferiana.
chlorina, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 88.—Madag.
cubensis, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 41. t. 246. f. 3.—Cuba.
cymosa, *Hook. Ic. Pl.* t. 591.—Afr. austr.
cymulosa, *Oliver, Fl. Trop. Afr.* i. 340.—Afr. trop.
deflexa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 109.—Afr. trop.
deflexifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 611.—Malaya.
Dichapetalum, *DC. Prod.* ii. 57.—Madag.
discolor, *Baker, in Journ. Linn. Soc.* xx. (1883) 119.—Madag.
edulis, Kurz, in Journ. As. Soc. Beng. xli. (1872) ii. 298 = Sarcostigma edule.
erecta, G. Don, in Edinb. Phil. Journ. xi. (1824) 348 = toxicaria.
fasciculata, *Spreng. Syst.* i. 931.—Madag.
flavicans, *Klotzsch, in Schomb. Fauna & Fl. Gui.* 1184, nomen.—Guiana.

CHAILLETIA :—
flexuosa, *Oliver, Fl. Trop. Afr.* i. 340.—Afr. trop.
floribunda, *Planch. in Hook. Ic. Pl.* t. 792.—Afr. trop.
gelonioides, *Hook. f. Fl. Brit. Ind.* i. 570.—Ind. or.; Malaya.
Griffithii, *Hook. f. l. c.* 571.—Malacca.
hainanensis, *Hance, in Journ. Bot.* xxiii. (1885) 322.—China.
Helferiana, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 297.—Burma.
Heudelotii, *Planch. ex Oliver, Fl. Trop. Afr.* i. 344.—Afr. trop.
hirtella, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 88.—Ins. Mascar.
hispida, *Oliver, Fl. Trop. Afr.* i. 343.—Afr. trop.
lanuginosa, Maing. ex Hook. f. Fl. Brit. Ind. i. 572 = Griffithii.
Lauro-cerasus, *Planch. ex Hook. f. l. c.*—Ins. Penang.
Leucosia, *Spreng. Syst.* i. 931.—Madag.
longipetala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 611.—Burma.
macrophylla, *Oliver, Fl. Trop. Afr.* i. 343.—Afr. trop.
mossambicensis, *Klotzsch, in Peters, Reise Mossamb. Bot.* 108.—Afr. trop.
nigrescens, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 85.—Madag.
oblonga, *Hook. f. ex Benth. in Hook. Niger Fl.* 277.—Afr. trop.
Pachypus, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 87.—Madag.
pallida, *Oliver, Fl. Trop. Afr.* i. 343.—Afr. trop.
paniculata, *Benth. in Hook. Niger Fl.* 279.—Afr. trop.
papuana, *Becc. Malesia*, i. (1877) 176.—N. Guinea.
Parkerii, *Planch. ex Triana & Planch. in Ann. Sc. Nat.* Sér. V. xv. (1872) 379.—Venezuela.
pedunculata, *DC. in Ann. Mus. Par.* xvii. (1811) 154.—Guiana.
reflexa, *Klotzsch, in Peters, Reise Mossamb. Bot.* 109. t. 20.—Afr. trop.
rufa, *Tul. in Ann. Sc. Nat.* Sér. IV. viii. (1857) 86.—Madag.
rufipilis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 611.—Afr. trop.
sessiliflora, DC. in Ann. Mus. Par. xvii. 1811) 154 = Tapura guianensis.
subauriculata, *Oliver, Fl. Trop. Afr.* i. 344.—Afr. trop.
subcordata, *Hook. f. ex Benth. in Hook. Niger Fl.* 277.—Afr. trop.
sumatrana, Miq. Fl. Ind. Bat. Suppl. 328 = gelonioides.
Thomsoni, *Oliver, Fl. Trop. Afr.* i. 342.—Afr. trop.
timoriensis, *DC. Prod.* ii. 57.—Ins. Timor.
toxicaria, *G. Don, in Edinb. Phil. Journ.* xi. 1824 348.—Afr. trop.
vestita, *Benth. in Hook. Kew Journ.* iii. (1851) 372.—Bras.
Virchowii, *O. Hoffm. & Hildeb. ex O. Hoffm. Sert. Madag.* 11.—Madag.
viticnsis, *Seem. Viti*, 434, nomen; *Fl. Vit.* 38.—Ins. Fiji.

CHAITAEA, Soland. ex Seem. Fl. Vit. 102 (1865) = **Tacca**, Forst.
Tacca, Soland. ex Seem. l. c. = T. pinnatifida.

CHAITURUS, Willd. Prod. Fl. Berl. 200 (1787) = **Leonurus**, Linn. (Labiat.).
leonuroides, Willd. l. c. 201 = L. Marrubiastrum.
Marrubiastrum, Spenn. in T. Nees, Gen. Fl. Germ. fasc. 18. t. 353 = L. Marrubiastrum.

CHAIXIA, Lapeyr. Hist. Pl. Pyr. Suppl. 38 (1818) = **Ramondia**, Rich. (Gesnerac.).
Myconi, Lapeyr. l. c. 39 = R. pyrenaica.

CHAKIATELLA, DC. Prod. v. 563 (1836) = CHATIAKELLA, Cass. = **Wulffia**, Neck. (Compos.).

CHALARIUM, Poit. ex DC. Prod. v. 546, in syn. (1836) = OGIERA, Cass. = **Eleutheranthera**, Poit. (Compos.).

CHALCANTHUS, Boiss. Fl. Orient. i. 211 (1867) = **Hesperis**, Linn. (Crucifer.).
renifolius, Boiss. l. c. 212 = H. renifolius.

CHALCAS, Linn. Mant. 68 (1767) = **Murraya**, Koen. (Rutac.).
Cammuneng, Burm. f. Fl. Ind. 104 = M. exotica.
crenulata, F. Muell. Phyt. N. Hebr. 7 = M. crenulata.
intermedia, Roem. Syn. Hesper. i. 48 = M. exotica.
japanensis, Lour. Fl. Cochinch. 271 = M. exotica.
Koenigii, Kurz, in Journ. As. Soc. Beng. xliv. (1875) II. 132 = M. Koenigii.
paniculata, Linn. Mant. i. 68 = M. exotica.
sumatrana, Roem. Syn. Hesper. i. 49 = M. exotica.

CHALEPOA, Hook. f. Ic. Pl. t. 1082 (1871) = **Tribeles**, Phil. (Pittospor.).
magellanica, Hook. f. l. c. = T. australis?

CHALEPOPHYLLUM, Hook. f. Ic. Pl. t. 1148 (1873). *RUBIACEAE*, Benth. & Hook. f. ii. 50.
guyanense, *Hook. f. l. c.*—Guiana.

CHALYBEA, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 99 = **Pachyanthus**, A. Rich. (Melastomac.).
corymbifera, Naud. l. c. 100 = P. corymbosus.
corymbosa, Triana, in Trans. Linn. Soc. xxviii. (1871) 97 = P. corymbosus.

CHAMABAINIA, Wight, Ic. vi. 11. t. 1981 (1853). *URTICACEAE*, Benth. & Hook. f. iii. 387.
cuspidata, *Wight, l. c.*—Ind. or.
squamigera, Wall. Cat. n. 4592; Wedd. in DC. Prod. xvi. I. 218 = cuspidata.

CHAMAEBATIA, Benth. Pl. Hartw. 308 (1850). *ROSACEAE*, Benth. & Hook. f. i. 617.
foliolosa, *Benth. l. c.*—Calif.
foliolosa, Newberry, in Pacif. Rail. Rep. vi. 73 = Spiraea Millefolium.

CHAMAEBATIARIA, Maxim. in Act. Hort. Petrop. vi. (1879) 225 = **Spiraea**, Linn. (Rosac.).
Millefolium, Maxim. l. c. = S. Millefolium.

CHAMAEBETULA, Opiz, in Lotos, v. (1855) 258 = **Betula**, Tourn. (Cupulif.).
acutifolia, Opiz, l. c. 259.—Europ.
glandulosa, Opiz, l. c. = B. glandulosa.
Gmelinii, Opiz, l. c. = B. fruticosa.
Hookeri, Opiz, l. c. = B. glandulosa.
humilis, Opiz, l. c. = B. humilis.
nana, Opiz, l. c. = B. nana.
pumila, Opiz, l. c. = B. pumila.
rotundifolia, Opiz, l. c. = B. glandulosa.

CHAMAEBUXUS, [Tourn.] Spach, Hist. Vég. Phan. vii. 125 (1839) = **Polygala**, Linn.
alpestris, Spach, l. c. 127 = P. Chamaebuxus.
arillata, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 153 = P. arillata.
coriacea, Opiz, Seznam, 28 = P. Chamaebuxus.
karensium, Kurz, For. Fl. Brit. Burma, i. 79 = P. karensium.
paniculata, Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat. i. 154 = P. arillata.
polygaleoides, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 11 = P. Chamaebuxus.
pulchra, Hassk. Pl. Jav. Rar. 294, in textu = P. pulchra.
Vayredae, Willk. Illustr. Pl. Hisp. 33. t. 23 = P. Vayredae.
venenosa, Hassk. Pl. Jav. Rar. 294.—Java.
vulgaris, Schur, Enum. Pl. Transs. 90 = P. Chamaebuxus.

CHAMAECALAMUS, Meyen, Reise, i. 456 (1834) = **Deyeuxia**, Clar. (Gramin.).
spectabilis, Meyen, l. c.; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 156 (= *Bromidium spectabile*).—Peruv.

CHAMAECASSIA, [Breyn] Link, Handb. ii. 139 (1831) = **Cassia**, Tourn. (Legumin.).
grandiflora, Link, l. c. = Cassia laevigata.
laevigata, Link, l. c. = Cassia laevigata.

CHAMAECERASUS, [Tourn.] Medic. Phil. Bot. i. 126 (1789) = **Lonicera**, Linn. (Caprifol.).
alpigena, Medic. l. c. = L. alpigena.
coerulea, Delarb. Fl. Auv. ed. II. 131 = L. coerulea.
dumetorum, Delarb. l. c. = L. Xylosteum.
fruticosa, Ser. in DC. Prod. ii. 537, sphalm. = Prunus Chamaecerasus.

CHAMAECERASUS:—
nigra, Medic. Phil. Bot. i. 126; Pfl. Anat. 390 = L. nigra.
Standishii, Lescuyer, in L'Horticult. Franç. (1862) 138. t. 10.—Hab.?
tatarica, Medic. Phil. Bot. i. 126; Pfl. Anat. 390 = Lonicera tatarica.
Xylosteum, Medic. Phil. Bot. i. 126 = L. Xylosteum.

CHAMAECISTUS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 401 (1821) = **Loiseleuria**, Desv. (Ericac.).
serpyllifolius, S. F. Gray, l. c. = L. procumbens.

CHAMAECISTUS, Regel, Gartenfl. (1874) 60 = **Rhododendron**, Linn. (Ericac.).
austriacus, Regel, l. c. = R. Chamaecistus.
kamtschaticus, Regel, l. c. = R. kamtschaticum.

CHAMAECLADON, Miq. in Bot. Zeit. xiv. (1856) 564. *AROIDEAE*, Benth. & Hook. f. iii. 983.
angustifolium, *Schott, in Bonplandia*, vi. (1858) 369.—Ins. Penang.
consobrinum, *Schott, in Ann. Mus. Bot. Lugd. Bat.* i. 126.—Sumatra.
Griffithii, *Schott, in Bonplandia*, vi. (1858) 369.—Malaya.
humile, *Miq. Fl. Ind. Bat.* iii. 213.—Ins. Penang.
lanceolatum, *Miq. in Bot. Zeit.* xiv. (1856) 564.—Java.
metallicum, *N. E. Brown, in Illustr. Hortic.* xxxi. (1884) t. 539.—Borneo.
nigrescens, *Schott, in Ann. Mus. Bot. Lugd. Bat.* i. 126.—Sumatra.
obliquatum, *Schott, in Bonplandia*, vi. (1858) 369.—Ins. Penang.
ovalifolium, *Schott, in Ann. Mus. Bot. Lugd. Bat.* i. 281.—Borneo.
ovatum, *Schott, in Bonplandia*, vii. (1859) 30.—Malaya.
purpurascens, Schott, l. c. vi. (1858) 369 = lanceolatum.
pygmaeum, Engl. in DC. Monog. Phan. ii. 345 = lanceolatum.
rubescens, *Schott, Gen. Aroid.* sub t. 60, in obs.—Hab.?
sanguinolentum, *Schott, Prod. Aroid.* 316.—Malaya.
saxorum, *Schott, in Ann. Mus. Bot. Lugd. Bat.* i. 280.—Borneo.
subcordatum, *Schott, l. c.* 281.—Borneo.
truncatum, *Schott, in Bonplandia*, vi. (1858) 369.—Burma.

CHAMAECLEMA, Moench, Meth. 393 (1794) = **Nepeta**, Tourn. (Labiat.).
hederacea, Moench, l. c. = N. Glechoma.

CHAMAECNIDE, Nees & Mart. ex Miq. in Mart. Fl. Bras. iv. I. 203 (1853) = **Pilea**, Lindl. (Urticac.).
microphylla, Nees, ex Miq. l. c. = P. muscosa.

CHAMAECRISTA, Moench, Meth. 272 (1794) = **Cassia**, Tourn. (Legumin.).
Burmanni, Eckl. & Zeyh. Enum. 260 = Cassia capensis.
capensis, E. Mey. in Linnaea, vii. (1832) 172 = Cassia capensis.
comosa, E. Mey. Comm. Pl. Afr. Austr. 160 = Cassia comosa.
nictitans, Moench, Meth. 272 = Cassia nictitans.
plumosa, E. Mey. Comm. Pl. Afr. Austr. 159 = Cassia mimosoides.
stricta, E. Mey. l. c. = Cassia mimosoides.

CHAMAECYPARIS, Spach, Hist. Vég. Phan. xi. (1842) 329 = **Thuya**, Tourn. (Conif.).
andelyensis, Hort. ex Gord. Pinet. ed. II. 365 = *leptoclada*.
Boursieri, Carr. Conif. ed. II. 125 = *Lawsoniana*.
Boursieri, Decne. in Bull. Soc. Bot. Fr. i. (1854) 70 = Juniperus occidentalis.
breviramea, Maxim. in Bull. Acad. Pétersb. x. (1866) 489 = T. obtusa.
decussata, Hort. ex Carr. Conif. ed. II. 140 = T. orientalis.
ericoides, Carr. l. c. = T. orientalis, occidentalis.
excelsa, Fisch. ex Carr. l. c. 127 = T. excelsa.
glauca, Hort. ex Carr. l. c. ed. I. 141 = T. orientalis.
Keteleri, Stand. ex Parl. in DC. Prod. xvi. II. 466 = T. obtusa.
kewensis, Hort. ex Carr. Conif. ed. I. 133 = T. sphaeroidalis.

CHAMAECYPARIS:—
Lawsoniana, Parl. in Ann. Mus. Stor. Nat. Fir. i. (1864) 181.—Am. bor. occ.
leptoclada, Henk. & Hochst. Nadelhoelz. 257.—Japon.
nana, Hort. ex Parl. in DC. Prod. xvi. II. 464 = Thuya sphaeroidalis.
nootkaënsis, Carr. Conif. ed. II. 127 = seq.
nutkaënsis, Lindl. & Gord. in Journ. Hort. Soc. v. (1850) 207 = T. excelsa.
nutkatensis, Spach, Hist. Vég. Phan. xi. 333 = praec.
obtusa, Sieb. & Zucc. in Endl. Conif. 63 = T. obtusa.
pendula, Maxim. in Bull. Acad. Pétersb. x. (1866) 489 = T. obtusa.
pisifera, Sieb. & Zucc. in Endl. Conif. 64 = T. pisifera.
pseudo-squarrosa, Parl. in DC. Prod. xvi. II. 467 = T. sphaeroidalis.
pumila, Hort. ex Carr. Conif. ed. II. 124 = T. sphaeroidalis.
sphaeroidea, Spach, Hist. Vég. Phan. xi. 331 = T. sphaeroidalis.
squarrosa, Hort. ex Carr. Conif. ed. I. 65 = T. sphaeroidalis.
squarrosa, Sieb. & Zucc. in Endl. Conif. 65 = T. pisifera.
thujaeformis, R. Smith, ex Gord. Pinet. ed. II. 429.—Hab.?
thurifera, Endl. Synop. Conif. 62 = Cupressus thurifera.
variegata, Hort. ex Carr. Conif. ed. I. 133 = T. sphaeroidalis.

CHAMAECYTISUS, Link, Handb. ii. 154 (1831) = **Cytisus**, Linn. (Legumin.).
austriacus, Link, l. c. 155 = Cytisus austriacus.
biflorus, Link, l. c. 154 = Cytisus biflorus.
capitatus, Link, l. c. 155 = Cytisus capitatus.
elongatus, Link, l. c. = Cytisus biflorus.
hirsutus, Link, l. c. = Cytisus hirsutus.
leucanthus, Link, l. c. 154 = Cytisus austriacus.
proliferus, Link, l. c. = Cytisus proliferus.
purpureus, Link, l. c. = Cytisus purpureus.
supinus, Link, l. c. 155 = Cytisus biflorus.

CHAMAECYTISUS, Vis. Fl. Dalm. iii. 272 (1852) = **Argyrolobium**, Eckl. & Zeyh. (Legumin.).
dalmaticus, Vis. l. c. = A. Linnaeanum.

CHAMAEDACTYLIS, T. Nees, Gen. Fl. Germ. Monocot. i. n. 66 (1840?) = **Aeluropus**, Trin. (Gramin.).
maritima, Nees, l. c. = A. littoralis.

CHAMAEDAPHNE, Mitchell, in Act. Phys. Med. Acad. Nat. Cur. viii. (1748) 222, App. = **Mitchellia**, Linn. (Rubiac.).

CHAMAEDAPHNE, Moench, Meth. 457 (1794) = **Cassandra**, D. Don (Ericac.).
calyculata, Moench, l. c. = Cass. calyculata.
crispa, Spach, Hist. Vég. Phan. ix. 477 = Cass. calyculata.

CHAMAEDOREA, Willd. Sp. Pl. iv. 638 (1805); et in Mém. Acad. Berl. 1804 (1807) 40. *PALMAE*, Benth. & Hook. f. iii. 910.
DASYSTACHYS, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 25.
KUNTHIA, Humb. & Bonpl. Pl. Aequin. ii. 128. t. 122 (1809).
MORENIA, Ruiz & Pav. Prod. 150 (1794).
NUNNEZHARIA, Ruiz & Pav. l. c. 147 (1794).
NUNNEZIA, Willd. Sp. Pl. iv. 1154 (1805).
SPATHOSCAPHE, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 29.
STACHYOPHORBE, Liebm. in Overs. Dansk. Vidensk. Selsk. (1846) 8.
STEPHANOSTACHYS, Klotzsch, ex Oerst. in Kjoeb. Vidensk. Meddel. (1858) 26.
affinis, *Liebm. in Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
alternans, *H. Wendl. in Regel, Gartenfl.* (1880) 104.—Mexic.
amazonica, *J. Linden, in Illustr. Hortic.* xxviii. (1881) 16, nomen.—Bras.
Andreana, *J. Linden, l. c.*, nomen.—N. Granat.
Arenbergiana, *H. Wendl. Ind. Palm.* (1854) 66.—Guatemala?

CHAMAEDOREA:—
atrovirens, Hort. ex H. Wendl. in Kerch. Palm. 239 = Martiana.
atrovirens, *Mart. in Flora*, xxxv. (1852) 721.—Mexic.
aurantiaca, *A. Brongn. in Rev. Hortic.* Sér. III. i. (1847) 86.—Mexic.
Bartlingiana, *H. Wendl. Ind. Palm.* (1854) 60.—N. Granat.
biloba, H. Wendl. l. c. 11 = Bartlingiana.
bifurcata, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 13.—Costa Rica.
brachyclada, *H. Wendl. in Regel, Gartenfl.* (1880) 101.—Panama.
bracteata, *H. Wendl. in Bot. Zeit.* xvii. (1859) 29.—Am. centr.
brevifrons, *H. Wendl. Ind. Palm.* (1854) 61.—N. Granat.
Casperiana, *Klotzsch, in Otto & Dietr. Allg. Gartenz.* xx. (1852) 363.—Guatemala.
cataractarum, *Mart. Hist. Nat. Palm.* iii. 309.—Mexic.
concolor, *Mart. l. c.* 160.—Mexic.
conocarpa, *Mart. Palmet. Orb.* 6.—Bolivia.
costaricana, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 19.—Costa Rica.
Deckeriana, *Hemsl. Biol. Centr. Am. Bot.* iii. 404.—Guatemala.
Deppeana, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 362 = elegans.
desmoncoides, *H. Wendl. in Otto & Dietr. l. c.* xxi. (1853) 177.—Mexic.
elatior, *Mart. in Linnaea*, v. (1830) 205.—Mexic.
elegans, *Mart. l. c.* 204.—Mexic.
elegantissima, *Hort. ex H. Wendl. in Kerch. Palm.* 239, nomen.—Hab. ?
Ernesti-Augusti, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 73.—Mexic.
fenestrata, Hort. ex H. Wendl. Ind. Palm. (1854) 28 = Malortiea gracilis.
fenestrata, *J. Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Mexic.
fibrosa, H. Wendl. Ind. Palm. (1854) 57 = Synechanthus fibrosus.
flavo-virens, *H. Wendl. l. c.* 60.—Hab. ?
flexuosa, Hort. ex H. Wendl. l. c. 14 = Martiana.
formosa, *Hort.—Cf Gard. Chron.* (1876) i. 724.—Hab. ?
fragrans, *Mart. Hist. Nat. Palm.* ii. 4. t. 3. f. 1, 2.—Peruv.
geonomaeformis, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 1.—Guatemala.
geonomoides, *Drude, in Mart. Fl. Bras.* iii. ii. 531.—Peruv.
Ghiesbreghtii, *Hort. ex H. Wendl. in Kerch. Palm.* 239, nomen.—Hab. ?
glauca, *J. Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Venezuela.
glaucifolia, *H. Wendl. Ind. Palm.* (1854) 64.—Mexic.
gracilis, *Willd. Sp. Pl.* iv. 800.—Am. austr.
graminifolia, *H. Wendl. Ind. Palm.* (1854) 62.—Guatemala.
Helleriana, Klotzsch, in Otto & Dietr. Allg. Gartenz. xx. (1852) 362 = elegans.
humilis, *Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
Karwinskyana, H. Wendl. in Otto & Dietr. Allg. Gartenz. xxi. (1853) 179 = elatior.
Klotzschiana, *H. Wendl. Ind. Palm.* (1854) 63.—Mexic.
lanceolata, *Kunth, Enum. Pl.* iii. 172.—Peruv.
latifrons, H. Wendl. Ind. Palm. (1854) 11 = Arenbergiana.
lepidota, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 138.—Mexic.
Liebmanni, *Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
Lindeniana, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 139.—N. Granat.
linearis, *Mart. Hist. Nat. Palm.* ii. 5.—Peruv.
lunata, Liebm. in Mart. l. c. iii. 307 = oblongata.
macrospadix, *Oerst. in Kjoeb. Vidensk Meddel.* (1858) 20.—Costa Rica.
Martiana, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 137.—Mexic.
membranacea, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 22.—Nicaragua.

CHAMAEDOREA:—
mexicana, Hort. Belg. ex Heynh. Nom. ii. 135 = Sartorii.
microphylla, *H. Wendl. in Bot. Zeit.* xvii. (1859) 102.—Panama.
montana, *Liebm. in Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
oblongata, *Mart. l. c.* 160.—Bras.
obovoidea, *H. Wendl. in Kerch. Palm.* 239, nomen.—Hab. ?
oreophila, *Mart. Hist. Nat. Palm.* iii. 309.—Mexic.
Pacaya, *Oerst. in Kjoeb. Vidensk. Meddel.* (1858) 12.—Costa Rica.
paradoxa, *H. Wendl. in Bot. Zeit.* xvii. (1859) 29.—Mexic.
pauciflora, *Mart. Hist. Nat. Palm.* ii. 5. t. 3. f. 3.—Bras.
pinnatifrons, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 14 = gracilis.
pochutlensis, *Liebm. in Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
pulchella, *Linden, Cat.* n. 117, p. 4.—Hab. ?
pygmaea, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xx. (1852) 217, 249.—N. Granat.
radicalis, *Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
regia, H. Wendl. Ind. Palm. (1854) 11 = desmoncoides.
repens, H. Wendl. l. c. = desmoncoides.
resinifera, *H. Wendl. in Otto & Dietr. Allg. Gartenz.* xxi. (1853) 179.—Mexic.
robusta, *Hort. ex H. Wendl. in Kerch. Palm.* 239, nomen.—Hab. ?
Sartorii, *Liebm. in Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
scandens, Liebm. in Mart. l. c. = elatior.
Schiedeana, *Mart. in Linnaea*, v. (1830) 204.—Mexic.
simplicifrons, Hort. Belg. ex Heynh. Nom. ii. 135 = Ernesti-Augusti.
speciosa, H. Wendl. Ind. Palm. (1854) 15 = Schiedeana.
spectabilis, *Hort. ex H. Wendl. in Kerch. Palm.* 240, nomen.—Hab. ?
tenella, *H. Wendl. in Regel, Gartenfl.* (1880) 102.—Mexic.
Tepejilote, *Liebm. in Mart. Hist. Nat. Palm.* iii. 308.—Mexic.
velutina, Hort ex H. Wendl. Ind. Palm. (1854) 13 = lepidota.
Verschaffelti, *Hort. ex H. Wendl. in Kerch. Palm.* 240, nomen.—Hab. ?
Wallisi, *J. Linden, in Illustr. Hortic.* xxviii. (1881) 16, nomen.—Ecuador.
Warscewiczii, *H. Wendl. in Bonplandia*, x. (1862) 37.—Guatemala.
Wendlandi, *H. Wendl. in Kerch. Palm.* 240.—Mexic.
Wendlandiana, Hemsl. Biol. Centr. Am. Bot. iii. 407 = praec.
Wobstiana, *Linden, Cat.* n. 117, p. 4.—Hab. ?

CHAMAEDRYS, Moench, Meth. 383 (1794) = **Teucrium**, Tourn. (Labiat.).
Botrys, Moench, l. c. = T. Chamaedrys.
capitata, Rafin. Fl. Tellur. iii. 85 = T. Polium.
cinerascens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 47 = T. flavum.
cretica, Rafin. Fl. Tellur. iii. 85 = T. creticum.
flava, Moench, Meth. 383 = T. flavum.
glauca, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 48 = T. flavum.
laciniata, S. F. Gray, Nat. Arr. Brit. Pl. ii. 369 = T. Botrys.
latifolia, Rafin. Fl. Tellur. iii. 85 = T. fruticans.
lucida, Moench, Meth. 383 = T. lucidum.
Marum, Moench, l. c. 384 = T. Marum.
montana, Rafin. Fl. Tellur. iii. 85 = T. montanum.
Nissoliana, Rafin. l. c. = T. Pseudochamaepitys.
officinalis, Moench, Meth. 383 = T. Chamaedrys.
ovalifolia, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 48 = T. flavum.
palustris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 369 = T. Scordium.
Polium, Rafin. Fl. Tellur. iii. 85 = T. Polium.
pulverulenta, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 47 = T. flavum.
pumila, Rafin. Fl. Tellur. iii. 85 = T. pumilum.
quadrula, Rafin. l. c. = T. quadratulum ?
Scordium, Moench, Meth. 384 = T. Scordium.

CHAMAEDRYS:—
tetonicifolia, Rafin. Fl. Tellur. iii. 85 = Teucrium betonicum.
virescens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 47 = T. flavum.

CHAMAEFISTULA, G. Don, Gen. Syst. ii. 451 (1832) = **Cassia**, Tourn. (Legumin.).
astroites, G. Don, l. c. = Cass. villosa.
bacillaris, G. Don, l. c. = Cass. Sophera.
chinensis, G. Don. l. c. 452 = Cass. Sophera.
contorta, G. Don, l. c. = Cass. Tora.
coromandelina, G. Don, l. c. = Cass. Sophera.
corymbosa, G. Don, l. c. 451 = Cass. corymbosa.
crotalarioides, G. Don, l. c. 452 = Cass. crotalarioides.
elegans, G. Don, l. c. 451 = Cass. obliqua.
excelsa, G. Don, l. c. 452 = Cass. excelsa.
floribunda, G. Don, l. c. = Cass. laevigata.
gigantea, G. Don, l. c. = Cass. siamea.
Herbertiana, G. Don, l. c. = Cass. laevigata.
inaequilatera, G. Don, l. c. 451 = Cass. inacquilatera.
indecora, G. Don, l. c. 452 = Cass. bicapsularis.
inflata, G. Don, l. c. 451 = Cass. bicapsularis.
laevigata, G. Don, l. c. 452 = Cass. laevigata.
macrophylla, G. Don, l. c. 451 = Cass. macrophylla.
melanocarpa, G. Don, l. c. = Cass. viminea.
obliqua, G. Don, l. c. = Cass. obliqua.
oxyphylla, G. Don, l. c. = Cass. oxyphylla.
pendula, G. Don, l. c. 452 = Cass. bicapsularis.
prominens, G. Don, l. c. 451 = Cass. macranthera.
puberula, G. Don, l. c. = Cass. bacillaris.
Ruiziana, G. Don, l. c. = Cass. Ruiziana.
Sophera, G. Don, l. c. 452 = Cass. Sophera.
speciosa, G. Don, l. c. 451 = Cass. speciosa.
torosa, G. Don, l. c. 452 = Cass. Sophera.

CHAMAEGERON, Schrenk. in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 107. *COMPOSITAE.*
oligocephalum, *Schrenk. l. c.*—Soongaria.

CHAMAEIRIS, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 417 = **Iris**, Tourn.
angustifolia, Medic. l. c. 418 = I. pumila.
desertorum, Medic. l. c. = I. Gueldenstaedtiana.
foetida, Medic. l. c. = I. foetida.
graminea, Medic. l. c. = I. graminea.
spuria, Medic. l. c. = I. spuria.

CHAMAELAUCIUM, Desf. in Mém. Mus. Par. v. (1819) 39. t. 3. *MYRTACEAE*, Benth. & Hook. f. i. 698.
DECALOPHIUM, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 153.
affine, Meissn. in Journ. Linn. Soc. i. (1857) 45 = uncinatum.
axillare, *F. Muell. ex Benth. Fl. Austral.* iii. 38.—Austral.
brevifolium, *Benth. l. c.* 37.—Austral.
Brownii, Desf. in Mém. Mus. Par. v. (1819) 271. t. 19 = Verticordia Brownii.
ciliatum, *Desf. l. c.* 40.—Austral.
Drummondii, *Meissn. in Journ. Linn. Soc.* i. (1857) 44.—Austral.
gracile, *F. Muell. Fragm.* iv. 62.—Austral.
heterandrum, *Benth. Fl. Austral.* iii. 36.—Austral.
megalopetalum, *F. Muell. ex Benth. l. c.* 38.—Austral.
pauciflorum, *Benth. l. c.*—Austral.
plumosum, Desf. in Mém. Mus. Par. v. (1819) 42. t. 4 = Verticordia Fontanesii.
Thomasii, F. Muell. Fragm. iv. 137. t. 30 = Darwinia Thomasii.
uncinatum, *Schau. in Lehm. Pl. Preiss.* i. 97.—Austral.
verticordinum, F. Muell. Fragm. iv. 57 = Darwinia verticordina.
virgatum, *Endl. in Ann. Wien. Mus.* ii. (1840) 198.—Austral.

CHAMAELE, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 59 (1867). *UMBELLIFERAE.*
Tanakae, Franch. & Sav. Enum. Pl. Jap. i. 185 = Carum Tanakae.
tenera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 60.—Japon.

CHAMAELEA, [Tourn.] Adans. Fam. ii. 345 (1763) = **Cneorum**, Linn. (Simarub.).
tricoccos, Lam. Fl. Fr. ii. 682 = Cneorum tricoccum.

CHAMAELEDON, Link, Enum. Hort. Berol. i. 210 (1821) = **Loiseleuria**, Desv. (Ericac.).
procumbens, Link, l. c. 211 = L. procumbens.

CHAMAELEON, Cass. in Dict. Sc. Nat. xlvii. 509 (1827) = **Atractylis**, Linn. (Compos.).
caulescens, Cass. l. c. l. 60 = A. macrophylla.
gummifer, Cass. l. c. xlvii. 509 = A. gummifera.
megacephalus, Cass. l. c. l. 59 = A. macrocephala.

CHAMAELINUM, Host, Fl. Austr. ii. 224 (1831) = **Camelina**, Crantz (Crucif.).
austriacum, Host, l. c. 225 = Nasturtium austriacum.
paniculatum, Host, l. c. 224 = Neslia paniculata.
sativum, Host, l. c. 225 = Cam. sativa.

CHAMAELIRIUM, Willd. in Ges. Nat. Fr. Berl. Mag. ii. (1808) 18. *LILIACEAE*, Benth. & Hook. f. iii. 826.
DASURUS, Salisb. Gen. Pl. Fragm. 51 (1865).
DICLINOTRYS, Rafin. Neogen. 3 (1825).
OPHIOSTACHYS, [Delile] in Red. Lil. t. 464 (1816).
SIRAITOS, Rafin. Fl. Tellur. iv. 26 (1836).
carolinianum, *Willd. l. c.* 19.—Am. bor.
luteum, A. Gray, Man. Bot. N. U. States, ed. V. 527 = carolinianum.
luteum, Miq. Ann. Mus. Bot. Lugd. Bat. 144 = Chionographis japonica.

CHAMAELUM, Baker, in Journ. Linn. Soc. xvi. (1877) 120, sphalm. = **Chamelum**, Phil. (Irid.).

CHAMAEMELES, Lindl. in Trans. Linn. Soc. xiii. (1822) 104. t. 11. *ROSACEAE*, Benth. & Hook. f. i. 628.
coriacea, *Lindl. l. c.*—Ins. Madera.
mexicana, Baill. Adansonia, ix. (1869) 148 = Photinia mexicana.

CHAMAEMELUM, Tourn. ex Adans. Fam. ii. 128 (1763); Cass. in Dict. Sc. Nat. xxix. 179, 185 (1823) = **Anthemis**, Linn. (Compos.).
alpinum, All. Fl. Pedem. i. 186 = Achillea oxyloba.
arvense, Schreb. Spicil. 18 = A. arvensis.
austriacum, J. & C. Presl, Fl. Cech. 173 = A. austriaca.
Cota, All. Fl. Pedem. i. 184 = A. Cota.
Cota, C. A. Mey. Verz. Pfl. Cauc. 75 = A. altissima.
Cotula, All. Fl. Pedem. i. 186 = A. Cotula.
discoideum, All. l. c. 188 = A. tinctoria.
foetidum, Baumg. Enum. Stirp. Transs. iii. 144 = A. Cotula.
incrassatum, Hoffmgg. Fl. Port. ii. 348, ex DC. Prod. vi. 616 = Anacyclus clavatus.
maritimum, All. Fl. Pedem. i. 184 = A. maritima.
Marschallianum, C. A. Mey. Verz. Pfl. Cauc. 75 = A. Biebersteiniana.
mixtum, All. Fl. Pedem. i. 185 = A. mixta.
montanum, All. l. c. 187 = A. montana.
nobile, All. l. c. 185 = A. nobilis.
orientale, Schult. ex Steud. Nom. ed. II. i. 343 = A. montana.
pinnatifidum, Moench, Meth. 605 = A. mixta.
proliferum, Moench, l. c. Suppl. 257 = Cladanthus proliferus.
tinctorium, Schreb. Spicil. 145 = A. tinctoria.
tomentosum, All. Fl. Pedem. i. 184 = Anacyclus clavatus.
Triumfetti, All. l. c. 187 = A. austriaca.
valentinum, All. l. c. = Anacyclus valentinus.

CHAMAEMELUM, Vis. Osserv. Matric. 12 (1845); Fl. Dalm. ii. 84 (1847) = **Matricaria**, Linn. (Compos.).
ambiguum, Boiss. Diagn. Ser. I. xi. 20 (= *Pyrethrum ambiguum*).—Oriens.
arvense, Reichb. f. ex Willk. in Bot. Zeit. xxii. (1864) 251 = M. glabra.
auriculatum, Boiss. Diagn. Ser. I. xi. 23.—Arabia.
callosum, Boiss. & Heldr. l. c. 22.—As. Min.
caucasicum, Boiss. l. c. 20 = M. grandiflora.
conoclinium, Boiss. & Bal. l. c. Ser. II. vi. 98 = M. conoclinia.
daghestanicum, Boiss. Fl. Orient. iii. 334.—Daghestan.
decipiens, Boiss. Diagn. Ser. I. xi. 22 = M. decipiens.
disciforme, Vis. Fl. Dalm. ii. 85 = M. corymbifera.
elongatum, Boiss. Diagn. Ser. I. xi. 20 (= *Pyrethrum elongatum*).—Oriens.

CHAMAEMELUM :—
grandiflorum, Boiss. & Haussk. in Boiss. Fl. Orient. iii. 331.—Mesopotam.
heterocarpum, Boiss. l. c. 329 = Matricaria heterocarpa.
inodorum, Vis. Osserv. Matric. 13; Fl. Dalm. ii. 85 = M. inodora.
Kochii, Boiss. Fl. Orient. iii. 329 = M. microcephala.
Kotschyi, Boiss. Diagn. Ser. II. iii. 27 = M. Oreades.
lamellatum, Boiss. Fl. Orient. iii. 326 = M. lamellata.
limosum, Maxim. Prim. Fl. Amur. 156 = M. inodora.
maritimum, Boiss. Diagn. Ser. I. xi. 21 = M. maritima.
melanolepis, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 118 = M. Oreades.
microcephalum, Boiss. Diagn. Ser. I. xi. 23 = M. microcephala.
monticolum, Boiss. & Huet, l. c. Ser. II. iii. 27.—Armenia.
nanum, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xiv. (1856) 231.—Kurdistan.
olympicum, Boiss. ex Tchihat. Asie Min. Bot. ii. 258, nomen.—As. Min.
Oreades, Boiss. Diagn. Ser. I. xi. 21 = M. Oreades.
Pichleri, Boiss. Fl. Orient. iii. 332.—Bithynia.
praecox, Vis. Osserv. Matric. 13; Fl. Dalm. ii. 86 = M. praecox.
Reuterianum, Boiss. ex Tchihat. Asie Min. Bot. ii. 258, nomen.—As. Min.
rosellum, Boiss. & Orph. in Boiss. Fl. Orient. iii. 334 = M. rosella.
Szovitsii, Boiss. l. c. 332.—Reg. Caucas.
Tchihatchewi, Boiss. in Tchihat. Asie Min. Bot. ii. 256.—As. Min.
tetragonospermum, F. Schmidt, Reisen Amur. 148.—Reg. Amur.
trichophyllum, Boiss. Diagn. Ser. I. xi. 21 = M. trichophylla.
uniglandulosum, Vis. Osserv. Matric. 13; Fl. Dalm. ii. 85 = M. trichophylla.

CHAMAEMESPILUS, Medic. Phil. Bot. i. 138, 155 (1789) = **Pyrus**, Linn. (Rosac.).
ambigua, Gren. ex Nym. Consp. 242 = P. Chamaemespilus.
Aria, M. Roem. Syn. Rosifl. 132 = P. Aria.
humilis, M. Roem. l. c. 131 = P. Chamaemespilus.
Pseudaria, M. Roem. l. c. 132 = P. Aria.
sudetica, M. Roem. l. c. 131 = P. sudetica.

CHAMAEMYRRHIS, Endl. ex Heynh. Nom. ii. 136 (1846) = **Caldasia**, Lag. (Umbellif.).
eriopoda, Endl. l. c. = Cald. eriopoda.

CHAMAENERION, [Tourn.] Adans. Fam. ii. 85 (1763) = **Epilobium**, Linn. (Onagrar.).
americanum, Sweet, Hort. Brit. ed. II. 198 = E. palustre.
angustifolium, Schur, Enum. Pl. Transs. 213 = E. angustifolium.
angustifolium, Scop. Fl. Carn. ed. II. i. 271 = E. Dodonaei.
denticulatum, Schur, Enum. Pl. Transs. 214 = E. angustifolium.
Dodonaei, Schur, ex Fuss. Fl. Transs. 214 = E. Dodonaei.
grandiflorum, Moench, Meth. 677 = E. hirsutum.
halimifolium, Salisb. Parad. Lond. t. 58 = E. latifolium.
Halleri, Sweet, Hort. Brit. ed. II. 198 = E. Dodonaei.
hirsutum, Scop. Fl. Carn. ed. II. i. 270 = E. hirsutum.
latifolium, Sweet, Hort. Brit. ed. II. 198 = E. latifolium.
montanum, Scop. Fl. Carn. ed. II. i. 270 = E. montanum.
obscurum, Schreb. Spicil. 147 = E. tetragonum.
palustre, Schreb. l. c. 22 = E. palustre.
parviflorum, Schreb. l. c. 146 = E. parviflorum.
pubescens, Moench, Meth. 677 = E. parviflorum.
ramosissimum, Moench, l. c. 678 = E. tetragonum.
roseum, Schreb. Spicil. 147 = E. roseum.
rosmarinifolium, Moench, Meth. Suppl. 288 = E. Dodonaei.
speciosum, Lodd. ex Steud. Nom. ed. II. i. 343 = E. angustifolium.
spicatum, S. F. Gray. Nat. Arr. Brit. Pl. ii. 559 = E. angustifolium.
tetragonum, Scop. Fl. Carn. ed. II. i. 271 = E. tetragonum.

CHAMAENERIUM, Spach, Hist. Vég. Phan. iv. 396 (1835) = praec.

CHAMAEORCHIS, Koch, Syn. Fl. Germ. 692 (1837) = CHAMORCHIS, Rich. = **Herminium**, Willd. (Orchid.).

CHAMAEPEUCE, DC. Prod. vi. 657 (1837) = **Cnicus**, Tourn. (Compos.).
afra, DC. l. c. 659 = Cn. afer.
Alpini, Jaub. & Spach, Illustr. v. 26. t. 425 = Cn. Chamaepeuce.
atropurpurea, Boiss. & Heldr. Diagn. Ser. II. iii. 47 = Cn. strictus.
Casabonae, DC. Prod. vi. 658 = Cn. Casabonae.
cynaroides, DC. l. c. 659 = Cn. cynaroides.
diacantha, DC. l. c. = Cn. diacanthus.
echinocephala, DC. l. c. 660 = Cn. echinocephalus.
firma, DC. l. c. 659 = Cn. firmus.
fruticosa, DC. l. c. 658 = Cn. fruticosus.
gnaphalodes, DC. l. c. = Cn. gnaphalodes.
hispanica, DC. l. c. = Cn. hispanicus.
horrida, DC. l. c. 660 = Cn. horridus.
imbricata, Rafin. Fl. Tellur. iv. 118 = Cn. Chamaepeuce.
macrantha, Schrank, in Bull. Sc. Acad. Pétersb. x. (1842) 354 = Cousinia Hablizii.
macrostachya, Trautv. in Bull. Phys.-Math. Acad. Pétersb. xvi. (1858) 326.—Reg. Caucas.
mutica, DC. Prod. vi. 657 = Cn. Chamaepeuce.
nivea, DC. Prod. vi. 659 = Cn. niveus.
polyacantha, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 144 = Cn. Casabonae.
polycephala, DC. Prod. vi. 658 = Cn. Chamaepeuce.
sinuata, Trautv. in Act. Hort. Petrop. ii. (1873) 477 = Cn. cynaroides.
stellata, DC. Prod. vi. 658 = Cn. stellatus.
stricta, DC. l. c. 659 = Cn. strictus.

CHAMAEPEUCE, Zucc. in Endl. Enchir. 139 (1841) = **Thuya**, Tourn. (Conif.).
obtusa, Zucc. ex Gord. Pinet. Suppl. 93 = T. obtusa.

CHAMAEPHYTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 371 = **Potentilla**, Linn. (Rosac.).
supinum, Fourr. l. c. = P. supina.

CHAMAEPITYS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 220 (1745) = **Ajuga**, Tourn. (Labiat.).
arvensis, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 60 = A. Chamaepitys.
Iva, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 138 = A. Iva.
pseudoiva, Fourr. l. c. = A. Iva.
trifida, Dum. Fl. Belg. 42 = A. Chamaepitys.
vulgaris, Link, Handb. i. 453 = A. Chamaepitys.

CHAMAEPLIUM, Wallr. Sched. Crit. i. 376 (1822) = **Sisymbrium**, Tourn. (Crucifer.).
officinale, Wallr. l. c. 377 = S. officinale.
polyceratium, Wallr. l. c. = S. polyceratium.
supinum, Wallr. l. c. = S. supinum.

CHAMAERANTHEMUM, Nees, in Lindl. Nat. Syst. ed. II. 445 (1836). *ACANTHACEAE*, Benth. & Hook. f. ii. 1095.
Beyrichii, *Nees, l. c.* 445.—Bras.
Gaudichaudii, *Nees, in Mart. Fl. Bras.* ix. 155.—Bras.
igneum, *Regel, Gartenfl.* (1868) 353.—Peruv.
pictum, *Mast. in Gard. Chron.* (1878) i. 527.—Bras.

CHAMAERAPHIS, R. Br. Prod. 193 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1107.
abortiva, Poir. Encyc. Suppl. ii. 189 = spinescens.
aspera, *Nees, in Wall. Cat.* n. 8679 = spinescens.
compressa, Poir. ex Steud. Nom. ed. I. 183 = Pennisetum compressum.
depauperata, Nees, ex Steud. Syn. Pl. Gram. 49 = Panicum asperum, *Wight*.
gracilis, *Hack. in Engl. Bot. Jahrb.* vi. (1885) 236.—Malaya.
hordeacea, *R. Br. Prod.* i. 194.—Austral.
paradoxa, *Poir. Encyc. Suppl.* ii. 189.—Austral.
parvigluma, *Munro, ex Wright, in Sauv. Fl. Cub.* 200, nomen.—Cuba.

CHAMAERAPHIS:—
spineseens, *Poir. Encyc. Suppl.* ii. 189.—As. trop.; Austral.
spinosa, Beauv. ex Roem. & Schult. Mant. ii. 253 = spinescens.

CHAMAEREPES, Spreng. Syst. iii. 702 (1826) = **Herminium**, Willd. (Orchid.).
alpina, Spreng. l. c. = H. alpinum.

CHAMAERHODODENDROS, Tourn. ex Hall. in Rupp. Fl. Jen. ed. Hall. 249 (1745) = **Rhododendron**, Linn. (Ericac.).

CHAMAERHODOS, Bunge, in Ledeb. Fl. Alt. i. 429 (1829). *ROSACEAE*, Benth. & Hook. f. i. 621.
altaica, *Bunge, l. c.*—Sibir. alt. occ.
erecta, *Bunge, l. c.* 430.—Sibir.; Am. bor. occ.
grandiflora, *Ledeb. l. c.* 431, in nota.—Sibir.
mongholica, *Bunge, in Ann. Sc. Nat.* Sér. II. xix. (1843) 177.—Mongolia.
polygyna, G. Don, Gen. Syst. ii. 563 = erecta.
sabulosa, *Bunge, in Ledeb. Fl. Alt.* i. 431.—Sibir.
trifida, *Ledeb. Fl. Ross.* ii. 34.—Sibir.

CHAMAERIPHES, Ponted. ex Gaertn. Fruct. i. 25 (1789) = seq.

CHAMAEROPS, Linn. Mus. Cliff. 10; Gen. ed. I. 354 (1737). *PALMAE*, Benth. & Hook. f. iii. 924.
CHAMAERIPHES, Ponted. ex Gaertn. Fruct. i. 25 (1789).
acaulis, Michx. Fl. Bor. Am. ii. 207 = Sabal Adansoni.
Antillarum, Desc. Antill. i. t. 28 (Thrinacis sp. ?).—Ind. occ.
arborescens, Steud. Nom. ed. I. 183 = humilis.
arundinacea, Sm. in Rees, Cycl. vii. n. 3 = Rhapidophyllum Hystrix.
Biroo, Siebold, in Mart. Hist. Nat. Palm. 252 = Livistona rotundifolia.
cochinchinensis, Lour. Fl. Cochinch. ii. 657 = Rhapis cochinchinensis.
conduplicata, Kickx, in Bull. Acad. Brux. v. (1838) 61 = humilis.
excelsa, Thunb. Fl. Jap. 139 = Trachycarpus excelsus.
excelsior, *Boj. Hort. Maurit.* 307.—Ins. Maurit.
Fortunei, Hook. Bot. Mag. t. 5221 = Trachycarpus Fortunei.
Ghiesbreghti, Hort. ex H. Wendl. in Kerch. Palm. 240 (= Sabal sp.).—Hab. ?
glabra, Mill. Gard. Dict. ed. VIII. n. 2 = Sabal Adansoni.
gracilis, *Lodd. Cat.* (1823) *ex Schult. f. Syst.* vii. 1489, nomen.—Am. austr.
Griffithiana, Wall. ex Voigt, Hort. Suburb. Calc. 641, nomen = seq.
Griffithii, Lodd. Cat. (1841) ex Verlot, in Rev. Hortic. (1870) t. 276 = Trachycarpus khasyanus.
guianensis, *Lodd. Cat. ex Loud. Hort. Brit.* 413, nomen.—Guiana.
humilis, *Linn. Sp. Pl.* 1187.—Reg. Mediterr. occ.
Histrix, Desf. ex Steud. Nom. ed. I. 183 = Corypha Hystrix.
Hystrix, Fras. ex Pursh, Fl. Am. Sept. i. 240 = Rhapidophyllum Hystrix.
khasyana, Griff. in Calc. Journ. Nat. Hist. v. (1845) 341 = Trachycarpus khasyanus.
Kwanwortsik, Siebold, ex H. Wendl. Ind. Palm. (1854) 34 = Rhapis flabelliformis.
louisiana, Rafin. Fl. Ludov. 159 = Rhapidophyllum Hystrix?
macrocarpa, *Linden, Cat.* n. 87 (1871).—Hab. ?
macrocarpa, Tin. in Guss. Fl. Sic. Syn. ii. 883 = humilis.
Martiana, Wall. Pl. As. Rar. iii. 5. t. 211 = Trachycarpus Martianus.
mitis, *J. Mey. in Mém. Acad. Berl.* (1769) 29.—Hab. ?
Mocini, H. B. & K. Nov. Gen. et Sp. i. 300 = Acanthorhiza Mocinni.
nepalensis, Lodd. Cat. (1823) ex Schult. f. Syst. vii. 1489 = Trachycarpus Martianus.
Palmetto, Michx. Fl. Bor. Am. i. 206 = Sabal Palmetto.
reclinata, *Juss. ex Schult. f. Syst.* vii. 1489, nomen.—Hab. ?
Ritchiana, Griff. in Calc. Journ. Nat. Hist. v. (1845) 342 = Nannorrhops Ritchieana.
serrulata, Michx. Fl. Bor. Am. i. 239 = Brahea serrulata.

CHAMAEROPS:—
Sirotsik, Hort. ex H. Wendl. Ind. Palm. (1854) 17 = Rhapis humilis.
stauracantha, Hort. Belg. ex Heynh. Nom. ii. 136 = Acanthorhiza aculeata.
tomentosa, *Fulchiron, ex Schult. f. Syst.* vii. 1489, nomen.—Hab. ?
tomentosa, *Ch. Morr. in Ann. Gand*, i. (1845) 488.—Hab. ?

CHAMAESARACHA, A. Gray, ex Franch. & Sav. Enum. Pl. Jap. ii. 454 (1876). *SOLANACEAE*, Benth. & Hook. f. ii. 891.
conoides, *Benth. & Hook. f. Gen.* ii. 891.—Mexic.
Coronopus, *A. Gray, in Bot. Calif.* i. 540.—Calif.
japonica, *Franch. & Sav. Enum. Pl. Jap.* ii. 454.—Japon.
nana, *A. Gray, in Bot. Calif.* i. 540.—Calif.
physaloides, Greene, in Bull. Torrey Club, ix. (1882) 122 = Physalis lobata.
sordida, *A. Gray, Synop. Fl. N. Am.* ii. 1. 232.—Texas; Mexic.

CHAMAESARACHIA, Franch. & Sav. Enum. Pl. Jap. ii. 454 (1876) = praec.

CHAMAESCIADIUM, C. A. Mey. Verz. Pfl. Cauc. 122 (1831). *UMBELLIFERAE*, Benth. & Hook. f. i. 895.
acaule, *C. A. Mey. l. c.*—As. Min.; Reg. Caucas.
albiflorum, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 360.—Sibir.
flavescens, C. A. Mey. Verz. Pfl. Cauc. 122 = acaule.
gracile, Boiss. & Heldr. in Boiss. Diagn. Ser. II. ii. 79 = Carum rupestre.
Heldreichii, Boiss. l. c. 78 = Carum Heldreichii.

CHAMAESCILLA, F. Muell. Fragm. vii. 68 (1870). *LILIACEAE*, Benth. & Hook. f. iii. 791.
corymbosa, *F. Muell. l. c.*—Austral.
spiralis, *F. Muell. l. c.*—Austral.

CHAMAESPARTIUM, Adans. Fam. ii. 321 (1763) = **Cytisus**, Linn. et **Genista**, Linn. (Legumin.).

CHAMAESPARTON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 357 = **Genista**, Linn. (Legumin.).
pilosum, Fourr. l. c. = G. pilosa.

CHAMAESPHACOS, Schrenk, Enum. Pl. Nov. i. 27 (1841). *LABIATAE*, Benth. & Hook. f. ii. 1207.
ilicifolius, *Schrenk, l. c.*—Sibir.

CHAMAESPHAERION, A. Gray, in Hook. Kew Journ. iii. (1851) 176 = **Chthonocephalus**, Steetz (Compos.).
pygmaeum, A. Gray, l. c. 177 = Chthonocephalus pygmaeus.

CHAMAESPILUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 378 = **Pyrus**, Linn. (Rosac.).
sudetica, Fourr. l. c. = P. sudetica.

CHAMAESTEPHANUM, Willd. in Ges. Nat. Fr. Berl. Mag. i. (1807) 140 = **Schkuhria**, Roth (Compos.).

CHAMAESYCE, S. F. Gray, Nat. Arr. Brit. Pl. ii. 260 (1821) = **Euphorbia**, Linn.
maritima, S. F. Gray, l. c. = E. Peplis.

CHAMAEZELUM, Link, Handb. i. 719 (1829) = **Antennaria**, R. Br. (Compos.).
alpinum, Link, l. c. = A. alpina.
carpaticum, Link, l. c. = A. carpathica.
dioicum, Link, l. c. = A. dioica.
margaritaceum, Link. l. c. = Anaphalis margaritacea.

CHAMAEXEROS, Benth. Fl. Austral. vii. 110 (1878). *JUNCACEAE*, Benth. & Hook. f. iii. 864.
fimbriata, *Benth. l. c.* 111.—Austral.
Serra, *Benth. l. c.* 110.—Austral.

CHAMAEXIPHIUM, Hochst. in Flora, xviii. (1844) 102 = **Ficinia**, Schrad. (Cyperac.).
clandestinum, Hochst. l. c. = F. clandestina.
Dregeanum, Steud. Syn. Pl. Cyp. 59 = F. laevis.

CHAMAGROSTIS, Borkh. ex Wibel, Prim. Fl. Werth. 126 (1799) = **Mibora**, Adans. (Gramin.).
minima, Borkh. l. c. = M. verna.

CHAMALIUM, Juss. in Ann. Mus. Par. vi. (1805) 324 = **Cardopatium**, Juss. (Compos.).

CHAMALIUM, Cass. in Dict. Sc. Nat. xlvii. (1827) 498, 599 = CHAMAELEON, Cass. = **Atractylis**, Linn. (Compos.).

CHAMAREA, Eckl. & Zeyh. Enum. 346 (1837) = **Carum**, Linn. (Umbellif.).
caffra, Eckl. & Zeyh. l. c.—Afr. austr.
capensis, Eckl. & Zeyh. l. c. = Carum capense.

CHAMBEYRONIA, Vieill. in Bull. Soc. Linn. Normand. Sér. II. vi. 1872 (1873) 229. *PALMAE*.
Brébissonii, *Vieill. l. c.* 230, nomen.—N. Caled.
macrocarpa, *Vieill. l. c.* nomen.—N. Caled.
Morieri, *Vieill. l. c.* nomen.—N. Caled.

CHAMEDRYS, Rafin. Fl. Tellur. iii. 85 (1836) = CHAMAEDRYS, Tourn. = **Teucrium**, Tourn. (Scrophul.).

CHAMELUM, Phil. in Linnaea, xxxiii. (1864-65) 250. *IRIDEAE*, Benth. & Hook. f. iii. 700.
CHAMAELUM, Baker, in Journ. Linn. Soc. xvi. (1877) 120, sphalm.
andinum, *Benth. & Hook. f. Gen.* iii. 700.—Chili.
luteum, *Phil. in Linnaea*, xxxiii. (1864-65) 250.—Chili.

CHAMIRA, Thunb. Nov. Gen. ii. 48 (1782). *CRUCIFERAE*, Benth. & Hook. f. i. 81.
cornuta, *Thunb. l. c.* 49.—Afr. austr.

CHAMISME, Rafin. ex Steud. Nom. ed. II. i. 776 (in syn.) (1840) = **Houstonia**, Gron. (Rubiac.).

CHAMISSOA, H. B. & K. Nov. Gen. et Sp. ii. 158. t. 125 (1817). *AMARANTACEAE*, Benth. & Hook. f. iii. 27.
KOKERA, Adans. Fam. ii. 269 (1763).
acuminata, *Mart. in Nov. Act. Nat. Cur.* xiii. (1826) 286.—Bras.
albida, Mart. l. c. 287 = Allmania albida.
altissima, *H. B. & K. Nov. Gen. et Sp.* 197. t. 125.—Am. austr.
altissima, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 32 = Maximiliani.
angustifolia, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iv. 717 = Allmania nodiflora.
arabica, Spreng. Syst. i. 815 = Digera arvensis.
aspera, Wight, Ic. v. t. 1772 = Allmania nodiflora.
Blanchetii, *Moq. in DC. Prod.* xiii. II. 251.—Bras.
Brownii, Steud. Nom. ed. II. i. 344 = Allmania nodiflora.
celosioides, *Griseb. in Goett. Abh.* xix. (1874) 79.—Reg. Argent.
ciliata, Spreng. Syst. i. 815 = Digera arvensis.
commutata, Spreng. l. c. = Digera arvensis.
dichotoma, Moq. in DC. Prod. xiii. II. 249 = Allmania nodiflora.
esculenta, Moq. l. c. = Allmania nodiflora.
javanica, Hassk. Pl. Jav. Rar. 434 = Allmania nodiflora.
macrocarpa, *H. B. & K. Nov. Gen. et Sp.* ii. 197.—Am. austr.
Martii, Moq. in DC. Prod. xiii. II. 252 = altissima.
Maximiliani, *Mart. ex Moq. l. c.* 251.—Bras.
muricata, Spreng. Syst. i. 815 = Digera arvensis.
nodiflora, Mart. in Nov. Act. Nat. Cur. xiii. (1826) 287 = Allmania nodiflora.
ovata, Endl. ex Moq. in DC. Prod. xiii. II. 322 = Charpentiera ovata.
paniculata, Moq. ex Moq. l. c. 239 = Celosia Moquini.
populifolia, Hochst. ex Moq. l. c. = Celosia populifolia.
pyramidalis, Moq. l. c. 248 = Allmania albida.

CHAMISSONIA, Endl. Gen. 1190, in syn. (1840) = CAMISSONIA, Link = **Oenothera**, Linn. (Onagrar.).

CHAMITEA, Kern. in Verh. Zool.-Bot. Ges. Wien, x. (1860) 275 = **Salix**, Tourn.
reticulata, Kern. l. c. 277 = S. reticulata.

CHAMITIS, Banks, ex Gaertn. Fruct. i. 94. t. 22 (1788) = **Azorella**, Lam. (Umbell.).
complicata, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 285 = A. caespitosa.
integrifolia, Gaertn. Fruct. i. 94. t. 22. f. 4 = A. filamentosa.
tricuspidata, Gaertn. l. c. t. 22 = A. trifurcata.
trifurcata, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 283 = A. trifurcata.
trifurcata, Gaertn. Fruct. i. 95 = A. caespitosa.

CHAMOLETTA, Adans. Fam. ii. 60 (1763) = **Iris**, Linn.

CHAMOMILLA, [Hall.] S. F. Gray, Nat. Arr. Brit. Pl. ii. 454 (1821); C. Koch, in Linnaea, xvii. (1843) 45 = **Matricaria**, Linn. (Compos.).
brachyglossa, J. Gay, ex Benth. & Hook. f. Gen. ii. 428 = M. brachyglossa.
Courrantiana, C. Koch, in Linnaea, xxiv. (1851) 338, in obs. = M. Chamomilla.
discoidea, J. Gay, ex A. Br. in Bot. Zeit. x. (1852) 650 = M. discoidea.
glabrata, Sch.-Bip. Ueber Tanacet. 21 = M. glabrata.
inodora, C. Koch, in Linnaea, xvii. (1843) 45 = M. inodora.
meridionalis, C. Koch, l. c. = M. Chamomilla.
officinalis, C. Koch, l. c. = M. Chamomilla.
patens, Gilib. Fl. Lituan. iii. 220 = M. Chamomilla.
praecox, C. Koch, in Linnaea, xvii. (1843) 46 = M. inodora.
unilateralis, C. Koch, l. c. xxiv. (1851) 338.—Reg. Caucas.
vulgaris, S. F. Gray, Nat. Arr. Brit. Pl. ii. 454 = M. Chamomilla.

CHAMOMILLA, Godr. Fl. Lorr. ii. 19 (1843) = **Anthemis**, Linn. (Compos.).
aurea, J. Gay, in Bull. Soc. Bot. Fr. iv. (1857) 278 = Cotula aurea.
fuscata, Gren. & Godr. Fl. Fr. ii. 151 = A. fuscata.
mixta, Gren. & Godr. l. c. = A. mixta.
nobilis, Godr. Fl. Lorr. ii. 19 = A. nobilis.

CHAMORCHIS, Rich. in Mém. Mus. Par. iv. (1818) 49 = **Herminium**, Linn. (Orchid.).
albida, Dum. Fl. Belg. 133 = Habenaria albida.
alpina, Rich. in Mém. Mus. Par. iv. (1818) 57 = Herminium alpinum.
viridis, Dum. Fl. Belg. 133 = Habenaria viridis.

CHAMPACA, Adans. Fam. ii. 365 (1763) = **Michelia**, Linn. (Magnoliac.).
fasciculata, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 13.—Malaya.
Michelia, Noronha, l. c. 11 = M. Champaca.
turbinata, Noronha, l. c. 12.—Malaya.

CHAMPEREIA, Griff. in Calc. Journ. Nat. Hist. iv. (1843) 237. *SANTALACEAE*, Benth. & Hook. f. iii. 231.
MALULUCBAN, Blanco, Fl. Filip. ed. I. 188 (1837).
gnetocarpa, *Kurz, in Journ. Bot.* xiii. (1875) 325.—Ins. Nicobar.
Griffithiana, *Planch. ex Kurz, in Journ. As. Soc. Beng.* xliv. (1875) II. 154.—Burma; Malaya.
Perrottetiana, Baill. Adansonia, iii. (1862-63) 125 = Scleropyrum Wallichianum.

CHAMPIONIA, C. B. Clarke, Comm. et Cyrt. Beng. 98. t. 68 (1874) = **Leptobaea**, Benth. (Gesnerac.).
multiflora, C. B. Clarke, l. c. = L. multiflora.

CHAMPIONIA, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 485. *GESNERACEAE*, Benth. & Hook. f. ii. 1024.
reticulata, *Gardn. l. c.*—Zeylan.

CHAMULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 (Quid?).

CHAPELIERA, Meissn. Gen. 162; Comm. 116 (1838) = seq.

CHAPELIERIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 252. *RUBIACEAE*, Benth. & Hook. f. ii. 96.
madagascariensis, *A. Rich. l. c.* 253.—Madag.

CHAPELLIERA, Nees, in Linnaea, ix. (1834) 298 = **Cladium**, P. Br. (Cyperac.).
arthrophylla, Nees, in Lehm. Pl. Preiss. ii. 77 = Clad. arthrophyllum.
articulata, Nees, l. c. 76, in adnot. = Clad. articulatum.
glomerata, Nees, l. c. = Clad. glomeratum.
iridifolia, Nees, in Linnaea, ix. (1834) 298 = Clad. flexuosum.
juncea, Nees, in Lehm. Pl. Preiss. ii. 76, in adnot. = Clad. junceum.
laxa, Nees, l. c. 76 = Clad. laxum.
pauciflora, Nees, l. c. 76, in adnot. = Tricostularia pauciflora.
riparia, Nees, l. c. 76 = Clad. riparium.
teretifolia, Nees, l. c. 76, in adnot. = Clad. teretifolium.

CHAPMANNIA, Torr. & Gray, Fl. N. Am. i. 355 (1839). *LEGUMINOSAE*, Benth. & Hook. f. i. 517.
floridana, *Torr. & Gray, l. c.*—Florida.

CHAPTALIA, Royle, Illustr. Bot. Himal. t. 59. f. 2 (1839) = **Gerbera**, Gronov. (Compos.).
gossypina, Royle, l. c. 251 = G. lanuginosa.
maxima, D. Don, Prod. Fl. Nep. 166 = G. macrophylla.

CHAPTALIA, Vent. Jard. Cels, t. 61 (1800). *COMPOSITAE*, Benth. & Hook. f. ii. 498.
LERIA, DC. in Ann. Mus. Par. xix. (1812) 68.
LIEBERKUHNA, Cass. in Dict. Sc. Nat. xxvi. 286 (1823).
LOXODON, Cass. l. c. xxvii. 253 (1823).
OXYDON, Less. in Linnaea, v. (1830) 357.
THYRSANTHEMA, Neck. Elem. i. 6 (1790).
albicans, *Vent. ex Steud. Nom.* ed. II. i. 344.—Ind. occ.
Anandria, Spreng. Syst. iii. 504 = Gerbera Anandria.
araneosa, *Casar. in Atti 3ª Riun. Sc. Ital.* iii. (1841) 514.—Bras.
dentata, *Cass. in Dict. Sc. Nat.* xxvi. 104.—Ind. occ.
denticulata, Spreng. ex Baker, in Mart. Fl. Bras. vi. III. 310 = Senecio pulcher.
Ehrenbergii, *Hemsl. Biol. Centr. Am. Bot.* ii. 255.—Mexic.
exscapa, *Poepp. ex DC. Prod.* vii. 44, in syn.—Chili.
heterophylla, D. Don, in Trans. Linn. Soc. xvi. (1830) 244 = Trichocline heterophylla.
integrifolia, *Baker, in Mart. Fl. Bras.* vi. III. 377.—Am. trop.
integrifolia, Vent. ex Spreng. Syst. iii. 504 = tomentosa.
lyrata, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 243.—Mexic.
lyrata, Spreng. Syst. iii. 504 = Gerbera Anandria.
nutans, *Hemsl. Biol. Centr. Am. Bot.* ii. 255.—Mexic.
oblonga, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 240.—Peruv.
obovata, *Wright, in Sauv. Fl. Cub.* 83.—Ins. Cuba.
ovalis, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 241.—Peruv.
piloselloides, *Baker, in Mart. Fl. Bras.* vi. III. 378.—Bras.
piloselloides, *Vent. ex Steud. Nom.* ed. II. i. 344; *Baker, in Mart. Fl. Bras.* vi. III. 378.—Reg. Argent.
rotundifolia, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 242.—Peruv.
runcinata, *H. B. & K. Nov. Gen. et Sp.* iv. 6.—N. Granat.
sarmentosa, Vent. ex Steud. Nom. ed. II. i. 344 = Chevreulia stolonifera.
Seemannii, *Hemsl. Biol. Centr. Am. Bot.* ii. 255.—Mexic.
sinuata, *Baker, in Mart. Fl. Bras.* vi. III. 378.—Reg. Uruguay; Argent.
sinuata, *Vent. ex Steud. Nom.* ed. II. i. 344.—Ins. Jamaic.
spathulata, *Hemsl. Biol. Centr. Am. Bot.* ii. 255.—Mexic.
tomentosa, *Vent. Jard. Cels*, t. 61.—Am. bor.

CHAQUEPIRIA, Endl. Gen. 562, in syn. (1838) sphalm. = CAQUEPIRIA, J. F. Gmel. = **Gardenia**, Linn. (Rubiac.).

CHARACHERA, Forsk. Fl. Aegypt. Arab. 115 (1775). *ACANTHACEAE*?, Benth. & Hook. f. ii. 1071.
XERALIS, Rafin. Sylva Tellur. 82 (1838).

CHARACHERA:—
spicata, Lam. Encyc. i. 695 = viburnoides.
viburnoides, *Forsk. Fl. Aegypt. Arab.* 116.—Arabia.

CHARACIAS, S. F. Gray, Nat. Arr. Brit. Pl. ii. 259 (1821) = **Euphorbia**, Linn.
amygdaloides, S. F. Gray, l. c. 260 = E. amygdaloides.
purpurea, S. F. Gray, l. c. 259 = E. Characias.

CHARADRA, Scop. Introd. 270 (1777) = CHADARA, Forsk. = **Grewia**, Linn.

CHARDINIA, Desf. in Mém. Mus. Par. iii. (1817) 455. t. 21. *COMPOSITAE*, Benth. & Hook. f. ii. 464.
cylindrica, *Desv. Fl. Anj.* 216 = Xeranthemum cylindraceum.
macrocarpa, C. Koch, in Linnaea, xxiv. (1851) 382 = xeranthemoides.
xeranthemoides, *Desf. in Mém. Mus. Par.* iii. (1817) 456.—As. Min.; Persia.

CHARIANTHUS, D. Don, in Mem. Wern. Soc. iv. (1823) 327. *MELASTOMACEAE*, Benth. & Hook. f. i. 762.
CHAENANTHERA, Rich. ex DC. Prod. iii. 196 (1828).
TETRAZYGOS, Rich. ex DC. l. c. 197 (1828).
Berteroanus, Ser. ex DC. l. c. 197 = glaberrimus.
brasiliensis, Cham. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 118 = Miconia brasiliensis.
caeruleus, D. Dietr. Syn. Pl. ii. 1277 = Memecylon Spathandra.
ciliatus, *DC. Prod.* iii. 197.—Ins. Trinitat.
coccineus, *D. Don, in Mem. Wern. Soc.* iv. (1823) 328.—Am. austr.
crinitus, *Naud. in Ann. Sc. Nat.* Sér. III. xviii. (1852) 112.—Ind. occ.
Fadyeni, *Griseb. in Goett. Abh.* ix. (1861) 56.—Jamaic.
glaberrimus, *DC. Prod.* iii. 196.—Ind. occ.
nodosus, Triana, in Trans. Linn. Soc. xxviii. (1871) 99 = ciliatus.
obliquus, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 186 = Miconia obtusa.
obtusus, Triana, in Trans. Linn. Soc. xxviii. (1871) 103 = Miconia obtusa.
purpureus, *D. Don, in Mem. Wern. Soc.* iv. (1823) 329.—Ins. Montserrat.
tinifolius, *D. Don, l. c.*—Ind. occ.

CHARIDIA, Baill. Etud. Gén. Euph. 572. t. 22 (1858) = **Savia**, Willd. (Euphorb.).
mimosoides, Baill. l. c. 573 = S. mimosoides.
pulchella, Baill. l. c. = S. pulchella.

CHARIEIS, Cass. in Bull. Soc. Philom. (1817) 68; et in Dict. Sc. Nat. viii. 191; xxiv. 369 (1817). *COMPOSITAE*, Benth. & Hook. f. ii. 266.
KAULFUSSIA, Nees, Hor. Phys. Berol. 53. t. 11 (1820).
caerulea, Cass. in Dict. Sc. Nat. xxiv. 372 = heterophylla.
heterophylla, *Cass. in Bull. Soc. Philom.* (1817) 68.—Afr. austr.
Neesii, Cass. l. c. (1821) 13, ex eod. in Dict. Sc. Nat. xxiv. 370, 372 = heterophylla.
pilosella, Baill. Hist. Pl. viii. 139 = heterophylla.

CHARIESSA, Miq. Fl. Ind. Bat. i. I. 794 (1855) = **Villaresia**, Ruiz & Pav. (Olacin.).
Smythii, Becc. Malesia, i. (1877) 118.—Malaya.
suaveolens, Miq. Fl. Ind. Bat. i. I. 794.—Malaya.

CHARIOMMA, Miers, Apocyn. S. Am. 110 (1878) = **Echites**, Linn. (Apocyn.).
domingensis, Miers, l. c. 112 = Echites domingensis.
flaxa, Miers, l. c. 113 = Dipladenia flava.
mucronulata, Miers, l. c. 112.—Ins. S. Thom.
nobilis, Miers, l. c. 113 = Dipladenia nobilis.
scandens, Miers, l. c. 114.—Ind. occ.
surrecta, Miers, l. c. 111 = Echites Andrewsii.
verticillata, Miers, l. c. 113.—Jamaic.

CHARLWOODIA, Sweet, Fl. Austral. t. 18 (1827) = **Cordyline**, Comm. (Liliac.).
angustifolia, Goepp. Beitr. Kenn. Dracaen. 17 = Cord. Pumilio.
australis, G. Don, in Loud. Hort. Brit. 130 = Cord. australis.
congesta, Sweet, Fl. Austral. t. 18 = Cord. Pumilio.
ensata, Goepp. Beitr. Kenn. Dracaen. 17 = Dianella ensifolia?

CHARLWOODIA :—
fragrantissima, Lem. Jard. Fleur. iv. (1854) t. 399. Misc. 57 = Cordyline dracaenoides.
indivisa, G. Don, in Loud. Hort. Brit. 130 = Cord. indivisa.
longifolia, Goepp. Beitr. Kenn. Dracaen. 17.—Hab.?
rigidifolia, Koch & Bouché, Ind. Sem. Hort. Berol. (1855) 11 = Cord. Pumilio.
rubra, Planch. in Fl. des Serres, vi. (1850–51) 138 = Cord. rubra.
Sellowiana, Planch. l. c. 137 = Cord. Sellowiana.
spectabilis, Planch. l. c. 136 = Cord. dracaenoides.
stricta, Sweet, Fl. Austral. sub t. 18 = Cord. Pumilio.

CHARPENTIERA, Gaudich. in Freyc. Voy. Bot. 444. t. 47, 48 (1826). *AMARANTACEAE*, Benth. & Hook. f. iii. 26.
obovata, *Gaudich. l. c.* t. 48.—Ins. Sandvic.
ovata, *Gaudich. l. c.* t. 47.—Ins. Sandvic.

CHARPENTIERA, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 346 = **Pavetta**, Linn. (Rubiac.).
bracteata, Vieill. l. c. 347.—N. Caled.

CHARTACALYX, Maing. ex Mast. in Hook. f. Fl. Brit. Ind. i. 382 (1874). *TILIACEAE*.
accrescens, *Mast. l. c.*—Malacca.

CHARTOCALYX, Regel, in Act. Hort. Petrop. vi. (1879) 367. *LABIATAE*.
Olgae, *Regel, l. c.* 368.—Turkestan.

CHARTOLEPIS, Cass. in Dict. Sc. Nat. xliv. 36 (1826) **Centaurea**, Linn. (Compos.).
Biebersteinii, Jaub. & Spach, Illustr. iii. 11 = Cent. glastifolia (ex parte).
cappadocica, Fenzl, in Tchihat. Asie Min. Bot. ii. 328 = Cent. cappadocica.
cappadocica, Jaub. & Spach, Illustr. iii. 137. t. 291 = Cent. macrolopha.
Cassiniana, Jaub. & Spach, l. c. 9 = Cent. glastifolia.
cheirolopha, Fenzl, in Tchihat. Asie Min. Bot. ii. 324 = Cent. cheiracantha.
drabaefolia, Jaub. & Spach, Illustr. iii. 136 = Cent. drabifolia.
glastifolia, Cass. in Dict. Sc. Nat. xliv. 36 = Cent. glastifolia.
intermedia, Boiss. Diagn. Ser. II. iii. 64.—Ross. mer.; Sibir.; Soongar.
Kotschyi, Fenzl, in Tchihat. Asie Min. Bot. ii. 328 = Cent. macrolopha.
lancifolia, Fenzl, l. c. 327 = Cent. lancifolia.
libanotica, Jaub. & Spach, Illustr. iii. 136.—Syria.
lyrata, Boiss. & Haussk. ex Boiss. Fl. Orient. iii. 696. —Persia.
saligna, C. Koch, in Linnaea, xxiv. (1851) 421 (= *Phaeopappus salignus*).—Armenia.
schizolepis, Boiss. ex Boiss. Fl. Orient. iii. 596 (= *Phaeopappus salignus*).—Kurdist.
Tournefortii, Jaub. & Spach, Illustr. iii. 9 = Cent. glastifolia.

CHARTOLOMA, Bunge, in Bot. Zeit. ii. (1844) 249 = **Isatis**, Tourn. (Crucif.).
platycarpum, Bunge, l. c. 250 = I. platycarpa.

CHASALIA, Comm. ex DC. Prod. iv. 531 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 126.
POLYOZUS, Blume, Bijdr. 947 (1826).
amicorum, *A. Gray, in Proc. Am. Acad.* iv. (1860) 43. —Ins. Fiji.
Borboniae, Comm. ex DC. Prod. iv. 532 = divaricata.
Boryana, *DC. l. c.*—Ins. Mascar.
capitata, *DC. l. c.* 531.—Ins. Maurit.
chondrophylla, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 203.—Sumatra.
clusiaefolia, *DC. Prod.* iv. 532.—Ins. Maurit.
coffeoides, DC. l. c. 531 = divaricata?
Commersonii, Juss. ex Steud. Nom. ed. II. i. 345 = capitata.
curviflora, *Thw. Enum. Pl. Zeyl.* 150.—Ind. or.; Malaya.
divaricata, *DC. Prod.* iv. 532.—Ins. Maurit.
expansa, Miq. Fl. Ind. Bat. ii. 280 = Psychotria malayana.
Fontanesii, *DC. Prod.* iv. 531.—Ins. Mascar.
grandifolia, *DC. l. c.* 532.—Ins. Maurit.

CHASALIA:—
Hasseltiana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 204. —Java.
laxiflora, Benth. in Hook. Niger Fl. 416 = Psychotria Ansellii.
leucocarpa, Miq. Fl. Ind. Bat. ii. 281 = Psychotria leucocarpa.
lurida, Miq. l. c. 282 = curviflora.
montana, Miq. l. c. 281 = Psychotria montana.
Mooniac, *Baker, Fl. Maurit.* 155.—Ins. Mascar.
oxylepis, *Miq. Fl. Ind. Bat.* ii. 283.—Java.
parviflora, *Benth. in Hook. Niger Fl.* 417.—Afr. trop.
perforata, Miq. Fl. Ind. Bat. Suppl. 546 = Psychotria angulata.
psychotrioides, DC. Prod. iv. 531 = Gaertnera psychotrioides.
pyriformis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 44. —Ins. Samoa.
robusta, Miq. Fl. Ind. Bat. ii. 280 = Psychotria robusta.
rostrata, *Miq. l. c.* 281.—Malaya.
sangiana, Miq. Fl. Ind. Bat. Suppl. 546 = curviflora.
stipulacea, DC. Prod. iv. 532 = divaricata.
tetrandra, Miq. Fl. Ind. Bat. ii. 282 = curviflora.
umbraticola, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 230.—Afr. trop.

CHASALLIA, Comm. ex Poir. in Dict. Sc. Nat. viii. 198 (1817); et Juss. in Mém. Mus. Par. vi. (1820) 379 = praec.

CHASCANUM, E. Mey. Comm. Pl. Afr. Austr. 275 (1837) = **Bouchea**, Cham. (Verbenac.).
cernuum, E. Mey. l. c. 276 = B. cernua.
cuneifolium, E. Mey. l. c. = B. cuneifolia.
garipense, E. Mey. l. c. 277 = B. garepensis.
laetum, Fenzl, ex Schau. in DC. Prod. xi. 559 = B. pterygocarpa.
marrubiifolium, Fenzl, ex Schau. l. c. 558 = B. marrubiifolia.
pinnatifidum, E. Mey. Comm. Pl. Afr. Austr. 277 = B. pinnatifida.
pumilum, E. Mey. l. c. = B. pumila.

CHASCOLYTRUM, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190 = **Briza**, Linn. (Gramin.).
barbatum, Kunth, Enum. Pl. i. 373 = B. barbata.
coarctatum, Phil. in Linnaea, xxxiii. (1864–65) 293.—Chili.
elegans, Beauv. Agrost. 86 = B. elegans.
erectum, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190 = B. erecta.
nutans, Lindl. ex Heynh. Nom. ii. 137, nomen.—Chili.
rotundatum, Kunth, Rev. Gram. i. 121 = B. rotundata.
rufum, J. & C. Presl, Rel. Haenk. i. 258 = B. rufa.
spicigerum, J. & C. Presl, l. c. 282 = B. spicigera.
strictum, E. Desv. in C. Gay, Fl. Chil. vi. 384 = B. stricta.
subaristatum, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 190 = B. rotundata.
trilobum, E. Desv. in C. Gay, Fl. Chil. vi. 384 = B. triloba.

CHASMANTHERA, Hochst. in Flora, xxvii. (1844) 21. *MENISPERMACEAE*, Benth. & Hook. f. i. 34.
RHIGIOCARYA, Miers, in Ann. Nat. Hist. Ser. III. xiv. (1864) 100.
dependens, *Hochst. l. c.*—Abyssin.
nervosa, *Miers, Contrib. Bot.* iii. 39.—Afr. trop.
uviformis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 459.—Madag.

CHASMANTHIUM, Link, Hort. Berol. i. 159 (1827) = **Uniola**, Linn. (Gramin.).
gracile, Link, l. c. = U. gracilis.
ornithorynchа, Nees, ex Steud. Syn. Pl. Gram. 280 = U. ornithorynchа.

CHASME, Salisb. Parad. Lond. sub t. 67 (1807); et in Knight, Prot. 16 (1809) = **Leucadendron**, R. Br. (Proteac.).
comosa, Salisb. l. c.; Knight, l. c. 17 = L. comosum.
pinifolia, Knight, l. c.—Afr. austr.
ramentacea, Knight, l. c. = L. acmulum.
spiralis, Knight, l. c. 16.—Afr. austr.
teretifolia, Knight, l. c. = L. abietinum.

CHASMIA, Schott, ex Spreng. Syst. iv. Cur. Post. 409 (1827) = **Arrabidaea**, DC. (Bignoniac.).
ochroleuca, Schott, l. c. = A. Agnus-castus, vel A. Salzmanni.

CHASMONE, E. Mey. Comm. Pl. Afr. Austr. 71 (1835) = **Argyrolobium**, Eckl. & Zeyh. (Legum.).
Andrewsiana, E. Mey. l. c. 74 = A. Andrewsianum.
angustissima, E. Mey. l. c. 75 = A. filiforme.
apiculata, E. Mey. l. c. 73 = A. collinum.
argentea, E. Mey. l. c. 74 = A. Linnaeanum.
ascendens, E. Mey. l. c. 73 = A. ascendens.
baptisioides, E. Mey. l. c. 71 = A. baptisioides.
barbata, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 77 = A. barbatum.
calycina, E. Mey. Comm. Pl. Afr. Austr. 74, adnot. = A. calycinum.
crassifolia, E. Mey. l. c. 72 = A. crassifolinm.
crinita, E. Mey. l. c. 71 = A. crinitum.
cuneifolia, E. Mey. l. c. = A. polyphyllnm.
diversifolia, E. Mey. l. c. = A. speciosum.
goodioides, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 75 = A. goodioides.
holosericea, E. Mey. Comm. Pl. Afr. Austr. 72 = A. sericeum.
lanceolata, E. Mey. l. c. 75 = A. lanceolatum.
longifolia, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 74 = A. longifolium.
obcordata, E. Mey. Comm. Pl. Afr. Anstr. 72 = A. obcordatum.
petiolaris, E. Mey. l. c. 75 = A. petiolare.
pumila, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 77 = A. pumilum.
rupestris, E. Mey. Comm. Pl. Afr. Austr. 74 = A. rupestre.
sessiliflorum, E. Mey. l. c. 72 = A. candicans.
splendens, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 78 = A. splendens.
stricta, E. Mey. Comm. Pl. Afr. Austr. 75 = A. strictum.
tenuis, E. Mey. l. c. 75, adnot. = A. angustifolium.
tuberosa, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 73 = A. tuberosum.
venosa, E. Mey. Comm. Pl. Afr. Austr. 73 = A. molle.
verticillata, E. Mey. l. c. 72 = A. stipulaceum.

CHASMONIA, Presl, Fl. Sic. Praef. p. xxxvii (1826) = **Moluccella**, Linn. (Labiat.).
incisa, Presl, l. c. = M. spinosa.

CHASSALIA, Comm. ex Poir. in Dict. Sc. Nat. viii. 198 (1817) = **Chasalia**, Comm. (Rubiac.).

CHASSELOUPIA, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 101 = **Symplocos**, Linn. (Styrac.).
arborea, Vieill. l. c. 102 = S. arborea.
caerulescens, Vieill. l. c. 103 = S. caerulescens, Vieillardi.
gracilis, Vieill. ex Miers, in Journ. Linn. Soc. xvii. (1879) 305 = S. gracilis.
lucida, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 102 = S. glaucescens, lucida.
microphylla, Vieill. l. c. 104.—N. Caled.
montana, Vieill. l. c. = S. montana.
neo-caledonica, Vieill. l. c. 103 = S. stravadioides, Lenormandiana.
nitida, Vieill. ex Miers, in Journ. Linn. Soc. xvii. (1879) 305 = S. rotundifolia.
tinctoria, Vieill. in Bull. Soc. Linn. Normand. x. (1866) 104 = S. baptica.

CHASTENAEA, DC. Prod. iii. 102 (1828) = **Axinaea**, Ruiz & Pav. (Melastomac.).
affinis, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 120 = A. macrophylla.
coriacea, Naud. l. c. iv. (1845) 55 = A. lepidota.
floribunda, Naud. l. c. xviii. (1852) 122 = A. floribunda.
glandulosa, Naud. l. c. 123 = A. glandulosa.
grandifolia, Naud. l. c. 122 = A. grandifolia.
lanceolata, Naud. l. c. 123 = A. lanceolata.
lepidota, Benth. Pl. Hartw. 182 = A. lepidota.
longifolia, Naud. in Ann. Sc. Nat. Sér. III. iv. (1845) 55 = Meriania Karstenii.
macrophylla, Naud. l. c. = A. macrophylla.
Merianiae, DC. Prod. iii. 102 = A. Merianiae.
rigida, Naud. in Ann. Sc. Nat. Sér. III. xviii. (1852) 123 = Meriania rigida.

CHASTOLOMA, Lindl. Veg. Kingd. 355 (1847), sphalm. = CHARTOLOMA, Bunge = **Isatis**, Linn.

CHATAEA, Soland. ex Seem. Pl. Vit. 102 (1865) = CHAITAEA, Soland. = **Tacca**, Forst.
Tacca, Soland. l. c. = T. pinnatifida.

CHATELANIA, Neck. Elem. i. 53 (1790) = **Tolpis**, Adans. (Compos.).

CHATIAKELLA, Cass. in Dict. Sc. Nat. xxix. 491 (1823) = **Wulffia**, Neck. (Compos.).
platyglossa, Cass. l. c. xlvi. 402 = W. stenoglossa.
stenoglossa, Cass. l. c. 403 = W. stenoglossa.

CHAUBARDIA, Reichb. f. in Bot. Zeit. x. (1852) 671 = **Zygopetalum**, Hook. (Orchid.).
surinamensis, Reichb. f. l. c. 672.—Guiana.

CHAULMOOGRA, Roxb. Hort. Beng. 105 (1814); Fl. Ind. iii. 835 (1832) = **Gynocardia**, R. Br. (Bixin.).
odorata, Roxb. ll. cc. 105, 836 = G. odorata.

CHAUNOCHITON, Benth. in Benth. & Hook. f. Gen. i. 996 (1867). *OLACINEAE*, Benth. & Hook. f. l. c.
loranthoides, *Benth. in Hook. Ic. Pl.* t. 1005.—Bras.

CHAUVINIA, Steud. Syn. Pl. Gram. 362 (1855) = **Spartina**, Schreb. (Gramin.).
chilensis, Steud. l. c. = S. densiflora.

CHAVANNESIA, A. DC. in DC. Prod. viii. 444 (1844) = **Urceola**, Roxb. (Apocyn.).
esculenta, A. DC. l. c. = U. esculenta.
javanica, Miq. Fl. Ind. Bat. ii. 458.—Java.
lucida, A. DC. in DC. Prod. viii. 444 = U. lucida.

CHAVICA, Miq. Syst. Pip. 222 (1843) = **Piper**, Linn.
amboinensis, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 134 = P. amboinensis.
Arnottiana, Miq. Syst. Pip. 268 = P. Hapnium.
auriculata, Miq. l. c. 269 = P. Betle.
Benthamiana, Miq. l. c. 233 = P. Barclayanum.
Betle, Miq. l. c. 228 = P. Betle.
Blumei, Miq. l. c. 246 = P. Malamiri.
boehmeriaefolia, Miq. l. c. 265 = P. boehmeriaefolium.
canaliculata, Presl, Epim. Bot. 224 = P. canaliculatum.
celebica, Miq. Syst. Pip. 276 = P. celebicum.
celtidiformis, Miq. l. c. = P. parvifolium, celtidiforme.
ceramica, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 135 = P. ceramicum.
Chaba, Miq. Syst. Pip. 251 = P. Chaba, *Blume*.
chaerostachya, Pritz. Ic. Ind. ii. 75, sphalm. = *sphaerostachya*.
Chawya, C. DC. in DC. Prod. xvi. 1. 351 = seq.
Chuvya, Miq. Syst. Pip. 267 = P. Betle.
consanguinea, Miq. l. c. 232 = P. Labillardieri.
corylistachya, Miq. l. c. 281 = P. corylistachyon.
densa, Miq. l. c. 252 = P. Betle.
diffusa, Miq. in Zoll. Syst. Verz. Ind. Archip. 84 = P. attenuatum.
exasperata, Miq. Syst. Pip. 275 = P. exasperatum.
frustrata, Miq. Pl. Jungh. 16.—Java.
glandulosa, Presl, Epim. Bot. 223 = P. miniatum.
grandis, Presl, l. c. 224 = P. grande.
Guigual, Miq. Syst. Pip. 280 = P. brachystachyum, *Wall.*
Haenkeana, Presl, Epim. Bot. 224 = P. Haenkeanum.
Huegeliana, Miq. in Linnaea, xxvi. (1853) 218 = P. Novae-Hollandiae vel P. Hugelianum.
imperialis, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 134 = P. Forstenii.
Labillardieri, Miq. Syst. Pip. 263 = P. Chaba, *Hunter*.
lanceolata, Miq. l. c. 264 = P. miniatum.
leptostachya, Hance, in Journ. Bot. vi. (1868) 301.—China.
Lessertiana, Miq. Syst. Pip. 270 = P. Pseudo-chavica.
macrostachya, Miq. l. c. 236 = P. miniatum.
maculata, Miq. l. c. 256 = P. maculatum.
madagascariensis, Miq. l. c. 277 = P. madagascariense.
majuscula, Miq. l. c. 271 = P. majusculum.
maritima, Miq. l. c. 262 = P. Chaba, *Hunter*.
Melamiris, Miq. l. c. 245 = P. Sirium.
microstigma, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 138 = P. microstigma.
miniata, Miq. Syst. Pip. 234 = P. miniatum.

CHAVICA:—
Mullesua, Miq. Syst. Pip. 280 = Piper brachystachyum, *Wall.*
multinervia, Miq. Fl. Ind. Bat. Suppl. 473 = P. prianamense.
Neesiana, Miq. Syst. Pip. 249 = P. peepuloides.
officinarum, Miq. l. c. 256 = P. Chaba, *Hunter*.
parvifolia, Hassk. in Flora, xlvii. (1864) 59 = P. parvifolium.
peepuloides, Miq. Syst. Pip. 237 = P. peepuloides.
peepuloides, Wight, Ic. t. 1927 = P. Chaba, *Hunter*.
penangensis, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 433 = P. Lonchites.
penangensis, Miq. Syst. Pip. 279 = P. penangense.
petiolata, C. DC. in DC. Prod. xvi. 1. 389.—Ind. or.
populifolia, Miq. Syst. Pip. 248.—Ins. Philipp.
pothoides, Miq. l. c. 266 = P. pothoides.
pseudomelamiris, Miq. l. c. 244 = P. Sirium.
puberula, Benth. Fl. Hongk. 335.—China.
pyrifolia, Miq. Syst. Pip. 275 = P. pyrifolium.
radicans, Presl, Epim. Bot. 224 = P. radicans.
Rafflesii, Miq. Syst. Pip. 273 = P. Rafflesii.
Reinwardtiana, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 136 = P. Reinwardtianum.
retrofracta, Miq. Syst. Pip. 275 = P. retrofractum.
Roxburghii, Miq. l. c. 239 = P. longum.
rufinervis, Presl, Epim. Bot. 224 = P. rufinerve.
Rumphii, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 141 = P. Rumphii.
salicina, Presl, Epim. Bot. 224 = P. salicinum.
sarmentosa, Miq. in Hook. Lond. Journ. Bot. iv. (1845) 433 = P. longum.
sarmentosa, Miq. Syst. Pip. 242 = P. sarmentosum.
sinensis, Champ. ex Benth. in Hook. Kew Journ. vi. (1854) 116 = P. sinense.
Siriboa, Miq. Syst. Pip. 224 = P. Betle.
sphaerostachya, Miq. l. c. 278 = P. brachystachyum, *Wall.*
subcordata, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 138 = P. subcordatum.
subseptemnervia, Miq. Fl. Ind. Bat. Suppl. 473 = P. subseptemnervium.
Suipigua, Miq. Syst. Pip. 275 = P. Suipigua.
sundaica, Miq. l. c. 274 = P. sundaicum.
sylvatica, Miq. l. c. 248 = P. sylvaticum.
syringaefolia, Miq. l. c. 275 = P. syringaefolium.
Teysmanni, Miq. in Ann. Mus. Bot. Lugd. Bat. i. 141 = P. Teysmanii.
thermalis, Miq. Syst. Pip. 276 = P. thermale.
Thomsonii, C. DC. in DC. Prod. xvi. 1. 389.—Ind. or.
viminalis, Miq. in Linnaea, xx. (1847) 131 = P. viminale.
Wallichii, Miq. Syst. Pip. 254 = P. aurantiacum.

CHAYOTA, Jacq. Am. Select. ed. pict. t. 245 (1780) = **Sechium**, P. Br. (Cucurbitac.).
edulis, Jacq. l. c. = S. edule.

CHEILANTHOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 56 (genus indesc. *LABIATARUM*).

CHEILOCLINIUM, Miers, in Trans. Linn. Soc. xxviii. (1872) 420. *HIPPOCRATEACEAE*.
anomalum, *Miers, l. c.*—Bras.

CHEILOCOCCA, Salisb. ex Sm. Spec. Bot. N. Holl. 17 (1793), cum cit. falsa = **Platylobium**, Sm. (Legumin.).
apocynifolia, Salisb. ex Sm. l. c. = P. formosum.

CHEILODISCUS, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 36 = **Pectis**, Linn. (Compos.).
littoralis, Triana, l. c.—N. Granat.

CHEILOPSIS, Moq. in Ann. Sc. Nat. Sér. I. xxvii. (1832) 230 = **Acanthus**, Tourn.
arborea, Nees, in DC. Prod. xi. 272 = A. arboreus.
montana, Nees, l. c. = A. montanus.
polystachya, Moq. in Ann. Sc. Nat. Sér. I. xxvii. (1832) 230 = A. arboreus.

CHILOSA, Blume, Bijdr. 613 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 322.
montana, *Blume*, l. c. 614.—Java.

CHEILOSANDRA, Griff. ex Lindl. Veg. Kingd. 672 (1847) = **Rhynchotechum**, Blume (Gesnerac.).
leptopus, Griff. ex Benth. & Hook. f. Gen. ii. 1025 = Leptobaea glabra.

CHEILOTHECA, Hook. f. in Benth. & Hook. f. Gen. ii. 607 (1876). *MONOTROPEAE*, Benth. & Hook. f. l. c.
khasiana, *Hook. l. c.* 608.—Reg. Himal.

CHEILYCTIS, Benth. Lab. Gen. et Sp. 726 (1836) = **Monarda**, Linn. (Labiat.).

CHEIRADENIA, Lindl. Fol. Orch. Cheirad. (1853) 1. *ORCHIDEAE*, Benth. & Hook. f. iii. 544.
cuspidata, *Lindl. l. c.*—Guiana.

CHEIRANTHERA, Endl. Enchir. 515 (1841) = **Cheirostemon**, Humb. & Bonpl. (Sterculiac.).

CHEIRANTHERA, Brongn. Voy. Coq. Bot. t. 77 (1826); et A. Cunn. in Bot. Reg. sub t. 1719 (1834). *PITTOSPOREAE*, Benth. & Hook. f. i. 133.
brevifolia, F. Muell. Fragm. i. 97 = filifolia.
cyanea, Brongn. Voy. Coq. Bot. t. 77 = linearis.
filifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 364.—Austral.
linearis, *A. Cunn. in Bot. Reg.* sub t. 1719.—Austral.
parviflora, *Benth. Fl. Austral.* i. 128.—Austral.
Preissiana, *Putterl. in Lehm. Pl. Preiss.* i. 201 (sp. dub.).—Austral.
tortilis, F. Muell. Fragm. ii. 79 = filifolia.
volubilis, *Benth. Fl. Austral.* i. 128.—Austral.

CHEIRANTHODENDRON, Benth. & Hook. f. Gen. i. 212 (1862) = CHIRANTHODENDRON, Cerv. = **Cheirostemon**, Humb. & Bonpl. (Sterculiac.).

CHEIRANTHODENDRUM, Steud. Nom. ed. I. 186 (1821) = praec.

CHEIRANTHUS, Linn. Gen. ed. I. 195 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 68.
CHEIRI, Adans. Fam. ii. 418 (1763); [Clairv.] Man. Herb. 221 (1811).
DICHROANTHUS, Webb & Berth. Phytog. Canar. i. 65. t. 5, 6 (1836).
PHOENICAULIS, Nutt. ex Torr. & Gray, Fl. N. Am. i. 89 (1838).
SCHELHAMERIA, Heist. Syst. 9 (1748).
acaulis, Balb. ex Spreng. Nov. Prov. 10 = Matthiola acaulis.
africanus, Linn. Pl. Afr. 13 (Amoen. Acad. vi. 90) = Heliophila pilosa.
albiflorus, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 133. —Tibet. occ.
albus, Mill. Gard. Dict. ed. VIII. n. 8 = Matthiola incana.
alpinus, All. Fl. Pedem. iii. t. 20 = Erysimum lanceolatum.
alpinus, Falk, Beitr. ii. 216 = Erysimum altaicum.
alpinus, Jacq. Fl. Austr. i. 48. t. 75 = Erysimum canescens.
alpinus, Lam. Encyc. ii. 716 = Erysimum ochroleucum.
alpinus, *Linn. Mant.* 93 (excl. syn.).—Scandinav.
angulatus, Schult. Obs. 129 = Syrenia sessiliflora.
angustifolius, Gilib. Fl. Lituan. ii. 61 = Erysimum hieracifolium.
angustifolius, Mill. Gard. Dict. ed. VIII. n. 4 = Erysimum lanceolatum.
annuus, Georgi, Reise, i. 225 = Hesperis aprica.
annuus, *Linn. Sp. Pl.* 662.—Davuria.
apricus, Steph. ex Willd. Sp. Pl. iii. 518 = Hesperis aprica.
aquaticus, Lejeune, Fl. Spa, ii. 68 = Erysimum cheiranthoides.
arborescens, Sieber, ex Link, Enum. Hort. Berol. ii. 166, nomen = Matthiola arborescens.
arboreus, *Sweet, ex Steud. Nom.* ed. II. i. 345.—Aegypt.
arbuscula, *Lowe, in Hook. Kew Journ.* viii. (1856) 289. —Ins. Canar.
arenarius, Suffren, ex DC. Syst. ii. 171 = Matthiola varia.
armeniacus, Sims, Bot. Mag. t. 835 = Erysimum ibericum.
arvensis, Bernh. Syst. Verz. Erf. 196 = Erysimum hieracifolium.
asper, *Cham. & Schlecht. in Linnaea*, i. (1826) 14. —Am. bor. occ.
asper, Nutt. Gen. Am. ii. 436 = Erysimum asperum.
aurantiacus, *Bunge, Enum. Pl. Chin. Bor.* 5.—China.
aureus, Willd. ex DC. Syst. ii. 581 = Sterigma elychrysifolium.

CHEIRANTHUS:—

auriculatus, Lapeyr. Hist. Abr. Pl. Pyr. 362 = Cochlearia auriculata.
axillaris, Brouss. ex Webb & Berth. Phyt. Canar. i. 67 = scoparius.
bicolor, Hornem. Hort. Hafn. ii. 613 = Erysimum bicolor.
bicornis, Sibth. & Sm. Fl. Graec. Prod. ii. 26 = Matthiola bicornis.
bicuspidatus, Willd. Sp. Pl. iii. 519 = Hesperis bicuspidata.
bithynicus, Pers. Syn. ii. 200 = Erysimum cuspidatum.
Bocconi, All. Fl. Pedem. i. 272 = Erysimum canescens.
callosus, Linn. f. Suppl. 296 = Heliophila callosa.
canus, Piller & Mitterp. Iter Poseg. 146. t. 15 = Erysimum angustifolium.
capitatus, Dougl. in Hook. Fl. Bor. Am. i. 38 = asper.
carnosus, Thunb. Prod. Pl. Cap. 108 = Heliophila platysiliqua.
caspicus, Lam. in Pall. Voy. 8° App. 348 = Sterigma tomentosum.
Cheiri, *Linn. Sp. Pl.* 661.—Europ.
chius, Linn. l. c. = Malcolmia chia.
chius, Pall. Reise, i. 497 = Chorispora tenella.
cinereus, *Webb & Berth. Phyt. Canar.* i. t. 5.—Ins. Canar.
coccineus, Mill. Gard. Dict. ed. VIII. n. 7 = Matthiola incana.
collinus, Bieb. Fl. Taur. Cauc. ii. 119 = Erysimum collinum.
contortuplicatus, Steph. in Willd. Sp. Pl. iii. 521 = Sisymbrium contortuplicatum.
corinthius, *Boiss. Diagn.* Ser. II. i. 18.—Graecia.
cornutus, Lam. Encyc. ii. 717 = Syrenia sessiliflora.
coronopifolius, Sibth. & Sm. Fl. Graec. Prod. ii. 25 = Matthiola tristis.
crassicaulis, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 45 = Erysimum crassicaule.
Cumbrae, Link, in Buch, Beschr. Canar. Ins. 152 = scoparius.
cuspidatus, Bieb. Fl. Taur. Cauc. ii. 120 = Erysimum cuspidatum.
decumbens, Schleich. ex Willd. Enum. Hort. Berol. 680 = ochroleucus.
Delilianus, *A. Gouault, in Rev. Hortic.* Sér. IV. i. (1852) 401.—Hab.?
denticulatus, Willd. ex Steud. Nom. ed. II. i. 346 = Pallasii.
dubius, Hornem. Hort. Hafn. Suppl. 73 = Erysimum dubium.
dubius, Suter, Fl. Helv. ii. 65 = ochroleucus.
elongatus, Thunb. Prod. Pl. Cap. 108 = Heliophila linearifolia.
erucaefolius, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 793 = Malcolmia lacera.
erysimoides, Gueldenst. ex Ledeb. Fl. Ross. i. 188 = Erysimum leptostylum.
erysimoides, Huds. Fl. Angl. ed. II. 287 = Erysimum cheiranthoides.
erysimoides, Linn. Sp. Pl. 661 = Erysimum lanceolatum.
erysimoides, Pall. Reise, i. 192 = Erysimum Andreioskianum.
erysimoides, Vill. ex Willk. & Lange, Prod. Fl. Hisp. iii. 808 = Erysimum ochroleucum.
Farsetia, Buch.-Ham. in Wall. Cat. n. 4801 = Farsetia Hamiltonii.
Farsetia, Linn. Mant. i. 94 = Farsetia aegyptiaca.
fenestralis, Linn. Sp. Pl. 1198 = Matthiola fenestralis.
firmus, Schleich. ex Willd. Enum. Hort. Berol. 45 = Erysimum lanceolatum?
firmus, Willd. Enum. Hort. Berol. Suppl. 45 = Erysimum virgatum.
flexuosus, *Sibth. & Sm. Fl. Graec. Prod.* ii. 24; *Fl. Graec.* t. 634.—Ins. Cyprus.
fragrans, Fisch. Hort. Gorenk. (1812) 51 = Matthiola fragrans.
frutescens, Vent. ex Pers. Syn. ii. 201 = semperflorens.
fruticosus, L'Hérit. ex DC. Prod. i. 197 = C. Cheiri.
fruticulosus, Georgi, Beschr. Russ. Reich. iii. iv. 1134 = Sterigma tomentosum.
fruticulosus, L'Hérit. ex DC. Syst. ii. 495 = Erysimum suffruticosum.
fruticulosus, Linn. Mant. 94 = C. Cheiri.
fruticulosus, Linn. Sp. Pl. 662 = Matthiola tristis.
fruticulosus, Sibth. & Sm. Fl. Graec. Prod. ii. 23 = Erysimum Boryanum.

CHEIRANTHUS:—

glaber, Mill. Gard. Dict. ed. VIII. n. 9 = Matthiola glabra.
glaberrimus, Colla, Antol. Bot. v. 861 = Matthiola glabra.
graecus, [Juss.] Pers. Syn. ii. 201 = Matthiola incana.
gramineus, Thunb. Prod. Pl. Cap. 108 = Heliophila graminea.
Griffithii, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 137 = Erysimum Hookeri.
helveticus, Jacq. Hort. Vindob. iii. 9 = Erysimum murale.
hesperidioides, *Torr. & Gray, Fl. N. Am.* i. 72.—Am. bor.
hieracifolius, Lam. Encyc. ii. 717 = Erysimum hieracifolium.
hilaris, Willd. ex Steud. Nom. ed. II. i. 346 = Matthiola tristis.
himalaicus, Hook. f. & Thoms. in Journ. Linn. Soc. v. (1861) 137 = himalayensis.
himalayensis, *Cambess. in Jacquem. Voy. Bot.* 14.—Reg. Himal.
hirtus, Schlecht. fide Stev. ex DC. Prod. i. 189 = Hesperis aprica.
hispidus, Adams, in Nouv. Mém. Soc. Nat. Mosc. iii. (1834) 250 = Hesperis aprica.
hortensis, Lam. Fl. Fr. ii. 506 = Matthiola incana.
humilis, Clarke, Trav. iii. 42 = Matthiola humilis.
ibericus, Adams, in Weber & Mohr, Beitr. i. 61 = Erysimum ibericum.
ibericus, Willd. Enum. Hort. Berol. 681 = Barbarea arcuata.
incanus, Linn. Sp. Pl. 662 = Matthiola incana.
incrassatus, Bory, ex Steud. Nom. ed. II. i. 346 = Malcolmia flexnosa.
integrifolius, Mill. Gard. Dict. ed. VIII. n. 2 = Erysimum canescens.
integerrimus, Steud. Nom. ed. II. i. 346 = Erysimum canescens.
intermedius, Schleich. Cat. Pl. Helv. ed. IV. 12 = C. Cheiri?
Jacomellii, Clem. Sert. Olymp. 9. t. 4. f. 2 = Erysimum Kotschyanum.
junceus, Kit. Pl. Rar. Hung. iii. 259 = Sisymbrium junceum.
Keiri, Neck. Delic. Gallo-Belg. i. 283 = C. Cheiri.
Knowleanus, D. Dietr. Syn. Pl. iii. 683 = ochroleucus.
Knowlesii, Walp. Rep. i. 125 = ochroleucus.
lacerifolius, Pall. Reise Statth. Russ. Reich. i. 124 = Matthiola tatarica.
lacerus, Gouan, Illustr. 44 = Malcolmia parviflora.
lacerus, Linn. Sp. Pl. 662 = Malcolmia lacera.
laciniatus, Poir. Encyc. Snppl. ii. 780 = Hesperis laciniata.
laevigatus, Willd. ex DC. Prod. i. 141 = Barbarea arcuata.
lanatus, Gmel. ex DC. Syst. ii. 581 = Sterigma elychrysifolium.
lanceolatus, Willd. Sp. Pl. iii. 515 = Hesperis tristis.
leptophyllus, Bieb. Fl. Taur. Cauc. ii. 119 = Erysimum leptophyllum.
leptophyllus, Willd. ex DC. Syst. ii. 464 = Sisymbrium junceum.
leucanthemus, Steph. ex Willd. Sp. Pl. iii. 521 = Erysimum versicolor.
linariaefolius, *Vis. in Nuov. Sagg. Acc. Pad.* v. (1840) 267.—Afr. bor.
linearis, Forsk. Fl. Aegypt. Arab. 119 = Farsetia aegyptiaca.
linearis, Thunb. Prod. Pl. Cap. 108 = Heliophila linearis.
linearis, Vent. Jard. Malm. t. 83 = semperflorens.
linifolius, Pers. Syn. ii. 201 = Erysimum linifolium.
littoreus, All. Fl. Pedem. i. 273 = Malcolmia maritima.
littoreus, Linn. Sp. Pl. ed. II. 925 = Malcolmia littorea.
littoreus, Pall. Reise, ii. 741 = Sterigma tomentosum.
lividus, Delile, Fl. Aeg. Illustr. 67 = Matthiola livida.
lividus, Sieber, ex Steud. Nom. ed. II. i. 346 = Malcolmia aegyptiaca.
longifolius, Vent. Jard. Malm. t. 83 = mutabilis.
longipetalus, Vent. Jard. Cels, t. 93 = Matthiola oxyceras.
longisiliquus, Steud. Nom. ed. II. i. 346 = Erysimum longisiliquosum.
luteus, Dulac, Fl. Hautes-Pyr. 199 = C. Cheiri.

CHEIRANTHUS:—

lyratus, Sibth. & Sm. Fl. Graec. Prod. ii. 24 = Malcolmia lyrata.
maritimus, Linn. Cent. Pl. i. 19 (Amoen. Acad. iv. 280) = Malcolmia maritima.
Marshallii, Stark, ex Moore, Gard. Mag. Bot. i. (1850) 289 = ochroleucus.
Menziesii, *Benth. & Hook. f. Gen.* i. 68.—Am. bor. occ.
mollis, Hornem. Hort. Hafn. ii. 615 = Arabis albida.
montanus, Pall. Reise, i. 496 = Syrenia siliculosa.
muralis, Salisb. Prod. 271 = C. Cheiri.
muricatus, Lam. Fl. Fr. ii. 507 = Matthiola sinuata.
muricatus, Weinm. Cat. Hort. Dorpat. (1810) 41 = Dontostemon integrifolius.
mutabilis, Brouss. ex Spreng. Syst. ii. 896 = semperflorens.
mutabilis, *L'Hérit. Stirp. Nov.* 92.—Ins. Mader. et Canar.
nanus, Merk. ex DC. Syst. ii. 486 = Matthiola nana.
nitrarius, *Pall. Reise Statth. Russ. Reich.* i. 105, 124, 141.—Rossia.
nodosus, Lam. Fl. Fr. ii. 508 = Malcolmia littorea.
ochroleucus, DC. Syst. ii. 181 = Erysimum ochroleucum.
ochroleucus, *Hall. f. ex Schleich. Cat.* 16.—Europ.
odoratissimus, Bieb. Casp. 116 = Matthiola odoratissima.
odoratus, *Pall. ex Spreng. Syst.* ii. 896.—Persia.
olympicus, *Sibth. ex DC. Syst.* ii. 494.—Graecia.
Pallasii, Pursh, Fl. Am. Sept. ii. 436 = pygmaens.
pallens, Hall. f. ex DC. Syst. ii. 501 = Erysimum ochroleucum.
pallidus, Pall. Reise, iii. 230 = Erysimum canescens.
paniculatus, Cav. Desc. 439 = Erysimum Kunzeanum.
paniculatus, Lam. Encyc. ii. 717 = Erysimum repandum.
paniculatus, Stev. ex DC. Syst. ii. 210 = Arabis incarnata.
Parnassii, Boiss. Diagn. Ser. II. i. 18 = Erysimum Boryanum.
parryoides, *Kurz, ex Hook. f. Fl. Brit. Ind.* i. 132.—Tibet.
parviflorus, Schousb. Vextr. Marokko, 195 = Matthiola parviflora.
parviflorus, Thib. ex DC. Prod. i. 134 = Matthiola coronopifolia.
persicus, Pall. ex DC. l. c. 132 = Anchonium Tournefortii.
persicus, Willd. ex Ledeb. Fl. Ross. i. 109 = Matthiola tatarica.
pinnatifidus, Willd. Sp. Pl. iii. 523 = Dontostemon pectinatus.
planifolius, Waldst. & Kit. ex Bieb. Fl. Taur. Cauc. iii. 443 = Erysimum cuspidatum.
pulchellus, Willd. Sp. Pl. iii. 523 = Erysimum rupestre.
Pumilio, Sibth. & Sm. Fl. Graec. Prod. ii. 26 = Matthiola bicornis.
pumilus, Hornem. Hort. Hafn. ii. 613 = Erysimum pumilum.
pygmaeus, *Adams, in Mém. Soc. Nat. Mosc.* v. (1817) 114.—Sibir.; Am. bor. occ.
quadrangularis, Steud. Nom. ed. II. i. 346 = seq.
quadrangulus, L'Hérit. Stirp. Nov. 91. t. 44 = Syrenia sessiliflora.
ramosissimus, Lam. Encyc. ii. 717 = Erysimum repandum.
ramosissimus, *Bory & Chaub. Nouv. Fl. Pélop.* 43.—Graecia.
rhaeticus, Schleich. Cat. Pl. Helv. ed. III. 12 = Erysimum ochroleucum.
ruderalis, Pall. ex DC. Syst. ii. 580 = Sterigma tomentosum.
rupestris, Sibth. & Sm. Fl. Graec. Prod. ii. 23; Fl. Graec. t. 633 = Erysimum rupestre.
salinus, *Linn. Mant.* 93.—Sibir.; Taur.
salinus, Pall. ex DC. Syst. ii. 580 = Sterigma tomentosum.
salinus, Willd. ex DC. l. c. 210 = Arabis incarnata.
scaber, Moench, Meth. 254 = Malcolmia africana.
scaber, *Webb & Berth. Phyt. Canar.* i. 68.—Ins. Canar.
scapiger, Adams, in Mém. Soc. Nat. Mosc. v. (1817) 112 = Parrya macrocarpa.
scapigerus, Willd. Prod. Fl. Berol. 214. t. 5. f. 10 = Erysimum cheiranthoides.
scoparius, *Brouss. ex Willd. Enum. Hort. Berol.* 681.—Ins. Canar.

CHEIRANTHUS:—
semperflorens, Coss. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 395 = Erysimum longifolium.
semperflorens, *Schousb. Vextr. Marokko*, 195. — Marocco.
Senoneri, *Heldr. & Sart. Cat. Hort. Athen.* (1862) *ex Reut. Cat. Gram. Genève* (1865), 4.—Archip. Graec.
sibiricus, Juss. ex DC. Syst. ii. 492 = Erysimum angustifolium.
silenifolius, Willd. Enum. Hort. Berol. Suppl. 45 = Erysimum murale.
siliculosus, Bieb. Fl. Taur. Cauc. ii. 121 = Syrenia siliculosa.
sinuatus, Linn. Sp. Pl. ed. II. 926 = Matthiola sinuata.
sinuatus, Pall. Reise, ii. 329 = Sterigma tomentosum.
Stewartii, *T. Anders. in Hook. f. Fl. Brit. Ind.* i. 132. —Reg. Himal.
Stocksianus, Boiss. Diagn. Ser. II. i. 19 = Erysimum Stocksianum.
strictus, Linn. f. Suppl. 296 = Heliophila callosa.
strictus, Poir. Encyc. Suppl. ii. 781 = Heliophila scoparia.
strigosus, Ledeb. in Mém. Acad. Pétersb. v. (1812) 589 = Erysimum strigosum.
subulatus, Moench, Meth. 254 = Malcolmia chia.
sulphureus, [Soland. in] Russ. Aleppo, ed. II. ii. 257 = Sterigma sulphureum.
sylvestris, Bernh. ex Steud. Nom. ed. II. i. 346 = Erysimum hieracifolium.
sylvestris, Crantz, Stirp. Austr. ed. I. i. 48 ; Lam. Encyc. ii. 716 = Erysimum lanceolatum.
syriacus, DC. Syst. ii. 185 = Conringia orientalis.
syriacus, Willd. ex Steud. Nom. ed. II. i. 346 = Matthiola oxyceras.
taraxacifolius, Balb. Cat. Hort. Taur. App. (1814) 10 = Malcolmia taraxacifolia.
taraxacifolius, Biasol. ex Reichb. Fl. Germ. Excurs. 687 = Erysimum strictum.
taraxacifolius, Schrank, in Denkschr. Bot. Ges. Regensb. ii. (1818) 164 = Chorispora tenella.
taraxacifolius, *Steph. ex Willd. Sp. Pl.* iii. 525.—Sibir.
tataricus, Pall. ex DC. Syst. ii. 170 = Matthiola tatarica.
tauricus, Lepech. in Nov. Act. Petrop. xiii. (1802) 336. t. 9 = Matthiola odoratissima.
tenuifolius, *L'Hérit. Stirp. Nov.* 92.—Ins. Madera.
tomentosus, Willd. Sp. Pl. iii. 523 = Sterigma tomentosum.
torulosus, Bieb. Fl. Taur. Cauc. ii. 121 = Sterigma torulosum.
torulosus, Thunb. Prod. Pl. Cap. 108 = Matthiola torulosa.
tricuspidatus, Huds. Fl. Angl. ed. I. 250 = Matthiola sinuata.
tricuspidatus, Linn. Sp. Pl. 663 = Matthiola tricuspidata.
trilobus, Linn. l. c. 662 = Malcolmia littorea.
tristis, Curt. Bot. Mag. t. 729 = Matthiola varia.
tristis, Forsk. Fl. Aegypt. Arab. 119 = Matthiola livida.
tristis, Linn. Syst. Nat. ed. X. 1134 = Matthiola tristis.
turritoides, Lam. Encyc. ii. 716 = Erysimum cheiranthoides.
unguiculatus, Willd. ex Steud. Nom. ed. II. i. 347 = Malcolmia littorea.
vallesiacus, J. Gay, ex DC. Prod. i. 134 = Matthiola varia.
varius, Sibth. & Sm. Fl. Graec. Prod. ii. 25 ; Fl. Graec. t. 636 = Matthiola varia.
versicolor, Bieb. Fl. Taur. Cauc. ii. 119 = Erysimum versicolor.
versicolor, Pall. ex Steud. Nom. ed. II. i. 347 = Hesperis hyrcanica.
villosus, Forsk. Fl. Aegypt. Arab. 120 = Matthiola tricuspidata.
villosus, Spreng. ex DC. Prod. i. 188 = Hesperis laciniata.
virgatus, Patr. ex DC. Syst. ii. 462 = Sisymbrium junceum.
virgatus, Poir. Encyc. Suppl. ii. 781 = Erysimum angustifolium.
viridifolius, Steud. Nom. ed. II. i. 347 = Matthiola glabra.
viridis, Ehrh. Beitr. vii. 158 = Matthiola incana.

CHEIRI, Adans. Fam. ii. 418 (1763) = **Cheiranthus**, Linn. (Crucif.).
hieracifolium, [Clairv.] Man. Herb. 221 = Erysimum hieracifolium.
montanum, [Clairv.] l. c. = Cheiranthus Cheiri.
vulgare, [Clairv.] l. c. = Cheiranthus Cheiri.

CHEIRINIA, Link, Enum. Hort. Berol. ii. 170 (1822) = **Erysimum**, Tourn. (Crucif.).
altissima, Link, l. c. 171 = E. hieracifolium.
Andrzejowskiana, Hort. Berol. ex Steud. Nom. ed. II. i. 347 = E. canescens.
angustifolia, Link, Enum. Hort. Berol. ii. 170 = E. angustifolium.
canescens, Link, l. c. 171 = E. canescens.
cheiranthoides, Link, l. c. 170 = E. cheiranthoides.
crepidifolia, A. Dietr. Fl. Boruss. vii. 446 = E. crepidifolium.
cuspidata, Link, Enum. Hort. Berol. ii. 170 = E. cuspidatum.
erysimoides, Dum. Fl. Belg. 123 = E. lanceolatum.
firma, Link, Enum. Hort. Berol. ii. 171 = E. virgatum.
helvetica, Link, l. c. = E. ochroleucum.
hieracifolia, Link, l. c. 170 = E. hieracifolium.
juncea, Link, l. c. 172 = Sisymbrium junceum.
lanceolata, Link, l. c. 171 = E. lanceolatum.
leptostyla, Link, Handb. ii. 315 = E. leptostylum.
longifolia, Link, Enum. Hort. Berol. ii. 171 = E. longifolium.
longisiliqua, Link, Handb. ii. 315 = E. virgatum.
longisiliquosa, Link, Enum. Hort. Berol. ii. 170 = E. longisiliquosum.
odorata, Link, Handb. ii. 315 = E. odoratum.
repanda, Link, Enum. Hort. Berol. ii. 171 = E. repandum.
sessiliflora, Link, l. c. 170 = Syrenia sessiliflora.
stricta, Link, l. c = E. hieracifolium.
strictissima, Link, l. c. 171 = Sisymbrium strictissimum.
suffruticosa, Link, l. c. 170 = E. murale.
virgata, Link, l. c. = E. virgatum.

CHEIRISANTHERA, Hort. ex Regel, Gartenfl. ii. (1853) 353 = **Heppiella**, Regel (Gesnerac.).
atrosanguinea, Hort. ex Regel, l. c. = H. atrosanguinea.
coccinea, Hort. ex Regel, l. c. = H. atrosanguinea.

CHEIRODENDRON, Nutt. ex Seem. in Journ. Bot. v. (1867) 236 = **Panax**, Linn. (Araliac.).
Gaudichaudii, Seem. l. c. = P. Gaudichaudii.
laetevirens, Seem. l. c.—Chili.
platyphylla, Seem. l. c. = P. platyphyllum.
samoënse, Seem. l. c. 237.—Samoa.
valdiviense, Seem. l. c. 236.—Chili.

CHEIROLAENA, Benth. in Benth. & Hook. f. Gen. i. 222 (1862). *STERCULIACEAE*, Benth. & Hook. f. l. c.
linearis, *Benth. c.*—Madag.

CHEIROLEPIS, Boiss. Diagn. Ser. I. x. 106 (1849) = **Centaurea**, Linn. (Compos.).
cappadocica, Boiss. l. c. 107 = Cent. cappadocica.
drabifolia, Boiss. l. c. = Cent. drabifolia.
Kotschyi, Boiss. & Heldr. l. c. = Cent. macrolopha.
libanotica, Boiss. l. c. (= *Phaeopappus libanoticus*).—Syria.
persica, Boiss. l. c. 108 = Cent. macrolopha.

CHEIROLOMA, F. Muell. in Linnaea, xxv. (1852) 401 = **Calotis**, R. Br. (Compos.).
hispidulum, F. Muell. l. c. = Cal. hispidula.

CHEIROLOPHUS, Cass. in Dict. Sc. Nat. l. 250 (1827) = **Centaurea**, Linn. (Compos.).
lanceolatus, Cass. l. c. li. 56 = Cent. sempervirens.
pinnatifidus, Cass. l. c. = Cent. intybacea.
sempervirens, Pomel, Nouv. Mat. Fl. Atl. 32 = Cent. sempervirens.

CHEIROPETALUM, E. Fries, Ind. Sem. Hort. Upsal. (1857) ex Urban, Addit. Ind. Sem. Hort. Berol. (1881) 11 = **Silene**, Linn. (Caryoph.).
radians, E. Fries, l. c. = S. odoratissima.

CHEIROPSIS, Bercht. & Presl, Rostl. i. Ranunculac. 11 (1823) = **Clematis**, Tourn. (Ranunculac.).
balearica, Bercht. & Presl, l. c. = Clem. balearica.
cirrhosa, Bercht. & Presl, l. c. = Clem. cirrhosa.

CHEIROPSIS:—
elegans, Spach, Hist. Vég. Phan. vii. 261 = Clematis cirrhosa.
semitriloba, Bercht. & Presl, Rostl. i. Ranunculac. 11 = Clem. semitriloba.

CHEIROPTEROCEPHALUS, Rodrig. Orch. Nov. i. 28 (1877). *ORCHIDEAE.*
sertuliferus, *Rodrig. l. c.* 29.—Bras.

CHEIROSTEMON, Humb. & Bonpl. Pl. Aequin. i. 81. t. 24 (1808). *STERCULIACEAE*, Benth. & Hook. f. i. 212, 982.
CHIRANTHODENDRON, Cerv. ex Cav. in Anal. Cienc. Nat. vi. (1803) 303, adnot.
apetalum, Cerv. in Roem. Collect. Bot. 97 = platanoides.
platanoides, *Humb. & Bonpl. Pl. Aequin.* 82. t. 24.—Mexic.

CHEIROSTYLIS, Blume, Bijdr. 413. t. 16 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 600.
cochinchinensis, *Blume, Orch. Archip. Ind.* t. 39. f. 2.—Cochinch.
flabellata, *Wight, Ic.* v. 16. (t. 1727, Monochilus).—Ind. or.
grandiflora, *Blume, Orch. Archip. Ind.* 55.—N. Guinea.
Griffithii, *Lindl. in Journ. Linn. Soc.* i. (1857) 188.—Reg. Himal. ; Burma.
gymnochiloides, *Reichb. f. in Flora*, lxviii. (1885) 537. —Madag.
heterosepala, *Reichb. f. Otia Bot. Hamb.* 110.—Afr. trop.
Humblotii, *Reichb. f. in Flora*, lxviii. (1885) 537.—Ins. Comoro.
malleifera, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 141.—Burma.
marmorata, Lindl. ex Lem. in Fl. des Serres, Sér. I. iv. (1848) t. 370 = Dossinia marmorata.
montana, *Blume, Bijdr.* 413.—Java.
parvifolia, *Lindl. Bot. Reg.* (1839) *Misc.* 19.—Zeylan.
pubescens, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 141.—Burma.
pusilla, *Lindl. Gen. et Sp. Orch.* 489.—Reg. Himal.

CHELIDONIUM, Tourn. ex Linn. Syst. ed. I (1735). *PAPAVERACEAE*, Benth. & Hook. f. i. 53.
aurantiacum, Salisb. Prod. 377 = Glaucium corniculatum.
corniculatum, Donn, Hort. Cant. 100 = Glaucium flavum.
corniculatum, Linn. Sp. Pl. 506 = Glaucium corniculatum.
dahuricum, Hort. ex DC. Prod. i. 123 = majus.
diphyllum, Michx. Fl. Bor. Am. i. 309 = Stylophorum diphyllum.
dodecandrum, Forsk. Fl. Aegypt. Arab. 100 = Roemeria orientalis.
fulvum, Poir. Encyc. Suppl. v. 606 = Glaucium flavum.
glabrum, Mill. Gard. Dict. ed. VIII. n. 5 = Glaucium corniculatum.
Glaucium, Linn. Sp. Pl. 506 = Glaucium flavum.
glaucum, Hill, Veg. Syst. xi. 16 = Glaucium flavum.
grandiflorum, DC. Prod. i. 123 = majus.
haematodes, Moench, Meth. 249 = majus.
hybridum, Linn. Sp. Pl. 506 = Roemeria hybrida.
japonicum, Thunb. Fl. Jap. 221 = Stylophorum japonicum.
laciniatum, Mill. Gard. Dict. ed. VIII. n. 2 = majus.
littorale, Salisb. Prod. 377 = Glaucium flavum.
luteum, Gilib. Fl. Lituan. ii. 211 = majus.
majus, *Linn. Sp. Pl.* 505.—Europ. ; As. Min.
majus, Lour. Fl. Cochinch. i. 330 = sinense.
multifidum, Moç. & Sesse, ex DC. Prod. iii. 344 = Eschscholzia californica.
murale, Ren. Fl. Orne, 180 = majus.
petiolatum, Spreng. ex Steud. Nom. ed. II. i. 347 = Stylophorum diphyllum.
phoeniceum, Lam. Fl. Fr. iii. 169 = Glaucium phoeniceum.
quercifolium, Willemet, Phyt. Encyc. ii. 613 = majus.
rubrum, Poir. Encyc. Suppl. v. 606 = Glaucium corniculatum.
ruderale, Salisb. Prod. 377 = majus.
sinense, *DC. Syst.* ii. 100 (sp. dub.).—China.
umbelliferum, Stokes, Bot. Mat. Med. iii. 180 = majus.

CHELIDONIUM:—
uniflorum, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1846) 169 = Stylophorum japonicum.
violaceum, Lam. Fl. Fr. iii. 169 = Roemeria hybrida.
vitellinum, C. Muell. in Walp. Ann. vii. 87 = Glaucium vitellinum.

CHELIDOSPERMUM, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 162, in syn. (1850) = **Pittosporum**, Banks.
verticillatum, Zipp. ex Blume, l. c. = P. Chelidospermum.

CHELIDURUS, Willd. in Cothen. Disp. Meth. 4 (1790) (Gen. ignot.).
ignotus, J. F. Gmel, Syst. 105.—Hab. ?

CHELIUSIA, Sch. Bip. in Flora, xxiv. (1841) I. Intell. 26, nomen; A. Rich. Tent. Fl. Abyss. i. 380 = **Vernonia**, Schreb. (Compos.).
abyssinica, Sch. Bip. ll. cc. = V. amygdalina.

CHELONANTHERA, Blume, Bijdr. 384. t. 51 (1825) = **Coelogyne**, Lindl. (Orchid.).
gibbosa, Blume, l. c. 383 = Pholidota gibbosa.
incrassata, Blume, l. c. 384 = Coel. incrassata.
longifolia, Blume, l. c. 385 = Coel. longifolia.
miniata, Blume, l. c. = Coel. miniata.
speciosa, Blume, l. c. 384. = Coel. speciosa.
sulphurea, Blume, l. c. 383 = Coel. sulphurea.
ventricosa, Blume, l. c. = Pholidota ventricosa.

CHELONE, Linn. Syst. ed. I (1735). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 939.
CHLONANTHUS, Rafin. Prine. Somiol. 26 (1814).
alba, Moench, Meth. 442 = glabra.
albida, Spreng. Syst. ii. 813 = Pentstemon albidus.
alpina, Spreng. l. c. iv. Cur. Post. 235 = Pentstemon glaber.
angustifolia, H. B. & K. Nov. Gen. et Sp. ii. 365. t. 173 = Pentstemon campanulatus.
angustifolia, Steud. Nom. ed. I. 186 = Pentstemon caeruleus.
atropurpurea, Sweet, Brit. Flow. Gard. Ser. I. t. 235 = Pentstemon campanulatus.
barbata, Cav. Ic. iii. 22. t. 242 = Pentstemon barbatus.
Bradburii, Steud. Nom. ed. I. 186 = Pentstemon grandiflorus.
caerulea, Spreng. Syst. ii. 813 = Pentstemon caeruleus.
campanulata, Cav. Ic. i. 18. t. 29 = Pentstemon campanulatus.
campanuloides, Andr. Bot. Rep. t. 40 = Pentstemon campanulatus.
capitata, Rafin. Med. Fl. ii. 118 = glabra.
centranthifolia, Benth. in Trans. Hort. Soc. Lond. N. S. i. (1835) 481 = Pentstemon centranthifolius.
cheilanthifolia, Paxt. Mag. Bot. ii. t. 50 = praec.
cristata, Spreng. Syst. ii. 813 = Pentstemon cristatus.
Digitalis, Sweet, Brit. Flow. Gard. Ser. I. t. 120 = Pentstemon laevigatus.
elatior, Rafin. Med. Fl. ii. 118 = glabra.
elegans, H. B. & K. Nov. Gen. et Sp. ii. 364 = Pentstemon campanulatus.
erianthera, Steud. Nom. ed. I. 186 = Pentstemon cristatus.
filiformis, Buch.-Ham. ex C. B. Clarke, in DC. Monog. Phan. v. 134 = Boeica filiformis.
formosa, Wendl. Beobacht. 51 = Pentstemon barbatus.
frutescens, Steud. Nom. ed. I. 186 = Pentstemon frutescens.
gentianoides, H. B. & K. Nov. Gen. et Sp. ii. 364. t. 172 = Pentstemon gentianoides.
glabra, *Linn. Sp. Pl.* 611.—Am. bor.
gracilis, Spreng. Syst. ii. 813 = Pentstemon gracilis.
grandiflora, Spreng. l. c. = Pentstemon grandiflorus.
hirsuta, Linn. Sp. Pl. 611 = Pentstemon pubescens.
imberbis, H. B. & K. Nov. Gen. et Sp. ii. 363 = Pentstemon imberbis.
inermis, Poir. in Dict. Sc. Nat. xxxviii. (1825) 385 = Pentstemon imberbis.
laevigata, Pers. Syn. ii. 169 = Pentstemon laevigatus.
lanceolata, Rafin. Med. Fl. ii. 117 = glabra.
latifolia, Buch.-Ham. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 374 = Rhynchotechum latifolium.
latifolia, Muhl. Cat. ex Ell. Sketch, ii. 127 = obliqua.
Lyoni, *Pursh, Fl. Am. Sept.* ii. 737.—Am. bor.
maculata, Rafin. Med. Fl. ii. 117 = glabra.
major, Sims, Bot. Mag. t. 1864 = Lyoni.

CHELONE:—
nemorosa, *Dougl. in Lindl. Bot. Reg.* t. 1211.—Am. bor. occ.
obliqua, *Linn. Syst.* ed. XII. 408.—Am. bor. occ.
Penstemon, Linn. Mant. 415 = Pentstemon pubescens.
Pentstemon, Linn. Sp. Pl. 612, pro parte = Pentstemon laevigatus.
pubescens, Sweet, Hort. Brit. ed. I. 305 = Pentstemon pubescens.
purpurea, Mill. Gard. Dict. ed. VIII. n. 2 = obliqua.
purpurea, Rafin. Med. Fl. ii. 118 = glabra.
rosea, Cerv. ex Sweet, Brit. Flow. Gard. Ser. I. t. 230 = Pentstemon campanulatus.
rubicunda, Buch.-Ham. ex C. B. Clarke, in DC. Monog. Phan. v. 135 = Boeica Griffithii.
ruelloides, Andr. Bot. Rep. t. 34 = Pentstemon barbatus.
ruelloides, Linn. f. Suppl. 271 = Ourisia magellanica.

CHELONOPSIS, Miq. in Ann. Mus. Lugd. Bat. ii. 111 (1865). *LABIATAE*, Benth. & Hook. f. ii. 1204.
moschata, *Miq. l. c.*—Japon.

CHELROSTYLIS, Pritz. Ic. Ind. 255 (1855), sphalm. = **Cheirostylis**, Blume.

CHEMNICIA, Scop. Introd. 139 (1777) = **Strychnos**, Linn. (Loganiac.).

CHEMNIZIA, Steud. Nom. ed. II. i. 347 (1840) = **Strychnos**, Linn. (Loganiac.).

CHENOCARPUS, Neck. Elem. i. 202 (1790) = **Spermacoce**, Linn. (Rubiac.).

CHENOLEA, Thunb. Nov. Gen. 10 (1781). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 59.
BASSIA, All. Misc. Taur. iii. (1766) 177. t. 4.
ECHINOPSILON, Moq. in Ann. Sc. Nat. Sér. II. ii. (1834) 127.
ERIOCHITON, F. Muell. 2nd Gen. Report, 15 (1854), nomen.
LONDESIA, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 40 (1835).
WILLEMETIA, Maerkl. in Schrad. Journ. Bot. iii. (1800) I. 329.
arabica, *Boiss. Diagn.* Ser. I. xii. 97.—Syria ; Arabia.
asphaltica, Boiss. l. c. xiii. 84 = Suaeda asphaltica.
astrocarpa, *F. Muell. Fragm.* xii. 12.—Austral.
biflora, *F. Muell. l. c.* x. 91.—Austral.
canariensis, *Moq. Chenop. Enum.* 96.—Marocco ; Ins. Canar.
carnosa, *Benth. Fl. Austral.* v. 190.—Austral.
Dallachyana, *Benth. l. c.* 191.—Austral.
diacantha, *F. Muell. Fragm.* x. 91.—Austral.
diffusa, *Thunb. Nov. Gen.* 10.—Afr. austr.
echinopsila, *F. Muell. Fragm.* x. 92.—Austral.
enchylaenoides, *F. Muell. l. c.*—Austral.
eurotioides, *F. Muell. ex Benth. Fl. Austral.* v. 191.—Austral.
Forrestiana, *F. Muell. Fragm.* xii. 12.—Austral.
hirsuta, *Archang. Comp. Fl. Ital.* 595.—Italia.
incana, *Jacq. Fragm.* 76.—Afr. austr.
lanata, Jacq. ex Moq. in DC. Prod. xiii. II. 132 = Kochia prostrata.
lanata, Moq. l. c. 129 = canariensis.
Muelleri, *Benth. Fl. Austral.* v. 191.—Austral.
paradoxa, *F. Muell. Fragm.* x. 91.—Austral.
quinquecuspis, *F. Muell. l. c.*—Austral.
sclerolaenoides, *F. Muell. ex Benth. Fl. Austral.* v. 192.—Austral.
salsuginosa, *F. Muell. Fragm.* x. 92.—Austral.
tricornis, *Benth. Fl. Austral.* v. 191.—Austral.
triens pis, *F. Muell. Fragm.* x. 92.—Austral.
tridens, *F. Muell. l. c.* xii. 12.—Austral.

CHENOPODINA, Moq. in DC. Prod. xiii. II. 159 (1849) = **Suaeda**, Forsk. (Chenopod.).
acuminata, Moq. l. c. 161 = S. acuminata.
aestuaria, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 329 = S. maritima.
altissima, Moq. in DC. Prod. xiii. II. 162 = S. altissima.
anonyma, Moq. l. c. 164.—Aegypt.
asphaltica, Boiss. Diagn. Ser. I. xii. 98 = S. asphaltica.
australis, Moq. in DC. Prod. xiii. II. 163 = S. australis.

CHENOPODINA:—
baccifera, Moq. in DC. Prod. xiii. II. 160 = Suaeda baccifera.
bacciformis, Dum. in Bull. Soc. Bot. Belg. vii. (1868) 328 = S. maritima.
Biebersteiniana, Moq. in DC. Prod. xiii. II. 163 = S. microphylla.
dendroides, Moq. l. c. 164 = S. dendroides.
depressa, Moq. l. c. = S. plattensis.
filiformis, Moq. l. c. = S. maritima.
fruticulosa, Moq. l. c. 165 = S. fruticulosa.
glauca, Moq. l. c. 162 = S. glauca.
Hostii, Moq. l. c. 165.—Dalmat.
indica, Wight, Ic. v. t. 1793 = S. maritima.
leiosperma, Moq. in DC. Prod. xiii. II. 162 = S. altissima.
linearis, Moq. l. c. 164 = S. linearis.
linearis, Torr. in Stansbury, Rep. 394 = S. Torreyana.
maritima, Coss. ex Willk. & Lange, Prod. Fl. Hisp. i. 261 = S. setigera.
maritima, Moq. in DC. Prod. xiii. II. 161 = S. maritima.
microphylla, Moq. l. c. = S. microphylla.
microsperma, Moq. l. c. 165 = S. microsperma.
minima, Moq. l. c. 162 = S. minima.
Moquini, Torr. in Pacif. Rail. Rep. vii. 18 = S. Torreyana.
parviflora, Moq. in DC. Prod. xiii. II. 165 = S. microsperma ?
physophora, Moq. l. c. 164 = S. physophora.
prostrata, Moq. l. c. 162 = S. microsperma.
pycnantha, C. Koch, in Linnaea, xxiv. (1851) 90 = Seidlitzia florida.
pygmaea, Moq. in DC. Prod. xiii. II. 161 = S. microsperma ?
salsa, Moq. l. c. 160 = S. salsa.
sativa, Moq. l. c. 165 = S. maritima.
setigera, Moq. l. c. 160 = S. setigera.
spicata, Moq. l. c. 165 = S. maritima.
splendens, Gren. in Gren. & Godr. Fl. Fr. iii. 30 = S. setigera.
tortuosa, Moq. in DC. Prod. xiii. II. 162 = S. maritima ?
vera, Moq. l. c. 163 = S. vera.

CHENOPODIUM, [Tourn.] Linn. Syst. ed. I (1735) ; Gen. ed. I. 67 (1737). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 51.
AGATHOPHYTUM, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 291. t. 10 c.
AMBRINA, Spach, Hist. Vég. Phan. v. 295 (1836).
ANSERINA, Dum. Fl. Belg. 21 (1827).
BLITUM, [Tourn.] Linn. Gen. ed. I. 20 (1737).
BOTRYDIUM, Spach, Hist. Vég. Phan. v. 298 (1836).
LIPANDRA, Moq. Chenop. Enum. 19 (1840).
MOROCARPUS, Adans. Fam. ii. 261 (1763).
OLIGANDRA, Less. in Linnaea, ix. (1834) 199.
OLIGANTHERA, Endl. Gen. Suppl. i. 1377 (1841).
ORTHOSPERMUM, Opiz, Seznam, 70 (1852) ; Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 63.
ORTHOSPORUM, T. Nees, Gen. Fl. Germ. Monochl. n. 58 (1835).
OXYBASIS, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 738.
TELOXYS, Moq. in Ann. Sc. Nat. Sér. II. i. (1834) 289. t. 10.
acerifolium, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 124, nomen.—Rossia.
acuminatum, Schur, Enum. Pl. Transs. 902 = rubrum ?
acuminatum, *Willd. in Ges. Nat. Fr. Neue Schr.* ii. (1799) 124. t. 5. f. 2.—Soongaria.
acutifolium, Hook. & Arn. Bot. Beech. Voy. 207 = acuminatum.
acutifolium, Kit. in Schult. Oestr. Fl. ed. II. i. 458 ; Sm. Comp. Fl. Brit. 42 = polyspermum.
aegyptiacum, *Hasselq. It. Palaest.* 460 = Suaeda baccata.
album, Bosc, ex Moq. in DC. Prod. xiii. II. 61 = Boscianum.
album, *Linn. Sp. Pl.* 219.—Reg. temp. et trop.
alexandrinum, Desf. ex Moq. in DC. Prod. xiii. II. 155 = Suaeda vermiculata.
altissimum, Linn. Sp. Pl. 221 = Suaeda altissima.
ambiguum, K. Br. Prod. i. 407 = glaucum.
ambrosioides, Bert. ex Steud. Nom. ed. II. i. 348 = citriodorum.
ambrosioides, Hance, in Journ. Linn. Soc. xiii. (1872) 86 = acuminatum.
ambrosioides, *Linn. Sp. Pl.* 219.—Reg. temp. et trop.
americanum, Spreng. Syst. i. 922 = Suaeda plattensis.

CHENOPODIUM:—
angulatum, Curt. ex Steud. Nom. ed. I. 187 = hybridum.
angulosum, Lam. Fl. Fr. iii. 249 = hybridum.
angustifolium, Gilib. Exercit. ii. 437 = polyspermum.
angustifolium, Pav. ex Moq. in DC. Prod. xiii. II. 114 = Atriplex linifolia.
antarcticum, *Benth. & Hook. f. Gen.* iii. 52.—Reg. Magellan.
anthelminticum, Bert. ex Steud. Nom. ed. II. i. 348 = chilense.
anthelminticum, Linn. Sp. Pl. 220 = ambrosioides.
arcticum, Hort. ex Moq. in DC. Prod. xiii. II. 86 = Monolepis chenopodioides.
arenarium, Gaertn. Meyer & Scherb. Fl. Wett. i. 356 = Kochia arenaria.
aristatum, *Linn. Sp. Pl.* 221.—Sibir.
arrectum, Desmaz. Cat. Pl. Belg. 60 = polyspermum.
asphalticum, Boiss. Diagn. Ser. I. xii. 98 = Suaeda asphaltica.
astracanium, Ledeb. Cat. Hort. Dorpat. (1819) = rubrum.
atriplicinum, *F. Muell. Fragm.* vii. 11.—Austral.
Atriplicis, Linn. f. Suppl. 171 = purpurascens.
augustanum, All. Fl. Pedem. ii. 108. t. 38 = Kochia prostrata.
augustanum, Hort. Par. ex Moq. in DC. Prod. xiii. II. 135 = Kochia hyssopifolia.
auricomum, *Lindl. in Mitch. Journ. Trop. Austral.* 94. —Austral.
australasicum, Moq. Chenop. Enum. 20 = Rhagodia linifolia.
australe, R. Br. Prod. 407 = Suaeda australis.
baccatum, Labill. Nov. Holl. Pl. i. 71. t. 96 = Rhagodia Billardieri.
baryosmum, Roem. & Schult. Syst. vi. 269 = Salsola foetida.
benghalense, Spielm. ex Steud. Nom. ed. II. i. 348 = Atriplex hortensis?
Berlandieri, *Moq. Chenop. Enum.* 23.—Am. centr.
betaceum, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 125, nomen.—Rossia.
bicolor, Boj. ex Moq. in DC. Prod. xiii. II. 71 = album.
Biebersteinianum, Roem. & Schult. Syst. vi. 267 = Suaeda microphylla.
biforme, Nees, in Lehm. Pl. Preiss. i. 636 = album.
bipinnatifidum, *Moric. ex Moq. in DC. Prod.* xiii. II. 76. —Mexic.
bisaeriale, Menyh. Kalocsa Vidék. Növény. 157 = polyspermum.
blitoides, Lejeune, Fl. Spa, 126 = rubrum.
Blitum, F. Muell. Sel. Pl. Indust. Cult. 49 = capitatum.
bonariense, Hort. Par. ex Moq. in DC. Prod. xiii. II. 67 = purpurascens.
bonariense, Tenore, Ind. Sem. Hort. Neap. (1833) 13 = hircinum.
Bonus-Henricus, *Linn. Sp. Pl.* 218.—Europ.
Boscianum, *Moq. Chenop. Enum.* 21.—Texas.
botrydium, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Botrys.
botryoides, Rafin. ex Moq. in DC. Prod. xiii. II. 75 = C. Botrys.
botryodes, Sm. Engl. Bot. t. 2247 = rubrum.
Botrys, *Linn. Sp. Pl.* 219.—Reg. bor.
Browneanum, Roem. & Schult. Syst. vi. 275 = album.
bryoniaefolium, *Bunge, Del. Sem. Hort. Petrop.* (1876) 10; *ex Trautv. in Act. Hort. Petrop.* ix. (1884) 398. —Sibiria.
calceoliforme, Hook. Fl. Bor. Am. ii. 126 = Suaeda calceoliformis.
californicum, *S. Wats. in Bot. Calif.* ii. 48.—Calif.
camphorataefolium, Pourr. in Mém. Acad. Toul. iii. (1788) 311 = Kochia prostrata.
camphorosmoides, Hort. ex Moq. in DC. Prod. xiii. II. 132 = praec.
candicans, Lam. Fl. Fr. iii. 248 = album.
cantabricum, *Lag. Hort. Matrit.* (1815) 15, nomen.—Hispan.
capitatum, *Aschers. Fl. Brandenb.* 572.—Reg. bor. et austr.
carinatum, *R. Br. Prod.* 407.—Austral.; N. Zel.
carnosulum, *Moq. in DC. Prod.* xiii. II. 64.—Calif.
Caroxylon, D'Urban, in Journ. Linn. Soc. viii. (1865) 272 = Salsola Caroxylon.
carthagenense, Zuccagni, in Roem. Collect. 133 = murale.
catenulatum, Schleich. ex Steud. Nom. ed. II. i. 348 = album.

CHENOPODIUM:—
caudatum, Jacq. Coll. ii. 325; Ic. Pl. Rar. ii. 12. t. 344 = Amarantus viridis.
Chamrium, Buch.-Ham. in Wall. Cat. n. 6953 = murale.
Chaudana, *Buch.-Ham. in Wall. Cat.* n. 6957.—Ind. or.
chilense, Pers. Syn. i. 294 = C. Quinoa.
chilense, *Schrad. Ind. Sem. Hort. Gott.* (1832) 2.—Chili.
chysomelanospermum, Zuccagni, in Roem. Collect. 134 = urbicum.
cinereum, Moq. Chenop. Enum. 27 = Atriplex peruviana.
citriodorum, *Steud. Nom.* ed. II. i. 348.—Ins. Juan Fernand.
Clemente, Spreng. Syst. i. 922 = Suaeda setigera.
concatenatum, Hort. ex Willd. Enum. Hort. Berol. 291 = lanceolatum.
concatenatum, Thuill. Fl. Paris, ed. II. 125 = album.
congestum, Hook. f. in Lond. Journ. Bot. vi. (1847) 280 = Rhagodia Billardieri.
corispermifolium, *Hort. Amstel.* (1814) 14, nomen.—Hab.?
cornutum, *Benth. & Hook. f. ex S. Wats. in Bot. Calif.* ii. 482.—Mexic.
coronopodum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = seq.
Coronopus, *Moq. in DC. Prod.* xiii. II. 76.—Ins. Teneriff.
crassifolium, Bieb. Fl. Taur. Cauc. iii. 176 = Suaeda microphylla.
crassifolium, Desf. Tabl. ed. II. 51 = Sueda maritima.
crassifolium, Hornem. Hort. Hafn. 254 = rubrum.
cristatum, *F. Muell. Fragm.* vii. 11.—Austral.
cuneifolium, *Vahl, in West, Fl. S. Cruc.* 278.—Santa Crux.
cuneifolium, Vent. ex Moq. in DC. Prod. xiii. II. 73 = spathulatum.
deltoideum, Lam. Fl. Fr. iii. 249 = urbicum.
detestans, *T. Kirk, in Trans. N. Z. Inst.* ix. (1877) 550.—N. Zel.
divaricatum, *Fisch. Cat. Hort. Gorenk.* (1808) 24, nomen.—Rossia?
effusum, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 346 = foetidum.
elatum, Shuttl. ex Moq. in DC. Prod. xiii. II. 71 = album.
erosum, Bast. in Desv. Journ. Bot. iii. (1814) 20 = opulifolium.
erosum, R. Br. Prod. 407 = murale.
esculentum, Salisb. Prod. 151 = C. Bonus-Henricus.
exocarpum, *Griseb. in Goett. Abh.* xxiv. (1879) 37.—Reg. Argent.
ficifolium, *Sm. Fl. Brit.* i. 276.—Europ.
filifolium, Krock. Fl. Sil. Suppl. i. 302 = praec.
filiforme, Dum. Fl. Belg. 22 = Suaeda maritima.
filiforme, Moench, Meth. 330 = Suaeda altissima.
flavescens, Roem. & Schult. Syst. vi. 269 = Salsola vermiculata.
flavum, Forsk. Fl. Aegypt. Arab. Suppl. 205 = murale.
foetidum, Lam. Fl. Fr. iii. 244 = C. Vulvaria.
foetidum, *Schrad. in Ges. Naturf. Fr. Berl. Mag.* ii. (1808) 79.—Reg. trop.
foliosum, Aschers. Fl. Brandenb. 572 = capitatum.
Fremonti, *S. Wats. in Bot. King, Exp.* 287.—Am. bor. occ.
frigidum, *Phil. Fl. Atac.* 47.—Chili.
frutescens, *C. A. Mey. in Ledeb. Fl. Alt.* i. 408.—Sibir. altaic.
fruticosum, Bieb. Fl. Taur. Cauc. i. 181 = Suaeda dendroides.
fruticosum, Linn. Sp. Pl. 221 = Suaeda fruticosa.
furfuraceum, Moq. in DC. Prod. xiii. II. 64 = auricomum.
Gandhium, Buch.-Ham. in Wall. Cat. n. 6953 = murale.
giganteum, D. Don, Prod. Fl. Nep. 75 = album.
glandulosum, F. Muell. Fragm. vii. 11 = carinatum.
glaucum, *Linn. Sp. Pl.* 220.—Reg. bor. et austr.
glochidatum, Gill. ex Moq. in DC. Prod. xiii. II. 61 = papulosum.
glomerulosum, Reichb. Fl. Germ. Excurs. 579 = album.
graveolens, Lag. & Rodr. in Anal. Cienc. Nat. v. (1802) 70 = foetidum.

CHENOPODIUM:—
graveolens, *Willd. Enum. Hort. Berol.* i. 290.—Mexic.
guineense, Jacq. Coll. ii. 346; Ic. Pl. Rar. ii. 12. t. 345 = murale.
Guinoa, Krock. Fl. Sil. Suppl. i. 301 = C. Quinoa.
Halimus, Thunb. Prod. Pl. Cap. 48 = Atriplex Halimus.
halophilum, *Phil. in Anal. Univ. Chil.* (1861) I. 67.—Chili.
hastatum, Dum. in Bull. Soc. Bot. Belg. iv. (1865) 339 = Atriplex hastata.
hastatum, *Phil. Fl. Atac.* 47.—Chili.
hircinum, *Schrad. Ind. Sem. Hort. Gott.* (1833) 2.—Bras.
hirsutum, Linn. Sp. Pl. 221 = Kochia hirsuta.
Hookerianum, Moq. in DC. Prod. xiii. II. 68 = murale.
hortense, Raddi, ex Moq. l. c. 161 = Suaeda maritima.
hortense, Roem. & Schult. Syst. vi. 268 = Suaeda hortensis.
Hostii, Ledeb. Fl. Ross. iii. 788 = Suaeda maritima.
humifusum, Kitt. Taschenb. ed. II. 264 = crassifolium.
humifusum, *Zuccagni, in Roem. Collect.* 134.—Europ.
humile, Hook. Fl. Bor. Am. ii. 127 = rubrum.
hybridum, *Linn. Sp. Pl.* 219.—Reg. temp. bor.
hybridum, Lour. Fl. Cochinch. 174 = Loureirei.
ilicifolium, Griff. Notul. iv. 337 = C. Botrys.
incisum, *Poir. Encyc. Suppl.* i. 392.—Am. centr.
intermedium, Mert. & Koch, in Roehl. Deutschl. Fl. ed. III. ii. 297 = urbicum.
Jacquini, Tenore, Fl. Nap. iii. 258 = Suaeda maritima.
lacerum, *Krock. Fl. Sil. Suppl.* i. 304.—Silesia.
laciniatum, Roxb. Hort. Beng. 21; Fl. Ind. ii. 59 = album.
laciniatum, Thunb. Prod. Pl. Cap. 48 = Atriplex laciniata.
lanceolatum, R. Br. Prod. 407 = album.
lanceolatum, Muhl. ex Willd. Enum. Hort. Berol. 291 = album.
lanuginosum, Moench, Meth. 330 = Kochia hyssopifolia.
laterale, [Soland. in] Ait. Hort. Kew. ed. I. i. 313 = murale.
latifolium, *Pall. Reise Statth. Russ. Reich.* i. 280.—Rossia.
leiospermum, DC. Fl. Fr. iii. 390 = album.
leptophyllum, *Nutt. ex Moq. in DC. Prod.* xiii. II. 71. —Am. bor.
leucospermum, Schrad. Ind. Sem. Hort. Gott. (1834) 2 = purpurascens.
lineare, Hort. ex Steud. Nom. ed. II. i. 348 = Kochia prostrata.
linifolium, Roem. & Schult. Syst. vi. 271 = Suaeda linifolia.
littorale, *Moq. Chenop. Enum.* 24.—Austral.
littorale, Thunb. in Act. Soc. Sc. Upsal. vii. (1815) 142 = Atriplex littoralis.
Loureirei, *Steud. Nom.* ed. II. i. 348.—China.
lucidum, Gilib. Exercit. ii. 440 = murale.
macrocarpum, Desv. Journ. Bot. i. (1813) 48 = Suaeda maritima.
macrospermum, *Hook. f. Fl. Antarct.* ii. 341.—Ins. Falkland.
macrospermum, Hort. Ant. ex Moq. in DC. Prod. xiii. II. 84 = rubrum.
marginatum, Spreng. ex Hornem. Hort. Hafn. i. 256 = polyspermum.
maritimum, Friwald. ex Ledeb. Fl. Ross. iii. 783 = Suaeda altissima.
maritimum, Linn. Sp. Pl. 221 = Suaeda maritima.
maritimum, Torr. in Ann. Lyc. N. York, ii. (1828) 239 = Suaeda diffusa.
Matthioli, Bertol. ex Moq. in DC. Prod. xiii. II. 84 = rubrum.
melanospermum, Wallr. Sched. Crit. 112 = urbicum.
mexicanum, *Moq. in DC. Prod.* xiii. II. 70.—Mexic.
micranthum, *Trautv. in Bull. Soc. Nat. Mosc.* xli. (1868) I. 464.—Rossia.
microphyllum, Bieb. Fl. Taur. Cauc. iii. 175 = Suaeda dendroides.
microphyllum, *F. Muell. in Trans. Phil. Inst. Vict.* ii. (1858) 74.—Austral.
microphyllum, *Thunb. Prod. Pl. Cap.* 48.—Afr. austr.
microspermum, Wallr. Sched. Crit. 112 = urbicum.

CHENOPODIUM:—
mucronatum, *Thunb. Prod. Pl. Cap.* 48.—Afr. austr.
multifidum, Linn. Sp. Pl. 220 = Roubieva multifida.
multiflorum, Moq. in DC. Prod. xiii. II. 75 = C. Botrys.
murale, *Linn. Sp. Pl.* 219.—Reg. bor. et austr.
neglectum, *Dum. in Bull. Soc. Bot. Belg.* iv. (1865) 339.—Belgium.
nepalense, Hort. Monsp. ex Moq. in DC. Prod. xiii. II. 74 = C. Botrys.
nigrum, *Rafin. Atl. Journ.* 146.—Am. bor.
nitidum, Spreng. Syst. i. 921 = Salsola nitida.
nitrariaceum, *F. Muell. ex Benth. Fl. Austral.* v. 158.—Austral.
obovatum, *Moq. in DC. Prod.* xiii. II. 73.—Am. austr.
oleraceum, *Schur, Enum. Pl. Transs.* 573.—Transsylv.
olidum, Curt. Fl. Lond. fasc. v. t. 20 = C. Vulvaria.
olidum, *S. Wats. in Proc. Am. Acad.* ix. (1874) 95.—Am. bor.
oppositifolium, Linn. f. Suppl. 172 = Halogeton oppositiflorus.
oppositifolium, Willd. Sp. Pl. i. 1307 = Girgensohnia oppositiflora.
opulaceum, Neck. Delic. Gallo-Belg. i. 130 = album.
opulifolium, *Schrad. in DC. Fl. Fr.* v. 372.—Reg. bor.
paganum, Reichb. Fl. Germ. Excurs. 579 = album.
Pallasianum, Roem. & Schult. Syst. vi. 271 = Kochia hirsuta.
pallidum, Moq. Chenop. Enum. 30 = glaucum?
paniculatum, *Hook. Bot. Misc.* ii. (1831) 237.—Peruv.
paniculatum, Salzm. ex Moq. in DC. Prod. xiii. II. 18 = Microtea paniculata.
papulosum, *Moq. l. c.* 61.—Bras.
parvifolium, Roem. & Schult. Syst. vi. 266 = Suaeda dendroides.
patulum, Mérat, Nouv. Fl. Par. 96 = rubrum.
patulum, *Roth, Nov. Pl. Sp.* 180.—Ind. or.
pauciflorum, Herb. Vind. ex Moq. in DC. Prod. xiii. II. 89 = Exomis axyrioides.
pauciflorum, *Mart. Hort. Monac.* (1829) 56, nomen.—Hab.?
Payco, Molina, Sagg. Chile, ed. II. 118 = Roubieva multifida.
pedunculare, Bertol. Fl. Ital. iii. 32 = album.
petiolare, *H. B. & K. Nov. Gen. et Sp.* ii. 191.—Ecuador.
petiolare, Mart. & Gal. in Bull. Acad. Brux. x. II. (1843) 345 = Cryptocarpus globosus.
physophora, Moq. ex Boiss. Fl. Orient. iv. 939 = Suaeda physophora.
pinnatum, Hort. ex Moq. in DC. Prod. xiii. II. 93 = Atriplex laciniata.
polispermum, Neck. Delic. Gallo-Belg. i. 128 = polyspermum.
polyspermum, *Linn. Sp. Pl.* 220.—Europ.; As. bor.
populifolium, Hort. ex Moq. in DC. Prod. xiii. II. 66 = ficifolium.
portulacoides, Thunb. Fl. Cap. ii. 178 = Atriplex portulacoides.
precatorium, *Dum. in Bull. Soc. Bot. Belg.* iv. (1865) 340.—German.
procerum, *Hochst. ex Moq. in DC. Prod.* xiii. II. 75.—Abyssin.
prostratum, Hort. ex Moq. l. c. 132 = Kochia prostrata.
prostratum, Jacquem. ex Moq. l. c. 72 = glaucum?
prostratum, Roem. & Schult. Syst. vi. 272 = Suaeda prostrata.
pseudo-botrys, *Hort. ex Heynh. Nom.* ii. 138, nomen.—Hab.?
Pumilio, *R. Br. Prod.* i. 407.—Austral.
punctulatum, Scop. Delic. Insub. i. 26. t. 11 = purpurascens.
purpurascens, Buch.-Ham. in Wall. Cat. n. 6956 = album.
purpurascens, Jacq. Hort. Vindob. iii. 43. t. 80 = album?
pusillum, *Hook. f. Handb. New Zeal. Fl.* 231.—N. Zel.
pusillum, *Bertol. in Nov. Comm. Bonon.* iv. (1842) 413.—Fl. Euphrat.
pygmaeum, Menyh. Kalocsa Vidék. Növény. 154 = rubrum.
quercifolium, Hort. Monsp. ex Moq. in DC. Prod. xiii. II. 74 = graveolens.
Quinoa, *Willd. Sp. Pl.* i. 1301.—Am. austr.
radiatum, Schrad. Neues Journ. iii. III. (1809) 85 = Cycloloma platyphyllum.

CHENOPODIUM:—
retusum, *Juss. ex Moq. in DC. Prod.* xiii. II. 73.—Bras.
rhadinostachyum, *F. Muell. in Wing, South Sc. Record*, ii. (1882) 98.—Austral.
rhombifolium, Mnhl. ex Willd. Enum. Hort. Berol. 288 = urbicum.
riparium, Boeningh. ex Moq. in DC. Prod. xiii. II. 71 = album.
rubricaule, Schrad. ex Moq. l. c. 67 = purpurascens.
rubrum, Forsk. ex Steud. Nom. ed. I. 188 = murale.
rubrum, *Linn. Sp. Pl.* 218.—Europ.; As. bor.
ruderale, *Kit. ex Moq. in DC. Prod.* xiii. II. 77.—Hungaria.
ruderale, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 69 = C. Bonus-Henricus.
sagittatum, Lam. Fl. Fr. ii. 244 = C. Bonus-Henricus.
salsum, Guss. Fl. Sic. Prod. i. 299 = Suaeda maritima.
salsum, Hohen. ex Ledeb. Fl. Ross. iii. 782 = Suaeda altissima.
salsum, Linn. Sp. Pl. 221 = Suaeda salsa.
Sancta-Maria, Vell. Fl. Flum. 126; iii. t. 104 = ambrosioides.
sandwicheum, *Moq. Chenop. Enum.* 28.—Ins. Sandvic.
Schraderianum, Roem. & Schult. Syst. vi. 260 = foetidum.
scoparia, Linn. Sp. Pl. 221 = Kochia scoparia.
secundiflorum, Viv. Fl. Lib. Spec. 67 = aristatum.
sepium, *Mey. in Abh. Boehm. Ges. Wiss.* (1787) 320.—Bohemia.
sericeum, Spreng. Syst. i. 921 = Kochia sericea.
sericeum, Vitm. Summa Veg. ii. 98 = Kochia prostrata.
serotinum, *Forsk. Fl. Aegypt. Arab.* 205.—Arab.
serotinum, Ledeb. Fl. Alt. i. 405 = album.
serotinum, Linn. Cent. Pl. ii. 12 = ficifolium.
serotinum, Suter, Fl. Helv. i. 148 = hybridum.
setigerum, DC. Cat. Hort. Monsp. 94 = Suaeda setigera.
simplex, *Rafin. Atl. Journ.* 146.—Am. bor.
sinense, Hort. ex Moq. in DC. Prod. xiii. II. 60 = aristatum.
sinense, Willem. Ess. Med. iii. 233 = Kochia sericea.
sinuatum, Thunb. Fl. Cap. ii. 178 = Atriplex laciniata.
spathulatum, *Sieber, ex Moq. in DC. Prod.* xiii. II. 73.—Bras.
spicatum, *Lavy, Veg. Orig.* 10.—Hab.?
spicatum, Roem. & Schult. Syst. vi. 271 = Suaeda maritima.
spinacifolium, Stokes, Bot. Mat. Med. ii. 14 = C. Bonus-Henricus.
spinosum, Hook. Fl. Bor. Am. ii. 127 = Grayia polygaloides.
stellatum, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 146.—Mexic.
stramoniifolium, Cheval. Fl. Par. ii. 383 = hybridum.
strictum, *Roth, Nov. Pl. Sp.* 180.—Ind. or.
subspicatum, Nutt. Gen. Am. i. 199 = Atriplex hastata.
suffruticosum, Willd. Enum. Hort. Berol. 290 = ambrosioides.
tenerrimum, *Krock. Fl. Sil. Suppl.* i. 305.—German.
tenue, Colla, in Mem. Acc. Torin. xxxix. (1836) 9. t. 50 = Roubieva multifida.
tomentosum, *Thou. Fl. Tristan.* 38.—Ins. Tristan d'Acugn.
Tournefortianum, Hort. ex Moq. in DC. Prod. xiii. II. 158 = Suaeda altissima.
Tournefortii, Hort. ex Moq. l. c. = praec.
triandrum, *Forst. f. Prod.* 21.—N. Zel.
triangulare, *R. Br. Prod.* 407.—Austral.
triangulare, Dulac, Fl. Hautes-Pyr. 171 = C. Bonus-Henricus.
triangulare, Forsk. Fl. Aegypt. Arab. Suppl. 205 = murale.
triangularifolia, Gilib. Exercit. ii. 437 = C. Bonus-Henricus.
trifidum, Trev. Ind. Sem. Vratisl. (1829) = Monolepis chenopodioides.
trigonon, Roem. & Schult. Syst. vi. 275 = triangulare.
trigynum, Roem. & Schult. l. c. 268 = Suaeda altissima.
trilobatum, Hort. ex Moq. in DC. Prod. xiii. II. 76 = foetidum.
trilobum, Hort. ex Moq. l. c. = foetidum.
trilobum, Schult. ex Moq. l. c. 66 = ficifolium.
tripteris, Dum. Fl. Belg. 22 = Kochia tripteris.
tumidum, *Kit. Cat. Krocm.* (1814) *Suppl.* 3, nomen.—Hab.?

CHENOPODIUM:—
Tweedii, *Moq. in DC. Prod.* xiii. II. 63.—Bras.
urbicum, *Linn. Sp. Pl.* 218.—Europ.; As. bor.
Vachellii, Hook. & Arn. Bot. Beech. Voy. 269 = acuminatum.
variegatum, Gouan, in Cat. Hort. Taur. (1810) = ambrosioides.
vestitum, Thunb. Prod. Pl. Cap. 48 = Atriplex glauca.
villosum, Hort. ex Moq. in DC. Prod. xiii. II. 132 = Kochia prostrata.
villosum, Lam. Encyc. i. 196 = Kochia hyssopifolia.
villosum, Spreng. Syst. i. 924 = Suaeda microphylla.
virens, *Wender. Sem. Marb.* (1836), nomen.—Hab.?
virgatum, *Thunb. in Nov. Act. Soc. Sc. Ups.* vii. (1815) 143.—Japon.
virginicum, Linn. Sp. Pl. 222 = aristata.
viride, Curt. Fl. Lond. fasc. ii. t. 16 = ficifolium.
viride, Linn. Sp. Pl. 219 = album.
viride, Loisel. Fl. Gall. i. 145 = opulifolium.
vulgare, Gueldenst. ex Ledeb. Fl. Ross. iii. 700 = album.
vulpinum, Buch.-Ham. in Wall. Cat. n. 6954 = album, ambrosioides.
Vulvaria, *Linn. Sp. Pl.* 220.—Europ.; As. bor.
Wolffii, *Simk. in Termész. Füzetek*, iii. (1879) 164.—Hungar.
Ziegleri, Perrot. ex Moq. in DC. Prod. xiii. II. 24 = Limeum linifolium.
zosteraefolium, Hook. Fl. Bor. Am. ii. 127 = Atriplex Gmelini.

CHEOBULA, Vell. Fl. Flum. iv. t. 146 (1827) = **Cleobula**, Vell. (Inc. sed.).

CHERINA, Cass. in Dict. Sc. Nat. viii. 437 (1817) = **Chaetanthera**, Ruiz & Pav. (Compos.).
microphylla, Cass. l. c. 438 = Chaet. linifolia.

CHERLERIA, Hall. It. Helv. ii. n. 79. t. 1 (1740); ex Linn. Gen. ed. II. 196 (1742) = **Arenaria**, Rupp. (Caryophyll.).
bipartita, Fisch. ex Steud. Nom. ed. II. i. 349, nomen.—Sibir. baical.
bisulca, Bartl. in Presl, Rel. Haenk. ii. 12.—Peruv.
caespitosa, Lam. Fl. Fr. iii. 46 = A. Cherleria.
dicranoides, Cham. in Linnaea, i. (1826) 63 = Stellaria dicranoides.
grandiflora, D. Don, Prod. Fl. Nep. 214 = A. globiflora.
imbricata, Ser. in DC. Prod. i. 421 = A. aretioides.
juniperina, D. Don, Prod. Fl. Nep. 214 = A. densissima.
laevis, Bartl. in Presl, Rel. Haenk. ii. 12.—Peruv.
nitida, Bartl. l. c.—Peruv.
octandra, Sieber, ex Spreng. Syst. ii. 416 = A. aretioides.
pedunculatis, Bunge, ex Steud. Nom. ed. II. i. 349 = *bipartita*.
sedoides, Forsk. Fl. Aegypt. Arab. p. lxvi = A. geniculata.
sedoides, Linn. Sp. Pl. 425 = A. Cherleria.
sedoides, Sibth. & Sm. Fl. Graec. Prod. i. 306 = A. trichocalycina.
sedoides, Turcz. in Bull. Soc. Nat. Mosc. xv. (1842) 608 = Stellaria petraea.
sibirica, Regel & Tiling. in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 72.—Sibir. or.
stellata, Clarke, Trav. iv. 211 (= *Alsine parnassica*).—Graecia.

CHEROPHILUM, Nocca. in Usteri, Ann. Bot. v. (1793) 51 = **Chaerophyllum**, Linn. (Umbellif.).

CHERSYDRIUM, Schott, in Oestr. Bot. Zeitschr. xv. (1865) 72 = **Dracontium**, Linn. (Aroid.).
Jararaca, Schott, l. c. 73 = D. asperum.

CHESNEA, Scop. Introd. 119 (1777) = **Cephaëlis**, Sw. (Rubiac.).

CHESNEYA, Bertol. in Nov. Comm. Bonon. v. (1842) 427 (Misc. Bot. i. 17. t. 2) = **Pimpinella**, Linn. (Umbellif.).
daucoides, Bertol. l. c. = P. eriocarpa.

CHESNEYA, Lindl. in Endl. Gen. 1275 (1840) = **Calophaca**, Fisch. (Legumin.).
astragalina, Jaub. & Spach, Illustr. i. 94.—Persia.
euphratensis, [Lindl. in] Chesn. Narrat. Euph. (1868) t. ad p. 439 = *rytidosperma*.

CHESNEYA :—
Kotschyii, Boiss. Diagn. Ser. I. vi. 34.—Persia.
microphylla, Jaub. & Spach, Illustr. i. 94 = *rytidosperma*.
mongolica, Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 462.—Mongolia.
Oliverii, Jaub. & Spach, Illustr. i. 94.—Mesopotam.
parviflora, Jaub. & Spach, l. c. 95.—Persia.
rytidosperma, Jaub. & Spach, l. c. 93.—Armenia.
turkestanica, Franch. in Ann. Sc. Nat. Sér. VI. xv. (1883) 253.—Turkestan.
vaginalis, Jaub. & Spach, Illustr. i. 96. t. 48.—Mesopotam.
velutina, Jaub. & Spach, l. c. 94 = *rytidosperma*.

CHESNYA, Reichb. Nom. 148 (1841) = praec.

CHETARIA, Steud. Nom. ed. II. ii. 521 (1841) = seq.

CHETASTRUM, Neck. Elem. i. 108 (1790) = **Scabiosa**, Tourn. (Dipsac.).

CHETOCRATER, Rafin. Sylva Tellur. 149 (1838) = **Casearia**, Jacq. (Samydac.).
hirta, Rafin. l. c. = Cas. hirta.
javitensis, Rafin. l. c. = Cas. javitensis.
tinifolia, Rafin. l. c. = Cas. tinifolia.

CHETROPIS, Rafin. Fl. Tellur. iii. 80 (1836) = **Arenaria**, Rupp. (Caryophyll.).
setacea, Rafin. l. c. = A. rostrata.

CHEVALIERIA, Gaudich. Voy. Bonite Bot. t. 61, 62 (1852?) = **Aechmea**, Ruiz & Pav. (Bromel.).
crocophylla, E. Morr. in Belg. Hortic. (1885) 81.—Bras.
Germinyana, Carr. in Rev. Hortic. (1881) 230.—N. Granat.
grandiceps, Griseb. in Goett. Abh. xxiv. (1879) 329 = Ae. brasiliensis.
lingulata, Griseb. Fl. Brit. W. Ind. 591 = Ae. lingulata.
ornata, Gaudich. Voy. Bonite Bot. t. 62 = Ae. ornata.
sphaerocephala, Gaudich. l. c. t. 61 = Ae. sphaerocephala.
Veitchii, E. Morr. in Belg. Hortic. xxviii. (1878) 177. t. 9 = Ae. Veitchii.

CHEVALLIERA, Carr. in Rev. Hortic. (1881) 230 = praec.

CHEVREULIA, Cass. in Bull. Soc. Philom. (1817) 69; et Dict. Sc. Nat. viii. 516 (1817). *COMPOSITAE*, Benth. & Hook. f. ii. 303.
LEUCOPODUM, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 124.
acuminata, *Less. in Linnaea*, v. (1830) 261.—Am. austr.
filiformis, Hook. & Arn. in Hook. Comp. Bot. Mag. i. (1835) 102 = acuminata.
gnaphalioides, *D. Don, ex Hook. & Arn. in Hook. l. c.*—Chili.
lanceolata, *Remy, in C. Gay, Fl. Chil.* iii. 333.—Chili.
longipes, Wedd. Chlor. And. i. 157 = acuminata.
lycopodioides, *DC. Prod.* vii. 45.—Ins. Falkland.
nivea, *Phil. in Anal. Univ. Chil.* ii. (1862) 380.—Chili.
pusilla, DC. Prod. vii. 45 = stolonifera.
stolonifera, *Cass. in Bull. Soc. Philom.* (1817) 69.—Am. austr.
Thouarsii, Remy, in C. Gay, Fl. Chil. iii. 332. t. 37. f. 2 = stolonifera.
xeranthemoides, *D. Don, ex Hook. Comp. Bot. Mag.* i. (1835) 102.—Chili.

CHEYNIA, J. Drumm. ex Harv. in Hook. Kew Journ. vii. (1855) 56 = **Balaustion**, Hook. (Myrtac.).
pulchella, J. Drumm. l. c. = B. pulcherrimum.

CHIAZOSPERMUM, Bernh. in Linnaea, viii. (1833) 465 = **Hypecoum**, Linn. (Fumariac.).
erectum, Bernh. l. c. xii. (1838) 662 = H. erectum.
lactiflorum, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 142.—Soongaria.
pendulum, Bernh. ex Edgew. in Journ. Linn. Soc. vi. (1862) 183 = H. procumbens.

CHIBACA, Bertol. f. in Mem. Acc. Bolog. iv. (1853) 545, nomen (= Gen. dub. LAURIN.) Benth. & Hook. f. iii. 150.
salutaris, *Bertol. f. l. c.; et in Rosenth. Syn. Pl. Diaphor.* i. 238.—Afr. austr.

CHICHAEA, Presl, Rel. Haenk. ii. 140 (1836) = **Sterculia**, Linn.
acerifolia, Presl, l. c. 141 = S. acerifolia.

CHICHARRONIA, A. Rich. Ess. Fl. Cub. 529. t. 43 (1845) = **Terminalia**, Linn. (Combretac.).
intermedia, A. Rich. l. c.—Ins. Cuba.

CHICKRASSIA, A. Juss. in Mém. Mus. Par. xix. (1830) 251. t. 22 (Chukrasia). *MELIACEAE*, Benth. & Hook. f. i. 339.
PLAGIOTAXIS, Wall. Cat. (nomina) nn. 1269, 1270.
Nimmonii, R. Grah. in Wight, Illustr. 148 = tabularis.
tabularis, *A. Juss. in Mém. Mus. Par.* xix. (1830) 251.—Ind. or.; Malaya.
trilocularis, M. Roem. Syn. Hesper. i. 135 = tabularis.
velutina, Wight & Arn. Prod. 123 = tabularis.

CHICOINAEA, Comm. ex DC. Prod. iv. 462 (1830) = **Psathura**, Comm. (Rubiac.).

CHILECHIUM, Pfeiff. Nom. i. 1. 713 (1873) = CHILOCHIUM, Rafin. = **Echiochilon**, Desf. (Boragin.).

CHILERANTHEMUM, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 166. *ACANTHACEAE*, Benth. & Hook. f. ii. 1119.
trifidum, *Oerst. l. c.* 167. t. 3. f. 30–33.—Mexic.

CHILIADENUS, Cass. in Dict. Sc. Nat. xxxiv. 34 (1825) = **Jasonia**, Cass. (Compos.).
camphoratus, Cass. l. c. 35 = J. glutinosa.
glutinosus, Fourr. in Ann. Soc. Linn. Lyon, N.S. xvii. (1869) 93 = J. glutinosa.

CHILIANDRA, Griff. Notul. iv. 150 (1854) = **Rhynchotechum**, Blume (Gesnerac.).
obovata, Griff. l. c. = R. latifolium.

CHILIANTHUS, Burch. Trav. i. 94 (1822). *LOGANIACEAE*, Benth. & Hook. f. ii. 793.
SEMNOS, Rafin. Sylva Tellur. 161 (1838).
arboreus, Benth. in DC. Prod. x. 435 = oleaceus.
corrugatus, *Benth. l. c.* 436.—Afr. austr.
dysophyllus, *Benth. l. c.*—Afr. austr.
lobulatus, *Benth. l. c.*—Afr. austr.
oleaceus, *Burch. Trav.* i. 94.—Afr. austr.
triphyllus, E. Mey. Zwei Pfl. Docum. 172 = Nuxia floribunda.

CHILIOCEPHALUM, Benth. in Hook. Ic. Pl. t. 1137 (1873). *COMPOSITAE*, Benth. & Hook. f. ii. 300.
KRALIKIA, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 151 (1867).
myriocephalum, Baill. Hist. Pl. viii. 173 = Schimperi.
Schimperi, *Benth. in Hook. Ic. Pl.* t. 1137.—Afr. trop.
Schultzii, Buching. ex Baill. Hist. Pl. viii. 173 = Schimperi.

CHILIOPHYLLUM, DC. Prod. v. 554 (1836) = **Zaluzania**, Pers. (Compos.).
globosum, DC. l. c. = Z. globosa.

CHILIOPHYLLUM, Phil. in Linnaea, xxxiii. (1864–65) 132. *COMPOSITAE*, Benth. & Hook. f. ii. 258.
densifolium, *Phil. l. c.* 133.—Chili.

CHILIOTRICHUM, Cass. in Dict. Sc. Nat. viii. 576 (1817). *COMPOSITAE*, Benth. & Hook. f. ii. 276.
TROPIDOLEPIS, Tausch, in Flora, xii. (1829) 68.
amelloides, DC. Prod. v. 216 = seq.
amelloideum, *Cass. in Dict. Sc. Nat.* viii. 577.—Reg. Magellan.
Darwinii, Hook. f. Fl. Antarct. 305 = Nardophyllum Darwini.
Feliciae, Hombr. & Jacquin. ex Decne. Bot. Voy. Astrol. et Zél. 52 = amelloidum.
humile, Hook. f. Fl. Antarct. 304 = Nardophyllum humile.
Kingii, *Hook. f. l. c.* 305 = Nardophyllum Kingii.
ovatifolium, *Hombr. & Jacquin. ex Decne. Bot. Voy. Astrol. et Zél. Dicot. Phan.* t. 28.—Hab.?
rosmarinifolium, *Less. in Linnaea*, vi. (1831) 109.—Chili.
virgatum, *Phil. in Anal. Univ. Chil.* (1873) 485.—Chili.

CHILMORIA, Buch.-Ham. in Trans. Linn. Soc. xiii. (1822) 500 = **Gynocardia**, R. Br. (Bixin.).
dodecandra, Buch.-Ham. l. c.—G. odorata.
pentandra, Buch.-Ham. l. c. 501 = Hydnocarpus inebrians.

CHILOCALYX, Klotzsch, in Peters, Reise Mossamb. Bot. 154 (1861) = **Cleome**, Linn. (Capparid.).
macrophyllus, Klotzsch, l. c. 155 = Cleome Chilocalyx.
tenuifolius, Klotzsch, l. c. = Cleome Chilocalyx.

CHILOCALYX, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 588 = **Atalantia**, Correa (Rutac.).
elliptica, Turcz. l. c. = A. missionis.

CHILOCARPUS, Blume, Cat. Gew. Buitenz. 22 (1823); et Bijdr. 1025 (1826). *APOCYNACEAE*, Benth. & Hook. f. ii. 691.
atroviridis, *Blume, Mus. Bot. Lugd. Bat.* i. 153.—Burma; Malaya.
australis, *F. Muell. Fragm.* ii. 90.—Austral.
ceylanicus, *Wight, Ic.* t. 1288 = Willughbeia ceylanica.
compositus, *Blume, Mus. Bot. Lugd. Bat.* i. 152.—Java.
costatus, *Miq. Fl. Ind. Bat.* ii. 393.—Sumatra.
cuspidatus, Benth. & Hook. f. Gen. ii. 698, in nota sub Hunteria = Melodinus orientalis.
decipiens, *Hook. f. Fl. Brit. Ind.* iii. 627.—Malacca.
densiflorus, *Blume, Mus. Bot. Lugd. Bat.* i. 152.—Sumatra.
denudatus, Blume, Bijdr. 1025 = atroviridis.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 552.—Sumatra.
enervis, *Hook. f. Fl. Brit. Ind.* iii. 626.—Malacca.
flavescens, Kew Gard. Report (1880) 47, ex Fl. Brit. Ind. iii. 625 = Willughbeia flavescens.
globuliferus, *Blume, Mus. Bot. Lugd. Bat.* i. 152.—Borneo.
Maingayi, *Dyer, ex Hook. f. Fl. Brit. Ind.* iii. 627.—Malacca.
malabaricus, Bedd. Ic. Pl. Ind. Or. i. 33 = atroviridis.
suaveolens, *Blume, Cat. Gew. Buitenz.* 23; *et Bijdr.* 1025.—Java.
vernicosus, *Blume, Mus. Bot. Lugd. Bat.* i. 152.—Borneo.

CHILOCHIUM, Rafin. in Ann. Gén. Sc. Phys. viii. (1821) 269 = **Echiochilon**, Desf. (Boragin.).

CHILOCHLOA, Beauv. Agrost. 37. t. 7. f. 2 (1812) = **Phleum**, Linn. (Gramin.).
annua, Roem. & Schult. Syst. ii. 410 = P. asperum.
arenaria, Beauv. Agrost. 37 = P. arenarium.
aspera, Beauv. l. c. = P. asperum.
Boëmerii, Beauv. l. c. = P. Boehmeri.
collina, Schur, Enum. Pl. Transs. 729 = P. phalaroides.
cuspidata, Beauv. Agrost. 37 = P. Michelii.
cuspidata, Schur, Enum. Pl. Transs. 729 = P. phalaroides.
dentata, Trin. Gram. Unifl. 168 = Phalaris dentata.
explicata, Roem. & Schult. Syst. ii. 410 = Heleochloa alopecuroides.
Gerardi, Bess. ex Schult. Mant. ii. 206 = Alopecurus Gerardi.
hirsuta, S. F. Gray, Nat. Arr. Brit. Pl. ii. 139 = P. Michelii.
hispida, Beauv. Agrost. 158, nomen = Arthraxon ciliaris.
Michelii, Trin. Gram. Unifl. 167 = P. Michelii.
paniculata, Beauv. Agrost. 37 = P. asperum.
ventricosa, Beauv. ex Steud. Nom. ed. II. i. 350 = Gastridium australe?

CHILODIA, R. Br. Prod. 507 (1810) = **Prostanthera**, Labill. (Labiat.).
australis, Loud. Hort. Brit. Suppl. i. 585.—Austral.
scutellarioides, R. Br. Prod. 507 = P. empetrifolia.

CHILOGLOSSA, Oerst. in Kjoeb. Vidensk. Meddel. (1854) 160 = **Dianthera**, Linn. (Acanthac.).
glabra, Oerst. l. c. 161 = D. glabra.
secunda, Pritz. Ic. Ind. ii. 77 = praec.

CHILOGLOTTIS, R. Br. Prod. 322 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 614.
cornuta, *Hook. f. Fl. Antarct.* 69.—N. Zel.
diphylla, *R. Br. Prod.* 323.—Austral.
formicifera, *Fitzgerald, Austral. Orchids*, i. III. (1877), —Austral.
Gunnii, *Lindl. Gen. et Sp. Orch.* 387.—Austral.
trapeziformis, *Fitzgerald, Austral. Orchids*, i. III. 1877.—Austral.
trilabra, *Fitzgerald, in Journ. Bot.* xxi. (1883) 204.—Austral.
Traversii, F. Muell. Veg. Chath.-Isl. 51 = Caladenia bifolia.

CHILOPORUS, Naud. in Ann. Sc. Nat. Sér. III. iv. (1843) 57 = **Miconia**, Ruiz & Pav. (Melastomac.).
andinus, Naud. l. c. = M. andina.

CHILOPSIS, D. Don, in Edinb. Phil. Journ. ix. (1823) 261. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1041.
glutinosa, *Engelm. in Bot. Wislizen. Mexic. Exped.* 94.—Mexic. bor.
linearis, Sweet, Hort. Brit. ed. I. 283 = saligna.
saligna, *D. Don, in Edinb. Phil. Journ.* ix. (1823) 261. —Mexic.

CHILOSCHISTA, Lindl. Bot. Reg. sub t. 1522 (1832) = **Sarcochilus**, R. Br. (Orchid.).
usneoides, Lindl. l. c. = S. usneoides.

CHILOSTIGMA, Hochst. in Flora, xxiv. (1841) 372 = **Aptosimum**, Burch. (Scrophularin.).
pumilum, Hochst. l. c. = A. pumilum.

CHIMANTHUS, Rafin. Fl. Ludov. 26 (1817) = **Prunus**, Tourn. (Rosac.).
amygdalina, Rafin. l. c. = P. caroliniana.

CHIMAPHILA, Pursh, Fl. Am. Sept. i. 279 (1814). *ERICACEAE*, Benth. & Hook. f. ii. 603.
CHIMAZA, R. Br. ex DC. Prod. vii. 775 (1839).
PSEVA, Rafin. in Journ. Phys. Sc. lxxxix. (1819) 261, in obs.
astyla, *Maxim. in Bull. Acad. Pétersb.* xi. (1867) 434. —Japon.
corymbosa, Pursh, Fl. Am. Sept. i. 300 = umbellata.
cymosa, J. & C. Presl, Fl. Cech. 89 = umbellata.
Durandi, *Rafin. Atl. Journ.* 119.—Am. bor.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 165.—Japon.
maculata, *Pursh, Fl. Am. Sept.* i. 300.—Am. bor.
Menziesii, *Spreng. Syst.* ii. 317.—Am. bor. occ.
umbellata, *Nutt. Gen. Am.* i. 274.—Europ.; As. bor.; Am. bor.

CHIMARHIS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 84 = seq.

CHIMARRHIS, Jacq. Stirp. Am. 61 (1763). *RUBIACEAE*, Benth. & Hook. f. ii. 45.
cymosa, *Jacq. l. c.*—Ins. Martinic.
Goudotii, *Baill. Adansonia*, xii. (1879) 307.—N. Granat.
paraënsis, *Baill. l. c.* 308.—Bras.
turbinata, *DC. Prod.* iv. 404.—Guiana.

CHIMAZA, R. Br. ex DC. Prod. vii. 775 (1839) = **Chimaphila**, Pursh (Ericac.).

CHIMOCARPUS, Baill. Dict. i. 784 (1879) = CHYMOCARPUS, D. Don = **Tropaeolum**, Linn. (Geraniac.).

CHIMONANTHUS, Lindl. Bot. Reg. sub t. 404 (1819); et t. 451 (1820). *CALYCANTHACEAE*, Benth. & Hook. f. i. 16.
MERATIA, Loisel. Herb. Gén. Amat. iii. t. 173 (1819).
fragrans, *Lindl. l. c.* t. 451.—Japon.
grandiflorus, Steud. Nom. ed. II. i. 350 = fragrans.
luteus, Hort. ex Bielawski, in Ann. Soc. Linn. Maine-et-Loire, ix. (1867) 96 = fragrans.
parviflorus, Rafin. Alsog. Am. 6 = fragrans.
praecox, Lindl. Bot. Reg. sub t. 404 = fragrans.
smilacifolius, Steud. Nom. ed. II. i. 350, sphalm. = Myropyrum smilacifolium.
verus, Hort. ex Bielawski, in Ann. Soc. Linn. Maine-et-Loire, ix. (1867) 96 = fragrans.

CHIMOPHILA, Radius, Diss. Pyrol. 7, 33 (1821) = **Chimaphila**, Pursh (Ericac.).

CHINCHARRONIA, A. Rich. Fl. Cub. Fanerog. ii. 328 (1853) = CHICHARRONIA, A. Rich. = **Terminalia**, Linn. (Combretac.).

CHINCHONA, Howard, in Rep. Bot. Congr. London, (1866) 221 = **Cinchona**, Linn. (Rubiac.).

CHIOCOCCA, P. Br. ex Linn. Syst. ed. X. 917 (1759). *RUBIACEAE*, Benth. & Hook. f. ii. 105.
SIPHONANDRA, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 581.
alternifolia, Linn. Syst. ed. XII. 165 = Cestrum vespertinum.

CHIOCOCCA:—
anguifuga, Mart. Reise Bras. i. 306, 554; Mat. Med. Bras. 17. t. 5 = brachiata.
axillaris, Moç. ex DC. Prod. iv. 483 (= *Margaris nudiflora*).—Mexic.
barbata, *Forst. f. Prod.* 16.—Ins. Pacif.
barbigera, D. Dietr. Syn. Pl. i. 777 = Symphoricarpos microphyllus.
brachiata, *Ruiz & Pav. Fl. Per.* ii. 67.—Am. trop.
coriacea, *Mart. & Gal. in Bull. Acad. Brux.* xi. i. (1844) 231.—Mexic.
densifolia, Mart. Reise Bras. i. 544; Spec. Mat. Med. Bras. 17. t. 6 = brachiata.
floridana, Rafin. Alsog. Am. 75 = racemosa.
javanica, *Blume, Bijdr.* 968.—Java.
latifolia, Rafin. Alsog. Am. 75 = racemosa.
macrocarpa, *Mart. & Gal. in Bull. Acad. Brux.* xi. i. (1844) 230.—Mexic.
malabarica, *Dennst. Schluess. Hort. Malab.* 37.—Ind. or.
nitida, *Benth. in Hook. Journ. Bot.* iii. (1841) 236.—Guiana.
nocturna, Jacq. Enum. Pl. Carib. 16 = Cestrum nocturnum.
nudiflora, D. Dietr. Syn. Pl. i. 777 (= *Margaris nudiflora*).—Mexic.
odorata, *Hook. & Arn. Bot. Beech. Voy.* 65.—Ins. Societ.
opulina, Spreng. Syst. i. 756 = Pavetta opulina.
paniculata, Hoffm. ex Roem. & Schult. Syst. v. 202 = brachiata.
paniculata, Linn. f. Suppl. 145 = Psychotria paniculata.
parviflora, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 202 = brachiata.
parvifolia, Wullschl. ex Griseb. Fl. Brit. W. Ind. 337 = racemosa.
phaenostemon, *Schlecht. in Linnaea*, ix. (1834) 594.—Mexic.
pubescens, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 202 = brachiata.
racemosa, H. B. & K. Nov. Gen. et Sp. iii. 352 = brachiata.
racemosa, *Linn. Syst.* ed. X. 917; *Jacq. Select. Am.* 68.—Ind. occ.
racemosa, Schlecht. & Cham. in Linnaea, v. (1830) 166 = phaenostemon.
sambucina, Spreng. Syst. i. 756 = Pavetta sambucina.
spicata, Blume, Bijdr. 958 = Xanthophytum spicatum.
staminea, *Mart. & Gal. Bull. Acad. Brux.* xi. i. (1844) 231.—Mexic.
triflora, Spreng. Syst. i. 756 = Pavetta triflora.
trisperma, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 219.—Ins. Galapag.

CHIOGENES, Salisb. in Trans. Hort. Soc. ii. (1817) 94. *VACCINIACEAE*, Benth. & Hook. f. ii. 577.
GLYCIPHYLLA, Rafin. in Am. Monthly Mag. (1819) 192.
LASIERPA, Torr. in Geol. Rep. N. York (1839), 152.
PHALEROCARPUS, G. Don, Gen. Syst. iii. 641 (1834).
hispidula, Torr. & Gray, ex Torr. Fl. N. York, i. 450. t. 68 = serpyllifolia.
japonica, *A. Gray, Syn. Fl. N. Am.* ii. i. 26.—Japon.
serpyllifolia, *Salisb. in Trans. Hort. Soc.* ii. (1817) 94. —Am. bor.; Japon.

CHIONACHNE, R. Br. in Benn. Pl. Jav. Rar. 15 (1838). *GRAMINEAE*, Benth. & Hook. f. iii. 1113.
barbata, *R. Br. l. c.* 18.—Ind. or.
cyathopoda, *F. Muell. ex Benth. Fl. Austral.* vii. 516. —Austral.
Koenigii, Thw. Enum. Pl. Zeyl. 357 = barbata.
Wightii, *Munro, ex Benth. & Hook. f. Gen.* iii. 1113, nomen.—Ind. or.

CHIONANTHUS, Gaertn. Fruct. i. 189. t. 39 (1788) = **Linociera**, Sw. (Oleac.).
acuminata, Wall. ex DC. Prod. viii. 298 = L. acuminata.
acuminigera, F. Muell. Fragm. viii. 42.—Austral.
albidiflora, Thw. Enum. Pl. Zeyl. 189 = L. albidiflora.
axillaris, R. Br. Prod. 523 = L. albidiflora?
axillaris, Roxb. ex Wall. Cat. sub n. 2824 = L. macrophylla.
callophylla, Blume, Mus. Bot. Lugd. Bat. i. 319.—Malaya.
caribaea, Jacq. Coll. ii. 110 = L. compacta.
compacta, Sw. Prod. Veg. Ind. Occ. 13 = L. compacta.
cotinifolia, Willd. Sp. Pl. i. 47 = L. cotinifolia.

CHIONANTHUS:—
courtallensis, Bedd. For. Man. Bot. 155 = Linociera leprocarpa.
cuspidata, Blume, Mus. Bot. Lugd. Bat. i. 319.—Ins. Celebes.
densiflora, Zoll. & Morr. in Nat. en Geneesk. Arch. Neêrl. Indic, ii. (1845) 4 = L. densiflora.
dichotoma, Roxb. Hort. Beng. 3; Fl. Ind. i. 108 = L. purpurea.
diversifolia, Miq. Fl. Ind. Bat. Suppl. 559.—Sumatra.
domingensis, Lam. Tabl. Encyc. i. 30 = L. latifolia.
effusiflora, F. Muell. Fragm. iv. 83 = L. macrophylla.
elliptica, Blume, Mus. Bot. Lugd. Bat. i. 319.—Malaya.
Ghaeri, Gaertn. Fruct. i. 190 = L. cotinifolia.
glomerata, Blume, Mus. Bot. Lugd. Bat. i. 318.—Sumatra.
guianensis, Pers. Syn. i. 9.—Guiana.
incrassata, Sw. Prod. Veg. Ind. Occ. 13; Willd. Sp. Pl. i. 47 = L. tetrandra.
insignis, Miq. Fl. Ind. Bat. Suppl. 559 = L. insignis.
intermedia, Bedd. Fl. Sylv. t. 239 = L. intermedia.
laxiflora, Blume, Mus. Bot. Lugd. Bat. i. 319.—Borneo.
leprocarpa, Thw. Enum. Pl. Zeyl. 189 = L. leprocarpa.
ligustrina, Pers. Syn. i. 9 = L. ligustrina.
littorea, Miq. Fl. Ind. Bat. Suppl. 559.—Sumatra.
luzonica, Blume, Mus. Bot. Lugd. Bat. i. 319 = L. luzonica.
macrocarpa, Blume, l. c. = L. insignis.
macrophylla, Blume, l. c. 317; Kurz, in Journ. As. Soc. Beng. xlv. (1876) ii. 139 = L. macrophylla.
malabarica, Heyne, ex Wall. Cat. sub n. 2828 = L. malabarica.
Mayepea, Vahl, Symb. Bot. ii. 1 = L. tetrandra.
minutiflorus, Kurz, For. Fl. Brit. Burma, ii. 159 = L. minutiflora.
montana, Blume, Bijdr. 681 = L. montana.
montana, Kurz, For. Fl. Brit. Burma, ii. 159 = L. insignis.
monticola, Blume, Mus. Bot. Lugd. Bat. i. 317 = L. montana.
palembanica, Miq. Fl. Ind. Bat. Suppl. 558 = L. panciflora.
picrophloia, F. Muell. Fragm. iii. 139. t. 24 = L. intermedia.
pubescens, H. B. & K. Nov. Gen. et Sp. iii. 235.—N. Granat.
purpurea, Lam. Tabl. Encyc. i. 30 = L. purpurea.
quadristaminea, F. Muell. Fragm. viii. 41.—Austral.
ramiflora, Roxb. Hort. Beng. 3; Fl. Ind. i. 107 = L. macrophylla.
rostrata, Miq. Fl. Ind. Bat. ii. 553 = L. rostrata.
rostrata, Thw. Enum. Pl. Zeyl. 189 = L. albidiflora.
smilacifolia, Wall. in Roxb. Fl. Ind. ed. Carey, i. 108 = Myxopyrum smilacifolium.
spicata, Blume, Mus. Bot. Lugd. Bat. i. 318.—Malaya.
sumatrana, Blume, l. c.—Sumatra.
tenuiflora, Wall. ex DC. Prod. viii. 298 = L. macrophylla.
terniflora, Griff. Notul. iv. 740 = L. terniflora.
tetrandra, Vahl, Enum. i. 45 = L. tetrandra.
timorensis, Blume, Mus. Bot. Lugd. Bat. i. 318.—Timor.
zeylanica, Heyne, ex Wall. Cat. sub n. 2828 = L. malabarica.
zeylanica, Lam. Tabl. Encyc. i. 30 = L. cotinifolia.
zeylanica, Linn. Sp. Pl. 8 = L. purpurea.

CHIONANTHUS, Royen, ex Linn. Gen. ed. I. 335 (1737). *OLEACEAE*, Benth. & Hook. f. ii. 677.
chinensis, Maxim. in Bull. Acad. Pétersb. xx. (1875) 430 = retusa.
angustifolia, Rafin. New Fl. Am. iii. 88 = virginica.
fraxinifolia, Kellogg, in Proc. Calif. Acad. v. (1873) 18 = Fraxinus dipetala.
fragrans, Edwards, ex Steud. Nom. ed. II. i. 351 = virginica.
heterophyla, Rafin. New Fl. Am. iii. 87 = virginica.
latifolia, Ait. ex Steud. Nom. ed. II. i. 350 = virginica.
longifolia, Rafin. New Fl. Am. iii. 88 = virginica.
maritima, Sweet, Hort. Brit. ed. II. 351 = virginica.
montana, Rafin. New Fl. Am. iii. 87 = virginica.
obovata, Rafin. l. c. = virginica.
retusa, *Paxt. Flow. Gard.* iii. (1852–53) 88.—China.
trifida, Moench, Meth. 678 = virginica.
triflora, Stokes, Bot. Mat. Med. i. 19 = virginica.
vernalis, Salisb. Prod. 14 = virginica.
verna, Baill. Hist. Pl. i. 295 = virginica.
virginica, *Linn. Sp. Pl.* 8.—Am. bor.

CHIONE, DC. Prod. iv. 461 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 107.
CRUSEA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 204. t. 19. f. 1.
SACCONIA, Endl. Gen. Pl. 541 (1838).
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 22.—Ins. Cuba.
elliptica, *Griseb. Pl. Wright. Cub.* 507.—Ind. occ.
glabra, *DC. Prod.* iv. 461.—Ind. occ.
lucida, *Griseb. Pl. Wright. Cub.* 507.—Ind. occ.
myrtifolia, *Griseb. l. c.*—Ind. occ.
vaccinifolia, Griseb. Cat. Pl. Cub. 133 = Erithalis vacciniifolia.

CHIONE, Salisb. Gen. Pl. Fragm. 100 (1866) = **Narcissus**, Tourn. (Amaryllid.).
italica, Salisb. l. c. = N. italicus.
papyracea, Salisb. l. c. = N. papyraceus.

CHIONICE, Bunge, ex Ledeb. Fl. Ross. ii. 33 (1844) = **Potentilla**, Tourn. (Rosac.).
dioica, Bunge, l. c. = P. tetrandra.

CHIONODOXA, Boiss. Diagn. Ser. I. v. 61 (1844). *LILIACEAE*, Benth. & Hook. f. iii. 813.
cretica, *Boiss. & Heldr. l. c.* xiii. 24.—Ins. Creta.
cretica, Jaub. & Spach, Illustr. v. 46 t. 443 = nana.
Forbesii, Baker, in Journ. Linn. Soc. xi. (1871) 436 = Luciliae.
Luciliae, *Boiss. Diagn.* Ser. i. v. 61.—As. Min.
nana, *Boiss. & Heldr, l. c.* xiii. 24.—Ins. Creta.

CHIONOGRAPHIS, Maxim. in Bull. Acad. Pétersb. xi. (1867) 435. *LILIACEAE*, Benth. & Hook. f. iii. 826.
japonica, *Maxim. l. c.* 436.—Japon.

CHIONOLAENA, DC. Prod. v. 397 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 302.
arbuscula, *DC. l. c.*—Bras.
corymbosa, *Hemsl. Diagn. Pl. Nov.* 32.—Mexic.
Glaziovii, *Baker, in Mart. Fl. Bras.* vi. III. 130.—Bras.
glomerata, *Baker, l. c.*—Bras.
Isabellae, *Baker, l. c.*—Bras.
latifolia, Baker, l. c. 132 = Leucopholis latifolia.
lavandulaceum, Hemsl. Biol. Centr. Am. Bot. ii. 134 = seq.
lavandulifolium, *Benth. & Hook. f. Gen.* ii. 302.—Mexic.
longifolia, *Baker, in Mart. Fl. Bras.* vi. III. 131.—Bras.
lychnophorioides, *Sch. Bip. in Pollichia*, xx-xxi. (1863) 391.—Bras.
phylicoides, Baker, in Mart. Fl. Bras. vi. III. 131 = Leucopholis phylicoides.
Wittigiana, *Baker, l. c.* 129.—Bras.

CHIONOPAPPUS, Benth. in Benth. & Hook. f. Gen. ii. 485 (1873). *COMPOSITAE*, Benth. & Hook. f. l. c.

CHIONOPHILA, Benth. in DC. Prod. x. 331 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 942.
Jamesii, *Benth. l. c.*—Am. bor. occ.

CHIONOPHILA, Miers, ex Lindl. Veg. Kingd. 701 (1847) = **Boopis**, Juss. (Calycer.).

CHIONOPTERA, DC. Prod. vii. 14 (1838). *COMPOSITAE*, Benth. & Hook. f. ii. 495.
Gayophyta, *DC. l. c.*—Chili.

CHIONOTHRIX, Hook. f. in Benth. & Hook. f. Gen. iii. 33 (1880). *AMARANTACEAE*, Benth. & Hook. f. l. c.
somalensis, *Hook. f. l. c.*—Afr. trop.

CHIONOTRIA, Jack. in Malay. Misc. ii. (1822) VII. 53 = **Glycosmis**, Correa (Rutac.).
monogyna, Walp. Rep. i. 382 = seq.
rigida, Jack, in Malay. Misc. ii. (1822) VII. 54 = G. pentaphylla.

CHIOPHILA, Rafin. Fl. Tellur. iii. 23 (1836) = **Gentiana**, Linn.
nivalis, Rafin. l. c. = G. nivalis.

CHIRANTHODENDRON, Cerv. ex Cav. in Anal. Cienc. Nat. vi. (1803) 303. adnot. = **Cheirostemon**, Humb. & Bonpl. (Sterculiac.).
californicum, Baill. Hist. Pl. iv. 70 = Fremontia californica.
pentadactylon, Larréat, Desc. Chiranth. ed. Lesc. 17 = Cheirostemon platanoides.

CHIRATA, G. Don, Gen. Syst. iv. 659 (1837) = **Chirita**, Buch.-Ham.

CHIRATIA, Montr. in Mém. Acad. Lyon, x. (1860) 202 = **Sonneratia**, Linn. f. (Lythrar.).
leucantha, Montr. l. c. 203.—N. Caled.

CHIRITA, Buch.-Ham. in D. Don, Prod. Fl. Nep. 89 (1825). *GESNERACEAE*, Benth. & Hook. f. ii. 1022.
BABACTES, DC. Prod. ix. 260 (1845).
CALOSACME, Wall. Cat. nn. 800 ad 806 (1829).
CHIRATA, G. Don, Gen. Syst. iv. 659 (1837).
GONATOSTEMON, Regel, Gartenfl. xv. (1866) 353.
HYPOPTERON, Hassk. Cat. Hort. Bog. Alt. 154 (1844).
LIEBIGIA, Endl. Gen. 1407 (1841).
MORSTDORFFIA, Steud. Nom. ed. II. ii. 161 (1841).
TROMSDORFFIA, Blume, Bijdr. 762 (1826).
acuminata, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Reg. Himal.
acuminata, A. DC. Prod. ix. 269 = Didymocarpus acuminata.
amplectens, Steud. Nom. ed. II. i. 351 = bifolia.
anachoretica, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 231.—China.
bifolia, *D. Don, Prod. Fl. Nep.* 90.—Reg. Himal.
Bilabium, *C. B. Clarke, in A. DC. Monog. Phan.* v. 127.—Sumatra.
Blumei, *C. B. Clarke, l. c.* 122.—Java.
bracteosa, Miq. Fl. Ind. Bat. ii. 728 = Horsfieldii.
brevipes, *C. B. Clarke, in A. DC. Monog. Phan.* v. 120.—Ind. or.
caerulea, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Java.
caliginosa, *C. B. Clarke, in A. DC. Monog. Phan.* v. 122.—Malacca.
calva, *C. B. Clarke, l. c.* 116.—Reg. Himal.
Clarkei, *Hook. f. Fl. Brit. Ind.* iv. 359.—Reg. Himal.
communis, Gardn. in Calc. Journ. Nat. Hist. iv. (1846) 481 = zeylanica.
cortusifolia, *Hance, in Journ. Bot.* xxi. (1883) 324.—China.
diaphana, Royle, Illustr. Bot. Himal. 294 = pumila.
dimidiata, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Reg. Himal.
eburnea, *Hance, in Journ. Bot.* xxi. (1883) 168.—China.
Edgeworthii, A. DC. in Prod. ix. 269 = pumila.
Fauriei, *Franch. in Bull. Soc. Linn. Par.* i. (1885) 450.—China.
flava, R. Br. in Benn. Pl. Jav. Rar. 117 = pumila.
fusca, *C. B. Clarke, in A. DC. Monog. Phan.* v. 129.—Borneo.
glabra, C. B. Clarke, Comm. et Cyrt. Beng. t. 73 = calva.
glabra, *Miq. Fl. Ind. Bat.* ii. 729.—Malaya.
grandiflora, Wall. Pl. As. Rar. i. 43 = urticaefolia.
Griffithii, Hook. f. & Thoms. ex C. B. Clarke, in A. DC. Monog. Phan. v. 60 = Loxostigma Griffithii.
hamosa, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Ind. or.
Hookeri, *C. B. Clarke, Comm. & Cyrt. Beng.* t. 70.—Reg. Himal.
Horsfieldii, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Java.
Horsfieldii, De Vriese, Pl. Ind. Bat. Reinw. 11 = Blumei.
humilis, *Miq. Fl. Ind. Bat.* ii. 729.—Sumatra.
insignis, *C. B. Clarke, in A. DC. Monog. Phan.* v. 121.—Assam.
Juliae, *Hance, in Journ. Bot.* xxi. (1883) 168.—China.
Kurzii, *C. B. Clarke, in Journ. Linn. Soc.* xv. (1876) 145.—Reg. Himal.
lachenensis, *C. B. Clarke, in A. DC. Monog. Phan.* v. 118.—Reg. Himal.
lilacina, *Lem. Illustr. Hortic.* (1869) t. 608.—Ind. or.

CHIRITA:—
macrophylla, *Wall. Pl. As. Rar.* i. 56. t. 72.—Reg. Himal.
macrosiphon, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 231.—China.
mollis, *Miq. Fl. Ind. Bat.* ii. 728.—Sumatra.
monophylla, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 360.—Reg. Himal.
Moonii, *Gardn. in Calc. Journ. Nat. Hist.* vi. (1846) 479.—Zeylan.
obtusa, *C. B. Clarke, in A. DC. Monog. Phan.* v. 114.—China.
polycarpa, *Steud. Nom.* ed. II. i. 351.—Ind. or.
polyneura, Miq. Fl. Ind. Bat. ii. 728 = Horsfieldii.
primulacea, *C. B. Clarke, Comm. et Cyrt. Beng.* t. 82.—Reg. Himal.
pumila, *D. Don, Prod. Fl. Nep.* 90.—Reg. Himal.
scaberrima, *R. Br. in Benn. Pl. Jav. Rar.* 117.—Java.
scabra, Royle, ex C. B. Clarke, in A. DC. Monog. Phan. v. 113 = bifolia.
sinensis, *Lindl. Bot. Reg.* (1844) t. 59.—Hongkong.
speciosa, *Kurz, in Journ. Bot.* xi. (1873) 195.—China.
spectabilis, *Miq. Fl. Ind. Bat. Suppl.* 564.—Sumatra.
ternifolia, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 124 = Lysionotus serrata.
urticaefolia, *Buch.-Ham. ex D. Don, l. c.* 90.—Reg. Himal.
vulgaris, Hort. ex Belg. Hortic. iii. (1853) t. [36].—Cf. p. 238 = zeylanica.
Walkeri, *Gardn. in Calc. Journ. Nat. Hist.* vi. (1846) 480.—Zeylan.
Walkeriae, Hook. Bot. Mag. t. 4327 = Walkeri.
zeylanica, *Hook. l. c.* t. 4182.—Zeylan.
Zollingeri, *C. B. Clarke, in A. DC. Monog. Phan.* v. 125.—Java.

CHIROCALYX, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 97 = **Erythrina**, Linn. (Legumin.).
abyssinicus, Hochst. in Flora, xxix. (1846) 600 = E. abyssinica.
Candolleanus, Walp. in Flora, xxxvi. (1853) 148 = E. spathacea.
divaricatus, Walp. l. c. = E. divaricata.
indicus, Walp. l. c. = E. indica.
latifolius, Walp. l. c. = E. senegalensis.
mollissimus, Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 98 = E. tomentosa.
pictus, Walp. in Flora, xxxvi. (1853) 148 = E. indica.
pubescens, Walp. in Linnaea, xxiii. (1850) 741 = E. fissa.
rubrinervius, Walp. in Flora, xxxvi. (1853) 148 = E. rubrinervis.
tomentosus, Hochst. in Flora, xxix. (1846) 600 = E. tomentosa.
umbrosus, Walp. in Flora, xxxvi. (1853) 148 = E. umbrosa.
velutinus, Walp. l. c. = E. velutina.

CHIRONIA, Linn. Coroll. Gen. 5 (1737). *GENTIANEAE*, Benth. & Hook. f. ii. 805.
EUPODIA, Rafin. Fl. Tellur. iii. 29 (1836).
EVALTHE, Rafin. l. c. 77 (1836).
ONEFERA, Rafin. l. c. 30 (1836).
PLOCANDRA, E. Mey. Comm. Pl. Afr. Austr. 181 (1837).
ROESLINIA, Moench, Meth. Suppl. 211 (1802).
ROSLINIA, G. Don, Gen. Syst. iv. 203 (1837).
amoena, Rafin. in Med. Repos. N. York, v. (1808) 359 = Sabbatia gracilis ?
amoena, Salisb. Prod. 137 = linoides.
angularis, Linn. Sp. Pl. 190 = Sabbatia angularis.
angustifolia, Sims, Bot. Mag. t. 818 = frutescens.
arenaria, *E. Mey. Comm. Pl. Afr. Austr.* ii. 180.—Afr. austr.
baccata, Hoffmgg. Verz. Pfl. Nachtr. i. 211 = seq. ?
baccifera, *Linn. Sp. Pl.* 190.—Afr. anstr.
baccifera, Zeyh. ex Griseb. in DC. Prod. ix. 41 = linoides.
Barclayana, Hort. Berol. ex Griseb. Gen. et Sp. Gent. 100 = peduncularis.
brachiata, Willd. ex Griseb. l. c. 145 = Erythraea Roxburghii.
calycosa, Michx. Fl. Bor. Am. i. 147 = Sabbatia calycosa.
campanulata, Linn. Sp. Pl. 190 = Sabbatia gracilis ?

CHIRONIA :—
capsularis, Blanco, Fl. Filip. ed. I. 102 = Ipomoea peltata.
caryophylloides, Linn. Cent. ii. 12; Amoen. Acad. iv. 308 = frutescens.
Centaurium, Schmidt, Fl. Bohem. ii. 31 = Erythraea Centaurium.
centauroides, Roxb. Hort. Beng. 16; Fl. Ind. i. 584 = Erythraea Roxburghii.
chilensis, Willd. Sp. Pl. i. 1067 = Erythraea chilensis.
chloroides, Michx. Fl. Bor. Am. i. 147 = Sabbatia chloroides.
cymosa, *Burm. f. Fl. Cap. Prod.* 5.—Afr. austr.
cymosa, Lam. Encyc. i. 479 = Sabbatia lanceolata.
decandra, Walt. Fl. Carol. 95 = Sabbatia chloroides.
decussata, Vent. Jard. Cels, 31 = frutescens.
dichotoma, Walt. Fl. Carol. 95 = Sabbatia calycosa.
dodecandra, Linn. Sp. Pl. 190 = Sabbatia chloroides.
dubia, Willd. ex Griseb. Gen. et Sp. Gent. 146 = Erythraea Muehlenbergii.
emarginata, Jarosz, Pl. Nov. Cap. 11 = lychnoides.
Erythraea, Schousb. Vextr. Marok. 95 = Erythraea ramosissima.
exigua, *Oliver, in Hook. Ic. Pl.* t. 1229.—Afr. austr.
filiformis, Desf. Tabl. ed. I. 77, nomen = Mierocala filiformis.
Fischeri, Hort. ex Paxt. Mag. Bot. xi. (1844) 237 = floribunda.
floribunda, *Paxt. l. c.* 223.—Afr. austr.
frutescens, *Linn. Sp. Pl.* 190; *Berg. Desc. Pl. Cap.* 45. —Afr. austr.
Gerardi, Schmidt, Fl. Bohem. ii. 33 = Erythraea pulchella.
glutinosa, *Paxt. Mag. Bot.* xv. (1849) 245.—Austral.?
gracilis, Michx. Fl. Bor. Am. i. 146 = Sabbatia gracilis.
grandiflora, Salisb. Prod. 137 = frutescens.
helodea, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 125 = Erythraea elodes.
inaperta, Willd. Sp. Pl. i. 1069 = Erythraea ramosisisima.
intermedia, Mérat, Fl. Paris, 91 = Erythraea ramosissima.
jasminoides, [Ker-Gawl. in] Bot. Reg. t. 197 = tetragona.
jasminoides, Lam. Illustr. i. 479. t. 108. f. 2 = nudicaulis.
jasminoides, *Linn. Pl. Rar. Afr.* 9 (*Amoen. Acad.* vi. 84).—Afr. austr.
Krebsii, *Griseb. Gen. et Sp. Gent.* 98.—Afr. austr.
lanceolata, Walt. Fl. Carol. 95 = Sabbatia lanceolata.
lanosanthera, Blanco, Fl. Filip. ed. II. 71 = Ipomoea peltata.
latifolia, *E. Mey. Comm. Pl. Afr. Austr.* ii. 178.—Afr. austr.
linarifolia, Loisel. Notes, 155 = Erythraea linarifolia.
linearifolia, Steud. Nom. ed. II. i. 352, in syn. sphalm. = praec.
linifolia, Steud. l. c. 351 = Erythraea linarifolia.
linoides, *Linn. Sp. Pl.* 189.—Afr. austr.
littoralis, Turn. in Turn. & Dillw. Bot. Guide, ii. 469 = Erythraea linarifolia.
lutea, Bertol. Rar. It. Pl. dec. ii. 32 = Erythraea maritima.
lychnoides, *Berg. Desc. Pl. Cap.* 45.—Afr. austr.
madagascariensis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 273.—Madag.
maritima, *Eckl. in S. Afr. Quart. Journ.* ii. (1830); *ex Griseb. Gen. et Sp. Gent.* 100.—Afr. austr.
maritima, Georgi, Beschr. Russ. Reich. iii. iv. 791 = Erythraea ramosissima.
maritima, Willd. Sp. Pl. i. 1069 = Erythraea maritima.
melampyrifolia, *Lam. Illustr.* ii. 479.—Afr. austr.
minima, Thuill. Fl. Par. ii. 116 = Cicendia pusilla.
minor, Desf. Cat. Hort. Par. ed. III. 126 = Erythraea ramosissima.
nana, Bast. ex Griseb. in DC. Prod. ix. 57 = Erythraea ramosissima.
nudicaulis, *Linn. f. Suppl.* 151.—Afr. austr.
nummularifolia, Willd. ex Griseb. Gen. et Sp. Gent. 144 = Erythraea diffusa.
occidentalis, DC. Fl. Fr. Suppl. 2782 = Erythraea maritima.
orthostylis, Reichb. Exot. t. 245 = frutescens.
ovata, Spreng. ex Griseb. in DC. Prod. ix. 41 = serpyllifolia.
palustris, *Burch. Trav.* ii. (1821) 226.—Afr. austr.
paniculata, Michx. Fl. Bor. Am. i. 146 = Sabbatia paniculata.
parviflora, Salisb. Prod. 136 = baccifera.

CHIRONIA :—
parvifolia, *E. Mey. Comm. Pl. Afr. Austr.* 180.—Afr. austr.
pedunculatis, *Lindl. Bot. Reg.* t. 1803.—Afr. austr.
perfoliata, *Eckl. in S. Afr. Quart. Journ.* ii. (1830), *ex Griseb. Gen. et Sp. Gent.* 104.—Afr. austr.
perfoliata, Salisb. Prod. 137 = Chlora perfoliata.
pubescens, *Baker, in Journ. Bot.* xx. (1882) 172.—Madag.
pulchella, Willd. Sp. Pl. i. 1067 = Erythraea ramosissima.
pusilla, Steud. Nom. ed. II. i. 351 = Microcala filiformis.
ramosissima, Ehrh. ex Griseb. Gen. et Sp. Gent. 137 = Erythraea ramosissima.
scabrida, *Griseb. l. c.* 103.—Afr. austr.
scabriuscula, Willd. ex Griseb. l. c. 143 = Erythraea linarifolia.
serpyllifolia, *Lehm. Ind. Sem. Hort. Hamb.* (1828).—Afr. austr.
speciosa, *E. Mey. Comm. Pl. Afr. Austr.* 178.—Afr. austr.
spicata, Willd. Sp. Pl. i. 1069 = Erythraea spicata.
stellata, Muhl. Cat. ex A. Gray, Syn. Fl. N. Am. ii. 1. 115 = Sabbatia stellaris.
suffruticosa, Salzm. ex Griseb. Gen. et Sp. Gent. 141 = Erythraea Centaurium.
tabularis, *Page, ex Steud. Nom.* ed. II. i. 352.—Afr. austr.
tenuiflora, *Link, ex Steud. l. c.*—Hab.?
tetragona, *Linn. f. Suppl.* 151.—Afr. austr.
trinervia, Linn. Sp. Pl. 189 = Exacum zeylanicum.
trinervis, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1306 = = pedunculatis.
uliginosa, Lapeyr. Hist. Abr. Pl. Pyr. Suppl. 39 = Erythraea elodes.
uliginosa, Waldst. & Kit. Pl. Rar. Hung. iii. 287. t. 259 = Erythraea linarifolia.
uniflora, Eckl. ex Griseb. in DC. Prod. ix. 41 = linoides.
uniflora, *Lam. Encyc.* i. 737.—Afr. austr.
Vaillantii, F. W. Schmidt, Fl. Bohem. ii. 34 = Erythraea ramosissima.
virescens, Willd. ex Steud. Nom. ed. II. i. 352 = Erythraea maritima.
viscosa, Zeyher, ex Griseb. Gen. et Sp. Gent. 105 = tetragona.
vulgaris, Cham. in Linnaea, vi. (1831) 343 = linoides, lychnoides, melampyrifolia.

CHIROPETALUM, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 21 = **Argithamnia**, Sw. (Euphorbiac.).
Berterianum, Schlecht. in Linnaea, xxvi. (1853–55) 637 = A. Berteroana.
canescens, Phil. Fl. Atac. 49 = A. canescens.
griseum, Griseb. in Goett. Abh. xxiv. (1879) 57.—Reg. Argent.
lanceolatum, C. Gay, Fl. Chil. v. 344 = A. tricuspidata.
lanceolatum, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 21 = A. Berteroana.
lineatum, Klotzsch, in Wiegm. Archiv, vii. (1841) 199 = A. lineata.
molle, Klotzsch, l. c. = A. mollis.
ovatum, Phil. in Linnaea, xxix. (1857–58) 42 = A. trieuspidata.
peruvianum, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 22 = A. quinquecuspidata.
triandrum, Griseb. in Goett. Abh. xxiv. (1879) 56.—Reg. Argent.
tricuspidatum, A. Juss. in Ann. Sc. Nat. Sér. I. xxv. (1832) 22 = A. tricuspidata.

CHIROSTEMUM, Cerv. in Anal. Cienc. Nat. vi. (1803) 303 = **Cheirostemon**, Humb. & Bonpl. (Sterculiac.).

CHISOCHETON, Blume, Bijdr. 168 (1825). *MELIACEAE*, Benth. & Hook. f. i. 333.
SCHIZOCHITON, Spreng. Syst. iv. Cur. Post. 251 (1827).
amabilis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 27.—As. or.
barbatus, *C. DC. in A. DC. Monog. Phan.* i. 536.—As. or.
canalensis, Baill. Adansonia, xi. (1873–76) 260 = Dysoxylum camaleuse.
ceramicus, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 29. —Ins. Ceram.

CHISOCHETON :—
costatus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 552.—Ind. or.
divergens, *Blume, Bijdr.* 169.—Java.
diversifolius, *Miq. Fl. Ind. Bat. Suppl.* i. 504.—Sumatra.
dysoxylifolius, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) ii. 49.—Burma.
erythrocarpus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 550. —Malacca.
fragrans, *Hiern, l. c.* 551.—Malacca.
glomeratus, *Hiern, l. c.*—Ins. Penang.
grandiflorus, *Kurz, in Journ. As. Soc.* xli. (1872) ii. 296.—Burma.
holocalyx, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 551.—Malaya.
Junghuhni, *Miq. in Ann. Mus. Bot Lugd. Bat.* iv. 30. —As. or.
paniculatus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 552.—Ind. or.; Burma.
patens, *Blume, Bijdr.* 169.—Java.
paucijugum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 30.—Sumatra.
penduliflorus, *Planch. ex Hiern, in Hook. f. Fl. Brit. Ind.* i. 550.—Malaya.
spectabilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 29. —As. or.
spicatus, *Hiern, in Hook. f. Fl. Brit. Ind.* i. 550.—Malacca.
tetrapetalus, *Turcz. in Bull. Soc. Nat. Mosc.* (1858) i. 411.—Java.
Vrieseanus, *C. DC. in A. DC. Monog. Phan.* i. 532.—As. or.

CHITHONANTHUS, Lehm. in Ind. Sem. Hort. Hamburg. (1842) 4; et Pl. Preiss. ii. 368 (1847) = **Acacia**, Willd. (Legumin.).
restiaceus, Lehm. ll. cc. = A. restiacea.

CHITONIA, D. Don, in Mem. Wern. Soc. iv. (1823) 317 = **Miconia**, Ruiz & Pav. (Melastomac.).
albicans, D. Don, ex Naud. in Ann. Sc. Nat. Sér. III. xv. (1851) 339 = Tetrazygia albicans.
aurea, D. Don, in Mem. Wern. Soc. iv. (1823) 319 = M. aurea.
bubalina, D. Don, l. c. = M. bubalina.
caudata, D. Don, l. c. = M. Donaeana.
Fothergilla, D. Don, l. c. 318 = M. Fothergilla.
macrophylla, D. Don, l. c. 319 = M. macrophylla.
pyramidalis, G. Don, in Loud. Hort. Brit. 174 = M. laevigata.
rubiginosa, Steud. Nom. ed. II. i. 352 = M. Fothergilla.
Tamonia, G. Don, in Loud. Hort. Brit. 174 = M. Fothergilla.

CHITONIA, Moç. & Sesse, ex DC. Prod. i. 707 (1824). *ZYGOPHYLLEAE*, Benth. & Hook. f. i. 268.
mexicana, *Moç. & Sesse, l. c.*—Mexic.

CHITONIA, Salisb. Gen. Pl. Fragm. 51 (1866) = **Zygadenus**, Michx. (Liliac.).

CHIZOCHETON, A. Juss. in Orb. Dict. Sc. viii. 80 (1849) = **Chisocheton**, Blume (Meliac.).

CHLAENANDRA, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 83 (1868–69). *MENISPERMACEAE*.
ovata, *Miq. l. c.* 84.—N. Guinea?

CHLAENOBOLUS, Cass. in Dict. Sc. Nat. xlix. 337 (1827) = **Pterocaulon**, Ell. (Compos.).
alopecuroides, Cass. l. c. 339 = P. virgatum.
pycnostachyus, Cass. l. c. 338 = P. pycnostachyum.
rugosus, Cass. l. c. 341 = P. virgatum.
spicatus, Cass. l. c. = P. virgatum.
virgatus, Cass. l. c. 340 = P. virgatum.

CHLAMYDANTHUS, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 355, 358 = **Thymelaea**, Endl.
argenteus, C. A. Mey. l. c. 358 = T. Tartonraira.
canescens, C. A. Mey. l. c. = T. canescens.
ellipticus, C. A. Mey. l. c. 359 = T. elliptica.
hirsutus, Griseb. Spicil. Fl. Rumel. ii. 54 = Passerina hirsuta.

CHLAMYDANTHUS:—
nitidus, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 358 = Thymelaea nitida.
Tartonraira, C. A. Mey. l. c. = T. Tartonraira.
thesioides, C. A. Mey. l. c. 359 = T. thesioides.
Thymelaea, C. A. Mey. l. c. = T. Sanamunda.
tinctorius, C. A. Mey. l. c. 358 = T. tinctoria.
villosus, C. A. Mey. l. c. = T. villosa.
virgatus, C. A. Mey. l. c. 359 = T. virgata.

CHLAMYDIA, Banks, ex Gaertn. Fruct. i. 71. t. 18 (1788) = **Phormium**, Forst. (Liliac.).
tenacissima, Gaertn. l. c. = P. tenax.

CHLAMYDOCARYA, Baill. Adansonia, x. (1872) 276. *OLACINEAE*.
capitata, *Baill. l. c.* 277.—Afr. trop.
Thomsoniana, *Baill. l. c.* 276.—Afr. trop.

CHLAMYDOPHORA, Ehrenb. ex Less. Syn. Comp. 265, 448 (1832) = **Cotula**, Linn. (Compos.).
pubescens, Coss. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 279 = Cot. pubescens.
tridentata, Ehrenb. ex Less. Syn. Comp. 266 = Cot. tridentata.

CHLAMYDOSPERMA, A. Rich. Ess. Fl. Cub. 631 (1845) = **Stegnosperma**, Benth. (Phytolacc.).

CHLAMYDOSTYLUS, Baker, in Journ. Bot. xiv. (1876) 185 = **Nemastylis**, Nutt. (Irid.).
bracteolatus, Baker, in Journ. Linn. Soc. xvi. (1877) 107.—Boliv.
cernuus, Baker, in Journ. Bot. xiv. (1876) 186 = N. triflora.
Medusa, Baker, l. c.—Bras.
multiflorus, Baker, l. c. = N. multiflora.
spathaceus, Griseb. in Goett. Abh. xxiv. (1879) 324 = N. spathacea.
tenuis, Baker, in Journ. Bot. xiv. (1876) 185 = N. tenuis.
triflorus, Baker, in Journ. Linn. Soc. xvi. (1877) 107 = N. triflora.

CHLAMYSPERMA, Less. Syn. Comp. 256 (1832) = **Villanova**, Lag. (Compos.).
arenaroides, Hook. & Arn. Bot. Beech. Voy. 300. t. 64 = V. pratensis.
polygama, Triana, in Ann. Sc. Nat. Sér. IV. ix. (1858) 40.—N. Granat.
pratense, Less. Syn. Comp. 256 = V. pratensis.

CHLAMYSPORUM, Salisb. Parad. Lond. t. 103 (1808) = **Thysanotus**, R. Br. (1810) (Liliac.).
dichotomum, Salisb. l. c. in nota sub t. 103 = T. dichotomus.
hispidulum, Salisb. ex Steud. Nom. ed. II. i. 352 = T. hispidulus.
juncifolium, Salisb. Parad. Lond. t. 103 = T. junceus.
multiflorum, Salisb. ex Steud. Nom. ed. II. i. 352 = T. multiflorus.
pauciflorum, Salisb. ex Steud. l. c. = T. pauciflorus.
triandrum, Salisb. ex Steud. l. c. = T. triandrus.

CHLANIS, Klotzsch, in Peters, Reise Mossamb. Bot. 144 (1861) = **Oncoba**, Forsk. (Bixin.).
macrophylla, Klotzsch, l. c. 145 = O. macrophylla.
tettensis, Klotzsch, l. c. = O. tettensis.

CHLAOTRACHELUS, Hook. f. Fl. Brit. Ind. iii. 234 (1881), sphalm. = CLAOTRACHELUS, Zoll. = **Vernonia**, Schreb. (Compos.).

CHLETERUS, Rafin. Princ. Somiol. 25 (1814) = **Boea**, Comm. (Gesnerac.).

CHLIDANTHUS, Herb. App. 46 (1821). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 723.
COLEOPHYLLUM, Klotzsch, in Otto & Dietr. Allg. Gartenz. viii. (1840) 185.
Cumingii, Presl, Bot. Bemerk. 115 = Hippeastrum advenum.
Ehrenbergii, *Kunth, Enum. Pl.* v. 654.—Mexic.
fragrans, *Herb. App.* 46.—Peruv.
luteus, D. Dietr. Syn. Pl. ii. 1175 = Stenomesson luteum.

CHLOAMNIA, Rafin. Neogenyt. 4 (1825) = **Festuca**, Tourn. (Gramin.).

CHLOANTHES, R. Br. Prod. 513 (1810). *VERBENACEAE*, Benth. & Hook. f. ii. 1140.
HEMISTEMON, F. Muell. Fragm. x. 73 (1876).
atriplicina, F. Muell. Fragm. i. 235 = Pityrodia atriplicina.
Bartlingii, Lehm. Ind. Sem. Hort. Hamb. (1844), ex Bartl. in Lehm. Pl. Preiss. i. 352 = Pityrodia Bartlingii.
Bonneyana, *F. Muell. Fragm.* x. 73.—Austral.
bullata, F. Muell. l. c. vi. 156 = Pityrodia uncinata.
coccinea, *Bartl. in Lehm. Pl. Preiss.* i. 352.—Austral.
cuneata, F. Muell. Census, 103 = Pityrodia cuneata.
dilatata, F. Muell. Fragm. vi. 157 = Pityrodia dilatata.
Elderi, F. Muell. l. c. x. 13 = Hemiphora Elderi.
glandulosa, *R. Br. Prod.* 514.—Austral.
halganiacea, *F. Muell. Fragm.* x. 14.—Austral.
hemigenioides, F. Muell. l. c. vi. 156 = Pityrodia hemigenioides.
lavandulaefolia, Sieber, ex Spreng. Syst. ii. 756 = Stoechadis.
lepidota, *F. Muell. in Wing, South. Sc. Record*, iii. (1883) 3.—Austral.
Lewellini, *F. Muell. Fragm.* viii. 50.—Austral.
loricata, *F. Muell. l. c.* x. 14.—Austral.
loxocarpa, F. Muell. l. c. ii. 22 = Pityrodia Drummondii.
Oldfieldii, F. Muell, l. c. i. 234 = Pityrodia Oldfieldii.
paniculata, F. Muell. l. c. iv. 80 = Pityrodia paniculata.
parviflora, *Walp. Rep.* iv. 58.—Austral.
rosmarinifolia, *A. Cunn. ex Steud. Nom.* ed. II. i. 352.—Austral.
salvifolia, F. Muell. Census, 103 = Pityrodia salvifolia.
stachyodes, F. Muell. Fragm. v. 50; vi. 158 = Pityrodia racemosa.
Stoechadis, *R. Br. Prod.* 514.—Austral.
uncinata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 194 = Pityrodia uncinata.
verbascina, F. Muell. Fragm. i. 233 = Pityrodia verbascina.

CHLOERUM, Willd. ex Spreng. Jahrb. i. III. 74 (1833) = **Abolboda**, Humb. & Bonpl. (Xyrid.).
gramineum, Willd. ex Kunth, Enum. Pl. iv. 27 = A. Poeppigii.
imberbe, Willd. ex Steud. Nom. ed. II. i. 352 = A. imberbis.

CHLOIDIA, Lindl. Gen. et Sp. Orch. 484 (1840) = **Corymbis**, Thou. (Orchid.).
decumbens, Lindl. l. c.—Bras.
flava, Reichb. f. in Walp. Ann. vi. 644 = Cor. flava.
polystachya, Reichb. f. l. c. = *vernalis*.
vernalis, Lindl. Gen. et Sp. Orch. 484.—Jamaica.

CHLONANTHES, Rafin. New Fl. Am. ii. 20 (1836) = **Chelone**, Linn. (Scrophul.).
tomentosa, Rafin. l. c.—Am. bor.

CHLONANTHUS, Rafin. Princ. Somiol. 26 (1814) = praec.

CHLOOPSIS, Blume, Enum. Pl. Jav. 14 (1827) = **Ophiopogon**, Ker-Gawl. (Haemodorac.).
acaulis, Blume, l. c. = O. japonicus.
caulescens, Blume, l. c. = O. japonicus.

CHLOOTHAMNUS, Buese, in Miq. Pl. Jungh. 386 (1851–55) = **Schizostachyum**, Nees (Gramin.).
chilianthus, Buese, l. c. 387.—Sumatra.

CHLORA, Ren. ex Adans. Fam. ii. 503 (1763). *GENTIANEAE*, Benth. & Hook. f. ii. 809.
BLACKSTONIA, Huds. Fl. Angl. ed. I. 146 (1762), nomen prius.
CHLOROSTIS, Rafin. Princ. Somiol. 26 (1814).
DISINSTYLIS, Rafin. Fl. Tellur. iii. 31 (1836).
PLEURIMARIA, Rafin. l. c. (1836).
acuminata, Koch & Ziz, Cat. Fl. Palat. 20 = serotina.
affinis, Willk. in Bot. Zeit. v. (1847) 874 = imperfoliata.
citrina, *Boiss. & Reut. Pugill. Pl. Nov.* 77.—Hispan.
crenulata, Salzm. ex Griseb. Gen. et Sp. Gent. 118 = imperfoliata.
dodecandra, Linn. Syst. ed. XII. 267 = Sabbatia chloroides.
dubia, Lam. Tabl. Encyc. ii. 434 = imperfoliata.
exaltata, Griseb. Gen. et Sp. Gent. 118 = Eustoma silenifolium.
grandiflora, Viv. Cors. App. Alt. 4 = perfoliata.

CHLORA:—
hybrida, *Tratt. Arch.* t. 229.—Europ.
imperfoliata, *Linn. f. Suppl.* 218.—Europ.
intermedia, Tenore, Syll. 565 = perfoliata.
lanceolata, Koch & Ziz, Cat. Pl. Palat. 20 = imperfoliata.
longifolia, *Lam. Tabl. Encyc.* ii. 434.—Hab.?
mascariensis, Desf. ex Steud. Nom. ed. I. 190; ed. II. i. 352 = perfoliata.
perfoliata, Gorter, ex Steud. Nom. ed. II. i. 352 = Saponaria Vaccaria.
perfoliata, *Linn. Syst.* ed. XII. 267.—Europ.; As. Min.
quadrifolia, *Linn. l. c.*—Europ. austr.
quadrifoliata, F. W. Schmidt, in Roem. Arch. i. I. 13 = quadrifolia.
serotina, *Koch, in Reichb. Pl. Crit.* iii. 6. t. 351.—Europ.; As. Min.
sessilifolia, Desv. in Trav. Soc. Amat. i. (1807) 74. t. 3. f. 2 = imperfoliata.
sessilis, Willd. Sp. Pl. ii. 341 (= Plantaginea?).—Reg. Argent.
sicula, Tineo, in Guss. Fl. Sic. Syn. ii. 819 = perfoliata.
triphylla, *Badaro, in Brugnat. Giorn. Fis.* II. ix. (1826) 166.—Italia.

CHLORADENIA, Baill. Etud. Gén. Euph. 471. t. 19 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 312.
ADENOGYNUM, Zoll. in Linnaea, xxviii. (1856) 325.
discolor, *Baill. l. c.* 472.—Java.

CHLORAEA, Lindl. in Brand. Quart. Journ. Roy. Instit. N. S. i. (1827) 47. *ORCHIDEAE*, Benth. & Hook. f. iii. 618.
ASARCA, Lindl. in Quart. Journ. Roy. Instit. i. (1827) 52.
BIENERIA, Reichb. f. in Bot. Zeit. xi. (1853) 3. t. 1.
GAVILLEA, Poepp. & Endl. ex Steud. Nom. ed. II. i. 666 (1840).
ULANTHA, Hook. Bot. Mag. lvii. sub t. 2990 (1830).
affinis, *Lindl. in Hook. Journ. Bot.* i. (1834) 4.—Chili.
alaris, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1827) 50.—Chili.
alpina, *Poepp. Fragm. Syn. Pl. Chil.* 17.—Chili.
aurantiaca, *Lindl. Gen. et Sp. Orch.* 403.—Chili.
aurea, *Phil. in Linnaea*, xxix. (1857–58) 48.—Chili.
Austinae, A. Gray, in Proc. Am. Acad. xii. (1877) 83 = Cephalanthera Oregana.
barbata, *Lindl. in Hook. Journ. Bot.* i. (1834) 5.—Chili.
Besseri, *Reichb. f. in Linnaea*, xxii. (1849) 862.—Chili.
biserialis, *Griseb. in Goett. Abh.* xxiv. (1879) 339.—Reg. Argent.
Bergii, *Hieron. in Bol. Acad. Nac. Cordova*, iii. (1879) 380.—Reg. Argent.
bletioides, Lindl. in Brand. Quart. Journ. Roy. Instit. N. S. i. (1827) 50 = ulanthoides.
calopogon, *Phil. in Linnaea*, xxix. (1857–58) 47.—Chili.
campestris, *Poepp. Fragm. Syn. Pl. Chil.* 15.—Chili.
chlorosticta, *Phil. in Linnaea*, xxix. (1857–58) 52.—Chili.
chrysantha, *Poepp. Fragm. Syn. Pl. Chil.* 15.—Chili.
chrysochlora, *Phil. in Linnaea*, xxix. (1857–58) 50.—Chili.
collina, *Phil. l. c.* 54.—Chili.
Commersonii, *Brongn. in Duperr. Voy. Bot.* 189.—Reg. Magell.
crispa, *Lindl. Gen. et Sp. Orch.* 401.—Chili.
cristata, *Lindl. in Hook. Journ. Bot.* i. (1834) 4.—Chili.
crocata, *Phil. in Linnaea*, xxix. (1857–58) 51.—Chili.
cuneata, *Lindl. Gen. et Sp. Orch.* 400.—Chili.
cygnaea, *Phil. in Linnaea*, xxxiii. (1864–65) 241.—Chili.
cylindrostachya, *Poepp. Fragm. Syn. Pl. Chil.* 15.—Chili.
dasypogon, *Phil. in Linnaea*, xxxiii. (1864–65) 243.—Chili.
decipiens, Poepp. Fragm. Syn. Pl. Chil. 17 = cristata, multiflora, Poeppigiana.
densa, *A. Rich. in C. Gay, Fl. Chil.* v. 454.—Chili.
disoides, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1827) 47.—Chili.
fimbriata, *Phil. in Linnaea*, xxix. (1857–58) 49.—Chili.

CHLORAEA:—
fimbriata, Pocpp. Fragm. Syn. Pl. Chil. 15 = Bipinnula mystacina.
Fonkii, *Phil. in Linnaea*, xxix. (1857–58) 55.—Chili.
galeata, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1827) 48.—Chili.
Gaudichaudii, *Brongn. in Duperr. Voy. Bot.* 189.—Reg. Magell.
Gavilu, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1827) 48.—Chili.
Gayana, *A. Rich. in C. Gay, Fl. Chil.* v. 439.—Chili.
grandiflora, *Poepp. Fragm. Syn. Pl. Chil.* 14.—Chili.
gymnoglossa, *Phil. in Linnaea*, xxix. (1857–58) 54.—Chili.
heteroglossa, *Reichb. f. l. c.* xxii. (1849) 863.—Chili.
homopetala, *Phil. l. c.* xxix. (1857–58) 48.—Chili.
incisa, *Poepp. Fragm. Syn. Pl. Chil.* 17.—Chili.
inconspicua, *Phil. in Anal. Univ. Chil.* (1865) II. 321.—Chili.
lamellata, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1827) 49.—Chili.
Lechleri, *Lindl. in Lechl. Berb. Am. Austr.* 53.—Chili.
Lindleyi, *Poepp. Fragm. Syn. Pl. Chil.* 16.—Chili.
litoralis, *Phil. in Linnaea*, xxxiii. (1864–65) 244.—Chili.
longebracteata, *Lindl. in Brand. Quart. Journ. Roy. Instit.* N. S. i. (1827) 48.—Peruv.
longipetala, *Lindl. Gen. et Sp. Orch.* 400.—Chili.
magellanica, *Hook. f. Fl. Antarct.* ii. 350.—Reg. Magellan.
membranacea, *Lindl. Gen. et Sp. Orch.* 401.—Reg. Argent.
multiflora, *Lindl. in Brand. Quart. Journ. Roy. Instit.* i. (1827) 49.—Chili.
nudilabia, *Poepp. Fragm. Syn. Pl. Chil.* 16.—Chili.
obovata, *Phil. in Linnaea*, xxix. (1857–58) 49.—Chili.
odontoglossa, *A. Rich. in C. Gay, Fl. Chil.* v. 455.—Chili.
papillosa, Phil. in Linnaea, xxix. (1857-58) 50 = cylindrostachys.
patagonica, *Phil. l. c.* xxxiii. (1864–65) 245.—Chili.
Pavonii, *Lindl. Gen. et Sp. Orch.* 404.—Chili.
Pearcei, *Phil. in Linnaea*, xxxiii. (1864–65) 243.—Chili.
penicillata, *Reichb. f. Otia Bot. Hamb.* 51.—Patagon.
Philippi, *Reichb. f. in Linnaea*, xxii. (1849) 863.—Chili.
Piquichen, *Lindl. Gen. et Sp. Orch.* 400.—Chili.
Poeppigiana, *A. Rich. in C. Gay, Fl. Chil.* v. 443.—Chili.
pogonata, *Phil. in Linnaea*, xxxiii. (1864–65) 240.—Chili.
prodigiosa, *Reichb. f. l. c.* xxii. (1849) 863.—Chili.
reflexa, *Phil. l. c.* xxix. (1857–58) 51.—Chili.
rypaloglossa, *A. Rich. in C. Gay, Fl. Chil.* v. 457.—Chili.
Sceptrum, *Reichb. f. in Linnaea*, xxii. (1849) 862.—Chili.
secunda, *Phil. l. c.* xxix. (1857–58) 52.—Chili.
semibarbata, *Lindl. Gen. et Sp. Orch.* 403.—Chili.
speciosa, *Poepp. Fragm. Syn. Pl. Chil.* 14.—Chili.
spectabilis, *Phil. in Linnaea*, xxxiii. (1864–65) 240.—Chili.
suaveolens, *Phil. l. c.* xxix. (1857–58) 46.—Chili.
ulanthoides, *Lindl. Gen. et Sp. Orch.* 404.—Chili.
Unguiscati, *Reichb. f. in Linnaea*, xxii. (1849) 864.—Chili.
venosa, *Reichb. f. l. c.*—Chili.
verrucosa, *Phil. l. c.* xxix. (1857–58) 53.—Chili.
virescens, *Lindl. in Brand. Quart. Journ. Roy. Instit.* i. (1827) 51.—Chili.
viridiflora, *Poepp. Fragm. Syn. Pl. Chil.* 14.—Chili.
volucris, *Lindl. in Hook. Journ. Bot.* i. (1834) 4.—Chili.

CHLORANTHUS, Sw. in Phil. Trans. lxxvii. (1787) 359. *CHLORANTHACEAE*, Benth. & Hook. f. iii. 134.
CREODUS, Lour. Fl. Cochinch. 88 (1790).
CRYPHAEA, Ham. in Brewst. Edinb. Journ. Sc. ii. (1825) 11.
NIGRINA, Thunb. Nov. Gen. 58 (1783).
PEPERIDIA, Reichb. Consp. 212ª (1828).
SAINTLEGERIA, Cordem. in Baill. Adansonia, iii. (1862–63) 300.

CHLORANTHUS:—
SARCANDRA, Gardn. in Calc. Journ. Nat. Hist. vi. (1846) 348.
STROPHA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4; ed. II. 66 (1827).
TRICERCANDRA, A. Gray, in Perry, Jap. Exped. ii. 318 (1857).
Blumeanus, Cordem. in Baill. Adansonia, iii. (1862–63) 296 = serratus.
brachystachys, *Blume, Fl. Jav. Chloranth.* 13. t. 2.—Ind. or.; Malaya; China.
ceylanicus, Miq. Fl. Ind. Bat. i. 1. 802 = brachystachyus.
denticulatus, Cordem. in Baill. Adansonia, iii. (1862–63) 296 = brachystachys.
elatior, R. Br. in Bot. Mag. sub t. 2190 = officinalis.
erectus, Sweet, Hort. Suburb. Lond. 28 = officinalis.
Fortunei, *Solms, in DC. Prod.* xvi. 1. 476.—China.
grandifolius, *Miq. Fl. Ind. Bat.* i. 1. 802.—Ind. or.
ilicifolius, Blume, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 129 = brachystachyus.
inconspicuus, Blanco, Fl. Filip. ed. II. 54 = officinalis.
inconspicuus, *Sw. in Phil. Trans.* lxxviii. (1787) 359. t. 15.—China; Japon.
indicus, Wight, Ic. t. 1945 = inconspicuus.
insignis, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 108.—Burma.
integrifolius, Schult. f. ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 130 = Cornus suecica.
japonicus, *Siebold, in Nov. Act. Nat. Cur.* xiv. II. (1829) 681.—China; Japon.
macranthera, *Schult. f. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 130.—Japon.
mandshuricus, Rupr. Dec. Pl. Amur. t. 2 = japonicus.
monander, R. Br. in Bot. Mag. sub t. 2190 = brachystachyus.
monostachyus, *R. Br. l. c.* t. 2190.—China.
montanus, Siebold, ex Miq. Ann. Mus. Bot. Lugd. Bat. iii. 129 = brachystachyus.
obtusifolius, Miq. Fl. Ind. Bat. i. 1. 802 = inconspicuus?
officinalis, *Blume, Enum. Pl. Jav.* 79.—Ind. or.; Malaya.
Oldhami, *Solms, in DC. Prod.* xvi. 1. 476.—Japon.
Pernyanus, *Solms, l. c.* 474.—China.
quadrifolius, A. Gray, in Miq. Ann. Mus. Bot. Lugd. Bat. iii. 129 = serratus.
salicifolius, Presl, Epim. Bot. 231 = officinalis.
serratus, *Roem. & Schult. Syst.* iii. 461.—Japon.
sumatranus, Miq. Fl. Ind. Bat. Suppl. 343 = officinalis.

CHLORASTER, Haw. in Phil. Mag. lxiii. 103 (1824) = **Narcissus**, Tourn. (Amaryllid.).
fissus, Haw. Monog. Narciss. 7 = N. viridiflorus.
integer, Haw. l. c. = N. viridiflorus.
oblitteratus, M. Roem. Syn. Ensat. 214 = N. Broussonetii.
viridiflorus, M. Roem. l. c. 213 = N. viridiflorus.

CHLORIS, Sw. Prod. Veg. Ind. Occ. 25 (1788). *GRAMINEAE*, Benth. & Hook. f. iii. 1165.
CHLOROIDES, Fisch. ex Regel, in Ind. Sem. Hort. Petrop. (1863) 28.
EUSTACHYS, Desv. in Nouv. Bull. Soc. Philom. ii. (1800) 188.
HETEROLEPIS, Ehrenb. ex Boiss. Fl. Orient. v. 554 (1881).
MACROSTACHYA, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 408, in nota (1851).
PHACELLARIA, Willd. ex Steud. Nom. ed. II. i. 353, in syn. (1840); ii. 313 (1841).
SCHULTESIA, Spreng. Pugill. ii. 17 (1815).
abyssinica, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 406.—Abyssin.
acicularis, *Lindl. in Mitch. Journ. Trop. Austral.* 31.—Austral.
acuminata, Trin. Sp. Gram. iii. t. 305 = distichophylla.
alba, J. & C. Presl, Rel. Haenk. i. 289 = elegans.
Alberti, *Regel, in Act. Hort. Petrop.* vii. (1880) 650.—Mongolia.
amethystea, *Hochst. in Flora*, xxxviii. (1855) 205.—Abyssin.
andropogonoides, *Fourn. Mex. Pl. Enum. Gram.* 143; *ex Hemsl. Biol. Centr. Am. Bot.* iii. 558.—Mexic.
anoplia, *Fourn. l. c.; ex Hemsl. l. c.*—Mexic.
Aristida, Salzm. ex Steud. Syn. Pl. Gram. 218 = Gymnopogon foliosus.
arundinacea, Nees, ex Steud. l. c. 207 = polydactyla.

CHLORIS:—
bahiensis, *Steud. Syn. Pl. Gram.* 208.—Bras.
barbata, *Sw. Fl. Ind. Occ.* i. 200.—Reg. trop.
Beyreichiana, Steud. Nom. ed. II. i. 353 = seq.
Beyrichana, Kunth, Rev. Gram. i. 89, 289. t. 56 = radiata.
brachystachys, *Anderss. in Peters, Reise Mossamb. Bot.* 556.—Afr. trop.
brevigluma, *Wright, in Sauv. Fl. Cub.* 191.—Ins. Cuba.
breviscta, *Benth. in Hook. Niger Fl.* 566.—Afr. trop.
campulodes, *Trin. ex Steud. Nom.* ed. II. i. 353, nomen.—Hab.?
caribaea, Spreng. Syst. i. 295 = barbata.
caudata, *Trin. ex Bunge, Enum. Pl. Chin. Bor.* 79.—China.
ciliata, *Sw. Prod. Veg. Ind. Occ.* 25.—Am. trop.
compressa, *DC. Cat. Pl. Hort. Monsp.* (1813) 94.—Am. austr.
confertifolia, Trin. in Spreng. Neue Entdeck. ii. 74 = distichophylla.
consanguinea, Kunth, Rev. Gram. i. 89 = polydactyla.
crinita, *Lag. Varied. Cienc.* ii. IV. (1805) 143.—Ins. Philipp.
crinita, Salzm. ex Doell, in Mart. Fl. Bras. ii. III. 64 = radiata.
cruciata, *Sw. Prod. Veg. Ind. Occ.* 25.—Ind. occ.
cryptostachya, *Steud. ex J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 148.—Ins. Cap. Verd.
ctenioides, *Steud. Syn. Pl. Gram.* 423.—Madag.
cucullata, *Bisch. in Ann. Sc. Nat.* Sér. III. xix. (1853) 357.—Mexic.
curtipendula, Michx. Fl. Bor. Am. i. 59 = Bouteloua racemosa.
Cynodon, Trin. Gram. Unifl. 229 = Cynodon Dactylon.
cynodontoides, *Balansa, in Bull. Soc. Bot. Fr.* xix. (1872) 318.—N. Caled.
decora, Nees, ex Steud. Syn. Pl. Gram. 205 = barbata.
Digitaria, H. B. & K. Nov. Gen. et Sp. i. 168 = Leptochloa Digitaria.
digitata, *Steud. Syn. Pl. Gram.* 207.—Ind. or.
distachya, Kunth, Rev. Gram. i. 291. t. 57 = monostachya.
distichophylla, *Lag. Gen. et Sp. Nov.* 4.—Am. austr.
divaricata, *R. Br. Prod.* i. 186.—Austral.
dolichostachya, Lag. Gen. et Sp. Nov. 5 = truncata.
dubia, H. B. & K. Nov. Gen. et Sp. i. 169 = Diplachne dubia.
Durandiana, Schult. Mant. ii. 341 = virgata.
elata, *Desv. Opusc.* 73.—Bras.
elegans, *H. B. & K. Nov. Gen. et Sp.* i. 166.—Mexic.
cleusinoides, *Griseb. Fl. Brit. W. Ind.* 539.—Ind. occ.
elongata, Poir. Encyc. Suppl. ii. 236 = truncata.
emarginata, *Beauv. Agrost.* 158, nomen.—Hab.?
equitans, *Trin. Sp. Gram.* iii. t. 309.—Hab.?
falcata, Sw. in Ges. Naturf. Freunde, Neue Schr. iii. (1801) 160 (= *Campulosus falcatus*).—Afr. austr.
fasciculata, Schrad. in Schult. Mant. ii. 339 = distichophylla.
filiformis, Hort. Par. ex Poir. Encyc. Suppl. ii. 237, n. 8 = Bouteloua tenuis.
filiformis, *Poir. l. c.* 238, n. 12.—Hab.?
floccifolia, Poir. l. c. = Eleusine floccifolia.
floridana, *Vasey, Cat. Gram. Un. St.* 61.—Florida.
foliosa, Willd. Sp. Pl. iv. 924 = Gymnopogon foliosus.
Gayana, *Kunth, Rev. Gram.* i. 89, 293. t. 58.—Afr. trop.
geminata, *Hochst. in Flora*, xxxviii. (1855) 205.—Abyssin.
glabrata, *Anderss. in Peters, Reise Mossamb. Bot.* 557.—Afr. trop.
glauca, *Vasey, Cat. Gram. Un. St.* 61.—Florida.
glaucescens, *Steud. Syn. Pl. Gram.* 206.—Peruv.
gracilis, P. Durand, Chlor. 10 = virgata.
gracilis, H. B. & K. Nov. Gen. et Sp. i. 168 = Leptochloa gracilis.
Gryllus, Honck. Syn. Pl. Germ. i. 487 = Chrysopogon Gryllus.
guineensis, *Schum. & Thonn. Beskr. Guin. Pl.* 55.—Afr. trop.
hispida, *P. Durand, Chlor.* 17.—Ins. Philipp.
Humboldtiana, *Steud. Syn. Pl. Gram.* 205.—Ins. Cuba.

CHLORIS:—
humilis, Kunth, Rev. Gram. i. 89; ii. 531. t. 180 = radiata.
imberbis, Desf. Cat. Hort. Par. ed. III. 15 = submutica.
incompleta, *Roth, Nov. Pl. Sp.* 60.—Ind. or.
inermis, *Trin. Gram. Unifl.* 232.—Mexic.
inflata, *Link, Enum. Hort. Berol.* i. 105.—Calif.; Ins. Philipp.
intermedia, *A. Rich. Tent. Fl. Abyss.* ii. 407.—Abyssin.
leptostachya, *Hochst. ex A. Rich. l. c.*—Abyssin.
longibarba, *Michx. ex Beauv. Agrost.* 158, nomen.—Am. bor.
longifolia, *Steud. Syn. Pl. Gram.* 205.—Ins. Philipp.
macrantha, *Desv. Opusc.* 73.—Arabia.
macrantha, Jaub. & Spach, Illustr. iv. 42 (= Tetrapogonis sp.).—Arabia.
macrostachya, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 408 = Enteropogon macrostachya.
madagascariensis, *Steud. Syn. Pl. Gram.* 206.—Madag.
maritima, Trin. Gram. Unifl. 236 = Cynodon Dactylon.
meccana, *Hochst. ex Steud. Syn. Pl. Gram.* 205.—Aegypt.; Arabia; Beluchist.
megastachya, Schrad. ex Schult. Mant. ii. 340 = truncata.
mendocina, *Phil. Sert. Mend. Alt.* 50.—Chili.
monostachya, Michx. Fl. Bor. Am. i. 59 = Ctenium carolinianum.
monostachya, *Poir. Encyc. Suppl.* ii. 238.—Ind. or.; Ins. Mascar.
montana, *Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 329.—Ind. or.
Moorei, F. Muell. in Linnaea, xxv. (1852) 444 = acicularis.
mucronata, Michx. Fl. Bor. Am. i. 59 = Eleusine aegyptiaca.
multiradiata, *Hochst. in Flora*, xxxviii. (1855) 204.—Abyssin.
myriostachya, *Hochst. l. c.*—Afr. trop.
notocoma, *Hochst. l. c.*—Abyssin.
obtusifolia, *Desv. Opusc.* 73.—Bras.
obtusifolia, *Trin. Gram. Unifl.* 233.—Ins. Philipp.
orthonoton, *Doell, in Mart. Fl. Bras.* ii. III. 64.—Bras.
pallida, *Hack. in Engl. Bot. Jahrb.* vi. (1885) 244.—Austral.
pallida, Willd. Sp. Pl. iv. 926 = radiata.
panicea, *Willd. l. c.* 923.—Ind. or.
paraguaiensis, Steud. Syn. Pl. Gram. 204 = barbata.
paspaloides, *Hochst. in Flora*, xxxviii. (1855) 206.—Abyssin.
paytensis, *Steud. Syn. Pl. Gram.* 207.—Peruv.
pectinata, *Benth. Fl. Austral.* vii. 612.—Austral.
pendula, Salzm. ex Steud. Syn. Pl. Gram. 208 = bahiensis.
penicillata, Jan, ex Trin. Gram. Unifl. 231 = compressa.
penicillata, *Pers. Syn.* i. 87.—Ind. or.
penicillata, Willd. ex Steud. Nom. ed. II. i. 353 = elegans.
peregrina, *P. Durand, Chlor.* 11.—Ins. Philipp.
petraea, *Sw. Prod. Veg. Ind. Occ.* 25.—Reg. trop.
pilosa, *Schum. & Thonn. Beskr. Guin. Pl.* 55.—Afr. trop.
piperita, Michx. ex Steud. Nom. ed. II. i. 353 = Ctenium carolinianum.
poaeformis, H. B. & K. Nov. Gen. et Sp. i. 169 = Leptochloa virgata.
polydactyla, P. Durand, Chlor. 14, 22 = virgata.
polydactyla, *Sw. Prod. Veg. Ind. Occ.* 26.—Am. austr.
polystachya, Lag. Gen. et Sp. Nov. 4 = inermis, Leptochloa virgata.
polystachya, *Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 330.—Ind. or.
Prieurii, *Kunth, Rev. Gram.* i. 89; ii. 441. t. 134.—Afr. trop.
procumbens, P. Durand, Chlor. 16 = Bouteloua tenuis.
propinqua, *Steud. Syn. Pl. Gram.* 204.—Guadaloupe.
prostrata, Poir. Encyc. Suppl. ii. 239 = Eleusine aegyptiaca.
pubescens, *Lag. Varied. Cienc.* ii. IV. (1805) 143.—Peruv.
pubescens, *Peyr. in Sitzb. Acad. Wien, Math. Nat.* xxxviii. (1860) 585.—Am. austr.

CHLORIS:—
pulchra, *Schum. & Thonn. Beskr. Guin. Pl.* 56.—Afr. trop.
Pumilio, *R. Br. Prod.* 186.—Austral.
punctulata, *Hochst. ex Steud. Syn. Pl. Gram.* 205.—Afr. trop.
pycnothrix, Trin. Gram. Unifl. 234 = radiata.
radiata, Heyne, ex Roth, Nov. Pl. Sp. 61 = incompleta.
radiata, *Sw. Prod. Veg. Ind. Occ.* 26.—Am. trop.
repens, Auct. ex Steud. Nom. ed. II. i. 353 = Eleusine indica.
repens, *Hochst. in Flora*, xxxviii. (1855) 204.—Abyssin.
retusa, *Lag. Gen. et Sp. Nov.* 5.—Reg. Argent.
rhachitricha, *Steud. Syn. Pl. Gram.* 205.—Ins. Philipp.
Roxburghii, Edgew. in Journ. As. Soc. Beng. xxi. (1853) 183 = digitata.
Roxburghiana, Schult. Mant. ii. 239 = polystachya.
rufescens, *Lag. Varied. Cienc.* ii. IV. (1805) 143.—Ins. Philipp.
rufescens, *Steud. Syn. Pl. Gram.* 206.—Ins. Maurit.
Sagraeana, *A. Rich. Fl. Cub. Fanerog.* ii. 315.—Ins. Cuba.
Salzmanni, Steud. Syn. Pl. Gram. 206 = radiata.
scariosa, *Beauv. Agrost.* 158, nomen.—Hab.?
scariosa, *F. Muell. Fragm.* vi. 85.—Austral.
sclerantha, Lindl. in Mitch. Journ. Trop. Austral. 31 = ventricosa.
scoparia, Desf. Tabl. ed. I. 14 = radiata.
simplex, *Schum. & Thonn. Beskr. Guin. Pl.* 54.—Afr. trop.
spathacea, *Hochst. ex Steud. Syn. Pl. Gram.* 204.—Abyssin.
subbiflora, *Steud. l. c.* 206.—Am. bor.
subdolichostachya, *C. Muell. in Bot. Zeit.* xix. (1861) 341.—Am. bor.
submutica, *H. B. & K. Nov. Gen. et Sp.* i. 167. t. 50.—Mexic.
subtriflora, *Steud. Syn. Pl. Gram.* 208.—Afr. trop.
Swartziana, *Doell, in Mart. Fl. Bras.* ii. III. 68.—Bras.
Swartzii, C. Muell. in Bot. Zeit. xix. (1861) 341 = petraea.
tacnensis, *Steud. in Lechl. Berb. Am. Austr.* 56.—Peruv.
tenella, *Koen. ex Roxb. Hort. Beng.* 82; *Fl. Ind.* i. 329.—Ind. or.
tenuior, Poir. ex Schult. Mant. ii. 321 = Leptochloa cynosuroides.
tenuis, Poir. Encyc. Suppl. v. 614 = Bouteloua tenuis.
tetrameris, Trin. Gram. Unifl. 235 = Cynodon elongatus.
Tetrapogon, Beauv. Agrost. 158 = Tetrapogon villosus.
triangulata, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 409.—Abyssin.
truncata, *R. Br. Prod.* 186.—Malaya; Austral.
unispicea, *F. Muell. Fragm.* vii. 118.—Austral.
ventricosa, *R. Br. Prod.* 186.—Austral.
villosa, Pers. Syn. i. 87 = Tetrapogon villosus.
virgata, P. Durand, Chlor. 10 = radiata.
virgata, *Sw. Fl. Ind. Occ.* i. 203.—Am. trop.
Wightiana, Nees, ex Steud. Syn. Pl. Gram. 206 = incompleta.

CHLORIZA, Salisb. Gen. Pl. Fragm. 21 (1866) = **Lachenalia**, Jacq. (Liliac.).
fragrans, Salisb. l. c. = L. versicolor.
lucida, Salisb. l. c. = L. lucida.
mediana, Salisb. l. c. = L. mediana.
pallida, Salisb. l. c. = L. pallida.
pustulata, Salisb. l. c. = L. pustulata.
unicolor, Salisb. l. c. = L. versicolor.

CHLOROCAULON, Klotzsch, in Endl. Gen. Suppl. iv. III. 89 (1850) = **Argithamnia**, Sw. (Euphorbiac.).

CHLOROCHLAMYS, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 142 (1868–69) = **Marsdenia**, R. Br. (Asclepiad.).
celebica, Miq. l. c.—Ins. Celebes.

CHLOROCODON, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 113 = **Erica**, Tourn.
scoparium, Fourr. l. c. = E. scoparia.

CHLOROCODON, Hook. f. Bot. Mag. t. 5898 (1871). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 745.
Whitei, *Hook. f. l. c.*—Afr. austr.

CHLOROCREPIS, Griseb. in Goett. Abh. v. (1852) 155 = **Hieracium**, Linn. (Compos.).
staticifolia, Griseb. l. c. = H. staticefolium.

CHLOROGALUM, Kunth, Enum. Pl. iv. 681 (1843). *LILIACEAE*, Benth. & Hook. f. iii. 785.
LAOTHOË, Rafin. Fl. Tellur. iii. 53 (1836), nomen prius.
angustifolium, *Kellogg, in Proc. Calif. Acad.* ii. (1863) 104. t. 30.—Calif.
divaricatum, Kunth, Enum. Pl. iv. 682 = pomeridianum.
Leichtlinii, *Baker, in Gard. Chron.* (1874) I. 689.—Am. bor. occ.
parviflorum, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 243.—Calif.
pomeridianum, *Kunth, Enum. Pl.* iv. 682.—Calif.

CHLOROIDES, Fisch. ex Regel, Ind. Sem. Hort. Petrop. (1863) 28 = **Chloris**, Sw. (Gramin.).
petraeum, Fisch. l. c. = Chloris petraea.

CHLOROMYRON, Pers. Syn. ii. 73 (1807) = **Rheedia**, Linn. (Guttifer.).
verticillatum, Pers. l. c. = R. acuminata.

CHLOROPHORA, Gaudich. in Freyc. Voy. Bot. 508, in nota (1826). *URTICACEAE*, Benth. & Hook. f. iii. 363.
tinctoria, *Gaudich. l. c.*—Mexic.

CHLOROPHYTUM, [Ker-Gawl.] in Bot. Mag. t. 1071 (1808). *LILIACEAE*, Benth. & Hook. f. iii. 788.
ASPHODELOPSIS, Steud. ex Baker, in Journ. Linn. Soc. xv. (1876) 321.
HARTWEGIA, Nees, in Nov. Act. Nat. Cur. xv. II. (1831) 372.
HOLLIA, Heynh. Nom. ii. 303 (1846).
SCHIDOSPERMUM, Griseb. in Lechl. Berb. Am. Austr. 56 (1857).
abyssinicum, Kotschy & Peyr. Pl. Tinn. 46 = laxiflorum.
acaule, *Baker, in Journ. Linn. Soc.* xv. (1876) 327.—Ind. or.
affine, *Baker, in Trans. Linn. Soc.* xxix. (1875) 160. t. 104.—Afr. trop.
Afzelii, *Baker, in Journ. Linn. Soc.* xv. (1876) 323.—Afr. trop.
alismaefolium, *Baker, l. c.* 324.—Afr. trop.
alpinum, *Benth. ex Baker, l. c.* 328.—Ins. Tasman.
amplexicaule, *Baker, l. c.* 325.—Afr. trop.
andongense, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 260.—Afr. trop.
anthericoideum, Dalz. in Hook. Kew Journ. ii. (1850) 141 = tuberosum.
arundinaceum, *Baker, in Journ. Linn. Soc.* xv. (1876) 323.—Ind. or.
attenuatum, *Baker, l. c.* 332.—Ind. or.
blepharophyllum, *Schweinf. ex Baker, l. c.* 327.—Abyssin.
Bowkeri, *Baker, in Saund. Refug. Bot.* t. 352.—Afr. austr.
bracteosum, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 260 = stenopetalum.
Brehmeanum, *Schult. f. Syst.* vii. 454.—Afr. austr.
breviscapum, *Dalz. in Hook. Kew Journ.* ii. (1850) 141.—Ind. or.
Burchellii, *Baker, in Journ. Linn. Soc.* xv. (1876) 330.—Afr. austr.
ciliatum, *Baker, in Journ. Bot.* xvi. (1878) 325.—Afr. trop.
comosum, Baker, in Gard. Chron. (1873) 75 = Sternbergianum.
crispum, *Baker, in Journ. Linn. Soc.* xv. (1876) 331.—Afr. austr.
debile, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 260.—Afr. trop.
decipiens, *Baker, in Journ. Linn. Soc.* xx. (1883) 275.—Madag.
dubium, Schult. f. Syst. vii. 455 = Anthericum Schultesii.
elatum, *R. Br. Prod.* 277.—Afr. austr.
falcatum, Baker, in Saund. Refug. Bot. t. 333 = laxiflorum.
filipendulum, *Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 260.—Afr. trop.
Finlaysonianum, Wall. Cat. n. 5061 (Dracaenae sp.).—Siam.

CHLOROPHYTUM:—
gallabatense, Schweinf. ex Baker, in Journ. Linn. Soc. xv. (1876) 325 = Orchidastrum.
glaucum, *Dalz. in Hook. Kew Journ.* ii. (1850) 143.—Ind. or.
graminifolium, Kunth, Enum. Pl. iv. 606 = Anthericum undulatum.
Hartmannianum, *Schweinf. Beitr. Fl. Aethiop.* 294, nomen.—Afr. trop.
Heyneanum, *Wall. Cat.* n. 5060.—Ind. or.
Heynei, Baker, in Journ. Linn. Soc. xv. (1876) 322 = praec.
inornatum, *Ker-Gawl. in Bot. Mag.* t. 1071.—Afr. trop.
juncifolium, *Baker, in Journ. Linn. Soc.* xv. (1876) 333.—Bras.
Kirkii, *Baker, in Gard. Chron.* (1882) i. 108.—Afr. trop.
lancifolium, *Welw. ex Baker, in Trans. Linn. Soc.* Ser. II. i. (1878) 260.—Afr. trop.
laxiflorum, Baker, in Journ. Linn. Soc. xv. (1876) 328 = laxum.
laxum, *R. Br. Prod.* 277.—Trop. As.; Afr.; Austral.
leucolepis, Welw. ex Baker, in Trans. Linn. Soc. Ser. II. i. (1878) 259 = macrophyllum.
longifolium, *Schweinf. Beitr. Fl. Aethiop.* 294, nomen.—Abyssin.
longipes, *Baker, in Journ. Bot.* xvi. (1878) 325.—Afr. trop.
macrophyllum, *Aschers. in Schweinf. Beitr. Fl. Aethiop.* 294.—Abyssin.
macrosporum, *Baker, in Journ. Linn. Soc.* xv. (1876) 330.—Afr. austr.
madagascariense, *Baker, in Journ. Bot.* xvi. (1878) 326.—Afr. trop.
malabaricum, *Baker, in Journ. Linn. Soc.* xv. (1876) 331.—Ind. or.
micranthum, *Baker, in Journ. Bot.* xvi. (1878) 325.—Afr. trop.
modestum, *Baker, in Journ. Linn. Soc.* xv. (1876) 329.—Afr. austr.
nepalense, *Baker, l. c.* 330.—Reg. Himal.
Nimmonii, Dalz. in Hook. Kew Journ. ii. (1850) 142 = Orchidastrum.
Orchidastrum, *Lindl. Bot. Reg.* t. 813.—Ind. or.; Afr. trop.
parviflorum, Dalz. in Hook. Kew Journ. ii. (1850) 141 = laxum.
petiolatum, *Baker, in Journ. Linn. Soc.* xv. (1876) 326.—Afr. trop.
polyrhizon, *Baker, in Gard. Chron.* (1878) ii. 396.—Afr. trop.
polystachys, *Baker, in Journ. Bot.* xvi. (1878) 326.—Afr. trop.
pubiflorum, *Baker, in Journ. Linn. Soc.* xv. (1876) 329.—Afr. trop.
pulchellum, Kunth, Enum. Pl. iv. 605 = Anthericum pulchellum.
pusillum, *Schweinf. ex Baker, in Journ. Bot.* xvi. (1878) 325.—Afr. trop.
rhizomatosum, *Baker, in Gard. Chron.* (1885) ii. 230.—Afr. trop.
rigidum, Kunth, Enum. Pl. iv. 604 = Anthericum rigidum.
Rutenbergianum, *Vatke, in Bremen Abh.* ix. (1885) 138.—Ins. Nossibé.
schidospermum, *Baker, in Journ. Linn. Soc.* xv. (1876) 326.—Peruv.
sparsiflorum, *Baker, l. c.* 325.—Afr. trop.
stenopetalum, *Baker, l. c.* 331.—Afr. trop.
Sternbergianum, *Steud. Nom.* ed. II. i. 354.—Afr. austr.
suffruticosum, *Baker, in Journ. Bot.* xvi. (1878) 326.—Afric.
tetraphyllum, *Baker, in Journ. Linn. Soc.* xv. (1876) 328.—Abyssin.
Tinneae, *Baker, l. c.* 333.—Afr. trop.
triflorum, Kunth, Enum. Pl. iv. 606 = Anthericum triflorum.
tuberosum, *Baker, in Journ. Linn. Soc.* xv. (1876) 332.—Ind. or.; Abyss.
undulatum, Wall. Cat. n. 5059 = nepalense.
vestitum, *Baker, in Journ. Linn. Soc.* xv. (1876) 326.—Afr. trop.
viscosum, Kunth, Enum. Pl. iv. 605 = Anthericum viscosum.
xerotinum, F. Muell. Fragm. i. 63 = laxum.

CHLOROPHYTUM, Pohl, ex DC. Prod. iv. 540 (1830) = **Spermacoce**, Dill. (Rubiac.).
nervosum, Pohl, l. c. 548 = S. Poaya.
pratense, Pohl, l. c. = S. Poaya.

CHLOROPYRON, Behr, in Proc. Calif. Acad. i. (1855) 61 = **Cordylanthus**, Nutt. (Scrophul.).
palustre, Behr, l. c. = Cordylanthus maritimus.

CHLOROSA, Blume, Bijdr. 420. t. 31 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 616.
gracilis, *Blume, l. c.*—Java.
latifolia, *Blume, l. c.*—Java.

CHLOROSPATHA, Engl. in Regel, Gartenfl. (1878) 97. t. 933. *AROIDEAE*, Benth. & Hook. f. iii. 977.
Kolbii, *Engl. l. c.*—N. Granat.

CHLOROSTEMMA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 398 = **Asperula**, Linn. (Rubiac.).
odoratum, Fourr. l. c. = A. odorata.

CHLOROSTIS, Rafin. Princ. Somiol. 26 (1814) = **Chlora**, Ren. (Gentian.).

CHLOROXYLON, Rumph. ex Scop. Introd. 208 (1777); DC. Prod. i. 625 (1824). *MELIACEAE*, Benth. & Hook. f. i. 340.
Dupada, Buch.-Ham. Journ. Mysore, i. 184 = Boswellia serrata.
Swietenia, *DC. Prod.* i. 625.—Ind. or.

CHLOROXYLUM, P. Br. Hist. Jamaic. 187. t. 7. f. 1 (1756); Adans. Fam. ii. 303 (1763) (Quid?).

CHLORYLLIS, E. Mey. Comm. Pl. Afr. Austr. 149 (1835) = **Dolichos**, Linn. (Legumin.).
pratensis, E. Mey. l. c. = D. Chloryllis.

CHNOANTHUS, Phil. in Anal. Univ. Chili (1862) ii. 405 = **Gomphrena**, Linn. (Amarantac.).
mendocinus, Phil. l. c.; et in Linnaea, xxxiii. (1864–65) 223.—Chili.
ovatus, Phil. Sert. Mend. Alt. 41.—Chili.

CHOCHO, Adans. Fam. ii. 500 (1763) = **Sechium**, P. Br. (Cucurbit.).

CHODONDENDRON, Bosc, Encycl. Agric. vii. 296 (1822), sphalm. = **Chondodendron**, Ruiz & Pav. (Menispermac.).

CHOENOMELES, Lindl. in Trans. Linn. Soc. xiii. (1822) 97 = **Pyrus**, Linn. (Rosac.).
alba, Carr. in Rev. Hortic. (1876) 410 = P. japonica.
cancellata, Pritz. Ic. Ind. i. 250, sphalm. = P. sinensis, *Poir.*
japonica, Lindl. in Trans. Linn. Soc. xiii. (1822) 97 = P. japonica.

CHOERADODIA, Herb. Amaryll. 87 (1837) = **Strumaria**, Jacq. (Amaryllid.).
chilensis, Herb. l. c. = S. chilensis.

CHOERADOPLECTRON, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 436 = **Habenaria**, Willd. (Orchid.).
Spiranthes, Schau. l. c. = H. laccrtifera.

CHOEROPHILLUM, Neck. Dclic. Gallo-Belg. i. 147 (1768) = **Chaerophyllum**, Hoffm. (Umbell.).

CHOEROSERIS, Link, Handb. i. 798 (1829) = **Picris**, Linn. (Compos.).
heterosperma, Link, l. c. 799 = Crepis Dioscoridis.
rhagadioloides, Link, l. c. = P. Sprengeriana.
Sprengeriana, Link, l. c. = P. Sprengeriana.

CHOETOPHORA, Franch. & Sav. Enum. Pl. Jap. ii. 548 (1879), sphalm. = CHAETOSPORA, R. Br. = **Schoenus**, Linn. (Cyperac.).

CHOISYA, H. B. & K. Nov. Gen. et Sp. vi. 4. t. 513 (1823). *RUTACEAE*, Benth. & Hook. f. i. 297.
JULIANIA, La Llave, in La Llave & Lex. Nov. Veg. Desc. fasc. ii. 4 (1825).
PLENCKIA, Moç. & Sesse, ex DC. Prod. i. 724 (1824).
grandiflora, Regel, Gartenfl. (1876) 257; in Act. Hort. Petrop. iv. (1876) 336 = ternata.
ternata, *H. B. & K. Nov. Gen. et Sp.* vi. 6. t. 513.—Mexic.

CHOMELIA, Jacq. Enum. Pl. Carib. 1 (1760). *RUBIACEAE*, Benth. & Hook. f. ii. 103.
ANISOMERIS, Presl, Symb. Bot. ii. 5. t. 54 (1833).
angustifolia, *Benth. in Hook. Journ. Bot.* iii. (1841) 235.—Guiana.
anisomeris, *Muell. Arg. in Flora*, lviii. (1875) 451.—Bras.
barbinervis, *Moric. ex Benth. in Linnaea*, xxiii. (1850) 446.—Bras.
brasiliana, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 183, partim.—Bras.
brasiliana, A. Rich. l. c. partim = Vauthieri.
estrellana, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
fasciculata, *Sw. Fl. Ind. Occ.* i. 238.—Jamaica.
filipes, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 41.—Nicaragua.
hirsuta, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 107.—Bras.
intercedens, *Muell. Arg. in Flora*, lviii. (1875) 451.—Bras.
malancoides, *Muell. Arg. l. c.* 452.—Bras.
Martiana, *Muell. Arg. l. c.* 451.—Bras.
obtusa, *Cham. & Schlecht. in Linnaea*, iv. (1829) 185.—Bras.
obtusa, Mart. in Flora, xxiv. (1841) ii. Beibl. 71 = Martiana.
occidentalis, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
odoratissima, *Teijsm. & Binn. Cat. Hort. Bog.* 384, nomen.—Hab.?
oligantha, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
parviflora, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 41.—Bras.
pedunculosa, *Benth. in Linnaea*, xxiii. (1850) 445.—Bras.
Pohliana, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
pubescens, *Cham. & Schlecht. in Linnaea*, iv. (1829) 187.—Bras.
ribesioides, *Benth. ex A. Gray, in Proc. Am. Acad.* iv. (1860) 38.—Bras.
sandvicensis, *A. Gray, l. c.*—Ins. Sandvic.
scricea, *Muell. Arg. in Flora*, lviii. (1875) 451.—Bras.
sessilis, *Muell. Arg. l. c.*—Bras.
spinosa, *Jacq. Enum. Pl. Carib.* 12.—Venezuela.
tenuiflora, *Benth. in Hook. Journ. Bot.* iii. (1841) 235.—Guiana.
transiens, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 457.—Bras.
triacantha, *Griseb. Cat. Pl. Cub.* 133.—Cuba.
tristis, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.
Vauthieri, *Muell. Arg. l. c.*—Bras.
versicolor, Spreng. Syst. i. 410 = Scolosanthus versicolor.
vulpina, *Muell. Arg. in Flora*, lviii. (1875) 452.—Bras.

CHOMELIA, Linn. Gen. ed. I. 55 (1737); cf. Richter, Codex Linn. 183 = **Tarenna**, Gaertn. (Rubiac.).

CHOMELIA, Vell. Fl. Flum. i. t. 107 (1827) = **Ilex**, Tourn.
amara, Vell. l. c.—Bras.

CHONA, D. Don, in Edinb. N. Phil. Journ. xvii. (1834) 155 = **Erica**, Linn.
sanguinea, D. Don, l. c. = E. Nivenia.

CHONAIS, Salisb. Gen. Pl. Fragm. 135 (1866) = **Hippeastrum**, Herb. (Amaryllid.).
vittata, Salisb. l. c. = H. vittatum.

CHONDODENDRON, Penth. & Hook. f. Gen. i. 34 (1862), sphalm. = **Odontocarya**, Miers (Menisperm.).

CHONDODENDRON, Ruiz & Pav. Prod. 132 (1794). *MENISPERMACEAE*, Benth. & Hook. f. i. 963.
BOTRYOPSIS, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 43.
acmulum, Miers, l. c. Ser. III. xix. (1867) 192 = platyphyllum.
cinerascens, Miers, l. c. = platyphyllum.
convolvulaceum, Poepp. & Endl. Nov. Gen. et Sp. ii. 65 = tomentosum.

CHONDODENDRON:—
cretosum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xix. (1867) 192.—Peruv.
hederifolium, Miers, l. c. Ser. II. vii. (1851) 44 = tomentosum.
nemophilum, *Miers, Contrib.* iii. 319.—Bras.
obscurum, *Miers, l. c.* 314.—Bras.
ovatum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. xix. (1867) 192.—Bras.
platyphyllum, *Miers, l. c.*—Bras.
scabrum, Miers, l. c. Ser. II. vii. (1851) 44 = tomentosum.
tamoides, *Miers, l. c.*—Guiana.
tomentosum, *Ruiz & Pav. Syst. Veg.* 261.—Peruv.

CHONDRACHNE, R. Br. Prod. 220 (1810) = **Lepironia**, Rich. (Cyperac.).
articulata, R. Br. l. c. = L. mucronata.

CHONDRACHYRUM, Nees, in Lindl. Introd. Nat. Syst. ed. II. 449 (1836) = **Melica**, Linn. (Gramin.).
scabrum, Nees, l. c. 450.—Bras.

CHONDRILLA, [Tourn.] Linn. Syst. ed. I (1735); Gen. ed. I. 241 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 524.
CALYCOCORSUS, F. W. Schmidt, Samml. Phys. Aufs. i. 271 (1795).
PELTIDIUM, Zollikof. Nat. Anz. (1820); Bartl. Ord. 145 (1830).
WIBELIA, Roehl. Deutschl. Flora, ii. 426 (1813).
WILLEMETIA, Neck. Elem. i. 50 (1790).
ZOLLIKOFERIA, Nees, in Bluff & Fingerh. Comp. Fl. Germ. ii. 305 (1825).
acanthifolia, Poir. Encyc. Suppl. ii. 330 = Lactuca acanthifolia.
acantholepis, Boiss. Diagn. Ser. I. xi. 48 = juncea.
acanthophylla, Borckh. ex Reichb. Fl. Germ. Excurs. 271 = juncea.
alba, Lam. Encyc. ii. 79 = Prenanthes alba.
altissima, Lam. l. c. 78 = Prenanthes altissima.
ambigua, Fisch. ex Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 398 = pauciflora.
angustissima, Hegetschw. Fl. Schw. 762 = juncea.
aspera, *Poir. Encyc. Suppl.* ii. 329.—Hab.?
asplenifolia, Poir. l. c. 330. = Launaea aspleniifolia.
auriculata, Wall. Cat. n. 3269 = Lactuca dissecta.
brevirostris, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 32.—Sibir. altaic.
canescens, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 397.—Soongar.
capitata, Sieber, ex DC. Prod. vii. 180 = Launaea glomerata.
chinensis, Poir. Encyc. Suppl. ii. 331 = Lactuca versicolor.
crepidinea, Poir. l. c. 331 = Prenanthes crepidinea.
crepoides, Lapeyr. Hist. Abr. Pl. Pyr. 462 = Lactuca saligna.
crepoides, *Murr. Syst. Veg.* ed. XIII. 595.—Mont. Pyren.
crepoides, Reich. Fl. Moeno-Francof. 59 = juncea.
cymosa, Poir. Encyc. Suppl. ii. 330 = Prenanthes cymosa.
debilis, Poir. l. c. 332 = Lactuca debilis.
Decandollei, Spreng. ex DC. Prod. vi. 158 = Crepis foetida.
dentata, Poir. Encyc. Suppl. ii. 328 = Lactuca denticulata.
erysimifolia, Poir. l. c. 331 = Lactuca muralis.
fontinalis, Wall. Cat. n. 3268 = Lactuca polycephala.
Gaudini, Hegetschw. in Suter, Fl. Helv. ed. II. ii. 162 = juncea.
glomerata, C. Koch, in Linnaea, xxiii. (1850) 673 = juncea.
gracilis, Wall. Cat. n. 3267 = Lactuca gracilis.
graminea, *Bieb. Fl. Taur. Cauc.* ii. 244.—Reg. Caucas. et Himal.; Sibiria.
graminea, Ledeb. Fl. Alt. iv. 148 = piptocoma.
graminifolia, Fisch. Cat. Hort. Gorenk. (1812) 33 = graminea.
hastata, Poir. Encyc. Suppl. ii. 332 = Lactuca denticulata.
hastata, Wall. Cat. n. 3271 = Lactuca hastata.
hieracioides, Roth, in Usteri, Ann. Bot. x. 50 = Tolpis coronopifolia.
hispida, Desf. Cat. Hort. Par. ed. III. 144 = juncea.
hispida, Poir. Encyc. Suppl. ii. 328 = Lactuca microcephala.
illinoiĕnsis, Poir. l. c. 331 = Prenanthes aspera.

CHONDRILLA:—
indica, Steud. Nom. ed. I. 191 = Lactuca Heyneana.
intybacea, Frivald. in Flora, xviii. (1835) 335 = juncea.
intybacea, Ledeb. Fl. Ross. ii. 809 = pauciflora.
japonica, Lam. Encyc. ii. 79 = Crepis japonica.
juncea, Ledeb. Fl. Alt. iv. 147 = brevirostris.
juncea, *Linn. Sp. Pl.* 796.—Europ.; As. med.
laciniata, Stev. in Bull. Soc. Nat. Mosc. xxix. (1856) II. 410 = juncea.
laevigata, Pursh, Fl. Am. Sept. ii. 497 = Pyrrhopappus carolinianus.
lanceolata, Poir. Encyc. Suppl. ii. 329 = Crepis integra.
latifolia, Bieb. Fl. Taur. Cauc. ii. 244 = juncea.
lejosperma, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 456.—Sibir. altaic.
longifolia, Buch.-Ham. in Wall. Cat. sub n. 3274 = Lactuca polycephala.
longifolia, Wall. Cat. n. 3273 = Lactuca longifolia.
lutea, Dulac, Fl. Hautes-Pyr. 495 = juncea.
lyrata, Poir. Encyc. Suppl. ii. 332 = Crepis japonica.
maracandica, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 380.—Turkestan.
multiflora, Poir. Encyc. Suppl. ii. 332 = Crepis japonica.
muralis, Lam. Fl. Fr. ii. 105 = Lactuca muralis.
nudicaulis, Linn. Mant. 278 = Launaea nudicaulis.
paniculata, Lam. Encyc. ii. 78 = prenanthoides.
pauciflora, D. Don, in Trans. Linn. Soc. xvi. (1833) 180 = Pyrrhopappus pauciflorus.
pauciflora, *Ledeb. Fl. Alt.* iv. 148.—Rossia; Sibiria.
Peltidium, *Moritz, Fl. Schw.* 361.—Europ.
phaeocephala, *Rupr. Sert. Tiansch.* 59.—As. centr.
piestocarpa, *Boiss. Fl. Orient.* iii. 793.—Persia.
pinnata, Lam. Encyc. ii. 79 = Sonchus leptocephalus.
pinnatifida, Poir. Encyc. Suppl. ii. 330 = Prenanthes cymosa.
piptocoma, *Fisch. Mey. & Avè-Lall. Ind. Sem. Hort. Petrop.* viii. 54.—Sibir. altaic.
Prenanthes, Steud. Nom. ed. I. 191 = seq.
prenanthoides, *Vill. Voy.* (1812) 16.—Europ.
pulchra, Lam. Fl. Fr. ii. 106 = Phaecasium lampsanoides.
purpurea, Lam. l. c. 105 = Prenanthes purpurea.
pusilla, *Boiss. Diagn.* Ser. I. vii. 11.—Persia; Turkestan.
racemosa, Poir. Encyc. Suppl. ii. 331 = Lactuca Heyneana.
ramosissima, *Sibth. & Sm. Fl. Graec. Prod.* ii. 128.—Graecia.
ramosissima Zeyher, ex DC. Prod. vii. 176 = Lactuca viminea.
Rauillieri, Walp. Rep. ii. 694, sphalm. = Rouillieri.
repens, Lam. Encyc. ii. 79 = Lactuca repens.
rhombifolia, Humb. ex Willd. Sp. Pl. iii. 1535-36 = Prenanthes rhombifolia.
rigens, Reichb. Fl. Germ. Excurs. 271 = juncea.
Rouillieri, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 456.—Sibir. altaic.
rubicunda, Poir. Encyc. Suppl. ii. 329 = Prenanthes alba.
ruderalis, Gaertn. ex Steud. Nom. ed. I. 191, 655 = Lactuca muralis.
runcinata, Wall. Cat. n. 3272 = Crepis japonica.
sagittata, Wall. l. c. n. 3270 = Lactuca sagittarioides.
sarmentosa, Poir. Encyc. Suppl. ii. 529 = Launaea pinnatifida.
Schraderi, Steud. Nom. ed. II. ii. 393, in syn. = aspera.
secunda, Royle, ex C. B. Clarke, Comp. Ind. 276 = Launaea secunda.
Sessaeana, D. Don, in Trans. Linn. Soc. xvi. (1833) 180 = Pyrrhopappus Sessaeanus.
sessiliflora, Lam. Fl. Fr. ii. 104 = Lactuca viminea.
setulosa, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iii. 402.—Himal. bor. occ.
sonchifolia, Poir. Encyc. Suppl. ii. 330 = Lactuca Heyneana.
soongarica, *Stschegl. in Bull. Soc. Nat. Mosc.* xxvii. (1854) I. 179.—Soongar.
squarrosa, Poir. Encyc. Suppl. ii. 332 = Lactuca brevirostris.
stipitata, *Sch. Bip. in Linnaea*, xv. (1841) 553.—Europ.
stricta, *Ledeb. Fl. Alt.* iv. 146.—Sibir. altaic.
tenuifolia, Lam. Encyc. ii. 78 = Prenanthes purpurea.
tenuis, Buch.-Ham. in Wall. Cat. n. 3274 = Lactuca polycephala.

CHONDRILLA:—
triquetra, Steud. Nom. ed. I. 191; ed. II. i. 354 = Lactuca triquetra.
versicolor, Sch. Bip. in Mus. Senckenb. iii. (1845) 49, in obs. 2 = Lactuca Fischeriana.
viminea, Lam. Encyc. ii. 27 = Lactuca viminea.
virgata, Poir. Encyc. Suppl. ii. 328 = Prenanthes virgata.
virgata, J. & C. Presl, Delic. Prag. 116 = juncea.
viscosa, Gilib. Fl. Lituan. i. 232 = juncea.

CHONDROCARPUS, Nutt. Gen. Am. ii. in corrigend. (1818) = **Hydrocotyle**, Linn. (Umbellif.).
asiaticus, Nutt. l. c.—Cf. i. 177 = H. asiatica.
erectus, Wats. Bibliog. Ind. N. Am. Bot. 425, sphalm. = H. repanda.
repandus, Nutt. Gen. Am. ii. in corrigend.—Cf. i. 177 = H. repanda.
sibthorpioides, Sweet, Hort. Brit. ed. I. 185 = H. rotundifolia.
triflorus, Nutt. Gen. Am. ii. in corrigend.—Cf. i. 177 = H. asiatica.

CHONDROCARPUS, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 268 = **Astragalus**, Tourn. (Legumin.).
brachycarpos, Stev. l. c. = A. brachycarpus.
clavatus, Stev. l. c. xxix. (1856) II. 149 = A. clavatus.
elongatus, Stev. l. c. iv. (1832) 268 = A. elongatus.
incurvus, Stev. l. c. = A. incurvus.
macrorhizus, Stev. l. c. = A. macrorrhizus.
nummularioides, Stev. l. c. xxix. (1856) II. 149 = A. nummularioides.
platyphyllus, Stev. l. c. = A. platyphyllus.

CHONDROCHILUS, Phil. in Linnaea, xxviii. (1856) 711 = **Chaetanthera**, Ruiz & Pav. (Compos.).
crenatus, Phil. l. c.—Chili.
involucratus, Phil. Fl. Atac. 27.—Chili.
parvifolius, Phil. in Linnaea, xxxiii. (1864-65) 114.—Chili.

CHONDROLAENA, Nees, Fl. Afr. Austr. Gram. 133 (1841) = **Prionachne**, Nees (Gramin.).
dentata, Steud. Nom. ed. II. i. 355, sphalm. = seq.
phalaroides, Nees, Fl. Afr. Austr. Gram. 134 = P. Ecklonii.

CHONDRODENDRON, Spreng. Syst. ii. 155, in syn. (1825) = **Chondodendron**, Ruiz & Pav. (Menisp.).

CHONDROLOMIA, Nees, in Mart. Fl. Bras. ii. 1. 173 (1842) = **Scleria**, Berg. (Cyperac.).
Selloviana, Nees, l. c. 174 = S. Sellowiana.

CHONDROPETALUM, Rottb. Progr. 12 (1772); Desc. 10. t. 3 (1773) = **Elegia**, Linn. (Restiac.).
deustum, Rottb. ll. cc. = E. deusta.
nudum, Rottb. ll. cc. 12, 11 = E. nuda.
tectorum, Rafin. Fl. Tellur. iv. 33 = E. deusta.

CHONDROPIS, Rafin. Fl. Tellur. iii. 29 (1836) = **Exacum**, Linn. (Gentian.).
trincrvis, Rafin. l. c. = E. zeylanicum.

CHONDRORHYNCHA, Lindl. Orch. Linden. 12 (1846). *ORCHIDEAE*, Benth. & Hook. f. iii. 548.
Chestertoni, *Reichb. f. in Gard. Chron.* (1879) II. 648.—N. Granat.
fimbriata, *Reichb. f. in Saund. Refug. Bot.* t. 107.—N. Granat.
rosea, *Lindl. Orch. Linden.* 13.—Venezuela.

CHONDROSEA, Haw. Saxifr. Enum. 10 (1821) = **Saxifraga**, Tourn.
Aizoon, Haw. l. c. 11 = S. Aizoon.
aizoopsis, Jord. & Fourr. Ic. Pl. Eur. ii. 5 = S. Aizoon.
alpicola, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 34 = S. Aizoon.
aretioides, Haw. Saxifr. Enum. 14 = S. aretioides.
beugesiaca, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 33 = S. Aizoon.
brevicaulis, Jord. & Fourr. Ic. Pl. Eur. ii. 3 = S. Aizoon.
Burseriana, Haw. Saxifr. Enum. 15 = S. Burseriana.
caesia, Haw. l. c. 14 = S. caesia.
calcarea, Jord. & Fourr. Ic. Pl. Eur. ii. 3 = S. Aizoon.
calyciflora, Haw. Saxifr. Enum. 13 = S. media.

CHONDROSEA:—
carthusiana, Jord. & Fourr. Ic. Pl. Eur. ii. 3 = Saxifraga Aizoon.
cartilaginea, Haw. Saxifr. Enum. 12 = S. cartilaginea.
diapensoides, Haw. l. c. 13 = S. diapensoides.
flexipes, Jord. & Fourr. Ic. Pl. Eur. ii. 3 = S. Aizoon.
glabrata, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 34 = S. Aizoon.
glareosa, Jord. & Fourr. Ic. Pl. Eur. ii. 4 = S. Aizoon.
gracilescens, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 35 = S. Aizoon.
intermedia, Haw. Saxifr. Enum. 11 = S. intermedia.
juniperina, Haw. l. c. 15 = S. juniperifolia.
laevigata, Jord. & Fourr. Ic. Pl. Eur. ii. 5 = S. Aizoon.
leptopetala, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 34 = S. Aizoon.
longifolia, Haw. Saxifr. Enum. 11 = S. longifolia.
luteo-purpurea, Haw. l. c. 13 = S. luteopurpurea.
mutata, Haw. l. c. = S. mutata.
nitida, Haw. l. c. 16 = S. nitida.
orophila, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 33 = S. Aizoon.
patens, Haw. Saxifr. Enum. 14 = S. caesia.
patulipes, Jord. & Fourr. Ic. Pl. Eur. ii. 4 = S. Aizoon.
petrophila, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 34 = S. Aizoon.
punctillata, Jord. & Fourr. l. c. 33 = S. Aizoon.
pusilla, Jord. & Fourr. Ic. Pl. Eur. ii. 3 = S. Aizoon.
pyramidalis, Haw. Saxifr. Enum. 10 = S. Aizoon.
refracta, Jord. & Fourr. Ic. Pl. Eur. ii. 4 = S. Aizoon.
rosella, Jord. & Fourr. l. c. = S. Aizoon.
rosularis, Haw. Saxifr. Enum. 12 = S. Aizoon.
tenella, Haw. l. c. 15 = S. tenella.
valida, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 32 = S. Aizoon.
Vandellii, Haw. Saxifr. Enum. 15 = S. Vandelli.
virgata, Jord. & Fourr. Ic. Pl. Eur. ii. 4 = S. Aizoon.
viridula, Jord. & Fourr. Brev. Pl. Nov. fasc. i. 33 = S. Aizoon.

CHONDROSIA, Benth. in Journ. Linn. Soc. xix. (1881) 104, err. typ. = seq.

CHONDROSIUM, Desv. Journ. Bot. iii. (1813) 68 = seq.

CHONDROSUM, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = **Bouteloua**, Lag. (Gramin.).
Aschenbornianum, Nees, in Linnaea, xix. (1847) 692 = B. oligostachya.
ciliatum, Beauv. Agrost. 41, nomen = Chloris ciliata.
eriopodum, Torr. in Emory, Notes Mil. Reconnois. 153 = B. eriopoda.
exile, Fourn. Mex. Pl. Enum. Gram. 137, ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
foeneum, Torr. in Emory, Notes Mil. Reconnois. 153. t. 12 = B. hirsuta.
gracile, H. B. & K. Nov. Gen. et Sp. i. 176. t. 58 = B. oligostachya.
hirsutum, Sweet, Hort. Brit. ed. I. 455, nomen = seq.?
hirtum, H. B. & K. Nov. Gen. et Sp. i. 176. t. 59 = B. hirsuta.
Humboldtianum, Kunth, Rev. Gram. i. 93 = B. ovata.
humile, Beauv. Agrost. 41 = B. prostrata.
Karwinskii, Fourn. Mex. Pl. Enum. Gram. 137, ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.
microstachyum, Fourn. l. c. 138, ex Hemsl. l. c. 562 = B. polystachya.
oligostachyum, Torr. in Marcy, Rep. 300 = B. oligostachya.
Parryi, Fourn. Mex. Pl. Enum. Gram. 150, ex Hemsl. Biol. Centr. Am. Bot. iii. 562 = B. polystachya.
polystachyum, Benth. Bot. Voy. Sulph. 56 = B. polystachya.
procumbens, Desv. in Nouv. Bull. Soc. Philom. ii. (1810) 188 = B. tenuis.
prostratum, Sweet, Hort. Brit. ed. I. 455 = B. prostrata.
scorpioides, Kunth, Rev. Gram. i. 94 = B. scorpioides.
simplex, Kunth, l. c. = B. prostrata.
subscorpioides, C. Muell. in Bot. Zeit. xiv. (1856) 347.—Calif.
tenue, Beauv. Agrost. 41 = B. tenuis.
Virletii, Fourn. Mex. Pl. Enum. Gram. 136, ex Hemsl. Biol. Centr. Am. Bot. iii. 564.—Mexic.

CHONDROSPERMUM, Wall. Cat. n. 2837 (1832) = **Myxopyrum**, Blume (Oleac.).
coriaceum, Wall. l. c. n. 2838 = M. smilacifolium.
laurifolium, Voigt, Hort. Suburb. Calc. 548 = M. smilacifolium.
smilacifolium, Wall. Cat. n. 2837 = M. smilacifolium.

CHONEMORPHA, G. Don, Gen. Syst. iv. 76 (1836).
APOCYNACEAE, Benth. & Hook. f. ii. 720.
EPICHYSIANTHUS, Voigt, Hort. Suburb. Calc. 523 (1845).
antidysenterica, G. Don, l. c. = Holarrhena antidysenterica.
bantamensis, G. Don, l. c. = Echites bantamensis.
convolvuloides, *G. Don, l. c.*—Ins. S. Thomas.
coriacea, G. Don, l. c. = Echites coriacea.
cristata, G. Don, l. c. = Aganosma marginata.
densiflora, G. Don, l. c. = Echites densiflora.
dichotoma, G. Don, l. c. = Aganosma caryophyllata.
grandiflora, G. Don, l. c. = macrophylla.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 662.—Reg. Himal.
inflata, G. Don, Gen. Syst. iv. 76 = Echites inflata.
macrophylla, *G. Don, l. c.*—Ind. or.; Malaya.
malabarica, *G. Don, l. c.*—Ind. or.
mollis, Miq. Fl. Ind. Bat. ii. 444 = macrophylla.
pubescens, G. Don, in Sweet, Hort. Brit. ed. III. 458 = Holarrhena antidysenterica.
reticulata, G. Don, Gen. Syst. iv. 76 = Cryptolepis Buchanani.
vestita, G. Don, l. c. = Wrightia tomentosa?

CHORETIS, Herb. Amaryll. 219. f. 35 (1837) = **Hymenocallis**, Salisb. (Amaryllid.).
galvestonensis, Herb. l. c. 221. t. 41. f. 34, 35 = H. galvestonensis.
glauca, Herb. l. c. 220. t. 35. f. 1 = H. glauca.

CHORETRUM, R. Br. Prod. 354 (1810). *SANTALACEAE*, Benth. & Hook. f. iii. 228.
Candollei, *F. Muell. ex Benth. Fl. Austral.* vi. 219.—Austral.
chrysanthum, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 23 = glomeratum.
glomeratum, *R. Br. Prod.* 354.—Austral.
lateriflorum, A. DC. in DC. Prod. xiv. 675 = Candollei.
lateriflorum, *R. Br. Prod.* 354.—Austral.
oxycladum, *F. Muell. Fragm.* i. 21.—Austral.
pauciflorum, A. DC. in DC. Prod. xiv. 676 = lateriflorum.
Preissianum, Miq. in Lehm. Pl. Preiss. i. 608 = Leptomeria Preissiana.
spicatum, *F. Muell. Fragm.* i. 21.—Austral.
spinosum, Miq. in Lehm. Pl. Preiss. i. 609 = Leptomeria spinosa.

CHORIBENA, Steud. Nom. ed. II. i. 355 (1840), sphalm. = **Chorilaena**, Endl. (Rutac.).

CHORICARPHA, Boeck. in Flora, xli. (1858) 19 = **Lepironia**, Rich. (Cyperac.).
aphylla, Boeck. l. c. 20 = L. mucronata.

CHORICERAS, Baill. Adansonia, xi. (1873) 119 = **Dissiliaria**, F. Muell. (Euphorbiac.).
australiana, Baill. l. c. 120 = D. tricornis?

CHORILAENA, Endl. Enum. Pl. Hueg. 17 (1837). *RUTACEAE*, Benth. & Hook. f. i. 295.
angustifolia, F. Muell. in Trans. Phil. Soc. Viet. i. (1855) 10 = Phebalium diosmeum.
hirsuta, *Benth. Fl. Austral.* i. 357.—Austral.
myoporoides, *Endl. ex Hcynh. Nom.* ii. 140.—Austral.
quercifolia, *Endl. Enum. Pl. Hueg.* 17.—Austral.

CHORIOPHYLLUM, Benth. in Hook. Ic. Pl. t. 1280 (1829). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 280.
malayanum, *Benth. l. c.*—Malaya.

CHORIOZANDRA, Steud. Nom. ed. II. i. 355 (1840), sphalm. = **Chorizandra**, R. Br. (Cyperac.).

CHORIPETALUM, A. DC. in Trans. Linn. Soc. xvii. (1834) 131 = **Embelia**, Burm. f. (Myrsin.).
aurantiacum, A. DC. l. c. = E. viridiflora.

CHORIPETALUM:—
australianum, F. Muell. Fragm. iii. 36 (= *Samara australiana*).—Austral.
Benthamii, Hance, in Walp. Ann. iii. 10 = *obovatum*.
obovatum, Benth. in Hook. Lond. Journ. Bot. i. (1842) 490.—Ins. Hongkong.
Reevesianum, Wall. ex Voigt, Hort. Suburb. Calc. 338, nomen = *obovatum*?
undulatum, A. DC. in Trans. Linn. Soc. xvii. (1837) 132 = E. Nagushia.
viridiflorum, A. DC. in DC. Prod. viii. 88 = E. viridiflora.

CHORISANDRA, Benth. & Hook. f. Gen. iii. 1057 (1883) err. typ. = **Chorizandra**, R. Br. (Cyperac.).

CHORISANDRA, Wight, Ic. vi. 13. t. 1994 (1853) = **Fluggea**, Willd. (Euphorbiac.).
pinnata, Wight, l. c. = F. microcarpa.

CHORISANTHERA, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 109 = **Pentarhaphia**, Lindl. (Gesnerac.).
tenera, Oerst. l. c. = P. tenera.

CHORISEMA, Fisch. Ind. Sem. Hort. Petrop. viii. 54 (1841) err. typ. = **Chorizema**, Labill. (Legumin.).

CHORISIA, H. B. & K. Nov. Gen. et Sp. v. 295. t. 485 (1821). *MALVACEAE*, Benth. & Hook. f. i. 210.
crispiflora, *H. B. & K. l. c.* 297.—Bras.
insignis, *H. B. & K. l. c.* t. 485. f. 1.—Peruv.
rosea, *Seem. Bot. Voy. Herald*, 84.—N. Granat.
speciosa, *St. Hil. Pl. Us. Bras.* t. 63.—Bras.
ventricosa, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 102. t. 9 = crispiflora.

CHORISIS, DC. Prod. vii. 177 (1838) = **Lactuca**, Tourn. (Compos.).
repens, DC. l. c. 178 = L. repens.

CHORISMA, D. Don, in Edin. N. Phil. Journ. (1828-29) 308 = CHORISIS, DC. = **Lactuca**, Linn. (Compos.).
repens, D. Don, l. c. = L. repens.

CHORISMA, Lindl. in Sweet, Geran. sub t. 79 (1820) = **Pelargonium**, L'Hérit. (Geraniac.).
flavescens, Eckl. & Zeyh. Enum. 71 = P. tetragonum.
tetragona, Lindl. in Sweet, Geran. sub t. 79 = P. tetragonum.

CHORISPERMUM, R. Br. in Ait. Hort. Kew. ed. II. iv. 129 (1812) = **Chorispora**, R. Br. (Crucifer.).
arcuatum, Andrz. ex DC. Syst. ii. 436 = Chorispora tenella.
ibericum, Andrz. ex DC. l. c. 437 = Chorispora iberica.
sibiricum, Andrz. ex DC. l. c. = Chorispora sibirica.
strictum, Andrz. ex DC. l. c. 436 = Diptychocarpus strictus.
tenellum, R. Br. in Ait. Hort. Kew. ed. II. iv. 129 = Chorispora tenella.

CHORISPORA, R. Br. ex DC. Syst. ii. 435 (1821). *CRUCIFERAE*, Benth. & Hook. f. i. 102.
CHORISPERMUM, R. Br. in Ait. Hort. Kew. ed. II. iv. 129 (1812).
Bungeana, Fisch. & Mey. in Enum. Pl. Nov. Sehrenk, 96 = exscapa.
compressa, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 385 = iberica?
elegans, Cambess. in Jacquem. Voy. Bot. 15. t. 14 = sabulosa.
exscapa, *Bunge, in Ledeb. Fl. Ross.* i. 169.—Sibir. altaic.
Greigi, *Regel, Gartenfl.* (1879) 257. t. 984.—As. temp.
hispida, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) i. 266.—As. temp.
iberica, *DC. Syst.* ii. 437.—As. Min.; Persia.
macropoda, *Trautv. in Bull. Soc. Nat. Mosc.* xxxiii. (1860) i. 109.—Soongaria.
persica, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 384.—Persia.
sabulosa, *Cambess. in Jacquem. Voy. Bot.* 15. t. 15.—Reg. Himal.; Afghan.
sibirica, *DC. Syst.* ii. 437.—Sibir. altaic.; Tibet.

CHORISPORA:—
sibirica, Regel & Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) i. 266 = macropoda.
songarica, *Fisch. & Mey. Enum. Pl. Nov. Schrenk*, ii. 57.—Soongar.
stenopetala, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 239.—Taschkent.
stricta, DC. Syst. ii. 436 = Diptychocarpus strictus.
syriaca, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 384.—Syria.
tenella, *DC. Syst.* ii. 435.—Reg. Caucas. et Himal.; Persia; Afghanist.

CHORISTEA, Thunb. Prod. Pl. Cap. 163 (1800); Nov. Gen. 179 (1801) = **Didelta**, L'Hérit. (Compos.).
carnosa, Thunb. ll. cc. 163, 180 = D. carnosa.
glabra, Soland. ex Poir. Encyc. v. 507 = D. spinosa.
spinosa, Thunb. Prod. Pl. Cap. 163; Nov. Gen. 180 = D. spinosa.

CHORISTES, Benth. Pl. Hartw. 63 (1840) = **Deppea**, Cham. & Schlecht. (Rubiac.).
cornifolia, Benth. l. c. 64 = D. cornifolia.
excelsa, Benth. l. c. = Psychotria excelsa.
obtusiflora, Benth. l. c. 63 = D. obtusiflora.

CHORISTYLIS, Harv. in Hook. Lond. Journ. Bot. i. (1842) 19. *SAXIFRAGEAE*, Benth. & Hook. f. i. 647.
rhamnoides, *Harv. l. c.*—Afr. austr.

CHORITAENIA, Benth. in Benth. & Hook. f. Gen. i. 907 (1867). *UMBELLIFERAE*, Benth. & Hook. f. l. c.
PAPPEA, Sond. & Harv. Fl. Cap. ii. 562 (1862).

CHORIZANDRA, Benth. & Hook. f. Gen. iii. 275, in textu (1880) = CHORISANDRA, Wight = **Fluggea**, Willd. (Euphorb.).

CHORIZANDRA, R. Br. Prod. 221 (1810). *CYPERACEAE*, Benth. & Hook. f. iii. 1057.
cymbaria, *R. Br. l. c.*—Austral.
enodis, *Nees, in Lehm. Pl. Preiss.* ii. 73.—Austral.
multiarticulata, *Nees, in Ann. Nat. Hist.* Ser. I. vi. (1841) 48.—Austral.
sphaerocephala, *R. Br. Prod.* 221.—Austral.

CHORIZANTHE, R. Br. ex Benth. in Trans. Linn. Soc. xvii. (1836) 416. t. 19. *POLYGONACEAE*, Benth. & Hook. f. iii. 93.
ACANTHOGONUM, Torr. in Pacif. Rail. Rep. iv. 132 (1856).
MUCRONEA, Benth. in Trans. Linn. Soc. xvii. (1836) 419. t. 20.
TRIGONOCARPUS, Bert. ex Steud. Nom. ed. II. ii. 710 (1841).
angustifolia, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 167.—Calif.
brevicornu, *Torr. in Bot. Mex. Bound.* 177.—Am. bor. occ.
Breweri, *S. Wats. in Proc. Am. Acad.* xii. (1877) 270.—Calif.
californica, *A. Gray, in Proc. Bost. Soc. Nat. Hist.* vii. (1861) 149.—Calif.
Clevelandi, *Parry, in Proc. Davenp. Acad.* iv. (1884) 62.—Am. bor. occ.
commisuralis, *Remy, in C. Gay, Fl. Chil.* v. 287.—Chili.
corrugata, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 198.—Calif.
dasyantha, *Phil. in Linnaea*, xxxiii. (1864–65) 227.—Chili.
diffusa, *Benth. Pl. Hartw.* 333.—Calif.
discolor, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 167.—Calif.
Douglasii, *Benth. in Trans. Linn. Soc.* xvii. (1836) 418.—Calif.
fernandina, *S. Wats. Bot. Calif.* ii. 481.—Calif.
fimbriata, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 168.—Calif.
flavescens, *Phil. in Linnaea*, xxxiii. (1864–65) 226.—Chili.
frankenioides, *Remy, in C. Gay, Fl. Chil.* v. 288.—Chili.
glabrescens, *Benth. in Trans. Linn. Soc.* xvii. (1836) 418.—Chili.
Kingii, *Phil. in Anal. Univ. Chil.* (1873) 536.—Chili.

CHORIZANTHE:—
laciniata, *Torr. in Pacif. Rail. Rep.* vii. 19.—Calif.
Lastarriaea, *Parry, in Proc. Davenp. Acad.* iv. (1884) 63.—Am. bor. occ.
leptoceras, *S. Wats. in Proc. Am. Acad.* xii. (1877) 269.—Calif.
Macraei, *Benth. in Trans. Linn. Soc.* xvii. (1836) 417.—Chili.
membranacea, *Benth. l. c.* 419.—Chili.
nudicaule, Benth. Pl. Hartw. 332 = Douglasii.
nudicaule, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 166.—Calif.
Orcuttiana, *Parry, in Proc. Davenp. Acad.* iv. (1884) 54.—Am. bor. occ.
Palmeri, *S. Wats. in Proc. Am. Acad.* xii. (1877) 271.—Calif.
paniculata, *Benth. in Trans. Linn. Soc.* xvii. (1836) 417.—Chili.
Parryi, *S. Wats. in Proc. Am. Acad.* xii. (1877) 271.—Calif.
pedunculatis, *Benth. in Trans. Linn. Soc.* xvii. (1836) 416.—Chili.
perfoliata, *A. Gray, in Proc. Bost. Soc. Nat. Hist.* vii. (1861) 148.—Calif.
polygonoides, *A. Gray, in Proc. Am. Acad.* viii. (1870) 197.—Calif.
procumbens, A. Gray, in Proc. Bost. Soc. Nat. Hist. vii. (1861) 148 = Xanti.
procumbens, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 167.—Calif.
pungens, *Benth. in Trans. Linn. Soc.* xvii. (1836) 419.—Calif.
ramosissima, *Benth. l. c.* 417.—Chili.
rigida, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 198.—Am. bor. occ.
rosea, *Phil. in Linnaea*, xxxiii. (1864–65) 225.—Chili.
spinosa, *S. Wats. Bot. Calif.* ii. 481.—Calif.
staticoides, *Benth. in Trans. Linn. Soc.* xvii. (1836) 418.—Calif.
stellulata, *Benth. Pl. Hartw.* 333.—Calif.
tenuis, *Phil. in Linnaea*, xxxiii. (1864–65) 227.—Chili.
Thurberi, *S. Wats. in Proc. Am. Acad.* xii. (1877) 269.—Am. bor. occ.
umbellata, *Phil. in Linnaea*, xxxiii. (1864–65) 225.—Chili.
uncinata, *Nutt. in Journ. Acad. Philad.* N. S. i. (1847) 167.—Calif.
uniaristata, *Torr. & Gray, in Proc. Am. Acad.* viii. (1870) 195.—Calif.
vaginata, *Benth. in Trans. Linn. Soc.* xvii. (1836) 417.—Chili.
valida, *S. Wats. in Proc. Am. Acad.* xii. (1877) 271.—Calif.
virgata, *Benth. in Trans. Linn. Soc.* xvii. (1836) 416. t. 19. f. 1.—Chili.
Watsoni, *A. Gray, in Proc. Am. Acad.* viii. (1870) 199.—Calif.
Wheeleri, *S. Wats. l. c.* xii. (1877) 272.—Calif.
Xanti, *S. Wats. l. c.*—Calif.

CHORIZEMA, Labill. Voy. i. 403. t. 21 (1799). *LEGUMINOSAE*, Benth. & Hook. f. i. 467.
ORTHOTROPIS, Benth. in Lindl. Swan River App. 16 (1839).
angustifolium, *Benth. Enum. Pl. Hueg.* 28.—Austral.
Baueri, Benth. in Ann. Wien. Mus. ii. (1838) 71 = Mirbelia pungens.
Baueri, Meissn. in Lehm. Pl. Preiss. i. 34; ii. 209 = Henchmanni.
Baxteri, R. Grah. in Edinb. Phil. Journ. (Apr.-June 1831) 187 = Oxylobium scandens.
callistachys, F. Muell. Fragm. iv. 18 = Oxylobium callistachys.
capillipes, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 255 = angustifolium.
Chandleri, Hort. ex Nichols. Dict. Gard. i. 318 = varium.
cordatum, *Lindl. Bot. Reg.* (1838) t. 10.—Austral.
cordifolium, F. Muell. Fragm. iv. 17 = Oxylobium cordifolium.
coriaceum, Sm. in Trans. Linn. Soc. ix. (1808) 254 = Oxylobium retusum.
costatum, Meissn. in Lehm. Pl. Preiss. i. 33 = Dicksoni.
cytisoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 256.—Austral.

CHORIZEMA:—
daviesioides, Meissn. in Lehm. Pl. Preiss. i. 34 = Mirbelia daviesioides.
denticulatum, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 252 = angustifolium.
Dicksoni, *R. Grah. in Maund, Botanist*, iii. t. 106.—Austral.
diversifolium, *A. DC. Pl. Rar. Jard. Genèv.* 7e *Not.* (1836) 44. t. 8.—Austral.
ellipticum, F. Muell. Pl. Vict. ii. 39 = Oxylobium ellipticum.
ericifolium, *Meissn. in Lehm. Pl. Preiss.* ii. 209.—Austral.
ericoides, Hort. Belg. ex Heynh. Nom. ii. 141 = praec. ?
flavum, Henfrey, in Moore & Ayres, Gard. Mag. Bot. i. 73. t. 7 = cordatum.
gomphocarpum, *Benth. ex A. Gray, in Bot. U. St. Expl. Exped.* i. 380.—Austral.
Henchmanni, *R. Br. in Bot. Reg.* t. 986.—Austral.
heterophyllum, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 255 = Oxylobium heterophyllum.
humile, *Turcz. l. c.* 254.—Austral.
ilicifolium, *Labill. Voy.* i. 405.—Austral.
ilicifolium, Steud. Nom. ed. II. i. 356, sphalm. = Oxylobium trilobatum.
Leichhardtii, F. Mucll. Fragm. iv. 20 = Isotropis filicaulis.
lineare, F. Muell. Fragm. iv. 17 = Oxylobium lineare.
magnifolium, F. Muell. l. c. 18 = Mirbelia racemosa.
nanum, Sims, Bot. Mag. t. 1032 = ilicifolium.
nervosum, *T. Moore, in Gard. Comp.* (1852) 122.—Austral.
ovatum, Lindl. Bot. Reg. t. 1528 = rhombeum.
parviflorum, *Benth. in Ann. Wien. Mus.* ii. (1838) 71.—Austral.
parvifolium, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 253 = nervosum.
platylobioides, DC. Prod. ii. 103 = Mirbelia grandiflora.
pubescens, Turcz. in Bull. Soc. Nat. Mosc. xxvi. (1853) i. 256 = Oxylobium carinatum.
Pultenaeae, F. Muell. Fragm. iv. 19 = parviflorum.
reticulatum, *Meissn. in Lehm. Pl. Preiss.* i. 34.—Austral.
rhombeum, *R. Br. in Ait. Hort. Kew.* ed. II. iii. 9.—Austral.
rhombeum, Lodd. Bot. Cab. t. 1619 = diversifolium.
rhynchotropis, Meissn. in Lehm. Pl. Preiss. ii. 209 = Henchmanni.
scandens, Sm. in Trans. Linn. Soc. ix. (1808) 253 = Oxylobium scandens.
sericeum, Sm. l. c. = Brachysema undulatum.
spartioides, Lodd. Bot. Cab. t. 1953 = Isotropis striata.
spectabile, Lindl. Bot. Reg. (1841) t. 45 = diversifolium.
superbum, Lem. Illustr. Hortic. i. (1854) t. 29 = cordatum.
triangulare, Lindl. Bot. Reg. t. 1513 = ilicifolium.
trigonum, *Turcz. in Bull. Soc. Nat. Mosc.* xxvi. (1853) i. 254.—Austral.
trilobatum, DC. Prod. ii. 103, sphalm. = seq.
trilobum, Sm. in Trans. Linn. Soc. ix. (1808) 253 = Oxylobium trilobatum.
varium, *Benth. ex Lindl. Bot. Reg.* (1839) t. 49; *Misc.* 45.—Austral.

CHORIZOTHECA, Muell. Arg. in Linnaea, xxxii. (1863) 76 = **Pseudanthus**, Sieber (Euphorbiac.).
micrantheoides, Muell. Arg. l. c. = P. virgatus.

CHOROBANCHE, Presl, Bot. Bemerk. 91 (1845) (Chorobane) = **Orobanche**, Linn.
indica, Presl, l. c. = O. indica.

CHOROSEMA, Brongn. Enum. Gen. 126 (1843) = **Chorizema**, Labill. (Legumin.).

CHOROZEMA, Sm. in Trans. Linn. Soc. ix. (1808) 251, sphalm. = **Chorizema**, Labill. (Legumin.).

CHORYZEMUM, Bosc, Encyc. Agric. vii. 297 (1822) = praec.

CHOTECKIA, Steud. Nom. ed. II. i. 356 (1840), sphalm. = seq.

CHOTEKIA, Opiz & Corda; in Flora, xiii. (1830) i. 33 = **Dysophylla**, Blume (Labiat.).
sericea, Opiz & Corda, l. c. 35 = D. quadrifolia.

CHOULETTIA, Pomel, Nouv. Mat. Fl. Atl. 81 (1874) = **Gaillonia**, A. Rich. (Rubiac.).
Reboudiana, Pomel, l. c. = G. Reboudiana.

CHRESTA, Vell. Fl. Flum. viii. t. 150, 151 (1827) = **Eremanthus**, Linn. (Compos.).
alpestris, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 239 = Vernonia alpestris.
angustifolia, Gardn. l. c. 240 = E. angustifolius.
cordata, Vell. Fl. Flum. viii. t. 150 = E. sphaerocephalus.
exsucca, DC. Prod. v. 85 = E. exsuccus.
intermedia, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 236 = E. sphaerocephalus.
lanceolata, Vell. Fl. Flum. viii. t. 151 = Vanillosmopsis erythropappa.
plantaginifolium, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 241 = E. plantaginifolius.
pycnocephala, DC. Prod. v. 85 = E. pycnocephalus.
scapigera, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 241 = E. scapigerus.
speciosa, Gardn. l. c. 240 = E. speciosus.
sphaerocephala, DC. in Deless. Ic. Sel. iv. 2. t. 6 = E. sphaerocephalus.
spathulaefolia, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 241 = E. plantaginifolius.

CHRISANTHEMUM, Neck. Delic. Gallo-Belg. ii. 358 (1768) = **Chrysanthemum**, Tourn. (Compos.).

CHRISOSPLENIUM, Neck. Delic. Gallo-Belg. i. 190 (1768) = **Chrysosplenium**, Tourn. (Saxifrag.).

CHRISTANNIA, Presl, Rel. Haenk. ii. 91. t. 67 (1831) = **Banara**, Aubl. (Samydac.).
salicifolia, Presl, l. c. = B. incana.

CHRISTANNIA, Walp. Rep. i. 360 (1842), sphalm. = **Christiana**, DC. (Tiliac.).

CHRISTIA, Moench, Meth. Suppl. 39 (1802) = **Lourea**, Neck. (Legumin.).
lunata, Moench, l. c. 40 = L. vespertilionis.

CHRISTIANA, DC. Prod. i. 516 (1824). *TILIACEAE*, Benth. & Hook. f. i. 232.
africana, *DC. l. c.*—Afr. trop.
cordifolia, Hook. f. in Niger Fl. 238 = africana.
madagascariensis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 542.—Madag.

CHRISTISONIA, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 153. *OROBANCHACEAE*, Benth. & Hook. f. ii. 982.
CAMPBELLIA, Wight; Ic. iv. 5. t. 1424, 1425 (1850).
OLIGOPHOLIS, Wight, l. c. t. 1422 (1850).
albida, *Thw. ex Hook. f. Fl. Brit. Ind.* iv. 323.—Zeylan.
aurantiaca, Wight, Ic. iv. t. 1486 = bicolor.
bicolor, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 160.—Ind. or.
calcarata, *Wight, Ic.* iv. t. 1426.—Ind. or.
grandiflora, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 155 = tricolor.
Hookeri, *C. B. Clarke, ex Hook. f. Fl. Brit. Ind.* iv. 321.—Reg. Himal.
Lawii, *Wight, Ic.* iv. t. 1427.—Ind. or.
neilgherrica, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 157.—Ind. or.
pallida, Gardn. l. c. 159 = bicolor.
Stocksii, Hook. Ic. Pl. t. 836 = calcarata.
subacaulis, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 162.—Ind. or.
Thwaitesii, *Trim. in Journ. Bot.* xxiii. (1885) 240.—Zeylan.
tricolor, *Gardn. in Calc. Journ. Nat. Hist.* viii. (1847) 156.—Zeylan.
tubulosa, *Benth. ex Hook. f. Fl. Brit. Ind.* iv. 321.—Ind. or.
unicolor, Gardn. in Calc. Journ. Nat. Hist. viii. (1847) 161 = neilgherrica.

CHRISTMANNIA, Dennst. Schluess. Hort. Malab. 30 (1818). 'Genus delendum,' Benth. & Hook. f. iii. 150.
CURONDIA, Rafin. Sylva Tellur. 154 (1838).
Courondi, Dennst. l. c.—Ind. or.

CHRISTOLEA, Cambess. in Jacquem. Voy. Bot. 17. t. 17 (1837). *CRUCIFERAE*, Benth. & Hook. f. i. 80.
crassifolia, *Cambess. l. c.*—Tibet. occ.

CHRISTOPHORIANA, Tourn. ex Rupp. Fl. Jen. ed. Hall. 94 (1745) = **Actaea**, Linn. (Ranunculac.).
spicata, Moench, Meth. 276 = A. spicata.
vulgaris, Rupr. Fl. Caucas. 44 = A. spicata.

CHRISTYA, Ward & Harv. in Hook. Journ. Bot. iv. (1842) 134. t. 21 = **Strophanthus**, DC. (Apocynac.).
speciosa, Ward & Harv. l. c. = S. capensis.

CHRITMUM, Brot. Fl. Lusit. i. 436 (1804), sphalm. = **Crithmum**, Linn. (Umbell.).

CHROILEMA, Bernh. Del. Sem. Hort. Erf. (1840); et in Linnaea, xv. (1841) Litt. 90 = **Aplopappus**, Cass.? (Compos.).
subcanescens, Bernh. ll. cc.—Chili.

CHROMANTHUS, Phil. Sert. Mend. Alt. 14 (1871). *PORTULACEAE*?

CHROMATOLEPIS, Dulac, Fl. Hautes-Pyr. 526 (1867) = **Carlina**, Linn. (Compos.).
acaulis, Dulac, l. c. 527 = Carl. acaulis.
lanata, Dulac, l. c. = Carl. lanata.
magna, Dulac, l. c. = Carl. acanthifolia.
vulgaris, Dulac, l. c. = Carl. vulgaris.

CHROMATOPOGON, F. W. Schmidt, Samml. Phys. Aufs. i. 275 (1795) = **Scorzonera**, Linn. (Compos.).
Jacquini, F. W. Schmidt, l. c. = S. hirsuta.

CHROMOCHITON, Cass. in Dict. St. Nat. lvi. 220 (1828) = **Cassinia**, R. Br. (Compos.).
aculeata, Cass. l. c. = Cassinia aculeata.
affinis, Cass. l. c. = Cassinia aculeata.
aurea, Cass. l. c. = Cassinia aurea.

CHROMOLAENA, DC. Prod. v. 133 (1836) = **Eupatorium**, Linn. (Compos.).
alternifolia, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 465 = E. stachyophyllum.
Candolleana, Gardn. l. c. 464 = E. horminoides.
decumbens, Gardn. l. c. 466 = E. decumbens.
epaleacea, Gardn. l. c. vi. (1847) 436 = E. lupulinum.
horminoides, DC. Prod. v. 133 = E. horminoides.
leucocephala, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 465 = E. horminoides.
pratensis, Gardn. l. c. i. (1842) 176 = E. macrocephalum.

CHROMOLEPIS, Benth. Pl. Hartw. 40 (1840). *COMPOSITAE*, Benth. & Hook. f. ii. 366.
heterophylla, *Benth. l. c.*—Mexic.

CHRONE, Dulac, Fl. Hautes-Pyr. 512 (1867) = **Eupatorium**, Linn. (Compos.).
heterophylla, Dulac, l. c. = E. cannabinum.

CHRONOBASIS, DC. ex Benth. & Hook. f. Gen. ii. 457 (1873) = **Ursinia**, Gaertn. (Compos.).

CHRONOPAPPUS, DC. Prod. v. 84 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 236.
bifrons, *DC. l. c.*—Bras.

CHROSPERMA, Rafin. Neogenyt. 3 (1825) = **Zygadenus**, Michx. (Liliac.).
laetum, Rafin. l. c. = Z. muscaetoxicum.

CHROZOPHORA, Neck. Elem. ii. 337 (1790). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 305.
CROZOPHORA, A. Juss. Tent. Euph. 27 (1824).
LEPIDOCROTON, Presl, Epim. Bot. 213 (1850).
RICINOIDES, Tourn. ex Moench, Meth. 286 (1794).
TOURNESOLIA, Nissol. ex Scop. Introd. 243 (1777).
Brocchiana, *Schweinf. Pl. Nilot.* 9.—Afr. trop.
Burmanni, Spreng. Syst. iii. 851 = plicata.

CHROZOPHORA:—
gracilis, *Fisch. & Mey. ex Kar. in Bull. Soc. Nat. Mosc.* (1839) 171, nomen; *Boiss. Fl. Orient.* iv. 1140.—As. centr.
hierosolymitana, Spreng. Syst. iii. 850 = verbascifolia.
integrifolia, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 490 = verbascifolia.
mollissima, A. Juss. Tent. Euph. 28 = Mallotus viscinoides.
obliqua, *A. Juss. l. c.*—Afr. bor.; As. occ.; Ind. or.
obliqua, Schweinf. Pl. Nilot. 13 = plicata.
oblongifolia, A. Juss. Tent. Euph. 28 = obliqua.
parvifolia, Klotzsch, ex Schweinf. Pl. Nilot. 11 = plicata.
peltata, Labill. Sert. Austr.-Caled. 74. t. 75 = Codiaeum inophyllum.
plicata, *A. Juss. Tent. Euph.* 28.—Europ. austr.; Afr. bor.; As. occ.; Ind. or.
prostrata, Dalz. & Gibs. Bomb. Fl. 233 = plicata.
Rottleri, A. Juss. Tent. Euph. 28 = plicata.
subulosa, Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 446 = gracilis.
senegalensis, *A. Juss. Tent. Euph.* 28.—Afr. trop.
Sieberi, Presl, Bot. Bemerk. 109 = verbascifolia.
tinctoria, *A. Juss. Tent. Euph.* 28. t. 7.—Europ. austr.; Orient; Ind. or.; Afr. bor.
tinctoria, Muell. Arg. in DC. Prod. xv. ii. 749 = obliqua.
verbascifolia, *A. Juss. Tent. Euph.* 28.—Reg. Mediterr.; Orient.

CHROZORRHIZA, Ehrh. Beitr. iv. 147 (1789) = **Asperula**, Linn. (Rubiac.) (A. tinctoria).

CHRYSA, Rafin. in Med. Repos. N. York, II. v. (1809) 350 = **Coptis**, Salisb. (Ranunculac.).

CHRYSACTINIA, A. Gray, in Mem. Am. Ac. N. S. iv. (1849) (Pl. Fendl.) 93. *COMPOSITAE*, Benth. & Hook. f. ii. 412.
mexicana, *A. Gray, l. c.*—Mexic.

CHRYSACTINIUM, Wedd. Chlor. And. i. 212. t. 39. (1856) = **Liabum**, Adans. (Compos.).
acaule, Wedd. l. c. = L. acaule.

CHRYSALIDOCARPUS, H. Wendl. in Bot. Zeit. xxxvi. (1878) 171. *PALMAE*, Benth. & Hook. f. iii. 882.
lutescens, *H. Wendl. l. c.*—Madag.

CHRYSANGIA, Link, Handb. i. 632 (1829) = **Muschia**, Dum. (Campanul.).
aurea, Link, l. c. = M. aurea.

CHRYSANTHELLINA, Cass. in Dict. Sc. Nat. xxv. 391 (1822) = **Chrysanthellum**, Rich. (Compos.).
fasciculata, Cass. l. c. = Chrysanthellum procumbens.
gracilis, Cass. l. c. 392 = Chrysanthellum procumbens.
Swartzii, Cass. l. c. = Chrysanthellum procumbens.

CHRYSANTHELLUM, Rich. in Pers. Syn. ii. 471 (1807). *COMPOSITAE*, Benth. & Hook. f. ii. 389.
ADENOSPERMUM, Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 318.
CHRYSANTHELLINA, Cass. in Dict. Sc. Nat. xxv. 391 (1828).
COLLAEA, Spreng. Syst. iii. 622 (1826).
HINTERHUBERA, Sch. Bip. ex Benth. & Hook. f. Gen. ii. 389 (1873).
NEURACTIS, Cass. in Dict. Sc. Nat. xxxiv. 496 (1825).
SEBASTIANIA, Bertol. Lucubr. (1822) 37.
UCACOU, Adans. Fam. ii. 131 (1763).
abyssinicum, Sch. Bip. in Flora, xxv. (1842) 440 = Microlecane abyssinica.
americanum, *Vatke, in Bremen Abh.* ix. (1885) 122.—Am. trop.
boliviense, *Sch. Bip. in Bull. Soc. Fr.* xii. (1865) 82, nomen.—Boliv.
crectum, *Anderss. in Vet. Akad. Handl. Stockh.* 1853 (1855) 188.—Ins. Galapag.
indicum, *DC. Prod.* v. 631.—As. et Afr. trop.
integrifolium, *Steetz, in Seem. Bot. Voy. Herald*, 160.—Panama.
procumbens, *A. Rich. in Pers. Syn.* ii. 471.—Am. trop.
pusillum, *Hook. f. in Trans. Linn. Soc.* xx. 1851 214.—Ins. Galapag.
senegalense, DC. Prod. v. 631 = indicum.
Swartzii, DC. l. c. = procumbens.

CHRYSANTHEMOIDES, Tourn. ex Medic. Phil. Bot. i. 159 (1789) = **Osteospermum**, Linn. (Compos.).
caerulea, Moench, Meth. 584 = O. caeruleum.
pisiformis, Medic. Phil. Bot. i. 159 = O. moniliferum.

CHRYSANTHEMUM, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 424.
AMMANTHUS, Boiss. & Heldr. Diagn. Ser. I. xi. 18 (1849).
ARGYRANTHEMUM, Webb, ex Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 258. t. 90 ad 96 (1836–50).
BALSAMITA, Desf. in Act. Soc. Hist. Nat. Par. i. (1792) t. 1.
BRACHANTHEMUM, DC. Prod. vi. 44 (1837).
CENTRACHAENA, Schott, Ind. Sem. Hort. Vindob. (1823), ex Reichb. Ic. Pl. Crit. iv. 47 (1827).
CENTROSPERMUM, Spreng. Nov. Prov. 19 (1818); Syst iii. 362 (1826).
COLEOSTEPHUS, Cass. in Dict. Sc. Nat. xli. 43 (1826).
DECANEURUM, Sch. Bip. Tanacet. 44 (1844).
DENDRANTHEMA, Des Moul. in Act. Soc. Linn. Bord. xx. (1855) 561.
ENDOPAPPUS, Sch. Bip. in Bonplandia, viii. (1860) 369.
GLEBIONIS, Cass. Dict. Sc. Nat. xli. 41 (1826).
GLOSSOPAPPUS, Kunze, in Flora, xxix. (1846) 748.
GYMNOCLINE, Cass. in Bull. Soc. Philom. (1816) 199; et in Dict. Sc. Nat. xx. 119 (1821).
HETERANTHEMIS, Schott, in Oken, Isis, (1818) 821.
HETEROMERA, Pomel, Nouv. Mat. Fl. Atl. 60 (1874).
HYMENOSTEMMA, Kunze, in Flora, xxix. (1846) 699.
ISMELIA, Cass. in Dict. Sc. Nat. xli. 40 (1826).
KREMERIA, Dur. in Duch. Rev. Bot. i. (1846) 364.
LEUCANTHEMUM, [Tourn.] DC. Prod. vi. 45 (1837).
MATRICARIA, Hall. ex Scop. Fl. Carn. ed. II. ii. 147 (1772).
MONOPTERA, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 253 (1836–50).
MYCONIA, Neck. Elem. i. 22 (1790).
OTOSPERMUM, Willk. in Bot. Zeit. xxii. (1864) 251.
PHALACROCARPUM, Willk. l. c. 252 (1864).
PHALACRODISCUS, Less. Syn. Comp. 253 (1832).
PHALACROGLOSSUM, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 276 (1855–60).
PINARDIA, Cass. in Dict. Sc. Nat. xli. 38 (1826).
PLAGIUS, L'Hérit. Diss. ex DC. Prod. vi. 135 (1837).
PONTIA, Bubani, in Nuov. Giorn. Bot. Ital. v. (1873) 318.
PREAUXIA, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 250 (1836–50).
PROLONGOA, Boiss. Voy. Bot. Espagne, 230. t. 29 a (1840).
PYRETHRUM, Hall. Enum. Stirp. Helv. ii. 720 (1742).
RICHTERIA, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1842) 126.
STIGMATOTHECA, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 255 (1836–50).
TRIDACTYLINA, Sch. Bip. Tanacet. 48 (1844).
XANTHOPHTHALMUM, Sch. Bip. l. c. 17 (1844).
absinthifolium, Fisch. ex DC. Prod. vi. 46 = sinuatum.
Achilleae, *Linn. Syst.* ed. XII. 262.—Italia.
achilleaefolium, *DC. Prod.* vi. 68.—Reg. Caucas.; Sibiria.
adustum, Fisch. ex Herd. in Bull. Soc. Nat. Mosc. xxxviii. (1865) i. 415 = arcticum.
alpestre, DC. Prod. vi. 68 = coronopifolium.
alpinum, Bieb. Casp. 211 = caucasicum.
alpinum, *Linn. Sp. Pl.* 889.—Europ.
anethifolium, *Brouss. ex Willd. Enum. Hort. Berol.* 904, in nota.—Ins. Canar.
angustifolium, *Rafin. Herb.* 25; *Atl. Journ.* 178.—Am. bor.
anomalum, *Lag. Varied. Cienc.* ii. iv. (1805) 40.—Hispan.
apetalum, *Hort. ex Steud. Nom.* ed. II. i. 356.—Europ.
aragonense, Asso, Syn. 123. t. 9. f. 1 = pulverulentum.
arcticum, *Linn. Sp. Pl.* 889.—Reg. arct.
argenteum, *Willd. Sp. Pl.* iii. 2146.—Armenia.
artemisiaefolium, Klatt, in Sitz. Akad. Muench. (1878) 88 = C. Richteria.

CHRYSANTHEMUM:—
arvense, Herb. Mex. ex DC. Prod. v. 311 = Xanthocephalum humile.
asteroides, *Desf. Tabl.* ed. II. 118, nomen.—Hab.?
Atkinsoni, *C. B. Clarke, Comp. Ind.* 147.—Reg. Himal.
atlanticum, *Ball, in Journ. Bot.* xi. (1873) 366.—Marocco.
atratum, Georgi, Beschr. Russ. Reich. iii. iv. 1252 = maximum?
atratum, *Linn. Sp. Pl.* ed. II. 1252.—Europ.
Balsamita, Baill. Hist. Pl. viii. 311 = Tanacetum Balsamita.
Balsamita, *Linn. Sp. Pl.* ed. II. 1252.—As. occ.
Barrelieri, DC. Prod. vi. 68 = coronopifolium.
Bauhini, Tausch, in Syll. Ratisb. ii. (1828) 251 = atratum.
bellidiflorum, Willd. ex Steud. Nom. ed. II. i. 356 = Bellis dubia.
Biebersteinianum, Adam, in Weber & Mohr, Beitr. i. 70 = Anthemis Biebersteiniana.
bipinnatum, *Linn. Sp. Pl.* 890.—Reg. arct.
Bocconi, *Pourr. ex DC. Prod.* vi. 61.—Hispan.
breviradiatum, Hort. ex DC. Prod. vi. 64 = coronarium.
Broussonetii, *Balb. ex Pers. Syn.* ii. 461.—Ins. Canar.
caespitosum, *Weinm. Cat. Hort. Dorp.* 43.—Hab.?
carinatum, *Schousb. Vextr. Marokko*, 198. t. 6.—Marocco.
carneum, Steud. Nom. ed. I. 192 = coccineum.
carnosulum, DC. Prod. vi. 65 = Phymaspermum carnosulum.
carnosum, Pers. Syn. ii. 460 = glabrum.
carolinianum, Walt. Fl. Carol. 204 = Boltonia asteroides.
Catananche, *Ball, in Journ. Bot.* xi. (1873) 366.—Marocco.
caucasicum, *Pers. Syn.* ii. 462.—Reg. Caucas.
cebennense, DC. ex Steud. Nom. ed. I. 192, 193 = monspeliense.
centrospermum, Bernh. ex Reichb. Ic. Pl. Crit. iv. 47 = viscosum.
ceratophylloides, All. Fl. Pedem. i. 190 = atratum.
Chamomilla, Bernh. Syst. Verz. Erf. 145 = Matricaria Chamomilla.
cinerariaefolium, *Vis. Fl. Dalmat.* ii. 88. t. 8.—Dalmatia.
cinerarium, Steph. in Ledeb. Fl. Ross. ii. 542 = sinuatum.
coccineum, Sims, Bot. Mag. t. 1080 = coronopifolium.
coccineum, *Willd. Sp. Pl.* iii. 2144.—Reg. Caucas.; Persia.
coronarium, *Linn. Sp. Pl.* 890.—Reg. Mediterr.
coronopifolium, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 208.—Ins. Canar.
coronopifolium, Steud. Nom. ed. II. i. 356 = grandiflorum.
coronopifolium, *Vill. Fl. Delphin.* 98; *Hist. Pl. Dauph.* iii. 202.—Corsica.
coronopifolium, Willd. Herb. ex Boiss. Fl. Orient. iii. 340 = coccineum.
corsicum, Sieber, ex DC. Prod. vi. 47 = coronopifolium, *Vill.*
corymbiferum, *Linn. Sp. Pl.* ed. II. 1251.—Helv.; Austr.; Germ.
corymbiferum, Pall. Reise, i. 196 = corymbosum.
corymbosum, *Linn. Sp. Pl.* 890.—Europ.; Afr. bor.; Reg. Caucas.
crithmifolium, Brouss. ex Willd. Enum. Hort. Berol. 904, in syn.; Buch, in Berl. Acad. Abh. 1816–17 (1819) 375 = anethifolium.
cuneatum, Roxb. Hort. Beng. [101]; Fl. Ind. iii. 436 = Boltonia indica.
daucifolium, *Pers. Syn.* ii. 462.—Hab.?
densum, *Steud. Nom.* ed. I. 192.—Syria.
dentatum, Gilib. Fl. Lituan. i. 219 = C. Leucanthemum.
diffusum, *Vell. Fl. Flum.* viii. t. 128.—Bras.
disciforme, C. A. Mey. Verz. Pfl. Cauc. 75 = Matricaria disciformis.
discoideum, *All. Fl. Pedem.* i. 190.—Italia.
dissectum, *Lowe, in Trans. Camb. Phil. Soc.* vi. (1838) reimpr. 17.—Ins. Canar.
elegans, Steud. Nom. ed. I. 192 = maritimum.
ericaefolium, *Burm. f. Fl. Cap. Prod.* 27.—Afr. austr.
floridum, Salisb. Prod. 202 = frutescens.
flosculosum, *Linn. Sp. Pl.* 890.—Reg. Mediterr.

CHRYSANTHEMUM:—
foeniculaceum, Steud. Nom. ed. I. 192; DC. Prod. vi. 66 = anethifolium.
foliosum, Brouss. Hort. Monsp. (1804); ex DC. Prod. vi. 66 = grandiflorum.
frutescens, *Linn. Sp. Pl.* 887.—Ins. Canar.
frutescens, Thunb. Prod. Pl. Cap. 161 = Thunbergii.
fruticosum, Buch, in Abh. Berl. Acad. 1816–17 (1819) 375 = frutescens?
fruticosum, *Vell. Fl. Flum.* viii. t. 130.—Bras.
fruticulosum, *Ledeb. Fl. Alt.* iv. 117.—Sibir. altaic.
fuscatum, Desf. Fl. Atlant. ii. 283 = Matricaria fuscata.
Gayanum, Ball, in Journ. Linn. Soc. xvi. (1878) 509 = Mawii.
glabratum, Thunb. Prod. Pl. Cap. 161 = Matricaria glabrata.
glabrum, DC. Prod. vi. 49, in syn. = paludosum.
glaucum, Pers. Syn. ii. 461 = multifidum.
Gmelini, Turcz. in Bull. Soc. Nat. Mosc. (1838) 94 = arcticum.
gracile, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 207.—Ins. Canar.
gracilicaule, Dufour, in Ann. Gén. Sc. Phys. vii. (1820) 306 = montanum.
graminifolium, *Linn. Sp. Pl.* 889.—Europ.
grandiflorum, Brouss. Elench. Pl. Hort. Monsp. (1804) 15 = coronopifolium.
grandiflorum, Hook. in Parry, App. Second Voy. 398 = Matricaria inodora.
grandiflorum, Hort. Par. (1819) ex DC. Prod. vi. 47 = lacustre.
grandiflorum, Lapeyr. Hist. Abr. Pl. Pyr. 527 = maximum.
grandiflorum, *Willd. Enum. Hort. Berol.* 904.—Ins. Canar.
Griffithii, C. B. Clarke, Compos. Ind. 148 = Stoliczkai.
haematomma, *Lowe, in Hook. Kew Journ.* viii. (1856) 296.—Ins. Madera.
Halleri, Suter, Fl. Helv. ii. 193 = coronopifolium, *Vill.*
Herminii, Hoffmgg. & Link, Fl. Port. ii. t. 102 = oppositifolium.
heterophyllum, *Willd. Sp. Pl.* iii. 2142.—Europ.
hirtum, Hort. ex Reichb. Ic. Pl. Crit. iv. 47 = viscidum.
hirtum, Thunb. Prod. Pl. Cap. 161 = Matricaria hirta.
hispidum, Rich. ex DC. Prod. vi. 47 = L. pallens.
humile, Spreng. Syst. iii. 584 = Xanthocephalum humile.
hybridum, Guss. Adnot. Cat. Pl. Bocc. 3 = C. Myconis.
incanum, Thunb. Prod. Pl. Cap. 161 = Pentzia virgata.
indicum, *Linn. Sp. Pl.* 889.—China; Japon.
indicum, Thunb. Fl. Jap. 320 = sinense.
inodorum, Linn. Sp. Pl. ed. II. 1253 = Matricaria inodora.
integrifolium, *Richards. in Frankl. 1st Journ. App.* ed. II. 33.—Reg. arct.
intermedium, Hort. Monsp. ex DC. Prod. vi. 66 = grandiflorum.
ircutianum, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) II. 177.—Sibiria.
italicum, *Linn. Mant.* i. 116.—Italia.
japonicum, Thunb. Fl. Jap. 321 = indicum.
lacerum, Auct. ex Desv. in Lam. Encyc. iii. 729, in syn. = pinnatifidum.
laciniatum, Gilib. Fl. Lituan. i. 218 = segetum.
lacustre, *Brot. Fl. Lusit.* i. 379.—Europ.
lanceolatum, *Pers. Syn.* ii. 460.—Europ.
lanceolatum, Vest, in Flora, iii. (1820) 4 = C. Leucanthemum.
lanuginosum, Geners. ex DC. Prod. vi. 57 = corymbosum.
larvatum, *Griseb. ex Pant. in Oestr. Bot. Zeitschr.* xxiii. (1873) 266.—Montenegro.
latifolium, DC. Prod. vi. 69 = grandiflorum.
leptophyllum, DC. l. c. 65 = nodosum.
Leucanthemum, *Linn. Sp. Pl.* 888.—Europ.; As. bor.
lidbeckioides, *Less. Syn. Comp.* 253.—Afr. austr.
macrocephalum, Viv. Fl. Lib. Spec. 56. t. 10. f. 4 = trifurcatum.
macrophyllum, *Waldst. & Kit. Pl. Rar. Hung.* i. 97.—Europ. or.; Reg. Caucas.
macrotum, *Ball, in Journ. Linn. Soc.* xvi. (1878) 509.—Hispan.; Marocco.
majus, Aschers. Fl. Brandenb. 329 = C. Leucanthemum.
Maresii, *Ball, in Journ. Bot.* xi. (1873) 366 (Maressii).—Marocco.
maritimum, *Pers. Syn.* ii. 462.—Europ. occ.

CHRYSANTHEMUM:—
Mawii, *Hook. f. Bot. Mag.* t. 5997.—Morocco.
maximum, *Ramond, in Bull. Soc. Philom.* ii. (1800) 140.—Mont. Pyren.
mexicana, *H. B. & K. Nov. Gen. et Sp.* iv. 299.—Mexic.
millefoliatum, *Linn. Syst.* ed. XII. 563.—Reg. Caucas.; As. Min.
minimum, Brot. Fl. Lusit. i. 379 = pulverulentum?
minimum, Vill. Hist. Pl. Dauph. iii. 202 = alpinum.
monspeliense, *Linn. Sp. Pl.* 889.—Europ.
monspeliense, Schkuhr, Handb. iii. 131. t. 253 = arcticum.
montanum, *Linn. Sp. Pl.* 888.—Europ.
montanum, Willd. Sp. Pl. iii. 2143 = C. Leucanthemum.
moschatum, *Willd. ex Steud. Nom.* ed. I. 192, nomen; ed. II. i. 357.—Hab. ?
multicaule, *Desf. Fl. Atlant.* ii. 282.—Afr. bor.
multifidum, *Desf. Cat. Hort. Par.* ed. III. 170.—Oriens.
Myconis, *Linn. Sp. Pl.* ed. II. 1254.—Reg. Mediterr.
nanum, Hook. Fl. Bor. Am. i. 320 = Blennosperma californicum.
nodosum, *DC. Prod.* vi. 65.—Afr. austr.
ochroleucum, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 208.—Ins. Canar.
odoratum, Lowe, Man. Fl. Mad. i. 462 = pinnatifidum.
oppositifolium, *Brot. Fl. Lusit.* i. 381.—Lusit.
orcastrum, *Hance, in Journ. Bot.* xi. (1878) 108.—China.
orientale, Pers. Syn. ii. 462 = Anthemis Biebersteiniana.
osmitoides, *Harv. Thes. Cap.* ii. 33. t. 152.—Natal.
pallens, *J. Gay, ex Perreym. in Guill. Arch. Bot.* ii. (1833) 545.—Europ.
pallens, Pers. Syn. ii. 461, in syn. = pulverulentum.
pallidum, Mill. Gard. Dict. ed. VIII. n. 12 = pulverulentum.
palmatum, Ledeb. Fl. Ross. ii. 552, sphalm. = parthenifolium.
paludosum, *Poir. Voy. Barb.* ii. 241.—Afr. bor.
palustre, *Pers. Syn.* ii. 461.—Armenia.
parthenifolium, Pers. l. c. 462 = praealtum.
Parthenium, *Bernh. Syst. Verz. Erf.* 145.—Europ.
parviflorum, Pers. Syn. ii. 462 = Matricaria praecox.
pectinatum, *Linn. Sp. Pl.* ed. II. 1255.—Hispan.
perpusillum, Loiscl. in Desv. Journ. Bot. ii. (1809) 369 = Nananthea perpusilla.
pinnatifidum, *Linn. f. Suppl.* 377.—Ins. Madera.
praealtum, *Vent. Jard. Cels*, t. 43.—Reg. Caucas.; As. Min.; Persia.
praecox, DC. Prod. vi. 69 = Matricaria praecox.
pratense, Salisb. Prod. 203 = C. Leucanthemum.
procumbens, Lour. Fl. Cochinch. ii. 499 = indicum.
prostratum, Pers. Syn. ii. 460 = Egletis domingensis.
ptarmicaefolium, Puschk. ex Willd. Sp. Pl. iii. 2151; Pers. Syn. ii. 460 = Achillea Ptarmica.
pubescens, Hort. ex Reichb. Ic. Pl. Crit. iv. 47 = viscidum.
pulverulentum, *Lag. ex Pers. Syn.* ii. 461.—Hispan.
pumilum, *Willd. Enum. Hort. Berol.* 903.—Hab.?
punctatum, Pers. Syn. ii. 461, in syn. = palustre.
purpureum, Pers. l. c. = indicum.
pusillum, *Viv. Fl. Lib. Spec.* 56.—Afr. bor.
radicans, *Pers. Syn.* ii. 462.—Hispan.
ramosum, *F. W. Schmidt, Samml. Phys. Aufs.* i. (1795) 231, nomen.—Europ.
Richteria, *Benth. ex Hook. f. Fl. Brit. Ind.* iii. 315.—Soongaria; Tibet. occ.
rigidum, Vis. Del. Sem. Hort. Patav. (1825) = cinerariaefolium.
roseum, Adam, in Weber & Mohr, Beitr. i. 70 = coccineum.
rotundifolium, *Waldst. & Kit. Pl. Rar. Hung.* iii. 262. t. 236.—Hungar.
Roxburghii, Desf. Cat. Hort. Par. ed. III. 170 = coronarium.
scandens, *Vell. Fl. Flum.* viii. t. 129.—Bras.
segetale, Salisb. Prod. 203 = segetum.
segetum, Forsk. ex DC. Prod. vi. 64 = coronarium.
segetum, *Linn. Sp. Pl.* 889.—Europ.; Afr. bor.; As. occ.
senecioides, Dun. ex DC. Prod. vi. 64 = coronarium.
sericeum, Adam, in Weber & Mohr, Beitr. i. 69 = argenteum.
sericeum, Hoffmgg. & Link, Fl. Port. ii. t. 101 = oppositifolium.
serotinum, *Linn. Sp. Pl.* 888.—Hungar.

CHRYSANTHEMUM:—
setabense, Dufour, ex DC. Prod. vi. 49 = glabrum.
sibiricum, Turcz. ex DC. l. c. 46 = arctieum.
sinense, *Sabine, in Trans. Linn. Soc.* xiv. (1825) 145.—Japon.
sinuatum, *Ledeb. Fl. Alt.* iv. 116.—Sibir. altaic.
spathulatum, *F. W. Schmidt, Samml. Phys. Aufs.* i. (1795) 231.—Europ.
speciosum, Brouss. ex Pers. Syn. ii. 463 = coronarium.
speciosum, Steud. Nom. ed. II. i. 357 = grandiflorum.
Stoliczkai, *C. B. Clarke, Comp. Ind.* 147.—Tibet. occ.
suaveolens, *Aschers. Fl. Brandenb.* 332.—Europ.
subcorymbosum, *Schur, in Verh. Siebenb. Ver. Naturw.* x. (1859) 146.—Transsylv.
sylvaticum, Hoffmgg. & Link, ex Link, Handb. i. 755 = C. Leucanthemum?
sylvaticum, Jauvy, ex DC. Prod. vi. 47 = pallens.
sylvestre, Brot. ex Steud. Nom. ed. I. 193; ed. II. i. 357, nomen.—Lusit.
sylvestre, Willd. Enum. Hort. Berol. Suppl. 60 = C. Leucanthemum.
tanacetifolium, Desf. Cat. Hort. Par. (1821); ex DC. Prod. vi. 66 = grandiflorum.
tanacetifolium, *Willd. Sp. Pl.* iii. 2145.—Oriens.
Tanacetum, Vis. Fl. Dalmat. ii. 89 = Tanacetum vulgare.
tenuifolium, *Kit. in Schult, Oester. Fl.* ed. II. ii. 498.—Hungaria.
tenuifolium, Tenore, ex Steud. Nom. ed. II. i. 357 = C. Achilleae.
tenuissimum, [*Soland. in*] *Russell, Aleppo*, ed. II. ii. 263.—Syria.
Thunbergii, *Harv. in Harv. & Sond. Fl. Cap.* iii. 162.—Afr. austr.
tibeticum, *C. B. Clarke, Comp. Ind.* 147.—Tibet. occ.
tingitanum, *Bernh. in Linnaea*, x. (1835) *Litt.* 74.—Marocc.
tomentosum, *Loisel. Fl. Gall.* 580. t. 18.—Corsica.
trichophyllum, *Boiss. Diagn.* Ser. I. iv. 10.—As. Min.
trichotomum, Walp. Ann. ii. 891 = trichophyllum.
tricolor, Andr. Bot. Rep. t. 109 = carinatum.
trifurcatum, *Desf. Fl. Atlant.* ii. 281.—Afr. bor.
tripartitum, Sweet, Brit. Flow. Gard. Ser. I. t. 193 = indicum.
Turreanum, Vis. Stirp. Dalm. Spec. 19 = cinerariaefolium.
uliginosum. *Pers. Syn.* ii. 460.—Hungar.
umbellatum, *Vell. Fl. Flum.* viii. t. 127.—Bras.
umbrosum, *Willd. Sp. Pl.* iii. 2149.—Graecia.
variabile, Rochel, Pl. Banat. Rar. Ind. Spec. = atratum.
viscosum, *Desf. Cat. Hort. Par.* ed. III. 170.—Reg. Mediterr.
Webbianum, *Ball. in Journ. Linn. Soc.* xvi. (1878) 509.—Marocc.
Webbii, *Masf. in Anal. Soc. Esp. Hist. Nat.* x. (1881) 207.—Ins. Canar.
Welwitschii, Sch. Bip. ex Nym. Consp. 370 = segetum.
Zawadskii, *Herbich, Add. Fl. Galic.* 43.—Galicia.

CHRYSEIS, Cass. in Dict. Sc. Nat. vii. 377; ix. 154 (1817) = **Centaurea**, Linn. (Compos.).
Centaurium, Kostel. Allg. Med. Pharm. Fl. ii. 606 = Cent. Centaurium.
glauca, Cass. in Dict. Sc. Nat. xliv. 39 = Cent. moschata.
moschata, Cass. l. c. = Cent. moschata.
odorata, Cass. l. c. ix. 154 = Cent. moschata.
suaveolens, Cass. l. c. xliv. 39 = Cent. moschata.

CHRYSEIS, Lindl. Bot. Reg. t. 1948 (1835) = **Eschscholzia**, Cham. (Papaverac.).
caespitosa, Torr. & Gray, Fl. N. Am. i. 63 = E. californica.
californica, Hook. & Arn. Bot. Beech. Voy. 319 = E. californica.
compacta, Lindl. Bot. Reg. t. 1948 = E. californica.
crocea, Lindl. l. c. = E. californica.
Douglasii, Hook. & Arn. Bot. Beech. Voy. 320 = E. californica.
hypecoides, Torr. & Gray, Fl. N. Am. i. 64 = E. californica.
tenuifolia, Torr. & Gray, l. c. = E. californica.

CHRYSION, Spach, Hist. Vég. Phan. v. 509 (1836) = **Viola**, Linn.
biflorum, Spach, l. c. 510 = V. biflora.

CHRYSIPHIALA, [Ker-Gawl. in] Bot. Reg. t. 778 (1824) = **Stenomesson**, Herb. (Amaryllid.).
aurantiaca, Schult. f. Syst. vii. 904 = S. aurantiacum.
coccinea, [Ker-Gawl. in] Bot. Reg. sub t. 778 = S. coccineum.
curvidentata, Herb. ex Steud. Nom. ed. II. i. 358 = S. flavum.
flava, [Ker-Gawl. in] Bot. Reg. t. 778 = S. flavum.
incarnata, Schult. f. Syst. vii. 908 = S. incarnatum.
latifolia, [Ker-Gawl. in] Bot. Reg. sub t. 778 = Urceolina latifolia.
pauciflora, Hook. Exot. Fl. t. 132 = S. flavum.
recurvata, [Ker-Gawl. in] Bot. Reg. sub t. 778 = S. recurvatum.
trichroma, Schult. f. Syst. vii. 907 = Pancratium trichromum.
variegata, Schult. f. l. c. = Pancratium variegatum.
viridiflora, Schult. f. l. c. 908 = S. viridiflorum.

CHRYSIS, Rencalm. ex DC. Prod. v. 585 (1836) = **Helianthus**, Linn. (Compos.).

CHRYSITRIX, Linn. Mant. ii. 165 (1771). *CYPERACEAE*, Benth. & Hook. f. iii. 1057.
capensis, *Linn. Mant.* ii. 304; *Linn. f. Gram.* 27. f. 1, 2.—Afr. austr.
junciformis, *Nees, in Linnaea*, x. (1836) 144.—Afr. austr.

CHRYSOBACTRON, Hook. f. Fl. Antarct. i. 72. t. 44 (1845) = **Bulbinella**, Kunth (Liliac.).
Hookeri, Colenso, in Hook. Ic. Pl. t. 817.—N. Zel.
Rossii, Hook. f. Fl. Antarct. i. 72. t. 44–45.—Ins. Auckland.

CHRYSOBALANUS, Linn. Gen. ed. I. 365 (1737). *ROSACEAE*, Benth. & Hook. f. i. 606.
Icaco, Adans. Fam. ii. 305 (1763).
cuspidatus, *Griseb. Fl. Brit. W. Ind.* 711.—Ind. occ.
ellipticus, *Soland. ex Sabine, in Trans. Hort. Soc.* v. (1824) 453.—Afr. trop.
guianensis, Klotzsch, in Schomb. Fauna et Fl. Brit. Guian. 1024 = ellipticus.
Icaco, *Linn. Sp. Pl.* 513.—Afr. et Am. trop.
incanus, Rafin. New Fl. Am. iii. 26 = oblongifolius.
luteus, Sabine, in Trans. Hort. Soc. v. (1824) 453 = C. Icaco?
macrophyllus, *Schott, in Spreng. Syst.* iv. *App.* 407.—Bras.
oblongifolius, *Michx. Fl. Bor. Am.* i. 283.—Am. bor.
orbicularis, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 232 = C. Icaco.
ovalifolius, *Schott, in Spreng. Syst.* iv. *App.* 406.—Bras.
pellocarpus, G. F. W. Mey. Prim. Fl. Esseq. 193 = C. Icaco.
prunifolius, Rafin. New Fl. Am. iii. 26 = oblongifolius.
purpureus, Mill. Gard. Dict. ed. VIII. n. 2 = C. Icaco.
racemosus, *Roxb. Hort. Beng.* [92]; *Fl. Ind.* ii. 506.—Ins. Molucc.
retusus, Rafin. New Fl. Am. iii. 26 = oblongifolius.
subundulatus, *Brign. in Mem. Soc. Ital. Mod.* Ser. II. i. (1862) 71.—Bras.

CHRYSOBAPHUS, Wall. Tent. Fl. Nep. 37. t. 27 (1826) = **Anoectochilus**, Blume (Orchid.).
Roxburghii, Wall. l. c. = A. Roxburghii.

CHRYSOBOTRYA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 18 = **Ribes**, Linn. (Saxifrag.).
intermedia, Spach, l. c. 19 = R. aureum.
Lindleyana, Spach, l. c. 20 = R. aureum.
revoluta, Spach, l. c. 19 = R. aureum.

CHRYSOCALYX, Guill. & Perr. Fl. Seneg. Tent. 157 t. 43 (1832) = **Crotalaria**, Linn. (Legumin.).
ebenoides, Guill. & Perr. l. c. 158. t. 43. f. 1 = Crot. ebenoides.
gracilis, Guill. & Perr. l. c. 159 = Crot. gracilis.
Perrottetii, Guill. & Perr. l. c. 160. t. 43. f. 2 = Crot. Perrottetii.
Petitianus, A. Rich. Tent. Fl. Abyss. i. 153 = Crot. macrocalyx.

CHRYSOCALYX :—
platycalyx, Steud. ex A. Rich. Tent. Fl. Abyss. 160 = Crotalaria platycalyx.
Quartinianus, A. Rich. l. c. 159 = Crot. Dilloniana.
rubiginosa, Guill. & Perr. Fl. Seneg. Tent. 158 = Crot. atrorubens.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 151 = Crot. incana.

CHRYSOCEPHALUM, Walp. in Linnaea, xiv. (1840) 503 = **Helichrysum**, Vaill. (Compos.).
apiculatum, Steetz, in Lehm. Pl. Preiss. i. 474 = H. apiculatum.
asperum, Steetz, l. c. 473 = H. semipapposum.
Behrianum, Sond. in Linnaea, xxv. (1852) 517 = H. Baxteri.
brevicilium, Walp. Rep. vi. 238 = H. semipapposum.
canescens, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 196 = H. apiculatum.
ciliatum, Steetz, in Lehm. Pl. Preiss. i. 474 = H. semipapposum.
flavispinum, Walp. Rep. i. 237, sphalm. = seq.
flavissimum, Steetz, in Lehm. Pl. Preiss. i. 472 = H. apiculatum.
glabratum, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 197 = H. apiculatum.
helichrysoides, Walp. in Linnaea, xiv. (1840) 503 = H. apiculatum.
odorum, Walp. Rep. vi. 238 = H. apiculatum.
pterochaetum, F. Muell. in Linnaea, xxv. (1852) 416 = H. pterochaetum.
semipapposum, Steetz, in Lehm. Pl. Preiss. i. 474 = H. semipapposum.
squarrulosum, Sond. in Linnaea, xxi. (1852) 515 = H. semipapposum.
squarrulosum, Steetz, in Lehm. Pl. Preiss. i. 472 = H. apiculatum.
vitellinum, Sond. & F. Mucll. in Linnaea, xxv. (1852) 514 = H. apiculatum.

CHRYSOCHAMELA, Boiss. Fl. Orient. i. 313 (1867) = **Cochlearia**, Tourn. (Crucifer.).
elliptica, Boiss. l. c. 314.—As. Min.
Noëana, Boiss. l. c.—Armenia.
velutina, Boiss. l. c. 313 = Coch. velutina.

CHRYSOCHLAMYS, Poepp. & Endl. Nov. Gen. et Sp. iii. 13. t. 211 (1842). *GUTTIFERAE*, Benth. & Hook. f. 172.
BERTOLONIA, Spreng. Neue Entdeck. ii. 110. t. 1. f. 1 (1821).
COMMIRHOEA, Miers, in Trans. Linn. Soc. xxi. (1855) 252. t. 26.
POECILOSTEMON, Planch. & Triana, in Ann. Soc. Nat. Sér. IV. xiv. (1860) 256.
TOVOMITOPSIS, Planch. & Triana, l. c. 261 (1860).
clusioides, *Griseb. Cat. Pl. Cub.* 37.—Cuba.
costa-ricana, *Hemsl. Biol. Centr. Am. Bot.* i. 87.—Costa Rica.
dependens, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 259.—Am. trop.
glauca, *Hemsl. Biol. Centr. Am. Bot.* i. 87.—Costa Rica.
Goudotii, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 259.—N. Granat.
laxa, *Planch. & Triana, l. c.* 258.—N. Granat.
membranacea, *Planch. & Triana, l. c.* 260.—N. Granat.
multiflora, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 13. t. 211.—Peruv.
myrcioides, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 260.—N. Granat.
nicaraguensis, *Hemsl. Biol. Centr. Am. Bot.* i. 87.—Nicaragua.
pachypoda, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 257.—Peruv.
Pavonii, *Planch. & Triana, l. c.* 258.—Peruv.
psychotriaefolia, *Hemsl. Biol. Centr. Am. Bot.* i. 87.—Costa Rica.

CHRYSOCOMA, Linn. Gen. ed. I. 248 (1737), partim. *COMPOSITAE*, Benth. & Hook. f. ii. 286.
acaulis, Walt. Fl. Carol. 196 = Vernonia oligophylla.
adpressa, Vell. Fl. Flum. 332; viii. t. 35 = Vernonia monticola.
Aitoni, *Spreng. Fl. Hal. Mant.* 50.—Hab.?
alata, *Vell. Fl. Flum.* 335; viii. t. 48 (Eupatorii sp.?).—Bras.
albida, *Vell. l. c.* 333; viii. t. 42.—Bras.

CHRYSOCOMA :—
alterna, *Vell. Fl. Flum.* 333; viii. t. 40 (Vernoniae sp.?).—Bras.
amara, Schum. & Thonn. Beskr. Guin. Pl. 383 = Vernonia senegalensis.
angustifolia, Steud. Nom. ed. II. i. 358 = Vernonia angustifolia.
aphylla, Vell. Fl. Flum. 324; viii. t. 1 = Baccharis aphylla.
arborea, Vell. l. c. 326; viii. t. 11 = Vernonia Tweedieana.
artemisifolia, Poepp. ex Steud. Nom. ed. II. i. 358 = Eupatorium coronopifolium.
aspera, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. 239 = Inula germanica.
aurea, Boj. ex DC. Prod. vi. 322 = Nidorella Chrysocoma.
aurea, Salisb. Prod. 118 = C. Coma-aurea.
biflora, Linn. Sp. Pl. 841 = Aster dracunculoides.
camphorata, Rob. & Cass. in Dict. Sc. Nat. xxxiv. 35 = Jasonia glutinosa.
candicans, Delile, Fl. Egypte, 271. t. 46 (= *Varthemia candicans*).—Aegypt.
capillacea, Michx. Fl. Bor. Am. ii. 101 = Eupatorium foeniculaceum.
cernua, Linn. Sp. Pl. 840 = C. Coma-aurea.
cernua, Thunb. Prod. Pl. Cap. 142, partim = tenuifolia.
ciliaris, Link, Enum. Hort. Berol. ii. 311 = Aster Ecklonis.
ciliaris, Willd. Sp. Pl. iii. 1789 = ciliata.
ciliata, *Linn. Sp. Pl.* 841.—Afr. austr.
ciliata, E. Mey. ex DC. Prod. v. 354 = oblongifolia.
cinerea, Labill. Nov. Holl. Pl. ii. 39. t. 182 = Helichrysum cinereum.
Coma-aurea, *Linn. Sp. Pl.* 840.—Afr. austr.
coronopifolia, Michx. Fl. Bor. Am. ii. 102 = Eupatorium coronopifolium.
crassa, *Vell. Fl. Flum.* 326; viii. t. 8.—Bras.
cruciata, *Vell. l. c.* 326; viii. t. 10 (Baccharidis sp.?).—Bras.
cuneifolia, *Jacq. Coll.* iii. 280.—Reg. Magell.
cymosa, Vell. Fl. Flum. 327; viii. t. 16 = Vernonia missionis.
decurrens, *DC. Prod.* v. 354.—Afr. austr.
decurrens, Harv. in Harv. & Sond. Fl. Cap. iii. 95 = Heteromma decurrens.
decussata, Vell. Fl. Flum. 326; viii. t. 9 = Baccharis camporum.
dentata, Vell. l. c. 334; viii. t. 47 = Baccharis orgyalis.
denticulata, Jacq. Hort. Schoenb. iii. 62 = Chrysanthemum flosculosum.
dichotoma, Jacq. Ic. Pl. Rar. i. 17. t. 171 = Allagopappus dichotomus.
divaricata, Fisch. Hort. Gorenk. ed. II. (1812) 37 = Aster divaricatus.
dracunculoides, Lam. Encyc. ii. 192 = Aster dracunculoides.
dracunculoides, Pursh, Fl. Am. Sept. ii. 517 = Bigelowia dracunculoides.
ferruginea, Spreng. Syst. iii. 424 = Helichrysum ferrugineum.
florida, *Vell. Fl. Flum.* 334; viii. t. 44.—Bras.
foetida, Lam. Encyc. ii. 193 = Nidorella foliosa.
fruticosa, J. F. Gmel. Syst. 1203 = cuneifolia.
fruticosa, Steud. Nom. ed. II. i. 358 = Vernonia rigida.
gigantea, Walt. Fl. Carol. 196 = Vernonia altissima.
gnaphaloides, *Burm. f. Fl. Cap. Prod.* 23.—Afr. austr.
graminifolia, Boeber, in Pall. N. Nord. Beitr. vi. 261 = Aster Linosyris.
graminifolia, Linn. Sp. Pl. 841 = Solidago lanceolata.
graminifolia, Walt. Fl. Carol. 196 = Vernonia angustifolia.
grandis, *Lepech. It.* iii. 38.—Sibir. ural.
graveolens, Nutt. Gen. Am. ii. 136 = Bigelowia dracunculoides.
graveolens, Torr. in Ann. Lyc. N. York, ii. (1828) 211 = Bigelowia pluriflora.
herbacea, Vell. Fl. Flum. 330; viii. t. 29 = Vernonia obovata.
heterophylla, Rafin. Fl. Ludov. 64 (= Baccharidis sp.?).—Am. bor.
hirsuta, *Vell. Fl. Flum.* 333; viii. t. 41.—Bras.
horizontalis, Vell. Fl. Flum. 330; viii. t. 28 = Vernonia obscura.
humilis, DC. Prod. vii. 280 = Aster angustus.
imberbis, *Vell. Fl. Flum.* 331; viii. t. 34.—Bras.
incana, Burm. f. Fl. Cap. Prod. 23 = Pteronia incana.

CHRYSOCOMA :—
incana, Léveillé, in Demid. Voy. Russ. Mér. ii. 237; iv. 496 (= *Linosyris villosa*).—Rossia.
indivisa, Vell. Fl. Flum. 326; viii. t. 12 = Baccharis brachylaenoides.
integerrima, *Vell. l. c.* 328; viii. t. 19.—Bras.
juncea, *Desf. Tabl.* ed. II. 115, nomen.—Hab.?
lappulina, Boj. ex DC. Prod. v. 323 = Nidorella mucronata.
lateralis, Vell. Fl. Flum. 329; viii. t. 21 = Baccharis cassinaefolia.
liburnica, Spreng. Syst. iii. 425 = Aster Linosyris.
linearis, *Vell. Fl. Flum.* 330; viii. t. 27.—Bras.
linifolia, Steud. Nom. ed. II. i. 358 = Aster subnlatus?
Linosyris, Linn. Sp. Pl. 841 = Aster Linosyris.
longifolia, *DC. Prod.* v. 354.—Afr. austr.
maculata, Vell. Fl. Flum. 325; viii. t. 6 = Eupatorium conyzoides.
maritima, Vell. l. c. 329; viii. t. 22 = Baccharis orgyalis.
medioserrata, *Vell. l. c.* 327; viii. t. 15.—Bras.
megapotamica, Spreng. Syst. iii. 424 = Eupatorinm ericoides.
microcephala, DC. Prod. v. 353 = tenuifolia.
microphylla, Thunb. Prod. Pl. Cap. 142 = tenuifolia.
monantha, *Vell. Fl. Flum.* 331; viii. t. 33.—Bras.
montana, Vahl. Symb. Bot. i. 70 (= *Varthemia montana*).—Arabia.
mucronata, Forsk. Fl. Aegypt. Arab. 147 = Iphiona juniperifolia.
nauseosa, Pursh, Fl. Am. Sept. ii. 517 = Bigelowia dracunculoides.
nivea, *Willd. Sp. Pl.* iii. 1790.—Afr. austr.
noveboracensis, Desf. Tabl. ed. I. 98 = Vernonia noveboracensis.
nuda, Vell. Fl. Flum. 335; viii. t. 50 = Baccharis aphylla.
nudata, Michx. Fl. Bor. Am. ii. 101 = Bigelowia nudata.
nupera, S. F. Gray, Nat. Arr. Brit. Pl. ii. 467 = Aster Linosyris.
oblongifolia, *DC. Prod.* 354.—Afr. austr.
obscura, *Vell. Fl. Flum.* 330; viii. t. 30.—Bras.
odoratissima, Raeusch. Nom. ed. III. 235 = Trilisia odoratissima.
oligophylla, Vell. Fl. Flum. 324; viii. t. 2. = Vernonia Cephalotes.
opposita, *Vell. l. c.* 334; viii. t. 46.—Bras.
oppositifolia, Linn. Pl. Rar. Afr. 18 = Pteronia divaricata.
oppositifolia, *Vell. Fl. Flum.* 328; viii. t. 18.—Bras.
ovata, *Forsk. Fl. Aegypt. Arab.* 147.—Arabia.
palustris, Savi, Pugill. 12; ex Bertol. Fl. Ital. ix. 91 = Aster Linosyris.
paniculata, J. F. Gmel. Syst. 1204 = Trilisia paniculata.
paniculata, Vell. Fl. Flum. 327; viii. t. 14 = Vernonia geminata.
patula, Berg. Desc. Pl. Cap. 234 = C. Coma-aurea.
pauciflora, Vell. Fl. Flum. 325; viii. t. 5 = Ageratum melissaefolium.
peduncularis, *DC. Prod.* v. 353.—Afr. austr.
pedunculata, Vell. Fl. Flum. 332; viii. t. 37 = Vernonia ammophila.
phosphorica, Vell. l. c. 325; viii. t. 4 = Vernonia polyanthes.
pilosa, J. F. Gmel. Syst. 1203 = Liatris pycnostachya.
pilosa, *Vell. l. c.* 328; viii. t. 17.—Bras.
pinifolia, *Burm. f. Fl. Cap. Prod.* 23.—Afr. austr.
pinifolia, Spreng. Syst. iii. 424 (= *Ozothamnus pinifolius*).—N. Zel.
pinnatifida, *DC. Prod.* v. 354.—Afr. austr.
praeacoa, Steud. Nom. ed. II. i. 358 = C. Preacaa.
praealta, Desf. Tabl. ed. I. 98 = Vernonia altissima?
Preacaa, *Vell. Fl. Flum.* 333; viii. t. 39.—Bras.
pterocaula, *DC. Prod.* v. 354.—Afr. austr.
pumila, Vell. Fl. Flum. 331; viii. t. 32 = Vernonia grandiflora.
punctata, Steud. Nom. ed. II. i. 358 = seq.
punctulata, Vell. Fl. Flum. 335; viii. t. 49 = Eupatorium laevigatum.
purpurascens, *Vell. l. c.* 331; viii. t. 31.—Bras.
purpurea, *Forst. f. Prod.* 54.—N. Hebrid.
purpurea, Rafin. Fl. Ludov. 64 (= Liatridis sp.).—Am. bor.
purpurea, Vell. l. c. 334; viii. t. 45 = Baccharis cassinaefolia.
quadrangularis, Vell. Fl. Flum. 329; viii. t. 25 = Piptocarpha quadrangularis.

CHRYSOCOMA:—
radiata, *Fisch. ex Steud. Nom.* ed. II. i. 359.—Hab.?
ramosa, *Raeusch. Nom.* ed. III. 235, nomen.—Carolin.
repanda, Vell. Fl. Flum. 327; viii. t. 13 = Vernonia scorpioides.
reticulata, Labill. Nov. Holl. Pl. ii. 40. t. 183 = Helichrysum reticulatum.
rosmarinifolia, Sprcng. Syst. iii. 424 = Helichrysum rosmarinifolium.
sancta, Vell. Fl. Flum. 324; viii. t. 3 = Baccharis Lundii.
saxatilis, DC. Fl. Fr. Suppl. 468 = Jasonia saxatilis.
scabra, Ledcb. in Mém. Acad. Pétersb. v. (1812) 563 = Inula germanica.
scabra, Linn. Sp. Pl. ed. II. 1177 = Felicia reflexa.
scoparia, Linn. Pl. Jam. Pug. 21; Amoen. Acad. v. 381 = Baccharis scoparia.
sericea, Linn. f. Suppl. 360 (= *Schizogyne sericea*).—Ins. Canar.
sernua, Hill, Veg. Syst. iii. 128, sphalm. = C. Comaaurca.
serrata, *Vell. Fl. Flum.* 329; viii. t. 23.—Bras.
sessilis, Vell. l. c. 328; viii. t. 20 = Vernonia maerophylla.
singularis, Vell. l. e. 325; viii. t. 7 = Vernonia nitidula et Baccharis cassinefolia.
spathulata, Forsk. Fl. Aegypt. Arab. 147 = Vernonia atriplicifolia.
sphacelata, Moench, Meth. 577 = Senecio paniculatus.
spicata, Forsk. Fl. Aegypt. Arab. p. lxxiii = Ifloga Fontanesii.
spinosa, Delile, Fl. Aegypt. 128. t. 46 = Iphiona juniperifolia.
squamata, Labill. Nov. Holl. Pl. ii. 40. t. 184 = Leptorhynchus squamatus.
tatarica, *Less. in Linnaea*, ix. (1834) 186.—Reg. . Casp.
tenuifolia, *Berg. Desc. Pl. Cap.* 235.—Afr. austr.
tenuifolia, Salisb. Prod. 188 = Aster Linosyris.
tomentosa, *Vell. Fl. Flum.* 332; viii. t. 38.—Bras.
tomentosa, *Linn. Syst.* ed. XII. 539.—Afr. austr.
tomentosa, Walt. Fl. Carol. 196 = Vernonia noveboracensis.
tridentata, *DC. Prod.* v. 354.—Afr. austr.
trinervis, *Vell. Fl. Flum.* 333; viii. t. 43.—Bras.
truncifolia, J. Shaw, in Journ. Linn. Soc. xiv. (1874) 206, sphalm. = tenuifolia.
tuberosa, *Vell. Fl. Flum.* 332; viii. t. 36.—Bras.
undulata, Thunb. Prod. Pl. Cap. 142 = Nidorella amplexicaulis.
uniflora, *Racusch. Nom.* ed. III. 235, nomen.—Am. bor.
uniflora, Spreng. Nov. Prov. (1818) 11 (= *Varthemia candicans*).—Aegypt.
verticalis, Lag. Gen. et Sp. Nov. 25 = Jasonia saxatilis.
verticillata, Vell. Fl. Flum. 330; viii. t. 26 = Piptocarpha oxyphylla.
villosa, Linn. Sp. Pl. 841 (= *Linosyris villosa*).—Europ.; Caue.; Sibir.
violacea, Schum. & Thonn. Beskr. Guin. Pl. 384 = Vernonia cinerea.
virgata, Nutt. Gen. Am. ii. 137 = Bigelowia nudata.
virginiana, Hort. ex DC. Prod. v. 341 = Solidago lanceolata.
volubilis, *Vell. Fl. Flum.* 329; viii. t. 24.—Bras.
vulgaris, *Gueldenst. ex Ledeb. Fl. Ross.* ii. 495 = Aster Linosyris.

CHRYSOCOPTIS, Nutt. in Journ. Acad. Philad. vii. (1834) 8. t. 1 = **Coptis**, Salisb. (Ranunculac.).
asplenifolia, Nutt. l. c. 9, in obs. = Coptis asplenifolia.
occidentalis, Nutt. l. e. 8 = Coptis occidentalis.

CHRYSOCORYNE, Endl. in Bot. Zeit. i. (1843) 457 = **Angianthus**, Wendl. (Compos.).
angianthoides, F. Muell. in Linnaea, xxv. (1852) 404, 488 = A. pusillus.
Drummondii, A. Gray, in Hook. Kew Journ. iii. (1851) 152 = A. tenellus.
Huegelii, A. Gray, l. e. 151 = A. tenellus.
myosuroides, A. Gray, l. e. 152 = A. myosuroides.
pusilla, Endl. in Bot. Zeit. i. (1843) 458 = A. pusillus.
tenella, F. Muell. in Trans. Viet. Inst. (1855) 130 = A. tenellus.
uniflora, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 188 = A. myosuroides.

CHRYSOCYATHUS, Falc. in Proc. Linn. Soc. i. (1839) 17 = **Calathodes**, Hook. f. & Thoms. (Ranunculac.).

CHRYSOCYCNIS, Linden & Reichb. f. in Bonplandia, ii. (1854) 280. *ORCHIDEAE*, Benth. & Hook. f. iii. 553.
Schlimii, *Linden & Reichb. f. l. c.*—N. Granat.

CHRYSODENDRON, Vaill. ex Meissn. in DC. Prod. xiv. 230 (1856–57) = **Protea**, Linn.

CHRYSODISCUS, Steetz, in Lehm. Pl. Preiss. i. 460 (1845) = **Athrixia**, Ker-Gawl. (Compos.).
niveus, Steetz, l. c. = A. stricta.

CHRYSOGLOSSUM, Blume, Bijdr. 337. t. 7 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 507.
DIGLYPHIS, Blume, Fl. Jav. Praef. p. vii. (1828).
DIGLYPHOSA, Blume, Bijdr. 336. t. 60 (1825).
latifolium, *Benth. & Hook. f. Gen.* iii. 508.—Java.
ornatum, *Blume, Bijdr.* 338.—Java.
vesicatum, *Reichb. f. in Seem. Fl. Vit.* 304.—Ins. Fiji.
villosum, *Blume, Bijdr.* 338.—Java.

CHRYSOGONUM, Linn. Hort. Cliff. 424 (1737); Coroll. Gen. 17 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 350.
CARGILLA, Adans. Fam. ii. 130 (1763).
DIOTOSTEPHUS, Cass. in Dict. Sc. Nat. xlviii. 543 (1827).
MOONIA, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 348.
PENTALEPIS, F. Muell. in Trans. Bot. Soc. Edinb. vii. (1863) 496.
Arnottianum, *C. B. Clarke, Compos. Ind.* 132.—Ind. or.
dichotomum, *Vahl, in West, Bidr. Stc. Croix*, 303; *DC. Prod.* v. 510.—Ins. S. Crucis.
Diotostephus, DC. l. e. = virginianum.
ecliptoides, *Benth. & Hook. f. ex F. Muell. Census*, 83.—Austral.
hetcrophyllum, *C. B. Clarke, Comp. Ind.* 132.—Ind or.
peruvianum, Linn. Sp. Pl. 920 = Zinnia multiflora.
procumbens, *Benth. & Hook. f. cx F. Muell. Census*, 83.—Austral.
trichodesmoides, *Benth. & Hook. f. l. c.*—Austral.
virginianum, *Linn. Sp. Pl.* 920.—Am. bor.

CHRYSOLIGA, Willd. ex DC. Prod. iii. 89, in syn. (1828) = **Nesaea**, Comm. (Lythrar.).

CHRYSOLINUM, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 349 = **Linum**, Tourn.
gallicum, Fourr. l. c. = L. gallicum.
strictum, Fourr. l. e. = L. strictum.

CHRYSOLYGA, Willd. ex Steud. Nom. ed. II. i. 359 (1840) = **Nesaea**, Comm. (Lythrar.).
salicifolia, Willd. ex Steud. l. c. = N. salicifolia.

CHRYSOMA, Nutt. in Journ. Acad. Philad. vii. (1834) 67 = **Solidago**, Vaill. (Compos.).
pumila, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 325 = S. pumila.
solidaginoides, Nutt. in Journ. Acad. Philad. vii. (1834) 67 = S. pauciflosculosa.
uniligulata, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 325 = S. negleeta.

CHRYSOMALLUM, Thou. Gen. Nov. Madag. 8 (1806) = **Vitex**, Linn. (Verbenae.).
madagascariense, Thou. ex Steud. Nom. ed. I. 194 = V. Chrysomallum.

CHRYSOMELEA, Tausch, Hort. Canal. fasc. i.—Cf. DC. Prod. v. 569 (1836) = **Coreopsis**, Linn. (Compos.).
auriculata, Tausch, l. c.; ex DC. l. c. 571 = Cor. pubescens.
lanccolata, Tausch, l. e.; ex DC. l. e. 570 = Cor. lanceolata.

CHRYSOMELON, Forst. ex A. Gray, Bot. U. St. Expl. Exped. i. 374, in syn. (1854) = **Spondias**, Linn. (Anacard.).
pomiferum, Forst, ex A. Gray, l. c. = S. dulcis.

CHRYSONIAS, Benth. ex Steud. Nom. ed. II. i. 259 (1840) err. typ. = **Chrysoscias**, E. Mey.

CHRYSOPHANIA, Kunth, ex Less. Syn. Comp. 224 (1832) = **Zaluzania**, Pers. (Compos.).
fastigiata, Kunth, l. c. = Z. angusta.

CHRYSOPHORA, Cham. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 94, 95. 147 = **Oxymeris**, DC. (Melastomae.).
cinnamomifolia, Cham. ex Triana, l. e. 147 = Ossaea cinnamomifolia.
cordigera, Cham. ex Triana, l. e. 94 = Oxym. cordigera.
hirsutior, Cham. ex Triana, l. e. 95 = Oxym. deflexa.
secundifolia, Cham. ex Triana, l. c. = Oxym. secundiflora.

CHRYSOPHTHALMUM, Phil. in Linnaea, xxix. (1857) 9 = **Grindelia**, Willd. (Compos.).
andinum, Phil. l. c. = G. andina.

CHYSOPHTHALMUM, Sch. Bip. in Walp. Rep. ii. 955 (1843) (Chrysophtalmum). *COMPOSITAE*, Benth. & Hook. f. ii. 341.
dichotomum, *Boiss. & Heldr. Diagn.* Sér. I. xi. 5.—As. Min.
montanum, Boiss. l. e. 4 = sternutatorium.
sternutatorium, *Sch. Bip. in Walp. Rep.* ii. 955.—As. Min.

CHRYSOPHYLLUM, Linn. Gen. ed. I. 361 (1737). *SAPOTACEAE*, Benth. & Hook. f. ii. 653.
CAINITO, Plum. ex Adans. Fam. ii. 166 (1763); Tuss. Fl. Antill. iii. 41. t. 9 (1824).
DACTIMALA, Rafin. Sylva Tellur. 153 (1838).
GUERSENTIA, Rafin. l. c. (1838).
NIEMEYERA, F. Muell. Fragm. vii. 114 (1869–71).
NYCTERISITION, Ruiz & Pav. Prod. 30. t. 5 (1794).
SCLEROXYLON, Bertol. in Mem. Acc. Sc. Bolog. viii. (1857) 237 (Misc. Bot. xiii. 15).
acuminatum, Boj. Hort. Maurit. 197 = Sideroxylon ferrugineum.
acuminatum, Lam. Tabl. Encyc. ii. 44 = monopyrenum.
acuminatum, Poir. Encyc. Suppl. ii. 15 = Sideroxylon guyanense.
acuminatum, Roxb. Hort. Beng. 17; Fl. Ind. i. 599 = Roxburghii.
acuminatum, Willd. ex Roem. & Schult. Syst. iv. 813 = Humboldtianum.
africanum, *A. DC. in DC. Prod.* viii. 163.—Afr. trop.
albidum, *G. Don, Gen. Syst.* iv. 32.—Afr. trop.
alnifolium, *Baker, in Oliver, Fl. Trop. Afr.* iii. 499.—Afr. trop.
amplifolium, *A. DC. in DC. Prod.* viii. 159.—Bras.
angustifolium, *Lam. Tabl. Encyc.* ii. 44.—Ins. S. Doming.
aquaticum, H. B. & K. Nov. Gen. et Sp. iii. 236 = C. Cainito.
argenteum, *Jacq. Enum. Pl. Carib.* 15.—Ins. Martinic.
auratum, *Miq. in Mart. Fl. Bras.* vii. 97.—Bras.
bancanum, *Miq. Fl. Ind. Bat. Suppl.* 579.—Ins. Banca.
Barbasco, Loefl. ex DC. Prod. viii. 149 = Jaequinia armillaris.
bicolor, *Poir. Encyc. Suppl.* ii. 15.—Ins. Porto Rico.
Bonplandii, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 94 = C. Cainito.
brasiliense, *A. DC. in DC. Prod.* viii. 156.—Bras.
bumelioides, *Mart. in Flora*, xxii. (1839) i. *Beibl.* 1.—Bras.
Buranhem, Riedel, in Taunay, Man. Agric. Bras. ed. II. 318 = Lucuma glycyphloea.
caeruleum, Jacq. Stirp. Select. Am. 52 = C. Cainito.
Cainito, *Linn. Sp. Pl.* 192.—Ind. occ.
carolinense, Jaeq. Obs. Bot. iii. 3 = Bumelia tenax.
cayennense, *A. DC. in DC. Prod.* viii. 160.—Guiana.
cuneifolium, *A. DC. l. c.*—Guiana.
decandrum, *Montr. in Mém. Acad. Lyon*, x. (1860) 228.—N. Caled.
dubium, *Steud. Nom.* ed. II. i. 359.—Ins. S. Doming.
dubium, *Panch. & Sebert, Not. Bois. Nouv. Caléd.* 195.—N. Caled.
ebenaceum, *Mart. in Flora*, xxi. (1838) ii. *Beibl.* 96.—Bras.
emarginatum, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 1087.—Guiana.
ferrugineum, Gaertn. Fruct. iii. 122 = monopyrenum.
ferrugineum, *Steud. Nom.* ed. II. i. 359.—Peruv.
firmum, *Miq. Fl. Ind. Bat. Suppl.* 579.—Ins. Banca.
flexuosum, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 98.—Bras.
fuscum, *Fisch. ex Steud. Nom.* ed. II. i. 359.—Ind. or.

CHRYSOPHYLLUM:—
Gardneri, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 102.—Bras.
glabrum, *Jacq. Enum. Pl. Carib.* 15.—Ins. Martinic.
glabrum, Juss. ex Lam. Encyc. i. 245 = Bumelia tenax.
glycyphloeum, Casar. in Atti 3zia Riun. Sc. Ital. Firenze (1841) 514; Dec. Stirp. Bras. 12 = Lucuma glycyphloea.
granatense, *Spreng. Syst.* i. 667.—N. Granat.
grandifolium, *Steud. Nom.* ed. II. i. 359.—Ins. Philipp.
guyanense, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 98 = sericeum.
Guilleminianum, A. DC. in DC. Prod. viii. 161 = ebenaceum.
Hostmannianum, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 98 = sericeum.
Humboldtianum, *Roem. & Schult. Syst.* iv. 813.—Am. austr.
Inophyllum, Mart. ex Miq. in Mart. Fl. Bras. vii. 105 = Lucuma Inophyllum.
inornatum, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 96.—Bras.
jamaicense, Jacq. Stirp. Select. Am. 52 = C. Cainito.
januariense, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 206.—Bras.
javanicum, *Steud. Nom.* ed. II. i. 359.—Java.
lanceolatum, Casar. Stirp. Bras. Dec. 64 = inornatum.
lanceolatum, *A. DC. in DC. Prod.* viii. 162.—Java.
leptocarpum, Spruce, ex Miq. in Mart. Fl. Bras. vii. 106 = Lucuma leptocarpa.
longifolium, *Mart. & Eichl. ex Miq. l. c.* 97.—Bras.
lucumifolium, *Griseb. in Goett. Abh.* xxiv. (1879) 223.—Reg. Argent.
ludovicianum, *Rafin. Fl. Ludov.* 53.—Louisiana.
Macoucou, *Aubl. Pl. Gui.* i. 233. t. 92.—Guiana.
macrophyllum, Desf. Cat. Hort. Par. ed. III. 132 = grandifolium.
macrophyllum, Gaertn. f. Fruct. iii. 122. t. 202. f. 1 = dubium.
macrophyllum, Lam. Tabl. Encyc. ii. 44 = Lucuma Rivicoa.
macrophyllum, Mart. in Flora, xxi. (1838) II. Beibl. 95 = brasiliense.
macrophyllum, Sabine, in Trans. Hort. Soc. v. (1824) 458 = africanum.
magalis-montana, *Sond. in Linnaea*, xxiii. (1850) 72.—Afr. austr.
maliforme, Linn. Syst. ed. X. 937 = C. Cainito.
Manglillo, Lam. Tabl. Encyc. ii. 45 = Myrsine Manglilla.
Martianum, A. DC. in DC. Prod. viii. 161 = ebenaceum.
maytenoides, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 1.—Bras.
melanophleum, Lam. Tabl. Encyc. ii. 45 = Myrsine mitis.
Michino, *H. B. & K. Nov. Gen. et Sp.* iii. 236.—N. Granat.
microcarpum, *Sw. Prod. Veg. Ind. Occ.* 49.—Ins. S. Doming.
microphyllum, A. DC. in DC. Prod. viii. 158 = monopyrenum.
microphyllum, Jacq. Stirp. Select. Am. 53. t. 37. f. 2 = C. Cainito.
Millerianum, Lam. Tabl. Encyc. ii. 45 = Myrsine mitis.
molle, Wall. Cat. n. 4161 = Payena Maingayi.
monopyrenum, Spreng. Syst. i. 666 = C. Cainito.
monopyrenum, *Sw. Prod. Veg. Ind. Occ.* 49.—Ind. occ.
myrsinodendron, F. Muell. Fragm. vi. 178 = Sideroxylon Brownii.
myrtifolium, Mart. in Flora, xxii. (1839) I. Beibl. 1 = maytenoides.
natalense, *Sond. in Linnaea*, xxiii. (1850) 72.—Afr. austr.
nitidissimum, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 98 = auratum.
nitidum, *G. F. W. Mey. Prim. Fl. Esseq.* 116.—Guiana.
obovatum, *Sabine, in Trans. Hort. Soc.* v. (1824) 458.—Afr. trop.
obovatum, Wall. ex A. DC. in DC. Prod. viii. 178 = Sideroxylon ferrugineum.
obtusatum, Willd. ex Roem. & Schult. Syst. iv. 813 = C. Cainito.
oleaefolium, *Spruce, ex Miq. in Mart. Fl. Bras.* vii. 101.—Bras.
Olfersii, Klotzsch, ex Miq. l. c. 91 = splendens.
oliviforme, Lam. Encyc. i. 552 = monopyrenum.
oliviforme, Linn. Syst. ed. X. 937 = C. Cainito.

CHRYSOPHYLLUM:—
Ottonis, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 94 = C. Cainito.
parviflorum, *Casar. Stirp. Bras. Dec.* 63.—Bras.
pauciflorum, Lam. Tabl. Encyc. ii. 44 = glabrum.
pentagonum, *Hance, in Journ. Bot.* xx. (1882) 78.—China.
Persicastrum, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 208.—Bras.
pomiforme, *Bert. ex Spreng. Syst.* i. 667.—Jamaic.
Prieurei, *A. DC. in DC. Prod.* viii. 161.—Guiana.
pruniferum, *F. Muell. Fragm.* vi. 26.—Austral.
prunifolium, *Baker, in Oliver, Fl. Trop. Afr.* iii. 499.—Afr. trop.
psilophyllum, *A. DC. in DC. Prod.* viii. 160.—Guiana.
pulcherrimum, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 92.—Bras.
pyriforme, Willd. Sp. Pl. i. 1084 = C. Macoucou.
ramiflorum, A. DC. in DC. Prod. viii. 158 = Ecclinusa ramiflora.
revolutum, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 104.—Bras.
rhodoneurum, *Hassk. in Flora*, xxxviii. (1855) 579.—Java.
Richardii, Klotzsch, ex Miq. in Mart. Fl. Bras. vii. 97 = auratum.
Roxburghii, *G. Don, Gen. Syst.* iv. 33.—As. trop.
rufum, *Mart. in Flora*, xxi. (1838) II. *Beibl.* 94.—Bras.
rugosum, Sw. Prod. Veg. Ind. Occ. 49 = Sideroxylon rugosum.
Schomburgkianum, *A. DC. in DC. Prod.* viii. 157.—Guiana.
Sebertii, *Panch. in Panch. & Sebert, Not. Bois Nouv. Caléd.* 194.—N. Caled.
sericeum, *A. DC. in DC. Prod.* viii. 158.—Guiana.
sericeum, Salisb. Prod. 138 = C. Cainito.
sessiliflorum, *Poir. Encyc. Suppl.* ii. 16.—Guiana.
sessilifolium, *Panch. & Sebert, Not. Bois Nouv. Caléd.* 195.—N. Caled.
sparsiflorum, *Klotzsch, ex Miq. in Mart. Fl. Bras.* vii. 90.—Guiana.
splendens, *Spreng. Syst.* i. 666.—Bras.; Peruv.
subnudum, *Baker, in Oliver, Fl. Trop. Afr.* iii. 499.—Afr. trop.
sumatranum, Miq. Fl. Ind. Bat. Suppl. 579 = Roxburghii.
sundaicum, *Miq. l. c.* 578.—Ins. Sunda.
viride, *Mart. & Eichl. ex Miq. in Mart. Fl. Bras.* vii. 102.—Bras.
Wakere, *Panch. & Sebert, Not. Bois Nouv. Caléd.* 193.—N. Caled.

CHRYSOPIA, Noronha, ex Thou. Gen. Nov. Madag. 14 (1806) = **Symphonia**, Linn. (Guttif.).
fasciculata, Thou. l. c. n. 48 = S. fasciculata.
gymnoclada, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 291.—Madag.
macrophylla, Cambess. Mém. Tern. et Gutt. 55. t. 4.—Madag.
microphylla, Cambess. & Boj. ex Cambess. in Mém. Mus. Par. xvi. (1828) 423. t. 4.—Madag.
urophylla, Decne. ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 291.—Madag.
verrucosa, Hilsenb. & Boj. ex Planch. & Triana, l. c. 290.—Madag.

CHRYSOPOGON, Trin. Fund. Agrost. 187 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1135.
CENTROPHORUM, Trin. l. c. 106. t. 5 (1820).
PORANTHERA, Rafin. in Ser. Bull. Bot. i. (1830) 221.
RAPHIS, Beauv. Agrost. 120 (1812).
RHAPHIS, Lour. Fl. Cochinch. 552 (1790).
TRIANTHIUM, Desv. Opusc. 69, obs. (1831).
aciculatus, *Trin. Fund. Agrost.* 188.—Reg. trop.
alternans, Trin. ex Steud. Nom. ed. II. i. 359 = violascens?
argutus, *Trin. ex Steud. l. c.* 360.—Austral.
Arnottianus, *Nees, ex Steud. Syn. Pl. Gram.* 396.—Ind. or.
avenaceus, *Benth. in Journ. Linn. Soc.* xix. (1881) 73.—Am. calid.
ciliolatus, Boiss. Fl. Orient. v. 458 = serrulatus.
filiformis, Voigt, Hort. Suburb. Calc. 704 = Dimeria ornithopoda.
elongatus, *Benth. Fl. Austral.* vii. 538.—Austral.

CHRYSOPOGON:—
Esenbeckii, Arn. ex Steud. Syn. Pl. Gram. 395 = serrulatus.
Francavillanus, Hemsl. Biol. Centr. Am. Bot. iii. 530 = avenaceus.
fuscus, Trin. ex Steud. Nom. ed. II. i. 360 = Andropogon serratus.
glabratus, Trin. in Mém. Acad. Pétersb. Sér. VI. ii. (1833) 318 = C. Gryllus.
Gryllus, *Trin. Fund. Agrost.* 188.—Reg. trop. et subtrop.; Europ. austr.; Afr.
increscens, Nees, ex Steud. Syn. Pl. Gram. 396 = serrulatus.
Minarum, Benth. in Journ. Linn. Soc. xix. (1881) 73 = stipoides.
montanus, *Trin. in Spreng. Neue Entdeck.* ii. 93.—Ind. or.
nutans, *Benth. in Journ. Linn. Soc.* xix. (1881) 73.—Am. calid.
pallidus, *Trin. ex Steud. Nom.* ed. II. i. 360.—Austral.
parviflorus, Benth. in Hook. Lond. Journ. Bot. ii. (1843) 411 = violascens.
pauciflorus, Benth. ex Vasey, Gram. U. St. 20 = Sorghum pauciflorum.
pictus, *Hance, in Ann. Sc. Nat.* Sér. V. v. (1866) 252.—As. trop.
quinqueplumis, *A. Rich. Tent. Fl. Abyss.* ii. 450.—Abyssin.
secundum, Benth. ex Vasey, Gram. U. St. 20 = Sorghum secundum.
serrulatus, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 318.—Ind. or.; Afr. trop.
stipoides, Benth. in Journ. Linn. Soc. xix. (1881) 73 = avenaceus.
stipoides, *Trin. in Bull. Sc. Acad. Pétersb.* i. (1836) 71.—Bras.
strictus, Nees, in Hook. Kew Journ. ii. (1850) 95 = Andropogon intermedius.
subtilis, *Miq. Fl. Ind. Bat.* iii. 492.—Java.
subulatus, Trin. ex Steud. Nom. ed. II. i. 360 = aciculatus.
tener, *Nees, ex Steud. Syn. Pl. Gram.* 397.—'Andor.'
trivialis, Arn. & Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 171 = aciculatus.
verticillatus, *Trin. ex Steud. Nom.* ed. II. i. 360.—Ind. or.
violascens, *Trin. in Mém. Acad. Pétersb.* Sér. VI. ii. (1833) 319.—As. trop.; Abyssin.; Austral.
Wightianus, *Thw. Enum. Pl. Zeyl.* 366.—Zeylan.
Wrightii, Munro, ex Vasey, Cat. Gram. U. St. 29 = Sorghum pauciflorum.
zeylanicus, Thw. Enum. Pl. Zeyl. 366 = Arnottianus.

CHRYSOPSIS, Ell. Sketch, ii. 333 (1824). *COMPOSITAE*, Benth. & Hook. f. ii. 252.
AMMODIA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 321.
DIPLOGON, Rafin. in Am. Monthly Mag. (1818) 268.
HECTOREA, DC. Prod. v. 95 (1836).
HEYFELDERA, Sch. Bip. in Flora, xxxvi. (1853) 35.
MACRONEMA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 322.
PITYOPSIS, Nutt. l. c. (1841) 317.
acaulis, Nutt. in Journ. Acad. Philad. vii. (1834) 33 = Aplopappus acaulis.
alba, DC. Prod. v. 264, in syn. = Aster ptarmicoides.
alpina, Nutt. in Journ. Acad. Philad. vii. (1834) 34. t. 3. f. 2 = Aster scopulorum.
amygdalina, DC. Prod. v. 272, in syn. = Aster umbellatus.
andicola, *Phil. in Anal. Univ. Chil.* (1862) ii. 398; *et in Linnaea*, xxxiii. (1864-65) 131.—Chili.
argentea, Ell. Sketch, ii. 334 = graminifolia.
Bolanderi, A. Gray, in Proc. Am. Acad. vi. (1864-65) 543 = villosa.
Breweri, *A. Gray, l. c.* 542.—Calif.
caespitosa, Nutt. in Journ. Acad. Philad. vii. (1834) 33 = Aplopappus acaulis.
canescens, DC. Prod. v. 328 = Erigeron filifolius.
canescens, Torr. & Gray, Fl. N. Am. ii. 256 = villosa.
coronopifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 34 = Aster tanacetifolius.
decumbens, Chapm. Fl. S. U. St. 217 = gossypina.
dentata, Ell. Sketch, ii. 337 = gossypina.
divaricata, Ell. l. c. 338 = Aster divaricatus.

CHRYSOPSIS:—
echioides, Benth. Bot. Voy. Sulph. 25 = villosa.
ericoides, Torr. & James, ex Eaton, N. Am. Bot. ed. VIII. 195 = Aster ericaefolius.
falcata, Ell. Sketch, ii. 336 = villosa.
foliosa, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 316 = villosa.
gossypina, *Ell. Sketch*, ii. 237.—Am. bor.
graminifolia, *Ell. l. c.* 334.—Am. bor.
hirtella, DC. Prod. v. 327 = Erigeron chrysopsidis.
hispida, DC. l. c. vii. 279 = villosa.
humilis, DC. l. c. v. 273, in syn. = Aster infirmus.
hyssopifolia, *Nutt. in Journ. Acad. Philad.* vii. (1834) 67.—Am. bor.
Lamarckii, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 315 = Aplopappus spinulosus.
linariifolia, DC. Prod. v. 277 = Aster linariifolius.
linifolia, DC. l. c. 255, in syn. = Aster acris.
Mariana, *Ell. Sketch*, ii. 335.—Am. bor.
mollis, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 316 = villosa.
obovata, Nutt. Gen. Am. ii. 152 = Aster reticulatus.
oligantha, Chapm. ex Torr. & Gray, Fl. N. Am. ii. 253 = graminifolia.
oregana, *A. Gray, in Proc. Am. Acad.* vi. (1864–65) 543.—Am. bor. occ.
pilosa, *Nutt. in Journ. Acad. Philad.* vii. (1834) 66. —Texas.
pinifolia, *Ell. Sketch*, ii. 335.—Am. bor.
scabra, Ell. l. c. 339 = Heterotheca Lamarckii.
scabra, Hook. & Arn. Bot. Beech. Voy. 434 = Oxypappus scaber.
scabrella, *Torr. & Gray, Fl. N. Am.* ii. 255.—Am. bor.
sessiliflora, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 317 = villosa.
trichophylla, *Ell. Sketch*, ii. 336.—Am. bor.
villosa, *DC. Prod.* v. 327; vii. 279.—Am. bor.
Wrightii, *A. Gray, Syn. Fl. N. Am.* i. II. 445.—Calif.

CHRYSORHOË, Lindl. in Hook. Comp. Bot. Mag. ii. (1836) 357 = **Verticordia**, DC. (Myrtac.).
nitens, Lindl. l. c. = V. nitens.
serrata, Lindl. Swan River App. 6 = V. serrata.

CHRYSOSCIAS, E. Mey. Comm. Pl. Afr. Austr. 139 (1835) = **Rhynchosia**, Lour. (Legumin.).
calycina, E. Mey. l. c. 140 = R. leucoscias.
floribunda, Lem. in Illustr. Hortic. i. (1854) Misc. 56; ii. (1855) t. 47.—Afr. austr.
grandiflora, E. Mey. Comm. Pl. Afr. Austr. i. 139 = R. Chrysoscias.
media, E. Mey. l. c.—Afr. austr.
parviflora, E. Mey. l. c. = R. microscias.

CHRYSOSPERMUM, Reichb. Consp. 166 (1828), nomen = **Anthospermum**, Linn. (Rubiac.).
repens, Pritz. Ind. Ic. 263, sphalm. = *Cryphiospermum repens*, Beauv. = Enhydra fluctuans, *Lour.* (Compos.).

CHRYSOSPLENIUM, Tourn. ex Linn. Syst. i. (1735). *SAXIFRAGEAE*, Benth. & Hook. f. i. 638.
adoxoides, Hook. f. & Thoms. ex Maxim. in Bull. Acad. Pétersb. xxiii. (1877) 342 = lanuginosum.
album, *Maxim. l. c.* xvii. (1872) 420.—Japon.
alpinum, Schur, in Verh. Siebenb. Ver. Naturw. v. (1852) 86 = oppositifolium.
alternans, Thunb. Fl. Jap. 182 = alternifolium.
alternifolium, *Linn. Sp. Pl.* 398.—Europ.; As. bor.; Am. bor.
americanum, *Schwein. ex Hook. Fl. Bor. Am.* i. 242.—Am. bor.
aulacocarpum, *Ernst, in Flora*, lvii. (1874) 214.—Venezuela.
auriculatum, Crantz, Inst. i. 133 = oppositifolium.
axillare, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 341.—China.
baicalense, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 21.—Sibir.
carnosulum, Maxim. in Bull. Acad. Pétersb. xxiii. (1877) 340 = carnosum.
carnosum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 73.—Reg. Himal.
Davidianum, *Decne. ex Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 343.—Tibet. or.
Delavayi, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 7.—As. or.

CHRYSOSPLENIUM:—
Dickinsii, *Franch. & Sav. Enum. Pl. Jap.* ii. 357.—Japon.
dubium, J. Gay, ex DC. Prod. iv. 48 = macrocarpum.
echinulatum, *Franch. & Sav. Enum. Pl. Jap.* ii. 358.—Japon.
Echinus, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 347.—Japon.
elevatum, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 281 = macranthum.
Fauriae, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 85.—Japon.
flagelliferum, *F. Schmidt, Reisen Amurl.* 134.—As. or.
flagelliforme, Franch. & Sav. Enum. Pl. Jap. ii. 355, lapsu = pracc.
geoides, Fisch. ex Ledeb. Fl. Ross. ii. 227 = alternifolium.
glaciale, Fuss, Fl. Transs. 247 = oppositifolium.
glechomaefolium, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 589.—Am. bor. occ.
Grayanum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 348.—Japon.
Griffithii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 74.—Reg. Himal.
kamtschaticum, *Fisch. ex DC. Prod.* iv. 48.—Kamtschat.
kamtschaticum, A. Gray, Bot. Jap. 389 = ramosum.
lanuginosum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 74.—Reg. Himal.
macranthum, *Hook. Lond. Journ. Bot.* i. (1842) 458. t. 16.—Patagon.
macrocarpum, *Cham. in Linnaea*, vi. (1831) 558.—Am. arct.
macrostemon, *Maxim. ex Franch. & Sav. Enum. Pl. Jap.* ii. 358.—Japon.
Maximowiczii, *Franch. & Sav. l. c.*—Japon.
multicaule, *Franch. & Sav. l. c.* 361.—Japon.
nepalense, *D. Don, Prod. Fl. Nep.* 210.—Reg. Himal.
nepalense, Maxim. Prim. Fl. Amur. 121, in nota = sulcatum.
nipponicum, *Franch. & Sav. Enum. Pl. Jap.* ii. 356.—Japon.
nivale, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 28 = alternifolium.
nudicaule, *Bunge, in Ledeb. Fl. Alt.* ii. 114.—Sibir.
octandrum, Caqué, ex Steud. Nom. ed. II. i. 360 = oppositifolium.
oppositifolium, Cham. in Linnaea, vi. (1831) 557 = kamtschaticum.
oppositifolium, *Linn. Sp. Pl.* 398.—Europ.; As. bor.; Ind. or.
oppositifolium, Trautv. & Mey. in Middend. Reise (Fl. Ochot). 42 = ramosum.
oppositifolium, Walt. Fl. Carol. 140 = americanum.
ovalifolium, Bertol. Fl. Ital. iv. 449 = macrocarpum.
ovalifolium, *Bieb. ex Ledeb. Fl. Alt.* ii. 115.—Sibir. altaic.
ovalifolium, Miq. Ann. Mus. Lugd. Bat. iii. 97 = macrostemon.
papillosum, Franch. & Sav. Enum. Pl. Jap. ii. 355 = alternifolium.
peltatum, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 273.—Sibir.
pilosum, *Maxim. Prim. Fl. Amur.* 122.—Manchur.
ramosum, *Maxim. l. c.* 121.—Manchur.
repens, Link, ex Steud. Nom. ed. II. i. 360 = oppositifolium.
rhabdospermum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 347.—As. or.
rosulare, Schott, ex Maxim. l. c. 345 = oppositifolium.
rotundifolium, Schlecht. ex Ledeb. Fl. Ross. ii. 227 = alternifolium.
Sedakowii, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 273.—Sibir.
sibiricum, Steph. ex DC. Prod. iv. 48 = alternifolium.
sinicum, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 348.—China.
sphacrospermum, *Maxim. l. c.* 349.—Japon.
sulcatum, Maxim. l. c. 347 = nepalense.
tenellum, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 73.—Reg. Himal.
tetrandrum, Fries, in Bot. Notiser (1858), 193 = alternifolium.
trachyspermum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 474.—China.
transsilvanicum, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 28.—Transsylvan.

CHRYSOSPLENIUM:—
trichospermum, *Edgew. ex Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 73.—Reg. Himal.
uniflorum, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 472.—China.
valdivicum, *Hook. Lond. Journ. Bot.* i. (1842) 459 t. 17.—Chili.
Vidalii, *Franch. & Sav. Enum. Pl. Jap.* ii. 360.—Japon.
Wrightii, *Franch. & Sav. l. c.* 356.—Sibir. or.
yesoënse, *Franch. & Sav. l. c.* 355.—Japon.
yunnanense, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) 7.—China.

CHRYSOSTACHYS, Pohl, Pl. Bras. ii. 65. t. 143 (1831) = **Combretum**, Linn.
ovatifolia, Pohl, l. c. 66.—Bras.

CHRYSOSTEMMA, Less. Syn. Comp. 227 (1832) = **Coreopsis**, Linn. (Compos.).
calendulacea, E. Mey. ex DC. Prod. vi. 501 = Gorteria calendulacea.
senifolium, Shuttl. ex Walp. Ann. ii. 857 = Cor. senifolia.
Tripteris, Less. Syn. Comp. 227 = Cor. Tripteris.

CHRYSOSTEMON, Klotzsch, in Lehm. Pl. Preiss. ii. 232 (1847) = **Pseudanthus**, Sieber (Compos.).
virgatus, Klotzsch, l. c. = P. virgatus.

CHRYSOSTOMA, Lilja, Fl. Suppl. i. 33 (1840); et in Linnaea, xv. (1841) 263 = **Mentzelia**, Linn. (Loas.).
aurea, Lilja, ll. cc. 33, 264 = M. oligosperma.

CHRYSOTHAMNUS, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 323 = **Bigelowia**, DC. (Compos.).
depressus, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 171 = B. depressa.
dracunculoides, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 324 = B. dracunculoides.
lanceolatus, Nutt. l. c. = B. viscidiflora.
pumilus, Nutt. l. c. 323 = B. viscidiflora.
speciosus, Nutt. l. c. = B. viscidiflora.
viscidiflorus, Nutt. l. c. 324 = B. viscidiflora.

CHRYSOTHEMIS, Decne. in Rev. Hortic. Sér. III. iii. (1849) 242 = **Tussacia**, Reichb. (Gesnerac.).
aurantiaca, Decne. l. c. iv. 1850) 381.—Ind. occ.
melissaefolia, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1402 = T. pulchella.
mellitifolia, G. Don, ex Loud. l. c. = pracc.
ovata, Klotzsch, in Schomb. Fauna et Fl. Gui. 108 — Guiana.
pulchella, Decne. in Rev. Hortic. Sér. III. iii. (1849) 242 = T. pulchella.
venosa, Decne. l. c. = Episcia melittifolia?

CHRYSOTHRIX, Roem. & Schult. Mant. ii. 6. 144 (1824) = **Chrysithrix**, Linn. (Cyperac.).

CHRYSOXYLON, Casar. Nov. Stirp. Bras. Dec. 59 (1843) = **Plathymenia**, Benth. (Legumin.).
Vinhatico, Casar. l. c. = P. reticulata.

CHRYSOXYLON, Wedd. Hist. Nat. Quinquin. 100 (1849) = **Pogonopus**, Klotzsch (Rubiac.).
febrifugum, Wedd. l. c.—Bolivia.

CHRYSURUS, Pers. Syn. i. 80 (1805) = **Lamarckia**, Moench (Gramin.).
aureus, Beauv. Agrost. 123 = L. aurea.
cynosuroides, Pers. Syn. i. 80 = L. aurea.
echinatus, Beauv. Agrost. 123 = Cynosurus echinatus.
effusus, Beauv. l. c. = Cynosurus elegans.
elegans, Beauv. l. c. = Cynosurus polybracteatus.
giganteus, Tenore, Fl. Nap. iv. 13. t. 210 = Cynosurus echinatus.
gracilis, Moris, Stirp. Sard. Elench. i. 50 = Cynosurus gracilis.

CHRYTOTHECA, G. Don, Gen. Syst. ii. 706 (1832) = CRYPTOTHECA, Blume = **Ammannia**, Houst. (Lythrar.).

CHRYZA, Rafin. in Med. Repos. N. York, v. (1808) 352 = **Coptis**, Linn. (Ranunculac.).
borealis, Rafin. l. c. = Coptis trifolia.

CHTHAMALIA, Decne. in DC. Prod. viii. 605 (1844) = **Gonolobus**, Michx. (Asclepiad.).
biflora, Decne. l. c. = G. biflorus.
humifusa, Fourn. in Mart. Fl. Bras. vi. IV. 313.—Bras.
major, Fourn. l. c. 314.—Bras.
nummularia, Decne. in DC. Prod. viii. 605 = G. nummularins.
pedunculata, Decne. l. c. = G. pedunculatus.
pubiflora, Decne. l. c. = G. pubiflorus.
purpurea, Decne. l. c.—Bras.

CHTHONIA, Cass. in Dict. Sc. Nat. ix. 173 (1817) = **Pectis**, Linn. (Compos.).
Bonplandiana, Cass. l. c. xxvii. 206 = P. Bonplandiana.
canescens, Cass. l. c. = P. canescens.
ciliaris, Cass. l. c. = P. ciliaris.
elongata, Cass. l. c. = P. elongata.
glaucescens, Cass. l. c. ix. 173 = P. ciliaris.
humifusa, Cass. l. c. = P. humifusa.
leptocephala, Cass. l. c. xxvii. 206 = P. Swartziana.
prostrata, Cass. l. c. ix. 173 = P. prostrata.
pygmaea, Cass. l. c. xxvii. 206 = P. pygmaea.
repens, Cass. l. c. = P. serpyllifolia.

CHTHONOCEPHALUS, Steetz, in Lehm. Pl. Preiss. i. 444 (1845). *COMPOSITAE*, Benth. & Hook. f. ii. 322.
CHAMAESPHAERION, A. Gray, in Hook. Kew Journ. iii. (1851) 176.
GYROSTEPHIUM, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) II. 76.
LACHNOTHALAMUS, F. Muell. Fragm. iii. 156 (1863).
Drummondii, *A. Gray, in Hook. Kew Journ.* iii. (1851) 178.—Austral.
pseudevax, *Steetz, in Lehm. Pl. Preiss.* i. 445.—Austral.
pygmaeus, *Benth. Fl. Austral.* iii. 582.—Austral.
tomentellus, *Benth. l. c.* 581.—Austral.

CHUKRASIA, A. Juss. in Mém. Mus. Par. xix. (1830) 251. t. 22, vide **Chickrassia**, A. Juss. (Meliac.).

CHULUSIUM, Rafin. Ann. Nat. 12 (1820) = **Polygonum**, Tourn.
acuminatum, Rafin. Fl. Tellur. iii. 14 = P. acuminatum.
amphibium, Rafin. l. c. = P. amphibium.
filiforme, Rafin. l. c. = P. tenue.
fluitans, Rafin. l. c. = P. amphibium?
natans, Rafin. l. c. = P. amphibium?

CHUNCHOA, Pers. Syn. i. 486 (1805) = seq.

CHUNCOA, Pav. ex Juss. Gen. 76 (1789) = **Terminalia**, Linn. (Combretac.).
actinophylla, Griseb. in Goett. Abh. xxiv. (1879) 133 = T. actinophylla.
amazonia, J. F. Gmel. Syst. 702.—Bras.
arbuscula, Griseb. Fl. Brit. W. Ind. 275 = T. arbuscula.
brasiliensis, Cambess. in St. Hil. Fl. Bras. Mer. ii. 244 = T. brasiliensis, *Eichl.*
Chicharronia, Griseb. Cat. Pl. Cub. 109 = T. Chicharronia.
diptera, F. G. Dietr. Vollst. Lex. Gaertn. Nachtr. ii. 244 = T. oblonga.
eriostachya, Griseb. Cat. Pl. Cub. 109 = T. eriostachya.
flavescens, Presl, Epim. Bot. 215.—Bras.
glabra, Buch.-Ham. ex Wall. Cat. sub n. 3979 = T. Arjuna.
lactiflua, F. Muell. Fragm. ii. 149 = T. bursarina.
oblonga, Pers. Syn. i. 486 = T. oblonga.
obovata, Pers. l. c. = T. obovata.
tomentosa, Buch.-Ham. ex Wall. Cat. sub n. 3978 = T. tomentosa.
triflora, Griseb. in Goett. Abh. xxiv. (1879) 132.—Reg. Argent.

CHUPALON, Adans. Fam. ii. 164 (1763) = **Thibaudia**, Pav. (Vacciniac.).

CHUPALONES, Nieremb. ex Steud. Nom. ed. II. i. 361 (1840) = praec.

CHUQUIRAGA, Juss. Gen. 178 (1789). *COMPOSITAE*, Benth. & Hook. f. ii. 488.
DASYPHYLLUM, H. B. & K. Nov. Gen. et Sp. iv. 17. t. 308 (1820).
DIACANTHA, Lag. Amen. Nat. i. 40 (1811).
FLOTOVIA, Spreng. Syst. iii. 359 (1826).
JOANNESIA, Pers. Syn. ii. 383 (1807).
JOHANNIA, Willd. Sp. Pl. iii. 1705 (1801).
PIPTOCARPA, Hook. & Arn. in Comp. Bot. Mag. i. (1835) 110.
acanthophylla, *Wedd. Chlor. And.* i. 5.—Bolivia.
acicularis, *D. Don, in Phil. Mag.* (1832) 392.—Chili.
alpina, Poepp. ex Less. Syn. Comp. 96 = oppositifolia.
anomala, *D. Don, in Phil. Mag.* (1832) 392.—Chili.
Avellanedae, *Lorentz, in Roca, Exped. Rio-Negro*, ii. 245.—Patagon.
Candolleana, *Baker, in Mart. Fl. Bras.* vi. III. 358.—Bras.
chrysantha, *Field. & Gardn. Sert. Pl.* t. 42.—Chili.
cryptocephala, *Baker, in Mart. Fl. Bras.* vi. III. 355.—Bras.
Doniana, *Baker, l. c.* 358.—Bras.
Echegarayi, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 47.—Reg. Argent.
erinacea, *D. Don, in Phil. Mag.* (1832) 392.—Chili; Patagon.
excelsa, *D. Don, l. c.*—Chili.
floribunda, *Baker, in Mart. Fl. Bras.* vi. III. 362.—Bras.
fodinarum, *Baker, l. c.* 356.—Bras.
fruticosa, Juss. ex Steud. Nom. ed. II. i. 361 = insignis.
glabra, *Baker, in Mart. Fl. Bras.* vi. III. 363.—Bras.
Grisebachii, *Hieron. in Bol. Acad. Nac. Cordova*, iv. (1881) 359.—Reg. Argent.
hispida, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 287.—Bras.
Hystrix, *D. Don, in Phil. Mag.* (1832) 392.—Chili; Patagon.
incana, D. Don, l. c. = ulicina.
infundibularis, *Baker, in Mart. Fl. Bras.* vi. III. 357.—Bras.
insignis, Humb. & Bonpl. Pl. Aequin. i. 153 = Jussieui.
juniperina, *Decne. in Orbign. Voy. Bot.* t. 11.—Am. austr.
Jussieui, *J. F. Gmel. Syst.* 1205.—Peruv.; Ecuador.
Kingii, *Ball, in Journ. Linn. Soc.* xxi. (1884) 224.—Patagon.
lancifolia, *Humb. & Bonpl. Pl. Aequin.* i. 153.—Peruv.; Ecuador.
latifolia, *Baker, in Mart. Fl. Bras.* vi. III. 357.—Bras.
latifolia, D. Don, in Trans. Linn. Soc. xvi. (1830) 288 = glabra.
leptacantha, *Baker, in Mart. Fl. Bras.* vi. III. 360.—Bras.
leucoxylon, *Poepp. ex Less. Syn. Comp.* 95, in syn.—Chili.
linearis, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) II. 94.—Chili.
macrocephala, *Baker, in Mart. Fl. Bras.* vi. III. 359.—Bras.
microphylla, Humb. & Bonpl. Pl. Aequin. i. 151. t. 43 = Jussieui.
oppositifolia, *D. Don, in Phil. Mag.* (1832) 392.—Chili.
orthacantha, *Baker, in Mart. Fl. Bras.* vi. III. 360.—Bras.
paniculata, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 289.—Bras.
patagonica, *Phil. in Linnaea*, xxxiii. (1864-65) 111.—Patagon.
peruviana, Jaume St. Hil. Fam. Nat. i. 394 = Jussieui.
racemosa, *Baker, in Mart. Fl. Bras.* vi. III. 363.—Bras.
Regnellii, *Baker, l. c.* 359.—Bras.
reticulata, *DC. Prod.* vii. 9.—Bras.
revoluta, *Field. & Gardn. Sert. Pl.* sub t. 43.—Chili.
rotundifolia, *Wedd. Chlor. And.* i. 4. t. 4.—Peruv.
ruscifolia, *D. Don, in Phil. Mag.* (1832) 392.—Chili.
spinescens, *Baker, in Mart. Fl. Bras.* vi. III. 362.—Bras.
spinosa, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 285.—Peruv.
Sprengeliana, *Baker, in Mart. Fl. Bras.* vi. III. 357.—Bras.
synacantha, *Baker, l. c.* 361.—Bras.
tomentosa, Baker, l. c. 360 = paniculata.
trichophylla, *Baker, l. c.* 356.—Bras.
ulicina, *Hook. Comp. Bot. Mag.* i. (1835) 110.—Chili.

CHUQUIRAGA:—
Unguis-cati, *Ces. in Atti Soc. Sc. Nap.* v. (1873) 11. n. 7.—Hab.?
vagans, *Baker, in Mart. Fl. Bras.* vi. III. 361.—Bras.
velutina, *Baker, l. c.* 358.—Bras.
vepreculata, *D. Don, in Trans. Linn. Soc.* xvi. (1830) 290.—Venezuela.

CHURUMAYA, Rafin. Sylva Tellur. 85 (1838) = **Piper**, Linn.
arborea, Rafin. l. c. = P. Churumaya.

CHUSQUEA, Kunth, Syn. Pl. Aequin. i. 254 (1822). *GRAMINEAE*, Benth. & Hook. f. iii. 1209.
DENDRAGROSTIS, Nees, in Linnaea, ix. (1834) 487.
MUSTELIA, Cav. ex Steud. Nom. ed. II. i. 361, in syn. (1840); ii. 168.
RETTBERGIA, Raddi, Agrost. Bras. 17 (1823).
abietifolia, *Griseb. Fl. Brit. W. Ind.* 529.—Ind. occ.
acuminata, *Doell, in Mart. Fl. Bras.* ii. III. 204.—Bras.
amplopaniculata, Steud. Syn. Pl. Gram. 337 = Dinochloa Tjankorreh.
andina, *Phil. in Linnaea*, xxix. (1857-58) 103.—Chili.
anelythra, *Nees, in Linnaea*, ix. (1834) 491.—Bras.
anelytroides, *Rupr. ex Doell, in Mart. Fl. Bras.* ii. III. 206.—Bras.
aristata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 61.—Ecuador.
Bilimeki, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 587.—Mexic.
breviglumis, Phil. in Linnaea, xxix. (1857-58) 103 = C. Culeou.
capitata, *Nees, in Linnaea*, ix. (1834) 489.—Bras.
capituliflora, *Trin. in Mém. Acad. Pétersb.* Sér. VI. iii. (1835) 613.—Bras.
carinata, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 587.—Mexic.
ciliata, Phil. in Linnaea, xxxiii. (1864-65) 299 = tenuiflora.
Culeou, *E. Desv. in C. Gay, Fl. Chil.* vi. 450.—Chili.
Cumingii, *Nees, in Linnaea*, ix. (1834) 487.—Chili.
Dombeyana, *Kunth, Rev. Gram.* ii. 553. t. 191.—Peruv.
fasciculata, *Doell, in Mart. Fl. Bras.* ii. III. 202.—Bras.
Fendleri, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 61.—Venezuela.
Fernandeziana, *Phil. in Anal. Univ. Chil.* (1873) 577.—Chili.
fimbriata, Steud. Syn. Pl. Gram. 338 = Merostachys capitata.
Galeottiana, *Rupr. in Bull. Acad. Brux.* ix. II. (1842) 246.—Mexic.
Gaudichaudii, *Kunth, Rev. Gram.* i. 138, 331. t. 78.—Bras.
glomerata, Munro, in Trans. Linn. Soc. xxvi. (1868) 50 = Merostachys capitata.
heterophylla, *Nees, in Linnaea*, ix. (1834) 488.—Bras.
humilis, Lechl. ex Munro, in Trans. Linn. Soc. xxvi. (1868) 60 = spicata.
intermedia, Steud. in Lechl. Berb. Am. Austr. 52, nomen = C. Quila.
Jamesoni, Steud. Syn. Pl. Gram. 337 = scandens.
leptophylla, Nees, in Linnaea, ix. (1834) 489 = Arthrostylidium leptophyllum, Trinii.
Liebmanni, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 587.—Mexic.
ligulata, Munro, in Trans. Linn. Soc. xxvi. (1868) 62 = Fernandeziana.
Lorentziana, *Griseb. in Goett. Abh.* xix. (1874) 249.—Reg. Argent.
Meyeriana, *Rupr. ex Doell, in Mart. Fl. Bras.* ii. III. 203.—Bras.
montana, *Phil. in Linnaea*, xxxiii. (1864-65) 298.—Chili.
Mulleri, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 65.—Mexic.
oligophylla, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 124.—Bras.
nigricans, *Phil. in Anal. Univ. Chil.* (1865) II. 323.—Chili.
pallida, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 65.—N. Granat.
parvifolia, Phil. in Linnaea, xxxiii. (1864-65) 299 = Cumingii.
pinifolia, *Nees, l. c.* ix. (1834) 490.—Bras.

CHUSQUEA :— .
pubescens, Steud. Syn. Pl. Gram. 337 = C. Quila.
Purdieana, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 56.—N. Granat.
Quila, E. Desv. in C. Gay, Fl. Chil. vi. 447 = tenuiflora.
Quila, *Kunth, Rev. Gram.* i. 138, 329. t. 77.—Chili.
scandens, *Kunth, Syn. Pl. Aequin.* i. 254.—Am. austr.
sclerophylla, *Doell, in Mart. Fl. Bras.* ii. III. 200.—Bras.
Sellowii, *Rupr. in Mém. Acad. Pétersb.* Sér. VI. v. (1839) 125.—Bras.
simpliciflora, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 54.—Panama.
simplicifolia, Hemsl. Biol. Centr. Am. Bot. iii. 587, sphalm. = simpliciflora.
Spencei, *Ernst, in Journ. Bot.* x. (1872) 262.—Venezuela.
spicata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 60.—Peruv.
spinosa, *Fourn. ex Hemsl. Biol. Centr. Am. Bot.* iii. 587.—Mexic.
tenella, *Nees, in Linnaea*, ix. (1834) 492.—Bras.
tenuiflora, *Phil. in Linnaea*, xxx. (1859–60) 206.—Chili.
tenuiglumis, *Doell, in Mart. Fl. Bras.* ii. III. 199.—Bras.
tessellata, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 60.—N. Granat.
uliginosa, *Phil. in Linnaea*, xxx. (1859–60) 207.—Chili.
uniflora, *Steud. Syn. Pl. Gram.* 337.—Ins. Cuba.
valdiviensis, E. Desv. in C. Gay, Fl. Chil. vi. 446 = C. Quila.
Venezuelae, *Steud. Syn. Pl. Gram.* 337.—Venezuela.
Wilkesii, *Munro, in Trans. Linn. Soc.* xxvi. (1868) 63.—Bras.

CHYDENANTHUS, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 111 = **Barringtonia**, Forst. (Myrtac.).
exeelsus, Miers, l. c. 112 = B. excelsa.

CHYLOCALYX, Hassk. in Flora, xxv. (1842) II. Beibl. 20 = **Polygonum**, Tourn.
perfoliatus, Hassk. l. c. = P. perfoliatum.
sentieosus, Meissn. in Miq. Ann. Mus. Bot. Lugd. Bat. ii. 65 = P. senticosum.

CHYLODIA, Rich. ex Cass. in Dict. Sc. Nat. xxix. 491 (1823) = **Wulffia**, Neck. (Compos.).
sarmentosa, Rich. l. c. 492 = W. stenoglossa.

CHYLOGALA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 150 = **Euphorbia**, Linn.
serrata, Fourr. l. c. = E. serrata.

CHYMOCARPUS. G. Don, in Trans. Linn. Soc. xvii. 13 (1834) = **Tropaeolum**, Linn. (Geraniac.).
brachyceras, Heynh. Nom. i. 195 = T. brachyceras.
Jarattii, Heynh. l. c. = T. Jarattii.
pentaphyllus, D. Don, in Trans. Linn. Soc. xvii. (1834) 14 = T. pentaphyllum.
polyphyllus, Heynh. Nom. i. 195 = T. polyphyllum.
speciosus, Walp. Ann. i. 142 = T. speciosum.
tricolor, Heynh. Nom. i. 195 = T. tricolorum.
tuberosus, Heynh. l. c. = T. tuberosum.

CHYMOCOCCA, Meissn. in DC. Prod. xiv. 565 (1857). *THYMELAEACEAE*, Benth. & Hook. f. iii. 194.
empetroides, *Meissn. l. e.*—Afr. austr.

CHYMOCORMUS, Harv. in Hook. Lond. Journ. Bot. i. (1842) 23 = **Fockea**, Endl. (Asclepiad.).
edulis, Harv. l. c. 24.—Afr. austr.

CHYSIS, Lindl. Bot. Reg. xxiii. t. 1937 (1837). *ORCHIDEAE*, Benth. & Hook. f. iii. 514.
aurea, *Lindl. l. e.*—Chili.
bractescens, *Lindl. l. e.* (1840) *Mise.* 61.—Mexic.
Bruennowiana, *Reichb. f. & Warse. in Bot. Zeit.* xv. (1857) 157.—Peruv.
Chelsoni ×, Reichb. f. in Gard. Chron. (1874) i. 535 (cf. p. 385).
laevis, *Lindl. Bot. Reg.* (1840) *Mise.* 61.—Mexic.
Limminghii, *Linden & Reichb. f. in C. Koch, Allg. Gartenz.* xxvi. (1858) 380.—Mexic.
Makoyi, *Heynh. Nom.* ii. 143.—Mexic.
Sedeni ×, Reichb. f. in Gard. Chron. (1880) i. 616.

CHYTRA, Gaertn. f. Fruct. iii. 184. t. 214 (1805) = **Buchnera**, Linn. ? (Scrophularin.).
anomala, Gaertn. f. l. c.—Hab. ?

CHYTRACULIA, P. Br. Hist. Jamaic. 239. t. 37 (1756) = **Calyptranthes**, Sw. (Myrtac.).

CHYTRALIA, Adans. Fam. ii. 80 (1763) = **Calyptranthes**, Sw. (Myrtac.).

CHYTRANTHUS, Hook. f. in Benth. & Hook. f. Gen. i. 403 (1863). *SAPINDACEAE*, Benth. & Hook. f. l. c.
Mannii, *Hook. f. l. c.*—Afr. trop.
Prieurianus, *Baill. Adansonia*, xi. (1873–76) 241.—Guiana.

CHYTROGLOSSA, Reichb. f. in Hamb. Gartenz. xix. (1863) 546. *ORCHIDEAE*, Benth. & Hook. f. iii. 569.
aurata, *Reichb. f. l. c.*—Hab. ?
Marileoniae, *Reichb. f. l. c.*—Bras.

CHYTROMA, Miers, in Trans. Linn. Soc. xxx. (1874) 229 = **Lecythis**, Loefl. (Myrtac.).
amara, Miers, l. c. 231 = L. amara.
apiculata, Miers, l. c. 245.—Bras.
basilaris, Miers, l. c. 239.—Guiana.
chartacea, Miers, l. c. 231 = L. chartacea.
eincturata, Miers, l. c. 237.—Bras.
eistella, Miers, l. c. 233.—Bras.
grandifolia, Miers, l. c. 237 = L. grandiflora.
Ibiriba, Miers, l. c. 236.—Bras.
Idatimon, Miers, l. c. 239 = L. Idatimon.
incarcerata, Miers, l. c. 232.—Bras.
marawynensis, Miers, l. c. 245 = L. marawynensis.
monosperma, Miers, l. c. 239 = L. monosperma.
ovalifolia, Miers, l. c. 244 = L. ovalifolia.
perspieua, Miers, l. c. 246.—Guiana.
pilocarpa, Miers, l. c. 238.—Bras.
retusa, Miers, l. c. 243 = L. retusa.
rorida, Miers, l. c. = L. rorida.
rosea, Miers, l. c. 242 = L. rosea.
rubriflora, Miers, l. c. 241.—Guiana.
salebrosa, Miers, l. c. 240 = L. salebrosa.
Schomburgkiana, Miers, l. c. 230 = L. Schomburgkii.
Spruceana, Miers, l. c. 235 = L. Spruceana.
turbinata, Miers, l. c. 234 = L. turbinata.
urceolata, Miers, l. c. 237.—Bras.
valida, Miers, l. c. 241.—N. Granat.

CIANITIS, Reinw. in Syll. Ratisb. ii. (1828) 10 = CYANITIS, Reinw. = **Dichroa**, Lour. (Saxifrag.).

CICCA, Linn. Mant. i. 124 (1767) = **Phyllanthus**, Linn. (Euphorbiac.).
aeidissima, Blanco, Fl. Filip. ed. I. 700 = P. acidissimus.
albizzioides, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 239.—Burma.
anomala, Baill. Etud. Gén. Euph. 619 = P. indicus.
antillana, A. Juss. Tent. Euph. 108. t. 4. f. 13 B = P. nobilis.
arborea, Wright, ex Griseb. in Goett. Nachr. (1865) 166 = P. nobilis.
brasiliensis, Baill. Etud. Gén. Euph. 618, nomen = P. nobilis.
chinensis, Baill. l. c.; Atlas, 45 = P. nobilis.
decandra, Blanco, Fl. Filip. ed. I. 701 = P. reticulatus.
discoidea, Baill. in Adansonia, i. (1860–61) 85 = P. discoideus.
disticha, Linn. Mant. i. 124 = P. distichus.
Emblica, Kurz, For. Fl. Brit. Burma, ii. 352 = P. Emblica.
flexuosa, Sieb. & Zucc. Fl. Jap. Fam. Nat. i. 35 = P. japonicus.
Gaertneriana, Muell. Arg. in DC. Prod. xv. II. 416, in syn. = P. cyanospermus.
Leucopyrus, Kurz, For. Fl. Brit. Burma, ii. 353 = Fluggia Leucopyrus.
maerocarpa, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 239.—Burma.
maerostachya, Benth. Bot. Voy. Sulph. 166 = Picramnia Antidesma.
mieroearpa, Penth. Fl. Hongk. 312 = P. reticulatus.
nodiflora, Lam. Encyc. ii. 1 = P. distichus.
obovata, Kurz, For. Fl. Brit. Burma, ii. 354 = Fluggia microcarpa.

CICCA :—
Pavoniana, Baill. Etud. Gén. Euph. 618 = Phyllanthus nobilis.
pentandra, Blanco, Fl. Filip. ed. I. 701 = Fluggia microcarpa.
racemosa, Lour. Fl. Cochinch. ii. 556 = P. acidissimus ?
retieulata, Kurz, For. Fl. Brit. Burma, ii. 354 = P. reticulatus.
rhomboidalis, Baill. Adansonia, ii. (1861–62) 51 = P. rhomboidalis.
scandens, Griseb. in Goett. Nachr. (1865) 165 = P. scandens.
siniea, Baill. Etud. Gén. Euph. 618. t. 24 = P. nobilis.
surinamensis, Miq. in Linnaea, xxi. (1848) 479 = P. nobilis.
virens, Wright, ex Griseb. in Goett. Nachr. (1865) 166 = P. virens.

CICENDIA, Adans. Fam. ii. 503 (1763). *GENTIANEAE*, Benth. & Hook. f. ii. 810.
campestris, Rafin. Fl. Tellur. iii. 26 = Gentiana campestris.
Candollei, Griseb. Gen. et Sp. Gent. 158 = pusilla.
exaltata, Griseb. l. c. 159 = Erythraea Douglasii.
fastigiata, Griseb. l. c. 158 = Hoppea fastigiata.
filiformis, Delarb. Fl. Auv. i. 20; ed. II. 20 = Microcala filiformis.
glacialis, Rafin. Fl. Tellur. iii. 26 = Gentiana tenella.
hyssopifolia, Wight & Arn. in Hook. Comp. Bot. Mag. ii. (1836) 249 = Enicostema littorale.
microphylla, Edgew. in Trans. Linn. Soc. xx. (1851) 83 = Sebaea khasiana.
Poeppigii, Griseb. in DC. Prod. ix. 62 (Microcalae sp).—Bras.
pulchella, Griseb. Gen. et Sp. Gent. 159 = Erythraea ramosissima.
pusilla, *Griseb. l. e.* 157.—Europ.
quadrangularis, Griseb. l. c. = Microcala quadrangularis.
quitensis, Griseb. in Linnaea, xxii. (1849) 33 = Erythraea quitensis.
Roxburghii, Griseb. Gen. et Sp. Gent. 160 = Hoppea dichotoma.
stricta, *Griseb. in Journ. Linn. Soc.* vi. (1862) 143.—Guatemala.
tenella, Rafin. Fl. Tellur. iii. 26 = Gentiana tenella.

CICER, [Tourn.] Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 524.
NOCHOTTA, S. G. Gmel. Reise, iii. 23. t. 3 (1774).
Adonis, Orph. ex Nym. Consp. 200 = ervoides.
anatolicum, *Alef. in Bonplandia*, ix. (1861) 349.—As. Min.
arietinum, *Linn. Sp. Pl.* 738.—Europ.; Oriens; Ind. or.
cuneatum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 195 = arietinum.
ervoides, Brign. Fasc. Pl. Forojul. 27 = Lens Lenticula.
ervoides, *Fenzl, in Russegg. Reise*, i. 894, in obs.—Graecia; As. Min.; Syria.
floribundum, *Fenzl, l. c.* 892. t. 9.—Cilicia.
glutinosum, Alef. in Bonplandia, ix. (1861) 349 = anatolicum.
graecum, *Orph. ex Boiss. Diagn.* Ser. II. ii. 43.—Graecia.
grossum, Salisb. Prod. 340 = arietinum.
Jacquemontii, Jaub. & Spach, in Ann. Sc. Nat. Ser. II. xviii. (1842) 231 = soongaricum.
judaicum, Boiss. Diagn. Ser. I. ix. 180 = pinnatifidum.
Lens, Willd. Sp. Pl. iii. 1114 = Lens esculenta.
microphyllum, Royle, Illustr. Bot. Himal. 200 = soongaricum.
minutum, *Boiss. & Hohen. Diagn.* Ser. I. ix. 180.—Persia.
Montbretii, *Jaub. & Spach, in Ann. Sc. Nat.* Ser. II. xviii. (1842) 229.—As. Min.
nigrum, Hort. ex Steud. Nom. ed. II. i. 361 = Lens esculenta.
numulariaefolium, Lam. Encyc. ii. 2 = Crotalaria biflora.
oxyodon, *Boiss. & Hohen. Diagn.* Ser. I. ix. 180.—Persia.
physodes, Reichb. Fl. Germ. Excurs. 532 = arietinum.
pimpinellifolium, Jaub. & Spach, in Ann. Sc. Nat. Ser. II. xviii. (1842) 228 = ervoides.

CICER :—
pinnatifidum, *Jaub. & Spach, in Ann. Sc. Nat.* Sér. II. xviii. (1842) 227.—As. Min.; Mesopotam.
punctatum, Hort. ex Steud. Nom. ed. II. i. 361 = Lens Lenticula.
punctulatum, Hort. ex Ser. in DC. Prod. ii. 366 = Lens esculenta.
pungens, *Boiss. Diagn.* Ser. II. ii. 44.—Afghanist.
sativum, Schkuhr, Handb. ii. 367. t. 202 = arietinum.
soloniense, Schrank, ex G. Don, Gen. Syst. ii. 312 = Lens Lenticula.
songaricum, Jaub. & Spach, Illustr. i. 84. t. 43 B = anatolicum.
songaricum, *Steph. ex DC. Mém. Lég.* viii. 349.—Soongar.; Himal. bor. occ.
spiroceras, *Jaub. & Spach, in Ann. Sc. Nat.* Sér. II. xviii. (1842) 233.—Persia.
subaphyllum, *Boiss. Diagn.* Ser. I. vi. 44.—Persia.
tragacanthoides, *Jaub. & Spach, in Ann. Sc. Nat.* Sér. II. xviii. (1842) 234.—Persia; Turkestan.

CICERBITA, Wallr. Sched. Crit. 433 (1822) = **Lactuca**, Tourn. (Compos.).
acuminata, Wallr. l. c. 434 = L. acuminata.
alpina, Wallr. l. c. = L. leucophaea.
borealis, Wallr. l. c. = L. leucophaea.
canadensis, Wallr. l. c. = L. canadensis.
corymbosa, Wallr. l. c. = L. quercina.
elongata, Wallr. l. c. = L. canadensis.
floridana, Wallr. l. c. = L. floridana.
intybacea, Wallr. l. c. = Brachyramphus intybaceus.
leucophaea, Wallr. l. c. = L. leucophaea.
macrophylla, Wallr. l. c. = L. macrophylla.
muralis, Wallr. l. c. 436 = L. muralis.
Plumieri, Kirschl. in Mém. Soc. Hist. Nat. Strassb. iv. (1855) [n. xv.] t. 1 = L. Plumieri.
quercina, Wallr. Sched. Crit. 435 = L. quercina.

CICERCULA, Medic. in Vorles. Churpf. Phys. Ges. i. (1787) 358 = **Lathyrus**, Tourn. (Legumin.).
alata, Moench, Meth. 163 = L. sativus.
alba, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358 = L. sativus.
anceps, Moench, Meth. 163 = L. Cicera.
caerulea, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358 = L. sativus.
Cicera, Alef. in Bonplandia, ix. (1861) 147 = L. Cicera.
sativa, Medic. Phil. Bot. i. 203 = L. sativus.

CICERELLA, DC. Prod. ii. 369 (1825) = praec.

CICHORIUM, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 506.
ACANTHOPHYTON, Less. Syn. Comp. 128 (1832).
alatum, Hochst. & Steud. ex DC. Prod. vii. 75 = Geigeria alata.
ambiguum, Schult. Obs. 170 = pumilum.
byzantinum, Clem. in Mem. Acc. Sc. Torin. Ser. II. xvi. (1857) 285 = C. Intybus.
caeruleum, Gilib. Fl. Lituan. i. 224 = C. Intybus.
callosum, *Pomel, Nouv. Mat. Fl. Atl.* 17.—Afr. bor.
calvum, *Sch. Bip. in Ind. Sem. Hort. Berol.* (1859) 7, nomen; *Schweinf. Beitr. Fl. Aethiop.* 143.—Afr. trop.
Casnia, C. B. Clarke, Comp. Ind. 250 = C. Intybus.
Cicorea, Dum. Fl. Belg. Prod. 64 = C. Intybus.
commune, Pall. Reise, iii. 655 = C. Intybus.
Cosnia, Buch.-Ham. in Wall. Cat. n. 3241 = C. Intybus.
crispum, Mill. Gard. Dict. ed. VIII. n. 4 = C. Endivia.
dichotomum, Link, Handb. i. 811, sphalm. = pumilum.
divaricatum, Heldr. ex Nym. Consp. 472 = C. Intybus.
divaricatum, Schousb. Vextr. Marok. 197 = pumilum.
Endivia, *Linn. Sp. Pl.* 813.—Europ.; Oriens.
esculentum, Salisb. Prod. 183 = C. Endivia.
glabratum, Presl, Fl. Sic. i. 32 = C. Intybus.
glandulosum, *Boiss. & Huet, Diagn.* Ser. II. iii. 87.—As. Min.; Mesopotam.
glaucum, *Hoffmgg. & Link, Fl. Port.* ii. 178.—Hispan.
hirsutum, Gren. in Acad. Sc. Besanç. Séanc. Publ. (1838) 137 = C. Intybus.
Intybus, *Linn. Sp. Pl.* 813.—Europ.; Oriens.

CICHORIUM:—
minimum, *Portenschl. Enum. Pl. Dalm.* 16.—Dalmatia.
nanum, Portenschl. ex Nym. Consp. 472 = pumilum.
Noëanum, *Boiss. Fl. Orient.* iii. 717.—Mesopot.
officinale, Gueldenst. ex Ledeb. Fl. Ross. ii. 774 = C. Intybus.
perenne, Stokes, Bot. Mat. Med. 133 = C. Intybus.
polystachyum, *Pomel, Nouv. Mat. Fl. Atl.* 18.—Afr. bor.
pumilum, *Jacq. Obs.* iv. 3. t. 80.—Reg. Mediterr.
rigidum, Salisb. Prod. 183 = C. Intybus.
spinosum, *Linn. Sp. Pl.* 813.—Italia; Graecia.
sylvestre, Lam. Fl. Fr. ii. 120 = C. Intybus.

CICLOSPERMUM, Lag. Amen. Acad. Madr. ii. (1821) 101 = **Apium**, Linn. (Umbellif.).
Ammi, Lag. l. c. = A. Ammi.

CICONIUM, Sweet, Geran. i. p. ix (1822) = **Pelargonium**, L'Hérit. (Geraniac.).
acetosellum, Hoffmgg. Verz. Pfl. 48 = P. acetosellum.
acetosum, Hoffmgg. l. c. = P. acetosum.
aequaliflorum, Sweet, ex Steud. Nom. ed. II. i. 362 = P. aequaliflorum.
albidum, Sweet, l. c. = P. albidum.
amabile, Hoffmgg. Verz. Pfl. 48 = P. amabile.
Bentinckianum, Sweet, Hort. Brit. ed. I. 85 = P. Bentinkianum.
Bluecherianum, Hoffmgg. Verz. Pfl. 48 = P. Bluecherianum.
bracteosum, Sweet, Hort. Brit. ed. I. 85 = P. bracteosum.
cerinum, Sweet, Geran. ii. t. 176 = P. inquinans.
clarum, Hoffmgg. Verz. Pfl. 48 = P. zonale.
coccineum, Hoffmgg. l. c. = P. hybridum.
coccini florum, Hoffmgg. l. c. = P. zonale.
crenatum, Hoffmgg. l. c. = P. zonale.
crenatum, Hoffmgg. l. c.; Sweet, Hort. Brit. ed. I. 85 = P. crenatum.
densiflorum, Eckl. & Zeyh. Enum. 75 = P. zonale.
Fothergillii, Sweet, Geran. iii. t. 226 = P. Fothergillii.
fulgens, Hoffmgg. Verz. Pfl. 48 = P. formosissimum.
fulgens, Sweet, Hort. Brit. ed. I. Add. 471 = P. fulgens.
glabrifolium, Sweet, Geran. iv. t. 363 = P. glabrifolium.
heterogamum, Sweet, Hort. Brit. ed. I. 84 = P. heterogamum.
hilare, Hoffmgg. Verz. Pfl. 49 = P. hilare.
homogamum, Hoffmgg. l. c. = P. heterogamum.
humile, Sweet, Hort. Brit. ed. I. 84 = P. scandens.
hybridum, Sweet, Geran. i. Ind. = P. hybridum.
inquinans, Hoffmgg. Verz. Pfl. 49 = P. inquinans.
lanuginosum, Hoffmgg. l. c. = P. lanuginosum.
lateritium, Hoffmgg. l. c. = P. malvaefolium.
lavateraefolium, Eckl. & Zeyh. Enum. 75 = P. lavateraefolium.
leptopetalum, Sweet, Hort. Brit. ed. I. 84 = P. leptopetalum.
leucanthum, Hoffmgg. Verz. Pfl. 49 = P. zonale.
lucidum, Hoffmgg. l. c. = P. lucidum.
malvaefolium, Sweet, Hort. Brit. ed. I. 84 = P. malvaefolium.
marginatum, Hoffmgg. Verz. Pfl. 49 = P. marginatum.
micranthum, Sweet, Geran. iii. t. 195 = P. micranthum.
monstrum, Sweet, Geran. i. Ind. = P. monstrum.
multiflorum, Hoffmgg. Verz. Pfl. 49 = P. multiflorum.
orbiculare, Hoffmgg. l. c. = P. obtusifolium.
oxyphyllum, Sweet, Hort. Brit. ed. III. 120 = P. hybridum.
placidum, Hoffmgg. Verz. Pfl. 49 = P. amoenum.
Porphyrio, Hoffmgg. l. c. 50 = P. Porphyrio.
propugnatum, Hoffmgg. l. c. = P. fulminans.
pumilum, Hoffmgg. l. c. = P. scandens.
puniceum, Hoffmgg. l. c. = P. puniceum.
purpuriflorum, Hoffmgg. l. c. = P. purpureum.
reticulatum, Sweet, Geran. ii. t. 143 = P. retinervium.
Rosula, Hoffmgg. Verz. Pfl. 50 = P. zonale.
scandens, Hoffmgg. l. c. = P. scandens.
siccifolium, Hoffmgg. l. c. = P. siccifolium.
Skelii, Hoffmgg. l. c. = P. Skelii.
stenopetalum, Hoffmgg. l. c. = P. zonale.
umbellatum, Hoffmgg. l. c. = P. umbellatum.
urenaefolium, Eckl. & Zeyh. Enum. 75 = P. urenaefolium.
velutinum, Hoffmgg. Verz. Pfl. 50 = P. ovale.
zonale, Hoffmgg. l. c. = P. zonale.

CICUTA, [Tourn.] Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 889.
KERASKOMION, Rafin. New Fl. Am. iv. 21 (1836).
africana, Lam. Encyc. ii. 4 = Capnophyllum africanum.
Amomum, Crantz, Cl. Umb. Emend. 96 = Sison Amomum.
angustifolia, Kit. in Schult. Oester. Fl. ed. II. i. 515 = virosa.
Bolanderi, *A. Gray, in Proc. Am. Acad.* xi. (1876) 139.—Am. bor. occ.
bulbifera, *Linn. Sp. Pl.* 255.—Am. bor.
californica, *A. Gray, in Proc. Am. Acad.* vii. (1868) 344.—Calif.
capillacea, Nutt. ex DC. Prod. iv. 107 = Discopleura Nuttallii.
cynapicum, Crantz, Cl. Umb. Emend. 98 = Aethusa Cynapium.
Cynapium, Targ. Ist. Bot. ii. 227 = Aethusa Cynapium.
cellulosa, Gilib. Fl. Lituan. ii. 36 = virosa.
dahurica, Fisch. Hort. Gorenk. ed. II. (1812) 45, nomen = Apium lineare.
dichotoma, Poir. Encyc. Suppl. ii. 261 = Capnophyllum dichotomum.
latifolia, Crantz, Cl. Umb. Emend. 97 = Sium latifolium.
maculata, Hook. & Arn. Bot. Beech. Voy. 142 = californica?
maculata, Lam. Fl. Fr. iii. 104 = Conium maculatum.
maculata, *Linn. Sp. Pl.* 256.—Am. bor.
major, Lam. Fl. Fr. iii. 456 = Conium maculatum.
Ninci, Crantz, Cl. Umb. Emend. 97 = Sium Ninsi.
nipponica, *Franch. in Bull. Soc. Bot. Fr.* xxvi. (1879) 85.—Japon.
nodiflora, Crantz, Cl. Umb. Emend. 97 = Apium nodiflorum.
occidentalis, *Dougl. ex Heynh. Nom.* ii. 144, nomen.—Am. bor.
officinalis, Crantz, Cl. Umb. Emend. 98 = Conium maculatum.
perennans, Walt. Fl. Carol. 116 = Cryptotaenia canadensis?
rigens, Lam. Encyc. ii. 4 (= *Bunium rigens*).—Afr. austr.
sicula, Crantz, Cl. Umb. Emend. 97 = Kundmannia sicula.
sinensis, Zuccagni, in Roem. Collect. 135 = Selinum Monnieri.
suffruticosa, Poir. Encyc. Suppl. ii. 261 = Capnophyllum Jacquini.
tenuifolia, Schrank, in Denkschr. Akad. Muench. vii. (1818?) 56. t. 4. f. 1 = virosa.
trachypleura, *S. Wats. Bibliog. Ind. N. Am. Bot.* 417.—Am. bor. occ.
venenata, Nutt. ex Steud. Nom. ed. II. i. 362 = Archangelica hirsuta.
venenosa, Greenw. in Trans. Am. Phil. Soc. iii. (1793) 234 = Archangelica hirsuta.
virosa, *Linn. Sp. Pl.* 255.—Europ.; As. bor.; Am. bor.

CICUTARIA, Moench, Meth. Suppl. 32 (1802) (Umbellif.).
dioica, Moench, l. c. = Trinia vulgaris.
verticillata, Moench, l. c. = Molopospermum cicutarium.

CICUTARIA, [Tourn.] Lam. Fl. Fr. iii. 445 (1778) = **Cicuta**, Tourn. (Umbellif.).
aquatica, Lam. Fl. Fr. iii. 445 = Cicuta virosa.
bulbifera, Lam. Encyc. ii. 3 = Cicuta bulbifera.
maculata, Lam. l. c. 2 = Cicuta maculata.
virosa, Delarb. Fl. Auv. ed. II. 415 = Cicuta virosa.

CIECA, Adans. Fam. ii. 355 (1763) = **Julocroton**, Mart. (Euphorb.) (J. argenteus).

CIECA, Medic. Malv. 97 (1787) = **Passiflora**, Linn.
angustifolia, M. Roem. Syn. Pepon. 143 = P. suberosa.
appendiculata, M. Roem. l. c. 145 = P. auriculata.
auriculata, M. Roem. l. c. 143 = P. auriculata.
bauhinaefolia, M. Roem. l. c. 145 = P. bauhinifolia.
Berteriana, M. Roem. l. c. = P. Berteriana.
bilobata, M. Roem. l. c. 146 = P. bilobata.
Cavanillesii, M. Roem. l. c. 140 = P. cuprea.
cinerea, M. Roem. l. c. 148 = P. cinerea.

CIECA:—
colubrina, M. Roem. Syn. Pepon. 143 = Passiflora triloba.
coriacea, M. Roem. l. c. 148 = P. coriacea.
cuprea, M. Roem. l. c. 139 = P. cuprea.
Dictamo, M. Roem. l. c. 146 = P. Dictamo.
difformis, M. Roem. l. c. 140 = P. coriacea.
discolor, M. Roem. l. c. = P. Maximiliana.
flexuosa, M. Roem. l. c. 148 = P. suberosa.
glabrata, M. Roem. l. c. 143 = P. lunata.
globosa, M. Roem. l. c. 144 = P. suberosa.
gracilis, M. Roem. l. c. 141 = P. gracilis.
guyanensis, Klotzsch, in Schomb. Fauna et Fl. Gui. 986, nomen.—Guian.
hedcracca, Klotzsch, l. c. 1090, nomen.—Guian.
hedcracca, M. Roem. Syn. Pepon. 141 = P. suberosa.
hcterophylla, Moench, Meth. Suppl. 101 = P. angustifolia.
limbata, M. Roem. Syn. Pepon. 148 = P. limbata.
littoralis, M. Roem. l. c. 145 = P. suberosa.
maculata, M. Roem. l. c. = P. maculata.
membranacea, M. Roem. l. c. 140 = P. membranacea.
mexicana, M. Roem. l. c. 146 = P. mexicana.
minima, Moench, Meth. Suppl. 102 = P. minima.
misera, M. Roem. Syn. Pepon. 140 = P. misera.
multiflora, M. Roem. l. c. 148 = P. multiflora.
nigra, Medic. Malv. 97 = P. suberosa.
normalis, M. Roem. Syn. Pepon. 144 = P. perfoliata.
olivaeformis, M. Roem. l. c. = P. suberosa.
pallida, M. Roem. l. c. 142 = P. subcrosa.
pannosa, M. Roem. l. c. 148 = P. pannosa.
peltata, M. Roem. l. c. 141 = P. suberosa.
porophylla, M. Roem. l. c. 147 = P. porophylla.
pseudo-suberosa, M. Roem. l. c. 146 = P. suberosa.
pubescens, M. Roem. l. c. 141 = P. capsularis.
suberosa, Moench, Meth. Suppl. 102 = P. suberosa.
Sururuca, M. Roem. Syn. Pepon. 141 = P. setacea.
trisetosa, M. Roem. l. c. 147 = P. fuscinata.
variolata, M. Roem. l. c. 140 = P. variolata.
Vellozii, M. Roem. l. c. 142 = P. Vellozii.
viridis, Medic. Malv. 97 = P. minima.
Warei, M. Roem. Syn. Pepon. 146 = P. Warei.

CIENFUEGIA, Willd. Sp. Pl. iii. 723 (1800) = seq.

CIENFUEGOSIA, Cav. Diss. 174. t. 72. f. 2 (1787) = **Fugosia**, Juss. (Malvac.).
digitata, Cav. l. c. = F. digitata.
hcterophylla, Garcke, in Bonplandia, viii. (1860) 150 = F. heterophylla.
Hildebrandtii, Garcke, in Eichl. Jahrb. Bot. Gard. Berl. ii. (1883) 337.—Afr. trop.
phlomidifolia, Garcke, in Bonplandia, viii. (1860) 150 = F. phlomidifolia.
sulphurea, Garcke, l. c. = F. sulfurea.
Welshii, Garcke, in Eichl. Jahrb. Bot. Gard. Berl. ii. (1883) 337.—Arabia.

CIENFUGOSIA, DC. Prod. i. 457 (1824) = praec.

CIENKOWSKIA, Regel & Rach, in Ind. Sem. Hort. Petrop. (1858) 48. 'Gen. dub.' Benth. & Hook. f. i. 997.
nitida, Regel & Rach, l. c.—Hab. ?

CIENKOWSKIA, Solms, in Schweinf. Beitr. Fl. Aethiop. t. 1. (1867) = seq.

CIENKOWSKYA, Solms, in Schweinf. Beitr. Fl. Aethiop. 197 (1867) = **Kaempferia**, Linn. (Scitamin.).
aethiopica, Solms, l. c.—Afr. trop.
Kirkii, Hook. f. Bot. Mag. t. 5994.—Afr. trop.

CILIARIA, Haw. Saxifr. Enum. 41 (1821) = **Saxifraga**, Tourn.
aspera, Haw. l. c. 42 = S. aspera.
bronchialis, Haw. l. c. 43 = S. bronchialis.
bryoides, Haw. l. c. 42 = S. aspera.

CIMBARIA, Hill, Veg. Syst. xx. 43 (1772) = **Cymbaria**, Linn. (Scrophul.).

CIMICIFUGA, Linn. Pl. Rar. Camsch. 21 (1750). *RANUNCULACEAE*, Benth. & Hook. f. i. 9.
Actinospora, Turcz. in Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 21 (1835).
Botrophis, Rafin. Med. Fl. i. 85 (1828).

CIMICIFUGA:—
Macrotrys, Rafin. in Med. Repos. N. York. II. v. (1808) 352.
Macrotys, DC. Syst. i. 383 (1818).
Megotrys, Rafin. in Am. Monthly Mag. (1818) 267.
Pityrosperma, Sieb. & Zucc. in Abh. Acad. Muench. iii. (1846) 734.
americana, *Michx. Fl. Bor. Am.* i. 316.—Am. bor.
americana, Muhl. Cat. ed. II. 54 = cordifolia.
arizonica, *S. Wats. in Proc. Am. Acad.* xx. (1885) 352.—Arizona.
biternata, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 9.—Japon.
cordifolia, *Pursh, Fl. Am. Sept.* ii. 373.—Am. bor.
elata, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 36.—Am. bor. occ.
foetida, *Linn. Syst.* ed. XII. 659.—Europ.; Sibir.
foetida, Pursh, Fl. Am. Sept. ii. 373 = elata.
frigida, Royle, Illustr. Bot. Himal. 57. t. 14 = foetida.
japonica, Spreng. Syst. ii. 628.—Japon.
laciniata, *S. Wats. in Proc. Am. Acad.* xx. (1885) 352.—Am. bor. occ.
obtusiloba, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 9.—Japon.
palmata, Michx. Fl. Bor. Am. i. 316 = Trautvetteria palmata.
podocarpa, Ell. Sketch, ii. 16 = americana.
racemosa, *Nutt. Gen. Am.* ii. 15.—Am. bor.
Serpentaria, Pursh, Fl. Am. Sept. ii. 372 = racemosa.
simplex, Wormsk. ex DC. Prod. i. 64 = foetida.

CIMINALIS, Adans. Fam. ii. 504 (1763) = **Gentiana**, Tourn.
acaulis, Borckh. in Roem. Arch. i. 1. 26 = G. acaulis.
algida, Bercht. & Presl, Rostl. i. Gentian. 10 = G. frigida.
decumbens, Bercht. & Presl, l. c. = G. decumbens.
frigida, Bercht. & Presl, l. c. = G. frigida.
glauca, Bercht. & Presl, l. c. 11 = G. glauca.
grandiflora, Mayrh. in Schrank, Fl. Monac. t. 37 = G. acaulis.
linearis, Bercht. & Presl, Rostl. i. Gentian. 11 = G. linearis.
longiflora, Moench, Meth. 514 = G. acaulis.
ochroleuca, Bercht. & Presl, Rostl. i. Gentian. 11 = G. ochroleuca.
Pneumonanthe, Borckh. in Roem. Arch. i. 1. 26 = G. Pneumonanthe.
pseudopneumonanthe, Bercht. & Presl, Rostl. i. Gentian. 11 = G. Pneumonanthe.
saponaria, Bercht. & Presl, l. c. = G. saponaria.
saxosa, Bercht. & Presl, l. c. = G. montana.
septemfida, Bercht. & Presl, l. c. 10 = G. septemfida.
vulgaris, Bercht. & Presl, l. c. = G. Pneumonanthe.

CIMINALIS, Rafin. Fl. Tellur. iii. 19 (1836) = **Voyria**, Aubl. (Gentian.).
alpina, angustifolia, grandiflora, Rafin. l. c. = V. aphylla.

CINARA, [Tourn.] Linn. Syst. ed. I. (1735) = **Cynara**, Linn. (Compos.).

CINCHONA, Linn. Gen. ed. II. 526 (1742). *RUBIACEAE*, Benth. & Hook. f. ii. 22.
Calisaya, Hort. ex Pav. Nucv. Quinol. sub t. 5. p. 6 (1862).
Cascarella, Ruiz, ex Steud. Nom. ed. II. i. 165 (1840).
Kinkina, Adans. Fam. ii. 147 (1763).
Quinquina, Condam. in 'Act. Acad. Par. (1738) 226.'
academica, Guibourt, Drogues Simples, iii. 98 = officinalis.
acuminata, Poir. Encyc. Suppl. iv. 640 = Cosmibuena acuminata.
acutifolia, Ruiz & Pav. Fl. Per. iii. 1. t. 225 = Cascarilla acutifolia.
affinis, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 8.—Peruv.
afroinda, Willem. in Usteri, Ann. Bot. xviii. (1796) 16 = Danais fragrans.
amygdalifolia, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 6.—Bolivia.
angustifolia, Ruiz, Quin. Suppl. 14. n. 17 = lancifolia.
angustifolia, Sw. Prod. Veg. Ind. Occ. 41 = Exostemma angustifolium.
asperifolia, *Wedd. in. Ann. Sc. Nat.* Sér. III. x. (1848) 7.—Bolivia.

CINCHONA:—
australis, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma australe.
australis, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 7.—Bolivia.
barbacoënsis, *Karst. in C. Koch & Fint. Wochensch.* ii. (1859) 31.—N. Granat.
Bergeniana, A. Braun, ex Mart. Reise Bras. iii. 1286 = Remijia Bergeniana.
bicolor, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma bicolor.
bogotensis, *Karst. in C. Koch & Fint. Wochenschr.* ii. (1859) 30.—N. Granat.
boliviana, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 7 = C. Calisaya.
Bonplandiana, *Klotzsch, in Hayne, Arzneigew.* vii. t. 37.—Peruv.
brachycarpa, Sw. Prod. Veg. Ind. Occ. 41 = Exostemma brachycarpum.
brasiliensis, Hoffmgg. ex Humb. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 119 = Machaonia brasiliensis.
caduciflora, Humb. & Bonpl. Pl. Aequin. i. 168, in obs. = Cascarilla caduciflora.
Calisaya, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 6.—Bolivia.
caloptera, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. (1868–69) 273.—Peruv.
Candollii, G. Don, Gen. Syst. iii. 479 = Remijia paniculata.
canescens, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma canescens.
capensis, Burm. ex DC. Prod. iv. 369 = Burchellia capensis.
capitata, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma capitatum.
carabayensis, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 9.—Peruv.
caribaea, Jacq. Enum. Pl. Carib. 16 = Exostemma caribacum.
caribbeana, W. Wright, in Phil. Trans. lxvii. (1778) 506 = pracc.
caroliniana, Poir. Encyc. vi. 40 = Pinckneya pubens.
Carua, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 275.—Bolivia.
Cattukambar, Koen. ex Retz, in Vet. Acad. Handl. Stockh. (1814) 32. t. 3 = Uncaria acida.
cava, Pav. ex Lamb. Illustr. Cinch. 8 = Cascarilla Pavonii.
Chahuarguera, Pav. ex DC. Prod. iv. 352 = officinalis.
China, DC. Prod. iv. 356, sphalm. = Cosmibuena obtusifolia.
chlorrhiza, Bory, ex DC. l. c. 361 = Danais rotundifolia.
Chomeliana, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 9.—Bolivia.
cladorrhiza, Bory, ex Wedd. Hist. Nat. Quinq. 72 = Danais rotundifolia.
coccinea, Pav. ex DC. Prod. iv. 353 = macrocalyx.
colorata, Laubert, in 'Bull. de Pharm. (1810) 294' = officinalis.
Condaminea, Humb. & Bonpl. Pl. Aequin. i. 33 = officinalis.
conglomerata, *Pav. Nuev. Quinol.* t. 15.—Ecuador.
cordifolia, *Mutis, ex Humb. in Ges. Naturf. Fr. Berl. Mag.* i. (1807) 117.—N. Granat.; Peruv.
coriacea, Poir. Encyc. vi. 38 = Exostemma coriaceum.
coronulata, *Miq. in Journ. Bot. Neerl.* i. (1861) 140.—Peruv.
corymbifera, Forst. in Act. Nov. Soc. Sc. Ups. iii. (1780) 176 = Exostemma corymbiferum.
corymbosa, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma corymbosum.
corymbosa, Karst. Fl. Columb. i. 19. t. 10 = pitayensis.
crassifolia, Pav. ex DC. Bibl. Univ. xli. (1829) 150 = Cascarilla calyptrata.
crispa, Tafalla, ex Howard, in Pav. Nuev. Quinol. t. 2 = officinalis.
cucumaefolia, Pav. ex Lamb. Illustr. Cinch. 2 = lancifolia.
cujabensis, Manso, ex Klotzsch, in Hayne, Arzneigew. xiv. sub t. 15 = Remijia cujabensis.
cuspidata, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma cuspidatum.
decurrentifolia, Pav. Nuev. Quinol. t. 23 = purpurascens.
Delondriana, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 7 = scrobiculata.

CINCHONA:—
densiflora, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Remijia densiflora.
dichotoma, Ruiz & Pav. Fl. Per. ii. 53. t. 197 = Ladenbergia dichotoma.
discolor, Klotzsch, in Hayne, Arzneigew. xiv. sub t. 14 = pubescens.
dissimiliflora, Mutis, ex Humb. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 120 = Exostemma dissimiliflorum.
elliptica, *Wedd. in Ann. Sc. Nat.* Sér. V. xii. (1869) 60.—Peruv.
erythrantha, *Pav. Nuev. Quinol.* t. 12.—Ecuador.
erythroderma, *Wedd. in Ann. Sc. Nat.* Sér. V. xi. (1869) 362.—Peruv.
euneura, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 265.—Bras.
excelsa, Roxb. Pl. Corom. ii. 3. t. 106 = Hymenodictyon excelsum.
ferruginea, St. Hil. in Mém. Mus. Par. ix. (1822) 314 = Remijia ferruginea.
firmula, Mart. ex Klotzsch, in Hayne, Arzneigew. xiv. sub t. 15 = Remijia firmula.
flaccida, Spreng. Syst. iv. Cur. Post. 73 = Hymenodictyon flaccidum.
floribunda, Sw. Prod. Veg. Ind. Occ. 41 = Exostemma floribundum.
Forbesiana, Howard, Quinol. E. Ind. Plant. 37* = lancifolia.
formosa, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma formosum.
fusca, Ruiz, ex Vitm. Summa, Suppl. i. 262 (= *Lasionema roseum*).—Am. trop.
glabra, Ruiz, Quinol. 64 = lancifolia.
glandulifera, *Ruiz & Pav. Fl. Per.* iii. 1. t. 224.—Peruv.
globifera, Pav. ex DC. Prod. iv. 345 = Nauclea Cinchonae.
govana, *Miq. in Journ. Bot. Néerl.* i. (1861) 142.—Peruv.
grandiflora, Ruiz & Pav. Fl. Per. ii. 54 = Cosmibnena obtusifolia.
grandifolia, Poir. Encyc. vi. 36 = Cascarilla magnifolia.
gratissima, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 154 = Luculia gratissima.
Haenkeana, Bartl. ex DC. Prod. iv. 530 = Palicourea Haenkeana.
Hasskarliana, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 266.—Peruv.
Henleana, *Karst. in C. Koch & Fint. Wochenschr.* ii. (1859) 31.—Venezucla.
herbacea, Linn. Mant. ii. 338 = Exostemma caribaeum ?
heterocarpa, *Karst. Fl. Columb.* i. 11. t. 6.—N. Granat.
heterophylla, Pav. ex DC. Prod. iv. 353 = cordifolia.
heterophylla, Pav. Nuev. Quinol. t. 18 = macrocalyx.
hexandra, G. Don, Gen. Syst. iii. 478 = Cascarilla hexandra.
hirsuta, *Ruiz & Pav. Fl. Per.* ii. 51. t. 192.—Peruv.
Howardiana, *O. Kuntze, Monog. Cinch.* (1878). 30.—Peruv.
Humboldtiana, *Lamb. Illustr. Cinch.* 7.—Peruv.
Humboldtiana, Roem. & Schult. Syst. v. 13 (= *Lasionema Humboldtianum*).—Peruv.
Humboldtiana, Wedd. Hist. Nat. Quinq. 67 = villosa.
jamaicensis, W. Wright, in Phil. Trans. lxvii. (1778) 506 = Exostemma caribaeum.
Josephiana, *Wedd. in Ann. Sc. Nat.* Sér. V. xii. (1869) 58.—Bolivia.
Kattukambar, Koen. ex DC. Prod. iv. 347 = Uncaria acida.
laccifera, Ruiz & Pav. ex Fée, Ess. Crypt. Ecorces Exot. Off. 8; Tafalla, ex Wedd. Hist. Nat. Quinq. 72 = Condaminea tinctoria ?
Lambertiana, A. Braun, ex Mart. Reise Bras. iii. 1286 = Cascarilla Lambertiana.
lanceolata, Ruiz & Pav. Fl. Per. ii. 51 = officinalis.
lancifolia, *Mutis, in Periodico de Santa Fé* (1793) 465, nomen.—Peruv.; N. Granat.
Lechleriana, *Schlecht. in Linnaea*, xxvi. (1853-55) 728.—Peruv.
Ledgeriana, *Moens, ex Trimen, in Journ. Bot.* xix. (1881) 323.—Bolivia.
legitima, *Ruiz, ex Laubert, in Bull. de Pharm.* (1810) 293.—Peruv. ?

CINCHONA:—
leucocalyx, *Brign. in Mem. Soc. Ital. Mod.* Ser. II. i. (1862) 64.—Bras.
lineata, Vahl, in Skriv. Naturh.-Selsk. i. (1790) 22 = Exostemma lineatum.
longiflora, Lamb. Desc. Cinch. 38. t. 12 = Exostemma longiflorum.
longiflora, Mutis, ex Steud. Nom. ed. I. 196 = Cosmibuena obtusifolia.
longifolia, Mutis, ex Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 62 = praec. ?
Luciana, Vitm. Summa, Suppl. i. 264 = Exostemma floribundum.
lucumaefolia, Pav. ex Lindl. Fl. Med. 416 = macrocalyx.
lutea, Pav. Nuev. Quinol. t. 14 = cordifolia.
lutescens, Ruiz, ex Vitm. Summa, Suppl. i. 262 = Cascarilla magnifolia.
macrocalyx, *Pav. ex DC. in Bibl. Univ. Sc. et Arts*, xli. (1829) 150.—Am. austr.
macrocarpa, Vahl, in Skriv. Naturh.-Selsk. i. (1790) 20 = Cascarilla macrocarpa.
Macrocnemia, A. Braun, ex Mart. Reise Bras. iii. 1286 = Remijia Macrocnemia.
macrophylla, *Karst. in Linnaea*, xxx. (1859-60) 151.—N. Granat.
magniflora, Pav. ex DC. Prod. iv. 354 = Cascarilla macrocarpa.
magnifolia, Humb. & Bonpl. Pl. Aequin. i. 136 = Cascarilla caduciflora.
magnifolia, Ruiz & Pav. Fl. Per. ii. 53. t. 196 = Cascarilla magnifolia.
maynensis, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma maynense.
micrantha, *Ruiz & Pav. Fl. Per.* ii. 52. t. 194.—Peruv.
microphylla, Mutis, ex Lamb. Illustr. Cinch. 9 = Mutisii.
microphylla, *Pav. Nuev. Quinol.* t. 17.—Peruv.
montana, Badier, in Journ. Phys. (1789) 129 = Exostemma floribundum.
Morado, Ruiz, Quinol. 67. n. 5 = purpurea.
Moritziana, *Karst. ex Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 275.—Peruv.
Mutisii, *Lamb. Illustr. Cinch.* 9.—Pernv.
muzonensis, Goudot, in Phil. Mag. (1828) 132 = Cascarilla muzonensis.
myrtifolia, Stokes, Bot. Mat. Med. i. 359 = Exostemma caribaeum.
nitida, Benth. Pl. Hartw. 109 = Cascarilla nitida.
nitida, *Ruiz & Pav. Fl. Per.* ii. 50.—Peruv.
oblongifolia, Lamb. Illustr. Cinch. 12 = Buena Riveroana.
oblongifolia, Mutis, in Periodico de Santa Fé (1793) 465, nomen; Humb. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 118 = Cascarilla oblongifolia.
obovata, Pav. Nuev. Quinol. sub t. 18. et t. 24 = pubescens.
obovata, Spreng. Syst. iv. Cur. Post. 73 = Hymenodictyon obovatum.
obtusifolia, D. Dietr. Syn. Pl. i. 721 = Cosmibuena obtusifolia.
obtusifolia, Pav. ex DC. Prod. iv. 353 = macrocalyx.
officinalis, Linn. Mant. ii. 338 = cordifolia.
officinalis, *Linn. Syst.* ed. X. 929.—Peruv.
ovalifolia, Humb. & Bonpl. Pl. Aequin. i. 65 (= *Lasionema Humboldtianum*).—Peruv.
ovalifolia, Mutis, ex Humb. in Ges. Naturf. Fr. Berl. Mag. i. (1807) 118 = Cascarilla macrocarpa.
ovata, *Ruiz & Pav. Fl. Per.* ii. 52. t. 195.—Peruv.
Pahudiana, *Howard, in Pav. Nuev. Quinol.* t. 21.—Am. austr.
pallescens, Ruiz, ex DC. Prod. iv. 353 = ovata.
Palton, Pav. Nuev. Quinol. t. 13 = macrocalyx.
paniculata, D. Dietr. Syn. Pl. i. 721 = Remijia paniculata.
parabolica, *Pav. Nuev. Quinol.* t. 16.—Peruv.
parviflora, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma parviflorum.
parviflora, Poir. Encyc. vi. 38 = micrantha.
pauciflora, *Tafalla, ex Steud. Nom.* ed. II. i. 363.—Am. austr.
Pavoniana, *O. Kuntze, Monog. Cinch.* (1878) 29.—Peruv.; Bolivia.
Pavonii, Lamb. Illnstr. Cinch. 8 = Cascarilla Pavonii.
pedunculata, *Karst. in C. Koch & Fint. Wochenschr.* ii. (1859) 30.—N. Granat.
Pelalba, *Pav. ex DC. Bibl. Univ.* xli. (1829) 152.—Am. austr.

CINCHONA:—
Pelletieriana, Wedd. in Ann. Sc. Nat. Sér. III. x. (1848) 8 = pubescens.
peruviana, *Howard, in Pav. Nuev. Quinol.* t. 27.—Andes.
peruviana, Poir. Encyc. Suppl. iv. 640 = Exostemma peruvianum.
philippica, Cav. Ic. iv. 15 = Exostemma philippicum.
pitayensis, *Wedd. in Ann. Sc. Nat.* Sér. III. xi. (1849) 269; et V. xii. (1862) 37.—N. Granat.
platyphylla, *Wedd. l. c.* Sér. V. xi. (1869) 362; xii. (1869) 73.—N. Granat.
prismatostylis, *Karst. Fl. Columb.* i. 13. t. 7.—N. Granat.
pubescens, Hort. Mack. ex Walp. Rep. ii. 943 = Rhyssocarpus pubescens.
pubescens, Lamb. Illustr. Cinch. 6 = ovata.
pubescens, Vahl, in Skriv. Naturh.-Selsk. i. (1790) 19 = cordifolia.
purpurascens, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 8.—Bolivia.
purpurea, *Ruiz & Pav. Fl. Per.* ii. 52. t. 193.—Peruv.
quercifolia, Pav. ex Lamb. Illustr. Cinch. 9 = Mutisii.
Quina, Ruiz, in Ruiz & Pav. Fl. Per. iii. 4 = Cosmibuena obtusifolia.
racemosa, Schrank, ex Steud. Nom. ed. II. i. 363 = Exostemma caribaeum.
Reicheliana, Howard, in Pav. Nuev. Quinol. t. 5 = micrantha.
Remijiana, St. Hil. Pl. Us. Bras. sub t. 2 = Remijia Hilarii.
Riedeliana, Casar. in Atti 3zia Riun. Scienz. Ital. (1841) 513 = Cascarilla Riedeliana.
robusta, *Howard, in Journ. Linn. Soc.* xx. (1884) 327, nomen.—Hab. ?
Roraimae, Benth. in Hook. Journ. Bot. iii. (1841) 214 = Cascarilla Roraimae.
rosea, Ruiz & Pav. Fl. Per. ii. 54. t. 199 (= *Lasionema roseum*).—Peruv.
rosulenta, *Howard, ex Wedd. in Ann. Sc. Nat.* Sér. V. xi. (1869) 362.—N. Granat.
rotundifolia, Pav. ex Lamb. Illustr. Cinch. 5 = cordifolia.
rubicunda, *Auct. ex Steud. Nom.* ed. II. i. 363.—Am. austr.
rubicunda, Tafalla, ex Wedd. in Ann. Sc. Nat. Sér. V. xi. (1869) 362 = ovata.
rufinervis, *Wedd. in Ann. Sc. Nat.* Sér. III. x. (1848) 8.—Peruv.
rugosa, Pav. ex DC. Prod. iv. 353 = cordifolia.
Sanctae-Luciae, Davids, ex Roem. & Schult. Syst. v. 19 = Exostemma floribundum.
scrobiculata, *Humb. & Bonpl. Pl. Aequin.* i. 165. t. 47.—Peruv.
souzana, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Exostemma souzanum.
spinosa, Vavass. in Journ. Phys. (Oct. 1790) 273 = Exostemma Vavassorii.
stenocarpa, Lamb. Illustr. Cinch. 13 = Cascarilla stenocarpa.
stupea, Pav. ex Lindl. Fl. Med. 416 = lucumaefolia.
subcordata, Pav. Nuev. Quinol. ed. Howard, ad calcem = cordifolia.
suberosa, Pav. l. c. = macrocalyx.
subsessilis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 272.—Peruv.
succirubra, *Pav. ex Klotzsch, in Abh. Akad. Berl.* 1857 (1858) 60.—Peruv.
Tarontaron, *Pav. Nuev. Quinol. ed. Howard*, ad calcem.—Peruv.
tenuiflora, Brign. in Mem. Soc. Ital. Mod. Ser. II. i. (1862) 63 = Remijia tenuiflora.
tenuis, Rniz, ex DC. Prod. iv. 353 = cordifolia.
thyrsiflora, Roxb. Hort. Beng. 15; Fl. Ind. i. 530 = Hymenodictyon excelsum.
timorana, Span. in Linnaca, xv. (1841) 315 = Hymenodictyon timoriense.
timorensis, Walp. Rep. ii. 509 = praec.
Trianae, Karst. in C. Koch & Fint. Wochenschr. ii. (1859) 31 = pitayensis.
triflora, W. Wright, in Lond. Med. Journ. viii. (1787) 240 = Exostemma triflorum.
tucujensis, *Karst. Fl. Columb.* i. 17. t. 9.—N. Granat.
nmbellulifera, *Pav. Nuev. Quinol.* t. 22.—Peruv.
undata, Karst. ex Wedd. in Journ. Linn. Soc. xi. (1869) 187 = Cascarilla undata.
undulata, Pav. Nuev. Quinol. 2. t. 6. = glandulifera.
Uritusinga, Pav. ex DC. Prod. iv. 353 = macrocalyx.

CINCHONA:—
Uritusinga, Pav. ex Howard, Nuev. Quinol. t. 19 = officinalis.
vanilliodora, *Auct. ex Steud. Nom.* ed. II. i. 363.—Am. austr.
Vellozii, St. Hil. Pl. Us. Bras. sub t. 2 = Remijia Vellozii.
villosa, Pav. ex Lamb. Illustr. Cinch. 7 = Humboldtiana.
violacea, *Pav. Nuev. Quinol. ed. Howard, ad calcem.*—Ecuador.
viridiflora, Pav. l. c. = pubescens.
Vritusino, Pav. ex DC. Prod. iv. 352 = C. Condaminca.
Weddelliana, *O. Kuntze, Monog. Cinch.* (1878) 29.—Bolivia.

CINCLIA, Hoffmgg. Preisverz. (1841).—Cf. Linnaea, xvi. (1842) Litt. 267 = **Ceropegia**, Linn. (Asclepiad.).

CINCLIDOCARPUS, Zoll. & Mor. in Nat. en Geneesk. Arch. Neerl. Indie, iii. (1846) 81 = **Caesalpinia**, Linn. (Legumin.).
nitidus, Zoll. & Mor. l. c. 82 = Caesalpinia Cinclidocarpa.

CINERARIA, Linn. Sp. Pl. ed. II. 1242 (1763). *COMPOSITAE*, Benth. & Hook. f. ii. 445.
Xenocarpus, Cass. in Dict. Sc. Nat. lix. 108 (1829).
abrotanifolia, *Berg. Desc. Pl. Cap.* 292.—Afr. austr.
abyssinica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 433.—Afr. trop.
acanthifolia, Reichb. Hort. Bot. t. 24 = Senecio Cineraria.
acutangula, Bertol. Fl. Guatimal. 35 = Senecio acutangulus.
Aitoniana, Spreng. Fl. Hal. Mant. 51 = canescens.
alata, Linn. f. Suppl. 374 = Othonna alata.
alchemilloides, *DC. Prod.* vi. 307.—Afr. austr.
alliarifolia, *Desf. Cat. Hort. Par.* ed. III. 403.—Hab.?
alpestris, Hoppe, Taschenb. (1806) 130 = Senecio alpestris.
alpestris, Pall. Reise, ii. 28 = Senecio campestris.
alpestris, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 111 = Senecio brachychaetus.
alpina, All. Fl. Pedem. i. 203. t. 38 = Senecio aurantiacus.
alpina, Huds. Fl. Angl. ed. II. 370 = Senecio campestris.
alpina, Linn. Sp. Pl. ed. II. 1243 = Senecio alpinus.
altaica, C. A. Mey. ex DC. Prod. vi. 315 = Senecio altaicus.
ambigua, Bivona, Manip. iii. 5 = Senecio ambiguus.
ambracea, Hornem. Hort. Hafn. Add. 972 = Senecio ambraceus.
amelloides, Linn. Sp. Pl. ed. II. 1245 = Aster rotundifolius.
americana, Linn. f. Suppl. 373 = Senecio denticulatus.
amoena, Salisb. Prod. 200 = Aster rotundifolius.
angulata, Alam. ex DC. Prod. vi. 431. n. 544 = Senecio sublobatus.
angulata, Mair, ex DC. l. e. n. 547 = Senecio angulifolius.
angulosa, Lam. Encyc. ii. 6 = humifusa, geifolia.
angulosa, E. Mey. ex DC. Prod. vi. 337 = Senecio longiflorus.
angustata, Schur, Enum. Pl. Transs. 346 = *transsilvanica*.
angustifolia, H. B. & K. Nov. Gen. et Sp. iv. 189 = Senecio salignus.
anomala, Sibth. & Sm. Fl. Graec. Prod. ii. 179; Fl. Graec. ix. 56 = Senecio Othonnae.
anthemoides, Lam. Encyc. ii. 9 = Senecio gallicus.
anthemoides, Spreng. ex DC. Prod. vi. 365 = Senecio ramosus.
appendiculata, Poir. Encyc. Suppl. ii. 263 = Senecio appendiculatus.
appendiculata, Steud. Nom. ed. II. i. 363 = Senecio populifolius.
arachnoidea, Reichb. in Moessl. Handb. ed. II. ii. 1501 = Senecio lanatus.
arachnoidea, Turcz. ex Ledeb. Fl. Ross. ii. 632 = Senecio frigidus.
arctotidea, *DC. Prod.* vi. 306.—Afr. austr.
arctotidea, Drège, ex Harv. & Sond. Fl. Cap. iii. 309 = mollis.
arvernensis, Rouy, in Bull. Soc. Bot. Fr. xxviii. (1880) 68 = Senecio campestris.

CINERARIA:—
aspera, *Thunb. Prod. Pl. Cap.* 155.—Afr. austr.
atriplicifolia, *DC. Prod.* vi. 308.—Afr. austr.
atropurpurea, Ledeb. in Mém. Acad. Pétersb. v. (1814) 574 = Senecio frigidus.
Aucheri, *C. Koch, in Linnaea*, xxiv. (1851) 358.—Oriens.
aurantiaca, Bess. Enum. Pl. Volh. 33 = Senecio campestris.
aurantiaca, Hoppe, ex Willd. Sp. Pl. iii. 2081 = Senecio aurantiacus.
aurea, Linn. Sp. Pl. ed. II. 1244 = Senecio auratus.
auriculata, Ledeb. Fl. Alt. iv. 105 = Senecio racemosus.
aurita, Andr. Bot. Rep. i. 24. t. 24 = Doronicum cruentum.
aurita, L'Hérit. Sert. Angl. 26 = Senecio maderensis.
Balbisiana, *Bertol. Fl. Ital.* ix. 290.—Italia.
Balsamita, Lam. Encyc. ii. 7 = Senecio aureus.
belbeisia, Spreng. Syst. iii. 552 = Senecio belbeysius.
Bergeriana, Spreng. l. c. 551 = Aster Bergerianus.
bicolor, Willd. Sp. Pl. iii. 2085 = Senecio bicolor.
bipinnata, Willd. l. c. 2075 = Senecio pinnatifidus.
bracteata, Chr. Sm. ex Link, in Buch, Beschr. Canar. Ins. 148 = Senecio malvaefolius.
brasiliensis, Spreng. Neue Entdeck. ii. 142 = Senecio brasiliensis.
cacaliformis, Lam. Fl. Fr. ii. 124 = Senecio Ligularia.
cacalioides, Linn. f. Suppl. 374 = Othonna carnosa.
Caledoniae, Spreng. Fl. Hal. Mant. 50 = Brachyglottis Caledoniae.
calvescens, Nym. Syll. Suppl. 1 = Senecio Cineraria.
campestris, DC. Fl. Fr. iv. 164 = Senecio campestris.
campestris, Fellm. ex Ledeb. Fl. Ross. ii. 647 = Senecio aurantiacus.
campestris, Zett. ex Willk. & Lange, Prod. Fl. Hisp. ii. 110 = Senecio pyrenaicus.
canadensis, Linn. Sp. Pl. ed. II. 1244 = Senecio Cineraria.
canadensis, Walt. Fl. Carol. 207 = Erechtites hieracifolia?
candida, J. & C. Presl, Delic. Prag. (1822) 95 = Senecio candidus.
canescens, Spreng. Syst. iii. 549 = Erigeron canescens.
canescens, *Wendl. Obs. ex Link, Enum. Hort. Berol.* ii. 332.—Afr. austr.
cantabrica, Nym. Syll. 2 = Senecio brachychaetus.
capillacea, Linn. f. Suppl. 375 = Steirodiscus capillaceus.
capitata, Ledeb. Fl. Alt. iv. 104 = Senecio campestris.
capitata, Wahlenb. Fl. Carp. 271 = Senecio aurantiacus.
carolinensis, Walt. Fl. Carol. 207 (sp. indeterminab.).—Am. bor.
caspica, Pers. Syn. ii. 439 = Senecio racemosus.
caucasica, Bieb. Fl. Taur. Cauc. ii. 313 = Senecio caucasicus.
caucasica, Hoffm. Hort. Mosq. (1808) n. 598 = Senecio platyphyllus.
caucasica, Schlecht. ex Ledeb. Fl. Ross. ii. 858 = Senecio renifolius, *Sch. Bip.*
ceratophylla, Tenore, Ind. Sem. Hort. Neap. 1825 = Senecio Cineraria.
cernua, Thore, Chl. Land. 344 = Arnica montana.
cernua, Wall. ex DC. Prod. vi. 314 = Senecio arnicoides.
chamaedrifolia, *Lam. Encyc.* ii. 9.—Afr. austr.
chinensis, Spreng. Syst. iii. 549 = Senecio scandens.
Cinia, Webb, in Webb & Berth. Phyt. Canar. ii. 339 = Doronicum papyraceum.
Clusiana, Host, Fl. Austr. ii. 482 = Senecio alpestris.
Coderi, Nym. Consp. 352 = Senecio brachychaetus.
collina, Salisb. Prod. 200 = Senecio campestris.
conformis, Rochel, Pl. Banat. Rar. Ind. = Senecio aurantiacus, brachychactus.
congesta, R. Br. in Parry, 1st Voy. App. 279 = Senecio palustris.
cordata, Jacq. Fl. Austr. ii. t. 176 = seq.
cordifolia, Gouan, Illustr. 69 = Senecio alpinus.
cordifolia, Jacq. Fl. Austr. t. 177 = Senecio subalpinus.
cordifolia, Lapeyr. ex Willk. & Lange, Prod. Fl. Hisp. ii. 115 = Senecio Doronicum.
coronata, Thunb. Prod. Pl. Cap. 154 = Senecio lasiorhizus.
coronata, Zeyh. ex DC. Prod. vi. 438 = Senecio junceus.
corymbosa, Moench, Meth. 588 = Erigeron annuus.

CINERARIA:—
crassiflora, Poir. Encyc. Suppl. ii. 267 = Senecio crassiflorus.
crassifolia, Host, [Fl. Austr. ii. 482?] ex DC. Prod. vi. 360.—Europ.
crassifolia, Kit. in Schult. Oestr. Fl. ed. II. ii. 514 = Senecio alpestris.
crenata, Spreng. ex DC. Prod. vi. 388 = Senecio erosus.
crispa, Jacq. Fl. Austr. ii. 48 = Senecio crispatus.
crocea, Tratt. Arch. i. 26 = Senecio crispatus.
cruenta, Mass. ex L'Hérit. Sert. Angl. 26 = Doronicum cruentum.
cuneata, E. Mey. ex DC. Prod. vi. 439 = Senecio diversifolius, *Harv.*
cymbalarifolia, Linn. Amoen. Acad. vi. 106 = Senecio cymbalarifolius.
dealbata, Spreng. ex DC. Prod. vi. 369 = Senecio Kunthianus.
decipiens, *Harv. in Harv. & Sond. Fl. Cap.* iii. 312.—Afr. austr.
decipiens, Spreng. ex DC. Prod. vi. 366 = Senecio chrysanthemoides.
degenerans, Kit. ex Kanitz, in Verh. Zool.-Bot. Ges. Wien, xiii. (1863) 523 = Senecio alpestris?
deltoidea, *Sond. in Linnaea*, xxiii. (1850) 68.—Afr. austr.
denticulata, DC. Prod. vi. 416, in syn. = Senecio denticulatus.
denticulata, Linn. f. Suppl. 375 (= *Brachyrhynchos eupatorioides*).—Afr. austr.
difformis, Rochel, Pl. Banat. Rar. Ind. = Senecio alpinus.
discoidea, *Kit. in Linnaea*, xxxii. (1863) 389.—Enrop.
discolor, Sw. Prod. Veg. Ind. Occ. 114 = Senecio discolor.
distans, Kunze, in Linnaea, xvii. (1843) 571 = Senecio geifolius.
divaricata, Spreng. ex DC. Prod. vi. 370 = Senecio scandens.
Doria, Spreng. ex DC. l. e. 364 = Senecio Wallichii.
dracunculoides, Moç. & Sesse, ex DC. l. c. 430 = Senecio salignus.
Dregeana, *DC. Prod.* vi. 305.—Afr. austr.
dubia, Spreng. Syst. iii. 550 = Senecio tomentosus.
elata, Steud. Nom. ed. II. i. 364 = Senecio Ligularia.
elatior, Bouché, in Link, Enum. Hort. Berol. ii. 333 = Senecio multiflorus.
elodes, Nym. Syll. 2 = Senecio elodes.
elongata, Linn. f. Suppl. 374 = Senecio diversifolium, *Harv.*
erodioides, *DC. Prod.* vi. 307.—Afr. austr.
erosa, Harv. in Harv. & Sond. Fl. Cap. iii. 309 = oxyodonta.
erosa, *Willd. Sp. Pl.* iii. 2073.—Afr. austr.
exilis, *DC. Prod.* vi. 305.—Afr. austr.
Farfara, Bernh. Syst. Verz. Erf. 146 = Tussilago Farfara.
filifolia, Thunb. Prod. Pl. Cap. 154 = Senecio longifolius.
Fischeri, Ledeb. Ind. Sem. Hort. Dorpat. (1820) 17 = Senecio Ligularia.
flammea, Turcz. ex DC. Prod. vi. 362 = Senecio flammeus.
Forsteri, Spreng. Pugill. i. 58 = Brachyglottis Forsteri.
fragrans, Hoffmgg. ex Steud. Nom. ed. II. i. 364 = Erigeron fragrans.
frigida, Richards. in Frankl. Narr. 1st Journ. App. 748 = Senecio frigidus.
fulva, Stev. in Mém. Soc. Nat. Mosc. iv. (1816) 64 = Senecio aurantiacus.
Fussii, Griseb. & Schenk, in Wiegm. Arch. xviii. (1852) 342 = Senecio campestris.
geifolia, *Linn. Sp. Pl.* ed. II. 1242.—Afr. austr.
geifolia, C. A. Mey. ex Steud. Nom. ed. II. i. 364 = Senecio geifolius.
geniculata, Wendl. ex Steud. Nom. ed. II. i. 364 = Senecio geniculatus.
geraniifolia, *DC. Prod.* vi. 308.—Afr. austr.
gibbosa, Guss. Adnot. Cat. Pl. Hort. Boce. (1821) 3 = Senecio gibbosus.
gigantea, Sm. Exot. Bot. ii. 11 = Senecio Smithii.
glabella, Turcz. ex DC. Prod. vi. 360 = Senecio campestris.
glaberrima, Spreng. ex DC. l. c. v. 25 = Vernonia cinerea.
glabra, Sw. Prod. Veg. Ind. Occ. 113 = Senecio Swartzii.
glabrata, Ledeb. Ic. Ross. t. 94 = Senecio racemosus.

CINERARIA:—
glabrata, Sw. Fl. Ind. Occ. iii. 1354 = Senecio Swartzii.
glandulifera, Vukot. in Rad Jugos. Akad. Zagreb. li. (1880) 34 = Senecio crispatus.
glastifolia, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 148 = Senecio glastifolius.
glauca, Linn. Sp. Pl. ed. II. 1242 (= *Senecillis glauca*). —Ross. austr.
gnaphalodes, *Nym. Consp.* 351.—Europ.
grandiflora, Spreng. ex DC. Prod. vi. 314 = Senecio arnicoides.
grandiflora, Vatke, in Linnaea, xxxix. (1875) 503 = abyssinica.
Gualtata, Gill. ex Hook. & Arn. in Hook. Journ. Bot. iii. (1841) 340 = Senecio Hualtata.
hastifolia, Linn. f. Suppl. 376 = Senecio cymbalariifolius.
Heldreichii, Nym. Syll. 2 = Senecio Heldreichii.
Helenitis, Georgi, Beschr. Russ. Reich. iii. IV. 1246 = Senecio campestris.
Helenitis, Steud. Nom. ed. II. i. 364 = Senecio brachychaetus.
heterophylla, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 75 = Senecio resedifolius.
heterophylla, Orteg. Hort. Matr. 71 = Senecio minutus.
heterophylla, Pursh, Fl. Am. Sept. ii. 528 = Senecio tomentosus.
Heuffelii, Hoppe, ex Nym. Consp. 351 = Senecio lugens.
Hinniculus ×, Hort. ex Marnock, Floric. Mag. iv. (1839) t. 39.
hirsuta, Vent. Jard. Malm. 95 = Aster hirsutus.
hirta, Vahl, ex DC. Prod. v. 226 = Aster elongatns.
Hualtata, Walp. Rep. ii. 658 = Senecio Hualtata.
humifusa, *L'Hérit. Sert. Angl.* 25.—Afr. austr.
humilis, Poepp. ex DC. Prod. vi. 417 = Senecio glaber.
hybrida, Bernh. Syst. Verz. Erf. 146 = Petasites officinalis.
hybrida, Willd. Enum. Hort. Berol. 893 = Doronicum cruentum.
hypoleuca, *Reichb. ex DC. Prod.* vi. 389.—Afr. austr.
hypoleuca, Sieber, ex Harv. & Sond. Fl. Cap. iii. 381 = Senecio verbascifolius.
hypoleuca, Spreng. ex DC. l. c. 369 = Senecio rufinervis.
incana, Sw. Fl. Ind. Occ. iii. 1356 = Senecio jamaicensis.
incisa, Hontt. ex DC. Prod. vi. 438 = Senecio incertus.
incisa, Thunb. Prod. Pl. Cap. 156 = Senecio incisus.
incisa, Willd. Sp. Pl. iii. 2074 = Senecio erosus.
integrifolia, Benth. ex Willk. & Lange, Prod. Fl. Hisp. ii. 110 = Senecio pyrenaicus.
integrifolia, Hook. & Arn. in Bot. Beech. Voy. 126 = Senecio lugens.
integrifolia, Jacq. ex Willd. Sp. Pl. iii. 2082 = Senecio alpestris.
integrifolia, Pursh, Fl. Am. Sept. ii. 528 = Senecio tomentosus.
integrifolia, Richards. in Frankl. Narr. 1st Journ. ed. II. 31 = Senecio Hookeri.
integrifolia, Schkuhr, Handb. t. 246 = Senecio crispatus.
integrifolia, Vill. ex Steud. Nom. ed. II. i. 364 = Senecio aurantiacns.
integrifolia, Wallr. Sched. Crit. 474 = Senecio spathulaefolius.
integrifolia, With. Bot. Arr. ed. II. 920; Sm. Eng. Bot. t. 152 = Senecio campestris.
japonica, Thunb. Fl. Jap. 317 = Senecio japonicus.
laciniata, Sw. Fl. Ind. Occ. iii. 1352 = Senecio Swartzianus.
lactea, Jacq. f. Eclog. t. 105 = Senecio cruentus.
lactea, Willd. Enum. Hort. Berol. Suppl. 59 = Senecio appendiculatus.
laevis, Spreng. ex DC. Prod. vi. 404 = Senecio angulatus.
lanata, Jacq. Coll. iii. 177 = lanosa.
lanata, Lam. Encyc. ii. 7 = Senecio Heritieri.
lanata, Link, Ennm. Hort. Berol. ii. 333 = Doronicum cruentum.
lanceolata, Lam. Fl. Fr. ii. 125 = Senecio campestris.
lanosa, *DC. Prod.* vi. 309.—Afr. austr.
laricifolia, Lam. Encyc. ii. 8 = Senecio pinifolius.
leucanthema, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 316 = Senecio Smithii, acanthifolius.
Lewisii, Richards. in Frankl. Narr. 1st Journ. App. 748 = Erigeron compositus.

CINERARIA:—
lineata, Linn. f. Suppl. 375 = Senecio lineatus.
lineata, Vahl, ex DC. Prod. vi. 395 = Senecio triplinervius.
linifolia, Linn. Amoen. Acad. vi. 106 = Euryops linifolius.
linifolia, Zeyh. ex DC. Prod. v. 224 = Aster linifolius.
littoralis, Vest, in Flora, iii. (1820) 2 = Senecio gibbosus.
lobata, *L'Hérit. Sert. Angl.* 26.—Afr. austr.
lobata, Mair. ex DC. Prod. vi. 431 = Senecio sublobatus.
longifolia, All. Fl. Pedem. i. 203 = Senecio Balbisianus.
longifolia, Bess. Enum. Pl. Volh. 33 = Senecio alpestris.
longifolia, Eichw. Casp. Cauc. 22 = Senecio cladobotrys.
longifolia, Jacq. Fl. Anstr. ii. 49. t. 181 = Senecio brachychaetus.
longifolia, Lapeyr. Hist. Abr. Pl. Pyr. 521 = Senecio pyrenaicus.
lucida, Sw. Fl. Ind. Occ. iii. 1360 = Senecio lucidus.
lyrata, *DC. Prod.* vi. 308.—Afr. austr.
lyrata, Ledeb. in Mém. Acad. Pétersb. v. (1818) 576 = Senecio resedifolius.
lyratifolia, Bluff & Fing. Comp. ii. 377 = Senecio alpinus.
macrophylla, Ledeb. Fl. Alt. iv. 108 = Senecio Ledebourii.
malvaefolia, L'Hérit. Sert. Angl. 26 = Senecio malvaefolius.
maritima, Linn. Sp. Pl. ed. II. 925 = Senecio Cineraria.
maritima, Wulf. ex Steud. Nom. ed. II. i. 364 = Senecio gibbosus.
megapotamica, Spreng. Syst. iii. 547 = Senecio pinnatus.
microglossa, *DC. Prod.* vi. 305.—Afr. austr.
microphylla, Vahl, ex DC. l. c. v. 225 = Aster aethiopicus.
microrhiza, Schur, in Verh. Siebenb. Naturw. Ver. (1859) 147 = Senecio campestris.
minuta, Cav. Ic. i. 21. t. 33. f. 3 = Senecio minutus.
mitellaefolia, *L'Hérit. Sert. Angl.* 25.—Afr. austr.
mollis, *E. Mey. ex DC. Prod.* vi. 306.—Afr. austr.
mongolica, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 199 = Senecio mongolicus.
montevidensis, *Spreng. Syst.* iii. 548 = Senecio chilensis.
multiflora, L'Hérit. Sert. Angl. 26 = Senecio multiflorus.
nebrodensis, Guss. Adnot. Cat. Pl. Bocc. (1821) 4 = Senecio candidus.
nivea, Willd. Sp. Pl. iii. 1974 = Senecio niveus.
nuda, Kit. in Linnaea, xxxii. (1863) 389 = Senecio alpestris.
oppositifolia, Moench, Meth. 588 = Aster rotundifolius.
Othonnitis, Linn. Sp. Pl. ed. II. 1244 = Othonna frutescens.
othonnoides, *Harv. in Harv. & Sond. Fl. Cap.* iii. 314.—Afr. anstr.
ovata, Spreng. ex DC. Prod. v. 344 = Aster asperulus.
ovirensis, Koch, in Flora, vi. (1823) 507 = Senecio alpestris.
oxyodonta, *DC. Prod.* vi. 306.—Afr. austr.
palmensis, Nees, Hort. Phys. Berol. 115. t. 24 = Senecio palmensis.
palustris, Linn. Sp. Pl. ed. II. 1243 = Senecio palustris.
pandurata, Thunb. Prod. Pl. Cap. 155 = Senecio panduratus.
papposa, Reichb. in Moessl. Handb. ed. II. ii. 1500 = Senecio papposus.
Parnassi, Nym. Consp. 352 = Senecio Parnassi.
parviflora, Bieb. Fl. Taur. Cauc. ii. 316 = Senecio lampsanoides.
parviflora, [R. Br. in] Ait. Hort. Kew. ed. II. v. 72 = canescens.
pauciflora, Spreng. ex DC. Prod. vi. 367 = Senecio nudicaulis.
pedunculosa, *DC. l. c.* 305.—Afr. austr.
perfoliata, Linn. f. Suppl. 375 = Othonna perfoliata.
Petasitis, Sims, Bot. Mag. t. 1536 = Senecio Petasitis.
phylicaefolia, Poepp. ex DC. Prod. vi. 415 = Senecio phylicaefolius.
pinnata, La Llave, in La Llave & Lex. Nov. Veg. Desc. i. 29 = Senecio pinnatisectus.

CINERARIA:—
pinnatifida, Willd. Sp. Pl. iii. 2074 = Senecio incertus.
platanifolia, Schrank, Pl. Rar. Hort. Monac. t. 95 = Senecio Petasites.
platycarpa, *DC. Prod.* vi. 305.—Afr. austr.
polycephala, *DC. l. c.* 307.—Afr. austr.
polyglossa, DC. l. c. 306 = mollis.
populifolia, Lam. Encyc. ii. 8 = Senecio appendiculatus.
populifolia, L'Hérit. Sert. Angl. 25 = Senecio populifolius.
praecox, Cav. Ic. iii. 23. t. 244 = Senecio praecox.
pratensis, Herder, in Bull. Soc. Nat. Mosc. xl. (1867) i. 400, partim = Senecio lugens.
pratensis, Hoppe, ex Reichb. Fl. Germ. Excurs. ii. 241 = Senecio campestris.
procera, *Griseb. Spicil. Fl. Rumel.* ii. 219.—Rumel.
pulchella, E. Mey. ex DC. Prod. vi. 438 = Senecio cymbalariaefolius.
pulchella, *Sweet, Hort. Brit.* ed. I. 232.—Ins. Canar.
pumila, Thunb. Prod. Pl. Cap. 155 = Senecio repandus.
purpurascens, Banks & Soland. ex Hook. f. Fl. Antarct. ii. 318 = Senecio acanthifolius.
purpurata, *Linn. Mant.* 285.—Afr. austr.
pyrenaica, Nym. Syll. 2 = Senecio pyrenaicus.
racemosa, Bieb. Fl. Taur. Cauc. ii. 314 = Senecio racemosus.
ramentosa, L'Hérit. Sert. Angl. 26 = Senecio echinatus.
ramosa, Vukot. in Rad Jugos. Akad. Zagreb. li. (1880) 35.—Europ.
renifolia, C. A. Mey. Verz. Pfl. Cauc. 82 = Senecio renifolius.
reniformis, Spreng. ex DC. Prod. vi. 315 = Cremanthodium reniforme.
repanda, Lour. Fl. Cochinch. ii. 501 = Senecio scandens.
repanda, Willd. Sp. Pl. iii. 2076 = Senecio Forsteri.
retusa, Wall. ex C. B. Clarke, Comp. Ind. 206 = Senecio retusus.
rivularis, Waldst. & Kit. Pl. Rar. Hung. iii. 265 = Senecio crispatus.
robusta, Ledeb. Fl. Alt. iv. 106 = Senecio robustus.
rotundifolia, J. F. Gmel. Syst. 1240; Willd. Sp. Pl. iii. 2076 = Senecio rotundifolius.
rupicola, Nym. Consp. 351 = *transsilvanica*.
salicifolia, H. B. & K. Nov. Gen. et Sp. iv. 188 = Senecio salignus.
samnitum, Huet, ex Nym. Consp. 352 = Senecio alpinus.
sarmentosa, Lesch. ex DC. Prod. vi. 369 = Senecio candicans.
Saxifraga, *DC. l. c.* 306.—Afr. austr.
scapiflora, L'Hérit. Sert. Angl. 25 = Senecio diversifolius, *Harv.*
scapigera, Banks, ex DC. Prod. vi. 439 = Senecio diversifolius, *Harv.*
Schimperi, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 439.—Abyssin.
Schkuhrii, Reichb. in Moessl. Handb. ed. II. ii. 1499 = Senecio crispatus.
scirpina, E. Mey. ex DC. Prod. vi. 438 = Senecio junceus.
scordifolia, Link, ex DC. l. c. 439 = Senecio diversifolius, *Harv.*
Selloi, Spreng. Syst. iii. 541 = Senecio Selloi.
senecifolia, Poir. Encyc. Suppl. ii. 265 = Senecio alpinus.
Senecionis, *L'Hérit. Sert. Angl.* 26.—Hab.?
serrata, Willd. Sp. Pl. iii. 2073 = Senecio Leontodontis.
serratifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 469 = Senecio paludosus.
sessilifolia, Jacquem. Voy. Bot. 91 = Werneria nana.
sibirica, Linn. Sp. Pl. ed. II. 1242 = Senecio Ligularia.
solidaginifolia, Fisch. ex Ledeb. Fl. Ross. ii. 858 = Senecio racemosus.
sonchifolia, *Linn. Sp. Pl.* ed. II. 1243.—Afr. austr.
spathulaefolia, C. C. Gmel. Fl. Bad. iii. 454 = Senecio spathulaefolius.
spathulata, Lam. Encyc. ii. 8 = Othonna alata.
speciosa, Fisch. ex Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) II. 11 (= *Ligularia speciosa*).—Sibiria.

CINERARIA:—
speciosa, Schrad. ex Link, Enum. Hort. Berol. ii. 334 = Senecio Ligularia.
spinulosa, *Lam. Encyc.* ii. 9.—Afr. austr.
subdentata, Bunge, Enum. Pl. Chin. Bor. 39 = Senecio campestris.
sudetica, Koch, in Flora, vi. (1823) 505 = Senecio crispatus.
taraxacifolia, Bieb. Fl. Taur. Cauc. ii. 313 = Senecio taraxacifolius.
taygetea, Nym. Syll. 2 = Senecio candidus.
tenella, Link, Enum. Hort. Berol. ii. 334 = Aster tenellus.
tenuifolia, Gaudin, Fl. Helv. v. 306 = Senecio brachychaetus.
thapsoides, Nym. Syll. 2 = Senecio thapsoides.
thyrsoidea, *Host, Fl. Austr.* ii. 481.—Austria.
thyrsoidea, Ledeb. Fl. Alt. iv. 107 = Senecio sibiricus.
tillandsiaefolia, Poepp. ex DC. Prod. vi. 415 = Senecio argenteus.
tomentosa, *Less. Syn. Comp.* 391.—Afr. austr.
tomentosa, Mill. Gard. Dict. ed. VIII. n. 5 = Senecio canadensis.
trachyphylla, Spreng. Neue Entdeck. iii. 43 = Aster Ecklonis.
transsilvanica, Schur, Enum. Pl. Transs. 345 = Senecio lugens.
trifurcata, Spreng. Syst. iii. 551 = Senecio trifurcatus.
tuberosa, E. Mey. ex DC. Prod. vi. 438 = Senecio incertus.
tussilaginea, Thunb. Fl. Cap. 671 = Senecio verbascifolius.
Tussilaginis, L'Hérit. Sert. Angl. 26 = Senecio Tussilaginis.
tussilaginoides, Webb, ex Loud. Hort. Brit. Suppl. ii. 621, nomen = Senecio Tussilaginis.
uliginosa, Ledeb. in Spreng. Nov. Prov. (1818) 11 = Senecio gallicus.
uncinata, Spreng. ex DC. Prod. vi. 369 = Senecio densiflorus.
unctuosa, Gilib. Fl. Lituan. i. 211 = Senecio palustris.
undulata, Willd. Sp. Pl. iii. 2071 = Senecio petiolaris.
verna, Mair. ex DC. Prod. vi. 430 = Senecio salignus.
verticillata, Mair. ex DC. l. c. = Senecio Andrieuxii.
vestita, Haw. in Phil. Mag. (1828) 187 = Othonna vestita.
vestita, Spreng. Syst. iii. 550 = Senecio crassiflorus.
viscosa, *L'Hérit. Sert. Angl.* 25.—Afr. austr.
Visianiana, Nym. Consp. 351 = Senecio thapsoides.
vulneraria, Alam. ex DC. Prod. vi. 428 = Senecio vulnerarius.
Waterhousiana, Paxt. Mag. Bot. iv. (1838) t. 219 = Senecio Tussilaginis?
Webberiana, Paxt. l. c. ix. (1842) t. 125.—Hab. ?
Wolffii, Schur, ex Nym. Consp. 352 = Senecio campestris.
Zahlbruckneri, Host, Fl. Austr. ii. 479 = Senecio lyratifolius.

CINGA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. iv. 2 (Quid?).

CINHONA, Linn. Gen. ed. VI. 91 (1764), lapsu = **Cinchona**, Linn. (Rubiac.).

CINNA, Linn. Sp. Pl. 5 (1753). *GRAMINEAE*, Benth. & Hook. f. iii. 1151.
Abola, Adans. Fam. ii. 31 (1763).
Blyttia, Fries, Nov. Fl. Succ. Mant. ii. 2 (1839).
agrostidea, Beauv. ex Steud. Nom. ed. I. 20, 198 = arundinacea.
alba, *Nees, ex Steud. Syn. Pl. Gram.* 182.—Chili.
arachnoidea, Kunth, Rev. Gram. i. 67 = arundinacea.
arundinacea, Hook. Fl. Bor. Am. ii. 238 = Colpodium latifolium.
arundinacea, *Linn. Sp. Pl.* 5.—Reg. bor.
arundinacea, Retz. ex Steud. Nom. ed. II. i. 365 = Muehlenbergia mexicana.
Brownii, Rupr. Beitr. Pfl. Russ. Reich. ii. 66 = Colpodium latifolium.
crinita, Trin. Fund. Agrost. 118 = Dichelachne crinita.
decipiens, Kunth, Rev. Gram. i. 67 = Deyeuxia scabra.
expansa, Link, Hort. Berol. ii. 236 = arundinacea.
filiformis, Link, Enum. Hort. Berol. i. 70 = Muehlenbergia mexicana.

CINNA:—
glomerata, Link, Hort. Berol. ii. 237 = Muehlenbergia glomerata.
glomerata, Walt. Fl. Carol. 59 = Andropogon macrourus.
japonica, Steud. Syn. Pl. Gram. 182 = Sporobolus indicus.
lanata, Kunth, ex Steud. Nom. ed. II. i. 365 = Epicampes lanata.
lateralis, Walt. Fl. Carol. 59 = Andropogon virginicus.
lateriflora, Kunth, Rev. Gram. i. 67 = Muehlenbergia mexicana.
latifolia, Griseb. in Ledeb. Fl. Ross. iv. 435 = arundinacea.
macroura, Kunth, Rev. Gram. i. 67 = Epicampes macroura.
macroura, Thurb. in S. Wats. Bot. Calif. ii. 276 = Epicampes rigens.
mexicana, Beauv. Agrost. 32 = Muehlenbergia mexicana.
ovata, Kunth, Rev. Gram. i. 67 = Echinopogon ovatus.
pendula, Trin. in Mém. Acad. Pétersb. Sér. VI. vi. (1845) II. 280 = arundinacea.
phleoides, *Kunth, Rev. Gram.* i. 67.—Venezuela.
pubescens, Kunth, l. c. = Epicampes pubescens.
Purshii, Kunth, l. c. = Calamagrostis canadensis.
racemosa, Kunth, l. c. = Muehlenbergia glomerata.
setifolia, Kunth, l. c. 64; et Enum. Pl. i. 208 = Epicampes macroura.
sobolifera, Link, Enum. Hort. Berol. i. 71 = Muehlenbergia sobolifera.
stricta, Kunth, Rev. Gram. i. 67 = Epicampes macroura.
suaveolens, Rupr. ex Ledeb. Fl. Ross. iv. 435 = arundinacea.
tenuiflora, Link, Enum. Hort. Berol. i. 71 = Muehlenbergia Willdenowii.
valdiviana, *Phil. in Anal. Univ. Chil.* (1873) 563.—Chili.

CINNAGROSTIS, Griseb. in Goett. Abh. xix. (1874) 256. *GRAMINEAE*, Benth. & Hook. f. iii. 1152.
polygama, *Griseb. l. c.* 257.—Reg. Argent.

CINNAMODENDRON, Endl. Gen. 1029 (1840). *CANELLACEAE*, Benth. & Hook. f. i. 121.
axillare, *Endl. ex Walp. Rep.* i. 398.—Bras.
corticosum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. i. (1858) 351.—Ind. occ.
macranthum, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 317.—Porto Rico.
rubrum, Griseb. Fl. Brit. W. Ind. 109 = corticosum.

CINNAMOMUM, [Tourn.] Linn. Syst. ed. I. (1735); Blume, Bijdr. 568 (1825). *LAURINEAE*, Benth. & Hook. f. iii. 155.
Camphora, [Bauh.] Linn. Syst. ed. I (1735); Nees, in Wall. Pl. As. Rar. ii. 61 (1831); Syst. Laurin. 87 (1836).
Cecicodaphne, Nees, Syst. Laurin. 202 (1836).
Parthenoxylon, Blume, Mus. Bot. Lugd. Bat. i. 322 (1851).
albiflorum, Hook. f. & Thoms. ex Meissn. in DC. Prod. xv. i. 21 = impressinervium.
albiflorum, Nees, in Wall. Pl. As. Rar. ii. 75 = C. Tamala.
angustifolium, Rafin. Sylva Tellur. 135 = Acrodiclidium salicifolium?
aromaticum, J. Grah. Cat. Bomb. Pl. 173 = zeylanicum.
aromaticum, Nees, in Wall. Pl. As. Rar. ii. 74 = C. Cassia.
aromaticum, Zoll. Syst. Verz. Ind. Arch. 112 = iners.
Basania, Nees, in Wall. Pl. As. Rar. ii. 73 = obtusifolium.
Bejolghota, Sweet, Hort. Brit. ed. I. 344 = obtusifolium.
borneense, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 260; *Meissn. in DC. Prod.* xv. i. 19.—Borneo.
brevifolium, *Miq. in Verh. Nederl. Inst.* III. (1852) v. 14.—Japon.
Burmanni, *Blume, Bijdr.* 569.—Java.
calophyllum, Nees, Syst. Laurin. 40 = camphoratum.
calyculatum, *Miq. Fl. Ind. Bat. Suppl.* 359.—Sumatra.

CINNAMOMUM:—
Camphora, *T. Nees & Eberm. Handb. Med.-Pharm. Bot.* ii. 430.—China; Japon.
camphoratum, *Blume, Bijdr.* 571.—Java.
camphoriferum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Camphora.
Cappara-Coronde, Blume, Rumphia, i. 34. t. 9. f. 2, 3 = iners.
Cassia, *Blume, Bijdr.* 570.—Burma; China.
Cassia, D. Don, Prod. Fl. Nep. 67 = C. Tamala.
Cassia, Siebold, in Verh. Batav. Gen. xii. (1830) 23 = Burmanni.
Cathia, D. Don, Prod. Fl. Nep. 66 = Phoebe paniculata.
caudatum, *Nees, in Wall. Pl. As. Rar.* ii. 76.—Reg. Himal.; Burma.
Cecicodaphne, *Meissn. in DC. Prod.* xv. i. 25.—Reg. Himal.
celebicum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 264.—Ins. Celebes.
chinense, Blume, Bijdr. 569 = Burmanni.
Chloroxylon, J. S. Presl, Rostl. ii. 37 = Ceanothus Chloroxylon.
citriodorum, *Thw. Enum. Pl. Zeyl.* 253.—Zeylan.
crassinervium, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 264.—Malaya.
Culiban, Rafin. Sylva Tellur. 135 = seq.?
Culilawan, *Blume, Bijdr.* 571.—Malaya; China.
Culitlawan, Nees, in Wall. Pl. As. Rar. ii. 75 = praec.
curvifolium, *Nees, Syst. Laurin.* 80.—China.
cuspidatum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 262.—Sumatra.
cyrtopodum, *Miq. Fl. Ind. Bat.* i. 1. 897.—Sumatra.
daphnoides, *Sieb. & Zucc. in Abh. Acad. Muench.* iv. III. (1846) 202.—Japon.
dasyanthum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 259.—Sumatra.
Doederleinii, *Engl. in Engl. Bot. Jahrb.* vi. (1885) 57.—Archip. Loo-Choo.
dubium, Nees, in Wall. Pl. As. Rar. ii. 73 = iners.
dulce, Nees, l. c. 75 = Burmanni.
eucalyptoides, Nees, l. c. 73 = iners.
glabrescens, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 264.—Malaya.
glanduliferum, *Meissn. in DC. Prod.* xv. i. 25.—Reg. Himal.
glaucum, J. S. Presl, Rostl. ii. 37; G. Don, ex Loud. Hort. Brit. 160 = Litsaea glauca.
gracile, Hort. ex Sweet, Hort. Brit. ed. I. 344, nomen = C. Camphora.
gracile, Miq. Ann. Mus. Bot. Lugd. Bat. i. 259 = iners.
Griffithii, Meissn. in DC. Prod. xv. i. 19 = iners.
Heyneanum, *Nees, in Wall. Pl. As. Rar.* ii. 76.—Ind. or.
impressinervium, *Meissn. in DC. Prod.* xv. i. 21.—Reg. Himal.; China.
iners, *Reinw. ex Blume, Bijdr.* 570.—Burma; Malaya.
iners, Wall. Cat. n. 2583 E = nitidum.
iners, Wight, Ic. t. 122, bis = zeylanicum.
inodorum, *Meissn. in DC. Prod.* xv. i. 26.—Malaya.
inunctum, *Meissn. l. c.* 24.—Burma.
japonicum, Siebold, in Verh. Gen. Batav. xii. 1830 23 = pedunculatum.
javanicum, *Blume, Bijdr.* 570.—Malaya.
Kiamis, Hassk. Cat. Hort. Bogor. Alt. 87 = pedunculatum.
Kiamis, Nees, in Wall. Pl. As. Rar. ii. 75 = Burmanni.
lampongum, *Miq. Fl. Ind. Bat. Suppl.* i. 358.—Sumatra.
Laubatii, F. Muell. Fragm. v. 165 = C. Tamala.
laxiflorum, *Meissn. in DC. Prod.* xv. i. 21.—Java.
litsaeaefolium, *Thw. Enum. Pl. Zeyl.* 253.—Zeylan.
Loureirii, *Nees, Syst. Laurin.* 65.—Cochinch.
lucens, Miq. Ann. Mus. Bot. Lugd. Bat. i. 261 = tavoyanum.
macrophyllum, *Miq. l. c.* 269.—Java.
Malabathrum, Batka, in Nov. Act. Nat. Cur. xvii. (1835) II. 618 = iners.
Malabathrum, Miq. Pl. Jungh. i. 174 = subtetrapterum.
malaccense, Meissn. in DC. Prod. xv. i. 27 = C. Parthenoxylon.
Maruba, *Meissn. in DC. l. c.* 15.—Malaya.
montanum, J. S. Presl, Rostl. ii. 36; Nees, Syst. Laurin. 81 = Phoebe montana.

CINNAMOMUM :—
multiflorum, *Wight, Ic.* t. 131.—Zeylan.
Neesianum, Meissn. in DC. Prod. xv. 1. 26 = C. Parthenoxylon.
neglectum, Blume, Rumphia, i. 38, in nota, t. 11. f. 2 = javanicum.
nitidum, *Blume, l. c.* 35. t. 13. f. 2; t. 16. f. 1, 2.—Burma; Malaya.
nitidum, Hook. Exot. Fl. iii. t. 176 = iners.
obscurum, *Meissn. in DC. Prod.* xv. 1. 26.—Sumatra.
obtusifolium, *Nees, in Wall. Pl. As. Rar.* ii. 73.—Reg. Himal.; Burma.
ochraceum, Blume, Rnmphia, i. 37 = iners.
officinarum, Nees, ex Steud. Nom. ed. II. i. 366 = C. Cassia?
ovalifolium, Gardn. ex Meissn. in DC. Prod. xv. 1. 11 = Wightii.
ovalifolium, *Wight, Ic.* t. 125.—Zeylan.
paraneuron, *Miq. Fl. Ind. Bat.* i. 1. 895.—Sumatra.
Parthenoxylon, *Meissn. in DC. Prod.* xv. 1. 26.—Burma; Malaya; China.
pauciflorum, *Nees, in Wall. Pl. As. Rar.* ii. 75.—Reg. Himal.
pedatinervium, *Meissn. in DC. Prod.* xv. 1. 15.—Ins. Fiji.
pedunculatum, *J. S. Presl, Rostl.* ii. 37; *Nees, Syst. Laurin.* 79.—Japon.
Perrottetii, *Meissn. in DC. Prod.* xv. 1. 22.—Ind. or.
politum, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 265.—Malaya.
pseudo-sassafras, Meissn. in DC. Prod. xv. 1. 27 = C. Parthenoxylon.
pseudo-sintok, Miq. Fl. Ind. Bat. i. 1. 902 = iners.
pyrifolium, *D. Don, Prod. Fl. Nep.* 66 (sp. dubia).—Nepal.
Rauwolfii, Blume, Rumphia, i. 36. t. 9. f. 4, 5 = iners.
recurvatum, Wight, Ic. t. 133 = pauciflorum.
Reinwardti, Miq. Fl. Ind. Bat. i. 1. 900 = iners.
Reinwardti, Nees, Syst. Laurin. 70 = C. Tamala.
rhynchophyllum, *Miq. Fl. Ind. Bat.* i. 1. 895.—Sumatra.
rubrum, Blume, in Hoev. & De Vriese, Tijdschr. i. (1834) 62 = C. Culilawan.
sericans, *Hance, in Journ. Bot.* xv. (1877) 336.—China.
sericeum, Siebold, in Batav. Gen. Verh. xii. (1830) 24 = Laurus sericea.
Sieboldii, Meissn. in DC. Prod. xv. 1. 18 = Burmanni.
Sintok, *Blume, Bijdr.* 571.—Malaya.
subavenium, *Miq. Fl. Ind. Bat.* i. 1. 902.—Sumatra.
subcuneatum, *Miq. l. c.* 895.—Sumatra.
subtetrapterum, *Miq. l. c.* 902.—Sumatra.
sulphuratum, Kurz, For. Fl. Brit. Burm. ii. 288 = tavoyanum.
sulphuratum, *Nees, in Wall. Pl. As. Rar.* ii. 74.—Ind. or.
sumatranum, *Meissn. in DC. Prod.* xv. 1. 25.—Sumatra.
Tamala, *T. Nees & Eberm. Handb. Med.-Pharm. Bot.* ii. 426.—Reg. Himal.
tavoyanum, *Meissn. in DC. Prod.* xv. 1. 20.—Burma.
tomentosum, D. Don, Prod. Fl. Nep. 66 = Phoebe paniculata.
validinerve, *Hance, in Journ. Bot.* xx. (1882) 80.—China.
verum, J. S. Presl, Rostl. ii. 36; Sweet, Hort. Brit. ed. I. 344 = zeylanicum.
villosum, Wight, ex Meissn. in DC. Prod. xv. 1. 15 = multiflorum.
villosum, Wight, Ic. t. 127 = iners.
vimineum, *Nees, in Wall. Pl. As. Rar.* ii. 76.—Ins. Penang.
Wightii, *Meissn. in DC. Prod.* xv. 1. 11.—Ind. or.
xanthoneurum, *Blume, in Hoev. & De Vriese, Tijdschr.* i. (1834) 63.—N. Guinea.
zeylanicum, *Nees, in Wall. Pl. As. Rar.* ii. 74; iii. 32.—Ind. or.; Malaya.

CINNAMOSMA, Baill. Adansonia, vii. (1866-67) 219. *CANELLACEAE*, Benth. & Hook. f. i. 970.
fragrans, *Baill. l. c.*—Malaya.

CINNASTRUM, Fourn. Gram. Mex. 90, ex Benth. & Hook. f. Gen. iii. 1153 (1883) = **Deyeuxia**, Clar. (Gramin.).

CINOGASUM, Neck. Elem. ii. 336 (1790) = **Croton**, Linn. (Euphorbiac.).

CIONANDRA, Griseb. Fl. Brit. W. Ind. 286 (1860) = **Cayaponia**, Silva Manso (Cucurbitac.).
angustiloba, Bello, in Anal. Soc. Esp. Hist. Nat. x. (1881) 274.—Porto Rico.
cuspidata, Griseb. Fl. Brit. W. Ind. 287 = Cay. americana.
graciliflora, Griseb. l. c. = Cay. americana.
racemosa, Griseb. l. c. 286 = Cay. racemosa.

CIONISACCUS, Breda, Orch. Kuhl et Hass. (1827) [t. 8] = **Goodyera**, R. Br. (Orchid.).
lanceolatus, Breda, l. c. = G. procera.

CIONOSICYOS, Benth. & Hook. f. Gen. i. 826 (1867) = seq.

CIONOSICYS, Griseb. Fl. Brit. W. Ind. 288 (1860). *CUCURBITACEAE*, Benth. & Hook. f. i. 826.
pomiformis, *Griseb. l. c.*—Ind. occ.

CIONURA, Griseb. Spicil. Fl. Rumel. ii. 69 (1844) = **Marsdenia**, R. Br. (Asclepiad.).
erecta, Griseb. l. c. = M. erecta.

CIPADESSA, Blume, Bijdr. 162 (1825). *MELIACEAE* [Benth. & Hook. f. i. 332].
MALLEA, A. Juss. in Mém. Mus. Par. xix. (1830) 221.
baccifera, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 6 = fruticosa.
Boiviniana, *Baill. Adansonia*, xi. (1874) 255.—Madag.
borneensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 6.—Borneo.
depauperata, *Baill. Adansonia*, xi. (1874) 256.—Ins. Comor.
fruticosa, *Blume, Bijdr.* 162.—Java.
subscandens, Miq. Ann. Mus. Bot. Lugd. Bat. iv. 7 = fruticosa.

CIPONIMA, Aubl. Pl. Gui. i. 566. t. 266 (1775) = **Symplocos**, Linn. (Styrac.).
guianensis, Aubl. l. c. 567 = S. Ciponima.
scabridula, Miers, in Journ. Linn. Soc. xvii. (1879) 288 = S. Ciponima.

CIPURA, Aubl. Pl. Gui. i. 38. t. 13 (1775). *IRIDEAE*, Benth. & Hook. f. iii. 694.
MARICA, Schreb. Gen. Pl. 37 (1789) (non Ker-Gawl.).
azurea, Griseb. & Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 387 = Calydorea azurea.
caerulea, Heynh. Nom. i. 197 = Marica caerulea.
coelestis, Heynh. l. c. = Nemastylis coelestinus.
cubensis, *Griseb. Cat. Pl. Cub.* 251.—Ins. Cuba.
gracilis, Heynh. Nom. i. 197 = Marica gracilis.
graminea, H. B. & K. Nov. Gen. et Sp. i. 320 = paludosa.
humilis, H. B. & K. l. c. = palndosa.
linearis, Klotzsch & Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 387 = Calydorea cipuroides.
longifolia, Heynh. Nom. i. 197 = Marica longifolia.
martinicensis, H. B. & K. Nov. Gen. et Sp. i. 320 = Trimezia lurida.
Northiana, *Endl. in Harting. Parad. Vindob.* t. 6.—Bras.
paludosa, *Aubl. Pl. Gui.* i. 38. t. 13.—Am. trop.
plicata, Griseb. Fl. Brit. W. Ind. 589 = Eleutherine plicata.
Sabini, Heynh. Nom. i. 197 = Northiana.
Sabiniana, Voigt, Hort. Suburb. Calc. 602 = Marica Sabini.
semiaperta, Heynh. l. c. = Trimezia lurida.

CIPURA, Klotzsch, ex Klatt, in Abh. Naturf. Ges. Halle, xii. (1882) 362 = **Herbertia**, Sweet (Irid.).

CIRCAEA, Tourn. ex Linn. Syst. ed. I (1735); Gen. ed. I. 3 (1737). *ONAGRARIEAE*, Benth. & Hook. f. i. 793.
REGMUS, Dulac, Fl. Hautes-Pyr. 328 (1867).
alpestris, Schur, Enum. Pl. Transs. 214 = C. lutetiana.
alpestris, Wallr. Beitr. Fl. Hercyn. 167 = alpina.
alpina, *Linn. Sp. Pl.* 9.—Europ.; As. bor.; Am. bor.
canadensis, Muhl. Cat. 2 = C. lutetiana.
cordata, *Royle, Illustr. Bot. Himal.* 211. t. 43. f. 1.—Reg. Himal.
cordifolia, Stokes, Bot. Mat. Med. i. 26 = alpina.
decumbens, Gilib. Fl. Lituan. ii. 127 = alpina.
erecta, Gilib. l. c. 126 = C. lutetiana.

CIRCAEA :—
ericetorum, *Martr. Pl. Crit. Tarn.* 29.—Gallia.
erubescens, *Franch. & Sav. Enum. Pl. Jap.* ii. 370.—Japon.
intermedia, *Ehrh. Beitr.* iv. 42.—Europ.
intermedia, Wall. Cat. n. 6342 = alpina.
lutetiana, Georgi, It. i. 195 = alpina.
lutetiana, *Linn. Sp. Pl.* 9.—Reg. bor.
major, Lam. Fl. Fr. iii. 473 = C. lutetiana.
minima, Lam. l. c. = C. alpina.
mollis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. (1847) 134.—Japon.
nemoralis, Salisb. Prod. 276 = C. lutetiana.
ovatifolia, Stokes, Bot. Mat. Med. i. 26 = C. lntetiana.
pacifica, *Aschers. & Magnus, in Bot. Zeit.* xxix. (1871) 392.—Am. bor. occ.
pubescens, Pohl, ex Steud. Nom. ed. II. i. 366 = C. lutetiana.
quadrisulcata, *Maxim. in Franch. & Sav. Enum. Pl. Jap.* i. 169.—Japon.
repens, Wall. Cat. n. 6341 = C. lutetiana.
vulgaris, Moench, Meth. 279 = C. lutetiana.

CIRCAEASTER, Maxim. in Bull. Acad. Pétersb. xxvii. (1881) 556. *CHLORANTHACEAE*?, Benth. & Hook. f. iii. 1220.
agrestis, *Maxim. l. c.* 557.—Reg. Himal.; As. centr.

CIRCINNUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 384 = **Hymenocarpus**, Savi [1798] (Legumin.).
vulnerarioides, Medic. l. c. = H. circinnata.

CIRCINUS, Medic. Phil. Bot. i. 208 (1789) = praec.

CIRCIS, Chapm. Fl. S. U. St. 114 (1860), lapsu = **Cercis**, Linn. (Legumin.).

CIRINOSUM, Neck. Elem. ii. 84 (1790) = **Cereus**, Haw. (Cactac.).

CIRIPEDIUM, Zumag. Fl. Pedem. i. 18 (1829) = **Cypripedium**, Linn. (Orchid.).

CIRRHAEA, Lindl. Bot. Reg. sub t. 930 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 585.
SARCOGLOSSUM, Beer, Prakt. Orch. 306 (1854).
SCLEROPTERIS, Scheidw. in Allg. Gartenz. (1839) 407.
adspersa, *Hoffm. ex Heynh. Nom.* ii. 145, nomen.—Bras.
dependens, Loud. Hort. Brit. 370 = Loddigesii.
dependens, Reichb. f. in Walp. Ann. vi. 496 = viridipurpurea.
fuscolutea, Hook. Bot. Mag. t. 3726 = saccata.
fuscolutea, *Lindl. Bot. Reg.* sub t. 1538.—Bras.
Hoffmanseggii, Heynh. ex Reichb. f. in Walp. Ann. vi. 496 = viridipurpurea.
Loddigesii, *Lindl. Bot. Reg.* t. 1538.—China?
obtusata, *Lindl. l. c.* xxiii. (1837) 2005.—Bras.
pallida, *Lindl. l. c.* sub t. 2005, nomen.—Hab.?
saccata, *Lindl. l. c.* (1839) *Misc.* 72.—Bras.
tristis, *Lindl. l. c.* t. 1889.—Mexic.
violaceo-virens, *Hoffmgg. Verz. Orch.*—*Cf. Bot. Zeit.* i. (1843) 831.—Hab.?
violascens, *Hoffmgg. ll. cc.*—Hab.?
viridipurpurea, *Lindl. Bot. Reg.* t. 1538.—Bras.
viridi-purpurea, Lodd. Bot. Cab. t. 1967 = Loddigesii.
Warreana, *Lindl. ex Reichb. f. in Walp. Ann.* vi. 497.—Bras.

CIRRHOPETALUM, Lindl. Bot. Reg. sub t. 832 (1824). *ORCHIDEAE*, Benth. & Hook. f. iii. 504.
EPHIPPIUM, Blume, Bijdr. 309. t. 65 (1825).
BOLBOPHYLLOPSIS, Reichb. f. in Bot. Zeit. x. (1852) 933.
HIPPOGLOSSUM, Breda, Orch. Knhl et Hass. (1827) [t. 14].
ZYGOGLOSSUM, Reinw. in Syll. Ratisb. ii. (1828) 4.
abbreviatum, *Reichb. f. in Gard. Chron.* (1881) II. 70.—Hab.?
albidum, *Wight, Ic.* t. 1653.—Ind. or.
Andersonii, *Kurz, Rep. Veg. Andaman.* 51.—Ins. Andaman.
antenniferum, *Lindl. Bot. Reg.* (1843) sub t. 49.—Ins. Philipp.
auratum, *Lindl. l. c.* (1840) *Misc.* 50.—Ins. Philipp.
Blumii, *Lindl. Gen. et Sp. Orch.* 59.—Java.
bootanensis, Griff. Notul. iii. 296 = maculosum.
caespitosum, *Wall. ex Lindl. Bot. Reg.* (1838) *Misc.* 35.—Reg. Himal.

CIRRHOPETALUM:—
capitatum, *Lindl. Bot. Reg.* (1843) sub t. 49.—Java.
carinatum, *Teijsm. & Binn. in Nederl. Kruidk. Arch.* iii. (1855) 397.—Java.
caudatum, Wight, Ic. t. 1658 = vaginatum.
chinense, *Lindl. Bot. Reg.* (1842) *Misc.* 37.—China.
clavigerum, *Fitzgerald, in Journ. Bot.* xxi. (1883) 204.—Austral.
compressum, *Lindl. Bot. Reg.* xxix. (1843) sub t. 49.—Java.
cornutum, *Lindl. l. c.* (1838) *Misc.* 75.—Reg. Himal.
Cumingii, *Lindl. l. c.* sub t. 49.—Ins. Philipp.
delitescens, Rolfe, in Gard. Chron. (1882) II. 461.—Hongkong.
elegans, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 310.—Malaya.
Elisae, F. Muell. Fragm. vi. 120. t. 57 = Bulbophyllum Elisae.
elongatum, *Lindl. Bot. Reg.* (1843) sub t. 49.—Reg. Himal.
fimbriatum, *Lindl. l. c.* (1839) *Misc.* 72.—Ind. or.
flagelliforme, Teijsm. & Binn. ex Reichb. in Walp. Ann. vi. 264 = Paludi.
gamosepalum, *Griff. Notul.* iii. 296.—Burma.
grandiflorum, *Wight, Ic.* v. t. 1658.—Zeylan.
lasiochilum, *Par. & Reichb. f. in Trans. Linn Soc.* xxx. (1874) 153 (Bulbophylli §).—Burma.
leopardinum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 309.—Java.
lineatum, *Teijsm. & Binn. l. c.* 311.—Malaya.
longiscapum, *Teijsm. & Binn. l. c.* 310.—Malaya.
Macraei, *Lindl. Gen. et Sp. Orch.* 59.—Zeylan.
Macraei, Wight, Ic. t. 1652 = Wightii.
maculosum, *Lindl. Bot. Reg.* (1841) *Misc.* 81.—Reg. Himal.
Makoyanum, Reichb. f. in Gard. Chron. (1879) I. 234 = Bulbophyllum Makoyanum.
maxillare, *Lindl. Bot. Reg.* (1843) sub t. 49.—Ins. Philipp.
Medusae, *Lindl. l. c.* (1842) t. 12.—Malaya.
neilgherrense, *Wight, Ic.* vi. t. 1654.—Ind. or.
nutans, *Lindl. Bot. Reg.* (1839) *Misc.* 71.—Manilla.
ornatissimum, *Reichb. f. in Gard. Chron.* (1882) II. 424.—Reg. Himal.
Pahudi, *De Vriese, Epim. Lugd.-Bat.* (1863) 2.—Java.
picturatum, *Lodd. in Bot. Reg.* (1840) *Misc.* 49.—Burma.
Pumilio, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 153 (Bulbophylli §).—Burma.
putidum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 311.—Java.
refractum, *Zoll. in Flora*, xxx. (1847) 456.—Reg. Himal.; Malaya.
retusiusculum, *Reichb. f. in Gard. Chron.* (1869) 1182.—Burma.
Roxburghii, *Lindl. Gen. et Sp. Orch.* 58.—Ind. or.
simillimum, *Reichb. f. Otia Hamburg.* 48.—Burma.
stramineum, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 310.—Java.
Thouarsii, *Lindl. Bot. Reg.* sub t. 832, in textu.—Ins. Masear.
trigonopus, *Reichb. f. in Gard. Chron.* (1881) II. 71.—Hab.?
tripudians, *Par. & Reichb. f. l. c.* (1876) I. 816.—Burma.
umbellatum, Linden, Illustr. Hortic. xxviii. (1881) 80. = maculosum?
vaginatum, *Lindl. in Wall. Cat.* n. 1979.—Malaya.
Walkerianum, Wight, Ic. v. t. 1657 = Macraei.
Wallichii, J. Grah. Cat. Pl. Bomb. 205 = fimbriatum?
Wallichii, Lindl. in Wall. Cat. n. 1980 = refractum.
Wallichii, *Lindl. in Wall. Pl. As. Rar.* i. 53. t. 67.—Reg. Himal.
Wightii, *Thw. Enum. Pl. Zeyl.* 299.—Zeylan.

CIRSELIUM, Brot. Fl. Lusit. i. 346 (1804) = seq.

CIRSELLIUM, Gaertn. Fruct. ii. 454. t. 163 (1791) = **Atractylis**, Linn. (Compos.).
cancellatum, Gaertn. l. c. = A. cancellata.
humile, Gaertn. l. c. 455 = A. humilis.
gummiferum, Brot. Fl. Lusit. i. 346 = A. gummifera.

CIRSIUM, [Tourn.] Adans. Fam. ii. 116 (1763); DC. Fl. Fr. iv. 110 (1805) = **Cnicus**, Tourn. (Compos.).
abyssinicum, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 456 = Cn. lanceolatus.
acanthifolium, Arv. in Bull. Soc. Dauph. iii. (1876) 73.—Gallia.
Acarna, Moench, Meth. Suppl. 226 = Cn. Acarna.
acaule, [Weber, ex] Wigg. Prim. Fl. Hols. 59 = Cn. acaulis.
acaulos, Scop. Fl. Carn. ed. II. ii. 131 = Cn. acaulis.
aciculare, Tausch, in Flora, xii. (1829) I. Erg. 40.—Hab.?
aduncum, Fisch. & Mey. ex DC. Prod. vi. 636 = *fallax*.
affine, Tausch, in Flora, xvi. (1833) 228.—Bohem.
afrum, DC. Cat. Hort. Monsp. 96 = Cn. afer.
aggregatum, Ledeb. Fl. Ross. ii. 737.—Reg. Caucas.
Alberti, Regel & Schmalh. in Act. Hort. Petrop. v. (1879) 318.—Turkestan.
albicans, Willk. in Linnaea, xxx. (1859–60) 109.—Hispan.
Allionii, Spenn. Fl. Friburg. iii. 1079 = Cn. acaulis.
Allionii, Thurm. ex Arduin. Fl. Alpes-Marit. 198 = Cn. pyrenaicus.
alpestre, Naegeli, Cirs. Schw. 84 = Cn. acaulis.
alpestre ×, Treuinf. in Zeitschr. Ferdinand. Tirol. III. xix. (1875) 228.—Tyrol.
alpinum, All. Fl. Pedem. i. 153 = Saussurea alpina, discolor.
altissimum, Hill, Hort. Kew. 63 = Cn. altissimus.
ambiguum, All. Auct. Pedem. 10 = Cn. heterophyllus.
anglicum, DC. Fl. Fr. iv. 118 = Cn. pratensis.
angulosum, Tausch, in Flora, xix. (1836) 402.—Hab.?
angustifolium, DC. Prod. vi. 648 = Carduus angustifolius.
angustifolium, Tausch, in Flora, xii. (1829) I. Erg. 39 = *Tauschii*.
angustissimum, DC. Prod. vi. 652 = Trichol epis strictophyllum.
apiculatum, DC. l. c. 642 = *libanoticum*.
appendiculatum, Griseb. Spicil. Fl. Rumel. ii. 250.—Rumel.; Hereeg.
arachnoideum, Bieb. Fl. Taur. Caue. iii. 557 = Cn. arachnoideus.
araneosum, Steud. Nom. ed. II. i. 367 = Cn. areneosus.
arctioides, Scop. Fl. Carn. ed. II. ii. 124 = Carduus arctioides.
arctotidifolium, Sch. Bip. in Schweinf. Beitr. Fl. Aethiop. 144 = Cn. Schimperi.
argentum, Peyer, ex Vest, in Flora, xii. (1829) I. Erg. 57 = Cn. arvensis.
arguncnse, DC. Prod. vi. 644.—Sibiria.
argyracanthum, DC. l. c. 640 = Cn. argyracanthus.
aristatum, DC. l. c.—Armenia.
armenum, DC. l. c. 647 = *simplex*.
arvense, Scop. Fl. Carn. ed. II. ii. 126 = Cn. arvensis.
Aschersonii ×, Čelak. in Arch. Naturfw. Landesd. Boehm. ii. (1873) 263.—Bohem.
atriplicifolium, Fisch. ex Steud. Nom. ed. II. i. 367 = Serratula atriplicifolia.
atticum, Sch. Bip. ex Nym. Consp. 406 = *crinitum*.
Ausserdorferi ×, Hausm. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 223 = Cn. ochroleucus?
autareticum, Mutel, Fl. Dauph. ii. 262 = Cn. heterophyllus.
baeticum, Spreng. Syst. iii. 377.—Hispan.
benacense, Treuinf. in Oestr. Bot. Zeitschr. xxiv. (1874) 172.—Tyrol.
Bertolonii, Spreng. Syst. iii. 377 = Cn. Bertolonii.
Bigelowii, DC. Prod. vi. 652 = Cn. muticus.
bipontinum, F. Schultz, in Arch. Fl. Fr. et Allem. 34 = Cn. acaulis.
Boujarti, Sch. Bip. in Oestr. Bot. Wochenbl. vi. (1856) 299 = Cn. ciliatus.
Bourgaeanum, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 191.—Hispan.
brachycephalum, Juratzka, in Verh. Zool. Bot. Ver. Wien, vii. (1857) 99 = *Chailleti*, Koch.
bracteatum, Link, in Linnaea, ix. (1834) 580 = Cn. syriacus.
bractcosum, DC. Prod. vi. 641 = Cn. strigosus.
Braunii, F. Schultz, ex Nym. Consp. 409 = Cn. oleraceus.
brevicaule, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 396 = Cn. japonicus.
brevifolium, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 421 = Cn. undulatus.
brevispinum, Sch. Bip. ex Nym. Consp. 410 = Cn. pungens.

CIRSIUM :—
britannicum, Scop. Ann. II. Hist. Nat. (1769) 60 = Cn. pratensis?
Brueggeri ×, Kill. ex Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii.–xxiv. (1880) 108.—Helvet.
Brunneri, A. Braun, ex Nym. Consp. 407 = Cn. tuberosus.
Buchingeri, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 284, nomen.—Abyss.
Buergeri, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 185 = Cn. Buergeri.
bulbosum, DC. Fl. Fr. iv. 118 = Cn. tuberosus.
bulbosum, Lam. Fl. Fr. ii. 22 = Cn. monspessulanus.
bulbosum, Willk. in Flora, xxxiv. (1851) 760 = *filipendulum*, Lange.
bulgaricum, DC. Prod. vi. 639 = Cn. giganteus.
byzantinum, Steud. Nom. ed. II. i. 367.—Rumel.
californicum, A. Gray, in Pacif. Rail. Rep. iv. 112 = Cn. californicus.
Candelabrum, Griseb. Spicil. Fl. Rumel. ii. 251.—Graecia; Rumel.
Candolleanum, Naeg. Cir. Schweiz. 98 = Cn. ochroleucus?
canescens, A. Gray, Pl. Fendl. 110 = Cn. neo-mexicanus.
canescens, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 420 = Cn. undulatus.
canum, All. Fl. Pedem. i. 151 = Cn. pratensis?
canum, Bieb. Fl. Taur. Caue. ii. 274 = Cn. canus.
Caput-Medusae, Schur, ex Nym. Consp. 408 = Cn. spinosissimus.
carlinoides, Fisch. in DC. Prod. vi. 661 = Cn. carlinoides.
carminans, Dum. Fl. Belg. 74.—Belgium.
carniolicum, All. Fl. Pedem. i. 149 = Cn. rivularis.
carniolicum, Scop. Fl. Carn. ed. II. ii. 128 = Cn. rufescens.
Carolorum ×, C. Jenner, ex Nym. Consp. 407 = Cn. heterophyllus.
carthamoides, Link, Enum. Hort. Berol. ii. 303 = Centaurea carthamoides.
Casabonae, DC. Fl. Fr. iv. 121 = Cn. Casabonae.
castellanum, Willk. in Linnaea, xxx. (1859–60) 109 = Cn. flavispinus.
cataonicum, Boiss. & Haussk. in Boiss. Fl. Orient. iii. 535.—Syria.
Čelakovskianum ×, Knaf, in Oestr. Bot. Zeitschr. xxii. (1872) 309 = Cn. arvensis.
Cephalotes, Boiss. Diagn. Ser. I. vi. 102 = Cn. horridus.
cernuum, Lag. Gen. et Sp. Nov. 24 = Cn. nivalis.
Cervini, Koch, Syn. Fl. Germ. ed. I. 399 = Cn. spinosissimus.
Chailleti, Gaudin, Fl. Helv. v. 182 = Cn. palustris.
Chailleti, Koch, Syn. Fl. Germ. ed. I. 393.—Europ.
Chamaepeuce, Tenore, Fl. Nap. v. 211 = Cn. gnaphalodes.
chelmeum, Orph. in Boiss. Diagn. Ser. II. iii. 41 = *Candelabrum*.
chinense, Gardn. & Champ. in Hook. Kew Journ. i. (1849) 323 = Cn. chinensis.
chium, DC. Prod. vi. 646 = Carduus chius.
chrysicum, Boiss. Fl. Orient. iii. 536.—Armenia.
cichoraceum, Spreng. Syst. iii. 370 = Serratula cichoracea.
ciliato-affine, Bess. Enum. Pl. Volh. 32 = Cn. serrulatus.
ciliatum, Bieb. Fl. Taur. Caue. iii. 556 = Cn. ciliatus.
ciliatum, Moench, Meth. Suppl. 227.—Hab.?
cilliacum, Tardent, Ess. Hist. Nat. Bessar. 62 = Cn. ciliatus.
compactum, Lam. Fl. Fr. ii. 24 = Cn. monspessulanus.
congestum, Fisch. & Mey. ex DC. Prod. vi. 641.—Persia.
consanguineum, DC. l. c. = *congestum*.
conspicuum, Sch. Bip. in Seem. Bot. Voy. Herald, 312 = Cn. conspicuus.
controversum, DC. Prod. vi. 654 = Cn. spinosissimus.
Cosmelii, Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 251 = Cn. lappaceus.
Coulteri, Harv. & Gray, Pl. Fendl. 110 = Cn. occidentalis.
creticum, Urv. in Mém. Soc. Linn. Par. i. (1822) 363.—Archip. Graec.
crinitum, Boiss. ex DC. Prod. vii. 305.—Hispan.
cryptocephalum, Sch. Bip. ex Boiss. Fl. Orient. iii. 592 = Centaurea pygmaea.
csepeliense, Borb. in Oestr. Bot. Zeitschr. xxviii. (1878) 392.—Europ.
cyanoides, Hill, Hort. Kew. 63 = Jurinea cyanoides.
cynaroides, Griseb. Spicil. Fl. Rumel. ii. 254 = Cn. atr.

CIRSIUM :—
cynaroides, Spreng. Syst. iii. 372 = Cnicus cynaroides.
dealbatum, Bieb. Fl. Taur. Cauc. iii. 560.—Reg. Caucas.
decipiens, Nym. Syll. Suppl. 4 = Cn. acaulis.
decoloratum, Koch, Syn. Fl. Germ. ed. I. 398 = Cn. acaulis, oleraceus.
decussatum, Janka, in Linnaea, xxx. (1859-60) 582 = Cn. ferox, *Sibth. & Sm.*
defloratum, Scop. Fl. Carn. ed. II. ii. 127 = Carduus defloratus.
dentatum, DC. Prod. vi. 654 = Kentrophyllum dentatum.
depilatum, Boiss. & Bal. in Boiss. Fl. Orient. iii. 543.—As. Min.
desertorum, Fisch. ex Link, Enum. Hort. Berol. ii. 300.—Sibiria.
desertorum, Janka, ex Nym. Consp. 405 = Cn. ciliatus.
Diacantha, DC. Cat. Hort. Monsp. 96 = Cn. Diacantha.
dioicum, Cass. in Dict. Sc. Nat. xxvii. 190 = Cn. arvensis.
discolor, Hort. ex Steud. Nom. ed. II. i. 368 = Cn. ciliatus.
discolor, Spreng. Syst. iii. 373 = Cn. altissimus.
dissectum, Hill, Hort. Kew. 63 = Cn. dissectus.
diversifolium, DC. Prod. vi. 649 = Cn. altissimus.
Dollinerii, Sch. Bip. ex Hausm. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 373 = Cn. pannonicus.
Douglasii, DC. Prod. vi. 643 = Cn. undulatus.
Drejeri, Reichb. f. Ic. Fl. Germ. xv. 60 = Cn. lanceolatus.
Drummondii, Torr. & Gray, Fl. N. Am. ii. 459 = Cn. Drummondii.
duriusculum, C. Koch, in Linnaea, xxiv. (1851) 410 = *elodes*, Cn. palustris.
echinatum, DC. Fl. Fr. Suppl. 465 = Cn. echinatus.
echinocephalum, Bieb. Fl. Taur. Cauc. iii. 559 = Cn. echinocephalus.
echinopifolium, Boiss. ex Tchihat. Asie Min. Bot. ii. 343.—As. Min.
edule, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 420 = Cn. edulis.
elatius, Link, Enum. Hort. Berol. ii. 299 = Cn. polyanthemus.
elatum, Sauter, in Flora, xxviii. (1845) 130.—Europ.
elatum, Tod. Ind. Sem. Hort. Panorm. (1858) 25; et in Ann. Sc. Nat. Sér. IV. xi. (1859) 378.—Sicilia.
elegans, Willd. ex Spreng. Syst. iii. 375 = *igniarium*.
elodes, Bieb. Fl. Taur. Cauc. iii. 555.—Rossia; Reg. Caucas.; Persia.
Empedocleanum, Lojac. in Natural. Sicil. iii. (1884) 284.—Sicil.
eriocephalum, A. Gray, in Proc. Acad. Sc. Philad. '1863' (1864) 69 = Cn. eriocephalus.
eriocephalum, Wallr. Beitr. Fl. Hercyn. 262 = Cn. eriophorus.
eriophorum, Scop. Fl. Carn. ed. II. ii. 130 = Cn. eriophorus.
Erisithales, Scop. Ann. II. Hist. Nat. 62 = Cn. ochroleucus?
erisithaloides ×, Huter, ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 271.—Tyrol.
erucagineum, DC. Fl. Fr. iv. 115 = Cn. rivularis.
erythrolepis, C. Koch, in Linnaea, xvii. (1843) 41.—Reg. Caucas.
esculentum, C. A. Mey. in Beitr. Pfl. Russ. Reich. iv. 43; et in Flora, xxxii. (1849) 747 = Cn. esculentus.
exaltatum ×, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 109.—Helvet.
exscapum, Fisch. Hort. Gorenk. ed. II. (1812) 35, nomen.—Hab.?
falcatum, Turcz. ex DC. Prod. vi. 650 = Cn. pendulus.
fallax, Fisch. & Mey. ex Hohen. in Bull. Soc. Nat. Mosc. (1838) 280.—Reg. Caucas.
fallax, Franch. in Billot, Annot. 109 = Cn. acaulis.
ferox, DC. Fl. Fr. iv. 120 = Cn. ferox.
ferox, Schur, ex Nym. Consp. 405 = Cn. ciliatus.
ficifolium, Fisch. in Mém. Soc. Nat. Mosc. iii. (1812) 69 = Serratula atriplicifolia.
filipendulum, Engelm. in A. Gray, Man. Bot. N. U. St. ed. V. 273 = Cn. altissimus.
filipendulum, Lange, in Kjoeb. Vidensk. Meddel. (1861) 92.—Hispan.
fimbriatum, Spreng. Syst. iii. 373 = Cn. fimbriatus.
Fischerianum, Andrz. ex Trautv. in Act. Hort. Petrop. viii. (1883) 501, nomen.—Rossia.
fissum ×, Ausserd. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 262.—Tyrol.
flavescens, Koch, Taschenb. 292 = Cn. ochroleucus.
flavescens Lam. Fl. Fr. ii. 28 = Cn. parviflorus.

CIRSIUM :—
flavispinum, Boiss. in DC. Prod. vii. 305 = Cnicus flavispinus.
foliosum, DC. l. c. vi. 654 = Cn. foliosus.
Forsteri, Loud. Hort. Brit. Suppl. i. 585 = Cn. palustris.
fraternum, DC. Prod. vi. 636 = Cn. strigosus.
Freyerianum, Koch, Taschenb. 288; Syn. Fl. Germ. ed. II. 454.—Dalmat.
Frickii, Fisch. & Mey. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. vi. (1849) 47 = *rhizocephalum*.
fulvispinum, Walp. Rep. ii. 676, sphalm. = Cn. flavispinus.
furiens, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) I. 348 = Cn. ciliatus.
Gaillardoti, Boiss. Diagn. Ser. II. iii. 42.—Syria.
Galactites, Hill, Hort. Kew. 64 = Galactites tomentosa.
Ganderi ×, Huter, ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 274.—Tyrol.
Gerhardti, Sch. Bip. in Flora, xxxii. (1849) 547 = Cn. lanceolatus.
giganteum, Spreng. Syst. iii. 375 = Cn. giganteus.
glabrum, DC. Fl. Fr. Suppl. 463 = Cn. spinosissimus?
glaucescens ×, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 227.—Tyrol.
glutinosum, Lam. Fl. Fr. ii. 27 = Cn. ochroleucus.
Gmelini, Tausch, in Flora, xi. (1828) 482 = Cn. acaulis.
gnaphalodes, Spreng. Syst. iii. 375 = Cn. gnaphalodes.
Godroni, Sch. Bip. ex Nym. Consp. 406 = Cn. lanceolatus.
gracile, Rostr. ex Nym. l. c. = Cn. lanceolatus.
gracile, Schur, Enum. Pl. Transs. 422 = Cn. rivularis.
Grahami, A. Gray, Pl. Wright. ii. 102 = Cn. Grahami.
grandiflorum ×, Kitt. Taschenb. ed. II. 551.—Europ.
gregarium, Willk. in Willk. & Lange, Prod. Fl. Hisp. ii. 189.—Hispan.
Gremblichii ×, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 213.—Tyrol.
Griffithii, Boiss. Fl. Orient. iii. 540.—Afghanist.
grumosum, Fisch. & Mey. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. vi. (1849) 48 = *rhizocephalum*.
grumosum, Willk. & Lange, Prod. Fl. Hisp. ii. 199 = Cn. grumosus.
Gutnickianum ×, Loehr, Enum. Fl. Deutsch. 365.—Europ.
Halleri, Tausch, in Flora, xix. (1836) 401.—Helvet.
Hallerianum, Gaudin, Fl. Helv. v. 188 = Cn. spinosissimus.
Hallerianum, Hill, Veg. Syst. iv. 27.—Helvet.
halophilum, Turcz. ex Herd. in Bull. Soc. Nat. Mosc. xliii. (1870) I. 83 = Cn. arvensis.
Haussknechtii, Boiss. Fl. Orient. iii. 535.—Kurdistan.
helenioides, All. Fl. Pedem. i. 152 = Cn. heterophyllus.
helenioides, Hill, Hort. Kew. 64 = Cn. helenioides.
hellenicum, Boiss. & Orph. ex Boiss. Fl. Orient. iii. 530 = Cn. ferox.
Hermonis, Boiss. Diagn. Ser. II. iii. 40 = Cn. lappaceus.
heterolepis, Benth. Pl. Hartw. 87 = Cn. subcoriaceus.
heteromallum, Spreng. Syst. iii. 472; Sweet, Hort. Brit. ed. I. 214 = Saussurea candicans.
heterophylloides ×, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 245.—Austr.
heterophyllum, Hill, Hort. Kew. 64; All. Fl. Pedem. i. 152 = Cn. heterophyllus.
heterotrichum, Panč. Élem. Fl. Bulg. 42.—Europ.
Hippolyti, Link, ex Steud. Nom. ed. II. i. 368 = Cn. pungens.
Hookerianum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 418 = Cn. Hookerianus.
horridulum, Michx. Fl. Bor. Am. ii. 90 = Cn. horridulus.
horridum, Fisch. Hort. Gorenk. ed. II. (1812) 35 = Cn. horridus.
horridum, Lag. ex DC. Prod. vi. 660.—Hispan.
Huguenini ×, Buegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 109.—Europ.
Huteri ×, Hausm. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 221.—Europ.
hybridum, Koch, in DC. Fl. Fr. Suppl. 463 = Cn. ochroleucus.
hygrophilum, Boiss. Diagn. Ser. I. x. 89.—Persia.
hypoleucum, DC. Prod. vi. 645.—Balkan; As. Min.
hypopsilum, Boiss. & Heldr. Diagn. Ser. I. vi. 101 = Cn. ferox.
igniarium, Spreng. Syst. iii. 373.—Sibir. altaic.
imereticum, Boiss. Fl. Orient. iii. 537.—Reg. Caucas.
incanum, Bieb. Fl. Taur. Cauc. iii. 561 = Cn. arvensis.
inclinatum, Lam. Fl. Fr. ii. 22 = Carduus medius.

CIRSIUM :—
inerme, G. Don, in Loud. Hort. Brit. 330; Reichb. Fl. Germ. Excurs. 287 = Cnicus tuberosus.
inerme, Link, Enum. Hort. Berol. ii. 301 = Cn. cynaroides.
insubricum, Moretti, ex Bertol. Fl. Ital. ix. 25 = Cn. eriophorus.
involucratum, DC. Prod. vi. 639 = Cn. involucratus.
italicum, DC. Cat. Hort. Monsp. 96 = Cn. italicus.
Jaegeri ×, F. Schultz, in Flora, xxxii. (1849) 228.—Europ.
japonicum, DC. Prod. vi. 640 = Cn. japonicus.
jorullense, Spreng. Syst. iii. 376 = Cn. jorullensis.
Joschii, Pacher, in Jahrb. Landesm. Kaernt. xv. (1882) 136.—Europ.
Juratzkae ×, Reichardt, in Verh. Zool.-Bot. Ges. Wien, xi. (1861) 380.—Austr.
kamtschaticum, Ledeb. ex DC. Prod. vi. 644 = Cn. kamtschaticus.
Kerneri ×, Ausserd. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 260.—Europ.
Killiasii, Bruegg. in Jahresb. Naturf. Ges. Graub. II. xxiii–xxiv. (1880) 108.—Helvet.
kirbense, Pomel, Nouv. Mat. Fl. Atl. 274.—Afr. bor.
Kochianum, Loehr, in Flora, xxv. (1842) 111 = Cn. palustris.
Kosmelii, Boiss. Fl. Orient. iii. 531, in syn. = Cn. lappaceus.
Kotschyanum, Boiss. Diagn. Ser. I. vi. 103 = *libanoticum*.
Lachenalii, DC. Prod. vi. 648, in syn. = Cn. rigens.
Lachenalii, Koch, Syn. Fl. Germ. ed. I. 397 = Cn. acaulis, oleraceus.
laciniatum, Doell, ex Nym. Consp. 409 = Cn. palustris.
lacteum ×, Ruhm. in Eichl. Jahrb. Bot. Gart. Berl. i. (1881) 241.—Europ.
lacteum, Schleich. Cat. (1821), ex Koch, Syn. Fl. Germ. ed. I. 394 = palustris.
laeve, Moench, Meth. Suppl. 227 = Cn. tataricus.
laevigatum, Tausch, in Flora, xi. (1828) 483 = Cn. arvensis.
lamprophyllum, Herbich, Stirp. Rar. Bucov. 54.—Bukovina.
lanatum, Spreng. Syst. iii. 373 = Cn. arvensis.
lanceolatum, Hill, Herb. Brit. i. 80; Scop. Fl. Carn. ed. II. ii. 130 = Cn. lanceolatus.
laniflorum, Bieb. Fl. Taur. Cauc. iii. 557 = Cn. laniflorus.
lanigerum, Naegeli, Cirs. Schweiz. 114. t. 3 = Cn. lanceolatus.
lappaceum, Fisch. Hort. Gorenk. ed. II. (1812) 35 = Cn. lappaceus.
lappaceum, Lam. Fl. Fr. ii. 24 = Carduus Personata.
lappoides, Sch. Bip. in Seem. Bot. Voy. Herald, 312 = Cn. lappoides.
latifolium, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 28.—Ins. Madera.
Lecontei, Torr. & Gray, Fl. N. Am. ii. 458 = Cn. Lecontei.
Ledebourii, Steud. Nom. ed. II. i. 368.—Sibiria.
leucanicum, Lojac. in Natural. Sicil. iii. (1884) 283.—Sicilia.
leucocephalum, C. Koch, in Linnaea, xxiv. (1851) 401 = *chynysicum*.
leucocephalum, Spreng. Syst. iii. 377 = Cn. leucocephalus.
leuconeurum, Boiss. & Heldr. in Boiss. Fl. Orient. iii. 534.—Cataonia.
leucophanum, Schur, Enum. Pl. Transs. 418 = Cn. laniflorus.
leucopsis, DC. Prod. vi. 644.—As. Min.
leucostylum, Moench, Meth. 556 = Carduus defloratus.
libanoticum, DC. Prod. vi. 647.—As. Min.; Syria.
ligulare, Boiss. Fl. Orient. iii. 529.—Thracia.
Linkii, Nym. Syll. 23.—Lusitan.
lineare, Sch. Bip. in Linnaea, xix. (1847) 335 = Cn. linearis.
litorale, Maxim. Prim. Fl. Amur. 173 = Cn. japonicus.
Linkianum ×, Loehr, Enum. Fl. Deutsch. 364.—Europ.
Lobelii, Tenore, Fl. Nap. v. 211. t. 189 = Cn. ferox.
longespinosum, Tod. ex Nym. Consp. 406 = Cn. lanceolatus.
longifolium, Boiss. Fl. Orient. iii. 553, sphalm. (= *Epitrachys longifolia*).—Reg. Caucas.
lucidum, DC. Prod. vi. 654.—Hab.?
lutescens, Rafin. in Med. Repos. N. York, II. v. (1808) 354, nomen.—Hab.?
lycaonicum, Boiss. & Heldr. Diagn. Ser. I. x. 90 = *libanoticum*.

CIRSIUM :—
lyratum, Bunge, Enum. Pl. Chin. Bor. 36 = Saussurea affinis.
Maackii, Maxim. Prim. Fl. Amur. 172 = Cn. japonicus.
macracanthum, Sch. Bip. in Flora, xxxv. (1852) 135 = Cn. Wallichii.
macrobotrys, Boiss. Fl. Orient. iii. 532.—Armenia.
macrocephalum, C. A. Mey. Verz. Pfl. Cauc. 70 = Cn. munitus.
macrochaetum, Tausch, in Flora, xix. (1836) 397.—Hab.?
macrostachyum, Boiss. Fl. Orient. iii. 550.—Reg. Caucas.
macrostylon, Reichb. Fl. Germ. Excurs. 856 = Cn. arvensis.
maculatum, Lam. Fl. Fr. ii. 22 = Tyrimnus leucographus.
maculatum, Moench, Meth. 557 = Cn. syriacus.
maculatum, Scop. Fl. Carn. ed. II. ii. 130 = Silybum Marianum.
mareoticum, Spreng. Syst. iii. 377 = Carthamus marcoticus.
matritense, Lag. ex Cut. Fl. Comp. Madrid, 409 = Cn. flavispinus.
maximum, Benth. Pl. Hartw. 289 = Cn. subcoriaceus.
medium, All. Fl. Pedem. i. 149 = Cn. tuberosum.
megacanthum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 421 = Cn. horridulus.
mexicanum, DC. Prod. vi. 636 = Cn. mexicanus.
micranthum ×, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 218.—Europ.
microcephalum, Lange, Ind. Sem. Hort. Havn. (1854) 22.—Hispan.
microcephalum, Moris, Fl. Sard. ii. 466 = Cn. microcephalus.
microcephalum, Sch. Bip. in Oestr. Bot. Zeitschr. viii. (1858) 246 = Cn. siculus.
Mielichhoferi, Sauter, in Flora, xxviii. (1845) 130 = Cn. heterophyllus.
misilmerense, Ces. Passer. & Gib. Comp. Fl. Ital. 483 = Cn. misilmerensis.
mite, Fisch. in Spreng. Nov. Prov. 13 = Cn. mitis.
molle, Scop. Fl. Carn. ed. II. ii. 126 = Jurinea mollis.
monspessulanum, Hill, Hort. Kew. 63; All. Fl. Pedem. i. 152 = Cn. monspessulanus.
montanum, Hill, Herb. Brit. i. 80 = Cn. montanus.
Morettianum, Nym. Syll. 24 = Cn. spathulatus.
morinaefolium, Boiss. & Heldr. ex Boiss. Fl. Orient. iii. 530.—Ins. Creta.
Morisianum, Reichb. f. Ic. Fl. Germ. xv. 59.—Europ.
munitum, Fisch. Hort. Gorenk, ed. II. (1812) 35 = Cn. munitus.
mutatum, Menyh. Kalocsa Vidék. Növény. 102 = Cn. arvensis.
muticum, Michx. Fl. Bor. Am. ii. 89 = Cn. muticus.
Naegelianum ×, Lochr, Enum. Fl. Deutsch. 362.—Europ.
neglectum, Fisch. ex Spreng. Syst. iii. 370 = Cn. arvensis.
nemorale, Reichb. Fl. Germ. Excurs. 286 = Cn. lanceolatus.
neo-mexicanum, A. Gray, Pl. Wright, ii. 101 = Cn neo-mexicanus.
nepalense, DC. Prod. vi. 642 = Cn. Wallichii.
nevadense, Willk. in Linnaea, xxx. (1859-60) 111.—Hispan.
nidulans, Regel, in Bull. Soc. Nat. Mosc. xl. (1867) ii. 160.—As. centr.
nivale, Sch. Bip. in Seem. Bot. Voy. Herald, 312 = Cn. nivalis.
niveum, Spreng. Syst. iii. 373 = Cn. niveus.
numidicum, Spreng. l. c. 377.—Afr. bor.
Nuttalii, DC. Prod. vi. 651 = Cn. Nuttallii.
oblongifolium, C. Koch, in Linnaea, xvii. (1843) 41.—Reg. Caucas.
obscurum ×, Kern. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 264.—Europ.
obvallatum, Bieb. Fl. Taur. Cauc. iii. 559 = Cn. obvallatus.
ochro-centrum, A. Gray, Pl. Fendl. 110 = Cn. ochrocentrus.
ochroleucum, All. Fl. Pedem. i. 150 = Cn. ochroleucus.
odontolepis, Boiss. ex DC. Prod. vii. 305.—Hispan.; Balkan.
oenanum ×, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 215.—Europ.
oenipontanum ×, Treuinf. l. c. 280.—Europ.
oleraceum, Scop. Fl. Carn. ed. II. ii. 124 = Cn. oleraceus.

CIRSIUM :—
onopordioides, Kotschy, ex Boiss. Fl. Orient. iii. 564 = Onopordum polycephalum.
oreithales, Hance, ex Walp. Ann. ii. 944 = Cnicus chinensis.
orgyale, Hort. ex Tausch, in Flora, xii. (1829) i. Erg. 41 = *aciculare*?
orgyale, Link, Enum. Hort. Berol. ii. 302 = Cn. orgyalis.
orientale, Spreng. Syst. iii. 373 = Galactites tomentosa.
palatinum, Sch. Bip. ex Nym. Consp. 409 = Cn. palustris.
pallens, DC. Prod. vi. 647 = Cn. oleraceus.
paludosum, G. Don, in Loud. Hort. Brit. 331 = Cn. paludosus.
palustre, Coss. ex Willk. & Lange, Prod. ii. 191 = *Bourgeanum*.
palustre, Scop. Fl. Carn. ed. II. ii. 128 = Cn. palustris.
paniculatum, Boiss. & Haussk. ex Boiss. Fl. Orient. iii. 535 = *cataonicum*?
paniculatum, Lam. Fl. Fr. ii. 25 = Carduus carlinoides.
paniculatum, Spreng. Syst. iii. 371.—Mont. Pyren.
paniculatum, Vahl, ex Willk. & Lange, Prod. Fl. Hisp. ii. 186 = Cn. flavispinus.
pannonicum, Link, Enum. Hort. Berol. ii. 301 = Cn. pannonicus.
paradoxum ×, Wallr. Beitr. Fl. Hercyn. 264.—Europ.
parviflorum, DC. Prod. vi. 647 = Cn. oleraceus.
parviflorum, Lange, ex Nym. Consp. 409 = Cn. palustris.
parviflorum, Schleich. ex Steud. Nom. ed. II. i. 369 = Cn. ochroleucus.
pauciflorum, Koch, Syn. Fl. Germ. ed. I. 395 = Cn. heterophyllus.
pauciflorum, Lam. Fl. Fr. ii. 22 = Carduus defloratus.
pauciflorum, Spreng. Syst. iii. 375 = Cn. pauciflorus.
pazcuarense, Spreng. l. c. 372 = Cn. pazcuarensis.
pectinellum, A. Gray, in Mem. Am. Acad. N. S. vi. (1858–59) 395 = Cn. pectinellus.
pendulum, Fisch. ex DC. Prod. vi. 650 = Cn. pendulus.
penicillatum, C. Koch, in Linnaea, xvii. (1843) 41 = Cn. lappaceus.
persicum, Boiss. Diagn. Ser. I. vi. 103 = Cn. strigosus.
phyllocephalum, Boiss. & Blanche, in Boiss. l. c. Ser. II. iii. 39.—Syria.
pinnatifidum, Spreng. Syst. iii. 395 = Serratula pinnatifida.
Pitcheri, Torr. & Gray, Fl. N. Am. ii. 456 = Cn. Pitcheri.
platycephalum, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1852) 110 = Cn. subcoriaceus.
polyacanthum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 456 = Cn. polyacanthus.
polyacanthum, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 450.—Sibir. altaic.
polyanthemon, Spreng. Syst. iii. 371 = Cn. polyanthemus.
polyanthos, Urv. in Mém. Soc. Linn. Par. i. (1822) 363 = Cn. pungens.
polycephalum, DC. Prod. vi. 639 = *byzantinum*.
polycephalum, Willd. ex Spreng. Syst. iii. 371.—Hab.?
polymorphum, Doll. ex Hausm. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 375 = *Linkianum*.
ponticum, C. A. Mey. Ind. Sem. Hort. Petrop. ix. 67 = *polycephalum*.
Portae, Hausm. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 373 = *Linkianum*.
praealtum, Cass. in Dict. Sc. Nat. xli. 333 = Cn. arvensis.
praemorsum, Koch, Syn. Fl., Germ. ed. I. 397 = Cn. ochroleucus?
pratense, DC. Fl. Fr. iv. 113 = Cn. monspessulanus.
proponticum, Griseb. Spicil. Fl. Rumel. ii. 254 = Cn. eriophorus.
pseudo-canum, Schur, Enum. Pl. Transs. 421 = Cn. pannonicus.
pseudo-lanceolatum, Schur, l. c. 418 = Cn. lanceolatus.
pseudo-oleraceum, Schur, l. c. 423 = Cn. pannonicus.
pseudo-palustre, Schur, l. c. 420 = *Chailleti*, Koch.
pseudo-pannonicum, Schur, l. c. = Cn. canus.
pseudo-personata, Boiss. & Bal. ex Boiss. Fl. Orient. iii. 545.—As. Min.
pubigerum, DC. Prod. vi. 650 = Cn. pubigerus.
Pumilio, Tausch, in Syll. Ratisb. ii. (1828) 249.—Austria.
pumilum, Spreng. Syst. iii. 375 = Cn. pumilus.

CIRSIUM :—
pungens, Schur, Enum. Pl. Transs. 420.—Europ.
pungens, Spreng. Syst. iii. 370 = Cnicus polyanthemus.
purpureum, All. Fl. Pedem. i. 150 = Cn. spinosissimus.
Pustariacum ×, Ausserd. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 258.—Europ.
pygmaeum, Scop. Fl. Carn. ed. II. ii. 123 = Saussurea pygmaea.
pyrenaicum, All. Fl. Pedem. i. 151 = Cn. montanus.
pyrenaicum, DC. Fl. Fr. iv. 112 = C. pyrenaicus.
racemosum, Boiss. in Tchihat. Asie Min. Bot. ii. 338 = *tenuilobum*.
radians, Benth. Pl. Hartw. 77 = Cn. radians.
Reichardtii ×, Jurotzka, in Verh. Zool.-Bot. Ges. Wien, ix. (1859) 317.—Europ.
Reichenbachianum ×, Lochr, Enum. Fl. Deutsch. 364 = Cn. arvensis.
remotifolium, DC. Prod. vi. 655 = Cn. remotifolius.
repandum, Michx. Fl. Bor. Am. iii. 28 = Cn. repandus.
replicatum, DC. Prod. vi. 655 = Cn. replicatus.
rheginum, Spreng. Pugill. ii. 77 = Cn. strictus.
rhizocephalum, C. A. Mey. Verz. Pfl. Cauc. 70.—As. Min.; Persia; Afghan.
rhodo-leucanthum, Walp. Ann. i. 453 = Cn. esculentus.
Richterianum, Gillot, in Bull. Soc. Bot. Fr. xxvii. (1880) p. li.—Europ.
rigens, Spreng. Neue Entdeck. iii. 38 = Cn. oleraceus.
rigens, Wallr. Sched. Crit. 446 = Cn. acaulis.
rigidum, DC. Prod. vi. 640 = Cn. arachnoideus.
rivulare, All. Auct. Pedem. 10 = Cn. montanus.
rivulare, Link, Enum. Hort. Berol. ii. 301 = Cn. rivularis.
Rosani, Tenore, Ind. Sem. Hort. Neap. (1830) 15; ex Syll. 414 = Cn. lanceolatus.
Rosenii, Vill. Hist. Pl. Dauph. iii. 14. t. 21 = Cn. acaulis?
rufescens, Ramond, ex DC. Fl. Fr. iv. 114 = Cn. rufescens.
ruthenicum, Fisch. Hort. Gorenk. ed. II. (1812) 35 = Cn. arvensis.
sabaudum ×, Lochr, Enum. Fl. Deutsch. 362.—Europ.
sabaudum ×, Ruhm. in Eichl. Jahrb. Bot. Gard. Berl. i. (1881) 240.—Europ.
salamanticum, Hill, Hort. Kew. 64 = Centaurea salmantica.
salinum, Bieb. ex Steud. Nom. ed. II. i. 370 = Centaurea salina.
salisburgense, G. Don, in Loud. Hort. Brit. 331 = Cn. rivularis.
samniticum, Steud. Nom. ed. II. i. 370 = Cn. italicus.
scariosum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 420 = Cn. scariosus.
schantarense, Trautv. & Mey. in Middend. Reise (Fl. Ochot. 58).—Sibir. or.
Schultzianum ×, Lochr, Enum. Fl. Deutsch. 365.
scleranthum, Bieb. Fl. Taur. Cauc. iii. 559.—As. Min.; Persia.
Scopolianum, Sch. Bip. ex Nym. Consp. 408 = Cn. Eristhales.
segetum, Bunge, Enum. Pl. Chin. Bor. 36 = Cn segetum.
Semenowi, Regel, in Bull. Soc. Nat. Mosc. xl. (1867) ii. 161.—As. centr.
semidecurrens, DC. Prod. vi. 646.—Hab.?
semidecurrens, Richt. Fl. Lips. 673 = Cn. palustris.
semipectinatum, Ledeb. ex Steud. Nom. ed. II. i. 370 = *Ledebourii*.
semipectinatum, Reichb. Fl. Germ. Excurs. 287 = Cn. rivularis.
semipectinatum, Spreng. Syst. iii. 378 = Cn. semipectinatus
serratuloides, Hill, Hort. Kew. 64 = Cn. serratuloides.
serratuloides, Scop. Fl. Carn. ed. II. ii. 127 = Cn. pannonicus.
serratum, Tausch, in Flora, xii. 1829) i. Erg. 39.—Europ.
serrulatum, Bieb. Fl. Taur. Cauc. iii. 557 = Cn. serrulatus.
serrulatum, Schur, Enum. Pl. Transs. 419 = Cn. fimbriatus.
sessiliflorum, Herbich, Stirp. Rar. Bucov. 54.—Turcia
setigerum, Ledeb. Ic. Pl. Fl. Koss. i. 9.—Sibir. altaic.
setosum, Bieb. Fl. Taur. Cauc. iii. 560 = Cn. arvensis.
siculum, Spreng. Neue Entdeck. iii. 36 = Cn pungens.
Sieboldi, Miq. Ann. Mus. Bot. Lugd. Bat. ii. 184 = Cn. Sieboldii.
Siegerti, Sch. Bip. ex Nym. Consp. 408 = Cn. rivularis.

CIRSIUM :—
silesiacum, Sch. Bip. ex Nym. Consp. 409 = Cnicus palustris.
silvaticum, Tausch, in Flora, xii. (1829) i. Erg. 38 = Cn. lanceolatus.
simplex, Hornem. Hort. Hafn. Snppl. 93 = Centaurea salina.
simplex, C. A. Mey. Verz. Pfl. Cauc. 70.—Reg. Caucas.
sinense, C. B. Clarke, Comp. Ind. 219 = Cn. chinensis.
sinuatum, Boiss. Fl. Orient. iii. 546 = Cn. cynaroides.
sinuatum, Boiss. & Kotschy, ex Boiss. l. c. 542 = *rhizocephalum*.
sordidum, Wallr. Beitr. Fl. Hercyn. 266 = Cn. arvensis.
sorocephalum, DC. Prod. vi. 636 = *fallax*, Fisch. & Mey.
sorocephalum, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 33.—Reg. Caucas.
spathulatum, Gaudin, Fl. Helv. v. 202 = Cn. eriophorus.
spectabile, DC. Prod. vi. 641.—Persia.
spinifolium ×, Beck, in Oestr. Bot. Zeitschr. xxxi. (1881) 312.—Europ.
spinosissimoides ×, Ausserd. ex Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 250.—Europ.
spinosissimum, Scop. Fl. Carn. ed. II. ii. 129 = Cn. spinosissimus.
Spitzelii, Sch. Bip. ex Nym. Consp. 408 = Cn. spinosissimus.
spurium, Delastre, in Ann. Sc. Nat. Sér. II. xviii. (1842) 149 = Cn. pratensis.
squarrosum, Fisch. ex Spreng. Syst. iii. 376 = Cn. squarrosus.
stellatum, All. Fl. Pedem. i. 153 = Cn. stellatus.
stenolepidum, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 419 = Cn. remotifolius.
stenophyllum, Boiss. Fl. Orient. iii. 534 = *tenuilobum*.
Stocksii, Boiss. Diagn. Ser. II. iii. 40 = Cn. arvensis.
strictum, Moris, Stirp. Sard. Elench. i. 27 = Cn. microcephalus.
strictum, Spreng. Syst. iii. 370 = Cn. strictus.
strigosum, Bieb. Fl. Taur. Cauc. iii. 558 = Cn. strigosus.
subalatum, Gaudin, Fl. Helv. v. 180 = Cn. oleraceus.
subalpinum, Gaudin, l. c. 182 = Cn. palustris.
subalpinum, Schleich. ex Reichb. Ic. Fl. Germ. xv. 73.—Helvetia.
subcoriaceum, Sch. Bip. ex Seem. Bot. Voy. Herald, 312 = Cn. subcoriaceus.
subinerme, Fisch. & Mey. in Mém. Acad. Pétersb. Sér. VI. Sc. Nat. vi. (1849) 55 = *elodes*.
subsagittatum, Schur, Enum. Pl. Transs. 421 = Cn. pannonicus.
succinctum, Ledeb. Fl. Ross. ii. 732 = Cn. lappaceus.
supertextum, Tausch, in Flora, xix. (1836) 397.—Hab. ?
sylvaticum, DC. Prod. vi. 656 = Cn. lanceolatus.
syriacum, Gaertn. Fruct. ii. 383. t. 163. f. 2 = Cn. syriacus.
Szovitsii, Boiss. Fl. Orient. iii. 527.—Reg. Caucas.
tartaricum, All. Fl. Pedem. i. 151 = Cn. tataricus.
tataricum, DC. Fl. Fr. iv. 114 = Cn. rigens.
tauricolum, Boiss. & Bal. ex Boiss. Fl. Orient. iii. 551.—Cataonia.
Tauschii, DC. Prod. vi. 656.—Hab. ?
tchefouënse, Debeaux, Fl. Tchef. 86 = Cn. tchefouënsis.
tenuilobum, Boiss. Fl. Orient. iii. 534.—Armenia.
texanum, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 460 = Cn. virginianus.
Thomasii, Naeg. Cirs. Schweiz. 93 = Cn. oleraceum.
tirolense ×, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 243.—Europ.
tomentosum, C. A. Mey. Verz. Pfl. Cauc. 69 = Cn. munitus.
tomentosum, Moench, Meth. Suppl. 227 = Jurinea cynaroides.
trachylepis, Boiss. Fl. Orient. iii. 528.—Armenia.
transsilvanicum, Schur, Enum. Pl. Transs. 419 = Cn. transsilvanicus.
tricephalodes, DC. Fl. Fr. iv. 116 = Cn. rivularis.
tricephalum, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = praec.
tricholoma, Fisch. & Mey. in Bull. Soc. Nat. Mosc. (1838) 251.—Reg. Caucas.
trispinosum, Moench, Meth. 556 = Cn. Casabonae.
tryphilinum ×, Treuinf. in Zeitschr. Ferdin. Tirol. III. xix. (1875) 276.—Europ.
tuberosum, All. Fl. Pedem. i. 151 = Cn. tuberosus.

CIRSIUM :—
ukranicum, Bess. ex DC. Prod. vi. 635 = Cnicus serrulatus.
uliginosum, Bieb. Fl. Taur. Cauc. iii. 555 = Cn. uliginosus.
uliginosum, Heldr. ex Boiss. Diagn. Ser. I. x. 90 = *libanoticum*.
undulatum, Spreng. Syst. iii. 374 = Cn. undulatus.
Vallis-Demonii, Lojac. in Natural. Sicil. iii. (1884) 267.—Sicilia.
variabile, Moench, Meth. 558 = Cn. oleraceus.
Verutum, Spreng. Syst. iii. 370 = Cn. argyracanthus.
virens, Timb. & Jeanb. ex Nym. Consp. 406 = Cn. lanceolatus.
virgatum, Tausch, in Flora, xix. (1836) 402.—Tauria.
virginianum, Hook. Comp. Bot. Mag. i. (1835) 48 = Cn. Lecontei.
virginianum, Michx. Fl. Bor. Am. ii. 90 = Cn. virginianus.
Vlassovianum, Fisch. ex DC. Prod. vi. 653 = Cn. Vlassovianus.
Wallichii, DC. l. c. 643 = Cn. Wallichii.
Wankelii ×, Reichardt, in Verh. Zool.-Bot. Ges. Wien, xi. (1861) 381.—Europ.
Welwitschii, Coss. Notes Crit. 118.—Hispania.
Weyrichii, Maxim. Prim. Fl. Amur. 174 = Cn. Weyrichii.
Wiedemanni, Fisch. Mey. & Avé-Lall. Ind. Sem. Hort. Petrop. vi. 48 = *hypoleucum*.
Willdenowii, Steud. Nom. ed. II. i. 370 = Cn. Willdenowii.
Wimmeri, Čelak. in Arch. Naturfw. Landesd. Boehm. ii. (1873) 263.—Bohemia.
Winklerianum ×, Čelak. l. c. 262.—Europ.
Wirtgenii ×, Loehr, Enum. Fl. Deutsch. 362.—Europ.
Woodwardii, H. C. Wats. ex Nym. Consp. 407 = Cn. tuberosus.
Wrightii, A. Gray, Pl. Wright. ii. 101 = Cn. Wrightii.
Zizianum, Koch, Syn. Fl. Germ. ed. I. 398 = Cn. tuberosus.

CISSAMPELOPSIS, Miq. Fl. Ind. Bat. ii. 102 (1856) = **Senecio**, Tourn. (Compos.).
volubilis, Miq. l. c. = S. araneosus.

CISSAMPELOS, Linn. Gen. ed. I. 368 (1737).
MENISPERMACEAE, Benth. & Hook. f. i. 37.
ANTIZOMA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 41.
CAAPEBA, Plum. ex Adans. Fam. ii. 357 (1763).
DISSOPETALUM, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xvii. (1866) 267.
Abutua, Vell. Fl. Flum. x. t. 140 = Chondodendron platyphyllum.
acuminata, Benth. Pl. Hartw. 58 = C. Pareira.
acuminata, *DC. Syst.* i. 538.—Ind. or.
acuta, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 43.—N. Granat.
amazonica, Miers, ex Benth. in Hook. Kew Journ. iii. (1851) 114 = ovalifolia.
andromorpha, *DC. Syst.* i. 539.—Guiana.
angustifolia, *Burch. Trav.* i. 389.—Afr. austr.
angustifolia, Drège, ex Harv. & Sond. Fl. Cap. i. 13 (= *Antizoma Miersiana*).—Afr. austr.
apiculata, Hochst. in Flora, xxviii. (1845) 93 = C. Pareira.
argentea, H. B. & K. Nov. Gen. et Sp. v. 67 = C. Pareira.
aristolochiaefolia, Fenzl, in Flora, xxvii. (1844) 312 = C. Pareira.
assimilis, Miers, ex Benth. in Hook. Kew Journ. iii. (1851) 114 = ovalifolia.
auriculata, *Miers, Contrib.* iii. 158.—Reg. Argent.
australis, St. Hil. Fl. Bras. Mer. i. 54 = C. Pareira.
barbata, Wall. Cat. n. 4978 = Cyclea peltata.
Benthamiana, *Miers, Contrib.* iii. 144.—Mexic.
Boivini, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 460.—Madag.
Bojeriana, *Miers, Contrib.* iii. 182.—Ins. Maurit.
Caapeba, Linn. Sp. Pl. 1032 = C. Pareira.
Caapeba, Vell. Fl. Flum. x. t. 139 = fasciculata ?
calcarifera, Burch. Trav. ii. 266 = fruticosa.
canescens, Miq. Sert. Exot. i. 7. t. 4 = C. Pareira.
capensis, *Thunb. Prod. Pl. Cap.* 110.—Afr. austr.
capitata, *Griff. Notul.* iv. 309.—Ind. or.
clematidea, Presl, Bot. Bemerk. 7 = C. Pareira.
Cocculus, Buch.-Ham. ex Steud. Nom. ed. II. i. 371 = C. Pareira.

CISSAMPELOS :—
Cocculus, *Poir. Encyc.* v. 9.—Ind. or.
Cocculus, Steud. Nom. ed. I. 198 = Anamirta paniculata.
comata, Miers, in Hook. Niger Flora, 215 = C. Pareira.
communis, St. Hil. Fl. Bras. Mer. i. 52 = ovalifolia.
consociata, *Miers, Contrib.* iii. 167.—Ind. occ.
convexa, Vell. Fl. Flum. x. t. 142 = Abuta rufescens.
convolvulacea, Willd. Sp. Pl. iv. 863 = C. Pareira.
cordifolia, *Boj. Ann. Sc. Nat.* Sér. II. xx. (1843) 54.—Afr. austr.
crenata, DC. Syst. i. 537 = ovalifolia.
Cumingiana, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 283.—Ins. Philipp.
delicatula, Miers, Contrib. iii. 197 = C. Pareira.
denudata, Miers, ex Benth. in Hook. Kew Journ. iii. (1851) 115 = fasciculata.
diffusa, *Miers, Contrib.* iii. 168.—Ind. occ.
discolor, DC. Syst. i. 534 = Stephania hernandifolia.
discolor, A. Gray, Bot. U. St. Expl. Exped. i. 38 = C. Pareira.
discolor, Wall. Cat. n. 4982 = Cyclea peltata.
diversa, Miers, Contrib. iii. 187 = C. Pareira.
ebracteata, St. Hil. Pl. Us. Bras. t. 35 = ovalifolia.
elata, Miers, Contrib. iii. 187 = C. Pareira.
eriantha, Miers, l. c. 192 = C. Pareira.
eriocarpa, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 41 = C. Pareira.
errabunda, *Miers, Contrib.* iii. 138.—Bras.
fasciculata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 361.—Bras.
floribunda, *Miers, Contrib.* iii. 164.—Peruv.
fluminensis, *Eichl. in Flora*, xlvii. (1864) 392 [*err. typ.* '382';] *et in Mart. Fl. Bras.* xiii. I. 191.—Bras.
fruticosa, Thunb. Prod. Pl. Cap. 110 = capensis.
glaberrima, *St. Hil. Fl. Bras. Mer.* i. 57.—Bras.
glabra, Buch.-Ham. ex Wight & Arn. Prod. i. 14 = Clypea hernandifolia et Cocculus Roxburghianus.
glabra, Roxb. Hort. Beng. 74; Fl. Ind. iii. 840 = Stephania rotunda.
glaucescens, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 41 = C. Pareira.
gracilis, St. Hil. Fl. Bras. Mer. i. 56 = C. Pareira.
grallatoria, Miers, Contrib. iii. 189 = C. Pareira.
grandifolia, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 44 = C. Pareira.
guayaquilensis, H. B. & K. Nov. Gen. et Sp. v. 67 = C. Pareira.
Haenkeana, Presl, Rel. Haenk. ii. 80 = ovalifolia.
hederacea, *Miers, Contrib.* iii. 159.—Reg. Argent.
Hernandia, *Vell. Fl. Flum.* x. t. 136.—Bras.
hernandifolia, Wall. Cat. 4977, partim = C. Pareira.
hernandifolia, Willd. Sp. Pl. iv. 861 = Stephania hernandifolia.
heterophylla, DC. Syst. i. 534 = C. Pareira.
hexandra, Roxb. Hort. Beng. 74; Fl. Ind. iii. 841 = Stephania hernandifolia.
hirsuta, Buch.-Ham. in DC. Syst. i. 535 = C. Pareira.
hirsutissima, Presl, Rel. Haenk. ii. 80 = ovalifolia.
hirta, Klotzsch, in Peters, Reise Mossamb. Bot. 174 = C. Pareira.
humilis, Poir. Encyc. v. 11 = capensis.
hypoglauca, *Schau. in Nov. Act. Acad. Nat. Cur.* xix. *Suppl.* i. (1843) 479.—China.
insolita, *Miers, Contrib.* iii. 179.—Afr. trop.
kohautiana, Presl, Rel. Haenk. ii. 81 = C. Pareira.
laurifolia, *Poir. Encyc.* v. 11.—Ind. occ.
limbata, *Miers, Contrib.* iii. 143.—Am. trop.
littoralis, St. Hil. Fl. Bras. Mer. i. 54 = C. Pareira.
longipes, *Miers, Contrib.* iii. 138.—Ind. occ.
macrostachya, Klotzsch, in Peters, Reise Mossamb. Bot. 172 = C. Pareira.
madagascariensis, *Miers, Contrib.* iii. 181.—Madag.; Ins. Borb.
mallophylla, *Miers, l. c.* 169.—Am. trop.
mauritiana, Thou. in Desv. Journ. Bot. ii. (1809) 67. t. 3, 4 = C. Pareira.
mauritiana, Wall. Cat. n. 4980 = Pericampylus incanus.
microcarpa, DC. Syst. i. 534 = C. Pareira.
monoica, St. Hil. Fl. Bras. Mer. i. 55 = C. Pareira.
mucronata, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 11 = C. Pareira.
myriocarpa, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 42 = C. Pareira.
nephrophylla, Boj. l. c. Sér. II. xx. (1843) 55 = C. Pareira.

CISSAMPELOS:—
nympheaefolia, R. Br. in Salt, Abyss. App. 65 = Stephania abyssinica.
obtecta, Wall. Cat. n. 4981 = C. Pareira.
oleracea, Wall. l. c. n. 4984 = Parabaena sagittata.
orbiculata, DC. Syst. i. 537 = C. Pareira.
orinocensis, H. B. & K. Nov. Gen. et Sp. v. 68 = C. Pareira.
ovalifolia, *DC. Syst.* i. 537.—Bras.
ovata, Poir. Encyc. v. 10 = Pachygone ovata.
ovata, Vell. Fl. Flum. x. t. 141 = Abuta Selloana?
owariensis, Beauv. ex DC. Prod. i. 100 = C. Pareira.
pannosa, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 283.—Cuba.
Pareira, *Linn. Sp. Pl.* 1031.—Reg. trop.
pareiroides, DC. Ess. Med. Pl. Comp. ed. II. 78 = C. Pareira.
Parriera, Vell. Fl. Flum. x. t. 138 = glaberrima.
Pata, Roxb. ex Wight & Arn. Prod. i. 15 = C. Pareira.
pauciflora, Nutt. in Journ. Acad. Philad. vii. (1834) 114 = Croomia pauciflora.
peltata, Ruiz, ex Eichl. in Mart. Fl. Bras. xiii. 1. 191 = tropaeolifolia.
psilophylla, *Presl, Rel. Haenk.* ii. 80.—Ins. Luzon.
ramiflora, *Miers, Contrib.* iii. 163.—Bras.
rotundata, Pohl, ex Eichl. in Mart. Fl. Bras. xiii. 1. 187 = ovalifolia.
sagittata, Buch.-Ham. ex Wall. Cat. n. 4983 = Parabaena sagittata.
Salzmanni, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 284.—Bras.
scutigera, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 42 = C. Pareira.
senensis, Klotzsch, in Peters, Reise Mossamb. Bot. i. 173 = C. Pareira.
smilacina, Linn. Sp. Pl. 1032 = Menispermum canadense.
subcrenata, *Klotzsch, in Schomb. Fauna, Fl. Gui.* 1162, nomen.—Guiana.
suborbicularis, St. Hil. Fl. Bras. Mer. i. 53 = ovalifolia.
subpeltata, Thw. Enum. Pl. Zeyl. 13, 399 = C. Pareira.
subreniformis, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 41 = C. Pareira.
subtriangularis, St. Hil. Fl. Bras. Mer. i. 51 = ovalifolia.
sympodiales, *Eichl. in Flora*, xlvii. (1864) 382 [err. typ. 392].—Bras.
tamnifolia, *Miers, Contrib.* iii. 185.—Afr. austr.
tamoides, Sagot, ex Eichl. in Mart. Fl. Bras. xiii. 1. 194 = fasciculata.
tamoides, *Willd. ex DC. Syst.* i. 536.—Bras.
testudinaria, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xvii. (1866) 134, nomen = seq.?
testudinum, *Miers, Contrib.* iii. 143.—Ins. Galapag.
tetrandra, Roxb. Hort. Beng. [105]; Fl. Ind. iii. 842 = C. Pareira.
tomentosa, DC. Syst. i. 535 = C. Pareira.
tomentosa, Vell. Fl. Flum. x. t. 143 = Abuta rufescens.
torulosa, *E. Mey. ex Harv. & Sond. Fl. Cap.* i. 11.—Afr. austr.
triloba, *Spreng. Neue Entdeck.* ii. 152.—Bras.
tropaeolifolia, *DC. Syst.* i. 532.—Am. austr.
umbellata, E. Mey. ex Harv. & Sond. Fl. Cap. i. 10 = Stephania hernandifolia.
velutina, St. Hil. Fl. Bras. Mer. i. 52 = ovalifolia.
vestita, Triana & Planch. in Ann. Sc. Nat. Sér. IV. xvii. (1862) 44 = ovalifolia.
Vitis, *Vell. Fl. Flum.* x. t. 137.—Bras.
Vogelii, Miers, in Hook. Niger Fl. 214 = C. Parcira.
zairensis, *Miers, Contrib.* iii. 180.—Afr. trop.

CISSABRYON, Meissn. Gen. 58; Comm. 42 (1837) = seq.?

CISSAROBRYON, Poepp. Fragm. Syn. Fl. Chil. 29 (1833) = **Viviania**, Cav. (Geraniac.).
elegans, Kunze, l. c. 30.—Chili.
macrophyllum, Phil. in Linnaea, xxxiii. (1864-65) 32.—Chili.
parvifolium, Phil. l. c. 33.—Chili.

CISSODENDRON, F. Muell. Census, 61 (1882) = KISSODENDRON, Scem. = **Hedera**, Tourn. (Araliac.).

CISSUS, Linn. Diss. Dass. 6 (1747); Amoen. Acad. i. 389 (1749) = **Vitis**, Tourn. (Ampelid.).
acetosa, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 24 = V. acetosa.
acida, Linn. Sp. Pl. ed. II. 170 = V. acida.
acida, Roxb. ex Wight & Arn. Prod. 127 = V. trifolia.
acida, Wall. Cat. n. 6009 B = V. setosa.
aconitifolia, Walp. Rep. i. 441 = V. serjaniaefolia.
aculeata, Span. in Linnaea, xv. (1841) 184 = V. aculeata.
acuminata, A. Gray, Bot. U. St. Expl. Exped. i. 273 = V. acuminata.
acuminata, Thw. Enum. Pl. Zeyl. 62 = V. repanda.
acutifolia, Poir. Encyc. Suppl. i. 106 = V. trifolia.
adenantha, Fresen. in Mus. Senckenb. ii. (1837) 283 = V. adenantha.
adenocaulis, Steud. ex A. Rich. Tent. Fl. Abyss. i. 111 = V. adenocaulis.
adnata, Roxb. Hort. Beng. 11; Fl. Ind. i. 405 = V. adnata.
adnata, Wight, ex Wall. Cat. sub n. 5988 = V. Heyneana.
aegirophylla, Bunge, in Arb. Naturf. Ver. Riga, i. (1847) 193 = V. aegirophylla.
alata, Blanco, Fl. Filip. ed. II. 51 = V. geniculata.
alata, Jacq. Select. Am. 23 = V. sulcicaulis?
albida, Cambess. in St. Hil. Fl. Bras. Mer. i. 344 = V. albida.
altissima, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 74 = V. papuana.
amazonica, Linden, in Belg. Hortic. xv. (1865) 99.—Bras.
amboinensis, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 76 = V. amboinensis.
Ampelopsis, Pers. Syn. i. 142 = V. indivisa.
anaimalaiensis, Bedd. in Madr. Journ. Sc. Ser. III. i. (1864) 41 = V. anamallayana.
anemonifolia, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 80 = V. anemonifolia.
angulata, Lam. Tabl. Encyc. i. 331 = V. Linnaei.
angustifolia, Roxb. Hort. Beng. 11; Fl. Ind. i. 408 = V. angustifolia.
antarctica, Vent. Choix, 21. t. 21 = V. antarctica.
aquosa, Buch.-Ham. ex Wall. Cat. n. 6000 = V. repanda.
arachnoidea, Hassk. Cat. Hort. Bog. Alt. 166 = V. indica.
araneosa, Miq. Fl. Ind. Bat. Suppl. i. 517 = V. araneosa.
arborea, Blanco, Fl. Filip. ed. II. 51 = Evodia Roxburghiana.
arborea, Desmoul. in Durand, Monog. Vit. 60 = V. arborea.
arborea, Forsk. Fl. Aegypt. Arab. 32 = Salvadora persica.
arborea, Willd. ex Roem. & Schult. Syst. iii. Mant. 248 = Cybianthus Humboldtii.
argenteus, Linden, in Belg. Hortic. xvii. (1867) 103.—Bras.
arguta, Hook. f. in Hook. Niger Fl. 261 = V. arguta.
aristata, Blume, Bijdr. 183 = V. adnata.
articulata, Guill. & Perr. Fl. Seneg. Tent. i. 135, in obs.—Afr. trop.
assimilis, Kurz, ex Laws. in Hook. f. Fl. Brit. Ind. i. 660 = V. lanceolaria.
auriculata, DC. Prod. i. 632 = V. auriculata.
auriculata, Roxb. Hort. Beng. 11; Fl. Ind. i. 411 = V. trifolia.
australasica, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 8 = V. hypoglauca.
Baudiniana, Brouss. ex DC. Prod. i. 629 = V. antarctica.
bifida, Schum. & Thonn. Beskr. Guin. Pl. 80 = V. quadrangularis.
bigemina, Harv. in Harv. & Sond. Fl. Cap. i. 253 = V. gracilis.
biglandulosa, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 75 = V. pubiflora.
bipinnata, Nutt. Gen. Am. i. 144 = V. arborea.
Blumeana, Hassk. in Flora, xxv. (1842) II. Beibl. 39 = V. indica.
Blumeana, Span. in Linnaea, xv. (1841) 184.—Java.
Blumeana, Steud. Nom. ed. II. i. 371 = V. pallida.
bonariensis, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 159 = V. striata.
bororensis, Klotzsch, in Peters, Reise Mossamb. Bot. 179 = V. bororensis.
brevipedunculata, Maxim. Prim. Fl. Amur. 68 = V. heterophylla.

CISSUS:—
bryonifolia, Regel, Fl. Ussur. t. 3. f. 3 = Vitis heterophylla.
caesia, Afzel. Rem. Guin. 55 = V. caesia.
canescens, Lam. Tabl. Encyc. i. 331.—Peruv.
cantoniensis, Hook. & Arn. Bot. Beech. Voy. 175 = V. cantoniensis.
capensis, Willd. Sp. Pl. i. 655 = V. capensis.
capreolata, Walp. Rep. i. 439 = V. capreolata.
carnosa, Lam. Encyc. i. 31 = V. trifolia.
caustica, Tussac, Fl. Ant. i. 116. t. 16 = V. trifoliata, *Baker*.
cerifera, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 418 = V. cerifera.
cinerea, Lam. Tabl. Encyc. i. 332 = V. trifolia.
cirrhiflora, Eckl. & Zeyh. Enum. 56 = *pauciflora*.
cirrhosa, Willd. Sp. Pl. i. 657 = V. cirrhosa.
clematidifolia, Zipp. ex Span. in Linnaea, xv. (1841) 184 = V. aculeata.
clematifolia, Walp. Rep. i. 439 = V. aculeata.
cochinchinensis, Spreng. Syst. i. 450.—Cochinchin.
compressa, Blume, Bijdr. 182 = V. adnata.
compressicaulis, Ruiz & Pav. Fl. Per. i. 64. t. 100.—Peruv.
connivens, Lam. Tabl. Encyc. i. 332.—Madag.
cordata, Roxb. Hort. Beng. 11; Fl. Ind. i. 407 = V. glauca, repens.
cordata, Wall. Cat. n. 6001 = V. adnata.
cordifolia, Linn. Sp. Pl. ed. II. 170.—Am. austr.
coriacea, DC. Prod. i. 632 = V. coriacea.
crenata, Blume, Bijdr. 186 = *Blumeana*.
crenata, Vahl, Symb. Bot. iii. 19 = V. trifolia.
cuneifolia, Eckl. & Zeyh. Enum. 56.—Afr. austr.
Currori, Hook. f. in Hook. Niger Fl. 265 = V. Currori.
cyanocarpa, Miq. Fl. Ind. Bat. i. II. 603 = V. japonica.
cymosa, Schum. & Thonn. Beskr. Guin. Pl. 82 = V. Thonningii.
cymosa, Steud. Nom. ed. II. i. 372 = V. tenuifolia.
cyphopetala, Fresen. in Mus. Senckenb. ii. (1837) 282 = V. cyphopetala.
Darik, Miq. Fl. Ind. Bat. Suppl. i. 515.—Sumatra.
Davidiana, Carr. in Rev. Hortic. (1868) 29.—China.
dealbata, Bedd. in Madr. Journ. Sc. Scr. II. xxii. (1861) 74.—Ind. or.
deficiens, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 160 = V. striata.
dendroides, Roem. & Schult. Syst. iii. Mant. 248 = Ardisia acuminata.
denticulata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) 1. 591 = V. Afzelii?
dichotoma, Blume, Bijdr. 186 = V. dichotoma.
dichrothrix, Miq. Fl. Ind. Bat. Suppl. i. 514.—Sumatra.
Diepenhorstii, Miq. l. c. 515 = V. Diepenhorstii.
digitata, Lam. Tabl. Encyc. i. 332.—Arabia.
dimidiata, Eckl. & Zeyh. Enum. 57.—Afr. austr.
dioica, Roxb. ex DC. Prod. i. 632 = V. lanceolaria.
discolor, Blume, Bijdr. 181 = V. discolor.
diversifolia, DC. Prod. i. 631 (= *heterophylla*, Link).—Hab.?
diversifolia, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 314 = V. cantoniensis.
Dregeana, Bernh. in Flora, xxvii. (1844) 297 = V. capensis.
Duarteana, Cambess. in St. Hil. Fl. Bras. Mer. i. 343 = V. Duarteana.
dubia, Steud. Nom. ed. II. i. 372.—China.
edulis, Dalz. in Hook. Kew Journ. ix. (1857) 248 = V. quadrangularis.
elegans, Hort. ex C. Koch, Dendrol. i. 555 = V. heterophylla.
elliptica, Cham. & Schlecht. in Linnaea, v. (1830) 221 = V. elliptica.
elongata, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) 110 = V. Miqueliana.
elongata, Roxb. Hort. Beng. 11; Fl. Ind. i. 411 = V. elongata.
emarginella, Sw. in Vet. Akad. Handl. Stockl. (1825) 427 = V. crenata.
Endresii, Hort.—Cf. Gard. Chron. (1876) i. 522.—Costa Rica.
enneaphylla, Vell. Fl. Flum. 40; i. t. 103 = V. enneaphylla.
epidendrica, Vell. l. c. 39; i. t. 98 = V. trifoliata.
Epidendron, Eichl. in Mart. Fl. Bras. xiv. II. 217 = prace.

CISSUS:—
erosa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = Vitis erosa.
erythrodes, Hochst. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 79 = V. erythrodes.
feminea, Roxb. Hort. Beng. 11; Fl. Ind. i. 410 = V. lanceolaria.
ferruginea, E. Mey. Zwei Pfl. Docum. 173 = *Thunbergii*.
ferruginea, Willd. ex Roem. & Schult. Syst. iii. Mant. 248 = Ardisia ferruginea.
Figariana, Webb, Fragm. Fl. Aethiop. 58 = V. Figariana.
flexuosa, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 415.—Ins. Luzon.
foemina, Steud. Nom. ed. II. i. 372 = V. lanceolaria.
fragariaefolia, Boj. Hort. Maurit. 60.—Zanzibar.
fragilis, E. Mey. Zwei Pfl. Docum. 173, nomen; Harv. & Sond. Fl. Cap. i. 249.—Afr. austr.
frutescens, Blanco, Fl. Filip. ed. I. 70 = Evodia Roxburghiana.
fuliginea, H. B. & K. Nov. Gen. et Sp. v. 224.—Am. austr.
Gardneri, Thw. Enum. Pl. Zeyl. 63 = V. Gardneri.
geniculata, Blume, Bijdr. 184 = V. geniculata.
gigantea, Bedd. in Madr. Journ. Sc. xxii. Ser. II. (1861) 75.—Ind. or.
glaberrima, Steud. Nom. ed. II. i. 372 = V. glaberrima.
glabra, Herb. Madr. ex Wall. Cat. sub n. 5988 = V. Heyneana.
glabra, E. Mey. Zwei Pfl. Docum. 173 = V. orientalis.
glabrata, Blume, Bijdr. 189 = V. serrulata.
glandulosa, J. F. Gmel. Syst. 256.—Arabia.
glandulosa, Poir. Encyc. Suppl. i. 105 = V. antarctica.
glandulosa, Roxb. Hort. Beng. 11, nomen.—Ind. or.
glauca, Roxb. l. c. 11; Fl. Ind. i. 406 = V. glauca.
glauca, Thw. Enum. Pl. Zeyl. 62 = V. pallida?
glaucophylla, Hook. f. in Hook. Niger Fl. 263 = V. glaucophylla.
glyptocarpa, Thw. Enum. Pl. Zeyl. 62 = V. glyptocarpa.
gongylodes, Burch. ex Baker, in Mart. Fl. Bras. xiv. ii. 209 = V. gongylodes.
gracilis, Guill. & Perr. Fl. Seneg. Tent. i. 134 = V. gracilis, *Baker*.
granulosa, Ruiz & Pav. Fl. Per. i. 62 = V. striata.
haematantha, Miq. in Linnaea, xxvi. (1853) 220.—Guiana.
Hahniana, Ernst, in Journ. Bot. xiv. (1876) 179.—Venezuela.
hastata, Miq. Fl. Ind. Bat. Suppl. i. 517 = V. hastata.
hederacea, Pers. Syn. i. 143 = V. hederacea.
hederacea, Royle, ex Walp. Rep. i. 439 = V. capreolata.
helleborifolia, Zipp. ex Span. in Linnaea, xv. (1841) 184 = V. auriculata.
heptaphylla, Retz. Obs. v. 22 = V. pedata.
heteroma, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 591.—Java.
heterophylla, Herb. Madr. ex Wall. Cat. n. 6036 = V. Rheedei.
heterophylla, Link, Enum. Hort. Berol. i. 143 = *diversifolia*, DC.
heterophylla, Poir. Encyc. Suppl. i. 107.—Java.
heterophylla, Walp. Rep. i. 441 = V. Landuk.
heterotoma, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 591.—Java.
Heyneana, Steud. Nom. ed. II. i. 372 = V. Heyneana.
himalayana, Walp. Rep. i. 441 = V. himalayana.
hirsuta, Steud. Nom. ed. 1. 199 = V. hederacea.
hirtella, Blume, Bijdr. 185 = V. geniculata.
humulifolia, Regel, Tent. Fl. Ussur. 36 = V. heterophylla.
hydrophora, Gaud. Bot. Voy. Bonite, t. 131.—Hab.?
hypoglauca A. Gray, Bot. U. St. Expl. Exped. i. 272 = V. hypoglauca.
hypoleuca, Harv. in Harv. & Sond. Fl. Cap. i. 252.—Afr. austr.
ibuensis, Hook. f. in Hook. Niger Fl. 265 = V. ibuensis.
inaequalis, Heyne, ex Wall. Cat. n. 6010 = V. inaequalis.
incarnata, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 417 = V. incarnata.
incisa, Desmoul. in Durand, Monog. Vit. 59, ex S. Wats. Bibliog. Ind. N. Am. Bot. 173 = V. incisa.
indica, Koen. ex Steud. Nom. ed. II. i. 371, 372 = V. Linnaei.

CISSUS:—
indica, Rottl. in Ges. Naturf. Freunde, Neue Schr. iv. (1803) 183 = Vitis repanda.
indica, Walp. Rep. i. 441 = V. indica.
indivisa, Desmoul. in Durand, Monog. Vit. 59, ex S. Wats. Bibliog. Ind. N. Am. Bot. 173 = V. indivisa.
intermedia, A. Rich. Ess. Fl. Cub. 311 = V. trifoliata, *Baker*.
involucrata, Spreng. Syst. iv. Cur. Post. 44 = Pterisanthes cissoides.
ipomaeifolia, Webb, Fragm. Fl. Aethiop. 57 = V. ipomoeifolia.
Jackii, Korth. ex Miq. in Ann. Mus. Bot. Lugd. Bat. i. 81 = V. geniculata.
japonica, Willd. Sp. Pl. i. 659 = V. japonica.
javana, DC. Prod. i. 628.—Java.
juncea, Webb, Fragm. Fl. Aethiop. 57 = V. juncea.
Kleinii, Laws. in Hook. f. Fl. Brit. Ind. i. 649, in syn. = V. adnata.
laciniata, Linn. Syst. ed. X. 897 = Sicyos laciniata.
laevigata, Blume, Bijdr. 191 = V. auriculata.
Lamarckiana, Roem. & Schult. Syst. iii. Mant. 239 = V. sicyoides.
lanceolaria, Roxb. Hort. Beng. 11; Fl. Ind. i. 412 = V. lanceolaria.
Landuk, Hassk. in Flora, xxv. (1842) ii. Beibl. 39 = V. Landuk.
lanigera, Harv. in Harv. & Sond. Fl. Cap. i. 252.—Afr. austr.
latifolia, Descourt. Fl. Med. Ant. v. 29. t. 11 = V. sicyoides.
latifolia, Lam. Encyc. i. 30 = V. glauca.
latifolia, Vahl, Symb. Bot. iii. 18 = V. adnata.
latifolia, Walp. Rep. i. 441 = V. latifolia.
leonensis, Hook. f. in Hook. Niger Fl. 264 = V. leonensis.
leucocarpa, Blume, Bijdr. 189 = V. japonica.
leucostaphyla, Steud. Nom. ed. II. i. 372 = V. lanceolaria.
Lindeni, André, Illustr. Hortic. xvii. (1870) 12. t. 2.—N. Granat.
Livingstoniana, Welw. in Journ. Linn. Soc. viii. (1865) 76.—Afr. trop.
lonchiphylla, Thw. Enum. Pl. Zeyl. 62 = V. lonchiphylla.
longepedunculata, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxv. (1863) 417 = V. longepedunculata.
lucida, Poir. Encyc. Suppl. i. 106 = V. erosa.
macrobotrys, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 269.—Sumatra.
macrophylla, Jungh. in Hoev. & De Vriese, Tijdschr. vii. (1840) 306.—Java.
macropus, Welw. in Journ. Linn. Soc. viii. (1865) 77 = V. macropus.
Mappia, Lam. Tabl. Encyc. i. 332 = V. Mappia.
membranacea, Hook. f. in Hook. Niger Fl. 266 = V. gracilis.
mexicana, Moç. & Sesse, ex DC. Prod. i. 631 = V. mexicana.
micrantha, Poir. Encyc. Suppl. i. 105.—Ins. S. Doming.
micrantha, Willd. ex Roem. & Schult. Syst. iii. Mant. 248 = *parviflora*.
microcarpa, Vahl, Eclog. Am. i. 16 = V. trifoliata, *Baker*.
microphylla, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 416.—Afr. austr.
mollis, Steud. ex Baker, in Oliver, Fl. Trop. Afr. i. 413 = V. serpens.
mucronata, Hook. & Arn. ex Steud. Nom. ed. II. i. 372 = V. lanceolaria.
multibracteata, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 75 = V. pubiflora.
multiflora, Willd. ex Roem. & Schult. Syst. iii. Mant. 249.—Am. austr.
muricata, Dalz. & Gibs. Bomb. Fl. 40; Thw. Enum. Pl. Zeyl. 63 = V. lanceolaria.
mutabilis, Blume, Bijdr. 190 = V. mutabilis.
napaulensis, DC. Prod. i. 632 = V. capreolata.
natalensis, Bernh. ex Krauss, in Flora, xxvii. (1844) 349 = Rhus natalensis.
neilgherrensis, Wight, Spicil. Neilgh. i. 32. t. 36 = V. himalayana.
nepalensis, Steud. Nom. ed. II. i. 372, vide *napaulensis*.
nilagirica, Miq. in Linnaea, xxvi. (1853) 221.—Ind. or.
nitida, Vell. Fl. Flum. 39; i. t. 100 = V. sicyoides.

CISSUS:—
nivea, Hochst. ex Schweinf. Beitr. Fl. Aethiop. 83.—Abyssin.
nodosa, Blume, Bijdr. 182 = Vitis nodosa.
obliqua, Ruiz & Pav. Fl. Per. i. 65.—Peruv.
obovata, Vahl, Symb. Bot. iii. 19 = V. trifoliata, *Baker*.
obscura, DC. Prod. i. 629 = V. sicyoides.
obtusata, Benth. Bot. Voy. Sulph. 77.—Panama.
obtusifolia, Lam. Encyc. i. 31 = V. trifolia.
ochracea, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 35 = V. ochracea.
officinalis, Klotzsch, in Bot. Zeit. iv. (1846) 110.—N. Granat.
opaca, F. Muell. in Trans. Phil. Inst. Vict. iii. (1859) 23 = V. opaca.
orientalis, Lam. Tabl. Encyc. i. 332. t. 84. f. 2 = V. orientalis.
ovata, Lam. l. c. 331 = V. sicyoides.
ovata, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106.—Guiana.
oxyodon, Planch. in Catal. Pl. Hort. Donat. 5.—Hab.?
pallida, Salisb. Prod. 66 = V. sicyoides.
pallida, Steud. Nom. ed. II. i. 372 = V. pallida.
palmata, Poir. Encyc. Suppl. i. 107 = V. Commersoni.
paniculata, Heyne, in Wall. Cat. n. 6022 = V. tenuifolia.
papillosa, Blume, Bijdr. 183 = V. pubiflora.
parviflora, Roem. & Schult. Syst. iii. Mant. 248.—Ind. or.
parvifolia, Salisb. Prod. 66 = V. acida.
paucidentata, Ernst, in Journ. Bot. viii. (1870) 374.—Venezuela.
paucidentata, Klotzsch, in Peters, Reise Mossamb. Bot. i. 179 = V. paucidentata.
pauciflora, DC. Prod. i. 630.—Afr. austr.
Pauli-Guilielmi, Schweinf. Rel. Kotsch. 46.—Afr. trop.
paullinifolia, Vell. Fl. Flum. 40; i. t. 102 = V. ternata?
pedata, Lam. Encyc. i. 31 = V. pedata.
pedata, Naves, ex Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 50 = V. lanceolaria.
peltata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) i. 590.—Java.
pentagona, Roxb. Hort. Beng. 11; Fl. Ind. i. 408 = V. pentagona.
pentandra, Willd. ex Roem. & Schult. Syst. iii. Mant. 248.—Am. austr.
pentaphylla, Buch.-Ham. ex Wall. Cat. sub n. 6016 = V. elongata.
pentaphylla, Herb. Madr. ex Wall. l. c. sub n. 6022 = V. tenuifolia.
pentaphylla, Willd. Sp. Pl. i. 659 = V. pentaphylla.
pergamacea, Blume, Bijdr. 183 = V. pergamacea.
petiolata, Hook. f. in Hook. Niger Fl. 262 = V. pallida.
pinnata, [Soland. in] Russell, Aleppo, ed. II. ii. 267.—Cilicia.
pisocarpa, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 79 = V. pisicarpa.
platanifolia, Carr. in Rev. Hortic. (1868) 39.—China.
populnea, Guill. & Perr. Fl. Seneg. Tent. i. 134 = V. pallida.
porphyrophylla, Lindl. in Proc. Hort. Soc. i. (1861) 225 = Piper porphyrophyllum.
producta, Afzel. Rem. Guin. 63 = V. producta.
pruinata, Weinm. in Syll. Ratisb. i. (1824) 224.—Bras.
pruinosa, Desf. Cat. Hort. Par. ed. III. 238.—Hab.?
pubescens, H. B. & K. Nov. Gen. et Sp. v. 226.—Am. austr.
pubescens, Walp. Rep. i. 441 = V. pubescens.
pubiflora, Miq. Fl. Ind. Bat. Suppl. i. 516 = V. pubiflora.
pubinervis, Blume, Bijdr. 185.—Java.
pulcherrima, Vell. Fl. Flum. 40; i. t. 101 = V. pulcherrima.
puncticulosa, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = V. sicyoides.
purpurea, Roxb. ex Steud. Nom. ed. II. i. 373 = V. glauca.
purpurascens, Zipp. ex Miq. Ann. Mus. Bot. Lugd. Bat. i. 85 = V. rostrata.
pyrrhodasys, Miq. Fl. Ind. Bat. Suppl. i. 517.—Sumatra.
quadrangula, Salisb. Prod. 66 = V. quadrangularis.
quadrangularis, Linn. Mant. i. 39 = V. quadrangularis.

CISSUS :—
quadrialata, H. B. & K. Nov. Gen. et Sp. v. 225 = Vitis erosa.
quinata, [Dryand. in] Ait. Hort. Kew. ed. II. i. 260 = V. eirrhosa.
quinquefolia, Desf. Cat. Hort. Par. ed. III. 238 = V. hederacea.
quinquefolia, Soland. in Sims, Bot. Mag. t. 2443 = V. Simsiana.
repanda, Vahl, Symb. Bot. iii. 18 = V. repanda.
repens, Lam. Encyc. i. 31 = V. repens.
repens, Thw. Enum. Pl. Zeyl. 62 = V. repanda.
reticulata, Willd. ex Roem. & Schult. Syst. iii. Mant. 248 = *canescens*.
reticulata, Thw. Enum. Pl. Zeyl. 63 = V. reticulata.
rhodocarpa, Blume, Bijdr. 185 = V. geniculata.
rhombifolia, Vahl, Eclog. Am. ii. 10 = V. rhombifolia.
rhomboidea, E. Mey. Zwei Pfl. Docum. 173, nomen ; Harv. & Sond. Fl. Cap. i. 252.—Afr. austr.
riparia, Laws. in Hook. f. Fl. Brit. Ind. i. 649, in syn. = V. repanda.
Rocheana, Planch. in Comptes-rendus, xciii. (1881) 369.—Afr. trop.
rosea, Walp. Rep. i. 437 = V. repanda.
rotundata, Heyne, ex Wall. Cat. sub n. 5988 = V. Heyneana.
rotundifolia, Blume, Bijdr. 480 = V. pallida.
rotundifolia, Vahl, Symb. Bot. iii. 19.—Arabia.
rubescens, Blanco, Fl. Filip. ed. I. 71 = V. geniculata.
rubricaulis, Carr. in Rev. Hortic. (1868) 39.—China.
rufescens, Guill. & Perr. Fl. Seneg. Tent. i. 133 = V. rufescens.
rugosa, DC. Prod. i. 629.—Ins. Jamaic.
salutaris, H. B. & K. Nov. Gen. et Sp. v. 225 = V. salutaris.
Sandersoni, Harv. in Harv. & Sond. Fl. Cap. i. 253.—Afr. austr.
scariosa, Blume, Bijdr. 191 = V. coriacea.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 114 = V. oxyphylla.
semicordata, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 481 = V. himalayana.
semiglabra, Harv. in Harv. & Sond. Fl. Cap. i. 250 = V. semiglabra.
sericea, Eckl. & Zeyh. Enum. 57.—Afr. austr.
serjaniaefolia, Walp. Rep. i. 441 = V. serjaniaefolia.
serpens, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 111 = V. serpens.
serratifolia, Rottl. ex Wight, in Hook. Bot. Misc. ii. (1831) 104 = V. pedata.
serrulata, Roxb. Hort. Beng. 11 ; Fl. Ind. i. 414 = V. capreolata.
setosa, Roxb. ll. cc. 83 ; 410 = V. setosa.
sexangularis, Ernst, in Journ. Bot. viii. (1870) 374.—Venezuela.
sicyoides, Klein, ex Steud. Nom. ed. II. i. 372, 373 = V. discolor.
sicyoides, Linn. Syst. ed. X. 897 = V. sicyoides.
simplex, Blanco, Fl. Filip. ed. I. 72 = V. adnata.
Simsiana, Roem. & Schult. Syst. iii. 246 = V. Simsiana.
smilacina, H. B. & K. Nov. Gen. et Sp. v. 224 = V. sicyoides.
spicifera, Griff. Notul. iv. 693 = V. macrostachys.
spinosa, Cambess. in St. Hil. Fl. Bras. Mer. i. 345 = V. spinosa.
stans, Pers. Syn. i. 143 = V. arborea.
stipulata, Vell. Fl. Flum. 39 ; i. t. 99 = V. sulcicaulis ?
striata, Ruiz & Pav. Fl. Per. i. 64. t. 100 = V. striata.
stricta, Bert. ex Steud. Nom. ed. II. i. 372, in syn. = V. striata.
subdiaphana, Steud. in A. Rich. Tent. Fl. Abyss. i. 110 = V. gracilis.
suberecta, Bedd. in Madr. Journ. Sc. Scr. II. xxii. (1861) 75.—Ind. or.
sylvatica, Cambess. in St. Hil. Fl. Bras. Mer. i. 345 = V. erosa.
tamoides, Cambess. l. c. 342 = V. sicyoides.
tenuicaulis, Hook. f. in Hook. Niger Fl. 266 = V. tenuicaulis.
tenuifolia, Heyne, in Wall. Cat. n. 6022 A = V. tenuifolia.
ternata, J. F. Gmel. Syst. 256.—Arabia.
tetragona, Harv. in Harv. & Sond. Fl. Cap. i. 249.—Afr. austr.
tetraptera, Hook. f. in Hook. Niger Fl. 263 = V. quadrangularis.

CISSUS :—
Teysmanni, Miq. Fl. Ind. Bat. Suppl. i. 516.—Sumatra.
Thunbergii, Eckl. & Zeyh. Enum. 56.—Afr. austr.
Thunbergii, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1846) 195 = Vitis inconstans.
thyrsiflora, Blume, Bijdr. 187 = V. thyrsiflora.
tiliacea, H. B. & K. Nov. Gen. et Sp. v. 222 = V. tiliacea.
timoriensis, DC. Prod. i. 630.—Ins. Timor.
tinctoria, Arruda, in Koster, Trav. Braz. 496.—Bras.
tinctoria, Mart. in Spix & Mart. Reise Bras. 368 = V. sicyoides.
tomentosa Lam. Tabl. Encyc. i. 330 = V. capensis.
triandra, Schum. & Thonn. Beskr. Guin. Pl. 81 = V. quadrangularis.
tricuspidata, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1846) 196.—Japon.
tricuspis, Burch. ex Baker, in Mart. Fl. Bras. xiv. II. 209 = V. tricuspis.
tridentata, Eckl. & Zeyh. Enum. 56 = *pauciflora*.
trifoliata, Buch.-Ham. in Wall. Cat. n. 6013 ? K = V. angustifolia.
trifoliata, Linn. Sp. Pl. ed. II. 170 = V. trifoliata, *Baker*.
trifoliata, Lour. Fl. Cochinch. 83 = *dubia*.
trigona, Willd. ex Roem. & Schult. Syst. iii. Mant. 248.—Bras.
trilobata, Lam. Encyc. i. 31 = V. Rheedei.
triternata, Miq. Fl. Ind. Bat. Suppl. i. 516.—Sumatra.
truncata, Laws. in Hook. f. Fl. Brit. Ind. i. 660 = V. tenuifolia.
tuberculata, Blume, Bijdr. 189 = V. tuberculata.
tuberculata, Jacq. Hort. Schoenb. i. 14.—Am. austr.
tuberosa, Moç. & Sesse, ex DC. Prod. i. 629 = V. tuberosa.
umbellata, Lour. Fl. Cochinch. 84.—China.
umbrosa, H. B. & K. Nov. Gen. et Sp. v. 223.—Am. austr.
unifoliata, Harv. in Harv. & Sond. Fl. Cap. i. 250.—Afr. austr.
uvifer, Afzel. Rem. Guin. 69 = V. uvifer.
vagans, Carr. in Rev. Hortic. (1875) 60 = V. japonica ?
Veitchii, Carr. l. c. (1877) 176.—Hab. ?
velutinus, Linden, ex Hook. Bot. Mag. t. 5207.—Hab. ?
venatorum, Descourt. Fl. Méd. Ant. v. 21. t. 309 = V. sicyoides.
verrucosa, Steud. Nom. ed. II. i. 373 = V. tuberculata.
vesicatoria, Blanco, Fl. Filip. ed. II. 50 = V. repens.
villosa, Roxb. Hort. Beng. 11, nomen.—Hab. ?
violacea, Meyen, Reise, i. 300.—Chili.
viticifolia, Sieb. & Zucc. in Abh. Akad. Muench. iv. II. (1846) 196 = V. serianaefolia.
vitiensis, A. Gray, Bot. U. St. Expl. Exped. i. 272 = V. vitiensis.
vitifolia, Boiss. Diagn. Ser. I. vi. 31 = V. persica.
vitifolia, Salisb. Prod. 66 = V. repanda.
vitiginea, Linn. Sp. Pl. 117 = V. repanda.
vitiginea, Roxb. Fl. Ind. i. 406 = V. lanata.
Vogelii, Hook. f. in Hook. Niger Fl. 267 = V. Vogelii.
Wallichiana, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) I. 416 = Hedera Helix.
Zollingeri, Turcz. l. c. xxxvi. (1863) I. 591.—Java.

CISTANCHE, Hoffmgg. & Link, Fl. Port. i. 318. t. 63 (1806). *OROBANCHACEAE*, Benth. & Hook. f. ii. 983.
HAEMODORON, Reichb. Consp. 212[b] (1828).
lutea, *Hoffmgg. & Link, l. c.* 319.—Reg. Mediterr. ; Arabia.
lutea, Wight, Illustr. ii. 180. t. 158. b. f. 4 = tubulosa.
tubulosa, *Wight, Ic.* t. 1420 bis.—Ind. bor. occ.

CISTANTHE, Spach, Hist. Vég. Phan. v. 229 (1836) = **Calandrinia**, H. B. & K. (Portulac.).
anceps, Spach, l. c. 230 = Cal. grandiflora.
discolor, Spach, l. c. 231 = Cal. discolor.
glauca, Spach, ex Heynh. Nom. ii. 146 = Cal. grandiflora.
speciosa, Lilja, ex Heynh. l. c. = Cal. discolor.

CISTELA, Blume, Fl. Jav. praef. p. vi (1828) = seq.

CISTELLA, Blume, Bijdr. 293. t. 55 (1825) = **Geodorum**, Jack Orchid.).
cernua, Blume, l. c. = G. dilatatum.

CISTICAPNOS, Adans. Fam. ii. 431 (1763) = CYSTICAPNOS, Boerh. = **Corydalis**, Vent. (Papaverac.).

CISTOCARPIUM, Spach, Hist. Vég. Phan. vi. 471 (1838) = **Vesicaria**, Tourn. (Crucifer.).
utriculatum, Spach, l. c. 472 = V. utriculata.

CISTOCARPUM, Pfeiff. Nom. i. 766 (1874) = seq.

CISTOCARPUS, Kunth, in Mém. Soc. Nat. Hist. Par. iii. (1827) 389 = **Balbisia**, Cav. (Geraniac.).

CISTOMORPHA, Caley, ex DC. Syst. i. 427, in syn. (1818) = **Hibbertia**, Andr. (Dilleniac.).
lanceolata, Caley, l. c. = H. saligna.

CISTRUM, Hill, Veg. Syst. iv. 36 (1762) = **Centaurea**, Linn. (Compos.).
asperum, Hill, l. c. = Cent. aspera.
Isnardi, Hill, l. c. = Cent. aspera.
napifolium, Hill, l. c. = Cent. napifolia.

CISTULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 (Quid ?).

CISTUS, [Tourn.] Linn. Syst. ed. I (1735). *CISTINEAE*, Benth. & Hook. f. i. 113.
HALIMIUM, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 365.
LADANIUM, Spach, l. c. 367 (1836).
LEDONIA, Spach, l. c. 369 (1836).
LIBANOTIS, Rafin. Sylva Tellur. 132 (1838).
RHODOCISTUS, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 367.
STEPHANOCARPUS, Spach, l. c. 368 (1836).
STROBON, Rafin. Sylva Tellur. 132 (1838).
acuminatus, Viv. Ann. Bot. i. II. (1804) 172 = Helianthemum variabile.
acutifolius, *Sweet, Cist.* t. 78.—Europ.
aegypticus, Linn. Sp. Pl. 527 = Helianthemum aegyptiacum.
aegyptius, Pall. ex Bieb. Fl. Taur. Cauc. ii. 8 = Helianthemum salicifolium.
affinis, Bertol. ex Guss. Fl. Sic. Prod. ii. 12 = monspeliensis.
albidus, *Linn. Sp. Pl.* 524.—Reg. Mediterr.
albissimus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
algarvensis, Sims, Bot. Mag. t. 627 = Helianthemum ocymoides.
alpestris, Crantz, Stirp. Austr. ed. I. fasc ii. 73 ; ed. II. 103 = Helianthemum cinereum.
alternifolius, Vahl, Symb. Bot. i. 38 = Helianthemum brasiliense.
alyssoides, Lam. Encyc. ii. 20 = Helianthemum alyssoides.
anglicus, Linn. Mant. ii. 245 = Helianthemum cinereum.
angustifolius, Jacq. Hort. Vindob. iii. 29 = Helianthemum variabile.
angustifolius, Lag. ex Willk. & Lange, Prod. Fl. Hisp. iii. 734 = Helianthemum leptophyllum.
angustifolius, Salisb. Prod. 368 = Helianthemum umbellatum.
annuus, Lamb. ex Dun. in DC. Prod. i. 272 = Helianthemum villosum.
anthurus, Link, in Schrad. Neues Journ. i. (1805) 112 = Helianthemum umbellatum.
appeninus, Linn. Sp. Pl. 529 = Helianthemum variabile.
arabicus, Linn. Cent. i. 14 = Helianthemum arabicum.
asperifolius, *Pomel, Nouv. Mat. Fl. Atl.* 355.—Afr. bor.
asperifolius, Sweet, Cist. t. 87 = longifolius.
atriplicifolius, Lam. Encyc. ii. 19 = Helianthemum atriplicifolium.
barbatus, Lam. l. c. 24 = Helianthemum vulgare.
Barrelieri, Steud. Nom. ed. I. 199 = Helianthemum glutinosum.
Bourgaeanus, *Coss. Notes Crit.* ii. 30.—Hispan.
brasiliensis, Lam. Encyc. ii. 22 = Helianthemum brasiliense.
brevifolius, *Mill. Gard. Dict.* ed. VIII. n. 3.—Hab. ?
bupleurifolius, Lam. Encyc. ii. 22 = Helianthemum guttatum.
calycinus, Linn. Mant. ii. 565 = Helianthemum Fumana.
canadensis, Linn. Sp. Pl. 526 = Helianthemum canadense.

CISTUS:—
canariensis, Jacq. Misc. ii. 339 = Helianthemum canariense.
candidissimus, Dun. ex DC. Prod. i. 264 = vaginatus.
canescens, Sweet, Cist. t. 45 = polymorphus.
canus, Georgi, Beschr. Russ. Reich. iii. IV. 1047 = Helianthemum oelandicum.
canus, Jacq. Fl. Austr. iii. 42. t. 277 = Helianthemum montanum.
canus, Linn. Sp. Pl. 525 = Helianthemum vineale.
capensis, *Linn. l. c.* ed. II. 736.—Afr. austr.
carolinianus, Walt. Fl. Carol. 152 = Helianthemum carolinianum.
cheiranthoides, Lam. Encyc. ii. 19 = Helianthemum lasianthum.
ciliatus, Desf. Fl. Atlant. i. 421. t. 109 = Helianthemum virgatum.
ciliatus, Rostung, ex Dun. in DC. Prod. i. 281 = Helianthemum obtusifolium.
cinereus, Cav. Ic. ii. 33. t. 141 = Helianthemum cinereum.
cistifolius, Steud. Nom. ed. I. 200 = Helianthemum variabile.
Clusii, *Dun. in DC. Prod.* i. 266.—Hispan.; Afr. bor.
collinus, Salisb. Prod. 368 = monspeliensis.
complicatus, Lam. Encyc. ii. 14 = parviflorus.
complicatus, Sprun. ex Nym. Consp. 70 = villosus.
corbariensis, Pourr. in Mém. Acad. Toul. iii. (1788) 312 = salvifolius.
cordifolius, *Mill. Gard. Dict.* ed. VIII. n. 12.—Hab. ?
cordifolius, Moench, Meth. 245 = populifolius.
coriaceus, Poir. Encyc. Suppl. ii. 278 (= *Hemistemma Aubertii*, *Commersonii*).—Madag.
corsicus, Loisel. Nonv. Not. 24 = villosns.
corymbosus, Poir. Encyc. Suppl. ii. 272 = Helianthemum corymbosum.
crassifolius, Poir. l. c. 277 = Helianthemum cinereum.
creticus, Linn. Sp. Pl. ed. II. 738 = polymorphus.
creticus, Sibth. & Sm. Fl. Graec. Prod. i. 364 = parviflorus.
crispus, *Linn. Sp. Pl.* 524.—Reg. Mediterr.
croceus, Desf. Fl. Atlant. i. 422. t. 110 = Helianthemum glaucum.
Cupanianus, Presl, Fl. Sic. i. 117 = polymorphus.
cymosus, Dun. in DC. Prod. i. 265 = parviflorus.
cyprius, *Lam. Encyc.* ii. 16.—Ins. Cyprus.
denticulatus, *Kit. ex Kanitz, in Verh. Zool.-Bot. Ges. Wien*, xiii. (1863) 529.—Hungar.
denticulatus, Poir. Encyc. Suppl. ii. 278 = Helianthemum salicifolium.
dichotomus, Cav. Ic. iii. 32 = Helianthemum origanifolium.
distachyus, Roth, Catalecta, i. 62 = Helianthemum distachyum.
dubius, *Pourr. in Mém. Acad. Toul.* iii. (1788) 312.—Mont. Pyren.
Dunalianus, Steud. Nom. ed. II. i. 374 = polymorphus.
echioides, Lam. Encyc. i. 21 = Helianthemum variabile.
ellipticus, Desf. Fl. Atlant. i. 418. t. 107 = Helianthemum Lippii.
elongatus, Vahl, Symb. Bot. i. 38 = Helianthemum lasianthum.
ericoides, Cav. Ic. ii. 56. t. 172 = Helianthemum Fumana.
eriocephalus, Viv. Fl. Cors. Sp. Nov. 8 = polymorphus.
fasciculatus, Steud. Nom. ed. I. 200 = Helianthemum fasciculatum.
fastigiatus, Guss. Fl. Sic. Prod. ii. 13 = Clusii.
feredjensis ×, Battand. in Bull. Soc. Bot. Fr. xxx. (1883) 263.—Afr. bor.
ferrugineus, Lam. Encyc. ii. 25 = Helianthemum arabicum.
florentinus, Lam. l. c. 17 = monspeliensis.
floribundus, *Tausch, in Flora*, xix. (1836) 417.—Gallia.
foetidus, Jacq. Misc. ii. 341 = Helianthemum variabile.
formosissimus, *Schneev. Ic. Pl. Rar.* t. 47.—Lusitan.
formosus, Curt. Bot. Mag. t. 264 = Helianthemum lasianthum.
Fumana, Linn. Sp. Pl. 525 = Helianthemum procumbens.
Fumana, Schousb. ex Ball, in Journ. Linn. Soc. xvi. (1877) 348 = Helianthemum glutinosum.

CISTUS:—
galactites, Schmidel, Ic. Pl. ed. II. 250. t. 65 = Helianthemum vulgare ?
garganicus, Tenore, Ind. Sem. Hort. Neap. (1829) 15, ex Syll. ed. fol. 74 = villosus.
glaucophyllus, Lam. Fl. Fr. iii. 162 = Helianthemum laevipes.
glaucus, Cav. Ic. iii. 31. t. 261 = Helianthemum glaucum.
glaucus, Desf. Fl. Atlant. i. 418 = Helianthemum crassifolium.
glaucus, *Pourr. in Mém. Acad. Toul.* iii. (1788) 311.—Gallia.
glaucus, Salisb. Prod. 368 = Helianthemum laevipes.
globularifolius, Lam. Encyc. ii. 22 = Helianthemum Tuberaria.
glomeratus, Lag. Gen. et Sp. Nov. 16 = Helianthemum glomeratum.
glutinosus, Linn. Mant. ii. 246 = Helianthemum glutinosum.
grandiflorus, Pourr. ex Willk. & Lange, Prod. Fl. Hisp. iii. 712 = ladaniferus.
grandiflorus, Scop. Fl. Carn. ed. II. i. 377 = Helianthemum vulgare.
guttatus, Linn. Sp. Pl. 526 = Helianthemum guttatum.
halimifolius, Linn. l. c. 524 = Helianthemum halimifolium.
helianthemoides, Crantz, Stirp. Anstr. ed. I. i. 69; ed. II. 101 = Helianthemum variabile.
helianthemoides, *Desf. Fl. Atlant.* i. 422.—Afr. bor.
Helianthemum, Linn. Sp. Pl. 528 = Helianthemum vulgare.
heterophyllus, Desf. Fl. Atlant. i. 411. t. 104 = polymorphus.
hirsutus, *Lam. Fl. Fr.* iii. 157.—Hispan.; Lusitan.
hirsutus, Lapeyr. Hist. Abr. Pl. Pyr. 303 = Helianthemum polymorphum.
hirtus, Chaub. ex Nym. Consp. 74 = Helianthemum glaucum.
hirtus, Gilib. Fl. Lituan. v. 225 = Helianthemum vulgare.
hirtus, Linn. Sp. Pl. 528 = Helianthemum hirtum.
hirtus, Sibth. & Sm. Fl. Graec. vi. 1. t. 501 = Helianthemum obtusifolium.
hirtus, Thib. ex Dun. in DC. Prod. i. 283 = Helianthemum asperum.
hispanicus, *Mill. Gard. Dict.* ed. VIII. n. 5.—Hispan.
hispidus, Lam. Encyc. ii. 26 = Helianthemum hirtum.
horridus, Lag. in Varied. Cienc. ii. IV. (1805) 214, sphalm. (= *Cirsium horridum*).
humilis, Salisb. Prod. 368 = Helianthemum vulgare.
hybridus, Pourr. in Mém. Acad. Toul. iii. (1788) 312 = salvifolius.
hybridus, *Vahl, Symb. Bot.* i. 37.—Hispan.
imbricatus, Poir. Encyc. Suppl. ii. 278 = Helianthemum ledifolium.
immaculatum, Steud. Nom. ed. I. 200 = Helianthemum guttatum.
incanus, Linn. Sp. Pl. 524 = polymorphus.
incanus, Sibth. & Sm. Fl. Graec. v. 76. t. 494 = parviflorus.
inconspicuus, Poir. Encyc. Suppl. ii. 278 = Helianthemum guttatum.
indicus, Linn. Syst. ed. XII. 367, sphalm. vide *italicus*.
intermedius, Poir. Encyc. Suppl. ii. 278 = Helianthemum intermedium.
involucratus, Lam. Encyc. ii. 20 = Helianthemum lasianthum.
italicus, Linn. Syst. ed. X. 1078 = Helianthemum montanum.
italicus, Pall. ex Bieb. Fl. Taur. Cauc. ii. 8 = Helianthemum oelandicum.
japonicus, *Fortune, in Journ. Agri.-Hort. Soc. Ind.* ix. (1855) 96.—Japon.
kahiricus, Steud. Nom. ed. I. 200 = Helianthemum kahiricum.
ladaniferus, Curt. Bot. Mag. t. 112 = cyprius.
ladaniferus, Gouan, ex Steud. Nom. ed. II. i. 374 = C. Ledon.
ladaniferus, *Linn. Sp. Pl.* 523.—Reg. Mediterr. occ.
ladaniferus, Stokes, Bot. Mat. Med. iii. 209 = polymorphus.
ladanosma, Hoffmgg. Verz. 51 = ladaniferus.
laevipes, Linn. Cent. i. 14 = Helianthemum laevipes.
laevis, Cav. Ic. ii. 35. t. 145. f. 1 = Helianthemum thymifolium.

CISTUS:—
lanceolatus, Vahl, Symb. Bot. ii. 62 = Helianthemum guttatum.
lanuginosus, Viv. Fl. Lib. Spec. 28. t. 14. f. 3 = Helianthemum lanuginosum.
lasianthus, Lam. Encyc. ii. 19 = Helianthemum lasianthum.
latifolius, Sweet, Cist. t. 15 = populifolius.
laurifolius, *Linn. Sp. Pl.* 523.—Europ. austr. occ.; As. Min.
lavandulaefolius, Lam. Encyc. ii. 25 = Helianthemum lavandulaefolium.
laxus, [Dryand. in] Ait. Hort. Kew. ed. II. iii. 305 = longifolius, *Lam.*
laxus, Brot. ex Nym. Consp. 71 = hirsutus.
ledifolius, Georgi, Beschr. Russ. Reich. Nachtr. 276 = Helianthemum salicifolium.
ledifolius, Linn. Sp. Pl. 527 = Helianthemum ledifolium.
ledonius, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = glaucus.
Ledon, Lam. Encyc. ii. 17 = glancus.
lepidotus, Amo, Fl. Iber. vi. 357 = Helianthemum halimifolium.
Libanotis, Desf. Fl. Atlant. i. 412 = Clusii.
Libanotis, Linn. Syst. ed. X. 1077 = Helianthemum Libanotis.
lilacinus, Hoffmgg. ex Steud. Nom. ed. II. i. 374 = parviflorus.
linearis, Cav. Ic. iii. 8. t. 216 = Helianthemum variabile.
Lippi, Linn. Mant. ii. 245 = Helianthemum Lippii.
longifolius, Hort. ex Steud. Nom. ed. II. i. 374 = grandiflorus.
longifolius, *Lam. Encyc.* ii. 16.—Hispan.; Gallia.
longifolius, *Mill. Gard. Dict.* ed. VIII. n. 18.—Hab. ?
lunulatus, All. Auct. Pedem. 30. t. 2. f. 3 = Helianthemum lunulatum.
lusitanicus, Mill. Gard. Dict. ed. VIII. n. 4 = Helianthemum variabile.
luteus, Gilib. Fl. Lituan. ii. 225 = Helianthemum variabile.
majoranifolius, Gouan, Herbor. 36 = Helianthemum hirtum.
marifolius, Bieb. Fl. Taur. Cauc. ii. 8 = Helianthemum oelandicum.
marifolius, Linn. Sp. Pl. 526 = Helianthemum vineale.
mauritanicus, Thib. ex Dun. in DC. Prod. i. 276 = Helianthemum glutinosum.
medius, *J. F. Gmel. Syst.* 822.—Italia.
micranthus, Viv. Fl. Lib. Spec. 27 = Helianthemum micranthum.
mollis, Cav. Ic. iii. 32. t. 262. f. 2 = Helianthemum origanifolium.
monspeliensis, *Linn. Sp. Pl.* 524.—Reg. Mediterr. occ.
multiflorus, Amo, Fl. Iber. vi. 359 = Helianthemum halimifolium.
Munbyi, *Pomel, Nouv. Mat. Fl. Atl.* 215.—Afr. bor.
mutabilis, Jacq. Misc. ii. 340 = Helianthemum variabile.
myrthifolius, Lam. Fl. Fr. iii. 161 = Helianthemum cinereum.
nervosus, Lam. l. c. 164 = Helianthemum Tuberaria.
nigricans, Pourr. in Mém. Acad. Toul. iii. (1788) 311 = longifolius.
niloticus, Linn. Mant. ii. 246 = Helianthemum ledifolium.
nudifolius, Lam. Fl. Fr. iii. 163 = Helianthemum Fumana.
nummularius, Cav. Ic. ii. 34. t. 142 = Helianthemum rubellum.
nummularius, Linn. Sp. Pl. 527 = Helianthemum vulgare.
oblongifolius, Sweet, Cist. t. 67 = lougifolius.
obscurus, Steud. Nom. ed. I. 201 = Helianthemum variabile.
obtusifolius, Sweet, Cist. t. 42 = longifolius.
occidentalis, Amo, Fl. Iber. vi. 352 = Helianthemum alyssoides.
ochreatus, *Chr. Sm. ex Buch, in Berl. Acad. Abh.* 1816–17 (1819) 380.—Ins. Canar.
ocymoides, Brot.—Cf. Nym. Consp. 72 = Helianthemum ocymoides.
ocymoides, Lam. Encyc. ii. 18 = polymorphns.

CISTUS:—
oelandicus, Lam. Encyc. ii. 20 = Helianthemum montanum.
oelandicus, Linn. Sp. Pl. 526 = Helianthemum italicum.
olbiensis, Huet & Hanry, in Bull. Soc. Bot. Fr. vii. (1860) 346 = monspeliensis.
oleaefolius, Mill. Gard. Dict. ed. VIII. n. 10 = monspeliensis.
oligophyllus, Clark, Trav. ii. 452 = Helianthemum oligophyllum.
origanifolius, Lam. Encyc. ii. 21 = Helianthemum origanifolium.
ovatus, Viv. Fl. Ital. Fragm. i. 6 = Helianthemum variabile.
parasiticus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
parviflorus, *Lam. Encyc.* ii. 14.—Reg. Mediterr. or.
Pechii, *Pourr. ex Timb. in Bull. Soc. Sc. Phys. et Nat. Toul.* ii. (1874) 90.—Gallia.
petiolatus, *Martr. Herbor. Midi Fr.—Cf. Bull. Soc. Bot. Fr.* ii. (1855) 122.—Gallia.
petiolatus, Poir. Encyc. Suppl. ii. 278 = Helianthemum petiolatum.
piliferus, Gmel. ex Willk. & Lange, Prod. Fl. Hisp. iii. 729 = Helianthemum pulverulentum.
piloselloides, Lapeyr. Hist. Abr. Pl. Pyr. 301 = Helianthemum montanum.
pilosus, All. Fl. Pedem. ii. 104 = Helianthemum polifolium.
pilosus, Linn. Sp. Pl. 528 = Helianthemum variabile.
piluliferus, Thib. ex Dun. in DC. Prod. i. 282 = Helianthemum variabile.
piperifolius, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
plantagineus, Willd. Sp. Pl. ii. 1197 = Helianthemum guttatum.
plantaginifolius, Salisb. Prod. 368 = Helianthemum Tuberaria.
platysepalus, *Sweet, Cist.* t. 47.—Ins. Creta.
polifolius, Linn. Sp. Pl. ed. II. 745 = Helianthemum pulverulentum.
polufolius, Huds. Fl. Angl. ed. I. 205 = praec.
polyanthos, Desf. Fl. Atlant. i. 420 = Helianthemum polyanthum.
polymorphus, *Willk. Ic. Pl. Hisp.* ii. 19 (sp. aggregat.).—Reg. Mediterr.
populifolius, *Linn. Sp. Pl.* i. 523.—Reg. Mediterr. occ.
porquerollensis, Hanry & Huet, in Bull. Soc. Bot. Fr. vii. (1860) 345 = monspeliensis.
Pouzolzii, Delile, Suppl. Cat. Hort. Monsp. (1839); ex Gren. & Godr. Fl. Fr. i. 163 = albidus.
psilosepalus, Sweet, Cist. t. 33 = hirsutus.
pulverulentus, Pourr. in Mém. Acad. Toul. iii. (1788) 312 = albidus.
punctatus, Willd. Sp. Pl. ii. 1199 = Helianthemum salicifolium.
purpureus, *Lam. Encyc.* ii. 14.—Oriens.
quinquenervis, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
quinquevulnerus, *Vis. in Act. Ist. Ven. Sc.* Ser. III. iv. (1858–59) 138.—Hab.?
racemosus, Cav. Ic. ii. 31. t. 140 = Helianthemum lavandulaefolium.
racemosus, Lam. Encyc. ii. 25 = Helianthemum virgatum.
racemosus, Linn. Mant. i. 76 = Helianthemum variabile.
ramuliflorus, Poir. Encyc. Suppl. ii. 274 = Helianthemum canadense.
retrofractus, Poir. l. c. 278 = Helianthemum retrofractum.
rorismarinifolius, Pourr. ex Timb. in Bull. Soc. Sc. Phys. & Nat. Toul. ii. (1874) 101 = Clusii.
roseus, All. Fl. Pedem. ii. 105 = Helianthemum vulgare.
roseus, Jacq. Hort. Vindob. iii. 37 = Helianthemum variabile.
rosmarinifolius, All. Fl. Pedem. ii. 105 = Helianthemum rosmarinifolium.
rosmarinifolius, Pourr. in Mém. Acad. Toul. iii. (1788) 313 = Clusii.
rosmarinifolius, Salisb. Prod. 368 = Helianthemum Libanotis.
rotundifolius, Sweet, Cist. t. 75 = vulgaris.
ruficomus, Viv. Fl. Cors. Sp. Nov. 8 = Helianthemum ruficomum.
salicifolius, Cav. Ic. ii. 35. t. 144 = Helianthemum intermedium.

CISTUS:—
salicifolius, Huds. Fl. Angl. ed. II. 233 = Helianthemum ledifolium.
salicifolius, Linn. Sp. Pl. 527 = Helianthemum salicifolium.
salviaefolius, Boiss. Fl. Orient. iii. 438 = seq.
salvifolius, *Linn. Sp. Pl.* 524.—Reg. Mediterr.
salvifolius, Pall. ex Bieb. Fl. Taur. Cauc. ii. 7 = polymorphus.
sampsueifolius, Cav. Ic. i. 65 = Helianthemum ocymoides.
sampsucifolius, Sims, Bot. Mag. xliii. (1816) t. 1803 = Helianthemum variabile.
sanguineus, *Lag. in Varied. Ciene.* ii. IV. (1805) 40.—Austr.
Savi, Bertol. ex Dun. in DC. Prod. i. 275 = Helianthemum arabicum.
scabrosus, [Soland. in] Ait. Hort. Kew. ed. I. ii. 236 = Helianthemum lasianthum.
scorpioides, Desf. ex Steud. Nom. ed. II. i. 375 = Helianthemum glaucum.
sedjera, *Pomel, Nouv. Mat. Fl. Atl.* 216.—Afr. bor.
Seguieri, Crantz, Stirp. Austr. ed. I. fasc. ii. 74, ed. II. 104 = Helianthemum cinereum.
Seguieri, Pourr. ex Dun. in DC. Prod. i. 277 = Helianthemum italicum.
sericeus, *Munby, Fl. Algér.* 53.—Algeria.
sericeus, *Vahl, Symb. Bot.* i. 37.—Hispan.
serpillifolius, Balb. ex Dun. in DC. Prod. i. 280 = Helianthemum variabile.
serpillifolius, Linn. Sp. Pl. 527 = Helianthemum vulgare.
serratifolius, Lamb. ex Dun. in DC. Prod. i. 271 = Helianthemum guttatum.
serratus, Cav. Ic. ii. 57. t. 175 = Helianthemum guttatum.
sessiliflorus, Desf. Fl. Atlant. i. 417 = Helianthemum Lippii.
Sideritis, *Presl, Fl. Sic.* i. 116.—Sicilia.
Skanbergi, *Lojac. in Natural. Sicil.* iv. (1885) 95.—Sicilia.
splendens, Lam. Fl. Fr. iii. 156 = Helianthemum variabile.
squamatus, Linn. Sp. Pl. 1196 = Helianthemum squamatum.
stellulatus, Steud. Nom. ed. I. 201 = Helianthemum stellulatum.
stenophyllus, Link, Enum. Hort. Berol. ii. 74 = Clusii, cyprius.
stipulatus, Forsk. Fl. Aegypt. Arab. 100 = Helianthemum kahiricum.
stoechadifolius, Brot. Fl. Lusit. ii. 270 = Helianthemum glaucum.
stoechadifolius, Hort. ex Dun. in DC. Prod. i. 280 = Helianthemum leptophyllum.
strictus, Cav. Ic. iii. 32. t. 263. f. 2 = Helianthemum virgatum.
strigosus, Fisch. ex Steud. Nom. ed. II. i. 375 = Helianthemum cinereum.
sulphureus, Steud. l. c. ed. I. 201 = Helianthemum variabile.
surrejanus, Linn. Sp. Pl. 527 = Helianthemum Chamaecistus.
symphytifolius, Lam. Encyc. ii. 15 = vaginatus.
syriaeus, Jacq. Ic. Pl. Rar. i. 10. t. 96 = Helianthemum lavandulaefolium.
syrticus, Viv. Fl. Lib. Spec. 27 = Helianthemum syrticum.
tauricus, *F. & C. Presl, Delic. Prag.* 24.—Tauria.
telephioides, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
teretifolius, Poir. Encyc. Suppl. ii. 279 = Helianthemum hirtum.
teretifolius, Thib. ex Dun. in DC. Prod. i. 281 = Helianthemum variabile.
Thibaudi, Poir. Encyc. Suppl. ii. 277 = Helianthemum lavandulaefolium.
thimifolius, Gouan, ex Vill. Prosp. 49 = Helianthemum thymifolium.
thymifolius, Falk, Beitr. ii. 197 = Helianthemum soongoricum.
thymifolius, Linn. Sp. Pl. 528 = Helianthemum glutinosum.
thymifolius, Sibth. & Sm. Fl. Graec. v. 81. t. 500 = Helianthemum thymifolium.
tomentosus, Lam. Fl. Fr. iii. 168 = polymorphus.
tomentosus, Scop. Ann. II. Hist. Nat. (1769) 53; Fl. Carn. ed. II. i. 376 = Helianthemum Chamaecistus.

CISTUS:—
tricuspis, Darwin, Variation Pl. & Anim. Domest. i. 377, err. typ. = Hibiscus tricuspis.
Tuberaria, Linn. Sp. Pl. 526 = Helianthemum Tuberaria.
umbellatus, Linn. l. c. 525 = Helianthemum umbellatum.
undulatus, Moench, Meth. 245 = polymorphus.
vaginatus, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. iii. 304.—Teneriffa.
valentinus, Pourr. ex Nym. Consp. 71 = monspeliensis.
varius, Pourr. in Mém. Acad. Toul. iii. (1788) 312 = albidus.
verticiliflorus, Steud. Nom. ed. II. i. 375 = Helianthemum anthurum.
verticillatus, Brot. Fl. Lusit. ii. 262 = Helianthemum umbellatum.
villosus, *Linn. Gen.* ed. VI. pag. ult. (nomen incertum).—Hispan.
vinealis, Willd. Sp. Pl. ii. 1195 = Helianthemum vineale.
violaceus, Cav. Ic. ii. 38. t. 147 = Helianthemum variabile.
virgatus, Desf. Fl. Atlant. i. 422 = Helianthemum virgatum.
virgatus, Thib. ex Dun. in DC. Prod. i. 269 = Helianthemum canadense.
viscosissimus, Tausch, in Flora, xix. (1836) 418.—Hab.?
viseosus, Stokes, Bot. Mat. Med. iii. 209 = ladaniferus.
vulgaris, Spach, in Ann. Sc. Nat. Sér. II. vi. (1836) 368 = polymorphus.

CITHARELLA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 = (Quid?).

CITHARELOMA, Bunge, in Cat. Hort. Dorp. (1843) 6; et in Linnaea, xviii. (1844) 149. *CRUCIFERAE*, Benth. & Hook. f. i. 67.
canescens, Bunge, in Linnaea, xviii. (1844) 150, in nota = vernum.
Lehmanni, *Bunge, Cat. Hort. Dorp.* (1843) 6; *et in Linnaea*, xviii. (1844) 149.—As. med.
vernum, *Bunge, in Linnaea*, xviii. (1844) 150, in nota.—As. med.

CITHAREXYLUM, Mill. Gard. Dict. ed. VI. 175 (1752); Linn. Sp. Pl. ed. I. 625 (1753). *VERBENACEAE*, Benth. & Hook. f. ii. 1149.
POEPPIGIA, Bert. in Bull. Férussac, xxiii. (Oct. 1830) 109.
RAUWOLFIA, Ruiz & Pav. Fl. Per. ii. 26. t. 152 (1799).
affine, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March 1831) 238.—Mexic.
affine, Mart. & Gal. in Bull. Acad. Brux. xi. II. (1844) 328 = reticulatum.
albicaule, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 208.—Cuba.
album, Mill. Gard. Dict. ed. VIII. n. 2 = caudatum.
alpinum, Poepp. ex Schau. in DC. Prod. xi. 545 = Verbena cinerascens.
barbinerve, *Cham. in Linnaea*, vii. (1832) 116.—Bras.
Berterii, *Spreng. Syst.* ii. 763.—Ins. Jamaica.
caudatum, Hort. ex Walp. Rep. vi. 646 = Buddleia melliodora.
caudatum, *Linn. Sp. Pl.* ed. II. 872.—Ins. Jamaic.
caudatum, Sw. Obs. Bot. 234 = quadrangulare.
Chamissonis, Walp. Rep. iv. 75 = caudatum.
cinereum, *Linn. Sp. Pl.* ed. II. 872.—Am. trop.
cinereum, Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ. (Jan. March, 1831) 238 = lucidum.
cinereum, Spreng. Syst. ii. 764, partim = myrianthum.
coriaceum, Desf. Cat. Hort. Par. ed. III. 302 = quadrangulare.
cyanocarpum, C. Gay, Fl. Chil. v. 34 = Rhaphithamnus parvifolius.
cyanocarpum, Hook. & Arn. Bot. Beech. Voy. 58. t. 11 = Rhaphithamnus cyanocarpus.
dentatum, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 237.—Peruv.
discolor, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 209.—Cuba.
elegans, Phil. ex Miers, in Trans. Linn. Soc. xxvii. (1870) 98 = Rhaphithamnus longiflorus.
ellipticum, *Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ.* (Jan. March, 1831) 238.—Mexic.
erectum, Sw. Prod. Veg. Ind. Occ. 91 = caudatum.
flexuosum, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March 1831) 237.—Peruv.

CITHAREXYLUM:—
fruticosum, Linn. Syst. ed. X. 1115 = cinereum.
ilicifolium, *H. B. & K. Nov. Gen. et Sp.* ii. 256.—Ecuador.
incanum, *Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 238.—Mexic.
laetum, *Hiern, in Kjoeb. Vidensk. Meddel.* (1877–78) 104.—Am. centr.
laevigatum, Hostm. ex Griseb. Fl. Brit. W. Ind. 497 = lucidum.
Lindenii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 208.—Cuba.
longiflorum, *Turcz. l. c.* 207.—Cuba.
lucidum, Cham. & Schlecht. in Linnaea, v. (1830) 97 = caudatum.
lucidum, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 238.—Mexic.
lycioides, *D. Don, l. c.* 237.—Mexic.
macrophyllum, *Poir. Encyc. Suppl.* ii. 367.—Guiana.
melanocardium, Sw. Prod. Veg. Ind. Occ. 91 = Petitia domingensis.
mendocinum, *Phil. Sert. Mend. Alt.* 35.—Chili.
Mocinni, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 238.—Mexic.
molle, H. B. & K. Nov. Gen. et Sp. ii. 257 = quitense.
molle, *Jacq. ex Spreng. Syst.* ii. 764.—Am. austr.
molle, Salisb. Prod. 108 = villosum.
myrianthum, *Cham. in Linnaea*, vii. (1832) 117.—Bras.
ovatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 207.—Chili.
paniculatum, *Gaertn. Fruct.* i. 270. t. 56.—Jamaica.
paniculatum, Poir. in Lam. Tabl. Encyc. iii. 95 = Premna integrifolia.
pentandrum, *Vent. Jard. Cels*, t. 47.—Porto Rico.
perforatum, Forst. f. Prod. 44 = Myoporum laetum.
Poeppigii, *Walp. Rep.* iv. 76.—Bras.
polystachyum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 209.—Cuba.
psilacanthum, *Turcz. l. c.* 207.—Hab. ?
pulverulentum, *Pers. Syn.* ii. 142.—Am. austr.
quadrangulare, Hort. Madr. ex Steud. Nom. ed. II. i. 375 = pulverulentum.
quadrangulare, *Jacq. Enum. Pl. Carib.* 26.—Ind. occ.
quadrangulare, Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ. (Jan.–March, 1831) 238 = Sessaei.
quitense, *Spreng. Syst.* ii. 763.—Ecuador.
reticulatum, *H. B. & K. Nov. Gen. et Sp.* ii. 257.—Peruv.
retusum, *D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 237.—Peruv.
Rugendasii, *Cham. in Linnaea*, vii. (1832) 120.—Bras.
scabrum, *Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ.* (Jan.–March, 1831) 238.—Mexic.
scabrum, Willd. ex Cham. in Linnaea, vii. (1832) 117 = myrianthum.
scariosum, Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ. (Jan.–March, 1831) 238 = reticulatum.
Sessaei, *D. Don, l. c.*—Mexic.
solanaceum, *Cham. in Linnaea*, vii. (1832) 119.—Bras.
spinosum, *H. B. & K. Nov. Gen. et Sp.* ii. 256.—Peruv.
spinosum, *Linn. Sp. Pl.* 625.—Ins. Barbad.
subserratum, *Sw. Prod. Veg. Ind. Occ.* 91.—Ins. S. Doming.
surrectum, *Griseb. Fl. Brit. W. Ind.* 497.—Ind. occ.
tenuifolium, *Forst. f. Prod.* 44 = Myoporum tenuifolium.
teres, Jacq. Enum. Pl. Carib. 46 = cinereum.
tomentosum, *H. B. & K. Nov. Gen. et Sp.* ii. 258.—Am. austr.
tomentosum, Moç. & Sesse, ex D. Don, in Edinb. N. Phil. Journ. (Jan.–March, 1831) 238 = Mocinni.
tomentosum, Poir. Encyc. Suppl. ii. 368 = pentandrum.
tristachyum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 209.—Cuba.
venustum, *Phil. in Bot. Zeit.* xiv. (1856) 646 = Rhaphithamnus longiflorus.
verticillatum, Klotzsch, ex Walp. Rep. iv. 73 = cyanocarpum.
villosum, *Jacq. Ic. Pl. Rar.* i. 12. t. 118; *Coll.* i. 72.—Ins. S. Doming.

CITINUS, All. Fl. Pedem. ii. 214 (1785) = **Cytinus**, Tourn.

CITREUM, Tourn. ex Rupp. Fl. Jen. ed. Hall. 141 (1745); Mill. Gard. Dict. ed. VI. (1752) = **Citrus**, Linn.

CITRIOBATHUS, A. Juss. in Orb. Dict. x. 228 (1849) = seq.

CITRIOBATUS, A. Cunn. in Loud. Hort. Brit. Suppl. i. 585 (1832). *PITTOSPOREAE*, Benth. & Hook. f. i. 132.
IXIOSPORUM, F. Muell. Fragm. ii. 76 (1860).
megacarpus, *F. Muell. Fragm.* ii. 76.—Austral.
multiflorus, *A. Cunn. in Loud. Hort. Brit. Suppl.* i. 585.—Austral.
pauciflora, *A. Cunn. l. c.*—Austral.

CITRONELLA, D. Don, in Edinb. N. Phil. Journ. [xiii.] (Oct. 1832) 243 = **Villaresia**, Ruiz & Pav. (Olacin.).
mucronata, D. Don, l. c. = V. mucronata.

CITROSENA, Bosc, ex Steud. Nom. ed. II. i. 375 (1840) = seq.

CITROSMA, Ruiz & Pav. Fl. Per. et Chil. Prod. 134. t. 29 (1794) = **Siparuna**, Aubl. (Monimiac.).
alternifolia, Spreng. Neue Entdeck. ii. 112 = S. alternifolia.
amazonum, Tul. in Arch. Mus. Par. viii. (1855–56) 332 = S. asperula.
andina, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 36 = S. andina.
apicifera, Tul. l. c. 38 = S. apicifera.
Apiosyce, Mart. ex Tul. l. c. 34 = S. Apiosyce.
aspera, Tul. in Arch. Mus. Par. viii. (1855–56) 324 = S. aspera.
asperula, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 35 = S. asperula.
bifida, Poepp. & Endl. Nov. Gen. et Sp. ii. 48 = S. bifida.
brasiliensis, Spreng. Neue Entdeck. ii. 113 = S. brasiliensis.
buddleiaefolia, Benth. Pl. Hartw. 250 = S. budleiaefolia.
camporum, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 39 = S. camporum.
chiridota, Tul. l. c. 37 = S. chiridota.
cristata, Poepp. & Endl. Nov. Gen. et Sp. ii. 47. t. 164 = S. cristata.
cujabana, Mart. ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 33 = S. cujabana.
cuspidata, Tul. l. c. 40 = S. cuspidata.
decipiens, Tul. in Arch. Mus. Par. viii. (1855–56) 368 = S. decipiens.
dentata, Poepp. ex A. DC. Prod. xvi. II. 653 = S. mollicoma.
dentata, Ruiz & Pav. Syst. Veg. i. 264 = S. limoniodora.
dimidiata, Sello, ex A. DC. Prod. xvi. II. 656 = S. paniculata.
discolor, Poepp. & Endl. Nov. Gen. et Sp. ii. 47 = S. discolor.
dubia, H. B. & K. Nov. Gen. et Sp. ii. 174 = S. mollis.
echinata, H. B. & K. l. c. 173 = S. echinata.
eriocalyx, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 35 = S. eriocalyx.
erythrocarpa, Mart. ex A. DC. Prod. xvi. II. 645 = S. estrellensis.
erythrocarpa, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 32 = S. erythrocarpa.
estrellensis, Tul. l. c. 33 = S. estrellensis.
foliosa, Tul. l. c. 38 = S. foliosa.
gesnerioides, H. B. & K. Nov. Gen. et Sp. ii. 172 = S. gesnerioides.
glabra, Spreng. Syst. ii. 545 = S. glabra.
glabrescens, Presl, Bot. Bemerk. 110 = S. glabrescens.
grandiflora, H. B. & K. Nov. Gen. et Sp. ii. 172 = S. Kunthii.
Gudotiana, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 38 = S. Goudotiana.
guianensis, Tul. in Arch. Mus. Par. viii. (1855–56) 361 = S. guianensis.
Kunthii, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 37 = S. Kunthii.
Lagopus, Tul. l. c. = S. Lagopus.
lanceolata, Tul. l. c. 33 = S. lanceolata.

CITROSMA:—
laurifolia, H. B. & K. Nov. Gen. et Sp. ii. 171 = Siparuna laurifolia.
lepidota, H. B. & K. l. c. 170 = S. lepidota.
limoniodora, Pav. ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 35 = S. limoniodora.
Lindeni, Seem. Journ. Bot. ii. (1864) 343 = S. Lindeni.
macrophylla, H. B. & K. Nov. Gen. et Sp. ii. 172 = S. macrophylla.
macrophylla, Mart. ex A. DC. Prod. xvi. II. 655 = S. cristata.
mollicoma, Mart. ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 38 = S. mollicoma.
mollis, H. B. & K. Nov. Gen. et Sp. ii. 173 = S. mollis.
muricata, Ruiz & Pav. Syst. Veg. 265 = S. muricata.
Mutisii, H. B. & K. Nov. Gen. et Sp. ii. 170 = S. Mutisii.
myristicoidea, Spruce, in Journ. Linn. Soc. v. (1861) 5.—Bras.
neglecta, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 36 = S. neglecta.
oblongifolia, Ruiz & Pav. Syst. Veg. i. 266 = S. laurifolia, oblongifolia.
obovata, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 343 = S. obovata.
oligandra, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 32 = S. obovata.
oligocarpa, Mart. ex A. DC. in DC. Prod. xvi. II. 654 = S. guianensis.
ovalis, Ruiz & Pav. Syst. Veg. i. 266 = S. ovalis.
paniculata, Spreng. Syst. ii. 545 = S. paniculata.
pauciflora, Beurling, in Vet. Akad. Handl. Stockh. 1854 (1856) 144 = S. pauciflora.
pellita, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 34 = S. pellita.
petiolaris, H. B. & K. Nov. Gen. et Sp. ii. 171 = S. petiolaris.
plebeia, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 33 = S. cujabana.
Poeppigii, Tul. l. c. 39 = S. Poeppigii.
polyantha, Tul. l. c. 31 = S. polyantha.
pyricarpa, Ruiz & Pav. Syst. Veg. i. 264 = S. macrophylla.
pyricarpa, Willd. ex Tul. in Arch. Mus. Par. viii. (1855–56) 358 = S. mollis.
radiata, Poepp. & Endl. Nov. Gen. et Sp. ii. 48 = S. radiata.
reginae, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 39 = S. reginae.
riparia, Tul. l. c. 36 = S. riparia.
ruficeps, Tul. l. c. 34 = S. ruficeps.
Schottiana, Spreng. Syst. iv. Cur. Post. 407 = Mollinedia brasiliensis.
Selloi, Spreng. Syst. ii. 545 = Mollinedia cinerea.
sessiliflora, H. B. & K. Nov. Gen. et Sp. ii. 174 = S. sessiliflora.
suaveolens, Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 36 = S. suaveolens.
subinodora, Ruiz & Pav. Syst. Veg. i. 265 = S. subinodora.
Thea, Seem. Journ. Bot. ii. (1864) 343 = S. Thea.
thecaphora, Poepp. & Endl. Nov. Gen. et Sp. ii. 48 = S. thecaphora.
tomentosa, Ruiz & Pav. Syst. Veg. i. 265 = S. aspera, tomentosa.
tomentosa, Willd. ex Tul. in Ann. Sc. Nat. Sér. IV. iii. (1855) 36 = S. neglecta.
triflora, Spreng. Syst. ii. 544 = Mollinedia triflora.
umbellata, Spreng. l. c. 545 = Mollinedia umbellata.

CITRULLUS, Forsk. Fl. Aegypt. Arab. 167 (1775); Neck. Elem. i. 240 (1790). *CUCURBITACEAE*, Benth. & Hook. f. i. 826.
amarus, Schrad. in Eckl. & Zeyh. Enum. 279 = vulgaris.
aquosus, Schur, Enum. Pl. Transs. 221 = vulgaris.
Battich, *Forsk. Fl. Aegypt. Arab.* 167.—Arabia.
caffer, Schrad. Ind. Sem. Hort. Gotting. (1834); ex Eckl. & Zeyh. Enum. 279 = vulgaris.
caffrorum, Schrad. in Linnaea, xii. (1838) 413 = vulgaris.
chodospermus, Fal. & Dun. in Bull. Soc. Agr. Hérault (1836), 264 = vulgaris.
Colocynthis, *Schrad. in Linnaea*, xii. (1838) 414.—Afr. calid.
edulis, Spach, Hist. Vég. Phan. vi. 214 = vulgaris.
fistulosus, Stocks, in Hook. Kew Journ. iii. (1851) 74. t. 3 = vulgaris.
Naudinianus, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 549.—Afr. trop.

CITRULLUS:—
Pasteca, Sageret, in Ann. Sc. Nat. Sér. I. viii. (1826) 312 = vulgaris.
Pseudo-Colocynthis, M. Roem. Syn. Pepon. 50 = C. Colocynthis.
variegatus, Schrad. ex M. Roem. l. c. 51 = Cucurbita, Pepo.
vulgaris, *Schrad. ex Eckl. & Zeyh. Enum.* 279.—Afr. trop.

CITRUS, Linn. Syst. ed. I (1735). *RUTACEAE*, Benth. & Hook. f. i. 305.
CITREUM, Tourn. ex Rupp. Fl. Jen. ed. Hall. 141 (1745); Mill. Gard. Dict. ed. VI (1752).
CITROPHORUM, Neck. Elem. ii. 401 (1790).
LIMON, Tourn. ex Mill. Gard. Dict. ed. VI (1752).
OXANTHERA, Montrous. in Mém. Acad. Lyon, x. (1860) 186.
PAPEDA, Hassk. in Flora, xxv. (1842) II. Beibl. 42.
PONCIRUS, Rafin. Sylva Tellur. 143 (1838).
PSEUDAEGLE, Miq. Ann. Mus. Lugd. Bat. ii. 83 (1865–66).
SARCODACTILIS, Gaertn. f. Fruct. iii. 39. t. 185 (1805).
acida, Pers. Syn. ii. 73 = C. Medica.
albida, *Boj. Hort. Maurit.* 49.—Ins. Maurit.
amara, *Hassk. in Flora*, xxv. (1842) *Beibl.* II. 43.—Java.
amara, Link, Handb. ii. 346 = C. Aurantium.
angulata, *Willd. Sp. Pl.* iii. 1426.—Ins. Amboin.
articulata, *Willd. ex Spreng. Syst.* iii. 334.—Afr. trop.
Aurantium, *Linn. Sp. Pl.* 783.—As. trop.
aurata, Risso, Hist. Nat. Or. Europ. Mérid. i. 409 = C. Aurantium.
australasica, *F. Muell. Fragm.* i. 26.—Austral.
australis, *Planch. in Hort. Donat.* 18.—Austral.
Bergamia, Risso, Hist. Nat. Or. Europ. Mérid. t. 53–56 = C. Aurantium.
Bergamota, Rafin. Sylva Tellur. 141 = praec.
Bigaradia, Loisel. in Duham. Arb. ed. nov. vii. 99 = C. Aurantium.
buxifolia, Hort. ex Ten. Atti 3ª Reun. Sc. Ital. (1841) 503 = Atalantia buxifolia.
buxifolia, Poir. in Lam. Encyc. iv. 580 = C. Aurantium.
Calot, Lag. Gen. et Sp. Nov. 17 = C. Aurantium.
cedra, Link, Handb. ii. 346 = C. Medica.
cedrata, Rafin. Sylva Tellur. 141 = C. Medica.
chilensis, Molina, Sagg. Chile, 171 = Villaresia mucronata.
Combara, Rafin. Sylva Tellur. 142.—Malaya.
costata, Rafin. l. c. = decumana?
crassa, *Hassk. Cat. Pl. Hort. Bog. Alt.* 217.—Java.
Daidai, *Bieb. ex Van Houtte, Cat.* n. 126 (1869).—Japon.
decumana, *Murr. Syst.* ed. XIII. 508.—As. trop.
deliciosa, Tenore, Sull' Arancio Mandarino, 11 = C. Aurantium.
dulcimedulla, Pritz. Ic. Bot. Ind. 275 = C. Aurantium?
dulcis, Pers. Syn. ii. 74 = C. Aurantium.
echinata, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 122 = C. Hystrix.
emarginata, *Desf. Cat. Hort. Par.* ed. III. 406.—Hab.?
florida, Salisb. Prod. 378 = C. Aurantium.
fragrans, Salisb. l. c. = C. Medica.
fusca, Lour. Fl. Cochinch. 467 = C. Aurantium.
fusiformis, Rafin. Sylva Tellur. 142.—Hab.?
Gongra, Rafin. l. c. = C. Medica?
grandis, *Hassk. in Flora*, xxv. (1842) *Beibl.* II. 44.—Java.
granulata, Rafin. Sylva Tellur. 143.—Hab.?
heterophyla, Rafin. l. c. 141.—Hab.?
Hillii, *F. Muell.* i. 26.—Austral.
humilis, *Poir. in Lam. Encyc.* iv. 580.—Ind. or.
Hystrix, *DC. Cat. Hort. Monsp.* 97.—Ind. or.; Malaya.
inermis, Roxb. Hort. Beng. 59; Fl. Ind. iii. 393 = japonica.
japonica, *Thunb. Fl. Jap.* 292.—Japon; China.
javanica, *Blume, Bijdr.* 140.—Java.
Karna, Rafin. Sylva Tellur. 142.—Hab.?
latipes, Hook. f. & Thoms. ex Hook. f. Fl. Brit. Ind. i. 515 = C. Hystrix.
Lima, Macfad. in Hook. Bot. Misc. i. (1830) 300 = C. Medica.
Limetta, Risso, in Ann. Mus. Par. xx. (1813) 195. t. 2. f. 1 = C. Medica.

CITRUS:—
Limonellus, Buch.-Ham. ex Wall. Cat. n. 6386 = C. Medica.
Limonum, Risso, in Ann. Mus. Par. xx. (1813) 201 = C. Medica.
lumia, Risso, Hist. Nat. Or. Europ. Mérid. i. 414 = C. Medica.
macracantha, *Hassk. Cat. Pl. Hort. Bog. Alt.* 218.—Java?
macroptera, *Montrous. in Mém. Acad. Lyon*, x. (1860) 187.—N. Caled.
madurensis, Lour. Fl. Cochinch. 467 = japonica.
Margarita, Lour. l. c. = japonica.
marginata, Steud. Nom. ed. II. i. 376 = japonica.
Medica, *Linn. Sp. Pl.* 782.—As. trop.
microcarpa, Bunge, in Mém. Sav. Etr. Pétersb. ii. (1835) 84 = japonica.
mitis, Blanco, Fl. Filip. ed. I. 610 = C. Medica.
myrtifolia, Rafin. Sylva Tellur. 141.—Hab.?
nobilis, Lour. Fl. Cochinch. 466 = C. Aurantium.
notissima, Blanco, Fl. Filip. ed. I. 607 = C. Medica.
obovata, Rafin. Sylva Tellur. 142.—Hab.?
obversa, *Hassk. Cat. Pl. Hort. Bog. Alt.* 218.—Java.
ovata, *Hassk. in Flora*, xxv. (1842 *Beibl.* II. 42.—Java.
paniculata, *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* iv. (1828) 152.—Afr. trop.
Papaya, *Hassk. in Cat. Pl. Hort. Bog. Alt.* 218.
Papedia, Miq. Fl. Ind. Bat. i. II. 530 = C. Hystrix.
papillaris, Blanco, Fl. Filip. ed. I. 610 = C. Aurantium.
Paradisi, Macfad. Fl. Jamaic. i. 131 = C. Aurantium.
Planchoni, *F. Muell. Sel. Pl. Indust. Cult.* 34.—Austral.
pyriformis, *Hassk. in Flora*, xxv. (1842) *Beibl.* II. 44.—Java.
reticulata, Blanco, Fl. Filip. ed. I. 610 = C. Aurantium.
rotundifolia, Rafin. Sylva Tellur. 141.—Hab.?
salicifolia, Rafin. l. c.—Hab.?
scandens, Griff. Notul. iv. 495 = Paramignya Griffithii.
sinensis, Pers. Syn. ii. 74 = C. Aurantium.
spinosa, S. G. Gmel. Reise Russl. iii. 278. t. 22.—Persia.
spinosissima, G. F. W. Mey. Prim. Fl. Esseq. 247 = C. Medica.
torosa, *Blanco, Fl. Filip.* ed. I. 609.—Ins. Philipp.
trifolia, Thunb. Fl. Jap. 294 = trifoliata.
trifoliata, Hort. ex M. Roem. Syn. Hesper. 39 = Glycosmis pentaphylla.
trifoliata, *Linn. Sp. Pl.* ed. II. 1101.—Japon.
triptera, Desf. Cat. Hort. Par. ed. III. 406 = trifoliata.
tuberosa, Mill. Gard. Dict. ed. VIII. n. 2 = C. Medica.
Vangasay, *Boj. Hort. Maurit.* 49.—Hab.?
Volkameriana, *Pasq. Cat. Orto Bot. Nap.* 29.—Hab.?
vulgaris, Risso, in Ann. Mus. Par. xx. (1813) 190 = C. Aurantium.

CITTA, Lour. Fl. Cochinch. 456 (1790) = **Mucuna**, Adans. (Legumin.).
nigricans, Lour. l. c. = M. nigricans.

CITTARONIUM, Reichb. Nom. 186 (1841) = **Viola**, Tourn.

CITTORHINCHUS, Willd. ex H. B. & K. Nov. Gen. et Sp. vi. 14 (1823) = **Gomphia**, Schreb. (Ochnac.).
javitensis, Willd. l. c. = G. aquatica.

CLADANTHUS, Cass. in Bull. Soc. Philom. (1816). *COMPOSITAE*, Benth. & Hook. f. ii. 421.
arabicus, *Cass. l. c.*; *et in Dict. Sc. Nat.* ix. 342. t. 87.—Arab. (Quid?).
canescens, Sweet, Hort. Brit. ed. III. 366.—Ins. Canar.
Geslini, Coss. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 397 = Mecomischus Geslini.
halimifolius, Coss. & Dur. l. c. = Mecomischus Geslini.
pedunculatus, Benth. & Hook. f. Gen. ii. 422, err. typ. = seq.
pedunculatus, Dur. in Bull. Soc. Bot. Fr. iv. (1857) 14 = Anthemis pedunculatus.
proliferus, *DC. Prod.* vi. 18.—Afr. bor.

CLADERIA, Rafin. Sylva Tellur. 12 (1838) = **Melia**, Linn.?
parviflora, Rafin. l. c.—Hab.?

CLADIUM, P. Br. Hist. Jamaic. 114 (1756). *CYPERACEAE*, Benth. & Hook. f. iii. 1065.

CLADIUM:—
AGYLLA, Phil. in Anal. Univ. Chil. (1865) I. 643.
BAUMEA, Gaudich. in Freyc. Voy. Bot. 416. t. 29 (1826).
CHAPELLIERA, Nees, in Linnaea, ix. (1834) 298
MACHAERINA, Vahl, Enum. ii. 228 (1806).
SCHOENOPSIS, Beauv. in Lestib. Essai Fam. Cyp. 341 (1819).
TEROBERA, Steud. Syn. Pl. Cyp. 164 (1855).
TRASI, Beauv. in Lestib. Essai Fam. Cyp. 32 (1819).
TRACHYRHYNCHIUM, Nees, in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 113.
TRACHYRHYNGIUM, Nees, ex Kunth, Enum. Pl. ii. 314 (1837).
VINCENTIA, Gaudich. in Freyc. Voy. Bot. 417 (1826).
acutum, Poir. in Dict. Sc. Nat. ix. 344 = schoenoides.
anceps, Hook. f. Handb. New Zeal. Fl. 305, in obs. = iridifolium.
ancistrophyllum, F. Muell. ex Benth. Fl. Austral. vii. 415 = Gahnia ancistrophylla.
aristatum, F. Muell. ex Benth. l. c. 416 = Gahnia aristata.
articulatum, *R. Br. Prod.* 237.—Austral.
asperum, F. Muell. Fragm. ix. 12 = Gahnia aspera.
athrophyllum, *F. Muell. l. e.* 49.—Austral.
bahiense, Nees, in Linnaea, ix. (1834) 301 = C. germanicum.
chinense, Nees, l. c.; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 116 = germanicum.
complanatum, *Bergg. in Minneskr. Fisiog. Sallks. Lund*, (1877) n. 8. p. 23.—N. Zel.
decompositum, R. Br. Prod. 237 = Gahnia decomposita.
deustum, R. Br. l. c. = Gahnia deusta.
dubium, Spreng. Syst. iv. Cur. Post. 21 = glomeratum
effusum, Torr. in Ann. Lyc. N. York, iii. (1836) 374 = germanicum.
elynanthoides, F. Muell. Fragm. ix. 31.—Austral.
ensifolium, *Benth. & Hook. f. Gen.* iii. 1066.—Bras.
ensigerum, *Hance, in Journ. Bot.* xxiii. (1885) 80.—China.
ficticum, *Hemsl. Bot. Challenger, Voy.* iii. 59, in obs. t. 60. f. 5–7.—Am. austr.
filifolium, *Benth. & Hook. f. Gen.* iii. 1065.—Cuba.
Filum, R. Br. Prod. 237, partim = Gahnia trifida.
floribundum, Presl, in Oken, Isis, xxi. (1828) 270 = germanicum.
gahnoides, Colenso, in Trans. N. Z. Inst. xvi. (1884) 340 = Sinclairi.
germanicum, *Schrad. Fl. Germ.* i. 75. t. 5. f. 7.—Reg. temp. et subtrop.
giganteum, Willk. in Willk. & Lange, Prod. Fl. Hisp. i. 136 = germanicum.
glomeratum, *R. Br. Prod.* 237.—As. or.; Australas.
Gunnii, *Hook. f. Fl. Tasm.* ii. 95. t. 148 B.—Austral.; N. Zeland.
Huttoni, *T. Kirk. in Trans. N. Z. Inst.* ix. (1877) 551.—N. Zeland.
insulare, *Benth. Fl. Austral.* vii. 403.—Austral.
iridifolium, *Baker, Fl. Maurit.* 424.—Ins. Mascar.; Bras.
jamaicense, Crantz, Inst. i. 362 = germanicum.
japonicum, Steud. Syn. Pl. Cyp. 152 = germanicum.
junceum, *R. Br. Prod.* 237.—Austral.
lanigerum, R. Br. l. c. = Gahnia lanigera.
latissimum, *F. Muell. Fragm.* ix. 15.—Austral.
laxiflorum, Hook. f. Fl. Tasm. ii. 95. t. 148 A = Gunnii.
laxum, *Benth. Fl. Austral.* vii. 405.—Austral.
leptostachyum, [Nees, in] Meyen, Reise, ii. 119; et in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 115 = germanicum.
mariscoides, Torr. in Ann. Lyc. N. York, iii. (1836) 372 = triglomeratum.
mariscoides, *F. Villar. in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 309.—Ins. Philipp.
Mariscus, R. Br. Prod. 236 = germanicum.
medium, R. Br. l. c. 237 = Gahnia lanigera.
melanocarpum, F. Muell. Fragm. ix. 13 = Gahnia melanocarpa.
Melleri, *Baker, in Journ. Linn. Soc.* xxi. (1885) 451 = Madag.
microstachyum, F. Muell. Census, 129 = Gahnia microstachya.
nudum, Boeck. in Linnaea, xxxviii. (1874) 236 = Gunnii.

CLADIUM :—
occidentale, Schrad. Fl. Germ. i. 76 = germanicum.
palustre, Poir. in Dict. Sc. Nat. ix. 344 = germanicum.
pantopodum, *Baker, in Journ. Linn. Soc.* xxi. (1885) 451.—Madag.
pauciflorum, R. Br. Prod. 237 = Tricostularia pauciflora.
polyphyllum, F. Muell. Census, 129 = Gahnia polyphylla.
Preissii, F. Muell. ex Benth. Fl. Austral. vii. 405 = latissimum.
psittacorum, F. Muell. Fragm. ix. 13 = Gahnia psittacorum.
quadrangulare, Nees, in Linnaea, ix. (1834) 301 = Lampocarya Gaudichaudii.
Radula, R. Br. Prod. 237 = Gahnia Radula.
rigidifolium, Presl, in Oken, Isis, xxi. (1828) 270 = teretifolium.
riparium, *Benth. Fl. Austral.* vii. 405.—Austral.
schoenoides, *R. Br. Prod.* 237.—Austral.
scirpoideum, *Benth. & Hook. f. ex Hemsl. Bot. Challenger Voy.* iii. 59. t. 60. f. 1–4.—Ins. Juan Fernand.
scleroides, F. Muell. Fragm. ix. 12 = Exocarya scleroides.
Sieberi, F. Muell. l. c. 14 = Gahnia Sieberi.
Sinclairii, *Hook. f. Handb. N. Zeal. Fl.* 304.—N. Zel.
teretifolium, *R. Br. Prod.* 237.—Austral.
tetragonocarpum, F. Muell. Census, 129 = Gahnia tetragonocarpa.
tetraquetrum, *Hook. f. Fl. Tasm.* ii. 95. t. 149.—Tasmania.
trifidum, F. Muell. in Papers Roy. Soc. Tasm. (1878) 117 = Gahnia trifida.
triglomeratum, *Nees, in Linnaea*, ix. (1834) 301.—Am. bor.
undulatum, *Thw. Enum. Pl. Zeyl.* 353.—Malaya; Austral.
vaginale, *Benth. Fl. Austral.* vii. 408.—Austral.
xanthocarpum, F. Muell. Fragm. ix. 13 = Gahnia xanthocarpa.
xipholepis, Baker, Fl. Maurit. 424 = Schoenus clongatus.

CLADOBIUM, Lindl. Introd. Nat. Syst. ed. II. 446 (1836) = **Scaphyglottis**, Poepp. (Orchid.).
violaceum, Lindl. l. c. = S. violacea.

CLADOCAULON, Gardn. in Hook. Ic. Pl. t. 528 (1843) = **Paepalanthus**, Mart. (Eriocauleae).
brasiliense, Gardn. l. c. = P. Gardnerianus.

CLADOCHAETA, DC. Prod. vi. 245 (1837) = **Helichrysum**, Vaill. (Compos.).
candidissima, DC. l. c. = H. candidissimum.

CLADODA et CLADODEA, Poir. in Dict. Sc. Nat. ix. 345 (1817) = seq.

CLADODES, Lour. Fl. Cochinch. ii. 574 (1790) = **Alchornea**, Sw. (Euphorbiac.).
ilicifolia, Baill. Adansonia, vi. (1866) 1 = A. ilicifolia.
rugosa, Lour. Fl. Cochinch. ii. 574 = A. rugosa.
Thozetiana, Baill. Adansonia, vi. (1866) 321 = A. Thozetiana.

CLADOGYNOS, Zipp. ex Span. in Linnaea, xv. (1841) 349. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 323.
orientalis, *Zipp. l. c.*—Malaya.

CLADOLEPIS, Moq. in DC. Prod. xiv. II. 203, in syn. (1849) = **Ofaiston**, Rafin. (Chenopod.).

CLADOMISCHUS, Klotzsch, ex A. DC. in DC. Prod. xv. I. 278 (1864) = **Begonia**, Linn.

CLADOPOGON, Sch. Bip. in Ind. Sem. Hort. Hamb. (1852) 8, ex Lehm. in Ann. Sc. Nat. Sér. III. xix. (1853) 359 = **Senecio**, Tourn. (Compos.).
aurantiacum, Lehm. ll. cc. 8; 360.—Mexic.?

CLADOSERIS, Spach, Hist. Vég. Phan. x. 35 (1841) = **Onoseris**, DC. (Compos.).

CLADOSICYOS, Hook. f. in Oliver, Fl. Trop. Afr. ii. 534 (1871) = **Cucumeropsis**, Naud. (Cucurbitac.).
edulis, Hook. f. l. c. = Cucum. edulis.

CLADOSTACHYS, D. Don, Prod. Fl. Nep. 76 (1825) = **Deeringia**, R. Br. (Amarantac.).
alternifolia, Sweet, Hort. Brit. ed. I. 336 = Digera arvensis.
arborescens, D. Dietr. Syn. Pl. i. 861 = Digera arvensis.
frutescens, D. Don, Prod. Fl. Nep. 76 = Deeringia celosioides.
muricata, Moq. in DC. Prod. xiii. II. 235 = Digera arvensis.

CLADOSTEMON, A. Br. & Vatke, in Monatsb. Acad. Berl. (1876) 866. *CAPPARIDEAE.*
paradoxus, *A. Br. & Vatke, l. c.*—Afr. trop.

CLADOSTIGMA, Radlk. in Bremen Abh. viii. (1883) 412. *CONVOLVULACEAE.*
dioicum, *Radlk. l. c.*—Abyssin.

CLADOSTYLES, Humb. & Bonpl. Pl. Aequin. i. 202. t. 57 (1808) = **Evolvulus**, Linn. (Convolvulac.).
ericoides, Nees, ex Steud. Nom. ed. II. i. 377 = E. glomeratus.
paniculata, H. B. & K. Pl. Aeqnin. i. 202 = E. paniculatus.

CLADOTHAMNUS, Bong. Veg. Sitcha, 37. t. 1, in Mém. Acad. Pétersb. Sér. VI. ii. (1833). *ERICACEAE*, Benth. & Hook. f. ii. 598.
TOLMIEA, Hook. Fl. Bor. Am. ii. 44 (1834).
pyroliflorus, *Bong. ll. cc.*—Am. bor. occ.

CLADOTHECA, Steud. Syn. Pl. Cyp. 178 (1855) = **Cryptangium**, Schrad. (Cyperac.).
vaginans, Steud. l. c. = Crypt. leptocladum.

CLADOTHRIX, Nutt. ex Moq. in DC. Prod. xiii. II. 359 (1849). *AMARANTACEAE*, Benth. & Hook. f. iii. 37.
lanuginosa, *Nutt. ex Moq. l. c.* 360.—Calif.
oblongifolia, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 376.—Am. bor. occ.

CLADOTRICHIUM, Vog. in Linnaea, xi. (1837) 401 = **Caesalpinia**, Linn. (Legumin.).
epunctatum, Vog. l. c. 403 = Caes. epunctata.
pilosum, Vog. l. c. = Caes. pilosa.
rubicundum, Vog. l. c. 401 = Caes. rubicnnda.
stipulare, Vog. l. c. 403 = Caes. stipularis.

CLADRASTIS, Rafin. Neogenyt. 1 (1825). *LEGUMINOSAE*, Benth. & Hook. f. i. 554.
BUERGERIA, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 53 (1867).
MAACKIA, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1856) 128, 143.
albiflora, *Rafin. New Fl. Am.* iii. 83.—Am. bor.
amurensis, *Benth. ex Maxim. in Bull. Acad. Pétersb.* xviii. (1873) 400.—Reg. Amur.
kentuckensis, Rafin. Neogenyt. (1825) 1 = tinctoria?
lutea, Rafin. l. c. = tinctoria.
secundiflora, Rafin. l. c. = Sophora secundiflora.
tinctoria, *Rafin. l. c.; New Fl. Am.* iii. 83.—Am. bor.

CLAIRISIA, Abat, ex Benth. & Hook. f. Gen. iii. 78 (1873) = CLARISIA, Abat = **Anredera**, Juss. (Chenopodiac.).

CLAIRVILLEA, DC. Prod. v. 636 (1836) = **Cacosmia**, H. B. & K. (Compos.).
quinquenervia, DC. l. c. = Cacosmia rugosa.

CLAMBUS, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xviii. (1866) 16. *MENISPERMACEAE*, Benth. & Hook. f. i. 962.
araneosus, *Miers, l. c.* 17.—Mexic.

CLAMYDANTHUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 147 = CHLAMYDANTHUS, C. A. Mey. = **Thymelaea**, Linn.

CLANDESTINA, Tourn. ex Adans. Fam. ii. 207 (1763) = **Lathraea**, Linn. (Orobanch.).
japonica, Miq. in Versl. en Med. Kon. Akad. Wetensch. II. ii. (1868) 85 = L. Miqueliana.
penduliflora, Lam. Fl. Fr. ii. 329 = L. Clandestina.
rectiflora, Lam. l. c. 328 = L. Squamaria.

CLANDESTINARIA, Spach, Hist. Vég. Phan. vi. 427 (1838) = **Nasturtium**, R. Br. (Crucif.).
indica, Spach, l. c. 429 = N. indicum.

CLAOTRACHELUS, Zoll. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 565 = **Vernonia**, Schreb. (Compos.).
rupestris, Zoll. & Mor. l. c.—Java.

CLAOXYLON, A. Juss. Euph. Tent. 43. t. 14 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 309.
ERYTHROCHILUS, Reinw. ex Flora, viii. (1825) I. 103.
affine, Baill. Adansonia, ii. (1861–62) 227 = insulanum.
affine, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie*, ii. (1845) 582.—Java.
africanum, Muell. Arg. in DC. Prod. xv. II. 777 = Barteri.
angolense, *Muell. Arg. in Journ. Bot.* ii. (1864) 333.—Afr. occ.
angustifolium, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 165.—Austral.
australe, *Baill. Etud. Gén. Euph.* 493.—Austral.
Barteri, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 21.—Afr. occ.
capense, *Baill. Etud. Gén. Euph.* 493.—Afr. austr.
columnare, *Muell. Arg. in Flora*, xlvii. (1864) 437.—Afr. trop.
cordifolium, Benth. in Hook. Niger Fl. 506 = Mallotus oppositifolius.
coriaceum, Baill. Etud. Gén. Euph. 493 = erythrophyllum.
crassifolium, *Baill. Adansonia*, i. (1860–61) 279.—Afr. trop.
digynum, Wight, Ic. t. 1884 = Macaranga digyna.
echinospermum, *Muell. Arg. in DC. Prod.* xv. II. 787.—Ins. Pacif.
erythrophyllum, *Miq. Fl. Ind. Bat.* i. II. 387.—Java.
fallax, *Muell. Arg. in DC. Prod.* xv. II. 780.—Ins. Pacif.
glabrifolium, *Miq. Fl. Ind. Bat.* i. II. 387.—Java.
glandulosum, *Baill. Adansonia*, i. (1860–61) 281.—Ins. Mascar.
graciliflorum, Miq. Fl. Ind. Bat. i. II. 386 = affine.
grandidentatum, Boiv. ex Baill. in Adansonia, i. (1860–61) 283 = racemiflorum.
grandifolium, Muell. Arg. in Linnaea, xxxiv. (1865–66) 163 = crassifolium.
hexandrum, *Muell. Arg. in Flora*, xlvii. (1864) 438.—Afr. occ.
Hillii, *Benth. Fl. Austral.* vi. 131.—Austral.
hirtellum, *Baill. Adansonia*, i. (1860–61) 284.—Madag.
indicum, *Hassk. Cat. Hort. Bogor. Alt.* 235.—As. trop.
insulanum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 164.—N. Caled.
Kirkii, *Muell. Arg. in Flora*, xlvii. (1864) 436.—Afr. occ.
leucocarpum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 244.—Burma.
linostachys, *Baill. Adansonia*, i. (1860–61) 281.—Ins. Maurit.
longifolium, Baill. Etud. Gén. Euph. 493, pro parte = oligandrum.
longifolium, *Muell. Arg. in DC. Prod.* xv. II. 781.—Malaya.
longipetiolatum, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 244.—Burma.
macranthum, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 164.—Ins. Madag.
macrophyllum, Boj. Hort. Maurit. 284 = indicum.
Mannii, *Hook. f. in Journ. Linn. Soc.* vi. (1862) 20.—Afr. trop.
marianum, *Muell. Arg. in DC. Prod.* xv. II. 783.—Ins. Marian.
medullosum, *Baill. Adansonia*, i. (1860–61) 283.—Madag.
membranaceum, *Muell. Arg. in Flora*, xlvii. (1864) 437.—Afr. trop.

CLAOXYLON:—
Mercurialis, *Thw. Enum. Pl. Zeyl.* 271.—As. trop.; Afr. trop.
Metzicrii, *Bouton, ex Boj. Hort. Maurit.* 285.—Ins. Maurit.
minus, Hassk. Pl. Jav. Rar. 251 = indicum.
molle, Miq. Fl. Ind. Bat. i. II. 386 = indicum.
monoicum, *Baill. Adansonia*, i. (1860-61) 283.—Madag.
muricatum, Wight, Ic. t. 1886 = Mallotus muricatus.
Neraudianum, *Baill. Adansonia*, i. (1860-61) 280.—Afr. trop.
occidentale, *Muell. Arg. in Flora*, xlvii. (1864) 438.—Afr. trop.
oligandrum, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 164.—Afr. trop.
parviflorum, Hook. & Arn. Bot. Beech. Voy. 212 = indicum.
pariflorum, *A. Juss. Tent. Euph.* 114. t. 14. f. 48.—Ins. Mascar.
parviflorum, Seem. in Bonplandia, ix. (1861) 258 = fallax.
pauciflorum, *Muell. Arg. in Journ. Bot.* ii. (1864) 333.—Afr. trop.
pedicellare, *Muell. Arg. in Flora*, xlvii. (1864) 437.—Afr. trop.
racemiflorum, *Baill. Etud. Gén. Euph.* 493.—Ins. Borbon.
reticulatum, *Boj. Hort. Maurit.* 285.—Ins. Maurit.
rivulare, *Muell. Arg. in Flora*, xlvii. (1864) 518.—Afr. trop.
rubescens, *Miq. Fl. Ind. Bat.* i. II. 337.—Ins. Pacif.
rubrinerve, Baill. Etud. Gén. Euph. 493, nomen = rubescens.
sandwicense, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 165.—Ins. Sandvic.
scabrum, Boj. Hort. Maurit. 285, nomen = linostachys.
spiciflorum, Baill. in Adansonia, i. (1860-61) 285 = indicum.
spiciflorum, A. Juss. Tent. Euph. 43 = Acalypha spiciflora.
stipulosum, Reichb. [f.] ex Zoll. in Verh. Naturk. Ver. Nederl. Indie, i. (1856) 20; et in Linnaea, xxviii. (1856) 323 = Mallotus stipulosus.
taitense, *Muell. Arg. in Linnaea*, xxxiv. (1865-66) 165.—Ins. Tahiti.
tenerifolium, *F. Muell. ex Baill. Adansonia*, vi. (1866) 323.—Austral.
trichogyne, *Muell. Arg. in Journ. Bot.* ii. (1864) 334.—Afr. trop.
triste, *Muell. Arg. l. c.*—Afr. trop.
Wallichianum, *Muell. Arg. in DC. Prod.* xv. II. 781.—Malaya.
Welwitschianum, *Muell. Arg. in Journ. Bot.* ii. (1864) 333.—Afr. trop.

CLAPPERTONIA, Meissn. Gen. Pl. 36 [28] (1837) = **Honckenya**, Willd. (Tiliac.).
ficifolia, Decne. in Deless. Ic. Select. Pl. v. 1. t. 1 = H. ficifolia.

CLAPPIA, A. Gray, in Torr. Bot. Mex. Bound. 93. (1858). *COMPOSITAE*, Benth. & Hook. f. ii. 413.
aurantiaca, Benth. in Hook. Ic. Pl. t. 1104 (= Dysodicae sp.).—Mexic.
suaedaefolia, *A. Gray, in Torr. Bot. Mex. Bound.* 93.—Mexic.

CLARA, Kunth, in Abh. Akad. Berl. (1848) 44 = **Herreria**, Ruiz & Pav. (Liliac.).
ophiopogonoides, Kunth, Enum. Pl. v. 296.—Bras.

CLARCKIA, Pursh, Fl. Am. Sept. i. 260 (1814) = **Clarkia**, Pursh (Onagrar.).

CLARIONA, Spreng. Syst. iii. 504 (1826) = seq.

CLARIONEA, Lag. ex DC. in Ann. Mus. Par. xix. (1812) 65 = **Perezia**, Lag. (Compos.).
acanthoides, D. Don, ex Hook. & Arn. in Comp. Bot. Mag. i. (1835) 34 = P. multiflora.
affinis, Phil. in Anal. Univ. Chil. (1873) 479.—Chili.
carthamoides, D. Don, in Phil. Mag. (1832) 328 = P. diversifolia.
ciliaris, D. Don, ex Hook. & Arn. in Comp. Bot. Mag. i. (1835) 34 = P. ciliaris.
dicephala, DC. Prod. vii. 62 = P. dicephala.
elegans, Phil. in Linnaea, xxviii. 1856) 717.—Chili.
Fonki, Phil. l. c. 718.—Chili.

CLARIONEA :—
glaberrima, Cass. Opusc. ii. 165 = Perezia lactucoides.
humilis, Phil. in Linnaea, xxviii. (1856) 717.—Chili.
laciniata, Phil. in Anal. Univ. Chil. (1873) 479.—Chili.
lactucoides, D. Don, in Trans. Linn. Soc. xvi. (1833) 206 = P. lactucoides.
Lechleri, Sch. Bip. in Flora, xxxviii. (1855) 122.—Peruv.
macrocephala, Sch. Bip. in Lechl. Berb. Am. Austr. 57.—Peruv.
magellanica, DC. in Ann. Mus. Par. xix. (1812) 66. t. 3 = P. magellanica.
multicapitata, Remy, in C. Gay, Fl. Chil. ii. 410 = P. multicapitata.
parvifolia, Phil. in Anal. Univ. Chil. (1873) 480.—Chili.
pedicularifolia, DC. Prod. vii. 61 = P. pedicularidifolia.
pilifera, D. Don, in Phil. Mag. (1832) 388 = P. pilifera.
pinnata, Phil. in Linnaea, xxviii. (1856) 718.—Chili.
pinnatifida, DC. Prod. vii. 62 = P. pinnatifida.
polycephala, Cass. Opusc. ii. 167 = P. multiflora.
pungens, DC. Prod. vii. 62 = P. pungens.
recurvata, D. Don, in Trans. Linn. Soc. xvi. (1830) 206 = P. Doniana.
runcinata, Lag. ex D. Don, l. c. 207 = P. runcinata.
spathulata, Lag. ex D. Don, l. c. 205 = P. viscosa.
variabilis, Phil. in Linnaea, xxxiii. (1864-65) 125.—Chili.
virens, D. Don, in Trans. Linn. Soc. xvi. (1833) 208 = P. Poeppigii.
Volckmanni, Phil. in Anal. Univ. Chil. (1873) 480.—Chili.

CLARIONELLA, DC. ex Steud. Nom. ed. II. i. 377 (1840) = CLARIONEA, Lag. = **Perezia**, Lag. (Compos.).
magellanica, Hombr. & Jacquin. ex Decne. Bot. Voy. Astrol. & Zél. 45 = P. magellanica.

CLARIONEMA, Phil. in Linnaea, xxviii. (1856) 717 = praec.

CLARIONIA, D. Don, in Trans. Linn. Soc. xvi. (1830) 204 = praec.

CLARISIA, Abat, in Mem. Soc. Med. Sev. x. (1792) 418.—Cf. Spreng. Gen. 202 (1830) = **Anredera**, Juss. (Chenopod.).
volubilis, Abat, l. c.; ex Moq. in DC. Prod. xiii. II. 230 = A. spicata.

CLARISIA, Ruiz & Pav. Prod. Fl. Per. & Chil. 158. t. 28 (1794). *URTICACEAE*, Benth. & Hook. f. iii. 377.
Soaresia, Fr. Allem. in Palestr. Scient. Rio Jan. (1858) 142.
biflora, *Ruiz & Pav. Syst. Veg.* 256.—Peruv.
racemosa, *Ruiz & Pav. l. c.* 255.—Peruv.

CLARKELLA, Hook. f. Fl. Brit. Ind. iii. 46 (1880). *RUBIACEAE*.
nana, *Hook. f. l. c.*—Reg. Himal.

CLARKIA, Pursh, Fl. Am. Sept. i. t. 11 (1814). *ONAGRARIEAE*, Benth. & Hook. f. i. 789.
Guaropsis, Presl, Epim. Bot. 219 (1849).
Opsianthes, Lilja, Fl. Sverig. Suppl. 25 (1840).
Phaeostoma, Spach, Hist. Veg. Phan. iv. 392 (1835).
Eisenеana, *Kellogg, in Proc. Calif. Acad.* vii. (1877) 94.—Calif.
elegans, *Dougl. in Lindl. Bot. Reg.* t. 1575.—Am. bor. occ.
elegans, Poir. in Dict. Sc. Nat. ix. 255 = pulchella.
gauroides, Dougl. ex Sweet, Brit. Flow. Gard. Scr. II. t. 379 = rhomboidea.
pulchella, *Pursh, Fl. Am. Sept.* i. t. 11.—Am. bor. occ.
pulcherrima, Hort. ex L'Horticul. Franç. (1861) 7 = pulchella.
rhomboidea, *Dougl. in Hook. Fl. Bor. Am.* i. 214.—Am. bor. occ.
unguiculata, Lindl. Bot. Reg. t. 1981 = elegans.
Xantiana, *A. Gray, in Proc. Bost. Soc. Nat. Hist.* vii. (1859-61) 145.—Calif.

CLASTA, Comm. ex Vent. Choix, sub t. 47 (1803) = **Casearia**, Jacq. (Samydac.).
fragilis, Comm. ex Vent. l. c. = Cas. fragilis.

CLASTILIX, Rafin. Sylva Tellur. 99 (1838) = **Miconia**, Ruiz & Pav. (Melast.).
mexicana, Rafin. l. c. = M. mexicana.
tunicata, Rafin. l. e. = M. tunicata.

CLASTOPUS, Bunge, ex Boiss. Fl. Orient. i. 261 (1867) = **Vesicaria**, Tourn. (Crucif.).
purpureus, Bunge, l. c.—Persia.
vestitus, Boiss. l. c.—Persia.

CLATHROSPERMUM, Planch. ex Benth. & Hook. f. Gen. i. 29 (1862). *ANONACEAE*, Benth. & Hook. f. l. c.
biovulatum, *S. Moore, in Journ. Bot.* xv. (1877) 65.—Afr. trop.
Mannii, *Oliver, Fl. Trop. Afr.* i. 25.—Afr. trop.
Vogelii, *Planch. ex Benth. in Trans. Linn. Soc.* xxiii. (1862) 479.—Afr. trop.

CLAUCENA, Burm. f. Fl. Ind. 87 (1768) = **Clausena**, Burm. f. (Rutac.).

CLAUDIA, Opiz, ex Panz. in Denkschr. Bot. Ges. (1808) II. 177 = **Melica**, Linn. (Gramin.).
ciliata, Opiz, l. c. = M. ciliata.

CLAUSENA, Burm. f. Fl. Ind. 243 (1768). *RUTACEAE*, Benth. & Hook. f. i. 304.
Aulacia, Lour. Fl. Cochinch. 273 (1790).
Cookia, Sonner. Voy. ii. 130. t. 131 (1782).
Fagarastrum, G. Don, Gen. Syst. ii. 87 (1832).
Gallesioa, M. Roem. Syn. Hesper. 45 (1846).
Glaucena, Vitm. Summa, ii. 441 (1789).
Myaris, Presl, Bot. Bemerk. 40 (1849).
Quinaria, Lour. Fl. Cochineh. 272 (1790).
Piptostylis, Dalz. in Hook. Kew Journ. iii. (1851) 33. t. 2.
Polycyema, Voigt, Hort. Suburb. Calc. 141 (1845).
anisata, *Hook. f. in Hook. Niger Fl.* 256.—Afr. trop.
brevistyla, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 31.—Austral.
ehrysogyne, *Miq. Fl. Ind. Bat. Suppl.* i. 502.—Malaya.
corymbiflora, *Linden, Cat.* n. 98 (1878), 3.—Ins. Loyalt.
crenulata, F. Muell. Phyt. New Hebr. 7 = Murraya crenulata.
dentata, M. Roem. Syn. Hesper. 44 = Willdenowii.
excavata, *Burm. f. Fl. Ind.* 89. t. 29.—Ind. or.; Malaya.
Halmaheirae, *Miq. Ann. Mus. Bot. Lugd. Bat.* i. 211.—Malaya.
heptaphylla, *Wight & Arn. Prod.* 95, in nota.—Ind. or.; Burma.
inequalis, *Benth. in Hook. Niger Fl.* 257.—Afr. trop.
indica, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 36.—Ind. or.
javanensis, Raeusch. ex DC. Prod. i. 538 = excavata.
javanica, *M. Roem. Syn. Hesper.* 45.—Java.
javensis, J. F. Gmel. Syst. 610 = excavata?
macrophylla, *Hook. f. Fl. Brit. Ind.* i. 504.—Burma.
nana, Wight & Arn. Prod. 96 = Willdenowii.
pentaphylla, *DC. Prod.* i. 538.—Reg. Himal.
pubescens, Wight & Arn. Prod. 96 = Willdenowii.
punctata, Wight & Arn. l. e. 95 = excavata.
simplicifolia, Dalz. in Hook. Kew Journ. iii. (1851) 180 = Acronychia laurifolia.
suffruticosa, *Wight & Arn. Prod.* 96, in nota.—Reg. Himal.
sumatrana, Wight & Arn. l. c. 95 = excavata.
timorensis, *M. Roem. Syn. Hesper.* 44.—Ins. Timor.
Wallichii, *Oliver, in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 35.—Burma.
Wampi, *Oliver, l. c.* 34.—China?
Wightii, M. Roem. Syn. Hesper. 44 = Willdenowii?
Willdenowii, *Wight & Arn. Prod.* 96.—Ind. or.; Malaya.

CLAUSIA, Trotzky, Ind. Sem. Hort. Casan. (1839) = **Hesperis**, Linn. (Crucif.).
aprica, Trotzky, l. c. = H. aprica.

CLAUSONIA, Pomel, Mat. Fl. Atl. 1 (1860) = **Asphodelus**, Tourn. (Liliac.).
acaulis, Pomel, l. c. = A. acaulis.

CLAVARIA, Steud. Nom. ed. I. 204 (1821), sphalm. = **Calvaria**, Gaertn. (Gen. dub.).

CLAVENA, DC. Prod. vi. 633 (1837) = **Carduus**, Tourn. (Compos.).
baecephala, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 373 = Card. baecephalus.
canariensis, DC. Prod. vi. 633 = Card. clavulatus.
squarrosa, DC. l. c. = Card. squarrosus.

CLAVENNA, Neck. Elem. ii. 145 (1790) = **Ammannia**, Houst. ? (Lythrar.).

CLAVIGERA, DC. Prod. v. 127 (1836) = **Brickellia**, Ell. (Compos.).
brachyphylla, A. Gray, Pl. Fendl. 63 = B. brachyphylla.
corymbosa, DC. Prod. v. 128 = B. corymbosa.
dentata, DC. l. c. = B. Riddellii.
pinifolia, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 461 = B. pinifolia.
Riddellii, Torr. & Gray, Fl. N. Am. ii. 77 = B. Riddellii.
scabra, Benth. Pl. Hartw. 19 = B. corymbosa.
scoparia, DC. Prod. v. 128 = B. scoparia.
spinulosa, A. Gray, Pl. Fendl. 63 = B. spinulosa.

CLAVIJA, Ruiz & Pav. Prod. 142. t. 30 (1794). *MYRSINEAE*, Benth. & Hook. f. ii. 649.
HORTA, Vell. Fl. Flum. 48 (1825); i. t. 124.
THEOPHRASTA, Linn. Gen. ed. I. 45 (1737).
ZACINTHA, Vell. Fl. Flum. 276 (1825); viii. t. 9.
biborrana, *Oerst. in Kjoeb. Vidensk. Meddel.* (1861) 117.—Costa Rica.
caloneura, *Mart. ex Miq. in Mart. Fl. Bras.* x. 277.—Bras.
clavata, *Decne. in Ann. Sc. Nat.* Sér. VI. iii. (1876) 144.—Hab. ?
grandis, *Decne. l. c.*—Hab. ?
fulgens, *Hook. f. Bot. Mag.* t. 5626.—Peruv.
Hookeri, *DC. Prod.* viii. 148.—Peruv.
integrifolia, *Mart. ex Miq. in Mart. Fl. Bras.* x. 277.—Bras.
lancifolia, *Desf. in Nouv. Ann. Mus. Par.* i. (1832) 402. t. 14.—Guiana.
latifolia, *C. Koch, in C. Koch & Fint. Wochenschr.* (1859) 108.—Ind. occ.
longifolia, *Ruiz & Pav. Syst. Veg.* i. 284.—Peruv.
macrocarpa, *Ruiz & Pav. l. c.*—Am. austr.
macrophylla, *Miq. in Mart. Fl. Bras.* x. 275.—Bras.
ornata, *D. Don, in Edinb. Phil. Journ.* (Jan. 1831) 236.—Am. austr.
pendula, *Ruiz & Pav. Syst. Veg.* i. 285.—Am. austr.
Riedeliana, *Regel, Gartenfl.* (1859) 245.—Bras.
Rodekiana, *Linden & André, Illustr. Hortic.* xxi. (1874) t. 188.—N. Granat.
sparsifolia, Miq. in Mart. Fl. Bras. x. 278 = Leonia glycycarpa.
spathulata, Hook. Ic. Pl. t. 140 = Hookeri.
spathulata, *Ruiz & Pav. Syst. Veg.* i. 285.—Am. austr.
speciosa, *C. Koch & Linden, Ind. Sem. Hort. Berol.* (1867) *App.* i. 8.—Am. trop.
umbrosa, *Regel, Ind. Sem. Hort. Petrop.* (1868) 86.—Bras.
undulata, *D. Don, in Edinb. Phil. Journ.* (Jan. 1831) 234.—Am. austr.

CLAVIMYRTUS, Blume, Mus. Bot. Lugd. Bat. i. 113. t. 49 (1849) = **Eugenia**, Linn. (Myrtac.).
claviflora, Blume, l. c. 114, in obs. = E. claviflora.
cylindrica, Blume, l. c. 115 = E. cylindrica.
firma, Blume, l. c. 116.—Java.
glabrata, Blume, l. c. 114.—Java.
latifolia, Blume, l. c. 117.—Borneo.
lineata, Blume, l. c. 116 = E. lineata, *Duthie*.
marginata, Blume, l. c. 115.—Java.
pauciflora, Blume, l. c. 114, in obs. = E. laeta.
ramosissima, Blume, l. c. 115 = E. ramosissima.
symphytocarpa, Blume, l. c. 117.—Sumatra.
virens, Blume, l. c. 114.—Java.

CLAVULA, Dum. Fl. Belg. 143 (1827) = **Eleocharis**, R. Br. & **Scirpus**, Linn. (Cyperac.).
acicularis, Dum. l. c. = E. acicularis.
Boeotryon, Dum. l. c. = S. pauciflorus.
caespitosa, Dum. l. c. = S. caespitosus.
comosa, Dum. l. c. = S. nanus ?
fluitans, Dum. l. c. = S. fluitans.
multiculmis, Dum. l. c. = E. multicaulis.
ovata, Dum. l. c. = E. ovata.
palustris, Dum. l. c. = E. palustris.
uniglumis, Dum. l. c. = E. palustris.

CLAVULIUM, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 407 = **Crotalaria**, Dill. (Legumin.).
mucronatum, G. Don, Gen. Syst. ii. 142 = Crot. mucronata.
pedunculatum, Steud. Nom. ed. II. i. 378 = seq.
pedunculosum, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 407 = Crot. laburnifolia.

CLAVULUM, G. Don, Gen. Syst. ii. 96, 142 (1832) = praec.

CLAYTONA, Linn. Hort. Ups. 52 (1748) = seq.

CLAYTONIA, Gronov. ex Linn. Gen. ed. I. 339 (1737). *PORTULACEAE*, Benth. & Hook. f. i. 158.
BELIA, Steller, ex S. G. Gmel. Fl. Sibir. iv. 88 (1769).
acutifolia, Ledeb. Fl. Alt. i. 253; Ic. t. 272 = arctica.
acutifolia, *Pall. ex Willd. in Roem. & Schult. Syst.* v. 436.—Sibir. or.
cautiflora, Sweet, Hort. Brit. ed. II. 220 = virginica.
alsinoides, Sims, Bot. Mag. t. 1309 = sibirica.
ambigua, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 365.—Am. bor. occ.
aquatica, Nutt. in Torr. & Gray, Fl. N. Am. i. 201 = Chamissonis.
arctica, *Adam, in Mém. Soc. Nat. Mosc.* v. (1817) 94.—As. et Am. arct.
assarifolia, Bong. Veg. Sitcha. 136 = sibirica ?
australasica, *Hook. f. in Hook. Ic. Pl.* t. 293.—Austral.
balonnensis, F. Muell. Census, 27 = Calandrinia balonnensis.
bifida, Willd. ex Roem. & Schult. Syst. v. 434 = sibirica.
brevipedata, F. Muell. Census, 27 = Calandrinia brevipedata.
bulbifera, *A. Gray, in Proc. Am. Acad.* xii. (1877) 54.—Am. bor. occ.
californica, *Herb. Petrop. ex Loud. Hort. Brit. Suppl.* ii. 621.—Calif.
calyptrata, F. Muell. Fragm. iii. 89 = Calandrinia calyptrata.
caroliniana, *Michx. Fl. Bor. Am.* i. 160.—Am. bor.
Chamissoi, Ledeb. ex Spreng. Syst. i. 790 = sarmentosa.
Chamissonis, *Eschscholtz, ex Cham. in Linnaea*, vi. (1831) 562.—Am. bor. occ.
composita, F. Muell. Census, 27 = Calandrinia composita.
cordifolia, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 365.—Am. bor. occ.
corrigioloides, F. Muell. Census, 27 = Calandrinia corrigioloides.
cubensis, Bonpl. in Ann. Mus. Par. vii. (1806) 82. t. 6 = perfoliata.
dichotoma, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 202.—Am. bor. occ.
diffusa, *Nutt. l. c.*—Calif.
Eschscholtzii, *Cham. in Linnaea*, vi. (1831) 561.—Am. arct.
exigua, *Torr. & Gray, Fl. N. Am.* i. 200.—Calif.
filicaulis, Dougl. in Hook. Fl. Bor. Am. i. 224. t. 72 = parvifolia.
flagellaris, Bong. Veg. Sitcha, 136 = Chamissonis.
gracilis, F. Muell. Census, 27 = Calandrinia gracilis.
grandiflora, Sweet, Brit. Flow. Gard. t. 216 = virginica.
granulifera, F. Muell. Census, 27 = Calandrinia granulifera.
gypsophiloides, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 33 = perfoliata.
Joanneana, Roem. & Schult. Syst. v. 434 = arctica.
lanceolata, Hook. & Arn. Bot. Beech. Voy. 123, 344 = sarmentosa.
lanceolata, Pursh, Fl. Am. Sept. i. 175. t. 3 = caroliniana.
Lehmanni, F. Muell. Census, 27 = Calandrinia Lehmanni.
linearis, *Dougl. in Hook. Fl. Bor. Am.* i. 224. t. 71.—Am. bor. occ.
liniflora, F. Muell. Census, 27 = Calandrinia liniflora.
media, Link, Handb. ii. 44 = virginica.
megarrhiza, *Parry, ex S. Wats. Bibliog. Ind. N. Am. Bot.* i. 118.—Am. bor. occ.
nemorosa, Willd. ex Roem. & Schult. Syst. v. 436 (= *Irlbachia Bonplandiana*).—N. Granat.
nevadensis, *S. Wats. Bot. Calif.* i. 77.—Calif.

CLAYTONIA :—
parviflora, Dougl. ex Hook. Fl. Bor. Am. i. 225. t. 73 = perfoliata.
parvifolia, *Moç. ex DC. Prod.* iii. 361.—Am. bor. occ.
perfoliata, *Donn, Ind. Hort. Cant.* 25, *ex Willd. Sp. Pl.* i. 1186.—Am. bor.
Pickeringi, F. Muell. Census, 27 = Calandrinia Pickeringii.
pleiopetala, F. Muell. l. c. = Calandrinia pleiopetala.
pogonophora, F. Muell. l. c. = Calandrinia pogonophora.
polyandra, F. Muell. l. c. = Calandrinia polyandra.
polypetala, F. Muell. l. c. = Calandrinia polypetala.
Portulacaria, Linn. Mant. ii. 211 = Portulacaria afra.
ptychosperma, F. Muell. Census, 27 = Calandrinia ptychosperma.
pumila, F. Muell. l. c. = Calandrinia pumila.
quadrivalvis, F. Muell. l. c. = Calandrinia quadrivalvis.
sarmentosa, *C. A. Mey. in Nouv. Mém. Soc. Nat. Mosc.* vii. (1829) 137. t. 3.—As. et Am. arct.
sibirica, *Linn. Sp. Pl.* 204.—As. bor.; Am. bor. occ.
sibirica, Pall. ex Steud. Nom. ed. II. i. 378 = arctica.
Simsii, Sweet, Brit. Flow. Gard. sub t. 216 = virginica.
spathulaefolia, Salisb. Parad. Lond. t. 71 = caroliniana.
spathulata, Dougl. in Hook. Fl. Bor. Am. i. 226. t. 74 = perfoliata.
spathulata, Hook. & Arn. Bot. Beech. Voy. 344 = exigua.
spatulata, Eaton, Man. ed. IV. 263 = caroliniana.
spergularina, F. Muell. Census, 27 = Calandrinia spergularina.
stolonifera, C. A. Mey. in Nouv. Mém. Soc. Nat. Mosc. vii. (1829) 139 = Chamissonis.
strophiolata, F. Muell. Fragm. xi. 82 = Calandrinia strophiolata.
tenuifolia, Torr. & Gray, Fl. N. Am. i. 201 = exigua.
triphylla, *S. Wats. in Proc. Am. Acad.* x. (1875) 345.—Am. bor. occ.
tuberosa, Moç. & Sesse, ex DC. Prod. iii. 357 = Talinum napiforme.
tuberosa, *Pall. ex Willd. in Roem. & Schult. Syst.* v. 436.—Sibir. or.
umbellata, *S. Wats. in Bot. King, Exp.* 43. t. 6.—Am. bor. occ.
unalaschkensis, Fisch. Hort. Gorenk. ed. II. (1812) 62; et in Roem. & Schult. Syst. v. 434 = sibirica.
uniflora, F. Muell. Census, 27 = Calandrinia uniflora.
Vancouverii, *A. Murr. Oreg. Circ.* 1.—Am. bor. occ.
Vestiana, Fisch. ex G. Don, Gen. Syst. iii. 82 = sarmentosa.
virginica, *Linn. Sp. Pl.* 204.—Am. bor.
volubilis, F. Muell. Census, 27 = Calandrinia volubilis.

CLEACHNE, Roland. ex Steud. Nom. ed. II. i. 378 (1840) = **Paspalum**, Linn. (Gramin.).

CLEANTHE, Salisb. in Trans. Hort. Soc. i. (1812) 312. *IRIDEAE*, Benth. & Hook. f. iii. 701.
bicolor, *Salisb. l. c.*—Afr. austr.
melaleuca, Baker, in Journ. Linn. Soc. xvi. (1877) 112 = bicolor.

CLEANTHES, D. Don, in Trans. Linn. Soc. xvi. (1830) 194 = **Trixis**, P. Br. (Compos.).
brasiliensis, D. Don, l. c. 195 = T. brasiliensis.
hieracioides, D. Don, l. c. 197 = T. ochroleuca.
othonnoides, Griseb. in Goett. Abh. xxiv. (1879) 217 = T. brasiliensis.

CLEGHORNIA, Wight, Ic. tt. 1310, 1312 (1850) = **Baissea**, DC. (Apocyn.).
acuminata, Wight, l. c. t. 1310 = B. acuminata.
cymosa, Wight, l. c. t. 1312 = B. acuminata.

CLEIDION, Blume, Bijdr. 612 (1825). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 320.
LASIOSTYLES, Presl, Bot. Bemerk. 149 (1844).
PSILOSTACHYS, Turcz. in Bull. Soc. Nat. Mosc. xvi. (1843) 58.
REDIA, Casar. Nov. Stirp. Bras. Dec. 51 (1842).
TETRAGLOSSA, Bedd. in Madr. Journ. Sc. Ser. II. xxii. (1861) 70.
castaneaefolium, *Muell. Arg. in Linnaea*, xxxiv. (1861–66) 184.—Peruv.
claoxyloides, *Muell. Arg. l. c.*—N. Caled.

CLEIDION :—
coriaceum, Baill. Adansonia, ii. (1861-62) 218 = Macaranga coriacea.
gabonicum, *Baill. l. c.* xi. (1874) 129.—Afr. trop.
javanicum, *Blume, Bijdr.* 613.—Ind. or.; Malaya.
macrophyllum, Baill. Adansonia, ii. (1861-62) 219 = Vieillardii.
nicaraguense, *Hemsl. Biol. Centr. Am. Bot.* iii. 130.—Nicaragua.
nitidum, *Thw. ex Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 245.—Ins. Andaman.
populifolium, *Zipp. ex Span. in Linnaea*, xv. (1841) 349, nomen.—Malaya.
spathulatum, *Baill. Adansonia*, ii. (1861-62) 221.—N. Caled.
tricoccum, *Baill. l. c.* iv. (1863-64) 370.—Bras.
ulmifolium, *Muell. Arg. in Flora*, xlvii. (1864) 481.—Ins. Loo-Choo.
verticillatum, *Baill. Adansonia*, ii. (1861-62) 221.—N. Caled.
Vieillardii, *Baill. l. c.* 220.—N. Caled.

CLEIEMERA, Rafin. Fl. Tellur. iv. 77 (1836) = **Ipomoea**, Linn. (Convolvulac.).
cuspidata, Rafin. l. c. 78 = Convolvulus Meyeri.
guinensis, Rafin. l. c. = I. capitata.
hederacea, Rafin. l. c. 77 = I. Nil.
hirsuta, Rafin. l. c. 78 = I. hederacea.
indica, Rafin. l. c. (= *Pharbitis violacea*).—As. trop.?
peruviana, Rafin. l. c. = I. longicuspis.

CLEISOCRATERA, Korth. Verh. Nat. Geschied. 256. t. 62 (1842). *RUBIACEAE*, Benth. & Hook. f. ii. 123.
elegans, Korth. l. c. 257. t. 62.—Borneo.

CLEISOSTOMA, Blume, Bijdr. 362 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 580.
Echinoglossum, Reichb. Nom. 54 (1841).
Echioglossum, Blume, Bijdr. 364. t. 28 (1825).
Pomatocalpa, Breda, Orch. Kuhl & Hass. (1827).
Synptera, Llanos, Fragm. Pl. Filip. 98 (1851).
acaule, Lindl. Gen. et Sp. Orch. 227 (= Saccolabii sp.).—Zeylan.
amabile, *Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind.* v. (1853) 493.—Malaya.
appendiculatum, *Benth. & Hook. f. Gen.* iii. 577.—Ind. or.
Armitii, F. Muell. Fragm. ix. 49 = Sarcochilus Armitii.
Beckleri, *F. Muell. ex Benth. Fl. Austral.* vi. 296.—Austral.
bicolor, *Lindl. & Paxt. Flow. Gard.* ii. (1851-52) 99.—Manilla.
bifidum, *Teijsm. & Binn. Cat. Hort. Bog.* 49, nomen.—Java.
brevilabre, *F. Muell. Fragm.* xi. 87.—Austral.
callosum, *Blume, Bijdr.* 364.—Java.
carinum, *Hance, in Journ. Bot.* xx. (1882) 359.—China.
complanatum, *Lindl.* (*Cf. Gen. et Sp. Orch.* 227. n. 8); *Steud. Nom.* ed. II. i. 378.—Ind. or.
crassifolium, *Lindl. & Paxt. Flow. Gard.* iii. (1852-53) t. 99, 126.—Burma?
cryptochilum, *F. Muell. in Wing, South Sc. Record*, N. S. i. (1885).—N. Guinea.
Cumingii, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 218.—Hab.?
Dawsonianum, Reichb. f. in Gard. Chron. (1868) 815 = Trichoglottis Dawsoniana.
dealbatum, *Lindl. Bot. Reg.* (1843) *Misc.* 5.—Manilla.
decipiens, *Lindl. l. c.* (1844) *Misc.* 11.—Zeylan.
discolor, *Lindl. l. c.* (1845) *Misc.* 59.—Ind. or.
emarginatum, *Blume, ex Teijsm. & Binn. Cat. Hort. Bog.* 49, nomen.—Java.
erectum, *Fitzg. Austral. Orch.* i. tv.—Austral.
expansum, *Reichb. f. Otia Bot. Hamb.* 52.—Ins. Mindanao.
Fordii, *Hance, in Journ. Bot.* xiv. (1876) 45.—Ins. Hongkong.
formosanum, *Hance, l. c.* xxii. (1884) 364.—China.
fuscum, Lindl. in Journ. Hort. Soc. v. (1850) 80 = latifolium.
galeatum, Thw. Enum. Pl. Zeyl. 305 = maculosum, *Lindl.*

CLEISOSTOMA :—
Guiberti, *Linden & Reichb. in Bot. Zeit.* xx. (1862) 375; *in Hamb. Gartenz.* xviii. (1862) 529.—Hab.?
ionosmum, *Lindl. Bot. Reg.* (1847) t. 41.—Manilla.
Keffordii, *F. M. Bailey, in Rep. Queensl. Acclim. Soc.* (Apr. 1884) 9.—Austral.
lanatum, *Lindl. in Journ. Hort. Soc.* iv. (1849) 164.—Burma.
latifolium, *Lindl. Bot. Reg.* (1840) *Misc.* 60.—Malaya.
longifolium, *Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind.* v. (1853) 494.—Java.
loratum, *Reichb. f. in Flora*, lv. (1872) 273.—Reg. Himal.
Macphersoni, *F. Muell. ex Benth. Fl. Austral.* vi. 297.—Austral.
macrostachyum, *Teijsm. & Binn. Cat. Hort. Bog.* 49.—Java.
maculosum, *Blume, ex Teijsm. & Binn. Cat. Hort. Bog.* 49, nomen.—Java.
maculosum, *Lindl. Gen. et Sp. Orch.* 227.—Zeylan.
maculosum, Thw. Enum. Pl. Zeyl. 304 = decipiens.
Mannii, *Reichb. f. in Flora*, lv. (1872) 274.—Reg. Himal.
roscum, *Lindl. Bot. Reg.* (1838) *Misc.* 80.—Manilla.
sagittatum, *Blume, Bijdr.* 363.—Java.
spathulatum, *Blume, l. c.* 364.—Java.
spicatum, *Lindl. Bot. Reg.* (1847) sub t. 32.—Malaya.
suaveolens, *Blume, Bijdr.* 363.—Java.
subulatum, *Blume, l. c.*—Java.
subviolaceum, *Reichb. f. in Bonplandia*, x. (1862) 335.—Ins. Philipp.
teretifolium, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 20.—Sumatra.
Thwaitesianum, Trim. in Journ. Bot. xxiii. (1885) 244 = decipiens.
tridentatum, *Lindl. Bot. Reg.* (1838) *Misc.* 33.—Austral.
undulatum, *Reichb. f. in Flora*, lv. (1872) 274.—Reg. Himal.
virginale, *Hance, in Journ. Bot.* xv. (1877) 38.—Ins. Hongkong.
vitellinum, *Reichb. f. in Linnaea*, xli. (1877) 76.—Ins. Philipp.
Wendlandorum, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 219.—Reg. Himal.; Burma.

CLEISOSTOMA, Rafin. Fl. Tellur. iv. 80 (1836) = **Convolvulus**, Tourn.
villosum, Rafin. l. c. = Conv. aggregatus.

CLEISSOCRATERA, Miq. Fl. Ind. Bat. ii. 383 (1856) = **Cleisocratera**, Korth. (Rubiac.).

CLEISTACHNE, Benth. in Journ. Linn. Soc. xix. (1881) 61. *GRAMINEAE*, Benth. & Hook. f. iii. 1120.
sorghoides, *Benth. in Hook. Ic. Pl.* t. 1379.—Afr. trop.; Ind. or.

CLEISTANTHIUM, Kunze, in Bot. Zeit. ix. (1851) 350 = **Gerbera**, Gronov. (Compos.).
nepalense, Kunze, l. c. = G. Kunzeana.

CLEISTANTHUS, Hook. f. ex Planch. in Hook. Ic. Pl. t. 779 (1848). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 268.
Lebidiera, Baill. Etud. Gén. Euph. Atlas, 50. t. 27 (1858).
Lebidieropsis, Muell. Arg. in Linnaea, xxxii. (1863) 79.
Leiopyxis, Miq. Fl. Ind. Bat. Suppl. 445 (1860).
Nanopetalum, Hassk. in Versl. Kon. Akad. Wetensch iv. (1855) 140.
acuminatus, *Muell. Arg. in DC. Prod.* xv. II. 508.—Zeylan.
angolensis, *Welw. ex Muell. Arg. in Journ. Bot.* ii. (1864) 339.—Afr. occ.
apodus, *Benth. Fl. Austral.* vi. 122.—Austral.
Blancoi, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 315.—Ins. Philipp.
Boivinianus, *Muell. Arg. in DC. Prod.* xv. II. 505.—Madag.
chartaceus, *Muell. Arg. l. c.* 507.—Reg. Himal.; Malaya.
collinus, *Benth. & Hook. f. Gen.* iii. 268.—Ind. or.

CLEISTANTHUS :—
Cunninghamii, *Muell. Arg. in DC. Prod.* xv. II. 506.—Austral.
Dallachyanus, *Baill. ex Benth. Fl. Austral.* vi. 122.—Austral.
diversifolius, *Muell. Arg. in DC. Prod.* xv. II. 509.—Madag.
ferrugineus, *Muell. Arg. in DC. Prod.* xv. II. 507.—Zeylan.
ferrugineus, F. Vill. in Blanco, Fl. Philipp. ed. III. Nov. App. 187. t. 353 = Blancoi.
lanceolatus, *Muell. Arg. l. c.*—Java.
malabaricus, *Muell. Arg. l. c.* 508.—Ind. or.
monoicus, *Muell. Arg. l. c.*—China.
myrianthus, *Kurz, For. Fl. Br. Burm.* ii. 370.—Burma; Malaya.
oblongifolius, Muell. Arg. in DC. Prod. xv. II. 506, pro parte = chartaceus.
pallidus, *Muell. Arg. l. c.* 508.—Zeylan.
patulus, *Muell. Arg. l. c.* 505.—Ind. or.
polystachyus, *Hook. f. ex Planch. in Hook. Ic. Pl.* t. 779.—Afr. trop.
robustus, *Muell. Arg. in DC. Prod.* xv. II. 504.—Zeylan.
semiopacus, *F. Muell. ex Benth. Fl. Austral.* vi. 123.—Austral.
stenophyllus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 242.—Burma.
stipitatus, *Muell. Arg. in DC. Prod.* xv. II. 506.—N. Caled.
stipularis, *Muell. Arg. l. c.* 509.—Ind. or.
subglaucus, Thw. ex Trimen, Cat. Pl. Ceyl. 78 = pallidus.
sumatranus, *Muell. Arg. in DC. Prod.* xv. II. 504.—Sumatra.
tomentosus, *Hance, in Journ. Bot.* xv. (1877) 337.—China.

CLEISTES, Rich. in Mém. Mus. Par. iv. (1818) 31 = **Pogonia**, Juss. (Orchid.).
Liliastrum, Reichb. f. in Linnaea, xxii. (1849) 816 = P. Liliastrum.
lutea, Lindl. Gen. et Sp. Orch. 409.—Guiana.
Miersii, Gardn. in Hook. Ic. Pl. sub t. 473-474 = P. Miersii.
montana, Gardn. l. c. = P. montana.
paludosa, Reichb. f. in Linnaea, xxii. (1849) 816 = P. paludosa.
parviflora, Lindl. Gen. et Sp. Orch. 410.—Guiana.
rosea, Lindl. l. c. = P. rosea.
speciosa, Gardn. in Hook. Ic. Pl. t. 473-474 = P. speciosa.

CLEISTOCACTUS, Lem. Illustr. Hortic. viii. (1861) Misc. 35 = **Cereus**, Mill. (Cact.).
Baumannii, Lem. l. c. = Cereus Baumannii.
colubrinus, Lem. l. c. = Cereus colubrinus.
rhodacanthus, Lem. l. c. = Echinopsis rhodacantha.

CLEISTOCALYX, Blume, Mus. Bot. Lugd. Bat. i. 84. t. 56 (1849) = **Eugenia**, Linn. (Myrtac.).
nervosus, Blume, l. c. 85 = E. nervosa.
nitidus, Blume, l. c. 84.—Java.

CLEISTOCALYX, Steud. Syn. Pl. Cyp. 151 (1855) = **Rynchospora**, Vahl (Cyperac.).
junciformis, Steud. l. c. = R. oligantha.

CLEISTOCHLAMYS, Oliver, in Journ. Linn. Soc. ix. (1867) 175. *ANONACEAE*, Benth. & Hook. f. i. 956.
Kirkii, *Oliver, l. c.*—Afr. trop.

CLEITHRIA, Steud. Nom. ed. II. ii. 746 (1841) = seq.

CLEITRIA, Schrad. Ind. Sem. Hort. Gotting. (1831); ex Linnaea, viii. (1832) Litt. 23 = **Venidium**, Less. Compos.).
hirta, Schrad. ll. cc. = V. Schraderi.

CLELIA, Casar. Nov. Stirp. Bras. Dec. 83 (1842) = **Calliandra**, Benth. (Legumin.).
ornata, Casar. l. c. 84 = Call. Harrisii.

CLEMANTHUS, Klotzsch, in Peters, Reise Mossamb. Bot. 143 (1861) = **Modecca**, Lam. Passiflor.).
senensis, Klotzsch, l. c. = M. senensis.

CLEMATIS, Dill. ex Linn. Gen. ed. I. 163 (1737). *RANUNCULACEAE*, Benth. & Hook. f. i. 3.
ATRAGENA, Linn. Syst. ed. I (1735); et Diss. Dass. 21 (1747); in Amoen. Acad. i. 132 (1749) (Atragene).
CHEIROPSIS, Spach, Hist. Vég. Phan. vii. 260 (1839).
CLEMATITIS, [Tourn.] Linn. Syst. ed. I (1735).
MECLATIS, Spach, Hist. Vég. Phan. vii. 272 (1839).
MURALTA, Adans. Fam. ii. 460 (1763).
SIEBOLDIA, Hoffmgg. ex Heynh. Nom. ii. 665 (1846).
TRIGULA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 4; ed. II. 67 (1827).
VIORNA, Reichb. Handb. 277 (1837).
VITICELLA, Dill. ex Moench, Meth. 296 (1794).
acapulcensis, *Hook. & Arn. Bot. Beech. Voy.* 410.—Mexic.
acerifolia, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) I. 2.—China.
acuminata, *DC. Syst.* i. 148.—Reg. Himal.
acutangula, *Hook. f. & Thoms. Fl. Ind.* 5.—Reg. Himal.
aethusiaefolia, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 181.—Mongolia.
affinis, *St. Hil. Fl. Bras. Mer.* i. 3.—Bras.
affinis, *Wight, Illustr.* i. 5.—Ind. or.
afoliata, *F. Buch. in Trans. N. Z. Inst.* iii. (1871) 211.—N. Zel.
ajanensis, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 176.—As. bor.
albida, Klotzsch, in Bot. Ergeb. Waldem. Reise 131 = orientalis.
alpina, *Mill. Gard. Dict.* ed. VIII. n. 9.—Europ.; Am. bor.
americana, Mill. l. c. n. 14 = dioica.
amplexicaulis, Edgew. in Trans. Linn. Soc. xx. (1851) 24 = connata.
anemoniflora, D. Don, Prod. Fl. Nep. 192 = montana.
anemonoides, O. Kuntze, in Verh. Bot. Ver. Brand. xxvi. (1885) 149 = florida.
anethifolia, *Hook. Ic. Pl.* t. 78.—Madag.
angustifolia, Jacq. Enum. Vindob. 310 = C. Flammula.
aphylla, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 146.—N. Zel.
apiculata, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 4.—Reg. Himal.
apiifolia, *DC. Syst.* i. 149.—Japon.
aquifolia, Steud. Nom. ed. II. i. 378, sphalm. = apiifolia.
aristata, *R. Br. in Bot. Reg.* t. 238.—Austral.
Armandi, *Franch. in Nouv. Arch. Mus. Par.* Sér. II. viii. (1885) 184.—As. or.
aromatica, *Lenné & C. Koch, in App. Ind. Sem. Hort. Berol.* (1855) 15; *ex Walp. Ann.* iv. 4.—Hab.?
asplenifolia, *Schrenk, Enum. Pl. Nov. Alt.* 68.—Soongaria; Afghanist.
azurea, *Hort. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 272 [bis].—Tauria (cult.).
azurea, Siebold, ex Steud. Nom. ed. II. i. 378 = florida.
baccata, Pers. Syn. ii. 99 = C. Viticella.
Baldwini, *Torr. & Gray, Fl. N. Am.* i. 8.—Am. bor.
balearica, Rich. in Journ. Phys. (Févr. 1779) 127 = cirrhosa.
bannatica, *Schur, in Verh. Siebenb. Ver. Naturw.* iv. (1853) 7.—Europ.
barbellata, *Edgew. in Trans. Linn. Soc.* xx. (1851) 25.—Reg. Himal.
Bergeroni, *A. Lavall. Clemat.* 35. t. 10.—Hab.?
bicolor, Steud. Nom. ed. II. i. 378 = florida.
Bigelovi, *Torr. in Pacif. Rail. Rep.* iv. 61.—Am. bor. occ.
biloba, Steud. Nom. ed. II. i. 379, vide *triloba*, St. Hil.
biternata, *DC. Syst.* i. 149.—Japon.; China.
blanda, Hook. Journ. Bot. i. (1834) 241 = aristata.
Bojeri, *Hook. Ic. Pl.* t. 10.—Madag.
bonariensis, *Juss. ex DC. Syst.* i. 145.—Reg. Argent.
brachiata, Ker-Gawl. in Bot. Reg. t. 97 = Kerrii.
brachiata, *Thunb. Prod. Pl. Cap.* 94.—Afr. austr.
brachyura, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 221.—Corea.
bracteata, DC. Syst. i. 142 = virginiana.
bracteata, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 43.—Ind. or.
bracteosa, Banks, ex Steud. Nom. ed. I. 204, 205 = recta.
brasiliana, *DC. Syst.* i. 143.—Bras.
brevicaudata, *DC. l. c.* 138.—China.

CLEMATIS:—
Bucamara, Buch.-Ham. ex DC. Syst. i. 140 = Buchaniana.
Buchanani, D. Don, Prod. Fl. Nep. 191 = seq.
Buchananiana, Wall. Cat. n. 4677 = seq.
Buchaniana, *DC. Syst.* i. 140.—Reg. Himal.
Cadmia, *Buch.-Ham. ex Wall. Cat.* n. 4669.—Reg. Himal.; Burma.
caerulea, Lindl. Bot. Reg. t. 1955 = patens.
caesariata, *Hance, in Journ. Bot.* viii. (1870) 71.—China.
caespitosa, Scop. Fl. Carn. ed. II. i. 389 = C. Flammula.
calycina, [Soland. in] Ait. Hort. Kew. ed. I. ii. 259 = cirrhosa.
campaniflora, *Brot. Fl. Lusit.* ii. 359.—Lusitan.
campaniflora, Lodd. ex Steud. Nom. ed. II. i. 379 = parviflora.
campestris, *St. Hil. Fl. Bras. Mer.* i. 4.—Bras.
cana, Wall. Cat. n. 4672 = Gouriana.
canadensis, Mill. Gard. Dict. ed. VIII. n. 5 = virginiana.
canaliculata, Lag. Gen. et Sp. Nov. 17 = C. Flammula.
capensis, Poir. Encyc. Suppl. ii. 296 = Anemone capensis.
caracassana, H. B. & K. ex DC. Syst. i. 141 = caripensis.
caripensis, *H. B. & K. Nov. Gen. et Sp.* v. 36.—Mexic.
Catesbyana, *Pursh, Fl. Am. Sept.* ii. 736.—Am. bor.
caudata, Hook. ex Hemsl. Biol. Centr. Am. Bot. i. 2 = Drummondii.
Chandleri, Hort. ex Lavall. Clemat. 40 = eriostemon.
chinensis, Osb. Iter, ed. Angl. i. 393 = recta.
chinensis, *Retz. Obs.* ii. 18. t. 2.—China.
chlorantha, Lindl. Bot. Reg. t. 1234 = grandiflora.
chrysocarpa, *Welw. ex Oliver, Fl. Trop. Afr.* i. 5.—Afr. trop.
cirrhosa, *Linn. Sp. Pl.* 544.—Reg. Mediterr.
clitorioides, DC. Syst. i. 158 = aristata.
coccinea, Engelm. ex A. Gray, Pl. Wright. ii. 7 = C. Viorna.
cocculifolia, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iv. (1840) 260.—N. Zel.
cognata, Steud. in Lehm. Pl. Preiss. i. 263 = aristata.
Colensoi, *Hook. f. Fl. N. Zeland.* i. 6.—N. Zel.
coloradoënsis, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 448 = Pitcheri.
columbiana, Torr. & Gray, Fl. N. Am. i. 11 = verticillaris.
commutata, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 128.—Afr. trop.
comosa, DC. Syst. i. 157 = triloba?
connata, *DC. Prod.* i. 4.—Reg. Himal.
contorta ×, Carr. in Rev. Hortic. (1879) 380.
cordata, Pursh, Fl. Am. Sept. ii. 384 = virginiana.
cordata, *Royle, Illustr. Bot. Himal.* 51.—Ind. or.
cordata, Sims, Bot. Mag. t. 1816 = crispa.
cordifolia, DC. Syst. i. 142 = virginiana.
coriacea, DC. l. c. 146 = aristata.
coriacea, *Korth. in Nederl. Kruidk. Arch.* i. (1848) 208.—Java.
corymbosa, Poir. Encyc. Suppl. ii. 297 = recta.
crassifolia, *Benth. Fl. Hongk.* 7.—China.
crenata, Jord. in Billot, Annot. Fl. Fr. Allem. 12 = C. Vitalba.
crispa, Lam. Encyc. ii. 44 = cylindrica.
crispa, *Linn. Sp. Pl.* 543.—Am. bor.
crispa, Thunb. Fl. Jap. 239 = paniculata.
cylindrica, Sims, Bot. Mag. t. 1160 = crispa.
dahurica, DC. Prod. i. 7 = daurica.
dasyoneura, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 123.—Borneo.
daurica, *Pers. Syn.* ii. 99.—As. temp.
Davidiana, Decne. ex Verlot, in Rev. Hortic. (1867) 90 = heracleaefolia.
davurica, Ledeb. Fl. Ross. i. 4 = daurica.
denticulata, Vell. Fl. Flum. 240; v. t. 134 = Hilarii.
depauperata, *Colenso, in Trans. N. Z. Inst.* i. (1868) 248, nomen.—N. Zel.
dioica, *Linn. Syst.* ed. X. 1084.—Jamaica.
dioica, Lour. Fl. Cochinch. 344 = Loureiriana.
dioica, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 273 = caripensis.
discolor, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 330.—Bras.
discolor, Steud. in Lehm. Pl. Preiss. i. 262 = aristata.
dissecta, *Baker, in Journ. Linn. Soc.* xx. (1883) 87.—Madag.

CLEMATIS:—
distorta, *Lavall. Clemat.* 37. t. 11.—Hab.?
divaricata, Jacq. Eclog. Pl. i. 51. t. 33 = crispa.
diversifolia, *DC. Syst.* i. 155.—Hab.?
diversifolia, Gilib. Exercit. i. 383 = C. Flammula.
dominica, *Lam. Encyc.* ii. 45.—Ind. occ.
Douglasii, *Hook. Fl. Bor. Am.* i. 1. t. 1.—Am. bor. occ.
Drummondii, *Torr. & Gray, Fl. N. Am.* i. 9.—Am. bor.
dumosa, Salisb. Prod. 371 = C. Vitalba.
edentata, *Baker, in Journ. Linn. Soc.* xxi. (1884) 318.—Madag.
elliptica, Endl. in Enum. Pl. Hueg. 1 = aristata.
elongata, Tratt. Archiv, t. 149 = integrifolia.
erecta, Linn. Syst. ed. XII. 377; All. Fl. Pedem. i. 296 = recta.
eriopoda, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 223.—Japon.
eriostemon, *Decne. in Rev. Hortic.* Sér. IV. i. (1852) 341.—Hab.?
excelsior ×, Van Houtte, in Fl. des Serres, Sér. II. x. (1874) 24. t. 2054-55.
Fawcettii, *F. Muell. Fragm.* x. 1.—Austral.
filifera, Benth. Pl. Hartw. 285 = reticulata.
Flammula, *Linn. Sp. Pl.* 544.—Reg. Mediterr.
Flammulastrum, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 153.—Cuba.
flava, DC. Syst. i. 135 = orientalis.
flavescens, *Rafin. Fl. Ludov.* 81.—Am. bor.
floribunda, *Kurz, in Journ. Bot.* v. (1867) 240.—Ind. or.
floribunda, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 9.—N. Granat.
florida, *Thunb. Fl. Jap.* 240.—Japon.
fluminensis, *Vell. Fl. Flum.* 240; v. t. 133.—Bras.
foetida, *Raoul, Choix Pl. Nouv. Zél.* 23.—N. Zel.
Forsteri, J. F. Gmel. Syst. 873 = hexasepala.
Fortunei, T. Moore, in Gard. Chron. (1863) 460, 676 = Williamsii.
fragrans, Salisb. Prod. 371 = virginiana.
fragrans, Tenore, Prod. Fl. Nap. p. xxxii = C. Flammula.
francofurtensis, Rinz. ex C. Koch, Dendrol. i. 431 = C. Viticella.
Fremonti, *S. Wats. in Proc. Am. Acad.* x. (1875) 339.—Am. bor. occ.
fruticosa, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 180.—China.
fulva, *Zoll. & Moritzi, Syst. Verz. Zoll.* 35.—Java.
fusca, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 60.—China; Japon.
Gebleriana, Bong. in Bull. Sc. Acad. Pétersb. viii. (1841) 338 = soongarica.
gentianoides, *DC. Syst.* i. 159.—Austral.
Gilbertiana, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 273 [bis] = aristata.
glabra, *DC. Syst.* i. 143.—Ins. S. Doming.
glandulosa, Blume, Bijdr. 1 = smilacifolia.
glauca, Claus, in Goebel, Reise, ii. 247 = orientalis.
glauca, Willd. Berl. Baumz. 65 = orientalis.
glaucescens, Presl, in Mus. Senckenb. ii. (1837) 268 = Thunbergii.
globosa, Royle, Illustr. Bot. Himal. 51 = orientalis.
glycinoides, *DC. Syst.* i. 145.—Austral.
Goudotiana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 10.—N. Granat.
Gouriana, *Roxb. Hort. Beng.* 43; *Fl. Ind.* ii. 670.—Ind. or.; Malaya.
gracilis, Edgew. in Trans. Linn. Soc. xx. (1851) 24 = connata.
Grahami, *Benth. Pl. Hartw.* 3.—Mexic.
grandiflora, *DC. Syst.* i. 151.—Afr. trop.
grata, O. Hoffm. ex Baker, in Journ. Linn. Soc. xx. (1883) 88 = Wightiana.
grata, *Wall. Pl. As. Rar.* i. 83. t. 98.—Reg. Himal.; China; Afr. trop.
graveolens, Hook. Bot. Mag. t. 4495 = orientalis.
graveolens, *Lindl. in Journ. Hort. Soc.* i. (1846) 307.—Reg. Himal.
grewiaeflora, *DC. Syst.* i. 140.—Reg. Himal.
grossa, *Benth. Pl. Hartw.* 33.—Mexic.
grossa, *Wall. Cat.* n. 4671.—Burma.
Guadeloupae, Pers. Syn. ii. 99 = americana.
Guascoi ×, Lem. & Versch. in Illustr. Hortic. iv. (1857) 117.
Haenkeana, *Presl, Rel. Haenk.* ii. 69.—Peruv.
hakonensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 263.—Japon.

CLEMATIS:—
Hancockiana, *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) i. 1.—China.
Hartwegi, Hort. ex C. Koch, Dendrol. i. 434 = reticulata.
havanensis, *H. B. & K. ex DC. Syst.* i. 152.—Cuba.
hedysarifolia, *DC. l. c.* 148.—Ind. or.
Hendersonii, Hort. ex Steud. Nom. ed. II. i. 379 = reticulata.
Henryi, *Hort.—Cf. Gard. Chron.* (1878) II. 439.—Hab.?
heracleaefolia, *DC. Syst.* i. 138.—China.
hexapetala, Linn. f. Suppl. 271 = hexasepala.
hexapetala, Pall, Reise, iii. 735 = C. Flammula.
hexasepala, *DC. Syst.* i. 146.—N. Zel.
Hilarii, *Spreng. Syst.* v. 177.—Bras.
hirsuta, Guill. & Perr. Fl. Seneg. Tent. i. 1 = Thunbergii.
hirsutissima, Pursh, Fl. Am. Sept. ii. 385 = Anemone patens.
hispanica, Mill. Gard. Dict. ed. VIII. n. 3 = recta.
holosericea, *Pursh, Fl. Am. Sept.* ii. 384.—Am. bor.
Hookeri, Decne. in Fl. des Serres, Sér. II. xiii. (1880) 282 = heracleaefolia.
Hothae, *Kurz, For. Fl. Brit. Burma*, i. 17.—Burma.
hysudrica, Munro, ex Hook. f. & Thoms. Fl. Ind. 9 = orientalis.
ibarensis, *Baker, in Journ. Linn. Soc.* xviii. (1881) 264.—Madag.
incisodentata, A. Rich. Tent. Fl. Abyss. i. 2 = grata.
inclinata, Scop. Fl. Carn. ed. II. i. 389 = integrifolia.
indica, Heyne, ex Roth, Nov. Pl. Sp. 252 = Gouriana.
indivisa, Steud. in Lehm. Pl. Preiss. i. 262 = aristata.
indivisa, *Willd. Sp. Pl.* ii. 1291.—N. Zel.
insidiosa, *Baill. in Bull. Soc. Linn. Par.* i. (1882) 331.—Madag.
insulensis, Hort. ex C. Koch, Dendrol. i. 433 = crispa.
integra, *Vell. Fl. Flum.* 241 ; v. t. 135.—Bras.
integrifolia, Forst. f. Prod. 42 = indivisa.
integrifolia, *Linn. Sp. Pl.* 544.—Eur. austr. ; As. bor.
intermedia, *Carr. in Rev. Hortic.* (1866) 339.—Hab.?
intricata, *Bunge, in Mém. Sav. Etr. Acad. Pétersb.* vi. (1835) 75.—China.
inversa, *Griff. Notul.* iv. 700.—Ind. or.
ispahanica, Boiss. Diagn. Ser. I. vi. 3 = orientalis.
italica, *Schnizl. Icon.* iii. *Ind.* 2.—Europ.
Jackmanni ×, Van Houtte, in Fl. des Serres, xvi. (1865) 39.
japonica, *Thunb. Fl. Jap.* 240.—Japon.
javana, DC. Syst. i. 152 = Gouriana.
javanica, Steud. Nom. ed. II. i. 379 = praec.
jubata, *Bisch. Del. Sem. Hort. Heidelb.* (1849) 4 ; *ex Walp. Ann.* ii. 4.—Hab.?
Junghuhniana, *De Vriese, in Miq. Pl. Jungh.* 75.—Java.
kamtschatica, *Bong. Verz. Saisang-Nor Pfl.* 10 = Kamtchatka.
Kerrii, *Steud. Nom.* ed. II. i. 379.—Afr. austr.
Kirilowi, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 210.—China.
Kirkii, *Oliver, Fl. Trop. Afr.* i. 5.—Afr. trop.
Kousabotan, *Decne. in Fl. des Serres*, xxiii. (1880) 284.—Japon.
laeta, Salisb. Prod. 370 = cirrhosa.
lanuginosa, *Lindl. & Paxt. Flow. Gard.* iii. (1852-53) 107.—China.
lasiandra, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 213.—Japon.
lasiantha, Fisch. Hort. Gorenk. ed. II. (1812) 47 = C. Flammula.
lasiantha, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 9.—Am. bor. occ.
lathyrifolia, Bess. ex Reichb. Fl. Germ. Excurs. 734 = recta.
Lavallei, *Decne. in Fl. des Serres*, xxiii. (1880) 284.—Japon.
laxiflora, *Baker, in Journ. Linn. Soc.* xxi. (1884) 317.—Madag.
Lechenaultiana, *DC. Syst.* i. 151.—Java ; Ind. or.
leptomera, *Hance, in Journ. Bot.* xviii. (1880) 257.—China.
ligusticifolia, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 9.—Am. bor.
linearifolia, Steud. in Lehm. Pl. Preiss. i. 262 = microphylla.
linearifloba, DC. Syst. i. 155 = crispa.
loasaefolia, DC. l. c. 140 = grewiaeflora?
loasifolia, D. Don, Prod. Fl. Nep. 191 = Buchaniana?

CLEMATIS:—
longecaudata, Ledeb. Fl. Ross. i. 3 = orientalis.
longicauda, *Steud. ex A. Rich. Tent. Fl. Abyss.* i. 2.—Abyssin.
longiloba, *DC. Syst.* i. 136.—Chili.
longipes, *Freyn, in Bremen Abh.* viii. (1880) 5.—Madag.
Loureiriana, *DC. Syst.* i. 144.—China.
lugubris, Salisb. Prod. 371 = C. Viticella.
Luloni, Hort. ex C. Koch, Dendrol. i. 435 = patens.
macropetala, *Ledeb. Ic. Pl. Fl. Ross.* i. 5. t. 11.—Sibiria.
magniflora, *Hoffmgg. Verz. Pfl. Nachtr.* i. 167.—Hab.?
mandschurensis, Hort. ex C. Koch, Dendrol. i. 429 = fusca.
mandschurica, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xv. (1857) 258 = recta.
marata, *Armst. in Trans. N. Z. Inst.* xiii. (1881) 335.—N. Zel.
maritima, Lam. Encyc. ii. 174 = C. Flammula.
maritima, Linn. Sp. Pl. 767 = recta.
Massoniana, DC. Syst. i. 135 = brachiata.
Massoniana, Maxim. Ind. Fl. Pekin. in Prim. Fl. Amur. 468 = Kirilowi.
mauritiana, *Lam. Encyc.* ii. 42.—Ins. Maurit. ; Madag.
Maximowicziana, *Franch. & Sav. Enum. Pl. Jap.* ii. 261.—Japon.
Mechowiana, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 171.—Afr. trop.
medusaea, *Planch. & Linden, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 11.—N. Granat.
mendocina, *Phil. in Anal. Univ. Chil.* (1862) II. 389.—Chili.
Meyeniana, *Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 297.—China.
microcuspis, *Baker, in Journ. Linn. Soc.* xxi. (1884) 317.—Madag.
microphylla, *DC. Syst.* i. 147.—Austral.
millefoliata, *Eichl. in Mart. Fl. Bras.* xiii. i. 150.—Peruv.
minor, *Lour. Fl. Cochinch.* i. 345.—China.
Mociniana, *G. Don, Gen. Syst.* i. 5.—Mexic.
monstrosa, Hort. ex C. Koch, Dendrol. i. 436 = patens.
montana, *Buch.-Ham. ex DC. Syst.* i. 164.—Reg. Himal.
montana, D. Don, Prod. Fl. Nep. 192 = napaulensis.
montevidensis, *Spreng. Syst.* ii. 667.—Reg. Argent.
Munroiana, Wight, Illustr. Ind. Bot. i. 5. t. 1 = smilacifolia.
nannophylla, *Maxim. in Bull. Acad. Pétersb.* xxiii. (1877) 305.—China.
napaulensis, *DC. Syst.* i. 164.—Nepal.
naravelioides, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 119.—Ind. or.
nepalensis, Royle, Illustr. Bot. Himal. 51 = barbellata.
nervata, Benth. Pl. Hartw. 5 = Drummondii.
Noronhiana, *DC. Syst.* i. 151.—Java.
nutans, Crantz, Stirp. Austr. ed. I. fasc. ii. 110 ; ed. II. 127 = integrifolia.
nutans, *Royle, Illustr. Bot. Himal.* 51.—Reg. Himal.
occidentalis, A. Cunn. ex Hook. in Mitch. Journ. Trop. Austral. 368 = microphylla.
occidentalis, *DC. Prod.* i. 10.—Hab.?
ochotensis, *Poir. Encyc. Suppl.* ii. 298.—Sibir. or.
ochroleuca, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 260.—Am. bor.
odora, *Wender.* '*Ind. Sem. Hort. Marb. App.* n. 4.'—Cf. *Steud. Nom.* ed II. i. 380.—Hab.?
odorata, *Wall. ex Steud. Nom.* ed. II. i. 380.—Ind. or.
oligophylla, *Hook. Ic. Pl.* t. 80.—Madag.
Oliveri, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 165.—Abyssin.
oreophila, Hance, in Walp. Ann. ii. 3 = Meyeniana.
orientalis, *Linn. Sp. Pl.* 543.—Reg. Himal. ; Oriens.
orientalis, Seem. ex Steud. Nom. ed. II. i. 380 = C. Flammula.
ovata, *Pursh, Fl. Am. Sept.* ii. 736.—Am. bor.
Orveniae, *Harv. in Harv. & Sond. Fl. Cap.* i. 2.—Afr. trop.
Pallasii, J. F. Gmel. Syst. 873 = C. Flammula.
pallida, *A. Rich. Ess. Fl. Cub.* 10.—Cuba.
panduana, Steud. Nom. ed. II. i. 380, sphalm. = montana.
paniculata, J. F. Gmel. Syst. 873 = indivisa.
paniculata, *Thunb. in Trans. Linn. Soc.* ii. (1794) 337.—Japon.
Parkinsoniana, *Colenso, in Trans. N. Z. Inst.* xii. (1880) 359.—N. Zel.

CLEMATIS:—
parviflora, *A. Cunn. in Ann. Nat. Hist.* Ser. I. iv. (1840) 259.—N. Zel.
parviflora, *DC. in Mém. Soc. Phys. Genèv.* i. (1823) 433.—Hab.?
parviflora, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 9.—Am. bor.
parvifolia, Edgew. in Trans. Linn. Soc. xx. (1846) 25 = orientalis.
parvifolia, Hook. f. & Thoms. Fl. Ind. 9 = graveolens.
parviloba, *Gardn. & Champ. in Hook. Kew Journ.* i. (1849) 241.—China.
patens, *Morr. & Decne. in Bull. Acad. Brux.* iii. (1836) 173.—Japon.
pauciflora, *Nutt. ex Torr. & Gray, Fl. N. Am.* i. 9.—Calif.
pedicellata, Sweet, Hort. Brit. ed. I. 2 ; Loud. Hort. Brit. 1 = cirrhosa.
pensylvanica, *Donn, Hort. Cantab.* ed. IV. 129 ; *Hort. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 273 [bis].—Am. bor.
perulata, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 153.—Bras.
peruviana, *DC. Syst.* i. 141.—Peruv.
Petersiana, Klotzsch, in Peters, Reise Mossamb. Bot. i. 170 = grata.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 1.—Ins. Fiji.
Pierotii, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 1.—Japon.
pilosa, Dulac, Fl. Hautes-Pyr. 211 = C. Vitalba.
pimpinellifolia, *Hook. Ic. Pl.* t. 77.—Madag.
pinnata, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 216.—China.
Pitcheri, Sarg. ex Lavall. Clemat. 60 = Sargenti.
Pitcheri, *Torr. & Gray, Fl. N. Am.* i. 10.—Am. bor. occ.
Plukenetii, DC. Syst. i. 153 = Catesbyana.
Poizati, Hort. ex Lavall. Clemat. 40 = eriostemon.
polycephala, *Bertol. Fl. Guat.* 23.—Guatemala.
polygama, *Jacq. Enum. Pl. Carib.* 24.—Ind. occ.
polymorpha, Viv. Fl. Cors. Sp. Nov. 9 = cirrhosa.
polypetala, DC. Syst. i. 167 = Anemone japonica.
polypetala, *Poir. Encyc. Suppl.* ii. 296.—Japon.
populifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 272.—Venezuela.
pseudoatragene, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 160.—Am. bor.
pseudograndiflora, *O. Kuntze, l. c.* 128.—Afr. trop.
pseudo-orientalis, *O. Kuntze, l. c.* 171.—Persia.
puberula, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 4.—Reg. Himal.
pubescens, *Benth. Pl. Hartw.* 5.—Mexic.
pubescens, Hueg. Enum. Pl. Hueg. 1 = aristata.
pulchella, Pers. Syn. ii. 99 = C. Viticella.
punduana, Wall. Cat. n. 4682 = montana.
Purshii, D. Dietr. Syn. Pl. iii. 345 = virginiana.
quadribracteolata, *Colenso, in Trans. N. Z. Inst.* xiv. (1882) 329.—N. Zel.
recta, *Linn. Sp. Pl.* 544.—Europ. austr.
reticulata, *Walt. Fl. Carol.* 156.—Am. bor.
revoluta, *Hort. Par. ex Steud. Nom.* ed. II. i. 380 = parviflora.
Robertsiana, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 29.—Afghan.
rosea, Hort. ex Lavall. Clemat. 35 = Bergeroni.
rosea, Sm. in Abbot, Nat. Hist. Lepidopt. 201. t. 101 = reticulata.
rubella, Pers. Syn. ii. 100 = C. Flammula.
rubro-violacea ×, Van Houtte, in Fl. des Serres, xvi. (1865) 37 = lanuginosa.
Sargenti, *Lavall. Clemat.* 60. t. 18.—Am. bor.
Savatieri, *Decne. in Fl. des Serres*, xxiii. (1880) 286.—Japon.
saxicola, *Hils. & Boj. ex Baill. Bull. Soc. Linn. Par.* i. (1882) 331, nomen.—Madag.
scabiosaefolia, *DC. Syst.* i. 154.—Ind. or.
scandens, Borkh. Handb. Forstb. ii. 1199 = C. Vitalba.
Schillingii, C. Koch, in App. Ind. Sem. Hort. Berol. (1855) 16 = cylindrica.
Scottii, *Porter, Syn. Fl. Colorado*, 1.—Am. bor. occ.
Seemanni, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 146.—Peruv.
semitriloba, Lag. Gen. et Sp. Nov. 17 = cirrhosa.
sepium, Lam. Fl. Fr. iii. 306 = C. Vitalba.
sericea, *H. B. & K. ex DC. Syst.* i. 144.—Am. austr.
sericea, Michx. Fl. Bor. Am. i. 319 = ochroleuca.

CLEMATIS:—
sibirica, Mill. Gard. Dict. ed. VIII. n. 12 = angustifolia.
Sieboldii, D. Don, ex Steud. Nom. ed. II. i. 380 = florida.
simensis, *Fresen. in Mus. Senckenb.* ii. (1837) 267.—Abyssin.
sinensis, Lour. Fl. Cochinch. 344 = chinensis.
Simsii, Sweet, Hort. Brit. ed. I. 1 = crispa.
smilacifolia, *Wall. in As. Res.* xiii. (1820) 402.—Ind. or.; Malaya.
smilacina, Blume, Bijdr. 1 = smilacifolia.
songarica, Siev. ex Steud. Nom. ed. II. i. 380 = glauca.
songorica, *Bunge, Del. Sem. Hort. Dorp.* (1839) 8.—*Cf. Linnaea*, xiv. (1840) *Litt.* 119.—Sibiria.
Sonnerati, Pers. Syn. ii. 99 = mauritiana.
Standishi, Van Houtte, in Fl. des Serres, xvi. (1865) 39 = patens.
Stanleyi, *Hook. Ic. Pl.* t. 589.—Afr. austr.
stans, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. (1846) II. 177.—Japon.
stenopetala, R. Br. ex DC. Syst. i. 147 = glycinoides.
stenophylla, Fraser, ex Hook. in Mitch. Trop. Austr. 368 = microphylla.
stenosepala, DC. Syst. i. 147 = glycinoides.
stipulata, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 154.—Mexic.
striata, *Rafin. Fl. Ludov.* 82.—Am. bor.
stricta, *Wender. in Ind. Sem. Marb.* (1836).—*Cf. Linnaea*, xii. (1838) *Litt.* 85.—Hab.?
strigillosa, *Baker, in Journ. Linn. Soc.* xviii. (1881) 265.—Madag.
Strobeliana, *Ces. in Atti Soc. Acc. Sc. Nap.* v. (1873) 3. n. 7.—Europ.
Stronachii, *Hance, in Journ. Bot.* xvi. (1878) 103.—China.
suaveolens, Salisb. Prod. 371 = C. Flammula.
subpeltata, Wall. Pl. As. Rar. i. 19. t. 20 = smilacifolia.
substipulata, *O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 147.—Ind. or.
subtriflora, Walp. Rep. i. 8, sphalm. = seq.
subtriloba, *Nees, ex G. Don, Gen. Syst.* i. 7.—Mexic.
subumbellata, *Kurz, For. Fl. Brit. Burma*, i. 16.—Burma.
sulcata, Wall. Cat. n. 4667 = C. Cadmia.
Tatarinowii, *Maxim. in Bull. Acad. Pétersb.* xxii. (1877) 214.—China.
taurica, Bess. ex Nym. Consp. 1 = C. Vitalba.
tenuiflora, DC. Prod. i. 3 = recta.
tenuifolia, Poir. Encyc. Suppl. ii. 298 = Anemone tenuifolia.
tenuifolia, Royle, Illustr. Bot. Himal. 51 = orientalis.
terniflora, DC. Syst. i. 138 = recta.
Tessen, Hassk. Cat. Hort. Bog. Alt. 179 = Noronhiana.
texensis, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 448 = C. Viorna.
thalictroides, *Steud. in Flora*, xxxix. (1856) 407.—Peruv.
Thunbergii, *Steud. Nom.* ed. II. i. 380.—Afr. austr.
tibetana, *O. Kuntze, in Verh. Bot. Ver Brand.* xxvi. (1885) 172.—Tibet.
tortuosa, Wall. Cat. n. 4675 = Buchaniana.
Tournefortii, *DC. Syst.* i. 167.—Oriens.
trifida, *Hook. Ic. Pl.* t. 79.—Madag.
triflora, Vahl, Symb. Bot. iii. 74 = mauritiana.
trifoliata, *Thunb. in Trans. Linn. Soc.* ii. (1794) 337.—Japon.
triloba, *Heyne, ex Roth, Nov. Pl. Sp.* 251.—Ind. or.
triloba, St. Hil. Fl. Bras. Mer. i. 3 = montevidensis.
triloba, Thunb. Prod. Pl. Cap. 94 = Thunbergii.
trinervis, Buch.-Ham. ex DC. Syst. i. 148 = acuminata.
triternata, *DC. l. c.* 150.—Hab.?
tubulosa, Hook. Bot. Mag. t. 6801 = heracleaefolia.
tubulosa, Turcz. in Bull. Soc. Nat. Mosc. (1837) n. VII. 148 = heracleaefolia.
tunbridgensis ×, Hort. ex Lavall. Clemat. 13.
umbraticola, Schur, in Verh. Naturf. Ver. Bruenn, xv. II. (1877) 8 = recta.
uncinata, *Champ. ex Benth. in Hook. Kew Journ.* iii. (1851) 255.—Hongkong.
uniflora, Balb. Cat. Hort. Taur. (1810) 21 = ochroleuca.
urophylla, *Franch. in Bull. Soc. Linn. Par.* i. (1884) 433.—China.
velutina, Edgew. in Trans. Linn. Soc. xx. (1859) 24 = connata.
venosa, Royle, Illustr. Bot. Himal. 51 = connata.
verticillaris, *DC. Syst.* i. 166.—Am. bor.
villosa, *DC. l. c.* 154.—Ind. or.
violacea, *A. DC. in Mém. Soc. Phys. Genèv.* xi. (1845) 67.—Hab.?
Viorna, Andr. Bot. Rep. i. t. 71 = crispa.
Viorna, *Linn. Sp. Pl.* 543.—Am. bor.
viornoides, *Schrad. ex Steud. Nom.* ed. II. i. 380.—Am. bor.
virginiana, *Linn. Cent.* i. 15; *Amoen. Acad.* iv. 275.—Am. bor.
virginiana, Lour. Fl. Cochinch. 345 = apiifolia.
virginica, Thunb. Fl. Jap. 240 = biternata.
viridiflora, Bert. Misc. Bot. xix. 7. t. 3 = grata.
virosa, *Sterl. Hort. Nymph. ex Steud. Nom.* ed. II. i. 380.—Hab.?
Vitalba, Buch.-Ham. ex Steud. l. c. 379, 380 = grewiaefolia.
Vitalba, *Linn. Sp. Pl.* 544.—Europ.; Afr. bor.; Reg. Caucas.
Viticella, *Linn. l. c.* 543.—Europ. or.; As. Min.; Persia.
vitifolia, Wall. Cat. n. 4676 = Buchaniana.
Walteri, Pursh, Fl. Am. Sept. ii. 384 = crispa.
Welwitschii, *Hiern, ex O. Kuntze, in Verh. Bot. Ver. Brand.* xxvi. (1885) 171.—Afr. trop.
Wenderothii, *Steud. Nom.* ed. II. i. 380.—Europ.
Wightiana, *Wall. Cat.* n. 4674.—Ind. or.
Williamsii, *A. Gray, ex Perry, Jap. Exp.* ii. 306.—Japon.
Wyethii, Nutt. ex Torr. & Gray, Fl. N. Am. i. 8 = Douglasii.
zanzebarica, Sweet, Hort. Brit. ed. II. 1 = seq.
zanzibarensis, *Boj. ex Loud. Hort. Brit.* 228.—Ins. Zanzibar.
zeylanica, Poir. Encyc. Suppl. ii. 296 = Naravelia zeylanica.
Zollingeri, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 274 [bis].—Java.

CLEMATITARIA, Bur. Monog. Bign. 44 (1864) = **Bignonia**, Linn.

CLEMATITIS, [Tourn.] Linn. Syst. ed. I (1735) = **Clematis**, Dill. (Ranunculac.).
bracteata, Moench, Meth. Suppl. 103 = Clematis virginiana.
cordifolia, Moench, l. c. 104 = Clematis virginiana.
crispa, Moench, Meth. 296 = Clematis crispa.
erecta, Steud. Nom. ed. I. 206 = Clematis erecta.
flava, Moench, Meth. 296 = Clematis orientalis.
integrifolia, Moench, l. c. = Clematis integrifolia.
Vitalba, Moench, l. c. = Clematis Vitalba.

CLEMATOPSIS, Boj. ex Hook. Ic. Pl. i. t. 10 (1837) = **Clematis**, Dill. (Ranunculac.).
suaveolens, Boj. l. c. = Clematis Bojeri.

CLEMENTEA, Cav. in Anal. Cienc. Nat. vii. (1804) 63 = **Canavalia**, Adans. (Legumin.).
nitida, Cav. l. c. t. 47 = Can. ensiformis.

CLEOBULA, Vell. Fl. Flum. 187 (1825). *INC. SED.* CHEOBULA, Vell. Fl. Flum. iv. t. 146 (1827).
pinnata, *Vell. l. c.*—Bras.

CLEOBULIA, Mart. ex Benth. in Ann. Wien. Mus. ii. (1838) 131. *LEGUMINOSAE*, Benth. & Hook. f. i. 537.
diocleoides, *Benth. in Mart. Fl. Bras.* xv. I. 168.—Am. trop.
leiantha, *Benth. l. c.*—Bras.
multiflora, *Mart. ex Benth. in Ann. Wien. Mus.* ii. (1838) 131.—Bras.

CLEOCHROMA, Miers, in Hook. Lond. Journ. Bot. vii. (1848) 349 = **Iochroma**, Benth. (Solanac.).
calycina, Micrs, l. c. 350 = I. calycina.
grandiflora, Miers, l. c. 351 = I. grandiflora.
macrocalyx, Micrs, l. c. 350 = I. macrocalyx.

CLEODORA, Klotzsch, in Wiegm. Archiv, vii. (1841) 196 = **Croton**, Linn. (Euphorbiac.).
Sellowiana, Klotzsch, l. c. 197 = Croton sphaerogynus.

CLEOME, Linn. Syst. ed. I (1735). *CAPPARIDEAE*, Benth. & Hook. f. i. 105.
ALEOME, Neck. Elem. iii. 68 (1790).
ANOMALOSTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 162 (1861).
ARIVELA, Rafin. Sylva Tellur. 110 (1838).
ATALANTA, Nutt. Gen. Am. ii. 73 (1818).
AUBION, Rafin. Sylva Tellur. 110 (1838).
BUHSIA, Bunge, Del. Sem. Hort. Dorpat. (1859) 4.
CHILOCALYX, Klotzsch, in Peters, Reise Mossamb. Bot. 154 (1861).
COALISIA, Rafin. Sylva Tellur. 114 (1838).
CORYNANDRA, Schrad. Cat. Sem. Hort. Gotting. (1846).
DECASTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 157 (1861).
DIANTHERA, Klotzsch, l. c. 160. t. 27 (1861).
DIORIMASPERMA, Rafin. Sylva Tellur. 111 (1838).
HEMISCOLA, Rafin. l. c. (1838).
ISEXINA, Rafin. l. c. 112 (1838).
LAGANSA, Rumpf, ex Rafin. l. c. 110 (1838).
MELIDISCUS, Rafin. l. c. (1838).
MICAMBE, Adans. Fam. ii. 407 (1763).
MITOSTYLIS, Rafin. Sylva Tellur. 114 (1838).
MOZAMBE, Rafin. l. c. 112 (1838).
ONCUFIS, Rafin. l. c. 114 (1838).
PERICLA, Rafin. l. c. 112 (1838).
PERITOMA, DC. Prod. i. 237 (1824).
PHYSOSTEMON, Mart. Nov. Gen. et Sp. i. 73. t. 45 (1824).
PODOGYNE, Hoffmgg. Verz. Pfl. 185 (1840).
POLANISIA, Rafin. in Am. Monthly Mag. (1818) 267; Journ. Phys. (Août 1819) 98.
RORIDA, J. F. Gmel. Syst. 260 (1791).
RORIDULA, Forsk. Fl. Aegypt. Arab. 35 (1775).
SCOLOSPERMA, Rafin. Sylva Tellur. 111 (1838).
SIERUELA, Rafin. l. c. 112 (1838).
SILIQUARIA, Forsk. Fl. Aegypt. Arab. 78 (1775).
SINAPISTRUM, Tourn. ex Hall. in Rupp. Fl. Jen. 269 (1745).
STYLISTA, Rafin. Sylva Tellur. 112 (1838).
SYMPHYOSTEMON, Klotzsch, in Peters, Reise Mossamb. Bot. 159 (1861).
TARENAYA, Rafin. Sylva Tellur. 111 (1838).
TETRATELEIA, Sond. in Harv. & Sond. Fl. Cap. i. 58 (1859).
aculeata, *Linn. Syst.* ed. XII. iii. 232.—Am. trop.
acuta, Schum. & Thonn. in Dansk. Vidensk. Selsk. Afh. iv. (1828) 67 = Gynandropsis pentaphylla.
affinis, *DC. Prod.* i. 241.—Bras.
affinis, Spreng. Syst. iv. Cur. Post. 138 = Blumeana.
albescens, *Franch. in Revoil, Comali*, 11.—Afr. trop.
alliacea, Blanco, Fl. Filip. ed. I. 522 = Gynandropsis pentaphylla.
alliodora, Blanco, l. c. ed. II. 363 = Gynandropsis pentaphylla.
angulata, *Schult. f. Syst.* vii. 45.—Java.
angustifolia, Forsk. Fl. Aegypt. Arab. 120 = tenella.
anomala, *H. B. & K. Nov. Gen. et Sp.* v. 85.—N. Granat.
aphylla, Thunb. Prod. Pl. Cap. 109 = Schepperia juncea.
arabica, *Linn. Cent. Pl.* i. 20.—Afr. bor.; Arab.
arborea, *H. B. & K. Nov. Gen. et Sp.* v. 86.—Venezuela
arborea, Schrad. in Goett. Gel. Anz. i. (1821) 707 = Crataeva Tapia.
arborea, Weinm. in Syll. Ratisb. i. (1824) 227 = dendroides.
armata, *Thunb. Prod. Pl. Cap.* 109.—Afr. austr.
Aschersoniana, Pfund, in Flora, lvii. (1874) 413 = trinervia.
aspera, *Koen. in DC. Prod.* i. 241.—Ind. or.
atropurpurea, Schott, in Schreib. Nachr. Oestr. Naturf. i. 129 = dendroides.
aurea, *Celak. in Oestr. Bot. Zeitschr.* xxxiv. (1884) 113.—Graecia.
aurea, Nutt. ex Torr. & Gray, Fl. N. Am. i. 122 = lutea.
bicolor, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 330.—Bras.
Blumeana, *D. Dietr. Syn. Pl.* ii. 1065.—Java.
bororensis, *Oliver, Fl. Trop. Afr.* i. 81.—Afr. trop.
brachycarpa, *Vahl, ex DC. Prod.* i. 240.—Afr. bor.; Arab.; Ind. bor. occ.
brasiliensis, *Weinm. in Syll. Ratisb.* i. (1824) 122.—Bras.
brevisiliqua, *Schult. f. Syst.* vii. 40.—Arab.

CLEOME:—
Bungei, *Steud. Nom.* ed. II. i. 381.—China.
Burmanni, *Wight & Arn. Prod.* 22.—Ind. or.
canadensis, Hort. ex Schult. f. Syst. vii. 38 = cardaminoides.
Candelabrum, *Sims, Bot. Mag.* t. 2656.—Hab. ?
canescens, Stev. in DC. Prod. i. 241 = ornithopodioides.
capensis, Linn. Sp. Pl. ed. II. 940 = Heliophila callosa.
capparoides, Willd. ex Eichl. in Mart. Fl. Bras. xiii. 1. 247 = anomala.
cardaminoides, *Hoffmgg. Verz. Pfl.* 93.—Afr. austr.
cardinalis, *DC. Prod.* i. 238.—Mexic.
Chelidonii, Boj. in Ann. Sc. Nat. Sér. II. xx. (1843) 56 = strigosa.
Chelidonii, *Linn. f. Suppl.* 300.—Ind. or.; Java.
chilensis, *DC. Prod.* i. 238.—Chili.
chilocalyx, *Oliver, Fl. Trop. Afr.* i. 81.—Afr. trop.
chrysantha, *Decne. in Ann. Sc. Nat.* Sér. II. iii. (1835) 274.—Arab.; Nubia.
ciliata, *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* iv. (1828) 67.—Afr. trop.
coluteoides, *Boiss. Diagn.* Ser. I. i. 3.—Persia.
cordata, Burch. ex DC. Prod. i. 239 = monophylla.
cordata, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 68 = Ehrenbergiana.
cordobensis, *Eichl. ex Griseb. in Goett. Abh.* xix. (1874) 73.—Am. austr.
cremoloba, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 318.—Mexic.
crenopetala, *DC. in Mém. Soc. Phys. Genèv.* vi. (1836) 220. t. 2.—Am. austr.
cubensis, A. Rich. Ess. Fl. Cub. 74 = Houstoni.
cuneifolia, *Muhl. ex Nutt. Gen. Am.* ii. 73.—Georgia.
cypria, *Čelak. in Oestr. Bot. Zeitschr.* xxxiv. (1884) 114.—Ins. Cyprus.
decipiens, *Triana & Planch. in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 74.—N. Granat.
deflexa, *Ruiz & Pav. ex DC. Prod.* i. 240.—Peruv.
dendroidea, *Schult. f. Syst.* vii. 28.—Bras.
denticulata, *Schult. f. l. c.* 26.—Afr. trop.
diandra, *Burch. Trav.* i. 548.—Afr. austr.
dichotoma, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 318.—Bras.
didynama, *Hochst. ex Oliver, Fl. Trop. Afr.* i. 79.—Abyssin.
diffusa, *Banks, ex DC. Prod.* i. 241.—Bras.
diffusa, Roxb. Hort. Beng. [96]; Fl. Ind. iii. 129 = aspera.
digitata, *Forsk. Fl. Aegypt. Arab.* 120.—Arabia.
Dilleniana, DC. Prod. i. 240 = ornithopodioides.
diversifolia, Hochst. & Steud. ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 5 = brachycarpa.
dodecandra, Banks, ex Wight & Arn. Prod. 22, partim = Burmanni.
dodecandra, Linn. Sp. Pl. ed. II. 939 = graveolens.
dodecaphylla, *Vell. Fl. Flum.* 271; vi. t. 110.—Bras.
droserifolia, *Delile, Fl. Egypt.* 250.—Aegypt.; Arab.
Eckloniana, Schrad. in Linnaea, x. (1836) Litt. 109 = Gynandropsis pentaphylla.
edulis, *Rafin. Fl. Ludov.* 86.—Am. bor.
Ehrenbergiana, *Schweinf. Beitr. Fl. Aethiop.* i. 68.—Nubia.
epilobioides, Jan, Elench. fide Schult. f. Syst. vii. 47 = graveolens.
Erucago, Mill. Gard. Dict. ed. VIII. n. 6 = heptaphylla, spinosa.
felina, *Linn. f. Suppl.* 300.—Ind. or.
filifolia, Vahl, Symb. Bot. i. 48 = tenella.
fimbriata, *Vicary, in Ann. Nat. Hist.* Ser. II. i. (1848) 425.—Scinde.
flava, Banks, ex DC. Prod. i. 241 = graveolens.
flexuosa, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. 296; ex Schult. f. Syst. vii. 27 = Gynandropsis pentaphylla.
flexuosa, *Griseb. in Goett. Abh.* xix. (1874) 73.—Am. austr.
foliolosa, *DC. Prod.* i. 240.—Persia.
foliosa, *Hook. f. in Hook. Niger Fl.* 219.—Afr. trop.
frutescens, *Aubl. Pl. Gui.* ii. 677.—Guiana.
fruticosa, Linn. Sp. Pl. 671 = Cadaba indica.
fugax, *Schrad. Ind. Sem. Hort. Gotting.* (1830).—*Cf. Linnaea*, vi. (1831) *Litt.* 72—Hab. ?
gigantea, Blanco, Fl. Filip. ed. II. 364 = Gynandropsis speciosa.
gigantea, *Linn. Mant.* 130.—Am. austr.
glandulosa, *Ruiz & Pav. ex DC. Prod.* i. 238.—Peruv.

CLEOME:—
glauca, *DC. Prod.* i. 239.—Mesopotam.
glaucescens, *DC. l. c.*—Mesopotam.; Persia.
gracilis, Edgew. in Journ. As. Soc. Beng. xvi. (1847) II. 1212 = papillosa.
gracilis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 74.—N. Granat.
Grandidieri, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 462.—Madag.
grandiflora, *Ehrenb. ex Schweinf. Beitr. Fl. Aethiop.* 71.—Afr. trop.
graveolens, *Rafin. in Journ. Phys.* (*Août* 1819) 98.—Am. bor.
guianensis, *Aubl. Pl. Gui.* ii. 675.—Guiana.
guineensis, Hook. f. in Hook. Niger Fl. 218 = ciliata.
gynandra, Linn. Sp. Pl. 671 = pentaphylla.
heptaphylla, *Linn. l. c.* ed. II. 987.—Am. trop.
heptaphylla, G. F. W. Mey. Prim. Fl. Esseq. 222 = Houstoni.
heptaphylla, Sw. Obs. 252 = pungens.
heratensis, *Bunge & Bien. ex Boiss. Fl. Orient.* i. 412.—Afghanist.
heterotricha, Burch. Trav. i. 537 = Gynandropsis heterotricha.
hexandra, Poir. ex Steud. Nom. ed. I. 206 = arabica.
hirsuta, *Ruiz & Pav. ex DC. Prod.* i. 238.—Peruv.
hirta, *Oliver, Fl. Trop. Afr.* i. 81.—Afr. trop.
horrida, Mart. in Schult. f. Syst. vii. 32 = spinosa.
Hostmanni, Miq. in Linnaea, xxii. (1849) 470 = latifolia.
Houstoni, *R. Br. in Ait. Hort. Kew.* iv. 131.—Jamaica.
Houstoni, Miq. l. c. xix. (1847) 433 = diffusa.
Houtteana, *Schlecht. in Linnaea*, xxiv. (1851) 669.—Am. austr.
Humboldtii, *DC. Prod.* i. 241.—Am. trop.
iberica, DC. l. c. 240 = ornithopodioides.
iberidella, *Welw. in Oliver, Fl. Trop. Afr.* i. 79.—Afr. trop.
icosandra, Linn. Sp. Pl. 672 = graveolens.
indecora, *Hornem. Ind. Sem. Hort. Hafn.* (1846); *ex Walp. Ann.* i. 60.—Mexic.
integrifolia, *Torr. & Gray, Fl. N. Am.* i. 122.—Am. bor.
juncea, Berg. Desc. Pl. Cap. 164 = Heliophila callosa.
juncea, Sparm. in Nov. Act. Soc. Sc. Upsal. iii. (1780) 192 = Schepperia juncea.
khorassanica, *Bunge & Bien. ex Boiss. Fl. Orient.* i. 412.—Persia.
Kotschyana, *Boiss. l. c.* 413.—Assyria.
laevigata, Soland. ex DC. Prod. i. 200 = Stanleya gracilis.
lateralis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 73.—N. Granat.
latifolia, *Vahl, ex DC. Prod.* i. 239.—Guiana.
laxa, *Thunb. Prod. Pl. Cap.* 109.—Afr. austr.
Lechleri, *Eichl. in Mart. Fl. Bras.* xiii. 1. 250.—Peruv.
Leschenaultii, Schult. f. Syst. vii. 49 = Chelidonii.
linearis, Stocks, ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 3 = papillosa.
longifolia, *Presl, Rel. Haenk.* ii. 84.—Peruv.
longifolia, Vahl, ex DC. Prod. i. 240 = Wormskioldia heterophylla.
longifolia, Willd. ex Schult. f. Syst. vii. 1617 = speciosa.
longipes, *DC. Prod.* i. 239.—Ecuador.
lusitanica, Mill. Gard. Dict. ed. VIII. n. 3 = graveolens.
lutea, *Hook. Fl. Bor. Am.* i. 70. t. 25.—Am. bor. occ.
lutea, *E. Mey. Zwei Pfl. Docum.* 173.—Afr. austr.
macrantha, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 319.—Mexic.
macrophylla, Roxb. Hort. Beng. [96], nomen (quid ?).—Ind. or.
macrorhiza, *Sauv. Fl. Cub.* 4.—Cuba.
macrothyrsus, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 72.—N. Granat.
malhadensis, Mart. ex Schult. f. Syst. vii. 43 = aculeata.
melilotoides, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 320.—Persia.
mexicana, *D. Dietr. Syn. Pl.* ii. 1068.—Mexic.
mexicana, *Hemsl. Diag. Pl. Nov.* 20.—Mexic.
micrantha, *Desv. in Ham. Prod. Fl. Ind. Occ.* 48.—Guiana.
micrantha, Schrad. ex Schult. f. Syst. vii. 30 = Dactylaena micrantha.

CLEOME:—
monandra, DC. Pl. Rar. Hort. Genèv. 54. t. 15 = Dactylaena micrantha.
monophylla, Law, in J. Grah. Cat. Bomb. Pl. 7 = simplicifolia.
monophylla, *Linn. Sp. Pl.* 672.—Ind. or.; Afr. trop.
Moritziana, *Klotzsch, ex Eichl. in Mart. Fl. Bras.* xiii. 1. 250.—Venezuela.
moschata, Stocks, ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 4 = brachycarpa.
multicaulis, *Moç. & Sesse, ex DC. Prod.* i. 240.—Mexic.
muricata, *Edgew. in Journ. As. Soc. Beng.* xvi. (1847) II. 1212 = paradoxa.
muricata, *Schult. f. Syst.* vii. 25.—Ind. or.
nivalis, Vahl, ex DC. Syst. ii. 245 = Macropodium nivale.
Noëana, Boiss. Diagn. Ser. II. i. 48 = quinquenervia.
nummularia, DC. Prod. i. 239 = rosea.
ofidocarpa, Balb. Hort. Lugd. Alob. (1826); ex Schult. f. Syst. vii. 51 = Gynandropsis muricata.
oleracea, Welw. Apont. 586 = pentaphylla.
ornithopodioides, Forsk. Fl. Aegypt. Arab. n. 402 = brevisiliqua.
ornithopodioides, *Linn. Sp. Pl.* 672.—As. Min.; Persia.
oxalidea, *F. Muell. Fragm.* i. 69.—Austral.
oxypetala, *Boiss. Diagn.* Ser. I. vi. 20.—Persia; Arab.
oxyphylla, *Burch. Trav.* ii. 226.—Afr. austr.
pallida, *Kotschy, in Sitzb. Acad. Wien. Math. Nat.* lii. *Abth.* 1. (1866) 262.—Arabia.
palmipes, *Spreng. Syst.* ii. 122.—Guiana.
paludosa, *Willd. ex Eichl. in Mart. Fl. Bras.* xiii. 1. 255.—Bras.
palustris, *Salzm. ex Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 317.—Bras.
papillosa, *Steud. Nom.* ed. II. i. 362.—Aegypt.; Arab.; Ind. bor. occ.
paradoxa, *R. Br. in Salt, Abyss. App.* 65.—Abyssin.
parviflora, R. Br. in Salt, Abyss. App. 65 = brachycarpa.
parviflora, H. B. & K. Nov. Gen. et Sp. v. 83 = psoraleaefolia.
pedunculata, Vell. Fl. Flum. 272; vi. t. 112 = gigantea.
pentaphylla, Blanco, Fl. Filip. ed. I. 523 = Gynandropsis speciosa.
pentaphylla, Hort. Van Houtte, ex Eichl. in Mart. Fl. Bras. xiii. 1. 251 = Houtteana.
pentaphylla, Linn. Sp. Pl. ed. II. 938 = Gynandropsis pentaphylla.
pentaphylla, Vell. Fl. Flum. 272; vi. t. 111 = rosea.
pilosa, *Benth. Bot. Voy. Sulph.* 65.—Am. austr.
pinnata, Pursh, Fl. Am. Sept. ii. 739 = Stanleya pinnatifida.
platycarpa, *Torr. in Bot. U. St. Expl. Exped.* ii. 235. t. 2.—Am. bor. occ.
Poiteauana, *Pacquet, in Journ. Hort. Prat.* i. (1843) 349.—Hab. ?
polygama, Linn. Sp. Pl. ed. II. 939; DC. Prod. i. 241 = serrata.
porphyrantha, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 71.—N. Granat.
procumbens, *Jacq. Enum. Pl. Carib.* 261; *Linn. Syst.* ed. XII. 448.—Ind. occ.
prostrata, Hort. ex DC. Prod. ii. 382 = Rothia trifoliata.
prostrata, Koen. ex Steud. Nom. ed. II. i. 382 = aspera.
pruinosa, *T. Anders. in Journ. Linn. Soc. v. Suppl.* i. (1860) 3.—Arabia.
pruriens, *Planch. & Triana. in Ann. Sc. Nat.* Ser. IV. xvii. (1862) 68.—N. Granat.
psoraleaefolia, *DC. Prod.* i. 239.—Bras.
puberula, *Planch. & Triana. in Ann. Sc. Nat.* Ser. IV. xvii. (1862) 71.—N. Granat.
pubescens, Sieber, ex Steud. Nom. ed. II. i. 382 = Gynandropsis pentaphylla.
pubescens, *Sims, Bot. Mag.* t. 1857.—Panama.
pulchella, *Schult. f. Syst.* vii. 1617.—Bras.
pungens, Willd. Enum. Hort. Berol. 689 = spinosa.
purpurea, *DC. Prod.* i. 239.—Peruv.
quinquenervia, *DC. l. c.*—Persia; Afghanist.; Ind. bor. occ.
Raddeana, *Trautv. in Act. Hort. Petrop.* i. (1871) 19.—Turcoman.
Radula, Fenzl, in Flora, xxvii. (1844) 312 = papillosa.

CLEOME:—
ramosissima, *Parl. ex Webb, Fragm. Fl. Aethiop.* 22.—Afr. trop.
raphanoides, DC. Syst. ii. 662 = Wormskioldia heterophylla.
recta, *Jacquem. Voy. Bot.* 19.—Ind. or.
Regnellii, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 190.—Am. austr.
Richii, *A. Gray, in Bot. U. St. Expl. Exped.* i. 67.—Bras.
roridula, R. Br. in Salt, Abyss. App. 65 = droserifolia.
rosea, Eckl. ex Steud. Nom. ed. II. i. 382 = Gynandropsis pentaphylla.
rosea, *Poit. in Rev. Hortic.* Sér. II. i. (1841–43) 69.—Texas.
rosea, *Vahl, ex DC. Prod.* i. 239.—Bras.
rubella, *Burch. Trav.* i. 543.—Afr. austr.
rubiginosa, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 69.—N. Granat.
rupestris, *Sond. in Linnaea*, xxiii. (1850) 6.—Afr. austr.
rupicola, *Vicary, in Ann. Nat. Hist.* Ser. II. i. (1848) 425.—Scinde.
Ruta, Cambess. in Jacquem. Voy. Bot. 19. t. 19 = brachycarpa.
rutidosperma, *DC. Prod.* i. 241.—Ind. occ.?
sandwicensis, *A. Gray, Bot. U. St. Expl. Exped.* i. 65.—Ins. Sandvic.
scabrella, *Eichl. in Kjoeb. Vidensk. Meddel.* (1870) 188.—Am. trop.
scandens, Ruiz, ex Eichl. in Mart. Fl. Bras. xiii. 1. 251 = longipes?
scaposa, DC. Prod. i. 239 = papillosa.
Schraderi, Schult. f. Syst. vii. 48 = Chelidonii.
secundiflora, *Schrad. in Schult. f. l. c.* 37.—Hab.?
Selloana, *Klotzsch, ex Eichl. in Mart. Fl. Bras.* xiii. 1. 252.—Bras.
semibetandra, *Sond. in Linnaea*, xxiii. (1850) 5.—Afr. austr.
serrata, *Jacq. Enum. Pl. Carib.* 26.—Ind. occ.
serrulata, Pursh, Fl. Bor. Am. ii. 441 = integrifolia.
sessilifolia, *Schult. f. Syst.* vii. 23.—Ind. or. occ.
siliculifera, *Eichl. in Mart. Fl. Bras.* xiii. 1. 260.—Bras.
siliquaria, R. Br. in Salt, Abyss. App. 65 = arabica.
simplicifolia, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 169.—Ind. or.
socotrana, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 501.—Ins. Socotra.
Sonorae, *A. Gray, Pl. Wright.* ii. 16.—N. Mexic.
sparsifolia, *S. Wats. in Bot. King, Exp.* 32. t. 5.—Am. bor. occ.
speciosa, H. B. & K. Nov. Gen. et Sp. v. 84. t. 436 = Gynandropsis speciosa.
speciosa, *Rafin. Fl. Ludov.* 86.—Am. bor.
speciosissima, *Deppe, ex Lindl. Bot. Reg.* t. 1312.—Mexic.
spinosa, *Jacq. Enum. Pl. Carib.* 26; *Linn. Sp. Pl.* ed. II. 939.—Am. austr.
spinosa, Sw. Obs. 252 = pungens.
stenophylla, Klotzsch, in Rich. Schomb. Reise, iii. 1164 = Physostemon lanceolatum.
Stevensiana, Schult. f. Syst. vii. 40 = ornithopodioides.
Stocksiana, *Boiss. Diagn.* Ser. II. i. 47.—Beluchistan.
strigosa, *Oliver, Fl. Trop. Afr.* i. 80.—Afr. trop.
stylosa, *Eichl. in Mart. Fl. Bras.* xiii. 1. 250.—Venezuela.
subcordata, Steud. ex Oliver, in Trans. Linn. Soc. xxix. (1872) 28 = monophylla.
surinamensis, *Miq. in Linnaea*, xviii. (1844) 239.—Guiana.
tenella, *Linn. f. Suppl.* 300.—Ind. or.; Afr. trop.
tenuifolia, *Le Conte, ex Torr. & Gray, Fl. N. Am.* i. 123.—Am. bor.
ternata, Willd. ex Eichl. in Mart. Fl. Bras. xiii. 1. 257 = polygama.
tetrandra, *Banks, ex DC. Prod.* i. 240.—Austral.
tetraphylla, Willd. ex Eichl. in Mart. Fl. Bras. xiii. 1. 257 = psoraleaefolia.
titubans, *Speg. in Anal. Soc. Cient. Argent.* xv. (1883) 97.—Reg. Argent.
trachycarpa, *Klotzsch, ex Eichl. in Mart. Fl. Bras.* xiii. 1. 252.—Bras.
trinervia, *Fresen. in Mus. Senckenb.* i. (1834) 177.—Aegypt.; Arabia.
triphylla, Descourt. Fl. Med. Ant. i. t. 44 = polygama.

CLEOME:—
triphylla, James, in Long, Exped. ii. 239 = integrifolia.
triphylla, Linn. Sp. Pl. ed. II. 938 = serrata.
triphylla, Vell. Fl. Flum. 272; vi. t. 113 = affinis.
uniglandulosa, *Cav. Ic.* iv. 3.—Mexic.
urens, *Brot. ex Heynh. Nom.* ii. 149, nomen.—Hab.?
Vahliana, Fresen. in Mus. Senckenb. ii. (1837) 110 = brachycarpa.
venusta, Fenzl, in Flora, xxvii. (1844) 312 = paradoxa.
villosa, *Gardn. in Hook. Lond. Journ. Bot.* i. (1842) 166.—Bras.
viminea, *Schrad. in Linnaea*, x. (1836) *Litt.* 109.—Hab.?
violacea, Linn. Sp. Pl. 672 = graveolens.
virgata, Stev. in DC. Prod. i. 240 = ornithopodioides.
virgata, *Thunb. Prod. Pl. Cap.* 109.—Afr. austr.
viridiflora, Schreb. in Nov. Act. Nat. Cur. iv. (1770) 136. t. 3 = gigantea.
viscosa, *Linn. Sp. Pl.* 672.—Geront. trop.

CLEOMELLA, DC. Prod. i. 237 (1824). *CAPPARIDEAE*, Benth. & Hook. f. i. 105.
Isopara, Rafin. Atl. Journ. 144 (1832).
angustifolia, *Torr. in Hook. Kew Journ.* ii. (1850) 255.—Mexic.
brevipes, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 365.—Calif.
Coulteri, Harv. ex A. Gray, Pl. Wright. i. 12, 129 = Wislizenia refracta.
Fremontii, *Torr. in Smithson, Contrib.* iii. (1852) 12, in nota.—Am. bor.
longipes, *Torr. in Hook. Kew Journ.* ii. (1850) 255.—Mexic.
medicaginea, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. 313.—Mexic.
mexicana, *Moç. & Sesse, ex DC. Prod.* i. 237.—Mexic.
mexicana, Torr. in Ann. Lyc. N. York, ii. (1828) 167 = angustifolia.
obtusifolia, *Torr. & Frém. in Frém. Second Report*, 311.—Am. bor. occ.
oocarpa, *A. Gray, in Proc. Am. Acad.* xi. (1876) 72.—Am. bor. occ.
parviflora, *A. Gray, l. c.* vi. (1864–65) 520.—Calif.
plocasperma, *S. Wats. in Bot. King, Exp.* 33.—Am. bor. occ.

CLEOMENA, Roem. & Schult. Syst. ii. 18, 383 (1817) = **Muehlenbergia**, Schreb. (Gramin.).

CLEONIA, Linn. Sp. Pl. ed. II. 837 (1762). *LABIATAE*, Benth. & Hook. f. ii. 1203.
lusitanica, *Linn. l. c.*—Europ. austr.

CLEOPHORA, Gaertn. Fruct. ii. 185. t. 120 (1791) = **Latania**, Comm. (Palm.).
dendriformis, Lodd. ex Baker, Fl. Maurit. 381 = L. Loddigesii.
lontaroides, Gaertn. Fruct. ii. 185. t. 120 = L. Commersonii.

CLERCIA, Vell. Fl. Flum. 29; i. t. 73, 74 (1825) = **Salacia**, Linn. (Celastr.).
clusiaefolia, Miers, in Trans. Linn. Soc. xxviii. (1872) 381 = S. sylvestris.
dispansa, Miers, l. c. 379.—Bras.
guianensis, Miers, l. c. 378.—Guiana.
lanceolata, Miers, l. c. 377.—Bras.
micrantha, Miers, l. c. 379 = S. Miersii.
ovata, Vell. Fl. Flum. 30; i. t. 73 = S. arborea.
Passiflora, Vell. l. c. t. 74.—Bras.
retusa, Miers, in Trans. Linn. Soc. xxviii. (1872) 380 = S. sylvestris.
Tweediana, Miers, l. c.—Bras.

CLERKIA, Neck. Elem. i. 251 (1790) = **Tabernaemontana**, Plum. (Apocyn.).

CLERMONTIA, Gaudich. in Freyc. Voy. Bot. 459. t. 71 ad 73 (1826). *CAMPANULACEAE*, Benth. & Hook. f. ii. 546.
grandiflora, *Gaudich. l. c.* t. 73.—Ins. Sandvic.
Kakeana, *Meyen, Reise*, ii. 139.—Ins. Sandvic.
macrocarpa, Gaudich. Bot. Voy. Bonite, t. 49 = Kakeana.
macrophylla, Nutt. Trans. Am. Phil. Soc. N. S. viii. (1843) 251 = Kakeana.
oblongifolia, *Gaudich. in Freyc. Voy. Bot.* 459. t. 71.—Ins. Sandvic.

CLERMONTIA:—
parviflora, *Gaudich. ex A. Gray, in Proc. Am. Acad.* v. (1861) 150.—Ins. Sandvic.
persicifolia, *Gaudich. in Freyc. Voy. Bot.* 459. t. 72.—Ins. Sandvic.

CLERODENDRON, Linn. Gen. ed. I. 186 (1737). *VERBENACEAE*, Benth. & Hook. f. ii. 1155.
Agricolaea, Schrank, in Denkschr. Akad. Muench. (1808) 98.
Cornacchinia, Savi, in Mem. Soc. Ital. Mod. xxi. (1837) 184. t. 7.
Cyclonema, Hochst. in Flora, xxv. (1842) 225.
Cyrtostemma, Kunze, in Bot. Zeit. i. (1843) 272.
Douglassia, Adans. Fam. ii. 200 (1763).
Egena, Rafin. Fl. Tellur. ii. 85 (1836).
Ovieda, Linn. Gen. ed. I. 59 (1737).
Rotheca, Rafin. Fl. Tellur. iv. 69 (1836).
Siphonanthus, Linn. Gen. ed. II. 526 (1742).
Spironema, Hochst. in Flora, xxv. (1842) 226.
Torreya, Spreng. Neue Entdeck. ii. 221 (1821).
Valdia, Plum. ex Adans. Fam. ii. 157 (1763).
Volkameria, Linn. Syst. ed. I (1735); Gen. ed. I. 347 (1737).
Volkmannia, Jacq. Hort. Schoenbr. iii. 48. t. 338 (1798).
Acerbiana, *Benth. & Hook. f. Gen.* ii. 1156.—Nubia.
aculeatum, *Griseb. Fl. Brit. W. Ind.* 500.—Jamaic.
acuminatum, Wall. Cat. n. 1792 = disparifolium.
affine, *Griff. Notul.* iv. 170.—Burma.
amicorum, *Seem. in Bonplandia*, x. (1862) 249.—Ins. Pacif.
amplius, Hance, in Ann. Sc. Nat. Sér. V. v. (1866) 233 = cyrtophyllum.
angustifolium, Salisb. Prod. 108 = fortunatum.
angustifolium, *Spreng. Syst.* ii. 758.—Jamaic.
attenuatum, R. Br. Prod. 511 = floribundum.
aurantium, *G. Don, in Edinb. Phil. Journ.* xi. (Oct. 1824) 349.—Afr. trop.
Arthur-Gordoni, Horne, ex Baker, in Journ. Linn. Soc. xx. (1883) 370 = Gordoni.
Balfouri, Hort. ex Flor. Mag. viii. (1869) sub t. 432 = Thomsonae.
Bethuneanum, *Low, Sarawak*, 378; *et in Bot. Mag.* t. 4485.—Borneo.
Blancoanum, Villar, in Blanco, Fl. Philipp. ed. III. Nov. App. 161 = Navesianum.
Blancoi, *Naves, ex Villar, l. c.*—Ins. Philipp.
Blumeanum, *Schau. in DC. Prod.* xi. 669.—Java.
brachyanthum, *Schau. l. c.* 668.—Ins. Philipp.
bracteatum, *Wall. Cat.* n. 1800.—Reg. Himal.
brunsvigioides, *Baker, in Journ. Linn. Soc.* xxi. (1885) 435.—Madag.
Buchananii, *Herb. Roxb. ex Wall. Cat.* 2653 (sp. dub.).—Ind. or.
Bungei, Steud. Nom. ed. II. i. 382 = foetidum.
buruanum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 252.—Malaya.
buxifolium, *Spreng. Syst.* ii. 758.—Hab.?
calamistratum, Hort. Belg. ex Lem. Jard. Fleur. iv. (1854) Misc. 48 = elegans.
calamitosum, *Linn. Mant.* i. 90.—Malaya.
calycinum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 222 = infortunatum.
canescens, *Wall. Cat.* n. 1800.—China.
capense, *D. Don, ex Steud. Nom.* ed. II. i. 382.—Afr. austr.
capense, Eckl. & Zeyh. ex Schau. in DC. Prod. xi. 661 = glabrum.
capitatum, Hook. Bot. Mag. t. 4355 = Whitfieldii.
capitatum, *Schum & Thou. in Dan. Vid. Selsk. Afh.* iv. (1828) 61.—Afr. trop.
capsulare, Blanco, Fl. Filip. ed. I. 509 = inerme.
cardiophyllum, F. Muell. Fragm. iii. 144 = floribundum.
castaneaefolium, Klotzsch, in Bot. Ergeb. Waldem. Reise, 102 = infortunatum.
castaneifolium, Hook. & Arn. Bot. Beech. Voy. 205 = fortunatum.
cauliflorum, *Vatke, in Linnaea*, xliii. (1880–82) 538.—Ins. Nossibé.
cernuum, *Wall. ex Steud. Nom.* ed. II. i. 382.—Ind. or.
coccineum, D. Dietr. Syn. Pl. iii. 616 = squamatum.
Colebrookianum, *Walp. Rep.* iv. 114.—Ind. or.; Malaya.
Commersonii, *Spreng. Syst.* ii. 758.—Ins. Philipp.
condensatum, *Miq. Fl. Ind. Bat.* ii. 874.—Malaya.

CLERODENDRON:—
cordatum, D. Don, Prod. Fl. Nep. 103 = infortunatum.
cordifolium, *A. Rich. Tent. Fl. Abyss.* ii. 170.—Abyssin.
coriaceum, R. Br. Prod. 511 = floribundum.
coriaceum, Poir. Encyc. Suppl. iv. 353 = javanicum.
coromandelianum, Sprcng. Syst. ii. 758 = ovatum.
costatum, *R. Br. Prod.* 511.—Austral.
cruentum, *Lindl. in Gard. Chron.* (1860) 456.—As. trop.
cubensis, *Schau. in DC. Prod.* xi. 658.—Cuba.
Cuningianum, *Schau. l. c.* 666.—Ins. Philipp.
cuneatum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 221 = serratum.
Cunninghamii, *Benth. Fl. Austral.* v. 64.—Austral.
cuspidatum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 221.—N. Granat.
cyrtophyllum, *Turcz. l. c.* 222.—China.
deflexum, *Wall. Pl. As. Rar.* iii. 10. t. 215.—Malaya.
densiflorum, *Griff. Notul.* iv. 170.—Burma.
dentatum, Wall. Cat. n. 1799 = squamatum.
depauperatum, Wall. ex Steud. Nom. ed. II. i. 382 = infortunatum.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* 568.—Malaya.
discolor, *Vatke, in Linnaea*, xliii. (1880-82) 536.—Afr. trop.
disparifolium, *Blume, Bijdr.* 809.—Malaya.
divaricatum, Jack, in Malay. Misc. i. (1820) n. v. p. 48 = serratum.
divaricatum, Sieb. & Zuec. in Abh. Akad. Mueneh. iv. III. (1846) 154 = Caryopteris divaricata.
diversifolium, Vahl, Symb. Bot. ii. 75 = emirnense?
elegans, *Manetti, ex Lem. Jard. Fleur.* iv. (1854) *Misc.* 47.—Hab.?
ellipticum, Zipp. ex Span. in Linnaea, xv. (1841) 329 = lacvifolium.
emirnense, *Boj. ex Hook. Bot. Mag.* t. 2925.—Madag.
eriosiphon, *Schau. in DC. Prod.* xi. 662.—Java.
fallax, *Lindl. Bot. Reg.* (1844) sub t. 19.—Java.
farinosum, Wall. Cat. n. 1810 = serratum.
ferrugineum, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) II. 221 = villosum.
fistulosum, *Becc. Malesia*, ii. (1884) 48.—Malaya.
floribundum, *R. Br. Prod.* 511.—Austral.
floribundum, Hort. Angl. ex Walp. Rep. iv. 111 = emirnense.
foctidum, *Bunge, Enum. Pl. Chin. Bor.* 52.—China.
foetidum, D. Don, Prod. Fl. Nep. 103 = Caryopteris grata.
foetidum, Hort. Par. ex Planch. in Fl. des Serres, Sér. I. ix. (1853-54) 17 = fragrans, *Vent.*
fortunatum, Buch.-Ham. ex Wall. Cat. n. 2652 (sp. dub.).—Assam.
fortunatum, *Linn. Cent. Pl.* ii. 23.—China.
fragrans, *Vent. Jard. Malm.* t. 70.—China; Sumatra; Java.
fragrans, *Willd. Enum. Hort. Berol.* 659.—Am. trop.
galcatum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 91.—Ins. Socotra.
glabrum, *E. Mey. Comm. Pl. Afr. Austr.* i 273.—Afr. austr.
glandulosum, Colebr. ex Wall. Cat. n. 1806 = Colebrookianum.
glandulosum, Lindl. Bot. Reg. (1844) sub t. 19 = praec.?
glaucum, *Wall. ex Steud. Nom.* ed. II. i. 383.—Ind. or.
Gordoni, *Baker, in Journ. Linn. Soc.* xx. (1883) 370.—Ins. Fiji.
grandiflorum, *Schau. in DC. Prod.* xi. 659.—Hab.?
grandifolium, Salisb. Prod. 108 = serrata.
gratum, Kurz, For. Fl. Brit. Burma, ii. 268 = Caryopteris paniculata.
gratum, Wall. Cat. n. 1813 = Caryopteris grata.
Griffithianum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 590.—Assam; Burma.
haematocalyx, Hance, in Walp. Ann. iii. 238 = canescens.
Harnierianum, *Schweinf. Beitr. Fl. Aethiop.* 119.—Afr. trop.
hastatum, *Lindl. Bot. Reg.* t. 1307.—Reg. Himal.
helianthemifolium, Wall. ex Steud. Nom. ed. II. i. 383, sphalm. = Caryopteris Wallichiana.

CLERODENDRON:—
hemiderma, *F. Muell. ex Benth. Fl. Austral.* v. 61.—Austral.
herbaceum, Wall. Cat. n. 1815 = serratum.
heterophyllum, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 64.—Ins. Mascar.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880-82) 536.—Afr. trop.
hircinum, *Schau. in DC. Prod.* xi. 661.—Madag.
hirsutum, G. Don, in Edinb. Phil. Journ. xi. (1824) 349 = umbellatum.
Horsfieldii, *Miq. Fl. Ind. Bat.* ii. 880.—Java.
Huegelii, *Hort. ex Regel, Gartenfl.* (1861) 51.—Austral.
illustre, *N. E. Br. in Gard. Chron.* (1884) II. 424.—Ins. Celebes.
imperialis, *Carr. in Rev. Hortic.* (1874) 110.—Hab.?
incisum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 257.—Afr. trop.
inerme, R. Br. Prod. 511 = neriifolium.
inerme, *Gaertn. Fruct.* i. 271.—Ind. or.
infortunatum, *Gaertn. l. c.*—Ind. or.; Malaya.
infortunatum, Dennst. Schluess. Hort. Malab. 27 = villosum.
intermedium, *Cham. in Linnaea*, vii. (1832) 150.—Ins. Luzon.
involucratum, *Vatke, in Linnaea*, xliii. (1880-82) 537.—Afr. trop.
ixoraeflorum, *Hassk. in Retzia*, 60.—Singapore.
Jackianum, Wall. in Hook. Bot. Misc. i. (1830) 284; Cat. n. 1794 = disparifolium.
japonicum, *Sweet, Hort. Brit.* ed. I. 322.—Japon.
javanicum, *Spreng. Syst.* ii. 759.—Java.
javanicum, Walp. Rep. iv. 113 = serratum.
Kaempferi, Fisch. ex Morr. Ann. Soc. Gand, i. (1845) 17, in syn. = squamatum.
laciniatum, *Balf. f. in Journ. Linn. Soc.* xvi. (1877) 19.—Ins. Mascar.
laevifolium, *Blume, Bijdr.* 808.—Malaya.
lanceolatum, *F. Muell. Fragm.* iii. 145.—Austral.
lanuginosum, *Blume, Bijdr.* 810.—Ins. Molucc.
lasiocephalum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 594.—Reg. Himal.
laxiflorum, *Baker, in Journ. Linn. Soc.* xx. (1883) 229.—Ins. Fiji.
Lehuntei, *Horne, ex Baker, l. c.* 369.—Ins. Fiji.
leucophloeum, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 91.—Ins. Socotra.
Leucosceptrum, D. Don, Prod. Fl. Nep. 103 = Leucosceptrum canum.
ligustrinum, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 64.—Mexic.
Lindemuthianum, *Vatke, in Linnaea*, xliii. (1880-82) 537.—Afr. trop.
Lindenianum, *A. Rich. Fl. Cub. Fanerog.* ii. 147.—Cuba.
Lindleyi, Decne. ex Planch. in Fl. des Serres, Sér. I. ix. (1853-54) 17 = fragrans, *Vent.*
Linnaei, F. Muell. Fragm. vi. 151 = hemiderma.
Linnaei, Thw. Enum. Pl. Zeyl. 243 = Glossocarya Linnaei.
lividum, Lindl. Bot. Reg. t. 945 = fortunatum.
Lobbii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 590.—Ins. Penang.
longicolle, *G. F. W. Mey. Prim. Fl. Esseq.* 217.—Guiana.
longiflorum, *Decne. in Nov. Ann. Mus. Par.* iii. (1834) 400.—Ins. Timor.
macradenium, *Miq. Fl. Ind. Bat.* ii. 874.—Java.
macroealyeinum, *Baker, in Journ. Linn. Soc.* xviii. (1881) 275.—Madag.
macrophyllum, *Blume, Bijdr.* 808.—Java.
macrophyllum, Sims, Bot. Mag. t. 2536 = serratum.
macrosiphon, *Hook. f. Bot. Mag.* t. 6695.—Afr. trop.
macrostachyum, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) II. 220.—Reg. Himal.; Burma.
macrostegium, *Schau. in DC. Prod.* xi. 666.—Ins. Philipp.
Madaccra, *Voigt, Hort. Suburb. Calc.* 467.—Ind. or.?
magnoliaefolium, *Baker, in Journ. Bot.* xx. (1882) 243.—Madag.
Manetti, *Vis. Sem. Hort. Patav.* (1848-49) n. 2.—Hab.?
medium, R. Br. Prod. 510 = floribundum.
Minahassae, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxv. (1863) 409.—Ins. Celebes.
mite, *Vatke, in Linnaea*, xliii. (1880-82) 537.—Afr. trop.
molle, *H. B. & K. Nov. Gen. et Sp.* ii. 244.—Ecuador.

CLERODENDRON:—
molle, Jack, in Malay. Misc. i. (1820) n. I. 15 = villosum.
mossambicense, *Klotzsch, in Peters, Reise Mossamb. Bot.* 259.—Afr. trop.
multiflorum, G. Don, in Edinb. Phil. Journ. xi. (1824) 350 = volubile?
myricoides, *R. Br. in Salt, Abyss. App.* p. lxv, nomen.—Abyssin.
Navesianum, *Vidal, Cat. Prov. Manila*, n. 398.—Ins. Philipp.
nereifolium, *Wall. Cat.* n. 1789.—Malaya.
Neumayeri, *Vatke, in Linnaea*, xliii. (1880-82) 535.—Afr. trop.
nutans, Jack, in Malay. Misc. i. (1820) n. I. 17 = Jackiannm.
nutans, *Wall. Cat.* n. 1793.—Reg. Himal.; Burma.
obovatum, *Walp. Rep.* iv. 112.—Ins. Molucc.
obtusidens, *Miq. Fl. Ind. Bat.* ii. 870.—Malaya.
odoratum, D. Don, Prod. Fl. Nep. 102 = Caryopteris Wallichiana.
ornatum, Wall. Cat. n. 1811 = serratum.
ovale, *Klotzsch, in Peters, Reise Mossamb. Bot.* 257.—Afr. trop.
ovalifolium, *A. Gray, in Proc. Am. Acad.* vi. (1862) 50.—Ins. Fiji.
ovatum, Poir. Encyc. Suppl. iv. 352 = coromandelinnm.
ovatum, R. Br. Prod. 511 = floribundum.
oxysepalum, *Miq. in Journ. Bot. Néerl.* i. (1861) 114.—China.
paniculatum, *Linn. Mant.* i. 90.—Burma; Malaya; China.
papuanum, *Scheff. in Ann. Jard. Buitenz.* i. (1876) 41.—N. Guin.
penduliflorum, *Wall. Cat.* n. 1795.—Burma.
pentagonum, Hance, in Walp. Ann. iii. 238 = fortunatum.
petunioides, *Baker, in Journ. Linn. Soc.* xx. (1883) 230.—Madag.
philippinum, *Schau. in DC. Prod.* xi. 667.—Malaya.
phlomoides, Hort. Ital. ex DC. l. e. v. 613 = Montanoa arboreseens.
phlomoides, *Linn. f. Suppl.* 292.—Ind. or.
phyllomega, Steud. Nom. ed. II. i. 383 = macrophyllum.
pubescens, *Lindl. in Bot. Reg.* t. 1035.—Ind. or.
pubescens, Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 380, lapsu = canescens.
pulchrum, *Fawc. in O. Forbes, Nat. Wand. East. Archip.* 514.—Ins. Timor.
pumilum, *Spreng. Syst.* ii. 759.—China.
putre, *Schau. in DC. Prod.* xi. 661.—Madag.
pyramidale, Andr. Bot. Rep. t. 628 = paniculatum.
pyrifolium, *Baker, in Journ. Linn. Soc.* xx. (1883) 228.—Madag.
ramosissimum, *Baker, l. c.*—Madag.
Riedelii, *Oliver, in Journ. Linn. Soc.* xv. (1876) 100.—Ins. Celebes.
robustum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 259.—Afr. trop.
roseum, *Poit. in Rev. Hortic.* (1876) 80.—Hab.?
rotundifolium, *Oliver, in Trans. Linn. Soc.* xxix. (1875) 132.—Afr. trop.
rubellum, *Baker, in Journ. Linn. Soc.* xx. (1883) 229.—Madag.
Rumphianum, *De Vriese, in Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 252.—Ins. Amboina.
sagittatum, Wall. Cat. n. 1786 = hastatum.
Sagraei, *Schau. in DC. Prod.* xi. 6591.—Ins. Cuba.
scandens, Beauv. Fl. Owar. i. 52. t. 32 = umbellatum.
seopiferum, *Miq. Fl. Ind. Bat.* ii. 881.—Malaya.
semiserratum, *Wall. Cat.* n. 1785.—Burma.
sericeum, Wall. l. e. n. 1814 = Hiptage sericea.
serotinum, Carr. in Rev. Hortic. (1867) 351 = trichotomum.
serratum, *Spreng. Syst.* ii. 758.—Ind. or.; Burma.
simplex, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 349.—Afr. trop.
singalense, *Miq. Fl. Ind. Bat. Suppl.* 568.—Sumatra.
sinuatum, *Hook. Bot. Mag.* t. 4255.—Afr. trop.
Siphonanthus, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 65.—Ind. or.; Malaya.
speciosissimum, Hort. Angl. ex Schau. in DC. Prod. xi. 672 = glandulosum.
speciosissimum, Paxt. Mag. Bot. iii. (1837) 217 et 271 = squamatum.
speciosum ×, Teijsm. & Binn. Cat. Hort. Bog. 386; et Lem. Illustr. Hortic. xvi. (1869) t. 593.
spicatum, *Thunb. Fl. Japan.* 22.—Java.
spinosum, *Spreng. Syst.* ii. 760.—Ins. Dominic.

CLERODENDRON:—
splendens, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 349.—Afr. trop.
splendidum, Wall. Cat. n. 1803 = paniculatum.
squamatum, *Vahl, Symb. Bot.* ii. 74.—Reg. Himal.; Burma; China.
stenanthum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 258.—Afr. trop.
ternifolium, *Baker, in Journ. Linn. Soc.* xx. (1883) 229.—Madag.
ternifolium, D. Don, Prod. Fl. Nep. 103 = serratum.
ternifolium, *H. B. & K. Nov. Gen. et Sp.* ii. 244.—Fl. Orinoco.
Thomsonae, *Balf. in Edinb. New Phil. Journ.* N. S. xv. (Jan.–Apr. 1862) 233.—Afr. trop.
tomentosum, *R. Br. Prod.* 510.—Austral.
Tracyanum, *F. Muell. ex Benth. Fl. Austri.* v. 62.—Austral.
trichotomum, *Thunb. Fl. Jap.* 256.—Japon.
trichotomum, Wall. Cat. n. 6315 B = villosum.
triflorum, *Vis. Illustr. Piante Nov. Ort Padov.* 19.—Nubia.
trifoliatum, *Steud. Nom.* ed. II. i. 383.—Ind. or.
tuberculatum, *A. Rich. Fl. Cub. Fanerog.* ii. 147.—Cuba.
umbellatum, *Poir. Encyc.* v. 166.—Afr. trop.
urticifolium, Wall. Cat. n. 1801 = squamatum.
velutinum, Wall. l. c. n. 1797 = villosum.
venosum, *Wall. l. c.* n. 1807.—Reg. Himal.
verrucosum, *Splitg. ex De Vriese, in Nederl. Kruidk. Arch.* i. (1848) 351.—Guiana.
verticillatum, D. Don, Prod. Fl. Nep. 102 = C. Siphonanthus.
vestitum, Wall. ex Steud. Nom. ed. II. i. 383 = infortunatum.
villosum, *Blume, Bijdr.* 811.—Malaya.
viscosum, Vent. Jard. Malm. t. 25 = infortunatum.
volubile, *Beauv. Fl. Owar.* i. 52. t. 32.—Afr. trop.
Whitfieldii, *Seem. in Bonplandia*, x. (1862) 250.—Afr. trop.

CLERODENDRUM, Linn. = praec.

CLETHRA, Bert. ex Steud. Nom. ed. II. i. 384 (1840) = **Viviania**, Cav.
mollis, Bert. ex Steud. l. c. = V. parvifolia.

CLETHRA, Gronov. ex Linn. Syst. ed. I (1735). *ERICACEAE*, Benth. & Hook. f. ii. 603.
Crossophrys, Klotzsch, in Linnaea, xxiv. (1851) 3.
Cuellaria, Ruiz & Pav. Prod. 59. t. 10 (1794); Syst. Veg. 103 (1798).
Gillena, Adans. Fam. ii. 166 (1763).
Junia, Adans. l. c. 165 (1763).
Kowalewskia, Turcz. in Bull. Soc. Nat. Mosc. (1859) i. 263.
Tinus, Linn. Syst. ed. X. 1010 (1759).
Volkameria, P. Br. Hist. Jamaic. 214. t. 21. f. 1 (1756).
acuminata, *Michx. Fl. Bor. Am.* i. 260.—Am. bor.
Alexandri, *Griseb. Fl. Brit. W. Ind.* 142.—Ind. occ.
alnifolia, Blanco, Fl. Filip. ed. II. 259 = canescens.
alnifolia, *Linn. Sp. Pl.* 396.—Am. bor.
angustifolia, *DC. Prod.* vii. 589.—Chili.
arborea, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 73.—Ins. Madera.
arborea, Vent. Jard. Malm. t. 40 = secundiflora.
barbinervis, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. iii. (1846) 128.—Japon.
bicolor, *H. B. & K. Nov. Gen. et Sp.* iii. 289.—N. Granat.
bracteata, *Griseb. Fl. Brit. W. Ind.* 141.—Ind. occ.
brasiliensis, *Cham. & Schlecht. in Linnaea*, viii. (1833) 510.—Bras.
brevifolia, *Benth. Pl. Hartw.* 143.—N. Granat.
byrsonimoides, *Griseb. Pl. Wright. Cub.* 170.—Cuba.
canescens, *Herb. Reinw. ex Blume, Bijdr.* 863.—Java.
castaneaefolia, *Meissn. in Mart. Fl. Bras.* vii. 169.—Peruv.
confertifolia, *Ernst, in Flora*, lvii. (1874) 213.—Venezuela.
cubensis, *A. Rich. Fl. Cub. Fanerog.* ii. 75.—Cuba.
dentata, [Soland. in] Ait. Hort. Kew. ed. I. ii. 73 = alnifolia.
divaricata, *Loisel. in Duham. Arb. ed. Nov.* v. 134.—Am. austr.
Fabri, *Hance, in Journ. Bot.* xxi. (1883) 130.—China.
fagifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 289.—Am. austr.
ferruginea, *Link, Enum. Hort. Berol.* i. 396, nomen.—Hab.?

CLETHRA:—
ferruginea, *Ruiz & Pav. Fl. Per.* iv. t. 380.—Peruv.
fimbriata, *H. B. & K. Nov. Gen. et Sp.* iii. 290. t. 264.—N. Granat.
Gardneri, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 233.—Bras.
glauca, Hort. ex Steud. Nom. ed. II. i. 383 = pubescens.
glauca, Pers. Syn. i. 483 = acuminata.
guianensis, *Klotzsch, ex Meissn. in Mart. Fl. Bras.* vii. 170.—Guiana.
incana, Pers. Syn. i. 483 = pubescens.
integrifolia, Moç. & Sesse, ex DC. Prod. vii. 590 = mexicana.
japonica, Thunb. ex Steud. Nom. ed. II. i. 383 = barbinervis.
javanica, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 232.—Java.
Kowalewskii, Turcz. l. c. 233 = mexicana.
laevigata, *Meissn. in Mart. Fl. Bras.* vii. 168.—Bras.
lanata, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 538.—Mexic.
lancifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 231.—Singapore.
macrophylla, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) 539.—Mexic.
macrostachya, *Loisel. in Duham. Arb. ed. Nov.* v. 155.—Hab.?
Mathewsii, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 232.—Peruv.
mexicana, *DC. Prod.* vii. 590.—Mexic.
Michauxi, *R. Court. in Mag. Hortic.* i. (1833) 249.—Am. bor.
micrantha, *Remy, in Ann. Sc. Nat.* Sér. III. viii. (1847) 233.—N. Granat.
montana, Fras. ex Loisel. in Duham. Arb. ed. Nov. v. 130 = acuminata.
nervosa, *Loisel. in Duham. l. c.* 133.—Peruv.
obovata, Hook. & Arn. Bot. Beech. Voy. 302 = quercifolia.
obovata, *Ruiz & Pav. Fl. Per.* iv. t. 381.—Peruv.
ocanensis, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 231.—N. Granat.
occidentalis, Auct. ex Steud. Nom. ed. II. i. 383 = tinifolia.
ovalifolia, Moç. & Sesse, ex DC. Prod. vii. 590 = mexicana.
ovalifolia, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 232.—Reg. Andin.
paniculata, [Soland. in] Ait. Hort. Kew. ed. I. ii. 73 = alnifolia.
pedicellaris, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 234.—Peruv.
pubescens, Willd. Enum. Hort. Berol. 455 = alnifolia.
punctata, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 235.—N. Granat.
quercifolia, *Schlecht. in Linnaea*, viii. (1833) 524; *Lindl. Bot. Reg.* (1842) t. 23.—Mexic.
repanda, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 231—Venezuela.
revoluta, *Ruiz & Pav. Fl. Per.* iv. t. 380.—Peruv.
scabra, Loisel. in Duham. Arb. ed. Nov. v. 132 = brasiliensis.
scabra, Pers. Syn. i. 482 = alnifolia.
secundiflora, *Decne. in Fl. des Serres*, Sér. II. xii. (1877) 171, in nota.—Hab.?
serrulata, Turcz. in Bull. Soc. Nat. Mosc. xxxvi. (1863) ii. 233 = mexicana.
spicigera, *Meissn. in Mart. Fl. Bras.* vii. 168.—Bras.
Sprucei, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) ii. 232.—Peruv.
suaveolens, *Turcz. l. c.* 230.—Mexic.
tinifolia, Schlecht. in Linnaea, viii. (1833) 524 = quercifolia.
tinifolia, *Sw. Prod. Veg. Ind. Occ.* 74.—Jamaic.
tomentosa, Lam. Encyc. ii. 46 = alnifolia.
virgata, *Ruiz & Pav. Fl. Per.* iv. t. 381.—Peruv.

CLETHROPSIS, Spach, in Ann. Sc. Nat. Sér. II. xv. (1841) 201 = **Alnus**, Tourn. (Cupulif.).
nepalensis, Spach, l. c. 202 = A. nepalensis.
nitida, Spach, l. c. = A. nitida.

CLETHROSPERUM, Planch. in Hook. Ic. Pl. sub t. 767 (1848), sphalm. = **Clathrospermum**, Planch. (Anonac.).

CLEYERA, Adans. Fam. ii. 224 (1763) = **Polypremum**, Linn. (Loganiac.).

CLEYERA, DC. in Mém. Soc. Phys. Genèv. i. (1822) 412; Prod. i. 524. (1824). *TERNSTROEMIACEAE*, Benth. & Hook. f. i. 183.
Mokof, Adans. Fam. ii. 50 (1763).
Tristylium, Turcz. in Bull. Soc. Nat. Mosc. xxxi. (1858) i. 247.
elegans, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 112.—Ins. Guadelup.
grandiflora, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 284.—Reg. Himal.
grandiflora, Wall. Cat. n. 1461 = ochnacea.
integrifolia, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 112.—Mexic.
japonica, Sieb. & Zucc. Fl. Jap. i. 153. t. 81 = ochnacea.
Lushia, G. Don, Gen. Syst. i. 566 = ochnacea.
Mertensiana, *Sieb. & Zucc. Fl. Jap.* i. 154.—Japon.
mexicana, *Planch. ex Hemsl. Biol. Centr. Am. Bot.* i. 93.—Mexic.
ochnacea, *DC. in Mém. Soc. Phys. Genèv.* i. (1822) 413.—Reg. Himal.
ochnoides, Wall. ex G. Don, Gen. Syst. i. 566 = ochnacea.
serrulata, *Choisy, in Mém. Soc. Phys. Genèv.* xiv. (1855) 112.—Mexic.
siphilitica, *Choisy, l. c.*—Mexic.
theoides, *Choisy, l. c.*—Ind. occ.
Wallichiana, Sieb. & Zucc. Fl. Jap. i. 154 = ochnacea.

CLEYERA, Thunb. Nov. Gen. 68 (1783) = **Ternstroemia**, Linn. f.
dubia, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 308 = T. japonica.
emarginata, Gardn. in Calc. Journ. Nat. Hist. vii. (1847) 447 = T. emarginata.
fragrans, Champ. ex Benth. in Hook. Kew Journ. iii. (1851) 307 = T. japonica.
gymnanthera, Wight & Arn. Prod. 87 = T. japonica.
japonica, Thunb. Nov. Gen. 69 = T. japonica.
lasiopetala, Wight, Illustr. i. 99 = Adinandra lasiopetala.
Millettii, Hook. & Arn. Bot. Beech. Voy. 171 = Adinandra Millettii.
pentapetala, Spreng. Syst. ii. 596 = Saurauja tristyla.
rubiginosa, Spreng. l. c.—Sumatra.

CLEYRIA, Neck. Elem. ii. 183 (1790) = **Dialium**, Linn. (Legumin.).

CLIANTHUS, Banks & Soland. ex G. Don, Syst. ii. 468 (1832). *LEGUMINOSAE*, Benth. & Hook. f. i. 503.
Donia, D. & G. Don, Gen. Syst. ii. 467 (1832).
Eremocharis, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 73 (1849).
Binnendyckianus, *Kurz, in Journ. As. Soc. Beng.* xl. (1871) 51.—Ins. Ceram.
carneus, Lindl. Bot. Reg. (1841) Misc. 2 = Streblorrhiza speciosa.
Dampieri, *A. Cunn. ex Lindl. in Trans. Hort. Soc.* Ser. II. i. (1835) 522.—Austral.
Dampieri, R. Br. in Sturt, Exped. Centr. Austral. ii. 71 = Oxleyi.
Oxleyi, *A. Cunn. ex Lindl. in Trans. Hort. Soc.* Ser. II. i. (1835) 522.—Austral.
pictus, *Endl. in Ann. Wien. Mus.* i. (1836) 185 = Streblorrhiza speciosa.
puniceus, *Banks & Soland. ex Lindl. in Trans. Hort. Soc.* Ser. II. i. (1835) 521.—N. Zel.
speciosus, Steud. Nom. ed. II. i. 381 = Streblorrhiza speciosa.

CLIBADIUM, Linn. Mant. ii. 161 (1771). *COMPOSITAE*, Benth. & Hook. f. ii. 345.
Baillieria, Aubl. Pl. Gui. ii. 804. t. 317 (1775).
Orsinia, DC. Prod. v. 104 (1836).
Oswalda, Cass. in Dict. Sc. Nat. lix. 322 (1829).
Trixis, Sw. Prod. Veg. Ind. Occ. 115 (1788).
acuminatum, *Benth. Bot. Voy. Sulph.* 114.—Ins. Cocos.
agreste, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 157 = Ichthyothera agrestis.
Alexandri, *Griseb. Fl. Brit. W. Ind.* 368.—Jamaic.
angustifolium, Mart. ex Baker, in Mart. Fl. Bras. vi. iii. 154 = lineare.
Armani, Sch. Bip. ex Baker, l. c. 152 = rotundifolium.
asperum, DC. Prod. v. 506 = surinamense.
Badieri, Griseb. Fl. Brit. W. Ind. 367 = surinamense.

CLIBADIUM:—
Barbasco, *DC. Prod.* v. 507.—Venezuela.
caracasanum, *DC. l. c.* 506.—Venezuela.
commelinoides, DC. l. c. = Icthyothera Cunabi.
erosum, *DC. l. c.*—Ind. occ.
foetidum, Alemand, ex Linn. Mant. ii. 294 = surinamense.
fragiferum, *Griseb. Fl. Brit. W. Ind.* 368.—Ind. occ.
havanense, *DC. Prod.* v. 506.—Cuba.
leiocarpum, *Steetz, in Seem. Bot. Voy. Herald*, 152.—Panama.
neriifolium, *DC. Prod.* v. 507.—Venezuela.
pedunculosum, *DC. l. c.*—Guiana.
peruvianum, Poepp. ex DC. l. c. 505 = surinamense.
rotundifolium, *DC. l. c.*—Bras.
Schomburgkii, *Sch. Bip. in Schomb. Fauna & Fl. Guian.* nomen, 940.—Guiana.
surinamense, *Linn. Mant.* ii. 294.—Am. austr.
sylvestre, Baill. Hist. Pl. viii. 307 = surinamense.
terebinthinaceum, *DC. Prod.* v. 506.—Ind. occ.
Trinitatis, DC. l. c. 505 = surinamense.
Vargassi, *DC. l. c.* 506.—Venezuela.
villosum, Benth. Pl. Hartw. 205 = surinamense.

CLIDANTHERA, R. Br. in Sturt, Exped. Centr. Austral. ii. App. 73 (1849) = **Glycyrrhiza**, Linn. (Legumin.).
psoralioides, R. Br. l. c. 74 = G. psoraloides.

CLIDEMIA, D. Don, in Mem. Wern. Soc. iv. (1823) 306. *MELASTOMACEAE*, Benth. & Hook. f. i. 766.
DANCERA, Rafin. Sylva Tellur. 93 (1838).
STAPHIDIUM, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 306.
STEPHANOTRICHUM, Naud. l. c. iv. (1845) 54.
acutiflora, Naud. l. c. xvii. (1852) 371 = Oxymeris acutiflora.
aggregata, D. Don, in Mem. Wern. Soc. iv. (1823) 309 = Henriettella aggregata.
agrestis, D. Don, l. c. = Oxymeris agrestis.
alata, G. Don, ex Loud. Hort. Brit. 174 = Miconia alata.
alpestris, Gardn. in Hook. Lond. Journ. Bot. iv. (1845) 101 = Oxymeris alpestris.
althaeoides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 363 = Octopleura confertiflora.
amblyandra, Naud. l. c. 353 = Oxymeris subscriata.
amygdaloides, DC. Prod. iii. 156 = Ossaea amygdaloides.
angustifolia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 358 = Oxymeris angustifolia.
anisophylla, DC. Prod. iii. 161 = Oxymeris anisophylla.
anisotrichum, Schlecht. in Linnaea, xiii. (1839) 427 = Miconia Berghesiana.
aphanantha, *Sagot, in Ann. Sc. Nat.* Sér. VI. xv. (1883) 327.—Guiana.
aptera, DC. Prod. iii. 161 = Oxymeris aptera.
aristigera, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 366 = Oxymeris aristigera.
asperifolia, Naud. l. c. 342 = Oxymeris asperifolia, Ossaea asperifolia.
astrotricha, DC. Prod. iii. 161 = Miconia astrotricha.
atrata, Spreng. in Flora, xx. (1837) Beibl. ii. 78 = Miconia atrata.
aurea, Cham. in Linnaea, x. (1836) 47 = Oxymeris aurea.
auricoma, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 375 = Leandra aurea.
australis, Cham. in Linnaea, x. (1836) 44 = Oxymeris australis.
barbata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 136.—N. Granat.
barbinervis, Benth. Bot. Voy. Sulph. 95 = Miconia barbinervis.
Benthamiana, Miq. in Linnaea, xviii. (1844) 276 = hirta.
berbiceana, DC. Prod. iii. 160 = Miconia criopoda.
Berterii, Griseb. in Goett. Abh. ix. (1861) 47 = Sagraea plumosa.
bifaria, Rich. ex Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 374 = Oxymeris heterobasis.
biseptena, DC. Prod. iii. 164 = Oxymeris dichotoma.
biseptena, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 171 = Oxymeris Nianga.
biserrata, *DC. Prod.* iii. 158.—Bras.
blepharodes, *DC. l. c.*—Bras.

CLIDEMIA:—
botryophora, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 350 = Oxymeris Nianga.
brachystachya, DC. Prod. iii. 156 = Ossaea brachystachya.
brachystephana, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 135.—N. Granat.
Brackenridgei, A. Gray, Bot. U. St. Expl. Exped. i. 591 = Oxymeris Brackenridgei.
bracteata, DC. Prod. iii. 162 = Miconia bracteata.
bullosa, *DC. l. c.* 158.—Bras.
calvescens, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 91 = Oxymeris calvescens.
campestris, Benth. in Hook. Journ. Bot. ii. (1840) 308 = Miconia campestris.
capillaris, D. Don, in Mem. Wern. Soc. iv. (1823) 309 = Octopleura leptopus.
capillaris, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 184 = Sagraea capillaris.
capitata, Benth. in Hook. Journ. Bot. ii. (1840) 307 = Sagraea capitata.
capitellata, D. Don, in Mem. Wern. Soc. iv. (1823) 310 = dependens.
carassana, DC. Prod. iii. 162 = Oxymeris carassana.
carassana, DC. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 111 = Miconia spondylantha.
cephalophora, Steud. ex Triana, l. c. = Miconia bracteata.
ceramicarpa, DC. Prod. iii. 160 = Miconia spondylantha.
cernua, *D. Don, in Loud. Hort. Brit.* 174.—Peruv.
chaetocalyx, A. Gray, Bot. U. S. Expl. Exped. i. 589. t. 74 = Oxymeris chaetocalyx.
Chaetodon, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 360 = Oxymeris Chaetodon.
Chamissois, Naud. l. c. 375 = Miconia ibaguensis.
chinatlana, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 135.—Mexic.
chrysopogon, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 127 = Miconia barbinervis.
ciliata, D. Don, in Mem. Wern. Soc. iv. (1824) 309 = Calophysa ciliata.
cinerea, Griseb. Cat. Pl. Cub. 97 = Oxymeris cinerea.
cinnamomifolia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 369 = Ossaea cinnamomifolia.
Claussenii, Naud. l. c. 357 = Oxymeris purpurascens.
coccinea, DC. Prod. iii. 162 = Miconia spondylantha.
cognata, Steud. ex Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 308 = hirta.
confertiflora, DC. Prod. iii. 156 = Ossaea confertiflora.
congestiflora, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 344 = Oxymeris congestiflora.
conglomerata, *DC. Prod.* iii. 156.—Guiana.
cordifolia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 361 = Oxymeris cordifolia.
cordigera, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 94 = Oxymeris cordigera.
coriacea, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 368 = Ossaea coriacea.
cornoides, *Griseb. in Goett. Abh.* ix. (1861) 50.—Reg. Argent.
crenata, DC. Prod. iii. 157 = hirta.
crenata, D. Don, in Mem. Wern. Soc. iv. (1823) 308 = Oxymeris crenata.
crossosepala, Griseb. Fl. Brit. W. Ind. 248 = Sagraea heteroneura, crossopetala.
Cruegeriana, Griseb. l. c. 249 = Sagraea salicifolia.
cuneata, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 375 = Oxymeris cuneata.
cyanocarpa, Benth. Bot. Voy. Sulph. 94 = Octopleura quinquenervia.
dasytricha, A. Gray, Bot. U. S. Expl. Exped. i. 588 = Oxymeris carassana.
debilis, Crueg. in Linnaea, xx. (1847) 104 = Sagraea rubra.
debilis, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 361 = Oxymeris debilis.
decurrens, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 127 = Octopleura quinquenervia.
deflexa, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 94 = Oxymeris deflexa.
deflexa, *Bunb. in Proc. Linn. Soc.* i. (1841) 108.—Bras.
dendroides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 367 = Oxymeris dendroides.

CLIDEMIA:—
dentata, *D. Don, in Mem. Wern. Soc.* iv. (1823) 308.—Peruv.
depauperata, DC. Prod. iii. 161 = Oxymeris agrestis.
dependens, *D. Don, in Mem. Wern. Soc.* iv. (1823) 307.—Peruv.
Deppeana, *Steud. Nom.* ed. II. i. 384.—Mexic.
desmantha, Benth. in Hook. Journ. Bot. ii. (1840) 309 = Miconia rhytidophylla.
dichotoma, D. Don, in Mem. Wern. Soc. iv. (1823) 307 = Oxymeris dichotoma.
dispar, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 345 = Oxymeris dispar.
divaricata, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 373 = Oxymeris secundiflora.
diversifolia, DC. Prod. iii. 159 = Octopleura diversifolia.
dolichantha, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 351 = Oxymeris dolichantha.
dolichostachya, Naud. l. c. 359 = Oxymeris dolichostachya.
Drosera, Sagot, l. c. Sér. VI. xv. (1883) 328 = aphanantha.
elegans, *DC. Prod.* iii. 157.—Am. trop.
elegans, D. Don, in Mem. Wern. Soc. iv. (1823) 309 = hirta.
Epibaterium, DC. Prod. iii. 157 = Octopleura Epibaterium.
erostrata, DC. l. c. 160 = Oxymeris erostrata.
erythropogon, *DC. l. c.* 157.—Jamaica.
euphorbioides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 370 = Ossaea euphorbioides.
fallax, Cham. in Linnaea, x. (1836) 41 = Oxymeris fallax.
fallax, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 344 = Oxymeris xantholasia.
fenestrata, Benth. Bot. Voy. Sulph. 94 = umbonata.
fenestrata, Seem. Bot. Voy. Herald, 124 = neglecta.
floribunda, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 349 = Oxymeris floribunda.
foveolata, DC. Prod. iii. 163 = Oxymeris foveolata.
glabrata, *Bunb. in Proc. Linn. Soc.* i. (1841) 108.—Bras.
glabrata, Steud. in Flora, xxvii. (1844) II. 721 = Miconia auriculata.
glabriflora, Presl, Bot. Bemerk. 70 = Miconia decussata.
glomerata, DC. Prod. iii. 163 = Oxymeris Nianga.
guadaloupensis, Griseb. in Goett. Abh. vii. (1857) 215 = Sagraea guadalupensis.
hematostemon, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 348 = Oxymeris aurea.
herbacea, Steud. ex Cogn. in Mart. Fl. Bras. xiv. III. 468 = Aciotis herbacea.
heterobasis, DC. Prod. iii. 164 = Oxymeris heterobasis.
heteroclita, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 365 = japurensis.
heteromalla, D. Don, in Mem. Wern. Soc. iv. (1823) 310 = Sagraea rubra.
heterophylla, *Steud. in Flora*, xxvii. (1844) II. 721.—Bras.
heterotricha, Griseb. Cat. Pl. Cub. 96 = Calycogonium heterotrichum.
hirsuta, *Griseb. in Mem. Am. Acad. N. S.* viii. (1861) 185.—Ind. occ.
hirsuta, Griseb. Fl. Brit. W. Ind. 248 = Ossaea hirsuta.
hirsuta, Macf. Fl. Jamaic. ii. 45 = Ossaea asperifolia.
hirta, *D. Don, in Mem. Wern. Soc.* iv. (1823) 309.—Jamaica.
hirtella, Griseb. in Goett. Abh. ix. (1861) 50 = Ossaea hirtella, Wrightii.
humilis, Miq. in Linnaea, xxii. (1849) 542 = Oxymeris Nianga.
holosericea, *D. Don, ex Loud. Hort. Brit.* 174.—Bras.
inaequalifolia, DC. Prod. iii. 164 = Oxymeris inaequalifolia.
inopogon, Mart. in Flora, xxiv. (1841) Beibl. ii. 37 = Oxymeris inopogon.
intermedia, DC. Prod. iii. 164 = Oxymeris intermedia.
interrupta, Poepp. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 111 = Miconia bracteata.
involucrata, *DC. Prod.* iii. 163.—Guiana
japurensis, *DC. l. c.* 159.—Bras.
lacera, DC. l. c. 164 = Miconia lacera.

CLIDEMIA :—
laevigata, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 94 = Oxymeris laevigata.
Lambertiana, DC. Prod. iii. 164 = Oxymeris crenata.
lanata, DC. l. c. 163 = Miconia lanata.
lanata, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 184 = Ossaea lanata.
lappacea, DC. Prod. iii. 160 = Miconia lappacea.
latifolia, DC. l. c. 159 = Sagraea latifolia.
latifolia, Griseb. Fl. Brit. W. Ind. 247 = Sagraea umbrosa.
laxiflora, *Schlecht. in Linnaea*, xiii. (1839) 426.—Mexic.
leandroides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 355 = Oxymeris purpurascens.
leptoclades, H. Crueg. in Linnaea, xx. (1847) 103 = hirta.
leptopus, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 147 = Ossaea leptopus.
leptostachya, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 172 = Ossaea brachystachya.
leucandra, Wright, ex Griseb. Cat. Pl. Cub. 96 = Sagraea domingensis.
Lima, DC. Prod. iii. 161 = Ossaea Lima.
lineata, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 356 = Oxymeris purpurascens.
longibarbis, DC. Prod. iii. 163 = Oxymeris longibarbis.
lugubris, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 127 = Oxymeris heterobasis.
lutescens, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 347 = Oxymeris crenata.
macrophylla, *Naud. l. c.* 375.—Ins. Cocos.
macrophylla, Seem. Bot. Voy. Herald, 124 = Miconia simplex.
macropora, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 94 = Oxymeris macropora.
maculata, Benth. in Hook. Journ. Bot. ii. (1840) 310 = Miconia auriculata.
marginata, DC. Prod. iii. 156 = Ossaea marginata.
Martiana, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 340 = Oxymeris cuneata.
Martii, Naud. l. c. 375 = Oxymeris ciliata.
melanodesma, Naud. l. c. 353 = Oxymeris melanodesma.
melanotricha, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 136.—Costa Rica.
Miconiastrum, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 352 = Oxymeris Miconiastrum.
miconioides, Benth. in Hook. Journ. Bot. ii. (1840) 310 = Miconia auriculata.
micrantha, *Sagot, in Ann. Sc. Nat.* Sér. VI. xv. (1883) 327.—Guiana.
micropetala, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 367 = Oxymeris micropetala.
microphylla, *D. Don, in Loud. Hort. Brit.* 174.—Am. austr.
microphylla, Griseb. Fl. Brit. W. Ind. 248 = Ossaea microphylla.
microstachya, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 357 = Oxymeris purpurascens.
monticola, Naud. l. c. 341 = Miconia erythrantha.
multiplinervis, Naud. l. c. 358 = Oxymeris multiplinervis.
mutabilis, DC. Prod. iii. 162 = Miconia mutabilis.
naevula, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 137.—Guiana.
neglecta, *D. Don, in Mem. Wern. Soc.* iv. (1823) 307.—Peruv.
nervosa, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 341 = Oxymeris nervosa.
Nianga, DC. Prod. iii. 163 = Oxymeris Nianga.
Nianga, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 344 = Oxymeris xantholasia.
nivea, *D. Don, in Loud. Hort. Brit.* 174.—Am. austr.
novemnervia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 136.—Am. trop.
oaxacensis, Steud. ex Triana, l. c. 92 = Oxymeris subseriata.
obscura, DC. Prod. iii. 162 = Miconia obscura.
octona, *Griseb. in Goett. Abh.* ix. (1861) 50.—Reg. Argent.
oligochaeta, Cham. in Linnaea, x. (1836) 45 = Oxymeris foveolata.
oocarpa, A. Gray, Bot. U. S. Expl. Exped. i. 590 = Oxymeris oocarpa.
ossaeoides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 360 = Oxymeris ossaeoides.

CLIDEMIA :—
oxyura, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 135.—Guiana.
paraguayensis, Steud. in Flora, xxvii. (1844) ii. 721 = Oxymeris paraguayensis.
parasitica, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 135.—Bras.
pauciflora, DC. Prod. iii. 157 = hirta.
penduliflora, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 369 = Oxymeris penduliflora.
penninervis, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 185 = Sagraea penninervis.
pennipilis, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 92 = Oxymeris pennipilis.
petiolaris, Triana, l. c. 135 = Deppeana.
petiolata, DC. Prod. iii. 157 = Sagraea petiolata.
phaeostaphis, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 343 = Oxymeris dichotoma.
phaeotricha, Naud. l. c. 35 = Oxymeris australis.
Pickeringii, A. Gray, Bot. U. S. Expl. Exped. i. 592 = Oxymeris Pickeringii.
pilosa, D. Don, in Mem. Wern. Soc. iv. (1823) 308 = Calophysa pilosa.
plumosa, DC. Prod. iii. 159 = Sagraea plumosa.
polyandra, Benth. Bot. Voy. Sulph. 95 = Miconia Benthamiana.
polystachya, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 347 = Oxymeris polystachya.
pulchra, A. Gray, Bot. U. St. Expl. Exped. i. 591 = Oxymeris pulchra.
purpurascens, DC. Prod. iii. 161 = Oxymeris purpurascens.
purpurea, DC. l. c. 158 = dentata.
purpurea, *D. Don, in Mem. Wern. Soc.* iv. (1823) 308.—Peruv.
pustulata, DC. Prod. iii. 159 = umbonata.
quadrisulca, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 364 = Oxymeris secundiflora.
quinquedentata, Naud. l. c. 375 = Oxymeris quinquedentata.
quinquenodis, Naud. l. c. 370 = Oxymeris quinquenodis.
quintuplinervia, Steud. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 137 = involucrata.
radulaefolia, Benth. in Hook. Journ. Bot. ii. (1840) 309 = Miconia radulaefolia.
rariflora, *Benth. l. c.* 308.—Guiana.
reclinata, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 125 = petiolaris.
Regnellii, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 91 = Oxymeris Regnellii.
Renggeri, Steud. in Flora, xxvii. (1844) 721 = Oxymeris inopogon.
repens, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 136.—N. Granat.
retropila, DC. Prod. iii. 156 = Ossaea retropila.
reversa, DC. l. c. 161 = Oxymeris reversa.
rhamnifolia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 373 = Oxymeris secundiflora.
rhodolasia, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 126 = Miconia lacera.
rhodopogon, DC. Prod. iii. 163 = Oxymeris rhodopogon.
ribesiiflora, Cham. in Linnaea, x. (1836) 38 = Oxymeris ribesiaeflora.
Riedeliana, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 92 = Oxymeris Riedeliana.
rubra, *Mart. Nov. Gen. et Sp.* iii. 152. t. 281.—Am. austr.
rubrinervis, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 185.—Cuba.
rufescens, Griseb. Cat. Pl. Cub. 97 = Ossaea rufescens.
sagittata, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 359 = Oxymeris subseriata.
salicifolia, Naud. l. c. 364 = Ossaea brachystachya.
scabrosa, *Griseb. in Mem. Am. Acad.* N. S. viii. (1861) 184.—Ind. occ.
scandens, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 346 = blepharodes.
secunda, D. Don, in Mem. Wern. Soc. iv. (1823) 308 = Oxymeris secunda.
secundiflora, DC. Prod. iii. 164 = Oxymeris secundiflora.
sericea, D. Don, in Mem. Wern. Soc. iv. (1823) 309 = Sagraea rubra.
serrulata, *Schlecht. in Linnaea*, xiii. (1839) 425.—Mexic.
silvestris, C. Muell. in Walp. Ann. iv. 760 = Oxymeris sylvestris.

CLIDEMIA :—
simplicicaulis, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 346 = Oxymeris simplicicaulis.
solearis, Naud. l. c. 339 = Octopleura macrophylla.
sparsiflora, *Griseb. in Goett. Abh.* vii. (1857) 218.—Reg. Argent.
spicaeformis, DC. Prod. iii. 160 = Miconia spondylantha.
spicata, DC. l. c. 159 = dependens.
spicata, D. Don, in Mem. Wern. Soc. iv. (1823) 308 = Miconia spondylantha.
spicata, Griseb. Cat. Pl. Cub. 96 = neglecta, Heterotrichum octonum.
spondylantha, Mart. ex Triana, in Trans. Linn. Soc. xxviii. (1871) 111 = Miconia spondylantha.
staphidioides, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 344 = Miconia staphidioides.
stenopetala, DC. Prod. iii. 163 = Oxymeris dichotoma.
strigilliflora, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 372 = Oxymeris strigilliflora.
strigillosa, *DC. Prod.* iii. 159.—Am. trop.
subseriata, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 354 = Oxymeris subseriata.
subspicata, Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 126 = dependens.
suffruticosa, O. Berg, ex Triana, in Trans. Linn. Soc. xxviii. (1871) 135.—Bras.
sulcicaulis, Poepp. ex Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 342 = Oxymeris dichotoma.
sulphurea, Naud. l. c. 371 = Oxymeris velutina.
surinamensis, Miq. Comm. Phytog. ii. 82 = umbonata.
Swartzii, Griseb. Fl. Brit. W. Ind. 248, pro parte = Sagraea pilosa.
sylvestris, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 356 = Oxymeris sylvestris.
tetraquetra, Cham. in Linnaea, x. (1836) 42 = Oxymeris tetraquetra.
tiliaefolia, *DC. Prod.* iii. 158.—Bras.
trichodes, DC. l. c. 161 = Miconia trichodes.
trichotoma, Wright, ex Griseb. Cat. Pl. Cub. 96 = Sagraea trichotoma.
triflora, Scheele, in Linnaea, xvii. (1843) 338 = Tibouchina viminea.
trinitensis, Griseb. Fl. Brit. W. Ind. 249 = Sagraea trinitensis.
tristis, D. Don, in Mem. Wern. Soc. iv. (1823) 307 = strigillosa.
umbonata, *DC. Prod.* iii. 158.—Bras.
urceolata, DC. l. c. = neglecta.
urticaefolia, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 354 = Miconia Berghesiana.
Valenzuelana, A. Rich. Ess. Fl. Cub. 552 = Heterotrichum octonum.
velutina, *D. Don, in Loud. Hort. Brit.* 174.—Ind. occ.
verticillata, DC. Prod. iii. 160 = Ossaea verticillata.
violacea, DC. l. c. 162 = Miconia spondylantha.
vittata, *Linden & André, Illustr. Hortic.* xxii. (1875) 152. t. 219.—Peruv.
Weddellii, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 346 = Oxymeris erostrata.
Wrightii, Griseb. in Mem. Am. Acad. N. S. viii. (1861) 185 = Sagraea Wrightii.
xanthocoma, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 352 = Oxymeris xanthocoma.
xantholasia, DC. Prod. iii. 163 = Oxymeris xantholasia.
xantholasia, Gardn. in Hook. Lond. Journ. Bot. ii. (1843) 344 = Oxymeris carassana.
xanthopogon, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 349 = Oxymeris xanthopogon.

CLIDEMIASTRUM, Naud. in Ann. Sc. Nat. Sér. III. xvii. (1852) 87 = **Oxymeris**, DC. (Melast.).
biseptum, Naud. l. c. 88 = Leandra reversa.
mexicanum, Naud. l. c. 87 = O. heterobasis.

CLIFFORTIA, Linn. Gen. ed. I. 311 (1737). *ROSACEAE*, Benth. & Hook. f. i. 624.
MONOGRAPHIDIUM, Presl, Epim. Bot. 202 (1850).
MORILANDIA, Neck. Elem. ii. 99 (1790).
acerosa, Banks, ex Gaertn. Fruct. i. 165 = Nenax acerosa.
arachnoidea, Lodd. Bot. Cab. t. 260 = ruscifolia.
baccans, *Harv. in Harv. & Sond. Fl. Cap.* ii. 303.—Afr. austr.
berberidifolia, Lam. Encyc. ii. 48 = ferruginea.
cinerea, Thunb. Prod. Pl. Cap. 93 (= *Ambraria hirta*).—Afr. austr.

CLIFFORTIA:—
complanata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 301 = serpyllifolia.
concavifolia, *Eckl. & Zeyh. Enum.* 268.—Afr. austr.
cordifolia, Lam. Encyc. ii. 47 = ilicifolia.
cordifolia, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 295.—Afr. austr.
crenata, *Linn. f. Suppl.* 430.—Afr. austr.
cuneata, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 413. Afr. austr.
dentata, Eckl. & Zeyh. Enum. 268 = triloba.
dentata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 299 = gracilis.
dentata, *Willd. Sp. Pl.* iv. 842.—Afr. austr.
Dregeana, *Presl, Bot. Bemerk.* 141.—Afr. austr.
drepanoides, *Eckl. & Zeyh. Enum.* 269.—Afr. austr.
ericaefolia, *Linn. f. Suppl.* 430.—Afr. austr.
eriocephalina, *Cham. in Linnaea*, vi. (1831) 349.—Afr. austr.
falcata, *Linn. f. Suppl.* 431.—Afr. austr.
falcata, Spreng. ex Eckl. & Zeyh. Enum. 269 = drepanoides.
ferruginea, *Linn. f. Suppl.* 429.—Afr. austr.
filicaulis, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 299 = tricuspidata.
filicaulis, *Schlecht. in Linnaea*, ii. (1827) 33.—Afr. austr.
filifolia, *Linn. f. Suppl.* 430.—Afr. austr.
flabellifolia, Sond. ex Harv. & Sond. Fl. Cap. ii. 597 = Myrothamnus flabellifolia.
flexuosa, E. Mey. ex Harv. & Sond. l. c. ii. 296 = ferruginea.
gracilis, *Harv. l. c.* 299.—Afr. austr.
graminea, *Linn. f. Suppl.* 429.—Afr. austr.
grandifolia, *Eckl. & Zeyh. Enum.* 266.—Afr. austr.
hirsuta, Eckl. & Zeyh. l. c. = odorata.
hirta, *Burm. f. Fl. Cap. Prod.* 31 (crr. typ. 27).—Afr. austr.
ilicifolia, *Linn. Sp. Pl.* 1038.—Afr. austr.
intermedia, *Eckl. & Zeyh. Enum.* 266.—Afr. austr.
juniperina, *Linn. f. Suppl.* 430.—Afr. austr.
laricina, E. Mey. ex Meissn. in Hook. Lond. Journ. Bot. ii. (1843) 528 = juniperina.
leptophylla, Eckl. & Zeyh. Enum. 271 = filifolia.
linearifolia, *Eckl. & Zeyh. l. c.* 270.—Afr. austr.
marginata, *Eckl. & Zeyh. l. c.* 269.—Afr. austr.
Meyeriana, *Presl, Epim. Bot.* 202.—Afr. austr.
obcordata, *Linn. f. Suppl.* 429.—Afr. austr.
obliqua, Spreng. Neue Entdeck. ii. 174 = obcordata.
obovata, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 297.—Afr. austr.
octandra, *Cham. in Linnaea*, vi. (1831) 350.—Afr. austr.
odorata, *Linn. f. Suppl.* 431.—Afr. austr.
phylicoides, Eckl. & Zeyh. Enum. 269 = eriocephalina.
polycephala, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 302.—Afr. austr.
polygonifolia, *Linn. Sp. Pl.* 1038.—Afr. austr.
polyphylla, Eckl. & Zeyh. Enum. 268 = serpyllifolia.
propinqua, Eckl. & Zeyh. l. c. = serpyllifolia.
pulchella, *Linn. f. Suppl.* 430.—Afr. austr.
pungens, *Presl, Epim. Bot.* 202.—Afr. austr.
reticulata, Eckl. & Zeyh. Enum. 266 = odorata.
rubricaulis, Presl, Bot. Bemerk. 141 = ilicifolia.
ruscifolia, *Linn. Sp. Pl.* 1038.—Afr. austr.
sarmentosa, *Linn. Mant.* ii. 299.—Afr. austr.
sericea, *Eckl. & Zeyh. Enum.* 269.—Afr. austr.
serpyllifolia, *Cham. & Schlecht. in Linnaea*, ii. (1827) 34.—Afr. austr.
serpyllifolia, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 299 = octandra.
serrata, *Reichb. ex Spreng. Syst.* iv. *Cur. Post.* 209.—Afr. austr.
serrata, Thunb. Prod. Pl. Cap. 93 = ferruginea.
spicata, *Reichb. ex Spreng. Syst.* iv. *Cur. Post.* 209.—Afr. austr.
strobilifera, *Murr. Syst.* ed. XIII. 749.—Afr. austr.
teretifolia, *Linn. f. Suppl.* 430.—Afr. austr.
ternata, Linn. f. l. c. = polygonifolia.
tricuspidata, *Harv. in Harv. & Sond. Fl. Cap.* ii. 299.—Afr. austr.
tridentata, Willd. Sp. Pl. iv. 838 = ruscifolia.
trifoliata, Linn. Sp. Pl. 1038 = polygonifolia.

CLIFFORTIA:—
triloba, *Harv. in Harv. & Sond. Fl. Cap.* ii. 298.—Afr. austr.

CLIFFORTIOIDES, Dryand. ex Hook. Ic. Pl. t. 639 (1844) = **Fagus**, Tourn. (Cupulif.).
oblongata, Dryand. l. c. 639, 673 = F. Solandri.

CLIFTONIA, Banks, ex Gaertn. f. Fruct. iii. 246. t. 225 (1805). *CYRILLEAE*, Benth. & Hook. f. ii. 1226.
Mylocaryum, Willd. Enum. Hort. Berol. 454 (1809).
Walteriana, Fras. ex Endl. Gen. 1413 (1841).
ligustrina, Sims, ex Spreng. Syst. ii. 316 = nitida.
nitida, *Gaertn. f. l. c.* 247.—Am. bor.

CLIMACANDRA, Miq. Pl. Jungh. 199 (1851–55) = **Ardisia**, Sw. (Myrsin.).
littoralis, Kurz, in Journ. As. Soc. Beng. xl. (1871) II. 68 = A. humilis.
multiflora, Miq. Pl. Jungh. 200 = A. humilis.
obovata, Miq. l. c. 199 = A. humilis.
pergamacea, Miq. Fl. Ind. Bat. Suppl. 578 = Embelia pergamacea.
salicifolia, Miq. Pl. Jungh. 200 = A. humilis.

CLIMEDIA, Rafarin, in Rev. Hortic. (1877) 38, sphalm. = **Clidemia**, D. Don (Melast.).

CLINACANTHUS, Nees, in DC. Prod. xi. 511 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1105.
angustus, Nees, l. c. = Isoglossa angusta.
Burmanni, *Nees, l. c.*—Java.

CLINANTHUS, Herb. App. 40 (1821) = **Stenomesson**, Herb. (Amaryll.).
luteus, Herb. l. c. 40 = S. luteum.

CLINHYMENIA, A. Rich. & Gal. in Comptes-Rendus, Acad. Sc. Par. xviii. (1844) 512 = **Cryptarrhena**, R. Br. (Orchid.).
pallidiflora, A. Rich. & Gal. l. c. = Crypt. pallidiflora.

CLINOGYNE, Salisb. in Trans. Hort. Soc. i. (1812) 276. *SCITAMINEAE*, Benth. & Hook. f. iii. 651.
Donax, Lour. Fl. Cochinch. 11 (1790).
dichotoma, *Salisb. l. c.*—Ind. or.; Malaya.
filipes, *Benth. & Hook. f. Gen.* iii. 651.—Afr. trop.
grandis, *Benth. & Hook. f. l. c.*—Burma; Malaya.
virgata, *Benth. & Hook. f. l. c.*—Ind. or.

CLINOPODIUM, Linn. Syst. ed. I (1735) = **Calamintha**, Tourn. (Labiat.).
aegyptiacum, Mill. Gard. Dict. ed. VIII. n. 6 = Calamintha Clinopodium.
albidum, Vell. Fl. Flum. 242; vi. t. 6 = Hyptis paludosa.
angustifolium, Roxb. Hort. Beng. 44, nomen (quid?).—Hab.?
arundanum, Nym. Syll. 101 = Calamintha Clinopodium.
arvense, Vell. Fl. Flum. 243; vi. t. 9.—Bras.
asiaticum, Lour. Fl. Cochinch. ii. 374 (= *Melissa asiatica*) (= Anisochili sp.).—Cochinch.
atropurpureum, Hort. Hafn. ex Steud. Nom. ed. II. i. 386 = Calamintha Clinopodium.
brasilicum, Vell. Fl. Flum. 241; vi. t. 3.—Bras.
capitatum, Sw. Prod. Veg. Ind. Occ. 88 = Hyptis capitata.
carolinianum, Mill. Gard. Dict. ed. VIII. n. 5 = Calamintha caroliniana.
Chamaedrys, Vahl, Symb. Bot. iii. 77 = Marsypianthes hyptoides.
cordatum, Vell. Fl. Flum. 243; vi. t. 8.—Bras.
fruticosum, Forsk. Fl. Aegypt. Arab. 107 = Otostegia scariosa.
glabrescens, Pomel, Nouv. Mat. Fl. Atl. 122.—Algeria.
grandiflorum, Gueldenst. ex Bieb. Fl. Taur. Cauc. iii. 399 = Stachys grandiflora.
humile, Mill. Gard. Dict. ed. VIII. n. 4.—Am. bor.

CLINOPODIUM:—
imbricatum, Vell. Fl. Flum. 242; vi. t. 5 = Hyptis pectinata.
incanum, Linn. Sp. Pl. 588 = Pycnanthemum incanum.
longicaule, Benth. in Wall. Pl. As. Rar. i. 66 = Calamintha longicaulis.
martinicense, Jacq. Enum. Pl. Carib. 25 = Leucas martinicensis.
origanifacie, Gilib. Fl. Lituan. i. 76 = Calamintha Clinopodium.
origanifolium, Labill. Ic. Pl. Syr. dec. iv. 14. t. 9 = Calamintha origanifolia.
plumosum, Sieber, in Flora, v. (1822) 242 = Calamintha Clinopodium.
repens, Roxb. Hort. Beng. 44; Fl. Ind. iii. 13 = Calamintha umbrosa.
repens, Vell. Fl. Flum. 242; vi. t. 7 = Peltodon radicans.
rugosum, Linn. Sp. Pl. 588 = Hyptis radiata.
variegatum, Hort. ex Steud. Nom. ed. II. i. 386 = Calamintha Clinopodium.
verticillatum, Vell. Fl. Flum. 242; vi. t. 4.—Bras.
villosum, Noé, in Bull. Soc. Bot. Fr. ii. (1855) 580 = Calamintha Clinopodium.
vulgare, Linn. Sp. Pl. 587 = Calamintha Clinopodium.

CLINOSTIGMA, H. Wendl. in Bonplandia, x. (1862) 196. *PALMAE*, Benth. & Hook. f. iii. 894.
Billardierii, Becc. Malesia, i. (1877) 41 = Microkentia Billardierii.
bractealis, Becc. l. c. 40 = Cyphokentia bractealis.
Deplanchei, Becc. l. c. 41 = Microkentia Deplanchei.
eriostachys, Becc. l. c. = Microkentia eriostachys.
Humboldtiana, Becc. l. c. 40 = Cyphokentia Humboldtiana.
macrostachya, Becc. l. c. = Cyphokentia macrostachya.
Moorianum, *F. Muell. Fragm.* viii. 235.—Austral.
Pancheri, Becc. Malesia, i. (1877) 40 = Microkentia Pancheri.
robusta, Becc. l. c. = Cyphokentia robusta.
samoënse, *H. Wendl. in Bonplandia*, x. (1862) 196.—Ins. Samoa.
surculosa, Becc. Malesia, i. (1877) 41 = Microkentia surculosa.
vaginata, Becc. l. c. = Cyphokentia vaginata.

CLINOSTYLIS, Hochst. in Flora, xxvii. (1844) 26 = **Gloriosa**, Linn. (Liliac.).
speciosa, Hochst. l. c. = G. abyssinica.

CLINTONIA, Dougl. ex Lindl. Bot. Reg. t. 1241 (1829) = **Downingia**, Torr. (Campanulac.).
Bergiana, Cham. in Linnaea, vii. (1832) 217 = Lobelia Bergiana.
corymbosa, DC. Prod. vii. 347 = D. elegans.
elegans, Dougl. ex Lindl. Bot. Reg. t. 1241 = D. elegans.
pulchella, Lindl. l. c. t. 1909 = D. pulchella.
pusilla, G. Don, Gen. Syst. iii. 718.—Chili.

CLINTONIA, Rafin. in Am. Monthly Mag. (1818) 266; Journ. Phys. lxxxix. (1819) 102. *LILIACEAE*, Benth. & Hook. f. iii. 832.
Hylocharis, Tiling, ex Regel & Tiling, in Nouv. Mém. Soc. Nat. Mosc. xi. (1859) 123.
Xeniatrum, Salisb. Gen. Pl. Fragm. 58 (1866).
Aitoni, Rafin. Atl. Journ. 120 = borealis.
alpina, *Kunth, Enum. Pl.* v. 159.—Reg. Himal.
Andrewsiana, *Torr. in Pacif. Rail. Rep.* iv. 150.—Calif.
Andrewsii, Wood, in Proc. Acad. Philad. (1868) 174 = Andrewsiana.
angustifolia, *Rafin. New Fl. Am.* ii. 79.—Am. bor.
biflora, Rafin. l. c. 78.—Am. bor.
biumbella, Rafin. l. c. 81 = borealis.
borealis, *Rafin. Atl. Journ.* 120.—Am. bor.
ciliata, Rafin. in Am. Monthly Mag. (1819) 357 = borealis.
decantha, *Rafin. Atl. Journ.* 120.—Am. bor.
falcata, *Rafin. New Fl. Am.* ii. 79.—Am. bor.
fulva, *Rafin. l. c.* 82.—Am. bor.
glomerata, *Rafin. l. c.* 83.—Am. bor.
latifolia, *Rafin. l. c.* 81.—Am. bor.
multiflora, Beck, Bot. 358 = umbellata.

CLINTONIA:—
multiflora, *Rafin. Atl. Journ.* 120.—Am. bor.
mutans, Rafin. Ann. Nat. 16 = borealis.
nutans, *Rafin. Med. Fl.* ii. 85 = praec.
odorata, Rafin. Ann. Nat. 16 = umbellata.
ophioglossoides, *Rafin. New Fl. Am.* ii. 80.—Am. bor.
parviflora, *Rafin. Ann. Nat.* 16.—Am. bor.
podanisia, *Rafin. l. c.*—Am. bor.
triflora, *Rafin. New Fl. Am.* ii. 79.—Am. bor.
udensis, *Trautv. & Mey. in Middend. Reise* (*Fl. Ochot.* 92).—Sibiria; Japon.
umbellata, *Torr. Fl. N. York*, ii. 301.—Am. bor.
undulata, Rafin. New Fl. Am. ii. 81 = borealis.
uniflora, *Kunth, Enum. Pl.* v. 159.—Am. bor. occ.

CLIOCARPUS, Miers, in Ann. Nat. Hist. Ser. II. iv. (1849) 141 = **Solanum**, Tourn.
didymus, Miers, l. c. xv. (1855) 204 = S. didymum.
eriocalyx, Miers, l. c. = S. trichocalyx.
Gardneri, Miers, l. c. iv. (1849) 141.—Bras.
megalochiton, Miers, l. c. xv. (1855) 203 = S. megalochiton.

CLIOCOCCA, Bab. in Proc. Linn. Soc. i. (1841) 90; et Trans. Linn. Soc. xix. (1842) 33. t. 3 = **Linum**, Tourn.
tenuifolia, Bab. in Trans. Linn. Soc. xix. (1845) 33, 34 = L. selaginoides.

CLIPTERIA, Rafin. New Fl. Am. ii. 44 (1836) = **Eclipta**?, Linn. (Compos.).
dichotoma, Rafin. l. c.—Am. bor.

CLISTANTHUS, Muell. Arg. in DC. Prod. xv. II. 1025 (1866) = CLISTRANTHUS, Poit. = **Pera**, Mutis (Euphorb.).

CLISTAX, Mart. Nov. Gen. et Sp. iii. 26 (1829). *ACANTHACEAE*, Benth. & Hook. f. ii. 1120.
CORYTHACANTHUS, Nees, in Lindl. Introd. Nat. Syst. ed. II. 444 (1836).
brasiliensis, *Mart. l. c.*—Bras.
speciosus, *Nees, in Mart. Fl. Bras.* ix. 14.—Bras.

CLISTRANTHUS, Poit. ex Baill. Etud. Gén. Euph. 434 (1858) = **Pera**, Mutis (Euphorb.).

CLITANDRA, Benth. in Hook. Niger Fl. 445 (1849). *APOCYNACEAE*, Benth. & Hook. f. ii. 692.
cirrhosa, *Radlk. in Bremen Abh.* viii. (1883) 400.—Afr. trop.
cymulosa, *Benth. in Hook. Niger Fl.* 445.—Afr. trop.

CLITANTHES, Herb. in Bot. Reg. (1839) Misc. 87 = **Chlidanthus**, Herb. & **Stenomesson**, Herb. (Amaryll.).
humilis, Herb. l. c. = Stenomesson humile.
lutea, Herb. l. c. = Chlidanthus fragrans.
Macleanica, Herb. l. c.—Peruv.

CLITANTHUM, Benth. & Hook. f. Gen. iii. 733 (1880–83) = **Chlidanthus**, Herb. (Amaryll.).

CLITOCYAMOS, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 128, 133 = **Ipomoea**, Linn. (Convolvulac.).
pinnatifidus, St. Lag. ll. cc. = I. Quamoclit.

CLITORIA, Linn. Gen. Pl. ed. I. 216 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 527.
MARTIA, Leandr. Sacr. in Denkschr. Akad. Muench. vii. (1821) 233. t. 12.
MARTIUSIA, Schult. Mant. i. 69 (1822).
NAUCHEA, Descourt. in Mém. Soc. Linn. Par. iv. (1825) 7. t. 1.
NEUROCARPUM, Desv. Journ. Bot. i. (1813) 119; iii. (1814) 75.
RHOMBIFOLIUM, Rich. ex DC. Prod. ii. 235 (1825).
TERNATEA, Tourn. ex Linn. Syst. ed. I. (1735); H. B. & K. Nov. Gen. et Sp. vi. 415 (1823).
VEXILLARIA, Rafin. in Am. Monthly Mag. (1818) 268.
acuminata, Benth. in Ann. Wien. Mus. i. (1838) 115 = amazonum.

CLITORIA:—
acuminata, R. Grah. in Wall. Cat. n. 5346 = Mariana.
alabamensis, Bertol. Misc. Bot. ix. 12 = virginiana.
alba, G. Don, Gen. Syst. ii. 215.—Ins. S. Thomas.
amazonum, *Mart. in Ann. Wien. Mus.* ii. (1838) 115.—Bras.
amoena, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) 24 = arborescens.
amoena, Roth, in Roem. Arch. i. III. 42 = Centrosema virginianum.
angustifolia, H. B. & K. Nov. Gen. et Sp. vi. 417 = Centrosema angustifolium.
angustifolia, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 29 = Galactis Martii.
aquilupiensis, Raeusch. Nom. ed. III. 210 = quadalupensis.
arborea, Benth. in Ann. Wien. Mus. ii. (1838) 115 = Hoffmanseggii.
arborescens, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 302.—Ins. Trinitat.
australis, *Benth. Fl. Austral.* ii. 242.—Austral.
Berteriana, DC. Mém. Lég. vi. 239 = Periandra Berteriana.
biflora, *Dalz. in Hook. Kew Journ.* ii. (1850) 35.—Ind. or.
brachystegia, *Benth. Bot. Voy. Sulph.* 84.—Ecuador.
bracteata, Poir. Encyc. Suppl. ii. 301 = C. Ternatea.
brasiliana, Linn. Sp. Pl. 753 = Centrosema brasilianum.
brasiliana, Vell. Fl. Flum. 312; vii. t. 129 = Canavalia ensiformis.
Broussonetii, Balb. Cat. Taur. (1813) 26 = Cologania Broussonetii.
cajanifolia, *Benth. in Mart. Fl. Bras.* xv. I. 121.—Reg. trop.
calcarata, L'Hérit. ex DC. Prod. ii. 235 = Centrosema Plumieri.
calcarigera, Salisb. Parad. Lond. t. 51 = Centrosema virginianum.
capitata, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 111.—Ind. occ.
coccinea, Schrad. in Goett. Gel. Anz. i. (1821) 717 = Periandra coccinea.
densiflora, *Benth. in Journ. Linn. Soc.* ii. (1858) 41.—Am. trop.
erecta, *Roxb. Hort. Beng.* 56.—Am. trop.
falcata, Lam. Encyc. ii. 51 = glycinoides.
falcata, Nees, in Flora, iv. (1821) 329 = Periandra coccinea.
flagellaris, *Benth. in Journ. Linn. Soc.* ii. (1858) 39.—Am. trop.
fluminensis, Vell. Fl. Flum. 312; vii. t. 128 = Centrosema Plumieri.
formosa, H. B. & K. Nov. Gen. et Sp. vi. 417 = Centrosema brasilianum.
fulgens, Paxt. Mag. Bot. xi. (1844) 121 (= *Collaea scarlatina*).—Bras.
Galactia, Linn. Sp. Pl. ed. II. 1026 = Galactia pendula.
gemina, Vell. Fl. Flum. 313; vii. t. 130 = Centrosema virginianum.
glabella, Desf. Tabl. ed. I. 192 = Galactia glabella.
gladiata, Schrank, in Syll. Ratisb. i. (1824) 229 = Centrosema virginianum.
glomerata, *Griseb. Cat. Pl. Cub.* 74.—Cuba.
glycinoides, *DC. Prod.* ii. 234.—Bras.
Grahami, Steud. Nom. ed. II. i. 386 = Mariana.
grandiflora, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) II. 189.—Mexic.
guadelupensis, vide quadelupensis.
guianensis, *Benth. in Journ. Linn. Soc.* ii. (1858) 40.—Am. trop.
heterophylla, *Lam. Encyc.* ii. 51.—Ind. or.
Hoffmanseggii, *Benth. in Journ. Linn. Soc.* ii. (1858) 43.—Am. trop.
insulana, Vell. Fl. Flum. 313; vii. t. 131 = Centrosema brasilianum.
javanica, Miq. Fl. Ind. Bat. i. I. 226 = macrophylla.
javitensis, *Benth. in Journ. Linn. Soc.* ii. (1858) 42.—Mexic.
lactescens, Linn. Syst. ed. X. 1172 = Galactia pendula.
lasciva, *Boj. Hort. Maurit.* 92, nomen; *Benth. in Ann. Wien. Mus.* ii. (1838) 114.—Madag.
laurifolia, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 411 = Centrosema vexillatum.
laurifolia, Poir. Encyc. Suppl. ii. 301 = cajanifolia.
leptostachya, *Benth. in Journ. Linn. Soc.* ii. (1858) 43.—Am. trop.
macrophylla, *Wall. Cat.* n. 5345.—Burma.

CLITORIA:—
Mariana, *Linn. Sp. Pl.* 753.—Ind. or.; Burma; Am. bor.
Mariana, Moç. & Sesse, ex DC. Prod. ii. 238 = Galactia radicata.
mexicana, Link, Enum. Hort. Berol. ii. 235 = Mariana.
micrantha, *Guill. & Perr. Fl. Seneg. Tent.* i. 190.—Afr. trop.
micrantha, Scop. Delic. Insub. ii. 1 = Galactia longifolia.
micrantha, Sm. ex Steud. Nom. ed. II. i. 386 = Centrosema virginianum.
mucronulata, *Benth. in Ann. Nat. Hist.* iii. (1839) 435.—Bras.
multiflora, *Mart. & Gal. in Bull. Acad. Brux.* x. II. (1843) 188.—Mexic.
multiflora, Sw. Prod. Veg. Ind. Occ. 106 = Vilmorinia multiflora.
nana, *Benth. in Journ. Linn. Soc.* ii. (1858) 40.—Am. trop.
oblonga, Hassk. in Flora, xxv. (1842) II. Beibl. 48 = Centrosema virginianum.
occidentalis, *Sweet, Hort. Brit.* ed. II. 140.—Ind. occ.
parviflora, *Rafin. Atl. Journ.* 147.—Am. bor.
pedunculata, *Boj. ex Benth. in Ann. Wien. Mus.* ii. (1838) 114.—Madag.
pedunculata, *Micheli, in Kjoeb. Vidensk. Meddel.* (1875) 77.—Bras.
Phyrne, Steud. Nom. ed. I. 208, 349 = Galactia sericea.
pilosula, Wall. Cat. n. 5347 = C. Ternatea.
Plumieri, *Turp. in Pers. Syn.* ii. 303 = Centrosema Plumieri.
Poitaei, Benth. in Ann. Nat. Hist. iii. (1839) 434 = arborescens.
Poiteaui, DC. Prod. ii. 234 = arborescens.
polyphylla, Poir. Encyc. Suppl. ii. 300 = Barbieria polyphylla.
polystachya, *Benth. Pl. Hartw.* 60.—Mexic.
portobellensis, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 119.—Panama.
pudica, *Steud. Nom.* ed. II. i. 386.—Cuba.
quadalupensis, *Scop. Delic. Insub.* ii. 4.—Ins. Guadalup.
racemosa, *G. Don, Gen. Syst.* ii. 215; *Benth. in Ann. Wien. Mus.* ii. (1838) 115.—Bras.
rubiginosa, Juss. ex Pers. Syn. ii. 312 = glycinoides.
rubiginosa, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 29 = Galactia Neesii.
rufescens, *Benth. in Journ. Linn. Soc.* ii. (1858) 39.—Bras.
Schiedeana, Schlecht. in Linnaea, xii. (1838) 284 = Centrosema pubescens.
Selloi, *Benth. in Journ. Linn. Soc.* ii. (1858) 42.—Am. trop.
simplicifolia, *Benth. l. c.* 40.—Am. trop.
sinuata, Desv. in Ann. Sc. Nat. Sér. I. ix. (1826) 120 = Centrosema sinuatum.
speciosa, *Cav. Desc.* 182.—Mexic.
spectabilis, Salisb. Prod. 336 = C. Ternatea.
stipularis, *Benth. in Journ. Linn. Soc.* ii. (1858) 41.—Am. trop.
Ternatea, *Linn. Sp. Pl.* 753.—Reg. trop.
ternatensium, Crantz, Inst. ii. 59 = C. Ternatea.
tetragona, Poir. ex DC. Prod. ii. 236 = glycinoides.
tomentosa, Mart. in Nov. Act. Nat. Cur. xii. (1824) 30 = Galactia tomentosa.
tristis, *Tenore, Cat. Hort. Neap.* (1845) 82.—Hab.?
vicioides, Nees & Mart. in Nov. Act. Nat. Cur. xii. (1824) 28 = Coursetia vicioides.
virginiana, Linn. Sp. Pl. 753 = Centrosema virginianum.
viridiflora, Bouton, ex Hook. Ic. Pl. t. 152 = Dolichos axillaris.
zanzibarensis, *Vatke, in Oestr. Bot. Zeitschr.* xxviii. (1878) 261.—Afr. trop.
zoophthalmum, Linn. Syst. ed. X. 1172 = Dolichos urens?

CLITORIASTRUM, Heist. Syst. 9 (1748) = **Clitoria**, Linn.? (Legumin.).

CLIVIA, Lindl. Bot. Reg. t. 1182 (1828). *AMARYLLIDEAE*, Benth. & Hook. f. iii. 729.
IMATOPHYLLUM, Hook. Bot. Mag. t. 2856 (1828).
Gardeni, *Hook. Bot. Mag.* t. 4895.—Afr. austr.
miniata, *Regel, Gartenfl.* (1864) 131.—Afr. austr.
nobilis, *Lindl. Bot. Reg.* t. 1182.—Afr. austr.

CLOANTHE, Nees, in R. Br. Verm. Schr. i. 78 (1825) = **Chloanthes**, R. Br. (Verbenac.).

CLOËZIA, Brongn. & Gris, in Bull. Soc. Bot. Fr. x. (1863) 576. *MYRTACEAE*, Benth. & Hook. f. i. 709.
BALLARDIA, Montr. in Mém. Acad. Lyon, x. (1860) 204.
MOORIA, Montr. l. c. (1860) 207.
buxifolia, *Brongn. & Gris, l. c.* 577.—N. Caled.
cancscens, *Brongn. & Gris, l. c.*—N. Caled.
Deplanchei, *Brongn. & Gris, l. c.*—N. Caled.
floribunda, *Brongn. & Gris, l. c.*—N. Caled.
ligustrina, *Brongn. & Gris, l. c.*—N. Caled.
sessilifolia, *Brongn. & Gris, l. c.*—N. Caled.

CLOMENA, Beauv. Agrost. 28. t. 7. f. 10 (1812) = **Muehlenbergia**, Schreb. (Gramin.).
peruviana, Beauv. l. c. = M. Clomena.

CLOMENOCOMA, Cass. in Dict. Sc. Nat. ix. 416 (1817) = **Dysodia**, Cav. (Compos.).
aurantia, Cass. l. c. = D. appendiculata.
montana, Benth. Pl. Hartw. 86 = D. grandiflora.
pinnata, DC. Prod. v. 641 = D. pubescens.

CLOMENOLEPIS, Cass. in Dict. Sc. Nat. iii. Suppl. 64 (1816), nomen (Compos.).

CLOMIUM, Adans. Fam. ii. 116 (1763) = **Carduus**, Tourn. (Compos.).

CLOMOPANUS, Steud. Nom. ed. II. i. 386 (1840) = seq.

CLOMPANUS, Aubl. Pl. Gui. ii. 773 (1775) = **Lonchocarpus**, Humb. & Bonpl. (Legumin.).
paniculata, Aubl. l. c.—Guiana.

CLOMPANUS, Rafin. Sylva Tellur. 73 (1838) = **Sterculia**, Linn.
molucanus, Rafin. l. c.—Ins. Molucc.

CLONODIA, Griseb. in Mart. Fl. Bras. xii. 1. 26 (1858). *MALPIGHIACEAE*, Benth. & Hook. f. i. 253.
verrucosa, *Griseb. l. c.*—Bras.

CLONOSTACHYS, Klotzsch, in Wiegm. Archiv, vii. (1841) 185 = **Sebastiania**, Spreng. (Euphorbiac.).

CLOSASCHIMA, Korth. Verh. Nat. Gesch. Bot. 139 (1842) = **Laplacea**, H. B. & K. (Ternstroemiac.).
marginata, Korth. l. c. 141 = L. marginata.
ovalis, Korth. l. c. 140 = L. ovalis.

CLOSIA, Remy, in C. Gay, Fl. Chil. iv. 119. t. 46. f. 2 (1849). *COMPOSITAE*, Benth. & Hook. f. ii. 404.
anthemoides, *Phil. Fl. Atac.* 31.—Chili.
catula, *Remy, in C. Gay, Fl. Chil.* iv. 120.—Chili.
chilensis, *Regel & Koern. Ind. Sem. Hort. Petrop.* (1857) 39.—Chili.
discoidea, *Phil. Fl. Atac.* 31.—Chili.
elata, *Phil. l. c.*—Chili.
pusilla, *Phil. l. c.*—Chili.

CLOSIROSPERMUM, Neck. Elem. i. 54 (1790) = **Crepis**, Linn. (Compos.).

CLOSTERANDRA, Boiv. ex Belang. Voy. Indes Or. t. absque num. (1839) = **Papaver**, Tourn.
minor, Boiv. l. c. = P. Belangeri.

CLOSTERANTHERA, Walp. Rep. i. 115 (1842) = praec.

CLOWESIA, Lindl. Bot. Reg. xxix. (1843) Misc. 25. t. 39 = **Catasetum**, Rich. (Orchid.).
rosea, Lindl. l. c. = Catasetum roseum, *Reichb. f.*

CLUACENA, Rafin. Sylva Tellur. 104 (1838) = **Myrtus**, Tourn.
myrsinoides, Rafin. l. c. = M. myrsinoides.
vaccinioides, Rafin. l. c. = M. vaccinioides, *H. B. & K.*

CLUERIA, Rafin. Fl. Tellur. iii. 87 (1836) = **Eremostachys**, Bunge (Labiat.).
laciniata, Rafin. l. c. = E. laciniata.

CLUGNIA, Comm. ex DC. Syst. i. 433 (1818) = **Wormia**, Rottb. (Dilleniac.).
volubilis, Comm. ex Steud. Nom. ed. I. 208 = W. madagascariensis.
volupis, Comm. ex DC. Syst. i. 433 = W. madagascariensis.

CLUSIA, Linn. Gen. ed. I. 344 (1737). *GUTTIFERAE*, Benth. & Hook. f. i. 170.
ANDROSTYLIUM, Miq. in Nat. Verh. Wet. Haarl. vii. (1851) 93.
ARRUDEA, St. Hil. Fl. Bras. Mer. i. 318. t. 66 (1827).
ASTHOTHECA, Miers, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 254.
BIROLIA, Rafin. Sylva Tellur. 163 (1838).
CAHOTIA, Karst. in Linnaea, xxviii. (1856) 448.
COCHLANTHERA, Choisy, Gutt. Ind. 46. t. 3 (1823).
ELWERTIA, Rafin. Sylva Tellur. 163 (1838).
FIRKEA, Rafin. l. c. (1838).
HAVETIA, H. B. & K. Nov. Gen. et Sp. v. 203 (1821).
ICOSTEGIA, Rafin. Sylva Tellur. 163 (1838).
LIPOPHYLLUM, Miers, in Trans. Linn. Soc. xxi. (1855) 251. t. 26.
OXYSTEMON, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiii. (1860) 314.
POLYTHECANDRA, Planch. & Triana, l. c. (1860).
QUAPOYA, Aubl. Pl. Gui. ii. 897 (1775).
SMITHIA, Scop. Introd. 322 (1777).
TRIPLANDRON, Benth. Bot. Voy. Sulph. 73. t. 38 (1844).
XANTHE, Schreb. Gen. 710 (1791).
acuminata, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 358.—Panama.
acuminata, *Spreng. Syst.* ii. 599.—Porto Rico.
alata, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 361.—N. Granat.
alba, Choisy, in Planch. & Triana, l. c. 326 = palmicida.
alba, *Jacq. Enum. Pl. Carib.* 34.—Ind. occ.
alba, Kunth, ex Griseb. Fl. Brit. W. Ind. 107 = rosea.
amazonica, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 358.—Bras.
bicolor, *Mart. Nov. Gen. et Sp.* iii. 164.—Bras.
Brongniartiana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 355.—Ind. occ.
Cambessedii, *Planch. & Triana, l. c.* 360.—Bras.
Candelabrum, *Planch. & Triana, l. c.* 348.—Am. trop.
cassinioides, *Planch. & Triana, l. c.* 369.—Peruv.
Couleti, Duchass. ex Planch. & Triana, l. c. 333 = minor.
crassifolia, *Planch. & Triana, l. c.* 351.—Am. trop.
Cruiva, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 317.—Bras.
cuneata, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 368.—Guiana.
decussata, *Ruiz & Pav. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 320, 345.—Peruv.
Dncu, *Benth. Pl. Hartw.* 126.—Peruv.
elliptica, *H. B. & K. Nov. Gen. et Sp.* v. 199.—Peruv.
eugenioides, *Planch. & Linden, ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 319, 328.—N. Granat.
flava, *Jacq. Enum. Pl. Carib.* 34.—Jamaica.
flava, Willd. ex Steud. Nom. ed. II. i. 387 = Hoffmannseggiana.
fluminensis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 349.—Bras.
Fockeana, *Miq. in Hoev. & De Vriese, Tijdschr.* x. (1843) 82.—Guiana.
fragrans, *Gardn. in Hook. Lond. Journ. Bot.* ii. (1843) 334.—Bras.
ganabarica, Casar. Nov. Stirp. Bras. Decad. 60 = Sellowiana.
Gardneri, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 319, 328.—Bras.
Gaudichaudii, *Cambess. in St. Hil. Fl. Bras. Mer.* i. 318.—Bras.
Gaudichaudii, Choisy, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiii. (1860) 359 = Sellowiana.
grandiflora, *Splitg. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 101.—Guiana.
guatemalensis, *Hemsl. Diag. Pl. Nov.* 2.—Am. centr.
Gundlachi, *Stahl, Estud. Fl. Puerto-Rico, foll.* ii. 122.—Ind. occ.
havetioides, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 368.—Ind. occ.
Hilariana, *Schlecht. in Linnaea*, viii. (1833) 181.—Bras.
Hoffmannseggiana, *Schlecht. l. c.* 185.—Bras.
Ildefonsiana, A. Rich. ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiii. (1860) 359 = Sellowiana.

CLUSIA:—
insignis, *Mart. Nov. Gen. et Sp.* iii. 164.—Bras.
lanceolata, *Cambess. in St. Hil. Pl. Bras. Mer.* i. 318.—Bras.
latipes, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 365.—N. Granat.
laurifolia, *Planch. & Triana, l. c.* 344.—N. Granat.
laxiflora, *D. Dietr. Syn. Pl.* iv. 866.—Peruv.
leprantha, *Mart. Nov. Gen. et Sp.* iii. 165.—Bras.
leprata, *Mart. l. c.*—Bras.
leucantha, *Schlecht. in Linnaea*, viii. (1833) 186.—Bras.
Lhotzkyana, *Schlecht. in Linnaea*, viii. (1833) 184.—Bras.
Lhotzskyana, Choisy, ex Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiii. (1860) 319, pro parte = Hilariana.
lincata, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 345.—N. Granat.
longifolia, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 113 = Micranthera clusioides.
loranthacea, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 346.—N. Granat.
macrocarpa, *Spreng. Syst.* ii. 599.—Guiana.
major, Linn. Sp. Pl. 509 = alba.
mammosa, *Casar. Nov. Stirp. Bras. Decad.* 60.—Bras.
Mangle, *Rich. ex Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 369.—Ins. Guadelup.
maxima, Rich. ex Planch. & Triana, l. c. 325 = grandiflora.
micrantha, Choisy, ex Planch. & Triana, l. c. 259 = Sellowiana.
microcarpa, *Spreng. Syst.* ii. 598.—Guiana.
microstemon, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 331.—Bras.
minor, *Linn. Sp. Pl.* 510.—Guatemala.
multiflora, *H. B. & K. Nov. Gen. et Sp.* v. 200.—N. Granat.
myriandra, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 332.—Guiana.
nemorosa, *G. F. W. Mey. Prim. Fl. Esseq.* 203.—Guiana.
nutans, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 373.—N. Granat.
odorata, *Seem. Bot. Voy. Herald*, 89.—Am. centr.
organensis, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 349.—Bras.
Orizabae, *Hemsl. Diag. Pl. Nov.* 3.—Mexic.
ovigera, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 354.—Mexic.
palmicida, *Rich. ex Planch. & Triana, l. c.* 326.—Guian.
Panapanari, Choisy, in DC. Prod. i. 559 = macrocarpa.
parviflora, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 976.—Ins. Margarita.
Pavonii, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 363.—Peruv.
pedicellata, *Forst. f. Prod.* 74. n. 390.—N. Caled.
pentarhyncha, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 365.—N. Granat.
petiolaris, *Planch. & Triana, l. c.* 373.—N. Granat.
Plumerii, *Planch. & Triana, l. c.* 340.—Ins. S. Doming.
popayenensis, *Planch. & Triana, l. c.* 368.—N. Granat.
pratensis, Seem. Bot. Voy. Herald, 89 = minor.
pseudochina, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 12.—Peruv.
pseudo-havetia, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 366.—Peruv.
pseudo-mangle, *Planch. & Triana, l. c.* 370.—Peruv.
Puat, *Montr. in Mém. Acad. Lyon*, x. (1860) 188.—N. Caled.
Quapoya, Choisy, in DC. Prod. i. 559 = microcarpa.
radicans, *Pav. ex Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 374.—Peruv.
renggerioides, *Planch. & Triana, l. c.* 350.—Bras.
retusa, Poir. Encyc. v. 183 = rosea.
rosaeflora, *Planch. & Linden, ex Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 343.—Venezuela.
rosea, *Jacq. Enum. Pl. Carib.* 34.—Ins. S. Doming.
rosea, Cambess. in St. Hil. Fl. Bras. Mer. i. 316 = Hilariana.
rupicola, *Casar. Nov. Stirp. Bras. Decad.* 60.—Bras.
Seemanni, *Planch. & Triana, in Ann. Sc. Nat.* Ser. IV. xiii. (1860) 347.—Panama.

CLUSIA :—
Sellowiana, *Schlecht. in Linnaea*, viii. (1833) 183.—Bras.
sessiliflora, *Poir. Encyc.* v. 183.—Madag.
sessilis, *Forst. f. Prod.* 74. n. 391.—Ins. Amicor.
Sieboldii, *Regel, Cat. Pl. Hort. Aksakov.* 38.—Hab.?
sphaerocarpa, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 362.—Peruv.
Spruceana, *Planch. & Triana, l. c.* 346.—Peruv.
subsessilis, *Benth. Bot. Voy. Sulph.* 72.—N. Granat.
sulphurea, *D. Dietr. Syn. Pl.* iv. 866.—Peruv.
tetrandra, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 978.—Ecuador.
thurifera, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiii. (1860) 364.—Peruv.?
venosa, *Jacq. Enum. Pl. Carib.* 34.—Ind. occ.
venosa, Linn. Sp. Pl. ed. II. 1495 = minor.
volubilis, *H. B. & K. Nov. Gen. et Sp.* v. 200.—N. Granat.
Weddelliana, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xiv. (1860) 233.—Bras.

CLUSIANTHEMUM, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 338 = **Garcinia**, Linn. (Guttifer.).
coriaceum, Vieill. l. c. 339 = *pedicellatum*.
densiflorum, Vieill. ex Pierre, Fl. For. Cochinch. fasc. v. 25 = G. densiflora.
pedicellatum, Vieill. in Bull. Soc. Linn. Normand. ix. (1865) 339.—N. Caled.

CLUSIELLA, Planch. & Triana, in Ann. Sc. Nat. Sér. IV. xiv. (1860) 253. *GUTTIFERAE*, Benth. & Hook. f. i. 172.
elegans, *Planch. & Triana, l. c.*—N. Granat.

CLUSIOPHYLLEA, Baill. Adansonia, xii. (1878) 197 = **Plectronia**, Linn. (Rubiac.).
Pervilleana, Baill. l. c. (= *Canthium Pervilleanum*).—Madag.

CLUSIOPHYLLUM, Muell. Arg. in Flora, xlvii. (1864) 518 = **Cunuria**, Baill. (Euphorb.).
Sprucei, Muell. l. c. 519 = Cun. Spruceana.

CLUYTIA, Boerh. ex Linn. Syst. ed. I (1735) (Clutia). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 302.
ALTORA, Adans. Fam. ii. 356 (1763).
CRATOCHWILIA, Neck. Elem. ii. 339 (1790).
abyssinica, *Jaub. & Spach, Illustr.* v. 77. t. 468.—Abyssin.
acuminata, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 125 = polifolia.
acuminata, *Linn. f. Suppl.* 432.—Afr. austr.
acuminata, Thunb. Prod. Pl. Cap. 53 = Lachnostylis hirta.
affinis, *Sond. in Linnaea*, xxiii. (1850) 126.—Afr. austr.
africana, Poir. Encyc. Suppl. ii. 302 = daphnoides.
alaternoides, *Linn. Sp. Pl.* 1042.—Afr. austr.
androgyna, Linn. Mant. i. 128 = Sauropus albicans.
benguelensis, *Muell. Arg. in Journ. Bot.* ii. (1864) 337.—Afr. trop.
brevifolia, Sond. in Linnaea, xxiii. (1850) 125 = polifolia.
Cascarilla, Linn. Sp. Pl. 1042 = Croton Cascarilla.
collina, Roxb. Pl. Corom. ii. 37 = Cleistanthus collinus.
cordata, *Bernh. in Flora*, xxviii. (1845) 81.—Afr. austr.
cotinifolia, Salisb. Prod. 390 = pulchella.
curvata, C. Mey. ex Sond. in Linnaea, xxiii. (1850) 121 = polygonoides.
daphnoides, Eckl. & Zeyh. ex Muell. Arg. in DC. Prod. xv. II. 1054 = polygonoides.
daphnoides, *Lam. Encyc.* ii. 54.—Afr. austr.
decandra, Crantz, Inst. i. 201 = Croton Eluteria.
diosmoides, *Sond. in Linnaea*, xxiii. (1850) 122.—Afr. austr.
diversifolia, Roxb. Hort. Beng. 104; Fl. Ind. iii. 731 = Cleistanthus diversifolius.
Dregeana, *Scheele, in Linnaea*, xxv. (1852) 583.—Afr. austr.
Eckloniana, *Muell. Arg. in DC. Prod.* xv. II. 1054.—Afr. austr.
Eluteria, Linn. Sp. Pl. 1042 = Croton Eluteria.

CLUYTIA :—
ericoides, Krebs, in Flora, xxviii. (1845) 82 = polygonoides.
ericoides, *Thunb. Prod. Pl. Cap.* 53.—Afr. austr.
floribunda, Baill. Etud. Gén. Euph. Atlas, 30. t. 16. f. 1–5 = alaternoides.
gracilis, Baill. Adansonia, iii. (1862–63) 151 = tenuifolia.
heterophylla, Sond. in Linnaea, xxiii. (1850) 128 = Sonderiana.
heterophylla, *Thunb. Prod. Pl. Cap.* 53.—Afr. austr.
hirsuta, Eckl. & Zeyh. ex Sond. in Linnaea, xxiii. (1850) 126 = daphnoides.
hirsuta, *Muell. Arg. in DC. Prod.* xv. II. 1046.—Afr. austr.
hirta, Linn. f. Suppl. 432 = Lachnostylis hirta.
hirta, *Vahl, Symb. Bot.* ii. 101.—Afr. austr.
humilis, Bernh. in Flora, xxviii. (1845) 81 = pubescens.
Jaubertiana, *Muell. Arg. in DC. Prod.* xv. II. 1044.—Arabia.
lanceolata, *Forsk. Fl. Aegypt. Arab.* 170.—Arabia.
lanceolata, Jaub. & Spach, Illustr. Pl. Or. v. t. 467 = Jaubertiana.
marginata, E. Mey. ex Sond. in Linnaea, xxiii. (1850) 130 = tomentosa.
Meyeriana, *Muell. Arg. in DC. Prod.* xv. II. 1055.—Afr. austr.
monoica, Lour. Fl. Cochinch. ii. 638 = Cleistanthus monoicus.
montana, Roxb. Pl. Corom. iii. 38. t. 171 = Briedelia montana.
myricoides, *Jaub. & Spach, Illustr. Pl. Or.* v. 73. t. 456, 466.—Arabia.
natalensis, *Bernh. in Flora*, xxviii. (1845) 81.—Afr. austr.
oblongifolia, Roxb. Hort. Beng. 70; Fl. Ind. iii. 730 = Cleistanthus chartaceus.
oblongifolia, Wall. Cat. n. 7886 = Trigonostemon semperflorens.
ovalis, Scheele, in Linnaea, xxv. (1852) 583 = Andrachne ovalis.
ovalis, *Sond. in Linnaea*, xxiii. (1850) 129.—Afr. austr.
patula, Herb. Roxb. ex Wall. Cat. n. 7877 = Cleistanthus collinus.
patula, Roxb. Pl. Corom. ii. 37. t. 170 = Cleistanthus patulus.
phyllanthifolia, Baill. Adansonia, iii. (1862–63) 153 = affinis.
polifolia, *Jacq. Hort. Schoenb.* ii. 67. t. 250.—Afr. austr.
polifolia, Sond. in Linnaea, xxiii. (1850) 124 = pterogona.
polygalaefolia, Salisb. Prod. 390 = alaternoides.
polygonoides, *Burm. f. Fl. Cap. Prod.* 31 (err. typ. 27).—Afr. austr.
polygonoides, *Linn. Sp. Pl.* ed. II. 1475.—Afr. austr.
polygonoides, Thunb. Prod. Pl. Cap. 53 = daphnoides.
polygonoides, Willd. Hort. Berol. t. 51 = alaternoides.
pterogona, *Muell. Arg. in DC. Prod.* xv. II. 1048.—Afr. austr.
pubescens, Eckl. & Zeyh. ex Sond. in Linnaea, xxiii. (1850) 126 = affinis.
pubescens, *Thunb. Prod. Pl. Cap.* 53.—Afr. austr.
pubescens, Willd. Sp. Pl. iv. 881 = daphnoides.
pulchella, *Linn. Sp. Pl.* 1042.—Afr. austr.
pulchella, Sparm. ex Sond. in Linnaea, xxiii. (1850) 126 = daphnoides.
retusa, Herb. Wight, ex Wall. Cat. n. 7877 = Cleistanthus collinus.
retusa, Linn. Sp. Pl. 1475 = Bridelia retusa.
retusa, Moon, Cat. Pl. Ceyl. 71 = Bridelia Moonii.
Richardiana, *Muell. Arg. in DC. Prod.* xv. II. 1044.—Afr. trop.
rubricaulis, Eckl. ex Sond. in Linnaea, xxiii. (1850) 128 = alaternoides.
scandens, Roxb. Pl. Corom. ii. 39. t. 173 = Bridelia stipularis.
semperflorens, Roxb. Hort. Beng. 104; Fl. Ind. iii. 730 = Trigonostemon semperflorens.
semperflorens, Wall. Cat. n. 7852 B, ex Baill. Etud. Gén. Euph. 474 = Symphyllia silhetiana.
sempervirens, Muell. Arg. in DC. Prod. xv. II. 764 = Symphyllia silhetiana, Trigonostemon Hookerianus.
sericea, *Muell. Arg. l. c.* 1053.—Afr. austr.
similis, *Muell. Arg. l. c.* 1046.—Afr. austr.
Sonderiana, *Muell. Arg. l. c.* 1051.—Afr. austr.

CLUYTIA :—
spinosa, Roxb. Pl. Corom. ii. 38. t. 172 = Briedelia retusa.
squarrosa, Lam. Encyc. ii. 54 = Briedelia retusa.
stipularis, Linn. Mant. 127 = Briedelia stipularis.
tabularis, Eckl. & Zeyh. ex Sond. in Linnaea, xxiii. (1850) 122 = polygonoides.
tenuifolia, *Willd. Sp. Pl.* iv. 880.—Afr. austr.
teretifolia, Sond. in Linnaea, xxiii. (1850) 124 = polifolia.
Thunbergii, Sond. l. c. 130 = daphnoides.
tomentosa, *Linn. Mant.* 299.—Afr. austr.
tomentosa, Thunb. Fl. Cap. ed. Schult. 271 = daphnoides.

CLUYTIA, Roxb. ex Steud. Nom. ed. II. i. 225 (1840) = **Bridelia**, Willd. (Euphorb.).

CLUYTIANDRA, Muell. Arg. in Journ. Bot. ii. (1864) 328. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 272.
trichopoda, *Muell. Arg. l. c.*—Afr. trop.

CLYBATIS, Phil. in Anal. Univ. Chil. (1872) 742. *COMPOSITAE*, Benth. & Hook. f. ii. 1237.
Volckmanni, *Phil. l. c.* 743.—Chili.

CLYMENUM, [Tourn.] Linn. Syst. ed. I (1735) = **Lathyrus**, Tourn. (Legumin.).
alatum, Link, Handb. ii. 188 = L. alatus.
articulatum, Medic. Phil. Bot. i. 204 = L. articulatus.
bicolor, Moench, Meth. 150 = L. articulatus.
dolichoides, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358 = L. articulatus.
Ochrus, Link, Handb. ii. 188; Alef. in Bonplandia, ix. (1861) 127 = L. Ochrus.
tenuifolium, Link, l. c. 494; Alef. l. c. 128 = L. tenuifolius.
tingitanum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 358 = L. tingitanus.
uncinatum, Moench, Meth. 150 = L. Clymenum.

CLYNHYMENIA, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 24, in syn. = CLINHYMENIA, A. Rich. & Gal. = **Cryptarrhena**, R. Br. (Orchid.).

CLYPEA, Blume, Bijdr. 26 (1825) = **Stephania**, Lour. (Menispermac.).
abyssinica, Dillon & A. Rich. in Ann. Sc. Nat. Sér. II. xiv. (1840) 263 = S. abyssinica.
acuminatissima, Blume, Bijdr. 28 = S. acuminata.
Burmanni, Wight & Arn. Prod. 14 = Cyclea Burmanni.
capitata, Blume, Bijdr. 28 = S. capitata.
consummata, Miers, Contrib. iii. 209.—Japon.
corymbosa, Blume, Bijdr. 27 = S. rotunda.
discolor, Blume, l. c. 26 = S. discolor.
effusa, Miers, Contrib. iii. 207.—Malabar.
Forsteri, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xvii. (1866) 270 = S. Forsteri.
glaucescens, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 423 t. 18 = S. glaucescens.
hernandifolia, Wight & Arn. Prod. 14 = S. hernandifolia.
longa, G. Don, Gen. Syst. i. 113 = S. hernandifolia.
oxyphylla, Miers, Contrib. iii. 206.—Nepal.
rotunda, Steud. Nom. ed. II. i. 387 = S. rotunda.
subovata, Miers, Contrib. iii. 209.—Japon.
tomentosa, Blume, Bijdr. 27 = S. tomentosa.
venosa, Blume, l. c. = S. venosa.

CLYPEOLA, Burm. ex DC. Prod. ii. 420 (1825) = **Pterocarpus**, Linn. (Legumin.).

CLYPEOLA, Linn. Gen. ed. I. 193 (1737). *CRUCIFERAE*, Benth. & Hook. f. i. 93.
BERGERETIA, Desv. Journ. Bot. iii. (1813) 161.
FOSSELINIA, Scop. Introd. 318 (1777).
IONTHLASPI, Adans. Fam. ii. 423 (1763).
ORIUM, Desv. Journ. Bot. iii. (1813) 161. t. 35. f. 10.
alliacea, Lam. Tabl. Encyc. iii. 113 = Peltaria alliacea.
altaica, Schangin, ex Ledeb. Fl. Ross. i. 135 = Alyssum spathulatum.
alyssoides, Linn. Sp. Pl. 652 = Alyssum calycinum.
ambigua, Jord. & Fourr. Brev. Pl. Nov. ii. 15 = microcarpa.

CLYPEOLA:—
auriculata, Crantz, Class. Crucif. 93 = Biscutella auriculata.
Bruhnsii, *Gruner, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 396.—Reg.. Casp.
calycina, All. ex Steud. Nom. ed. II. i. 387 = Alyssum calycinum.
capensis, Burm. ex DC. Prod. ii. 420 = Wiborgia fusca.
capensis, Poir. Encyc. Suppl. ii. 304 = Heliophila diffusa.
caroliniana, Walt. Fl. Carol. 173 = Lepidium virginicum.
chaetocarpa, Jaub. & Spach, Illustr. iii. 8. t. 206 = echinata.
ciliata, *Boiss. Fl. Orient.* i. 309.—Lycia.
edentula, Link, Handb. ii. 296 = Alyssum gemonense.
cyclodontea, *Delile, in Bull. Soc. Agr. Hérault*, (1830) 258.—Gallia.
dichotoma, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 175.—Persia; Armen.
didyma, Crantz, Stirp. Austr. ed. I. fasc. i. 20; ed. II. 22 = Biscutella laevigata.
echinata, Boiss. Fl. Orient. i. 310 = lappacea.
echinata, *DC. Syst.* ii. 328.—Syria; Persia; Afghan.
elegans, *Boiss. & Huet, Diagn.* Ser. II. v. 38.—Armenia.
eriocarpa, *Cav. Desc.* 401.—Hispan.
eriophora, DC. Syst. ii. 327 = eriocarpa.
Gaudini, Trachsel, in Flora, xiv. (1831) 743 = microcarpa.
glabra, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 173 = microcarpa.
gracilis, Planch. in Bull. Soc. Bot. Fr. v. (1858) 494 = microcarpa.
halimifolia, Link, Handb. ii. 295 = Alyssum halimifolium.
hispida, *Presl, Bot. Bemerk.* 9.—Arabia.
hispidula, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 15 = microcarpa.
indica, Poir. Encyc. Suppl. ii. 304 = Fortuynia Gareini.
Jonthlaspi, *Linn. Sp. Pl.* 652.—Reg. Mediterr.; Oriens.
laevigata, Jord. Brev. Pl. Nov. fasc. ii. 16 = microcarpa.
lapidicola, Jord. & Fourr. l. c. 14 = C. Jonthlaspi.
lappacea, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 174.—Mesopotam.; Persia.
lasiocarpa, Jaub. & Spach, Illustr. iii. 6. t. 205 = lappacea.
lasiocarpa, Juss. ex Pers. Syn. ii. 193 = echinata.
lomatotricha, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 194 = microcarpa.
macrocarpa, Link, Handb. ii. 295 = Alyssum macrocarpum.
maritima, Georgi, Beschr. Russ. Reich. iii. IV. 1126 = Alyssum canescens.
maritima, Linn. Sp. Pl. 652 = Alyssum maritimum.
microcarpa, *Boiss. Diagn.* Ser. I. i. 74.—Graecia; As. occ.
monosperma, Lam. Fl. Fr. ii. 462; 484 = C. Jonthlaspi.
montana, Crantz, Stirp. Austr. ed. I. fasc. i. 19; ed. II. 19 = Alyssum montanum.
perennis, Ard. Animad. Spec. 16. t. 6 = Peltaria alliacea.
petraea, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 14 = C. Jonthlaspi.
psilocarpa, Jord. & Fourr. l. c. = C. Jonthlaspi.
pyrenaica, Bordère, ex Dur. in Act. Soc. Linn. Bord. xxvi. (1866) 87 = C. Jonthlaspi.
pyrenaica, Link, Handb. ii. 295 = Alyssum pyrenaicum.
rupestris, Link, l. c. = Alyssum rupestre.
Schangini, Sievers, in Pall. N. Nord. Beitr. vi. 274 = Alyssum spathulatum.
semiglabra, Jord. & Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 194 = C. Jonthlaspi.
spathulaefolia, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 15 = microcarpa.
spinosa, Link, Handb. ii. 295 = Alyssum spinosum.
tomentosa, Linn. Mant. 92 = Alyssum orientale.

CLYTIA, Stokes, Bot. Mat. Med. iv. 543 (1812) = **Cluytia**, Boerh. (Euphorb.).

CLYTORIA, J. S. Presl, Rostl. iii. 196 (1835) = **Clitoria**, Linn. (Legumin.).

CLYTOSTOMA, Miers, ex Bur. in Adansonia, viii. (1868) 353 = **Pithecoctenium**, Mart. (Bignoniae.).
calystegioides, Bur.—Cf. Adansonia, viii. (1868) 353.—Am. austr.

CNEMIDIA, Lindl. Bot. Reg. sub t. 1618 (1833) = **Tropidia**, Lindl. (Orchid.).
angulosa, Lindl. l. c. = T. angulosa.
bambusifolia, Thw. Enum. Pl. Zeyl. 314.—Zeylan.
ctenophora, Reichb. f. Otia Bot. Hamb. 51.—Ins. Fiji.
semilibera, Lindl. Bot. Reg. sub t. 1618 = T. angulosa.

CNEMIDOSTACHYS, Mart. Nov. Gen. et Sp. i. 66. t. 40–44 (1824) = **Sebastiana**, Spreng. (Euphorb.).
acalyphoides, Mart. l. c. 71 = S. corniculata.
bidentata, Mart. l. c. 69 = S. virgata.
campestris, Mart. ex Baill. Etud. Gén. Euph. 516 = S. corniculata.
Chamaelaea, Spreng. Syst. iii. 835 = S. Chamaelea.
coriacea, Mart. Nov. Gen. et Sp. i. 68 = S. marginata.
crotonoides, Mart. l. c. 71 = S. corniculata.
daphnoides, Mart. l. c. = S. daphnoides.
ditassoides, F. Didr. in Kjoeb. Vidensk. Meddel. (1853) 88 = S. ditassoides.
dubia, Wawra, in Oestr. Bot. Zeitschr. xii. (1862) 241 = S. corniculata.
glabra, Mart. ex Baill. in Adansonia, v. (1864–65) 325 = S. multiramea.
glabrata, Mart. Nov. Gen. et Sp. i. 70 = S. corniculata.
glandulosa, Mart. l. c. 71 = S. corniculata.
hispida, Mart. l. c. = S. corniculata.
linearifolia, Miq. Fl. Ind. Bat. Suppl. 460 = S. Chamaelea.
longifolia, Mart. Nov. Gen. et Sp. i. 71 = S. corniculata.
madagascariensis, Boj. Hort. Maurit. 284 = Excoecaria melanosticta.
marginata, Mart. Nov. Gen. et Sp. i. 68. t. 41 = S. marginata.
myrtilloides, Mart. l. c. 67. t. 40 = S. daphnoides.
oleoides, Mart. l. c. 71 = S. oleoides.
patula, Mart. in Flora, xxiv. (1841) II. Beibl. 8 = S. corniculata.
prostrata, Mart. Nov. Gen. et Sp. i. 70 = S. corniculata.
salicifolia, Mart. l. c. = S. corniculata.
scoparia, Mart. l. c. = S. virgata.
Sellowiana, Klotzsch, ex Baill. Etud. Gén. Euph. 516 = S. corniculata.
serrulata, Mart. Nov. Gen. et Sp. i. 68 = S. serrulata.
stipulacea, Klotzsch, ex Baill. Etud. Gén. Euph. 516 = S. stipulacea.
tragioides, Mart. Nov. Gen. et Sp. i. 70 = S. corniculata.
Pahlii, Spreng. Syst. iii. 834 = S. corniculata.
velutina, Klotzsch, ex Baill. Etud. Gén. Euph. 516 = S. corniculata.

CNENAMUM, Tausch, in Flora, xi. (1828) I. Erg. 80 = CRENAMUM, Adans. = **Crepis**, Linn. (Compos.).

CNEORIDIUM, Hook. f. in Benth. & Hook. f. Gen. i. 312 (1862). *SIMARUBEAE*, Benth. & Hook. f. l. c.
PITAVIA, Nutt. ex Torr. & Gray, Fl. N. Am. 215 (1840).
dumosum, *Hook. f. l. c.*—Am. bor. occ.

CNEORUM, Linn. Syst. ed. I (1735). *SIMARUBEAE*, Benth. & Hook. f. i. 311.
pulverulentum, *Vent. Jard. Cels*, t. 77.—Ins. Canar.
tricoccon, Gueld. ex Bieb. Fl. Taur. Cauc. iii. 160 = Evonymus nanus.
tricoccon, *Linn. Sp. Pl.* 34.—Europ. austr.

CNESMOCARPUS, Zipp. ex Blume, Rumphia, iii. 115 (1847) = **Pometia**, Forst. (Sapind.).
excelsus, Zipp. ex Blume, l. c. (= *Irina Diplocardia*).—N. Guinea.

CNESMONE, Blume, Bijdr. 630 (1825) (Cnesmosa); Fl. Jav. Praef. p. vi (1828). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 330.
glabrata, *Kurz, in Flora*, lviii. (1875) 31.—Singapore.
javanica, *Blume, Bijdr.* 630.—Java.

CNESTIDIUM, Planch. in Linnaea, xxiii. (1850) 439. *CONNARACEAE*, Benth. & Hook. f. i. 433.
lasiocarpum, *Baker, in Mart. Fl. Bras.* xiv. II. 195.—Bras.
rufescens, *Planch. in Linnaea*, xxiii. (1850) 440.—Mexic.; Panama.

CNESTIS, Juss. Gen. 374 (1789). *CONNARACEAE*, Benth. & Hook. f. i. 433.
SPONDIOIDES, Smeathm. ex Lam. Encyc. iii. 23 (1789).
acuminata, Wall. Cat. n. 8533 = Rourea acuminata.
borboniensis, Raeusch. Nom. ed. III. 132 = glabra.
corniculata, Blanco, Fl. Filip. ed. I. 386 = Connarus ferrugineus.
corniculata, *Lam. Encyc.* iii. 23.—Afr. trop.
diffusa, Blanco, Fl. Filip. ed. I. 386 = Rourea rugosa.
emarginata, *Jack, in Malay. Misc.* ii. (1822) VII. 42.—Malaya.
ferruginea, *DC. Prod.* ii. 87.—Afr. trop.
flaminea, Griff. Notul. iv. 433 = ramiflora.
flammea, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) II. 76 = praec.
florida, *Jack, in Malay. Misc.* ii. (1822) VII. 43.—Sumatra.
foliosa, Planch. ex Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 216 = ramiflora.
fraterna, *Planch. in Linnaea*, xxiii. (1850) 440.—Afr. trop.
glabra, Blanco, Fl. Filip. ed. I. 387 = Connarus paniculatus.
glabra, *Lam. Encyc.* iii. 23.—Ins. Mascar.
glabra, Naves, in Blanco, Fl. Philipp. ed. III. t. 140 = Rourea multiflora.
grisea, *Baker, in Oliver, Fl. Trop. Afr.* i. 461.—Afr. trop.
ignea, Planch. ex Kurz, in Journ. As. Soc. Beng. xlv. (1876) II. 216 = ramiflora.
lasiocarpa, Mart. ex Baker, in Mart. Fl. Bras. xiv. II. 195 = Cnestidium lasiocarpum.
lurida, *Baill. Adansonia*, vii. (1866–67) 244. — Ins. Nossibé.
macrantha, *Baill. l. c.* 242.—Afr. trop.
madagascariensis, Raeusch. Nom. ed. III. 132 = polyphylla.
mimosoides, *Jack, in Malay. Misc.* ii. (1822) VII. 44.—Malaya.
monadelpha, Roxb. Hort. Beng. 34; Fl. Ind. ii. 454 = Rourea commutata.
natalensis, *Planch. & Sond. ex Harv. & Sond. Fl. Cap.* i. 528.—Afr. austr.
obliqua, *Beauv. Fl. Owar.* i. 97. t. 59.—Afr. trop.
obliqua, Boj. Hort. Maurit. 84 = Agelaea Lamarckii.
oblongifolia, *Baker, in Oliver, Fl. Trop. Afr.* i. 462.—Afr. trop.
pentaphylla, *Span. in Linnaea*, xv. (1841) 189.—Ins. Timor.
pinnata, Beauv. Fl. Owar. i. 98. t. 60 = Rourea Palisotii.
platantha, Griff. Notul. iv. 434 = ramiflora.
polyphylla, Blanco, Fl. Filip. ed. II. 270 = Rourea rugosa.
polyphylla, *Lam. Encyc.* iii. 23.—Maurit.; Madag.
racemosa, *G. Don, Gen. Syst.* ii. 91.—Afr. trop.
ramiflora, *Griff. Notul.* iv. 432.—Ind. or.
scandens, J. F. Gmel. Syst. 729 = glabra?
stenopetala, Griff. Notul. iv. 433 = Rourea stenopetala.
trifolia, Lam. Encyc. iii. 24 = Agelaea villosa.
vestita, Wall. ex Hook. f. Fl. Brit. Ind. ii. 47 = Agelaea vestita.
volubilis, Blanco, Fl. Filip. ed. I. 385 = Rourea heterophylla.

CNICOTHAMNUS, Griseb. in Goett. Abh. xix. (1874) 196. *COMPOSITAE*, Benth. & Hook. f. ii. 1237.
Lorentzii, *Griseb. l. c.* 197.—Reg. Argent.

CNICUS, Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 468.
ACARNA, Hill. Veg. Syst. iv. 13 (1762).
ANCATHIA, DC. in Guill. Arch. Bot. ii. (1833) 331.
BREEA, Less. Syn. Comp. 9 (1832).
CEPHANANOPLOS, Neck. Elem. i. 68 (1790).
CHAMAEPEUCE, DC. Prod. vi. 657 (1837).
CIRSIUM, [Tourn.] Adans. Fam. ii. 116 (1763); DC. Fl. Fr. iv. 110 (1805).
CLOMIUM, Adans. Fam. ii. 116 (1763).
ECHENAIS, Cass. in Bull. Soc. Philom. (1818) 33.
EPITRACHYS, C. Koch, in Linnaea, xxiv. (1851) 396.
ERIOLEPIS, Cass. in Dict. Sc. Nat. xxxv. 172 (1825); xli. 331 (1826).
ERYTHROLAENA, Sweet, Brit. Flow. Gard. Ser. I. t. 134 (1825).

CNICUS:—

Ixine, Hill, Veg. Syst. iv. 17 (1762).
Lamyra, Cass. in Dict. Sc. Nat. xxv. 218 (1822).
Lophiolepis, Cass. l. c. xli. 313 (1826).
Notobasis, Cass. l. c. xxxv. 170 (1825).
Onotrophe, Cass. l. c. xxxvi. 145 (1825).
Orthocentron, Cass. l. c. 480 (1825).
Picnomon, Adans. Fam. ii. 116 (1763).
Platyraphium, Cass. in Dict. Sc. Nat. xxxv. 173 (1825).
Polyacantha, Hill, Veg. Syst. iv. 12 (1762).
Ptilostemon, Cass. in Dict. Sc. Nat. xxxv. 173 (1825).
Pycnocomon, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 78, 84.
Spanioptilon, Less. Syn. Comp. 10 (1832).
Tetralix, Hill, Veg. Syst. iv. 18 (1762).
Xylanthema, Neck. Elem. i. 67 (1790).

acantholepis, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.
Acarna, *Linn. Syst.* ed. X. 1201.—Europ.
acaulis, *Willd. Prod. Fl. Berl.* 260.—Europ.
afer, *Sibth. & Sm. Fl. Graec.* ix. 20. t. 829.—Graecia; As. Min.
afer, Tenore, Prod. p. lxiii = niveus.
afer, Willd. Sp. Pl. iii. 1682 = Cn. Diacantha.
alpinus, Loisel. Fl. Gall. 542 = Saussurea alpina.
alsophilus, *Pollini, Fl. Veron.* ii. 620. t. 5.—Italia.
altaicus, Steph. ex DC. Prod. vi. 538 = Saussurea Frolowii.
altissimus, *Willd. Sp. Pl.* iii. 1671.—Am. bor.
ambiguus, Loisel. Fl. Gall. 540 = heterophyllus.
americanus, *A. Gray, in Proc. Am. Acad.* xix. (1884) 56.—Am. bor. occ.
Andersonii, *A. Gray, l. c.* x. (1874) 44.—Am. bor. occ.
Andrewsii, *A. Gray, l. c.* 45.—Calif.
anglicus, C. C. Gmel. Fl. Bad. iii. 372 = pratensis.
angustifolius, Buch.-Ham. in Wall. Cat. n. 2915 = Tricholepis Stictophyllum.
arabicus, *Roth, in Usteri, Ann. Bot.* viii. 8.—Arabia.
arachnoides, Hort. Par. ex DC. Prod. vi. 635 = fimbriatus.
arachnoides, Wall. Cat. n. 2891 = Wallichii.
arachnoideus, *Bieb. Fl. Taur. Cauc.* ii. 277.—Reg. Caucas.
araneosus, *Desf. ex DC. Prod.* vi. 635.—Reg. Caucas.
arenarius, Willd. Sp. Pl. iii. 1663 = Carduus arenarius.
Argemone, Lapeyr. Hist. Abr. Pl. Pyr. ii. 493 = medius.
argentatus, Roth, Catalect. i. 111 = Carduus argentatus.
argyracanthus, *C. B. Clarke, Comp. Ind.* 218.—Reg. Himal.
arizonicus, *A. Gray, in Proc. Am. Acad.* x. (1874) 44.—Arizona.
armatus, [*Soland. in*] *Russ. Aleppo*, ed. II. ii. 262.—Syria.
arvensis, *Hoffm. Deutschl. Fl.* ed. II. i. 11. 130.—Europ.; As. temp.
arvensis, Hook. Comp. Bot. Mag. i. (1835) 48 = virginianus.
asiaticus, *Sievers, in Pall. N. Nord. Beitr.* vii. 350.—Sibir. altaic.
auriculatus, Wall. Cat. n. 3289 = Saussurea hypoleuca.
autareticus, Chaix, in Vill. Hist. Pl. Dauph. i. 381; Willd. Sp. Pl. iii. 1676 = heterophyllus.
autareticus, Loisel. Fl. Gall. 541 = rivularis.
Bertolonii, *Bertol. Fl. Ital.* ix. 23.—Italia.
Boujarti, DC. Prod. vi. 635 = fimbriatus vel arachnoideus.
bracteatus, Gilib. Fl. Lituan. i. 184 = oleraceus.
Breweri, *A. Gray, in Proc. Am. Acad.* x. (1874) 43.—Calif.
Buergeri, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 502.—Japon.
californicus, *A. Gray, in Proc. Am. Acad.* x. (1874) 46.—Calif.
callosus, Roth, Catalect. iii. 76 = Carduus callosus.
candicans, Wall. Cat. n. 2893 = arvensis.
canus, *Roth, Tent. Fl. Germ.* i. 345.—Europ.
capitatus, *Krock. Fl. Sil. Suppl.* ii. 290.—Europ.
carlinoides, *Fisch. ex Schrank, Pl. Rar. Hort. Monac.* t. 11.—Reg. Caucas.
carniolicus, *Willd. Sp. Pl.* iii. 1677.—Europ.
carthamoides, Wall. Cat. n. 2896 = Saussurea affinis.
carthamoides, Willd. Sp. Pl. iii. 1686 = Centaurea carthamoides.

CNICUS:—

Casabonae, *Roth, in Usteri, Ann. Bot.* viii. 9.—Ins. Ital.; Marocco.
centauroides, *Linn. Sp. Pl.* 826.—Mont. Pyren.
centauroides, Willd. ex Ledeb. Fl. Ross. ii. 753 = Leuzea salina.
cernuus, A. Gray, in Proc. Am. Acad. x. (1874) 44 = subcoriaceus.
cernuus, Linn. Sp. Pl. 826 = Carduus cernuus.
cernuus, Wall. Cat. n. 2892 = Wallichii.
chamaecephalus, *Vatke, in Linnaea*, xxxix. (1875) 511.—Abyssin.
Chamaepeuce, *Desf. Arb.* i. 280.—Graecia; As. Min.
Chamaepeuce, Tenore, Prod. p. xlviii = gnaphalodes.
chinensis, *Benth. ex Maxim. in Bull. Acad. Pétersb.* xix. (1874) 509.—China.
chrysacanthus, *Ball, in Journ. Bot.* xi. (1873) 369.—Marocco.
ciliatus, *Roth, Catalect.* ii. 104.—Europ.
ciliatus, Vitm. Summa, iv. 447 = ochroleucus.
collinus, Hornem. Hort. Hafn. Suppl. 92 = Carduus collinus.
communis, Lam. Fl. Fr. ii. 15 = Cynara Cardunculus.
comosus, *Franch. & Sav. Enum. Pl. Jap.* ii. 409.—Japon.
conspicuus, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.
costaricensis, *Polak. in Linnaea*, xli. (1879) 581.—Costa Rica.
cyanoides, *Roth, in Usteri, Ann. Bot.* viii. 9.—Europ.
Cynara, Lam. Fl. Fr. ii. 14 = Centaurea cynaroides.
cynaroides, Bieb. Fl. Taur. Cauc. ii. 280 (= *Cirsium sinuatum*).—Reg. Caucas.
cynaroides, Sibth. & Sm. Fl. Graec. ix. 20. t. 828 = eriophorus.
cynaroides, *Willd. Sp. Pl.* iii. 1670.—Ins. Creta.
deltoideus, Wall. ex DC. Prod. vi. 541 = Saussurea deltoidea.
dentatus, Forsk. Fl. Aegypt. Arab. 217 = Kentrophyllum dentatum.
dentatus, Willd. Sp. Pl. iii. 1672 = pratensis.
Diacantha, *Desf. Cat. Hort. Par.* ed. III. 152.—As. Min.; Syria.
dipsacolepis, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 497.—Japon.
discolor, *Muhl. ex Willd. Sp. Pl.* iii. 1670.—Am. bor.
discolor, Roth, ex Steud. Nom. ed. II. i. 283, 388 = Carduus callosus.
dissectus, *Willd. Sp. Pl.* iii. 1665.—Europ.
Drummondii, *A. Gray, in Proc. Am. Acad.* x. (1874) 40.—Am. bor. occ.
dubius, Willd. Prod. Fl. Berl. 260 = acaulis.
Eatoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 56.—Am. bor. occ.
echinatus, *Willd. Sp. Pl.* iii. 1668.—Reg. Mediterr. occ.
echinocephalus, *Willd. l. c.* 1685.—Reg. Caucas.
edulis, *A. Gray, in Proc. Am. Acad.* x. (1874) 47.—Am. bor. occ.
effusus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 500.—Japon.
elongatus, Wall. ex C. B. Clarke, Comp. Ind. 239 = Tricholepis elongata.
eriocephalus, *A. Gray, in Proc. Am. Acad.* x. (1874) 46.—Am. bor. occ.
eriophoroides, *Hook. f. Fl. Brit. Ind.* iii. 363.—Reg. Himal.
eriophorus, C. B. Clarke, Comp. Ind. 217 = eriophoroides.
eriophorus, Juss. ex DC. Prod. vi. 635 = spathulatus.
eriophorus, *Roth, Tent. Fl. Germ.* i. 345.—Europ.; As. Min.
Erisithales, Linn. Sp. Pl. ed. II. 1157 = ochroleucus?
esculentus, *Sievers, in Pall. N. Nord. Beitr.* vii. 362.—Rossia.
Falconeri, *Hook. f. Fl. Brit. Ind.* iii. 363.—Reg. Himal.
ferox, *Linn. Mant.* i. 109; *Sibth. & Sm. Fl. Graec. Prod.* ii. 152.—Europ. or.; As. Min.
fimbriatus, *Bieb. Fl. Taur. Cauc.* ii. 276.—Reg. Caucas.
firmus, *J. & C. Presl, Delic. Prag.* 107.—Europ.
flavescens, Willd. Sp. Pl. iii. 1683 = Serratula flavescens.
flavispinus, *Ball, in Journ. Linn. Soc.* xvi. (1878) 522.—Hispan.; Marocco.
foliosus, *A. Gray, in Proc. Am. Acad.* x. (1874) 40.—Am. bor. occ.

CNICUS:—

fruticosus, *Desf. Arb.* i. 280.—Graecia.
Galactites, Loiscl. Fl. Gall. ed. I. 538 = Galactites tomentosa.
giganteus, *Willd. Sp. Pl.* iii. 1671.—Reg. Mediterr.
glaber, Ell. Sketch, ii. 270 = Nuttallii.
glaber, Nutt. Gen. Am. ii. 129 = muticus?
glutinosus, Bigel. Fl. Bost. ed. II. 291 = muticus.
glutinosus, Gilib. Fl. Lituan. i. 185 = ochroleucus?
Gmelini, Spreng. Hist. Rei Herb. ii. 270 = acaulis.
gnaphalodes, *Willd. ex Spreng. Syst.* iii. 375.—Italia.
gossypinus, Wall. ex DC. Prod. vi. 541, 542 = Saussurea gossypiphora.
Gouani, Willd. Sp. Pl. iii. 1665 = Carduus medius.
Grahami, *A. Gray, in Proc. Am. Acad.* xix. (1883) 57.—Arizona.
Griffithii, *Hook. f. Fl. Brit. Ind.* iii. 363.—Reg. Himal.
grumosus, *Hoffmgg. & Link, Fl. Port.* ii. 196.—Lusitan.
Hallii, *A. Gray, in Proc. Am. Acad.* xix. (1883) 56.—Am. bor. occ.
helenioides, Willd. Sp. Pl. iii. 1674 = heterophyllus.
heterolepis, A. Gray, in Proc. Am. Acad. x. (1874) 44 = subcoriaceus.
heteromallus, Steud. Nom. ed. II. i. 388 = Saussurea candicans.
heterophyllus, *Roth, Catalect.* i. 114.—Europ.
Hilgendorfi, *Franch. & Sav. Enum. Pl. Jap.* ii. 410.—Japon.
hispanicus, *Ball, in Journ. Linn. Soc.* xvi. (1878) 523.—Hispan.
Hookerianus, *A. Gray, in Proc. Am. Acad.* x. (1874) 46.—Am. bor. occ.
horridulus, *Pursh, Fl. Am. Sept.* ii. 507.—Am. bor.
horridus, Bertol. Amoen. Ital. 40 = Bertolonii.
horridus, *Bieb. Fl. Taur. Cauc.* ii. 278.—Reg. Caucas.
horridus, C. B. Clarke, Comp. Ind. 220 = Falconeri.
horridus, Forsk. Fl. Aegypt. Arab. 217 = Cardopatium corymbosum.
hybridus, Schleich. Cat. Pl. Helv. ed. III. 12 = ochroleucus.
igniarius, *Benth. & Hook. f. Gen.* ii. 469.—Sibir. altaic.
incomptus, *Franch. & Sav. Enum. Pl. Jap.* ii. 410.—Japon.
inermis, Willd. Sp. Pl. iii. 1672 = centauroides.
intermedius, *Heller, Fl. Wirc.* ii. 256.—Europ.
involucratus, *Wall. ex DC. Prod.* vi. 639.—Reg. Himal.
italicus, *Seb. & Mauri, Fl. Rom. Prodr.* ii. 282.—Italia.
japonicus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 503.—Japon.
jorullensis, *H. B. & K. Nov. Gen. et Sp.* iv. 22.—Mexic.
kamtschaticus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 495.—Japon.; Alaska.
Kosmelii, Willd. ex Ledeb. Fl. Ross. ii. 729 = lappaceus.
Kotschyi, Sch. Bip. in Ind. Sem. Hort. Berol. (1857) App. 24 = Carbenia benedicta.
Lachenalii, *C. C. Gmel. Fl. Bad.* iii. 380.—Europ.
lacteus, Schleich. Cat. Pl. Helv. ed. IV. 13 = palustris.
lanatus, Willd. Sp. Pl. iii. 1671 = arvensis.
lanceolatus, *Willd. Prod. Fl. Berl.* 259.—Europ.
laniflorus, *Bieb. Fl. Taur. Cauc.* ii. 276.—Reg. Caucas.
lappaceus, *Bieb. l. c.* 277.—Reg. Caucas.
lappaceus, Hort. Par. ex DC. Prod. vi. 639 = nivalis.
lappoides, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.
Lecontei, *A. Gray, in Proc. Am. Acad.* x. (1874) 39.—Am. bor.
leucocephalus, *Willd. Sp. Pl.* iii. 1668.—Creta; As. Min.
leucographus, Roth, in Usteri, Ann. Bot. viii. 8 = Tyrimnus leucographus.
leucophanus, Baumg. ex Schur, Enum. Pl. Transs. 418 = lanceolatus.
linearis, *Benth. & Hook. f. ex Franch. & Sav. Enum. Pl. Jap.* i. 261.—Japon.
lomatolepis, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.
longifolius, Lam. Fl. Fr. ii. 13 = Carduncellus monspeliensis.
lucidus, Wall. ex DC. Prod. vi. 622 = Carduus nutans.
macrostylus, *Moretti, in Brugnat. Giorn. Fis.* II. (1824) 102.—Italia.

CNICUS :—

medius, *Krock. Fl. Sil. Suppl.* ii. 278.—Silesia.
mexicanus, *Hemsl. Biol. Centr. Am. Bot.* ii. 251.—Mexic.
microcephalus, *Bertol. Fl. Ital.* ix. 36.—Ital.; Sardinia.
microcephalus, Boiss. Fl. Orient. iii. 705 = Carbenia benedicta.
misilmcrensis, *Tineo, ex Ces. Passer. & Gib. Comp. Fl. Ital.* 483.—Ital.
mitis, *Hornem. Hort. Hafn. Suppl.* 93.—Sibir.
mitissimus, Lam. Fl. Fr. ii. 13 = Carduncellus mitissimus.
mollis, DC. Prod. vi. 652, sphalm. = acaulis.
mollis, Roth, Tent. Fl. Germ. i. 346 = Jurinea mollis.
monspessulanus, *Roth, in Usteri, Ann. Bot.* viii. 9.—Gallia.
montanus, *Waldst. & Kit. ex Willd. Sp. Pl.* iii. 1676.—Europ.
multicaulis, Wall. ex DC. Prod. vi. 540 = Saussurea affinis.
munitus, *Bieb. Fl. Taur. Cauc.* ii. 279.—Reg. Caucas.
muticus, *Ell. Sketch*, ii. 268.—Am. bor.
neo-mexicanus, *A. Gray, in Proc. Am. Acad.* x. (1874) 45.—Am. bor. occ.
nipponicus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 496.—Japon.
nivalis, *H. B. & K. Nov. Gen. et Sp.* iv. 23.—Mexic.
niveus, *J. & C. Presl, Delic. Prag.* 108.—Europ.
niveus, Wall. Cat. n. 2898 = Saussurea deltoidea.
nudiflorus, *Schleich. Cat. Pl. Helv.* ed. IV. 14, nomen.—Helvet.
Nuttallii, *A. Gray, in Proc. Am. Acad.* x. (1874) 41.—Am. bor.
obvallatus, *Bieb. Fl. Taur.-Cauc.* ii. 279.—Reg. Caucas.
obvallatus, Salzm. ex DC. Prod. vi. 660 = Notobasis syriaca.
occidentalis, *A. Gray, in Proc. Am. Acad.* x. (1874) 45.—Am. bor. occ.
ochrocentrus, *A. Gray, l. c.* xix. (1883) 57.—Am. bor. occ.
ochroleucus, *Spreng. Nachr.* i. *Bot. Gart. Halle*, 19.—Europ.
odoratus, Muhl. ex Steud. Nom. ed. II. i. 388 = pumilus.
oleraceus, *Linn. Sp. Pl.* 826.—Europ.
oligophyllus, *Franch. & Sav. Enum. Pl. Jap.* ii. 412.—Japon.
orgyalis, *Willd. Enum. Hort. Berol. Suppl.* 56.—Hab.?
orientalis, Willd. Sp. Pl. iii. 1670 = Galactites tomentosa.
ornatus, *Ball, in Journ. Bot.* xi. (1873) 368.—Marocco.
ovalifolius, *Franch. & Sav. Enum. Pl. Jap.* ii. 412.—Japon.
paludosus, *Loisel. Fl. Gall.* 542.—Europ.
palustris, Sibth. & Sm. Fl. Graec. Prod. ii. 151 = pungens.
palustris, *Willd. Prod. Fl. Berl.* 260.—Europ.; As. bor.
pannonicus, *Host, Fl. Austriac.* ii. 444.—Europ.
Parryi, *A. Gray, in Proc. Am. Acad.* x. (1874) 47.—Am. bor. occ.
parviflorus, *Heller, Fl. Wirc. Suppl.* 76.—Europ.
pauciflorus, *Waldst. & Kit. Pl. Rar. Hung.* ii. 175.—Europ. or.
pazcuarensis, *H. B. & K. Nov. Gen. et Sp.* iv. 22. t. 310.—Mexic.
pectinellus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 494.—Japon.
pedemontanus, *Willd. Sp. Pl.* iii. 1675.—Europ.
pendulus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 510.—Sibir. or.
pexus, *Franch. & Sav. Enum. Pl. Jap.* ii. 410.—Japon.
pinnatifidus, Willd. Sp. Pl. iii. 1684 = Serratula pinnatifida.
Pitcheri, *Torr. in Eaton, Man.* ed. V. 180.—Am. bor.
polyacanthus, *Vatke, in Linnaea*, xxxix. (1875) 510.—Afr. trop.
polyanthemus, *Bertol. Amoen. Ital.* 41.—Italia.
Pontederae, Pollini, in Brugnat. Giorn. Fis. ix. (1816) 98 = rivularis.
praemorsus, Michl, in Flora, iii. (1820) 317 = ochroleucus?
pratensis, Lam. Fl. Fr. ii. 14 = oleraceus.
pratensis, *Willd. Sp. Pl.* iii. 1672.—Europ.

CNICUS :—

pseudo-benedictus, Hort. Dorpat. ex Aschers. Ind. Sem. Hort. Berol. (1861) 5 = Carbenia benedicta.
pubigerus, *Desf. Cat. Hort. Par.* ed. III. 399.—As. Min.
pumilus, *Torr. Compend.* 282.—Am. bor.
pungens, Bivona, Sic. Pl. Cent. i. 26 = siculus.
pungens, Sebast. Pl. Rom. fasc. i. 9. t. 2 = polyanthemus.
pungens, *Willd. Sp. Pl.* iii. 1663.—Europ. or.; As. Min.
purpuratus, *Maxim. in Bull. Acad. Pétersb.* xix. (1879) 491.—Japon.
purpureus, Bertol. Fl. Ital. ix. 21 = spinosissimus.
pycnocephalus, Roth, in Usteri, Ann. Bot. viii. 8 = Carduus pycnocephalus?
pygmaeus, Linn. Sp. Pl. ed. II. 1156 = Saussurea pygmaea.
pyrenaicus, *Willd. Sp. Pl.* iii. 1664.—Europ.
quercetorum, *A. Gray, in Proc. Am. Acad.* x. (1874) 40.—Calif.
radians, *Hemsl. Biol. Centr. Am. Bot.* ii. 252.—Guatemala.
rhaphilepis, *Hemsl. l. c.*—Mexic.
rectus, Hort. Par. ex DC. Prod. vi. 659 = strictus.
Reinii, *Franch. & Sav. Enum. Pl. Jap.* ii. 413.—Japon.
remotifolius, *A. Gray, in Proc. Am. Acad.* x. (1874) 47.—Am. bor. occ.
repandus, *Ell. Sketch*, ii. 269.—Am. bor.
replicatus, *Hornem. Hort. Hafn. Suppl.* 93.—Hungaria.
rigens, *Willd. Sp. Pl.* iii. 1675.—Europ.
rivularis, *Willd. l. c.* 1676.—Europ.
Rothrockii, *A. Gray, in Proc. Am. Acad.* xvii. (1881–82) 220.—Arizona.
rufescens, *Loisel. Fl. Gall.* 541.—Gallia.
ruthenicus, Henning, in Mém. Soc. Nat. Mosc. vi. (1823) 78 = arvensis.
saanensis, Schlecht. ex DC. Prod. vi. 648 = ochroleucus.
salinus, Willd. ex Ledeb. Fl. Ross. ii. 752 = Leuzea serratuloides.
salisburgensis, Willd. Sp. Pl. iii. 1675 = rivularis.
samniticus, Tenore, ex Steud. Nom. ed. II. i. 389 = italicus.
scariosus, *A. Gray, Syn. Fl. N. Am.* i. II. 402.—Am. bor. occ.
Schimperi, *Vatke, in Linnaea*, xxxix. (1875) 511.—Afr. trop.
segetum, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 511.—China.
seminudus, Schleich. ex DC. Prod. vi. 650 = rivularis.
semipectinatus, *Desf. Cat. Hort. Par.* ed. III. 151.—As. centr.
semipectinatus, *Hornem. & Ledeb. ex DC. Prod.* vi. 647.—Hab.?
semipectinatus, Thom. ex Steud. Nom. ed. II. i. 389 (= *Cirsium subalpinum*).—Europ.
serratuloides, *Roth, Tent. Fl. Germ.* i. 346.—Sibiria.
serrulatus, *Bieb. Fl. Taur.-Cauc.* ii. 275.—Reg. Caucas.
setosus, Bess. Prim. Fl. Galic. ii. 172 = arvensis?
siculus, Guss. Fl. Sic. Syn. ii. 442 = pungens.
Sieboldii, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 503.—Japon.
sinensis, C. B. Clarke, Comp. Ind. 219 = chinensis.
spathulatus, *Moretti, in Brugnat. Giorn. Fis.* II. v. (1822) 111.—Italia.
spicatus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 501.—Japon.
spinosissimus, Darling. Fl. Cestr. ed. II. 438 = horridulus.
spinosissimus, Habl. Taur. 158, ex Bieb. Pl. Taur.-Cauc. ii. 281 = echinocephalus.
spinosissimus, Lapeyr. Hist. Abr. Pl. Pyr. ii. 496 = seq.?
spinosissimus, *Linn. Sp. Pl.* 826.—Europ.
spurius, Hoffm. Deutschl. Fl. ed. II. ii. 128 = pratensis.
squarrosus, *Hornem. Hort. Hafn.* ii. 782.—Sibiria.
stellatus, *Roth, in Usteri, Ann. Bot.* viii. 9.—Europ.
strictus, *Tenore, Prod.* p. xlviii.—Italia.
strigosus, *Bieb. Fl. Taur.-Cauc.* ii. 277.—Reg. Caucas.; Persia.
subcoriaceus, *Hemsl. Biol. Centr. Am. Bot.* ii. 252.—Mexic.
suffultus, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 498.—Japon.

CNICUS :—

syriacus, *Roth, in Usteri, Ann. Bot.* viii. 10.—Reg. Mediterr.
Tanakae, *Franch. & Sav. Enum. Pl. Jap.* ii. 411.—Japon.
tartaricus, *Willd. Sp. Pl.* iii. 1677.—As. temp.
tchefouensis, *Franch. in Mém Soc. Sc. Nat. Cherbourg*, xxiv. (1884) 229.—China.
transsylvanicus, *Schur, in Oestr. Bot. Zeitschr.* x. (1860) 179.—Transsylvan.
tuberosus, *Roth, Tent. Fl. Germ.* i. 345.—Europ.
uliginosus, *Bieb. Fl. Taur.-Cauc.* ii. 274.—Reg. Caucas.
undulatus, *A. Gray, in Proc. Am. Acad.* x. (1874) 42.—Am. bor.
uniflorus, Linn. Mant. ii. 572 = Centaurea monanthos.
uniflorus, Sievers, in Pall. N. Nord. Beitr. vii. 346 = Leuzea carthamoides.
Verutus, D. Don, Prod. Fl. Nep. 167 = argyracanthus.
virginianus, Hook. Comp. Bot. Mag. i. (1835) 48 = Lecontei.
virginianus, *Pursh, Fl. Am. Sept.* ii. 506.—Am. bor.
Vlassovianus, *Maxim. in Bull. Acad. Petersb.* xix. (1874) 509.—As. temp.
Wallichii, *Hook. f. Fl. Brit. Ind.* iii. 363.—Reg. Himal.
Weyrichii, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 495.—Ins. Sachalin.
Wheeleri, *A. Gray, in Proc. Am. Acad.* xix. (1883) 56.—Arizona.
Willdenowii, *F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr.* ii. 331.—Hab.?
Wrightii, *A. Gray, in Proc. Am. Acad.* x. (1874) 41.—Texas; Arizona.
yezoënsis, *Maxim. in Bull. Acad. Pétersb.* xix. (1874) 508.—Japon.

CNICUS, [Vaill.] Gaertn. Fruct. ii. 385. t. 162 (1791) = **Carbenia**, Adans. (Compos.).
benedictus, Linn. Sp. Pl. 826; Gaertn. l. c. = Carb. benedicta.

CNIDIUM, Cusson, in Mém. Soc. Méd. Paris (1782) 280 = **Selinum**, Linn. (Umbell.).
affine, C. A. Mey. in Beitr. Pfl. Russ. Reich. vi. 38.—Reg. Caucas.
alatum, Spreng. ex Link, Handb. i. 329 = Ligusticum alatum.
alsaticum, Spreng. Pl. Umbellif. Prod. 40 = Peucedanum alsaticum.
anomalum, Ledeb. Fl. Alt. i. 330; Ic. Fl. Ross. t. 311.—Sibir. altaic.
apioides, Spreng. Pl. Umbellif. Prod. 40 = Carum rigidulum.
argenteum, Cesati, in Linnaea, xi. (1837) 324 = Peucedanum involucratum?
athoum, Griseb. Spicil. Fl. Rumel. i. 363 = Carum rigidulum.
atropurpureum, Spreng. in Roem. & Schult. Syst. vi. 418 = Thaspium trifoliatum.
canadense, Spreng. l. c. 415 = S. canadense.
carvifolium, Bieb. Fl. Taur.-Cauc. iii. 212 = Carum caucasicum.
cenolophioides, Benth. & Hook. f. Gen. i. 914 = S. sylvestre.
chinense, Spreng. ex Steud. Nom. ed. II. i. 389 = S. canadense.
confertum, Moench, Meth. 98 = S. Monnieri.
coniifolium, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 299.—As. Min.
cuneatum, Ledeb. Fl. Alt. i. 331; et Ic. Fl. Ross. t. 312.—Sibir. altaic.
cuneifolium, Griseb. Spicil. Fl. Rumel. i. 363, in obs. = Ligusticum cuneifolium.
dahuricum, Fisch. & Mey. in Ann. Sc. Nat. Sér. II. v. (1836) 188.—Dahuria.
diffusum, DC. Prod. iv. 153 = Seseli indicum.
divaricatum, Ledeb. Fl. Alt. i. 324 = Silaus longifolius.
Fischeri, Spreng. Nov. Prov. 12 = Silaus longifolius.
Fontanesii, Spreng. Sp. Umbellif. Prod. 41.—Numidia.
intermedium, Tchihat. Asie Min. Bot. i. 430.—As. Min.
japonicum, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 60.—Japon.
Kraussianum, Sond. in Harv. & Sond. Fl. Cap. ii. 552 = Pimpinella caffra.

CNIDIUM :—
meifolium, Bieb. Fl. Taur. Cauc. iii. 213 = Carum armenum.
microcarpum, Turcz. ex Steud. Nom. ed. II. i. 389.—Sibir. baical.
Monnieri, Cusson, in Mém. Soc. Méd. Par. (1782) 280 = Selinum Monnieri.
multicaule, Ledeb. Fl. Ross. ii. 284.—Sibiria.
myrrhidifolium, Bieb. Fl. Taur. Cauc. iii. 212 = Ligusticum alatum.
orientale, Boiss. in Ann. Sc. Nat. Sér. III. i. (1844) 299 = S. orientale.
palustre, Reichb. Fl. Germ. Excurs. 469 = S. sylvestre.
Petroselinum, DC. Prod. iv. 152 = Carum Petroselinum.
peucedanoides, H. B. & K. Nov. Gen. et Sp. v. 15 = Eulophus peucedanoides.
pyrenaeum, Spreng. in Roem. & Schult. Syst. vi. 416 = Ligusticum pyrenaeum.
salinum, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 733.—Dahuria.
Silaus, Bieb. Fl. Taur. Cauc. iii. 212 = Silaus Besseri.
Silaus, Spreng. Pl. Umbellif. Prod. 40 = Silaus pratensis.
striatum, Turcz. ex Fisch. & Mey. in Ann. Sc. Nat. Sér. II. v. (1836) 188 = *dahuricum*.
suffruticosum, Cham. & Schlecht. in Linnaea, i. (1826) 387 = S. suffruticosum.
tenuifolium, Moench, Meth. 98 = Apium leptophyllum.
tolucense, Spreng. Syst. i. 888 = Arracacia tolucensis.
venosum, Koch, Umbellif. 109 = S. sylvestre.

CNIDONE, E. Mey. ex Endl. Gen. Suppl. ii. 76 (1842) = **Kissenia**, R. Br. (Loasac.).
mentzelioides, E. Mey. ex Presl, Bot. Bemerk. 73 = K. spathulata.

CNIDOSCOLUS, Pohl, Pl. Bras. Ic. i. 56. t. 49 (1827) = **Jatropha**, Linn. (Euphorbiac.).
angustidens, Torr. Bot. Mex. Bound. 198 = J. augustidens.
cnicodendron, Griseb. in Goett. Abh. xxiv. (1879) 53.—Reg. Argent.
fragrans, Pohl, Pl. Bras. i. 63 = J. fragrans.
hamosus, Pohl, l. c. 57 = J. hamosa.
lobatus, Pohl, l. c. 62 = J. phyllacantha.
Marcgravii, Pohl, l. c. 58. t. 50 = J. urens.
Michauxii, Pohl, l. c. 58 = J. urens.
napaefolius, Pohl, l. c. 63 = J. aconitifolia.
neglectus, Pohl, l. c. 60. t. 51 = J. urens.
obtusifolius, Pohl, ex Baill. in Adansonia, iv. (1863-64) 275 = J. obtusifolia.
osteocarpus, Pohl, Pl. Bras. i. 63 = J. osteocarpa.
palmatus, Pohl, l. c. = J. palmata.
pubescens, Pohl, l. c. 62 = J. obtusifolia.
quercifolius, Pohl, ex Baill. in Adansonia, iv. (1863-64) 275 = J. phyllacantha.
quinquelobus, Pohl, Pl. Bras. i. 63 = J. urens.
repandus, Pohl, l. c. 62 = J. phyllacantha.
stimulosus, Engelm. & A. Gray, in Bost. Journ. Nat. Hist. v. (1845) 234 = J. texana.
stimulosus, A. Gray, Man. Bot. 389 = J. urens.
surinamensis, Miq. in Linnaea, xviii. (1844) 749 = Croton lobatus.
vitifolius, Pohl, Pl. Bras. i. 61 = J. vitifolia.

CNOPOS, Rafin. Fl. Tellur. iii. 13 (1836) = **Polygonum**, Tourn.
ramosissimum, Rafin. l. c. = P. ramosissimum.

COA, Adans. Fam. ii. 445 (1763) = **Hippocratea**, Linn. (Celastr.).

COALISIA, Rafin. Sylva Tellur. 114 (1838) = **Cleome**, Linn. (Capparid.).
angustifolia, Rafin. l. c. = Cleome angustifolia.

COATESIA, F. Muell. Fragm. iii. 26 (1862) = **Geijera**, Schott (Rutac.).
paniculata, F. Muell. l. c. = G. Muelleri.

COBAEA, Cav. Ic. i. 11. t. 16, 17 (1791). *POLEMONIACEAE*, Benth. & Hook. f. ii. 824.
ROSENBERGIA, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 30.
gracilis, *Hemsl. in The Garden*, xvii. (1880) 352.—Costa Rica.

COBAEA :—
lutea, D. Don, in Edinb. Phil. Journ. x. (1824) 4 = macrostemma.
macrostemma, *Pav. ex D. Don*, *l. c.* 112.—Guatemala.
minor, *Mart. & Gal. in Bull. Acad. Brux.* xii. (1845) II. 276.—Mexic.
penduliflora, *Hook. f. Bot. Mag.* t. 5757.—Venezuela.
scandens, *Cav. Ic.* i. 11. tt. 16, 17.—Mexic.
stipularis, *Benth. Pl. Hartw.* 45.—Mexic.

COBAEA, Neck. Elem. i. 129 (1790) = **Lonicera**, Linn. (Caprifol.).

COBAMBA, Blanco, Fl. Filip. ed. I. 510 (1837) = **Canscora**, Lam. (Gentian.).
Blancoi, Blanco, l. c. ed. II. Suppl. 591 = Cans. decussata.
dichotoma, Blanco, l. c. ed. I. 510 = Cans. diffusa.

COBRESIA, Pers. Syn. ii. 534 (1807) = **Kobresia**, Willd. (Cyperac.).

COBURGIA, Sweet, Brit. Flow. Gard. Ser. II. t. 17 (1838) = **Stenomesson**, Herb. (Amaryllid.).
acuta, Herb. in Bot. Mag. t. 3865 = S. incarnatum.
angusta, Herb. in Bot. Reg. (1842) Misc. 55 = S. recurvatum.
Belladonna, Herb. in Bot. Mag. sub tt. 2113, 2114 = Amaryllis Belladonna.
blanda, Herb. l. c. = Amaryllis Belladonna.
chachapoyensis, Herb. in Bot. Reg. (1842) Misc. 53 = S. incarnatum.
ciliaris, Herb. in Trans. Hort. Soc. iv. (1822) 181 = Buphane disticha.
coccinea, Herb. in Bot. Reg. (1842) Misc. 54 = S. coccineum.
discolor, Herb. Amaryll. 197 = S. recurvatum.
flava, Hort. Belg. ex Heynh. Nom. ii. 151 = seq.
fulva, Herb. in Bot. Reg. t. 1497 = S. incarnatum.
humilis, Herb. l. c. (1842) Misc. 55 = S. humile.
incarnata, Sweet, Brit. Flow. Gard. Ser. II. t. 17 = S. incarnatum.
Josephinae, Herb. in Trans. Hort. Soc. iv. (1822) 181 = Brunsvigia Josephinae.
laeta, Herb. in Bot. Reg. (1842) Misc. 53 = S. incarnatum.
langensis, Herb. l. c. = S. incarnatum.
lutea, Herb. l. c. 55 = S. recurvatum.
Macleanica, Herb. l. c. = S. recurvatum.
miniata, Lindl. Bot. Reg. (1844) Misc. 13.—Peruv.
multiflora, Herb. in Bot. Mag. sub tt. 2113, 2114 = Brunswigia multiflora.
obragillensis, Herb. in Bot. Reg. (1842) Misc. 54 = S. recurvatum.
pallida, Herb. in Trans. Hort. Soc. iv. (1822) 181 = Amaryllis Belladonna.
praecipitata, Herb. in Bot. Reg. (1842) Misc. 54 = S. coccineum.
pudica, Herb. in Bot. Mag. sub tt. 2113, 2114 = Amaryllis Belladonna.
quitensis, M. Roem. Syn. Ensat. 165.—Ecuador.
Radula, Herb. ex Steud. Nom. ed. II. i. 390 = Brunsvigia Radula.
recurvata, Herb. in Bot. Reg. (1842) Misc. 54 = S. recurvatum.
reticulata, Herb. App. 34 = Hippeastrum reticulatum.
splendens, Herb. Amaryll. 400. t. 47. f. 5 = S. incarnatum.
striatifolia, Herb. App. 34 = Hippeastrum reticulatum.
stylosa, Lindl. in Journ. Hort. Soc. ii. (1847) 310.—Peruv.
taenisphylla, Salisb. Gen. Pl. Fragm. 134.—Hab. ?
trichroma, Herb. Amaryll. 196 = S. incarnatum.
variegata, Herb. l. c. = S. incarnatum.
venusta, Herb. in Bot. Reg. (1842) Misc. 54 = S. coccineum.
versicolor, Herb. l. c. 53. t. 66 = S. incarnatum.

COCCANTHERA, C. Koch & Hanst. Ind. Sem. Hort. Berol. (1855) 17 = **Codonanthe**, Hanst. (Gesnerac.).
Devosiana, C. Koch & Hanst. l. c. = Cod. Devosiana.
Hookeriana, C. Koch & Hanst. l. c. = Cod. Hookerii.

COCCINIA, Wight & Arn. Prod. 347 (1834). *CUCURBITACEAE*, Benth. & Hook. f. i. 827.
CEPHALANDRA, Schrad. in Eckl. & Zeyh. Enum. ii. 280 (1836).
abyssinica, *Cogn. in DC. Monog. Phan.* iii. 536.—Abyssin.
adoënsis, *Cogn. l. c.* 538.—Afr. trop. et austr.
cordifolia, Cogn. l. c. 529 = indica.
decipiens, *Cogn. l. c.* 539.—Afr. trop..
diversifolia, Cogn. l. c. 536 = C. Hartmanniana.
grandis, Voigt, Hort. Suburb. Calc. 59 = indica.
Hartmanniana, *Schweinf. Reliq. Kotschy.* 42. t. 27.—Abyssin.
indica, *Wight & Arn. Prod.* 347.—As. trop.
jatrophaefolia, *Cogn. in DC. Monog. Phan.* iii. 538.—Abyssin.
Loureiriana, M. Roem. Syn. Pepon. 93 = indica.
Mac-Kennii, *Cogn. in DC. Monog. Phan.* iii. 541.—Afr. austr.
Moghadd, Aschers. in Schweinf. Beitr. Fl. Aethiop. 267 = C. Moimoi.
Moimoi, *M. Roem. Syn. Pepon.* 93.—Afr. trop.; Arabia.
Monteroi, *Hort.—Cf. Cat. Gr. Jard. Bot. Bordeaux* (1866).—Afr. trop. ?
palmata, *Cogn. in DC. Monog. Phan.* iii. 540.—Afr. austr.
palmata, M. Roem. Syn. Pepon. 93 = Bryonopsis laciniosa.
palmatisecta, *Kotschy, in Sitzb. Acad. Wien, Math.-Nat.* li. (1865) 360. t. 4.—Abyssin.
quinqueloba, *Cogn. in DC. Monog. Phan.* iii. 533.—Afr. austr.
Schimperi, Naud. in Ann. Sc. Nat. Sér. IV. viii. (1857) 366 = C. Moimoi.
senensis, *Cogn. in DC. Monog. Phan.* iii. 535.—Afr. trop.
sessilifolia, *Cogn. l. c.* 534.—Afr. austr.
Wightiana, M. Roem. Syn. Pepon. 93 = indica.

COCCOBRYON, Klotzsch, in Miq. Syst. Pip. 343 (1843) = **Piper**, Linn.
capense, Klotzsch, l. c. = P. capensis, radicans.

COCCOCERAS, Miq. Fl. Ind. Bat. Suppl. 455 (1860). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 318.
muticum, *Muell. Arg. in Flora*, xlvii. (1864) 470.—Malaya.
plicatum, *Muell. Arg. l. c.* 539.—Burma.
sumatranum, *Miq. Fl. Ind. Bat. Suppl.* 456.—Sumatra.

COCCOCYPSELUM, P. Br. Hist. Jamaic. 144 (1756) (Coccocipsilum). *RUBIACEAE*, Benth. & Hook. f. ii. 73.
BELLARDIA, Schreb. Gen. i. 790 (1789).
CONDALIA, Ruiz & Pav. Prod. 11. t. 2 (1794).
LIPOSTOMA, D. Don, in Edinb. Phil. Journ. (1830) i. 168.
SICELIUM, P. Br. Hist. Jamaic. 144 (1756).
TONTANEA, Aubl. Pl. Gui. i. 108. t. 42 (1775).
aureum, *Cham. & Schlecht. in Linnaea*, iv. (1829) 139.—Bras.
biflorum, Willd. Sp. Pl. i. 618 = Manettia Lygistum.
buxifolium, Spreng. Syst. i. 416 = Fernelia buxifolia.
campanuliflorum, *Cham. & Schlecht. in Linnaea*, iv. (1829) 140.—Bras.
canescens, *Willd. ex Roem. & Schult. Syst.* iii. *Mant.* 130.—Am. austr.
capitatum, Willd. ex DC. Prod. iv. 396 = umbellatum.
ciliatum, *Cham. & Schlecht. in Linnaea*, vi. (1831) 414.—Mexic.
Condalia, Pers. Syn. i. 132 = umbellatum.
cordifolium, *Nees & Mart. in Nov. Act. Nat. Cur.* xii. (1824) 14.—Bras.
dichroolasium, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 76.—Bras.
discolor, *Van Houtte, ex Kunze, in Linnaea*, xxiv. (1851) 192.—Bras. ?
erythrocephalum, *Cham. & Schlecht. in Linnaea*, iv. (1829) 144.—Bras.
geophiloides, *Wawra, in Oestr. Bot. Zeitschr.* xxxi. (1881) 70.—Bras.
glabrum, *Bartl. ex DC. Prod.* iv. 397.—Panama.
herbaceum, Aubl. Pl. Gui. i. 68 = repens.
hirsutum, *Bartl. ex DC. Prod.* iv. 396.—Hab. ?
lanceolatum, *Pers. Syn.* i. 132.—Peruv.

COCCOCYPSELUM:—
metallicum, *Linden, in Belg. Hortic.* xv. (1865) 99.—Guiana.
microphyllum, *Willd. ex Roem. & Schult. Syst.* iii. *Mant.* 130.—Ins. Maurit.
montanum, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 76.—Bras.
nummularifolium, *Cham. & Schlecht. in Linnaea*, iv. (1829) 145.—Bras.; Ind. occ.; Mexic.
obovatum, *Pers. Syn.* i. 132.—Peruv.
ovatum, *Cham. & Schlecht. in Linnaea*, iv. (1829) 141.—Bras.
pedunculare, *Cham. & Schlecht. l. c.* 142.—Bras.
Pseudo-Tontanea, *Griseb. Fl. Brit. W. Ind.* 322.—Ind. occ.
pulchellum, *Cham. in Linnaea*, ix. (1834) 238.—Bras.
repens, H. B. & K. Nov. Gen. et Sp. iii. 403 = canescens.
repens, *Sw. Prod. Veg. Ind. Occ.* 31.—Ind. occ.
sessile, *Pers. Syn.* i. 132.—Peruv.
spicatum, H. B. & K. Nov. Gen. et Sp. iii. 406 = Gonzalea spicata.
Tontanea, *H. B. & K. l. c.*—Am. austr.
umbellatum, *Poir. Encyc. Suppl.* ii. 307.—Peruv.
uniflorum, Willd. Sp. Pl. i. 618 = Fernelia obovata.
veronicoides, *Presl, Bot. Bemerk.* 87.—Am. trop.
virgatum, *Lam. Illustr.* i. 270. n. 1457 = Manettia picta.
violaceum, Desf. Cat. Hort. Par. ed. III. 404 = capituliflorum.

COCCODERMA, Miers, in Lindl. Veg. Kingd. 309 b (1847), nomen (Menisperm.).

COCCOLOBA, Linn. Syst. ed. X. 1007 (1759). *POLYGONACEAE*, Benth. & Hook. f. iii. 102.
LYPERODENDRON, Willd. ex Meissn. in DC. Prod. xiv. 168, in syn. (1857).
NAUCOREPHES, Rafin. Fl. Tellur. ii. 34 (1836).
SCHLOSSERIA, Mill. ex Steud. Nom. ed. II. ii. 531 (1841).
acrostichoides, *Cham. in Linnaea*, viii. (1833) 132.—Bras.
acuminata, *H. B. & K. Nov. Gen. et Sp.* ii. 176.—N. Granat.
alagoënsis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 260.—Bras.
alnifolia, Casar. Nov. Stirp. Bras. Dec. viii. 71 = populifolia.
appressa, Meissn. ex Steud. Nom. ed. II. i. 390 = Polygonum adpressum.
armata, *Wright, ex Griseb. Cat. Pl. Cub.* 283.—Cuba.
asiatica, Lour. Fl. Cochinch. 239 = Polygonum asiaticum.
australis, Forst. f. Prod. 29. n. 176 = Muehlenbeckia australis.
barbadensis, *Jacq. Enum. Pl. Carib.* 37; *Linn. Syst.* ed. XII. 277.—Am. trop.
barbadensis, Willd. Sp. Pl. ii. 459 = caracasana.
Blanchetiana, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 257 = ochreolata.
bracteolosa, *Meissn. in Mart. Fl. Bras.* v. I. 30.—Bras.
brasiliensis, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) I. 30.—Bras.
brasiliensis, Spreng. Syst. ii. 252 (= Hedyosmi sp.).—Bras.
bullata, Hort. ex Meissn. in DC. Prod. xiv. 152 = rugosa.
calobotrys, Meissn. l. c. 157 = coriacea.
Candolleana, *Meissn. in Mart. Fl. Bras.* v. I. 41.—Bras.
caracasana, *Meissn. in DC. Prod.* xiv. 157.—Venezuela.
carinata, Ruiz, ex Meissn. l. c. 150 = Muehlenbeckia leptobotrys.
cordata, *Cham. in Linnaea*, viii. (1833) 133.—Bras.
cordifolia, Meissn. in Mart. Fl. Bras. v. I. 37 = laevis.
coriacea, *A. Rich. Fl. Cub. Fanerog.* ii. 184.—Cuba.
coriacea, Willd. ex Meissn. in DC. Prod. xiv. 155 = obovata.
coronata, *Jacq. Enum. Pl. Carib.* 19.—Ind. occ.
costata, Mart. ex Meissn. in DC. Prod. xiv. 153 = populifolia.
costata, *Wright, in Sauv. Fl. Cub.* 139.—Cuba.
crescentiaefolia, *Cham. in Linnaea*, viii. (1833) 134.—Bras.
crispata, Buch.-Ham. in Roxb. Fl. Ind. ii. 292 = Polygonum chinense.

COCCOLOBA:—
Cruegerii, *Griseb. Fl. Brit. W. Ind.* 710.—Ind. occ.
cubensis, *Meissn. in DC. Prod.* xiv. 162.—Cuba.
cujabensis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 259.—Bras.
cymosa, Lour. Fl. Cochinch. 240 = Polygonum cymosum.
declinata, *Mart. ex Meissn. in Mart. Fl. Bras.* v. I. 30.—Bras.
densifrons, *Mart. ex Meissn. l. c.* 26.—Bras.
dioica, Steud. Nom. ed. II. i. 390 = Muehlenbeckia sagittifolia.
diversifolia, *Jacq. Enum. Pl. Carib.* 19.—Ins. S. Doming.
emarginata, *Jacq. l. c.* 37 (sp. dub.).—Am. austr.
excelsa, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 624.—Am. trop.
excoriata, *Linn. Syst.* ed. X. 1007.—Ind. occ.
fagifolia, *Jacq. Hort. Schoenb.* iii. 55. t. 352.—Am. austr.
fasciculata, Meissn. in Kjoeb. Vidensk. Meddel. (1870) 128 = longependula.
fasciculata, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 258 = crescentiaefolia.
fastigiata, *Meissn. in Mart. Fl. Bras.* v. I. 34.—Bras.
ferruginea, Endl. Cat. i. 274 = excoriata.
flavescens, *Jacq. Enum. Pl. Carib.* 19.—Ins. S. Doming.
floridana, Meissn. in DC. Prod. xiv. 165 = laurifolia.
Gardneri, Meissn. in Mart. Fl. Bras. v. I. 36 = crescentiaefolia.
Goudotiana, Meissn. l. c. 35 = fastigiata.
Goudotiana, Wedd. in Ann. Sc. Nat. Sér. III. xiii. (1849) 260 = obovata.
gracilis, *H. B. & K. Nov. Gen. et Sp.* ii. 176.—Peruv.
grandifolia, *Jacq. Enum. Pl. Carib.* 19.—Am. trop.
grandis, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 624 = latifolia.
guatemalensis, *Teijsm. & Binn. Cat. Hort. Bog.* 383, nomen.—Hab.?
guianensis, *Meissn. in Linnaea*, xxi. (1848) 254.—Guiana.
Humboldtii, *Meissn. in DC. Prod.* xiv. 163.—Bras.
ilheensis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 258.—Bras.
indica, Wight, ex Meissn. in DC. Prod. xiv. 131 = Polygonum chinense.
japurana, Meissn. in Mart. Fl. Bras. v. I. 25 = (Alsodeiae sp.).—Bras.
Klotzschiana, *Meissn. in DC. Prod.* xiv. 155.—Ins. S. Thomas.
Kunthiana, Meissn. l. c. 166 = pyrifolia.
laevis, *Casar. Nov. Stirp. Bras. Dec.* viii. 71.—Bras.
latifolia, *Lam. Encyc.* vi. 61.—Am. austr.
latifolia, Poepp. ex Meissn. in DC. Prod. xiv. 151 = polystachya.
laurifolia, *Jacq. Hort. Schoenbr.* iii. 9. t. 267.—Am. trop.
leoganensis, Jacq. Enum. Pl. Carib. 19 = uvifera.
leptostachya, *Benth. Bot. Voy. Sulph.* 159.—Am. austr.
longependula, *Mart. ex Meissn. in Mart. Fl. Bras.* v. I. 27.—Bras.
longifolia, *Fisch. Hort. Gorenk.* 25.—Ind. occ.
lucidula, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 627.—Guiana.
macrantha, Hook. Bot. Mag. sub t. 4536 = seq.
macrophylla, Desf. Tabl. ed. I. 38 = rugosa.
magnifolia, Hort. ex Meissn. in DC. Prod. xiv. 152 = rugosa.
manzinellensis, *Beurl. in Vet. Akad. Handl. Stockl.* 1854 (1856) 142.—Panama.
marginata, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 626.—Bras.
Martii, Meissn. in Mart. Fl. Bras. v. I. 37 = nitida.
membranacea, Klotzsch, in Linnaea, xiv. (1831) 289, nomen = ilheensis.
microneura, *Meissn. in DC. Prod.* xiv. 163.—Demerara.
microphylla, *Griseb. Cat. Pl. Cub.* 62.—Cuba.
microstachya, *Willd. Sp. Pl.* ii. 459.—Ind. occ.
mollis, Casar. Nov. Stirp. Bras. Dec. viii. 72 = polystachya.
monoica, Ruiz, ex Meissn. in DC. Prod. xiv. 149 = . = Muehlenbeckia tamnifolia.
Moritzii, Klotzsch, ex Meissn. in Mart. Fl. Bras. v. I. 28 = ovata.

COCCOLOBA:—
nitida, Desf. Tabl. ed. II. 46 = fagifolia.
nitida, *H. B. & K. Nov. Gen. et Sp.* ii. 176.—N. Granat.
nitida, Mart. ex Meissn. in DC. Prod. xiv. 164 = ovata.
nivea, Jacq. Enum. Pl. Carib. 19 = excoriata.
nutans, *H. B. & K. Nov. Gen. & Sp.* ii. 176.—Peruv.
nymphaeifolia, Linden, Cat. n. 13 (1858) = peltata.
obovata, *H. B. & K. Nov. Gen. et Sp.* ii. 175.—N. Granat.
obtusifolia, Jacq. Enum. Pl. Carib. 19 = microstachya.
ochreolata, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 259.—Am. austr.
orbicularis, Lodd. Cat. ex Loud. Hort. Brit. 159 = Muehlenbeckia orbicularis.
ovata, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 627.—Guiana.
padiformis, *Meissn. in DC. Prod.* xiv. 166.—Venezuela.
pallida, *Wright, ex Griseb. Cat. Pl. Cub.* 61.—Cuba.
paniculata, Meissn. in Mart. Fl. Bras. v. I. 43 = polystachya.
paraënsis, *Meissn. l. c.* 38.—Bras.
parimensis, Benth. in Hook. Lond. Journ. Bot. iv. (1845) 626 = excelsa.
parvifolia, *Nutt. N. Am. Sylv.* iii. 25. t. 89 (sp. dub.).—Am. bor.
parvifolia, Poir. Encyc. vi. 64 = microstachya.
peltata, *Schott, in Spreng. Syst.* iv. *Cur. Post.* 405.—Bras.
peltigera, Meissn. in Mart. Fl. Bras. v. I. 39 = peltata.
pendula, Salzm. ex Meissn. in DC. Prod. xiv. 159 = nitida.
Persicaria, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 256.—Bolivia.
peruviana, Willd. ex Meissn. in DC. Prod. xiv. 163 = gracilis.
pipericarpa, *Mart. ex Meissn. l. c.* 161.—Bras.
plantaginea, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 257.—Bras.
platyclada, F. Muell. ex Hook. Bot. Mag. t. 5382 = Muehlenbeckia platyclada.
Plumieri, *Griseb. Fl. Brit. W. Ind.* 162.—Ind. occ.
polystachya, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 261.—Bras.
populifolia, *Wedd. l. c.* 257.—Bras.
pubescens, Linn. Syst. ed. X. 1007 = grandifolia.
punctata, Linn. Sp. Pl. ed. II. 523 = coronata.
pyrifolia, *Desf. Tabl.* ed. II. 46.—Ind. occ.
racemulosa, *Meissn. in Mart. Fl. Bras.* v. I. 30.—Bras.
ramosisissima, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 258.—Bras.
retusa, *Griseb. Cat. Pl. Cub.* 61.—Ind. occ.
rheifolia, Desf. Tabl. ed. II. 46 = latifolia.
rigida, Meissn. in Mart. Fl. Bras. v. I. 29 = parvifolia.
rigida, Willd. ex Meissn. in DC. Prod. xiv. 163 = Humboldtii.
rosca, *Meissn. in Mart. Fl. Bras.* v. I. 33.—Bras.
rotundifolia, *Meissn. in DC. Prod.* xiv. 154.—Ins. S. Doming.
rubescens, Linn. Sp. Pl. ed. II. 523 = grandifolia.
rubiginosa, Mart. ex Meissn. in DC. Prod. xiv. 160 = acrostichoides.
rufescens, *Wright, in Sauv. Fl. Cub.* 138.—Cuba.
rugosa, *Desf. Cat. Hort. Par.* ed. III. 389.—Ind. occ.
sagittata, Poir. Encyc. vi. 64 = Muehlenbeckia sagittifolia.
sagittifolia, Ortega, Hort. Matr. v. 60 = Muehlenbeckia sagittifolia.
salicifolia, *Wedd. in Ann. Sc. Nat.* Ser. III. xiii. (1849) 259.—Bras.
scandens, Casar. Nov. Stirp. Bras. Dec. viii. 70 = sticticaulis.
scandens *Forsyth, ex Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 625.—Ind. occ.
scandens, Poepp. ex Meissn. in DC. Prod. xiv. 154 = peltata.
Schomburgkiana, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 628.—Guiana.
Schomburgkii, *Meissn. in Linnaea*, xxi. (1848) 265.—Guiana.

COCCOLOBA:—
sticticaulis, *Wedd. in Ann. Sc. Nat.* Sér. III. xiii. (1849) 261.—Bras.
striata, *Benth. in Hook. Lond. Journ. Bot.* iv. (1845) 626.—Bras.
stricta, Klotzsch, in Schomb. Faun. et Fl. Guian. 934 = lucidula.
strobilulifera, *Meissn. in Mart. Fl. Bras.* v. 1. 25.—N. Granat.
Swartzii, *Meissn. in DC. Prod.* xiv. 159.—Ind. occ.
tenuifolia, Griseb. Fl. Brit. W. Ind. 162 = retusa.
tenuifolia, *Linn. Syst.* ed. X. 1007; *Amoen. Acad.* v. 397 (sp. dub.).—Jamaic.
Totnea, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 74 = Polygonum molle.
uvifera, *Linn. Syst.* ed. X. 1007.—Am. trop.
Vellosiana, Casar. Nov. Stirp. Bras. Dec. viii. 70 = crescentiaefolia.
venosa, Linn. Syst. ed. X. 1007 = coronata.
virens, Lindl. Bot. Reg. t. 1816 = coronata.
Warmingii, *Meissn. in Kjoeb. Vidensk. Meddel.* (1870) 128.—Bras.
Zebra, *Griseb. Fl. Brit. W. Ind.* 162.—Jamaic.

COCCOLOBIS, P. Br. Hist. Jamaic. 209 (1756) = **Coccoloba**, Linn.

COCCOMELIA, Reinw. Cat. Gew. Buitenz, 110 (1823); et in Syll. Ratisb. ii. (1828) 5 = **Baccaurea**, Lour. (Euphorbiac.).
racemosa, Reinw. ll. cc. 110, 6 = B. racemosa.

COCCONERION, Baill. Adansonia, xi. (1873) 87. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 323.
Balansae, *Baill. l. c.* 88.—N. Caled.
minus, *Baill. l. c.*—N. Caled.

COCCOS, Gaertn. Fruct. i. 15 (1788) = **Cocos**, Linn. (Palm.).

COCCOSIPSILUM, Sw. Prod. Veg. Ind. Occ. 31 (1788) = **Coccocypselum**, P. Br. (Rubiac.).

COCCOSPERMA, Klotzsch, in Linnaea, xii. (1838) 215 = **Salaxis**, Salisb. (Ericac.).
Forbesianum, Klotzsch, l. c. = S. hexandra.
rugosum, Klotzsch, l. c. = S. rugosa.

COCCULIDIUM, Spach, Hist. Vég. Phan. viii. 16 (1839) = **Cocculus**, DC. (Menisp.).
populifolium, Spach, l. c. 17 = Cocculus carolinus.

COCCULUS, DC. Syst. i. 515 (1818). *MENISPERMACEAE*, Benth. & Hook. f. i. 36.
ADENOCHETON, Fenzl, in Flora, xxv. (1842) 312.
ANDROPHYLAX, Wendl. Beobacht. 38 (1798).
BAUMGARTIA, Moench, Meth. 650 (1794).
CEBATHA, Forsk. Fl. Aegypt. Arab. 171 (1775).
COCCULIDIUM, Spach, Hist. Vég. Phan. viii. 16 (1839).
COLUMBRA, Comm. ex Endl. Gen. 826 (1840).
DIPLOCLISIA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 42.
EPIBATERIUM, Forst. Char. Gen. 107. t. 54 (1776).
EPIBIASTRUM, Scop. Introd. 253 (1777).
FERRANDIA, Gaudich. Freyc. Voy. Bot. 477 (1826).
GALLOA, Hassk. Cat. Hort. Bog. Alt. 172 (1844).
HOLOPEIRA, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 42.
LEAEBA, Forsk. Fl. Aegypt. Arab. 172 (1775).
LEGNEPHORA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. xix. (1867) 88.
NEPHROIA, Lour. Fl. Cochinch. 565 (1790).
QUINIO, Schlecht. in Linnaea, xxvi. (1854) 732.
SELWYNIA, F. Muell. Fragm. iv. 153 (1864).
WENDLANDIA, Willd. Sp. Pl. ii. 275 (1799).
Abuta, Kostel. Allg. Med.-Pharm. Fl. ii. 501 = Abuta rufescens.
acuminatus, DC. Syst. i. 527 = Tiliacora racemosa.
amazonum, *Mart. ex Herberger, in Buchn. Rep. Pharm.* xxxvi. (1830) 353.—Bras.
angustifolius, Hassk. Cat. Pl. Hort. Bog. Alt. 172 = laurifolius.
Aristolochiae, DC. Syst. i. 520 = villosus.
Bakis, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. i. 12. t. 4 = Tinospora Bakis.

COCCULUS:—
Balfourii, *Schweinf. ex Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 500.—Ins. Socotra.
banisteriaefolius, A. Rich. ex St. Hil. & Tul. in Ann. Sc. Nat. Sér. II. xvii. (1842) 136 = Pachygone oblongifolia.
bantamensis, *Blume, Bijdr.* 26.—Java.
Blumeanus, Wall. Cat. n. 4971, partim = Coscinium fenestratum.
brachystachyus, *DC. Syst.* i. 528.—Ins. Timor.
Burmanni, DC. l. c. 517 = Cyclea Burmanni.
calophyllus, Wall. ex Voigt, Hort. Suburb. Calc. 331, nomen; et ex Mast. in Hook. f. Fl. Brit. Ind. i. 578 = Erythropalum scandens.
carolinus, *DC. Syst.* i. 524.—Am. bor.
Cebatha, *DC. l. c.* 526.—Arabia.
Chondodendrum, *DC. l. c.* 522.—Am. austr.
cinerascens, St. Hil. Fl. Bras. Mer. i. 59 = Chondodendron platyphyllum.
cinereus, *Zoll. & Moritzi, Syst. Verz. Zoll.* 38.—Java.
convolvulaceus, DC. Syst. i. 518 = Tinospora cordifolia.
convolvulaceus, *Endl. ex Walp. Rep.* i. 95.—Peruv.
cordifolius, DC. Syst. i. 517 = Tinospora cordifolia.
coriaceus, Blume, Bijdr. 25.—Java.
corymbosus, *Blume, l. c.* 24.—Java.
Cotoneaster, *DC. Syst.* i. 525.—Am. austr.
crispus, DC. l. c. 521 = Tinospora crispa.
cuneatus, *Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 50.—China.
cuspidatus, Wall. Cat. n. 4960 = Limacia cuspidata.
cynanchoides, *Presl, Rel. Haenk.* ii. 79.—Zeylan.
diantherus, Hook. & Arn. Bot. Beech. Voy. 167 = Thunbergii.
dichopetalus, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) II. 280.—Japon.
dichrous, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 46.—Bras.
diversifolius, *DC. Syst.* i. 523.—Mexic.
domingensis, DC. l. c. 528 = Pachygone domingensis.
ellipticus, DC. l. c. 526 = C. Leaeba.
enneandra, Eichl. in Flora, xlvii. (1864) 391.—Peruv.
Epibaterium, *DC. Syst.* i. 530.—Ins. Pacif.
Ferrandianus, *Gaudich. in Freyc. Voy. Bot.* 477.—Ins. Sandvic.
Fibraurea, DC. Syst. i. 525 = Fibraurea tinctoria.
filipendula, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 43.—Bras.
Finlaysonianus, Wall. Cat. n. 4974 = Stephania rotunda.
flavescens, *DC. Syst.* i. 520.—Ins. Molucc.
flavicans, *Wall. Cat.* n. 4976.—Ins. Penang.
Forsteri, *DC. Syst.* i. 517.—Hab.?
glaber, Wight & Arn. Prod. i. 13 = C. Leaeba.
glaucescens, *Blume, Bijdr.* 25.—Burma; Java.
glaucus, *DC. Syst.* i. 521.—Ins. Molucc.
gomphioides, *DC. l. c.* 530.—Madag.
hastatus, DC. l. c. 522 = villosus.
hexagynus, *Colebr. in Trans. Linn. Soc.* xiii. (1822) 63.—China.
Hookerianus, *F. Muell. ex Benth. in Journ. Linn. Soc.* v. *Suppl.* ii. (1861) 52.—Austral.
Imene, Mart. in Flora, xxiv. (1841) II. *Beibl.* 44 = Anomospermum japurense.
incanus, Colebr. in Trans. Linn. Soc. xiii. (1822) 57. t. 17 = Pericampylus incanus.
japonicus, *DC. Syst.* i. 516.—Japon.
japurensis, Mart. in Flora, xxiv. (1841) II. Beibl. 44 = Anomospermum japurense.
lacunosus, DC. Syst. i. 519 = Anamirta Cocculus.
laevigatus, Mart. in Flora, xxiv. (1841) II. Beibl. 45 = Abuta concolor.
laevis, Wall. Cat. n. 4975 = C. Leaeba.
lanuginosus, *Blume, Bijdr.* 24.—Java.
laurifolius, *DC. Syst.* i. 520.—Reg. Himal.; Japon.
Leaeba, *DC. l. c.* i. 529.—As. trop. occ.; Afr. trop.
leptostachyus, *DC. l. c.* 528.—Ins. Timor.
Limacia, *DC. l. c.* 526.—Cochinch.
Linnaeanus, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 62 (err. typ. 61)—Burma.
lobata, Mart. ex Eichl. in Mart. Fl. Bras. xiii. 1. 169 = Disciphania lobata.
longifolius, *Decne. in Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 84.—Ins. Timor.
lucidus, Teijsm. & Binn. in Nat. Tijdschr. Nederl. Ind. iv. (1853) 397 = Antitaxis fasciculata.
macranthus, Hook. f. in Hook. Ic. Pl. t. 759 = Jateorhiza strigosa.

COCCULUS:—
macrocarpus, *Wight & Arn. Prod.* 13.—Ind. or.; Burma; China.
macrophyllus, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 134.—Bras.
malabaricus, DC. Syst. i. 518 = Tinospora malabarica.
Martii, *St. Hil. & Tul. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 135.—Bras.
megaspermus, *Voigt, Hort. Suburb. Calc.* 331.—Silhet.
membranaceus, Wall. Cat. n. 4967 = Cissampelos Pareira.
milleflorus, *DC. Syst.* i. 530.—Madag.
mollis, *Wall. Cat.* n. 4973.—Reg. Himal.
Moorei, F. Muell. Fragm. i. 162 = Pericampylus incanus.
Nephroia, *DC. Syst.* i. 531.—Cochinch.
nudiflorus, *Griff. Notul.* iv. 307.—Burma.
oblongifolius, *DC. Syst.* i. 529.—Am. bor. occ.
oblongifolius, Mart. in Flora, xxiv. (1841) II. Beibl. 43 = Pachygone oblongifolia.
oblongus, Wall. Cat. n. 4963 = Limacia oblonga.
orbiculatus, DC. Syst. i. 523 = Cissampelos Pareira.
ovalifolius, Blume, Bijdr. 25 = umbellatus.
ovalifolius, DC. Syst. i. 526 = Thunbergii.
Pahni, Mart. in Flora, xxiv. (1841) II. Beibl. 45 = Anomospermum japurense.
palmatus, DC. Syst. i. 522 = Jateorhiza Columba.
palmatus, Hook. Bot. Mag. tt. 2970–71 = Jateorhiza Miersii.
paniculiger, Mart. in Flora, xxiv. (1841) II. Beibl. 43 = Pachygone domingensis.
pauper, *Griseb. in Goett. Abh.* vii. (1856–57) 162.—Am. trop.
peltatus, DC. Syst. i. 516 = Cyclea peltata.
petiolaris, Wall. Cat. n. 4964 = Tinomiscium petiolare.
planiflorus, Wall. l. c. n. 4961 = Pycnarrhena planiflora.
platyphyllus, St. Hil. Fl. Bras. Mer. i. 59 = Chondodendron platyphyllum.
Plukenetii, DC. Syst. i. 520 = Pachygone ovata.
polycarpus, Wall. Cat. n. 4958 = Tiliacora racemosa.
populifolius, DC. Syst. i. 519 = Anamirta Cocculus.
radiatus, DC. l. c. 527 = Tiliacora racemosa.
recisus, Miers, Contrib. iii. 258 = C. Leaeba.
reticulatus, Mart. in Flora, xxiv. (1841) II. Beibl. 44 = Anomospermum reticulatum.
rimosus, *Blume, Bijdr.* 24.—Java.
rotundifolius, *DC. Syst.* i. 517.—Hab.?
Roxburghianus, DC. l. c. 516 = Stephania rotunda.
sagittaefolius, *Miers, Contrib.* iii. 255.—Am. bor.
Selwynii, F. Muell. Fragm. iv. 153 = Selwynia laurina.
sepium, Colebr. in Trans. Linn. Soc. xiii. (1822) 58. t. 6 = villosus.
smilacinus, F. Muell. ex Benth. in Journ. Linn. Soc. v. Suppl. ii. (1861) 52 = Tinospora smilacina.
suberosus, DC. Syst. i. 519 = Anamirta paniculata.
tamoides, DC. l. c. 521 = Chondodendron tamoides.
tetrandrus, *Zipp. ex Span. in Linnaea*, xv. (1841) 164, nomen.—Malaya.
Thunbergii, *DC. Syst.* i. 524.—Japon.
tomentosus, Colebr. in Trans. Linn. Soc. xiii. (1822) 59 = Tinospora tomentosa.
tomentosus, Mart. ex Eichl. in Mart. Fl. Bras. xiii. 1. 175 = Abuta rufescens.
triandrus, Colebr. in Trans. Linn. Soc. xiii. (1822) 64 = Limacia triandra.
triangularis, *Zipp. ex Span. in Linnaea*, xv. (1841) 164, nomen.—Malaya.
triflorus, *DC. Syst.* i. 529.—Java.
trilobus, *DC. l. c.* 522.—Japon.
umbellatus, *Steud. Nom.* ed. II. i. 392.—Java.
urophyllus, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 45.—Bras.
velutinus, Wall. Cat. n. 4970 = Limacia velutina.
verrucosus, Wall. l. c. n. 4966 A, B = Tinospora crispa.
verrucosus, Wall. l. c. n. 4966 C, D, E = Tinospora cordifolia.
villosus, *DC. Syst.* i. 525.—Ind. or.; Afr. trop.
villosus, Wall. Cat. n. 4957, partim = Cissampelos Pareira.
Wightianus, Wall. l. c. n. 4959 A = Pachygone ovata.

COCCUS, Linn. Mus. Cliff. 11 (1736); Gen. ed. I. 355 (1737) = **Cocos**, Linn. (Palm.).

COCCYGANTHE, Reichb. Fl. Germ. Excurs. xvi. 55 (1844) = **Lychnis**, Linn. (Caryophyll.).
Flos-cuculi, Reichb. l. c. = L. Flos-cuculi.
pratensis, Schur, Enum. Pl. Transs. 107 = L. Flos-cuculi.

COCHLANTHERA, Choisy, in Mém. Soc. Phys. Genèv. xii. (1851) 426 = **Clusia**, Linn. (Guttifer.).
lanceolata, Choisy, l. c.—Venezuela.

COCHLANTHUS, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 78. *ASCLEPIADACEAE.*
socotranus, *Balf. f. l. c.* 79.—Ins. Socotra.

COCHLEANTHES, Rafin. Fl. Tellur. 45 (1836) = **Zygopetalum**, Hook. (Orchid.).
fragrans, Rafin. l. c. = Z. cochleare.

COCHLEARIA, Tourn. ex Linn. Syst. ed. I (1735). *CRUCIFERAE*, Benth. & Hook. f. i. 75.
ARMORACIA, Riv. ex Gaertn. Mey. & Scherb. Fl. Wett. ii. 426 (1800).
CHRYSOCHAMELA, Boiss. Fl. Orient. i. 313 (1867).
GONYCLISIA, Dulac, Fl. Hautes-Pyr. 191 (1867).
IONOPSIDIUM, DC. ex Steud. Nom. ed. II. i. 392 (1840).
KERNERA, Medic. Pflanzeng. 71 (1792).
RAPHANIS, Dod. ex Moench, Meth. 267 (1794).
RHIZOBOTRYA, Tausch, in Flora, xix. (1836) 33.
TAPHROSPERMUM, C. A. Mey. in Ledeb. Fl. Alt. iii. 172 (1831).
TROPHOSPERMUM, Walp. Rep. i. 171 (1842).
acaulis, *Desf. Fl. Atlant.* ii. 69.—Lusitan.; Afr. bor.
alpina, Sweet, Hort. Brit. ed. I. 22 = officinalis.
altaica, Schlecht. ex DC. Prod. i. 173 = microcarpa.
alyssoides, DC. l. c. 172 = flava.
amphibia, *Ledeb. Fl. Ross.* i. 160.—Soongaria.
anglica, *Linn. Syst.* ed. X. 1128.—Reg. bor. et arct.
aphragmodes, *Boiss. Diagn.* Ser. I. vi. 15.—Persia.
aquatica, Eaton, Man. ed. V. 181 = Nasturtium lacustre?
aquatica, *C. A. Mey. in Ledeb. Fl. Alt.* iii. 88.—As. bor.
arctica, Schlecht. in DC. Syst. ii. 367 = anglica.
Armoracia, *Linn. Sp. Pl.* t. 648.—Enrop.
Armoracia, Pall. ex DC. Syst. ii. 362 = microcarpa.
Aucheri, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 168.—Armenia.
auriculata, *Lam. Encyc.* ii. 165.—Europ.
austriaca, *Ledeb. Fl. Ross.* i. 160.—Rossia.
austriaca, Vis. Fl. Dalm. iii. 122 = Nasturtium austriacum.
batava, *Dum. Fl. Belg. Prod.* 120.—Batavia.
brevicaulis, *Facch. in Zeitschr. Ferdin. Tirol.* III. (1854) 79.—Tyrol.
Caisia, Wall. Cat. n. 4802 = flava.
Camelinae, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 172.—Persia.
Camelinae, *Ledeb. Fl. Ross.* i. 160, in nota.—Manchuria.
campylocarpa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 171.—Persia.
cordifolia, *Turcz. ex Bunge, in Mém. Sav. Etr. Acad. Pétersb.* ii. (1835) 579.—Sibir. baical.
Coronopus, Linn. Sp. Pl. 648 = Senebiera Coronopus.
Crantzii, *Crantz, Class. Crucif. Emend.* 100.—Europ.
crassifolia, Willd. ex DC. Prod. i. 173 = macrocarpa.
danica, Gunn. Fl. Norv. 72. n. 197 = officinalis.
danica, *Linn. Sp. Pl.* 647.—Reg. bor. et arct.
decipiens, *Willk. Sert. Fl. Hispan.* 12.—Mont. Pyren.
Draba, Linn. Syst. ed. X. 1129 = Lepidium Draba.
drabaccarpa, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 169.—Persia.
fenestrata, R. Br. in App. Voy. Ross, (1819) p. cxliii = officinalis.
flagrans, Gilib. Fl. Lituan. ii. 51 = officinalis.
flava, *Buch.-Ham. ex Roxb. Hort. Beng.* 48, nomen.—Ind. or.
foetida, Schkuhr, Baier. Fl. ii. 185 = Camelina foetida.
glastifolia, Falk, Beitr. ii. 215 = Hymenophysa pubescens.
glastifolia, *Linn. Sp. Pl.* 648.—Europ.
glastifolia, Pall. ex DC. Syst. ii. 369 = Draba salsa.
glaucophylla, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 169.—Persia.
globosa, *Ledeb. Fl. Ross.* i. 159.—Dahuria.
grandiflora, *DC. Syst.* ii. 363.—Sibiria.
groenlandica, Gunn. Fl. Norv. 72. n. 196 = anglica.

COCHLEARIA:—
groenlandica, Linn. Sp. Pl. 647 = officinalis.
groenlandica, Willd. ex DC. Prod. i. 173 = lenensis.
hastata, Moench, Meth. Suppl. 87 = danica.
hederacea, S. F. Gray, Nat. Arr. Brit. Pl. ii. 696 = danica.
heterophylla, Cav. Desc. 405 = Camelina dentata.
heterophylla, Fisch. Hort. Gorenk. ed. II. (1812) 50, nomen = seq.
heterophylla, Schlecht. in DC. Syst. ii. 372 = sisymbrioides.
himalaica, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 154.—Reg. Himal.
humifusa, Michx. Fl. Bor. Am. ii. 27 = Senebiera didyma.
integrifolia, DC. Syst. ii. 368 = Eutrema alpestre.
kamtschatica, Schlecht. ex DC. Prod. i. 173 = officinalis.
lancifolia, Stokes, Bot. Mat. Med. iii. 437 = C. Armoracia.
lapathifolia, Gilib. Fl. Lituan. iv. 53 = C. Armoracia.
lenensis, *Adams, ex Fisch. in DC. Syst.* ii. 367.—Sibiria.
Linnaei, Griewank, ex Aschers. in Fl. Prov. Brandenb. 54 = officinalis.
longifolia, *Medic. in Usteri, Neu. Ann.* ii. 41.—Europ.
lyrata, *Sibth. & Sm. Fl. Graec. Prod.* ii. 8; *Fl. Graec.* t. 619.—Sicilia.
macrocarpa, *Waldst. & Kit. Pl. Rar. Hung.* ii. 101. t. 184.—Hungar.; Transsylv.
mexicana, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 320.—Mexic.
microcarpa, *DC. Syst.* ii. 362.—Sibir.
minima, Steud. Nom. ed. II. i. 392 = danica.
natans, C. A. Mey. in Ledeb. Fl. Alt. iii. 90 = amphibia.
nilotica, Delile, Fl. Egypte, 245 = Senebiera nilotica.
oblongifolia, DC. Syst. ii. 363 = officinalis.
officinalis, Lapeyr. Hist. Abr. Pl. Pyr. 368 = pyrenaica.
officinalis, *Linn. Sp. Pl.* 647.—Reg. bor. et arct.
olyssiponensis, Brot. Fl. Lusit. i. 571 = acaulis.
ovalifolia, Stokes, Bot. Mat. Med. iii. 436 = anglica.
planisiliqua, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 169.—Persia.
polysperma, Costa, ex Willk. in Linnaea, xxx. (1859–60) 86 = Kernera polysperma.
Pseud-Armoracia, Czern. ex Trautv. in Act. Hort. Petrop. viii. (1883) 93 = macrocarpa?
pusilla, Brot. Phyt. 100 = acaulis.
pyrenaica, DC. Syst. ii. 365 = officinalis.
renifolia, Stokes, Bot. Mat. Med. iii. 435 = officinalis.
repanda, *Medic. in Usteri. Neu. Ann.* ii. 42.—Europ.
repens, Lam. Fl. Fr. ii. 473 = Senebiera Coronopus.
rhizobotrya, *Walp. Rep.* i. 153.—Austria.
rotundifolia, S. F. Gray, Nat. Arr. Brit. Pl. ii. 695 = officinalis.
rusticana, Lam. Fl. Fr. ii. 471 = C. Armoracia.
sagittaefolia, *Desv. Journ. Bot.* iii. (1812) 182.—Hab.?
sagittata, Crantz, Class. Crucif. Emend. 99 = Neslia paniculata.
salsa, *Schlecht. ex DC. Syst.* ii. 369.—Sibiria?
sativa, Cav. Desc. 405 = Camelina sativa.
saxatilis, *Linn. Sp. Pl.* 648.—Europ.
saxifragaefolia, DC. Syst. ii. 370 = Graellsia saxifragaefolia.
scapillora, *Hook. f. & Thoms. in Journ. Linn. Soc.* v. (1861) 154.—Reg. Himal.
Sempervivum, *Boiss. & Bal. Diagn.* Ser. II. v. 28.—As. Min.
septentrionalis, DC. Prod. i. 174 = Draba hyperborea.
septentrionalis, Schlecht. in DC. Syst. ii. 355 = officinalis.
sibirica, Willd. Sp. Pl. iii. 450 = Sobolewskia lithophila.
siliquosa, Schlecht. in DC. Syst. ii. 369 = Draba hyperborea.
sisymbroides, *DC. l. c.* 368.—Sibir. arct.
spathulata, *Schlecht. in DC. l. c.* 369 = Draba hyperborea.
Szovitzii, *Boiss. Fl. Orient.* i. 248.—Persia.
tenuifolia, Salisb. Prod. 267 = Senebiera Coronopus.
tridactylites, *Banks, ex DC. Syst.* ii. 367.—Am. arct.
variifolia, Salisb. Prod. 267 = C. Armoracia.
velutina, *DC. Syst.* ii. 370.—Mesopot.
violacea, *Boiss. in Ann. Sc. Nat.* Ser. II. xvii. (1842) 171.—Persia.

COCHLEARIA:—
Wahlenbergii, Rupr. Beitr. Pfl. Russ. Reich. ii. 21 = anglica.
Wunderlichii, *C. A. Mey. ex Claus, in Beitr. Pfl. Russ. Reich.* viii. 203.—Rossia.

COCHLEATA, Riv. ex Rupp. Fl. Jen. ed. Hall. 264 (1745) = **Medicago**, Linn. (Legum.).
intertexta, Medic. in Vorles. Churf. Ges. ii. (1787) 385 = M. intertexta.
scutellata, Medic. Phil. Bot. i. 210 = M. scutellata.

COCHLIA, Blume, Bijdr. 320. t. 59 (1825) = **Bulbophyllum**, Thou. (Orchid.).
violacea, Blume, l. c. = B. violaceum.

COCHLIANTHUS, Benth. in Miq. Pl. Jungh. i. 234, in nota (1851?). *LEGUMINOSAE*, Benth. & Hook. f. i. 533.
gracilis, *Benth. l. c.*—Reg. Himal.

COCHLIDIOSPERMUM, Reichb. Consp. 121 (1826) = **Veronica**, Tourn. (Scrophularin.).
agreste, Opiz, Seznam, 31 = V. agrestis.
Buxbaumii, Opiz, l. c. = V. Tournefortii.
digitatum, Opiz, l. c. = V. triphyllos.
Frieseanum, Opiz, l. c. = V. opaca.
hederaefolium, Opiz, l. c. = V. hederaefolia.
Lappago, Opiz, in Lotos, iv. (1854) 158 = V. hederaefolia.
opacum, Opiz, Seznam, 31 = V. opaca.
praecox, Opiz, l. c. = V. praecox.

COCHLIODA, Lindl. Fol. Orchid. (1853). *ORCHIDEAE*, Benth. & Hook. f. iii. 560.
densiflora, *Lindl. l. c.*—Reg. Andin.

COCHLIOPETALUM, Beer, in Flora, xxxvii. (1854) 347; Bromel. 68 (1857) = **Pitcairnia**, L'Hérit. (Bromeliac.).
albiflos, Beer, Bromel. 68 = P. albiflos.
flavescens, Beer, l. c. 69 = P. xanthocalyx.
odoratum, Hemsl. Biol. Centr. Am. Bot. iii. 317 = P. odorata.
Schuechii, Beer, Bromel. 69 = P. albiflos.
stamineum, Beer, in Flora, xxxvii. (1854) 347 = P. staminea.

COCHLIOSPERMUM, Lag. Mem. Pl. Barill. 58 (1817) = **Suaeda**, Forsk. (Chenopod.).
altissimum, Lag. l. c. 57 = S. altissima.
Cavanillesii, Lag. l. c. 60 = S. maritima.
Clementei, Lag. l. c. 62 = S. splendens.
fruticosum, Lag. l. c. 59 = S. fruticosa.
hispanicum, Lag. l. c. 58 = S. altissima.
salsum, Lag. l. c. 55 = S. maritima.

COCHLIOSTEMA, Lem. Illustr. Hortic. vi. (1859) Misc. 70 et t. 217. *COMMELINACEAE*, Benth. & Hook. f. iii. 850.
Jacobianum, *C. Koch & Linden, in C. Koch & Fint. Wochenschr.* x. (1867) 322.—Ecuador.
odoratissimum, *Lem. Illustr. Hortic.* vi. (1859) *Misc.* 70. t. 217.—Ecuador.
odoratum, C. Koch, in C. Koch & Fint. Wochenschr. ii. (1859) 340 = odoratissimum.

COCHLOSPERMUM, Kunth, Malv. 6 (1822). *BIXINEAE*, Benth. & Hook. f. i. 124.
AZEREDIA, Arruda, ex Allem. Desenb. (1846).
MAXIMILIANA, Mart. in Flora, ii. (1819) 451.
WITTELSBACHIA, Mart. Nov. Gen. et Sp. i. 80. t. 55 (1824).
angolense, *Welw. Apont.* 566; *Oliver, Fl. Trop. Afr.* i. 113.—Afr. trop.
Cadinae, *Eichl. in Mart. Fl. Bras.* xiii. t. 431.—Bras.
Fraseri, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 307.—Austral.
Gillivraei, *Benth. Fl. Austral* i. 106.—Austral.
Gossypium, *DC. Prod.* i. 527.—Ind. or.
Gregorii, *F. Muell. Fragm.* i. 71.—Austral.
heteronemum, *F. Muell. in Hook. Kew Journ.* ix. (1857) 15 (= seq.?).—Austral.
heteroneurum, *F. Muell. ex Benth. Fl. Austral.* i. 106.—Austral.
hibiscoides, *Kunth, Syn. Pl. Aeq.* iii. 214.—Mexic.
insigne, *St. Hil. Pl. Us. Bras.* t. 57.—Bras.
niloticum, *Oliver, Fl. Trop. Afr.* i. 113.—Afr. trop.

COCHLOSPERMUM:—
orinoccense, *Steud. Nom.* ed. II. i. 393.—Venezuela.
Parkeri, *Planch. in Hook. Lond. Journ. Bot.* vi. (1847) 310.—Guiana.
parkiaefolium, *Planch. l. c.* 311.—Guiana.
Planchoni, *Hook. f. ex Planch. l. c.* 309.—Afr. trop.
serratifolium, DC. Prod. i. 537 = hibiscoides.
tinctorium, *Perr. in Ann. de la Marine et des Colonies* (1830); *ex A. Rich. in Guill. & Perr. Fl. Seneg. Tent.* i. 99.—Afr. trop.
vitifolium, Spreng. Syst. ii. 596 = serratifolium.
vitifolium, *Spreng. l. c.* iv. *Cur. Post.* 206.—Mexic.

COCHRANEA, Miers, Trav. Chile, ii. 529 (1826) nomen; et in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 125. *BORAGINEAE*, Benth. & Hook. f. ii. 845.
chenopodiacea, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. ii. (1868) 132.—Chili.
conferta, *Miers, Trav. Chile*, ii. 529, nomen; *et l. c.* 125.—Chili.
corymbosa, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. ii. (1868). 126.—Chili.
ericoidea, *Miers, l. c.* 130.—Chili.
filifolia, *Miers, l. c.* 131.—Chili.
florida, *Miers, l. c.* 129.—Chili.
hebecula, *Miers, l. c.* 130.—Chili.
hispidula, *Miers, l. c.* 132.—Chili.
myosotifolia, *Miers, l. c.* 128.—Chili.
sinuata, *Miers, l. c.* 127.—Chili.
stenophylla, *Miers, l. c.* 128.—Chili.

COCKBURNIA, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 90. *SELAGINEAE.*
socotrana, *Balf. f. l. c.*—Ins. Socotra.

COCOCIPSILUM, Jaume St. Hil. Expos. i. 434 (1805) = **Coccocypselum**, P. Br. (Rubiac.).

COCOLOBA, Rafin. Fl. Tellur. ii. 34 (1836) = **Coccoloba**, Linn. (Polygal.).

COCOS, Linn. Sp. Pl. ed. I. 1188 (1753). *PALMAE*, Benth. & Hook. f. iii. 945.
GLAZIOVA, Mart. ex H. Wendl. in Flor. et Pom. (1871) 116.
LANGSDORFFIA, Raddi, in Mem. Soc. Ital. Mod. xviii. (1820) 347. t. 1.
LITHOCARPUS, Targ. Tozz. ex Steud. Nom. ed. II. i. 173, in syn. (1840).
PLATENIA, Karst. in Linnaea, xxviii. (1856) 250.
SYAGRUS, Mart. Hist. Nat. Palm. ii. 129 (1824).
acaulis, *Drude, in Mart. Fl. Bras.* iii. II. 426.—Bras.
acicularis, Sw. Prod. Veg. Ind. Occ. 58 = Bactris minor.
acrocomioides, *Drude, in Mart. Fl. Bras.* iii. II. 409.—Bras.
aculeata, Jacq. Select. Am. 278 = Acrocomia sclerocarpa.
aequatorialis, Drude, in Mart. Fl. Bras. iii. II. 407, sphalm. = C. Inajai.
amara, *Jacq. Select. Am.* 277.—Ind. occ.
arenaria, Gomez, in Mem. Ac. Lisboa, iii. (1812) Mem. dos Corresp. 61 = Diplothemium littorale.
argentea, *Engel, in Linnaea*, xxxiii. (1864–65) 690.—N. Granat.
Aricui, Neuwied, Reise Bras. i. 272 = schizophylla.
australis, *Mart. Palmet. Orbign.* 95.—Paraguay.
Bonneti, *Hort. ex H. Wendl. in Kerch. Palm.* 240, nomen.—Hab.?
botryophora, *Mart. Hist. Nat. Palm.* ii. 118. t. 83, 84.—Bras.
butyracea, *Linn. f. Suppl.* 454.—Am. austr.
butyracea, Mart. ex H. Wendl. in Kerch. Palm. 140 = Scheelea butyracea.
butyrosa, H. Wendl. l. c. = Attalea humilis.
campestris, *Mart. Hist. Nat. Palm.* ii. 121.—Bras.
capitata, *Mart. l. c.* 114.—Bras.
chilensis, Molina, ex H. B. & K. Nov. Gen. et Sp. i. 309, in obs. = Jubaea spectabilis.
comosa, *Mart. Hist. Nat. Palm.* ii. 121. t. 88. f. 1–2.—Bras.
coronata, *Mart. l. c.* 115. tt. 80, 81.—Bras.
crispa, *H. B. & K. Nov. Gen. et Sp.* i. 302.—Cuba.
Datil, *Drude & Griseb. in Goett. Abh.* xxiv. (1879) 283.—Reg. Argent.
elegantissima, Hort. ex H. Wendl. in Kerch. Palm. 241 = Weddelliana.

COCOS:—
equatorialis, Barb.-Rodr. Enum. Palm. Nov. 38 = C. Inajai.
eriospatha, *Mart. ex Drude, in Mart. Fl. Bras.* iii. II. 424.—Bras.
flexuosa, *Mart. Hist. Nat. Palm.* ii. 120.—Bras.
frigida, *Linden, in Illustr. Hortic.* xxviii. (1881) 16, nomen.—Bras.
fusiformis, Sw. Fl. Ind. Occ. i. 616 = Acrocomia sclerocarpa.
Geriba, Barb.-Rodr. Prot. App. 43 = botryophora.
graminifolia, *Drude, in Mart. Fl. Bras.* iii. II. 415.—Bras.
Guacuyule, *Liebm. ex Mart. Hist. Nat. Palm.* iii. 323.—Mexic.
guineensis, Linn. Mant. i. 137 = Bactris minor.
Inajai, *Spruce, ex Trail, in Journ. Bot.* xv. (1877) 79.—Bras.
insignis, *Mart. ex H. Wendl. in Kerch. Palm.* 241.—Bras.
Kotchoubeyi, *Linden, Illustr. Hortic.* xxviii. (1881) 16, nomen.—Bras.
lapidea, *Gaertn. Fruct.* i. 16. t. 6. f. 1.—Bras.?
leiospatha, *Barb.-Rodr. Prot. App.* 44.—Bras.
maldioica, J. F. Gmel. Syst. 569 = Lodoicea sechellarum.
mamillaris, *Blanco, Fl. Filip.* ed. I. 722.—Ins. Philipp.
maritima, Comm. ex H. Wendl. in Kerch. Palm. 241 = Lodoiceae sp.
Martiana, *Drude & Glaz. ex Drude, in Mart. Fl. Bras.* iii. II. 418.—Bras.
melanococca, Hort. Belg. ex Heynh. Nom. ii. 152, nomen.—Hab.?
Mikaniana, *Mart. Hist. Nat. Palm.* iii. 128.—Bras.
Naja, *Arruda, ex Kunth, Enum. Pl.* iii. 288.—Am. trop.
nana, Griff. Notul. iii. 166 = nucifera.
Nolaia-assu, *Neuwied, Reise Bras.* i. 271.—Bras.
Normanbyi, W. Hill, Rep. Brisb. Bot. (1876) 6; ex F. Muell. Fragm. viii. 235 = Areca Normanbyi.
nucifera, *Linn. Sp. Pl.* 1188.—Reg. trop.
Nypa, Lour. Fl. Cochinch. 567 = Nipa fruticans.
oleracea, *Mart. Hist. Nat. Palm.* ii. 117. t. 82.—Bras.
orinocensis, *Spruce, in Journ. Linn. Soc.* xi. (1871) 161.—Am. trop.
ovata, *Hort. Belg. ex Heynh. Nom.* ii. 152, nomen.—Hab.?
petraea, *Mart. Palmet. Orbign.* 100.—Bolivia.
pityrophylla, *Mart. l. c.* 99.—Bolivia.
plumosa, *Hook. Bot. Mag.* t. 5180.—Bras.
plumosa, Lodd. ex Loud. Hort. Brit. 381 = comosa.
Procopiana, *Glaz. ex Drude, in Mart. Fl. Bras.* iii. II. 412.—Bras.
regia, *Liebm. in Mart. Hist. Nat. Palm.* iii. 323.—Mexic.
Romanzoffiana, *Cham. in Choris, Voy. Pittor. (Chili)* 5. t. 6 & 6 [bis].—Bras.
rupestris, Barb.-Rodr. Prot. App. 45 = petraea.
schizophylla, *Mart. Hist. Nat. Palm.* ii. 119. tt. 84, 85.—Bras.
speciosa, *Barb.-Rodr. Enum. Palm. Nov.* 38.—Bras.
simplicifolia, *Hort. Belg. ex Heynh. Nom.* ii. 152.—Hab.?
Syagrus, *Drude, in Mart. Fl. Bras.* iii. II. 406.—Bras.
sylvestris, *Hort. ex H. Wendl. in Kerch. Palm.* 241, nomen.—Hab.?
Urucuru, *Linden, in Illustr. Hortic.* xxviii. (1881) 16, nomen.—Bras.
ventricosa, *Arruda, in Koster, Trav. Braz.* 485.—Bras.
venatorum, Poepp. ex H. Wendl. in Kerch. Palm. 241 = Maximiliana venatorum.
vinifera, Mart. Hist. Nat. Palm. iii. 324 = Gaussia vinifera.
Weddelliana, *H. Wendl. in Flor. et Pom.* (1871) 114.—Bras.
Weddellii, *Drude, in Mart. Fl. Bras.* iii. II. 411.—Bras.
Yatay, *Mart. Palmet. Orb.* 93.—Reg. Argent.
Yurumaguas, *Hort. ex H. Wendl. in Kerch. Palm.* 241.—Peruv.

CODANTHERA, Rafin. Fl. Tellur. iii. 90 (1836) = **Salvia**, Tourn. (Labiat.).
biflora, Rafin. l. c. = S. hebiflora.
glabra, Rafin. l. c. = S. strictiflora.

CODARIA, Linn. ex Benn. Pl. Jav. Rar. 99 (1838) = **Lerchea**, Kalm. (Rubiac.).

CODARIOCALYX, Hassk. in Flora, xxv. (1842) II. Beibl. 48 = **Desmodium**, Desv. (Legumin.).
capitatus, Hassk. l. c. 49 = D. capitatum.
conicus, Hassk. l. c. 49, 62 = D. gyroides.
gyrans, Hassk. l. c. = D. gyrans.
gyroides, Hassk. l. c. 49 = D. gyroides.

CODARIUM, Soland. ex Vahl, Enum. i. 302 (1806) = **Dialium**, Linn. (Legumin.).
acutifolium, Afzel. in Schrad. N. Journ. ii. (1807) 233 = D. guineense.
discolor, DC. Prod. ii. 520 = D. guineense.
nitidum, Vahl, Enum. i. 302; ii. 400 = D. guineense.
obtusifolium, Afzel. in Schrad. N. Journ. ii. (1807) 233 = D. guineense.
Solandri, Vahl, Enum. i. 302 = D. guineense.

CODAZZIA, Karst. & Triana, in Linnaea, xxviii. (1856) 426 = **Delostoma**, D. Don (Bignoniac.).
dentata, Karst. Fl. Columb. ii. 37. t. 119 = D. rosea.
rosea, Karst. in Linnaea, xxviii. (1856) 427 = D. rosea.
speciosa, Karst. l. c.—N. Granat.

CODDAMPULI, Adans. Fam. ii. 445 (1763) = **Garcinia**, Linn. (Guttif.).

CODDA-PANA, Adans. Fam. ii. 25 (1763) = **Bentinckia**, Berry (Palm.).

CODIA, Forst. Char. Gen. 59. t. 30 (1776). *SAXIFRAGEAE*, Benth. & Hook. f. i. 648.
ferruginea, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 77.—N. Caled.
floribunda, *Brongn. & Gris, l. c.*—N. Caled.
montana, *Forst. Char. Gen.* 68.—N. Caled.
montana, Labill. ex D. Don, in Edinb. N. Phil. Journ. (Apr.–June 1830) 94 = Callicoma Billardieri.
obcordata, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 77.—N. Caled.
serratifolia, Ser. ex DC. Prod. iv. 7 = Callicoma serratifolia.
spathulata, *Brongn. & Gris, in Bull. Soc. Bot. Fr.* ix. (1862) 77.—N. Caled.

CODIAEUM, Rumph. ex A. Juss. Euph. Tent. 33. t. 9 (1824). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 299.
CROZOPHYLA, Rafin. Sylva Tellur. 64 (1838).
JUNGHUHNIA, Miq. Fl. Ind. Bat. i. II. 412 (1855).
PHYLLAUREA, Lour. Fl. Cochinch. 575 (1790).
SYNASPISMA, Endl. Gen. 1110 (1840).
alternifolium, *Muell. Arg. in DC. Prod.* xv. II. 1117.—N. Caled.
andamanicum, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 246 (= Blachiae sp.).—Ins. Andaman.
aurantiacum, *Muell. Arg. in DC. Prod.* xv. II. 1118.—Ins. Banca.
Balansae, *Baill. Adansonia*, xi. (1873–76) 77.—N. Caled.
Brongniartii, *Baill. l. c.* 76.—N. Caled.
Bureavi, *Baill. l. c.* 74.—N. Caled.
carunculatum, *Muell. Arg. in DC. Prod.* xv. II. 1117.—N. Caled.
chrysosticton, Spreng. Syst. iii. 866 = variegatum.
cuneifolium, Zipp. ex Span. in Linnaea, xv. (1841) 348 = variegatum.
Deplanchei, *Baill. Adansonia*, xi. (1873–76) 75.—N. Caled.
drimiflorum, *Baill. l. c.*—N. Caled.
inophyllum, *Muell. Arg. in DC. Prod.* xv. II. 1120.—N. Caled.
lutescens, Kurz, in Journ. As. Soc. Beng. xlii. (1873) II. 246 = Sphyranthera capitellata.
medium, Baill. Adansonia, i. (1860–61) 251 = variegatum.
moluccanum, Decne. in Nouv. Ann. Mus. Par. iii. (1834) 485 = variegatum,
obovatum, Zoll. & Morr. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 582 = variegatum.
Pancheri, Muell. Arg. in DC. Prod. xv. II. 1117 = Baloghia Pancheri.
Pentzii, Muell. Arg. l. c. 1118 = Blachia Pentzii.
pictum, Hook. Bot. Mag. t. 3051 = variegatum.
roseo-pictum, *André, Illustr. Hortic.* xxvi. (1879) 170. t. 364.—Hab.?
timorense, A. Juss. Tent. Euph. 34, nomen = variegatum.
umbellatum, Muell. Arg. in DC. Prod. xv. II. 1118 = Blachia umbellata.
variegatum, *Blume, Bijdr.* 606.—Malaya.

CODIAMINUM, Rafin. Fl. Tellur. iv. 21 (1836) = **Narcissus**, Tourn. (Amaryllid.) (*N.* Bulbocodium).

CODIPHUS, Rafin. Fl. Tellur. ii. 77 (1836) = **Prismatocarpus**, L'Hérit. (Campanulac.).
nitidus, Rafin. l. c. = P. nitidus.

CODIVALIA, Rafin. Fl. Tellur. iii. 40 (1836) = **Pupalia**, Juss. (Amarant.).
lappacea, Rafin. l. c. = P. lappacea.
patula, Rafin. l. c. = P. lappacea.

CODOCHONIA, Dun. in DC. Prod. xiii. 1. 482 (1852) = **Acnistus**, Schott (Solanac.).
solanoides, Dun. l. c. = A. umbellatus.

CODON, Linn. Syst. ed. XII. 292 (1767). *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 830.
aculeatum, Gaertn. Fruct. ii. 88 = Royeni.
Dregei, *E. Mey. Zwei Pfl. Docum.* 174, nomen.—Afr. austr.
Royeni, *Linn. Syst.* ed. XII. 292.—Afr. austr.

CODONACANTHUS, Nees, in DC. Prod. xi. 103 (1847). *ACANTHACEAE*, Benth. & Hook. f. ii. 1098.
acuminatus, Nees, l. c. = pauciflorus.
pauciflorus, *Nees, l. c.*—Hongkong.

CODONACHNE, Wight & Arn. ex Steud. Nom. ed. II. i. 393 (1840) = **Chloris**, Sw. (Gramin.).
Neesiana, Wight & Arn. l. c. = Chloris tenella.

CODONANDRA, Karst. Fl. Columb. ii. 43. t. 122 (1862 ?) = **Calliandra**, Benth. (Legumin.).
coroënsis, C. Muell. in Walp. Ann. vii. 841, sphalm. = Call. purpurea.
purpurea, Karst. Fl. Columb. ii. 43. t. 122 = Call. Codonandra.

CODONANTHE, Hanst. in Linnaea, xxvi. (1853) 209. *GESNERACEAE*, Benth. & Hook. f. ii. 1011.
Coccanthera, C. Koch & Hanst. Ind. Sem. Hort. Berol. (1855) 17.
aggregata, *Hanst. in Linnaea*, xxvi. (1853-55) 209.—Bras.
calcarata, *Hanst. l. c.* xxxiv. (1865-66) 416.—Am. trop.
carnosa, *Hanst. in Mart. Fl. Bras.* viii. 418.—Bras.
ciliosa, *Lem. in Illustr. Hortic.* ii. (1855) sub t. 56.—Bras.
Devosiana, *Lem. l. c.* t. 56.—Bras.
gracilis, *Hanst. in Linnaea*, xxvi. (1853-55) 209.—Bras.
Hookerii, *Lem. in Illustr. Hortic.* ii. (1855) sub t. 56.—Bras.
picta, Lem. l. c. iv. (1857) t. 144 = Devosiana.

CODONANTHE, Mart. ex Stend. Nom. ed. II. i. 791, in syn. (1840) = **Hypocyrta**, Mart. (Gesnerac.).

CODONANTHEMUM, Klotzsch, in Linnaea, xii. (1838) 240 = **Simocheilus**, Klotzsch (Ericac.).
discolor, Benth. in DC, Prod. vii. 708.—Afr. austr.
parviflorum, Klotzsch, in Linnaea, xii. (1838) 240.—Afr. austr.
puberulum, Klotzsch, l. c.—Afr. austr.
tenue, Benth. in DC. Prod. vii. 708.—Afr. austr.

CODONANTHES, Rafin. Fl. Tellur. iv. 24 (1836) = **Pitcairnia**, L'Hérit. (Bromeliac.).
albiflos, Rafin. l. c. = P. albiflos.

CODONANTHUS, G. Don, Gen. Syst. iv. 166 (1836) = **Breweria**, R. Br. (Convolvulac.).
africana, G. Don, l. c. = B. africana.
alternifolia, Planch. in Hook. Ic. Pl. t. 796.—Afr. trop.

CODONANTHUS, Hassk. in Flora, xxv. (1842) II. Beibl. 24 = **Physostelma**, Wight (Asclepiad.).
campanulatus, Hassk. l. c. = P. Wallichii.

CODONEMMA, Miers, Apocyn. S. Am. 72 (1878) = **Tabernaemontana**, Plum. (Apocynac.).
calycinum, Miers, l. c. 73 = T. Muelleriana.
macrocalyx, Miers, l. c. = T. macrocalyx.

CODONIUM, Vahl, in Skriv. Naturh. Selsk. ii. (1793) 1. 206; Symb. Bot. iii. 36 (1794) = **Schoepfia**, Schreb. (Olacin.).
arborescens, Vahl, ll. cc. 207; 36 = S. arborescens.
flexuosum, M. Roem. Syn. Hesper. 21 = S. flexuosa.

CODONOCALYX, Klotzsch, ex Baill. Etud. Gén. Euph. 369 (1858) = **Croton**, Linn. (Euphorb.).
divaricatus, Klotzsch, l. c. 370 = Crot. nitrariaefolius.
lanatus, Klotzsch, l. c. = Crot. lanuginosus.
longifolius, Klotzsch, l. c. = Crot. montevidensis.
montevidensis, Klotzsch, l. c. = Crot. montevidensis.
polymorphus, Klotzsch, l. c. = Crot. nitrariaefolius.
velleriflorus, Klotzsch, l. c. = Crot. nitrariaefolius.

CODONOCALYX, Miers, in Lindl. Veg. Kingd. 764 (1847) = **Suteria**, DC. (Rubiac.).

CODONOCARPUS, A. Cunn. ex Hook. Bot. Misc. i. (1830) 244, in syn. *PHYTOLACCACEAE*, Benth. & Hook. f. iii. 86.
Hymenotheca, F. Muell. Fragm. i. 201 (1859).
australis, *A. Cunn. l. c.*—Austral.
cotinifolia, *F. Muell. Pl. Vict.* i. 200.—Austral.
pyramidalis, *F. Muell. l. c.* 201.—Austral.

CODONOCEPHALUM, Fenzl, in Flora, xxvi. (1843) 397. *COMPOSITAE*, Benth. & Hook. f. ii. 330.
Sprunnera, Sch. Bip. in Walp. Rep. ii. 954 (1843).
inuloides, *Fenzl, l. c.*—Kurdistan.

CODONOCRINUM, Willd. ex Schult. f. Syst. vii. 718 (1829) = **Yucca**, Dill. (Liliac.).
agavoides, Willd. ex Schult. f. l. c. = Y. acaulis.

CODONOPHORA, Lindl. Bot. Reg. sub t. 1110 (1827) = **Paliavana**, Vand. (Gesnerac.).
grandiflora, Lindl. l. c. = P. prasinata.
lanceolata, Lindl. l. c. = Rytidophyllum tomentosum.
prasinata, Lindl. l. c. = P. scriciflora.
Selloana, Hanst. in Mart. Fl. Bras. viii. 396 = P. prasinata.

CODONOPRASUM, Reichb. Consp. 66 (1828) = **Allium**, Tourn. (Liliac.).
alpicolum, Jord. & Fourr. Brev. Pl. Nov. fasc. ii. 127 = A. oleraceum.
carinatum, Reichb. Fl. Germ. Excurs. 114 = A. carinatum.
complanatum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159 = A. oleraceum.
consimile, Fourr. l. c. = A. carinatum.
flavum, Reichb. Fl. Germ. Excurs. 115 = A. flavum.
flexifolium, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159 = A. carinatum.
flexum, Reichb. Fl. Germ. Excurs. 114 = A. flavum.
fuscum, Reichb. l. c. 115 = A. paniculatum.
intermedium, Reichb. l. c. 114 = A. oleraceum.
longispathum, Reichb. l. c. 115 = A. paniculatum.
oleraceum, Reichb. l. c. 114 = A. oleraceum.
pallens, Reichb. l. c. 115 = A. flavum.
paniculatum, Reichb. l. c. = A. pulchellum.
pulchellum, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 159 = A. pulchellum.
viridiflorum, Schur, Enum. Pl. Transs. 671 = A. oleraceum.

CODONOPSIS, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 103 (1824). *CAMPANULACEAE*, Benth. & Hook. f. ii. 557.
Glosocomia, D. Don, Prod. Fl. Nep. 158 (1824).
affinis, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 12.—Reg. Himal.
albiflora, Griff. Notul. iv. 279 = Campanumoea celebica.
Benthamii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 14.—Reg. Himal.
celebica, Miq. Fl. Ind. Bat. ii. 566 = Campanumoea celebica.
clematidea, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iii. 433 = ovata.
convolvulacea, *Kurz, in Journ. Bot.* xi. (1873) 195.—Burma.
cordata, Hassk. in Retzia, i. 9 = Campanumoea javanica.
foetens, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 16.—Reg. Himal.
gracilis, Hook. f. & Thoms. in Illustr. Himal. Pl. t. 16 A = Leptocodon gracilis.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iii. 431.—Reg. Himal.
inflata, Hook. f. & Thoms. in Journ. Linn. Soc. ii. (1858) 13 = Campanumoea inflata.
japonica, *Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 192.—Japon.

CODONOPSIS:—
javanica, *Hook. f. & Thoms. in Illustr. Himal. Pl.* t. 16 B = Campanumoea javanica.
lanceolata, *Benth. & Hook. f. Gen.* ii. 558; *Trautv. in Act. Hort. Petrop.* vi. (1879) 46.—As. temp.
leucocarpa, Miq. Fl. Ind. Bat. ii. 565 = Campanumoea celebica.
lurida, Lindl. Bot. Reg. (1839) Misc. 82 = rotundifolia.
ovata, *Benth. in Royle, Illustr. Bot. Himal.* 253. t. 69. f. 3.—Reg. Himal.
parviflora, Wall. Cat. n. 1300 = Campanumoea parviflora.
purpurea, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 105.—Reg. Himal.
rotundifolia, *Royle, Illustr. Bot. Himal.* 254. t. 62.—Reg. Himal.
subsimplex, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 16.—Reg. Himal.
thalictrifolia, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 106.—Reg. Himal.
truncata, Wall. Cat. n. 1301 = Campanumoea celebica.
viridiflora, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 496.—As. or.
viridis, *Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 103.—Reg. Himal.

CODONORAPHIA, Oerst. in Vidensk. Selsk. Skr. V. v. (1859) 82; (Gesn. Centr. Am. 8) = **Pentarhaphia**, Lindl. (Gesnerac.).

CODONORCHIS, Lindl. Gen. et Sp. Orch. 410 (1840) = **Pogonia**, Juss. (Orchid.).
Lessonii, Lindl. l. c. 411 = P. tetraphylla.
Poeppigii, Lindl. l. c. 410 = P. tetraphylla.

CODONOSTIGMA, Klotzsch, ex Benth. in DC. Prod. vii. 709 (1839) = **Scyphogyne**, Brongn. (Ericac.).
Erinus, Klotzsch, ex Benth. l. c.—Afr. austr.

CODORIOCALYX, Hassk.—Cf. Linnaea, xv. (1841) Lit. 80, 81 = CODARIOCALYX, Hassk. = **Desmodium**, Desv. (Legumin.).

CODYLIS, Rafin. in Am. Monthly Mag. (1819) 191 = **Nicotiana**, Linn. (Solanac.).

COELACHNE, R. Br. Prod. 187 (1810). *GRAMINEAE*, Benth. & Hook. f. iii. 1156.
brachiata, *Munro, ex Benth. Fl. Austral.* vii. 626, in obs.—Zeylan.; China.
infirma, *Buese, in Miq. Pl. Jungh.* 350.—Java.
perpusilla, *Thw. Enum. Pl. Zeyl.* 373.—Zeylan.
pulchella, *R. Br. Prod.* 187.—Burma; Austral.
simpliciuscula, Munro, ex Benth. in Journ. Linn. Soc. xix. (1881) 93 = brachiata.
subulifolia, F. Muell. Fragm. v. 208 = Micraira subulifolia.

COELACHYRUM, Nees, in Linnaea, xvi. (1842) 221 = **Eragrostis**, Beauv. (Gramin.).
brevifolium, Nees, l. c. = E. Coelachyrum.

COELANDRIA, Fitzg. Austral. Orch. i. pt. VII. t. 2 (1882) = **Dendrobium**, Sw. (Orchid.).
Smilliae, Fitzg. l. c. = D. Smilliae.

COELANTHE, Griseb. Gen. et Sp. Gent. 297. in syn. (1839) = COILANTHA, Borckh. = **Gentiana**, Linn.

COELANTHIUM, Sond. in Harv. & Sond. Fl. Cap. i. 147 (1859) = seq.

COELANTHUM, E. Mey. ex Fenzl, in Ann. Wien. Mus. i. (1836) 353; ii. (1838) 267. *FICOIDEAE*, Benth. & Hook. f. i. 857.
grandiflorum, *E. Mey. ex Fenzl, l. c.*—Afr. austr.
parviflorum, *Fenzl, l. c.* 268.—Afr. austr.

COELANTHUS, Willd. ex Schult. f. Syst. vii. p. xlvi (1830) = **Lachenalia**, Jacq. (Liliac.).
complicatus, Willd. l. c. in adnot. = L. reflexa.

COELAS, Dulac, Fl. Hautes-Pyr. 303 (1867) = **Potentilla**, Tourn. (Rosac.).
procumbens, Dulac, l. c. = P. Sibbaldi.

COELEBOGYNE, J. Sm. in Proc. Linn. Soc. i. (1839) 41; Trans. Linn. Soc. xviii. (1841) 511. t. 36 = **Alchornea**, Sw. (Euphorb.).
ilicifolia, J. Sm. l. c. 512. t. 36 = A. ilicifolia.

COELESTINA, Cass. in Bull. Soc. Philom. (Jan. 1817) 10 = CAELESTINA, Cass. = **Ageratum**, Linn. (Compos.).

COELESTINA, Hill, Hort. Kew. 8 (1769); Veg. Syst. ii. 60** = **Aster**, Tourn. (Compos.).

COELESTINIA, Endl. Gen. 366 (1838) = CAELESTINA, Cass. = **Ageratum**, Linn. (Compos.).

COELIA, Lindl. Gen. et Sp. Orch. 36 (1830). *ORCHIDEAE*, Benth. & Hook. f. iii. 508.
BOTHRIOCHILUS, Lem. Illustr. Hortic. iii. (1856) Misc. 30.
Baueriana, *Lindl. Gen. et Sp. Orch.* 36.—Mexic.
bella, *Reichb. f. in Walp. Ann.* vi. 218.—Guatemala.
Galeottiana, *Van Houtte, ex Heynh. Nom.* ii. 152, nomen.—Mexic.
glacialis, *Van Houtte, l. c.*, nomen.—Am. austr.
guatemalensis, *Reichb. f. in Walp. Ann.* vi. 219.—Guatemala.
macrostachya, *Lindl. in Benth. Pl. Hartw.* 92.—Mexic.
triptera, G. Don, ex Steud. Nom. ed. II. i. 394 = Baueriana.

COELIDIUM, Vog. ex Walp. in Linnaea, xiii. (1839) 472. *LEGUMINOSAE*, Benth. & Hook. f. i. 473.
Bowei, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 454.—Afr. austr.
bullatum, *Benth. l. c.* 453.—Afr. austr.
ciliare, *Vog. ex Walp. in Linnaea*, xiii. (1839) 472.—Afr. austr.
muraltioides, *Benth. in Hook. Lond. Journ. Bot.* ii. (1843) 455.—Afr. austr.
roseum, *Benth. l. c.* 454.—Afr. austr.
Thunbergii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 24.—Afr. austr.
Vogelii, *Walp. in Linnaea*, xiii. (1839) 472.—Afr. austr.

COELINA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 (Quid ?).

COELIOPSIS, Reichb. f. in Gard. Chron. (1872) 9. *ORCHIDEAE*, Benth. & Hook. f. iii. 477.
hyacinthosma, *Reichb. f. l. c.*—Panama.

COELOCARPUM, Balf. f. in Proc. Roy. Soc. Edinb. xii. (1884) 90. *VERBENACEAE.*
socotranum, *Balf. f. l. c.* 91.—Ins. Socotra.

COELOCHLOA, Hochst. ex Steud. Nom. ed. II. i. 394 (1840) = COELACHYRUM, Nees = **Eragrostis**, Beauv. (Gramin.).
brevifolia, Hochst. l. c. = E. Coelachyrum.

COELOCLINE, A. DC. in Mém. Soc. Phys. Genèv. v. (1832) 208 = **Xylopia**, Linn. (Anonac.).
acutiflora, A. DC. l. c. = X. acutiflora.
lucida, A. DC. l. c. 209 = X. longifolia.
oxypetala, A. DC. l. c. = X. acutiflora ?
parviflora, A. DC. l. c. = X. parviflora.
polycarpa, A. DC. l. c. = X. polycarpa.

COELOCOCCUS, H. Wendl. in Bonplandia, x. (1862) 199 = **Metroxylon**, Rottb. (Palm.).
vitiensis, H. Wendl. l. c. = M. vitiense.

COELODEPAS, Hassk. in Flora, xl. (1857) 531. *EUPHORBIACEAE*, Benth. & Hook. f. iii. 313.
KOILODEPAS, Hassk. in Versl. en Med. Kon. Akad. Wetensch. iv. (1855) 139.
bantamense, *Hassk. in Versl. en Med. Kon. Acad. Wetensch.* iv. (1855) 140 (Koilodepas).—Ins. Bantam.
calycinum, *Bedd. Forest. Man. Bot.* 207.—Ind. or.
Wallichianum, *Benth. in Hook. Ic. Pl.* t. 1288.—Ins. Penang.

COELODISCUS, Baill. Etud. Gén. Euph. 293 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 317.
dispar, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 244.—Ind. or.
eriocarpoides, Kurz, l. c. = Mallotus decipiens.
eriocarpus, *Kurz, l. c.*—Ind. or.
glabriusculus, *Kurz, For. Fl. Brit. Burma*, ii. 393.—Burma.

COELODISCUS:—
hirsutulus, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) II. 243.—Burma.
lappaceus, *Kurz, l. c.* 244; *et For. Fl. Brit. Burma*, ii. 392.—Burma.
longipes, *Kurz, ll. cc.* 244, 393.—Burma.
montanus, *Muell. Arg. in DC. Prod.* xv. II. 759.—Ins. Penang.
speciosus, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 154.—Ind. or.
Thunbergianus, *Muell. Arg. in DC. Prod.* xv. II. 758.—Zeylan.

COELOGLOSSUM, Hartm. Handb. Skand. Fl. ed. I. 329 (1820) = **Habenaria**, Willd. (Orchid.).
acuminatum, Lindl. Gen. et Sp. Orch. 302 = H. lacertifera.
albidum, Hartm. Handb. Skand. Fl. ed. I. 329 = H. albida.
alpinum, Schur, in Verh. Siebenb. Ver. Naturw. ii. (1851) 169; iv. (1853) 72 = H. cucullata.
bracteatum, Parl. Fl. Ital. iii. 409 = H. viridis.
brevifolium, Lindl. Gen. et Sp. Orch. 302 = H. cubitalis.
cernuum, Reichb. f. in Bonplandia, iii. (1855) 250 = H. stenostachya.
cordatum, Nym. Syll. 359 = H. cordata.
densiflorum, Hartm. ex Willk. & Lange, Prod. Fl. Hisp. i. 164 = H. intacta.
densum, Lindl. Gen. et Sp. Orch. 302 = H. stenostachya.
Erdingeri, Kerner, in Oestr. Bot. Zeitschr. xiv. (1864) 140 = H. viridis.
lacertiferum, Lindl. Gen. et Sp. Orch. 302 = H. lacertifera.
luteum, Dalz. in Hook. Kew Journ. ii. (1850) 263.—Ind. or.
Mannii, Reichb. f. in Linnaea, xli. (1877) 54.—Reg. Himal.
peristyloides, Reichb. f. in Bonplandia, iv. (1856) 321 = H. stenostachya.
purpureum, Schur, Enum. Pl. Transs. 646 = H. cucullata.
satyrioides, Nym. Syll. 359.—Tauria.
secundum, Lindl. Bot. Reg. sub t. 1701.—Ind. or.
tenue, Lindl. l. c. = H. stenostachya.
Vaillantii, Schur, Enum. Pl. Transs. 645 = H. viridis.
viride, Hartm. Handb. Skand. Fl. ed. I. 329 = H. viridis.

COELOGYNE, Lindl. Collect. sub t. 33 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 518.
ACANTHOGLOSSUM, Blume, Bijdr. 381 (1825).
BOLBORCHIS, Zoll. & Mor. Verz. Pfl. 89 (1846).
GOMPHOSTYLIS, Wall. ex Lindl. Gen. et Sp. Orch. 43 (1830).
NEOGYNA, Reichb. f. in Bot. Zeit. x. (1852) 931.
PLEIONE, D. Don, Prod. Fl. Nep. 36 (1825).
advena, *Par. & Reichb. f. Otia Bot. Hamb.* 47.—Burma.
alba, Reichb. f. in Walp. Ann. vi. 236 = Otochilus alba.
angustifolia, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 16. t. 6 = odoratissima.
angustifolia, Wight, Ic. t. 1641 = breviscapa.
apiculata, Reichb. f. in Walp. Ann. vi. 225 = Panisea apiculata.
Arthuriana, Reichb. f. in Gard. Chron. (1881) I. 40 = maculata.
articulata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota articulata.
asperata, *Lindl. in Journ. Hort. Soc.* iv. (1849) 221.—Malaya.
assamica, Linden & Reichb. f. in C. Koch, Allg. Gartenz. xxv. (1857) 403 = fuscescens.
barbata, *Lindl. ex Griff. Itin. Not.* 72.—Reg. Himal.
biflora, *Parish, ex Reichb. f. in Gard. Chron.* (1865) 1035.—Burma.
bilamellata, *Lindl. Fol. Orchid. Coelog.* 14.—Ins. Philipp.
birmanica, *Reichb. f. in Gard. Chron.* (1882) II. 840.—Burma.
brachyptera, *Reichb. f. l. c.* (1881) II. 6.—Burma.
brevifolia, *Lindl. Fol. Orchid. Coelog.* 7.—Reg. Himal.
breviscapa, *Lindl. l. c.* 4.—Ind. or.
brunnea, Lindl. in Gard. Chron. (1848) 71 = fuscescens.
calceata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota calceata.

COELOGYNE:—
camelostalix, Reichb. f. in Walp. Ann. vi. 238 = Pholidota camelostalix.
carnea, Reichb. f. l. c. 237 = Pholidota carnea.
caulescens, *Griff. Notul.* iii. 282.—Malacca.
chinensis, Reichb. f. in Walp. Ann. vi. 237 = Pholidota chinensis.
chloroptera, *Reichb. f. in Gard. Chron.* (1883) I. 466.—Ins. Philipp.
ciliata, *Boxall, ex Naves, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 237.—Ins. Philipp.
clypeata, Reichb. f. in Walp. Ann. vi. 237 = Pholidota clypeata.
conchoidea, Reichb. f. l. c. = Pholidota conchoidea.
conferta, *Hort.—Cf. Gard. Chron.* (1875) I. 314.—Ind. or.
Convallariae, *Par. & Reichb. f. in Flora*, lv. (1872) 277.—Reg. Himal.; Burma.
corniculata, *Reichb. f. in Gard. Chron.* (1865) 746.—Assam ?
coronaria, Lindl. Bot. Reg. (1841) Misc. 83 = Trichosma suavis.
corrugata, *Wight, Ic.* t. 1639.—Ind. or.
corymbosa, *Lindl. Fol. Orchid. Coelog.* 7.—Reg. Himal.
cristata, *Lindl. Collect. Bot.* sub t. 33.—Reg. Himal.
Croockewitii, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* v. (1853) 488.—Java.
crotalina, Reichb. f. in Walp. Ann. vi. 238 = Pholidota crotalina.
Cumingii, *Lindl. Bot. Reg.* (1843) *Misc.* 76.—Malaya.
Cycnoches, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 147.—Burma.
cymbidioides, *Reichb. f. in Walp. Ann.* vi. 239.—Ins. Molucc.
Dayana, *Reichb. f. in Gard. Chron.* (1884) I. 826.—Borneo.
decora, *Wall. ex Voigt, Hort. Suburb. Calc.* 621, nomen.—Hab. ?
diphylla, *Lindl. Fol. Orchid. Coelog.* 15.—Ind. or.
elata, *Lindl. in Wall. Cat.* n. 1959.—Reg. Himal.
fimbriata, *Lindl. Bot. Reg.* t. 868.—Reg. Himal.
flaccida, *Lindl. in Wall. Cat.* n. 1961.—Reg. Himal.
flavida, *Hook. f. ex Lindl. Fol. Orchid. Coelog.* 10.—Reg. Himal.
fuliginosa, *Lodd. ex Hook. Bot. Mag.* t. 4440.—Reg. Himal.
fusca, Reichb. f. in Walp. Ann. vi. 236 = Otochilus fusca.
fuscescens, *Lindl. in Wall. Cat.* n. 1962.—Malaya.
fusco-lutea, *Teijsm. & Binn. Cat. Hort. Bog.* 378, nomen.—Hab. ?
Gardneriana, *Lindl. in Wall. Pl. As. Rar.* i. 33. t. 38.—Reg. Himal.
gibbosa, Reichb. f. in Walp. Ann. vi. 237 = Pholidota gibbosa.
glandulosa, *Lindl. Fol. Orchid. Coelog.* 6.—Ind. or.
globosa, Reichb. f. in Walp. Ann. vi. 236 = Pholidota globosa.
Goweri, *Reichb. f. in Gard. Chron.* (1869) 443.—Reg. Himal.
graminifolia, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 146.—Burma.
Hookeriana, *Lindl. Fol. Orchid. Coelog.* 14.—Reg. Himal.
Huettneriana, *Reichb. f. in Flora*, lv. (1872) 277.—Burma.
humilis, *Lindl. Collect. Bot.* sub t. 37.—Reg. Himal.
imbricata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota imbricata.
incrassata, *Lindl. Gen. et Sp. Orch.* 40.—Java.
interrupta, *Lindl. ex Heynh. Nom.* ii. 153, nomen.—Ind. or.
javanica, *Lindl. Fol. Orchid. Coelog.* 17.—Java.
khasyana, Reichb. f. in Walp. Ann. vi. 238 = Pholidota articulata.
lactea, *Reichb. f. in Gard. Chron.* (1885) I. 692.—Burma.
Lagenaria, *Lindl. Fol. Orchid. Coelog.* 15.—Reg. Himal.
lentiginosa, *Lindl. l. c.* 3.—Burma.
ligulata, *Teijsm. & Binn. Cat. Hort. Bog.* 46.—Java.
longeciliata, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvii. (1864) 16.—Ind. or.
longifolia, *Lindl. Gen. et Sp. Orch.* 42.—Java.
longipes, *Lindl. Fol. Orchid. Coelog.* 10.—Reg. Himal.

COELOGYNE:—
loricata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota loricata.
Lourii, Paxt. Mag. Bot. xvi. (1849) 225 = asperata.
macrophylla, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxix. (1867) 241.—Sumatra.
maculata, *Lindl. Gen. et Sp. Orch.* 43.—Reg. Himal.
marmorata, *Reichb. f. in Linnaea*, xli. (1877) 116.—Ins. Philipp.
Massangeana, *Reichb. f. in Gard. Chron.* (1878) II. 684.—Malay.
Maycriana, *Reichb. f. l. c.* (1877) II. 134.—Hab.?
media, *Wall. ex Voigt, Hort. Suburb. Calc.* 621, nomen.—Ind. or.
micrantha, *Lindl. in Gard. Chron.* (1855) 173.—Reg. Himal.; Burma.
miniata, *Lindl. Gen. et Sp. Orch.* 42.—Java.
nervillosa, Reichb. f. in Walp. Ann. vi. 236 = Pholidota nervosa.
nervosa, A. Rich. in Ann. Sc. Nat. Sér. II. xv. (1841) 16 = corrugata.
nervosa, Wight, Ic. t. 1638 = glandulosa.
nigrescens, *P. Don, ex Loud. Hort. Brit. Suppl.* iii. 520, nomen.—Ind. or.
nitida, *Lindl. in Wall. Cat.* n. 1954.—Reg. Himal.
ocellata, *Lindl. Gen. et Sp. Orch.* 40.—Ind. or.
ocellata, Lindl. in Wall. Cat. n. 1953 = nitida, ochracea, cristata.
ochracea, *Lindl. Bot. Reg.* (1846) t. 69.—Reg. Himal.
odoratissima, *Lindl. in Wall. Cat.* n. 1960.—Ind. or.
ovalis, *Lindl. Bot. Reg.* (1838) *Misc.* 91.—Reg. Himal.
pallida, Reichb. f. in Walp. Ann. vi. 238 = Pholidota imbricata.
pandurata, *Lindl. in Gard. Chron.* (1853) 791.—Malaya.
Papagena, Reichb. f. in Bot. Zeit. xx. (1862) 214 = micrantha.
Parishii, *Hook. f. in Bot. Mag.* t. 5323.—Burma.
parviflora, Lindl. Gen. et Sp. Orch. 44 = Panisea parviflora.
peltastes, *Reichb. f. in Gard. Chron.* (1880) II. 296.—Borneo.
Pholas, Reichb. f. in Walp. Ann. vi. 237 = Pholidota Pholas.
pilosissima, Planch. Hort. Donat. 144 = ovalis.
plantaginea, *Lindl. in Gard. Chron.* (1855) 20.—Ind. or.
porrecta, Reichb. f. in Walp. Ann. vi. 236 = Otochilus porrecta.
praecox, *Lindl. Collect. Bot.* sub t. 37.—Reg. Himal.
prolifera, *Lindl. in Wall. Cat.* n. 1956.—Reg. Himal.
psittacina, *Reichb. f. Xenia Orch.* ii. 141.—Ins. Molucc.
pulverula, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxiv. (1862) 306.—Sumatra.
pumila, Reichb. f. in Walp. Ann. vi. 236 = Dendrochilum pumilum.
punctulata, Lindl. Collect. Bot. sub t. 33 = nitida.
recurva, Reichb. f. in Walp. Ann. vi. 237 = Pholidota recurva.
Reichenbachiana, *T. Moore, in Gard. Chron.* (1868) 1210.—Burma.
Rhodeana, Reichb. f. l. c. (1867) 901 = Rossiana.
rhombophora, *Reichb. f. in Linnaea*, xli. (1877) 116.—Ins. Manilla.
rigida, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 146.—Burma.
Rochusseni, *De Vriese, Illustr. Orch.* [t. 2. et t. 11. f. 6].—Java.
Rossiana, *Reichb. f. in Gard. Chron.* (1884) II. 808.—Burma.
rubra, Reichb. f. in Walp. Ann. vi. 238 = Pholidota rubra.
Rumphii, *Lindl. Fol. Orchid. Coelog.* 14.—Malaya.
salmonicolor, *Reichb. f. in Gard. Chron.* (1883) II. 328.—Ins. Sundaic.
Schilleriana, *Reichb. f. in C. Koch, Allg. Gartenz.* xxvi. (1858) 189.—Burma.
simplex, *Lindl. Fol. Orchid. Coelog.* 13.—Ind. or.
sparsa, *Reichb. f. in Gard. Chron.* (1883) I. 306.—Ins. Philipp.
speciosa, *Lindl. Gen. et Sp. Orch.* 39.—Malaya.
sulphurea, *Reichb. f. in Bonplandia*, v. (1857) 43.—Malaya.

COELOGYNE:—
testacea, *Lindl. Bot. Reg.* (1842) *Misc.* 38.—Malaya.
Thuniana, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiii. (1855) 145.—Nepal.
tomentosa, *Lindl. Fol. Orchid. Coelog.* 3.—Malaya.
trifida, Reichb. f. in Hamb. Gartenz. xix. (1863) 546 = odoratissima.
trinervis, *Lindl. in Wall. Cat.* n. 1955.—Burma.
triotos, *Reichb. f. in Hamb. Gartenz.* xviii. (1862) 34.—Ins. Philipp.
triplicatula, Reichb. f. in Bot. Zeit. xxii. (1864) 415 = fuliginosum?
triptera, A. Brongn. in Duperr. Voy. Coq. Bot. 201 = Epidendrum caespitosum, *Poepp. & Endl.*
trisaccata, Griff. Notul. iii. 400 = Gardneriana.
undulata, Reichb. f. in Walp. Ann. vi. 238 = Pholidota rubra.
undulata, *Wall. ex Voigt, Hort. Suburb. Calc.* 621, nomen.—Ind. or.
uniflora, *Lindl. in Wall. Cat.* n. 1966.—Reg. Himal.; Burma.
ustulata, *Par. & Reichb. f. in Trans. Linn. Soc.* xxx. (1874) 146.—Burma.
ventricosa, Reichb. f. in Walp. Ann. vi. 237 = Pholidota ventricosa.
viscosa, *Reichb. f. in Otto & Dietr. Allg. Gartenz.* xxiv. (1856) 218.—Reg. Himal.
Wallichiana, Lindl. Gen. et Sp. Orch. 43 = praecox.

COELONEMA, Maxim. in Bull. Acad. Pétersb. xxvi. (1880) 423. *CRUCIFERAE.*
draboides, *Maxim. l. c.* 424.—China.

COELOPLEURUM, Ledeb. Fl. Ross. ii. 361 (1844) = **Archangelica**, Hoffm. (Umbellif.).
Gmelini, Ledeb. l. c. = A. Gmelini.

COELOPYRUM, Jack, in Malay. Misc. ii. (1822) VII. 65. *INC. SED.*
coriaceum, *Jack, l. c.*—Malaya.

COELORACHIS, Brongn. in Duperr. Voy. Coq. Bot. 64. t. 14 (1829) = **Rottboellia**, Linn. (Gramin.).
hirsuta, Brongn. l. c. 65, in obs. = R. hirsuta.
muricata, Brongn. l. c. 65. t. 14 = R. glandulosa.

COELOSPERMUM, Blume, Bijdr. 994 (1826) (Caelospermum). *RUBIACEAE*, Benth. & Hook. f. ii. 119.
POGONOLOBUS, F. Mucll. Fragm. i. 55 (1858).
Balansaeanum, *Baill. Adansonia*, xii. (1876–79) 236.—N. Caled.
barbatum, *Span. in Linnaea*, xv. (1841) 318.—Ins. Timor.
corymbosum, *Blume, ex DC. Prod.* iv. 468.—Java.
decipiens, *Baill. in Bull. Soc. Linn. Par.* i. (1879) 218.—Austral.
Gmelini, Miq. Ann. Mus. Bot. Lugd. Bat. iii. 62, sphalm. = Archangelica Gmelini.
paniculatum, *F. Muell. Fragm.* v. 19.—Austral.
reticulatum, *Benth. Fl. Austral.* iii. 425.—Austral.
scandens, *Blume, Bijdr.* 994.—Malaya.

COELOSTEGIA, Benth. in Benth. & Hook. f. Gen. i. 213 (1862). *MALVACEAE*, Benth. & Hook. f. l. c.
Griffithii, *Benth. l. c.*—Malacca.

COELOSTELMA, Fourn. in Mart. Fl. Bras. vi. IV. 320 (1885). *ASCLEPIADEAE.*
refractum, *Fourn. l. c.*—Bras.

COELOSTYLIS, Torr. & Gray, ex Endl. Nov. Stirp. Dec. 33 (1839) = **Spigelia**, Linn. (Loganiac.).
loganioides, Torr. & Gray, l. c. = S. loganioides.
texana, Torr. & Gray, Fl. N. Am. ii. 44 = S. texana.

COEMANSIA, March. in Bull. Acad. Belg. Sér. II. xlvii. (1879) 94. *ARALIACEAE.*
COUDENBERGIA, March. l. c. 514 (1879).
Warmingiana, *March. l. c.* 95.—Am. austr.

COENOLOPHIUM, Reichb. Consp. 143 (1828) = CENOLOPHIUM, Koch = **Selinum**, Linn. (Umbell.).

COENOTUS, Benth. & Hook. f. Gen. ii. 281 (1873), lapsu = CAENOTUS, Rafin. = **Erigeron**, Linn. (Compos.).

COERULINIA, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 128 = **Veronica**, Tourn. (Scrophular.).
montana, Fourr. l. c. = V. montana.

COETOCAPNIA, Link & Otto, Ic. Pl. Rar. 35. t. 18 (1828) = **Bravoa**, La Llav. (Amaryllid.).
geminiflora, Link & Otto, l. c. = B. geminiflora.

COFFEA, Linn. Syst. ed. I (1735). *RUBIACEAE*, Benth. & Hook. f. ii. 114.
HEXEPTA, Rafin. Sylva Tellur. 164 (1838).
LACHNASTOMA, Korth. in Ned. Arch. Kruidk. ii. II. (1851) 201.
acuminata, *Ruiz & Pav. Fl. Per.* ii. 64.—Peruv.
Afzelii, *Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 174.—Afr. trop.
alpestris, Wight, Ic. t. 1040 = Webera lucens.
angustifolia, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 541.—Malaya.
arabica, Benth. in Hook. Niger Fl. 413 = liberica, stenophylla.
arabica, *Linn. Sp. Pl.* 172.—Arabia; Afr. trop.
australis, Vell. Fl. Flum. 62; ii. t. 13 = Faramea australis.
bengalensis, *Roxb. Hort. Beng.* 15; *Fl. Ind.* i. 540.—Ind. or.; Malaya.
bidentata, D. Dietr. Syn. Pl. i. 776 = microcarpa.
biflora, Vell. Fl. Flum. 63; ii. t. 16 = Coussarea biflora.
Boryana, D. Dietr. Syn. Pl. i. 777 = Chasalia Boryana.
brachyphylla, *Radlk. in Bremen Abh.* viii. (1883) 390.—Ins. Nossibé.
brasiliensis, Walp. Rep. ii. 477 = Rudgea lanceolata.
brevipes, *Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 172.—Afr. trop.
calycina, *Benth. in Hook. Journ. Bot.* iii. (1841) 232.—Guiana.
capitata, D. Dietr. Syn. Pl. i. 777 = Chasalia capitata.
capitata, Sieber, ex DC. Prod. iv. 531 = Chasalia divaricata?
Chamissonis, Hook. & Arn. Bot. Beech. Voy. 86 = Straussia kaduana.
chasalioides, D. Dietr. Syn. Pl. i. 777 = Chasalia divaricata?
ciliata, *Ruiz & Pav. Fl. Per.* ii. 65. t. 216.—Peruv.
clusiaefolia, D. Dietr. Syn. Pl. i. 777 = Chasalia clusiaefolia.
coriacea, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 201 = laurifolia.
corymbulosa, *Bertol. Fl. Guatimal.* 10.—Guatemala.
crassiloba, Benth. in Hook. Journ. Bot. iii. (1841) 233 = Rudgea Schomburgkiana.
cymosa, Wilid. ex Roem. & Schult. Syst. v. 201 = Chasalia Fontanesii.
densiflora, *Blume, Bijdr.* 965.—Java.
Deppeana, *Steud. Nom.* ed. II. i. 394.—Mexic.
didymocarpa, Bartl. ex DC. Prod. iv. 481 = Declieuxia psychotroides.
divaricata, Tausch, ex DC. l. c. 532 = Chasalia divaricata.
elliptica, Thw. Enum. Pl. Zeyl. 154 = Byrsophyllum ellipticum.
elongata, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 254.—Sumatra.
eriantha, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 534 = Rudgea eriantha.
flavicans, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 201 = Faramea jasminoides.
floribunda, Mart. in Flora, xxiv. (1841) II. Beibl. 85 = Ixora densiflora.
floribunda, *Miq. in Linnaea*. xxii. (1849) 804.—Bras.
Fontanesii, D. Dietr. Syn. Pl. i. 777 = Chasalia Fontanesii.
foveolata, *Ruiz & Pav. Fl. Per.* ii. 65. t. 216.—Peruv.
fragrans, *Wall. ex Hook. f. Fl. Brit. Ind.* iii. 154.—Ind. or.; Burma.
gardenioides, Cham. in Linnaea, ix. (1834) 230 = Rudgea gardenioides.
glabra, *Korth. in Nederl. Kruidk. Arch.* ii. II. (1851) 254.—Borneo.
grandifolia, Boj. ex Baker, Fl. Maurit. 152 = macrocarpa.
grandifolia, D. Dietr. Syn. Pl. i. 777 = Chasalia grandifolia.
grumelioides, Wight, Ic. t. 1041 = Webera lucens.
guianensis, *Aubl. Pl. Gui.* i. 150. t. 57.—Guiana.
hirsuta, G. Don, Gen. Syst. iii. 581 = Cremaspora africana.

COFFEA :—
Horsfieldiana, Miq. Fl. Ind. Bat. ii. 308 = bengalensis.
Humblotiana, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 514.—Ins. Comor.
hypoglauca, *Welw. ex Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 173.—Afr. trop.
indica, *Poir. Encyc. Suppl.* ii. 14.—Java.
jasminoides, Cham. in Linnaea, ix. (1834) 222 = Rudgea jasminoides.
jasminoides, *Welw. ex Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 175.—Afr. trop.
javanica, Blume, Cat. Gew. Buitenz. 46 = Chiococca javanica.
Jenkinsii, *Hook. f. Fl. Brit. Ind.* iii. 155.—Reg. Himal.
kaduana, Cham. & Schlecht. in Linnaea, iv. (1829) 33 = Straussia kaduana.
khasiana, *Hook. f. Fl. Brit. Ind.* iii. 154.—Reg. Himal.
Kraussiana, Hochst. in Flora, xxv. (1842) 237 = Kraussia floribunda.
lanceolata, Cham. in Linnaea, ix. (1834) 232 = Rudgea lanceolata.
lanceolata, Cham. & Schlecht. l. c. vi. (1831) 412 = Deppeana.
laurifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 372.—Venezuela.
laurifolia, Salisb. Prod. 62 = arabica.
laurina, Poir. in Lam. Encyc. Suppl. ii. 14 = Craterispermum laurinum.
laurina, *Smeathm. ex DC. Prod.* iv. 499.—Afr. trop.
lepidophloia, *Miq. Fl. Ind. Bat. Suppl.* 548.—Ins. Banca.
liberica, *Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 171. t. 24.—Afr. trop.
longifolia, *Ruiz & Pav. Fl. Per.* ii. 66. t. 218.—Peruv.
luçoniensis, Cham. & Schlecht. in Linnaea, iv. (1829) 32 = Psychotria luzoniensis.
macrocarpa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 168.—Ins. Mascar.
macrophylla, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. ii. 344 = subsessilis.
madurensis, *Teijsm. & Binn. Cat. Hort. Bog.* 112.—Ins. Madur.
magnoliaefolia, Cham. in Linnaea, ix. (1834) 225 = Rudgea magnoliaefolia.
major, Cham. l. c. 226 = Rudgea major.
marginata, *Benth. Pl. Hartw.* 193.—N. Granat.
Mariniana, Cham. & Schlecht. in Linnaea, iv. (1829) 35 = Straussia Mariniana.
mauritiana, *Lam. Encyc.* i. 550.—Ins. Mascar.
melanocarpa, *Welw. ex Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 173.—Afr. trop.
meridionalis, Vell. Fl. Flum. 62; ii. t. 14 = Coussarea meridionalis.
mexicana, *DC. Prod.* iv. 501.—Mexic.
microcarpa, DC. l. c. 499 = Perrottetii.
microcarpa, *Ruiz & Pav. Fl. Per.* ii. 66. t. 218.—Peruv.
minor, Cham. in Linnaea, ix. (1834) 227 = Rudgea Clausseniana.
Moka, Hort. ex Heynh. Nom. ii. 153 = arabica.
mozambicana, DC. Prod. iv. 500 = racemosa, *Lour.*
myrtifolia, *Roxb. Hort. Beng.* 15, nomen.—Ins. Maurit.
neurophylla, *Miq. Fl. Ind. Bat.* ii. 1079.—Java.
nitida, *Ruiz & Pav. Fl. Per.* ii. 65. t. 217.—Peruv.
nodosa, Cham. in Linnaea, ix. (1834) 233 = Rudgea nodosa.
novoguineensis, *Miq. in Ann. Mus. Bot. Lugd. Bat.* iv. 259.—N. Guinea.
obovata, *Cham. & Schlecht. in Linnaea*, vii. (1832) 412.—Mexic.
obovata, D. Dietr. Syn. Pl. i. 777 = Chasalia divaricata.
occidentalis, Jacq. Enum. Pl. Carib. 16 = Faramea odoratissima.
occidentalis, Vell. Fl. Flum. 62; ii. t. 12 = Faramea occidentalis.
odorata, *Forst. f. Prod.* 16.—Ins. Amicor.
oleifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 372.—N. Granat.
opulina, Forst. f. Prod. 16 = Pavetta opulina.
paniculata, *Aubl. Pl. Gui.* i. 152. t. 58.—Guiana.
parquioides, Cham. in Linnaea, ix. (1834) 224 = Rudgea parquioides.
parvifolia, Cham. l. c. 231 = Rudgea parvifolia.
pedunculata, *Roxb. Hort. Beng.* 86; *Fl. Ind.* i. 541.—Ins. Molucc.
Perrottetii, *Steud. ex Buek, DC. Prod. Index*, i. praef. p. ix.—Afr. trop.
porophylla, Vell. Fl. Flum. 62; ii. t. 11 = Coussarea porophylla.

COFFEA :—
psychotrioides, D. Dietr. Syn. Pl. i. 777 = Gaertnera psychotrioides.
racemosa, *Lour. Fl. Cochinch.* 145.—Afr. trop.
racemosa, *Ruiz & Pav. Fl. Per.* ii. 64. t. 214.—Peruv.
rachiformis, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 514.—Ins. Comor.
ramosa, Roem. & Schult. Syst. v. 198 = racemosa, *Lour.*
rosea, *Moç. & Sesse, ex DC. Prod.* iv. 499.—Mexic.
rupestris, *Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 174.—Afr. trop.
salicifolia, *Mig. Fl. Ind. Bat.* ii. 307.—Java.
sambucina, Forst. f. Prod. 16 = Pavetta sambucina.
semiexserta, *Colebr. ex Wall. in Roxb. Fl. Ind. ed. Carey*, ii. 195.—Ind. or.
sessilis, Vell. Fl. Flum. 64; ii. t. 20 = Psychotria axillaris.
spicata, *H. B. & K. Nov. Gen. et Sp.* iii. 371. t. 286.—N. Granat.
stenophylla, *G. Don, Gen. Syst.* iii. 581.—Abyssin.
stipulacea, *DC. Prod.* iv. 499.—Guiana.
stipulacea, Steud. Nom. ed. II. i. 395 = seq.
stipulata, Vell. Fl. Flum. 63; ii. t. 17 = Bathysa stipulata.
subcordata, *Hiern, in Trans. Linn. Soc.* Ser. II. i. (1876) 174.—Afr. trop.
subsessilis, Benth. in Hook. Journ. Bot. iii. (1841) 232 = Mapouria subsessilis.
subsessilis, *Ruiz & Pav. Fl. Per.* ii. 64. t. 215.—Peruv.
sundana, *Mig. Fl. Ind. Bat.* ii. 306.—Java.
sylvestris, Willd. ex Roem. & Schult. Syst. v. 201 = mauritiana.
tenuiflora, *Benth. in Hook. Journ. Bot.* iii. (1841) 232.—Guiana.
tetrandra, Roxb. Hort. Beng. 85; Fl. Ind. i. 538 = Prismatomeris albidiflora.
travancorensis, *Wight & Arn. Prod.* 435.—Ind. or.
triflora, Forst. f. Prod. 16 = Pavetta triflora.
triflora, Moon, Cat. 15 = travancorensis.
truncata, Vell. Fl. Flum. 63; ii. t. 15 = Faramea truncata.
umbellata, *Ruiz & Pav. Fl. Per.* ii. 64. t. 215.—Peruv.
umbellata, Vell. Fl. Flum. 63; ii. t. 19 = Faramea salicifolia.
verticillata, *Ruiz & Pav. Fl. Per.* ii. 66. t. 217.—Peruv.
verticillata, Vell. Fl. Flum. 63; ii. t. 18 = Ixora verticillata.
viburnoides, Cham. in Linnaea, ix. (1834) 228 = Rudgea viburnoides.
volubilis, Blanco, Fl. Filip. ed. I. 157 = Morinda tinctoria.
vulgaris, Moench, Meth. 504 = arabica.
Wightiana, *Wall. Cat.* n. 6246; *Wight & Arn. Prod.* 436.—Ind. or.
Zanguebariae, *Lour. Fl. Cochinch.* i. 145.—Afr. trop.

COGNIAUXELLA, Baill. in Bull. Soc. Linn. Par. i. (1884) 424 = **Cogniauxia**, Baill. (Cucurb.).

COGNIAUXIA, Baill. in Bull. Soc. Linn. Par. i. (1884) 423. *CUCURBITACEAE.*
COGNIAUXELLA, Baill. l. c. 424.
podolaena, *Baill. l. c.* 424.—Madag. ?

COGSWELLIA, Schult. in Roem. & Schult. Syst. vi. p. xlviii (1820) = **Peucedanum**, Tourn. (Umbellif.).
villosa, Schult. l. c. 588 = P. foeniculaceum.

COGYLIA, Molina, Sagg. Chil. ed. II. 300 (1810) = **Lardizabala**, Ruiz & Pav. (Berberid.).
biternata, Molina, l. c. = L. biternata.
ternata, Molina, l. c. = L. trifoliata.
triternata, Molina, l. c. = L. triternata.

COHAUTIA, Endl. Ench. 273 (1841) = KOHAUTIA, Cham. & Schlecht. = **Oldenlandia**, Linn. (Rubiac.).

COHIBA, Rafin. Fl. Tellur. iii. 75 (1836) = **Wigandia**, H. B. & K. (Hydrophyllac.).
urens, Rafin. l. c. = W. pruritiva.

COHNIA, Kunth, Enum. Pl. v. 35 (1850) = **Cordyline**, Comm. (Liliac.).
floribunda, Kunth, l. c. 36 = Cord. floribunda,
macrophylla, Kunth, l. c. 37 = Cord. macrophylla.
mauritiana, Baker, in Journ. Linn. Soc. xiv. (1875) 546 = Cord. macrophylla.
neocaledonica, Baker, l. c. = Cord. neocaledonica.
parviflora, Kunth, Enum. Pl. v. 37 = Cord. floribunda.

COHNIA, Reichb. f. in Bot. Zeit. x. (1852) 928. *ORCHIDEAE*, Benth. & Hook. f. iii. 477.
quekettioides, *Reichb. f. l. c.*—Nicaragua.

COILADENA, Rafin. Fl. Tellur. ii. 12 (1836) = **Ipomoea**, Linn. (Convolvulac.).
hyemalis, Rafin. l. c. = I. Horsfalliae.

COILANTHA, Borckh. in Roem. Arch. i. 1. (1796) 25 = **Gentiana**, Tourn.
asclepiadea, G. Don, Gen. Syst. iv. 186 = G. asclepiadea.
biloba, Bercht. & Presl, Rostl. i. Gentian. 6 = G. lutea.
Burseri, Bercht. & Presl, l. c. = G. Burseri.
campanulata, G. Don, Gen. Syst. iv. 185 = G. punctata.
glauca, G. Don, l. c. = G. glauca.
Mocinni, D. Don, ex G. Don, l. c. = G. calyculata.
pannonica, G. Don, l. c. = G. pannonica.
punctata, G. Don, l. c. = G. punctata.
purpurea, Borckh. in Roem. Arch. i. 1. (1796) 25 = G. purpurea.
Sessaei, D. Don, ex G. Don, Gen. Syst. iv. 185 = G. Sessaei.

COILANTHERA, Rafin. Sylva Tellur. 38 (1838) = **Cordia**, Linn. (Boragin.).
rotundifolia, Rafin. l. c. = Cordia rotundifolia.

COILMEROA, Endl. Gen. Suppl. iii. 99 (1843) = COLMEIROA, Reut. = **Securinega**, Juss. (Euphorbiac.).

COILONOX, Rafin. Fl. Tellur. ii. 28 (1836) = **Ornithogalum**, Tourn. (Liliac.).
albucoides, Rafin. l. c. = O. suaveolens.

COILOSPERMA, Rafin. Fl. Tellur. iii. 43 (1836) = **Deeringia**, R. Br. (Amarantac.).
cordata, Rafin. l. c. = D. baccata.

COILOSTIGMA, Klotzsch, in Linnaea, xii. (1838) 234 = **Scyphogyne**, Brongn. (Ericac.).
Dregeanum, Klotzsch, l. c. 235.—Afr. austr.
glabrum, Benth. in DC. Prod. vii. 708.—Afr. austr.
puberulum, Benth. l. c. = S. puberula.
tenuifolium, Klotzsch, in Linnaea, xii. (1838) 234.—Afr. austr.
Zeyherianum, Klotzsch, l. c.—Afr. austr.

COILOSTYLIS, Rafin. Fl. Tellur. iv. 37 (1836) = **Epidendrum**, Linn. (Orchid.).
emarginata, Rafin. l. c. = E. ciliare.
obtusifolia, Rafin. l. c. = E. lineare ?

COILOTAPALUS, P. Br. Hist. Jamaic. 111 (1756) = **Cecropia**, Loefl. (Urticac.).

COINOCHLAMYS, T. Anders. ex Benth. & Hook. f. Gen. ii. 1091 (1876). *ACANTHACEAE*, Benth. & Hook. f. l. c.
angolana, *S. Moore, in Journ. Bot.* xiv. (1876) 322.—Afr. trop.
hirsuta, *T. Anders. ex Benth. & Hook. f. Gen.* ii. 1091.—Afr. trop.

COINOGYNE, Less. in Linnaea, vi. (1831) 520 = **Jaumea**, Pers. (Compos.).
carnosa, Less. l. c. 521 = J. carnosa.

COIX, Linn. Gen. ed. I. 280 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1112.
LACRYMA, Medic. Phil. Bot. i. 177 (1789).
LACRYMA-JOB, Tourn. ex Linn. Gen. ed. I 280 (1737).
LACHRYMARIA, Heist. Syst. 12 (1748).
LITHAGROSTIS, Gaertn. Fruct. i. 7. t. 1 (1788).
SPHAERIUM, Linn. Syst. ed. I (1735).
agrestis, *Lour. Fl. Cochinch.* 551.—Amboina; China.
angulata, Mill. Gard. Dict. ed. VIII. n. 2 = Tripsacum dactyloides.
aquatica, *Roxb. Hort. Beng.* 66; *Fl. Ind.* iii. 571.—Ind. or.
arundinacea, Koen. ex Willd. Sp. Pl. iv. 203 = Chionachne barbata.
arundinacea, Lam. Encyc. iii. 422 = C. Lacryma-Jobi.
barbata, Roxb. Hort. Beng. 66; Fl. Ind. iii. 569 = Chionachne barbata.

COIX:—
crypsoides, *C. Muell. in Bot. Zeit.* xix. (1861) 334.—Ind. or.
dactyloides, Linn. Sp. Pl. 972 = Tripsacum dactyloides.
exaltata, Jacq. ex Spreng. Syst. i. 228 = C. Lachryma-Jobi.
gigantea, Koen. ex Roxb. Hort. Beng. 66; Fl. Ind. iii. 570 = C. Lachryma-Jobi.
heteroclita, Roxb. ll. cc. 66; 572 = Polytoca bracteata.
Koenigii, Spreng. Syst. i. 228 = Chionachne barbata.
Lacryma, Linn. Syst. ed. X. 1261 = C. Lachryma-Jobi.
Lacryma-Jobi, *Linn. Sp. Pl.* 972.—As. trop.
Ma-Yûên, *Romanet, in Bull. Soc. Acclimat.* Sér. II. viii. (1881) 442.—Cochinch.; China; Burma.
ovata, Stokes, Bot. Mat. Med. iv. 342 = C. Lachryma-Jobi.
pendula, Salisb. Prod. 28 = C. Lachryma-Jobi.
pumila, *Roxb. Hort. Beng.* 66; *Fl. Ind.* iii. 572.—Ins. Maurit.
stigmatosa, *C. Koch & Bouché, Ind. Sem. Hort. Berol.* (1855) 9.—Hab. ?

COLA, Schott & Endl. Meletem. 33 (1832). *STERCULIACEAE*, Benth. & Hook. f. i. 218.
BICHEA, Stokes, Bot. Mat. Med. ii. 564 (1812).
COLARIA, Rafin. Sylva Tellur. 73 (1838).
COURTENIA, R. Br. in Benn. Pl. Jav. Rar. 236 (1844).
EDWARDIA, Rafin. Specch. i. 158 (1824); ex DC. Prod. ii. 92 (1825).
LUNANEA, DC. Prod. ii. 92 (1825).
SIPHONIOPSIS, Karst. Fl. Columb. i. 139. t. 69 (1858).
acuminata, *Schott & Endl. Meletem.* 33.—Afr. trop.
Afzelii, *Mast. in Oliver, Fl. Trop. Afr.* i. 223.—Afr. trop.
argentea, *Mast. l. c.* 224.—Afr. trop.
cauliflora, *Mast. l. c.* 221.—Afr. trop.
clavata, *Mast. l. c.* 222.—Afr. trop.
cordifolia, [*R. Br. in*] *Benn. Pl. Jav. Rar.* 237.—Java.
digitata, *Mast. in Oliver, Fl. Trop. Afr.* i. 224.—Afr. trop.
ficifolia, *Mast. l. c.* 223.—Afr. trop.
gabonensis, *Mast. l. c.* 222.—Afr. trop.
grandiflora, *Schott & Endl. Meletem.* 33.—Hab. ?
heterophylla, *Schott & Endl. l. c.; Mast. in Oliver, Fl. Trop. Afr.* i. 223.—Afr. trop.
laurifolia, *Mast. in Oliver, l. c.* 222.—Afr. trop.
macrocarpa, Schott & Endl. Meletem. 33 = acuminata.
natalensis, *Oliver, in Hook. Ic. Pl.* t. 1390.—Afr. austr.
nitida, Schott & Endl. Meletem. 33 = acuminata.
quinqueloba, *Garcke, in Peters, Reise Mossamb. Bot.* 130.—Afr. trop.
tomentosa, Schott & Endl. Meletem. 33 = Sterculia tomentosa.

COLARIA, Rafin. Sylva Tellur. 73 (1838) = **Cola**, Schott & Endl. (Stercul.).
acuminata, Rafin. l. c. = Cola acuminata.
heterophyla, Rafin. l. c. = Cola heterophylla.

COLAX, Lindl. ex Spreng. Syst. iii. 727 (1826); et Bot. Reg. (1843) Misc. 50 = **Lycaste**, Lindl. (Orchid.).
aromaticus, Spreng. Syst. iv. Cur. Post. 307 = L. aromatica.
Barringtoniae, Lindl. ex Spreng. Syst. iii. 727 = L. Barringtoniae.
grandiflorus, Rafin. Fl. Tellur. ii. 85 = Bifrenaria Harrisoniae.
Harrisoniae, Lindl. ex Spreng. Syst. iii. 727 = Bifrenaria Harrisoniae.
jugosus, Lindl. Bot. Reg. (1843) Misc. 51.—Bras.
modestior, Reichb. f. in Hamb. Gartenz. xvi. (1860) 14.—Bras.
pallidiflorus, Spreng. f. Tent. 29 = Maxillaria pallidiflora.
palmifolius, Lindl. ex Spreng. Syst. iii. 727 = Maxillaria palmifolia.
Parkeri, Spreng. f. Tent. 28 = Maxillaria Parkeri.
parvulus, Spreng. Syst. iv. Cur. Post. 307 = Bifrenaria parvula.
placantherus, Lindl. Bot. Reg. (1843) Misc. 50.—Bras.
Puydtii, Linden & André, in Illustr. Hortic. xxvii. (1880) t. 369.—Bras.
racemosus, Spreng. f. Tent. 29 = Bifrenaria racemosa.
viridis, Lindl. Bot. Reg. (1843) Misc. 50 = L. viridis.

COLBERTIA, Salisb. Parad. Lond. sub t. 73 (1807) = **Dillenia**, Linn.
augusta, Wall. Cat. n. 948 = D. pentagyna.
coromandelina, DC. Syst. i. 435 = D. pentagyna.
floribunda, Wall. Cat. n. 950 = D. floribunda.
minor, Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Indie, ii. (1845) 579.—Java.
obovata, Blume, Bijdr. 6 = D. aurea.
scabrella, D. Don, Prod. Fl. Nep. 226 = D. scabrella.

COLCHICUM, Linn. Syst. ed. I (1735); Gen. ed. I. 107 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 821.
FOUHA, Pomel, Mat. Fl. Atl. 2 (1860).
HERMODACTYLOS, Reichb. Consp. 64 (1828).
MONOCARYUM, R. Br. in App. Voy. Denh. et Clapp. 243 (1826).
PALUDARIA, Salisb. Gen. Pl. Fragm. 53 (1865).
SYNSIPHON, Regel, in Act. Hort. Petrop. vi. (1879) 490.
aegyptiacum, Boiss. Diagn. Ser. I. v. 66 = montanum.
aestivale, Bor. Fl. Centr. Fr. ed. III. ii. 612 = byzantinum.
aetnense, Tineo, in Guss. Fl. Sic. Syn. ii. 818 = neapolitanum.
agrippinum, *Baker, in Journ. Linn. Soc.* xvii. (1879) 425.—Hab. ?
Alberti, *Regel, in Act. Hort. Petrop.* viii. (1884) 647.—As. centr.
alpinum, *DC. Fl. Fr.* iii. 195.—Europ.
alpinum, Schimp. ex Nym. Consp. 742 = montanum.
amabile, *Heldr. in Atti Congr. Bot. Firenze*, 1874 (1876) 227.—Graecia.
arenarium, Koch, Syn. Fl. Germ. ed. I. 836 = neapolitanum.
arenarium, *Waldst. & Kit. Pl. Rar. Hung.* ii. 195. t. 179.—Ital.; Hungar.
atticum, Sprun. ex Tommas. in Flora, xxiii. (1840) 730 = Merendera attica.
autumnale, *Linn. Sp. Pl.* 341.—Europ.
Balansae, Planch. in Ann. Sc. Nat. Sér. IV. iv. (1855) 145 = laetum.
Bertolonii, Stev. in Nouv. Mém. Soc. Nat. Mosc. vii. (1829) 268 = montanum.
Bisignani, Tenore, fide Janka, ex Baker, in Journ. Linn. Soc. xvii. (1879) 427 = Tenorii.
Bivonae, *Guss. Adnot. Cat. Pl. Bocc.* (1821) 4.—Europ. austr.
Bivonae, Tenore, Fl. Nap. Prod. App. v. 11 = Tenorii.
Bivonae, Willk. & Lange, Prod. Fl. Hisp. i. 194, ex parte = lusitanum.
Boissieri, *Orph. in Atti Cong. Bot. Firenze*, 1874 (1876) 31.—Graecia.
brachyphyllum, *Boiss. & Haussk. ex Boiss. Fl. Orient.* v. 164.—Syria.
bulbocodioides, Bieb. Fl. Taur. Cauc. i. 293; iii. 281 = montanum.
bulbocodioides, Brot. Fl. Lusit. i. 597 = Merendera Bulbocodium.
Bulbocodium, Ker-Gawl. in Bot. Mag. sub t. 1028 = Bulbocodium vernum.
byzantinum, *Ker-Gawl. l. c.*—Graecia; As. Min.
byzantinum, Tenore, Fl. Nap. iii. 397 = Tenorii.
castrense, Laramb. in Bull. Soc. Bot. Fr. ii. (1855) 688 = neapolitanum.
candidum, Schott & Kotschy, ex Baker, in Journ. Linn. Soc. xvii. (1879) 429 = laetum.
catacuzenium, Heldr. ex Boiss. Fl. Orient. v. 165 = montanum.
caucasicum, Spreng. Syst. i. 143 = Merendera caucasica.
chionense, Haw. ex Kunth, Enum. Pl. iv. 139 = variegatum?
Clementei, Graells, in Mem. Acad. Cienc. Madr. ii. [1854] 483 = bulbocodioides.
commune, Neck. Delic. Gallo-Belg. i. 176 = autumnale.
corsicum, *Baker, in Journ. Linn. Soc.* xvii. (1879) 431.—Ins. Corsica.
crociflorum, *Sims, Bot. Mag.* t. 2673.—Europ. austr.
crociflorum, *Regel, in Act. Hort. Petrop.* vii. (1880) 385.—Kokan.
crociflorum, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 97 = montanum.
crocifolium, Boiss. Diagn. Ser. I. v. 67 = montanum.
Cupani, Guss. Fl. Sic. Prod. i. 452 = montanum.
Decaisnei, *Boiss. Fl. Orient.* v. 157.—Syria.
cubocum, Orph. in Atti Cong. Bot. Firenze, 1874 (1876) 28 = Bivonae.
fasciculare, *Boiss. Fl. Orient.* v. 163.—Syria.
fasciculare, R. Br. in Denh. & Clapp. Trav. 243 = montanum.

COLCHICUM:—
falcifolium, *Stapf, in Denkschr. Acad. Wien*, l. (1885) 19.—Persia.
floribundum, Lawson, Cat. 6; ex Salisb. in Trans. Hort. Soc. i. (1812) 329 = byzantinum.
Haussknechtii, *Boiss. Fl. Orient.* v. 157.—Persia.
Haynaldi, Heuff. in Verh. Zool.-Bot. Ges. Wien, viii. (1858) 213 = autumnale.
hexapetalum, Pourr. ex Schult. f. Syst. vii. 1521 = Merendera Bulbocodium.
hololophum, Coss. & Dur. in Bull. Soc. Bot. Fr. iv. (1857) 489 = montanum.
illyricum, Frivald. ex Kunth, Enum. Pl. iv. 140 = speciosum.
illyricum, *Stokes, Bot. Mat. Med.* ii. 331.—As. Min.
Jankae, Freyn, in Oestr. Bot. Zeitschr. xxvii. (1877) 361 = neapolitanum.
Kochii, Parl. Fl. Ital. iii. 188 = neapolitanum.
Kotschyi, Boiss. Diagn. Ser. I. xiii. 38 = laetum.
lactum, *Stev. in Nouv. Mém. Soc. Nat. Mosc.* vii. (1829) 262. t. 13.—Orient.
latifolium, Griseb. Spicil. Fl. Rumel. ii. 378 = speciosum.
latifolium, Sibth. & Sm. Fl. Graec. iv. 43. t. 350 = Sibthorpii, byzantinum.
Levieri, *Janka, in Oestr. Bot. Zeitschr.* xxv. (1875) 82.—Italia.
libanoticum, *Ehrenb. ex Boiss. Fl. Orient.* iii. 166.—Syria.
lingulatum, *Boiss. & Sprun. Diagn.* Ser. I. v. 66.—Europ.
longifolium, Castag. Cat. Pl. Marseille, 135 = neapolitanum.
lusitanum, *Brot. Phyt. Lusit.* ii. 211. t. 173, 174.—Lusitania.
luteum, *Baker, in Gard. Chron.* (1874) 33.—Reg. Himal.
Merendera, Ker-Gawl. in Bot. Reg. sub t. 541 = Merendera Bulbocodium.
micranthum, *Boiss. Fl. Orient.* v. 162.—Byzant.
minimum, Elwes, ex Baker, in Journ. Linn. Soc. xvii. (1879) 433 = montanum.
monogynum, Bieb. ex Schult. f. Syst. vii. 796 = Sternbergia caucasica.
montanum, All. Fl. Pedem. i. 117 = alpinum.
montanum, *Linn. Sp. Pl.* 342.—Europ. austr.
montanum, Linn. l. c. partim = Merendera Bulbocodium.
montanum, Schousb. Vextr. Marok. 185 = autumnale.
multiflorum, Brot. Fl. Lusit. i. 597 = autumnale.
neapolitanum, Orph. ex Nym. Consp. 743 = parnassicum.
neapolitanum, *Tenore, Fl. Neap. Prod. App.* v. 11.—Italia.
nivale, Boiss. & Huet, ex Baker, in Journ. Linn. Soc. xvii. (1879) 433 = montanum.
orientale, Frivald. ex Kunth, Enum. Pl. iv. 143 = neapolitanum.
pannonicum, Griseb. & Schenk, in Wiegm. Archiv, xviii. (1852) i. 359 = autumnale.
Parkinsoni, Hook. f. Bot. Mag. t. 6090 = variegatum.
Parlatoris, *Orph. in Atti Congr. Bot. Firenze*, 1874 (1876) 32.—Graecia.
parnassicum, *Sart. Orph. & Heldr. in Boiss. Diagn.* Ser. II. iv. 122.—Graecia.
parviflorum, Bivona, ex Parl. Fl. Ital. iii. 191 = montanum.
parvulum, Janka, in Oestr. Bot. Zeitschr. xxiii. (1873) 243 = micranthum.
parvulum, Tenore, Viag. in Basilic. 120 = alpinum.
patens, Schultz, in Flora, ix. (1826) 132 = autumnale.
persicum, *Baker, in Journ. Linn. Soc.* xvii. (1879) 430.—Persia.
polyanthon, Ker-Gawl. in Bot. Mag. sub t. 1028 = autumnale.
polymorphum, Orph. in Atti Congr. Bot. Firenze, 1874 (1876) 29 = neapolitanum.
polyphyllum, Boiss. & Heldr. Diagn. Ser. II. iv. 121 = Steveni.
praecox, Spenn. Fl. Friburg. i. 215 = autumnale.
provinciale, H. Loret, in Bull. Soc. Bot. Fr. vi. (1859) 459 = neapolitanum.
pulchrum, *Herb. ex Baker, in Journ. Linn. Soc.* xvii. (1879) 425.—Graecia.
pusillum, Sieber, in Flora, v. (1822) 248 = montanum.
pyrenaicum, Pourr. in Mém. Acad. Toul. iii. (1788) 316 = Merendera Bulbocodium.

COLCHICUM:—
Ritchii, R. Br. in App. Denh. & Clapp. Trav. 241 = montanum.
Sibthorpii, *Baker, in Journ. Linn. Soc.* xvii. (1879) 427.—Graecia.
speciosum, *Stev. in Nouv. Mém. Soc. Nat. Mosc.* vii. (1829) 265. t. 15.—Reg. Caucas.
Steveni, *Kunth, Enum. Pl.* iv. 144.—Syria; Arab.
Szovitsii, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. 24.—Reg. Caucas.
Tenorii, *Parl. Fl. Ital.* iii. 176.—Italia.
tessellatum, Hort. ex Baker, in Journ. Linn. Soc. xvii. (1879) 425 = agrippinum.
tesselatum, Link.—Cf. Kunth, Enum. Pl. iv. 139 = Bivonae.
tessellatum, Salisb. Prod. 236 = variegatum.
tessulatum, Mill. Gard. Dict. ed. VIII. n. 4 = variegatum.
Todarii, Parl. Fl. Ital. iii. 178 = lusitanum.
trapezunticum, Boiss. ex Baker, in Journ. Linn. Soc. xvii. (1879) 430 = polyphyllum.
transsilvanicum, Schur, Enum. Pl. Transs. 679 = byzantinum.
triphyllum, Kunze, in Flora, xxix. (1846) 755 = montanum.
Troodi, *Kotschy, in Unger & Kotschy, Cyprus*, 190.—Ins. Cyprus.
turcicum, *Janka, in Oestr. Bot. Zeitschr.* xxiii. (1873) 242.—Europ. or.
umbrosum, *Stev. in Nouv. Mém. Soc. Nat. Mosc.* vii. (1829) 268. t. 14.—Tauria.
Valery, Tineo, in Guss. Fl. Sic. Syn. ii. 818 = montanum.
variegatum, Bivona, Sic. Pl. Cent. i. 27 = Bivonae.
variegatum, *Linn. Sp. Pl.* 342.—Europ. austr.
variegatum, Sieber, ex Baker, in Journ. Linn. Soc. xvii. (1879) 427 = Sibthorpii.
variopictum, Janka, in Oestr. Bot. Zeitschr. xxv. (1875) 83 = amabile.
vernale, Hoffm. Deutschl. Fl. ed. II. i. 174 = autumnale.
vernum, Kunth, Enum. Pl. iv. 141 = autumnale.
vernum, Kunth, l. c. 146 = Bulbocodium vernum.
vernum, Pall. ex Bieb. Fl. Taur. Cauc. i. 293 = montanum.
versicolor, Ker-Gawl. in Bot. Reg. t. 571 = Bulbocodium versicolor.
Visianii, Parl. Fl. Ital. iii. 175 = Bivonae.

COLDENIA, Linn. Diss. Dass. 9 (1747); Amoen. Acad. i. 393 (1749). *BORAGINEAE*, Benth. & Hook. f. ii. 841.
EDDYA, Torr. & A. Gray, in Pacif. Rail. Rep. ii. 170 (1855).
GALAPAGOA, Hook. f. in Trans. Linn. Soc. xx. (1847) 196.
GOLDENIA, Raeusch. Nom. ed. III. 43 (1797).
LOBOPHYLLUM, F. Muell. in Hook. Kew Journ. ix. (1857) 21.
MONOMESIA, Rafin. Fl. Tellur. iv. 87 (1836).
PTILOCALYX, Torr. & A. Gray, in Pacif. Rail. Rep. ii. 170. t. 8 (1855).
STEGNOCARPUS, Torr. l. c. 169 (1855).
TIQUILIA, Pers. Syn. i. 157 (1805).
angolensis, *Welw. Apont.* 591.—Afr. trop.
atacamensis, *Phil. Fl. Atac.* 37.—Chili.
canescens, *DC. Prod.* ix. 559.—Am. bor. occ.
dichotoma, *Lehm. Asperifol.* 9.—Peruv.
Dombeyana, Buek, Ind. DC. Prod. iii. 108 = dichotoma.
Greggii, *A. Gray, Syn. Fl. N. Am.* ii. 1. 182.—Am. bor. occ.
hispidissima, *A. Gray, in Proc. Am. Acad.* v. (1862) 340.—Am. bor. occ.
litoralis, *Phil. Fl. Atac.* 37.—Chili.
mexicana, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 119.—Mexic.
Nuttallii, *Hook. in Hook. Kew Journ.* iii. (1851) 296.—Am. bor. occ.
Palmeri, *A. Gray, in Proc. Am. Acad.* viii. (1870) 292.—Am. bor. occ.
pentandra, Juss. ex Steud. Nom. ed. II. i. 396 = dichotoma.
procumbens, *Linn. Sp. Pl.* 125.—Ind. or.
tomentosa, *S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 120.—Am. bor. occ.

COLEA, Boj. Hort. Maurit. 220 (1837). *BIGNONIACEAE*, Benth. & Hook. f. ii. 1051.

COLEA:—
TRIPINNA, Lour. Fl. Cochinch. 391 (1790).
TRIPINNARIA, Pers. Syn. ii. 173 (1807).
ULOMA, Rafin. Fl. Tellur. ii. 62 (1836).
cauliflora, DC. Prod. ix. 241 = floribunda.
Chapelieri, DC. l. c. 242 = decora.
Commersonii, DC. l. c. = floribunda.
decora, *Boj. Hort. Maurit.* 220, nomen; *DC. l. c.* 241.—Madag.
discolor, *Seem. in Trans. Linn. Soc.* xxiii. (1860) 8.—Madag.
floribunda, *Boj. Hort. Maurit.* 220, nomen; *Lindl. Bot. Reg.* (1841) t. 19.—Madag.
hispidissima, *Seem. in Trans. Linn. Soc.* xxiii. (1860) 9.—Madag.
involucrata, *Boj. Hort. Maurit.* 221, nomen; *DC. Prod.* ix. 242.—Madag.
mauritiana, *Boj. l. c.* 220.—Ins. Mascar.
nitida, DC. Prod. ix. 242 = decora.
obtusifolia, DC. l. c. 241 = mauritiana.
parviflora, *Baker, in Journ. Linn. Soc.* xxi. (1885) 428.—Madag.
pedunculata, *Baker, Fl. Maurit.* 244.—Ins. Seychelles.
purpurascens, *Seem. in Trans. Linn. Soc.* xxiii. (1860) 8.—Madag.
ramiflora, DC. Prod. ix. 241 = mauritiana.
Seychellarum, *Seem. in Trans. Linn. Soc.* xxiii. (1860) 8.—Ins. Seychelles.
Sieberi, *A. DC. in DC. Prod.* ix. 240, in nota.—Hab.?
Telfairii, *Boj. Hort. Maurit.* 220.—Madag.
tetragona, *DC. Prod.* ix. 242.—Madag.
tripinnata, *Seem. in Bonplandia*, iv. (1856) 128.—Cochinch.
undulata, *Regel, Gartenfl.* (1870) 322.—Madag.?

COLEACHYRON, J. Gay, ex Boiss. Fl. Orient. v. 408 (1881) = **Carex**, Dill. (Cyperac.).
Oliverianum, J. Gay, l. c. = Carex illegitima.

COLEANTHERA, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) 1. 4. *EPACRIDEAE*, Benth. & Hook. f. ii. 611.
MICHIEA, F. Muell. Fragm. iv. 95. t. 27 (1864).
caelophylla, *Benth. Fl. Austral.* iv. 150.—Austral.
myrtioides, *Stschegl. in Bull. Soc. Nat. Mosc.* xxxii. (1859) 1. 4.—Austral.
virgata, *Stschegl. l. c.* 5.—Austral.

COLEANTHUS, Seid. in Roem. & Schult. Syst. ii. 11, 276 (1817). *GRAMINEAE*, Benth. & Hook. f. iii. 1147.
SCHMIDTIA, Tratt. Fl. Austr. i. 12. t. 451 (1811).
WILIBALDA, Sternb. in Flora, ii. (1819) 6.
subtilis, *Seid. in Roem. & Schult. Syst.* ii. 276.—Europ.

COLEATAENIA, Griseb. in Goett. Abh. xxiv. (1879) 308 = **Panicum**, Linn. (Gramin.).
gynerioides, Griseb. l. c.—Reg. Argent.

COLEBROOKEA, Sm. Exot. Bot. ii. 111. t. 115 (1806). *LABIATAE*, Benth. & Hook. f. ii. 1180.
oppositifolia, Lodd. Bot. Cab. t. 487 = Elsholtzia polystachya.
oppositifolia, *Sm. Exot. Bot.* ii. 111. t. 115.—Ind. or.; Burma.
ternifolia, Roxb. Hort. Beng. 45; Fl. Ind. iii. 25 = oppositifolia.

COLEBROOKIA, Donn, Cat. Hort. Cantab. i. (1796) = **Globba**, Linn. (Zingib.).
bulbifera, Donn, l. c. = G. marantina.

COLEBROOKIA, Spreng. Syst. ii. 713 = **Colebrookea**, Sm. (Labiat.).

COLEMA, Rafin. New Fl. Am. iii. 49 (1836) = **Corema**, D. Don (Empetrac.).
arenaria, Rafin. l. c. = Cor. Conradii.

COLENSOA, Hook. f. Fl. N. Zel. i. 156 (1853) = **Pratia**, Gaudich. (Campanulac.).
physaloides, Hook. f. l. c. 157.—N. Zel.

COLEOCOMA, F. Muell. in Hook. Kew Journ. ix. (1857) 19. *COMPOSITAE*, Benth. & Hook. f. ii. 293.
Centaurea, *F. Muell. l. c.*—Austral.

COLEOGYNE, Torr. in Smithson. Contrib. vi. (Plant. Frémont.) 8. t. iv. (1854). *ROSACEAE*, Benth. & Hook. f. i. 617.
ramosissima, *Torr. l. c.*—Calif.

COLEONEMA, Bartl. & Wendl. f. Diosm. 55. t. A (1824). *RUTACEAE*, Benth. & Hook. f. i. 289.
album, *Bartl. & Wendl. f. l. c.* 56.—Afr. austr.
album, E. Mey. Zwei Pfl. Docum. 174 = Diosma vulgaris.
aspalathoides, *A. Juss. in Mém. Mus. Par.* xii. (1825) 471.—Afr. austr.
Dregeanum, Presl, Bot. Bemerk. 31 = pulchrum.
filiforme, A. Juss. in Mém. Mus. Par. xii. (1825) 471 = pulchrum.
gracile, Eckl. & Zeyh. Enum. 106 = pulchrum.
juniperifolium, *Eckl. & Zeyh. l. c.*—Afr. austr.
juniperinum, Sond. in Harv. & Sond. Fl. Cap. i. 379 = juniperifolium.
pulchrum, *Hook. Bot. Mag.* t. 3340.—Afr. austr.
virgatum, Eckl. & Zeyh. Enum. 106 = pulchrum.
virginianum, Eckl. & Zeyh. l. c. = juniperifolium.

COLEOPHORA, Miers, in Ann. Nat. Hist. Ser. II. vii. (1851) 197 = **Daphnopsis**, Mart. (Thymel.).
gemmiflora, Miers, l. c. = D. Martii.

COLEOPHYLLUM, Klotzsch, in Otto & Dietr. Allg. Gartenz. viii. (1840) 185 = **Chlidanthus**, Herb. (Amaryll.).
Ehrenbergii, Klotzsch, l. c. = Chlidanthus Ehrenbergii.

COLEOSANTHUS, Cass. in Bull. Soc. Philom. (1817) 67; et Dict. Sc. Nat. x. (1818) 36 = **Brickellia**, Ell. (Compos.).
Cavanillesii, Cass. in Dict. Sc. Nat. x. 37 = B. Cavanillesii.
tiliaefolius, Cass. l. c. xxiv. 519 = Eupatorium macrophyllum.

COLEOSPADIX, Becc. in Ann. Jard. Buitenz. ii. (1885) 90. *PALMAE*.
litigiosa, *Becc. l. c.*—N. Guinea.
oninensis, *Becc. l. c.*—N. Guinea.

COLEOSTACHYS, A. Juss. in Ann. Sc. Nat. Sér. II. xiii. (1840) 329. *MALPIGHIACEAE*, Benth. & Hook. f. i. 253.
genipaefolia, *A. Juss. l. c.*—Bras.
hypoleuca, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 125 = Blepharandra hypoleuca.
spicata, Juss. in Arch. Mus. Par. iii. (1843) t. 5 = genipaefolia.
vestita, Benth. in Hook. Lond. Journ. Bot. vii. (1848) 124 = Diacidia vestita.

COLEOSTEPHUS, Cass. in Dict. Sc. Nat. xli. 43 (1826) = **Chrysanthemum**, Linn. (Compos.).
Clausonis, Pomel, Nouv. Mat. Fl. Atl. 59.—Algeria.
hybridus, Lange, in Kjoeb. Vidensk. Meddel. (1861) 77 (Pugill. 127) = Chrys. Myconis.
macrotus, Dur. in Duch. Rev. Bot. i. (1845–46) 363 = Chrys. macrotum.
multicaulis, Dur. l. c. 364 = Chrys. multicaule.
Myconis, Cass. in Dict. Sc. Nat. xli. 43 = Chrys. Myconis.

COLEOSTYLES, Benth. & Hook. f. Gen. ii. 535 [bis] (1873) = seq.

COLEOSTYLIS, Sond. in Lehm. Pl. Preiss. i. 391 (1845) = **Levenhookia**, R. Br. (Stylid.).
Preissii, Sond. l. c. = L. Preissii.
Sonderi, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 46 = L. Sonderi.
umbellulata, Sond. in Lehm. Pl. Preiss. i. 391 = L. stipitata.

COLEOTRYPE, C. B. Clarke, in DC. Monog. Phan. iii. 238. t. 8 (1881). *COMMELINACEAE*, Benth. & Hook. f. iii. 851.
Goudotii, *C. B. Clarke, l. c.* 240.—Madag.
madagascarica, *C. B. Clarke, l. c.* 239.—Madag.
natalensis, *C. B. Clarke, l. c.*—Afr. austr.

COLETIA, Vell. Fl. Flum. 32 (1825) = **Mayaca**, Aubl.
madida, Vell. l. c. 33; i. t. 79 = M. Sellowiana.

COLEUS, Lour. Fl. Cochinch. ii. 372 (1790). *LABIATAE*, Benth. & Hook. f. ii. 1176.
MITSA, Chap. ex Benth. Lab. Gen. et Sp. 52 (1832).
acuminatus, *Benth. in Linnaea*, vi. (1831) 81.—Ins. Philipp.
africanus, *Benth. Lab. Gen. et Sp.* 54.—Afr. trop.
albidus, *Vatke, in Linnaea*, xxxvii. (1871–73) 321.—Abyss.
alpinus, *Vatke, l. c.* 322.—Abyss.
amboinicus, Lour. Fl. Cochinch. ii. 372 = aromaticus.
anamallayensis, *Bedd. in Madr. Journ. Sc.* Ser. II. xxii. (1861) 73.—Ind. or.
arabicus, *Benth. in DC. Prod.* xii. 79.—Arab.
aromaticus, *Benth. in Wall. Pl. As. Rar.* ii. 16.—Ind. or.
atropurpureus, *Benth. l. c.*—Malaya; Pacific.
barbatus, *Benth. l. c.* 15.—Ind. or.; Afr. trop.
Benthamianus, Arn. in Nov. Act. Nat. Cur. xviii. (1836) 354 = inflatus.
bicolor, *Benth. Lab. Gen. et Sp.* 55.—Java.
Blancoi, *Benth. in DC. Prod.* xii. 79.—Ins. Philipp.
Blumei, *Benth. Lab. Gen. et Sp.* 56.—Java.
Bojeri, *Benth. l. c.* 52.—Madag.
caninus, *Vatke, in Linnaea*, xxxvii. (1871–73) 318.—Abyss.
carnosus, *Hassk. in Flora*, xxv. (1842) II. *Beibl.* 25.—Java.
comosus, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 183 = spicatus.
crassifolius, Benth. in Wall. Pl. As. Rar. ii. 15 = aromaticus.
edulis, *Vatke, in Linnaea*, xxxvii. (1871–73) 319.—Abyss.
flaccidus, *Vatke, in Linnaea*, xliii. (1880–82) 90.—Afr. trop.
fruticosus, Benth. in DC. Prod. xii. 78 = Plectranthus fruticosus.
fruticosus, *Wight, ex Benth. in DC. Prod.* xii. 78.—Ind. or.
galeatus, *Benth. Lab. Gen. et Sp.* 56.—Java.
Garckeanus, *Vatke, in Linnaea*, xxxvii. (1871–73) 323.—Abyss.
Gibsonii, *Verlot, in Rev. Hortic.* (1866) 279.—Hab.?
glabratus, Benth. Lab. Gen. et Sp. 58 = Plectranthus coleoides.
glandulosus, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 211.—Afr. trop.
grandifolius, *Benth. Lab. Gen. et Sp.* 53.—Ins. Timor.
grandifolius, Blanco, Fl. Filip. ed. I. 482 = Blancoi.
Heynei, Benth. Lab. Gen. et Sp. 50 = spicatus.
Huberi, *Regel, in Act. Hort. Petrop.* vii. (1880) 543.—Abyss.
igniarius, *Schweinf. Beitr. Fl. Aethiop.* 121.—Afr. trop.
inflatus, *Benth. Lab. Gen. et Sp.* 58.—Zeylan.
ingratus, *Benth. l. c.* 53.—Java.
laciniatus, *Benth. l. c.* 56.—Java.
lactiflorus, *Vatke, in Linnaea*, xliii. (1880–82) 89.—Afr. trop.
lanuginosus, *Hochst. ex Benth. in DC. Prod.* xii. 79.—Abyss.
latifolius, *Hochst. ex Benth. l. c.* 74.—Abyss.
leptostachys, *Benth. l. c.* 77.—Zeylan.
Macraei, Benth. Lab. Gen. et Sp. 58 = malabaricus.
macrophyllus, *Benth. l. c.* 55.—Java.
macropus, *Miq. Fl. Ind. Bat.* ii. 956.—Java.
macrostachys, *Benth. Lab. Gen. et Sp.* 57.—Java.
malabaricus, *Benth. in Wall. Pl. As. Rar.* ii. 16.—Ind. or.
Mannii, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 211.—Afr. trop.
micranthus, *Maxim. in Act. Hort. Petrop.* iii. (1875) 108.—Abyss.
mollis, Benth. in DC. Prod. xii. 77 = malabaricus.
multiflorus, *Benth. Lab. Gen. et Sp.* 55.—Ins. Philipp.
ovatus, Benth. l. c. 57 = malabaricus.
palustris, *Vatke, in Linnaea*, xxxvii. (1871–73) 319.—Abyss.
paniculatus, Benth. in Wall. Pl. As. Rar. ii. 16 = Plectranthus coleoides.
parviflorus, *Benth. in DC. Prod.* xii. 72.—Ind. or.
Persoonii, *Benth. Lab. Gen. et Sp.* 55.—Madag.
Petersianus, *Vatke, in Linnaea*, xl. (1876) 180.—Afr. trop.
puberulus, *Miq. Fl. Ind. Bat.* ii. 955.—Malaya.
pumilius, Blanco, Fl. Filip. ed. I. 482 = acuminatus.
remotiflorus, *Miq. Fl. Ind. Bat.* ii. 954.—Java.
rivularis, *Vatke, in Linnaea*, xxxvii. (1871–73) 320.—Abyss.

COLEUS:—
rugosus, Benth. in Wall. Pl. As. Rar. ii. 15 = Plectranthus rotundifolius.
Schimperi, Vatke, in Linnaea, xxxvii. (1871–73) 320 = barbatus.
Schweinfurthii, *Vatke, l. c.* 323, in textu.—Afr. trop.
scutellarioides, *Benth. in Wall. Pl. As. Rar.* ii. 16.—Malaya; Austral.
secundiflorus, *Benth. Lab. Gen. et Sp.* 55.—Ins. Timor.
spectabilis, *Miq. Fl. Ind. Bat.* ii. 955.—Java.
spicatus, *Benth. in Wall. Pl. As. Rar.* ii. 15.—Ind. or.
suborbicularis, *Zoll. & M oritzi, Syst. Verz. Zoll.* 4.—Java.
subspicatus, Walp. Rep. vi. 658 = Plectranthus spicatus.
Suganda, Blanco, Fl. Filip. ed. I. 483 = aromaticus.
tenuicaulis, *Hook. f. in Journ. Linn. Soc.* vii. (1864) 211.—Afr. trop.
tenuiflorus, *Vatke, in Linnaea*, xliii. (1880–82) 92.—Afr. trop.
Tryoni ×, *Hort. ex Flor. Mag.* N. S. i. (1872) t. 34.
tuberosus, Benth. Lab. Gen. et Sp. 59 = parviflorus.
tuberosus, *A. Rich. Tent. Fl. Abyss.* ii. 185.—Abyss.
umbrosus, *Vatke, in Linnaea*, xliii. (1880–82) 91.—Afr. trop.
urticifolius, Benth. in DC. Prod. xii. 78 = Plectranthus urticifolius.
Verschaffeltii, *Lem. Illustr. Hortic.* viii. (1861) t. 293.—Java.
Walkeri, Benth. in DC. Prod. xii. 77 = malabaricus.
Wightii, Benth. Lab. Gen. et Sp. 58 = Plectranthus coleoides.
Zatarhendi, Benth. l. c. 50 = spicatus?

COLICODENDRON, Mart. in Flora, xxii. (1839) I. Beibl. 25 = **Capparis**, Linn.
anceps, Shuttl. ex Chapm. Fl. S. U. St. 32 = Capp. jamaicensis.
avicenniaefolium, Seem. Bot. Voy. Herald, 78 = Capp. avicennifolia.
Breynia, Endl. ex Heynh. Nom. ii. 154 = Capp. Breynia.
lepidotum, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 327.—Venezuela.
longifolium, Mart. in Flora, xxii. (1839) I. Beibl. 26.—Bras.
obliquifolium, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 328.—N. Granat.
obovatum, Turcz. l. c. 327.—Venezuela.
pulchellum, Seem. Bot. Voy. Herald, 78 = Capp. pulcherrima.
scabridum, Seem. l. c. = Capp. scabrida.
subbilobum, Seem. l. c. = Capp. subbiloba.
Yco, Mart. in Flora, xxii. (1839) I. Beibl. 25 = Capp. Yco.

COLIGNONIA, Endl. Gen. 311 (1837). *NYCTAGINEAE*, Benth. & Hook. f. iii. 8.
COLLIGNONIA, Endl. Ench. 193 (1841).
biumbellata, *Ball, in Journ. Linn. Soc.* xxii. (1885) 54.—Peruv.
glomerata, *Griseb. in Goett. Abh.* xix. (1874) 87.—Reg. Argent.
parviflora, *Endl. Gen.* 311; *Choisy, in DC. Prod.* xiii. II. 439.—Peruv.
scandens, *Benth. Pl. Hartw.* 148.—Peruv.

COLIQUEA, Steud. ex Bibra, in Denkschr. Wien. Acad. v. (1853) 115 = **Chusquea**, Kunth (Gramin.).
Quila, Steud. ex Bibra, l. c. = Chus. Cumingii.

COLLABIUM, Blume, Bijdr. 357 (1825). *ORCHIDEAE*, Benth. & Hook. f. iii. 508.
nebulosum, *Blume, l. c.* 358.—Java.
simplex, *Reichb. f. in Gard. Chron.* (1881) I. 462.—Borneo.

COLLADEA, Pers. Syn. i. 107 (1805) = seq.

COLLADOA, Cav. Ic. v. 37. t. 460 (1799) = **Ischaemum**, Linn. (Gramin.).
distachia, Cav. l. c. = I. rugosum.
monostachya, Pers. Syn. i. 107 = Anthephora elegans.

COLLADONIA, Spreng. Syst. i. 516 (1825) = **Palicourea**, Aubl. (Rubiac.).
tinifolia, Spreng. l. c. 757 = P. tinifolia.

COLLADONIA, DC. Prod. iv. 240 (1830) = **Prangos**, Lindl. (Umbellif.).
alata, Boiss. Fl. Orient. ii. 945.—Syria.
anatolica, Boiss. l. c.—As. Min.
angustifolia, Bertol. Fl. Ital. iii. 408 = P. angustifolia.
anisoptera, Boiss. Fl. Orient. ii. 946.—Syria.
cilicica, Boiss. l. c. 947.—Cilicia.
crenata, Boiss. l. c. 946.—Syria.
heptaptera, Boiss. l. c. 945 = P. colladonioides.
microcarpa, Boiss. l. c. 946 = P. microcarpa.
syriaca, Boiss. in Ann. Sc. Nat. Sér. III. ii. (1844) 86 = Smyrniopsis cachroides.
triquetra, DC. Prod. iv. 240 = P. triquetra.

COLLAEA, DC. Mém. Leg. vi. 244 (1825); et in Ann. Sc. Nat. Sér. I. iv. (1825) 96 = **Galactia**, P. Br. (Legumin.).
angustifolia, Benth. in Ann. Wien. Mus. ii. (1838) 129 = G. angustifolia.
argentina, Griseb. in Goett. Abh. xix. (1874) 125.—Reg. Argent.
bullata, Benth. in Mart. Fl. Bras. xv. I. 149.—Bras.
cinerascens, R. Grah. in Wall. Cat. n. 5575 = Atylosia mollis.
crassifolia, Benth. in Ann. Wien. Mus. ii. (1838) 128.—Bras.
decumbens, Benth. in Linnaea, xxii. (1849) 515.—Bras.
diversifolia, Benth. in Mart. Fl. Bras. xv. I. 147.—Am. trop.
formosa, Griseb. in Goett. Abh. xix. (1874) 126.—Reg. Argent.
gibba, R. Grah. in Wall. Cat. n. 5572 = Dunbaria Heynei, ferruginea.
glaucescens, Benth. in Ann. Wien. Mus. ii. (1838) 128 = G. glaucescens.
grewiaefolia, Benth. l. c. 128.—Bras.
guianensis, Klotzsch, ex Schomb. Faun. & Fl. Gui. 1203.—Guiana.
lancifolia, Benth. in Mart. Fl. Bras. xv. I. 150.—Bras.
latisiliqua, Benth. in Ann. Wien. Mus. ii. (1838) 129 = G. latisiliqua.
longiflora, Benth. l. c. 128.—Ins. Trinit.
longifolia, Benth. in Mart. Fl. Bras. xv. I. 151.—Am. trop.
macrophylla, Benth. in Ann. Wien. Mus. ii. (1838) 128.—Bras.
Martii, Benth. l. c. 129 = G. Martii.
mollis, R. Grah. in Wall. Cat. n. 5574 = Atylosia mollis.
Neesii, Benth. in Ann. Wien. Mus. ii. (1838) 129 = G. Neesii.
obtusa, Benth. l. c. 128.—Bras.
pascuorum, Mart. ex Benth. l. c.—Bras.
pedicellatum, Benth. in Hook. Journ. Bot. ii. (1840) 60.—Bras.
peduncularis, Benth. in Ann. Wien. Mus. ii. (1838) 128.—Bras.
pendula, Benth. l. c. = G. pendula.
rosea, Benth. in Ann. Nat. Hist. iii. (1839) 437 = G. Jussiaeana.
rotundifolia, Benth. in Mart. Fl. Bras. xv. I. 149.—Bras.
rugosa, Benth. in Ann. Wien. Mus. ii. (1838) 128.—Bras.
scarlatina, Mart. ex Benth. l. c. 129.—Bras.
speciosa, DC. in Mém. Lég. vi. 245.—Peruv.
stenophylla, Benth. in Mart. Fl. Bras. xv. I. 146 = G. stenophylla.
trinervia, DC. Mém. Lég. vi. 247 = Atylosia Candollei.
tomentosa, Benth. in Mart. Fl. Bras. xv. I. 149 = G. tomentosa.
velutina, Benth. in Ann. Wien. Mus. ii. (1838) 129 = G. velutina.
venosa, R. Grah. in Wall. Cat. n. 5573 = Dunbaria ferruginea.
virgata, Benth. in Mart. Fl. Bras. xv. I. 146.—Bras.

COLLAEA, Endl. Gen. Suppl. ii. 20 (1842) in syn. COLLEA, Lindl. = **Pelexia**, Lindl. (Orchid.).

COLLAEA, Spreng. Syst. iii. 622 (1826) = **Chrysanthellum**, Rich. (Compos.).
procumbens, Spreng. l. c. = Chrys. procumbens.

COLLANDRA, Lem. in Fl. des Serres, iii. (1847) sub t. 223 = **Columnea**, Linn. (Gesnerac.).
atronitens, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1402 = Columnea aureo-nitens.
aureo-nitens, Hanst. in Linnaea, xxvi. (1853) 209 = Columnea aureo-nitens.
petiolaris, Griseb. Fl. Brit. W. Ind. 463.—Ind. occ.
phoenicea, G. Don, ex Loud. Encyc. Pl. Suppl. ii. 1402.—N. Granat.
picta, Lem. Jard. Fleur. ii. (1852) t. 214 = Columnea picta.
pilosa, Lem. in Fl. des Serres, iii. (1847) sub t. 223 = Columnea aureo-nitens.
sanguinea, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 526 = Columnea sanguinea.

COLLANIA, Herb. Amaryll. 103. t. 8 (1837) = **Bomarea**, Mirb. (Amaryllid.).
andimarcana, Herb. l. c. 105 = B. andimarcana.
dulcis, Herb. l. c. 104 = B. glaucescens.
glaucescens, Herb. l. c. = B. glaucescens.
involucrosa, Herb. l. c. 103. t. 9 = B. glaucescens.
puberula, Herb. l. c. 105. t. 11. f. 1 = B. glaucescens.

COLLANIA, Schult. f. Syst. vii. II. p. liii, 893 (1830) = **Urceolina**, Reichb. (Amaryllid.).
dubia, Schult. f. l. c.—Ecuador.
fulva, M. Roem. Syn. Ensat. 55 = U. fulva.
urceolata, Schult. f. Syst. vii. 893 = U. pendula.

COLLEA, Lindl. Bot. Reg. sub t. 760 (1823) = **Pelexia**, Lindl. (Orchid.).
adnata, Lindl. l. c. = P. spiranthoides.
calcarata, Lindl. l. c. = P. setacea.

COLLEMA, Anders. ex DC. Prod. vii. 512 (1839) = **Goodenia**, Sm.

COLLETIA, Comm. ex Juss. Gen. 380 (1789). *RHAMNEAE*, Benth. & Hook. f. i. 383.
SCYPHARIA, Miers, in Ann. & Mag. Nat. Hist. Ser. III. vi. (1860) 8.
aciculata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 214.—Peruv.
armata, Miers, l. c. 211 = spinosa.
articulata, *Phil. in Linnaea*, xxviii. (1856) 679.—Chili.
atrox, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 206.—Reg. Argent.
bictonensis, Lindl. in Journ. Hort. Soc. v. (1850) 31 = cruciata.
brevispina, *Phil. in Anal. Univ. Chil.* (1872) 690.—Chili.
cataphracta, *Miers, in Ann. & Mag. Mat. Hist.* Ser. III. v. (1860) 210.—Chili.
Chacaye, G. Don, Gen. Syst. ii. 35 = Discaria serratifolia.
crenata, *Clos, in C. Gay, Fl. Chil.* ii. 35.—Chili.
crenata, Regel, Ind. Sem. Hort. Petrop. (1856) 37 = Discaria crenata.
cruciata, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 152. t. 43.—Chili.
Cruzerillo, *Bert. in 'Merc. Chil. 1829'.—Cf. Bull. Férussac*, xx. (1830) 109, nomen.—Chili.
Cunninghami, Fenzl, in Enum. Pl. Hueg. 23 = Discaria australis.
discolor, *Hook. Ic. Pl.* t. 538.—Patagon.
disperma, *Moç. & Sesse, ex DC. Prod.* ii. 29.—Mexic.
Doniana, *Clos, in C. Gay, Fl. Chil.* ii. 36.—Chili.
dumosa, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 212.—Chili.
Ephedra, Bert. ex Steud. Nom. ed. II. i. 397 = Trevoa Trutilla.
Ephedra, *Vent. Choix*, 16. t. 16.—Peruv.
exserta, *Klotzsch, ex Reiss. in Mart. Fl. Bras.* xi. 1. 100.—Bras.
ferox, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 154.—Chili.
horrida, Brongn. ex Drap. Encyc. & Fl. des Serres Angl. iii. (Juill. 1835) f. 6 = ferox.
horrida, Willd. Sp. Pl. i. 1113 = spinosa.
Hystrix, *Clos, in C. Gay, Fl. Chil.* ii. 32.—Chili.
infesta, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 366 = Adolphia infesta.
insidiosa, *Reiss. in Mart. Fl. Bras.* xi. 1. 100.—Bras.
intricata, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 204.—Chili.
invicta, *Miers, l. c.* 205.—Chili.
Kunthiana, *Miers, l. c.* 208.—Peruv.

COLLETIA :—
longispina, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 173.—Chili.
longissima, Steud. Nom. ed. II. i. 397 = praec.
maytenoides, *Griseb. in Goett. Abh.* vi. (1854) 117.—Reg. Argent.
montana, *Phil. in Linnaea*, xxxiii. (1864-65) 37.—Chili.
multiflora, Moç. & Sesse, ex DC. Prod. ii. 29 = Celastrus aphyllus.
nana, *Clos, in C. Gay, Fl. Chil.* ii. 37.—Chili.
obcordata, *Vent. Jard. Cels*, t. 92.—Peruv.
polyacantha, Humb. & Bonpl. ex Roem. & Schult. Syst. v. 513 = obcordata, spinosa.
pubescens, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 366 = Discaria australis.
pungens, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 209.—Chili.
serratifolia, Vent. Choix, 15. t. 15 = Discaria serratifolia.
spartioides, *Bert. ex Colla, in Mem. Acc. Torin.* xxxvii. (1834) 52.—Ins. J. Fernand.
spicata, *Humb. & Bonpl. ex Roem. & Schult. Syst.* v. 513.—Am. austr.
spinosa, *Lam. Tabl. Encyc.* ii. 91. t. 129.—Am. austr.
spinosissima, J. F. Gmel. Syst. 408 = praec.
stipellacea, *Phil. in Linnaea*, xxxiii. (1864-65) 36.—Chili.
tenuicola, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 209.—Reg. Argent.?
tetragona, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 366.—Peruv.
tetrandra, *Clos, in C. Gay, Fl. Chil.* ii. 31.—Chili.
tomentosa, *Phil. in Linnaea*, xxviii. (1856) 618.—Chili.
Tralhuen, Bert. ex Colla, in Mem. Acc. Torin. xxxvii. (1834) 53 = Trevoa quinquenervia.
Treba, Bert. ex Colla, l. c. = Trevoa trinervia.
ulicina, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 155.—Chili.
valdiviana, *Phil. in Linnaea*, xxxiii. (1864-65) 35.—Chili.
Valenzuela, Bert. ex Steud. Nom. ed. II. i. 397 = ulicina.
velutina, *Spreng. Syst.* i. 771.—Bras.
veprecula, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. v. (1860) 213.—Chili.
Weddelliana, *Miers, l. c.* 207.—Bolivia.

COLLETIA, Scop. Introd. 207 (1777) = **Celtis**, Linn. (Urticac.).

COLLIGNONIA, Endl. Ench. 193 (1841) = **Colignonia**, Endl. (Nyctag.).

COLLIGUAJA, Molina, Sagg. Chil. 158, 354 (1782). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 338.
brasiliensis, *Klotzsch, ex Baill. Etud. Gén. Euph.* 535.—Bras.
Bridgesii, *Muell. Arg. in Linnaea*, xxxii. (1863) 125.—Chili.
Dombeyana, C. Gay, Fl. Chil. 342 = salicifolia.
Dombeyana, *A. Juss. in Ann. Sc. Nat.* Sér. I. xxv. (1832) 23.—Chili.
integerrima, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 140.—Chili.
obtusa, *Regel, Ind. Sem. Hort. Petrop.* (1855) 18.—Chili.
odorifera, *Molina, Sagg. Chil.* 158, 354.—Chili.
salicifolia, *Gill. & Hook. in Hook. Bot. Misc.* i. (1830) 141.—Chili.
triquetra, *Gill. & Hook. l. c.* 142.—Chili.

COLLINARIA, Ehrh. Beitr. iv. (1780) 147 = **Koeleria**, Pers. (Gramin.).

COLLINIA, Liebm. in Oefers. Danske Vid. Selsk. 1845 (1846) 8 = **Chamaedorea**, Willd. (Palm.).
elatior, Liebm. l. c. = Cham. Liebmanni.
elegans, Liebm. l. c. = Cham. elegans.
fibrosa, Oerst. in Kjoeb. Vidensk. Meddel. (1858) 5 = Synechanthus fibrosus.
humilis, Liebm. in Oefers. Danske Vid. Selsk. 1845 (1846) 8 = Cham. humilis.

COLLINIA, & COLLINIANA, Rafin. in Am. Monthly Mag. (1819) 194 = seq.

COLLINSIA, Nutt. in Journ. Acad. Philad. i. (1817) 190. t. 9. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 941.
barbata, Bosse, in Verh. Berl. Gartenb. Ver. N. Reihe, i. (1853) 399 = tinctoria.
bartsiaefolia, *Benth. in DC. Prod.* x. 318.—Calif.
bicolor, *Benth. in Trans. Hort. Soc.* N. S. i. (1831-35) 480.—Calif.
Childii, *Parry, ex A. Gray, Syn. Fl. N. Am.* ii. 1. 257.—Calif.
corymbosa, *Herder, Ind. Sem. Hort. Petrop.* (1857) 32; *Regel, Gartenfl.* (1868) 33. t. 568.—Calif.
divaricata, *Kellogg, in Proc. Calif. Acad.* iii. (1863) 36.—Calif.
grandiflora, *Dougl. ex Lindl. Bot. Reg.* sub t. 1166 et t. 1107.—Am. bor. occ.
Greenii, *A. Gray, in Proc. Am. Acad.* x. (1874) 75.—Calif.
heterophylla, R. Grah. in Hook. Bot. Mag. t. 3695 = bicolor.
hirsuta, Kellogg, in Proc. Calif. Acad. ii. (1863) 110. t. 33 = bartsiaefolia.
linearis, *A. Gray, in Proc. Am. Acad.* xv. (1880) 50.—Am. bor. occ.
minima, Nutt. in Journ. Acad. Philad. vii. (1834) 47 = parviflora.
multicolor, *Lindl. & Paxt. Flow. Gard.* ii. (1851-52) 89. t. 55.—Calif.
Parryi, *A. Gray, Syn. Fl. N. Am.* ii. 1. 257.—Calif.
parviflora, *Lindl. Bot. Reg.* t. 1082.—Am. bor.
pauciflora, Hook. Fl. Bor. Am. ii. 94, sphalm. = parviflora.
purpurea, *Rafin. Atl. Journ.* 149.—Am. bor.
Rattani, *A. Gray, in Proc. Am. Acad.* xv. (1880) 50.—Am. bor. occ.
septemnervia, Kellogg, in Proc. Calif. Acad. ii. (1863) 224 = tinctoria.
solitaria, Kellogg, l. c. 10 = sparsiflora.
sparsiflora, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. (1836) 33.—Calif.
tenella, Benth. in DC. Prod. x. 593 = Tonella collinsioides.
tinctoria, *Hartw. ex Benth. Pl. Hartw.* 328.—Calif.
Torreyi, *A. Gray, in Proc. Am. Acad.* vii. (1868) 378.—Calif.
tricolor, Rafin. Cat. 13 = verna.
verna, *Nutt. in Journ. Acad. Philad.* i. (1817) 190. t. 9.—Am. bor.
violacea, *Nutt. in Trans. Am. Phil. Soc.* v. (1837) 179.—Am. bor.

COLLINSONIA, Linn. Gen. ed. I. 334 (1737). *LABIATAE*, Benth. & Hook. f. ii. 1181.
DIALLOSTEIRA, Rafin. Neogenyt. 2 (1825).
HYPOGON, Rafin. Fl. Ludov. 148 (1817).
PLEURADENIA, Rafin. Neogenyt. 2 (1825).
anisata, *Sims, Bot. Mag.* t. 1213.—Am. bor.
canadensis, *Linn. Sp. Pl.* 28.—Am. bor.
cuneata, Wender. in Schr. Nat. Ges. Marb.? ex Benth. in DC. Prod. xii. 253 = canadensis.
decussata, Moench, Meth. 379 = canadensis.
ovalis, Pursh, Fl. Am. Sept. i. 21 = scabriuscula.
praecox, Walt. Fl. Carol. 65 = scabriuscula.
punctata, Ell. Sketch, i. 36 = canadensis.
scabra, Pers. Syn. i. 29 = scabriuscula.
scabriuscula, *Ait. Hort. Kew.* ed. I. i. 47.—Am. bor.
serotina, Walt. Fl. Carol. 65 = canadensis.
tuberosa, Michx. Fl. Bor. Am. i. 17 = scabriuscula.
urticifolia, Salisb. Prod. 75 = canadensis.
verticillaris, Rafin. Fl. Ludov. 41 = seq.?
verticillata, *Baldw. ex Ell. Sketch*, i. 36.—Am. bor.

COLLOCOCCUS, P. Br. Hist. Jamaic. 167 (1756) (Gen. dub.).

COLLOMIA, Nutt. Gen. Am. i. 126 (1818). *POLEMONIACEAE*, Benth. & Hook. f. ii. 822.
COURTOISIA, Reichb. Ic. Exot. iii. 4. t. 208 (1827).
aggregata, T. C. Porter, ex A. Gray, Syn. Fl. N. Am. ii. 1. 394 = Gilia aggregata.
Cavanillesiana, *G. Don, Gen. Syst.* iv. 247.—Mexic.; Texas.
Cavanillesii, Hook. & Arn. Bot. Beech. Voy. 34 = coccinea.
coccinea, *Lehm. ex Benth. in Bot. Reg.* t. 1622.—Chili.
eritrichoides, *Griseb. in Goett. Abh.* vi. (1853-55) 130.—Chili.
erythraeoides, *Griseb. l. c.*—Chili.

COLLOMIA:—
gilioides, *Benth. in Bot. Reg.* sub t. 1622.—Calif.
glutinosa, Benth. l. c. = gilioides.
gracilis, *Dougl. ex Benth. l. c.*—Am. bor. occ.
grandiflora, *Dougl. in Lindl. Bot. Reg.* t. 1174.—Am. bor. occ.
heterophylla, *Hook. Bot. Mag.* t. 2895.—Am. bor. occ.
lateritia, D. Don, in Sweet, Brit. Flow. Gard. Ser. II. iii. t. 206 = coccinea.
leptalea, *A. Gray, in Proc. Am. Acad.* viii. (1870) 261.—Calif.
linearis, *Nutt. Gen. Am.* i. 126.—Am. bor. occ.
linoides, Nutt. in Journ. Acad. Philad. N. S. i. (1847) 159 = Gilia minutiflora.
longiflora, *A. Gray, in Proc. Am. Acad.* viii. (1870) 261.—Am. bor. occ.
micrantha, Kellogg, in Proc. Calif. Acad. iii. (1863) 18. f. 3 = gracilis.
myotica, *Griseb. in Goett. Abh.* vi. (1853–55) 129.—Chili.
Navarretia, G. Don, Gen. Syst. iv. 247 = Gilia Navarettia.
nudicaulis, *Hook. & Arn. Bot. Beech. Voy.* 368.—Am. bor. occ.
stenosiphon, *Kunze, in Linnaea*, xxiv. (1851) 228.—Chili.
tenella, *A. Gray, in Proc. Am. Acad.* viii. (1870) 259.—Am. bor. occ.
Thurberi, *A. Gray, l. c.* 261.—N. Mexic.
tinctoria, Kellogg, in Proc. Calif. Acad. iii. (1863) 17. f. 2 = lincaris.

COLLOMIA, Sieber, ex Steud. Nom. ed. II. i. 398 (1840) = **Felicia**, Cass. (Compos.).
setosa, Sieber, ex Steud. l. c. = F. reflexa.

COLLOPHORA, Mart. in Buchn. Rep. Pharm. (1828).—Cf. Bull. Férussac (Jan. 1831) 64 = **Couma**, Aubl. (Apocyn.).
utilis, Mart. l. c. = Couma utilis.

COLLYRIS, Vahl, in Skrift. Nat. Selsk. Kiobenh. vi. (1810) 110 = **Dischidia**, R. Br. (Asclepiad.).
major, Vahl, l. c. = D. Rafflesiana, D. Collyris.
minor, Vahl, l. c. 111 = D. nummularia.

COLMEIROA, F. Muell. Fragm. vii. 149 (1871). *SAXIFRAGEAE.*
carpodetoides, *F. Muell. l. c.*—Ins. D. Howe.

COLMEIROA, Reut. in Mém. Soc. Phys. Genèv. x. (1843) 240. t. 6 = **Securinega**, Comm. (Euphorb.).
buxifolia, Reut. l. c. = S. buxifolia.

COLOBACHNE, Beauv. Agrost. 22. t. 6. f. 9 (1812) = **Alopecurus**, Linn. (Gramin.).
brachystachys, Nym. in Bot. Notiser (1851) 69 = A. brachystachys.
Gerardi, Link, Hort. Berol. i. 74 = Phleum Gerardi.
pontica, Nym. in Bot. Notiser (1851) 69 = A. glacialis.
vaginata, Beauv. Agrost. 22 = A. vaginatus.

COLOBANDRA, Bartl. in Lehm. Pl. Preiss. i. 357 (1845) = **Hemigenia**, R. Br. (Labiat.).
canescens, Bartl. l. c. 358 = H. canescens.
lanata, Bartl. l. c. 359 = H. canescens?
mollis, Bartl. l. c. 358 = H. canescens.
platyphylla, Bartl. l. c. = H. platyphylla.
robusta, Bartl. l. c. 357 = H. polystachya.
subvillosa, Bartl. l. c. 359 = H. platyphylla?

COLOBANTHUS, Bartl. in Presl, Reliq. Haenk. ii. 13. t. 49 (1830). *CARYOPHYLLEAE*, Benth. & Hook. f. i. 151.
acicularis, *Hook. f. Handb. New Zeal. Fl.* 25.—N. Zel.
affinis, Hook. f. in Hook. Journ. Bot. ii. (1840) 410 = Billardieri.
arctioides, *Gill. ex Hook. Bot. Misc.* iii. (1833) 336.—Chili.
Benthamianus, Fenzl, in Ann. Wien. Mus. i. (1836) 49 = subulatus.
Billardieri, *Fenzl, l. c.*—Austral.
cherlerioides, *Hook. f. Fl. Antarct.* 249.—Patagon.
crassifolius, *Hook. f. l. c.* 248.—Chili; Patagon.
diffusus, *Hook. f. l. c.* 249.—Ins. Amsterdam.
kerguelensis, *Hook. f. l. c.*—Ins. Kerguel.
Lechleri, *Phil. in Linnaea*, xxviii. (1856) 689.—Chili.

COLOBANTHUS:—
lycopodioides, *Griseb. in Goett. Abh.* vi. (1853–55) 116.—Reg. Magellan.
muscoides, *Hook. f. Fl. Antarct.* 14.—Ins. Antarct.
polycnemoides, *Hieron. in Bol. Acad. Nac. Córdova*, iii. (1879) 334.—Reg. Argent.
pulvinatus, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 101 = subulatus.
quitensis, *Bartl. in Presl, Reliq. Haenk.* ii. 13. t. 49. f. 2.—Ecuador.
saginoides, *Bartl. l. c.*—Chili.
subulatus, *Hook. f. Fl. Antarct.* 13.—Ins. Antarct.

COLOBANTHUS, Trin. in Mém. Acad. Pétersb. Sér. VI. i. (1831) 66 = **Avellinia**, Parl. (Gramin.).

COLOBATUS, Walp. in Linnaea, xiii. (1839) 478 = COLOBOTUS, E. Mey. = **Buchenroedera**, Eckl. & Zeyh. (Legumin.).

COLOBIUM, Roth, in Roem. Archiv, i. (1796) t. 36 = **Leontodon**, Linn. (Compos.).
hirtum, Roth, l. c. = L. hirtum.
hispidum, Roth, l. c. 37 = L. Rothii.

COLOBOGYNIUM, Schott, in Oestr. Bot. Zeitschr. xv. (1865) 34 = **Schismatoglottis**, Zoll. (Aroid.).
tecturatum, Schott, l. c. = S. variegata.

COLOBOTUS, E. Mey. Comm. Pl. Afr. Austr. 156 (1835) = **Buchenroedera**, Eckl. & Zeyh. (Legumin.).
ochreatus, E. Mey. l. c.—Afr. austr.

COLOCASIA, Link, Diss. Bot. Suerin (1795) 77; et Handb. i. 267 = **Richardia**, Kunth (Aroid.).
aethiopica, Link, ll. cc. = R. aethiopica.

COLOCASIA, Neck. Elem. iii. 289 (1790) (Gen. dub. Aroidearum).

COLOCASIA, Schott, Meletem. i. 18 (1832). *AROIDEAE*, Benth. & Hook. f. iii. 974.
LEUCOCASIA, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 34.
acris, Schott, Meletem. i. 18 = antiquorum.
affinis, *Schott, in Bonplandia*, vii. (1859) 28.—Reg. Himal.
alba, Hort. ex Engl. in DC. Monog. Phan. ii. 501 = Alocasia alba.
argyroneura, Hort.—Cf. Gard. Chron. (1875) i. 589 = Caladium Schomburgkii.
antiquorum, *Schott, Meletem.* i. 18.—As. trop.
Boryi, Kunth, Enum. Pl. iii. 41 = Alocasia indica.
cochleata, Miq. Epim. Sem. Hort. Amstelod. (1853) = Aloeasia cucullata.
cucullata, Schott, Meletem. i. 18 = Alocasia cucullata.
esculenta, Schott, l. c. = antiquorum.
euchlora, C. Koch & Linden, in App. Sem. Hort. Bot. Berol. (1854) 4 = antiquorum.
fallax, *Schott, in Bonplandia*, vii. (1859) 28.—Ind. or.
Fontanesii, Schott, in Oestr. Bot. Wochenbl. iv. (1854) 409 = antiquorum.
fornicata, Kunth, Enum. Pl. iii. 41 = Alocasia fornicata.
gracilis, *Engl. Bot. Jahrb.* i. (1881) 185.—Sumatra.
heterophylla, *Kunth, Enum. Pl.* iii. 40.—Ins. Philipp.
humilis, *Hassk. Cat. Hort. Bogor. Alt.* 56.—Java.
indica, *Hassk. in Hoev. & De Vriese, Tijdschr.* ix. (1842) 160.—Java.
indica, Kunth, Enum. Pl. iii. 39 = Alocasia indica.
macrorrhiza, Schott, Meletem. i. 18 = Alocasia macrorhiza.
Marchalii, *Engl. in DC. Monog. Phan.* ii. 494.—Hab.?
montana, Kunth, Enum. Pl. iii. 40 = Aloeasia montana.
mucronata, Kunth, l. c. = Alocasia macrorhiza.
navicularis, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1853) 13 = Alocasia navicularis.
neo-guineensis, *André, in Illustr. Hortic.* xxvii. (1880) 68.—N. Guinea.
nitida, Walp. Ann. i. 1041, sphalm. = Coprosma nitida.
nymphaeifolia, Kunth, Enum. Pl. iii. 37 = antiquorum.
obtusiloba, *Kunth, l. c.* 41—Hab.?
odora, Brongn. in Nouv. Ann. Mus. Par. iii. (1834) 145. t. 7 = Alocasia odora.
peregrina, *Rafin. Fl. Tellur.* iii. 65.—Hab.?
prunipes, C. Koch & Bouché, Ind. Sem. Hort. Berol. (1854) 4 = indica.
pubera, Hassk. Cat. Hort. Bogor. Alt. 302 = Alocasia pubera.

COLOCASIA:—
pumila, Kunth, Enum. Pl. iii. 40 = Gonatanthus sarmentosus.
ramosa, Schott, ex Hcynh. Nom. ii. 154 = Arum ramosum.
rapiformis, *Kunth, Enum. Pl.* iii. 40.—Burma.
rugosa, Kunth, l. c. 41 = Alocasia cucullata.
virosa, *Kunth, l. c.*—Ind. or.
vivipara, *Thw. Enum. Pl. Zeyl.* 336.—Zeylan.
vulgaris, Rafin. Fl. Tellur. iii. 65 = antiquorum.

COLOCOCCA, Rafin. Sylva Tellur. 40 (1838) = **Cordia**, Linn. (Boragin.).
macrophyla, Rafin. l. c. = Cordia macrophylla.

COLOCYNTHIS, [Tourn.] Linn. Syst. ed. I (1735); Schrad. Ind. Sem. Hort. Gotting. (1833) 2; Linnaea, x. (1836) Lit. 70 = **Citrullus**, Neck. (Cucurbit.).
amarissima, Schrad. ll. cc. = Cit. vulgaris.
officinalis, Schrad. in Linnaea, xii. (1838) 421 = Cit. Colocynthis.
officinarum, Cogn. in DC. Monog. Phan. iii. 511, sphalm. = praec.
vulgaris, Schrad. Ind. Sem. Hort. Gotting. (1833) 2 = Cit. Colocynthis.

COLOGANIA, Kunth, Mim. 205. t. 57 (1823). *LEGUMINOSAE*, Benth. & Hook. f. i. 529.
affinis, *Mart. & Gal. in Bull. Acad. Brux.* x. (1843) II. 188.—Mexic.
angustifolia, *Kunth, Mim.* 209. t. 58.—Am. bor. occ.
australis, *Griseb. in Goett. Abh.* xix. (1874) 124.—Reg. Argent.
biloba, *Nichols. Dict. Gard.* i. 363.—Mexic.
Bronssonetii, *DC. Prod.* ii. 237.—Mexic.
heterophylla, Gill. ex Hook. Bot. Misc. iii. (1833) 181 = Galactia marginalis.
humifusa, *Hemsl. Diagn. Pl. Nov.* 47.—Mexic.
intermedia, *H. B. & K. Nov. Gen. et Sp.* vi. 414.—Mexic.
Lemmoni, *A. Gray, in Proc. Am. Acad.* xix. (1883) 74.—Arizona.
longifolia, *A. Gray, Pl. Wright.* ii. 35.—Am. bor. occ.
Martia, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 345.—Mexic.
obovata, *Schlecht. in Linnaea*, xii. (1838) 257.—Mexic.
ovalifolia, H. B. & K. Nov. Gen. et Sp. vi. 412 = pulchella.
procumbens, *Kunth, Mim.* 205. t. 57.—N. Granat.
pulchella, *H. B. & K. Nov. Gen. et Sp.* vi. 413.—Am. bor. occ.

COLOGYNE, Griff. Notul. iii. 280 (1851) err. typ. = **Coelogyne**, Lindl. (Orchid.).

COLOMANDRA, Neck. Elem. ii. 142 (1790) = **Aiouea**, Aubl. (Laurin.).

COLONA, Cav. Ic. iv. 47. t. 370 (1797) = **Columbia**, Pers. (Tiliac.).
serratifolia, Cav. l. c. = Columbia americana.

COLONNA, Jaume St. Hil. Expos. ii. 95 (1805) = praec.

COLONNEA, Endl. Gen. 421, in syn. (1838) = CALONNEA, Buchoz = **Gaillardia**, Foug. (Compos.).

COLOPHONIA, Comm. ex Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 352 = **Canarium**, Linn. (Burserac.).
mauritiana, DC. Prod. ii. 79 = Canarium mauritianum.
tetramera, Boiv. ex Engl. in DC. Monog. Phan. iv. 27 = Commiphora tetramera.

COLPIAS, E. Mey. ex Benth. in Hook. Comp. Bot. Mag. ii. (1836) 53. *SCROPHULARINEAE*, Benth. & Hook. f. ii. 931.
mollis, *E. Mey. l. c.*—Afr. austr.

COLPODIUM, Trin. Fund. Agrost. 119 (1820). *GRAMINEAE*, Benth. & Hook. f. iii. 1196.
altaicum, *Trin. in Ledeb. Fl. Alt.* i. 100.—Sibir. alt
aquaticum, Trin. Fund. Agrost. 136 = Catabrosa aquatica.
arundinaceum, *Hook. Fl. Bor. Am.* ii 238.—Am. bor.
bulbosum, Trin. in Mem. Acad. Petersb. Ser. VI. i. (1831) 394 = Catabrosa humilis.
Calvertii, Boiss. Diagn. Ser. II. iv. 133 = Catabrosa humilis.

COLPODIUM:—
compressum, Trin. in Spreng. Neue Entdeck. ii. 58 = Sporobolus compressus.
fibrosum, Trautv. in Act. Hort. Petrop. ii. (1873) 486 = Catabrosa fibrosa.
filifolium, Trin. in Bull. Sc. Acad. Pétersb. i. (1836) 69 = Catabrosa humilis.
fulvum, Griseb. in Ledeb. Fl. Ross. iv. 385 = Poa fulva.
humile, Griseb. l. c. 384 = Catabrosa humilis.
junceum, Trin. in Spreng. Neue Entdeck. ii. 37 = Sporobolus junceus.
latifolium, *R. Br. in Parry*, 1*st Voy. Suppl.* p. clxxxvi. —Reg. arct.
Malmgrenii, *Anderss. in Ofvers. Vet. Akad. Stockh.* xxiii. (1866) 121.—Ins. Spitzberg.
minutum, *Steud. Syn. Pl. Gram.* 177.—Hispan.
monandrum, Trin. Fund. Agrost. 119 = Phippsia algida.
parviflorum, Boiss. & Buhse, Diagn. Ser. II. iv. 133 = Catabrosa parviflora.
pauciflorum, *Hook. Fl. Bor. Am.* ii. 238.—Am. bor.
pendulinum, Griseb. in Ledeb. Fl. Ross. iv. 386 = Poa Laestadii.
pusillum, *Nees, Fl. Afr. Austr.* 149.—Afr. austr.
Steveni, Hohen. in Bull. Soc. Nat. Mosc. vi. (1833) 213 = Catabrosa humilis.
Steveni, *Trin. Fund. Agrost.* 119. t. 7.—Reg. Caucas.
Tilesii, *Griseb. in Ledeb. Fl. Ross.* iv. 385.—Sibir. or.
variegatum, *Boiss. ex Griseb. l. c.* 384, in textu.—As. Min.
violaceum, *Griseb. l. c.*—Persia.

COLPOON, Berg. Desc. Pl. Cap. 38. t. 1 (1767).
SANTALACEAE, Benth. & Hook. f. iii. 225.
Fusanus, Murr. Syst. ed. XIII. 765 (1774).
Hamiltonia, Harv. Gen. Pl. Cap. ed. I. 298 (1838).
Rhoicarpos, A. DC. Prod. xiv. 634 (1857).
compressum, *Berg. Desc. Pl. Cap.* 38—Afr. austr.

COLPOTHRINAX, Griseb. & H. Wendl. in Bot. Zeit. xxxvii. (1879) 147. *PALMAE*, Benth. & Hook. f. iii. 927.
Wrightii, *H. Wendl. in Kerch. Palm.* 241.—Cuba.

COLQUHOUNIA, Wall. in Trans. Linn. Soc. xiii. (1822) 608. *LABIATAE*, Benth. & Hook. f. ii. 1208.
coccinea, *Wall. l. c.* 608.—Reg. Himal.
elegans, *Wall. Pl. As. Rar.* i. 65.—Burma.
mollis, *Schlecht. in Linnaea*, xxiv. (1851) 681.—Reg. Himal.?
tenuiflora, *Hook. f. Fl. Brit. Ind.* iv. 674.—Burma.
tomentosa, Houll. in Rev. Hortic. (1873) 131 = coccinea.
vestita, *Wall. Tent. Fl. Nep.* i. 14.—Reg. Himal.

COLSMANNIA, Lehm. in Ges. Nat. Fr. Berl. Mag. viii. (1814) 92. t. 4 = **Onosma**, Linn. (Boragin.).
flava, Lehm. l. c. = O. flavum.

COLUBRINA, Rich. ex Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 368. t. 15. f. 3. *RHAMNEAE*, Benth. & Hook. f. i. 379.
Diplisca, Rafin. Sylva Tellur. 31 (1838).
Marcorella, Neck. Elem. iii. 122 (1790).
Tubanthera, Comm. ex DC. Prod. ii. 30 (1825).
acuminata, *Griseb. Cat. Pl. Cub.* 34.—Cuba.
Alamani, *G. Don, Gen. Syst.* ii. 37.—Mexic.
americana, Nutt. Sylva, ii. 58 = ferruginosa.
asiatica, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 369.—As. et Afr. trop.
buxifolia, Schlecht. in Linnaea, xv. (1841) 469 = Ceanothus buxifolius.
celtidifolia, *Schlecht. l. c.* 471.—Mexic.
cordifolia, *Reiss. in Mart. Fl. Bras.* xi. 1. 98.—Bras.
cubensis, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 369 = Ceanothus cubensis.
Ehrenbergii, *Schlecht. in Linnaea*, xv. (1841) 469.—Mexic.
excelsa, Fenzl, in Enum. Pl. Hueg. 20 = Alphitonia excelsa.
fermenta, *Rich. ex Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 369.—Guiana.
ferruginosa, *Brongn. l. c.*—Am. bor.
glomerata, *Hemsl. Biol. Centr. Am. Bot.* i. 200.—Mexic.
granulosa, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 369.—Mexic.
Greggii, *S. Wats. in Proc. Am. Acad.* xvii. (1881-82) 336.—Am. bor. occ.
guineensis, G. Don, Gen. Syst. ii. 37 = Chailletia toxicaria?

COLUBRINA:—
infesta, Schlecht. in Linnaea, xv. (1841) 468 = Adolphia infesta.
javanica, Miq. Fl. Ind. Bat. i. 1. 648 = asiatica.
Leschenaultii, G. Don, Gen. Syst. ii. 37 = asiatica?
macrocarpa, G. Don, l. c. 36 = Ceanothus macrocarpus.
Mociniana, G. Don, l. c. = Ceanothus Mocinianus.
mystacina, G. Don, l. c. 37 = Helinus scandens.
nepaulensis, G. Don, l. c. 36 = Rhamnus nepaulensis.
oppositifolia, *Brongn. ex H. Mann, in Proc. Am. Acad.* vii. (1867) 161.—Ins. Sandvic.
pubescens, G. Don, Gen. Syst. ii. 36 = Ceanothus pubescens.
pubescens, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) ii. 301.—Burma.
pubiflora, G. Don, Gen. Syst. ii. 37 = Zizyphus pubiflorus.
reclinata, Brongn. in Ann. Sc. Nat. Sér. I. x. (1827) 369 = Ceanothus reclinatus.
rufa, *Reiss. in Mart. Fl. Bras.* xi. 1. 98.—Bras.
texensis, *A. Gray, in Bost. Journ. Nat. Hist.* vi. (1850) 169.—Texas.
travancorica, *Bedd. Ic. Pl. Ind. Or.* i. 43.—Ind. or.
triflora, *Brongn. in Ann. Sc. Nat.* Sér. I. x. (1827) 369.—Mexic.
triquetra, G. Don, Gen. Syst. ii. 37 = Rhamnus triquetra.
vitiensis, Seem. Syst. List Vit. Pl. 4 = Rhamnus vitiensis.

COLUMBARIA, J. & C. Presl, Fl. Čech. 32 (1819) = **Scabiosa**, Tourn. (Dipsac.).
canescens, J. & C. Presl, l. c. = S. suaveolens.
crenata, Presl, ex Steud. Nom. ed. II. i. 398 = S. crenata.
glabrescens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = S. lucida.
gramuntea, Fourr. l. c. = S. gramuntea.
minor, S. F. Gray, Nat. Arr. Brit. Pl. ii. 476 = S. Columbaria.
norica, J. & C. Presl, Fl. Čech. 32 = S. lucida.
ochroleuca, J. & C. Presl, l. c. = S. ochroleuca.
patens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = S. Columbaria.
pratensis, Fourr. l. c. = Knautia arvensis.
rubella, Opiz, ex Steud. Nom. ed. II. i. 398 = S. Columbaria.
suaveolens, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 401 = S. suaveolens.
tenuisecta, Fourr. l. c. = S. gramuntia.
vulgaris, J. & C. Presl, Fl. Čech. 32 = S. Columbaria.

COLUMBEA, Salisb. in Trans. Linn. Soc. viii. (1807) 317 = **Araucaria**, Juss. (Conifer.).
angustifolia, Bertol. in Bologn. Opusc. Sc. iii. (1819) 411 = excelsa.
angustifolia, Raddi, ex Steud. Nom. ed. II. i. 398 = A. brasiliana.
Bidwilli, Carr. Conif. ed. II. 601 = A. Bidwillii.
brasiliensis, Carr. l. c. 596 = A. brasiliana.
excelsa, Spreng. Syst. iv. Cur. Post. 215 = A. excelsa.
imbricata, Carr. Conif. ed. II. 598 = A. imbricata.
quadrifaria, Salisb. in Trans. Linn. Soc. viii. (1807) 317 = A. imbricata.

COLUMBIA, Pers. Syn. ii. 66 (1807). *TILIACEAE*, Benth. & Hook. f. i. 233.
Colona, Cav. Ic. iv. 47. t. 370 (1797).
americana, *Pers. Syn.* ii. 66.—Ins. Philipp.
Anilao, *Blanco, Fl. Filip.* ed. I. 654.—Ins. Philipp.
Blancoi, *Rolfe, in Journ. Linn. Soc.* xxi. (1884) 308. —Ins. Philipp.
celebica, *Blume, Bijdr.* 117.—Ins. Celebes.
floribunda, Naves, in Blanco, Fl. Philipp. ed. III. t. 312 = Blancoi.
floribunda, *Kurz, in Journ. As. Soc. Beng.* xlii. (1873) ii. 63.—Burma.
gamosepala, *Turcz. in Bull. Soc. Nat. Mosc.* xxxvi. (1863) i. 575.—Hab.?
inaequilatera, Turcz. l. c. xxxi. (1858) i. 233 = serratifolia.
javanica, *Blume, Bijdr.* 117.—Java.
merguensis, *Planch. ex Mast. in Hook. f. Fl. Brit. Ind.* i. 394.—Burma.
serratifolia, Blanco, Fl. Filip. ed. II. 427 = Grewia columnaris.
serratifolia, *DC. Prod.* i. 512.—Ins. Philipp.

COLUMELLA, Comm. ex DC. Prod. i. 443, in syn. (1824) = **Pavonia**, Cav. (Malvac.).
Malvinda, Comm. ex DC. l. c. = P. Columella.

COLUMELLA, Lour. Fl. Cochinch. i. 85 (1790) = **Vitis**, Linn. (Ampelid.).
pedata, Lour. l. c. 86 = V. cochinchinensis.

COLUMELLA, Vahl, Enum. i. 300 (1805) = **Columellia**, Ruiz & Pav.

COLUMELLA, Vell. Fl. Flum. 155 (1825); iv. t. 17 = **Pisonia**, Linn. (Nyctagin.).
rustica, Vell. l. c. = P. subcordata.

COLUMELLEA, Jacq. Hort. Schoenbr. iii. 28. t. 301 (1798) = **Nestlera**, Spreng. (Compos.).
biennis, Jacq. l. c. 29 = N. biennis.

COLUMELLIA, Ruiz & Pav. Prod. 3. t. 1 (1794).
COLUMELLIACEAE, Benth. & Hook. f. ii. 989.
Uluxia, Juss. in Dict. Sc. Nat. x. 103 (1818).
arborescens, Pers. Syn. i. 13 = oblonga.
frutescens, Pers. l. c. = oblonga.
oblonga, *Ruiz & Pav. Fl. Per.* i. 28.—Peruv.
obovata, *Ruiz & Pav. l. c.*—Peruv.
sericea, *H. B. & K. Nov. Gen. et Sp.* ii. 388.—Ecuador.

COLUMNEA, Plum. ex Linn. Gen. ed. I. 373 (1737).
GESNERACEAE, Benth. & Hook. f. ii. 1009.
Aponoa, Rafin. Sylva Tellur. 84 (1838).
Collandra, Lem. in Fl. des Serres, iii. (1847) sub t. 223.
Dalbergaria, Tussac, Fl. Antill. i. 141. t. 19 (1808).
Eusynetra, Rafin. Fl. Tellur. ii. 57 (1836).
Glycanthes, Rafin. Sylva Tellur. 83 (1838).
Ortholoma, Hanst. in Linnaea, xxvi. (1853) 209.
Pentadenia, Hanst. l. c. 211 (1853).
Stenanthus, Oerst. ex Hanst. l. c. 209 (1853).
Stygnanthe, Hanst. l. c. (1853).
acuminata, *Benth. Pl. Hartw.* 231.—N. Granat.
anisophylla, *DC. Prod.* vii. 542.—Am. austr.
argentea, *Griseb. Fl. Brit. W. Ind.* 465.—Ind. occ.
aurantiaca, *Decne. ex Planch. in Fl. des Serres*, vi. (1850–51) 45.—Reg. Andin.
aurea, *Warcz. in Otto & Dietr. Allg. Gartenz.* xiii. (1845) 253.—N. Granat.?
aureonitens, *Hook. Bot. Mag.* t. 4294.—N. Granat.
balsamea, Roxb. Hort. Beng. 47; Fl. Ind. iii. 97 = Limnophila gratioloides.
bilabiata, *Seem. Bot. Voy. Herald*, 186.—Panama.
Billbergiana, *Beurl. in Vet. Akad. Handl. Stockh.* 1854 (1856) 135.—Panama.
campanulata, *Benth. Pl. Hartw.* 232.—N. Granat.
capitata, Walp. Rep. vi. 405, sphalm. = Besleria capitata.
coccinea, Hort. ex Hanst. in Linnaea, xxxiv. (1865–66) 405 = rotundifolia.
cochinchinensis, Poir. Encyc. Suppl. ii. 315 (= *Diceros cochinchinensis*).—Cochinch.
consanguinea, *Hanst. in Linnaea*, xxxiv. (1865–66) 383.—Costa Rica.
crassifolia, *Hook. Bot. Mag.* t. 4330.—Mexic.
divaricata, Walp. Rep. vi. 405, sphalm. = Besleria divaricata.
diffusa, Roxb. ex Hook. f. Fl. Brit. Ind. iv. 277 = Torenia vagans.
erecta, Lam. Encyc. ii. 66 = Achimenes coccinea.
eriantha, *Hanst. in Linnaea*, xxxiv. (1865–66) 391.—Ecuador.
erythrocalyx, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 133 = querceti.
erythrophaea, *Decne. ex Houll. in Rev. Hortic.* (1867) 172.—Mexic.
erythrophylla, *Hanst. in Linnaea*, xxxiv. (1865–66) 389.—N. Granat.
flaccida, *Seem. Bot. Voy. Herald*, 186.—Panama.
flava, *Mart. & Gal. in Bull. Acad. Brux.* ix. (1842) ii. 39.—Mexic.
glabra, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 136. —Costa Rica.
grandiflora, Hort. ex Hanst. in Mart. Fl. Bras. viii. 414 = Nematanthus longipes.
guttata, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 1.—Peruv.
henkelioides, Spreng. ex Heynh. Nom. ii. 154 = Streptocarpus Rexii.

COLUMNEA:—
heterophylla, *Hanst. in Linnaea*, xxxiv. (1865-66) 390.—Costa Rica.
heterophylla, Roxb. Hort. Beng. 47; Fl. Ind. iii. 97 = Limnophila heterophylla.
hirsuta, Klotzsch, ex Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 135 = hirta.
hirsuta, *Sw. Prod. Veg. Ind. Occ.* 94.—Jamaica.
hirta, *Klotzsch & Hanst. in Linnaea*, xxxiv. (1865-66) 403.—Costa Rica.
hispida, *Sw. Prod. Veg. Ind. Occ.* 94.—Jamaica.
humilis, *Lam. ex Steud Nom.* ed. I. 213; ed. II. i. 399.—Hab.?
inaequilatera, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 1.—Peruv.
Kalbreyeri, Hook. f. Bot. Mag. t. 6633 = Kalbreyeriana.
Kalbreyeriana, *Mast. in Gard. Chron.* (1882) i. 44 217.—N. Granat.
Kienastiana, *Regel, in Act. Hort. Petrop.* viii. (1883) 274.—N. Granat.
labellosa, *Karst. Fl. Columb.* ii. 105. t. 154.—N. Granat.
lepidocaula, *Hanst. in Linnaea*, xxxiv. (1865-66) 411.—Costa Rica.
Lindleyana, *Pasq. Cat. Orto Bot. Nap.* 31.—Hab.?
linearis, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 133.—Costa Rica.
longepedunculata, Hort. ex Hanst. in Mart. Fl. Bras. viii. 414 = Nematanthus longipes.
longifolia, Linn. Mant. 90 = Artanema sesamoides.
lucida, Walp. Rep. vi. 405, sphalm. = Besleria lucida.
macrantha, *Benth. Pl. Hartw.* 232.—Ecuador.
magnifica, *Klotzsch & Hanst. ex Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 134.—Costa Rica.
major, *Hanst. in Linnaea*, xxxiv. (1865-66) 388.—N. Granat.
microcalyx, *Hanst. l. c.* 408.—Costa Rica.
microphylla, *Klotzsch & Hanst. ex Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 137.—Costa Rica.
minor, *Hanst. in Linnaea*, xxxiv. (1865-66) 387.—N. Granat.
minuta, Roxb. Hort. Beng. 47; Fl. Ind. iii. 98 = Vandellia scabra.
mocsta, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 1. t. 206.—Peruv.
nervosa, *Klotzsch & Hanst. in Linnaea*, xxxiv. (1865-66) 400.—Costa Rica.
nicaraguensis, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 136.—Nicaragua.
ochrolcuca, *Klotzsch & Hanst. in Linnaea*, xxxiv. (1865-66) 393.—Costa Rica.
Oerstediana, *Klotzsch, ex Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 135.—Costa Rica.
ovata, *Cav. Ic.* iv. 62. t. 391.—Ins. Chiloe.
oxyphylla, *Hanst. in Linnaea*, xxxiv. (1865-66) 405.—Costa Rica.
pendula, *Klotzsch & Hanst. l. c.* 397.—Costa Rica.
pichinchensis, *Hanst. l. c.* 398.—Ecuador.
picta, *Karst. Fl. Columb.* ii. 105. t. 154.—N. Granat.
pilosa, Lem. in Fl. des Serres, iii. (1847) t. 223 = aureonitens.
praetexta, *Hanst. in Linnaea*, xxxiv. (1865-66) 394.—Costa Rica.
purpurata, *Hanst. l. c.* 386.—Costa Rica.
querceti, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 133.—Costa Rica.
repens, *Hanst. in Linnaea*, xxxiv. (1865-66) 396.—Costa Rica.
repens, Rafin. Sylva Tellur. 84 = scandens?
ringens, *Regel, in Act. Hort. Petrop.* viii. (1883) 273.—N. Grenat.
rotundifolia, *Salisb. Parad. Lond.* t. 29.—Ins. Trinit.
rutilans, *Sw. Prod. Veg. Ind. Occ.* 94.—Jamaica.
sanguinea, *Hanst. in Linnaea*, xxxiv. (1865-66) 384.—Ins. S. Doming.
sanguinolenta, *Hanst. l. c.* 389.—Costa Rica.
scabrosa, Soland. ex DC. Prod. ix. 277 = Rhabdothamnus scabrosus.
scandens, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 136 = Billbergiana.
scandens, *Linn. Sp. Pl.* 938.—Am. austr.
scandens-hirsuta, Schott, ex Hanst. in Mart. Fl. Bras. viii. 417 = Codonanthe Hookerii.
Schiedeana, *Schlecht. in Linnaea*, viii. (1833) 249.—Mexic.
serrata, *Hanst. in Linnaea*, xxxiv. (1865-66) 390.—Costa Rica.

COLUMNEA:—
speciosa, Presl, Bot. Bemerk. 145 = rotundifolia.
speciosa, Wendl. ex Steud. Nom. ed. II. i. 399 = Rytidophyllum prasinatum.
splendens, Paxt. Mag. Bot. x. (1843) 5 = Nematanthus longipes.
stellata, *Lour. Fl. Cochinch.* 384.—Cochinch.
strigosa, *Benth. Pl. Hartw.* 232.—N. Granat.
tenuis, *Klotzsch, ex Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 137.—Costa Rica.
tincta, *Griseb. in Mem. Am. Acad.* N. S. viii. (1863) 526.—Ins. Cuba.
tomentosa, *Oerst. in Vidensk. Selsk. Skr.* V. v. (1861) 138.—Nicaragua.
tomentosa, Roxb. Hort. Beng. 47; Fl. Ind. iii. 98 = Mazus rugosus.
trifoliata, Link, Enum. Hort. Berol. ii. 145 = Stemodia trifoliata.
umbellata, *Bertol. Fl. Guatim.* 26.—Guatemala.
vestita, Hanst. in Linnaea, xxxvi. (1865-66) 392 = Warszewicziana.
villosa, Willd. Enum. Hort. Berol. Suppl. 43 = scandens.
violacea, Jacq. f. ex Steud. Nom. ed. II. i. 399 = Stemodia trifoliata.
Warscewicziana, *Klotzsch & Hanst. in Linnaea*, xxxiv. (1865-66) 392.—Costa Rica.
Wendlandiana, *Hanst. l. c.* 402.—Costa Rica.
Wullschlaegeliana, *Hanst. l. c.* 414.—Jamaica.
zebrina, Hort. ex Loud. Encyc. Pl. Suppl. ii. 1402 = Alloplectus dichrous.

COLUPPA, Adans. Fam. ii. 268 (1763) = **Gomphrena**, Linn. (Amarantac.).

COLURIA, R. Br. in Parry, Voy. App. 276 (1823). *ROSACEAE*, Benth. & Hook. f. i. 619.
LAXMANNIA, Fisch. ex Ledeb. Fl. Alt. ii. 262 (1830).
geoides, Ledeb. Fl. Alt. ii. 263 = potentilloides.
mongolica, *Maxim. in Bull. Acad. Pétersb.* xxvii. (1881) 466.—As. or.
potentilloides, *R. Br. in Parry, 1st Voy. Suppl.* p. cclxxvi.—Sibir.

COLUS, Racusch. Nom. ed. III. 171 (1797) = **Coleus**, Lour. (Labiat.).

COLUTEA, [Tourn.] Linn. Syst. ed. I (1735). *LEGUMINOSAE*, Benth. & Hook. f. i. 505.
aeschinomenoides, *Scop. Delic. Insub.* iii. 22. t. 12.—Ins. Baham.
alpina, Lam. Encyc. i. 354 = Astragalus alpinus, membranaceus.
americana, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
annua, Murr. in Nov. Comm. Gotting. 1774 (1775) 40 = halepica.
aperta, Moench, Verz. 24 = orientalis.
arborescens, *Linn. Sp. Pl.* 723.—Europ.; Oriens.
arenaria, Poir. Encyc. Suppl. i. 562 = Astragalus chorinensis.
armena, *Boiss. & Huet. Diagn.* Sér. II. v. 83.—Armenia.
astragalina, Poir. Encyc. Suppl. i. 561 = Astragalus alpinus.
australis, Lam. Encyc. i. 354 = Astragalus australis.
baetica, Poir. Encyc. Suppl. i. 561 = Astragalus lusitanicus.
brevialata, *Lange, Ind. Sem. Hort. Haun.* (1861) 30.—Hab.?
caspica, Bieb. Fl. Taur. Cauc. ii. 169 = Sphaerophysa salsula.
cilicica, *Boiss. & Bal. Diagn.* Sér. II. v. 83.—Cilicia.
cruenta, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 55.—Reg. Caucas.
davurica, Spreng. Syst. iii. 242 = Sphaerophysa salsula.
excisa, Thunb. Prod. Pl. Cap. 134 = Lessertia excisa.
fistulosa, Retz. Obs. iii. 40 = Lessertia perennans.
floribunda, Poir. Encyc. Suppl. i. 562 = Sesbania vesicaria.
florida, Salisb. Prod. 337 = arborescens.
frigida, Poir. Encyc. Suppl. i. 561 = Astragalus frigidus.

COLUTEA:—
frutescens, Linn. Sp. Pl. 723 = Sutherlandia frutescens.
galegifolia, Sims, Bot. Mag. t. 792 = Swainsona galegifolia.
glabra, Poir. Encyc. Suppl. i. 561 = Astragalus australis.
grandiflora, Salisb. Prod. 338 = Sutherlandia frutescens.
halepica, *Lam. Encyc.* i. 353.—Arab.; Abyssin.
halicacaba, Poir. Encyc. Suppl. i. 562 = Astragalus Vulneraria.
herbacea, Linn. Sp. Pl. 723 = Lessertia linearis.
hirsuta, Roth, Tent. Fl. Germ. i. 305 = arborescens.
humilis, Scop. Delic. Insub. ii. 23 = orientalis.
incana, Poir. Encyc. Suppl. i. 562 = Astragalus anthylloides.
istria, Mill. Gard. Dict. ed. VIII. n. 2 = halepica.
linearis, Thunb. Prod. Pl. Cap. 135 = Lessertia linearis.
media, *Willd. Enum. Hort. Berol.* 771.—Hab.?
melanocalyx, Boiss. & Heldr. Diagn. Ser. I. ix. 35 = arborescens.
microphylla, Delile, Ind. Sem. Hort. Monsp. (1847) 7 = halepica.
nepalensis, Sims, Bot. Mag. t. 2622 = arborescens.
Novae-Hollandiae, Walp. Ann. ii. 368 = Clianthus Dampieri.
obtusata, Thunb. Prod. Pl. Cap. 134 = Lessertia obtusata.
orientalis, Lam. Encyc. i. 353 = cruenta.
pallida, Salisb. Prod. 337 = halepica.
perennans, Jacq. Enum. Vindob. App. 311 = Lessertia perennans.
persica, *Boiss. Diagn.* Ser. I. vi. 33.—Persia.
Pocockii, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 55 = halepica.
procumbens, Mill. Gard. Dict. ed. VIII. n. 7 = Lessertia procumbens.
prostrata, Thunb. Prod. Pl. Cap. 134 = Lessertia prostrata.
pubescens, Thunb. l. c. = Lessertia pubescens.
rigida, Thunb. l. c. = Lessertia fruticosa.
rubra, Medic. Beobacht. (1782) 359 = arborescens.
salsola, Poir. Encyc. Suppl. i. 562 = Sphaerophysa salsula.
sanguinea, Pall. Fl. Ross. ii. 88 = cruenta.
spinosa, Forsk. Fl. Aegypt. Arab. 131 = Astragalus Russellii.
tomentosa, Thunb. Prod. Pl. Cap. 135 = Lessertia tomentosa.
triflora, Poir. Encyc. Suppl. i. 562 = Astragalus triflorus.
trifoliata, Poir. l. c. (= *Phaca trifoliata*).—China.
triphylla, *Bunge, ex Boiss. Fl. Orient.* ii. 196.—Persia.
vesicaria, Thunb. Prod. Pl. Cap. 135 = Lessertia vesicaria.
versicolor, Salisb. Prod. 337 = cruenta.
Wightiana, Wall. ex Steud. Nom. ed. II. i. 399 = Sutherlandia frutescens.
wolgarica, Lam. Encyc. i. 353 = Calophaca wolgarica.

COLUTEOCARPUS, Boiss. in Ann. Sc. Nat. Sér. II. xxii. (1842) 162. *CRUCIFERAE*, Benth. & Hook. f. i. 73.
LAGOWSKIA, Trautv. in Bull. Phys.-Math. Acad. Pétersb. xvi. (1858) 321.
reticulatus, *Boiss. l. c.*—As. Min.

COLUTIA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 366 = **Sutherlandia**, R. Br. (1812) (Leguminos.).
amoena, Medic. l. c. = S. frutescens.
frutescens, Medic. Phil. Bot. i. 210 = S. frutescens.
rubra, Moench, Meth. 160 = S. frutescens.

COLVILLEA, Boj. ex Hook. Bot. Mag. tt. 3325, 3326 (1834). *LEGUMINOSAE*, Benth. & Hook. f. i. 569.
racemosa, *Boj. l. c.*—Madag.

COLYMBADA, Hill, Veg. Syst. iv. 31 (1762) = **Centaurea**, Linn. (Compos.).
centauroides, Hill, Hort. Kew. 65 = Cent. Rhapontica.
sicula, Hill, l. c. = Cent. sicula.

COLYMBEA, Steud. Nom. Bot. ed. II. i. 399 (1840) = COLUMBEA, Salisb. = **Araucaria**, Juss. (Conif.).

COLYRIS, Endl. Gen. 596 (1838) = COLLYRIS, Vahl = **Dischidia**, R. Br. (Asclepiad.).

COLYTHRUM, Schott, Rutac. 13. t. 7 (1834) = **Esenbeckia**, Kunth (Rutac.).
febrifugum, Schott, l. c. 14 = E. febrifuga.
grandiflorum, Steud. Nom. ed. II. i. 399 = E. grandiflora.
latifolium, Schott, Rutac. 14 = E. pumila.
maurioides, Schott, l. c. = E. maurioides.
puberulum, Schott, l. c. 13. t. 7.—Bras.
pumilum, Schott, l. c. 14 = E. pumila.

COMACEPHALUS, Klotzsch, in Linnaea, xii. (1838) 224 = **Grisebachia**, Klotzsch (Ericac.).
incurvus, Klotzsch, l. c.—Afr. austr.

COMACLINIUM, Scheidw. & Planch. in Fl. des Serres, viii. (1852) 19. t. 756 = **Dysodia**, Cav. (Compos.).
aurantiacum, Scheidw. & Planch. l. c. = D. grandiflora.

COMACUM, Adans. Fam. ii. 345 (1763) = **Myristica**, Linn.

COMANDRA, Nutt. Gen. Am. i. 157 (1818).
SANTALACEAE, Benth. & Hook. f. iii. 224.
HAMILTONIA, Spreng. Syst. i. 831 (1825), pro parte.
cuneifolia, Rafin. New Fl. Am. ii. 33 = umbellata.
Darbya, *A. DC. in DC. Prod.* xiv. 637.—Am. bor.
elegans, *Reichb. f. Ic. Fl. Germ.* xi. 11. t. 547. f. 1162.—Europ. or. austr.
elliptica, Rafin. New Fl. Am. ii. 33 = umbellata.
livida, *Richards. in Frankl. Narr. 1st Journ. App.* 734.—Am. bor.
media, Rafin. New Fl. Am. ii. 33 = umbellata.
obovata, Rafin. l. c. = umbellata.
obtusifolia, Rafin. l. c. = umbellata.
pallida, *A. DC. in DC. Prod.* xiv. 636.—Am. bor.
rigida, Steud. Nom. ed. II. i. 399 = livida.
umbellata, *Nutt. Gen. Am.* i. 157.—Am. bor.
umbellulata, Rafin. New Fl. Am. ii. 33 = praec.

COMANTHOSPHACE, S. Moore, in Journ. Bot. xv. (1877) 293. *LABIATAE.*
barbinervis, *S. Moore, l. c.*—Japon.
japonica, *S. Moore, l. c.*—Japon.
stellipila, *S. Moore, l. c.*—Japon.
sublanceolata, *S. Moore, l. c.*—Japon.

COMAROPSIS, Rich. in Nestl. Monog. Potent. 16. t. 1 (1816) = **Waldsteinia**, Willd. (Rosac.).
Doniana, Ser. in DC. Prod. ii. 555 = W. fragarioides.
fragarioides, Ser. l. c. = W. fragarioides.
pedata, Ser. l. c. = Rubus pedatus.
radicans, Ser. l. c. = Rubus radicans.
sibirica, Ser. l. c. = W. sibirica.

COMAROSTAPHYLIS, Zucc. in Abh. Akad. Muench. ii. (1837) 331 = **Arctostaphylos**, Adans. (Ericac.).
angustifolia, Klotzsch, in Linnaea, xxiv. (1851) 74 = A. angustifolia.
arbutoides, Lindl. Bot. Reg. (1843) t. 30 = A. arbutoides.
arguta, Zucc. in Linnaea, xii. (1838) Litt. 86 = A. arguta.
attenuata, Klotzsch, l. c. xxiv. (1851) 75 = A. attenuata.
bicolor, Klotzsch, l. c. 78 = A. bicolor.
formosa, Lem. in Illustr. Hortic. v. (1858) sub t. 162 = Pieris formosa.
glauca, Buckl. in Proc. Acad. Sc. Philad. '1861' (1862) 461 = Arbutus Menziesii.
glaucescens, Zucc. ex Klotzsch, in Linnaea, xxiv. (1851) 76 = A. glaucescens.
Hartwegiana, Klotzsch, in Linnaea, xxiv. (1851) 77 = A. Hartwegiana.
latifolia, Klotzsch, l. c. 76.—Mexic.
longifolia, Klotzsch, l. c. 75 = A. longifolia.
mucronata, Klotzsch, l. c. 77 = A. mucronata.
mucronifera, Klotzsch, l. c. 76 = A. mucronifera.
oaxacana, Klotzsch, l. c. 77 = A. oaxacana.
polifolia, Zucc. ex Klotzsch, l. c. = A. polifolia.
rubescens, Klotzsch, l. c. = A. rubescens.

COMAROUNA, Carr. in Rev. Hortic. (1873) 232 = COUMAROUNA, Aubl. = **Dipteryx**, Schreb. (Legumin.).

COMARUM, Linn. Syst. ed. I (1735) = **Potentilla**, Linn. (Rosac.).
angustifolium, Rafin. Fl. Tellur. ii. 56 = P. Comarum.
digitatum, Rafin. l. c. = P. Comarum.
flavum, Buch.-Ham. ex Roxb. Hort. Beng. 39; Fl. Ind. ii. 521 = P. supina.
fragarioides, Roth, Tent. Fl. Germ. ii. 577 = P. Fragariastrum.
palustre, Linn. Sp. Pl. 502 = P. Comarum.
rubrum, Gilib. Fl. Lituan. ii. 255 = P. Comarum.
Salessowii, Bunge, Del. Sem. Dorp. (1839) 8.—Cf. Linnaea, xiv. (1840) Lit. 119 = P. Salessoviana.
supinum, Alef. in Bot. Zeit. xxiv. (1866) 262 = P. supina?

COMASPERMUM, Pers. Syn. ii. 275 (1807) = **Comesperma**, Labill. (Polygal.).

COMATOCROTON. Karst. in C. Koch & Fint. Wochenschr. (1859) 6 = **Croton**, Linn. (Euphorbiac.)
ovalifolia, Karst. l. c. = C. ovalifolius.

COMATOGLOSSUM, Karst. & Triana, in Linnaea, xxviii. (1856) 436 = **Talisia**, Aubl.? (Sapindac.).
strictum, Karst. & Triana, l. c. 437.—N. Granat.

COMBESIA, A. Rich. Tent. Fl. Abyss. i. 307 (1847) = **Tillaea**, Linn. (Crassulac.).
abyssinica, A. Rich. l. c. = T. pharnaceoides.
Schimperi, Schweinf. in Beitr. Fl. Aethiop. 80 = T. pentandra.

COMBRETOCARPUS, Hook. f. in Benth. & Hook. f. Gen. i. 683 (1865). *RHIZOPHOREAE*, Benth. & Hook. f. l. c.
Motleyi, *Hook. f. l. c.*—Borneo.

COMBRETUM, Linn. Gen. ed. I. App. 308 (1737).
COMBRETACEAE, Benth. & Hook. f. i. 688.
AETIA, Adans. Fam. ii. 84 (1763).
BUREAVA, Baill. Adansonia, i. 71 (1860).
CALOPYXIS, Tul. in Ann. Sc. Nat. Sér. IV. vi. (1856) 86.
CHRYSOSTACHYS, Pohl, Pl. Bras. ii. 65. t. 143 (1831).
CRISTARIA, Sonner. Voy. Ind. iii. 284 (1782).
EMBRYOGONIA, Blume, Mus. Bot. Lugd. Bat. ii. 122 (1854).
FORSGARDIA, Vell. Fl. Flum. 152 (1825); iv. t. 13 (1827).
GONOCARPUS, Ham. Prod. Fl. Ind. Occ. 39 (1825).
PEVRAEA, Comm. ex Juss. Gen. 320 (1789).
POIVREA, Comm. ex Thou. Obs. Pl. Afr. Austr. 28 (1806).
SEGUIERA, Reichb. ex Oliver, Fl. Trop. Afr. ii. 424 (1871).
SCHEADENDRON, Bertol. f. in Mem. Acc. Sc. Bologn. ii. (1850) 572.
SHEADENDRON, Bertol. f. l. c. t. 40 (1850).
THILOA, Eichl. in Flora, xlix. (1866) 149.
accedens, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 234.—Guiana.
aculeatum, *Vent. Choix*, 58.—Afr. trop.
acuminatum, *Roxb. Hort. Beng.* [28]; *Fl. Ind.* ii. 228.—Ind. or.; Malaya.
acutum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 424.—Afr. trop.
adenogonium, Steud. ex A. Rich. Tent. Fl. Abyss. i. 266 = reticulatum.
adenophyllum, Mart. in Flora, xxiv. (1841) II. Beibl. 1 = Jacquini.
Afzelii, G. Don, in Trans. Linn. Soc. xv. (1827) 437 = grandiflorum.
Afzelii, Hort. ex A. Juss. in Walp. Rep. v. 338 = Hiraea Wiedeana.
albidum, G. Don, in Trans. Linn. Soc. xv. (1827) 429 = ovalifolium.
album, Pers. Syn. i. 411 = decandrum.
Alfredi, *Hance, in Journ. Bot.* ix. (1871) 131.—China.
alternifolium, Herb. Madr. ex Wight & Arn. Prod. 317 = Lumnitzera racemosa.
alternifolium, *Pers. Syn.* ii. 412.—Panama.
alternifolium, Spreng. Syst. ii. 331 = aculeatum.

COMBRETUM:—
altum, *Guill. & Perr. ex DC. Prod.* iii. 20.—Afr. trop.
anfractuosum, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 3.—Bras.
angolense, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 428.—Afr. trop.
anisopterum, *Welw. ex Laws. l. c.* 429.—Afr. trop.
apetalum, *Wall. Cat.* n. 3990.—Burma.
apiculatum, *Sond. in Linnaea*, xxiii. (1850) 45.—Afr. austr.
arboreum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 115.—Malaya.
argenteum, Bertol. Fl. Guatimal. 12 = erianthum.
argyrotrichum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 432.—Afr. trop.
assimile, *Eichl. in Mart. Fl. Bras.* xiv. II. 109.—Bras.
attenuatum, Wall. Cat. n. 3989 = quadrangulare.
Aubletii, *DC. Prod.* i. 19.—Guiana.
aurantiacum, *Benth. in Hook. Journ. Bot.* ii. (1840) 222.—Guiana.
auratum, Steud. Nom. ed. II. i. 400 = seq.
aureum, *Wall. Cat.* n. 3988.—Burma.
barbatum, *Newm. ex G. Don, in Loud. Hort. Brit.* 157.—Bras.
bellum, Steud. Nom. ed. II. i. 400 = trifoliatum.
Benthamianum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 220.—N. Granat.
Binderianum, *Kotschy, in Sitzb. Wien. Akad.* li. *Abth.* II. (1865) 363.—Afr. trop.
Blanchetii, *Eichl. in Mart. Fl. Bras.* xiv. II. 114.—Bras.
bracteatum, Herb. Madr. ex Wall. Cat. n. 4009 = decandrum.
brevistylum, *Eichl. in Mart. Fl. Bras.* xiv. II. 118.—Bras.
Bugi, Cambess. in St. Hil. Fl. Bras. Mer. ii. 247. t. 130 = Jacquini.
butyrosum, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 87, in nota.—Afr. trop.
capituliflorum, *Fenzl, ex Schweinf. in Reliq. Kotsch.* 33.—Afr. trop.
celastroides, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 422.—Afr. trop.
chinense, *Roxb. Hort. Beng.* [28]; *Fl. Ind.* ii. 230.—As. trop.
chrysophyllum, Guill. & Perr. Fl. Seneg. Tent. i. 289 = tomentosum.
coccineum, *Lam. Encyc.* i. 734.—Madag.; Ins. Maurit.
collinum, *Fresen. in Mus. Senckenb.* ii. (1837) 153.—Abyssin.
comosum, *G. Don, in Edinb. Phil. Journ.* (1824) 344.—Afr. trop.
confertum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 422.—Afr. trop.
constrictum, *Laws. l. c.* 423.—Afr. trop.
cordatum, *G. Don, in Trans. Linn. Soc.* xv. (1827) 440.—Ins. S. Doming.
corylifolium, *Griseb. Fl. Brit. W. Ind.* 711.—Ind. occ.
corymbosum, Schum. in Schum. & Thonn. Beskr. Guin. Pl. 185 = racemosum.
costatum, Roxb. Hort. Beng. [28]; Fl. Ind. ii. 227 = acuminatum.
cuspidatum, *Planch. ex Benth. in Hook. Niger Fl.* 340.—Afr. trop.
cyclophyllum, Steud. Nom. ed. II. i. 400 = extensum.
dasystachyum, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 187.—Reg. Himal.; Burma.
decandrum, Jacq. Enum. Pl. Carib. 19 = alternifolium.
decandrum, Ruiz & Pav. ex G. Don, Gen. Syst. ii. 665 = Pavonii.
decandrum, *Roxb. Pl. Corom.* i. 43. t. 59.—Ind. or.; Burma.
dipterum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 429.—Afr. trop.
distillatorium, Blanco, Fl. Filip. ed. I. 295 = ovalifolium.
Duarteanum, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 248.—Bras.
elaeagnifolium, Planch. in App. Speke, Journ. Disc. Nile, 634 = collinum.
elaeagnoides, *Klotzsch, in Peters, Reise Mossamb. Bot.* 73.—Afr. trop.
elegans, *Cambess. in St. Hil. Fl. Bras. Mer.* ii. 247.—Bras.
elegans, H. B. & K. Nov. Gen. et Sp. vi. 109. t. 538 = Aubletii.

COMBRETUM :—
elegans, Wall. Cat. n. 4003 = trifoliatum.
erianthum, *Benth. Pl. Hartw.* 73.—Guatemala.
eriopetalum, *G. Don*, *Gen. Syst.* ii.665.—Ins. S. Doming.
erythrophyllum, *Sond. in Linnaea*, xxiii. (1850) 43.—Afr. austr.
extensum, *Roxb. Hort. Beng.* [28]; *Fl. Ind.* ii. 229.—Ind. or.; Malaya.
farinosum, *H. B. & K. Nov. Gen. et Sp.* vi. 110.—Mexic.
ferrugineum, *G. Don, in Trans. Linn. Soc.* xv. (1827) 440.—Am. austr.
ferrugineum, Hoffmgg. ex Walp. Rep. ii. 65 = Jacquini.
ferrugineum, A. Rich. Tent. Fl. Abyss. i. 267 = trichanthum.
flagrocarpum, *Herb. Calc. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 455.—Ind. or.; Burma.
floccosum, *Eichl. in Mart. Fl. Bras.* xiv. II. 119.—Bras.
formosum, *G. Don, in Trans. Linn. Soc.* xv. (1827) 420.—Bras.
formosum, Griff. Notul. iv. 682 = extensum.
frangulaefolium, *H. B. & K. Nov. Gen. et Sp.* vi. 109.—Venezuela.
fuscum, *Planch. ex Benth. in Hook. Niger Fl.* 339.—Afr. trop.
gallabatense, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 664.—Afr. trop.
Geheebii, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 223.—Bras.
glabrum, *DC. Prod.* iii. 19.—Guiana.
glaucocarpum, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 3.—Bras.
glomeruliflorum, *Sond. in Linnaea*, xxiii. (1850) 42.—Afr. austr.
glutinosum, *Guill. & Perr. Fl. Seneg. Tent.* 288. t. 68.—Afr. trop.
glutinosum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 268 = lepidotum.
Goldieanum, *F. Muell. Desc. Papuan Pl.* iv. 66.—N. Guinea.
gracile, *Schott, in Spreng. Syst.* iv. *App.* 406.—Bras.
grandiflorum, *G. Don, in Edinb. Phil. Journ.* (1824) 346.—Afr. trop.
Griffithii, Heurck & Muell. Arg. in Heurck, Obs. Bot. 231 = chinense.
Gueinzii, *Sond. in Linnaea*, xxiii. (1850) 43.—Afr. austr.
guianense, Miq. in Ann. Mag. Nat. Hist. xi. (1843) 12 = Aubletii.
Hartmannianum, *Schweinf. Beitr. Fl. Aethiop.* 24. t. 3.—Afr. trop.
herbaceum, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 347.—Afr. trop.
Heyneanum, Wall. Cat. n. 4001 = ovalifolium.
Hilariana, *D. Dietr. Syn. Pl.* ii. 1303.—Bras.
hispidum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 421.—Afr. trop.
holosericeum, *Sond. in Linnaea*, xxiii. (1850) 44.—Afr. austr.
Horsfieldii, Miq. Fl. Ind. Bat. i. 1. 609 = extensum.
imberbe, *Wawra, in Sitzb. Acad. Wien, Math.-Nat.* xxxviii. (1860) 556.—Afr. trop.
insigne, Heurck & Muell. Arg. in Heurck, Obs. Bot. 247 = pilosum.
intermedium, G. Don, in Edinb. Phil. Journ. xi. (1824) 345 = comosum.
Jacquini, *Griseb. Fl. Brit. W. Ind.* 275.—Ind. occ.
Kirkii, *Laws. in Oliver, Fl. Trop. Afr.* ii. 429.—Afr. trop.
Klotzschii, *Welw. ex Laws. l. c.* 422.—Afr. trop.
Kraussii, *Hochst. in Flora*, xxvii. (1844) II. 424.—Afr. austr.
laetum, Wall. Cat. n. 4004 = pilosum.
lanceolatum, *Pohl, ex Eichl. in Mart. Fl. Bras.* xiv. II. 110.—Bras.
lanuginosum, *G. Don, in Trans. Linn. Soc.* xv. (1827) 422.—Abyssa.
latifolium, Blume, Bijdr. 641 = extensum.
latifolium, *G. Don, in Trans. Linn. Soc.* xv. (1827) 432.—Ind. or.
laurifolium, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 64.—Bras.
laxiflorum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 428.—Afr. trop.
laxum, Aubl. Fl. Gui. 351 = Aubletii.
laxum, Jacq. Enum. Pl. Carib. 19 = Jacquini.

COMBRETUM :—
laxum, Loefl. It. 248, 308 = Loeflingii.
laxum, Roxb. Hort. Beng. [88]; Fl. Ind. ii. 231 = ovalifolium.
lepidotum, Presl, Bot. Bemerk. 142 = squamosum.
lepidotum, *A. Rich. Tent. Fl. Abyss.* i. 268.—Abyss.
leprosum, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 1.—Bras.
leptostachyum, *Mart. l. c.* 2.—Bras.
leucanthum, Heurck & Muell. Arg. in Heurck, Obs. Bot. 240 = extensum.
leucophyllum, G. Don, in Edinb. Phil. Journ. xi. (1824) 346 = racemosum.
Loeflingii, *Eichl. in Mart. Fl. Bras.* xiv. II. 110.—Am. trop.
lucidum, Blume, Bijdr. 641 = trifoliatum.
macrocarpum, Beauv. Fl. Owar. ii. 90. t. 118 = racemosum.
macrophyllum, Roxb. Hort. Beng. [88]; Fl. Ind. ii. 231 = latifolium.
macrostachyum, Wall. Cat. n. 3997 = extensum.
Madolata, Buch.-Ham. ex Wall. Cat. n. 4009 = decandrum.
magnificum, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 4.—Bras.
Malulaea, Wall. Cat. n. 3991 = squamosum.
Mechowianum, *O. Hoffm. in Linnaea*, xliii. (1880–82) 131.—Afr. trop.
mellifluum, *Eichl. in Mart. Fl. Bras.* xiv. II. 113.—Bras.
mexicanum, *Humb. & Bonpl. Pl. Aequin.* ii. 159. t. 132.—Mexic.
micranthum, G. Don, in Edinb. Phil. Journ. xi. (1824) 347 = altum.
micropetalum, DC. Prod. iii. 19 = Loeflingii.
micropetalum, Llanos, in Mem. Acad. Cienc. Madr. iv. (1856) 502 = extensum.
microphyllum, *Klotzsch, in Peters, Reise Mossamb. Bot.* 74.—Afr. trop.
molle, *R. Br. ex G. Don, in Trans. Linn. Soc.* xv. (1827) 431.—Abyss.
Monetaria, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 2.—Bras.
mucronatum, *Schum. & Thonn. Beskr. Guin. Pl.* 184.—Afr. trop.
myrtifolium, *Laws. in Oliver, Fl. Trop. Afr.* ii. 431.—Afr. trop.
nanum, *Buch.-Ham. in D. Don, Prod. Fl. Nep.* 219.—Reg. Himal.
neurophyllum, Miq. Fl. Ind. Bat. i. 1. 608 = acuminatum.
nigricans, *Leprieur, ex Guill. & Perr. Fl. Seneg. Tent.* i. 290.—Afr. trop.
nitidum, *Spruce, ex Eichl. in Mart. Fl. Bras.* xiv. II. 118.—Bras.
obscurum, *Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 83.—Madag.
obtusifolium, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 108.—Guian.
obtusum, Walp. Rep. ii. 67 = prace.?
occidentalis, *Linn. Syst.* ed. X. 999.—Am. austr.
odoratum, *Pav. ex G. Don, in Trans. Linn. Soc.* xv. (1827) 430.—Ecuador.
ovale, R. Br. ex G. Don, l. c. 434 = aculeatum.
ovalifolium, *Roxb. Hort. Beng.* [28]; *Fl. Ind.* ii. 226.—Ind. or.
oxypetalum, G. Don, in Trans. Linn. Soc. xv. (1827) 420 = Loeflingii.
oxystachyum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 422.—Afr. trop.
pachycladum, *Baker, in Journ. Linn. Soc.* xviii. (1881) 270.—Madag.
paniculatum, *Vent. Choix*, t. 58.—Afr. trop.
paradoxum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 430.—Afr. trop.
parviflorum, *Eichl. in Mart. Fl. Bras.* xiv. II. 114.—Bras.
parviflorum, Reichb. ex DC. Prod. iii. 21 = altum.
Pavonii, *G. Don, in Trans. Linn. Soc.* xv. (1827) 436.—Ecuador.
pedicellare, *DC. Prod.* iii. 20.—Bras.
pentagonum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 424.—Afr. trop.
Petitianum, *A. Rich. Tent. Fl. Abyss.* i. 268.—Abyss.
phaeocarpum, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 1.—Bras.
pilosum, *Roxb. Hort. Beng.* [28]; *Fl. Ind.* ii. 231.—Ind. or.; Burma.

COMBRETUM :—
Pincianum, *Hook. Bot. Mag.* (1846) sub t. 4262.—Afr. trop.
pisoniaeflorum, *Klotzsch, in Peters, Reise Mossamb. Bot.* t. 14.—Afr. trop.
platypetalum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 433.—Afr. trop.
platyphyllum, Heurck & Muell. Arg. in Heurck, Obs. Bot. 242 = extensum.
polyandrum, Hoffmgg. ex Walp. Rep. ii. 6 = Moquilia paraënsis.
Porterianum, Wall. Cat. n. 4000 = chinense.
puberum, *Rich. in Act. Soc. Hist. Nat. Par.* i. (1792) 108.—Guiana.
pulchellum, Mart. in Flora, xxii. (1839) I. Beibl. 64 = Jacquini.
punctatum, A. Rich. Tent. Fl. Abyss. i. 266 = lepidotum.
punctatum, *Blume, Bijdr.* 640.—Java.
punctatum, Steud. in Flora, xxvi. (1843) II. 761 = Aubletii.
purpureum, Vahl, Symb. Bot. iii. 51 = coccineum.
pyramidatum, *Desv. in Ham. Prod. Fl. Ind. Occ.* 35.—Guiana.
pyrifolium, *Kurz, in Journ. As. Soc. Beng.* xliii. (1874) II. 188.—Burma.
quadrangulare, *Kurz, l. c.*—Burma.
Quartinianum, A. Rich. Tent. Fl. Abyss. i. 266 = lepidotum.
racemosum, *Beauv. Fl. Owar.* 87. t. 118.—Afr. trop.
reticulatum, *Fresen. in Mus. Senckenb.* ii. (1837) 154.—Abyss.
reticulatum, *Presl, Reliq. Haenk.* ii. 25.—Mexic.
Richardianum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 234.—Abyss.
riparium, *Sond. in Linnaea*, xxiii. (1850) 47.—Afr. austr.
Rochetianum, *A. Rich. ex A. Juss. in Ann. Sc. Nat.* Sér. III. xv. (1851) 369.—Abyss.
rotundifolium, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 108 = Aubletii.
rotundifolium, Roxb. Hort. Beng. [88]; Fl. Ind. ii. 226 = extensum.
Roxburghii, G. Don, in Trans. Linn. Soc. xv. (1827) 429 = ovalifolium.
Roxburghii, Spreng. Syst. ii. 331 = decandrum.
rubiginosum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 428.—Afr. trop.
Rueppellianum, *A. Rich. Tent. Fl. Abyss.* i. 267.—Abyss.
salicifolium, *E. Mey. ex Hook. Ic. Pl.* t. 592.—Afr. austr.
sarcopterum, Thw. Enum. Pl. Zeyl. 415 = acuminatum.
Schimperianum, *A. Rich. Tent. Fl. Abyss.* i. 265.—Abyss.
secundum, G. Don, in Trans. Linn. Soc. xv. (1827) 419 = Aubletii.
secundum, Jacq. Enum. Pl. Carib. 19 = Jaequini.
semiadnatum, *Heurck & Muell. Arg. in Heurck, Obs. Bot.* 244.—Ind. or.
sericeum, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 347.—Afr. trop.
sericeum, Wall. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 450 = Calycopteris floribunda.
Smeathmanni, G. Don, in Trans. Linn. Soc. xv. (1827) 425 = mucronatum.
Sonderi, *Gerr. ex Sond. in Harv. & Sond. Fl. Cap.* ii. 511.—Afr austr.
spinescens, Wall. Cat. n. 4006 = pilosum.
spinosum, G. Don, in Edinb. Phil. Journ. xi. (1824) 345 = paniculatum.
spinosum, Humb. & Bonpl. Pl. Aequin. ii. 161 = alternifolium.
Sprucei, *Eichl. in Mart. Fl. Bras.* xiv. II. 115.—Bras.
squamosum, *Roxb. Hort. Beng.* [88]; *Fl. Ind.* ii. 231.—Ind. or.; Malaya.
stelligerum, *Presl, Epim. Bot.* 215.—Ind. or.
stenopetalum, Heurck & Muell. Arg. in Heurck, Obs. Bot. 225 = acuminatum.
stigmarium, *Mart. in Flora*, xxiv. (1841) II. *Beibl.* 3.—Bras.
subalternans, Wall. Cat. n. 4008 = trifoliatum.
sundaicum, *Miq. Fl. Ind. Bat. Suppl.* 327.—Sumatra.
terminalioides, *Steud. in Flora*, xxvi. 1843 II. 762.—Guiana.
ternatum, Wall. Cat. n. 4002 = chinense.
tetragonocarpum, *Kurz, in Journ. As. Soc. Beng.* xli. (1872) II. 306.—Burma.

COMBRETUM :—
tetragonum, *Laws. in Oliver, Fl. Trop. Afr.* ii. 430.—Afr. trop.
tetragonum, *Presl, ex Steud. Nom.* ed. II. i. 400.—Mexic.
tetralophum, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 454.—Malaya.
Theuschii, *O. Hoffm. in Linnaea*, xliii. (1880–82) 132.—Afr. trop.
Thwaitesianum, Heurck & Muell. Arg. in Heurck, Obs. Bot. 238 = ovalifolium.
tinctorum, Welw. ex Laws. in Oliver, Fl. Trop. Afr. ii. 430 = Kirkii.
tomentosum, *G. Don, in Edinb. Phil. Journ.* xi. (1824) 346.—Afr. trop.
tomentosum, Willd. ex Eichl. in Mart. Fl. Bras. xiv. II. 111 = frangulaefolium.
trichanthum, *Fresen. in Mus. Senckenb.* ii. (1837) 155.—Abyssin.
trifoliatum, *Vent. Choix*, t. 58.—Hab. ?
trigonoides, Perr. ex DC. Prod. iii. 20 = racemosum.
truncatum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 427.—Afr. trop.
undulatum, Wall. Cat. n. 3993 = trifoliatum.
Vanheurckii, *Muell. Arg. in Heurck, Obs. Bot.* 227.—Ind. or.
variabile, *Mart. in Flora*, xxii. (1839) I. *Beibl.* 62.—Bras.
velutinum, *DC. Prod.* iii. 20.—Bras.
vernicosum, Fenzl, in Flora, xxvii. (1844) 312 = lepidotum.
villosum, *Boj. ex Tul. in Ann. Sc. Nat.* Sér. IV. vi. (1856) 80, in syn.—Madag.
violaceum, *Boj. ex Tul. l. c.* 79.—Madag.
virgatum, Wall. Cat. n. 3992 = apetalum.
virgatum, *Welw. ex Laws. in Oliver, Fl. Trop. Afr.* ii. 425.—Afr. trop.
viscidum, *Griseb. Cat. Pl. Cub.* 109.—Cuba.
Wallichii, *DC. Prod.* iii. 21.—Reg. Himal.
Wallichii, Kurz, For. Fl. Burm. i. 465 = flagrocarpum.
Wightianum, Thw. Enum. Pl. Zeyl. 103, partim = ovalifolium.
Wightianum, Wall. Cat. n. 4007 = extensum.
Zeyheri, *Sond. in Linnaea*, xxiii. (1850) 46.—Afr. austr.

COMESPERMA, Labill. Nov. Holl. Pl. ii. 21. t. 159–163 (1806). *POLYGALEAE*, Benth. & Hook. f. i. 138.
acerosum, *Steetz, in Lehm. Pl. Preiss.* ii. 299.—Austral.
acutifolium, Steetz, l. c. 296 = ericinum.
aemulum, Steud. l. c. i. 208 = virgatum.
anthocarpum, Walp. Rep. v. 66, sphalm. = *xanthocarpum*.
aphyllum, *R. Br. ex Benth. Fl. Austral.* i. 143.—Austral.
calymega, *Labill. Nov. Holl. Pl.* ii. 23.—Austral.
ciliatum, *Steetz, in Lehm. Pl. Preiss.* ii. 304.—Austral.
compactum, Sieber, ex Steud. Nom. ed. II. i. 400 = retusum.
confertum, *Labill. Nov. Holl. Pl.* ii. 23. t. 161.—Austral.
contractum, Steud. in Lehm. Pl. Preiss. i. 208 = virgatum.
coridifolium, A. Cunn. in Field, N. S. Wales, 337 = ericinum.
corniculatum, Steud. in Lehm. Pl. Preiss. i. 206 = virgatum.
defoliatum, *F. Muell. Vict. Pl.* i. 189.—Austral.
Drummondii, *Steetz, in Lehm. Pl. Preiss.* ii. 301.—Austral.
ericinum, *DC. Prod.* i. 334.—Austral.
flavum, *DC. l. c.*—Austral.
floribundum, St. Hil. Fl. Bras. Mer. ii. 55. t. 91 = Bredemeyera floribunda.
gracile, Paxt. Mag. Bot. v. (1838) 145 = volubile.
herbaceum, Steud. in Lehm. Pl. Preiss. i. 211 = calymega.
hirtulum, Steud. l. c. 209 = confertum.
integerrimum, *Endl. in Enum. Pl. Hueg.* 7.—Austral.
isocalyx, Spreng. Syst. iii. 172 = calymega.
Kunthianum, St. Hil. Fl. Bras. Mer. ii. 54. t. 90 = Bredemeyera Kunthiana.
lanceolatum, *R. Br. ex Benth. Fl. Austral.* i. 148.—Austral.
latifolium, Steetz, in Lehm. Pl. Preiss. ii. 295 = ericinum.
laurifolium, St. Hil. Fl. Bras. Mer. ii. 56 = Bredemeyera laurifolia.
laxiusculum, Steud. in Lehm. Pl. Preiss. i. 210 = virgatum.
Lhotzkyanum, Manso, ex Klotzsch, in Peters, Reise Mossamb. Bot. 117, nomen (= Bredemeyerae sp.).—Afr. trop.
linariaefolium, A. Cunn. ex Steetz, in Lehm. Pl. Preiss. ii. 297 = ericinum.
longebracteatum, Steud. l. c. i. 207 = virgatum.
longifolium, Steud. l. c. 206 = confertum.
lucidum, Baill. Hist. Pl. v. 82 = Bredemeyera lucida.
Mansonii, Lhotsky, ex Klotzsch, in Peters, Reise Mossamb. Bot. 117 = Bredemeyera lucida.
megapteryga, Steud. in Lehm. Pl. Preiss. i. 207 = nudiusculum.
nudiusculum, *DC. Prod.* i. 334.—Austral.
nudiusculum, Steetz, in Lehm. Pl. Preiss. ii. 308 = defoliatum.
parviflorum, Steud. l. c. i. 210 = calymega.
patentifolium, F. Muell. Fragm. i. 48 = ericinum.
paucifolium, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 352 = volubile.
planifolium, *Hassk. in Miq. Ann. Mus. Bot. Lugd. Bat.* i. 185.—Austral.
polygaloides, *F. Muell. in Trans. Phil. Soc. Vict.* i. (1855) 7.—Austral.
praecelsum, *F. Muell. Fragm.* xi. 2.—Austral.
pubescens, Gardn. ex A. W. Benn. in Mart. Fl. Bras. xiii. III. 49 = Bredemeyera brevifolia.
ramosissimum, Steud. in Lehm. Pl. Preiss. i. 209 = nudiusculum.
retusum, *Labill. Nov. Holl. Pl.* ii. 22. t. 160.—Austral.
rhadinocarpum, *F. Muell. Fragm.* xi. 1.—Austral.
roseum, Steud. in Lehm. Pl. Preiss. i. 207 = virgatum.
scandens, Steud. l. c. 211 = integerrimum.
scoparium, *Drum. in Hook. Journ. Bot.* ii. (1840) 369; *Steetz, in Lehm. Pl. Preiss.* ii. 309.—Austral.
secundum, *Banks, ex DC. Prod.* i. 334.—Austral.
selaginoides, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 352 = virgatum.
silvestre, F. Muell. Fragm. i. 49 = sylvestre.
simplex, Endl. in Enum. Pl. Hueg. 7 = virgatum.
spathulatum, Turcz. in Bull. Soc. Nat. Mosc. xxvii. (1854) II. 352 = calymega.
sphaerocarpum, *Steetz, in Lehm. Pl. Preiss.* ii. 314.—Austral.
spinosum, *F. Muell. Fragm.* i. 144.—Austral.
strictum, Endl. in Enum. Pl. Hueg. 7 = calymega.
subherbaceum, Walp. Ann. ii. 84, sphalm. = calymega.
sylvestre, *Lindl. in Mitch. Journ. Trop. Austral.* 342.—Austral.
tenue, Steud. in Lehm. Pl. Preiss. i. 208 = calymega.
tortuosum, Steetz, l. c. ii. 303 = volubile.
variabile, Mart. ex Steud. Nom. ed. II. i. 400 (= Bredemeyerae sp.).—Bras.
varians, Steud. in Lehm. Pl. Preiss. i. 210 = calymega.
virgatum, *Labill. Nov. Holl. Pl.* ii. 21. t. 159.—Austral.
viscidulum, *F. Muell. Fragm.* x. 4.—Austral.
volubile, *Labill. Nov. Holl. Pl.* ii. 24.—Austral.
xanthocarpum, Steud. in Lehm. Pl. Preiss. i. 209 = flavum.

COMETES, Linn. Mant. i. 4 (1767). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 18.
CERATONYCHIA, Edgew. in Journ. As. Soc. Beng. xvi. (1847) 1215.
SALTIA, R. Br. in Salt, Abyss. App. 64 (1814), nomen.
abyssinica, *R. Br. in Wall. Pl. As. Rar.* i. 18.—Abyssin.
alternifolia, Linn. Syst. ed. XII. 127 = surattensis.
apiculata, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 244 = surattensis.
surattensis, *Linn. Mant.* i. 39; *Burm. f. Fl. Ind.* 39. t. 15.—Ind. bor. occ.; Arab.

COMETIA, Thou. ex Baill. Etud. Gén. Euph. 642 (1858). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 283.
lucida, *Baill. Adansonia*, ii. (1861) 55.—Madag.
Thouarsii, *Baill. Etud. Gén. Euph.* 643.—Madag.

COMEURYA, Baill. Adansonia, x. (1872) 329 = **Dracontomelon**, Blume (Anacard.).
Cumingiana, Baill. l. c. 330 = D. Cumingianum.

COMMARUM, Schrank, Prim. Fl. Salisb. 132 (1792) = COMARUM, Linn. = **Potentilla**, Linn. (Rosac.).

COMMELINA, Plum. ex Linn. Syst. ed. I. (1735). *COMMELINACEAE*, Benth. & Hook. f. iii. 847.
ANANTHOPUS, Rafin. Fl. Ludov. 20 (1817).
DISECOCARPUS, Hassk. in Flora, xlix. (1866) 211.
ERXLEBIA, Medic. in Act. Acad. Theod. Palat. vi. Phys. (1790) 494.
HEDWIGIA, Medic. l. c. 495 (1790).
HETEROCARPUS, Wight, Ic. vi. 29. t. 2067 (1853).
LECHEA, Lour. Fl. Cochinch. 60 (1790).
OMPHALOTHECA, Hassk. in Bull. Congr. Bot. Amst. (1865) 103.
SPATHODITHYROS, Hassk. in Flora, xlix. (1866) 211.
TRITHYROCARPUS, Hassk. l. c. (1866).
ALLOSPERMA, ALLOTRIA, DITELESIA, DIRTEA, EUDIPETALA, LARNALLES, NEPHRAELES, et OVIDIA, Rafin. Fl. Tellur. iii. 68–70, 122 (1836).
acuminata, *H. B. & K. Nov. Gen. et Sp.* i. 258.—Mexic.
acuminata, Poir. Encyc. Suppl. ii. 324 = Aneilema acuminatum.
acuminata, R. Br. in Salt, Abyss. App. 62 = pyrrhoblepharis.
aequinoctialis, Roem. & Schult. Syst. i. 538 = Aneilema aequinoctiale.
aequinoxialis, Beauv. Fl. Owar. i. t. 38 = praec.
aethiopica, *C. B. Clarke, in DC. Monog. Phan.* iii. 189.—Afr. trop.
affinis, Poir. Encyc. Suppl. ii. 324 = Aneilema gramineum.
africana, *Linn. Sp. Pl.* 41.—Afr. austr.
africana, Mirb. in Ann. Mus. Par. xvi. (1810) 456 = communis.
agraria, Kunth, Enum. Pl. iv. 38 = nudiflora.
agrostophylla, *F. Muell. Fragm.* viii. 59.—Austral.
alba, Buch.-Ham. ex C. B. Clarke, in DC. Monog. Phan. iii. 186 = appendiculata.
albescens, *Hassk. in Schweinf. Beitr. Fl. Aethiop.* 210.—Afr. trop.
ambigua, Beauv. Fl. Owar. i. 26. t. 15 = Palisota ambigua.
amplexicaulis, *Hassk. in Schweinf. Beitr. Fl. Aethiop.* 208.—Afr. trop.
angolensis, *C. B. Clarke, in DC. Monog. Phan.* iii. 167.—Afr. trop.
angustifolia, Hassk. in Peters, Reise Mossamb. Bot. 528 = Gerrardi.
angustifolia, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = graminifolia.
angustifolia, Michx. Fl. Bor. Am. i. 24 = virginica.
angustifolia, Reichb. Ic. Bot. Exot. ii. 17. t. 143 = graminifolia.
anthericoides, Poir. Encyc. Suppl. ii. 324 = Aneilema gramineum.
appendiculata, *C. B. Clarke, Commel. et Cyrt. Beng.* t. 13.—Ind. or.
aspera, *G. Don, ex Benth. in Hook. Niger Fl.* 542.—Afr. trop.
attenuata, Heyne, ex Steud. Nom. ed. II. i. 401 = subulata.
attenuata, *Koen. ex Vahl, Enum.* ii. 168.—Ind. or.
auriculata, *Blume, Enum. Pl. Jav.* i. 2.—Sumatra.
auriculata, E. Mey. in Presl, Rel. Haenk. i. 137 = virginica.
avenaefolia, *J. Grah. Cat. Pl. Bombay*, 224.—Ind. or.
axillaris, Linn. Sp. Pl. 42 = Cyanotis axillaris.
bahiensis, Hoffmgg. ex Spreng. Jahrb. i. III. 73 = virginica.
Bainesii, *C. B. Clarke, in DC. Monog. Phan.* iii. 184.—Afr. austr.
barbata, Boj. Hort. Maurit. 359 = communis.
barbata, *Lam. Tabl. Encyc.* i. 129.—Am. austr.
barbata, Sieber, ex C. B. Clarke, in DC. Monog. Phan. iii. 247 = Cyanotis cristata.
benghalensis, Forsk. Fl. Aegypt. Arab. 12 = divaricata.
benghalensis, *Linn. Sp. Pl.* 41.—Geront. trop.
benghalensis, Wall. Cat. n. 8980 = persicariaefolia.
beniniensis, Beauv. Fl. Owar. ii. 49. t. 87 = Aneilema beninense.
bicolor, Poepp. ex Kunth, Enum. Pl. iv. 89 = Tradescantia gracilis.
biflora, Poir. Encyc. Suppl. ii. 324 = Aneilema biflorum.
Blumii, A. Dietr. Sp. Pl. ii. 400 = auriculata.
Boissieriana, *C. B. Clarke, in DC. Monog. Phan.* iii. 161.—Afr. trop.
bracteolata, Lam. Encyc. ii. 69 = Aneilema spiratum.

COMMELINA :—

bracteosa, *Hassk. in Flora*, xlvi. (1863) 386.—Afr. trop.
Brownii, A. Dietr. Sp. Pl. ii. 415 = Aneilema acuminatum.
bulbosa, Heyne, ex Steud. Nom. ed. II. i. 401 = Aneilema dimorphum.
caerulea, Salisb. Prod. 216 = virginica.
caespitosa, Roxb. Hort. Beng. 5; Fl. Ind. i. 174 = nudiflora.
cajannensis, *Sieber, ex Steud. Nom.* ed. II. i. 401.—Ins. Martinic.
cajennensis, Kunth, Enum. Pl. iv. 38, in syn. = nudiflora.
Callisia, Steud. Nom. ed. II. i. 401 = Ancilema Hamiltonianum.
canariensis, Ch. Sm. in Buch, Canar. Ins. 140 = nudiflora.
canescens, Vahl, Enum. ii. 173 = benghalensis.
capitata, *Benth. in Hook. Niger Fl.* 541.—Afr. trop.
capitata, Klotzsch, ex C. B. Clarke, in DC. Monog. Phan. iii. 178 = obliqua.
cardiocephala, Kunze, ex C. B. Clarke, l. c. 157 = quitensis.
cardiosepala, Kunze, in Linnaea, xx. (1847) 9 = quitensis.
caripensis, H. B. & K. Nov. Gen. et Sp. i. 260 = virginica.
carnea, Schlecht. Ind. Sem. Hort. Halens. (1839) adn. 9 = scabra.
caroliniana, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = stricta.
caroliniana, Walt. Fl. Carol. 68 = nudiflora.
caroliniana, Willd. ex Kunth, Enum. Pl. iv. 54 = angustifolia.
cayennensis, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 105 = nudiflora.
Chamissonis, *Klotzsch, ex C. B. Clarke, in DC. Monog. Phan.* iii. 186.—Austral.
Chantransia, Roem. & Schult. Mant. i. Add. 1. 376 = Floscopa africana.
clandestina, Hort. Berol. (1832) ex Kunth, Enum. Pl. iv. 46 = graminifolia.
clandestina, *Mart. ex Steud. Nom.* ed. II. i. 401.—Bras.
clavata, *C. B. Clarke, Commel. et Cyrt. Beng.* t. 5.—Ind. or.; Malaya.
coelestis, *Willd. Enum. Hort. Berol.* 69.—Mexic.
commelinoides, *Forsk. Fl. Aegypt. Arab.* 12.—Arab.
communis, Engelm. ex Kunth, Enum. Pl. iv. 38 = agraria.
communis, F. Muell. Fragm. viii. 59 = cyanea.
communis, *Linn. Sp. Pl.* 40.—China.
communis, Roxb. Hort. Beng. 5; Fl. Ind. i. 171 = obliqua.
communis, Wall. Cat. n. 8978 = salicifolia, Hasskarlii.
communis, Walt. Fl. Carol. 68 = nudiflora.
comosa, E. Mey. Zwei Pfl. Docum. 174 = Aneilema Dregeanum.
condensata, *C. B. Clarke, in DC. Monog. Phan.* iii. 190.—Afr. trop.
congesta, *C. B. Clarke, l. c.* 160.—Afr. trop.
conspicua, Blume, Enum. Pl. Jav. i. 4 = Aneilema conspicuum.
conspicua, Zoll. Syst. Verz. Ind. Arch. 64 = Aneilema monadelphum.
cordifolia, *A. Rich. Tent. Fl. Abyss.* ii. 341.—Abyss.
crassicaulis, *C. B. Clarke, in DC. Monog. Phan.* iii. 149.—Afr. trop.
crispata, Poir. Encyc. Suppl. ii. 324 = Pollia crispata.
cristata, Linn. Sp. Pl. 42 = Cyanotis cristata.
cucullata, Linn. Mant. ii. 176 = benghalensis.
cyanea, *R. Br. Prod.* 269.—Austral.
cymosa, Blume, Enum. Pl. Jav. i. 4 = Floscopa scandens.
debilis, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = graminifolia.
debilis, Ledeb. in Roem. & Schult. Mant. i. 342 = virginica.
decumbens, E. Mey. in Presl, Rel. Haenk. i. 136 = pallida.
deficiens, *Herb. ex Steud. Nom.* ed. II. i. 401.—Bras.
deficiens, Hook. Bot. Mag. t. 2644 = virginica.
deficiens, Van Houtte, Fl. des Serres, xviii. (1869) t. 1824 = nudiflora.
delicatula, Schlecht. in Gartenfl. (1854) 349. t. 104 = benghalensis.
densiflora, Blume, Enum. Pl. Jav. i. 4 = Floscopa scandens.
diandra, Steud. Nom. ed. II. i. 401 = Ancilema nudiflorum.
dianthifolia, [*Delile, in*] *Red. Lil.* vii. t. 390.—Mexic.

COMMELINA :—

didyma, Steud. Nom. ed. II. i. 401 = Pollia sorzogonensis.
diffusa, Burm. f. Fl. Ind. 18. t. 7. f. 2 = nudiflora.
diffusa, Willd. ex Kunth, Enum. Pl. iv. 39, 40 = salicifolia, barbata.
diffusa, Zoll. ex C. B. Clarke, in DC. Monog. Phan. iii. 171 = clavata.
divaricata, *Vahl, Enum.* ii. 169.—Arab.
Donii, A. Dietr. Sp. Pl. ii. 395 = obliqua.
dubia, [*Delile, in*] *Red. Lil.* t. 359.—Mexic.
dubia, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = coelestis.
dubia, Jacq. Obs. Bot. iii. 9 = Heteranthera graminea.
ebracteata, Ehrenb. ex Schweinf. Beitr. Fl. Aethiop. 210 = Ancilema Ehrenbergii.
Eckloniana, *Kunth, Enum, Pl.* iv. 57.—Austral.
edulis, *A. Rich. Tent. Fl. Abyss.* ii. 341.—Abyss.
edulis, Stokes, Bot. Mat. Med. i. 184 = tuberosa.
Ehrenbergiana, Link, Klotzsch & Otto, Ic. Pl. Rar. 75. t. 30 = scabra.
elata, Vahl, Enum. ii. 178 = Aneilema herbaceum.
elegans, H. B. & K. Nov. Gen. et Sp. i. 259 = virginica.
elliptica, *H. B. & K. l. c.*—Am. trop.
ensifolia, F. Muell. Fragm. viii. 60 = undulata.
ensifolia, *R. Br. Prod.* 269.—Austral.
equinoctialis, Beauv. Fl. Owar. i. 65 = Aneilema aequinoctiale.
erecta, Chapm. Fl. S. U. St. 497 = hirtella.
erecta, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = coelestis.
erecta, *Linn. Sp. Pl.* 41.—Am. bor.
esculenta, Herb. Madr. ex Wall. Cat. sub n. 5208 = Ancilema esculentum.
exilis, Steud. Nom. ed. II. i. 401 = Ancilema nudiflorum.
falcata, Hassk. in Flora, xlvi. (1863) 387 = Forskalaei.
fasciculata, *Ruiz & Pav. Fl. Per.* i. 44. t. 72. f. 6.—Peruv.
filiformis, Steud. Nom. ed. II. i. 401 = Aneilema vaginatum.
firma, Welw. ex C. B. Clarke, in DC. Monog. Phan. iii. 180 = aspera.
flava, Salisb. Prod. 215 = africana.
flexuosa, Welw. ex C. B. Clarke, in DC. Monog. Phan. iii. 258 = Cyanotis nodiflora.
floribunda, H. B. & K. Nov. Gen. et Sp. i. 260 = Tradescantia cumanensis.
foecunda, Hochst. ex C. B. Clarke, in DC. Monog. Phan. iii. 255 = Cyanotis foecunda.
formosa, R. Grah. in Edinb. N. Phil. Journ. (July–Sept. 1830) 366 = nudiflora.
Forskailii, Hochst. ex C. B. Clarke, in DC. Monog. Phan. iii. 173 = Kotschyi.
Forskâlaei, *Vahl, Enum.* ii. 172.—Geront. trop.
Gambiae, *C. B. Clarke, in DC. Monog. Phan.* iii. 146.—Afr. trop.
gelatinosa, *Edgew. in Journ. As. Soc. Beng.* xxi. (1853) 177.—Ind. or.
geniculata, *Ham. Prod. Fl. Ind. Occ.* 4.—Ins. S. Doming.
Gerrardi, *C. B. Clarke, in DC. Monog. Phan.* iii. 183.—Afr. austr.
gigantea, Vahl, Enum. ii. 177 = Aneilema giganteum.
glabra, Baker, in Saund. Refug. Bot. ii. t. 166 = fasciculata.
glabra, *C. B. Clarke, in DC. Monog. Phan.* iii. 163.—Ind. or.
glabra, G. F. W. Mey. Prim. Fl. Esseq. 22 = nudiflora.
gracilis, Hook. Bot. Mag. t. 3047 = fasciculata.
gracilis, Ruiz & Pav. Fl. Per. i. 44. t. 72. f. a = nudiflora.
graminea, Poir. Encyc. Suppl. ii. 324 = Ancilema gramineum.
graminifolia, *H. B. & K. Nov. Gen. et Sp.* i. 258.—Mexic.
graminifolia, Pav. ex C. B. Clarke, in DC. Monog. Phan. iii. 151 = dianthifolia.
grandiflora, *Tenore, Cat. Hort. Bot. Nap.* (1845) 22.—Hab.?
guianensis, Klotzsch, in Schomb. Faun. et Fl. Guian. 1064 = Phaeospherion persicariaefolium.
Hamiltonii, Spreng. Syst. v. 186 = Floscopa scandens.
hamipila, *Wright, in Sauv. Fl. Cub.* 157.—Cuba.
Harcocuskii, *Hort. Pillnitz. ex Tenore, Cat. Ort. Bot. Nap.* (1845) 100.—Hab.?
Hasskarlii, *C. B. Clarke, Commel. et Cyrt. Beng.* t. 5.—Ind. or.
herbacea, Roxb. Hort. Beng. [5]; Fl. Ind. i. 175 = Aneilema herbaceum.

COMMELINA :—

Heudelotii, *C. B. Clarke, in DC. Monog. Phan.* iii. 184.—Afr. trop.
hexandra, Aubl. Pl. Gui. i. 35. t. 12 = Dichorisandra Aubletiana.
hirsuta, R. Br. in Salt, Abyss. App. 62 = benghalensis.
hirsuta, *C. B. Clarke, in DC. Monog. Phan.* iii. 163.—Ind. or.
hirsuta, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 344 = Cyanotis hirsuta.
hirsuta, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 660 = elliptica.
hirsuta, Willd. in Spreng. & Link, Jahrb. i. III. 73 = virginica.
hirtella, *Vahl, Enum.* ii. 166.—Am. bor.
hispida, Ham. ex Spreng. Syst. iv. Cur. Post. 25 = Floscopa scandens.
hispida, *Ruiz & Pav. Fl. Per.* i. 43. t. 73. f. a.—Peruv.
Hookeri, A. Dietr. Sp. Pl. ii. 404 = Aneilema giganteum.
Hookeriana, Spreng. Syst. v. 186 = Aneilema giganteum.
huillensis, *Welw. ex C. B. Clarke, in DC. Monog. Phan.* iii. 175.—Afr. trop.
humilis, Klotzsch, ex C. B. Clarke, l. c. 154 = C. elliptica.
ignorata, *Kunth, Enum. Pl.* iv. 60.—Am. bor.
imberbis, Ehrenb. ex Hassk. in Schweinf. Beitr. Fl. Aethiop. 209 = latifolia.
intermedia, Schlecht. in Linnaea, xxiv. (1851) 655 = coelestis.
involucrosa, *A. Rich. Tent. Fl. Abyss.* ii. 342.—Abyss.
Jamesoni, *C. B. Clarke, in DC. Monog. Phan.* iii. 155.—Ecuador.
japonica, Hort. Berol. (1841) ex Kunth, Enum. iv. 659 = coelestis.
japonica, Thunb. in Trans. Linn. Soc. ii. (1794) 332 = Ancilema japonicum.
karooica, *C. B. Clarke, in DC. Monog. Phan.* iii. 166.—Afr. austr.
Karwinskiana, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 660 = seq.
Karwinskii, Mart. ex Kunth, l. c. = elliptica.
Kirkii, *C. B. Clarke, in DC. Monog. Phan.* iii. 167.—Afr. trop.
Koenigii, Steud. Nom. ed. II. i. 401 = Aneilema herbaceum, Koenigii.
Kotschyi, *Hassk. in Schweinf. Beitr. Fl. Acthiop.* 207.—Aegypt.
Krebsiana, Kunth, Enum. Pl. iv. 40 = africana.
Kurzii, *C. B. Clarke, in Journ. Linn. Soc.* xi. (1871) 444.—Ind. or.
lanceolata, Benth. Fl. Austral. vii. 84 = agrostophylla.
lanceolata, *R. Br. Prod.* 269.—Austral.
lanuginosa, Heyne, ex Wall. Cat. n. 5221 = Aneilema lanuginosum.
latifolia, *Hochst. ex A. Rich. Tent. Fl. Abyss.* ii. 340.—Abyss.
latifolia, Hochst. ex C. B. Clarke, in DC. Monog. Phan. iii. 160, 169 = benghalensis, uncata.
laxa, Poir. Encyc. Suppl. ii. 324 = Aneilema laxum.
leiandra, *C. B. Clarke, in DC. Monog. Phan.* iii. 150.—Mexic.
leiocarpa, Benth. in Bot. Voy. Sulph. 176 = Athyrocarpus leiocarpus.
linearifolia, Kunth, Enum. Pl. iv. 43 = subulata.
linearis, Benth. Pl. Hartw. 27 = dianthifolia.
lineolata, Blume, Enum. Pl. Jav. i. 3 = Aneilema herbaceum.
Livingstoni, *C. B. Clarke, in DC. Monog. Phan.* iii. 190.—Afr. austr.
Loddigesii, Steud. Nom. ed. II. i. 401 = tuberosa.
longicapsa, *C. B. Clarke, in DC. Monog. Phan.* iii. 176.—Afr. occ.
longicaulis, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = coelestis.
longicaulis, Jacq. Coll. iii. 234 = nudiflora.
longifolia, *Lam. Tabl. Encyc.* i. 129.—Java.
longifolia, Michx. Fl. Am. Bor. i. 23 = hirtella.
longifolia, Spreng. Syst. iv. Cur. Post. 25 = Aneilema giganteum.
longifolia, Thw. Enum. Pl. Zeyl. 322 = Kurzii.
Loureiri, *Kunth, Enum. Pl.* iv. 60.—China.
ludens, Miq. in Journ. Bot. Neerl. i. (1861) 88 = communis.
lunata, Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 188 = ensifolia.
lutea, Moench, Meth. 287 = africana.
macrophylla, Poir. Encyc. Suppl. ii. 324 = Pollia macrophylla.

COMMELINA :—
maculata, Edgew. in Trans. Linn. Soc. xx. (1851) 89 = obliqua.
madagascarica, *C. B. Clarke, in DC. Monog. Phan.* iii. 174.—Madag.
Mannii, *C. B. Clarke, l. c.* 167.—Afr. trop.
Martiana, Seub. in Mart. Fl. Bras. iii. 1. 265. t. 37. f. 1 = virginica.
mascarenica, *C. B. Clarke, in DC. Monog. Phan.* iii. 174.—Ins. Maurit.
medica, Lour. Fl. Cochinch. 40 = Aneilema medicum.
mexicana, E. Mey. in Presl, Rel. Haenk. i. 137 = Tradescantia elongata.
micrantha, Vahl, Enum. ii. 178 = Aneilema micranthum.
minuta, Blume, Cat. Gew. Buitenz. 34 = Aneilema nudiflorum.
mollis, Jacq. Coll. iii. 235 = benghalensis.
moluccana, Roxb. Hort. Beng. [81]; Fl. Ind. i. 172 = Cyanotis uniflora.
monadelpha, Blume, Enum. Pl. Jav. i. 4 = Aneilema monadelphum.
montana, Steud. Nom. ed. II. i. 401 = Aneilema montanum.
monticola, *Seub. in Mart. Fl. Bras.* iii. 1. 264.—Bras.
multicaulis, Hochst. ex C. B. Clarke, in DC. Monog. Phan. iii. 184 = albescens.
multiflora, Martens & Gal. in Bull. Acad. Brux. ix. (1842) II. 374 = Callisia Martensiana.
nana, Roxb. Hort. Beng. 5; Fl. Ind. i. 173 = Aneilema spiratum.
nervosa, Burm. f. Fl. Ind. 18 = benghalensis.
nervosa, Ruiz & Pav. Fl. Per. i. 44 = fasciculata.
nigritana, Baker, in Trans. Linn. Soc. xxix. (1875) 163 = subulata.
nigritana, Benth. in Hook. Niger Fl. 541 = umbellata.
nilagirica, Steud. ex C. B. Clarke, in DC. Monog. Phan. iii. 163, 171 = hirsuta, clavata.
Nimmoniana, *J. Grah. Cat. Pl. Bombay*, 224.—Ind. or.
nitens, *Hort. Pillnitz. ex Tenore, Cat. Ort. Bot. Nap.* (1845) 100.—Hab.?
nudicaulis, Burm. f. Fl. Ind. 17. t. 8. f. 1 = Aneilema nudiflorum.
nudiflora, Linn. Mant. ii. 177 = Aneilema nudiflorum.
nudiflora, *Linn. Sp. Pl.* 4.—Reg. trop.
obliqua, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 45.—Ind. or.; Malaya.
obliqua, Vahl, Enum. ii. 173 = erecta.
oblongifolia, Hassk. Commel. Ind. 27 = auriculata.
obtusifolia, Vahl, Enum. ii. 168 = nudiflora.
ochreata, Schau. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 447 = nudiflora.
oligotricha, *Miq. Fl. Ind. Bat.* iii. 532.—Malaya.
orchioides, Booth, ex Lindl. Bot. Reg. (1838) Misc. 53 = elliptica.
orchioides, *Roth, ex Steud. Nom.* ed. II. i. 402.—Mexic.
ovata, Steud. l. c. = Aneilema ovatum.
ovato-oblonga, Roem. & Schult. Mant. i. Add. 1. 376 = Aneilema ovato-oblongum.
pacifica, Vahl, Enum. ii. 168 = nudiflora.
paleata, *Hassk. in Miq. Pl. Jungh.* 139.—Ind. or.; Malaya.
Palisoti, A. Dietr. Sp. Pl. ii. 414 = Aneilema ovato-oblongum.
pallida, *Willd. Hort. Berol.* ii. 87. t. 87.—Mexic.
pallida, Schlecht. in Linnaea, xxvi. (1853-55) 454 = Athyrocarpus leiocarpus.
paludosa, Blume, Enum. Pl. Jav. i. 2 = obliqua.
paludosa, Steud. Nom. ed. II. i. 402 = persicariaefolia.
paniculata, *Hill, Veg. Syst.* xxiii. 4.—Ind. or.
paniculata, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = coelestis.
paniculata, Soland. ex C. B. Clarke, in DC. Monog. Phan. iii. 223 = Aneilema laxum.
paniculata, Vahl, Enum. ii. 179 = Aneilema Forskalei
papilionacea, Burm. f. Fl. Ind. 17 = Cyanotis papilionacea.
parviflora, Link, Enum. Hort. Berol. i. 61 = graminifolia.
parviflora, Reichb. Fl. Exot. ii. 17. t. 142 = tuberosa.
pedunculosa, Spreng. & Link, Jahrb. i. III. 74 = salicifolia.
persicariaefolia, [Delile, in] Red. Lil. t. 472 = Athyrocarpus persicariaefolius.
persicariaefolia, *Wight, in Wall. Cat.* n. 8984.—Burma.
Petersii, *Hassk. in Flora*, xlvi. (1863) 385.—Afr. trop.

COMMELINA :—
phalangoides, *Crantz, Inst.* i. 459.—Hab.?
pilosa, Pers. Syn. i. 55 = nudiflora.
pilosa, Steud. Nom. ed. II. i. 402 = Aneilema ochraceum.
pilosula, Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 106 = nudiflora.
platyphylla, *Klotzsch, in Schomb. Faun. et Fl. Guiana*, 897.—Guiana.
Pohliana, Seub. in Mart. Fl. Bras. iii. 1. 265. t. 37. f. 1 = erecta.
poligama, Blanco, Fl. Filip. ed. Naves (Ic. sine num.) = benghalensis.
polyclada, Welw. ex C. B. Clarke, in DC. Monog. Phan. iii. 165 = africana.
polygama, Roth, in Roem. & Ust. Mag. iv. x. (1790) 14 = communis.
polyspatha, Wight, Ic. t. 2066 = obliqua.
procurrens, Schlecht. in Linnaea, xxv. (1852) 183 = benghalensis.
prostrata, H. B. & K. Nov. Gen. et Sp. i. 259 = nudiflora.
prostrata, Poepp. ex Kunth, Enum. Pl. iv. 38 = nudiflora.
prostrata, Regel, Gartenfl. (1868) 289. t. 592. f. 1 = benghalensis.
protensa, Steud. Nom. ed. II. i. 402 = Aneilema protensum.
puberula, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv 659; et in Linnaea, xxiv. (1851) 197 = dianthifolia.
pubescens, Klotzsch, ex C. B. Clarke, in DC. Monog. Phan. iii. 152 = acuminata.
pubinervis, *Hassk. in Flora*, xxv. (1842) II. *Beibl.* 4.—Java.
pumila, Royle, ex C. B. Clarke, in DC. Monog. Phan. iii. 207 = Aneilema spiratum.
pyrrho-blepharis, *Hassk. in Schweinf. Beitr. Fl. Aethiop.* 209.—Abyssin.
quitensis, *Benth. Pl. Hartw.* 258.—Ecuador.
radicans, Spreng. Syst. iv. Cur. Post. 25 = Aneilema nudiflorum.
radiciflora, R. Br. ex C. B. Clarke, in DC. Monog. Phan. iii. 159 = benghalensis.
Rafinesquei, *Kunth, Enum. Pl.* iv. 60.—Am. bor.
rajmahlensis, C. B. Clarke, in Journ. Linn. Soc. xi. (1871) 444 = attenuata.
rara, *Roxb. ex Steud. Nom.* ed. II. i. 402.—Ind. or.
reniformis, Steud. l. c. = Pollia subumbellata.
repens, Roxb. & Heyne, ex C. B. Clarke, in DC. Monog. Phan. iii. 172 = attenuata.
rhizocarpa, Afzel. ex C. B. Clarke, l. c. 159 = benghalensis.
robusta, *Kunth, Enum. Pl.* iv. 52.—Bras.
rorigera, *Fisch. ex Tenore, Cat. Ort. Bot. Nap.* (1845) 100.—Hab.?
rosea, *Schlecht. Ind. Sem. Hort. Hal.* (1838) 7.—Mexic.
rubens, Hort. Berol. (1841) ex Kunth, Enum. Pl. iv. 659 = graminifolia.
rubens, [Delile, in] Red. Lil. vii. t. 367 = pallida.
rufipes, Seub. in Mart. Fl. Bras. iii. 1. 265 = Phaeospherion persicariaefolium.
rugulosa, C. B. Clarke, in Journ. Linn. Soc. xi. (1871) 446 = suffruticosa.
Rumphii, *Kostel. Allg. Med.-Pharm. Fl.* i. 126.—As. trop.
Sabatieri, *C. B. Clarke, in DC. Monog. Phan.* iii. 146.—Afr. trop.
sagittifolia, Hassk. in Schweinf. Beitr. Fl. Aethiop. 206 = latifolia.
salicifolia, Boj. Hort. Maurit. 360 = communis.
salicifolia, *Roxb. Hort. Beng.* 5; *Fl. Ind.* i. 172.—Ind. or.
salicifolia, Thw. Enum. Pl. Zeyl. 321 = clavata.
Saltiana, Steud. Nom. ed. II. i. 402 = pyrrho-blepharis.
scaberrima, Blume, Enum. Pl. Jav. i. 4 = Aneilema protensum.
scabra, *Benth. Pl. Hartw.* 26.—Mexic.
scabrata, Seub. in Mart. Fl. Bras. iii. 1. 266 = Phaeospherion persicariaefolium.
scandens, *Welw. ex C. B. Clarke, in DC. Monog. Phan.* iii. 146.—Afr. trop.
scapiflora, Roxb. Hort. Beng. [5]; Fl. Ind. i. 175 = Aneilema scapiflorum.
scapigera, Hort. Berol. (1832) ex Kunth, Enum. Pl. iv. 46 = elliptica.
Schimperiana, Hochst. ex C. B. Clarke, in DC. Monog. Phan. iii. 184 = albescens.
Schomburgkiana, Klotzsch, in Schomb. Faun. et Fl. Guiana, 1064 = nudiflora.

COMMELINA :—
Schoubertii, Zucc. ex Tenore, Cat. Hort. Bot. Nap. (1845) 22 = hispida.
Schweinfurthii, *C. B. Clarke, in DC. Monog. Phan.* iii. 158.—Afr. trop.
secundiflora, Blume, Enum. Pl. Jav. i. 3 = Pollia sorzogonensis.
Sellowiana, Kunth, Enum. Pl. iv. 44 = nudiflora.
Sellowii, Schlecht. Ind. Sem. Hort. Hal. (1850) 7 = nudiflora.
semi-ovata, Buch.-Ham. in Wall. Cat. n. 8985 = obliqua, suffruticosa.
senegalensis, Tenore, Cat. Ort. Bot. Nap. (1845) 22 = benghalensis.
serrulata, *Vahl, Eclog. Am.* ii. 4.—Guiana.
setosa, Wight, in Wall. Cat. n. 8981 = undulata.
sikkimensis, *C. B. Clarke, Commel. & Cyrt. Beng.* t. 6.—Reg. Himal.
siliculosa, Poir. Encyc. Suppl. ii. 324 = Aneilema siliculosum.
simplex, Vahl, Enum. ii. 177 = Ancilema sinicum.
Simsoni, C. B. Clarke, in Journ. Linn. Soc. xi. (1871) 446 = suffruticosa.
singularis, *Vell. Fl. Flum.* 31; i. t. 76.—Bras.
sinica, Roem. & Schult. Mant. i. App. 1. 376 = Aneilema sinicum.
speciosa, Thunb. Prod. Pl. Cap. 58 = Cyanotis nodiflora.
spectabilis, *C. B. Clarke, in DC. Monog. Phan.* iii. 175.—Afr. trop.
spicata, Steud. Nom. ed. II. i. 402 = Aneilema Loureirii.
spirata, Koen. ex Steud. l. c. = Aneilema herbaceum.
spirata, Linn. Mant. ii. 176 = Aneilema spiratum.
striata, Edgew. in Trans. Linn. Soc. xx. (1851) 89 = obliqua.
striata, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 340 = subulata.
striata, Wall. Cat. n. 8981 = albescens, Kurzii, ensifolia.
stricta, Desf. Cat. Hort. Par. ed. III. 388 = graminifolia.
subamplectens, Hassk. Commel. Ind. 27 = africana.
subaurantiaca, Hochst. in Kunth, Enum. Pl. iv. 658 = subulata.
subulata, *Roth, Nov. Pl. Sp.* 23.—Geront. trop.
suffruticosa, *Blume, Enum. Pl. Jav.* i. 3.—Java.
sulcata, Benth. in Hook. Niger Fl. 541 = Vogelii.
sulcata, Hoffmgg. in Spreng. & Link, Jahrb. i. III. 74 = virginica.
tenuis, Roth, Nov. Pl. Sp. 22 = Aneilema acuminatum.
tinctoria, Nees, in Linnaea, xix. (1847) 699 = dianthifolia.
triquetra, Steud. Nom. ed. II. i. 402 = Aneilema triquetrum.
truncata, Willd. Enum. Hort. Berol. Suppl. 4 = virginica.
tuberosa, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 110 = virginica.
tuberosa, Burm. ex Steud. Nom. ed. II. i. 402 = salicifolia.
tuberosa, [DC. in] Red. Lil. ii. t. 108 = coelestis.
tuberosa, Forsk. Fl. Aegypt. Arab. 12 = Aneilema Forskalei.
tuberosa, *Linn. Sp. Pl.* 41.—Mexic.
tuberosa, Lour. Fl. Cochinch. 40 = Aneilema Loureirii.
turbinata, Vahl, Enum. ii. 171 = benghalensis.
umbellata, *Schum. & Thonn. Beskr. Guin. Pl.* 21.—Afr. trop.
umbrosa, Vahl, Enum. ii. 179 = Aneilema ovato-oblongum.
uncata, *C. B. Clarke, in DC. Monog. Phan.* iii. 169.—Abyssin.
undulata, Benth. Fl. Austral. vii. 83 = Chamissonis.
undulata, Lodd. Bot. Cab. t. 1553 = tuberosa.
undulata, *R. Br. Prod.* 270.—Austral.
vaginata, Heyne, ex Steud. Nom. ed. II. i. 402 = Aneilema esculentum.
vaginata, Linn. Mant. 177 = Aneilema vaginatum.
vaginata, Sonn. ex C. B. Clarke, in DC. Monog. Phan. iii. 188 = ensifolia.
variabilis, Schlecht. Ind. Sem. Hort. Hal. (1838) 7 = elliptica.
vestita, Seub. in Mart. Fl. Bras. iii. 1. 264 = monticola.
villosiuscula, Soland. ex C. B. Clarke, in DC. Monog. Phan. iii. 159 = benghalensis.
virginica, Forst. f. Prod. 6 = nudiflora.
virginica, *Linn. Sp. Pl.* ed. II. 61.—Am. bor.
Vogelii, *C. B. Clarke, in DC. Monog. Phan.* iii. 189.—Afr. trop.

COMMELINA:—
vulgaris, [DC. in] Red. Lil. iv. t. 206 = communis.
Wallichiana, Steud. Nom. ed. II. i. 402 = Aneilema divergens.
Welwitschii, *C. B. Clarke, in DC. Monog. Phan.* iii. 175.—Afr. trop.
Werneana, Hassk. in Schweinf. Beitr. Fl. Aethiop. 206 = nudiflora.
Willdenowii, Kunth, Enum. Pl. iv. 37 = communis.
zambesica, *C. B. Clarke, in DC. Monog. Phan.* iii. 161.—Afr. trop.
Zanonia, Linn. Sp. Pl. 41 = Campelia Zanonia.
Zebrina, Hort. ex C. B. Clarke, in DC. Monog. Phan. iii. 318 = Zebrina pendula.
zeylanica, *Falkenberg, in Just, Jahresb.* iv. (1876) 406.—Zeylan.

COMMELYNA, Hoffmgg. ex Endl. Gen. 125 (1837) = **Commelina**, Linn.

COMMERÇONIA, F. Muell. Census, 16 (1882) = **Commersonia**, Forst. (Stercul.).

COMMERSONA, Sonner. Voy. t. 89 (1782) = **Barringtonia**, Forst. (Myrtac.).
speciosa, Salisb. Prod. 355 = B. Butonica.

COMMERSONIA, Comm. ex DC. Prod. ii. 10, in syn. (1825) = **Polycardia**, Juss. (Celastrin.).

COMMERSONIA, Forst. Char. Gen. 43. t. 22 (1776). *STERCULIACEAE*, Benth. & Hook. f. i. 226.
aspera, G. Don, Gen. Syst. i. 523 = Buettneria aspera.
cinerea, Steud. in Lehm. Pl. Preiss. i. 238 = Rulingia grandiflora.
craurophylla, F. Muell. Fragm. ix. 59 = Rulingia craurophylla.
crispa, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) i. 501.—Austral.
cuneata, F. Muell. Census, 16 = Rulingia cuneata.
cygnorum, Steud. in Lehm. Pl. Preiss. i. 237 = Rulingia malvaefolia.
dasyphylla, Andr. Bot. Rep. t. 603 = Rulingia pannosa.
dasyphylla, Hort. ex Steud. Nom. ed. II. i. 402 = Rulingia pannosa.
densiflora, F. Muell. Fragm. x. 21 = Rulingia densiflora.
diphylla, Andr. Bot. Rep. t. 603 = Rulingia pannosa.
echinata, Blume, Bijdr. 86 = platyphylla.
echinata, *Forst. Char. Gen.* 43. t. 22.—Malaya.
Fraseri, *J. Gay, in Mém. Mus. Par.* x. (1823) 215. t. 15.—Austral.
Fraseri, Sieber, ex Steetz, in Lehm. Pl. Preiss. ii. 352 = Rulingia pannosa.
Gaudichaudii, *J. Gay, in Mém. Mus. Par.* x. (1823) 213. t. 14.—Austral.
herbacea, G. Don, Gen. Syst. i. 523 = Buettneria herbacea.
hermanniaefolia, *J. Gay, in H. B. & K. Nov. Gen. et Sp.* v. 311, in nota.—Hab.?
hermanniaefolia, F. Muell. Census, 16 = Rulingia hermanniaefolia.
Jackiana, G. Don, Gen. Syst. i. 523 = Buettneria Jackiana.
javensis, G. Don, l. c. = platyphylla.
Kempeana, F. Muell. Fragm. xi. 113 = Rulingia Kempeana.
laxophylla, F. Muell. l. c. x. 22 = Rulingia laxophylla.
Leichardtii, *Benth. Fl. Austr.* i. 242.—Austral.
magniflora, F. Muell. Fragm. viii. 223 = Rulingia magniflora.
melanopetala, *F. Muell. l. c.* x. 21.—Austral.
microphylla, *Benth. Fl. Austral.* i. 244.—Austral.
parviflora, F. Muell. Census, 16 = Rulingia parviflora.
pilosa, G. Don, Gen. Syst. ii. 524 = Buettneria pilosa.
platycalyx, F. Muell. Census, 16 = Rulingia platycalyx.
platyphylla, *Andr. Bot. Rep.* t. 603 adn. et t. 519.—Malaya.
Preissii, Steud. in Lehm. Pl. Preiss. i. 238 = Rulingia corylifolia.
pulchella, *Turcz. in Bull. Soc. Nat. Mosc.* xix. (1846) i. 502.—Austral.
rotundifolia, F. Muell. Census, 16 = Rulingia rotundifolia.
rugosa, F. Muell. l. c. = Rulingia rugosa.
salvifolia, F. Muell. l. c. = Rulingia salvifolia.

COMMIA, Lour. Fl. Cochin. 605 (1790) = **Excoecaria**, Linn. (Euphorb.).
cochinchinensis, Lour. l. c. 606 = E. Agallocha.

COMMIANTHUS, Benth. in Hook. Journ. Bot. iii. (1841) 223 = **Retiniphyllum**, Humb. & Bonpl. (Rubiac.).
concolor, Spruce, ex Benth. in Hook. Kew Journ. v. (1853) 235 = R. concolor.
discolor, Spruce, ex Benth. l. c. 234 = R. discolor.
pilosus, Spruce, ex Benth. l. c. 233 = R. pilosum.
Schomburgkii, Benth. in Hook. Journ. Bot. iii. (1841) 223 = R. Schomburgkii.
speciosus, Spruce, ex Benth. in Hook. Kew Journ. v. (1853) 234 = R. speciosum.

COMMIDENDRON, Lem. in Orb. Dict. iv. 147 (1849) = seq.

COMMIDENDRUM, Burch. ex DC. in Guill. Archiv. Bot. ii. (1833) 334. *COMPOSITAE*, Benth. & Hook. ii. 277.
Burchellii, *Benth. & Hook. f. ex Hemsl. Bot. Chall. Voy.* i. II. 71.—Ins. S. Helena.
gummiferum, DC. Prod. v. 344 = robustum.
robustum, *DC. l. c.*—Ins. S. Helena.
rotundifolium, DC. l. c. = Psiadia rotundifolia.
rugosum, *DC. l. c.* 345.—Ins. S. Helena.
spurium, *DC. l. c.* 344.—Ins. S. Helena.

COMMILOBIUM, Benth. in Ann. Wien. Mus. ii. (1838) 110 = **Pterodon**, Vog. (Legumin.).
abruptum, Moric. Pl. Rar. Nouv. Am. 98. t. 62 = P. abruptus.
polygalaeflorum, Benth. in Ann. Wien. Mus. ii. (1838) 111 = P. polygalaeflorus.
pubescens, Benth. l. c. = P. pubescens.

COMMIPHORA, Jacq. Hort. Schoenb. ii. 66 (1797). *BURSERACEAE*.
BALSAMEA, Gled. in Berl. Ges. Naturf. Fr. Schr. iii. (1782) 127.
BALSAMODENDRON, Kunth, in Ann. Sc. Nat. Sér. I. ii. (1824) 348.
BALSAMOPHLOEOS, O. Berg, in Bot. Zeit. xx. (1862) 163.
BALSAMUS, Stackh. De Liban. 11 (1814).
HEMPRICHIA, Ehrenb. in Linnaea, iv. (1829) 396.
HEUDELOTIA, A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 150. t. 39 (1832).
HITZERIA, Klotzsch, in Peters, Reise Mossamb. Bot. 89 (1861).
NIOTOUTT, Adans. Voy. 162 (1759).
PROTIONOPSIS, Blume, Mus. Bot. Lugd. Bat. i. 229 (1850).
PROTIUM, Wight & Arn. Prod. 176 (1834).
abyssinica, *Engl. in DC. Monog. Phan.* iv. 10.—Abyss.
africanum, *Endl. l. c.* 14.—Abyss.
Agallocha, *Engl. l. c.* 11.—Ind. or.
angolensis, *Engl. l. c.* 24.—Afr. trop.
Berryi, *Engl. l. c.* 17.—Ind. or.
Boiviniana, *Engl. l. c.* 21.—Afr. trop.
capensis, *Engl. l. c.* 18.—Afr. austr.
caudata, *Engl. l. c.* 27.—Ind. or.
edulis, *Engl. l. c.* 22.—Afr. trop.
erythraea, *Engl. l. c.* 20.—Afr. trop.
grandifolia, *Engl. l. c.* 25.—Madag.
Harveyi, *Engl. l. c.*—Afr. austr.
Hildebrandtii, *Engl. l. c.* 15.—Afr. trop.
Kataf, *Engl. l. c.* 19.—Arab.
longebracteata, *Engl. l. c.*—Afr. trop.
madagascariensis, *Jacq. Hort. Schoenb.* ii. 66.—Madag.
Marchandi, *Engl. in DC. Monog. Phan.* iv. 26.—Madag.
mollis, *Engl. l. c.* 23.—Afr. trop.
mossambicensis, *Engl. l. c.* 26.—Afr. trop.
Mukul, *Engl. l. c.* 12.—Ind. or.
Myrrha, *Engl. l. c.* 10.—Arab.
Olivieri, *Engl. l. c.* 24.—Afr. trop.
Opobalsamum, *Engl. l. c.* 15.—Arab.
orbicularis, *Engl. l. c.* 18.—Madag.
parvifolia, *Engl. l. c.* 536.—Ins. Socotra.
pedunculata, *Engl. l. c.* 23.—Africa.
Pervilleana, *Engl. l. c.* 29.—Madag.
pilosa, *Engl. l. c.* 12.—Afr. trop.
planifrons, *Engl. l. c.* 536.—Ins. Socotra
pubescens, *Engl. l. c.* 21.—Ind. or.
Rehmanni, *Engl. l. c.* 15.—Afr. austr.
Schimperi, *Engl. l. c.* 13.—Abyssin.

COMMIPHORA:—
serrata, *Engl. in DC. Monog. Phan.* iv. 24.—Afr. trop.
socotrana, *Engl. l. c.* 536.—Ins. Socotra.
Stocksiana, *Engl. l. c.* 17.—Beluchist.
tetramera, *Engl. l. c.* 27.—Madag.
Welwitschii, *Engl. l. c.* 22.—Afr. trop.
zanzibarica, *Engl. l. c.* 28.—Afr. trop.

COMMIRHOEA, Miers, in Trans. Linn. Soc. xxi. (1855) 252 = **Chrysochlamys**, Poepp.? (Guttif.).
mecocarpa, Miers, l. c. 258.—Hab.?

COMOCLADIA, P. Br. Hist. Jamaic 124 (1756). *ANACARDIACEAE*, Benth. & Hook. f. i. 419.
DODONAEA, Plum. ex Adans. Fam. ii. 342 (1763).
acuminata, *Moç. & Sesse, ex DC. Prod.* ii. 65.—Mexic.
angulosa, Willd. Sp. Pl. i. 188 = ilicifolia.
Brasiliastrum, *Poir. Encyc. Suppl.* ii. 325.—Am. austr.
dentata, *Jacq. Enum. Pl. Carib.* 12.—Ind. occ.
Ehrenbergii, *Engl. in Bot. Jahrb.* i. (1881) 420.—Ins. S. Doming.
glabra, *Spreng. Syst.* i. 176.—Cuba.
ilicifolia, *Sw. Prod. Veg. Ind. Occ.* 17.—Ind. occ.
integrifolia, *Jacq. Enum. Pl. Carib.* 12.—Am. trop.
intermedia, *Wright, ex Engl. in DC. Monog. Phan.* iv. 366.—Cuba.
loxensis, *H. B. & K. Nov. Gen. et Sp.* vii. 16.—N. Granat.
mollissima, *H. B. & K. l. c.* 16. t. 607.—Mexic.
pinnatifida, Linn. Syst. ed. X. 861 = integrifolia.
platyphylla, *A. Rich. ex Griseb. Cat. Pl. Cub.* 68.—Cuba.
propinqua, *H. B. & K. Nov. Gen. et Sp.* vii. 16.—Cuba.
pubescens, *Engl. in Bot. Jahrb.* i. (1881) 420.—Jamaic.
serrata, Blanco, Fl. Filip. ed. I. 30 = Salacia oblonga.
Tapaculo, H. B. & K. Nov. Gen. et Sp. vii. 16 = Tapiria guianensis.
tomentosa, Willd. ex Schult. Mant. i. 350 = mollissima.
tricuspidata, Lam. in Mém. Acad. Sc. Par. 1784 (1787) 347 = ilicifolia.

COMOLIA, DC. Prod. iii. 114 (1828). *MELASTOMACEAE*, Benth. & Hook. f. i. 741.
HOSTMANNIA, Steud. ex Naud. in Ann. Sc. Nat. Sér. III. xiii. (1849) 25.
LEIOSTEGIA, Benth. in Hook. Journ. Bot. ii. (1840) 294.
PACHYLOMA, DC. Prod. iii. 120 (1828).
TETRAMERIS, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850) 120. t. 4.
TRICENTRUM, DC. Prod. iii. 123 (1828).
amazonica, *Cogn. in Mart. Fl. Bras.* xiv. III. 422.—Bras.
Aubletii, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 37.—Guiana.
berberifolia, *DC. Prod.* iii. 115.—Bras.
denudata, *Naud, in Ann. Sc. Nat.* Sér. III. xiii. (1850) 25.—Bras.
hirtella, *Naud. l. c.* 26 = microphylla.
lanceaeflora, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 37.—Bras.
latifolia, Cogn. in Mart. Fl. Bras. xiv. III. 433 = Aubletii.
leptophylla, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1850) 27.—Guiana.
lythrarioides, *Naud. l. c.*—Guiana.
microphylla, *Benth. in Hook. Journ. Bot.* ii. (1840) 295.—Guiana.
neglecta, *Cogn. in Mart. Fl. Bras.* xiv. III. 423.—Guiana.
numnularioides, *Naud. in Ann. Sc. Nat.* Sér. III. xiii. (1850) 27.—Bras.
ovalifolia, Triana, in Trans. Linn. Soc. xxviii. (1871) 37. t. 2 = denudata.
purpurea, Miq. in Linnaea, xviii. (1844) 617 = veronicaefolia.
Sertularia, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 37. t. 2. f. 27 f.—Bras.
sessilis, *Triana, l. c.* f. 27 e.—Bras.
stenodon, *Triana, l. c.* 37.—Bras.
surinamensis, Miq. Surp. Surin. 55 = lythrarioides.
tetraptera, *Cogn. in Mart. Fl. Bras.* xiv. III. 426.—Bras.

COMOLIA:—
vernicosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 37. t. 2. f. 27 g.—Bras.
veronicaefolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 295.—Guiana.
veronicaefolia, Triana, in Trans. Linn. Soc. xxviii. (1871) 36 = purpurea.
villosa, *Triana, l. c.* 37.—Guiana.
violacea, *Triana, l. c.* —Bras.

COMOMYRSINE, Hook. f. in Benth. & Hook. f. Gen. ii. 643 (1876). *MYRSINEAE*, Benth. & Hook. f. l. c.
Schlimii, *Hook. f. l. c.* 644.—Am. austr.
simplex, *Hook. f. l. c.*—Am. austr.
Sprucei, *Hook. f. l. c.*—Am. austr.

COMOSPERMA, Poir. in Dict. Sc. Nat. x. (1818) 128 = **Comesperma**, Labill. (Polygal.).

COMOSTEMUM, Nees, in Linnaea, ix. (1834) 283 = **Androtrichum**, Brongn. (Cyperac.).
globuliferum, Nees, in Mart. Fl. Bras. ii. 1. 5 = A. polycephalum.
impolitum, Nees, in Hook. Journ. Bot. ii. (1840) 397 = Cyperus impolitus.
laetum, Nees, l. c. = Cyperus laetus, *Presl.*
montevidense, Nees, in Linnaea, ix. (1834) 283 = A. polycephalum.
prolixum, Nees, in Hook. Journ. Bot. ii. (1840) 397 = Cyperus prolixus.
Schottii, Nees, in Linnaea, ix. (1834) 283 = Cyperus prolixus.

COMPARETTIA, Poepp. & Endl. Nov. Gen. et Sp. i. 42. t. 73 (1835). *ORCHIDEAE*, Benth. & Hook. f. iii. 558.
PLECTROPHORA, H. C. Focke, in Tijdschr. Natuurk. Wetensch. i. (1848) 212.
coccinea, *Lindl. Bot. Reg.* xxiv. (1838) t. 68.—Bras.
cryptocera, *Morr. in Belg. Hortic.* ii. (1852) 310.—Hab.?
falcata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 42. t. 73.—Peruv.
macroplectron, *Reichb. f. & Triana, in Gard. Chron.* (1878) ii. 524; (1879) i. 398.—N. Granat.
rosea, Lindl. Bot. Reg. (1840) Misc. 78 = falcata.
saccata, *Poepp. & Endl. Nov. Gen. et Sp.* i. 42. t. 72.—Peruv.
speciosa, *Reichb. f. in Gard. Chron.* (1878) ii. 524.—Ecuador.

COMPERIA, C. Koch, in Linnaea, xxii. (1849) 287 = **Orchis**, Linn.
taurica, C. Koch, l. c. 288 = O. Comperiana.

COMPHOROPSIS, Moq. in DC. Prod. xiii. ii. 200, in syn. (1849) = CAMPHOROPSIS, Moq. = **Nanophytum**, Less. (Chenopod.).

COMPHRENA, Aubl. Pl. Gui. i. 280 (1775), sphalm. = **Gomphrena**, Linn. (Amarant.).

COMPSANTHUS, Spreng. Syst. iv. Cur. Post. 137 (1827) = **Tricyrtis**, Wall. (Liliac.).
maculatus, Spreng. l. c. = T. pilosa.

COMPSOA, D. Don, Prod. Fl. Nep. 50 (1825) = **Tricyrtis**, Wall. (1826) (Liliac.).
maculata, D. Don, l. c. 51 = T. pilosa.

COMPTONIA, Banks, ex Gaertn. Fruct. ii. 58. t. 90 (1791) = **Myrica**, Linn.
asplenifolia, Banks, l. c. = M. asplenifolia.
Ceterach, Mirb. in Duham. Arb. ed. nov. ii. t. 11 = M. asplenifolia.

COMUS, Salisb. Gen. Pl. Fragm. 24 (1866) = **Muscari**, Tourn. (Liliac.).
carnosus, Salisb. l. c. 25 = M. comosnm.

CONAMI, Aubl. Pl. Gui. ii. 926. t. 354 (1775) = **Phyllanthus**, Linn. (Euphorb.).
brasiliensis, Aubl. l. c. 927 = P. Conami.

CONANDRON, Sieb. & Zucc. in Abh. Akad. Wiss. Muench. iii. iii. (1843) 729. t. 3. *GESNERACEAE*, Benth. & Hook. f. ii. 1024.
ramondioides, *Sieb. & Zucc. l. c.*—Japon.

CONANTHERA, Ruiz & Pav. Fl. Per. iii. 68. t. 301 (1802). *HAEMODORACEAE*, Benth. & Hook. f. iii. 679.
CUMMINGIA, D. Don, in Sweet, Brit. Flow. Gard. Ser. I. iii. t. 257 (1828).
albiflora, Cham. & Schlecht. in Linnaea, vi. (1831) 50 = Echeandia terniflora.
bifolia, *Ruiz & Pav. Fl. Per.* iii. 68. t. 301.—Chili.
bifolia, Sims, in Bot. Mag. t. 2496 = Simsii.
campanulata, Lindl. in Trans. Hort. Soc. vi. (1826) 283 = Simsii.
Echeandia, Pers. Syn. i. 370 = Echeandia terniflora.
Forsteri, *Spreng. Syst.* ii. 91.—N. Caled.
Simsii, *Sweet, Hort. Brit.* ed. I. 423.—Chili.

CONANTHODIUM, A. Gray, in Hook. Kew Journ. iv. (1852) 272 = **Helichrysum**, Vaill. (Compos.).
Drummondii, A. Gray, l. c. = H. argyroglottis.

CONANTHUS, S. Wats. Bot. King's Exped. 256 (1871). *HYDROPHYLLACEAE*, Benth. & Hook. f. ii. 828.
aretioides, *S. Wats. l. c.*—Am. bor. occ.

CONCEVEIBA, Aubl. Pl. Gui. ii. 923. t. 353 (1775). *EUPHORBIACEAE*, Benth. & Hook. f. iii. 316.
africana, *Muell. Arg. in Flora*, xxxiii. (1864–65) 530.—Afr. trop.
cordata, A. Juss. Tent. Euph. 43. t. 13 = Alchornea cordata.
guianensis, *Aubl. Pl. Gui.* ii. 924. t. 353.—Guiana.
Hostmani, *Benth. in Hook. Kew Journ.* vi. (1854) 332.—Guiana.
javanensis, Blume, Bijdr. 614 = Alchornea rugosa.
latifolia, Zipp. ex Span. in Linnaea, xv. (1841) 349 = Alchornea rugosa.
latifolia, *Benth. in Hook. Kew Journ.* vi. (1854) 332.—Bras.
macrophylla, Klotzsch, ex Benth. l. c. 333 = Alchornea cordata.
Martiana, *Baill. in Adansonia*, v. (1864–65) 221.—Bras.
megalophylla, Muell. Arg. in Linnaea, xxxiv. (1865–66) 167 = Alchornea Martiana.
ovata, Rich. ex A. Juss. Tent. Euph. 43 = guyanensis.
rhytidocarpa, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 372.—Peruv.
terminalis, *Muell. Arg. in Linnaea*, xxxiv. (1865–66) 167.—Bras.
tomentosa, Span. l. c. xv. (1841) 349 = Cladogynos orientalis.
trigonocarpa, *Muell. Arg. in Mart. Fl. Bras.* xi. ii. 372.—Bras.

CONCEVEIBUM, Rich. ex A. Juss. Tent. Euph. 42 (1824) = praec.

CONCHIDIUM, Griff. Notul. iii. 321. t. 310 (1851) = **Eria**, Lindl. (Orchid.).
pusillum, Griff. l. c. = E. pusilla.
sinicum, Lindl. ex Benth. in Hook. Kew Journ. vii. (1855) 34 = E. sinica.

CONCHIUM, Sm. in Trans. Linn. Soc. iv. (1798) 215 = **Hakea**, Schrad. (Proteac.).
aciculare, Sm. ex Vent. Jard. Malm. t. 111 = H. acicularis.
ceratophyllum, Sm. in Trans. Linn. Soc. ix. (1808) 124 = H. ceratophylla.
clavatum, Willd. Enum. Hort. Berol. 141 = H. clavata.
compressum, Sm. in Trans. Linn. Soc. ix. (1808) 121 = H. acicularis.
corniculatum, Sm. ex Willd. Enum. Hort. Berol. 141 = H. pugioniformis.
cornutum, Gaertn. f. Fruct. iii. 216. t. 219 = H. gibbosa.
dactyloides, Vent. Jard. Malm. t. 110 = H. dactyloides.
drupaceum, Gaertn. f. Fruct. iii. 217. t. 219 = H. leucoptera.
ellipticum, Sm. in Trans. Linn. Soc. ix. (1808) 123 = H. elliptica.

CONCHIUM:—
epiglottis, Willd. Enum. Hort. Berol. 141 = H. epiglottis.
gibbosum, Sm. in Trans. Linn. Soc. ix. (1808) 119 = H. gibbosa.
longifolium, Sm. l. c. 121 = H. pugioniformis.
mucronatum, Cels, ex Spreng. Syst. i. 480 = H. varia.
nervosum, Sm. ex Willd. Enum. Hort. Berol. 141 = H. dactyloides.
oleifolium, Sm. in Trans. Linn. Soc. ix. (1808) 124 = H. oleifolia.
pubescens, Willd. Enum. Hort. Berol. 141 = H. gibbosa.
pugioniforme, Sm. in Trans. Linn. Soc. ix. (1808) 122 = H. pugioniformis.
pyriforme, Willd. Enum. Hort. Berol. 141 = Xylomelum pyriforme.
ruscifolium, Willd. l. c. = H. ruscifolia.
salicifolium, Gaertn. f. Fruct. iii. 217 = H. saligna.
salignum, Sm. in Trans. Linn. Soc. ix. (1808) 124 = H. saligna.
sphaeroideum, Sm. l. c. 120 = H. gibbosa.
teretifolium, Gaertn. f. Fruct. iii. 217. t. 219 = E. epiglottis.
trifurcatum, Sm. in Trans. Linn. Soc. ix. (1808) 122 = H. trifurcata.

CONCHOCARPUS, Mikan, Delect. fasc. i. t. 2 (1820) = **Cusparia**, Humb. (Rutac.).
cuneifolius, Nees & Mart. in Nov. Act. Nat. Cur. xi. (1823) 162 = Cusp. cuneifolia.
macrophyllus, Mikan, Delect. fasc. i. t. 2 = Cusp. macrophylla.

CONCHOCHILUS, Hassk. in Hoev. & De Vriese, Tijdschr. ix. (1842) 146 = **Appendicula**, Blume, (Orchid.).
distichus, Hassk. l. c. = A. pendula.
oppositiflorus, Hassk. l. c. 147 = A. cristata.

CONCHOPHYLLUM, Blume, Bijdr. 1060 (1826) = **Dischidia**, R. Br. (Asclepiad.).
imbricatum, Blume, l. c. 1061 = D. Collyris.

CONCILIUM, Rafin. Fl. Tellur. ii. 78 (1836) = **Lightfootia**, L'Hérit. (Campan.).
pedunculare, Rafin. l. c. = L. subulata.

CONDAEA, Steud. Nom. ed. II. i. 403 (1840) = CONDEA, Adans. = **Hyptis**, Jacq. (Labiat.).

CONDALIA, Cav. in Anal. Cienc. Nat. i. (1799) 39. t. 4. *RHAMNEAE*, Benth. & Hook. f. i. 376.
buxifolia, *Reiss. in Mart. Fl. Bras.* xi. i. 89.—Bras.
coriacea, Reichb. in Flora, xiii. (1830) i. 130 = Rhamnus integrifolius.
ferrea, Griseb. Fl. Brit. W. Ind. 100 = Scutia ferrea.
infectoria, *Reiss. in Mart. Fl. Bras.* xi. i. 90.—Bras.
lineata, *A. Gray, Bot. U. St. Expl. Exped.* i. 275.—Patagon.
maytenoides, *Miers, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 14.—Chili.
mexicana, *Schlecht. in Linnaea*, xv. (1841) 471.—Mexic.
microphylla, *Cav. in Anal. Cienc. Nat.* i. (1799) 40.—Chili.
obovata, *Hook. Ic. Pl.* t. 287.—Texas.
paradoxa, Spreng. Syst. i. 825 = Colletia cruciata.
spathulata, *A. Gray, Pl. Wright.* i. 32.—Am. bor. occ.
spinosa, *Spreng. Syst.* iv. *Cur. Post.* 108.—Bras.

CONDALIA, Rniz & Pav. Prod. 11. t. 2 (1794) = **Coccocypselum**, P. Br. (Rubiac.).
lanceolata, Ruiz & Pav. Fl. Per. i. 54 = Cocc. lanceolatum.
obovata, Ruiz & Pav. l. c. = Cocc. obovatum.
repens, Ruiz & Pav. l. c. = Cocc. umbellatum.
sessilis, Ruiz & Pav. l. c. = Cocc. sessile.

CONDAMINEA, DC. Prod. iv. 402 (1830). *RUBIACEAE*, Benth. & Hook. f. ii. 42.
corymbosa, *DC. l. c.*—Am. austr.
glabrata, *DC. l. c.*—Peruv.
macrophylla, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 30.—Bras.
microcarpa, *DC. Prod.* iv. 402.—Pernv.
tinctoria, *DC. l. c.*—Venezuela.
utilis, *Goudot, in Comptes Rendus*, xviii. (1844) 260.—N. Granat.
venosa, *DC. Prod.* iv. 402.—Peruv.

CONDEA, Adans. Fam. ii. 504 (1763) = **Hyptis**, Jacq. (Labiat.).

CONDYLICARPUS, Steud. Nom. ed. II. ii. 691 (1841) err. typ. = CONDYLOCARPUS, Hoffm. = **Tordylium**, Tourn. (Umbellif.).

CONDYLOCARPON, Desf. in Mém. Mus. Par. viii. (1822) 119. t. 11. *APOCYNACEAE*, Benth. & Hook. f. ii. 701.
HORTSMANIA, Miq. in Nat. Verh. Wet. Haarl. vii. 167 (1850).
MAYCOCKIA, A. DC. Prod. viii. 324 (1844).
brasiliense, Mart. ex Muell. Arg. in Mart. Fl. Bras. vi. 1. 64 = Rauwolfiae.
breviarticulatum, *Muell. Arg. l. c.* 65.—Bras.
ciliatum, *Muell. Arg. l. c.* 66.—Bras.
gracile, *Miers, Apocyn. S. Am.* 28.—Am. austr.
guyanense, *Desf. in Mém. Mus. Par.* viii. (1822) 120. t. 8.—Guiana.
intermedium, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 65.—Bras.
isthmicum, *A. DC. Prod.* viii. 381.—Bras.
laxum, *Muell. Arg. in Mart. Fl. Bras.* vi. 1. 66.—Bras.
myrtifolium, *Muell. Arg. l. c.* 65, in obs.—Guiana.
obtusiusculum, *Muell. Arg. l. c.*—Bras.
pubiflorum, *Muell. Arg. l. c.* 67.—Bras.
Rauwolfiae, *Muell. Arg. l. c.* 64.—Bras.

CONDYLOCARPUS, Hoffm. Gen Umb. ed. II. 202 (1816) = **Tordylium**, Tourn. (Umbellif.).
apulus, Hoffm. l. c. 203 = T. apulum.
humilis, Koch, in Nov. Act. Nat. Cur. xii. 1. (1824) 87 = T. apulum.
officinalis, Koch, l. c. = T. officinale.

CONDYLOCARPUS, Salisb. ex Lamb. Pinus, ed. 8vo, 120, sub t. 64 (1832) = **Sequoia**, Endl. (Conif.).
sempervirens, Salisb. ex Carr. Conif. ed. I. 163 = S. sempervirens.

CONFORATA, Caesalp. ex Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 91 = **Achillea**, Linn. (Compos.).
Ageratum, Fourr. l. c. = A. Ageratum.

CONGDONIA, Muell. Arg. in Flora, lix. (1876) 437. *RUBIACEAE.*
cacrulea, *Muell. Arg. l. c.*—Bras.

CONGEA, Roxb. Pl. Corom. iii. 90. t. 293 (1819). *VERBENACEAE*, Benth. & Hook. f. ii. 1159.
CALOCHLAMYS, Presl, Bot. Bemerk. 148 (1844).
azurea, Wall. Cat. n. 1733 = Sphenodesma pentandra.
barbata, Wall. l. c. n. 1738 = Sphenodesma barbata.
ferruginea, Wall. l. c. n. 1737 = Sphenodesma unguiculata.
involucratum, Wall. l. c. n. 1740 = Symphorema polyandrum.
Jackiana, Wall. l. c. n. 1735 = Sphenodesma pentandra, triflora.
paniculata, Wall. l. c. n. 1739 = Symphorema involucratum, Sphenodesma paniculata.
pentandra, Wall. l. c. n. 1734 = Sphenodesma pentandra.
tomentosa, *Roxb. Pl. Corom.* iii. 90.—Burma.
unguiculata, Wall. Cat. n. 1736 = Sphenodesma unguiculata.
velutina, *Wight, Ic.* t. 1479/3.—Ind. or.
vestita, *Griff. Notul.* iv. 174.—Burma.
villosa, *Wight, Ic.* t. 1479 B.—Burma.

CONGHAS, Wall. Cat. n. 8080; ex Hiern, in Hook. f. Fl. Brit. Ind. i. 681 (1875) = **Schleichera**, Willd. (Sapindac.).
seylonensis, Wall. l. c. = S. trijuga.

CONIANDRA, Schrad. in Eckl. & Zeyh. Enum. 275 (1836) = **Kedrostis**, Medic. (Cucurb.).
africana, Sond. in Harv. & Sond. Fl. Cap. ii. 483 = K. africana.
corallina, Fenzl, ex Naud. in Ann. Sc. Nat. Sér. IV. xvi. (1862) 180 = Corallocarpus Fenzlii.
digitata, Sond. in Harv. & Sond. Fl. Cap. ii. 483 = K. digitata.
dissecta, Schrad. in Eckl. & Zeyh. Enum. 276 = K. africana.
glauca, Schrad. l. c. 275 = K. grossulariaefolia.

CONIANDRA :—
grossulariaefolia, Arn. in Hook. Journ. Bot. iii. (1841) 273 = Kedrostis grossulariaefolia.
mollis, Sond. in Harv. & Sond. Fl. Cap. ii. 485 = K. mollis.
pinnatisecta, Schrad. in Eckl. & Zeyh. Enum. 276 = K. africana.
punctulata, Sond. in Harv. & Sond. Fl. Cap. ii. 484 = K. punctulata.
Thunbergi, Sond. l. c. = K. nana.
Zeyheri, Schrad. in Eckl. & Zeyh. Enum. 275 = K. Zeyheri.

CONIOGETON, Blume, Bijdr. 1156 (1826) = **Buchanania**, Roxb. (Anacard.).
arborescens, Blume, l. c. = B. florida.

CONIOSELINUM, Fisch. ex Hoffm. Gen. Umb. ed. I. p. 180 et xxviii. (1814); ed. II. 185 (1816) = **Ligusticum**, Linn. (Umbellif.).
altaicum, Rupr. in Beitr. Pfl. Russ. Reich. xi. 22.—Sibir.
canadense, Torr. & Gray, Fl. N. Am. i. 619 = Selinum canadense.
cenolophioides, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 736 = Selinum cenolophioides.
Czernaevia, Fisch. & Mey. Ind. Sem. Hort. Petrop. ii. 33 = Peucedanum latifolium.
Fischeri, Wimm. & Grab. Fl. Siles. (1827) i. 266 = Selinum Benthami.
gayoides, Less. in Linnaea, ix. (1834) 178.—Sibir.
Gmelini, Steud. Nom. ed. II. i. 403 = Ligusticum Gmelini.
humile, Turcz. ex Ledeb. Fl. Ross. ii. 313 = Peucedanum salinum.
ingricum, Fisch. Ind. Sem. Hort. Petrop. (1824) ex Ledeb. l. c. 290 = Selinum Benthami.
kamtschaticum, Rupr. in Beitr. Pfl. Russ. Reich. xi. 22.—Kamtsch.
latifolium, Rupr. Sert. Tiansch. 48.—Am. centr.
longifolium, Turcz. in Bull. Soc. Nat. Mosc. xvii. (1844) 736.—Sibir.
neglectum, Fisch. ex Steud. Nom. ed. II. i. 403, 404 = Selinum Benthami.
tataricum, Hoffm. Gen. Umb. ed. II. 185 = Selinum Benthami.
univittatum, Turcz. ex Kar. & Kir. in Bull. Soc. Nat. Mosc. xv. (1842) 363 = Selinum Benthami.

CONIOTHELE, DC. Prod. v. 531 (1836) = **Blennosperma**, Less. (Compos.).
californica, DC. l. c. = B. californicum.

CONIROSTRUM, Dulac, Fl. Hautes-Pyr. 195 (1867) = **Brassica**, Tourn. (Crucif.).
obtusangulum, Dulac, l. c. = B. Erucastrum.

CONISA, Desf. ex Steud. Nom. ed. II. i. 404 (1840) = **Conyza**, Linn. (Compos.).

CONIUM, Linn. Syst. ed. I (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 883.
africanum, Hort. ex DC. Prod. iv. 108 = Carum copticum.
africanum, Jacq. Hort. Vind. ii. 91. t. 194 = Capnophyllum Jacquini.
africanum, Linn. Sp. Pl. 243 = Capnophyllum africanum.
Arracacia, Hook. Exot. Bot. t. 152 = Arracacia xanthorrhiza.
chaerophylloides, Eckl. & Zeyh. Enum. 355 = Seseli chaerophylloides.
Cicuta, Neck. Delic. Gallo-Belg. i. 142 = maculatum.
croaticum, Waldst. & Kit. ex Willd. Enum. Hort. Berol. 305 = maculatum.
dichotomum, Desf. Fl. Atlant. i. 246. t. 66 = Capnophyllum dichotomum.
divaricatum, Boiss. & Orph. Diagn. Ser. II. v. 103 = maculatum.
filifolium, Vahl, Symb. Bot. iii. 49 = Oenanthe filiformis.
Jacquini, D. Dietr. Syn. Pl. ii. 988 = Capnophyllum Jacquini.
maculatum, *Linn. Sp. Pl.* 243.—Europ.; Oriens.
maculosum, Pall. Reis. Statth. Russ. Reich. i. 478 = maculatum.
Monieri, *Hort. ex Opiz, Natural.* x. (1825) 260, nomen.—Europ.

CONIUM:—
moschatum, H. B. & K. Nov. Gen. et Sp. v. 14. t. 420 = Arracacia moschata.
nepalense, D. Dietr. Syn. Pl. ii. 988 = Vicatia coniifolia.
nodosum, Fisch. ex Steud. Nom. ed. I. 215 = maculatum.
rigens, Linn. Mant. ii. 352, 512 = Bunium rigens.
rigidum, Mert. ex DC. Prod. iv. 140 (= *Sclerosciadium humile*).—Marocco.
Royeni, Linn. Sp. Pl. 243 = Caucalis daucoides.
rugosum, Thunb. Prod. Pl. Cap. 50 = Capnophyllum Jacquini.
sibiricum, Hort. ex Steud. Nom. ed. II. i. 404 = maculatum.
strictum, Tratt. Arch. Gew. i. t. 21 = maculatum.
suffruticosum, Berg. Desc. Pl. Cap. 77 = Selinum suffruticosum.
tenuifolium, Mill. Gard. Dict. ed. VIII. n. 2 = maculatum.
tenuifolium, Vahl, Symb. Bot. iii. 49 = Oenanthe filiformis.
verrucosum, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 324 = Pimpinella simensis.

CONIZA, Neck. Delic. Gallo-Belg. ii. 348 (1768) = **Conyza**, Linn. (Compos.).

CONNAROPSIS, Planch. ex Hook. f. in Trans. Linn. Soc. xxiii. (1860) 166. *GERANIACEAE*, Benth. & Hook. f. i. 277.
diversifolia, Kurz, in Journ. As. Soc. Beng. xxxix. (1870) ii. 69 = Griffithii.
glauca, *Hook. f. in Linn. Soc. Trans.* xxiii. (1860) 166.—Borneo.
Griffithii, *Planch. l. c. et ex Hook. f. Fl. Brit. Ind.* i. 440.—Malaya.
monophylla, *Planch. ll. cc.*—Malacca.
philippica, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 33.—Ins. Philipp.

CONNARUS, Linn. Diss. Dass. 23 (1747); in Amoen. Acad. i. 406 (1749). *CONNARACEAE*, Benth. & Hook. f. i. 432.
ANISOSTEMON, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) 1. 152.
CANICIDIA, Vell. Fl. Flum. 184 (1825); iv. t. 139 (1827).
CASTANOLA, Llanos, in Mem. Acad. Cienc. Madr. ii. (1859) 503.
ERYTHROSTIGMA, Hassk. in Flora, xxv. (1842) ii. Beibl. 45.
KALAWAEL, Adans. Fam. ii. 344 (1763).
MATHRANCIA, Stend. Nom. ed. II. i. 404, in syn. (1840).
OMPHALOBIUM, Gaertn. Fruct. i. 217. t. 46 (1788).
TALI, Adans. Fam. ii. 319 (1763).
TAPOMANA, Adans. Fam. ii. 343 (1763).
africanus, *Lam. Encyc.* ii. 95.—Afr. trop.
africanus, G. F. W. Mey. Prim. Fl. Esseq. 228 = guianensis.
Afzelii, R. Br. ex Planch. in Linnaea, xxiii. (1850) 418 = Rourea santaloides.
asiaticus, Willd. Sp. Pl. iii. 692 = monocarpus.
Beyrichii, *Planch. in Linnaea*, xxiii. (1850) 430.—Bras.
Blanchetii, *Planch. l. c.* 431.—Bras.
Championii, *Thw. Enum. Pl. Zeyl.* 80.—Zeylan.
conchocarpus, *F. Muell. Fragm.* v. 105.—Austral.
confertiflorus, *Baker, in Mart. Fl. Bras.* xiv. ii. 193.—Guiana.
crassifolius, Benth. ex Baker, l. c. 186 = favosus.
cuneifolius, *Baker, l. c.* 194.—Bras.
cymosus, *Planch. in Linnaea*, xxiii. (1850) 430.—Bras.
decumbens, *Thunb. in Roem. Arch.* i. 1. (1796) 1.—Afr. austr.
detersus, *Planch. in Linnaea*, xxiii. (1850) 435.—Bras.
Diepenhorstii, *Miq. Fl. Ind. Bat. Suppl.* i. 529.—Sumatra.
Duparquetianus, *Baill. Adansonia*. vii. (1866–67) 236.—Afr. trop.
erianthus, *Benth. ex Baker, in Mart. Fl. Bras.* xiv. ii. 191.—Bras.
falcatus, *Blume, Mus. Bot. Lugd. Bat.* i. 266.—Malaya.
fasciculatus, Planch. in Linnaea, xxiii. (1850) 432 = erianthus.
favosus, *Planch. l. c.* 434.—Bras.

CONNARUS:—
fecundus, *Baker, in Mart. Fl. Bras.* xiv. II. 185.—Bras.
ferrugineus, *Jack, in Malay. Misc.* ii. (1822) n. VII. 37. —Malaya.
floribundus, Planch. in Linnaea, xxiii. (1850) 435 = Perrottetii.
floribundus, *Schum. & Thonn. Beskr. Guin. Pl.* 299.— Afr. trop.
floribundus, Wall. ex Hook. f. Fl. Brit. Ind. ii. 52 = semidecandrus.
florulentus, Thonn. ex DC. Prod. ii. 86 = floribundus.
fluminensis, Gardn. in Hook. Lond. Journ. Bot. i. (1842) 529 = Bernardinia fluminensis.
foetens, Blanco, Fl. Filip. ed. I. 525 = Murraya exotica.
foliolosus, Jack, in Wall. Cat. n. 8529 = Cnestis ramiflora.
fulgens, Wall. Cat. n. 8524 = Rourea fulgens.
fulvus, *Planch. in Linnaea*, xxiii. (1850) 434.—Bras.
furfuraceus, *Blume, Mus. Bot. Lugd. Bat.* i. 268.— Sumatra.
Gaudichaudii, *Planch. in Linnaea*, xxiii. (1850) 429.— Ins. Molucc.
gibbosus, *Wall. Cat.* n. 8541, *ex Planch. l. c.* 436.— Burma; Malaya.
glaber, DC. Prod. ii. 85 = Rourea glabra.
grandifolius, *Planch. in Linnaea*, xxiii. (1850) 432.— Ins. S. Doming.
grandis, *Jack, in Malay. Misc.* ii. (1822) n. VII. 40. —Burma; Malaya.
Griffithii, *Hook. f. Fl. Brit. Ind.* ii. 52.—Burma.
Griffonianus, *Baill. Adansonia*, vii. (1866–67) 235.— Afr. trop.
guianensis, *Lamb. ex DC. Prod.* ii. 85.—Am. trop.
haemorrhaeus, *Karst. Fl. Columb.* ii. 73. t. 137.— N. Granat.
Hasseltii, *Blume, Mus. Bot. Lugd. Bat.* i. 266.—Java.
igneus, Wall. Cat. n. 8528 = Cnestis ramiflora.
incomptus, *Planch. in Linnaea*, xxiii. (1850) 433.— Guiana.
Jackianus, Wall. Cat. n. 8552 = Cupania Jackiana.
javanicus, *Blume, Bijdr.* 1166.—Java.
juglandifolius, Hook. & Arn. Bot. Beech. Voy. 179 = Rhus succedanea.
kavaiensis, *H. Mann, in Proc. Am. Acad.* vii. (1867) 162.—Ins. Sandvic.
latifolius, *Wall. Cat.* n. 5837.—Burma.
laurifolius, *Baker, in Mart. Fl. Bras.* xiv. II. 186.— Bras.
lucidus, *Hassk. in Hoev. & De Vriese, Tijdschr.* x. (1843) 144.—Malaya.
lucidus, *Jack, in Malay. Misc.* ii. (1822) n. VII. 41. —Malaya.
macrophyllus, Poepp. & Endl. Nov. Gen. et Sp. iii. 76 = Rourea macrophylla.
Maingayi, *Hook. f. Fl. Brit. Ind.* ii. 53.—Malaya.
Manni, *Baker, in Oliver, Fl. Trop. Afr.* i. 459.—Afr. trop.
marginatus, *Planch. in Linnaea*, xxiii. (1850) 429.— Bras.
microphyllus, Hook. & Arn. Bot. Beech. Voy. 179 = Rourea microphylla.
mimosoides, Jack, in Malay. Misc. ii. (1822) n. VII. 44 = Rourea concolor.
mimosoides, *Vahl, Symb. Bot.* iii. 87 (sp. dubia).—Ins. Nicobar.
mimosoides, Wall. Cat. n. 8526 = Rourea Wallichiana, similis.
moluccanus, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 266 = Gaudichaudii.
monadelphus, *Wall. Cat.* n. 8547 B.—Ind. or.
monocarpus, *Linn. Sp. Pl.* 675.—Ind. or.
monocarpus, Wight & Arn. Prod. 143 = Rourea santaloides.
monophyllus, Wall. Cat. n. 8551 = Ellipanthus tomentosus.
mutabilis, *Blume, Mus. Bot. Lugd. Bat.* i. 269.— Malaya.
nemorosus, *Vahl, ex Schum. & Thonn. Beskr. Guin. Pl.* 299.—Afr. trop.?
neurocalyx, *Planch. in Linnaea*, xxiii. (1850) 248.— Ins. Philipp.
nitidus, *Hassk. in Flora*, xxvii. (1844) II. 616.—Java.
nitidus, *Roxb. Hort. Beng.* 49; *Wall. Cat.* n. 8540 C. —Ind. or.
nodosus, *Baker, in Mart. Fl. Bras.* xiv. II. 190.—Bras.
obliquus, *Presl, in Walp. Ann.* iii. 844.—Ins. Luzon.
oblongifolius, Mart. Herb. Fl. Bras. n. 1267 = Rourea glabra.

CONNARUS:—
obtusifolius, *Planch. in Linnaea*, xxiii. (1850) 428.— Ins. Luzon.
odoratus, *Hook. f. in Trans. Linn. Soc.* xxiii. (1862) 172.—Borneo.
oligophyllus, *Planch. in Linnaea*, xxiii. (1850) 427.— Penang.
panamensis, *Griseb. in Bonplandia*, vi. (1858) 6.— Panama.
panamensis, Turcz. in Bull. Soc. Nat. Mosc. xxxii. (1859) I. 277 = Turczaninowii.
paniculatus, *Roxb. Hort. Beng.* 49; *Fl. Ind.* iii. 139.— Ind. or.
Patrisii, *Planch. in Linnaea*, xxiii. (1850) 432.— Guiana.
pentagynus, Lam. Encyc. ii. 95 = Agelaea Lamarckii.
pentandrus, Roxb. Hort. Beng. 49; Fl. Ind. iii. 140 = paniculatus.
Perrottetii, *Planch. in Linnaea*, xxiii. (1850) 432.— Guiana.
Pickeringii, *A. Gray, Bot. U. St. Expl. Exped.* i. 375. —Ins. Fiji.
pinnatus, Hasselt, ex Blume, Mus. Bot. Lugd. Bat. i. 266 = Hasseltii.
pinnatus, Lam. Encyc. ii. 95 = monocarpus.
polyanthus, *Planch. in Linnaea*, xxiii. (1850) 428.— Ins. Luzon.
polyphyllus, *Miq. Fl. Ind. Bat. Suppl.* 529.— Sumatra.
pubescens, *Baker, in Oliver, Fl. Trop. Afr.* i. 458.— Afr. trop.
pubescens, DC. Prod. ii. 85 = Rourea revoluta.
punctatus, *Planch. in Linnaea*, xxiii. (1850) 432.— Guiana.
pyrrhocarpus, *Miq. Fl. Ind. Bat. Suppl.* i. 530.— Sumatra.
reticulatus, *Griseb. Cat. Pl. Cub.* 84.—Cuba.
Ritchiei, *Hook. f. Fl. Brit. Ind.* ii. 51.—Ind. or.
Rolfei, *Vidal, Phan. Cuming. Philipp.* 23, 106.—Ins. Philipp.
Roxburghii, Hook. & Arn. Bot. Beech. Voy. 179 = Rourea Millettii.
ruber, *Planch. in Linnaea*, xxiii. (1850) 436.—Am. austr.
rugosus, Wall. Cat. n. 8527 = Rourea rugosa.
santaloides, Blanco, Fl. Filip. ed. II. 366 = Murraya exotica.
santaloides, Vahl, Sym. Bot. iii. 87 = Rourea santaloides.
Schomburgkii, *Planch. in Linnaea*, xxiii. (1850) 431. —Guiana.
semidecandrus, *Jack, in Malay. Misc.* ii. (1822) n. VII. 39.—Burma; Malaya.
Smeathmanni, *Planch. in Linnaea*, xxiii. (1850) 436; *Baker, in Oliver, Fl. Trop. Afr.* i. 458.—Afr. trop.
Spanoghei, *Blume, Mus. Bot. Lugd. Bat.* i. 267.—Ins. Timor.
Sprucei, *Baker, in Mart. Fl. Bras.* xiv. II. 187.—Bras.
stictophyllus, *Kurz, For. Fl. Brit. Burma*, i. 548.— Burma.
suberosus, *Planch. in Linnaea*, xxiii. (1850) 433.— Bras.
Thomsoni, *Baker, in Oliver, Fl. Trop. Afr.* i. 458.— Afr. trop.
Turczaninowii, *Triana & Planch. in Ann. Sc. Nat.* Sér. V. xvi. (1872) 364.—Panama.
turgidus, *Wall. Cat.* n. 8545, *ex Planch. in Linnaea*, xxiii. (1850) 436.—Ind. or.
unifoliolatus, Thw. Enum. Pl. Zeyl. 80 = Ellipanthus Thwaitesii.
velutinus, Wall. Cat. n. 8536 = Ellipanthus tomentosus.
venezuelanus, *Baill. Adansonia*, ix. (1868–70) 151.— Venezuela.
venosus, Smeathm. ex DC. Prod. ii. 85 = africanus.
villosus, *Jack, in Malay. Misc.* ii. (1822) n. VII. 38. —Sumatra.
villosus, Planch. in Linnaea, xxiii. (1850) 429 = grandis.
Wallichii, Planch. l. c. 426 = grandis.
Wightii, *Hook. f. Fl. Brit. Ind.* ii. 51.—Ind. or.

CONOBAEA, Bert. ex Steud. Nom. ed. II. i. 404 (1840) = **Muehlenbeckia**, Meissn. (Polygon.).
sagittaefolia, Bert. ex Steud. l. c. = M. sagittifolia.

CONOBEA, Aubl. Pl. Gui. ii. 639. t. 258 (1775). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 951.
BLANCKIA, Neck. Elem. i. 347 (1790).
LEUCOSPORA, Nutt. in Journ. Acad. Philad. vii. (1834) 87.

CONOBEA:—
SCHISTOPHRAGMA, Benth. in Endl. Gen. 679 (1838).
SPHAEROTHECA, Cham. & Schlecht. in Linnaea, ii. (1827) 605.
alata, R. Grah. in Edinb. N. Phil. Journ. (1830) 3 = Leucocarpus alatus.
aqnatica, *Aubl. Pl. Gui.* ii. 639. t. 258.—Guiana.
aquatica, Billb. ex Beurl. in Vet. Akad. Handl. Stockh. 1854 (1856) 141 = Herpestis lanigera.
borealis, Spreng. Neue Entdeck. iii. 26 = Gratiola virginiana.
indica, Spreng. Syst. ii. 771 = Limnophila polystachya.
intermedia, *A. Gray, in Torr. Bot. Mex. Bound.* 117. —Mexic.
multifida, *Benth. in DC. Prod.* x. 391.—Am. bor.
ovata, Spreng. ex Schrank, in Syll. Ratisb. ii. (1828) 62 = Stemodia parviflora?
pumila, Spreng. Nov. Prov. 13 = Stemodia parviflora.
punctata, *Nees & Mart. in Nov. Act. Nat. Cur.* xi. (1823) 143.—Bras.
pusilla, *Benth. & Hook. f. Gen.* ii. 951.—Mexic.
scoparioides, *Benth. in DC. Prod.* x. 391.—Mexic.
vandellioides, *Benth. l. c.*—Bras.
verticillaris, Hort. Berol. ex Spreng. Nov. Prov. 13 = Stemodia durantifolia.
viscosa, Schrank, in Syll. Ratisb. ii. (1828) 61 = Stemodia durantifolia.
viscosa, *Spreng. Neue Entdeck.* i. 263.—Bras.

CONOCALPIS, Boj. ex Decne. in DC. Prod. viii. 621 (1844) = **Gymnema**, R. Br. (Asclepiad.).
umbellata, Boj. ex Decne. l. c. = G. rufescens.

CONOCARPUS, Adans. Fam. ii. 284 (1763) = **Leucadendron**, R. Br. (Proteac.).

CONOCARPUS, Linn. Gen. ed. I. 376 (1737); Gaertn. Fruct. ii. 470. t. 177 (1791). *COMBRETACEAE*, Benth. & Hook. f. i. 686.
RHIZAERIS, Rafin. Sylva Tellur. 90 (1838).
RUDBECKIA, Adans. Fam. ii. 80 (1763).
acuminatus, Buch.-Ham. ex Wall. Cat. n. 4014, 4015 = Anogeissus latifolia.
acuminatus, Roxb. Hort. Beng. 34; Fl. Ind. ii. 443 = Anogeissus acuminata.
acutifolius, Willd. in Roem. & Schult. Syst. v. 574 = erectus.
erectus, *Linn. Sp. Pl.* 147; *Jacq. Select. Am.* 78. t. 52. —Ind. occ.
hirtus, Buch.-Ham. ex Wall. Cat. n. 4016 = Anogeissus acuminata.
lanceolatus, Heyne, ex Wall. l. c. n. 4014 = Anogeissus acuminata.
latifolius, Roxb. Hort. Beng. 34; Fl. Ind. ii. 442 = Anogeissus latifolia.
leiocarpus, DC. Prod. iii. 16 = Anogeissus leiocarpa.
monocarpus, Steud. Nom. ed. I. 216 = Rhus Rhadaelijavel.
myrtifolius, Buch.-Ham. ex Wall. Cat. n. 4017 = Anogeissus pendula.
niloticus, *Delile, ex Steud. Nom.* ed. II. i. 404.— Aegypt.
Novae-Zeylandiae, *Cels, ex Tenore, Cat. Hort. Neap.* (1845) 82.—N. Zel.
parvifolius, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 269 = Anogeissus leiocarpa.
procumbens, Linn. Sp. Pl. 148 = erectus.
pubescens, *Schum. & Thonn. Beskr. Guin. Pl.* 115.— Afr. trop.
racemosus, Linn. Syst. ed. X. 930; Jacq. Select. Am. 80 = Laguncularia racemosa.
Schimperi, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 269 = Anogeissus leiocarpa.
sericea, Forst. ex G. Don, Gen. Syst. ii. 662 = erectus.
supina, Crantz, Inst. i. 355 = erectus.

CONOCEPHALUS, Blume, Bijdr. 483 (1825). *URTICACEAE*, Benth. & Hook. f. iii. 380.
acuminatus, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 91.—Ins. Philipp.
azureus, *Teijsm. & Binn. in Tijdschr. Nederl. Ind.* xxvi. (1864) 26.—Java; Sumatra.
Blumei, *Gaudich. Bot. Voy. Bonite*, t. 96.—Malaya.
borneensis, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 210. —Borneo.
concolor, *Dalz. in Dalz. & Gibs. Bomb. Fl.* 239.— Ind. or.
ellipticus, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 91.—Penang.

CONOCEPHALUS:—
erectus, *Villar, in Blanco, Fl. Philipp.* ed. III. *Nov. App.* 203.—Ins. Philipp.
gratus, Miq. Pl. Jungh. i. 43 = suaveolens.
lanceolatus, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 88.—Ind. or.
micranthus, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 210.—Borneo.
microphyllus, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 92.—Ins. Philipp.
naucleiformis, Lindl. Bot. Reg. t. 1203 = suaveolens.
niveus, Wight, Ic. vi. 7. t. 1959 = Debregeasia velutina.
ovatus, Tréc. in Ann. Sc. Nat. Sér. III. viii. (1847) 88 = suaveolens.
pubescens, Tréc. l. c. 90 = suaveolens.
Roxburghii, Tréc. l. c. 88 = suaveolens.
suaveolens, *Blume, Bijdr.* 483.—Ind. or.; Malaya.
subtrinervius, *Miq. Fl. Ind. Bat. Suppl.* 417.—Malaya.

CONOCLINIUM, DC. Prod. v. 135 (1836) = **Eupatorium**, Tourn. (Compos.).
acuminatum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 361 = E. organense.
adenolepis, Sch. Bip. ex Baker, l. c. = E. Selloi.
affine, Gardn. in Hook. Lond. Journ. Bot. v. (1846) 466 = E. betonicaeforme.
album, Mart. Del. Sem. Hort. Monac. (1848) 4.—Mexic.
ballotaefolium, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 360 = E. ballotaefolium.
betonicaeforme, DC. Prod. v. 135 = E. betonicaeformc.
betonicum, DC. Prod. v. 135 = E. betonicum.
cerasifolium, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 308 = E. cerasifolium.
coelestinum, DC. Prod. v. 135 = E. coelestinum.
dichotomum, Chapm. in Coult. Bot. Gaz. iii. (1878) 5 = E. coelestinum.
dissectum, A. Gray, Pl. Wright. i. 88 = E. Greggii.
hirsutissimum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 311 = E. hirsutissimum.
humile, Benth. Pl. Hartw. 199.—N. Granat.
ianthinum, Morr. in Ann. Soc. Gand, v. (1849) 173 = E. ianthinum.
Lasseauxii, Dur. Ind. Sem. Hort. Burdig. (1872) 15.—Uruguay.
maritimum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 362 = E. apiculatum.
microcephalum, Sch. Bip. ex Baker, l. c. 364 = E. Candollcanum.
microphyllum, DC. Prod. v. 135 = E. microphyllum.
mikanioides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 306 = E. sordescens.
oligolepis, Kunze, Del. Sem. Hort. Lips. (1840) iv. 2 = E. oligolcpis.
organense, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 361 = E. organense.
palustre, Sch. Bip. ex Baker, l. c. 363 = E. palustrc.
platylepis, Sch. Bip. ex Baker, l. c. 355 = E. platylcpis.
prasiifolium, DC. in Deless. Ic. Sel. iv. t. 15 = E. nepetoides.
resinosum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 306 = E. sordescens.
rigidum, Chapm. in Coult. Bot. Gaz. iii. (1878) 6 = E. hcteroclinum.
rugosum, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) I. 168.—Ecuador.
scandens, Gardn. in Hook. Lond. Journ. Bot. vi. (1847) 437.—Bras.
semistriatum, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 318 = E. semistriatum.
subglutinosum, A. Gray, in Proc. Am. Acad. v. (1861) 115; Sch. Bip. ex Baker, l. c. 362 = E. apiculatum.
subviscosum, Baker, l. c., sphalm. = E. apiculatum.
Vindex, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. II. 312 = E. Vindex.

CONOHORIA, Aubl. Pl. Gui. i. 239. t. 95 (1775) = **Alsodeia**, Thou. (Violar.).
alternifolia, Spreng. Neue Entdeck. ii. 151 = A. integrifolia.
Aubletii, D. Dietr. Syn. Pl. i. 831 = Cascaria dentata.
castaneaefolia, St. Hil. in Mém. Mus. Par. xi. (1824) 495 = A. guianensis.

CONOHORIA:—
Cuspa, H. B. & K. Nov. Gen. et Sp. vii. 242 = Alsodeia Cuspa.
dentata, D. Dietr. Syn. Pl. i. 831 = A. guianensis.
flavescens, Aubl. Pl. Gui. i. 239 = A. flavescens.
Loboloba, St. Hil. in Mém. Mus. Par. xi. (1824) 494 = A. physiphora.
megapotamica, G. Don, Gen. Syst. i. 341 = A. megapotamica.
owariensis, D. Dietr. Syn. Pl. i. 831 = A. owariensis.
Passoura, DC. Prod. i. 312 = A. flavescens.
Perrini, D. Dietr. Syn. Pl. i. 832 = A. Perrini.
Regnellii, Miq. in Linnaea, xxii. (1849) 555.—Bras.
Riana, DC. Prod. i. 312 = A. prunifolia.
Rinorea, St. Hil. Pl. Rem. 321 = A. paniculata.
ulmifolia, H. B. & K. Nov. Gen. et Sp. v. 387. t. 491 = A. ulmifolia.

CONOMITRA, Fenzl, in Endl. Nov. Stirp. Dec. 65 (1839) = **Glossonema**, Decne. (Asclepiad.).
linearis, Fenzl, l. c. = G. lineare.

CONOMORPHA, A. DC. in Trans. Linn. Soc. xvii. (1834) 102. *MYRSINEAE*, Benth. & Hook. f. ii. 644.
CONOSTYLUS, Pohl, ex A. DC. l. c. (1834).
bumelioides, *Griseb. Cat. Pl. Cub.* 163.—Cuba.
guianensis, A. DC. in Ann. Sc. Nat. Sér. II. xvi. (1841) 92 = Cybianthus guyanensis.
heterantha, *Benth. ex Miq. in Mart. Fl. Bras.* x. 304.—Bras.
jacquinioides, *Griseb. Cat. Pl. Cub.* 163.—Cuba.
laxiflora, *DC. in Trans. Linn. Soc.* xvii. (1837) 103.—Bras.
macrophylla, *Mart. Herb. Fl. Bras.* n. 260.—Bras.
multipunctata, *Miq. Stirp. Surin. Sel.* 111. t. 34.—Guiana.
nemoralis, *Mart. & Miq. in Mart. Fl. Bras.* x. 304.—Bras.
oblongifolia, *A. DC. in Trans. Linn. Soc.* xvii. (1837) 103.—Bras.
peruviana, *A. DC. in Ann. Sc. Nat.* Sér. II. xvi. (1841) 92.—Peruv.
reticulata, *Benth. ex Miq. in Mart. Fl. Bras.* x. 303.—Bras.
robusta, *Klotzsch, in Faun. & Fl. Gui.* 974.—Guiana.

CONOPHALLUS, Schott, Syn. Aroid. 34 (1856) = **Amorphophallus**, Blume (Aroid.).
Blumei, Schott, in Ann. Mus. Bot. Lugd. Bat. i. 124.—Java.
bulbifer, Schott, Syn. Aroid. 34 = A. bulbifer.
commutatus, Schott, in Bonplandia, vii. (1859) 28 = A. commutatus.
giganteus, Schott, ex Miq. Fl. Ind. Bat. iii. 198 = A. giganteus.
Gigas, Miq. Ann. Mus. Bot. Lugd. Bat. i. 285 = A. Gigas.
Konjak, Schott, Syn. Aroid. 34 = A. Rivieri.
lyratus, Schott, l. c. 35 = A. lyratus.
Muelleri, Miq. Ann. Mus. Bot. Lugd. Bat. 286, in obs. = A. Mulleri.
planus, Miq. l. c. 286 = A. planus.
punctulatus, Schott, Prod. Aroid. 35 = A. punctulatus.
sativus, Schott, l. c. = A. sativus.
spectabilis, Miq. Fl. Ind. Bat. iii. 199 = A. spectabilis.
Titanum, Becc. in Bull. Soc. Tosc. Ortic. iii. (1878) 271, 291 = A. Titanum.
tuberculiger, Schott, in Bonplandia, vii. (1859) 28 = A. tuberculiger.

CONOPHARYNGIA, G. Don, Gen. Syst. iv. 94 (1837) = **Plumeria**, Tourn. (Apocyn.).
longifolia, G. Don, l. c. 95 = P. longifolia.
retusa, G. Don, l. c. = P. retusa.

CONOPHOLIS, Wallr. Orob. Diask. 78 (1825). *OROBANCHACEAE*, Benth. & Hook. f. ii. 985.
americana, *Wallr. l. c.*—Am. bor.
americana, Endl. Iconogn. t. 81 = mexicana.
mexicana, *A. Gray, ex S. Wats. in Proc. Am. Acad.* xviii. (1882–83) 131.—N. Mexic.

CONOPHYTA, Schum. ex Hook. f. in Trans. Linn. Soc. xxii. (1856) 42 = **Thonningia**, Vahl (Balanoph.).

CONOPODIUM, Koch, in Nov. Act. Nat. Cur. xii. (1824) 118. *UMBELLIFERAE*, Benth. & Hook. f. i. 896.
BULBOCASTANUM, Lag. Amen. Acad. Madr. 99 (1821).
BUTINIA, Boiss. Elench. 53 (1838).
ELAEOSTICTA, Fenzl, in Flora, xxvi. (1843) 458.
HETEROTAENIA, Boiss. Voy. Bot. Espagne, 267. t. 80 A (1840).
SCALIGERIA, DC. Prod. iv. 248 (1830).
SPHALLEROCARPUS, Bess. in DC. Prod. iv. 230 (1830).
allioides, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 588.—Turkestan.
boetica, *Boiss. Voy. Bot. Espagne*, ii. 738.—Hispan.
Bourgaei, *Coss. Notes Crit.* 110.—Hispan.
brachycarpum, *Boiss. ex Lange, in Kjoeb. Vidensk. Meddel.* (1865) 44.—Hispan.
butinoides, *Boiss. & Reut. ex Boiss. Voy. Bot. Espagne*, ii. 736.—Hispan.
canadense, Koch, in Nov. Act. Nat. Cur. xii. (1824) 119 = Cryptotaenia canadensis.
capillifolium, *Boiss. Voy. Bot. Espagne*, ii. 736.—Lusitan.
Cyminum, *Benth. & Hook. f. Gen.* i. 896.—Dahuria; Volh.
denudatum, *Koch, in Nov. Act. Nat. Cur.* xii. (1824) 118.—Europ.
hirtulum, *Regel & Schmalh. in Pl. Nov. Fedsch.* 28.—As. centr.
Marianum, *Lange, in Kjoeb. Vidensk. Meddel.* (1877–78) 232.—Hispan.
mutabile, Miégev. in Bnll. Soc. Bot. Fr. xxi. (1874) . p. xxxii = denudatum.
pumilum, Nym. Consp. 303 = Bourgaei.
pyrenaicum, Miégev. in Bull. Soc. Bot. Fr. xxi. (1874) p. xxxii = denudatum.
ramosum, *Costa, Fl. Catal.* 105.—Hispan.
subcarneum, *Boiss. & Reut. ex Boiss. Voy. Bot. Espagne*, ii. 736.—Hispan.
subcarneum, Bourg. in Bull. Soc. Bot. Fr. ii. (1855) 643 = Bourgaei.

CONOPSIDIUM, Wallr. in Linnaea, xiv. (1840) 147 = **Habenaria**, Willd. (Orchid.).
platantherum, Wallr. l. c. = H. bifolia.
sthenantherum, Wallr. l. c. = H. bifolia.

CONORIA, Juss. Gen. 287 (1789) = **Alsodeia**, Thou. (Violar.).

CONOSAPIUM, Muell. Arg. in Linnaea, xxxii. (1863) 87 = **Sapium**, P. Br. (Euphorb.).
madagascariensis, Muell. Arg. l. c.—Madag.

CONOSILENE, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344 = **Silene**, Linn. (Caryophyll.).
conica, Fourr. l. c. = S. conica.
conoidea, Fourr. l. c. = S. conoidea.

CONOSIPHON, Poepp. & Endl. Nov. Gen. et Sp. iii. 27 (1842); Endl. Gen. Pl. Suppl. ii. 54 (1842) = **Sphinctanthus**, Benth. (Rubiac.).
aureus, Poepp. & Endl. l. c. = S. rupestris.
polycarpus, Karst. in Linnaea, xxx. (1859–60) 155.—N. Granat.
striiflorus, Walp. Rep. ii. 520 = S. rupestris.

CONOSPERMUM, Sm. in Trans. Linn. Soc. iv. (1798) 213. *PROTEACEAE*, Benth. & Hook. f. iii. 174.
acerosum, *Lindl. Swan River App.* 30.—Austral.
acinacifolium, *R. Grah. in Edinb. N. Phil. Journ.* (Apr.-July, 1826) 171.—Austral.
affine, *Roem. & Schult. Syst.* iii. *Mant.* 274.—Austral.
amoenum, *Meissn. in Lehm. Pl. Preiss.* i. 522.—Austral.
brachyphyllum, *Lindl. Swan River App.* 31.—Austral.
bracteosum, *Meissn. in Lehm. Pl. Preiss.* i. 518; ii. 248.—Austral.
Brownii, *Meissn. l. c.* ii. 248.—Austral.
canaliculatum, Meissn. l. c. 250 = Stoechadis.
capitatum, *R. Br. in Trans. Linn. Soc.* x. (1811) 155.—Austral.
coerulescens, F. Muell. Fragm. i. 157 = amoenum.
coeruleum, *R. Br. in Trans. Linn. Soc.* x. (1811) 154.—Austral.
commutatum, *Roem. & Schult. Syst.* iii. *Mant.* 275.—Austral.

CONOSPERMUM:—
crassinervium, *Meissn. in Hook. Kew Journ.* iv. (1852) 184.—Austral.
Dallachyi, F. Muell. Ann. Rep. (1858) fide Benth. Fl. Austral. v. 371 = Mitchellii.
debile, *Kipp. ex Meissn. in Hook. Kew Journ.* vii. (1855) 70.—Austral.
densiflorum, *Lindl. Swan River App.* 32.—Austral.
diffusum, *Benth. Fl. Austral.* v. 367.—Austral.
distichum, Meissn. in Lehm. Pl. Preiss. i. 522 = floribundum.
distichum, *R. Br. in Trans. Linn. Soc.* x. (1811) 155.—Austral.
ellipticum, *Sm. in Rees, Cycl.* ix. n. 2.—Austral.
ephedroides, *Kipp. ex Meissn. in Hook. Kew Journ.* vii. (1855) 70.—Austral.
erectum, Hort. ex R. Grah. in Edinb. N. Phil. Journ. (Apr.-June 1828) 172 = ericifolium.
ericifolium, *Sm. in Rees, Cycl.* ix. n. 4.—Austral.
falcifolium, [Salisb. in] Knight, Prot. 95 = taxifolium.
filifolium, Meissn. in Lehm. Pl. Preiss. i. 523 = brachyphyllum.
flexuosum, *R. Br. Prod. Suppl.* 11.—Austral.
floribundum, *Benth. Fl. Austral.* v. 373.—Austral.
glumaceum, *Lindl. Swan River App.* 30.—Austral.
Huegelii, *R. Br. ex Endl. Nov. Stirp. Dec.* 58.—Austral.
imbricatum, *Sieber, ex Spreng. Syst.* iv. *Cur. Post.* 46.—Austral.
incurvum, *Lindl. Swan River App.* 30.—Austral.
intricatum, Lindl. ex Meissn. in DC. Prod. xiv. 323 = Huegelii.
lanceolatum, *R. Br. Prod. Suppl.* 10.—Austral.
laniflorum, *Endl. in Ann. Wien. Mus.* ii. (1838) 208.—Austral.
lavandulaefolium, *A. Cunn. ex Meissn. in DC. Prod.* xiv. 319.—Austral.
linifolium, *A. Cunn. l. c.* 320.—Austral.
longifolium, *Sm. Exot. Bot.* ii. 45. t. 82.—Austral.
lupulinum, Endl. ex Meissn. in Lehm. Pl. Preiss. ii. 249 = glumaceum.
marginatum, *Meissn. l. c.* 248.—Austral.
Mitchellii, *Meissn. in DC. Prod.* xiv. 320.—Austral.
nervosum, *Meissn. in Hook. Kew Journ.* vii. (1855) 71.—Austral.
patens, *Schlecht. in Linnaea*, xx. (1847) 587.—Austral.
petiolare, *R. Br. Prod. Suppl.* 11.—Austral.
polycephalum, *Meissn. in Lehm. Pl. Preiss.* ii. 249.—Austral.
procerum, F. Muell. Fragm. i. 157 = distichum.
propinquum, R. Br. Prod. Suppl. 10 = taxifolium.
repens, Sieber, ex Roem. & Schult. Syst. iii. Mant. 276 = tenuifolium.
reticulatum, Sm. in Rees, Cycl. ix. n. 5 = Synaphea petiolaris.
rigidum, [Salisb. in] Knight, Prot. 95 = ellipticum.
scaposum, *Benth. Fl. Austral.* v. 369.—Austral.
sclerophyllum, *Lindl. Swan River App.* 30.—Austral.
Smithii, Pers. Syn. i. 113 = longifolium.
sphacelatum, *Hook. in Mitch. Trop. Austral.* 342.—Austral.
spicatum, R. Br. Prod. Suppl. 10 = taxifolium.
Stoechadis, *Endl. in Ann. Wien. Mus.* ii. (1838) 208.—Austral.
taxifolium, Sieber, ex Roem. & Schult. Syst. iii. Mant. 274 = affine.
taxifolium, *Sm. in Rees, Cycl.* ix. n. 3.—Austral.
tenuifolium, *R. Br. in Trans. Linn. Soc.* x. (1811) 154.—Austral.
tenuifolium, Sieber, ex Roem. & Schult. Syst. iii. Mant. 275 = commutatum.
teretifolium, *R. Br. in Trans. Linn. Soc.* x. (1811) 155.—Austral.
Toddii, *F. Muell. Fragm.* x. 20.—Austral.
triplinervium, *R. Br. Prod. Suppl.* 11.—Austral.
undulatum, *Lindl. Swan River App.* 31.—Austral.

CONOSTEGIA, D. Don, in Mem. Wern. Soc. iv. (1823) 316. *MELASTOMACEAE*, Benth. & Hook. f. i. 763.
BRUGUIERA, Rich. ex DC. Prod. iii. 173, in syn. (1828).
CALYCOTOMUS, Rich. ex DC. l. c. (1828).
SYNODON, Rafin. Sylva Tellur. 95 (1838).
acuminata, Steud. in Flora, xxvii. (1844) 722 = Miconia axilliflora.
acutidentata, A. Rich. Ess. Fl. Cub. 558 = xalapensis.
alpina, Macf. Fl. Jamaic. ii. 72 = montana.

CONOSTEGIA:—
alternifolia, Macf. Fl. Jamaic. ii. 71 = superba.
arborea, *Steud. Nom.* ed. II. i. 405; *Schau. in Linnaea*, xx. (1847) 733.—Mexic.
attenuata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 98.—N. Granat.
Balbisiana, Ser. ex DC. Prod. iii. 174 = procera.
bracteata, *Triana, in Journ. Bot.* v. (1867) 209.—Nicaragua.
calyptrata, *D. Don, in Mem. Wern. Soc.* iv. (1823) 317.—Ind. occ.
clidemioides, *Wright, ex Griseb. Cat. Pl. Cub.* 98.—Cuba.
cornifolia, Ser. ex DC. Prod. iii. 175 = Miconia cinnamomifolia.
cucullata, D. Don, in Mem. Wern. Soc. iv. (1823) 317 = Calyptrella cucullata.
dentata, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 99.—N. Granat.
discolor, DC. Prod. iii. 174 = Charianthus coccineus.
extinctoria, *D. Don, in Mem. Wern. Soc.* iv. (1823) 317.—N. Granat.
formosa, Macf. Fl. Jamaic. ii. 70 = rufescens.
Galeottii, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 107 = arborea.
glabra, D. Don, in Mem. Wern. Soc. iv. (1823) 317 = Astronia Forsteri.
gloriosa, Macf. Fl. Jamaic. ii. 68 = procera.
holosericea, D. Don, in Mem. Wern. Soc. iv. (1823) 317 = xalapensis.
lasiopoda, *Benth. Bot. Voy. Sulph.* 96.—N. Granat.
lutescens, Ser. ex DC. Prod. iii. 175 = Miconia cinnamomifolia.
macrantha, *O. Berg, ex Triana, in Trans. Linn. Soc.* xxviii. (1871) 97.—Costa Rica.
macrophylla, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 112 = superba.
mexicana, Ser. ex DC. Prod. iii. 175 = Miconia mexicana.
montana, *D. Don, in Mem. Wern. Soc.* iv. (1823) 317.—Jamaic.
Mutisii, Ser. ex DC. Prod. iii. 174 = Centronia Mutisii.
myriasporoides, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 99.—N. Granat.
Oerstediana, *O. Berg, ex Triana, l. c.* 98.—Costa Rica.
parviflora, *DC. Prod.* iii. 175.—Ins. S. Doming.
polyandra, *Benth. Bot. Voy. Sulph.* 96. t. 35.—N. Granat.
procera, *D. Don, in Mem. Wern. Soc.* iv. (1823) 317.—Jamaic.
pulverulenta, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851) 110 = extinctoria.
purpurea, *Griseb. in Bonplandia*, vi. (1858) 6.—Panama.
quadrangularis, *Schlecht. ex Steud. Nom.* ed. II. i. 405.—Mexic.
rufescens, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 107.—Ind. occ.
semicrenata, Ser. ex DC. Prod. iii. 175 = Miconia semicrenata.
setosa, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 99.—N. Granat.
speciosa, *Naud. in Ann. Sc. Nat.* Sér. III. xvi. (1851) 109.—Panama.
sphaerica, *Triana, in Trans. Linn. Soc.* xxviii. (1871) 98.—Mexic.
subcrustulata, Triana, l. c. = purpurea.
subhirsuta, *DC. Prod.* iii. 174.—Cuba.
superba, *D. Don, in Mem. Wern. Soc.* iv. (1823) 317.—Am. austr.
tunicata, Ser. ex DC. Prod. iii. 175 = Miconia tunicata.
xalapensis, *D. Don, in Mem. Wern. Soc.* iv. (1823) 317.—Mexic.

CONOSTEMUM, Kunth, Enum. Pl. ii. 557 (1837), sphalm. = COMOSTEMUM, Nees = **Androtrichum**, Brongn. (Cyperac.).

CONOSTEPHIOPSIS, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 5 = **Conostephium**, Benth. (Epacrid.).
Drummondi, Stschegl. l. c. 6 = Conostephium planifolium.
minor, Stschegl. l. c. = Conostephium minus.
Preissii, Stschegl. l. c. = Conostephium Preissii.

CONOSTEPHIUM, Benth. in Enum. Pl. Hueg. 76 (1837). *EPACRIDEAE*, Benth. & Hook. f. ii. 611.
CONOSTEPHIOPSIS, Stschegl. in Bull. Soc. Nat. Mosc. xxxii. (1859) i. 5.
minus, *Lindl. Swan River App.* 25.—Austral.
nitens, *Lindl. ex Sond. in Lehm. Pl. Preiss.* i. 303.—Austral.
pendulum, *Benth. in Enum. Pl. Hueg.* 76.—Austral.
pendulum, Deless. Ic. Sel. v. t. 23 = minus.
planifolium, *F. Muell. Fragm.* vi. 30.—Austral.
Preissii, *Sond. in Lehm. Pl. Preiss.* i. 304.—Austral.
Roei, *Benth. Fl. Austral.* iv. 160.—Austral.

CONOSTYLIS, R. Br. Prod. 300 (1810). *HAEMODORACEAE*, Benth. & Hook. f. iii. 676.
ANDROSTEMMA, Lindl. Swan River App. 46 (1839).
aculeata, *R. Br. Prod.* 300.—Austral.
aemula, Lindl. Swan River App. 45 = setigera.
albicans, A. Cunn. ex Benth. Fl. Austral. vi. 436 = candicans.
americana, Pursh, Fl. Am. Sept. i. 224. t. 6 = Lophiola aurea.
androstemma, *F. Muell. Fragm.* viii. 19.—Austral.
assimilis, Endl. in Lehm. Pl. Preiss. ii. 20 = setigera.
aurea, *Lindl. Swan River App.* 44.—Austral.
Bealiana, *F. Muell. Fragm.* ix. 50.—Austral.
bracteata, *Endl. in Lehm. Pl. Preiss.* ii. 16.—Austral.
bracteata, Lindl. Swan River App. 45 = dealbata.
breviscapa, *R. Br. Prod.* 301.—Austral.
bromelioides, *Endl. in Lehm. Pl. Preiss.* ii. 18.—Austral.
candicans, *Endl. Nov. Stirp. Dec.* 20.—Austral.
canescens, F. Muell. Fragm. viii. 19 = Blancoa canescens.
caricina, *Lindl. Swan River App.* 45.—Austral.
cymosa, *F. Muell. ex Benth. Fl. Austral.* vi. 439.—Austral.
dealbata, *Lindl. Swan River App.* 45.—Austral.
discolor, Endl. in Lehm. Pl. Preiss. ii. 20 = setigera.
Drummondi, *Benth. Fl. Austral.* vi. 433.—Austral.
ensifolia, Endl. in Lehm. Pl. Preiss. ii. 21 = serrulata.
festucacea, Endl. l. c. 18 = Preissii.
filifolia, *F. Muell. Fragm.* viii. 18.—Austral.
gladiata, *Benth. Fl. Austral.* vi. 434.—Austral.
graminea, Endl. in Lehm. Pl. Preiss. ii. 19 = caricina.
involucrata, *Endl. l. c.* ii. 23.—Austral.
juncea, *Endl. Nov. Stirp. Dec.* 19.—Austral.
juncea, *Lindl. Bot. Reg.* (1839) *Misc.* 54.—Austral.
laxiflora, *Benth. Fl. Austral.* vi. 439.—Austral.
longifolia, Endl. in Lehm. Pl. Preiss. ii. 22 = serrulata.
melanopogon, *Endl. l. c.* ii. 18.—Austral.
minima, Endl. l. c. ii. 21 = C. Psyllium.
misera, Endl. l. c. 22 = serrulata.
occulta, Endl. l. c. = serrulata.
petrophiloides, *F. Muell. ex Benth. Fl. Austral.* vi. 431.—Austral.
Preissii, *Endl. in Lehm. Pl. Preiss.* ii. 18.—Austral.
prolifera, *Benth. Fl. Austral.* vi. 436.—Austral.
propinqua, Endl. in Lehm. Pl. Preiss. ii. 17 = candicans.
Psyllium, *Endl. l. c.* 21.—Austral.
pusilla, Endl. l. c. 20 = psyllium.
racemosa, *Benth. Fl. Austral.* vi. 436.—Austral.
scorsiflora, *F. Muell. Fragm.* i. 158.—Austral.
serrulata, *R. Br. Prod.* 300.—Austral.
setigera, *R. Br. l. c.*—Austral.
setosa, *Lindl. Swan River App.* 44. t. 6 A—Austral.
spathacea, Endl. in Lehm. Pl. Preiss. ii. 22 = serrulata.
spinuligera, *F. Muell. ex Benth. Fl. Austral.* vi. 438.—Austral.
stylidioides, *F. Muell. Fragm.* viii. 17.—Austral.
sulphurea, Endl. in Lehm. Pl. Preiss. ii. 17 = aurea.
teretiuscula, *F. Muell. Fragm.* viii. 18.—Austral.
vaginata, *Endl. in Lehm. Pl. Preiss.* ii. 23.—Austral.
villosa, *Benth. Fl. Austral.* vi. 433.—Austral.

CONOSTYLUS, Pohl, ex A. DC. in Trans. Linn. Soc. xvii. (1834) 102 = **Conomorpha**, A. DC (Myrsin.).
oblongifolius, Pohl, ex A. DC. l. c. 103 = Conom. oblongifolia.

CONOTHAMNUS, Lindl. Swan River App. 9 (1839). *MYRTACEAE*, Benth. & Hook. f. i. 705.
divaricatus, *Benth. Fl. Austral.* iii. 164.—Austral.
trinervis, *Lindl. Swan River App.* 9.—Austral.

CONOTRICHIA, A Rich. in Mém. Soc. Hist. Nat. Par. (1830) 277. t. 24. f. 1 = **Manettia**, Mutis (Rubiac.).
alba, A. Rich. l. c. = M. picta.

CONRADIA, Mart. Nov. Gen. et Sp. iii. 38 (1829) = **Pentarhaphia**, Lindl. (Gesnerac.).
calycina, Mart. l. c. 38 = P. calycina.
calycosa, Hook. Ic. Pl. t. 689-690 = P. calycosa.
cassioides, Benth. in DC. Prod. x. 511, lapsu, in syn. = Seymeria tenuifolia.
celsoides, Griseb. Cat. Pl. Cub. 200.—Cuba.
clandestina, Griseb. Fl. Brit. W. Ind. 461.—Jamaica.
corrugata, Griseb. in Mem. Am. Acad. N. S. viii. (1863) 526.—Cuba.
Craniolaria, Mart. Nov. Gen. et Sp. iii. 38 = P. Craniolaria.
cuneifolia, DC. Prod. vii. 526 = P. cuneifolia.
depressa, Griseb. Cat. Pl. Cub. 200.—Cuba.
exserta, Mart. Nov. Gen. et Sp. iii. 38 = P. exserta.
floribunda, Paxt. Mag. Bot. xv. (1849) 99 = P. libanensis.
gloxinioides, Griseb. Cat. Pl. Cub. 200.—Cuba.
hispida, Benth. Pl. Hartw. 264 = P. hispida.
humilis, Mart. Nov. Gen. et Sp. iii. 38 = P. humilis.
libanensis, Griseb. Fl. Brit. W. Ind. 462 = P. libanensis.
mimuloides, Griseb. l. c. 461.—Jamaica.
neglecta, Hook. Ic. Pl. t. 874 = P. negleeta.
pedunculosa, DC. Prod. vii. 525 = P. pedunculosa.
pumila, Mart. Nov. Gen. et Sp. iii. 38 = P. pumila.
reticulata, Griseb. Cat. Pl. Cub. 201.—Cuba.
scabra, Mart. Nov. Gen. et Sp. iii. 38 = P. scabra.
Sloanei, DC. Prod. vii. 526 = P. Sloanei.
ventricosa, Mart. Nov. Gen. et Sp. iii. 38 = P. longiflora.
verrucosa, Scheidw. in Otto & Dietr. Allg. Gartenz. xv. (1847) 226 = P. verrucosa.

CONRADIA, Nutt. in Journ. Acad. Philad. vii. (1834) 88. t. 12 = **Macranthera**, Torr. (Scrophul.).
fuchsioides, Nutt. l. c. = M. fuchsioides.

CONRADIA, Rafin. Neog. 3 (1825) = **Tofieldia**, Huds. (Liliac.).

CONRADINA, A. Gray, in Proc. Am. Acad. viii. (1870) 294. *LABIATAE*, Benth. & Hook. f. ii. 1191.
canescens, *A. Gray, l. c.* 295.—Am. bor.

CONRINGIA, Heist. ex Linn. Syst. ed. I (1735); Link, Enum. Hort. Berol. ii. 172 (1822). *CRUCIFERAE*, Benth. and Hook. f. i. 78.
GORINKIA, J. & C. Presl, Fl. Čech. 140 (1819).
alpina, *Link, Enum. Hort. Berol.* ii. 172.—Europ.
austriaca, *Sweet, Hort. Brit.* ed. I. 25.—Austria; Reg. Caucas.
clavata, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 84.—As. Min.; Syria.
grandiflora, *Boiss. & Heldr. Diagn.* Ser. I. viii. 25.—As. Min.
nana, Boiss. l. c. Ser. II. v. 25 = Malcolmia conringioides.
orientalis, *Dum. Fl. Belg.* 123.—Europ.; Oriens.
perfoliata, Link, Enum. Hort. Berol. ii. 172 = orientalis.
persica, *Boiss. Diagn.* Ser. I. vi. 12.—Persia.
planisiliqua, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* iii. 32.—As. occ.
Thaliana, Reichb. Ic. Fl. Germ. ii. t. 60. fig. 4380 = Sisymbrium Thalianum.

CONSANA, Adans. Fam. ii. 420 (1763) = **Draba**, Linn. (Crucif.).

CONSOLEA, Lem. Rev. Hortic. (1862) 174 = **Opuntia**, Tourn. (Cacteae).
catacantha, Lem. l. c. = O. catocantha.
ferox, Lem. l. c. = O. ferox.
leucacantha, Lem. l. c. = O. leucacantha.
rubescens, Lem. l. c. = O. rubescens.
spinosissima, Lem. l. c. = O. spinosissima.

CONSOLIDA, S. F. Gray, Nat. Arr. Brit. Pl. ii. 711 (1821) = **Delphinium**, Tourn. (Ranunculac.).
Aconiti, Lindl. in Journ. Hort. Soc. vi. (1851) 55 = D. Aconiti.
Ajacei, Schur. in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = D. Ajacis.
arvensis, Opiz, Seznam, 32 = D. Consolida.

CONSOLIDA:—
paniculata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = Delphinium paniculatum.
regalis, S. F. Gray, Nat. Arr. Brit. Pl. ii. 711 = D. Consolida.
segetum, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 47 = D. Consolida.

CONSOLIDA, Riv. ex Rupp. Fl. Jen. ed. Hall. 8 (1745) = **Symphytum**, Tourn. (Boragin.).
major, Gilib. Fl. Lituan. i. 24 = S. officinale.

CONSTANTIA, Rodrig. Orch. Nov. i. 78 (1877). *ORCHIDEAE*.
rupestris, *Rodrig. l. c.* 79.—Bras.

CONSUEGRA, Spreng. Neue Entdeck. ii. 303 (1821) = seq.

CONSUEGRIA, Mutis, ex Caldas, in Nuev. Seman. Gran. III. (1810) 22 (Quid?).

CONTARENA, Adans. Fam. ii. 120 (1763) = **Corymbium**, Linn. (Compos.).

CONTARENIA, Vand. Fl. Lusit. et Bras. Spec. 42. t. 3. f. 20 (1788)(Gen. delend.). Benth. & Hook. f. ii. 1137.
trinervia, Vand. ex Roem. & Schult. Syst. iii. 294.—Bras.

CONTORTUPLICATA, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 378 = **Astragalus**, Tourn. (Legumin.).
astragaloides, Medic. l. c. = A. contortuplicatus.

CONTRARENIA, Jaume St. Hil. Expos. ii. 346 (1805) = CONTARENIA, Vand.

CONULEUM, A. Rich. in Mém. Soc. Hist. Par. i. (1823) 391. t. 25. *MONIMIACEAE*, Benth. & Hook. f. iii. 143.
guianensis, *A. Rich. l. c.*—Guiana.

CONVALLARIA, Linn. Syst. ed. I (1735); Gen. ed. I. 96 (1737). *LILIACEAE*, Benth. & Hook. f. iii. 770.
AXILLARIA, Rafin. in Journ. Phys. lxxxix. (1819) 102, 261.
GLOBERIA, Rafin. Med. Fl. ii. 84 (1830).
LILIUM-CONVALLIUM, Tourn. ex Moench, Meth. 636 (1794).
ambigua, Schrank, Baier. Fl. i. 606 = Polygonatum multiflorum.
angulosa, Lam. Fl. Fr. iii. 268 = Polygonatum officinale.
angustifolia, Poir. Encyc. Suppl. v. 737 = Polygonatum biflorum.
biflora, Walt. Fl. Carol. 122 = Polygonatum biflorum.
bifolia, Linn. Sp. Pl. 316 = Smilacina bifolia.
borealis, Poir. Encyc. Suppl. v. 737 = Clintonia borcalis.
bracteata, Dulac, Fl. Hautes-Pyr. 135 = majalis.
bracteata, Thomas, in Gaud. Fl. Helv. ii. 531 = Polygonatum multiflorum.
Broteri, Guss. Fl. Sic. Syn. i. 416 = Polygonatum Gussonii.
canadensis, Poir. Encyc. Suppl. iv. 30 = Smilacina bifolia.
canaliculata, Willd. Enum. Hort. Berol. 45 = Polygonatum biflorum.
ciliata, Poir. Encyc. Suppl. iv. 30 = Smilacina racemosa.
cirrhifolia, Wall. in As. Res. xiii. (1820) 382 = Polygonatum sibiricum.
commutata, Schult. f. Syst. vii. 1671 = Polygonatum latifolium.
compressa, Steud. Nom. ed. II. i. 406 = Polygonatum japonicum.
cordifolia, Stokes, Bot. Mat. Med. ii. 265 = Smilacina bifolia.
dichotoma, Thib. ex Pers. Syn. i. 373 = Streptopus distortus.
diffusa, Vell. Fl. Flum. 143; iii. t. 160 = Dichorisandra pubescens.
fragrans, Salisb. Prod. 254 = majalis.
fruticosa, Linn. Syst. Nat. ed. X. 984 = Cordyline terminalis.
gigantea, Hort. ex Kunth, Enum. Pl. v. 136 = Polygonatum latifolium.
glabra, *Kit. in Linnaea*, xxxii. (1863) 347.—Europ.
Govaniana, Wall. Cat. n. 5137 = Polygonatum multiflorum.

CONVALLARIA:—
graminifolia, Salisb. Prod. 254 = Ophiopogon japonicus.
hirta, Bosc, ex Poir. Encyc. iv. 369 = Polygonatum latifolium.
hybrida, March. in Van Hall, Bijdr. iv. 343 = Smilacina stellata.
japonica, Linn. f. Suppl. 204 = Ophiopogon japonicus.
Keiskei, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 148.—Japon.
latifolia, Jacq. Fl. Austr. iii. 18. t. 232 = Polygonatum latifolium.
latifolia, Lam. ex Steud. Nom. ed. II. i. 406 = Polygonatum multiflorum.
latifolia, Mill. Gard. Dict. ed. VIII. n. 2 = majalis.
latifolia, Savi, Bot. Etrusc. ii. 222 = Polygonatum officinale.
leptophylla, D. Don, Prod. Fl. Nep. 47 = Polygonatum verticillatum.
majalis, *Linn. Sp. Pl.* 314.—Reg. bor. temp.
Mappii, C. C. Gmel. Fl. Bad. ii. 52 = majalis.
monophylla, Oed. in Fl. Dan. t. 291 = Smilacina bifolia.
multiflora, Linn. Sp. Pl. 315 = Polygonatum multiflorum.
obtusifolia, Guenth. ex Steud. Nom. ed. II. i. 406 = Polygonum officinale.
odorata, Mill. Gard. Dict. ed. VIII. n. 4 = Polygonatum officinale.
oppositifolia, Wall. in As. Res. xiii. (1820) 380 = Polygonatum oppositifolium.
orientalis, Poir. Encyc. Suppl. iv. 30 = Polygonatum polyanthemum.
parviflora, Poir. l. c. 29 = Polygonatum officinale.
polyanthema, Bieb. Fl. Taur. Cauc. iii. 272 = Polygonatum polyanthemum.
Polygonatum, Linn. Sp. Pl. 315 = Polygonatum officinale.
Polygonatum, Muhl. Cat. 35 = Polygonatum latifolium.
Polygonatum, Thunb. Fl. Jap. 142 = Polygonatum japonicum.
pubescens, Willd. Hort. Berol. i. 45 = Polygonatum biflorum.
pulchella, Salisb. Prod. 255 = Smilacina bifolia.
punctata, Wall. Cat. n. 5133 = Polygonatum punctatum.
quadrifida, Lam. Fl. Fr. iii. 269 = Smilacina bifolia.
racemosa, Forsk. Fl. Aegypt. Arab. 73 = Smilacina Forskaliana.
racemosa, Linn. Sp. Pl. 315 = Smilacina racemosa.
racemosa, Vell. Fl. Flum. 142; iii. t. 159 = Dichorisandra Gaudichaudiana.
rosea, Ledeb. Fl. Alt. ii. 41 = Polygonatum roseum.
rupestris, Salisb. Prod. 254 = Polygonatum officinale.
scaposa, Gilib. Exercit. ii. 460 = majalis.
scepusiensis, Kit. in Linnaea, xxxii. (1863) 347 = glabra.
sibirica, Schult. f. Syst. vii. 1, 297 = Polygonatum sibiricum.
spicata, Thunb. Fl. Jap. 141 = Liriope graminifolia.
stellata, Linn. Sp. Pl. 316 = Smilacina stellata.
stellifolia, Peterm. in Flora, xxvii. (1844) 363 = Polygonatum verticillatum.
tetrapetala, Gilib. Exercit. ii. 461 = Smilacina bifolia.
Thunbergii, Steud. Nom. ed. II. i. 406 = Polygonatum latifolium.
trifolia, Linn. Sp. Pl. 316 = Smilacina trifolia.
umbellata, Bunge, ex Baker, in Journ. Linn. Soc. xiv. (1875) 553 = Polygonatum umbellatum.
umbellulata, Michx. Fl. Bor. Am. i. 202 = Clintonia umbellata.
verticillata, Linn. Sp. Pl. 315 = Polygonatum verticillatum.

CONVOLVULOIDES, Moench, Meth. 451 (1794) = **Ipomoea**, Linn. (Convolv.).
elongata, Moench, l. c. 452 = I. sibirica.
leucosperma, Moench, l. c. = I. purpurea.
palmata, Moench, l. c. = I. Pes-tigridis.
pilosa, Moench, l. c. = I. pubescens.
purpurea, Moench, l. c. = I. purpurea.
triloba, Moench, l. c. = I. hederacea.

CONVOLVULUS, [Tourn.] Linn. Syst. ed. I (1735). *CONVOLVULACEAE*, Benth. & Hook. f. ii. 874.
BUCHAREA, CALIXNOS, CAULOTULIS, CLEISOSTOMA, DITEREIA, EMPLINA, EXALLOSIS, EXOCROA, IDALIA, LATRIENDA, LIZERON, NEMOSTIMA, PENTANTHUS, PERIPHAS, PODALETRA, RHODOXYLON, SANTIUM, SCIADIARA, SYMETHUS, THYELLA, Rafin. Fl. Tellur. 70–84 (1836).

CONVOLVULUS:—

MERREMIA, Dennst. Schluess. Hort. Malab. 34 (1818).

PANTOČSEKIA, Griseb. in Oestr. Bot. Zeitschr. (1873) 267.

RHODORHIZA, Webb, in Bot. Reg. (1841) Misc. 69.

STEVOGTIA, Neck. Elem. ii. 23 (1790).

abruptus, Spreng. Syst. i. 596 = Ipomoea abrupta.

abutiloides, H. B. & K. Nov. Gen. et Sp. iii. 106 = Ipomoea abutiloides.

acanthocladus, *Boiss. & Kotschy, Diagn.* Ser. I. vii. 27. —Persia.

acaulis, Choisy, in DC. Prod. ix. 406 = erubescens.

acetosaefolius, Steud. Nom. ed. II. i. 400 = Ipomoea martinicensis.

acetosaefolius, Turcz. in Bull. Soc. Nat. Mosc. (1840) 73 = Calystegia hederacea.

acetosaefolius, Vahl, Eclog. Am. i. 18 = Ipomoea carnosa.

acetosellaefolius, Desr. in Lam. Encyc. iii. 564 = Ipomoea acetosellaefolia.

acicularis, *Vatke, in Linnaea*, xliii. (1880–82) 518.— Afr. trop.

aculeatus, D. Dietr. Syn. Pl. i. 667 = Ipomoea Bona-nox.

aculeatus, *Hill, Veg. Syst.* vii. 10 (= seq.?).—Am. merid.

culeatus, *Linn. Sp. Pl.* 155.—Am. trop.

acuminatus, Salzm. ex Meissn. in Mart. Fl. Bras. vii. 267 = Ipomoea fastigiata.

acuminatus, Vahl, Symb. Bot. iii. 26 = Ipomoea acuminata.

acutangulus, Spreng. Syst. i. 605 = Ipomoea coccinea.

Adansonii, Desr. in Lam. Encyc. iii. 560 = Ipomoea aquatica.

adpressus, Wall. Cat. n. 1424 = Lettsomia adpressa.

adscendens, De Vriese, in Lehm. Pl. Preiss. i. 346 = erubescens.

adscendens, Stend. ex Boiss. Fl. Orient. iv. 100 = reticulatus.

aegyptius, *Linn. Syst.* ed. X. 923.—Aegypt.

affghanus, *Vatke, in Oestr. Bot. Zeitschr.* xxv. (1875) 168.—Afghan.

affinis, *Scheele, in Linnaea*, xxi. (1848) 749.— Texas.

affinis, C. Sm. ex Steud. Nom. ed. II. i. 407 = Tuckeyanus.

affinis, Wall. Cat. n. 1385 = Ipomoea polyantha.

africanus, Nickols, Hort. St. Doming. 260 = Ipomoea cathartica.

aggregatus, Lour. Fl. Cochinch. 109 = Ipomoea aggregata.

aggregatus, Steud. Nom. ed. II. i. 407 = Lettsomia aggregata.

agrestis, Mart. ex Choisy, in DC. Prod. ix. 405 = Jacquemontia agrestis.

agrophilus, C. Koch, in Linnaea, xxii. (1849) 745 = galaticus.

agrostopolis, Vell. Fl. Flum. 71; ii. t. 51 = Breweria Burchellii.

Aitchisoni, *C. B. Clarke, in Journ. Linn. Soc.* xix. (1882) 179.—Afghan.

alatus, Ham. Prod. Fl. Ind. Occ. 24 = Ipomoea Turpethum.

alatus, Spreng. Syst. i. 596 = Ipomoea alata.

albicans, Wall. Cat. n. 1401 = Ipomoea pilosa.

albiflorus, West, Bidr. Ste Croix, 271 = nodiflorus.

albivenius, Lindl. Bot. Reg. t. 1116 = Ipomoea albivenia.

albus, [*Nocca*] *Onom. Hort. Ticin.* (1813) 19.—Hab.?

alceaefolius, Bory & Chaub. Fl. Pélop. 14 = althaeoides.

alceaefolius, *Lam. Tabl. Encyc.* i. 461.—Afr. austr.

alsinoides, Linn. Sp. Pl. 157 = Evolvulus alsinoides.

althaeoides, *Linn. l. c.* 156.—Reg. Mediterr.

althaeoides, Tenore, Syll. Fl. Neap. 92 = tenuissimus.

althaeoides, Thunb. Fl. Cap. ii. 18 = Thunbergii.

altissimus, Spreng. Syst. i. 613 = Ipomoea staphylina?

amblyocalyx, Steud. Nom. ed. II. i. 407 = fatmensis.

Ammannii, *Desr. in Lam. Encyc.* iii. 549.—Sibir.

ammocharis, *Boiss. & Haussk. Pl. Or. Nov. Dec.* i. 6. —Persia.

ammophilus, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 123 = ammocharis.

amoenus, D. Dietr. Syn. Pl. i. 670 = Ipomoea amoena.

amoenus, C. Koch, in Linnaea, xix. (1846) 19 = hirsutus.

ampelopsifolius, Cham. & Schlecht. l. c. v. (1830) 118 = Ipomoea quinquefolia.

CONVOLVULUS:—

ampliatus, Herb. Brit. Mus. ex Choisy, in DC. Prod. ix. 361 = Ipomoea ampliata.

amplus, Wall. Cat. n. 1420 = Argyreia Roxburghii.

ampullaceus, Vahl, ex Choisy, Conv. Or. 61 = Lepistemon Wallichii.

anceps, Linn. Mant. i. 43 = Ipomoea Turpethum.

andinus, *Phil. in Linnaea*, xxxiii. (1864–65) 184.— Chili.

angularis, Burm. f. Fl. Ind. 46 = Ipomoea vitifolia.

angulatus, Buch.-Ham. ex Choisy, Conv. Or. 52 = Ipomoea coccinea.

angulatus, Salzm. ex Choisy, in Mém. Soc. Phys. Genèv. ix. (1837) 137 = Ipomoea bahiensis.

angulatus, Spreng. Syst. i. 594 = Ipomoea angulata.

angustifolius, Desr. in Lam. Encyc. iii. 547 = Ipomoea angustifolia.

angustissimus, R. Br. Prod. 482 = erubescens.

aphyllus, Viv. Ann. Bot. i. 95 = Ipomoea pentaphylla.

apocynoides, Cham. & Schlecht. in Linnaea, v. (1830) 117 = Jacquemontia hirsuta.

aquaticus, Walt. Fl. Carol. 94 = Breweria aquatica.

arborescens, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 204 = Ipomoea arborescens.

arboreus, Balb. ex Steud. Nom. ed. II. i. 407 = praec.

arcuatus, Presl, Fl. Sic. p. xxxiii = pentapetaloides.

arenarius, Vahl, Symb. Bot. i. 18 = Ipomoea carnosa.

argenteus, Heyne, ex Wall. Cat. n. 1394 = Argyreia fulgens.

argenteus, Lam. Fl. Fr. ii. 266; Desr. in Lam. Encyc. iii. 552 = C. Cneorum.

argenteus, Pourr. in Mém. Acad. Toul. iii. (1788) 316 = lanuginosus.

argenteus, Wall. Cat. n. 1364 = Argyreia argentea.

argyraeus, DC. Fl. Fr. Suppl. 423 = althaeoides.

argyrophyllus, Hoffmgg. Verz. Pfl. 53 = althaeoides.

argyrophyllus, Wall. Cat. n. 1395 = Argyreia lanceolata.

aristolochiaefolius, H. B. & K. Nov. Gen. et Sp. iii. 102 = Ipomoea aristolochiaefolia.

aristolochiaefolius, Mill. Gard. Dict. ed. VIII. n. 9 = Ipomoea umbellata.

aristolochiaefolius, Willd. ex Roem. & Schult. Syst. iv. 302 = Humboldtianus.

armatus, Delile, Fl. Egypte, 189. t. 18 = C. Hystrix.

armenus, *Boiss. & Kotschy, ex Boiss. Fl. Orient.* iv. 105. —Armenia; Syria.

arvensis, *Linn. Sp. Pl.* 153.—Geront. temp.

asarifolius, Desr. in Lam. Encyc. iii. 562 = Ipomoea asarifolia.

asarifolius, Salisb. Prod. 125 = Calystegia Soldanella.

ascendens, Steud. ex Choisy, in DC. Prod. ix. 400 = reticulatus.

ascendens, Walp. Rep. vi. 540 = erubescens.

asclepiadeus, Wall. Cat. n. 1396 = parviflorus.

asper, Wall. l. c. n. 1388 = Ipomoea grandiflora.

assyricus, *Griseb. Spicil. Fl. Rumel.* ii. 75, in nota.— As. Min.; Mesopotam.

asyrensis, *Kotschy, in Sitzb. Acad. Wien, Math. Nat.* lii. *Abth.* i. (1866) 260.—Oriens.

atriplicifolius, Poir. Encyc. Suppl. iii. 467 = hirsutus.

atropurpureus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 54 = Lettsomia atropurpurea.

attenuatus, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 261.—Mexic.

aturensis, H. B. & K. Nov. Gen. et Sp. iii. 96 = Ipomoea aturensis.

Aucheri, *Choisy, in DC. Prod.* ix. 402.—Syria.

aureus, D. Dietr. Syn. Pl. i. 672 = Jacquemontia hirsuta.

aureus, *Thunb. Mus. Nat. Ups.* 168.—Hab.?

auriculatus, Desr. in Lam. Encyc. iii. 540 = arvensis.

auritus, *Willd. ex Roem. & Schult. Syst.* iv. 301.— Ins. Porto Rico.

azureus, Desr. in Lam. Encyc. iii. 554; Rich. in Act. Soc. Hist. Nat. Par. i. (1792) 107 = Jacquemontia hirsuta.

babylonicus, *Tenore, Ind. Sem. Hort. Nap.* (1829) 16. —'Afr.'

bahiensis, Spreng. Syst. i. 598 = Ipomoea bahiensis.

barbarus, *Pomel, Nouv. Mat. Fl. Atl.* 294.—Afr. bor.

barbatus, Wall. Cat. n. 1391; Pl. As. Rar. ii. 45. t. 155 = Lettsomia barbata.

barbiger, Wall. Cat. n. 1404 = Lettsomia barbigera.

barlerioides, Buch.-Ham. ex Wall. l. c. n. 2256 = Ipomoea barlerioides.

Batata, Vell. Fl. Flum. 73; ii. t. 59(= seq.?).—Bras.

Batatas, Linn. Sp. Pl. 154; Amoen. Acad. vi. 121 = Ipomoea Batatas.

CONVOLVULUS:—

batataefolius, Boiss. & Haussk. ex Boiss. Fl. Orient. iv. 106 = armenus.

Batatilla, H. B. & K. Nov. Gen. et Sp. iii. 106 = Ipomoea Batatilla.

bauhiniaefolius, Salisb. Prod. 125 = Ipomoea Pes-caprae.

Beladambu, Spreng. Syst. i. 608 = Ipomoea Beladamboe.

bellus, Spreng. l. c. 590 = Ipomoea palmata.

benehoavensis, *C. Bolle, in Bonplandia*, ix. (1861) 54. —Ins. Canar.

Benthami, Wall. Cat. n. 1393 = Ipomoea linifolia.

Bentira, Buch.-Ham. ex Wall. l. c. sub n. 1358 = Ipomoea martinicensis.

Besseri, Spreng. Syst. i. 610 = lineatns.

betonicaefolius, Mill. Gard. Dict. ed. VIII. n. 20 = hirsutus.

bibracteatus, Poepp. ex Meissn. in Mart. Fl. Bras. vii. 270 = Ipomoea luxurians.

bicolor, Desr. in Lam. Encyc. iii. 564 = Ipomoea pilosa.

bicolor, Vahl, Symb. Bot. iii. 25 = Hewittia bicolor.

bicuspidatus, Fisch. ex Link, Enum. Hort. Berol. i. 201 = arvensis.

bidentatus, *Bernh. in Flora*, xxvii. (1844) II. 829.— Afr. austr.

bifidus, Vahl, Symb. Bot. iii. 30 = Ipomoea cymosa.

bifidus, Vell. Fl. Flum. 71; ii. t. 52 (= *Aniseia Velloziana*).—Bras.

biflorus, Forsk. Fl. Aegypt. Arab. 203 = Ipomoea acetosaefolia.

biflorus, *Georgi, Beschr. Russ. Reich. Nachtr.* 261 (quid?).—Tauria.

biflorus, Linn. Sp. Pl. ed. II. 1668 = Ipomoea biflora.

biglandulosus, Stokes, Bot. Mat. Med. i. 326 = Ipomoea biloba.

bignonioides, Spreng. Syst. iv. Cur. Post. 60 = Ipomoea bignonioides.

bilobatus, Roxb. Hort. Beng. 14; Fl. Ind. i. 485 = Ipomoea biloba.

binectariferus, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 47 = Lepistemon Wallichii.

Blanchetii, Steud. Nom. ed. II. i. 407 = Jacquemontia Blanchetii.

blandus, Roxb. Hort. Beng. [13]; Fl. Ind. i. 470 = Ipomoea cymosa.

Blumii, D. Dietr. Syn. Pl. i. 675 = Ipomoea angustifolia.

Boerhavia, [*Spreng.*] *Bot. Gart. Halle*, 28.—Hab.?

boerhavioides, Blanco, Fl. Filip. ed. II. 67 = Breweria valerianoides.

bogotensis, Humb. & Bonpl. ex Willd. Enum. Hort. Berol. 203 = Ipomoea bogotensis.

Boissieri, Steud. Nom. ed. II. i. 407 = nitidus.

Bona-nox, Spreng. Syst. i. 600 = Ipomoea Bona-nox.

bonariensis, Cav. Ic. v. 54. t. 480 = incanus.

Bonplandianus, Roem. & Schult. Syst. iv. 884 = Ipomoea sidaefolia.

Bonplandii, Meissn. in Mart. Fl. Bras. vii. 825 = praec.

Bourgaei, *C. Bolle, in Bonplandia*, ix. (1861) 54.— Ins. Teneriffa.

brachyphyllus, Boiss. Diagn. Ser. II. iii. 122 = scindicus.

bracteatus, Bert. & Balb. ex Choisy, in DC. Prod. ix. 359 = Ipomoea setifera.

bracteatus, Vahl, Symb. Bot. iii. 25 = Hewittia bicolor.

brasiliensis, Linn. Sp. Pl. 159 = Ipomoea Pes-caprae.

breviacuminatus, Mart. & Choisy, in DC. Prod. ix. 409 = Jacquemontia racemosa.

breviflorus, Spreng. Syst. i. 606 = Ipomoea setifera.

brevipes, *Pomel, Nouv. Mat. Fl. Atl.* 86.—Afr. bor.

brevispinus, Jaub. & Spach, Illustr. iv. 106. t. 369 = fruticosus.

Brownii, Spreng. Syst. i. 590 = Ipomoea quinata.

Brownii, Spreng. l. c. 612 = Ipomoea heterophylla.

bryoniaefolius, Reichb. Fl. Germ. Excurs. 348 = althaeoides.

bryoniaefolius, Salisb. Prod. 125 = Ipomoea Pestigridis.

bryoniaefolius, *Sims, Bot. Mag.* t. 943.—Reg. Mediterr.

bufalinus, Lour. Fl. Cochinch. 109 = Ipomoea bufalina.

Burmanni, *Choisy, in DC. Prod.* ix. 405.—Afr. austr.

caddoënsis, Buckl. in Proc. Acad. Sc. Philad. 1862 (1863) 6 = Ipomoea leptophylla.

caeruleus, Auct. vide coeruleus.

caespitosus, Roxb. Hort. Beng. 14; Fl. Ind. i. 483 = Hewittia linifolia.

cairicus, Linn. Syst. ed. X. 922 = Ipomoea palmata.

CONVOLVULUS:—
californicus, Benth. Pl. Hartw. 326 = luteolus.
californicus, *Choisy, in DC. Prod.* ix. 405.—Am. bor. occ.
caliginosus, Koen. ex Choisy, in Mém. Soc. Phys. Genèv. vi. (1833) 461 = Ipomoea cymosa.
Calverti, *Boiss. Diagn.* Ser. II. iii. 124.—Tauria; As. Min.
calycinus, *Drège, ex Choisy, in DC. Prod.* ix. 408.—Afr. austr.
calycinus, H. B. & K. Nov. Gen. et Sp. iii. 109 = Ipomoea cissoides.
calycinus, Roxb. Hort. Beng. [13]; Fl. Ind. i. 471 = Ipomoea calycina.
calyculatus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Malaya.
calystegioides, *Choisy, Conv. Or.* 98.—China.
campanulatus, Spreng. Syst. i. 608 = Ipomoea campanulata.
campestris, Vell. Fl. Flum. 74; ii. t. 66 = Ipomoea polymorpha.
canariensis, *Linn. Sp. Pl.* 155.—Ins. Canar.
candicans, Roem. & Schult. Syst. iv. 302 = Rivea hypocrateriformis.
candicans, Roth, Nov. Pl. Sp. 106 = Rivea ornata.
candicans, Soland. ex Sims, in Bot. Mag. t. 1603 = Ipomoea fastigiata.
canescens, H. B. & K. Nov. Gen. et Sp. iii. 99 = Jacquemontia violacea.
canescens, *Phil. in Linnaea*, xxxiii. (1864–65) 182.—Chili.
canescens, Roth, Nov. Pl. Sp. 107 = Lettsomia aggregata.
cantaber, Pall. Phys. Topogr. Gem. Taur. 99 = C. Cantabrica?
Cantabrica, *Linn. Sp. Pl.* 158.—Europ.; Oriens.
capensis, *Burm. f. Fl. Cap. Prod.* 5.—Afr. austr.
capensis, Willd. Sp. Pl. i. 877 = hastatus.
capillaceus, H. B. & K. Nov. Gen. et Sp. iv. 260 = Ipomoea muricata.
capitatus, Cav. Ic. ii. 72. t. 189 = lanuginosus.
capitatus, Desv. in Lam. Encyc. iii. 554 = Jacquemontia tamnifolia.
capitatus, Schimp. ex T. Anders. in Journ. Linn. Soc. v. Suppl. i. (1860) 25 = glomeratus.
capitatus, Vahl, Symb. Bot. iii. 28 = Lettsomia strigosa.
capitellatus, Buch.-Ham. ex Wall. Cat. n. 2250 = Ipomoea Pes-tigridis.
capitiformis, Poir. Encyc. Suppl. iii. 469 = Lettsomia strigosa.
capitulifcrus, *Franch. in Révoil, Çomali*, 41.—Afr. or. trop.
capripes, Stokes, Bot. Mat. Med. i. 327 = Ipomoea Pes-caprae.
Caput-Medusae, *Lowe, in Ann. & Mag. Nat. Hist.* Ser. III. vi. (1860) 155.—Ins. Madera.
caracasanus, Willd. ex Roem. & Schult. Syst. iv. 301 = Ipomoea umbellata.
cardiosepalus, *Boiss. Fl. Orient.* iv. 96.—Cilicia.
carneus, Spreng. Syst. i. 602 = Ipomoea carnea.
carnosus, Spreng. l. c. 609 = Ipomoea carnosa.
carolinus, Linn. Sp. Pl. 154 = Ipomoea commutata.
cataonicus, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov. dec.* i. 5.—As. Min.
Catesbaei, Spreng. Syst. i. 603 = Calystegia sepium.
catharticus, Blanco, Fl. Filip. ed. I. 94 = Ipomoea grandiflora.
catharticus, *Crantz, Inst.* ii. 338.—Am. trop.?
Cavanillesii, Spreng. Syst. i. 590 = Ipomoea Cavanillesii.
cephalanthus, Wall. Cat. n. 1402 = Lepistemon Wallichii.
cephalophorus, *Boiss. Diagn.* Ser. I. vii. 22.—Persia.
cephalopodus, Boiss. l. c. 24 = sericeus.
cespitosus, Ehrenb. ex Boiss. Fl. Orient. iv. 97 = libanoticus.
chenopodioides, *Desr. in Lam. Encyc.* iii. 544.—Hab.?
Cherleri, Agardh, ex Roem. & Schult. Syst. iv. 261 = arvensis.
chilensis, Pers. Syn. i. 180 = dissectus.
chinensis, Ker-Gawl. in Bot. Reg. t. 322 = arvensis.
Choisyanus, Wall. Cat. n. 2259 = Argyreia Leschenaultii.
cholulensis, Spreng. Syst. i. 599 = Ipomoea coccinea.
chondrilloides, *Boiss. Diagn.* Ser. I. xi. 83.—Persia.
Choysianus, Steud. Nom. ed. II. i. 408, lapsu = *Choisyanus*.

CONVOLVULUS:—
chryseides, Spreng. Syst. i. 598 = Ipomoea chryseides.
chrysorrhizus, Schrad. (vel Soland.) ex Steud. Nom. ed. II. i. 108, 109 = Ipomoea mammosa?
ciliatus, Roth, Catalect. fasc. i. 39 = undulatus.
ciliatus, Vahl, Eclog. Am. ii. 13 = Jacquemontia tamnifolia.
ciliolatus, Michx. Fl. Bor. Am. i. 137 = Ipomoea pandurata.
cinerascens, Spreng. Syst. i. 609 = Ipomoea incisa.
circinnatus, Willd. ex Roem. & Schult. Syst. iv. 302 = Ipomoea parasitica.
cirrhosus, *R. Br. in Salt, Abyss. App.* p. lxiv; *ed. Germ.* 475.—Abyss.
cissoides, Lam. Tabl. Encyc. i. 462; Vahl, Eclog. Am. ii. 15 = Ipomoea cissoides.
cladotrichus, *Mart. Del. Sem. Hort. Monac.* (1846) 4.—Mexic.
Clappertonii, Spreng. Syst. iv. 62 = Ipomoea Clappertonii.
clavatus, Pav. ex Choisy, in DC. Prod. ix. 46 (= *Calonyction clavatum*).—Bras.
Cneorum, Forsk. Fl. Aegypt. Arab. p. lxiii = lanatus.
Cneorum, Habl. Taur. 147, ex Bieb. Fl. Taur. Cauc. i. 146 = Calverti.
Cneorum, *Linn. Sp. Pl.* 157.—Europ. austr.
Cneorum, Sprun. ex Nym. Consp. 505 = oleaefolius.
coccineus, Salisb. Prod. 124 = Ipomoea coccinea.
cochlearis, *Griseb. Spicil. Fl. Rumel.* ii. 76, in nota.—Graec.; As. Min.
coelestis, Forst. f. Prod. 14 = Ipomoea hederacea.
coelesyriacus, *Boiss. Diagn.* Ser. I. xi. 85.—Syria.
coeruleus, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 259 = Jacquemontia parviflora.
coeruleus, Schum. & Thonn. Beskr. Guin. Pl. 101 = Ipomoea ovalifolia.
coeruleus, Spreng. Syst. i. 593 = Ipomoea hederacea.
coluhrinus, Blanco, Fl. Filip. ed. II. 66 = Ipomoea muricata.
commatophyllus, Steud. ex Choisy, in DC. Prod. ix. 354 = Ipomoea commatophylla.
Commersonii, Steud. Nom. ed. II. i. 408 = Evolvulus incanus.
commutatus, *Boiss. Diagn.* Ser. I. xi. 81.—Mesopot.
compactus, *Boiss. l. c.* iv. 40.—As. Min.
condensatus, Bertol. Misc. Bot. xv. 18 = Jacquemontia tamnifolia.
confertus, Hort. Hamb. ex Steud. Nom. ed. II. i. 408 = Argyreia setosa.
congestus, *R. Br. in Salt, Abyss. App.* p. lxiv.—Abyss.
congestus, Spreng. Syst. i. 601 = Ipomoea congesta.
contortus, Vell. Fl. Flum. 76; ii. t. 48 = Ipomoea glabra.
copticus, Linn. Mant. ii. 559 = Ipomoea dissecta.
cordatifolius, Vell. Fl. Flum. 73; ii. t. 60 = Ipomoea Batatas.
cordifolius, Bacle, ex Choisy, in DC. Prod. ix. 381 = Ipomoea Baclii.
cordifolius, Juss. ex Choisy, l. c. 412 = farinosus.
cordifolius, Salzm. ex Meissn. in Mart. Fl. Bras. vii. 267 = Ipomoea fastigiata.
cordifolius, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
coriaceus, Herb. Brit. Mus. ex Choisy, in DC. Prod. ix. 385 = Ipomoea pentaphylla.
Correjuela, *Steud. Nom.* ed. II. i. 408.—Chili.
corsicus, Roem. & Schult. Syst. iv. 256 = arvensis.
corymbosus, Forst. f. Prod. 14 = parviflorus.
corymbosus, Linn. Syst. ed. X. 923 = Ipomoea Burmanni.
costatus, Meyen, Reise, i. 264 (= *Aniseia costata*).—Chili.
crassifolius, *Schcele, in Linnaea*, xxi. (1849) 752.—Bras.
crassipes, Kunze, ex Choisy, in DC. Prod. ix. 433 = Calystegia sepium.
crenatifolius, *Ruiz & Pav. Fl. Per.* ii. 10.—Peruv.
crenatus, Jacq. Coll. ii. 277 = Hermanniae.
crenatus, Vahl, Symb. Bot. iii. 31 = Vahlii.
crinitus, Desr. in Lam. Encyc. iii. 568 = Jacquemontia tamnifolia.
crispatulus, Wall. Cat. n. 1403 = Ipomoea petaloidea.
crispifolius, *F. Muell. in Linnaea*, xxv. (1852) 423.—Austral.
crispus, *Thunb. Fl. Cap.* ii. 15.—Afr. austr.
cumanensis, H. B. & K. Nov. Gen. et Sp. iii. 99 = Ipomoea ferruginea.

CONVOLVULUS:—
cuneatus, Willd. Sp. Pl. i. 873 = Argyreia cuneata.
cuneiformis, Buch.-Ham. ex Wall. Cat. p. 84. n. 1366 = Argyreia cuneata.
Cupanianus, Tod. in Ann. Sc. Nat. Sér. IV. xx. (1863) 304 = tricolor.
Curtana, Buch.-Ham. ex Wall. Cat. p. 83, n. 1344 = Ipomoea eriocarpa.
cuspidatus, Willd. ex Spreng. Syst. i. 597 = Meyeri.
cymosus, Desr. in Lam. Encyc. iii. 556 = Ipomoea cymosa.
cymosus, Herb. Madr. ex Wall. Cat. p. 84, n. 1366 = Argyreia cuneata.
cymosus, Roem. & Schult. Syst. iv. 303 = Senegambiae.
cymosus, Ruiz & Pav. Fl. Per. ii. 9 = Ipomoea ramosissima.
cynanchifolius, Wall. Cat. n. 1399 = Ipomoea cynanchifolia.
cynanchifolius, Willd. ex Roem. & Schult. Syst. iv. 301 = mollis, Ipomoea Bona-nox.
cyprius, *Boiss. Fl. Orient.* iv. 93.—Ins. Cyprus.
dahuricus, Herb. in Bot. Mag. t. 2609 = Calystegia dahurica.
damascenus, Boiss. & Gaill. Diagn. Ser. II. vi. 122 = stachydifolius.
dasycephalus, Pall. ex Choisy, in DC. Prod. ix. 401 = lanuginosus.
dasyspermus, Spreng. Syst. i. 591 = Ipomoea dasysperma.
decumbens, Orteg. ex Steud. Nom. ed. II. i. 408 = undulatus.
defloratus, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 136, 132.—Malaya.
demissus, *Choisy, in DC. Prod.* ix. 404.—Chili.
densiflorus, Hook. & Arn. Bot. Beech. Voy. 303 = Ipomoea umbellata.
dentatus, Blanco, Fl. Filip. ed. I. 89 = Ipomoea Blancoi.
dentatus, Vahl, Symb. Bot. iii. 25 = Ipomoea chryseides.
denticulatus, Desr. in Lam. Encyc. iii. 540 = Ipomoea denticulata.
denticulatus, Spreng. Syst. i. 603 = Ipomoea angustifolia.
denticulatus, Steud. Nom. ed. II. i. 408 = Ipomoea chryseides.
deserti, Hochst. & Steud. ex Steud. l. c. = microphyllus.
dianthoides, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 708 = C. Pseudo-Cantabrica.
Dianthus, J. F. Gmel. Syst. 343 = Ipomoea littoralis.
dichrous, Roem. & Schult. Syst. iv. 263 = Ipomoea pilosa.
digitatus, Buch.-Ham. ex Wall. Cat. p. 85. n. 1423 = Ipomoea cissoides.
digitatus, Roxb. Hort. Beng. 14; Fl. Ind. i. 479 = Ipomoea palmata.
digitatus, Spreng. Syst. i. 592 = Ipomoea digitata.
Dillenii, Desr. in Lam. Encyc. iii. 544 = Ipomoea hederacea.
discolor, H. B. & K. Nov. Gen. et Sp. iii. 105 = Ipomoea discolor.
dissectus, Cav. Ic. v. 54 = incanus.
dissectus, Herb. Madr. ex Wall. Cat. p. 84, n. 1351 = Ipomoea dissecta.
dissectus, Jacq. Obs. Bot. ii. 4 = Ipomoea sinuata.
distillatorius, Blanco, Fl. Filip. ed. I. 95 = Ipomoea campanulata.
divaricatus, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 337, 338.—Turkestan.
divaricatus, Wall Cat. n. 1422 = arvensis.
diversifolius, Kunze, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 367 = incanus.
diversifolius, Schum. & Thonn. Beskr. Guin. Pl. 94 = Ipomoea sagittata.
diversifolius, Spreng. Syst. i. 592 = Ipomoea diversifolia.
divisus, Vell. Fl. Flum. 71; ii. t. 49 = Jacquemontia menispermoides.
domingensis, Desr. in Lam. Encyc. iii. 554 = Ipomoea tubata.
domingensis, Spreng. Neue Entdeck. iii. 29 = Ipomoea Hispaniolae.
dorycnioides, *Den. Rep. Lig.* 283.—Ligur.; Ins. Palm.
Dorycnium, *Linn. Syst.* ed. X. 923.—Graecia.
Dorycnium, Pall. ex Steud. Nom. ed. II. i. 408 = lanuginosus?

CONVOLVULUS:—
Dregeanus, *Choisy, in DC. Prod.* ix. 411.—Afr. austr.
Dryandri, Spreng. Syst. i. 597 = Massoni.
Dudia, Buch.-Ham. ex Wall. Cat. p. 83. n. 1328 = Calystegia hederacea.
dumetorum, H. B. & K. Nov. Gen. et Sp. iii. 101 = Ipomoea dumetorum.
Durandoi, *Pomel, Nouv. Mat. Fl. Atl.* 85.—Afr. bor.
ebracteatus, Desr. in Lam. Encyc. iii. 541 = Ipomoea ebracteata.
echioides, Hochst. ex Choisy, in DC. Prod. ix. 332 = Argyreia abyssinica.
edulis, Thunb. Fl. Jap. 84 = Ipomoea Batatas.
edulis, *Vell. Fl. Flum.* 72 ; ii. t. 55.—Bras.
Eichwaldi, Boiss. Diagn. Ser. I. vii. 27, in nota = fruticosus.
elegans, Wall. Cat. n. 1392 = Breweria elegans.
elegantissimus, Mill. Gard. Dict. ed. VIII. n. 22 = althaeoides.
ellipticus, Spreng. Syst. i. 613 = Lettsomia elliptica.
elongatus, Salisb. Prod. 123 = C. Scammonia.
elongatus, *Willd. Enum. Hort. Berol.* 205.—Ins. Canar.
emarginatus, *Link, Enum. Hort. Berol.* i. 201.—Hab. ?
emarginatus, Vahl, Symb. Bot. iii. 23 = Ipomoea martinicensis.
equitans, Benth. Pl. Hartw. 16 = incanus.
erectus, Spreng. Syst. i. 612 = Ipomoea erecta.
eremophilus, *Boiss. & Buhse, in Nouv. Mem. Soc. Nat. Mosc.* xii. (1860) 148.—Persia.
erianthus, Willd. ex Spreng. Syst. i. 610 = Breweria aquatica.
erinaceus, *Ledeb. in Eichw. Casp. Cauc.* 11. t. 7.—Reg. Casp.
erinaceus, C. A. Mey. Verz. Pfl. Cauc. 102 = eremophilus.
eriocarpus, Spreng. Syst. i. 598 = Ipomoea eriocarpa.
eriocaulos, Willd. ex Roem. & Schult. Syst. iv. 301 = Ipomoea purpurea.
eriospermus, Desr. in Lam. Encyc. iii. 567 = Ipomoea Desrousseauxii.
erosus, Desr. l. c. 558 = Hermanniae.
erubescens, *Sims, Bot. Mag.* t. 1067.—Austral.
erythrocarpus, Wall. Cat. n. 1413 = Argyreia lanceolata.
esculentus, Salisb. Prod. 123 = Ipomoea Batatas.
esculentus, Vell. Fl. Flum. 73 ; ii. t. 58 = septangularis.
essequebensis, Spreng. Syst. i. 600 = Ipomoea fastigiata.
evolvuloides, *Boiss. Diagn.* Ser. I. vii. 25.—Persia.
evolvuloides, Desf. Fl. Atlant. i. 176. t. 49 = undulatus.
excisus, Zipp. ex Span. in Linnaea, xv. (1841) 338 = Ipomoea reniformis.
Falckia, Thunb. Prod. Pl. Cap. 35 = Falkia repens.
Falkia, *Jacq. Hort. Schoenb.* ii. 38. t. 198.—Afr. austr.
farinosus, *Linn. Mant.* ii. 203.—Afr. trop.
fastigiatus, Herb. Roxb. ex Wall. Cat. n. 2258 = Argyreia populifolia.
fastigiatus, Roxb. Hort. Beng. [13] ; Fl. Ind. i. 468 = Ipomoea fastigiata.
fatmensis, *Kunze, in Flora*, xxiii. (1840) i. 172.—Afr. bor.
ferrugineus, Steud. Nom. ed. II. i. 408 = Jacquemontia ferruginea.
ferrugineus, Vahl, Eclog. Am. i. 17 = Ipomoea ferruginea.
festivus, Wall. Cat. n. 1414 = Argyreia argentea, populifolia.
filicaulis, Vahl, Symb. Bot. iii. 24 = Ipomoea angustifolia.
filifolius, Hook. & Arn. Bot. Beech. Voy. 35 = Ipomoea Cruckshanksii.
filiformis, Desr. in Lam. Encyc. iii. 555 = Ipomoea filiformis.
filiformis, Thunb. Fl. Cap. ii. 16 = Ipomoea angustifolia.
filipes, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 82.—Ins. Socotr.
Finlaysonianus, Wall. Cat. n. 6607 (Argyreiae sp.).—Cochinch.
firmus, Spreng. Syst. i. 613 = Ipomoea stans.
flagelliformis, Roxb. Hort. Beng. 14 ; Fl. Ind. i. 481 = Ipomoea Beladamboe.
flavescens, D. Dietr. Syn. Pl. i. 680 = Ipomoea flavescens.
flavus, Heyne, ex Wall. Cat. sub n. 1349 = Ipomoea chryseides.
flavus, Salisb. Prod. 124 = Ipomoea umbellata.

CONVOLVULUS:—
flavus, *Willd. Sp. Pl.* i. 852.—Ind. or.
flexuosus, *Pomel, Nouv. Mat. Fl. Atl.* 84.—Algeria.
flexuosus, Spreng. Syst. i. 612 = Ipomoea erecta.
floribundus, *H. B. & K. Nov. Gen. et Sp.* iii. 96.—Am. austr.
floridus, *Linn. f. Suppl.* 136.—Ins. Teneriff.
formosus, J. F. Gmel Syst. 343 = Ipomoea sagittata.
formosus, Roxb. ex Hook. f. Fl. Brit. Ind. iv. 214 = Ipomoea kentrocaulos.
Forskalei, Delile, Fl. Egypte, 190. t. 18. f. 3 = sericeus.
Forskolei, Spreng. Syst. i. 596 = Ipomoea verticillata.
Frankeanus, Schlecht. in Linnaea, vi. (1831) 613 = Jacquemontia menispermoides.
frondosus, Willd. ex Roem. & Schult. Syst. iv. 303 = havanensis.
frutescens, Mill. Gard. Dict. ed. VIII. n. 13 = Ipomoea operculata ?
frutescens, Pall. ex Ledeb. Fl. Ross. iii. 88 = fruticosus.
fruticans, Pall. Reise, ii. 474 = seq.
fruticosus, *Pall. l. c.* ii. 734.—Orient.
fruticulosus, *Desr. in Lam. Encyc.* iii. 541.—Ins. Canar.
fugacissimus, Hochst. ex Choisy, in DC. Prod. ix. 449 = Evolvulus alsinoides.
fulcratus, Greene, in Bull. Calif. Acad. i. (1885) 208 = villosus.
fulgens, Wall. Cat. n. 1394 = Argyreia fulgens.
galapagensis, *Anderss. in Vet. Acad. Handl. Stockh.* 1853 (1855) 213.—Ins. Galap.
galaticus, *Rost. ex Choisy, in DC. Prod.* ix. 408.—As. Min.
gangeticus, Linn. Cent. Pl. ii. 9 ; Amoen. Acad. iv. 306 = Evolvulus gangeticus.
gangeticus, Roxb. Hort. Beng. [13] ; Fl. Ind. i. 467 = Argyreia tiliaefolia.
gangeticus, Wall. Cat. n. 1340 = Ipomoea staphylina.
Garberi, Chapm. in Coult. Bot. Gaz. iii. (1878) 11 = havanensis.
Gaudichaudii, Choisy, Conv. Or. 68 = Ipomoea Turpethum.
gemellus, Burm. f. Fl. Ind. 46 ; Vahl, Symb. Bot. iii. 27 = Ipomoea obscura.
geniculatus, Lehm. Sem. Hort. Hamb. (1826) 17 = erubescens.
geniculatus, *Munby, ex Pomel, Nouv. Mat. Fl. Atl.* 85.—Afr. bor.
genistoides, Jaub. & Spach, Illustr. iv. 107. t. 370 = spinosus.
geranioides, *Phil. in Anal. Univ. Chil.* (1884) reimp. 7.—Chili.
Gerardi, Roem. & Schult. Syst. iv. 294 = lineatus.
Germaniciae, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 104.—As. Min.
giganteus, Silva Manso, Enum. Subst. Braz. 18 = Calystegia palmato-pinnata.
glaber, Aubl. Pl. Gui. i. 138 = Ipomoea glabra.
glaber, *Mill. Gard. Dict.* ed. VIII. n. 11.—Ind. occ.
glandulifer, Spreng. Syst. i. 599 = Ipomoea purpurea.
glandulosus, Herb. Buch.-Ham. ex Wall. Cat. n. 2252 = Lettsomia rubens.
glaucescens, H. B. & K. Nov. Gen. et Sp. iii. 101 = Ipomoea dumetorum.
glaucifolius, Choisy, in DC. Prod. ix. 412 = incanus.
glaucifolius, Spreng. Syst. i. 604 = Ipomoea glaucifolia.
glomeratus, *Choisy, in DC. Prod.* ix. 401.—Arab.
glomeratus, *Thunb. in Mus. Nat. Ups.* 168.—Hab. ?
glomeratus, Wall. Cat. n. 2257 = Argyreia cymosa.
gonatodes, Steud. ex A. Rich. Tent. Fl. Abyss. ii. 69 = Ipomoea obscura.
gonocladus, Boiss. Diagn. Ser. I. vii. 22 = Haussknechtii.
Gortschakovii, *Schrenk, Enum. Pl. Nov. Song.* 18.—Soongaria.
gossypifolius, H. B. & K. Nov. Gen. et Sp. iii. 107 = Ipomoea tuberosa.
gossypinus, Wall. Cat. n. 1407 = Ipomoea Wightii.
gracilis, Salisb. Prod. 124 = Argyreia mollis.
gracilis, Spreng. Syst. i. 604 = Ipomoea gracilis.
gramineus, Spreng. l. c. 607 = Ipomoea graminea.
grandiflorus, Desr. in Lam. Encyc. iii. 543 = Ipomoea Tuba.
grandiflorus, Hort. Brit. ex Steud. Nom. ed. II. i. 409 = Argyreia tiliaefolia.
grandiflorus, Jacq. Hort. Vindob. iii. 39 = Ipomoea latiflora.

CONVOLVULUS:—
grandiflorus, Linn. f. Suppl. 136 = Ipomoea grandiflora.
grandiflorus, Wall. Cat. n. 1370 = Ipomoea Bona-nox.
griseus, *Rafin. Atl. Journ.* 177.—Am. bor.
guadaloupensis, *Steud. Nom.* ed. II. i. 409.—Ins. Guadalup.
guianensis, Aubl. Pl. Gui. i. 136 = Jacquemontia guyanensis.
guineensis, Schum. & Thonn. Beskr. Guin. Pl. 90 = Ipomoea capitata.
guinensis, D. Dietr. Syn. Pl. i. 673 = Ipomoea pileata.
Hamiltonii, Spreng. Syst. iv. Cur. Post. 61 = Ipomoea Hamiltonii.
Hardwickii, Spreng. l. c. 60 = Ipomoea calycina.
hastatus, Desr. in Lam. Encyc. iii. 542 = Ipomoea angustifolia.
hastatus, Forsk. Fl. Aegypt. Arab. 203 = arvensis.
hastatus, Heyne, ex Wall. Cat. n. 1384 = Ipomoea sepiaria.
hastatus, Nutt. in Am. Phil. Trans. N. S. v. (1837) 194 = incanus.
hastatus, Sieber, ex Choisy, in DC. Prod. ix. 412 = fatmensis.
hastatus, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
hastifolius, Poir. Encyc. Suppl. iii. 467 = arvensis.
hastigerus, Spreng. Syst. i. 594 = Ipomoea coccinea.
Haussknechtii, *Boiss. Fl. Orient.* iv. 102.—Persia.
havanensis, *Jacq. Obs. Bot.* ii. 25.—Ind. occ.
havanensis, Willd. ex Steud. Nom. ed. II. i. 409 = Ipomoea umbellata.
hederaceus, Blanco, Fl. Filip. ed. I. 90 = Hewittia bicolor.
hederaceus, Linn. Sp. Pl. 154 = Ipomoea hederacea.
hederaceus, Mill. Gard. Dict. ed. VIII. n. 17 = Ipomoea commutata.
hederaefolius, Salisb. Prod. 123 = Ipomoea hederacea.
helianthemifolius, *Poir. Encyc. Suppl.* iii. 465.—Am. austr.
hepaticaefolius, Spreng. Syst. i. 594 = Ipomoea Pestigridis.
hepaticaefolius, Willd. ex Roem & Schult. Syst. iv. 303 = Ipomoea trifida.
heptaphyllus, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 196 = Ipomoea palmata.
Hermanniae, Choisy, in DC. Prod. ix. 408 = hermannioides.
Hermanniae, *L'Hérit. Stirp. Nov.* 67. t. 33.—Peruv.
hermannioides, *A. Gray, Syn. Fl. N. Am.* ii. 1. 216.—Texas.
heterophyllus, Ehrenb. ex Boiss. Fl. Orient. iv. 109 = coelesyriacus.
heterophyllus, Spreng. Syst. i. 592 = Ipomoea heterophylla.
Heynii, Spreng. l. c. 699 = Lettsomia aggregata.
Heynii, Wall. Cat. n. 1387 = Ipomoea sepiaria.
Hildebrandtii, *Vatke, in Linnaea*, xliii. (1880–82) 519.—Afr. trop.
hirsutissimus, Wall. Cat. n. 1400 = Lettsomia hirsutissima.
hirsutus, *Bieb. Fl. Taur. Cauc.* i. 422.—Graecia ; As. Min.
hirsutus, Roem. ex Choisy, in DC. Prod. ix. 344 = Ipomoea hederacea.
hirsutus, Roxb. Hort. Beng. 14 ; Fl. Ind. i. 479 = Ipomoea pentaphylla.
hirsutus, Tenore, Fl. Nap. i. 60. t. 15 = althaeoides.
hirtus, Herb. Madr. ex Wall. Cat. n. 2254 = Ipomoea rumicifolia.
hirtus, *Linn. Sp. Pl.* 159.—Ind. or.
hirtus, Wight, ex Steud. Nom. ed. II. i. 409 = Ipomoea eriocarpa.
Hispaniolae, Spreng. Syst. i. 590 = Ipomoea Hispaniolae.
hispidus, Vahl, Symb. Bot. iii. 29 = Ipomoea eriocarpa.
holosericeus, Bélang. ex Choisy, in DC. Prod. ix. 402 = Aucheri.
holosericeus, *Bieb. Fl. Taur. Cauc.* i. 147.—As. Min.
Hookerianus, D. Dietr. Syn. Pl. i. 666 = Ipomoea ficifolia.
Horsfalliae, D. Dietr. l. c. 664 = Ipomoea Horsfalliae.
Houstoni, Steud. Nom. ed. II. i. 408, in syn. = glaber.
Huegelii, De Vriese, in Lehm. Pl. Preiss. i. 346 = erubescens.
Humboldtianus, *Roem. & Schult. Syst.* iv. 884.—Venezuela.
humilis, Jacq. Coll. iv. 209. t. 32 = undulatus.
humilis, Salisb. Prod. 125 = lineatus.
humistratus, Walt. Fl. Carol. 94 = Breweria humistrata.

CONVOLVULUS:—
hybridus, *Zoll. & Morr. in Nat. Geneesk. Arch. Neêrl. Ind.* ii. (1845) 6.—Malaya.
hyoscyamoides, *Vatke, in Linnaea*, xliii. (1880–82) 520.—Afr. trop.
hypocrateriformis, Desr. in Lam. Encyc. iii. 561 = Rivea hypocrateriformis.
hypoleucus, Roem. & Schult. Syst. iv. 884 = Rivea hypocrateriformis.
Hystrix, *Vahl, Symb. Bot.* i. 16.—Aegypt.; Arab.
ianthinus, Hance, ex Walp. Ann. iii. 113 = parviflorus.
igneus, Schrad. in Goett. Gel. Anz. i. (1821) 716 = Ipomoea flammea.
imbricatus, Spreng. Syst. i. 613 = Lettsomia aggregata.
Imperati, Vahl, Symb. Bot. i. 17 = Ipomoea littoralis.
incanus, *Vahl, l. c.* iii. 23.—Am. bor. et austr.
incarnatus, Vahl, Eclog. Am. ii. 12 = Ipomoea incarnata.
incisus, *Herb. Mus. Par. ex Choisy, in DC. Prod.* ix. 409.—Peruv.?
incisus, Spreng. Syst. i. 609 = Ipomoea incisa.
incrassatus, Wall. Cat. n. 1386 = Ipomoea sepiaria.
incurvus, Schum. & Thonn. Beskr. Guin. Pl. 99 = Ipomoea Clappertonii.
indicus, Mill. Gard. Dict. ed. VIII. n. 5 = Ipomoea violacea.
indivisus, Vell. Fl. Flum. 71; ii. t. 50 = Ipomoea coccinea.
inflatus, Desf. Tabl. ed. I. 74 = Calystegia silvatica.
insignis, Spreng. Syst. i. 592 = Ipomoea digitata.
insuavis, D. Dietr. Syn. Pl. i. 668 = Ipomoea obscura.
intermedius, Loisel. in Journ. Bot. ii. (1809) 264 (Not. 40) = lineatus.
intermedius, Roem. & Schult. Syst. iv. 264 = Ipomoea purpurea.
involucellatus, *Klotzsch, in Peters, Reise Mossamb. Bot.* 241.—Afr. trop.
involucratus, Ker-Gawl. in Bot. Reg. t. 318 = Hewittia bicolor.
involucratus, Spreng. Syst. i. 690 = Ipomoea pileata.
involucratus, Willd. Sp. Pl. i. 845 = Ipomoea teretistigma.
Ipomoea, Vell. Fl. Flum. 72; ii. t. 56 = Ipomoea triloba.
italicus, Roem. & Schult. Syst. iv. 266 = althaeoides.
jacobacus, *C. Sm. in Tuckey, Congo*, 249, nomen.—Ins. Cap. Verd.
Jalapa, Linn. Mant. i. 42 = Ipomoea Jalapa.
jamaicensis, Jacq. Obs. iii. 6 = havanensis?
jamaicensis, Spreng. Syst. i. 595 = Ipomoea jamaicensis.
japonicus, Thunb. Fl. Jap. 85 = Ipomoea angustifolia.
Jatiauca, J. F. Gmel. Syst. 339 = Ipomoea Jalapa.
javanensis, Burm. f. Fl. Ind. 43 = Ipomoea denticulata.
javanicus, Burm. f. l. c. 50 = Ipomoea sagittaefolia.
javanicus, Spreng. Syst. i. 605 = Ipomoea hastata.
jemensis, *Kotschy, in Sitzb. Wien. Acad.* lii. (1866) t. 2.—Arab.
kentrocaulos, Steud. ex Choisy, in DC. Prod. ix. 362 = Ipomoea kentrocaulos.
Kleinii, Spreng. Syst. i. 606 = Ipomoea staphylina.
Kleinii, Wall. Cat. n. 1377 = Ipomoea sinuata.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 337, 338.—Turkestan.
Kotschyanus, *Boiss. Diagn.* Ser. I. vii. 23.—Persia.
Krauseanus, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 339.—Turkestan.
lachnaeus, Spreng. Syst. i. 606 = Ipomoea lachnaea.
lachnospermus, Hochst. in Ind. Sem. Hort. Monsp. (1847) 7 = Ipomoea lachnosperma.
laciniatus, *Desr. in Lam. Encyc.* iii. 546.—Reg. Argent.
lactescens, Gronov. ex Choisy, in DC. Prod. ix. 433 = Calystegia sepium.
lacteus, Wall. in J. Grah. Cat. Bomb. Pl. 133 = Ipomoea lactea.
lacunosus, Spreng. Syst. i. 597 = Ipomoea lacunosa.
lactus, Salisb. Prod. 123 = Calystegia sepium.
laevicaulis, Willd. ex Roem. & Schult. Syst. iv. 303 = Ipomoea sidaefolia.
laevigatus, Herb. Mus. Brit. ex Steud. Nom. ed. II. i. 409 = Ipomoea laevigata.
lanatus, Vahl, Symb. Bot. i. 16 = sericeus.
laniger, *Willd. ex Spreng. Syst.* i. 593.—Hab.?
lanuginosus, Aitch. Cat. Punjab Pl. 98 = Aitchisoni.
lanuginosus, *Desr. in Lam. Encyc.* iii. 551.—Hispan.

CONVOLVULUS:—
lanuginosus, Heyne, ex Wall. Cat. n. 1407 = Ipomoea Wightii.
lanuginosus, Ledeb. Fl. Ross. iii. 88 = Calvertii.
lanuginosus, Vahl, Symb. Bot. iii. 23 = hirsutus.
lapathifolius, *Schlecht. in Linnaea*, xix. (1847) 736.—Mexic.
lapathifolius, Spreng. Syst. i. 604 = Ipomoea acetosellaefolia.
lasianthus, *Cav. Ic.* v. 53. t. 479.—Chili.
lasiophloeus, Jaub. & Spach, Illustr. iv. 105. t. 368 = leiocalycinus.
lasiospermus, Vis. Pl. Aeg. Nub. 13. t. 1. f. 2 = Ipomoea rumicifolia.
latiflorus, Desr. in Lam. Encyc. iii. 561 = Ipomoea Bona-nox.
laurifolius, Roxb. Hort. Beng. [13]; Fl. Ind. i. 470 = Lettsomia elliptica.
laurifolius, Wight, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 186 = Argyreia obtecta.
leiocalycinus, *Boiss. Diagn.* Ser. I. vii. 28.—Persia.
leptocladus, *Boiss. l. c.* 25.—Persia.
leucanthus, Desr. in Lam. Encyc. iii. 541 = Ipomoea leucantha.
leucanthus, Herb. Madr. ex Wall. Cat. p. 83, n. 1340 = Argyreia tiliaefolia.
leucophyllus, Willd. ex Roem. & Schult. Syst. iv. 301 = Ipomoea discolor.
leucotrichus, *Pomel, Nouv. Mat. Fl. Atl.* 87.—Afr. bor.
libanoticus, *Boiss. Diagn.* Ser. I. xi. 82.—Syria.
liberalis, Lhotsky, ex Choisy, in Mém. Soc. Phys. Genèv. ix. (1837) 132 = Ipomoea echioides.
lilacinus, D. Dietr. Syn. Pl. i. 669 = Ipomoea lilacina.
limphaticus, Vell. Fl. Flum. 70 = Ipomoea palmata?
Lindleyi, Steud. Nom. ed. II. i. 409 = Ipomoea hederacea.
linearifolius, Mill. Gard. Dict. ed. VIII. n. 28 = pilosellaefolius.
linearis, Curt. Bot. Mag. t. 289 = oleaefolius.
linearis, DC. Fl. Fr. Suppl. 424 = lanuginosus.
linearis, Lam. Fl. Fr. ii. 267 = cantabrica.
lincatus, *Linn. Syst.* ed. X. 923.—Europ. austr.; As. temp.
lineatus, Sibth. & Sm. Fl. Graec. ii. 81. t. 199 = oleaefolius.
linifolius, D. Dietr. Syn. Pl. i. 675 = Ipomoea linifolia.
linifolius, Linn. Cent. Pl. ii. 10 = Evolvulus alsinoides.
linifolius, Wall. Cat. n. 1389 = Ipomoea angustifolia.
Linki, F. G. Dietr. Vollst. Lexik. Gaertn. Nachtr. II. iii. 94 = emarginatus.
littoralis, D. Dietr. Syn. Pl. i. 680, in syn. = Ipomoea littoralis.
littoralis, Linn. Syst. ed. X. 924 = Ipomoea carnosa.
littoralis, *Vatke, in Linnaea*, xliii. (1880–82) 519.—Afr. trop.
lividus, Moç. ex Steud. Nom. ed. II. i. 409 = Ipomoea Jalapa.
lobatus, Engelm. & Gray, Pl. Lindheim. i. 44 = incanus.
longiflorus, Heyne, ex Steud. Nom. ed. II. i. 409 = Ipomoea palmata.
longiflorus, Spreng. Syst. i. 595 = Ipomoea grandiflora?
longipes, *S. Wats. in Am. Natural.* vii. (1873) 302.—Am. bor. occ.
Lotnoha, Buch.-Ham. ex Wall. Cat. p. 83, n. 1342 = Ipomoea cymosa.
Loureiri, G. Don, Gen. Syst. iv. 290 = calystegioides.
lucanus, Tenore, Fl. Neap. Prod. App. v. 9 = Calystegia sepium.
lupulifolia, Griff. Notul. iv. 284 = Ipomoea palmata.
luteolus, *A. Gray, in Proc. Am. Acad.* xi. (1876) 90.—Calif.
luteolus, Spreng. Syst. i. 599 = Ipomoea coccinea.
luteolus, Wight, ex Wall. Cat. p. 84. n. 1349 = Ipomoea chryseides.
luteus, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 260.—Mexic.
lycioides, Boiss. Diagn. Ser. I. vii. 28 = leiocalycinus.
lycioides, Wall. Cat. n. 1390 = Blinkworthia lycioides.
lymphaticus, Vell. Fl. Flum. ii. t. 47 = Ipomoea palmata.
macranthus, H. B. & K. Nov. Gen. et Sp. iii. 95 = Ipomoea murucoides.
macrocalyx, Ruiz & Pav. Fl. Per. ii. 10 = Ipomoea glabra.
macrocarpus, *Linn. Syst.* ed. X. 923.—Ind. occ.

CONVOLVULUS:—
macrocarpus, Spreng. Syst. i. 592 = Ipomoea tuberosa.
macrorhizus, Ell. Sketch, i. 252 = Ipomoea Jalapa.
macrorhizus, A. Gray, ex Choisy, in DC. Prod. ix. 381 = Ipomoea fastigiata.
macrorhizus, Linn. Syst. ed. X. 923 = Ipomoea Jalapa, macrorrhiza.
macrorrhizus, Heyne, ex Wall. Cat. n. 1350 = Ipomoea paniculata.
macrosolen, Spreng. Syst. i. 603 = Ipomoea Bona-nox.
macrospermus, Willd. Sp. Pl. i. 860 = Maripa passifloroides.
macrostegius, *Greene, in Bull. Calif. Acad.* i. (1885) 208.—Calif.
Mahur, Buch.-Ham. ex Wall. Cat. p. 85, n. 1422 = arvensis.
major, Gilib. Fl. Lituan. i. 43 = Calystegia sepium.
major, Hort. ex Nichols, Dict. Gard. i. 370 = Ipomoea purpurea.
malabaricus, Linn. Sp. Pl. 155 = Argyreia malabarica.
malabaricus, Roxb. Hort. Beng. [13]; Fl. Ind. i. 469 = Ipomoea staphylina.
malabaricus, Wall. Cat. n. 1414 B = Argyreia Hookeri.
Malcolmi, Roxb. Hort. Beng. [14]; Fl. Ind. i. 474 = arvensis.
malvaceus, *Oliver, in Trans. Linn. Soc.* xxix. (1874) 117.—Afr. trop.
mammosus, Lour. Fl. Cochinch. 108 = Ipomoea mammosa.
marginalis, Buch.-Ham. ex Wall. Cat. p. 83, n. 1340 = Argyreia tiliaefolia.
marginatus, Desr. in Lam. Encyc. iii. 558 = Ipomoea sepiaria.
marginatus, Spreng. Syst. i. 603 = Calystegia marginata.
maritimus, Desr. in Lam. Encyc. iii. 550 = Ipomoea biloba.
maritimus, D. Dietr. Syn. Pl. i. 680 = Ipomoea littoralis.
maritimus, Gouan, Fl. Monsp. 27 = Calystegia sepium.
maritimus, Lam. Fl. Fr. ii. 265 = Calystegia Soldanella.
maritimus, Pall. ex Ledeb. Fl. Ross. iii. 89 = C. Cantabrica.
Martii, Spreng. Syst. i. 598 = Calystegia heterantha.
martinicensis, Jacq. Select. Am. 26 = Ipomoea martinicensis.
mascatensis, Boiss. Diagn. Ser. I. vii. 25 = virgatus.
Massoni, *A. Dietr. Gaert. Nachr.* ii. 377.—Ins. Canar.
mataxocarpus, *Spreng. Syst.* i. 606.—N. Granat.
matutinus, *West, Bidrag. Ste-Croix*, 271.—Ind. occ.
mauritanicus, *Boiss. Voy. Esp.* 418. t. 122.—Afr. bor.
Maximiliani, Nees, in Pr. Neuwied, Reise, 17 = Calystegia sepium.
maximus, Buch.-Ham. ex Wall. Cat. n. 1341 = Ipomoea staphylina.
maximus, Linn. f. Suppl. 137; Vahl, Symb. Bot. iii. 26 = Ipomoea sepiaria.
Mechoacana, Vitm. Summa Pl. i. 434 = Ipomoea Jalapa.
Medium, Linn. Sp. Pl. 156 = Ipomoea denticulata.
Medium, Roxb. Hort. Beng. 14; Fl. Ind. i. 474 = Ipomoea angustifolia.
megapotamicus, Spreng. ex Meissn. in Mart. Fl. Bras. vii. 259 = Ipomoea megapotamica.
melanostictus, Schlecht. in Linnaea, vi. (1831) 737 = Argyreia tiliaefolia.
meonanthus, Hoffmgg. & Link, Fl. Port. i. 369 = tricolor.
meslatanus, *Thunb. Mus. Nat Ups.* 168.—Hab.?
Meyeri, *Spreng. Syst.* i. 597.—Hab.?
micranthus, Riddell, Syn. Fl. W. St. 70 = Ipomoea lacunosa.
micranthus, *Roem. & Schult. Syst.* iv. 276.—Ind. occ.; Mexic.
micranthus, Willd. ex Spreng. Syst. i. 601 = Sprengelii.
microcalyx, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 220.—Assam.
microcalyx, *Pelletan, ex Steud. Nom.* ed. II. i. 410.—Mexic.
microphyllus, *Sieber, ex Spreng. Syst.* i. 611.—Aegypt.
Millerianus, Roem. & Schult. Syst. iv. Index, 821 = Ipomoea umbellata.
minimus, *Aubl. Pl. Gui.* i. 141.—Guiana.
minimus, Vitm. Summa Pl. i. 434 = Evolvulus incanus.

CONVOLVULUS:—
minor, Gilib. Fl. Lituan. i. 43 = arvensis.
minor, Hort. ex Mill. Gard. Dict. ed. VIII. sub n. 24 = tricolor.
minutiflorus, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 262 —Mexic.
modestus, Boiss. Diagn. Ser. I. xi. 82 = commutatus.
mollis, Burm. f. Fl. Ind. 44 = Argyreia mollis.
mollis, H. B. & K. Nov. Gen. et Sp. iii. 104 = Ipomoea mollis.
mollis, *Meissn. in Mart. Fl. Bras.* vii. 314.—Bras.
mollissimus, *Bertol. Misc. Bot.* xx. 11. t. 3.—Guatemala.
mollissimus, Pall. ex Choisy, in DC. Prod. ix. 403 = holosericeus.
monospermus, [*Tin.*] *Elench. Pl. Hort. Parm.* (1802) 12.—Hab.?
montevidensis, *Spreng. Syst.* i. 604.—Reg. Argent.
mucronatus, Benth. Pl. Hartw. 120 = nodiflorus.
mucronatus, Forst. f. Prod. 14 = Ipomoea palmata.
mucronifer, Choisy, in DC. Prod. ix. 414 = Jacquemontia guyanensis.
multibracteatus, Wall. Cat. n. 1408 = Argyreia tomentosa.
multifidus, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
multiflorus, Banks, ex Steud. Nom. ed. II. i. 410 = Ipomoea congesta.
multiflorus, Bess. ex Choisy, in DC. Prod. ix. 407 = arvensis.
multiflorus, H. B. & K. Nov. Gen. et Sp. iii. 100 = Ipomoea tubata.
multiflorus, Mill. Gard. Dict. ed. VIII. n. 15 = Ipomoea multiflora.
multiflorus, Poit. ex Choisy, in DC. Prod. ix. 367 = Ipomoea quinquepartita.
multivalvis, R. Br. Prod. 483 = parviflorus.
munitiflorus, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 262, sphalm. = minutiflorus.
munitus, Wall. Cat. n. 1354 = Ipomoea quinata, pentaphylla.
muricatus, Linn. Mant. i. 44 = Ipomoea muricata.
mutabilis, Blanco, Fl. Filip. ed. I. 92 = Ipomoea Bona-nox.
mutabilis, Salisb. Prod. 123 = Ipomoea purpurea.
mutabilis, Spreng. Syst. i. 593 = Ipomoea acuminata.
muticus, Steud. Nom. ed. II. i. 410 = Ipomoea grandiflora.
natalensis, *Bernh. in Flora*, xxvii. (1844) II. 829.—Afr. austr.
Neei, Spreng. Syst. i. 593 = Ipomoea peduncularis.
nemorosus, Willd. ex Roem. & Schult. Syst. iv. 303 = Ipomoea pentaphylla.
nervosus, Burm. f. Fl. Ind. 48 = Argyreia speciosa.
nervosus. *Rafin. Fl. Ludov.* 46.—Am. bor.
nevisiensis, Ham. Prod. Fl. Ind. Occ. 25 = Jacquemontia hirsuta.
Nil, Linn. Sp. Pl. ed. II. 219 = Ipomoea hederacea.
nitens, C. Koch, in Linnaea, xxii. (1849) 743 = lineatus.
nitidus, *Boiss. Elench.* 65.—Hispan.
nitidus, Desr. in Lam. Encyc. iii. 544 = Argyreia nitida.
nodiflorus, *Desr. l. c.* 557.—Mexic.
nolanaeflorus, Zipp. ex Span. in Linnaea, xv. (1841) 341 = Ipomoea pumila.
nummularius, Linn. Sp. Pl. 157 = Evolvulus alsinoides.
nummularius, Vahl, Eclog. Am. ii. 13 = Jacquemontia nummularia.
nymphaeaefolius, D. Dietr. Syn. Pl. i. 670 = Ipomoea peltata.
obdentatus, Steud. Nom. ed. II. i. 410 = Ipomoea chryseides.
oblongus, Wall. Cat. n. 1382 = Ipomoea barlerioides.
obovatus, Wall. l. c. n. 1381 = Lettsomia setosa.
obscurus, Linn. Sp. Pl. ed. II. 220 = Ipomoea obscura.
obtectus, Wall. Cat. n. 1416 = Argyreia obtecta.
obtusilobus, Michx. Fl. Bor. Am. i. 139 = Ipomoea acetosaefolia.
obvallatus, Spreng. Syst. i. 595 = Ipomoea bracteata.
occidentalis, *A. Gray, in Proc. Am. Acad.* xi. (1876) 89.—Am. bor. occ.
occidentalis, S. Wats. l. c. 118 = macrostegius.
ocellatus, *Hook. Bot. Mag.* t. 4065.—Afr. austr.
ochraceus, Lindl. Bot. Reg. t. 1060 = Ipomoea ochracea.
ochraceus, Vahl, ex Choisy, in DC. Prod. ix. 380 = Ipomoea afra.
Oenotherae, *Vatke, in Linnaea*, xliii. (1880–82) 520.—Afr. trop.
oenotheroides, Linn. f. Suppl. 157 = Ipomoea argyreioides?

CONVOLVULUS:—
officinalis, Pelletan, ex Steud. Nom. ed. II. i. 410 = Ipomoea Purga.
oleaefolius, *Desr. in Lam. Encyc.* iii. 552.—Graecia.
Olgae, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 55.—As. centr.
oligodontus, *Baker, in Journ. Linn. Soc.* xx. (1883) 212.—Madag.
operculatus, Gomez, in Mem. Ac. Lisboa, iii. (1812) Mem. Corresp. 27 = Ipomoea operculata.
orinoccensis, Willd. ex Steud. Nom. ed. II. i. 410 = Ipomoea cissoides.
orixensis, *Wall. Cat.* n. 1362 = Lettsomia aggregata.
orizabensis, Pelletan, ex Steud. Nom. ed. II. i. 410 = Ipomoea orizabensis.
ornatus, Wall. Cat. n. 1369 = Rivea ornata.
oronocensis, Willd. ex Roem. & Schult. Syst. iv. 303, vide *orinoccensis*.
osyrensis, Roth, Nov. Pl. Sp. 405 = Lettsomia aggregata.
Otites, *Willd. ex Fisch. in Cat. Hort. Gorenk.* 28.—Hab.?
Ottonis, *Meissn. in Mart. Fl. Bras.* vii. 311.—Bras.
ovalifolius, Vahl, Eclog. Am. ii. 16 = Ipomoea ovalifolia.
ovalifolius, West, Bidrag. Ste-Croix, 271 = Ipomoea quinquepartita.
ovatus, Moench, Meth. 450 = siculus.
ovatus, Reinw. ex Choisy, in DC. Prod. ix. 332 = Argyreia Guichenotii.
owariensis, Spreng. Syst. i. 594 = Ipomoea owariensis.
Owsjanianus, *Smirnow, ex Trautv. in Act. Hort. Petrop.* ix. (1884) 25, nomen.—Rossia.
oxyphyllus, *Boiss. Diagn.* Ser. I. vii. 26.—Persia.
oxysepalus, *Boiss. l. c.* 23.—Persia.
palaestinus, *Boiss. l. c.* xi. 84.—Palaestin.
pallidus, Salisb. Prod. 123 = canariensis.
pallidus, Wall. Cat. n. 1418 = Argyreia pallida.
palmatus, Mill. Gard. Dict. ed. VIII. n. 8 = Ipomoea quinquefolia.
palustris, *Cav. Ic.* vi. 80. t. 600. f. 2.—Mexic.
palustris, Herb. Ham. ex Wall. Cat. p. 85, n. 1422 = arvensis.
panduratus, Linn. Sp. Pl. 153 = Ipomoea fastigiata.
panduratus, Lour. Fl. Cochinch. 107 = Ipomoea panduraeformis.
paniculatus, Blanco, Fl. Filip. ed. I. 96 = Ipomoea tuberosa.
paniculatus, Linn. Sp. Pl. 156 = Ipomoea digitata.
paniculatus, Wall. Cat. n. 1350 = Ipomoea kentrocaulos.
pannifolius, [*Dryand. in*] *Ait. Hort. Kew.* ed. II. i. 328.—Ins. Canar.
pannosus, Spreng. Syst. i. 612 = Ipomoea erecta.
Papiria, Spreng. l. c. 592 = Ipomoea pubescens.
paradoxus, Poir. Encyc. Suppl. v. 720 = sagittatus.
parasiticus, H. B. & K. Nov. Gen. et Sp. iii. 163 = Ipomoea parasitica.
pareiraefolius, Bert. ex Spreng. Syst. i. 613 = Ipomoea pareiraefolia.
parnassicus, Boiss. & Orph. Diagn. Ser. II. iii. 125 = cochlearis.
parviflorus, Arn. ex Steud. Nom. ed. II. i. 410 = C. Retzia.
parviflorus, Desr. in Lam. Encyc. iii. 556 = micranthus.
parviflorus, Salisb. Prod. 125 = siculus.
parviflorus, *Vahl, Symb. Bot.* iii. 29.—Geront. trop.
parvifolius, Spreng. Syst. i. 611 = pluricaulis.
patens, *Clarke, Trav.* i. ed. II. *Add.* p. xxii.—Oriens.
patens, Desr. in Lam. Encyc. iii. 547 = Breweria humistrata.
pauciflorus, Willd. ex Roem. & Schult. Syst. iv. 302 = Ipomoea fastigiata.
paulistanus, Silva Manso, Enum. Subst. Braz. 17 = Ipomoea echioides.
pedatus, Forsk. ex Choisy, in DC. Prod. ix. 410 = althaeoides.
pedatus, Roxb. Hort. Beng. 14; Fl. Ind. i. 478 = Ipomoea dasysperma.
peduncularis, Boiss. Diagn. Ser. I. xi. 84 = hirsutus.
peduncularis, Wall. Cat. n. 1417 = Argyreia elliptica.
pedunculatus, Walp. Ann. iii. 112, sphalm. = hirsutus.
pellitus, Ledeb. Fl. Alt. i. 223 = Calystegia pellita.
peltatus, Linn. Sp. Pl. 1194 = Ipomoea peltata.
penangianus, Wall. Cat. n. 1425 = Lettsomia penangiana.
pendulus, Silva Manso, Enum. Subst. Bras. 17 = Ipomoea utilis.

CONVOLVULUS:—
pendulus, Spreng. Syst. i. 590 = Ipomoea palmata.
penicillatus, *A. Rich. Tent. Fl. Abyss.* ii. 74.—Abyssin.
pennatifolius, Salisb. Prod. 124 = Ipomoea Quamoclit.
pennatus, Desr. in Lam. Encyc. iii. 567 = praec.
pennifolius, Drapiez, Encycl. iii. (1835) = praec.
pensylvanicus, Schrank, in Syll. Ratisb. i. (1824) 197 = spithamaeus.
pentadactylis, Wall. Cat. n. 1367 = Ipomoea quinata.
pentagonus, Roxb. Hort. Beng. 14; Fl. Ind. i. 485 = Ipomoea cymosa.
pentanthus, Jacq. Coll. iv. 210 = Jacquemontia violacea.
pentapetaloides, *Linn. Syst.* ed. XII. 229.—Europ. austr.; As. Min.
pentaphylloides, *Schrank, in Flora*, vii. (1824) II. *Beil.* 50.—Bras.
pentaphyllus, Linn. Sp. Pl. ed. II. 223 = Ipomoea pentaphylla.
pentaphyllus, Salzm. ex Meissn. in Mart. Fl. Bras. vii. 287 = Ipomoea glabra.
perfoliatus, Schum. & Thonn. Beskr. Guin. Pl. 89 = Ipomoea involucrata.
persicus, *Linn. Sp. Pl.* 158.—As. Min.; Persia.
peruvianus, Spreng. Syst. i. 593 = Ipomoea cuspidata.
Pes-caprae, Linn. Sp. Pl. 159 = Ipomoea Pes-caprae.
Pes-tigridis, Spreng. Syst. i. 502 = Ipomoea Pestigridis.
petiolaris, Kunth, ex Spreng. I. c. 598 = Ipomoea Bona-nox.
petraeus, *Lee, ex Choisy, in DC. Prod* ix. 413.—Afr. austr.
philomega, Vell. Fl. Flum. 74; ii. t. 63 = Ipomoea capparoides.
phoeniceus, Pav. ex Meissn. in Mart. Fl. Bras. vii. 214 (= *Operculina pteropus*).—Bras.
phoeniceus, Spreng. Syst. i. 596 = Ipomoea coccinea.
physoides, *Pomel, Nouv. Mat. Fl. Atl.* 294.—Algeria.
Pickeringii, M. A. Curt. in Bost. Journ. Nat. Hist. i. (1837) 130 = Breweria Pickeringii.
pileatus, Spreng. Syst. iv. 61 = Ipomoea pileata.
piliferus, *Steud. Nom.* ed. II. i. 410.—Abyssin.
pilosella, *Liebm. Ind. Sem. Hort. Haun.* (1853) 22; *in Ann. Sc. Nat.* Sér. IV. i. (1854) 33.—Hab.?
pilosellaefolius, *Desr. in Lam. Encyc.* iii. 551.—As. Min.; Persia.
piloselloides, D. Dietr. Syn. Pl. i. 676 = pilosellaefolius.
pilosus, R. Br. in Salt, Abyss. App. p. lxiv = piliferus.
pilosus, *Rottl. in Ges. Naturf. Fr. Neue Schr.* iv. (1803) 196.—Ind. or.
pilosus, Roxb. Hort. Beng. 14; Fl. Ind. i. 473 = Ipomoea pilosa.
pilosus, Wickstr. in Vet. Acad. Handb. Stockh. (1827) 60 = Ipomoea cissoides.
pinnatifidus, *H. B. & K. Nov. Gen. et Sp.* iii. 168.—Cuba.
pinnatus, Baker, Fl. Maurit. 212, sphalm. vide pennatus.
Pinto, *Vand. Fasc. Pl.* 9.—Bras.
plantagineus, Choisy, in DC. Prod. ix. 405 = Ipomoea simplex.
platanifolius, Vahl, Sym. Bot. iii. 26 = Ipomoea fastigiata.
platensis, Spreng. Syst. i. 591 = Ipomoea digitata.
platycarpus, Cav. Ic. v. 55 = Phacelia platycarpa.
platypeltis, Zipp. ex Span. in Linnaea, xv. (1841) 338 = Ipomoea campanulata.
platyphyllus, Juss. ex Steud. Nom. ed. II. i. 410; ii. 460 = Rivea tiliaefolia.
plebejus, Spreng. Syst. i. 604 = Ipomoea plebeia.
plicatus, *Desr. in Lam. Encyc.* iii. 558.—Afr. austr.
Plumerii, *Spreng. Syst.* i. 602.—Ins. S. Doming.
pluricaulis, *Choisy, Conv. Or.* 95.—Ind. or.
polyanthus, Cham. & Schlecht. in Linnaea, v. (1830) 117 = Jacquemontia violacea.
polyanthus, Wall. Cat. n. 1378 = Ipomoea polyantha, staphylina.
polycarpus, H. B. & K. Nov. Gen. et Sp. iii. 98 = micranthus.
polyrhizos, Silva Manso, Enum. Subst. Braz. 18 = Ipomoea polyrhizos.
pomaceus, Wall. Cat. n. 1419 = Argyreia pomacea, nellygheria.
portoricensis, Spreng. Syst. i. 595 = Ipomoea cathartica.
pratensis, Buch.-Ham. ex Choisy, Conv. Or. 105 = Ipomoea linifolia.

CONVOLVULUS:—
Preissii, De Vriese, in Lehm. Pl. Preiss. i. 346 = erubescens.
procumbens, Lund, ex Choisy, in DC. Prod. ix. 352 = Ipomoea polymorpha.
procumbens, Pall. ex Roem. & Schult. Syst. iv. 295 = holosericeus.
proliferus, *Vahl, Eclog. Am.* i. 18.—Am. austr.
proliferus, Willd. ex Roem. & Schult. Syst. iv. 302 = Ipomoea tubata.
prostratus, Forsk. Fl. Aegypt. Arab. 203 = pentapetaloides.
prostratus, F. W. Schmidt, Fl. Boëm. ii. 93 = arvensis.
pruinosus, Steud. Nom. ed. II. i. 411 = Jacquemontia glaucescens.
Pseudo-Cantabrica, Schrenk, Enum. Pl. Nov. i. 21 = dianthoides.
Pseudo-Scammonia, C. Koch, in Linnaea, xxii. (1849) 746 = C. Scammonia.
Pseudo-siculus, Cav. Desc. 97 = elongatus.
Pseudo-Soldanella, Mcr. in App. Loisel. ex Bull. Férussac (1828) 111 = Calystegia Soldanella.
Pseudo-tricolor, Viv. Ann. Bot. i. II. 164 = tricolor.
pterocarpus, Bert. ex Coll. Hort. Ripul. 37 = Ipomoea martinicensis.
pterygocaulos, Steud. ex Choisy, in DC. Prod. ix. 381 = Ipomoea pterygocaulos.
pubescens, [Soland. in] Russ. Aleppo, ed. II. ii. 246 = hirsutus.
pubescens, Willd. Enum. Hort. Berol. 203 = Ipomoea pubescens.
pudibundus, Lindl. Bot. Reg. t. 999 = Ipomoea cathartica.
puellaris, *Thunb. Mus. Nat. Ups.* 168.—Hab.?
pulchellus, H. B. & K. Nov. Gen. et Sp. iii. 101 = Ipomoea fastigiata.
pulcher, D. Dietr. Syn. Pl. i. 667 = Argyreia Guichenotii.
pulcherrimus, Vell. Fl. Flum. 72; ii. t. 54 = Ipomoea Bona-nox.
pungens, Kar. & Kir. in Bull. Soc. Nat. Mosc. (1841) 709 = Gortschakovii.
puniceus, Silva Manso, Enum. Subst. Braz. 18 = Ipomoea punicea.
Purga, Wender. in Pharm. Centralb. i. (1830) 457 = Ipomoea Purga.
purpureus, Linn. Sp. Pl. ed. II. 219 = Ipomoea purpurea.
pusillus, Pall. ex Roem. & Schult. Syst. iv. 295 = Calverti.
pycnanthus, Hochst. ex Choisy, in DC. Prod. ix. 365 Ipomoea capitata.
pyrrhotrichus, Boiss. Diagn. Ser. II. iii. 122 = Haussknechtii.
quadriflorus, *Hochst. in Lorent, Wanderungen*, 335.—As. Min.
Quamoclit, Spreng. Syst. i. 591 = Ipomoea Quamoclit.
quinatus, Spreng. l. c. 590 = Ipomoea quinata.
quinqueflorus, *Vahl, Symb. Bot.* iii. 31.—Ins. Mascar.
quinquefolius, Linn. Syst. ed. X. 923 = Ipomoea quinquefolia.
quinquelobus, Lindem. in Bull. Soc. Nat. Mosc. xxiii. (1850) II. 508 = arvensis.
quinquelobus, Vahl, Symb. Bot. iii. 32 = Ipomoea palmata.
quinquenervis, Buch.-Ham. ex Wall. Cat. p. 85, n. 1396 = parviflorus.
quinquepartitus, Vahl, Eclog. Am. ii. 14 = Ipomoea quinquepartita.
racemosus, Roem. & Schult. Syst. iv. 302 = Ipomoea staphylina.
radicans, *Thunb. Fl. Cap.* ii. 15.—Afr. austr.
radicosus, *Heldr. & Sart. in Boiss. Diagn.* Ser. II. iii. 124.—Graecia.
ramosissimus, Poir. Encyc. Suppl. iii. 468 = Ipomoea ramosissima.
Rampania, Buch.-Ham. ex Wall. Cat. n. 2251 = Ipomoea eriocarpa.
refractus, *Pomel, Nouv. Mat. Fl. Atl.* 84.—Afr. bor.
Reinwardtianus, D. Dietr. Syn. Pl. i. 675 = Argyreia Guichenotii.
remotus, R. Br. Prod. 483 = erubescens.
reniformis, Cat. Hort. Carlsr. (1811) 76 (Quid?).
reniformis, Poir. Encyc. Suppl. iii. 475 = Calystegia Soldanella.
reniformis, Roxb. Hort. Beng. 14; Fl. Ind. i. 481 = Ipomoea reniformis.

CONVOLVULUS:—
repandus, Desr. in Lam. Encyc. iii. 555 = Ipomoea repanda.
repens, Linn. Sp. Pl. 158 = Calystegia sepium.
repens, Sw. Obs. 64 = Ipomoea acetosaefolia.
repens, Vahl, Symb. Bot. i. 17 = Ipomoea aquatica.
repens, Willd. Sp. Pl. i. 874 = Ipomoea reptans.
reptans, Linn. Sp. Pl. 158 = Ipomoea aquatica.
reticulatus, Buch.-Ham. ex Steud. Nom. ed. II. i. 411 = Ipomoea petaloidea.
reticulatus, *Choisy, in DC. Prod.* ix. 399.—Syria.
retusus, Colla, App. Hort. Ripul. iii. 31 = Ipomoea biloba.
Retzia, *Steud. Nom.* ed. II. i. 410.—Ind. or.
Rheedii, Wall. in Roxb. Fl. Ind. ed. Carey, ii. 70 = Ipomoea uniflora.
rhyniospermus, *Hochst. ex Choisy, in DC. Prod.* ix. 405.—Afr. trop.
Riedlei, Choisy, Conv. Or. 68 = Ipomoea Turpethum.
riparius, H. B. & K. Nov. Gen. et Sp. iii. 109 = Ipomoea cissoides.
Robertianus, Spreng. Syst. v. 192 = Ipomoea heterophylla.
roseus, H. B. & K. Nov. Gen. Sp. iii. 108 = Ipomoea digitata.
roseus, Mill. Gard. Dict. ed. VIII. n. 18 = Ipomoea fastigiata.
roseus, Pall. Reise, iii. 317 = lineatus.
rostratus, Zipp. ex Span. in Linnaea, xv. (1841) 339 = Ipomoea aquatica.
Rothii, Spreng. Syst. i. 600 = Ipomaea eymosa.
Rottleri, Spreng. Syst. iv. Cur. Post. 61 = Argyreia malabarica.
Rottlerianus, *Choisy, Conv. Or.* 95.—Ind. or.
rotundifolius, Schum. & Thonn. Fl. Beskr. Guin. Pl. 102 = Ipomoea biloba.
Roxburghii, Wall. Cat. n. 1415 = Argyreia Roxburghii.
Royeni, Steud. Nom. ed. II. i. 410, in syn. = Ipomoea pentaphylla.
rubens, Wall. Cat. n. 1421 = Lettsomia rubens.
ruber, Vahl, Eclog. Am. ii. 12 = Calystegia setifera.
rubescens, Poir. Encyc. Suppl. iii. 466 = erubescens.
rubicundus, Wall. Cat. n. 1409 = Lettsomia rubicunda.
rubrocoeruleus, D. Dietr. Syn. Pl. i. 670 = Ipomoea rubrocoerulea.
ruderarius, H. B. & K. Nov. Gen. et Sp. iii. 96 = havanensis.
rufescens, Choisy, Conv. Or. 97 = flavus.
rugosus, Rottl. in Ges. Naturf. Fr. Neue Schr. iv. (1803) 196 = Ipomoea Beladamboe.
Ruizii, Spreng. Syst. i. 594 = Ipomoea villosa.
rupestris, Buch, Beschr. Canar. Ins. 193 = Massoni.
rupestris, Pall. Reise, iii. 723 = Ipomoea sibirica.
rupestris, *Willd. Sp. Pl.* i. 846.—Sibir.
Ruprechti, *Boiss. Fl. Orient.* iv. 96.—Reg. Caucas.
sabatius, Viv. Fl. Lib. Spec. 67 = elongatus.
sagittaefolius, Salisb. Prod. 123 = arvensis.
sagittatus, Dulac, Fl. Hautes-Pyr. 439 = arvensis.
sagittatus, A. Gray, in Syn. Fl. N. Am. ii. 1. 434, = hirsutus?
sagittatus, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
sagittifer, H. B. & K. Nov. Gen. et Sp. iii. 100 = Ipomoea umbellata.
sagittifolius, Fisch. ex Choisy, in DC. Prod. ix. 407 = arvensis.
sagittifolius, Michx. Fl. Bor. Am. i. 138 = Ipomoea sagittata.
sagittifolius, *Schlecht. in Linnaea*, xxi. (1848) 747.—Texas.
sagittifolius, Sibth. & Sm. Fl. Graec. Prod. i. 133; Fl. Graec. t. 193 = hirsutus.
salicifolius, Desr. in Lam. Encyc. iii. 543 = Ipomoea martinicensis.
salicifolius, Roxb. ex Steud. Nom. ed. II. i. 411 = Ipomoea salicifolia.
Saltii, *Steud. l. c.* = congestus.
salvifolius, Sieber, ex Link, Enum. Hort. Berol. i. 203 = secundus.
sanguineus, Spreng. Syst. i. 594 = Ipomoea hederifolia.
sanguineus, Willd. ex Roem. & Schult. Syst. iv. 302 = Ipomoea superba.
sarcocarpus, Silva Manso, Enum. Subst. Braz. 49 = Ipomoea operculata.
sarmentosus, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 83.—Ins. Socotra.

CONVOLVULUS:—
sarothrocladus, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iv. 92.—Assyria.
saxatilis, Bieb. Fl. Taur. Cauc. i. 146 = Calverti.
saxatilis, Salisb. Prod. 124 = Massoni.
saxatilis, Vahl, Symb. Bot. iii. 33 = lanuginosus.
scamensis, Wall. ex Steud. Nom. ed. II. i. 411 = siamensis.
Scammonia, *Linn. Sp. Pl.* 153.—As. Min.
Scammonia, Lour. Fl. Cochinch. 106 = calystegioides.
scandens, *Delile, Ind. Sem. Hort. Monsp.* (1847) 7.—Aegypt.
Schiedeanus, D. Dietr. Syn. Pl. i. 670 = Ipomoea Schiedeana.
Schimperi, *Boiss. Diagn.* Ser. I. xi. 81.—Arabia.
schirazianus, *Boiss. l. c.* 82.—Persia.
Schottianus, *Dehnh. Cat. Pl. Hort. Camald.* ed. II. 19.—Hab.?
Schultesii, Roem. & Schult. ex Steud. Nom. ed. II. i. 411 = Ipomoea purpurea.
Schultesii, Weinw. ex Steud. l. c. = Hewittia bicolor.
scindicus, Boiss. Diagn. Ser. II. iii. 123 = microphyllus.
scindicus, *Stocks, in Hook. Kew Journ.* iv. (1852) 173.—Ind. bor. occ.
scoparius, *Linn. f. Suppl.* 135.—Ins. Teneriff.
scrobiculatus, Lindl. Bot. Reg. t. 1076 = Ipomoea scrobiculata.
secundus, *Desr. in Lam. Encyc.* iii. 553.—Aegypt.; Syria.
secundus, Ruiz & Pav. Fl. Per. ii. 10 = Jacquemontia secunda.
Selloi, *Meissn. in Mart. Fl. Bras.* vii. 314.—Bras.
Semenovii, *Herder, in Bull. Soc. Nat. Mosc.* xli. (1868) I. 74.—Sibir.
semidigynus, Roxb. Hort. Beng. [13]; Fl. Ind. i. 468 = Breweria cordata.
Senegambiae, Spreng. Syst. i. 610 = Ipomoea Senegambiae.
sepiarius, Wall. Cat. n. 1384 = Ipomoea sepiaria.
sepincola, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 123 = Calystegia sepium.
sepium, Linn. Sp. Pl. 153 = Calystegia sepium.
sepium, Pall. Reise, iii. 270 = arvensis.
septangularis, Steud. Nom. ed. II. i. 411 = Ipomoea Batatas.
septenatus, Spreng. Syst. i. 591 = Ipomoea umbellifera.
Ser, Spreng. l. c. 598 = Ipomoea sinensis.
sericeus, *Burm. f. Fl. Ind.* 49. t. 19. f. 3.—Persia.
sericeus, Choisy, in DC. Prod. ix. 400 = lanatus.
sericeus, Forsk. Fl. Aegypt. Arab. 203 = tenuissimus.
sericeus, Linn. Mant. 43 = Argyreia mollis.
sericophyllus, *T. Anders. in Journ. Linn. Soc.* v. *Suppl.* i. (1860) 25.—Arab.
serotinus, DC. Cat. Hort. Monsp. 97 = Ipomoea serotina.
serpens, Linn. Syst. ed. X. 923 = Ipomoea pentaphylla.
serpiculans, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art IV. 10.—Malaya?
serpyllifolius, H. B. & K. Nov. Gen. et Sp. iii. 95 = Ipomoea serpyllifolia.
serpylloides, *Griseb. Cat. Pl. Cub.* 207.—Cuba.
sessiliflorus, Spreng. Syst. i. 599 = Ipomoea eriocarpa.
setifer, Spreng. l. c. 597 = Calystegia setifera.
setosus, Spreng. l. c. 594 = Argyreia setosa.
setosus, Wall. Cat. n. 1412 = Lettsomia setosa.
Sherardi, Pursh, Fl. Am. Sept. ii. 30 = micranthus.
Shumardianus, *Torr. Bot. Expl. Red River* (1852) 291.—Am. bor.
siamensis, *Wall. Cat.* n. 6668.—Siam.
sibiricus, Linn. Mant. ii. 203 = Ipomoea sibirica.
Sibthorpii, Roem. & Schult. Syst. iv. 285 = hirsutus.
siculus, *Linn. Sp. Pl.* 156.—Reg. Mediterr.
sidaefolius, H. B. & K. Nov. Gen. et Sp. iii. 99 = Ipomoea tubata.
simplex, Buch. Ind. DC. Prod. iii. 118 = plantagineus.
simplex, Pers. Syn. i. 178 = Ipomoea angustifolia.
simplex, Spreng. Syst. i. 607 = Ipomoea simplex.
sinensis, Desr. in Lam. Encyc. iii. 557 = Ipomoea sinensis.
sinuatus, Nees, ex G. Don, Gen. Syst. iv. 276 = Ipomoea Deppeana.
sinuatus, Petagna, ex Steud. Nom. ed. II. i. 411 = Ipomoea carnosa.
sinuatus, *Scheele, in Linnaea*, xxi. (1848) 748.—Texas.
Sloanei, Spreng. Syst. i. 593 = Ipomoea triloba.

CONVOLVULUS:—
smilacifolius, Salisb. Prod. 124 = Ipomoea muricata.
sogdianus, Bunge, Reliq. Lehm. Bot. 219 = pilosellaefolius.
solanifolius, Lowe, in Trans. Camb. Phil. Soc. iv. (1831) 22 = Massoni.
solanifolius, Spreng. Syst. i. 596 = Ipomoea solanifolia.
Soldanella, Despr. ex Choisy, in DC. Prod. ix. 345 (= *Pharbitis Preauxii*).—Ins. Canar.
Soldanella, Linn. Sp. Pl. 159 = Calystegia Soldanella.
somalensis, *Franch. in Révoil, Çomali*, 43.—Afr. trop.
somalensis, *Vatke, in Linnaea*, xliii. (1880–82) 517.—Afr. trop.
speciosus, Linn. f. Suppl. 137 = Argyreia speciosa.
speciosus, Walt. Fl. Carol. 93 = Ipomoea sagittata.
spectabilis, Hort. Brit. ex Steud. Nom. ed. II. i. 411 = Rivea tiliaefolia.
sphaerocephalus, Roxb. Hort. Beng. 14; Fl. Ind. i. 472 = eriocarpa.
sphaerostigma, Cav. Ic. v. 54. t. 481 = Jacquemontia hirsuta.
spicaefolius, Desr. in Lam. Encyc. iii. 549 = lineatus.
spicaefolius, Buch.-Ham. ex Wall. Cat. p. 83. n. 1316 = pluricaulis.
spinosus, *Burm. f. Fl. Ind.* 47. t. 19. f. 4.—Persia.
spinosus, Desr. in Lam. Encyc. iii. 548 = fruticosus.
spinosus, Forsk. Fl. Aegypt. Arab. p. cvi = Hystrix.
spiralis, Willd. ex Roem. & Schult. Syst. iv. 302 = nodiflorus.
spithamaeus, *Linn. Sp. Pl.* 158.—Am. bor.
splendens, Hornem. Hort. Hafn. Suppl. 123 = Argyreia splendens.
Sprengelii, *Choisy, in DC. Prod.* ix. 416.—Hab.?
stachydifolius, *Choisy, l. c.* 408.—Syria; Mesopotam.
stans, H. B. & K. Nov. Gen. et Sp. iii. 96 = Ipomoea stans.
stans, Michx. Fl. Bor. Am. i. 136 = spithamaeus.
stellaris, Salisb. Prod. 124 = Hermanniae.
stellatus, *Brot. ex Heynh. Nom.* ii. 158, nomen.—Hab.?
stenophyllus, *Boiss. Fl. Orient.* iv. 106.—Syria.
stipulaceus, Roxb. Hort. Beng. 14; Fl. Ind. i. 484 = Ipomoea sepiaria.
stipulatus, Desr. in Lam. Encyc. iii. 546 = Ipomoea dissecta.
Stocksii, *Boiss. Fl. Orient.* iv. 110.—Beluchist.
stoloniferus, Cyrilli, Pl. Rar. Neap. i. 14. t. 5 = Ipomoea carnosa.
striatus, Vahl, Symb. Bot. iii. 28 = Ipomoea sepiaria.
strictus, Lehm. Ind. Sem. Hort. Hamb. (1823) 17 = undulatus.
strictus, Willd. ex Steud. Nom. ed. II. i. 412 = Ipomoea murucoides.
strigosus, Russell, ex Wall. Cat. n. 2259 = Argyreia Leschenaultii.
strigosus, Spreng. Syst. i. 600 = Lettsomia setosa.
strigosus, Wall. Cat. n. 1365 = Lettsomia strigosa.
strigulosus, Boiss. Diagn. Ser. I. xi. 83 = assyricus.
suaveolens, Mart. & Gal. in Bull. Acad. Brux. xii. II. (1845) 261 = Ipomoea suaveolens.
subacaulis, Buch.-Ham. ex Wall. Cat. sub n. 2256 = Ipomoea barlerioides.
subangularis, Buch.-Ham. ex Steud. Nom. ed. II. i. 412 = Ipomoea obscura.
suberosus, Willd. Enum. Hort. Berol. Suppl. 10 = Massoni.
subglobosus, *G. Don, Gen. Syst.* iv. 293.—Cochinch.
subhirsutus, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 338, 339.—Turkestan.
sublobatus, Linn. f. Suppl. 135 = Hewittia bicolor.
subpinnatifidus, De Vriese, in Lehm. Pl. Preiss. i. 347 = erubescens.
subpubescens, Meyen, ex Walp. in Nov. Act. Nat. Cur. xix. Suppl. i. (1843) 367 = incanus.
subsericeus, *Schrenk, Enum. Pl. Nov.* i. 19.—Sibir. altaic.
subsessilis, Steud. Nom. ed. II. i. 412 = Jacquemontia subsessilis.
subspathulatus, *Vatke, in Linnaea*, xliii. (1880–82) 520.—Afr. trop.
subulatifolius, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 10.—Malaya.
subviscidus, *Steud. Nom.* ed. II. i. 412.—Chili.
subvolubilis, Ledeb. Fl. Alt. i. 222, in nota = Calystegia subvolubilis.
suffruticosus, [Dryand. in] Ait. Hort. Kew. ed. II. i. 331 = Massoni.

CONVOLVULUS:—
suffruticosus, *Desf. Fl. Atlant.* i. 175. t. 48.—Afr. bor.
suffultus, H. B. & K. Nov. Gen. et Sp. iii. 102. t. 211 = Ipomoea suffulta.
sulphureus, La Llave, in Nov. Veg. Desc. fasc. i. 17 = Ipomoea corralinensis.
superbus, H. B. & K. Nov. Gen. et Sp. iii. 103 = Ipomoea superba.
supinus, *Coss. & Kral. in Bull. Soc. Bot. Fr.* iv. (1857) 397, 487, nomen.—Algeria.
sylvaticus, Spreng. Syst. i. 602 = Calystegia sylvatica.
sylvestris, Waldst. & Kit. Pl. Rar. Hung. iii. 290 = Calystegia sylvatica.
tamnifolius, G. F. W. Mey. Prim. Fl. Esseq. 95 = Jacquemontia tamnifolia.
tamnifolius, Willd. ex Roem. & Schult. Syst. iv. 301 = Ipomoea eriocarpa.
Tarita, Buch.-Ham. ex Wall. Cat. n. 2253 = Rivea ornata.
tenellus, Desr. in Lam. Encyc. iii. 559 = Breweria humistrata.
tenellus, Stocks, in Hook. Kew Journ. iv. (1852) 172 = Stocksii.
tenuiflorus, Steud. Nom. ed. II. i. 412 = Jacquemontia parviflora.
tenuifolius, Buch.-Ham. ex Wall. Cat. p. 84, n. 1354 = Ipomoea pentaphylla.
tenuifolius, Heyne, ex Wall. l. c. n. 1353 = Ipomoea palmata.
tenuifolius, *Mart. & Gal. in Bull. Acad. Brux.* xii. II. (1845) 260.—Mexic.
tenuifolius, Vahl, Symb. Bot. iii. 33 = Ipomoea quinata.
tenuirostris, Steud. ex Choisy, in DC. Prod. ix. 380 = Ipomoea tenuirostris.
tenuissimus, *Sibth. & Sm. Fl. Graec. Prod.* i. 134; *Fl. Graec.* t. 195.—Europ. austr.; Afr. bor.
terminalis, Salisb. Prod. 125 = cantabrica.
ternatus, Spreng. Syst. i. 590 = Ipomoea ternata.
ternifolius, Spreng. l. c. 613 = Ipomoea ternifolia.
terrestris, Linn. Sp. Pl. ed. II. 224 = cantabrica.
thalassicus, Steud. Nom. ed. II. i. 412 = Ipomoea littoralis.
Thonningii, Schum. & Thonn. Beskr. Guin. Pl. 98 = Ipomoea dissecta.
Thunbergii, *Roem. & Schult. Syst.* iv. 268.—Afr. austr.
tiliaceus, Willd. Enum. Hort. Berol. 203 = Ipomoea fastigiata.
tiliaefolius, Desr. in Lam. Encyc. iii. 544 = Argyreia tiliaefolia.
timorensis, D. Dietr. Syn. Pl. i. 667 = Hewittia bicolor.
tomentosus, Choisy, in DC. Prod. ix. 413 = hirsutus.
tomentosus, *Linn. Sp. Pl.* 156.—Jamaica.
tomentosus, Vell. Fl. Flum. 74; ii. t. 65 = Ipomoea acuminata.
Tournefortii, Sieber, ex Spreng. Syst. i. 611 = oleaefolius.
tragacanthoides, *Turcz. in Bull. Soc. Nat. Mosc.* v. (1832) 201.—Mongolia.
translucens, *Seem. in Journ. Bot.* vii. (1869) 165.—Mongolia.
trianthus, Spreng. Syst. i. 598 = Ipomoea sepiaria?
trichocalyx, Schum. & Thonn. Beskr. Guin. Pl. 91 = Ipomoea ochracea.
trichocalyx, Zoll. & Mor. in Nat. en Geneesk. Arch. Nêerl. Indie, ii. (1845) 6 = Ipomoea pileata.
trichocarpus, Spreng. Syst. i. 593 = Ipomoea commutata.
trichophorus, Wall. Cat. n. 1423 = Ipomoea cissoides.
trichosanthes, Michx. Fl. Bor. Am. i. 137 = Breweria humistrata.
trichotosus, D. Dietr. Syn. Pl. i. 670 = Argyreia capitata.
tricolor, *Linn. Sp. Pl.* 158.—Europ. austr.
tricornutus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 10.—Malaya.
tridentatus, Linn. Sp. Pl. 157 = Ipomoea tridentata.
trifidus, H. B. & K. Nov. Gen. et Sp. iii. 107 = Ipomoea trifida.
triflorus, Pav. ex Choisy, in DC. Prod. ix. 340 = Ipomoea quinquefolia.
triflorus, *Phil. in Linnaea*, xxxiii. (1864–65) 183.—Chili.
triflorus, Vahl, Symb. Bot. iii. 30 = Ipomoea sepiaria.
trigonus, Buch.-Ham. ex Wall. Cat. p. 84, n. 1383 = Ipomoea obscura.

CONVOLVULUS:—
trilobus, Desr. in Lam. Encyc. iii. 564 = Ipomoea triloba.
trilobus, *Thunb. Prod. Pl. Cap.* 35.—Afr. austr.
trinervis, *Thunb. Fl. Jap.* 85.—Japon.
triqueter, Rehm. ex Boiss. Fl. Orient. iv. 94 = Calverti.
triqueter, Vahl, Symb. Bot. iii. 30 = Ipomoea Turpethum.
triquetrus, Vell. Fl. Flum. 71; ii. t. 53 = Ipomoea batatoides.
truncatus, *Vell. l. c.* 74; ii. t. 74.—Bras.
Tuba, Schlecht. in Linnaea, vi. (1831) 735 = Ipomoea grandiflora.
tuberculatus, Desr. in Lam. Encyc. iii. 545 = Ipomoea palmata.
tuberifer, Steud. Nom. ed. II. i. 412 = Ipomoea Batatas.
tuberosus, Spreng. Syst. i. 591 = Ipomoea tuberosa.
tuberosus, Vell. Fl. Flum. 72; ii. t. 57 = Ipomoea Batatas.
Tuckeyanus, *Steud. Nom.* ed. II. i. 407. lin. 1.—Afr. trop.
tuguriorum, Forst. f. Prod. 14 = Calystegia tuguriorum.
turbinifolius, *Hort. Bonon. ex Pasq. Cat. Orto Bot. Nap.* 81, nomen.—Hab.?
Turpethum, Linn. Sp. Pl. 155 = Ipomoea Turpethum.
ulicinus, *Boiss. Diagn.* Ser. I. vii. 26.—Arabia.
umbellatus, Linn. Sp. Pl. 155 = Ipomoea umbellata.
umbellatus, Wall. Cat. n. 2329 = Ipomoea cymosa.
undulatus, *Cav. Ic.* iii. 39. t. 277. f. 1.—Reg. Mediterr.
uniflorus, Burm. f. Fl. Ind. 47 = Ipomoea uniflora.
unilateralis, Roem. & Schult. Syst. iv. 284 = Jacquemontia secunda.
urbicus, Salzm. ex Choisy, in DC. Prod. ix. 350 = Ipomoea asarifolia.
urceolatus, Spreng. Syst. i. 601 = Ipomoea urceolata.
Vahlii, *Roem. & Schult. Syst.* iv. 280.—Bras.
Vahlii, Spreng. Syst. i. 611 = Ipomoea chryseides.
valentinus, *Cav. Ic.* ii. 65. t. 180. f. 2.—Hispan.
valentinus, Hort. ex Steud. Nom. ed. II. i. 412 = Ipomoea sagittata.
valenzuelanus, A. Rich. Fl. Cub. Fanerog. ii. 134 = havanensis.
valerianoides, Blanco, Fl. Filip. ed. I. 90 = Evolvulus alsinoides.
variabilis, Cham. & Schlecht. in Linnaea, v. (1830) 116 = Ipomoea variabilis.
variabilis, Weinm. ex Steud. Nom. ed. II. i. 412 = dissectus.
variifolius, Steud. l. c. = Ipomoea hederacea.
varius, Vell. Fl. Flum. 73; ii. t. 61 = Ipomoea Batatas.
velutinus, Spreng. Syst. i. 601 = Ipomoea velutina.
venenatus, *West, Bidr. Ste-Croix*, 271.—Ins. S. Crux.
venosus, Desr. in Lam. Encyc. iii. 566 = Ipomoea venosa.
venosus, Herb. Madr. ex Wall. Cat. p. 84, n. 1353 = Ipomoea palmata.
ventricosus, Bert. ex Colla, Hort. Ripul. 37 = Ipomoea ventricosa.
ventricosus, Silva Manso, Enum. Subst. Braz. 20 = Ipomoea silvana.
venulosus, Ehrenb. ex Boiss. Fl. Orient. iv. 107 = stachydifolius.
venustus, Spreng. Syst. i. 600 = Ipomoea tricolor.
verrucipes, Tenore, Ind. Sem. Hort. Neap. (1840) = Ipomoea verrucipes.
verrucosus, D. Dietr. Syn. Pl. i. 670 = Ipomoea verrucosa.
versicolor, Salisb. Prod. 125 = tricolor.
verticillatus, Linn. Sp. Pl. ed. II. 220 = Jacquemontia hirsuta.
vestitus, Wall. Cat. n. 1411 = Lettsomia bella?
villosulus, D. Dietr. Syn. Pl. i. 667 = Ipomoea villosula.
villosus, *A. Gray, in Proc. Am. Acad.* xi. (1876) 90.—Calif.
villosus, Pers. Syn. i. 177 = Jacquemontia tamnifolia.
violaceus, Spreng. Syst. i. 599 = Ipomoea violacea.
violaceus, Vahl, Symb. Bot. iii. 29 = Jacquemontia violacea.
virgatus, *Boiss. Diagn.* Ser. I. vii. 24.—Persia; Arab.
viscidus, Roxb. Hort. Beng. 14, nomen = Ipomoea tridentata.
vitifolius, Burm. f. Fl. Ind. 45 = Ipomoea vitifolia.
vittatus, Zipp. ex Span. in Linnaea, xv. (1841) 340 = Ipomoea palmata.

CONVOLVULUS :—
volubilis, Brouss. ex Link, in Buch, Beschr. Canar. Ins. 145 = Massoni.
Wallichianus, Spreng. Syst. iv. Cur. Post. 61 = Calystegia hederacea.
waltherioides, *Boiss. & Haussk. ex Boiss. Pl. Or. Nov.* dec. i. 6.—Persia.
Wheleri, Vahl, Symb. Bot. ii. 36 = Ipomoea sagittata.
Wightii, Wall. Pl. As. Rar. ii. 55. t. 171 = Ipomoea Wightii.
Willdenowii, Steud. Nom. ed. II. i. 412 = Ipomoea heterophylla.
zeylanicus, Hook. ex Steud. l. c. = Argyreia populifolia.
zeylanicus, Moon, ex Wall. Cat. n. 1379 = Ipomoea sidaefolia.
Zollingeri, *Choisy, in Zoll. Syst. Verz. Ind. Archip.* 130, 132.—Ins. Saraja.

CONYSA, Burm. f. Fl. Ind. 180 (1768) = **Conyza**, Linn. (Compos.).

CONYSTYLUS, Pritz. Ic. Index, ii. 90 (1866) = **Gonystylus**, Teijsm. & Binn. (Thymel.).

CONYZA, Linn. Gen. ed. I. 251 (1737); Less. Syn. Comp. 203 (1832). *COMPOSITAE*, Benth. & Hook. f. ii. 283.
DIMORPHANTHES, Cass. in Bull. Soc. Philom. (1818) 30; et in Dict. Sc. Nat. xiii. 254 (1819).
EDEMIAS, Rafin. Fl. Tellur. ii. 49 (1836).
ESCHENBACHIA, Moench, Meth. 573 (1794).
FIMBRILLARIA, Cass. in Bull. Soc. Philom. (1818) 31; et in Dict. Sc. Nat. xvii. 54 (1820).
FRAGMOSA, Rafin. Fl. Tellur. ii. 50 (1836).
LAENNECIA, Cass. in Dict. Sc. Nat. xxv. 91 (1822).
MARSEA, Adans. Fam. ii. 122 (1763).
MUSTERON, Rafin. Fl. Tellur. ii. 50 (1836).
PINARDIA, Neck. ex Steud. Nom. ed. II. i. 412 (1840).
TUBIFERA, DC. ex Steud. l. c. (1840).
WEBBIA, Sch. Bip. in Walp. Rep. ii. 970 (1843).
abbreviata, Wall. Cat. n. 3048 = Vernonia cinerea.
absinthifolia, DC. in Wight, Contrib. 16 = stricta.
abyssinica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 390.—Geront. trop.
acrocephala, *Miq. Fl. Ind. Bat.* ii. 43.—Malaya.
acuminata, Wall. Cat. n. 3034 = Vernonia arborea.
acutata, Miq. Fl. Ind. Bat. ii. 48 = Blumea laciniata.
adenocarpa, *Dalz. & Gibs. Bomb. Fl.* 125.—Ind. or.
adnata, *H. B. & K. Nov. Gen. et Sp.* iv. 74.—Mexic.
aegyptiaca, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 183.—Aegypt.
aegyptiaca, Hook. f. & Thoms. ex Fl. Brit. Ind. iii. 254 = Erigeron asteroides.
ageratoides, *DC. Prod.* v. 385.—Madag.
alata, Baumg. Enum. Transsyl. iii. 102 = Inula bifrons.
alata, Roxb. Hort. Beng. 61; Fl. Ind. iii. 430 = Laggera alata.
albicans, Wall. Cat. n. 3072 = Vernonia albicans.
albida, Willd. ex Spreng. Syst. iii. 514 = Erigeron bonariensis.
alopecuroides, Lam. Encyc. ii. 93; Sw. Fl. Ind. Occ. iii. 1344 = Pterocaulon virgatum.
altaica, DC. Prod. v. 380 = Aster angustus.
ambigua, DC. Fl. Fr. Suppl. 468 = Erigeron unifolius.
ambigua, Hook. & Arn. in Bot. Beech. Voy. 57 = chilensis.
amboinica, *Crantz, Inst.* i. 272.—Amboina.
amoena, Link, Enum. Hort. Berol. ii. 322 = Blumea lacera.
amorphoglossa, Boiss. Diagn. Ser. I. vi. 81 = Erigeron amorphoglossus.
amplectens, Wall. Cat. n. 3096 = Blumea amplectens.
amplexicaulis, Lam. Encyc. ii. 84 = Blumea amplectens.
amplexicaulis, Michx. Fl. Bor. Am. ii. 126 = Pluchea bifrons.
amplexicaulis, Wall. Cat. n. 3104 = Blumea bifoliata.
amygdalina, *Lam. Encyc.* ii. 90.—Ins. Mascar.
amygdalina, Spreng. ex DC. Prod. vi. 446 = Blumea densiflora.
anagallidifolia, Wall. Cat. n. 3098 = Blumea bifoliata.
anchusaefolia, *Poir. Encyc. Suppl.* ii. 339.—Ins. Mascar.
andina, *Remy, in C. Gay, Fl. Chil.* iv. 72.—Chili.

CONYZA :—
andryaloides, DC. Prod. v. 377 = Erigeron andryaloides.
angustifolia, Hort. Par. ex DC. l. c. 420 = Baccharis Pingraea.
angustifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 109 = Pluchea bifrons?
angustifolia, *Roxb. Hort. Beng.* 61; *Fl. Ind.* iii. 429.—Reg. Himal.
angustifolia, Steud. Nom. ed. I. 221 = Monarrhenus salicifolius.
angustifolia, Thw. ex Hook. f. Fl. Brit. Ind. iii. 254 = Erigeron linifolius.
anthelmintica, Linn. Sp. Pl. ed. II. 1207 = Vernonia anthelmintica.
appendiculata, Blume, Bijdr. 895 = Blumea balsamifera.
appendiculata, Lam. Encyc. ii. 88 = Senecio appendiculatus.
apurensis, H. B. & K. Nov. Gen. et Sp. iv. 73 = Erigeron apurensis.
arabica, *Willd. Sp. Pl.* iii. 1949.—Arab.
arabidaefolia, *Remy, in C. Gay, Fl. Chil.* iv. 76.—Chili.
arborea, Wall. Cat. n. 3074 = Vernonia arborea.
arborescens, DC. Prod. v. 344, in syn. cum cit. falsa = Commidendron spurium?
arborescens, Linn. Syst. ed. X. 1213 = Vernonia arborescens.
arbutifolia, Lam. Encyc. ii. 92 = Baccharis arbutifolia.
argentea, *Lam. l. c.* 87.—Ins. Mascar.
argentea, Perr. & Lepyr. ex DC. Prod. v. 449 = Laggera aurita.
argentea, Wall. Cat. n. 3014 = Inula Cappa.
arguta, *Less. in Linnaea*, vi. (1831) 138.—Bras.
argutidens, *Miq. Fl. Ind. Bat.* ii. 58.—Malaya.
aromatica, Wall. Cat. n. 3054 = Blumea aromatica.
artemisiaefolia, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 262.—Peruv.
artemisioides, Sch. Bip. ex Baker, in Mart. Fl. Bras. vi. III. 63 = Baccharis rufescens.
articulata, Lam. Encyc. ii. 94 = Baccharis articulata.
arvensis, *Phil. in Linnaea*, xxxiii. (1864-65) 143.—Chili.
aspera, *Bory, Voy.* iii. 174 = melastomoides.
aspera, Wall. ex Steud. Nom. ed. II. i. 413 = Vernonia Roxburghii.
asperifolia, *Benth. & Hook. f. Gen.* ii. 284.—Guatemala.
asperula, *Phil. in Linnaea*, xxviii. (1856) 736.—Chili.
asteroides, Desf. Tabl. ed. II. (1815) 114 = Pluchea subdecurrens.
asteroides, Host, ex Beck, in Denkschr. Acad. Wien, Math. Nat. xliv. (1882) 332 = Inula bifrons.
asteroides, Linn. Sp. Pl. 861 = Sericocarpus conyzoides.
asteroides, Wall. Cat. n. 3052 = aegyptiaca.
asteroides, Wall. ex Steud. Nom. ed. II. i. 413 = Chrysopsis gossypina.
attenuata, *DC. Prod.* v. 384.—Madag.
attenuata, Wall. Cat. n. 3020 = Vernonia attenuata.
Aucheri, DC. Prod. vii. 281 = Erigeron Aucheri.
aurea, Perr. ex DC. l. c. v. 68 = Vernonia senegalensis.
auriculata, Moench, Meth. Suppl. 235 = Phagnalon rupestre.
aurita, Heyne, ex DC. Prod. v. 441 = Blumea lacera.
aurita, *Linn. f. Suppl.* 367.—Nubia.
axillaris, *Hornem. ex Willd. Enum. Hort. Berol.* 871.—Afr. occ.
axillaris, Lam. Encyc. ii. 84 = Blumea lacera.
Baccharis, *Mill. Gard. Dict.* ed. VIII. n. 16.—Honduras.
baccharoides, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 390 = Pluchea Dioscoridis.
balsamica, Wall. ex DC. Prod. v. 319 = Psiadia balsamica.
balsamifera, Linn. Sp. Pl. ed. II. 1208 = Blumea balsamifera.
barbata, Wall. Cat. n. 3099 = Blumea barbata.
bellidifolia, *Baker, in Journ. Bot.* xx. (1882) 169.—Madag.
bellidifolia, Wall. Cat. n. 3048 = Vernonia cinerea.
Berteroana, *Phil. in Linnaea*, xxviii. (1856) 737.—Chili.
bidentata, Steud. Nom. ed. I. 222 = calocephala.
bifoliata, Cham. & Less. in Linnaea, vi. (1831) 135 = Blumea Wightiana.
bifoliata, Linn. Sp. Pl. 862 = Blumea bifoliata.

CONYZA :—
bifoliata, Walt. Fl. Carol. 204 = Sericocarpus tortifolius.
bifoliolata, DC. Prod. v. 435 = Blumea Wightiana.
bifrons, Linn. Sp. Pl. 861; Gouan, Hort. Reg. 436 = Inula bifrons.
bifrons, Pursh, Fl. Am. Sept. ii. 524 = Pluchea bifrons.
Bilbaoana, Remy, in C. Gay, Fl. Chil. iv. 76 = Erigeron canadensis.
Blanchetii, *Baker, in Mart. Fl. Bras.* vi. III. 36.—Bras.
blanda, Wall. Cat. n. 3033 = Vernonia blanda.
Bojeri, *DC. Prod.* v. 384.—Madag.
Bovei, DC. in Ann. Sc. Nat. Sér. II. i. (1834) 261 = Erigeron Bovei.
bovina, Buch.-Ham. ex DC. Prod. v. 444 = Blumea oxyodonta.
bracteolata, Rottl. ex DC. l. c. 434 = Blumea bifoliata.
bracteolata, Wall. Cat. n. 3036 = Vernonia subsessilis.
brasiliensis, Steetz, in Peters, Reise Mossamb. Bot. 455 = Epaltes brasiliensis.
britannica, Rupr. Fl. Ingric. 569 = Inula britannica.
bryoides, Lam. Encyc. ii. 91 = Baccharis bryoides.
burmeana, *Miq. Fl. Ind. Bat.* ii. 44.—Burma.
Bustilloziana, *Remy, in C. Gay, Fl. Chil.* iv. 72.—Chili.
buxifolia, Lam. Encyc. ii. 92 = Baccharis buxifolia.
caffra, DC. Prod. v. 381 = Blumea natalensis.
calocephala, *Bory, Voy.* ii. 395.—Ins. Borbon.
callosa, *Klatt, in Decken, Reise Ost-Afr.* iii. 76.—Afr. trop.
calycina, Cav. in Anal. Cienc. Nat. iv. (1801) 87 = Phagnalon rupestre.
camphorata, Pursh, Fl. Am. Sept. ii. 523 = Pluchea camphorata.
canariensis, Willd. Sp. Pl. iii. 1937 = Inula candida.
candicans, Wall. Cat. n. 2999 = Leucomeris spectabilis.
candida, Linn. Sp. Pl. 862 = Inula candida.
Candolleana, Boiss. Fl. Orient. iii. 217 = Pluchea Wallichiana.
canescens, Bory, ex DC. Prod. v. 473 (= *Schizogyne sericea*).—Ins. Canar.
canescens, Linn. f. Suppl. 367 = Vernonia pinifolia.
canescens, *Phil. in Linnaea*, xxviii. (1856) 734.—Chili.
canescens, Thunb. Fl. Cap. 655 = Vernonia pinifolia.
capitata, Spreng. Neue Entdeck. ii. 141 = Vanillosmopsis capitata.
Cappa, Buch.-Ham. in D. Don, Prod. Fl. Nep. 176 = Inula Cappa.
caprariaefolia, *Willd. ex Steud. Nom.* ed. II. i. 413.—Hab.?
cardaminaefolia, *H. B. & K. Nov. Gen. et Sp.* iv. 71.—Ecuador.
carolinensis, Jacq. Ic. Rar. iii. t. 585 = Pluchea odorata.
cataphracta, Willd. ex DC. Prod. v. 427 = Baccharis cataphracta.
cayennensis, *Spreng. Syst.* iii. 512.—Guian.
celebica, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 122.—Ins. Celebes.
cernua, Wall. Cat. n. 3012 = Laggera alata.
chenopodiifolia, Lam. Encyc. ii. 85 = Gynura mauritiana.
chenopodioides, DC. Prod. v. 379 = Erigeron bonariensis.
chilensis, *Spreng. Nov. Prov.* 14.—Chili.
chinensis, Lam. Encyc. ii. 83 = Vernonia chinensis.
chinensis, Linn. Sp. Pl. 862 = Blumea chinensis.
chinensis, Pers. Syn. ii. 426 = Lourieiii.
Chingoyo, H. B. & K. Nov. Gen. et Sp. iv. 76. t. 328 = Pluchea Chingoyo.
chrysocoma, *Vatke, in Bremen Abh.* ix. (1885) 120.—Madag.
chrysocomoides, Desf. Fl. Atlant. ii. 269. t. 282 = Nolletia chrysocomoides.
chrysocomoides, Desf. Hist. Arb. i. 292 = Felicia angustifolia.
chrysocomoides, Tenore, ex Steud. Nom. ed. II. i. 413 = procumbens.
cichoriifolia, Miq. Fl. Ind. Bat. ii. 47 = Blumea laciniata.
ciliaris, *Spreng. Nov. Prov.* 14.—Hab.?
cinerarioides, Boj. ex DC. Prod. v. 345 = Rochonia cinerarioides.
cinerascens, Wall. Cat. n. 3062 = Vernonia cinerea.
cinerea, Linn. Sp. Pl. 862 = Vernonia cinerea.
clarenceana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 316.—Afr. trop.
clethraefolia, Willd. ex Baker, in Mart. Fl. Bras. vi. II. 100 = Vernonia scabra.

CONYZA:—
collina, *Phil. in Linnaea*, xxxiii. (1864–65) 144.—Chili.
communis, Wall. Cat. n. 3018 = Blumea hieracifolia.
concava, Spreng. ex Steud. Nom. ed. II. i. 413 = Baccharis concava.
congesta, Spreng. ex DC. Prod. v. 383 = japonica.
conica, *Poir. Encyc. Suppl.* ii. 341.—Hab.?
consanguinea, *Phil. in Linnaea*, xxxiii. (1864-65) 142.—Chili.
conspicua, Wall. Cat. n. 3066 = Microglossa albescens.
coronipifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 70.—Mexic.
Coronopus, Lam. Encyc. ii. 89 = Psiadia Coronopus.
Cortesii, *H. B. & K. Nov. Gen. et Sp.* iv. 75.—Mexic.
corymbosa, *Mill. Gard. Dict.* ed. VIII. n. 7.—Mexic.
corymbosa, Roxb. Hort. Beng. 61; Fl. Ind. iii. 426 = Pluchea indica.
costata, *Harv. in Harv. & Sond. Fl. Cap.* iii. 114.—Afr. austr.
Cotoneaster, Willd. ex Spreng. Syst. iii. 509 = Vernonia Cotoneaster.
Coulteri, *A. Gray, in Proc. Am. Acad.* vii. (1868) 355.—Am. bor. occ.
Coulteri, Rothr. in Wheeler, Rep. 155 = Erigeron subdecurrens.
crispa, Rupr. in Bull. Phys.-Math. Acad. Pétersb. xiv. (1856) 235 = Erigeron linifolius.
crispata, *Vahl, Symb. Bot.* i. 71.—Arabia.
ctenoptera, Kunth, Ind. Sem. Hort. Berol. (1845) 12 = Laggera pterodonta.
cuneata, Boj. ex DC. Prod. v. 345 = Rochonia cuneata.
cuneata, Buch.-Ham. ex DC. l. c. 384 = absinthifolia.
cuneifolia, Lam. Encyc. ii. 91 = Baccharis cuneifolia.
cuneifolia, Raeusch. Nom. ed. III. 240 = Commidendrum spurium.
cuneifolia, Wall. Cat. n. 3101 = Blumea lacera.
cupressiformis, Lam. Encyc. ii. 91; Illustr. t. 697 = Lepidophyllum cupressiforme.
Dampieri, A. Cunn. ex DC. Prod. v. 266 = Olearia axillaris.
dasycoma, Miq. Fl. Ind. Bat. ii. 56 = Blumea densiflora.
decurrens, *Linn. Sp. Pl.* ed. II. 1206.—Ind. or.
densiflora, Heyne, in Wall. Cat. n. 2997 = Blumea densiflora.
dentata, Blanco, Fl. Filip. ed. I. 629 = Inula Cappa.
dentata, *Willd. Sp. Pl.* iii. 1928.—Afr. trop.
dichotoma, Pers. Syn. ii. 428 = Jasonia dichotoma.
diffusa, Poir. Encyc. Suppl. ii. 338 = Gouani.
diffusa, Roxb. Hort. Beng. 61; Fl. Ind. iii. 429 = Blumea membranacea.
dilatata, Steud. Nom. ed. I. 222 = laurifolia.
Dioscoridis, Desf. Tabl. ed. II. 114 = Pluchea Dioscoridis.
discolor, Bory, ex DC. Prod. vi. 376 = Senecio appendiculatus.
disticha, Heyne, in Wall. Cat. n. 3024 = viscidula.
disticha, *Spreng. Syst.* iii. 509.—Am. austr.
divaricata, Spreng. l. c. 508 = Vernonia diffusa.
divergens, Wall. Cat. n. 3027 = Vernonia divergens.
diversifolia, Weinm. in Flora, iii. (1820) 611 = chilensis.
domingensis, Willd. ex DC. Prod. v. 389 = lucens.
dongolensis, *Delile, Pl. Afr. Voy. Méroé*, i. 43.—Nubia.
dumetorum, *Phil. in Linnaea*, xxxiii. (1864–65) 144.—Chili.
echinoides, A. Rich. Tent. Fl. Abyss. i. 388 = aegyptiaca.
elaeagnifolia, Wall. Cat. n. 3041 = Vernonia elaeagnifolia.
elata, *Kunth & Bouché, Ind. Sem. Hort. Berol.* (1848) 14.—Bras.
elegantula, Wall. Cat. n. 3049 = Vernonia cinerea.
elongata, Wall. l. c. n. 3078 = Blumea elongata.
ericoides, Lam. Encyc. ii. 92 = Liabum ericoides.
erigeroides, DC. Prod. v. 378 = Erigeron bonariensis.
eriophora, Wall. Cat. n. 3013 = Inula Cappa.
erosa, Blanco, Fl. Filip. ed. II. 439 = Blumea manillensis.
eucoma, *Miq. Pl. Jungh.* 500.—Java.
eupatorioides, Wall. Cat. n. 2993 = Inula eupatorioides.
excisa, Wall. l. c. n. 3011 = Blumea densiflora.
extensa, Wall. l. c. n. 3016 = Vernonia extensa.
falcata, *Buch.-Ham. ex Spreng. Syst.* ii. 513.—Reg. Himal.
falcigera, Wall. Cat. n. 2048 = Blumea densiflora.
fasciculata, Wall. l. c. n. 3019 = Laggera flava.
fasciculata, Wall. l. c. n. 3094 (ex Steud.) = Blumea glomerata.

CONYZA:—
fastigiata, *Willd. Enum. Hort. Berol. Suppl.* 57.—Afr. bor.
fistulosa, Roxb. Hort. Beng. [101]; Fl. Ind. iii. 429 = Blumea glomerata.
flexilis, *DC. Prod.* v. 385.—Madag.
floresiana, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 121.—Malaya.
floribunda, H. B. & K. Nov. Gen. et Sp. iv. 73 = Erigeron bonariensis.
foetida, Lam. Encyc. ii. 85 = Pluchea camphorata.
foetida, Willd. Sp. Pl. iii. 1952 = Nidorella foliosa.
foliolosa, Buch.-Ham. in Wall. Cat. n. 3281 = Pulicaria foliolosa.
foliolosa, Wall. ex DC. Prod. v. 451 = Pluchea indica.
foliolosa, Wight, in Wall. Cat. n. 3092 = Blumea bifoliata.
foliolosa, *Willd. Sp. Pl.* iii. 1919.—Ind. or.
fruticosa, Linn. Sp. Pl. ed. II. 1209 = Vernonia rigida.
Garnieri, *Klatt, in Linnaea*, xxxvii. (1871–73) 509.—Madag.
Gayana, *Phil. l. c.* xxviii. (1856) 737.—Chili.
geminiflora, Tenore, Cat. Hort. Neap. (1819) 75 = Phagnalon rupestre.
genistelloides, Lam. Encyc. ii. 93 = Baccharis genistelloides.
glabra, Willd. Sp. Pl. iii. 1940 = Vernonia Vahliana.
glabrata, Wall. ex Steud. Nom. ed. II. i. 414 = Eupatorium glabrum.
glomerata, Link, Enum. Hort. Berol. ii. 322 = decurrens.
glomerata, Wall. Cat. n. 3055 = Blumea glomerata.
glutinosa, Lam. Encyc. ii. 89 = Psiadia glutinosa.
gnaphalioides, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 389 = Gouani.
gnaphalodes, *H. B. & K. Nov. Gen. et Sp.* iv. 73. t. 327.—Mexic.
gnaphalodes, Sieber, Reise Kreta, i. 352; ii. 322 = Senecio gnaphalodes.
Gouani, Blanco, Fl. Filip. ed. I. 627 = Blumea manillensis.
Gouani, *Willd. Sp. Pl.* iii. 1928.—Ins. Canar.
gracilis, Heyne, in Wall. Cat. n. 3064 = Blumea membranacea.
gracilis, Hoffmgg. & Link, Fl. Port. ii. 253 = Pulicaria gracilis.
gracilis, Wall. Cat. n. 3064 = Blumea glomerata.
graeca, Nym. Syll. 16 = Phagnalon graecum.
graminifolia, Spreng. Syst. iii. 515 = Aster divaricatus.
Grahami, *DC. Prod.* v. 380.—Reg. Argent.
grandis, Wall. Cat. n. 3051 = Blumea densiflora.
graveolens, Steud. Nom. ed. I. 222 = Inula graveolens.
graveolens, Wall. ex DC. Prod. v. 292 = Erigeron asteroides.
guineensis, Willd. Sp. Pl. iii. 1930 = Laggera aurita.
gummifera, Roxb. in Beats. St. Hel. Tracts, App. 304 = Commidendrum robustum.
halimifolia, Desf. Tabl. ed. II. 114 = Baccharis halimifolia.
Hamiltoni, Steud. Nom. ed. II. i. 414 = Blumea hieracifolia.
heliotropifolia, Lam. Encyc. ii. 90 = Monarrhenus rufescens.
heterophylla, Lam. Encyc. ii. 84 = Vernonia cinerea.
heterotricha, Wall. Cat. n. 3225 = Inula eupatorioides.
Heudelotii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 317.—Afr. trop.
Heyneana, Wall. Cat. n. 3089 = Blumea lacera.
hieracifolia, Spreng. Syst. iii. 514 = Blumea hieracifolia.
hirsuta, Comm. ex DC. Prod. v. 458 = Cylindrocline Commersonii.
hirsuta, Linn Sp. Pl. i. 863 = Pluchea hirsuta.
hirsuta, Wall. ex DC. Prod. v. 292; et Cat. n. 3007 B, partim = aegyptiaca.
hirtella, *DC. Prod.* v. 385.—Madag.
hispida, *H. B. & K. Nov. Gen. et Sp.* iv. 71.—Mexic.
Hochstetterii, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 387.—Abyssin.
holosericea, Wall. Cat. n. 3103 = Blumea glomerata.
Huberti, *Boj. ex Sweet, Hort. Brit.* ed. III. 349.—Madag.
humifusa, *Miq. Fl. Ind. Bat.* ii. 41.—Java.
hymenophylla, Wall. Cat. n. 3038 = Blumea membranacea.
hypoleuca, *A. Rich. Tent. Fl. Abyss.* i. 391.—Abyssin.
iliensis, *Trautv. in Bull. Soc. Nat. Mosc.* xxxix. (1866) II. 342.—As. centr.

CONYZA:—
incana, Heyne, ex DC. Prod. v. 24 = Vernonia cinerea?
incana, Wall. Cat. n. 3046 = Vernonia cinerea.
incana, *Willd. Sp. Pl.* iii. 1937.—Arabia.
incisa, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 184.—Afr. austr.
indica, Blume, ex DC. Prod. v. 451 = Pluchea indica.
indica, Wall. Cat. n. 3042 = Vernonia indica.
intermedia, Lag. Gen. et Sp. Nov. 28 = Phagnalon saxatile.
inuloides, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 182 = Jasonia dichotoma.
ivaefolia, *Burm. f. Fl. Ind.* 180.—Ind. or.
ivaefolia, *Less. in Linnaea*, vi. (1831) 138.—Afr. austr.
japonica, *Less. Syn. Comp.* 204.—Reg. Himal.; China; Japon.
javanica, Blume, Bijdr. 897 = Blumea lacera.
Jerdoni, C. B. Clarke, Comp. Ind. 62 = Erigeron asteroides.
Junghuhniana, *Miq. Fl. Ind. Bat.* ii. 55.—Java.
Kotschyi, Boiss. Diagn. Ser. I. vi. 83 = Erigeron Aucheri.
Kotschyi, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 307 = Vicoa auriculata.
Korthalsiana, *Miq. Fl. Ind. Bat.* ii. 51.—Malaya.
Krausseana, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 616.—Turkestan.
Kraussii, *Sch. Bip. ex Walp. Rep.* ii. 972.—Afr. austr.
Kunthiana, *DC. Prod.* v. 379.—Ecuador.
lacera, Burm. f. Fl. Ind. 180 = Blumea lacera.
laciniata, Roxb. Hort. Beng. 61; Fl. Ind. iii. 427 = Blumea laciniata.
lactucaefolia, Wall. Cat. n. 3088 = Blumea Wightiana.
laeta, Hort. Calc. ex C. B. Clarke, Comp. Ind. 14 = Vernonia saligna.
lanceolaria, Roxb. Hort. Beng. 61; Fl. Ind. iii. 432 = Blumea myriocephala.
lanceolata, Wall. l. c. n. 2991 = Pluchea lanceolata.
lanceolata, Wall. Cat. n. 3059 = Vernonia divergens.
lanuginosa, Wall. l. c. n. 2922 = Inula Cappa.
Larrainiana, *Remy, in C. Gay, Fl. Chil.* iv. 71.—Chili.
laurifolia, *Lam. Encyc.* ii. 89.—Ins. Mascar.
laxa, *Spreng. Syst.* iii. 513.—Bras.
laxiflora, *Desf. Cat. Hort. Par.* ed. III. 400.—Hab.?
leucanthema, Buch.-Ham. ex Steud. Nom. ed. II. i. 414 = Blumea leucanthema.
leucodasys, *Miq. in Journ. Bot. Néerl.* i. (1860) 103.—Malaya.
leucophylla, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 387 = incana.
limonifolia, Sibth. & Sm. Fl. Graec. Prod. ii. 174 = Inula candida.
linearifolia, Lam. Encyc. ii. 92 = Psiadia linearifolia.
linariloba, DC. Prod. v. 385 = aegyptiaca.
linearis, *DC. l. c.* 378.—Chili.
linifolia, *Burm. f. Prod. Fl. Cap.* 26.—Afr. austr.
linifolia, Linn. Sp. Pl. 861 = Sericocarpus solidagineus.
linifolia, *Phil. in Linnaea*, xxviii. (1856) 734.—Chili.
linifolia, Wall. Cat. n. 3063 = Vernonia cinerea.
lithospermifolia, *Lam. Encyc.* ii. 90.—Ins. Mascar.
lobata, Juss. ex Steud. Nom. ed. II. i. 414, 521 = Verbesina diversifolia.
lobata, Linn. Sp. Pl. 862 = Neurolaena lobata.
longicaulis, Wall. Cat. n. 3073 = Eupatorium Reevesii.
longifolia, Desf. Cat. Hort. Par. ed. III. 164 = chilensis.
longifolia, Heyne, in Wall. Cat. n. 3026 = Blumea spectabilis.
longispina, Hook. f. Fl. Brit. Ind. iii. 268, sphalm. = seq.
longissima, Zoll. & Moritzi, Syst. Verz. Zoll. 68 = Blumea chinensis.
Lorentzii, *Griseb. in Goett. Abh.* xxiv. (1879) 176.—Reg. Argent.
Loureirii, *Poir. Encyc. Suppl.* ii. 339.—China.
lucens, *Poir. l. c.* 341.—Ins. S. Doming.
lucida, Spreng. Neue Entdeck. ii. 142 = Piptocarpha lucida.
lurida, *F. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 186.—Ins. Madera.
lycopodioides, Lam. Encyc. ii. 91 = Eriothrix juniperifolia.
lyrata, *H. B. & K. Nov. Gen. et Sp.* iv. 70.—Ecuador.
macrophylla, *Spreng. Syst.* iii. 513.—Reg. Argent.
macrorhiza, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 387.—Abyssin.

CONYZA:—
macrostachya, Wall. Cat. n. 3053 = Blumea hieracifolia.
madagascariensis, Lam. Encyc. ii. 85 = Psiadia madagascariensis.
magellanica, Lam. l. c. 91 = Baccharis magellanica.
manillensis, Less. in Linnaea, vi. (1831) 135 = Blumea manillensis.
marginata, Edgew. in Trans. Linn. Soc. xx. (1851) 67 = rufa.
marylandica, Michx. Fl. Bor. Am. ii. 126 = Pluchea camphorata.
mauritiana, *Boj. ex Sweet, Hort. Brit.* ed. III. 349.—Ins. Maurit.
mauritiana, Mérat, ex Steud. Nom. ed. II. i. 414 = Nidorella compressa.
mauritiana, Willd. ex Steud. l. c. = Nolletia chrysocomoides.
maxima, *Zoll. & Mor. in Nat. en Geneesk. Arch. Neêrl. Ind.* ii. (1845) 236.—Java.
megapotamica, Spreng. Syst. iii. 516 = Stenachaenium megapotamicum.
melastomoides, *Lam. Encyc.* ii. 90.—Ins. Mascar.
membranacea, Wall. ex DC. Prod. v. 441; Cat. n. 3019 = Blumea membranacea.
membranifolia, *Klatt, in Leopoldina*, xx. (1884) 94.—Bras.
micrantha, Spreng. ex DC. Prod. v. 384 = stricta.
microcephala, *Hemsl. Biol. Centr. Am. Bot.* 126.—Mexic.
modatensis, Sch. Bip. ex Schweinf. Beitr. Fl. Aethiop. 144 = Pluchea Dioscoridis.
modesta, *Phil. in Linnaea*, xxxiii. (1864-65) 142.—Chili.
modesta, *Kunth, Ind. Sem. Hort. Berol.* (1845) 11.—Hab. ?
mollis, Willd. Sp. Pl. iii. 1924 = Vernonia cinerea.
moluccana, *Gaudich. Bot. Voy. Bonite*, 465.—Ins. Moluce.
monocephala, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 122.—Malaya.
montana, Heyne, ex DC. Prod. v. 384 = stricta.
montevidensis, *Spreng. Syst.* iii. 510.—Reg. Argent.
multicaulis, DC. Prod. v. 383 = japonica.
multiflora, Wall. Cat. n. 3032 = Vernonia divergens.
muralis, Buch.-Ham. in Wall. Cat. n. 3077 = Blumea membranacea.
Mursa, Buch.-Ham. l. c. n. 3095 = Blumea lacera.
myosotifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 69.—N. Granat.
myriocephala, Remy, in C. Gay, Fl. Chil. iv. 76 = Erigeron canadensis.
myrsinites, Lam. Encyc. ii. 92 = Baccharis myrsinites.
nana, *Sch. Bip. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 315.—Abyss.
napifolia, Wall. Cat. n. 3075 = Blumea lacera.
natalensis, *Sch. Bip. ex Walp. Rep.* ii. 971.—Afr. austr.
natans, Buch.-Ham. ex Hook. f. Fl. Brit. Ind. iii. 263 = Blumea hieracifolia.
Naudini, *Bonnet, in Bull. Soc. Bot. Fr.* xxv. (1878) 208.—Hab. ?
necessaria, Buch.-Ham. ex DC. Prod. v. 439 = Blumea hieracifolia.
nereifolia, L'Hérit. ex DC. l. c. 430 = Brachylaena nereifolia.
Neuri, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 317.—Afr. trop.
nitida, Miq. Fl. Ind. Bat. ii. 54 = Blumea Wallichii.
nitida, *Sch. Bip. in Zoll. Syst. Verz. Ind. Archip.* 121.—Malaya.
nivalis, Boiss. Diagn. Ser. I. vi. 82 = Erigeron chionophilus.
notobellidiastrum, *Griseb. in Goett. Abh.* xxiv. (1879) 177.—Reg. Argent.
nutans, Blume, Bijdr. 896 = Laggera alata.
nutans, Buch.-Ham. ex DC. Prod. v. 436 = Blumea cernua.
nutans, Wall. Cat. n. 3080 = Blumea hieracifolia.
obliqua, Buch.-Ham. ex DC. Prod. v. 439 = Blumea flava.
obliqua, Willd. Sp. Pl. iii. 1930 = Blumea amplectens.
oblonga, Wall. ex DC. Prod. v. 469 = Inula Cappa.
oblongifolia, *Spreng. Syst.* iii. 512.—Bras.
obovata, Wall. Cat. n. 3022 = Blumea obovata.
obscura, *DC. Prod.* v. 387.—Am. austr.
obtusa, *H. B. & K. Nov. Gen. et Sp.* iv. 71.—Mexic.

CONYZA:—
odontophylla, Boiss. Fl. Orient. iii. 217 = Pluchea arguta.
odontoptera, *Webb, in Hook. Niger Fl.* 135.—Ins. Cap. Verd.
odora, *Forsk. Fl. Aegypt. Arab.* 148.—Aegypt
odorata, Linn. Syst. ed. X. 1213 = Pluchea odorata.
odorata, Willd. ex Steud. Nom. ed. II. i. 414 = Vernonia chinensis.
oleaefolia, Lam. Encyc. ii. 86 = Linosyris villosa.
oligandra, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 179.—Java.
oligocephala, Miq. Fl. Ind. Bat. ii. 41 = Blumea bifoliata.
orientalis, *Willd. Sp. Pl.* iii. 1931.—Armen.
ovata, Wall. Cat. n. 3097 = Blumea cinerea.
oxyodonta, Wall. ex DC. Prod. v. 444 = Blumea oxyodonta.
pallida, *Zipp. ex Span. in Linnaea*, xv. (1841) 322 nomen.—Malaya.
panamensis, *Willd. Sp. Pl.* iii. 1942.—Panama.
paniculata, Desf. Tabl. ed. II. 114 = fastigiata.
paniculata, Wall. Cat. n. 3090 = Blumea Wightiana.
paniculata, Willd. Sp. Pl. iii. 1920 = Blumea membranacea.
pannosa, *Webb, in Hook. Niger Fl.* 135.—Afr. trop.
parvifolia, Wall. Cat. n. 3004 = Blumea Wightiana.
patula, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 184 = Vernonia chinensis.
pauciflora, Willd. Sp. Pl. iii. 1927 = Vernonia pauciflora.
paucifolia, Wall. Cat. n. 3037 = Blumea membranacea.
pectinata, *Sch. Bip. ex Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 315.—Afr. trop.
pedunculata, *Mill. Gard. Dict.* ed. VIII. n. 15.—Hab. ?
persica, Boiss. Diagn. Ser. I. vi. 81 = Erigeron persicus.
persicaefolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 312.—Afr. trop.
petiolaris, Wall. Cat. n. 3282 = japonica.
pinifolia, Bory, Voy. ii. 395 = Faujasia pinifolia.
pinifolia, Lam. Encyc. ii. 86 = Vernonia pinifolia.
pinnatifida, Buch.-Ham. ex Roxb. Hort. Beng. 61; Fl. Ind. iii. 430 = stricta.
pinnatifida, *Less. Syn. Comp.* 204.—Afr. austr.
pinnatilobata, *DC. Prod.* v. 387.—Afr. austr.
platensis, Spreng. Syst. iii. 509 = Vernonia platensis.
platylepis, Spreng. ex DC. Prod. v. 31 = Vernonia Roxburghii.
podocephala, *DC. l. c.* 387.—Afr. austr.
polycephala, Edgew. in Trans. Linn. Soc. xx. (1851) 66 = viscidula.
polygonata, Wall. Cat. n. 3010 = Inula polygonata.
Ponae, Sieber, ex Tausch, in Flora, xii. (1829) i. Erg. 37 = Cylindrocline Commersonii.
populifolia, Lam. Encyc. ii. 87 = Vernonia populifolia.
portoricensis, Bert. ex DC. Prod. v. 52 = Vernonia remotiflora ?
pratensis, *Phil. in Linnaea*, xxviii. (1856) 736.—Chili.
procera, Desf. Cat. Hort. Par. ed. III. 164 = chilensis.
procera, Wall. Cat. n. 3050 = Blumea procera.
procumbens, *Balb. Cat. Hort. Taur.* 28.—Ins. Creta.
prolifera, Lam. Encyc. ii. 84 = Vernonia cinerea.
prostrata, Tenore, Cat. Hort. Neap. (1819) 75 = procumbens.
proteiformis, Perr. ex DC. Prod. v. 376 = Pluchea lanceolata.
pterocaulon, *C. Bolle, in Bonplandia*, vii. (1859) 295.—Ins. Canar.
pubescens, DC. Prod. v. 381 = Brachyactis robusta.
pubigera, *Linn. Mant.* i. 113.—Amboina.
pulchella, H. B. & K. Nov. Gen. et Sp. iv. 73 = sophiaefolia.
pulchra, Lam. ex Steud. Nom. ed. II. i. 415 = amygdalina.
pumila, *Lam. ex Pers. Syn.* ii. 427.—Hab. ?
pumila, Sibth. & Sm. Fl. Graec. Prod. ii. 173 = Phagnalon pumilum.
punctata, Willd. Sp. Pl. iii. 1937 = Aplopappus Chamissonis.
punctulata, Wall. Cat. n. 2995 = Vernonia chinensis.
punduana, Wall. l. c. n. 3035 = Vernonia volkameriaefolia.
pungens, Lam. Encyc. ii. 86 = Iphiona juniperifolia.
purpurascens, Sw. Prod. Veg. Ind. Occ. 112 = Pluchea purpurascens.
purpurea, Buch.-Ham. in Wall. Cat. n. 3076 = Blumea glomerata.

CONYZA:—
pusilla, *Houtt. Handleid.* x. 618. t. 69.—Afr. austr.
pusilla, H. B. & K. Nov. Gen. et Sp. iv. 69 = Kunthiana.
pycnostachya, Michx. Fl. Bor. Am. ii. 126 = Pterocaulon pycnostachyum.
pygmaea, Sieber, Reise Kreta, ii. 322 = Phagnalon pumilum.
pyrifolia, Lam. Encyc. ii. 89 = Microglossa volubilis.
pyrrhopappa, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 389.—Abyss.
racemosa, Spreng. Syst. iii. 508 = Baccharis aviceniaefolia.
racemosa, Wall. Cat. n. 3002 = Blumea glomerata.
ramiflora, Spreng. Syst. iii. 509 = Piptocarpha ramiflora.
redolens, Willd. Sp. Pl. iii. 1951 (= *Monenteles redolens*).—N. Caled.
repanda, Roxb. Hort. Beng. 61; Fl. Ind. iii. 431 = Blumea procera.
retusa, *Lam. Encyc.* ii. 90.—Ins. Mascar.
rigida, Sw. Prod. Veg. Ind. Occ. 113 = Vernonia rigida.
riparia, Blume, Bijdr. 899 = Blumea chinensis.
riparia, H. B. & K. Nov. Gen. et Sp. iv. 76 = Tessaria integrifolia.
rivularis, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 124.—Bras.
robusta, Roxb. in Beats. St. Hel. Tracts, App. 304 = Commidendrum robustum.
Roylei, DC. Prod. v. 381 = Brachyactis menthodora.
rubra, Buch.-Ham. ex DC. l. c. 376 = Pluchea lanceolata.
rudis, Willd. ex Steud. Nom. ed. II. i. 415 = lyrata.
rufa, *Wall. Cat.* n. 3079.—Ind. or.
rufescens, Hoffmgg. & Link, Fl. Port. ii. 253 = Pulicaria rufescens.
rufidula, *Spreng. Neue Entdeck.* ii. 141.—Bras.
rugosa, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 184 = Commidendrum rugosum.
rugosa, Vahl, Symb. Bot. i. 71 = Pterocaulon virgatum.
runcinata, Wall. Cat. n. 3087 = Blumea laciniata.
rupestris, Linn. Mant. 113 = Phagnalon rupestre.
rutilans, Poir. Encyc. Suppl. ii. 341 = Vernonia senegalensis.
sagittalis, Lam. Encyc. ii. 94 = Baccharis sagittalis.
salicifolia, Lam. Encyc. ii. 89 = Monarrhenus salicifolius.
salicifolia, *Mill. Gard. Dict.* ed. VIII. n. 6.—Ind. occ.
salicina, *Kellogg, in Proc. Calif. Acad.* iii. (1863) 36.—Calif.
salicina, Rupr. Fl. Ingric. 568 = Inula salicina.
saligna, Wall. Cat. n. 3061 = Vernonia saligna.
salsoloides, Turcz. in Bull. Soc. Nat. Mosc. v. (1832) 197 = Inula ammophila.
salvifolia, Bory, Voy. iii. 176 = Blumea salvifolia.
saxatilis, Linn. Sp. Pl. ed. II. 1206 = Phagnalon saxatile.
saxatilis, Sibth. & Sm. Fl. Graec. t. 862 = Phagnalon graecum.
saxatilis, Tenore, Fl. Neap. ii. 213 = Phagnalon rupestre.
saxatilis, Zoll. ex C. B. Clarke, Comp. Ind. 90 = Blumea balsamifera.
scabiosaefolia, *Remy, in C. Gay, Fl. Chil.* iv. 72.—Chili.
scabra, *Linn. Mant.* 113.—Ind. or.
scabrida, *DC. Prod.* v. 387.—Afr. austr.
scandens, *Mill. Gard. Dict.* ed. VIII. n. 11.—Mexic.
scandens, Wall. Cat. n. 3060 = Vernonia scandens.
Schimperi, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 386 = stricta.
Schlechtendalii, *C. Bolle, in Bonplandia*, vii. (1859) 294.—Ins. Canar.
scorpioides, Lam. Encyc. ii. 88 = Vernonia scorpioides.
semipinnatifida, *Wall. Cat.* n. 3058.—Ind. or.
semivestita, Wall. l. c. n. 2996 = Blumea procera.
senegalensis, Sieber, ex DC. Prod. v. 449 = Laggera aurita.
senegalensis, *Willd. Sp. Pl.* iii. 1928.—Afr. trop.
sericea, *Bory*, Voy. ii. 376.—Ins. Borbon.
sericea, [Soland. in] Ait. Hort. Kew. ed. I. iii. 183 (= *Schizogyne sericea*).—Ins. Canar.
sericea, Wall. Cat. n. 3021 = Blumea glomerata.
serici-vestita, Wall. l. c. n. 2996 = Blumea chinensis.
serpentaria, *Griseb. in Goett. Abh.* xxiv. (1879) 176.—Reg. Argent.

CONYZA:—
serrata, Bory, ex DC. Prod. vi. 376 = Senecio salicifolius.
serrulata, *C. B. Clarke, Comp. Ind.* 64.—Ind. or.
serrulata, Lam. Encyc. ii. 85 = Baccharis serrulata.
serrulata, Wall. Cat. sub n. 3024 = viscidula.
sessifolia, Blume, Bijdr. 897 = Pluchea sessifolia.
sessiliflora, *Miq. Fl. Ind. Bat.* ii. 48.—Ins. Timor.
sessiliflora, *Salisb. Prod.* 195.—Ins. Bourbon.
sessilifolia, Steud. Nom. ed. II. i. 415 = Pluchea sessifolia.
sessilifolia, Wall. Cat. n. 3029 = Pluchea Wallichiana.
sicula, Willd. Sp. Pl. iii. 1931 = Pulicaria sicula.
sinensis, J. F. Gmel. Syst. 1221 = Blumea chinensis.
sinuata, Ell. Sketch, ii. 323 = Erigeron linifolius.
solidaginoides, Wall. Cat. n. 3066 = Aster ignoratus.
sonchifolia, Buch.-Ham. in Wall. l. c. n. 3085 = Blumea laciniata.
sophiaefolia, *H. B. & K. Nov. Gen. et Sp.* iv. 73. t. 326.—Mexic.
sordida, Linn. Mant. 466 = Phagnalon sordidum.
spathulata, *Hornem. Hort. Hafn.* ii. 805.—Hab. ?
spathulifolia, Maxim. ex Regel, in Act. Hort. Petrop. iii. (1874) 109 = stricta.
spatulata, Link, Enum. Hort. Berol. ii. 322 = Vernonia pauciflora.
spicata, Lam. Encyc. ii. 93 = Pterocaulon virgatum.
spicata, *Mill. Gard. Dict.* ed. VIII. n. 14.—Hab. ?
spinellosa, *Miq. Fl. Ind. Bat.* ii. 45.—Burma.
spinidens, Miq. l. c. 44 = Blumea membranacea.
spinosa, *Sch. Bip. ex Oliver & Hiern, in Fl. Trop. Afr.* iii. 312.—Afr. trop.
spinulosa, Spreng. Syst. iii. 510 = Sommerfeldtia spinulosa.
splendens, Spreng. l. c. 509 = Piptocarpa lucida.
Sprengelii, *Sch. Bip. ex Walp. Rep.* ii. 958.—Afr. austr.
squamata, Spreng. Syst. iii. 515 = Aster divaricatus.
squamosa, Bory, Voy. iii. 110 = Senecio squamosus.
squarrosa, *Georgi, It.* i. 229.—Sibir. baical.
squarrosa, Linn. Sp. Pl. 861 = Inula vulgaris.
squarrosa, Wall. Cat. n. 3025 = Blumea myriocephala.
stachyoides, Spreng. ex DC. Prod. v. 442 = Blumea glomerata.
Steetzii, *Griseb. App. Ind. Sem. Hort. Gotting.* (1878) 8.—Afr. austr.
stellata, Spreng. Neue Entdeck. ii. 142 = Vernonia oppositifolia.
Steudelii, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 388.—Abyssin.
stipulacea, *Desf. Cat. Hort. Par.* ed. III. 164.—Hab. ?
stipulata, Vahl, ex DC. Prod. v. 97 = Liabum Jussieui.
striata, Wall. Cat. n. 3065 = viscidula.
stricta, Wall. l. c. n. 3044 = japonica.
stricta, *Willd. Sp. Pl.* iii. 1922.—Ind. or.; Afr. trop.
subcapitata, Wall. Cat. n. 3056 = Blumea subcapitata.
subdecurrens, DC. Prod. v. 379 = Erigeron subdecurrens.
subdecurrens, A. Gray, Pl. Fendl. 78; Pl. Wright. i. 102 = Coulteri.
subracemosa, *Miq. Fl. Ind. Bat.* ii. 41.—Java.
subsessilis, Wall. Cat. n. 3000 = Vernonia subsessilis.
subsimplex, Wall. l. c. n. 3003 = Vernonia cinerea.
subspicata, *Phil. in Linnaea*, xxviii. (1856) 735.—Chili.
suffruticosa, *Phil. l. c.*—Chili.
sumbavensis, *Miq. Fl. Ind. Bat.* ii. 46.—Malaya.
sylvatica, Blume, Bijdr. 898 = Blumea sylvatica.
symphytifolia, *Mill. Gard. Dict.* ed. VIII. n. 10.—Hab. ?
syringaefolia, *Meyen & Walp. in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) 263.—China.
tenella, *Miq. Fl. Ind. Bat.* ii. 40.—Ins. Timor.
tenera, Wall. Cat. n. 3023 = Blumea oxyodonta.
Tenorii, Presl, ex Spreng. Pugill. i. 55 = Phagnalon rupestre.
tenuifolia, Salisb. Prod. 194 (= *Schizogyne sericea*).—Ins. Canar.
ternatensis, *Miq. Fl. Ind. Bat.* ii. 49.—Ins. Ternate.
tetraptera, Turcz. in Bull. Soc. Nat. Mosc. xxiv. (1851) i. 178 = Blumea tetraptera.
thapsoides, Bieb. Fl. Taur. Cauc. ii. 304 = Inula thapsoides.
thermarum, *Phil. in Anal. Univ. Chil.* (1862) ii. 381.—Chili.
thesiifolia, *H. B. & K. Nov. Gen. et Sp.* iv. 75.—Mexic.

CONYZA:—
thuyoides, Lam. Encyc. ii. 90 = Baccharis thyoides.
thyrsifera, C. B. Clarke, Comp. Ind. 76, sphalm. = seq.
thyrsiflora, Wall. Cat. n. 3082 = Blumea lacera.
thyrsoidea, Pers. Syn. ii. 426. n. 8 = Blumea lacera.
tigrensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 316.—Afr. trop.
timorensis, *Miq. Fl. Ind. Bat.* ii. 47.—Ins. Timor.
tomentosa, Burm. f. Fl. Cap. Prod. 26 = Alciope lanata.
tomentosa, Forsk. Fl. Aegypt. Arab. 148 = Phagnalon rupestre.
tomentosa, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
tomentosa, Wall. Cat. n. 3031 = Pluchea tomentosa.
tortuosa, *Linn. Sp. Pl.* 862.—Am. austr.
travancorensis, Wall. Cat. n. 3067 = viscidula.
tricuneata, Willd. Sp. Pl. iii. 1946 = Baccharis conferta.
tridentata, Poir. Encyc. Suppl. ii. 342 = Baccharis tridentata ?
triloba, Decne. in Ann. Sc. Nat. Sér. II. ii. (1834) 261 = Erigeron trilobus.
trinervia, *Mill. Gard. Dict.* ed. VIII. n. 12.—Mexic.
trinervis, Lam. Encyc. ii. 85 = Baccharis trinervis.
triplinervia, *Less. in Linnaea*, vi. (1831) 137.—Bras.
trisulca, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 172 = stricta.
tuberosa, Lam. ex Steud. Nom. ed. II. i. 415 = Jasonia radiata.
uliginosa, Pers. Syn. ii. 426 = Pluchea bifrons.
umbrosa, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 379.—Sibiria.
undulata, *Poir. Encyc. Suppl.* ii. 339.—Afr. trop.
uniflora, Banks & Soland. ex Hook. f. Fl. N. Zel. i. 140 = Helichrysum filicaule.
uniflora, *Mill. Gard. Dict.* ed. VIII. n. 13.—Mexic.
vagans, Wall. Cat. n. 3040 = Vernonia vagans.
variegata, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 389.—Abyssin.
Vatkeana, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 314.—Afr. trop.
verbascifolia, Bieb. ex Georgi, Beschr. Russ. Reich. iv. 1239 = Inula thapsoides.
verbascifolia, Bory, Voy. ii. 377 = anchusaefolia.
verbascifolia, Frivald. ex Nym. Consp. 393 = Inula Aschersoniana.
verbascifolia, Willd. Sp. Pl. iii. 1924 = Inula candida.
veronicaefolia, Wall. Cat. n. 3005 = japonica.
verrucosa, Houtt. ex DC. Prod. vii. 13 = Printzia Bergii.
vestita, Wall. Cat. n. 2998 = Blumea balsamifera.
villosa, Herb. Madr. ex Wall. l. c. n. 3105 = Blumea amplectens.
villosa, Willd. Sp. Pl. iii. 1929 = Laggera aurita.
viminea, *Miq. Fl. Ind. Bat.* ii. 49.—Malaya.
virens, Wall. Cat. n. 3037 = Blumea virens.
virgata, *DC. Prod.* v. 385.—Madag.
virgata, Linn. Sp. Pl. ed. II. 1206 = Pterocaulon virgatum.
viscidula, *Wall. Cat.* n. 3006.—As. trop.; Austral.
viscosa, *Boj. ex Sweet, Hort. Brit.* ed. III. 349.—Ins. Maurit.
viscosa, Hort. ex Steud. Nom. ed. II. i. 415 = Inula viscosa.
viscosa, *Mill. Gard. Dict.* ed. VIII. n. 8.—Mexic.
viscosula, Wall. ex DC. Prod. v. 441 = Blumea membranacea.
volkameriaefolia, Wall. Cat. n. 3001 = Vernonia volkameriaefolia.
volubilis, Wall. l. c. n. 3057 = Microglossa volubilis.
vulgaris, Lam. Fl. Fr. ii. 78 = Inula vulgaris.
vulgaris, *Phil. in Linnaea*, xxviii. (1856) 735.—Chili.
Wallichii, DC. Prod. v. 384 = viscidula.
wangtuensis, *C. B. Clarke, Comp. Ind.* 61.—Reg. Himal.
Wightiana, Wall. Cat. n. 3028 = Vernonia arborea.
Wightiana, Wall. ex DC. Prod. v. 435 = Blumea Wightiana.
Wulfeni, Host, ex Beck, in Denkschr. Akad. Wien, Math. Nat. xliv. (1882) 332 = Inula bifrons.
Zollingeriana, *Turcz. in Bull. Soc. Nat. Mosc.* xxiv. (1851) i. 179.—Java.

CONYZELLA, Rupr. in Mém. Acad. Pétersb. Sér. VII. xiv. n. 4 (1869); Sert. Tiansch. 51 = **Erigeron**, Linn. (Compos.).
canadensis, Rupr. l. c. = E. canadensis.

CONYZOIDES, Tourn. ex DC. Prod. vi. 281 (1837) = **Carpesium**, Linn. (Compos.).

COOCKIA, Batsch, Tab. 53 (1802) = COOKIA, Sonner.

COOKIA, J. F. Gmel. Syst. 24 (1791) = **Pimelea**, Banks & Soland. (Thymel.).
Gnidia, J. F. Gmel. l. c. = P. Gnidia.
pilosa, J. F. Gmel. l. c. = P. pilosa.
prostrata, J. F. Gmel. l. c. = P. prostrata.

COOKIA, Sonner. Voy. Ind. iii. 258. t. 130 (1782) = **Clausena**, Burm. f. (Rutac.).
anisata, Desf. Cat. Hort. Par. ed. III. 415, 406.—Hab. ?
anisodora, Blanco, Fl. Filip. ed. II. 253 = Clausena excavata.
Anisum-olens, Blanco, l. c. ed. I. 359 = Clausena excavata.
australis, F. Muell. Fragm. i. 25 = Pentaceras australis.
chlorosperma, Blume, Bijdr. 135 = Glycosmis chlorosperma.
cyanocarpa, Blume, l. c. 136 = Glycosmis cyanocarpa.
dulcis, Bedd. in Madr. Journ. Sc. Ser. II. xxii. (1861) 73 = Clausena Willdenowii.
falcata, DC. Prod. i. 537.—China.
glabrescens, Miq. Fl. Ind. Bat. i. ii. 524.—Ins. Philipp.
graveolens, Wight & Arn. Prod. 95 = Clausena excavata.
macrophylla, Lindl. in Wall. Cat. n. 6367 = Clausena heptaphylla.
pentaphylla, Wight & Arn. ex M. Roem. Syn. Hesper. 45 = Clausena pentaphylla.
punctata, Hassk. Cat. Hort. Bog. Alt. 215 = Micromelum pubescens.
punctata, Retz. Obs. vi. 29 = Clausena excavata.
punctata, Sonner. Voy. Ind. iii. 258. t. 130 = Clausena Wampi.
Sonneratii, Montr. in Mém. Acad. Lyon, x. (1860) 184.—N. Caled. ?
Wampi, Blanco, Fl. Filip. ed. I. 358 = Clausena Wampi.

COOPERIA, Herb. in Bot. Reg. t. 1835 (1836).
AMARYLLIDEAE, Benth. & Hook. f. iii. 722.
SCEPTRANTHUS, R. Grah. in Edinb. N. Phil. Journ. xx. (1838) 413.
chlorosolen, Herb. in Bot. Reg. t. 1835 = Drummondi.
Drummondii, *Herb. l. c.*—Texas.
mexicana, Herb. Amaryll. 182 = Drummondii.
pedunculata, *Herb. l. c.* 179. t. 42. ff. 3–5.—Texas.

COPAIBA, Adans. Fam. ii. 341 (1763) = seq.

COPAIFERA, Linn. Sp. Pl. ed. II. 557 (1762).
LEGUMINOSAE, Benth. & Hook. f. i. 585.
COPAIBA, Adans. Fam. ii. 341 (1763).
COPAIVA, Jacq. Enum. Pl. Carib. 4 (1760).
GORSKIA, Bolle, in Peters, Reise Mossamb. Bot. i. 15. t. 3 (1861).
GUIBOURTIA, J. J. Benn. in Journ. Linn. Soc. i. (1857) 149.
Beyrichii, *Hayne, Arzneig.* x. t. 12.—Bras.
bijuga, *Willd. ex Hayne, l. c.* t. 16.—Bras.
bracteata, *Benth. in Hook. Journ. Bot.* ii. (1840) 101.—Guiana.
coleosperma, *Benth. in Trans. Linn. Soc.* xxv. (1866) 316. f. 43.—Afr. trop.
confertiflora, *Benth. in Mart. Fl. Bras.* xv. ii. 241.—Am. trop.
copallina, Baill. Hist. Pl. ii. 142, 163 = Guibourtiana.
cordifolia, *Hayne, Arzneig.* x. t. 21.—Bras.
coriacea, *Mart. Reise Bras.* i. 285; *et in Hayne, l. c.* t. 20.—Bras.
disperma, *Willem. in Usteri, Ann. Bot.* xviii. (1796) 34.—Ins. Maurit.
elliptica, *Mart. in Flora*, xx. (1837) ii. *Beibl.* 127.—Bras.
glabra, *Vogel, in Linnaea*, xi. (1837) 410.—Bras.
Gorskiana, *Benth. in Trans. Linn. Soc.* xxv. (1866) 317.—Afr. trop.
guyanensis, *Desf. in Mém. Mus. Par.* vii. (1821) 376.—Guiana.
Guibourtiana, *Benth. in Trans. Linn. Soc.* xxv. (1866) 317.—Afr. trop.

COPAIFERA :—
hymenaeifolia, *Moric. in Mem. Soc. Phys. Genèv.* vi. (1833) 529.—Cuba.
Jacquini, Desf. in Mém. Mus. Par. vii. (1821) 376 = officinalis.
Jacquiniana, G. Don, Gen. Syst, ii. 455 = officinalis.
Jussieui, *Hayne, Arzneig.* x. sub t. 17.—Peruv.
Lansdorfii, *Desf. in Mém. Mus. Par.* vii. (1821) 377. —Bras.
laxa, *Hayne, Arzneig.* x. t. 18.—Bras.
Mannii, Baill. Adansonia, vi. (1866) 202 = Hardwickia Mannii.
marginata, *Benth. in Mart. Fl. Bras.* xv. II. 243.—Bras.
Martii, *Hayne, Arzneig.* x. t. 15.—Bras.
Mopane, J. Kirk, ex Benth. in Trans. Linn. Soc. xxv. (1866) 317.—Afr. trop.
multijuga, *Hayne, Arzneig.* x. sub t. 17.—Bras.
nitida, *Mart. ex Hayne, l. c.* x. t. 17.—Bras.
oblongifolia, *Mart. ex Hayne, l. c.* t. 23 ; *et in Mart. Reise Bras.* i. 545.—Bras.
officinalis, *Linn. Sp. Pl.* ed. II. 557.—Am. trop.
pubiflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 101. —Guiana.
rigida, *Benth. in Mart. Fl. Bras.* xv. II. 243.—Bras.
Sellowii, *Hayne, Arzneig.* x. t. 22.—Bras.
trapezifolia, *Hayne, l. c.* sub t. 23.—Bras.

COPAIVA, Jaeq. Enum. Pl. Carib. 4 (1760) = **Copaifera**, Linn. (1762).
officinalis, Jacq. l. c. 21 = Copaifera officinalis.

COPERNICIA, Mart. Hist. Nat. Palm. ii. 151. t. 50 A ; iii. 242 (1837 ?). *PALMAE*, Benth. & Hook. f. iii. 927.
CRYOSOPHILA, Blume, Rumphia, ii. 53, in nota (1836).
CRYSOPHILA, Benth. & Hook. f. Gen. iii. 928 (1883).
cerifera, *Mart. Hist. Nat. Palm.* iii. 242. t. 50 A.—Am. trop.
hospita, *Mart. l. c.*—Cuba.
maeroglossa, *H. Wendl. in Kerch. Palm.* 241, nomen. —Hab. ?
maritima, *Mart. Hist. Nat. Palm.* iii. 243. t. 50 A. —Cuba.
Miraguana, Mart. l. c. = Thrinax Miraguana.
nana, *Liebm. ex Hemsl. Biol. Centr. Am. Bot.* iii. 411. —Mexie.
Pumos, *Mart. Hist. Nat. Palm.* iii. 243,—Cuba.
robusta, *Hort. ex Linden, Illustr. Hortic.* xxviii. (1881) 16.—Cuba.
tectorum, *Mart. Hist. Nat. Palm.* iii. 243.—N. Granat.
Wrightii, *Griseb, Cat. Pl. Cub.* 220.—Cuba.

COPIANTHUS, Hill, Decade t. 1 (1773). Amaranthac. ? monstr.
indica, Hill, l. c.—Ind. or.

COPIOGLOSSA, Miers, in Proc. Roy. Hort. Soc. iii. (1863) 294 = **Ruellia**, Linn. (Acanthac.).
pilosa, Miers, l. c. (= *Dipteracanthus Neesianus*).—Bras.

COPISMA, E. Mey. Comm. Pl. Afr. Austr. 132 (1835) = **Rhynchosia**, Lour. (Legumin.).
caribaeum, E. Mey. l. c. 133 = R. caribaea.
diversifolium, E. Mey. l. c. 135 = R. hirsuta.
effusum, E. Mey. l. c. = R. adenodes.
falcatum, E. Mey. l. c. 136 = R. minima.
gibbum, E. Mey. l. c. 137 = R. caribaea.
glabrum, E. Mey. l. c. 134 = R. glandulosa.
glandulosum, E. Mey. l. c. 133 = R. glandulosa.
grandifolium, E. Mey. l. c. 138 = R. grandifolia.
nitidum, E. Mey. l. c. 136 = R. nitida.
paniculatum, E. Mey. l. c. 134 = R. totta.
pictum, E. Mey. l. c. 135 = R. caribaea.
pilosum, E. Mey. l. c. 133 = R. pilosum.
rotundifolium, E. Mey. l. c. 137 = R. rotundifolia.
secundum, Walp. in Linnaea, xiii. (1839) 537 = R. secunda.
simplicifolium, E. Mey. Comm. Pl. Afr. Austr. i. 138 = R. singulifolia.
singulifolium, Steud. Nom. ed. II. ii. 454, sphalm. = R. singulifolia.
subsericeum, Sond. in Linnaea, xxiii. (1850) 34 = Galactia tenuiflora.
tenue, E. Mey. Comm. Pl. Afr. Austr. 136 = R. minima.
tottum, E. Mey. l. c. 133 = R. puberula.
trichodes, E. Mey. l. c. 134 = R. totta.
viscidulum, E. Mey. l. c. = R. viscidula.

COPPENSIA, Dum. in Mém. Acad. Brux. ix. (1835) (Sur Maelen.) 10, in nota. = **Oncidium**, Sw. (Orchid.).
bifolia, Dum. l. c. = O. bifolium.

COPPOLERIA, Todaro, in Atti Acc. Se. Palermo, N. S. i. (1845) 14 = **Vicia**, Tourn. (Legumin.).
monantha, Todaro. l. c. = V. monantha.

COPROSMA, Forst. Char. Gen. 137. t. 69 (1776). *RUBIACEAE*, Benth. & Hook. f. ii. 139.
EUARTHRONIA, Nutt. ex A. Gray, in Proc. Am. Acad. iv. (1860) 49, in syn.
EURYNOME, DC. Prod. iv. 475 (1830).
MARQUISIA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 192.
PELAPHIA, Banks & Soland. ex A. Cunn. in Ann. Nat. Hist. ii. (1839) 206.
acerosa, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 207. —N. Zel.
acutifolia, *Hook. f. in Journ. Linn. Soc.* i. (1857) 128. —N. Zel.
acutifolia, F. Muell. ex Benth. Fl. Austral. iii. 429 = Morinda acutifolia.
affinis, Hook. f. Fl. Antarct. i. 21. t. 14 = foetidissima.
arborea, *T. Kirk, in Trans. N. Z. Inst.* x. (1878) 420. —N. Zel.
arcuata, *Colenso, Excurs. North. Isl.* 84.—N. Zel.
Baueri, *Endl. Iconog.* t. 111.—Ins. Norfolc. ; N. Zel.
Baueriana, Hook. f. Fl. N. Zel. i. 104 = Baueri.
Billardieri, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 465 [bis].—Austral.
calycina, A. Gray, in Proc. Am. Acad. iv. (1860) 306 (= *Leptostigma Arnottianum*).—Chili.
ciliata, *Hook. f. Fl. Antarct.* i. 22.—Ins. Auckland.
Colensoi, *Hook. f. Handb. New Zeal. Fl.* 117.—N. Zel.
coneinna, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 330.—N. Zel.
conferta, A. Cunn. in Ann. Nat. Hist. ii. (1839) 206 = Cunninghamii.
crassifolia, *Colenso, Excurs. North. Isl.* 75.—N. Zel.
cuneata, *Hook. f. Fl. Antarct.* i. 21. t. 15.—N. Zel.
Cunninghamii, *Hook. f. Handb. New Zeal. Fl.* 113.—N. Zel.
cuspidifolia, DC. Prod. iv. 578 = hirtella.
depressa, *Colenso, ex Hook. f. Fl. N. Zel.* 110.—N. Zel.
divaricata, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 207. —N. Zel.
ernodeoides, *A. Gray, in Proc. Am. Acad.* iv. (1860) 49.—Ins. Pacif.
foetidissima, A. Cunn. in Ann. Nat. Hist. ii. (1839) 206 = Cunninghamii.
foetidissima, *Forst. Char. Gen.* 138.—N. Zel.
foliosa, *A. Gray, in Proc. Am. Acad.* iv. (1860) 48.—Ins. Sandvic.
gracilis, A. Cunn. in Ann. Nat. Hist. ii. (1839) 206 = rhamnoides ?
grandifolia, *Hook. f. Fl. N. Zel.* 104.—N. Zel.
hirtella, *Labill. Nov. Holl. Pl.* i. 70. t. 95.—Austral.
lanceolaris, *F. Muell. Fragm.* ix. 70.—Austral.
ligustrina, *Kunth, Ind. Sem. Hort. Berol.* (1846) 13. —Hab. ?
linariifolia, *Hook. f. Handb. New Zeal. Fl.* 118.—N. Zel.
longifolia, *A. Gray, in Proc. Am. Acad.* iv. (1860) 48. —Ins. Sandvic.
lucida, *Forst. Char. Gen.* 138.—N. Zel.
Menziesii, *A. Gray, in Proc. Am. Acad.* iv. (1860) 49.—Ins. Pacific.
microcarpa, *Hook. f. Fl. N. Zel.* 110.—N. Zel.
microphylla, A. Cunn. ex Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 465 [bis] = Billardieri.
myrtifolia, *Noronha, in Ver. Batav. Gen.* v. (1790) ed. I. Art. IV. 10.—Java.
myrtillifolia, Hook. f. Fl. Antarct. i. 21 = parviflora.
Nertera, F. Muell. Fragm. ix. 186 = Nertera depressa.
nitida, *Hook. f. in Hook. Lond. Journ. Bot.* vi. (1847) 465 [bis].—Austral.
parviflora, *Hook. f. Fl. N. Zel.* 106.—N. Zel.
persicaefolia, *A. Gray, in Proc. Am. Acad.* iv. (1860) 50.—Ins. Fiji.
petiolata, *Hook. f. in Journ. Linn. Soc.* i. (1857) 128. —Ins. Kermadec.
pilosa, *Endl. Fl. Ins. Norf. Prod.* 60.—Ins. Norfolc.
propinqua, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 206.—N. Zel.
pubens, *A. Gray, in Proc. Am. Acad.* iv. (1860) 49.—Ins. Sandvic.

COPROSMA :—
pumila, *Hook. f. Fl. N. Zel.* 111.—Austral. ; N. Zel.
pusilla, Forst. f. Prod. 90 ; A. Rich. Voy. Astrol. Bot. i. 265 = foetidissima.
putida, *C. Moore & F. Muell. Fragm.* vii. 45.—Ins. D. Howe.
repens, A. Rich. Voy. Astrol. Bot. i. 264 = foetidissima.
repens, *Hook. f. Fl. Antarct.* i. 22. t. 16.—N. Zel.
reptans, *F. Muell. Fragm.* ix. 186.—Austral.
retusa, Banks & Soland. ex Hook. f. in Hook. Lond. Journ. Bot. iii. (1844) 415 = Baueri.
rhamnoides, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 206.—N. Zel.
rhynchocarpa, *A. Gray, in Proc. Am. Acad.* iv. (1860) 48.—Ins. Sandvic.
robusta, *Raoul, in Ann. Sc. Nat.* Sér. III. ii. (1844) 121.—N. Zel.
rotundifolia, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 206.—N. Zel.
rubra, *Petrie, in Trans. N. Z. Inst.* xvii. (1885) 269.—N. Zel.
serrulata, *Hook. f. ex J. Buch. l. c.* iii. (1871) 212.—N. Zel.
spathulata, *A. Cunn. in Ann. Nat. Hist.* ii. (1839) 207.—N. Zel.
Stocki Hort. ex J. Barbier, in Rev. Hortic. Belg. iii. (1877) t. 12 = Baueri.
sundana, *Miq. Fl. Ind. Bat.* ii. 327.—Malaya.
taitensis, *A. Gray, in Proc. Am. Acad.* iv. (1860) 49. —Ins. Tahiti.
tenuicaulis, *Hook. f. Fl. N. Zel.* 106.—N. Zel.
triflora, *Benth. & Hook. f. Gen.* ii. 139.—Ins. Juan Fernand.
venosa, A. Cunn. ex Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 465 [bis] = hirtella.
vireseens, *Petrie, in Trans. N. Z. Inst.* xi. (1879) 426. —N. Zel.
Waimeae, *Wawra, in Flora*, lvii. (1874) 327.—Ins. Pacif.

COPROSMANTHUS, Kunth, in Abh. Akad. Berl. (1848) 35 ; et Enum. Pl. v. 263 (1850) = **Smilax**, Linn. (Liliac.).
consanguineus, Kunth, Enum. Pl. v. 268 = S. herbacea.
herbaceus, Kunth, l. c. 265 = S. herbacea.
japonicus, Kunth, l. c. 268 = S. China.
lasioneuron, Kunth, l. c. 265 = S. herbacea.
pedunculatis, Kunth, l. c. 264 = S. herbacea.
tamnifolius, Kunth, l. c. 267 = S. tamnifolia.

COPTIDIUM, Nym. Consp. 13 (1878) = **Ranunculus**, Tourn.
ranunculoides, Nym. l. c. = R. lapponieus.

COPTIS, Salisb. in Trans. Linn. Soc. viii. (1807) 305 *RANUNCULACEAE*, Benth. & Hook. f. i. 8.
CHRYSOCOPTIS, Nutt. in Journ. Acad. Philad. vii. (1834) 9. t. 1.
anemonaefolia, *Sieb. & Zucc. in Abh. Math. Phys. Kl. Muench.* iv. II. (1843) 180.—Japon.
asplenifolia, *Salisb. in Trans. Linn. Soc.* viii. (1807) 306.—Am. bor.
brachypetala, *Sieb. & Zucc. in Abh. Math. Phys. Kl. Muench.* iv. II. (1843) 180.—Japon.
occidentalis, *Torr. & Gray, Fl. N. Am.* i. 28.—Am. bor. occ.
orientalis, *Maxim. in Bull. Acad. Pétersb.* xii. (1868) 60.—Japon.
quinquefolia, *Miq. Ann. Mus. Bot. Lugd. Bat.* iii. 7.—Japon.
Teeta, *Wall. in Linnaea*, xii. (1838) *Litt.* 227.—Reg. Himal.
trifolia, *Salisb. in Trans. Linn. Soc.* viii. (1807) 305.—Reg. bor. temp. et arct.

COPTOCHEILE, Hoffmgg. Verz. Pfl. Nachtr. (1841) ex Linnaea, xvi. (1842) Litt. 279. *GESNERACEAE* ?
maerorhiza, *Hoffmgg. ll. cc.*—Bras. ?

COPTOPHYLLUM, Korth. in Ned. Kruidk. Arch. ii. (1851) 161. *RUBIACEAE*, Benth. & Hook. f. ii. 68.
bracteatum, *Korth. l. c.* 162.—Sumatra.
capitatum, *Miq. Fl. Ind. Bat.* ii. 348.—Malaya.
pilosum, *Miq. Ann. Mus. Bot. Lugd. Bat.* iv. 230.—Java.

COPTOSAPELTA, Korth. in Ned. Kruidk. Arch. ii. ii. (1851) 112. *RUBIACEAE*, Benth. & Hook. f. ii. 35.
flavescens, *Korth. l. c.* 113.—Malaya.
Griffithii, *Hook. f. Ic. Pl.* t. 1059.—Malaya.

COPTOSPERMA, Hook. f. in Benth. & Hook. f. Gen. ii. 86 (1873). *RUBIACEAE*, Benth. & Hook. f. l. c.
nigrescens, *Hook. f. l. c.* 87.—Afr. trop.

COQUEBERTIA, Brongn. in Ann. Sc. Nat. Sér. I. xxx. (1833) 111 = **Zollernia**, Maximil. & Nees (Legumin.).
ilicifolia, Brongn. l. c. = Z. ilicifolia.

CORALLIORRHIZA, Aschers. Fl. Brandenb. 697 (1864) = **Corallorrhiza**, Hall. (Orchid.).

CORALLOBOTRYS, Hook. f. in Benth. & Hook. f. Gen. ii. 575 (1876). *VACCINIACEAE*, Benth. & Hook. f. l. c.
acuminata, *Hook. f. Fl. Brit. Ind.* iii. 455.—Reg. Himal.

CORALLOCARPUS, Welw. ex Benth. & Hook. f. Gen. i. 831 (1867). *CUCURBITACEAE*, Benth. & Hook. f. l. c.
conocarpus, *Benth. & Hook. f. l. c.*—Ind. or.; Afr. trop.
corallinus, Cogn. in DC. Monog. Phan. iii. 647 = Fenzlii.
Courbonii, Cogn. l. c. 655 = velutinus.
Ehrenbergii, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 566.—Nubia.
epigaeus, *Benth. & Hook. f. Gen.* i. 831.—Ind. or.
erostris, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 567.—Afr. trop.
etbaicus, Hook. f. l. c. = velutinus.
Fenzlii, *Hook. f. l. c.* 565.—Afr. trop.
Gijef, *Hook. f. l. c.* 566.—Nubia.
glomeratus, *Cogn. in DC. Monog. Phan.* iii. 658.—Venezuela.
gracilipes, *Cogn. l. c.* 650.—Ind. or.
palmatus, *Cogn. l. c.* 648.—Ind. or.
parvifolius, *Cogn. l. c.* 658.—Arabia.
pedunculosus, *Cogn. l. c.* 654.—Abyssin.
Poissonii, *Cogn. l. c.* 651.—Madag.
Schimperi, *Hook. f. in Oliver, Fl. Trop. Afr.* ii. 567.—Abyssin.
velutinus, *Benth. & Hook. f. Gen.* i. 831.—Ind. or.
Welwitschii, *Benth. & Hook. f. l. c.*—Afr. trop.

CORALLODENDRON, Tourn. ex Rupp. Fl. Jen. ed. Hall. 257 (1745) = **Erythrina**, Linn. (Legumin.).
aculeatum, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 341 = E. Corallodendron.

CORALLOPHYLLUM, H. B. & K. Nov. Gen. et Sp. vii. 275. t. 660 bis (1825) = **Lennoa**, La Llave & Lex.
caeruleum, H. B. & K. l. c. 276 = L. caerulea.

CORALLORRHIZA, Hall. Enum. Stirp. Helv. i. 278 (1742) (Corallorhiza); R. Br. in Ait. Hort. Kew. ed. II. v. 209 (1813). *ORCHIDEAE*, Benth. & Hook. f. iii. 497.
RHIZOCORALLON, Hall. in Rupp. Fl. Jen. 301 (1745).
arizonica, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 379.—Arizona.
Bigelovii, *Wats. l. c.* xii. (1877) 275.—Calif.
bulbosa, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 19 = mexicana.
dentata, Host, Fl. Austr. ii. 547 = innata.
Ehrenbergii, *Reichb. f. in Linnaea*, xxii. (1849) 833.—Mexic.
ericetorum, Drej. in Fl. Dan. t. 2364 = innata.
foliosa, *Lindl. Gen. et Sp. Orch.* 535.—Ind. or.
grandiflora, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 19.—Mexic.
Halleri, Rich. in Mém. Mus. Par. iv. (1818) 61 = innata.
hiemalis, *Nutt. Gen. Am.* ii. 198.—Am. bor.
indica, *Lindl. in Journ. Linn. Soc.* iii. (1859) 26.—Ind. or.
innata, *R. Br. in Ait. Hort. Kew.* ed. II. v. 209.—Europ.; Am. bor.
intacta, Cham. & Schlecht. in Linnaea, iii. (1828) 35 = innata.
Jacquemontii, *Decne. in Jacquem. Voy. Bot.* 165. t. 163.—Reg. Himal.
Macraei, A. Gray, Man. Bot. N. U. St. (1856) 453 = stricta.
Mertensiana, *Bong. Veg. Ins. Sitcha*, 165.—Alaska.
mexicana, *Lindl. Gen. et Sp. Orch.* 534.—Mexic.
micrantha, *Chapm. Fl. S. U. St.* 454.—Am. bor.
multiflora, *Nutt. in Journ. Acad. Philad.* iii. (1823) 138. t. 7.—Am. bor.
nemoralis, Sw. ex Nym. Consp. 686 = innata.
Neottia, Scop. Fl. Carn. ed. II. ii. 207 = innata.
odontorhiza, *Nutt. Gen. Am.* ii. 197.—Am. bor.
patens, Lindl. Gen. et Sp. Orch. 535 = Oreorchis patens.
squamata, Poir. in Dict. Sc. Nat. x. 375 = Dipodium squamatum.
striata, *Lindl. Gen. et Sp. Orch.* 534.—Am. bor.
verna, Nutt. in Journ. Acad. Philad. iii. (1823) 136 = innata.
virescens, Drej. ex Heynh. Nom. ii. 159, nomen = innata.
Wisteriana, Conrad, in Journ. Acad. Philad. vi. (1829) 145 = odontorhiza.

CORALLOSPARTIUM, Armst. in Trans. N. Z. Inst. xiii. (1881) 333. *LEGUMINOSAE*.
crassicaule, *Armst. l. c.*—N. Zel.

CORALLUMA. Schrank, ex Haw. Syn. Pl. Succ. ed. II. 47 (1819) = **Caralluma**, R. Br. (Asclep.).

CORBICHONIA, Scop. Introd. 264 (1777) = **Orygia**, Forsk. (Ficoid.).

CORBULARIA, Salisb. in Trans. Hort. Soc. i. (1812) 349 = **Narcissus**, Tourn. (Amaryllid.).
albicans, Haw. Revis. 122 = N. Bulbocodium.
aurea, Haw. Monog. (in Sweet, Brit. Flow. Gard. iv.) 1. n. 5 = N. Bulbocodium.
Bulbocodium, Haw. Revis. 123 = N. Bulbocodium.
cantabrica, Haw. Monog. 1. n. 1 = N. Bulbocodium.
conspicua, Haw. l. c. n. 8 = N. Bulbocodium.
filifolia, M. Roem. Syn. Ensat. 190 = N. Bulbocodium.
Gigas, Haw. Monog. 1. n. 10 = N. Bulbocodium.
Graëllsii, Webb. ex Willk. & Lange, Prod. Fl. Hisp. i. 150 = N. Bulbocodium.
hedraeantha, Webb & Heldr. Cat. Pl. Hisp. 220 = N. Bulbocodium.
lobulata, Haw. Revis. 122 = N. Bulbocodium.
monophylla, Dur. in Duch. Rev. Bot. ii. (1846–47) 425 = N. Bulbocodium.
nivalis, Nym. Consp. 713 = N. Bulbocodium.
obesa, Salisb. in Trans. Hort. Soc. i. (1812) 350 = N. Bulbocodium.
Schultesii, M. Roem. Syn. Ensat. 190 = N. Bulbocodium.
serotina, Haw. Monog. 1. n. 9 = N. Bulbocodium.
tenuifolia, Salisb. in Trans. Hort. Soc. i. (1812) 349 = N. Bulbocodium.
turgida, Salisb. l. c. 351 = N. Bulbocodium.

CORCHOROPSIS, Sieb. & Zucc. in Abh. Akad. Muench. iii. (1843) 737. t. 4. *TILIACEAE*, Benth. & Hook. f. i. 235.
crenata, *Sieb. & Zucc. l. c.*—Japon.

CORCHORUS, [Tourn.] Linn. Syst. ed. I (1735). *TILIACEAE*, Benth. & Hook. f. i. 235.
ANTICHORUS, Linn. Mant. 9. n. 1257 (1767).
CARICHERIA, Scop. Introd. 104 (1777).
GANJA, Reichb. Handb. 303 (1837).
MAERLENSIA, Vell. Fl. Flum. v. t. 112 (1827).
PALLADIA, Lam. Illustr. t. 285 (1792).
acutangulus, *Lam. Encyc.* ii. 104.—Cosmop. trop.
aesticans, Hill, Veg. Syst. xiv. 23 = aestuans.
aestuans, Blanco, Fl. Filip. ed. I. 441 = acutangulus.
aestuans, Forsk. Fl. Aegypt. Arab. 101 = trilocularis.
aestuans, Herb. Madr. ex Wall. Cat. p. 237. n. 1074 = urticaefolius.
aestuans, *Linn. Syst.* ed. X. 1079.—Am. trop.
alatus, *G. Don, Gen. Syst.* i. 542.—Afr. trop.
americanus, Schrank, ex Steud. Nom. ed. II. i. 416 = Sida erubescens.
angustifolius. *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* x. (1828) 18.—Afr. trop.
Antichorus, *Raeusch. Nom.* ed. III. 158.—Geront. trop.
arenarius, *H. B. & K. Nov. Gen. et Sp.* v. 339.—Mexic.
argutus, *H. B. & K. Nov. Gen. et Sp.* v. 337.—N. Granat.
asplenifolius, *Burch. Trav.* i. 400.—Afr. austr.
asplenifolius, E. Mey. ex Harv. & Sond. Fl. Cap. i. 229 = trilocularis.
australis, A. Cunn. ex F. Mucll. Fragm. iii. 8 = Cunninghami.
bifurcatus, *Mill. Gard. Dict.* ed. VIII. n. 5.—Jamaic.
brachycarpos, *Rich. ex Steud. Nom.* ed. II. i. 416.—Afr. trop.
brachycarpus, Guill. & Perr. Fl. Seneg. Tent. i. 89 = fascicularis.
bullatus, Jacq. Eclog. Pl. t. 163 = pilolobus.
Burmanni, DC. Prod. i. 505 = tridens.
campestris, Macfad. Fl Jamaic. i. 107 = aestuans.
capsularis, *Linn. Sp. Pl.* 529.—Cosmop. trop.
caryophylloides, Linn. Syst. ed. X. 1079 = siliquosus?
catharticus, Blanco, Fl. Filip. ed. I. 442 = olitorius.
cordifolius, Salisb. Prod. 366 = capsularis.
Coreta, Linn. Pl Jam. Pugill. 14 = siliquosus.
Cunninghami, *F. Muell. Fragm.* iii. 8.—Austral.
decemangularis, Roxb. Hort. Beng. 42; Fl. Ind. ii. 582 = olitorius.
depressus, Stocks, in Proc. Linn. Soc. i. (1848) 367 = C. Antichorus.
echinatus, *Benth. Fl. Austral.* i. 276.—Austral.
elachocarpus, *F. Muell. Fragm.* viii. 6.—Austral.
erodioides, *Balf. f. in Proc. Roy. Soc. Edinb.* xi. (1882) 504.—Ins. Socotra.
fascicularis, *Lam. Encyc.* ii. 104.—Geront. trop.
flexuosus, *Thunb. in Trans. Linn. Soc.* ii. (1794) 335.—Japon.
foliosus, *Spreng. Syst.* ii. 583.—Cuba.
frutescens, Lam. Encyc. ii. 105 = hirsutus.
fruticulosus, Vis. in Flora, xx. (1837) 443 = trilocularis.
furcatus, G. Don, Gen. Syst. i. 544 = seq.
fuscus, Roxb. Hort. Beng. 42; Fl. Ind. ii. 582 = acutangulus.
gracilis, *R. Br. in Salt, Abyss. App.* 65.—Abyssin.
grandiflorus, Spring, ex Mart. in Flora, xx. (1837) II. Beibl. 102 = Turnera serrata.
Grevеanus, *Baill. in Bull. Soc. Linn. Par.* i. (1885) 543.—Madag.
guadalupensis, *Spreng. Syst.* ii. 584.—Ins. Guadalup.
hirsutus, *Linn. Sp. Pl.* 530.—Ind. occ.
hirtus, *Linn. l. c.* ed. II. 747.—Am. calid.
hirtus, Thunb. Fl. Jap. 228 = Zelkova acuminata.
humilis, Munro, ex Stocks, in Proc. Linn. Soc. i. (1848) 367 = C. Antichorus.
humilis, *A. St. Hil. Fl. Bras. Mer.* i. 280.—Bras.
hygrophilus, *A. Cunn. ex Benth. Fl. Austral.* i. 276.—Austral.
japonicus, Thunb. Fl. Jap. 227 = Kerria japonica.
javanicus, *Burm. f. Fl. Ind.* 123. t. 36. f. 3.—Ind. or.
lanceolatus, *G. Don, Gen. Syst.* i. 543.—Guinea.
lanuginosus, Macfad. Fl. Jamaic. i. 108 = hirsutus.
lasiolobus, Spreng. Syst. ii. 583 = pilolobus.
leptocarpus, *A. Cunn. ex Benth. Fl. Austral.* i. 278.—Austral.
linearis, *Mill. Gard. Dict.* ed. VIII. n. 5.—Mexic.
linearis, Wall. Cat. n. 1070 = tridens.
longicarpus, *G. Don, Gen. Syst.* i. 543.—Guinea.
longipedunculatus, *Mast. in Oliver, Fl. Trop. Afr.* i. 262.—Afr. trop.
Marua, Buch.-Ham. in Wall. Cat. n. 6311 = capsularis.
microphyllus, Fresen. in Mus. Senckenb. ii. (1837) 156 = C. Antichorus.
mompoxensis, *H. B. & K. Nov. Gen. et Sp.* v. 338.—N. Granat.
muricatus, *Hochst. ex A. Rich. Tent. Fl. Abyss.* i. 81.—Abyssin.
muricatus, *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* x. (1828) 20.—Afr. trop.
olitorius, *Linn. Sp. Pl.* 529.—Cosmop. trop.
oppositiflorus, Hassk. in Hoev. & De Vriese, Tijdschr. xii. (1845) 126 = acutangulus.
orinocensis, *H. B. & K. Nov. Gen. et Sp.* v. 337.—Venezuela.
patens, Lehm. Sem. Hort. Hamb. (1823) ex Linnaea, iii. (1828) Lit. 9 = tridens.
pilolobus, *Link, Enum. Hort. Berol.* ii. 72.—Am. calid.
pilosus, *H. B. & K. Nov. Gen. et Sp.* v. 338. t. 487.—N. Granat.
polygonus *Schum. & Thonn. in Dan. Vid. Selsk. Afh.* x. (1828) 20.—Afr. trop.

CORCHORUS:—
prismatocarpus, *St. Hil. Fl. Bras. Mer.* i. 282.—Bras.
procumbens, *Boj. Hort. Maurit.* 43.—Afr. trop.
pseudocapsularis, *Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 670.—Afr. trop.
Pumilio, *R. Br. ex Benth. Fl. Austral.* i. 277.—Austral.
pumilus, *Buch.-Ham. ex Wall. Cat.* n. 6310.—Ind. or.
quadrangularis, *J. A. Schmidt, Beitr. Fl. Cap. Verd. Ins.* 293.—Ins. Cap. Verd.
quinquelocularis, Moench, Meth. 248 = olitorius.
quinquenervis, Hochst. ex A. Rich. Tent. Fl. Abyss. i. 80 = urticifolius.
scandens, *Thunb. in Trans. Linn. Soc.* ii. (1794) 335.—Japon.
secundiflorus, Moç. & Sessé, ex DC. Prod. i. 504 = siliquosus.
senegalensis, Juss. ex Steud. Nom. ed. II. i. 417 = tridens.
serraefolius, *Burch. Trav.* i. 537.—Afr. austr.
serraefolius, DC. Prod. i. 504 = trilocularis.
serratus, *Thunb. in Trans. Linn. Soc.* ii. (1794) 335.—Japon.
sidoides, *F. Muell. Fragm.* iii. 9.—Austral.
siliquosus, *Linn. Sp. Pl.* 529.—Am. calid.
siliquosus, Poepp. ex Steud. Nom. ed. II. i. 417 = foliosus.
siliquosus, Torr. & Gray, Fl. N. Am. i. 239 = pilolobus.
spinosus, Sieber, ex Steud. Nom. ed. II. i. 417 = Pavonia spinifex.
subscandens, Wall. Cat. n. 1074 = urticaefolius.
tetragona, *Mill. Gard. Dict.* ed. VIII. n. 4.—Hab.?
tetrangulus, Langsd. in Syll. Ratisb. ii. (1828) 168 = pilolobus
tomentillus, *F. Muell. Fragm.* iii. 10.—Austral.
tomentosus, *Thunb. Fl. Jap.* 228.—Japon.
Torresianus, *Gaudich. Bot. Voy. Bonite*, 477.—Ins. Marian.
tortipes, A. St. Hil. Fl. Bras. Mer. i. 281 = hirtus.
tridens, *Linn. Mant.* 566.—Cosmop. trop.
triflorus, Boj. Hort. Maurit. 43 = trilocularis.
trilocularis, Burm. f. Fl. Ind. 123. t. 37. f. 2 = tridens.
triloculatis, *Linn. Mant.* 77.—Geront. trop.
urticaefolius, *Wight & Arn. Prod.* 73.—Ind. or.; Afr. trop.
vermicularis, *F. Muell. Fragm.* iii. 10.—Austral.
villosissimus, *A. St. Hil. Fl. Bras. Mer.* i. 280.—Bras.
Walcottii, *F. Muell. Fragm.* iii. 9.—Austral.

CORDA, St. Lag. in Ann. Soc. Bot. Lyon, viii. (1881) 177 = **Cordia**, Linn.

CORDAEA, Spreng. Gen. ii. 581 (1831) = **Cyamopsis**, DC. (Legumin.).
fabaeformis, Spreng. l. c. = Cyamopsis psoralioides.

CORDEMOYA, Baill. Adansonia, i. 255 (1861) = **Mallotus**, Lour. (Euphorb.).
acuminata, Baill. l. c. = M. Baillonianus.
integrifolia, Baill. l. c. = M. integrifolius.

CORDIA, Linn. Gen. ed. I. 52 (1737). *BORAGINEAE*, Benth. & Hook. f. ii. 838.
Acnadena, Carpiphea, Coilanthera, Colococca, Ectemis, Novella, Piloisa, Quarena, Topiaris, Toquera, Rafin. Sylva Tellur. 39–43 (1838).
Borellia, Neck. Elem. i. 275, 276 (1790).
Cordiada, Vell. Fl. Flum. 98 (1825); ii. t. 156.
Cerdiana, Ruiz & Pav. Prod. 37. t. 6 (1794).
Firensia, Neck. Elem. i. 275, 276 (1790).
Gerascanthus, P. Br. Hist. Jamaic. 170. t. 29. f. 3 (1756).
Gynaion, A. DC. Prod. ix. 468 (1845).
Hemigymnia, Griff. in Calc. Journ. Nat. Hist. iii. (1843) 363.
Macielia, Vand. Fl. Lusit. et Bras. Spec. 14 (1788).
Macria, Tenore, in Mem. Soc. Ital. Moden. xxiv. (1848) 366.
Myxia, Caesalp. ex Linn. Hort. Cliff. 63 (1737) in syn.
Paradigma, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 30. t. 8.
Pavonia, Domb. ex Lam. Tabl. Encyc. i. 421 (1800).
Plethostephia, Miers, in Trans. Linn. Soc. Ser. II. i. (1875) 32. t. 8.
Rhabdocalyx, Lindl. Veg. Kingd. 629 (1847).
Sebestena, [Dill.] Gaertn. Fruct. i. 364. t. 76. f. 1 (1788).
Varronia, P. Br. Hist. Jamaic. 172 (1756); L. n. Syst. ed. X. 916 (1759).
abyssinica, R. Br. in Salt, Abyss. App. 65; ed. Germ. 473 = C. Myxa.
acuminata, Wall. Cat. n. 896; et in Roxb. Fl. Ind. ed. Carey, ii. 339 = Ehretia Wallichiana.
acutifolia, *Fresen. in Mart. Fl. Bras.* viii. 11.—Am. trop.
adnata, *DC. Prod.* ix. 493.—Cuba?
affinis, *Fresen. in Mart. Fl. Bras.* viii. 22.—Am. trop.
africana, Lam. Illustr. n. 1806 = C. Myxa.
alba, *Roem. & Schult. Syst.* iv. 466.—Ind. occ.
alliodora, *Cham. in Linnaea*, viii. (1833) 121.—Am. trop.
alnifolia, *Hornem. Hort. Hafn.* i. 227.—Hab.?
alnifolia, Roem. & Schult. Syst. iv. 801 = serratifolia.
ambigua, *Cham. in Linnaea*, v. (1830) 115.—Mexic.
ambigua. Cham. l. c. viii. (1833) 125 = anabaptista.
amphifolia, *A. DC. in DC. Prod.* ix. 481.—Maurit.?
anabaptista, *Cham. in Linnaea*, viii. (1833) 512.—Bras.
angiocarpa, *A. Rich. Fl. Cub. Fanerog.* ii. 110.—Cuba
angustifolia, *Roem. & Schult. Syst.* iv. 460.—Ind. occ.
angustifolia, Roxb. Hort. Beng. 17; Fl. Ind. i. 595 = Rothii.
aspera, *Forst. f. Prod.* 18. n. 109.—Austral.; Ins. Pacif.
asperrima, *DC. Prod.* ix. 498.—Jamaic.
asperrima, Spreng. Syst. i. 649 = Hyptis membranacea.
asterophora, *Mart. ex Fresen. in Mart. Fl. Bras.* viii. 5.—Am. trop.
avicularis, Salisb. Prod. 111 = nodosa.
Aubletii, *DC. Prod.* iv. 490.—Guiana.
bahiensis, DC. l. c. ix. 489 = cylindristachya.
Banalo, Blanco, Fl. Filip. ed. I. 124 = subcordata.
bantamensis, *Blume, Bijdr.* 843.—Java.
Berterii, Spreng. ex Griseb. Fl. Brit. W. Ind. 480 = salvifolia.
Beurreria, Willd. Sp. Pl. i. 1078, in syn. = Bourreria succulenta.
bicolor, *A. DC. ex DC. Prod.* ix. 485.—Guiana.
bifurcata, *Roem. & Schult. Syst.* iv. 466.—Peruv.
Blanchetii, *DC. Prod.* ix. 477.—Am. trop.
bogotensis. *Benth. Pl. Hartw.* 240.—N. Granat.
Boissieri, *A. DC. in DC. Prod.* ix. 478.—Mexic.
Bonplandiana, H. B. & K. Nov. Gen et Sp. iii. 77 = seq.
Bonplandii, *Roem. & Schult. Syst.* iv. 462.—Am. austr.
Bourreria, Linn. Syst. ed. X. 936 = Bourreria succulenta.
brachypoda, *DC. Prod.* ix. 487.—Am. trop.
brachytricha, *Fresen. in Mart. Fl. Bras.* viii. 13.—Am. trop.
bracteata, *DC. Prod.* ix. 472.—Cuba.
brevispicata, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 331.—Mexic.
Brownii, DC. Prod. ix. 499 = C. Myxa.
bullata, Roem. & Schult. Syst. iv. 462 = globosa.
buxifolia, *Juss. in Poir. Encyc.* vii. 47.—Ind. occ.
caffra, *Sond. in Linnaea*, xxiii. (1850) 81.—Afr. austr.
Callococca = Collococca.
calocephala, *Cham. in Linnaea*, iv. (1829) 488.—Bras.
calocoma, *Miq. Fl. Ind. Bat.* ii. 916.—Java.
calophylla, *Vahl, Eclog. Am.* iii. 5.—Guiana.
calyptrata, Bert. ex Spreng. Syst i. 649 = alba.
campanulata, Roxb. Hort. Beng. 17; Fl. Ind. i. 593 = subcordata.
campestris, *Warm. in Kjoeb. Vidensk. Meddel.* (1867) 12.—Bras.
cana, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 331.—Mexic.
candida, *Vell. Fl. Flum.* 98; ii. t. 155.—Bras.
canescens, *H. B. & K. Nov. Gen. et Sp.* iii. 73.—N. Granat.
canescens, Willd. ex Roem. & Schult. Syst. iv. 799 = polystachya.
caracasana, *DC. Prod.* ix. 489.—Venezuela.
Cerdana, Roem. & Schult. Syst. iv. 467 = alliodora.
chamaedrifolia, DC. Prod. ix. 499, in syn. = mirabiliflora.
chamaedryoides, *Roem. & Schult. Syst.* iv. 801.—Hab.?
Chamissoniana, *Steud. Nom.* ed. II. i. 417.—Bras.
cinerascens, DC. Prod. ix. 492 = cylindristachya.
coffeoides, *Warm, in Kjoeb. Vidensk. Meddel.* (1867) 4.—Bras.
Collococa, Aubl. Pl. Gui. i. 219 = nodosa.
Collococca, *Linn. Sp. Pl.* ed. II. 274.—Jamaic.
Collococca, Sw. ex Griseb. Fl. Brit. W. Ind. 479 = nitida.
copulata, *Poir. in Dict. Sc. Nat.* x. 410.—Java.
corchorifolia, *A. DC. in DC. Prod.* ix. 496.—Am. trop.
cordata, Steud. Nom ed. II. i. 417 = grandis.
cordifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 70.—N. Granat.
cordifolia, Wall. Cat. n. 894 = grandis.
coromandeliana, Retz. ex DC. Prod. ix. 509 = Ehretia buxifolia.
corylifolia, Willd. ex Roem. & Schult. Syst. iv. 461 = alba.
corymbosa, G. Don, Gen. Syst. iv. 383 = monosperma.
corymbosa, Willd. ex Steud. Nom. ed. II. i. 417 = alba.
crenata, *Delile, Fl. Egypte*, 195. t. 20.—Aegypt.
crenata, Roem. & Schult. Syst. iv. 463 = lantanoides.
crenulata, *A. DC. in DC. Prod.* ix. 492.—Mexic.
crispiflora, *DC. l. c.* 476.—Mexic.
cujabensis, *Manso & Lhotzky, ex Cham. in Linnaea*, viii. (1833) 12.—Bras.
Cumingiana, *Vidal, Phan. Cuming. Philipp.* 187.—Ins. Philipp.
cuneata, Heyne, ex DC. Prod. ix. 480 = Rothii.
cuneiformis, *DC. l. c.* 492.—Venezuela.
curassavica, *Roem. & Schult. Syst.* iv. 460.—Am. austr.
cylindristachya, *Roem. & Schult. l. c.* 459.—Am. trop.
dasycephala, *H. B. & K. Nov. Gen. et Sp.* iii. 76.—Mexic.
decandra, *Hook. & Arn. Bot. Beech. Voy.* i. 38.—Chili.
dentata, Vahl, Eclog. Am. iii. 5 = alba.
dichotoma, Forst. f. Prod. 18. n. 110 = C. Myxa.
digynia, *Vell. Fl. Flum.* 97; ii. t. 153.—Bras.
Dillenii, *Spreng. Syst.* i. 649.—Ins. Baham.
dioica, *Boj. ex DC. Prod.* ix. 481.—Madag.
dioica, Thonn. ex DC. l. c. 480 = guineensis.
diospyrifolia, *Cham. in Linnaea*, iv. (1829) 477.—Bras.
discolor, *Cham. l. c.* 482.—Bras.
divaricata, *H. B. & K. Nov. Gen. et Sp.* iii. 74.—Am. austr.
diversa, Thw. Enum. Pl. Zeyl. 214 = monoica.
diversifolia, *Pav. ex DC. Prod.* ix. 474.—Mexic.
dodecandra, *DC. l. c.* 478.—Mexic.
domestica, Bouton, ex Boj. Hort. Maurit. 235 = amplifolia.
domestica, Roth, Nov. Sp. Pl. 123 = C. Myxa.
domingensis, *Lam. Illustr.* n. 1900.—Ins. S. Doming.
dubiosa, *Blume, Bijdr.* 844.—Java.
ecalyculata, *Vell. Fl. Flum.* 96; ii. t. 149.—Bras.
egyptiaca, *Rafin. Sylva Tellur.* 37.—Aegypt.
ehretioides, Lam. Illustr. n. 1902 = C. Collococca vel nitida.
elaeagnoides, *DC. Prod.* ix. 474.—Mexic.
elliptica, Sieber, ex Griseb. Fl. Brit. W. Ind. 479 = laevigata.
elliptica, *Sw. Prod. Veg. Ind. Occ.* 47.—Ind. occ.
erythrococca, *Griseb. Cat. Pl. Cub.* 208.—Cuba.
exaltata, *Lam. Illustr.* n. 1919.—Guiana.
excelsa, *A. DC. in DC. Prod.* ix. 473.—Am. trop.
ferruginea, *Roem. & Schult. Syst.* iv. 458; *H. B. & K. Nov. Gen. et Sp.* iii. 71.—Mexic.; N. Granat.
Finlaysoniana, *Wall. Cat.* n. 7010.—Ind. or.
flavescens, *Aubl. Pl. Gui.* i. 226. t. 89.—Guiana.
floribunda, *Spreng. Syst.* i. 652.—Am. austr.
floridana, Nutt. Sylva, iii. 107 = Bourreria havanensis.
foliosa, *Mart. & Gal. in Bull. Acad. Brux.* xi. II. (1844) 330.—Mexic.
formicarum, Roem. & Schult. Syst. iv. 800 = nodosa.
fragrantissima, *Kurz, For. Fl. Brit. Burma*, ii. 207.—Ind. or.; Burma.
Francisci, *Tenore, Cat. Orto Nap.* (1845) 83.—Hab.?
frondosa, Schott, in Spreng. Syst. iv. App. 403 = alliodora.
fulvosa, *Wight, Ic.* iv. t. 1380.—Ind. or
Galeottiana, *A. Rich. Fl. Cub. Fanerog.* ii. 109.—Cuba.
Geraseanthus, H. B. & K. Nov. Gen. et Sp. iii. 69 = tinifolia.
Gerascanthus, *Linn. Syst.* ed. X. 936.—Mexic.
Gerascanthus, Sw. ex Griseb. Fl. Brit. W. Ind. 478 = geraschcanthoides.
geraschanthoides, *H. B. & K. Nov. Gen. et Sp.* iii. 69.—Cuba.
geraschanthoides, Rich. ex Griseb. Fl. Brit. W. Ind. 478 = A. Gerascanthus.

CORDIA :—
Geraschanthus, Jacq. Select. Am. 43. t. 175 = A. Geraschanthus.
glabra, *Cham. in Linnaea*, viii. (1833) 124.—Bras.
glabra, Linn. Sp. Pl. 191 = C. Collococca.
glabrata, *A. DC. Prod.* ix. 473.—Am. trop.
glandulosa, *Fresen. in Mart. Fl. Bras.* viii. 19.—Am. trop.
globosa, Andr. ex DC. Prod. ix. 497 = oaxacana.
globosa, *H. B. & K. Nov. Gen. et Sp.* iii. 76.—Ind. occ.; Am. centr.
glomerata, *Lem. Jard. Fleur.* iv. (1854) t. 326.—Bras.
grandiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 77.—Am. austr.
grandifolia, *DC. Prod.* ix. 475.—Am. trop.
grandis, Cham. in Linnaea, iv. (1829) 473 = grandifolia.
grandis, *Roxb. Hort. Beng.* 17; *Fl. Ind.* i. 593.—Ind. or.
grandis, Wall. Cat. n. 892 B = obliqua.
graveolens, *H. B. & K. Nov. Gen. et Sp.* iii. 74.—Am. austr.
Greggii, *Torr. in Bot. Mex. Bound.* 135.—Calif.; Mexic.
Griffithii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 139.—Malaya.
guayaquilensis, *DC. Prod.* ix. 496.—Ecuador.
guazumaefolia, Roem. & Schult. Syst. iv. 463 = urticifolia.
guianensis, *Roem. & Schult. l. c.* 460.—Guiana.
guineensis, *Thonn. in Schum. Beskr. Guin. Pl.* 128.—Afr. trop.
Hartwissiana, *Regel, Gartenfl.* (1857) 342.—Mexic.
hebecarpa, *DC. Prod.* ix. 488.—Am. trop.
hermanniaefolia, *Cham. in Linnaea*, iv. (1829) 484.—Am. trop.
heterophylla, *Poir. in Dict. Sc. Nat.* x. 409; *Roem. & Schult. Syst.* iv. 800.—Guiana.
hexandra, Willd. ex Roem. & Schult. l. c. 799 = subcordata.
hirsuta, Willd. Sp. Pl. i. 1076 = nodosa.
hispida, *Benth. Bot. Voy. Sulph.* 139.—Mexic.
hispidissima, *DC. Prod.* ix. 475.—Am. trop.
Humboldtii, Willd. ex Roem. & Schult. Syst. iv. 801 = floribunda.
hypoleuca, *DC. Prod.* ix. 472.—Am. trop.
ignota, Blanco, Fl. Filip. ed. II. 88 = subcordata.
incana, Royle, Illustr. Bot. Himal. 306, nomen = vestita.
indica, Lam. Encyc. vii. 49.; Illustr. n. 1913 = C. Myxa.
insignis, *Cham. in Linnaea*, viii. (1833) 122.—Bras.
integrifolia, *Roem. & Schult. Syst.* iv. 461.—Ins. S. Doming.
intermedia, *Fresen. in Mart. Fl. Bras.* viii. 8.—Am. trop.
interrupta, *DC. Prod.* ix. 491.—Guiana.
intricata, *Wright, in Sauv. Fl. Cub.* 109.—Cuba.
ipomoeaefolia, *Hook. Bot. Mag.* t. 5027.—Hab. ?
ixiocarpa, F. Muell. Fragm. i. 59 = C. Myxa.
juglandifolia, *Jacq. Enum. Pl. Carib.* 14.—Ins. S. Doming.
lacerata, F. Muell. Fragm. v. 193 = aspera.
laevigata, *Lam. Illustr.* n. 1912.—Ind. occ.
laevis, *Jacq. Hort. Schoenb.* l. t. 40.—N. Granat.
lanata, H. B. & K. Nov. Gen. et Sp. iii. 72 = tomentosa.
lanceolata, *H. B. & K. l. c.* 75.—N. Granat.
lantanoides, *Spreng. Syst.* i. 658.—Peruv.
lapensis, *Warm. in Kjoeb. Vidensk. Meddel.* (1867) 9.—Bras.
latifolia, *Cham. in Linnaea*, viii. (1833) 126.—Bras.
latifolia, Roxb. Hort. Beng. 17; Fl. Ind. i. 588 = obliqua.
latifolia, Wall. Cat. n. 893 = C. Myxa.
Latora, *Buch.-Ham. ex Royle, Illustr. Bot. Himal.* 306.—Ind. or.
laxiflora, *H. B. & K. Nov. Gen. et Sp.* iii. 72.—N. Granat.
Leschenaultii, DC. Prod. ix. 482 = oblongifolia.
leucocalyx, *Fresen. in Mart. Fl. Bras.* viii. 22.—Am. trop.
leucocephala, *Moric. Pl. Nouv. Am.* 148. t. 88.—Bras.
leucocoma, *Miq. Fl. Ind. Bat.* ii. 916.—Malaya.
leucophlyctis, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 199.—Ins. Galap.
leucosebestera, *Griseb. Cat. Pl. Cub.* 208.—Cuba.
Lima, *Roem. & Schult. Syst.* iv. 475.—Ins. S. Doming.
linearis, *A. DC. in DC. Prod.* ix. 493.—Mexic.
linearis, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 199.—Ins. Galap.

CORDIA :—
lineata, Roem. & Schult. Syst. iv. 464 = ulmifolia.
longifolia, *A. DC. in DC. Prod.* ix. 495.—Am. trop.
Loureiri, *Roem. & Schult. Syst.* iv. 465.—China.
lutea, *Lam. Tabl. Encyc.* i. 421.—Peruv.
Macleodii, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 128.—Ind. or.
macrocephala, *H. B. & K. Nov. Gen. et Sp.* iii. 77.—Peruv.
macrophylla, *Linn. Sp. Pl.* ed. II. 273.—Ind. occ.
macrophylla, Vahl, ex Griseb. Fl. Brit. W. Ind. 480 = sulcata.
macrostachya, *Roem. & Schult. Syst.* iv. 461.—N. Granat.
macrostachya, Spreng. Syst. i. 652 = cylindristachya.
magnoliaefolia, *Cham. in Linnaea*, iv. (1829) 476.—Bras.
mariquitensis, *H. B. & K. Nov. Gen. et Sp.* iii. 75.—N. Granat.
Martii, DC. Prod. ix. 471 = insignis.
martinicensis, Link, Enum. Hort. Berol. i. 162 = curassavica.
martinicensis, *Roem. & Schult. Syst.* iv. 461.—Ins. Martinic.
martinicensis, Sieber, ex Griseb. Fl. Brit. W. Ind. 480 = cylindristachya.
McLeodii, vide Macleodii.
melanoneura, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 960.—Guiana.
membranacea, *A. DC. ex DC. Prod.* ix. 484.—Ecuador.
micrantha, *Sw. Prod. Veg. Ind. Occ.* 47.—Jamaica.
micrantha, Sw. l. c. pro parte, ex Griseb. Fl. Brit. W. Ind. 479 = C. Collococca.
microcephala, *Willd. in Roem. & Schult. Syst.* iv. 801.—Venezuela.
microphylla, *Roem. & Schult. l. c.* 463.—Ins. S. Doming.
mirabiliflora, *A. DC. in DC. Prod.* ix. 499.—Cuba.
mirabiloides, *Roem. & Schult. Syst.* iv. 465.—Ins. S. Doming.
miranda, *DC. Prod.* ix. 475.—Am. trop.
moluccana, *Roxb. Hort. Beng.* 84; *Fl. Ind.* ii. 594.—Ins. Molucc.
monoica, Boj. Hort. Maurit. 235 = ovalis.
monoica, *Roxb. Pl. Corom.* i. 43. t. 58.—Ind. or.
monosperma, *Roem. & Schult. Syst.* iv. 463.—Venezuela.
mucronata, *Fresen. in Mart. Fl. Bras.* viii. 9.—Am. trop.
multispicata, *Cham. in Linnaea*, iv. (1829) 499.—Bras.
Munneco, *H. B. & K. Nov. Gen. et Sp.* vii. 207.—Venezuela.
Myxa, Forsk. Fl. Aegypt. Arab. p. lxiii = crenata.
Myxa, *Linn. Sp. Pl.* 190.—As. trop.; Austral.
Myxa, Thw. Enum. Pl. Zeyl. 213 = obliqua.
Neowediana, *A. DC. in DC. Prod.* ix. 498.—Am. trop.
nervosa, *Lam. Illustr.* n. 1906.—Guiana.
nitida, *Vahl, in West, Bidrag Ste-Croix*, 275.—Ind. occ.
nitida, *Willd. in Roem. & Schult. Syst.* iv. 800.—Hab. ?
nivea, *Fresen. in Mart. Fl. Bras.* viii. 26.—Am. trop.
nodosa, *Lam. Illustr.* n. 1905.—Guiana.
oaxacana, *DC. Prod.* ix. 497.—Mexic.
obliqua, Heyne, ex DC. l. c. 479 = Wallichii.
obliqua, H. B. & K. Nov. Gen. et Sp. iii. 74 = peruviana.
obliqua, *Vell. Fl. Flum.* 97; ii. t. 150.—Bras.
obliqua, *Willd. Sp. Pl.* i. 1072.—Ind. or.
oblongifolia, *Hochst. ex DC. Prod.* ix. 480, nota.—Abyssin.
oblongifolia, *Thw. Enum. Pl. Zeyl.* 214.—Zeylan.
obovata, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 80.—Ins. Socotra.
obscura, *Cham. in Linnaea*, xiii. (1839) 480.—Bras.
obtusa, *Balf. f. in Proc. Roy. Soc. Edinb.* xii. (1884) 80.—Ins. Socotra.
ochnacea, *DC. Prod.* ix. 477.—Am. trop.
octandra, *A. DC. l. c.*—Ind. or.
officinalis, Lam. Illustr. ii. 420 = C. Myxa.
olitoria, *Blanco, Fl. Filip.* ed. I. 123.—Ins. Philipp.
oncocalyx, *F. Allem. in Trab. Comm. Sc. Expl. Bot.* i. (1862) 11.—Bras.
orientalis, R. Br. Prod. 498 = subcordata.
ovalis, *R. Br. in Salt, Abyss. App.* 65.—Abyssin.
oxyphylla, *DC. Prod.* ix. 492.—Guiana.
panicularis, *Rudge, Pl. Guian.* 30. t. 46.—Guiana.
paniculata, Roth, Nov. Pl. Sp. 125 = C. Myxa.

CORDIA :—
paniculata, Sieber, ex Presl, in Oken, Isis, xxi. (1828) 273 = Wilbrandia paniculata.
parviflora, Orteg. Hort. Matr. 85 = serratifolia.
parvifolia, *A. DC. in DC. Prod.* ix. 498.—Mexic.
parvifolia, Hemsl. Biol. Centr. Am. Bot. ii. 369, in syn. = serratifolia.
Patagonula, Ait. Hort. Kew. i. 259 = Patagonula americana.
patens, *H. B. & K. Nov. Gen. et Sp.* iii. 75.—Am. austr.
patens, Miq. ex Griseb. Fl. Brit. W. Ind. 480 = ulmifolia.
paucidentata, *Fresen. in Mart. Fl. Bras.* viii. 25.—Bras.
pedunculosa, *Griseb. Cat. Pl. Cub.* 209.—Cuba.
Pelida, Perr. ex DC. Prod. ix. 480 = senegalensis.
pellita, Willd. ex Roem. & Schult. Syst. iv. 800 = Tournefortia hirsutissima.
Perrottetii, *A. DC. Prod.* ix. 482.—Ind. or.
peruviana, *Roem. & Schult. Syst.* iv. 459.—Peruv.
petiolata, Hort. ex Steud. Nom. ed. II. i. 418 = Ehretia petiolaris.
Petta-pelioporet, Heyne, ex Roth, Nov. Pl. Sp. 125 = paniculata.
piauhiensis, *Fresen. in Mart. Fl. Bras.* viii. 9.—Bras.
pilulifera, Willd. ex Roem. & Schult. Syst. iv. 801 = globosa.
platyphylla, Steud. Nom. ed. II. i. 418 = latifolia.
podocephala, *Torr. in Bot. Mex. Bound.* 135.—Texas; Mexic.
Poeppigii, *DC. Prod.* ix. 492.—Am. austr.
poliophylla, *Fresen. in Mart. Fl. Bras.* viii. 26.—Am. trop.
polyantha, *Benth. Pl. Hartw.* 121.—Ecuador.
polygama, Roxb. Hort. Beng. 17; Fl. Ind. i. 594 = monoica.
polygama, Wall. Cat. n. 892, partim = fulvosa.
polystachya, *H. B. & K. Nov. Gen. et Sp.* iii. 73.—Venezuela.
portoricensis, Spreng. Nene Entdeck. ii. 127 = cylindristachya.
prionodes, Voigt, Hort. Suburb. Calc. 441 = octandra.
pubescens, *Willd. ex Roem. & Schult. Syst.* iv. 803.—Venezuela.
pyrifolia, Willd. ex Roem. & Schult. l. c. 802 = ferruginea.
quercifolia, *Klotzsch, in Peters, Reise Mossamb. Bot.* 247.—Afr. trop.
Radula, Spreng. Syst. i. 654 = Lima.
rariflora, *A. Rich. Fl. Cub. Fanerog.* ii. 110.—Cuba.
reflexa, *Willd. ex Roem. & Schult. Syst.* iv. 800.—Ind. or.
reticulata, Roth, Nov. Pl. Sp. 124 = Rothii.
reticulata, *Vahl, Eclog. Am.* iii. 5.—Ins. Montserrat.
retusa, Vahl, Symb. Bot. ii. 42 = Ehretia buxifolia.
revoluta, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 199.—Ins. Galap.
riparia, *H. B. & K. Nov. Gen. et Sp.* iii. 71.—N. Granat.
rotata, *Moç. ex DC. Prod.* ix. 483.—Mexic.
Rothii, *Roem. & Schult. Syst.* iv. 798.—Ind. or.
rotundifolia, Heyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 139 = fragrantissima.
rotundifolia, *Ruiz & Pav. Fl. Per.* ii. 24. t. 148.—Panama.
Roxburghii, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 138.—Zeylan.
rubra, Hochst. ex A. Rich. Tent. Fl. Abyss. ii. 82 = C. Myxa.
rufa, *Klotzsch, in Schomb. Fauna et Fl. Gui.* 960.—Guiana.
rufescens, *A. DC. in DC. Prod.* ix. 476.—Am. trop.
rugosa, Willd. ex Roem. & Schult. Syst. iv. 801 = canescens.
Rumphii, Blume, Bijdr. 843 = subcordata.
salicifolia, *Cham. in Linnaea*, iv. (1829) 481.—Bras.
salicina, *DC. Prod.* ix. 492.—Am. trop.
salviaefolia, H. B. & K. Nov. Gen. et Sp. iii. 72 = cylindristachya.
salvifolia, *Juss. in Poir. Encyc.* vii. 46.—Ins. Guadalup.
Salzmanni, [Benth. ex] Griseb. Fl. Brit. W. Ind. 480 = ulmifolia.
Salzmanni, *DC. Prod.* ix. 494.—Bras.
sarmentosa, Lam. Illustr. n. 1907 = flavescens.
scaberrima, *H. B. & K. Nov. Gen. et Sp.* iii. 71.—Peruv.

CORDIA:—
scabra, *Desf. Tabl.* ed. I. 71.—Ind. or.
scabrida, *Mart. ex Fresen. in Mart. Fl. Bras.* viii. 11.—Bras.
scabrifolia, [Benth. ex] Griseb. Fl. Brit. W. Ind. 479 = paniculata.
scabrifolia, *A. DC. in DC. Prod.* ix. 485.—Guiana.
scandens, *Martin, ex Poir. in Dict. Sc. Nat.* x. 410.—Guiana.
Schomburgkii, *DC. Prod.* ix. 490.—Guiana.
Schottiana, *Fresen. in Mart. Fl. Bras.* viii. 7.—Am. trop.
Scouleri, *Hook. f. in Trans. Linn. Soc.* xx. (1851) 200.—Ins. Galap.
Sebestena, Andr. Bot. Rep. iii. t. 157 = scabra.
Sebestena, Forsk. Fl. Aegypt. Arab. p. lxiii = C. Myxa.
Sebestena, Forst. f. Prod. 18 = subcordata.
Sebestena, *Linn. Sp. Pl.* 190.—Am. trop.
Sebestena, Willd. ex Spreng. Syst. i. 649 = Dillenii.
Sellowiana, *Cham. in Linnaea*, iv. (1829) 479.—Bras.
senegalensis, *Juss. in Poir. Encyc.* vii. 47.—Afr. occ.
sericicalyx, *A. DC. in DC. Prod.* ix. 485.—Guiana.
serrata, *Juss. ex Poir. Encyc.* vii. 41.—Ind. or.
serrata, Roxb. Hort. Beng. 17; Fl. Ind. i. 591 = octandra.
serratifolia, *H. B. & K. Nov. Gen. et Sp.* iii. 76.—Mexic.
sessilifolia, *Cham. in Linnaea*, iv. (1829) 488.—Bras.
sinensis, *Lam. Illustr.* n. 1914.—China.
speciosa, Salisb. Prod. 111 = C. Sebestena.
spectabilis, *Steud. Nom.* ed. II. i. 419.—Ind. or.
sphaerocephala, Willd. ex Roem. & Schult. Syst. iv. 801 = dasycephala.
spicata, Willd. ex Roem. & Schult. l. c. 799 = graveolens.
spinescens, *Linn. Mant.* 206.—Ind. or.
Sprengelii, *DC. Prod.* iv. 473.—Ins. Fiji.
striata, *Fresen. in Mart. Fl. Bras.* viii. 23.—Am. trop.
strigosa, *Spreng. Neue Entdeck.* iii. 30.—Ind. occ.
suaveolens, *Blume, Bijdr.* 843.—Java.
subcordata, *Lam. Illustr.* n. 1899.—As. et Austral. trop.
subdentata, *Miq. Fl. Ind. Bat. Suppl.* 571.—Sumatra.
subopposita, DC. Prod. ix. 480 = Rothii.
subpubescens, *Decne. Herb. Timor.* 67.—Ins. Timor.
sulcata, *DC. Prod.* ix. 488.—Ind. or.
superba, *Cham. in Linnaea*, iv. (1829) 474.—Bras.
taguahyensis, *Vell. Fl. Flum.* 98; ii. t. 154.—Bras.
tenuifolia, *Bertol. Misc. Bot.* xxi. 13. t. 1.—Guatemala.
tetrandra, *Aubl. Pl. Gui.* i. 222. t. 87.—Guiana.
tetraphylla, *Aubl. l. c.* 224. t. 88.—Guiana.
Thibaudiana, *DC. Prod.* ix. 489.—Am. austr.
thyrsiflora, Sieb. & Zucc. Fl. Jap. Fam. Nat. ii. 26 = Ehretia acuminata.
timorensis, Span. in Linnaea, xv. (1841) 334 = subpubescens.
tinifolia, *Willd. ex Roem. & Schult. Syst.* iv. 800.—Am. austr.
tomentosa, Cham. & Schlecht. in Linnaea, iv. (1829) 472 = Chamissoniana.
tomentosa, *Roem. & Schult. Syst.* iv. 459.—Am. austr.
tomentosa, Wall. in Roxb. Fl. Ind. ed Carey, ii. 339 = obliqua.
Toqueve, *Aubl. Pl. Gui.* i. 228. t. 90.—Am. trop.
Toqueve, Sieber, ex Griseb. Fl. Brit. W. Ind. 480 = sulcata.
trachyphylla, *Mart. Herb. Fl. Bras.* n. 412.—Bras.
tremula, *Griseb. Fl. Brit. W. Ind.* 479.—Ind. occ.
trichoclada, *DC. Prod.* ix. 474.—Am. trop.
trichostemon, *DC. l. c.* 482.—Ins. Timor.
trichotoma, *Vell. ex Steud. Nom.* ed. II. i. 419.—Bras.
truncata, *Fresen. in Mart. Fl. Bras.* viii. 25.—Am. trop.
ulmifolia, *Juss. in Dum.-Cours. Bot. Cult.* ed. I. ii. 148.—Panama; Ind. occ.
ulmifolia, Spreng. Syst. i. 653 = monosperma.
umbraculifera, *DC. Prod.* ix. 484.—Am. trop.
urticifolia, *Cham. in Linnaea*. iv. (1829) 483.—Bras.
Valenzuelana, *A. Rich. Fl. Cub. Fanerog.* ii. 110.—Cuba.
velutina, Mart. Herb. Fl. Bras. n. 268 = cujabensis.
verbenacea, *DC. Prod.* ix. 491.—Am. trop.
vestita, *Hook. f. & Thoms. in Journ. Linn. Soc.* ii. (1858) 128.—Reg. Himal.
viburnoides, *Zipp. ex Span. in Linnaea*, xv. (1841) 335, nomen.—Malaya.

CORDIA:—
villicaulis, *Fresen. in Mart. Fl. Bras.* viii. 24.—Am. trop.
villosa, *Spreng. Neue Entdeck.* iii. 31.—Ins. S. Doming.
Wallichii, G. Don, Gen. Syst. iv. 379 = obliqua.
Wickstroemii, *Steud. Nom.* ed. II. i. 419.—Ins. Guadalup.

CORDIADA, Vell. Fl. Flum. 98 (1825); ii. t. 156 = **Cordia**, Linn. (Boragin.).
trichotoma, Vell. ll. cc. = Cordia trichotoma.

CORDIERA, A. Rich. in Mém. Soc. Hist. Nat. Par. v. (1830) 222. t. 20 = **Alibertia**, A. Rich. (Rubiac.).
acuminata, Benth. in Hook. Journ. Bot. iii. (1841) 221.—Guiana.
latifolia, Benth. l. c.—Bras.
triflora, DC. Prod. iv. 445.—Guiana.
trifolia, Steud. Nom. ed. II. i. 419, sphalm. = praec.

CORDIOPSIS, Desv. in Ham. Prod. Fl. Ind. Occ. 23 (1825) = **Cordia**, Linn. (Boragin.).
mirabiliflora, DC. Prod. ix. 499 = Cordia mirabiliflora.
mirabiloides, Desv. in Ham. Prod. Fl. Ind. Occ. 23 = Cordia mirabiliflora.
monosperma, Ham. ex Heynh. Nom. ii. 160 = Cordia monosperma.

CORDULA, Rafin. Fl. Tellur. iv. 46 (1836) = **Cypripedium**, Linn. (Orchid.).
insignis, Rafin. l. c. = Cyprip. insigne.

CORDYLA, Blume, Bijdr. 416 (1825) = **Pogonia**, Juss. (Orchid.).
concolor, Blume, l. c. = P. concolor.
discolor, Blume, l. c. = P. discolor.

CORDYLA, Lour. Fl. Cochinch. 411 (1790). *LEGUMINOSAE*, Benth. & Hook. f. i. 562.
CALYCANDRA, Lepr. ex A. Rich. in Guill. & Perr. Fl. Seneg. Tent. 30; et 232. t. 9 (1832).
africana, *Lour. l. c.* 412.—Afr. trop.
Richardi, Planch. in App. Speke, Journ. Disc. Nile, 632 = africana.

CORDYLANTHUS, Blume, Mus. Bot. Lugd. Bat. ii. 27. t. 3. 1852 [1856] = **Homalium**, Jacq. (Samydac.).
frutescens, Blume, l. c. 28 = H. Cordylanthus.

CORDYLANTHUS, Nutt. ex Benth. in DC. Prod. x. 597 (1846). *SCROPHULARINEAE*, Benth. & Hook. f. ii. 974.
ADENOSTEGIA, Benth. in Lindl. Nat. Syst. ed. II. 445 (1836).
CHLOROPYRON, Behr, in Proc. Calif. Acad. i. (1855) 61.
canescens, *A. Gray, in Proc. Am. Acad.* vii. (1868) 383.—Am. bor. occ.
capitatus, *Nutt. ex Benth. in DC. Prod.* x. 597.—Am. bor occ.
filifolius, *Nutt. ex Benth. l. c.*—Calif.
Kingii, *S. Wats. in Bot. King's Exp.* 233, 459. t. 22.—Am. bor. occ.
laxiflorus, *A. Gray, in Torr. Bot. Mex. Bound.* 120.—Arizona.
maritimus, *Nutt. ex Benth. in DC. Prod.* x. 598.—Calif.
mollis, *A. Gray, in Proc. Am. Acad.* vii. (1868) 384.—Calif.
Nevinii, *A. Gray, l. c.* xvii. (1881–82) 229.—Calif.
Orcuttianus, *A. Gray, l. c.* xix. (1883) 95.—Calif.
Parryi, S. Wats. in Am. Natural. ix. (1875) 346 = canescens.
pilosus, *A. Gray, in Proc. Am. Acad.* vii. (1868) 382.—Calif.
Pringlei, *A. Gray, l. c.* xix. (1883) 94.—Calif.
ramosus, *Nutt. ex Benth. in DC. Prod.* x. 597.—Am. bor. occ.
tenuis, *A. Gray, in Proc. Am. Acad.* vii. (1868) 383.—Am. bor. occ.
Wrightii, *A. Gray, in Torr. Bot. Mex. Bound.* 120.—Texas; N. Mexic.

CORDYLESTYLIS, Falc. in Hook. Journ. Bot. iv. (1842) 74 = **Goodyera**, R. Br. (Orchid.).
foliosa, Falc. l. c. 75 = G. procera.

CORDYLIA, Pers. Syn. ii. 260 (1807) = **Coryda**, Lour. (Legumin.).

CORDYLINE, Royen, ex Adans. Fam. ii. 54 (1763). *LILIACEAE*, Benth. & Hook. f. iii. 799.
CALODRACON, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 137.
CHARLWOODIA, Sweet, Fl. Austral. t. 18 (1827).
COHNIA, Kunth, Enum. Pl. v. 35 (1850).
DRACAENOPSIS, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 110.
TAETSIA, Medic. Theodora, 82 (1786).
angustifolia, Kunth, Enum. Pl. v. 32 = stricta.
arborea, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55.—Afr. trop.
atropurpurea, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 132 = Dracaena elliptica.
australis, Endl. Prod. Fl. Ins. Norf. 29 = Baueri.
australis, *Hook. f. in Gard. Chron.* (1860) 792.—N. Zel.
australis, Nadeaud & Jouan, in Mém. Soc. Sc. Nat. Cherb. xi. (1865) 127 = terminalis.
Banksii, *Hook. f. in Gard. Chron.* (1860) 792.—N. Zel.
Baueri, *Hook. f. l. c.*—Ins. Norfolc.
Betschleriana, Goepp. in Nov. Act. Nat. Cur. xxv. (1885) 55 = Dracaena concinna.
Beuckelaerii, C. Koch, Wochenschr. viii. (1865) 91 = Banksii.
brasiliensis, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 136.—Bras.
calocoma, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 542 = australis.
cannaefolia, R. Br. Prod. 280 = terminalis.
Casanovae, *Linden, ex André, Illustr. Hortic.* xxi. (1874) 137. t. 181.—N. Hebrid.
cernua, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 111 = Dracaena reflexa.
coerulea, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55.—Hab. ?
congesta, Endl. Gen. 151 = stricta.
congesta, *Steud. Nom.* ed. II. i. 419.—Austral.
Dennisoni, *Hort. ex André, Illustr. Hortic.* xix. (1872) 360.—Hab. ?
densicoma, *Linden & André, l. c.* (1874) 185. t. 190.—Austral.
diffusa, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 330.—N. Zel.
dracaenoides, *Kunth, in Abh. Acad. Berl.* (1842) 30.—Bras.
ensifolia. Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 132 = Dianella ensifolia.
erythrorachis, Hort. ex Baker, in Journ. Linn. Soc. xiv. (1875) 541 = Banksii.
Eschscholziana, Mart. in Schult. f. Syst. vii. 347, 1677 = terminalis.
ferrea, C. Koch, Wochenschr. xiv. (1871) 12 = terminalis.
flabelliformis, *C. Koch, Allg. Gartenz.* (1858) 254.—Ins. Maurit.
flexuosa, *Miq. Fl. Ind. Bat.* iii. 558.—Malaya.
floribunda, *C. Koch, Allg. Gartenz.* (1858) 254.—Ins. Maurit.
Fontanesiana, Goepp. in Nov. Act. Nat. Cur. xxv. (1855) 54 t. 3 = Dracaena Fontanesiana.
Forsteri, F. Muell. Sel. Pl. Indust. Cult. 58 = australis.
fragrans, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 111, 136.—Afr. trop.
fruticosa, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 53.—Java.
gloriosa, Linden & Andre, Illustr. Hortic. xx. (1873) t. 125–6 = terminalis.
Guilfoylei, Hort. Linden, ex Lem. Illustr. Hortic. xvi. (1869) t. 600 terminalis.
Haageana, *C. Koch, Wochenschr.* (1867) 195.—Austral.
hedychioides, F. Muell. Fragm. v. 196 = terminalis.
heliconiaefolia. Otto & Dietr. in Allg. Gartenz. (1835) 32 = terminalis.
hemichrysa, *Comm. ex Lam. Encyc.* ii. 276.—Ins. Maurit.
Hookbrenkeana, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55.—N. Zel.
Hookeri, *T. Kirk, in Trans. N. Z. Inst.* vi. (1874) 215.—N. Zel.
humilis, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55.—Hab. ?

CORDYLINE:—
indivisa, Regel, Gartenfl. (1859) 331 = australis.
indivisa, *Steud. Nom.* ed. II. i. 419.—N. Zel.
Jacquinii, Kunth, in Abh. Acad. Berl. (1842) 30 = terminalis.
javanica, Klotzsch, ex Kunth, Enum. Pl. v. 26 = terminalis.
lentiginosa, Linden & André, Illustr. Hortic. xvii. (1870) t. 35 = australis.
linearifolia, *Miq. Fl. Ind. Bat. Suppl.* 610.—Sumatra.
longifolia, Benth. Pl. Hartw. 53 = Nolina Hartwegiana.
macrophylla, *C. Koch, Allg. Gartenz.* (1858) 254.—Ins. Mascar.
maculata, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 132 = Dracaena elliptica.
madagascariensis, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 53.—Madag.
Manners-Suttoniae, F. Muell. Fragm. v. 195 = terminalis.
marginata, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55.—Madag.; Ins. Borb.
metallica, Hort. Dallière, in Belg. Hortic. xxii. (1872) 353. t. 26 = terminalis.
micrantha, Baker, in Gard. Chron. (1880) II. 243 = Brocchinia cordylinoides.
Murchisoniae, F. Muell. Fragm. v. 195 = Haageana.
neo-caledonica, *Linden, Cat.* n. 97 (1877) 4.—N. Caled.
nobilis, C. Koch, Allg. Gartenz. (1858) 254 = terminalis.
nutans, *Hort. ex Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 55, nomen.—Hab.?
obtecta, Baker, in Journ. Linn. Soc. xiv. (1875) 543 = Baueri.
odorata, C. Koch, in C. Koch & Fint. Wochenschr. (1858) 97 = dracaenoides.
ovata, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 111, 136 = Dracaena ovata.
oxyphylla, *Miq. Fl. Ind. Bat. Suppl.* 609.—Sumatra.
parviflora, H. B. & K. Nov. Gen. et Sp. i. 269 = Nolina parviflora.
Pumilio, *Hook. f. in Gard. Chron.* (1860) 792.—N. Zel.
Reali, *Nichols. Dict. Gard.* i. 374.—Hab.?
reflexa, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 109.—Afr. trop.
Regelii, *C. Koch, Wochenschr.* viii. (1865) 90.—Hab.?
Regina, Veitch, ex Regel, Gartenfl. (1871) 149 = terminalis.
rigida, *Miq. Fl. Ind. Bat. Suppl.* 610.—Sumatra.
rigidifolia, C. Koch, in C. Koch & Fint. Wochenschr. (1858) 359 = stricta.
rubra, *Hueg. ex Kunth, Enum. Pl.* v. 34.—N. Zel.
Rumphii, *Hook. Bot. Mag.* t. 4297 = Dracaena angustifolia, Hookeriana.
Rumphii, F. Muell. Fragm. v. 194 = Dracaena angustifolia.
salicifolia, *Goepp. in Nov. Act. Nat. Cur.* xxv. (1855) 53.—Java.
Sellowiana, Kunth, in Abh. Acad. Berl. (1842) 30 = dracaenoides.
Sepiaria, Seem. Fl. Vit. 311. t. 94 = terminalis.
Sieberi, Kunth, in Abh. Acad. Berl. (1842) 30 = terminalis.
Sieboldii, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 109. sub t. 569 = Dracaena elliptica.
spectabilis, Kunth & Bouché, in Ind. Sem. Hort. Berol. (1848) = dracaenoides.
spicata, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 132.—Hab.?
stricta, *Endl. in Ann. Wien. Mus.* i. (1836) 162.—Austral.
stricta, Hook. f. in Fl. N. Zel. i. 257. t. 50 = Pumilio.
Sturmii, *Colenso, in Trans. N. Z. Inst.* xv. (1883) 331.—N. Zel.
superbiens, C. Koch, in C. Koch & Fint. Wochenschr. (1859) 381 = australis.
terminalis, *Kunth, in Abh. Acad. Berl.* (1820) 30.—As. trop.; Austral.
Ti, Schott, in Flora, xi. (1828) 575 = terminalis.
timorensis, *Planch. in Fl. des Serres*, Sér. I. vi. (1850–51) 132.—Ins. Timor.
umbraculifera, Planch. l. c. 111 = Dracaena umbraculifera.

CORDYLINE:—
Veitchii, Regel, Gartenfl. (1871) 149 = australis.
violascens, Regel, Ind. Sem. Hort. Petrop. (1858) 25; et Gartenfl. (1859) 331 = rubra.
vivipara, Hort. ex Steud. Nom. ed. II. i. 419 = Chlorophytum Sternbergianum.
Wallichii, Planch. in Fl. des Serres, Sér. I. vi. (1850–51) 132 (Dracaenae sp.).—Ind. or.

CORDYLOBLASTE, Hensch. ex Moritzi, in Bot. Zeit. vi. (1848) 604 = **Symplocos**, Linn. (Styrac.).
Henschelii, Moritzi, l. c. = S. Henscheli.

CORDYLOCARPUS, Desf. Fl. Atlant. ii. 79. t. 152 (1800) = **Rapistrum**, Tourn. (Crucif.).
crassifolius, Steud. Nom. ed. I. 226 = Erucaria crassifolia.
glaber, Dietr. ex Steud. l. c. = R. glabrum.
laevigatus, Willd. Sp. Pl. iii. 563 = Erucaria aleppica.
muricatus, Desf. Fl. Atlant. ii. 79. t. 152.—Algeria.
pubescens, Sibth. & Sm. Fl. Graec. Prod. ii. 33 = Brassica adpressa.
ramosissimus, Pomel, Nouv. Mat. Fl. Atl. 357.—Algeria.
tenuifolius, Sibth. & Sm. Fl. Graec. Prod. ii. 33 = Erucaria tenuifolia.

CORDYLOGNE, Lindl. Veg. Kingd. 626 (1848), sphalm. = seq.

CORDYLOGYNE, E. Mey. Comm. Pl. Afr. Austr. 218 (1837). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 759.
PERIGLOSSUM, Decne. in DC. Prod. viii. 520 (1844).
globosa, *E. Mey. Comm. Pl. Afr. Austr.* 218.—Afr. austr.

COREMA, D. Don, in Edinb. N. Phil. Journ. ii. (1826–27) 63. *EMPETRACEAE*, Benth. & Hook. f. iii. 414.
ENDAMMIA, Rafin. New Fl. Am. iii. 49 (1836).
EULEUCUM, Rafin. Fl. Tellur. iii. 56 (1836).
OAKESIA, Tuckerm. in Hook. Lond. Journ. Bot. i. (1842) 445.
TUCKERMANNIA, Klotzsch, in Wiegm. Archiv, vii. (1841) 248.
album, *D. Don, in Edinb. N. Phil. Journ.* ii. (1827) 63.—Lusitan.
Conradi, *Torr. ex Loud. Encyc. Trees*, 1092.—Am. bor.
febrifugum, Boiss. ex Willk. & Lange, Prod. Fl. Hisp. iii. 512 = album.

COREOCARPUS, Benth. Bot. Voy. Sulph. 28. t. 16 (1844). *COMPOSITAE*, Benth. & Hook. f. ii. 384.
ACOMA, Benth. Bot. Voy. Sulph. 29. t. 17 (1844).
heterocarpus, *A. Gray, in Proc. Am. Acad.* v. (1861) 162.—Calif.
parthenioides, *Benth. Bot. Voy. Sulph.* 28. t. 16.—Calif.

COREOPIS, Gunn. Fl. Norveg. ii. 87 (1772), sphalm. = seq.

COREOPSIS, Linn. Gen. ed. I. 263 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 385.
ACISPERMUM, Neck. Elem. i. 34 (1790).
AGARISTA, DC Prod. v. 569 (1836).
ANACIS, Schrank, in Denkschr. Acad. Muench. v. (1817) 5.
CALLIOPSIS, Reichb. Ic. et Descr. Pl. t. 10 (1824).
CAMPYLOTHECA, Cass. Dict. Sc. Nat. li. 476 (1827).
CHRYSOMELEA, Tausch, Hort. Canal. fasc. i. (1823).
CHRYSOSTEMMA, Less. Syn. Comp. 227 (1832).
COREOPSOIDES, Moench, Meth. 594 (1794).
DIATONTA, Walp. Rep. ii. 614 (1843).
DIODONTA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 360.
DIPLOSASTERA, Tausch, Hort. Canal. fasc. i. t. 4 (1823).
DOLICHOTHECA, Cass. Dict. Sc. Nat. li. 476 (1827).
ELECTRA, DC. Prod. v. 630 (1836).
EPILEPIS, Benth. Pl. Hartw. 17 (1839).

COREOPSIS:—
HETERODONTA, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 361.
LEACHIA, Cass. in Dict. Sc. Nat. xxv. 388 (1822).
LERCHIA, Reichb. Consp. 109 (1828).
PERAMIBUS, Rafin. ex DC. Prod. v. 568 (1836).
PRESTINARIA, Sch. Bip. ex Steud. Nom. ed. II. ii. 393 (1841).
abyssinica, *Sch. Bip. ex Walp. Rep.* vi. 163.—Abyssin.
acuta, Pursh, Fl. Am. Sept. ii. 569 = Actinomeris squarrosa.
alata, Cav. Ic. iii. 30. t. 260 = Actinomeris tetraptera.
alata, Pursh, Fl. Am. Sept. ii. 567 = Verbesina occidentalis.
alba, Linn. Sp. Pl. 908 = Bidens pilosa.
alternifolia, Linn. l. c. 909 = Actinomeris squarrosa.
ambigua, Nutt. in Journ. Acad. Philad. vii. (1834) 75 = aurea.
amplexicaulis, Cav. Desc. 226 = Encelia amplexicaulis.
angustifolia, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 253.—Am. bor.
angustifolia, Linn. Sp. Pl. 908 = Helianthus angustifolius.
angustifolia, Pav. ex DC. Prod. v. 603 = Bidens procera.
anthemoides, *DC. Prod.* v. 573.—Mexic.
arguta, Pursh, Fl. Am. Sept. ii. 517 = aurea.
aristata, Willd. Sp. Pl. iii. 2253 = aristosa.
aristosa, *Michx. Fl. Bor. Am.* ii. 140.—Am. bor.
artemisiaefolia, Jacq. Coll. Suppl. 155 = Cosmus sulphureus.
aspera, *Pursh, Fl. Am. Sept.* ii. 570.—Am. bor.
Atkinsoniana, *Dougl. in Lindl. Bot. Reg.* t. 1376.—Am. bor. occ.
aurea, [*Dryand. in*] *Ait. Hort. Kew.* ed. I. iii. 252.—Am. bor.
aurea, Lindl. Bot. Reg. t. 1228 = trichosperma.
auriculata, *Linn. Sp. Pl.* 908.—Am. bor.
auriculata, Schkuhr, Handb. t. 260 = pubescens.
Bacana, Heyne, ex DC. Prod. v. 632 = Glossogyne pinnatifida.
baccata, Linn. Pl. Surin. n. 111; Amoen. Acad. viii. 262 = Wulffia stenoglossa.
baccata, *Linn. f. Suppl.* 380.—Guiana.
Barteri, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
bicolor, *Bosse, ex Buchenau, in Linnaea*, xxv. (1852) 630.—Hab.?
Bidens, Linn. Sp. Pl. 908 = Bidens cernua.
Bidens, Walt. Fl. Carol. 215 = Bidens chrysanthemoides.
bidentoides, *Nutt. ex Torr. & Gray, Fl. N. Am.* ii. 339.—Am. bor.
biternata, *Lour. Fl. Cochinch.* 508.—China.
bituminosa, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 11.—Malaya.
Borianiana, *Sch. Bip. ex Schweinf. in Verh. Zool.-Bot. Ges. Wien*, xviii. (1868) 684.—Afr. trop.
Boykiniana, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 358 = grandiflora.
calliopsidea, A. Gray, in Torr. Bot. Mex. Bound. 90 = Leptosyne calliopsidea.
callosa, *Bertol. Misc. Bot.* vii. 42.—Am. bor.
capillacea, *H. B. & K. Nov. Gen. et Sp.* iv. 230.—Peruv.
cardaminefolia, *Torr. & Gray, Fl. N. Am.* ii. 346.—Am. bor. occ.
chrysantha, Linn. Sp. Pl. ed. II. 1282 = Bidens chrysantha.
chrysantha, Spreng. Syst. iii. 614 = Bidens rubifolia.
chrysantha, *Vatke, in Linnaea*, xxxix. (1875) 499.—Afr. trop.
coronata, Bert. ex Steud. Nom. ed. II. i. 420 = Cosmus caudatus.
coronata, *Hook. Bot. Mag.* t. 3460.—Texas.
coronata, Linn. Sp. Pl. ed. II. 1282 (ex syn. Plum.) = Bidens pilosa.
coronata, Walt. Fl. Carol. ii. 15 = aurea.
corymbifolia, Buch.-Ham. ex DC. Prod. v. 602 = Bidens pilosa.
cosmoides, *A. Gray, in Proc. Am. Acad.* v. (1861) 126.—Ins. Sandvic.
crassifolia, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 252 = lanceolata.
cuspidata, *Bertol. Misc. Bot.* vii. 44.—Am. bor.
delphinifolia, *Lam. Encyc.* ii. 108.—Am. bor.
dichotoma, Michx. Fl. Bor. Am. ii. 137 = gladiata.

COREOPSIS:—
discoidea, *Torr. & Gray, Fl. N. Am.* ii. 339.—Am. bor.
discolor, *Link, Enum. Hort. Berol.* ii. 353.—Am. bor.
diversifolia, DC. Prod. v. 571. (excl. syn.) = auriculata.
diversifolia, Hook. Bot. Mag. t. 3474 = Drummondii.
diversifolia, Jacq. Eclog. 80. t. 54 = Bidens grandiflora.
Drummondii, *Torr. & Gray, Fl. N. Am.* ii. 345.—Texas.
emarcida, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. IV. 11.—Malaya.
fasciculata, *Wedd. Chlor. And.* i. 71.—Peruv.
ferulaefolia, Jacq. Hort. Schoenb. iii. 65 = Bidens ferulaefolia.
filifolia, Hook. Bot. Mag. t. 3505 = Thelesperma filifolium.
Flammula, Banks, ex Steud. Nom. ed. I. 226, 108 = Bidens chrysanthemoides.
flexicaulis, *Rafin. in Med. Repos. N. York*, v. (1808) 361.—Am. bor.
foeniculacea, Moç. & Sesse, ex DC. Prod. v. 603 = Bidens foeniculifolia.
foetida, Cav. Ic. i. 55 = Encelia foetida.
foliosa, *A. Gray, in Proc. Am. Acad.* v. (1861) 125.—Peruv.
formosa, *Bonato, Memoria* 2; *ex Pritz. Ic. Ind.* i. 301.—Hab. ?
formosa, Hort. ex Steud. Nom. ed. II. i. 420 = Cosmos bipinnatus.
fruticosa, *Forst. f. Prod.* 91.—Ins. Sandvic.
fruticosa, *Soland. ex Seem. Fl. Vit.* 143.—Ins. Pacif.
fruticosa, Vest, in Flora, iii. (1820) 7 = Bidens fruticosa.
Galeottii, *Hemsl. Biol. Centr. Am. Bot.* ii. 195.—Mexic.
Georgina, Cass. in Dict. Sc. Nat. xviii. 441 = Dahlia variabilis.
gladiata, *Walt. Fl. Carol.* 215.—Am. bor.
glaucescens, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 389.—Afr. trop.
gracilis, Blanco, Fl. Filip. ed. II. Suppl. 591 = Cosmos caudatus.
grandiflora, *Nutt. ex Chapm. Fl. S. U. St.* 335.—Am. bor.
Grantii, *Oliver, in Trans. Linn. Soc.* xxix. (1874) 98. t. 65.—Afr. trop.
guineensis, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
Harveyana, *A. Gray, Syn. Fl. N. Am.* i. II. 292.—Am. bor. occ.
heterophylla, *Bertol. Misc. Bot.* vii. 43.—Am. bor.
heterophylla, Cav. Ic. iii. 34. t. 268 = Iostephane heterophylla.
heterophylla, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 358 = grandiflora.
hirta, *Rafin. Ann. Nat.* 15.—Am. bor.
incisa, Ker-Gawl. in Bot. Reg. t. 7 = Bidens rubifolia.
incurva, Moench, Meth. Suppl. 245 = Bidens ferulaefolia.
integrifolia, *Poir. Encyc. Suppl.* ii. 352 (1811).—Am. bor.
involucrata, *Nutt. in Journ. Acad. Philad.* vii. (1834) 74.—Am. bor.
involucrata, *Sch. Bip. ex Walp. Rep.* vi. 163.—Abyss.
jasminifolia, *Bertol. Misc. Bot.* vii. 44.—Am. bor.
Kirkii, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
lanceolata, *Linn. Sp. Pl.* 908.—Am. bor.
latifolia, *Michx. Fl. Bor. Am.* ii. 137.—Am. bor.
Leavenworthii, *Torr. & Gray, Fl. N. Am.* ii. 346.—Am. bor.
leucantha, Linn. Sp. Pl. ed. II. 1282 = Bidens pilosa.
leucanthema, Linn. Cent. Pl. i. 29 = praec.
leucorrhiza, Lour. Fl. Cochinch. ii. 508 = Bidens leucorrhiza.
Liebmannii, *Sch. Bip. ex Hemsl. Biol. Centr. Am. Bot.* ii. 196.—Mexic.
limensis, Jacq. Coll. ii. 299 = Encelia canescens.
linearifolia, *DC. Prod.* v. 570.—Mexic.
linearifolia, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
linifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 75 = angustifolia.
longipes, Hook. Bot. Mag. t. 3586 = grandiflora.
lucida, Cav. Desc. 224 = Bidens heterophylla.
Macraei, *A. Gray, in Proc. Am. Acad.* v. (1861) 126.—Ins. Sandvic.
macrantha, *Sch. Bip. ex Walp. Rep.* vi. 163.—Abyss.

COREOPSIS:—
macrocarpa, *A. Gray, in Proc. Am. Acad.* v. (1861) 126.—Ins. Sandvic.
major, Walt. Fl. Carol. 214 = senifolia.
maritima, *Hook. f. Bot. Mag.* t. 6241.—Calif.
Menziesii, *A. Gray, in Proc. Am. Acad.* v. (1861) 127.—Ins. Sandvic.
mexicana, *Hemsl. Biol. Centr. Am. Bot.* ii. 196.—Mexic.
micrantha, *A. Gray, in Proc. Am. Acad.* v. (1861) 127.—Ins. Sandvic.
minima, *Hort. ex Steud. Nom.* ed. II. i. 420.—Am. bor.
mitis, Michx. Fl. Am. Bor. ii. 140 = aurea.
monticola, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 390.—Afr. trop.
multifida, *DC. Prod.* v. 573.—Peruv.
mutica, *DC. l. c.* 571.—Mexic.
Neaei, *DC. l. c.* 573.—Am. austr.
nudata, *Nutt. Gen. Am.* ii. 180.—Florida.
oblongifolia, Nutt. in Journ. Acad. Philad. vii. (1834) 76 = lanceolata.
odorata, Lam. Illustr. t. 704. f. 1 = Bidens odorata.
odoratissima, Cav. ex Pers. Syn. ii. 477 = Cosmos tenellus.
Oemleri, Ell. Sketch, ii. 435 = senifolia.
Oerstediana, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1852) 93.—Bias.
ovata, Cav. Ic. iii. 41. t. 280 = Actinomeris ovata.
pachyloma, *Oliver & Hiern, in Oliver, Fl. Trop. Afr.* iii. 391.—Afr. trop.
palmata, *Nutt. Gen. Am.* ii. 180.—Am. bor.
parviflora, Jacq. Hort. Schoenb. iii. 65 = Cosmos parviflorus.
pauciflora, Lehm. Ind. Sem. Hort. Hamb. (1833); Linnaea, x. (1836) Litt. 76 = palmata.
perfoliata, Walt. Fl. Carol. 214 = Bidens chrysanthemoides ?
Pickeringii, *A. Gray, in Proc. Am. Acad.* v. (1861) 124.—Peruv.
praecox, Fresen. in Linnaea, xiii. (1839) Litt. 93 = palmata.
Prestinaria, *Sch. Bip. ex Walp. Rep.* vi. 163.—Abyss.
prestinariaeformis, *Vatke, in Linnaea*, xxxix. (1875) 499.—Afr. trop.
procera, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 253 = Actinomeris squarrosa.
pubescens, *Ell. Sketch*, ii. 441.—Am. bor.
purpurea, Moç. & Sesse, ex DC. Prod. v. 604 = Cosmos purpureus.
quadricornis, Krock. Fl. Siles. ii. II. 481 = Bidens cernua.
radiata, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. bor.
reptans, Linn. Syst. ed. X. 1228; Amoen. Acad. v. 507 = Bidens rubifolia.
ridens, *Gunn. Fl. Norveg.* ii. 87.—Norvegia.
rigida, Nutt. Gen. Am. ii. 180 = delphinifolia.
rosea, *Nutt. l. c.* 179.—Am. bor.
rudis, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 196.—Mexic.
Rueppellii, Sch. Bip. ex Walp. Rep. vi. 163 = chrysantha
sambucifolia, Cav. Desc. 225 = Bidens sambucifolia.
scabra, *Rafin. Fl. Ludov.* 72.—Louisiana.
Schaffneri, *A. Gray, in Proc. Am. Acad.* xix. (1883) 15.—Mexic.
senifolia, *Michx. Fl. Bor. Am.* ii. 138.—Am. bor.
serrulata, Poir. Encyc. Suppl. iii. 352 = Bidens grandiflora.
setigera, *Sch. Bip. ex Walp. Rep.* vi. 163.—Abyss.
spectabilis, *A. Gray, in Proc. Am. Acad.* v. (1861) 125.—Peruv.
stellata, Banks, ex DC. Prod. v. 572 = senifolia.
stellata, *Nutt. in Journ. Acad. Philad.* vii. (1834) 76.—Am. bor.
Steppia, *Steetz, in Peters, Reise Mossamb. Bot.* 496.—Afr. trop.
Tanna, Steud. Nom. ed. II. i. 420 = Glossogyne tenuifolia.
tannensis, Spreng. Fl. Hal. Mant. 53 = Glossogyne tenuifolia.
tenuifolia, Ehrh. Beitr. vii. 168 = verticillata.
tetragona, Cerv. ex La Llave & Lex. Nov. Veg. Desc. fasc. i. 31 = Bidens tetragona.
tinctoria, *Nutt. in Journ. Acad. Philad.* ii. (1821) 114.—Am. bor.
trichosperma, *Michx. Fl. Bor. Am.* ii. 139.—Am. bor.
trifida, Poir. Encyc. Suppl. ii. 353 = Thelesperma filifolium.

COREOPSIS:—
trifoliata, *Bertol. Fl. Guatimal.* 36.—Guatemala.
Tripteris, *Linn. Sp. Pl.* 908.—Am. bor.
variifolia, Salisb. Prod. 206 = Bidens rubifolia.
venusta, *H. B. & K. Nov. Gen. et Sp.* iv. 229.—Peruv.
verticillata, Ehrh. Beitr. vii. 168 = delphinifolia.
verticillata, Lam. Encyc. ii. 108 = tenuifolia.
verticillata, *Linn. Sp. Pl.* 907.—Am. bor.
Wrayi, Nutt. in Journ. Acad. Philad. vii. (1834) 76 = senifolia.

COREOPSOIDES, Moench, Meth. 594 (1794) = **Coreopsis**, Linn. (Compos.).
lanceolata, Moench, l. c. = Coreopsis lanceolata.

COREOSMA, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 2 = **Ribes**, Linn. (Saxifrag.).
affinis, Spach, l. c. 24 = R. campanulatum.
Dombeyana, Spach, l. c. 25 = R. viscosum.
florida, Spach, l. c. 22 = R. floridum.
multiflora, Spach, l. c. 24 = R. Kunthii.
sanguinea, Spach, Hist. Vég. Phan. vi. 155 = R. sanguineum.
viscosissima, Spach, in Ann. Sc. Nat. Sér. II. iv. (1835) 23 = R. viscosissimum.

CORESANTHA, Alef. in Bot. Zeit. xxi. (1863) 298 = **Iris**, Tourn.
alata, Klatt, in Linnaea, xxxiv. (1865–66) 575 = I. stylosa.
caucasica, Klatt, l. c. = I. caucasica.
persica, Alef. in Bot. Zeit. xxi. (1863) 298 = I. persica.

CORESANTHE, Baker, in Journ. Linn. Soc. xvi. (1877) 122 = praec.

CORETHRODENDRON, Fisch. & Basin. in Mém. Sav. Etr. Pétersb. vi. (1851) 90 (Monogr. Hedys. 48. t. 2). *LEGUMINOSAE*, Benth. & Hook. f. i. 512.
scoparium, *Basin. l. c.* 91.—Soongaria.

CORETHROGYNE, DC. Prod. v. 215 (1836). *COMPOSITAE*, Benth. & Hook. f. ii. 270.
californica, *DC. l. c.*—Calif.
detonsa, Greene, in Bull. Torrey Club, x. (1883) 41 Diplostephium canum.
filaginifolia, *Nutt. in Trans. Am. Phil. Soc.* Ser. II. vii. (1841) 290.—Calif.
incana, Nutt. l. c. = californica.
obovata, *Benth. Bot. Voy. Sulph.* 22.—Calif.
spathulata, A. Gray, in Proc. Am. Acad. vii. (1868) 351 = obovata.
tomentella, Torr. & Gray, Fl. N. Am. ii. 99 = filaginifolia.
virgata, Benth. Bot. Voy. Sulph. 23 = filaginifolia.

CORETHROSTYLES, Benth. & Hook. f. Gen. i. 228 (1862), sphalm. = seq.

CORETHROSTYLIS, Endl. Nov. Stirp. Dec. i. (1839) 1 = **Lasiopetalum**, Sm. (Sterculi.).
bracteata, Endl. l. c. = L. bracteatum.
cordifolia, Steetz, in Lehm. Pl. Preiss. ii. 344 = L. cordifolium.
coriacea, Steud. l. c. i. 236 = L. bracteatum.
membranacea, Steud. l. c. = L. membranaceum.
microphylla, Turcz. in Bull. Soc. Nat. Mosc. xxv. (1852) II. 148 = L. cordifolium.
oppositifolia, F. Muell. Fragm. ii. 6 = L. floribundum.
parviflora, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) I. 174 = L. floribundum.
Schulzenii, F. Muell. in Trans. Phil. Soc. Vict. i. (1855) 36 = L. Schulzenii.

CORETHRUM, Vahl, in Skr. Naturk. Selsk. Kiobenh. vi. (1810) 85 = **Bouteloua**, Lag. (Gramin.).
bromoides, Vahl, l. c. = B. bromoides.

CORIANDRUM, [Tourn.] Linn. Syst. ed. I. (1735). *UMBELLIFERAE*, Benth. & Hook. f. i. 926.
KERAMOCARPUS, Fenzl, in Russegg. Reise ii. 966 (1843).
americanum, Nutt. ex DC. Prod. iv. 250 = Bifora americana.
Cicuta, Crantz, Stirp. Austr. ed. I. fasc. iii. 100; ed. II. 211 = Conium maculatum.
Cicuta, Roth, Tent. Fl. Germ. i. 130 = Cicuta virosa.

CORIANDRUM:—
Cynapium, Crantz, Stirp. Austr. ed. I. fasc. iii. 99; ed. II. 211 = Aethusa Cynapium.
didymum, Stokes, Bot. Mat. Med. ii. 121 = Bifora testiculata.
diversifolium, Gilib. Fl. Lituan. ii. 26 = sativum.
globosum, Salisb. Prod. 166 = sativum.
latifolium, Crantz, Stirp. Austr. ed. I. fasc. iii. 101; ed. II. 211 = Sium latifolium.
maculatum, Roth, Tent. Fl. Germ. i. 130 = Conium maculatum.
majus, Goüan, Hort. Monsp. 145 = sativum.
melphitense, *Tenore & Guss. Ind. Sem. Hort. Neap.* (1837) 3.—Italia.
sativum, *Linn. Sp. Pl.* 256.—Europ. austr.; Oriens.
seselifolium, DC. ex DC. Prod. iv. 250 = Astoma seselifolium.
tordylioides, *Boiss. Fl. Orient.* ii. 921.—Oriens.
testiculare, Salisb. Prod. 166 = Bifora testiculata.
testiculatum, Bieb. Fl. Taur. Cauc. i. 228 = Bifora radians.
testiculatum, Linn. Sp. Pl. 256 = Bifora testiculata.
testiculatum, Lour. Fl. Cochinch. 225 = Bifora Loureiri.

CORIARIA, Niss. ex Linn. Gen. ed. I. 310 (1737). *CORIARIEAE*, Benth. & Hook. f. i. 429.
HETEROCLADUS, Turcz. in Bull. Soc. Nat. Mosc. xx. (1847) i. 152.
HETEROPHYLLEIA, Turcz. l. c. (1848) i. 591.
angustissima, *Hook. f. Handb. New Zeal. Fl.* 47.—N. Zel.
arborea, *Lindsay, Contrib. N. Z. Bot.* 84.—N. Zel.
atropurpurea, Moç. & Sesse, ex DC. Prod. i. 740 = thymifolia.
hermaphrodita, Turra, Farset. Nov. Gen. 13 = myrtifolia.
japonica, *A. Gray, in Mem. Acad. Am.* N. S. vi. (1858–59) 383.—Japon.
microphylla, *Poir. Encyc.* vi. 87.—Peruv.
myrtifolia, *Linn. Sp. Pl.* 1037.—Reg. Mediterr.
nepalensis, *Wall. Pl. As. Rar.* iii. 67.—Reg. Himal.; China.
phylicifolia, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 819.—Peruv.
procumbens, *Whitl. ex Hoffmgg. Verz. Pfl. Nachtr.* iii. 9, nomen.—Hab.?
ruscifolia, *Linn. Sp. Pl.* 1037.—Peruv.
sarmentosa, *Forst. f. Prod.* 71. n. 377.—N. Zel.
sinica, *Maxim. in Mém. Acad. Pétersb.* Sér. VII. xxix. (1881) n. III. 9.—China.
thymifolia, *Humb. & Bonpl. ex Willd. Sp. Pl.* iv. 819.—Peruv.
tinctoria, Dulac, Fl. Hautes-Pyr. 243 = myrtifolia.
Tutu, *Lindsay, Contrib. N. Z. Bot.* 84.—N. Zel.

CORIDOCHLOA, Nees, in Edinb. N. Phil. Journ. xv. (1833) 381, in nota = **Panicum**, Linn. (Gramin.).
cimicina, Nees, l. c. = P. cimicinum.
semialata, Nees, ex Benth. Fl. Austral. vii. 473 = P. semialatum.

CORIDOTHYMUS, Reichb. f. in Oestr. Bot. Wochenbl. vi. (1857) 160 = **Thymus**, Tourn. (Labiat.).
capitatus, Reichb. f. l. c. 161 = T. capitatus.

CORILUS, Nocca, in Usteri, Ann. Bot. ii. (1793) 50 = **Corylus**, Tourn. (Cupul.).

CORINDUM, Adans. Fam. ii. 388 (1763) = **Paullinia**, Linn. (Sapind.).

CORINDUM, Tourn. ex Medic. Malv. 110 (1787) = **Cardiospermum**, Linn. (Sapind.).
Halicacabum, Medic. l. c. = Card. Halicacabum.

CORINGIA, J. & C. Presl, 'Fl. Čech. 141'; Fl. Sicula, 79 (1826) = **Conringia**, Heist. (Crucif.).

CORINOCARPUS, Poir. in Dict. Sc. Nat. x. 441 (1818) = **Corynocarpus**, Forst. (Anacard.).

CORION, Hoffmgg. & Link, Fl. Port. ii. 457 (1820) = **Bifora**, Hoffm. (Umbellif.).
esticulatum, Hoffmgg. & Link, l. c. = B. testiculata.

CORION, Mitch. in Act. Nat. Cur. viii. (1748) 208, App. = **Spergularia**, J. & C. Presl (Caryophyll.).

CORIS, Tourn. ex Linn. Gen. ed. I. 338 (1737). *PRIMULACEAE*, Benth. & Hook. f. ii. 638.
hispanica, *Lange, in Kjoeb. Vidensk. Meddel.* (1863) 53 (*Pugill.* 219).—Hispan.
monspeliensis, *Linn. Sp. Pl.* 177.—Europ. austr.

CORISANTHERA, C. B. Clarke, in Hook. f. Fl. Brit. Ind. iv. 373 (1884) = CORYSANTHERA, Wall. = **Rhynchotechum**, Blume (Gesnerac.).

CORISANTHES, Steud. Nom. ed. II. i. 474, in syn. (1840) = CRIOSANTHES, Rafin. = **Cypripedium**, Linn. (Orchid.).

CORISPERMUM, B. Juss. ex Linn. Syst. ed. I (1735). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 57.
apetalum, *Hort. ex Steud. Nom.* ed. II. i. 420.—Hab.?
borysthenicum, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 129, nomen.—Rossia.
bracteatum, Viv. Lett. Corisp. 2 = Marschallii.
canescens, *Kit. ex Schult. Oestr. Fl.* ed. II. i. 7.—Europ. occ.
coloratum, *Andrz. ex Trautv. in Act. Hort. Petrop.* ix. (1884) 129, nomen.—Rossia.
confertum, *Bunge, in Maxim. Prim. Fl. Amur.* 225.—Reg. Amur.
crassifolium, *Turcz. in Bull. Soc. Nat. Mosc.* xxv. (1852) 428.—Sibiria.
declinatum, Steph. ex Stev. in Mém. Soc. Nat. Mosc. v. (1814) 334 = hyssopifolium.
elatum, Host, Fl. Austr. i. 319 = Marschallii.
elongatum, *Bunge, ex Maxim. Prim. Fl. Amur.* 224.—Manchur.
filifolium, *Becker, in Bull. Soc. Nat. Mosc.* xxxi. (1858) i. 13.—Rossia.
glomeratum, *Lag. Gen. et Sp. Nov.* 1.—Hab.?
Gmelini, *Bunge, in Bull. Acad. Pétersb.* xxv. (1879) 357.—Sibiria.
heteracanthum, *Ledeb. ex Heynh. Nom.* ii. 161, nomen.—Rossia.
hybridum, Bess. ex Moq. in DC. Prod. xiii. II. 141 = Marschallii.
hyssopifolium, *Linn. Sp. Pl.* 4.—Europ.; As. temp.; Am. bor.
hyssopifolium, Pall. Reise, i. 484; Fl. Ross. t. 98 = orientale.
intermedium, Reichb. Fl. Germ. Excurs. 584 = Marschallii.
intermedium, *Schweig. in Koenigsb. Arch. Naturw.* i. (1812) 211.—Europ.
latifolium, Link, Handb. ii. 408 = orientale.
laxiflorum, Schrenk, in Bull. Phys.-Math. Acad. Pétersb. i. (1843) 361 = Marschallii.
Lehmannianum, *Bunge, in Mém. Sav. Etr. Pétersb.* vii. (1851) 458.—Turkestan.
macrocarpum, *Bunge, in Maxim. Prim. Pl. Amur.* 226.—Manchur.
macrospermum, Trautv. in Act. Hort. Petrop. ix. (1884) 130 = macrocarpum.
marginatum, Steph. ex Moq. in DC. Prod. xiii. II. 140 = Marschallii.
Marschallii, *Stev. in Mém. Soc. Nat. Mosc.* v. (1814) 336.—Europ.
microspermum, Host. Fl. Austr. i. 319 = nitidum.
nitidum, *Kit. ex Schult, Oestr. Fl.* ed. II. i. 7.—Europ.; As. bor.
orientale, *Lam. Encyc.* ii. 111; *Illustr.* i. 15.—Rossia; As. bor.; Persia.
Pallasii, Stev. in Mém. Soc. Nat. Mosc. v. (1814) 336 = orientale.
patens, Fisch. in Roem. & Schult. Syst. i. 579 = hyssopifolium.
patens, Link, in Spreng. Jahrb. Gewaechsk. i. III. 28 = Redowskii.
patens, *Roem. & Schult. Syst.* i. 579.—Sibir.
pentandrum, Willd. ex Link, in Spreng. Jahrb. Gewaechsk. i. III. 28 = orientale.
pilosum, Rafin. New Fl. Am. iv. 46 = hyssopifolium?
polygaloides, Fisch. & Mey. Ind. Sem. Hort. Petrop. i. 24 = Anthochlamus polygaloides.
pungens, Savi, sec. Uechtr. ex Steud. Nom. ed. II. i. 421 = squarrosum.
pungens, Vahl, Enum. i. 117 = Agriophyllum arenarium, *Bieb.*
purpurascens, Host, Fl. Austr. i. 318 = nitidum.
Redofski, Fisch. & Stev. in Mém. Soc. Nat. Mosc. v. (1814) 336 = seq.

CORISPERMUM:—
Redowskii, [*Fisch.*] *Hort. Gorenk.* ed. I. (1808) 25.—Sibir. baical.
sabulosum, *Ledeb. ex Sweet, Hort. Brit.* ed. III. 257.—Sibir.
salsoloides, Hort. ex Steud. Nom. ed. II. i. 420 = glomeratum.
spicatum, Willd. Enum. Hort. Berol. Suppl. 1 = orientale.
squarrosum, Bieb. Fl. Taur. Cauc. i. 3 = Marschallii.
squarrosum, Link, Enum. Hort. Berol. i. 7 = orientale.
squarrosum, Linn. Sp. Pl. 4 = hyssopifolium.
squarrosum, Pall. Fl. Ross. i. II. 113. t. 99 = Agriophyllum arenarium, *Bieb.*
squarrosum, *Vahl, Enum.* i. 16.—Europ.; As. temp.
squarrosum, Willd. Enum. Hort. Berol. 8 = Marschallii.
Stauntonii, *Moq. Chenop. Enum.* 104.—China.
subpentandrum, Pall. Illustr. t. 59 = orientale.
tenue, Link, in Spreng. Jahrb. Gewaechsk. i. III. 29 = nitidum.
tylocarpum, *Hance, in Journ. Bot.* vi. (1868) 47.—China.
ulopterum, *Fenzl, ex Ledeb. Fl. Ross.* iii. 765.—Sibir.

CORISTOSPERMUM, Bertol. Fl. Ital. iii. 466 (1837) = **Ligusticum**, Linn. (Umbellif.).
cuneifolium, Bertol. l. c. 467 = L. cuneifolium.

CORIZOSPERMUM, Zipp. ex Blume, Mus. Bot. Lugd. Bat. i. 255 (1850) = **Casearia**, Jacq. (Samyd.).
clutiaefolium, Zipp. ex Blume, l. c. = Cas. clutiaefolia.

CORMIGONUS, Rafin. in Ann. Gén. Sc. Phys. vi. (1820) 83 = **Bikkia**, Reinw. (Rubiac.).

CORMONEMA, Reissek, ex Endl. Gen. 1098 (1840). *RHAMNEAE*, Benth. & Hook. f. i. 379.
CAESIA, Vell. Fl. Flum. iii. t. 23 (1827).
spinosum, *Reissek, in Mart. Fl. Bras.* xi. I. 96.—Bras.

CORMUS, Spach, Hist. Vég. Phan. ii. 96 (1834) = **Pyrus**, Linn. (Rosac.).
domestica, Spach, l. c. 97 = P. Sorbus.

CORNA, Noronha, in Verh. Batav. Gen. v. (1790) ed. I. Art. IV. 2 (quid?).

CORNACCHINIA, Savi, in Mem. Soc. Ital. Moden. xxi. (1837) 184. t. 7 = **Clerodendron**, Linn. (Verbenac.).
fragiformis, Savi, l. c. 185 = Clerod. Acerbiana.

CORNELIA, Ard. Spec. ii. 9 (1764) = **Ammannia**, Houst. (Lythrar.).
verticillata, Ard. l. c. = A. salicifolia.

CORNICINA, Boiss. Voy. Bot. Espagne, ii. 162 (1840?) = **Anthyllis**, Linn. (Legumin.).
circinata, Boiss. l. c. 163 = Hymenocarpus circinnatus.
hamosa, Boiss. l. c. 162 = A. hamosa.
Loeflingii, Boiss. l. c. = A. Cornicina.
lotoides, Boiss. l. c. = A. lotoides.
nummularia, Boiss. l. c. 163 = Hymenocarpus nummularius.
pseudocytisus, Boiss. Diagn. Ser. I. ii. 15 = Cytisopsis dorycniifolia.

CORNIDIA, Ruiz & Pav. Prod. 63. t. 35 (1794) = **Hydrangea**, Linn. (Saxifrag.).
integerrima, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 344 = H. scandens, *Poepp.*
radiata, Oerst. in Kjoeb. Vidensk. Meddel. (1856) 42 = H. peruviana.
serratifolia, Hook. & Arn. in Hook. Bot. Misc. iii. (1833) 344 = H. serratifolia.
umbellata, Ruiz & Pav. Syst. Veg. 91.—Peruv.

CORNIOLA, Adans. Fam. ii. 321 (1763) = **Genista**, Linn. (Legumin.).
anxanthica, Presl, Bot. Bemerk. 137 = G. anxantica.
corsica, Presl, l. c. 136 = G. corsica.
mantica, Presl, l. c. = G. mantica.
microphylla, Presl, l. c. 137 = G. microphylla.
ovata, Presl, l. c. 136 = G. ovata.
scariosa, Presl, l. c. = G. triangularis.
Scorpius, Presl, l. c. 137 = G. Scorpius.
sibirica, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 342 = G. sibirica.
tinctoria, Medic. l. c. = G. tinctoria.
triangularis, Presl, Bot. Bemerk. 136 = G. triangularis.
virgata, Presl, l. c. = G. virgata.

CORNUCOPIAE, Linn. Gen. ed. I. 13 (1737). *GRAMINEAE*, Benth. & Hook. f. iii. 1139.
alopecuroides, Linn. Mant. i. 28 = Alopecurus utriculatus.
altissimum, Walt. Fl. Carol. 74 = Agrostis elata?
cucullatum, *Linn. Sp. Pl.* 54.—As. Min.; Mesopot.
hyemale, Walt. Fl. Carol. 73 = Agrostis scabra.
perennans, Walt. l. c. 74 = Agrostis scabra.

CORNULACA, Delile, Fl. Egypte, 206. t. 22. f. 3 (1813). *CHENOPODIACEAE*, Benth. & Hook. f. iii. 69.
amblyacantha, *Bunge, Anabas. Rev.* 88.—Persia.
Aucheri, *Moq. Chenop. Enum.* 163.—Persia.
Ehrenbergii, *Aschers. in Schweinf. Beitr. Fl. Aethiop.* 184.—Afr. trop.
monacantha, *Delile, Fl. Egypte*, 206.—Aegypt.; Persia.
muricata, Delile, Fl. Egypte Illustr. 57 = Kochia muricata.
setifera, Moq. in DC. Prod. xiii. II. 218 = tragacanthoides.
tragacanthoides, *Moq. Chenop. Enum.* 163.—Syria; Mesopot.

CORNUS, [Tourn.] Linn. Syst. ed. I (1735). *CORNACEAE*, Benth. & Hook. f. i. 950.
BENTHAMIA, Lindl. Bot. Reg. t. 1579 (1833).
CYNOXYLON, Rafin. Alsog. Am. 59 (1838).
EUKRANIA, Rafin. l. c. (1838).
SVIDA, Opiz, Seznam, 94 (1852).
TELUKRAMA, Rafin. Alsog. Am. 64 (1838).
alba, Hook. & Arn. Bot. Beech. Voy. 142 = californica.
alba, Lam. Encyc. ii. 115, partim (et auct. Amer.) = stolonifera.
alba, *Linn. Mant.* i. 40.—Sibir.
alba, Thunb. Fl. Jap. 62 = macrophylla.
alba, Walt. Fl. Carol. 88 = C. Amomum.
albida, Ehrh. Beitr. iv. 16 = candidissima.
alterna, Marsh. Arbust. 35 = alternifolia.
alternifolia, *Linn. f. Suppl.* 125.—Am. bor.
Amomum, *Mill. Gard. Dict.* ed. VIII. n. 5.—Am. bor.
asperifolia, *Michx. Fl. Bor. Am.* i. 93.—Am. bor.
atrata, *Rafin. Alsog. Am.* 61.—Am. bor.
australis, C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb. iii. (1845) 372 = sanguinea.
biramis, Stokes, Bot. Mat. Med. i. 221 = suecica.
borealis, Krasch. Fl. Ingr. 24 = suecica.
brachypoda, C. A. Mey. in Mém. Acad. Pétersb. vii. (1845) II. 223 = macrophylla.
caerulea, Auct. vide *coerulea*.
californica, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 373.—Am. bor. occ.
canadensis, Cham. & Schlecht. in Linnaea, iii. (1828) 139 = unalaschkensis.
canadensis, *Linn. Sp. Pl.* 118.—Am. bor.
canadensis, Willd. ex Ledeb. Fl. Ross. ii. 377 = suecica.
candissima, Bisch. Ind. Sem. Hort. Heidelb. (1851) = stolonifera.
candidissima, *Mill. Gard. Dict.* ed. VIII. n. 6.—Am. bor.
capitata, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 434.—Reg. Himal.
caudata, Zoll. in Flora, xxx. (1847) 603 = Polyosma integrifolia.
chilensis, Molina, Sagg. Chil. 173 = Aristotelia Maequi.
cinerea, *Rafin. Alsog. Am.* 59.—Am. bor.
circinata, *L'Hérit. Cornus*, 7.—Am. bor.
circinnata, Cham. & Schlecht. in Linnaea, iii. (1828) 139 = californica.
citrifolia, Hort. ex Lam. Encyc. ii. 116 = candidissima.
citrifolia, Wahlenb. in Isis, xxi. (1828) 982 = sanguinea.
coerulea, Lam. Encyc. ii. 116 = C. Amomum.
coerulea, Meerb. Pl. Sel. t. 3 = stricta.
comosa, *Rafin. Alsog. Am.* 63.—Am. bor.
crispula, Hance, in Journ. Bot. xix. (1881) 216 = macrophylla.
cyananthus, *Rafin. Atl. Journ.* 151.—Am. bor.
cyanocarpus, J. F. Gmel. Syst. 257 = stricta.
cyanocarpus, Moench, Verz. 27 = C. Amomum.
davurica, Laxm. ex Ledeb. Fl. Ross. ii. 386 = Viburnum davuricum.
dichotoma, *Rafin. Alsog. Am.* 60.—Am. bor.
disciflora, *Moç. & Sesse, ex DC. Prod.* iv. 273.—Mexic.
Drummondi, *C. A. Mey. in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 372.—Am. bor.

CORNUS:—
erythrocarpa, St. Lager, ex Bull. Soc. Bot. Fr. xxxi. (1883) Bibl. 201 = C. Mas.
excelsa, *H. B. & K. Nov. Gen. et Sp.* iii. 430.—Mexic.
fastigiata, Michx. Fl. Bor. Am. i. 92 = stricta.
femina, Auct. vide *foemina*.
ferruginea, Hort. ex Lam. Encyc. ii. 116 = C. Amomum.
ferulacea, Jacq. ex Roem. & Schult. Syst. iii. 325 = ferulaefolia.
ferulaefolia, *Nocca, Syn. Pl.* 38.—Hab.?
flava, Steud. Nom. ed. I. 227 = C. Mas.
florentina, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 157.—Hab.?
florida, Hook. Fl. Bor. Am. i 277, partim = Nuttallii.
florida, *Linn. Sp. Pl* 117.—Am. bor.
foemina, Mill. Gard. Dict. ed. VIII. n. 4 = sanguinea.
glabrata, *Benth. Bot. Voy. Sulph.* 18.—Calif.
grandis, *Cham. & Schlecht. in Linnaea*, v. (1830) 171.—Mexic.
herbacea, Steller, in Pall. N. Nord. Beitr. ii. 300 = suecica.
ignorata, *C. Koch, Ind. Sem. Hort. Berol.* (1867) *App.* i. 8.—Japon.
ilicifolia, Hassk. & Zoll. ex Hassk. Cat. Hort. Bog. Alt. 168, 311 = Polyosma ilicifolia.
japonica, *Thunb. Fl. Jap.* 63.—Japon.
Kousa, *Buerger, ex Miq. Ann. Mus. Bot. Lugd. Bat.* ii. 159, in syn.—Japon.
lancifolia, *Rafin. Alsog. Am.* 60.—Am. bor.
lanuginosa, Michx. Fl. Bor. Am. i 92 = C. Amomum.
latifolia, Bray, in Denkschr. Bot. Ges. Regensb. i. II. (1818) 35 = sanguinea.
longifolia, *Medic. Beobacht.* (1782) 308.—Hab.?
macrophylla, *Wall. in Roxb. Fl. Ind. ed Carey*, i. 433.—Reg. Himal.; China; Japon.
Mas, *Linn. Sp. Pl.* 117.—Europ.; As. Min.
mascula, Linn. Syst. ed. X. 897 = C. Mas.
mexicana, Carr. in Rev. Hortic. (1875) 460 = C. Amomum.
minor, Hort. ex Steud. Nom. ed. II. i. 421 = sericea.
nudiflora, Dum. Fl. Belg. 83 = C. Mas.
Nuttallii, *Audubon, Birds Am.* t. 367; *ex Torr. & Gray, Fl N. Am.* i. 652.—Am. bor.
obliqua, Rafin. Ann. Nat. (1820) 13 = C. Amomum.
oblonga, *Wall. in Roxb. Fl. Ind. ed. Carey*, i. 432.—Reg. Himal.
oblongifolia, Rafin. ex DC. Prod. iv. 272 = sericea.
officinalis, *Sieb. & Zucc. Fl. Jap.* i. 100. t. 50.—Japon.
paniculata, Buch.-Ham. ex D. Don, Prod. Fl. Nep. 140 = oblonga.
paniculata, L'Hérit. Cornus, 9. t. 5 = candidissima.
parvifolia, *Rafin. Alsog. Am.* 61.—Am. bor.
paucinervis, *Hance, in Journ. Bot.* xix. (1881) 216.—China.
plicata, Tausch, in Flora, xxi. (1838) 733 = alternifolia.
polygama, Rafin. Fl. Ludov. 78 = C. Amomum?
praecox, Stokes, Bot. Mat. Med. i. 222 = C. Mas.
pubescens, Nutt. Sylva, iii. 54 = C. Amomum.
pubescens, Torr. Bot. Mex. Bound. 71 = californica.
pubescens, Willd. ex Roem. & Schult. Syst. iii. 252 = excelsa.
punctata, *Rafin. Alsog. Am.* 62.—Am. bor.
purpurea, Tausch, in Flora, xxi. (1838) 731 = tartarica.
Purshii, G. Don, Gen. Syst. iii 399 = stolonifera.
racemosa, Lam. Encyc. ii. 116 = candidissima.
riparia, *Rafin. Alsog. Am.* 62.—Am. bor.
rotundifolia, *Rafin. l. c.*—Am. bor.
rubiginosa, Ehrh. Beitr. iv. 15 = C. Amomum.
rugosa, Lam. Encyc. ii. 115 = circinata
sanguinea, Forsk. Fl. Aegypt. Arab. 33 = Cordia Myxa.
sanguinea, *Linn. Sp. Pl.* 117.—Europ.; As. bor.
sanguinea, Thunb. Fl. Jap. 63 = macrophylla.
sanguinea, Walt. Fl. Carol. 88 = stricta.
sericea, Linn. Mant. ii. 199; L'Hérit. Cornus, 5. t. 2 = C. Amomum.
serotina, *Rafin. Alsog. Am.* 60.—Am. bor.
serrulata, Hassk. & Zoll. ex Hassk. Cat. Bog. Alt. 168 = Polyosma serrulata.
sessilis, *Torr. ex Durand, in Journ. Acad. Nat. Sc. Philad.* Ser. II. iii. (1855) 89.—Am. bor. occ.
sibirica, Lodd. in Loud. Hort. Brit. 50 = tartarica.
stolonifera, *Michx. Fl. Bor. Am.* i. 92.—Am. bor.

CORNUS:—
striata, DC. Prod. iv. 272 = foemina.
stricta, A. Gray, in Hall, Pl. Texas, 11, ex S. Wats. Bibliog. Ind. N. Am. Bot. i. 438 = Drummondii.
stricta, L'Hérit. Cornus, 8 = foemina.
stricta, *Lam. Encyc.* ii. 116.—Am. bor.
stricta, Zoll. & Morr. in Nat. en Geneesk. Arch. Neêrl. Indïe, ii. (1845) 10 = Polyosma integrifolia.
suecica, A. Gray, in Proc. Am. Acad. viii. (1873) 387 = canadensis.
suecica, *Linn Sp. Pl.* 118.—Reg. bor. et arct.
suffruticosa, *Rafin. Atl. Journ.* 151.—Am. bor.
tartarica, *Mill. Gard. Dict* ed. VIII. n. 7.—Sibir.
tolucensis, *H. B. & K. Nov. Gen. et Sp.* iii. 430.—Mexic.
tomentosa, Steud. Nom. ed. II. i. 422 = circinata.
tomentosula, DC. Prod. iv. 272 = circinata.
tomentulosa, Michx. Fl. Bor. Am. i. 91 = circinata.
Torreyi, *S. Wats. in Proc. Am. Acad.* xi. (1876) 145.—Calif.
trilobata, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 157.—Hab.?
unalaschkensis, *Ledeb. Fl. Ross.* ii. 378.—Alaska.
undulata, *Rafin. Alsog. Am.* 61.—Am. bor.
vernalis, Salisb. Prod. 66 = C. Mas.
virginiana, Hort. ex Lam. Encyc. ii. 115 = circinata.

CORNUTA, Linn. Gen. ed. VI. 316 (1764) = **Cornutia**, Plum. (Verbenac.).

CORNUTIA, Plum. ex Linn. Gen. ed. I. 366 (1737). *VERBENACEAE*, Benth. & Hook. f. ii. 1153.
AGNANTHUS, Vaill. ex Linn. Gen. ed. I. 366 (1737).
CORNUTA, Linn. l. c. ed. VI. 316 (1764).
HOSTA, Jacq. Hort. Schoenb. i. 60. t. 114 (1797).
HOSTANA, Pers. Syn. i. 143 (1805).
cayennensis, *Schau. in DC. Prod.* xi. 681.—Guiana.
corymbosa, Burm. f. Fl. Ind. 132 = Premna corymbosa.
corymbosa, Lam. Encyc. i. 54 = Callicarpa lanata et Premna tomentosa.
crenata, Medic. Beobacht. (1783) 112 = pyramidata.
grandiflora, Steud. Nom. ed. II. i. 422 = sep.
grandifolia, *Schau. in DC. Prod.* xi. 682.—Mexic.
longifolia, *Spreng. Syst.* i. 39.—Mexic.
odorata, *Poepp. ex Schau. in DC. Prod.* xi. 681.—Am. trop.
pubescens, *Gaertn. f. Fruct.* iii. 172. t. 213.—Guiana.
punctata, *Willd. Sp. Pl.* iii. 322.—Am. trop.
pyramidata, Ait. Hort. Kew. ii. 353 = punctata.
pyramidata, *Linn. Sp. Pl.* 628.—Mexic.
quinata, Lour. Fl. Cochinch. ii. 387 = Vitex Loureirii.

CORNUTIOIDES, Linn. Fl. Zeyl. 195 (1747) (Gen. dub.).

COROCEPHALUS, D. Dietr. Syn. Pl. i. 424, 540 (1839), sphalm. = **Conocephalus**, Blume (Urtic.).

CORONA-IMPERIALIS, Tourn. ex Rupp. Fl. Jen. ed. Hall. 148 (1745) = **Fritillaria**, Linn. (Liliac.).

COROKIA, A. Cunn. in Ann. Nat. Hist. iii. (1839) 249. *CORIACEAE*, Benth. & Hook. f. i. 949.
buddleioides, *A. Cunn. l. c.*—N. Zel.
Cotoneaster, *Raoul, Choix Pl. N. Zel.* 22.—N. Zel.

CORONANTHERA, Vieill. ex C. B. Clarke, in DC. Monog. Phan. v. 170 (1883). *GESNERACEAE*.
aspera, *C. B. Clarke, l. c.* 172.—N. Caled.
barbata, *C. B. Clarke, l. c.*—N. Caled.
deltoidifolia, *Vieill. ex C. B. Clarke, l. c.* 171.—N Caled.
glabra, *C. B. Clarke, l. c.* 174.—N. Caled.
Pancheri, *C. B. Clarke, l. c.*—N. Caled.
pedunculosa, *C. B. Clarke, l. c.* 173.—N. Caled.
pinguior, *C. B. Clarke, l. c.* 171.—N. Caled.
pulchra, *C. B. Clarke, l. c.* 173.—N. Caled.
sericea, *C. B. Clarke, l. c.* 172.—N. Caled.

CORONARIA, Linn. Gen. ed. I. 135 (1737) = **Lychnis**, Linn. (Caryophyll.).
Coeli-rosa, Fries, ex Heynh. Nom. ii. 161 = L. Coeli-rosa.
Flos-cuculi, A. Br. in Flora, xxvi. (1843) 365 = L. Flos-cuculi.
Flos-Jovis, A. Br. l. c. = L. Flos-Jovis.
tomentosa, A. Br. l. c. = L. Coronaria.

CORONE, Hoffmgg. ex Steud. Nom. ed. II. i. 422 (1840) = **Silene**, Linn. (Caryophyll.).
agrestina, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 344 = S. sylvestris.
gallica, Fourr. l. c. = S. gallica.
nocturna, Fourr. l. c. = S. nocturna.
quinquevulnera, Hoffmgg. ex Steud. Nom. ed. II. i. 422 = S. quinquevulnera.
vespertina, Hoffmgg. ex Steud. l. c = S. vespertina.

CORONILLA, Tourn. ex Linn. Gen. ed. I. 227 (1737). *LEGUMINOSAE*, Benth. & Hook. f. i. 509.
AVIUNCULUS, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvi. (1868) 367.
CALLISTEPHANA, Fourr. l. c. (1868).
EMERUS, Tourn. ex Mill. Gard. Dict. ed. VI. (1752).
SCORPIUS, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 369.
aculeata, Willd. Sp. Pl. iii. 1147 = Sesbania aculeata.
argentea, Burm. f. Cap. Prod. 22; Thunb. Prod. ii. 131 = Lessertia argentea.
argentea, *Linn. Sp. Pl.* 743.—Ins. Creta.
atlantica, *Boiss. & Reut. Pugill. Pl. Nov.* (1852) 42, in syn.—Marocco.
Balansae, Boiss. & Bal. in Boiss. Fl. Orient. ii. 181 = cappadocica.
buxifolia, *Hance, in Ann. Sc. Nat.* Sér. IV. xviii. (1862) 219.—China.
cannabina, Willd. Sp. Pl. iii. 1147 = Sesbania aculeata.
cappadocica, *Willd. l. c.* 1152.—As. Min.
chia, Orph. ex Boiss. Fl. Orient. ii. 183 = parviflora.
Clusii, Dufour, in Ann. Sc. Gén. Phys. vii. (1820) 307 = minima.
coccinea, Willd. Sp. Pl. iii. 1145 = Sesbania grandiflora.
cochinchinensis, *Lour. Fl. Cochinch.* ii. 452.—Cochinch.
coronata, DC. Prod. ii. 310 = minima.
coronata, Jacq. Fl. Austr. i. 59. t. 95 = montana.
coronata, *Linn. Syst.* ed. X. 1168.—Europ. austr.
cretica, *Linn. Sp. Pl.* 743.—Europ. or.; As. Min.
dura, *Boiss. Diagn.* Ser. II. ii. 34.—Lusitan.
elegans, Panč. ex Nym. Consp 185 = varia?
emera, St. Lag. in Ann. Soc. Bot. Lyon, vii. (1880) 123 = C. Emerus.
emeroides, *Boiss. & Sprun. Diagn.* Ser. I. ii. 100.—Graecia; As. Min.
Emerus, *Linn. Sp. Pl.* 742.—Europ. austr.
Emerus, Sibth. & Sm. Fl. Graec. viii. 7. t. 710 = emeroides.
eriocarpa, Boiss. Elench 37 = Hippocrepis squamata.
extensa, Jord. Pugill. 61 = minima.
florida, Salisb. Prod. 341 = C. Emerus.
fruticans, *Jord. Pugill.* 62.—Europ.
gibraltarica, *Brouss. ex Ball, in Journ. Linn. Soc.* xvi. (1878) 428.—Hispan.; Afr. bor.
glauca, Asso, ex Nym. Consp. 185 = coronata.
glauca, *Linn. Cent. Pl.* i. 23.—Reg. Mediterr.
globosa, *Lam. Encyc.* ii. 122.—Ins. Creta.
Glycyrrhiza, Spreng. Syst. iii. 324 = Glycyrrhiza asperrima.
grandiflora, *Boiss. Diagn.* Ser. I. v. 88.—As. Min.
grandiflora, Willd. Sp. Pl. iii. 1145 = Sesbania grandiflora.
Haussknechtii, Reut. ex Boiss. Fl. Orient. ii. 182 = varia.
Heineri, *Regel, Cat. Pl. Hort. Aksakov.* 41.—As. temp.
hirsuta, *DC. Prod.* ii. 310.—Afr. austr.
hirsuta, Vell. Fl. Flum. 311; vii. t. 122 = Isodesmia tomentosa.
hirta, Bunge, ex Boiss. Fl. Orient. ii. 182 = varia.
hispanica, Mill. Gard. Dict. ed. VIII. n. 4 = glauca.
iberica, Bieb. Fl. Taur. Cauc. ii. 171 = cappadocica.
juncea, *Linn. Sp. Pl.* 742.—Europ. austr.
libanotica, Boiss. Diagn. Ser. I. ix. 104 = varia.
lotoides, Koch, in Roehl. Deutschl. Fl. ed. Mert. & Koch, v. 199 = minima.
minima, Jacq. Fl. Austr. iii. 39. t. 271 = virginalis.
minima, *Linn. Cent. Pl.* ii. 28.—Europ.
monilis, Linn. Pl. Surin. n. 102; Amoen. Acad. viii. 261 = Muellera moniliformis?
montana, *Scop. Pl. Carn.* ed. II. ii. 72.—Reg. Caucas.; Persia.
multiflora, *DC. Prod.* ii. 310.—Hispan.
myrtifolia, Burm. f. Fl. Cap. Prod. 22 = Tephrosia myrtifolia.
occidentalis, Willd. Sp. Pl. iii. 1147 = Sesbania occidentalis.
orbiculata, Moench, Meth. Suppl. 40 = valentina.

CORONILLA:—
orientalis, *Mill. Gard. Dict.* ed. VIII. n. 8.—Iberia; As. Min.
pallida, Salisb. Prod. 341 = glauca.
parviflora, Moench, Meth. 121 = cretica.
parviflora, *Willd. Sp. Pl.* iii. 1155.—Graecia; As. Min.
pauciflora, Lam. Fl. Fr. ii. 661 = C. Emerus.
pendula, *Kit. in Linnaea*, xxxii. (1863) 634.—Europ.
pentaphylla, *Desf. Fl. Atlant.* ii. 171.—Afr. bor.
picta, Willd. Sp. Pl. iii. 1148 = Sesbania aegyptiaca.
pubescens *Schum. & Thonn. in Kon. Dan. Vid. Sel. Afh.* iv. (1828) 128.—Afr. trop.
pulchra, Ball, in Journ. Bot. xi. (1873) 307 = viminalis.
ramosissima, *Ball, l. c.* 306.—Marocco.
repanda, *Boiss. Diagn.* Ser. II. ii. 35.—Hispan.
rostrata, Boiss. & Sprun. Diagn. Ser. I. ii. 100 = parviflora.
sanguinea, *Roxb. Hort. Beng.* 56, nomen.—Hab.?
scandens, *Linn. Sp. Pl.* 743.—Ind. occ.
scandens, Vell. Fl. Flum. 310; vii. t. 121 = Chaetocalyx brasiliensis.
scorpioides, *Koch, in Roehl. Deutschl. Fl. ed Mert. & Koch*, v. 201.—Reg. Mediterr.; Persia.
Securidaca, Linn. Sp. Pl. 743 = Securigera Coronilla.
Securigera, Salisb. Prod. 341 = Securigera Securidaca.
sericea, Willd. Enum. Hort. Berol. 773 = Sesbania sericea.
Sesban, Moench, Meth. 120 = Sesbania aegyptiaca.
squamata, Cav. Ic. ii. 43 = Hippocrepis squamata.
stipularis, Lam. Encyc. ii. 120 = valentina.
stipulata, *Buch.-Ham. ex D. Don, Prod. Fl. Nep.* 246.—Reg. Himal.
suaveolens, Salisb. Prod. 340 = valentina.
surattensis, Garc. ex DC. Prod. ii. 323 = Smithia sensitiva.
thymifolia, *Burm. f. Fl. Cap. Prod.* 22.—Afr. austr.
vaginalis, *Lam. Encyc.* ii. 121.—Europ.
vaginalis, Ledeb. Fl. Ross. i. 625 = emeroides.
valentina, Lam. Fl. Fr. ii. 663 = coronata.
valentina, *Linn. Sp. Pl.* 742.—Europ. austr.
valentina, Pall. ex Bieb. Fl. Taur. Cauc. ii. 173 = parviflora.
varia, *Linn. Sp. Pl.* 743.—Europ.
viminalis, *Salisb. Parad. Lond.* t. 13.—Afr. bor.
virgata, Willd. Sp. Pl. iii. 1148 = Coursetia virgata.

CORONOCARPUS, Schum. & Thonn. Beskr. Guin. Pl. 393 (1827) = **Aspilia**, Thou. (Compos.).
Gayanus, Benth. in Hook. Niger Fl. 434 = A. helianthoides.
helianthoides, Benth. l. c. = A. latifolia.
helianthoides, Schum. & Thonn. Beskr. Guin. Pl. 393 = A. helianthoides.
Kotschyi, Benth. in Hook. Niger Fl. 433 = A. Kotschyana.
Prieurianus, Benth. l. c. = A. Kotschyana.

CORONOPUS, Reichb. Handb. 202 (1837) = **Plantago**, Tourn.
vulgaris, Fourr. in Ann. Soc. Linn. Lyon, N. S. xvii. (1869) 140 = P. Coronopus.

CORONOPUS, Rupp. ex Linn. Syst. ed. I (1735); Gaertn. Fruct. ii. 293 (1791) = **Senebiera**, DC. (1799) (Crucif.).
anomalus, Spreng. Syst. ii. 853 = Brachycarpaea varians.
coadunatus, Stokes, Bot. Mat. Med. iii. 440 = S. Coronopus.
cristatus, [Clairv.] Man. Herb. 212 = S. Coronopus.
depressus, Moench, Meth. 220 = S. Coronopus.
didymus, Sm. Fl. Prit. ii. 691 = S. didyma.
fungosus, Sm. ex DC. Syst. ii. 522 = S. integrifolia.
Helenae, Spreng. Syst. ii. 853 = S. Heleniana.
incisus, Hornem. Hort. Hafn. 599 = S. didyma.
integrifolius, Spreng. Syst. ii. 853 = S. integrifolia.
linoides, Spreng. l. c. = S. integrifolia.
myricaefolius, Sm. ex DC. Syst. ii. 526 = S. serrata.
niloticus, Spreng. Syst. ii. 853 = S. nilotica.
pinnatifidus, Dulac, Fl. Hautes-Pyr. [186] = S. didyma.
pinnatus, Hornem. Hort. Hafn. 599 = S. didyma.
procumbens, Gilib. Fl. Lituan. ii. 52 = S. Coronopus.
Raddii, Savi, in Mem. G. Raddi, (1830) 33.—Aegypt.
Ruellii, All. Fl. Pedem. i. 256 = S. Coronopus.
serratus, Desv. Journ. Bot. iii. (1814) 163 = S. serrata.
squamatus, Aschers. Fl. Brandenb. 62 = S. Coronopus.
vulgaris, Desf. Tabl. ed. I. 132 = S. Coronopus.

COROTHAMNUS, Presl (Bot. Bemerk. 137); in Abh. Boehm. Ges. V. iii. (1844) 567 = **Genista**, Linn. (Legumin.).
decumbens, C. Koch, Dendrol. i. 44 = Cytisus decumbens.
diffusus, Presl, Bot. Bemerk. 137 = Cytisus decumbens.
Halleri, Presl, l. c. = Cytisus decumbens.
myrtifolius, Presl, l. c.—Hungar.
procumbens, Presl, l. c. = G. procumbens, *Waldst.*

COROZO, Jacq. ex Giseke, Prael. 42 (1792) = **Elaeis**, Jacq. (Palm.).

CORPODETES, Reichb. Consp. 61 (1828) = CARPODETES, Herb. = **Stenomesson**, Herb. (Amaryll.).

CORRAEA, Sm. in Trans. Linn. Soc. iv. (1798) 219 = seq.

CORREA, Andr. Bot. Rep. t. 18 (1798). *RUTACEAE*, Benth. & Hook. f. i. 294.
ANTOMMARCHIA, Colla, Hort. Ripul. App. ii. 345 (1826).
DIDIMERIA, Lindl. in Mitch. Three Exp. Austral. ed. II. ii. 198 (1839).
MAZEUTOXERON, Labill. Voy. ii. 11 (1798).
acmula, *F. Muell. Fragm.* i. 3.—Austral.
alba, *Andr. Bot. Rep.* t. 18.—Austral.
Backhousiana, Hook. Journ. Bot. i. (1834) 253 = speciosa.
Bauerlenii, *F. Muell. in Proc. Linn. Soc. N. S. W.* ix. (1885) 960.—Austral.
bicolor, Paxt. Mag. Bot. ix (1842) 267 = speciosa.
cardinalis, F. Muell. ex Hook. Bot. Mag. t. 4912 = speciosa.
cordifolia, Lindl. in Mitch. Three Exp. Austral. ii. 233 = speciosa.
cotinifolia, Salisb. Parad. Lond. t. 100 = alba.
decumbens, *F. Muell. in Trans. Vict. Inst.* i. (1855) 30.—Austral.
ferruginea, Hook. Comp. Bot. Mag. i. (1835) 276 = Lawrenciana.
glabra, Lindl. in Mitch. Three Exp. Austral. ii. 48 = speciosa.
Latrobeana, F. Muell. in Dietr. Fl. Univ. N. Ser. t. 11 = Lawrenciana,
Lawrenciana, *Hook. Journ. Bot.* i. (1834) 254.—Austral.
leucoclada, Lindl. in Mitch. Three Exp. Austral. ii. 39 = speciosa.
pulchella, Mackay, ex Sweet, Fl. Austral. t. 1 = speciosa.
reflexa, Vent. Jard. Malm. sub t. 13 = speciosa.
revoluta, Vent. l. c. sub t. 13 = speciosa.
rotundifolia, Lindl. in Mitch. Three Exp. Austral. ii. 219 = alba.
rubra, Sm. Exot. Bot. ii. 26 = speciosa.
rufa, Vent. Jard. Malm. t. 13 adnot. = alba.
Schlechtendalii, Behr, in Linnaea, xx. (1847) 630 = speciosa.
speciosa, *Ait. Epit. Hort. Kew.* 366.—Austral.
virens, *Hook. Journ. Bot.* i. (1834) 253.—Austral.
virens, Sm. Exot. Bot. ii. 25. t. 72 = speciosa.
viridiflora, Andr. Bot. Rep. t. 436 = speciosa.

CORREAS, Hoffmgg. Verz Pfl. Nachtr. i. 168 (1824) = **Correa**, Andr. (Rutac.).

CORREIA, Vell. in Roem. Script. 106. t. 6 (1796) = **Gomphia**, Schreb. (Ochnac.).

CORRIGIOLA, Linn. Gen ed. I. 340 (1737). *ILLECEBRACEAE*, Benth. & Hook. f. iii. 17.
POLYGONARIA, Heist. Syst. 6 (1748).
POLYGONIFOLIA, Adans. Fam. ii. 272 (1763).
albella, Forsk. Fl. Aegypt. Arab. 207 = Paronychia arabica.
andina, *Planch. & Triana, in Ann. Sc. Nat.* Sér. IV. xvii. (1862) 146.—N. Granat.
capensis, *Willd. Sp. Pl.* i. 1507.—Afr. austr.
deltoidea, Hook. & Arn. Bot. Beech. Voy. 24 = Monocosmia corrigioloides.
glomeruliflora, *Steud. Nom.* ed. II. i. 423.—Chili.
imbricata, Lapeyr. Fl. Pyr. i. 83; Abr. 169 = littoralis.
latifolia, *C. Gay, Fl. Chil.* ii. 517.—Chili.
littoralis, *Linn. Sp. Pl.* 271.—Europ.
littoralis, Thunb. Prod. Pl. Cap. 55 = capensis.
propinqua, *C. Gay, Fl. Chil.* ii. 517.—Chili.
psammatrophoides, *Baker, in Journ. Linn. Soc.* xx. (1883) 238.—Madag.

CORRIGIOLA :—
repens, Forsk. Fl. Aegypt. Arab. 207 = Polycarpaea fragilis.
squamosa, *Hook. & Arn. in Hook. Bot. Misc.* iii. (1833) 337.—Chili.
telephiifolia, Pourr. in Mém. Acad. Toul. iii. (1788) 316; (Chl. Narb. 20) = littoralis.

CORSIA, Becc. Malesia, i. 238. t. 9 (1878). *BURMANNIACEAE*, Benth. & Hook. f. iii. 460.
ornata, *Becc. l. c.* 239.—N. Guin.

CORTESIA, Cav. Ic. iv. 53. t. 377 (1797). *BORAGINEAE*, Benth & Hook. f. ii. 841.
cuncata, Miers, in Ann. & Mag. Nat. Hist. Ser. IV. ii. (1868) 429 = cuneifolia.
cuneifolia, *Cav. Ic.* iv. 53. t. 377.—Reg. Argent.
cuneiformis, Poir. in Dict. Sc. Nat. x. 668 = cuneifolia.
microphylla, *Miers, in Ann. & Mag. Nat. Hist.* Ser. IV. ii. (1868) 430.—Am. austr.

CORTHUMIA, Reichb. Nom. 201 (1841) = **Pelargonium**, L'Hérit. (Geran.).

CORTHUSA, Reichb. Consp. 128 (1828) = **Cortusa**, Linn. (Primul.).

CORTIA, DC. Prod. iv. 186 (1830). *UMBELLIFERAE*, Benth. & Hook. f. i. 915.
SCHULTZIA, Wall. Cat. n. 589 (1830).
elata, Edgew. in Trans. Linn. Soc. xx. (1851) 55 = Ligusticum elatum.
Hookeri, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 702.—Reg. Himal.
Lindleyi, *DC. Prod.* iv. 187.—Reg. Himal.
vaginata Edgew. in Trans. Linn. Soc. xx. (1851) 55 = Selinum vaginatum.

CORTUSA, Linn. Gen. ed. I. 40 (1737). *PRIMULACEAE*, Benth. & Hook. f. ii. 633.
Gmelini, Linn. Sp. Pl. 144 = Androsace Gmelini.
gradissima, Schur. Enum. Pl. Transs. 556 = Matthioli.
Matthioli, *Linn. Sp. Pl.* 144.—Europ.; As. bor.
pubens, *Schott, Nyman & Kotschy, Analect. Bot.* 17. —Transsylv.
Semenovii, *Herd. in Bull. Soc. Nat. Mosc.* xli. (1868) i. 63.—As. centr.
sibirica, *Andrz. ex Steud. Nom.* ed. II. i. 423.—Sibir. baical.

CORTUSINA, Eckl. & Zeyh. Enum. 77 (1835) = **Pelargonium**, L'Hérit. (Geraniac.).
odoratissima, Eckl. & Zeyh. l. c. = P. odoratissimum.
reniformis, Eckl. & Zeyh. l. c. = P. reniforme.
rubropurpurea, Eckl. & Zeyh. l. c. = P. echinatum.
sidaefolia, Eckl. & Zeyh. l. c. = P. sidoides.
velutina, Eckl. & Zeyh. l. c. = P. velutinum.

CORVINA, Steud. Nom. ed. II. i. 256 (1840) sphalm. = **Muricaria**, Desv. (Crucif.).
prostrata, Steud. l. c. = M. prostrata.

CORVINIA, Stadtm. ex Willem. in Usteri, Ann. Bot. xviii. 28 (1796) = **Nephelium**, Linn. (Sapindac.).
Litschi, Stadm. ex Willem. l. c. = N. Litchi.

CORVISARTIA, Mérat, Fl. Par. ed. I. 328 (1812) = **Inula**, Linn. (Compos.).
caucasica, G. Don, ex Loud. Hort. Brit. 350 = Buphthalmum speciosum.
Helenium, Mérat, Fl. Par. ed. I. 328 = I. Helenium.
indica, Lindl. Bot. Reg. (1842) Misc. 58 = I. Royleana.

CORYANTHES, Hook. Bot. Mag. t. 3102 (1831). *ORCHIDEAE*, Benth. & Hook. f. iii. 549.
CORYTHANTHES, Lem. in Orb. Dict. iv. 259 (1849).
PANSTREPIS, Rafin. Fl. Tellur. iv. 41 (1836).
MELICLIS, Rafin. l. c. ii. 99 (1836).
Albertinae, Karst. Ausw. Neuer Gew. Vencz. 5 = maculata.
Barkerii, Beer, Prakt. Orch. 222, sphalm. = *Parkeri*.
bifiora, *Rodrig. Orch. Nov.* i. 103.—Bras.
Bruchmuelleri, *Reichb. f. in Linnaea*, xli. (1877) 74.—N. Granat.
elegantissima, Mart. in Gard. Chron. (1885) i. 144 = elegantium.
elegantium, *Linden & Reichb. f. in Gard. Chron.* (1868) 1114.—Bras.

CORYANTHES :—
Feildingii, *Lindl. in Journ. Hort. Soc.* iii. (1848) 15. —Hab.?
macrantha, *Hook. Bot. Mag.* t. 3102, in textu.—Venezuela.
macrantha, [Mast. in] Gard. Chron. (1882) i. 592 = elegantium.
maculata, *Hook. Bot. Mag.* t. 3102.—Am. austr.
Parkeri, Endl. in Hartinger. Parad. Vindob. 19. t. 32. f. 1 = speciosa.
pieturata, *Reichb. f. in Bot. Zeit.* xxii. (1864) 332.—Honduras.
punctata, Beer, Prakt. Orch. 222 = maculata.
speciosa, *Hook. Bot. Mag.* t. 3102. in textu.—Bras.
splendens, *Rodrig. Orch. Nov.* i. 103.—Bras.
Sumneriana, *Lindl. in Gard. Chron.* (1856) 452.—Hab.?

CORYBAS, Salisb. Parad. Lond. t. 83 (1805) = **Corysanthes**, R. Br. (1810) (Orchid.).
aconitifolius, Salisb. l. c. = Corysanthes bicalcarata.
fimbriatus, Reichb. f. Beitr. Syst. Pfl. 42 = Corysanthes fimbriata.
fornicatus, Reichb. f. l. c. 67 = Corysanthes fornicata.
macranthus, Reichb. f. l. c. = Corysanthes macrantha.
mucronatus, Reichb. f. l. c. = Corysanthes mucronata.
oblongus, Reichb. f. l. c. = Corysanthes oblonga.
pictus, Reichb. f. l. c. = Corysanthes picta.
pruinosus, Reichb. f. l. c. 43 = Corysanthes fimbriata.
rivularis, Reichb. f. l. c. 67 = Corysanthes rivularis.
rotundifolius, Reichb. f. l. c. = Corysanthes rotundifolia.
trilobus, Reichb. f. l. c. = Corysanthes triloba.
unguiculatus, Reichb. f. l. c. 43 = Corysanthes unguiculata.

CORYCARPUS, Zea, ex Spreng. Syst. i. 123 (1825) = **Diarrhena**, Rafin. (Gramin.).
arundinaceus, Zea, ex Spreng. l. c. = D. americana.

CORYCIUM, Sw. in Vet. Acad. Nya Handl. Stockh. xxi. (1800) 220. *ORCHIDEAE*, Benth. & Hook. f. iii. 633.
bicolorum, *Sw. l. c.* 222.—Afr. austr.
bifidum, *Sond. in Linnaea*, xix. (1847) 111.—Afr. austr.
erispum, *Sw. in Vet. Acad. Nya Handl. Stockh.* xxi. (1800) 222.—Afr. austr.
excisum, *Lindl. Gen. et Sp. Orch.* 368.—Afr. austr.
ligulatum, Reichb. f. in Linnaea, xix. (1847) 375 = bifidum.
microglossum, *Lindl. Gen. et Sp. Orch.* 369.—Afr. austr.
nigrescens, *Sond. in Linnaea*, xix. (1847) 110.—Afr. austr.
orobanchoides, *Sw. in Vet. Acad. Nya Handl. Stockh.* xxi. (1800) 222.—Afr. austr.
vestitum, *Sw. l. c.*—Afr. austr.

CORYDALIS, Vent. Choix, 19 (1803), in nota; DC. Fl. Fr. iv. 636 (1805). *PAPAVERACEAE*, Benth. & Hook. f. i. 55.
BORCKHAUSENIA, Gaertn. Mey. & Scherb. Fl. Wett. iii. 4 (1801).
BULBOCAPNOS, Bernh. in Linnaea, viii. (1833) 469.
CAPNOIDES, Adans. Fam. ii. 431 (1763).
CERATOCAPNOS, Dur. in Parl. Giorn. Bot. Ital. i. 1. (1844) 336.
CRYPTOCERAS, Schott & Kotschy, in Oestr. Bot. Wochenbl. iv. (1854) 121.
CYSTICAPNOS, Boerh. ex Adans. Fam. ii. 431 (1763).
NECKERIA, Scop. Introd. 313 (1777).
PHACOCAPNOS, Bernh. in Linnaea, xii. (1838) 664.
PISTOLOCHIA, Bernh. Syst. Verz. Erf. 57 (1800).
PSEUDOFUMARIA, Medic. Phil. Bot. i. 110 (1789).
SOPHOROCAPNOS, Turcz. in Bull. Soc. Nat. Mosc. xxi. (1848) i. 570.
acaulis, Pers. Syn. ii. 270 = ochroleuca.
adiantifolia, *Hook. f. & Thoms. Fl. Ind.* i. 271.—Reg. Himal.
aduncа, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 29.—China.
albicaulis, *Franch. Pl. David.* i. (1884) 30.—As. or.
alliflora, *Kit. in Linnaea*, xxxii. (1863) 491.—Hungar.
alpestris, C. A. Mey. Verz. Pfl. Cauc. 176 = pauciflora.
alpina, J. Gay, in Ann. Sc. Nat. Ser. I. xxvi. (1832) 252 = fabacea.

CORYDALIS :—
alpina, C. Koch, in Linnaea, xv. (1841) 252 = rutaefolia.
ambigua, *Cham. & Schlecht. in Linnaea*, i. (1826) 558.—Kamtschat.
angustifolia, *DC. Syst.* ii. 120.—Reg. Caucas.; Persia.
annua, Hoffmgg. ex Steud. Nom. ed. II. i. 423 = glauca.
astragalina, Hook. f. & Thoms. Fl. Ind. i. 270 = stricta.
aurea, *Willd. Enum. Hort. Berol.* 740.—Am. bor.
biaurita, Hornem. Hort. Hafn. ii. 668 = Dicentra formosa.
bicalcarata, Haw. ex G. Don, in Loud. Hort. Brit. Suppl. i. 585 = rutaefolia.
Bidwelliae, *S. Wats. in Bot. Calif.* ii. 429.—Calif.
blanda, Schott, in Oestr. Bot. Wochenbl. vii. (1857) 149 = tuberosa.
bracteata, *Pers. Syn.* ii. 269.—Sibiria.
bracteosa, Spreng. Syst. iii. 162 (= *Diclytra bracteosa*). —Am. bor.?
Brandegei, *S. Wats. in Bot. Calif.* ii. 430.—Calif.
breviflora, *DC. Syst.* ii. 127.—Kamtschat.
bulbosa, *DC. Fl. Fr.* iv. 637.—Europ.
bulbosa, Pers. Syn. ii. 269 = tuberosa.
bulbosa, Turcz. ex Steud. Nom. ed. II. i. 423 = Turtschaninovii.
Bungeana, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 62. —Sibir.
Burmanni, *Eckl. & Zeyh. Enum.* 4.—Afr. austr.
canadensis, Goldb. in Edinb. Phil. Journ. vi. (1822) 329 = Dicentra canadensis.
capnoides, Pers. Syn. ii. 270 = lutea.
Caseana, *A. Gray, in Proc. Am. Acad.* x. (1874) 69.—Calif.
cashmeriana, *Royle, Illustr. Bot. Himal.* 69. t. 16. f. 1. —Reg. Himal.
caucasica, Adams, ex Steud. Nom. ed. II. i. 423 = angustifolia.
caucasica, *DC. Syst.* ii. 119.—Reg. Caucas.
caudata, Pers. Syn. ii. 269 = longiflora.
cava, Schweigg. & Kort. Fl. Erlang. ii. 44; Wahlenb. Veg. Helv. 126 [bis] = tuberosa.
chaerophylla, *DC. Prod.* i. 128.—Reg. Himal.
chinensis, Franch. Pl. David. i. 28 = edulis.
clavicula, S. F. Gray, Nat. Arr. Brit. Pl. ii. 700 = seq.
claviculata, *DC. Fl. Fr.* iv. 638.—Europ.
conorhiza, *Ledeb. Fl. Ross.* i. 99.—Reg. Caucas.
cornuta, *Royle, Illustr. Bot. Himal.* 69.—Reg. Himal.
Cracca, *Cham. & Schlecht. in Linnaea*, i. (1826) 567. —Afr. austr.
crassifolia, *Royle, Illustr. Bot. Himal.* 68.—Reg. Himal.
crassissima, Jacquem. Voy. Bot. 12. t. 11 = crassifolia.
crithmifolia, *Royle, Illustr. Bot. Himal.* 68.—Reg. Himal.
crystallina, *Engelm. ex A. Gray, Man. U. St. Bot.* ed. V. 62.—Am. bor.
Cucullaria, Pers. Syn. ii. 269 = Dicentra Cucullaria.
curvisiliqua, *Engelm. ex A. Gray, in Man. Bot. U. St.* ed. V. 62.—Am. bor. occ.
Cusickii, *S. Wats. in Bot. Calif.* ii. 430.—Calif.
dasyptera, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 28.—China.
debilis, Edgew. in Trans. Linn. Soc. xx. (1851) 30 = cornuta.
decipiens, Schott, Nyman & Kotschy, Analect. Bot. 42 = pumila.
decumbens, *Pers. Syn.* ii. 269.—Japon.
densiflora, J. & C. Presl, Delic. Prag. 10 = bulbosa.
depauperata, Schur, in Verh. Siebenb. Ver. Naturw. iv. (1853) 51 = bulbosa.
digitata, Pers. Syn. ii. 269 = bulbosa.
diphylla, Wall. Tent. Fl. Nep. 54 = rutaefolia.
edulis, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 30. —China.
elegans, *Wall. Cat.* n. 1435.—Reg. Himal.
Emanueli, C. A. Mey. Verz. Pfl. Cauc. 176 = pauciflora.
enneaphylla, DC. Fl. Fr. Suppl. 587 = Sarcocapnos enneaphylla.
epithymifolia, Walp. Rep. i. 120, sphalm. = crithmifolia.
Erdelii, Zucc. in Abh. Akad. Muench. iii. (1843) 251 = rutaefolia.
eximia, Link, Enum. Hort. Berol. ii. 218 = Dicentra eximia.
fabacea, *Pers. Syn.* ii. 269.—Europ.

CORYDALIS:—
Falconeri, *Hook. f. & Thoms. in Fl. Brit. Ind.* i. 124 —Reg. Himal.
Fedtschenkoana, *Regel, Pl. Nov. Fedsch.* 3.—As. centr.
fibrosa, *Maxim. in Bull. Phys.-Math. Acad. Pétersb.* xv. (1857) 227, nomen.—Manchur.
filiformis, Royle, Illustr. Bot. Himal. 68 = sibirica.
flabellata, *Edgew. in Trans. Linn. Soc.* xx. (1851) 30.—Reg. Himal.
flaccida, *Hook. f. & Thoms. Fl. Ind.* i. 260.—Reg. Himal.
flavidula, Chapm. Fl. S. U. St. Suppl. 604 = flavula.
flavula, *DC. Prod.* i. 129.—Am. bor.
formosa, Pursh, Fl. Am. Sept. ii. 462 = Dicentra canadensis. eximia.
formosa, Spreng. Syst. iii. 162 = Dicentra formosa.
fumariaefolia, *Maxim. in Prim. Fl. Amur.* 39.—Manchur.
fungosa, Vent. Choix, t. 19 = Adlumia cirrhosa.
gamosepala, Maxim. Prim. Fl. Amur. 38 = bulbosa.
Gebleri, Ledeb. Ind. Hort. Dorpat. (1823) 3 = capnoides.
geraniifolia, Hook. f. & Thoms. Fl. Ind. i. 269 = chaerophylla.
gigantea, *Trautv. & Mey. in Middend. Reise (Fl. Ochot.* 13).—Manchur.
glauca, *Pursh, Fl. Am. Sept.* ii. 463.—Am. bor.
glaucescens, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) i. 253.—As. temp.
Gortschakovii, *Schrenk, Enum. Pl. Nov.* i. 100.—Reg. Himal.; Soongar.
Govaniana, *Wall. Tent. Fl. Nep.* 55.—Reg. Himal.
gracilipes, S. Moore, in Journ. Bot. xiii. (1875) 226 = decumbens.
gracilis, *Ledeb. Fl. Ross.* i. 101.—Reg. Caucas.
Griffithii, Boiss. Diagn. Ser. II. i. 15 (*Griffithsii*) = rutaefolia.
Halleri, Willd. Enum. Hort. Berol. 740 = bulbosa.
Hamiltoniana, G. Don, Gen. Syst. i. 143 = rutaefolia.
heterocarpa, *Ball, in Journ. Linn. Soc.* xvi. (1877) 314.—Afr. bor. occ.
heterocarpa, Sieb. & Zucc. in Abh. Akad. Muench. iv. (1843) ii. 173 = pallida.
Hoffmeisteri, Klotzsch, in Bot. Ergeb. Waldem. Reise, 129 = meifolia.
Impatiens, Fisch. in DC. Syst. ii. 124 = sibirica.
incisa, *Pers. Syn.* ii. 269.—Japon.
inconspicua, *Bunge, ex Ledeb. Fl. Ross.* i. 104.—Sibiria.
intermedia, Mérat, Fl. Par. 272 = fabacea.
juncea, *Wall. Tent. Fl. Nep.* 54. t. 42.—Reg. Himal.
Kareliniana, *Pritz. ex Walp. Rep.* ii. 750.—As. centr.
kaschmiriana, Auct. = cashmeriana.
kashgarica, *Rupr. Sert. Tiansch.* 38.—As. centr.
Kolpakowskiana, *Regel, in Act. Hort. Petrop.* v. (1877) 633.—Turkestan.
lachenaliaeflora, Fisch. ex DC. Syst. ii. 111 = Dicentra lachenaliaeflora.
laevigata, E. Mey. ex Bernh. in Linnaea, xii. (1838) 664 = C. Cracca.
latiflora, *Hook. f. & Thoms. Fl. Ind.* i. 270.—Reg. Himal.
laxa, *Franch. & Sav. Enum. Pl. Jap.* ii. 274.—Japon.
laxa, *Fries. Summa Veg. Scand.* 145.—Fennia.
Ledebouriana, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* (1841) 377.—Sibiria.
leptocarpa, *Hook. f. & Thoms. Fl. Ind.* i. 260.—Reg. Himal.
libanotica, Hochst. in Lorent, Wanderungen, 339 = rutaefolia.
linarioides, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 27.—China.
linearibloba, Sieb. & Zucc. in Abh. Akad. Muench. iv. ii. (1843) 174 = bulbosa.
Lobelii, Tausch, ex Hoppe, in Flora, xxii. (1839) 379 = pumila.
longiflora, *Pers Syn.* ii. 269.—Reg. Casp.; Sibir.
longipes, DC. Prod. i. 128 = sibirica.
longipes, D. Don, Prod. Fl. Nep. 198 = rutaefolia.
lutea, *DC. Fl. Fr.* iv. 638.—Europ.
Maackii, *Rupr. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 71.—Manchur.
macrocentra, *Regel, l. c.* viii. (1884) 694.—As. centr.
macrophylla, *Nutt. in Torr. & Gray, Fl. N. Am.* i. 69. —Am. bor. occ.
macrosepala, *Ssred. ex Trautv. in Act. Hort. Petrop.* viii. (1883) 71.—Reg. Caucas.
Marschalliana, *Pers. Syn.* ii. 269.—Rossia; As. Min.

CORYDALIS:—
meifolia, *Wall. Tent. Fl. Nep.* 55.—Reg. Himal.
melanochlora, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 26.—China.
micropoda, *Franch. Pl. Davidian.* i. (1884) 29.—As. or.
montana, Engelm. ex A. Gray, Man. U. St. Bot. ed. V. 62 = aurea.
Moorcroftiana, *Wall. Cat.* n. 1432.—Reg. Himal.
nana, *Royle, Illustr. Bot. Himal.* 68.—Reg. Himal.
nivalis, Boiss. & Huet, in Diagn. Ser. II. v. 16 = pauciflora.
nobilis, *Pers. Syn.* ii. 269.—Sibir.
nudicaulis, *Regel, in Act. Hort. Petrop.* viii. (1884) 695.—As. centr.
ochotensis, *Turcz. in Bull. Soc. Nat. Mosc.* (1840) 62. —Sibir.
ochroleuca, *Koch, in Flora*, xiv. (1831) 708.—Italia.
ophiocarpa, *Hook. f. & Thoms. Fl. Ind.* i. 259.—Reg. Himal.
oppositifolia, DC. Syst. ii. 114 = rutaefolia.
orthoceras, *Sieb. & Zucc. in Abh. Akad. Muench.* iv. (1843) ii. 175.—Japon.
paeoniaefolia, *Pers. Syn.* ii. 269.—Sibir.; Am. bor. occ.
pallida, *Pers. l. c.* 270.—China; Japon.
paniculigera, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 3.—As. centr.
parnassica, *Orph. & Heldr. in Boiss. Diagn.* Ser. II. vi. 9.—Graecia.
pauciflora, Edgew. in Trans. Linn. Soc. xx. (1851) 30 = rutaefolia.
pauciflora, *Pers. Syn.* ii. 269.—As. bor.; Am. bor. occ.
persica, *Cham. & Schlecht. in Linnaea*, i. (1826) 567. —Persia.
physocarpa, Cambess. in Jacquem. Voy. Bot. 13. t. 12 = crassifolia.
polygalina, *Hook. f. & Thoms. Fl. Ind.* i. 263.—Reg. Himal.
pruinosa, *E. Mey. ex Bernh. in Linnaea*, xii. (1838) 604, in syn.—Afr. austr.
pseudo-cava, Pantoč. in Verh. Ver. Natur- u. Heilk. Presb. ii. (1871–72) 88 = tuberosa.
pseudo-pumila, Schur, Enum. Pl. Transs. 37 = pumila.
pulchella, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 151.—Afghan.
pumila, *Koch, Syn. Fl. Germ.* ed. II. 34.—Europ.
racemosa, Bunge, Enum. Pl. Chin. Bor. 5 = Bungeana.
racemosa, *Pers. Syn.* ii. 270.—Japon.
Raddeana, *Regel, in Bull. Soc. Nat. Mosc.* xxxiv. (1861) ii. 143.—Manchur.
ramosa, *Wall. Cat.* n. 1434.—Reg. Himal.
remota, Fisch. ex Maxim. Prim. Fl. Amur. 37 = bulbosa.
rosea, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 28. —China.
rosea, *Zeyh. ex Steud. Nom.* ed. II. i. 424.—Afr. austr.
rupestris, *Kotschy, ex Boiss. Diagn.* Ser. I. vi. 8.—Persia.
rutacea, *T. M. Fr. in Bot. Notiser* (1854) 39.—Sibir.?
rutaefolia, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 13 = persica.
rutaefolia, *DC. Syst.* ii. 115.—As. Min.; Persia; Reg. Himal.
scandens, Spreng. Syst. iv. Cur. Post. 265 = Dicentra spectabilis.
Schelesnowiana, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 4.—As. centr.
Scouleri, *Hook. Fl. Bor. Am.* i. 36. t. 14.—Am. bor. occ.
Semenowii, *Regel & Herd. in Bull. Soc. Nat. Mosc.* xxxvii. (1864) i. 407.—Sibir.
sempervirens, Pers. Syn. ii. 269 = glauca.
senanensis, *Franch. & Sav. Enum. Pl. Jap.* ii. 273.—Japon.
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) i. 252.—As. centr.
Sheareri, *S. Moore, in Journ. Bot.* xiii. (1875) 225.—China.
sibirica, *Pers. Syn.* ii. 270.—Sibir.; Reg. Himal.
solida, [Sw. in] Svensk Bot. viii. 531 (1819) = bulbosa.
speciosa, Maxim. Prim. Fl. Amur. 39 = pallida.
spectabilis, Pers. Syn. ii. 269 = Dicentra spectabilis.
streptocarpa, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 30.—China.
stricta, *Steph. ex DC. Syst.* ii. 123.—Sibir.; Reg. Himal.
Stummeri, *Pantoč. in Magyar Növént. Lapok*, v. (1881) 150.—Hungar.

CORYDALIS:—
suaveolens, *Hance, in Journ. Bot.* xviii. (1880) 258.—China.
tenella, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 143.—As. centr.
tenella, *Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb.* ii. (1837) 313.—Reg. Caucas.
tenuifolia, Pursh, Fl. Am. Sept. ii. 462 = Dicentra lachenaliaeflora.
tenuis, Schott, Nyman & Kotschy, Analect. Bot. 43 = bulbosa.
thalictrifolia, *Jameson, ex Regel, in Bull. Soc. Nat. Mosc.* xxxiv. (1861) ii. 148.—China.
tibetica, *Hook. f. & Thoms. Fl. Ind.* i. 265.—Reg. Himal.
trachycarpa, *Maxim. in Bull. Acad. Pétersb.* xxiv. (1878) 27.—China.
triternata, Zucc. in Abh. Akad. Muench. iii. (1843) 251 = bulbosa.
tuberosa, *DC. Fl. Fr.* iv. 637.—Europ.
Turtschaninovii, *Bess. in Flora*, xvii. (1834) i. *Beibl.* 6. —Rossia.
uniflora, Nym. Syll. 185 = rutaefolia.
uralensis, *Fisch. ex DC. Syst.* ii. 127.—Mont. Ural.
vaginans, *Royle, Illustr. Bot. Himal.* 68.—Reg. Himal.
Vernyi, *Franch. & Sav. Enum. Pl. Jap.* ii. 273.—Japon.
verticillaris, DC. Syst. ii. 114 = rutaefolia.
vesicaria, *Pers. Syn.* ii. 269.—Afr. austr.
Wilfordi, Regel, in Bull. Soc. Nat. Mosc. xxxiv. (1861) ii. 148 = pallida.

CORYDALLIS, Aschers. Fl. Brandenb. 28 (1864) = praec.

CORYDANDRA, Reichb. Nom. 53 (1841) = **Galeandra**, Lindl. (Orchid.).

CORYLOPSIS, Sieb. & Zucc. Fl. Jap. i. 45. tt. 19, 20 (1835). *HAMAMELIDEAE*, Benth. & Hook. f. i. 667.
glabrescens, *Franch. & Sav. Enum. Pl. Jap.* ii. 367.—Japon.
himalayana, *Griff. in Journ. As. Soc. Beng.* xxiii. (1854) 641.—Reg. Himal.
Kesakii, *Sieb. & Zucc. Fl. Jap.* i. 49.—Japon.
multiflora, *Hance, in Ann. Sc. Nat.* Sér. IV. xv. (1861) 224.—China.
pauciflora, *Sieb. & Zucc. Fl. Jap.* i. 48. t. 20.—Japon.
spicata, *Sieb. & Zucc. l. c.* 45. t. 19.—Japon.

CORYLUS, [Tourn.] Linn. Syst. ed. I (1735). *CUPULIFERAE*, Benth. & Hook. f. iii. 406.
alba, Ait. ex Steud. Nom. ed. II. i 421 = C. Avellana.
americana, Siebold, in Verh. Bat. Gen. xii. (1830) 26 = heterophylla.
americana, *Walt. Fl. Carol.* 236.—Am. bor.
americana, Willd. Sp. Pl iv. 471 = humilis.
arborea, Hort. ex Steud. Nom. ed. II. i. 424 = C. Avellana.
arborescens, Gaertn. Mey. & Scherb. Fl. Wett. iii. 376 = tubulosa.
arborescens, Muench. Hausv. iii. 826 = C. Colurna.
atropurpurea, Hort. ex A. DC. in DC. Prod. xvi. ii. 132 = tubulosa.
Avellana, *Linn. Sp. Pl.* 998.—Europ.; As. Min.
Avellana, Thunb. Fl. Jap. 160 = heterophylla.
bizantina, Desf. Tabl. ed. II. 245 = C. Colurna.
californica, Hort. ex C. Koch, Dendrol. ii. ii. 15 = rostrata.
Colurna, *Linn. Sp. Pl.* 999.—Europ. or.; As. Min.; Himal.
cornuta, Duroi, ex Steud. Nom. ed. I. 229 = rostrata.
Davidiana, *Baill. Hist. Pl.* vi. 224.—China.
ferox, *Wall. Cat.* n. 2797.—Reg. Himal.
filicifolia, Hort. ex A. DC. in DC. Prod. xvi. ii. 131 = C. Avellana.
grandis, [Dryand. in] Ait. Hort. Kew. ed. I. iii. 363 = C. Avellana.
Hasibani, *Siebold, Kruidk. Naaml.* 27.—Japon.
heterophylla, Fisch. ex Bess. in Flora, xvii. (1834) i. Beibl. 24 = C. Avellana.
humilis, Willd. Enum. Hort. Berol. 983 = americana.
intermedia, Fingerh. in Linnaea, iv. (1829) 385 = C. Colurna.
Jacquemontii, Decne. in Jacquem. Voy. Bot. 160. t. 169 = C. Colurna.
lacera, Wall. Cat. n. 2798 = C. Colurna.

CORYLUS:—
laciniata, Hort. ex A. DC. in DC. Prod. xvi. II. 131 = C. Avellana.
mandshurica, Maxim. in Bull. Phys.-Math. Acad. Pétr sb. xv. (1856) 137 = rostrata.
maxima, Mill. Gard. Dict. ed. VIII. n. 2 = C. Avellana.
mexicana, Hort. ex C. Koch, Dendrol. ii. II. 15 = rostrata.
mongolica, Hort. ex C. Koch, l. c. 13 = C. Avellana.
ovata, Lam. ex Steud. Nom. ed. II. i. 424 = C. Avellana.
pontica, Dochmahl, Fuehr. Obstkunde, iv. 38 = C. Avellana.
pontica, C. Koch, in Linnaea, xxii. (1849) 329 = C. Colurna.
pumila, Lodd. ex Loud. Hort. Brit. Suppl. iii. 523 = C. Avellana.
purpurea, Hort. ex A. DC. in DC. Prod. xvi. II. 132 = tubulosa.
quercifolia, Hort. ex A. DC. l. c. 131 = C. Avellana.
rostrata, *Ait. Hort. Kew.* iii. 364.—Am. bor.
Serényiana, Pluk. in Oestr. Bot. Wochenbl. ii. (1852) 394 = C. Avellana.
Sieboldiana, Blume, Mus. Bot. Lugd. Bat. i. 310 = heterophylla.
sylvestris, Salisb. Prod. 392 = C. Avellana.
tetraphylla, Ledeb. in Denkschr. Baier. Bot. Ges. iii. (1841) II. 58. sphalm. = C. Avellana.
Thunbergii, Hort. ex C. Koch, Dendrol. ii. II. 13 = heterophylla.
tiliacea, Jacquem. Voy. Bot. 160 = C. Colurna.
tubulosa, *Willd. Sp. Pl.* iv. 470.—Europ. austr.; As. Min.
Turtschaninovii, Steud. Nom. ed. II. i. 425, sphalm. = Corydalis Turtschaninovii.

CORYMBIS, Thou. Orch. Iles Afr. t. 37, 38 (1822). *ORCHIDEAE*, Benth. & Hook. f. iii. 591.
CHLOIDIA, Lindl. Gen. et Sp. Orch. 484 (1840).
CORYMBORCHIS, Thou. ex Blume, Orch. Arch. Ind. 125 (1855).
HYSTERIA, Reinw. in Blume, Cat. Gew. Buitenz. 99 (1823).
MACROSTYLIS, Breda Orch. Kuhl. et Hass. (1827 ?).
RHYNCHANTHERA, Blume, Bijdr. Tabell. 78 (1826).
TOMOTRIS, Rafin. Fl. Tellur. ii. 89 (1836).
angustifolia, *Miq Fl. Ind. Bat. Suppl.* 617.—Sumatra.
corymbosa, *Thou. Orch. Iles Afr.* tt 37, 38.—Madag.
disticha, Lindl. Fol. Orchid. Corymbis = corymbosa.
flava, *Hemsl. Biol. Centr. Am. Bot.* iii. 297.—Mexic.
galipaensis, *Reichb. f. in Flora*, xlviii. (1865) 184.—Am. trop.
Thouarsii, Reichb. f. in Bot. Zeit. vii. (1849) 886 = corymbosa.
veratrifolia, *Reichb. f. in Flora*, xlviii. (1865) 184.—Ind. or.; Malaya.
Wewitschii, *Reichb. f. l. c.* 183.—Afr. occ.

CORYMBIUM, Linn. Coroll. Gen. 14 (1737). *COMPOSITAE*, Benth. & Hook. f. ii. 234.
CONTARENA, Adans. Fam. ii. 120 (1763).
africanum, *Linn. Sp. Pl.* 928.—Afr. austr.
congestum, *E. Mey. ex Steud. Nom.* ed. II. i. 425.—Afr. austr.
cymosum, *E. Mey. ex Steud. l. c.*—Afr. austr.
filiforme, *Linn. f. Suppl.* 392.—Afr. austr.
glabrum, *Linn. Syst.* ed. XII. 582.—Afr. austr.
glabrum, Thunb. Prod. Fl. Cap. 170 = filiforme ?
gramineum, Burm. f. Fl. Cap. Prod. 29 = filiforme.
gramineum, Lam. Encyc. ii. 129 = filiforme, glabrum.
hirsutum, Eckl. ex DC. Prod. v. 89 = villosum.
hirsutum, Thunb. Fl. Cap. 729, 730 = seq.
hirtum, *Thunb. Prod. Pl. Cap.* 170.—Afr. austr.
latifolium, *Harv. in Harv. & Sond. Fl. Cap.* iii. 55.—Afr. austr.
luteum, *E. Mey. ex Steud. Nom.* ed. II. i. 425.—Afr. austr.
nervosum, *Thunb. Prod. Pl. Cap.* 170.—Afr. austr.
scabridum, Berg. Descr. Pl. Cap. 341 = africanum.
scabrum, Linn. Mant. i. 120 = praec.
villosum, *Linn. f. Suppl.* 392.—Afr. austr.

CORYMBORCHIS, Thou. ex Blume, Orch. Ind. Arch. 125 (1855) = **Corymbis**, Thou. (Orchid.).
assamica, Blume, Orchid. Archip. Ind. 126. t. 43 = Corymbis veratrifolia.
Thouarsii, Blume, l. c. 126. t. 44. f. 1. a-c = Corymbis corymbosa.
veratrifolia, Blume, l. c. 125. t. 42 = Corymbis veratrifolia.

CORYMBULA, Rafin. New Fl. Am. iv. 88 (1836) = **Polygala**, Linn.

CORYNAEA, Hook. f. in Trans. Linn. Soc. xxii. (1856) 31. t. 13, 14. *BALANOPHOREAE*, Benth. & Hook. f. iii. 237.
crassa, *Hook. f. l. c.* 31, 54. t. 13.—N. Granat.
Purdiei, *Hook. f. l. c.* 31, 55.—N. Granat.
sphaerica, *Hook. f. l. c.* 31, 54. t. 14.—N. Granat.
Sprucei, *Eichl. in DC. Prod.* xvii. 137.—N. Granat.

CORYNANDRA, Schrad. ex Spreng. Syst. iv. Cur. Post. 201 (1827) = **Cleome**, Linn. (Cappar.).
pulchella, Schrad. ex Spreng. l. c. 204 = Cleome Chelidonii.

CORYNANTHE, Welw. in Trans. Linn. Soc. xxvii. (1869) 37. t. 14. *RUBIACEAE*, Benth. & Hook. f. ii. 36.
paniculata, *Welw. l. c.*—Afr. trop.

CORYNANTHELIUM, Kunze, in Linnaea, xx. (1847) 19 = **Mikania**, Willd. (Compos.).
Moronoa, Kunze, l. c.—Bras.

CORYNANTHES, Schlecht. in Bot. Zeit. vi. (1848) 65 = **Coryanthes**, Hook. (Orchid.).

CORYNELIA, Reichb. Nom. 148 (1841) = seq.

CORYNELLA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 93. *LEGUMINOSAE*, Benth. & Hook. f. i. 500.
CORYNITIS, Spreng. Syst. iv. Cur. Post. 263 (1827).
TOXOTROPIS, Turcz. in Bull. Soc. Nat. Mosc. xix. (1846) 506.
gracilis, *Griseb. Cat. Pl. Cub.* 71.—Cuba.
immarginata, *Wright, in Sauv. Fl. Cub.* 26.—Cuba.
paucifolia, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 93.—Ins. S. Doming.
polyantha, *DC. l. c.*—Ind. occ.

CORYNELOBOS, R. Roem. ex Willk. in Linnaea, xxv. (1852) 7 = **Brassica**, Linn. (Crucif.).
boeticus, R. Roem. ex Willk. l. c. = Brassica coryneloba.

CORYNEPHORUS, Beauv. Agrost. 90. t. 18. f. 2 (1812). *GRAMINEAE*, Benth. & Hook. f. iii. 1157.
WEINGAERTNERIA, Bernh. Syst. Verz. Erf. 23, 51 (1800), nomen prius.
aetnensis, Schult. Mant. iii. 641 = articulatus.
articulatus, *Beauv. Agrost.* 90.—Europ.; As. Min.
canescens, *Beauv. l. c.* t. 18. f. 2.—Europ. austr.
fasciculatus, *Boiss. & Reut. Pugill. Pl. Nov.* 123.—Algieria.
macrantherus, Boiss. & Reut. l. c. 124 = fasciculatus.

CORYNITIS, Spreng. Syst. iv. Cur. Post. 263, 280 (1827) = **Corynella**, DC. (Legum.).
domingensis, Spreng. l. c. 280 = Corynella paucifolia.
polyantha, Spreng. l. c. = Corynella polyantha.

CORYNOCARPUS, Forst. Char. Gen. 31. t. 16 (1776). *ANACARDIACEAE*, Benth. & Hook. f. i. 425.
laevigata, *Forst. l. c.* 32.—N. Zel.

CORYNOPHALLUS, Schott, in Oestr. Bot. Wochenbl. (1857) 389 = **Amorphophallus**, Blume (Aroid.).
Afzelii, Schott, l. c. (= *Hydrosme leonensis*).—Afr. trop.
leonensis, Engl. in DC. Monog. Phan. ii. 326 = A. leonensis.

CORYNOPHORUS, Kunth, in Mém. Mus. Par. ii. (1815) 72 = **Corynephorus**, Beauv. (Gramin.)

CORYNOSICYOS, F. Muell. Sel. Pl. Indust. Cult. 59 (1876) = **Cucumeropsis**, Naud. (Cucurbit.).
edulis, F. Muell. l. c. = Cucum. edulis.

CORYNOSTIGMA, Presl, Epim. Bot. 218 (1849) = **Jussieua**, Linn. (Onagrar.).
jussiaeoides, Presl, l. c. = J. nervosa.

CORYNOSTYLIS, Mart. Nov. Gen. et Sp. i. 25. t. 17, 18 (1824). *VIOLARIEAE*, Benth. & Hook. f. i. 116.
CALYPTRION, Ging. in DC. Prod. i. 288 (1824).
albiflora, Linden, ex T. Moore, in Flor. st (1872) 9.—Bras.
Benthami, Walp. Rep. i. 223 = C. Hybanthus.
Berterii, *Spreng Syst* i. 805.—Mexic.
carthagenensis, Karst. Fl. Columb. ii. 53. t. 127. f. 9-14 = C. Hybanthus.
diandra, *Spreng. Syst.* i, 805.—Am. austr.
guayanensis, Karst. Fl. Columb. ii. 53. t. 127. f. 1-8 = C. Hybanthus.
Hyhanthus. *Mart. & Zucc. Nov. Gen. et Sp.* i. 26. t. 17.—Mexic.
Loeflingii, *Spreng. Syst.* i. 805.—Am. austr.

CORYNOTHECA, F. Muell. Fragm. vii. 68. (1870). *LILIACEAE*, Benth. & Hook. f. iii. 792.
acanthoclada, *F. Muell. l. c.*—Austral.
dichotoma, *F. Muell. l. c.*—Austral.
lateriflora, *F. Muell. l. c.*—Austral.

CORYNULA, Hook. f. Ic. Pl. t. 1123 (1872). *RUBIACEAE*, Benth. & Hook. f. ii. 138.
pilosa, *Hook. f. l. c.*—Am. austr.

CORYPHA, Linn. Mus. Cliff. 11 (1736); Gen. ed. I. 354 (1737). *PALMAE*, Benth. & Hook. f. iii. 922.
BESSIA, Rafin. Sylva Tellur. 13 (1838).
CODDA-PANA, Adans. Fam. ii. 25 (1763).
DENDREMA, Rafin. Sylva Tellur. 14 (1838).
GEMBANGA, Blume, in Flora, viii. (1825) 580, 678.
TALIERA, Mart. Progr. Palm. Fam. et Gen. 10 (1824).
africana, Lour. Fl. Cochinch. 213 = Hyphaene coriacea.
australis, R. Br. Prod. 267 = Livistona australis.
cerifera, Arruda, in Koster, Trav. Braz. 494; Mart. Hist. Nat. Palm ii. 56. t. 49 = Copernicia cerifera.
dulcis, H. B. & K. Nov. Gen. et Sp. i. 300 = Brahea dulcis.
elata, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 176.—Ind. or.
Gebang, *Mart. Hist. Nat. Palm.* iii. 233.—Bras.
Gebanga, *Blume, Rumphia*, ii. 59. t. 97, 98, 105.—Malaya.
glaucescens, Lodd. ex Loud. Hort. Brit. 125 = Sabal glaucescens?
guineensis, Linn. Mant. i. 137 = umbraculifera?
Hystrix, Desf. Tabl. ed. I. 19 = Rhapidophyllum Hystrix.
Licuala, Lam. Encyc. ii. 131 = Licuala Rumphii.
macropoda, *Kurz, ex Linden, Cat.* n. 87 (1871).—Ins. Andam.
maritima, H. B. & K. Nov. Gen. et Sp. i. 298 = Copernicia maritima.
minor, *Blanco, Fl. Filip.* ed. I. 229.—Ins. Philipp.
minor, Jacq. Hort. Vindob. iii. 8 = Sabal Adansoni.
Miraguama, H. B. & K. Nov. Gen. et Sp. i. 298 = Thrinax Miraguama.
nana, H. B. & K. l. c. 299 = Copernicia nana.
palmacea, Steud. Nom. ed. I. 229, 834 = Thrinax parviflora.
Palmetto, Walt. Fl. Carol. 119 = Sabal Palmetto.
pilearia, Lour. Fl. Cochinch. 213 = Licuala pilearia.
pumila, Walt. Fl. Carol. 119 = Sabal Adansoni.
Pumos, H. B. & K. Nov. Gen. et Sp. i. 298 = Copernicia Pumos.
repens, Bertram, Trav. 61 = Rhapidophyllum Hystrix.
rotundifolia, Lam. Encyc. ii. 131 = Livistona rotundifolia.
Saribus, Lour. Fl. Cochinch. 212 = Livistona cochinchinensis.
sylvestris, *Mart. Hist. Nat. Palm.* iii. 233.—Ins. Molucc.
Taliera, *Roxb. Hort. Beng.* 25; *Fl. Ind.* ii. 174.—Ind. or.
tectorum, H. B. & K. Nov. Gen. et Sp. i. 299 = Copernicia tectorum.
thebaica, Linn. Sp. Pl. 1187 = Hyphaene cucifera.
umbraculifera, Jacq. Fragm. Bot. 12 = Sabal Blackburniana.
umbraculifera, *Linn. Sp. Pl.* 1178.—Afr. trop.; Ind. or.
Utan, Lam. Encyc. ii. 131 = sylvestris.

CORYPHANTHA, Lem. Cact. 32 (1868) = **Mammillaria**, Haw (Cact.).
acanthostephes, Lem. l. c. 35 = M. acanthostepha.
ancistracantha, Lem. l. c. 34 = M. raphidacantha.

CORYPHANTHA :—
aulacothele, Lem. Cact. 34 = Mammillaria aulacothele.
brevimamma, Lem. ex Foerst. Handb. Cact. ed. II. 394 = M. brevimamma.
calcarata, Lem. Cact. 35 = M. calcarata.
Clava, Lem. l. c. 34 = M. Clava.
conspicua, Lem. l. c.—Hab. ?
cornifera, Lem. l. c. 35 = M. cornifera.
daimonoceras, Lem. l. c. = M. scolymoides.
elephantidens, Lem. l. c. = M. elephantidens.
Engelmanni, Lem. l. c. 34.—Hab. ?
erecta, Lem. l. c. = M. erecta.
exsudans, Lem. ex Foerst. Handb. Cact. ed. II. 395 = M. exudans.
glandulifera, Lem. Cact. 34 = M. glandulifera ?
heterophylla, Lem. l. c.—Hab. ?
Hookeri, Lem. l. c.—Hab. ?
impexicoma, Lem. ex Foerst. Handb. Cact. ed. II. 414 = M. cornifera.
Lehmanni, Lem. Cact. 34 = M. Lehmanni.
loricata, Lem. l. c. 35 = M. loricata.
macromeris, Lem. l. c. = M. macromeris.
Nuttallii, Engelm. ex Foerst. Handb. Cact. ed. II. 407 = M. Nuttallii.
Ottonis, Lem. Cact. 34 = M. Ottonis.
pycnacantha, Lem. l. c. 35 = M. pycnacantha.
raphidacantha, Lem. l. c. 34 = M. raphidacantha.
Scheerii, Lem l. c. 35 = M. Scheerii.
Schlechtendalii, Lem. l. c. 34 = M. Schlechtendalii.
sublanata, Lem. l. c. 35 —Hab. ?
sulcolanata, Lem. l. c. = M. sulcolanata.

CORYSADENIA, Griff. in Proc. Linn. Soc. i. (1846) 281 = **Illigera**, Blume (Combret.).

CORYSANTHERA, Wall. Cat. n. 6411 (1832) = **Rhynchotechum**, Blume (Gesnerac.).
elliptica, Wall. l. c. = R. ellipticum.
vestita, Griff. Itin. Notes, 124 = R. vestitum.

CORYSANTHES, R. Br. Prod. 328 (1810). *ORCHIDEAE*, Benth. & Hook. f. iii. 609.
CALCEARIA, Blume, Bijdr. 417. t. 33 (1825).
CORYBAS, Salisb. Parad. Lond. t. 83 (1805), nomen prius.
NEMATOCERAS, Hook. f. Fl. N. Zel. i. 249. t. 57 (1853).
Betchei, *F. Muell. in Wing, South Sc. Record*, i. (1881) 171.—Ins. Samoa.
bicalcarata. *R. Br. Prod.* 328.—Austral.
callosa, *Blume, Orch. Archip. Ind.* 173. t. 63.—Malaya.
Cheesemanii, *Hook. f. ex T. Kirk, in Trans. N. Z. Inst.* iii. (1871) 180.—N. Zel.
diemenica, Lindl. Gen. et Sp. Orch. 393 = fimbriata.
fimbriata, *R. Br. Prod.* 328.— Austral.
fornicata, *Lindl. Gen. et Sp. Orch.* 394.—Malaya.
hypogea, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 336.—N. Zel.
limbata, *Hook. f. Bot. Mag.* t. 5357.—Java.
macrantha, *Hook. f. Handb. N. Zeal. Fl.* 266.—N. Zel.
maculata, Heynh. Nom. i. 222 = Coryanthes maculata.
mucronata, *Blume, Orch. Archip. Ind.* 175.—Malaya.
oblonga, *Hook f. Handb. N. Zeal. Fl.* 266.—N. Zel.
papillosa, *Colenso, in Trans. N. Z. Inst.* xvi. (1884) 337.—N. Zel.
picta, *Lindl. Gen. et Sp. Orch.* 394.—Malaya.
pruinosa, R. Cunn. in N. S. Wales Mag. (1833) n. 1 = fimbriata.
rivularis, *Hook. f. Handb. N. Zeal. Fl.* 266.—N. Zel.
rotundifolia, *Hook. f. l. c.*—N. Zel.
triloba, *Hook. f. l. c.* 265.—N. Zel.
undulata, *R. Cunn. in N. S. Wales Mag.* (1833), nomen.—Austral.
unguiculata, *R. Br. Prod.* 328.—Austral.

CORYTHACANTHUS, Nees in Lindl. Introd. Nat. Syst. ed II. 444 (1836) = **Clistax**, Mart. (Acanthac.).
speciosus, Nees, l. c.—Bras.

CORYTHANTHES, Lem in Orb. Dict. iv. 259 (1849) = **Coryanthes**, Hook. (Orchid.).

CORYTHOLOBIUM, Mart. ex Benth. in Ann. Wien. Mus. ii (1838) 93 = **Securidaca**, Linn. (Polygal.).
macrophyllum, Benth. l. c. = S. Corytholobium.

CORYTHOLOMA, Decne. in Rev. Hortic. Sér. III. ii. (1848) 466 = **Gesneria**, Mart.
aggregatum, Decne. l. c. 467 = G. aggregata.
caracasanum, Decne. l. c. = G. caracasana.
chelonoides, Regel, Gartenfl. vii. (1858) 374 = G. chelonoides.
ellipticum, Decne. in Rev. Hortic. Sér. III. ii. (1848) 466 = G. elliptica.
fragile, Decne. l. c. 467 = G. Sceptrum.
gracile, Brongn. ex Decne. l. c. 466.—Hab. ?
igneum, Decne. l. c. 467 = G. Sceptrum.
Lindleyi, Decne. l. c. = G. Lindleyi.
Marchii, Decne. l. c. = G. Marchii.
melittifolium, Brongn. ex Decne. l. c. = G. melittifolia.
Merckii, Decne. ex Regel, Gartenfl. iv. (1855) 247 = G. Merckii.
pendulinum, Decne. in Rev. Hortic. Sér. III. ii. (1848) 467 = G. pendulina.
rupicolum, Decne. l. c. = G. rupicola.
rutilum, Decne. l. c. = G. rutila.
Sceptrum, Decne. l. c. = G. Sceptrum.
strictum, Decne. l. c. 466 = G. stricta.
tubiflorum, Decne. l. c. 467 = Isoloma tubiflorum.

CORYTOPLECTUS, Oerst. in Vidensk. Selsk. Skr. V. v. (1861) 119 = **Alloplectus**, Mart. (Gesnerac.).
capitatus, Oerst. l. c. 120 = A. capitatus ?

CORYZADENIA, Griff. Notul. iv. 356 (1854) = **Illigera**, Blume (Combretac.).
trifoliata, Griff. l. c. = I. Coryzadenia.

COSARIA, J. F. Gmel. Syst. 22, 71 (1791) = KOSARIA, Forsk. = **Dorstenia**, Linn. (Urticac.).
Forskălii, J. F. Gmel. l. c. = D. radiata.

COSBAEA, Lem. Illustr. Hortic. ii. (1855) Misc. 71 = **Schisandra**, Michx. (Magnoliac.)
coccinea, Lem. l. c. = S. Hanceana.

COSCINIUM, Colebr. in Trans. Linn. Soc. xiii. (1822) 51. *MENISPERMACEAE*, Benth. & Hook. f. i. 35
PEREIRIA, Lindl. Fl. Med. 370 (1838).
Blumeanum, *Miers, in Ann. & Mag. Nat. Hist.* Ser. II. vii. (1851) 37.—Ind. or.
fenestratum, *Colebr. in Trans. Linn. Soc.* xiii. (1822) 65.—Ind. or.
Maingayi, *Pierre, Fl. For. Cochin.* sub t. 112 (err. Mangayi).—Malaya.
usitatum, *Pierre, l. c.*—Cambodia.
Wallichianum, Miers, in Ann. & Mag. Nat. Hist. Ser. II. vii. (1851) 37 = fenestratum.
Wightianum, Miers, l. c. = fenestratum.

COSMANTHUS, Nolte, ex A. DC. Prod. ix. 296 (1838) = **Phacelia**, Juss. (Hydrophyll.).
fimbriatus, Nolte. ex A. DC. l. c. 297 = P. fimbriata.
grandiflorus, A. DC. l. c. = P. grandiflora.
mexicanus, A. DC. l. c. = P. pimpinelloides.
nemophiloides, Kunth, Ind. Sem. Hort. Berol. (1846) 12.—Texas.
parviflorus, A. DC. in DC. Prod. ix. 297 = P. parviflora.
pectinatus, E. Mey. in Ann. Sc. Nat. Sér. III. v. (1846) 266.—Am. bor.
viscidus, A. DC. in DC. Prod. ix. 296 = P. grandiflora.

COSMARIUM, Dulac, Fl. Hautes-Pyr. 215 (1867) [non Corda, quod Algar. gen.] = **Adonis**, Linn. (Ranunc.).
aestivale, Dulac, l. c. = A. aestivalis.
autumnale, Dulac, l. c. = A. autumnalis.
pyrenaicum, Dulac, l. c. = A. pyrenaica.

COSMEA, Willd. Sp. Pl. iii. 2250 (1803) = **Cosmos**, Cav. (Compos.).
bipinnata, Willd. l. c. = Cosmos bipinnatus.
caudata, Spreng. Syst. iii. 615 = Cosmos caudatus.
chrysanthemifolia, Spreng. l. c. = Cosmos chrysanthemifolius.
crithmifolia, Spreng. l. c. = Cosmos crithmifolius.
diversifolia, Heynh. Nom. ii. 163 = Cosmos diversifolius.
lutea, Sims, Bot. Mag. t. 1689 = Bidens grandiflora.

COSMEA :—
parviflora, Willd. Sp. Pl. iii. 2250 = Cosmos parviflorus.
pilosa, Spreng. Syst. iii. 615 = Cosmos pilosus.
Scabiosae, Spreng. l. c. = Cosmos scabiosoides.
sulphurea, Willd. Sp. Pl. iii. 2250 = Cosmos sulphureus.
tenella, Spreng. Syst. iii. 615 = Cosmos tenellus.

COSMELIA, R. Br. Prod. 553 (1810). *EPACRIDEAE*, Benth. & Hook. f. ii. 617.
angustifolia, DC. Prod. vii. 766 = rubra.
rubra, *R. Br. Prod.* 553.—Austral.

COSMIA, Domb. ex Juss. Gen. 312 (1789) = **Calandrinia**, H. B. & K. (Portulac.).
caulescens, Domb. ex Spreng. Syst. ii. 453 = Calandrinia caulescens.
montana, Domb. ex DC. Prod. iii. 359 = Calandrinia caulescens.
prostrata, Domb. l. c. = Calandrinia acaulis.

COSMIBUENA, Ruiz & Pav. Fl. Per. iii. 3. t. 226 (1802). *RUBIACEAE*, Benth. & Hook. f. ii. 40.
BUENA, Pohl, Pl. Bras. Ic. i. 8, partim (1827).
acuminata, *Ruiz & Pav. Fl. Per.* iii. 3. t. 236.—Peruv.
dichotoma, G. Don, Gen. Syst. iii. 479 = Ladenbergia dichotoma.
latifolia, *Klotzsch, ex Walp. Rep.* vi. 69.—N Granat.
macrocarpa, *Klotzsch, ex Walp. l. c.*—Ins. Gorgona.
obtusifolia, *Ruiz & Pav. Fl. Per.* iii. 3.—Peruv.
ochracea, *Endl. Iconogr.* t. 90.—Hab. ?
quinqueflora, *Klotzsch, in Hayne, Arzneig.* xiv. sub t. 15.—Venezuela.
Skinneri, *Hemsl. Biol. Centr. Am. Bo* ii. 12.—Nicaragua.
triflora, *Klotzsch, in Hayne, Arzneig.* xiv. sub t. 15.—Guiana.

COSMIBUENA, Ruiz & Pav. Prod. Fl. Per. 10. t. 2 (1794) = **Hirtella**, Linn. (Rosac.).

COSMIDIUM, Nutt. in Trans. Am. Phil. Soc. Ser. II. vii. (1841) 361 = **Thelesperma**, Less. (Compos.).
Burridgeanum ×, Hort. ex Regel, Ind. Sem. Hort. Petrop. (1857) 40 = T. filifolium.
filifolium, Nutt. in Trans. Am. Phil. Soc. N. S. vii. (1841) 362 = T. filifolium.
gracile, Nutt. l. c. = T. gracile.
simplicifolium, A. Gray, Pl. Fendl. 86 = T. subsimplicifolium.

COSMIUSA, Alef. in Bot. Zeit. xxiv. (1866) 146 = **Parochetus**, Buch.-Ham. (Legumin.).
repens, Alef. l. c. = P. communis ?

COSMIZA, Rafin. Fl. Tellur. iv. 110 (1836) = **Polypompholyx**, Lehm. (Lentibul.).
coccinea, Rafin. l. c. = P. multifida.

COSMOPHYLLUM, C. Koch, Ind. Sem. Hort. Berol. (1854) 12 = **Podachaenium**, Benth. (Compos.).
cacaliaefolium, C. Koch, l. c.—Guatemala.

COSMOS, Cav. Ic. i. 9. tt. 14, 79 (1791). *COMPOSITAE*, Benth. & Hook. f. ii. 387.
ADENOLEPIS, Less. in Linnaea, vi. (1831) 510.
COSMEA, Willd. Sp. Pl. iii. 2250 (1803).
COSMUS, Pers. Syn. ii. 477 (1807).
bipinnatus, *Cav. Ic.* i. 10. t. 14.—Am. bor.; Mexic.
carvifolius, *Benth. Bot. Voy. Sulph.* 117.—Mexic.
caudatus, *H. B. & K. Nov. Gen. et Sp.* iv. 240.—Ind. occ.
chrysanthemifolius, *H. B. & K. l. c.* 239. t. 382.—Mexic.
crithmifolius, *H. B. & K. l. c.* 242.—Mexic.
diversifolius, *Otto, in Knowles & Westc. Flor. Cab.* ii. 3. t. 47.—Am. bor.
integrifolius, *Wedd. Chlor. And.* i. 70, nota.—Bolivia.
linearifolius, *Hemsl. Biol. Centr. Am. Bot.* ii. 200.—Mexic.
marginatus, *Klatt, in Abh. Naturf. Ges. Halle*, xv. (1882) 328.—Peruv.
parviflorus, *Pers. Syn.* ii. 477; *H. B. & K. Nov. Gen. et Sp.* iv. 241.—N. Mexic.
peucedanifolius, *Wedd. Chlor. And.* i. 70.—Bolivia.
pilosus, *H. B. & K. Nov. Gen. et Sp.* iv. 241.—Mexic.
pinnatus, Jacq. ex Steud. Nom. ed. II. i. 426 = Coreopsis coronata.

COSMOS :—
pulcherrimus, Sch. Bip. in Bull. Soc. Bot. Fr. xii. (1865) 79 = diversifolius.
purpureus, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 200.—Mexic.
reptans, Benth. Pl. Hartw. 40 = diversifolius.
roseus, *Mart. ex Steud. Nom.* ed. II. i. 426.—Bras.
scabiosoides, *H. B. & K. Nov. Gen. et Sp.* iv. 242.—Mexic.
subpubescens, *Wedd. Chlor. And.* i. 70.—Peruv.
sulphureus, *Cav. Ic.* i. 56. t. 79.—Mexic.
tenellus, *H. B. & K. Nov. Gen. et Sp.* iv. 240.—Mexic.
tenuifolius, *Lindl. Bot. Reg.* t. 2007.—Mexic.
Uhdeanus, *Kunth, Ind. Sem. Hort. Berol.* (1846) 12.—Mexic.

COSMOSTIGMA, Wight, Contrib. Bot. Ind. 41 (1834). *ASCLEPIADEAE*, Benth. & Hook. f. ii. 774.
acuminata, Wight, Ic. t. 1270 = racemosa.
racemosa, *Wight, Contrib. Bot. Ind.* 42.—Ind. or.

COSSIGNEA, Willd. Sp. Pl. ii. 226 (1799) = seq.

COSSIGNIA, Comm. ex Juss. Gen. 248 (1789) = seq.

COSSIGNYA, Baker, Fl. Maurit. 58 (1877) = seq.

COSSINIA, Comm. ex Lam. Encyc. ii. 132 (1786). *SAPINDACEAE*, Benth. & Hook. f. i. 397.
MELICOPSIDIUM, Baill. Adansonia, xi. (1874) 243.
borbonica, DC. Prod. i. 614 = pinnata.
Commersonii, Raeusch. Nom. ed. III. 103 = pinnata.
madagascariensis, *Baill. Adansonia*, xi. (1874) 247.—Madag.
pinnata, *Comm. ex Lam. Encyc.* ii. 132. t. 256.—Ins. Maurit.
splendens, Sieber, ex Steud. Nom. ed. II. i. 426 = pinnata.
ternata, Baker, Fl. Maurit. 59 = pinnata.
trifoliata, *Radlk. in Sitzb. Math.-Phys. Acad. Muench.* viii. (1878) 272.—N. Caled.
triphylla, Comm. ex Lam. Encyc. ii. 132 = pinnata.

COSSONIA, Dur. in Ann. Sc. Nat. Sér. III. xx. (1853) 82. t. 6 = **Raffenaldia**, Godr. (Crucifer.).
africana, Dur. l. c. = R. primuloides.

COSTA, Vell. Fl. Flum. 19 (1825); i. t. 48 (1827) = **Galipea**, Aubl. (Rutac.).
aromatica, Vell. l. c. 20; i. t. 48 = Galipea Arrabidae.

COSTAEA, A. Rich. Fl. Cub. Fanerog. ii. 75 (1853) = **Purdiaea**, Planch. (Cyrill.).
cubensis, A. Rich. l. c. 76 = P. monodynama.

COSTEA, A. Rich. Fl. Cub. Fanerog. ii. 328 (1853) = praec.

COSTIA, Willk. in Bot. Zeit. xvi. (1858) 377 = **Agropyron**, J. Gaertn. (Gramin.).
cristata, Willk. l. c. = A. cristatum.
imbricata, Willk. l. c. = A. imbricatum.
orientalis, Willk. l. c. = A. orientale.
squarrosa, Willk. l. c. = A. patulum.

COSTIA, Willk. in Bot. Zeit. xviii. (1860) 131 = **Iris**, Tourn.
caucasica, Willk. l. c. = I. caucasica.
persica, Willk. l. c. = I. persica.
scorpioides, Willk. l. c. = I. alata.

COSTUS, Linn. Mus. Cliff. 2 (1736); Gen. ed I. 331 (1737). *SCITAMINEAE*, Benth. & Hook. f. iii. 646.
ACINAX, Rafin. Fl. Tellur. iv. 53 (1836).
BANKSEA, Kocn. in Retz. Obs. iii. 75 (1783).
GISSANTHE, Salisb. in Trans. Hort. Soc. i. (1812) 279.
HELLENIA, Retz. Obs. vi. 18 (1791).
TSIANA, J. F. Gmel. Syst. 9 (1791).
JACUANGA, Lestib. in Ann. Sc. Nat. Sér. II. xv. (1841) 329, 341.
PLANERA, Gieske, Pracl. 205 (1792).
afer, *Ker-Gawl. in Bot. Reg.* t. 683.—Afr. trop.
Anachiri, Jacq. Fragm. 55 = spicatus.
angustifolius, Ker-Gawl. in Bot. Reg. t. 665 = speciosus.

COSTUS :—
appendiculatus, *Humb. & Bonpl. ex Roem. & Schult. Mant.* i. 32.—Am. austr.
arabicus, Aubl. Pl. Gui. i. 2 = spicatus.
arabicus, Linn. Sp. Pl. 2 = speciosus.
arabicus, Rosc. ex Spreng. Syst. i. 13 = glabratus.
arabicus, Vell. Fl. Flum. 2; i. t. 5 = Arrabidae.
argenteus, *Ruiz & Pav. Fl. Per.* i. 3. t. 4.—Peruv.
argyrophyllus, Wall. Cat. n. 6555 G, H = speciosus.
Arrabidae, *Steud. Nom.* ed. II. i. 426.—Bras.
cernuus, *Sw. ex Roem. & Schult. Syst.* i. 25.—Am. trop.
ciliatus, *Humb. & Bonpl. ex Roem. & Schult. Mant.* i. 32.—Am. trop.
ciliatus, *Miq. in Linnaea*, xviii. (1844) 73.—Guian.
comosus, *Rosc. in Trans. Linn. Soc.* viii. (1807) 350.—Am. austr.
conicus, Stokes, Bot. Mat. Med. i. 75 = spicatus.
crispiflorus, Stokes, l. c. = speciosus.
cylindricus, Jacq. Fragm. 54 = spiralis.
cylindricus, *Rosc. Scitam.* t. 78.—Ins. Trinit.
discolor, *Rosc. l. c.* t. 81.—Bras.
echinatus, Pers. Syn. i. 3 = Amomum echinatum.
foeniculaceus, *Noronha, in Verh. Batav. Gen.* v. (1790) ed. I. Art. iv. 12.—Malaya.
glabratus, Reichb. in Mocssl. Handb. ed. II. i. 8 = speciosus.
glabratus, *Sw. Prod. Veg. Ind. Occ.* 11.—Ind. occ.
globosus, *Blume, Enum. Pl. Jav.* 62.—Java.
hirsutus, *Presl, Rel. Haenk.* i. 112.—Mexic.
igneus, *N. E. Br. in Illustr. Hortic.* xxxi. (1884) 25. t. 511.—Bras.
laevis, *Ruiz & Pav. Fl. Per.* i. 3.—Peruv.
linearis, Spreng. Syst. i. 13 = Kaempferia secunda.
Loureiri, *Horan. Prod. Scitam.* 38.—Sumatra.
luteus, Blanco, Fl. Filip. ed. I. 4 = Roscoea lutea.
maculatus, *Rosc. Scitam.* t. 82.—Afr. trop.
malaccensis, Retz. Obs. iii. 71 = Alpinia malaccensis.
Malortieanus, *H. Wendl. in Hamb. Gartenz.* xix. (1863) 30.—Am. centr.
nepalensis, Rosc. Scitam. t. 80 = speciosus.
nigricans, Blanco, Fl. Filip. ed. II. 3 = Roscoea nigro-ciliata.
niveopurpureus, *Jacq. Fragm.* 55.—Ins. Martinic.
niveus, *G. F. W. Mey. Prim. Fl. Esseq.* 1.—Guian.
pictus, *D. Don, ex Lindl. Bot. Reg.* t. 1594.—Mexic.
Pisonis, Lindl. l. c. t. 899 = spiralis.
Potierae, *F. Muell. Fragm.* iv. 164.—Austral.
pulverulentus, *Presl, Rel. Haenk.* i. 111.—Mexic.
pungens, Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxix. (1867) 244 = Tapeinocheilos pungens.
quartus, Roem. & Schult. Syst. i. 26 = cylindricus.
quintus, Roem. & Schult. l. c. = spicatus.
ruber, *Wright, ex Griseb. Cat. Pl. Cub.* 256.—Cuba.
sarmentosus, *Boj. in Ann. Sc. Nat.* Sér. II. iv. (1835) 262. t. 8.—Ins. Maurit.
scaber, *Ruiz & Pav. Fl. Per.* i. 2. t. 3.—Peruv.
secundus, Roem. & Schult. Syst. i. 22 = Renealmia Pasco-Secora.
secundus, Spreng. Syst. i. 13 = Kaempferia secunda.
septimus, Roem. & Schult. Syst. i. 26 = villosissimus.
sericeus, Blume, Enum. Pl. Jav. 62 = speciosus.
sextus, Roem. & Schult. Syst. i. 26 = niveo-purpureus.
speciosus, *Sm. in Trans. Linn. Soc.* i. (1791) 249.—Ind. or.; Malaya.
spicatus, *Sw. Prod. Veg. Ind. Occ.* 11.—Ind. occ.
spiralis, *Rosc. in Trans. Linn. Soc.* viii. (1807) 350.—Venezuela.
vaginalis, Salisb. in Trans. Hort. Soc. i. (1812) 277 = speciosus.
Verschaffeltianus, *Lem. in Jard. Fleur.* iv. (1854) t. 381.—Bras.
villosissimus, *Jacq. Fragm.* 55.—Ins. S. Vincent.
Zerumbet, Pers. Syn. i. 3 = Alpinia nutans.

COTA, J. Gay, in Guss. Fl. Sic. Syn. ii. 866 (1844) = **Anthemis**, Mich. (Compos.).
altissima, J. Gay, l. c. 867 = A. Cota.
austriaca, Sch. Bip. in Oestr. Bot. Wochenbl. iv. (1854) 155 = A. austriaca.
Bourgaei, Boiss. Fl. Orient. iii. 284 = A. coclopoda.
brachycentros, J. Gay, in Guss. Fl. Sic. Syn. ii. 867 = A. brachycentros.
Brachmanni, Boiss. Diagn. Scr. II. iii. 21 = A. Brachmanni.
coclopoda, Boiss. l. c. = A. coclopoda.
Cossoniana, Willk. & Lange, Prod. Fl. Hisp. ii. 85 = A. austriaca.

COTA :—
lyonnetioides, Boiss. & Kotschy, Diagn. Ser. II. v. 109 = A. lyonnetioides.
ormenioides, Boiss. l. c. iii. 21 = A. Wiedemanniana.
oxylepis, Boiss l. c. v. 109 = A. oxylepis.
palaestina, Reut. ex Unger & Kotschy, Ins. Cyp. 240 = A. palestina?
parnassica, Boiss. & Heldr. Diagn. Scr. II. iii. 20 = A. tinctoria.
Pestalozzae, Boiss. l. c. 21 = A. Pestalozzae.
Thirkei, Sch. Bip. in Oestr. Bot. Wochenbl. iv. (1854) 163 = A. brachycentros.
tinctoria, J. Gay, in Guss. Fl. Sic. Syn. ii. 867 = A. tinctoria.
Triumfetti, J. Gay, l. c. = A. austriaca.

COTINUS, [Tourn.] Linn. Syst. ed. I (1735) = **Rhus**, Linn. (Anacard.).
americanus, Nutt. Sylva, ii. t. 81 = R. cotinoides.
coggygria, Scop. Fl. Carn. ed. II. i. 220 = R. Cotinus.
Coriaria, Duham. Arb. ed. I. i. 191 = R. Cotinus.

COTONEASTER, Rupp. Fl. Jen. ed. Hall. 137 (1745); Medik. Phil. Bot. i. 155 (1789). *ROSACEAE*, Benth. & Hook. f. i. 627.
GYMNOPYRENIUM, Dulac, Fl. Hautes-Pyr. 316 (1867),
NAEGELIA, Lindl. Veg. Kingd. 560 (1847).
acuminata, *Lindl. in Trans. Linn. Soc.* xiii. (1822) 101.—Reg. Himal.
acutifolia, Maxim. Prim. Fl. Amur. 171 = integerrima.
acutifolia, *Turcz. in Bull. Soc. Nat. Mosc.* iv. (1832) 190.—Mongolia.
aestivalis, Wenzig, in Linnaea, xxxviii. (1874) 202 = Crataegus aestivalis.
affinis, DC. Prod. ii. 632 = buxifolia.
affinis, Hohen. ex Hook. f. Fl. Brit. Ind. ii. 385 = frigida?
affinis, Lindl. in Trans. Linn. Soc. xiii. (1822) 101 = Stranvaesia glaucescens.
arborescens, Wenzig, in Linnaea, xxxviii. (1874) 203 = Crataegus arborescens.
bacillaris, *Wall. ex Lindl. Bot. Reg.* sub t. 1229.—Reg. Himal.
buxifolia, Baker, in Saund. Refug. Bot. i. t. 52 = microphylla.
buxifolia, *Wall. Cat.* n. 661; *et Lindl. Bot. Reg.* sub t. 1229.—Reg. Himal.
californica, *A. Murr. Oreg. Circ.* 1, nomen.—Calif.
coccinea, Steud. Nom. ed. II. i. 426 = tomentosa.
comptus, *Lem. in Fl. des Serres*, Sér. I. iv. (1848) 338 b.—Mexic.
congesta, Baker, in Saund. Refug. Bot. i. t. 51 = microphylla.
crenulata, C. Koch, Dendrol. i. 175 = Crataegus crenulata.
Cuila, Lee, ex C. Koch, l. c. 172 = nummularia.
denticulata, *H. B. & K. Nov. Gen. et Sp.* vi. 214. t. 556.—Mexic.
disticha, *Lange, in Bot. Tidssk.* xiii. (1882) 19.—Nepal?
elliptica, Hort. ex Loud. Encyc. Pl. 1208 = nummularia.
emarginata, Hort. ex C. Koch, Dendrol. i. 178 = microphylla.
Fontanesii, *Spach, Hist. Vég. Phan.* ii. 77.—Hab.?
Fortunei, Wenzig, in Linnaea, xxxviii. (1874) 200 = Photinia Fortuneana.
frigida, *Wall. Cat.* n. 657; *et Lindl. Bot. Reg.* t. 1229.—Reg. Himal.
Grammontii, Hort. ex C. Koch, Dendrol. i. 172 = nummularia.
granatensis, *Boiss. Elench.* 41.—Hispan.
horizontalis, *Decne. in Fl. des Serres*, Sér. II. xii. (1877) 168.—China.
hymalaica, *Carr. in Rev. Hortic.* (1875) 160.—Reg. Himal.
integerrima, *Medic. Gesch.* 85.—Europ.; As. temp.
intermedia, Medic. ex Trautv. in Act. Hort. Petrop. viii. (1883) 337, sphalm. = praec.
laevis, Hort. ex Steud. Nom. ed. II. i. 426 = nummularia.
lanata, Hort. ex Regel, Gartenfl. ix. (1860) 59 = buxifolia.
laxiflora, Jacq. ex Lindl. Bot. Reg. t. 1305 = integerrima.
Lindleyi, Steud. Nom. ed. II. i. 426 = nummularia.

COTONEASTER:—
lucida, Schlecht. in Linnaea, xxvii. (1854) 541 = integerrima.
marginata, Hort. ex C. Koch, Dendrol. i. 177 = buxifolia.
melanocarpa, Fisch. ex Loud. Hort. Brit. Suppl. ii. 624 = integerrima.
microphylla, *Wall. ex Lindl. Bot. Reg.* t. 1114; *Cat.* n. 662 A.—Reg. Himal.
microphylla, Wall. Cat. n. 662 B = thymifolia.
multiflora, *Bunge, in Ledeb. Fl. Alt.* ii. 220.—Sibir. alt.
multiflora, C. A. Mey. Verz. Pfl. Cauc. 171 = nummularia.
nebrodensis, *Nym. Syll.* 267.—Sicil.
nepalensis, Hort. ex C. Koch, Dendrol. i. 177 = frigida.
nervosa, *Decne. in Nouv. Arch. Mus. Par.* Sér. I. x. (1874) 177.—Hab.?
nigra, Wahlenb. Fl. Gothob. 53 = integerrima.
nummularia, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 34.—Afr. bor.; Oriens.
obtusa, Wall. Cat. n. 659; et Lindl. Bot. Reg. sub t. 1229 = bacillaris.
orientalis, *Kerner, in Oestr. Bot. Zeitschr.* xix. (1869) 270.—Europ.
parnassica, *Boiss. & Heldr. Diagn.* Ser. II. ii. 48.—Graecia.
pedunculares, Boiss. l. c. Ser. I. iii. 8 = integerrima.
polonica, *Jastrz. ex Rostaf. Fl. Polon. Prod.* 121.—Polonia.
prostrata, Baker, in Saund. Refug. Bot. i. t. 53 = rotundifolia.
Pyracantha, Spach, Hist. Vég. Phan. ii. 73 = Crataegus Pyracantha.
racemiflora, C. Koch, Dendrol. i. 170 = nummularia.
reflexa, *Carr. in Rev. Hortic.* (1870–71) 520.—China.
rosea, Edgew. in Trans. Linn. Soc. xx. (1851) 46 = bacillaris.
rotundifolia, *Wall. Cat.* n. 663; *et Lindl. Bot. Reg.* sub t. 1229.—Reg. Himal.
Royleana, Hort. ex C. Koch, Dendrol. i. 171 = acuminata.
Simonsii, *Hort. ex Baker, in Saund. Refug. Bot.* i. t. 55.—Reg. Himal.
spathulata, *Wenzig, in Linnaea*, xxxviii. (1874) 201.—Am. bor.
spicata, Hort. ex C. Koch, Dendrol. i. 166 = tomentosa.
thymaefolia, *Hort. ex Baker, in Saund. Refug. Bot.* i. t. 50.—Reg. Himal.
thymifolia, Hort. ex Loud. Encyc. Pl. 1384 = microphylla.
tomentosa, *Lindl. in Trans. Linn. Soc.* xiii. (1822) 101.—Europ.
tomentosa, C. A. Mey. Verz. Pfl. Cauc. 171 = nummularia.
uniflora, *Bunge, in Ledeb. Fl. Alt.* ii. 220.—Sibir. alt.
Uva-ursi, G. Don, in Loud. Hort. Brit. 480 = rotundifolia.
vulgaris, Lindl. in Trans. Linn. Soc. xiii. (1822) 101 = integerrima.

COTTAEA, Endl. Gen. 92 (1841) = seq.

COTTEA, Kunth, Rév. Gram. i. 84, 281. t. 52 (1830). *GRAMINEAE*, Benth. & Hook. f. iii. 1174.
pappophoroides, *Kunth, l. c.*—Peruv.
sarmentosa, *Nees, ex Steud. Syn. Pl. Gram.* 201.—Ins. S. Vinc.

COTTENDORFIA, Schult. f. Syst. vii. 64, 1193 (1830). *BROMELIACEAE* [Benth. & Hook. f. iii. 667].
albicans, *Griseb. in Goett. Abh.* xxiv. (1879) 330.—Reg. Argent.
florida, *Schult. f. Syst.* vii. II. 1193.—Bras.
guianensis, *Klotzsch, ex Beer, Bromel.* 44.—Guiana.

COTTONIA, Wight, Ic. v. I. 21. t. 1755 (1852). *ORCHIDEAE*, Benth. & Hook. f. iii. 572.
Championii, *Lindl. ex Benth. in Hook. Kew Journ.* vii. (1855) 35.—Hong Kong.
macrostachya, *Wight, Ic.* t. 1755.—Ind or.
pedunculares, Thw. Enum. Pl. Zeyl. 303 = macrostachya.

COTULA, [Tourn.] Linn. Syst. ed. I (1735). *COMPOSITAE*, Benth. & Hook. f. ii. 428.
BALDINGERIA, Neck. Elem. i, 88? (1790).
BROCCHIA, Vis. Pl. Aeg. et Nub. 35 (1836).
CENOCLINE, C. Koch, in Bot. Zeit. i. (1843) 41.
CHLAMYDOPHORA, Ehrenb. ex Less. Syn. Comp. 265, 448 (1832).
COTULINA, Pomel, Nouv. Mat. Fl. Atl. 56 (1874).
CTENOSPERMA, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 115.
GYMNOGYNE, Steetz, in Lehm. Pl. Preiss. i. 431 (1845).
LANCISIA, Adans. Fam. ii. 126 (1763).
LEPTINELLA, Cass. in Bull. Soc. Philom. (1822) 127.
LEPTOGYNE, Less. in Linnaea, vi. (1831) 151.
MACHLIS, DC. Prod. vi. 140 (1837).
OTOGLYPHIS, Pomel, Nouv. Mat. Fl. Atl. 56 (1874).
PLEIOGYNE, C. Koch, in Bot. Zeit. i. (1843) 40.
STRONGYLOSPERMA, Less. Syn. Comp. 261 (1832).
SYMPHYOMERA, Hook. f. in Hook. Lond. Journ. Bot. vi. (1847) 116.
abrotanifolia, Willd. Sp. Pl. iii. 2167 = Anacyclus creticus.
abyssinica, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 419.—Afr. trop.
africana, Steud. Nom. ed. I. 230 = Matricaria capensis.
alba, Linn. Syst. ed. XII. ii. 564 = Eclipta erecta.
alpina, *Hook. f. Fl. Tasm.* i. 192. t. 51 A.—Austral.
anthemoides, *Linn. Sp. Pl.* 891.—Hisp.; Afr. bor. et austr.; Ind. or.
anthemoides, Willd. Sp. Pl. iii. 2165 = microcephala.
atrata, *Hook. f. Handb. N. Zeal. Fl.* 142.—N. Zel.
aurea, *Loefl. Iter*, 163.—Reg. Mediterr.; Oriens.
aurea, Sibth. & Sm. Fl. Graec. ix. 61 = Matricaria decipiens.
australis, *Hook. f. Fl. Nov. Zel.* 128.—Austral.; N. Zel.
barbata, DC. Prod. vi. 79 = Cenia barbata.
bicolor, Roth, Catalect. fasc. ii. 116 = Dichrocephala latifolia.
bipinnata, *Thunb. Prod. Pl. Cap.* 162.—Afr. austr.
bracteolata, *E. Mey. ex DC. Prod.* vi. 78.—Afr. austr.
capensis, Linn. Mant. ii. 287 = Matricaria capensis.
ceniaefolia, *DC. Prod.* vi. 79.—Afr. austr.
chamaemelifolia, *Ehrenb. ex C. Koch, in Bot. Zeit.* i. (1843) 41.—Africa?
chrysanthemifolia, Blume, Bijdr. 918 = Dichrocephala chrysanthemifolia.
cinerea, *Delile, Fl. Egypte*, 275. t. 47. f. 4.—Afr. bor.; Arab.
cinerea, Kotschy, ex Benth. & Hook. f. Gen. ii. 420 = Kotschyi.
complanata, Sibth. & Sm. Fl. Graec. Prod. ii. 187 = Anthemis pectinata.
conica, Wall. Cat. n. 3185 = Spilanthes Acmella.
coronopifolia, *Linn. Sp. Pl.* 892.—Afr. austr. (Europ. alien.).
crinita, E. Mey. ex DC. Prod. vi. 144 = Hippia gracilis.
cryptocephala, *Sch. Bip. ex A. Rich. Tent. Fl. Abyss.* i. 420.—Abyss.
cuneifolia, Willd. Sp. Pl. iii. 2169 = Centipeda orbicularis.
Cunninghami, F. Muell. Fragm. viii. 143 = Centipeda Cunninghami.
debilis, Balb. ex Spreng. Neue Entdeck. ii. 137 = Eclipta pusilla.
decumbens, Pers. Syn. ii. 464 = Centipeda orbicularis.
dichotoma, Pers. l. c. = C. Pyrethraria.
dichrocephala, Sch. Bip. ex A. Rich. Tent. Fl. Abyss. i. 419 = anthemoides.
dichrocephaloides, C. B. Clarke, Comp. Ind. 150 = Dichrocephala Hamiltoni.
dioica, *Hook. f. Handb. N. Zeal. Fl.* 143.—N. Zel.
divaricata, Wall. ex DC. Prod. v. 375 = Thespis divaricata.
Drummondii, *Benth. Fl. Austral.* iii. 550.—Austral.
ebracteata, Walp. Rep. ii. 991 = bracteolata.
ebracteolata, C. Koch, in Bot. Zeit. i. (1843) 39 = bracteolata.
elongata, *C. B. Vogel, in Trew, Pl. Rar.* iii. 11. t. 26.—Java.
Featherstonei, *F. Muell. ex J. Buch. in Trans. N. Z. Inst.* vii. (1875) 337.—Ins. Chatham.
Filicula, *Hook. f. ex Benth. Fl. Austral.* iii. 551.—Austral.
filifolia, *Thunb. Prod. Pl. Cap.* 161.—Afr. austr.
filiformis, *Hook. f. Handb. N. Zeal. Fl.* 142.—N. Zel.

COTULA:—
fimbriata, Spreng. Neue Entdeck. iii. 41 = Hippia gracilis.
foetida, Poepp. ex DC. Prod. vi. 139 (= *Myriogyne elatinoides*).—Chili.
globifera, Licht. ex Steud. Nom. ed. II. i. 427 = Pentzia globosa.
globifera, Thunb. Prod. Pl. Cap. 162 = Matricaria globifera.
grandis, Jacq. Obs. Bot. iv. 4. t. 81 = Chrysanthemum flosculosum.
grandis, Linn. Sp. Pl. ed. II. 1257 (= *Plagius grandiflorus*).—Afr. bor.
gymnogyne, *F. Muell. ex Benth. Fl. Austral.* iii. 549.—Austral.
hemisphaerica, *Wall. Cat.* n. 3236.—Ind. or.
heterocarpa, *DC. Prod.* vi. 80.—Afr. austr.
hispida, Harv. in Harv. & Sond. Fl. Cap. iii. 183 = Cenia hispida.
humifusa, Willd. ex Steud. Nom. ed. II. i. 427 = Egletes humifusa.
integrifolia, Burch. Trav. i. 51 = bracteolata.
integrifolia, *Hook. f. Fl. Tasm.* i. 192. t. 50 B.—Tasmania.
Kotschyi, *Benth. & Hook. f. Gen.* ii. 429.—Afr. trop.
lanata, *Hook. f. Handb. N. Zeal. Fl.* 141.—N. Zel.
latifolia, Pers. Syn. ii. 464 = Dichrocephala latifolia.
laxa, *DC. Prod.* vi. 78.—Afr. austr.
leptalea, *DC. l. c.* 80.—Afr. austr.
linearifolia, *Cheesem. in Trans. N. Z. Inst.* xv. (1883) 299.—N. Zel.
linifolia, Burm. f. Fl. Cap. Prod. 27 = Felicia fragilis.
lyrata, Boj. ex DC. Prod. v. 372 = Dichrocephala lyrata.
maderaspatana, Willd. Sp. Pl. iii. 2170 = Grangea maderaspatana.
Maniotota, *Petrie, in Trans. N. Z. Inst.* xiv. (1882) 362.—N. Zel.
matricarioides, Bong. Veg. Ins. Sitcha, 29 = Matricaria discoidea.
microcephala, *DC. Prod.* vi. 79.—Afr. austr.
minima, *DC. l. c.* 80.—Hab.?
minima, Willd. Sp. Pl. iii. 2170 = Centipeda orbicularis.
minor, Caruel, ex Boiss. Fl. Orient. iii. 358 = cinerea.
minor, *Hook. f. Handb. N. Zeal. Fl.* 142.—N. Zel.
minuta, Forst. f. Prod. 57 = Centipeda orbicularis.
montevidensis; Spreng. Syst. iii. 497 = coronopifolia.
Moseleyi, *Hemsl. Bot. Voy. Challenger*, i. II. 152.—Tristan d'Acugna.
multifida, DC. Prod. vi. 80 = villosa.
myriophylloides, *Hook. Ic. Pl.* t. 335.—Afr. austr.
nudicaulis, *Thunb. Prod. Pl. Cap.* 162.—Afr. austr.
Oederi, *Murr. Hort. Bot. Stirp.* 228.—Hab.?
oxydonta, DC. Prod. vi. 78 = bipinnata.
pectinata, *Hook. f. Handb. N. Zeal. Fl.* 142.—N. Zel.
perpusilla, *Hook. f. l. c.* 143.—N. Zel.
pilulifera, Linn. f. Suppl. 378 = Matricaria globifera.
Piper, *Vell. Fl. Flum.* viii. t. 136.—Bras.
plumosa, *Hook. f. Handb. N. Zeal. Fl.* 141.—N. Zel.
poekilophylla, *C. Koch, in Bot. Zeit.* i. (1843) 39.—Abyss.
prostrata, Linn. Syst. ed. XII. 564 = Eclipta erecta.
pterocarpa, *DC. Prod.* vi. 80.—Afr. austr.
pubescens, *Desf. Fl. Atlant.* ii. 284.—Afr. bor.
pubescens, Viv. ex Coss. in Bull. Soc. Bot. Fr. xii. (1865) 278 = aurea.
pumila, Houtt. Handleid. x. 772 = Cenia turbinata.
pusilla, *Thunb. Prod. Pl. Cap.* 162.—Afr. austr.
pygmaea, *Benth. & Hook. f. ex Hemsl. Biol. Centr. Am. Bot.* ii. 230.—Mexic.
pygmaea, Poir. Encyc. Suppl. ii. (1810) 371 = Nananthea perpusilla.
Pyrethraria, *Linn. Mant.* i. 116.—Am. austr.
pyrethrifolia, *Hook. f. Handb. N. Zeal. Fl.* 142.—N. Zel.
quinquefida, Thunb. Prod. Fl. Cap. 161 = Pentzia quinquefida.
quinqueloba, Blanco, Fl. Filip. ed. I. 626 = Centipeda orbicularis.
quinqueloba, Linn. f. Suppl. 377 = Lidbeckia lobata.
reptans, *Benth. Fl. Austral.* iii. 551.—Austral.
reptans, Boj. ex DC. Prod. v. 371 = Sphaeranthus cotuloides.
rosea, *Boj. ex Less. in Linnaea*, vi. (1831) 152.—Madag.
Russeliana, Wall. Cat. n. 3240 = Sphaeromorphea Russeliana.
sericea, Linn. f. Suppl. 377 = Cenia sericea.

COTULA :—
sinapifolia, Buch.-Ham. ex Roxb. Hort. Beng. 62; Fl. Ind. iii. 437 = Dichrocephala latifolia.
sinapifolia, Wall. ex DC. Prod. v. 375 = Thespis divaricata.
sonchifolia, Bieb. Fl. Taur. Cauc. ii. 328 = Dichrocephala latifolia.
sororia, *DC. Prod.* vi. 79.—Afr. austr.
Sphaeranthus, Link, Enum. Hort. Berol. ii. 344 = Grangea maderaspatana.
Spilanthus, Linn. Mant. i. 116 = Spilanthes urens.
squalida, *Hook. f. Handb. N. Zeal. Fl.* 143.—N. Zel.
stenophylla, *C. Koch, in Bot. Zeit.* i. (1843) 40.—Afr. austr.
sternutatoria, Wall. Cat. n. 3259 = Centipeda orbicularis.
stolonifera, Harv. in Harv. & Sond. Fl. Cap. iii. 182 = ceniaefolia.
stricta, Linn. Mant. 287 = Lidbeckia pectinata.
tanacetifolia, Linn. Syst. ed. XII; Murr. Syst. Veg. ed. XIII. 644 n. 10 = Matricaria multiflora.
Teesdaliae, *DC. Prod.* vi. 80.—Afr. austr.
tenella, *E. Mey. ex DC. l. c.* 80.—Afr. austr.
thespidioides, F. Muell. Fragm. viii. 143 = Centipeda thespidioides.
Thunbergii, Harv. in Harv. & Sond. Fl. Cap. iii. 183 = Cenia Thunbergii.
tridentata, *Benth. & Hook. f. Gen.* ii. 420.—Aegypt.
tripinnata, Thunb. Prod. Pl. Cap. 162 = Matricaria grandiflora.
turbinata, Linn. Sp. Pl. 892 = Cenia turbinata.
umbellata, Linn. f. Suppl. 378 = Peyrousea calycina.
valparadisea, *Phil. in Anal. Univ. Chil.* (1873) 503.—Chili.
Verbesina, *Linn. Amoen. Acad.* v. 407 = Adenostemma Swartzii.
villosa, *DC. Prod.* vi. 79.—Afr. austr.
viscosa, Linn. Sp. Pl. 892 = Egletes viscosa.
Zeyheri, *Fenzl, ex Harv. & Sond. Fl. Cap.* iii. 180.—Afr. austr.

COTULINA, Pomel, Nouv. Mat. Fl. Atl. 56 (1874) = **Cotula**, Tourn. (Compos.).
aurea, Pomel, l. c. = Cotula aurea.

COTYLANTHERA, Blume, Bijdr. 707 (1825).
GENTIANEAE, Benth. & Hook. f. ii. 803.
EOPHYLON, A. Gray, in Journ. Linn. Soc. xi. (1871) 22.
paucisquama, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* iv. 94.—Reg. Himal.
tenuis, *Blume, Bijdr.* 708.—Java.

COTYLARIA, Rafin. Princ. Somiol. 25 (1814) = **Cotyledon**, Tourn. (Crassulac.).

COTYLEDON, Tourn. ex Linn. Syst. ed. I (1735).
CRASSULACEAE, Benth. & Hook. f. i. 659.
COTYLARIA, Rafin. Princ. Somiol. 25 (1814).
COTYLIPHYLLUM, Link, Handb. ii. 22 (1831).
COURANTIA, Lem. Jard. Fleur. i. (1851) Misc. 92.
ECHEVERIA, DC. Prod. iii. 401 (1828).
OROSTACHYS, [Fisch.] Hort. Gorenk. ed. I. 99 (1808).
ORTHOSTACHYS, Spach, Hist. Vég. Phan. ix. 32 (1840).
PACHYPHYTUM, Klotzsch, in Otto & Dietr. Allg. Gartenz. ix. (1841) 9.
PISTORINIA, DC. Prod. iii. 399 (1828).
UMBILICARIA, Pers. Syn. i. 510 (1805).
UMBILICUS, DC. in Bull. Soc. Philom. (1801) 49.
acaulon, Walt. ex Steud. Nom. ed. I. 230 = spuria.
acutifolia, *Baker, in Saund. Refug. Bot.* i. sub t. 71. n. 34.—Mexic.
aduncа, *Baker, l. c.* t. 60.—Mexic.
aegyptiaca, Lam. Encyc. ii. 142 = Kalanchoë crenata.
africana, Burm. f. ex Steud. Nom. ed. I. 230 = spuria.
agavoides, *Baker, in Saund. Refug. Bot.* t. 67.—Mexic.
albiflora, *Hemsl. Diagn. Pl. Nov.* 9.—Mexic.
alternans, Haw. Suppl. Pl. Succ. 26 = maculata.
alternans, Vahl, Symb. Bot. ii. 51 = Kalanchoë alternans.
ambigua, Salisb. Prod. 307 = orbiculata.
amplexicaulis, Hcyne, ex C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 415 = Kalanchoë floribunda.

COTYLEDON :—
arborescens, Mill. Gard. Dict. ed. VIII. n. 7 = Crassula arborescens.
aristata, Walp. Rep. ii. 258, sphalm. = cristata.
atropurpurea, *Baker, in Saund. Refug. Bot.* t. 198.—Mexic.
Batesii, *Hemsl. Diag. Pl. Nov.* 9.—Mexic.
bifida, *Hemsl. Biol. Centr. Am. Bot.* i. 388.—Mexic.
bracteolata, *Baker, in Saund. Refug. Bot.* sub t. 62.—N. Granat.
brasilica, Vell. Fl. Flum. 197; iv. t. 184 = Kalanchoë brasiliensis.
cacalioides, Eckl. & Zeyh. Enum. 307 = Eckloniana.
cacalioides, *Linn. f. Suppl.* 242.—Afr. austr.
caespitosa, *Haw. Syn. Pl. Succ.* 107.—Am. bor. occ.
californica, *Baker, in Saund. Refug. Bot.* 70.—Am. bor. occ.
calycina, Roth, Nov. Pl. Sp. 217 = Bryophyllum calycinum.
canaliculata, *Baker, in Saund. Refug. Bot.* i. sub t. 58.—Mexic.
canaliculata, Haw. Suppl. Pl. Succ. 22 = ungulata.
canalifolia, Haw. in Phil. Mag. (1825) 327 = coruscans.
carnicolor, *Baker, in Saund. Refug. Bot.* iii. t. 199.—Mexic.
caryophyllacea, *Burm. f. Prod. Fl. Cap.* 13.—Afr. austr.
cespitosa, Haw. Misc. 180 = caespitosa.
chiclensis, *Ball, in Journ. Linn. Soc.* xxii. (1885) 38.—Peruv.
clavifolia, *Haw. in Phil. Mag.* (1827) 274.—Afr. austr.
coccinea, *Cav. Ic.* ii. 54. t. 170.—Mexic.
Cooperi, *Baker, in Saund. Refug. Bot.* t. 72.—Afr. austr.
Corderoyi, *Baker, in Gard. Chron.* (1874) 599.—Mexic.
coruscans, *Haw. Suppl. Pl. Succ.* 21.—Afr. austr.
corymbosa, Rottl. ex Wight & Arn. Prod. 360 = Kalanchoë floribunda.
Cossoniana, *Ball, in Journ. Bot.* xi. (1873) 332.—Marocc.
crassifolia, Haw. in Phil. Mag. (1827) 273 = orbiculata?
crassifolia, Salisb. Prod. 307 = hemisphaerica.
crenata, Vent. Jard. Malm. 49 = Kalanchoë crenata.
cristata, *Haw. in Phil. Mag.* (1827) 274.—Afr. austr.
cuneata, E. Mey. ex Harv. & Sond. Fl. Cap. ii. 372 = Meyeri.
cuneata, *Thunb. Prod. Pl. Cap.* 83.—Afr. austr.
cuneiformis, *Haw. in Phil. Mag.* (1828) 185.—Afr. austr.
curviflora, *Sims, Bot. Mag.* t. 2044.—Afr. austr.
cymosa, *Baker, in Saund. Refug. Bot.* t. 68.—Mexic.
decipiens, *Baker, l. c.* 200.—Peruv.
decussata, *Sims, Bot. Mag.* t. 2518.—Afr. austr.
deficiens, Forsk. Fl. Aegypt. Arab. 89 = Kalanchoë crenata.
deficiens, Hochst. & Steud. ex A. Rich. Tent. Fl. Abyss. i. 310 = Kalanchoë Schimperiana.
Desmetiana, *Hemsl. Biol. Centr. Am. Bot.* i. 389.—Mexic.
dichotoma, *Haw. Suppl. Pl. Succ.* 27.—Afr. austr.
Eckloniana, *Harv. in Harv. & Sond. Fl. Cap.* ii. 374.—Afr. austr.
edulis, *Brewer, in Bot. Calif.* i. 211.—Calif.
elata, Haw. Suppl. Pl. Succ. 20 = orbiculata, triflora.
erecta, D. Dietr. Syn. Pl. ii. 1626 = lusitanica.
farinosa, *Baker, in Saund. Refug. Bot.* i. t. 71.—Calif.
farinulenta, *Hemsl. Biol. Centr. Am. Bot.* i. 389.—Mexic.
fasciculariis, [*Soland. in*] *Ait. Hort. Kew.* ed. I. ii. 106; *Bot. Mag.* t. 5602.—Afr. austr.
filicaulis, Eckl. & Zeyh. Enum. 307 = mamillaris.
fimbriata, *Turcz. in Bull. Soc. Nat. Mosc.* xvii. (1844) 241.—China; As. bor.
fulgens, *Baker, in Saund. Refug. Bot.* t. 64.—Mexic.
Galeottiana, *Hemsl. Diag. Pl. Nov.* i. 9.—Mexic.
gibbiflora, *Moç. & Sessé, ex DC. Prod.* iii. 401.—Mexic.
glauca, *Baker, in Saund. Refug. Bot.* t. 61.—Mexic.

COTYLEDON :—
globulariacefolia, *Baker, in Saund. Refug. Bot.* t. 201.—Syria; As. Min.
gracilis, *Harv. in Harv. & Sond. Fl. Cap.* ii. 373.—Afr. austr.
gracilis, *Haw. Suppl. Pl. Succ.* 26.—Afr. austr.
grandiflora, Burm. f. Fl. Cap. Prod. 13 = tuberculosa.
Grayii, *Baker, in Saund. Refug. Bot.* sub t. 71. n. 33.—N. Mexic.
hemisphaerica, *Linn. Sp. Pl.* 429.—Afr. austr.
heterophylla, Roxb. Hort. Beng. 34; Fl. Ind. ii. 456 = Kalanchoë floribunda.
hirsuta, Herb. Heyne, ex C. B. Clarke, in Fl. Brit. Ind. 414 = Kalanchoë glandulosa.
hispanica, *Linn. Sp. Pl.* 1196.—Hispan.; Marocc.
hispida, Lam. Encyc. ii. 141 = C. Mucizonia.
horizontalis, Guss. Ind. Sem. Hort. Boccad. (1826) 4 = C. Umbilicus.
hybrida, Hort. Par. ex DC. Prod. iii. 395 = Kalanchoë spathulata.
incarum, *Ball, in Journ. Linn. Soc.* xxii. (1885) 37 (err. typ. *incanum*).—Peruv.
integra, Medic. in Act. Acad. Theod. Palat. iii. Phys. (1775) 200. t. 9 = Kalanchoë crenata.
interjecta, *Haw. in Phil. Mag.* (1828) 185.—Afr. austr.
japonica, *Maxim. in Bull. Acad. Pétersb.* xxix. (1883) 122.—Japon.
jasminiflora, *Salm-Dyck, Obs.* 5.—Afr. austr.
Jurgensenii, *Hemsl. Diag. Pl. Nov.* 9.—Mexic.
laciniata, Linn. Sp. Pl. 430 = Kalanchoë laciniata.
lanceolata, *Benth. & Hook. f. ex S. Wats. Bot. Calif.* i. 211.—Calif.
lanceolata, Blanco, Fl. Filip. ed. I. 382 = spathulata.
lanceolata, Forsk. Fl. Aegypt. Arab. 89 = Kalanchoë lanceolata.
laxa, Benth. & Hook. f. ex Bot. Calif. i. 212 = californica.
leucantha, *Ledeb. Fl. Alt.* ii. 198.—As. bor.; Sibir. altaic.
libanotica, Labill. Ic. Pl. Syr. Dec. iii. 3.—Kurdist.; Syria.
Lievenii, *Ledeb. Fl. Alt.* ii. 197.—Persia; Sibiria.
linguaefolia, *Lem. Illustr. Hortic.* x. (1863) *Misc.* 81, in syn.—Mexic.
linguaeformis, [R. Br. in] Ait. Hort. Kew. ed. II. iii. 109 = caespitosa.
Lingula, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 293.—Am. bor. occ.
lurida, *Baker, in Saund. Refug. Bot.* t. 59.—Mexic.
lusitanica, *Lam. Encyc.* ii. 140.—Europ.
lutea, Hort. ex Lam. l. c. 172 = Crassula portulacea.
lutea, Huds. Fl. Angl. ed. II. 194 = lusitanica.
maculata, *Salm-Dyck, Obs.* 5.—Afr. austr.
malacophylla, *Pall. Reise*, iii. 729.—Sibiria.
mamillaris, *Linn. f. Suppl.* 242.—Afr. austr.
margiuatum, *Rafin. Atl. Journ.* 79.—Hab.?
mexicana, *Hemsl. Biol. Centr. Am. Bot.* i. 390.—Mexic.
Meyerii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 372.—Afr. austr.
Mucizouia, *Ortega, Muciz.* (1772).—Hispan.; Marocc.
mucronata, *Baker, in Saund. Refug. Bot.* sub t. 55. n. 4.—Mexic.
mucronata, *Lam. Encyc.* ii. 142.—Afr. austr.
nevadensis, *S. Wats. Bot. Calif.* i. 212.—Am. bor. occ.
nodulosa, *Baker, in Saund. Refug. Bot.* t. 56.—Mexic.
nuda, *Baker, l. c.* t. 57.—Mexic.
nudicaulis, Lam. Encyc. ii. 142; Vahl, Symb. Bot. ii. 51 = Kalanchoë crenata.
oblonga, Haw. Syn. Pl. Succ. 106 = orbiculata.
ombilicus, Lam. Encyc. ii. 140 = C. Umbilicus.
oppositifolia, *Ledeb. ex Nordm. in Bull. Sc. Acad. Pétersb.* ii. (1837) 313.—Reg. Caucas.
orbiculata, Forsk. Fl. Aegypt. Arab. p. cxii = Kalanchoë alternans.
orbiculata, *Linn. Sp. Pl.* 429.—Afr. austr.
Oreades, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 416.—Ind. or.
oregonensis, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 373.—Am. bor. occ.
ovata, Haw. Syn. Pl. Succ. 105 = orbiculata.
ovata, Mill. Gard. Dict. ed. VIII. n. 8 = Crassula portulacea.
Pachyphytum, *Baker, in Saund. Refug. Bot.* t. sub t. 59. n. 12.—Mexic.
Palmeri, *S. Wats. in Proc. Am. Acad.* xiv. (1879) 293.—Am. bor. occ.

COTYLEDON :—
paniculata, Linn. f. Suppl. 242 = orbiculata.
paniculata, Rottl. ex Wight & Arn. Prod. 360 = Kalanchoë floribunda.
paniculata, Thunb. Prod. Pl. Cap. 83 = fascicularis.
pannosa, *Baker, in Journ. Linn. Soc.* xviii. (1881) 269. —Madag.
papillaris, Haw. Suppl. Pl. Succ. 21 = decussata.
papillaris, *Linn. f. Suppl.* 242.—Afr. austr.
papillosa, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 58.—Afghan.
parviflora, *Desf. in Ann. Mus. Par.* xi. (1808) 444; *Sibth. & Sm. Fl. Graec. Prod.* i. 308.—Europ.
parviflora, *Hemsl. Diag. Pl. Nov.* 9.—Mexic.
parvula, *Burch. Trav.* i. 219.—Afr. austr.
Peacockii, *Baker, in Gard. Chron.* (1874) II. 258.—Am. bor.
peltata, Wendl. Beobacht. 49 = C. Umbilicus.
peruviana, *Baker, in Saund. Refug. Bot.* sub t. 58.—Peruv.
Pestalozzae, *Mast. in Gard. Chron.* (1877) II. 456.—Oriens.
pinnata, Lam. Encyc. ii. 141 = Bryophyllum calycinum.
Pistorinia, *Ortega, Muciz.* (1772).—Hispan.
pubescens, *Baker, in Saund. Refug. Bot.* i. sub t. 55.—Mexic.
pubescens, C. A. Mey. Verz. Pfl. Cauc. 150 = Sedum pilosum.
pulverulenta, *Baker, in Saund. Refug. Bot.* i. t. 66.—Calif.
pumila, *Baker, l. c.* t. 62.—Mexic.
punctata, Hort. ex Lam. Encyc. ii. 173 = Crassula arborescens.
purpurea, Haw. Suppl. Pl. Succ. 23 = curviflora.
purpurea, *Thunb. Prod. Pl. Cap.* 83.—Afr. austr.
quitensis, *Baker. in Saund. Refug. Bot.* sub t. 55.—Ecuador.
racemosa, *E. Mey. ex Harv. & Sond. Fl. Cap.* ii. 375.—Afr. austr.
ramosa, Haw. Suppl. Pl. Succ. 24 = orbiculata.
ramosissima, Mill. Gard. Dict. ed. VIII, n. 6 = orbiculata.
ramosissima, *Salm-Dyck, ex Haw. l. c.* 25.—Afr. austr.
reflexa, Willd. Enum. Hort. Berol. Suppl. 24 = caespitosa.
reticulata, *Thunb. Prod. Pl. Cap.* 83.—Afr. austr.
retusa, *Baker, in Saund. Refug. Bot.* sub t. 64. n. 22.—Mexic.
rhizophylla, Roxb. Hort. Beng. 34; Fl. Ind. ii. 456 = Bryophyllum calycinum.
rhombifolia, Eckl. & Zeyh. Enum. 307 = hemisphaerica.
rhombifolia, *Haw. in Phil. Mag.* (1825) 33.—Afr. austr.
rosea, *Less. in Linnaea*, ix. (1834) 177.—Sibiria.
roseata, *Baker, in Saund. Refug. Bot.* sub t. 55. n. 3.—Mexic.
rotundifolia, Eckl. & Zeyh. Enum. 307 = hemisphaerica.
rotundifolia, Haw. in Phil. Mag. (1827) 273 = orbiculata.
rupestris, Salisb. Prod. 307 = C. Umbilicus.
Rusbyi, *Greene, in Bull. Torrey Club*, x. (1883) 125.—Am. bor. occ.
Salzmannii, Hook. f. Bot. Mag. t. 5801 = hispanica.
Samium, *Urv. in Mém. Soc. Linn. Par.* i. (1822) 306.—As. Min.
Schaffneri, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 354.—Am. bor. occ.
Scheerii, *Baker, in Saund. Refug. Bot.* sub t. 62. n. 19.—Mexic.
secunda, *Baker, l. c.* sub t. 60. n. 14.—Mexic.
sediformis, Lapeyr. Hist. Abr. Pl. Pyr. 257 = sedoides.
sedoides, *DC. Rapp. Jard. Bot. Genèv.* ii. 79.—Europ.
Sempervivum, *Bieb. Beschr. Casp.* 176.—Reg. Casp.
serrata, *Linn. Sp. Pl.* 429.—As. Min.
simensis, *Britten, in Oliver, Fl. Trop. Afr.* ii. 399.—Afr. trop.
spathulata, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* ii. 416.—Reg. Himal.
spathulata, Poir. Encyc. Suppl. ii. 373 = Kalanchoë spathulata.
spinosa, C. B. Clarke, in Hook. f. Fl. Brit. Ind. ii. 416 = leucantha.
spinosa, *Linn. Sp. Pl.* 429.—As. bor.
Sprucei, *Baker, in Saund. Refug. Bot.* sub t. 71.—Ecuador.

COTYLEDON :—
spuria, *Linn. Sp. Pl.* ed. II. 614.—Afr. austr.
stolonifera, *Baker, in Saund. Refug. Bot.* t. 63.—Mexic.
strictiflora, *Baker, l. c.* sub t. 62. n. 17.—N. Mexic.
subspicata, *Baker, l. c.* sub t. 71.—N. Granat.
subulata, C. A. Mey. Verz. Pfl. Cauc. 150 = Sedum subulatum.
subulifolia, *Baker, in Saund. Refug. Bot.* sub t. 71. n. 32.—Mexic.
tardiflora, Bonpl. Pl. Rar. Malm. 93. t. 37 = fasciculatis.
tenuicaulis, *Aitch. & Hemsl. in Journ. Linn. Soc.* xviii. (1880) 57.—Afghan.
teretifolia, Lam. Encyc. ii. 139 = spuria.
teretifolia, *Thunb. Prod. Pl. Cap.* 83.—Afr. austr.
thyrsiflora, *D. Dietr. Syn. Pl.* ii. 1627.—Sibir.
thyrsiflora, Maxim. in Bull. Acad. Pétersb. xxix. (1884) 123 = leucantha.
tomentosa, *Harv. in Harv. & Sond. Fl. Cap.* ii. 373.—Afr. austr.
tricuspidata, Harv. in Phil. Mag. (1825) 32 = papillaris.
triflora, *Thunb. Prod. Pl. Cap.* 83.—Afr. austr.
trigyna, *Burch. Trav.* ii. 226.—Afr. austr.
tuberculosa, *Lam. Encyc.* iii. 139.—Afr. austr.
tuberosa, *Hort. Par. ex Webb & Berth. Phyt. Canar.* i. 177.—Ins. Canar.
umbilicata, Lam. Fl. Fr. iii. 59 = C. Umbilicus.
umbilicifolia, Stokes, Bot. Mat. Med. ii. 545 = C. Umbilicus.
Umbilicus, *Linn. Sp. Pl.* 429.—Europ.; As. Min.
undulata, Eckl. & Zeyh. Enum. 306 = cuneata.
undulata, *Haw. Suppl. Pl. Succ.* 20.—Afr. austr.
ungulata, *Lam. Encyc.* ii. 139.—Afr. austr.
velutina, *Hook. f. Bot. Mag.* t. 5602.—Afr. austr.
ventricosa, *Burm. f. Fl. Cap. Prod.* 13.—Afr. austr.
Verea, Jacq. Hort. Schoenb. iv. 17 = Kalanchoë crenata.
viridis, Haw. in Phil. Mag. (1827) 273 = orbiculata?
viscida, *S. Wats. in Proc. Am. Acad.* xvii. (1881–82) 372.—Am. bor. occ.
viscosa, Vahl, Symb. Bot. ii. 51 = C. Mucizonia.
Wallichii, *Harv. in Harv. & Sond. Fl. Cap.* ii. 374.—Afr. austr.
Zeyheri, *Harv. l. c.* 377.—Afr. austr.

COTYLEPHORA, Meissn. Gen. Comm. 28 (1837) = **Neesia**, Blume (Malvac.).
altissima, Meissn. l. c. = N. altissima.

COTYLIPHYLLUM, Link, Handb. ii. 23 (1831) = **Cotyledon**, Tourn. (Crassulac.).
erectum, Link, l. c. = Cotyledon Umbilicus.
horizontale, Link, l. c. = Cotyledon Umbilicus
Mucizonia, Link, l. c. = Cotyledon Mucizonia.
Umbilicus, Link, l. c. = Cotyledon Umbilicus.

COTYLISCUS, Desv. Journ. Bot. iii. (1813) 164, 175 = **Senebiera**, DC. (Crucif.).
niloticus, Desv. l. c. = S. nilotica.

COTYLODISCUS, Radlk. in Sitzb. Math.-Phys. Acad. Muench. viii. (1878) 334. *SAPINDACEAE.*
stelechanthus, *Radlk. l. c.*—Madag.

COTYLOPLECTA, Alef. in Oestr. Bot. Zeitschr. xiii. (1863) 14 = **Hibiscus**, Linn. (Malvac.).
corymbosa, Alef. l. c. 15 = H. corymbosus.
dongolensis, Alef. l. c. = H. lunariifolius.
macrantha, Alef. l. c. = H. lunariifolius.

COUBLANDIA, Aubl. Pl. Gui. ii. 937. t. 356 (1775) = **Muellera**, Linn. (Legumin.).
frutescens, Aubl. l. c. = M. moniliformis.
mexicana, Baill. Adansonia, ix. (1870) 238.—Mexic.

COUDENBERGIA, March. in Bull. Acad. Belg. Sér. II. xlvii. (1879) 514 = **Coemansia**, March. (Araliac.).

COUEPIA, Aubl. Pl. Gui. i. 519. t. 221 (1775). *ROSACEAE*, Benth. & Hook. f. i. 608.
ACIA, Willd. Sp. Pl. iii. 717 (1800).
ACIOA, Aubl. Pl. Gui. ii. 698. t. 280 (1775).
DULACIA, Neck. Elem. ii. 414 (1790).
bracteosa, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Guiana.

COUEPIA :—
canomensis, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Bras.
chrysocalyx, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 42.—Bras.
comosa, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Guiana.
eriantha, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. II. 45.—Bras.
glandulosa, *Miq. Stirp. Surin. Sel.* 28.—Guian.
glaucescens, *Spruce, ex Hook. f. in Mart. Fl. Bras.* xiv. II. 49.—Bras.
grandiflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Bras.
guianensis, *Aubl. Pl. Gui.* i. 519, 207.—Guiana.
hypoleuca, *Miq. in Linnaea*, xix. (1847) 438.—Mexic.
Kunthiana, *Benth. in Hook. Journ. Bot.* ii. (1840) 216.—Mexic.
leptostachya, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 44.—Bras.
macrophylla, *Spruce, ex Hook. f. l. c.* 41.—Peruv.
magnoliaefolia, *Benth. ex Hook. f. l. c.* 43.—Bras.
Martiana, *Hook. f. l. c.* 47.—Bras.
multiflora, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Guiana.
myrtifolia, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 44.—Bras.
ovatifolia, *Benth. in Hook. Journ. Bot.* ii. (1840) 216.—Bras.
paraënsis, *Benth. l. c.*—Bras.
Parillo, *DC. Prod.* ii. 526.—Guiana.
racemosa, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 43.—Bras.
Steudeliana, *Miq. Stirp. Surin. Sel.* 28.—Guiana.
subcordata, *Benth. ex Hook. f. in Mart. Fl. Bras.* xiv. II. 46.—Bras.
thyrsiflora, *Hook. f. l. c.* 45.—Bras.
vismioides, Klotzsch, ex Hook. f. in Mart. Fl. Bras. xiv. II. 48 = paraënsis.
Uiti, *Benth. in Hook. Journ. Bot.* ii. (1840) 215.—Bras.

COULA, Baill. Adansonia, iii. 61, 64. t. 3 (1862). *OLACINEAE*, Benth. & Hook. f. i. 995.
edulis, *Baill. l. c.* 64.—Afr. trop.

COULEJIA, Dennst. Schluess. Hort. Malab. 31 (1818) = **Antidesma**, Linn. (Euphorb.).
amentacea, Dennst. l. c.—Ind. or.

COULTERIA, H. B. & K. Nov. Gen. et Sp. vi. 328. t. 568 = **Caesalpinia**, Linn. (Legumin.).
africana, Guill. & Perr. Fl. Seneg. Tent. i. 256 = Prosopis oblonga.
chilensis, DC. Prod. ii. 481 = Caesalp. tinctoria.
horrida, H. B. & K. Nov. Gen. et Sp. vi. 330 = Caesalp. horrida.
mexicana, DC. Prod. ii. 481 = Caesalp. vesicaria.
mollis, H. B. & K. Nov. Gen. et Sp. vi. 330 = Caesalp. mollis.
tinctoria, H. B. & K. l. c. 331 = Caesalp. tinctoria.

COUMA, Aubl. Pl. Gui. Suppl. 39. t. 392 (1775). *APOCYNACEAE*, Benth. & Hook. f. ii. 693.
COLLOPHORA, Mart. in Buchn. Rep. Pharm. xxxv. (1830) 196, ex Bull. Férussac (Janv. 1831), 64.
dulcis, Spruce, ex Muell. Arg. in Mart. Fl. Bras. vi. I. 19 = utilis.
guianensis, *Aubl. Pl. Gui. Suppl.* 39. t. 392.—Guiana.
rigida, *Muell. Arg. in Mart. Fl. Bras.* vi. I. 20.—Bras.
utilis, *Muell. Arg. l. c.* 19.—Bras.

COUMAROUNA, Aubl. Pl. Gui. ii. 740. t. 296 (1775) = = **Dipteryx**, Schreb. (1791) (Legumin.).
odora, Aubl. l. c. t. 296 = seq.
odorata, Aubl. l. c. 740 = D. odorata.

COUPIA, G. Don, Gen. Syst. ii. 44 (1832) sphalm. = **Goupia**, Aubl. (Celastrin.).

COUPOUI, Aubl. Pl. Gui. Suppl. 16. t. 377 (1775). *APOCYNACEAE* [Benth. & Hook. f. i. 696].
CUPIRANA, Miers, Apocyn. S. Am. 15 (1878).
CUPUIA, Rafin. Princ. Somiol. 29 (1814).
aquatica, *Aubl. l. c.*—Guiana.

COURALIA, Splitg. in Hoev. & De Vriesc, Tijdschr. ix. (1842) 14. *BIGNONIACEAE*, Benth. & Hook. f. ii. 1042.
fluviatilis, *Splitg. l. c.* 15.—Guiana.

COURANTIA, Lem. Jard. Fleur. i. (1851) Misc. 92 = **Cotyledon**, Tourn. (Crassulac.).
echeverioides, Lem. l. c. = Cotyl. roscata.
rosea, Lem. l. c. iii. (1853) 244, in nota = Cotyl. rosea.

COURATARI, Aubl. Pl. Gui. ii. 723. t. 290 (1775). *MYRTACEAE*, Benth. & Hook. f. i. 722.
AMPHORICARPUS, Spruce, ex Miers, in Trans. Linn. Soc. xxx. (1874) 288.
CARINIANA, Casar. Nov. Stirp. Bras. 35 (1842).
aulacocarpa, *Mart. ex Berg, in Mart. Fl. Bras.* xiv. i. 509.—Bras.
coriacea, *Mart. ex Berg, l. c.* 510.—Bras.
dictyocarpa, *Mart. ex Berg, l. c.*—Bras.
domestica, *Mart. in Flora*, xx. (1837) II. *Beibl.* 127, in textu.—Bras.
estrellensis, *Raddi, in Mem. Mod.* xviii. *Fis.* (1820) 403.—Bras.
guianensis, *Aubl. Pl. Gui.* ii. 724. t. 290.—Am. trop.
Lecythopsis, *Mart. in Flora*, xx. (1837) II. *Beibl.* 89.—Bras.
legalis, *Mart. l. c.* 88.—Bras.
lineata, *Mart. ex Berg, in Mart. Fl. Bras.* xiv. i. 508.—Bras.
macrocarpa, *Mart. ex Berg, l. c.* 509.—Bras.
Martiana, *Miers, in Trans. Linn. Soc.* xxx. (1874) 282.—Bras.
paraënsis, *Mart. ex Berg, in Mart. Fl. Bras.* xiv. i. 510.—Bras.
rubra, *Gardn. ex Miers, in Trans. Linn. Soc.* xxx. (1874) 288.—Bras.
Tauari, *Berg, in Mart. Fl. Bras.* xiv. i. 509.—Bras.
uaupensis, *Spruce, ex Berg, l. c.* 508.—Bras.
Vriesii, *Miers, in Trans. Linn. Soc.* xxx. (1874) 284.—Guiana.

COURATARI, Cambess. in St. Hil. Fl. Bras. Mer. ii. 379 (1829) = **Lecythopsis**, Schrank (Myrtac.).
glabra, Cambess. l. c. = L. glabra.
rufescens, Cambess. l. c. 380 = L. rufescens.

COURBARI, Adans. Fam. ii. 317 (1763) = **Hymenaea**, Linn. (Legumin.).

COURBARIL, Plum. ex Endl. Gen. 1317 (1841) = praec.

COURBONIA, Brongn. in Bull. Soc. Bot. Fr. vii. (1860) 901. *CAPPARIDEAE*, Benth. & Hook. f. i. 969.
PHYSANTHEMUM, Klotzsch, in Peters, Reise Mossamb. Bot. 167. t. 29 (1861).
decumbens, *Brongn. l. c.*—Afr. trop.
virgata, *Brongn. l. c.*—Afr. trop.

COURIMARI, Aubl. Pl. Gui. Suppl. 27 (1775) (Gen. dub. affin.).
guianensis, *Aubl. l. c.*—Guiana.

COURINGIA, Adans. Fam. ii. 418 (1763) = **Conringia**, Heist. (Crucif.).

COUROUPITA, Aubl. Pl. Gui. ii. 708. t. 282 (1775). *MYRTACEAE*, Benth. & Hook. f. i. 722.
ELSHOLTZIA, Neck. Elem. ii. 256 (1790).
PONTOPIDANA, Scop. Introd. 195 (1777).
antillana, *Miers, in Trans. Linn. Soc.* xxx. (1874) 191.—Ind. occ.
crenulata, *Miers, l. c.* 195.—Bras.
guianensis, *Aubl. Pl. Gui.* ii. 708. t. 282.—Guian.
guianensis, Hook. Bot. Mag. t. 3158 = antillana.
lentula, *Miers, in Trans. Linn. Soc.* xxx. (1874) 194.—Bras.
membranacea, *Miers, l. c.*—N. Granat.
nicaraguarensis, *DC. Prod.* iii. 294.—Am. centr.
odoratissima, *Seem. Bot. Voy. Herald*, 126.—Panama.
peruviana, *Berg, in Linnaea*, xxxi. (1861-62) 261.—Peruv.
surinamensis, *Mart. ex Berg, in Mart. Fl. Bras.* xiv. i. 476.—Guian.

COURRANTIA, Sch. Bip. in Webb & Berth. Phyt. Canar. ii. 276. t. 89 (cir. 1845) = **Matricaria**, Linn. (Compos.).
chamomilloides, Sch. Bip. l. c. 278 = M. Courrantia.

COURSETIA, DC. in Ann. Sc. Nat. Sér. I. iv. (1825) 92. *LEGUMINOSAE*, Benth. & Hook. f. i. 501.
POISSONIA, Baill. Adansonia, ix. (1870) 295.
arborea, *Griseb. Fl. Brit. W. Ind.* 183.—Ins. Trinit.
dubia, *DC. Prod.* ii. 264.—Am. austr.
criantha, *Benth. in Hook. Ic. Pl.* sub t. 1065.—Peruv.
glandulosa, *A. Gray, in Proc. Am. Acad.* v. (1861) 156.—Calif.
grandiflora, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 10.—Peruv.
microphylla, *A. Gray, in Proc. Am. Acad.* xvii. (1881-82) 201.—Arizona.
orbicularis, *Benth. in Hook. Ic. Pl.* t. 1065.—Peruv.
rostrata, *Benth. in Mart. Fl. Bras.* xv. i. 44.—Am. trop.
tomentosa, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 92.—Peruv.
vicioides, *Benth. in Mart. Fl. Bras.* xv. i. 44.—Am. trop.
virgata, *DC. in Ann. Sc. Nat.* Sér. I. iv. (1825) 92.—Mexic.

COURTENIA, R. Br. in Benn. Pl. Jav. Rar. 236 (1844) = **Cola**, Schott (Sterculiac.).
Afzelii, R. Br. l. c. = Cola Afzelii.
heterophylla, R. Br. l. c. = Cola heterophylla.
triloba, R. Br. l. c. = Cola heterophylla.

COURTOISIA, Nees, in Linnaea, ix. (1834) 286. *CYPERACEAE*, Benth. & Hook. f. iii. 1046.
cyperoides, *Nees, l. c.*—Ind. or.
olivacea, *Boeck. in Flora*, xliv. (1861) 331.—Guiana.

COUSINIA, Cass. in Dict. Sc. Nat. xlvii. 503 (1827). *COMPOSITAE*, Benth. & Hook. f. ii. 467.
AUCHERA, DC. Prod. vi. 557 (1837).
acicularis, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 319.—Turkestan.
Actinia, *Boiss. Diagn.* Ser. II. iii. 55.—Afghan.
actinocephala, Jaub. & Spach, Illustr. ii. 93. t. 175 = xiphiolepis.
adnata, *Bunge, Gatt. Cous.* 50.—Persia.
affinis, *Schrenk, Enum. Pl. Nov.* i. 41.—Soongaria.
aggregata, *DC. Prod.* vi. 553.—Persia.
aintabensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 503.—Syria.
alata, *Schrenk, Enum. Pl. Nov.* i. 40.—Reg. Casp.; Soongaria.
Alberti, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 315.—Turkestan.
albicaulis, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 128.—Persia.
albida, *DC. Prod.* vi. 552.—Persia.
Alepideae, *Boiss. Diagn.* Ser. II. iii. 53.—Beluchist.
aleppica, *Boiss. l. c.* Ser. I. x. 101.—Syria.
alpicola, Borszczow, ex Trautv. in Act. Hort. Petrop. viii. (1883) 484 = seq.
alpina, *Bunge, Del. Sem. Hort. Dorp.* (1845) 3.—Turkest.
amplissima, *Boiss. Fl. Orient.* iii. 462.—Persia.
anisoptera, Jaub. & Spach, Illustr. ii. 88. t. 170 = calolepis.
anomala, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 316.—Turkestan.
anthacantha, Regel, in Bull. Soc. Nat. Mosc. xl. (1867) II. 156, sphalm. *vide* xanthacantha.
aptera, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 170.—Afghan.
arachnoidea, *Fisch. & Mey. ex DC. Prod.* vi. 553.—Soongar.
aralensis, Bunge, in Mém. Sav. Etr. Pétersb. vii. (1851) 354 = minuta.
araneosa, *DC. Prod.* vi. 556.—Persia.
arctioides, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* ii. (1844) 115.—Soongaria.
arctotidifolia, *Bunge, Gatt. Cous.* 42.—Persia.
arenaria, *Bunge, l. c.* 29.—Persia.
assyriaca, Jaub. & Spach, Illustr. ii. 87. t. 179 = leucochlora.
atropatana, *Bunge, Gatt. Cous.* 17.—Persia.
Aucheri, *DC. Prod.* vi. 555.—Cappadoc.
[illegible]ulata, *Boiss. Diagn.* Ser. II. iii. 58.—Ind. bor.
ma[illegible]
pur[illegible]*lata*, Hook. f. Fl. Brit. Ind. iii. 360 = buphthalmoides.
sent[illegible]s.
tinct[illegible]*oiss. Diagn.* Ser. II. iii. 53.—Afghan.
villos[illegible]*a, Boiss. & Haussk. in Boiss. Fl. Orient.* iii.
virgin[illegible]ersia.

COUSINIA :—
Belangeri, *DC. Prod.* vi. 553.—Persia.
Bienerti, *Bunge, Gatt. Cous.* 27.—Persia.
bipinnata, *Boiss. Diagn.* Ser. II. iii. 52.—Afghan.
Boissieri, *Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 127.—Persia.
Bonvaleti, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 320.—Turkestan.
brachyptera, *DC. Prod.* vi. 554.—Armenia.
Brandisii, C. B. Clarke, Comp. Ind. 214 = Thomsoni.
Buhsei, Bunge, Gatt. Cous. 22 = hamosa.
bulgarica, *C. Koch, in Linnaea*, xxiv. (1851) 385.—Europ. or.
Bungeana, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 318.—Turkestan.
buphthalmoides, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 155.—Afghan.; Ind. bor. occ.
cacsarca, *Boiss. & Bal. Diagn.* Ser. II. vi. 116.—Cappadoc.
calcitrapaeformis, Jaub. & Spach, Illustr. ii. 76. t. 158 = minuta.
Calcitrapa, *Boiss. Fl. Orient.* iii. 490.—Persia.
calocephala, *Jaub. & Spach, Illustr.* ii. 96. t. 178.—Persia.
calolepis, *Boiss. Diagn.* Ser. I. vi. 116.—Persia.
calolepis, Boiss. Fl. Orient. iii. 480 = commutata.
Candolleana, *Jaub. & Spach, Illustr.* ii. 74. t. 156.—Persia.
canescens, *DC. Prod.* vi. 556.—Armenia.
canescens, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 323.—Turkestan.
Capusi, *Franch. l. c.* 317.—Turkestan.
carduiformis, *Cass. in Dict. Sc. Nat.* xlvii. 503.—Reg. Caucas.
carduncelloidea, *Regel & Schmalh. in Pl. Nov. Fedsch.* 50.—As. centr.
carlinoides, Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc. xii. (1860) 126 = obovata.
carlinoides, *DC. Prod.* vi. 554.—Persia.
carthamoides, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 171.—Afghan.
cataonica, *Boiss. & Haussk. Fl. Orient.* iii. 483.—As. Min.
centauroides, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1839) 159.—Persia.
centauroides, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 152.—As. centr.
Chamaepeuce, *Boiss. Diagn.* Ser. I. vi. 120.—Persia.
chlorocephala, *C. A. Mey. ex DC. Prod.* vi. 554.—Armenia.
chrysacantha, *Jaub. & Spach, Illustr.* ii. 78. t. 160.—Persia.
cirsioides, *Boiss. & Bal. Diagn.* Ser. II. vi. 117.—Cappadoc.
commutata, *Bunge, Gatt. Cous.* 24.—Persia.
concinna, *Boiss. & Haussk. in Boiss. Fl. Orient.* iii. 513.—Persia.
concolor, *Bunge, Gatt. Cous.* 33.—Persia.
congesta, *Bunge, l. c.* 12.—Persia.
coronata, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 321.—Turkestan.
crispa, *Jaub. & Spach, Illustr.* ii. 81. t. 163.—Persia.
cylindracea, *Boiss. Diagn.* Ser. I. vi. 115.—Persia.
cylindrocephala, *Jaub. & Spach, Illustr.* ii. 84. t. 166.—Persia.
cymbolepis, *Boiss. Diagn.* Ser. I. vi. 119.—Kurdist.
cynaroides, *C. A. Mey. Verz. Pfl. Cauc.* 230.—Reg. Caucas.
decipiens, *Boiss. & Buhse, in Nouv. Mém Soc. Nat. Mosc.* xii. (1860) 125.—Persia.
decurrens, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 154.—Afghan.
deserti, *Bunge, Gatt. Cous.* 29.—Persia.
dichotoma, *Bunge, Del. Sem. Hort. Dorpat.* (1845) 3.—Turkest.
discolor, *Bunge, Gatt. Cous.* 34.—Persia.
dissecta, *Kar. & Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) 391.—Turcoman.
dolicholepis, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 107.—Soongaria.
clata, *Boiss. & Buhse, in Nouv. Mém. Soc. Nat. Mosc.* xii. (1860) 126.—Persia.
elegans, *Aitch. & Hemsl. in Journ. Linn. Soc.* xix. (1882) 172.—Afghan.
erinacea, *Jaub. & Spach, Illustr.* ii. 83. t. 165.—Persia.
eriobasis, *Bunge, Gatt. Cous.* 37.—Persia.
eriocephala, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 504.—As. Min.

COUSINIA :—
eriophora, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 313.—Turkestan.
eryngioides, *Boiss. Diagn.* Ser. I. x. 101.—Persia.
Falconeri, *Hook. f. Fl. Brit. Ind.* iii. 360.—Reg. Himal.
flavispina, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 315.—Turkestan.
foliosa, *Boiss. & Bal. Diagn.* Ser. II. vi. 116.—Cappadoc.
gracilis, *Boiss. l. c.* Ser. I. vi. 117.—Persia.
grandiceps, *Bunge, Gatt. Cous.* 54.—Persia.
grandis, *C. A. Mey. ex DC. Prod.* vi. 557.—Persia.
Griffithiana, *Boiss. Diagn.* Ser. II. iii. 55.—Afghan.
Gundelia, C. Koch, in Linnaea, xxiv. (1851) 387 = brachyptera.
Hablitzii, *C. A. Mey. ex DC. Prod.* vi. 554.—Persia.
hamosa, *C. A. Mey. l. c.* 555.—Persia.
hamulosa, *C. A. Mey. ex Boiss. Fl. Orient.* vi. 483 = hamosa.
heliantha, *Bunge, Gatt. Cous.* 39.—Persia.
Hermonis, *Boiss. Diagn.* Ser. I. x. 102.—Syria.
heterolepis, C. Koch, in Linnaea, xxiv. (1851) 386 = brachyptera.
heterophylla, *Boiss. Diagn.* Ser. II. iii. 54.—Afghan.
Hohenackeri, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* ii. 34.—Persia.
hololeuca, *Bunge, Gatt. Cous.* 27.—Persia.
humilis, *Boiss. in Ann. Sc. Nat.* Sér. IV. ii (1854) 248.—Cappadoc.
hypoleuca, *Boiss. Diagn.* Ser. I. x. 103; *Fl. Orient.* iii. 464.—Persia.
Hystrix, C. B. Clarke, Comp. Ind. 214 = microcarpa.
Hystrix, *C. A. Mey. Ind. Sem. Hort. Petrop.* ii. 34.—Persia.
ilicifolia, *Jaub. & Spach, Illustr.* ii. 82. t. 164.—Persia.
incompta, *DC. Prod.* vi. 553.—Persia.
inflata, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 512.—Kurdistan.
integrifolia, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 322.—Turkestan.
intermedia, *C. A. Mey. ex DC. Prod.* vi. 553.—Armenia.
involucrata, *Boiss. Diagn.* Ser. I. vi. 114.—Persia.
karatavica, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 317.—Turkestan.
Karelini, Less. ex Kar. in Bull. Soc. Nat. Mosc. (1839) 159 = onopordioides.
kokanica, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 49.—As. centr.
Kornhuberi, *Heimerl, in Denkschr. Acad. Wien*, l. (1885) 62.—Persia.
Korolkowi, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 315.—Turkestan.
Kotschyi, *Boiss. Diagn.* Ser. I. vi. 120.—Persia.
Krauseana, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 48.—As. centr.
lachnosphaera, *Bunge, Gatt. Cous.* 36.—Persia.
lappacea, *Schrenk, Enum. Pl. Nov.* i. 42.—Soongaria.
lasiandra, *Bunge, Gatt. Cous.* 40.—Persia.
lasiolepis, *Boiss. Diagn.* Ser. I. vi. 118.—Persia.
lepida, *Bunge, Gatt. Cous.* 26, in nota.—Persia.
leptocephala, *Fisch. & Mey. in Bull. Soc. Nat. Mosc.* (1839) 159.—Turcoman.
leucochlora, *Bunge, Gatt. Cous.* 24, in nota.—Persia.
libanotica, *DC. Prod.* vi. 556.—Syria.
lucida, *DC. l. c.* 553.—Assyria.
lyrata, *Bunge, Gatt. Cous.* 53.—Persia.
macrocephala, *C. A. Mey. Verz. Pfl. Cauc.* 231.—Reg. Caucas.
macrolepis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 505.—Persia.
macroptera, *C. A. Mey. ex DC. Prod.* vi. 555.—Armenia.
microcarpa, *Boiss. Diagn.* Ser. II. iii. 59.—Persia; Reg. Himal.
microcephala, *C. A. Mey. ex DC. Prod.* vi. 552.—Persia.
minuta, *Boiss. Diagn.* Ser. I. vi. 117.—Persia; Ind. bor. occ.
mollis, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 108.—Soongaria.
monocephala, *Bunge, Gatt. Cous.* 52.—Persia.
multiloba, *DC. Prod.* vi. 554.—Persia.
neurocentra, *Bunge, Gatt. Cous.* 29.—Persia.
Noëana, *Boiss. Diagn.* Ser. II. iii. 57.—Persia.
obovata, *Boiss. Fl. Orient.* iii. 491.—Persia.
odontolepis, *DC. Prod.* vi. 556.—Assyria.

COUSINIA :—
odontolepis, Sch. Bip. ex Boiss. l. c. 512 = cymbolepis.
Olgae, *Regel & Schmalh. in Pl. Nov. Fedsch.* 48.—As. centr.
oligocephala, *Boiss. Diagn.* Ser. I. vi. 116.—Persia.
oligocephala, Haussk. ex Boiss. Fl. Orient. iii. 475 = bachtiarica.
Olivieri, *DC. Prod.* vi. 552.—Persia?
Olivieri, Jaub. & Spach, Illustr. t. 172 = serratuloides.
onopordioides, *Ledeb. in Eichw. Pl. Casp. Cauc.* 40. t. 37.—Turcoman.
orientalis, *C. Koch, in Linnaea*, xxiv. (1851) 386.—Oriens.
outichaschensis, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 322.—Turkestan.
palmatiloba, Jaub. & Spach, Illustr. ii. 80. t. 162 = multiloba.
pauciflora, Bunge, Gatt. Cous. 12 = aggregata.
pentacantha, *Regel & Schmalh. in Act. Hort. Petrop.* vi. (1879) 315.—As. centr.
pergamacea, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 513.—Persia.
Pestalozzae, *Boiss. l. c.* 471.—Syria.
pinarocephala, *Boiss. Diagn.* Ser. I. vi. 121.—Persia.
piptocephala, *Bunge, Gatt. Cous.* 26.—Persia.
platyacantha, Bunge, l. c. 37 = centauroides.
platylepis, *Fisch. Mey. & Avé-Lall. Ind. Sem. Hort Petrop.* ix. *Suppl.* 10.—Turkestan.
polycephala, *Rupr. Sert. Tiansch.* 54.—As. centr.
prasina, *Jaub. & Spach, Illustr.* ii. 86. t. 168.—Persia.
Princeps, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 324.—Turkestan.
prolifera, *Jaub. & Spach, Illustr.* ii. 74. t. 157.—Persia.
pterocarpa, *Boiss. Fl. Orient.* iii. 501.—Afghan.
pugionifera, *Jaub. & Spach, Illustr.* ii. 92. t. 174.—Persia.
pulchella, *Bunge, Del. Sem. Hort. Dorpat.* (1841) 3.—As. centr.
purpurea, *C. A. Mey. ex DC. Prod.* vi. 555.—Persia.
pycnoloba, Boiss. Diagn. Ser. II. iii. 57 = multiloba.
pyrrhocephala, Jaub. & Spach, Illustr. ii. 85. t. 167 = Olivieri.
racemosa, *Boiss. Diagn.* Ser. II. iii. 56.—Afghan.
radians, *Bunge, Del. Sem. Hort. Dorpat.* (1845) 3.—As. centr.
ramosissima, *DC. Prod.* vi. 552.—Cappadoc.
recurvata, *DC. l. c.* 555.—Persia.
sakawensis, *Boiss. & Haussk. ex Boiss. Fl. Orient.* iii. 488.—Persia.
Scala, *Aitch. & Hemsl. in Journ. Linn. Soc.* ix. (1882) 172.—Afghan.
scariosa, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 154.—Afghan.
Schtschurowskiana, *Regel & Schmalh. ex Regel, Pl. Nov. Fedsch.* 50.—As. centr.
Schultziana, Jaub. & Spach, Illustr. ii. 95. t. 177 = cymbolepis.
Seidlitzii, *Bunge, Gatt. Cous.* 20.—Persia.
Semenovii, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 147.—As. centr.
serratuloides, *Boiss. Diagn.* Ser. I. vi. 118.—Persia.
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 151.—Turkestan.
silyboides, Jaub. & Spach, Illustr. ii. 94. t. 175 = Kotschyi.
Smirnowii, *Trautv. in Act. Hort. Petrop.* viii. (1883) 488.—As. centr.
sphaerocephala, Jaub. & Spach, Illustr. ii. 79. t. 161 = C. Chamaepeuce.
squarrosa, *Boiss. Diagn.* Ser. I. x. 102.—Persia.
squarrosa, C. Koch, in Linnaea, xxiv. (1851) 387 = brachyptera.
stenocephala, *Boiss. Diagn.* Ser. I. vi. 115.—Assyria.
stenolepis, *Regel & Schmalh. in Pl. Nov. Fedsch.* 49.—As. centr.
submutica, *Franch. in Ann. Sc. Nat.* Sér. VI. xvi. (1883) 314.—Turkestan.
sylvicola, Bunge, Del. Sem. Hort. Dorp. (1845) 3 = dichotoma.
tabrisiana, *Bunge, Gatt. Cous.* 16.—Persia.
tenella, *Fisch. & Mey. Ind. Sem. Hort. Petrop.* i. [illegible] Armenia.
tenuifolia, *C. A. Mey. ex DC. Prod.* vi. 554.—P[illegible]
thamnodes, *Boiss. & Haussk. in Boiss. Fl. Or[illegible]* 471.—Persia.
Thomsoni, *C. B. Clarke, Comp. Ind.* 240) 215. Himal.

COUSINIA :—
trachylepis, *Bunge, Gatt. Cous.* 22.—Persia.
Trautvetteri, *Regel, in Act. Hort. Petrop.* v. (1877) 257.—Turkestan.
triflora, *Schrenk, in Bull. Phys.-Math. Acad. Pétersb.* iii. (1845) 108.—Soongaria.
umbrosa, *Bunge, Gatt. Cous.* 10.—Persia.
uncinata, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 145.—Soongaria.
verbascifolia, *Bunge, Gatt. Cous.* 52.—Persia.
verticillaris, *Bunge, in Mém. Sav. Etr. Pétersb.* vi. (1851) 357.—As. centr.
Verutum, *Bunge, Gatt. Cous.* 31.—Persia.
wolgensis, *C. A. Mey. ex DC. Prod.* vi. 555.—Rossia.
xanthacantha, *Regel, in Bull. Soc. Nat. Mosc.* xl. (1867) II. 156 (*sphalm.* anthacantha).—Oriens.
xeranthemoides, Bunge, ex Boiss. Fl. Orient. iii. 497 = microcarpa.
xiphiolepis, *Boiss. Diagn.* Ser. I. vi. 121.—Persia.

COUSSAPOA, Aubl. Pl. Gui. ii. 955. t. 362, 363 (1775). *URTICACEAE*, Benth. & Hook. f. iii. 379.
acutifolia, *Klotzsch, in Linnaea*, xx. (1847) 529.—Peruv.
angustifolia, *Aubl. Pl. Gui.* ii. 956. t. 363.—Guiana.
asperifolia, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 96.—Guiana.
cuneata, *Miq. in Mart. Fl. Bras.* iv. I. 138.—Guiana.
dealbata, *André, Illustr. Hortic.* xvii. (1870) 16.—Bras.
fagifolia, *Klotzsch, in Linnaea*, xx. (1847) 528.—Guiana.
ferruginea, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 93.—Guiana.
Fontanesiana, *A. Brongn. ex Tréc. l. c.* 94.—Bras.
hirsuta, *Tréc. l. c.* 97.—Peruv.
intermedia, *Mart. ex Miq. in Mart. Fl. Bras.* iv. I. 133.—Bras.
laevigata, *Poepp. & Endl. Gen. et Sp.* ii. 33.—Bras.
latifolia, *Aubl. Pl. Gui.* ii. 955. t. 362.—Guiana.
magnifolia, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 98.—Peruv.
Martiana, *Miq. in Mart. Fl. Bras.* iv. I. 132.—Bras.
microcephala, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 96.—Guiana.
nitida, *Miq. in Mart. Fl. Bras.* iv. I. 133.—Bras.
obovata, *Miq. in Versl. en Medel. Kon. Nederl. Inst.* 1842 (1843) 200.—Hab.?
ovalifolia, *Tréc. in Ann. Sc. Nat.* Sér. III. viii. (1847) 95.—Peruv.
puberula, *Klotzsch, in Linnaea*, xx. (1847) 529.—Peruv.
Ruizii, *Klotzsch, l. c.* 529.—Peruv.
Schottii, *Miq. in Mart. Fl. Bras.* iv. I. 137.—Bras.
setosa, *Klotzsch, in Linnaea*, xx. (1847) 528.—Peruv.
subincana, *Mart. ex Miq. in Mart. Fl. Bras.* iv. I. 134.—Bras.
vellerea, *Klotzsch, in Linnaea*, xx. (1847) 527.—Peruv.
villosa, *Poepp. & Endl. Nov. Gen. et Sp.* ii. 33. t. 147.—Peruv.

COUSSAREA, Aubl. Pl. Gui. i. 98. t. 38 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 120.
BILLARDIERIA, Vahl, Eclog. Am. i. 13. t. 10 (1796).
FROELICHIA, Vahl, l. c. Praef. 3 (1796).
PECHEYA, Scop. Introd. 143 (1777).
accedens, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
albescens, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 87, in nota.—Guiana.
ampla, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
bahiensis, *Muell. Arg. l. c.* 467.—Bras.
biflora, *Muell. Arg. l. c.* 468.—Bras.
capitata, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
Catingana, *Muell. Arg. in Flora*, lviii. (1875) 468.—Bras.
coffeoides, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 89.—Bras.
congestiflora, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
contracta, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
corcovadensis, *Muell. Arg. in Flora*, lviii. (1875) 468.—Bras.
cornifolia, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
flava, *Poepp. & Endl. Nov. Gen. et Sp.* iii. 26. t. 231.—Peruv.

COUSSAREA:—
Froelichia, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 177.—Ins. Trinit.
graciliflora, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
grandis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 100.—Bras.
hydrangeaefolia, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
ilheotica, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
lagoënsis, *Muell. Arg. l.c.*—Bras.
lanceolata, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 104.—Bras.
leptoloba, *Muell. Arg. in Flora*, lviii. (1875) 468.—Bras.
leptophragma, *Muell. Arg. l. c.* 466.—Bras.
leptopus, *Muell. Arg. l. c.* 465.—Bras.
longiflora, *Muell. Arg. l.c.* 468.—Bras.
longifolia, *Muell. Arg. l.c.*—Bras.
macrocarpa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 177.—Guiana.
macrophylla, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
Martini, *Benth. & Hook. f. Gen.* ii. 121.—Am. trop.
meridionalis, Muell. Arg. in Mart. Fl. Bras. vi. v. 85 = nemoralis.
nemoralis, *Benth. & Hook. f. Gen.* ii. 121.—Bras.
nodosa, *Benth. & Hook. f. l. c.*—Bras.
obscura, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
pentamera, *Karst. Fl. Columb.* ii. 13. t. 107.—Costa Rica.
platyphylla, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
porophylla, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 87.—Bras.
procumbens, *Muell. Arg. l.c.* 104.—Bras.
racemosa, *A. Rich. in Mém. Soc. Hist. Nat. Par.* v. (1834) 177.—Guiana.
Regnelliana, *Muell. Arg. in Flora*, lviii. (1875) 466.—Bras.
Schomburgkiana, Benth. & Hook. f. Gen. ii. 121 = violacea.
squamosa, *Lam. Tabl. Encyc.* i. 281.—Ind. occ.
strigosipes, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
triflora, *Muell. Arg. l. c.* 48.—Bras.
uniflora, *Gardn. in Hook. Lond. Journ. Bot.* iv. (1845) 107.—Bras.
verticillata, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
violacea, *Aubl. Pl. Gui.* i. 48. t. 38.—Guiana.
virens, *Muell. Arg. in Flora*, lviii. (1875) 467.—Bras.
viridis, *Muell. Arg. in Mart. Fl. Bras.* vi. v. 104.—Bras.

COUTAREA, Aubl. Pl. Gui. i. 314. t. 122 (1775). *RUBIACEAE*, Benth. & Hook. f. ii. 42.
alba, *Griseb. in Goett. Abh.* xxiv. (1879) 153.—Reg. Argent.
Campanilla, *DC. Prod.* iv. 350.—Venezuela.
corymbosa, *Brign. in Mem. Soc. Ital. Mod.* Ser. II. i. (1862) 69.—Bras.
diervilloides, *Planch. & Linden, Cat.* n. 9 (1854) 3.—N. Granat.
flavescens, *Moç. & Sesse, ex DC. Prod.* iv. 350.—Mexic.
latifolia, *Moç. & Sesse, ex DC. l. c.*—Mexic.
Lindeniana, *Baill. Adansonii*, xii. (1879) 300.—Venezuela.
Lumaeana, *Baill. l. c.* 301.—Guatemala.
mexicana, *Zucc. & Mart. ex DC. Prod.* iv. 350.—Mexic.
mollis, *Cham. in Linnaea*, ii. (1834) 259.—Bras.
pubescens, *Pohl, Pl. Bras. Desc.* ii. 148. t. 200.—Bras.
Scherffiana, *André, in Illustr. Hortic.* xxv. (1878) 120. t. 321.—N. Granat.
speciosa, *Aubl. Pl. Gui.* i. 314. t. 122.—Guiana.

COUTHOVIA, A. Gray, in Proc. Am. Acad. iv. (1858) 324; v. (1861) 320. *LOGANIACEAE*, Benth. & Hook. f. ii. 797.
corynocarpa, *A. Gray, l. c.* iv. (1858) 324.—Ins. Sandvic.
Seemanni, *A. Gray, l. c.* v. (1861) 320.—Ins. Fiji.

COUTOUBAEA, Ham. Pl. Ind. Occ. 18 (1825) = seq.

COUTOUBEA, Aubl. Pl. Gui. i. 72. t. 27, 28 (1775) *GENTIANEAE*, Benth. & Hook. f. ii. 812.
LIMBORCHIA, Scop. Introd. 139 (1777).
PICRIUM, Schreb. Gen. 791 (1791).
alba, Lam. Encyc. ii. 162 = spicata.
arenaria, Willd. ex Griseb. Gen. et Sp. Gent. 132 = racemosa.
densiflora, Mart. Nov. Gen. et Sp. ii. 111. t. 185 = spicata.
lutea, Steud. in Flora, xxvi. (1843) 765 = spicata.
minor, H. B. & K. Nov. Gen. et Sp. iii. 179 = spicata.
purpurea, Lam. Encyc. ii. 162 = ramosa.
racemosa, *G. F. W. Mey. Prim. Fl. Esseq.* 86.—Guiana.
ramosa, *Aubl. Pl. Gui.* i. 74. t. 28.—Am. trop.
reflexa, *Benth. in Ann. Nat. Hist.* ii. (1839) 442.—Guiana.
spicata, *Aubl. Pl. Gui.* i. 72.—Am. trop.
spicata, Griseb. in Journ. Linn. Soc. vi. (1862) 141 = reflexa.
spicata, H. B. & K. Nov. Gen. et Sp. iii. 179 = densiflora.
ternifolia, Cav. Ic. iv. 14 = densiflora.
verticillata, G. Don, in Loud. Hort. Brit. 48 = Enicostema littorale.
volubilis, *Mart. Nov. Gen. et Sp.* ii. 112.—Ins. S. Doming.

COUTUBEA, Steud. Nom. ed. II. i. 429 (1840) = **Coutoubea**, Aubl. (Gentian.).

COVALIA, Reichb. Consp. 94 (1828) = COVOLIA, Neck. = **Spermacoce**, Linn. (Rubiac.).

OVELIA, Endl. Gen. 527, in syn. (1838) = praec.

OVELLIA, Gasp. Nov. Gen. Fic. 10 (1844) = **Ficus**, Tourn. (Urticac.).
albipila, Miq. Fl. Ind. Bat. Suppl. 434 = F. mollis.
assamica, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 464 = F. hispida.
aurata, Miq. Fl. Ind. Bat. Suppl. 433 = F. chrysocarpa.
Barclayana, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 461. t. 7 B = F. Barclayana.
composita, Miq. Fl. Ind. Bat. i. II. 324 = F. rapiformis.
congesta, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 463 = F. congesta.
conglomerata, Miq. l. c. 459 = F. Cunia.
costata, Miq. l. c. 468 = F. costigera.
courtallensis, Miq. l. c. 463 = F. hispida.
cuneata, Miq. l. c. 466. t. 8 B = F. cuneata.
Cunia, Miq. l. c. 459 = F. Cunia.
cyrtophylla, Miq. l. c. 460 = F. cyrtophylla.
daemonum, Miq. l. c. 462 = F. hispida.
dasycarpa, Miq. l. c. 465 = F. hispida.
dasycaula, Miq. l. c. 460 = F. obscura.
didynama, Miq. Pl. Jungh. i. 65 = F. lepicarpa.
glomerata, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 465 = F. glomerata.
grandifolia, Miq. Fl. Ind. Bat. Suppl. 434 = F. rapiformis.
Griffithii, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 467 = F. heterophylla.
guttata, Wight, Ic. t. 1966 = F. guttata.
hispida, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 462 = F. hispida.
inaequiloba, Miq. l. c. 459 = F. Cunia.
lanceolata, Miq. l. c. 465 = F. lanceolata.
lepicarpa, Miq. Fl. Ind. Bat. i. II. 328 = F. lepicarpa.
leucopleura, Miq. l. c. 326 = F. leucatoma.
macrophylla, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 465 = F. Roxburghii.
microcarpa, Miq. l. c. 466. t. 9 A = F. Ribes.
mollis, Miq. l. c. 466 = F. glomerata.
oppositifolia, Gasp. Nov. Gen. Fic. 85 = F. hispida.
paniculata, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 467 = F. Ribes.
prostrata, Miq. l. c. 465 = F. prostrata.
semifera, Miq. l. c. = F. variegata.
septica, Miq. Fl. Ind. Bat. i. II. 328 = F. septica.
rapiformis, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 464 = F. rapiformis.
Ribes, Miq. Fl. Ind. Bat. i. II. 325 = F. Ribes.
roxburghii, Kurz, ex Teijsm. & Binn. in Tijdschr. Nederl. Ind. xxvii. (1864) 28 = F. botryocarpa.
scabrosa, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 462 = F. hispida.
botryocarpa, Miq. Pl. Jungh. 65 = F. leucantatoma.

COVELLIA:—
subdenticulata, Miq. Fl. Ind. Bat. i. II. 324 = Ficus subdenticulata.
subopposita, Miq. Pl. Jungh. 66 = F. fistulosa.
ternatana, Miq. Fl. Ind. Bat. i. II. 323 = F. ternatana.
tuberculata, Miq. in Zoll. Syst. Verz. Ind. Archip. 94, 99 = F. fistulosa.
umbonata, Miq. Fl. Ind. Bat. i. II. 323 = F. umbonata.
venosa, Miq. in Hook. Lond. Journ. Bot. vii. (1848) 468 = F. leucatoma.
volkameriaefolia, Miq. in Hook. l. c. 464. t. 8 A = F. lepicarpa.
Webbiana, Miq. l. c. 467 = F. Webbiana.
Wightiana, Miq. l. c. 463 = F. hispida.
Zollingeriana, Miq. l. c. 460 = F. coronata.

COVILHAMIA, Korth. in Nederl. Kruidk. Arch. i. (1848) 307. *STERCULIACEAE*?
ovata, *Korth. l. c.*—Borneo.

COVOLA, Medic. in Staatsw. Vorles. Churpf. Phys.-Oek. Ges. i. (1791) 227 = **Salvia**, Tourn. (Labiat.).
napifolia, Medic. Phil. Bot. ii. 67 = S. napifolia.
verticillata, Medic. l. c. = S. verticillata.

COVOLIA, Neck. Elem. i. 201 (1790) = **Spermacoce**, Linn. (Rubiac.).

COWANIA, D. Don, in Trans. Linn. Soc. xiv. (1825) 574. t. 22. *ROSACEAE*, Benth. & Hook. f. i. 618.
GREGGIA, Engelm. in Bot. Wisliz. Exped. 30 (1848).
ericaefolia, *Torr. ex A. Gray, Pl. Wright.* ii. 106.—Calif.
Havardia, *S. Wats. in Proc. Am. Acad.* xx. (1885) 364.—Texas.
mexicana, *D. Don, in Trans. Linn. Soc.* xiv. (1825) 575. t. 22. f. 1–6.—Am. bor. occ.
plicata, *D. Don, in Sweet, Brit. Flow. Gard.* Ser. II. t. 400.—Am. bor. occ.
plicata, Torr. in Frém. Rep. 314 = mexicana.
purpurea, Zucc. in Abh. Akad. Muench. iv. II. (1843) 7. t. 2 = plicata.
Stansburiana, Torr. in Stansb. Exped. Utah, 386 = mexicana.

COXIA, Endl. Gen. 733 (1839) = **Lysimachia**, Linn. (Primulac.).

CRABBEA, Harv. Gen. S. Afr. Pl. 276 (1838) = **Barleria**, Linn. (Acanthac.).
pungens, Harv. l. c. = B. pungens.

CRABBEA, Harv. in Hook. Lond. Journ. Bot. i. (1842) 26. *ACANTHACEAE*, Benth. & Hook. f. ii. 1092.
angustifolia, *Nees, in DC. Prod.* xi. 163.—Afr. austr.
cirsioides, Nees, l. c. 163 = nana.
hirsuta, *Harv. in Hook. Lond. Journ. Bot.* i. (1842) 27.—Afr. austr.
nana, *Nees, in DC. Prod.* xi. 162.—Afr. austr.
ovalifolia, *Ficalho & Hiern, in Trans. Linn. Soc.* Ser. II. ii. (1881) 24.—Afr. trop.

CRABOWSKIA, G. Don, Gen. Syst. iv. 480 (1836), sphalm. = **Grabowskia**, Schlecht. (Solan.).

CRACCA, Benth. ex Oerst. in Kjoeb. Vidensk. Meddel. (1853) 8. *LEGUMINOSAE*, Benth. & Hook. f. i. 501.
caribaea, *Benth. ex Oerst. l. c.* 9.—Ind. occ.
Edwardsii, *A. Gray, Pl. Wright.* ii. 35.—Mexic.
glabrescens, *Benth. ex Oerst. in Kjoeb. Vidensk. Meddel.* (1853) 9.—Ecuador.
glandulifera, *Benth. l. c.* 8.—Ecuador.
mollis, *Benth. l. c.* 9.—Peruv.
ochroleuca, *Benth. l. c.*—Peruv.
sericea, *A. Gray, in Proc. Am. Acad.* xix. (1883) 74.—N. Mexic.

CRACCA, Linn. Fl. Zeyl. 139 (1747) = **Tephrosia**, Pers. (1807) Leguminos.
maxima, Linn. Sp. Pl. ed. I. 752 = T. purpurea.
purpurea, Linn. l. c. = T. purpurea.
senticosa, Linn. l. c. = T. senticosa.
tinctoria, Linn. l. c. = T. tinctoria.
villosa, Linn. l. c. = T. villosa.
virginiana, Linn. l. c. = T. virginiana.

CRACCA, [Riv.] Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 359 = **Vicia**, Tourn. (Legum.).
acutifolia, Alef. in Bonplandia, ix. (1861) 118 = Vicia acutifolia.
agrigentina, Alef. l. c. 119 = V. Bivonea.
albicans, Alef. l. c. 120 = V. atropurpurea.
atropurpurea, Gren. & Godr. Fl. Fr. i. 471 = V. atropurpurea.
benghalensis, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 359 = V. benghalensis.
Bertolonii, Gren. & Godr. Fl. Fr. i. 470 = V. Pseudocracca.
Bivonea, Alef. in Bonplandia, ix. (1861) 121 = V. Bivonea.
Boissieri, Alef. l, c. = V. villosa.
Broteriana, Alef. l. c. 120 = V. atropurpurea.
calcarata, Gren. & Godr. Fl. Fr. i. 472 = V. calcarata.
caroliniana, Alef. in Bonplandia, ix. (1861) 124 = V. caroliniana.
cassubica, Opiz, Seznam, 34 = V. cassubica.
ciceroidea, Alef. in Bonplandia, ix. (1861) 121 = V. ciceroidea.
cirrhosa, Alef. l. c. 119 = V. cirrhosa.
corsica, Gren. & Godr. Fl. Fr. i. 473 (= *Ervum corsicum*).—Corsica.
dasycarpa, Alef. in Bonplandia, ix. (1861) 121 = V. varia.
disperma, Gren. & Godr. Fl. Fr. i. 472 = V. disperma.
dumetorum, Opiz, Seznam, 34 = V. dumetorum.
elegans, Alef. in Bonplandia, ix. (1861) 124 = V. tenuifolia.
erotanthos, Alef. l. c. 118 = V. Leavenworthii.
exigua, Alef. l. c. 119 = V. exigua.
Gerardi, Gren. & Godr. Fl. Fr. i. 469 = V. Gerardi.
Gerhardi, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 359 = V. Gerardi.
hypoleuca, Alef. in Bonplandia, ix. (1861) 123 = V. alpestris.
Kotschyana, Alef. l. c. 119 = V. Kotschyana.
linifolia, Alef. l. c. 118 = V. atropurpurea.
longifolia, Alef. l. c. 123 = V. longifolia.
ludoviciana, Alef. l. c. 119 = V. ludoviciana.
major, Alef. l. c. 125 = V. Cracca.
microphylla, Alef. l. c. 121 = V. microphylla.
minor, Gren. & Godr. Fl. Fr. i. 473 = V. hirsuta.
monanthos, Gren. & Godr. l. c. 471 = V. monanthos.
Monardi, Alef. in Bonplandia, ix. (1861) 119 = V. Monardi.
ochroleuca, Alef. l. c. 123 = V. ochroleuca.
pauciflora, Genneri, in Nuov. Giorn. Bot. ii. (1870) 136 = V. pauciflora?
pineta, Alef. in Bonplandia, ix. (1861) 123 = V. pinetorum.
pisiformis, Opiz, Seznam, 34 = V. pisiformis.
plumosa, Timb. in Bull. Soc. Sc. Phys. et Nat. Toulouse, iii. (1875-76) 143 = V. Cracca.
polyphylla, Alef. in Bonplandia, ix. (1861) 124 = V. tenuifolia?
Pseudocracca, Alef. l. c. 121 = V. Pseudocracca.
pulchella, Alef. l. c. 119 = V. pulchella.
Sibthorpii, Alef. l. c. 120 = V. Sibthorpii.
sylvatica, Opiz, Seznam, 34 = V. sylvatica.
syriaca, Medic. in Vorles. Churpf. Phys. Ges. ii. (1787) 359.—Syria.
tenuifolia, Opiz, Seznam, 34; Gren. & Godr. Fl. Fr. i. 469 = V. tenuifolia.
tenuissima, Alef. in Bonplandia, ix. (1861) 120 = V. gracilis.
Timbaliana, Debeaux.—Cf. Bull. Soc. Bot. Fr. xxviii. (1881) Bibl. 73 = V. cuneata.
tridentata, Alef. in Bonplandia, ix. (1861) 119, 125 = V. ludoviciana.
varia, Gren. & Godr. Fl. Fr. i. 469 = V. varia.
villosa, Gren. & Godr. l. c. 470 = V. villosa.
vulgaris, Opiz, Seznam, 34 = V. Cracca.

CRACCINA, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 266 = **Astragalus**, Tourn. (Legumin.).
austriaca, Stev. l. c. = A. austriacus.
bifidus, Stev. l. c. xxix. (1856) II. 145 = A. multicaulis.
Cracca, Stev. l. c. iv. (1832) 266 = A. Cracca.
depressa, Stev. l. c. = A. depressus.
ervoides, Stev. l. c. xxix. (1856) II. 145 = A. ervoides.
falcata, Stev. l. c. iv. (1832) 266 = A. falcatus.
gracilis, Stev. l. c. = A. gracilis.
leptocaulis, Stev. l. c. xxix. (1856) II. 145 = A. leptocaulis.

CRACCINA:—
leptostachys, Stev. in Bull. Soc. Nat. Mosc. iv. (1832) 266 = Astragalus sulcatus.
leucophaea, Stev. l. c. = A. leucophaeus.
longipes, Stev. l. c. xxix. (1856) II. 145 = A. macropterus.
multicaulis, Stev. l. c. = A. multicaulis.
puberulus, Stev. l. c. = A. puberulus.
racemosa, Stev. l. c. iv. (1832) 266 = A. racemosus.
reptans, Stev. l. c. = A. reptans.
scopaeformis, Stev. l. c. xxix. (1856) II. 145 = A. tauricus.
sulcata, Stev. l. c. iv. (1832) 266 = A. sulcatus.
taurica, Stev. l. c. = A. tauricus.
triangularis, Stev. l. c. = A. triangularis.
tunetana, Stev. l. c. = A. tunetanus.
vaginata, Stev. l. c. = A. vaginatus.
versicolor, Stev. l. c. = A. versicolor.
vicioides, Stev. l. c. xxix. (1856) II. 145 = A. macropterus.

CRAEPALIA, Schrank, Baier. Fl. i. 102, 382 (1789) = **Lolium**, Linn. (Gramin.).
temulenta, Schrank, l. c. 382 = L. temulentum.

CRAEPALOPRUMNON, Karst. Fl. Columb. i. 123 t. 61, 62 (1859?) = **Xylosma**, Forst. (Bixin.).
heterophyllum, Karst. l. c. 123. t. 61.—N. Granat.
cordatum, Karst. l. c. 124 = Flacourtia cordata.
obovatum, Karst. l. c. 123. t. 61.—N. Granat.
prunifolium, Karst. l. c. 124 = Flacourtia prunifolia.
racemosum, Karst. l. c. = Flacourtia racemosa.
rubicundum, Karst. l. c. 125. t. 62.—Venezuela.

CRAFORDIA, Rafin. Specch. i. 156 (1814) = **Tephrosia**, Pers.? (Legumin.).
bracteata, Rafin. l. c. = T. spicata?

CRAMBE, Tourn. ex Linn. Syst. ed. I (1735 *CRUCIFERAE*, Benth. & Hook. f. i. 98.
amplexicaulis, [Soland. in] Russell, Aleppo, ed. ii. 258 = Calepina Corvini.
aspera, Bieb. Fl. Taur. Cauc. ii. 90 = tatarica.
Aucheri, Boiss. in Ann. Sc. Nat. Sér. II. xvii. (1842) 3 = orientalis.
Biebersteinii, *Janka, in Termész. Füzetek*, viii. (18 36.—Europ.
bursaefolia, L'Hérit. ex DC. Syst. ii. 649 = Calep Corvini.
Buxbaumii, Willd. ex Ledeb. Fl. Ross. i. 222 orientalis.
caspica, Raeusch. Nom. ed. III. 183 = tatarica.
chlorocarpa, *Kit. in Linnaea*, xxxii. (1863) 502. Austria.
cordata, Willd. Enum. Hort. Berol. Suppl. 4 cordifolia.
cordifolia, Duf. in Ann. Gén. Sc. Phys. vii. (1820) 3 = glabrata.
cordifolia, *Stev. in Mém. Soc. Nat. Mosc.* iii. (18 267.—Reg. Caucas.
Corvini, All. Fl. Pedem. i. 256 = Calepina Corvini.
edentula, *Fisch. & Mey. ex Kar. in Bull. Soc. N Mosc.* (1839) 145, nomen.—Turcoman.
filiformis, Boiss. Voy. Bot. Espagne, 43 = hispanica.
filiformis, *Jacq. Coll. Suppl.* 120.—Patagon.; Chili
fruticosa, *Linn. f. Suppl.* 299.—Ins. Madera.
gibberosa, *Rupr. Fl. Cauc.* 136, in textu.—R Caucas.
gigantea, Kit. ex Janka, in Oestr. Bot. Zeitschr. (1871) 65 = maritima.
glabrata, *DC. Prod.* i. 226.—Hispan.
grandiflora, *DC. Syst.* ii. 652.—As. Min.
hispanica, *Linn. Sp. Pl.* 671.—Reg. Mediterr.
juncea, *Bieb. Fl. Taur. Cauc.* iii. 421.—Reg. Cauc Himal.
Kotschyana, Boiss. Diagn. Ser. I. vi. 19 = cordifolia.
Kralikii, *Coss. in Bull. Soc. Bot. Fr.* iv. (1857) 4, nomen.—Afr. bor.?
laciniata, Desf. ex Steud. Nom. ed. II. i. 429 = pinnatifida.
laciniata, Lam. Encyc. ii. 163 = tatarica.
laevis, Kit. in Linnaea, xxxii. (1863) 502 = chlorocarpa.
macrocarpa, Bieb. Fl. Taur. Cauc. ii. 90 = Sobolewskia lithophila.
maritima, *Linn. Sp. Pl.* 671.—Europ.
orientalis, Jacq. Ic. Rar. i. t. 128 = pinnatifida.
orientalis, *Linn. Sp. Pl.* 671.—As. Min.; Persia.
orientalis, Reich. Syst. iii. 290 = tatarica.

CRAMBE:—
palmatifida, *Regel & Schmalh. in Act. Hort. Petrop.* v. (1877) 243.—Turkestan.
paniculata, All. Fl. Pedem. i. 256 = Neslia paniculata.
pannonica, Hort. Par. ex Lam. Encyc. ii. 163 = tatarica.
persica, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 389.—Persia.
pinnatifida, [*R. Br. in*] *Ait. Hort. Kew.* ed. II. iv. 72.—Reg. Caucas.
pontica, *Stev. ex Rupr. Fl. Cauc.* 293.—Tauria.
Pritzelii, *C. Bolle, Ind. Sem. Hort. Berol.* (1861) 10.—Ins. Canar.
quadricostata, *Boiss. in Ann. Sc. Nat.* Sér. II. xvii. (1842) 388 = orientalis.
reniformis, Desf. Fl. Atlant. ii. 78. t. 151 = hispanica.
scabra, Lam. Encyc. ii. 163 = strigosa.
Sewerzowi, *Regel, in Bull. Soc. Nat. Mosc.* xliii. (1870) I. 282.—Turkestan.
sinuato-dentata, *Hochst. & Schimp. Beitr. Fl. Aethiop.* 238.—Abyss.
Steveniana, *Rupr. Fl. Cauc.* 136, in textu.—Reg. Caucas.
strigosa, *L'Hérit. Stirp. Nov.* 151. t. 72.—Ins. Canar.
suecia, Mill. Gard. Dict. ed. VIII. n. 2 = pinnatifida.
tatarica, *Jacq. Misc.* ii. 274.—Europ. or.; As. bor.

CRAMERIA, Murr. Syst. ed. XIII. 138 (1774) = **Krameria**, Linn. (Polygal.).

CRANICHIS, Sw. Prod. Veg. Ind. Occ. 120 (1788). *ORCHIDEAE*, Benth. & Hook. f. iii. 593.
OCAMPOA, A. Rich. & Gal. in Ann. Sc. Nat. Sér. III. iii. (1845) 31.
aphylla, *Sw. l. c.*—Jamaica.
apiculata, *Lindl. in Benth. Pl. Hartw.* 92.—Guatemala.
ciliata, *Kunth, Syn. Pl. Aeq.* i. 324.—Venezuela.
diphylla, *Sw. Prod. Veg. Ind. Occ.* 120.—Jamaica.
Engelii, *Reichb. f. in Linnaea*, xli. (1877) 19.—Venezuela.
foliosa, *Lindl. Gen. et Sp. Orch.* 451.—Peruv.
gibbosa, *Lindl. in Ann. Nat. Hist.* xv. (1845) 385.—Jamaica.
glandulosa, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 30.—Mexic
Lehmanni, *Reichb. f. Otia Bot. Hamb.* 4.—Ecuador.
luteola, Sw. Fl. Ind. Occ. iii. 1433 = Polystachya luteola.
macroblepharis, *Reichb. f. Otia Bot. Hamb.* 4.—Ecuador.
micrantha, *Griseb. in Gott. Abh.* xxiv. (1879) 337.—Reg. Argent.
micrantha, Spreng. Syst iii. 700 = Prescottia Myosurus.
monophylla, *Lindl. Orchid. Linden*, 27.—N. Granat.
multiflora, Ell. ex Nutt Gen. Am. ii. 191 = Ponthieva glandulosa.
muscosa, *Sw. Prod. Veg. Ind. Occ.* 120.—Jamaic.
nudifolia, *Pers. Syn.* ii. 511.—China.
oligantha, Sw. Prod. Veg. Ind. Occ. 120 = Prescottia Myosurus.
ovata, *Wickstr. in Vet. Akad. Handl. Stockh.* (1827) 73.—Ins. Guadalup.
parvilabris, *Lindl. Orchid. Linden.* 27.—N. Granat.
pauciflora, *Sw. Fl. Ind. Occ.* iii. 1431.—Jamaic.
picta, *Reichb. f. in Linnaea*, xli. (1877) 52.—Ecuador.
reticulata, *Reichb. f. Beitr. Orch. Centr. Am.* 62.—Costa Rica.
Schaffneri, *Reichb. f. in Bonplandia*, iii. (1855) 238.—Mexic.
Schlimii, *Reichb. f. in Linnaea*, xli. (1877) 19.—Venezuela.
similis, *Reichb. f. Otia Bot. Hamb.* 83.—Bras.
speciosa, *La Llave & Lex. Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 5.—Mexic.
stachyoides, Sw. Prod Veg. Ind. Occ. 120 = Ponthieva glandulosa.
subumbellata, *A. Rich. & Gal. in Ann. Sc. Nat.* Sér. III. iii. (1845) 30.—Mexic.
sylvatica, *A. Rich. & Gal. l. c.*—Mexic.
tenuiflora, *Griseb. Cat. Pl. Cub.* 268.—Cuba.
tenuis, *Reichb. f. in Flora*, xlviii. (1865) 274.—Ind. occ.
tubularis, *La Llave & Lex Nov. Veg. Desc. fasc.* ii. (*Orch. Opusc.*) 6.—Mexic.
tubulosa, *Lindl. Gen. et Sp Orch.* 451.—Mexic.
ventricosa, *Griseb. Fl. Brit. W. Ind.* 639.—Ind. occ.
Wageneri, *Reichb. f. in Linnaea*, xli. (1877) 19.—Venezuela.