

Epiperipatus puri sp. nov., a new velvet worm from Atlantic Forest in Southeastern Brazil (Onychophora, Peripatidae)

Cristiano Sampaio Costa¹, Amanda Cruz Mendes² and Alessandro Ponce de Leão Giupponi³

¹ Departament of Biology and Zoology, Universidade Federal de Mato Grosso—UFMT, Cuiabá, Mato Grosso, Brazil

² Departament of Zoology, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

³ Fundação Oswaldo Cruz—FIOCRUZ, Instituto Oswaldo Cruz—IOC, Collection CAVAIISC, LAC, Rio de Janeiro, Brazil

ABSTRACT

Epiperipatus ohausi (Bouvier, 1900) is the first species known from Rio de Janeiro, and more than 120 years later a new species is described in the state of Rio de Janeiro (RJ). In this study, we describe the second species in the state of Rio de Janeiro, which we are naming in honor of the indigenous population called *puri* who resided in southeastern coastal Brazil. The species can be diagnosed mainly by large dorsal primary papillae close to the insertion of the legs drawing a light band from the anterior to the posterior region of the body, and large dorsal primary papillae alternating on the dorsal plicae. Moreover, they are recognized *in vivo* by the color of the diamond-shaped marks brownish orange on the dorsal portion of the body.

Epiperipatus puri sp. nov. morphologically seems to be related to *Epiperipatus acacioi* (Marcus & Marcus, 1995) by the shape of the primary papillae apical piece and to *E. ohausi* by the resemblance of dorsal papillae. The phylogeny shows a close relationship between the new species and *E. ohausi* in a clade with a still undescribed species from Rio de Janeiro, Brazil located within the Atlantic Forest, one of the most threatened biomes in the world.

Submitted 10 March 2022

Accepted 18 April 2023

Published 2 October 2023

Corresponding author

Alessandro Ponce de Leão Giupponi,
agiupponi@gmail.com

Academic editor

Joseph Gillespie

Additional Information and
Declarations can be found on
page 12

DOI 10.7717/peerj.15384

© Copyright
2023 Costa et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Biodiversity, Molecular Biology, Taxonomy, Zoology

Keywords New species, Neotropical, Neopatida, Taxonomy, Biodiversity, Threatened species

INTRODUCTION

Currently, the *circa* 200 recent species of Onychophora are distributed in two families, Peripatidae Evans, 1901 and Peripatopsidae Bouvier, 1905. The phylum Onychophora (velvet worms) has received more attention in the last decade than ever, especially the clade Neopatida (=the Neotropical Peripatidae), which resulted in the recent description of five species ([Baker et al., 2021](#)).

Although velvet worms are among the most fascinating terrestrial groups of invertebrates, their biodiversity is poorly understood, and the taxonomy is elusive ([Sedgwick, 1888](#); [Read, 1988](#); [Giribet et al., 2018](#)). Some causes are the poor sampling of individuals in the field, low numbers of specimens available in museums often accompanied by inadequate preservation, poor access to historical collections, scarce

type-locality data, and the conservative morphology of the group (*Froehlich, 1968; Peck, 1975; Read, 1988; Chagas-Jr & Costa, 2014; le Bras et al., 2015; Costa & Giribet, 2016; Costa, Chagas-Jr & Pinto-da-Rocha, 2018; Giribet et al., 2018*). The lack of distinguishable external characters capable of clearly delimiting the boundaries of the taxa has led to the search for new methodologies such as scanning electron microscopy (e.g., *Read, 1988; Morera-Brenes & Monge-Nájera, 1990; de Oliveira, Wieloch & Mayer, 2010; Costa & Giribet, 2016*) and DNA sequences analyses which have led to the description of many species for the phylum in the last decade (e.g., *de Oliveira et al., 2013; Giribet et al., 2018*).

Epiperipatus Clark, 1913 is the most diverse of the 11 recent genera within Peripatidae, with 36 valid species distributed in the Antilles, Central, and South America, with 15 species from Brazil (*Peck, 1975; Brito et al., 2010; de Oliveira et al., 2011; de Oliveira, Read & Mayer, 2012; Costa, Chagas-Jr & Pinto-da-Rocha, 2018; Costa, Giribet & Pinto-da-Rocha, 2021*).

Epiperipatus was erected by *Clark (1913)* as a subgenus of *Peripatus* Guilding, 1826 along with other two new subgenera (*Plicatoperipatus* Clark, 1913 and *Macroperipatus* Clark, 1913), designating *Peripatus edwardsii* Blanchard, 1847 as the type species and transferring more nine species to this subgenus (Clark, 1913: 18). Since *Peck (1975)*, *Epiperipatus* is treated as genus, although this author did not explicitly elevate the rank from the subgenera of *Peripatus*. Peck cited it under an identification key for families and genera of Onychophora and made the combinations with the species without using *Peripatus* in the binomina (*Peck, 1975*: 345). After Peck's work *Epiperipatus* counted with 17 species. Subsequent works described new species as *Epiperipatus* (*Brito et al., 2010; de Oliveira et al., 2011; Costa, Chagas-Jr & Pinto-da-Rocha, 2018*) or reassigned extant ones to the genus (*de Oliveira, Wieloch & Mayer, 2010; Chagas-Jr & Costa, 2014; Costa, Chagas-Jr & Pinto-da-Rocha, 2018; Costa, Giribet & Pinto-da-Rocha, 2021; Morera-Brenes & Monge-Nájera, 2010*).

A recent revision of *Epiperipatus* intended to improve the delimitation of the genus based on morphological and molecular data, aiming to test the information of 33 morphological characters applied in the taxonomy of Peripatidae (*Costa, Giribet & Pinto-da-Rocha, 2021*). In this article, although *Epiperipatus* appears as non-monophyletic, a core monophyletic group emerges including the type species *Epiperipatus edwardsii*, which was recently redescribed under modern parameters (*Costa, Chagas-Jr & Pinto-da-Rocha, 2018*).

An undescribed species occurs in Cachoeiras de Macacu, Rio de Janeiro, Brazil, and according to *Costa, Giribet & Pinto-da-Rocha (2021)* belongs to *Epiperipatus*. The area where the types were collected is a fragment of the Atlantic Forest (AF, hereafter), an extremely threatened biome (*Ribeiro et al., 2011*). The AF is one of the 36 conservation hotspots of the world (*Myers et al., 2000; CEPF, 2019*), and one of the most important given its conservation value in the area (*Laurance, 2009*).

In the present work, this species is newly described as *Epiperipatus puri* sp. nov. Its type specimens are deposited at Museu Nacional, Universidade Federal do Rio de Janeiro, (MNRJ), as part of the rebuilding of the burned MNRJ zoological collections. *Epiperipatus puri* sp. nov. is the first species of onychophorans deposited in MNRJ after its main

building was lost in a ruthless fire. The collection of Onychophora was held together with the collections of Arachnida and Myriapoda, under the curatorship of Adriano B. Kury. Most of the three collections were lost in the fire except for material under loan and all the data, which were safe due to the routine backup policy of the curator ([Kury, Giupponi & Mendes, 2018](#)).

MATERIALS AND METHODS

The type-series is composed of six specimens deposited in Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro (MNRJ) (curator A. B. Kury) and Museu de Zoologia da Universidade de São Paulo, São Paulo (MZUSP) (curator R. Pinto-da-Rocha), both in Brazil. The specimens were collected between 2012 and 2018 in a small patch of tropical humid forest ([IBGE, 2019](#)) (Fig. 1) around the Metropolitan Region of the state of Rio de Janeiro, under grass roots (Fig. 2F), in a private conservation area (Reserva Ecológica de Guapiaçu). License code number: INEA (Brasil) n° 005/2020 (Alessandro Ponce de Leão Giupponi). Specimens were preserved in 70% and 100% EtOH. We examined their morphology in detail and compared it with specimens of *Epiperipatus ohausi* (Bouvier, 1900) from Nova Iguaçu municipality, Rio de Janeiro, Brazil to detect diagnostic features of the new species: MNRJ 0056; 1♀; BRAZIL, Rio de Janeiro, Nova Iguaçu, Reserva Particular do Patrimônio Natural dos Petroleiros; 23.XII.2009; Costa, C.S., Giupponi, A.P.L. leg. MNRJ 0058; 1♂; same locality; 11.III.2010; Costa, C.S., Chagas-Jr, A., Giupponi, A.P.L., Kury, A.B. leg.

We studied one of the specimens (MNRJ 0093, voucher 065) using scanning electron microscopy (SEM) following [Chagas-Jr & Costa \(2014\)](#) and [Costa, Chagas-Jr & Pinto-da-Rocha \(2018\)](#). We dissected out one mandible, the fifth oncopod of the left side, and a small rectangular section of the dorsal integument from the dorsomedian furrow to the base of the oncopods. The structures were critical point dried and mounted in SEM stubs with bi-adhesive carbon tape. A 5-nm gold layer was applied. Samples were imaged with a JEOL JSM-6390LV at the SEM Platform Rudolf Barth at Instituto Oswaldo Cruz—Fundação Oswaldo Cruz (IOC-FIOCRUZ).

Photographs *in vivo* were taken with a SONY Cybershot DSC-HX1 with built-in flash, or Canon EOS Rebel XS with a macro lens and flash circular camera. Images were edited using Adobe Photoshop CS5 (Figs. 2A–2E). For color descriptions, we followed the standard names of the 267 Color Centroids of the NBS/ISCC Color System (see [Kelly, 1958](#); also available at [Centore, 2016](#), and [W3Schools, 2023](#)). Also, we combined the stereomicroscopy and Scanning Electron Microscopy (SEM) studies of the external morphology of the specimens for descriptions. The morphological descriptive nomenclature follows the terminologies of [Read \(1988\)](#), [Morera-Brenes & Monge-Nájera \(2010\)](#), [de Oliveira, Wieloch & Mayer \(2010\)](#), and [Costa, Chagas-Jr & Pinto-da-Rocha \(2018\)](#). All measurements are given in millimeters (mm). The species we described here was part of a more detailed study of *Epiperipatus*, combining morphological and molecular data for specimens from the Neotropics ([Costa, Giribet & Pinto-da-Rocha, 2021](#)).

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature




Figure 1 Updated distribution of onychophorans in Brazil assembling the results of Costa, Chagas-Jr & Baptista (2009), Costa, Chagas-Jr & Pinto-da-Rocha (2018), and Costa, Giribet & Pinto-da-Rocha (2021). The records of specimens of onychophorans in Guapiaçu represented by red stars in all images of this plate, state of Rio de Janeiro. We do not include *Epiperipatus tucupi* (Froehlich, 1968) in the maps due to the imprecise location information provided on the label, which mentions only "Pará." The Reserva Ecológica de Guapiaçu (type-locality of *Epiperipatus puri* sp. nov.) is represented by a yellow star (image from Google Earth).

Full-size DOI: 10.7717/peerj.15384/fig-1

(ICZN). This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved, and the associated information viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSID for this publication is urn:lsid:zoobank.org:pub:A8971D25-1C9D-4807-BE2B-730FB010B717. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central and CLOCKSS.

RESULTS

Taxonomic results

Family **Peripatidae** Evans, 1901

Genus ***Epiperipatus*** Clark, 1915

Epiperipatus puri sp. nov.

Epiperipatus [sp6]: Costa: 2016; Costa, Giribet and Pinto-da-Rocha: 2021: 6, 8, 25. (Figs. 1–4)

Diagnosis (based on a combination of characters as follows). *Epiperipatus* species with large dorsal primary papillae close to the insertion of the legs drawing a narrow light band from the anterior to posterior regions of the body. The distribution of large pale dorsal




Figure 2 Photos of female paratype. *Epiperipatus puri* sp. nov., body background and papillae of female paratype MZUSP 0122, and the environment where the specimens were found. Body background in (A and B) dorsal side, (C and D) lateral side and (E) ventral side. (F) Site where the specimens were collected, under the roots of the grass, at Reserva Ecológica de Guapiaçu (REGUA), Cachoeiras de Macacu, Rio de Janeiro, Brazil. Scales bars 2, 4 and 5 = 1 mm; 3, 6 = 2 mm. [Full-size](#) DOI: 10.7717/peerj.15384/fig-2

primary papillae alternates the dorsal plicae with clear differences in the number in two sequential plicae. Moreover, they are recognized *in vivo* by the color the diamond-shape marks brownish-orange (Figs. 2A–2D).

Description of female holotype (MNRJ 0093).

Measurements. Length 43; width 3.0, height 3.9.

Color (for living specimen Figs. 2A–2E). Background color of body dark reddish-brown, overlaid by blurry diamond-shaped marks brownish orange. A broad dashed-line (close to the insertion of the legs) similarly colored as the diamond-shape marks. Dorsomedian




Figure 3 SEM of holotype. *Epiperipatus puri* sp. nov., female holotype MNRJ 0093, scanning electron micrographs. (A) Dorsal plicae. The dorsomedian furrow position indicated by the arrow head, Additionally, two primary papillae are marked in purple, and three accessory papillae are highlighted in yellow. (B) shape of primary papilla, in lateral view, with the posterior view depicted on the left side of the image. (C) Foot of the fifth oncopodid in prolateral view, (D and E) on the top in inner and outer jaws blades respectively. Scales bars 3A = 500 µm; 3B = 50 µm; 3C-E = 100 µm.

[Full-size](#) DOI: 10.7717/peerj.15384/fig-3

furrow dark reddish-brown. Anterior portion of head moderate reddish-brown and antennae strong brown. Color of the dorsal portion of the legs grayish reddish orange. Legs and ventral surface display the same color, moderate reddish-brown. Ventral and preventral organs moderate brown.

Description of the body (Figs. 2A–2E and 3). Conspicuous dorsomedial furrow and hyaline organs along the main body axis (Figs. 2B and 3A). Twelve plicae per segment, two




Figure 4 Total evidence analysis adapted from [Costa, Giribet & Pinto-da-Rocha \(2021\)](#) of *Epiperipatus puri* sp. nov. and related species. Our species, represented by three vouchers, is in a clade in shades of blue with a so far unnamed species (*Epiperipatus* sp5: MNRJ 0059) and *E. ohausi* (MNRJ 0056, 0058), all from Rio de Janeiro (RJ), Brazil. The RJ clade is sister-group to a clade in orange composed by *E. machadoi* (MNRJ 0043) from Minas Gerais, Brazil and three unnamed species from Espírito Santo, Brazil (*Epiperipatus* sp13: MNRJ 0018, 0020, 0023, 0026; *Epiperipatus* sp.: MNRJ 0016, 0100; *Epiperipatus* sp11: MNRJ 0042, 0046).

[Full-size](#) DOI: [10.7717/peerj.15384/fig-4](https://doi.org/10.7717/peerj.15384/fig-4)

incompletes as broad as the diamond-shape marks, and seven crossing to the ventral side (Fig. 3A).

Almost all dorsal papillae on the plicae, except for the smaller accessory papillae on the furrow between the plicae. Primary dorsal papillae aligned on top of folds; two primary papillae separated by one to five accessory papillae, the former do not occur close together (Figs. 2D and 3A). Dorsal primary papillae with a conical basal piece composed of scales that never overlap each other at the base of the papillae (Figs. 3A and 3B). Primary papillae as the largest dorsal papillae, with roundish dome insertion and asymmetrical regular spherical apical piece (Fig. 3B). Basal piece larger than apical, with a range of at least seven scale ranks (Fig. 3B). Apical piece with three posterior scale ranks (Fig. 3B).

Narrow scales both on base and apical piece (Fig. 3B). Needle-shaped sensory bristle directed posteriorly (Fig. 3B). Small and large primary papillae with a conspicuous constriction between the base and the apical piece (Figs. 3A and 3B). Two-sized dorsal primary papillae: the largest are on the top of the body in continuous plicae (close to the dorsomedian furrow and drawing the diamond-shaped marks) and close to the legs. Lateral papillae in alternated dorsal plicae. Accessory papillae are the smallest dorsal, with roundish insertion similar in shape to the base of the primary papillae; they are more abundant per plicae and differ in position in relation to primary papillae (Figs. 2D and 3A).

Head. No evident structures or patterns on the head. Antennae (Figs. 2A–2C) composed of 40 rings. Antennal tip composed of seven broad rings, excluding the disc on the top, followed by a sequence of narrow and broad rings alternating until at least ring 20. Eyes and frontal organs present ventrolateral to the antennal base. Conspicuous frontal organs as long as six fused antennal papillae. Mouth opening surrounded by a small, anterior, unique lobe, and seven flanked lobes decreasing in size from the anterior to the posterior end of the mouth. Dental formula of inner and outer jaws (Figs. 3D and 3E), respectively: 1/1 and 1/2/10. The accessory tooth is thinner in the outer jaw. The second accessory tooth is reduced.

Legs. 28 pairs of legs in the holotype (Fig. 2E). Nephridial tubercle on fourth and fifth pairs of legs, between third and fourth spinous pads (Fig. 2E), connected by the top to the third spinous pad. On the fourth and the fifth pairs of legs, four spinous complete pads are present and no evidence of a fifth one.

Sexual dimorphism. Two pairs of pregenital legs with one crural papilla (male) each, absent in females. Anal glands are inconspicuous and represented only by two pores on the anterior margin of the anal aperture, absent in females.

Type material. Holotype: MNRJ 0093, 1♀, BRAZIL, Rio de Janeiro, Cachoeiras de Macacu, Reserva Ecológica de Guapiaçu (REGUA), X.2012, A.P.L. Giupponi, J.S. Silva leg. Paratypes: MNRJ 0087, 1 unsexed specimen, MNRJ 0088, 1♂, same locality, 28.II – 02. III.2012, A.P.L. Giupponi, J.S. Silva leg; MNRJ 0107, 1 unsexed specimen, same locality, 19. III.2018, R.L.C. Baptista leg; MZUSP 0122, 1♀, same locality, 21.XII.2014, A. Ferreira, A.P.L. Giupponi, A. Rezende, C.S. Costa leg.

Distribution. Only known from the type locality (Fig. 1).

Etymology. The epithet *puri* (in apposition) refers to the Puri indigenous group belonging to the *Macro-Jê* linguistic group. They inhabited, among other places, the mountain region of the Rio de Janeiro state where specimens of this species were collected. Noun in apposition.

Remarks. Paratype. Length 12 to 22; width 1.0 to 2.5. Legs. 26 (female) and 27 (male) pairs of legs.

DISCUSSION

Classification of *Epiperipatus puri* sp. nov.

The fuzzy generic limits due to the lack of clear morphological characters is a major issue of Neopatida taxonomy. Splitting species of Peripatidae in several genera might be a problem, especially in the systematics of Brazilian fauna (Giribet et al., 2018: 860; Costa, 2016). *Epiperipatus* is one of the most speciose genera of Peripatidae (de Oliveira, Read & Mayer, 2012; Costa, Chagas-Jr & Pinto-da-Rocha, 2018) with 36 described species after this article.

However recent studies based on the molecular data regards the genus as non-monophyletic (Costa, 2016; Giribet et al., 2018). Phylogenetically, Costa, Giribet & Pinto-da-Rocha (2021) based on the study of four molecular markers besides morphological data distinguished two clades of Brazilian species, the smallest with species from the state of Pará and the largest with the remaining species of the country included in the study. *Epiperipatus puri* sp. nov. appears nested in the largest clade closely related with species from the state of Rio de Janeiro: *Epiperipatus ohausi* (Bouvier, 1900) and a potentially undescribed species (see in Costa, Giribet & Pinto-da-Rocha (2021): 23, fig. 3, clade S, and Fig. 4 here). This clade is sister-group to another composed by *Epiperipatus machadoi* (Oliveira & Wieloch, 2005) and three other unnamed species from the state of Espírito Santo, Brazil, which is geographically close to the state of Rio the Janeiro and located within the Atlantic Forest.

Epiperipatus puri sp. nov. is characterized as a new species by the roundish insertion of dorsal papillae, the three posterior scale ranks, and two prolateral and one retrolateral foot

papillae in the feet of the fourth and fifth oncopods (Fig. 3C). The presence of incomplete folds differs *E. puri* sp. nov. from *Epiperipatus brasiliensis* (Bouvier, 1899), *Epiperipatus tucupi* Froehlich, 1968 and *Epiperipatus cratensis* Brito et al., 2010. The new species differs from *Epiperipatus diadenoprotus* de Oliveira et al., 2011 by the inconspicuous anal glands in *E. puri* sp. nov.

In *Epiperipatus paurognostus* de Oliveira et al., 2011 the background color of the body is reddish-brown (*in vivo*) and the fourth spinous pad can be complete or incomplete, in *E. puri* sp. nov. the background color of the body is dark reddish-brown (*in vivo*) and the fourth spinous pad is complete. The apical piece is conical in *Epiperipatus adenocryptus* (de Oliveira et al., 2011) and *E. machadoi*, conical/cylindrical in *Epiperipatus lucerna* Costa, Chagas-Jr & Pinto-da-Rocha, 2018 and *Epiperipatus marajoara* (Costa, Chagas-Jr & Pinto-da-Rocha, 2018), and spherical in *E. puri* sp. nov. The apical piece is conical and reduced in *Epiperipatus beckeri* (Costa, Chagas-Jr & Pinto-da-Rocha, 2018) and *Epiperipatus titanicus* (Costa, Chagas-Jr & Pinto-da-Rocha, 2018), robust in *Epiperipatus hyperbolicus* (Costa, Chagas-Jr & Pinto-da-Rocha, 2018), while the apical piece is regular in *E. puri* sp. nov. The new species seems to be closely related to *Epiperipatus acacioi* (Marcus & Marcus, 1955) by the shape of the apical piece of the primary papillae, however, *E. puri* sp. nov. primary papillae are lighter than other papillae. *E. ohausi* and *E. puri* sp. nov. bear dorsal papillae with similar shape and size, but the latter also bear accessory papillae on the flanks and uniform background color of body and oncopods.

Additionally, the results of Costa, Giribet & Pinto-da-Rocha (2021) include the species in a clade containing the type species of the genus, although the clade also includes species of other genera. Giribet et al. (2018) suggested that Caribbean species of *Epiperipatus* are closer to *Peripatus* than to the remaining “*Epiperipatus*”, however, this could be confirmed only by the inclusion of *Peripatus juliformis* Guilding, 1826 (type species of *Peripatus*) in Peripatidae analyses, neither of both studies included this species due to the lack of specimens in the collections adequate for the analyses.

Although the boundaries between *Epiperipatus* and *Peripatus* remain unclear, we cautiously preferred keeping the new species as *Epiperipatus*, as their putative closer species *E. ohausi* is currently classified. According to Bouvier (1905), the new species could not be a *Peripatus* due to the presence of only two pairs of pregenital oncopods with crural papillae, in *Peripatus* should be more than three. Also, according to Read (1988) it could not be classified as *Peripatus* because it has three ranks of scales in the apical piece, *Peripatus* should have more ranks.

CONSERVATION

Epiperipatus puri sp. nov. is the 16th *Epiperipatus* species described from Brazil (Chagas-Jr & de Oliveira, 2019). Besides *E. puri* sp. nov., only *E. ohausi* is nominally recorded from Rio de Janeiro (See Costa, Chagas-Jr & Baptista, 2009; Costa, Chagas-Jr & Pinto-da-Rocha, 2018), a Brazilian state entirely within the Atlantic Forest domain.

The Atlantic Forest is the second largest rainforest in South America and one of the most distinctive biogeographic units in the Neotropical Region with high levels of endemism and biodiversity (Ribeiro et al., 2011). However, this biome has experienced

large habitat losses since European colonization and currently, only 12.59% of its original area remains (*Ribeiro et al., 2009, 2011*) making it one of the “hottest hotspots” for conservation (*Mittermeier et al., 2004; Laurance, 2009*).

Our species was collected in a private conservation area, the RPPN Reserva Ecológica de Guapiaçu (REGUA) (Fig. 1). The reserve is in the upper Guapiaçu River Valley and started as a group of farms registered as a non-governmental organization in the early 2000s. Today part of its area (in a total of 357 ha of two areas) is officially recognized as one of the private conservation areas of the state of Rio de Janeiro, part of the state program of Private Reserves of Natural Patrimony (Portuguese acronym: RPPN) (see *Guagliardi, 2018*). The total area of the reserve (official and nonofficial) encompasses 7,500 ha of forest in different stages of forest cover, with an altitudinal range from 0 to above 2,000 m.a.s.l. (*Soares et al., 2011*). The region is in a mountain range where the Guapi-Macacu Basin belongs, which contributes to the water supply of 2.5 million inhabitants of five municipalities (*Rodríguez Osuna et al., 2014*). In this watershed, the degradation of aquatic resources is caused by urbanization, intense agriculture, and conversion of riparian vegetation (*Rodríguez Osuna et al., 2014*). The forest cover of the landscape in the Guapi-Macacu Basin is *circa* 40%, and it is a mosaic of different ages; unfortunately fragments anterior to 1976 occupy only 12% of the landscape (*Costa et al., 2017*).

Although composed mainly of secondary forests, REGUA is important to the conservation of local fauna. The area is known for its rich fauna of birds (*Pimentel & Olmos, 2011*), butterflies (*Soares et al., 2011*), mosquitoes (*Silva et al., 2014*), and for dragonflies and damselflies, with more than 200 species (*Kompier, 2015; Pinto & Kompier, 2018*). *E. puri sp. nov.* is the first Onychophora to REGUA. This demonstrates the high value of this reserve for the recovery of the endangered area it is embedded.

For more than 30 years, scientists advocated for the importance of invertebrates and their conservation, but the perspective has not changed much (*Wilson, 1987; Collen et al., 2012*). The velvet worms are at risk given their distribution in threatened biomes, such as the Atlantic Forest itself, and because they seem to occur in small-sized populations, although the amount of data available on population dynamics is scarce (*New, 1995*). Sometimes the newly described species are already critically endangered (e.g., *de Oliveira et al., 2015*).

Currently 80% of Brazilian species of onychophorans are in the Livro Vermelho da Fauna Brasileira Ameaçada de Extinção (*ICMBio, 2018*). One of the species considered endangered in Brazil’s Red Book is *E. ohausi* (see *Costa, Cordeiro & Chagas-Jr, 2018*), the only named species from Rio de Janeiro state before this work, considered here to be a close species to *E. puri sp. nov.*

Epiperipatus ohausi is known from Petrópolis (type-locality) and Nova Iguaçu (*Chagas-Jr & Costa, 2014*), forests from both localities suffer from pressures of urbanization. The population of the species is severely fragmented since it occupies humid shaded habitats, with an extent amount of litter, typical of forested areas (*Costa, Cordeiro & Chagas-Jr, 2018*). Although *E. puri sp. nov.* is distributed in a close area, also with high pressures of urbanization and agriculture, and probably its distribution extends to the area

of Parque Estadual dos Três Picos, a State Reserve contiguous to REGUA. This reinforces the need for the preservation of those reserves and encourages the expansion of their areas.

After a few years of political stability and economic growth (1995-2014), with policies prioritizing the fight against poverty, environmental destruction, and historical deficit in science and education, Brazil is passing through severe economic, political, and social turmoil ([Dobrovolski et al., 2018](#)). Recently the indexes of deforestation are skyrocketing in all Brazilian biomes ([Instituto Nacional de Pesquisas Espaciais \(INPE\), 2023](#)), which seems to be related to recent policies favoring livestock ranching and agribusiness, and the weakening of the Brazilian system of protection of the environment and Indigenous lands ([Ferrante, Gomes & Fearnside, 2020](#)).

Although fires in Amazonian Forest, Cerrado, and Pantanal are usually related to the replacement of natural vegetation by cattle ranching and soy crops, in the AF the deforestation is related to urbanization (see [Joly, Metzger & Tabarelli, 2014](#)) and pressure of the real estate market. The increment in deforestation occurs in association with the negligence towards the scientific institutions (the destruction of Museu Nacional and its collections being an emblematic symbol), which suffered significant budget cuts ([Martelli-Jr et al., 2019](#); [Escobar, 2019](#)), staff shortage, and direct federal political intervention in their management, which undermines the protection of fragile biota.

CONCLUSIONS

Our description of *Epiperipatus puri sp. nov.* contributes to the knowledge of the biodiversity in a hotspot for conservation, the Atlantic Forest. We characterized the species morphologically with the use of SEMs and photographs, including *in vivo* (important for recognizing the species in the field). All the type material was collected in a private reserve that is contiguous to a State Protected Area, demonstrating the importance of this type of initiative. *E. puri sp. nov.* was assigned to *Epiperipatus* such as the putative closer species, *E. ohausi*, but future studies could reveal the actual boundaries of the genera for there is molecular evidence that they could belong to *Peripatus* ([Giribet et al., 2018](#)). One of the known obstacles to the conservation of invertebrates is the poor state of knowledge of the species, many still unnamed. In the case of velvet worms, the difficulty to describe a species is notorious, and one recent proposed solution to deal with this problem is to connect information about undescribed species to common names ([Sosa-Bartuano, Monge-Nájera & Morera-Brenes, 2018](#)).

ACKNOWLEDGEMENTS

We are grateful to Plataforma de Microscopia Eletrônica Rudolf Barth (FIOCRUZ-IOC) and to their staff, Wendell Girard Dias, Roger Magno Macedo Silva, Rômulo Custódio dos Santos, and Taíssa Ribeiro Adriano de Oliveira, for the help with the SEM images. We are grateful to Adriano B. Kury (MN RJ) who provided support with the Latin grammatical forms. We are in debt to Júlia dos Santos Silva (FIOCRUZ) Nicholas Locke, and Raquel Locke who assisted with SEM images. Jorge Bizarro for valuable information about REGUA. We also thank Flávio Uemori Yamamoto for providing beautiful photographs of

the new species. We also thank Alicia Santos and Jackie Thai (PeerJ Editorial Board) and two anonymous reviewers for their valuable contributions that enhanced the article.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work and fieldwork for Cristiano Sampaio Costa was funded by the São Paulo Research Foundation (FAPESP, who was supported by fellowships from FAPESP #2011/20211-0, 2012/02969-6 and 2014/20557-2). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

São Paulo Research Foundation: FAPESP #2011/20211-0, 2012/02969-6 and 2014/20557-2.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Cristiano Sampaio Costa conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Amanda Cruz Mendes analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Alessandro Ponce de Leão Giupponi analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.

Field Study Permissions

The following information was supplied relating to field study approvals (*i.e.*, approving body and any reference numbers):

INEA (Brazil) #005/2020 to Alessandro Ponce de Leão Giupponi.

Data Availability

The following information was supplied regarding data availability:

The sequences are available at NCBI:

- Epiperipatus [sp5] MNRJ 0059 ony-012: [MN905635-MN544110](#).
- Epiperipatus [sp11] MNRJ 0046 ony-003: [MN905628-MN544104](#).
- Epiperipatus [sp11] MNRJ 0042 ony-027a: [MN905643](#), [MN639368](#), [MN544117](#).
- Epiperipatus [sp11] MNRJ 0042 ony-027b: [MN905644](#), [MN639369](#), [MN544118](#).
- Epiperipatus [sp11], MNRJ 0042 ony-027c: [MN905645](#), [MN639370](#), [MN544119](#).
- Epiperipatus [sp13] MNRJ 0020 ony-029: [MN639371](#), [MN544120](#).
- Epiperipatus [sp13] MNRJ 0018 ony-030a: [MN639372](#), [MN544121](#).
- Epiperipatus [sp13] MNRJ 0018 ony-030b: [MN639373](#).

- Epiperipatus [sp13] MNRJ 0023 ony-048: [MN905657](#), [MN639383](#), [MN544133](#).
- Epiperipatus [sp13] MNRJ 0026 ony-049: [HG531958](#), [MN933779](#), [MN933788](#).
- Epiperipatus [sp14] MNRJ 0016 ony-032: [MN544122](#).
- Epiperipatus machadoi MNRJ 0043 ony-008: [MN905632](#), [MN639360](#), [MN544108](#).
- Epiperipatus ohausi MNRJ 0058 ony-010: [MN905633](#).
- Epiperipatus ohausi MNRJ 0056 ony-011: [MN905634](#), [MN639361](#), [MN544109](#), [MN705441](#).
- Epiperipatus sp. MNRJ 0100 ony-153: [MN639449](#), [MN544194](#), [MN705490](#).

New Species Registration

The following information was supplied regarding the registration of a newly described species:

Publication LSID: urn:lsid:zoobank.org:pub:A8971D25-1C9D-4807-BE2B-730FB010B717.

Epiperipatus puri species LSID: urn:lsid:zoobank.org:act:5F941DB2-F54D-4E32-9337-0A90B26D99E7.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.15384#supplemental-information>.

REFERENCES

- Baker CM, Buckman-Young RS, Costa CS, Giribet G. 2021. Phylogenomic analysis of velvet worms (Onychophora) uncovers an evolutionary radiation in the neotropics. *Molecular Biology and Evolution* **38**(12):5391–5404 DOI [10.1093/molbev/msab251](https://doi.org/10.1093/molbev/msab251).
- Bouvier E-L. 1905. Monographie des Onychophores. *Annales des Sciences Naturelles, Zoologie* **2**:1–383.
- Brito SV, Pereira JC, Ferreira FS, Vasconcellos A, de Almeida WO. 2010. Epiperipatus cratensis sp. nov. (Onychophora: Peripatidae) from northeastern Brazil. *Neotropical Biology and Conservation* **5**:47–52 DOI [10.4013/nbc.2010.51.07](https://doi.org/10.4013/nbc.2010.51.07).
- Centore P. 2016. sRGB Centroids for the ISCC-NBS Colour System. Available at <https://www.munsellcolourscienceforpainters.com/ISCCNBS/ISCCNBSSystem.html> (accessed 17 July 2019).
- CEPF. 2019. Explore the Biodiversity Hotspots | CEPF. Available at <https://www.cepf.net/our-work/biodiversity-hotspots>.
- Chagas-Jr A, Costa CS. 2014. *Macroperipatus ohausi*: redescription and taxonomic notes on its status (Onychophora: Peripatidae). *Revista de Biología Tropical* **62**:977–985 DOI [10.15517/rbt.v62i3.11643](https://doi.org/10.15517/rbt.v62i3.11643).
- Chagas-Jr A, de Oliveira IS. 2019. Onychophora Grube, 1853. Available at <http://fauna.jbrj.gov.br/fauna/faunadobrasil/19> (accessed 25 August 2023).
- Clark AH. 1913. A revision of the American species of *Peripatus*. *Proceedings of the Biological Society Washington* **26**:15–19.
- Collen B, Böhm B, Kemp R, Baillie JEM. 2012. *Spineless: status and trends of the world's invertebrates*. London: Zoological Society of London, United Kingdom.

- Costa CS.** 2016. Sistemática e análise filogenética de *Epiperipatus* Clark, 1913 baseada em dados moleculares e morfológicos (Onychophora: Peripatidae). Ph.D. Thesis, Universidade de São Paulo.
- Costa CS, Chagas-Jr A, Baptista RLC.** 2009. Brazilian species of Onychophora with notes on their taxonomy and distribution. *Zoologia (Curitiba)* **26**(3):553–561
DOI [10.1590/S1984-46702009005000004](https://doi.org/10.1590/S1984-46702009005000004).
- Costa CS, Chagas-Jr A, Pinto-da-Rocha R.** 2018. Redescription of *Epiperipatus edwardsii*, and descriptions of five new species of *Epiperipatus* from Brazil (Onychophora: Peripatidae). *Zoologia* **35**(2):1–15 DOI [10.3897/zoologia.35.e23366](https://doi.org/10.3897/zoologia.35.e23366).
- Costa CS, Cordeiro LM, Chagas-Jr A.** 2018. *Epiperipatus ohausi* (Bouvier, 1900). In: *Instituto Chico Mendes de Conservação da Biodiversidade*. Vol. VII. Brasília: ICMBio, 577–586.
- Costa CS, Giribet G.** 2016. Taxonomic notes on *Mesoperipatus tholloni* (Onychophora: Peripatidae), an Elusive Velvet Worm from Gabon. *Breviora* **552**(1):1–10
DOI [10.3099/MCZ30.1](https://doi.org/10.3099/MCZ30.1).
- Costa CS, Giribet G, Pinto-da-Rocha R.** 2021. Morphological and molecular phylogeny of *Epiperipatus* (Onychophora: Peripatidae): a combined approach. *Zoological Journal of the Linnean Society* **192**(3):763–793 DOI [10.1093/zoolinnean/zlaa100](https://doi.org/10.1093/zoolinnean/zlaa100).
- Costa RL, Prevedello JA, de Souza BG, Cabral DC.** 2017. Forest transitions in tropical landscapes: a test in the Atlantic Forest biodiversity hotspot. *Applied Geography* **82**(6):93–100
DOI [10.1016/j.apgeog.2017.03.006](https://doi.org/10.1016/j.apgeog.2017.03.006).
- de Oliveira IS, Lacorte GA, Fonseca CG, Wieloch AH, Mayer G.** 2011. Cryptic speciation in Brazilian *Epiperipatus* (Onychophora: Peripatidae) reveals an underestimated diversity among the peripatid velvet worms. *PLOS ONE* **6**(6):e19973 DOI [10.1371/journal.pone.0019973](https://doi.org/10.1371/journal.pone.0019973).
- de Oliveira IS, Lacorte GA, Weck-Heimann A, Cordeiro LM, Wieloch AH, Mayer G.** 2015. A new and critically endangered species and genus of Onychophora (Peripatidae) from the Brazilian savannah—a vulnerable biodiversity hotspot. *Systematics and Biodiversity* **13**(3):211–233 DOI [10.1080/14772000.2014.985621](https://doi.org/10.1080/14772000.2014.985621).
- de Oliveira IS, Read VMSJ, Mayer G.** 2012. A world checklist of Onychophora (velvet worms), with notes on nomenclature and status of names. *ZooKeys* **211**:1–70
DOI [10.3897/zookeys.211.3463](https://doi.org/10.3897/zookeys.211.3463).
- de Oliveira IS, Schaffer S, Kvartalnov PV, Galoyan EA, Palko IV, Weck-Heimann A, Geissler P, Ruhberg H, Mayer G.** 2013. A new species of *Eoperipatus* (Onychophora) from Vietnam reveals novel morphological characters for the South-East Asian Peripatidae. *Zoologischer Anzeiger* **252**:495–510 DOI [10.1016/j.jcz.2013.01.001](https://doi.org/10.1016/j.jcz.2013.01.001).
- de Oliveira IS, Wieloch AH, Mayer G.** 2010. Revised taxonomy and redescription of two species of the Peripatidae (Onychophora) from Brazil: a step towards consistent terminology of morphological characters. *Zootaxa* **2493**:16–34 DOI [10.5281/zenodo.195686](https://doi.org/10.5281/zenodo.195686).
- Dobrovolski R, Loyola R, Rattis L, Gouveia SF, Cardoso D, Santos-Silva R, Gonçalves-Souza D, Bini LM, Diniz-Filho JAF.** 2018. Science and democracy must orientate Brazil's path to sustainability. *Perspectives in Ecology and Conservation* **16**(3):121–124
DOI [10.1016/j.pecon.2018.06.005](https://doi.org/10.1016/j.pecon.2018.06.005).
- Escobar H.** 2019. ‘We can’t take another hit like this’: Brazilian scientists lament big budget freeze. *Science AAAS* DOI [10.1126/science.aax6227](https://doi.org/10.1126/science.aax6227).
- Ferrante L, Gomes M, Fearnside PM.** 2020. Amazonian indigenous peoples are threatened by Brazil’s Highway BR-319. *Land Use Policy* **94**(2):104548 DOI [10.1016/j.landusepol.2020.104548](https://doi.org/10.1016/j.landusepol.2020.104548).
- Froehlich CG.** 1968. On some Brazilian Onychophores. *Beiträge zur Neotropischen Fauna* **5**(3):160–171 DOI [10.1080/01650526809360404](https://doi.org/10.1080/01650526809360404).

- Giribet G, Buckman-Young RS, Sampaio Costa C, Baker CM, Benavides LR, Branstetter MG, Daniels SR, Pinto-da-Rocha R.** 2018. The “Peripatos” in Eurogondwana?—Lack of evidence that South-east Asian onychophorans walked through Europe. *Invertebrate Systematics* 32(4):842–865 DOI [10.1071/IS18007](https://doi.org/10.1071/IS18007).
- Guagliardi R.** 2018. *RPPN—private natural heritage reserves state program: 10 years in support of biodiversity conservation.* Instituto Estadual do Ambiente: Rio de Janeiro.
- IBGE.** 2019. *Biomes and coastal-marine system of Brazil: compatible with the 1:250,000 scale [Bacias e sistema costeiro-marinho do Brasil: compatível com a escala 1:250 000].* Rio de Janeiro: Instituto Brasileiro de Geografia e Estatística. (In Portuguese).
- ICMBio.** 2018. *Red Book of Brazilian Fauna Threatened with Extinction 2018 [Livro Vermelho da Fauna Brasileira Ameaçada de Extinção].* Brasília: Instituto Chico Mendes de Conservação da Biodiversidade. (In Portuguese).
- Instituto Nacional de Pesquisas Espaciais (INPE).** 2023. Programa Queimadas. Available at <http://terrabrasilis.dpi.inpe.br/queimadas> (accessed 25 August 2023).
- Joly CA, Metzger JP, Tabarelli M.** 2014. Experiences from the Brazilian Atlantic Forest: ecological findings and conservation initiatives. *New Phytologist* 204(3):459–473 DOI [10.1111/nph.12989](https://doi.org/10.1111/nph.12989).
- Kelly KL.** 1958. Central notations for the revised iscc-nbs color-name blocks. *Journal of Research of the National Bureau of Standards* 61:427 DOI [10.6028/JRES.061.035](https://doi.org/10.6028/JRES.061.035).
- Kompiér T.** 2015. *A guide to the dragonflies and damselflies of the Serra dos Orgaos [sic] South-eastern Brazil.* Brussels: REGUA Publications.
- Kury AB, Giupponi API, Mendes AC.** 2018. Immolation of Museu Nacional, Rio de Janeiro—unforgettable fire and irreplaceable loss. *The Journal of Arachnology* 46(3):556–558 DOI [10.1636/JoA-S-18-094.1](https://doi.org/10.1636/JoA-S-18-094.1).
- Laurance WF.** 2009. Conserving the hottest of the hotspots. *Biological Conservation* 142(6):1137 DOI [10.1016/j.biocon.2008.10.011](https://doi.org/10.1016/j.biocon.2008.10.011).
- le Bras G, Geoffroy J-J, Albenga L, Mauriès J-P.** 2015. The Myriapoda and Onychophora collection (MY) of the Muséum national d’Histoire naturelle (MNHN, Paris). *ZooKeys* 518(1):139–153 DOI [10.3897/zookeys.518.10223](https://doi.org/10.3897/zookeys.518.10223).
- Martelli-Jr H, Martelli DR, Silva ACS e, Oliveira MCL, Oliveira EA.** 2019. Brazil’s endangered postgraduate system. *Science* 363(6424):240 DOI [10.1126/science.aav9015](https://doi.org/10.1126/science.aav9015).
- Mittermeier RA, Gil PR, Hoffmann M, Pilgrim J, Brooks T, Mittermeier CG, Lamoreux J, Fonseca GABD.** 2004. *Hotspots revisited: earths biologically richest and most endangered terrestrial ecoregions.* Mexico City: CEMEX.
- Morera-Brenes B, Monge-Nájera J.** 1990. *Epiperipatus hilkae m sp from Costa Rica* (Onychophora: Peripatidae). *Revista de Biología Tropical* 38:449–455.
- Morera-Brenes B, Monge-Nájera J.** 2010. A new giant species of placented worm and the mechanism by which onychophorans weave their nets (Onychophora: Peripatidae). *Revista de Biología Tropical* 58:1127–1142 DOI [10.15517/rbt.v58i4.5398](https://doi.org/10.15517/rbt.v58i4.5398).
- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J.** 2000. Biodiversity hotspots for conservation priorities. *Nature* 403(6772):853–858 DOI [10.1038/35002501](https://doi.org/10.1038/35002501).
- New TR.** 1995. Onychophora in invertebrate conservation: priorities, practice and prospects. *Zoological Journal of the Linnean Society* 114:77–89 DOI [10.1006/zjls.1995.0017](https://doi.org/10.1006/zjls.1995.0017).
- Peck SB.** 1975. A review of the New World Onychophora with the description of a new cavernicolous genus and species from Jamaica. *Psyche: A Journal of Entomology* 82(3–4):341–358 DOI [10.1155/1975/98614](https://doi.org/10.1155/1975/98614).

- Pimentel L, Olmos F.** 2011. The birds of Reserva Ecológica Guapiaçu (REGUA), Rio de Janeiro, Brazil. *Cotinga* 33:8–24.
- Pinto AP, Kompiér T.** 2018. In honor of conservation of the Brazilian Atlantic Forest: description of two new damselflies of the genus *Forcepsioneura* discovered in private protected areas (Odonata: Coenagrionidae). *Zoologia* 35(2):1–19 DOI 10.3897/zootaxa.35.e21351.
- Read VMSJ.** 1988. The application of scanning electron microscopy to the systematics of the neotropical Peripatidae (Onychophora). *Zoological Journal of the Linnean Society* 93(3):187–223 DOI 10.1111/j.1096-3642.1988.tb01361.x.
- Ribeiro MC, Martensen AC, Metzger JP, Tabarelli M, Scarano F, Fortin M-J.** 2011. The Brazilian Atlantic Forest: a shrinking biodiversity hotspot. In: Zachos FE, Habel JC, eds. *Biodiversity Hotspots: Distribution and Protection of Conservation Priority Areas*. Heidelberg [Germany]. New York: Springer, 405–434.
- Ribeiro MC, Metzger JP, Martensen AC, Ponzoni FJ, Hirota MM.** 2009. The Brazilian Atlantic Forest: how much is left, and how is the remaining forest distributed? Implications for conservation. *Biological Conservation* 142(6):1141–1153 DOI 10.1016/j.biocon.2009.02.021.
- Rodríguez Osuna V, Börner J, Nehren U, Prado RB, Gaese H, Heinrich J.** 2014. Priority areas for watershed service conservation in the Guapi-Macacu region of Rio de Janeiro, Atlantic Forest, Brazil. *Ecological Processes* 3(1):16 DOI 10.1186/s13717-014-0016-7.
- Sedgwick A.** 1888. A monograph on the species and distribution of the genus *Peripatus*. *Quarterly Journal of Microscopic Science* 28:431–493 DOI 10.1242/jcs.s2-28.112.431.
- Silva JS, Souto Couri M, de Giupponi APL, Alencar J.** 2014. Mosquito fauna of the Guapiaçu Ecological Reserve, Cachoeiras de Macacu, Rio de Janeiro, Brazil, collected under the influence of different color CDC light traps. *Journal of Vector Ecology* 39(2):384–394 DOI 10.1111/jvec.12114.
- Soares A, Bizarro JMS, Bastos CB, Tangerini N, Silva NA, da Silva AS, Silva GB.** 2011. Preliminary analysis of the diurnal Lepidoptera fauna of the Três Picos State Park, Rio de Janeiro, Brazil, with a note on *Parides ascanius* (Cramer, 1775). *Tropical Lepidoptera Research* 22:66–79.
- Sosa-Bartuano Á, Monge-Nájera J, Morera-Brenes B.** 2018. A proposed solution to the species problem in velvet worm conservation (Onychophora). *UNED Research Journal* 10(1):193–197 DOI 10.22458/urj.v10i1.2027.
- Wilson EO.** 1987. The Little Things That Run the world* (The Importance and Conservation of Invertebrates). *Conservation Biology* 1(4):344–346 DOI 10.1111/j.1523-1739.1987.tb00055.x.
- W3Schools.** 2023. Colors—ISCC/NBS. Available at https://www.w3schools.com/colors/colors_nbs.asp (accessed 25 August 2023).