


1902

USNM

EX LIBRIS

William Healey Dall

Division of Mollusks
Sectional Library



Redue

Division of Mollusks
Sectional Library

CONCHOLOGIA ICONICA:

OR,

ILLUSTRATIONS

OF THE

SHELLS OF MOLLUSCOUS ANIMALS.

VOL. XIV.

CONTAINING MONOGRAPHS OF THE GENERA

HALIA.	CHONDROPOMA.	CHAMOSTREA.
CONCHOLEPAS.	ADAMSIELLA.	VENUS.
ZIZYPHINUS.	ANASTOMA.	DIONE.
TEREBELLUM.	TRIDACNA.	CIRCE.
PALUDINA.	HIPPOPUS.	CY THEREA.
CYCLOTUS.	ANATINA.	TAPES.
PTERO CYCLOS.	TUGONIA.	MEROË.

BY

LOVELL AUGUSTUS REEVE, F.L.S., F.G.S.,

CORRESPONDING MEMBER OF THE NATURAL HISTORY SOCIETY OF WÜRTEMBERG, OF THE LYCEUM OF NATURAL HISTORY OF NEW YORK, OF THE ACADEMY OF NATURAL SCIENCES OF PHILADELPHIA, AND OF THE ZOOLOGICO-BOTANICAL SOCIETY OF VIENNA.

“O, quantum Natura varia est! componit, coagmentat, separat,
iterum jungit, ludit quoque, ut fallere videatur; sed manet simplicissima,
manet veridica natura:—post Biblia sacra, nullus mihi earior libro
naturæ.”—*Klein.*

LONDON:

LOVELL REEVE & CO., HENRIETTA STREET, COVENT GARDEN.

1864.



MONOGRAPH

OF THE GENUS

H A L I A.


"Meditation here
May think down hours to moments." — *Corper.*

EX LIBRIS



123

JOHN BROOKS HENDERSON



H A L I A.

PLATE I.

Genus HALIA, Riso.

Testa oblongo-ovata, ventricosa, tenuicula, spirā subexsertā, apice obtusā, anfractibus quinque ad sex, lēvigatis, subcorneis, supernè obtuse declivi-angulatis, columellā arcuatā, versus basin caloso-involutā, truncatā; aperiturā amplā, supra et infra sinuatā, labro simplici.

Shell oblong-ovate, ventricose, rather thin, spire somewhat exserted, obtuse at the apex, whorls five to six, smooth, slightly horny, obtusely slopingly angled, columella arched, callously involute towards the base, truncated; aperture large, sinuated above and below, lip simple.

The very characteristic shell, of which back and front views of two specimens are given in the accompanying plate, appears to have been first known about the same time in Germany and England, between eighty and ninety years ago. In Germany it was named by Meuschen, in the Catalogue of the Gronovian Museum (1778), *Helix Priamus*, indicative of its belonging to a terrestrial mollusk. In England its generic character and habitat were more correctly designated, for it was named by Martyn in his 'Universal Conchologist,' *Buccinum fucus*, from Spain, and illustrated by two very beautiful drawings.

Various conjectures have been made by naturalists as to its place in the system, some referring it to *Bulla*, but the greater number, including all the higher authorities,—Bruguière, Lamarck, Féruccac,—have inclined to the opinion that it must belong to a land snail of the *Glandina* section of *Achatina*. After the lapse of more than half a century, the notion of its being the shell of a sea mollusk was revived by the Danish conchologist, Dr. Beck, who became possessed of some new information on the subject, which he communicated to M. Deshayes while engaged upon his edition of Lamarck's *Anim. sans vert.* In vol. viii. of that work, in a note to p. 300, published in 1838, M. Deshayes announced that the shell under consideration could no longer remain in *Achatina*, Dr. Beck having informed him that it belonged to a marine operculate mollusk living in the seas of Spain and Portugal, and that it must henceforth rank as a new genus allied to *Buccinum*, for which the old specific name of *Priamus* was proposed. This information helped to confirm the original statement of Martyn, which has been overlooked by all subsequent writers, that the shell was a "*Buccinum* from Spain;" but the intimation conveyed by Dr. Beck, that the animal is operculate, has been since shown to be erroneous.

In 1846 it was discovered by Herrmannsen that the *Priamus* is identical with a fossil of the miocene beds of Turin, named by Brocchi *Bulla helicoides* (Conch. Foss. Subapp. 1814, p. 283), for which Risso had in 1826 (Hist. Nat. Prod. Europe Mérid. vol. iv.) proposed a new genus with the name of *Halalia*. No doubt is now entertained of the recent *Helix Priamus* of Meuschen being identical with the fossil *Bulla helicoides* of Brocchi, and now that our knowledge of the animal justifies us in making it the type of a separate genus the generic name of *Priamus* gives way to that of *Halalia*, proposed twelve years before by Risso.

In 1858 all doubt as to the character and habits of this mollusk were set at rest in an able monograph by M. P. Fischer, of Paris, published in the third volume of the second series of the 'Journal de Conchyliologie.' M. Fischer already possessed a fragment of the animal, collected by a French botanist, M. Picard, travelling on the coast of Spain, when a more perfect specimen, preserved in spirits, sent to him by a Spanish gentleman, M. de Elizalde, enabled him to complete the anatomy. The result of M. Fischer's dissection went to show that there were no traces of an opereuligerous lobe nor of any operculum, and that in the principal features of its organization the animal is nearest allied to the inoperculated *Desfrancia* section of *Pleurotomæ*.

Another specimen of *Halalia*, with the animal, lately received from Vigo Bay by Mr. Darbshire, of Manchester, has, I believe, been submitted for dissection to the care of Mr. Albany Hancock, of Newcastle.

Species 1. (Fig. a, b, Mus. Cuming; Fig. c, d, Mus. Newman Smith.)

HALIA PRIAMUS. *Hal. testā oratā, tenuiculā, ventricosā, fulvo-spadiceā, subcorneā, maculis rubro-castaneis quadratis subremolis, nisi circa anfractuum partem superiorem, fascialim aspersā, anfractibus lēvibus, nitentibus, interdum quasi encausticis, vel longitudinaliter arcuatim plicato-striatis, columelli arcuatā, callosā, subcontortā; aperiturā amplā, utrinque sinuatā, labro simplici.*

THE PRIAM HALIA. Shell ovate, rather thin, ventricose, fulvous-fawn, rather horny, sprinkled except about the upper part of the whorls with bands of distant square reddish-chestnut spots, whorls smooth, shi-

HALIA.—PLATE I.

ming, sometimes as though enamelled, or longitudinally areately plicately striated, columella arched, callous, slightly twisted; aperture large, sinuated at both ends, lip simple.

Helix Priamus, Menschen, Cat. Mus. Gronov. no. 1355.

Buccinum fucus, Martyn, Univ. Conch. vol. iii. pl. 81.

Buccinum stercus-pulicium, Chemnitz.

Bulla stercus pulicium, Gmelin.

Bulimus Priamus, Brugnière.

Bulla helicoides, Brocchi.

Bulla Priamus, Dillwyn.

Impulla Priamus, Bolten.

Ichatina Priamus, Lamarek.

Helix (Cochlicopa) Priamus, Féussae.

Halia helicoides, Risso.

Priamus stercus-pulicium, Beck.

Ichatina maculata, Swainson.

Halia stercus-pulicium, Hermannsen.

Priamus helicoides, Sismonda.

Halia Priamus, Woodward.

Hab. Atlantic coast of Spain.

The glairy enamel-like surface of this shell shows that it is largely enveloped by the mantle of the animal; in most specimens the permanent contact of the mantle to the under surface being widely marked. It is always of a peculiar fulvous-fawn or nankeen colour, sprinkled in the suture and below the angle of the whorls with bands of small square distant reddish-chestnut spots. The plicate sinuation of the lines of growth at the suture indicates a development of shell-matter somewhat analogous to the growth of the shells of the family to which M. Fischer inclines to refer it.

MONOGRAPH
OF THE GENUS
CONCHOLEPAS.

"Where rose the mountains, there to him were friends :
Where roll'd the ocean, thereon was his home."—*Byron*.



CONCHOLEPAS.

PLATE I.

Genus CONCHOLEPAS, Lamarck.

Testa ovata, ventricosissima, spirā minutā, anfractibus tribus, rapidissimè accrescentibus, ultimo peramplo, aperturā patulā, ad basin modicè canaliculatā, labro continuo, juxta canalem bidentato. Operculum cornuum, parviusculum.

Shell ovate, extremely ventricose, spire minute, whorls three, increasing very rapidly, last whorl very large, aperture wide open, moderately channelled, lip continuous, two-toothed next the canal. Operculum horny, rather small.

Concholepas is a form of *Purpura* inhabiting the western coast of South America, from Ecuador to Cape Horn, round to the eastern shores of Tierra del Fuego. The growth of the shell is limited to three whorls, and so small, consequently, is the spire that it has very much the appearance of a Limpet. The sculpture radiates from the apex in ribs and ridges, and these are mostly imbricated with concentric frills of scales. As in *Purpura* and *Monoceros*, the aperture is notched and toothed at the base, the lip being continuous in an expanded form over the columella. I venture to characterize two species.

Species 1. (Fig. *a* to *d*, Mus. Cuming.)

CONCHOLEPAS PERUVIANUS. *Couch. testā ovalā, ventricosā, rufescente-brunneā, in testā juniore interdum albo-maculatā, radiatim costatā et liratā, costis lirisque subirregularibus, concentricè plus minus fimbriatosquamatis.*

THE PERUVIAN CONCHOLEPAS. Shell ovate, ventricose, reddish-brown, sometimes blotched in the young shell with white, radiately ribbed and ridged, ribs and ridges rather irregular, concentrically more or less fimbriately scaled.

LAMARCK, Anim. sans vert. vol. 10. p. 126.

Buccinum concholepas, Bruguière.

Patella lepas, Gmelin.

Buccinum lepas, Burrows.

Purpura Peruviana, De Blainville.

Hab. Peru.

The typical state of this species is with the outer surface elaborately imbricated with concentric frills of scales. Specimens are, however, found with the scales almost obsolete. In the young shell there are mostly blotches of white.

June, 1863.



2b.



2c.



2a.

CONCHOLEPAS.

PLATE II.

Species 2. (Fig. *a*, *b*, *c*, Mus. Cuming.)

CONCHOLEPAS OBLONGUS. *Couch. testá oblongo-ovatá, ad basin subattenuatá, nigricante-brunnea, radiatim liratá, liris subirregularibus, rudibus, obsoletè nodulatis.*
THE OBLONG CONCHOLEPAS. Shell oblong-ovate, somewhat attenuated towards the base, blackish-brown, radiately ridged, ridges rather irregular, rude, obsoletely noduled.


Hab. Cape Horn, and Bay of Good Success, East Tierra del Fuego.

Mr. Cuming has long kept this oblong form of *Concholepas* separate from his numerous varieties of the Peruvian species. It is from Cape Horn and the East side of Tierra del Fuego. Time will show whether its characteristic proportions and rude blunted sculpture are permanent.

June, 1863.

MONOGRAPH
OF THE GENUS
ZIZYPHINUS.

"O Nature! by impassioned hearts alone
Thy genuine charms are felt. The vulgar mind
Sees but the shadow of a power unknown."—*Pringle*.



ZIZYPHINUS.

PLATE I.

Genus ZIZYPHINUS, Leach.

*Tesla subacutè conica, rarò umbilicata, variè colorata, sæ-
pissime grano-lirata, interdum lœvis, aperturā obliquè
subquadratā, labro simplici, columellā plus minus cal-
losā, argenteā, rarò dentata. Operculum cornuum,
tenue, circulariter spirale.*

Shell rather sharply conical, rarely umbilicated, variously coloured, mostly grain-ridged, sometimes smooth, aperture obliquely subquadrate, lip simple, columella more or less callous, silvery, seldom toothed. Operculum horny, thin, circularly spiral.

The genus *Zizyphinus* was proposed by Dr. Leach for the reception of the British species of *Trochus*, six or seven in number, partaking of the general typical character of the well-known Linnaean *Trochus zizyphinus*. At New Zealand and Western Australia this particular generic type is still more characteristically developed; there is, indeed, good ground for believing that the above-named species is possessed by the very remote seas of Europe and Australia in common. The shells are rather sharply conical, mostly sculptured with rows of grains or grained ridges, and the prevailing colour of the species is that of the *Zizyphum*, or Jujube-fruit, a dull flesh-tinged yellow, or yellowish-brown. The columellar axis of the shell is callous and silvery, arched, but rarely ending in a tooth, and still more rarely umbilicated. The operculum is horny, thin, and regularly circularly spiral.

Of about fifty species of *Zizyphius* inhabiting the eastern hemisphere, at least half are natives of the seas of West Australia and New Zealand, eight are European, one North African, and two Japanese. Eight inhabit the Philippine Islands, Tranquebar, and Malacca, and one the shores of Bombay. Only seven species are referred to this genus from the New World, five from California, Honduras, and Panama, one from Puget Sound, and one from Lord Hood's Island in the Pacific.

Species 1. (Mus. Cuming.)

*ZIZYPHINUS SELECTUS. Zizyph. testa subdepresso-cono-
dea, tumida, versus apicem attenuata, anfractibus de-
cliri-rotundatis, spiraliter grano-liralis, carneo-alba
aut flavescente, granis irregulariter ustulato-rufo tinctis,
colore obscurissimè biseriatim maculato.*

THE CHOICE ZIZYPHINUS. Shell rather depressedly conoid, swollen, attenuated towards the apex, whorls slopingly rounded, spirally grain-ridged, flesh-white or yellowish, grains irregularly stained with burnt red, the colour being very obscurely disposed in two rows of blotches.

Trochus selectus, Chemnitz, Conch. Cab. vol. xi. p. 168.
pl. 196. f. 1896, 1897.

Hab. New Zealand.

Distinguished from *Z. granatus* by its depressedly swollen proportions and the promiscuous burnt-red colouring of the grains, which stand out more conspicuously on a paler ground. Next the suture, and at the periphery of the whorls, the colour is very obscurely disposed in blotches.

Species 2. (Mus. Cuming.)

*ZIZYPHINUS GRANATUS. Zizyph. testa regulariter conoidea,
anfractibus declici-concexis, densè grano-liratis, rufes-
cente-alba vel carneâ, infra suturas subobscure rufo
strigato-maculata, granis alternatim, vel in interstitiis,
rufis.*

THE GRAINED ZIZYPHINUS. Shell regularly conoid, whorls slopingly convex, densely grain-ridged, reddish white or flesh-tinged, rather obscurely streak-spotted beneath the sutures, grains red alternately or in the interstices.

Trochus granatus, Chemnitz, Conch. Cab. vol. v. p. 100.
pl. 170. f. 1654, 1655.

Hab. Tasmania.

Although this shell is much smaller than that represented by Chemnitz, there is every reason to believe that it is an average example of the species. It is of a more regularly conical form than the preceding species, differently coloured. Martyn's *Trochus tigris*, which Deshayes refers to this species, is quite another shell, for which see Sp. 4.

Species 3. (Mus. Cuming.)

*ZIZYPHINUS ARTICULATUS. Zizyph. testa depresso-conoidea,
versus apicem attenuata, anfractibus depresso-concexis,
ad peripheriam subangulatis, spiraliter grano-liratis,*

ZIZYPHINUS.—PLATE I.

liris alternatim majoribus, linearibus, continuis, carneá, liris albido et rufo nitidè articulatis, basi lærigatá, circulariter incisá, rufo remotè articulatá.

THE ARTICULATED ZIZYPHINUS. Shell depressedly conoid, attenuated towards the apex, whorls depressedly convex, slightly angled at the periphery, spirally grain-ridged, ridges alternately large, linear, continuous, flesh-colour, ridges neatly articulated with red; base smooth, circularly incised, distantly articulated with red.

Hab. —?

A delicate depressedly-conoid species in which the ridges are alternately large and smaller, and uniformly narrow, linear, and continuous. The colouring is disposed in neat articulations, which are present in a very characteristic manner at the base of the shell.

Species 4. (Mus. Cuming.)

ZIZYPHINUS TIGRIS. *Zizyph. testá inflato-conicá, versus apicem attenuato-elevatá, anfractibus declivi-convergis, densè grano-liratis, liris læriusculis, superficiariis, albidá, rufo-fusco undique consertim obliquè promiscue strigata.*

THE TIGER ZIZYPHINUS. Shell inflatedly conical, attenuately raised towards the apex, whorls slopingly convex, densely grain-ridged, ridges rather smooth, superficial, whitish, everywhere closely obliquely promiscuously streaked with red-brown.

Trochus tigris, Martyn, Univ. Conch. pl. 75.

Zizyphinus tigris, Adams.

Hab. New Zealand.

This fine species, the largest of the genus, is at once distinguished by its handsome turban-like form, slopingly convex and rather ventricose round the periphery, peculiarly attenuately raised towards the apex.

Species 5. (Fig. a, b, Mus. Cuming.)

ZIZYPHINUS SPECTABILIS. *Zizyph. testó regulariter conoidea, solidá, subponderosa, anfractibus declivi-convergis, grano-liratis, liris subdistantibus, granis grandibus, lirarum interstiliis læribus, excavatis, carneo aut flavescente, liris rosaceo-rubo punctatis.*

THE REMARKABLE ZIZYPHINUS. Shell regularly conoid, solid, rather ponderous, whorls slopingly convex, grain-ridged, ridges rather distant, grains large, interstices between the ridges smooth, excavated, flesh-colour or yellowish, ridges dotted with pinkish-red.

ADAMS, Pro. Zool. Soc. 1854, p. 37.

Hab. New Zealand.

This fine species is conspicuously distinguished from any other of the genus by its rather ponderous growth, pinkish-red colour, and bold sculpture. The ridges are almost squamately grained, and the interstices between them are smooth and excavated.

Species 6. (Mus. Cuming.)


ZIZYPHINUS CUNNINGHAMII. *Zizyph. testá latè depresso-conoidea, apice attenuatá, anfractibus plano-declinibns, parum convexis, ultimo ad basin subacutè angulato, grano-liratis, liris subtilibus, inconspicuis, alternatim minoribus, livido-carneá, liris rufescente punctatis.*

CUNNINGHAM'S ZIZYPHINUS. Shell broadly depressedly conoid, attenuated at the apex, whorls flatly sloping, but little convex, the last rather sharply angled at the base, grain-ridged, ridges fine, inconspicuous, alternately smaller; livid flesh-colour, ridges dotted with pale-red.

GRAY, Griffith's Cuv. Anim. Kingdom.

Hab. New Zealand.

A shell of rather delicate substance, depressed, and unusually wide-spread at the base.



ZIZYPHINUS.

PLATE II.

Species 7. (Fig. a, b, Mus. Cuming.)

ZIZYPHINUS ORNATUS. *Zizyph. testā obliquē conoideā, basi dilatata, anfractibus convexis, dense grano-striatis, ultimo decliviore, ad peripheriam obtuse angulato, grainis fortioribus; flavescente, carneo-rufo oblique areuatim nebulata et flammata, basi subaurantiā, interdum rufo punctato-articulata.*

THE ORNATE ZIZYPHINUS. Shell obliquely conoid, dilated at the base, whorls convex, densely grain-striated, last whorl more sloping, obtusely angled at the periphery, grains stronger; yellowish, obliquely areuatedly clouded and flamed with flesh-red, base tinged with orange, sometimes dot-articulated with red.

Trochus ornatus, Lamarek, Anim. sans vert. vol. ix. p. 146.

Zizyphinus ornatus and *nebulosus*, A. Adams.

Hab. Torres Straits, North Australia; Inc.

This species resembles the following in general aspect, but it is well characterized by Lamarek in being grain-striated and dilated at the base. In *L. unicus* the last whorl is conspicuously rounded.

Species 8. (Mus. Cuming.)

ZIZYPHINUS UNICUS. *Zizyph. testā obliquē conoideā, basi ventricoso-rotundata, anfractibus rotundatis, dense lirato-striatis, parum granatis, basi levigata, circulatiter incisa; albidā vel flavescente, sanguineo-rufo sub-remotè flammata, ad peripheriam conspicue articulata.*

THE UNIQUE ZIZYPHINUS. Shell obliquely conoid, ventricosely rounded at the base, whorls rounded, densely ridge-striated, but little grained, base smooth, circularly engraved; whitish or yellowish, rather distantly flamed with blood-red, conspicuously articulated at the periphery.

DUNKER, Moll. Japonica.

Hab. Japan; A. Adams.

Very closely allied, and very similar in painting, to the Australian *Z. ornatus*. The surface is much less grained, the last whorl of the shell is ventricosely rounded, and the periphery is encircled by a well-defined articulated band of rather distant red spots.

Species 9. (Fig. a, b, Mus. Cuming.)

ZIZYPHINUS SPECIOSUS. *Zizyph. testā erecto-conicā, sub-pyramidalis, ad basin depresso-rotundata, anfractibus subconcavo-declivibus, undique grano-liratis, liris tenibus, lirā parrā levigata interveniente; flavescente-albidā, purpureo-rufo obscurè tessellata et punctato-gemmatā, basi purpureo-punctata, callo centrali ustulato-aurantio tingita.*

THE BEAUTIFUL ZIZYPHINUS. Shell erectly conical, somewhat pyramidal, depresso roinded at the base, whorls rather concavely sloping, grain-ridged throughout, ridges fine, with a rather smaller smooth ridge coming between; yellowish white, obscurely tessellated and dot-gemmed with purple, base purple-dotted, central callosity tinged with burnt orange.

ADAMS, Pro. Zool. Soc. 1854, p. 38.

Hab. Moreton Bay, Australia; Strange.

The colouring of this fine species is very promiscuously distributed. Faint tessellated blotches, accompanied by a general sprinkling of dots in livid purple, are seen throughout, sparse and irregular.

Species 10. (Mus. Cuming.)

ZIZYPHINUS NOBILIS. *Zizyph. testā acutē conicā, ad basin angulata, anfractibus plano-declivibus, validē grano-liratis, lirā basali majore, subprominente; flavescente, ferrugineo-rufo tessellata, liris alternatim purpureo-roseis.*

THE NOBLE ZIZYPHINUS. Shell sharply conical, angled at the base, whorls flatly sloping, strongly grain-ridged, basal ridge larger, rather prominent; yellowish, tessellated with rust-brown, ridges alternately purple-rose.

Trochus nobilis, Philippi, Kuster, Conch. Cab. p. 86. pl. 15. f. 6.

Hab. Swan River.

An exquisitely gemmed species, in which the ridges are alternately coloured purple-rose, both on the upper surface of the shell and at the base.

Species 11. (Mus. Cuming.)

ZIZYPHINUS SPLENDIDUS. *Zizyph. testá subacutè conicā, tumidiusculā, anfractibus convexo-declivibus, undique regulariter grano-liratis; albidiā, aurantio-flavo et purpureo-roseo undique variegatā et maculatā.*

THE SPLENDID ZIZYPHINUS. Shell rather sharply conical, somewhat swollen, whorls convexly sloping, regularly grain-ridged throughout; whitish, variegated and spotted throughout with orange-yellow and purple-rose.

PHILIPPI, MS. in Mus. Cuming.

Hab. Australia.

In colour this species partakes very much of the character of the preceding, the purple-rose being more conspicuous, and broken up into blotches. The shell is of rather a lighter, more tumid growth, with the basal ridge no more prominent than the rest.

Species 12. (Mus. Cuming.)

ZIZYPHINUS JUJUBINUS. *Zizyph. testá pyramidali-conicā, solidiusculā, anfractibus concavo-declivibus, deinde tumido-rotundatis, undique granatis, granis parvis, subdistantibus; figlinā, unfractuum parte inferiori salutatā, albo-flavamatā, basi minutè albo-punctatā.*

THE JUJUBE ZIZYPHINUS. Shell pyramidal conical, rather solid, whorls concavely sloping, then tumidly rounded, everywhere grained, grains small, rather distant; brick-red, lower part of the whorls darker, flamed with white, base minutely white-dotted.

Trochus jujubinus, Gmelin, Syst. Nat. p. 3570.

Trochus Zizyphinus, Chemnitz.

Zizyphinus jujubinus, Adams.

Hab. Swan River.

Of rather solid growth, minutely granulated, the lower part of each whorl tumidly bulging, and flamed with white.

Species 13. (Mus. Cuming.)

ZIZYPHINUS ANNULATUS. *Zizyph. testá subacutè conicā, tenuiculā, anfractibus convexo-planatis, eximie granoliratis, interstitiis subexcavatis, aureo-flavā, ad marginem basalem anfractuum vividè cœruleo-purpureā, liris purpureo-sanguineo minutè punctatis.*

THE RINGED ZIZYPHINUS. Shell rather sharply conical, somewhat thin, whorls convexly flattened, delicately grain-ridged, interstices rather excavated; golden yellow, brightly stained with blue-purple round the basal edge of each whorl, ridges minutely dotted with purple-blood.

Trochus annulatus, Martyn, Univ. Coneh. vol. i. f. 33.

Trochus virgineus, Chemnitz.

Zizyphinus annulatus, Adams.

Hab. Monterey, California.

This well-known species is remarkable for the clear definition and constancy of its painting—a bright golden yellow, encircled with minutely red-dotted grained ridges, bordered with a clear blue-purple band winding round the basal margin of each whorl, from the apex to the aperture.

Species 14. (Mus. Cuming.)

ZIZYPHINUS CASTRA. *Zizyph. testá latè conoideā, depresso-acutangulā, tenuiculā, anfractibus planō-declivibus, spiraliter linearī-incisis, infra suturas obliquè plicato-crenulatis, ad marginem basalem funiculatis; pallidè carneā, obscurè tessellatā et articulatā.*

THE CAMP ZIZYPHINUS. Shell broadly conoid, depressed, sharp-angled, rather thin, whorls flatly sloping, spirally linearly engraved, obliquely plicately crenulated below the sutures, corded at the basal margin; pale flesh-colour, obscurely tessellated and articulated.

Hab. Swan River.

Chiefly remarkable for its light, broadly depressed form and almost total absence of sculpture.




15




16a.



17.



18



16b.



16c.



19



20



21a.



21b

ZIZYPHINUS.

PLATE III.

Species 15. (Mus. Cuming.)

ZIZYPHINUS TRANQUEBARICUS. *Zizyph.* testā sublatē conoideā, anfractibus plano-declivibus, spiraliter linearī-incisis, ad marginem basalem conspicuē annuloso-funiculatis, funi interdum dupli, flavescente, pallidē lividā, maculis parvis carneo-rubidis undique lentiginosā et radiatā.

THE TRANQUEBAR ZIZYPHINUS. Shell rather broadly conoid, whorls flatly sloping, spirally linearly engraved, conspicuously annulosely corded at the basal margin, cords sometimes double, yellowish, faintly livid, freckled and rayed throughout with flesh-red small spots.

Trochus Tranquebaricus, Chemnitz, Conch. Cab. vol. v. p. 68. pl. 166, ff. 1595, 1596.

Hab. Tranquebar.

A smooth, glossy, flesh-spotted shell, encircled round the basal edge of the whorls with a very conspicuous double cord.

Species 16. (Fig. a, b, c, Mus. Cuming.)

ZIZYPHINUS CONULOIDES. *Zizyph.* testā subobliquē conicā, ad basin tumidiusculā, anfractibus pluno-declivibus, interdum convexiusculis, spiraliter linearī-incisis vel liratis, ad marginem basalem modicē funiculatis, flavescente vel lividā, carneo-ruso plus minus maculatā.

THE CONE-LIKE ZIZYPHINUS. Shell rather obliquely conical, a little swollen at the base, whorls flatly sloping, sometimes rather convex, spirally linearly engraved or ridged, moderately corded at the basal edge of the whorl, yellowish or livid, more or less blotched with flesh-red.

Trochus zizyphiinus and *conuloides*, Lamarek, Anim. sans. vert. vol. ix. p. 142.

Trochus discrepans, Brown.

Trochus Lyonsii, Leach.

Trochus albus, Wood.

Trochus ciliaris, Menke.

Trochus agrestis, Philippi.

Hab. European and Australian Seas.

It is a curious feature in the geographical distribution of this well-known species that it appears both in European and Australian waters. The specimen represented

at Fig. 10a is *T. ciliaris* from the north-west coast of Australia, described by Menke, Moll. Nov. Holland. p. 17.

Species 17. (Mus. Cuming.)

ZIZYPHINUS EUGLYPTUS. *Zizyph.* testā conicā, solidiusculā, anfractibus superne concavis, deinde tumido-rotundatis, undique densē linearī-granatis; flavescente vel rosaceo-carneā, plus minus obscurē albo-flammata, apice atro-purpureo.

THE WELL-ENGRAVED ZIZYPHINUS. Shell conical, rather solid, whorls concave round the upper part, then tumidly rounded, closely linearly grained throughout, yellowish or rose-tinted flesh; more or less obscurely white-flamed, apex dark-purple.

ADAMS, Pro. Zool. Soc. 1854, p. 38.

Hab. Tasmania.

A solid, tumidly bulging shell, minutely linearly grained throughout, of a roseate-flesh hue, obscurely flamed with white, and possessing a peculiar dark-purple apex.

Species 18. (Mus. Cuming.)

ZIZYPHINUS CANALICULATUS. *Zizyph.* testā acutē latē conicā, ad basin concavo-planatā, anfractibus declivibus, undique annuloso-liratis, liris lăvibus, interstitiis conspicuē excavatis, præcipue ad suturas; olivaceo-luteā, in interstitiis saturatiore.

THE CHANNELLED ZIZYPHINUS. Shell sharply broadly conical, concavely flattened at the base, whorls sloping, annulosely ridged throughout, ridges smooth, interstices conspicuously excavated; especially at the sutures, olive-yellow, darker in the interstices.

Trochus canaliculatus, Martyn, Univ. Conch. vol. i. pl. 32.

Trochus doliarus, Chemnitz.

Trochus doliarus, Chemnitz.

Zizyphiinus canaliculatus, Gray.

Zizyphiinus doliarus, Adams.

Hab. Australia, New Zealand.

Both the sharply-defined ringed sculpture and the buff colouring of this species are constant.

ZIZYPHINUS.—PLATE III.

Species 19. (Mus. Cuming.)

ZIZYPHINUS ARMILLATUS. *Zizyph. testā tumido-conicā, tenuiculā, versus apicem acutā, anfractibus plano-convexis, undique grano-liratis, granis creberrimis, crenatis; flavescente, aureo-rufescente pallidē lentiginosā.*

THE JEWELLED ZIZYPHINUS. Shell tumidly conical, rather thin, sharp towards the apex, whorls flatly convex, strongly grain-ridged throughout, grains very close, crenated; yellowish, faintly freckled with golden-red.

Trochus armillatus, Wood, Ind. Test. Suppl. pl. 9. f. 5.

Hab. Tasmania.

An exquisitely jewelled shell, freckled with light golden-red, allied to the smaller Australian *Z. splendidus*.

Species 20. (Mus. Cuming.)

ZIZYPHINUS BICINGULATUS. *Zizyph. testā latē depresso-conoideā, anfractibus concavo-declivibus, deinide tumido-rotundatis, spiraliter liratis, liris duabus basaliibus valde prominentibus, omnibus densē plicato-crenatis; flavescente, aureo-rubo et albo profusè lentiginosā.*

THE TWO-GIRT ZIZYPHINUS. Shell broadly depressedly conoid, whorls concavely sloping round the upper part, then tumidly rounded, spirally ridged, two of the basal ridges very prominent, all densely plicately crenated; yellowish, profusely freckled with golden-red and white.

Trochus bicingulatus, Lamarek, Anim. sans vert. vol. ix.

p. 147.

Trochus vinctus, Philippi.

Hab. Raine's Island, Torres Straits; Ince.

Beautifully sculptured and freckled, the shell being conically depressed with two of the ridges towards the basal margin developed into conspicuous cords.

Species 21. (Fig. a, b, Mus. Cuming.)

ZIZYPHINUS ZONAMESTUS. *Zizyph. testā umbilicatā, acutē conicā, nunc depresso-conica, nunc pyramidali, ad basin acutangulā, anfractibus concavo-declivibus, minutē subobsoletē grano-liratis, interstitiis obliquē subtilissimē crenulatis; liris olivaceo-luteis, interstitiis ustulato-brunneis.*

THE NEATLY-ZONED ZIZYPHINUS. Shell unbilicated, conical, sometimes depressed, sometimes pyramidal, sharply angled at the base, whorls concavely sloping, minutely rather obsoletely grain-ridged; interstices obliquely very finely crenulated, ridges olive-yellow, interstices burnt-brown.

ADAMS, Pro. Zool. Soc. 1851, p. 166.

Hab. Honduras; Dyson.

In this species the ridges of fine granules are of a peculiar olive-yellow colour, while the interstices, minutely obliquely crenulated, are a rich burnt sienna.




22a



23



24



23b



25a



25b



26a



27



28



26b

ZIZYPHINUS.

PLATE IV.

Species 22. (Fig. *a, b*, Mus. Cuming.)

ZIZYPHINUS CONULUS. *Zizyph. testā conicā, ad basin lumen midiusculā, anfractibus convexo-declivibus, levigatis, nitentibus, ad apicem interdum granulatis, ad marginem basalem funiculatis; aureo-flavescente aut violaceā, rufa et albo nebulatā et articulata.*

THE LITTLE-CONE ZIZYPHINUS. Shell conical, rather tumid at the base, whorls convexly sloping, smooth, shining, sometimes granulated at the apex, corded at the basal margin; golden-yellow or violet, clouded and articulated with red and white.

Trochus conulus, Linnaeus, Syst. Nat. p. 1230.

Zizyphinus conulus, Adams.

Trochus violaceus, Risso.

Trochus Laugieri, Payraudeau.

Hab. British and Mediterranean Seas.

The specimen given at Fig. 22 *a* is Risso's *Trochus violaceus*.

Species 23. (Mus. Cuming.)

ZIZYPHINUS GEMMOSUS. *Zizyph. testā subtumido-conicā, solidā, anfractibus declivi-concavis, densè validè granatis, hic illic linearī-incisis, lineis obliquè minutè crenulatis, margine basali obtusè rotundato; aureo-stramineā, lineis incisis chocolato-purpureis.*

THE GEMMED ZIZYPHINUS. Shell rather tumidly conical, solid, whorls slopingly convex, densely strongly grained, here and there linearly engraved, lines obliquely minutely crenulated, basal margin obtusely rounded; golden straw-colour, engraved lines chocolate-purple.

Trochus gemmosus, Reeve, Pro. Zool. Soc. 1842, p. 184.
Conch. Syst. vol. ii. p. 165. pl. 218. f. 9.

Hab. Puerto Galero, island of Mindanao, Philippines (in sandy mud, at a depth of about six fathoms); Cuming.

Strongly grained upon a bright golden straw-colour, enlaced with engraved lines of a deep chocolate-purple.

Species 24. (Mus. Cuming.)

ZIZYPHINUS LURIDUS. *Zizyph. testā conicā, anfractibus plano-declivibus, ad basin tumidiusculis, levigatis vel malleatis, interdum spiraliter linearī-incisis; ustulato-fuscā, obscurissimè lurido-flammatā.*

THE LURID ZIZYPHINUS. Shell conical, whorls flatly sloping, rather tumid at the base, smooth or malleated, sometimes spirally linearly engraved; burnt-brown, very obscurely luridly flamed.

NUTTALL, MS. in Mus. Cuming.

Hab. California.

A smooth shell, spirally engraved towards the apex, of a peculiar burnt-brown colour, faintly obliquely flamed.

Species 25. (Fig. *a, b*, Mus. Cuming.)

ZIZYPHINUS EXIMIUS. *Zizyph. testā subdepresso-conicā, solidā, anfractibus supernè concavo-declivibus, deinude obtusè tumidis, spiraliter liratā, liris lævibus, ad marginem basalem angulari-prominentibus; albida, purpureo vel cœruleo nebulatā et articulatā.*

THE PRETTY ZIZYPHINUS. Shell somewhat depressly conical, solid, whorls concavely sloping round the upper part, then obtusely swollen, spirally ridged, ridges smooth, angularly prominent at the basal margin; whitish, clouded and articulated with purple or blue.

Trochus eximus, Reeve, Pro. Zool. Soc. 1842, p. 185.
Conch. Syst. vol. ii. p. 165. pl. 218. f. 12.

Hab. Panama (in sandy mud at ten fathoms); Cuming.

The painting of this species is mostly disposed in dots and articulated fillets.

Species 26. (Fig. *a, b*, Mus. Cuming.)

ZIZYPHINUS CHLOROSTOMA. *Zizyph. testā subpyramidalis-conicā, anfractibus concavo-declivibus, subtilissimè liratis, intersticiis minutè obliquè reticulatis, anfractuum margine basali compressè angulato-productis; purpureo aut virente, saturatè nebulatā et albo-maculatā, apice sanguineo.*

THE GREEN-MOUTH ZIZYPHINUS. Shell somewhat pyramidalis conical, whorls concavely sloping, very finely ridged, interstices minutely obliquely reticulated, basal margin of the whorls compressly angularly produced; purple or green, darkly clouded and white-spotted, apex blood-red.

Trochus chlorostoma, Menke, Moll. Nov. Holland, p. 17.
Zizyphinus chlorostoma, Adams.

Hab. West coast of New Holland; Menke. South Australia; Angas.

I have received specimens of this species from Mr. Augas, of Adelaide, South Australia, collected by him at Yorke's Peninsula.

Species 27. (Mus. Cuming.)

ZIZYPHINUS FILOSUS. *Zizyph. testā subobtusē conicā, solidā, anfractibus rotundatis, undique validē liratis, liris levibus, interstitiis excavatis, præcipue ad suturas; violaceo-fuscā, immaculatā, liris flarescentibus.*

THE CORDED ZIZYPHINUS. Shell rather obtusely conical, solid, whorls rounded, strongly ridged throughout, ridges smooth, interstices excavated, especially at the sutures; violet-brown, unspotted, ridges yellowish.

Trochus filosus, Wood, Ind. Test. Supp. pl. 5. f. 23.

Trochus ligatus, Gould.

Trochus castaneus, Nuttall.

Hab. Puget Sound; Gould.

A solid shell, regularly encircled throughout with smooth

ring-like ridges, excavated in the interstices. Wood gives no descriptive characters with this species, but he figures it with unmistakeable accuracy.

Species 28. (Mus. Cuming.)

ZIZYPHINUS DECORATUS. *Zizyph. testā subpyramidalis conicā, ad basin tumidiusculā, anfractibus concavo-declivibus, undique rugoso-granatis, granis irregularibus, interstitiis sub lente minutè obliquè striatis; albida, viridi aut purpureo-griseo profuse variegatā.*

THE EMBELLISHED ZIZYPHINUS. Shell somewhat pyramidal conical, rather swollen at the base, whorls concavely sloping, rugosely grained throughout, grains irregular, interstices beneath the lens obliquely minutely striated; whitish, profusely variegated with green or purple-grey.

Trochus decoratus, Philippi.

Hab. Brisbane waters, East Australia; Strange.

Of a more rugose-grained sculpture than usual, minutely obliquely striated in the interstices.



28.



29.



30.



31.



32a.



32b.



33.



34.



35.



36.

ZIZYPHINUS.

PLATE V.

Species 28. (Mus. Cuming.)

ZIZYPHINUS INCERTUS. *Zizyph. testá abbreviato-conicá, sinistrali, flavescente-albá, anfractibus declivi-convexis, spiraliter dense linearí-liratis, anfractu ultimo basi obtusè angulato.*

THE DOUBTFUL ZIZYPHINUS. Shell shortly conical, sinistral, yellowish white, whorls slopingly convex, spirally closely linearly ridged, last whorl obtusely angled at the base.

Hab. Tasmania.

This sinistrally coiled shell appears to me to be distinct from any other described species. Its normal growth is probably dextral.

Species 29. (Mus. Cuming.)

ZIZYPHINUS SCOBINATUS. *Zizyph. testá erecto-conicá, ad basin tumidiusculá, flavescente-albá, rufo obscurè flammatá et punctatá, anfractibus concavo-declivibus, undique dense papillari-granatis, anfractuum margine inferiori producto.*

THE FILED ZIZYPHINUS. Shell erectly conical, rather tumid at the base, yellowish white, obscurely flamed and dotted with red, whorls concavely sloping, densely papillary-grained throughout, lower margin of the whorls bulging.

ADAMS, MS. in Mus. Cuming.

Hab. Bombay.

The grains in this elegantly-formed shell are closely serially developed, and they are so large as to have a papillary appearance.

Species 30. (Mus. Cuming.)

ZIZYPHINUS MILLEGRANUS. *Zizyph. testá subaculè conicá, ad basin tumidiusculá, flavescente-albá, purpureo-roseo sparsim punctatá et articulatá, anfractibus plano-declivibus, undique oblique reticulatim striaturatis, anfractuum margine inferiori crenato-funiculato.*

THE THOUSAND-GRAINED ZIZYPHINUS. Shell rather sharply conical, slightly tumid at the base, yellowish white, sparsely dotted and articulated with purple-rose, whorls flatly sloping, obliquely reticulately fretted throughout, lower margin of the whorls crenately corded.

Trochus millegranus, Philippi, Enum. Moll. Sicil. vol. i.

p. 183. pl. 10. f. 25.

Trochus Clelandi, Wood.

Trochus Martini, Smith.

Trochus miliaris, Scacchi.

Hab. European Seas, from Norway to the Mediterranean.

The granular sculpture is in this species contorted, as it were, by oblique ridges into a netted fretwork. It is usually pale in colour, more or less delicately dotted and articulated with purple-rose.

Species 31. (Mus. Cuming.)

ZIZYPHINUS ANTONII. *Zizyph. testá sublatè conicá, solidá, argenteo-albá, atro-purpureo sparsim strigatá et punctatá, anfractibus concavo-declivibus, spiraliter linearí-granatis, granorum seriebus alternatim majoribus.*

ANTON'S ZIZYPHINUS. Shell rather broadly conical, solid, silvery white, sparsely streaked and dotted with dark purple, whorls concavely sloping, spirally linearly grained, rows of grains alternately larger.

Trochus Antonii, Koeh, Phil. Abbild. pl. i. f. 4.

Zizyphinus Antonii, Adams.

Hab. Australia.

Of a peculiar silvery-grey colour, sparingly dotted and streaked with dark purple.

Species 32. (Fig. a, b, Mus. Cuming.)

ZIZYPHINUS SIMILARIS. *Zizyph. testá subacuminato-conicá, flavescente, purpureo-roseo nebulatá et articulatá, anfractibus plano-declivibus, densè spiraliter grano-liratis, interstitiis eximiè reticulatis, anfractuum margine inferiori angulato-bifuniculato.*

THE SIMILAR ZIZYPHINUS. Shell rather acuminately conical, yellowish, clouded and articulated with purple rows, whorls flatly sloping, densely spirally grain-ridged, interstices faintly reticulated, lower margin of the whorls angularly double-corded.

Hab. Lizard Island, Torres Straits.

We have already noticed that the European *Z. conulus* appears in North-West Australia; the present species

is a very near representative, if not identical, of the European *Z. millegranus*.

Species 33. (Mus. Cuming.)

ZIZYPHINUS FLAVUS. *Zizyph.* testā latē depressē conicā, aurantio-flavā, rufescente pallidissimē punctatā, anfractibus plano-declivibus, rudē spiraliter liratis, liris lœvibus, circa apicem granulatis.

THE YELLOW ZIZYPHINUS. Shell broadly depressedly conical, orange yellow, very faintly dotted with red, whorls flatly sloping, rudely spirally ridged, ridges smooth, granulated round the apex.

Trochus flavus, Anton, Zeitschr. für Malak. 1848. p. 105.
Hab. Australia.

It is quite possible that this may be a somewhat obscure variety of a more characteristic species not yet discovered in a typical state.

Fig. 34. (Mus. Cuming.)

The shell here figured, set apart in Mr. Cuming's collection as a distinct species, is a rounded form of *Z. scobinatus* represented at Fig. 29. Both are from Bombay.

Species 35. (Mus. Cuming.)

ZIZYPHINUS INDISTINCTUS. *Zizyph.* testā angustē conicā, pyramidalī, olivaceo-flavā, purpureo-rufō tessellatū punctatā, anfractibus subconvexo-declivibus, spiraliter linearī-liratis, liris lœvibus, conferitusculis, intersticiis subtilissimē obliquē clathratis.

THE INDISTINCT ZIZYPHINUS. Shell narrowly conical, pyramidal, olive-yellow, tessellately dotted with purple-red, whorls somewhat convexly sloping, spirally linearly ridged, ridges smooth, rather close-set, interstices very finely obliquely latticed.

Trochus indistinctus, Wood, Ind. Test. Supp. pl. 6. f. 41.
Zizyphinus indistinctus, Adams.

Hab. New Holland.

An olive-yellow erectly conical shell, sprinkled in a tessellated manner with red dots.

Fig. 36. (Mus. Cuming.)

The shell here figured is a specimen of *Z. decoratus* in which the grains are more distinct and papillary, the interstices between them being still obliquely striated or wrinkled, whilst the ground-colour is a rose-lake, articulated with purple-black.



37



38



39



40



41a



41b



42



43.



44



45

ZIZYPHINUS.

PLATE VI.

Species 37. (Mus. Cuming.)

ZIZYPHINUS STRIATUS. *Zizyph. testā subpyramidalis-conicā, cinereo-griseā, cærneo-nigro longitudinaliter linearī-strigatā, anfractibus concavo-declivibus, spiraliter tenuiliratis, interstitiis et interdum liris obliquè striatis.*

THE STRIATED ZIZYPHINUS. Shell somewhat pyramidal conical, ash-grey, longitudinally linearly striped with blue-black, whorls concavely sloping, spirally finely ridged, interstices and sometimes the ridges obliquely striated.

Trochus striatus, Linnaeus, Syst. Nat. p. 1230.

Trochus erythrolencns and *punctulatus*, Gmelin.

Trochus parvus, Da Costa.

Trochus conicus, Donovan.

Trochus depictus, Deshayes.

Trochus Sartorii, Aradas and Maggiore.

Hab. South British and Mediterranean Seas.

The painting of this little species, though variable, is mostly disposed in dark, longitudinally waved, linear streaks.

Species 38. (Mus. Cuming.)

ZIZYPHINUS ELEGANTULUS. *Zizyph. testā acutē conicā, versus basin depresso tumido-rotundatā, lutescente, anfractibus concavo-declivibus, granoso-liratis, liris distantibus, granis etiam subdistantibus, interstitiis latē excavatis, præcipue ul suturas, liris violaceo-fuscis.*

THE ELEGANT ZIZYPHINUS. Shell sharply conical, depressedly tumidly rounded towards the base, yellowish, whorls concavely sloping, granosely ridged, ridges distant, grains also rather distant, interstices broadly excavated, especially at the sutures, ridges violet-brown.

ADAMS, Pro. Zool. Soc. 1851, p. 167.

Hab. Malacea (in coral sand from a depth of ten fathoms); Cuming.

The grains, which on the upper part of the whorls are rather distant from each other, gradually diminish in size, and approximate towards the base, where they amalgamate into smooth cords.

Species 39. (Mus. Cuming.)

ZIZYPHINUS MONILE. *Zizyph. testā erecto-conicā, ad basin*

tumidiusculā, pellucido-albā, maculis violaceis monili-formi-cinctā, anfractibus concavo-declivibus, spiraliter liratis, liris levibus, duabus basalibus prominentioribus.

THE NECKLACE ZIZYPHINUS. Shell erectly conical, rather swollen at the base, transparent white, encircled by a necklace of violet spots, whorls concavely sloping, spirally ridged, ridges smooth, the two basal ridges more prominent.

Hab. Port Curtis, East Australia.

The necklace of violet spots has an exceedingly pretty appearance on the delicate transparent ground of the shell.

Species 40. (Mus. Cuming.)

ZIZYPHINUS POLYCHROMA. *Zizyph. testā angustē pyramidalis-conicā, olivaceā vel brunnearā, strigis lineisque tenuibus albis longitudinaliter floccati, anfractibus concavo-planatis, spiraliter lineari-incisis, anfractuum margine basali bifuniculato.*

THE MANY-COLOURED ZIZYPHINUS. Shell narrowly pyramidal conical, olive or brown, longitudinally flaked with white streaks and fine lines, whorls concavely flattened, spirally linearly engraved, basal margin of the whorls double-corded.

ADAMS, Pro. Zool. Soc. 1851, p. 168.

Hab. Island of Masbate, Philippines (in sandy mud at a depth of about seven fathoms); Cuming.

There is no indication of any grained sculpture on this prettily-painted species.

Species 41. (Fig. a, b, Mus. Cuming.)

ZIZYPHINUS PYRAMIS. *Zizyph. testā acutē pyramidalis-conicā, varie colorata et maculata, anfractibus concavo-declivibus, spiraliter minutē grano-liratis, interstitiis interdum obliquè striatis, anfractuum margine basali conspicue tumido-producto.*

THE PYRAMID ZIZYPHINUS. Shell narrowly pyramidal conical, variously coloured and spotted, whorls concavely sloping, spirally minutely grain-ridged, interstices sometimes obliquely striated, basal margin of the whorls conspicuously tumidly bulging.

Trochus pyramis, Gmelin; Chemn. Conch. Cab. vol. v. pl. 170. f. 1652, 1653.

Hab. Australia.

In this narrow pyramidal shell, the lower margin of the whorls is more than usually tumidly produced. The painting is too variable to afford the means of definition.

Species 42. (Mus. Cuming.)

ZIZYPHINUS PULCHELLUS. *Zizyph.* *testā abbreviato-conicā,*
ad basin tumidā, carneo-albū, roseo profusè punctatā et
interdum strigatā, anfractibus concavo-declivibus, spirali-
ter liratis, liris laceribus, interstitiis spiraliter striatis,
anfractuum margine basali depresso angulato-producto.

THE PRETTY ZIZYPHINUS. Shell shortly conical, swollen at the base, flesh-white, profusely dotted, and sometimes streaked, with rose, whorls eoneavely sloping, spirally ridged, ridges smooth, interstices spirally striated, basal edge of the whorls depresso angularly produced.

Trochus pulchellus, Philippi, Zeitschr. für Malac. 1846,
p. 101.

Hab. —?

The sculpture in the interstices between the ridges runs parallel with the ridges themselves.

Species 43. (Mus. Cuming.)

ZIZYPHINUS TICAONICUS. *Zizyph.* *testā subabbreviato-*
conicā, versus basin subventricosā, pallidē carneo-roseā,
obscure punctato-articulatā, anfractibus convexis, spi-
raliter tenuiliratis, liris levibus, alternatim minori-
būs.

THE TICAO ZIZYPHINUS. Shell rather shortly conical, rather ventricose towards the base, pale flesh-rose, faintly dot-articulated, whorls convex, spirally finely ridged, ridges smooth, alternately smaller.

ADAMS, Pro. Zool. Soc. 1851, p. 167.

Hab. Island of Ticao, Philippines (in sandy mud at a depth of about six fathoms); Cuming.

Remarkable only for the simplicity of its characters.

Species 44. (Mus. Cuming.)

ZIZYPHINUS SCITULUS. *Zizyph.* *testā umbilicatā, tumido-*
conicā, tenuiculā, aurantio-fulvā, punctis sanguineis
minutè articulatā, hic illic albo-floccatā, anfractibus
convexo-declivibus, spiraliter striato-liratis, lirā juxta
suturas majore.

THE NEAT ZIZYPHINUS. Shell umbilicated, tumidly conical, rather thin, orange fulvous, minutely articulated with crimson dots, and here and there flaked with white, whorls convexly sloping, spirally striately ridged, ridge next the suture larger.

ADAMS, Pro. Zool. Soc. 1854, p. 38.

Hab. New Zealand; Strange.

The painting of this species as above described is characteristic.

Species 45. (Mus. Cuming.)

ZIZYPHINUS LEANUS. *Zizyph.* *testā subacutē conicā, li-*
vido-ærueā, fusco obscure flammatū, anfractibus supernè
concavo-declivibus, deinde tumido-productis, undique
validē spiraliter granulis.

LEA'S ZIZYPHINUS. Shell rather sharply conical, livid bronze, obscurely flamed with brown, whorls eoneavely sloping round the upper part, then tumidly bulging, strongly spirally grained throughout.

Trochus Leanus, C. A. Adams. *fig. 2*

Hab. —?

A strongly-grained shell of a peculiar livid bronze hue, faintly brown-streaked.



46



47



48



49



50



51.



52



53.



54.



55

ZIZYPHINUS.

PLATE VII.

Species 46. (Mus. Cuming.)

ZIZYPHINUS ALABASTRUM. *Zizyph. testá sublatè conicá, tenui, sordidè albá, anfractibus convexis, spiraliter carinatis, carinis læribus, circa apicem granulatis, distantibus, intersticiis latè excavatis, præcipue ad suturas, anfractu ultimo tumido-producto.*

THE ALABASTER ZIZYPHINUS. Shell rather broadly conical, thin, dull white, whorls convex, spirally keeled, keels smooth, granulated round the apex, distant, interstices broadly excavated, especially at the sutures, last whorl tumidly bulging.

Margarita alabastrum, Beek, Lovén, Index Moll. Scan. p. 20.

Trochus formosus, Forbes.

Trochus alabastrum, Forbes and Hanley.

Zizyphinus alabastrum, Adams.

Hab. North European Seas.

Some doubt exists as to whether this interesting prominently-keeled *Zizyphinus*, which is a boreal European species, is not the same as a North American species described by Michels and Adams in the 'Journal of the Natural History Society of Boston' for 1841, with the name *Trochus occidentalis*.

Species 47. (Mus. Cuming.)

ZIZYPHINUS FRAGUM. *Zizyph. testá conicá, sordidè albá, fuscescente pallidè tessellatá, anfractibus convexo-declivibus, spiraliter densè granatis, granis amplis, papillaribus.*

THE STRAWBERRY ZIZYPHINUS. Shell conical, dull white, faintly tessellated with brown, whorls convexly sloping, spirally closely grained, grains large, papillary.

Trochus fragum, Philippi, Zeitschr. für Malac. 1848. p. 106.

Hab. Australia.

A large-grained shell, faintly tessellated with violet-brown.

Species 48. (Mus. Cuming.)

ZIZYPHINUS COMPTUS. *Zizyph. testá erecto-conicá, sordidè albá, ad marginem basalem fusco sparsim punctatá, anfractibus declivi-convexis, ad suturas impressis, undique seriatim granulatis.*

THE NEAT ZIZYPHINUS. Shell erectly conical, dull white sparsely dotted with red-brown at the basal margin, whorls slopingly convex, impressed at the sutures, encircled throughout with rows of granules.

ADAMS, Pro. Zool. Soc. 1854, p. 38.

Hab. New Caledonia.

A rather solid, dull, semitransparent shell, with indications of brown dots about the basal margin.

Species 49. (Mus. Cuming.)

ZIZYPHINUS JAPONICUS. *Zizyph. testá elevato-conicá, purpureá et rufá, flammis albidis variegatá vel reticulatá, basi punctatá, anfractibus plano-declivibus, læribus, prope suturas linearis-incisis.*

THE JAPAN ZIZYPHINUS. Shell elevately conical, purple and red, variegated or reticulated with white flames, base dotted, whorls flatly sloping, smooth, linearly engraved next the suture.

ADAMS, Pro. Zool. Soc. 1851. p. 167.

Hab. Japan.

Variously reticulated and articulated with white, of the same smooth richly-painted type as *Z. polychroma*.

Species 50. (Fig. 50, 51, Mus. Cuming.)

ZIZYPHINUS MAC-ANDREÆ. *Zizyph. testá conicá, solidiusculá, albida, ustulato-purpureo tessellato-strigata, anfractibus supernè concavo-impressis, deinde tumido-rotundatis, undique spiraliter grano-liratis.*

MAC-ANDREW'S ZIZYPHINUS. Shell conical, rather solid, white, tessellately streaked with burnt-purple, whorls concavely impressed round the upper part, then tumidly rounded, spirally grain-ridged throughout.

CARPENTER, Cat. Reigen Coll. p. 232.

Hab. Mazatlan, Panama.

Tessellately streaked in a characteristic manner with burnt-purple or blood-colour.

Species 52. (Mus. Cuming.)

ZIZYPHINUS FLAMMIGER. *Zizyph. testá acutè elevato-*

conicā, fulvo-rufescēte, albo flammatā, anfractibus convexo-declivibus, lāevis.

THE FLAMED ZIZYPHINUS. Shell sharply elevately conical, fulvous red, flamed with white, whorls convexly sloping, smooth.

DUNKER, MS. in Mus. Cuming.

Hab. —?

A perfectly smooth shell, flamed with white, upon a peculiar fulvous-red ground.

Species 53. (Mus. Cuming.)

ZIZYPHINUS PICTURATUS. *Zizyph. testā angustē erecto-conicā, vividē virente, albo flammatā, purpureo-rufō variè tinctā et punctatā, anfractibus plano-declivibus, linearī-incisio, ad marginem basalem funiculatis.*

THE EMBROIDERED ZIZYPHINUS. Shell narrowly erectly conical, bright green, flamed with white, variously stained and dotted with purple-red, whorls flatly sloping, linearly engraved, corded round the basal margin.

ADAMS, Pro. Zool. Soc. 1851, p. 168.

Hab. Island of Negros, Philippines (in sandy mud at the depth of seven fathoms); Cuming.

Exquisitely painted in the most variable manner, with green and purple-red, the former colour predominating.

Species 54. (Mus. Cuming.)

ZIZYPHINUS CRENULATUS. *Zizyph. testā subacutē conicā, albida, roseo tinctā et articulatā, aut variè coloratā, anfractibus plano-declivibus, minulē grano-liratis, intersticiis subtilissimè crenulatis, anfractuum margine basali conspicuè tumido-funiculato.*

THE CRENULATED ZIZYPHINUS. Shell rather sharply conical, whitish, stained and articulated with rose, or variously coloured, whorls flatly sloping, minutely grain-ridged, interstices very finely crenulated, basal margin of the whorls conspicuously tumidly corded.

Trochus crenulatus, Brocchi, Conch. Foss. Subapp. p. 354. pl. 6. f. 2.

Trochus punctatus, Renier.

Trochus punctatus, Dillwyn.

Trochus minimus, Chemnitz.

Trochus pyramidatus, Lamarck.

Trochus Matoni, Payraudeau.

Hab. Mediterranean.

This pretty species, allied to *Z. millegranus*, is encircled with a stout tumid cord, on which the colour is conspicuously articulated throughout.

Species 55. (Mus. Cuming.)

ZIZYPHINUS DUPLICATUS. *Zizyph. testā elevato-conicā, sordidē albā, anfractibus ad suturas impressis, deinde tumido-rotundatis, undique densè granatis.*

THE DUPLICATE ZIZYPHINUS. Shell elevately conical, dull white, whorls impressed at the sutures, then tumidly rounded, closely grained throughout.

ADAMS, Pro. Zool. Soc. 1851, p. 168.

Hab. —?

This is a very unsatisfactory species. Although the sculpture is well defined, the shell is dead and colourless, and nothing is known of its habitat.

Zyphinus Pl. VIII.



56.



57.



58.



59.



60.



61.



62.



63.



64.



65.

ZIZYPHINUS.

PLATE VIII.

Species 56. (Mus. Cuming.)

ZIZYPHINUS RUBRO-PUNCTATUS. *Zizyph. testá abbreviato-conicá, lutescente, anfractibus convexis, ad suturas impressis, liris squamato-spinulosis cinctis, interstitiis profundis, minutis, rubris.*

THE RED-DOTTED ZIZYPHINUS. Shell shortly conical, yellowish, whorls convex, impressed at the sutures, encircled with squamately spined ridges, interstices deep, minute, red.

ADAMS, Pro. Zool. Soc. 1851, p. 167.

Hab. —?

The sculpture of this little species is developed throughout in erect prickly scales, the minute interstices between which are red.

Species 57. (Mus. Cuming.)

ZIZYPHINUS MONTAGUI. *Zizyph. testá conicá, flavescente-albá, fuscescente punctatá, anfractibus convexis, spiraliiter linearí-liratis, liris lœvibus, interstitiis latè profundè excavatis, minutissimè longitudinaliter striatis.*

MONTAGU'S ZIZYPHINUS. Shell conical, yellowish-white, dotted with pale-brown, whorls convex, spirally linearly ridged, ridges smooth, interstices broadly deeply excavated, very minutely longitudinally striated.

Trochus Montagui, Gray, Wood, Ind. Test. Suppl. pl. 6. f. 43.

Trochus striatus, Forbes.

Zizyphinus striatus, Adams.

Hab. British Seas. Vigo Bay, Galicia; M'Andrew.

Encircled with narrow smooth ridges, the interstices between which are neatly excavated.

Species 58. (Mus. Cuming.)

ZIZYPHINUS UNICINCTUS. *Zizyph. testá subfusciforme-conicá, flavescente-albá, purpureo-rufo sparsim articulatá, anfractibus spiraliter tenuistriatis, supernè declivibus, rudè plicatis, medio angulato-tumidis, ad angulum nodatis, anfractu ultimo versus basin attenuato, unifuniculato.*

THE ONE-GIRT ZIZYPHINUS. Shell somewhat fusiformly conical, yellowish-white, sparingly articulated with

purple-red, whorls spirally finely striated, sloping and rudely plicated round the upper part, angularly swollen in the middle, noduled at the angle, last whorl attenuated towards the base, one-corded.

ADAMS, Pro. Zool. Soc. 1851, p. 167.

Hab. Lord Hood's Island (on pearl oysters, at a depth of from eight to ten fathoms); Cuming.

A curiously formed species, of which the character is well given in the accompanying figure.

Species 59. (Mus. Cuming.)

ZIZYPHINUS VEXILLUM. *Zizyph. testá conicá, albidú, viridi latistrigatá, anfractibus ruditè convexis, lœvibus, spiraliter linearí-incisis.*

THE FLAG ZIZYPHINUS. Shell conical, white, broadly striped with green, whorls rudely convex, smooth, spirally linearly engraved.

Hab. Australia; Strange.

A shell of simple form, with no sculpture excepting faintly engraved spiral lines.

Species 60. (Mus. Cuming.)

ZIZYPHINUS CALICHROUS. *Zizyph. testá subabbreviato-conicá, versus basin latè angulato-ventricosá, purpureo-roseá, lineis nigris distantibus cinctá, fusco oblique flammatá et articulatá, anfractibus plano-convexis, spiraliter linearí-incisis, circa apicem obscurè granulatis.*

THE PRETTY-COLOURED ZIZYPHINUS. Shell rather shortly conical, broadly angularly ventricose towards the base, purple-rose, encircled with distant black lines, obliquely flamed and articulated with brown, whorls flatly convex, spirally linearly engraved, obscurely granulated round the apex.

Trochus callichrous, Philippi, Zeitschr. für Malak. 1849, p. 149.

Zizyphinus strigosus, Adams.

Hab. Morocco.

The evidence of this species being *Trochus strigosus* of Gmelin is insufficient; and the figure in Chennitz, to

which Mr. Adams refers in illustration of it, represents a *Z. crenulatus*.

Species 61. (Mus. Cuming.)

ZIZYPHINUS OCELLATUS. *Zizyph. testā sublatē conicā, rufescētē-fūlvā, maculis albīs fuscō-umbratīs ocellatā, anfractibūs supērē concavō-impressis, deinde rotundatis, undique spiraliter grano-liratis.*

THE EYED ZIZYPHINUS. Shell rather broadly conical, reddish fulvous, ocellated with brown-shaded white spots, whorls concavely impressed round the upper part, then rounded, spirally grain-ridged throughout.

Hab. —?

Rather constrictedly impressed below the sutures, then rounded and ocellated with shaded opake-white spots.

Species 62. (Mus. Cuming.)

ZIZYPHINUS EXIGUUS. *Zizyph. testā subpyramidalē-conicā, cinereo-fuscescente, obliquē nebulatā, apice intensē roseo, anfractibūs concavō-declivibus, spiraliter obtuse liratis, interstītiis crenulato-striatis, anfractuum marginē basali tumido-funiculato.*

THE LITTLE ZIZYPHINUS. Shell somewhat pyramidal conical, ash-brown, clouded obliquely, apex deep-rose, whorls concavely sloping, spirally obtusely ridged, interstices crenulately striated, basal margin of the whorls tumidly corded.

Trochus exiguis, Pulteney, Hutchins' Hist. Dorset, p. 44.

Trochus conulus, Da Costa.

Trochus minutus, Dillwyn.

Trochus erythrolencus, Lamarck.

Trochus interruptus, Wood.

Zizyphinus exiguis, Adams.

Hab. South Britain and Mediterranean shores.

Though variable in colour this species is mostly of a light tawny or ash-brown, the apex being always a deep-rose.

Species 63. (Mus. Cuming.)

ZIZYPHINUS ZEBUENSIS. *Zizyph. testā obeso-conicā, albidā, pallidē cærulescētē-viridi variegatā, anfractibus rotundatis, undique confertim tenuiliratis, infra suturas minutē crenulatis, liris lœvibus, subacutis.*

THE ZEBU ZIZYPHINUS. Shell stoutly conical, whitish, variegated with pale blue-green, whorls rounded, closely finely ridged throughout, minutely crenulated below the sutures, ridges smooth, rather sharp.

ADAMS, MS. in Mus. Cuming.

Hab. Island of Zebu, Philippines; Cuming.

The ridges of this little species are unusually fine and sharp, with no indications of granules.

Species 64. (Mus. Cuming.)

ZIZYPHINUS INTERRUPTUS. *Zizyph. testā erecto-conicā, griseo-albidā, lineis tenuibus cæruleo-nigricantibus interruptis cinctā, ad suturas minutē articulatā, anfractibus plano-declivibus, spiraliter tenuiliratis, interstītiis minutē crenulato-striatis.*

THE INTERRUPTED ZIZYPHINUS. Shell erectly conical, greyish-white, encircled with fine blue-black interrupted lines, minutely articulated at the sutures, whorls flatly sloping, spirally finely ridged, interstices minutely crenulately striated.

Trochus interruptus, Wood, Ind. Test. Suppl. pl. vi. f. 42.
Zizyphinus interruptus, Adams.

Hab. —?

A very distinct erectly-conical shell, painted in an elaborately tessellated manner.

Species 65. (Mus. Cuming.)

ZIZYPHINUS DECUSSATUS. *Zizyph. testā subacutē conicā, ad basin tumidiusculā, albidā, vividē viridi maculatā et variegatā, anfractibus plano-declivibus, spiraliter tenuē grano-liratis, ad marginem basalem decussatis et squamatis, suturis angustē impressis.*

THE DECUSSATED ZIZYPHINUS. Shell rather sharply conical, a little swollen at the base, white, spotted and variegated with bright-green, whorls flatly sloping, spirally finely grain-ridged, decussated and crenulately scaled at the basal margin, sutures narrowly impressed.

ADAMS, Pro. Zool. Soc. 1851, p. 167.

Hab. Island of Mindoro, Philippines (in coarse gravel, at the depth of twelve fathoms); Cuming.


A very elegant little species, in which the granules at the basal margin of each whorl and on the under surface of the shell are transformed into scales.

Z I Z Y P H I N U S.

Plate,	Species.	Plate	Species.
<i>ngrestis</i> (<i>Trochus</i>), Philippi	III. 16	<i>luridus</i> , <i>Nuttall</i>	IV. 24
<i>alabastrum</i> (<i>Margarita</i>), Beck	VII. 46	<i>Lyonsii</i> (<i>Trochus</i>), Leach	III. 16
<i>albus</i> (<i>Trochus</i>), Wood	III. 16	<i>MacAndreae</i> , <i>Carpenter</i>	VII. 50
<i>annulatus</i> (<i>Trochus</i>), Martyn	II. 13	<i>Martini</i> (<i>Trochus</i>), Smith	V. 30
<i>Autonii</i> (<i>Trochus</i>), Koch	V. 31	<i>Matoni</i> (<i>Trochus</i>) <i>Payraudeau</i>	VII. 54
<i>armillatus</i> (<i>Trochus</i>), Wood	III. 19	<i>miliaris</i> (<i>Trochus</i>), <i>Seacchi</i>	V. 30
<i>articulatus</i> , <i>Reeve</i>	I. 3	<i>millegranus</i> (<i>Trochus</i>), Philippi	V. 30
<i>bicinctulus</i> (<i>Trochus</i>), Lamarck	III. 20	<i>minimus</i> (<i>Trochus</i>), Chemnitz	VII. 54
<i>callichrous</i> (<i>Trochus</i>), Philippi	VIII. 60	<i>minutus</i> (<i>Trochus</i>), Dillwyn	VIII. 62
<i>canaliculatus</i> (<i>Trochus</i>), Martyn	III. 18	<i>monile</i> , <i>Reeve</i>	VI. 39
<i>vastaneus</i> (<i>Trochus</i>), Gould	IV. 27	<i>Montagui</i> (<i>Trochus</i>), Gray	VIII. 57
<i>castra</i> , <i>Reeve</i>	II. 14	<i>nebulosus</i> , Adams	II. 7
<i>chlorostoma</i> (<i>Trochus</i>), Menke	IV. 26	<i>nobilis</i> (<i>Trochus</i>), Philippi	II. 10
<i>ciliaris</i> (<i>Trochus</i>), Menke	III. 16	<i>ocellatus</i> , <i>Reeve</i>	VIII. 61
<i>Oleandi</i> (<i>Trochus</i>), Wood	V. 30	<i>ornatus</i> (<i>Trochus</i>), Lamarck	II. 7
<i>comptus</i> , Adams	VII. 48	<i>parbus</i> (<i>Trochus</i>), Da Costa	VI. 37
<i>conicus</i> , (<i>Trochus</i>), Donovan	VI. 37	<i>picturatus</i> , Adams	VII. 53
<i>conuloides</i> (<i>Trochus</i>), Lainarek	III. 16	<i>polychroma</i> , Adams	VI. 40
<i>conulus</i> (<i>Trochus</i>), Linnaeus	IV. 22	<i>pulchellus</i> (<i>Trochus</i>), Philippi	VI. 42
<i>conulus</i> (<i>Trochus</i>), Da Costa	VIII. 62	<i>punctatus</i> (<i>Trochus</i>) Renieri	VII. 54
<i>crenulatus</i> (<i>Trochus</i>), Brocchi	VII. 54	<i>punctulatus</i> (<i>Trochus</i>), Dillwyn	VII. 54
<i>Cunninghamii</i> , Gray	I. 6	<i>punctulatus</i> (<i>Trochus</i>), Gmelin	VI. 37
<i>decoratus</i> (<i>Trochus</i>), Philippi	IV. 28	<i>pyramidalis</i> (<i>Trochus</i>), Lamarck	VII. 54
<i>decussatus</i> , Adams	VIII. 65	<i>pyramis</i> (<i>Trochus</i>), Gmelin	VI. 41
<i>depietus</i> (<i>Trochus</i>), Deshayes	VI. 37	<i>rubro-punctatus</i> , Adams	VIII. 56
<i>discrepans</i> (<i>Trochus</i>), Brown	III. 16	<i>Sartorii</i> (<i>Trochus</i>), Aradas	VI. 37
<i>doliarius</i> (<i>Trochus</i>), Chemnitz	III. 18	<i>scitulus</i> , Adams	VI. 44
<i>duplicatus</i> , Adams	VII. 55	<i>scobinatus</i> , Adams	V. 29
<i>eleganthulus</i> , Adams	VI. 38	<i>selectus</i> (<i>Trochus</i>), Chemnitz	I. 1
<i>erythroleucus</i> (<i>Trochus</i>), Gmelin	VI. 37	<i>similaris</i> , <i>Reeve</i>	V. 32
<i>erythroleucus</i> (<i>Trochus</i>), Lamarck	VIII. 62	<i>speciosus</i> , Adams	II. 9
<i>englyptus</i> , Adams	III. 17	<i>spectabilis</i> , Adams	I. 5
<i>exiguus</i> (<i>Trochus</i>), Pulteney	VIII. 62	<i>splendidus</i> , Philippi	II. 11
<i>eximius</i> , <i>Reeve</i>	IV. 25	<i>striatus</i> (<i>Trochus</i>), Forbes	VIII. 57
<i>filosus</i> (<i>Trochus</i>), Wood	IV. 27	<i>striatus</i> (<i>Trochus</i>), Linnaeus	VI. 37
<i>flammeiger</i> , Dunker	VII. 52	<i>strigosus</i> , Adams	VIII. 60
<i>flavus</i> (<i>Trochus</i>), Anton	V. 33	<i>Ticaonicus</i> , Adams	VI. 43
<i>formosus</i> (<i>Trochus</i>), Forbes	VII. 46	<i>tigris</i> (<i>Trochus</i>), Martyn	I. 4
<i>fragum</i> (<i>Trochus</i>), Philippi	VII. 47	<i>Tranquebaricus</i> (<i>Troch.</i>), Chemnitz	III. 15
<i>gemmosus</i> , <i>Reeve</i>	IV. 23	<i>unicinetus</i> , Adams	VIII. 58
<i>granatus</i> (<i>Trochus</i>), Chemnitz	I. 2	<i>unicus</i> , Dunker	II. 8
<i>incertus</i> , <i>Reeve</i>	V. 28	<i>vexillum</i> , Adams	VIII. 59
<i>indistinctus</i> (<i>Trochus</i>), Wood	V. 35	<i>vinetus</i> (<i>Trochus</i>), Philippi	III. 20
<i>interruptus</i> (<i>Trochus</i>), Wood	VIII. 64	<i>virgineus</i> (<i>Trochus</i>), Chemnitz	II. 13
<i>Japonicus</i> , Adams	VII. 49	<i>vigilaceus</i> (<i>Trochus</i>), Risso	IV. 22
<i>jujubinus</i> (<i>Trochus</i>), Gmelin	II. 12	<i>Zebuensis</i> , Adams	VIII. 63
<i>Langieri</i> (<i>Trochus</i>), Payraudeau	IV. 22	<i>zezypinus</i> (<i>Trochus</i>), Chemnitz	II. 12
<i>Leamus</i> (<i>Trochus</i>), C. B. Adams	VI. 45	<i>zizyphinus</i> (<i>Trochus</i>), Lamarck	III. 16
<i>ligatus</i> (<i>Trochus</i>), Gould	IV. 27	<i>zouanestus</i> , Adams	III. 21

M O N O G R A P H
OF THE GENUS
T E R E B E L L U M.

"And I have felt
A presence that disturbs me with the joy
Of elevated thoughts ; a sense sublime
Of something far more deeply interfused,
Whose dwelling is the light of setting suns,
And the round ocean."—*Wordsworth*.



TEREBELLUM.

PLATE I.

Genus TEREBELLUM, Lamarek.

Testa cylindraceo-subulata, angustè leviter convoluta, spirâ prominente, subobtusa, suturâ linearis-incisâ, supernâ encausticâ, aperturâ longitudinali, supernâ tenuâ canaliculatâ, ad basin abrupte decurtatâ et truncatâ, columellâ subrectâ, encausticâ, in mucronem desinente.

Shell cylindrically subulate, narrowly lightly convoluted, spire prominent, rather obtuse, suture linearly cut, enamelled along the upper margin, aperture longitudinal, thinly channelled at the upper part, abruptly cut short and truncated at the base, columella nearly straight, enamelled, ending in a point.

In 1846, when publishing the third Number of my 'Elements of Conchology,' I took occasion to notice that although the *Terebellum* shell had been known for a century and more, and is not unfrequently collected on the shores of the Eastern Seas, no specimen had yet been found with its molluscous inhabitant. Mr. Cuming, however, remembered seeing living specimens, seven years before, at the Philippine Islands, having remarked its activity of habit in leaping from his hand. In 1845 the animal of *Terebellum* was well observed by Mr. Arthur Adams during the voyage of the 'Samarang' in the China Sea and among the islands of the Sooloo Archipelago, and an excellent drawing and description of it was published in 1850 in the Zoology of that expedition. Agreeably with the conjectures of Sowerby and De Blainville, the soft parts of *Terebellum* were found to resemble those of *Strombus*, and, as in *Oliva*, which De Blainville placed after it in the series, the mantle is provided with a long filamentary cord winding into the sutures of the shell.

"The *Terebellum*," says Mr. Adams, "is extremely shy in its movements. Poising its shell in a vertical position,

and cautiously protruding its long telescope-eye from the truncature in front of the shell, it will remain stationary until assured of security. It will then use its pointed foot as a lever, and roll its shell over and over, progressing by a series of irregular leaps. When removed from the water, before dying, it will jump several inches from the ground."

Species 1. (Fig. *a* to *g*, Mus. Cuming.)

TEREBELLUM PUNCTATUM. *Ter. testâ cylindraceo-subulatâ, levigatâ, nitente, spirâ conspicuâ, suturâ linearis-incisâ, encausticâ, fulvescente-albâ, spadiceo-fusco varie lentiginosâ vel undatâ, interdum castaneo-punctatâ vel acutâ undatâ.*

THE DOTTED TEREBELLUM. Shell cylindrically subulate, smooth, shining, spire conspicuous, suture linearly cut, enamelled, fulvous-white, variously freckled or waved with fawn-brown, sometimes dotted or sharply waved with chestnut.

Bulla terebellum, Linnaeus, Syst. Nat. p. 1185.

Conus terebellum, Linnaeus.

Terebellum punctatum, pullum, and album, Klein.

Strombus terebellum, Rumphius.

Terebra alata, Argenville.

Terebellum punctatum, Chemnitz.

Terebellum subulatum, Lamarek.

Hab. China Sea, and Indian and Philippine Archipelagoes.

All the specimens of *Terebellum* hitherto collected appear to be referable to one species. It will be seen by the varieties assembled in the accompanying plate, that the colouring is fawn or chestnut-brown upon a fulvous ground, disposed in freckles, dots, or zigzag lines.

MONOGRAPH
OF THE GENUS
P A L U D I N A.

"How green you are, and fresh in this old world!"—*Shakespeare*.



2c




1



2b



3a



4a



4b



3b

PALUDINA.

PLATE I.

Genus PALUDINA, Lamarck.

Testa ovato-conica, plerumque umbilicata, interdum umbilico fere clauso, anfractibus quinque ad ser, levigatis aut spiraliter fuscinalatis, epidermide fusco vel olivaceo-virente, interdum ciliata, indutis; aperturâ sepiissimè pyriformi-rotundata, labro simplici. Operculum cornuum, concentricum, nucleo sublaterali.

Shell ovately conical, generally umbilicated, sometimes with the umbilicus almost closed, whorls five to six, smooth or spirally corded, covered with a brown or olive-green epidermis, which is sometimes ciliated; aperture most frequently pyriformly rounded, lip simple. Operculum horny, concentric, with the nucleus sublateral.

In studying the genus *Paludina* it is necessary to give especial attention to the animal. Dr. Gray noticed, as long back as 1821, in a paper in the 'London Medical Repository,' that two out of the four mollusks of our own streams hitherto regarded as *Paludinæ*, were distinguished by characters peculiarly their own, and the distinction has now been established in a very numerous series of foreign species. *Bithynia*, the genus proposed for their reception, is more numerous in species than *Paludina*. The animal differs, as we shall presently see, in being oviparous, in having the eyes sessile, the mantle comparatively simple, and the tentacles slender and thread-like, while the shells are uniformly smaller.

The *Paludinæ* are ovo-viviparous, the eggs being hatched in the ovary, whence, at the end of about two months, the young are ejected, three or four at a time, alive. The animal has a largely dilated foot, broad in front and attenuately rounded behind; the head is produced into a proboscis, and the tentacles, unlike those of *Bithynia*, are rather stout and cylindrical, with the eyes raised on short conjoined stalks. This modification in the position of the eyes, is apparently designed to make room for an organ at the outer base of the right tentacle, used for conveying water to the branchial chamber. It is in the form of a tubular lobe, and on the left side of the neck there is a corresponding lobe, to which no particular use is assigned. In the large *Ampullariæ* of the tropics, affecting situations where the water is more liable to be dried up, the animal is provided with a double system of respiration, having, in addition to the branchial chamber on the right side, a pulmonary chamber on the left side. The dormant lobe of *Paludina* is then developed into an elongated siphonic

tube, for the passage to this chamber of the air. In most of the marine water-breathing Cephalis, the siphonic lobes, which appear in such a rudimentary form in *Paludina*, are combined into a conspicuous central tube adapted to the same use, and the shell is either notched in front for its reception, as in *Buccinum*, or extended into a long canal for its special protection, as in *Murex*.

The general distribution of *Paludina* over the globe, does not range with that of the inoperculated freshwater genera *Planorbis*, *Lymnaea*, or *Physa*. We have no record of its existence in South Australia, New Zealand, or Polynesia; nor does it appear in the West Indies or South America. About seventy species are known, of which rather more than two-thirds belong to the Eastern Hemisphere, and the remainder to North America. India and China, the chief area of habitation of this genus in the Old World, contribute about thirteen species, embracing several very characteristic varieties of form, sculpture, and painting. In some, the shells are conspicuously keeled, in others they are prettily banded, but green or olive-green is in all the predominant ground-colour. The Philippine Island rivers contribute only half-a-dozen comparatively insignificant species of *Paludina*, their place being occupied to a great extent by *Melania*, remarkable for the size and exquisitely acuminate proportions of their shells; and in Ceylon the rapids, chiefly those flowing from Adam's Peak, are inhabited by a very characteristic group having peculiarly globose shells, *Tanalia* and *Paludomus*.

The remainder of the Old World *Paludinæ*, so far as they are at present known, are six from Borneo, Java, Celebes, six from Burmah and Siam (in which latter country the land and freshwater mollusks all present strikingly new specific forms), and nine from more northern latitudes, of which two are European, two Siberian, and five Japanese. Temperature, which has so marked an influence in the development of mollusean life among the land snails, as may be seen by comparing the West African and Philippine *Helices* and *Bulimi* with the European, or those of Bolivia and Venezuela with the North American, has comparatively little influence among the water snails. We have no such large *Physæ* in Britain as inhabit Australia and the West Indies, but in *Lymnaea stagnalis*, which ranges eastward and northward to Siberia, we have the largest species known of that genus; and we have also in Britain the largest known *Planorbis*. It is the same with *Paludina* in Russian Asia. The largest variety of the *P. gigantea* of Bengal is exceeded in size by *P. Ussuriensis*, a

species collected by Gerstfeldt at the mouth of the Ussuri, a tributary of the Amoor, in Siberia, which is in the isothermal latitude of Iceland, little south of the line of permanent ground frost. The European *Paludina vivipara*, at least a light inflated form of similar structure and colour referred to that species, appears in North America, but the *Paludinae* of that continent consist of fifteen species, chiefly of a peculiarly solid type, forming the genus *Melanthon* of Bowditch.

Species 1. (Mus. Cuming.)

PALUDINA DOLIARIS. *Pal. testā ovatā, ventricosā, ferrugineo-olivaceā, liris nigricantibus funiculatā, anfractibus supernē oblongis angulatis, deinde modicē converxis; aperturā pyriformi-rotundatā.*

THE TONGUE PALUDINA. Shell ovate, ventricose, rusty-olive, corded with blackish ridges, whorls obtusely angled round the upper part, then moderately convex; aperture pyriformly rounded.

GOULD, Otia Conch. p. 191.

Hab. Burmah.

A bold rust-olive shell, corded throughout with dark raised ridges and lines.

Species 2. (Fig. *a, b*, Mus. Cuming.)

PALUDINA ANGULATA. *Pal. testā subfusciformi-ovatā, fuscā, virescente-olivaceo fasciatā, spirā elevatā, subpyramidalē, suturis angustē canaliculatis, anfractibus supernē concavo-declivibus, ad peripheriam tuberculis uni-biseriatim cinctis; aperturā trapezoideā.*

THE ANGLED PALUDINA. Shell somewhat fusiformly ovate, fuscous, banded with greenish-olive, spire raised, somewhat pyramidal, with the sutures narrowly channelled, whorls concavely sloping at the upper part, encircled at the periphery with a row of tubercles, sometimes with two rows; aperture trapezoid.

LEA, Trans. Amer. Phil. Soc. vol. ix. p. 22.

Vivipara magnifica (young), Binney.

Hab. Coosa River, Alabama.

Mr. Binney considers this species to have been founded on the young of *P. magnifica*.

Species 3. (Fig. *a, b*, Mus. Cuming.)

PALUDINA CONTECTA. *Pal. testā scalariformi-turbinatā, profundē umbilicatā, intensē olivaceo-virente, fasciis tribus purpureo-rufis cingulatā, anfractibus subangustis, ad suturas appressis et constrictis, deinde tumido-converxis, striatis et malleatis; aperturā subpyriformi-rotundatā.*

THE COVERED PALUDINA. Shell scalariformly turbinated, deeply umbilicated, dark olive-green, encircled with three purple-red bands, whorls rather narrow, appressed and constricted at the sutures, then tumidly convex, striated and malleated; aperture somewhat pyriformly rounded.

Cyclostoma contectum, Millet, Moll. Maine et Loire, p. 5.

Nerita vivipara, Müller (not *Helix vivipara*, Linnaeus).

Cyclostoma viviparum, Draparnaud.

Natica vivipara, Féussae.

Lymnaea vivipara, Fleming.

Paludina vivipara, Studer.

Paludina crystallina, Gray.

Vivipara communis, Dupuy.

Paludina Listeri, Forbes and Hanley.

Hab. Central Europe.

This well-known species is distinguished from *P. vivipara* by its more scalariform growth, constricted at the sutures, and deep umbilicus.

Species 4. (Fig. *a, b*, Mus. Cuming.)

PALUDINA VIVIPARA. *Pal. testā obtusē ovatā, minutē compressē umbilicatā, subobesā, fulvo-olivaceā, fasciis tribus purpureo-rufis cingulatā, anfractibus latiusculis, converxis, confertim plicato-striatis; aperturā subpyriformi-ovatā.*

THE VIVIPAROUS PALUDINA. Shell obtusely ovate, minutely compressly umbilicated, rather stout, fulvous-olive, encircled with three purple-red bands, whorls rather broad, convex, closely plicately striated; aperture somewhat pyriformly ovate.

Helix vivipara, Linnaeus, Syst. Nat. ed. 10. p. 772.

Nerita fasciata, Müller.

Helix fasciata, Gmelin.

Helix ventricosa, Olivi.

Helix compactilis, Pulteney.

Bulinus viviparus, Poiret.

Cyclostoma achatinum, Draparnaud.

Paludina achatina, Studer.

Paludina vulgaris, Gray.

Turbo achatinus, Sheppard.

Paludina vivipara, Say.

Paludina lineata, Valenciennes.

Paludina fasciata, Deshayes.

Paludina Georgiana, Lea.


Vivipara fasciata, Dupuy.

Vivipara vivipara, Binney.

Vivipara vera, Frauenfeld.

Hab. Central Europe.

Composed of a whorl less than the preceding species, the whorls being broader and more compactly coiled.



PALUDINA.

PLATE II.

Species 5. (Fig. a, b, Mus. Cuming.)

PALUDINA BENGALENSIS. *Pal. testā elevato-conicā, tenui, virescente, nitente, saturatē viridi lineatā et angustē fasciatā; anfractibus convexis, læribus, minutissimè decussatim striatis.*

THE BENGAL PALUDINA. Shell elevately conical, thin, greenish, shining, lineated and narrowly banded with dark-green; whorls convex, smooth, very minutely decussately striated.

LAMARCK, Anim. sans vert. vol. vi. part 2. p. 174.
Delessert, Rec. Coq. pl. 1. f. 2.

Paludina elongata, Swainson.

Hab. North-west Provinces of Hiudostan; Benson.

A shining pale-green shell, banded and lineated throughout with dark-green, sometimes turning to rusty-brown. The surface is very minutely engraved with impressed spiral striae, decussating slightly-raised longitudinal striae, which are striae of growth.

Species 6. (Mus. Cuming.)

PALUDINA COSTATA. *Pal. testā elevato-conicā, ustulato-olivaceā, anfractibus supernē plano-declivibus, deinde acutē angulatis, costis lirisque tenuibus filiformibus undique funiculatis, anfractu ultimo subventricoso, ad basin multistriato.*

THE BIBBED PALUDINA. Shell elevately conical, swarthy-olive, whorls flatly sloping round the upper part, then sharply angled, corded throughout with fine thread-like ribs and ridges, last whorl rather ventricose, many-striated at the base.

QUOY, Moll. Voy. de l'Astrol. pl. 80. f. 1 to 3.

Hab. Borneo.

Of a peculiar swarthy-olive colour, sharply angled at the shoulder, corded throughout with fine thread-like ribs and ridges, which are rather paler in colour, yellowish. It comes very near to *P. angularis*.

Species 7. (Mus. Cuming.)

PALUDINA GIGANTEA. *Pal. testā elevato-conicā, subven-tricosā, virescente, saturatē viridi plus minus obscurè*

fasciatā et lineatā, anfractibus supernē obtusè tumidis, deinde rotundatis, decussatim minutissimè striatis, striis longitudinalibus numerosis, confertis.

THE GIGANTIC PALUDINA. Shell elevately conical, rather ventricose, pale-green, more or less obscurely banded and lineated with dark-green, whorls obtusely swollen round the upper part, then rounded, decussately very minutely striated, longitudinal striae numerous, close-set.

VON DEM BUSCH, MS. in Mus. Cuming.

Hab. Bengal; Central Africa.

It is somewhat doubtful whether this is not a light inflated form of *P. Bengaleensis*; yet Mr. Cuming possesses a smaller brighter-coloured shell, from Central Africa, which has these peculiarities developed in a manner which would lead to the conclusion that it is distinct.

Species 8. (Fig. a, b, Mus. Cuming.)

PALUDINA USSURIENSIS. *Pal. testā subgloboso-conicā, flavescente, viridi tinctā, rufo plerunque trifasciatā, anfractibus supernē planulatis, deinde tumido-convexus, spiraliter carinatis, longitudinaliter irregulariter pli-cato-liratis, interstitiis malleatis et striatis.*

THE USSURI PALUDINA. Shell somewhat globosely conical, yellowish, tinged with green, generally encircled with three red bands, whorls a little flattened at the upper part, then tumidly convex, spirally keeled, longitudinally irregularly plicately ridged, interstices malleated and striated.

GERSTFELDT, Land und Süss. Moll. Sib. und Amur-Geb. part 1. p. 2. pl. 1. f. 1 to 4.

Hab. Mouth of the Ussuri, a tributary of the River Amoor, Siberia; Gerstfeldt.

A fine and very characteristic Siberian species, in which the whorls are keeled, but the keels are serrated by irregular longitudinal ridges and striae, which are scarcely developed until the shell approaches maturity, and seem to indicate a departure from the typical symmetry of the plan of growth.

PALUDINA.—PLATE II.

Species 9. (Mus. Cuming.)

PALUDINA OXYTROPIS. *Pal. testā pyramidali-conicā, tenui, ventricosā, ferrugineaā, ustulato-fusco fasciatā, anfractibus declivi-convexit, costis tribus distantibus regulariter funiculatis, infernē inflato-angulatis.*


THE STRONGLY-KEELED PALUDINA. Shell pyramidal-conical, thin, ventricose, rust-stained, banded with burnt-brown, whorls slopingly convex, regularly

corded with three distant ribs, inflatedly angled at the lower part.

BENSON, Journ. Asiat. Soc. 1836, vol. v.

Paludina pyramidata, Von dem Busch.
Hab. Bengal.


A pyramidal-conical shell, of light inflated growth, encircled throughout with three strongly-developed distant cord-like keels.



118



119.



119.



120.



120.



121.



122.

PALUDINA.

PLATE III.

Species 10. (Mus. Cuming.)

PALUDINA PREROSA. *Pal. testā conico-turbinatā, crassā, solidā, olivaceā, apice erosā, anfractibus subangustis, ad suturas constrictis, deinde declivi-rotundatis, undique obliquè plicato-striatis, striis rudibus, valde irregularibus.*

THE ERODED PALUDINA. Shell conically turbinated, thick, solid, olive, eroded at the apex, whorls rather narrow, constricted at the sutures, then slopingly rounded, everywhere obliquely plicately striated, striae rude, very irregular.

GERSTFELDT, Land und Siiss. Moll. Sib. und Amur-Geb. Part I. p. 5, pl. 1, f. 5 to 7.

Hab. Mouth of the Ussuri, a tributary of the river Amoor, Siberia; Gerstfeldt.

A shell of remarkably solid growth, unusually constricted at the sutures.

Species 11. (Fig. a, b, Mus. Cuming.)

PALUDINA CUMINGII. *Pal. testā ovato-conicā, olivaceā; anfractibus subangustis, rotundatis, superne impresso-planis, undique minutè decussatim striatis et malleatis.*

CUMING'S PALUDINA. Shell ovately conical, olive; whorls rather narrow, rounded, impressly flattened at the upper part, everywhere minutely decussately striated and malleated.

HANLEY, MS. in Mus. Cuming.

Hab. Calanang, Province of Baic, Island of Luzon, Philippines.

Distinguished from the European *P. contexta*, which it much resembles in the roundness of the whorls and in the decussately-striated malleated character of the surface, by the peculiar impressed flatness of their upper edge next the sutures. *Paludina Cumingii*, D'Orbigny, is a species of *Paludestrina*.

Species 12. (Mus. Cuming.)

PALUDINA AMPULLACEA. *Pal. testā subgloboso-conicā, flavescens-olivaceā, anfractibus subangustis, tumido-rotundatis, longitudinaliter densè striatis et malleatis.*

THE BOTTLE PALUDINA. Shell somewhat globosely conical, yellowish-olive, whorls rather narrow, tumidly rounded, longitudinally densely striated and malleated.

CHARPENTIER, MS. in Mus. Cuming.

Hab. Italy.

This fine species is not striated spirally, but only longitudinally, and the whorls are unusually widely convoluted. It may prove to be a gigantic form of *P. contexta*.

Species 13. (Fig. a, b, Mus. Cuming.)

PALUDINA JAPONICA. *Pal. subpyramidalis-ovata, virescente-olivaceā, nitente, anfractibus convexo-declivibus, infernè obtuse angulatis et carinatis, longitudinaliter plicato-striatis, striis versus aperturam conserioribus rugosis.*

THE JAPAN PALUDINA. Shell somewhat pyramidalis-ovata, greenish-olive, shining, whorls convexly sloping, obtusely angled and keeled round the lower part, longitudinally plicately striated, striae closer and rough towards the aperture.

MARTENS, Malak. Blätt. vol. vii. p. 44.

Hab. Japan.

A very distinct form of *Paludina*, in which the whorls slope to an obtuse angle at the lower part, the angle falling in the place of the suture.

Species 14. (Mus. Cuming.)

PALUDINA ANGULARIS. *Pal. testā elevato-conicā, saturatē olivaceā, anfractibus subangustis, fortiter decussatim striatis, superne angulatis, costis tenuibus tribus, liris duabus in interstitiis, spiraliter funiculatis.*

THE ANGULAR PALUDINA. Shell elevately conical, dark-olive, whorls rather narrow, strongly decussately striated, angled round the upper part, spirally corded with three fine ribs, having two faint ridges in the interstices.


Nerita angularis, Müller, Hist. Verm. part 2. p. 187.

Helix angularis, Gmelin.

Paludina angularis, Philippi.

Hab. Manila, Island of Luzon, Philippines; Cuming.

Closely allied to *P. costata*, but it is a more delicately-formed shell, of a brighter olive colour.



P A L U D I N A.

PLATE IV.

Species 15. (Mus. Cuming.)

PALUDINA INTERTEXTA. *Pal. testā abbreviato-globosā, intensē viridi, spirā parum exsertā, anfractibus subangustis, tumido-rotundatis, striis minutis crispato-punctatis undique decussatis.*

THE WOVEN PALUDINA. Shell shortly globose, dark-green, spire but little exserted, whorls rather narrow, tumidly rounded, decussated throughout with minute crisply-punctured striae.

SAY, Amer. Coneh. pl. 30. f. 3, 4.

Hab. New Orleans (in lakes and marshes); Say.

Of a curious, shortly-globose form, very dark green in colour, with the surface minutely superficially sculptured with decussating crisply-punctured striae. *P. sinistrorsa*, Lea, may be a sinistral state of this species.

Species 16. (Mus. Cuming.)

PALUDINA BURROUGHIANA. *Pal. testā conico-oratā, flavescente, anfractibus rotundatis, subobsoletè spiraliter multiliratis, undique malleatis et decussatim rugoso-striatis.*

BURROUGH'S PALUDINA. Shell conically ovate, yellowish, whorls rounded, rather obsoletely spirally many-ridged, everywhere malleated and decussately rugosely striated.

LEA, Trans. Amer. Phil. Soc. vol. v. pl. 19. f. 80.

Hab. Manilla, Island of Luzon, Philippines.

The only specimen I have been able to procure is characterized by a variety of sculpture, confusedly developed, and more or less obsolete. Mr. Lea figures one of much larger size, and darker olive-green colour, with the spiral ridges more defined.

Species 17. (Mus. Cuming.)

PALUDINA QUADRATA. *Pal. testā pyramidalis-conicā, solidiusculā, fusco-olivaceā, anfractibus declivibus, plano-convexis, inferne obtusè angulatis et carinatis, longitudinaliter obliquè striatis, liris duabus vel tribus plus minus obsoletis interdum cinctis.*

THE SQUARED PALUDINA. Shell pyramidal conical, rather solid, brown-olive, whorls sloping, flatly convex, obtusely angled and keeled round the lower part,

longitudinally obliquely striated, sometimes encircled with two or three more or less obsolete ridges.

BENSON, Ann. and Mag. Nat. Hist. 1842. Phil. Abbild. vol. ii. p. 135. pl. 2. f. 5.

Hab. China.

Well distinguished by its solid pyramidal-conical form, obtusely angled at the base.

Species 18. (Mus. Cuming.)

PALUDINA CHINENSIS. *Pal. testā ovato-ventricosū, fusco-olivaceā, anfractibus latiusculis, inflatis, obliquè longitudinaliter plicato-striatis, versus apicem malleatis.*

THE CHINA PALUDINA. Shell ovately ventricose, brown-olive, whorls rather broad, inflated, obliquely longitudinally plicately striated, malleated towards the apex.

GRAY, Griffith's Cuv. Anim. King. Moll. pl. 1. f. 5.

Hab. Chusan.

The whorls of this species are broader and more ventricose than in any other of the genus.

Species 19. (Mus. Cuming.)

PALUDINA INFLATA. *Pal. testā conico-turbanatā, flavescente-olivaceā, anfractibus subangustis, rotundatis, ad suturas profundè impressis, undique rugoso-malleatis et decussatim plicato-liratis et minutissimè striatis; apertura marginē purpureo-nigro.*

THE INFLATED PALUDINA. Shell conically turbinated, yellowish-olive, whorls rather narrow, rounded, deeply impressed at the sutures, roughly malleated and decussately plicately ridged, and very minutely striated; margin of the aperture purple-black.

VILLA, Archiv. Skandinav. 1850, p. 330.

Hab. Sweeden; Austria.

The whorls of this species are rather more than usually pressed in at the sutures.

Species 20. (Mus. Cuming.)

PALUDINA VIRIDIS. *Pal. testā elevato-conicā, intensē viridi, anfractibus rotundatis, liribus, ad suturas subconstrictis, sub lente undique minutè decussatim striatis.*

PALUDINA.—PLATE IV.

THE GREEN PALUDINA. Shell elevately conical, dark green, whorls rounded, smooth, rather constricted at the sutures, beneath the lens minutely decussately striated throughout.

HANLEY, MS. in Mus. Cuming.

Hab. —?

An apparently smooth shell, of very dark green colour, but elaborately decussately striated beneath the lens.

Species 21. (Mus. Cuming.)

PALUDINA LECYTHOIDES. *Pal. testá conico-ovatá, sub-*

ventricosá, flavescente-olivacea, obscurè pallidissimè fasciatá, anfractibus rotundatis, tumidiusculis, ad suturas subconstrictis, longitudinaliter irregulariter pliato-striatis.

THE YOLK-LIKE PALUDINA. Shell conically ovate, rather ventricose, yellowish-olive, obscurely very faintly banded, whorls rounded, rather swollen, slightly constricted, longitudinally irregularly plicately striated.

BENSON, Phil. Abbild. vol. ii. p. 133. pl. 2. f. 1.

Hab. Chusan.

A shell of very simple character, quite devoid of spiral striae.



23.



23.



24.




25.a.



25.a.



25.b.



25.b.



27.



28.



29.

PALUDINA.

PLATE V.

Species 22. (Mus. Cuming.)

PALUDINA COOSAENSIS. *Pal. testā subglobosā, sordidē olivaceā, spirā scalariformi, suturis valdē impressis, anfractibus subangustis, rotundatis, longitudinaliter corrugatim crispato-striatis, spiraliter lineari-incisis; aperturā lunato-circulari.*

THE COOSA PALUDINA. Shell subglobose, dusky-olive, spire scalariform, sutures very much impressed, whorls rather narrow, rounded, longitudinally corrugately crisply striated, spirally linearly engraved; aperture lunar-circular.

LEA, Trans. Amer. Phil. Soc. vol. ix. p. 23.

Vivipara Coosaensis, Binney.

Hab. Coosa River, Alabama, United States.

Distinguished by an impressively channelled suture, and a peculiar sculpture of longitudinal crisply wrinkled striae.

Species 23. (Mus. Cuming.)

PALUDINA WAREANA. *Pal. testā subgloboso-conicā, solidiusculā, olivaceā, anfractibus convexis, copiosè constricto-striatis; aperturā luanari-ovatā, parviusculā.*

THE WARE PALUDINA. Shell subglobosely conical, rather solid, olive, whorls convex, copiously constrictedly striated; aperture lunar-ovate, rather small.

SHUTTLEWORTH, Küster, Conch. Cab. p. 21. pl. iv. f. 10. 11.

Vivipara Wareana, Binney.

Hab. Lake Ware, East Florida, United States.

According to the specimen preserved in Mr. Cuming's collection as a type of this species, it is a shell of rather solid growth, densely constrictedly striated throughout, but yet thin and transparent towards the aperture.

Species 24. (Mus. Cuming.)

PALUDINA TEXANA. *Pal. testā subpyramidalī-conicā, solidā, virescente, suturis impressis, anfractibus planodeclivibus, medio subconcavis, ad basin rotundatis; aperturā pyriformi-circulari, parviusculā.*

THE TEXAS PALUDINA. Shell somewhat pyramidalily conical, solid, light green, sutures impressed, whorls flatly sloping, slightly concave in the middle, rounded

at the base; aperture pyriformly circular, rather small.

Vivipara Texana, Tryon, Pro. Acad. Nat. Sci. Phil. 1862.

Vivipara subpurpurea, var., Binney.

Hab. Texas, United States.

This species has been lately described by Mr. Tryon. Mr. Binney considers it a variety of *P. subpurpurea*; but if the specimen here figured can be relied upon as a type, and I know not to what other species it can be referred, *P. Texana* should, I think, be retained.

Species 25. (Fig. a, b, Mus. Cuming.)

PALUDINA MALLEATA. *Pal. testā globoso-conicā, subinflatā, vividē olivaceā, anfractibus rotundatis, longitudinaliter copiosè plicato-striatis, spiraliter obscurè linearī-puncturatis, ubique malleatis; aperturā pyriformi-ovatā, labro nigro-marginato.*

THE MALLEATED PALUDINA. Shell globosely conical, rather inflated, bright olive, whorls rounded, longitudinally copiously plicately striated, spirally obscurely linearly punctured, everywhere malleated; aperture pyriformly ovate, lip black-edged.

Hab. Japan.

This and the following Japanese species differ from the American species of similar form, *P. intertexta* for example, in being malleated throughout.

Species 26. (Fig. a, b, Mus. Cuming.)

PALUDINA ABBREVIATA. *Pal. testā globosā, nigrante olivaceā, spirā abbreviatā, anfractibus rotundatis, ad suturas appressis, infernè obsoletè angulatis, longitudinaliter copiosè constricto-striatis, spiraliter obscurè linearī-puncturatis, ubique malleatis; aperturā pyriformi-ovatā, labro nigro-marginato.*

THE SHORTENED PALUDINA. Shell globose, dark olive, spire shortened, whorls rounded, appressed at the sutures, obsoletely angled round the lower part, longitudinally copiously constrictedly striated, spirally obscurely linearly punctured, everywhere malleated; aperture pyriformly ovate, lip black-edged.

Hab. Japan.

Similar to the preceding species in detail of character,

but of different form. The spire is abbreviated, impressing the sutures into a canal, and they are obscurely obtusely angled round the lower part.

Species 27. (Mus. Cuming.)

PALUDINA MELANOSTOMA. *Pal. testā globoso-conicā, flavescente-olivaceā, infernē pallidē fasciatā, anfractibus convexis, subinflatī, ubique malleatis et minulē puncturatis; aperturā pyriformi-circulari, parviusculā.*

THE BLACK-MOUTHED PALUDINA. Shell globosely conical, yellowish-olive, faintly banded round the lower part, whorls convex, rather inflated, everywhere malleated and minutely punctured; aperture pyriformly circular, rather small.

BENSON, MS.

Hab. Bengal.

A faintly banded shell, with a malleated surface, crowded throughout with minute punctures.

Species 28. (Mus. Anthony.)

PALUDINA BERMONDIANA. *Pal. testā profundē angustē*

umbilicatā, globosā, flavescente-olivaceā, purpureo conspicuē trifasciatā, spirā brevi, anfractibus rotundatis, ad suturas impressis, densē longitudinaliter striatis; aperturā ovatā.

BERMOND'S PALUDINA. Shell deeply narrowly umbilicated, globose, yellowish-olive, encircled with three conspicuous purple bands, spire short, whorls rounded, impressed at the sutures, closely longitudinally striated; aperture ovate.


D'ORBIGNY, Moll. Cuba, p. 7. pl. 10. f. 5.

Hab. Cuba.

A compact, closely convoluted shell, with a small spire, and narrow deep umbilicus, encircled with three very distinct purple bands, the upper one of which is in the suture.

Fig. 29. (Mus. Cuming.)

The shell here figured, *Paludina nucleus* of Mousson, inhabiting the vicinity of Constantinople, appears to me to be a variety of *P. rivipara*.



P A L U D I N A.

PLATE VI.

Species 30. (Mus. Cuming.)

PALUDINA AMPULLARIOIDES. *Pal. testā subgloboso-conicā, solidiusculā, fuscescente-olivaceā, purpureo irregulariter lineuri-fasciatā, anfractibus rotundatis, ad suturas subangulato-appressis, ubique minutē puncturatis; aperturā pyriformi-ovatā.*

THE AMPULLARIA-LIKE PALUDINA. Shell somewhat globosely conical, rather solid, dusky olive, irregularly linearly banded with purple, whorls rounded, rather angularly appressed at the sutures, minutely punctured throughout; aperture pyriformly ovate.

HANLEY, MS. in Mus. Cuming.

Hab. —?

A very distinct minutely punctured species, of rather solid growth, encircled very irregularly with purple bands, some of which are quite linear.

Species 31. (Mus. Cuming.)

PALUDINA FILOSA. *Pal. testā ovato-conicā, flavescente-olivaceā, nitente, fasciis quatuor linearibus aequidistantibus nigris cingulatā, anfractibus convexis, lāvibus, sub lente obsoletē minutē puncturatis; aperturā subcirculari.*

THE THREAD-ENCIRCLED PALUDINA. Shell ovately conical, yellowish-olive, shining, encircled with four equidistant black linear bands, whorls convex, smooth, beneath the lens obsoletely minutely punctured; aperture nearly circular.

HANLEY, MS. in Mus. Cuming.

Hab. —?

Another very characteristic species, with a shining yellowish-olive surface neatly ruled with black linear bands.

Species 32. (Mus. Cuming.)

PALUDINA CEYLONICA. *Pal. testū orato-conicā, solidiusculā, virescente-olivaceā, anfractibus plano-convexis, supernē et infernē declivi-angulatis, ubique densē spiraliiter lineari-striatis et puncturatis, ad angulos obtusē carinatis; aperturā pyriformi-ovatā, nigro-marginali.*

THE CEYLON PALUDINA. Shell ovately conical, rather solid, greenish-olive, whorls flatly convex, flatly angled

above and below, everywhere closely spirally linearly striated and punctured, obtusely keeled at the angles aperture pyriformly ovate, black-edged.

DOHRN, Pro. Zool. Soc. 1857, p. 123.

Hab. Ceylon.

Readily recognized by its obtusely-angled form and dense sculpture of spiral lines and punctures.

Species 33. (Fig. a, b, Mus. Cuming.)

PALUDINA CRASSA. *Pal. testā orato-turbinalatā, solidā, flavescente-olivaceā, anfractibus rotundatis, ad suturas constrictis, obliquē plicato-striatis, spiraliiter minutissimē undato-striatis et puncturatis; aperturā ferè circulari, nigro-marginatā.*

THE THICK PALUDINA. Shell ovately turbinate, solid, yellowish-olive, whorls rounded, constricted at the sutures, obliquely plicately striated, spirally very minutely wave-striated and punctured; aperture nearly circular, black-edged.

HUTTON, Benson, Journ. Asiatic. Soc. 1836, p. 745.

Paludina obtusa, Troschel.

Hab. Jaunpore, India; Burmah.

Of a peculiarly solid constricted growth, copiously minutely punctured, like most Eastern species.

Species 34. (Mus. Cuming.)

PALUDINA ZONATA. *Pal. testā angustē subprofundē umbilicatā, ovato-conicā, viridē virente-olivaceā, nitente, fasciis nigris linearibus nitidis cingulatā, anfractibus lāvibus, sub lente minutē puncturatis; aperturā pyriformi-circulari, nigro-marginali.*

THE ZONED PALUDINA. Shell narrowly rather deeply umbilicated, ovately conical, bright greenish-olive, shining, encircled with neat black linear bands, whorls convex, smooth, beneath the lens minutely punctured; aperture pyriformly circular, black-edged.

HANLEY, MS. in Mus. Cuming.

Hab. Mauritius.

Another shining black-banded species of the same bright porcelain-like type as *P. filosa* figured above it.

Species 35. (Mus. Cuming.)

PALUDINA PRÆMORSA. *Pal. testā ovato-conicā, olivaceā, anfractibus convexis, longitudinaliter minutē decussatim crissato-striatis, spiraliter dense minutissimē puncturatis; aperturā pyriformi-circulari, nigro-marginatā.*

THE ERODED PALUDINA. Shell ovately conical, olive, whorls convex, longitudinally minutely decussately crisply striated, spirally densely very minutely punctured; aperture pyriformly circular, black-edged.

BENSON MS., Philippi (misprinted *Remossii*), Abbild. und Besch. Conch. vol. ii. p. 134. pl. 2. f. 3.

Hab. Bengal.

I have the authority of Mr. Benson for stating that the name *Remossii* published by Philippi, and adopted by Küster and Frauenfeld, is a mistaken reading of the name *præmorsa*, communicated by Mr. Benson to Dr. Philippi in manuscript. As there is no such person as Remosse, that name must be cancelled.

Species 36. (Fig. a, b, Mus. Cuming.)

PALUDINA CILIATA. *Pal. testā profundē angustē umbilicatā, ovato-conicā, solidā, olivaceo-viridi, versus apicem purpurascente tinctā, anfractibus convexis, undique spiraliter lineari-ciliatis, juxta suturas angustē angulatis, ungulo versus aperturam exanido, anfractu ultimo circa umbilicum excavato et umbilicato; aperturā pyriformi-circulari, viridē nigro-marginatā.*

THE EYE-LASH PALUDINA. Shell deeply narrowly umbilicated, ovately conical, solid, olive-green, purple-tinged towards the apex, whorls convex, spirally linearly eyelashed throughout, narrowly angled next the sutures, angle disappearing towards the aperture, last whorl excavated and keeled around the umbilicus; aperture pyriformly circular, with a shining black edge.

Hab. Siam; Mouhot.

This very remarkable and beautiful species, encircled throughout with minute bristly hair-lashes, was collected the year before last by the unfortunate Mouhot.

Species 37. (Mus. Cuming.)

PALUDINA HAMILTONI. *Pal. testā angustē profundē umbilicatā, attenuatim conico-ovatā, intensē olivaceā, spirā elatā, anfractibus rotundatis, juxta suturas appressis, minutē decussatim striatis, ultimo circa umbilicum subexcavato; aperturā pyriformi-circulari.*

HAMILTON'S PALUDINA. Shell narrowly deeply umbilicated, attenuately conically ovate, dark olive, spire raised, whorls rounded, appressed next the suture, minutely decussately striated, last whorl a little excavated around the umbilicus; aperture pyriformly circular.

METCALFE, Pro. Zool. Soc. 1851, p. 73.

Hab. Borneo.

A species apparently of an ordinary character, but partaking of the Siam type in being angularly excavated, though in a slight degree, around the umbilicus.



P A L U D I N A.

PLATE VII.

Species 38. (Fig. *a*, *b*, *c*, Mus. Cuming.)

PALUDINA POLYZONATA. *Pal. testā subpyramidalī-conicā, flavescente-olivaceā, nigricante-fusco linearī-fasciatā, anfractibns declivi-converis, ad basin angulatis, carinis subdistantibns undique funiculatis; aperturā ovatā, margine columellari subcalloso, carneō tincto.*

THE MANY-ZONED PALUDINA. Shell somewhat pyramidal-conical, yellowish-olive, linearly banded with dark brown, whorls slopingly convex, angled at the base, eorded throughout with rather distant keels; aperture ovate, columellar margin rather callous, flesh-tinted.

Vivipara polyzonata, Frauenfeld, Ueb. die Gatt. p. 4.

Hab. Philippine Islands.

Of the same peculiar type as *P. oxytropis* of Bengal, but of more solid growth and smaller. The cords in each species are conspicuously raised.

Species 39. (Fig. *a*, *b*, Mus. Cuming.)

PALUDINA INGALLSIANUS. *Pal. testā obliquē pyramidalī-conicā, cinereo-viridi, olivaceo striatā et variegatā, spirā elatā, acutā, anfractibns declivi-convexis, obliquē plicato-corrugatis et crispato-striatis, spiraliter minutē puncturatā, anfractu ultimo infra peripheriam unicarinate; aperturā pyriformi-ovatā, fance callosā, albā.*

INGALLS' PALUDINA. Shell obliquely pyramidal-conical, ash-green, streaked and variegated with olive, spire elevated, sharp, whorls slopingly convex, obliquely plicately wrinkled and crisply striated, spirally minutely punctured, last whorl encircled with a keel below the periphery; aperture pyriformly ovate, interior callous, white.

LEA, Pro. Acad. Nat. Sci. Phil. June, 1856.

Hab. Japan.

This fine and remarkable species, of which there are many varieties known, was described by Mr. Lea, from specimens collected in Japan by Mr. Ingalls, an American missionary.

Species 40. (Fig. *a*, *b*, Mus. Cuming.)

PALUDINA UMBILICATA. *Pal. testā subobtectē umbilicatā, pyramidalī-conicā, subabbreviatā, solidā, flarescente-olivaceā, versns apicem purpureo-cinereā, anfractibus supernē appressē rotundatis, deinde plano-converis, ad basin carinatis et angulatis, undique obliquē arcuatim striatis, spiraliter minutē puncturatā; aperturā subquadrato-ovatā, fance callosā, albā.*

THE UMBILICATED PALUDINA. Shell with a covered umbilicus, pyramidal-conical, rather shortened, solid, yellowish-olive, purple-ash towards the apex, whorls appressly rounded at the upper part, then flatly convex, keeled and angled at the base, everywhere obliquely arcuately striated, spirally minutely punctured; aperture somewhat squarely ovate, interior callous, white.

LEA, Pro. Acad. Nat. Sci. Phil. June, 1856.

Hab. Siam; Mouhot.

Curiously allied in typical character to the Japanese *P. Ingallsiana*, but of a peculiar shortened subquadrate form.


Species 41. (Fig. *a*, *b*, *c*, Mus. Cuming.)

PALUDINA ERUGINOSA. *Pal. testā subangustē pyramidalī-conicā, intensē cæruleo-viridi, anfractibus declivi-convexis, interdum versus basin obtusē carinatis, ubique obliquē plicato-striatis, sub lente spiraliter obsolete minutissimē puncturatā; aperturā pyriformi-ovatā.*

THE BRONZE PALUDINA. Shell rather narrowly pyramidal-conical, dark blue-green, whorls slopingly convex, sometimes obtusely keeled towards the base, everywhere obliquely plicately striated, beneath the lens spirally obsolete very minutely punctured; aperture pyriformly ovate.

Hab. China.

Very closely allied to *P. quadrata*, from the same locality, the specimens are uniformly lighter, and of a peculiar bronze blue-green colour.



P A L U D I N A.

PLATE VIII.

Species 42. (Mus. Cuming.)

PALUDINA HAINESIANA. *Pal. testā obliquē globosā, solidā, olivaceā, anfractibus ad suturas impressis, deinde declivi-convexis, versus basin gibbosis, ubique obliquē plicato-striatis, spiraliter minutissimē punctatis; aperturā ovatā.*

HAINES' PALUDINA. Shell obliquely globose, solid, olive, whorls impressed at the sutures, then slopingly convex, gibbous towards the base, everywhere obliquely plicately striated, spirally very minutely punctured; aperture ovate.

LEA, MS. in Mus. Cuming.

Hab. Sian.

A solid globose shell, still bearing a crowded profusion of the very minute punctures characteristic of the Eastern species.

Species 43. (Mus. Cuming.)

PALUDINA CYCLOSTOMATIFORMIS. *Pal. testā cylindraceo-turrīta, solidā, olivaceā, anfractibus rotundatis, lēvibus, ad suturas constricto-impressis; aperturā obliquē ovatā, parvā.*

THE CYCLOSTOMA-SHAPED PALUDINA. Shell cylindrically turreted, solid, olive, whorls rounded, smooth, constrictedly impressed at the sutures; aperture obliquely ovate, small.

LEA, Trans. Amer. Phil. Soc. vol. ix. part i. p. 23.

Paludina contorta, Shuttleworth.

Paludina Elliotti, Lea.

Ticipara Cyclostomatiformis, Binney.

Hab. Coosa River, Alabama, United States.

Well distinguished from all other species of the genus by its narrow cylindrically-turreted form and small aperture.

Species 44. (Mus. Cuming.)

PALUDINA SUBCARINATA. *Pal. testā globoso-turrīta, flavescente-olivaceā, spirā ad apicem acutā, carneotinctū, suturis impresso-canaliculatis, anfractibus rotundatis, minutē decussatim undato-striatis, superne obtusè carinatis; aperturā obliquē ovatā.*

THE FAINTLY-KEELED PALUDINA. Shell globosely turreted, yellowish-olive, spire sharp at the apex, flesh-tinted, sutures impressly channelled, whorls rounded,

minutely decussately wave-striated, obtusely keeled round the upper part, aperture obliquely ovate.

SAY. Nicholson Eney. pl. i. f. 7.

Paludina sulclosa, Menke.

Paludina bicarinata, Potiez and Michaud.

Helix decisā, Wood.

Hab. Alabama, Ohio, etc., United States.

Sharp-pointed and flesh-tinged at the apex, whorls rudely tumidly rounded next the sutures, and presenting one or two obtusely developed keels.

Species 45. (Fig. a, c, d, Mus. Anthony; Fig. b, Mus. Cuming.)

PALUDINA DECISA. *Pal. testā ovatā, plus minus conica, solidā, vivide virescente-olivaceā, anfractibus concavis, sub leute niantissimē reticulato-striatis, ad suturas ruāe appressis; aperturā pyriformi-ovatā, fance callosā, cārulecente-albā.*

THE ERODED PALUDINA. Shell ovate, more or less conical, solid, bright greenish-olive, whorls concave, beneath the lens very minutely reticulately striated, rudely appressed at the sutures; aperture pyriformly ovate, eallous, bluish-white within.

SAY. Nicholson Eney. pl. 3. f. 5.

Paludina integra and *ponderosa*, Say.

Paludina genicula, Conrad.

Paludina limosa, *cornea*, and *ovalaris*, Valenciennes.

Paludina heterostropha, Kirtland.

Paludina rufa, Haldeman.

Helix dissimilis, Wood.

Paludina subsolida, Anthony.

Ticipara ponderosa and *decisa*, Binney.

Hab. Alabama, Georgia, Ohio, etc., United States.

Far too many species have been made of this well-known and somewhat variable species. The extreme forms are Say's *P. integra* (Fig. 45 a), in which the spire is elevated, and Conrad's *P. genicula* (Fig. 45 c, d), in which it is contracted. Kirtland's *P. heterostropha* is the same form reversed. Say's *P. ponderosa* (Fig. 45 b) is an intermediate form of rudely-developed thickened growth.

Species 46. (Fig. 46 a, Mus. Anthony.)

PALUDINA COARCTATA. *Pal. testā pyramidali-conica,*

solidā, virescente-olivaceā, anfractibus ad suturas constrictis, deinde declivi-converis, planiusculis, sub lente minutē decussatim undulato-striatis; aperturā subtri-gono-ovatā, parviusculā.

THE CONTRACTED PALUDINA. Shell pyramidal conical, solid, greenish-olive, whorls constricted at the sutures, then slopingly convex, rather flat, beneath the lens minutely deenssately wave-striated; aperture somewhat triangularly ovate, rather small.

LEA, Trans. Amer. Phil. Soc. vol. ix. p. 30.

Paludina exilis, Anthony.

Vivipara coarctata, pars, Binney.

Hab. Mississippi, United States.

I follow Mr. Binney in referring Mr. Anthony's *P. exilis* to Mr. Lea's *P. coarctata*, but cannot agree with him in referring Mr. Anthony's *P. lima* to the same species. I am indebted to the assiduous conchologist of Cincinnati for the loan of the same typical specimens of which Mr. Binney has given outline sketches in his monograph. Here they are represented in finished drawings, and no figures could be more characteristic. It will be seen by the numbering of the figures that I determined in the first instance to follow Mr. Binney in giving Mr. Anthony's *P. exilis* and *lima* as varieties of one and the same species, but on a further examination of their characters in detail, I revert to the opinion that they are distinct.

Species 47. (Mus. Anthony.)

PALUDINA SUBPURPUREA. *Pal. testā subcylindraceo-tur-ritā, solidā, olivaceā, purpurascente pallidissimē tinctā,*

anfractibus ad suturas rudē constrictis, supernē de-clivi-contractis, deinde rotundatis, undique plicato-striatis; aperturā pyriformi-circulari, parciusculā.

THE PURPLE-TINGED PALUDINA. Shell somewhat cylindrically turreted, solid, olive, very faintly tinged with purple, whorls rudely constricted at the sutures, slopingly contracted round the upper part, then rounded, plicately striated throughout; aperture pyriformly circular, rather small.

SAY, Amer. Conch. vol. iii. pl. 30. f. 2.

Hab. Southern and western United States.

This species may be readily distinguished by the slopingly contracted growth of the upper part of the whorls, and their constriction at the sutures.

Species 48. (Fig. 46 b, Mus. Anthony.)

PALUDINA LIMA. *Pal. testā oblongo-ovalā, virescente-olivaceā, anfractibus convexis, undique spiraliter sub-remotē obtusē liratis, interstitiis concaviusculis, striis minutis undulatis decussatis; aperturā ovatā.*

THE FILE PALUDINA. Shell oblong-ovate, greenish-olive, whorls convex, spirally obtusely rather distantly ridged throughout, interstices slightly concave, deenssated with minute waved striae; aperture ovate.

ANTHONY, Acad. Nat. Sci. Phil. 1860, p. 70.

Vivipara coarctata, pars, Binney.

Hab. South Carolina, United States.

Obviously distinguished from *P. coarctata* by its light oblong-ovate form and characteristic sculpture.



49



50a



50b



51



52.



53



54



55



56

57

PALUDINA.

PLATE IX.

Species 49. (Mus. Cuming.)

PALUDINA THERSITES. *Pal. testá globosá, fulvo-olivaceá, spirá brevi, subimmersá, suturis canaliculatis, anfractibus supernè obtusè multicarinatis; aperturá parviusculá.*

THE THERSITES PALUDINA. Shell globose, fulvous-olive, spire short, somewhat immersed, sutures channeled, whorls obtusely many-keeled round the upper part; aperture rather small.

HANLEY, MS. in Mus. Cuming.

Hab. —?

The spire is in this species curiously immersed, but it is not a deformity of growth. There is another equally characteristic specimen in the British Museum.

Species 50. (Fig. a, b, Mus. Cuming.)

PALUDINA LINEOLATA. *Pal. testá ovato-conicá, virescente, nigrante-olivaceo multilineatá, anfractibus levibus, supernè declivibus, deinde convexis; aperturá parviusculá, pyriformi-rotundatá.*

THE LINEATED PALUDINA. Shell ovately conical, greenish, many-lineated with dark-olive, whorls smooth, sloping round the upper part, then convex; aperture rather small, pyriformly rounded.

MOUSSON, MS. in Mus. Cuming.

Hab. Siam.

A smooth linearly-painted shell, of the same type as *P. Bengalensis*, uniformly smaller and of a more slopingly conical form.

Species 51. (Mus. Cuming.)

PALUDINA FALLAX. *Pal. testá ovato-conicá, subinflatá, virescente-olivaceá, anfractibus supernè subangulatis, obscurissimè carinatis et malleatis, undique minutè grano-striatis; aperturá pyriformi-ovatá.*

THE DECEPTIVE PALUDINA. Shell ovately conical, rather inflated, greenish-olive, whorls faintly angled round the upper part, very obscurely keeled and malleated, everywhere minutely grain-striated; aperture pyriformly ovate.

Vivipara fallax, Frauenfeld, Ueber die Gattung *Vivipara*, November, 1862.

Hab. Brazils? Madras?

Having compared this shell in Mr. Cuming's collection with Frauenfeld's original type of *P. fallax*, I can affirm, without any doubt, that they are the same. The habitat "Brazils," which Mr. Cuming has with it, may be erroneous. Frauenfeld gives Madras.

Species 52. (Mus. Cuming.)

PALUDINA JAVANICA. *Pal. testá ovato-conicá, solidiusculá, nigricante-olivaceá, anfractibus convexis, obscurissimè carinatis, undique validè linearis-striatis, longitudinaliter plicato-striatis, anfractu ultimo circa umbilicum angulato-excavato; aperturá pyriformi-ovatá.*

THE JAVA PALUDINA. Shell ovately conical, rather solid, dark-olive, whorls convex, very obscurely keeled, strongly linearly striated throughout, plicately striated longitudinally, last whorl angularly excavated around the umbilicus; aperture pyriformly ovate.

VON DEM BUSCH, Albild. und Besch. Conch. pl. 1. f. 11, 12.

Hab. Java.

In this species the granular striae assume the form of linear ridges, and longitudinal striae descend in a plied manner from the sutures.

Species 53. (Mus. Cuming.)

PALUDINA CARINATA. *Pal. testá subgloboso-conicá, fulvá, obscurissimè fasciatá, anfractibus rotundatis, primo ad marginem inferiorem carinatis, deinde striis undulatis sericeis undique confertim decussatis; aperturá ovatá.*

THE KEELED PALUDINA. Shell somewhat globosely conical, fulvous, very faintly banded, whorls rounded, at first keeled at the lower margin, then very closely decussated throughout with waved silken siraæ; aperture ovate.

Hab. Philippine Islands.

Of a peculiar fulvous colour, keeled round the lower edge of all but the last whorl, in which the keel fades away.

PALUDINA.—PLATE IX.

Species 54. (Mus. Anthony.)

PALUDINA MAGNIFICA. *Pal. testā subpyramidalis-ovatā, solidā, fusco-olivaceā, suturis profundē canaliculatis, anfractibus supernē plicato-marginatis, deinde concavis, deinde tuberculatis biseriatim cinctis; aperturā parviusculā, trapezoideā.*

THE MAGNIFICENT PALUDINA. Shell somewhat pyramidal ovate, solid, brown-olive, sutures deeply channelled, whorls plicately margined round the upper part, then concave, then encircled with two rows of tubercles; aperture rather small, trapezoid.

CONRAD, New Freshwater Shells, p. 48. pl. 8. f. 4.

Paludina bimoniifera, Lca.

Vivipara magnifica, Binney.

Hab. Alabama and Georgia, United States.

This fine species is remarkably distinct from any other form of the genus, and partakes largely of the generic character of *Melania*.

Species 55. (Mus. Cuming.)

PALUDINA AMPLIOR. *Pal. testā ovato-conicā, subinflatā, virescente-olivaceā, anfractibus convexis, undique minutē grano-striatis, ultimo infernē obscurissimē angulato; aperturā ovalā.*

THE LARGER PALUDINA. Shell ovately conical, rather inflated, greenish-olive, whorls convex, minutely grain-striated throughout, the last very obscurely angled round the lower part; aperture ovate.

MOUSSON, MS. in Mus. Cuming.

Hab. —?

It is somewhat doubtful whether this is not a variety of *P. Javanica*.

Species 56. (Mus. Cuming.)

PALUDINA DISSIMILIS. *Pal. testā acuminato-ovatā, cinereo-olivaceā, anfractibus supra et infra angulatis, subtiliter linearis-carinatis et grano-striatis, ultimo circa umbilicum angulato-excavato; aperturā parviusculā, pyriformi-ovatā.*

THE DISSIMILAR PALUDINA. Shell acuminate ovate, ash-olive, whorls angled above and below, finely linearly keeled and grain-striated, last whorl angularly excavated around the umbilicus; aperture rather small, pyriformly ovate.

MÜLLER, MS. in Mus. Cuming.

Hab. —?

I give this shell as I find it in Mr. Cuming's collection. It is an unsatisfactory species, not well authenticated, and has much the appearance of a young *P. Ingallsiana*.

Species 57. (Mus. Cuming.)

PALUDINA INTERMEDIA. *Pal. testā globoso-conicā, subinflatā, fuscescente-olivaceā, anfractibus supernē subangulato-impressis, infernē obscurè carinatis, undique minutē grano-striatis, longitudinaliter plicato-striatis; aperturā pyriformi-ovatā.*

THE INTERMEDIATE PALUDINA. Shell globosely conical, rather inflated, fuscous-olive, whorls subangularly impressed round the upper part, obscurely keeled at the lower, minutely grain-striated throughout, pliately striated longitudinally; aperture pyriformly ovate.

HANLEY, MS. in Mus. Cuming.

Hab. —?

Allied to *P. Javanica*, but of a more depressly globose form, reticulately decussated with the characteristic plicate and granular striae.



59



58



60



62



61



63



64



65a



65b



66

PALUDINA.

PLATE X.

Species 58. (Mus. Cuming.)

PALUDINA VARIATA. *Pal. testā ovalo-conicā, sordidē olivaceā, anfractibus rotundatis, subangustis, supernē subimpressis, striis sericeis undulatis confertim decussatis; aperturā parviusculā, ovatā.*

THE VARIED PALUDINA. Shell ovately conical, dull-olive, whorls rounded, rather narrow, faintly impressed round the upper part, closely decussated with waved silken striae; aperture rather small, ovate.

Vivipara variata, Frauenfeld, Ueber die Gattung *Vivipara*, November, 1862.

Hab. Pondicherry.

Of very simple character, known to me only by a single specimen received from M. Frauenfeld.

Species 59. (Mus. Cuming.)

PALUDINA NITENS. *Pal. testā conico-ovatā, virescente-olivaceā, nitente, anfractibus declivi-convexitatis, minutiē decussatim striatis, infernē angulatis, angulo in anfractu ultimo evanido; aperturā ovatā.*

THE SHINING PALUDINA. Shell conically ovate, greenish-olive, shining, whorls slopingly convex, minutely decussately striated, angled at the lower part, angle in the last whorl fading away; aperture ovate.

Hab. Japan.

A minutely-sculptured shining olive shell, obscurely angled around a small umbilicus.

Species 60. (Mus. Cuming.)

PALUDINA AETHIOPS. *Pal. testā globoso-ovatā, subcylindraceā, tenui, semipellucidā, virescente-olivaceā, anfractibus obtusè convexis, undique minutiē striatis et malleatis, anfractu ultimo infernē exiliter angulato; aperturā pyriformi-ovatā, labro nitente-nigro.*

THE NEGRO PALUDINA. Shell globosely ovate, subcylindrical, thin, semitransparent, greenish-olive, whorls obtusely convex, minutely striated and malleated throughout, last whorl finely angled round the lower part; aperture pyriformly ovate, lip shining-black.

Hab. Central Africa.

An interesting semitransparent dark-green shell, with a conspicuous shining-black lip, of a subcylindrical pupoid form.

Fig. 61. (Mus. Cuming.)

This shell proves to be a variety of Swainson's *P. carinata*, described in the preceding Plate, with a conspicuous shining-black lip.

Species 62. (Mus. Frauenfeld.)

PALUDINA FORMOSULA. *Pal. testā obeso-ovatā, sordidē olivaceā, anfractibus supernē nitidē declivi-angulatis, deinde plano-convexitatis, infernē exiliter angulatis, spiraliiter grano-striatis, striis subdistantibus, umbilico sub-excavato; aperturā ovatā.*

THE BEAUTIFUL PALUDINA. Shell stoutly ovate, dingy-olive, whorls neatly slopingly angled round the upper part, then flatly convex, faintly angled round the lower part, spirally grain-striated, striae rather distant, umbilicus somewhat excavated; aperture ovate.

Vivipara formosula, Frauenfeld, Ueber die Gattung *Vivipara*, November, 1862.

Hab. —?

Chiefly distinguished from *P. Javanica* by its more angular form and more widely separated spiral striae.

Fig. 63. (Mus. Cuming.)

The specimen here figured appears to be one of the United States species; probably *P. lima*.

Species 64. (Mus. Cuming.)

PALUDINA FULVA. *Pal. testā ovato-conicā, solidiusculā, fulvā, anfractibus rotundatis, obscurè minutē decussatim striatis, umbilico obtecto; aperturā pyriformi-ovatā.*

THE FULVOUS PALUDINA. Shell ovately conical, rather solid, fulvous, whorls rounded, obscurely minutely decussately striated, umbilicus covered; aperture pyriformly ovate.

Hab. Camboja; Mouhot.

This shell resembles, in general appearance, the *P. carinata* of Ceylon and Philippine Islands, but there is no keel at the base of the upper whorls, and the striation is different.

Species 65. (Mus. Cuming.)

PALUDINA SUMATRENSIS. *Pal. lestā ovato-conicā, solidusculā, flavescente-olivaceā, lineis nigris conspicue funiculatā, anfractibus supernè declivibus, deinde plano-convexis, ultimo versus basin subangulato-inflato; aperiturā parciusculā.*

THE SUMATRA PALUDINA. Shell ovately conical, rather solid, yellowish-olive, conspicuously corded with black lines, whorls sloping round the upper part, then flatly convex, last whorl rather angularly inflated towards the base; aperture rather small.

DUNKER, Zeitschr. für Malak. 1852, p. 128.

Hab. Sumatra.

The prettily-painted linear bands are smooth in the earlier whorls, and rise into relief with the growth of the shell.

Species 66. (Mus. Cuming.)

PALUDINA PUNCTURATA. *Pal. globoso-conicā, subinflatā, purpureo-olivaceā, ad suturas pallidā, anfractibus converis, supernè subappressis, spiraliter minutissimè punctato-striatis; aperturā ovalā.*

THE PUNCTURED PALUDINA. Shell globosely conical, rather inflated, purple-olive, pale at the sutures, whorls convex, rather appressed round the upper part, spirally very minutely puncture-striated; aperture ovate.

Hab. —?

A dark purple-olive shell, smooth and shining, sculptured with rather distant spiral rows of minute punctures.



67



68a



68b



69



70



71



72



73



74



75

PALUDINA.

PLATE XI.

Fig. 67. (Mus. Cuming.)

The shell here represented is the *Paludina Boissieri*, Charpentier, inhabiting the vicinity of Naples; but according to Frauenfeld, who has lately made an especial study of this Family, it is a *Bithynia*.

Species 68. (Fig. a, b, Mus. Anthony.)

PALUDINA SCALARIS. *Pal. testā subtrigono-ovatā, albā, sinistrā, spirā latē planatā, interdum concavo-immersā, anfractibus longitudinaliter striatis, supernē acutē angulatis, deinde convexis; aperturā oblongā, amplā.*

THE WINDING-STAIR PALUDINA. Shell somewhat triangularly ovate, white, sinistral, spire broadly flattened, sometimes concavely immersed, whorls longitudinally striated, sharply angled round the upper part, then convex; aperture oblong, large.

JAY, Catalogue, pl. 1. f. 8, 9.

Hab. Near Tampa Bay, Florida.

A curious sinistrally coiled shell, with the spire broadly flattened or concavely immersed.

Species 69. (Mus. Anthony.)

PALUDINA REGULARIS. *Pal. testā globoso-ovatā, crassiusculā, nigricante-olivaceā, versus apicem plus minus erosā, spirā brevi, subacutā, anfractibus convexis, lœvibus; aperturā pyriformi-oblongā, subamplā.*

THE REGULAR PALUDINA. Shell globosely ovate, rather thick, blackish-olive, more or less eroded towards the apex, spire short, rather sharp, whorls convex, smooth; aperture pyriformly oblong, rather large.

LEA, Trans. Amer. Phil. Soc., vol. ix. p. 13.

Vivipara regularis, Bimney.

Hab. Illinois and Ohio, United States.

Of a much shorter obovate growth, with a smaller spire than any other United States species.

Species 70. (Mus. Cuming.)

PALUDINA VIRESSENS. *Pal. testā ovato-conicā, subinflatā, pellucido-virescente, obscurē fasciatā, versus apicem rubidā, anfractibus convexis, lœvibus, minutē granostriatis, striis subdistantibus, anfractu ultimo infernē exiliter angulato; aperturā ovatā.*

THE GREENISH PALUDINA. Shell ovately conical, rather inflated, light transparent green, obscurely banded, ruddy towards the apex, whorls convex, smooth, minutely grain-striated, striae rather distant, last whorl delicately angled at the lower part; aperture ovate.
Hab. Macassar.

A delicately-tinted faintly banded shell of the variable *P. Javanica* type.

Species 71. (Mus. Cuming.)

PALUDINA AUSTRALIS. *Pal. testā ovato-conicā, subyellowosā, fusco-olivaceā, lineis fuscouscentibus subdistantibus pallidē cinctā, anfractibus supernē subappressis, deinde rotundatis, undique minutissimē striatis; aperturā rotundato-ovatā.*

THE SOUTHERN PALUDINA. Shell ovately conical, rather globose, fuscous-olive, faintly encircled with rather distant brownish lines, whorls slightly appressed round the upper part, then rounded, everywhere very minutely striated; aperture rotundately ovate.

Hab. Victoria River, North Australia; Dring.

A faintly lineated shell, of peculiar fuscous-olive colour.

Species 72. (Mus. Cuming.)

PALUDINA MAHEYANA. *Pal. testā ovato-conicā, subobliquā, pallidē virescente-olivaceā, rufescente interdum fusciatā, anfractibus supernē declivi-convexis, ad suturas constrictis, longitudinaliter plicato-striatis, interdum conspicuē malleatis; aperturā ovatā.*

THE MAHEY PALUDIN. Shell ovately conical, rather oblique, pale greenish-olive, sometimes banded with red, whorls slopingly convex round the upper part, constricted at the sutures, longitudinally plicately striated, sometimes conspicuously malleated; aperture ovate.

GRATELOUP, MS. in Mus. Cuming.

Hab. Madagascar?

Of a peculiar pale sickly-green colour, sometimes banded with red after the manner of *P. vivipara*.

Species 73. (Mus. Cuming.)

PALUDINA POLITA. *Pal. testā obtusē ovatā, subobliquā,*

*solidā, pallidē olivaceo-cinereā, anfractibus rotundatis,
laevibus, politis; aperturā parvā, rotundato-ovalā.*

THE POLISHED PALUDINA. Shell obtusely ovate, rather oblique, solid, pale olive-ash, whorls rounded, smooth, polished; aperture small, rotundately ovate.

Vivipara polita, Frauenfeld, Gattung *Vivipara*, Nov. 1862.
Hab. South Africa?

The shell here figured is larger and more sharply conical than the true *P. polita*, of which a type specimen has been received from M. Frauenfeld.

Species 74. (Mus. Cuming.)

This appears to be a variety from Nicobar of the Macassar *P. virescens*, Sp. 70.

Species 75. (Mus. Anthony.)

PALUDINA DECAPITATA. *Pal. testā ovato-globosā, pal-
lidē cinereo-virente, spirā erosā, anfractibus plano-
convexis; aperturā pyriformi-ovalā.*

THE DECAPITATED PALUDINA. Shell ovately globose, pale ash-green, spire eroded, whorls flatly convex; aperture pyriformly ovate.

ANTHONY, Pro. Acad. Nat. Sci. Phil. 1860, p. 71.
Hab. Tennessee, United States.

I give Mr. Anthony's original type of this species, kindly forwarded to me for examination, doubting, however, equally with Mr. Binney whether this and *P. regularis* are anything more than varieties of *P. decisa*.

PALUDINA.


	Plate.	Species.		Plate.	Species.
abbreviata, <i>Reeve</i>	V.	26	<i>integra</i> , <i>Say</i>	VIII.	45
achalina, <i>Studer</i>	I.	4	<i>intermedia</i> , <i>Hanley</i>	IX.	57
æruginea, <i>Reeve</i>	VII.	41	<i>intertexta</i> , <i>Say</i>	IV.	15
æthiops, <i>Reeve</i>	X.	60	<i>Japonica</i> , <i>Martens</i>	III.	13
amplior, <i>Monsson</i>	IX.	55	<i>Javanica</i> , <i>Busch</i>	IX.	52
ampullacea, <i>Charpentier</i>	III.	12	<i>lecythoides</i> , <i>Benson</i>	IV.	21
Ampullarioides, <i>Hanley</i>	VI.	30	<i>lima</i> , <i>Anthony</i>	VIII.	48
angularis (<i>Nerita</i>), <i>Müller</i>	III.	14	<i>limosa</i> , <i>Valenciennes</i>	VIII.	45
angulata, <i>Lea</i>	I.	2	<i>lineata</i> , <i>Valenciennes</i>	I.	4
australis, <i>Reeve</i>	XI.	71	<i>lineolata</i> , <i>Mousson</i>	IX.	50
Bengalensis, <i>Lamarck</i>	II.	5	<i>Listeri</i> , <i>Forbes and Hanley</i>	I.	3
Bermondiana, <i>D'Orbigny</i>	V.	28	<i>magnifica</i> , <i>Conrad</i>	IX.	54
bicarinata, <i>Potiez</i> and <i>Michaud</i>	VIII.	44	<i>Maheyana</i> , <i>Grateloup</i>	XI.	72
bimarginata, <i>Lea</i>	IX.	54	<i>mallectata</i> , <i>Reeve</i>	V.	25
Boissieri, <i>Charpentier</i>	XI.	67	<i>melanostoma</i> , <i>Benson</i>	V.	27
Burroughiana, <i>Lea</i>	IV.	16	<i>nitens</i> , <i>Reeve</i>	X.	59
carinata, <i>Swainson</i>	IX.	53	<i>nucleus</i> , <i>Mousson</i>	V.	29
Ceylanica, <i>Dohrn</i>	VI.	32	<i>obtusa</i> , <i>Trochel</i>	VI.	33
Chimensis, <i>Gray</i>	IV.	18	<i>ovularis</i> , <i>Valenciennes</i>	VIII.	45
ciliata, <i>Reeve</i>	VI.	36	<i>oxytropis</i> , <i>Benson</i>	II.	9
coaretata, <i>Lea</i>	VIII.	46	<i>polit</i> , <i>Frauenfeld</i>	XI.	73
conectata (<i>Cyclostoma</i>), <i>Millet</i>	I.	3	<i>polyzonata</i> , <i>Frauenfeld</i>	VII.	38
contorta, <i>Shuttleworth</i>	VIII.	43	<i>ponderosa</i> , <i>Say</i>	VIII.	45
Coosaensis, <i>Lea</i>	V.	22	<i>præmorsa</i> , <i>Benson</i>	VI.	35
cornuta, <i>Valenciennes</i>	VIII.	45	<i>prærosa</i> , <i>Gerstfeldt</i>	III.	10
costata, <i>Quoy</i>	II.	6	<i>puncturata</i> , <i>Reeve</i>	X.	66
crassa, <i>Hutton</i>	VI.	33	<i>pyramidata</i> , <i>Buseh</i>	II.	9
Cumingii, <i>Hanley</i>	III.	11	<i>quadrata</i> , <i>Benson</i>	IV.	17
Cyclostomatiformis, <i>Lea</i>	VIII.	43	<i>regularis</i> , <i>Lea</i>	XI.	69
decapitata, <i>Anthony</i>	XI.	75	<i>Remossii</i> , <i>Philippi</i>	VI.	35
deceisa, <i>Say</i>	VIII.	45	<i>rufa</i> , <i>Haldemann</i>	VIII.	45
dissimilis, <i>Müller</i>	IX.	56	<i>sealaris</i> , <i>Jay</i>	XI.	68
doliaris, <i>Gould</i>	I.	1	<i>sinistrorsa</i> , <i>Lea</i>	IV.	15
Elliotti, <i>Lea</i>	VIII.	43	<i>subcarinata</i> , <i>Say</i>	VIII.	44
elongata, <i>Swainson</i>	II.	5	<i>subpurpurea</i> , <i>Say</i>	VIII.	47
exilis, <i>Anthony</i>	VIII.	46	<i>subsolida</i> , <i>Anthony</i>	VIII.	45
fallax, <i>Frauenfeld</i>	IX.	51	<i>sulculosa</i> , <i>Menke</i>	VIII.	44
fasciata, <i>Deshayes</i>	I.	4	<i>Sumatrensis</i> , <i>Dunker</i>	X.	65
filosa, <i>Hanley</i>	VI.	31	<i>Texana</i> , <i>Tryon</i>	V.	24
formosula, <i>Frauenfeld</i>	X.	62	<i>Thersites</i> , <i>Hanley</i>	IX.	49
fulva, <i>Reeve</i>	X.	64	<i>umbilicata</i> , <i>Lea</i>	VII.	40
genicula, <i>Conrad</i>	VIII.	45	<i>Ussuriensis</i> , <i>Gerstfeldt</i>	II.	8
Georgiana, <i>Lea</i>	I.	4	<i>variata</i> , <i>Frauenfeld</i>	X.	58
gigantea, <i>Busch</i>	II.	7	<i>virescens</i> , <i>Reeve</i>	XI.	70
Hainesiana, <i>Lea</i>	VIII.	42	<i>viridis</i> , <i>Hanley</i>	IV.	20
Hamiltoni, <i>Metcalfe</i>	VI.	37	<i>vicipara</i> , <i>Studer</i>	I.	3
heterostrophia, <i>Kirtland</i>	VIII.	45	<i>vulgaris</i> , <i>Gray</i>	I.	4
inflata, <i>Filla</i>	IV.	19	<i>Wareana</i> , <i>Shuttleworth</i>	V.	23
Ingallsianus, <i>Lea</i>	VII.	39	<i>zonata</i> , <i>Hanley</i>	VI.	34

MONOGRAPH
OF THE GENUS
CYCLOTUS.

"If thou art worn and hard beset
With sorrows that thou wouldst forget,
If thou wouldst read a lesson that will keep
Thy heart from fainting and thy soul from sleep,
Go to the woods and hills! No tears
Dim the sweet look that nature wears." —*Longfellow.*



1a



2a



2b



3a



4a




4b



3b.



5a



7b



5b

C Y C L O T U S.

PLATE 1.

Genus CYCLOTUS, *Guilding.*

Testa plus minus depresso-turbinata vel discoidea, sc̄pissimè latè umbilicata, aufractibus subangustis, aperturā parviusculā, marginibus continuis vel callo sinuato junctis. Operculum orbiculare, testaceum, arctispiram, exstus concurvatus, anfractuum margine in laminam incurvum interdum elevato.

Shell more or less depresso-turbinate, or discoid, mostly broadly umbilicated, whorls rather narrow, aperture rather small, margins continuous, or joined by a sinuated callus. Operculum orbicular, testaceous, closely spired, rather concave outwardly, edge of the whorls sometimes raised into an incurved lamina.

In this division of the Lamarekian genus *Cyclostoma* there are two well-marked groups, each numbering about a dozen species. The species of the first of these, the largest and most conspicuously coloured, are natives of the woods of New Granada, West Columbia, Bolivia and Brazil. The shell is elegantly convoluted in a very discoid manner, largely openly umbilicated, and enveloped with a very characteristic epidermis of a rich burnt-brown, banded with olive-green colour. The species of the second group are natives of Jamaica, of rather smaller size and stouter growth, chiefly of a rose-brown hue, more or less pliately wrinkled. Half-a-dozen species from India and China, and the same number from the Philippine Islands, together with four from New Guinea, two from Bombay, and one from the Lao Mountains, have been referred to *Cyclotus*, but they are much smaller than the *Cycloti* of the Western Hemisphere, and of quite a different typical character. We have a small species of *Cyclotus* from Mauritius, and one also from Natal.

Species 1. (Fig. a, b, Mus. Cuming.)

CYCLOTUS GRANULATUS. *Cycl. testa modice umbilicata, depresso-orbiculari, fulvā, roseo-brunneō spiraliiter latifasciatā, aufractibus subangustis, supernè depresso, undique corrugato-granatis.*

THE GRANULATED CYCLOTUS. Shell moderately umbilicated, depresso-orbicular, fulvous, spirally broadly banded with rose-brown, whorls rather narrow, depressed at the upper part, everywhere wrinkle-grained.

CHITTY, MS. in Mus. Cuming.

Hab. Jamaica.

The last whorl of this species is not excavately angled and ridged around the umbilicus as in its nearest ally *C. Jamaicensis*.

Species 2. (Fig. a, b, Mus. Cuming.)

CYCLOTUS SEMINUDUS. *Cycl. testa angulatum excavato-umbilicata, subdepressa, fulvā, roseo-brunneō spiraliter latifasciatā, aufractibus undique rugatis, rugis partim obsoletis.*

THE SEMINUDUS CYCLOTUS. Shell angularly excavately umbilicated, slightly depressed, fulvous, encircled with a broad rose-brown band, whorls everywhere wrinkled, wrinkles partially obsolete.

Cyclostoma seminudum, C. B. Adams, Contr. Conch. No. viii. p. 143.

Cyclotus Jamaicensis, var., Pfeiffer.

Hab. Jamaica.

Dr. Pfeiffer formerly considered this a distinct species from *C. Jamaicensis*.

Species 3. (Mus. Cuming.)

CYCLOTUS GIGANTEUS. *Cycl. testa ampliter aperte umbilicata, depressa, solidiuscula, astnato-fusco olivaceo-viridi latè fasciata et tincta, ad peripheriam fascia angusta fulvā cingulata, aufractibus angustis, ad suturas plano-impressis, supra concentricè filo-striatis, infra corrugato-striatis.*

THE GIGANTIC CYCLOTUS. Shell largely openly umbilicated, depressed, rather solid, burnt-brown, broadly banded and stained with olive-green, encircled with a narrow fulvous band at the periphery, whorls narrow, flatly impressed at the sutures, concentrically thread-striated above, wrinkle-striated below.

Cyclostoma giganteum, Sowerby, Pro. Zool. Soc. 1843, p. 30.

Cyclotus gigantens, Gray.
Hab. Panama (in the neighbouring woods).

This fine species should be regarded as the type of the genus.

CYCLOTUS.—PLATE I.

Species 4. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS LAXATUS. *Cycl. testā latissimē apertē umbili-
catā, subdiscoideā, intensē olivaceo-viridi, versus apicem
ustulato-fuscescente, ad peripheriam angustē fasciatā,
anfractibus perangustis, supernē densē reticulato-rugatis
et malleatis, infra concentricē corrugato-striatis.*

THE LOOSELY-COILED CYCLOTUS. Shell very broadly openly umbilicated, nearly discoid, dark olive-green, burnt-brown towards the apex, narrowly banded at the periphery, whorls very narrow, densely reticulately wrinkled and malleated at the upper part, concentrically wrinkle-striated below.

Cyclostoma laxatum, Sowerby, Thes. Conch. Supp. p. 159.
pl. 31 A. f. 302.

Cyclotus laxatus, Sowerby.

Hab. Columbia.

Characterized by an unusually discoid growth, and an exquisitely reticulated and malleated sculpture.

Species 5. (Mus. Cuming.)

CYCLOTUS JAMAICENSIS. *Cycl. testū angulatim excavato-
umbilicatā, subdepressā, fulvā, roseo-brunneo lineare
multifasciatā, anfractibus declivi-convexis, dense ob-
liquè striatis, ad suturas plicato-malleatis.*

THE JAMAICA CYCLOTUS. Shell angularly excavately umbilicated, slightly depressed, fulvous, linearly many-banded with rose-brown, whorls slopingly convex, densely obliquely striated, pliately malleated at the sutures.

Turbo Jamaicensis, Chemnitz, vol. ii. p. 277. pl. 209.
f. 2057.

Cyclostoma Jamaicense, Gray.

Cyclotus lineatus, Gray.

Hab. Jamaica.

An extremely variable species, sometimes smooth, often more or less wrinkled.



7a



6a



7b




8a



9a



9b.



6b



10a



10b



6b.

C Y C L O T U S.

PLATE II.

Species 6. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS JUGOSUS. *Cycl. testā apertē umbilicatā, depresso, sub epidermide fulvā albā, anfractibus quinque, subangustis, concentricè filoso-liratis, ruderiter nodoso-corrugatis, rugis infra obliquè continuis.*

THE RUGGED CYCLOTUS. Shell openly umbilicated, depressed, white, beneath a fulvous epidermis, whorls four to five, rather narrow, concentrically threadridged, rudely nodosely wrinkled, wrinkles obliquely continuous beneath.

Cyclostoma jugosum, C. B. Adams, Contr. Conch. n. 8. p. 143.

Cyclotus jugosus, Pfeiffer.

Hab. Jamaica.

A bright fulvous shell, peculiarly strongly nodosely wrinkled, the wrinkles falling at the base into oblique furrows.

Species 7. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS PALLESCENS. *Cycl. testā angulatè excavato-umbilicatā, subdepressā, solidiusculā, ustulato-fulvā, versus apicem rosaceā, anfractibus quinque, prope suturam planatis, ad basin angulatis, concentricè filoso-striatis, obliquè validè ruderiter corrugatis.*

THE PALE CYCLOTUS. Shell angularly excavately umbilicated, moderately depressed, rather solid, burnt-fulvous, tinged with rose towards the apex, whorls five, flattened next the suture, angled at the base, concentrically thread-striated, obliquely strongly rudely wrinkled.

Cyclostoma pallescens, C. B. Adams, Contr. Conch. n. 8. p. 143.

Hab. Jamaica.

Of a peculiar burnt-fulvous colour, tinged with rose at the apex. The sculpture consists mainly of coarse oblique wrinkles.

Species 8. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS INCOMPTUS. *Cycl. testā ampliter apertē umbilicatā, depressā, subdiscoidē, sub epidermide fuscā corneā deciduā albā, ad peripheriam saturatè fasciatā, anfractibus concentricè ruderiter striatis, ultimo supernè ad aperturam dilatato-alato.*

THE ROUGH CYCLOTUS. Shell largely openly umbilicated, depressed, somewhat discoid, white, beneath a

deciduous brown horny epidermis, darkly banded at the periphery, whorls concentrically rudely striated, the last dilately winged at the upper part of the aperture.

Cyclostoma incomptum, Sowerby, Thes. Conch. p. 160. pl. 31. f. 298, 299.

Pterocyclos incomplus, Pfeiffer.

Cyclotus incomptus, Pfeiffer.

Hab. Brazil.

This fine species possesses a modification of the sinuated wing-like development of *Pterocyclos*.

Species 9. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS DUNKERI. *Cycl. testā subprofundè umbilicatā, conico-depressā, fulvescente-fuscā, infra peripheriam saturatè latè fasciatā, anfractibus ad suturas concavo-impressis, striis elevatis undique densè concentricè fusciculatis.*

DUNKER'S CYCLOTUS. Shell rather deeply umbilicated, conically depressed, fulvous-brown, broadly dark-banded below the periphery, whorls conceavely impressed at the sutures, concentrically densely corded throughout with raised striae.

PFEIFFER, Malak. Blatt. 1856, p. 256.

Hab. New Granada.

A beautifully-sculptured shell, of a bright fulvous-brown colour, rather broadly dark-banded below the periphery.

Species 10. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS INCA. *Cycl. testā subprofundè umbilicatā, subdepressā, viridè olivaceo-virente, ad peripheriam fulvescente, deinde saturatā, anfractibus ad suturas impresso-concavis, undique concentricè filoso-striatis.*

THE INCA CYCLOTUS. Shell rather deeply umbilicated, moderately depressed, bright olive-green, yellow at the periphery, then darker, whorls impressly concave at the sutures, concentrically thread-striated throughout.

Cyclostoma Inca, D'Orbigny, Voy. Amér. Mérid. p. 361. pl. 16. f. 21 to 23.

Cyclotus Inca, Gray.

Hab. Bolivia; La Guayra.

Chiefly recognizable by its bright olive-green banded colouring.



11a



12a



12b



13a



14a



14b



13b



15a



15b



11b

C Y C L O T U S.

PLATE III.

Species 11. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS CORRUGATUS. *Cycl. testā angulatē excavato-umbilicatā, subdepressā, solidā, fulvo-fuscescente, unfractibus validissimè undulato-corrugatis, ad basin angulatis et crassè marginatis.*

THE WRINKLED CYCLOTUS. Shell angularly excavately umbilicated, slightly depressed, solid, fulvous-brown, whorls very strongly undulately wrinkled, angled and thickly margined at the base.

Cyclostoma corrugatum, Sowerby, Pro. Zool. Soc. 1843, p. 30.

Cyclotus corrugatus, Pfeiffer.

Hab. Jamaica.

The wrinkled sculpture in this species has a conspicuous waved character, varying, however, very much in strength.

Fig. 12. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS VARIANS. For description of this extremely variable species see Plate VII., where a more characteristic specimen of it is figured.

Species 13. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS CINGULATUS. *Cycl. testā modice profundē umbilicatā, subdepressā, usculatē olivaceo-fuscā, ad peripheriam saturatē fasciatā, anfractibus ad suturas concavo-impressis, deinde declivi-convexis, concentricè filoso-striatis, obliquè reticulato-incisis.*

THE GIRDLED CYCLOTUS. Shell moderately deeply umbilicated, rather depressed, burnt olive-brown, dark-banded at the periphery, whorls concavely impressed at the sutures, then slopingly convex, concentrically thread-striated, obliquely reticulately engraved.

Cyclostoma cingulatum, Sowerby, Thes. Conch. part ix. p. 93. pl. 29. f. 213, 214.
Cyclotus cingulatus, Gray.

Hab. New Granada.

This beautiful species is distinguished by a sloping convexity of the whorls, which are sculptured by a loose oblique network of superficially-engraved grooves.

Species 14. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS QUITENSIS. *Cycl. testā subapertè umbilicatā, modicè depressā, vividè fulvescente-castaneā, versus apicem rosaceā, ad peripheriam pallidè fasciatā, anfractibus ad suturas concavo-impressis, concentricè rudē striatis, striis subtus validis.*

THE QUITO CYCLOTUS. Shell rather openly umbilicated, moderately depressed, bright fulvous-chestnut, rose towards the apex, faintly banded at the periphery, whorls concavely impressed at the sutures, concentrically rudely striated, striae strong beneath.

Cyclostoma Quiteense, Pfeiffer, Pro. Zool. Soc. 1851.

Cyclotus Quiteensis, Pfeiffer.

Hab. Quito.

A shell of rather solid growth, pink about the apex, with the striae much increasing in strength towards the umbilicus.

Species 15. (Fig. *a, b*, Mus. Cuming.)

CYCLOTUS CRASSUS. *Cycl. testā angulatē excavato-umbilicatā, crassā, sordidè fuscā, anfractibus declivi-convexis, supernè rudē corrugatā et malleatā.*

THE THICK CYCLOTUS. Shell angularly excavately umbilicated, thick, dull-brown, whorls slopingly convex, rudely wrinkled and malleated at the upper part.

Cyclostoma crassum, C. B. Adams, Contr. Conch. n. viii. p. 143.

Hab. Jamaica.

A small species, possessing the typical character of the wrinkled Jamaica forms obscurely developed.




16



17a



17b.



18a.



19a



19b.



18b



20a



20b



21

C Y C L O T U S.

PLATE IV.

Species 16. (Mus. Cuming.)

CYCLOTUS SEMISTRIATUS. *Cycl. testā profundē subangustē umbilicatā, depressa-canicā, albida, rufescente tinctā, aurantio-fusco interdum vittatā, anfractibus subangustis, superne spiraler liratis.*

THE HALF-STRIPED CYCLOTUS. Shell deeply rather narrowly umbilicated, depressedly conical, white, tinged with red, sometimes filleted with orange-brown, whorls rather narrow, spirally ridged round the upper part.

Cyclotoma semistriatum, Sowerby, Pro. Zool. Soc. 1843, p. 29.

Cyclotus semistriatus, Gray.

Hab. India.

Of quite a different type from the Western species of the genus. The whorls have neither concentric striae nor wrinkles, but are sculptured round the upper part with spiral ridges, while the lower part is smooth.

Species 17. (Fig. a, b, Mus. Cuming.)

CYCLOTUS FORTUNEI. *Cycl. testā subamplicher profundē umbilicatā, fulvescente-fuscā, sericeā, interdum obscurè fasciatā, anfractibus angustis, rotundatis, concentricè tenuistriatis.*

FORTUNE'S CYCLOTUS. Shell rather largely deeply umbilicated, fulvous-brown, silky, sometimes obscurely banded, whorls narrow, rounded, concentrically finely striated.

Cyclotoma Fortunei, Pfeiffer, Pro. Zool. Soc. 1852, p. 146.

Cyclotus Fortunei, Pfeiffer.

Hab. Shanghai, China; Fortune.

A small silky shell of narrow rounded whorls, with much of the aspect of a *Helix*.

Species 18. (Fig. a, b, Mus. Cuming.)

CYCLOTUS IRREGULARIS. *Cycl. testā modicè profundē umbilicatā, subdepressā, crassā, pallidē fulvescente-castaneā, infra peripheriam saturatē fasciatā, anfractibus ad suturas concavo-impressis, striis elevatis concentricis undique densè funiculatis.*

THE IRREGULAR CYCLOTUS. Shell moderately deeply

umbilicated, rather depressed, thick, pale fulvous-chestnut, dark-banded below the periphery, whorls concavely impressed at the sutures, densely corded throughout with raised concentric striae.

Cyclotoma (Cyclotus) irregularis, Pfeiffer, Pro. Zool. Soc. 1855, p. 117.

Hab. Costa Rica, Central America.

Pale in colour, of rather solid growth, beautifully corded throughout with raised concentric striae.

Species 19. (Fig. a, b, Mus. Cuming.)

CYCLOTUS VOLVULOIDES. *Cycl. testā modicè profundē umbilicatā, subdepressā, albida, purpureo-violacea obscurè fasciatā, anfractibus subangustis, rotundatis, spiraler liratis, ad peripheriam laevigatis, labro callosoreflexo.*

THE WHORL-LIKE CYCLOTUS. Shell moderately deeply umbilicated, slightly depressed, whitish, obscurely banded with purple-violet, whorls rather narrow, rounded, spirally ridged, smooth at the periphery, lip callously reflected.

Cyclotoma volvuloides, Sowerby, Thes. Couch. p. 162. pl. 31 b. f. 313, 314.

Cyclatus volvuloides, Pfeiffer.

Hab. —?

This is a very characteristic species of the Eastern type represented by the smaller Indian *C. semistriatus*. It has no concentric sculpture, the ridges being of spiral growth.

Species 20. (Fig. a, b, Mus. Cuming.)

CYCLOTUS BOGOTENSIS. *Cycl. testā modicè profundē umbilicatā, conico-depressā, vividè virente varie fasciatā, anfractibus ad suturam concavo-impressis, deinde rotundatis, striis concentricis undique densè funiculatis.*

THE BOGOTA CYCLOTUS. Shell moderately deeply umbilicated, conically depressed, bright green, variously banded, whorls concavely impressed at the suture, then rounded, densely corded throughout with concentric striae.

Cyclotoma Bogotense, Pfeiffer, Pro. Zool. Soc. 1855, p. 117.

Hab. New Granada.

CYCLOTUS.—PLATE IV.

Scarcely distinguishable from *C. Inca*, and probably a variety of that species.

— — — — —
Species 21. (Mus. Cuming.)

CYCLOTUS SUBDISCOIDEUS. *Cycl. testā latē profundē umbilicatā, depressā, lacteā, aurantio-carneo lineatā et tinctā, anfractibus angustis, rotundatis, ad suturas excavato-impressis, deinde liris spiralibus validē funiculatis, labro calloso-reflexo.*

THE SLIGHTLY-DISCOID CYCLOTUS. Shell broadly deeply umbilicated, depressed, cream color, lineated and tinged with orange-flesh, whorls narrow, rounded, excavately impressed at the suture, then strongly corded with spiral ridges, lip callously reflected.

Cyclostoma subdiscoideum, Sowerby, Thes. Conch. p. 161.
pl. 31 b. p. 304, 305.

Hab. —?

Another characteristic species of the spirally-ridged Eastern type.



22.



23a



23b



27a



25a



25b



27b



26a



26b



27.

C Y C L O T U S.

PLATE V.

Species 22. (Mus. Cuming.)

CYCLOTUS SUBSTRIATUS. *Cycl. testā subampliter umbilicatā, depressā, fulvā, anfractibus angustis, rotundatis, ad suturas impressis et plicato-striatis.*

THE SLIGHTLY-STRIATED CYCLOTUS. Shell rather largely umbilicated, depressed, fulvous, whorls narrow, rounded, impressed and plicately striated at the sutures.

Cyclostoma substriatum, Sowerby, Pro. Zool. Soc. 1843, p. 61.

Cyclotus substriatus, Gray.

Hab. Island of Siquijor, Philippines; Cuming.

Sculptured with concentric striae, which are more prominent and pucker'd next the sutures.

Species 23. (Fig. a, b, Mus. Cuming.)

CYCLOTUS DISCOIDEUS. *Cycl. testā latissimè ampliter umbilicatā, plano-depressā, subdiscoideā, pallidē fulvescente, subpellucidā, anfractibus angustis, rotundatis, parum striatis, labro continuo, reflexo.*

THE DISCOID CYCLOTUS. Shell very broadly largely umbilicated, flatly depressed, somewhat discoid, light fulvous, somewhat transparent, whorls narrow, rounded, but little striated, silky, lip continuous, reflected.

Cyclostoma discoideum, Sowerby, Thes. Conch. p. 111. pl. 25. f. 87, 88.

Cyclotus discoideus, Gray.

Hab. Demerara; Sowerby.

A narrow-whorled subdiscoidal shell, with a delicate silky surface.

Species 24. (Fig. a, b, Mus. Cuming.)

CYCLOTUS POPAYANUS. *Cycl. testā modicè umbilicatā, subdepressā, sub epidermide fulvā deciduā albā, versus apicem rosaceā, ad peripheriam fasciatā, anfractibus superne concarinseculis, deinde declivi-rotundatis, concentricè densè filoso-striatis.*

THE POPAYAN CYCLOTUS. Shell moderately umbilicated, slightly depressed, white beneath a deciduous fulvous epidermis, rose towards the apex, banded at the periphery, whorls slightly concurved round the upper part, then slopingly rounded, concentrically densely thread-striated.

Cyclostoma Popayanum, Lea, Obs. vol. ii. p. 94. pl. 23. f. 76.

Cyclostoma inconspicuum, Sowerby.

Cyclotus inconspicuus, Gray.

Cyclotus Popayanus, Pfeiffer.

Hab. Popayan, New Granada; Gibbon. Chimborazo; Linden.

The whorls of this species, rather slopingly inflated, are characteristically tinged with rose towards the apex.

Species 25. (Fig. a, b, Mus. Cuming.)

CYCLOTUS STRAMINEUS. *Cycl. testā modicè profundè umbilicatā, subdepressā, eximiè stramineā, anfractibus declivi-rotundatis, pulcherrimè diagonaliter crispato-corrugatis.*

THE STRAW-COLOURED CYCLOTUS. Shell moderately deeply umbilicated, rather depressed, delicate straw-colour, whorls slopingly rounded, very beautifully diagonally crisply wrinkled.

Cyclostoma straminenm, Reeve, Pro. Zool. Soc. 1843, p. 46.

Cyclotus straminens, Gray.

Hab. Merida, West Columbia.

Remarkable for the semitransparent delicacy of its colour and curious diagonally-wrinkled sculpture.

Fig. 26. (Fig. a, b, Mus. Cuming.)

This shell, which is Mr. Sowerby's *C. subrugosus*, is merely a variety of Professor C. B. Adams's *C. crassus*.

Species 27. (Mus. Cuming.)

CYCLOTUS MUCRONATUS. *Cycl. testā subampliter apertè umbilicatā, pallidē luteā, rufescente tinctā, apice submucronato, anfractibus angustis, rotundatis, ad suturas concentricè plicato-striatis, sub lente superficialiter spiraliter striatis, labro reflexo.*

THE POINTED CYCLOTUS. Shell rather largely openly umbilicated, pale yellow, tinged with red, apex rather pointed, whorls narrow, rounded, plicately concentrically striated at the sutures, faintly spirally striated beneath the lens, lip reflected.

Cyclostoma mucronatum, Sowerby, Pro. Zool. Soc. 1843, p. 63.

Cyclotus mucronatus, Gray.

Hab. Island of Luzon, Philippines; Cuming.

Pucker'd with concentric striae next the suture, while the general surface of the whorls is faintly spirally striated.



28



29a



29b



30a



31b



34a



30b



32a



33b



33.

C Y C L O T U S.

PLATE VI.

Species 28. (Mus. Cuming.)

CYCLOTUS PROMINULUS. *Cycl. testā ampliter apertē umbilicatā, subdepressā, albida, anfractibus angustis, densè concentricè filoso-striatis, subtubulosis, ultimo plus minus soluto.*

THE PRODUCED CYCLOTUS. Shell largely openly umbilicated, rather depressed, whitish, whorls narrow, densely concentrically thread-striated, subtubular, the last more or less unrolled.

Cyclostoma prominulum, Féruccac; D'Orbigny, Voy. Amér. Mérid. p. 362.

Cyclostoma Brasiliense, Sowerby.

Cyclotus Brasiliensis, Sowerby.

Cyclotus prominulus, Pfeiffer.

Hab. Rio Janeiro, Minas Geraes; Brazil.

A delicate subtubular shell, in which the last whorl is partially uncoiled.

Species 29. (Fig. a, b, Mus. Cuming.)

CYCLOTUS VARIEGATUS. *Cycl. testā latissimè apertē umbilicatā, plano-discoideā, fulvescente, castaneo-brunneo reticulato-pictā, anfractibus angustis, rotundatis, concentricè densè filoso-striatis, labro reflexo, supernè subalato.*

THE VARIEGATED CYCLOTUS. Shell very broadly openly umbilicated, flatly discoid, fulvous, reticulately painted with chestnut-brown, whorls narrow, rounded, concentrically densely thread-striated, lip reflected, slightly winged at the upper part.

SWAINSON, Treat. Malacol. p. 336.

Cyclostoma planorbolum, Sowerby (not of Lamarck).

Cyclostoma cornu-venatorium, Petit.

Hab. Philippine Islands; Cuming.

This species partakes of the characters of *Cyclophorus* and even of *Pterocyclos*.

Species 30. (Fig. a, b, Mus. Cuming.)

CYCLOTUS TRANSLUCIDUS. *Cycl. testā angustè umbilicatā, subconoideā, pallidè luteā, fulvo interdum multi-fasciatā, anfractibus declivi-rotundatis, densè concentricè rugoso-striatis.*

THE TRANSPARENT CYCLOTUS. Shell narrowly umbili-

cated, somewhat conoid, light yellow, sometimes many-banded with fulvous, whorls slopingly rounded, densely concentrically wrinkle-striated.

Cyclostoma translucidum, Sowerby, Pro. Zool. Soc. 1813, p. 29.

Cyclotus translucidus, Gray.

Hab. Columbia.

A pale-yellow semitransparent shell, sometimes many-banded with fulvous, in which the whorls are more closely convoluted than usual, and the shell is accordingly more narrowly umbilicated.

Species 31. (Fig. a, b, Mus. Cuming.)

CYCLOTUS DYSONI. *Cycl. testā profundè modicè umbilicatā, depresso-conoideā, luteā, fulvo strigatā et fasciatā, anfractibus prope suturas subplanatis, deinde declivi-rotundatis, undique striis elevatis concentricis decussatim undato-corrugatis.*

DYSON'S CYCLOTUS. Shell deeply moderately umbilicated, depresso-conoid, yellow, streaked and banded with fulvous, whorls rather flat next the sutures, then slopingly rounded, everywhere decussately wave-wrinkled with concentric raised striae.

Cyclostoma Dysoni, Pfeiffer, Pro. Zool. Soc. 1851, p. 243.

Cyclophorus? Dysoni, Pfeiffer.

Cyclotus Dysoni, Pfeiffer.

Hab. Honduras; Dyson.

The concentric thread-like striae are strongly wrinkled in this species and incline to cross one another.

Species 32. (Fig. a, b, Mus. Cuming.)

CYCLOTUS HEBRAICUS. *Cycl. testā modicè profundè umbilicatā, depresso, albida, castaneo-brunneo fasciatā et plus minus variegatā, anfractibus subangustis, rotundatis, levigatis, vel minutissimè concentricè striatis, labro tenuè dilatato.*

THE HEBREW CYCLOTUS. Shell moderately deeply umbilicated, depressed, whitish, banded and more or less variegated with chestnut-brown, whorls rather narrow, rounded, smooth, or very minutely concentrically striated, lip thinly dilated.

Valvata Hebraicus, Lesson, Voy. de la Coquille, p. 347.
pl. 13. f. 8.

CYCLOTUS.—PLATE VI.

Cyclostoma papua, Quoy.

Cyclostoma distomella, Sowerby.

Cyclotus Hebraicus, Pfeiffer.

Hab. New Guinea.

As in the case of *C. variegatus*, this species partakes very much of the generic character of *Cyclophorus*.

—
Species 33. (Mus. Cuming.)

CYCLOTUS ASPERULUS. *Cycl. testá modicè profundè umbilicatá, depressá, fulvo-fuscescente, anfractibus superne subconcavis, deinde declivi-rotundalis, concentricè pliato-striatis et obscurè retuse corrugatis.*

THE ROUGH CYCLOTUS. Shell moderately deeply umbilicated, depressed, fulvous-brown, whorls slightly concave round the upper part, then slopingly rounded, concentrically plicately striated and obscurely bluntly wrinkled.

Cyclostoma rugulosum, Sowerby (not of Pfeiffer), Pro. Zool. Soc. 1843, p. 29.

Cyclostoma asperulum, Sowerby.

Cyclotus asperulus, Gray.

Hab. Jamaica.

The wrinkling of this species is blunt and obscure, the interstices having the appearance of punctures.



34



35a



35b



36a



37a



37b



36b



38a



38b



39

C Y C L O T U S.

PLATE VII.

Species 34. (Mus. Cuming.)

CYCLOTUS DISTINCTUS. *Cycl. testū latē apertē umbilicatā, discoideā, sub epidermide oliraceo-luteā albida, anfractibus angustis, tubulosis, spiraliter tenuiliratis, liris regularibus, alternis interdum minoribus.*

THE DISTINCT CYCLOTUS. Shell broadly openly umbilicated, discoid, whitish, beneath an olive-yellow epidermis, whorls narrow, tubular, spirally finely ridged, ridges regular, the alternate ones sometimes smaller.

Cyclostoma distinctum, Sowerby, Thes. Conch. p. 106. pl. 24. f. 38.

Cyclotus distinctus, Pfeiffer.

Hab. West Columbia; Cuming. Haiti; Sallé.

A bright *Planorbis*-like species, regularly spirally ridged throughout.

Species 35. (Fig. a, b. Mus. Cuming.)

CYCLOTUS GLAUCOSTOMA. *Cycl. testā modice subprofundē umbilicatā, depresso-conoideā, ustulato-fuscā, subtus fulvescente, anfractibus declivi-rotundatis, spiraliter decussatim corrugatis.*

THE BLUE-MOUTH CYCLOTUS. Shell moderately rather deeply umbilicated, depresso conoid, burnt-brown, fulvous beneath, whorls slopingly rounded, spirally decessantly wrinkled.

Cyclostoma (Cyclotus) glaucostoma, Pfeiffer, Pro. Zool. Soc. 1855, p. 103.

Hab. Venezuela.

The wrinkled sculpture of this shell runs in the spiral direction in a decussating manner, of quite a different type from the concentric wrinkling of the Jamaica species.

Species 36. (Fig. a, b. Mus. Cuming.)

CYCLOTUS VARIANS. *Cycl. testā angulatē excavato-umbilicatā, erassā, ustulato-fuscā, rufō tintatā, anfractibus declivi-rotundatis, superne plicato-rugosis et malleatis, obsoletē spiraliter liratis, aperture fance rufā.*

THE VARYING CYCLOTUS. Shell angularly excavately umbilicated, thick, burnt-brown, tinged with red, whorls slopingly rounded, plicately rugose and malleated round the upper part, obsoletely spirally ridged, interior of the aperture red.

Cyclostoma varians, Contr. Coneh. no. 8. p. 143.

Cyclotus varians, Pfeiffer.

Hab. Jamaica.

A shell of stout growth, rudely plicated and malleated next the sutures, with faint indications of spiral ridges beneath.

Species 37. (Fig. a, b. Mus. Cuming.)

CYCLOTUS BLANCHETIANUS. *Cycl. testā umpliter apertē umbilicatā, depressā, subdiscoideā, ustulato-fuscā, ad peripheriam pallidā, deinde castaneo latifasciatā, anfractibus ad suturas concavo-impressis, undique dense concentricè filoso-striatis.*

BLANCHET'S CYCLOTUS. Shell largely very openly umbilicated, depressed, somewhat discoid, burnt-fulvous, pale, and then broadly chestnut-banded at the periphery, whorls coneavely impressed at the sutures, densely concentrically thread-striated throughout.

Cyclostoma Blanchetianum, Morieand, Mém. Soc. Phys. Genève, vol. vii. p. 442. pl. 2. f. 21 to 23.

Cyclotus Inca, var., Pfeiffer.

Hab. Brazil.

This differs from the typical form of *C. Inca* in being more thinly flatly coiled, while it is uniformly of a different colour.

Species 38. (Fig. a, b. Mus. Cuming.)

CYCLOTUS SPURCUS. *Cycl. testā subangustē umbilicatā, subgloboso-conoideā, sordidē fulvescente-albā, anfractibus subcontractis, declivi-rotundatis, plus minus obscurè concentricè filoso-striatis.*

THE DIRTY CYCLOTUS. Shell rather narrowly umbilicated, subglobosely conoid, dull fulvous-white, whorls rather contracted, slopingly rounded, more or less obscurely concentrically thread-striated.

Cyclostoma spurcum, Grateilou, Act. Soc. Linn. Bordeaux, vol. ii. p. 444. pl. 3. f. 18.

Otopoma spurcum, Gray.

Cyclotus spurcus, Pfeiffer.

Hab. Malabar, Bombay.

This is the first Indian *Cyclotus* that has come before me in which the characters partake of those of the Western

CYCLOTUS.—PLATE VII.

species. It comes very near to the Columbian *C. translucidus* of the preceding plate.

Species 39. (Mus. Cuming.)

CYCLOTUS PUSILLUS. *Cycl. testā latissimē apertē umbilicatā, subdiscoideā, pallidē fulvescente-fuscā, anfractibus subangustis, prope suturas impressis et tenuē platicatis, deinde rotundatis, tævibus, labro expanso-reflexo.*

THE SMALL CYCLOTUS. Shell very broadly openly um-

bilicated, somewhat discoid, pale fulvous-brown, whorls rather narrow, impressed and faintly plicated next the sutures, then rounded, smooth, lip expandedly reflected.

Cyclostoma pusillum, Sowerby, Pro. Zool. Soc. 1843, p. 59.

Cyclotus pusillus, Gray.

Hab. Philippine Islands; Cuming.

Belonging to the *Cyclophorus* form of the genus, widely openly convoluted.

Cydetus Pl. VIII.



40.



41.



42.



43.



44.



45.



46.



47.



48.



49.

(Moderately enlarged.)

C Y C L O T U S.

PLATE VIII.

Species 40. (Mus. Cuming.)

CYCLOTUS SUTURALIS. *Cycl. testā perampliter discoideā, lutescente-fuscā, anfractibus levigatis, angustis, ad suturam conspicuē lirato-marginatis.*

THE SUTURAL CYCLOTUS. Shell very largely umbilicated, discoid, yellowish-brown, whorls smooth, narrow, conspicuously ridge-margined at the suture.

Cyclostoma suturale, Sowerby, Pro. Zool. Soc. 1843, p. 29.

Aperostoma suturale, Pfeiffer.

Cyclotus suturalis, Gray.

Hab. Jamaica, Demerara.

Characterized by a distinct marginal ridge next the suture.

Species 41. (Mus. Cuming.)

CYCLOTUS DUBIOSUS. *Cycl. testā subampliter umbilicata, helicoideā, sordidē albā, epidermide tenui deciduā induitā, anfractibus subangustis, undique transversim filo-striatis.*

THE DOUBTFUL CYCLOTUS. Shell rather largely umbilicated, helicoid, dull white, covered with a thin deciduous epidermis, whorls rather narrow, transversely thread-striated throughout.

Cyclostoma dubiosum, C. B. Adams, Contr. to Conch., no. ix. p. 157.

Cyclotus dubiosus, Pfeiffer.

Hab. Jamaica.

Closely sculptured throughout with thread-like striae.

Species 42. (Mus. Cuming.)

CYCLOTUS ORBELLUS. *Cycl. testā subampliter umbilicata, subdiscoideā, aurantio-fulgā, diaphanā, anfractibus angustis, ad suturam canaliculato-impressis, spiraliter filo-liratis, liris distantibus.*

THE LITTLE ORB CYCLOTUS. Shell rather largely umbilicated, somewhat discoid, orange-fulvous, diaphanous, whorls narrow, canaliculately impressed at the suture, spirally thread-ridged, ridges distant.

Cyclostoma orbella, Lamarck, Anim. sans vert. vol. viii. p. 360.

Cyclostoma variabile, Féussac.

Species 43. (Mus. Cuming.)

Cyclostoma Lamarekii, Petit.

Cyclotus orbellus, Gray.

Hab. Madagascar, Guadalupe?

A delicate spirally corded semi-transparent shell, partaking somewhat of the generic character of *Cyclophorus*.

Species 44. (Mus. Cuming.)

CYCLOTUS SUBFLAMMULATUS. *Cycl. testā modicē umbilicata, fulvescente-corneā, fusco pallidē flammatā, anfractibus declivi-concavis, lēvibus, aperturā tumidiusculā, labro tenuē reflexo.*

THE FAINTLY-FLAMED CYCLOTUS. Shell moderately umbilicated, fulvous horny, faintly flamed with brown, whorls slopingly convex, smooth, aperture rather swollen, lip thinly reflected.

PFEIFFER, MS. in Mus. Cuming.

Hab. —?

A semi-transparent shell with a faint indication of radiating flames.

Species 45. (Mus. Cuming.)

CYCLOTUS PATERA. *Cycl. testā ampliter umbilicata, discoideā, livido-corneā, epidermide intensē fuscā biseriatim cristatā induitā, anfractibus angustis, spiraliter lirato-striatis.*

THE GOBLET CYCLOTUS. Shell largely umbilicated, discoid, livid-horny, covered with a dark-brown biserially tufted epidermis, whorls narrow, spirally ridge-striated.

PFEIFFER, MS. in Mus. Cuming.

Hab. Lao Mountains; Mouhot.

The chief peculiarity of this species consists in its double-tufted epidermis.

Species 46. (Mus. Cuming.)

CYCLOTUS LINDSTEDTI. *Cycl. testā subampliter umbilicata, lutescente, castaneo-fusco profusē acutē undulati, anfractibus rotundatis, transversim filo-striatis.*

CYCLOTUS.—PLATE VIII.

LINDSTEDT'S CYCLOTUS. Shell rather largely umbilicated, yellowish, profusely sharply waved with chestnut-brown, whorls rounded, transversely thread-striated.

Cyclotoma (Cyclotus) Lindstedti, Pfeiffer, Pro. Zool. Soc. 1856, p. 391.

Cyclotus Lindstedti, Pfeiffer.

Hab. Mount Ophir, Malacca; Lindstedt.

Beautifully painted with sharply zigzag lines and flames.

Species 46. (Mus. Cuming.)

CYCLOTUS BATCHIANENSIS. *Cycl. testā ampliter apertē umbilicatā, lutescente-corneā, ustulato-castaneo copiosē tinctā et flammatā, anfractibus exiliter spiraliter lirato-striatis, ad suturam subimpressis.*

THE BATCHIAN CYCLOTUS. Shell largely openly umbilicated, yellowish-horny, copiously stained and flamed with burnt-chestnut, whorls faintly spirally ridge-striated, rather impressed at the suture.

PFEIFFER, MS. in Mus. Cuming.

Hab. Island of Batchian, Indian Ocean.

This shell has very much the character of a *Cyclophorus*.

Species 47. (Mus. Cuming.)

CYCLOTUS ILLOTUS. *Cycl. testā apertē umbilicatā, versus aperturam laxē convolutā, solidiusculā, sordidē albā, spirā subimmersā, anfractibus transversim corrugato-striatis.*

THE DIRTY CYCLOTUS. Shell openly umbilicated, loosely coiled towards the aperture, rather solid, dull white, spire rather immersed, whorls transversely wrinkled-striated.

GOULD, Otia Conch. p. 104.

Hab. Loo Choo.

A rather loosely coiled shell, striated across in a rough obtusely wrinkled manner.

Species 48. (Mus. Cuming.)

CYCLOTUS FLOCCOSUS. *Cycl. testā excavatē perspectivē umbilicatā, subdiscoideā, pellucido-stramineā, anfractibus ad suturas lirato-marginatis, undique spiraliter filo-liratis, liris granatis.*

THE FLAKED CYCLOTUS. Shell excavately perspectively umbilicated, slightly discoid, transparent straw-colour, whorls ridge-margined at the sutures, spirally thread-ridged throughout, ridges grained.

Cyclotoma (Cyclotus) floccosus, Shuttleworth, Journ. Conch. vol. iii. p. 268.

Cyclotus distinctus, pars, Pfeiffer.

Cyclotus floccosus, Pfeiffer.

Hab. Island of Haiti; Sallé.

The operculum of this beautiful little species is circumlaterally ridged, the ridges being furnished with curious flakes of epidermis.

Species 49. (Mus. Cuming.)

CYCLOTUS DAUCINUS. *Cycl. profundē excavato-umbilicatā, pellucidē aeneo-rufā, anfractibus declivi-convexis, spiraliter lirato-striatis, ad peripheriam carinatis, versus aperturam transversim plicato-rugatis.*

THE CARROT-COLOURED CYCLOTUS. Shell deeply excavately umbilicated, transparent bronze-red, whorls slopingly convex, spirally ridge-striated, keeled at the periphery, transversely plicately wrinkled towards the aperture.

Cyclotoma (Cyclotus) daucinum, Pfeiffer, Pro. Zool. Soc. 1856, p. 337.

Hab. Salomon's Island.

Another beautiful little species, of a bright transparent bronze-red colour.



50.



51.



52.



53.



54.



55.



56.



57.



58.



59.

(Moderately enlarged.)

C Y C L O T U S .

PLATE IX.

Species 50. (Mus. Cuming.)

CYCLOTUS FILOCINCTUS. *Cycl. testū profundē excavato-umbilicatā, pellucido-albā, epidermide corneā deciduā indutā, anfractibus rotundatis, undique spiraliter filo-striatis.*

THE THREAD-GIRT CYCLOTUS. Shell deeply excavately umbilicated, transparent-white, covered with a deciduous horny epidermis, whorls rounded, spirally thread-striated throughout.

Cyclostoma filocinctum, Benson, Ann. Nat. Hist. Second Series, vol. viii. p. 188.

Cyclotus filocinctus, Benson.

Hab. Neilgherry Hills, India.

This delicate little *Cyclotus* is of a raised obtusely-conical form.

Species 51. (Mus. Cuming.)

CYCLOTUS SCALARIS. *Cycl. testū angustē profundē umbilicatā, acutē elevato-convolutā, fulvescente-corneā, unfractibus rotundatis, levigatis, labro subreflexo.*

THE LADDER-LIKE CYCLOTUS. Shell narrowly deeply umbilicated, sharply elevated coiled, fulvous horny, whorls rounded, smooth, lip slightly reflexed.

Cyclostoma scalare, Pfeiffer, Pro. Zool. Soc. 1851, p. 250.

Cyclotus scalaris, Pfeiffer.

Hab. Philippine Islands; Cuming.

The spire is in this species rather sharply, and somewhat loosely, elevated.

Species 52. (Mus. Cuming.)

CYCLOTUS SORDIDUS. *Cycl. testū subangustē profundē umbilicatā, sordidē albā, epidermide tenui corneā deciduā indutā, spirā elevatiusculā, anfractibus spiraliter filo-striatis.*

THE DULL CYCLOTUS. Shell rather narrowly deeply umbilicated, dull-white, covered with a thin deciduous horny epidermis, spire slightly raised, whorls spirally thread-striated.

Cyclostoma (Cyclotus) sordidus, Pfeiffer, Pro. Zool. Soc. 1855, p. 103.

Cyclotus sordidus, Pfeiffer.

Hab. China.

Closely sculptured throughout in the spiral direction, with very fine striae.

Species 53. (Mus. Cuming.)

CYCLOTUS RECLUZIANUS. *Cycl. testū modicē umbilicatā, tenui, fulvescente-fuscā, unfractibus declivi-convexis, spiraliter filo-striatis, ad peripheriam carinatis, ad suturam exiliter plicato-rugatis.*

RECLUZ' CYCLOTUS. Shell moderately umbilicated, thin, fulvous-brown, whorls slopingly convex, spirally thread-striated, keeled at the periphery, faintly plicately wrinkled at the suture.

Cyclostoma (Cyclotus) Recluzianum, Pfeiffer, Pro. Zool. Soc. 1853, p. 51.

Cyclotus Recluzianus, Pfeiffer.

Hab. Salomon's Island.

A thin shell, plicately wrinkled next the suture, across the spiral striae.

Species 54. (Mus. Cuming.)

CYCLOTUS NATALENSIS. *Cycl. testū angustē umbilicatā, cornico-turbanatā, fuscescente, fuscescente undique pallidē linearī-andulatā, castaneo infra peripherium conspicuē fasciatā, anfractibus lāvibūs.*

THE NATAL CYCLOTUS. Shell narrowly umbilicated, conically turbinate, fulvous, faintly linearly waved with brown throughout, conspicuously banded with chestnut below the periphery, whorls smooth.

PFEIFFER, MS. in Mus. Cuming.

Hab. Natal.

In form and colouring this shell partakes of the character of the prevailing Madagascar type of *Cyclostoma*.

Species 55. (Mus. Cuming.)

CYCLOTUS PLEBEJUS. *Cycl. testū angustē umbilicatā, acutē depresso turbinatā, opaco-albā, sordidē livido tinctā, anfractibus rotundatis, obsoletē transversim rugatā.*

THE POOR CYCLOTUS. Shell narrowly umbilicated, sharply depressly turbinated, opake-white, stained with dull

CYCLOTUS.—PLATE IX.

livid-colour, whorls rounded, obsoletely transversely wrinkled.

Cyclostoma plebejum, Sowerby, Pro. Zool. Soc. 1843, p. 60.

Aperostoma plebejum, Pfeiffer.

Cyclotus plebejus, Gray.

Hab. Island of Luzon, Philippines; Cuming.

A dull opaque-white shell, stained with livid-colour, showing traces on its surface of obtuse transverse wrinkles.

Species 56. (Mus. Cuming.)

CYCLOTUS TRAILLII. *Cycl. testā modicē profundē umbilicatā, depressē orbiculari, pallidē spadiceā, purpurascente-castaneo tinctā et fasciatā, anfractibus ad suturam impressis, undique spiraliter tenuē filo-liratis.*

TRAILL'S CYCLOTUS. Shell moderately deeply umbilicated, depressly orbicular, light fawn-colour, tinged and banded with purple-chestnut, whorls impressed at the suture, spirally finely thread-ridged throughout.

PFEIFFER, MS. in Mus. Cuming.

Hab. —?

A characteristic species, but it comes rather near to *C. subdiscoideus*.

Species 57. (Mus. Cuming.)

CYCLOTUS MACGILLIVRAYI. *Cycl. testā profundē perspectivē umbilicatā, obtusē conicā, solidiusculā, vivide anrantio-rufā, anfractibus declivi-convexis, undique confertim filo-liratis, ad peripheriam carinatis.*

MACGILLIVRAY'S CYCLOTUS. Shell deeply perspectively umbilicated, obtusely conical, rather solid, bright orange-red, whorls slopingly convex, closely thread-ridged throughout, keeled at the periphery.

Cyclostoma (Cyclotus) Macgillivrayi, Pfeiffer, Pro. Zool. Soc. 1855, p. 103.

Cyclotus Macgillivrayi, Pfeiffer.

Hab. New Hebrides and New Georgia; Macgillivray.

A rather solid obtusely conoid shell of a bright bronzy orange-red colour.

Species 58. (Mus. Cuming.)

CYCLOTUS MONTANUS. *Cycl. testā angustē profundē umbilicatā, depresso-conicā, tenui, albā, rosaceo-spadicē obscurē punctatā et fasciatā, anfractibus declivi-rotundatis, undique spiraliter filo-liratis.*

THE MOUNTAIN CYCLOTUS. Shell narrowly deeply umbilicated, depresso-conical, thin, white, obscurely dotted and banded with pinkish fawn-colour, whorls slopingly rounded, spirally thread-ridged throughout.

Cyclostoma montanum, Pfeiffer, Pro. Zool. Soc. 1854, p. 300.

Cyclotus montanus, Pfeiffer.

Hab. Neilgherry Hills, India.

The colouring above described may probably be more distinct in some specimens.

Species 59. (Mus. Cuming.)

CYCLOTUS CONOIDEUS. *Cycl. testā angustē umbilicatā, conoideā, fulvā, purpureo-griseo tinctā, fasciā angusta castaneā infra peripheriam cingulatā, anfractibus spiraliter filo-striatis, ad peripheriam carinatis, ad suturam plicatis.*

THE CONOID CYCLOTUS. Shell narrowly umbilicated, conoid, fulvous, tinged with purple-grey, encircled below the periphery with a narrow chestnut band, whorls spirally thread-striated, keeled at the periphery, plicated at the suture.

Cyclostoma spurcum, Sowerby (not of Grataloup), Pro. Zool. Soc. 1843, p. 60.

Cyclostoma conoideum, Pfeiffer.

Tropidophora conoidea, Pfeiffer.

Cyclotus conoideus, Gray.

Hab. Mauritius and Seychelle Islands.

The typical characters of the group to which this little species belongs are developed in it with interesting fullness.

C Y C L O T U S.

	Plate.	Species.		Plate.	Species
asperulus, <i>Sowerby</i>	VI.	33	Lindstedti, <i>Pfeiffer</i>	VIII.	45
Batehianensis, <i>Pfeiffer</i>	VIII.	46	Macgillivrayi, <i>Pfeiffer</i>	IX.	57
Blanchetianus, <i>Moricand</i>	VII.	37	montanus, <i>Pfeiffer</i>	IX.	58
Bogotensis, <i>Pfeiffer</i>	IV.	20	mucronatus, <i>Sowerby</i>	V.	27
Brasiliensis, <i>Sowerby</i>	VI.	28	Nataleensis, <i>Pfeiffer</i>	IX.	54
cingulatus, <i>Sowerby</i>	III.	13	orbellus, <i>Lamarek</i>	VIII.	42
conoideus, <i>Pfeiffer</i>	IX.	59	pallescens, <i>C. B. Adams</i>	II.	7
cornu-venatorius, <i>Petit</i>	VI.	29	Papua, <i>Quoy</i>	VI.	32
corrugatus, <i>Sowerby</i>	III.	11	patera, <i>Pfeiffer</i>	VIII.	44
crassus, <i>C. B. Adams</i>	III.	15	plebejus, <i>Sowerby</i>	IX.	55
daunicus, <i>Pfeiffer</i>	VIII.	49	Popayanus, <i>Lea</i>	V.	24
discoideus, <i>Sowerby</i>	V.	23	prominulus, <i>Férussac</i>	VI.	28
distinctus, <i>Sowerby</i>	VII.	34	pusillus, <i>Sowerby</i>	VII.	39
distomella, <i>Sowerby</i>	VI.	32	Quitensis, <i>Pfeiffer</i>	III.	14
dubiosus, <i>C. B. Adams</i>	VIII.	41	Recluzianus, <i>Pfeiffer</i>	IX.	53
Dunkeri, <i>Pfeiffer</i>	II.	9	sealaris, <i>Pfeiffer</i>	IX.	51
Dysoni, <i>Pfeiffer</i>	VI.	31	seminudus, <i>C. B. Adams</i>	I.	2
filocinetus, <i>Benson</i>	IX.	50	semistriatus, <i>Sowerby</i>	IV.	16
floccosus, <i>Shuttleworth</i>	VIII.	48	sordidus, <i>Pfeiffer</i>	IX.	52
Fortunei, <i>Pfeiffer</i>	IV.	17	spureum (Cyclost.), <i>Sowerby</i>	IX.	59
giganteus, <i>Sowerby</i>	I.	3	spureus, <i>Grateloup</i>	VII.	38
glaucostoma, <i>Pfeiffer</i>	VII.	35	stramineus, <i>Reeve</i>	V.	25
granulatus, <i>Chilly</i>	I.	1	subdiscoideus, <i>Sowerby</i>	IV.	21
Hebraicus (<i>Talca</i>), <i>Lesson</i>	VI.	32	subflammulatus, <i>Pfeiffer</i>	VIII.	13
illotus, <i>Gould</i>	VIII.	47	subrugosus, <i>Sowerby</i>	V.	26
Inea, <i>D'Orbigny</i>	II.	10	substriatus, <i>Sowerby</i>	V.	22
ineomptus, <i>Sowerby</i>	II.	8	suturalis, <i>Sowerby</i>	VIII.	40
inconspicuus, <i>Sowerby</i>	V.	24	Traillii, <i>Pfeiffer</i>	IX.	56
irregularis, <i>Pfeiffer</i>	IV.	18	translucidus, <i>Sowerby</i>	VI.	30
Jamaicensis (<i>Turbo</i>), <i>Chemnitz</i>	I.	5	varians, <i>C. B. Adams</i>	VII.	36
jungosus, <i>C. B. Adams</i>	II.	6	variegatus, <i>Swainson</i>	VI.	29
Lamarekii, <i>Petit</i>	VIII.	42	volvuloides, <i>Sowerby</i>	IV.	19
laxatus, <i>Sowerby</i>	I.	4			

MONOGRAPH
OF THE GENUS
P T E R O C Y C L O S.


"Ay, of a snail; for though he comes slowly, he carries his house on his head; he brings his destiny with him."—*Shakespeare*.



2a



2b




2c



3a



3b



4a



4b



5a



5b



7a

PTERO CYCLOS.

PLATE I.

Genus PTEROCYCLOS, *Benson.*

Testa perampliter umbilicata, plus minus discoidea, fulvo-scente, castaneo vel ustulato-brunneo varie fasciata et undulato-maculata, spirâ depresso, anfractibus angustis, rotundatis, ultimo interdum soluto, tubulo parvo in suturam pone aperturam saepe munito; apertura circulari, labro duplicate, peritremate externo nunc in alam sinuosam nunc in tubulum producto.

Operculum calcareum, utrinque laminatum, inter laminas erectè spiraliter lamellatum, lateribus interdum sub-equaliter concaviusculis, interdum latere externo planiusculo, interno valde concavo.

Shell very largely umbilicated, more or less discoid, fulvous, variously banded and wave-spotted with chestnut or burnt-brown, spire depressed, whorls narrow, rounded, the last sometimes uncoiled, often furnished with a small tube in the suture behind the aperture; aperture circular, lip duplicate, outer peritreme produced, sometimes into a sinuous wing, sometimes into a tube.

Operculum calcareous, laminated on each side, and erectly spirally lamellated between the laminae, sides sometimes equally slightly concave, sometimes rather flat on the outer side, very concave on the inner.

The shells associated in this genus are distinguished by the curious feature of a winged sinus being developed at the upper margin of the lip on arriving at maturity (*Pterocyclos* proper). In some the marginal sinus is modified into a spouted tube (*Rhiostoma*); in others, a spouted tube is developed at the marginal sinus before the shell is mature (*Opisthoporus*), and remains at a greater or less distance behind the aperture. All three forms of *Pterocyclos* have been introduced and minutely described under the foregoing generic names by Mr. Benson. They are natives chiefly of Bengal, Burmah, Siam, Ceylon, and the islands of the Eastern Archipelago. What especial filaments of the animal correspond with this sinus or tube, no one has yet been able to show. Mr. Arthur Adams has described the external characters of the animal from a specimen collected as far north as Tsu-Sima, in the Korea Strait, but no mention is made of the relation between the soft parts and the particular structure which characterizes the genus.

Along with the modification of the sinns, there is also a modification of the operculum. It is composed in both

groups of two laminae, between which is an erect spiral lamella. In *Pterocyclos* proper the spire is pushed out as it were from the inner side, which is thus rendered exceedingly convex; in the *Opisthoporus* and *Rhiostoma* groups the inner and outer laminae are almost equally slightly concave, and as each lamina reaches at its circumference beyond the intervening erect lamelle, the operculum has the appearance of a pulley (Recluz) or vertebra of a fish (Benson).

Species 1. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS BIFRONS. *Pter. testâ perampliter apertâ umbilicatâ, discoideâ, supra peripheriam purpureo-castaneâ et fulvâ, infra violaceâ, immaculatâ, anfractibus rotundatis, supernâ arcuato-striatis; apertura obliquâ circulari, sinu latiusculo, tumido.*

THE TWO FACED PTEROCYCLOS. Shell very largely openly umbilicated, discoid, above the periphery purple-chestnut and fulvous, below violet, unsotted, whorls rounded, arcuately striated at the upper part; aperture obliquely circular, sinus rather broad, swollen.

PFEIFFER, Pro. Zool. Soc. 1855, p. 117.

Hab. Ceylon.

Named *bifrons* from the circumstance of the chestnut colouring terminating abruptly at a line encircling the shell at its periphery.

Species 2. (Fig. a, b. Mus. Cuming.)

PTERO CYCLOS PARILIS. *Pter. testâ modicè perspectivè umbilicatâ, subgloboso-turbanatâ, solidiusculâ, albidi, fulvo-castaneo infra peripheriam fasciatâ, supra undulato-flammatâ, anfractibus declivi-rotundatis, ad suturas marginatis; apertura subamplâ, labro crassè reflexo, sinu brevissimo, subobsoleto.*

THE LIKE PTEROCYCLOS. Shell moderately perspectively umbilicated, rather solid, somewhat globosely turbinated, whitish, banded below the periphery with fulvous-chestnut, undulately flamed above it, whorls slopingly rounded, margined at the sutures, aperture rather large, lip thickly reflected, sinus very short, almost obsolete.

PFEIFFER, MS. in Mus. Cuming.

Hab. —?

In this species, lately named by Dr. Pfeiffer, the shell is subglobosely turbinated, with the umbilicus rather constricted, as in *P. brevis*. Immediately below the periphery it is encircled with a conspicuous chestnut band, above it the colouring is disposed in promiseous zigzag splashes.

Species 3. (Fig. *a, b*, Mus. Cuming.)

PTEROZYCLOS PLANORBULUS. *Pter. testā perampliter apertē umbilicatā, discoideā, fulvo-castaneo tinctā et obscurē fasciatā, anfractibus rotundatis, ad suturas linearē-impressis, supernē tumidē plicato-corrugatis, deinde diagonaliter malleatis; aperturā obliquē circulari, labro duplicito, sinu parviusculo.*

THE COILED PTEROCYCLOS. Shell very largely openly umbilicated, discoid, stained and obscurely banded throughout with fulvous chestnut, whorls rounded, linearly impressed at the sutures, tumidly plicately wrinkled round the upper part, then diagonally malleated; aperture obliquely circular, lip duplicate, sinus rather small.

Cyclostoma planorbulum, Lamarck, Enc. Méth. pl. 461, f. 3.
Cyclotus planorbulus, Swainson.

Pterocyclus planorbulus, Pfeiffer.
Hab. Pulo Condore.

The swollen wrinkles throughout the sutural margin of the whorls show indications of the presence of the auricular sinus, or of the development of some approach to the formation of one, from an early stage of growth.

Species 4. (Fig. *a, b*, Mus. Cuming.)

PTEROZYCLOS BREVIS. *Pter. testā subampliter excavato-umbilicatā, solidā, fulvo-castaneo tinctā et obscurē fasciatā, versus apicem strigatā, anfractibus rotundatis, ad suturas impressis et marginatis; aperturā circulari, labro crassē reflexo, sinu continuo-producto.*

THE SHORT PTEROCYCLOS. Shell rather largely excavately umbilicated, solid, stained and obscurely banded with fulvous chestnut, streaked towards the apex, whorls rounded, impressed and margined at the sutures, aperture circular, lip thickly reflected, sinus continuously produced.

Lituus brevis, Martyn, Univ. Conch. pl. 28.

Turbo Petiverianus, Wood.

Cyclostoma Petiverianum, Gray.

Cyclostoma breve, Pfeiffer.

Myxostoma Petiverianum, Troschel.

Cyclophorus brevis, Benson.

Hab. Pulo Condore.

In this species the sinus is continuously produced, as the shell advances in growth, resting on the body whorl. The circumference of the umbilicus is angularly excavated.

Species 5. (Fig. *a, b*, Mus. Cuming.)

PTEROZYCLOS TENUILABIATUS. *Pter. testā perampliter apertē umbilicatā, angustē discoideā, fulvo-castaneo undique undato-marmorata, anfractibus rotundatis, subobscure diagonaliter malleatis, densē annulato-striatis; aperturā parviusculā obliquā, labro plano-reflexo, sinu tenuē latē dilatato.*

THE THIN-LIPPED PTEROCYCLOS. Shell very largely openly umbilicated, narrowly discoid, wave-marbled throughout with fulvous chestnut, whorls rounded, somewhat obscurely diagonally malleated, closely ring-striated; aperture rather small, oblique, lip flatly reflexed, sinus thinly broadly dilated.

Cyclostoma tenuilabiatum, Metcalfe, Pro. Zool. Soc. 1851, p. 71.

Hab. Borneo.

An elegant, narrowly discoid species, in which the sinus, continuous with the outer expanded lip, is thinly broadly dilated, wing-like.



6a.



7a.



7c.



8a.



9a.



9c.



8b.



10a.



10b.



6b.

PTERO CYCLOS.

PLATE II.

Species 6. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS BATCHIANENSIS. *Pter. testā perampliter apertē umbilicatā, subdiscoideā, fulvo-castaneo tinctā et undatā, anfractibus rotundatis, densē striatis, supernē obscurē spiraliter liratis et malleatis; apertura parvinsculā, sinn sublatē dilatato.*

THE BATCHIAN PTEROCYCLOS. Shell very largely openly umbilicated, subdiscoid, stained and waved with fulvous chestnut, whorls rounded, closely striated, spirally obscurely ridged and malleated at the upper part; aperture rather small, sinus rather broadly dilated.

PFEIFFER, MS. in Mus. Cuming.

Hab. Isle of Batchian.

The whorls in this species present indications of spiral equidistant ridges on the upper part.

Species 7. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS BLANDI. *Pter. testā subampliter apertē umbilicatā, subdiscoideā, fulvo-castaneo obliquè strigatā et variegatā, infra peripheriam angustē fasciatā, anfractibus ad suturas impressis, rūdē striatis; apertura obliquè circulari, labro crassè reflexo, duplice, sinn flexuoso auriculo-dilatato.*

BLAND'S PTEROCYCLOS. Shell rather largely openly umbilicated, somewhat discoid, obliquely streaked and variegated with fulvous chestnut, narrowly banded below the periphery, whorls impressed at the sutures, rudely striated; aperture obliquely circular, lip thickly reflected, duplicate, sinus flexuously auricularly dilated.

BENSON, Ann. Nat. Hist. ser. 2. vol. viii. p. 5. f. 1.

Hab. Island of Pulo Sussan, near Pulo Penang.

I have no authority for regarding this shell as *P. Blandi* beyond the circumstance of its differing from the other described species of the genus, and agreeing pretty nearly with Mr. Benson's outline figure.

Species 8. (Fig. a, b, Mus. Benson.)

PTERO CYCLOS RUFESTRIS. *Pter. testā latissimē per-*

apertē umbilicatā, angustē discoideā, ustulato-fusco tinctā et obscurē fasciatā, anfractibus rotundatis, ad suturas linearē-impressis, deinde plicato-striatis; apertura obliquē circulari, sinn tumido et auriculato-contorta, margine inferiore ad extremitatem nucinato.

THE ROCK-INHABITING PTEROCYCLOS. Shell very broadly, very openly umbilicated, narrowly discoid, stained and obscurely banded with burnt brown, whorls rounded, linearly impressed at the sutures, then plicately striated, aperture obliquely circular, sinus swollen and auricularly twisted, lower margin hooked at the extremity.

BENSON, Journ. Asiat. Soc. 1832, p. 11. pl. 2.

Pterocyclos pictus, Troschel.

Steganotoma pictum, Troschel.

Cyclostoma pictum, Petit.

Hab. Cuttack, India.

The sinus in this very interesting species is tumidly auricularly contorted on the body whorl, and the lower margin curls into it at its extremity in the form of a sharp involute hook.

Species 9. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS ALBERSI. *Pter. testā ampliter apertē umbilicatā, subdiscoideā, fulvā, castaneo undatā, infra peripheriam et circa umbilicum latifasciatā, anfractibus rotundatis, ad suturas plicato-marginatis, deinde rūdē striatis; apertura parvinsculā, sinn flexuoso-auriculato, margine dilatato descendente.*

ALBERS' PTEROCYCLOS. Shell largely openly umbilicated, subdiscoid, fulvous, waved with chestnut, broadly banded below the periphery and around the umbilicus, whorls rounded, plicately margined at the sutures, then rudely striated, aperture rather small, sinus flexuously auriculated, the dilated margin descending.

PFEIFFER, Zeitschr. für Malak. 1847, p. 151.

Cyclostoma Albersi, Petit.

Hab. Khasia Hills, India.

Encircled with two characteristic dark bands. At the aperture the dilated wing-like process descends forward over the sinus.

Species 10. Fig. *a*, *b*, (Mus. Cuming.)

PTEROCYCLOS ANGULIFERUS. *Pter. testā latissimē per-
aperlē umbilicatā, angustē discoideā, fulvo-castaneo-
undatā et tinctā, anfractibus ad suturas impressis,
deinde rotundatis, undique densē ruditē striatis; aper-
turā oblique circulari, labro calloso-duplicato, reflexo,
sinu continuē angulato-producto.*

THE ANGULAR PTEROCYCLOS. Shell very broadly, very
openly umbilicated, narrowly discoid, waved and
stained with fulvous brown, whorls impressed at the

sutures, then rounded, closely rudely striated through-
out, aperture obliquely circular, lip callously dupli-
cate, reflected, sinus continuously angularly pro-
duced.

Cyclostoma anguliferum, Souleyet, Voy. de la Bonite,
Moll. pl. 30, f. 6–11.

Pterocyclos anguliferus, Pfeiffer.

Hab. Cochin-China.

A narrowly discoid species, with the sinus very con-
spicuously angularly produced upon the body whorl.



H.



12 a.



12 b.



13 a.



14 a.



14 b.



13 b



15 a.



15 b.



16.

PTERO CYCLOS.

PLATE III.

Species 11. (Mus. Benson.)

PTERO CYCLOS CETRA. *Pter. testā sublatē apertē umbilicatā, subdiscoideā, fulvo-albidā, rufo-fusco arcuato-maculatā, anfractibus angustis, rotundatis, tenuistriatis; aperturā parvā, sinu brevi, labro parum dilatato.*

THE BUCKLER PTEROCYCLOS. Shell rather broadly openly umbilicated, subdiscoid, fulvous-white, arcuately blotched with red-brown, whorls narrow, rounded, finely striated; aperture small, sinus short, lip but little dilated.

BENSON, Ann. Nat. Hist. scr. 2, vol. xvii. p. 228.
Hab. Moulmein, India.

Very like *P. nanus* which follows, but more discoid and more broadly umbilicated, with the lip less flexuously dilated over the sinus.

Species 12. (Fig. a, b, Mus. Benson.)

PTERO CYCLOS NANUS. *Pter. testā aperlē umbilicatā, subdiscoideā, pallidē fulvā, ad peripheriam rufo-fusco linearī-fasciatā, supra arcuato-maculatā, anfractibus angustis, rotundatis, tenuissimē striatis; aperturā parvā, sinu tumidē flexuoso-ditata.*

THE DWARF PTEROCYCLOS. Shell openly umbilicated, subdiscoid, light fulvous, linearly banded with red-brown at the periphery, arcuately blotched above it, whorls narrow, rounded, very finely striated; aperture small, sinus tumidly flexuously dilated.

BENSON, Ann. Nat. Hist. vol. viii. p. 450.
Hab. Neilgherry Hills.

A small species, in which the wing-like dilation of the lip is flexuously swollen and bent forwards, somewhat after the manner of *P. rupestris* and *Albersi*.

Species 13. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS BILABIATUS. *Pter. testā ampliter subexcavato-umbilicatā, rix discoideā, albida, rufo-fusco undato-strigatā et fasciatā, anfractibus rotundatis, densē striatis; aperturā obliquā, labro peculiariter undato-crispato, ad sinu tumido-cucullato.*

THE TWO-LIPPED PTEROCYCLOS. Shell largely somewhat excavately umbilicated, scarcely discoid, whitish,

wave-streaked and banded with red-brown, whorls rounded, closely striated; aperture oblique, lip peculiarly crimped, tumidly chaubered over the sinus.

BENSON, Zool. Journ. 1834, vol. v. p. 462.

Cyclostoma bilabiatum, Sowerby.

Hab. Salem, near Madras.

Remarkably distinguished by a peculiar crimping of the lip.

Species 14. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS CUMINGII. *Pter. testā latissimē perapertē umbilicatā, discoideā, fulvescente-albā, castaneo undique undato-strigatā, anfractibus angustis, rotundatis, tenuē annulato-striatis; aperturā obliquē circulari, labro calloso-reflexo, sinu conspicuē flexuoso-auriculato.*

CUMING'S PTEROCYCLOS. Shell very broadly, very openly umbilicated, discoid, fulvous-white, sharply waved throughout with chestnut, whorls narrow, rounded, finely ring-striated; aperture obliquely circular, lip callously reflexed, sinus conspicuously flexuously auriculated.

PFEIFFER, Zeitschr. für Malac. 1851, p. 4; Pro. Zool. Soc. 1852, p. 158.

Pterocyclos Cingalensis, Benson.

Hab. Ceylon.

A depresso discoid, very openly umbilicated species, exquisitely painted in a zigzag manner throughout.

Species 15. (Fig. a, b, Mus. Benson.)

PTERO CYCLOS PARVUS. *Pter. testā latē umbilicatā, subdiscoideā, fulvo-albidā, rufo-castaneo fasciatā et undato-strigatā, anfractibus rotundatis, rudē striatis; aperturā obliquē circulari, sinu tumido-auriculato.*

THE SMALL PTEROCYCLOS. Shell broadly umbilicated, subdiscoid, fulvous-white, banded and wave-streaked with reddish chestnut, whorls rounded, rudely striated; aperture obliquely circular, sinus tumidly auriculated.

Spiraculum parvum, Pearson, Journ. Asiat. Soc. 1833, p. 392, pl. 20.

Pterocyclos parvus, Benson.

Cyclostoma spiraculum, var., Sowerby.

Hab. Khasia Hills.

PTEROCYCLOS.—PLATE III.

Very closely allied to *P. cetera* and *nanus*, with the sinus more distinctly auriculated.

Species 16. (Mus. Benson.)

PTEROCYCLOS PULLATUS. *Pter. testá subampliter excavato-umbilicatá, vix discoideá, pullá, obscurè fasciatá et undatá, infra peripheriam saturatiore, anfractibus rotundatis, rudè striatis; aperturá circulari, sinu angustè auriculato.*

THE DARK PTEROCYCLOS. Shell rather largely excavately umbilicated, scarcely discoid, dark-brown, obscurely banded and waved, darker below the periphery, whorls rounded, rudely striated; aperture circular, sinus narrowly auriculated.

BENSON, Ann. Nat. Hist. ser. 2. vol. xvii. p. 227.

Hab. Near the Irawadi, Burmah.

More contractedly umbilicated, and of darker colour than its allies of the Khasia and Neilgherry Hills.



17.



18a



18 b



19 a



20 a



20 b



19 b.



21a.



21 b.



22

PTERO CYCLOS.

PLATE IV.

Species 17. (Mus. Cuming.)

PTERO CYCLOS BICILIATUS. *Pter. testā latē apertē umbilicatā, discoideā, fulvā, castaneo fasciatā et acutē undato-strigatā, anfractibus rotundatis, densē striatis, supra et infra peripheriam seriatim ciliatis, anfractu ultimo pone aperturam tubulo recurvo munito; aperturā obliquē circulari, labro supernē tenuē flexuoso-dilatato.*

THE TWO-LASHED PTEROCYCLOS. Shell broadly openly umbilicated, discoid, fulvous, broadly waved with chestnut, whorls rounded, densely striated, encircled above and below the periphery with a row of lashes, last whorl furnished behind the aperture with a re-curved tube; aperture obliquely circular, lip thinly flexuously dilated at the upper part.

MOUSSON, Benson, Ann. Nat. Hist. ser. 2. vol. xv. p. 14.

Cyclostoma biciliatum, Pfeiffer.

Cyclostoma Charbonnierii, Recluz.

Cyclotus Taylorianus, Pfeiffer.

Opisthoporus biciliatus, Benson.

Hab. Sarawak, Borneo.

The species represented in this and the following plates, characterized by the presence of a tubular process emerging from behind the aperture, constitute the genera *Opisthoporus* and *Rhiostoma* of Benson. When it is a little removed from the aperture the species belong to the former group, when arising from a tubular sinuation of the lip, to the latter. *P. ciliatus* is distinguished as a species in being encircled on either side of the periphery, with a neatly-defined row of bristly lashes.

Species 18. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS BIROSTRIS. *Pter. testā latissimē apertē umbilicatā, discoideā, fulvā, castaneo fasciatā et acutē undato-strigatā, anfractibus rotundatis, undique annulato-striatis, anfractu ultimo subsoluto, pone aperturam tubulo incurvo munito; aperturā circulari, labro concavo-inflecto, supernē tubulo-sinuato.*

THE TWO-BEAKED PTEROCYCLOS. Shell very broadly openly umbilicated, discoid, fulvous, banded and sharply wave-streaked with chestnut, whorls rounded, ring-striated throughout, last whorl rather uncoiled, furnished with an incurved tube behind the aperture; aperture circular, lip concavely inflected, tubularly sinuated at the upper part.

Cyclotus birostris, Pfeiffer, Pro. Zool. Soc. 1854, p. 300.

Opisthoporus birostris, Pfeiffer.

Hab. Sarawak, Borneo.

The tube behind the aperture is not curved backwards, as in the last species, but forwards, and a second tube is formed by an inflected sinuation of the lip.

Species 19. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS HAINESII. *Pter. testā ampliter perspectivē umbilicatā, subdiscoideā, fulvescente-fuscā, castaneo fasciatā et marmoratā, anfractibus ad suturas subappressis, deinde rotundatis, densē striatis, anfractu ultimo conspicuē soluto; aperturā circulari, labro crassè reflexo, supernē tubulo recurvo prominente munito.*

HAINES' PTEROCYCLOS. Shell largely perspectively umbilicated, somewhat discoid, fulvous-brown, banded and mottled with chestnut, whorls a little appressed at the sutures, then rounded, densely striated, last whorl conspicuously unrolled; aperture circular, lip thickly reflected, furnished at the upper part with a prominent recurved tube.

Rhiostoma Hainesii, Pfeiffer, MS. in Mus. Cuming.

Hab. Camboja; Mouhot.

A very striking species, in which the last whorl is uncoiled on arriving at maturity, to make room for a conspicuous recurved tube, emerging from the upper margin of the aperture.

Species 20. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS SIMPLICILABRIS. *Pter. testā latē apertē umbilicatā, discoideā, fulvā, rufo-fusco fasciatā et undulato-strigatā, anfractibus rotundatis, undique densē annulato-striatis, anfractu ultimo soluto; aperturā circulari, labro calloso-reflexo, sinu in tubulum producto.*

THE SIMPLE-LIP PTEROCYCLOS. Shell broadly umbilicated, discoid, fulvous, banded and wave-streaked with brown, whorls rounded, closely ring-striated throughout, last whorl uncoiled; aperture circular, lip callously reflected, sinus produced into a tube.

PFEIFFER, MS. in Mus. Cuming.

Hab. Camboja.

In this and the species which follows, the last whorl is

also curiously pushed from its spire of convolution on arriving at maturity, in consequence of the sinus, characteristic of this genus, being developed into a tube.

Species 21. (Fig. *a, b*, Mus. Cuming.)

PTEROCYCLOS HOUSEI. *Pter. testā subampullē apertē umbilicatā, discoideā, albida, strigis angustis fulvescente-fuscis pictā, anfractibus ad suturas impressis et marginatis, deinde rotundatis, densē tenuistriatis, anfractu ultimo soluto; aperturā circulari, labro tenuē expanso, sinu in tubulum angustum elongatum producto.*

HOUSE'S PTEROCYCLOS. Shell rather largely openly umbilicated, discoid, whitish, painted with narrow fulvous-brown angular streaks, whorls impressed and margined at the sutures, then rounded, closely finely striated, last whorl uncoiled; aperture circular, lip thinly expanded, sinus produced into a narrow elongated tube.

Cyclostoma Housei, Haines, Ann. Lyc. New York, vol. vi.

p. 157. pl. 5. f. 12 to 15.

Rhiostoma Housei, Benson.

Hab. Siam; House.

A more slenderly convoluted species than the preceding with the tube rather narrower and longer.

Species 22. (Mus. Cuming.)

PTEROCYCLOS COCHINCHINENSIS. *Pter. testā sublate apertē umbilicatā, discoideā, sub epidermide fulvā albā, anfractibus rotundatis, undique tenuistriatis, anfractu ultimo subsoluto; aperturā circulari, labro simplici, sinu in tubulum parvum producto.*

THE COCHIN-CHINA PTEROCYCLOS. Shell rather broadly openly umbilicated, discoid, white, beneath a fulvous epidermis, whorls rounded, finely striated throughout, last whorl slightly uncoiled; aperture circular, lip simple, sinus produced into a small tube.

Cyclostoma Cochinchinense, Pfeiffer, Pro. Zool. Soc. 1856. p. 337.

Opisthoporus Cochinchinensis, Pfeiffer.

Hab. Cochin-China.

Better specimens than are at present known of this species are required to ascertain its colouring. The shell here figured is entirely denuded of its epidermis.



23



24



25



26a



27a.



27b.



26b



28



29



30

PTERO CYCLOS.

PLATE V.

Species 23. (Mus. Cuming.)

PTERO CYCLOS BERNARDII. *Pter. testā ampliter umbilicatā, subdiscoideā, fulvescente, anfractibus rotundatis, tenuē annulato-striatis; aperturā circulari, labro crassè reflexo, sinu in tubulum parvum incurvum producto.*

BERNARDI'S PTEROCYCLOS. Shell largely umbilicated, somewhat discoid, fulvous, whorls rounded, finely ring-striated; aperture circular, lip thickly reflected, sinus produced into a small incurved tube.

PFEIFFER, MS. in Mus. Cuming.

Hab. —?

Of simple form and semitransparent fulvous colour, with a small incurved tubular sinus.

Species 24. (Mus. Cuming.)

PTERO CYCLOS TUBULIFERUS. *Pter. testā latē apertē umbilicatā, subdiscoideā, fulvescente, rufo-fusco marmoratā, anfractibus rotundatis, densē tenuistriatis; aperturā obliquē circulari, labro duplicito, concavodilatato, sinu in tubulum minutum producto.*

THE TUBE-BEARING PTEROCYCLOS. Shell broadly openly umbilicated, subdiscoid, fulvous, mottled with red-brown, whorls rounded, closely finely striated; aperture obliquely circular, lip duplicate, concavely dilated, sinus reduced into a minute tube.

Cyclostoma tubuliferum, Pfeiffer, Pro. Zool. Soc. 1853, p. 51.

Cyclotus tubuliferus, Pfeiffer.

Opisthoporus tubuliferus, Benson.

Hab. —?

The tubular sinus is smaller in this species than in any other, almost concealed from view within the sutures.

Species 25. (Mus. Cuming.)

PTERO CYCLOS ROSTELLATUS. *Pter. modicē excavato-umbilicatā, subdiscoideā, fulvā, rufo-fusco undato-marmoratā, anfractibus rotundatis, tenuistriatis, anfractu ultimo tubulo minuto erecto pone aperturam; aperturā circulari, labro expanso-reflexo.*

THE BEAKED PTEROCYCLOS. Shell moderately excavately umbilicated, subdiscoid, fulvous, wave-marbled

with red-brown, whorls rounded, finely striated, last whorl with a small erect tube behind the aperture; aperture circular, lip expandedly reflected.

Cyclostoma rostellatum, Pfeiffer, Zeitschr. für Malak. 1851, p. 8.

Cyclotus rostellatus, Pfeiffer.

Opisthoporus rostellatus, Pfeiffer.

Hab. —?

In this species, which is less openly convoluted, the minute tube, accidentally omitted in our figure, is a little removed from the aperture.

Species 26. (Fig. a, b, Mus. Cuming.)

PTERO CYCLOS HISPIDUS. *Pter. testā latissimè perapertē umbilicatā, plano-discoideā, solidā, fulvā, suprēmè usculato-fusco tinctā et marmoratā, onfractibus rotundatis, spiraliter multiliratis, concentricè densē striatis, epidermide hispidulā indutis, anfractu ultimo tubulo recurvo subremotè pone aperturam; aperturā obliquē circulari, labro densē callosō-reflexo, sinu latē angulatē continuo-producto.*

THE HAIRY PTEROCYCLOS. Shell very broadly very openly umbilicated, flatly discoid, solid, fulvous, stained and marbled at the upper part with burnt brown, whorls rounded, spirally many-ridged, concentrically closely striated, covered with a slightly hairy epidermis, last whorl with a recurved tube at some distance behind the aperture; aperture obliquely circular, lip thickly callously reflected, sinus broadly angularly continuously produced.

Spiraculum hispidum, Pearson, Journ. Asiat. Soc. 1853, p. 391, pl. 20.

Cyclostoma spiraculum, Sowerby.

Steganotoma Princepsi, Von dem Busch.

Pterocyclos hispidus, Benson.

Pterocyclos Princepsi, Pfeiffer.

Hab. Khasia Hills, India.

This is the only species of the genus in which there is any close well-defined spiral sculpture; and the recurved tube at a point of the suture removed from the aperture, characteristic of the group *Opisthoporus*, is here combined with the produced angular sinus chiefly typical of *Pterocyclos* proper.

Species 27. (Fig. a, b, Mus. Cuming.)

PTEROXYCLOS ANOMALUS. *Pter. testā latissimē apertē umbilicatā, discoideā, fulvescente-albā, rufo-fusco undique nitidē undato-strigatā, anfractibus rotundatis, decussatim striatis, anfractu ultimo tubulo suturali recumbente pone aperturam; aperturā obliquē circulari, labro latē plano-reflexo, supernē subdilatato, sinus fere obsolete.*

THE ANOMALOUS PTEROCYCLOS. Shell very broadly openly umbilicated, discoid, fulvous-white, neatly wave-streaked throughout with red-brown, whorls rounded, decussately striated, last whorl with a recumbent sutural tube behind the aperture; aperture obliquely circular, lip broadly flatly reflected, a little dilated at the upper part, sinus almost obsolete.

Cyclostoma pterocyclodes, Pfeiffer, Pro. Zool. Soc. 1854, p. 300.

Cyclotus pterocyclodes, Pfeiffer.

Opisthoporus pterocyclodes, Pfeiffer.

Hab. Borneo.

The tube of this species is reeumbent in the suture.

Species 28. (Mus. Cuming.)

PTEROXYCLOS TENER. *Pter. testā ampliter umbilicatā, subdiscoideā, pellucido-luteā, anfractibus rotundatis, undique tenuē annulato-striatis; aperturā circulari, labro tenui, duplicato, sinus in tubulo minuto recurvo producto.*

THE DELICATE PTEROCYCLOS. Shell largely umbilicated, subdiscoid, transparent yellow, whorls rounded, finely ring-striated throughout; aperture circular, lip thin, duplicate, sinus produced into a small recurved tube.

MENKE, Malak. Blat. 1856, p. 69.

Rhiostoma tener, Benson.

Hab. Cochin-China.

A delicate semitransparent shining yellow shell, ringed throughout with thread-like striae.

Species 29. (Mus. Cuming.)

PTEROXYCLOS EURYOMPHALUS. *Pler. testā latissimē perapertē umbilicatā, plano-discoideā, fulvā, castaneo undique aculē undatā, spirā subimmersā, anfractibus angustis, rotundatis, tenuistriatis, anfractu ultimo tubulo parvo recurvo subremotē pone aperturam; aperturā obliquē circulari, labro duplicato, supernē tenuē tumido-dilatato.*

THE WIDELY-UMBILICATED PTEROCYCLOS. Shell very widely very openly umbilicated, flatly discoid, fulvous, sharply waved throughout with chestnut, spire rather immersed, whorls narrow, rounded, finely striated, last whorl with a small reeurved tube at some distance behind the aperture; aperture obliquely circular, lip duplicate, thinly tumidly dilated at the upper part.

Cyclostoma euryomphalus, Pfeiffer, Pro. Zool. Soc. 1856, p. 337.

Opisthoporus euryomphalus, Pfeiffer.

Hab. Borneo.

Very slenderly discoidly coiled, with a crowded painting of sharply zigzag chestnut streaks.

Species 30. (Mus. Benson.)

PTEROXYCLOS HAUGHTONI. *Pter. testā ampliter umbilicatā, subdiscoideā, fulvo-albidā, supernē rufo-fusco tinctā, ad peripheriam fasciatā et undulatā, anfractibus declivi-rotundatis, densē plicato-striatis, anfractu ultimo soluto; aperturā circulari, labro vix reflexo, sinus erecto-auriculato, subrecurvo.*

HAUGHTON'S PTEROCYCLOS. Shell largely umbilicated, subdiscoid, fulvous-white, stained on the upper part with red-brown, banded and delicately waved at the periphery, whorls slopingly rounded, closely plicately striated, last uneoiled; aperture circular, lip scarcely reflected, sinus erectly auriculated, a little reeurved.

Rhiostoma Haughtoni, Benson, Ann. Nat. Hist. ser. 3, vol. 5, p. 96.

Hab. Moulmein, India.

A very distinct dark-stained species, in which the sinus assumes the form of a reeurved section of a tube.

P T E R O C Y C L O S.

	Plate.	Species.		Plate.	Species.
Albersi, <i>Pfeiffer</i>	II.	9	Haughtoni (<i>Rhiostoma</i>), Benson	V.	30
anguliferus (<i>Cyclostoma</i>), Souleyet ..	II.	10	lispidus (<i>Spiraculum</i>), Pearson	V.	26
anomalus, <i>Reeve</i>	V.	27	Housei, <i>Haines</i>	IV.	21
Batchianensis, <i>Pfeiffer</i>	II.	6	nanus, <i>Benson</i>	III.	12
Bernardii, <i>Pfeiffer</i>	V.	23	parilis, <i>Pfeiffer</i>	I.	2
biciliatus (<i>Opisthoporus</i>), Pfeiffer	IV.	17	parvus (<i>Spiraculum</i>), Pearson	III.	15
bifrons, <i>Pfeiffer</i>	I.	1	pictus, Troschel	II.	8
bilabiatus, <i>Benson</i>	III.	13	planorbulus (<i>Cyclostoma</i>), Lamarck ..	I.	3
birostris (<i>Opisthoporus</i>), Pfeiffer	IV.	18	Princepsi, <i>Pfeiffer</i>	V.	26
Blandi, <i>Benson</i>	II.	7	Pterocycloides (<i>Opisthoporus</i>), Pfeiffer	V.	27
brevis (<i>Litus</i>), Martyn	I.	4	pullatus, <i>Benson</i>	III.	16
cetra, <i>Benson</i>	III.	11	rostellatus (<i>Opisthoporus</i>), Pfeiffer ..	V.	25
Cingalensis, <i>Benson</i>	III.	14	rupestris, <i>Benson</i>	II.	8
Cochinehinensis (<i>Opisthoporus</i>), Pfeiffer	IV.	22	simplicilabris, <i>Pfeiffer</i>	IV.	20
Cuningii, <i>Pfeiffer</i>	III.	14	tenuilabiatus (<i>Cyclostoma</i>), Metcalfe	I.	5
euryomphalus (<i>Opisthoporus</i>), Pfeiffer	V.	29	tubuliferus (<i>Opisthoporus</i>), Pfeiffer ..	V.	24
Hainesii (<i>Rhiostoma</i>), Pfeiffer	IV.	19	tener, <i>Menke</i>	V.	28

MONOGRAPH
OF THE GENUS
C H O N D R O P O M A .

“I would not for a world of gold
That Nature’s lovely face should tire;
Fountain of blessings yet untold;
Pure source of intellectual fire!”—*Bloomfield.*



Slightly Enlarged

CHONDROPOMA.

PLATE I.

Genus CHONDROPOMA, Pfeiffer.

Testa oblongo-turrita, rarius globoso-turbanata, plerumque densè filo-lirata vel reticulata, ad suturas crenulata; apertura parvissima, ovalis, labro nunc simplici vel expanso, nunc fimbriato. Operculum ovale, subcartilagineum, planum, anfractibus paucis, rapidè accrescentibus, nucleo plerumque valde excentrico.

Shell oblong-turreted, rarely globosely turbinated, generally densely thread-ridged or reticulated, and crenulated at the sutures; aperture rather small, oval, with the lip sometimes simple or expanded, sometimes fimbriated. Operculum oval, somewhat cartilaginous, flat, whorls few, increasing rapidly, nucleus mostly far removed from the centre.

This genus, in which I include *Cistula* of Gray, and of Pfeiffer's monograph, is characterized by the operculum being of a flat cartilaginous substance, of a few rapidly increasing whorls, with the nucleus removed from the centre. It comprises about ninety species, all inhabiting the islands of the West Indies and the adjacent lands of Venezuela, Guatemala, and Florida; but the Continental species are comparatively few in number. The shells are, for the most part, characterized by a remarkable similarity of form and sculpture, oblong-turreted, densely crowded with raised thread-like striae or ridges, which in some species run longitudinally only, in others both longitudinally and transversely, producing a more or less elaborately woven reticulation. The aperture of the shell is generally small, circularly ovate, and fimbriated, and when the frill is auricled at the upper part, which is not unfrequently the case, the auricles are deposited within the suture, as the whorl advances in growth, in tooth-like erenules. In a few species the longitudinal striae assume the form of erect lamellæ or membranaceous threads, and they are often rendered scabrous by decussating incised striae. The colouring of the shells is mostly fulvous, variously banded, streaked, or freckled with reddish brown.

Species 1. (Fig. a, b, c, Mus. Cuming.)

CHONDROPOMA MAGNIFICA. *Chond. testa ovato-conica, subcylindracea, plus minus umbilicata, diaphano-alba, strigis ritterisque rufescente-fuscis cingulata, anfractibus tumido-convervis, dense longitudinaliter filo-liratis, liris*

diaphanis: apertura ovata, labro latè fimbriato-reflexo, fusco lineari-radiato.

THE MAGNIFICENT CHONDROPOMA. Shell ovately conical, subcylindrical, more or less umbilicated, diaphanous white, encircled with red-brown streaks and fillets, whorls tumidly convex, densely longitudinally thread-ridged, ridges diaphanous; aperture ovate, lip broadly fimbriately reflexed, lineally rayed with brown.

Cyclotoma magnificum, Pfeiffer, Pro. Zool. Soc. 1852, p. 142. pl. 13. f. 3.

Chondropoma magnificum, Pfeiffer.

Chondropoma Gutierrezi, Poey.

Hab. Island of Haiti, West Indies; Sallé.

Of a delicate diaphanous white, finely stringed throughout with thread-like ridges of similarly diaphanous substance, the whorls being encircled with red-brown fillets and bands of streaks, conspicuously rayed in the aperture.

Species 2. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA WEINLANDI. *Chond. testa vix umbilicata, elongato-cylindracea, albida vel rufescente-aurantia, castaneo-fusco varie striata, fasciata, vel punctata, anfractibus modice convexis, creberrime pliato-striatis; apertura obliquè ovata, parva, supernè carinata, subsoluta, labro simplici, continuo.*

WEINLAND'S CHONDROPOMA. Shell scarcely umbilicated, elongately cylindrical, white or reddish orange, variously streaked, banded or dotted with chestnut-brown, whorls moderately eonvex, very closely pliately striated; aperture obliquely ovate, small, keeled at the upper part, a little detached, lip simple, continuous.

PFEIFFER, Malak. Blat. 1862.

Chondropoma semilabre, Pfeiffer (not *Cyclotoma semi-labre*, Lamarck).

Hab. Island of Haiti, West Indies; Sallé.

Extremely variable in painting, but constant in form; a somewhat elongately cylindrical form, with a small semi-detached aperture.

Species 3. (Mus. Cuming.)

CHONDROPOMA EGREGIUM. *Chond. testa coarctatè umbilicata, ovato-conica, sericeo-aurantia, rufescente-fusco*

obscure punctato-fasciatā et prope apicem strigatā, anfractibus convexis, dense minutē reticulatis; aperturā obliquè ovatā, latissimē fimbriatā.

THE CHOICE CHONDROPOMA. Shell contractedly umbilicated, ovately conical, silky orange, obscurely dot-banded with red-brown, and near the apex streaked, whorls convex, densely minutely reticulated; aperture obliquely ovate, very broadly fimbriated.

Cyclostoma egregium, Gundlach, in Poey, Mém. vol. ii. p. 13. pl. 1. f. 13.

Chondropoma egregium, Pfeiffer.

Hab. Cuba; Gundlach.

The specimen of this beautiful species here figured is a bright silky orange colour. Dr. Pfeiffer describes a dark violet variety, and there are probably many others.

Species 4. (Mus. Cuming.)

CHONDROPOMA SHUTTLEWORTHI. *Chond. testā clausè umbilicatā, tumido-conicā, subcylindracea, fulvā, diaphanā, rufescens-fusco interruplē tenuiatā, pone labrum radiatā, anfractibus rudē convexis, spiraliter filo-striatis, versus apicem minutē reticulatis; aperturā obliquè ovatā, subsolutā, latissimē fimbriatā, fuscō-radialitā.*

SHUTTLEWORTH'S CHONDROPOMA. Shell with closed umbilicus, tumidly conical, subcylindrical, fulvous, diaphanous, interruptedly filleted with red-brown, rayed behind the lip, whorls rudely convex, spirally thread-striated, minutely reticulated towards the apex; aperture obliquely ovate, slightly detached, very broadly fimbriated, frill rayed with brown.

Cyclostoma Shuttleworthi, Pfeiffer, Pro. Zool. Soc. 1851, p. 246.

Chondropoma Shuttleworthi, Pfeiffer.

Cyclostoma verecundum, Poey.

Chondropoma verecundum, Pfeiffer.

Hab. Cuba.

This is a very thin semitransparent shell, finely decussately sculptured, but the longitudinal sculpture soon disappears, leaving the spiral sculpture well defined throughout.

Species 5. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA PICTUM. *Chond. testā compressè umbilicatā, oblongo-conicā, parum cylindracea, fulvo-albidā, diaphanā, punctis ustulato-fusca promiscue strigatā, interdum nigro-violaceū, labro aurantio, anfractibus subtumido-convexis, liris tenuibus spiraliter confertim funiculatis; aperturā obliquè ovatā, labro subincrassato, continuo.*

THE PAINTED CHONDROPOMA. Shell compressly umbilicated, oblong conical, but little cylindrical, fulvous-white, diaphanous, promiscuously streaked with burnt-brown dots, sometimes dark violet, with the lip orange, whorls rather tumidly convex, closely spirally cored with fine ridges; aperture obliquely ovate, lip rather thick, continuous.

Cyclostoma pictum, Pfeiffer, Wieg. Arch. für Nat. 1839, vol. i. p. 356.

Cyclostoma Sagra, D'Orbigny.

Cyclostoma Malogani, Gould.

Chondropoma pictum, Pfeiffer.

Hab. Cuba.

Very definitely and regularly cored throughout, with fine ridges, the interstices between which are not reticulated, but striately punctured. This species has been sometimes mistaken for *C. Weinlandi*, in which the ridged sculpture runs in the opposite direction.

Species 6. (Mus. Cuming.)

CHONDROPOMA HABICII. *Chond. testā compressè umbilicatā, ovato-conicā, diaphano-albā, ferrugineo tinctā, fusco sparsim punctato-fasciatā, anfractibus subrotundatis, longitudinaliter filo-liratis, liris creberrinis, subfimbriatis; aperturā obliquè ovatā, labro latè caloso-expanso.*

HABICHE'S CHONDROPOMA. Shell compressly umbilicated, ovately conical, diaphanous white, tinged with rust, sparsely dot-banded with brown, whorls rather rounded, longitudinally thread-ridged, ridges very close set, slightly fimbriated; aperture obliquely ovate, lip broadly callously expanded.

WEINLAND, Malak. Blat. 1862.

Hab. Island of Haiti; Weinland.

Sculptured with close-set longitudinal thread-striae of semi-diaphanous substance, like *C. magnificum*, a little fimbriately waved.



ab

Slightly Enlarged

CHONDROPOMA.

PLATE II.

Species 7. (Fig. 7, 9, Mus. Cuming.)

CHONDROPOMA TENEBROSUM. *Chond. testā modicē umbilicatā, ovato-turritā, ferruginea vel purpureo-fuscā, obscurè punctato-fasciatā, anfractibus rotundatis, densē scabrose lirato-reticulatis; aperturā circulari, latissimè fimbriatā.*

THE DARK CHONDROPOMA. Shell moderately umbilicated, ovately turreted, rust or purple-brown, obscurely dot-banded, whorls rounded, densely sebrously ridge-reticulated; aperture circular, very broadly fimbriated.

Cyclostoma tenebrosum, Morelet, Test. Noviss. p. 23.

Chondropoma tenebrosum, Pfeiffer.

Cyclostoma claudicans, Pfeiffer.

Hab. Cuba.

The surface of this species is quite sebrous, with a crowded reticulation of thin sharp lamella-like ridges. The specimen represented at Fig. 9 is the type of *C. claudicans*.

Species 8. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA IRRADIANS. *Chond. testā compressē umbilicatā, acuminato-turritā, diaphano-albā, maculis quadratis rufescente-fuscis, quadrifasciatā, anfractibus convexis, minutè scabroso-reticulatis; aperturā parvā, subcirculari, labro latissimè fimbriato, conspicuè radiato.*

THE SHINING CHONDROPOMA. Shell compressly umbilicated, acuminately turreted, diaphanous-white, encircled with four bands of square red-brown spots, whorls convex, minutely sebrously reticulated; aperture small, nearly circular, lip very broadly fimbriated, conspicuously rayed.

Cyclostoma iradians, Shuttleworth, MS.; Pfeiffer, Monog. Pneum. vol. i. p. 294.

Chondropoma iradians, Pfeiffer.

Hab. Cuba.

A beautifully reticulated shell, of delicate subdiaphanous substance, in which the red-brown bands are mostly broken up into square spots.

Species 10. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA SALLEANUM. *Chond. testā compressē umbilicatā, acuminato-turritā, tenuiculā, albā vel pal-*

lidē aurantiā, castaneo-fusco lentiginosā et fasciatā anfractibus declivi-rotundatis, densē lirato-reticulatis, liris ad suturas plicato-auriculatis; aperturā obliquè ovatā, subsolutā, supernè carinatā, labro tenuè reflexo.

SALLE'S CHONDROPOMA. Shell compressly umbilicated, acuminate turreted, rather thin, white or pale orange, freckled and banded with chestnut-brown, whorls slopingly rounded, densely ridge-reticulated, ridges plicately auriculated at the sutures; aperture obliquely ovate, a little detached, keeled at the upper part, lip thinly reflected.

Cyclostoma Salleanum, Pfeiffer, Zeitschr. für Malac. 1850, p. 78.

Chondropoma Salleanum, Pfeiffer.

Hab. Island of Haiti.

A shell of very delicate substance, very elegantly turreted in a sloping manner.

Species 11. (Mus. Cuming.)

CHONDROPOMA LATILABRE. *Chond. testā latē umbilicatā, globoso-turbinatā, diaphano-albā, maculis quadratis ferrugineo-fuscis fasciatā, anfractibus rotundatis, confertim longitudinaliter filo-striatis; aperturā circulari, fauce livido-purpureā, labro latē expanso, albo.*

THE BROAD-LIP CHONDROPOMA. Shell broadly umbilicated, globosely turbinated, diaphanous-white, encircled with bands of square rust-brown spots, whorls rounded, closely longitudinally thread-striated; aperture circular, interior livid-purple, lip broadly expanded, white.

Cyclostoma latilabris, D'Orbigny, Moll. Cuba, vol. i. p. 225. pl. 21. f. 12.

Choanapoma latilabre, Gray.

Chondropoma latilabre, Pfeiffer.

Hab. Cuba.

Distinguished by the tubular form of the whorls and by their globose manner of turbination, leaving a large central umbilicus.

Species 12. (Mus. Cuming.)

CHONDROPOMA GLABRATUM. *Chond. testā imperforatā, oblongo-cylindraceā, fulvescente, castaneo-fusco fasciatā et vittatā, anfractibus tumido-convexit, liris tenuibus*

glabris spiraliiter funiculatis; aperturā obliquè ovatā, labro simplici.

THE SMOOTH CHONDROPOMA. Shell imperforate, oblong-cylindrical, fulvous, banded and filleted with chestnut-brown, whorls tumidly convex, spirally corded with fine smooth ridges; aperture obliquely ovate, lip simple.

WEINLAND, Malak. Blat. 1862.

Hab. Bahamas.

Of a rather tumid oblong form, encircled throughout with fine smooth cord-like ridges.

Species 13. (Mus. Cuming.)

CHONDROPOMA VIOLACEUM. *Chond. testā compressè umbilicatā, ovato-turritatā, violaceā vel ferrugineo-tinctā, fusco obscurè strigatā, anfractibus convexis, densissimè scabroso-reticulatis; aperturā ovatā, parviusculā, labro scabroso-reflexo, albo.*

THE VIOLET CHONDROPOMA. Shell compressly umbilicated, ovately turreted, violet or rust-tinged, obscurely streaked with brown, whorls convex, very densely seaborously reticulated; aperture ovate, rather small, lip seaborously reflected, white.

Cyclostoma violaceum, Pfeiffer, Pro. Zool. Soc. 1851, p. 245.

Tudora violacea, Pfeiffer.

Cistula violacea, Adams.

Chondropoma violaceum, Pfeiffer.

Hab. Cuba.

The surface of this species is remarkably closely reticulated with erect lamella-like striæ.

Species 14. (Mus. Cuming.)

CHONDROPOMA MÆSTUM. *Chond. testā compressè umbilicatā, cylindraceo-ovatā, diaphano-albā, inter liras rufo-fusco articulatā, anfractibus tumido-converxis, liris confertis funiculatis; aperturā obliquè ovatā, supernè subcarinatā, labro parum reflexo.*

THE SAD CHONDROPOMA. Shell compressly umbilicated, cylindrically ovate, diaphanous-white, articulated with red-brown between the ridges, whorls tumidly convex, corded with close-set ridges; aperture obliquely ovate, slightly keeled at the top, lip but little reflected.

Cyclostoma mæstum, Shuttleworth, Pfeiffer, Malak. Blat. 1854, p. 97.

Cyclostoma Poeyanum, Pfeiffer (not of D'Orbigny).

Chondropoma Poeyanum, Pfeiffer.

Cyclostoma Hellenicum, Gray.

Cyclostoma Charpentieri, Shuttleworth.

Chondropoma mæstum, Pfeiffer.

Hab. Cuba.

Well distinguished by the colouring being in the interstices between the corded ridges.



15



16a



16b



17



18



19



20



21



22



23

Slightly Enlarged

CHONDROPOMA.

PLATE III.

Species 15. (Mus. Cuming.)

CHONDROPOMA POEYANUM. *Chond. testā compressē umbilicatā, oblongo-cylindraceā, solidisculnā, albida, anfractibus convexis, spiraliter confertim filo-liratis; aperturā obliquē ovatā, supernē subcarinatā, labro parum reflexo.*

POEY'S CHONDROPOMA. Shell compressly umbilicated, oblong cylindrical, rather solid, whitish, whorls convex, spirally closely thread-ridged; aperture obliquely ovate, slightly keeled at the upper part, lip but little reflected.

Cyclostoma Poeyanum, D'Orbigny, Moll. Cub. p. 264. pl. 22. f. 24 to 27.

Chondropoma Poeyanum, Pfeiffer.

Hab. Cuba.

A simple uncoloured corded shell, supposed at one time to be the *Turbo elongatus* of Wood.

Species 16. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA NEWCOMBIANUM. *Chond. testā compressē umbilicatā, subacuminato-turrītā, fulvescente, purpureo-castaneo variè punctato-fasciatā et lentiginosā, anfractibus rotundatis, longitudinaliter densē filo-striatis et hic illic plicatis, plicis ad suturas auriculatis; aperturā parvā, fere circulari, dissoluta, supernē carinatā, labro simplici.*

NEWCOMB'S CHONDROPOMA. Shell compressly umbilicated, rather acuminate turreted, fulvous, variously dot-banded and freckled with purple chestnut, whorls rounded, longitudinally densely thread-striated, and here and there plicated, plaits auriculated at the sutures; aperture small, nearly circular, detached, keeled at the upper part, lip simple.

Cyclostoma Newcombianum, Adams.

Cyclostoma Troscheli, Pfeiffer.

Cyclostoma Swiftianum, Newcomb.

Chondropoma Nereombianum, Pfeiffer.

Hab. Island of St. Thomas, West Indies.

An elegantly-turreted prettily-freckled shell, which, in addition to the close-set thread-like striae, is sculptured with occasional plaits, which appear as crenules in the sutures.

Species 17. (Mus. Cuming.)

CHONDROPOMA JUCUNDUM. *Chond. testā compressē umbilicatā, ovato-turrītā, rubicundo-aurantiā, tenui, semi-pellucidā, anfractibus rotundatis, striis filiformibus, in suturas hic illuc auriculatis, confertim reticulatis; aperturā obliquē ovatā, labro sublate reflexo.*

THE PLEASANT CHONDROPOMA. Shell compressly umbilicated, ovately turreted, ruddy orange, thin, semi-transparent, whorls rounded, closely reticulated with thread-like striae, here and there auriculated in the sutures; aperture obliquely ovate, lip rather broadly reflected.

Cyclostoma jucundum, Pfeiffer, Pro. Zool. Soc. 1852, p. 66.
Hab. ——?

An extremely delicate transparent shell, of a peculiar ruddy orange, closely reticulated throughout.

Species 18. (Mus. Cuming.)

CHONDROPOMA HEMIOTUM. *Chond. testā compressē umbilicatā, oblongo-turrītā, albida, fusco minutē punctulatā, ad basin fasciatā, anfractibus convexis, striis filiformibus ad suturas auriculatis, confertim decussatis, quarum spiralibus fere obsolete; aperturā obliquē ovatā, subsolutā, supernē carinatā, labro subreflexo.*

THE DOTTED CHONDROPOMA. Shell compressly umbilicated, oblong turreted, whitish, minutely dotted with brown, banded at the base, whorls convex, closely decussated with thread-like striae, auriculated at the sutures, of which the spiral are nearly obsolete; aperture obliquely ovate, a little detached, keeled at the upper part, lip slightly reflected.

PFEIFFER, Pro. Zool. Soc. 1852, p. 143.

Hab. Island of Haiti; Sallé.

In this species, encircled by a chestnut band, which is only seen in the last whorl, the spiral striae of the decussating sculpture are nearly obsolete.

Species 19. (Mus. Cuming.)

CHONDROPOMA BLAUNERI. *Chond. testā compressē umbilicatā, ovato-turrītā, tenuiculā, violaceo-fuscā, strigis undulatis fasciisque castaneis obscurē pictā, anfracti-*

bus decliei-converxis, longitudinaliter minutè filo-striatis, striis ad suturas auriculatis; aperturā obliquè oratā, supernè carinatā, labro vix reflexo.

BLAUNER'S CHONDROPOMA. Shell compressly umbilicated, ovately turreted, rather thin, violet-brown, obscurely marked with chestnut-brown waved streaks and bands, whorls slopingly convex, minutely longitudinally thread-striated, striae auriculated at the sutures; aperture obliquely ovate, keeled at the upper part, lip scarcely reflected.

Cyclotoma Blauneri, Shuttleworth, Diag. Moll. No. 7. p. 151.

Hab. Island of Porto Rico.

Of a peculiar swarthy violet-brown, obscurely painted with chestnut neatly-waved streaks and bands.

Species 20. (Mus. Cuming.)

CHONDROPOMA PLICATULUM. *Chond. testā compressē umbilicatā, subacuminato-turritatē, diaphano-albā, castaneo-fusco interruptē strigatā et fasciatā, anfractibus rotundatis, longitudinaliter densē filo-liratis; aperturā obliquè circulari, subsolutā, supernè carinatā, labro vix reflexo.*

THE FINELY-PLAITED CHONDROPOMA. Shell compressly umbilicated, rather acuminate turreted, diaphanous-white, interruptedly streaked and banded with chestnut-brown, whorls rounded, longitudinally densely thread-ridged; aperture obliquely circular, a little detached, keeled at the upper part, lip scarcely reflected.

Cyclotoma plicatum, Pfeiffer, Zeitschr. für Malac. 1846, p. 48.

Chondropoma plicatum, Pfeiffer.
Hab. Venezuela.

Rather more slenderly tubularly turreted than any of the previous species, painted with chestnut bands and streaks in a very characteristic interrupted manner.

Species 21. (Mus. Cuming.)

CHONDROPOMA SWIFTI. *Chond. testā compressē umbilicatā, orato-conicā, albida, rufo-fusco copiosè punctato-fasciatā, anfractibus concexis, longitudinaliter dense filo-liratis, liris ad suturas hic illic auriculatis; aperturā obliquè ovatā, supernè carinatā, labro tenui, vix reflexo.*

SWIFT'S CHONDROPOMA. Shell compressly umbilicated,

ovately conical, whitish, copiously dot-banded with red-brown, whorls convex, longitudinally closely thread-ridged, ridges here and there auriculated at the sutures; aperture obliquely ovate, keeled at the upper part, lip thin, scarcely reflected.

Cyclotoma Swifti, Shuttleworth, Diag. Moll. n. 7. p. 151.
Hab. Porto Rico; Swift.

A thin semidiaphanous shell, copiously encircled with bands of pale red-brown dots.

Species 22. (Mus. Cuming.)

CHONDROPOMA OTTONIS. *Chond. testā compressē umbilicatā, ovato-conicā, sordidè albā, subdiaphanā, pallidè fuscescente obscurè fasciatā, anfractibus rotundatis, minutè scabroso-decussatis; aperturā parvinsulā, rotundatā, latè fimbriatā.*

OTTO'S CHONDROPOMA. Shell compressly umbilicated, ovately conical, dull-white, subdiaphanous, faintly banded with light-brown, whorls rounded, minutely sebrously decussated; aperture rather small, rounded, broadly fimbriated.

Cyclotoma Ottonis, Pfeiffer, Zeitschr. für Malak. 1846, p. 45.

Chondropoma Ottonis, Pfeiffer.

Cyclotoma petricosum, Morelet.

Hab. Cuba.

A dull obscurely-banded shell, of scabrous surface, the whorls being rather tubular.

Species 23. (Mus. Cuming.)

CHONDROPOMA OBESUM. *Chond. testā compressē umbilicatā, oblongo-conicā, fulvo-luteā vel rubicundā, anfractibus rotundatis, liris filiformibus cingulatis, liris striis minutis scabrosis longitudinaliter decussatis; aperturā parvā, obliquè ovatā, labro simplici.*

THE STOUT CHONDROPOMA. Shell compressly umbilicated, oblong-conical, fulvous yellow or ruddy, whorls rounded, encircled with thread-like ridges, ridges crossed longitudinally by minute scabrous striae; aperture small, obliquely ovate, lip simple.

Truncatella obesa, Menke, Syn. ed. 2. p. 137.

Cyclotoma obesum, Pfeiffer.

Chondropoma obesum, Pfeiffer.

Hab. Cuba.

A very delicate ruddy fulvous shell, most exquisitely sculptured with minute scabrous striae.



25a



27.



25b



26.



27a



27b.



28.



29.



30.



31.

Slightly Enlarged

CHONDROPOMA.

PLATE IV.

Species 24. (Mus. Cuming.)

CHONDROPOMA CORDOVANUM. *Chond. testá compressè umbilicatá, turritá, pallidè fulvá, fasciis castaneis interruptis longitudinalibus tinctá, anfractibus rotundatis, densè longitudinaliter filo-striatis; aperturá circulari, parviusculá, labro scabroso-fimbriato.*

THE CORDOVA CHONDROPOMA. Shell compressly umbilicated, turreted, pale fulvous, stained with interrupted longitudinal chestnut bands, whorls rounded, densely longitudinally thread-striated; aperture circular, rather small, lip seaborously fimbriated.

PFEIFFER, Pro. Zool. Soc. 1856, p. 323.

Hab. Cordova, Vera Cruz, Mexico; Sallé.

A rather slenderly turreted species, imperfectly marked with interrupted longitudinal chestnut bands.

Species 25. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA INTEGRUM. *Chond. testá compressè umbilicatá, turritá, fulvá, lineis fuscis longitudinaliter undulatis rotatá, ad basin latifasciatá, anfractibus convexis, densè decussatim filatis, ad suturas crenatis; aperturá circulari, parviusculá, labro fimbriato.*

THE ENTIRE CHONDROPOMA. Shell compressly umbilicated, turreted, fulvous, marked with brown longitudinally waved lines, broad-banded at the base, whorls convex, closely decussately threaded, crenated at the sutures; aperture circular, rather small, lip fimbriated.

Cyclostoma integrum, Pfeiffer, Pro. Zool. Soc. 1851, p. 247.
Chondropoma integrum, Pfeiffer.

Hab. West Indies.

Very prettily painted throughout with delicate longitudinal brown lines, waved with great regularity.

Species 26. (Mus. Cuming.)

CHONDROPOMA LITTURATUM. *Chond. testá compressè umbilicata, cylindraceo-turritá, albida, purpurea punctata-fasciatá et lentiginosá, anfractibus declivi-convexis, undique reticulata-granosis, ad suturas crenatis; aperturá obliquè ovalá, parvá, dissolutá, labro simplici, vix reflexa.*

THE BLOTTED CHONDROPOMA. Shell compressly umbi-

licated, cylindrically turreted, whitish, dot-banded and freckled with purple, whorls slopingly convex, reticulately grained throughout, crenated at the sutures; aperture obliquely ovate, small, detached, lip simple, scarcely reflected.

Cyclostoma litturatum, Pfeiffer, Zeitsehr. für Malac. 1850, p. 78.

Chandropoma litturatum, Pfeiffer.

Hab. Island of Haiti; Sallé.

The sculpture of this species is a fine thimble-like grain, the colour a copious sprinkling of minute purple dots, disposed at the sutures in small flames and at the base in a band.

Species 27. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA DISSOLUTUM. *Chand. testá compressè umbilicata, fulvá, turritá, lineis ferrugineis longitudinaliter undulatá, ad basin fasciatá, anfractibus declivi-concisis, longitudinaliter densè filo-striatis; aperturá circulari, dissolutá, supernè carinatá, madicè fimbriata.*

THE DETACHED CHONDROPOMA. Shell compressly umbilicated, turreted, fulvous, longitudinally waved with rust-lines, banded at the base, whorls slopingly convex, longitudinally densely thread-striated; aperture circular, detached, keeled at the upper part, moderately fimbriated.

PFEIFFER, Malak. Blat. 1854, p. 158.

Hab. Cuba.

This species is probably a variety of *C. integrum*.

Species 28. (Mus. Cuming.)

CHONDROPOMA RUDE. *Chond. testá compressè umbilicata, solidiusculá, aurantio-carnea, ad apicem rosacea, intus rufo-aurantia, anfractibus decliri-canvexis, spiraliiter liratis, longitudinaliter densè subtilissimè filo-striatis, subscabrosis; aperturá obliquè ovalá, dissolutá, supernè carinatá, labro calloso, vix reflexo.*

THE RUDE CHONDROPOMA. Shell compressly umbilicated, rather solid, orange-flesh, pink at the apex, reddish-orange within, whorls slopingly convex, spirally ridged, longitudinally densely very finely thread-striated, rather scabrous; aperture obliquely ovate,

detached, keeled at the upper part, lip callous, scarcely reflected.

WEINLAND, Malak. Blat. 1862.

Hab. Bahama Islands.

Very distinct from any of the preceding species, but of precisely the same generic type. It is of rather solid growth and very characteristic in colour—a bright flesh-orange, without any marking, pink towards the apex, orange-red in the interior.

Species 29. (Mus. Cuming.)

CHONDROPOMA ANGUSTATUM. *Chond. testā compressē umbilicatā, subangustē turritā, fulvā, maculis linearibus purpureo-castaneis fasciatā, anfractibus convexis reticulatum filo-liratis, ad suturas auriculato-crenatis; aperturā parvā, obliquè circulari, labro subreflexo.*

THE NARROWED CHONDROPOMA. Shell rather narrowly turreted, fulvous, banded with linear blotches of purple-chestnut, whorls convex, reticulately thread-ridged, auriculately crenated at the sutures; aperture small, obliquely circular, lip a little reflected.

PFEIFFER, Pro. Zool. Soc. 1854, p. 128.

Hab. —?

The habit of *C. angustum* is not known, but there can be little doubt of its being a West Indian species.

Species 30. (Mus. Cuming.)

CHONDROPOMA PERLATUM. *Chond. testā compressē umbilicatā, subangustē turritā, margaritacea, diaphanā, maculis fuscis inter liras strigatā, anfractibus rotundatis, liris tenuibus nitide spiraliter funiculatis; aperturā obliquè ovatā, scabroso-fimbriatā.*

THE BROUGHT CHONDROPOMA. Shell compressly umbilicated, rather narrowly turreted, nearly white, diaphanous, streaked with brown spots between the ridges, whorls rounded, neatly spirally corded with fine ridges; aperture obliquely ovate, seaborously fimbriated.

Cyclostoma perlatum, Gundlach, Malak. Blat. 1857, p. 41.

Chondropoma perlatum, Pfeiffer.

Hab. Cuba.

An extremely delicate shining diaphanous shell, stringed transversely with very fine pellucid ridges.

Species 31. (Mus. Cuming.)

CHONDROPOMA TORTOLENSE. *Chond. testā compressē umbilicatā, orato-turritā, albida, maculis rufo-fuscis inter liras sparsim strigatā, anfractibus rotundatis, ad suturas minutē crenulatis, liris tenuibus spiraliter funiculatis, in interstitiis puncturatis; aperturā obliquè ovatā, subdissolutā, supernē carinatā, labro simplici, rix reflexo.*

THE TORTOLA CHONDROPOMA. Shell compressly umbilicated, ovately turreted, whitish, sparsely streaked with red-brown spots between the ridges, whorls rounded, minutely crenulated at the sutures, spirally corded with fine ridges, punctured in the interstices; aperture obliquely ovate, a little detached, keeled at the upper part, lip simple, scarcely reflected.

PFEIFFER, Malak. Blat. 1857, p. 158.

Hab. Island of Tortola.

The sculpture and marking of this shell is not much unlike that of the preceding, but coarser and more opake. The lip is of quite another character, not being fimbriated.



(Slightly Enlarged)

Sowerby, del et lith

Vincent Brooks, imp

CHONDROPOMA.

PLATE V.

Species 32. (Mus. Cuming.)

CHONDROPOMA DIAPHANUM. *Chond. testā compressē umbilicatā, oblongo-turritā, candidā, anfractibus convexis, subangustis, pulcherrimē reticulatim filo-striatis, ad suturas promiscuē crenulatis; aperturā obliquē circulari, dissolutā, subfimbriatā.*

THE DIAPHANOUS CHONDROPOMA. Shell compressly umbilicated, oblong-turreted, white, whorls convex, rather narrow, beautifully reticulately thread-striated, promiscuously crenulated at the sutures; aperture obliquely circular, detached, slightly fimbriated.

Cyclostoma diaphanum, Pfeiffer, Pro. Zool. Soc. 1851.

Chondropoma diaphanum, Pfeiffer.

Hab. —?

A pure white *Scalaria*-like shell, exquisitely reticulately sculptured with thread-striæ.

Species 33. (Mus. Cuming.)

CHONDROPOMA DELATREANUM. *Chond. testā compressē umbilicatā, subangustē turritā, fulvā, purpureo-castaneo strigatā et fasciatā, anfractibus modicē convexis, densē reticulatim filo-striatis, ad suturas conspicuē crenulatis; aperturā parvā, obliquē circulari, labro fimbriato, radiato.*

DELATRE'S CHONDROPOMA. Shell compressly umbilicated, rather narrowly turreted, fulvous, streaked and banded with purple-chestnut, whorls moderately convex, closely reticulately thread-striated, conspicuously crenulated at the sutures; aperture small, obliquely circular, lip fimbriated, rayed.

Cyclostoma Delatreanum, D'Orbigny, Moll. Cub. p. 262.

Chondropoma Delatreanum, Pfeiffer.

Hab. Cuba.

A rather narrowly turreted shell, neatly reticulately sculptured and conspicuously cross-banded.

Species 34. (Mus. Cuming.)

CHONDROPOMA PETITIANUM. *Chond. testā compressē umbilicatā, angustē turritā, sericeo-fulvā, pallidē rufō-fusco lentiginosā, anfractibus convexis, densē longitudinaliter filo-striatis, striis hic illuc partim opaco-*

albis; aperturā parvā, obliquē circulari, labro fimbriato, conspicuē radiato.

PETIT'S CHONDROPOMA. Shell compressly umbilicated, narrowly turreted, silky-fulvous, freckled with pale red-brown, whorls convex, densely longitudinally thread-striated, striae here and there partially opake-white; aperture small, obliquely circular, lip fimbriated, conspicuously rayed.

Cyclostoma Petitianum, Pfeiffer, Zeitschr. für Malae. 1850, p. 78.

Chondropoma Petitianum, Pfeiffer.

Hab. Island of Haiti; Sallé.

A bright transparent silky fulvous shell, freckled with red-brown, in which the longitudinal thread-striae have a partially opake-white aspect, as seen in some of the *Clausiliæ*.

Species 35. (Mus. Cuming.)

CHONDROPOMA PFEIFFERIANUM. *Chond. testā modicē umbilicatā, oblongo-turritā, sordidē albida, rufō-fusco inter liras strigatā, anfractibus declivi-convexis, undique spiraliter filo-liratis, versus apicem minutē obliquē striatis; aperturā obliquē ovatā, supernē carinatā, labro vix reflexo.*

PFEIFFER'S CHONDROPOMA. Shell moderately umbilicated, oblong turreted, dull white, streaked between the ridges with red-brown, whorls slopingly convex, spirally thread-ridged throughout, minutely obliquely striated towards the apex; aperture obliquely ovate, keeled at the upper part, lip scarcely reflected.

Cyclostoma Pfeifferianum, Poey, Mem. p. 419.

Chondropoma Pfeifferianum, Pfeiffer.

Hab. Cuba.

The longitudinal striae of this species, which are somewhat crowded towards the apex, do not cross the spiral ridges, but appear only in the interstices.

Species 36. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA CUMANENSE. *Chond. testā angustē umbilicatā, oblongo-turritā, pellucido-fulvā, castaneo interrupte strigatā et fasciatā, anfractibus declivi-convexis, longitudinaliter densē filo-liratis; aperturā obliquē circulari, parvā, labro vix reflexo.*

THE CUMANA CHONDROPOMA. Shell narrowly umbilicated, oblong-turreted, transparent fulvous, interruptedly streaked and banded with chestnut, whorls slopingly convex, longitudinally densely threadridged; aperture obliquely circular, small, lip scarcely reflected.

Cyclostoma Cumanense, Pfeiffer, Pro. Zool. Soc. 1851.

Chondropoma Cumanense, Pfeiffer.

Hab. Cumana, Venezuela; Dyson.

An extremely pretty species, of bright diaphanous fulvous substance, richly stained with interrupted bands and streaks of chestnut.

Fig. 37. (Mus. Cuming.)

The shell here figured proves to be a variety, more distinctly coloured, of *C. Poeyorum*, represented at Plate III. Fig. 15.

Species 38. (Fig. *a, b*, Mus. Cuming.)

CHONDROPOMA DENTATUM. *Chond. testā compressē umbilicatā, fulvā, maculis rufo-fuscis parvis equidistantibus, tessellatā, anfractibus convexis, densē granoso-reticulatis, ad suturas dentato-crenatis; aperturā parvā, obliquē circulari, labro fimbriato-reflexo, radiato.*

THE TOOTHED CHONDROPOMA. Shell compressly umbilicated, fulvous, tessellated with small equidistant red-brown spots, whorls convex, densely granosely reticulated, dentately crenated at the sutures; aper-

ture small, obliquely circular, lip fimbriately reflected, rayed.

Cyclostoma dentatum, Say, Philad. Journ. vol. v. p. 125.

Cyclostoma lineolatum, Anton.

Cyclostoma Auberianum, D'Orbigny.

Cyclostoma lunulatum, Murch.

Chondropoma dentatum, Pfeiffer.

Hab. Cuba and Florida, United States.

The sutural crenulations of this species are unusually prominent, having the appearance of teeth.

Species 39. (Mus. Cuming.)

CHONDROPOMA RUFO-PICTUM. *Chond. testā vix umbilicatā, ovato-conicā, subventricosā, fulvā, rufo-castaneo linearī-fasciatā, anfractibus declivi-convexis, longitudinaliter plicato-striatis, striis subdistantibus, irregularibus; aperturā obliquē ovalā, labro parum reflexo.*

THE RED-PAINTED CHONDROPOMA. Shell scarcely umbilicated, ovately conical, rather ventricose, fulvous, linearly banded with reddish-chestnut, whorls slopingly convex, longitudinally plicately striated, striae rather distant, irregular; aperture obliquely ovate, lip but little reflected.

GUNDLACH, MS. in Mus. Cuming.

Hab. Cuba.

A rather ventricose species, in which the longitudinal thread-like striae assume the form of promiscuous rather distant folds.



40.



41.



42.



43.



44a.



44b.



45.



46a.



46b.



47.

(Slightly Enlarged.)

CHONDROPOMA.

PLATE VI.

Species 40. (Mus. Cuming.)

CHONDROPOMA CANDEANUM. *Chond. testā vix umbilicatā, oblongo-turritatā, rufescente-fulgā, anfractibus convexis, validē densē filo-reticulatis; aperturā obliquè ovalā, labro subfimbriato.*

THE CANDE CHONDROPOMA. Shell scarcely umbilicated, oblong-turreted, reddish fulvous, whorls convex, strongly closely thread-reticulated; aperture obliquely ovate, lip slightly fimbriated.

Cyclostoma Candeanum, D'Orbigny, Moll. Cuba, p. 261. pl. 22. f. 15 to 17.

Cyclostoma truncatum, Pfeiffer.

Cistula truncata, Gray.

Cistula Candeania, Pfeiffer.

Chondropoma Candeanum, Pfeiffer.

Hab. Cuba.

A rather stout reddish-fulvous shell, with the surface very strongly finely reticulated.

Species 41. (Mus. Cuming.)

CHONDROPOMA PUNCTICULATUM. *Chond. testā compressè umbilicatā, ovato-oblongā, candidā, punctis parvis rufofuscis tessellatā, anfractibus convexis, nitidē densē filo-reticulatis; aperlurā parvā, obliquè circulari, subdisolutā, supernè carinatā, labro subreflexo.*

THE FINELY-DOTTED CHONDROPOMA. Shell compressly umbilicated, ovately oblong, white, tessellated with small red-brown dots, whorls convex, neatly closely thread-reticulated; aperture small, obliquely circular, a little detached, keeled at the upper part, lip somewhat reflected.

PFEIFFER, Pro. Zool. Soc. 1854, p. 128.

Hab. West Indies.

A strongly finely reticulated white shell, tessellated with distant small red-brown dots.

Species 42. (Mus. Cuming.)

CHONDROPOMA SIMPLEX. *Chond. testā minutè compressè umbilicatā, ovato-oblongā, fulvescente-albā, subdiaphanā, maculis rufescente-fuscis in seriebus longitudinalibus piclā, anfractibus tumido-convexitatis, validē densē*

filo-reticulatis; aperturā parvā, obliquè ovalā, labro simplici.

THE SIMPLE CHONDROPOMA. Shell minutely compressly umbilicated, ovately oblong, fulvous white, somewhat diaphanous, painted with longitudinal rows of red-brown spots, whorls tumidly convex, strongly densely thread-reticulated; aperture small, obliquely ovate, lip simple.

PFEIFFER, Pro. Zool. Soc. 1852.

Hab. Island of Haiti; Sallé.

The surface sculpture of this species has a beautifully delicate pearly aspect.

Species 43. (Mus. Cuming.)

CHONDROPOMA ABNATUM. *Chond. testā minutè compressè umbilicatā, oblongo-turritatā, solidiusculā, sordidē albā, anfractibus declivi-convexitatis, subtiliter scabroso-reticulatis, ad suturas canaliculatis et crenulatis; aperturā parvā, obliquè circulari, labro auriculato-fimbriato.*

THE ABNATE CHONDROPOMA. Shell minutely compressly umbilicated, oblong-turreted, rather solid, dead-white, whorls slopingly convex, finely scabrously reticulated, channelled and crenulated at the sutures; aperture small, obliquely circular, lip auriculately fimbriated.

GUNDLACH, MS. in Mus. Cuming.

Hab. Cuba.

This is the first species in which the suture is distinctly channeled throughout.

Species 44. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA LOWEANUM. *Chond. testā compressè umbilicatā, subventricoso-ovatā, albida, pallide fusco marmorata et strigata, ad basin fasciatā, anfractibus tumido-convexitatis, longitudinaliter pulcherrimè densē filo-striatis; aperturā obliquè ovalā, dissolutā, supernè carinatā et crenulatā, labro vix reflexo.*

LOWE'S CHONDROPOMA. Shell compressly umbilicated, somewhat ventricosely ovate, white, marbled and streaked with pale-brown, banded at the base, whorls tumidly convex, longitudinally very beautifully densely thread-striated; aperture obliquely ovate, detached,

keeled and crenulated at the upper part, lip scarcely reflected.

PFEIFFER, Pro. Zool. Soc. 1852.

Hab. Island of Haiti; Sallé.

An exquisitely sculptured species, more ovately ventricose than any which has preceded it.

Species 45. (Mus. Cuming.)

CHONDROPOMA VENEZUELENSE. *Chond. testá modicè umbilicatá, acuminato-turritá, ustulato-fulvá, rufo-fusco interrupte linearí-fasciatá, anfractibus convexis, longitudinaliter densè filo-striatis; aperturá circulari-ovata, labro patente-expanso.*

THE VENEZUELA CHONDROPOMA. Shell moderately umbilicated, acuminately turreted, swarthy fulvous, interruptedly linearly banded with red-brown, whorls convex, longitudinally densely thread-striated; aperture circularly ovate, lip openly expanded.

PFEIFFER, Pro. Zool. Soc. 1853, p. 52.

Hab. Venezuela.

A thin acuminately turreted shell, of swarthy hue, in which the aperture is openly expanded at the margin.

Species 46. (Fig. *a, b*, Mus. Cuming.)

CHONDROPOMA REVINCTUM. *Chond. testá compressè umbilicatá, oblongo-turritá, livido-fuscescente, maculis punctisque opaco-albis, purpureo-umbratis, cingulatá et tentiginosá, anfractibus convexis, densè filo-reticulatis, ad suturas crenulatis; aperturá circulari, labro subfimbriato.*

THE BOUND CHONDROPOMA. Shell compressly umbilicated, oblong-turreted, livid brown, encircled and freckled with purple-shaded opake-white spots and dots, whorls convex, closely thread-reticulated, crenulated at the sutures; aperture circular, lip slightly fimbriated.

Cyclostoma revinctum, Poey, Mem. p. 99. pl. 5. ff. 24 to 27.

Chondropoma revinctum, Pfeiffer.

Hab. Cuba.

Painted with very characteristic purple-shaded opake-white spots and dots upon a dingy fuscous ground.

Species 47. (Mus. Cuming.)

CHONDROPOMA RUBICUNDUM. *Chond. testá vix umbilicatá, acuminato-turritá, solidiusculá, pellucidá, vinoso-rubicundá, anfractibus rotundatis, striis filiformibus annulatis; aperturá circulari, labro fimbriato, intensè roseo.*

THE RUDDY CHONDROPOMA. Shell scarcely umbilicated, acuminately turreted, rather solid, transparent, vinous-red, whorls rounded, ringed with thread-like striae; aperture circular, lip fimbriated, deep rose-colour.

Cyclostoma rubicundum, Morelet, Test. Noviss. p. 22.

Chondropoma rubicundum, Pfeiffer.

Hab. Vera Paz, Guatemala.

This Guatemala species is of quite a distinct type from any other. The shell, though solid, is of a rich subglistening transparency, of a vinous-red colour, while the fimbriated lip is a deep crimson-rose.



48.



49a.



49b.



50



51a



51b.



52



53.



54



55

(Slightly Enlarged.)

CHONDROPOMA.

PLATE VII.

Species 48. (Mus. Cuming.)

CHONDROPOMA ARGUTUM. *Chond. testā minutē compressē umbilicatā, angustē turritā, solidiusculā, sordidē albū, anfractibus convexis, conspicuè spiraliter carinato-costatis, interstītiis excavatis; aperturā parvā, circulari, subdissolutā, labro simplici.*

THE NEAT CHONDROPOMA. Shell minutely compressly umbilicated, narrowly turreted, rather solid, dead-white, whorls convex, conspicuously spirally keel-ribbed, with the interstices excavated; aperture small, circular, a little detached, lip simple.

PFEIFFER, Monog. Pneum. Suppl. I. p. 138.

Hab. —?

This shell is of quite a distinct type. The sculpture is like that of a sea shell, strong encircling keel ribs, excavated in the interstices like those of many species of *Pleurotoma*.

Species 49. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA CRENULATUM. *Chond. testā compressē umbilicatā, oblongo-turritā, fulvā, usculato-castaneo varie strigata et nebulata, anfractibus convexis, longitudinaliter sebrosè filo-striatis, ad suturas crenulatis; aperturā obliquè circulari, labro tenui, subexpanso.*

THE CRENULATED CHONDROPOMA. Shell compressly umbilicated, oblong-turreted, fulvous, variously streaked and clouded with burnt chestnut, whorls convex, longitudinally sebrously thread-striated, crenulated at the sutures; aperture obliquely circular, lip thin, a little expanded.

Cyclostoma crenulatum, Féussac, Pot. et Mich. Gal. Douai, vol. i. p. 235. pl. 24. ff. 3, 4.

Cyclostoma nodulosum, Gray.

Cyclostoma Guadaloupense, Pfeiffer.

Chondropoma Guadaloupense, Pfeiffer.

Chondropoma crenulatum, Pfeiffer.

Hab. Guadaloape; Petit.

A swarthy clouded shell, in which the sutural crenulations are white, and therefore rather conspicuous.

Species 50. (Mus. Cuming.)

CHONDROPOMA SANTACRUZENSE. *Chond. testā compressē umbilicatā, oblongo-turritā, tenuiculatā, aurantio-*

rubellā, castaneo minutē punctato-fasciatā, anfractibus convexis, subventricosis, densē sebrosè lamelli-striatis; aperturā obliquè circulari, labro tenui, subexpanso.

THE SANTA CRUZ CHONDROPOMA. Shell completely umbilicated, oblong-turreted, rather thin, orange-red, minutely dot-banded with chestnut, whorls convex, slightly ventricose, closely sebrously lamella-striated; aperture obliquely circular, lip thin, a little expanded.

PFEIFFER, Malak. Blat. 1855, p. 101.

Hab. Island of Santa Cruz, West Indies.

Beautifully sculptured with a close succession of longitudinal lamella-like striæ, sharp and sebrous at the edge.

Species 51. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA TEXTUM. *Chond. testā compressē umbilicatā, conico-turritā, solidiusculā, pallidē stramineā, sparsē obscurè rufo-punctatā, anfractibus declivi-convexis, densē reticulatim filo-striatis, ad suturas crenulatis; aperturā circulari-oratā, labro calloso-fimbriato.*

THE WOVEN CHONDROPOMA. Shell compressly umbilicated, conically turreted, rather solid, pale straw-colour, sparsely obscurely red-dotted, whorls slopingly convex, closely reticulately thread-striated, crenulated at the sutures; aperture circularly ovate, lip callously fimbriated.

GUNDLACH, Malak. Blat. 1862.

Hab. Cuba.

A very characteristic straw-coloured shell, of rather solid growth, with a prominent callously fimbriated aperture.

Species 52. (Mus. Cuming.)

CHONDROPOMA TURRITUM. *Chond. testā compressē umbilicatā, acuminato-turritā, fulvā, fasciis rufo-fuscis linearibus interruptis ciectā, anfractibus rotundatis, minutē reticulatim filo-striatis, ad suturas crenulatis; aperturā obliquè circulari, labro tenui, subexpanso.*

THE TURRETED CHONDROPOMA. Shell compressly umbilicated, acuminately turreted, fulvous, encircled with interrupted linear red-brown bands, whorls rounded, minutely reticulately thread-striated, crenulated at

the sutures; aperture obliquely circular, lip thin, a little expanded.

Cyclotoma turritum, Pfeiffer, Pro. Zool. Soc. 1851.

Chondropoma turritum, Pfeiffer.

Hab. Honduras; Dyson.

Of an obvious acuminate turreted form, with the characteristic net-sculpture of the genus unusually finely woven.

Species 53. (Mus. Cuming.)

CHONDROPOMA REVOCATUM. *Chond. testā modicē umbilicatā, globoso-conicā, flavescente, versus apicem carneā, punctis rufis obscurè radiatā, anfractibus rotundatis, spiraliter granoso-scabrosē filo-striatis; aperturā pyriformi-ovatā, subamplā, labro tenuē expanso.*

THE RECALLED CHONDROPOMA. Shell moderately umbilicated, globosely conical, yellowish, flesh-tinged towards the apex, obscurely rayed with red dots, whorls rounded, encircled with granosely seaborous thread-ridges; aperture pyriformly ovate, rather large, lip thinly expanded.

Cyclotoma revocatum, Gundlach, Malak. Blat. 1857, p. 178.

Chondropoma revocatum, Pfeiffer.

Hab. Cuba.

Distinguished by a light globosely inflated form, pale-yellow or straw-colour, passing towards the apex into flesh-rose.

Species 54. (Mus. Cuming.)

CHONDROPOMA EUSARCUM. *Chond. testā compressē umbilicatā, ovatā, subventricosā, rubello-stramineā, fasciis pallidē rufis interruptis cinctā, anfractibus convexis,*

subtiliter longitudinaliter filo-striatis; aperturā obliquē ovalatā, subdissolutā, supernē carinatā, labro tenui, rix reflexo.

THE DELICATE CHONDROPOMA. Shell compressly umbilicated, ovate, rather ventricose, ruddy straw-colour, encircled with faint linear red bands, whorls convex, finely longitudinally thread-striated; aperture obliquely ovate, a little detached, keeled at the upper part, lip thin, scarcely reflected.

PFEIFFER, Pro. Zool. Soc. 1852.

Hab. Island of Haiti; Sallé.

The most characteristic feature of this species is its light ovate form, with the last whorl ventricose, and proportionately larger than usual.

Species 55. (Mus. Cuming.)

CHONDROPOMA HARPA. *Chond. testā modicē umbilicatā, subtubuloso-turritā, pallidē cinnamomeā, rufescente punctato-fasciatā, anfractibus rotundatis, longitudinaliter membranacius filo-striatis, striis subdistantibus, irregularibus; aperturā obliquē circulari, labro reflexo, rosaceo.*

THE HARP CHONDROPOMA. Shell moderately umbilicated, somewhat tubularly turreted, pale cinnamon-colour, dot-banded with red-brown, whorls rounded, longitudinally membranaceously thread-striated, striae rather distant, irregular; aperture obliquely circular, lip reflected, rose-tinted.

Cyclotoma harpa, Pfeiffer, Pro. Zool. Soc. 1851.

Chondropoma harpa, Pfeiffer.

Hab. Havana.

A very delicate cinnamon-coloured shell, of rather tubular growth, in which the striae are like membranaceous threads, irregular and rather distant from each other.



(Slightly Enlarged)

Sowerby, del. & lith.

Vincent Brooks, Imp.

CHONDROPOMA.

PLATE VIII.

Species 56. (Mus. Cuming.)

CHONDROPOMA LATUM. *Chond. testā ampliter excavato-umbilicatā, depresso-globosā, stramineā, anfractibus rotundatis, obliquē eximiē filo-striatis, ad umbilicum conspicue carinatis; aperturā circulari, partim fimbriatā.*

THE BROAD CHONDROPOMA. Shell largely excavately umbilicated, depressly globose, straw-colour, whorls rounded, obliquely delicately thread-striated, conspicuously keeled at the umbilicus; aperture circular, partially fimbriated.

GUNDLACH, MS. in Mus. Cuming.

Hab. Cuba.

A very distinct form of the genus, in which the shell is depressly globose and largely deeply umbilicated, the umbilicus being prominently keeled throughout.

Species 57. (Fig. a, Mus. Cuming.)

CHONDROPOMA MARINUM. *Chond. testū subcompressē umbilicatā, oblongo-turrīta, fulvī, punctis castaneis in seriebus duplicitibus longitudinaliter strigatā, ad basin fasciatā, anfractibus convexis, longitudinaliter dense filo-liratis, ad suturas crenulatis; aperturā obliquē ovatā, subexpansa, radiatā.*

THE MARINE CHONDROPOMA. Shell rather compressly umbilicated, oblong-turreted, fulvous, longitudinally streaked with double rows of chestnut dots, banded at the base, whorls convex, longitudinally densely thread-ridged, crenulated at the sutures; aperture obliquely ovate, a little expanded, rayed.

WEINLAND, Malak. Blat. 1862.

Hab. Island of Haiti; Weinland.

Probably named *marinum* from the circumstance of its being found in the vicinity of the sea.

Species 58. (Mus. Cuming.)

CHONDROPOMA BASICARINATUM. *Chond. testū compressē umbilicatā, oblongo-turrīta, sordidē albā, semifossili, anfractibus plano-convexis, spiraliter superficialiter liratis, striis obliquis scabrosis transgredientibus, anfractu ultimo busi carinato; aperturā obliquē circulari, dissolutā, supernē carinata, labro subfimbriato.*

THE BASE-KEELED CHONDROPOMA. Shell compressly

umbilicated, oblong-turreted, dead white, semifossil. whorls flatly convex, spirally superficially ridged, with oblique seaborous striae passing over them, last whorl keeled at the base; aperture obliquely circular, detached, keeled at the upper part, lip slightly fimbriated.

Cyclostoma basicarinatum, Pfeiffer, Malak. Blat. 1855, p. 101. pl. 4. f. 2, 3.

Hab. Cuba.

A dead-white semifossil shell, belonging to a type of the genus, of which there are many other species of smaller size.

Species 59. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA MARGINALBUM. *Chond. testū subclavide-umbilicatā, cylindraceo-oblongā, solidiusculū, pellucido-rosaceū, anfractibus plano-convexis, unilique minutē filo-reticulatis, ad suturas subcrenulatis; aperturā obliquē circulari-ovatā, labro callosō-fimbriato, albo.*

THE WHITE-EDGED CHONDROPOMA. Shell with a nearly-closed umbilicus, cylindrically oblong, rather solid, transparent rose-tinted, whorls flatly convex, minutely thread-reticulated throughout, slightly crenulated at the sutures; aperture obliquely circularly ovate, lip callously fimbriated, white.

GUNDLACH, MS. in Mus. Cuming.

Hab. Cuba.

A very distinct form of *Chondropoma*, of solid growth, but yet of transparent rose-colour, white at the lip.

Species 60. (Mus. Cuming.)

CHONDROPOMA QUATERNATUM. *Chond. testū subclavide-umbilicatā, cylindraceo-oblongā, flavescente-albā, interdum rosaceo tinctā, interdum purpureo-fuscā supra suturas, anfractibus plano-convexis, longitudinaliter dense costulato-striatis; aperturā parvā, obliquē ovata, labro callosō-reflexo.*

THE QUATERNATE CHONDROPOMA. Shell with a nearly-closed umbilicus, cylindrically oblong, yellowish-white, sometimes tinged with rose, sometimes purple-brown above the sutures, whorls flatly convex, longitudinally closely rib-striated, aperture small, obliquely ovate, lip callously reflexed.

Cyclostoma quaternatum, Lamarek, Anim. sans vert. vol. viii. p. 358.
Cistula quaternata, Gray.
Tudora quaternata, Pfeiffer.
Hab. —?

Allied to the preceding species in typical form.

Species 61. (Mus. Cuming.)

CHONDROPOMA HJALMERSONI. *Chond. testā obtusē cylindraceā, fulvescente, obscurē rufo-punctatā, anfractibus convexis, decussatim obsoletē liratis et malleatis; aperturā obliquē ovatā, labro subexpanso.*

HJALMERSON'S CHONDROPOMA. Shell obtusely cylindrical, fulvous, faintly red-dotted, whorls convex, decussately obsoletely ridged and malleated, aperture obliquely ovate, lip a little expanded.

PFEIFFER, MS. in Mus. Cuming.

Hab. —?

A peculiar *Pupina*-like shell, dull in colour, with the surface obsoletely decussately ridged and malleated.

Species 62. (Mus. Cuming.)

CHONDROPOMA BLANDUM. *Chond. testā subclaudente-umbilicatā, ovato-turritā, fusco-violaceā, punctis rufis obscurē strigatā, anfractibus convexis, longitudinaliter densē costulato-striatis, versus apicem scabrosis; aperturā obliquē ovatā, auriculato-fimbriatā.*

THE FAIR CHONDROPOMA. Shell with a nearly-closed umbilicus, ovately turreted, brownish-violet, obscurely streaked with red dots, whorls convex, longitudinally densely rib-striated, subealous towards the apex, aperture obliquely ovate, auriculately fimbriated.

PFEIFFER, Pro. Zool. Soc. 1852.

Hab. Island of Haiti; Sallé.

This and the three preceding species partake in nearly equal proportion of the characters of *Chondropoma*, *Cistula*, and *Tudora*, and might be referred to either genus.

Species 63. (Mus. Cuming.)

CHONDROPOMA PAPYRACEUM. *Chond. testā compressē umbilicatā, ovato-turritā, tenuiculā, pallidē fuscā, castaneo punctatā et undatā, anfractibus convexis, longitudinaliter densē filo-striatis, ad suturas crenulatis: aperturā obliquē ovatā, labro expanso, reflexo.*

THE PAPER CHONDROPOMA. Shell compressly umbilicated, ovately turreted, rather thin, light fuscous, dotted and waved with chestnut, whorls convex, longitudinally densely thread-striated, crenulated at the sutures, aperture obliquely ovate, lip expandedly reflected.

Cyclostoma papyraceum, C. B. Adams, Contr. no. 6. p. 92.

Chondropoma papyraceum, Pfeiffer.

Tudora papyracea, Pfeiffer.

Hab. Jamaica.

A very elaborately thread-striated shell, rather more inflated in the last whorl than usual.




65a.



65b.



66.



67a.




67b.



68.



69.



70.



71.

CHONDROPOMA.

PLATE IX.

Species 64. (Mus. Cuming.)

CHONDROPOMA AGASSIZII. *Chond. testā subprofundē umbilicatā, globoso-turbinatā, rnbello-stramineā, castaneo irregulariter maculatā, suturis angustē canaliculatis, anfractibus rotundatis, spiraliter liratis; aperturā pyriformi-circulari, labro subexpanso.*

AGASSIZ' CHONDROPOMA. Shell rather deeply umbilicated, globosely turbinated, reddish straw-colour, irregularly spotted with chestnut, sutures narrowly channelled, whorls rounded, spirally ridged; aperture pyriformly circular, lip a little expanded.

Cyclostoma Agassizii, Charpentier, Küster, Conch. Cab. pl. 38. f. 1, 2.

Cistula Agassizii, Pfeiffer.

Hab. Cuba.

Of a bright, warm straw-colour, variously freckled with chestnut spots and dots.

Species 65. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA TERNATUM. *Chond. testā subampliter profundē umbilicatā, globoso-turbinatā, stramineā, hyalinā, rno-fusco maculatā, suturis angustē canaliculatis, anfractibus declivi-rotundatis, spiraliter tenui-liratis, ultimo subinflato, aperturā circulari, labro latē plano-fimbriato.*

THE TERNATE CHONDROPOMA. Shell rather largely deeply umbilicated, globosely turbinated, straw-colour, hyaline, spotted with reddish-brown, sutures narrowly channelled, whorls slopingly rounded, spirally finely ridged, last whorl rather inflated, aperture circular, lip broadly fimbriated.

Cyclostoma ternatum, Gould, MS. in Mus. Cuming.

Hab. —?

Very like the preceding species in sculpture and colouring, but a more inflatedly convoluted hyaline shell, with the lip broadly fimbriated.

Species 66. (Mus. Cuming.)

CHONDROPOMA SOLARE. *Chond. testā imperforatā, conico-turritā, purpureo-cinereā, albo tenuifuniculata, anfractibus convexis, spiraliter angustē liratis, inter-*

stitiis longitudinaliter minutē crenulatis, apertura ovato-circulari, labro latissimē plicato-fimbriato.

THE SOLAR CHONDROPOMA. Shell imperforate, conically turreted, purple-ash, finely corded with white, whorls convex, spirally narrowly ridged, interstices longitudinally minutely crenulated, aperture ovately circular, lip encircled with a very broad puckered frill.

PFEIFFER, MS. in Mus. Cuming.

Hab. —?

This fine species, lately named by Dr. Pfeiffer, is finely corded throughout with white ridges upon a purple-ash ground, which is most exquisitely crenulated in the opposite direction. The bold plated frill encircling the aperture, except on the side where it would be obstructed by the body whorl, is an equally striking character.

Species 67. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA SAGITTIFERUM. *Chond. testā umbilicatā, ovato-turritā, sordidē albā, ferrugineo-fusco longitudinaliter undatā, anfractibus declivi-convexit, liris plano-obtusis densissimē reticulatis; aperturā pyriformi-ovatā, labro plus minus fimbriato.*

THE ARROW-BEARING CHONDROPOMA. Shell umbilicated, ovately turreted, dull white, longitudinally waved with rust-brown, whorls slopingly convex, very densely reticulated with flatly obtuse ridges, aperture pyriformly ovate, lip more or less fimbriated.

Cyclostoma sagittiferum, C. B. Adams, Contr. Conch. no. 8. p. 137.

Cistula? sagittifera, Pfeiffer.

Hab. Jamaica.

The sculpture of this shell is a close obtuse reticulation, with the interstices mere punctures.

Species 68. (Mus. Cuming.)

CHONDROPOMA GRUNERI. *Chond. testā angustē umbilicatā, oblongo-conicā, sordidē flavescente, subhyalinā, pose aperturam sparsē fusco-fusciatā, anfractibus declivi-convexit, spiraliter elevato-striatis; aperturā ovato-circulari, labro latissimē concavo-fimbriato.*

GRUNER'S CHONDROPOMA. Shell narrowly umbilicated,

oblong-conical, dull yellowish, subhyaline, sparsely banded with brown behind the aperture, whorls slopingly convex, spirally elevately striated, aperture ovately-circular, lip very broadly concavely frilled.

Cyclostoma Grunerii, Pfeiffer, Zeitschr. für Malac. 1846, p. 47.

Cistula Grunerii, Gray.

Hab. Honduras.

Of a peculiar dull, delicately transparent substance, faintly sculptured.

Species 69. (Mus. Cuming.)

CHONDROPOMA RADIOSUM. *Chond. testā vix umbilicatā, angustē turbinatā, rubello-stramineā, pallidē fusco interruptē lineatā, anfractibus convexis, longitudinaliter densē crispato-striatis; aperturā parvā, labro latē concavo-fimbriato, fimbriā plicatā, fusco-radiatā.*

THE RAYED CHONDROPOMA. Shell scarcely umbilicated, narrowly turbinated, reddish straw colour, faintly interruptedly lined with brown, whorls convex, longitudinally densely crisply striated; aperture small, lip broadly concavely frilled, frill puckered, rayed with brown.

Cyclostoma radiosum, Morelet, Test. Noviss. p. 22.

Cistula radiosum, Pfeiffer.

Hab. Guatemala.

Exquisitely sculptured longitudinally, and beautifully rayed with brown behind the aperture.

Species 70. (Mus. Cuming.)

CHONDROPOMA ROSTRATUM. *Chond. testā umbilicatā, subangustē turbinatā, albida, subhyalinā, fusco longitudinaliter alternatim undata et punctata, anfractibus*

convexit, creberrimè longitudinaliter funiculatis, in suturas denticulatis; aperturā ovato-circulari, labro calloso-fimbriato.

THE BEAKED CHONDROPOMA. Shell umbilicated, rather narrowly turbinated, whitish, subhyaline, alternately longitudinally waved and dotted with brown, whorls convex, very closely longitudinally corded, dentieulated in the sutures, aperture ovately circular, lip callously frilled.

Cyclostoma rostratum, Pfeiffer, Pro. Zool. Soc. 1852, p. 66.

Cistula rostrata, Pfeiffer.

Hab. —?

Neatly painted throughout with curiously regular waved lines and distant dots, ranging alternately.

Species 71. (Mus. Cuming.)

CHONDROPOMA BILABIATUM. *Chond. testā umbilicatā, ovato-turritatā, livido-cinereā, anfractibus convexis, lineis spiralibus lamellisque scabrosis longitudinalibus decussatis; aperturā pyriformi-ovatā, obliquā, labro subfimbriato.*

THE TWO-LIPPED CHONDROPOMA. Shell umbilicated, ovately turreted, livid ash-colour, whorls convex, decussated with spiral lines and longitudinal seaceous lamellæ, aperture pyriformly ovate, oblique, lip slightly fimbriated.

Cyclostoma bilabiatum, D'Orbigny, Moll. Cub. p. 258. pl. 22. f. 3 to 5.

Cyclostoma D'Orbignianum, Petit.

Cyclostoma salebrosum, Morelet.

Choanopoma bilabiatum, Gray.

Ctenopoma bilabiatum, Pfeiffer.

Hab. Cuba.

Chiefly distinguished by its close-set longitudinal seaceous lamellæ.



73a.



72.



73b.



74.



75.



76.



77.



78.



79.



80.

Slightly Enlarged

CHONDROPOMA.

PLATE X.

Species 72. (Mus. Cuming.)

CHONDROPOMA ANTIGUENSE. *Chond. testā vix umbilicatā, oblongo-turrītā, albida, fuscescente pallidē punctatā, anfractibus convexis, longitudinaliter densē lamellatis, in suturas subdenticulatis; aperturā circulari, parvā, labro minutē fimbriato.*

THE ANTIGUA CHONDROPOMA. Shell scarcely umbilicated, oblong-turreted, whitish, faintly dotted with brown, whorls convex, longitudinally densely lamellated, slightly denticulated in the sutures; aperture circular, small, lip minutely fimbriated.

Cyclostoma Antiguense, Pfeiffer, Mon. Pneum. Suppl. 1. p. 131.

Cistula Antiguensis, Pfeiffer.

Hab. Antigua.

Beautifully sculptured longitudinally with thin close-set lamellæ.

Species 73. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA RUFILABRE. *Chond. testā minutissimē umbilicatā, oblongo-cylindraceā, solidiusculā, rubello-stramineā, roseo tinctā, fusco obscurē punctatā, anfractibus convexis, spiralerter tenuiratis, longitudinaliter crenuloso-lamellatis; aperturā circulari, parvā, labro sanguineo-rufo, obtusē fimbriato.*

THE RED-LIP CHONDROPOMA. Shell very minutely umbilicated, oblong-cylindrical, rather solid, reddish straw-colour, tinged with rose, obscurely dotted with brown, whorls convex, spirally finely ridged, longitudinally crenulously lamellated; aperture circular, small, lip blood-red, obtusely fimbriated.

Cyclostoma rufilabre, Beck, Pfeiffer, Mon. Pneum. Suppl. 1. p. 133.

Cyclostoma bilabre, Meuke.

Cistula rufilabris, Gray.

Tudora rufilabris, Murch.

Hab. Santa Cruz, West Indies.

A small solid cylindrical shell, with a callous blood-red lip.

Species 74. (Mus. Cuming.)

CHONDROPOMA INCULTUM. *Chond. testā vix umbilicatā, oblongo-turrītā, flavescente, castaneo interruptē lineatā,*

anfractibus convexis, obscurē reticulatis, in suturas valde denticulatis; aperturā circulari, parvā, labro expanso-reflexo.

THE NEGLECTED CHONDROPOMA. Shell scarcely umbilicated, oblong-turreted, yellowish, interruptedly lined with chestnut, whorls convex, obscurely reticulated, strongly denticulated in the sutures; aperture circular, small, lip expandedly reflected.

Cyclostoma incultum, Poey, Mém. 1. p. 98. pl. 8. f. 4, 5.

Chondropoma incultum, Pfeiffer.

Cistula inculta, Pfeiffer.

Hab. Cuba.

Sculpture is very obscure on the surface of this shell, but strongly developed in sutural denticles.

Species 75. (Mus. Cuming.)

CHONDROPOMA TAMSIANA. *Chond. testā conico-turrītā, rubello-stramineā, purpureo-fusco tinctā et interruptē fasciatā, anfractibus rotundatis, longitudinaliter densē tenuilamellatis; aperturā pyriformi-circulari, labro tennē expanso-reflexo.*

TAMS' CHONDROPOMA. Shell conically turreted, reddish straw-colour, stained and interruptedly banded with purple-brown, whorls rounded, longitudinally densely finely lamellated; aperture pyriformly circular, lip thinly expandedly reflected.

Cyclostoma Tamsiana, Pfeiffer, Zeitschr. für Malac. 1850, p. 77.

Hab. Venezuela; Dr. Tams.

The conically turreted form of this species is peculiar; the sculpture as in many of the more oblong Cuban and Jamaica species.

Species 76. (Mus. Cuming.)

CHONDROPOMA LUGUBRE. *Chond. testā vix umbilicatā, subangustē turrītā, flavescente, violaceo-cinereo latifasciatā, anfractibus plano-convexis, longitudinaliter conspicuē lamellatis, in suturas denticulatis; aperturā circulari, labro expanso-fimbriato, fimbriā purpureo-radiatā.*

THE MOURNFUL CHONDROPOMA. Shell scarcely umbilicated, rather narrowly turreted, yellowish, broadly

banded with violet-ash, whorls flatly convex, longitudinally conspicuously lamellated, denticulated in the sutures; aperture circular, lip expaudedly fimbriated, frill rayed with purple.

Cyclostoma lugubre, Pfeiffer, Pro. Zool. Soc. 1851, p. 249.

Cistula lugubris, Pfeiffer.

Hab. Jamaica.

The longitudinal lamellæ are rather more than usually conspicuous in this species, which is especially distinguished by its violet-ash-banded colouring.

Species 77. (Mus. Cuming.)

CHONDROPOMA PALLIDUM. *Chond. testā modicē umbilicatā, conico-turritā, sordidē stramineā, subhyalinā, pallidē fusco sparsim punctatā, anfractibus convexis, longitudinaliter minutissimē densē lamellatā; aperturā circulari, labro sublatè fimbriato.*

THE PALE CHONDROPOMA. Shell moderately umbilicated, conically turreted, dull straw-colour, subhyaline, sparsely dotted with pale-brown, whorls convex, longitudinally very minutely closely lamellated; aperture circular, lip rather broadly fringed.

Cyclostoma pallidum, Pfeiffer, Pro. Zool. Soc. 1851, p. 248.

Cistula pallida, Pfeiffer.

Hab. Havana.

Of the conically turreted type, with the longitudinal lamellar sculpture extremely finely developed.

Species 78. (Mus. Cuming.)

CHONDROPOMA SHEPARDIANUM. *Chond. testā angustē turritā, rubello-luteā, fusco punctato-strigatā, anfractibus rotundatis, longitudinaliter conspicuè lamellatis, in suturis denticulatis, anfractu ultimo subsoluto; aperturā circulari, labro subfimbriato.*

SHEPARD'S CHONDROPOMA. Shell narrowly turreted, reddish yellow, dot-streaked with brown, whorls rounded, longitudinally conspicuously lamellated, denticulated in the sutures, last whorl slightly uncoiled; aperture circular, lip a little fimbriated.

Cyclostoma Shepardiana, C. B. Adams, Contr. Conch. no. 6. p. 92.

Cistula Shepardiana, Pfeiffer.

Hab. Jamaica.

The longitudinal lamellar sculpture strongly predominates; the last whorl is slightly uncoiled.

Species 79. (Mus. Cuming.)

CHONDROPOMA MORDAX. *Chond. testā compressè umbilicatā, angustē turritā, flavescente, castaneo varie lentiginosā, anfractibus rotundatis, longitudinaliter densē scabroso-lamellatis, anfractu ultimo castaneo unifasciato; aperturā pyriformi-circulari, labro tenuè erpanso.*

THE CUTTING CHONDROPOMA. Shell compressly umbilicated, narrowly turreted, yellowish, variously freckled with chestnut, whorls rounded, longitudinally densely seaborously lamellated, last whorl encircled with a chestnut band; aperture pyriformly circular, lip thinly expanded.

Cyclostoma mordax, C. B. Adams, Contr. Conch. no. 1. p. 12.

Cistula mordax, Pfeiffer.

Hab. Jamaica.

The lamellæ of this species are more than usually seaborous, sharp and cutting.

Species 80. (Mus. Cuming.)

CHONDROPOMA RIISEI. *Chond. testā vix umbilicatā, conico-turritā, flavescente, pallidē fusco promiscue punctatā, anfractibus declivi-convexis, longitudinaliter densē filoso-lamellatis, lamellis in suturas denticulatis; aperturā obliquè ovalā, parvā, labro planoreflexo, radiale.*

RIISE'S CHONDROPOMA. Shell scarcely umbilicated, conically turreted, yellowish, promiscuously dotted with pale-brown, whorls slopingly convex, longitudinally densely thread-lamellated, lamellæ denticulated in the sutures; aperture obliquely ovate, small, lip flatly reflexed, rayed.

PFEIFFER, Mon. Phemu. p. 418.

Hab. Porto Rico.

A conically turreted species, in which the longitudinal lamellæ have more the character of fine threads.



81.



82.



83.



84.



85a.



85b.



86.



87.



88.



89.

Slightly Enlarged.

CHONDROPOMA.

PLATE XI.

Species 81. (Mus. Cuming.)

CHONDROPOMA CINCLIDODES. *Chond. testá imperforatá, cylindraceo-oblongá, albidá, fusco sparsim punctatá, anfractibus plano-convexis, scabroso-reticulatis, in suturis minutè denticulatis; aperturá obliquè ovata, parvá, labro rix reflexo.*

THE DELICATE CHONDROPOMA. Shell imperforate, cylindrically oblong, whitish, sparsely dotted with brown, whorls flatly convex, scarcely reticulated, minutely denticulated in the sutures; aperture obliquely ovate, small, lip scarcely reflected.

Cyclotoma cinclidodes, Pfeiffer, Pro. Zool. Soc. 1852, p. 142.

Cistula cinclidodes, Pfeiffer.

Hab. Haiti; Sallé.

A small cylindrical species, lamellarly sculptured after the prevailing type of the genus, but more reticulated with raised spiral lines.

Species 82. (Mus. Cuming.)

CHONDROPOMA TROCHLEARE. *Chond. testá minutè umbilicatá, angustè turritá, flavescente, fusco interruptrè fasciatá, anfractibus rotundatis, obliquè tenuilamellatis; aperturá circulari, parvá, labro fimbriato, fimbriae fusco radiatae.*

THE COILED CHONDROPOMA. Shell minutely umbilicated, narrowly turreted, yellowish, interruptedly banded with brown, whorls rounded, obliquely finely lamellated; aperture circular, small, lip fimbriated, frill rayed with brown.

Cyclotoma trochlearia, Pfeiffer, Pro. Zool. Soc. 1851, p. 249.

Cistula trochlearis, Pfeiffer.

Hab. Chiapa, Mexico.

The slender longitudinal lamellæ are not so close-set in this species as usual.

Species 83. (Mus. Cuming.)

CHONDROPOMA LIVIDUM. *Chond. testá compressè umbilicatá, conico-turritá, lividá, fusco obscurè punctatá, anfractibus convexis, striis densè filoso-reticulatis; aperturá circulari, labro subfimbriato.*

THE LIVID CHONDROPOMA. Shell compressly umbilicated, conically turreted, livid, obscurely dotted with brown, whorls convex, closely reticulated with thread-like striae; aperture circular, lip slightly fimbriated.

Hab. Cuba.

This species appears to have been mistaken for D'Orbigny's *C. bilabialum*.

Species 84. (Mus. Cuming.)

CHONDROPOMA GRATELOUPI. *Chond. testá oblonga, solidiuscula, flavescente, pallidè fusco punctatá et fasciatá, anfractibus convexis, longitudinaliter confertim obtuse tiratis, intersticiis minutè clathratis, suturis denticulatis; aperturá circulari, labro concavo-fimbriato.*

GRATELOUP'S CHONDROPOMA. Shell oblong, rather solid, yellowish, dotted and banded with pale brown, whorls convex, longitudinally closely obtusely ridged, interstices minutely latticed, sutures denticulated; aperture circular, lip concavely fimbriated.

Cyclotoma Grateloupi, Pfeiffer, Pro. Zool. Soc. 1851, p. 246.

Cistula Grateloupi, Pfeiffer.

Hab. Yucatan.

The longitudinal sculpture, commonly of a fine lamellar character, is here developed in obtuse ridges.

Species 85. (Fig. a, b, Mus. Cuming.)

CHONDROPOMA SAULIÆ. *Chond. testá subapertè umbilicatá, gracili-turritá, albida, subhyalina, punctis paucis pallidè fusci tincta, anfractibus rotundatis, longitudinaliter exilissimè lamellatis; aperturá rotundata, labro callosò-fimbriato, fimbriæ albæ, castaneo conspicuè radiatae.*

SAUL'S CHONDROPOMA. Shell rather openly umbilicated, slenderly turreted, whitish, subhyaline, stained with a few pale-brown dots, whorls rounded, longitudinally very delicately lamellated; aperture round, lip callously fimbriated, frill white, conspicuously rayed with chestnut.

Cyclotoma Sauliae, Sowerby, Thes. Conch. p. 145, pl. 28, f. 189.

Choanopoma Sauliae, Gray.*Cistula Sauliae*, Pfeiffer.*Hab.* Jamaica.

An extremely delicate species, with a pretty contrast of dark radiated painting on the frill.

Species 86. (Mus. Cuming.)

CHONDROPOMA LARGILLIERTI. *Chond. testā minutē umbilicatā, oblongo-cylindraceā, livido-albidā, fusco sparsim punctatā, anfractibus plano-convexis, longitudināliter confertim obtusè liratis, interstitiis minutē clathratis, suturis denticulatis; aperturā circulari, labro concavo-fimbriato.*

LARGILLIERT'S CHONDROPOMA. Shell minutely umbilicated, oblong-cylindrical, livid white, sparsely dotted with brown, whorls flatly convex, longitudinally closely obtusely ridged, interstices minutely latticed, sutures dentieulated, aperture circular, lip concavely fimbriated.

Cyclostoma Largillierti, Pfeiffer, Zeitschr. für Malac. 1846, p. 46.

Choanopoma Largillierti, Gray.*Cistula Largillierti*, Pfeiffer.*Hab.* Yucatan; Largilliert.

A dull, obtusely ridged shell, very closely allied to *C. Grateloupi*.

Species 87. (Mus. Cuming.)

CHONDROPOMA KUSTERI. *Chond. testā minutē umbilicatā, conico-turritā, rubello-stramineā, fusco obscurē punctatā, anfractibus rotundatis, striis scabrosis confertim decussatis; aperturā circulari, labro concavo-fimbriato, fimbriā subplicata.*

KUSTER'S CHONDROPOMA. Shell minutely umbilicated, conically turreted, reddish straw-colour, obscurely dotted with brown, whorls rounded, closely decussated with seaceous striae; aperture circular, lip concavely fimbriated, frill a little pnekered.

Cyclostoma Kusteri, Pfeiffer, Pro. Zool. Soc. 1851, p. 249.*Hab.* Honduras; Dyson.

Sculptured with a decussation of a very uniform character.

Species 88. (Mus. Cuming.)

CHONDROPOMA IGNEA. *Chond. testā compressē umbilicatā, cylindraceo-conicā, solidiusculā, albā, igneo-roseo tintatā, anfractibus convexis, minutissimē lamellato-decussatis, in suturis subtiliter denticulatis; aperturā pyriformi-circulatā, labro vix reflexo.*

THE FIERY CHONDROPOMA. Shell compressly umbilicated, cylindrically conical, rather solid, whitish, tinged with fiery rose, whorls convex, very minutely lamellarly decussated, finely denticulated in the sutures, aperture pyriformly circular, lip scarcely reflected.

Hab. —?

An extremely minutely lamellated species, remarkably tinged towards the apex with a bright, bronze-like, fiery rose-colour.

Species 89. (Mus. Cuming.)

CHONDROPOMA RADULA. *Chond. testā minutē umbilicatā, conico-turritā, flavescente, fusco sparsim punctato-fasciatā, anfractibus rotundatis, decussatim serrato-lamellatis; aperturā circulari, parvā, labro latē plano-fimbriato, fimbriā fusco-radiatā.*

THE RASP CHONDROPOMA. Shell minutely umbilicated, conically turreted, yellowish, sparsely dot-banded with brown, whorls rounded, decussately serrately lamellated; aperture circular, small, lip broadly flatly fimbriated, frill rayed with brown.

Cyclostoma radula, Pfeiffer, Pro. Zool. Soc. 1851, p. 246.
Cistula radula, Pfeiffer.

Hab. Havana.

In this species the decussating lamellæ are more serrated, like a rasp to the touch.

CHONDROPOMA.

	Plate	Species.		Plate	Species
abnatum, <i>Gundlach</i>	VI.	43	littoratum, <i>Pfeiffer</i>	IV.	26
Agassizii (<i>Cistula</i>), Pfeiffer . . .	IX.	64	lividum, <i>Reeve</i>	XI.	83
angustatum, <i>Pfeiffer</i>	IV.	29	Loweanum, <i>Pfeiffer</i>	VI.	44
Antiguense (<i>Cistula</i>), Pfeiffer . .	X.	72	lugubre (<i>Cistula</i>), Pfeiffer . . .	X.	76
argutum, <i>Pfeiffer</i>	VII.	48	lunulatum, <i>Morch</i>	V.	38
Auberianum, D'Orbigny	V.	38	magnificum, <i>Pfeiffer</i>	I.	1
basicarinatum, <i>Pfeiffer</i>	VIII.	58	Malogani, Gould	I.	5
bilabiatum (<i>Cyclos.</i>), D'Orbigny	IX.	71	marginalbum, <i>Gundlach</i>	VIII.	59
blandum, <i>Pfeiffer</i>	VIII.	62	marinum, <i>Weinland</i>	VIII.	57
Blameri, <i>Shuttleworth</i>	III.	19	mæstum, <i>Shuttleworth</i>	II.	14
Candeanum, D'Orbigny	VI.	40	mordax (<i>Cistula</i>), Pfeiffer . . .	X.	79
Charpentieri, Shuttleworth	II.	14	Newcombianum, <i>Adams</i>	III.	16
cincliodes (<i>Cistula</i>), Pfeiffer . .	XI.	81	nodulosum, Gray	VII.	49
claudicans, <i>Pfeiffer</i>	II.	7	obesum (<i>Trunc.</i>), Menke	III.	23
Cordovanum, <i>Pfeiffer</i>	IV.	24	Ottonis, <i>Pfeiffer</i>	III.	22
crenulatum (<i>Cycl.</i>), Féussac . .	VII.	49	pallidum (<i>Cistula</i>), Pfeiffer . . .	X.	77
Cumanense, <i>Pfeiffer</i>	V.	36	papyraceum, <i>C. B. Adams</i> . . .	VIII.	63
Delatreamum, D'Orbigny	V.	33	perlatum, <i>Gundlach</i>	IV.	30
dentatum (<i>Cycl.</i>), Say	V.	38	Petitianum, <i>Pfeiffer</i>	V.	34
diaphanum, <i>Pfeiffer</i>	V.	32	petricosum, Morelet	III.	22
dissolutum, <i>Pfeiffer</i>	IV.	27	Pfeifferianum, <i>Poey</i>	V.	35
egregium, <i>Gundlach</i>	I.	3	pictum, <i>Pfeiffer</i>	I.	5
eusareum, <i>Pfeiffer</i>	VII.	54	plicatulum, <i>Pfeiffer</i>	III.	20
glabratum, <i>Weinland</i>	II.	12	Poeyanum (<i>Cycl.</i>), D'Orbigny .	III.	15
Grateloupi (<i>Cistula</i>), Pfeiffer . .	XI.	84	Poeyanum, <i>Pfeiffer</i>	II.	14
Grumeri (<i>Cistula</i>), Pfeiffer . . .	IX.	68	puncticulatum, <i>Pfeiffer</i>	VI.	41
Gnadaloupense, <i>Pfeiffer</i>	VII.	49	quaternatum (<i>Cycl.</i>), Lamarek .	VIII.	60
Gutiérrezi, Poey	I.	1	radiosum, Morelet	X.	69
Habichii, <i>Weinland</i>	I.	6	radula (<i>Cistula</i>), Pfeiffer	XI.	89
harpa, <i>Pfeiffer</i>	VII.	55	revinctum (<i>Cycl.</i>), Poey	VI.	46
Hellenicum (<i>Cycl.</i>), Gray	II.	14	revocatum, <i>Gundlach</i>	VII.	53
hemiotum, <i>Pfeiffer</i>	III.	18	Rüsei, <i>Pfeiffer</i>	X.	80
Hjalmersoni, <i>Pfeiffer</i>	VIII.	61	rostratum (<i>Cistula</i>), Pfeiffer . .	IX.	70
igneus, <i>Reeve</i>	XI.	88	rubicundum (<i>Cycl.</i>), Morelet . .	VI.	47
incultum (<i>Cyclos.</i>), Poey	X.	74	rude, <i>Weinland</i>	IV.	28
integrum, <i>Pfeiffer</i>	IV.	25	rufilabre (<i>Cyclos.</i>), Beck	X.	73
irradians, <i>Shuttleworth</i>	II.	8	rufopictum, <i>Gundlach</i>	V.	39
jucundum, <i>Pfeiffer</i>	III.	17	sagittiferum, <i>Pfeiffer</i>	IX.	67
Küsteri, <i>Pfeiffer</i>	XI.	87	Sagra, D'Orbigny	I.	5
Largillierti (<i>Cistula</i>) Pfeiffer . .	XI.	86	Salleanum, <i>Pfeiffer</i>	II.	10
latilabre (<i>Cycl.</i>), D'Orbigny . .	II.	11	Santaeruzense, <i>Pfeiffer</i>	VII.	50
latum, <i>Gundlach</i>	VIII.	56	Sauliae (<i>Cyclos.</i>), Sowerby . . .	XI.	85
lineolatum, Anton	V.	38	semilabre, <i>Pfeiffer</i>	I.	2

CHONDROPOMA.—INDEX.

Plate.	Species.	Plate.	Species.	
Shepardianum (<i>Cycl.</i>), C. B. Adams X.	78	Tortolense, <i>Pfeiffer</i>	IV.	31
Shuttleworthi, <i>Pfeiffer</i>	I.	trochleare (<i>Cistula</i>), <i>Pfeiffer</i>	XI.	82
simplex, <i>Pfeiffer</i>	VI.	<i>Troscheli</i> , <i>Pfeiffer</i>	III.	16
solare, <i>Pfeiffer</i>	IX.	<i>truncatum</i> , <i>Pfeiffer</i>	VI.	40
Swifti, <i>Shuttleworth</i>	III.	<i>turritum</i> , <i>Pfeiffer</i>	VII.	52
<i>Swiftianum</i> , Newcomb.	III.	Venezuelense, <i>Pfeiffer</i>	VI.	45
Tamsiana (<i>Cyclos.</i>), <i>Pfeiffer</i> . . .	X.	<i>verecundum</i> , Poey	I.	4
tenebrosum (<i>Cycl.</i>), Morelet . . .	II.	<i>violaceum</i> , <i>Pfeiffer</i>	II.	13
ternatum, <i>Gould</i>	IX.	Weinlandi, <i>Pfeiffer</i>	I.	2
textum, <i>Gundlach</i>	VII.			

M O N O G R A P H
OF THE GENUS
A D A M S I E L L A.

He's walk'd the way of nature ;
And, to our purposes, he lives no more."—*Shakespeare*.



2



3



4



5a



5b



6



7a.



7b



8

(Slightly Enlarged)

A D A M S I E L L A .

PLATE 1.

Genus ADAMSIELLA, Pfeiffer.

Testa oblongo-turrata vel pupaformis, plerumque minutè reticulata, ad suturas raro crenulata, anfractu ultimo sepe plus minus soluto; apertura parviuscula, labro plerumque calloso-fimbriato. Operculum circulare, tenue, subcartilagineum, anfractibus paucis, sensim crescentibus, margine sublibero, nucleo subcentrali.

Shell oblong-turreted or pupa-shaped, generally minutely reticulated, rarely crenulated at the sutures, last whorl often more or less uncoiled; aperture rather small, lip generally callously fimbriated. Operculum circular, thin, somewhat cartilaginous, whorls few, increasing gradually, with the edge somewhat free, nucleus nearly central.

In this genus of *Cyclostomacea*, named in honour of the late Professor C. B. Adams, the operculum is circular, formed of a few gradually increasing whorls, slightly free at the edge, with the nucleus nearly central. The species, of which there are scarcely more than a dozen well defined, are composed of rather narrow whorls, the last being generally more or less uncoiled. The sculpture is minute, delicately thread-like, including in some instances a few crenulations in the sutures. The aperture is small and subtubular, mostly encircled with a flatly expanded callous frill. All the species, excepting one from Cuba, are from Jamaica, one ranging into Demerara. A shell, unknown to me, it should however be added, was described in 1861 (Pro. Zool. Soc. p. 232) by the Rev. H. B. Tristram, under the name *A. Osberti*, collected by Mr. Salvin in Guatemala.

Species 1. (Mus. Cuming.)

ADAMSIELLA IGNILABRIS. Adams. *testa angustè profundè umbilicatâ, turritatâ, fuscâ, labro vividè miniato; anfractibus rotundatis, striis elevatis undique dense annulatis; labro dupliciti, calloso-reflexo.*

THE FIRE-LIP ADAMSIELLA. Shell narrowly deeply umbilicated, turreted, fuscous, lip bright scarlet-red; whorls rounded, closely ringed throughout with raised striae; lip duplex, callously reflexed.

Cyclostoma ignilabre, Adams, Contr. to Conch. No. 1. p. 9.

Adamsiella ignilabris, Pfeiffer.

Cyclostoma chlorostoma, Sowerby.

Choanopoma chlorostoma, Gray.

Adamsiella chlorostoma, Pfeiffer.

Hab. Jamaica. Demerara.

A closely ring-striated species, in which the fiery-red colouring of the lip contrasts in a striking manner with the dull brown of the shell.

Fig. 2. (Mus. Cuming.)

This shell, which is Mr. Sowerby's *Cyclostoma articulatum*, proves to be a variety of Professor C. Adams's *Cyclostoma variable* (*Adamsiella variabilis*, Pfeiffer).

Species 3. (Mus. Cuming.)

ADAMSIELLA GRAYANA. Adams. *testa subangustè profundiè umbilicatâ, ovato-pupaformi, violaceo-fuscâ, sericeâ, anfractibus rotundatis, striis elevatis undique confertim decussatis, labro dupliciti, latè calloso-expanso, fulvescente.*

GRAY'S ADAMSIELLA. Shell rather narrowly deeply umbilicated, ovately pupa-shaped, violet-brown, silky, whorls rounded, closely decussated throughout with raised striae, lip duplex, broadly callously expanded, yellowish.

Cyclostoma Grayanum, Pfeiffer, Zeitsch. für Malac. 1846.

Cyclostoma obscurum, Gray (not of Draparnaud).

Choanopoma Grayanum, Pfeiffer.

Adamsiella Grayanum, Pfeiffer.

Cyclostoma concentricum, C. B. Adams.

Cyclostoma serriferum, C. B. Adams.

Hab. Jamaica.

Of a rather short pupa-like form, with the lip broadly callously expanded.

Species 4. (Mus. Cuming.)

ADAMSIELLA MONSTROSA. Adams. *testa subaperitè umbilicatâ, oblongo-turritatâ, fulvescente, semipellucidâ, rufo-fusco exiùtâ maculatâ et marmoratâ, anfractibus angustis, rotundatis, decussatim striatis, ultimo soluto; apertura minutâ, labro latissimè calloso-reflexo.*

THE MONSTROUS ADAMSIELLA. Shell rather openly umbilicated, oblong-turreted, yellowish, semitransparent, lightly spotted and marbled with red-brown, whorls

narrow, rounded, decussately striated, last whorl uncoiled; aperture very small, lip very broadly callously reflected.

Cyclostoma monstrosum, Adams, Contr. to Conch. no. 1. p. 5.

Adamsiella monstrosa, Pfeiffer.

Hab. Jamaica.

A semitransparent delicately-painted shell of narrow subtubular whorls, of which the last is partially unrolled, with the lip unusually broadly flatly reflected.

Species 5. (Fig. a, b, Mus. Cuming.)

ADAMSIELLA VARIABILIS. *Adams. testā profundē umbilicatā, ovato-turritā, fulvescente, fusco varie fasciatā, juxta suturas articulatā, anfractibus subangustis, rotundatis, striis elevatis minutē decussatis, labro duplice, modicē expanso.*

THE VARIABLE ADAMSIELLA. Shell deeply umbilicated, ovately turbinated, fulvous, variously banded with brown, articulated next the sutures, whorls rather narrow, rounded, minutely decussated with raised striae, lip duplex, moderately expanded.

Cyclostoma articulatum, Gray (not of Draparnaud).

Cyclostoma variabile, Adams.

Cyclostoma mutatum, Adams.

Choanopoma articulatum, Pfeiffer.

Adamsiella variabilis, Pfeiffer.

Hab. Jamaica.

This species, as may be seen by our figures, varies in form, and the decussating striae are stronger in some specimens than in others. The painting consists almost invariably of bands, with an articulation of spots below the sutures.

Species 6. (Fig. a, b, Mus. Cuming.)

ADAMSIELLA PEARMANEANA. *Adams. testā opertē profundē umbilicatā, oblongo-turritā, pellucido-albā, rufo-fusco minutē lentiginosā et radiatā, anfractibus sub-augustis, rotundatis, striis pulchre decussatis, longitudinalibus lamelliformibus, anfractu ultimo soluto; aperturā parvā, labro eleganter fimbriato.*

PEARMAN'S ADAMSIELLA. Shell openly deeply umbilicated, oblong-turreted, transparent-white, minutely freckled and rayed with red-brown, whorls rather narrow, rounded, beautifully decussated with striae, of which the longitudinal are raised into lamellæ, last whorl uncoiled; aperture small, lip elegantly fimbriated.

Cyclostoma Pearmaneum, Chitty, Contr. to Conch. no. 1.

p. 2.

Cyclostoma longicolle, Pfeiffer.

Adamsiella Pearmaneana, Pfeiffer.

Hab. Jamaica.

In this most elegant species the longitudinal striae are raised throughout into fine thread-like lamellæ, while the frill encircling the aperture is removed some little distance back from the lip.

Species 7. (Fig. a, b, Mus. Cuming.)

ADAMSIELLA INTERMEDIA. *Adams. testā subapertē profundē umbilicatā, oblongo-turritā, fulvescente-albā, semipellucidā, rufo-fusco interruptē fasciatā et articulatā, anfractibus rotundatis, striis elevatis reticulato-decussatis; aperturā parviusculā, labro calloso-fimbriato.*

THE INTERMEDIATE ADAMSIELLA. Shell rather openly deeply umbilicated, oblong turreted, fulvous white, semitransparent, interruptedly banded and articulated with red-brown, whorls rounded, reticulately decussated with raised striae; aperture rather small, lip callously fimbriated.

Cyclostoma intermedium, Adams, Contr. to Conch. no. 1.

p. 5.

Adamsiella intermedia, Pfeiffer.

Hab. Jamaica.

Characterized among other features by the interrupted slightly-waved style of the painting.

Species 8. (Mus. Cuming.)

ADAMSIELLA BINNEYANA. *Adams. testā profundē umbilicatā, ovato-turritā, subcylindraceā, fulvescente-albā, medio fasciū latā castaneā cingulatā, anfractibus convexis, sub lente minutē decussatis, labro calloso-reflexo, albo.*

BINNEY'S ADAMSIELLA. Shell deeply umbilicated, ovately turreted, subcylindrical, yellowish-white, encircled round the middle with a broad chestnut band, whorls convex, minutely decussated beneath the lens, lip callously reflected, white.

Cyclostoma Binneyanum, Adams, Pfeiffer, Zeitschr. für Malac. 1846, p. 47.

Cyclostoma pulchrius, Adams.

Choanopoma Binneyanum, Gray.

Adamsiella pulchrior, Pfeiffer.

Hab. Jamaica.

This apparently smooth shell is especially distinguished by its single broad chestnut band.



(Slightly Enlarged.)

A D A M S I E L L A.

PLATE II.

Species 9. (Mus. Cuming.)

ADAMSIELLA CHORDATA. Adams. testū minutè umbilicatā, subangustē turritū, fulvā, rividē castano linearifasciatā, anfractibus convexis, striis tenuibus decussatis, striis longitudinalibus subtiliter filosis, ad suturas crenulatis, labro tenuè reflexo.

THE STRINGED ADAMSIELLA. Shell minutely umbilicated, rather narrowly turreted, fulvous, linearly banded with bright-chestnut, whorls convex, decussated with fine striae, the longitudinal striae raised in fine threads, crenulated at the sutures, lip thinly reflected.

GUNDLACH, MS. in Mus. Cuming.

Hab. Cuba.

The delicate longitudinal thread-like striae are paler in colour than the ground of the shell, which is encircled throughout with from three to four linear bands of bright-chestnut.

Species 10. (Mus. Cuming.)

The shell here represented is the type of Mr. Sowerby's *Cyclostoma chlorostoma* (*Adamsiella chlorostoma*, Pfeiffer). It appears to me to be a yellow-lipped variety of *A. ignobilis*.

Species 11. (Mus. Cuming.)

ADAMSIELLA PINGUIS. Adams. testā subampliter umbilicata, acuminate-turritū, cinnamoneo-fuscā, subminante, anfractibus angustis, rotundatis, liris spiralibus obtusis, striisque longitudinalibus minutis, sculptis.

THE GREASY ADAMSIELLA. Shell rather largely umbilicated, acuminate turreted, cinnamon-brown, rather shining, whorls narrow, rounded, sculptured with obtuse spiral ridges and minute longitudinal striae.

Cyclostoma pingue, Pfeiffer, Pro. Zool. Soc. 1851, p. 248.

Adamsiella pinguis, Pfeiffer.

Hab. —?

The oily shining aspect of this shell, after which it is named, can hardly be relied on as a specific character.

Species 12. (Mus. Cuming.)

ADAMSIELLA MORIBUNDA. Adams. testū apertè profundè umbilicata, oblonga, solidiuscula, lutescente, anfrac-

tibus rotundatis, minutissimè decussatim striatis, striis longitudinalibus validioribus, ad suturas crenatis, labro angustè plano-fimbriato.

THE DEATH-LIKE ADAMSIELLA. Shell openly deeply umbilicated, oblong, rather solid, yellowish, whorls rounded, very minutely decussately striated, longitudinal striae the stronger, crenated at the sutures, lip narrowly flatly fimbriated.

Cyclostoma moribundum, Adams, Contr. to Conch. no. 1. p. 5.

Adamsiella moribunda, Pfeiffer.

Hab. Jamaica.

A pale-yellowish shell, minutely decussated throughout, the longitudinal striae, which are the stronger, having the appearance of delicate silken cords.

Species 13. (Fig. a, b, Mus. Cuming.)

ADAMSIELLA MIRABILIS. Adams. testā apertè profundè umbilicata, oblongo-conicā, tenuiculā, albida, rufo-fusco interruptè lineatim trifasciatā, infra suturas conspicuè maculata, anfractibus rotundatis, striis elevatis granoso-decussatis, labro plano-fimbriato, subcalloso.

THE WONDERFUL ADAMSIELLA. Shell openly deeply umbilicated, oblong-conical, rather thin, whitish, interruptedly linearly three-banded with red-brown, conspicuously spotted beneath the sutures, whorls rounded, granosely decussated with raised striae, lip flatly fimbriated, rather callous.

Turbo mirabilis, Wood, Ind. Test. Supp. pl. 6. f. 22.

Cyclostoma mirabile, Gray.

Cyclostoma decussatum, Adams.

Adamsiella mirabilis, Pfeiffer.

Hab. Jamaica.

In this species the decussation has almost the granular roughness of a thimble.

Species 14. (Mus. Cuming.)

ADAMSIELLA MIRANDA. Adams. testā apertè profundè umbilicata, oblongo-conicā, albida, rufo-fusco interruptè trifasciatā, infra suturas conspicuè maculata, anfractibus rotundatis, minutè decussatim striatis, striis longitudinalibus validioribus, anfractu ultimo paululum soluto, labro conspicuè plano-fimbriato.

THE ADMIRED ADAMSIELLA. Shell openly deeply umbilicated, oblong-eombieal, whitish, interruptedly three-banded with red-brown, conspicuously spotted beneath the sutures, whorls rounded, minutely decussately striated, longitudinal striae the stronger, last whorl a little uneoiled, lip conspicuously flatly frilled.
Cyclostoma mirandum, Couth. to Conch. no. 1. p. 4.

Adamsiella miranda, Pfeiffer.

Hab. Jamaica.

Resembling the preceding species in colouring, but with the last whorl more loosely coiled, while the decussation is different. Here the spiral striae are almost obsolete, and the longitudinal striae have the appearance of fine silken cords.

Fig. 15 and 16. (Mus. Cuming.)

The shells here figured represent respectively Professor C. B. Adams' *Cyclostoma (Adamsiella) serriferum* and *concentricum*, both of which alleged species Dr. Pfeiffer refers to *A. Grayana*.

Species 17. (Mus. Cuming.)

ADAMSIELLA XANTHOSTOMA. *Adams. testā apertē profundē umbilicatā, subacuminato-turrilā, cinereo-fuscescente, anfractibus rotundatis, spiraliter obtusē tiratis, longitudinaliter densē minutissimē striatis, labro callosē plano-fimbriato, aurantio-tuteo.*

THE YELLOW-MOUTH ADAMSIELLA. Shell openly deeply umbilicated, rather acuminate turreted, light ash-brown, whorls rounded, spirally obtusely ridged, longitudinally densely very minutely striated, lip callously flatly fimbriated, orange-yellow.

Cyclostoma xanthostoma, Sowerby, Thesaurus Conch. p. 144. pl. 29. f. 175.

Cyclostoma tenuistriatum, Adams.

Choanopoma xanthostoma, Gray.

Adamsiella xanthostoma, Pfeiffer.

Hab. Jamaica.

Of a more acuminate turreted form than usual in the genus; this shell is further distinguished by the prominence of its spiral ridges and orange-yellow lip.

MONOGRAPH
OF THE GENUS
A N A S T O M A.

"In her most green recesses, to be found
By all who seek her truly."—*Barry Cornwall.*



1a



2a



2b




3a



4a



4b



3b



5a



5b



7b.

ANASTOMA.

PLATE I.

Genus ANASTOMA, Fischer.

Testu depresso-globosa, lenticularis, aufractu ultimo abrupte transvolvuto, aperturā semilunari, supra spiram reversā, utrinque intro dentatā, dentibus plicato-laminatis, exterioribus quatuor ad quinque, columellaribus duobus, labro expanso-reflexo, interdum supernè perforato.

Shell depressly globose, lens-shaped, last whorl abruptly turned across, aperture semilunar, turned upwards towards the spire, toothed within on both sides, teeth plicately laminate, outer teeth four to five, columellar teeth two, lip expandedly reflected, sometimes perforated at the upper part.

It is a curious feature in the natural history of Brazil that that country should give birth to a form of snail differing from all other snails in a habit of reversing the position of its shell on arriving at maturity. Up to the formation of the first half of the last whorl the growth of the shell proceeds as usual, but from this point the whorl is suddenly turned across in the direction of the apex. By this new effort of the animal the shell is overturned, and the aperture is brought on a plane with the spire. The snail thenceforward crawls with its spire downwards, and base upwards. Two very distinct species of *Anastoma* were known to Féussac and Lamarck, *A. ringens* and *globulosum*, the first having been known to Linnaeus. Three more species have been added by modern writers, but they are of much inferior value. *A. Deshayesianum* and *Terreauxianum* differ little from *A. ringens*; and *A. carinatum* is probably only a variety of *A. globulosum*.

Species 1. (Fig. a, b, Mus. Cuming.)

ANASTOMA CARINATUM. *Anast. testū orbiculari, depresso-sinuata, ad peripheriam plus minus angulato-varinata, livido-olivaceā, supra peripheriam ustulato-purpureo fasciatā, infra purpureo-rufō maculatā, aufractibus supra concentricē arcuato-striatis, infra reticulato-corrugatis, labro supernè perforato.*

THE KEELED ANASTOMA. Shell orbicular, rather depressed, more or less angularly keeled at the periphery, livid olive, banded above the periphery with burnt-purple, beneath spotted with purple-red, whorls concentrically arcuately striated above, reticulately wrinkled below, lip perforated above.

PFEIFFER, Pro. Zool. Soc. 1853. p. 57.

Hab. Brazil.

This and *A. globulosum*, which are probably varieties of one and the same species, are curiously characterized by the lip being perforated at its upper junction with the body whorl.

Species 2. (Fig. a, b, Mus. Cuming.)

ANASTOMA GLOBULOSUM. *Anast. testū depresso-globosā, ad peripheriam nunc rotundatā, nunc subcarinatā, livido-olivaceā, supra peripheriam ustulato-purpureo fasciatā, infra purpureo-rufō lentiginosā, aufractibus supra concentricē arcuato-striatis, infra reticulato-corrugatis.*

THE GLOBULAR ANASTOMA. Shell depressly globose, now rounded, now slightly keeled at the periphery, livid olive, banded above the periphery with burnt-purple, freckled beneath with purple-red, whorls concentrically arcuately striated above, reticulately wrinkled below.

LAMARCK, Anim. sans vert. vol. 8. p. 153.

Helix ringula, Féussac.

Hab. Brazil.

The examples set apart for this species are uniformly smaller and rounder than those of the preceding.

Species 3. (Fig. a, b, Mus. Cuming.)

ANASTOMA RINGENS. *Anast. testū depresso-orbiculari, subrudē tumidā, ad peripherium sēpe subcarinatō, albida, rufō-castaneo inquinatā et maculatā, juxta suturas fasciatā, supra lēvigatā, infra obscurē corrugato-striatā.*

THE GRINNING ANASTOMA. Shell depressly orbicular, rather rudely swollen, often slightly keeled at the periphery, whitish, daubed and spotted with reddish-chestnut, banded next the sutures, smooth above, obscurely wrinkle-striated beneath.

Helix ringens, Linnaeus, Syst. Nat. p. 1243.

Tomogerus ringens, De Montford.

Anostoma depressa, Lamarck.

Anostoma ringens, Deshayes.

Hab. Brazil.

The largest of the genus, rather rudely tumidly convoluted, the aperture being white and conspicuously expanded.

Species 4. (Fig. *a, b*, Mus. Cuming.)

ANASTOMA VERREAUXIANUM. *Anast. testā depressē orbiculari, subruditē tumidō, ad peripheriam sēpe subcarinatā, fulvescente, livido-purpurascente tinctā, rufo-castaneo sparsim maculatā, supra arcuato-striatā, infra corrugato-striatā, labro livido-purpurascente.*

VERREAUX'S ANASTOMA. Shell depresso orbicular, somewhat rudely swollen, often keeled at the periphery, fulvous, tinged with livid purple, sparingly blotched with reddish-chestnut, areately striated above, wrinkle-striated beneath, lip livid purple.

HUPÉ, Journ. Conch. vol. 5. p. 352.

Hab. Brazil.

Distinguished by its livid purple colouring, especially about the aperture.

Species 5. (Fig. *a, b*, Mus. Cuming.)

ANASTOMA DESHAYESIANUM. *Anast. testā depressē orbiculari, tumidiusculā, ad peripheriam parum canaliculatā, albida, rufo-castaneo maculatā et fasciatā, supra lăvigatā, infra corrugato-striatā, aperturā subcontractā.*


DESHAYES' ANASTOMA. Shell depresso orbicular, rather swollen, but little keeled at the periphery, whitish, blotched and banded with reddish-chestnut, smooth above, wrinkle-striated beneath, aperture rather contracted.

FISCHER, Journ. Conch. vol. 5. p. 350. pl. 12. f. 1, 2.
Hab. Brazil.

This has very much the appearance of being a dwarf variety of *A. ringens*.

MONOGRAPH
OF THE GENUS
T R I D A C N A.

"It was a garden still beyond all price,
Even yet it was a place of Paradise
And here were coral bowers
And grotos of madrepores."—*Southern*



TRIDACNA.

PLATE I.

Genus TRIDACNA, Bruguière.

Testa ovata vel oblongo-ovala, æquivalvis, inæquilateralis, antice ampliter hians, radiatim pauci-costata, costis plerumque squamatis. Cardo dentibus duobus in utrâque valvâ compressis, inæqualibus, posticis, insertis. Ligamentum submarginale, externum.

Shell ovate or oblong-ovate, equivalve, inequilateral, largely gaping anteriorly, radiately few-ribbed, ribs generally sealed. Hinge with two posterior compressed interlocking teeth in each valve. Ligament submarginal, external.

The *Tridacna*, or Furbelowed Clams, one of which, *T. gigas*, is the largest of all mollusks, live in beds of some extent in lagunes among coral reefs, among the islands of the Eastern and Pacific seas. The shell is generally white, sometimes tinged with red and saffron or lemon-yellow, but the animal is brilliantly coloured. M. Quoy describes the beautiful iridescent glare of blue, violet, and yellow, variegated with fantastic markings, that is presented by these submarine parterres as seen through the clear blue water; and Mr. Cuming speaks with enthusiasm of passing over a mass of them nearly a mile in extent, which resembled nothing so much as a beautiful bed of tulips.

With the exception of *T. gigas*, from Navigators' Islands, all the species of which I have the habitat, are assigned to the Molueas or Philippine Islands. The shells are mostly furbelowed with large high-vaulted scales, and all are white, occasionally tinged with red and saffron or lemon-yellow. The lumle of the shell is widely open for

the passage of a foot capable of spinning a powerful tendinous byssus.

Species I. (Pl. I. Fig. 1 *a*; Pl. II. Fig. 1 *b, c*, Mus. Cuming.)

TRIDACNA GIGAS. *Trid. testâ maximâ, antice subobliquè productâ, quadricostatâ, costis densè squamatis, squamis numerosis, creberrimis, ætate brevibus, plus minus obsoletis, intersticiis subobscure radialiter liratis; albâ, intus interdum roseâ.*

THE GIANT TRIDACNA. Shell very large, rather obliquely produced anteriorly, four-ribbed, ribs densely scaled, scales numerous, close-set, short and more or less obsolete with age, interstices rather obscurely radiately ridged; white, sometimes tinged within with rose.


Chama gigas, Linnaeus, Syst. Nat. Gmel. edit. p. 3299.

Tridacna gigas and *multica*, Lamarck.


Hab. Indian and Pacific seas.

This colossal species, which attains the enormous size of from six to seven hundred pounds weight, may be distinguished in all its stages of growth by a tendency to radiate anteriorly in an oblique direction, and by the closely-sealed character of the ribs. The ribs are broad, only four in number, and as the shell increases in age the scales become shorter, closer-set, and more obscure. In the shell represented in the opposite plate the interior is tinged with deep rose-colour.


June, 1862.




1b



1c



2a



2b

TRIDACNA.

PLATE II.

Species 1. (Fig. *b*, *c*, Mus. Cuming.)

The shells here figured represent the great *T. gigas* in an early stage of growth. At first, Fig. 1 *b*, the scales are lofty, very regular, and neatly-defined. Presently they become eroded and worn down, whilst the successive additions of scales, as shown at Fig. 1 *c*, are shorter and closer set. Lastly, as in the specimen represented in Plate I., the development of the scales is reduced to a mere crowded incrustation of rude layers of growth.

Species 2. (Fig. *a*, *b*, Mus. Cuming.)

TRIDACNA ELONGATA. *Trid. testā ovato-oblongā, anticē elongato-productā, quinque- ad sex-costatā, costis convexis, regulariter et subconfertim squamatis, inter-*

stiliis inter costas biliratis, inter squamas lævibus; albā vel citrinā.


THE ELONGATED TRIDACNA. Shell ovately oblong, anteriorly elongately produced, five- to six-ribbed, ribs convex, regularly and rather closely sealed, interstices between the ribs two-ridged, between the scales smooth; white or lemon-yellow.

LAMARCK, Anim. sans vert. vol. vii. p. 9.

Hab. Philippine Islands; Cuming.

The elongately-produced growth of this species gives a peculiarly gibbous form to the shell; and it retains this characteristic contour in all stages. In the British Museum there is a specimen of quite colossal dimensions, of precisely the same form. The scales are very regularly set throughout the ribs in neatly-defined furrows.

June, 1862.



TRIDACNA.

PLATE III.

Species 3. (Fig. *a*, Mus. Brit., and Pl. IV. Fig. *b, c, d*,
Mus. Cuming.)

TRIDACNA SQUAMOSA. *Trid. testá ovatá, quadri- ad quinque-costatá, costis elevato-convexus, distantibus, fimbriato-squamatis, squamis peramplis, distantibus, erectis, fornicatis, costarum interstitiis decussatim multistriatis; albá vel croceá, interdum rubente tinctá.*

THE SCALED TRIDACNA. Shell ovate, four- to five-ribbed, ribs elevately convex, distant, fimbriately scaled, scales very large, distant, erect, vaulted, interstices


between the ribs decussately many-striated; white or saffron-yellow, sometimes tinged with red.

LAMARCK, Anim. sans vert. vol. vii. p. 10.


Hab. Moluccas.

This well-known species is characterized by a much more symmetrical growth than *T. gigas*. It has no tendency to radiate obliquely, and the scales are developed throughout with amplitude and precision. The ribs are raised and tumidly convex, whilst the interstices between them are spacious and decussated throughout with impressed striae.


June, 1862.



3b



3c



3d

TRIDACNA.


PLATE IV.

Species 3. (Fig. *b*, *c*, *d*, Mus. Cuming.)

TRIDACNA SQUAMOSA. In this Plate are examples of three coloured varieties of the species described at Plate III. The specimen, Fig. 3 *b*, is white, tinged concentrically, not in rays, with pale orange-red. At Fig. 3 *c* the shell is of a dark saffron-yellow, with a mixture of red

that gives it an orange tone. At Fig. 3 *d* the shell is white, with a tinge of lemon-yellow without red. In all the scales or furbelows are largely developed, and in the shells of the upper and lower figures they are white; the colour described in those specimens being confined to the ground of the shell.

June, 1862.



TRIDACNA.

PLATE V.

Species 4. (Fig. *a*, *b*, *c*, Mus. Cuming.)

TRIDACNA RUDIS. *Trid. testā gibboso-ovatā, rudi, quadri-
ad quinque-costatā, costis latiusculis, parum elevatis,
radiatum liratis, squamatis, squamis numerosis, valde
irregularibus, hic illie continuis, costarum interstitiis
angustis, radiatum liratis; albā.*

THE RUDE TRIDACNA. Shell gibbously ovate, rude, four-
to five-ribbed, ribs rather broad, but little raised,
radiately ridged, scaled, scales numerous, very ir-
regular, here and there continuous, interstices between
the ribs narrow, radiately ridged; white.

Hab. Philippine Islands; Cuming.

The shells figured in this plate represent a very charac-
teristic form of *Tridacna*, certainly distinct from *T. squa-*

mosa, and apparently distinct from *T. gigas*. They have
not the anterior obliquity which is peculiar to *T. gigas*, as
shown both in the large specimens, Plate I., and the small
specimens given in Plate II. Another feature in *T. rудis* is
that the ribs are conspicuously radiately ridged through-
out between the scales. The scales are very irregular,
running in some places continuously over the intercostal
furrows, and the whole shell is singularly uncouth and
unsymmetrical.

In the species represented at Plate II., Fig. 1 *b*, *c*, which
I take to be the young of *T. gigas*, there are no radiating
ridges between the scales; and the shell is characterized
by the same obliquity of the anterior portion seen in the
larger shell figured in Plate I.

June, 1862.



TRIDACNA.

PLATE VI.

Species 5. (Fig. 5 a; and Pl. VII. Fig. 5 b, c.)

TRIDACNA COMPRESSA. *Trid. testā ovatā, compressā, sex-costatā, costis latiusculis, radiatim liratis, densè squamatis, squamis parvulis, erectis, costarum interstitiis radiatim liratis, liris tribus centralibus prominentioribus; albā, interdum pallidē citrino tinctā.*

THE COMPRESSED TRIDACNA. Shell ovate, compressed, six-ribbed, ribs rather broad, radiately ridged, densely scaled, scales rather small, erect, interstices between the ribs ridged, the three central ridges being the more prominent; white, sometimes tinged with pale lemon-colour.

Hab. —?

Either this is a species or it is a non-elongated form of *T. elongata*. In a genus like *Tridacna*, where the species are so numerous and the varieties so variable, some licence in naming is really useful to distinguish them. In the present instance, the shell is broader from the umboes to the margin and more compressed than that of the typical *T. elongata*, and the scales are smaller in proportion to the width of the ribs, more distant, and more erect. All this, it may nevertheless be argued, is a correlation of growth arising out of the compression of the valves. At Plate

VII., Fig. 5 b and 5 c, are represented younger specimens of the same form.

Species 6. (Mus. Cuming.)


TRIDACNA SERRIFERA. *Trid. testā latē ovatā, compressissimā, tenuiculā, tri- quadri-costatā, costis interstitiisque amplis, nudis, radiatim sulcato-striatis, striis in interstitiis fortioribus, marginibus anticis squamoso-serratis; albā vel pallidē citrinā.*

THE SERRATED TRIDACNA. Shell broadly ovate, rather compressed, somewhat thin, three- to four-ribbed, ribs and interstices large, scaleless, radiately groove-striated, striae stronger in the interstices, anterior margins scale-serrated; white or pale lemon-colour.


LAMARCK, Anim. sans vert. vol. vii. p. 11.

Hab. Moluccas.


The character of this species is to be without scales, except on the anterior margins, where they are short and often nearly obsolete. But it not unfrequently happens, as in the specimen figured, that all the ribs are serrated with minute rudely-developed scales in the vicinity of the umboes. The shell is very copiously groove-striated throughout, and the striae are much stronger in the interstices between the ribs than upon them.




7a



5b



5c



7b

T R I D A C N A .

PLATE VII.

Species 7. (Fig. *a, b*, Mus. Cuming.)

TRIDACNA CUMINGII. *Trid. testá ovatá vel oblongo-ovatá, gibbosiusculá, quadri- ad quinque-costatá, costis conspicuè radiatim liratis, nitidè regulariter squamatis, squamis erectis, costarum interstitiis copiosè corrugato-liratis; costis squamisque citrinis, costarum interstitiis vividè rufescente-croceis.*

CUMING'S TRIDACNA. Shell ovate or oblong-ovate, rather gibbous, four- to five-ribbed, ribs conspicuously radiately ridged, neatly regularly sealed, scales erect, interstices between the ribs copiously wrinkle-ridged; ribs and scales lemon-yellow, interstices between the ribs bright reddish-saffron.

Hab. Philippine Islands; Cuming.


This very beautiful *Tridacna*, of which specimens are

represented in our Plate, in different stages of growth, appears to be very distinct from any hitherto described. The ribs and scales are of a delicate lemon-colour throughout, while the interstices are of a rich reddish-saffron. It is very neatly sealed, and the radiating ridges in the interstices between the ribs have a peculiar interruptedly-wrinkled character.


Fig. 5 *b, c*, (Mus. Cuming.)

The specimens here figured are intended to illustrate *T. compressa*, as distinguished from *T. elongata*, in an earlier stage of growth than the specimen described and figured in the preceding Plate. That at Fig. 5 *b* is a pale lemon-colour.

June, 1862.




8a



8b



9a



9b

TRIDACNA.

PLATE VIII.

Species 8. (Fig. *a*, *b*, Mus. Cuming.)

TRIDACNA FERRUGINEA. *Trid. testā suboblongo-ovatā, quinque-costatā, costis parum elevatis, ad latera sub-obsolete, densē squamato-serratis, interstitiis triliratis; sordidē albā, versus marginem ferrugineo tinctā.*

THE RUST-STAINED TRIDACNA. Shell rather oblong-ovate, five-ribbed, ribs but little raised, almost obsolete at the sides, densely scale-serrated, interstices two- to three-ridged; dead white, rust-stained towards the margin.

Hab. —?

This appears to me to be very generally distinguished from *T. crocea*, to which it is the nearest allied, by its more oblong form and rust-brown colouring. The same distinctions of character may be observed in young specimens of the respective species as given at Fig. 8 *b* and 9 *a*.

Species 9. (Fig. *a*, *b*, Mus. Cuming.)

TRIDACNA CROCEA. *Trid. testā latè oratā, versus umbones gibbosiusculā, quinque-costatā, costis parum elevatis, densē squamato-serratis, interstitiis subobscnè triliratis; albida, croceo-rubente tinctā et concentricè undatā.*

THE SAFFRON TRIDACNA. Shell broadly ovate, rather gibbose towards the umboes, five-ribbed, ribs but little elevated, densely scale-serrated, interstices rather obscurely three-ridged; whitish, stained and concentrically waved with saffron-red.

LAMARCK, Anim. sans vert. vol. vii. p. 10.


Hab. Philippine Islands; Cuming.

In consequence of the lunule of this species gaping more largely than usual, the shell is very gibbose about the umboes, and attenuately pinched at the margin. The ribs are but superficially developed, and the scales are merely stunted imbrications.

June, 1862.

MONOGRAPH
OF THE GENUS
HIPPOPSIS.

“O happy living things! no tongue
Their beauty might declare:
A spring of love gushed from my heart,
And I bless’d them unaware.”—*Coleridge*.



HIPPONUS.

PLATE I.

Genus HIPPOPUS, Meuschen.

Testa aequivalvis, regularis, inaequilateralis. Lunulā sere clausā. Cardo dentibus duobus compressis inaequalibus posticis insertis. Ligamentum marginale, externum.

Shell equivalve, regular, inequilateral, with the lunule almost closed. Hinge composed of two unequal, compressed, posterior teeth, interlocking with one another. Ligament marginal, external.

The genus *Hippopus*, of which the well-known *Chama hippopus* of Linnaeus is still the only species known, differs from *Triducna*, the nearest allied form, in having the lunule, the anterior concave depression of the shell, nearly closed. It will be readily understood, therefore, that the animal has very little, if any, byssus. It seems strange that no second species of this well-defined generic type should have been discovered, but such is the fact. *Hippopus maculatus* has been known for more than a century, and is collected and imported in great abundance from the Eastern seas.

Species 1. (Mus. Brit.)

HIPPOPUS MACULATUS. *Hipp. testā subtrigono-ovatā, trapeziformi, anticē latē lunulatā, medio ventricosā, versus latera constrictā, radiatim costatā et liratā, profusè squamatā, squamis pareis, foliolatis, spinoso-imbricatis; marmoreo-albā, purpureo-roseo vividè maculata.*

THE SPOTTED HIPPOPUS. Shell somewhat triangularly ovate, trapeziform, anteriorly broadly lunulate, ventricose in the middle, constricted towards the sides, radiately ribbed and ridged, profusely scaled, scales small, leaf-like, spinosely imbricated; marble-white, richly spotted with purple-rose.

Chama hippopus, Linnaeus, Syst. Nat. p. 1137.

Hippopus maculatus, Lamarek.

Hab. Eastern seas.

This very elegant shell is too well known to need any comment on its specific characters, which are in all respects unique. No other species of the genus has been collected, and there is no variation even of the species worth calling attention to.

June, 1862.

MONOGRAPH
OF THE GENUS
A N A T I N A.

"Therefore bear you the lantern."—*Shakespeare*.



A N A T I N A.

PLATE I.

Genus ANATINA, Lamarck.

Testa æquivalvis, inæquilateralis, tenuis, hyalina, argenteo-alba, laevigata, interdum circa umbones concentricè plicata, versus margines plus minus granulata, granulis ad latus posticum in lineam de umboibus abruptè cessantibus, latere antico rotundato, postico in rostrum expansum plerumque producto, lateribus ambobus hiantibus. Cardo processu cochleariformi, septo obliquo sustentato, in utrâque valvâ, ligamentum continente, compitus.

Shell equivalve, inequilateral, thin, hyaline, silvery white, smooth, sometimes concentrically plicated around the umboes, more or less granulated towards the margins, granules abruptly ceasing at the posterior side in a line from the umboes, anterior side rounded, posterior generally produced into a pouted beak; both sides gaping. Hinge composed of a spoon-shaped process, supported by an oblique septum, in each valve, containing the ligament.

An interesting genus of the *Mya* tribe, ranging along the eastern shores of the eastern hemisphere from Japan to Tasmania, and even to the Antarctic Ocean, with very little variation in the general typical character of the shell. There are, however, two well-marked forms of *Anatina*, one in which the posterior side of the shell is broadly truncated, as in *A. truncata* and *elliptica*, the other, much the more numerous in species, in which the posterior side is produced into a pouted beak, gaping, of course, in both forms for the passage of the siphons, which are united throughout their length in one sheath. The shell of *Anatina* is invariably thin and hyaline, and of a silvery whiteness, crowded towards the margins with fine grains, which abruptly cease on the posterior side where the shell commences to be beaked, that is to say, on a constricted line sloping from the umboes.

The hinge of *Anatina* is composed of a prominent spoon-shaped process in each valve, developed rather obliquely in a direction posterior to centre, and supported by a firm septum. There is also a fine linear septum in front of it, and both radiate towards the interior. The hinge processes work upon each other with the aid and action of a contained elastic ligament.

Of the thirty species described in the following monograph the habitats of twenty-two are known. Two are from Japan, six from the Philippines and Moluccas, one from Borneo, one from Ceylon, eleven from Australia and

Tasmania, and one, a very remarkable species, from the island of New South Shetland, in the Antarctic Ocean.

Species 1. (Mus. Cuming.)

ANATINA CUMINGII. *Anat. testâ ovatâ, tumidinsculâ, tenui, hyalinâ, versus marginem posticum profusè granulatâ, latere antico compressè rotundato, postico constricto-rostrato, concentricè corrugato, expanso-hiantâ.*

CUMING'S ANATINA. Shell ovate, rather swollen, thin, hyaline, profusely granulated towards the posterior margin, anterior side compressly rounded, posterior constrictedly beaked, concentrically wrinkled, not granulated, expandedly gaping.

VALENCIENNES, MS. in Mus. Cuming.

Hab. —?

This elegant species is chiefly distinguished by its constrictedly pouted posterior extremity being concentrically wrinkled.

Species 2. (Mus. Cuming.)

ANATINA SIPHONATA. *Anat. testâ subabbreviato-ovatâ, medio tumidâ, tenui, hyalinâ, circa umbones concentricè plicatâ, versus marginem posticum profusè granulatâ, latere antico rotundato, postico attenuato et constricto-rostrato, remotè concentricè corrugato, hand granulato, subtubuloso-hiantâ.*

THE SPOUTED ANATINA. Shell rather shortly ovate, swollen in the middle, thin, hyaline, concentrically plicated around the umboes, profusely granulated towards the posterior margin, anterior side rounded, posterior attenuated, and constrictedly beaked, distantly concentrically wrinkled, not granulated, somewhat tubularly gaping.

Hab. Borneo.

Allied to the preceding species, but a shorter form, with the posterior extremity even more attenuated and constrictedly pouted.

Species 3. (Mus. Cuming.)

ANATINA BULLATA. *Anat. testâ subquadrato-ovatâ, gibbosâ, subhyalinâ, concentricè subrugoso-striatâ, latere antico latè tumido-truncato, postico attenuato, flexuoso-rostrato.*

THE BUBBLE ANATINA. Shell somewhat squarely ovate, gibbous, subhyaline, concentrically rather rugosely striated, anterior side broadly tumidly truncated, posterior side attenuated, flexuously beaked.

VALENCIENNES, MS. in Mus. Cuming.

Hab. Sibonga, Island of Zebu, Philippines (in fine sand, at the depth of thirty fathoms); Cuming.

Of a characteristic flexuous growth, broadly tumidly truncated in front.

Species 4. (Mus. Cuming.)

ANATINA VALENCIENNESII. *Anat. testā ovatā, subhyalinā, versus margines profusè granulatā, lateribus latè rotundatis, postico subconstricto, hiante, marginibus vir expansis.*

VALENCIENNES' ANATINA. Shell ovate, subhyaline, profusely granulated towards the margins, sides broadly rounded, posterior a little constricted, gaping, margins scarcely expanded.

Hab. China.

In this fine species there is only a faint constriction of the posterior extremity, the granules, which are very crowded towards the margins, cease, as usual, very abruptly at the line of constriction.

Species 5. (Mus. Cuming.)

ANATINA FLEXUOSA. *Anat. testā subelongato-oratā, tenui, hyalinā, crassè granulatā, circa umbones plicato-corrugatā, latere antico rotundato, subproducto, postico flexuoso-attenuato, rostrato, modicè hiante.*

THE FLEXUOUS ANATINA. Shell rather elongately ovate, thin, hyaline, coarsely granulated, plicately wrinkled round the umboes, anterior side rounded, rather produced, posterior flexuously attenuated, beaked, moderately gaping.

Hab. Japan.

Well distinguished in being plicately wrinkled around the umboes, while the posterior extremity is peculiarly flexuously attenuated.

Species 6. (Mus. Cuming.)

ANATINA SUBROSTRATA. *Anat. testā oblongo-ovatā, tenui, hyalinā, plus minus granulatā, latere antico rotundato, postico latè expanso-rostrato.*

THE SLIGHTLY-BEAKED ANATINA. Shell oblong-ovate, thin, hyaline, more or less granulated, anterior side rounded, posterior broadly expandedly beaked.

LAMARCK, Anim. sans vert. vol. vi. p. 78.

Solen anatinus, Linnaeus.

Hab. —?

In this shell, which M. Valenciennes, from a comparison probably with Lamarck's type, set apart in Mr. Cuming's collection as the species under consideration, the posterior extremity is peculiarly largely beaked.

Species 7. (Mus. Cuming.)

ANATINA LATERNA. *Anat. testā subquadrato-oblongā, compressiusculā, hyalinā, profusè granulatā, latere antico rotundato, postico flexuoso-truncato, vix rosstrato.*

THE LANTERN ANATINA. Shell somewhat squarely oblong, rather compressed, hyaline, profusely granulated, anterior side rounded, posterior flexuously truncated, scarcely beaked.

LAMARCK, Anim. sans vert. vol. vi. p. 77.

Hab. North Australia.

A beautifully delicate shell of firm growth, rather compressed, with only a moderately beaked flexuous gaping of the posterior extremity.




8



9



10



11



12



13



14

ANATINA.

PLATE II.

Species 8. (Mus. Cuming.)

ANATINA ANSERIFERA. *Anat. testā subquadrato-oblongā, hyalinā, concentricē modicē rugoso-lineatā, plus minus granulatā, lateribus subæqualibus, hiantibus, antico vir longiore, subquadrato-rotundato, postico obliquē rotundato.*

THE GOOSE-BILL ANATINA. Shell somewhat squarely oblong, hyaline, concentrically rugosely lined, more or less granulated, sides nearly equal, moderately gaping, the anterior scarcely the longer, squarely rounded, the posterior obliquely rounded.

Mya anserifera, Spengler, Schrift. Nat. Ges. zu Copenhagen, vol. iii. p. 32. n. 8; Chemn. Conch. Cab. vol. xi. p. 193. vign. 26. A, B.

Solen Spengleri, Gmelin.

Hab. Tasmania; R. C. Gunn.

A comparatively smooth shell with the sides nearly equal.

Species 9. (Mus. Cuming.)

ANATINA GRACILIS. *Anat. testā angustē oblongā, subcylindraceā, hyalinā, lavigatā, copiosē minutē granulatā, granulis ad latus anticū gradatim evanidis, lateribus subæqualibus, ambobus rotundatis, modicē hiantibus, antico subattenuato.*

THE SLENDER ANATINA. Shell narrowly oblong, somewhat cylindreal, hyaline, smooth, copiously minutely granulated, granules gradually fading on the anterior side, sides nearly equal, both rounded, moderately gaping, the anterior rather attenuated.

Hab. Moreton Bay, Australia.

The sides of this species are nearly equal, the valves are narrowly gibbous, comparatively free from concentric wrinkles, but copiously minutely granulated. The granules, though disappearing on the anterior side, do not end, as in most species, in an abrupt linear angle.

Species 10. (Mus. Cuming.)

ANATINA MARILINA. *Anat. testā suboblongo-quadratā, hyalinā, submargaritaceā, versus marginem conspicue lirato-rugatā et granulatā, valvis subæqualibus, utrinque compressē rotundatis, parum hiantibus.*

THE ASH ANATINA. Shell rather oblong-square, hyaline, rather pearly, conspicuously wrinkle-ridged and granulated towards the margin, valves nearly equal, compressly rounded at each end, but little gaping.

VALENCIENNES, MS. in Mus. Cuming.

Hab. Australia.

Compared with the allied species, the valves of this are unusually equal, and but little gaping, the shell has, moreover, a more pearly aspect.

Species 11. (Mus. Cuming.)

ANATINA TRUNCATA. *Anat. testā elongato-oblongā, sub-compressā, hyalinā, solidiusculā, concentricē rudē rugatā, granulatā, granulis ad latus anticū abruptē evanidis, lateribus subæqualibus, antico quadrato-truncato, valde hante, postico compressē rotundato, modicē hante.*

THE TRUNCATED ANATINA. Shell elongately oblong, rather compressed, hyaline, rather solid, concentrically rudely wrinkled, granulated, granules abruptly fading on the anterior side, sides nearly equal, anterior squarely truncated, largely gaping, posterior compressly rounded, moderately gaping.

LAMARCK, Anim. sans. vert. vol. vi. p. 77.

Anatina olorina, Valenciennes.

Hab. Bay of Manila, Philippines (at low water); Cuming.

This well-marked species, which I take to be Lamarck's *A. truncata*, was collected in great abundance by Mr. Cuming in the Bay of Manila.

Species 12. (Mus. Cuming.)

ANATINA CRECCINA. *Anat. testā subelongato-oblongā, cylindraceā, tenui, hyalinā, minutē granulatā, prope umbones sublatē obliquē rugatā, latere antico attenuato-rostratā, rotundato-hante, extremitatibus reflexis, postico compressiusculo, modicē hante, extremitatibus concentricē rugatis.*

THE ROUGH-BILL ANATINA. Shell rather elongately oblong, cylindrical, thin, hyaline, minutely granulated, rather broadly obliquely wrinkled near the umboes, anterior side attenuately beaked, rotundately gaping,

extremities reflected, posterior side rather compressed, moderately gaping, extremities concentrically wrinkled.

VALENCIENNES, MS. in Mus. Cuming.

Hab. Adelaide, South Australia (in sandy mud, at a depth of ten fathoms).

Younger specimens of this species in Mr. Cuming's collection, show its characteristics better than the adult. An apparently slight but important specific character consists in the presence of a few faint, oblique, not concentric, wrinkles near the umbo of each valve; and the species inclines to be of a rather narrow cylindrically oblong form, with the anterior extremities beaked in a rather confined rotundately pouting manner.

Species 13. (Mus. Cuming.)

ANATINA BOSCHIASINA. *Anal. testā subtrigono-oblongā, inflatā, hyalinā, concentricè corrugato-liratā, granulata, granulis ad latus anticū abruptè evanidis, latere antico attenuato, modicè hiante, extremitatibus subreflexis, postico late, subgloboso, vix hiante.*

BOSCH'S ANATINA. Shell somewhat triangularly oblong, inflated, hyaline, concentrically wrinkle-ridged, granulated, granules abruptly disappearing on the anterior side, anterior side attenuated, moderately gaping, extremities slightly reflected, posterior side broad, rather globose, scarcely gaping.

VALENCIENNES, MS. in Mus. Cuming.

Hab. Island of Negros, Philippines (found in mud at low water); Cuming.

Rather more attenuated on the anterior side than usual, which attenuation contrasts very perceptibly with the somewhat globosely inflated posterior side.

Species 14. (Mus. Cuming.)

ANATINA ELLIPTICA. *Anal. testā abbreviato-oblongā, subtrapeziformi, crassiusculā, opaco-albā, rudē concentricè rugosā, intus pallidè prismaticā, latere antico latissime truncato, valde hiante, postico rotundato, parum hiante.*


THE ELLIPTIC ANATINA. Shell shortly oblong, subtrapeziform, rather thick, opaque white, rudely concentrically wrinkled, faintly prismatic in the interior, anterior side very broad, largely gaping, posterior rounded, slightly gaping.

KING and BRODERIP, Zool. Journ. vol. v. p. 335.

Anatina prismatica, Sowerby.

Hab. New South Shetland, Antarctic Ocean; Lieut. Kendall, Dr. Eights.

An interesting broadly truncated *Panopaea*-like species, with a remarkably large anterior gape. The specimen from which our figure is taken, was found on the beach at New South Shetland after a gale of wind, by Dr. Eights of Philadelphia, and described by Mr. Sowerby in the Proceedings of the Zoological Society for 1834. A specimen had, however, been previously brought to England by Captain King, to whom it had been presented by Lieutenant Kendall, of H.M.S. Chanticleer, who found it in the very same locality. This specimen, which is in the British Museum, was described by King and Broderip in the Zoological Journal for 1830-31, and figured by Dr. Gray in Griffith's Cuvier's Animal Kingdom, pl. xxii. f. 3.



A N A T I N A.

PLATE III.

Species 15. (Mus. Cuming.)

ANATINA CISTELLA. *Anat. testā quadrato-oblongā, sub-compressā, tenuiculā, hyalinā, nisi ad latus anticū copiosē granulatā, latere postico compressē subrostrato, modicē hiantē, antico latē rotundato, vix hiantē.*

THE CASKET ANATINA. Shell squarely oblong, somewhat compressed, rather thin, hyaline, copiously granulated except at the posterior side, anterior side compressly slightly beaked, moderately gaping, anterior side broadly rounded, scarcely gaping.

Hab. Moluccas.

Of a distinct squarely oblong form, broad and rather compressed.

Species 16. (Mus. Cuming.)

ANATINA ATTENUATA. *Anat. testā elongato-oblongā, sub-angustā, tenui, hyalinā, nisi ad latus anticū copiosē granulatā, latere postico attenuato, subtubuloso, antico latiore, rotundato.*

THE ATTENUATED ANATINA. Shell elongately oblong, rather narrow, thin, hyaline, copiously granulated, except at the anterior side, posterior side attenuated, somewhat tubulous, anterior side broader, rounded.

Hab. Sydney.

Of a rather narrow, elongated growth, the posterior side being attenuated and inclined to be tubular.

Species 17. (Mus. Cuming.)

ANATINA BLAINVILLEI. *Anat. testā oblongo-ovatā, subflexuosa, tenui, hyalinā, minutē granulatā, latere postico subproducto, subrostrato, tubulos-hiantē, antico subattenuato, rotundato.*

BLAINVILLE'S ANATINA. Shell oblong-ovate, rather flexuous, thin, hyaline, minutely granulated, posterior side rather produced, slightly beaked, tubularly gaping, anterior side rather attenuated, rounded.

VALENCIENNES, MS. in Mus. Cuming.

Hab. Island of Burias, Philippines (in sandy mud at a depth of from ten to twenty fathoms); Cuming.

Very thin and transparent, of a rather narrow subflexuous form.

Species 18. (Mus. Cuming.)

ANATINA CONSTRICTA. *Anat. testā subquadrato-oblongā, compressiusculā, hyalinā, minutissimē granulatā, latere postico breviusculo, curvato, constricto-rostrato, tubulos-hiantē, antico producto, latē rotundato, vix hiantē.*

THE CONSTRICTED ANATINA. Shell somewhat squarely oblong, rather compressed, hyaline, very minutely granulated, posterior side rather short, curved, constrictedly beaked, tubularly gaping, anterior side produced, broadly rounded, scarcely gaping.

Hab. North Australia.

An oblong beaked form, in which the beak is more than usually constricted.

Species 19. (Fig. a, b, Mus. Cuming.)

ANATINA LABIATA. *Anat. testā oblongo-ovatā, tenui, eximiē hyalinā, minutissimē granulatā, latere postico abrupte flexuoso-attenuato, marginibus tubuloso-expansis, antico rotundato.*

THE POUTING ANATINA. Shell oblong-ovate, thin, delicately hyaline, very minutely granulated, posterior side abruptly flexuously attenuated, with the margins tubularly pouted, anterior side rounded.

Hab. Ceylon; E. L. Layard.

An extremely delicate species, very abruptly tubularly attenuated at the posterior end, with the margins expensively pouted, varying in character as in the two specimens figured.

Species 20. (Mus. Cuming.)

ANATINA TASMANICA. *Anat. testā subaequilateralē, oblongā, compressiusculā, subhyalinā, subobsoletē granulatā, lateribus rotundatis, parum hiantibus, postico subangustiore.*

THE TASMANIAN ANATINA. Shell nearly equilateral, oblong, rather compressed, subhyaline, rather obsoletely granulated, sides rounded, but little gaping, posterior rather the narrower.

Hab. Tasmania.

A shell of very simple character, in which the sides are nearly equal.



21



22



23




24



25



26



27



28



29



30

A N A T I N A.

PLATE IV.

Species 21. (Mus. Cuming.)

ANATINA NAVICULA. *Anat. subæquilateralis, oblonga, gibbosæ, hyalinæ, granoso-seabrosa, lateribus rotundatis, parum hiantibus, antico sublatiore, obtuse tumido-truncato.*

THE BOAT ANATINA. Shell nearly equilateral, oblong, gibbous, hyaline, granosely seaborous, sides rounded, but little gaping, anterior side rather the broader, obtusely tumidly truncated.

Hab. —?

An oblong gibbous shell, with very much the form of *Novaculina*.

Species 22. (Mus. Cuming.)

ANATINA FABA. *Anat. testa ovata, utrinque attenuata, tenuissimæ, hyalinæ, concentricè striato-lirata, latere postico flexuoso-rostrato, vix hiante.*

THE BEAN ANATINA. Shell ovate, attenuated at each end, very thin, hyaline, concentrically striately ridged, posterior side flexuously beaked, scarcely gaping.

Hab. Brisbane, east coast of New Holland.

A small *Nucula*-shaped species, very thin and transparent, but distinctly concentrically striately ridged, the posterior side is attenuately flexuously beaked, scarcely gaping.

Species 23. (Mus. Cuming.)

ANATINA AMPHORA. *Anat. testa orata, medio ventricosa, lenii, hyalinæ, concentricè striato-lirata, latere postico abruptè rostrato, tubuloso-hiante, marginibus exsertis, antico attenuato-rotundato.*

THE PITCHER ANATINA. Shell ovate, ventriose in the middle, thin, hyaline, concentrically striately ridged, posterior side abruptly beaked, tubularly gaping, margins pouted, anterior side attenuately rounded.

Hab. Island of Burias, Philippines.

A delicately transparent, concentrically ridged species, broadly ventriose in the middle, in which the posterior side is very abruptly beaked and pouted.

Species 24. (Mus. Cuming.)

ANATINA RECTA. *Anat. testa subæquilateralis, elongato-oblonga, recta, sordidè albæ, lateribus rotundatis, postico subobsoletè concentricè corrugato, antico latiore; subexpanso, lateribus ambobus modice hiantibus.*

THE STRAIGHT ANATINA. Shell nearly equilateral, elongately oblong, straight, dull white, sides rounded, posterior rather obsoletely concentrically wrinkled, anterior broader, slightly expanded, both sides moderately gaping.

Hab. Port Phillip, Australia.

A straight, nearly equilateral species, gaping at both ends, with the margins inclined to expand.

Species 25. (Mus. Cuming.)

ANATINA CORRUGATA. *Anat. testa oblonga, subangustata, tenni, hyalinæ, latere postico concentricè corrugato, attenuato-truncato, modice hiante, antico subventricoso, rotundato, modice hiante.*

THE WRINKLED ANATINA. Shell oblong, rather narrow, thin, hyaline, posterior side concentrically wrinkled, attenuately truncated, moderately gaping, anterior rather ventriose, rounded, moderately gaping.

Hab. Island of Ticao, Philippines (in coral sand at a depth of about six fathoms); Cuming.

The posterior side is in this species attenuately truncated in form, and faintly but distinctly wrinkled.

Species 26. (Mus. Cuming.)

ANATINA VAGINA. *Anat. testa oblongo-ovata, curvata, tenuissimæ, hyalinæ, copiosè minute granulata, latere postico breviusculo, constricto-rostrato, tubuloso-hiante, marginibus expansis, antico compressè rotundato, parum hiante.*

THE SHEATH ANATINA. Shell oblong-ovate, curved, very thin, hyaline, copiously minutely grained, posterior side rather short, constrictedly beaked, tubularly gaping, margins pouted, anterior compressly rounded, but little gaping.

Hab. Moreton Bay, Australia.

A very delicate hyaline shell, of characteristic form, the

posterior side being short, constrictedly beaked and pouted, but not angled.

Species 27. (Mus. Cuming.)

ANATINA LIMICOLA. *Anat. testá subaequilateralí, cylindraceo-oblongá, argenteo-hyaliná, levigatá, latere postico abrupte truncato, ampliter hiante, antico latiore, tumido, rotundato, parum hiante.*

THE MUD-DWELLING ANATINA. Shell nearly equilateral, cylindrically oblong, silvery hyaline, smooth, posterior side abruptly truncated, largely gaping, anterior side broader, swollen, rounded, but little gaping.

Hab. Japan.

Closely allied to *A. truncata*, of the Philippine Islands, and equally abundant, perforating the mud in all directions.

Species 28. (Mus. Cuming.)

ANATINA GRACILIS. *Anat. elongato-pyriformi, tenuissimá, hyaliná, concentricè striato-liratá, latere postico constricto-rostrato, rostro attenuato-producto, antico tumido-expanso, rotundato.*

THE SLENDER ANATINA. Shell elongately pear-shaped, very thin, hyaline, concentrically striately, ridged, posterior side constrictedly beaked, beak attenuately produced, anterior tumidly expanded, rounded.

Hab. Port Curtis, Australia.

A very thin, transparent shell, remarkable for the attenuated prolongation of the beaked posterior side.

Species 29. (Mus. Cuming.)

ANATINA ARGENTEA. *Anat. testá ovalá, tenui, lœrigatá, argenteá, hyaliná, latere postico constricto-attenuato, vix rostrato, modicè hiante, antico rotundato, parum hiante.*

THE SILVERY ANATINA. Shell ovate, thin, smooth, silvery, hyaline, posterior side constrictedly attenuated, scarcely beaked, moderately gaping, anterior side rounded, but little gaping.

Hab. —?

This species differs from all others in having the posterior side constrictedly attenuately produced, with very little inclination to become beaked.

Species 30. (Mus. Cuming.)

ANATINA EXIMIA. *Anat. testá ovalá, curvatá, tenui, hyaliná, circa umbones eximiè plicatá, latere postico constricto-attenuato, rostrato, hiante, marginibus subexpansis, antico arcuato-rotundato, modicè hiante.*

THE DELICATE ANATINA. Shell ovate, curved, thin, hyaline, neatly plicated round the umbones, posterior side constrictedly attenuated and beaked, gaping, margins a little pouted, anterior side arcuately rounded, moderately gaping.

Hab. —?

An elegantly curved shell, attenuately beaked on the posterior side, with the margins a little pouted.

A N A T I N A .


	Plate.	Species.		Plate.	Species.
amphora, <i>Reeve</i>	IV.	23	<i>gracilis</i> , <i>Reeve</i>	II.	9
anserifera (<i>Mya</i>), Spengler	II.	8	<i>labiata</i> , <i>Reeve</i>	III.	19
argentea, <i>Reeve</i>	IV.	29	<i>laterna</i> , <i>Lamarck</i>	I.	7
attenuata, <i>Reeve</i>	III.	16	<i>limicola</i> , <i>Reeve</i>	IV.	27
Blainvillei, <i>Valenciennes</i>	III.	17	<i>maritima</i> , <i>Valenciennes</i>	II.	10
Boschasma, <i>Valenciennes</i>	II.	13	<i>navicula</i> , <i>Reeve</i>	IV.	21
bullata, <i>Valenciennes</i>	I.	3	<i>olorina</i> , <i>Valenciennes</i>	II.	11
cistella, <i>Reeve</i>	III.	15	<i>prismatica</i> , <i>Sowerby</i>	II.	14
constricta, <i>Reeve</i>	III.	18	<i>prolongata</i> , <i>Reeve</i>	IV.	28
corrugata, <i>Reeve</i>	IV.	25	<i>recta</i> , <i>Reeve</i>	IV.	24
creccina, <i>Valenciennes</i>	II.	12	<i>siphonata</i> , <i>Reeve</i>	I.	2
Cumingii, <i>Valenciennes</i>	I.	1	<i>subrostrata</i> , <i>Lamarck</i>	I.	6
elliptica, <i>King and Broderip</i> ...	II.	14	<i>Tasmanica</i> , <i>Reeve</i>	III.	20
eximia, <i>Reeve</i>	IV.	30	<i>truncata</i> , <i>Lamarck</i>	II.	11
faba, <i>Reeve</i>	IV.	22	<i>vagina</i> , <i>Reeve</i>	IV.	26
flexuosa, <i>Reeve</i>	I.	5	<i>Valenciennesii</i> , <i>Reeve</i>	I.	4

ERRATUM.

Plate IV. Sp. 28. For *A. gracilis*, Reeve, read *A. prolongata*, Reeve.

MONOGRAPH
OF THE GENUS
T U G O N I A.

"My instruction may serve to naturalize thee." — *Shakespeare.*




1a



2



3



1b



4a



4b.



5

TUGONIA.

PLATE I.

Genus TUGONIA, Gray.

Testa æquivalvis, valde inæquilateralis, plus minus globosa, opaco-alba, solidiuscula, radiatim tenuilirata vel striata, liris ad aream anticom interdum desinuentibus, latere antico perampliter gibboso, postico brevissimo, coarctato-rostrato, hante, orificio rotundo, marginibus callosis. Cardo processu parvo cochleariformi, septo brevi pero obliquo sustentato, in utrâque valvâ, ligamentum continente, compositus.

Shell equivalve, very inequilateral, more or less globose, opake-white, rather solid, radiately finely ridged or striated, ridges sometimes wanting on the anterior area, anterior side very largely gibbous, posterior very short, contractedly beaked, gaping, orifice round, with the margins callous. Hinge composed of a small spoon-shaped process, supported by a short very oblique septum in each valve containing the ligament.

The genus *Tugonia* was founded by Dr. Gray for the reception of a characteristic form of *Anatina* inhabiting the west coast of Africa, originally described by Adanson, in his 'Natural History of Senegal,' with the name *Le Tugon*. The shell is of a rather solid opake substance, curiously gibbously globose, with the posterior extremity contracted, where the siphonic sheath of the animal is exerted, into a conspicuous subtubularly-beaked orifice. Adanson mistook it for the shell of a *Pholas*. Chemnitz, who, as shown by his admirable figures (Conch. Cab. vol. vi. pl. 2. f. 13–16), possessed specimens, described it as living in abundance near the mouth of the Niger, and named it "The Guinea *Mya analina*." The genus is, however, rare in collections, and Mr. Cuming has obtained no precise information of its habitat. The specimens in his collection given in the accompanying Plate, appear to represent six species, varying in form and convexity and in the details of sculpture. Some are radiately ridged throughout; in some the radiating ridges abruptly terminate on the posterior side at a line sloping from the umbo. Five of the species are from West Africa, one from Ceylon.

Species 1. (Fig. a, b, Mus. Cuming.)

TUGONIA GUINEENSIS. *Tug. testâ globosâ, gibbosissimâ, albâ, radiatim tenuiliratâ, liris striis concentricis subobsoletè fenestratis, striis anticis versus marginem evanidis, latere antico peramplio, postico abruptè rostrato.*

THE GUINEA TUGONIA. Shell globose, very gibbous, white, radiately finely ridged, ridges somewhat obsoletely latticed with concentric striae, anterior striae fading towards the margin, anterior side very large, posterior abruptly beaked.

Le tugon, Adanson, Hist. Nat. Coq. du Senegal, pl. 19. f. 2.

Mya analina, pars, Gmelin.

Mya analina Guineensis, Chemnitz.

Anatina globulosa, Lamarck.

Mya tugon, Deshayes.

Hab. West coast of Africa.

M. Deshayes proposed to adopt the French appellative of Adanson for the name of this species, but such a compound is unclassical. Chemnitz was the first to distinguish the shell specifically from those included by the older writers under the general term of *Mya analina*.

Species 2. (Mus. Cuming.)

TUGONIA DIVARICATA. *Tug. testâ subquadrato-ovatâ, candidâ, radiatim minutissimè linearè incisâ, versus marginem posticum nitidè divaricatum corrugatâ, latere antico rotundato, postico latè subangulato-rostrato.*

THE DIVARICATE TUGONIA. Shell somewhat squarely ovate, white, radiately very minutely linearly engraved, towards the posterior margin prettily divaricately wrinkled, anterior side rounded, posterior broadly subangularly beaked.

Hab. Ceylon; E. L. Layard.

Of quite a distinct type from the West African species, the shell being apparently smooth, but under the lens minutely radiately engraved, while towards the posterior and posterior-ventral margins it is sculptured with very characteristic divaricating wrinkles.

Species 3. (Mus. Cuming.)

TUGONIA COMPRESSA. *Tug. testâ suborbiculari-ovatâ, versus marginem compressâ, lutescente-albâ, undique radiatim liratâ, liris irregularibus, striis concentricis clathratis et tenè crenulatis, latere antico peramplio, postico brevissimo, latè rostrato.*

THE COMPRESSED TUGONIA. Shell suborbiculary ovate, compressed towards the margin, yellowish-white, ra-

diately ridged throughout, ridges irregular, latticed and finely crenulated with concentric striae, anterior side very large, posterior very short, broadly beaked.
Hab. West Africa.

The shell is in this species comparatively rather compressed, while the radiating ridges are stronger, crenulated on the summit, finely latticed in the interstices.

Species 4. (Fig. 4 *a*, Mus. Cuming.)

TUGONIA SEMISULCATA. *Tug. testā obliquē globosā, gibbosā, albā, radiatim tenuiliratā, liris striis concentricis minutē dense clathratis, areā anticā sine liris radianibus, latere postico modice rostrato.*

THE HALF-GROOVED TUGONIA. Shell obliquely globose, gibbous, white, radiately finely ridged, ridges minutely closely latticed with concentric striae, anterior area without radiating striae, posterior side moderately beaked.

ADAMS, Gen. Rec. Moll. vol. ii. p. 355.

Hab. West Africa.

The anterior area of this and the following species is sculptured with concentric ridges, but the radiating ridges are wanting. They cease at the side, in a line sloping from the umbo. The grooved contraction of the specimen here represented, as seen in the upper part of the figure, is merely an accident of growth.

Species 5. (Mus. Cuming.)

TUGONIA NOBILIS. *Tug. testā globoso-oblongā, antīcē gibbosā, albā, radiatim tenuiliratā, liris dense clathratis ad latera obsoletis, latere antīco obtusē angulato, postico truncato-rostrato.*

THE NOBLE TUGONIA. Shell globosely oblong, anteriorly gibbous, white, radiately finely ridged, ridges closely latticed, obsolete at the sides, anterior side obtusely angled, posterior truncately beaked.

ADAMS, Gen. Rec. Moll. vol. ii. p. 355.

Hab. West Africa.

This species scarcely differs from *T. anatina*, except in being of a more oblong form.

Species 6. (Fig. 4 *b*, Mus. Cuming.)

TUGONIA SIPHONATA. *Tug. testā obliquē globosā, albā, undique radiatim liratā, liris confertis, irregularibus, minutē crenulatis, latere antīco gibboso, peramplo, postico subproducto, expanso-rostrato.*


THE SPOUTED TUGONIA. Shell obliquely globose, white, everywhere radiately ridged, ridges close, irregular, minutely crenulated, anterior side gibbous, very large, posterior rather produced, pointedly beaked.

Hab. West Africa.

It will be observed by the numbering of this species that it was taken originally to be a variety of *T. semisulcata*, but the radiating ridges, as in *T. compressa*, range throughout; they do not stop short of the anterior area.

MONOGRAPH
OF THE GENUS
CHAMOSTREA.

"They that go down to the sea in ships, and occupy
their business in great waters ;
These men see the works of the Lord, and his wonders
in the deep."—*Psalms*.



1a.




1b.



1d.



1c.



1e.

CHAMOSTREA.

PLATE I.

Genus CHAMOSTREA, *De Roissy.*

Testa valde inaequivalvis, valde inaequilateralis, umboibus involutis, latere postico acutè trigono, antico plano, adhærente; valvā dextrā perampliter trigono-convexā, sinistrā planiusculā, operculoideā. Cardo dente parvo solido erecto in valvā sinistrā in fossulam valvæ dextræ intrante, fossulāque alterā ligamentum cum ossiculo continente, ligamento elongato etiam ad marginem.

Shell very inequivalve, very inequilateral, with the umbones involute, posterior side sharply triangular, anterior flat, adhering; right valve very large, triangularly convex, the left rather flat, lid-like. Hinge composed of a small solid erect tooth in the left valve, entering into a pit in the right valve, behind which there is another pit containing elastic ligament, with an ossicle attached; there is also a marginal ridge of ligaments.

The animal of this curious Australian bivalve has been described by Albany Hancock, but it is not at all clear that the plate assigned to it in the family of *Anatinidae* is correct. The shell is of very solid growth, flatly adhering to foreign bodies by the right valve, which is deeply triangular convex, while the left valve, which forms a lid to it, is of very peculiar growth. The posterior portion of this valve is developed, and corresponds with, the free posterior portion of the right valve, and is of the same concentrically striated growth, but the anterior portion of the left valve, which is developed with the adhering portion of the right valve, presents an abrupt change of growth, having a crimped wrinkled surface, with the lines of increase ranging obliquely.

The hinge of *Chamostrea* is unique in structure. In the left valve is a blunt erect tooth, fitting into a pit in the right valve, beside which, in each valve, there is another pit containing ligament with a shelly appendage or ossicle, quite distinct from the shell, attached to it. The valves open and shut with the aid of this moving ossicle, and in

opening dried specimens of the shell it will be found adhering to ligament either in the ligamentary pit of the right or left valve as accident may direct. In the specimen represented in the accompanying Plate, at Fig. 1 *b* and 1 *c*, the ossicle may be seen adhering to the ligament in the left valve. At Fig. 1 *d* it is again given by itself. Along the margin above the hinge is an auxiliary more elongated portion of ligament. In Fig. 1 *c* it may be seen commencing in a dark curve above the tooth-pit.

Lamarek, on the authority of M. Leschenault, gives Java as the habitat of this shell, but this must be a mistake. The only satisfactorily authenticated habitats are east coast of New Holland and New Hebrides. Fig. 1 *e* shows the mode in which specimens adhere to one another.

Species 1. (Fig. *a*, *b*, *c*, *d*, *e*, Mus. Cuming.)

CHAMOSTREA ALBIDA. *Cham. testā trigono-globosā, solidā, purpureo-roseā et fulvā, valvā dextrā concentricè rugoso-striatā, valvē sinistre diuidio postico concentricè rugoso-striato, altero obliquè peculiariter crispato-corrugato.*

THE WHITISH CHAMOSTREA. Shell triangularly globose, solid, purple-rose and fulvous, right valve concentrically roughly striated, posterior half of the left valve concentrically roughly striated, the other half obliquely peculiarly crisplately wrinkled.

Chama albida, Lamarek, Anim. sans vert. vol. vi. p. 585.

Chamostrea albida, De Roissy.

Cleidothærus Chamoides, Stutchbury.

Cleidothærus albidus, Deshayes.

Hab. Sydney, New Hebrides.

The Australian specimens of this genus are strongly hinged and rayed with purple-rose. The specimens in Mr. Cuming's collection from New Hebrides, one of which is given at Fig. 1 *a*, are uniformly pale and livid.

MONOGRAPH
OF THE GENUS
V E N U S.

"And in the symmetry of their parts is found
A power, like that of harmony in sound."—*Johastou*.




1



?a



?b



3.

V E N U S.

PLATE I.

Genus VENUS, *Linnæus.*

Testa subtrigono-ovata, vel orbicularis, crassiuscula, sèpissimè turgida, concentricè lirata vel laminata, interdum, autem raro, radiatim lirata vel striata, ad margines plerumque crenulata. Cardo incrassatus, in utrâque valvâ tridentatus, dentibus cardinalibus, divergentibus, interdum bifidis. Impressio pallii posticè sinuoso, sinu breviusculo.

Shell somewhat triangularly ovate or orbicular, rather thick, most frequently turgid, concentrically ridged or laminated, sometimes, but seldom, radiately ridged or striated, generally crenulated at the margins. Hinge thickened, three-toothed in each valve, teeth cardinal, diverging, sometimes bifid. Impression of the mantle situated posteriorly, sinus rather short.

The species referred to this genus are conspicuously distinguished from the rest of the family by their elaborate sculpture of concentric ridges variously developed in crenulated and reflected laminae, sometimes crossed in the interstices with radiating striae, and by their turgid growth. They are about equally distributed in the two hemispheres, in one chiefly on the west coast of America, especially Central America, and the West Indies, in the other chiefly among the Philippine Islands, Australia, and New Zealand. Seven species inhabit European seas, of which five are natives of our own shores. The laminated species, of which the beautiful Tasmanian *V. lamellata*, the *V. plicata*, *paphia*, *thiara*, and others are characteristic examples, are separated, however, by some authors under the generic name of *Chione*. The United States possess a very large species, *V. mercenaria*, which has been subdivided into four species, and then set apart as a genus of itself. A genus has also been made, *Gemma*, of a small pellucid species, which inhabits the same locality, and has been regarded as the fry of *V. mercenaria*. Lastly a small group of half-a-dozen species, of a peculiar flexuous form, inhabiting the West Indies and Central America, have been raised somewhat unnecessarily, for the animal is not known, to the rank of a separate genus, with the name *Anomalogardia*.

The *Veneres* are not remarkable for colour. A few of the species, chiefly those referred to *Chione*, are gaily painted, but the predominant tone is fulvous cream-colour, more or less indistinctly marked with rust-brown.

Species 1. (Mus. Cuming.)

VENUS OBLONGA. *Ven. testâ oblongo-ovatâ, subobliquâ, valde inaequilaterali, subdepressâ convexâ, squalidè fuscenscente-albâ, concentricè densè lamellato-liratâ, liris plus minus reflexis.*

THE OBLONG VENUS. Shell oblong-ovate, rather oblique, very inequilateral, rather depresso convex, dull fuscous-white, concentrically densely lamellately ridged, ridges more or less reflected.

Venus oblonga, Hanley, Ind. Test. Supp. pl. 16. f. 1.

Venus Zelandica, Gray.

Hab. New Zealand.

Varying occasionally in form in being more or less convex and lamellated, but always of the same dull colourless hue.

Species 2. (Fig. a, b, Mus. Cuming.)

VENUS LIMA. *Ven. testâ pectinato-ovatâ, albida, aurantio-fusco sparsim variegata, radiatim modice costata, costis rotundatis, imbricato-squamatis, lunula fulvo-fuscenscente.*

THE FILE VENUS. Shell pectinately ovate, whitish, sparingly variegated with orange-brown, radiately moderately ribbed, ribs rounded, imbricately scaled, lunule fulvous-brown.

Tapes lima, Sowerby, Thes. Conch. vol. i. p. 698. pl. 151. f. 144.

Chione lima, Deshayes.

Hab. New Zealand.

A seabrous radiately-ribbed species belonging to the pectinate division of the genus.

Species 3. (Mus. Cuming.)

VENUS CLATHRATA. *Ven. testâ cordato-ovatâ, subobliquâ, crassâ, ventricosa, lactea, rufescente-spadiceo pallide maculata et radiata, concentricè densè lirata, liris irregularibus, prope umbones tenuibus, deinde callosis et planatis, sulcis linearibus rudimentibus undique clathratis, liris ad aream posticam serrato-tumidis.*

THE LATTICED VENUS. Shell cordately ovate, rather


VENUS.—PLATE I.

oblique, thick, ventricose, cream-coloured, faintly blotched and rayed with reddish-fawn, concentrically densely ridged, ridges irregular, near the umbones thin, then callous and flattened, latticed throughout with radiating linear grooves, ridges on the posterior area serrately laminated.

DESHAYES, Pro. Zool. Soc. 1853, p. 3.

Venus Listeri, var., Sowerby.
Hab. —?


It is somewhat doubtful whether M. Deshayes is correct in separating this from *V. Listeri*. There is strong evidence of its being a large callously-sculptured form of that species.



4a.



5.



4b.



6.

VENUS.

PLATE II.

Species 4. (Fig. a, b, Mus. Cuming.)

VENUS MERCENARIA. *Ven. testā subtrigono-cordatā, compressiusculd, sordidē fulvo-ferruginea, in testā juniore fusco acutē undulato concentricē fasciatā, intus interdum nigricante-violacea, densē concentricē striatā, præcipue ad latera, latere postico obscurè flexuoso, obliquè curvato, antico brevi, lunulā conspicuā, sœpè pureo-castaneā.*

THE TRADE VENUS. Shell somewhat triangularly heart-shaped, rather compressed, dirty fulvous rust-colour, concentrically banded in the young shell with zigzag brown, sometimes deep-violet within, densely concentrically striated, especially at the sides, posterior side obscurely flexuous, obliquely curved, anterior short, lunule conspicuous, often purple-chestnut.

LINNÆUS, Syst. Nat. ed. 12, p. 1131.

Venus notata, Say.

Venus præparca, Say.

Venus obliqua, Anton.

Venus cyprinoides, Anton.

Hab. North America.

This species inhabits the Massachusetts shores of North America in greater or less abundance, and, as its name denotes, is used as an article of commerce. Dr. Gould mentions that in the markets of New York and Philadelphia, where *Venus mercenaria* is known by the name of Quahog, it is even more in request as an article of food than the *Mya arenaria*. Mr. Say's *V. notata* is the shell in a young state, marked with bands of zigzag brown, as

represented at Fig. 4 a. His *V. præparca* is the same without markings.

Species 5. (Mus. Cuming.)

VENUS SCULPTA. *Ven. testā orbiculari, subglobosā, albidā, fusco sparsim maculatā, areā ligamenti fuscā, concentricē liratā, liris regularibus, crispato-crenatis, subreflexis.*

THE SCULPTURED VENUS. Shell orbicular, rather globbose, whitish, sparingly spotted with brown, ligament area brown, concentrically ridged, ridges regular, crisplately crenated, slightly reflected.

DESHAYES, Pro. Zool. Soc. 1853, p. 3.

Hab. —?

The ridges of this species are a little reflected and curiously crisplately crenated.

Species 6. (Mus. Cuming.)

VENUS SPHERISULCA. *Ven. testā subtrigono-globosā, crassā, calcareo-albā, concentricē liratā, liris latiusculis, plicato-planatis, valle irregularibus.*



THE SPHERE-GROOVED VENUS. Shell somewhat triangularly globose, thick, chalk-white, concentrically ridged, ridges rather broad, plicately flattened, very irregular.

DESHAYES, MS. in Mus. Cuming.

Hab. —?

The ridges range concentrically over the shell in very irregular flattened folds.


April, 1863.



8a.



8b.



g

V E N U S.

PLATE III.

Species 7. (Mus. Cuming.)

VENUS ALBORADIATA. *Ven. testá ovatá, subcentricosá, fulvo-griseá, carneo el albo radiatá, concentricè irregulariter liratá, ad latera densè tenui-lamellatá, latere postico subflexuoso.*

THE WHITE-RAYED VENUS. Shell ovate, rather ventricose, fulvous-grey, rayed with flesh and white, concentrically irregularly ridged, at the sides densely thinly laminated, posterior side slightly flexuous.

SOWERBY, Thesaurus Conch. part xiv. p. 732. pl. 161. f. 199.

Hab. Gulf of Mexico.

A rather ventricose shell, faintly rayed with white and flesh-colour, which is more ruddy and broken up into spots towards the umboes.

Species 8. (Fig. a, b, Mus. Cuming.)

VENUS GRATA. *Ven. testá subquadrato-ovatá, albida, cinereo-griseo et fusco variè nebulatá, strigatá, et testellatá, intus interdum nigricante-violaceá, decussatim liratá et sulcatá, liris ad latera nodulatis, liris medianis posticè angulatis, latere postico subcompressè truncato, antico brevi, rotundato.*

THE WELCOME VENUS. Shell somewhat squarely ovate, whitish, variously clouded, streaked, and tessellated with ash-grey and brown, sometimes dark-violet in the interior, decussately ridged and grooved, ridges noduled at the sides, middle ridges angled poste-

riorly, posterior side rather compressly truncated, anterior short, rounded.

SAY, American Conchology, pl. 36.

Tapes grata, Sowerby.

Tapes tricolor, Sowerby.

Hab. Gulf of Mexico.

In this prettily-painted species the middle ridges radiating from the umboes to the margin are pressed to an angle posteriorly.

Species 9. (Mus. Cuming.)

VENUS MULTICOSTATA. *Ven. testá elliptico-ovatá, subventricosá, crassá, fulvo-spadiceá, carneo-fuscescente obscurè radiatá, densè concentricè costatá, costis lamello-recurvis, noduloso-crenatis, costis ad latera undulatis, nodulis evanidis, interstitiis costarum excavatis et crenatis.*

THE MANY-RIBBED VENUS. Shell elliptically ovate, rather ventricose, thick, fulvous-drab, obscurely rayed with flesh-brown, densely concentrically ribbed, ribs lamellarly recurved, nodulously crenated, ribs waved at the sides, nodules fading away, interstices between the ribs excavated and crenated.

SOWERBY, Pro. Zool. Soc. 1835, p. 22.

Hab. Bay of Panama (in coarse sand at low water); Cuming.

I doubt if this is anything more than a variety of *V. Listeri*, in which the ribs are more tumidly thickened and recurved.



VENUS.

PLATE IV.

Species 10. (Mus. Cuming.)

VENUS PUPEREA. *Ven. testā orbiculari-cordatā, ventri-
cosā, crassā, albida, griseo radiatā, ferrugineo-fusco
præcipue ad latū posticum tinctā et maculatā, radi-
atim densē obtusè costatā, concentricè laminatā, lamine
subdistantibus, sēpe irregularibus, latere postico sub-
flexuoso-truncato, antico brevissimo.*

THE VENTRICOSE VENUS. Shell orbicularly heart-shaped, ventricose, thick, whitish rayed with grey, spotted and stained, especially on the posterior side, with rust-brown; radiately closely obtusely ribbed, concentrically laminated, laminae rather distant, often irregular, posterior side slightly flexuously truncated, anterior very short.

LINNAEUS, Mantissa, p. 545.

Hab. Philippine Islands; Cuming.

The rust-stained colouring of this well-known species is not an unimportant character.

Species 11. (Mus. Cuming.)

VENUS FOVEOLATA. *Ven. testā suborbiculari, compres-
siusculā, fulvescente-albā, ad latera carneo tinctā, con-
centricè densē lamellatā, lamellis rugosis, ralde irregu-
laribus, lunula subamplā, areā ligamenti elongato-
excavatā.*

THE PITTED VENUS. Shell somewhat orbicular, rather compressed, yellowish-white, flesh-tinged at the sides, concentrically densely lamellated, lamelle rough, very irregular, lunule rather large, ligamentary area elongately excavated.

SOWERBY, Thes. Conch. part xiv. p. 730.

Hab. Martinique.

The name of this species is apt to mislead; it does not

refer to any sculpture of small pits, but to a characteristic elongated excavation of the ligamentary area.

Species 12. (Mus. Cuming.)

VENUS TENUILAMELLATA. *Ven. testā oratā, subcunei-
formi, soridē calcareo-albā, concentricè lamellatā,
lamellis subregularibus, recurvinsculis, ad latū anti-
cum corrugatis, latere postico oblongo, subtruncato.*

THE THINLY-LAMELLATED VENUS. Shell ovate, somewhat wedge-shaped, dull chalk-white, concentrically lamellated, lamelle rather regular, a little curved, wrinkled at the anterior side, posterior side oblong, slightly truncated.

SOWERBY, Thes. Conch. part xiv. p. 733.

Hab. Valparaiso (in sandy mud at twenty fathoms); Cuming.

Of a dull chalk-white character, with the ridges rather inclined to form irregular wrinkles on the anterior side.


Species 13. (Mus. Cuming.)

VENUS EXALBIDA. *Ven. testā subelliptico-ovatā, soridē
calcareo-albā, concentricè densē elevato-striatā, striis
rugibus, valde irregularibus, latere postico curvato et
flexuoso-truncato.*


THE WHITE VENUS. Shell somewhat elliptically ovate, dull chalk-white, concentrically densely elevately striated, striae rude, very irregular, posterior side curved and flexuously truncated.

CHEMNZITZ, Conch. Cab. vol. xi. p. 225. pl. 202. f. 1974.
Hab. Strait of Magellan.


A densely rudely striated chalk-white shell, peculiarly curved and flexuously striated down the posterior side.




14.



15.



16.



17.

VENUS.

PLATE V.

Species 14. (Mus. Cuming.)

VENUS LISTERI. *Ven. testā subquadrato-ovatā, ventri-
cosā, fulvescente-albā, carneo-fuscescente obscurè tenti-
ginosā, concentricè laminatā, laminis ad latūs posticum
tenuibus, erectis, ad anticum tumidis, recurvis, laminis
vannibus interstiliisque costellis radiantibus crenulatis,
latere postico subtruncato.*

LISTER'S VENUS. Shell somewhat squarely ovate, ventricose, fulvous-white, obscurely freckled with flesh-brown, concentrically laminated, laminae thin and erect on the posterior side, swollen and recurved on the anterior, all the laminae and interstices crenulated with small radiating ribs, posterior side slightly truncated.

Dosina Listeri, Gray, Analyst, 1838.

Venus Listeri, Sowerby.

Hab. Philippine Islands; Cuming.

The laminae, which from being crowded together on the anterior side are swollen and recurved, are on the posterior side, thin, free, and erect.

Species 15. (Mus. Cuming.)

VENUS CASINA. *Ven. testā subquadrato-orbiculari, cal-
careo-albā, interdum spadiceo-roseo maculatā et bira-
diatā, concentricè costatā et liratā, costis lirisque valde
irregularibus, plus minus plicato-recurrens, ad latūs posticum
erecto-lamellatis.*

THE CHAMBERED VENUS. Shell somewhat squarely orbicular, chalk-white, sometimes spotted and two-rayed with fawn-rose, concentrically ribbed and ridged, ribs and ridges very irregular, more or less pliately recurved, cretely lamellated at the posterior side.

Venus casina, Linnaeus, Syst. Nat. p. 1130.

Pectunculus membranaceus, Da Costa.

Venus reflexa, Montagu.

Venus lactea, Donovan.

Venus discina, Lamarek.

Venus Rustensis, Payraudeau.

Cytherea reflexa, Couch.

Hab. European Seas.

The British specimens of this well-known species are rarely coloured, excepting in an early stage of growth. They are mostly a dull stone-white, with the ribs and ridges more or less reflected.

Species 16. (Mus. Cuming.)

VENUS ROSALINA. *Ven. testā subtrigono-ovatā, gibbosin-
culā, fulcescente-albā, spadiceo-roseo tinctā et radiatim
promiscue grandimaculatā, concentricè costatā, costis
latinsculpis, solidis, plano-obtusis, ad extremitatem pos-
ticam plicato-anrichulatis.*

THE ROSY VENUS. Shell somewhat triangularly ovate, rather gibbous, fulvous-white, tinged and radiately promiscuously blotched with fawn-rose, concentrically ribbed, ribs rather broad, solid, flatly obtuse, pliately auriculated at the posterior end.

RANG, Guérin, Mag. de Zool. 1834, class v. pl. 42.

Hab. Island of Gorce, West Africa.

This shell is not of the same quadrate form as *V. casina*, and the colouring and sculpture are of a distinct character.

Species 17. (Mus. Cuming.)


VENUS MAGNIFICA. *Ven. testā globoso-cordatā, crussā,
ponderosā, fulvescente-albā, ferrugineo-fusco tinctā et
circa umbones maculatā, nigricante-violaceo concentricè
fasciatā, radiatim confertim linearis-sulcatā, concentricè
liratā, liris distantibus, vix laminatis, ad latūs anticum
oblusè subfimbriatis.*

THE MAGNIFICENT VENUS. Shell globosely heart-shaped, thick, heavy, fulvous-white, stained and spotted about the umbones with rust-brown, concentrically banded with dark-violet, radiately closely linearly grooved, concentrically ridged, ridges distant, scarcely laminated, slightly obtusely fimbriated on the anterior side.


SOWERBY, Thes. Conch. part xiv. p. 704. pl. 153. f. 5.

Hab. Island of Tieao, Philippines; Cuming.


A globose violet-banded shell, quite distinct from any form of *V. puerpera*.




18.



19 a



19 b



20

V E N U S.

PLATE VI.

Species 18. (Mus. Cuming.)

VENUS LACERATA. *Ven. testā subquadrato-ovatā, valde inaequilateralē, ventricosā, lacteā, violaceo-fusco obscurē interruptē radiatā, liris parvis obtusis concentricis et radiantibus undique fimbriato-decussatā.*

THE TORN VENUS. Shell somewhat squarely ovate, very inequilateral, ventricose, cream-white, obscurely interruptedly rayed with violet-brown, fimbriately decussated throughout with concentric and radiating small obtuse ridges.

HANLEY, Pro. Zool. Soc. 1844, p. 161.

Hab. Island of Zebu, Philippines; Cuming.

Closely allied to *V. puerpera*, lighter and more delicate in colour and sculpture, elaborately decussated.

Species 19. (Fig. a, b, Mus. Cuming.)

VENUS ASPERRIMA. *Ven. testā ovatā, depressiusculā, superne subtrigonā, calcareo-albā, violaceo-fusco interdum radiatim maculatā, radiatim acutē multiliratā, lirā minutā in intersticiis, liris radiantibus liris tenuibus concentricis elaborate clathratis.*

THE ROUGH VENUS. Shell ovate, rather depressed, somewhat triangular at the upper part, chalk-white, sometimes radiately blotched with violet-brown, radiately

sharply many-ridged, with a minute ridge in the interstices, radiating ridges elaborately latticed with fine concentric ridges.

SOWERBY, Pro. Zool. Soc. 1835 p. 41.

Hab. Guacamayo, Central America (in sandy mud, at a depth of thirteen fathoms); Cuming.

A form of *V. pectorina*, in which the shell is of lighter substance, broader and more depressed, with the sculpture more elaborately and definitely latticed.

Species 20. (Mus. Cuming.)

VENUS LAQUEATA. *Ven. testā subgloboso-ovatā, ventricosā, fulvo-spadiceā, violaceo-fusco pallidē radiatō, concentricē valde laminatā, laminis erectis, undique arcuatae laqueatis, intersticiis subtiliter radiatim liratis, liris planis, duplicitis.*

THE FLUTED VENUS. Shell somewhat globosely ovate, ventricose, fulvous-drab, faintly rayed with violet-brown, concentrically strongly laminated, laminae erect, arcuately fluted throughout, interstices finely radiately ridged, ridges flat, duplicate.

SOWERBY, Thes. Conch. part xiv. p. 706. pl. 153. f. 15.

Hab. Swan River, Australia; Jukes.

A solid ventricose shell, crowded throughout with arcuately fluted concentric laminae.

April, 1863.



21



22a



22b.



23

VENUS.

PLATE VII.

Species 21. (Mus. Cuming.)

VENUS LENTICULARIS. *Ven. testā quadrato-orbiculari, valde inaequilaterali, crassā, depressā, sordidè calcareo-albā, concentricè densè ruditè striatā, striis irregularibus, latere postico subflexuoso.*

THE LENS-SHAPED VENUS. Shell squarely orbicular, very incquilateral, thick, depressed, dull chalk-white, densely concentrically rudely striated, striae irregular, posterior side slightly flexuous.

SOWERBY, Pro. Zool. Soc. 1835, p. 42.

Hab. Valparaiso Bay (in coarse sand, at low water); Cuming.

A dull-white heavy shell, almost smooth, with the peculiar depressed quadrately orbicular form of *Artemis*.

Species 22. (Fig. *a, b*, Mus. Cuming.)

VENUS DISCORS. *Ven. testā obliquè ovatā, depressiusculā, albida, violaceo-fusco plus minus tinctā, radiatim profusè plano-tiratis, liris posticè angulatis, liris anticis decussatim crenatis.*

THE DIFFERING VENUS. Shell obliquely ovate, rather depressed, whitish, more or less stained with violet-brown, radiately profusely flatly ribbed, ridges angled along the posterior edge, anterior ridges decussately crenated.

SOWERBY, Pro. Zool. Soc. 1835, p. 42.

Tapes discors, Sowerby Junr.

Hab. St. Elena and Guacamayo, Central America (in sandy mud at six to nine fathoms); Cuming.

The concentric ridges by which the anterior radiating ridges are crenated cease rather abruptly at a distance, from the interior extremity, of rather more than a third of the shell.

Species 23. (Mus. Cuming.)

VENUS RUGOSA. *Ven. testā subgloboso-cordatā, ventricosā, fulvo-spadiceā, roseo-fuscescente obscurè tinctā et maculatā, concentricè undique validè costatā, costis regularibus, laminato-recurvis, interstitiis excavatis, crispato-indentatis et striatis.*

THE WRINKLED VENUS. Shell somewhat globosely heart-shaped, ventricose, fulvous-drab, obscurely tinged and spotted with rose-brown, strongly concentrically ribbed throughout, ribs regular, laminately recurved, interstices excavated, crisply indented and striated.

GMELIN, Syst. Nat. p. 3276.



Venus cincta, Chemnitz.

Venus rigida, Dillwyn.

Hab. West Indies.

A globose heart-shaped shell, ribbed with great regularity, the ribs being smooth, laminately curved over towards the umboes.

April, 1863.



VENUS.

PLATE VIII.

Species 24. (Mus. Cuming.)

VENUS SOWERBYI. *Ven. testá ovatá, ventricosá, crassá, flavescente-albá, rosaceo-fusco interrupté radiatú, concentricè irregulariter laminatá, laminis fimbriato-crenatis, interstiliis elaboratè radiatim liratis.*

SOWERBY'S VENUS. Shell ovate, ventricose, thick, yellowish-white, interruptedly rayed with rose-brown, concentrically irregularly laminated, laminae fimbriately crenulated, interstices elaborately radiately ridged.

DESHAYES, Pro. Zool. Soc. 1853, p. 2.

Hab. Philippine Islands; Cuming.

A very solid ventricose shell, in which the concentric laminae are obtusely fimbriately crenulated.

Species 25. (Mus. Cuming.)

VENUS PECTORINA. *Ven. testá ovato-cordatá, supernè subtrigoná, fulvo-ferruginea, violaceo-fusco plus minus posticè maculatá, radiatim multiliratá, lira minutá interveniente, liris tenuibus concentricis elaboratè clathratá.*

THE HEART-SHAPED VENUS. Shell ovately heart-shaped, slightly triangular at the upper part, fulvous-rust, more or less blotched on the posterior side with violet-brown, radiately many-ridged, with a minute ridge intervening, elaborately latticed with fine concentric ridges.

LAMARCK, Anim. sans vert. p. 344.

Venus cardioidea, Lamarek.

Hab. Central America.

Of rather a cockle-like form, tinged with a fulvous-rust colour and elaborately cancellated.

Species 26. (Mus. Cuming.)

VENUS CINGULATA. *Ven. testá trigono-ovatá, fulvescente, lineis acutè undulatis fusco picta, areá ligamenti fusco conspicue strigata, concentricè costatá, costis primis laminatis et subtiliter crenulatis, cæteris incrassatis, obtusis.*

THE BELTED VENUS. Shell triangularly ovate, fulvous, painted with zigzag brown lines, ligamentary area conspicuously striped with brown, concentrically ribbed, the first ribs laminated and finely crenulated, the rest thickened and obtuse.

LAMARCK, Anni. sans vert. p. 345.

Venus pulicaria, Broderip.

Hab. West Columbia; Cuming.

The concentric ribs, which in the early growth of the shell are thin and laminated, and crenulated, become gradually more and more thickened and obtuse.

Species 27. (Mus. Cuming.)


VENUS RESTICULATA. *Ven. testá subquadrato-cordatá, ventricosá, crassá, albida, circa umbones ferrugineo tintata et fusco maculata, concentricè densè laminata, laminis tumido-crenatis et subrecurvis, interstiliis radiatim obtuse crenatis.*

THE CORDED VENUS. Shell somewhat squarely heart-shaped, ventricose, thick, whitish, rust-tinged and spotted with brown about the umboes, concentrically densely laminated, laminae tunnily crenated and a little recurved, interstices radiately obtusely ridged.


SOWERBY, Thes. Conch. part xiv. p. 706.

Hab. Philippine and Society Islands and Madagascar.

A solid globosely cordate shell, sculptured with stout concentric laminae, having almost the consistency of ribs, crenated and fluted throughout in a rather callous manner.



28



29



30.



31

V E N U S.

PLATE IX.

Species 28. (Mus. Cuming.)

VENUS COR. *Ven. testā trapeziformi-globosā, tenuiculā, inflatā, sordidē albā, plus minus ferrugineo tinctā, concentricè liratā, liris obtusis, valde irregularibus, hic illie undulatis.*

THE HEART VENUS. Shell trapeziformly globose, rather thin, inflated, dull-white, more or less rust-stained, concentrically ridged, ridges obtuse, numerous, very irregular, here and there undulated.

SOWERBY, Thes. Conch. part xiv. p. 727. pl. 160. f. 184.
Hab. Kurachee, mouth of the Indus.

A light trapeziformly globose shell, sculptured with concentric ridges, which are very irregular and undulating.

Species 29. (Mus. Cuming.)

VENUS DOMBEIJI. *Ven. testā orbiculari-ovatā, subdepressā, sordidē fuscescente-albā, fuscescente radiatim undulatā et lentiginosā, undique obtusè crenato-decussatā.*

DOMBEY'S VENUS. Shell orbicularily ovate, rather depressed, dull brownish-white, radiately waved and freckled with light-brown, everywhere obtusely crenately decussated.

LAMARCK, Anim. sans vert. p. 346.

Venus Chilensis, Sowerby.

Tapes Dombeii, Sowerby, Jun.

Hab. Valparaiso Bay; Cuming.

This shell has very much the form and sculpture, but not the whiteness, of a *Lucina*.

Species 30. (Mus. Cuming.)

VENUS COSTELLATA. *Ven. testā subquadrato-ovatā, sordidē albā, ferrugineo tinctā, radiatim tenuicostatā, costis planis, confertis, liris obtusis concentricis irregularibus decussatis.*

THE FINELY-RIBBED VENUS. Shell somewhat squarely ovate, dull-white, rust-stained, radiately finely ribbed, ribs flat, close-set, decussated with obtuse irregular concentric ridges.

SOWERBY, Pro. Zool. Soc. 1835, p. 42.
Dosina antiqua, Gray.

Hab. Chili and Peru (in coarse sand from six to fifteen fathoms); Cuming.

A dull stone-coloured shell very rudely irregularly decussated.

Species 9. (Mus. Cuming.)


VENUS CRISPATA. *Ven. testā oratā, crassiusculā, albida, rufo-fusco undatā et radiatim fasciatā, concentricè crenato-laminatā, laminis tumidiusculis, confertis, interstitiis radiatim liratis, liris subdistantibus.*

THE CRISPED VENUS. Shell ovate, rather thick, whitish, waved and radiately banded with red-brown, concentrically crenately laminated, laminae rather swollen, close-set, interstices radiately ridged, ridges rather distant.


DESHAYES, Pro. Zool. Soc. 1853, p. 2.

Hab. —?


Of a peculiar regularly convex oval form, with the concentric laminae close-set and very crowded on the anterior side.




32a.



32b



33.



34.

V E N U S.

PLATE X.

Species 32. (Fig. *a*, *b*, Mus. Cuming.)

VENUS CHEMNITZII. *Ven. testā ovatā, fulvescente-albā, rufo-fusco acutē undatā et radiatim fasciatā, concentricē densē laminatā, laminis tenuibus, fimbriato-laqueatis, interstitiis radiatim liratis, liris subdistantibus.*

CHEMNITZ'S VENUS. Shell ovate, yellowish-white, sharply waved and radiately banded with red-brown, concentrically densely laminated, laminæ thin, fimbriately fluted, interstices radiately ridged, ridges rather distant.

HANLEY, Pro. Zool. Soc. 1844, p. 160.

Hab. Island of Ticao, Philippines; Cuming.

In this species the concentric lamellæ are extremely thin and delicately fimbriately fluted. The colour is disposed in sharply zigzag waves.

Species 33. (Mus. Cuming.)

VENUS CRENULATA. *Ven. testā trigono-ovatā, subrentricosā, albida, rufo-fusco profusè maculatā et undatā, concentricē laminato-liratā, liris confertis, elevatisculnis, crenulatis, ad latus anticū subrecurvis, ad posticum tenuibus, erectis, laqueatis, interstitiis radiatim crispato-striatis.*

THE CRENULATED VENUS. Shell triangularly ovate, rather ventricose, whitish, profusely spotted and waved with red-brown, concentrically laminately

ridged, ridges close-set, rather elevated, crenulated, a little recurved on the anterior side, on the posterior thin, erect, fluted, interstices radiately crisplately striated.

CHEMNITZ, Coneh. Cab. vol. vi. p. 370. pl. 36. f. 385.

Venus crenata, Gmelin.

Hab. West Indies.

A very elaborately sculptured shell, of a triangularly ovate form, profusely wave-painted with red-brown.

Species 34. (Mus. Cuming.)

VENUS RETICULATA. *Ven. testā quadra-to-ovatā, suboblongā, albā vel fuscescente, rufo-fusco linearī-undatā et maculatā, undique densē crenato-reticulatā, latere pos-tico supernē subalato.*


THE RETICULATED VENUS. Shell squarely ovate, rather oblong, white or brownish, spotted and linearly waved with red-brown, everywhere closely crenately reticulated, posterior side slightly winged at the upper part.

LINNÆUS, Mus. Ulr. p. 503.


Venus corbis, Lamarck.

Hab. Philippine Islands, etc.

A constant peculiarity of this well-known species consists in the hinge teeth being of a bright orange-red colour.




35



36^a



36^b



37

VENUS.

PLATE XI.

Species 35. (Mus. Cuming.)

VENUS CALIFORNIENSIS. *Ven. testá orbiculari-cordatá, subobliquá, valde inaequilaterali, pallidè fulvo-cinereá, concentricè erecto-laminatá, laminis versus marginem obtusis, confertis, intersticiis longitudinaliter liratis, liris irregularibus, transversim corrugato-striatis, areá ligamenti ampliter concavo-excavatá.*

THE CALIFORNIAN VENUS. Shell orbicularly heart-shaped, rather oblique, very inequilateral, pale fulvous-ash, concentrically erectly laminated, laminae obtuse and crowded towards the margin, interstices longitudinally ridged, ridges irregular, transversely wrinkle-striated, ligamentary area largely concavely excavated.

BRODERIP, Pro. Zool. Soc. 1835, p. 43.

Venus leucodon, Sowerby.

Hab. Guaymas, Gulf of California (in sandy mud at low water); Cuming.

This fine species, it will be seen, is of remarkably cor- date growth, but at the same time orbicularly depressed. The ligamentary area is unusually large and rather sharply concavely excavated.

Species 36. (Fig. a, b, Mus. Cuming.)

VENUS AMATHUSIA. *Ven. testá subtrigono-ovatá, fulvo-spadiceá, concentricè laminatá, laminis nunc confertis, nunc distantibus, crenatis et erectè muricato-squamatis, intersticiis longitudinaliter liratis, liris binis, latere postico subflexuoso, lunatá et areá ligamenti vio- laceo-tinctis.*

THE AMATHUSIA VENUS. Shell somewhat triangularly

ovate, fulvous-drab, concentrically laminated, laminae sometimes close-set, sometimes distant, crenated and erectly prickly-sealed, interstices longitudinally ridged, ridges in pairs, posterior side rather flexuous, lunule and ligamentary area tinged with violet.

PHILIPPI, Abbild. und Besch. Conch. pl. II. f. 4.

Hab. Gulf of California.

Chiefly distinguished from *V. Guidia* by its more tri- angularly ovate form, and by the more flexuously produced growth of the posterior extremity. The radiating ridges in the interstices between the laminae are moreover more distinctly set in pairs without any intermediate raised stria or smaller ridge.

Species 37. (Mus. Cuming.)


VENUS GNIDIA. *Ven. testá subtrigono-ovalá, ventrico- siusculá, fulvescente-spadiceá, concentricè laminatá, laminis crenatis et erectè muricato-squamatis, intersti- ciis radiatim liratis, liris vix binis, lira minutá in- terveniente.*

THE GNIDIA VENUS. Shell somewhat triangularly ovate, rather ventricose, fulvous drab, concentrically laminated, laminae crenated and erectly prickly-sealed, interstices radiately ridged, ridges scarcely in pairs, with a small ridge intervening.


BRODERIP and SOWERBY, Zool. Journ. vol. iv. p. 364.

Hab. San Blas, Mexico.


The scales with which the concentric laminae of this species are so elegantly festooned, are more prickly pointed in the smaller varieties of the species.




38.




39a




39b




40a



41



42



40b

VENUS.

PLATE XII.

Species 38. (Mus. Cuming.)

VENUS AFFINIS. *Ven. testā subtrigono-ovatā, crassā, albida, rufo-brunneo interrupte radiatā et maculatā, concentricē taminato-costatā, costis conspicuē reflexis, ad latus posticum fornicatis, deinde interrupte sulcatis, areā ligamenti latā, planā, fusco minutē linearī-lenticīginosā.*

THE ALLIED VENUS. Shell somewhat triangularly ovate, thick, whitish, interruptedly blotched and rayed with red-brown, concentrically laminately ribbed, ribs conspicuously reflected, vaulted, then interruptedly grooved on the posterior side, ligamentary area broad, flat, minutely linearly freckled with brown.

SOWERBY, Thes. Conch. part xiv. p. 720. pl. 155. f. 62.

Hab. Senegal.

Allied to *V. paphia*, but of a more oblong form, with somewhat of the colouring of *V. rosalina*.

Species 39. (Fig. a, b, Mus. Cuming.)

VENUS LAMARCKII. *Ven. testā oratā, arcuatē subtrigonā, albida, roseo-fusco maculatā et strigatā, concentricē fimbriato-costatā, fimbriis eleganter plicatis et recurvis, radiatim liratā, latere postico subattenuato; intus rosacea.*

LAMARCK'S VENUS. Shell ovate, areuately subtriangular, whitish, blotched and streaked with rose-brown, concentrically fimbriately ribbed, frills elegantly puckered and recurved, radiately ridged, posterior side slightly attenuated; tinged with rose within.

GRAY, Sowerby, Thes. Conch. part xiv. p. 707. pl. 153. f. 20, 21.

Venus cancellata, var., Chemnitz.

Hab. China. Wide Bay, East Australia; Strange.

This well-known China species was figured by Chemnitz as a variety of *V. cancellata*, and a figure of it in the 'Encyclopédie Méthodique' was erroneously referred to by Lamarck in illustration of his *V. subrostrata*, which is a West Indian species. The sculpture is most elaborately fretted. In the interior the valves are tinged towards the umboes with rose.

Species 40. (Fig. a, b, Mus. Cuming.)

VENUS VERRUCOSA. *Ven. testā subcordato-ovatā, solidusculā, fuscescente-albā, rosaceo-fusco plus minus obscurè tinctā et linearī-undatā, concentricē costatā et tiratā, costis lirisque confertis, irregularibus, tuberculato-nodatis, areā ligamenti plano-excavatā, vividē fusco strigatā.*

THE WARTY VENUS. Shell somewhat cordately ovate, rather solid, brownish-white, more or less obscurely stained and linearly waved with rose-brown, concentrically ribbed and ridged, ribs and ridges close-set, irregular, tubercularly noduled, ligamentary area flatly excavated, streaked with deep brown.

Linnaeus, Syst. Nat. 12th edit. p. 1130.

Venus erycina, Pennant.

Pectunculus strigosus, Da Costa.

Venus cancellata, Donovan.

Venus Lemanii, Payraudeau.

Hab. South European and North African seas, including Britain and the Channel Islands.

In this well-known species the concentric ribs and ridges are broken up, more especially on the posterior side, into warty nodules.

Species 41. (Mus. Cuming.)

VENUS KENNERLEYI. *Ven. testā oblongo-ovali, subquadratā, sordidē opaco-albā, concentricē plicato-costatā, costis confertis, subrecurvis, subirregularibus, lunulā et areā ligamenti angustis.*

KENNERLEY'S VENUS. Shell oblong oval, somewhat square, dull opake white, concentrically plicately ribbed, ribs close-set, a little recurved, rather irregular, lunule and ligamentary area narrow.

Carpenter, MS. in Mus. Cuming.

Hab. —?

A very characteristic chalk-white shell, strongly plicately concentrically ribbed, the ribs being very crowded and irregular.

Species 42. (Mus. Cuming.)

VENUS CRASSICOSTATA. *Ven. testā subquadrato-ovatā, calcareo-albā, ferrugineo tinclā, radiatim costatā, costis subplanatis, irregularibus, confertis, costis anticus nodoso-crenatis, posticis ruditè corrugatis.*

THE THICK-RIBBED VENUS. Shell somewhat squarely ovate, chalk-white, tinged with rust, radiately ribbed, ribs rather flattened, irregular, close-set,

anterior ribs nodosely crenated, posterior rudely wrinkled.


QUOY and GAIMARD, Voy. de l'Astrolabe, Moll. pl. 84.
f. 1, 2.

Tapes crassicostata, Sowerby.
Hab. Otago, New Zealand.

An opake rust-tinged chalk-white shell, having a semi-fossil appearance.



44




43




45.




46



47



48



49

VENUS.

PLATE XIII.

Species 43. (Mus. Cuming.)

VENUS SUGILLATA. *Ven. testá subtrigono-ovatá, fulvescente-griseá, purpureo sparsim undato-lineatá, umbonibus vividè purpureis, liris tenuibus, undique subacutè clathratá, lunulá amplá, areá ligamenti augustá, ambabus intèse castaneis.*

THE BRUISED VENUS. Shell somewhat triangularly ovate, fulvous-grey, sparingly wave-lineated with purple, umboes deep purple, everywhere rather sharply latticed with fine ridges, lunule large, ligamentary area narrow, both dark-chestnut.

Hab. California.

I cannot identify this species with any hitherto described. Its shining purple umboes, finely latticed sculpture, and dark-stained lunule and ligamentary area are quite peculiar.

Species 44. (Mus. Cuming.)

VENUS SIMILLIMA. *Ven. testá subcordato-ovatá, depresso-sculpulá, sordidè fuscescente-albá, concentricè laminato-liratá, liris subconfertis, plicatis, subreflexis, posticè appressis, radiatim tenuiliratis, liris duplicitibus.*

THE VERY LIKE VENUS. Shell somewhat cordately ovate, rather depressed, dull brownish-white, concentrically laminately ridged, ridges rather close-set, puckered, a little reflected, posteriorly appressed, radiately finely ridged, ridges duplicate.

SOWERBY, Thes. Conch. part xiv. p. 708. pl. 153. f. 17, 18.

Hab. San Diego, California.

Resembling the beautiful *V. compta* in detail of sculpture, but of a distinct form, without any of its characteristic colouring.

Fig. 45. (Mus. Cuming.)

The shell here represented was drawn as a new species; but it appears to be an unusually gibbous form of *V. casina*, with the concentric sculpture more slenderly laminated and crowded.

Fig. 46. (Mus. Cuming.)

A very distinct variety, from the Gulf of California, of *V. crenulata*, of which a typical specimen of the West Indian form is represented at Plate X. f. 33. In the Californian variety the shell is more globose, and beautifully coloured in the interior with purple-rose.

Species 47. (Mus. Cuming.)

VENUS GIBBOSULA. *Ven. testá pectinato-orbiculari, albidá, violaceo-griseo tinctá et obscurè fasciatá, undique obtuse corrugato-reticulatá, præcipue ad latus anticum, latere postico radiatim sulcato-striato, lunulá et areá ligamenti subindistinctis.*

THE GIBBOS VENUS. Shell pectinately orbicular, whitish, stained and obscurely banded with violet-grey, everywhere obtusely wrinkle-reticulated, especially on the anterior side, posterior side radiately groove-striated, lunule and ligamentary area rather indistinct.

DESHAYES, MS. in Mus. Cuming.

Hab. ——?

An interesting, delicately tinted species, of rather solid growth, sculptured with a network of obtuse wrinkled ridges.

Species 48. (Mus. Cuming.)

VENUS COMPTA. *Ven. testá abbreviato-cordatá, subtriangularis, depressá, crassí, sordidè fulvo-albá, rinfo-castaneo radiatim linearis-trigatá, concentricè laminato-costatá, costis distantibus, areá ligamenti amplá, acutè excavatá.*

THE NEAT VENUS. Shell shortly cordate, somewhat triangular, depressed, thick, dull fulvous-white, radiately linearly streaked with reddish-chestnut, concentrically laminately ribbed, ribs distant, ligamentary area large, sharply excavated.

BRODERIP, Pro. Zool. Soc. 1838, p. 43.

Hab. Bay of Sechura, Peru (in coarse sand and mud at the depth of seven fathoms); Cuming.

VENUS.—PLATE XIII.

The radiated linear colouring of this species is a characteristic feature.

Species 49. (Mus. Cuming.)

VENUS NUTTALLI. *Ven. testā pectinato-cuneiformi, crassā, albidā, versus umbones ferrugineo tinctā, radiatim nitidē liratā, liris medianis angustis, tenuibus, concentricē liratā, liris laminato-reflexis, versus marginem anticum numerosis, valde irregularibus, lunulā violaceā, areā ligamenti latē plano-excavatā.*


NUTTALL'S VENUS. Shell pectinately wedge-shaped,

thick, whitish, stained towards the umboes with rust, radiately neatly ridged, middle ridges narrow, thin, concentrically ridged, ridges laminae reflected, crowded towards the anterior margin, very irregular, lunule violet, ligamentary area broadly flatly excavated.

CONRAD, Acad. Nat. Sci. Philadelphia, vol. vii. p. 250.
pl. 19. f. 15.

Hab. California.

As the neat radiating ridges on approaching the margin become more distinct, the concentric anterior laminar ridges are more crowded.



VENUS.

PLATE XIV.

Species 50. (Mus. Cuming.)

VENUS EBURNEA. *Ven. testā suboblongo-ovatā, eburnea, ad lunulam pallidē roseo-fuscescente, concentricè corrugato-liratis, liris ad latus anticū irregularibus, interruptis.*

THE IVORY VENUS. Shell somewhat oblong-ovate, ivory-white, pale rose-brown at the lunule, concentrically wrinkle-ridged, ridges on the anterior side irregular, interrupted.

Hab. —?

A neatly-defined ivory-white species, allied to *V. oblonga*, in which the ridges cease to be concentric on the anterior side, but form irregular interrupted wrinkles.

Species 51. (Mus. Cuming.)

VENUS MUNDULUS. *Ven. testā subquadrato-ovatā, tenuiculā, fulvescente-albā, violaceo-cinereo interrupte flammatā, undique minutè crenato-reticulatā, lunulā sub-indistinctā, areā ligamenti angustā.*

THE TRIM VENUS. Shell somewhat squarely ovate, rather thin, fulvous-white, interruptedly flamed with violet-ash, everywhere minutely crenately reticulated, lunule rather indistinct, ligamentary area narrow.

Hab. —?

A neatly-sculptured shell of rather light substance, interruptedly flamed throughout with violet-ash.

Species 52. (Mus. Cuming.)

VENUS INTERSECTA. *Ven. testā pectinato-ovali, tenui, subventricosā, albida, violaceo et pallidē ferrugineo plus minus radiatim tinctā, liris subtilibus filosis undique reticulatum decussatā, lunulā et areā ligamenti violaceo-fuscis.*

THE INTERSECTED VENUS. Shell pectinately oval, thin, rather ventricose, whitish, more or less radiately tinged with violet and pale-rust, everywhere reticulately decussated with fine thread-like ridges, lunule and ligamentary area violet-brown.

SOWERBY, Pro. Zool. Soc. Thes. Conch. p. 714. pl. 155. f. 59, 60.

Hab. Puerto Puerto, Central America; Cuming.

A delicately-coloured shell of thin substance, very elaborately minutely sculptured throughout.

Species 53. (Mus. Cuming.)

VENUS MONILIFERA. *Ven. testā quadrato-ovatā, subcompressā, fulvescente-albā, purpureo-fusco profusè undatā et maculatā, undique liris acutè reticulatā, latere postico compressè producto.*

THE NECKLACE VENUS. Shell squarely ovate, rather compressed, fulvous-white, profusely waved and blotched with purple-brown, everywhere sharply reticulated with ridges, posterior side compressly reduced.

SOWERBY, Thes. Conch. part xiv. p. 707. pl. 153. f. 19.

Hab. Island of Mindoro, Philippines; Cuming.

A characteristic form of *V. reticulata*, in which the shell is compressly squarely produced, with the painting much waved, especially on the posterior side.

Species 54. (Fig. a, b, Mus. Cuming.)

VENUS SUBROSTRATA. *Ven. testā cordato-subtrigonā, positiè declivi-truncatā, crassā, albā, fusco reticulato-maculatā, concentricè fimbriato-liratā, liris numerosis, brevibus, radiatim costellatā, costellis versus marginem obsolete, lunulā intensè fuscā, areā ligamenti acutè excavatā, fusco strigatā.*

THE SLIGHTLY-BEAKED VENUS. Shell cordately subtriangular, posteriorly slopingly truncated, thick, white, reticulately blotched with brown, concentrically fimbriately ridged, ridges numerous, short, radiately finely ribbed, ribs soon becoming obsolete towards the margin, lunule dark-brown, ligamentary area sharply excavated, striped with brown.

LAMARCK, Anim. sans vert. vol. vi. p. 343.

Venus neglecta, Sowerby (in Beechey's Voyage, p. 151. pl. 41. f. 8).

Hab. Mazatlan and West Indies.

In consequence of Lamarck having referred for the illustration of this species to a figure in the 'Encyclopédie

VENUS.—PLATE XIV.

Méthodique,' which represents a well-known China species,
V. Lamarekii, it was lost sight of, until Sowerby named
it in the 'Zoology of Beechey's Voyage' *V. neglecta*.


Species 55. (Mus. Cuming.)

VENUS ALBINA. *Ven. testa suboblongo-ovalis, albâ, concen-*
tricè densè scabroso-striatâ, striis irregularibus, lunula-
fusca.

THE WHITE VENUS. Shell rather oblong-ovate, white, con-
centrically densely sebrously striated, striae irregular,
lunule dark-brown.

SOWERBY, Thes. Conch. part xiv. p. 732. pl. 161. f. 201.
Hab. China.

A peculiarly soft white shell, in which the concentric
sculpture is scarcely more than sebrous striae.



V E N U S.

PLATE XV.

Species 56. (Mus. Cuming.)

VENUS LYRA. *Ven. testá orbiculari-ovatá, subventricosá, fulvescente-albá, rufescente-fusco undatá et maculatá, concentricè liratá, liris numerosis, subcallosis, lævibus, subreflexis.*

THE HARP VENUS. Shell orbicularly ovate, rather ventricose, fulvous-white, waved and spotted with red-brown, concentrically ridged, ridges numerous, rather callous, smooth, a little reflected.

HANLEY, Pro. Zool. Soc. 1844, p. 161. Recent Shells, App. 360. pl. 16. f. 21.

Hab. Gulf of Guinea.

I give this species, but doubt exceedingly whether it is not the same as *V. crenulata*. It is an established fact in the geographical distribution of mollusks that West Africa possesses many species common to the West Indies and Mazatlan.

Species 57. (Mus. Cuming.)

VENUS NODOSA. *Ven. testá globosá, sordidè albá aut fulvescente, concentricè costatá, costis subdistantibus, obtusè tumidis, verrucoso-tuberculatis, lunulá rufo-fusca, areá ligamenti rufo-fusco latè strigatá.*

THE NODULED VENUS. Shell globose, dull-white or yellowish, concentrically ribbed, ribs rather distant, obtusely swollen, wart-tubercled, lunule red-brown, ligamentary area broadly striped with red-brown.

DUNKER, Moll. Guin. p. 57.

Venus tuberosa, Deshayes.

Hab. St. Vincent, West Africa.

This apparently characteristic species is probably a strongly warted form of the European *V. verrucosa*.

Species 58. (Mus. Cuming.)

VENUS PILULA. *Ven. testá orbiculari-ovalá, tumido-ventricosá, fulvescente-albá, carneo-fusco promiscuè tinctá, concentricè liratá, liris subobtusis, interstitiis concavis, concentricè elevato-striatis, lunulá parvá, areá ligamenti rufo-fusco strigatá.*

THE LITTLE-BALL VENUS. Shell orbicularly ovate, tumidly ventricose, fulvous-white, promiscuously stained

with flesh-brown, concentrically ridged, ridges rather obtuse, interstices concave, sculptured with raised striae, lunule small, ligamentary area striped with red-brown.

Hab. —?

Closely allied to Dr. Gould's *V. toreuma*, but the concentric ridges are not warted, and the lunule is smaller.

Species 59. (Mus. Cuming.)

VENUS STUTCHBURII. *Ven. testá obliquè cordatá, posticè subproductá, albidá, ferrugineo-tinctá posticè, præsertim intus, violaceo tinctá, radiatim tenuiliratá, concentricè corrugato-striatá, lunulá et areá ligamenti subindistinctis.*

STUTCHBURY'S VENUS. Shell obliquely cordate, rather produced posteriorly, whitish, tinged with rust, posteriorly tinged with violet, especially within, radiately finely ridged, concentrically corrugately striated, lunule and ligamentary area rather indistinct.

GRAY, Wood, Ind. Test. Supp. pl. 2. f. 4.

Hab. Sandwich Islands.

A shell of solid growth, tinged within on the posterior side with deep violet. It comes very near to the Californian *V. callosa*, Conrad, of which specimens have been found also at the Sandwich Islands.

Species 60. (Mus. Cuming.)

VENUS MUSCARIA. *Ven. testá pectinato-oralá, subcompressá, fulvescente, cinereo-violaceo radiatim aspersa, radiatim crebriliratá, concentricè, densè crenato-lirata, lunulá subindistincta.*

THE FLY-DOTTED VENUS. Shell pectinately ovate, rather compressed, fulvous, radiately sprinkled with ash-violet, closely radiately ridged, concentrically densely crenately ridged, lunule rather indistinct.

Hab. —?

An extremely interesting species, most elaborately sculptured throughout.

Species 61. (Fig. *a*, *b*, Mus. Cuming.)

VENUS COLUMBIENSIS. *Ven. testá pectinato-ovalá, fulvescente, cinereo-griseo radiatim aspersá et maculatá, interdum plus minus cinereo-fuscá, radiatim costatá, costis obtuse planatis, interstitiis subprofundè incisis, costis anticis corrugato-crenatis.*

THE COLUMBIAN VENUS. Shell pectinately ovate, fulvous, radially sprinkled and blotched with ash-grey,

sometimes more or less with ash-brown, radially ribbed, ribs obtusely flattened, interstices rather deeply cut, anterior ribs corrugately crenated.

SOWERBY, Pro. Zool. Soc. 1835, p. 21.

Hab. St. Elena, West Columbia.

This very elegant species varies considerably in its pattern of colouring, but little in form and sculpture.



63




62



64 a.




64 b.




65.



66



67



68



69



70.

V E N U S.

PLATE XVI.

Species 62. (Mus. Cuming.)

VENUS GRANULATA. *Ven. testā pectinato-ovatā, solidiusculā, albā, ferrugineo tinctā, nigricante-cinereo versus marginem reticulato-pictā, radiatim costatā, costis obtusè planatis, posticē angulatis, liris nodulosis concentricis decussatis.*

THE GRANULATED VENUS. Shell pectinately ovate, rather solid, whitish, rust-tinged, reticulately painted towards the margin with dark ash, radiately ribbed, ribs obtusely flattened, posteriorly angled, decussated with concentric nodulous ridges.

GMELIN, Syst. Nat. p. 3277.

Venus entobapta, Jonas.

Hab. West Indies.

A substantially-grained shell, boldly patched here and there with dark colour.

Species 63. (Mus. Cuming.)

VENUS LACUNATUS. *Ven. testā subtrigono-ovatā, fulvescente, roseo tinctā, rufō-fusco obscurè radiatā, concentricè laminato-liratā, liris erectis, extus lucunatis, areā ligamenti rufō-fusco maculatā.*

THE FRETTED VENUS. Shell somewhat triangularly ovate, fulvous, tinged with rose, obscurely rayed with red-brown, concentrically laminately ridged, ridges erect, supported outwardly by fretwork, ligamentary area blotched with red-brown.

Hab. —?

Of a delicate fulvous-brown colour, tinged with rose, most elaborately fretted throughout with punctured lamellæ.

Species 64. (Fig. a, b, Mus. Cuming.)

VENUS TOREUMA. *Ven. testā pectinato-orbiculari, gibbosulā, albidā, rufō-castaneo radiatim trigono-maculatā, concentricè liratā et striatā, interstitiis regulariter concavis, striis elevatis, lateribus verrucoso-corrugatis.*

THE EMBOSSED VENUS. Shell pectinately orbicular, rather gibbous, white, radiately triangularly blotched with reddish-chestnut, concentrically ridged and

striated, interstices regularly concave, with the striae raised, sides wart-wrinkled.

GOULD, Otia Conch. p. 84.

Venus crebrisulca, Sowerby (not of Lamarek).

Venus Jukesii, Deshayes.

Hab. Mangsi Island; Gould.

A prettily-painted globose shell, in which the concentric ridges are wrinkled into warts towards the sides, and ventral margin.

Species 65. (Mus. Cuming.)

VENUS PLUMBEA. *Ven. testā ovatā, valde inaequilaterali, sordidè fulvā, plumbeo-violaceo radiatā et tinctā, radiatim costatā, anticis latiusculis, posticis angustis, obtusè angulatis, omnibus liris concentricis undique crenatis, lunulā et areā ligamenti latiusculis.*

THE LEADEN VENUS. Shell ovate, very inequilateral, dull fulvous, rayed and stained with leaden-violet, radiately ribbed, anterior ribs rather broad, posterior narrow, obtusely angled, all crenated throughout with concentric ridges, lunule and ligamentary area rather broad.

Hab. St. John's, West Indies.

This shell was labelled in Mr. Cuming's collection as a variety of *V. tessellata*, but it is of a different form, less oblong, with a larger lunule, and broader, more excavated, ligamentary area.

Species 66. (Mus. Cuming.)

VENUS TESSELLATA. *Ven. testā oblongo-ovatā, fulvā, plumbeo-violaceo radiatā et tinctā, radiatum densè callosō-liratā, postice costatā, costis planis, obtusè angulatis, lunulā et areā ligamenti angustis.*

THE TESSELLATED VENUS. Shell oblong-ovate, fulvous, rayed and stained with leaden-violet, radiately densely callously ridged, ribbed posteriorly, ribs flat, obtusely angled, lunule and ligamentary area narrow.

ADAMS and REEVE, Moll. Voy. Samarang, p. 79. pl. 22. f. 11.

Tapes tessellata, Sowerby.

Hab. Philippine Islands; Cuming.

Of a characteristic oblong form, with a very short anterior side.

Species 67. (Mus. Cuming.)

VENUS CASINA. *Ven. testā lenticulari, fulvā, ferrugineo-fusco tinctā, concentricē laminato-costatā, costis rubris, subirregularibus, laminā parvā interveniente.*

THE CASINA VENUS. Shell lens-shaped, fulvous, tinged with rust-brown, concentrically lamellately ribbed, ribs rude, rather irregular, with a small lamina in the interstices.

LINNÆUS, Syst. Nat. p. 1130.

Venus reflexa, Montagu.

Venus lactea, Donovan.

Venus discina, Lamarek.

Venus Rusterncii, Payraudeau.

Venus intermedia, Gray.

Venus casinula, Deshayes.

Hab. European Seas.

This well-known species appears to have been named after a town in Campania, on the shores of the Mediterranean.

Species 68. (Mus. Cuming.)

VENUS UNDATELLA. *Ven. testā ovatā, subcompressā, cærulescente-albā, violaceo-fusco lineari-undatā et maculatā, radiatim regulariter liratā, liris versus marginem obsoletis, concentricē liratā, liris versus marginem undatis, confertis et validioribus, areā ligamenti sublatè excavatā, conspicuē strigatā.*

THE WAVED VENUS. Shell ovate, rather compressed, bluish-white, lineally waved and blotched with violet-brown, radiately regularly ridged, ridges obsolete towards the margin, concentrically ridged, ridges waved, close-set and stronger towards the margin, ligamentary area rather broadly excavated, conspicuously striped.

SOWERBY, Pro. Zool. Soc. 1835. p. 22.

Hab. Gulf of California.

A rather pale-marked species, with the sculpture much changing in its development towards the margin.

Species 69. (Mus. Cuming.)

VENUS FUSCO-LINEATA. *Ven. testā oblongo-ovatā, subquadratā, albida, lineis nigrante-violaceis undatis pictā, radiatim costatā, costis obtusè planatis, posticè angulatis, anticis subacutè crenatis.*

THE BROWN-LINED VENUS. Shell oblong-ovate, somewhat square, whitish, painted with dark-violet waved lines, radiately ribbed, ribs obtusely flattened, angled posteriorly, anterior ribs rather sharply crenated.

SOWERBY, Pro. Zool. Soc. 1835. p. 41.

Tapes fusco-lineata, Sowerby, jun.

Hab. Guacomayo, Central America (in sandy mud, at the depth of thirteen fathoms); Cuming.

Crossed throughout with elegantly-defined pencilings of dark violet.

Species 70. (Mus. Cuming.)

VENUS HISTRIONICA. *Ven. testā subquadrato-ovatā, compresiusculā, albā, cinereo-fusco conspicuē maculatā, undique nitidè crenato-reticulatā, lunulā intensè fuscā.*

THE PATCHED VENUS. Shell somewhat squarely ovate, rather depressed, white, conspicuously blotched with ash-brown, neatly crenately reticulated throughout, lunule dark-brown.

SOWERBY, Pro. Zool. Soc. 1835. p. 41.

Hab. Guacomayo, Central America (in sandy mud, at the depth of thirteen fathoms); Cuming.

Very closely allied to *V. granulata*, though more compressed and delicately sculptured.




72




71




73




74



75



76



77

V E N U S.

PLATE XVII.

Species 71. (Mus. Cuming.)

VENUS MACTRACEA. *Ven. testā ovatā, calcareo-albā, concentricē liratā et striatā, liris striisque ratde irregulribus, tunnla excavato-elongatā.*

THE MACTRA-LIKE VENUS. Shell ovate, chalk-white, concentrically ridged and striated, ridges and striae very irregular, lunule excavately lengthened.

BRODERIP, Pro. Zool. Soc. 1833, p. 44.

Hab. Valparaiso (from sandy mud at a depth of twenty fathoms); Cuming.

An opaque chalk-white species, with very much the appearance of a tertiary fossil.

Species 72. (Mus. Cuming.)

VENUS DISCREPANS. *Ven. testā subpectinato-ovatā, fulvescente-albā, concentricē costatā, costis tumidis, rotundatis, confertiusculis, subitus crispato-crenulatis, ferrugineo-fusco punctatis, areā ligamenti latā.*

THE DIFFERING VENUS. Shell subpectinately ovate, fulvous-white, concentrically ribbed, ribs swollen, rounded, rather close-set, crisplately crenulated beneath, dotted with rust-brown, ligamentary area broad.

SOWERBY, Pro. Zool. Soc. 1833, p. 22.

Hab. Islay, Peru; Cuming.

Very like *V. Peruviana*, with the posterior foliations of the concentric ribs decorticated.

Species 73. (Mus. Cuming.)

VENUS APHRODINOIDES. *Ven. testā obliquē cuneiformi, plano-compressā, fulvescente-albā, griseo-punctatā, undique concentricē validē liratā, liris acutis, intersticiis profundē excavatis, latere antico brevissimo, postico elongato producto.*

THE APHRODINE-LIKE VENUS. Shell obliquely wedge-shaped, flatly compressed, fulvous-white, dotted with grey, concentrically strongly ridged throughout, ridges sharp, interstices deeply excavated, anterior side very short, posterior elongately produced.

Hab. St. Vincent's Gulf, South Australia; Augas.

More inequilateral than *V. Aphrodina* or *Peronii*, all of which are very closely allied, with the umbones more sharply angular, and, above all, with the ridges, as Lamarck describes them, more eminent.

Species 74. (Mus. Cuming.)

VENUS EGROTA. *Ven. testā subquadrate-ovatā, ventricosā, tenuiculā, albida, pallidē rufescente-fusco sparsim flammatā, undique densē irregulariter reticulata.*

THE SICKLY VENUS. Shell somewhat squarely ovate, ventricose, rather thin, whitish, sparingly flamed with pale red-brown, densely irregularly reticulated throughout.

Hab. Singapore; Cuming.

This appears to me to differ from any variety of *V. reticulata*, and it has no indication of the red colouring so characteristic of the internal hinge margin of that species.

Species 75. (Mus. Cuming.)

VENUS STRIATULA. *Ven. testā pectinato-orbiculari, albidū, cinereo-griseo linearē-punctatā et radiatā, concentricē liratā, liris nunc acutis nunc obtusis, subreflexis, interdum undato-corrugatis, confertis.*

THE FINELY-STRIATED VENUS. Shell pectinately orbicular, whitish, linearly dotted and rayed with ashy-grey, concentrically ridged, ridges now sharp now obtuse, a little reflected, sometimes wave-wrinkled, close-set.

Pectunculus striatus, Da Costa, Brit. Conch. p. 191. pl. 12. f. 2.

Venus striatula, Donovan.

Venus gallina, Maton and Rackett.

Venus Pennantii, Forbes.

Venus laminosa, Turton.

Venus rugosa, Pennant.

Venus Prideauxii, Couch.

Venus sulcata and *costata*, Brown.

Hab. European Seas.

The large specimen here figured is the Mediterranean

form of the species. For British examples as originally described by Da Costa, see Plate XX.

Species 76. (Mus. Cuming.)

VENUS APHRODINA. *Ven. testā obliquē cuneatā, fulvo-ferrugineā, pallidē fusco obscurē reticulatā, sublævigatā, posticē concentricē corrugato-sulcatā.*

THE APHRODINE VENUS. Shell obliquely wedge-shaped, fulvous-rust, obscurely reticulated with pale-brown, nearly smooth, concentrically wrinkle-grooved down the posterior side.

LAMARCK, Anim. sans vert. vol. vi. p. 368.

Hab. Australia.

A comparatively smooth species, concentric grooves being scarcely distinguishable except on the posterior side.


Species 77. (Mus. Cuming.)

VENUS ADAMSII. *Ven. testā quadrato-oblongā, tenuiculā, compressā, sordidē albā, ferrugineo tinctā, concentricē laminato-liratā, liris eximie fimbriatis, interstitiis liris tenuibus numerosis radiantibus pulcherrimē clathratis.*


ADAMS' VENUS. Shell squarely oblong, rather thin, compressed, dead-white, tinged with rust, concentrically lamellately ridged, ridges delicately fimbriated, interstices beautifully latticed with numerous fine radiating ridges.

Hab. Japan.


A shell of singularly elliptic form, thin and depressed, most elaborately and delicately sculptured throughout.




78.




79.




80.




81.



82.



84.



83.

VENUS.

PLATE XVIII.

Species 78. (Mus. Cuming.)

VENUS LAMELLATA. *Ven. testā oblongo-ovatā, posticē obliquē trigonā, fulvescente-albā, roseo tintā, concentricē lamellatā, lamellis distantibus, crispato-striatis et reflexis, ad angulum posticum fimbriatis et lacunatis, interstitiis læribus.*

THE LAMELLATED VENUS. Shell oblong-ovate, posteriorly obliquely triangular, fulvous-white, tinged with rose, concentrically lamellated, lamellæ distant, crisply striated and reflected, frilled and spouted at the posterior angle, interstices smooth.

LAMARCK, Anim. sans vert. vol. 6. p. 349.

Hab. Tasmania; R. C. Gunn.

The spouted foliations of the concentric lamellæ are quite remarkable in this very beautiful species.

Species 79. (Mus. Cuming.)

VENUS PERUVIANA. *Ven. testā subangustē ovatā, calcareo-albā, concentricē costatā, costis tumidis, reflexis, ferrugineo-tinctis, posticē laminato-appressis.*

THE PERUVIAN VENUS. Shell rather narrowly ovate, chalk-white, concentrically ribbed, ribs swollen, reflected, rust-dotted, posteriorly laminately appressed.

SOWERBY, Pro. Zool. Soc. 1835. p. 22.

Hab. Ancon, Peru; Cuming.

In this species the concentric laminae, instead of being erectly spouted, are appressed.

Species 80. (Mus. Cuming.)

VENUS ORNATISSIMA. *Ven. testā pectinato-globosā, tenuiculā, sordidē albā, radiatim crebricostatā, costis regularibus, versus marginem duplicibus, concentricē laminato-fimbriatā, fimbriā erectā.*

THE HIGHLY-ADORNED VENUS. Shell pectinately globose, rather thin, dead-white, radiately closely ribbed, ribs regular, duplicate towards the margin, concentrically laminately frilled, frill erect.

BRODERIP, Pro. Zool. Soc. 1835, p. 44.

Hab. Panama (dredged from sandy mud, at a depth of ten fathoms); Cuming.

This elegantly-frilled shell, described nearly thirty years ago, is still unique.

Species 81. (Mus. Cuming.)

VENUS PLICATA. *Ven. testā pectinato-subtrigonā, planiusculā, concentricē plicato-laminatā, laminis erectis, supernē subreflexis, posticē bifariam fimbriatis et lacunatis, interstitiis læribus.*

THE PLAITED VENUS. Shell pectinately subtriangular, rather flat, concentrically plicately laminated, laminæ erect, a little reflected at the top, posteriorly double-frilled and spouted, interstices smooth.

GMELIN, Syst. Nat. p. 1130.

Hab. West Africa.

Some authors have assigned this well-known species to the *V. dysera* of Linnaeus, but so much uncertainty prevails as to which species Linnaeus intended to denote by *V. dysera*, that the name has been expunged from the series altogether, first by Lamarck, and notably so by Deshayes.

Species 82. (Mus. Cuming.)

VENUS KELLETTII. *Ven. testā subangustē pectinato-ovatā, calcareo-albā, ferrugineo-fusco tintā et radiatā, concentricē lineari-sulcatā, sulcis distantibus, prope umboes subtiliter radiatim liratis, areis intermediis læribus, lateribus laminato-alatis.*

KELLETT'S VENUS. Shell rather narrowly pectinately ovate, chalk-white, stained and rayed with rust-brown, concentrically linearly grooved, grooves distant, near the umboes finely radiately ridged, intermediate spaces smooth, sides laminately winged.

HINDS, Moll. Voy. Sulphur, p. 65, pl. 19. f. 5.

Hab. West coast of Veragua, Central America.

Distinguished by foliations on either side, without any intervening lamellæ.

Species 83. (Mus. Cuming.)

VENUS ALATUS. *Ven. testā subtrigon.o-ovatā, fulvescente-albā, spadiceo-fuscescente radiatā, umbonibus carneo*

VENUS.—PLATE XVIII.

*tinctis, lævigatâ vel subirregulariter striatâ, ad latus
anticum laminato-alatâ.*

THE WINGED VENUS. Shell somewhat triangularly ovate, fulvous-white, rayed with light fawn-brown, imboes flesh-tinted, smooth or somewhat irregularly striated, lamellately winged on the anterior side.

Hab. Swan River.

A new form of the genus, entirely destitute of lamellæ, with foliations, few in number, on the anterior side only.

Species 84. (Mus. Cuming.)

VENUS YATESII. *Ven. testâ subtrigono-oval longâ, fulves-*

*cente-croceâ, lævigatâ, concentricè lamellatâ, lamellis
brevibus, subdistantibus, irregularibus, posticè muricato-
fimbriatis.*

YATES'S VENUS. Shell somewhat triangularly oblong, fulvous saffron-colour, smooth, concentrically lamellated, lamellæ short, rather distant, irregular, posteriorly prickly-frilled.


GRAY, Yates' New Zealand, Appendix.

Hab. New Zealand.


The lamellæ are here comparatively short, but curiously prickly spouted on the posterior side.



85a.




85b.



85c.




86



87.



88



89

VENUS.

PLATE XIX.

Species 85. (Fig. a, b, c, Mus. Cuming.)

VENUS SUBIMBRICATA. *Ven. testā pectinato-ovatā, crassā, albā, rufo-fusco copiosè radiatim maculatā et punctatā, concentricè costatā, costis latis, obtusè tumidis, radiatim liratis, liris regularibus, interdum duplicitibus, lunulā et areā ligamenti latè excavatis.*

THE IMBRICATED VENUS. Shell pectinately ovate, thick, white, copiously radiately blotched and dotted with red-brown, concentrically ribbed, ribs broad, obtusely swollen, radiately ridged, ridges regular, sometimes duplicate, lunule and ligamentary area broadly excavated.

SOWERBY, Pro. Zool. Soc. 1835, p. 21.

Hab. Puerto Portrero, Central America (in fine sand at a depth of thirteen fathoms); Cuming.

A solid richly-coloured species, with somewhat of the general external aspect of a *Pectunculus*.

Species 86. (Mus. Cuming.)

VENUS SUBRUGOSA. *Ven. testā subtrigono-ovalā, fulcescente-albā, griseo et olivaceo-nigrante conspicuè quadri-radiatā, lavigatā, versus umbones plus minus corrugato-costatā, interstitiis crispato-suleatis et punctatis.*

THE WRINKLED VENUS. Shell somewhat triangularly ovate, fulvous-white, conspicuously four-rayed with grey and olive-black, smooth, more or less wrinkle-ribbed towards the umboes, the interstices being crisply grooved and punctured.

SOWERBY, Genera of Shells, *Venus*, f. 2.

Hab. West Columbia; Cuming.

Almost invariably characterized by four conspicuous dark rays.

Species 87. (Mus. Cuming.)

VENUS CALLOSA. *Ven. testā pectinato-ovatā, posticè subtrigono-productā purpureo-griseo et ferrugineo tinctā et variegatā, radiatim duplicito-liratā, concentricè irregulariter rude plicato-striatā.*

THE CALLOUS VENUS. Shell pectinately ovate, posteriorly somewhat triangularly produced, stained and variegated with purple-grey and rust, radiately dupli-

cately ridged, concentrically irregularly rudely pli-cately striated.

CONRAD, Journ. Acad. Nat. Sci. Phil. vol. vii. p. 252.

Hab. Sandwich Islands and California.

Richly stained within with violet on the posterior side. The lunule and ligamentary area are almost obsolete.

Species 88. (Mus. Cuming.)

VENUS CANCELLOTA. *Ven. testā pectinato-ovatā, fulvō, brunneo maenlatā et aspersā, concentricè laminatā, laminis brevibus, distantibus, versus marginem irregularibus, radiatim liratā, liris duplicitibus, areā ligamenti latè excavatā, brunneo strigata.*

THE CANCELLED VENUS. Shell pectinately ovate, fulvous, blotched and sprinkled with brown, concentrically laminated, laminae short, distant, irregular towards the margin, radiately ridged, ridges duplicate, ligamentary area broadly excavated, striped with brown.

LINNÆUS, Syst. Nat. p. 1130.

Venus dysera Linnei, Chemnitz.

Hab. Honduras.

So much confusion attends the synonymy of the Linnaean *V. dysera* that the species, as already stated, is expunged from the catalogue.

Species 89. (Mus. Cuming.)

VENUS PAPHIA. *Ven. testā trigono-ovatā, crassā, carneo-albā, violaceo, fuscescente undique undato-maculatā, concentricè costatā, costis latissimis callosis, posticè erectè laminatis, areā ligamenti latissimè excavatā.*

THE PAPHIA VENUS. Shell triangularly ovate, thick, flesh-white, wave-marked throughout with violet-brown, concentrically ribbed, ribs very broad and callous, posteriorly erectly laminated, ligamentary area very broadly excavated.

LINNÆUS, Syst. Nat. p. 1129.

Hab. West Indies.

The type of an exceedingly interesting group, of which some conspicuous examples occur in Plate XXIII.




91 a.



90 a.



91 b.




92.




93.



94.




95.



96 a.



90 b.



96 b.

VENUS.

PLATE XX.

Species 90. (Fig. a, b, Mus. Cuming.)

VENUS SPURCA. *Ven. testā ovatā, plus minus crassā, fulvescente-spadiceā, violaceo-fusco interdum linearis-undatā, concentricē costatā, costis tunidis, obtusis, confertis, lunulā et areā ligamenti angustis.*

THE DIRTY VENUS. Shell ovate, more or less thick, fulvous-drab, sometimes linearly waved with violet-brown, concentrically ridged, ribs swollen, obtuse, close-set, lunule and ligamentary area narrow.

SOWERBY, Pro. Zool. Soc. 1835, p. 23.

Venus crassa, denticulata, mesodesma, spissa, and violacea, Quoy.

Venus cælala, Menke.

Hab. New Zealand; Quoy. Valparaiso and Philippine Islands; Cuming.

A well-marked species, possessing a singularly wide geographical distribution.

Fig. 91. (Fig. a, b, Mus. Cuming.)

Having figured only the large Mediterranean form of *V. striatula* (*V. gallina*, Maton and Rackett), we give here two typical British varieties.

Species 92. (Mus. Cuming.)

VENUS AÆQUILATERA. *Ven. testā trigono-oblongā, compressā, subæquilaterali, fulvescente, carneo-fuscescente latè triradiatā, versus marginem fusco lentiginosā, concentricē liratā, liris subtilibus, plano-obtusis, creberrimis, plus minus obsoletis, lunulā et areā ligamenti angustis.*

THE EQUAL-SIDED VENUS. Shell triangularly oblong, compressed, nearly equal-sided, fulvous, broadly three-rayed with flesh-brown, freckled towards the margin with brown, concentrically ridged, ridges fine, flatly obtuse, very close-set, more or less obsolete, lunule and ligamentary area narrow.

Donax aequilatera, Sowerby, Tusk. Cat. p. 12.

Venus aequilatera, Sowerby, jun.

Hab. Japan.

This species when catalogued in the Tankerville collection, was referred by Mr. Sowerby, with a note of doubt, to the *Venus donacina* of Chemnitz, figured below it in the accompanying plate.

Species 93. (Mus. Cuming.)

VENUS PERONII. *Ven. testā ovato-enneiformi, latere antico brevissimo, subventricosā, fulvescente, griseo linearis-reticulatā, concentricē liratā, liris confertis, plano-obtusis.*

PERON'S VENUS. Shell ovately wedge-shaped, anterior side very short, rather ventricose, fulvous, linearly reticulated with grey, concentrically ridged, ridges close-set, flatly obtuse.

LAMARCK, Anim. sans vert. vol. vi. p. 369.

Hab. Australia.

Of the same peculiar type as *V. Aphrodinoides*, but more ovate and gibbous, with the concentric ridges almost obsolete.

Species 94. (Mus. Cuming.)

VENUS STRIGOSA. *Ven. testā orato-enneiformi, posticē subattenuatā, fulvescente-spadiceā, concentricē liratā et striatā, liris striisque confertis, valde irregularibus, ad latera corrugatis, lunulā et areā ligamenti angustis.*

THE FULLY-STRRIATED VENUS. Shell ovately wedge-shaped, rather attenuated posteriorly, fulvous-rust, concentrically ridged and striated, ridges and striae crowded, very irregular, wrinkled at the sides, lunule and ligamentary area narrow.

LAMARCK, Anim. sans vert. vol. vi. p. 368.

Hab. Australia.

As intimated by M. Deshayes, this may prove to be a ridged variety of *V. Aphrodina*.

Species 95. (Mus. Cuming.)

VENUS DONACINA. *Ven. testā trigono-ovatā, compressā, subæquilaterali, crassā, latere postico flexuoso-angulato,*

antico rotundato, fulvescente, purpureo-griseo undato-lentiginosā, lærigatā, vel obscurè concentricè planoliratā.

THE DONAX-LIKE VENUS. Shell triangularly ovate, compressed, nearly equilateral, thick, posterior side flexuously angled, anterior rounded, fulvous, wave-freeckled with purple-grey, smooth, or obscurely concentrically flatly ridged.

CHEMNITZ, Conch. Cab. vol. xi. p. 231. pl. 202. f. 1983.
1984.

Donax Venusiformis, Lamarck.
Venus semicancellata, Koch.

Hab. Japan.

Cheznitz truly noticed that this peculiar species combined the form of *Donax* with the hinge of *Venus*.

Species 96. (Fig. a, b, Mus. Cuming.)

VENUS SCALARINA. *Ven. testa ovato-cuneiformi, nunc compressa nunc ventricosinclusa, fulvescente-alba, griseo lineari-undata, concentrica tirata, liris valde irregularibus, obtusis, ad latera corrugato-productis, area ligamenti subexcavata.*

THE LADDER-MARKED VENUS. Shell ovately wedge-shaped, sometimes compressed, sometimes rather ventricose, fulvous-white, linearly waved with grey, concentrically ridged, ridges very irregular, obtuse, corrugately produced at the sides, ligamentary area slightly excavated.

LAMARCK, Anim. sans vert. vol. vi. p. 359.

Venus conularis, Lamarck.

Hab. Australia.

Very closely allied to *V. Aphrodisoides* and *strigosa*, but of a different form with the ligamentary area more defined.




97 a.




97 b.



98 a.



98 b.




98 c.



99.



98 d.



100.



101.



102.

V E N U S.

PLATE XXI.

Species 97. (Mus. Cuming.)

VENUS SCABRA. *Ven. testā oratā, posticē subflexuosa, tenui, sordidē spadiceā, radiatim tenuiratā, concentricē densē seabroso-fimbriatā, lunulā subamplā, areā ligamenti augustā.*

THE ROUGH VENUS. Shell ovate, slightly flexuous posteriorly, thin, dull fawn-colour, radiately finely ridged, concentrically densely seaborously fimbriated, lunule rather large, ligamentary area narrow.

HANLEY, Pro. Zool. Soc. 1844. p. 161. Recent Shells, App. p. 361. pl. 16. f. 24.

Hab. Australia.

A shell of light substance, elaborately seaborously sculptured throughout.

Species 98. (Fig. *a* to *d*, Mus. Cuming.)

VENUS MACRODON. *Ven. testā oblique trigono-ovatā, posticē flexuoso-productā, crassā, fulvescente-albā, griseo interdum lenticinosā, interdum omnino violaceo-fuscā, concentricē rudē rugatā, rugis ad angulum posticum parvis, numerosis, areā ligamenti amplā, latissimā.*

THE LARGE-TOOTH VENUS. Shell obliquely triangularly ovate, posteriorly flexuously produced, thick, fulvous-white, sometimes freckled with grey, sometimes altogether violet-brown, concentrically rudely wrinkled, wrinkles small and numerous at the posterior angle, ligamentary area large, very broad.

Cytherea macrodon, Lamarek, Anim. sans vert. vol. 6. p. 327.

Cytherea lunulata, Lamarek.

Venus macrodon, Philippi.

Hab. Brazil.

This species, as may be seen from the four specimens selected for illustration, is extremely variable both in form and colour.

Species 99. (Mus. Cuming.)

VENUS FLEXUOSA. *Ven. testā oratā, gibbosā, posticē flexuoso-attenuatā, fulvescente-spadiceā, lirido-griseo*

tinctā et radiatā, concentricē subrugatā, rugis sulcis linearibus radiantibus crenatis, lunulā amplā, areā ligamenti subangustā.

THE FLEXUOUS VENUS. Shell ovate, gibbous, posteriorly flexuously attenuated, fulvous-drab, stained and radiated with livid-grey, concentrically rather rudely wrinkled, wrinkles crenated by radiating linear grooves, lunule large, ligamentary area rather narrow.

LINNÆUS, Syst. Nat. p. 1133.

Venus impressa, Anton.

Hab. China.

A more gibbous shell than the preceding, more attenuately produced on the posterior side, and having, therefore, a narrower ligamentary area.

Species 100. (Mus. Cuming.)

VENUS DECORATA. *Ven. testā cordato-oratā, gibbosā, posticē attenuato-productā, fulvescente-albā, ferrugineo-fusco tinctā et radiatā, concentricē costatā, liris radiantibus crenato-clathratā, liris posticis seaborosis, lunulā et areā ligamenti nitidē impressis, intensē fuscis.*

THE ADORNED VENUS. Shell cordately ovate, gibbous, posteriorly attenuately produced, fulvous-white, stained and radiated with rust-brown, concentrically ribbed, crenately latticed with radiating ridges, posterior ridges seaborous, lunule and ligamentary area neatly impressed, dark brown.

BRODERIP and SOWERBY, Zool. Journ. vol. 5. p. 49.

Venus bella, Jonas.

Hab. West Indies.

The latticed sculpture is in strong relief, imparting an elaborately carved aspect to the shell.

Species 101. (Mus. Cuming.)

VENUS SQUAMOSA. *Ven. testā cordato-oratā, gibbosinsculā, posticē flexuoso-acuminata, sordidē fulvescente, radiatim lirata, liris confertis, liris tenuibus, seaborosis, rudē fimbriato-clathratis, lunulā amplā, areā ligamenti amplā, subcompressā.*

THE SCALED VENUS. Shell cordately ovate, rather gib-

bous, posteriorly flexuously acuminate, dull-fulvous, radiately ridged, ridges close-set, rudely fimbriately latticed with thin scarrous ridges, lunule large, ligamentary area large but rather compressed.

LINNÆUS, Syst. Nat. p. 1133.

Cytherea squamosa, Lamarck.

Hab. Philippine Islands, Bay of Manilla (in sandy mud); Cuming.

The latticed sculpture of this species is not in sharp relief as in *V. decorata*, but crowded and obtuse.

Species 102. (Mus. Cuming.)

VENUS SUBNODULOSA. *Ven. testā suboblongo-ovatā, subaequilaterali, fulvescente-albā, roseo-fusco tinctā et radiatim trigono-maculatā, radiatim plicato-costatā, cos-*

tis et intersticiis undique regulariter lineari-sulcatis, lunulā impressā, intense fuscā, areā ligamenti pareā, excavatā, maculatā.

THE SLIGHTLY-NODULED VENUS. Shell rather oblong-ovate, nearly equilateral, fulvous-white, stained and triangularly radiately blotched with rose-brown, radiately plicately ribbed, ribs and interstices regularly linearly grooved throughout, lunule impressed, dark-brown, ligamentary area small, excavated, spotted.

HANLEY, Pro. Zool. Soc. 1844. p. 160. Recent Shells, App. p. 360. pl. 16. f. 19.

Hab. Philippine Islands.

Encircled with great regularity by fold-like ribs, which together with the interstices, are beautifully linearly grooved throughout. The grooves can scarcely, however, be said to render the ribs nodulous.




103.




104a.




104b



104c.



105a




105b.



106



107a



107b.



108.

V E N U S.

PLATE XXII.

Species 103. (Mus. Cuming.)

VENUS COSTELLIFERA. *Ven. testā transversē oratā, sub-aquilateralē, fulvescente-albā, rufo-fusco sparsim punctatā, undique radiatim costellatā, costellis confertis, concentricē imbricatis, lunulā obsoletā, areā ligamenti parrā, angustā.*

THE FINELY-RIBBED VENUS. Shell somewhat transversely ovate, nearly equilateral, fulvous-white, sparingly dotted with red-brown, finely radiately ribbed throughout, ribs close-set, concentrically imbricated, lunule obsolete, ligamentary area small, narrow.

ADAMS and REEVE, Moll. Voy. Samarang, p. 79. pl. 21. f. 18.

Hab. Philippine Islands; Beleher.

Allied to *V. Marica*, but destitute of any posterior foliations, and with a much smaller ligamentary area.

Species 104. (Fig. a, b, c, Mus. Cuming.)

VENUS MARICA. *Ven. testā transversē ovatā, sub-aquilateralē, utrinque umbones subangulatā, fulvescente-albā, interdum roseo tinctā, spadiceo-fusco radiatim maculatā, radiatim costellatā, costellis laminis scabrosis concentricis imbricatis, laminis ad angulum posticum foliaceis, lunulā et areā ligamenti subamplis, modicē impressis, saturatē fuscis.*

THE MARICA VENUS. Shell transversely ovate, nearly equilateral, rather angular on either side of the umbones, fulvous-white, sometimes tinged with rose, radiately blotched with fawn-brown, radiately finely ribbed, ribs imbricated with concentric seaceous laminæ, laminæ foliaceous on the posterior angle, lunule and ligamentary area rather large, moderately impressed, dark-brown.

LINNAEUS, Syst. Nat. p. 1130.

Hab. Philippine Islands; Cuming.

It will be seen that a great deal of interesting character is developed in this species.

Species 105. (Fig. a, b, Mus. Cuming.)

VENUS BILINEATA. *Ven. testā cordato-orbiculari, interdum subtriangulari, crassā, fulvescente, olivaceo-fusco undique radiatim bilineatā, radiatim tenuiliratā, con-*

centricē liratā, liris irregularibus, nunc laminatis, nunc obtusis, lunulā et areā ligamenti intensē fuscis, acutē excavatis.

THE TWO-LINED VENUS. Shell cordately orbicular, sometimes rather triangular, thick, fulvous, radiately two-lined throughout with olive-brown, radiately finely ridged, concentrically ridged, ridges irregular, sometimes laminated, sometimes obtuse, lunule and ligamentary area dark-brown, sharply excavated.

Hab. Gulf of California.

A strongly-marked species, partaking of the characters of *V. compta* and *subimbricata*; all three may, indeed, be different states of one and the same species.

Species 106. (Mus. Cuming.)

VENUS GALLINULA. *Ven. testā transversē ovatā, sub-oblongā, fulvescente, spadiceo-fusco undique punctatā, concentricē laminato-costatā, costis numerosis, sub-reflexis, suberulatis, lunulā et areā ligamenti modicē excavatis.*

THE HEN VENUS. Shell transversely ovate, rather oblong, fulvous, dotted throughout with fawn-brown, concentrically laminate ribbed, ribs numerous, a little reflected, slightly crenulated, lunule and ligamentary area moderately excavated.

LAMARCK, Añim. sans vert. vol. vi. p. 348.

Hab. Australia.

Concentrically reflectedly ridged with great regularity, the ridges being dotted throughout with fawn-brown. I follow Mr. Sowerby in assigning this shell to Lamarek's *V. gallinula*, but it should be stated that great doubt attaches to that opinion. Lamarek describes his species as being "cordately elliptic," which this shell is not.

Species 107. (Fig. a, b, Mus. Cuming.)

VENUS AUSTRALIS. *Ven. testā transversē ovatā, utrinque umbones subtriangulari, fulvescente-albā, ferrugineo-roseo interdum tinctā, spadiceo-fusco interrupte triradiatā, undique confertim imbricato-clathratā, lunulā et areā ligamenti subangustis.*

THE SOUTHERN VENUS. Shell transversely ovate, some-

VENUS.—PLATE XXII.

what triangular on either side the umboes, fulvous-white, sometimes tinged with rust-rose, interruptedly three-rayed with fawn-brown, closely imbricately latticed throughout, lunule and ligamentary area rather narrow.

SOWERBY, Pro. Zool. Soc. 1835. p. 22.

Hab. Australia.

Exquisitely sculptured throughout, and almost invariably characterized by three interrupted rays of colour.

Species 108. (Mus. Cuming.)

VENUS FASCIATA. *Ven. testá cordato-orbiculari, compressá, crassá, fulcescente-carneá vel aurantiá, rufo irregulariter radiatum fasciatá, radiatim corrugato-costatá, costis versus marginem calloso-incrassatis et*

obtusis, lunula parum impressá, areá ligamenti subprofundè excavatá.

THE BANDED VENUS. Shell cordately orbicular, compressed, thick, fulvous flesh-colour, or orange, irregularly radiately banded with red, radiately wrinkle-ribbed, ribs callously thickened and obtuse towards the margin, lunule but little impressed, ligamentary area rather deeply excavated.


Pectunculus fasciatus, Da Costa, Brit. Conch. p. 188. pl. 13. f. 3.

Venus fasciata, Donovan.

Venus Brouniarti, Payraudeau.

Hab. European seas, from the Mediterranean to the shores of Norway (at depths of from ten to fifty fathoms).


The colour of this well-known species varies from bright orange-red to deep purple-red, and sometimes the pattern is not only rayed but reticulately flamed.



109 b




109 a.



110




111.




112




113



114.



115.



116 b



116 a

VENUS.

PLATE XXIII.

Species 109. (Fig. 109, *a*, *b*, and 110, Mus. Cuming.)

VENUS THIARA. *Ven. testā cordato-orbiculari, subdepressā, fulvescente-albā, nitente, purpureo modicē radiatā, concentricē laminatā, laminis latis, interdum callosiusculis, subreflexis, posticē interruptis et foliaceis.*

THE TURBAN VENUS. Shell cordately orbicular, rather depressed, fulvous-white, shining, moderately rayed with purple, concentrically laminated, laminae broad, sometimes rather callous, a little reflected, interrupted and foliaceous on the posterior side.

DILLWYN, Desc. Cat. vol. i. p. 162.

Venus foliacea, Philippi.

Hab. —?

An elegantly laminated shell, more or less stained with purple.

Species 111. (Mus. Cuming.)

VENUS DECLIVIS. *Ven. testā orbiculari-ovalā, crassiusculā, fulvescente-albā, purpureo-fusco variè lentiginosā, concentricē laminato-liratā, laminis erectis, regularibus, ad latū antīcum appressis, areā ligamenti modicē excavatā.*

THE SLOPING VENUS. Shell orbicularly ovate, rather thick, fulvous-white, variously freckled with purple-brown, concentrically lamination ridged, laminae erect, regular, appressed on the posterior side, ligamentary area moderately excavated.

SOWERBY, Thes. Conch. part 14. p. 730. pl. 157. f. 123, 124.

Hab. —?

The laminae are unusually regular, rather appressed on the posterior side.

Species 112. (Mus. Cuming.)

VENUS ISABELLINA. *Ven. testā cordato-orbiculari, solidiusculā, nitente-albā, carneo-purpureo sparsim tinctū, concentricē laminatā, laminis subdistantibus, erectis, ad latū antīcum excavato-interruptis, lunulā purā, areā ligamenti subamplā, latā.*

THE ISABELLINE VENUS. Shell cordately-orbicular, rather solid, shining white, sparingly tinged with flesh-purple, concentrically laminated, laminae rather distant, erect, excavately interrupted on the anterior side, lunule small, ligamentary area rather large, broad.

PHILIPPI, MS. in Mus. Cuming.

Hab. Sydney.

This may be regarded as the Australian representative of the China *V. calophylla*. It is a more solid shell, of rounder form, with no posterior foliations of the laminae.

Species 113. (Mus. Cuming.)

VENUS ROBORATA. *Ven. testā subcordato-orbiculari, crassā, depressā, eburnea, concentricē costatā, costis conspicue reflexis, ad extremitatem antīcum radiatim excavatis, areā ligamenti modicē excavatā.*

THE STRENGTHENED VENUS. Shell subcordately orbicular, thick, depressed, ivory-white, concentrically ribbed, ribs conspicuously reflected, radiately excavated at the anterior extremity, ligamentary area moderately excavated.

HANLEY, Pro. Zool. Soc. 1844. p. 161. Recent Shells, App. p. 361. pl. 16. f. 25.

Venus placida, Philippi.

Hab. Tasmania.

An excellent species of shining ivory-white substance, with the ribs developed in bold relief and obtusely reflected.

Species 114. (Mus. Cuming.)

VENUS CALOPHYLLA. *Ven. testā cordato-orbiculari, calcareo-albā, concentricē laminatā, laminis latis, distantibus, declivi-erectis, ad latū antīcum radiatim excavatis, ad posticum interrupte foliaceis, areā ligamenti modicē excavatā.*

THE BEAUTIFUL-LEAVED VENUS. Shell cordately orbicular, chalk-white, concentrically laminated, laminae broad, distant, slopingly erect, radiately excavated on the anterior side, interruptedly foliaceous on the posterior, ligamentary area moderately excavated.

HANLEY, Recent Shells, App. p. 361. pl. 16. f. 26.

Venus Cumingii, Sowerby.

Hab. China.

For a long time this well-known China shell was regarded by collectors as the *V. thiara*.

Species 115. (Mus. Cuming.)

VENUS ALTA. *Ven. testá cordato-trigoná, crassiusculá, compressá, fulvescente-albá, carneo-purpurascente tati-radiatá, concentricè laminatá, laminis callosiusculis, paululum reflexis, areá ligamenti amplá, latè excavatá, intense violaceo-purpureá.*

THE CHERISHED VENUS. Shell cordately triangular, rather thick, compressed, fulvous-white, broadly rayed with flesh-purple, concentrically laminated, laminae rather callous, a little reflected, ligamentary area large, broadly excavated, deep violet-purple.

SOWERBY, Thes. Conch. part 14. p. 724. pl. 158. f. 131 to 133.

Hab. Philippine Islands; Cuming.

A solid compressly-pinched shell, with a broadly-excavated burnt purple-stained ligamentary area.

Species 116. (Mus. Cuming.)


VENUS CYPRIA. *Ven. testá trigono-ovatá, subangustá, posticè attenuatá, fulvescente-albá, carneo-purpurascente radiatá et aspersá, concentricè costatá, costis plus minus incrassatis, reflexis, posticè sublucunatis, lunulá et areá ligamenti subampolis.*

THE CYPRIA VENUS. Shell triangularly ovate, rather narrow, posteriorly attenuated, fulvous-white, rayed and sprinkled with flesh-purple, concentrically ribbed, ribs more or less thickened, reflected, posteriorly slightly spouted, lunule and ligamentary area rather large.

SOWERBY, Pro. Zool. Soc. 1852. p. 43.

Hab. Isle of Plata, West Columbia.

This is not a satisfactory species. It has all the appearance of being an attenuately-produced form of the West Indian *V. Papilio*.



VENUS.

PLATE XXIV.

Figure 117. (Mus. Cuming.)

This shell, which was thought to be an undescribed species, appears to be a rather gibbons form of *V. undulosa*.

Species 118. (Mus. Cuming.)

VENUS IMBRICATA. *Ven. testā ovato-subcordatā, posticē angulatā, albida, ferrugineo-fusco tinctā, ad umbones livido-violaceā, radialim costatā, costis crenato-imbricatis, interstitiis nitidē excavatis, pallidis.*

THE IMBRICATED VENUS. Shell ovately subcordate, posteriorly angled, whitish, tinged with rust-brown, livid-violet at the umboes, radiately ribbed, ribs crenately imbricated, interstices neatly excavated, pale.

SOWERBY, Thes. Conch. vol. ii. p. 175. pl. 156. f. 81, 82.

Venus Cochinensis, Sowerby.

Chione imbricata and *Cochinensis*, Deshayes.

Hab. China Seas.

A radiately ribbed *Cardita*-like shell, the ribs being neatly imbricated throughout with equidistant scales, and the ribs are coloured with a rust-brown hue, while the interstices are cleanly excavated and paler in colour. The umboes are livid-violet.

Species 119. (Fig. *a, b*, Mus. Cuming.)

VENUS FLUCTUOSA. *Ven. testā subtrigono-ovalatā, tenui, compressiuscula, posticē subproductā, albida, epidermide lutescente nitente indutā, ad umbones plus minus erosā, concentricē irregulariter corrugato-liratā.*

THE WAVY VENUS. Shell somewhat triangularly ovate, thin, rather compressed, posteriorly slightly produced, whitish, covered with a shining yellowish epidermis, more or less eroded at the umboes, concentrically irregularly wrinkle-ridged.

GOULD, Invert. Mass. p. 87. f. 50.

Venus Astartoides, Beck.

Chione Astartoides, Deshayes.

Hab. Greenland; Newfoundland; Bay of Massachusetts, United States.

Of quite a different type from any other species of the genus.

Species 120. (Mus. Cuming.)

VENUS CUNEIFORMIS. *Ven. testā obliquē cuneiformi, solidū, compressiuscula, albida, fusco pallide linearis-nudatā, ad umbones livido-purpurascente, concentricē irregulariter rudē plicato-striatā, striis circa umbones et ad marginem posticum tirutis.*

THE WEDGE-SHAPED VENUS. Shell obliquely wedge-shaped, solid, rather compressed, whitish, faintly linearly waved with brown, livid-purple at the umboes, concentrically irregularly rudely plicately striated, striae ridged round the umboes and along the posterior margin.

Hab. —?

This shell approximates to the Australian *V. Aphrodisoides*, but it differs materially in sculpture.

Species 121. (Mus. Cuming.)

VENUS TASMANICA. *Ven. testā subtrigono-oblonga, compressa, posticē subtruncata, albida, livido-purpurascente plus minus tincta et fasciata, concentricē plicato-costatis, costis obtuse reflexis, ad basin crenulatis, lunula parva, oblongo-cordata, marginibus denticulatis, intus livido-purpurea.*

THE TASMANIAN VENUS. Shell somewhat triangularly oblong, compressed, posteriorly rather truncated, whitish, more or less tinged and banded with livid-purple, concentrically plicately ribbed, ribs obtusely reflected, crenulated at the base, lunule small, oblong-cordate, margins denticulated, livid-purple within.

Hab. Tasmania.

An interesting little species of the same peculiar type as M. Deshayes' *V. mitis*, one of its most conspicuous features being the basal crenulation of the ribs.

Species 122. (Mus. Cuming.)

VENUS BERRII. *Ven. testā cordato-trigonatā, solidū, compressa, valde inaequilateralis, albida, livido-cinereo antifuscescente tincta, concentricē plicato-lirata, areæ ligamenti ampliæ, curvata, concavo-excavata.*

BERRY'S VENUS. Shell cordately triangular, solid, compressed, very inequilateral, whitish, tinged with livid-

ash or pale-brown, concentrically plicately ridged, ligamentary area large, curved, concavely excavated.

GRAY, Ind. Test. Supp. pl. 2. f. 2.

Chione Berrii, Deshayes.

Hab. Australia.

A solid well-marked species, of a peculiarly livid tone of colour.

Species 123. (Mus. Cuming.)

VENUS EXILIS. *Ven. testā ovatā, modicē convexā, cærulescente-albā, epidermide corneā lævigatā tenui indutā, lunulā nullā.*

THE SLENDER VENUS. Shell ovate, moderately convex, bluish-white, covered with a thin smooth horny epidermis, no lunule.

LAMARCK, Annn. sans vert. vol. vi. p. 359.

Hab. Ceylon.

It is not without considerable doubt that I refer this shell to Lamarek's *Venus exilis*. It is of very simple character, covered with a smooth horny epidermis.

Species 124. (Mus. Cuming.)

VENUS SALLEI. *Ven. testā trigono-Donaciformi, versus margines compressā, albida, cinereo-griseo latiradiatā,*

latere antico lævigato, vel exiliter plicato, postico densè clathrato.

SALLÈ'S VENUS. Shell triangularly *Donax*-shaped, compressed towards the margins, whitish, broadly rayed with ash-grey, anterior side smooth or faintly concentrically plicated, posterior side densely latticed.

Hab. Bird's Island, Caribbean Sea; Sallè.

A very characteristic little species, with the general aspect of a *Donax*.

Species 125. (Mus. Cuming.)

VENUS VARICOSA. *Ven. testā obliquè trigonā, subcordatā, albida, fulvescente-ryfo obscurè radiatā et linearī-marmoratā, concentricè costalā, costis crassissimis, latissimis.*

THE VARICOSE VENUS. Shell obliquely triangular, somewhat heart-shaped, whitish, obscurely rayed and linearly mottled with fulvous-red, concentrically ribbed, ribs very thick and very broad.

SOWERBY, Thes. Conch. vol. ii. p. 723. pl. 155. f. 67.

Chione varicosa, Deshayes.

Hab. Bay of Mexico.

Remarkable for the very broad and solidly-developed character of its concentric ribs.



VENUS.

PLATE XXV.

Species 126. (Fig. *a, b*, Mus. Cuming.)

VENUS UNDULOSA. *Ven. testā trigonā, subaequilaterati, compressè convexā, albida, rufo-fusco plus minus lineariter undulata, concentricè tenuistriatā.*

THE WAVED VENUS. Shell triangular, nearly equilateral, compressly convex, whitish, more or less linearly waved with red-brown, concentrically finely striated.

LAMARCK, Anim. sans vert. vol. vi. p. 370.

Venus undulosa and *variabilis*, Sowerby.

Chione undulosa, Deshayes.

Hab. Swan River, New Holland.

Of an unusual equilaterally triangular form, more or less copiously waved with red-brown lines upon a white ground.

Species 127. (Mus. Brit.)

VENUS MITIS. *Ven. testā subtrigono-ovatā, posticè subdeclivi-acuminatā, opaco-albā, umbonibus roseis, concentricè laminato-costulatā, costis calloso-reflexis, in interstītiis radiatim crenatis, intus purpurascente.*

THE MILD VENUS. Shell somewhat triangularly-ovate, posteriorly rather slopingly acuminate, opaque-white, umboes rose-coloured, concentrically lamellately ribbed, ribs callously reflected, radiately crenated in the interstices, purple within.

Chione mitis, Deshayes, Cat. Conch. Brit. Mus. p. 142.

Hab. Swan River, New Holland.

A delicate opaque-white *Corbula*-like shell, rose-coloured at the umboes, delicately crenulated in the interstices between the concentric ribs.

Species 128. (Mus. Cuming.)

VENUS GEMMA. *Ven. testā subtrigono-orbiculari, subobliquā, pellucido-albā, livido-purpurascente tinctā, concentricè tennè lirato-striatā.*

THE GEM VENUS. Shell somewhat triangularly orbicular, rather oblique, pellucid white, tinged with livid purple, concentrically finely ridge-striated.

TOTTEN, Sillim. Journ. vol. xxvi. p. 367. f. 2. *a* to *d*.

Gemma gemma, Deshayes.

Hab. Massachusetts Bay, United States.

"This beautiful little amethystine gem, as it has been

appropriately called," says Dr. Gould, "is found in great abundance on all the sandy shores of Massachusetts Bay. Colonel Totten found it in Newport Harbour, beyond which its range is not known."

Species 129. (Mus. Belcher.)

VENUS QUADRANGULARIS. *Ven. testā subquadrato-ovatā, compressinsculā, pallidè stramineā, nitente, umbonibus roseis, concentricè tenuè irregulariter striatā, lunulā parum distinctā.*

THE QUADRANGULAR VENUS. Shell somewhat squarely ovate, rather compressed, pale straw-colour, shining, pink at the umboes, concentrically irregularly striated, lunule but little distinct.

ADAMS and REEVE, Moll. Voy. Samarang, p. 79. pl. 24. f. 7.

Chione quadrangularis, Deshayes.

Hab. Corean Archipelago.

Peculiarly square-formed, of a light shining straw-colour, with pink umboes.

Species 130. (Mus. Brit.)

VENUS RUDERATA. *Ven. testā latè obliquè ovatā, sordidè albā, tenuè radiatim liratā, liris binis, concentricè pli-ecatā, plicis valde irregularibus, ad latius posticum rudi laminatis, lunulā nullā.*

THE RUDE VENUS. Shell broadly obliquely ovate, dead-white, finely radiately ridged, ridges in pairs, concentrically plicated, plaits very irregular, rudely lamellated on the posterior side, no lunule.

Chione ruderata, Deshayes, Cat. Conch. Brit. Mus. p. 136. Hab. California.

Of peculiarly rude growth, with much of the general aspect of a *Venerupis*.

Species 131. (Mus. Cuming.)

VENUS OBESA. *Ven. testā obliquè ovatā, posticè subacuminatā, solidā, gibbosiusculā, lacteā, violaceo-griseo-nitidè obscurè radiatā, lavigatā, epidermide tenui cornue partim indutā.*

THE STOUT VENUS. Shell obliquely ovate, posteriorly

rather acuminate, solid, rather gibbous, cream-colour, delicately obscurely rayed with violet-grey, smooth, partially covered with a thin horny epidermis.
Cytherea obesa, Sowerby, Thes. Conch. vol. ii. p. 626. pl. 131. f. 83, 84.

Dione obesa, Deshayes.

Hab. — ?

A solid gibbous shell belonging more strictly to the genus *Dione*.

Species 132. (Mus. Brit.)

VENUS OBOVALIS. *Ven. testá ovatá, opaco-albá, concentricè densè minutè striatá, striis plus minus erosis, hic illuc obsoletis, ad marginem posticum rudè plicato-corrugatis, lunulá parvá.*

THE OVAL VENUS. Shell ovate, opaque-white, concentrically densely minutely striated, striae more or less eroded, here and there obsolete, rudely plicately wrinkled on the posterior margin, lunule small.

Chione obovalis, Deshayes, Cat. Conch. Brit. Mus. p. 154.
Hab. — ?

A peculiarly soft opaque-white shell, distinguished by a characteristic style of sculpture.

Species 133. (Mus. Belcher.)

VENUS ELEGANS. *Ven. testá oblongo-ovatá, compressè cordatá, candidá, posticè subangulatá, concentricè cos-*

tatá, costis subdistantibus, irregularibus, laminato-reflexis, praesertim ad angulum posticum, lunulá parvá.

THE ELEGANT VENUS. Shell oblong-ovate, compressly heart-shaped, white, posteriorly faintly angled, concentrically ribbed, ribs rather distant, irregular, lamellately reflected, especially on the posterior angle, lunule small.

ADAMS and REEVE, Moll. Voy. Samarang, p. 79. pl. 24. f. 13.

Chione elegans, Deshayes.

Hab. China Seas.

A delicate white shell, neatly sculptured with irregular concentric lamellæ, which are stronger on the posterior angle.

Species 134. (Mus. Cuming.)

VENUS IRREGULARIS. *Ven. testá cordato-ovatá, calcareo-albá, concentricè laminato-liratá et striatá, laminis striisque versus marginem numerosis gradatim convertoribus et irregularibus.*

THE IRREGULAR VENUS. Shell cordately ovate, chalk-white, concentrically lamellately ridged and striated, lamellæ and striae numerous, gradually more close-set, and irregular towards the margin.

Hab. Month of the Gaboon, West Africa.

The concentric lamellæ are tolerably equidistant around the umboes, but soon they become particularly crowded and irregular.



135



136



137 o



137 b



138 a



138 b



138 c



139



140



141

V E N U S.

PLATE XXVI.

Species 135. (Mus. Cuming.)

VENUS STRIATISSIMA. *Ven. testā suboblongo-ovatā, antice subventricosā, posticē subacuminatā, fulvescente-albā, rufescē-fusco maculatā, liris radiantibus striisque lamelliformib⁹ concentricis undique nitidē clathratā.*

THE MUCH-STRIATED VENUS. Shell rather oblong-ovate, anteriorly somewhat ventricose, posteriorly somewhat acuminated, fulvous-white, mottled with red-brown, neatly latticed throughout with radiating ridges and concentric lamelliform striae.

SOWERBY, Thes. Conch. vol. ii. p. 718, pl. 157. f. 103-105.
Chione striatissima, Deshayes.

Hab. New Holland.

A beautifully-sculptured shell, belonging to the *V. marica* section of the genus.

Species 136. (Mus. Cuming.)

VENUS LAYARDI. *Ven. testā suboblongo-ovatā, posticē flexuoso-acuminatā, fulvescente purpureo-fusco tiuctā, concentricē costatā, costis callosis, ad marginem posteriorum bifarium obtusè squamatis, costarum interstitiis radiatim crenulatis, marginibus denticulatis.*

LAYARD'S VENUS. Shell rather oblong-ovate, posteriorly flexuously acuminated, fulvous, tinged with purple-brown, concentrically ribbed, ribs callous, obtusely doubly scaled on the posterior margin, interstices between the ribs radiately crenulated, margins denticulated.

Hab. Ceylon; E. L. Layard.

An interesting flexuously-winged species, of the same crenulated type as the Australian *V. mitis* and the Tasmanian *V. Tasmanica*.

Species 137. (Fig. a, b, c, Mus. Cuming.)

VENUS OVATA. *Ven. testū oratā vel oblongo-ovatā, posticē declivi-truncatā, fulvescente aut griseā, fusco sparsim maculatā, radiatim costatā, costis confertis, undique densē plicato-crenatis, lunulā rie nullā.*

THE OVATE VENUS. Shell ovate or oblong-ovate, posteriorly slopingly truncated, fulvous or grey, sparingly blotched with brown, radiately ribbed, ribs

close-set, densely plicately crenated throughout, scarcely any lunule.

PENNANT, Brit. Zool. vol. iv. p. 206, pl. 59. f. 3.

Venus spadicea, Renier.

Venus pectinula, Lamarck.

Venus radiata, Scacchi.

Cytherea ovata, Thompson.

Chione ovata, Deshayes.

Hab. European Seas.

This pretty species of our own shores has a wide range throughout the European Seas passing into the Atlantic. The little specimen given at Fig. 137 a is from one of the Cape de Verd Islands.

Species 138. (Fig. a, b, c, Mus. Cuming.)

VENUS PYGM.EA. *Ven. testā cordato-oblongā, compressinerculā, valde inaequilateralē, albida, fusco vel purpureo-roseo maculatā, interdum viridē radiatā, liris radiantibus lamellis concentricis fimbriatis undique confertim clathratā.*

THE PYGMY VENUS. Shell cordately oblong, rather compressed, very inequilateral, white, spotted with brown or purple-rose, sometimes brightly rayed, closely latticed throughout with radiating ridges and concentric frilled lamellæ.

LAMARCK, Anim. sans vert. vol. vi. p. 337.

Chione pygmaea, Deshayes.

Hab. West Indies.

A beautifully-sculptured latticed species, in which the concentric lamellæ are furrowed.

Figure 139. (Mus. Cuming.)

This proves to be a specimen of *V. exilis*, already described and figured in Plate XXIV.

Species 140. (Mus. Cuming.)

VENUS CERINA. *Ven. testā subquadrato-cordatā, solidiusculā, fulvescente-albā, ad lunulam et ad aream ligamenti fusco sparsim linearē, concentricē densē striatā.*

VENUS.—PLATE XXVI.

striis hic illic confluentibus, in interstitiis decussatim crenulatā, ad latera tenuē radiatim plicatā, umbonibus subrostratis.

THE WAXEN VENUS. Shell somewhat squarely heart-shaped, rather solid, fulvous-white, sparingly lineated with brown at the lunule and ligamentary area, concentrically densely striated, striae here and there confluent, decussately crenulated in the interstices, finely radiately plicated at the sides, umboes rather beaked.

C. B. ADAMS, MS. in Mus. Cuming.

Hab. —?

This little species will be found, on examination with the lens, to be full of character.

Species 141. (Mus. Cuming.)

VENUS GLOBULUS. *Ven. testā globoso-cordatā, posticē subtruncatā, solidiusculā, fulvescente-albā, undique densē calloso-liratā, liris subirregularibus, hic illic subreflexis, lunulā amplā, subindistinctā.*

THE GLOBULE VENUS. Shell globosely heart-shaped, a little truncated on the posterior side, rather solid, fulvous-white, densely callously ridged throughout, ridges rather irregular, here and there a little reflected, lunule large, rather indistinct.

Hab. —?

A solid globose shell, elaborately concentrically ridged from the umboes throughout, with very much of the general appearance of a *Lucina*.

V E N U S.

	Plate.	Species.		Plate.	Species.
Adamsii, <i>Reeve</i>	XVII.	77	<i>crenata</i> , <i>Gmelin</i>	X.	33
agrota, <i>Reeve</i>	XVII.	74	<i>crenulata</i> , <i>Chemnitzi</i>	X.	33
æquilatera (<i>Donax</i>), <i>Sowerby</i> . .	XX.	92	<i>crispata</i> , <i>Deshayes</i>	IX.	31
affinis, <i>Sowerby</i>	XII.	38	<i>Cumingii</i> , <i>Sowerby</i>	XXIII.	114
alatus, <i>Reeve</i>	XVIII.	83	<i>cuneiformis</i> , <i>Reeve</i>	XXIV.	120
albina, <i>Sowerby</i>	XIV.	55	<i>Cypria</i> , <i>Sowerby</i>	XXIII.	116
alboradiata, <i>Sowerby</i>	III.	7	<i>Cyprinoides</i> , <i>Anton</i>	II.	4
alta, <i>Sowerby</i>	XXIII.	115	<i>declivis</i> , <i>Sowerby</i>	XXIII.	111
Amithusia, <i>Philippi</i>	XI.	36	<i>decorata</i> , <i>Broderip and Sowerby</i> . .	XXI.	100
antiqua (<i>Dosina</i>), <i>Gray</i>	IX.	30	<i>denticulata</i> , <i>Quoy</i>	XX.	90
Aphrodina, <i>Lamarck</i>	XVII.	76	<i>discors</i> , <i>Sowerby</i>	VII.	22
Aphrodinoides, <i>Lamarck</i>	XVII.	73	<i>discina</i> , <i>Lamarck</i>	V.	15
asperrima, <i>Sowerby</i>	VI.	19	<i>diserepans</i> , <i>Sowerby</i>	XVII.	72
Astartoides, <i>Beck</i>	XXIV.	119	<i>donacina</i> , <i>Chemnitzi</i>	XX.	95
australis, <i>Sowerby</i>	XXII.	107	<i>Dombeii</i> , <i>Lamarck</i>	IX.	29
bella, <i>Jonas</i>	XXI.	100	<i>dysera</i> , <i>Chemnitzi</i>	XIX.	88
Berryi, <i>Gray</i>	XXIV.	122	<i>eburnea</i> , <i>Reeve</i>	XIV.	50
bilineata, <i>Reeve</i>	XXII.	105	<i>elegans</i> , <i>Adams and Reeve</i>	XXV.	133
Brongniarti, <i>Payraudeau</i>	XXII.	108	<i>entobapta</i> , <i>Jonas</i>	XVI.	62
cælata, <i>Menke</i>	XX.	90	<i>erycina</i> , <i>Pennant</i>	XII.	40
Californiensis, <i>Broderip</i>	XI.	35	<i>exalbida</i> , <i>Chemnitzi</i>	IV.	13
callosa, <i>Conrad</i>	XIX.	87	<i>exilis</i> , <i>Lamarck</i>	XXIV.	123
calophylla, <i>Hanley</i>	XXIII.	114	<i>fasciata</i> (<i>Pectunculus</i>), <i>Da Costa</i> . .	XXII.	108
cancellata, <i>Donovan</i>	XII.	40	<i>flexuosa</i> , <i>Linnæus</i>	XXI.	99
cancellata, <i>Linnæus</i>	XIX.	88	<i>fluctuosa</i> , <i>Gould</i>	XXIV.	119
cardioides, <i>Lamarck</i>	VIII.	25	<i>foliacea</i> , <i>Philippi</i>	XXIII.	109
casina, <i>Linnæus</i>	V.	15	<i>foveolata</i> , <i>Sowerby</i>	IV.	11
casinula, <i>Deshayes</i>	XVI.	67	<i>fusco-lineata</i> , <i>Sowerby</i>	XVI.	69
ceerina, <i>C. B. Adams</i>	XXVI.	140	<i>gallina</i> , <i>Maton and Rackett</i>	XVII.	75
Chemnitzi, <i>Hanley</i>	X.	32	<i>gallinula</i> , <i>Lamarck</i>	XXII.	106
Chilensis, <i>Sowerby</i>	IX.	29	<i>gemma</i> , <i>Totten</i>	XXV.	125
cincta, <i>Chemnitzi</i>	VII.	23	<i>gibbosula</i> , <i>Deshayes</i>	XIII.	47
cingulata, <i>Lamarck</i>	VIII.	26	<i>globulus</i> , <i>Reeve</i>	XXVI.	141
elathrata, <i>Deshayes</i>	I.	3	<i>Gnidia</i> , <i>Broderip and Sowerby</i> . .	XI.	37
Cochinensis, <i>Sowerby</i>	XXIV.	118	<i>granulata</i> , <i>Gmelin</i>	XVI.	62
Columbiensis, <i>Sowerby</i>	XV.	61	<i>grata</i> , <i>Say</i>	III.	8
compta, <i>Broderip</i>	XIII.	48	<i>histriomica</i> , <i>Sowerby</i>	XVI.	70
conularis, <i>Lamarck</i>	XX.	96	<i>imbricata</i> , <i>Sowerby</i>	XXIV.	118
cor, <i>Sowerby</i>	IX.	28	<i>impressa</i> , <i>Anton</i>	XXI.	99
corbis, <i>Lamarck</i>	X.	34	<i>intermedia</i> , <i>Gray</i>	XVI.	67
costata, <i>Brown</i>	XVII.	75	<i>intersecta</i> , <i>Sowerby</i>	XIV.	52
costellata, <i>Sowerby</i>	IX.	30	<i>irregularis</i> , <i>Reeve</i>	XXV.	134
costellifera, <i>Adams and Reeve</i> . .	XXII.	103	<i>Isabellina</i> , <i>Philippi</i>	XXIII.	112
crassa, <i>Quoy</i>	XX.	90	<i>Jukesii</i> , <i>Deshayes</i>	XVI.	64
crassicostata, <i>Quoy and Gaimard</i>	XII.	42	<i>Kelletii</i> , <i>Hinds</i>	XVIII.	82
crebrisulca, <i>Sowerby</i>	XVI.	64	<i>Kennerleyi</i> , <i>Carpenter</i>	XII.	41

VENUS.—INDEX.


	Plate.	Species.		Plate.	Species.
<i>lacerata</i> , Hanley.....	VI.	18	<i>quadrangularis</i> , Adams and Reeve	XXV.	129
<i>lactea</i> , Donovan.....	V.	15	<i>radiata</i> , Sacchi.....	XXVI.	137
<i>laevinatus</i> , Reeve	XVI.	63	<i>reflexa</i> , Montagu	V.	15
<i>Lamarckii</i> , Gray	XII.	39	<i>resticulata</i> , Sowerby	VIII.	27
<i>laminosa</i> , Turton	XVII.	75	<i>reticulata</i> , Linnaeus	X.	34
<i>lamellata</i> , Lamarck	XVIII.	78	<i>rigida</i> , Dillwyn	VII.	23
<i>laqueata</i> , Sowerby	VI.	20	<i>robusta</i> , Henley	XXIII.	113
<i>Lemanii</i> , Payraudeau.....	XII.	40	<i>rosalina</i> , Rang	V.	16
<i>Layardi</i> , Reeve	XXVI.	136	<i>ruderata</i> (<i>Chione</i>), Deshayes....	XXV.	130
<i>lencodon</i> , Sowerby.....	XI.	35	<i>rugosa</i> , Gmelin	VII.	23
<i>lenticularis</i> , Sowerby	VII.	21	<i>rugosa</i> , Pennant.....	XVII.	75
<i>lima</i> (<i>Tapes</i>), Sowerby	I.	2	<i>Rusterusii</i> , Payraudeau	V.	15
<i>Listeri</i> (<i>Dosina</i>), Gray	V.	14	<i>Sallei</i> , Reeve	XXIV.	124
<i>lunulata</i> (<i>Cytherea</i>), Lamarek ..	XXI.	98	<i>scabra</i> , Hanley	XXI.	97
<i>lyra</i> , Hanley	XV.	56	<i>scalarina</i> , Lamarck	XX.	96
<i>macrodon</i> (<i>Cyltherea</i>), Lamarck ..	XXI.	98	<i>sculpta</i> , Deshayes	II.	5
<i>Mactracea</i> , Broderip	XVII.	71	<i>semicancellata</i> , Koch	XX.	95
<i>magnifica</i> , Sowerby	V.	17	<i>simillima</i> , Sowerby	XIII.	44
<i>Marica</i> , Linnaeus	XXII.	104	<i>Sowerbyi</i> , Deshayes	VIII.	24
<i>mercenaria</i> , Linnaeus	II.	4	<i>spadicea</i> , Renieri	XXVI.	137
<i>mesodesma</i> , Quoy	XX.	90	<i>sphaerisulca</i> , Deshayes	II.	6
<i>mitis</i> (<i>Chione</i>), Deshayes	XXV.	127	<i>spissa</i> , Quoy	XX.	90
<i>monilifera</i> , Sowerby	XIV.	53	<i>spurca</i> , Sowerby	XX.	90
<i>multicostata</i> , Sowerby	III.	9	<i>squamosa</i> (<i>Cytherea</i>), Lamarek ..	XXI.	101
<i>mundulus</i> , Reeve	XIV.	51	<i>striata</i> (<i>Pectunculus</i>), Da Costa ..	XVII.	75
<i>muscaria</i> , Reeve	XV.	60	<i>striatissima</i> , Sowerby	XXVI.	135
<i>neglecta</i> , Sowerby	XIV.	54	<i>strigosa</i> , Lamarck	XX.	94
<i>nodos-a</i> , Dunker.....	XV.	57	<i>Stuchburii</i> , Gray	XV.	59
<i>notata</i> , Say	II.	4	<i>subimbricata</i> , Sowerby	XIX.	85
<i>Nuttallii</i> , Conrad	XIII.	49	<i>subrostrata</i> , Lamarck	XIV.	54
<i>obesa</i> (<i>Cytherea</i>), Sowerby	XXV.	131	<i>subrugosa</i> , Sowerby	XIX.	86
<i>obliqua</i> , Anton	II.	4	<i>subnodulosa</i> , Hanley	XXI.	102
<i>obovalis</i> (<i>Chione</i>), Deshayes ..	XXV.	132	<i>sugillata</i> , Reeve	XIII.	43
<i>oblonga</i> , Hanley	I.	1	<i>sulcata</i> , Brown	XVII.	75
<i>ornatissima</i> , Broderip	XVIII.	80	<i>Tasmanica</i> , Reeve	XXIV.	121
<i>ovata</i> , Pennant	XXVI.	137	<i>tenuilamellata</i> , Sowerby	IV.	12
<i>Paphia</i> , Linnaeus	XIX.	89	<i>tessellata</i> , Adams and Reeve	XVI.	66
<i>pectinula</i> , Lamarck	XXVI.	137	<i>thiara</i> , Dillwyn	XXIII.	109
<i>pectorina</i> , Lamarck	VIII.	25	<i>tricolor</i> (<i>Tapes</i>), Sowerby	III.	8
<i>Pennantii</i> , Forbes	XVII.	75	<i>torcuma</i> , Gould	XVI.	64
<i>Teronii</i> , Lamarck	XX.	93	<i>tuberosa</i> , Deshayes	XV.	57
<i>Peruviana</i> , Sowerby	XVIII.	79	<i>undatella</i> , Sowerby	XVI.	68
<i>pilula</i> , Reeve	XV.	58	<i>undulosa</i> , Lamarck	XXV.	126
<i>placido</i> , Philippi	XXIII.	113	<i>variabilis</i> , Sowerby	XXV.	126
<i>plicata</i> , Gmelin	XVIII.	81	<i>varicosa</i> , Sowerby	XXIV.	125
<i>plumbea</i> , Reeve	XVI.	65	<i>Veneriformis</i> (<i>Donax</i>), Lamarek ..	XX.	95
<i>proparca</i> , Say	II.	4	<i>verrucosa</i> , Linnaeus	XII.	40
<i>Prideauxii</i> , Couch	XVII.	75	<i>violacea</i> , Quoy	XX.	90
<i>puerpera</i> , Linnaeus	IV.	10	<i>Yatesii</i> , Gray	XVIII.	84
<i>pubicaria</i> , Broderip	VIII.	26	<i>Zelandica</i> , Gray	I.	1
<i>pygmæa</i> , Lamarck	XXVI.	138			

MONOGRAPH
OF THE GENUS
CYTHEREA.

"Dost thou love pictures? we will fetch thee straight
Adonis, painted by a running brook,
And Cytherea all in sedges hid."—*Shakspeare*



1.



2.



3.



4.a.

Sowerby, del et lith



4.b.

Vincent Brooks, Im

C Y T H E R E A.

PLATE I.

Genus CY THEREA, Lamarck.

Testa subrigono-ovata, plus minus turgida, crassiuscula, levigata, rurissimè sculpta, marginibus integris. Cardo in utrâque valvâ tridentatus, dente antico subremoto, lateraliter compresso. Impressio pallii posticè vix sinuata.

Shell somewhat triangularly ovate, more or less turgid, rather thick, smooth, very rarely sculptured, margins entire. Hinge in each valve three-toothed, anterior tooth rather distant, laterally compressed. Impression of the mantle scarcely sinuated posteriorly.

Unwilling to give up the well-known generic name of *Cytherea*, it is here used for the characteristic group of smooth *Veneridae*, of which the Lamarckian *C. petechialis*, *impudica*, and *meretrix* are examples, constituting the genus *Meretrix* of Dr. Gray, of M. Deshayes. With these are included the species, typified by the Linnaean *Venus tripla*, and separated by those authors under the head of *Trigona*. They are chiefly natives of China and the Philippine Islands. All of which we have any record of their habitation in the New World are six from Central America, two from West Indies, and two from California. None are reported from Australia or New Zealand. *C. tripla* is a native of Senegal. The only European *Cytherea* is *C. nitidula* of the Mediterranean.

Species 1. (Mus. Cuming.)

CY THEREA GRAPHICA. *Cyth. testâ subrigono-cordatâ, posticè subproductâ, flavescente-griseâ, utois fuscis hieroglyphicis profusè obliquè radiatâ, concentricè rugosâ.*

THE GRAPHIC CY THEREA. Shell somewhat triangularly heart-shaped, posteriorly somewhat produced, yellowish-grey, profusely obliquely rayed with brown hieroglyphic marks, concentrically rugose.

LAMARCK, Anim. sans vert. vol. vi. p. 300.

Meretrix graphica, Deshayes.

Hab. Sumatra.

Distinguished among other characters by the numerous rays of hieroglyphic marks.

Species 2. (Mus. Cuming.)

CY THEREA DOLABELLA. *Cyth. testâ acutè trigonâ, compressâ, violaceo-lacteâ, epidermide corneâ fulvo-olivaceâ*

nitente indutâ, latere postico conspicuè areuato-producto.

THE LITTLE HATCHET CY THEREA. Shell sharply triangular, compressed, violet cream-colour, covered with a shining fulvous-olive horny epidermis, posterior side conspicuously areately produced.

SOWERBY, Thes. Conch. vol. i. p. 619. pl. 127. f. 19.
Hab. Red Sea.

This and the preceding species represent the extreme forms of the genus as indicated by the subgeneric titles *Trigona* and *Meretrix*. It will be seen that in *C. graphicâ* the posterior margin is convex, while in *C. dolabella* it is absolutely concave.

Species 3. (Mus. Cuming.)

CY THEREA CRASSATELLOIDES. *Cyth. testâ obliquè ovatâ, suboblongâ, modice trigonâ, crassâ, ponderosâ, ventricosâ, epidermide corneâ crassâ indutâ, epidermide lacteâ, purpureo-violaceo radiatâ.*

THE CRASSATELLA-LIKE CY THEREA. Shell obliquely ovate, rather oblong, moderately triangular, thick, heavy, ventricose, covered with a thick horny epidermis, epidermis cream-colour, rayed with purple-violet.

Trigona Crassatelloides, Conrad, Ann. Nat. Sci. Philadelphia, vol. vii. p. 253. pl. 19. f. 17.

Cytherea Crassatelloides, Hanley.

Cytherea (Trigonella) Crassatelloides, Hinds.

Hab. Bay of California.

The colouring of this species resides in the horny epidermis, with which the shell is entirely enveloped.

Species 4. (Fig. a, b, Mus. Cuming.)

CY THEREA DAMAOIDES. *Cyth. testâ compressè coniformi, posticè areuato-acuminatâ, solidiusculâ, flavescente-spadiceâ, interdum carueo-fusco radialâ, interdum griseo præfusâ undato-punctatâ.*

THE DEER-LIKE CY THEREA. Shell compressly wedge-shaped, posteriorly areately acuminate, rather solid, yellowish fawn-colour, sometimes rayed with flesh-brown, sometimes profusely wave-dotted with grey.

Venus damaoides, Gray, Wood Ind. Test. Supp. pl. 2. f. 17.

Cytherea damaoides, Hanley.

Trigona damaoides, Gray.

Hab. — ?

A solid compressed *Donax*-shaped shell, in which the colouring is extremely variable.



C Y T H E R E A .

PLATE II.

Species 5. (Mus. Cuming.)

CY THEREA PONDEROSA. *Cyth. testā subquadrato-trigonā, crassā, ponderosā, sordidē lacteā, roseo-purpurascente pallidē radiatā, lærivculā, nitente, concentricē rugosā, areā ligamenti latiusculā.*

THE HEAVY CY THEREA. Shell somewhat square triangular, thick, heavy, dull cream-colour, faintly rayed with rose-purple, nearly smooth, shining, concentrically rugose, ligamentary area rather broad.

HANLEY, Deser. Cat. p. 107.

Trigona damaoides, var., Deshayes.

Hab. —?

M eshayes quotes this species as a variety of *C. da-*
maoides, which is a distinct form of uniformly smaller size.

Species 6. (Mus. Cuming.)

ERE A CASTANEA. *Cyth. testā ovato-cordatā, posticē obliquē productā, crassā, lærigatā, nitente, pallidē castaneā, violaceo pallidē radiatā, latere postico violaceo tincto, marginibus albis.*

THE CHESTNUT CY THEREA. Shell ovately heart-shaped, posteriorly obliquely produced, thick, smooth, shining, light chestnut, faintly rayed with violet, posterior side stained with violet, margins white.

LAMARCK, Anim. sans vert. vol. vi. p. 299.

Meretrix castanea, Deshayes.

Hab. China.

There is a variety of this species in which the chestnut colouring is replaced by pale yellow.

Species 7. (Mus. Cuming.)

CY THEREA PETECHIALIS. *Cyth. testā subtrigono-ovatā, fulvescente-lacteā, maculis undatis nigricante-fuscis concentricē nebulatā, latere postico obliquē curvato, lærigatā, epidermide corneā nitente indutā.*

THE PETECHIAL CY THEREA. Shell somewhat triangularly ovate, fulvous cream-colour, concentrically clouded with dark-brown zigzag marks, posterior side obliquely curved, smooth, covered with a shining horny epidermis.


LAMARCK, Anim. sans vert. vol. vi. p. 229.

Meretrix petechialis, Deshayes.

Hab. China.

Lamarek's French translation of the specific name given by him to this species is "petechiale," but I have sought in vain to discover its meaning, unless it refers to the shore of the province of Pe-te-hi, China, as one of the places of habitation.

January, 1864.



8.



9a



9.b.



10.

C Y T H E R E A .

PLATE III.

Species 8. (Mus. Cuming.)

CY THEREA LAMARCKII. *Cyth. testā trigono-ovatā, suboblongā, crassā, concentricē rugosā, flavescente-albā, violaceo pallidissimē latē radiatā, epidermide corneā fulvā partim indutā, lunulā rix nullā.*

LAMARCK'S CY THEREA. Shell triangularly ovate, rather oblong, thick, concentrically rugose, yellowish-white, very faintly broadly rayed with violet, partially covered with a fulvous horny epidermis, scarcely any lunule.

Meretrix Lamarckii, Deshayes, Cat. Conch. Brit. Mus. p. 39.
Hab. China Seas.

Of a characteristic, oblong-oval form, with the faint violet rays very much diffused over the shell.

Species 9. (Fig. *a, b*, Mus. Cuming.)

CY THEREA ZONARIA. *Cyth. testā trigono-ovatā, posticē subacuminato-productā, lærigatā, albida, castaneo profusē concentricē linearī-undatā, epidermide corneā fulvo indutā, areā ligamenti concavinsculā, interdum lividā.*

.. GIRDLED CY THEREA. Shell triangularly ovate, posteriorly somewhat acuminately produced, smooth, whitish, profusely concentrically linearly waved with chestnut, covered with a fulvous horny epidermis, ligamentary area slightly concave, sometimes livid.

LAMARCK, Anim. sans vert. vol. vi. p. 299.

Meretrix zonaria, Deshayes.

Hab. China and Japan.

This comes very near to certain forms of *C. petechialis*, and is possibly a variety of that species.

Species 10. (Mus. Cuming.)

CY THEREA IMPUDICA. *Cyth. testā subtrigono-ovalā, posticē arcuato-productā, crassā, lærigatā, nitente, aurantio-lactea, posticē livido-cinereā, circa umbones fuso punctatā, marginibus albis.*

THE IMMODEST CY THEREA. Shell somewhat triangularly ovate, arcuately produced on the posterior side, thick, smooth, shining, orange cream-colour, livid-ash posteriorly, dotted with brown about the umboes, margins white.

LAMARCK, Anim. sans vert. vol. vi. p. 299.

Cytherea meretrix, Lamarck.

Meretrix impudica, Deshayes.

Hab. China and the Philippine Islands.

The warm and comparatively colourless tint of this shell is no unimportant character of the species. The livid-ash ligamentary area presents a conspicuous contrast.



C Y T H E R E A.

PLATE IV.

Species 11. (Mus. Cuming.)

CY THEREA ARGENTINA. *Cytl. testā trigonā, medio tu-*
midā, ad basin rotundatā, utrinque subacuminatā, le-
vigatā, flavescente-albā, epidermide fulvā corneā in-
dutā.

THE SILVERY CY THEREA. Shell triangular, tumid in the middle, rounded at the base, somewhat acuminate on either side, smooth, yellowish-white, covered with a fulvous horny epidermis.

SOWERBY, Pro. Zool. Soc. 1835, p. 46.

Trigona argentina, Deshayes.

Hab. Central America.

The shell of this species is not unfrequently covered by a soft, velvety, chalk-white tissue, resembling a fungus.

Species 12. (Mus. Cuming.)

CY THEREA MORPHINA. *Cytl. testā subtrigono-cordatā,*
posticē acuminato-productā, crassā, lærusculā, nitente,
concentricē rugosā, cinereo-albā et aurantio-carneā,
violaceo radiatim tinctā, præsertim ad latus posticum.

THE MORPHINE CY THEREA. Shell somewhat triangularly heart-shaped, posteriorly acuminate produced, thick, nearly smooth, shining, concentrically rugose, ash-white and orange-flesh colour, radiately tinged with violet, especially on the posterior side.

LAMARCK, Anim. sans vert. vol. vi. p. 300.

Meretrix morphina, Deshayes.

Hab. China.

A solid delicately-coloured shell, of almost porcellanous substance.

Species 13. (Mus. Cuming.)

CY THEREA LYRATA. *Cytl. testā subtrigono-rotundatā rel-*
ovalā, ventricosā, solidā, sordidē lacteā, epidermide
tenui pelucido-corneā indutā, ad aream ligamenti
livido-nigrante, transversim concentricē corrugato-
liratā, liris latis, obtusis, irregularibus, interdum con-
fluentibus.

THE HARP-STRINGED CY THEREA. Shell somewhat triangularly rounded or ovate, ventricose, solid, dull cream-colour, covered by a thin transparent horny epidermis, livid-black at the ligamentary area, transversely concentrically wrinkle-ridged, ridges broad, obtuse, irregular, sometimes flowing together.

SOWERBY, Thes. Conch. vol. i. p. 621, pl. 129. f. 61.

Meretrix lyrata, Deshayes.

Hab. Philippine Islands; Cuming.

A very striking species, sculptured throughout with irregularly confluent, obtuse, concentric ridges.

Species 14. (Mus. Cuming.)

CY THEREA FORMOSA. *Cytl. testā oblongo-ovatā, posticē*
subobliquè productā, lacteā, pallidē violaceo concentricē
fasciatā, castaneo ab umboibus remotè biradiatā.

THE BEAUTIFUL CY THEREA. Shell oblong-ovate, posteriorly rather obliquely produced, cream-colour, concentrically banded with faint-violet, distantly two-rayed from the umboes with chestnut.


SOWERBY, Thes. Conch. vol. i. p. 620, pl. 129. f. 47.

Hab. —?

As in many other species of this genus, differing little from each other in form, the colouring is very characteristic.




15




16a




16b



17



18a



18c



18b

C Y T H E R E A.

PLATE V.

Species 15. (Mus. Cuming.)

CY THEREA LÆVIGATA. *Cyth. testā ovato-trigonā, subæquilaterali, tenuicnlā, lutescente-albā, violaceo tintā et concentricè fasciatā, umboibus violaceis, lærigatā, nictē.*

THE SMOOTH CY THEREA. Shell ovately triangular, nearly equilateral, rather thin, yellowish-white, tinged and concentrically banded with violet, umboes violet, smooth, shining.

Trigona lærigota, Gray, Cat. Cyth. Anal. vol. viii. p. 304.
Hab. —?

A delicate species, of nearly equilateral triangular form.

Species 16. (Fig. a, b, Mus. Cuming.)

CY THEREA TRIPLOA. *Cyth. testā trigono-cuneiformi, medio gibbosā, utrinque peculiariter latè compressā, sordidè flavescente-albā, nunc aurantio tintā, interdum cinereo-griseo radiatā, solidā, lærigatā, vel densè concentricè striatā, areis lateralibus latè excavatis.*

THE TRIPLE CY THEREA. Shell triangularly wedge-shaped, gibbous in the middle, peculiarly broadly compressed on each side, dull yellowish-white, sometimes tinged with orange, sometimes rayed with ash-grey, solid, smooth or densely concentrically striated, lateral areas broadly excavated.

Venus tripla, Linnaeus, Mant. p. 545.

Cytherea tripla, Lamarck.

Trigona tripla, Gray.

. Senegal.

A solid triangular shell, extremely varied in colour, wing the lunule and ligament areas pinched, as it were, into broad excavations.

Species 17. (Mus. Cuming.)

CY THEREA VENTRICOSA. *Cyth. testā cordato-trigonā, gib-*

bosissimā, griseo-albā, epidermide cornē fulvā partim indatā, latere antico immerso, latere poslico arcuato, latè abruptè truncato, areā ligamenti peramplā, excavatā.

THE VENTRICOSE CY THEREA. Shell cordately triangular, very gibbous, greyish-white, partially covered with a fulvous horny epidermis, anterior side immersed, posterior side arched, broadly abruptly truncated, ligamentary area very large, excavated.

Trigona ventricosa, Gray, Cat. Cyth. Anal. vol. viii. p. 304.
Cytherea ventricosa, Sowerby.

Hab. China.

This fine species is the most gibbous and abruptly truncated of the genus, and none of similar type approaches it in size.

Species 18. (Fig. a, b, c, Mus. Cuming.)

CY THEREA MACTROIDES. *Cyth. testā trionā, ventricosā, subæquilaterali, utrinque subcompressā, albida, rufescente-castaneo plus minus tintā et radiata, epidermide cornē fulvā interdum partim indatā, lærigatā.*

THE MACTRA-LIKE CY THEREA. Shell triangular, ventricose, nearly equilateral, rather compressed on each side, whitish, more or less rayed and stained with reddish-chestnut, sometimes partially covered with a fulvous horny epidermis, smooth.

Venus Mactroides, Born, Test. Mus. Vind.

Venus corbicula, Gmelin.

Cytherea corbicula, Lamarck.

Trigona corbicula, Gray.

Trigona Mactroides, Deshayes.

Hab. West Indies.

The colouring is mostly reddish-chestnut, though extremely variable in design. The lateral areas are rather broader than in *C. radiata* and other allied species.



19.a



19.b.



19.c



20.



21



22



23

C Y T H E R E A.

PLATE VI.

Species 19. (Fig. *a, b, c*, Mus. Cuming.)

CY THEREA OVUM. *Cytl. testā ovatā, solidissimā, gibbā, antieē cordatā, posticē subproductā, albidā, epidermide griseā vel rufescente-aurantiā indutā, latere postico intus extusque violaceo-tinecto.*

THE EGG CY THEREA. Shell ovate, very solid, gibbous, anteriorly heart-shaped, posteriorly a little produced, whitish, covered with a grey or reddish-orange epidermis, posterior side stained within and without with violet.

HANLEY, Pro. Zool. Soc. 1845, p. 21.

Meretrix ovum, Deshayes.

I'ob. Madras.

A solid, gibbous shell, varying in form, as in the specimens figured, from subglobose to oval, of a grey or reddish-orange colour, deeply stained with violet, chiefly in the interior, at the posterior extremity.

Species 20. (Mus. Cuming.)

CY THEREA LUSORIA. *Cytl. testā subtrigono-ovalā, solidā, subaequilaterā, albā, spadiceo-rufo nebulatā et fusciatā, infra epidermide pallidē fulvā, lavigatā, nitente, areā ligamenti lividā.*

THE SPORTIVE CY THEREA. Shell somewhat triangularly ovate, solid, nearly equilateral, white, clouded and banded with fawn-red, beneath a pale fulvous epidermis, smooth, shining, ligamentary area livid.

Meretrix lusoria, Rumphius, Amb. pl. 42. f. G.

Veans lusoria, Chemnitz.

Cytherea lusoria, Lamarck.

Hab. China Sea.

The fawn-red, or iron-mould colouring of this species, is mostly constant.

Species 21. (Mus. Cuming.)

CY THEREA VESTITA. *Cytl. testā ovalā, subtrigonā, modicē depressā, solidiusculā, fulvescente-lacteā, violaceo pallidē multiradiatā, epidermide intensē olivaceo-fuscā indutā, concentricē rulē striatā, areā ligamenti lividā.*

THE CLOTHED CY THEREA. Shell ovate, slightly trian-

gular, moderately depressed, rather solid, fulvous cream-colour, faintly many-rayed with violet, covered with a dark olive-brown epidermis, concentrically rudely striated, ligamentary area livid.

Hab. —?

I cannot refer this to any described species. Its colouring and epidermis are quite peculiar.

Species 22. (Mus. Cuming.)

CY THEREA STULTORUM. *Cytl. testā oblongo-ovalā, sub-Douaciformi, modicē depressā, albido-griseā, flavescente-roseo pallidē fasciatā, lavigatā, nitente, latere antico compressè rotundato, postico exilissimè flexuoso-angulato.*

THE FOOLS' CY THEREA. Shell oblong-ovate, somewhat Donax-shaped, moderately depressed, whitish-grey, faintly rayed with yellowish-rose, smooth, shining, anterior side compressly rounded, posterior very faintly flexuously angled.

Douac stultorum, Gray, Wood, Ind. Test. Supp. pl. 2. f. 2.
Trigona stultorum, Gray.

Hab. Indian Ocean.

A very delicately tinged, shining shell, of rather compressed growth.

Species 23. (Mus. Cuming.)

CY THEREA BICOLOR. *Cytl. testā acentē trigonā, depressē cuneiformi, albā, epidermide pallidē sulcā indutā, lavigatā, lunata et arei ligamenti peramplis, planis, areā violaceo-castaneā.*

THE TWO-COLOURED CY THEREA. Shell sharply triangular, depresso wedge-shaped, white, covered with a pale fulvous epidermis, smooth, lunule and ligamentary area very large, flat, area violet-chestnut.

Trigona bicolor, Gray, Anal. 1838, vol. viii. p. 304.

Cytherea bicolor, Hanley.

Hab. Senegal.

A sharply-angled shell, of a violet-chestnut colour on the largely-expanded ligamentary area, while all the rest of the shell is white.



C Y T H E R E A.

PLATE VII.

Species 24. (Mus. Cuming.)

CY THEREA DILLWYNI. *Cyth. testā trigono-cuneiformi, compressinsculā, subaequilaterali, spadiceo-fusca, vio-laceo-fusco pallidē radiatā et concentricē fasciatā, lunulā subampulā, oblongo-ovali.*

DILLWYN'S CY THEREA. Shell triangularly wedge-shaped, rather compressed, nearly equilateral, fawn-brown, faintly rayed and concentrically banded with violet-brown, lunule rather large, oblong-oval.

Cytherea mactroides, Sowerby (not of Lamarck), Thes. vol. i. p. 615. pl. 128. f. 36.

Trigona Dillwyni, Deshayes.

Hab. Indian Ocean.

A solid, Donax-shaped shell, neatly cross-banded and rayed with violet-brown.

Species 25. (Mus. Cuming.)

CY THEREA CASTA. *Cyth. testū ovatā, vix cordatā, sub-depressā, solidiusculā, glaucescente-griseā, pallidissimē radiatā, levigatā, nitente.*

THE CHASTE CY THEREA. Shell ovate, scarcely heart-shaped, somewhat depressed, rather solid, glaucous-grey, very faintly rayed, smooth, shining.

Venus casta, Chemnitz, Conch. C.b. vol. vi. p. 349. pl. 33. f. 346.

Cytherea casta, Lamarck.

Merytrix casta, Deshayes.

Hab. Ceylon.

The shining glaucous-grey colour, indistinctly rayed, is a characteristic feature of this species.

Species 26. (Fig. a, b, Mus. Cuming.)

CY THEREA ASTRICTA. *Cyth. testā subtrigono-cordatā, gibbā, solidā, posticē subproductā, olivaceo-spadiceā, rufo linctā, radiis duobus subdistantibus nigris aut fuscis notatā, levigatā, nitente, lunulā indistinctā.*

THE TIED CY THEREA. Shell somewhat triangularly heart-shaped, gibbous, solid, slightly produced posteriorly, olive-fawn colour, tinged with red, marked

with two rather distant black or brown rays, smooth, shining, lunule indistinct.

Hab. —?

Allied to *C. casta*, but of a more gibbous, heart-shaped form, distinctly produced at the posterior side. The two distant rays of black or brown are striking, but they may not be permanent.

Species 27. (Mus. Cuming.)

CY THEREA RADIATA. *Cyth. testā trigono-ovatā, posticē subproductā, solidā, gibbosā, fulvescente, rufescente-brunneō profusē radiatā et concentricē fasciatā, levigatā, haud nitente, antīcē subplicato-corrugatā, epidermide corueā partim indutā, lunulā amplā.*

THE RAYED CY THEREA. Shell triangularly ovate, rather produced posteriorly, solid, gibbous, fulvous, profusely rayed and concentrically banded with reddish-brown, smooth, not shining, obscurely plicately wrinkled on the anterior side, partially covered with a horny epidermis, lunule large.

SOWERBY, Pro. Zool. Soc. 1835, p. 23.

Venus Salangensis, D'Orbigny.

Cytherea gracilior, Sowerby.

Trigona Byronensis, Gray.

Trigona radiata, Deshayes.

Hab. Salango and Xipixapi, West Columbia (in sandy mud, at the depth of nine fathoms); Cuming.

Of a solid, triangular form, obscurely plicately wrinkled on the anterior side, profusely rayed with light red-brown, not shining, but partially covered with a horny epidermis.

Species 28. (Mus. Cuming.)

CY THEREA NUBILA. *Cyth. testā trigono-cordatā, gibba, præsertim ad latū posticum, fulcā, pallidē fusco profusē undatā et fasciatā, levigatā, nitente, areā posticā perampulā, latā, planā.*

THE CLOUDY CY THEREA. Shell triangularly heart-shaped, gibbous, especially on the posterior side, fulvous, profusely waved and banded with light

CY THEREA.—PLATE VII.

brown, smooth, shining, posterior area very large, broad, flat.

Hab. —?

Of a much more gibbous triangular form than *C. radiata*, the posterior area being peculiarly large, broad, and flat. The surface is moreover smooth and glossy.

Species 29. (Mus. Cuming.)

CY THEREA TRIGONA. *Cyth. testá abbreviato-trigoná, ad rumbones acuminatá, solidá, compressá, fulvá, rufo-*

fusco concentricè undulatá, latere postico subangulato, lunulá oblongá, indistinctá.

THE TRIANGULAR CY THEREA. Shell shortly triangular, acuminate at the umboes, solid, compressed, fulvous, concentrically waved with red-brown, posterior side rather angled, lunule oblong, indistinct.

DUNKER, MS. in Mus. Cuming.

Hab. —?

This appears to be a very distinct species, of solid, acuminate triangular growth. It is finely waved throughout with faint red lines.




30.



31.



32.



33a.



34a.



34b.



33b.

C Y T H E R E A.

PLATE VIII.

Species 30. (Mus. Cuming.)

CY THEREA VIRGINEA. *Cyth. testā oblongo-ovatā, latē triangulā, subaequilateralī, crassiusculā, cinerascente-albā, obscurē radiatā, lavigatā vel concentricē rugosā, latere postico subquadratim flexuoso-rostrato.*

THE VIRGIN CY THEREA. Shell oblong-ovate, widely triangular, nearly equilateral, rather thick, ash-white, obscurely rayed, smooth, or concentrically rugose, posterior side somewhat squarely flexuously beaked.

ADAMS and REEVE, Moll. Voy. Samarang, p. 78. pl. 24. f. 10.

Hab. Moluccas.

A very characteristic species, in which the posterior side is flexuously beaked.

Species 31. (Mus. Cuming.)

CY THEREA INTERMEDIA. *Cyth. testā latē ovatā, versus umbones trigonā, albida, violaceo interdum pallidē radiatā, epidermide fulvā partim indutā, solidā, lavigatā, concentricē rugosā, latere postico subtruncato.*

THE INTERMEDIATE CY THEREA. Shell broadly ovate, triangular towards the umboes, whitish, sometimes faintly rayed with violet, partially covered with a fulvous epidermis, solid, smooth, concentrically rugose, posterior side sometimes truncated.

SOWERBY, Thes. Conch. vol. i. p. 612. pl. 128. f. 39.
Trigona intermedius, Deshayes.

Hab. California.

Of a somewhat truncheately wedge-shaped form, faintly rayed with violet upon a dull cream-coloured ground.

Species 32. (Mus. Cuming.)

CY THEREA COMPRESSA. *Cyth. testā latē ovatā, trigonā, subcompressā, fulvescente-lacteā, fasciis rubescentibus plus minus profuse radiatā, concentricē densē striatā, subsericeā, latere postico subtruncato.*

THE COMPRESSED CY THEREA. Shell broadly ovate, triangular, rather compressed, fulvous cream-colour, more or less profusely rayed with reddish bands, concentrically closely striated, rather silky, posterior side slightly truncated.

SOWERBY, Thes. Conch. vol. i. p. 616. pl. 128. f. 33, 34.
Trigona compressa, Deshayes.

Hab. Cape of Good Hope.

This and the remaining species of the plate are charac-

terized by an unusually broadly ovate-triangular form, all more or less compressed in growth.

Species 33. (Fig. a, b, Mus. Cuming.)

CY THEREA UNDULATA. *Cyth. testā latē ovalā, subaequilateralī, versus umbones acutē trigonā, plano-compressā, fulvescente-lucteā, violaceo-rubente plus minus radiatā, radius interdum interruptē undulatis, lavigatā, concentricē densē striatā, ad latus anticum plicato-corrugatā, latere postico subabruptē abbreviato.*

THE WAVED CY THEREA. Shell broadly ovate, nearly equilateral, sharply triangular towards the umboes, flatly compressed, fulvous cream-colour, more or less rayed with violet-red, rays sometimes interruptedly waved, smooth, concentrically closely striated, plicately wrinkled on the anterior side, posterior side rather abruptly shortened.

SOWERBY, Thes. Conch. vol. i. p. 618. pl. 27. f. 12.

Trigona undulata, Deshayes.

Hab. Columbia.

It will be seen by the variety given at Fig. 33 b that the interruptedly waved rays after which the *C. undulata* is named, is by no means a constant character of the species.

Species 34. (Fig. a, b, Mus. Cuming.)

CY THEREA PLANULATA. *Cyth. testā latē ovatā, trigono-cuneiformi, subaequilateralī, plano-compressā, fulvescente lacteā, violaceo-rubente variè radiatā, epidermide cornea olivacea interdum indutā, concentricē densē striatā, ad latus anticum sēpe plicato-corrugatā, latere postico subabruptē abbreviato.*

THE FLAT CY THEREA. Shell broadly ovate, triangularly wedge-shaped, nearly equilateral, flatly compressed, fulvous cream-colour, variously rayed with violet-red, sometimes covered with an olive horny epidermis, concentrically closely striated, often plicately wrinkled on the anterior side, posterior side rather abruptly shortened.

BRODERIP and SOWERBY, Zool. Journ. vol. 5. p. 48.

Donax Lessoni, Deshayes.

Trigona planulata, Gray.

Venus planulata, D'Orbigny.

Hab. Central America.

This and the preceding species run so closely into each other that with a variety of specimens it is scarcely possible to define them.



35.



36.



37.



38.



39a.



39b.



40.

C Y T H E R E A.

PLATE IX.

Species 35. (Mus. Brit.)

CY THEREA EXILIS. *Cyth. testā trigono-ovatā, compressiusculā, ad umboes acentā, posticē subangulatā, albida, maculis purpureo-rufis sagittiformibus sparsim notatā, epidermide tenui corneā indutā, lavigatā, ad latus posticum plicato-striatā.*

THE DELICATE CY THEREA. Shell triangularly ovate, rather compressed, sharp at the umboes, posteriorly slightly angled, whitish, sparingly marked with purple-red arrow-headed spots, covered with a thin, horny epidermis, smooth, plicately striated on the posterior side.

Meretrix exilis, Deshayes, Cat. Conch. Brit. Mus. p. 40.

Hab. — ?

The arrow-headed dashes, with which this species is promiscuously sparingly marked, are quite peculiar.

Species 36. (Mus. Brit.)

CY THEREA POLITA. *Cyth. testā subtrigono-ovatā, subarcuatā, solidā, lavigatā, candidā, purpureo-carneo radiatā, nitente, latere postico subangulato, curvato.*

THE POLISHED CY THEREA. Shell somewhat triangularly ovate, slightly curved, solid, smooth, white, rayed with purple flesh-colour, shining, posterior side slightly angled, curved.

SOWERBY, Thes. Conch. vol. i. p. 618. pl. 127. f. 14.

Trigona polita, Deshayes.

Hab. Natal.

A very delicately tinted shell, smooth and shining.

Species 37. (Mus. Brit.)

CY THEREA AMBIGUA. *Cyth. testā subtrigono-ovatā, solidinuscūlā, depresso, albida, purpureo-carneo varie radiatā, lunulā et areā ligamenti aurantio-castaneis, lavigatā, nitente.*

THE DOUBTFUL CY THEREA. Shell somewhat triangularly ovate, rather solid, depressed, whitish, variously rayed with purple flesh-colour, lunule and ligamentary area orange-chestnut, smooth, shining.

Trigona ambigua, Deshayes, Cat. Conch. Brit. Mus. p. 47.

Hab. — ?

Very like the preceding species, although of a more triangular form. The orange-chestnut colouring of the lunule and ligamentary area is certainly peculiar.

Species 38. (Mus. Brit.)

CY THEREA CORPULENTA. *Cyth. testā subtrigono-orbiculari, solidā, cordato-tumidā, posticē obtusè angulatā, lavigatā, albā, lineis aurantio-castaneis undatis undique pictā, epidermide tenui corneā indutā, areā ligamenti lividā.*

THE STOUT CY THEREA. Shell somewhat triangularly orbicular, solid, cordately tumid, posteriorly obtusely angled, smooth, white, painted throughout with waved orange-chestnut lines, covered with a thin, horny epidermis, ligamentary area livid.

GRAY, Cat. Cyth. Anal. vol. viii. p. 303.

Meretrix corpulenta, Deshayes.

Hab. — ?

Described from a single specimen in the British Museum, which approximates rather closely to certain varieties of *C. zonaria*.

Species 39. (Fig. a, Mus. Brit. Fig. b, Mus. Cuming.)

CY THEREA HINDSII. *Cyth. testā trigonā, gibbosissimā, tenuiculā, ad umboes tumidā, utrinque impressā, fulvescente-lacteā, lineis aurantio-brunneis acentē undulatis conspicue concentricè fasciatā, versus marginem ustulato-brunneō plus minus tinctā, lavigatā.*

HINDS'S CY THEREA. Shell triangular, very gibbous, rather thin, tumid at the umboes, impressed on either side, fulvous cream-colour, conspicuously concentrically banded with sharply waved orange-brown lines, more or less stained with burnt-brown towards the margin, smooth.

HANLEY, Pro. Zool. Soc. 1844, p. 110.

Trigona Hindsii, Hanley.

Hab. Bay of Guayaquil; Hinds.

An excellent species, collected in the above-named locality by the assiduous conchologist after whom it is named.

CY THEREA.—PLATE IX.

Species 40. (Mus. Brit.)

CY THEREA FULMINATA. *Cyth. testā subrhombeā, aequilateralī, albā, lineis rufō-castaneis copiose undato-fasciatā, areā ligamenti castaneo tinctā, levigatā vel rugoso-striatā.*

THE LIGHTNING-MARKED CY THEREA. Shell somewhat rhomboid, equilateral, white, copiously wave-

banded with reddish-chestnut lines, ligamentary area stained with chestnut, smooth, or rugosely striated.

PHILIPPI, Abbild. und Besch. Conch. p. 198, pl. 3, f. 4.
Trigona fulminata, Deshayes.

Hab. —?

Of a peculiar rhomboidal form.



41.



42.



43.



44.



45.



46.



47^a.



47.b.



48.



49.

C Y T H E R E A.

PLATE X.

Species 41. (Mus. Cuming.)

CYTHERA TRIGONELLA. *Cyth. testá trigono-ovatá, compresiusculá, fulvá, livido-purpureo varie tinctá, lavigatá, latere postico flexuose truncata.*

THE TRIANGULAR CYTHERA. Shell triangularly ovate, rather compressed, fulvous, variously stained with livid-purple, smooth, posterior side flexuously truncated.

LAMARCK, Anim. sans vert. vol. vi. p. 307.

Trigona angulifera, Gray.

Hab. West Indies.

The livid-purple colouring of this little species is generally conspicuous on the umboes.

Species 42. (Mus. Cuming.)

CYTHERA INCERTA. *Cyth. testá trigono-ovatá, gibbosá, utrinque umbones subexcavatá, albá, lavigatá, nitente, rufescente-fusco obscurissimè maculatá.*

THE UNCERTAIN CYTHERA. Shell triangularly ovate, gibbous, slightly excavated on each side of the umboes, white, smooth, shining, very obscurely spotted with red-brown.

SOWERBY, Thes. Conch. vol. i. p. 617. pl. 128. f. 17.

Trigona incerta, Deshayes.

Hab. San Juan.

A small species of the gibbous form of the genus, with the lunule and ligament areas a little excavated.

Species 43. (Mus. Cuming.)

CYTHERA GOULDII. *Cyth. testá obliquè trigoná, posteriè subacuminatá, gibbosá, solidá, atbá, concentricè undique densè filoso-liratá.*

GOULD'S CYTHERA. Shell obliquely triangular, posteriorly rather acuminate, gibbous, solid, white, concentrically densely thread-ridged throughout.

Hab. Malacca.

A very distinct, conspicuously-sculptured shell, in Mr. Cuming's collection, with the above name appended to it in manuscript.

Species 44. (Mus. Cuming.)

CYTHERA NITIDULA. *Cyth. testá suboblongo-ovatá, obtuse trigoná, albida, fulvo-rubente undique undato-marmorata et strigata, lavigatá, nitente.*

THE SHINING CYTHERA. Shell rather oblong-ovate, obtusely triangular, whitish, streaked and wave-mottled throughout with fulvous-red, smooth, shining.

LAMARCK, Anim. sans vert. vol. vi. p. 305.

Trigona nitidula, Deshayes.

Hab. Mediterranean.

A prettily-painted species, nearly equilateral, smooth and shining.

Species 45. (Mus. Hanley.)

CYTHERA HANLEYANA. *Cyth. testá trigono-ovatá, gibbosá, posticè suboblique producta, albida, versus umbones livido-cærulescente tincta, rufescente-fusco versus marginem interruptè radiatá.*

HANLEY'S CYTHERA. Shell triangularly ovate, gibbous, posteriorly rather obliquely produced, whitish, tinged towards the umboes with livid-blue, interruptedly rayed towards the margin with red-brown.

SOWERBY, Thes. Conch. vol. i. p. 616. pl. 128. f. 23, 24.

Trigona Hanleyana, Deshayes.

Hab. —?

Distinguished by a characteristic, interruptedly rayed painting of red-brown on a blue-white ground, passing into a deep livid colour at the umboes.

Species 46. (Mus. Cuming.)

This proves to be a larger specimen of *C. Gouldii*, described at Species 43, and figured above it in the Plate.

Species 47. (Fig. a, b, Mus. Cuming.)

CYTHERA PHILIPPINARUM. *Cyth. testá cordato-ovatá, inaequilaterali, subveetricea, crassá, albida, violaceo-purpureo plus minus profuse linearis-undulata, rugulata et tincta, concentricè undique dense plicato-costata, lunula intense purpureo-rosea.*

THE PHILIPPINE CYTHERA. Shell cordately ovate, in-equalateral, slightly ventricose, thick, whitish, more

CY THEREA.—PLATE X.

or less profusely linearly waved, rayed and stained with violet-purple, concentrically plicately densely ribbed throughout, lunule deep purple-rose.

HANLEY, Pro. Zool. Soc. 1844, p. 110.

Meretrix Philippinarum, Deshayes.

Hab. Philippine Islands, Cuming.

A closely ribbed species, elaborately painted with fine zigzag linear waves of violet-purple, and sometimes stained throughout with that colour.

Species 48. (Mus. Cuming.)

CY THEREA FUMEA. *Cyth. testá subtrigono-ovatá, compresá, fumeo-spadiceá, ad umboes subacuminatá, posticè obliquè productá, fulvescente-spadiceá, concentricè perconspicuè plicato-costatá, costis subdistantibus, intersticiis excavatis.*

THE SMOKY CY THEREA. Shell somewhat triangularly ovate, compressed, smoky fawn-colour, livid at the umboes, then sharply concentrically waved with red-brown, smooth, shining.

Hab. —?

This little shell appears to differ in form from any other species, while it is characterized by a distinct style of painting.

Species 49. (Mus. Cuming.)

CY THEREA GRATA. *Cyth. testá cordato-ovatá, ad umboes subacuminatá, posticè obliquè productá, fulvescente-spadiceá, concentricè perconspicuè plicato-costatá, costis subdistantibus, intersticiis excavatis.*

THE PLEASING CY THEREA. Shell cordately ovate, rather acuminate at the umboes, obliquely produced posteriorly, fulvous fawn-colour, concentrically very conspicuously plicately ribbed, ribs rather distant, with the interstices excavated,

Meretrix grata, Deshayes, Pro. Zool. Soc. 1853.

Hab. China Sea.


Remarkable on account of its general external resemblance to an *Astarte*.

CYTHEREA.

	Plate.	Species		Plate.	Species
ambigua (<i>Trigona</i>), Deshayes	IX.	37	Lamarekii (<i>Meretrix</i>), Deshayes	III.	8
angulifera (<i>Trigona</i>), Gray	X.	41	laevigata (<i>Trigona</i>), Gray	V.	15
argentina, Sowerby	IV.	11	Lessoni (<i>Douar</i>), Deshayes	VIII.	34
astricta, Reeve	VII.	26	lusoria (<i>Meretrix</i>), Rumphius	VI.	20
bicolor (<i>Trigona</i>), Gray	VI.	23	lyrata, Sowerby	IV.	13
Byronensis (<i>Trigona</i>), Gray	VII.	27	Maetroides (<i>Venus</i>), Born	V.	18
casta (<i>Venus</i>), Chemnitz	VII.	25	meretrix, Lamarek	III	10
eastanea, Lamarck	II.	6	morphina, Lamarek	IV.	12
compressa, Sowerby	VIII.	32	nitidula, Lamarek	X.	44
corbicula (<i>Venus</i>), Born	V.	18	nubila, Reeve	VII.	28
corpulenta, Gray	IX.	38	ovum, Hanley	VI.	19
Crassatelloides (<i>Trigona</i>), Conrad	I.	3	petechialis, Lamarek	II.	7
damaoides (<i>Venus</i>), Gray	I.	4	Philippinarum, Hanley	X.	47
Dillwyni (<i>Trigona</i>), Deshayes	VII.	24	planulata, Broderip and Sowerby	VIII.	34
dolabella, Sowerby	I.	2	polita, Sowerby	IX.	36
exilis (<i>Meretrix</i>), Deshayes	IX.	35	ponderosa, Hanley	II.	5
formosa, Sowerby	IV.	14	radiata, Sowerby	VII.	27
fulminata, Philippi	IX.	40	Salangensis (<i>Venus</i>), D'Orbigny	VII.	27
fumea, Reeve	X.	48	stultorum (<i>Trigona</i>), Gray	VI.	22
Gouldii, Reeve	X.	43	trigona, Dunker	VII.	29
gracilior, Sowerby	VII.	27	trigonella, Lamarek	X.	41
graphica, Lamarck	I.	1	tripla (<i>Venus</i>), Linnaeus	V.	16
grata (<i>Meretrix</i>), Deshayes	X.	49	undulata, Sowerby	VIII.	33
Hanleyana, Sowerby	X.	45	ventricosa (<i>Trigona</i>), Gray	V.	17
Hindsii, Hanley	IX.	39	vestita, Reeve	VI.	21
impudica, Lamarck	III.	10	virginea, Adams and Reeve	VIII.	30
incerta, Sowerby	X.	42	zonaria, Lamarek	III.	9
intermedia, Sowerby	VIII.	31			

MONOGRAPH
OF THE GENUS
DIONE.


"Call me goddess, nymph, divine, and rare,
Precious, celestial."—*Shakespere*.




1a




1b




2.




3.



4a



4b.



5.

DIONE.

PLATE I.

Genus DIONE, Megerle.

Testa ovata, turgidiuscula, nunc subincerassata, saepissime concentricè costata, nunc tenuis, levigata, valde inaequilateralis, marginibus integris. Cardo in utrâque valvâ tridentatus, dentibus eis divergentibus. Impressio pallii posticè sinnatus, sinn subproundo.

Shell ovate, rather swollen, sometimes rather thick, and mostly concentrically ribbed, sometimes thin and smooth, very inequilateral, margins entire. Hinge in each valve three-toothed, teeth scarcely divergent. Impression of the mantle situated posteriorly, sinus rather deep.

Two very distinct forms, represented respectively by the Linnæan *Venus Erycina* and *V. lœta*, are referred to the genus *Dione*, even by those who have subdivided the family of *Veneridae* into more genera than it has been deemed advisable to adopt in this work. In the first of these groups the shells are large, sometimes smooth and highly polished, sometimes concentrically ribbed, and all more or less conspicuous for their brilliancy of colouring; in the second the shell is thin, inflated, and almost colourless. The most remarkable species are a group, of which *D. Veneris* is the type, inhabiting Central America and the West Indies, in which the concentric ridges are prolonged on the posterior angle of the shell into spines.

The general distribution of *Dione* is the same as that of *Venus*, though not half so numerous in species. Central America, the Philippine Islands, and Australia are the principal places of habitation. There are, however, three important species, *D. Hagenowi*, *florida*, and *umbonella*, from the Red Sea, and we have a remarkably fine species for so northern a latitude as our own in *D. Chione*. *D. umbonella* is quoted as being a native also of Brazil; and another instance is presented of the same species being found in both hemispheres in *D. squalida*, which was certainly collected by Mr. Cuming at the Philippine Islands, and is reported to inhabit California also.

Species 1. (Fig. 1 a, b, Mus. Cuming.)

DIONE FLORIDA. *Dion. testâ ovatâ, versus marginem compressiusculâ, lacteâ vel cinereâ, fulvo-fuscenscente plerumque minutâ undatâ et variegatâ, versus marginem nigricante-violaceo conspicue latè radiatâ, latere pos-*

tico obliquè producto, antico concentricè linearisulcato, sulcis subdistantibus, lunulâ nigricante-violaceâ, area ligamenti nitide strigata.

THE FLORID DIONE. Shell ovate, rather compressed towards the margin, cream-coloured, or ash, generally minutely waved and variegated with fulvous-brown, conspicuously broadly rayed towards the margin with dark-violet, posterior side obliquely produced, anterior side concentrically linearly grooved, grooves rather distant, lunule dark-violet, ligamentary area neatly streaked.

Cytherea florida, Lamarek, Anim. sans vert. vol. vi. p. 305.

Cytherea pulchra, Gray.

Chione pulchra, Gray.

Cytherea Erycinella, Lamarek.

Dione florida, Deshayes.

Hab. Red Sea.

This very pretty species is sculptured in a very characteristic manner on the anterior side by a series of rather distant concentric linear grooves, the posterior being smooth.

Species 2. (Mus. Cuming.)

DIONE FESTIVA. *Dion. testâ suboblongo-ovatâ, posticè subattenuatâ, solidiusculâ, fulvescente-spadiceâ, violaceo-roseo radiatâ et fasciatâ, latere antico concentricè linearisulcato.*

THE PRETTY DIONE. Shell rather oblong-ovate, posteriorly a little attenuated, rather solid, fulvous-fawn, rayed and banded with violet-pink, anterior side concentrically linearly grooved.

Cytherea festiva, Sowerby, Thes. Conch. vol. ii. p. 625. pl. 130. f. 72.

Dione festiva, Deshayes.

Hab. Eastern Seas.

Conspicuously distinguished by its pretty violet-rose colouring.

Species 3. (Mus. Cuming.)

DIONE ERYCINA. *Dion. testâ subcordato-ovatâ, valde inaequilaterali, crassisculâ, spadiceo-fuscâ, versus marginem aurantio-rufâ tinctâ, chocolato-fusco interru te radiatâ, concentricè undique linearisulcata.*

DIONE.—PLATE I.

THE ERYCINA DIONE. Shell somewhat cordately ovate, very inequilateral, rather thick, fawn-brown, stained with orange-red towards the margin, interruptedly rayed with chocolate-brown, concentrically linearly grooved throughout.

Venus Erycina, Linnæus, Syst. Nat. p. 1131.

Cytherea Erycina, Lamarek.

Chione Erycina, Gray.

Dione Erycina, Deshayes.

Hab. Indian Ocean.

This very common species is quite remarkable for the richness and brilliancy of its colouring.

Species 4. (Fig. *a, b*, Mus. Cuming.)

DIONE CHINENSIS. *Dion. testā subobliquā ovatā, posticē subattenuatā, depressā, spadiceā, violaceo-vel fuscō-cescente variè radiatā et fasciatā, præsertim versus marginem, latere antico concentricè linearī-sulcatō.*

THE CHINA DIONE. Shell rather obliquely ovate, posteriorly somewhat attenuated, depressed, fawn-coloured, variously rayed and banded with violet or light-brown, especially towards the margin, anterior side concentrically linearly grooved.

Venus Chinensis, Chenmitz, Coneh. vol. ii. p. 227. pl. 202. f. 1976.

Cytherea Sinensis, Sowerby.
Hab. China Seas; New Holland.

Very closely allied to *D. festiva*, but more depressed and more attenuated posteriorly.

Species 5. (Mus. Cuming.)

DIONE LILACINA. *Dion. testā subcordato-ovatā, valde ineq̄ilaterali, posticē subattenuatā, spadiceo-fuscō-cente, obscurè linearī-radiatā, versus marginem lila-ciūd, concentricè undique linearī-sulcatā, costis inter-medīis convexiusculis.*

THE LILAC DIONE. Shell somewhat cordately ovate, very inequilateral, posteriorly rather attenuated, fawn-brown, obscurely linearly rayed, stained towards the margin with lilac, concentrically linearly grooved throughout, the intermediate ribs rather convex.

Cytherea lilacina, Lamarek, Anim. sans vert. vol. vi. p. 303.

Chione lilacina, Gray.

Dione lilacina, Deshayes.

Hab. Australia.

Coloured towards the margin with a very peculiar tone of lilac.



DIONE.

PLATE II.

Species 6. (Fig. 6 a, Mus. Cuming.)

DIONE SEMISULCATA. *Dion. testá suboblongo-ovalá, gibbosíscula, lactea, violaceo-cinereo variegatá, latere postico subattenuato, lavigato, antico rotundato, concentricè regulariter sulcato.*

THE HALF-GROOVED DIONE. Shell rather oblong-ovate, somewhat gibbous, cream-coloured, variegated with violet-ash, posterior side rather attenuated, smooth, anterior rounded, concentrically regularly grooved.

Cytherea semisulcata, Sowerby, Thes. Conch. p. 624. pl. 131. f. 52.

Dione florida, var., Deshayes.

Hab. Australia.

M. Deshayes considers this species a variety of *D. florida*. If so, it is a very distinct form of it, more gibbous, and attenuated posteriorly.

It will be observed that the numbering of the figures in this plate, owing to some confusion of the lithographer and carelessness of the author, does not agree with the numbering of the species.

Species 7. (Fig. 6 b, Mus. Cuming.)

DIONE HAGENOWI. *Dion. testá subtrigono-cordatá, compressá, fulro-spadicea, fuscescente tincta et interrupte radiatá, concentricè plicato-sulcatá, latere postico subobliquo, violaceo marginali, lunulá et areá ligamenti castaneo strigatis.*

HAGENOW'S DIONE. Shell somewhat triangularly cordate, compressed, fulvous-fawn, tinged and interruptedly rayed with pale-brown, concentrically pliately grooved, posterior side rather oblique, edged with violet, lunule and ligamentary area streaked with chestnut.

Cytherea Hagenowi, Dunker, Zeitschr. für Malak. 1848, p. 184.

Cytherea spathulata, Sowerby.

Dione Hagenori, Gray.

Hab. Red Sea.

The most characteristic features in the painting of this

species are its dark-violet posterior margin and the bold chestnut zigzag streaks of the lunule and posterior area.

Species 8. (Fig. 6 c and 8 a, b, Mus. Cuming.)

DIONE IMPAR. *Dion. testú oblongo-oralí, crassiuscula, gibbosá, lavigatá, nitente, albida, violaceo-cinereo aut fusco varie radiatá, fasciatá et maculatá, latere postico lavigato, antico concentricè regulariter sulcato.*

THE UNEVEN DIONE. Shell oblong-oval, rather thick, gibbous, smooth, shining, whitish, variously rayed, banded, and spotted with violet-ash or brown, posterior side smooth, anterior concentrically regularly grooved.

Cytherea impar, Lamarck, Anim. sans vert. vol. vi. p. 304.

Dione impar, Deshayes.

Hab. Swan River.

This very beautifully painted porcelain-like shell belongs to the same well-marked type as *D. florida* and *semisulcata*, in which the shell is prominently concentrically grooved on the anterior side only.

Species 9. (Fig. 7 and 9, Mus. Cuming.)

DIONE COSTATA. *Dion. testá oratá, versus marginem subproductá, turgidá, albida vel spadicea, fuscescente interrupte exiliter radiatá, ad marginem sepe nigricante-violacea, concentricè undique densè plicato-costatá.*

THE RIBBED DIONE. Shell ovate, somewhat produced towards the margin, swollen, white or fawn-colour, faintly interruptedly rayed with light-brown, often dark-violet at the margin, concentrically densely pliately ribbed throughout.

Venus costata, Chevallier, Conch. Cab. vol. ii. pl. 226. pl. 202. f. 1975.

Dione costata, Deshayes.

Hab. Philippine Islands; Cuming. Swan River.

This very fine species partakes very much of the characters of the old *Dione Erycina*, but they are perfectly distinct.



10.



11a



11b



12.

D I O N E.

PLATE III.

Species 10. (Mus. Cuming.)

DIONE SQUALIDA. *Dion. testā subcordatā, posticē subattenuatā, infra epidermide cornē nitente sordidē fuscā, exilissimē radiatā, versus umbones marmoratā et undato-maculatā, levigatā.*

THE UGLY DIONE. Shell subcordate, slightly attenuated posteriorly, dull fuscous beneath a shining horny epidermis, very faintly radiated, mottled, and wave-blottedched towards the umboes, smooth.

Cytherea squalida, Sowerby, Pro. Zool. Soc. 1835, p. 23.

Venus squalida, D'Orbigny.

Cytherea biradiata, Sowerby.

Cytherea elegans, Koch.

Hab. California; Philippine Islands.

The mottled pattern of this otherwise dull species is very characteristic.

Species 11. (Fig. a, b, Mus. Cuming.)

DIONE MACULATA. *Dion. testā ovali, compressiusculā, posticē obliquē subproductā, spadiceo-albā, violaceo-fusco maculatā vel undatā, epidermide nitente cornē indutā.*

THE SPOTTED DIONE. Shell oval, rather compressed, posteriorly obliquely somewhat produced, fawn-white, blotched or waved with violet-brown, enveloped with a shining horny epidermis.

Pectunculus maculatus, Lister, Conch. p. 270. f. 106.

Le Jouret, Adamson.

Venus maculata, Linnaeus.

Cytherea maculata, Lamarck.

Dione maculata, Gray.

Hab. West Indies; Brazil; Pacific Ocean.

An ancient and well-known species, widely distributed in both hemispheres.

Species 12. (Mus. Cuming.)

DIONE AURANTIA. *Dion. testā globoso-ovatā, crassā, turgidā, carneo-aurantiā, epidermide levigatā cornē in-dutā, concentricē obscurē plicato-striatā, posticē vix angulatā.*

THE ORANGE DIONE. Shell globosely ovate, thick, swollen, flesh-tinged orange, covered with a smooth horny epidermis, concentrically obscurely plicately striated, scarcely angled on the posterior side.

Chione aurantia, Gray, Analyst, vol. viii. p. 305.


Cytherea aurantiaca, Sowerby.

Dione aurantia, Deshayes.


Hab. Aeapuleo, South America.

The delicate flesh-orange colouring of this fine species is reduced to a more or less dull horny hue by the overlying epidermis. It is of a bold globose form, indistinctly angled on the posterior side.

August, 1863.




13



14 a



14 b.



15

D I O N E.

PLATE IV.

Species 13. (Mus. nob.)

DIONE CHIONE. *Dion. testū ovali, solidū, compressiusculā, concentricē plicato-striatā, fulvo-spadiceā, fusco-rubente obscurè fasciatā et maculato-radiatā, margine dorsali subelevato.*

THE CHIONE DIONE. Shell oval, solid, rather compressed, concentrically plicately striated, fulvous-fawn, obscurely banded and radiately spotted with fulvous-red, dorsal margin rather elevated.

Venus Chione, Linnaeus, Syst. Nat. 12th edit. p. 1131.

Pectunculus glaber, Da Costa.

Cytherea Chione, Lamarck.

Chione vulgaris, Gray.

Hab. European and Mediterranean Seas.

This fine species is a very conspicuous object among British bivalves; it is, however, comparatively rare on our shores. It is abundant in the Mediterranean.

Species 14. (Fig. a, b, Mus. Cuming.)

DIONE GRATA. *Dion. testā transversē oratā, subpectinatā, flavescente-lacteā, violaceo-castaneo radiatā et undatā, concentricē crebricostatā, lunulā et areā ligamenti castaneo strigatis.*

THE PRETTY DIONE. Shell transversely ovate, somewhat pectinate, yellowish cream-colour, rayed and

waved with violet-chestnut, concentrically closely ribbed, lunule and ligamentary area streaked with chestnut.

Cytherea pectoralis, Sowerby (not of Lamarck), Thes. Conch. p. 625. pl. 130. f. 75.

Dione grata, Deshayes, Cat. Conch. Brit. Mus. p. 62.

Hab. Philippine Islands; Cuming.

A well-marked beautifully painted species.

Species 15. (Mus. Cuming.)

DIONE NOBILIS. *Dion. testā orbiculari-cordatā, crassā, candidā, epidermide tenui corneā nitente indutā, concentricē plicato-costatā, costis valde irregularibus, corrugatis, hic illuc intermissis, non parallelis.*

THE NOBLE DIONE. Shell orbicularly heart-shaped, thick, white, covered with a thin shining horny epidermis, concentrically plicately ribbed, ribs very irregular, wrinkled, here and there intermitted, not parallel.

Cytherea nobilis, Reeve, Pro. Zool. Soc. 1849, part ii. p. 126.

Dione nobilis, Deshayes.


Hab. California.

This fine shell has very much the form of an *Artemis*. The sculpture is a mass of concentric wrinkles.


August, 1863.



16a



16b



17



18



19

DIONE.

PLATE V.

Species 16. (Fig. *a*, *b*, Mus. Cuming.)

DIONE VULNERATA. *Dion. testā orbiculari, lacteā, aurantio-fusco tinctā tenuiradiatā et marmoratā, intensè violaceo concentricè lineatā, undique concentricè plicato-striatā.*

THE WOUNDED DIONE. Shell orbicular, cream-colour, tinged, finely rayed and mottled with orange-brown, concentrically lineated with deep violet, concentrically plicately striated throughout.

Cytherea vulnerata, Broderip, Pro. Zool. Soc. 1835, p. 46.

Dione vulnerata, Deshayes.

Hab. Real Llejos, Central America.

An exquisitely-painted shell, delicately pencilled with orange-brown and violet on a ground of warm cream-colour.

Species 17. (Mus. Cuming.)

DIONE GIGANTEA. *Dion. testā oblongo-ovali, compressā, spadiceo-albā, cæruleo-griseo radiatā, epidermide cornueā castaneā versus marginem indutā, lævigatā, vel concentricè plicato-striatā.*

THE GIGANTIC DIONE. Shell oblong-oval, compressed, fawn-white, rayed with blue-grey, covered towards the margin with a chestnut horny epidermis, smooth, or concentrically plicately striated.

Venus gigantea, Chemnitz, Conch. Cab. vol. x. p. 354. pl. 171. f. 1661.

Cytherea gigantea, Lamarck.

Dione gigantea, Deshayes.

Hab. Bay of Mexico.

This well-known species is one of the few shells in which the radiated painting is of a blue-grey colour. To-

wards the margin it is covered by a thin chestnut horny epidermis.

Species 18. (Mus. Cuming.)

DIONE RUTILA. *Dion. testā subquadrato-ovatā, crassā, fulvo-lacteā, rufo-fusco copiosè interruptè radiatā, undique concentricè plicato-striatā, lunulā et areā ligamenti conspicuè undulato-maculatis.*

THE RUDDY DIONE. Shell somewhat squarely ovate, thick, fulvous cream-colour, copiously interruptedly rayed with red-brown, concentrically plicately striated throughout, lunule and ligamentary area conspicuously wave-blotted.

Cytherea rutila, Sowerby, Thes. Conch. p. 743. pl. 163. f. 205.

Dione rutila, Deshayes.

Hab. —?

A compressed solid shell, conspicuously painted, especially about the margin of the lunule and ligamentary area.

Species 19. (Mus. Cuming.)

DIONE STRIATA. *Dion. testā globoso-cordatā, tumidā, flarescente, posticè fumeo-violaceo abruptè radiatā, concentricè densè plicato-striatā.*


THE STRIATED DIONE. Shell globosely heart-shaped, swollen, yellowish, posteriorly abruptly rayed with smoky violet, concentrically densely plicately striated.

Chione striata, Gray, Analyst, vol. viii. p. 306.


Dione citrina, var. Deshayes.

Hab. Philippine Islands; Cuming.

Compared with *D. citrina* this species is of a more globose cordate form, less solid, and more swollen.



21




20.a.




22




23



20.b.



24



20.c

DIONE.

PLATE VI.

Species 20. (Fig. *a, b, c*, Mus. Cuming.)

DIONE SEMILAMELLOSA. *Dion. testā subtrigono-ovatā, ventricosiusculā, albā, purpureo-violaceo plus minus tinctā, concentricè corrugato-liratā, liris ad latū antīcum lamellatis, per marginem posticum hic illic longissimè spinosis, spinis interdum biserialibus.*

THE HALF-LAMELLATED DIONE. Shell somewhat triangularly ovate, rather ventricose, white, more or less tinged with purple-violet, concentrically wrinkle-ridged, ridges lamellated on the anterior side, here and there very long-spined along the posterior margin, spines sometimes in two rows.

Cytherea lamellosa, Gaudichaud; Delessert, Recq. de Coq. pl. 19. f. 2.

Cytherea lupanaria, Lesson.

Venus lupanaria, D'Orbigny.

Dione lupanaria, Deshayes.

Hab. Central America.

For reasons which are obvious I think it better to abandon the foul name given to this lovely species by Lesson.

Species 21. (Mus. Cuming.)

DIONE BREVISPINATA. *Dion. testā subtrigono-ovatā, ventricosiusculā, albā, concentricè erecto-liratā, liris ad latū antīcum lamellatis, per marginem posticum brevispinatis.*

THE SHORT-SPINED DIONE. Shell somewhat triangularly ovate, rather ventricose, white, concentrically erectly ridged, ridges lamellated on the anterior side, short-spined along the posterior margin.

Cytherea brevispina, Sowerby, Thes. Conch. p. 632. pl. 132. f. 109.

Hab. California.

No actual spines are developed on the posterior margin of this species.

Species 22. (Mus. Cuming.)

DIONE MULTISPINOSA. *Dion. testā subtrigono-ovatā, ventricosiusculā, albā, purpurascente-violaceo interdum pallidè tinctā, concentricè lamellato-liratā, liris ad latū antīcum lamellatioribus, latere postico spinis tenuibus copiosè biserialiter munito.*

THE MANY-SPINED DIONE. Shell somewhat triangularly ovate, rather ventricose, white, sometimes faintly tinged with purple-violet, concentrically lamellately

ridged, ridges more lamellate on the anterior side, posterior furnished with a double row of numerous slender spines.

Cytherea multispinosa, Sowerby, Thes. Conch. p. 632. pl. 132. f. 112.

Hab. Peru.

A small form of this peculiar type, in which the concentric ridges are more thinly lamellate, and the spines more slender and numerous.

Species 23. (Mus. Cuming.)

DIONE VENERIS. *Dion. testā subtrigono-ovatā, ventricosiusculā, albā, purpurascente-violaceo plus minus tinctā, concentricè lamellato-liratā, liris ad latū antīcum lamellatioribus, per marginem posticum modicè spinatis.*

VENUS'S DIONE. Shell somewhat triangularly ovate, rather ventricose, white, more or less tinged with purple-violet, concentrically lamellately ridged, ridges more lamellate on the anterior side, moderately spined along the posterior margin.

Concha Veneris, D'Argenville, Conch. pl. 21. f. 1.

Venus Dione, Linnaeus.

Cytherea Dione, Lamarek.

Dione Veneris, Deshayes.

Hab. West Indies and Central America.

The concentric sculpture in this form of the group is not obtusely wrinkled, as in *D. semilamellosa*, but erect and lamellate throughout; and the spines are only moderately developed.

Species 24. (Mus. Cuming.)

DIONE EXSPINATA. *Dion. testā subtrigono-ovatā, ventricosiusculā, albā, purpurascente-violaceo plus minus tinctā, concentricè corrugato-liratā, liris ad latū antīcum lamellatis, per marginem posticum muricalato-tuberculatis.*

THE SPINELESS DIONE. Shell somewhat triangularly ovate, rather ventricose, white, more or less tinged with purple-violet, concentrically wrinkle-ridged, ridges lamellate on the anterior side, prickly tubercled along the posterior margin.

Hab. Central America.

If the other alleged species of this Plate are distinct from one another, so also is this. Excepting that the spines are wanting, it comes nearest to *D. semilamellosa*.




25.b




25.a.



26.



27.



28.a



28.b



29.

DIONE.

PLATE VII.

Species 25. (Fig. *a, b*, Mus. Cuming.)

DIONE CIRCINATA. *Dion. testā rotundato-ovatā, interdum subtrigonā, rufescente-purpureā, nonnunquam radiatā vel albā, liris erectis concentricis undique densē fasciulatā, liris ad latera laminatis.*

THE ROUNDED DIONE. Shell rotundately ovate, sometimes rather triangular, reddish-purple, sometimes rayed or white, densely corded throughout with erect concentric ridges, ridges laminate at the sides.

Venus circinata, Born, Test. Mus. p. 61, pl. 4. f. 8.

Venus rubra, Gmelin.

Venus Guineensis, Gmelin.

Cytherea Guineensis, Lamarck.

Cytherea alternata, Broderip.

Cytherea circinata, Sowerby.

Dione circinata, Deshayes.

Hab. Mazatlan.

Of a uniform reddish-purple throughout, rusty towards the margin.

Species 26. (Mus. Cuming.)

DIONE RUFESCENS. *Dion. testā trigono-ovatā, sublongitā, crassā, ventricosā, fulvescente-castaneā, ad lunulam et aream ligamenti rufescente, concentricè densē striatā, versus umbones levigatā.*

THE REDDISH DIONE. Shell triangularly ovate, rather oblong, thick, ventrieose, fulvous-chestnut, reddish at the lunule and ligamentary area, concentrically closely striated, smooth towards the umboes.

Dione rufescens, Deshayes, Pro. Zool. Soc. 1853, p. 2. Cat. Mus. Brit. Conch. p. 67.

Hab. Philippine Islands; Cuming.

A stout shell of a peculiar fulvous-chestnut colour, reddish at the lunule and ligamentary area, with somewhat of the aspect of a *Crassatella*.

Species 27. (Mus. Cuming.)

DIONE UMBONELLA. *Dion. testā cordato-ovatā, crassā, ventricosā, nitente violacea, versus umbones pallidiorē,*

levigatā, latere antico regulariter sulcato, sulcis linearibus, subdistantibus.

THE UMBONED DIONE. Shell cordately ovate, thick, ventrieose, shining-violet, paler towards the umboes, smooth, anterior side regularly grooved, grooves linear, rather distant.

Cytherea umbonella, Lamarck, Anim. sans vert. vol. vi. p. 319.

Cytherea nivea, Hanley.

Dione umbonella, Deshayes.

Hab. Red Sea; Brazil.

In this beautifully-tinted shell the anterior linear grooves are engraved with great regularity.

Fig. 28 *a, b* (Mus. Cuming).

The two shells here figured are the types of Mr. Broderip's *Cytherea alternata*. M. Deshayes refers them in his monograph of the *Veneridae*, in the British Museum Catalogue, to *D. circinata*, and on comparing them with that species the affinity is obvious. They represent a variety in which the reddish-purple is either rayed or altogether wanting.

Speciees 29. (Mus. Cuming.)

DIONE ROSEA. *Dion. testā obliquè cordatā, compressā, roseo-purpurascente, obscurè fasciatā, concentricè densē liratā, liris ad latus auticum hic illic lamellatis, ad posterium per marginem muricato-lamellatis, vic spinosis.*

THE ROSE DIONE. Shell obliquely heart-shaped, compressed, rose-purple, obscurely banded, concentrically closely ridged, ridges here and there lamellated on the anterior side, on the posterior prickly lamellated along the the margin, scarcely spined.


Cytherea rosea, Broderip and Sowerby, Zool. Journ. vol. iv. p. 364.

Cytherea lepida, Chenu.

Dione rosea, Deshayes.

Hab. Panama.


This species partakes largely of the characters of the group represented in the preceding Plate.




3t




3t a




3t b



32.



33 a



33 b.



34

DIONE.

PLATE VIII.

Species 30. (Mus. Cuming.)

DIONE AFFINIS. *Dion. testā ovato-oblongā, obtusē triangulari, subcompressā, albā, purpureo-violaceo radiatā, lunulā et areā ligamenti violaceis, concentricē liratā, liris subreflexis, hic illic corrugatis, latere postico flexuoso.*

THE ALLIED DIONE. Shell ovately oblong, obtusely triangular, rather compressed, white, rayed with purple-violet, lunule and ligamentary area violet, concentrically ridged, ridges slightly reflected, here and there wrinkled, posterior side flexuous.

Cytherea affinis, Broderip, Pro. Zool. Soc. 1825, p. 45.

Venus Paytensis, D'Orbigny.

Dione affinis, Deshayes.

Hab. Xipixapi, West Columbia; Cuming.

Scarcely distinguishable from the next species except in being rather more flexuously compressed on the posterior side, and in having the lunule and ligamentary area violet.

Species 31. (Fig. a, b, Mus. Cuming.)

DIONE CONCINNA. *Dion. testā oblongo-cordatā, subcompressā, castaneo-rufā vel albā, purpureo-roseo radiatā, concentricē liratā, liris subreflexis, hic illic corrugatis, latere postico subflexuoso.*

THE ELEGANT DIONE. Shell oblong heart-shaped, rather compressed, chestnut-red or white, rayed with purple rows, concentrically ridged, ridges slightly reflected, here and there wrinkled, posterior side a little flexuous.

Cytherea concinna, Sowerby, Pro. Zool. Soc. 1835, p. 23.

Dione concinna, Deshayes.

Hab. Panama, Mazatlan.

The details of sculpture are precisely the same in this as in the preceding species.

Species 32. (Mus. Cuming.)

DIONE PURPURATA. *Dion. testā rotundato-cordatā, crassā, violaceo-purpureā, pallidē fasciatā, marginibus albis, lacerigatā, concentricē irregulariter plicato-sulcatā.*

THE PURPLE DIONE. Shell rotundately heart-shaped,

thick, violet-purple, pale banded, margins white, smooth, concentrically irregularly pliately grooved.

LAMARCK, Anim. sans vert. vol. vi. p. 301.

Hab. —?

A fine solid porcelain-like shell, distinguished by a charming tone of colour.

Species 33. (Fig. a, b, Mus. Cuming.)

DIONE UNICOLOR. *Dion. testā trigono-cordatā, subcompressā, crassā, albida vel castaneo-spadicea, concentricē liratā, liris subobtusis, hic illic corrugatis, versus latus posticum interdum obsoletis.*

THE ONE-COLOURED DIONE. Shell triangularly heart-shaped, rather compressed, thick, whitish or chestnut-bay, concentrically ridged, ridges rather obtuse, here and there wrinkled, sometimes partially obsolete towards the posterior side.

Cytherea unicolor, Sowerby, Pro. Zool. Soc. 1835, p. 23.

Chione badia, Gray.

Cytherea ligula, Anton.

Dione unicolor, Deshayes.

Hab. West Columbia.

There is a tinge of the characteristic chestnut-bay colouring of this species in the pale variety of it.

Species 34. (Mus. Cuming.)

DIONE CYGNUS. *Dion. testā trigono-oblongā, subcordata, posticē obliquē productā, calcareo-albā, concentricē liratis, liris lamellatis, precipitē ad latus anticanum, crenulatis et exasperatis.*


THE SWAN DIONE. Shell triangularly oblong, somewhat heart-shaped, posteriorly obliquely produced, chalk-white, concentrically ridged, ridges lamellate, especially on the anterior side, crenulated, and broken.

Cytherea cygnus, Lamarck, Anim. sans vert. vol. vi. p. 328.

Dione cygnus, Deshayes.

Hab. —?


A dead-white shell, in which the ridges are finely crenulated, and more or less broken.



35 a.



36 a



36 b



35 b



37 a



37 b



38

DIONE.

PLATE IX.

Species 35. (Fig. *a*, *b*, Ms. Cuming.)

DIONE LÆTA. *Dion. testā subquadrato-ovalā, inflatā, fulvescente-albā, ferrugineo-rufō variè radialē vel aspersā, concentricè irregulariter striatā, latere antico brevissimo, contracto.*

THE CHEERFUL DIONE. Shell somewhat squarely ovate, inflated, fulvous white, variously rayed or sprinkled with rust-red, concentrically irregularly striated, anterior side very short, contracted.

Venus lœta, Linnaeus, Syst. Nat. p. 1132.

Venus affinis, Gmelin.

Cytherea lœta, Lamarek.

Venus lumescens, Blainville.

Cytherea cordiformis, Dunker.

Dione lœta, Deshayes.

Hab. Philippine Islands; Cuming.

The painting of this inflated squarely oblong species is extremely variable in pattern, but always of the same peculiar rust-red.

Species 36. (Fig. *a*, *b*, Mus. Cuming.)

DIONE KINGII. *Dion. testā ovatā, tenui, compressinsculā, fulvescente-albū, ferrugineo-rufō variè tinctā et radiatō, concentricè irregulariter striatā, latere antico brevissulo, rotundato.*

KING'S DIONE. Shell ovate, thin, rather compressed, fulvous white, variously stained and rayed with rust-red, concentrically irregularly striated, anterior side rather short, rounded.

Venus Kingii, Gray, Ind. Test. Supp. pl. 2. f. 9.

Cytherea Kingii, Sowerby.

Dione Kingii, Deshayes.

Hab. Island of Nicobar.

Of a more compressly oval form than *D. lœta*, less abruptly shortened on the anterior side.

Species 37. (Fig. *a*, *b*, Mus. Cuming.)

DIONE INFLATA. *Dion. testā subquadrato-orbiculari, tenuiculā, inflatā, fulvescente-lacteā, ferrugineo-rufō, variè tinctā vel radialē aspersā, concentricè irregulariter striatā, lateribus subæqualibus.*

THE INFLATED DIONE. Shell somewhat squarely orbicular, rather thin, inflated, fulvous cream-colour, variously stained or radiately sprinkled with rust-red, concentrically irregularly striated, sides somewhat equal.

Cytherea inflata, Sowerby, Thes. Conch. vol. i. p. 637. pl. 133. f. 127, 128.

Dione inflata, Deshayes.

Hab. Philippine and Molucca Islands, and Port Curtis, Australia.

This species is also closely allied to *D. lœta*; an orbicular form of it, in which the sides are more equal.

Species 38. (Mus. Cuming.)

DIONE PRORA. *Dion. testā oxato-trapezoidalī, solidiusculā, albū vel lacteā, immaculatā, epidermide tenui sericeā versus marginem, levigatā, lateribus ad marginem flexuoso-tumidis, lunulā subampullā, excavato-curvatā.*

THE PROW DIONE. Shell ovately trapezoidal, rather solid, white or cream-colour, unspotted, with a slight silky epidermis towards the margin, smooth, sides flexuously swollen at the margin, lunule rather large, excavately curved.


Cytherea prora, Conrad, Ann. Acad. Nat. Sci. Philadelphia, vol. vii. p. 253. pl. 19. f. 18.

Hab. Cape St. Lucas, Xanthis, California; Carpenter.

A colourless shell, distinguished by a peculiar trapezoid form.



40



39




41




42



43



44



45

D I O N E.

PLATE X.

Species 39. (Mus. Cuming.)

DIONE ALBIDA. *Dion. testā oblongo-cuneiformi, valde inaequilateralī, albida, versus umbones ferrugineo tinctā, concentricē rudē striatā, latere antico brevissimo.*

THE WHITE DIONE. Shell oblong-wedge-shaped, very inequilateral, whitish, tinged with rust towards the umboes, concentrically rudely striated, anterior side very short.

Pectunculus albidus, Lister, Conch. pl. 273. f. 109.

Venus albida, Gmelin.

Cytherea albida, Hanley.

Dione albida, Sowerby.

Hab. Jamaica?

A rather solid compressed shell, of wedge-like form, with scarcely any tinge of colour.

Species 40. (Mus. Cuming.)

DIONE CONVEXA. *Dion. testā ovato-cuneiformi, compressulā, sordidē albā, versus umbones ferrugineo pallide tinctā, concentricē subrugoso-striatā.*

THE CONVEX DIONE. Shell ovately wedge-shaped, rather compressed, dead-white, faintly tinged towards the umboes with rust, concentrically rather roughly striated.

Cytherea convexa, Say, Journ. Acad. Nat. Sci. Philadelphia, vol. iv. pl. 12. f. 3.

Cytherea Sayana, Conrad.

Dione convexa, Deshayes.

Hab. Bay of Massachusetts; Rhode Island.

A dead-white shell, of a more ovate form than that of the preceding species, more coarsely striated.

Species 41. (Mus. Cuming.)

DIONE ROSTRATA. *Dion. obliquē cuneiformi, solidiusculā, compressulā, sordidē albā, densē concentricē striatā, latere postico arcuato-declivē, antico brevissimo, abrupte truncato, umbonibus rostratis.*

THE BEAKED DIONE. Shell obliquely wedge-shaped, rather solid, compressed, dull white, densely concentrically striated, posterior side arcuately sloping, anterior very short, abruptly truncated, with the umboes beaked.

Cytherea rostrata, Koch, Phil. Abbild. Conch. p. 150.

pl. 1. f. 3.

Cytherea rostrata, Sowerby.

Tenus Tehuelcha, D'Orbigny.

Dione rostrata, Deshayes.

Hab. Brazil.

Distinguished from the preceding species by its very abrupt wedge-like form, the anterior side being so short as to impart a beaked character to the umboes.

Species 42. (Mus. Cuming.)

DIONE ALABASTRUM. *Dion. testā orato-cordatā, tenuiculā, inflatā, candidā, ferrugineo vix tinctā, concentricē irregulariter corrugato-striatā, lunulā et umbo-nibus subimmersis.*

THE ALABASTER DIONE. Shell ovately cordate, rather thin, inflated, white, scarcely tinged with rust, concentrically irregularly wrinkle-striated, lunule and umboes rather immersed.

Hab. —?

An elegantly-formed inflated species, with the lunule and umboes more cordately immersed than usual.

Species 43. (Mus. Cuming.)

DIONE CITRINA. *Dion. testā subtrigono-oratā, solidā, citrino-luteā, purpureo-violaceo posticē tinctā, umbo-nibus carneis, concentricē irregulariter corrugato-striatā.*

THE CITRON DIONE. Shell somewhat triangularly ovate, solid, lemon-yellow, tinged with purple-violet posteriorly, umboes flesh-tinged, concentrically irregularly wrinkle-striated.

Cytherea citrina, Lamarek, Anim. sans vert. vol. vi. p. 306.

Dione citrina, Deshayes.

Hab. Australia and Philippine Islands.

Of a peculiarly delicate lemon colour, stained about the posterior end with dark purple-violet.

Species 44. (Mus. Cuming.)

DIONE STRIATA. *Dion. testā orbiculari-cuneiformi, solidā,*

DIONE.—PLATE X.

gibbosiuscula, pallide spadicea, ad umboes roseo tincta, concentricè dense sericeo-striata, margine postico obtuse tumido-angulato, umbonibus incurvis.

THE STRIATED DIONE. Shell orbicularly wedge-shaped, solid, rather gibbous, pale fawn-colour, tinged with rose at the umboes, concentrically densely silk-striated, posterior margin obtusely tumidly angled, umboes incurved.

Chione striata, Gray, Anal. vol. viii. p. 306.

Dione citrina, var. Deshayes.

Hab. Philippine Islands; Cuming.

A comparison of the above description with that of the preceding species will show that they differ in detail throughout.

Fig. 45. (Mus. Smithsonian Institute of North America.)

This shell, from Cape St. Lucas, Xantus, California, kindly placed in my hands by Mr. Carpenter, as being probably undescribed, proves to be the *Dione prora* (*Cytherea prora*, Conrad) of our preceding Plate.



46




47.




48.




49 a



50




51



49 b.



52



53



54

DIONE.

PLATE XI.

Fig. 46. (Mus. Cuming.)

This little shell appears in Mr. Cuming's collection with the name *Cytherea consanguinea*, C. B. Adams (Panama Shells, p. 273). It is apparently a small specimen of a variety of *C. laeta*.

Species 47. (Mus. Cuming.)

DIONE AUSTRALICA. *Dion. testā ovotā, subæquilateralē, tenui, compressiusculā, albida, ferrugineo-rufo copiosè radialē, versus umbones lividis, concentricē densē tenuē lirato-striatā.*

THE AUSTRALIAN DIONE. Shell ovate, nearly equilateral, thin, rather compressed, white, copiously rayed with rust-red, livid toward the umboes, concentrically densely finely ridge-striated.

Hab. Torres' Straits, North Australia.

A delicate and very characteristic species, nearly equilateral in form.

Species 48. (Mus. Cuming.)

DIONE PELLUCIDA. *Dion. testā ocatā, subæquilateralē, tenuiculā, fulvescente-albā, ferrugineo-rufo radiatim maculatā, lavigatā, vel tenuē concentricē striatā.*

THE PELLUCID DIONE. Shell ovate, nearly equilateral, rather thin, fulvous white, radiately blotched with rust-red, smooth, or finely concentrically striated.

Cytherea pellucida, Lamarck, Anim. s. vert. vol. vi. p. 313.

Dione pellucida, Deshayes.

Hab. —?

This, and the shell which I have called *D. Australica*, have a strong general resemblance, but the umboes are more central and tumid in *D. pellucida*, which is altogether of a different character and composition.

Species 49. (Fig. a, b, Mus. Cuming.)

DIONE USTULATA. *Dion. testā globoso-cunciformi, solidā, pallidē ustulato-fuscā, purpureo-violaceo posticē et versus marginem ventralem tintatā, concentricē subrude striatā, lunulā amplā, superficiariā.*

THE SCORCHED DIONE. Shell globosely wedge-shaped, solid, pale burnt-brown, stained posteriorly and

towards the ventral margin with purple-violet, concentrically rather rudely striated, lunule large, superficial.

Hab. Swan River and New Caledonia.

Allied in general aspect to *D. citrina*, but of a more gibbous, wedge-like form.

Species 50. (Mus. Cuming.)

DIONE VIRGO. *Dion. testā subtrigono-ovatā, subcordatā, solidā, sordidē albā, ad umbones saepe roseo-tinctā, concentricē rugoso-striatā.*

THE VIRGIN DIONE. Shell somewhat triangularly ovate, solid, dull white, often tinged with rose at the umboes, concentrically roughly striated.

Chione virgo, Gray, Cat. Cyth. Anal. vol. viii. p. 306.

Dione virgo, Deshayes.

Hab. Java.

Of simple character, mostly distinguished, however, by a blush of rose about the umboes, which is more conspicuous in the interior of the shell.

Species 51. (Mus. Cuming.)

DIONE TORRESICA. *Dion. testā globoso-cordata, tenui, inflatā, fulvescente-albā, ferrugineo-rufo radiatim aspersā, circa marginem posticam purpureo-violaceo tintatā, lunulā et areā ligamenti purpureo-violaceis.*

THE TORRES DIONE. Shell globosely cordate, thin, inflated, fulvous white, radiately sprinkled with rust-red, tinged with purple-violet about the posterior margin, lunule and ligamentary area purple-violet.

Hab. Torres' Straits, North Australia.

There is a close resemblance between the general detail of character of this and the species from the same locality which I have named *D. Australica*, but these are widely different in form.

Species 52. (Mus. Cuming.)

DIONE SUBPELLUCIDA. *Dion. testā suboblique ovatā, compressiusculā, tenui, fulvescente-albā, ferrugineo-*

rufa copiosè radiatâ, concentricè densè striatâ, margine ventrali, lunulâ et areâ ligamenti ustulato-purpureis.

THE SUBPELLUCID DIONE. Shell somewhat obliquely ovate, rather compressed, thin, fulvous white, copiously rayed with rust-red, concentrically densely striated, ventral margin, lunule, and ligamentary area scorched purple.

Cytherea subpellucida, Sowerby, Thes. Conch. vol. i. p. 639. pl. 133. f. 136.

Dione subpellucida, Deshayes.

Hab. Island of Luzon, Philippines; Cuming.

Rather broad in dimensions from the umboes to the ventral margin, which, together with the lunule and ligamentary area, are strongly tinged with scorched livid purple.

Species 53. (Mus. Cuming.)

DIONE CROCEA. *Dion. testâ globoso-subcordatâ, tenui-ctâ, eximiè croceâ, aut vitellinâ, versus umbones roseo-tinctâ, lævigatâ, nitente, concentricè densè tenui-striatâ.*

THE SAFFRON DIONE. Shell globosely subcordate, rather

thin, delicate saffron, or yolk, tinged with rose towards the umboes, smooth, shining, concentrically densely finely striated.

DESHAYES, MS. in Mus. Cuming.

Hab. Philippine Islands; Cuming.

Coloured both within and without with a rose-tinged saffron, inclining to a warm yolk-yellow.


Species 54. (Mus. Cuming.)

DIONE ERUBESCENS. *Dion. testâ globoso-cordatâ, tenui-inflatâ, lacteâ, versus umbones roseo-tinctâ, nitente, concentricè densè lirato-striatâ, lunulâ amplâ, indistinctâ.*

THE BLUSHING DIONE. Shell globosely cordate, thin, inflated, cream colour, tinged with rose towards the umboes, shining, concentrically densely ridge-striated, lunule large, indistinct.

Hab. Ceylon; E. L. Layard.

Somewhat like the preceding species in general aspect, but more heart-shaped, while, on examination with the lens, the surface will be found to be sculptured concentrically with distinct ridges.



DIONE.

PLATE XII.

Species 55. (Mus. Cuming.)

DIONE ANGULIFERA. *Dion. oblongo-ovali, posticè angulato-productā, fulvescente-albā, maculis angularibus griseo-purpureis radiatā, concentricè rugoso-striatā.*

THE ANGULAR DIONE. Shell oblong-oval, angularly produced on the posterior side, fulvous white, rayed with angular greyish-purple spots, concentrically roughly striated.

Cytherea angulifera, Sowerby, Thes. Conch. vol. i. p. 635. pl. 136. f. 189.

Dione angulifera, Deshayes.

Hab. —?

This small species is chiefly distinguished by its angularly prolonged growth on the posterior side.

Species 56. (Mus. Cuming.)

DIONE PHASIANELLA. *Dion. testā oblongo-ovatā, solidisculā, versus marginem compressā, fulvescente-albā, violaceo-castaneo variè tinctā, undulatā et biradiatā, lunulā et areā ligamenti violaceo-castaneis, concentricè lineari-sulcatā, sulcis distinctis, aequidistantibus.*

THE LITTLE PHEASANT DIONE. Shell oblong-ovate, rather solid, compressed towards the margin, fulvous white, variously stained, waved and two-rayed with violet-chestnut, lunule and ligamentary area violet-chestnut, concentrically linearly grooved, grooves distinct, equidistant.

Venus phasianella, Deshayes, Pro. Zool. Soc. 1853, p. 2.

Dione phasianella, Deshayes.

Hab. —?

Characterized by a concentric sculpture of equidistant, linear grooves, rather widely separated from each other.

Species 57. (Mus. Cuming.)

DIONE AFRICANA. *Dion. subelongato-ovali, posticè acuminato-productā, fulvescente-albā, nigricante-griseo concentricè irregulariter undulatā, levigatā, margine postico dense corrugato-sulcata.*

THE AFRICAN DIONE. Shell somewhat elongately oval, acuminately produced on the posterior side, fulvous white, concentrically irregularly waved with dark grey, smooth, posterior margin densely wrinkled-grooved.

Cytherea Africana, Philippi, MS. in Mus. Cuming.

Hab. Australia?

The posterior side of this shell is peculiar, not only in being acuminately produced, but in being conspicuously wrinkled along the margin with close-set grooves.

Species 58. (Mus. Cuming.)

DIONE PALLESCENS. *Dion. testā subcordato-ovatā, sor-didè albā, concentricè densè striatā, striis numerosissi-mis, filiformibus.*

THE PALE DIONE. Shell subcordately ovate, dull white, concentrically densely striated, striae very numerous and thread-like.

Cytherea pallescens, Sowerby, Pro. Zool. Soc. 1835, p. 47.

Dione pallescens, Deshayes.

Hab. Isle of Annaa; Cuming.

A shell of simple form, without any indication of colour, sculptured with very numerous, fine, thread-like striae.

Species 59. (Mus. Cuming.)

DIONE PIPERITA. *Dion. testā subelongato-ovatā, solidisculā, fulvescente-lucteā, punctis maculisque hieroglyphicis nigris promiscue aspersā, levigatā, latere pos-tico concentricè lineari-sulcata, sulcis aequidistantibus.*

THE PEPPERED DIONE. Shell somewhat elongately ovate, rather solid, fulvous cream-colour, promiscuously sprinkled with black dots and hieroglyphical marks, smooth, posterior side concentrically linearly grooved, grooves equidistant.

Cytherea piperita, Sowerby, Thes. Conch. vol. i. p. 625. pl. 136. f. 175.

Dione piperita, Deshayes.

Hab. Island of Burias, Philippines.

Sprinkled in a very characteristic manner with black dots and small hieroglyphic marks.

Species 60. (Mus. Cuming.)

DIONE MULTISTRIATA. *Dion. testā oblongo-ovatā, solidā, fulvo-spadiceā, purpureo-cinereo radiatā, concentricè irregulariter rugosè multistriatā.*

THE MANY-STRIATED DIONE. Shell oblong-ovate, solid,

fulvous drab, rayed with purple-ash, concentrically irregularly roughly many-striated.

Cytherea multistriata, Sowerby, Thes. Conch. vol. i. p. 628. pl. 136. f. 177.

Dione multistriata, Deshayes.

Hab. New Zealand.

Rather variable in character, but generally more or less rayed with purple-ash upon a fulvous bay ground.

Species 61. (Mus. Cuming.)

DIONE TORTUOSA. *Dion. testā subangustē obliquē cordatā, solidā, eburneā, ad umbones fulvescente tinctā, latere antico breviusculo, rotundato, postico flexuose attenuato-producto, concentricē obtusè corrugato-liratā, tiris hic illic confluentibus.*

THE TWISTED DIONE. Shell rather narrowly obliquely cordate, solid, ivory-white, tinged with fulvous at the umboes, anterior side rather short, rounded, posterior flexuously attenuately produced, concentrically obtusely wrinkle-ridged, ridges here and there flowing together.

Cytherea tortuosa, Broderip, Pro. Zool. Soc. 1835, p. 45.

Venus tortuosa, D'Orbigny.

Dione tortuosa, Deshayes.

Hab. Nipixapi, Central America; Cuming.

The confluent ridges of this attenuately cordate species and the interstices are of a uniform delicate ivory-white.

Species 62. (Fig. *a, b, c*, Mus. Cuming.)

DIONE PANNOSA. *Dion. testā subtrigono-ovatā, solidiusculā, fulvā, violaceo-fusco varie radiatā, tinctā vel reticulatā, levigatā, vel concentricē subrugoso-striatā, sub teste radiatim minutissimè striatā.*

THE RAGGED DIONE. Shell somewhat triangularly ovate, rather solid, fulvous, variously stained, rayed, or reticulated with violet-brown, smooth, or concentrically rather roughly striated, beneath the lens radiately very minutely striated.

Cytherea pannosa, Sowerby, Pro. Zool. Soc. 1835, p. 47.

Cytherea lutea, Koch.

Venus pannosa, D'Orbigny.

Dione pannosa, Deshayes.

Callista puella, Carpenter.

Hab. Chili, Peru, Mazatlan.

The colouring of this species, though extremely variable, is mostly disposed in irregular mottled reticulations, more especially about the lunule and ligamentary area.

D I O N E .

Plate.	Species.	Plate.	Species.
affinis (<i>Cytherea</i>), Broderip	VIII. 30	Inpanaria (<i>Cytherea</i>), Lesson	VI. 20
affinis (<i>Venus</i>), Gmelin	IX. 35	Iteata (<i>Cytherea</i>), Koch	XII. 62
Africana (<i>Cytherea</i>), Philippi	XII. 57	maeulata (<i>Pectunculus</i>), Lister	III. 11
alabastrum, Reeve	X. 42	multispinosa (<i>Cytherea</i>), Sowerby	VI. 22
albida (<i>Pectunculus</i>), Lister	X. 39	multistriata (<i>Cytherea</i>), Sowerby	XII. 60
alternata (<i>Cytherea</i>), Broderip	VII. 25	nivea (<i>Cytherea</i>), Hanley	VII. 27
angulifera (<i>Cytherea</i>), Sowerby	XII. 55	nobilis (<i>Cytherea</i>), Reeve	IV. 15
aurantia (<i>Chione</i>), Gray	III. 12	Pacifica (<i>Venus</i>), Dillwyn	I. 5
Australica, Reeve	XI. 47	pallidescens (<i>Cytherea</i>), Sowerby	XII. 58
badia (<i>Chione</i>), Gray	VIII. 33	pannosa (<i>Cytherea</i>), Sowerby	XII. 62
biradiata (<i>Cytherea</i>), Sowerby	III. 10	Paytensis (<i>Cytherea</i>), D'Orbigny	VIII. 30
brevispinata (<i>Cytherea</i>), Sowerby	VI. 21	pectoralis (<i>Cytherea</i>), Sowerby	IV. 14
Chiensis (<i>Venus</i>), Chemnitz	I. 5	pellucida (<i>Cytherea</i>), Lamarck	XI. 48
Chione (<i>Venus</i>), Linnaeus	IV. 13	phasianella (<i>Venus</i>), Deshayes	XII. 56
circinata (<i>Venus</i>), Born	VII. 25	piperita (<i>Cytherea</i>), Sowerby	XII. 59
citrina (<i>Cytherea</i>), Lamarck	X. 43	prora (<i>Cytherea</i>), Conrad	IX. 38
coeninna (<i>Cytherea</i>), Sowerby	VIII. 31	puella (<i>Callista</i>), Carpenter	XII. 62
consanguinea (<i>Cyth.</i>), C. B. Adams	XI. 46	pulchra (<i>Cytherea</i>), Gray	I. 1
convexa (<i>Cytherea</i>), Say	X. 40	purpurata, Lamarck	VIII. 32
cordiformis (<i>Cytherea</i>), Dunker	IX. 35	rosea (<i>Cyth.</i>), Brod. and Sowerby	VII. 29
costata (<i>Venus</i>), Chemnitz	II. 9	rostrata (<i>Cytherea</i>), Koch	X. 41
erocea, Deshayes	XI. 53	rubra (<i>Venus</i>), Gmelin	VII. 25
eygnus (<i>Cytherea</i>), Lamarck	VIII. 34	rufescens, Deshayes	VII. 26
Dione (<i>Venus</i>), Linnaeus	VI. 23	rutila (<i>Cytherea</i>), Sowerby	V. 18
elegans (<i>Cytherea</i>), Koch	III. 10	Sayana (<i>Cytherea</i>), Conrad	X. 40
erubescens, Reeve	XL 54	semilamellosa (<i>Cyth.</i>), Gaudichaud	VI. 20
Erycinia (<i>Venus</i>), Favanne	I. 3	semisuleata (<i>Cytherea</i>), Sowerby	II. 6
Erycinella (<i>Cytherea</i>), Lamarck	I. 1	Sinensis (<i>Cytherea</i>), Sowerby	I. 5
exspinata, Reeve	VI. 24	spathulata (<i>Cytherea</i>), Sowerby	II. 7
festiva (<i>Cytherea</i>), Sowerby	I. 2	squalida (<i>Cytherea</i>), Sowerby	III. 10
florida (<i>Cytherea</i>), Lamarck	I. 1	striata (<i>Chione</i>), Gray	X. 44
gigantea (<i>Venus</i>), Chemnitz	V. 17	subpellueida (<i>Cytherea</i>), Sowerby	XI. 52
glaber (<i>Pectunculus</i>), Da Costa	IV. 13	Tehuelcha (<i>Venus</i>), D'Orbigny	X. 41
grata, Deshayes	IV. 14	Torresica, Reeve	XI. 51
Guineensis (<i>Venus</i>), Gmelin	VII. 25	tortuosa (<i>Cytherea</i>), Broderip	XII. 61
Hagenowi (<i>Cytherea</i>), Dunker	II. 7	tumescens (<i>Venus</i>), Blainville	IX. 35
impar (<i>Cytherea</i>), Lamarck	II. 8	umbonella (<i>Cytherea</i>), Lamarck	VII. 27
inflata (<i>Cytherea</i>), Sowerby	IX. 37	unicolor (<i>Cytherea</i>), Sowerby	VIII. 33
Kingii (<i>Venus</i>), Gray	IX. 36	ustulata, Reeve	XI. 49
laeta (<i>Venus</i>), Linnaeus	IX. 35	Veneris (<i>Concha</i>), D'Argenville	VI. 23
leptida (<i>Cytherea</i>), Chenu	VII. 29	virgo (<i>Chione</i>), Gray	XI. 50
ligula (<i>Cytherea</i>), Anton	VIII. 33	rulgarris (<i>Chione</i>), Gray	IV. 13
lilacina (<i>Cytherea</i>), Lamarck	I. 4	vulnerata (<i>Cytherea</i>), Broderip	V. 16

ERRATA.

Plate VI., Sp. 20, for "*Cytherea lamellosa*, Gaudichaud," read "*Cytherea semilamellosa*, Gaudichaud."

Plate VI., Fig. 23 on the plate should be Fig. 24, and Fig. 24 should be Fig. 23.

MONOGRAPH
OF THE GENUS
C I R C E.


“Read Nature; Nature is a friend to truth.
Nature is Christian, preaches to mankind,
And bids dead matter aid us in our creed.”—*Young.*



2.a.




1.a




2.b.



1.b.



3.a.



3.b.



1.c.

C I R C E.

PLATE I.

Genus CIRCE, Schumacher.

Testa nunc plano-lenticularis, umboibus compressis, nunc ovala, turgida, umboibus cordatis. Cardo tridentatus, dentibus inaequalibus, divergentibus, antico lateraliter elongato. Impressio pallii simplex, haud sinuata.

Shell sometimes flatly lens-shaped with the umboes compressed, sometimes ovate and turgid with the umboes cordate. Hinge three-toothed, teeth unequal, diverging, the anterior laterally elongated. Impression of the mantle simple, not sinuated.

In this group of about fifty species, all of which, so far as the habitats are at present known, are, with a single exception (*C. nummulina*), inhabitants of the Eastern Hemisphere, we have two apparently very dissimilar forms. In the first of these, of which the Linnaean *Venus scripta* (*Cytherea scripta*, Lamarck) is the type, the shell is singularly flat, compressly pinched at the umboes; in the second, of which the Linnaean *Venus castrensis* (*Cytherea castrensis*, Lamarck) is the type, the shell is ovately turgid, with the umboes cordate. And, intermediate between these forms, a very characteristic link is presented in the Linnaean *Venus pectinata* (*Cytherea pectinata*, Lamarck), with the sculpture peculiarly divaricate, which divarication is present in *C. intermedia*, a member of the *C. castrensis* group. All are distinguished generically by the absence of any posterior sinus in the pallial impression. *C. castrensis* and its congeners have been separated as a subgenus by Morech, under the title *Lioconcha*.

With the exception of the single Central American species already mentioned, and the two small European species *C. rufis* and *minima*, the *Circe* inhabit exclusively the Indian Ocean (including the Philippine Islands), Australia, and the Red Sea, and *C. gibba*, *pectinata*, and *divaricata*, belonging to the intermediate group, inhabit curiously enough, all of these three localities in common. It should be added, however, that it is only of about one-half of the *Circe* that the habitats are as yet known.

Species 1. (Fig. a, b, c, Mus. Cuming.)

CIRCE SCRIPTA. *Circ. testa subquadrato-lentiformi, compressa, posticè angulatè truncata, fulvescente-alba, castaneo-fusco linearis-undata, plerumque biradiatim maculata, concentricè crebrilirata, umboibus plano-compressis.*

THE WRITTEN CIRCE. Shell somewhat squarely lens-

shaped, compressed, posteriorly angularly truncated, fulvous-white, linearly waved with chestnut-brown, generally biradiately blotched, concentrically closely ridged, umboes flatly compressed.

Venus scripta, Linnaeus, Syst. Nat. p. 1135.

Cytherea scripta and *undata*, Lamarck.

Hab. Eastern Seas.

The shell represented at Fig. 1 c is, I believe, the variety named *Cytherea undata* by Lamarck. It is somewhat conspicuously removed from the normal type of the species, Fig. 1 b, but this species is one of the most variable of the genus.

Species 2. (Fig. a, b, Mus. Cuming.)

CIRCE PLICATINA. *Circ. testa obliquè lenticulari, crassiuscula, posticè rotundato-truncata, fulvescente-alba, nigricante-castaneo reticulato-undata, concentricè validè corrugato-plicata.*

THE PLEATED CIRCE. Shell obliquely lens-shaped, rather thick, posteriorly rotundately truncated, fulvous-white, reticulately waved with dark chestnut, concentrically strongly wrinkle-pleated.

Cytherea plicatina, Lamarck, Anim. sans vert. vol. vi. p. 326.

Hab. New Holland.

This is a stouter and more convex species than *C. scripta*, of a peculiar oblique, instead of square growth, very strongly concentrically pleated.

Species 3. (Fig. a, b, Mus. Cuming.)

CIRCE RIVULARIS. *Circ. testa subquadrato-lentiformi, depressissimam, fulvescente-alba, spadiceo-rubo radiatum nebulatum, circa umbones levigata, deinde concentricè rufè lirata, ad latera exiliter diagonaliter rugata.*

THE RILLED CIRCE. Shell somewhat squarely lens-shaped, very depressed, fulvous-white, radiately clouded with fawn-red, smooth around the umboes, then concentrically rudely ridged, at the sides faintly diagonally wrinkled.


Venus rivularis, Born, Test. Mus. Vindob. p. 72. pl. 5. f. 7.

Cytherea rivularis, Menke.

Circe rivularis, Sowerby.

Hab. New Holland.

A pale, very depressed species, radiately clouded with fawn-red.



C I R C E .

PLATE II.

Species 4. (Mus. Cuming.)

CIRCE CORRUGATA. *Circ. testā suborbiculari, convexā, posticē obliquē angulatā, compressiusculā, fulvescente-lacteā, lineis tenuibus rufo-fuscis reticulatā, concentricē ratiōne plicato-liratā, circa umbones conspicuē undato-rugosā.*

THE WRINKLED CIRCE. Shell suborbicular, eouvex, posteriorly obliquely angled, rather compressed, fulvous cream-colour, reticulated with fine red-brown lines, concentrically strongly plicately ridged, conspicuously wave-wrinkled round the umboes.

Venus corrugata, Chemnitz, Conch. Cab. vol. vii. p. 25, pl. 39. f. 410, 411.

Cytherea rugifera, Lamarck.

Circe corrugata, Gray.

Circe rugifera, Sowerby.

Hab. Red Sea.

The fine network of red-brown lines becomes obsolete with the thickening of the crowded concentric ridges.

Species 5. (Mus. Cuming.)

CIRCE FULGURATA. *Circ. testā ovatā, posticē subquadratā, compressā, livido-albidente, lineis rufo-fuscis acutē angulatis copiosè notatā, concentricē densè sulcatā et lirato-striatā, posticē lăviusculā.*

THE LIGHTNING-MARKED CIRCE. Shell ovate, posteriorly somewhat square, compressed, livid-white, copiously marked with sharply zigzag red-brown lines, concentrically densely grooved and ridge-striated, posteriorly rather smooth.

Hab. —?

An oblong-ovate form of *C. scripta*, in which the marking is crowded and sharply angled.

Species 6. (Fig. a, b, c, Mus. Cuming.)

CIRCE PERSONATA. *Circ. testū suborbiculari, posticē subquadratā, valde compressā, livido-fulvescente, rufo-castaneo radiatā et nebulatā, concentricē densè sulcatā et lirato-striatā, posticē lăviusculā.*

THE MASKED CIRCE. Shell nearly orbicular, posteriorly somewhat square, very compressed, livid-fulvous, rayed and clouded with reddish-chestnut, concentrically closely grooved and ridge-striated, posteriorly rather grooved.

DESHAYES, Pro. Zool. Soc. 1853, p. 6.

Hab. Philippine Islands; Cuming.

This also is a form of *C. scripta*, in which the clouded rays of colour give fancied resemblances at their margin of the human profile.

Species 7. (Fig. a, b, Mus. Cuming.)


CIRCE LENTICULARIS. *Circ. testā circulari-ovatā, posticē subproductā, solidā, convexā, fulvescente-albā, lineis purpureo-nigris obliquē diagonaliter undatā, concentricē rudē plicato-liratā, versus umbones radiatim divaricato-striatā.*

THE LENS-SHAPED CIRCE. Shell circularly ovate, a little produced posteriorly, solid, convex, fulvous-white, diagonally sharply waved with purple-black lines, concentrically rudely plicately ridged, towards the umboes radiately divaricately striated.

DESHAYES, Pro. Zool. Soc. 1853, p. 7.

Hab. —?

A very characteristic species, swollen in the middle, compressed towards the periphery, conspicuously marked with diagonally-waved sharply zigzag purple-black lines.



9.



8.




10. a.



10. b.



11.



13.



12.

C I R C E .

PLATE III.

Species 8. (Mus. Cuming.)

CIRCE ORBICA. *Circ. testā orbiculari, lenticulari, versus umbones subtriangulari, lacteā, purpureo-nigro transversim undulatā, concentricē confertim plicato-liratā, ad umbones nitidē divaricatum plicatā.*

THE CIRCULAR CIRCE. Shell orbicular, lens-shaped, slightly triangular towards the umboes, cream-colour, transversely waved with purple-black, concentrically closely plicately ridged, neatly divaricately plaited at the umboes.

Hab. —?

Of the same typical character as *C. scripta*, but of a solid circular lens-shape, beautifully painted and sculptured.

Species 9. (Mus. Cuming.)

CIRCE TUMEFACTA. *Circ. testā suborbiculari-ovalā, posticē declivi-lobatā, medio tumidiusculā, fulvescente-albā, nigricante-castaneo acutē linearī-undatā, concentricē rudē plicato-liratā, posticē levigatā, ad umbones divaricatum plicatā.*

THE SWOLLEN CIRCE. Shell suborbicularly ovate, posteriorly slopingly lobed, rather swollen in the middle, fulvous-white, sharply linearly waved with dark-chestnut, concentrically rudely plicately ridged, smooth posteriorly, divaricately plated at the umboes.

SOWERBY, Thes. Conch. vol. i. p. 652. pl. 138. f. 28 to 30.

Hab. —?

Differing from the preceding species in being slopingly lobed on the posterior side, while the concentric sculpture is more rude and irregular.

Species 10. (Fig. a, b, Mus. Cuming.)

CIRCE CROCEA. *Circ. testā obliquē suborbiculari, nunc compressā, nunc tumidiusculā, fulvescente-lacteā, purpurascente pallidē tinctā, ad umboes purpurascente-ferrugineo biradiatā, exiliter concentricē striatā, ad umboes divaricatum plicato-rugatā, intus croceā.*

THE SAFFRON CIRCE. Shell obliquely suborbicular, sometimes compressed, sometimes rather swollen, fulvous cream-colour, faintly tinged with purple, two-rayed with purple-rust at the umboes, finely concentrically striated, divaricately plicately wrinkled at the umboes, saffron-colour in the interior.

GRAY, Cat. Cyth. Anal. vol. viii. p. 307.

Cytherea crocea, Hanley.

Hab. Red Sea.

Mostly distinguished by two purple-rust rays at the umboes. The interior is invariably tinged with saffron-colour.

Species 11. (Mus. Cuming.)

CIRCE SUGILLATA. *Circ. testā circulari-ovalā, posticē subproductā, livido-albicante, castaneo-brunneo copiosè radiatim tinctā, concentricē rudē liratā, ad umboes exiliter divaricatum plicatā.*

THE BRUISED CIRCE. Shell circularly ovate, slightly produced posteriorly, livid-white, copiously radiately stained with chestnut-brown, concentrically rudely ridged, faintly divaricately plicated at the umboes.

Hab. —?

This richly-stained chestnut-brown shell approaches *C. trigona*, but it has not the triangular form characteristic of that species.

Species 12. (Mus. Cuming.)

CIRCE TRIGONA. *Circ. testā orbiculari, versus umbones conspicuē trigonā, planō-compressā, fulvescente-albā, concentricē confertim rudē liratā, ad umboes divaricatum corrugato-plicatā.*

THE TRIANGULAR CIRCE. Shell orbicular, conspicuously triangled towards the umboes, flatly compressed, fulvous-white, concentrically closely rudely ridged, divaricately corrugately plicated at the umboes.

Hab. —?

Distinguished by a flatly compressed triangular growth, though still rudely concentrically ridged throughout.

CIRCE.—PLATE III.

Species 13. (Mus. Cuming.)

CIRCE ALBIDA. *Circ. testā ovatā, posticē subquadrato-lobatā, valde compressā, lacteā, immaculatā, concentricē striato-liratā, striis ad latus posticum tenuioribus.*

THE WHITISH CIRCE. Shell ovate, posteriorly somewhat squarely lobed, very compressed, cream-colour,

unspotted, concentrically striately ridged, striae finer on the posterior side.

DESHAYES, Cat. Conch. Brit. Mus. p. 84.

Hab. Australia.

Very closely allied to *C. scripto*; remarkably thinly compressed, without any indication of colour or marking.



14 a



14 b.



14 c.



15



16 a



16 b



17



16 c.



18



19

C I R C E .

PLATE IV.

Species 14. (Fig. *a, b, c*, Mus. Cuming.)

CIRCE MINIMA. *Circ. testū orbiculari-cordatā, interdum ventricosā, fulvescente-albā, rufo-fusco lentiginosā vel radiatim maculatā, concentricè conspicuè striatā, striis plus minus irregularibus.*

THE VERY SMALL CIRCE. Shell orbicularly heart-shaped, sometimes ventricose, fulvous white, freckled or radiately blotched with red-brown, concentrically conspicuously striated, striae more or less irregular.

Venus minima, Montagu, Test. Brit. p. 121. pl. 3. f. 3.

Venus triangularis, Montagu.

Cyprina minima, Turton.

Cyprina triangularis, Turton.

Cytherea triangularis, Macgillivray.

Cytherea apicalis, Philippi.

Circe triangularis, King.

Cytherea Cyrilli, Philippi.

Cytherea minima, Brown.

Cytherea minuta, Brown.

Circe minima, Forbes and Hanley.

Hab. European Seas.

This well-known species, the smallest of the genus, varies curiously in convexity and in marking. It is very generally distributed throughout the British Isles.

Species 15. (Mus. Cuming.)

CIRCE ARTEMIS. *Circ. testā orbiculato-lenticulari, depressā, candidā, rufo-fusco obsoletē sparsim radiatā, lunulā et areā ligamenti rufo maculatā, concentricè tirato-striatā, intersticiis crispato-punctatis.*

THE ARTEMIS CIRCE. Shell orbicularly lens-shaped, depressed, white, obsoletely sparingly rayed with red-brown, lunule and ligamentary area red-spotted, concentrically ridge-striated, interstices crisply punctured.

DESHAYES, Cat. Conch. Mus. Brit. p. 86.

Hab. —?

Bright opaque white, peculiarly sparingly rayed with red-brown dots or broken spots of the same colour.

Species 16. (Fig. *a, b, c*, Mus. Cuming.)

CIRCE SULCATA. *Circ. testā subtrigono-orbiculari, lenti-*

culari, depressā, fulvescente-albā, rufescente-castaneo radiatā vel plus minus copiosè tinctā, concentricè tenui-liratā, sulcis interstitialibus crispato-punctatis.

THE GROOVED CIRCE. Shell somewhat triangularly orbicular, lens-shaped, depressed, fulvous white, rayed or more or less copiously stained with reddish-chestnut, concentrically finely ridged, interstitial grooves crisply punctured.

GRAY, Cat. Cyth. Anal. vol. viii. p. 107.

Cytherea sulcata, Hanley.

Cytherea erythræa, Jonas.

Hab. Philippine Islands; Cuming.

Extremely variable in marking, but the marks are generally of the same uniform reddish-chestnut colour.

Species 17. (Mus. Cuming.)

CIRCE PLEBEIA. *Circ. testū obliquè orbiculari, depressā, albida, rufo-fusco sparsissimè obsoletè linearī-undatā, concentricè tenuiliratā, sulcis interstitialibus crispato-punctatis.*

THE POOR CIRCE. Shell obliquely orbicular, depressed, whitish, very sparingly obsoletely linearly waved with red-brown, concentrically finely ridged, interstitial grooves crisply punctured.

Cytherea plebeia, Hanley, Pro. Zool. Soc. 1844, p. 109.

Index Test. Supp. pl. 15. f. 37.

Circe plebeia, Sowerby.

Circe sulcata, var., Deshayes.

Hab. Philippine Islands; Cuming.

We give a figure of this shell as being the type of Mr. Hanley's *Cytherea plebeia*, but it may prove, as suggested by M. Deshayes, to be a large obliquely produced variety of the preceding species. The details of sculpture are the same.

Fig. 18. (Mus. Cuming.)

This interesting small shell, marked throughout with an angular network of red-brown lines, proves to be *C. corrugata* in an early stage of growth.

CIRCE.—PLATE IV.

Species 19. (Mus. Cuming.)

CIRCE AUSTRALIS. *Circ. testā subtrigono-ovatā, postice subquadratā, depressā, cærulescente-albā, punctis rufescente-fuscis pallidè aspersā, tunulā elongatā, intensē livido-fuscā, concentricè confertim liratā, liris subrecurvis, interstitiis excavatis.*


THE SOUTHERN CIRCE. Shell somewhat triangularly ovate, posteriorly rather square, depressed, bluish-white, faintly sprinkled with red-brown dots, lunule

elongated, dark livid-brown, concentrically closely ridged, ridges slightly recurved, interstices excavated.

SOWERBY, Thes. Conch. vol. i. p. 651. pl. 137. f. 16,
17.

Hab. Australia.


A very characteristic triangularly ovate species, in which the dark livid-brown lunule contrasts strongly in colour with the faint sprinkling of red dots on the general surface of the shell.




20 a



20 b.




20 c.




21 a



21 b.



21 c.



21 d

CIRCE.

PLATE V.

Species 20. (Fig. *a*, *b*, *c*, Mus. Cuming.)

CIRCE PECTINATA. *Circ. testā oblongo-ovali, compressā, posticē subquadratā, albida, rufo-fusco concentricè maculatā et punctatā, radiatim costatā, costis irregularibus, nodoso-granatis, versus marginem bifurcatis, latere postico divaricatum bifurcatum plicato.*

THE COMB-LIKE CIRCE. Shell oblong-oval, compressed, posteriorly somewhat square, whitish, concentrically blotched and dotted with red-brown, radiately ribbed, ribs irregular, nodosely grained, bifurcated towards the margin, posterior side divaricately bifurcate plaited.

Tenus pectinata, Linnaeus, Syst. Nat. p. 1135.

Cytherea pectinata, Lamarck.

Circe pectinata, Gray.

Cytherea Sarignyi, Jonas.

Hab. Red Sea, Philippine Islands.

This well-known species is typically of a compressed form, but it varies considerably and passes into the gibbosity of the next species. The sculpture is finer and more elaborate.

Species 21. (Fig. *a*, *b*, *c*, *d*, Mus. Cuming.)

CIRCE GIBBA. *Circ. testā oblongo-ovatā, crassā, valde gibbosā, posticē parum quadratā, albida, interdum ferrugineo tinctā, rufo-fusco aut nigro posticē et versus umbones maculatā, radiatim costatā, costis irregularibus, rudē nodoso-granatis, ad marginem interdum bifurcatis, latere postico divaricatum bifurcatum plicato.*

THE GIBBOUS CIRCE. Shell oblong-ovate, thick, very gibbous, posteriorly but little square, whitish, sometimes tinged with rust, spotted posteriorly and towards the umboes with red-brown, or black, radiately ribbed, ribs irregular, rudely nodosely grained, sometimes bifurcated at the margin, posterior side divaricately bifurcate plaited.

Cytherea gibba, Lamarck, Anim. sans vert. vol. v. p. 577

Cytherea ranella, Lamarck.

Circe gibba, Gray.

Hab. Red Sea, Philippine Islands, Australia.

Of a very solid gibbous growth compared with the preceding species, much more rudely sculptured, the ribs being strongly nodosely grained, deeply excavated in the interstices.

October, 1863.



22.a.



22.b.




22.c.




23.a.



23.b.



23.c.



23.d.



24.a



24.b.



25

C I R C E .

PLATE VI.

Species 22. (Fig. *a, b, c*, Mus. Cuming.)

CIRCE ÆQUIVOCATA. *Circ. testá obliquè ocatá, compressiusculá, ad latus posticum subtruncatá, fulvescente-albá, cinereo-fusco undulatá aut varie tinctá, versus marginem plerumque reticulato-maculatá, concentricè rudè rugoso-liratá, antice posticèque validè divaricatum plicatá.*

THE EQUIVOCAL CIRCE. Shell obliquely ovate, rather compressed, slightly truncated on the posterior side, fulvous white, waved or variously stained with ash-brown, generally reticulately spotted towards the margin, concentrically rudely wrinkle-ridged, strongly divaricately plicated on the anterior and posterior sides.

x)

Venus æquivoca, Chemnitz, Conch. Cab. vol. ii. p. 229. pl. 202. f. 1980.

Circe æquivoca, Deshayes.

Circe transversaria, Deshayes.

Hab. Philippine Islands; Cuming.

The divaricate plications of this prettily painted species, crossing the concentric ridges nearly throughout, give the shell a crisply granular appearance.

Species 23. (Fig. *a, b, c, d*, Mus. Cuming.)

CIRCE DIVARICATA. *Circ. testá oblongo-ovali, ad latus posticum obliquè productá vel attenuatá, convexá, interdum gibbosiusculá, crassá, fulvescente-albá, cinereo et rufo-castaneo varie angulato-maculatá, maculis interrupte radiantibus, concentricè irregulariter obtuse liratá, undique validè divaricatum plicatá, ad decussationem plus minus conspicuè granulatá.*

THE DIVARICATE CIRCE. Shell oblong-oval, obliquely produced or attenuated on the posterior side, convex, sometimes rather gibbous, thick, fulvous white, variously angularly blotched with ash and reddish-chestnut, blotches interruptedly radiating, concentrically irregularly obtusely ridged, strongly divaricately plicated throughout, more or less conspicuously granulated at the crossing.

Venus divaricata, Chemnitz, Conch. Cab. vol. vi. p. 317. pl. 30. f. 316.

Cytherea divaricata, Lamarck.

Cytherea testudinalis, Lamarck.

Venus discors, Schröter.

Circe divaricata, Gray.

Hab. Red Sea; Philippine Islands.

The four specimens of this well-known species here figured tolerably represent the range of its numerous varieties. It may be regarded as the type of the divaricately plicated forms of the genus.

Species 24. (Fig. *a, b*, Mus. Cuming.)

CIRCE DISPAR. *Circ. testá oblongo-ovali, compressiusculá, fulvescente-albá, maculis rufo-aurantiis vel purpurascientibus subprofuse marmoratá, concentricè irregulariter rudè liratá, divaricatum plicatá, plicis posticis multo validioribus.*

THE UNLIKE CIRCE. Shell oblong-oval, rather compressed, fulvous white, rather profusely mottled with reddish-orange or purple spots, concentrically irregularly rudely ridged, divaricately plicated, posterior plaits much the stronger.

Venus dispar, Chemnitz, Conch. Cab. vol. ii. p. 230. pl. 202. f. 1981, 1982.

Cytherea muscaria, Lamarck.

Cytherea pulicaris, Lamarck.

Cytherea dispar, Hanley.

Circe dispar, Gray.

Hab. Red Sea.

One of the most obvious characteristics of this species is its mottled painting of reddish-orange or purple dots.

Species 25. (Mus. Cuming.)

CIRCE NUMMULINA. *Circ. testá obliquè trigono-orbiculari, subcuneiformi, crassá, fulvescente-lacteá, lunulá et areá ligamenti livido-purpurascientibus, concentricè tenuiliratá, versus umbones subcompressá et indistinctè divaricatum plicatá.*

THE COIN-LIKE CIRCE. Shell obliquely triangularly orbicular, somewhat wedge-shaped, thick, fulvous cream-colour, lunule and ligamentary area livid purple, concentrically finely ridged, compressed and faintly divaricately plicated towards the umboes.

Cytherea nummulina, Lamarck, Anim. sans vert. vol. vi. p. 321.

Circe nummulina, Gray.

Hab. Central America.


Of an obtuse wedge-like form, rather obsoletely sculptured, and but slightly coloured.




27. a



26.



27. b.



28. b.



28. a



28. c.



28. d.

C I R C E.

PLATE VII.

Species 26. (Mus. Taylor.)

CIRCE INTERMEDIA. *Circ. testā subobliquē globosā, crassā, ad umboes compressā, fulvescente-albā, rufo-castaneo trigono-reticulatā, maculis versus lunulam et aream ligamenti in strigis conspicuis decurrentibus, concentricē subrugosē plicato-striatā, ad umboes subobsoletē divaricatim plicatā.*

THE INTERMEDIATE CIRCE. Shell rather obliquely globose, thick, compressed at the umboes, fulvous white, triangularly reticulated with reddish-chestnut, the spots running towards the lunule and ligamentary area into conspicuous streaks, concentrically somewhat roughly plicately striated, at the umboes rather obsoletely divaricately plicated.

This appears to me to be a new species, and one of importance in the genus, as being intermediate between the types of the two groups in this and the preceding Plates. My attention was first called to the shell by Mr. Sowerby, who was preparing to send it to Mr. Taylor. On receipt of it, Mr. Taylor found in his collection another undoubted specimen in an earlier stage of growth. It helps to confirm in a singular manner the propriety of removing to this genus the Lamarekian *Cytherea castrensis* and its allies. Along with the reticulated camp-like painting of *C. castrensis*, it possesses the divaricate umbonal wrinkles of *C. divaricata* and its numerous allies, to say nothing of the absence of any angular production of the pallial impression in the interior.

Species 27. (Fig. a, b, c, Mus. Cuming.)

CIRCE ORNATA. *Circ. testā subtrigono-rotundatā, gibbosā, subcordatā, cinereā, versus umboes albida, lineis nigris ant rufis acutē undatis undique profuse radiati, concentricē irregulariter striatā, striis ad latera fortioribus.*

THE ADORNED CIRCE. Shell somewhat triangularly rounded, gibbous, somewhat heart-shaped, ash-co-

loured, whitish towards the umboes, profusely rayed throughout with black or red zigzag lines, concentrically irregularly striated, striae stronger at the sides.

Cytherea ornata, Lamarek, Anim. sans vert. vol. vi. p. 309.

Circe castrensis var., Deshayes.

Lioconcha ornata, Mörch.

Hab. Indian Ocean.

M. Deshayes is probably correct in recording the Lamarekian *C. ornata* as a variety of *C. castrensis*, but it is a very characteristic variety and it is of a less solid porcelain-like substance.

Species 28. (Fig. a, b, c, Mus. Cuming.)

CIRCE CASTRENSIS. *Circ. testā subtrigono-globosā, vel cordatā, albā, nitente, interdum cinereo vel croceo tinctā, lineis vel maculis castaneis ant nigris acutē angulatis plus minus suffusis pictā, concentricē irregulariter striatā, interdum læviusculā, striis ad latera fortioribus.*

THE CAMP CIRCE. Shell somewhat triangularly globose or heart-shaped, white, shining, sometimes ash- or saffron-tinged, painted with more or less suffused chestnut or black sharply angled lines or spots, concentrically irregularly striated, sometimes almost smooth, striae stronger at the sides.

Tenus castrensis, Linnaeus, Syst. Nat., p. 1132.

Tenus australis, Chemnitz.

Tenus Lorenziana, Chemnitz.

Cytherea castrensis, Lamarek.

Circe castrensis, Gray.

Lioconcha castrensis, Mörch.

Hab. Indian Ocean.

Among the varieties of this beautiful species, the shell represented at Fig. 28 b is that which passes into *C. ornata*. It is broader from the umboes to the ventral margin, and has less of the smooth porcelain surface which is so characteristic a feature of the other varieties.




30.



29.a



31.




32.a.




29.b.



32.b.



33.



29.c.



34.



35.

CIRCE.

PLATE VIII.

Species 29. (Fig. *a, b*, Mus. Cuming.)

CIRCE SOWERBYI. *Circ. testā trigono-rotundatā rel-*
ovatā, turgidā, fulvescente-albā, castaneo-rufo tenuē
acutē undatā, concentricē confertim striato-liratā, liris
obtusis.

SOWERBY'S CIRCE. Shell triangularly rounded or ovate,
 swollen, fulvous white, finely sharply waved with
 chestnut-red, concentrically closely striately ridged,
 ridges obtuse.

DESHAYES, Cat. Conch. Brit. Mus. p. 95.

Cytherea Hebræa, Sowerby (not of Lamarck).

Hab. Philippine Islands.

This pretty species runs very closely into *C. tigrina*.
 If a variety, it is a very characteristic one.

Species 30. (Mus. Cuming.)

CIRCE MARMORATA. *Circ. testā ovatā, subcompressā,*
livido-cærulescente, olivaceo-rufo reticulato-marmoratā,
lunulā et areā ligamenti plumbeis, umboibus purpu-
rascente radialis, concentricē obtusē striato-liratā, pos-
ticē et circa umbones profusē tenuē divaricatim pli-
cataū.

THE MARBLED CIRCE. Shell ovate, rather compressed,
 livid bluish, reticulately marbled with olive-red,
 lunule and ligamentary area lead-colour, umboes
 rayed with faint purple, concentrically obtusely
 striately ridged, posteriorly and around the umboes
 profusely finely divaricately plicated.

Hab. —?

Apparently a new species, fully characterized, as may
 be seen by the foregoing description, both in respect of
 sculpture and colour.

Fig. 31. (Mus. Cuming.)

The shell here figured proves to be an interesting va-
 riety of *C. elliptica*, described at Plate X. Species 48.

Species 32. (Fig. *a, b, c*, Mus. Cuming.)

CIRCE TIGRINA. *Circ. testā subtrigono-ovatā, posticē at-*
tennato-productā, versus umbones turgidā, albida, cas-
taneo-rufo aut nigricante-brunneo profusē reticulato-
marmoratā, medio laeviscentia, ad latera concentricē
striato-liratā, liris obtusis, irregulariter corrugatis.

THE TIGER CIRCE. Shell somewhat triangularly ovate,
 attenuately produced on the posterior side, swollen
 towards the umboes, whitish, profusely reticulately
 mottled with chestnut-red or dark-brown, rather
 smooth in the middle, concentrically striately ridged
 at the sides, ridges obtuse, irregularly wrinkled.

Cytherea tigrina, Lamarck, Anim. sans vert. vol. vi.
 p. 310.

Circe tigrina, Deshayes.

Hab. Molucca and Philippine Islands.

Strikingly characterized by its reticulately mottled painting. The shell, which has invariably a tendency to become attenuated on the posterior side, is comparatively smooth in the middle, the sides being obtusely irregularly wrinkle-striated.

Species 33. (Mus. Cuming.)

CIRCE TRIMACULATA. *Circ. testā obliquē cordatā, posticē*
curratā, albā, castaneo-brunneo profusē tinctā et ra-
diatā, lunulā et areā ligamenti intensē brunneis, undique
concentricē lirato-striatā, striis obtusis, irregulariter
corrugatis.

THE THREE-SPOTTED CIRCE. Shell obliquely heart-
 shaped, curved posteriorly, white, profusely stained
 and rayed with chestnut-brown, lunule and ligamen-
 tary area dark-brown, concentrically ridge-striated
 throughout, striae obtuse, irregularly wrinkled.

Cytherea trimaculata, Lamarck, Anim. sans vert. vol. vi.
 p. 313.

Circe trimaculata, Deshayes.

Hab. Philippine Islands; Cuming.

Of a more curved heart-shaped form than the preceding
 species, sculptured throughout.

Species 34. (Mus. Cuming.)

CIRCE HEBREA. *Circ. testā oratā, turgidā, posticē sub-*

CIRCE.—PLATE VIII.

truncatā, fulvescente-albā, rufo-castaneo sparsim reticulatā, concentricè subobsoletè striatā.

THE HEBREW CIRCE. Shell ovate, swollen, posteriorly slightly truncated, fulvous white, sparingly reticulated with reddish-chestnut, concentrically rather obsoletely striated.

Cytherea Hebræa, Lamarck, Anim. sans vert. vol. vi. p. 308.

Dione Hebræa, Deshayes.

Hab. ——?

This shell partakes very much of the characters of *Dione*, and should, perhaps, be referred to that genus.

Species 35. (Mus. Cuming.)

CIRCE FUMATA. *Circ. testā obtongo-ovatā, compressè convexā, fulvescente-albā, ustulato-castaneo copiosè reticulatā et nebulatā, concentricè densè filo-striatā, striis versus marginem obtusè corrugatis.*

THE SMOKED CIRCE. Shell oblong-ovate, compressly convex, fulvous white, copiously reticulated and clouded with burnt-chestnut, concentrically densely thread-striated, striae obtusely wrinkled towards the margin.

Hab. ——?

Of a distinct convexly oval form, peculiarly clouded and reticulately streaked with burnt-chestnut.



36.



37




38



39.




40.a




40.b




41



42.a



42.b



43

CIRCE.

PLATE IX.

Species 36. (Mus. Cuming.)

CIRCE FLUCTUATA. *Circ. testā globoso-cordatā, tenuiculā, turgidā, albida, rufo-fusco prope umboes pallidē reticulatā, undique concentricē densē filo-striatā.*

THE WAVED CIRCE. Shell globosely heart-shaped, rather thin, swollen, whitish, faintly reticulated near the umboes with red-brown, concentrically densely thread-striated throughout.

Cytherea fluctuata, Sowerby, Thes. Coneh. vol. i. p. 634. pl. 136. f. 185, 186,

Dione fluctuata, Deshayes.

Hab. St. Helena.

A small thread-striated swollen globose shell, partaking of the generic characters both of *Dione* and *Circe*.

Species 37. (Mus. Cuming.)

CIRCE MANILLE. *Circ. testā obliquē cordatā, turgidā, posticē flexuoso-attenuatā, albida, concentricē densē filo-striatā, lunula amplā.*

THE MANILLA CIRCE. Shell obliquely heart-shaped, swollen, posteriorly flexuously attenuated, white, concentrically densely thread-striated, lunule large.

Cytherea Manillæ, Sowerby, Thes. Coneh. vol. i. p. 634. pl. 136. f. 180, 181.

Dione Manillæ, Deshayes.

Hab. Manilla, Island of Luzon, Philippines; Cuming.

Allied to the preceding species, but with the posterior side flexuously attenuated.

Species 38. (Mus. Cuming.)

CIRCE RUDIS. *Circ. testā obliquē cordatā, turgidā, fulvescente-albida, ferrugineo-rufō tinctā et radiatā, concentricē tenuē filo-striatā, striis in medio plus minus obsoletis.*

THE RUDE CIRCE. Shell obliquely heart-shaped, swollen, fulvous white, stained and rayed with rust-red, concentrically finely thread-striated, striae more or less obsolete in the middle.

Venus rufis, Poli, Test. utr. Sieil. vol. ii. p. 40. pl. 20. f. 15, 16.

Cytherea Venetiana, Lamarek.

Dione rufis, Deshayes.

Hab. Mediterranean.

The thread-like striae of this species are comparatively obsolete in the middle of the shell towards the umboes.

Species 39. (Mus. Cuming.)

CIRCE VARIANS. *Circ. testā suboblongo-ovatā, antice cordatā, posticē subproductā, tenui, ventricosā, fulvescente-lactea, maculis ferrugineo-rufis radia, lunula et areā ligamenti maculatis, concentricē densē filo-striatā.*

THE VARYING CIRCE. Shell rather oblong-ovate, anteriorly heart-shaped, posteriorly rather produced, thin, ventricose, fulvous cream-colour, rayed with rust-red spots, lunule and ligamentary area spotted, concentrically densely thread-striated.

Cytherea varians, Wood, Ind. Test. Supp. pl. 15. f. 33.

Cytherea rubiginosa, Philippi.

Dione varians, Deshayes.

Hab. —?

This and the three preceding species represent by their affinity to *Dione laeta* on the one side and *Circe sulcata* on the other, the connection between the two genera.

Species 40. (Fig. a, b, Mus. Cuming.)

CIRCE SULCATA. *Circ. testā trigono-cordatā, plus minus turgidā, solidusculā, fulvescente-lactea, castaneo-rufō cariē picta, medio levigata, ad latera corrugato-striata.*

THE FINELY-GROOVED CIRCE. Shell triangularly heart-shaped, more or less swollen, rather solid, fulvous cream-colour, variously painted with chestnut-red, smooth in the middle, wrinkle-striated at the sides.

Cytherea sulcata, Lauarek, Anim. sans vert. vol. vi. f. 308.

Circe sulcata, Deshayes.

Hab. Philippine Islands; Cuming.

Extremely variable in painting, which is sometimes disposed in rays sometimes in concentrically waved lines.

CIRCE.—PLATE IX.

Species 41. (Mus. Cuming.)

CIRCE FASTIGIATA. *Circ. testá sublrigono-ovalá, posticè subattenuato-productá, fulvá vel rufescente-albá, nigricante-castaneo conspicuè angulatim maculatá, concentricè corrugato-striatá, striis rudibus, irregularibus.*

THE POINTED CIRCE. Shell somewhat triangularly ovate, posteriorly slightly attenuately produced, fulvous or reddish-white, conspicuously angularly blotched with dark-chestnut, concentrically wrinkle-striated, striae rude, irregular.

Cytherea fastigiata, Sowerby, Thes. Conch. vol. i. p. 643. pl. 135. f. 158, 159.

Circe fastigiata, Deshayes.

Hab. Sydney, New Holland.

Strongly resembling *C. castrensis* in painting, but constantly distinct in form and uniform size.

Species 42. (Fig. *a, b*, Mus. Cuming.)

CIRCE PICTA. *Circ. testá trigono-ovalá, posticè attenuato-productá, albá, lineis nigricante-castaneis angulatim reticulatá, concentricè filo-striatá, striis ad latera interdum corrugatis.*

THE PAINTED CIRCE. Shell triangularly ovate, posteriorly attenuately produced, white, angularly reti-

culated with dark-chestnut lines, concentrically thread-striated, striae sometimes wrinkled at the sides.

Cytherea picta, Lamarck, Anim. sans vert. vol. vi. p. 569.

Circe picta, Gray.

Hab. Indian Ocean.

Elegantly reticulated with an angular textile of dark-chestnut, more or less open in its distribution.

Species 43. (Mus. Cuming.)

CIRCE HIEROGLYPHICA. *Circ. testá trigono-ovalá, compressá, posticè attenuato-productá, albá, maculis nigricante-castaneis triangularibus aspersá, medio lavigatá, ad latera corrugato-striatá.*

THE HIEROGLYPHIC CIRCE. Shell triangularly ovate, compressed, posteriorly attenuately produced, white, sprinkled with triangular blotches of dark-chestnut, smooth in the middle, wrinkle-striated at the sides.

Meretrix hieroglyphica, Conrad, Journ. Aead. Nat. Sci. Philadelphia, vol. vii. p. 253. pl. 19. f. 22.

Cytherea hieroglyphica, Sowerby.

Hab. — ?


This is closely allied to *C. fastigiata*, but it is characterized by a much more delicate growth, while the painting is peculiar.



44.a




44.b




44.c.




45.a



45.b.



46.



45.c.



47.



49.



48.

CIRCE.

PLATE X.

Species 44. (Fig. *a, b, c*, Mus. Cuming.)

CIRCE ARABICA. *Circ. testā ovalā vel obliquē orbiculari, posticē subquadratā, crassā, fulvā aut cārūlescente-albā, rufō-fusco obscurē biradiatā et punctato-aspersā, concentricē irregulariter callosē lirato-striatā.*

THE ARABIC CIRCE. Shell ovate or obliquely orbicular, posteriorly somewhat square, thick, fulvous or bluish-white, obscurely two-rayed and dot-sprinkled with red-brown, concentrically irregularly callously ridge-striated.

Venus Arabica, Chemnitz, Conch. Cab. vol. ii. p. 224. pl. 201. f. 1968-70.

Cytherea Arabica, Lamarck.

Circe Arabica, Gray.

Hab. Red Sea; Madagascar.

Variable in form, but mostly distinguished by a dot-sprinkled colouring.

Species 45. (Fig. *a, b, c*, Mus. Cuming.)

CIRCE LENTIGINOSA. *Circ. testā ovalā vel oblongo-ovatā, crassā, albida, griseo et ferrugineo-fusco variè lentiginosā, concentricē rudē corrugato-liratā, liris in medio interdum obsoletis.*

THE FRECKLED CIRCE. Shell ovate or oblong-ovate, thick, whitish, variously freckled with grey and rust-brown, concentrically rudely wrinkle-ridged, ridges sometimes obsolete in the middle.

Venus lentiginosa, Chemnitz, Conch. Cab. vol. ii. p. 223. pl. 202. f. 1963, 1964.

Cytherea Pfeifferi, Philippi.

Circe liturata, Gray.

Circe lentiginosa, Deshayes.

Hab. Red Sea; Indian Ocean.

This species varies greatly in form, colour, and sculpture. The three specimens selected for illustration represent its extremest variations.

Species 46. (Mus. Cuming.)

CIRCE PULCHRA. *Circ. testā oblongo-ovatā, crassā, convexā, posticē subangustā, violaceo-griseā, ustulato-castaneo latē biradiatā, medio lēvigatā, ad latera corrugato-striatā.*

THE BEAUTIFUL CIRCE. Shell oblong-ovate, thick, convex, posteriorly rather narrow, violet-grey, broadly two-rayed with burnt-chestnut, smooth in the middle, wrinkle-striated at the sides.

Deshayes, Pro. Zool. Soc. 1853, p. 6.

Cytherea elegans, Sowerby (not of Koch).

Hab. Red Sea.

The colouring of this species is strongly marked; if permanent, it is highly characteristic.

Species 47. (Mus. Cuming.)

CIRCE ABBREVIATA. *Circ. testā rotundato-ovatā, anticē abbreviatā, fulvescente-lacteā, ferrugineo-rufō sparsim undatā, concentricē undique regulariter liratā et striatā.*

THE SHORTENED CIRCE. Shell rotundately ovate, shortened on the anterior side, fulvous cream-colour, sparingly waved with rust-red, concentrically regularly ridged and striated throughout.

Cytherea abbreviata, Lamarck, Anim. sans vert. vol. vi. p. 322.

Circe abbreviata, Deshayes.

Hab. —?

This passes very close to the variety of *C. lentiginosa*, represented above it at Fig. 45 *b*.

Species 48. (Mus. Cuming.)

CIRCE ELLIPTICA. *Circ. testā elliptico-ovatā, posticē subquadratā, fulvescente-spadiceā, livido-griseo tinctā, purpureo-castaneo sparsim maculatā, umboibus carneotinctis, ad latera lirato-corrugatā.*

THE ELLIPTIC CIRCE. Shell elliptically ovate, posteriorly somewhat square, fulvous bay, tinged with livid grey, sparingly blotched with purple chestnut, umboes flesh-tinged, wrinkle-ridged at the sides.

Cytherea elliptica, Sowerby, Thes. Conch. vol. i. p. 645. pl. 135. f. 174, 175.

Circe Callipyga, var., Deshayes.

Hab. —?

This shell and the shell represented at Plate VIII, Fig. 31, appear to me to confirm Mr. Sowerby's view, that it is a species distinct from *C. Callipyga*.

CIRCE.—PLATE X.

Species 49. (Mus. Cuming.)

CIRCE CALLIPYGA. *Circ. testá oblongo-ovatá, compres-
sinsculá, fulvescente-albá, rufescente-fusco variè nebu-
latá, concentricè rudè irregulariter corrugato-liratá.*

THE CALLIPYGA CIRCE. Shell oblong-ovate, rather compressed, fulvous white, variously clouded with red-brown, concentrically irregularly rudely wrinkled.

Venus Callipyga, Born, Mus. Vind. p. 68. pl. 5. f. 1.

Cytherea Callipyga, Sowerby.

Cytherea Adeneensis, Philippi.

Circe Callipyga, Deshayes.

Hab. Red Sea.


This, like *C. lentiginosa*, is also extremely variable. The two species almost pass, indeed, one into the other.

C I R C E .


	Plate.	Species.		Plate.	Species.
abbreviata (<i>Cytherea</i>), Lamarek ..	X.	47	minima (<i>Venus</i>), Montagu	IV.	14
<i>Adenensis</i> (<i>Cytherea</i>), Philippi ..	X.	49	<i>muscaria</i> (<i>Cytherea</i>), Lamarek ..	VI.	24
<i>aequivoca</i> (<i>Venus</i>), Chemnitz	VI.	22	<i>nummulina</i> (<i>Cytherea</i>), Lamarek ..	VI.	25
<i>albida</i> , <i>Deshayes</i>	III.	13	<i>orbica</i> , <i>Reeve</i>	III.	8
<i>apicalis</i> (<i>Cytherea</i>), Philippi	IV.	14	<i>ornata</i> (<i>Cytherea</i>), Lamarek	VII.	27
<i>Arabica</i> (<i>Venus</i>), Chemnitz	X.	14	<i>pectinata</i> (<i>Venus</i>), Linnaeus	V.	20
<i>Artemis</i> , <i>Deshayes</i>	IV.	15	<i>personata</i> , <i>Deshayes</i>	II.	6
<i>australis</i> , <i>Sowerby</i>	IV.	19	<i>Pfeifferi</i> (<i>Cytherea</i>), Philippi	X.	45
<i>australis</i> (<i>Venus</i>), Chemnitz	VII.	28	<i>picta</i> (<i>Cytherea</i>), Lamarek	IX.	42
<i>Callipyga</i> (<i>Venus</i>), Born	X.	49	<i>plicebia</i> (<i>Cytherea</i>), Hanley	IV.	17
<i>caestrensis</i> (<i>Venus</i>), Linnaeus	VII.	28	<i>plicatina</i> (<i>Cytherea</i>), Lamarek	I.	2
<i>corrugata</i> (<i>Venus</i>), Chemnitz	II.	4	<i>pulchra</i> , <i>Deshayes</i>	X.	46
<i>erocea</i> , <i>Gray</i>	III.	16	<i>pnlicaris</i> (<i>Cytherea</i>), Lamarek	VI.	24
<i>Cyrilli</i> (<i>Cytherea</i>), Philippi	IV.	14	<i>ranella</i> (<i>Cytherea</i>), Lamarek	V.	21
<i>dispar</i> (<i>Venus</i>), Chemnitz	VI.	24	<i>rivularis</i> (<i>Venus</i>), Born	I.	3
<i>discors</i> (<i>Venus</i>), Schröter	VI.	23	<i>rubicinosa</i> (<i>Cytherea</i>), Philippi	IX.	39
<i>divaricata</i> (<i>Venus</i>), Chemnitz	VI.	23	<i>rudis</i> (<i>Venus</i>), Poli	IX.	38
<i>elegans</i> (<i>Cytherea</i>), Sowerby	X.	46	<i>rugifera</i> (<i>Cytherea</i>), Lamarek	II.	4
<i>elliptica</i> (<i>Cytherea</i>), Sowerby	X.	48	<i>Savignyi</i> (<i>Cytherea</i>), Jonas	V.	20
<i>erythrea</i> (<i>Cytherea</i>), Jonas	IV.	16	<i>scripta</i> (<i>Venus</i>), Linnaeus	I.	1
<i>fastigiata</i> (<i>Cytherea</i>), Sowerby	IX.	41	<i>Sowerbyi</i> , <i>Deshayes</i>	VIII.	29
<i>fluctuata</i> (<i>Cytherea</i>), Sowerby	IX.	36	<i>sugillata</i> , <i>Reeve</i>	III.	11
<i>fulgorata</i> , <i>Reeve</i>	II.	5	<i>suleata</i> , <i>Gray</i>	IV.	16
<i>fumata</i> , <i>Reeve</i>	VIII.	35	<i>sulevata</i> (<i>Cytherea</i>), Lamarek	IX.	40
<i>gibba</i> (<i>Cytherea</i>), Lamarek	V.	21	<i>testudinalis</i> (<i>Cytherea</i>), Lamarek	VI.	23
<i>Hebreæa</i> (<i>Cytherea</i>), Lamarek	VIII.	34	<i>tigrina</i> (<i>Cytherea</i>), Lamarek	VIII.	32
<i>hieroglyphica</i> (<i>Meretrix</i>), Conrad	IX.	43	<i>transversaria</i> , <i>Deshayes</i>	VI.	22
<i>intermedia</i> , <i>Reeve</i>	VII.	26	<i>triangularis</i> (<i>Venus</i>), Montagu	IV.	14
<i>lenticularis</i> , <i>Deshayes</i>	II.	7	<i>trigona</i> , <i>Reeve</i>	III.	12
<i>lentiginosa</i> (<i>Venus</i>), Chemnitz	X.	45	<i>trimaenulata</i> (<i>Cytherea</i>), Lamarek	VIII.	33
<i>liturata</i> , <i>Gray</i>	X.	45	<i>tumefacta</i> , <i>Sowerby</i>	III.	9
<i>Lorenziana</i> (<i>Venus</i>), Chemnitz	VII.	28	<i>undatina</i> (<i>Cytherea</i>), Lamarek	I.	1
<i>Manille</i> (<i>Cytherea</i>), Sowerby ...	IX.	37	<i>varians</i> (<i>Cytherea</i>), Wood	IX.	39
<i>marmorata</i> , <i>Reeve</i>	VIII.	30	<i>Venetiana</i> (<i>Cytherea</i>), Lamarek	IX.	38

MONOGRAPH
OF THE GENUS
T A P E S.


"He who casts himself
On Nature's fair full bosom, and draws food,
Drunks from a fountain that is never dry."—*Longfellow*.



1.



2.a.



2.b



2.c

T A P E S.

PLATE I.

Genus TAPES, Megerle.

Testa ovata vel subquadrato-oblonga, saepissimè vulnè inaequalateralis, concentricè sulcata vel striata, interdum decussata vel corrugata, marginibus integris. Cardo tridentatus, dentibus nunc divergentibus, nunc approximatis. Impressio pallii posticè subampliter sinuata.

Shell ovate or somewhat squarely oblong, mostly very inequilateral, concentrically grooved or striated, sometimes decussated or wrinkled, margins entire. Hinge three-toothed, teeth sometimes diverging, sometimes approximating. Impression of the mantle rather largely situated on the posterior side.

Tapes is one of the most characteristic genera of the *Veneridae*. Not only is the animal distinguished, in its typical state, by the presence of a byssal groove in the foot, from which a byssus is often spun for its attachment, but the shells are marked by considerable uniformity of sculpture and colour. Four-fifths of the species are concentrically grooved and striated, and of a peculiar fulvous-drab, painted with a fine textile or with sharply angled blotches of olive-brown or black. The remaining fifth constitute a very distinct section, in which the colouring is of a duller rust-colour, and the sculpture closely decussated or wrinkled.

Tapes is essentially an Eastern genus. With the exception of eight species inhabiting the Mediterranean, of which four are moderately abundant on our own shores, and four from Senegal, the Red Sea, Natal, and Senegambia, they are all natives of the Islands of the Indian Archipelago, reaching northward to Japan and southward to Australia and New Zealand. They are most abundant at the Philippines, and several of the species inhabit the Philippine Islands and Australia in common. A further exception arises from a species believed to be from Guadalupe, which is now added to the genus.

Species 1. (Mus. Cuming.)

TAPES SULCARIA. *Tap. testa subquadrato-oblonga, sub-*

angustia, vulnè inaequalaterali, aurantio-spadicea, castaneo-lineari-reticulata et radiatim maculata, maculis violaceo-cinereo umbratis, concentricè validè sulcata et latilirata.

THE GROOVED TAPES. Shell somewhat squarely oblong, rather narrow, very inequilateral, orange-fawn, linearly reticulated and radiately blotched with chestnut, blotches shaded with violet-ash, concentrically strongly grooved and broadly ridged.

Venus sulcaria, Lamarek, Anim. sans vert. vol. vi. p. 254.

Venus retifera, Hanley.

Tapes sulcaria, Sowerby.

Tapes sulcaria and *retifera*, Deshayes.

Hab. Indian Ocean.

This species is characterized not only by its strong concentric groove, but by a peculiar violet-ash clouding of the rays.

Species (Fig. 2, a, b, c, Mus. Cuming.)

TAPES LITERATA. *Tap. testa subquadrato-oblonga, posticè declivi-truncata, spadiceo-alba, fusco-nigricante acutè lineari-reticulata, interdum tautum punctata, interdum inquinato-umbulata, concentricè levilirata.*

THE LETTERED TAPES. Shell somewhat squarely oblong, posteriorly slopingly truncated, fawn-white, sharply linearly reticulated with fuscous-black, sometimes only dotted, sometimes smearingly clouded, concentrically finely ridged.


Venus literata, Linneus, Syst. Nat. p. 1135.

Pullastra literata, Sowerby.

Tapes literata, Sowerby.

Hab. Indian Ocean; Australia.

Of this fine species the sculpture is comparatively delicate. In form its chief distinguishing feature is the arched truncated slope of the posterior side.



T A P E S.

PLATE II.

Species 3. (Mus. Cuming.)

TAPES TEXTRIX. *Tap. testā oblongo-ovatā, posticē subtenuatā, pallidē fulvā, lineis undatis violaceo-griseis undique reticulatō, glaberrimā.*

THE WEAVER TAPES. Shell oblong-ovate, posteriorly somewhat attenuated, pale fulvous, everywhere reticulated with waved violet-grey lines, very smooth.

Venus textrix, Chemnitz, Conch. Cab. vol. vii. p. 48. pl. 42. f. 442.

Venus textile, Gmelin.

Venus undulata, Dillwyn.

Pullastra vulgaris, Sowerby.

Tapes textile, Sowerby.

Tapes textrix, Deshayes.

Hab. Indian Ocean; New Holland.

This well-known species may always be recognized by its posteriorly-attenuated oblong form, and textile painting.

Species 4. (Fig. a, b, Mus. Cuming.)

TAPES DESHAYESII. *Tap. testā subquadrato-oblongā, modicē convexā, valde inaequilaterali, fulvescente-lacteā, castaneo-brunneo vel griseo varie oblique reticulatū, concentricē obtuse liratā, liris ad latus posterius plicatis.*

DESHAYES' TAPES. Shell somewhat squarely oblong, moderately convex, very inequilateral, fulvous cream-colour, variously obliquely reticulated with chestnut-brown or grey, concentrically obtusely ridged, ridges plicated on the posterior side.

Venus Deshayesii, Hanley, Wood, Ind. Test. App. pl. 16. f. 35.

Venus radiata, Anton (not of Chemnitz or Brocchi).

Tapes Deshayesii, Sowerby.

Hab. Philippine Islands; Cuming.

Of a peculiar oblong-square form, strongly plicately ridged on the posterior angle.

Species 5. (Fig. a, b, Mus. Cuming.)

TAPES ASPERSA. *Tap. testā oblongo-ovatā, valde inaequilaterali, subdepressē convexā, latere postico obliquē*

rhomboideo, rufescente-lacteā, maculis castaneis interrūptē triradiatā, deinde punctis castaneis aspersā, interdum lineis violaceo-griseis obscurē reticulatā, concentricē obtuse liratā, liris sape confluentibus.

THE SPRINKLED TAPES. Shell oblong-ovate, very inequilateral, rather depressly convex, posterior side obliquely rhomboid, reddish cream-colour, interruptedly three-rayed with chestnut spots, then sprinkled with chestnut dots, sometimes obscurely reticulated with violet-grey lines, concentrically obtusely ridged, ridges often flowing together.

Venus aspersa, Chemnitz, Conch. Cab. vol. vii. p. 44. pl. 42. f. 438.

Venus literata, var., Gmelin.

Tapes aspersa, Sowerby.

Hab. Australia and Philippine Islands.

The rays of chestnut blotches are generally very irregular in this species, whilst the sprinkling of dots is rather conspicuous. In most specimens there are traces of a textile pattern of violet-grey.

Species 6. (Mus. Cuming.)

TAPES QUADRIRADIATA. *Tap. testā suboblongo-ovatā, subventricosā, fulvescente-lacteā, maculis nigrantibus quadriradiatā, liris griseis undique reticulatā, concentricē dense obtuse liratā, margine ventrali tumidiusculo.*

THE FOUR-RAYED TAPES. Shell rather oblong-ovate, slightly ventricose, fulvous cream-colour, four-rayed with blackish spots, everywhere reticulated with grey lines, concentrically densely obtusely ridged, ventral margin rather tumid.

Deshayes, Pro. Zool. Soc. 1853. p. 9. pl. 19. f. 6.

Hab. Philippine Islands; Cuming.

Of a distinct, subventricose form, rather tumidly arched at the ventral margin.

Species 7. (Mus. Cuming.)

TAPES ROTUNDATA. *Tap. testā subgracili-ovatā, utrinque rotundatā, rufescente-spadiceā, maculis castaneis qua-*

TAPES.—PLATE II.

divirradiatā, lineis griseis undique pallidissimè reticulatā, concentricè obtusè liratā, liris planis.

THE ROUNDED TAPES. Shell rather slenderly ovate, rounded on each side, reddish fawn-colour, four-rayed with chestnut blotches, very faintly reticulated throughout with grey lines, concentrically obtusely ridged, ridges flat.

Venus rotundata, Linnaeus, Syst. Nat. p. 1135.

Venus ala-papilionis, Chemnitz.


Venus papilionacea, Lamarek.

Pullastra papilionacea, Sowerby.

Tapes papilionacea, Sowerby.

Hab. Indian Ocean; New Holland.

We are indebted to M. Deshayes for having identified this species, so well known to conchologists by the characteristic name of *papilionacea*, with Linnaeus' *Venus rotundata*.




8.



9.




10.




11.



13.a



12.



13.b

T A P E S.

PLATE III.

Species 8. (Mus. Cuming.)

TAPES UNDULATA. *Tap. testā oblongo-ovatā, subangustā, fulvā, purpureo-violaceo reticulato-undulatā, præsertim ad marginem ventralēm et ad lunulam areamque ligamenti, latere postico concentricè rugoso, antico diuagonaliter obtusè corrugato.*

THE WAVED TAPES. Shell oblong-ovate, rather narrow, fulvous, reticulately waved with purple-violet, especially at the ventral margin and at the lunule and ligamentary area, posterior side concentrically rugose, anterior diagonally obtusely wrinkled.

Venus undulata, Born, Test. Mus. Vind. p. 67.

Venus rimosa, Philippi.

Tapes rimosa, Sowerby.

Tapes undulata, China.

Hab. China.

This species might be taken at first sight to be a variety of *T. testrix*, but it will be found on inspection to be peculiarly characterized on the anterior side by a sculpture of irregular diagonal wrinkles.

Species 9. (Mus. Cuming.)

TAPES GRATA. *Tap. testā suboblongo-ovatā, modicè inæquilaterali, rufescente-spadiceā, maculis castaneis sparsim quadriradiatā, ad umbones purpurascente tinctā, undique concentricè liratā, liris obtusis, eruberrimis, interstitiis lineariter incisis.*

THE PLEASING TAPES. Shell rather oblong-ovate, moderately inequilateral, reddish fawn-colour, sparingly four-rayed with chestnut blotches, tinged with purple at the umboes, concentrically ridged throughout, ridges obtuse, very close-set, with the interstices linearly incised.

DESHAYES, Pro. Zool. Soc. p. 10.

Hab. Philippine Islands; Cuming.

Peculiarly closely ridged with great regularity.

Species 10. (Mus. Cuming.)

TAPES SIMILIS. *Tap. testā ovato-cunneiformi, valde inæquilaterali, fulvescente-alba, lineis cinnereo-griseis undique reticulatā, concentricè confertim obtusè liratā, liris anticis convexis, posticis planutis.*

THE LIKE TAPES. Shell ovately wedge-shaped, very inæquilateral, fulvous-white, reticulated throughout with ash-grey lines, concentrically closely obtusely ridged, anterior ridges convex, posterior flattened.

DESHAYES, Pro. Zool. Soc. 1853, p. 8.

Hab. —?

Rather more than usually curved in growth, with a copious textile of light grey, sometimes faintly rayed.

Species 11. (Mus. Cuming.)

TAPES TURGIDA. *Tap. testā subquadrato-oblongā, posticè turgidā et obtusè angulatā, fulvo-spadiceā, castaneo pallidè quadriradiatā, lineisque griseis pallidis undatā, concentricè undique dense liratā, liris interdum confluentibus, interstitiis excavatis.*

THE TURGID TAPES. Shell somewhat squarely oblong, posteriorly turgid and obtusely angled, fulvous fawn-colour, faintly four-rayed with chestnut, and waved with pale-grey lines, concentrically closely ridged throughout, ridges sometimes flowing together, with the interstices excavated.

Venus turgida, Lamarek, Anim. sans vert. vol. vi. p. 353.

Venus dorsata, Lamarek.

Venus ovulaea, Lamarek.

Tapes turgida, Sowerby.

Hab. New Holland.

Characterized by a broadly truncated, obtusely angled, turgid growth on the posterior side.

Species 12. (Mus. Cuming.)

TAPES OBSCURATA. *Tap. testā ovato-cunneiformi, suboblongā, valde inæquilaterali, squalidè fulvo-rufescente, griseo obscurè quadriradiatā, solidiusculā, regulariter convexā, densè concentricè liratā, liris anticis elevatis, posticis planutatis.*

THE OBSCURED TAPES. Shell ovately wedge-shaped, rather oblong, very inæquilateral, dull fulvous-red, obscurely four-rayed with grey, rather solid, regularly convex, closely concentrically ridged, anterior ridges slightly raised, posterior flattened.

DESHAYES, Pro. Zool. Soc. 1853, p. 9.

Hab. —?

TAPES.—PLATE III.

An evenly convex wedge-shaped shell, with dull obscure traces of colouring.

— — — — —
Species 13. (Fig. *a*, *b*, Mus. Cuming.)

TAPES DURA. *Tap. testā subangustē ovatā, plerumque turgidā, fulvā vel rufescente-spadiceā, maculis castaneis plus minus radiatā, lineis griseis undatis pallidē reticulatā, concentricè obtusè liratā, liris creberrimis.*

THE ROUGH TAPES. Shell rather narrowly ovate, gene-

rally turgid, fulvous or reddish-fawn, more or less rayed with chestnut blotches, faintly reticulated with waved grey lines, concentrically obtusely ridged, ridges very close-set.

Venus dura, Gmelin, Syst. Nat. p. 3292.

Venus rariplamma, Lamarek.

Tapes rariplamma, Sowerby.

Tapes dura, Deshayes.

Hab. Senegal.

An extremely variable species, from Senegal, Le Pegen of Adanson.



15.a



15.b




16.a



17.a



17.b



16.b

T A P E S.

PLATE IV.

Species 14. (Mus. Cuming.)

TAPES ALBA. *Tap. testā oblongo-ovatā, posticē subtruncato-angulatā, sordidē albā, concentricē rugatā, rugis valde irregularibus, ad angulum posticum plicatā.*

THE WHITE TAPES. Shell oblong-ovate, posteriorly somewhat truncalely angled, dull white, concentrically wrinkled, wrinkles very irregular, plicated on the posterior angle.

DESHAYES, Pro. Zool. Soc. 1853, p. 9.

Hab. Swau River, Western Australia.

A very characteristic, opaque-white species, of solid growth, sculptured with irregularly concentric wrinkles.

Species 15. (Fig. a, b, Mus. Cuming.)

TAPES VITULATA. *Tap. testā ovato-cordatā, posticē obliquē attenuatā, gibbosiusculā, albida, fulvo-rubente undique copiosè variegatā, concentricē ruditē liratā, liris anticis corrugato-elevatis, posticis plicato-planatis.*

THE VARIEGATED TAPES. Shell ovately-cordate, posteriorly obliquely attenuated, rather gibbous, whitish, copiously variegated throughout with fulvous-red, concentrically rudely ridged, anterior ridges corrugately raised, posterior plicately flattened.

DESHAYES, MS. in Mus. Cuming.

Hab. —?

I give this species as I find it named by M. Deshayes in Mr. Cuming's cabinet. It comes very near to certain varieties of our British *T. virginea*.

Species 16. (Fig. a, b, Mus. Cuming.)

TAPES RADIATA. *Tap. testā subquadrato-oblongā, posticē compressē flabelliformi, truncato-angulatā, fulvescente-lacteā, nigro obscurē radiatā et copiosē punctatā, interdum usculato-castaneo tinctā, concentricē densē striato-liratā.*

THE RAYED TAPES. Shell somewhat squarely oblong, posteriorly compressly fan-shaped, truncately angled, fulvous cream-colour, obscurely rayed and copiously dotted with black, sometimes tinged with burnt-chestnut, concentrically densely striately ridged.

Venus literata radiata, Chenmitz, Conch. Cab. vol. vii. p. 45. pl. 42. f. 439.

Tapes literata, var., Sowerby.

Tapes radiata, Deshayes.

Hab. Indian Ocean.

Many will hesitate to follow M. Deshayes in separating this from *T. literata*; still it is a very characteristic form of it, always readily recognizable.

Species 17. (Fig. a, b, Mus. Cuming.)

TAPES VIRGINEA. *Tap. testā orato-oblongā, posticē subobliquē truncatā, lacteā, spadiceo-roseo copiosē variegatā, concentricē crebriliratā, liris irregularibus, medio planatis, ad latera plicatis.*

THE VIRGIN TAPES. Shell ovately oblong, posteriorly rather obliquely truncated, cream-colour, copiously variegated with fawn-rose, concentrically closely ridged, ridges irregular, flattened in the middle, plicated at the sides.

Venus virginea, Linnaeus, Syst. Nat. p. 1136.

Venus rhomboides, Pennant.

Cunens fasciatus, Da Costa.

Venus Sarniensis, Turton.

Venerupis Sarniensis, Fleming.

Venerupis virginica, Fleming.


Pullastra virginaea, Brown.

Venus virago, Lovén.


Tapes virginaea, Forbes and Hanley.

Hab. European Seas.


This well-known species of our own shores presents a beautiful variegation of delicate fawn-rose colour.




18.




19.



20.




21.



22.



23.



24.

T A P E S .

PLATE V.

Species 18. (Mus. Cuming.)

TAPES SINUOSA. *Tap. testā subtrigono-ovatā, subaequilaterali, fulvā, purpureo radiatim strigatā et variegatā, concentricè undique confertim lamellato-liratā, latere postico sinuosè contorto.*

THE SINUOUS TAPES. Shell somewhat triangularly ovate, nearly equilateral, fulvous, radiately streaked and variegated with purple, concentrically closely lamellately ridged throughout, posterior side sinuously contorted.

Venus sinuosa, Lamarck, Anim. sans vert. vol. vi. p. 366.

Tapes sinuosa, Sowerby.

Hab. —?

The general aspect of this species is singularly like that of a *Tellina*.

Species 19. (Mus. Cuming.)

TAPES MEROEFORMIS. *Tap. testā trigono-oblongā, subaequilaterali, compressiusculā, lacteā, circa umbones roseo-spadiceo reticulato-nebulatā, concentricè plicato-liratā, liris obtusis, interdum confluentibus, areā ligamenti subexcavatā, latere postico truncato.*

THE MEROE-SHAPED TAPES. Shell triangularly-oblong, nearly equilateral, rather compressed, cream-colour, reticulately clouded about the umboes with rose-fawn, concentrically plicately ridged, ridges obtuse, sometimes flowing together, ligamentary area slightly excavated, posterior side truncated.

SOWERBY, Thes. Conch. vol. i. p. 745. pl. 163. f. 160.

Hab. Australia.

Another abnormal and beautiful species, partaking in some degree of the characters of *Meroë*.

Species 20. (Mus. Cuming.)

TAPES LIRATA. *Tap. testā subelongato-ovatā, solidā, rufescente-spadiceā, maculis castaneis quadriradiatā, lido-griseo in interstitiis undatā, concentricè liratā, liris prominentibus, confertis.*

THE RIDGED TAPES. Shell rather elongately-ovate, solid, reddish-fawn, four-rayed with chestnut blotches,

waved in the interstices with livid-grey, concentrically ridged, ridges prominent, close-set.

Venus lirata, Philippi, Abbild. und Besch. Conch. p. 24. pl. 7. f. 5.

Tapes lirata, Sowerby.

Hab. Philippine Islands; Cuming.

The ridges are of an obtusely callous character, prominent and close-set.

Species 21. (Mus. Cuming.)

TAPES AMABILIS. *Tap. testā orato-oblongā, roseo-spadiceā vel carneā, lineis griseis obscurè reticulato-undatā, concentricè liratā, liris latiusculis, obtusis.*

THE LOVELY TAPES. Shell ovately-oblong, rose fawn-colour, or flesh-colour, obscurely reticulately waved with grey lines, concentrically ridged, ridges rather broad, obtuse.

Venus amabilis, Philippi, Zeitschr. für Malac. 1847, p. 90.

Tapes amabilis, Sowerby.

Hab. —?

The concentric ridges are in this species very conspicuously defined, and the colouring is peculiar.

Species 22. (Mus. Cuming.)

TAPES SULCOSA. *Tap. testā subcordato-oblongā, postice subtruncatā, carneo-lacteā, maculis parviusculis castaneis quadriradiatā, lineis pallidè griseis in interstitiis reticulatā, concentricè densè corrugato-liratā.*

THE GROOVED TAPES. Shell somewhat cordately oblong, posteriorly slightly truncated, flesh-tinted cream-colour, four-rayed with rather small chestnut spots, very faintly reticulated in the interstices with grey lines, concentrically densely wrinkle-ridged.

Venus sulcosa, Philippi, Zeitschr. für Malac. 1847, p. 89.

Tapes sulcosa, Sowerby.

Hab. Australia.

The truncated posterior end of this species is peculiar; the ridges very conspicuous and plicately confluent.

Species 23. (Mus. Cuming.)

TAPES DECLIVIS. *Tap. testā elongato-ovatā, subtrigonā, subæquilaterali, ad latus posticum flexuoso-declivi, fulvo-spadiceā, cinereo-griseo biradiatā et copiosè reticulato-variegatā, concentricè densè liratā, liris ad latera flexuoso-confluentibus.*

THE SLOPING TAPES. Shell elongately ovate, somewhat triangular, nearly equilateral, flexuously sloping at the posterior side, fulvous fawn-colour, two-rayed, and copiously reticulately variegated with ash-grey, concentrically densely ridged, ridges flexuously confluent at the sides.

Tapes declivis, Sowb. Th. Conch. vol. i. p. 680. pl. 145. f. 9.
Hab. Philippine Islands; Cuming.

Peculiarly flexuously extended on the posterior side.


Species 24. (Mus. Cuming.)

TAPES SINENSIS. *Tap. testā ovatā, rufescente-spadiceā, castaneo obscurè radiatā et nebulatā, levigatā vel rugosā, ad latera plicato-corrugatā, rugis versus medium confluentibus, umboibus rosaceis.*

THE CHINA TAPES. Shell ovate, reddish fawn-colour, obscurely rayed and clouded with chestnut, smooth or rugose, plicately wrinkled at the sides, wrinkles confluent towards the middle, umboes pink.

Hab. China.

A characteristic buff-coloured shell, finely plicately wrinkled on each side, the wrinkles being curiously confluent towards the middle of the shell.



TAPE S.

PLATE VI.

Species 25. (Mus. Cuming.)

TAPES LENTIGINOSA. *Tap. testā sublatē ovatā, rufescente-spadiceā, maculis custaneis parvis obscurè radiatā, cinereo-griseo copiosè reticulato-undatā, concentricè dense liratā, liris ad latera flexuose laminatis.*

THE FRECKLED TAPES. Shell rather broadly ovate, reddish-fawn, obscurely rayed with small chestnut spots, copiously reticulately waved with ash-grey, concentrically densely ridged, ridges flexuously lamellated at the sides.

Hab. China; E. L. Layard.

A rather broadly ovate species, in which the ridges are conspicuously defined, and at the sides rise into flexuous laminae.

Species 26. (Fig. *a, b*, Mus. Cuming.)

TAPES MARMORATA. *Tap. testā oblongo-ovatā, posticè subobliquè productā, rufescente-fulvā, livido-griseo varie nebulatā et tinctā, posticè lacerigata, anticè densè plicato-corrugata, umboibus interdum rosaceis.*

THE MARBLED TAPES. Shell oblong-ovate, rather obliquely produced posteriorly, reddish-fulvous, variously clouded and stained with livid-grey, posteriorly smooth, anteriorly densely plicately wrinkled, umboes sometimes tinged with rose.

Cytherea marmorata, Lamarek, Anim. sans vert. vol. vi. p. 261.

Venus variabilis, Philippi.

Tapes laterisulca, Sowerby.

Chione marmorata, Deshayes.

Hab. Bombay.

An extremely variable species; but the colouring is peculiar, and the form and anterior lateral sculpture constant.

Species 27. (Mus. Cuming.)

TAPES MALABARICA. *Tap. testā cordato-ovatā, areuatā, fulvescente-lucteā, lineis cinereo-griseis undatis pallidè*

radiatā et lentiginosā, concentricè obtusè liratā, liris confertis, ad latū posticum flexuosis.

THE MALABAR TAPES. Shell cordately ovate, arched, fulvous cream-colour, faintly waved and freckled with ash-grey lines, concentrically obtusely ridged, ridges close-set, and rather flexuous on the posterior side.

Venus Malabarica, Chemnitz, Conch. Cab. vol. vi. p. 323. pl. 31. f. 324, 325.

Venus gallus, Spengler.

Tapes Malabarica, Sowerby.

Hab. Malabar.

Chiefly distinguished by its arched cordate form, and somewhat flexuous posterior side.

Species 28. (Fig. *a, b*, Mus. Cuming.)

TAPES HANTINA. *Tap. testā subquadrato-ovatā, posticè obliquè truncatā, tumidā, fulvescente-lacteā, olivaceo-griseo varie reticulato-flammatā, concentricè tenuiliratā, liris obtusis vel plicato-planatis, umboibus lividis.*

THE GAPPING TAPES. Shell somewhat squarely ovate, posteriorly obliquely truncated, tumid, fulvous cream-colour, variously reticulately flamed with olive-grey, concentrically finely ridged, ridges obtuse or plicately flattened, umboes livid.

Venus hantina, Lamarek, Anim. sans vert. vol. vi. p. 350.

Venus flammiculata, Lamarek.

Tapes flammiculata, Sowerby.

Chione hantina, Deshayes.

Hab. Australian Seas.

This interesting species varies considerably, both in form and colour, yet there is nevertheless a peculiarity about the flamed painting which renders it easy of recognition.

Species 29. (Mus. Cuming.)

TAPES RIMULARIS. *Tap. testā subquadrato-ovatā vel globosā, sordidè fulvā, livido-griseo tinctā, concentricè liratā, liris ad latū posticum planatis et plicato-confluentibus, latere postico truncato-angulato.*

TAPES.—PLATE VI.

THE CLEFTED TAPES. Shell somewhat squarely ovate or globose, dull fulvous, tinged with livid-grey, concentrically ridged, ridges flattened, and plicately confluent on the posterior side, posterior side truncately angled.

Venus rimularis, Lamarek, Annn. sans vert. vol. vi. p. 367.

Tapes rimularis, Sowerby.

Chione rimularis, Deshayes.

Hab. — ?

This is a form of the preceding species, without marking, in which the ridges are peculiarly flattened and pliately confluent.



T A P E S.

PLATE VII.

Species 30. (Fig. *a, b*, Mus. Cuming.)

TAPES CEYLONENSIS. *Tap. testā ovali, interdum oblongā, crassā, gibbosā, fulvescente-carnea, cinereo-griseo interdum linearī-undulatā et obscurē radiatā, laevigatā, nitente.*

THE CEYLON TAPES. Shell oval, sometimes oblong, thick, gibbous, fulvous flesh-colour, sometimes linearly waved and obscurely rayed with ash-grey, smooth, shining.

SOWERBY, Thes. Conch. vol. i. p. 683. pl. 146. f. 24, 25.
Chione Ceylonensis, Deshayes.

Hab. Ceylon.

The species grouped together in this Plate partake largely of the characters of *Cytherea*, and may be referred to either genus. This and *T. pinguis* are especially connected with *Dione* and *Cytherea*; but, as being arranged in Mr. Cuming's collection by Mr. Sowerby under *Tapes*, I have not thought it worth while to displace them.

Species 31. (Fig. *a, b*, Mus. Cuming.)

TAPES FUMIGATA. *Tap. testā anticē cordatā, posticē obliquē acuminate, fulvescente-lactea vel cinereo-grisea, laevigata, nitente, margine postico concentricē linearisulcato.*

THE SMOKED TAPES. Shell anteriorly heart-shaped, posteriorly obliquely acuminate, fulvous cream-colour or ash-grey, smooth, shining, posterior margin concentrically linearly grooved.

Venus fumigata, Sowerby, Thes. Conch. vol. i. p. 737. pl. 159. f. 152 to 155.

Chione fumigata, Deshayes.

Hab. Australia; Strange.

A smooth shell, of a smoky fulvous or ash-grey colour, delicately carved with concentric lines down the posterior margin. Mr. Sowerby refers this shell to *Venus*, but it partakes more of the character of *Tapes* than the preceding species.

Species 32. (Mus. Cuming.)

TAPES TURGIDULA. *Tap. testā trigono-cordatā, elongato-curratā, inaequilaterali, concentricē irregulariter corrugato-liratā, fulvescente-spudicea, griseo obscurē radiatā et reticulatā, latere antico brevi, subacuminato, postico ample, flexoso, turgido.*

THE SWOLLEN TAPES. Shell triangularly heart-shaped, elongately curved, inequilateral, concentrically irregularly wrinkle-ridged, fulvous fawn-colour, obscurely rayed and reticulated with grey, anterior side short, rather acuminate, posterior large, flexuous, swollen.

DESHAYES, Pro. Zool. Soc. 1853, p. 8.

Hab. Philippine Islands; Cuming.

This is a true *Tapes*, of a characteristic curved form, rather acuminate anteriorly, whilst the posterior side is comparatively large and swollen.

Species 33. (Fig. *a, b*, Mus. Cuming.)

TAPES PINGUIS. *Tap. testā ovato-cordatā, interdum globosā, crassā, conspicue turgida, fulvescente-luctea nata curvata, nigricante vel fusco nebulata et radiatum variegata, laevigata, nitente.*

THE FAT TAPES. Shell ovately cordate, sometimes globose, thick, conspicuously turgid, fulvous cream-colour or flesh-colour, clouded and radiately variegated with black or brown, smooth, shining.

Tenus pinguis, Chemnitz. Conch. Cab. vol. vi. p. 355. pl. 34. f. 355 to 357

Venus triradiata, Chemnitz.

Venus nebulosa, Chemnitz.

Venus opima, Gmelin.

Pallastria nebulosa, Sowerby.

Tapes pinguis, Sowerby.

Chione pinguis, Deshayes.

Hab. Indian Ocean.

Of a conspicuous globosely turgid form, solid and smooth.




34 b




34 a



35.



36 a




37 a



37 b



36 b



38 a.



38 b.



39.

T A P E S.

PLATE VIII.

Species 34. (Fig. a, b, Mus. Cuming.)

TAPES ORIENTALIS. *Tap. testā ovatā, subventricosā, fulvescente-spadiceā, fumeo-fusco plus minus tinctā et interrupte reticulatā, concentricē corrugato-liratā, liris obtusis, hic illuc confluentibus, posticē plicato-expansis.*

THE EASTERN TAPES. Shell ovate, slightly ventricose, fulvous-fawn, more or less stained and interruptedly reticulated with smoky-brown, concentrically corrugately ridged, ridges obtuse, here and there confluent, posteriorly plicately expanded.

Hab. Bombay.

Of this apparently undescribed species Mr. Cuming possesses several varieties, closely allied in sculpture and general aspect.

Species 35. (Mus. Cuming.)

TAPES OCCIDENTALIS. *Tap. testā ovatā, posticē subflexuosā, fulvescente-lacteā, nigricante interrupte reticulato-nebulatā, concentricē corrugato-liratā, liris obtusis, irregulariter confluentibus, posticē plicato-expansis.*

THE WESTERN TAPES. Shell ovate, posteriorly slightly flexuous, yellowish cream-colour, interruptedly reticulately clouded with black, concentrically corrugately ridged, ridges obtuse, irregularly confluent, posteriorly plicately expanded.

Hab. Guadalupe, West Indies.

Singularly like the preceding species in design of sculpture, though the two species are from opposite hemispheres. In form and colour they are obviously distinct.

Species 36. (Fig. a, b, Mus. Cuming.)

TAPES PHILIPPIN. *Tap. testā trigono-ovatā, compressā, posticē subacuminatā, rufescente-spadiceā, griseo pallide reticulatā, concentricē liratā, liris confertis, obtusis, hic illuc confluentibus.*

PHILIPPI'S TAPES. Shell triangularly ovate, compressed, posteriorly rather acuminated, reddish fawn-colour, faintly reticulated with grey, concentrically ridged, ridges close-set, obtuse, here and there confluent.

Venus striata, pars, Sowerby, Thes. Conch. vol. i. p. 735.
Chione Philippin, Deshayes.

Hab. China. Philippine Islands.

This species, which M. Deshayes refers to his genus *Chione*, appears to me to be an undoubted *Tapes*.

Species 37. (Fig. a, b, Mus. Cuming.)

TAPES AUREA. *Tap. testā oblongo-cuneiformi, albida vel fulvescente, griseo vel fusco promiscue flammata vel tenuireticulatā, concentricē tenuiliratā, liris confertis, hic illuc confluentibus, latere antico abrupte abbreviato, medio turgido, postico plus minus acuminato.*

THE GOLDEN TAPES. Shell oblong wedge-shaped, white or fulvous, promiscuously flamed and finely reticulated with grey or brown, concentrically finely ridged, ridges close-set, here and there confluent, anterior side abruptly shortened, middle swollen, posterior side more or less acuminate.

Venus aurea, Gmelin, Syst. Nat. p. 3288.

Tellina fasciata, Lister.

Venus edulis, Chemnitz.

Venus aenea, Turton.

Venus nitens, Turton.

Pullastra aurea, Brown.

Tapes aurea, Forbes and Hanley.

Hab. European Seas.

This well-known *Tapes* of our own shores may be readily distinguished by its peculiarly turgid wedge-like form.

Species 38. (Fig. a, b, Mus. Cuming.)

TAPES KOCHII. *Tap. testā anticē globoso-cordatā, turgida, posticē flexuoso-acuminatā, albida, cæruleo-griseo vel fumeo-fusco reticulato-variegata aut nebulata, variegata, nivata, obscure concentricē striatā.*

KOCH'S TAPES. Shell anteriorly globosely cordate, swollen, posteriorly flexuously acuminated, whitish, reticulately variegated or clouded with blue-grey or smoky-brown, smooth, shining, obscurely concentrically striated.

Venus Kochii, Philippi, Abbild. und Besch. Conch. vol. i. p. 3, pl. 1, f. 5.

Chione Kochii, Deshayes.

Hab. —?

TAPES.—PLATE VIII.

More flexuously acuminate than the preceding species, with the anterior portion of the shell swollen into an abrupt convexity.

Species 39. (Mus. Cuming.)

TAPES TABA. *Tap. testā ovato-trigonā, subcuneiformi, versus marginem compressā, pallidē spadiceā, fuscescente interrupte reticulato-triradiatā, tærigatā, nitente, latere antico rotundato, postico subacuminato.*

THE BEAN TAPES. Shell ovately triangular, somewhat wedge-shaped, compressed towards the margin, light fawn-colour, interruptedly reticulately three-rayed with light-brown, smooth, shining, anterior side rounded, posterior rather acuminate.

Hab. —?

This species partakes of the characters of *T. Kochii* and *fumigata*, without being satisfactorily referable to either species.



T A P E S .

PLATE IX.

Species 40. (Mus. Cuming.)

TAPES LUZONICA. *Tap. testā oratā, fulvescente, purpu-*
rascente-fusco copiosè reticulato-aspersā et obscurē
radiatā, concentricè liratā, liris prouinentibus, corrugato-
confuentibus, latere postico fulvescente-maculato.

THE LUZON TAPES. Shell ovate, fulvous, copiously re-

ticulately sprinkled, and obscurely rayed with purple-
brown, concentrically ridged, ridges prominent, cor-
rugately confluent, posterior side fulvous-spotted.

SOWERBY, Thes. Conch. vol. i. p. 687. pl. 149. f. 100, 101.
Hab. Island of Luzon, Philippines; Cuming.

A species in which the ridges are prominent and the
colouring unusually copious.

Species 41. (Mus. Cuming.)

TAPES KRAUSSI. *Tap. testā abruptè cuneiformi, crassā,*
gibbosiusculā, spadiceā, griseo radiatione pallidè tinctā,
concentricè obtuse corrugetā, rugis latè planulatis, ferè
obsoletis.

KRAUSS'S TAPES. Shell abruptly wedge-shaped, thick,
rather gibbous, fawn-colour, faintly radiately tinged
with grey, concentrically obtusely wrinkled, wrinkles
broadly flattened, almost obsolete.

Chione Kraussi, Deshayes, Cat. Conch. Brit. Mus. p. 154.
Venus paupercula, var., Krauss.

Hab. Natal.

Of a peculiarly shortened wedge-like form, faintly
coloured, and rather obscurely obtusely sculptured.

Species 42. (Mus. Cuming.)

TAPES BICOLORATA. *Tap. testā oblongo-ovali, valde inae-*
quilateralī, sordidè lacteū, nigrante-fusco et auranti-
o-rufo sparsim reticulato-maculata, concentricè li-
ratā, liris plicato-planatis, hic illuc confluentibus.

THE TWO-COLOURED TAPES. Shell oblong-oval, very in-
equilateral, dull cream-colour, sparingly reticulately
blotched with blackish-brown and orange-red, con-
centrically ridged, ridges plicately flattened, here and
there confluent.

Hab. —?

The curious mingling of blackish-brown and orange-red

in the colouring of this shell appears to be of specific im-
portance.

Species 43. (Fig. 43 and 46 a, Mus. Cuming.)

TAPES JAPONICA. *Tap. testā subquadrato-trigonā, medio*
turgidā, posticè flexuoso-angulatā, fulvescente-lactea,
griseo lineari-undata, interdum obscurē radiatā, con-
centricè liratā, liris callosis, subacutè reflexis.

THE JAPAN TAPES. Shell somewhat squarely triangular,
turgid in the middle, flexuously angled posteriorly,
fulvous cream-colour, linearly waved with grey, some-
times obscurely rayed, concentrically ridged, ridges
callous, rather sharply reflected.

Venus Japonica, Gmelin, Syst. Nat. p. 3279.

Venus striata, pars, Sowerby.

Chione Japonica, Deshayes.

Hab. China, Japan.

Of a peculiar callous ivory-like substance, more or less
profusely painted with zigzag waves of ash-grey.

Species 44. (Fig. a, b, Mus. Cuming.)

TAPES TRISTIS. *Tap. testā obliquū subcordatā, subcom-*
pressā, posticè flexuoso-productā, fulvā, violaceo-cinereo
undique tinctā, presertim versus marginem, concentri-
cè liratā, liris latè obtusis, reflexis, intersticiis exca-
vatis.

THE DULL TAPES. Shell obliquely subcordate, rather
compressed, posteriorly flexuously produced, fulvous,
tinged throughout with violet-ash, especially towards
the margin, concentrically ridged, ridges broadly ob-
tuse, reflected, with the interstices excavated.

Venus tristis, Lamarek, Anim. sans vert. vol. vi. p. 366.

Chione tristis, Deshayes.

Chione regularis, Deshayes.

Hab. Australia and Philippine Islands.

After a careful examination of the type of M. Deshayes'
Chione regularis, I have no hesitation in referring it to
this comparatively colourless species.

Species 45. (Fig. a, b, Mus. Cuming.)

TAPES STRIATA. *Tap. testā anticè tumido-cordatā, posticè*

TAPES.—PLATE IX.

flexuoso-acuminatā, fulvā, pallidē griseo sc̄epissimē li-
neari-undatā et lentiginosā, concentricē densē liratā,
liris subcallosis, hic illic confluentibus.

THE STRIATED TAPES. Shell anteriorly tumidly cordate, posteriorly flexuously acuminate, fulvous, frequently linearly waved and freckled with pale-grey, densely concentrically ridged, ridges rather callous, here and there confluent.

Venus striata, Chemnitz, Conch. Cab. vol. vi. p. 358. pl. 34.
f. 365, 366.

Venus Labuana, Adams and Reeve.

Chione striata, Deshayes.

Hab. Eastern Seas.

Resembling *T. Kochii* in form; but instead of being smooth, the shell is densely concentrically ridged throughout.

Species 46. (Fig. 46 b, Mus. Cuming.)

TAPES BIRADIATA. *Tap. testā suboblongo-ovatā, lacteā,*
pallidē griseo linearī-reticulatā, et castaneo conspicuē
biradiatā, concentricē liratā, liris prominulis, callosis,
subreflexis.

THE TWO-RAYED TAPES. Shell rather oblong-ovate, cream-colour, linearly reticulated with pale-grey, and conspicuously two-rayed with chestnut, concentrically ridged, ridges rather prominent, callous, slightly reflected.

DESHAYES, Pro. Zool. Soc. 1853, p. 9.

Hab. Philippine Islands; Cuming.

A characteristic oval species, blotched on a faint grey-waved ground, with two broad rays of chestnut blotches.

The shell represented at Fig. 46a, originally placed with this in Mr. Cuming's collection, is a variety of *T. Japonica*.



47



48



49



50.



51



52



53.a



53.b



54



55.

T A P E S.

PLATE X.

Species 47. (Mus. Cuming.)

TAPES LIVIDA. *Tap. testā suboblongo-ovatā, tenui, compressā, livido-fulvū, radiatim tenuiliratā, extremitate posticā lavigatā, subangulatā.*

THE LIVID TAPES. Shell rather oblong-ovate, thin, compressed, livid fulvous, radiately thinly ridged, posterior end smooth, slightly angled.

Hab. —?

A small, rather fragile shell, ridged in a direction radiating from the umboes.

Species 48. (Mus. Cuming.)

TAPES VERNICOSA. *Tap. testā elongato-ovatā, crassā, compressā, fulvescente-lacteā, aurantio-rubo lentiginosā, maculis castaneis interruptis quadriradiatā, lavigatā, nitente, ad marginem posticum plicato-sulcatā.*

THE VARNISHED TAPES. Shell elongately ovate, thick, compressed, fulvous cream-colour, freckled with orange-red, and four-rayed with interrupted chestnut blotches, smooth, shining, plicately grooved at the posterior margin.

Hab. —?

This species, with a smooth porcellaneous surface, is characterized by a style of painting which belongs commonly to ridged species.

Species 49. (Mus. Cuming.)

TAPES POLITA. *Tap. testā subelongato-ovatā, ventricosā, rubescens-falcā, pallidē griseo apertē linearī-undulatā, umbonibus sanguineis, lavigatā, nitente, versus marginem obscurē plicato-liratā.*

THE POLISHED TAPES. Shell somewhat elongately ovate, ventricose, reddish-fulvous, openly linearly waved with pale-grey, umboes blood-red, smooth, shining, obscurely plicately ridged towards the margin.

SOWERBY, Thes. Conch. vol. i. p. 682. pl. 145. f. 15, 16.
Hab. Sydney, New Holland.

At first sight this species has very much the appearance of a young *T. textrix*.

Species 50. (Mus. Cuming.)

TAPES AMPHIDESMOIDES. *Tap. testā suborbiculari-ovatā, medio turgido-convergatā, fulvo-ferruginea, cinereo-fusco punctato-lentiginosā, umbonibus carneo-roseis, circa umboes densē striatā, deinde corrugato-liratā, liris planatis et obliquē decliri-confluentibus, sub lente radiatā undique minutissimè striatā.*

THE AMPHIDESMA-LIKE TAPES. Shell somewhat orbicularly ovate, turgidly convex in the middle, fulvous-rust, dot-freckled with ash-brown, umboes flesh-pink, around the umboes densely striated, then corrugately ridged, ridges flattened, and obliquely slopingly confluent, beneath the lens radiately very minutely striated throughout.

Hab. Red Sea.

It will be seen by an observance of the above characters, that this is a species of quite a distinct type.

Species 51. (Mus. Cuming.)

TAPES FERRUGINEA. *Tap. testā ovato-oblongā, compressiusculā, lacteā, ferrugineo promiscuè undulato-nublatā, concentricè liratā, liris plicato-planatis, valde irregularibus, hic illuc bifurcatis et confluentibus.*

THE RUST-STAINED TAPES. Shell ovately oblong, rather compressed, cream-colour, promiscuously wave-clouded with rust, concentrically ridged, ridges plicately flattened, very irregular, here and there bifurcate and confluent.

Hab. Philippine Islands; Cuming.

Approximating in character to the Bombay *T. orientalis*, but certainly distinct.

Species 52. (Mus. Cuming.)

TAPES ARCTICA. *Tap. testā subcuneiformi-ovatā, compressā, anticē abruptē declivi, posticē attenuato-productā, calcareo-albā, epidermide cornea cinerea lavigatā nitente indutā, ad umboes erosā, deinde irregulariter subuplicatā.*

THE ARCTIC TAPES. Shell somewhat cuneiformly ovate,

TAPES.—PLATE X.

compressed, anteriorly abruptly sloping, posteriorly attenuately produced, chalk-white, covered with a smooth shining ash-coloured horny epidermis, eroded at the umbones, then irregularly slightly plicated.

Hab. Arctic Seas.

A distinct and quite abnormal form of *Tapes*, if *Tapes* at all.

Species 53. (Fig. *a, b*, Mus. Cuming.)

TAPES FLORIDELLA. *Tap. testā subangustē ovatā, utrinque subobliquè attenuatā, medio turgidā, fulvescente-lacteā, griseo profusè reticulato-lentiginosā, plus minus distinctè intensè sex-radiatā, concentricè densè subtiliter corrugato-liratā.*

THE BLOOMING TAPES. Shell rather narrowly ovate, somewhat obliquely attenuated on either side, swollen in the middle, fulvous cream-colour, profusely reticulately freckled with grey, more or less distinctly darkly six-rayed, concentrically densely finely wrinkle-ridged.

Venus floridella, Deshayes, Anim. sans vert. vol. vi. p. 365.

Tapes floridella, Deshayes.

Hab. —?

Of a peculiar form, minutely elaborated in respect both of colour and sculpture.

Species 54. (Mus. Cuming.)

TAPES PETALINA. *Tap. testā subangustē ovatā, ventri-*

cosā, fulvescente-lacteā, ferrugineo et violaceo-griseo variè nebulatā et reticulatā, pallidè radiatā, latere antico concentricè tenuè lirato et sulcato, postico lirato-gato.

THE PETAL-LIKE TAPES. Shell rather narrowly ovate, ventricose, fulvous cream-colour, variously clouded and reticulated with rust and violet-grey, faintly rayed, anterior side concentrically finely ridged and grooved, posterior smooth.

Venus petalina, Lamarck, Anim. sans vert. vol. vi. p. 365.

Venus Beudanti, Payraudeau.

Tapes petalina, Sowerby.

Hab. Mediterranean Sea.

Resembling the preceding species in general aspect, but it will be found on comparison that *T. petalina* is smooth on the posterior half of the shell.

Species 55. (Mus. Cuming.)


TAPES TENUISTRATI. *Tap. testā abruptè cuneiformi, posticè subexpansā, albida, pallidè rosaceo-cinereo latiradiatā, concentricè liratā, liris confertis, hic illic confluentibus, ad latus posticum obtusè planatis.*

THE FINELY-STRIPED TAPES. Shell abruptly wedge-shaped, posteriorly rather expanded, whitish, broadly rayed with pale pinkish ash-colour, concentrically ridged, ridges close-set, here and there confluent, obtusely flattened on the posterior side.

SOWERBY, Thes. Conch. vol. i. p. 687. pl. 148. f. 78, 79.

Hab. —?

A very delicately-coloured species, without any indication of marking.



T A P E S.

PLATE XL.

Species 56. (Fig. *a, b*, Mus. Cuming.)

TAPES INDICA. *Tap. testā subquadrato-oblongā vel rhomboidē, fulvā, ferrugineo tinctā, cinereo-fusco variè radiatim flammatā, umboibus carneo-rosaceis, undique confertim decussatā, ad latera nodoso-crenulatā.*

THE INDIAN TAPES. Shell somewhat squarely oblong or rhomboid, fulvous, tinged with rust, variously radiately flamed with ash-brown, umboes tinged with flesh-pink, closely decussated throughout, nodosely crenulated at the sides.

SOWERBY, Thes. Conch. vol. i. p. 694. pl. 151. f. 146, 147.

Hab. Indian Ocean.

The European *T. decussata* is very closely represented in the Indian Ocean by this species. In sculpture the two species are almost identical, the present having, however, a more tropical look in being more sharply crenulated, with the colour a deeper pink at the umboes.

— — — — —
Species 57. (Fig. *a, b*, Mus. Cuming.)

TAPES DECUSSATA. *Tap. testā subquadrato-oblongā, fulvo-ferruginea, cinerea-fusco promiscue radiatim lenticulosā, undique confertim decussatā, ad latera obtuse crenulatā.*

THE DECUSSATED TAPES. Shell somewhat squarely oblong, fulvous-rust, promiscuously radiately freckled with ash-brown, closely decussated throughout, obtusely crenulated at the sides.

Venus decussata, Linnaeus, Syst. Nat. p. 1135.

Venus literata, Pennant.

Venus deflorata, Born.

Cuneus reticulatus, Da Costa.

Venus obscura, Gmelin.

Venus sanguinolenta, Gmelin.

Venus florida, Poli.

Pullastra decussata, Anton.

Venerupis decussata, Maegillivray.

Tapes decussata, Forbes and Hanley.

Hab. European Seas.

Of a duller rusty hue than the preceding species, with the sculpture more obtuse.

— — — — —
Species 58. (Fig. *a, b*, Mus. Cuming.)

TAPES PULLASTRA. *Tap. testā quadrato-oblongā, interdum latiusculā, albida, ferrugineo tinctā, maculis*

rinereis posticè radiatā, undique minutè obsoletè decussatā, hic illie quasi lirigatā, areā posticā plicato-corrugatā.

THE DUSKY TAPES. Shell squarely oblong, sometimes rather broad, white, tinged with rust, posteriorly rayed with ash blotches, minutely obsoletely deensated throughout, here and there as if smooth, posterior area plicately wrinkled.

Venus pullastræ, Montagu, Test. Brit. p. 125.

Venus perforans, Montagu.

Venus Senegalensis, Dillwyn.

Venerupis perforans, Lamarek.

Venerupis nucleus, Lamarek.

Venus palustris, Mawe.

Venerupis pullastræ, Fleming.

Venerupis vulgaris, Sowerby.

Venus vulgaris, Broderip.

Pullastra vulgaris, Brown.

Pullastra perforans, Brown.

Venus plagia, Jeffreys.

Tapes pullastræ, Forbes and Hanley.

Hab. European and North African Seas.

In this well-known species of our shores the decussation is very elaborate, though nearly obsolete in places. The posterior area of the shell is especially characterized by a textile of wavy wrinkles.

— — — — —
Species 59. (Mus. Cuming.)

TAPES INTERMEDIA. *Tap. testā obliquè quadratā, posticè obtuse declivi-angulatā, fulvescente-ferruginea, concentricè densè liratā, liris anticis subreflexis, medianis profusè bifurcali confluentibus, posticis obtuse planitis.*

THE INTERMEDIATE TAPES. Shell obliquely quadrate, posteriorly obtusely slopingly angled, fulvous rust-colour, concentrically densely ridged, anterior ridges somewhat reflected, middle ridges profusely bifurcately confluent, posterior ridges obtusely flattened.


Venus intermedia, Quoy and Gaimard, Voy. de l'Astrolabe, Moll. pl. 84. f. 9, 10.

Venus Largillierti, Philippi.

Tapes intermedia, Sowerby.

Hab. New Zealand.

A remarkably sculptured species possessing the general character of this section of the genus, but yet wanting the decussation.




60.



61




62.



63



64a.



64b



65.

T A P E S .

PLATE XII.

Species 60. (Mus. Cuming.)

TAPES PHILIPPINARUM. *Tap. testā oblongo-ovatā, gibbā, valde inaequilaterali, antice abbreviatā, truncatā, fulvā, fuscō variegatā et obscurē radiatā, radiatim liratā, liris posticē angulatis, ad latera crispato-decussatis.*

THE PHILIPPINE TAPES. Shell oblong-ovate, gibbous, very inequilateral, anteriorly very short, truncated, fulvous, variegated and obscurely rayed with brown, radiately ridged, ridges posteriorly angled, crisply decussated at the sides.

Venus Philippinarum, Adams and Reeve, Moll. Voy. Samarang, p. 79, pl. 22. f. 10.

Tapes Philippinarum, Sowerby.

Hab. Philippine Islands; Beleher.

Remarkable for its very short truncated growth on the anterior side.

Species 61. (Mus. Cuming.)

TAPES PUNICEA. *Tap. testā ovatā, convexiusculā, latere antice brevi, obuso, deinde attenuato, postico latiore, fulvā, fuscō irregulariter punctulā, obscurē radiatā, maculis parris albis irregulariter aspersā, intus purpureo-roseā, circa umbones violaceā, undique dense largigato-decussatā, liris posticis majoribus, distantioribus.*

THE SCARLET TAPES. Shell ovate, slightly convex, anterior side short, obtuse, then attenuated, posterior broader, fulvous, irregularly dotted with brown, obscurely rayed, irregularly sprinkled with small white spots, purple-rose within, violet about the umbones, densely smoothly decussated throughout, the posterior ridges being larger and more distant.

DESHAYES, Pro. Zool. Soc. 1853, p. 10.

Tapes variegata, Sowerby.

Hab. Philippine Islands; Cuming.

I give M. Deshayes' *T. punicea*, but it is much to be doubted whether the peculiarly broadly ovate form here noted is sufficient to distinguish it, as a species, from *T. variegata*, with which it agrees in other respects.

Species 62. (Mus. Cuming.)

TAPES CINEREA. *Tap. testā subaugustē ovatā, posticē obliquē truncatā, fulvescente-lacteā, griseo et ferrugineo-fuscō variegatā, undique densē decussatā, liris posticē angulatis, ad latera tenuē serrato-nodulatis.*

sparsim posticē radiatā, undique densē decussatā, liris lateralibus subasperis.

THE ASH-COLOURED TAPES. Shell rather narrowly ovate, posteriorly obliquely truncated, ash-white, sparingly rayed on the posterior side with purple-grey blotches, densely decussated throughout, lateral ridges rather rough.

DESHAYES, Pro. Zool. Soc. 1853, p. 11.

Hab. —?

The radiated painting on the posterior side of this shell assumes the importance of a specific character.

Species 63. (Mus. Cuming.)

TAPES DENTICULATA. *Tap. testā ovato-rhomboideā, crassiusculā, sordidē fulvā, purpureo-griseo obscurē radiatā et punctulā, undique irregulariter rugoso-decussatā, liris lateralibus distantioribus, serrato-nodulatis.*

THE TOOTHED TAPES. Shell ovately rhomboid, rather thick, dull fulvous, obscurely rayed and dotted with purple-grey, irregularly rugosely decussated throughout, lateral ridges more distant, serrately noduled.

SOWERBY, Thes. Conch. vol. i. p. 694. pl. 150. f. 114.

Hab. —?

Distinguished by a rough irregular kind of decussation, which at the sides is serrately noduled.

Species 64. (Fig. a, b, Mus. Cuming.)

TAPES VARIEGATA. *Tap. testā oblongo-ovatā, posticē obliquē truncatā, fulvescente-lacteā, griseo et ferrugineo-fuscō variegatā, undique densē decussatā, liris posticē angulatis, ad latera tenuē serrato-nodulatis.*

THE VARIEGATED TAPES. Shell oblong-ovate, posteriorly obliquely truncated, fulvous cream-colour, variegated with grey and rust-brown, closely decussated throughout, ridges posteriorly angled, finely serrately noduled at the sides.

SOWERBY, Thes. Conch. vol. i. p. 696. pl. 151. f. 133 to 138.

Hab. Philippine Islands; Cuming.

Variously painted, but often leaving the cream-coloured

TAPES.—PLATE XII.

ground with the appearance of a sprinkling of small spots.
The interior is generally tinged with rose.

Species 65. (Mus. Cuming.)

TAPES GALACTITES. *Tap. testā subelongato-oblongā, antīcē obtusē angulatā, candidā, undique lāvigato-decussatā, lateribūs ambobus rotundatis.*

THE MILKY TAPES. Shell rather elongately oblong, anteriorly obtusely angled, white, smoothly decussated throughout, both sides rounded.

Venus galactites, Lamarck, Anim. sans vert. vol. vi. p. 359.

Tapes galactites, Sowerby.

Hab. New Holland; New Zealand.

A solid oblong white species, with very much of the general aspect of a *Cypricardia*.



T A P E S.

PLATE XIII.

Species 66. (Mus. Cuming.)

TAPES FABAGELLA. *Tap. testū oblongo-ovatū, compressiusculū, valde inaequilateralī, antīcē brevi, subattenuatā, trapezoideo-productā, truncatā, candidā, concentricē exilissimē striatā, striis hic illic confluentibas, ad aream posticam corrugato-plicatis.*

THE LITTLE BEAN TAPES. Shell oblong-ovate, rather compressed, very inequilateral, anteriorly short, somewhat attenuated, posteriorly trapezoidally produced, truncated, white, concentrically very finely striated, striae here and there confluent, wrinkle-plaited on the posterior area.

DESHAYES, Pro. Zool. Soc. 1853, p. 10.

Hab. New Zealand.

Of a peculiar trapezoidal form, colourless, but exquisitely sculptured.

Species 67. (Mus. Cuming.)

TAPES SEMIDECCUSSATA. *Tap. testū subquadrato-ovatā, valde inaequilateralī, posticē latē truncatā, fulvescente-lacteā, maculis magnis irregularibus rufis posticē maculatā, medio radiatim plano-liratā, ad latera nodoso-decussatā.*

THE HALF-DECUSSATED TAPES. Shell somewhat squarely ovate, very inequilateral, posteriorly broadly truncated, fulvous cream-colour, marbled on the posterior side with large irregular red blotches, radiately flatly ridged in the middle, nodosely decussated at the sides.

Tapes semidecussata, Deshayes, MS. in Mus. Cuming.

Tapes Japonica, Deshayes (not *Tenus Japonica*, Gmelin, *Chioue Japonica*, Deshayes, *Tapes Japonica*, nobis), Pro. Zool. Soc. 1853, p. 10.

Hab. Japan; Dr. Siebold.

This runs very closely into some varieties of that very variable species, *T. variegata*.

Species 68. (Mus. Cuming.)

TAPES VIOLASCENS. *Tap. testū suboblongo-ovatā, antīcē brevissimā, posticē declivi-rotundatā, albida, violaceo-griseo aut fulvescente radiatā, circa umbones purpureā, undique decussatim liratā, interstitiis longitudinalibus subprofundē incisis.*

THE VIOLET TAPES. Shell somewhat oblong-ovate, anteriorly very short, posteriorly slopingly rounded, whitish, rayed with violet-grey or light-brown, purple about the umboes, denticulately ridged throughout, longitudinal interstices rather deeply engraved.

DESHAYES, Pro. Zool. Soc. 1853, p. 10.

Hab. —?

A very characteristic species, especially in respect of form and colouring.

Species 69. (Mus. Cuming.)

TAPES DACTYLOIDES. *Tap. testā oblongā, gibbosusculā, valde inaequilateralī, rufescente-ferruginea, intus posticē violacea, concentricē irregulariter plicata-corrugata, rugis ad latus posticum crispato-serratis.*

THE FINGER-LIKE TAPES. Shell oblong, rather gibbose, very inequilateral, reddish rust-colour, posteriorly violet within, concentrically irregularly plicately wrinkled, wrinkles on the posterior side, crisply serrated.

SOWERBY, Thes. Conch. vol. i. p. 691, pl. 150. f. 129.

Hab. Senegambia.

The four species figured in the centre of this plate (Sp. 66, 69, 72, 75) are of a distinct, peculiarly finely wrinkled type.

Species 70. (Mus. Cuming.)

TAPES TEXTURATA. *Tap. testā subtrigono-ovatā, fulva, rufa linearī-reticulatā, concentricē liratā, liris obtusis, levibus, ad latera confertis, undulatis.*

THE WOVEN TAPES. Shell somewhat triangularly ovate, fulvous, linearly reticulated with red, concentrically ridged, ridges obtuse, smooth, crowded and a little waved at the sides.

Tenus texturata, Lamarck, Anim. sans vert. vol. vi. p. 355.

Tapes texturata, Sowerby.

Hab. Mediterranean.

Closely concentrically sculptured with smooth obtuse ridges, which at the sides, especially the posterior side, are crowded and a little waved.

TAPES.—PLATE XIII.

Species 71. (Mus. Cuming.)

TAPES GEOGRAPHICA. *Tap. testā oblongā, compressā, posticē rhomboideā, truncatā, fulvescente-lacteā, castaneo-apertē linearī-reticulatā, undique subtilissimē decussatā, ad aream posticam rugosā.*

THE GEOGRAPHIC TAPES. Shell oblong, compressed, posteriorly rhomboid, truncated, fulvous cream-colour, openly linearly reticulated with chestnut, very finely decussated throughout, rough on the posterior area.

Venus geographica, Chemnitz, Conch. Cab. vol. vii. p. 45. pl. 42. f. 410.

Venus literata, Poli (not of Linnaeus).

Tapes geographica, Sowerby.

Hab. Mediterranean.

A pretty delicately-sculptured species, distinguished by the open character of its reticulated painting.

Species 72. (Mus. Cuming.)

TAPES CORRUGATA. *Tap. testā oblongo-ovatā, ad latera rotundatā, crassā, calcareo-albā, intus posticē violaceā, concentricē rūdē corrugatā, rugis posticē irregulariter plicatis.*

THE WRINKLED TAPES. Shell oblong-ovate, rounded at the sides, thick, chalk-white, posteriorly violet within, concentrically rudely wrinkled, wrinkles posteriorly irregularly plaited.

Venus corrugata, Deshayes, Anim. sans vert. vol. vi. p. 351.
Tapes corrugata, Deshayes.

Hab. —?

A solid opaque chalk-white shell, more or less roughly concentrically wrinkled.

Species 73. (Mus. Cuming.)

TAPES BRUGUIERI. *Tap. testā suboblongo-ovatā, tenui, albā, rufescente-fulvo latiradiatā, obscurē punctatā, radiatum plano-liratā, liris lateralibus decussatim crenatis.*

BRUGUIÈRE'S TAPES. Shell rather oblong-ovate, thin, white, broadly rayed with reddish-fulvous, obscurely

dotted, radiately flatly ridged, lateral ridges decussately crenated.

Venus Bruguieri, Hanley, Pro. Zool. Soc. 1845, p. 21.

Tapes Bruguieri, Sowerby.

Hab. Ceylon.

A very delicate neatly-sculptured species.

Species 74. (Mus. Cuming.)

TAPES FLORIDA. *Tap. testā subtrigono-ovatā, ventricosā, fulvo-spadiceā, fuscescente obscurē linearī-reticulatā, præsertim ad latūs posticum, umbonibus rosaceis, concentricē tenuē lirato-striatā, striis hic iltic confluentibus.*

THE FLORID TAPES. Shell somewhat triangularly ovate, ventricose, fulvous-drab, obscurely linearly reticulated with brown, especially on the posterior side, umboes tinged with rose, concentrically finely ridge-striated, striae here and there confluent.

Venus florida, Lamarck, Anim. sans vert. vol. vi. p. 364.

Venus lœta, Poli (not of Linnaeus).

Venus virginea, var., Philippi.

Tapes florida, Sowerby.

Hab. Mediterranean.

A comparatively smooth shell, faintly coloured, but tinged with flesh-rose about the umboes.

Species 75. (Mus. Cuming.)

TAPES CUMINGII. *Tap. testā subelongatā, compressiusculā, posticē declivi-attenuatā, calcareo-albā, violaceo interrūptè radiatā, undique minutè divaricato-corrugatā.*

CUMING'S TAPES. Shell rather elongated, rather compressed, posteriorly slopingly attenuated, chalk-white, interruptedly rayed with violet, everywhere minutely divaricately wrinkled.

SOWERBY, Thes. Conch. vol. i. p. 690. pl. 150. f. 128.

Hab. Philippine Islands; Cuming.

An extremely pretty species, minutely divaricately wrinkled and delicately rayed with violet.

T A P E S .

	Plate.	Species.		Plate.	Species.
<i>aenea</i> (<i>Venus</i>), Turton	VIII.	37	<i>laeta</i> (<i>Venus</i>), Poli	XIII.	74
<i>ala-papilionis</i> (<i>Venus</i>), Chemnitz ..	II.	7	<i>laterisnlea</i> (<i>Tapes</i>), Sowerby	VI.	26
<i>alba</i> , <i>Deshayes</i>	IV.	14	<i>Largillierti</i> (<i>Venus</i>), Philippi	XI.	59
<i>anabilis</i> (<i>Venus</i>), Philippi	V.	21	<i>lentiginosa</i> , <i>Reeve</i>	VI.	25
<i>Amphidesmoides</i> , <i>Reeve</i>	X.	50	<i>lirata</i> (<i>Venus</i>), Philippi	V.	20
<i>Arctica</i> , <i>Reeve</i>	X.	52	<i>literata</i> (<i>Venus</i>), Linnaeus	I.	2
<i>aspersa</i> (<i>Venus</i>), Chemnitz	II.	5	<i>literata</i> (<i>Venus</i>), Pennant	XI.	57
<i>aurea</i> (<i>Venus</i>), Gmelin	VIII.	37	<i>literata</i> (<i>Venus</i>), Poli	XIII.	71
<i>Beudanti</i> (<i>Venus</i>), Payraudeau	X.	54	<i>livida</i> , <i>Reeve</i>	X.	47
<i>bicolorata</i> , <i>Reeve</i>	IX.	42	<i>Luzonica</i> , <i>Sowerby</i>	IX.	40
<i>biradiata</i> , <i>Deshayes</i>	IX.	46	<i>Malabarica</i> (<i>Venus</i>), Chemnitz	VI.	27
<i>Bruguieri</i> (<i>Venus</i>), Hanley	XIII.	73	<i>marmorata</i> (<i>Cytherea</i>), Lamarek	VI.	26
<i>Ceylonensis</i> , <i>Sowerby</i>	VII.	30	<i>Meroëformis</i> , <i>Sowerby</i>	V.	19
<i>cinercea</i> , <i>Deshayes</i>	XII.	62	<i>nebulosa</i> (<i>Venus</i>), Chemnitz	VII.	33
<i>corrugata</i> , <i>Deshayes</i>	XIII.	72	<i>nitens</i> (<i>Venus</i>), Turton	VIII.	37
<i>Cuningii</i> , <i>Sowerby</i>	XIII.	75	<i>nucleus</i> (<i>Venerupis</i>), Lamarek	XI.	58
<i>daetyloides</i> , <i>Sowerby</i>	XIII.	69	<i>obscura</i> (<i>Venus</i>), Gmelin	XI.	57
<i>declivis</i> , <i>Sowerby</i>	V.	23	<i>obscurata</i> , <i>Deshayes</i>	III.	12
<i>decussata</i> (<i>Venus</i>), Linnaeus	XI.	57	<i>occidentalis</i> , <i>Reeve</i>	VIII.	35
<i>deflorata</i> (<i>Venus</i>), Born	XI.	57	<i>opina</i> (<i>Venus</i>), Gmelin	VII.	33
<i>Deshayesii</i> (<i>Venus</i>), Hanley	II.	4	<i>orientalis</i> , <i>Reeve</i>	VIII.	34
<i>denticulata</i> , <i>Sowerby</i>	XII.	63	<i>ovalea</i> (<i>Venus</i>), Lamarek	III.	11
<i>dorsata</i> (<i>Venus</i>), Lamarek	III.	11	<i>palustris</i> (<i>Venus</i>), Mawe	XI.	58
<i>dura</i> (<i>Venus</i>), Gmelin	III.	13	<i>papilionacea</i> (<i>Venus</i>), Lamarek	II.	7
<i>edulis</i> (<i>Venus</i>), Chemnitz	VIII.	37	<i>perforans</i> (<i>Venerupis</i>), Lamarek	XI.	58
<i>faba</i> , <i>Reeve</i>	VIII.	39	<i>perforans</i> (<i>Venus</i>), Montagu	XI.	58
<i>fabagella</i> , <i>Deshayes</i>	XIII.	66	<i>petalina</i> (<i>Venus</i>), Lamarek	X.	54
<i>fasciata</i> (<i>Tellina</i>), Lister	VIII.	37	<i>Philippii</i> (<i>Chione</i>), Deshayes	VIII.	36
<i>fasciatus</i> (<i>Cuneus</i>), Da Costa	IV.	17	<i>Philippinarum</i> (<i>Venus</i>), Ad. & <i>Reeve</i>	XII.	60
<i>ferruginea</i> , <i>Reeve</i>	X.	51	<i>pinguis</i> (<i>Venus</i>), Chemnitz	VII.	33
<i>flammeolata</i> (<i>Venus</i>), Lamarek	VI.	28	<i>playia</i> (<i>Venus</i>), Jeffreys	XI.	58
<i>florida</i> (<i>Venus</i>), Lamarek	XIII.	74	<i>politata</i> , <i>Sowerby</i>	X.	49
<i>florida</i> (<i>Venus</i>), Poli	XI.	57	<i>pullastra</i> (<i>Venus</i>), Montagu	XI.	58
<i>floridella</i> (<i>Venus</i>), Deshayes	X.	53	<i>pullastra</i> (<i>Venerupis</i>), Fleming	XI.	58
<i>fumigata</i> (<i>Venus</i>), Sowerby	VII.	31	<i>pumicea</i> , <i>Deshayes</i>	XII.	61
<i>galactites</i> (<i>Venus</i>), Lamarek	XII.	65	<i>quadriradiata</i> , <i>Deshayes</i>	II.	6
<i>gallus</i> (<i>Venus</i>), Spengler	VI.	27	<i>radiata</i> (<i>Venus</i>), Anton	II.	4
<i>geographica</i> (<i>Venus</i>), Chemnitz	XIII.	71	<i>radiata</i> (<i>Venus</i>), Chemnitz	IV.	16
<i>grata</i> , <i>Deshayes</i>	III.	9	<i>rariplana</i> (<i>Venus</i>), Lamarek	III.	13
<i>hiantina</i> (<i>Venus</i>), Lamarek	VI.	28	<i>regularis</i> (<i>Chione</i>), Deshayes	IX.	44
<i>Indica</i> , <i>Sowerby</i>	XI.	56	<i>reticulatus</i> (<i>Cuneus</i>), Da Costa	XI.	57
<i>intermedia</i> (<i>Venus</i>), Quoy	XI.	59	<i>retifera</i> (<i>Venus</i>), Hanley	I.	1
<i>Japonica</i> , <i>Deshayes</i>	XIII.	67	<i>rhomboides</i> (<i>Venus</i>), Pennant	IV.	17
<i>Japonica</i> (<i>Venus</i>), Gmelin	IX.	43	<i>rimosa</i> (<i>Venus</i>), Philippi	III.	8
<i>Kochii</i> (<i>Venus</i>), Philippi	VIII.	38	<i>rimularis</i> (<i>Venus</i>), Lamarek	VI.	29
<i>Krausii</i> (<i>Chione</i>), Deshayes	IX.	41	<i>rotundata</i> (<i>Venus</i>), Linnaeus	II.	7
<i>Labuana</i> (<i>Venus</i>), Adams and <i>Reeve</i>	IX.	45	<i>sanguinolenta</i> (<i>Venus</i>), Gmelin	XI.	57

TAPES.—INDEX.

Plate.	Species.	Plate.	Species.
<i>Sarniensis</i> (<i>Venus</i>), Turton	IV. 17	turgida (<i>Venus</i>), Lamarek	III. 11
semidecussata, <i>Deshayes</i>	XIII. 67	turgidula, <i>Deshayes</i>	VII. 32
<i>Senegaleensis</i> (<i>Venus</i>), Dillwyn	XI. 58	undulata (<i>Venus</i>), Born	III. 8
similis, <i>Deshayes</i>	III. 10	undulata (<i>Venus</i>), Dillwyn	II. 3
<i>Sinensis</i> , <i>Reeve</i>	V. 24	variabilis (<i>Venus</i>), Philippi	VI. 26
sinuosa (<i>Venus</i>), Lamarek	V. 18	variegata, <i>Sowerby</i>	XII. 64
striata (<i>Venus</i>), Chemnitz	IX. 45	vernacosa, <i>Reeve</i>	X. 48
sulearia (<i>Venus</i>), Lamarek	I. 1	violascens, <i>Deshayes</i>	XIII. 68
suleosa (<i>Venus</i>), Philippi	V. 22	virago (<i>Venus</i>), Lovén	IV. 17
tenuistriata, <i>Sowerby</i>	X. 55	virginea (<i>Venus</i>), Linnaeus	IV. 17
textile (<i>Venus</i>), Gmelin	II. 3	vitulata, <i>Deshayes</i>	IV. 15
textrix (<i>Venus</i>), Chemnitz	II. 3	vulgaris (<i>Pullastra</i>), <i>Sowerby</i>	II. 3
texturata (<i>Venus</i>), Lamarek	XIII. 70	vulgaris (<i>Venerupis</i>), <i>Sowerby</i>	XI. 58
triradiata (<i>Venus</i>), Chemnitz	VII. 33	vulgaris (<i>Venus</i>), Broderip	XI. 58
tristis (<i>Venus</i>), Lamarek	IX. 44		

MONOGRAPH
OF THE GENUS
M E R O E.

"Various as beauteous, Nature, is thy face;
All that grows has grace; all are appropriate." — *Crabbe*



M E R O È.

PLATE I.

Genus MEROÈ, Schumacher.

Testa plerumque trigono-orata, interdum depresso cordato-orbicularis, saepissimè compressa, interdum tumida, nunc lœrigata, nunc concentricè lirata et sulcata, marginibus tenuè denticulatis, latere antico plerumque longiore. Area ligamenti peculiariter profundè excavata. Cardo bi-tridecalvis, dente subamplo antico. Impressio pallii posticè sinnata.

Shell generally triangularly ovate, sometimes depresso cordately orbicular, mostly compressed, sometimes tumid, smooth or concentrically ridged and grooved, margins finely denticulated, anterior side generally the longer. Ligamentary area peculiarly deeply excavated. Hinge two- or three-toothed, with a rather large anterior tooth. Impression of the mantle situated posteriorly.

This genus, less popularly known by the generic names *Cunens* and *Sunella*, is founded on the Linnaean *Venus Meroë*, a highly characteristic form, differing from all other genera of *Veneridæ* in the anterior side of the shell being mostly the longer, and in the ligamentary area being sunk into a deeply excavated pit. About a dozen species are known; two inhabiting Japan, three Australia, and seven the intermediate ocean from the Philippine Islands to China and Bombay. The Japanese and Australian species belong, curiously enough, to the same peculiar type,—depresso cordately orbicular and smooth, with painting of richly mottled violet; whilst the Indian Ocean species are of a quite distinct type,—triangularly ovate, the anterior side being attenuately produced, mostly concentrically ridged and linearly grooved, with the painting almost invariably represented by a zigzag textile of livid-purple. All are partially covered by a thin transparent horny epidermis, and all have finely denticulated margins, with the deeply excavated ligamentary pit.

Species 1. (Fig. *a* to *e*, Mus. Cuming.)

MEROÈ SEMINUDA. *Mer. testa orato-trigona, compressa, subæquilaterali, utrinque rotundata, latere antico delicii-rotundato, carneo-lactea, lirido-purpureo reticulato-maculata, in medio et posticè concentricè sulcata, antice lœrigata.*

THE SEMINUDE MEROÈ. Shell ovately triangular, compressed, nearly equilateral, rounded on both sides, anterior side slopingly attenuated, flesh-tinted cream-colour, reticulately blotched with livid-purple, in the middle, and posteriorly, concentrically grooved, anteriorly smooth.

Cytherea seminuda, Philippi, Abbild. und Besch. Conch. p. 97. pl. 3. f. 4.

Donax semiunda, Anton.

Venus semiunda, Anton.

Meroë semiunda, Sowerby.

Cunens semiundus, Deshayes.

Hab. Molucca Islands.

The chief peculiarity of this very delicately painted species consists in the concentric grooves suddenly ceasing on reaching the anterior side of the shell, which is consequently smooth.

Species 2. (Fig. *a*, *b*, Mus. Cuming.)

MEROÈ MAGNIFICA. *Mer. testa orbiculari, lentiformi, compressa, crassâ, antice abbreviatâ, lœrigatâ, violaceo-cinereâ, pallidè fusco aut violaceo acutè linearitereticulatâ, intensè purpureo-violaceo plus minus concentricè fasciatâ et interrupè radiatâ, epidermide tenui corneari versus marginem indutâ.*

THE FINE MEROÈ. Shell orbicular, lens-shaped, compressed, thick, rather shortened anteriorly, smooth, violet-tinged ash, sharply linearly reticulated with pale-brown or violet, more or less concentrically banded and interruptedly rayed with deep purple-violet, covered towards the margin with a thin horny epidermis.

Hab. Japan.

This fine species, pre-eminent in size and brilliancy of colouring, might be regarded as a colossal form of *M. excavata*, from the same locality, but no specimens intermediate in size and general form are at present known to connect them. It remains to be seen whether the compressly attenuated character of the anterior portion of the shell of the latter species is indicative of an early stage of growth.




3a




3b.




3c.




4




5a.



5c



5b.



6a.



6c.



6b.

M E R O E.

PLATE II.

Species 3. (Fig. *a*, *b*, *c*, Mus. Cuming.)

MEROE TRUNCATA. *Mer. testa trigono-ovata, compressa, antice subattenuato-producta, fulvescente-alba, livido-purpureo concentricè acutè linearis-undata, levigata. sub lente exiliter decussatim striata.*

THE TRUNCATED MEROE. Shell triangularly ovate, compressed, anteriorly somewhat attenuately produced, fulvous-white, concentrically sharply linearly waved with livid-purple, smooth, beneath the lens finely decussately striated.

Cuneus truncatus, Deshayes, Pro. Zool. Soc. 1853, p. 1.

Hab. Philippine Islands; Cuming.

Though apparently smooth, this species, like *M. scripta*, which it very closely resembles, will be found, on examination with the lens, to be minutely decussately striated.

Species 4. (Mus. Cuming.)

MEROE EFFOSA. *Mer. testa trigono-ovata, subangusta, subaequilaterali, albida, livido-purpureo profuse reticulato-maculata, undique creberimè concentricè costellata, interstitiis conspicue excavatis, area ligamenti profundè effossa.*

THE PITTED MEROE. Shell triangularly ovate, rather narrow, nearly equilateral, whitish, profusely reticulately blotched with livid-purple, very closely concentrically ribbed throughout, the interstices being conspicuously excavated, ligamentary area deeply pitted.

Cytherea effossa, Hanley, Pro. Zool. Soc. 1842, p. 123.

Meroe effossa, Sowerby.

Cuneus effossus, Deshayes.

Hab. Philippine Islands; Cuming.

Conspicuously distinguished from *M. picta*, which it resembles in painting, by its narrower and more equilateral form, deeply-cut ribs, and correspondingly deeply-cut ligamentary area.

Species 5. (Fig. *a*, *b*, *c*, Mus. Cuming.)

MEROE PICTA. *Mer. testa trigono-ovala, compressa, late antico elongato-producta, postico declivi-truncata,*

albida, livido-purpureo reticulato-maculata, interdum omnino purpurea, concentricè plano-lirata, interstitiis linearis-sulcata.

THE PAINTED MEROE. Shell triangularly ovate, compressed, anterior side elongately produced, posterior slopingly truncated, whitish, reticulately blotched with livid-purple, sometimes altogether purple, concentrically flatly ridged, interstices linearly grooved.

Venus Meroe, Linnaeus, Syst. Nat. p. 1132.

Cuneus costatus, Megerle.

Donax Meroe, Lamarek.

Cytherea Meroe, Sowerby.

Meroe picta, Schumacher.

Cuneus Meroe, Deshayes.

Hab. Indian Ocean.

In this species, which is the original type of the group, the anterior side is much produced, whilst the posterior is more than usually slopingly truncated. The painting is exceedingly variable.

Species 6. (Fig. *a*, *b*, *c*, Mus. Cuming.)

MEROE SCRIPTA. *Mer. testa trigono-ovata, antice attenuato-producta, postice declivi-truncata, carneo-alba, livido-purpureo linearis-undata vel reticulato-maculata, interdum omnino purpurea, levigata, sub lente minute decussatim striata.*

THE WRITTEN MEROE. Shell triangularly ovate, anteriorly attenuately produced, posteriorly slopingly truncated, flesh-white, linearly waved or reticulately blotched with livid-purple, sometimes altogether purple, smooth, beneath the lens minutely decussately striated.

Cuneus scriptus, Rumphius, Mus. Amb. pl. 43, f. 1. m.

Donax scripta, Linnaeus.


Meroe scripta, Gray.

Hab. Indian Ocean.

An apparently smooth species, which on examination with the lens will be found to be minutely decussately striated.



8.



9.




10a



11a



11b.



10b



12a.



12b.



12c

MEROË.

PLATE III.

Species 7. (Mus. Cuming.)

MEROË VAGINALIS. *Mer. testá ovalá, parum trigoná, anticé subabbreviato-attenuatá, lœvigatá, flavescente-albá, purpureo-violaceo radiatim marmoratá et nebulatá.*

THE SHEATHED MEROË. Shell ovate, but little triangular, anteriorly rather shortly attenuated, smooth, yellowish-white, radiately mottled and clouded with purple-violet.

Cytherea vaginalis, Menke, Moll. Nov. Holl. p. 42.

Cuneus vaginalis, Deshayes.

Hab. West coast of New Holland.

Unlike the typical species of the Indian Ocean, the anterior side of the Australian and Japanese species of *Meroë* is the shorter.

Species 8. (Mus. Cuming.)

MEROË ALICIE. *Mer. testá cordato-orbiculari, depresso-sinuclá, lœvigatá, flavescente-albá, purpureo-roseo reticulato-lentiginosá et obscurè radiatá.*

ALICE'S MEROË. Shell cordately orbicular, rather depressed, smooth, yellowish-white, reticulately freckled and obscurely rayed with purple-rose.

Sanetta Aliciae, Adams and Angas, Pro. Zool. Soc. 1863.
Hab. Port Elliott, South Australia; Angas.

This species, dredged at Port Elliott, South Australia, by Mr. Angas, comes very near to the preceding; it is, however, rounder and more gibbous.

Species 9. (Mus. Cuming.)

MEROË MENSTRUALIS. *Mer. testá orbiculari-ovatá, medio-gibbosinserlá, latere antico brevi, albida, livido-purpureo linearis-undulatá, lœvigatá, sub lente obscurè radiatim striatá.*

THE MENSTRUAL MEROË. Shell orbicularly ovate, rather gibbous in the middle, anterior side short, whitish, linearly waved with livid-purple, smooth, obscurely radiately striated beneath the lens.

Cytherea menstrualis, Menke, Moll. Nov. Holl. p. 43.

Cuneus menstrualis, Deshayes.

Hab. New Holland?

This appears to be an important species, intermediate in character between the smooth Australian form, *M. vaginalis*, and *M. truncatus* of the Philippine Islands.

Species 10. (Fig. a, b, Mus. Cuming.)

MEROË SOLANDRI. *Mer. testá subtrigono-ovatá, tumidá, subaequilaterali, laevid, livido-purpureo undulatim maculatá aut lineatá, epidermide tenui cornéa induitá, lœrigatá.*

SOLANDER'S MEROË. Shell somewhat triangularly ovate, swollen, nearly equilateral, cream-colour, undulately blotched or lined with livid-purple, covered with a thin horny epidermis, smooth.

Cytherea Solandri, Gray, Wood, Ind. Test. Suppl. pl. 2. f. 11.

Meroë Solandri, Gray.

Cuneus Solandri, Deshayes.

Hab. China.

Of a swollen gibbous form, nearly equilateral, but somewhat variable.

Species 11. (Fig. a, b, Mus. Cuming.)

MEROË EXCAVATA. *Meroë testá orbiculari-ovatá, anticé compressá, cordato-attenuatá, posticé latè rotundatá, albida, fulvo-violaceo plus minus lineatá, purpureo interrupitè fascialá, lœvigatá, epidermide tenui cornéa partim induitá.*

THE EXCAVATED MEROË. Shell orbicularly ovate, anteriorly compressed and cordately attenuated, posteriorly broadly rounded, whitish, more or less stained with fulvous-violet, interruptedly banded with purple, smooth, partially covered with a thin horny epidermis.

Cytherea excavata, Hanley, Pro. Zool. Soc. 1842, p. 123.

Meroë excavata, Sowerby.

Cuneus excavatus, Deshayes.

Hab. Japan.

The painting of this species is similar to that of the large species from the same locality, which I have named *M. magnifica*. It differs in form in having the posterior side broadly rounded and the anterior compressly attenuated.

MEROË.—PLATE III.

Species 12. (Fig. *a*, *b*, *c*, Mus. Cuming.)

MEROE HIANS. *Mer. testá subangustè ovatâ, conspicuè tumidâ, subæquilaterali, latere antico longiore, albida, livido-purpureo plus minus tinctâ ant angulato-flammatâ, lævigatâ, epidermide temni corneâ partim indutâ.*

THE GAPING MEROË. Shell rather narrowly ovate, conspicuously tumid, nearly equilateral, anterior side the longer, whitish, more or less stained or angularly

flamed with livid-purple, smooth, partially covered with a thin horny epidermis.

Hab. Bombay.

The three specimens here figured occur, together with a few others, in Mr. Cuming's cabinet, with the name *M. hians* in manuscript. They run very closely into the China *M. Solandri*, but are all of the same peculiarly tumid narrowly ovate form.

