



MANUAL

BACTERIOLOGY

EDGAR N. LITTLE

MANUAL
OF
BACTERIOLOGY

QR41

C8

1890

BIOLOGY
LIBRARY
G

BACTERIOLOGY



RIDGAR M. CROOKSHANK, M.B.

LONDON :

H. K. LEWIS, 136 GOWER STREET, W.C.

REVISIONS AND CORRECTIONS
BY THE AUTHOR

LONDON :

H. K. LEWIS, 136 GOWER STREET, W.C.

To

SIR JOSEPH LISTER, BART., M.B., F.R.S.,

WHO HAS CREATED A NEW EPOCH IN

MEDICINE AND SURGERY,

BY APPLYING A KNOWLEDGE OF MICRO-ORGANISMS

TO THE TREATMENT OF DISEASE,

This Work is Dedicated

AS A

TOKEN OF ADMIRATION AND RESPECT

BY THE AUTHOR.

The author trusts that this manual will continue to be of use to students and practitioners in both the medical and veterinary professions.

For references to recent literature, the author again takes the opportunity of calling attention to Professor Baumgarten's invaluable annual.

EDGAR M. CROOKSHANK.

24, MANCHESTER SQUARE, LONDON, W.

October 28th, 1890.

PREFACE

TO THE SECOND EDITION.

THE fact that a new edition of this manual was called for a few months after its publication, has induced the author to extend its scope in the hope of adding to its usefulness.

The work has not only been revised throughout and brought up to date, but, in order to admit of a more concise arrangement of the species, the Systematic part has been recast.

Additional chapters have been written upon the General Morphology and Physiology of Bacteria, upon Antiseptics and Disinfectants, and Immunity.

Seventy-three illustrations have been added. Those not duly acknowledged as coming from other sources were drawn on the wood by the author from his own preparations.

A list of references to works on Bacteriology, which was not ready for the first edition, has now been completed and extended. It has no pretension to be a complete bibliography, but being arranged

as much as possible in accordance with the chapters, and in chronological order, may be useful to those seeking further details. No doubt Professor Baumgarten's *Jahresbericht*, the first number of which has been issued this year, will be found a valuable guide to current literature in the future.

The author desires again to express his acknowledgments to Professor Gerald Yeo and Mr. Herroun, of King's College, London.

EDGAR M. CROOKSHANK.

24, MANCHESTER SQUARE, W.

December, 1886.

CONTENTS.

INTRODUCTION.

	PAGE
<i>THE GERM THEORY</i>	I

PART I.

GENERAL METHODS.

CHAPTER I.

<i>APPARATUS, MATERIAL, AND REAGENTS EMPLOYED IN A BACTERIOLOGICAL LABORATORY.</i>	23
<i>a.</i> Histological apparatus	23
<i>b.</i> Reagents and materials employed in the processes of hardening, decalcifying, embedding, fixing, and cutting of tissues	25
<i>c.</i> Reagents for examining and staining microscopical preparations	27
<i>d.</i> Reagents for mounting and preserving preparations	34
<i>e.</i> Drawing and photographic apparatus	34
<i>f.</i> Sterilisation apparatus	36
<i>g.</i> Apparatus and material for preparing and storing nutrient gelatine and nutrient agar-agar	38
<i>h.</i> Apparatus for employment of nutrient jelly in test-tube and plate- cultivations	41
<i>i.</i> Apparatus for preparation of potato-cultivations	45
<i>j.</i> Apparatus for preparation of solidified sterile blood serum	46
<i>k.</i> Apparatus for storing, and for cultivations in, liquid media	48
<i>l.</i> Apparatus for incubation	49
<i>m.</i> Inoculating and dissecting instruments and apparatus in common use	58
<i>n.</i> General laboratory requisites	59

CHAPTER II.

	PAGE
<i>MICROSCOPICAL EXAMINATION OF BACTERIA IN LIQUIDS, IN CULTIVATIONS, ON SOLID MEDIA, AND IN TISSUES</i>	62
<i>a.</i> Examination in the fresh state	63
<i>b.</i> Cover-glass preparations: methods of Ehrlich, Babes, and His	65
<i>c.</i> Cover-glass impressions	69

CHAPTER III.

<i>PREPARATION AND STAINING OF TISSUE SECTIONS</i>	71
<i>a.</i> Methods of hardening and decalcifying preparations	71
<i>b.</i> Methods of embedding, fixing, and cutting	72
<i>c.</i> General principles of staining bacteria in tissue sections: methods of Weigert, Gram, and Weigert-Ehrlich	75

CHAPTER IV.

<i>PREPARATION OF NUTRIENT MEDIA AND METHODS OF CULTIVATION</i>	80
<i>SOLID MEDIA:</i>	
<i>a.</i> Preparation of nutrient gelatine and nutrient agar-agar	82
<i>b.</i> Methods of employing nutrient jelly in test-tube and plate-cultivations	87
<i>c.</i> Preparation and employment of sterilised potatoes, potato-paste, bread-paste, vegetables, fruit, and white of egg	99
<i>d.</i> Preparation and employment of sterile blood serum	104
<i>LIQUID MEDIA:</i>	
<i>e.</i> Preparation of sterilised bouillon, liquid blood serum, urine, milk, vegetable infusions, and artificial nourishing liquids	106
<i>f.</i> Methods of storing and employing liquid media; Lister's flasks, Aitken's test-tubes, Sternberg's bulbs, Pasteur's apparatus, Miquel's bulbs; Drop-cultures; Warm stages	109

CHAPTER V.

	PAGE
<i>EXAMINATION OF AIR, SOIL, AND WATER</i>	125
Air	125
Soil	130
Water	131

CHAPTER VI.

<i>EXPERIMENTS UPON THE LIVING ANIMAL</i>	137
<i>a.</i> Inhalation of micro-organisms	137
<i>b.</i> Administration with food	137
<i>c.</i> Cutaneous and subcutaneous inoculation	138
<i>d.</i> Special operations	139

CHAPTER VII.

<i>EXAMINATION OF ANIMALS EXPERIMENTED UPON, AND THE METHODS OF ISOLATING MICRO-ORGANISMS FROM THE LIVING AND DEAD SUBJECT</i>	141
<i>a.</i> Method of dissection and examination	141
<i>b.</i> Isolation of micro-organisms from the living subject	144

PART II.

GENERAL BIOLOGY OF BACTERIA.

CHAPTER VIII.

<i>GENERAL MORPHOLOGY AND PHYSIOLOGY</i>	147
--	-----

CHAPTER IX.

<i>ANTISEPTICS AND DISINFECTANTS</i>	180
--	-----

CHAPTER X.

<i>IMMUNITY</i>	192
---------------------------	-----

PART III.

*SYSTEMATIC AND DESCRIPTIVE, WITH SPECIAL
MICROSCOPICAL METHODS.*

CHAPTER XI.

	PAGE
<i>CLASSIFICATION OF BACTERIA</i>	205

CHAPTER XII.

<i>SYSTEMATIC AND DESCRIPTIVE</i>	224
GROUP I. COCCACEÆ	224
Genus I. Streptococcus	225
„ II. Merismopedia	241
„ III. Sarcina	243
„ IV. Micrococcus	245
„ V. Ascococcus	257
GROUP II. BACTERIACEÆ	260
Genus I. Bacterium	261
„ II. Spirillum	282
„ III. Leuconostoc	296
„ IV. Bacillus	299
„ V. Vibrio	349
„ VI. Clostridium	350
GROUP III. LEPTOTRICHEÆ	353
Genus I. Crenothrix	354
„ II. Beggiatoa	356
„ III. Phragmidiothrix	360
„ IV. Leptothrix	361
GROUP IV. CLADOTRICHEÆ	362
Genus I. Cladothrix	362

APPENDICES.

A. Yeasts and moulds	369
B. Actinomyces	379
C. Flagellated protozoa in the blood	397
D. Hæmatozoa of malaria (Laveran)	408
E. Chronological bibliography	414
F. Table showing the magnifying power of Zeiss' objectives	452



LIST OF ILLUSTRATIONS.

FIG.	WOOD ENGRAVINGS.	PAGE
1.	Koch's Steam-Steriliser	37
2.	Hot-air Steriliser	38
3.	Section of Hot-air Steriliser	38
4.	Hot-water Filtering Apparatus, with Ring Burner	39
5.	Wire Cage for Test-tubes	42
6.	Platinum Needles	42
7.	Damp Chamber for Plate-cultivations	43
8.	Apparatus employed for Plate-cultivations	43
9.	Box for Glass Plates	44
10 and 11.	Glass Benches for Glass Plates or Slides	44
12.	Israel's Case	45
13.	Damp Chamber for Potato-cultivations	45
14.	Serum Steriliser	46
15.	Serum Inspissator	47
16.	D'Arsonval's Incubator	50
17.	Schlösing's Membrane Regulator	51
18.	Gas-burner protected with Mica-cylinder	52
19.	Moitessier's Gas-pressure Regulator	53
20.	Koch's Safety Burner	53
21.	Babes' Incubator	54
22.	Reichert's Thermo-regulator	56
23.	Meyer's Thermo-regulator	57
24.	Siphon Bottle, with Flexible Tube, Glass Nozzle, and a Mohr's Pinchcock	59
25.	Dessicator	60
26.	Method of making a Folded Filter	85
27.	Method of Inoculating a Test-tube, containing Sterile Nutrient Jelly	88
28.	Method of Inoculating Test-tubes in the preparation of Plate-cultivations	93
29.	Method of dividing Potatoes	101
30.	Method of Forming a Simple Moist Chamber	114
31.	Simple Warm Stage	115
32.	Simple Warm Stage shown in Operation	116
33.	Schäfer's Warm Stage	117

FIG.	PAGE
34. Stricker's Warm Stage	118
35. Section of Israel's Warming Apparatus and Drop-culture Slide	118
36. Israel's Warming Apparatus	119
37. Israel's Warming Apparatus in Operation	120
38. Simple Gas Chamber	121
39. Gas Chamber in use with Apparatus for generating Carbonic Acid	121
40. Stricker's Combined Gas Chamber and Warm Stage	122
41. Simple Moist Chamber adapted for transmission of Electricity	122
42. Apparatus arranged for transmitting Electricity	123
43. Slide with Gold-leaf Electrodes	124
44. Hesse's Apparatus	128
45. Apparatus employed for counting the Colonies on a Plate-cultivation	134
46. From a Preparation of <i>Bacillus anthracis</i>	149
47. <i>Ascococcus Billrothii</i> , $\times 65$ (after Cohn)	152
48. <i>Streptococcus</i> and <i>Sarcinacoccus</i> from a Drop-cultivation, $\times 1200$	153
49. <i>Streptococcus</i> in the Blood of a Rabbit, $\times 1200$	153
50. <i>Streptococcus</i> of Progressive Tissue Necrosis in Mice (after Koch)	153
51. <i>Spirochæta</i> from Sewage Water, $\times 1200$	155
52. Bacteria showing Flagella	157
53. <i>Bacillus megaterium</i>	159
54. <i>Clostridium butyricum</i> , $\times 1020$	160
55. A Thread of <i>Bacillus anthracis</i> with Spores in a Drop-cultivation, $\times 1400$	161
56. <i>Leuconostoc Mesenteroides</i> ; Cocci-chains with Arthrospores (after Van Tieghem and Cienkowski)	161
57. Spores of <i>Bacillus anthracis</i> , stained with Gentian-violet, after passing the Cover-glass twelve times through the Flame, $\times 1200$	163
58. Spore-bearing Threads of <i>Bacillus anthracis</i> , Double-stained with Fuch sine and Methylene-blue, $\times 1200$	163
59. Tubercle Bacilli in Sputum, $\times 2500$ (from photographs)	165
60. Leprosy Bacilli from a Section of Skin, $\times 1200$	165
61. Glanders Bacilli from a Section of a Glanders Nodule . $\times 1200$	165
62. Bacterium of Chicken-cholera from Blood of Infected Hen, $\times 1200$	166
63. Bacterium of Chicken-cholera from Muscle Juice of an Infected Hen, $\times 2500$ (from a photograph)	166
64. Comma Bacilli in Sewage Water stained with Gentian-violet, $\times 1200$	166
65. <i>Vibrios</i> in Water contaminated with Sewage, $\times 1200$	167
66. <i>Spirillum undula</i> , $\times 1200$	167
67. <i>Cladothrix dichotoma</i>	214
68. <i>Bacterium pneumoniæ croupose</i> , $\times 1500$ (after Zopf)	219
69. Emmerich's Bacterium, $\times 700$ (after Emmerich)	220
70. Colonies on Nutrient Gelatine, $\times 60$	220
71. Colonies on Nutrient Agar-agar, $\times 60$	221
72. Colony of <i>Bacillus anthracis</i> , $\times 60$	221
73. Bacterium of Rabbit Septicæmia	222
74. <i>Streptococcus</i> of Progressive Tissue Necrosis in Mice	237

FIG.	PAGE
75. Micrococcus of Pyæmia in Rabbits; Vessel from the Cortex of the Kidney, × 700	252
76. <i>Ascococcus Billrothii</i> (after Cohn)	257
77. Bacterium Pneumoniæ Crouposæ, from Pleural Cavity of a Mouse, × 1500 (after Zopf)	261
78. Bacterium Neapolitanum, × 700 (after Emmerich)	264
79. Bacteria of Rhinoscleroma, × 1400 (after Cornil)	264
80. Bacterium of Chicken Cholera; Blood of Inoculated Hen, × 1200	268
81. Bacterium of Chicken Cholera, from Muscle Juice of Inoculated Hen, × 2500	268
82. Bacterium of Rabbit Septicæmia; Blood of Sparrow, × 700 (after Koch)	270
83. Bacterium Indicum; Colonies on Nutrient Agar-agar, × 60	275
84. Bacterium Zopfii; Successive Changes in the same Thread, × 740	279
85. Cover-glass Preparation of the Edge of a Drop of Meat Infusion, × 600 (after Koch)	284
86. Colonies of Comma Bacilli on Nutrient Gelatine, natural size (after Koch)	285
87. Colonies of Koch's Comma Bacilli, × 60	285
88. Cover-glass Preparation from the Contents of a Cholera Intestine, × 600 (after Koch)	286
89. Cover-glass Preparation of Cholera Dejecta on Damp Linen (two days old), × 600 (after Koch)	286
90. Section of the Mucous Membrane of a Cholera Intestine, × 600 (after Koch)	287
91. Pure Cultivation of Finkler's Bacillus, twenty-four hours old	288
92. " " " " two days old	288
93. " " Koch's Cholera Bacillus, twenty-four hours old	288
94. " " " " " two days old	288
95. Comma-shaped Organisms with other Bacteria in Sewage-contaminated Water, × 1200	289
96. Pure Cultivation of Spirillum Finkleri in twenty-four hours	292
97. " " " " thirty-six hours	292
98. <i>Spirillum sputigenum</i> , × 1200	293
99. <i>Spirillum tyrogenum</i> , × 1200	293
100. <i>Spirillum plicatile</i> (Marsh Spirochæte), × 1200	294
101. <i>Spirillum undula</i> , × 1500	295
102. <i>Leuconostoc mesenteroides</i>	297
103. Leprosy Bacilli from a Section of Skin, × 1200	300
104. <i>Bacillus typhosus</i> from a Potato-cultivation, × 1500	303
105. <i>Bacillus tuberculosis</i> from Tubercular Sputum, stained by Ehrlich's method, × 2500	306
106. Cultivation of the Tubercle Bacillus on Glycerine Agar-agar	308
107. <i>Bacillus anthracis</i> , × 1200	315
108. Pure Cultivation of the <i>Bacillus anthracis</i> in Nutrient Gelatine	318

FIG.	PAGE
109. Colonies in a Plate-cultivation, $\times 70$.	319
110. Cover-glass Impression-preparation, $\times 70$.	320
111. Spores of <i>Bacillus anthracis</i> unstained, $\times 1500$.	324
112. Spores of <i>Bacillus anthracis</i> , $\times 1200$.	325
113. From a Double-stained Preparation of <i>Bacillus anthracis</i> , $\times 1200$.	325
114. <i>Bacillus mallei</i> , $\times 1200$.	326
115 and 116. Pure Cultivations of the Bacillus of Septicæmia of Mice.	331
117. From a preparation of Bronchial Mucus of a Pig (after Klein).	333
118. Blood of Fresh Spleen of a Mouse, after inoculation with Swine Fever (after Klein).	333
119. Bacilli from an Artificial Culture with Spores (after Klein).	333
120. <i>Bacillus cyanogenus</i> , $\times 650$ (after Neelsen).	338
121. <i>Bacillus megaterium</i> (after De Bary).	343
122. Pure Cultivation of <i>Bacillus figurans</i> in Nutrient Agar-agar.	344
123. <i>Bacillus saprogenes</i> , No. 1 (after Rosenbach).	347
124. <i>Vibrio rugula</i> , $\times 1020$ (after Prazmowski).	349
125. <i>Clostridium butyricum</i> (after Prazmowski).	352
126. <i>Crenothrix Kühniana</i> (after Zopf).	355
127. <i>Beggiatoa alba</i> (after Zopf).	357
128. Several Phase-forms of <i>Beggiatoa Roseopersicina</i> (after Warming).	358
129. <i>Cladothrix dichotoma</i> (after Zopf).	363
130. Parasites in the Blood of Rats (after Lewis).	397
131. <i>Hæmatomonas cobitis</i> (after Mitrophanow).	399
132. Organisms in the Blood of the Carp (after Mitrophanow).	400
133. "Surra" Parasites occurring singly and fused, $\times 1200$.	402
134. A Monad in Rat's Blood, $\times 3000$.	404
135. Monads in Rat's Blood, $\times 1200$.	405
136. " " " " (stained).	406
137. Hæmatozoa of Malaria; Non-pigmented Forms (after Marchiafava and Celli).	409
138. Pigmented Amœboid Forms (after Golgi).	409
139. Semilunar Bodies of Laveran (after Golgi).	410
140. Rosette Forms with Segmentation (after Golgi).	411
141. Flagellated Forms (after Vandyke Carter).	411

DESCRIPTION OF PLATES.

PLATE I. (Facing page 18.)

Bacteria, Schizomycetes, or Fission fungi.

- FIG.
1. Cocci singly and varying in size.
 2. Cocci in chains or rosaries (streptococcus).
 3. Cocci in a mass or swarm (zoogloea).
 4. Cocci in pairs (diplococcus).
 5. Cocci encapsuled (*Bacterium pneumoniae crouposa*).
 6. Cocci in groups of four (merismopedia).
 7. Cocci in packets (sarcina).
 8. *Bacterium termo*.
 9. *Bacterium termo*, $\times 4000$ (after Dallinger and Drysdale).
 10. *Bacterium lineola*.
 11. *Bacillus subtilis*.
 12. *Bacillus tuberculosis*.
 13. *Bacillus leprae*.
 14. *Bacillus malariae* (after Klebs).
 15. *Bacillus typhosus* (after Eberth).
 16. *Spirillum undula* (after Cohn).
 17. *Spirillum volutans* (after Cohn).
 18. *Spirillum cholerae Asiatica*: from an artificial cultivation.
 19. *Spirillum Obermeieri* (after Koch).
 20. *Spirochete plicatilis* (after Flügge).
 21. *Vibrio rugula* (after Prazmowski).
 22. *Cladothrix Færsteri* (after Cohn).
 23. *Cladothrix dichotoma* (after Cohn).
 24. *Monas Okenii* (after Cohn).
 25. *Monas Warmingii* (after Cohn).
 26. *Rhabdomonas rosea* (after Cohn).
 27. From a cover-glass preparation of blood from the spleen of a mouse which had died of anthrax; stained with fuchsine (Zeiss' $\frac{1}{8}$ o.i. Oc. 4).
 28. From a drop-cultivation of *Bacillus anthracis* (Zeiss' $\frac{1}{8}$ o.i. Oc. 4).
 29. From a cover-glass impression-preparation of a potato-cultivation of *Bacillus anthracis* (Zeiss' $\frac{1}{8}$ o.i. Oc. 4).
 30. From a preparation of *Bacillus anthracis*, cultivated in nutrient gelatine (torula-form).
 31. Involution-form of *Crenothrix* (after Zopf).

FIG.

32. Involution-forms of *Vibrio serpens* (after Warming).
33. Involution-forms of *Vibrio rugula* (after Warming).
34. Involution-forms of *Clostridium Polymyxa* (after Prazmowski).
35. Involution-forms of the *Spirillum cholerae Asiaticæ*, from an artificial cultivation.
36. Involution-forms of *Bacterium acetii* (after Zopf and Hansen).
37. Spirulina-form of *Beggiatoa alba* (after Zopf).
38. Various thread-forms of *Bacterium merismopedioides* (after Zopf).
39. False-branching of *Cladothrix* (after Zopf).

PLATE II. (*Facing page 76.*)

1. *Micrococcus tetragonus*.—From a section of a kidney of a mouse which had died in eight days, after inoculation subcutaneously with a pure cultivation. Encapsuled tetrads, isolated and in masses, were found in the kidneys, lungs, and other organs. Stained with Gram's method (gentian-violet) without a contrast stain. × 1500.
2. *Micrococcus pyogenes aureus* (*Staphylococcus pyogenes aureus*).—From a section of the liver of a rabbit. A small vessel is shown plugged with cocci. From small abscesses in the liver, cultivations were obtained of the characteristic yellow coccus of pus. Stained with Gram's method (gentian-violet) without a contrast stain. × 1500.

PLATE III. (*Facing page 78.*)

1. *Bacillus tuberculosis*.—From a cover-glass preparation of tubercular pus. Stained with Ehrlich's method (fuchsine and methylene-blue). × 1500.
2. *Bacillus lepræ*.—From a section of a kidney from a case of leprosy. Stained with Ehrlich's method (fuchsine and methylene-blue). In the centre of the field is a glomerulus with a collection of the leprosy bacilli. × 400.

PLATE IV. (*Facing page 88.*)

1. *Spirillum cholerae Asiaticæ* (*Comma-bacillus of Koch*).—Tube inoculated from a plate-cultivation. The growth in this case was very striking. The funnel-shaped area of liquefaction, enclosing an air-bubble, and the white thread along the needle-track, are in marked contrast to the appearances, under similar conditions, of the comma-bacillus in *Cholera nostras* (p. 288).
2. *Bacterium cholerae gallinarum* (*Micrococcus cholerae gallinarum*; *Microbe du choléra des poules*).—Tube inoculated from the blood of a hen which had died of so-called chicken-cholera. After several days the growth forms a very delicate, finely beaded thread.
3. *Micrococcus cereus albus* (*Staphylococcus cereus albus*).—Tube inoculated from the pus of a subcutaneous abscess in a rabbit. The growth assumed a nodular appearance along the needle-track.

FIG.

PLATE V. (*Following Plate IV.*)

1. *Micrococcus tetragonus*.—Tube inoculated from a plate-cultivation of bacteria in sputum. The cultivation consisted of a milk-white growth heaped up on the surface of the gelatine, and growing freely along the upper part of the needle-track.
2. *Bacterium pneumoniae crouposæ* (*Micrococcus pneumoniae crouposæ*; *Friedländer's pneumo-coccus*).—Tube inoculated from pneumonic exudation. The growth, in nutrient gelatine, in the form of a round-headed nail, is not by itself distinctive.
3. *Saccharomyces niger* (*Black torula*).—Tube inoculated from an old contaminated nutrient-gelatine cultivation. The growth, isolated and reinoculated, formed a black crust on the surface of the gelatine. In some of the tubes little separate centres of growth occurred in the upper part of the track of the needle.

PLATE VI. (*Following Plate V.*)

1. *Bacillus pyocyaneus* (*Bacterium æruginosum*; *Bacillus fluorescens*).—Tube inoculated from pus. The gelatine was liquefied, and appeared green by transmitted and orange by reflected light.
2. *Sarcina lutea*.—Tube inoculated from a colony which occurred on potato exposed to the air. The gelatine was partially liquefied, and a canary-yellow growth had subsided to the bottom of the liquefied layer.
3. *Bacillus anthracis*.—Tube inoculated from the blood of a mouse which had died of anthrax. The typical growth which occurs in a few days is shown on p. 318. In this figure the appearance after three weeks is represented. The gelatine was completely liquefied, and a flocculent mass had subsided to the bottom of the tube.

PLATE VII. (*Following Plate VI.*)

1. *Sarcina lutea*.—In this tube and the two adjacent ones, the inoculations were made by thrusting the needle into the nutrient agar-agar. In all three cases the growth on the surface, freely exposed to air, developed a characteristic pigment, while the growth in the track of the needle was scanty and colourless.
2. *Bacterium indicum* (*Micrococcus indicus*, Koch; *Bacillus inaiicus*, Flügge).
3. *Saccharomyces rosaceus* (*Pink torula*).

PLATE VIII. (*Following Plate VII.*)

1. *Bacterium indicum* (*Micrococcus indicus*, Koch; *Bacillus indicus*, Flügge).—Tube inoculated from a nutrient agar-agar plate-cultivation. By plate-cultivation, or by successive cultivation on potatoes, a *pure* cultivation can be obtained. The growth has then the colour of red

FIG.

sealing-wax, and a peculiar crinkled appearance. After some days the growth loses its bright colour, and becomes purplish, like an old cultivation of *Bacterium prodigiosum*.

2. **Bacillus cyanogenus** (*Bacterium syncyanum* ; *Bacillus of blue milk*).—Tube inoculated from a potato-cultivation. The bacillus forms a whitish layer, and colours the nutrient agar-agar a smoky brown.
3. **Bacterium prodigiosum** (*Monas prodigiosa* ; *Micrococcus prodigiosus*, Cohn ; *Bacillus prodigiosus*, Flügge ; *Blood rain*).—Tube inoculated from a potato-cultivation. The bacterium grows very rapidly, forming a blood-red growth, which gradually acquires a purplish colour.

PLATE IX. (Following Plate VIII.)

1. **Sarcina lutea**.—Tube of nutrient agar-agar inoculated from a plate-cultivation. The canary-yellow colour forms a strong contrast to the colour of the growth in the adjacent tube.
2. **Micrococcus pyogenes aureus** (*Staphylococcus pyogenes aureus*).—Tube inoculated from an abscess in a rabbit.
3. **Bacillus pyocyaneus** (*Bacillus fluorescens* ; *Bacterium æruginosum*).—Tube inoculated from a colony on a plate-cultivation. The growth formed a whitish, transparent layer composed of slender bacilli. The pigment diffused itself throughout the nutrient jelly. The growth appears green by transmitted light, owing to the colour of the medium behind it. The bacillus is now regarded as identical with the bacillus of green-blue pus.

PLATE X. (Following Plate IX.)

1. **Bacterium lineola**.—Tube inoculated from putrid blood-serum which swarmed with *Bacterium lineola*. The same bacteria were found in the cultivation.
2. **Micrococcus rosaceus**.—Tube inoculated from an old cultivation on nutrient agar-agar which had become contaminated. The cultivation, macroscopically resembling pink yeast, consisted of a pure cultivation of micrococci.
3. **Micrococcus pyogenes citreus**.—Tube inoculated with cocci isolated from a subcutaneous abscess in a mouse.

PLATE XI. (Following Plate X.)

1. **Bacillus anthracis** (*Bactériidie du charbon* ; *Bacillus of splenic fever, wool-sorter's disease, or malignant pustule*).—Tube of nutrient agar-agar inoculated with the blood of a sheep which had died of anthrax. White flocculent patches developed, which were entirely composed of threads and spores of the bacilli.

FIG.

2. *Bacillus subtilis*.—Tube inoculated with bacilli, isolated by plate-cultivation, from dust. The bacilli appeared to be identical with the hay-bacillus, but in this case formed a peculiar crinkled layer along the track of the needle.
3. *Micrococcus cereus albus* (*Staphylococcus cereus albus*).—Tube inoculated from the discharge of a subcutaneous abscess in a rabbit.

PLATE XII. (Facing page 96.)

Spirillum of Finkler and Prior (*Comma-bacillus* of Finkler and Prior).—This figure represents the appearance of a plate-cultivation of the comma-bacillus from Cholera nostras, when examined over a slab of blackened plate-glass. The colonies differ very markedly from the colonies of Koch's comma-bacilli (see p. 285). The drawing was made from a typical result of thinning out or attenuating* the colonies by the process of plate-cultivation. At this stage they were completely isolated one from the other; but later they became confluent, and produced complete liquefaction of the gelatine.

PLATE XIII. (Following Plate XII.)

Spirillum of Finkler and Prior (*Comma-bacillus* of Finkler and Prior).—This figure represents the result obtained by a still further thinning out of the organisms than in the preceding case. The attenuation had been so far carried out that several of the colonies remained completely isolated for days.

PLATE XIV. (Facing page 102.)

1. *Bacterium prodigiosum* (*Monas prodigiosa*; *Micrococcus prodigiosus*, Cohn; *Bacillus prodigiosus*, Flügge; *Blood rain*).—Growth on potato after three days.
2. *Penicillium glaucum*.—A potato which had been freely exposed to the air was covered in three weeks by a growth of *Penicillium glaucum*. The surface of the growth is studded with dew-like drops of moisture.

PLATE XV. (Following Plate XIV.)

1. *Sarcina lutea*.—Growth on sterilised potato five days after inoculation from a tube-cultivation. Potatoes, especially old ones, have sometimes a tendency to become discoloured, and the brown appearance in this figure has nothing to do with the growth of the organisms.
2. *Saccharomyces rosaceus* (*Pink torula*).—Growth on sterilised potato which had been inoculated from a colony contaminating a plate-cultivation. This yeast develops a coral-pink colour, but does not grow so luxuriantly as the chromogenic bacteria.

* The term "attenuation" is applied also to a virus, in the sense of weakening or modifying its effect. To avoid confusion the term *mitigation* might be employed exclusively to express this, and *attenuation* used only in the sense indicated above.

FIG.

PLATE XVI. (*Following Plate XV.*)

1. *Bacillus anthracis* (*Bactériidie du charbon*; *Bacillus of splenic fever, wool-sorter's disease, or malignant pustule*).—The bacillus of anthrax grows very rapidly on sterilised potato, especially when placed in the incubator at the temperature of the blood. The growth forms a creamy-yellow layer, with copious spore-formation.
2. *Bacterium indicum* (*Micrococcus indicus*, Koch; *Bacillus indicus*, Flügge).—Sterilised potato inoculated with a pure growth obtained by successive cultivations. Unless the growth is quite free from the presence of other bacteria, the brilliant red colour is not obtained.

PLATE XVII. (*Following Plate XVI.*)

- 1 and 2. *Bacillus cyanogenus* (*Bacterium syncyanum*; *Bacillus of blue milk*).—Potato inoculated from a cultivation in nutrient gelatine. In three days a peculiar bluish-green growth develops on the surface of the potato, and in nine days it has a heaped-up margin of a bluish-green colour, while the central portion has turned almost black.

PLATE XVIII. (*Facing page 104.*)

1. *Bacillus tuberculosis*.—Pure cultivation of the tubercle-bacillus on blood-serum solidified obliquely.
2. Pure cultivation on solid blood-serum in a glass capsule.
3. The same as Fig. 2, examined under a low power of the microscope. $\times 80$.
4. Impression-preparation showing the peculiar serpentine growth of the colonies on blood-serum. $\times 700$.

(After Koch, *Mittheil. a. d. Kaiserl. Gesundheitsamt.*)PLATE XIX. (*Facing page 228.*)

1. *Streptococcus pyogenes*.—From a microscopical specimen prepared from pus from a pyæmic abscess. Stained with gentian-violet by the method of Gram, and contrast-stained with eosin. $\times 1200$. Powell and Lealand's apochromatic $\frac{1}{2}$ Hom. imm. E. P. 10.
2. From a microscopical specimen of an artificial cultivation of the streptococcus in broth. Complex chains with elements dividing both longitudinally and transversely, and varying considerably in size in different lengths of the same chain. $\times 1200$. Powell and Lealand's apochromatic $\frac{1}{2}$ Hom. imm. E. P. 10.

PLATE XX. (*Facing page 300.*)

1. *Bacillus lepræ*.—From a section of the skin of a leper. The section is, almost in its entirety, stained red, and, with moderate amplification, has a finely granular appearance. Stained with Ehrlich's method (fuchsine and methylene-blue). $\times 200$.

FIG.

2. Part of the same preparation with high amplification, showing that the appearances described above are due entirely to an invasion of the tissue by the bacilli of leprosy. $\times 1500$.

PLATE XXI. (*Facing page 308.*)

1. *Bacillus tuberculosis*.—From a section of a lymphatic gland of a foetal calf. The preparation was stained by the Ehrlich-Koch method (methyl-violet and bismarck-brown), and eosin. The giant cell takes the eosin stain, the nuclei are stained brown, and the bacilli blue. In the interior of the giant cell are numerous coloured grains, the significance of which is not known, and a number of tubercle-bacilli. $\times 1500$.

For the material from which this preparation was made the author is indebted to Professor Johne, by whom an account of this case was published, "Ein Zweifelloser Fall von Congenitaler Tuberkulose," *Fortsch. d. Med.*, 1885, No. 7, p. 198.

2. From a section of a lung of a rabbit after inoculation with tubercular sputum. Caseous areas are seen, and masses of bacilli showing distinct beading. Stained by the Ehrlich-Koch method (methyl-violet) without a contrast stain. $\times 1500$.

PLATE XXII. (*Following Plate XXI.*)

1. *Bacillus tuberculosis*.—From a section of the liver of a tubercular hen. With a moderate power the areas of caseation and the topographical distribution of the bacilli can be studied. Stained with the Ehrlich-Koch method (methyl-violet and bismarck-brown). $\times 400$.
2. From the same preparation with high amplification, showing that the parts stained blue consist entirely of bacilli. $\times 1500$.

PLATE XXIII. (*Facing page 316.*)

1. *Bacillus anthracis* (*Bactériidie du charbon; Bacillus of splenic fever, wool-sorter's disease, or malignant pustule*).—From a section of the mucous membrane of the stomach of a mouse. The glandular capillaries are mapped out by the bacilli. Stained by the method of Gram (gentian-violet), and eosin. $\times 500$.
2. From a section of a kidney of a mouse. Under a low power the preparation has exactly the appearance of an injected specimen. Under higher amplification the bacilli are seen to have threaded their way along the capillaries between the tubules, and to have collected in masses in the glomeruli. Stained with Gram's method (gentian-violet), and eosin. $\times 500$.

PLATE XXIV. (*Following Plate XXIII.*)

FIG.

1. *Bacillus anthracis* (*Bactériidie du charbon; Bacillus of splenic fever, wool-sorter's disease, or malignant pustule*).—From a section of the liver of a mouse which had died after inoculation with a pure cultivation of the bacillus. The bacilli are seen to have threaded their way between the liver-cells. The preparation is triple-stained by combining the methods of Weigert and Orth. $\times 500$.
2. *Bacillus anthracis* and *Micrococcus tetragonus*.—From a section of a lung of a mouse which had been inoculated with anthrax three days after inoculation with *Micrococcus tetragonus*. A double or mixed infection resulted. Anthrax-bacilli occurred in vast numbers, completely filling the small vessels and capillaries, and in addition there were great numbers of the characteristic tetrads. Stained with Gram's method (gentian-violet), and eosin. $\times 500$.

PLATE XXV. (*Facing page 332.*)

1. *Bacillus of septicæmia of mice*.—From a section of a kidney of a mouse which had died after inoculation with a pure cultivation of the bacillus. With moderate amplification, the white blood-corpuscles have a granular appearance, and irregular granular masses scattered between the kidney tubules are seen. Stained with Gram's method and eosin. $\times 200$.
2. Part of the same preparation with high amplification. The granular appearances are found to be due to the presence of great numbers of extremely minute bacilli. $\times 1500$.

PLATE XXVI. (*Facing page 334.*)

1. *Bacillus of swine-erysipelas* (*Bacillus of German swine-fever*).—Pure cultivation in nutrient gelatine. The growth is stated to be identical with that of the bacillus of septicæmia in mice.
2. Colonies on a plate-cultivation.
3. Cover-glass preparation of blood from an inoculated pigeon.
(After Schütz, *Arb. a. d. Kais. Gesundh. Amt*, Bd. I.)

PLATE XXVII. (*Facing page 344.*)

1. *Bacillus figurans* (*Bacterium Zopfii*).—From an impression-preparation of a growth on the surface of nutrient gelatine. $\times 50$.
2. Part of the same preparation with high amplification, showing that the coils and filaments of the growth are due to a peculiar and regular arrangement of the individual bacilli. $\times 1500$.

PLATE XXVIII. (*Facing page 368.*)

1. Yeast fungi, or saccharomycetes; and mould fungi, or hyphomycetes.—*Actinomyces* teased out in the fresh state and stained with eosin.

FIG.

2. *Torula cerevisiae* (after Rees).
3. *Saccharomyces mycoderma*, or *oidium albicans*, from an artificial cultivation (after Grawitz).
4. *Saprolegnia* (after Sachs).
5. *Oidium lactis* (after Flügge).
6. Fungi of *favus*, or *oidium lactis* (after Neumann).
7. *Fenicillium glaucum* (after Flügge).
8. *Aspergillus niger*, from a preparation mounted in glycerine.
9. *Aspergillus niger*, from the same preparation (Zeiss $\frac{1}{2}$ o.i.).
10. *Aspergillus glaucus* (after De Bary).
11. *Botrytis Bassiana* (after De Bary)

PLATE XXIX. (Facing page 384.)

1. *Actinomyces* (bovis).—From a section of a maxillary tumour in a cow. Stained by Weigert's method (orseille and gentian-violet). $\times 900$.
2. From a section of the lung of a cow. The rosettes are much smaller, possibly owing to their being more confined by their surroundings than when growing in the soft pulpy tissue of the maxillary tumour. They are here shown with high amplification, but under a power of about 50 diam. (Zeiss A.A. Oc. 2) the section of a lung resembles miliary tuberculosis, and in the centre of a neoplasm the rosette appears about the size of a pin's head. Stained with Weigert's method (orseille and gentian-violet.) $\times 500$.

PLATE XXX. (Following Plate XXIX.)

1. *Actinomyces* (bovis).—From a section of a maxillary tumour in a cow. Stained by Plaut's method (magenta and picric acid). $\times 90$.
Note.—Neelsen's solution may be used instead of magenta solution.
2. Part of the same preparation, with higher amplification. The fungoid masses are very deeply stained by this method. The component club-shaped elements and their radiate arrangement are clearly shown. $\times 500$.

PLATE XXXI. (Facing page 390.)

1. *Actinomyces* (hominis).—From a preparation of the grains from an actinomycotic abscess in a boy. Examined in glycerine. The drawing has been made of a complete rosette, examined by focussing successively the central and peripheral portions. Towards the centre the extremities of the clubs are alone visible; they vary in size; and if pressed upon by the cover-glass, give the appearance of an irregular mosaic. Towards the periphery the clubs are seen in profile, and their characteristic form recognised. At one part there are several elongated elements, composed of separate links. $\times 1200$.

FIG.

2. Different forms of clubs from preparations in which the rosettes have been flattened out by gentle pressure on the cover-glass. $\times 2500$.
- (a) Single club.
 - (b) Bifid club.
 - (c) Club giving rise to four secondary clubs.
 - (d) Four clubs connected together, recalling the form of a branch of bananas.
 - (e) Mature club with a lateral bud.
 - (f) Apparently a further development of the condition represented at (e).
 - (g) Club with a lateral bud and transverse segmentation.
 - (h) Single club with double transverse segmentation.
 - (i) Club with oblique segmentation.
 - (j) Collection of four clubs, one with lateral gemmation, another with oblique segmentation.
 - (k) Club with lateral buds on both sides, and cut off square at the extremity.
 - (l) Club with a daughter club which bears at its extremity two still smaller clubs.
 - (m) Club divided by transverse segmentation into four distinct elements.
 - (n) Elongated club composed of several distinct elements.
 - (o) and (p) Clubs with terminal gemmation.
 - (q) Palmate group of clubs.
 - (r) Trilobed club.
 - (s) Club with apparently a central channel.
 - (t) Filament bearing terminally a highly refractive oval body.

PLATE XXXII. (*Facing page 392.*)

1. *Actinomyces (hominis)*.—From cover-glass preparations of the fungus teased out of the new growths produced by inoculation of a calf with pus from a boy suffering from pulmonary actinomycosis. Stained by Gram's method and orange-rubin. The threads are stained blue and the clubs crimson. In the younger clubs the thread can be traced into the interior of the club. In some of the older clubs the central portion takes a yellowish stain, and in others the protoplasm is not continued as a thread, but is collected into a spherical, or ovoid, or pear-shaped mass. In others, again, irregular grains stained blue are scattered throughout the central portion.
2. From a section of a portion of the growth removed from a boy during life. The tissue was hardened in alcohol, and cut in celloidin. The section was stained with Gram's method and orange-rubin. There are several rosettes, surrounded by granulation tissue.
3. A mass of extremely fine filaments occupies the central part of the rosette. Many of the filaments have a terminal enlargement. The marginal part shows a palisade of clubs stained by the orange-rubin. $\times 500$.

BACTERIOLOGY.



INTRODUCTION.

THE GERM THEORY.

THE researches of Pasteur into the *rôle* played by micro-organisms in the processes of fermentation and putrefaction, and in diseases such as anthrax, the silkworm malady, pyæmia, septicæmia, and chicken cholera, have invested the science of Bacteriology with universal interest and vast importance. The further researches of the practical mind of Lister, with the resulting evolution of antiseptic surgery, have demonstrated the necessity for a more intimate acquaintance with the life-history of these micro-organisms; while the more recent investigations which have established the intimate connection between bacteria and infective diseases, and more especially the discovery by Koch of the tubercle and cholera bacilli, have claimed the attention of the whole thinking world.

Those bacteria which are connected with disease, and more especially those which have been proved

to be the *causa*, if not the actual *materies morbi*, are of predominant interest and importance.

The first attempt to demonstrate the germ theory of disease dates back almost to the discovery of the microscope. Athanasius Kircher, nearly two and a half centuries ago, expressed his belief that there were definite micro-organisms to which diseases were attributable. The microscope had revealed that all decomposing substances swarmed with countless micro-organisms which were invisible to the naked eye, and Kircher sought for similar organisms in diseases which he considered might be due to their agency. The microscope which he described* obviously could not admit of the possibility of studying, or even detecting, the micro-organisms which are now known to be associated with certain diseases; and it is not surprising that his teachings did not at the time gain much attention. They were destined, however, to receive a great impetus from the discoveries which emanated from "the father of microscopy."

Antony van Leeuwenhoek had learned as a youth to grind and polish lenses, and later in life employed his spare time in constructing microscopes, and in conducting those researches which have made for him a name which is familiar to all microscopists. His researches were published in a series of letters to the Royal Society. In 1675 he described extremely minute organisms in

* *Ars Magna Lucis et Umbræ*, 1646.

rain-water, well-water, infusions of pepper, hay, and other vegetable and animal substances, in saliva, and in scrapings from the teeth ; and, further, he was able to differentiate these minute living things by their size, their form, and the character of their movements. In 1683 these discoveries were illustrated by means of woodcuts, and there can be little doubt, from the drawings of these micro-organisms, that they are intended to represent leptothrix filaments, vibrios, and spirilla. Indeed, we can almost recognise these micro-organisms as bacteria from Leeuwenhoek's graphic descriptions, apart from his figures. They were described as moving in the most characteristic manner, progressing with great rapidity, or spinning round like a top, and as so excessively minute that they were only perceived with great difficulty. The smallest forms could hardly be examined individually ; but, viewed *en masse*, they closely resembled a swarm of gnats or flies. In another communication, published in 1692, he gives some idea of the size of these animalcules by stating that they were a thousand times smaller than a grain of sand. Others which, comparatively speaking, were of considerable length, were characterised by their peculiar mode of progression, bending and rolling on themselves, movements which, he adds, created both delight and astonishment in the mind of the observer. Leeuwenhoek himself was not disposed to believe in the possibility of such organisms

being found in the blood in disease; but as soon as he had proved the actual existence of such minute creatures, theoretical physicians were not wanting who at once attributed various maladies to their agency. Among these, Nicholas Andry is made conspicuous by his work published in 1701.* Andry classed the minute organisms discovered by Leeuwenhoek as worms.

In 1718 Lancisi believed that the deleterious effect of the air of malarial districts depended upon animalcules, and others considered that the plague in Toulon and Marseilles in 1721 arose from a similar cause. In fact, by some, all diseases were attributed to vermicules, and this led to the theory being ridiculed and discredited.

In spite of adverse criticism, the theory of contagium vivum survived, and Linnæus acknowledged it by placing the micro-organisms discovered by Leeuwenhoek, the contagia of specific fevers, and the causes of putrefaction and fermentation, into one order—"chaos." The theory was further supported by the writings of Plenciz, who, in 1762,† very ably discussed the nature of contagium, as well as the relation of animalcules to putrefaction and disease. However, no proofs in support of these theories were forthcoming, and gradually the

* *De la Génération des Vers dans le Corps de l'Homme.*

† *Opera Medico-Physica, Tractatio de Contagio, de Lue Bovina, de Variolis; de Scarlatina.*

idea of *contagium vivum* fell into obscurity, and indeed came to be regarded by some as an absurd hypothesis.

Though a causal relation of animalcules to diseases was for a time discredited, the natural history of these micro-organisms was studied with increasing interest. In 1778 Baron Gleichen* described and figured a great number of micro-organisms which he had discovered in various vegetable infusions. Joblot, Lesser, Réaumur, Hill, and many others worked at the same subject. Hill† remarked that there was hardly the least portion of matter or the least drop of fluid of any kind naturally found in the earth which was not inhabited by multitudes of animalcules. But these observers inclined rather to searching for new forms than to studying more thoroughly those which had been already discovered; and, as a result, but little scientific progress was made until the time of Müller, of Copenhagen. Müller, in 1786, criticised the work of these previous writers, and pointed out that they had been too much occupied with merely finding new micro-organisms. Müller took into account the form of the micro-organism, its mode of progression, and other biological characters, and on such data based a classification. Thus the scientific knowledge of

* *Dissertation sur la Génération, les Animalcules Spermatiques et ceux d'Infusions.*

† *Essays in Natural History and Philosophy.*

these minute beings was considerably advanced by his writings and his illustrations.

The subject which now eclipsed all others in interest was the origin of these micro-organisms. Two rival theories were widely discussed—spontaneous generation and development from pre-existing germs; and the researches that were made in the course of this discussion, and the discoveries which resulted, indirectly materially advanced the germ theory of disease, and explain many of the phenomena in the life-history of the pathogenic microbes which have been brought to light in recent years.

Spontaneous development of micro-organisms in putrescible infusions was believed in by many, but was supported by no one with greater persistency than Needham. Needham found, if meat infusion were boiled and transferred to a well-stoppered flask, that animalcules readily developed, and he could only explain this by supposing that they originated spontaneously from the material of the infusion. In 1768 Bonnet* strenuously opposed these conclusions on purely theoretical grounds, and maintained that it was far more probable that the ova of the animalcules were present in the infusions or were suspended in the air enclosed in the flask.

Spallanzani was the first to demonstrate by experiment the correctness of Bonnet's arguments

* *Considérations sur les Corps Organisés.*

It occurred to him to boil the infusion in flasks, and *to seal the vessels during the process of boiling*. As a result the flasks remained free from putrefaction, and animalcules only developed when the infusion was exposed to the air by making a hole in the flask. That Spallanzani's experiments were reliable, and his conclusions correct, was evidenced by the fact that his simple precaution led to great practical results, as François Appert introduced on this principle the method of preserving meats, vegetables, and other provisions.

The disciples of Needham nevertheless brought forward counter objections. Treviranus urged that a certain *quantity* and *quality* of air was necessary for the spontaneous development of these infusoria, and that by sealing the flasks too small a quantity of air was in contact with the infusion, and, further, that this air had become changed in quality by the process of boiling.

Spallanzani argued against these objections, but did not support his opinions by further experiments, so that the question remained for a time undecided.

In 1836 Francis Schulze devised an experiment which brought still further evidence against Needham's theory. Schulze filled a glass vessel half full with distilled water and different animal and vegetable substances. This was plugged with a doubly bored cork, and through each perforation a glass tube was introduced, bent at a right angle.

On boiling the flask, steam issued freely from each tube, and all parts were thoroughly sterilised. Each tube was then connected with a bulbed tube, one bulb containing concentrated sulphuric acid, and the other a solution of potash. By aspiration fresh air was drawn into the flask, and this was deprived of any germs which might be present by its passage through the sulphuric acid. The result was that the infusion remained without any development of micro-organisms. If, on the other hand, air were admitted without first being drawn through the sulphuric acid, the infusion in a short time teemed with animalcules. In other words, Schulze demonstrated that in spite of free access to air, *which had not been heated*, the infusions remained free from germs.

Schwann, in 1837, arrived at similar results. He found that putrescible substances remained sterile if exposed to an abundant supply of air which was heated by being passed through a melted mixture of metals. This convinced him that the cause of the decomposition which would otherwise have occurred must exist in the air.

The objection remained that in the experiments of Schulze and Schwann, the air which was admitted to the flasks had undergone either a *chemical* or a *thermal* change, and therefore the theory of Needham was not yet entirely disposed of.

In 1854 the final blow was dealt by Schroeder and Van Dusch. These investigators demonstrated

that decomposition could be obviated without resorting either to thermal or chemical treatment of the air, as simple filtration of the air through cotton-wool was shown to be efficacious in excluding germs. Finally, Hoffman, in 1860, and, independently, Chevreuil and Pasteur, in 1861, showed that even cotton-wool could be dispensed with, as a sterile solution might be kept sterile if the neck of the vessel were bent with an S-shaped curve. Micro-organisms in the air entering the flask, were deposited by gravitation in the bend of the tube.

The advocates of spontaneous generation were ready with fresh objections. They now urged that the medium lost its power of undergoing decomposition by being boiled. This objection was at once set aside by the fact that if unfiltered air were admitted to the infusion decomposition set in. Additional evidence was brought against spontaneous generation by the experiments of Pasteur, Burdon Sanderson, Lister, and others, in which it was shown that blood, urine, and milk would remain without decomposition, if all precautions were adopted to avoid contamination in filling the sterilised flasks.

Even at this stage of this great scientific controversy fresh difficulties arose, for it was found that in certain solutions which had been boiled and hermetically sealed in flasks micro-organisms made their appearance. In 1872 Charlton Bastian

published a research* which was to prove that spontaneous generation actually took place.

Decoctions of turnip and cheese which had been filtered, neutralised, and boiled for ten minutes, and hermetically sealed during the boiling, were found after a time to contain micro-organisms. These results, however, were before long explained by the fact that in milk, infusions of hay, and certain other decoctions, the spores of bacilli are present, which are much more resistant than the bacilli themselves. In such cases mere scalding or boiling for a few minutes will not sterilise the solution. The bacilli are destroyed, but not their spores; and if the latter remain unhurt, they will germinate, increase, and multiply. But if, as Tyndall found, the boiling be repeated a second and a third time, all the spores will be destroyed; for in the intervals between the boilings the spores sprout into bacilli, and the bacilli at the next boiling perish; so that after three or four repeated boilings the infusion is rendered perfectly free from germs.

While this discussion was occupying the attention of the whole scientific world, some investigators had been again following up the theory of a connection between micro-organisms and disease.

In 1837 Cagniard Latour and Schwann independently made the discovery that the yeast plant was a living organism, and the true cause of the fermentation. The close analogy between the

* *Proceedings of the Royal Society.*

processes of fermentation and of certain diseases had long been held ; and, therefore, when it was proved that fermentation was due to a micro-organism, fresh advocates appeared in support of the theory that diseases were produced by similar agencies. Boehm, in 1838, described certain organisms in cholera which was at that time raging in Europe ; but the publication of Bassi, who a year previously had discovered the cause of a disease of silkworms, attracted much greater attention.

Bassi discovered that in this disease extremely minute spores existed on the bodies of the worms, which were conveyed from the sick to the healthy. They destroyed the healthy worms by germinating in their skins and growing into their bodies. These discoveries may be said to have brought the theory of *contagium vivum* to life again ; and Henle, in reviewing the facts of the case in 1840, came to the conclusion that the cause of all contagious diseases must be of a living nature, and this he maintained, although he had searched in vaccine and small-pox lymph, in the desquamation of scarlet fever, and in other diseases without success.

Bassi's discovery and Henle's doctrine encouraged a number of investigators, and remarkable results followed. In favus, in herpes tonsurans, in pityriasis versicolor, fungus threads and spores were found, and were regarded as being of etiological importance, inasmuch as the morbid lesions corresponded with the growth of the particular fungus.

Cholera became especially a subject for research. Swaine, Brittan, and Budd found micro-organisms in choleraic dejecta. Davaine described certain monads in the intestinal contents, but no causal connection was established between these organisms and the disease; and when the cholera disappeared the interest in *contagium vivum* waned, and was eclipsed by the question of fermentation. The discoveries which followed in this subject had, as we shall see, a very important bearing on the micro-parasitic origin of communicable diseases.

Pasteur followed up the researches of Cagniard Latour and Schwann, and in 1857 demonstrated that the lactic, acetic, and butyric fermentations were also produced by micro-organisms.

Previous to this, in 1850, Davaine and Rayer had noted the existence of little rod-like or filamentous bodies about the size of a blood corpuscle in the blood of a sheep that had died of splenic fever. Pollender had seen similar bodies in the blood of cows.

Davaine did not pay much heed to his discovery; but in 1863 he thoroughly reinvestigated the subject, and conducted a series of experiments which led him to the conclusion that the actual cause of splenic fever was an organised being whose presence and multiplication in the blood produced changes in that fluid of the nature of fermentation, resulting in the death of the animal.

These conclusions were not accepted by all; and, indeed, the evidence was so far incomplete that the sceptic was justified in considering that these experiments afforded only a working hypothesis. But Davaine's comparison between this disease and fermentation attracted the attention of Pasteur whose mind had been fully trained for entering upon this investigation by the researches which he had been carrying on in the interval between Davaine's publications of 1857 and 1863.

Pasteur, as already mentioned, had been working at fermentation, and his attention was next directed to studying the so-called diseases of wines, and subsequently to a contagious disease which committed ravages among silkworms. By laborious researches Pasteur was able to confirm the belief that the disease was due to the presence of microorganisms only discernible with the aid of the microscope. These oval shining bodies in the moth, worm, and eggs had been previously observed by Cornalia, and described by Naegeli as *Nosema bombycis*, and by Lebert as *Panhistophyton*. But it was reserved for Pasteur to introduce a means of combating the disease. Pasteur showed that if a silkworm, whose body contained these microorganisms, were pounded up with water in a mortar, and the mixture painted with a brush on the leaves on which healthy worms were fed, they would all without fail succumb to the disease.

As the contagious particles were transmitted to

the eggs, a method for preventing the spread of the disease suggested itself. Each female moth was kept separate from the others, and allowed to deposit her eggs on a small linen cloth. The moth was then pinned to the corner of the cloth, and left for future examination. When the time for this arrived, the moth was crushed up with water in a mortar, and a drop examined under the microscope. If any trace of corpuscular matter were found to be present, the cloth with its collection of eggs was burnt; and if not, the eggs were set aside for use.

Complete as this appears to be as a demonstration of a causal connection between the micro-organisms and the disease, it could obviously be objected that there was no distinct proof that the corpuscular bodies constituted the actual contagium. There was no isolation of the organisms, no artificial cultivation of them apart from the diseased moth or worm, and subsequent production of the disease by means of the isolated organisms. The same objection was applicable to Davaine's investigations. Davaine found the rods in association with the disease, and maintained that they were causally related; but others stated that it was possible to inoculate animals with anthrax blood containing rods, and to produce the disease without being able to detect the rods again in the blood of the animal experimented upon. It was also urged that it was possible to infect with anthrax blood after the rods

had disappeared, and to find a reappearance of the bacilli in the blood of the inoculated animal.

What appeared also to lend great support to these objections was the well-known fact that anthrax or splenic fever was especially prevalent in certain seasons and certain localities. The disease, in fact, was regarded by some as originating from peculiar conditions of climate and soil. The fallacies in these objections were, however, rapidly dispelled. Bollinger, in 1872, pointed out that the blood, from which the rods had disappeared, was still virulent owing to the presence of the *spores* of the bacillus, and that it was owing to the soil being impregnated with these spores that the disease broke out in certain localities. But there still remained many who refused to regard these particles as living bodies, some looking upon them simply as crystals, and the question of their importance remained undecided for several years.

In 1877 Robert Koch published a memoir in which he fully described the life-history of the anthrax or splenic fever bacillus, and gave a complete demonstration of the life-history of the micro-organism, and the definite proofs of its pathogenic properties. He pointed out how quickly the rods grew in the blood and tissues by lengthening and by cross division. Further, that in the blood or in serum or in aqueous humour they grew into long leptothrix filaments, and formed numerous shining spores. He traced, by continuous observation on

the warm stage, the whole life cycle from the fission of the rods to the formation of spores and the sprouting of the spores into fresh rods. Further, he carried on the disease by inoculating from mouse to mouse for several generations, and observed that in the blood of the animal and in the swollen spleen the glass-like rods were always to be found.

Pasteur also studied the microbe of splenic fever, and amply confirmed and extended the observations of Koch by his researches on the attenuation of the anthrax virus.

Pasteur also met with adverse criticism. Paul Bert argued that the bacilli were of no importance, because he could destroy them by exposing material containing them to great pressure, and yet the material produced the disease on inoculation. But such measures did not destroy the *spores*; and finally Paul Bert was convinced of his error when Pasteur demonstrated cultures of the anthrax bacillus in urine, from which successive generations were started, and that with such cultivations the disease could always be produced.

It was, however, principally the researches of Koch which placed the doctrine of *contagium vivum* on a scientific basis.

Koch's improvements in the methods of cultivation, his recommendation of the necessary microscopical apparatus, his histological methods for examining these minute organisms, and his famous postulates for proving beyond controversy the existence of

specific pathogenic micro-organisms, elevated the theory of *contagium vivum* to a demonstrated and established fact. The chain of evidence regarded by Koch as essential for proving the existence of a pathogenic organism was composed as follows:—

1. The micro-organism must be found in the blood, lymph, or diseased tissue of man or animal suffering from or dead of the disease.

2. The micro-organisms must be isolated from the blood, lymph, or tissues, and cultivated in suitable media, *i.e.*, outside the animal body. These pure cultivations must be carried on through successive generations of the organism.

3. A pure cultivation thus obtained must, when introduced into the body of a healthy animal, produce the disease in question.

4. Lastly, in the inoculated animal the same micro-organism must again be found.


It is, however, impossible by localising one's knowledge to pathogenic species to thoroughly understand the life-history of these particular forms, or to be able to grasp and appreciate the various arguments and questions that arise in comparing their life-history with the progress of disease.

Nor is it sufficient to know only how to recognise and artificially cultivate a bacterium associated with disease; we must endeavour to establish the exact relationship of the bacterium to the disease in question.


The determination of the true pathogenic microbe is beset with fallacies. In many diseases bacteria have been regarded as the actual contagia, until a searching inquiry by other investigators has shown that the evidence was most unsatisfactory or entirely misleading. For example, in diseases with lesions of the external or internal linings of the body, extraneous micro-organisms may get into the circulation and be swept into the internal organs, where they either perish in the battle with the healthy tissues which are opposed to their existence, or they may gain the upper hand and set up destructive processes. Such organisms, when found in association with these diseases, may be discovered in the blood and internal organs; and though accidental epiphytes, often associated with septic complication, they may only too readily be accepted by the enthusiast as the actual contagium of the disease in question.

It is only when such fallacies are exposed that we are brought once more face to face with the fact that the nature of the contagium in many diseases, such as hydrophobia, variola, vaccinia, scarlet fever, and measles, is at the present day undetermined.

Such cases invite further research; but these fallacies must be borne in mind, or the application of bacteriology to the elucidation of the germ theory of disease will not only be misleading, but tend to retard rather than assist the progress of science.



BACTERIA, SCHIZOMYCETES OR FISSION-FUNGI.



Koch's postulates naturally suggest a sequence in the various processes which must be adopted in a practical study of micro-organisms associated with disease. Inasmuch, however, as these processes embrace those which are employed in the isolation, cultivation, etc., of non-pathogenic species, we shall, in studying the bacteria as a whole, adopt the order suggested. After an introduction to the apparatus commonly employed in a bacteriological laboratory, we shall describe the methods of examining liquids, tissues, etc., and the means of recognising micro-organisms. Then will follow the methods of isolating these micro-organisms from such liquids, tissues, etc., and of carrying on pure cultivations in nutrient media. Lastly, we shall refer briefly to experimental researches on the living animal, and the means of isolating micro-organisms from the liquids and tissues of the body after death.

In Part II. will be found chapters upon the General Biology of bacteria, and in Part III. a chapter upon their classification, followed by a description of each species, more particularly of those of pathological interest, with a detailed account of the special methods of examination and of staining employed for demonstrating the different species.

In the Appendix a descriptive list of important yeasts and moulds will be given, with any special technique required in their case. Yeasts and moulds are constantly encountered in the investigation of

the bacteria in air, soil, and water ; and several are of interest in being, like many bacteria, micro-organisms associated with disease. An account is given of Actinomycosis, of Flagellated Protozoa which have been found in the blood of animals, and of the micro-organisms associated with malaria.



PART I.
GENERAL METHODS.

CHAPTER I.

APPARATUS, MATERIAL, AND REAGENTS EMPLOYED IN A BACTERIOLOGICAL LABORATORY.

(A) HISTOLOGICAL APPARATUS.

Microscope.—For the investigation of micro-organisms a good microscope with oil-immersion system and a condenser, such as Abbé's, is essential. Such instruments are supplied by Leitz, Zeiss, and Hartnack in Germany, and Powell & Lealand, and Swift & Son, in England. Zeiss' microscope, with $\frac{1}{12}$ and $\frac{1}{18}$ oil-immersion lenses, or Powell & Lealand's, with $\frac{1}{12}$ and $\frac{1}{25}$, is recommended for investigators; while Swift's or Leitz', with $\frac{1}{12}$, is a serviceable and economical one for students.* In addition to the usual microscopic fittings, Zeiss supplies a micrometer eyepiece, with directions for use. Some such arrangement is essential for the measurement of bacteria. Other accessories to the microscope are—

A large bell-glass for covering the microscope when not in use.

About a foot square of blackened plate-glass.

* Leitz', with $\frac{1}{12}$, costs about £15; Zeiss', with the same, £30, and with $\frac{1}{18}$, £20 more. Refer to foot-note on p. 61.

A white porcelain slab of the same size.

Glass bottles with ground-glass stoppers, for alcoholic solutions of aniline dyes, etc.

Glass bottles with funnels, for aqueous solutions of the dyes, and others provided with pipettes.

A small rod-stoppered bottle of cedar oil. This is recommended by Zeiss in preference to other oils for his immersion lenses.

Set of small glass dishes or capsules, and watch-glasses for section staining, etc.

Stock of best glass slides, in packets of fifty.

Several boxes of round and square thin cover-glasses, in various sizes, of the best quality.

Needle-holders, with a couple of platinum needles, and a packet of ordinary sewing-needles.

Glass rods drawn out to a fine point; useful for manipulating sections when acids are employed.

Copper lifters, preferably plated.

One pair of small brass or spring-steel platinum-pointed forceps, for holding cover-glasses.

One pair of brass tongs.

Collapsible tubes for containing Canada balsam; very serviceable for transport and general use.

Turn-table, for sealing cover-glass preparations with rings of cement.

Boxes for preparations, book-form.

Tickets and labels, various sizes.

Soft rags or old pocket-handkerchiefs, for removing cedar oil after use of immersion lens, cleaning cover-glasses, etc.

Chamois leather for wiping lenses.

Microtome.—Schanze's is much in favour in Germany; but Jung's, of Heidelberg,* though a

* Price lists may be obtained from any of the above-mentioned firms, from which an idea of the instruments can be formed, and a comparison of the prices made.

somewhat cumbrous instrument, is preferred by others. Smaller accessories, which should be within reach, are—

A small can of sewing-machine oil.

A soft rag and chamois leather, for wiping the knives immediately after use.

Stone and leather for setting and sharpening the same.

Two or three camel's-hair brushes.

A Freezing Microtome is very useful. The method of embedding in celloidin may be used with advantage in combination with the ordinary process of freezing.

(B) REAGENTS AND MATERIAL EMPLOYED IN THE PROCESSES OF HARDENING, DECALCIFYING, EMBEDDING, FIXING, AND CUTTING OF TISSUES.

Alcohol, absolute.

Bergamot oil.

Celloidin.

Dissolved in equal parts of ether and alcohol.

Cork, or stock of ready-cut corks.

Ebner's solution. A mixture in the following proportions :—

Hydrochloric acid	5
Alcohol	100
Distilled water	20
Chloride of sodium	5

Gelatine.

Melted in a small porcelain capsule, and set aside ready to be re-melted when required for use.

Glycerine gelatine (Klebs).

Best well-washed gelatine 10

Add distilled water, allow gelatine to swell up, pour off excess of water, melt gelatine with gentle heat, add

Glycerine 10

Lastly, a few drops of phenol for preservation.

Gum.**Kleinenberg's solution.**

Saturated watery solution of picric acid 100

Strong sulphuric acid 2

Filter and add

Distilled water 300

Müller's fluid.

Bichromate of potash 2

Sulphate of sodium 1

Distilled water 100

Osmic acid.

Distilled water 100

Osmic acid 5

Paper trays (small or glass-capsules).

Paraffine.

Spermaceti.**Xylol.**

Hardening and decalcifying solutions should be kept in stock in quantities according to requirement. A jacket of brown paper should be pasted round a well-stoppered bottle to contain osmic acid to efficiently protect it from light, and it should be kept in a cool place. The celloidin solution may be kept in stock in a wide-mouthed glass bottle, from which small wide-mouthed bottles may be filled according to the number required. To put several pieces of different tissues in the same bottle leads to confusion.

(c) REAGENTS FOR EXAMINING AND STAINING
MICROSCOPICAL PREPARATIONS.

1. **Acetic acid, strong.**
2. **Alcohol, absolute.**
3. **Alcohol, 60 per cent.**
4. **Alcohol, acidulated.**

Alcohol	100
Hydrochloric acid	1

5. **Alum carmine (Grenacher).**

Carmine	1
Five per cent. solution of alum	100

Boil twenty minutes, filter when cold.

6. **Ammonia, strong.**7. **Aniline.**8. **Aniline water.**

Distilled water	100
Aniline	5

Shake well and filter emulsion.

9. **Bismarck brown.**

(a) Concentrated solution in equal parts of glycerine and water.

(b) Aqueous solution.

Bismarck brown	2
Alcohol	15
Distilled water	85

10. **Borax carmine (Grenacher).**

Borax	2
Carmine	5
Distilled water	100

To the dark purple solution add a 5 per cent. solution of acetic acid until a red colour is produced; set aside twenty-four hours, filter, and add a drop of carbolic acid.

11. **Cedar oil.**12. **Eosin.**

(a) Saturated alcoholic solution.

(b) Aqueous solution.

Distilled water	100
Eosin	5

13. Ether.

14. Fuchsine.

(a) Saturated alcoholic solution.

(b) Aqueous solution.

Fuchsine	2
Alcohol	15
Water	85

15. Gentian violet.

a) Saturated alcoholic solution.

Aqueous solution.

Gentian violet	2.25
Distilled water	100

16. Gibbes' solution, for double staining.

Take of

Rosaniline hydrochlorate	2
Methylene blue	1

Triturate in a glass mortar,

Dissolve aniline oil	3
In rectified spirit	15

and add slowly to the above.

Lastly, slowly add distilled water 15

Keep in stoppered bottle.

17. Glycerine, pure.

18. Hæmatoxylin solution.

Hæmatoxylin	2
Alcohol	100
Distilled water	100
Glycerine	100
Alum	2

19. Iodine solution.

Iodine, pure	1
Iodide of potassium	2
Distilled water	50

20. Iodine solution (Gram).

Iodine	1
Iodide of potassium	2
Distilled water	300

21. Lithium-carmin solution (Orth).

Saturated solution of carbonate of lithium	100
Carmin	2·5

22. Magenta solution (Gibbes).

Magenta	2
Aniline oil	3
Alcohol (Sp. Gr. ·830)	20
Distilled water	20

23. Methylene blue.

(a) Concentrated alcoholic solution.

(b) Aqueous solution.

Methylene blue	2
Alcohol	15
Water	85

(c) Koch's solution.

Concentrated alcoholic solution of methylene blue	1
Ten per cent. potash solution	2
Distilled water	200

(a) Löffler's solution.

Concentrated alcoholic solution of methylene blue	30
Solution of potash 1—10,000	100

24. Methyl violet.

(a) Concentrated alcoholic solution.*(b)* Aqueous solution.

Methyl violet	2.25
Distilled water	100

(c) Koch's solution.

Aniline water	100
Alcoholic solution of methyl violet	11
Absolute alcohol	10

25. Neelsen's solution.

Dissolve fuchsine	1
In alcohol	10
Add a 5 per cent. watery solution of carbolic acid.	100

26. Nitric acid, pure.

27. Orseille (Wedl).

Dissolve pure ammonia-free orseille in

Absolute alcohol	20
Acetic acid	5
Distilled water	40

until a dark red liquid results : filter.

28. Picric acid.

(a) Concentrated alcoholic solution.

(b) Saturated aqueous solution.

29. Picro-carmine (Ranvier).

Carmine	1
Distilled water	10
Solution of ammonia	3
Triturate, add cold saturated solution of picric acid	200

30. Picro-lithium-carmine (Orth).

To above-mentioned lithium-carmine
solution add

Saturated solution of picric acid	2'3
---	-----

31. Potash solution.

(a) 1 to 3 per cent.

(b) 10 " "

(c) 33 " "

32. **Safranine.**

(a) Concentrated alcoholic solution.

(b) Watery solution 1 per cent.

33. **Sulphuric acid, pure.**34. **Salt solution 0·8 per cent.**35. **Turpentine.**36. **Vesuvium.**

(a) Concentrated alcoholic solution.

(b) Watery solution.

Water, distilled.**Water, sterilised.**

Distilled water can be kept for use in a wash bottle, or far better in a siphon apparatus. Sterilised water is convenient in plugged sterile test-tubes, which may be kept close at hand in a beaker, or tumbler, with a pad of cotton wool at the bottom. The *numbered* reagents can be conveniently arranged on shelves within easy reach. Alcoholic solutions of the aniline dyes and other special preparations should be kept in bottles with ground-glass stoppers. Aqueous solutions of the dyes may be kept in bottles with funnel filters, and the solution filtered before use. To both aqueous and alcoholic solutions a few drops of phenol, or a

crystal of thymol, should be added as a preservative. For the rapid staining of cover-glass preparations, it is convenient also to have the most frequently used stains (fuchsine, methyl violet) in bottles provided with pipette-stoppers.

(D) REAGENTS FOR MOUNTING AND PRESERVING PREPARATIONS.

Acetate of potash.

Concentrated solution.

Asphalte lac.

Canada balsam.

Dissolved in xylol.

Glycerine gum (Farrant's solution).

Glycerine.

Water.

Saturated solution of arsenious acid.

Equal parts, mix and add of picked gum arabic half a part.

Hollis' glue.

Zinc-white.

(E) DRAWING AND PHOTOGRAPHIC APPARATUS.

Camera Lucida.—The camera lucida of Zeiss is an excellent instrument, though many prefer

the pattern made by Nachet of Paris. Combined with the use of a micro-millimeter objective, it affords also a simple method for the measurement of bacteria.

For drawing macroscopical appearances, and for illustrating microscopical specimens with or without the use of a camera lucida, the following materials should be within reach :—

Pencils.

Etching Pens.

Prepared Indian Ink.

Water-colour Paints and Brushes.

Ordinary and tinted drawing paper and other usual accessories.

Photo-micrographic Apparatus.—Zeiss of Jena, Seibert & Kraft of Wetzlar, Nachet of Paris, and Swift & Son of London, may all be recommended for constructing an arrangement in which the photographic camera is combined with the microscope.

For illumination either sunlight or artificial light may be employed. In the case of sunlight a heliostat is necessary to procure the best results; but as sunlight is not always available by day, and it is also more convenient for many to work at night, it is better to have recourse altogether to artificial light. Excellent results may be obtained with an ordinary paraffine lamp, or with magnesium, oxycalcium, or electric light. Specimens are preferably stained yellow, brown, or black, and for mounting

the preparations Koch* recommends a saturated solution of acetate of potash; but there is little or no objection to the use of Canada balsam dissolved in xylol. Hauser,† who employed the electric light, obtained some excellent pictures of preparations mounted in balsam. Van Ermengem‡ first recommended the isochromatic dry plates, and produced most successful results with the lime-light from objects stained with fuchsine and methyl violet. The author also has investigated the applicability of photographic processes for illustrating micro-organisms. Numerous preparations have been satisfactorily depicted by means of the isochromatic plates without any reference to the staining reagents employed. For a full description of the apparatus and methods employed the reader is referred to the author's publication.

(F) STERILISATION APPARATUS.

Steam-steriliser.—A cylindrical vessel of tin about half a metre or more in height, jacketed with thick felt, and provided with a conical cap or lid (Fig. 1). The lid is also covered with felt, has handles on either side, and is perforated

* Koch, *Verfahren zur Untersuchung zum Conserviren und Photographiren der Bacterien*. 1877.

† Hauser, *Über Fäulniss Bacterien und deren Beziehungen zur Septicämie*. 1885.

‡ Van Ermengem, *Bull. de la Soc. Belge de Microscopie*, No. X., pp. 170-2. 1884.

§ *Photography of Bacteria*. 1887.

at the apex to receive a thermometer. Inside the vessel is an iron grating or diaphragm about two-thirds the way down, which divides the interior into two chambers—the upper or “steam chamber,” and the lower or “water-chamber.” A gauge outside marks the level of the water in the lower chamber; this should be kept about two-thirds full. The apparatus stands upon three legs, and is heated from below with two or three Bunsen, or better, a Fletcher’s burner. It is employed for sterilising nutrient media in tubes or flasks, for cooking potatoes, or hastening the filtration of agar-agar. When the thermometer indicates 100° C. the lid is removed, and test-tubes are lowered in a wire basket by means of a hook and string, and the lid quickly replaced. Potatoes or small flasks are lowered into the cylinder in a tin receiver with a perforated bottom, which rests upon the grating and admits of its contents being exposed to the steam.



FIG. 1.—Koch's
STEAM-STERILISER.

Hot-air Steriliser.—A cubical chest of sheet iron with double walls, supported on four legs; it may also be suspended on the wall of the laboratory, with a sheet of asbestos intervening (Figs. 2 and 3).

It is heated with a rose gas-burner from below,

and the temperature of the interior indicated by a thermometer inserted through a hole in the roof; in a second opening a gas regulator can be fixed. Test-tubes, flasks, funnels, cotton wool, etc., may be sterilised by exposure to a temperature of 150° C. for an hour or more.




FIG. 2.—HOT-AIR STERILISER.




FIG. 3.—SECTION OF HOT-AIR STERILISER.

(G) APPARATUS AND MATERIAL FOR PREPARING AND STORING NUTRIENT GELATINE AND NUTRIENT AGAR-AGAR.

Water-bath.—A water-bath on tripod stand is required for boiling the ingredients of nutrient jellies and for general purposes. The lid may be conveniently composed of a series of concentric rings, so that the mouth of the vessel may be graduated to any size required.

Test-tube Water-bath.—This consists of a circular rack for test-tubes within a water-bath. It is sometimes employed instead of the steam cylinder for sterilising nutrient jelly in tubes, by boiling for an hour for three successive days.

Hot-water Filter.—A copper funnel with double walls, the interspace between which is filled with hot water. A glass funnel fits inside the copper cone, the stem of the glass funnel passing through and being tightly gripped by a perforated caoutchouc plug, which fits in the opening at the apex of the cone. The water in the cone is heated by applying the flame of a burner to a tubular prolongation of the water chamber. In a more recent model, as represented in Fig. 4, this prolongation is dispensed with, and the temperature is maintained by means of a circular burner which acts at the same time as a funnel ring.




FIG. 4.
HOT-WATER FILTERING APPARATUS
WITH RING BURNER.

Glass Vessels.—A number of glass vessels should be kept in stock according to requirements.

Bohemian hard glass flasks are employed in several sizes, for boiling nutrient media. The conical forms are especially used in the larger sizes for storing nutrient jelly.

Glass funnels large and small are necessary, not only in the processes of preparing nutrient jelly, but for filtering solutions of aniline dyes and for general purposes.

A liberal supply of test-tubes should always be kept in stock, as they are not only employed for the tube-cultivations, but can be conveniently used for storing bouillon, sterilised water, etc.

Cylindrical glasses graduated in cubic centimetres, 10 ccm., 100 ccm., 500 ccm., are required for measuring the liquid ingredients of nutrient jelly, and also in preparing the various staining solutions.

A large wide-mouthed glass jar, with a glass cover, is extremely useful. It must be padded at the bottom with cotton wool for containing a stock of tubes of sterilised nutrient jelly, and should be placed within reach on the working table.

Balance and Weights.—A balance, with large pans and set of gramme weights, is constantly required.

Cotton Wool.—The best or “medicated” cotton wool should be procured.

Gelatine.—The gelatine for bacteriological purposes must be of the very best quality (gold label).

Agar-agar.—This is also called Japanese Isin-glass; it consists of the shrivelled filaments of certain Algæ (*Gracilaria lichenoides* and *Gigartina speciosa*).*

Peptonum Siccum (German).

Table Salt.—Prepared table salt can be obtained in tins or packets.

Litmus Papers.—Blue or red litmus paper in cheque books, for testing the gelatine mixture, etc.

Carbonate of Soda.—A bottle, containing a saturated solution of carbonate of soda, and provided with a pipette-stopper, may be kept, especially for use in the preparation of nutrient jelly.

Lactic Acid.

Filter Paper.—For filtering gelatine stout Swedish filter paper of the best quality is recommended.

Flannel or Frieze.—This is employed as a substitute for, or combined with, filter paper in the preparation of nutrient agar-agar.

(H) APPARATUS FOR EMPLOYMENT OF NUTRIENT JELLY
IN TEST-TUBE AND PLATE-CULTIVATIONS.

Wire Cages.—These cages or crates are used for containing test-tubes, especially when they are

* Hueppe, *Die Methoden der Bakterien Forschung*. 1885.

to be sterilised in the hot-air steriliser; or for lowering tubes of nutrient jelly into the steam steriliser, etc. (Fig. 5).




FIG. 5.—WIRE CAGE FOR TEST-TUBES.

Test-tube Stands.—The ordinary wooden pattern, or the metallic folding stands, are called into use for holding cultivations. Pegged racks are also recommended for draining test-tubes

after washing.

Caoutchouc Caps.—These are caps for fitting over the cotton-wool plugs, and may be used in different sizes for test-tubes and stock-flasks.

Platinum Needles.—A platinum needle for inoculating nutrient media, examining cultivations,




FIG. 6.—PLATINUM NEEDLES; STRAIGHT, HOOKED, LOOPED.

etc., consists of two or three inches of platinum wire, fixed to the end of a glass rod. Several of these needles should be made, with platinum wire of various thicknesses. A piece of glass rod, about seven inches long, is heated at the extreme point in the flame of a Bunsen burner, and a piece

of platinum wire, held near one extremity with forceps, is then fused into the end of the rod. Some needles should be perfectly straight, and kept especially for inoculating test-tubes of




FIG. 7.—DAMP CHAMBER FOR PLATE-CULTIVATIONS.

nutrient jelly. For other purposes the needles may be bent at the extremity into a small hook, and others provided with a loop (Fig. 6).

Tripod Levelling-stand.—A triangular wooden




FIG. 8.—APPARATUS EMPLOYED FOR PLATE-CULTIVATIONS.

Tripod Stand; Glass Dish, filled with cold or iced water; Sheet of Plate-glass; Spirit Level, and Glass Bell.

frame supported upon three screw-feet which enable it to be raised or lowered to adjust the level.

Large Glass Plate.—A piece of plate-glass, or a pane of ordinary window glass, about a foot square.

Spirit Level.

Glass Bells and Dishes.—Shallow glass bells

and dishes, for making a dozen or more damp chambers (Fig. 7), and for completing the apparatus for pouring out liquefied nutrient jelly on glass plates or slides (Fig. 8).

Iron Box.—A box of sheet-iron (Fig. 9) for containing glass plates during their sterilisation in the hot-air steriliser, and for storing them until required for use.






FIG. 9.
BOX FOR GLASS
PLATES.

Glass Plates.—Small panes of glass, about six inches by four. Not less than three dozen are required for a dozen damp chambers.

Glass Benches.—These are necessary for arranging the glass plates or slides in tiers in the damp chambers (Fig. 10). Metal shelves may be substituted for them, but the former are to be preferred.



FIGS. 10, 11.—GLASS BENCHES FOR GLASS PLATES OR SLIDES.

They can be easily made, in any number required, by cementing a little piece of plate-glass at either end of a glass slip (Fig. 11).

Glass Rods.—One dozen or more glass rods, twelve to eighteen inches in length. They are employed for smoothly spreading out the liquefied nutrient gelatine or agar-agar on the glass plates, etc.

Thermometers.—Two or three centigrade thermometers.

(I) APPARATUS FOR PREPARATION OF POTATO-CULTIVATIONS.

Israel's Case.—Sterilising instruments in the flame of a Bunsen burner is most destructive. It is better, therefore, to have a sheet-iron case (Fig. 12) to contain potato-knives, scalpels, and other instruments, and to




FIG. 12.—ISRAEL'S CASE.

sterilise them by placing the case in the hot-air steriliser for an hour at 150° C. The box can be opened at the side, and each instrument withdrawn with a pair of sterilised forceps when required for use.

Glass Dishes.—Several shallow glass dishes are required for preparing damp chambers for potato-cultivations (Fig. 13). The upper, being the larger, fits over the lower, and having no handle, admits of these damp chambers being placed, if necessary,




FIG. 13.

DAMP CHAMBER FOR POTATO-CULTIVATION.

in the incubator in tiers. The large size may also be used in the same way for plate-cultivations.

Potato Knives.—A common broad smooth-

bladed knife set in a wooden handle is sold for this purpose.

Scalpels.—Half a dozen scalpels, preferably with metal handles, may be kept especially for inoculating sterilised potatoes.

Brush.—A common stout nail-brush, or small scrubbing brush, is essential for cleansing potatoes.

(J) APPARATUS FOR PREPARATION OF SOLIDIFIED
STERILE BLOOD SERUM.

Glass Jar.—A tall cylindrical glass jar, on foot, with a broad ground stopper, for receiving blood.

Pipette.—An ordinary or graduated pipette for transferring the serum from the jars to sterile test-tubes or glass capsules.




FIG. 14.—SERUM STERILISER.

Serum Steriliser.—A cylindrical case, with double walls forming an interspace to contain water, closed with a lid, also double walled and provided with a tubular prolongation of the enclosed water chamber (Fig. 14). The water in the cylinder is heated

from below, and that in the lid by means of the prolongation.

In the centre of the cylinder is a column which communicates with the water chamber of the cylinder, and from it pass four partitions, which serve to support the test-tubes.

In the lid are three openings, one of which communicates with the water chamber in the lid by which the latter is filled, and into which a thermometer is then fixed. In the centre an opening admits a thermometer, which passes into the central pipe of the cylinder; through a third opening a thermometer passes to the cavity of the cylinder. The cylinder and cover are jacketed with felt, and the apparatus is supported on iron legs.

Serum Inspissator.—A shallow tin case with glass cover, both case and cover jacketed with felt (Fig. 15). The case is double walled, and the water contained in the interspace is heated from below. It is supported on four legs, and the two front ones move in grooves in the




FIG. 15.—SERUM INSPISSATOR.

case, so that the latter can be placed obliquely at the angle required, and secured in position by screw-clamps. It is employed for coagulating sterile liquid serum, and for solidifying nutrient agar-agar so as to give them a sloping surface.

Glass Capsules.—Small capsules or hollowed-out cubes of crystal glass are employed for cultivations on solid blood serum, on nutrient gelatine, and on agar-agar. They may be procured of white and blackened glass, and are provided with glass slips as covers.

(K) APPARATUS FOR STORING, AND FOR CULTIVATIONS IN, LIQUID MEDIA.

Lister's Flasks.—Lister devised a globe-shaped flask with two necks; a vertical and a lateral one. The lateral one is a bent spout, tapering towards its constricted extremity. When the vessel is restored to the erect position after pouring out some of its contents, a drop of liquid remains behind in the end of the nozzle, and prevents the regurgitation of air through the spout. A cap of cotton wool is tied over the orifice, and the residue in the flask kept for future use. The vertical neck of the flask is plugged with sterilised cotton wool in the ordinary way.

Sternberg's Bulbs.—Sternberg advocates the use of a glass bulb, provided with a slender neck drawn out to a fine point and hermetically sealed.*

Aitken's Test-tube.—This is an ingenious device for counteracting the danger of entrance of atmospheric germs on removal from the ordinary

* Magnin and Sternberg, *Bacteria*. 1884.

test-tube of the cotton-wool plug. Each test-tube is provided with a lateral arm tapering to a fine point, which is hermetically sealed.

Drop culture Slides.—About a dozen or more thick glass slides with a circular excavation in the centre are required for drop-cultures.

Vaseline.—A small pot of vaseline with a camel's-hair brush should be reserved especially for use in the preparation of drop-cultures.

Bulbed Tubes.—Glass vessels such as test-tubes, flasks, and pipettes, which are used in dealing with liquid media, have already been mentioned under other headings; but bulbed tubes, Pasteur's bulbs, and various other forms are also required for special experiments.

(L) APPARATUS FOR INCUBATION.

There are several forms of incubator, each of which has its advocates. They are mostly rectangular chests, with glass walls, front and back, or in front only. A cylindrical model is preferred by some. Two only will be described here—D'Arsonval's and Babes'. The former admits of very exact regulation of temperature, and the latter is a very practical form for general use.

D'Arsonval's Incubator.—The "*Étuve D'Arsonval*" (Fig. 16) is a very efficient apparatus, and is provided with a heat regulator, which enables the temperature to be maintained with a minimum varia-

tion. It consists of a cylindrical copper vessel, with double walls, enclosing a wide interspace for containing a large volume of water. The roof of the




FIG. 16.—D'ARSONVAL'S INCUBATOR.

water-chamber is oblique, so that the wall rises higher on one side than on the other. This admits of the interspace being completely filled with water. At the highest point is an opening fitted with a perforated caoutchouc stopper, through which a glass tube passes. The mouth of the cylinder itself is horizontal, and is closed by a lid, which is also double-walled to contain water. In the lid are four openings; one serves for filling its water-chamber, and the others for thermometers and for regulating the air supply in the cavity of the cylinder. The cylinder is continued below by a cone, also double-walled, and there is a perforated grating at the line of junction of the cylinder and cone. The cone

terminates in a projecting tube provided with an adjustable ventilator. The apparatus is fixed on three supports united to one another below. One of them is utilised for adjusting the height of the heating apparatus. Situated above this leg is the heat-regulating apparatus (Fig. 17), attached to a circular, lipped aperture in the outer wall of the incubator.

To the lip is fixed with six screws the corresponding lip of a brass box, with a tightly stretched diaphragm of india-rubber intervening. Thus the diaphragm separates the cavity of the box from the water in the interspace of




FIG. 17.—SCHLOSING'S MEMBRANE REGULATOR.

the incubator. The cap of the box, which screws on, is bored in the centre for the screw-pipe, by which the gas is supplied. Another pipe entering the box from below is connected with the gas-burners. Around the end of the screw-pipe a collar loosely fits, and is pressed against the diaphragm by means of a spiral wire spring. Close to the mouth of the screw-pipe a small opening exists, so that the gas supply to the burners is not entirely cut off even when the diaphragm completely occludes the mouth of the screw-pipe.

To work the apparatus the tube and plug must be removed, and the water-chamber filled completely with distilled or rain water at the temperature required. The

caoutchouc plug is replaced and the tube placed in position. Gas enters through *d* (Fig. 17), and passes through the opening at its extremity into the chamber of the box. Thence it passes through the vertical exit which is connected with the gas-burners. As the temperature rises the water rises in the tube, and at the same time exercises a pressure on every part of the walls of the incubator, and hence on the diaphragm. In consequence of this, the diaphragm bulging outwards approaches the end of the tube *d*, and gradually diminishes the gas supply. As a result the temperature falls, the water contracts and sinks in the tube, and the diaphragm receding from *d*, the gas supply is again increased. By adjusting the position of the tube *d* to the diaphragm, any required temperature within the limits of the working of the apparatus can be regulated to the tenth of a degree; *provided*, (1) that the gas supply is




FIG. 18.
GAS-BURNER PROTECTED
WITH MICA CYLINDER.

rendered independent of fluctuations of pressure, by means of a gas-pressure regulator, (2) that the height of the water in the tube is controlled daily by the withdrawal or addition of a few drops of distilled water, and (3) that the apparatus is kept in a place with as even a temperature as possible, and sheltered from currents of air.

The burners in Fig. 16 are protected with mica cylinders similar to the burner represented in Fig. 18.

The flames of these burners can be turned down to the smallest length without danger of extinction, and the temperature may be regulated very satisfactorily without using the heat regulator just described, if the gas first passes through a pressure regulator (Fig. 19). To provide against the danger resulting from accidental extinction of

the gas, Professor Koch has devised a self-acting apparatus (Fig. 20), which, simultaneously with the extinction of the flame of the burner, shuts off the supply of gas.

Babes' Incubator.—The pattern of Dr. Babes is very simple, and is recommended by the author in preference to all others (Fig. 21).




FIG. 19.
MOITESSIER'S GAS-PRESSURE REGULATOR.

It consists of a double-walled chest with sides and




FIG. 20.—Koch's SAFETY BURNER.

roof jacketed with felt. Water fills the interspace

between the walls, and on the roof are two apertures—one for a gas regulator, and the other for a thermometer. In front, the chest is closed in by a sheet of felt, a glass door, and a sliding glass panel. The apparatus can be suspended on the wall or supported on legs, and is heated from below by means of protected burners.




FIG. 21.—BABES' INCUBATOR.

The gas should pass first through a pressure regulator, and then through a thermo-regulator to the burners.

Moitessier's Gas-pressure Regulator.—This apparatus is best explained by reference to the diagram (Fig. 19). In the bottom of the cylinder A are the entrance (*k*) and exit (*l*) gas tubes. The tap (*m*) regulates the size of the flame. The cover (*nn*) roofs in the cylinder A. The bell (B) supports by means of *e* and *f* the ball valve *d*, which lies in the cover *cc*. The gas, entering by *k*, passes through the valve *d*, and is thence conducted by the tube *a* to the tube *l*. The bell B and the weighted dish *h* are screwed on to the connecting rod *g*. To diminish as much as possible the friction of *g* in *i*, *g* only touches *i* by three projecting ridges.

Section of *i* and *g* is shown at *s*. To put the apparatus in use it is first levelled, then *h* is screwed off and the cover *nn* removed. A mixture of two parts of pure acid-free glycerine to one of distilled water is poured into the cylinder until it flows out at *g*, which is then closed, and the cover *nn* replaced. The manometers are filled with coloured water, and *k* and *l* connected with the entrance and exit gas tubing respectively. The pressure of the incoming gas raises the bell *B*; and with it the valve *d* is raised towards the opening at *cc*. The weight *h*, which is replaced on *g*, by its downward pressure counteracts this upward pressure of the gas and opens the valve *cc*. Thus the flame is best regulated in the morning, when the pressure is at a minimum; then supposing an increase of pressure occurs, the weight of *h* is overbalanced, *B* is raised and with it *d*, and the gas supply proportionately diminished by the gradual closing of the valved opening.

Reichert's Thermo-regulator.—This regulator (Fig. 22) consists of three parts—a hollow T piece, a stem, and a bulb. The T piece fits like a stopper in the upper widened portion of the stem. One arm of the T is open, and connected with the gas supply; the vertical portion terminates in a small orifice, and is also provided with a minute lateral opening. The stem is provided with a lateral arm, and this arm, the stem, and the bulb contain mercury. The regulator is fixed in the

roof of the incubator, so that the bulb projects either into the interior of the incubator or into the water chamber. When the incubator reaches the required temperature, the mercury is forced up by means of the screw in the lateral arm, until it closes the orifice, at the extremity of the vertical portion of the T. The gas which passes through the lateral orifice is sufficient to maintain the apparatus at the required temperature. If the temperature of the incubator falls the mercury contracts, and gas passing



FIG. 22.
REICHERT'S
THERMO-
REGULATOR.

through the terminal orifice of the T, increases the flame of the burner, and the temperature is restored.

Page's Thermo-regulator resembles the above, but instead of the T piece there are two pieces of glass tubing. The outer tubing envelops the upper part of the stem of the regulator, and admits of being raised or lowered. The upper end of this tubing is closed by a cork, which is perforated to admit the narrow glass tubing, which represents the vertical arm of the T passing within the stem of the regulator. This has a terminal and a lateral opening, and is the means of entrance for the gas. This regulator is adjusted by noting when the thermometer indicates the desired temperature, and then pushing down the outer tube until the terminal opening of the inner tube, which is carried down with it, is obstructed by the mercury.

Meyer's Thermo-regulator is represented in Fig. 23. No. I. shows the construction of the regulator; its inner tube terminates in an oblique




FIG. 23.—MEYER'S THERMO-REGULATOR.

opening, and is also provided with a minute lateral aperture, which prevents the complete shutting off of the gas supply. No. II. illustrates the method of introducing the mercury by suction through a filling tube, which is substituted for the

inner tube of the regulator. No. III. represents Fränkel's modification of the same instrument.

(M) INOCULATING AND DISSECTING INSTRUMENTS AND APPARATUS IN COMMON USE.

Mouse Cages.—As mice are the animals most frequently employed for experimental purposes, mouse cages have been especially introduced, consisting simply of a cylindrical glass jar with a weighted wire cover.

Dressing Case.—A small surgical dressing case, with its usual accessories—forceps, knives, small straight and curved scissors, needles, silk, and so forth—will serve for most purposes.

Pravaz' Syringe.—Koch's modification of Pravaz' syringe admits of sterilisation by exposure to 150° C. for a couple of hours.

Special Instruments and Material.—Instruments required for special operations and the materials necessary for strict antiseptic precautions need not be detailed here.*

Dissecting Boards.—Slabs of wood in various sizes, or gutta-percha trays, provided with large-headed pins, are employed for ordinary purposes.

Dissecting Case.—A dissecting case fitted with scalpels, scissors, hooks, etc., should be reserved entirely for post-mortem examinations.

* *Vide* Cheyne, *Antiseptic Surgery*. 1882.

(N) GENERAL LABORATORY REQUISITES.

Siphon Apparatus.—Two half-gallon or gallon glass bottles, with siphons connected with long flexible tubes provided with glass nozzles and pinchcocks (Fig. 24), should be employed for the




FIG. 24.

SIPHON BOTTLE, WITH FLEXIBLE TUBE, GLASS NOZZLE, AND
A MOHR'S PINCHCOCK.

following purposes :—One is used to contain distilled water, with the nozzle hanging down conveniently within reach of the working table ; the other is to contain a solution of corrosive sublimate (1 in 1000), and may be placed so that the nozzle hangs close to the lavatory sink or basin. The former replaces the use of the ordinary wash bottle, in washing off

surplus stain from cover-glasses, etc., and the latter is conveniently placed for disinfection of vessels and hands after cleansing with water. They should be placed on the top of a cupboard, or on a high shelf.

Desiccator.—The desiccator (Fig. 25) consists of a porcelain pan containing concentrated sulphuric acid, and covered over with a bell-glass receiver.




FIG. 25.—DESICCATOR.

The sheet of plate-glass upon which the pan rests is ground upon its upper surface, and the rim of the glass bell is also ground and well greased. In the centre of the pan is a column supporting a circular frame, which is covered with wire gauze. Slices of potatoes, upon which micro-organisms have been cultivated, are rapidly dried by the action of sulphuric acid in confined air. A cultivation of *Micrococcus prodigiosus*, for example,

may be dried in this way, and preserved for subsequent experiments.

Other items commonly in use in a research laboratory cannot be detailed here, and a description of air-pumps, refrigerators, etc., access to which is nevertheless necessary for some special investigations, must be sought for elsewhere.*

* All bacteriological apparatus, as employed by Professor Koch, may be obtained from Dr. Muencke, 58, Louisen Strasse, Berlin. Nearly all the figures of apparatus here given are from blocks, kindly lent to me by Dr. Muencke. Griffin & Son, 22, Garrick Street, Covent Garden, W.C., will make *to order* any bacteriological apparatus required, and from them all glass vessels and chemical apparatus of home manufacture can be obtained. All histological instruments and material, such as microscopes, microtomes, aniline dyes, celloidin, gelatine, agar-agar, etc., are supplied by G. König, Berlin, N.W., 35, Dorotheen Strasse. Chemicals, staining reagents, and ready-prepared nutrient gelatine can also be obtained from Dr. Georg Grübler, Leipzig, 17, Dufour Strasse. Solutions of lithium carmine, picro-lithium carmine (Orth), picro-carmine (Weigert), alum and borax carmine (Grenacher), etc., ready for use, are prepared by Becker & Co., 34, Maiden Lane, Covent Garden, London, W.C. The latter firm also keep in stock bacteriological apparatus and glass ware of the German pattern.

Mr. Baker, of High Holborn, W.C., is recommended for the supply of microscopes and the ordinary objectives by continental makers, and the new apochromatic objectives recently introduced by Zeiss. Objectives made of the new glass are also constructed by Powell and Leland, but though invaluable to the specialist their expense places them beyond the reach of the general student. Messrs. Swift & Son have recently introduced an excellent $\frac{1}{2}$ oil. imm. for five guineas, and are prepared to supply a microscope completely equipped for bacteriological work at a very low price.

CHAPTER II.

MICROSCOPICAL EXAMINATION OF BACTERIA IN LIQUIDS, IN CULTIVATIONS ON SOLID MEDIA, AND IN TISSUES.

Preliminary Remarks.—In conducting bacteriological researches, the importance of absolute cleanliness cannot be too strongly insisted upon. All instruments, glass vessels, slides, and cover-glasses should be thoroughly cleansed before use. A wide-mouthed glass jar should always be close at hand, containing refuse alcohol for the reception of rejected slide preparations or dirty cover-glasses. When required again for use, slides can be easily wiped clean with a soft rag. Cover-glasses require further treatment, for, unless they are perfectly clean, it is difficult to avoid the presence of air bubbles when mounting specimens. They should be left in strong acid (hydrochloric, sulphuric, or nitric) for some hours; they are then washed, first with water and then with alcohol, and carefully wiped with a soft rag. The same principle applies in the preparation and employments of culture media; any laxity in the processes of sterilisation, or insufficient attention to minute technical details,

will surely be followed with disappointing results in the contamination of the cultures, resulting in the loss of much time. When using platinum needles, either for inoculating fresh tubes in carrying on a series of pure cultures, or in transferring a small portion of a cultivation to a cover-glass for examination under the microscope, the careful sterilisation of the needle by heating the platinum wire till it is white hot in every part, and heating also as much of the glass rod as is made to enter the test-tube, must be carried out with scrupulous care. Indeed it is a good plan to let it become a *force of habit* to sterilise the needle before and after use on every occasion, whatever may be the purposes for which it is employed.

(A) EXAMINATION IN THE FRESH STATE.

Liquids containing micro-organisms, such as pus, blood, juices, culture-fluids, can be investigated by transferring a drop with a looped platinum needle or a capillary pipette to a slide, covering it with a clean cover-glass, and examining without further treatment. If it is desirable to keep the specimen under prolonged observation, a drop of sterilised water or salt solution must be run in at the margin of the cover-glass to counteract the tendency to dry. *Cultures on solid media* can be examined by transferring a small portion with a sterilised needle to a drop of sterilised water on a slide, thinning

it out, and covering with a cover-glass as already described. A more satisfactory method, by which one can keep micro-organisms under observation and study their movements, spore-formation, etc., will be described under "Drop-cultures." *Tissues in the fresh state* may be teased out with needles in sterilised salt solution, and pressed out into a sufficiently thin layer between the slide and cover-glass. Glycerine may in many cases be substituted for salt solution, especially for the examination of micro-organisms such as *Actinomyces*, *Aspergilli*, etc.

There is, as a rule, no difficulty in recognising the larger micro-organisms such as those just mentioned; but where we have to deal with very small bacilli, bacteria, and micrococci, they may possibly be mistaken for granular detritus or fat-crystals, or *vice versâ*. They are distinguished by the fact that fatty and albuminous granules are altered or dispersed by acetic acid, and changed by solution of potash; alcohol, chloroform, and ether dissolve out fat-crystals or fatty particles; on the other hand, micro-organisms remain unaffected by these reagents. This micro-chemical reaction is made the basis of Baumgarten's method.

(B) COVER-GLASS PREPARATIONS.

The method next to be described is the most commonly employed ; in addition to its value as a means of examining liquids, etc., it affords the additional advantage of enabling one to make, if necessary, a large number of preparations which when dried can be preserved, stained or unstained, in ordinary cover-glass boxes ; they are then in a convenient form for transport, and can be mounted permanently at leisure.

The method is as follows :—A cover-glass is smeared with the cut surface of an organ, or pathological growth, or with sputum ; or a drop of blood, pus, or culture-fluid is conveyed to it with a looped platinum needle. By means of another cover-glass, the juice, or fluid, is squeezed out between them into a thin layer, and on sliding them apart each cover-glass bears on one side a thin film of the material to be examined. They are then placed with the prepared side upwards and allowed to dry. After a few minutes, they are held with a pair of flat-bladed or spring forceps, with the prepared side uppermost, and passed rapidly three times through the flame of a spirit lamp or Bunsen burner. To stain them, put two or three drops of an aqueous solution of fuchsine or methyl violet over the film, and after a minute or two wash off the surplus stain with distilled water by means of the siphon apparatus or a

wash-bottle. Turn the cover-glass on to a slide, remove excess of water with filter-paper, and wipe the exposed surface; examine with a power of about 250 diams., and if a higher power be required, which is usually the case, place a droplet of cedar oil on the cover-glass, and examine with an immersion lens.

If the specimen is to be made permanent, fix the cover-glass at one corner with the thumb, and with a soft rag carefully wipe off the cedar oil; then float off the cover-glass by running in distilled water at its margin, and having made a little ledge with a strip of filter-paper, place the cover-glass up against it upon one of its edges and leave it to dry. When perfectly dry mount in Canada balsam, or put it away in a cover-glass box provided with a label of contents.

A minute portion of a culture may be stained and examined in the same way after spreading it out with a hooked platinum needle into a thin film, with or without the addition of a droplet of sterilised water.

In many cases it is necessary or preferable to apply the stain for a much longer period. This is effected by pouring some of the staining solution into a watch-glass, and allowing the cover-glasses to swim on the surface, with their prepared side, of course, downwards. Throughout all these manipulations it is necessary to bear in mind which is the prepared surface of the cover-glass.

Double Coloration of Cover-glass Preparations can also be obtained as in Ehrlich's method for staining tubercular sputum, or by staining with eosin after treatment by the method of Gram.

Ehrlich's Method is as follows:—Five parts of aniline oil are shaken up with one hundred parts of distilled water, and the emulsion filtered through moistened filter-paper. A saturated alcoholic solution of fuchsine, methyl violet, or gentian violet, is added to the filtrate in a watch-glass drop by drop until precipitation commences. Cover-glass preparations are floated in this mixture for fifteen minutes to half an hour, then washed for a few seconds in diluted nitric acid (one part nitric acid to two of water), and then rinsed in distilled water. The stain is removed from everything except the bacilli; but the ground substance can be after-stained, brown if the bacilli are violet, or blue if they have been stained red (Plate III., Fig. 1).

Double staining with eosin after the method of Gram is described under tissue staining. The cover-glass preparations are treated by the same processes as employed with sections; superfluous oil of cloves can be removed by gently pressing the cover-glass between double layers of filter-paper.

Babes' Method affords a very rapid means of examining cultivations, etc. A little of the growth, removed by means of a sterilised platinum hook or small loop, is spread out on a cover-glass into as thin a film as possible: when almost dry, a drop or

two of a weak aqueous solution of methyl violet is allowed to fall from a pipette upon the film. The cover-glass with the drop of stain is, after a minute, carefully turned over on to a slide, and the excess of stain gently and gradually removed by pressure with a strip of filter-paper. It affords a rapid means of demonstration—for example, of such a cultivation as Koch's comma bacilli in nutrient gelatine—enabling the microbes to be seen in some parts of the preparation both stained and in active movement.

His' Method.—The staining of fresh preparations, especially those with no coagulable albumen to fix them, may be also carried out by His' method. A slide is prepared as already described in the examination of micro-organisms in the fresh state. The reagents are then applied by placing them with a pipette drop by drop at one margin of the cover-glass, and causing them to flow through the preparation by means of a strip of filter-paper placed at the opposite margin.

To stain Spores, the method described on p. 65 is somewhat modified. The cover-glass preparations may be either passed as many as twelve times through the flame, or heated to a temperature of 210° for half an hour, or exposed to the action of strong sulphuric acid for a few seconds, and then stained with a watery solution of the dye.

To double-stain Spore-bearing Bacilli.—The cover-glass preparations may be floated for twenty

minutes on a fuchsine aniline-water solution, as used in Ehrlich's method, which has been heated to boiling-point. The fuchsine is removed from the bacilli either by simply rinsing in water, in alcohol, or in weak acid, according to the species, and then the preparations are floated for a few minutes on solution of methylene blue, rinsed in water, dried, and mounted.

To stain Flagella.—Koch recommends floating the cover-glasses on a concentrated watery solution of hæmatoxylin. From this they are transferred to a 5 per cent. solution of chromic acid or to Müller's fluid, by which the flagella obtain a brownish-black coloration. The author has succeeded in demonstrating and photographing flagella, by staining with a drop of a saturated solution of gentian violet in absolute alcohol. Before the alcohol has time to evaporate, the cover-glass is rinsed in water, and then allowed to dry, and finally mounted in balsam. A very intense staining of the whole preparation results.

(c) COVER-GLASS IMPRESSIONS.

One of the most instructive methods for examining micro-organisms is to make an *impression-preparation*. This enables us, in many cases, to study the relative position of individual micro-organisms one to another in their growth on solid cultivating media, and in some cases produces the

most exquisite preparations for the microscope. A perfectly clean, usually small-sized, cover-glass is carefully deposited on a plate- or potato-culture, and gently and evenly pressed down. One edge is then levered up carefully, with a needle, and the cover-glass lifted off by means of forceps. It is then allowed to dry, passed through the flame three times, and stained as already described. In the case of plate-cultures, especially where no liquefaction has taken place, the growth is bodily transferred to the cover-glass, and a vacant area left on the gelatine or agar-agar, corresponding exactly with the form and size of the cover-glass employed (Plate XXVII., Figs. 1 and 2).

CHAPTER III.

PREPARATION AND STAINING OF TISSUE SECTIONS.

(A) METHODS OF HARDENING AND DECALCIFING PREPARATIONS.

To harden small organs, such as the viscera of a mouse, they must be placed on a piece of filter-paper at the bottom of a small, wide-mouthed glass jar, and covered with about twenty times their volume of absolute alcohol. Larger organs, pathological growths, etc., are treated in the same way, but must first be cut into small pieces, or cubes, varying from a quarter of an inch to an inch in size. Müller's fluid may also be employed, and methylated spirit may be substituted for alcohol, from motives of economy. Tissues hardened in absolute alcohol are ready for cutting in two or three days, and those hardened in Müller's fluid in as many weeks.

Teeth, or osseous structures, must first be placed in a decalcifying solution, such as Kleinenberg's. When sufficiently softened, they are allowed to soak

in water, to wash out the picric acid, and then transferred through weak spirit to absolute alcohol. Ebner's solution also gives excellent results, especially when the structures to be decalcified are placed in fresh solution from time to time.

(B) METHODS OF EMBEDDING, FIXING, AND CUTTING.

Material to be cut with the freezing microtome, if hardened in spirit, must be well soaked in water before being frozen; if hardened in Müller's fluid, it can be frozen at once.

If Williams' microtome is employed, the hardened tissues must first be well soaked in gum mucilage, then frozen, and cut.

For cutting with Jung's microtome, the tissues are embedded in paraffine, or celloidin, and mounted on cork; or, if firm enough, they may be fixed upon cork without any embedding material at all. Paraffine, dissolved in chloroform, will be found very serviceable as an embedding material, but celloidin is more commonly employed now. The pieces of tissue to be embedded are placed, after the process of hardening is completed, in a mixture of ether and alcohol for an hour or more. They are then transferred to a solution of celloidin in equal parts of ether and alcohol, and left there, usually for several hours. Meanwhile, corks ready cut for the clamp of the microtome are smeared

over with the solution of celloidin ; this is applied with a glass rod to the surface which is to receive the piece of tissue. The corks are then set aside for the film of celloidin to harden. The pieces of tissue are allowed to remain in the celloidin solution for from one to twenty-four hours, the time varying according to the structure of the specimen. Better results are obtained in the case of lung, or degenerated broken-down tissue, if left for a much longer time than is found to be sufficient for firmer structures. The specimen, when ready, is removed from the celloidin solution with forceps and placed upon a prepared cork. A little of the solution, which is of syrupy consistence, is allowed to fall on the piece of tissue to cover it completely, and the mounted specimen is finally placed in 60 to 80 per cent. alcohol to harden the celloidin. The specimen will be ready for cutting next day.

The specimen may be more neatly embedded by fixing it with a pin in a small paper tray, pouring the celloidin solution over it, and then placing the tray in alcohol to harden the celloidin. The embedded specimen is then fixed on a cork, which has been cut for the clamp of the microtome. The celloidin in the section disappears in the process of clearing with clove-oil.

Material infiltrated with paraffine must be cut perfectly dry, and the sections prevented from rolling up by gentle manipulation with a camel's-hair brush. They must then be picked off the blade of

the knife with a clean needle, and dropped into a watch-glass containing xylol. This dissolves out the paraffine; the sections are then transferred to alcohol to get rid of the xylol, and then to the staining solution.

In the case of specimens embedded in celloidin, or mounted directly on a cork, the tissue, as well as the blade of the knife, should be kept constantly bathed with alcohol, and the sections transferred from the blade with a camel's-hair brush, and floated in alcohol.

For fixing small organs and pieces of firm tissue directly on cork, such as the kidneys of a mouse, or liver, we may employ gelatine, or glycerine gelatine, liquefied over a Bunsen burner in a porcelain capsule. The cork, with specimen affixed, is placed in alcohol, and is ready for cutting sections next day.

The advantage of glycerine gelatine consists in that it may be used for fixing irregular pieces of tissue, as it does not become of a consistency that would injure the edge of the knife.

Embedding in Celloidin and Freezing.—This method gives excellent results. Celloidin is used as already described. The piece of tissue is then placed in a glass capsule, and some of the celloidin solution poured over it. After hardening in spirit, the tissue is cut out and put in water until it sinks. It is then transferred to gum, and frozen and cut in the ordinary way.

(C) GENERAL PRINCIPLES OF STAINING BACTERIA
IN TISSUE SECTIONS: METHODS OF WEIGERT,
GRAM, AND WEIGERT-EHRLICH.

Sections of fresh tissues made with the freezing microtome are to be floated and well spread out in .8 per cent. salt solution, and then carefully transferred, well spread out on the copper lifter, to a watch-glass containing absolute alcohol. Similarly, sections selected from those cut with Jung's microtome may be transferred from the spirit to absolute alcohol. The sections may be then stained by any of the methods to be described.

It is often advisable to employ some method which will enable one to study the structure of the tissue itself. In the same way with sections, however prepared, one should always examine with a low power (one inch) first; this enables one to recognise the tissue under examination in most cases, and even to examine in many cases the topographical distribution of masses of bacteria. With a power of about 250 diams. (one sixth), very many bacteria can be distinguished; and with the oil immersion lenses the minutest bacilli and micrococci can be recognised, and the exact form of individual bacteria accurately determined. As most good modern instruments, like Zeiss' microscopes, are provided with a triple nose-piece, there

is no loss of time in examining a preparation successively with these different powers.

Weigert's Method.—A very useful method for staining both the tissue and the bacteria is as follows :—Place the sections for from six to eighteen hours in a 1 per cent. watery solution of any of the basic-aniline dyes (methyl violet, gentian violet, fuchsine, Bismarck brown). To hasten the process, place the capsule containing the solution in the incubator, or heat it to 45° C. A stronger solution may also be employed, in which case the sections are far more rapidly stained, and are easily overstained. In the latter case they must be treated with a half-saturated solution of carbonate of potash. In either case the sections are next washed with distilled water, and passed through 60 per cent. alcohol into absolute alcohol. When almost decolorised, spread out the section carefully on a copper lifter and transfer it to clove-oil, or stain with picro-carmin solution (Weigert's) for half an hour, wash in water, alcohol, and then treat with clove-oil. After the final treatment with clove-oil, transfer with the copper lifter to a clean glass slide. Dry the preparation by pressure with a piece of filter-paper folded four times, and preserve in Canada balsam dissolved in xylol.

Gram's Method.—In the method of Gram the sections are stained for three minutes in aniline-gentian-violet solution. This is prepared by shaking up one ccm. of pure aniline with twenty-four




Fig 1.





Fig 2.



parts of water, and filtering the emulsion. Half a gramme of the best finely powdered gentian violet is dissolved in the clear filtrate, and the solution filtered before use. The sections are then transferred to a solution of iodine in iodide of potassium till they become dark brown in colour, and then decolorised in absolute alcohol. The time required for complete decolorisation in alcohol varies from a few minutes to twenty-four hours. They are then treated with clove-oil and mounted in Canada balsam. It is much better, however, to employ the aniline-gentian-violet solution quite freshly prepared, and the following useful method is invariably used by the author:—Place four or five drops of pure aniline in a test-tube, fill it three-quarters full with distilled water, close the mouth of the tube with the thumb and shake it up thoroughly. Filter the emulsion twice, and pour the filtrate into a watch-glass or glass capsule. To the perfectly clear aniline water thus obtained add drop by drop a concentrated alcoholic solution of gentian violet till precipitation commences. Stain sections in this solution from ten minutes to half an hour, then transfer to iodine-potassic-iodide solution, and decolorise in alcohol. The process of decolorisation may be hastened by placing the section in clove-oil and returning it to alcohol, and again to clove-oil. If examined, after it has been finally treated with clove-oil and mounted in Canada balsam, the tissue appears colourless or

tinged faintly yellow, while micro-organisms, *e.g.*, bacilli and micrococci, are stained blue or blue-black (Plate II.). Double staining is obtained by transferring the sections after decolorisation to a solution of eosin, Bismarck brown, or vesuvin, again rinsing in alcohol, clearing in clove-oil, and mounting in balsam. Another instructive method is to place the decolorised sections in picro-carminate of ammonia for three or four minutes, and then treat with alcohol, oil of cloves, and balsam. In this way the nuclei are well stained. A somewhat similar result is obtained by placing the sections for a few minutes in Orth's solution (picro-lithium carmine); transferring to acidulated alcohol, then to an alcoholic solution of picric acid, and treating with clove-oil and balsam (Plate XXIV., Fig. 1).

Weigert-Ehrlich Method.—This is a method in which nitric acid is employed as a decolorising agent. It is as follows:—Filtered, saturated, watery solution of aniline, one hundred parts; saturated alcoholic solution of a basic aniline dye (methyl violet, gentian violet, fuchsine), eleven parts; are mixed and filtered. Rapid staining is obtained by warming the solution. Decolorise with nitric acid (1 in 2), and stain with a contrast colour. As delicate sections are apt to be injured by immersion in the nitric acid, they may be transferred from the fuchsine solution to distilled water, then rinsed a few minutes with alcohol, and finally placed in the following contrast stain for one or two hours:—





Fig 1.



Fig 2.



Distilled water	100 cc.
Saturated alcoholic solution of		
methylene blue	20 cc.
Formic acid	10 mm.*

The method of Weigert-Ehrlich may be employed for staining both leprosy and tubercle bacilli (Plate III.). The aniline-fuchsine solution may also be prepared by the simple plan described for Gram's method (p. 77). The more special methods for staining will be given with the description of those species of micro-organisms to which they apply.

* Watson Cheyne, *Practitioner*. 1883.

CHAPTER IV.

PREPARATION OF NUTRIENT MEDIA AND METHODS OF CULTIVATION.

To cultivate micro-organisms artificially, and, in the case of the pathogenic bacteria, to fulfil the second of Koch's postulates, they must be supplied with nutrient material free from pre-existing micro-organisms. Hitherto various kinds of nutrient liquids have been employed, and in many cases they still continue to be used with advantage, but as a general rule they have been in a great measure supplanted by the methods of cultivation on sterile solid media about to be described. The advantages of the latter method are obvious. In the first place, in the case of liquid media, in spite of elaborate precautions and the expenditure of much labour and time, it was almost impossible or extremely difficult to obtain a pure culture. If a drop of liquid containing several kinds of bacteria be introduced into a liquid medium, we have a mixed cultivation from the very first. If in the struggle for existence some bacteria were unable to develop in the presence of others, or a change of temperature and soil allowed

one form to predominate over another, then one might be led to the conclusion that many bacteria were but developmental forms of one and the same micro-organism ; while possibly the contamination of such cultures might lead to the belief in the transformation of a harmless into a pathogenic bacterium. In the case of solid cultivating media, on the other hand, the chance of contamination by gravitation of germs from the air is avoided by the fact that test-tubes, flasks, etc., can be inverted and inoculated from below. The secret of the success of Koch's method, however, depends upon the possibility, in the case of starting with a mixture of micro-organisms, of being able to isolate them completely one from another, and to obtain an absolutely pure growth of each cultivable species. When sterile nutrient gelatine has been liquefied in a tube and inoculated with a mixture of bacteria in such a way that the individual micro-organisms are distributed throughout it, and the liquid is poured out on a plate of glass and allowed to solidify, the individual bacteria, instead of moving about freely as in a liquid medium, are fixed in one spot, where they develop individuals of their own species. In this way colonies are formed each possessing its own characteristic biological and morphological appearances. If an adventitious germ from the air fall upon the culture, it also grows exactly upon the spot upon which it fell, and can be easily recognised as a stranger. To maintain the individuals isolated from one another

during their growth, and free from contamination, it is only necessary to thin out the cultivation, and to protect the plates from the air. The slower growth of the micro-organisms in solid media, affording much greater facility for examining them at various intervals and stages of development, is an additional point in favour of these methods; and the characteristic macroscopical appearances so frequently assumed are, more especially in the case of morphological resemblance or identity, of the greatest importance. The colonies on nutrient gelatine (examined with a low power) of *Bacillus anthracis* and *Proteus mirabilis*, the naked eye appearances in test-tubes of nutrient gelatine of the bacillus of mouse-septicæmia (Figs. 114, 115), and of anthrax (Fig. 107), and the brilliant and curious growth of *Micrococcus indicus* (Koch) upon nutrient agar-agar (Plate VIII., Fig. 1), may be quoted as examples in which the appearances in solid cultivations are pathognomic.

SOLID MEDIA.

(A) PREPARATION OF NUTRIENT GELATINE AND NUTRIENT AGAR-AGAR.

Nutrient Gelatine is prepared as follows:—
Take half a kilogramme of beef (one pound), as free as possible from fat. Chop it up finely, transfer it to a flask or cylindrical vessel, and shake it up well with a litre of distilled water. Place the vessel in

an ice-pail, ice-cupboard, or in winter in a cold cellar, and leave for the night. Next morning commence with the preparation of all requisite apparatus. Thoroughly wash, rinse with alcohol, and allow to dry, about 100 test-tubes. Plug the mouth of the test-tubes with cotton-wool, taking care that the plugs fit firmly but not too tightly. Place them in their wire cages in the hot-air steriliser, to be heated for an hour at a temperature of 150° C. In the same manner cleanse and sterilise several flasks and a small glass funnel. In the meantime the meat infusion must be again well shaken, and the liquid portion separated by filtering and squeezing through a linen cloth. The red juice thus obtained must be brought up to a litre by again transferring it to a large measuring glass and adding distilled water. It is then poured into a sufficiently large and strong beaker, and set aside after the addition of

10 grammes of peptonum siccum.

5 grammes of common salt.

100 grammes of best gelatine.

In about half an hour the gelatine is sufficiently softened, and subsequent gently heating in a water-bath causes it to be completely dissolved. The danger of breaking the beaker may be avoided by placing a cloth several times folded at the bottom of the water-bath.

The next process requires the greatest care and

attention. Some micro-organisms grow best in a slightly acid, others in a neutral or slightly alkaline medium. For example, for the growth and characteristic appearances of the comma bacillus of Asiatic cholera a faintly alkaline soil is absolutely essential. This slightly alkaline medium will be found to answer best for most micro-organisms, and may be obtained as follows :—

With a clean glass rod dipped into the mixture, the reaction upon litmus-paper may be ascertained, and a concentrated solution of carbonate of soda must be added drop by drop, until red litmus-paper becomes faintly blue. If it has been made too alkaline, it can be neutralised by the addition of lactic acid.

Finally, the mixture is heated for an hour in the water-bath. Ten minutes before the boiling is completed, the white of an egg beaten up with the shell is added, and the liquid is then filtered while hot. For the filtration, the hot-water apparatus can be used with advantage, furnished with a filter of Swedish paper made in the following way :—

About eighteen inches square of the best and stoutest filtering paper is first folded in the middle, and then, as in Fig. 26, creased into sixteen folds. The filter is made to fit the glass funnel by gathering up the folds like a fan, and cutting off the superfluous part. The creasing of each fold should be made firmly to within half an inch of the apex of the filter, which part is to be gently

inserted into the tube of the funnel. To avoid bursting the filter at the point, the broth, when poured out from the flask, should be directed against the side of the filter with a glass rod. During filtration the funnel should be covered over with a circular plate of glass, and the process of filtration must be repeated, if necessary, until a pale, straw-coloured, perfectly transparent filtrate results.

The sterilised test-tubes are filled for about a third of their depth by pouring in the gelatine




FIG. 26.—METHOD OF MAKING A FOLDED FILTER.

carefully and steadily, or by employing a small sterilised glass funnel. The object of this care is to prevent the mixture touching the part of the tube with which the plug comes into contact; otherwise, when the gelatine sets, the cotton-wool adheres to the tube and becomes a source of embarrassment in subsequent procedures. As the tubes are filled they are placed in the test-tube basket, and must then be sterilised. They are either lowered into the steam steriliser, when the thermometer indicates 100° C., for twelve minutes

for four or five successive days; or they may be transferred to the test-tube water-bath and heated for an hour a day for three successive days.

If the gelatine show any turbidity after these processes, it must be poured back from the test-tubes into a flask, boiled up for ten minutes, and once more filtered. The processes of sterilisation just described must be repeated.

Nutrient Agar-agar.—Agar-agar has the advantage of remaining solid up to a temperature of about 45° . The preparation of a sterile nutrient jelly is conducted on much the same principles as those already described. Instead, however, of 100 grammes of gelatine, only about 20 grammes of agar-agar are employed (1.5—2 per cent.), and to facilitate its solution it must be allowed to soak in salt-water overnight. For the filtration, flannel is substituted for filter-paper, or may be used in combination with the latter. The hot-water apparatus is invariably employed, unless, to accelerate the process, the glass funnel and receiver are bodily transferred to the steam steriliser. If the conical cap cannot be replaced, cloths laid over the mouth of the steriliser must be employed instead. It may be necessary to repeat the process of filtration, but it must not be expected that such a brilliant transparency can be obtained as with gelatine. The final result, when solid, should be colourless and clear; but if only slightly milky, it may still be employed.

A little liquid gradually collects in the tubes, being expressed by the contraction of the agar-agar.

Glycerine Agar-agar.—This is prepared by adding 5 per cent. of glycerine to the mixture of nutrient agar-agar after the boiling and before the filtration.

After the final treatment in the steam steriliser some of the tubes of gelatine and agar-agar may be placed in the blood-serum apparatus, and left to gelatinise with an oblique surface.

(B) METHODS OF EMPLOYING NUTRIENT JELLY IN
TEST-TUBE- AND PLATE-CULTIVATIONS.

Test-tube-cultivations.—To inoculate test-tubes containing nutrient jelly, the cotton-wool plug must be *twisted* out, by which means any adhesions that may exist are broken down. A sterilised needle charged with blood, pus, etc., containing the micro-organisms, or with a colony from a plate-culture, is steadily thrust once, and once only, into the nutrient jelly. The tube should be held with its mouth downwards, to avoid, as far as possible, accidental contamination from the gravitation of germs in the air; and the plug, which has been removed with the thumb and index finger of the right hand and held between the fourth and fifth fingers of the left, is replaced as rapidly as possible (Fig. 27).

The chances of error arising from contamination of the cultivations are further reduced by avoiding

draughts at the time of inoculation, and it is best that these manipulations should be carried on in a quiet room in which the tables and floor are wiped with damp cloths, rather than in a laboratory in which the air becomes charged with germs through




FIG. 27.
METHOD OF INOCULATING A
TEST-TUBE CONTAINING
STERILE NUTRIENT JELLY.

constant sweeping and dusting, and the entrance and exit of classes of students. In conducting any investigation a dozen or more tubes should be inoculated, and if by chance an adventitious germ, in spite of these precautions, gain an entrance, the contaminated tube can be rejected and the experiments continued with the remaining pure cultivations.

Where, however, one tube is inoculated from another containing a liquid medium, as in the process of preparing plate-cultures, or where a culture is made from a tube in which the growth has liquefied the gelatine, it is obvious that the tubes cannot be inverted, and they must then be held and inoculated as in Fig. 28. To inoculate those tubes of nutrient media which have been gelatinised obliquely, the sterilised needle with the material to be cultivated is streaked over the surface from below upwards.



Fig 1.



Fig 2.




Fig 3.





Fig 1.



Fig 2.



Fig 3.






Fig 1.




Fig 2.



Fig 3.






Fig 1.

Fig 2.

Fig 3.






Fig 1.



Fig 2.



Fig 3.



Fig 1.

Fig 2.

Fig 3.




Fig 1.

Fig 2.

Fig 3.

Examination of Test-tube-cultivations.—

The appearances produced by the growths in test-tubes can be in most cases sufficiently examined with the naked eye. The illustrations in the accompanying plates (Plates IV.—XI., and Plate XVIII., Fig. 1) are given as examples of the various changes produced in the nutrient media. In some cases the jelly is partially or completely liquefied, while in others it remains solid. The growths may be abundant or scanty, coloured or colourless. The nutrient jelly may itself be tinged or stained with products resulting from the growth of the organisms. When liquefaction slowly takes place in the needle track, or the organism grows without producing this change, the appearances which result are often very delicate, and in some cases very characteristic. The appearance of a simple white thread, of a central thread with branching lateral filaments, of a cloudiness, or of a string of beads in the track of the needle, may be given as examples. In such cases much may be learnt by examining the growth with a magnifying glass. Here, however, a difficulty is encountered, for the cylindrical form of the tube so distorts the appearance of its contents, that the examination is rendered somewhat difficult. To obviate this, the following very simple contrivance may be employed with advantage.

Cheshire's Trough.—This consists of a rectangular vessel, four inches in height, two inches in

width, and one inch in depth. It may be easily constructed by cementing together two slips of glass to form the back and front, with three slips of stout glass with ground edges forming the sides and base. The front may be constructed of thin glass, and the base of the vessel made to slope so that the test-tube when placed in the trough has a tendency to be near the front. The trough is filled with a mixture of the same refractive index as the nutrient gelatine. The latter has a refractive index rather higher than water, which is about 1.333; alcohol has a refractive index of 1.374. The trough is filled with water, and alcohol is then added until the proper density is reached. The test-tube is placed in the trough, and held in position by means of a clip. The trough can be fixed on the inclined stage of the microscope, and the contents of the tube conveniently examined with low-power objectives.

PLATE-CULTIVATIONS.

The key to the success of Koch's method of cultivation on solid media consists in the employment of plate-cultivations. By this means, as has already been mentioned (p. 81), a mixture of bacteria, whether it be in fluids, excreta, or in artificial cultivations, can be so treated that the different species are isolated one from the other, and perfectly pure cultivations of each of the cultivable bacteria in the original mixture established in various nutrient media. We are enabled

also to examine under a low power of the microscope the individual colonies of bacteria, and to distinguish, by their characteristic appearances, micro-organisms which, in their individual form, closely resemble one another, or are even identical. The same process, with slight modification, is also employed in the examination of air, soil, and water, as will be referred to later.

The preparation of plate-cultivations, therefore, must be described in every detail; and, to take an example, we will suppose that a series of plates are to be prepared from a test-tube-cultivation.

Arrangement of Levelling Apparatus.—In order to spread out the liquid jelly evenly on the surface of a glass plate, and hasten its solidification, it is necessary to place the glass plate upon a level and cool surface. This is obtained in the following manner:—Place a large shallow glass dish upon a tripod stand, and fill it to the brim with cold water; carefully cover the dish with a slab of plate-glass, or a pane of window-glass, and level it by placing the spirit-level in the centre and adjusting the screws of the tripod. Replace the level by a piece of filter-paper, the size of the glass plates to be employed, and cover it with a shallow bell-glass (Fig. 8).

Sterilisation of Glass Plates.—The glass plates are sterilised by filling the iron box (p. 44), and placing it in the hot-air steriliser, at 150° C., from one to two hours. As these plates are used

also for other purposes, a quantity ready sterilised should always be kept in the box.

Preparation of Damp Chambers.—The damp chambers for the reception of the inoculated plates are prepared thus :—Thoroughly cleanse and wash out with (1-1000) sublimate solution a shallow glass dish and bell (Fig. 7). Cut a piece of filter-paper to line the bottom of the glass dish, and moisten it with the same solution.

In a glass beaker or an ordinary glass tumbler, with a pad of cotton-wool at the bottom, place the tube containing the cultivation, the three tubes to be inoculated, and three glass rods which have been sterilised by heating in the flame of a Bunsen burner. Provide yourself with a strip of paper of a large label, a pencil, a pair of forceps, and inoculating needles. All is now ready at hand to commence the inoculation of the tubes.

Method of Inoculating the Test-tubes.—Liquefy the gelatine in the three tubes by placing them in a beaker containing water at 30° C., or by gently warming them in the flame of the Bunsen burner. Keep the tubes, both before and after the inoculation, in the warm water, to maintain the gelatine in a state of liquefaction. Hold the tube containing the cultivation and a tube of the liquefied gelatine (to be called the "original") as nearly horizontal as possible between the thumb and index finger of the left hand. With the finger and thumb of the right hand loosen the plugs of

the tubes. Take the looped platinum needle in the right hand and hold it like a pen. Remove the plug from the culture-tube by using the fourth and fifth fingers of the right hand as forceps, and place it between the fourth and fifth fingers of the left. Remove the plug of the "original" in the same way, placing it between the third and fourth fingers of the left hand (Fig. 28). With the loop take up a droplet



FIG. 28.—METHOD OF INOCULATING TEST-TUBES IN THE PREPARATION OF PLATE-CULTIVATIONS.

of the cultivation and inoculate the "original," twisting the loop several times in the liquid gelatine. Replace the plugs and set aside the cultivation. Hold the freshly inoculated tube between the fore-finger and thumb of either hand, almost horizontally, then raise it to the vertical, so that the liquid gelatine gently flows back. By repeating this motion and rolling the tube between the fingers and thumbs the micro-organisms which have been introduced are

distributed throughout the gelatine. Any violent shaking, and consequent formation of bubbles, must be carefully avoided. From the so-called "original" inoculate in the same manner a fresh tube of liquefied gelatine, introducing into it three droplets with a sterilised loop. This tube is then called the "first attenuation," or No. 1. After treating No. 1 as has been already described in the case of the original, the same process is repeated with a third tube, which is inoculated in the same way from No. 1. This is the "second attenuation," or No. 2; and in some cases a third attenuation" is carried out from No. 2. The last tube must be inoculated in different ways, according to experience, for different micro-organisms. Sometimes a sufficient separation of the micro-organisms is attained by inoculating the last tube with a fine straight needle, dipped from one into the other from three to five times.

The next process consists in pouring out the gelatine on a glass plate and allowing it to solidify.

Preparation of the Gelatine-plates.—The directions to be observed in pouring out the gelatine are as follows :—

Place the box containing sterilised plates horizontally, and so that the cover projects beyond the edge of the table; remove the cover, and withdraw a plate with sterilised forceps; hold it between the finger and thumb by opposite margins, rapidly transfer it to the filter-paper under the bell-glass and

quickly replace the cover of the box. On removing the plug from "the original," an assistant raises the bell-glass, and the contents of the tube are poured on to the plate; with a glass rod the gelatine must be then rapidly spread out in an even layer within about half an inch of the margin of the plate. The assistant replaces the bell-glass, and the gelatine is left to set. Meanwhile a glass bench or metallic shelf is placed in the damp chamber, ready for the reception of the plate-cultivation, and when the gelatine is quite solid the plate is quickly transferred from under the bell-glass to the damp chamber; precisely the same process is repeated with tubes 1 and 2, and the damp chamber, labelled with the details of the experiment, is set aside for the colonies to develop. Not only plate-cultures should be carefully labelled with date and description, but the same remark applies equally to all preparations—tube-cultures, potato-cultures, drop-cultures, etc. In plate-cultivations write the source of the material, the date, and the number of inoculations; for example, thus:—

Finkler's comma-bacilli.

From tube-cultivation on "agar-agar," 5th February, 1885.

Lower plate (Orig.) . . . from cultivation.

Middle plate, No. 1 . . . ,, Orig.

Upper plate, No. 2 . . . ,, No. 1.


Corresponding with the fractional cultivation of the micro-organisms obtained in this manner, the colonies will be found to develop in the course of a day or two, the time varying with the temperature of the room. The lower plate will contain a countless number of colonies which, if the micro-organism liquefies gelatine, speedily commingle, and produce, in a very short time, a complete liquefaction of the whole of the gelatine. On the middle plate, with the first attenuation, the colonies will also be very numerous, but retain their isolated position for a longer time; while on the uppermost plate, the second attenuation, the colonies are completely isolated from one another, with an appreciable surface of gelatine intervening (Plates XII. and XIII).


Examination of Plate-cultivations.—The macroscopical appearances of the colonies are best studied by placing the plate on the slab of blackened glass, or on the porcelain slab if the colonies are coloured.


To examine the microscopical appearances, a selected plate is placed upon the stage of the microscope; it is better to have a larger stage than usual for this purpose. The smallest diaphragm is employed, and the appearances studied principally with a low power. These appearances should be carefully noted, and a rapid sketch of the colony made. The morphological characteristics of the micro-organisms of which the colony is formed can then be examined in the following way:—A small




PLATE 12
faang p 98.









looped platinum needle or one bent at the extremity into a miniature hook, is held like a pen, and the hand steadied by resting the little finger on the stage of the microscope. The extremity of the needle is steadily directed to the space between the lens and the gelatine without touching the latter, until, on looking through the microscope, it can be seen in the field, above or by the side of the colony under examination. The needle is then dipped into the colony, steadily raised, and withdrawn. Without removing the eye from the microscope this manipulation can be seen to be successful by the colony being disorganised or completely removed from the gelatine. It is, however, not easy to be successful at first, but with practice this can be accomplished with rapidity and precision. A cover-glass preparation is then made in the manner already described, viz., by rubbing the extremity of the needle on a perfectly clean cover-glass and examining by Babes' rapid method, or by thinning out the micro-organisms in a droplet of sterilised water previously placed on the cover glass, drying, passing three times through the flame, and staining with a drop of fuchsine.

Inoculations should be made in test-tubes of nutrient gelatine and agar-agar, from the micro-organisms transferred to the cover-glass before it is dried and stained, from any remnants of the colony which was examined, or from other colonies bearing exactly similar appearances. In this way

pure cultivations are established, and the macroscopical appearances of the growth in test-tubes can be studied. The plates should be replaced in the damp chambers as soon as possible; drying of the gelatine, or contamination with micro-organisms gravitating from the air during their exposure, may spoil them for subsequent examination. Nutrient agar-agar can also be employed for the preparation of plate-cultivations, but it is much more difficult to obtain satisfactory results. The test-tubes of nutrient agar-agar must be placed in a beaker with water and heated until the agar-agar is completely liquefied. The gas is then turned down and the temperature of the water allowed to fall until the thermometer stands just above 50° C. The water must be maintained at this temperature, and the test-tubes must be in turn rapidly inoculated and poured out upon the glass plates, as already described. Glass plates may also be employed in a much simpler method. The nutrient jelly is liquefied, poured out, and allowed to set. A needle charged with the material to be inoculated is then streaked in lines over the surface of the jelly. This method is of especial value in inoculating different organisms side by side, and watching the effect of one upon the other, or a micro-organism in this way may be sown upon gelatine which has been already altered by the growth of another micro-organism; the change produced in the gelatine, as in the case of the

fluorescing bacillus, extending far beyond the limits of the growth itself (Plate IX., Fig. 3).

Nutrient jelly may also be spread out on sterilised microscopic slides and inoculated as just described, or cultivations may be made in shallow glass dishes, glass capsules, etc., which must be sterilised on the principles already laid down, and after inoculation placed in damp chambers for the growths to develop.

(C) PREPARATION AND EMPLOYMENT OF STERILISED POTATOES, POTATO-PASTE, BREAD-PASTE, VEGETABLES, FRUIT, AND WHITE OF EGG.

Potato-cultivations.—Sterilised potatoes form an excellent medium for the cultivation of many micro-organisms, more especially the chromogenous species. Potato-cultivations also give in some cases very characteristic appearances, which are of value in distinguishing bacteria which possess morphological resemblances (Plates XIV.—XVII).

Preparation of Sterilised Potatoes.—Potatoes, preferably smooth-skinned, which are free from “eyes” and rotten spots should be selected. If they cannot be obtained without eyes and spots, these must be carefully picked out with the point of a knife with as little destruction of the surface as possible. The potatoes are well scrubbed with a stiff brush and allowed to soak in sublimate solution for half an hour. They are then transferred to the

potato-receiver and steamed in the steam steriliser for twenty minutes to half an hour, varying according to the size of the potatoes. When cooked, the potato-receiver is withdrawn and left to cool, the potatoes being retained in it until required for use.

Damp chambers are prepared ready for their reception, the vessels being cleansed and washed with sublimate as described for plate-cultivations. Small glass dishes of the same pattern as the large ones (Fig. 13) may be employed for single halves of potatoes. Potato-knives and several scalpels which have been sterilised in an Israel's case by heating them in the hot-air steriliser at 150° for one hour must be ready to hand. The potato-knives may also be sterilised by heating them in the flame of a Bunsen burner and placing them on their backs with their blades projecting over the edge of the table. Scalpels may be sterilised in the same way and laid upon a sterilised glass plate and covered with a bell-glass. It must not be forgotten, however, that heating the blades in the flame destroys the temper of the steel, and therefore knives and other instruments should preferably be sterilised in the hot-air steriliser, enclosed in an Israel's case, or simply enveloped in cotton-wool.

Inoculation of Potatoes.—The coat-sleeves should be turned back, and the hands, after a thorough washing with good lathering soap, be dipped in sublimate solution. An assistant opens

the potato-receiver, and a potato is selected and held between the thumb and index finger of the left hand (Fig. 29). With the knife held in the right hand, the potato is almost completely divided in the direction which will give the largest surface. The assistant raises the cover of the damp chamber,




FIG. 29.—METHOD OF DIVIDING POTATOES.

and the potato is introduced, and while withdrawing the knife, allowed to fall apart. The cover is quickly replaced, and another potato treated in the same way is placed in the same damp chamber. The four halves are then quite ready for inoculation. As an extra precaution the left hand is again dipped in sublimate, and one half of a potato is taken up between the tips of the thumb and index finger,

care being taken to avoid touching the cut surface. Holding it with its cut surface vertical, a small portion of the substance to be inoculated is placed on the centre with a sterilised platinum needle. With a sterilised scalpel the inoculated substance is rapidly spread over the surface of the potato with the flat of the blade to within a quarter of an inch of the margin, and the potato is then as quickly as possible replaced in the moist chamber. With another sterilised scalpel a small portion of the potato from the inoculated surface of the first half is in the same way spread over the surface of the second half, this forming as in plate-cultivations a "first attenuation." Exactly the same is repeated with a third potato, and even a fourth, so that a still further attenuation or fractional cultivation of the micro-organisms may be obtained. In some cases it is necessary to place the cultures in the incubator; others grow very well at the temperature of the room. As in plate-cultivations, the potato may also be inoculated by simply streaking it in lines with a needle charged with the material to be inoculated. In that case it is better to place a wedge-shaped slice of raw potato in a sterilised test-tube. A number of these tubes may be prepared ready for use. The tubes are plugged with cotton-wool, and the potato steamed in the steam steriliser.

POTATO-PASTE.

Potato-paste is sometimes employed where it is



Fig 1.



Fig 2.



Fig 1.



Fig 2.



Fig 1.



Fig 2.



Fig 1.



Fig 2.

desirable to obtain an extensive growth of certain bacteria. The potatoes are boiled for an hour, and the floury centre squeezed out of their skins. This is then mashed up with sufficient sterilised water to produce a thick paste, and is heated in the steam steriliser for half an hour for three successive days.

BREAD-PASTE.

Some micro-organisms, more especially mould fungi, grow very well on bread-paste. This is prepared by removing the crust from a stale loaf of bread and allowing it to dry in the oven. It is then broken up, and reduced to a fine powder with a pestle and mortar. Small, carefully cleansed, conical, or globe-shaped flasks are plugged with cotton-wool and sterilised in the oven. When cool a small quantity of the powder is placed in them, and sterilised water added in the proportion of one part for every four of the powder. The paste is sterilised by steaming in the steriliser at 100° C. for half an hour for three successive days. The flasks can be reversed, and are inoculated in the usual way with a platinum needle.

CULTIVATIONS ON VEGETABLES, FRUIT, WHITE OF EGG, ETC.

Boiled carrots and other vegetables, and various kinds of stewed fruit, are also occasionally employed for the cultivation of bacteria. The sterilisation of

these media must be carried out on the principles already explained. White of egg may be poured out on sterilised glass plates, or in shallow glass dishes, boiled in the steam steriliser, and after inoculation placed in a damp chamber.

(D) PREPARATION AND EMPLOYMENT OF STERILE BLOOD SERUM.

The tubercle-bacillus, the bacillus of glanders and many other micro-organisms thrive well when cultivated on solid blood serum. This medium has the additional advantage of remaining solid at all temperatures. The technique required for its preparation and sterilisation is as follows:—Several cylindrical vessels, about 20 cm. high, are thoroughly washed with sublimate solution (1-1000), and then with alcohol, and finally rinsed out with ether. The ether is allowed to evaporate, and the vessels are then ready for use. The skin of the animal selected—calf, sheep, or horse—is washed with sublimate at the seat of operation, and the bleeding is performed with a sterilised knife. The first jet of blood from the vein is rejected, and that which follows is allowed to flow into the vessels until they are almost full. The ground-glass stoppers, greased with vaseline, are replaced, and the vessels set aside in ice, as quickly as possible, for from twenty-four to thirty hours. By that time the separation of the clot is completed, and the clear serum can then be trans-



Fig 1



Fig 2.



Fig 3.



Fig 4.



ferred to plugged sterile test-tubes. These should be filled with a sterilised pipette for about a third of their length, and are then placed in Koch's slow steriliser with the temperature maintained for an hour at 58° C. The same process is repeated for six successive days, the temperature on the last day being gradually raised to 60° . This completes the sterilisation, but to solidify the serum it is necessary to arrange the tubes in the inspissator at the angle required. The temperature of this apparatus is kept between 65° and 68° C. Directly solidification takes place the tubes must be removed, and they should then present the character of being hard, solid, of a pale straw colour, and transparent. A little liquid collects at the lowest point, and the serum is sometimes milky in appearance at its thickest part. The serum may not only be employed in test-tubes, but also in small flasks, glass capsules, or other vessels, all of which must be cleansed and sterilised in the usual way. Hydrocele fluid and other serous effusions may be prepared in the same manner, or gelatine may be added to the serum in the proportion of 5 per cent.

Inoculation of the Tubes.—A small portion of the material to be inoculated is taken up with a sterilised platinum needle, and drawn in lines over the sloping surface of the serum; or a minute piece of tissue, tubercle, etc., may be introduced into the tube and deposited on the surface of the nutrient medium. The precautions that are to be observed

in isolating the material to be inoculated will be referred to later (p. 141).

LIQUID MEDIA.

(E) PREPARATION OF STERILISED BOUILLON, LIQUID BLOOD SERUM, URINE, MILK, VEGETABLE INFUSIONS, AND ARTIFICIAL NOURISHING LIQUIDS.

Nutrient liquids are still largely employed, and by some observers even in preference to the solid media advocated by Koch. It must not be supposed, however, that the methods of cultivation in liquids are discarded entirely by the modern school, for there is no more instructive method than the employment of so-called drop-cultures. For inoculation experiments where the presence of gelatine is undesirable, for studying the physiology and chemistry of bacteria and where for any object a rapid growth of micro-organisms is necessary, the employment of liquid media is not only advisable, but is absolutely necessary. Liquid media comprise two distinct groups—*natural* and *artificial*. The natural group includes meat broths, blood, urine, milk, and vegetable infusions; the artificial are solutions built up from a chemical formula representing the essential food constituents.

NATURAL MEDIA.

Bouillon.—A broth of bouillon of beef, pork, or chicken may be made in the same manner as

described for the preparation of nutrient gelatine, with simply omission of the gelatine. After the neutralisation with carbonate of soda solution drop by drop, the flask of broth is placed in the steam steriliser for half an hour at 100° C. A clear liquid results on filtration, which is transferred to plugged sterilised flasks or test-tubes, and sterilisation effected by exposing them in the steam steriliser for half an hour at 100° C. for two or three successive days.

Liquid Blood Serum.—The preparation of sterile blood serum has already been described. It may be used for cultivation, especially in the form of drop-cultivations, before the final treatment by which it is solidified. Hydrocele fluid, peritonic and pleuritic effusions, can also be employed after sterilisation in the steam steriliser. The fluid should be withdrawn with a sterilised trocar and canula, and received into plugged sterilised flasks.

Urine.—In order to obtain urine free from micro-organisms the following precautions must be observed :—The orifice of the urethra must be thoroughly cleansed with sublimate solution. The first jet of urine should be rejected, and the rest received into sterilised vessels, which must be quickly closed with sterile plugs. If these precautions be not attended to the urine must be rendered sterile by the means described for the sterilisation of bouillon.

Milk.—If milk has been drawn into sterile flasks after thoroughly cleansing and disinfecting the teats

and hands, it may be kept without change. If procured without these precautions, it must be steamed in the steriliser for half an hour for five successive days.

Vegetable Infusions. — Infusions of hay, cucumber, and turnip are used for special purposes, and more rarely decoctions of plums, raisins, malt, and horse-dung. They are mostly prepared by boiling with distilled water, after maceration for several hours. The filtrate is received into sterile flasks and sterilised in the usual way in the steam steriliser.

ARTIFICIAL FLUIDS.

Pasteur's Fluid.—This solution is prepared by mixing the ingredients in the following proportions :—

Distilled water	100
Pure cane sugar	10
Ammonium tartrate	1
Ash of yeast	·075

Cohn-Mayer Fluid.—Mayer's modification of the nourishing fluid employed by Cohn is as follows :—

Distilled water	20
Phosphate of potassium	·1
Sulphate of magnesium	·1
Tribasic calcium phosphate	·01
Ammonium tartrate	·2

- (F) METHODS OF STORING AND EMPLOYING LIQUID MEDIA ; LISTER'S FLASKS, AITKEN'S TEST-TUBES, STERNBERG'S BULBS, PASTEUR'S APPARATUS, MIQUEL'S BULBS ; DROP-CULTURES.

Cultivations in liquid media can be carried on in test-tubes, but it is more satisfactory to employ special forms of flasks, bulbs, U tubes, etc. As test-tubes and flasks containing liquid media cannot be inverted, inoculation with a sterilised needle must be carried out as rapidly as possible, with the additional precaution of closed windows and doors.

Lister's Flasks.—These flasks (p. 48) were especially introduced by Lister as a means of storing liquid nutrient media. They are so constructed that after removal of a portion of the contents, on restoring the vessel to the vertical position, a drop of liquid always remains in the extremity of the nozzle, which prevents the regurgitation of unfiltered air.

Sternberg's Bulbs.—The method of introducing liquid into the bulbs employed by Sternberg, and of sterilising and inoculating it, is as follows:—The bulb is heated slightly over the flame, and the extremity of the neck, after breaking off the sealed point, is plunged beneath the surface of the liquid. As the air cools the liquid is drawn into the bulb, usually filling it to about one-third of its capacity. The neck of the flask is again sealed up, and the

liquid which has been introduced is sterilised by repeatedly boiling the flasks in the water-bath. They should then be placed in the incubator for two or three days; and if the contents remain transparent and free from film, they may be set aside as stock-bulbs, to be used when required.

To inoculate the liquid in the bulb the end of the neck is heated to sterilise the exterior, the bulb is gently warmed, and the extremity of the neck nipped off with a pair of sterilised forceps. The open extremity is plunged into the liquid containing the micro-organism, a minute quantity enters the tube and mingles with the fluid in the bulb, without fear of contamination by atmospheric germs. The extremity of the neck is once more sealed up in the flame of a Bunsen burner.

Aitken's Tubes.—These tubes are plugged and sterilised, and the nutrient medium introduced as into ordinary test-tubes. Instead of withdrawing the cotton-wool plug, they are inoculated by means of the lateral arm. The sealed extremity of the arm is nipped off with sterilised forceps, and the inoculating needle is carefully introduced through the opening thus made. It is directed along the arm until it touches the opposite side of the test-tube, where it deposits the material with which it was charged. The needle is withdrawn, and the end of the lateral arm again sealed up in the flame; the test-tube is then tilted until the liquid touches the deposited material; on restoring the tube to

the vertical the material is washed down into the body of the nutrient liquid.

Pasteur's Apparatus.—Special forms of tubes, bulbs, and pipettes are employed by the school of Pasteur. The tubes are provided with lateral or with curved arms drawn out to a fine point, and with slender necks plugged with cotton-wool. A double form shaped like a tuning-fork, each limb with a bent arm, is a convenient form for storing sterilised bouillon. The sealed end of an arm is nipped off with sterilised forceps, the sterile bouillon aspirated into each limb, and the arm again sealed in the flame; a series of such tubes can be arranged upon a rack on the working table.*

Bulbs with a vertical neck drawn out to a fine point, others with a neck bent at an obtuse angle, plugged with cotton-wool, and a lateral curved arm drawn out to a fine point, are also employed. For a description of these various vessels and their special advantages, the works of Pasteur and Duclaux must be consulted.

Miquel's Bulbs.—The *tube à boule* of Miquel† is also a very useful form. It consists of a bulb of 50 cc. capacity blown in the middle of a glass tube. The part of the tube above the bulb is contracted about half-way between the bulb and its extremity, and can either be left quite straight or

* Duclaux, *Ferments et Maladies*. 1882.

† Miquel, *Les Organismes Vivants de l'Atmosphère*. 1883.

can be made to curve slightly. On either side of the contraction the tube is plugged with asbestos. The portion of the tube below the bulb is S shaped, and drawn out at its extremity into a fine point. The bulbs are charged with nutrient liquid and inoculated by aspiration, and the point of the S tube sealed in the flame of a Bunsen burner.

Drop-cultures. — This method of cultivation has already been referred to as a particularly instructive one. It enables us to study many of the changes which take place during the life history of micro-organisms. This is illustrated, for example, by the anthrax bacillus, where we can watch the gradual growth of a single bacillus into a long filament, and the subsequent development of bright oval spores. It is necessary carefully to observe the minutest details to maintain the cultivation pure. An excavated slide is thoroughly cleaned, and then sterilised by being held with the cupped side downwards in the flame of the Bunsen burner. A ring of vaseline is painted round the excavation, and the slide is then placed under a glass bell. Meanwhile a carefully cleansed cover-glass is also sterilised by passing it through the flame, and should be deposited on the plate of blackened glass. With a sterilised looped needle, a drop of sterile bouillon is transferred to the cover-glass, and this is inoculated by touching it with another sterilised needle charged with the material, without disturbing the form of the drop. It is quite sufficient just to

touch the drop instead of transferring a visible quantity of blood, juice, or growth, as the case may be. The slide is then inverted and placed over the cover-glass, so that the drop will come exactly in the centre of the excavation, and is gently pressed down. On turning the slide over again the cover-glass adheres, and an additional layer of vaseline is painted round the edges of the cover-glass itself. The slide must be labelled, and, if necessary, placed in the incubator, and the results watched from time to time. Instead of bouillon, liquid blood serum may also be employed in this form of cultivation. If it be required to preserve the drop-cultivation as a microscopic preparation, the cover-glass is gently lifted off and allowed to dry. Any vaseline adhering to the cover-glass should be wiped off, and the cover-glass can then be passed through the flame and stained in the usual manner.

Moist Chambers.—Unless drop-cultures are very carefully prepared, they are liable to dry up, if kept for examination for several days. Many therefore prefer employing a moist chamber. There are several different forms in use.

The drop-culture slide may be converted into a moist chamber by having a deep groove cut round the circumference of the concavity. This groove is filled with sterilised water by means of a pipette. A ring of vaseline is painted with the camel's-hair brush outside the groove, and the cover-glass, with the drop-cultivation, is inverted and placed over the

concavity. This form is very useful, as the slide can be easily cleansed and effectually sterilised by holding it in the flame of the Bunsen burner.

A very simple form of moist chamber, which may be used in some cases, but possesses the disadvantage of not admitting of sterilisation by heat, may be constructed as follows*:

A small piece of putty or modelling wax is rolled into a cord about two inches long and $\frac{1}{8}$ inch thick.




FIG. 30.—METHOD OF FORMING A SIMPLE MOIST CHAMBER.

By uniting the ends a ring is formed, which is placed on the middle of a clean glass slide (Fig. 30). A drop of water is placed in the centre of the ring, and the cell roofed in by applying the cover-glass.

A somewhat similar cell in form, which has the advantage of permitting of thorough cleansing, may be constructed by cementing a glass ring with flat surfaces to an ordinary slide. Vaseline is applied with a camel's-hair brush to the upper surface of

* Schäfer's *Course of Practical Histology*. 1877.

the ring, and one or two drops of water placed with a pipette at the bottom of the cell. The cover-glass, with the preparation, is then inverted over the cell and gently pressed down upon the glass ring. The vaseline renders the cell air-tight, and, to a certain extent, fixes the cover-glass to the ring.

Warm Stages.—To apply warmth while a preparation is under continuous observation, we must either place the microscope bodily within an incubator, with the eyepiece protruding through an




FIG. 31.—SIMPLE WARM STAGE.

opening, so that we may observe what is going on without moving the preparation, or we must employ some means of applying heat directly to the preparation.

A simple warm stage may be made of an oblong copper plate, two inches long by one inch wide, from one side of which a rod of the same material projects. The plate has a round aperture in the middle, half an inch in diameter, and is fastened to an ordinary slide with sealing wax. The drop to be examined is placed on a large-sized cover-glass and covered with a smaller one. Olive oil or vaseline

is painted round the edge of the smaller cover-glass to prevent evaporation, and the preparation is placed over the hole in the plate (Fig. 31).

The slide bearing the copper plate is clamped to the stage of the microscope (Fig. 32). The flame




FIG. 32.—SIMPLE WARM STAGE SHOWN IN OPERATION.

of a spirit-lamp is applied to the extremity of the rod, and the heat is conducted to the plate and thence transmitted to the specimen. That the temperature of the copper plate may be approximately that of the body, the lamp is so adjusted that a fragment of cacao butter and wax placed close to the preparation is melted.

For more accurate observations, the apparatus shown in Fig. 33 may be employed. The vessel *f*, filled with water which has been boiled to expel the air, is heated by means of a gas-flame at *g*. The warmed water ascends the india-rubber tube *c* to the brass box *a*. The box is pierced by a tubular aperture to admit light to the object, and has an




FIG. 33.—SCHÄFER'S WARM STAGE.

exit tube *c'*, by which the cooled water from the stage returns to be reheated by the flame *g*. At *d* is a gas-regulator, so that a constant temperature at any desired point can be maintained.

Another form, in which warm water or steam can be used for heating, and by the employment of iced water also used for observing the effects of cold, is shown in Fig. 34. It consists of a hollow rectangular box, with a central opening (*C*) permitting

the passage of light. The water makes its exit and entrance at the side tubes *a, a*, and the temperature is indicated by a thermometer in front.




FIG. 34.—STRICKER'S WARM STAGE.

Israel's Warming Apparatus.—It is obvious that in employing very high powers a difficulty will be presented by the warm stages just described,




FIG. 35.—SECTION OF ISRAEL'S WARMING APPARATUS AND DROP-CULTURE SLIDE.

owing to their interference with the illumination. To overcome this an apparatus has been constructed by which the slide is warmed from above.*

* Israel, *Zeitsch. f. Wiss. Mikrosk.*, ii., pp. 459-63. 1885.

The drop-culture slides are provided with a shallow groove 1 mm. deep and 1 mm. broad, cut round the concavity. Into this the cover-glass fits, so that its upper surface is flush with that of the slide. The heating apparatus consists of a flat disk-shaped box with a central conical aperture (Fig. 35).

The entrance and exit pipes are fixed on at a




FIG. 36.—ISRAEL'S WARMING APPARATUS.

right angle to the side (Fig. 36). The former, *z*, is of metal, and the latter, *a*, of glass fitted with a thermometer, the bulb of which, *k*, is contained within the box. A partition, *s*, keeps up a current between the openings of the pipes, which are supported on a stand and connected by tubing with the hot-water supply (Fig. 37).

A mixture of paraffine and vaseline is recommended for indicating the temperature of the chamber, and experience has shown that if a

temperature of 37° C. be required the temperature of the water in the box must range between 42° and 47° C.

Gas Chambers.—To investigate the action of




FIG. 37.—ISRAEL'S WARMING APPARATUS IN OPERATION.

gases or vapours upon micro-organisms, a modification of the simple moist chamber (Fig. 30) may be employed (Fig. 38).

A piece of glass tubing is first fixed to the slide by means of sealing wax, and the ring of putty is

so placed as to include the end of this, leaving a small interval at the side, or a little notch is




FIG. 38.—SIMPLE GAS CHAMBER.

made in the putty opposite, so as to afford an exit for the gas or vapour (Fig. 39).




FIG. 39.—GAS CHAMBER IN USE WITH APPARATUS FOR GENERATING CARBONIC ACID.

A more complicated apparatus, combining both a warm stage and a gas chamber, is shown in Fig. 40. This consists of a rectangular piece of ebonite (EE)

fixed to a brass plate which rests on the stage of the microscope. On the upper surface of the ebonite is




FIG. 40.—STRICKER'S COMBINED GAS CHAMBER AND WARM STAGE.

another brass plate (p), with an aperture (c) leading into a brass tube closed below by a piece of glass. To heat the apparatus the copper wire B is placed




FIG. 41.—SIMPLE MOIST CHAMBER ADAPTED FOR TRANSMISSION OF ELECTRICITY.

on the tube a , and its extremity heated by the flame of the lamp. The nearer the lamp to the stage the higher the temperature, which is indicated by the thermometer t . To employ it as a gas chamber

the wire B is laid aside, and the gas is conducted into the chamber by the tube a' , and escapes by the tube a .

Application of Electricity.—To study the effect of electricity we may prepare a drop-culture in the moist chamber (Fig. 41). The cover-glass to be used is provided with two strips of tinfoil,




FIG. 42.—APPARATUS ARRANGED FOR TRANSMITTING ELECTRICITY.

which are isolated from the brass of the microscope, and so arranged that a current of electricity may be passed through them (Fig. 42).

A much simpler plan, which may also be employed, is to take an ordinary glass slide and coat the surface with gold-size. The slide is then pressed firmly down on gold-leaf or tinfoil and allowed to dry. When dry, the metal is scraped away, leaving

two triangles with a small interval between them, as in Fig. 43.




FIG. 43.—SLIDE WITH GOLD-LEAF ELECTRODES.

The liquid containing the micro-organisms is placed between the electrodes, covered with a cover-glass, and then subjected to the electric current.

CHAPTER V.

EXAMINATION OF AIR, SOIL, AND WATER.

AIR.

THE air, as is well known, contains in suspension mineral, animal, and vegetable substances. The mineral world is represented by such substances as silica, silicate of aluminium, carbonate and phosphate of calcium, which may be raised from the soil by the wind, and particles of carbon, etc., which gain access from accidental sources. Belonging to the animal kingdom we find the *débris* of perished creatures, as well as sometimes living animals. The vegetable world supplies micrococci, bacilli, and other forms of the great family of bacteria, spores of other fungi, pollen seeds, parts of flowers, and so forth. The air of hospitals and sick rooms has been found to be especially rich in vegetable forms, *e.g.*, fungi and spores have been observed as present in particularly large numbers in cholera wards, spores of trichophyton have been discovered in the air of hospitals for diseases of the skin, and achorion in wards with cases of favus. The tubercle-bacillus is said to have been detected in the breath of patients suffering from phthisis.

These points indicate that, in addition to the interest for the microbiologist, considerable importance, from a hygienic point of view, must be attached to the systematic examination of the air. Especially a knowledge of the microbes which are found in the air of marshy and other unhealthy districts, and in the air of towns, dwellings, hospitals, workshops, factories, and mines, will be of practical value.

Miquel,* who has particularly studied the bacteria in the air, has found that their number varies considerably. The average number per cubic metre of air for the autumn quarter at Montsouris is given at 142, winter quarter 49, spring quarter 85, and summer quarter 105. In air collected 2,000 to 4,000 metres above the sea-level, not a single bacterium or fungus spore was furnished, while in 10 cubic metres of air from the Rue de Rivoli (Paris) the number was computed at 55,000.

The simplest method for examining the organisms in air consists in exposing plates of glass or microscopic slides coated with glycerine, or a mixture of glycerine and glucose, which is stable, colourless, and transparent. Nutrient gelatine spread out on glass plates (p. 91), may be exposed to the air for a certain time, and then put aside in damp chambers for the colonies to develop. Sterilised potatoes, prepared in the usual way (p. 99), may be similarly exposed. In both the last-mentioned methods

* Miquel, *Organismes Vivantes de l'Atmosphère*.

separate colonies develop, which may be isolated as already described, and pure cultivations carried on in various other nutrient media (p. 97). Nutrient gelatine has also been employed in the special methods of Koch and Hesse.

Koch's Apparatus.—This consists of a glass jar, about six inches high, the neck of which is plugged with cotton-wool. In the interior is a shallow glass capsule, which can be removed by means of a brass lifter. The whole is sterilised by exposure to 150° C. for an hour in the hot-air steriliser. The nutrient gelatine in a stock-tube is liquefied, and the contents emptied into the glass capsule. The jar is exposed to the air to be examined for a definite time, the cotton-wool plug replaced, and the apparatus set aside for the colonies to develop.

Hesse's Apparatus (Fig. 44).—The advantage of this apparatus consists in that a known volume of air can be examined. A glass cylinder, 70 cm. long and 3.5 cm. in diameter, is closed at one end by an india-rubber cap, perforated in the centre. Over this fits another cap, which is not perforated. The opposite end of the cylinder is closed with a caoutchouc stopper, perforated to admit a glass tube plugged with cotton-wool. The tube can be connected by means of india-rubber tubing with an aspirating apparatus. This apparatus consists of a couple of litre-flasks, suspended by hooks from the tripod-stand which supports the whole apparatus.

The cylinder, caps, and plug are washed with solution of corrosive sublimate, and then with alcohol. After being thus cleansed, 50 ccm. of nutrient gelatine are introduced, and the whole sterilised by steaming for half an hour for three successive days.




FIG. 44.—HESSE'S APPARATUS.

After the final sterilisation, the cylinder is rotated on its long axis, so that the nutrient medium solidifies in the form of a coating over the whole of the interior. When required for use, the cotton-wool plug is removed from the small glass tube, and the latter connected with the upper flask by means of the india-rubber tubing.

The apparatus is placed in the air which is to be examined, the outer india-rubber cap removed from the glass cylinder, and the upper flask tilted until the water begins to flow into the lower one. The emptying continues by syphon action, and air is drawn in along the cylinder to replace the water. When the upper flask is empty, the position of the two is reversed, and the flow again started. When a sufficient volume has been drawn through the cylinder, the outer cap and the cotton-wool plug are replaced, and it is set aside for the colonies to develop. As an example, twenty-five litres of air from an open square in Berlin gave rise to three colonies of bacteria and sixteen moulds; on the other hand, two litres from a schoolroom just vacated by the scholars gave thirty-seven colonies of bacteria and thirty-three moulds.

Various forms of "aeroscopes" and "aeroniscopes" have from time to time been employed. Pouchet's aeroscope consists of a small funnel, drawn out to a point below which is a glass slip coated with glycerine. The end of the funnel and the glass slip are enclosed in an air-tight chamber, from which a small glass tube passes out, connected by india-rubber tubing with an aspirator. The air passing down the funnel strikes upon the glycerine, which arrests any solid particles. For a description of the more exact apparatus employed by Maddox, Cunningham, and Miquel, reference should be made to the writings of these

authors, and particularly to the treatise published by the last-named.

EXAMINATION OF SOIL.

Surface soil, or mould, is exceedingly rich in bacteria. Miquel, *e.g.*, has computed that there exists in a gramme of soil an average of 750,000 germs at Montsouris, 1,300,000 in the Rue de Rennes, and 2,100,000 in the Rue de Monge. As agents of putrefaction and fermentation they play a very important *rôle* in the economy of nature; but there exist in addition bacteria in the soil which are pathogenic in character. Pasteur has succeeded in isolating the bacillus of anthrax from the earth, and sheep, sojourning upon a plot of ground where animals with anthrax had been buried, succumbed to the disease. Pasteur considered that the spores were conveyed by worms from buried beasts to the surface soil. The bacillus of malignant œdema is also present in soil, and Nicolaier has cultivated a bacillus from earth which produces tetanus in mice, rabbits, guinea-pigs, and other animals.

To obtain a cultivation of the microbes in soil a sample of the latter must be first dried and then triturated. It may then be shaken up with distilled water, and from this a drop transferred to sterilised bouillon. The employment of solid media is, however, much more satisfactory: a sample of earth is collected, dried, and triturated, and a small quantity

sprinkled over the surface of nutrient gelatine prepared for a plate-cultivation. In another method the gelatine is liquefied in a test-tube, the powder added, and, in the usual way, distributed throughout the medium, which is then poured out upon a glass plate. Just in the same way the dust which settles from the air in houses and hospitals, or food substances in powder, may be distributed over nutrient gelatine, and the micro-organisms which develop studied, both as to their morphological and biological characteristics.

EXAMINATION OF WATER.

As in the case of air, so, too, in that of water a knowledge of the micro-organisms which may be present is not only of interest to the microbiologist, but of the greatest importance in practical hygiene. Common putrefactive bacteria and vibrios may not be hurtful in themselves, but they indicate the probability of the presence of organic matter in some of which there may be danger.*

The Microzyme Test, which was introduced for their detection, consisted in adding three or four drops of the sample of water to 1 or 2 ccm. of Pasteur's fluid, the nourishing fluid having been previously boiled in a sterilised test-tube. If the microzymes or their germs existed in the water, the liquid in a few days became milky from the

* Parkes, *Manual of Practical Hygiene*. 1883.

presence of countless bacteria. This test is of no real value, for it does little more than indicate that bacteria are present, which we know to be the case in all ordinary water, and even in ice. On the other hand, the bacteriological test of Koch is a most valuable addition to the usual methods of water-analysis. It enables us not only to detect the presence of bacteria, but to ascertain approximately their number, and to study very minutely their morphological and biological characteristics. The importance of a thorough acquaintance with the life-history of the individual micro-organisms cannot be too strongly insisted upon. For example, by such means the spirillum of Asiatic cholera can be distinguished from other comma-shaped organisms, and inasmuch as its presence may be an indication of contamination with choleraic discharges, such water should be condemned for drinking purposes, even though we may not yet be in a position to affirm that the microbe is the cause of the disease. The test, in short, consists in making plate-cultivations of a known volume of water, counting the colonies which develop, isolating the micro-organisms, and studying the characters of each individual form.

Collection and Transport of Water Samples.—Erlenmeyer's conical flasks of about 100 ccm. capacity may be employed with advantage for collecting the samples of water. They are cleansed, plugged, and sterilised in the hot-air

steriliser. When required for use, the plug is removed and held between the fingers, which must not touch the part which enters the neck of the flask. About 30 ccm. of the water to be examined are introduced into the flask, and the plug must be quickly replaced and covered with a caoutchouc cap. If collected from a tap, the water should first be allowed to run for a few minutes, and the sample should be received into the flask without the neck coming into contact with the tap. From a reservoir or stream the flasks may be filled by employing a sterilised pipette. During transport contact between the water and cotton-wool plug must be avoided, and if likely to occur the sample must be collected and forwarded in a Sternberg's bulb.

Examination by Plate-cultivation. — The apparatus for plate-cultivation should be arranged as already described. Crushed ice may be added to the water in the glass dish to expedite the setting of the gelatine, so that the plate may be transferred as quickly as possible to the damp chamber. The caoutchouc cap is removed from the flask, and the cotton-wool plug singed in the flame to prevent contamination from adventitious germs on the outside of the plug. The flask is then held slantingly in the hand, and the plug twisted out and retained between the fingers. With a graduated pipette a drop of the sample is transferred to a tube of liquefied nutrient gelatine, and the plugs of the flask

and of the tube quickly replaced. If the water be very impure, it may be necessary to first dilute the sample with sterilised water. The inoculated tube must be gently inclined backwards and forwards and rolled as already explained, to distribute the germs throughout the gelatine, and the gelatine finally poured on a plate. When the gelatine has set, the plate is transferred to a damp chamber, which should be carefully labelled and set aside in a place of moderate temperature. In about two or three days the cultiva-




FIG. 45.—APPARATUS FOR ESTIMATING THE NUMBER OF COLONIES IN A PLATE-CULTIVATION.

tion may be examined. In some cases the colonies may be counted at once; more frequently they are so numerous that the plate must be placed on a dark background, and a special process resorted to. A glass plate, ruled by horizontal and vertical lines into centimetre squares, some of which are again subdivided into ninths, is so arranged on a wooden frame that it can cover the nutrient-gelatine plate without touching it (Fig. 45). A lens is added to assist in discovering minute colonies. If then the colonies are very numerous, the number in some

small divisions is counted, if less in some large ones, and an average is obtained from which the number of colonies on the entire surface is calculated. A separate calculation of the *liquefied* colonies should be also made, and their number, as well as the total number of colonies present in 1 ccm. of the sample, recorded. Any peculiar macroscopical appearances, colour, etc., should be noted, and then the microscopical appearances of the colonies studied. Lastly, examination of the individual organisms should be made by cover-glass preparations, and by inoculation of nutrient gelatine, potatoes, and other media.

Examination by Test-tube-cultivation.—A drop of the sample of water may also be added to liquefied nutrient gelatine in a tube, the organisms distributed as already explained (p. 93), and the gelatine allowed to solidify in the tube. A rough comparison of water samples may be made in this way.

Microscopic Examination.—A drop of the water may be mounted and examined in the way described under drop-cultivations (p. 112), or a drop is allowed to evaporate on a cover-glass placed under a bell-glass. This is then passed three times through the flame, and stained in the usual manner. The examination of rain water, drinking water, tap water, sea water, various liquids and infusions, etc., by these methods, opens up a wide field for research. Pettenkofer has shown that impregnation with

carbonic acid of water containing many bacteria diminishes the number of the latter. The examination of waters before and after filtration, or after addition of chemical substances, are matters which require further investigation.

CHAPTER VI.

EXPERIMENTS UPON THE LIVING ANIMAL.

To carry out the last of Koch's postulates, and so complete the chain of evidence in favour of the causal relation of micro-organisms to disease, and to study the mode of action of a pathogenic bacterium, it is necessary to introduce into a living animal a pure cultivation of the micro-organism in question. For this purpose various animals are employed—such as mice, guinea-pigs, rabbits, pigeons, and fowls.

Inhalation of Micro-organisms.—The animals may be made to inhale an atmosphere impregnated with micro-organisms by means of a spray. In this way Friedländer succeeded in administering the bacteria of pneumonia to mice, and the production of tuberculosis by experimental inhalation has thrown light upon the clinical records of cases reported as instances of the infectiousness of phthisis.

Administration with Food.—A sheep fed upon potatoes which have been the medium for the cultivation of the anthrax bacillus dies in a

few days. Similarly, animals fed upon the nodules of bovine tuberculosis become tubercular, and even the flesh and milk of tuberculous animals will occasionally set up tuberculosis.

Cutaneous and Subcutaneous Inoculation.—Cutaneous inoculation may be carried out by making a superficial wound, and inoculating it with a sterilised platinum needle, charged with the micro-organisms to be inoculated. Another simple method is to take a sterilised knife, infect the point with the material to be inoculated, and then make a minute wound or incision. In either case a situation should be selected, such as the root of the ear, which cannot be licked by the animal after the operation.

Subcutaneous inoculation is very simple and effectual, and consequently the method most frequently employed. The animal selected—for example, a guinea-pig—is held by an assistant, who covers it with a towel, leaving only the hinder extremities exposed. By so doing, and gently laying it upon its back, with its head low, a guinea-pig passes apparently into a state of hypnotism, and the trivial operation can be performed with little or no movement on the part of the animal. From a spot on the inner side of the thigh the hair is cut close with a small pair of scissors curved on the flat, and the skin must be thoroughly purified with sublimate solution. A small fold of skin is then pinched up

with a pair of sterilised forceps, and with a pair of sharp sterilised scissors, or with a tenotomy knife, a minute incision is made. A sterilised platinum loop is charged with the material to be inoculated, and the loop is gently inserted under the skin, forming a small pocket in the subcutaneous tissue. The needle is then withdrawn, and the sides of the wound gently pressed into apposition. In a mouse the same process is adopted, with the exception that the root of the tail is the usual site of the operation. In a method suggested by Koch an assistant can be dispensed with: a glass bell reversed is placed as a cover to a wide-mouthed glass jar, in which a mouse is held by the tail with a pair of forceps, while the cover is so placed over the mouth of the jar as to leave a small interval near the rim uncovered. The mouse rests with its head downwards and with its feet against the inner wall of the jar, and in the interval between the cover and the rim the root of the tail is exposed, and must be cleansed and treated as already described.

Special Operations.—In many cases it is absolutely necessary to perform an operation of greater severity. After the administration of an anæsthetic, infective material may be introduced into the peritoneal cavity by the performance of abdominal section, or injected into the duodenum in the manner employed in the case of Koch's

comma bacilli by Nicati and Rietsch. In such cases antiseptic precautions must be rigidly followed, and use made of iodoform and other antiseptic dressings. The disinfection of the skin of the animal, of the instruments employed, and of the hands of the operator, are details essential to secure success. To inoculate tubercular matter, sputum may be rubbed up with distilled water, filtered, and the filtrate injected into a tracheal fistula, or the first steps of the operation of iridectomy may be performed, and tubercular material inserted in the anterior chamber of the eye. The advantage of the latter method consists in that it enables the results and changes to be observed from day to day. A cultivation of micro-organisms may also be mixed with sterilised water, and then injected with a syringe directly into the circulation. In rabbits this may be performed without difficulty by injecting the large vein at the base of the ear with a Pravaz' syringe. Before every inoculation the instruments must be sterilised, as already explained, by employing an Israel's case, and after each operation all instruments should be placed in strong sublimate solution (1 in 100), wiped dry, and sterilised in the hot-air steriliser, before they are put away. If these precautions be not observed, instances of accidental infection are sure to occur.

CHAPTER VII.

EXAMINATION OF ANIMALS EXPERIMENTED UPON AND THE METHODS OF ISOLATING MICRO- ORGANISMS FROM THE LIVING AND DEAD SUBJECT.

METHOD OF DISSECTION AND EXAMINATION.

ALL animals that die after an experimental inoculation should be examined immediately after death. Every precaution must be taken, in conducting the dissection, to exclude extraneous micro-organisms, and all instruments employed must have been sterilised in the hot-air steriliser, or heated in the Bunsen burner. If a mouse, for example, has died after an inoculation, it should be at once pinned out by its feet on a slab of wood or in a gutta-percha tray, and bathed with sublimate solution. In the same way, before examining a dead rabbit, a stream of sublimate should be directed over it to lay the fur, which otherwise interferes with the dissection. The hair should be cut away with sterilised scissors from the seat of inoculation, which is the first part to be examined, and any suppuration, hæmorrhage, œdema, or other pathological

change should carefully be noted. From any pus or exudation that may be present, material for inoculations should at once be taken, and cover-glass-preparations made for microscopical examination.

To examine the internal organs and to make inoculations from the blood of the heart or spleen, the skin is cut through from below upwards in the median line of the abdominal and thoracic regions. The abdominal cavity is then opened, and the walls pinned back on either side of the animal. Any abnormal appearances should be noted, and especially the state of the spleen should be examined, by turning the intestines aside. After noting its appearances it should be removed with sterilised forceps and scissors, and deposited upon a sterilised glass slide. After washing it with sublimate solution by means of a camel's hair brush or strip of filter paper, it should be incised with sterilised scissors; the pulp may be squeezed out from the cut surface, and test-tubes of nutrient gelatine and agar-agar can be inoculated from it, and, if necessary, potato- and drop-cultivations also established. Precisely the same care must be taken in examining lymphatic glands, tubercles, or pathological nodules; any chance putrefactive micro-organisms on the surface are destroyed by the sublimate solution, and a section is then made, and a minute fragment snipped out of the centre of the nodule, to be examined or transferred to the nutrient medium. The examination of the thorax is made by cutting

through the ribs on either side of the sternum with sterilised scissors, and turning the sternum up where it will be out of the way. The pericardium is then opened, and the right auricle or ventricle pierced with the point of a sterilised scalpel, and inoculations and cover-glass-preparations are made from the blood which escapes.

The lungs also require to be especially studied. They should be incised with a sterilised scalpel, and inoculations and cover-glass-preparations made from the cut surface. It may be necessary to embed a piece of lung or fragment of spleen, so that it shall be free from air. This may be done by isolating a fragment with the precautions just described and depositing it upon the surface of a test-tube of nutrient agar-agar. The contents of another tube, which have been liquefied, and allowed to cool almost to the point of gelatinisation, must then be poured over it. From a potato a little cube must be cut, the tissue deposited in the trough thus formed, and the cube replaced. Blood may also be taken directly from a vein by laying it bare by dissection, making a small section with sterilised scissors, and inserting a looped platinum needle, the needle of a Pravaz' syringe, a capillary tube, or the extremity of the capillary neck of a Sternberg's bulb. If the cultivation be contaminated by the presence of putrefactive or other micro-organisms, they must be isolated subsequently by carrying out a series of plate-cultivations.

Having completed the dissection, the organs of such a small animal as a mouse may be removed *en masse* and transferred to absolute alcohol for subsequent examination. In other cases it may be only necessary to reserve portions of each organ. In any case it should be remembered that with a virulent micro-organism, *e.g.*, anthrax, any remaining part of the animal should be cremated, and the hands and all instruments should be thoroughly disinfected.

Isolation of Micro-organisms from the Living Subject.—Micro-organisms in the living subject may be isolated from pus of abscesses, or other discharges, and from the blood and tissues. Abscesses should be opened, and other operations performed, when practicable, with Listerian precautions, and a drop of the discharge taken up with a looped needle or capillary pipette, as already explained.

To make a cultivation from the blood of a living person, the tip of a finger must be well washed with soap and water and bathed with strong sublimate, or 1 in 20 carbolic, solution. Venous congestion is produced by applying an elastic band or ligature to the finger, which is pricked with a sterilised sewing needle. From the drop of blood which exudes the necessary inoculations and examinations can be made.

CHAPTER VIII.

PART II.

GENERAL BIOLOGY OF BACTERIA.

CHAPTER VIII.

GENERAL MORPHOLOGY AND PHYSIOLOGY.

BACTERIA may be considered as minute vegetable cells destitute of nuclei. They are distinguished from animal cells by being able to derive their nitrogen from ammonia compounds, and they differ from the higher vegetable cells in being unable to split up carbonic acid into its elements, owing to the absence of chlorophyll. Von Engelmann and Van Tieghem include among the bacteria certain organisms, named by them *Bacterium chlorinum*, *Bacterium viride*, and *Bacillus virens*, which are coloured green by this substance; but further researches are required before any conclusions are definitely arrived at as to the place of these particular organisms in the vegetable kingdom. It is quite possible that they may be Algæ, and they will, therefore, find no place in the classification which will be here adopted.

Chemical composition.—For our knowledge of the composition of bacteria we are chiefly indebted to Nencki. Their constituents are found on

analysis to vary slightly, according to whether the bacteria are in zooglœa or in the active state. In the latter condition they are said to consist of 83·42 per cent. of water. In one hundred parts of the dried constituents there are the following:—

A nitrogenous body	84·20
Fat	6·04
Ash	4·72
Undetermined substances	5·04

This nitrogenous body is called *Mycoprotein*, and consists of

Carbon	52·32
Hydrogen	7·55
Nitrogen	14·75

but no sulphur or phosphorus.

The nitrogenous body appears to vary in different species, for in *Bacillus anthracis* a substance has been obtained which does not give the reactions of mycoprotein, and, therefore, is distinguished as anthraxprotein.

Considering bacteria as cells, we may speak of the cell-wall and the cell-contents.

Cell-wall.—The cell-wall consists of cellulose, or according to Nencki in the putrefactive bacteria of mycoprotein. It may be demonstrated by the action of iodine, which contracts the protoplasmic contents, and renders the cell-wall visible. The author has taken advantage of the action of iodine to differentiate by staining the sheath of the *Bacillus*

anthracis from its contents. If we stain cover-glass preparations of this bacillus by the method of Gram, we get the following results. By the first solution the rods are uniformly stained blue; by subjecting them to the iodine solution, the protoplasmic contents are contracted, while the next solution, alcohol, decolorises the sheath, which may be then stained in contrast with eosin.

The cell-wall may be either pliable or rigid. Pliability is observed in the long filaments, which are endowed with a slow vermicular movement, while rigidity accounts for the maintenance of the characteristic form of several species, such as spirilla.

Cell-contents.—The cell-protoplasm yields mycoprotein. In some it is homogeneous, and in others granular. The action of the aniline dyes indicates a close relation to nuclear protoplasm, though all nuclear stains are not suitable for bacteria. In some cases also, the bacteria remain stained under the influence of a reagent, which removes the colour from nuclei. The power of fixing the stain is not always present, and indicates a difference in the protoplasm of different species. Thus in staining phthisical sputum, the nitric acid removes the stain from all bacteria and bacilli




FIG. 46.

FROM A PREPARATION OF *BACILLUS ANTHRACIS*, DOUBLE STAINED WITH GENTIAN VIOLET AND EOSIN. The Sheath was stained pink, and the cell-contents blue. $\times 1200$.

present, with the exception of the tubercle bacillus. This difference in the protoplasm of different species is also illustrated by the necessity in many cases of using special processes, owing to the ordinary methods being unsatisfactory or not producing any result.

The protoplasm of some bacteria contains starch granules; thus *Clostridium butyricum* gives the starch reaction with iodine. Sulphur granules are present in some species of *Beggiatoa* which thrive in sulphur springs. The colouring matter of the pigment bacteria is probably external to the cell as a rule; for example, in *micrococcus prodigiosus* the pigment granules are distinctly between the cells; on the other hand, in *Beggiatoa roseo-persicina*, or the peach-coloured bacterium, the special pigment *bacterio-purpurin* appears to be dissolved in the cell protoplasm. In *Bacillus pyocyaneus* the pigment is certainly not localised entirely in the cell; for it becomes rapidly diffused in the surrounding medium, considerably beyond the confines of the growth itself.

Gelatinous envelope.—In several species, either as a result of a secretion from the cell, or of the absorption of moisture and swelling up of the outer layer of the cell-wall, a mucinous or gelatinous envelope develops around them. This envelope may form a capsule, such as we meet with in certain bacteria found in the rusty sputum of pneumonia, and in *Micrococcus tetragonus*; or it may

occur as a continuous sheath around a chain of bacteria, which by its disappearance sets the individual links free. The capsule is soluble in water, and under some circumstances is difficult to demonstrate. In the pneumo-coccus of Friedländer the capsule disappears on cultivation, but reappears in preparations made from an inoculated animal. In the pleuritic fluid of a mouse these cocci are often found with a strikingly well-marked capsule, and in other capsuled cocci the extent of the envelope has been observed to vary considerably in the same species of bacterium.

When this gelatinous material forms a matrix, in which numbers of bacteria are congregated in an irregular mass, we have what is termed a zooglœa. The zooglœan stage is a resting stage, often preceded or followed by a motile stage. Thus bacteria may be present in a solution in an active state, and after a time a scum or pellicle forms on the surface of the liquid, which consists of zooglœa. At the edges of the zooglœa individuals may be seen again to become motile, detaching themselves from the edges of the mass, and swimming off in the surrounding fluid.

The same may be observed sometimes in cultivations started in nutrient gelatine. The inoculated bacteria grow and multiply, and liquefy the gelatine, and after a time a zooglœan film appears on the surface of the liquefied layer. On potatoes the appearances are very varied. In a bacillus which

readily develops on unsterilised potatoes, the zoogloea may spread over the cut surface, forming a pellicle which can be raised *en masse* like a delicate veil. Another bacillus forms a zoogloea, consisting of a tenacious layer which can be drawn out in long stringy threads. In *Ascococcus Billrothii* the gelatinous envelope develops to such an enormous extent that it forms the characteristic feature of the species (Fig. 47).




FIG. 47.—ASCOCOCCUS BILLROTHII, $\times 65$. [After Cohn.]

Form.—The individual cells vary in form, and may either remain isolated or attached to each other. Round cells and egg-shaped cells are called *cocci*. The spherical form is the most common, but cocci are occasionally exclusively ovoid, as in *Streptococcus bombycis*. The giant cocci of some species are spoken of as *megacocci*, to distinguish them from the ordinary cocci, such as *micrococci*. The fission by which the cocci increase may take place in one direction only, and if the two resulting

cells remain attached to each other, they form a *diplococcus*. If these two cells again divide, and the resulting cells remain linked together, we get a chain or rosary, or *streptococcus* (Figs. 48, 49, 50).



FIG. 48.—STREPTOCOCCUS AND SARCINOCOCCUS FROM A DROP-CULTIVATION, X 1200.



FIG. 49.—STREPTOCOCCUS IN THE BLOOD OF A RABBIT, X 1200.

These chains may consist of a few individuals linked together, or of several hundreds, in which case the chains are generally curved or twisted. If the division occur in two directions, so that




FIG. 50.—STREPTOCOCCUS OF PROGRESSIVE TISSUE NECROSIS IN MICE. [After Koch.]

four cocci result, a tetrad or *merismopedia* is formed. If the division occur in three directions, one coccus divides into eight, and we get a packet form or *sarcinacoccus*. Immediately after division, the daughter cells are not perfectly

circular, but are flattened or faceted where they are opposite to each other. They gradually become rounded off, and each daughter cell is then ready to divide in its turn. In other cases the cocci after division only form irregular heaps or collections like bunches of grapes. This form is sometimes distinguished as *staphylococcus*, but it cannot be considered an important feature. Where we find irregular masses or balls embedded in a copious gelatinous matrix, the extent of the latter affords a characteristic condition described as *ascococcus*.

Another type is the rod, characteristic of bacterium and bacillus. The rods may vary considerably in length. The very short rods with rounded ends are very difficult to distinguish from the oval cocci, but differ in that a rod, however short it may be, must have at least two sides parallel. The *vibrio* or bent rod may be considered as the connecting link between the rods and the corkscrew forms or *spirilla*. Lastly we have the filamentous forms, which may be straight, *leptothrix*, or wavy, *spirochæta* (Fig. 51), or the wavy thread may be looped and entwined on itself, *spirulina* (Plate I., Fig. 37).

By *involution* forms we signify certain irregular shapes which result especially in exhausted cultivations. They are peculiar, oval, pear-shaped, or irregular enlargements (Plate I, Figs. 31 to 36).

Movement.—Many bacteria are devoid of move-

ment throughout the whole of their life history. Others, during certain stages of their life cycle, and possibly some forms always, are endowed with locomotive power. The character of the movement is very varied, and ranges from a slow undulatory motion to one of extreme rapidity. Many appear to progress in a definite direction. Others move continuously, first in one direction and then in another, and others again seem to hesitate before altering their course. They may either glide along smoothly or progress with a tremulous action. They appear to be able to avoid obstacles, and to




FIG. 51.—SPIROCHÆTA FROM SEWAGE WATER, $\times 1200$.

set themselves free from objects with which they have accidentally come into contact. Vibrios have a peculiar serpentine movement, but other forms, such as the commonly-known *Bacterium termo* and segments of spirilla, such as comma-bacilli, revolve around their long axis as well as make distinct progression. The complete spirilla are characterised by the familiar corkscrew movement. With regard to cocci there is some doubt as to whether they are endowed with independent movement; any quivering or oscillation is generally regarded as only brownian or molecular. In some

straight thread-forms, which are motile, the movement is very slow and vermicular in character, but in wavy threads, such as the *Spirochæte plicatilis*, there is not only an undulatory motion, with rapid progression across the field of the microscope, but if they are confined by more or less *débris*, they give very peculiar and characteristic spasmodic movements.

The rod-forms of *Proteus vulgaris* exhibit very extraordinary movements on the surface of solid nutrient gelatine. Groups of rods may be observed to pass each other in opposite directions. Single individuals meet and progress side by side, or one or more individuals may part from a group and glide away independently. Occasionally a number of rods progress in single file. It is, however, difficult to believe that these movements can occur on a solid surface. The author is inclined to believe that there is an almost inappreciable layer of liquid on the surface of the gelatine, which is expressed after the gelatine sets. In tubes of nutrient agar-agar gelatinised obliquely and then kept upright the liquid so expressed collects at the bottom of the sloping surface.

What the means are by which bacteria are endowed with the power of spontaneous movement and of progression may still be said to be unsettled. The author has watched the movement of long slender threads in sewage-contaminated water, which could only be explained by the inherent

contractility of the protoplasmic contents; for if any drawing or propelling organ existed in proportion to the length of the organism, it would probably have been visible. But in many cases the organism is provided with a vibratile lash or




FIG. 52.

1. Coccus with flagellum. 2. Similar coccus dividing with two flagella
3. Colony of flagellated macrococci of *Beggiatoa roseopersicina*. 4. Short rod from the same *Beggiatoa* with flagella [all after Zopf]. 5. Bacillus with flagella [from a photograph by Koch]. 6. *Bacillus subtilis* [after Brefeld]. 7, 8. Short rod-forms of *Beggiatoa roseopersicina* with one flagellum [after Zopf]. 9. Very long rod of the same, with flagellum at both ends [after Warming]. 10. *Vibrio*, with double flagellum at each end [after Warming]. 11. *Vibrio*, with flagella [from a photograph by the author]. 12. *Spirillum* with flagella [after a photograph by Koch]. 13. *Spirillum* with flagella [after Zopf]. 14. *Spirillum* with double flagella [after Zopf]. 15. *Beggiatoa roseopersicina*, with a triple flagellum at one end; and 16, with a double flagellum at both ends [after Warming].

flagellum at one end, or with one or more at both ends (Fig. 52).

Some observers believe that the movement of cocci is due to the existence of a flagellum. In *Bacterium termo* the existence of a lash at either end was first determined by the researches of Dallinger and Drysdale. In motile bacilli, such as the hay bacillus and *Bacillus ulna*, and in vibrios and spirilla, the flagella can be readily recognised by expert microscopists with the employment of the best lenses, and, what is of equal importance, proper illumination. They are objects of extreme delicacy, and tenuity, and in stained preparations may be absent from retraction or injury. Koch succeeded in photographing them after staining with logwood, which turned them a brown colour. They may also be stained with the aniline dyes, for the author has observed them in vibrios in preparations stained with gentian violet, from which also they have been photographed, in spite of the violet colour, by the use of isochromatic dry plates.

It is not certain whether the flagella are extensions of the cell-wall, or derived from the internal protoplasm. Van Tieghem holds the first view, and does not regard them as motile organs at all. Zopf, on the other hand, adheres to the second view, and moreover believes that they can be retracted within the cell-wall.

Reproduction.—Bacteria multiply by fission, and by processes which may be considered as representing fructification. The bacteria exhibiting the latter processes have been divided into two

groups, distinguished by the formation of endospores in the one, and of arthrospores in the other. In the process of fission the cell first increases in size, and a transverse septum forms from the cell-wall, dividing the internal protoplasm into two equal parts; these may separate and lead an independent existence, or remain linked together. In chains of




FIG. 53.—*BACILLUS MEGATERIUM*.

- a.* A chain of rods, $\times 250$. The rest $\times 600$.
b. Two active rods.
d to *f.* Successive stages of spore-formation.
h and *l.* Successive stages of germination.

[After De Bary.]

cocci the individual cells are easily visible and distinct, but in the thread-forms resulting from the linking together of rods, as in the anthrax bacillus, the composition of the thread is only demonstrated by the action of reagents.

Endospore formation may be conveniently studied in *Bacillus anthracis*, *Bacillus megaterium*, or *Bacillus subtilis*. The protoplasm becomes granular,

and at certain points in the thread a speck appears, which gradually enlarges and develops into a circular or egg-shaped, sharply defined, highly refractive body. The spore grows at the expense of the protoplasm of the cell, which in time, together with the cell-wall, entirely disappears, and the spore is set free. These phenomena are best seen in an immotile bacillus in a drop-cultivation




FIG. 54.—*CLOSTRIDIUM BUTYRICUM*, X 1020.

B. Stages of spore-formation.

C. Stages of germination.

[After Prazmowski.]

on a warm stage, the whole process may then be observed continuously from beginning to end. Spores may form in each link of the thread, so that a regular row results, or they may occur at irregular intervals. Spore-formation also occurs in free rods in the centre or at one end. Occasionally a spore develops at the extreme end, giving a bacillus the appearance of a drum-stick. The spore may be

considerably wider, but is never longer than the parent cell.

Arthrospore formation is illustrated in *Leuconostoc mesenteroides*. Certain elements in the chain of cocci, apparently not differing from the rest,




FIG. 55.—A THREAD OF BACILLUS ANTHRACIS WITH SPORES, IN A DROP-CULTIVATION, $\times 1400$.

become larger, with tougher walls, and more refractive (Fig. 56). The remaining cells die, and these cells having acquired the properties of spores are set free, and can reproduce a new growth in any




FIG. 56.—LEUCONOSTOC MESENTEROIDES ; COCCI-CHAINS WITH ARTHROSPORES (after Van Tieghem and Cienkowski).

fresh nourishing soil. That this occurs in all species which do not form endospores is at present only a supposition.

Spores are invested by a thick membrane, which is believed to consist of two layers. To this they probably owe the property they possess of retaining vitality when desiccated, and of offering a greater

resistance to the action of chemical reagents and heat than the parent cells.

Spore-formation has been regarded by some as occurring when the nourishing soil is exhausted, thus providing for the perpetuation of the species. For instance, anthrax bacilli do not form spores in the living body, but when the animal dies it has been stated that development of spores takes place, and hence the danger of contaminating the soil if the body is disposed of by burial. Klein, however, has pointed out that if mice and guinea-pigs which have died of anthrax are kept unopened, the bacilli simply degenerate and ultimately disappear. Thus there is good reason for believing that spore-formation is not due to exhaustion of the pabulum, but probably free access to oxygen constitutes an important factor in inducing this condition. If we inoculate a potato with anthrax, copious spore-formation occurs, though we cannot consider that the nourishing soil has been exhausted. But we have in this case the surface of the potato freely exposed to the air in the damp chamber. In the same way, in cultivation on agar-agar solidified obliquely, so as to get a large surface, spore-formation readily takes place. Contamination of a burial-ground must result, therefore, from bodies in which a post-mortem examination has been made, by which the blood and organs have been freely exposed to the air, or from animals which have not been examined, owing to their hides being soiled with

excretions, and with blood which issues from the mouth and nostrils before death.

When spores are introduced into a suitable medium at a favourable temperature, they develop again into rods. The spore loses its sharp contour, and, at one pole or on one side, a pale process bursts through the membrane, gradually growing into a rod from which the empty capsule is thrown off (Figs. 53 and 54).

Spores differ from the parent cells in their be-




FIG. 57.—SPORES OF *BACILLUS ANTHRACIS*, STAINED WITH GENTIAN VIOLET, AFTER PASSING THE COVER-GLASS TWELVE TIMES THROUGH THE FLAME, $\times 1200$.




FIG. 58.—SPORE-BEARING THREADS OF *BACILLUS ANTHRACIS*, DOUBLE-STAINED WITH FUCHSINE AND METHYLENE BLUE, $\times 1200$.

haviour to staining reagents. Like them, they can be stained with aniline dyes, but not by the ordinary processes. They require to be specially treated. This is probably due to the tough capsule, which must first be altered or softened by heat or strong acid, until it allows the stain to penetrate.

Once stained, they again differ from the parent cells in resisting decolorisation; this fact is taken advantage of to double-stain spore-bearing bacilli.

In staining micro-organisms, the protoplasm is

sometimes broken up into irregular segments or granules, as in many spirilla, and we may perhaps add the bacilli of tuberculosis and leprosy. The beaded appearance of the tubercle bacillus is well known. Some observers have regarded the beads, others the bright spaces between them, as spores. But spores in unstained preparations appear as glistening bodies with sharp contour, and do not stain at all, or very little, by the ordinary processes. It appears, therefore, very doubtful whether either the clear spaces or the beads are spores, especially as the tubercle bacillus, when unstained, is a slightly curved hyaline rod, without any differentiation into granules. These considerations led the author to stain and examine tubercular sputum from various sources under careful illumination, and with such lenses as Powell and Lealand's $\frac{1}{25}$ in. Hom. imm. The tubercle bacillus may then be frequently seen to consist of a very delicate sheath, holding together a number of deeply stained granules, for the most part round or cylindrical, with irregular contour, and differing considerably in size, while the light interspaces are seen to vary in form according to the shape of the granules. In some preparations more distinct, and clearly ovoid, bodies may be observed which are sometimes terminal. They can be readily demonstrated by taking a photograph with a $\frac{1}{25}$ in. Hom. imm., and subsequently enlarging the negative to from 2,500 (Fig. 59) to 6,000 diameters.

It is not impossible that these ovoid bodies are spores, which, in their behaviour towards staining reagents, thus form an exception to the general rule. But there can be little doubt that a tubercle bacillus consists, for the most part, of a very delicate sheath, with protoplasmic contents which have a great tendency to be broken up or coagulated into little



FIG. 59.—TUBERCLE BACILLI IN SPUTUM, X 2500 (from photographs).




FIG. 60.
LEPROSY BACILLI, FROM A SECTION OF SKIN, X 1200.

segments or roundish granules, owing possibly to the treatment they are subjected to in making a microscopical preparation. This, however, does not always occur, for the bacilli at times are not beaded, but are stained in their entirety. In the leprosy bacilli a similar appearance occurs. In stained sections the rods have a beaded appearance, but the intervals between the granules are sometimes very long, and occasionally the protoplasm appears to have collected only at the extreme ends of the rod (Fig. 60). Very probably the appearances in the case of the bacillus of glanders (Fig. 61) and the bacterium of chicken-cholera (Figs. 62 and 63) may be similarly explained.



FIG. 61.
GLANDERS BACILLI, FROM A SECTION OF A GLANDERS NODULE, X 1200.

The fact that tubercular sputum preserves its virulence for several months, even after desicca-



FIG. 62.

BACTERIUM OF CHICKEN-
CHOLERA, FROM BLOOD
OF INFECTED HEN, \times
1200.

tion, has been attributed to the formation of spores, and Babes has drawn attention to ovoid grains in old cultivations of the bacilli, which he succeeded in staining red, while the bacilli were stained blue.

In his definition of spirilla Zopf gives the spore-formation as absent or unknown. In comma-bacilli in sewage water, the author has often noted appearances very suggestive of refractive




FIG. 63.

BACTERIUM OF CHICKEN-
CHOLERA,
FROM MUSCLE JUICE OF AN IN-
FECTED HEN, \times 2500 (from a
photograph).



FIG. 64.

COMMA-BACILLI IN SEWAGE
WATER, STAINED WITH
GENTIAN VIOLET, \times 1200.

spores (Fig. 64). The same also may be observed in vibrios, differing by their regular contour from the irregular spaces occasionally observed in stained preparations (Figs. 65 and 66). They are possibly only vacuoles.

Respiration and Nutrition.—Like all a-chlorophyllous vegetables, bacteria require for their nutrition oxygen, nitrogen, carbon, water, and certain mineral salts. Many require free access to oxygen, others can derive it from the oxidised

compounds in the medium in which they grow. Pasteur divided bacteria into two great classes—the ærobic and anærobic; and considered that the latter not only had no need for oxygen, but that its presence was actually deleterious. Though this view must be considerably modified, the terms are convenient, and are still retained. They are well illustrated by the bacillus of anthrax, and the bacillus of malignant œdema; and a simple plan of demonstration has been employed by the author. A fragment of tissue from the spleen, for example,



FIG. 65.

VIBRIOS IN WATER CONTAMINATED
WITH SEWAGE, X 1200.



FIG. 66.

SPIRRILLUM UNDULA,
X 1200.

known to contain anthrax bacilli, is deposited with a sterilised inoculating needle, with the necessary precautions, on the surface of nutrient agar-agar in a test-tube; another tube of nutrient agar-agar is liquefied, and when cooled down almost to the point of gelatinisation, a part is poured into the first tube, so that when it sets the piece of tissue is completely embedded. A piece of tissue from an animal suffering from malignant œdema is treated in the same way, and the tubes are placed in the incubator. If then we examine

them after two or three days, we shall find no change in the anthrax tube; the bacillus being eminently ærobic, no growth whatever has occurred. In the tube containing the bacilli of malignant œdema there will be a more or less characteristic cultivation.

The nitrogen which is essential for building up their protoplasm can be obtained either from albumins, or from ammonia and its derivatives. That the albumins can be dispensed with was shown by Pasteur, who employed an artificial nourishing solution built upon a formula representing the essential food constituents (p. 108).

Carbon is derived from such substances as cane sugar, milk sugar, and glycerine, and, in some cases, by the splitting up of complex proteid bodies.

Water is essential for their growth, but deprivation of water does not kill all bacteria. Desiccation on potato is employed for preserving some micro-organisms, as a new growth can be started, when required, by transferring some of the dried potato to fresh nourishing ground. Comma-bacilli, on the other hand, are destroyed by drying. Sugar, by abstracting water, prevents the development of micro-organisms in preserves.

Mineral or inorganic substances, such as compounds of sodium and potassium, and different phosphates and sulphates, are necessary in small proportions.

Circumstances affecting their growth.

Nature of the Soil.—Though we know the elements necessary, we are, nevertheless, as yet unable to provide a pabulum suitable for all kinds of bacteria. Thus we are quite unable to cultivate some species artificially. Others will only grow upon special media. Many grow upon nutrient gelatine; but some species only if it be acid or alkaline respectively. Whether in the latter case this is due purely to the reaction or to the presence of the particular ingredients is an unsettled point. Though the comma-bacillus of Koch, like the majority of organisms, grows best on an alkaline medium, yet the surface of a potato is acid, and on this it is well known to flourish at the temperature of the blood.

Effect of Temperature.—In their behaviour towards temperature bacteria vary considerably, but still for the majority we may distinguish a *maximum*, *optimum*, and *minimum* temperature.

Many grow best at the temperature of the blood, and hence the value of nutrient agar-agar, which is not liquefied at 37° C. The tubercle bacillus will only grow at a temperature varying between 30° and 41° C. On the other hand, many forms grow between the limits of 5° and 45° C. At these temperatures their functional activity is paralysed, but they are not destroyed, for by removal to favourable conditions they spring again to life. Bacteria seem to have a special power

of resisting the effects of cold. It has been stated that comma-bacilli exposed to a temperature of -10° for an hour, and bacilli of anthrax after exposure to a temperature of -110° C., still retained their vitality. Temperatures over 50° to 60° C. destroy most bacteria, but not their spores; spores of anthrax retain their vitality after immersion in boiling water, but are destroyed by prolonged boiling. Roughly speaking, all pathogenic bacteria grow best at the temperature of the blood, and non-pathogenic bacteria at the ordinary temperature of the room.

Effect of Movement.—Bacteria probably grow best when left undisturbed. Violent agitation of a vessel in which they are growing certainly retards their growth, but a steady movement is stated not to affect it; at any rate anthrax bacilli grow with enormous rapidity in the blood-vessels, in spite of the circulation.

Effect of Compressed Air.—Paul Bert maintained that a pressure of twenty-three to twenty-four atmospheres stopped all development of putrefactive bacteria. Oxygen, under a pressure of five or six atmospheres, is stated to stop their growth. Other observers have, however, obtained different results.

Effect of Gases.—Hydrogen and carbonic acid are stated to stop the movements of the motile bacteria. Chloroform is believed to arrest the changes brought about by the zymogenic species.

Electricity.—Cohn and Mendelsohn found that a constant galvanic current produced a deleterious effect owing to electrolysis. At the positive pole the liquid became distinctly acid, and at the negative pole distinctly alkaline. With a weak current there appeared to be no effect, two powerful cells at the very least being necessary.

Light.—Downes has shown that sunlight is fatal to putrefactive bacteria. This is believed to be due to a process of induced hyper-oxidation, from which living organisms ordinarily are shielded by protective developments of the cell-wall, or of colouring matter, which cut off injurious rays. Duclaux has investigated the same subject, and observed that micrococci were more sensitive to sunlight than the spore-bearing bacilli. Engelmann has described a bacterium whose movements cease in the dark, and Zopf states that in his cultures of *Beggiatoa roseo-persicina* the growth was much more strongly developed on the side of the vessel facing the light.


Chemical Reagents.—Many substances, such as carbolic acid, corrosive sublimate, chlorine, bromine, etc., have a marked effect upon the growth of bacteria. This will be more fully described in another chapter. In several cases the bacteria themselves secrete a substance which is injurious to their future development.

Products of Growth. — Bacteria may be grouped together according to the changes pro-


duced in the media in which they grow. Thus we have pigment-forming, fermentative, putrefactive, and pathogenic bacteria.

Chromogenic or pigment-forming bacteria elaborate during their growth definite colour stuffs. Such species are exemplified by *Bacillus ianthinus*, which produces a striking purple growth; *Bacillus pyocyaneus*, which secretes pyocyanin, a substance which has been isolated and obtained in a crystalline form; *Micrococcus prodigiosus*, which produces a pigment allied to fuchsine; *Beggiatoa roseo-persicina*, which is characterised by the presence of bacterio-purpurin; *Sarcina lutea*, *Bacillus cyanogenus*, and many others.

Zymogenic or ferment bacteria produce their changes in non-nitrogenised media. *Bacterium aceti*, by its growth produces the acetic fermentation in wine by which alcohol taking up atmospheric oxygen is converted into vinegar—




The fermentation of urine, by which urea is converted into carbonate of ammonia, can be brought about by several micro-organisms, but notably by the *Bacterium ureæ*. The change produced is represented by the following formula:—




Clostridium butyricum converts the salts of lactic acid into butyric acid, producing the butyric


fermentation in solutions of starch, dextrine, and sugar. These bacteria are agents in the ripening of cheese, and the production of sauerkraut. Thus, in a solution neutralised with calcium carbonate :—



In the so-called viscous fermentation the *Streptococcus viscosus* produces a gummy substance in wines. According to Pasteur, the change may be thus represented :—



And as another example may be mentioned the *Bacillus acidi lactici*, through whose agency sugar of milk is converted into lactic acid :—



Saprogenic or putrefactive bacteria play a most important rôle in the economy of nature. They produce changes allied to fermentation in complex organic substances. The nitrification of soil has been attributed to their agency. Their action on proteids, according to Hoppe-Seyler, may be compared to digestion; bodies like peptones are first produced, then leucin, tyrosin, and fatty acids; lastly indol, phenol, sulphuretted hydrogen, ammonia, carbonic acid, and water. They abstract the elements they require, and the remainder enter into new combinations. Associated with the formation of these substances are certain bodies, which have a poisonous effect when introduced into

animals. These poisonous alkaloids, *ptomaines*, produce a septic poisoning, which must be distinguished from septic infection. The effects of septic poisoning depend on the dose, whereas the effects of septic infection are, to a certain extent, independent of the dose. A small quantity of a septic poison may produce only transient effects, and a relatively large quantity may be necessary to produce vomiting, rigors, and death. Septic infection, on the other hand, may result equally from a small dose, because the poison introduced is a living organism which is capable of propagation and multiplication. Our knowledge of these alkaloids is greatly attributable to the researches of Selmi, Gautier, and also Brieger and others. Brieger has isolated ptomaines from the human cadaver, putrid meat, fish, and cheese. These substances—cadaverin, putrescin, saprin, peptotoxin, and many others—vary in their toxic properties.

Pathogenic bacteria are those which are genetically related to disease. Many organisms have been supposed to be pathogenic, or have been described in connection with diseases, which are only saprophytic associates. By the latter we mean organisms which feed upon dead organic matter. Such are many forms which are found on the skin, in the intestinal canal, and sometimes in the internal organs, especially the liver and kidneys; the tissues have lost their vitality, and the organisms, through some lesion, have been carried into the circulation.

That many organisms are causally related to disease, there is strong evidence in proof; for no organism can be considered to be productive of disease unless it fulfils the conditions which have been laid down by Koch (p. 17). Great stress must be laid upon the importance of successive cultivation through many generations, as the objection that a chemical virus may be carried over from the original source is thus overcome. Any hypothetical chemical poison carried over from one tube to another would, after a great number of such cultivations, be diluted to such an immense extent as to be inappreciable and absolutely inert.

Though we may accept as a fact the existence of pathogenic organisms, we are not yet in a position to assert the means by which they produce their deleterious or fatal effects. Many theories have been propounded. It has been suggested that the organisms—micrococci, for example—may be compared to an invading army. The phagocytes arrayed against them endeavour to assimilate and destroy them, but perish themselves in the attempt. This might explain the breaking down of tissue, and the formation of local lesions, but does not assist us in understanding the fatal result in thirty-six to forty-eight hours produced by the inoculation of the bacilli of anthrax. Another view is that the invading army seizes upon the commissariat, appropriating the general pabulum, which

is so essential to the life of the tissues. But this would hardly account for so acute and fatal a result as in anthrax, but would lead one to expect symptoms of inanition and gradual exhaustion. Moreover, against this theory we have the fact that death may result, for example, from anthrax, with the occasional presence of comparatively few bacilli; and, again, the blood may teem with parasites such as the flagellated monads in well-nourished, healthy-looking rats, without apparently causing any symptoms whatever. In the same category may be placed the theory that eminently ærobic organisms seize upon the oxygen of the blood and produce death by asphyxia. Another explanation is afforded by the suggestion of interference with the functions of the lung and kidney by mechanical blocking of the capillaries. Here the same objection is met with in the case of anthrax, the same fatal result may occur with only a few bacilli, while other cases yield very beautiful sections, looking like injected preparations from the mapping out of the capillaries with the countless crowds of bacilli (Plates XXIII. and XXIV).

The most satisfactory explanation is probably afforded by analogy with the putrefactive bacteria. We have seen that they derive their necessary elements from complex organic substances, and accompanying the residue we find the presence of poisonous alkaloids. Do pathogenic bacteria act in the same way? Does the anthrax bacillus

produce a ptomaine *anthracin*, which in a certain dose produces death, independently of the number of bacilli, provided there are sufficient present to develop that dose? Though this is possible, observers as yet have failed to extract from cultivations of the anthrax bacillus any alkaloid with virulent properties.

Lastly, it has been suggested that possibly a special ferment is secreted by the organisms, and that by the changes ultimately wrought by the action of this ferment, the symptoms and phenomena of disease arise. We have an analogy with this theory in the alkaline fermentation of urine by means of the *Torula ureæ*. By the researches of Musculus, and later of Sheridan Lea, it has been shown that a ferment is secreted by the cells which can be isolated in aqueous solution, and is capable of rapidly inducing an active fermentation of urea. Either of the two last theories assists us in understanding how it is that in anthrax or in tuberculosis we may find the presence of only a few bacilli, or that, assuming both tetanus and hydrophobia to be due to microbes, we can have such a violent disturbance of the system produced by the presence of very few micro-organisms. We may conceive that different species of bacilli may vary greatly in their power of producing an alkaloid or secreting a ferment, just as the elaboration of pigment is much more marked in some species than in others; thus it need not follow that the number of

micro-organisms bears any relation to the virulence or activity of the substance they produce. There is, however, yet another factor in the production of disease. We know that in health we are proof against most of these micro-organisms; if it were not so, we should all rapidly fall victims to the tubercle bacillus or some others, which we in health inhale with impunity. We know that a microbe may only cause a local lesion in one animal, and death in another. It is still more striking that the same micro-organism, as is the case with anthrax, may have no effect whatever upon certain species of animals, though it is deadly to others. Again, an animal naturally susceptible to the effect of a pathogenic organism may be rendered proof against it. These matters will be discussed in a future chapter.

Distribution of Bacteria.—Bacteria are commonly described as ubiquitous. They are ever present in the air, though not in such exaggerated numbers as is commonly supposed. In nutrient media exposed to the air one is often astonished at times at the comparatively few bacteria which develop in comparison to the amount of floating matter, such as mineral particles, scales, spores of fungi, and *débris* known to be present. In water they are also present in considerable numbers, though of course varying according to the character of the water. Wherever there is putrefaction, they are present in vast numbers. In the soil, in

sewage, in the intestines ; and in uncleanly persons especially, on the skin and between the teeth, various species may always be found, but in the healthy blood and healthy tissues bacteria are never present. In a previous chapter the method of examining the blood of living persons has been described, and there is, by this means, ample opportunity for satisfying oneself that bacteria are never to be found in the blood in health. The organs removed from a perfectly healthy animal, with the necessary precautions, into sterilised media can be kept indefinitely without undergoing putrefaction, or giving any development of bacteria. This has been established by many observers, notably Cheyne and Hauser ; and the results of former observers to the contrary must be attributed to imperfect methods admitting of accidental contamination.

CHAPTER IX.

ANTISEPTICS AND DISINFECTANTS.

IN the previous chapter several conditions were alluded to which affected the growth of bacteria, such as the nature of the nutrient soil, temperature, light, and electricity. The effect of certain chemical substances, and of excessive heat and cold, was also mentioned; but this constitutes a subject of such practical importance that it must be considered more fully.

Agents which retard the growth of bacteria are generally spoken of as *antiseptics*, as distinguished from *disinfectants*, which altogether destroy their vitality.

Though chemical disinfectants, or germicides, when diluted, act as efficient antiseptics, the converse, that an antiseptic in a sufficiently concentrated form will act as a disinfectant, is not the case. The term "antiseptic," indeed, should be restricted to those substances or agents which arrest the changes bacteria produce, but which do not prevent their springing into activity when removed to favourable conditions. Thus excessive heat, which destroys

bacteria and their spores, is a true disinfectant ; and excessive cold, which only benumbs them, retarding their development without killing them, is an antiseptic.

Spores have a greater power of resisting the action of these various agents than the parent cells, and many species of micro-organisms differ from each other in their resisting power. An exact knowledge of the subject can, therefore, only be based upon investigations which will determine the effect of these agents upon pure cultivations of the different micro-organisms causally related to putrefaction and disease. In the latter case, especially, this is not possible in the present state of our knowledge. In some cases of communicable disease there is considerable doubt as to the etiological importance of the organisms which have been described ; in other cases no organisms have as yet been discovered, or the organisms cannot be artificially cultivated, or the disease is not reproduced by inoculation, so that there is no means of testing whether the agents have had any effect. One can, therefore, only draw general conclusions by selecting some well-known pathogenic and non-pathogenic micro-organisms, and considering the influence of chemicals, of hot air, and of steam upon them, as representing the effect upon the various contagia of disease and the causes of putrefaction.

Such knowledge must necessarily prove of the greatest importance,—to the sanitarian, who is

concerned in preventing the spreading of disease and in the disposal of putrefactive matter,—to the surgeon, who is anxious to exclude micro-organisms during surgical operations and to arrest the development in wounds of bacteria which have already gained an entrance,—to the physician in the treatment of micro-parasitic diseases. The sanitarian and the surgeon must profit directly by such experiments, for in the disinfection of clothes and the sick-room by the one, and in the application of antiseptic dressings and lotions by the other, the micro-organisms are encountered as in the test experiments apart from the living body. The physician, on the other hand, is principally concerned in dealing with micro-parasites when circulating in the blood, or carrying on their destructive processes in the internal tissues. So far as our knowledge at present goes, the physician can avail himself but little of the effect of the direct application of the substances which have been found to retard or destroy the growth of the organisms in artificial cultivations, for the concentrated form in which they would have to be administered would prove as deleterious or as fatal to the host as to the parasites. Thus Koch has stated that to check the growth of the anthrax bacillus in man it would be necessary that there should be twelve grammes of iodine constantly in circulation; and that the dose of quinine necessary to destroy the spirilla of relapsing fever would be

from twelve to sixteen grammes. The retarding influence, however, of certain substances when diluted, and the fact that disinfectants are sometimes equally efficacious in a diluted form when their application is prolonged, seem to indicate measures which may be adopted, in some cases, with chances of success, such as the inhalation of antiseptic vapours in phthisis. For the most part the physician must look rather to combating the effects of micro-organisms by restoring to its normal standard the lowered vitality which enabled the bacteria to get a footing.

There is no wider field for research than the determination of the real effect of disinfectants and antiseptics. Painstaking and laborious as the researches are which have been hitherto made, the subject is so beset with fallacies, leading, in some cases, to totally erroneous conclusions, that it is not surprising that one meets on all sides with conflicting statements. The author has no intention of analysing these results, but a general idea will be given of the methods which have been employed, and for further details reference must be made to the original papers mentioned in the bibliography.

Chemical Substances.—It was customary to judge of the power of a disinfectant or antiseptic by adding it to some putrescent liquid. A small portion of the latter was, after a time, transferred to some suitable nourishing medium, and the efficacy

of the substance estimated by the absence of cloudiness, odour, or other sign of development of bacteria in the inoculated fluid. Koch pointed out the errors that might arise in these experiments from accidental contamination, or from there being no evidence of the destruction of spores; and we are indebted to him for a complete and careful series of observations with more exact methods.

Instead of employing a mixture of bacteria, Koch's plan was to subject a pure cultivation of some well-known species with marked characteristics to the reagent to be tested. A small quantity was then transferred to fresh, nourishing soil, under favourable conditions, side by side with nutrient material inoculated from a cultivation without treatment with the disinfectant. The latter constituted a control test, which is most essential in all such experiments. To test the resistant power of bacteria which are easily destroyed, two species were selected, *micrococcus prodigiosus*, and the bacillus of blue pus. These were cultivated on potatoes, the surface of which was sliced off and dried. A fragment transferred to freshly prepared potato gave rise to a growth of the particular micro-organism; but if after treatment with some reagent no growth occurred, the conclusion was drawn that the agent was efficacious in destroying the vitality of the bacteria.

Anthrax bacilli in blood withdrawn from an animal just killed were taken to represent spore-

less bacteria, while silk threads steeped in an artificial cultivation of the bacilli and dried, afforded a means of testing the vitality of spores.

Even by employing pure cultivations on solid media, great precautions were necessary to avoid mistakes. If, for instance, a large quantity of the growth which had been subjected to some chemical solution were carried over to the fresh tube containing the nutrient medium, or if a silk thread, which had been dipped in a solution, were directly transferred to the new soil, enough of the supposed disinfectant might be mechanically carried over to retard the development of the bacteria, though it was ineffectual in destroying them. From a growth not appearing, the conclusion might be drawn that the spores or the bacteria had been affected, and so a mistake occurs. To avoid this Koch made a point of transferring a minimum of the disinfected growth to as large a cultivation area as possible, so that any chemical substance mechanically carried over, would be so diluted as to be inert. For the same reason threads, after withdrawal from the disinfecting solution, were rinsed in sterilised water, or weak alcohol, and then transplanted; or, instead of judging from the development on nutrient gelatine, the effect of inoculation in a healthy animal was made the test.

A few examples may be quoted in illustration. Silk threads, impregnated with anthrax spores, were placed in bottles containing carbolic acid of

various strengths. A thread was removed from each on successive days, and transferred to nutrient gelatine, and the result noted. It was found that immersion of the thread in a 5 per cent. solution of carbolic acid was sufficient in two days to effect complete sterilisation, and seven days in a 3 per cent. solution was equally efficacious. Since for practical purposes a strength should be selected which would be effectual in twenty-four hours, Koch recommended that for general use, allowing for deterioration by keeping, a solution containing not less than 5 per cent. should be employed, and for complex fluids probably a still higher percentage would be necessary. In the case of sporeless bacilli the results were very different. Blood, containing the bacilli, from an animal just killed, was dried on threads, and after exposure for two minutes to a 1 per cent. solution, was completely sterilised. Fresh blood mixed with a 1 per cent. carbolic solution produced no effect on inoculation. If, on the other hand, the blood was mixed with a .5 per cent. solution, the virulence was not destroyed. The facility with which the bacilli are destroyed, compared with their spores, illustrates how easily errors may occur, if mere arrest of growth or loss of motility be regarded as a sign of the efficacy of disinfection.

To test vapours, Koch exposed anthrax spores or the spores which occur in garden earth by suspending them over solutions, *e.g.*, of bromine or

chlorine in a closed vessel. After a time they were transferred to a nutrient medium to test their vitality. To test the power of sulphurous acid gas, the spores were spread about in a room in which the gas was generated by burning sulphur in the ordinary way for disinfecting a room. To test chemicals which might be recommended for disinfecting vans and railway carriages, spores were laid on boards which were then washed or sprayed, and the spores then transferred to the nutrient gelatine.

By such simple methods Koch investigated a long list of chemical reagents, and according to these experiments the salts of mercury, and the chloride especially, proved most valuable. Where heat is not admissible, these compounds were therefore highly recommended, though their poisonous nature is a drawback to their indiscriminate use. Koch states, for disinfecting a ship's bilge, where a 5 per cent. solution of carbolic acid must be left for forty-eight hours, a 1 in 1000 solution of mercuric chloride would only require a few minutes.

There is, on the other hand, reason for doubting the efficacy of very dilute solutions; for, though anthrax spores subjected to a 1 in 20,000 solution of mercuric chloride for ten minutes, and then washed in alcohol, gave no growth in nutrient gelatine, silk threads exposed for ten minutes to a 1 in 20,000 solution, or even 1 in 10,000, still proved fatal to mice.

Herroun considers that the value of mercuric chloride as an antiseptic is much over-rated, as he has cultivated ordinary septic bacteria in albuminous filtrates, containing 1 in 2000. It is precipitated by albumins if used of greater strength, and is readily converted by the sulphur of albuminous bodies into mercuric sulphide,—a compound which has practically no antiseptic properties.

Sternberg has also made an elaborate series of experiments with regard to the action of germicides. In this case cultivations of well-known pathogenic organisms in liquid media were employed. The supposed germicide was added to the liquid cultivation, and after two hours a fresh flask of sterilised culture was inoculated from the disinfected cultivation, and placed in the incubator. In twenty-four to forty-eight hours, if the chemical were not efficient, there was evidence of a growth of bacteria. Blyth has investigated the disinfection of cultivations of *Bacterium termo*, of sewage, and typhoid excreta, and, in conjunction with Klein, the effect of well-known disinfectant materials on anthrax spores. Miquel, Laws, and others have also contributed to our knowledge of the effect of antiseptics and disinfectants upon micro-organisms. In spite of all that has been done, there is room for many workers; a great deal of ground must be gone over again to rectify discrepancies, examine conflicting results, and thus determine what

observations may be relied upon for practical application.

Hot Air and Steam.—Koch, in conjunction with Wolfhügel, also tested the value of hot air. A similar plan was adopted as in disinfection with chemicals. Bacteria and spores were subjected for a certain time to a known temperature in a hot-air chamber, and then were transferred to a nourishing soil, or animals were inoculated.

Paper parcels, blankets, bags, and pillows, containing samples of micro-organisms wrapped up inside, were also placed in the hot-air chamber, to test the power of penetration of heat.

The conclusions from such experiments were as follows:—

Sporeless micro-organisms at a little over 100° C. are destroyed in one hour and a half.

Spores of bacilli require three hours at 140° C.

If enclosed in pillows and blankets, exposure from three to four hours to 140° C. is necessary.

Spores of fungi require one and a half hours at 110° C. to 115° C.

Further experiments showed that at the temperature necessary for the destruction of spores of bacilli almost all fabrics are more or less injured.

Koch, in conjunction with Gaffky and Löffler, also investigated the effect of steam under pressure and at the atmospheric pressure.

Rolls of flannel with anthrax spores or earth spores, and a thermometer wrapped up inside, were

subjected to steam, and the results compared with the effect obtained with hot air.

Thus in hot air four hours' exposure to a temperature of 130° C. to 140° C. brought the temperature inside the roll to 85° C., and the spores were not injured; on the other hand, exposure to steam under pressure at 120° C. for one and a half hours, raised the internal temperature to 117° C. and killed the spores.

By such experiments the superior penetrative power of steam-heat was established.

To test steam-heat at the atmospheric pressure, water was boiled in a glass flask with its neck prolonged by means of a glass tube, the temperature in which was found to be uniform throughout. Anthrax and earth spores placed in the tube were found to be unable to withstand steam at 100° C. even for a few minutes. It was, therefore, concluded that disinfection by steam at atmospheric pressure was superior to hot air from its greater efficiency, and to steam under pressure from the simplicity of the necessary apparatus.

Parsons and Klein made some experiments which were more in favour of dry heat than the above. These observers state that anthrax bacilli are destroyed by an exposure of five minutes to from 100° C. to 103° C., and that anthrax spores are destroyed in four hours at 104° C., or in one hour at 118° C. Guinea-pigs inoculated with tuberculous pus which had been exposed for five minutes to 104° C.,

remained unaffected. They concluded that as none of the infectious diseases, for which disinfecting measures are in practice commonly applied, are known to depend upon the presence of bacilli in a spore-bearing condition, their contagia are not likely to retain their activity after being heated for an hour to 105° C. (220° F.).

In experiments with steam, the results were in accordance with those already given, and complete penetration of an object by steam-heat for more than five minutes was deemed sufficient. They also arrived at the same result as in Koch's experiments, that steam-chambers are preferable to those in which dry heat is employed, though it must be borne in mind that some articles, such as leather, are injured by exposure to steam.

CHAPTER X.

IMMUNITY.

THE condition of being insusceptible to an infective disease may be either natural or acquired. In the description of the pathogenic organisms several examples of natural immunity will be encountered. The bacillus of septicæmia, so fatal to house mice, has been shown to have no effect upon field mice. The bacillus of anthrax is innocuous to cats, white rats, and, it is commonly asserted, to adult dogs and asses. The bacterium of rabbit septicæmia is equally inert in dogs, rats, and guinea-pigs. The immunity may be as in these cases complete, or only partial. Ordinary sheep are very easily affected with anthrax, but Algerian sheep only succumb to large doses of the virus. Natural immunity may not only be characteristic of certain species, but it may occur in certain individuals of a susceptible species. The same occurs in man, for certain individuals, though equally exposed during an epidemic of small-pox, may escape where others readily fall victims to the disease.

Acquired immunity is illustrated by the protection afforded by one attack of the exanthemata against subsequent attacks. Thus one attack of measles or small-pox, as a rule, affords complete protection. A knowledge of the immunity resulting in the latter case led to the introduction of inoculation of small-pox as a protection against natural small-pox.

Immunity may be acquired by acclimatization, for the inhabitants of tropical climates are less susceptible to the diseases of the country, malarial fevers for instance, than strangers.

In civilised communities also there appears to be a degree of acquired immunity, for the infectious diseases introduced among savages or isolated communities have assumed the most virulent properties.

The immunity acquired by protective inoculation constitutes, in connection with the study of pathogenic micro-organisms, a subject of pre-eminent interest and importance. Pasteur, in his researches upon fowl-cholera, observed that after non-fatal cases the disease either did not recur, or the severity of a subsequent attack was in inverse proportion to the severity of the first attack. It occurred to him to endeavour to obtain the virus of this disease in a form which would provoke a mild attack of the disease, and thus give protection against the virulent form. This attenuation or mitigation of the virus was

successfully attained in the following manner: Cultivations of the microbe, in chicken-broth, were allowed to remain with a lapse of several months between the carrying on of successive cultivations in fresh media. The new generations which were then obtained were found to have diminished in virulence, and ultimately a virus was obtained which produced only a slight disorder; on recovery the animal was found to be proof against inoculation with virulent matter. The explanation given by Pasteur of this change was, that prolonged contact with the oxygen of the air was the influence which diminished the virulence, and he endeavoured to prove this by showing that if broth were inoculated in tubes which could be sealed up, so that only a small quantity of air was accessible to the microbe, the virulence of the cultures was retained.

Toussaint investigated the possibility of attenuating the virus of anthrax. Sheep injected with 3 ccm. of defibrinated blood, containing anthrax bacilli, which had been exposed to 55° C. for ten minutes, recovered, and were afterwards insusceptible. Pasteur subsequently argued that this method did not admit of practical application; difficulties would arise in dealing with infective blood in quantity, and artificial cultivations started from this blood could not be relied upon, as they proved sometimes as virulent as ever.

Pasteur endeavoured to apply the same method

for obtaining an attenuated virus of anthrax, as he had successfully employed in chicken-cholera. A difficulty was soon encountered, for in cultivations of the bacillus with free access of air spore-formation readily takes place, and the spores are well known to have an extraordinary power of retaining their virulence. Pasteur found that the bacilli ceased to develop at 45°C ., and he believed that spore-formation ceased at 42° — 43°C ., the bacilli continuing to develop by fission only. The cultivations were, therefore, kept at this temperature, and at the end of eight days the bacilli were found to have lost their virulence, and were quite inert when inoculated in guinea-pigs, sheep, or rabbits. This total destruction was, however, preceded by a gradual mitigation, so that a virus could be obtained, by taking it at the right time, which only gave a mild disease, and afforded subsequent protection.

At Melun, in 1881, the protective inoculation against anthrax was put to a practical test. Sheep and oxen were inoculated with the mitigated virus, and then with a virulent form; at the same time other sheep and oxen were inoculated with the virulent form without previous vaccination as a control experiment. The unprotected sheep died without exception; the unprotected oxen suffered from œdematous swellings at the seat of inoculation, and a rise of temperature; but all the protected animals remained healthy.

As a result of these experiments an idea arose that by preventive inoculation with attenuated virus all communicable diseases would in time be eradicated; but this does not follow, for all communicable diseases do not confer immunity after a first attack, and in some cases the very reverse is believed to occur. Thus erysipelas of the face leads to an increased liability to subsequent attacks of the same disease. Again, the occurrence of one disease is stated to induce a liability to others; small-pox and typhoid fever are regarded as predisposing to tuberculosis; so that the principle of preventive inoculation does not apply in these cases, and its effect would probably tend rather to deleterious results than otherwise. Even with regard to the prevention of anthrax, Pasteur's researches were opposed and criticised. Koch investigated the subject, and came to the conclusion that the process did not admit of practical application, chiefly on the ground that as immunity only lasted a year, the losses from the vaccination process would be as great or even greater than from the spontaneous disease; further, there was danger in disseminating a vaccine of the strength required to be effectual. Chauveau proved that the attenuation was due to the temperature, and not to the prolonged effect of oxygen. By keeping cultivations at 42° — 43° C. *in vacuo*, the virulence was found to disappear in twenty-four hours, and by keeping cultivations at a low temperature with free

access of air the virulence was retained. Chauveau considered, therefore, not only that oxygen was not the agent, but that the mitigation was much more easily effected in its absence. In spite of these adverse criticisms, these researches nevertheless confirmed the principle of Pasteur's conclusion, that immunity could be induced by experimental measures, and further showed that he had considerably advanced the method by which this could be effected.

Chauveau succeeded also in attenuating the virus by a modification of Toussaint's method. Sterilised broth was inoculated with the bacilli, and placed in the incubator at 42°—43° C. After the lapse of twenty hours it was removed to another incubator at 47° C. According to the time of exposure to this increased temperature, the mitigation varied in degree. Thus inoculation with the virus, before it was exposed to 47° C., was fatal to guinea-pigs; but after one hour at 47° C. the virulence was diminished, and, though ultimately fatal, life was prolonged; after two hours' exposure at 47° C. only half the animals died; and after three hours' exposure they recovered and were rendered refractory to subsequent inoculation.

Attenuation of the virus has also been induced by chemical means. Chamberland and Roux stated that a fresh growth started from a cultivation of bacilli which had been subjected for twenty-nine days to $\frac{1}{500}$ of carbolic acid was found to

be inert in guinea-pigs and rabbits. Bichromate of potash added to a cultivation in the proportion of $\frac{1}{12000}$ — $\frac{1}{5000}$ gave, after three days, a new growth, which killed rabbits, guinea-pigs, and half the sheep inoculated; after ten days, rabbits and guinea-pigs, but not sheep; and after a longer time even guinea-pigs were unaffected.

In other diseases similar results have been obtained.

Arloing, Cornevin, and Thomas found that by inoculating a small quantity of the virus of symptomatic anthrax anywhere in the subcutaneous connective tissue, or a moderate quantity at the root of the tail, and even by intravenous injection, immunity was obtained from a virulent dose.

In swine-erysipelas, Pasteur and Thuillier obtained attenuated virus upon quite another principle. They discovered that by passing the virus through pigeons the virulence was increased, but by passing it through rabbits it was progressively diminished. Thus a virus was obtained from the rabbit, which produced only a mild disease in pigs, and after recovery complete immunity. Similarly in rabies Pasteur finds that passage of the virus through various animals considerably modifies its properties. By inoculating a monkey from a rabid dog, and then passing the virus through other monkeys, the virulence is diminished; but by inoculating a rabbit from the dog, and passing the virus from rabbit to rabbit, the virulence is increased. More recently

Pasteur has employed another method of attenuating the virus of rabies. The spinal cord of inoculated rabbits is removed with all possible precautions, and portions a few centimetres in length are suspended in flasks in which the air is dried by fragments of potash. By this process the virulence is found to gradually diminish and finally disappear.

Animals inoculated with portions of these cords, after suspension for a certain time, are rendered refractory to inoculation with virulent cords. Having rendered dogs, which had been previously bitten, free from the supervention of symptoms of hydrophobia by means of protective inoculation, Pasteur proceeded to apply the same treatment to persons bitten by rabid animals, with results which tend to the belief that a prophylactic for rabies has been found, though this must still be considered to be *sub judice*.

The question as to what constitutes immunity is a vexed one.

Raulin has shown that *Aspergillus niger* develops a substance which is prejudicial to its own growth in the absence of iron salts in the nutrient soil. Pasteur has suggested that in rabies side by side with the living and organised substance there is some other substance which has, as in Raulin's experiment, the power of arresting the growth of the first substance. If we accept the theory of arrest by some chemical substance, we must suppose that in the acquired immunity afforded by

one attack of an infectious disease this chemical substance is secreted, and, remaining in the system, opposes the onset of the micro-organism at a future time. In the natural immunity of certain species and individuals we must suppose that this chemical substance is normally present.

Another theory is, that the micro-organisms assimilate the elements which they require for their nutrition from the blood and tissues, and render the soil impoverished or otherwise unsuitable for the development of the same micro-organisms hereafter; this condition may be permanent, or the chemical constitution of the tissues may be restored to normal, when immunity ceases. If, however, we explain acquired immunity by the result of the growth of a previous invasion of micro-organisms, we are still confronted with the difficulty of explaining natural immunity.

A third theory is that the tissues are endowed with some power of vital resistance to the development of micro-organisms, similar to the vital resistance to the coagulation of the blood, which is supposed to exist in the lining membrane of the healthy blood-vessel; that in some species and individuals this exists to a high degree, and hence their natural immunity. But this does not explain how one attack confers immunity from a subsequent one. One would expect that the vital resistance would invariably be lowered by a previous attack, and increased liability be the constant result.

Lastly, that leucocytes appear to have the power of destroying bacteria in some cases, has been demonstrated by the researches of Metschnikoff.

If anthrax bacilli are inoculated in the frog, the white blood-cells are observed to incorporate and destroy them until they entirely disappear, and the animal is not affected. But if the animal, after inoculation, is kept at a high temperature, the bacilli increase so rapidly that they gain the upper hand over the leucocytes, and the animal succumbs.

In septicæmia of mice the white blood-cells are attacked and disintegrated by the bacilli in a similar way. It is difficult, however, to accept any explanation of immunity from these observations, —to suppose, for example, that immunity depends upon the micro-organisms being unable to cope with the leucocytes in certain species. It is difficult to conceive that the leucocytes in the blood and tissues in the field mouse are differently constituted from those in the house mouse, so that they form an effectual barrier in the one case, though so readily destroyed in the other.

PART III.

SYSTEMATIC AND DESCRIPTIVE, WITH SPECIAL
MICROSCOPICAL METHODS.

PART III

SYNOPSIS AND DESCRIPTIONS OF THE SPECIES
MICROSCOPICALLY

CHAPTER XI.

CLASSIFICATION OF BACTERIA.

IN reviewing the history of the various classifications which have from time to time been proposed, we shall see that the gradual improvements in the means of studying such minute objects, the methods of cultivating them artificially, and of studying their chemistry and physiology, and the ever-increasing revelations of the microscope, have resulted in establishing these microscopic objects as members of the vegetable kingdom, ranking among the lowest forms of fungi. While enabling us to settle their position as a whole, these improved methods have further given us so great an insight into the life-history of individual forms, that, with regard to the division into genera and species, we are, up to the present time, still in a position of doubt and uncertainty.

Müller, in 1773, was the first to suggest a classification. He established two genera, *Monas* and *Vibrio*, and grouped them with the *Infusoria*. In

1824 Bory de Saint Vincent also attempted a classification; but it was not until Ehrenberg in 1838, and Dujardin in 1841, worked at the subject, that a scientific distinction of species was attempted.

Ehrenberg described four genera :—

- | | |
|----------------------|-----------------------------------|
| I. Bacterium . . . | . filaments straight, rigid. |
| II. Vibrio . . . | . filaments snake-like, flexible. |
| III. Spirillum . . . | . filaments spiral, rigid. |
| IV. Spirochæte . . . | . filaments spiral, flexible. |

Dujardin united *Spirillum* and *Spirochæte*, and classed them thus :—

- | | |
|----------------------|-----------------------------------|
| I. Bacterium . . . | . filaments rigid, vacillating. |
| II. Vibrio . . . | . filaments flexible, undulatory. |
| III. Spirillum . . . | . filaments spiral, rotatory. |

Up to that time, bacteria were still considered as *Infusoria*; but the year 1853 marked the commencement of a new era in their history, for Robin then pointed out the affinity of the *Bacteria* and *Vibrios* to *Leptothrix*. Davaine, in 1859, still more definitely insisted that the *Vibrios* were vegetables, and that they were in fact allied to the *Algæ*.

Since that time a flood of light has poured in upon this subject through the writings of Hoffmann, Pasteur, Cohn, Rabenhorst, Hallier, Billroth, Warming, Nägeli, Magnin, Marchand, Sternberg, Van Tieghem, Koch, Flügge, De Bary, Zopf,

Cornil, Babes, and many other workers in the recent widespread revival of bacteriological research.

Of all these writers we are most indebted to Cohn,* not only on account of his researches, which extended over very many years, but also for his system of classification, which has since been almost universally adopted.

In his first classification, published in 1872, Cohn considered the Bacteria as a distinct group belonging to the *Algæ*, and divisible into four tribes, including six genera :—

- I. Sphærobacteria . globules (Micrococcus).
- II. Microbacteria . short rods (Bacterium).
- III. Desmobacteria . long rods (Bacillus and Vibrio).
- IV. Spirobacteria . spirals (Spirochæte and Spirillum).

Cohn noted, in spite of placing them with the *Algæ*, that the absence of chlorophyll connected the *Bacteria* to *Fungi*, and we find Nägeli subsequently adopting this view, and employing the term *Schizomycetes*.

Billroth, in 1874, disputed the division into species, and considered that all the forms described by Cohn were but developmental forms of one micro-organism, *Coccobacteria septica*. In the following year Cohn answered the criticism of Billroth, and produced a second classification, in which he still

* Cohn, *Beiträge zur Biologie der Pflanzen*, 1872, *et seq.*

maintained that distinct genera and species existed. The genera Cohn considered to be distinguished by definite differences in shape, which were adhered to throughout life, while some special feature, as a difference in size or physiological action, or some minute difference in form, determined the various species. Cohn illustrated, by his well-known comparison of a sweet and a bitter almond, the appearances of which are similar, but the properties very different, that a distinction into species might depend upon a difference in physiological action only. Others strongly support Cohn's views. By cultivating various micro-organisms through several generations, we conclude that a micrococcus cannot be transformed into a bacterium, or a bacterium into a bacillus or spirillum. There is also no evidence to show that a bacillus can change its nature, and be converted from a harmless into a pathogenic form, as asserted by Büchner.*

The second classification of Cohn (1875) only differed from the first in that, instead of keeping the bacteria as a separate group, he placed them, from their closer relationship with the *Fhycocchro-maceæ*, under a new group, the *Schizophytes*, and added the genera *Leptothrix*, *Beggiatoa*, *Crenothrix*, *Sarcina*, *Ascococcus*, *Streptococcus*, *Myconostoc*, and *Streptothrix*.

Nägeli maintained that *Bacteria* were allied to

* Büchner, *Ueber d. experim. Erzeugung d. Milzbrandcontagiums aus d. Heupilzen.*

Yeasts, and should be included in the class of *Fungi*. In fact, he divided the *fungi-producing decomposition* into :—

- Mucorini moulds
- Saccharomycetes yeasts
- Schizomycetes fission-fungi


(This last class comprising bacteria.)

Flügge,* following Rabenhorst, maintained the term *Schizomycetes*, and divided them as shown in the table on the following pages :—

* Flügge, *Fermente und Mikroparasiten*. 1883.

SCHIZOMYCETES (FLÜGGE'S ORIGINAL CLASSIFICATION).

<p>Cells round or ovoid.</p> <p>Isolated, or in chains, or united in amorphous gelatinous families</p>	<p><i>Micrococcus.</i></p>
<p>Forming gelatinous fami- lies of definite form.</p>	<p>In large numbers, in irregular colo- nies</p> <p><i>Ascococcus.</i></p>
<p>Colonies solid, filled with cells</p>	<p>In small but definite numbers, in regular groups</p> <p><i>Sarcina.</i></p>
<p>Colonies, with simple layer of cells at the periphery</p>	<p><i>Clathrocystis.</i></p>



The belief is nevertheless rapidly gaining ground that the lowest forms of vegetable life cannot be divided by a hard-and-fast line into a series with chlorophyll (*Algæ*), and a series without it (*Fungi*), and the tendency now is to solve the difference of opinion between Cohn and Nägeli by following the example of Sachs, and amalgamating the two series into one group, the *Thallophytes*.

Researches by competent observers have quite recently clearly demonstrated that several micro-organisms in their life cycle exhibit successively the shapes characteristic of the orders of Cohn.

This doctrine of *pleomorphism*, now widely accepted, was distinctly foreshadowed in a publication* by Lister in 1873, though this memoir contained certain conclusions which have since been abandoned. Lister described forms of cocci, bacteria, bacilli, and streptothrix in milk, which he regarded as phases of the same micro-organism, *Bacterium lactis*. As a result of his observations, Lister remarks that "any classification of bacteria hitherto made from that of Ehrenberg to that of Cohn based upon absolute morphological characters is entirely untrustworthy." To Lankester, however, belongs the credit of having definitely and precisely formulated this doctrine. In a paper,† also published in 1873, Lankester observed that the series of form-phases which he had discovered in the case of a

* Lister, *Quart. Journ. Microscop. Sci.*, 1873, pp. 380-408.

† Lankester, *Ibid.*, pp. 408-425, and 1876, pp. 278-283.

peach-coloured bacterium led him to suppose that the natural species of these plants were "within the proper limits protean, and that the existence of true species of bacteria must be characterised, not by the simple form-features used by Cohn, but by the ensemble of their morphological and physiological properties as exhibited in their complete life-histories." Lankester inferred that these phase-forms were genetically connected from their all possessing the common characteristic of a special pigment bacterio-purpurin. These conclusions were vigorously opposed by Cohn, and doubt still remains in the minds of some as to whether the different forms are really only stages in the life-history of a single species. Nevertheless the theory of pleomorphism has steadily gained ground ever since.

Among the recent observers Cienkowski and Neelsen have worked out the different forms assumed by the *bacillus of blue milk*; Zopf has in a similar manner investigated *Cladothrix*, *Beggiatoa* and *Crenothrix*, and traced out various forms (Fig. 67); Van Tieghem has investigated *Bacillus amylobacter* with a similar result; Hauser has quite recently described bacillar, spirillar, and spirular, and various other forms in the *Proteus mirabilis* and *Proteus vulgaris*. These facts obviously shake the very foundation of Cohn's classification, and we are left without possessing a sound basis for classification into genera or species. The mode of




FIG. 67.

CLADOTRICH DICHOTOMA.—A Branched Schizomycete: (a) Vibrio-form (b) Spirillum-form [slightly magnified]. B. Screw-form at the ends: (a) Spirillum-form; (b) Vibrio-form. C. Very long Spirochæta-form. D. Branch fragment, at one end Spirillum-form, at the other Vibrio-form. E. Screw-form: (a) Continuous; (b) Composed of rods, and (c) Cocci. F. Spirochæta-form: (a) Continuous; (b) Composed of long rods, (c) Short rods, and (d) Cocci [after Zopf].

reproduction is not sufficiently known to afford a better means for distinction than the other morphological appearances taken alone, nor can we depend upon physiological action, which is held by many to vary with the change of form, according to altered surroundings.

Zopf, who has warmly supported the pleomorphism of bacteria, has suggested as a result of his investigations a division of the *Schizomycetes*, *Spaltpilze*, or *Fission-fungi*, into the following four groups* :—

1. Coccaceæ.—Possessing (so far as our knowledge at present reaches) only cocci, and thread-forms resulting from the juxtaposition of cocci. The fission occurs in one or more directions.

Genera :—Streptococcus, Micrococcus, Merismopedia, Sarcina, Ascococcus.

2. Bacteriaceæ.—Possessing mostly cocci, rods (straight or bent), and thread-forms (straight or spiral). The first may be absent, and the last possess no distinction between base and apex.

Division (as far as is known) occurs only in one direction.

Genera :—Bacterium, Spirillum, Vibrio, Leuconostoc, Bacillus, Clostridium.

3. Leptotricheæ.—Possessing cocci, rods, and thread-forms (which show a distinction between base and apex). The last straight or spiral.

* Zopf, *Die Spaltpilze*, 1885.

Genera :—Leptothrix, Beggiatoa, Crenothrix, Phragmidiothrix.

4. **Cladotricheæ**.—Possessing cocci, rods, threads, and spirals. Thread-forms provided with false branchings.

Genus :—Cladotrix.

Zopf, however, does not assert that all the fission-fungi exhibit this pleomorphism, nor does he pretend that his classification will include all the micro-organisms described. Cohn, on the other hand, was ready to admit that all the forms described by him were not truly independent species. Quite recently De Bary, Hueppe, Baumgarten, and Flügge have expressed new views with regard to the classification of bacteria.

De Bary divides them into two great groups—bacteria which form endospores, and bacteria which form arthrospores. This affords but little practical assistance, though regarded by botanists, from a scientific standpoint, as a step in the right direction.

Hueppe, acknowledging that the fructification must eventually be made the basis for classification, suggests an arrangement for provisional use in which this view is introduced.

HUEPPE'S CLASSIFICATION.

Cocci forms .	Arranged in chains .	Zooglyea moderate .	Spores endogenous (?) .	<i>Streptococcus</i> (?) .	
		Zooglyea strongly marked .			Arthrospores or spores unknown
	Arranged in fours .	With small chains	<i>Merista</i> .
		Without small chains	<i>Sarcina</i> .
Arranged in eights .	Zooglyea indeterminate	<i>Micrococcus</i> .	
	In irregular heaps .	Zooglyea united in balls	<i>Ascococcus</i> .	
Rod forms .	Smaller or longer threads, without, distinction of base and apex. Threads flexible or rigid.	Threads straight or wavy, no endogenous spores	<i>Artho-bacterium</i> .	
		Threads straight, wavy, or screw form, no endogenous spores	<i>Spirulina</i> .	
	Threads with distinctions of base and apex	Threads straight or wavy, spores endogenous .	{ Without change of the cells	<i>Bacillus</i> .
		Threads without divisions .	{ With change of the cells, spindle rods	<i>Clostridium</i> .
	Screw-like threads, flexible or rigid .	Threads with divisions .	Without presence of sulphur	<i>Leptothrix</i> .
			With presence of sulphur	<i>Beggiatoa</i> .
	Screw rods .	Formation of arthrospores or fructification unknown .	Unbranched	<i>Crenothrix</i> .
			Branched	<i>Cladothrix</i> .
	Screw rods .	Endogenous spores	With change of form .	Without change of form	<i>Spirochæta</i> .
					<i>Vibrio</i> .
				<i>Spirillum</i> .	

It has already been mentioned that the production of arthrospores is only established in a very few species. Therefore, we are hardly justified in assuming that all bacteria, the spore-formation of which is quite unknown, are to be included with those in which this kind of fructification has been observed, and consequently to distinguish genera on the same grounds may be considered, to say the least, somewhat premature. In Baumgarten's classification the genus bacterium is dispensed with, and the genera divided into two groups, the monomorphic and the pleomorphic.

GROUP I.—MONOMORPHIC.

Genera.—Coccus.

Bacillus.

Spirillum.

GROUP II. PLEOMORPHIC.

Genera.—Spirulina.

Leptothrix.

Cladotrix.

Flügge also, in his recent classification, includes the genus bacterium in the genus bacillus. The new classification differs also from the original one in the grouping together of the different species according to the character and behaviour of the colonies in nutrient gelatine. The abolition, in Flügge's and Baumgarten's classification, of the genus bacterium is no doubt owing to confusion having arisen from the distinction between a

bacterium and a bacillus, being a question of length. Observers differed as to whether a rod of a certain length ought to be considered a bacterium or a bacillus. To meet this difficulty a rough-and-ready rule was suggested, viz., that a rod less than twice its breadth in length should be considered as a bacterium, and otherwise a bacillus. But this purely arbitrary division was inadequate, from the fact that a rod at one stage of its growth or under certain




FIG. 68.—*BACTERIUM PNEUMONIÆ CROUPOSÆ*, $\times 1500$ (after Zopf).

conditions might, as far as length went, truly be a bacterium, and under other circumstances be of such a length as to entitle its being considered a bacillus. We avoid such confusion if we follow Zopf, and acknowledge as a difference between a bacterium and a bacillus the presence or absence of that form of spore-formation now distinguished as endogenous spore-formation. We can then most conveniently retain this generic term, to include that group of rod-forms in which this spore-formation is as yet unknown; moreover, we shall find that by so

doing, with one or two exceptions, we get collected together those short rod-forms (Fig. 68), which appear to link the simple cocci to the spore-bearing rods or bacilli.

This must surely lead to less confusion than




FIG. 69.—EMMERICH'S BACTERIUM, $\times 700$ (after Emmerich).

regarding all rod-forms as bacilli, and massing them together into one genus. For by those who adopt the latter plan, not only are very short rods with rounded ends included as bacilli, e.g., *Bacillus*



FIG. 70.—COLONIES ON NUTRIENT GELATINE, $\times 60$.

Neapolitanus or Emmerich's bacterium (Fig. 69), but even cells which are described as ovoid are also regarded as bacilli, as in Löffler's so-called *Bacillus parvus ovatus*.

The grouping together of the different species according to the character of the colonies in nutrient gelatine (Figs. 70, 71) is also of questionable advisability. These characters can hardly




FIG. 71.—COLONIES ON NUTRIENT AGAR-AGAR, $\times 60$.

be considered to be of sufficient importance, or indeed in many cases to be sufficiently constant, to serve by themselves for this purpose. In many cases a slight variation in the composition of the



FIG. 72.—COLONY OF *BACILLUS ANTHRACIS*, $\times 60$. From a cover-glass impression-preparation, stained with gentian-violet.

nutrient medium may considerably affect the appearances of the colonies. At the same time, the appearances are very characteristic of certain species of bacteria (Fig. 72), and in other

cases the characters of the colonies, together with the characters of the growth in test-tubes, assist us in distinguishing species which are morphologically similar, as in the case of the comma bacilli of Finkler and of Koch.

The classification here given will be found to be a convenient form for arranging the microorganisms for reference, and it may lead the investigator to work upon the same lines as Zopf, and by tracing the life-history of individual forms in pure cultivations, either to extend the work of establishing protean species or to restrict the



FIG. 73.—BACTERIUM OF RABBIT SEPTICÆMIA.

doctrine of pleomorphism to a very few forms. For though the author adheres to the lines of classification proposed by Zopf, he is not prepared to accept his teachings in their entirety; thus, to embrace all described species, and to be consistent with the author's views, it has been necessary not only to add to Zopf's classification, but in many cases to modify his arrangement of species. For instance, Zopf regards the bacterium of rabbit septicæmia (Fig. 73) as a micrococcus. Of some species alteration in the nomenclature is justified by necessity.

Any arrangement at present can only be con-

sidered provisional, and, therefore, that arrangement which is of most practical assistance, and which leads to a clear description of the important characteristics on which the final classification will depend, must be considered to be the best. For example, if we abolish the genus bacterium, any rod-form may be at once classed as a bacillus; on the other hand, in the plan here adopted, we must determine the presence or absence of endogenous spore-formation before we can decide whether it be a bacterium or a bacillus. This necessarily leads to a more thorough study of their life-history. In the systematic description which follows, stress is laid upon the morphological appearances of bacteria, upon the absence or presence of spore-formation, and upon the appearances under cultivation, in addition to other characteristics, such as the changes produced by their growth. The determination of species rests upon the accumulated evidence afforded by a thorough knowledge of their life-history. The form of the organism, the changes it effects, and the microscopical appearances under cultivation must be collectively taken into account.

CHAPTER XII.

SYSTEMATIC AND DESCRIPTIVE.

THE *Schizomycetes*, *Spaltpilze*, or *Fission-fungi* have already been described as divisible, according to Zopf, into four groups: *Coccaceæ*, *Bacteriaceæ*, *Lep-totricheæ*, and *Cladotricheæ*. They comprise the following genera and species:—

GROUP I.—COCCACEÆ.

Genus I. *Streptococcus* (Chain-cocci).—Division in one or more directions. Individual cocci remain united together to form chains.

Genus II. *Merismopedia* (Plate-cocci).—Division in two directions, forming lamellæ or plates.

Genus III. *Sarcina* (Packet-cocci).—Division in three directions, forming colonies in cubes or packets.

Genus IV. *Micrococcus* (Mass-cocci).—Division in one direction, cocci after division remain aggregated in irregular clusters, or singly, or in pairs or in chains of three or four elements.

Genus V. *Ascococcus* (Pellicle-cocci).—Like micrococcus, but the cocci grow in characteristic gelatinous pellicles.

Genus I.—Streptococcus.

Streptococcus pyogenes, Rosenbach.—Cocci occurring singly, in chains, and in zooglœa. The individual cocci are small spherical cells, with a special tendency after fission for the resulting elements to remain attached to each other, forming chains or rosaries. There may be a few, three or more elements, linked together, or a great number, forming straight, serpentine, and twisted chains.

Preparations, stained preferably by the method of Gram (Plate XIX.), should be studied minutely with the best lenses and the best method of illumination in order to make out all the minute details. The individual elements composing the chains will be found to vary considerably in size; here and there in a preparation will be found a chain composed of excessively small elements, in another part the elements are all on a larger scale, and again in another part the elements will be peculiarly conspicuous on account of their size. Very characteristic appearances result from the fact that the elements enlarge and divide both longitudinally and transversely, and, indeed, the largest elements, for the most part, clearly show a division in two directions, resulting in the formation of tetrads. So great is the diversity in the size of the elements of some of the chains, that one might imagine that

there was more than one kind of streptococcus present in a preparation, until on examining some of the longest chains it is observed that various sizes are represented in different lengths of the same chain. In addition to the forms resulting from the fission of the elements, there are here and there in a chain, and sometimes terminally, larger elements, which are spherical, spindle-shaped, or in the form of a lemon. In the length of the chains, as in the size of the individual elements, there is usually a great diversity. In some cases they are composed of only a few, three or four elements, or in others eight, ten, or twenty. Here and there an exquisite rosary will extend in a straight line across the field of the microscope, or be twisted, curved, or serpentine; in some preparations twisted or entangled strands are observed which are composed of several hundred elements. Such chains will be found to be much thicker in one part than another. Another characteristic appearance is produced by separation of the elements resulting from fission in the long direction of the chain, by which lateral twigs or branches are formed. Another character, which is very striking, may be seen when the individuals in a chain have become separated; an unstained or faintly stained membrane may be found bridging across the interval. This will become still more visible in some of the preparations contrast-stained with eosin.

Cultivations.—In plate-cultivations the appear-

ances of the colonies are not very striking. They appear to the naked eye after three or four days as extremely minute, greyish-white, translucent dots, which under the microscope have a slightly yellowish-brown colour. They are finely granular and well defined. They do not liquefy the gelatine, and after several weeks do not exceed the size of a pin's head.

If the surface of nutrient gelatine solidified obliquely be traced over once or twice with a platinum needle bent at the extremity into a little hook charged with the cocci, a ribbon-shaped film develops in two or three days. This film is composed of minute, greyish-white, translucent dots or droplets, which can be more easily recognised with the aid of a pocket lens. According to the number of organisms sown on the jelly, the dots or colonies may be completely isolated, or form a more or less continuous film. The film by reflected light has an iridescent appearance like mother-of-pearl, but has a bluish or bluish-grey tint by transmitted light, and with a pocket lens appears distinctly brownish. The gelatine is not liquefied, and even after several weeks the cultivation is limited to the inoculated area, and the individual colonies are not larger than pins' heads. In gelatine-cultivations of the same age, but kept in the incubator at 18° C., the colonies get irregular in form, especially at the margin of the film, and give the growth an arborescent, fringed, or serrated appearance. Cultivated on the oblique surface of nutrient

agar-agar at 37° C. the growth is very similar, forming a film composed of minute dot-like colonies like grains of sand. But the film appears less transparent, is whiter, and the colonies have a greater tendency to get irregular in form. If inoculated with one tracing of the needle the growth is scanty, but tends to get thicker in the centre than towards the margins, which may have a terraced appearance. Inoculated in the depth of gelatine, there appears after a day or two at 18° C. a thread-like growth along the track of the inoculating needle. This delicate growth is found on examination with a pocket lens to consist of a linear series of extremely minute granules. In a few days more the beads or granules become more marked, but even after weeks the cultivation only appears like a string of minute white, compact, globular masses or grains. In broth at 37° C. the cocci in twenty-four hours create a turbidity, and gradually develop beautiful chains varying in length according to the age of the cultivations. Even in forty-eight hours there may be chains of eight, ten, twenty, or a hundred elements. After a few days the growth settles down at the bottom of the tube in the form of a white deposit, while the supernatant liquid becomes clear again.

Inoculated subcutaneously in the ear of rabbits, they produce in two days an inflammatory thickening with erysipelatous redness without suppuration.

They occur in pus and in diseases associated with septic complication.




Fig 1.




Fig 2.

The diseases and conditions in which streptococci have been found will be enumerated. In some cases the streptococci are identical with the first named, and in others there is need for further investigation; some being regarded as varieties, and others as distinct species.

Abscesses.—In abscesses a chain forming micrococcus was first described by Ogston. It was later studied by the methods of cultivation introduced by Koch and named by Rosenbach *Streptococcus pyogenes*. According to Flügge, after subcutaneous inoculation of mice with a small quantity of a cultivation, there is no result in 80 per cent. of the animals experimented on. Sometimes there is limited pus-formation at the seat of inoculation, sometimes the animals die without any very striking pathological appearances.

Rosenbach examined six cases of pyæmia. From Case 1 cultivations of *Streptococcus pyogenes* were obtained from the blood of the patient during life. The blood was stroked over the surface of the culture medium, and in two tubes the *Staphylococcus pyogenes aureus* was also present. In Case 2, with suppurative pleuritis, the pleura was tapped during life, and cultivations of *Streptococcus pyogenes* were obtained. In Case 3 streptococci were found in the metastases in the kidneys and in the other suppurations that were examined. In Case 4 the pleura was opened during life, and *Streptococcus pyogenes* and *Staphylococcus pyogenes aureus* were isolated from the fluid which escaped. In Case 5 pure cultivations of *Streptococcus pyogenes* were obtained from pus from the knee, which was punctured during life. This case was one of erysipelas and pyæmia after removal of a carcinoma of the breast. In Case 6, a case of whitlow with metastatic abscesses, the patient recovered, and no streptococci were found. Thus in six cases of

metastatic pyæmia *Streptococcus pyogenes* was found five times, partly in the blood and partly in the metastatic deposits, and twice in company with *Staphylococcus pyogenes aureus*.

Baumgarten, also, found the *Streptococcus pyogenes* in the internal organs in pyæmic cases, and Eiselsberg found *Streptococcus pyogenes* in company with *Staphylococcus pyogenes aureus* in the blood of cases of septicæmia.

Erysipelas.—In erysipelas Fehleisen isolated a streptococcus and described the appearances on cultivation. Fehleisen regarded the organism as quite distinctive, with special characters on cultivation, and claimed that it produced erysipelas when inoculated in the human subject. Rosenbach pointed out the extremely close resemblance in every respect to the *Streptococcus pyogenes*, though he described an apparent difference on cultivation on nutrient agar-agar. Other observers maintain that these organisms cannot be distinguished with certainty by either their morphological appearances or by their characters on cultivation. Moreover, the marked differences which have been described after inoculation were not obtained by those who repeated the experiments. Passet found that the results of inoculating *Streptococcus pyogenes* in the rabbit's ear induced a very similar condition to the result of inoculation of *Streptococcus erysipelatis*; and both organisms, by subcutaneous inoculation, and by inoculation in the cornea of rabbits and other animals, induced results without any constant difference. Passet showed that the inoculation of the cornea produced the same form of keratitis. Hoffa and Hajek described minute differences at the seat of inoculation, but Biondi and Eiselsberg repeated the experiments, and failed to establish the alleged differences. Baumgarten also investigated this subject, and failed to prove any essential difference, and, indeed, found much more often than might have

been expected from the publications of previous authors that no marked result at all was obtained on inoculation; and he concluded that *Streptococcus pyogenes* and *Streptococcus erysipelatis* in their form, cultivation, and their effects on animals were identical. Passet, Biondi, Eiselsberg, Baumgarten, and Fränkel have definitely accepted the identity of the streptococcus associated with suppuration, and the streptococcus associated with erysipelas.

Spreading Gangrene.—From a case of spreading gangrene, which was identical with Ogston's erysipelatoid wound gangrene, and regarded by him as the most intense and dangerous form of erysipelas, Rosenbach obtained pure cultivations of a streptococcus by incising the skin of the limb, and inoculating tubes from the turbid reddish fluid which escaped. That the streptococcus was identical with *Streptococcus pyogenes* was ascertained by comparison with a cultivation derived from pus, of the mode of growth, and of the effect on animals.

Surgical Fever.—Eiselsberg proved the presence of a streptococcus in the blood of several cases of surgical fever in Billroth's clinic. The organism was identified by cultivation with *Streptococcus pyogenes*.

Diphtheria.—In three cases of typical diphtheria Löffler found a streptococcus. He isolated it by cultivation, found that it was similar in form, characters on cultivation, and effects after inoculation, to Fehleisen's streptococcus of erysipelas. Löffler was not inclined to regard them as identical, because Fehleisen never found his cocci in the blood-vessels. Flügge named the organism *Streptococcus articulorum*, and states that, after subcutaneous inoculation or injection of a cultivation in mice, a large proportion of the animals die, and in the sections of the spleen and other organs the streptococci are again seen. Baumgarten investigated the same subject, and decided that the streptococcus was identical with *Streptococcus pyogenes*.

Puerperal Fever.—Fränkel isolated a streptococcus from puerperal fever, which he at first called *Streptococcus puerperalis*, but subsequently identified with *Streptococcus pyogenes*. Winkel obtained a pure cultivation of a streptococcus from the blood of the heart in a case of puerpera peritonitis. It produced erysipelatous redness when inoculated in the rabbit's ear, and in form and in cultivation was similar to the streptococcus in erysipelas. Cushing found *Streptococcus pyogenes* associated with puerperal infection. The cocci were found in endometritis diphtheritica as well as in secondary puerperal inflammation. These observations were still further confirmed by Baumgarten, and Bumm isolated the same organism in puerperal mastitis.

Scarlet Fever.—The occurrence of a streptococcus in certain cases of scarlet fever has been observed by several investigators—Croke, Löffler, Babes, Heubner, and Bahrdt, and notably by Fränkel and Freudenberg.

Croke, in cases of scarlet fever with severely affected throat, found bacilli, cocci, and streptococci in the organs of the throat, and cocci in the internal organs. Croke left it an open question whether these cocci were the specific organisms of scarlet fever, or to be regarded as diphtheritic or septic associates. He inclined on clinical grounds to the latter view.

Löffler, in cases of scarlatinal diphtheria, found the same chain-forming micrococcus which he had found in typical diphtheria.

Babes was able to constantly prove in inflammatory products secondary to scarlatina the presence of a streptococcus greatly resembling that in pus.

Heubner and Bahrdt, in a fatal case of scarlet fever in a boy, complicated with suppuration of the finger- and knee-joints and with pericarditis, found a streptococcus identical in form, from his description, with *Streptococcus pyogenes*. Cultivations were not made. The secondary

infection started from diphtheritically affected tonsils, which were followed by retro-pharyngeal abscesses.

Fränkel and Freudenberg examined for micro-organisms three cases of scarlatina with well-marked affection of the throat. In all three cases they obtained cultivations of cocci from the submaxillary lymphatic glands, spleen, liver, and kidney. These cocci could in no way be distinguished from *Streptococcus pyogenes* derived from pus, nor from the undoubtedly identical streptococcus which one of them (A. Fränkel) had already repeatedly cultivated in large numbers from puerperal affections. In two of the cases *Streptococcus pyogenes* was the only organism present, and in all three cases it was far in excess of other colonies which developed. The organisms were also found in sections of the organs by microscopical examination. The identity of this streptococcus with *Streptococcus pyogenes* and *Streptococcus puerperalis* was established by comparison of their macroscopical and microscopical appearances in cultivations on nutrient agar-agar, nutrient gelatine, and in broth, both at the ordinary and at higher temperatures, and also by experiments on animals. They concluded that it could be stated with certainty that the organisms in question did not stand in causal relation to scarlet fever. They considered that special methods of microscopical and biological research were apparently needed for demonstrating the true scarlet fever contagium, which probably was especially present in the skin. They considered that the presence of the streptococcus was due to a secondary infection, to which the door was opened by the lesions of the throat—a view which was supported by the fact that the organisms were found in the submaxillary lymphatic glands. They preferred to use the term “secondary” to “complicated” or “combined” infection, because this expresses the fact that by the effect of the scarlatinal virus the soil is rendered suitable for this ubiquitous microbe when it has once gained an entrance.

Klein found the streptococcus in five out of eleven cases of scarlet fever, twice in association with other cocci, and three times alone, and regarded it as the contagium of scarlet fever. The cases from which the organism was obtained were all cases with ulcerated throat. An analysis of these cases is shown in the following table :—

	Name of patient.	Condition of tonsils.	Source of blood.	Micro-organisms isolated.	Death or recovery.
1	Lilian Fuller, aged 5	Severely ulcerated	Finger	Streptococcus	Ultimately recovered.
2	Kate Fuller, aged 2	Much ulcerated	"	{ Staphylococcus pyogenes aureus Liquefying micrococcus Streptococcus Staphylococcus Streptococcus	} Died of pyæmia.
3	Henry Lampson, aged 8	Ulcerated	"		
4	— (a woman), aged 40	Much ulcerated	Arm		
5	— (a girl), aged 19	"	"	"	"
6	Blanche Moody, aged 15	Ulcerated	Finger	Streptococcus	Recovered.
7	Ellen Warr, aged 22	Much ulcerated	"	None	[Not stated.]
8	Robert Hughes, aged 8	Ulcerated	"	"	"
9	Florence Glossop, aged 2½	"	Heart	Streptococcus	Died.
10	Edward Fenton, aged 3	—	"	None	"
11	R. Boucher, aged 20 months	Advanced ulceration	"	"	"

But the evidence is sufficient to show that in any disease accompanied with a lesion of the skin or mucous membrane, in which the blood and tissues are profoundly affected by the virus of that disease, septic micro-organisms gain an entrance into the circulation, and may not only escape destruction, but may gain the upper hand and set up destructive processes. And it is interesting to observe in connection with this, that the streptococcus was isolated in all cases at or about the

time at which the temperature was highest; in other words, at a time when the tissue resistance to an invasion of micro-organisms would be lowered. The occurrence of streptococci, in certain cases of diphtheria, scarlet fever, cow-pox, small-pox, measles, typhoid fever, must be regarded as a secondary result associated with suppuration or with septic or pyæmic complication; in fact, with the exception of negative evidence, bacteriology has not assisted us in the least in the determination of the real nature of the morbid agent or actual contagium of scarlet fever.

Small-pox.—Hlava has established the presence of *Streptococcus pyogenes* in the pustules of variola, and Garré found streptococci in the internal organs in a case of variola hæmorrhagica.

In a fatal case of variola complicated with pemphigus, Garré found a streptococcus in the pemphigus vesicles. Whether it was identical with *Streptococcus erysipelatis* Garré left an open question.

Foot - and - mouth Disease.—In 1868 Brown figured a streptococcus in the milk of cows affected with foot-and-mouth disease. From the vesicles of this disease in sheep, Klein isolated a streptococcus which he regarded as the contagium of the disease. From Klein's description of its morphological features and characters on cultivation, it closely corresponds with *Streptococcus pyogenes*. Baumgarten regards Klein's alleged contagium as most probably the *Streptococcus pyogenes*. Against the fact of this organism being the virus of this disease, and in favour of its being a septic organism, we have the results of inoculation experiments. A great number of subcutaneous inoculations in sheep were without any result. One of several guinea-pigs fed with the organism had an abscess, another an ulcer. In sheep fed with the organism, two out of nine, after a number of repeated administrations, developed a disease, regarded by Klein as foot-and-mouth disease. There is the

possibility of two fallacies—one that the disease, if really foot-and-mouth disease, may have occurred incidentally; the other, that the result which occurred may have been a spurious form of foot-and-mouth disease. The control experiment of exposing a number of sheep with these two sheep to see if the disease would spread was wanting.

Cattle Plague.—Semner cultivated streptococci from the blood and lymphatic glands of a sheep suffering from cattle plague. A calf inoculated with a cultivation was stated to have died in seven days from cattle plague. The cocci were stated to lose their virulence by successive cultivation, and the weakened cultivation to protect against the virulent disease. It is extremely likely that this was another manifestation of septic streptococci, and that the animal died very probably of septic infection. Klein is also of opinion that the specific nature of these micro-organisms cannot be considered to be established.

Strangles.—Schutz has described a streptococcus as the contagium of glanders.

Contagious Mammitis.—Nocard has cultivated a streptococcus in contagious mammitis which he regards as specific.

Yellow Fever.—Babes observed the presence of streptococci in the vessels of the kidney and liver in yellow fever. Cultivation experiments are wanting. It was probably secondary infection with *Streptococcus pyogenes*.

Bilious Fever.—Babes, in a case of fièvre bilieuse typhoïde, found masses of streptococci filling the vessels of the liver, kidney, and spleen. Probably another instance of secondary infection with *Streptococcus pyogenes*.

Measles.—From the blood and from inflammatory post-products in measles Babes isolated a streptococcus, which he describes as closely resembling the *Streptococcus pyogenes*.

Ulcerative Endocarditis.—Wyssokowitsch found cocci in the internal organs in ulcerative endocarditis, and produced the disease in animals after injury to the valves

by injection of *Streptococcus pyogenes* and other organisms. Wechselbaum, by microscopical research and by cultivation experiments, proved the presence of *Streptococcus pyogenes* in acute verrucous endocarditis. Baumgarten confirmed this. He found *Streptococcus pyogenes* alone in one case and accompanied by *Staphylococcus aureus* in another.

Typhoid Fever.—Senger found a streptococcus in a case of typhoid with secondary infection. This was probably *Streptococcus pyogenes*. Dunin found the well-known pyogenic organisms in post-typhoid suppuration, mostly as staphylococci, but sometimes streptococci.

Pneumonia.—Wechselbaum found a *Streptococcus pneumoniae*, which resembled *Streptococcus pyogenes* and *erysipelatis* morphologically and on cultivation. It was found in twenty-one cases of pneumonia by microscopical research, and cultivated in nineteen. It had no effect on the rabbit's ear. Baumgarten, nevertheless, regards this as *Streptococcus pyogenes*, and is of opinion that it is only a matter of further research to establish that view.

Neumann found streptococci by microscopical research in the lungs, and by cultivation isolated a streptococcus in pneumonia after typhoid. It corresponds with Wechselbaum's streptococcus in not affecting the rabbit's ear.

Empyema.—Rosenbach obtained a pure cultivation of *Streptococcus pyogenes* in a case of empyema. Wechselbaum, in two cases, also established its presence.

Broncho-pneumonia.—Thaon found a streptococcus in the lungs of children in fatal cases of broncho-pneumonia, complicating measles, diphtheria, and whooping-cough. It was regarded as identical with the streptococcus isolated by Löffler from diphtheria. Fränkel discovered a streptococcus in the lungs of a case of true croup complicated with broncho-pneumonia, and by cultivation established its identity with *Streptococcus pyogenes*.

Progressive Tissue Necrosis in Mice.—Koch produced a disease in mice by subcutaneous injection of

putrid blood. In tissue sections a chain coccus was found, and Baumgarten is of opinion that it is very probably identical with the *Streptococcus pyogenes*; but cultivations are still wanting.




FIG. 74.—STREPTOCOCCUS OF PROGRESSIVE TISSUE NECROSIS IN MICE. (a) Necrotic cartilage cells, and (b) Chains in masses; (c) Chains isolated [after Koch].

Anthrax.—Charrin found cocci in rabbits, examined some hours after death from anthrax. These, when isolated, produced death in rabbits from septicæmia without suppuration. Chains composed of from fifteen to twenty elements were found in all the organs. This was probably another instance of *Streptococcus pyogenes*.

Syphilis.—Kassowitz and Hochsinger found the presence of a streptococcus in the tissues and internal organs, and especially in the blood-vessels, in fatal cases of congenital syphilis. These observers regarded their discovery as having an important bearing on the etiology of syphilis, but Kolisko pointed out that it was only the result of septic infection with presence of *Streptococcus pyogenes*, as had already been established in scarlet fever.

Cerebro-spinal Meningitis.—From the meningeal exudation of a case of apparently idiopathic cerebro-meningitis, Banti found by Koch's methods the presence of *Streptococcus pyogenes* and *Staphylococcus aureus* and

albus. The cocci probably entered through an abscess of the jejunum.

Blepharadenitis and Dacryocystis.—Widmark isolated by cultivations *Streptococcus pyogenes* and other organisms from cases of blepharadenitis and phlegmonous dacryocystis. In phlegmonous dacryocystis Widmark found *Streptococcus pyogenes* almost exclusively.

Leukæmia.—Flügge cultivated a streptococcus from necrotic patches in the spleen of a fatal case of leukæmia. Cultures corresponded very closely with *Streptococcus pyogenes*. Inoculation in the ears of rabbits produced similar results to *Streptococcus pyogenes* or *erysipelatis*. Flügge calls it *Streptococcus pyogenes malignus*, but concludes that it is probably identical with the streptococcus from pus.

Soil.—Nicolaier, and later Guarneri, isolated a streptococcus from soil. Microscopically it could not be distinguished from other streptococci. Baumgarten is of opinion that it is neither in form nor in cultivation to be distinguished with certainty from *Streptococcus pyogenes*.

Air.—Emmerich succeeded in proving the presence of streptococci in the air of a hospital where erysipelas had broken out. These cocci, in their form, their characters on cultivation, and in inoculation results, were identified with the *Streptococcus erysipelatis*.

Putrefaction.—Baumgarten points out that streptococci are found in the most various substances undergoing putrefaction.

EXAMINATION AND CULTIVATION OF STREPTOCOCCI.

Cover-glass preparations can be stained with the watery solutions of the aniline dyes. In some cases very beautiful preparations can be obtained by using Neelsen's solution, and removing excess of stain by rinsing in alcohol. To examine pus, milk, or broth, take an ordinary platinum inoculating needle bent at the extremity

into a hooklet. Dip it into the liquid to be examined, and spread it on a cover-glass into as thin a film as possible ; the preparation is treated in the ordinary way, that is to say, the film is allowed to dry, and the cover is taken up with forceps, and passed three times through the flame with its prepared side uppermost.

Gram's Method of Eosin.—In this way the streptococci are stained blue, and stand out in marked contrast to the rest of the preparation. Use freshly prepared solution. Float the cover-glasses on the solution for ten minutes to half an hour, then transfer them to iodine-potassic-iodide solution, until they assume the colour of a tea leaf ; then immerse them in alcohol until they are decolourised ; dip them in an alcoholic solution of eosin for a few moments, and then transfer them to clove oil to clarify the film ; to remove the clove oil gently press the cover between two layers of clean filter paper, then mount in xylol balsam (Plate XIX.).

A good method for cultivating streptococci is to employ a sterilised looped platinum wire, and to spread a droplet, for example, of pus or blood over the surface of nutrient agar-agar solidified obliquely. The tubes are then placed in the incubator at 37° C. ; the streptococci will appear in the course of two or three days in the form of minute dotted colonies. If present alone, and in considerable quantities, the inoculated surface will exhibit a pure cultivation consisting of a number of such colonies, whilst a flocculent mass is observed in the liquid which collects at the bottom of the agar-agar tubes ; this flocculent mass will be found to be composed of chains. From such a tube inoculate a number of the small flasks employed in Pasteur's laboratory for cultivations in liquids. In this way a number of pure cultivations in milk and broth are established, which can be readily examined from time to time. From a pure cultivation in broth or agar-agar tubes of nutrient gelatine can be inoculated. Cover-glass-preparations from the growths on solid media can be

made in the usual way, and stained with either a watery solution of fuchsine or gentian violet; but to stain preparations made from milk or broth, or from the liquid in agar-agar tubes, use the method of Gram; the stain will then be removed, except from the streptococci, and very beautiful preparations result.

Genus II.—*Merismopedia*.

SPECIES.

ASSOCIATED WITH DISEASE:—

<i>Merismopedia gonorrhœæ</i>	.	.	Pathogenic in man.
<i>Micrococcus tetragonus</i>	.	.	{ Saprophytic in man.
<i>Diplococcus albicans tardissimus</i>	.	.	} Pathogenic in animals.
			. Saprophytic in man.

UNASSOCIATED WITH DISEASE:—

<i>Micrococcus citreus conglomeratus</i>	.	.	} Simple saprophytes.
<i>Micrococcus subflavus</i>	.	.	
<i>Micrococcus albicans amplus</i>	.	.	

***Merismopedia gonorrhœæ* (*Coccus of Gonorrhœa*).**—Cocci 0.83μ in diam., singly, in pairs, in tetrads, and zooglœa groups. They are found in gonorrhœal pus adhering to the pus corpuscles and epithelial scales. Artificial cultivations have been carried out,* and it has been claimed that the pathogenic character of the cocci has been established by inoculation.

***Micrococcus tetragonus*.**—Cocci about 1μ in diam., in groups of four (tetrads), surrounded by a hyaline capsule. They are found in the sputa of phthisical patients and in the walls of tubercular cavities. In a test-tube of nutrient gelatine they form an irregular white growth, more especially in the upper part of the needle track

* Bockhart, *Sitzungsberichte der Phys. Med. Gesell. Würzburg*. 1882.

(Plate V., Fig. 1). On the sloping surface of nutrient agar-agar thick, whitish, heaped-up masses develop. Guinea-pigs and mice inoculated with a minute quantity of a pure cultivation die in from two to ten days, and the groups of the characteristic tetrads may be found in the capillaries throughout the body, especially in the spleen, lung, and kidney (Plate II., Fig. 1).

Double infection can be produced by inoculating a mouse with a pure cultivation of *Bacillus anthracis* two or three days after inoculation with *Micrococcus tetragonus*. On examination after death, the capillaries of the lungs, liver, and kidney are filled with both anthrax bacilli and masses of tetrads* (Plate XXIV., Fig. 2).

Micrococcus citreus conglomeratus, Bumm.—Cocci 1.5μ in diam., similar to gonococci. They form lemon-yellow colonies on plate-cultivations. Isolated from blennorrhœic pus and from dust from the air.

Diplococcus albicans tardissimus, Bumm.—Cocci morphologically identical with gonococci. They grow extraordinarily slowly on gelatine. They were isolated from urethral pus.

Micrococcus subflavus, Bumm.—Cocci morphologically resembling gonococci. Cultivated on nutrient gelatine, they form whitish dots which become gradually greyish and then yellow in colour, and confluent. They were observed in lochial discharges and vaginal secretions.

Micrococcus albicans amplius, Bumm.—Cocci in pairs and tetrads similar to gonococci, but considerably larger. Found in vaginal secretions.

* The Author, *Lancet*, 1885.

Genus III.—Sarcina.

UNASSOCIATED WITH DISEASE:—

<i>Sarcina lutea</i> . . .	}	Chromogenic saprophytes.
<i>Sarcina aurantiaca</i> . . .		
<i>Sarcina ventriculi</i> . . .		
<i>Sarcina intestinalis</i> . . .	}	Simple saprophytes.
<i>Sarcina urinæ</i> . . .		
<i>Sarcina litoralis</i> . . .		
<i>Sarcina Reitenbachii</i> . . .		
<i>Sarcina hyalina</i> . . .		
<i>Sarcina alba</i> . . .		

***Sarcina lutea*.**—Cocci singly, in pairs, in tetrads, and in packets. A single individual in a tetrad may be divided into two, or into four, so that a tetrad within a tetrad results. Cultivated in nutrient agar-agar in a test-tube, they form a colourless growth along the track of the needle, and a bright canary-yellow layer upon the surface, where they have access to the air (Plate VII., Fig. 1; Plate IX., Fig. 1). In plate-cultivations the colonies are round, slightly granular in appearance, and yellow. Cultivated in a test-tube containing nutrient gelatine, they grow rapidly; the gelatine becoming liquid, the yellow growth forms a wad about the middle of the tube (Plate VI., Fig. 2), or, liquefying the whole of the gelatine, subsides to the bottom of the test-tube. Cultivated on sterilised potatoes, they form a yellow layer (Plate XV., Fig. 1). In drop-cultures in bouillon the subdivision into tetrads within tetrads and formation of groups of 8, 16, and 24 can be studied (Plate I., Fig. 7). Inoculation of mice produces negative results. The cocci are occasionally present in the air.

Sarcina aurantiaca.—Cocci singly, in pairs, in tetrads, and in packets. They form small orange-yellow colonies on plate-cultivations, and in test-tubes slowly liquefy the gelatine along the whole needle track, forming on the surface an orange-yellow growth. On potatoes they slowly develop the same pigment.

Sarcina ventriculi, Goodsir.*—Cocci reaching $4\ \mu$ in diam., united in groups of four, or multiples of four, producing cubes or packets with rounded-off corners. Contents of the cells are greenish or yellowish-red. They occur in the stomach of man and animals in health and disease, and were first detected in vomit.

Sarcina intestinalis, Zopf.†—Cocci in groups of four or eight. Very regular in form; never in the large packets which occur in *Sarcina ventriculi*. They are found in the intestinal canal, especially the cæcum, of poultry, particularly fowls and turkeys.

Sarcina urinæ, Welcker.—Very small cocci, $1.2\ \mu$ in diam., united in families of 8 to 64. Observed in the bladder.

Sarcina litoralis, Oersted.—Cocci 1.2 — $2\ \mu$ in diam., bound together in 4 to 8 families, which, in their turn, may unite and include as many as 64 tetrads. Plasma colourless; in each cell 1—4 sulphur granules. Discovered in sea-water containing putrefying matter.

Sarcina Reitenbachii, Caspary.—Cocci about 1.5 to $2.5\ \mu$ in diam., at the time of division lengthened to $4\ \mu$. Mostly united together from 4 to 8 in number; occasionally 16 or more. Colourless cell-wall, lined with

* Goodsir, *Edinburgh Med. and Surg. Journal*. 1842.

† Zopf, *Die Spaltpilze*. 1885.

rose-red layer of plasma. Found on rotting water-plants.

Sarcina hyalina, Kützing.—Cocci round, 2.5μ in diam., almost colourless. United in families of 4 to 24 cells, reaching 15μ in diam. In marshes.

Sarcina alba.—Small cocci. They form small white colonies on nutrient gelatine. In test-tube cultivations they grow slightly along the needle track, but are heaped up on the surface without liquefying the gelatine. They are present in the air.

Genus IV.—Micrococcus.

SPECIES.

ASSOCIATED WITH DISEASE :—

	Micrococcus pyogenes aureus	} Pyogenic in man (?). } Pathogenic in animals.
	Micrococcus pyogenes albus	
	Micrococcus pyogenes citreus	} Non-pathogenic.
	Micrococcus cereus albus	
	Micrococcus cereus flavus	} Non-pathogenic.
	Micrococcus in scarlatina	
	Micrococcus in measles	} Possibly only saprophytic.
	Micrococcus in whooping-cough	
In man .	Micrococcus in hæmophilia neonatorum	} Pathogenic (?).
	Micrococcus in typhus	
	Micrococcus in acute yellow atrophy	} Pathogenic (?).
	Micrococcus in dental caries	
	Micrococcus in gangrene	} Pathogenic (?).
	Micrococcus pyogenes tenuis	
	Micrococcus in rabies	} Pathogenic.
	Micrococcus of septicæmia in rabbits	
	Micrococcus of pyæmia in rabbits	} Pathogenic (?).
	Micrococcus of progressive suppuration in rabbits	
In animals .	Micrococcus parvus ovatus	} Pathogenic (?).
	Micrococcus of pyæmia in mice	
	Micrococcus perniciosus	} Pathogenic (?).
	Micrococcus bombycis	
In plants .	Micrococcus insectorum	} Pathogenic (?).
	Micrococcus amyliovorus	

UNASSOCIATED WITH DISEASE :—

Micrococcus cyaneus	} Chromogenic saprophytes.
Micrococcus aurantiacus	
Micrococcus chlorinus	
Micrococcus violaceus	
Micrococcus luteus	
Micrococcus rosaceus	} Simple saprophytes.
Micrococcus hæmatodes	
Micrococcus candidus	
Micrococcus candicans	
Micrococcus fœtidus	
Micrococcus crepusculum	
Micrococcus cinnabareus	
Micrococcus flavus liquefaciens	
Micrococcus flavus tardigradus	
Micrococcus versicolor	
Micrococcus viticulosus	
Micrococcus lacteus faviformis	
Micrococcus fulvus	
Micrococcus viscosus	
Micrococcus coronatus	
Micrococcus radiatus	
Micrococcus flavus desidens	

Micrococcus pyogenes aureus. (*Staphylococcus pyogenes aureus*, Rosenbach. *Yellow coccus in pus*.* *Coccus of acute infectious osteomyelitis*).—Cocci singly, in pairs, very short chains, and irregular masses. Cultivated on nutrient agar-agar an orange-yellow culture develops, looking like a streak made with oil paint† (Plate IX., Fig. 2). Cultivated in a test-tube of nutrient gelatine, the gelatine is rapidly liquefied, and the growth subsides as an orange-yellow sediment. On potatoes and blood serum a similar orange-yellow culture grows luxuriantly.

The micro-organisms injected into the pleura or knee of a rabbit produce, as a rule, a fatal result on the following day; but if it survives longer, it eventually dies of severe phlegmon. If injected

* Ogston, *Brit. Med. Journ.* 1881.

† Rosenbach.

into the knee of a dog, suppuration occurs, followed by disintegration of the joint. The cocci do not cause any septic odour in pus, nor does any gas develop. Albumen is converted by their action into peptones.

They occur in the pus of boils and in the abscesses of pyæmia, puerperal fever, and acute osteomyelitis. Injected into the peritoneal cavity of animals, they set up peritonitis, and introduced into the jugular vein they produce septicæmia and death. When a small quantity of a cultivation was introduced into the jugular vein after previous fracture or contusion of the bones of the leg, the animal died in about ten days, and abscesses were found in and around the bones, and in some cases in the lungs and kidneys. Similar cocci were found in the blood and pus of the animals.*

Micrococcus pyogenes albus (*Staphylococcus pyogenes albus*, Rosenbach). Cocci microscopically indistinguishable from the above. In cultivations also they resemble the *Micrococcus pyogenes aureus*, but the growth consists of opaque white masses. They liquefy nutrient gelatine rapidly, and subside to the bottom as a white sediment. They are also similar to the above-mentioned in their pathogenic action. Pure cultivations of the organism were obtained from a case of acute suppuration of the knee-joint.

Micrococcus pyogenes citreus (*Staphylo-*

* Becker, *Deutsche Med. Wochenschr.* Nov., 1883.

coccus pyogenes citreus, Passet).* Cocci singly, in pairs, very short chains, and irregular masses. If cultivated on nutrient gelatine or nutrient agar-agar, a sulphur or lemon-yellow growth develops (Plate X., Fig. 3). When inoculated under the skin of mice, guinea-pigs, or rabbits, an abscess forms after a few days, from which a fresh cultivation of the micro-organism can be obtained. They are frequently present in pus.

Micrococcus cereus albus (*Staphylococcus cereus albus*, Passet).* Cocci, morphologically similar to the above, but distinguished by forming on nutrient gelatine a white, slightly shining layer, like drops of stearine or wax, with somewhat thickened, irregular edge. The needle track develops into a greyish-white, granular thread. In plate-cultivations, on the first day, white points are observed, which spread themselves out on the surface to spots of 1—2 mm. When cultivated on blood serum a greyish-white, slightly shining streak develops, and on potatoes the cocci form a layer which is similarly coloured. (Plate IV., Fig. 3, XI. Fig. 3.)

Micrococcus cereus flavus (*Staphylococcus cereus flavus*, Passet).*—Cocci which also occur in pus. If cultivated in nutrient jelly, the growth, which is at first white, becomes lemon-yellow, somewhat darker in colour than *Micrococcus pyogenes citreus*. Microscopically *Micrococcus cereus flavus* corresponds with *Micrococcus cereus albus*, and they both form

* Passet, *Fortschritte der Medicin*, Jan. 15th and Feb. 1st, 1885.

zooglœa of medium-sized cocci (diam. $1\cdot16\ \mu$). Inoculation experiments with both kinds gave negative results. Among the micro-organisms present in pus a coccus has been described as occurring occasionally which is almost identical with *Bacterium pneumoniæ crouposæ*; refer also to *Streptococcus pyogenes* (p. 225) and to *Micrococcus pyogenes aureus* (p. 246).

In the following diseases cocci have been described. On cultivation they have proved to be in many cases identical with organisms commonly found in pus, etc. Others require further investigation.

Scarlet fever.—Cocci have been found in the blood,* in the scales of the desquamating epidermis,† and in the discharges and ulcerated tissue of the throat (see p. 232).

Measles.—Round cocci and diplococci have been observed in the catarrhal exudation, in the papules and in the capillary vessels of the skin, and in the blood of patients attacked with measles.‡

Whooping-cough.—Elliptical cocci are said to be constantly present in the expectoration of persons suffering from whooping-cough.§

Hæmophilia neonatorum, Klebs.—A coccus, which has been named *Monas hæmorrhagicum*, is stated to be characteristic of this disease.

Typhus.—Actively motile dumb-bell cocci have been described in the blood, and plugs of cocci in the lymphatics of the heart, in cases of typhus fever.||

* Coze and Feltz, *Malad. Infect.* 1872.

† Pohl-Pincus, *Centralblatt f. d. Med. Wiss.* 1883.

‡ Keating, *Phil. Med. Times.* 1882. Cornil and Babes, *Les Bactéries.* 1885.

§ Bürger, *Berl. Klin. Woch.* 1883.

|| Mott, *Brit. Med. Journal.* 1883.

Acute yellow atrophy.—Cocci have been observed in the vessels of the liver in this disease.*

Gangrene.—Oval and round cocci are found, which form zooglœa in the depth of gangrenous tissues. From gangrene of the lung cocci have been isolated, which form greyish-white colonies in plate-cultivations of nutrient gelatine. In a test-tube of nutrient gelatine a growth results chiefly on the surface; the cultivations yield a penetrating odour.

Micrococcus pyogenes tenuis, Rosenbach.—Cocci, which, cultivated on agar-agar, form an opaque streak in the track of the needle with a transparent glassy growth extending from it. Occasionally found in the pus of closed abscesses.

Rabies.—Cocci have been described in connection with hydrophobia. The cocci were observed in sections of the spinal chord of rabid dogs. The descriptions given by different observers † vary considerably, and it is not yet ascertained whether any particular coccus is constantly associated with the disease. Nor have the organisms observed in stained preparations been cultivated apart from the diseased animal. By many, however, hydrophobia is believed to be due to the presence of a micro-organism, and researches which are being carried on in connection with attenuation of the virus still continue to excite the keenest interest ‡ (p. 198).

Micrococcus of septicæmia in rabbits, Koch.§—Ellipsoidal cocci $\cdot 8$ — 1μ in largest diam. The disease was produced by the injection of putrid

* Eppinger, *Prager Vierteljahrsschrift*. 1875.

† Hermann Fol, *Académie des Sciences*. 1885. Babes, *Les Bactéries*. 1886. Dowdeswell, *Journ. Royal Microscop. Soc.* 1886.

‡ Pasteur, *Comptes Rendus*. 1882. *Congrès de Copenhague*, 1884. *Académie des Sciences*, 1885, 1886. Vignal, "Report on M. Pasteur's Researches on Rabies and the Treatment of Hydrophobia by Preventive Inoculation," *Brit. Med. Journal*. 1886.

§ Koch, *Wundinfect. Krankheit*. 1878.

meat infusion. After death slight œdema was noted at the site of injection, slight extravasation of blood, and great enlargement of the spleen. No emboli or peritonitis resulted. Masses of cocci were found in the capillaries of different organs, especially in the glomeruli of the kidneys. Rabbits and mice inoculated with blood from the heart proved susceptible to the disease.

Micrococcus of pyæmia in rabbits, Koch.*—Round cocci and diplococci $\cdot 25 \mu$ in diam. The disease was produced by the subcutaneous injection in a rabbit of distilled water in which the skin of a mouse had been macerated. At the autopsy there were found great infiltration around the site of injection, peritonitis, and accumulations in the liver and lungs; in short, the appearances of pyæmia. In the capillaries of the organs examined, masses of cocci were observed enclosing blood-corpuscles (Fig. 75). Fresh inoculations in rabbits with exudation-fluid, or blood from the heart, reproduced the same disease.

Micrococcus of progressive suppuration in rabbits, Koch.*—Cocci only about $\cdot 15 \mu$ in diam., principally in thick zooglœa. The disease was induced by the injection into rabbits of decomposing blood. At the place of injection a spreading abscess formed, which was fatal to the animal in about twelve days. No bacteria were observed in the blood, but in the walls of the abscess thick

* Koch, *Wundinfect. Krankheit.* 1878.

masses of cocci were found. The pus is infectious, causing the same disease in healthy rabbits.

Micrococcus parvus ovatus (*Bacillus parvus ovatus*, Löffler).—Small ovoid cocci, similar to the coccus of rabbit septicæmia. Cultivated on gelatine they develop readily a greyish-white growth at the entrance of the inoculating needle. They are pathogenic in mice and rabbits. They proved also fatal in a pig after two days, producing œdema of the




FIG. 75.—MICROCOCCLUS OF PYÆMIA IN RABBITS; VESSEL FROM THE CORTEX OF THE KIDNEY, $\times 700$. (a) Nuclei of the vascular wall; (c) Masses of micrococci adherent to the wall and enclosing blood-corpuscles.

skin, inflammation of the mucous membrane of the stomach, but no effect on the general intestinal tract or mesentery. They were isolated from a pig suffering from a fatal disease simulating swine-erysipelas.

Micrococcus of pyæmia in mice, Klein.—Certain cocci which were present in pork broth proved fatal to mice in about a week, producing purulent inflammation and abscess in the lungs. Fresh inoculations in mice again produced a fatal result with pyæmic symptoms.

Micrococcus perniciosus (*Parrot disease*).—Cocci, singly, in chains, and in zoogloea have been

described in connection with the disease of the grey parrot (*Psittacus erithacus*).^{*} This disease is fatal to about 80 per cent. of these parrots imported to Europe. They suffer from diarrhœa and general weakness; their feathers are ruffled, their wings hang loosely, and their eyelids close; convulsions set in, and death follows. At the autopsy greyish nodules are found in the lungs, liver, spleen, and kidney; in and around the capillaries of these nodules, and in the blood of the heart, the cocci are found in great numbers in zooglœa, and more rarely in chains. Inflammatory change in the surrounding tissue is absent.

Micrococcus bombycis, Béchamp (*Microzyma bombycis*).—Oval cocci $\cdot 5 \mu$ in diam., singly, in pairs, and chains. They occur in the contents of the alimentary canal, and in the gastric juice of silkworms suffering from "*flacherie*" ("*Maladie de morts blancs*," "*flaccidezza*," "*schlafsucht*").

Micrococcus insectorum, Burrill.—Obtusely oval cells, $\cdot 7$ — 1μ long and $\cdot 55 \mu$ broad, singly, in pairs, chains, or zooglœa. They were detected in the digestive organs of the chinck-bug (*Blissus leucopterus*) when suffering from a certain contagious disease.

Micrococcus amyliivorus, Burrill.—Oval cells, 1 — $1\cdot 4 \mu$ long, $\cdot 7 \mu$ broad, singly, in pairs, and rarely in fours, never in chains, are found embedded in an abundant mucilage which is very soluble in water. They have been described as producing the so-called "fire blight" of the pear tree and other plants.

Micrococcus cyaneus, Cohn (*Bacteridium cyaneum*, Schröter).—Elliptical cells, growing upon cooked potato, and producing a blue colour. In nutrient solutions they form zooglœa, at first colourless, then bluish-green, and finally intense blue.

Micrococcus aurantiacus, Schröter.—Cocci, oval, $1\cdot 5 \mu$ in diam., singly or in pairs, or in zooglœa. They occur as orange-yellow spots which coalesce into patches. A golden-yellow pellicle develops when they are cultivated

^{*} Wolff, *Virchow's Archiv.* 1883.

in nutrient liquids. The colouring matter is soluble in water. They were observed on boiled potatoes and white of egg.

Micrococcus chlorinus, Cohn.—Cocci occur in the form of a finely granular zooglœa, causing a yellowish-green or sap-green layer on boiled eggs and nourishing solutions. The colouring matter is soluble in water, and is decolorised by acids.

Micrococcus violaceus, Schröter.—Cocci or elliptical cells, described as uniting into violet-blue gelatinous spots, which again unite to form larger patches. They were observed on boiled potatoes exposed to the air.

Micrococcus luteus, Schröter.—Cocci similar in size to the above, elliptical, with highly refractive cell contents. They form yellow drops of 1—3 mm. diam. on boiled potato, and a thick, wrinkled, yellow skin on nutrient liquids. The colouring matter is insoluble in water, and unchanged by sulphuric acid or alkalies.

Micrococcus rosaceus.—Cocci forming pink colonies, and a rose-coloured growth on the surface of nutrient agar-agar. Observed contaminating an old cultivation (Plate X., Fig. 2).

Micrococcus hæmatodes, Zopf.—Cocci, which, cultivated on boiled white of egg in a damp chamber in the incubator, form a red layer. The reaction of the colouring matter is similar to that produced by *Micrococcus prodigiosus*. They have been observed in human sweat, especially from the axilla, colouring the surrounding parts and the linen an intense brick or blood-red colour.*

Micrococcus candidus, Cohn.—Cocci forming snow-white points and spots, upon slices of cooked potato. Possibly identical with the following :—

Micrococcus candicans, Flügge.—Cocci which collect in masses. In plate-cultivations they form in two or three days milk-white colonies. Cultivated in test-tubes they form a white nail-shaped cultivation. They were isolated from contaminated plate-cultivations.

* Babes, "Vom Rothen Schweiss," *Biol. Centrabl.*, Bd. 2. 1882.

Micrococcus fœtidus, Rosenbach.—Small oval cocci. Cultivated in agar-agar they develop gas-bubbles and a fœtid odour. Isolated from carious teeth. Possibly closely allied to, if not identical with, the following :—

Micrococcus crepusculum, Cohn (*Monus crepusculum*, Ehrenberg. *Mikrokokken in faulenden Substraten*, Flügge).—Round or short oval cells, scarcely 2μ in diam. ; singly or in zooglœa. They occur in various infusions and putrefying fluids in company with *Bacterium termo*.

Micrococcus cinnabareus, Flügge. — Large cocci occurring in twos, threes, and fours. In plate-cultivations they grow very slowly, forming punctiform colonies, in colour bright red at first, and afterwards reddish-brown. In test-tube cultivations they form on the surface of the gelatine a heaped-up, red-coloured growth. Found contaminating old cultivations.

Micrococcus flavus liquefaciens, Flügge.—Cocci, diplococci, and zooglœa. On plate-cultivations they form yellowish colonies, and in test-tubes yellowish beads, which become confluent and rapidly liquefy the gelatine.

Micrococcus flavus tardigradus, Flügge.—Cocci forming chrome-yellow colonies. Cultivated in test-tubes they form yellowish beads in the needle track, which remain isolated, and do not liquefy the gelatine. Isolated from contaminated cultures.

Micrococcus versicolor, Flügge.—Small cocci, which form iridescent colonies. In test-tubes they grow in the form of yellowish beads, and develop an iridescent layer on the surface.

Micrococcus viticulosus, Flügge.—Oval cocci, 1.2μ in length, and 1μ in width. In plate-cultivations the colonies differ when embedded in the nutrient medium and when growing on the surface. The characteristic appearance consists of a delicate network, which is visible also in test-tube cultivations in the track of the needle. On the surface of the medium they form a viscous layer. They were isolated from contaminated cultivations.

Micrococcus lacteus faviformis, Bumm and Bockart.—Cocci 1.25μ in diam. Cultivated in gelatine they form milk-white confluent colonies, and preparations made from the cultivations have a characteristic honeycomb appearance. Isolated from vaginal secretions, and from sputum.

Micrococcus fulvus, Cohn.—Cocci round 1.5μ in diam., frequently in pairs. They form rusty-red conical drops of a firm consistency, and about $.5$ mm. diam., on horse-dung.

Micrococcus viscosus, Pasteur.—Globular cells $.2 \mu$ in diam., singly or in chains. These and allied forms have been considered to be the cause of mucoid fermentation in wine and beer* (*vin filant, bière malade*).

Micrococcus coronatus (*Streptococcus coronatus*, Flügge).—Cocci 1μ in diam., singly, in short chains, and in zooglœa. In plate-cultivations the colonies have a characteristic halo, formed by the liquefaction of the gelatine around the colony. Isolated from the air.

Micrococcus radiatus (*Streptococcus radiatus*, Flügge).—Cocci less than 1μ in diam., singly, and in short chains. They rapidly form whitish colonies with a yellowish-green sheen. They liquefy the gelatine, the colonies sinking down, and after one or two days developing a circlet of rays. A peculiar ray-like appearance is characteristic also of the growth in test-tubes. Isolated from contaminated plate-cultivations.

Micrococcus flavus desidens (*Streptococcus flavus desidens*, Flügge).—Cocci, diplococci, and short chains. They form yellowish-white colonies, which gradually sink down in the gelatine. In test-tubes they form china-white, confluent masses in the tract of the needle, and on the surface a yellowish-brown slimy layer. Isolated from contaminated plate-cultivations.

* Pasteur, *Etudes sur le Vin; sur la Bière*. 1866; 1876.

Genus V.—Ascococcus.

SPECIES.

UNASSOCIATED WITH DISEASE:—

Ascococcus Billrothii . Zymogenic saprophyte.

Ascococcus Billrothii.—Small globular cocci, united into characteristic colonies. They form on the surface of nourishing fluids a cream-like skin, divisible into an enormous number of globular or oval families. Each family is surrounded by a thick capsule of cartilaginous consistency. In a




FIG. 76.—ASCOCOCCUS BILLROTHII [after Cohn].

solution containing acid tartrate of ammonia the fungi generate butyric acid, and change the originally acid fluid into an alkaline one. They were first observed on putrid broth, and later on ordinary nourishing solutions; they also readily develop upon damp slices of boiled roots, carrots, beetroots, etc.

METHODS OF STAINING COCCI.

Cocci stain well with watery solutions of gentian-violet, methyl-violet, fuchsine, methylene blue, and bismarck-brown. For examining cocci in liquids such as pus or blood, or in cultivations in solid media, a little of the material should be spread out on a cover-glass (page 65), and stained with a drop or two of a watery solution of fuchsine or methyl-violet. The former is especially recommended for staining *Merismopedia gonorrhœæ*.

For a zooglœa, or pellicle of micrococci, Klein recommends transference bodily to a watch-glass containing the dye, leaving it there till deeply tinted, then taking it out with a needle, washing in water, and then in alcohol till excess of colour is removed. It must then be transferred to a glass slide, spread well out, and a drop of clove-oil placed on it; after a minute or two the clove-oil is drained off, a drop of Canada balsam added, and covered with a cover-glass.*

Cocci in the tissues may be stained by immersing the sections in an aqueous solution of gentian-violet, or in aniline-gentian-violet solution, then rinsing in water, decolorising in alcohol, treating with clove-oil, and preserving in balsam (p. 76); or, after washing with alcohol, they may be rinsed with water, and stained for half an hour with Weigert's picrocarmine. From this they are again removed to water, then to alcohol, clove-oil, and Canada balsam.

The method of Gram is much more satisfactory (p. 76, Plate II., Figs. 1 and 2). Sections should be examined with and without a contrast stain. The *after-stain* most commonly employed is eosin. The sections after the process of decolorisation should be placed in a weak alcoholic solution of eosin (two or three drops of a concentrated alcoholic solution added to a watch-glassful of alcohol), till stained a delicate pink. They are then rinsed in

* Klein, *Micro-organisms and Disease*. 1885.

fresh alcohol, treated with clove-oil, and preserved in Canada balsam.

Sections containing cocci of osteomyelitis may be *after-stained* with weak solution of vesuvin. Safranin and picrolithium-carmin may also be used as contrast stains (p. 78).

Nuclear stains, such as carmine, hæmatoxylin, may also be employed. Sections may be left one minute in Grenacher's solution, then washed out in weakly acidulated alcohol (2—1000); and finally treated in the usual way, with alcohol, oil of cloves, and balsam.

Sections containing *Micrococcus tetragonus* are best stained with Gram's method and eosin (Plate XXIV., Fig. 2), but they may also be treated by the method of Friedländer, to demonstrate their capsules (p. 263).

To stain the cocci of *rabbit-septicæmia* in the tissues, place the sections twenty-four hours in Löffler's solution, wash in water faintly acidulated with acetic acid, then treat with alcohol, oil of cloves, and balsam.

GROUP II.—BACTERIACEÆ.

Genus I. *Bacterium*.—Cocci and rods, or only rods, which are joined together to form threads. Spore-formation absent or unknown.

Genus II. *Spirillum*.—Threads screw-form, made up of rods (long or short) only, or of rods and cocci. Spore-formation absent or unknown.

Genus III. *Leuconostoc*.—Cocci and rods. Spore-formation present in cocci.

Genus IV. *Bacillus*.—Cocci and rods, or rods only, forming straight or twisted threads. Spore-formation present either in rods or cocci.

Genus V. *Vibrio*.—Threads screw-form in long or short links. Spore-formation present.

Genus VI. *Clostridium*.—Same as bacillus, but spore-formation takes place in characteristically enlarged rods.

Genus I.—Bacterium.

SPECIES.

ASSOCIATED WITH DISEASE :—

In man . . .	}	Bacterium pneumoniae crouposae . . .	Pathogenic (?); possibly only saprophytic in man, pathogenic in animals.
		Bacterium pseudo-pneumonicum . . .	Saprophytic in man, pathogenic in animals.
		Bacterium Neapolitanum . . .	" "
		Bacterium in rhinoscleroma . . .	Pathogenic in man (?).
		Bacterium in diphtheria . . .	Saprophytic in man (?), pathogenic in animals.
In animals . . .	}	Bacterium saprogenes . . .	Saprophytic in man, pathogenic in animals.
		Bacterium decalvans . . .	Saprophytic.
		Bacterium in diphtheria of calves . . .	Pathogenic (?).
		Bacterium of diphtheria of pigeons . . .	Pathogenic.
		Bacterium cholerae gallinarum . . .	"
		Bacterium septicum agrigenum . . .	"
		Bacterium of septicæmia in rabbits . . .	"
		Bacterium of Davaine's septicæmia . . .	"
		Bacterium septicum sputigenum . . .	"
		Bacterium crassum sputigenum . . .	"
		Bacterium pneumonicum agile . . .	"
		Bacterium oxytocum perniciosum . . .	"
In plants . . .	}	Bacterium cavicida . . .	"
		Bacterium coli commune . . .	"
		Bacterium lactis aërogenes . . .	"
		Panhistophyton ovatum . . .	"
		Bacterium hyacinthi . . .	Pathogenic (?).

UNASSOCIATED WITH DISEASE :—

}	Bacterium synxanthum . . .	} Chromogenic saprophytes.
	Bacterium indicum . . .	
	Bacterium rubrum . . .	
	Bacterium prodigiosum . . .	
	Bacterium luteum . . .	
	Bacterium violaceum . . .	
	Bacterium brunneum . . .	
	Bacterium fluorescens putidum . . .	
	Bacterium fluorescens liquefaciens . . .	
	Bacterium ureæ . . .	
}	Bacterium aceti . . .	} Zymogenic saprophytes.
	Bacterium Pasteurianum . . .	
	Bacterium liodermos . . .	
	Bacterium multipediculum . . .	
	Bacterium ramosum liquefaciens . . .	
	Bacterium Zopfii . . .	
	Bacterium merismopedioides . . .	
	Bacterium Pflügeri . . .	
	Bacterium photometricum . . .	
	Bacterium litoreum . . .	
}	Bacterium fusiforme . . .	} Simple saprophytes.
	Bacterium navicula . . .	
	Proteus vulgaris . . .	
	Proteus mirabilis . . .	
	Proteus Zenkeri . . .	
	Bacterium termo . . .	
	Bacterium lineola . . .	

Bacterium pneumoniae crouposae (*Pneumococcus*, Friedländer).—Cocci ellipsoidal and round, singly, or in pairs (diplococci), rods and thread-forms. The cell-membrane thickens, and develops into a gelatinous capsule, which is round if the coccus is single, and ellipsoidal if the cocci occur in pairs or in rod-forms (Fig. 77, Plate I., Fig. 5). Cultivated in a test-tube of nutrient gelatine they grow along the needle track in the form of a round-headed nail (Plate V., Fig. 2), without liquefaction




FIG. 77.—BACTERIUM PNEUMONIAE CROUPOSÆ, FROM PLEURAL CAVITY OF A MOUSE, $\times 1500$. A, B. Thread-forms. C, D, E. Short rod-forms. G. Diplococci. H. Cocci. I. Streptococci. [After Zopf.]

of the gelatine. The cocci when artificially cultivated have no capsule, but it again appears after their injection into animals. The cocci can also be cultivated on blood serum and on boiled potatoes. They occur in pneumonic exudation.* Inoculation of dogs with a cultivation of the cocci occasionally gave positive results; but in rabbits no results

* Friedländer, *Fortschr. d. Med.* 1883.

followed. Guinea-pigs proved to be susceptible in some cases; but thirty-two mice, after injection of a cultivation diffused in sterilised water into the lungs, died without exception. The lungs were red and solid, and contained the cocci, which were also present in the blood, and in enormous numbers in the pleural exudation. Inhalation experiments by spraying the cocci diffused in water into mouse cages succeeded in producing pneumonia and pleurisy in three out of ten mice. The nail-shaped cultivation is not always produced, nor are these conclusions accepted by all investigators.*

METHODS OF STAINING THE BACTERIA OF PNEUMONIA.

(*Pneumonie-Coccen*, Friedländer.)

Cover-glass preparations (p. 65) of pneumonic sputum or exudation may be treated as follows:—

(a) Stain by the method of Gram, and after-stain with eosin (p. 76).

(b) Treat with acetic acid, then stain with gentian-violet or bismarck-brown. Examine in distilled water, or dry and preserve in Canada balsam.

(c) Float them on weak solutions of the aniline dyes twenty-four hours; differentiation between coccus and capsule is thus obtained.

(d) Stain with osmic acid; the contour of the capsules is brought out.

Sections of pneumonic lung should be stained by—

(a) Method of Gram.

(b) *Method of Friedländer*. This method is employed to demonstrate the capsules in tissue sections. It consists

* Klein, *Micro-organisms and Disease*. 1885.

in placing the sections twenty-four hours in the following solution :—

Fuchsine	1
Distilled water	100
Alcohol	5
Glacial acetic acid	2

They are then rinsed with alcohol, transferred for a couple of minutes to a 2 per cent. solution of acetic acid, and in the usual way treated with alcohol and oil of cloves, and preserved in Canada balsam.

Bacterium pseudo-pneumonicum (*Bacillus pseudo-pneumonicus*, Passet).—Cocci round, oval, and occasionally elongated, similar to the bacterium of pneumonia. The oval forms are $.87 \mu$ in width, and 1.16μ in length. The colonies on plates appear in twenty-four hours as white dots; in test-tubes the growth develops as a greyish-white layer. If injected into the pleura they set up pleuritis, and into the abdomen peritonitis, in mice, rats, and guinea-pigs. Subcutaneous inoculation produces septicæmia in mice, and abscesses in rats, guinea-pigs, and rabbits. Inhalation experiments gave no results. They were isolated from pus.

Bacteria Neapolitanum (*Bacillus Neapolitanus*, Emmerich).—Short rods with rounded ends. In width $.9 \mu$ (Fig. 78). They form circular colonies, which later become irregular, granular, strongly refractive, and of a yellowish-brown colour. By introducing a large quantity into small animals changes were produced in the intestines with an analogy to the post-mortem appearances of cholera. They are probably identical with bacteria found in healthy

fæces. They were isolated from some cases of cholera at Naples.

Bacterium of Rhinoscleroma (*Bacillus of Rhinoscleroma*, Cornil and Alvarez*).—Cocci and




FIG. 78.—BACTERIUM NEAPOLITANUM, $\times 700$. (a) From intestinal contents in a case of cholera; (b) From peritoneal fluid of an inoculated guinea-pig [after Emmerich].

short rods, $1.5-3 \mu$ in length, $.5-.8 \mu$ thick. Deeply coloured points or granules may occur in the course of the rods when stained, but it is very doubtful whether these can be considered as spores. The bacteria are encapsuled, the capsule being round when enclosing a coccus, and ovoid when enclosing



FIG. 79.—BACTERIA OF RHINOSCLEROMA, $\times 1400$. Encapsuled cocci, diplococci, and short and long rod-forms [after Cornil].

a rod (Fig. 79). The capsule is composed of a tough resisting substance; two or more capsules may unite by fusion, enclosing two or three, or a great number of rods. The bacilli were observed in sections of a tumour, *rhinoscleroma*, which develops

* Cornil and Babes, *Les Bactéries*. 1885.

on the lip and on the nasal and pharyngo-laryngeal regions.

METHOD OF STAINING THE BACILLUS OF RHINO-SCLEROMA.

Method of Cornil and Alvarez :—

Sections are immersed in a solution of methyl-violet (B) for twenty-four to forty-eight hours, with or without the addition of aniline-water; are then decolorised after treatment with the solution of iodine in iodide of potassium. If the sections are left to decolorise in alcohol for forty-eight hours, the capsule is rendered visible.

Bacterium in diphtheria of man (*Bacillus of diphtheria*, Löffler).—Rods about the same length as the tubercle bacillus, but about twice as thick; the longer ones consist of single individuals linked together. Spores not observed. They were cultivated in a mixture consisting of three parts of calf's or lamb's blood serum, to which was added one part of neutralised veal broth, containing 1 per cent. peptone, 1 per cent. grape sugar, $1\frac{1}{2}$ per cent. common salt. Cultivated in 5 per cent. gelatine at 20—22° C., the rods developed into irregular involution-forms. Inoculation gave doubtful results. The bacillus was isolated from diphtheritic membrane.* They were particularly noticeable in those typical cases characterised by a thick false membrane, extending over the fauces, larynx, and trachea. They occupied

* Löffler, *Mittheil. a.d. K. Gesundheitsamte* (*Microparasites in Disease*, New Syd. Society).

the deeper layers below the masses of bacteria which are found on the surface, such as the streptococci already described (p. 231).

METHOD OF STAINING THE BACTERIA IN DIPHTHERIA.

Löffler's Method:—

Sections are placed in Löffler's solution for a few minutes, and excess of stain removed by $\frac{1}{2}$ per cent. solution of acetic acid. They are then treated with alcohol and cedar-oil, and mounted in Canada balsam.

Bacterium saprogenes (*Bacillus saprogenes* No. 3, Rosenbach).—Rods isolated from the putrid marrow of a case of compound fracture. Cultivated on nutrient agar-agar, an ash-grey, almost liquid culture is developed, with a strong characteristic odour of putrefaction. Injected into the knee-joint or abdomen of a rabbit, an opaque, yellowish-green infiltration resulted (*vide Bacillus saprogenes*, p. 346).

Bacterium decalvans, Thin.—Cocci, singly or in pairs, 1.6μ in length. Observed in the roots of the hair in cases of *Alopecia areata*.

Bacterium in diphtheria of calves (*Bacillus vitulorum*, Löffler).—Rods about five or six times as long as wide, mostly united in long threads. A piece of tissue placed on blood serum developed a white layer composed of the bacteria. Successive generations were not obtainable. Mice inoculated directly from the calf died of a characteristic illness, and the same long bacteria were again found in the inoculated animals accompanying widespread infiltration, starting from the point of inoculation. Inoculation of guinea-pigs and rabbits gave doubtful

results. The bacteria were found in the deeper stratum of the diphtheritic patches.

Bacterium of diphtheria of pigeons (*Bacillus columbarum*, Löffler).—Short rods with rounded ends, mostly in irregular masses. In plate-cultivations on nutrient gelatine they formed whitish patches on the surface, and compact, ball-like masses when embedded in the gelatine. They were also cultivated on blood serum and potatoes. Subcutaneous inoculations in pigeons with a pure cultivation produced local inflammation and necrosis; inoculation in the mucous membrane of the mouth gave the appearances of the original disease. Other animals were only locally affected, except mice, in which characteristic symptoms and death resulted. They were isolated from the diphtheritic exudations in pigeons, and in sections were found in the vessels of the lungs and liver.

Bacterium cholerae gallinarum (*Micrococcus cholerae gallinarum*, Zopf. *Bacterium of Fowl-cholera*. *Microbe du choléra des poules*).—Cocci 2—3 μ in diam., short rods staining deeply at either pole, and longer beaded rods (Figs. 80, 81). In the tissues they appear mostly as rods 2—3 μ in length and 5 μ in diam., with their extremities stained more deeply than their middle* (*vide* p. 165). When cultivated by introducing a drop of the infected blood into sterile chicken broth, a number of round bodies,

* Cornil and Babes, *Les Bactéries*.

undergoing rapid movement and as a rule united as diplococci, or elongated and contracted in the middle, appear in the broth, which is at first slightly milky, but becomes limpid, and the microbes at the same time pass into a finely granular state. From this,




FIG. 80.—BACTERIUM OF CHICKEN CHOLERA; BLOOD OF INOCULATED HEN, X 1200.

however, fresh cultures can still be started. Cultivated in a test-tube of nutrient gelatine, after from three days to a week there develops along the needle track a fine, almost imperceptible, greyish thread



FIG. 81.—BACTERIUM OF CHICKEN CHOLERA, FROM MUSCLE JUICE OF INFECTED HEN, X 2500 [from a Photograph].

without liquefaction of the gelatine (Plate IV., Fig. 2). The growth is exceedingly scanty, even after several weeks.

Fowls suffering from the disease usually die very rapidly. In the less acute cases they are somnolent, weak in their legs, and their wings trail. They suffer from diarrhoea, and pass into a state of sopor and die. The micro-organisms are found in large

numbers in the blood and organs after death, and in the intestinal discharges.

A drop of the broth injected into the connective tissue in the region of the pectoral muscles causes the death of the fowl the following day, with characteristic pathological changes.* If a culture be kept for some time, and a fowl be then inoculated with it, instead of death only local changes are produced, and the fowl is protected against the action of a virulent culture; thus affording an example of so-called *mitigation of the virus*.† The microbe is ærobic, and its toxic effect has been supposed to be due to the abstraction of oxygen from the blood producing asphyxia.

Bacterium septicum agrigenum (*Bacillus septicus agrigenus*, Nicolaier).—Cells morphologically similar to the microbe of chicken cholera. The colonies on plate-cultivations have a yellowish-brown centre, with a greyish-yellow zone. In test-tube cultivations the appearances are not characteristic. They are pathogenic in mice and in rabbits. The organs show no characteristic post-mortem appearances, but the bacteria abound in the blood. They were isolated from earth.

Bacterium of septicæmia in rabbits (*Bacillus cuniculicida*, Koch).—Short rods, slightly pointed at both ends; in width $\cdot 6 \mu$ — $\cdot 7 \mu$, in length $1\cdot 4 \mu$. They stain deeply at the ends, leaving an uncoloured

* Cornil, "Observ. Hist. sur les Lésions des Muscles déterminées par l'injection du Microbe du Choléra des Poules" (*Archives de Physiologie*. 1882); Cornil and Babes, *Les Bactéries*. 1885.

† Pasteur, "Sur le Choléra des Poules," *Compt. Rendus*. 1880.

interval in the middle (Fig. 82), an appearance which must be distinguished from a diplococcus or figure of eight. Two or more bacteria may be linked together in a chain. They may be cultivated in bouillon, blood serum, and nutrient gelatine. In plate-cultivations of the latter, they produce dot-like colonies, and in test-tubes little spherical masses in the needle track, and a layer on the free surface. The smallest quantity inoculated subcutaneously or in the cornea of a rabbit produces a rise of temperature and laboured breathing after 10—12 hours, and death in 16—20 hours. The



FIG. 82.—BACTERIUM OF RABBIT SEPTICÆMIA; BLOOD OF SPARROW, X 700 [after Koch].

spleen and lymphatic glands are found to be enlarged, and the lungs congested, but no extravasations, and no peritonitis. In the blood the characteristic rods abound, and in sections they are found in the vessels and capillaries. Mice and birds are very susceptible; guinea-pigs and white rats have an immunity. The disease was produced by inoculating rabbits with contaminated water (River Panke) and with putrid meat infusion.

Bacterium of Davaine's septicæmia.—Rods similar to the bacteria described by Koch. They were also found in the blood of rabbits suffering

from septicæmia, which, however, differed from Koch's septicæmia in that guinea-pigs were susceptible and pigeons immune.

Bacterium septicum sputigenum (*Microbe de salive*, Pasteur. *Micrococcus Pasteuri*, Sternberg.* *Bacillus septicus sputigenus*, Fränkel).—Cocci oval, singly, in pairs, and in chains; often lanceolate or rod-shaped; encapsuled. They grow well in broth, and on agar-agar at 30° to 35° C. On the solid media they form a superficial, nearly transparent deposit of gelatinous consistence. They are pathogenic in rabbits, producing typical "sputum septicæmia." Fowls and dogs have an immunity, and guinea-pigs are less susceptible than rabbits. Mice die within forty-eight hours after being inoculated. The blood of an infected rabbit just dead is more potent than a liquid culture or than saliva containing the coccus. An animal which recovers after an injection of saliva is stated to be protected from the potent virus. The pathogenic power is modified by cultivation at a temperature between 39·5° and 40·5°. The organism has been supposed to be intimately associated with croupous pneumonia, but any exact relation cannot be considered as established. The organism differs from the *Bacterium pneumoniae crouposæ* in that it is pathogenic in rabbits, it can be directly isolated from rusty sputum, and it requires a temperature for

* Sternberg, *Stud. Biol. Lab., John Hopkins, Univ.* 11, No. 2, 1882. *Journal Royal Microscop. Society.* 1886.

its growth at which nutrient gelatine is liquefied. The cocci were first observed in the blood of a rabbit inoculated with healthy saliva, and again found in a rabbit which died after inoculation with the saliva of a child suffering from rabies. Later they were isolated from the blood of rabbits inoculated with the buccal secretions of different individuals, and were found to be constantly present in the rusty sputum of pneumonic patients.

Bacterium crassum sputigenum, Kreibohm.—Short thick rods with rounded ends. Colonies on plate-cultivations appear as clear grey-white points, which ultimately form greyish slimy drops. In test-tubes they develop very quickly a nail-shaped growth. They are fatal to mice, and after death are found in the blood, and in sections, more especially in the capillaries of the liver. Rabbits die of septicæmia after intravenous injection. A large quantity of a cultivation injected into the circulation sets up fatal gastro-enteritis in rabbits and dogs in 3—10 hours. They were isolated from sputum.

Bacterium pneumonicum agile (*Bacillus pneumonicus agilis*, Schou).—Short thick rods, or almost elliptical cells, often two to four linked together. They form dark granular colonies, which after twenty-four hours commence to liquefy the gelatine; a movement is then visible in the centre of the colony, and an appearance of circumferential rays results. They grow also on blood

serum, bouillon, and potatoes. Cultures injected through the chest wall, or into the trachea, or administered by inhalation, set up pneumonia. They were isolated from pneumonic lungs of rabbit.

Bacterium oxytocum perniciosum (*Bacillus oxytocus perniciosus*, Wyssokowitsch).—Short rods with rounded ends, somewhat shorter and thicker than the bacterium of sour milk. They form yellowish colonies, and in test-tubes develop a nail-shaped growth. Cultivated in milk they produced curdling, and an acid reaction. They were sometimes pathogenic in rabbits. Isolated from sour milk.

Bacterium cavicida (*Bacillus cavicida*, Brieger).—Very small rods, about twice as long as broad. They form colonies in the form of whitish concentric rings. On potatoes they develop dirty yellow tufts. They are very fatal to guinea-pigs. Isolated from human fæces.

Bacterium coli commune, Escherich.—Short slightly curved rods, 1.5μ in length, $.3$ — $.4 \mu$ thick, colonies yellowish and granular. They develop a white scum on agar-agar and blood serum. Fatal to guinea-pigs and rabbits, when inoculated intravenously. Isolated from fæces of infants fed exclusively on mother's milk.

Bacterium lactis aërogenes, Escherich.—Short rods with rounded ends, 1.4 — 2μ long, $.5 \mu$ wide. Cultivations in gelatine resemble the bacterium of pneumonia. They produce fermentation in milk and in solution of grape-sugar. Pathogenic effects similar to the above. Isolated from the same source.

Panhistophyton ovatum, Lebert (*Nosema bombycis*, *Micrococcus ovatus*, *Corpuscles du ver à soie*).—Shining oval cocci, 2 — 3μ long, 2μ wide, singly and in pairs, or masses;* or rods, 2.5μ thick, and twice as long.† They multiply by subdivision. They were experimentally proved

* Flüge, *Fermente und Mikro-Parasiten*. 1883.

† Zopf, *Die Spaltpilze*.

to be the cause of *pébrine*, *gattine*, *maladie des corpuscles* or *Flecksucht*; and were discovered in the organs of diseased silkworms, as well as in the pupæ, moths, and eggs.

Bacterium hyacinthi, Wakker.—Cells resembling *Bacterium termo*. Observed in the yellow slime of diseased hyacinth bulbs.

Bacterium synxanthum, Ehrenberg (*Bacterium xanthinum*. *Bacterium of yellow milk*).—Cocci $\cdot 7$ — $1\ \mu$ in length, and rod-forms.* They produce a yellow colour in boiled milk, which at first becomes acid, and then strongly alkaline. They also occur on boiled potatoes, carrots, etc., where they form small lemon-yellow masses. The colouring matter soluble in water, insoluble in ether and alcohol, unchanged by alkalis, decolorised by acids. It is similar to yellow aniline colours both spectroscopically and in ordinary reactions.

Bacterium indicum (*Micrococcus indicus*, Koch. *Bacillus indicus*, Flügge).—Very short rods with rounded ends. In plate-cultivations on nutrient agar-agar, the colonies have a scarlet tint. They are round, ovoid, or spindle-shaped, and have characteristic granular margins (Fig. 83). Grown upon nutrient agar-agar in a test-tube, the appearances are very characteristic. In a pure cultivation a brilliant, vermilion-coloured reticulated pellicle develops on the surface (Plate VIII., Fig. 1). In the track of the needle beneath the surface no pigment is formed (Plate VII., Fig. 2). Cultivated in nutrient gelatine they liquefy the medium, and colour it crimson. The growth, of a darker crimson hue, subsides to the bottom of the tube. Upon sterilised potato they form a vermilion layer (Plate XVI., Fig. 2).

* Zopf, *Die Spaltpilze*.

Bacterium rubrum, Frank.—Minute motile rods, singly, in twos, and fours. They were observed on boiled rice, where they develop a brick-red pigment.

Bacterium prodigiosum (*Micrococcus prodigiosus*, *Bacillus prodigiosus*, "Blood-rain," "Bleeding host").—Very short rods with rounded ends, and thread-forms, $\cdot 5-1 \mu$ in width, forming at first rose-red, and then blood-red zoogloea. They grow luxuriantly when cultivated on sterilised potatoes (Plate XIV., Fig. 1), and on the sloping surface of




FIG. 83.—BACTERIUM INDICUM; COLONIES ON NUTRIENT AGAR-AGAR, X 60.

nutrient agar-agar (Plate VIII., Fig. 3). They appear occasionally on bread, boiled rice, and starch-paste, and more rarely on boiled white of egg and meat. Milk sometimes becomes coloured blood-red by the growth of this fungus, an appearance formerly attributed to a disease of the cow.

In Paris, in 1843, the fungus was peculiarly prevalent, attacking especially the bread produced in the military bakehouses.

The cells themselves are colourless. The colouring matter resembles fuchsine; it is insoluble in water, but soluble in alcohol. The addition of

acids changes it to a carmine red, and of alkalies to a yellow colour.

Bacterium luteum (*Bacillus luteus*, Flügge).—Short immotile rods. Colonies irregular in form, appear brownish under a low power, but macroscopically yellow. In test-tube cultivations they form a yellow growth without liquefying the gelatine. They occur contaminating plate-cultivations.

Bacterium violaceum, Bergonzini.—Cells similar to *Bacterium termo*, $0.6-1 \mu$ thick, $2-3 \mu$ long. They occur on white of egg, forming a violet pigment.

Bacterium brunneum, Schröter.—Motile rods, producing a brown colour. They were observed on a rotting infusion of maize.

Bacterium fluorescens putidum (*Bacillus fluorescens putidus*, Flügge).—Short rods with rounded ends; motile; spore-formation not known. They form small dark colonies with a greenish sheen and penetrating odour. In test-tubes they produce a pale-grey turbidity, and after three days colour the medium with a greenish tinge spreading down from above. On potatoes they rapidly develop a brownish layer. They occur on decomposing substances, producing a greenish coloration.

Bacterium fluorescens liquefaciens (*Bacillus fluorescens liquefaciens*, Flügge).—Short rods with rounded ends. Colonies on plates develop an iridescence around them. In test-tubes a similar iridescent sheen is produced. On potatoes they develop a brownish layer.

Bacterium ureæ (*Micrococcus ureæ*, Cohn).—Cocci $1.25-2 \mu$ in diam., singly or in chains, and rods. The rods split up by division into chains of cocci, and the latter are finally set free. The cocci increase further by subdivision, and a jelly-like membrane develops around them. Masses of cocci

exist in the form of irregular or roundish lumps. Cultivations, after twenty-four hours, consist exclusively of rods; after forty-eight hours, of cocci chains; and in fourteen days, of zooglœa; the cocci transplanted into fresh nourishing solution again grow into rods. These observations point to the existence of a pleomorphic species, *Bacterium ureæ*; and the former nomenclature, *Micrococcus ureæ*, must be regarded as untenable. They are ærobic; occurring in urine they set up ammoniacal fermentation, converting urea into carbonate of ammonia.* Rods, 2 μ long and 1 μ wide, have been isolated from stale urine (*Bacillus ureæ*, Leube), which also most energetically cause the ammoniacal fermentation of urine.

Bacterium aceti.—Cocci, short rods, long rods, leptothrix-forms, and zooglœa. Cocci and short rods may occur in the same thread. The long rods and threads may develop irregular swellings, so-called involution-forms, which have a thickened membrane and a grey colour. The effect of the action of this microbe is to oxidise alcohol in wine and other fruit juices into vinegar. The masses of zooglœa united together form a membranous layer which must not be mistaken for the pellicle formed by *Saccharomyces mycoderma*. The latter prepares the medium for the action of the *Bacterium aceti*.

Bacterium Pasteurianum, Hansen.—Morphologically similar to *Bacterium aceti*, but the cells contain

* Zoph, *Die Spaltpilze*. 1885.

a starch-like substance, which is turned blue by iodine. They occur in beer-wort.

Bacterium liodermos (*Bacillus liodermos*, Flügge. *Potato bacterium*).—Short rods with rounded ends, motile. On plate-cultivations the colonies appear as small white pellicles floating on liquefied gelatine. In test-tubes the gelatine is liquefied, and the growth sinks down in flocculent masses. They occur on potatoes, forming a smooth shining layer, which ultimately becomes crumpled.

Bacterium multipedunculatum (*Bacillus multipedunculatus*, Flügge).—Long slender rods. They form peculiar insect-like colonies on plate-cultivations. In test-tubes the appearance is less characteristic. They occur on potatoes, forming a dirty yellow growth.

Bacterium ramosum liquefaciens (*Bacillus ramosus liquefaciens*, Flügge).—Rods, slowly motile. They form characteristic colonies on plate-cultivations. The colonies gradually sink down, forming a well-marked funnel with later an appearance of concentric rings. In test-tubes the funnel-shaped liquefaction sends off rays into the surrounding gelatine. They occur occasionally, contaminating cultivations.

Bacterium Zopfii, Kurth.—Cocci, 1—1.25 μ in diameter; rods and threads. Cultivated in a streak on nutrient gelatine spread out on a glass slide, a peculiar development takes place. In twenty-four hours after inoculation threads have developed; in forty-eight hours windings of the threads are observed, and in six days the threads have broken up into cocci (Fig. 84). They were observed in the intestine of fowls, especially in the contents of the vermiform appendix. Inoculation of rabbits was followed by negative results. Identical with *Bacillus figurans* (Crookshank).

Bacterium merismopedioides, Zopf.—Forms threads 1—1.5 μ in thickness; these subdivide into long rods, short rods, and finally into cocci. The cocci divide first in one and subsequently in two directions, forming characteristic groups, which




FIG. 84.—BACTERIUM ZOPFII. SUCCESSIVE CHANGES IN THE SAME THREAD, $\times 740$: (a) A thread-form, (b) breaking up into rod-forms, (c) into cocci [after Kurth].

appear like merismopedia. These groups may eventually consist of 64×64 cells or more, and ultimately form zooglœa. The cocci develop again into rods and threads. They were observed in water containing putrefying substances (River Panke Berlin).*

* Zopf, *Die Spaltpilze*. 1885.

Bacterium Pflügeri, Ludwig.—Large, round cocci, mostly in zooglœa, and thread-forms composed of rods. They can be cultivated on boiled white of egg and potatoes. They were observed to produce phosphorescence in putrid fish and meat. Gelatine not liquefied.

Bacterium photometricum, Engelmann.—Cells slightly reddish in colour, motile. The movements are stated to depend on light.

Bacterium litoreum, Warming.—Cells ellipsoidal, 2—6 μ long, 1·2—2·4 μ wide, occur singly in sea-water, never as chains or zooglœa.

Bacterium fusiforme, Warming.—Cells spindle-shaped, with pointed ends, 2·5 μ long and ·5—·8 μ thick. Observed as a spongy layer on the surface of sea-water.

Bacterium navicula, Reinke and Berthold.—Cells spindle-form or ellipsoidal, including motile and non-motile forms. They have one or more dark spots, which may be coloured blue by iodine. They have been observed in rotting potatoes.

Proteus vulgaris.—This and the two following species have been isolated* from putrefying meat infusion, and are stated to be intimately connected with the process of putrefaction. In the history of their development coccoid, bacterioid, spindle-form, spirular, and involution forms have been described. In *Proteus vulgaris* the bacteria vary in size; some measure 4 μ in length, and are almost as broad as long, and others vary from ·94—1·25 μ long and ·42—·63 μ wide. They are actively motile, and cultivated on nutrient gelatine they convert it into a turbid, greyish-white liquid. If cultivated in a capsule containing 5 per cent. of nutrient gelatine, a few hours after inoculation the most characteristic

* Hauser, *Ueber Fäulniss-Bakterien*. 1885.

movements of the individual bacilli are observed on the surface of the nutrient gelatine, although at this early stage no superficial liquefaction can be detected. Probably the movements depend upon the existence of a thin liquid layer, as they are not observed if the nutrient medium contains 10 per cent. of gelatine.

Proteus mirabilis.—Cocci $\cdot 4 \mu$ — $\cdot 9 \mu$. They occur singly and in zooglœa, and sometimes in tetrads, pairs, chains, or as short rods in twos resembling *Bacterium termo*; in fact, in all conceivable transition forms. Cultivated on nutrient gelatine they form a thick, whitish layer in concentric circles, which in time liquefies the medium. Similar movements are observed in capsule-cultivations as in *Proteus vulgaris*.

Proteus Zenkeri.—Cocci, $\cdot 4 \mu$ in twos like *Bacterium termo*, and short rods $1\cdot65 \mu$ long. Cultivated on nutrient gelatine no liquefaction results, but a thick, whitish-grey layer is formed. The bacilli are motile, and the same phenomena are observed on the solid medium as in the other forms. In cover-glass impressions most varied groupings of the bacilli are seen, and also developmental and involution-forms.

The two following forms are only provisionally regarded as distinct species. They are both probably phase-forms of protean species.

Bacterium termo, Dujardin.—Short cylindrical

or oblong cells, 1.5μ long, $.5$ — $.7$ broad, generally occurring as dumb-bells. The cells have dark contents, invested by a thick membrane, and are provided with flagella, to which the characteristic movements are due (Plate I., Fig. 8). They are associated with putrefaction, invariably appearing in decomposing albuminous substances and liquids. A growth can be readily started by placing a piece of meat in water in a warm place. Cultivated in broth, they produce a turbidity, and on sterilised potatoes a slimy grey layer.

Bacterium lineola.—Cells 3.8μ — 5.2μ long, 1.5μ wide. They occur singly or in pairs, occasionally in zooglœa, but never in chains. The cells are provided with flagella, and contain strongly refringent contents. They resemble *Bacterium termo* in form and in movement, but are considerably larger. They occur in well water and stagnant water, and form slimy heaps on rotting potatoes, and zooglœa and pellicles on various infusions. Cultivated on nutrient agar-agar they form a semi-transparent growth (Plate X., Fig. 1).

Genus II.—*Spirillum*.

SPECIES.

ASSOCIATED WITH DISEASE :—

In man	{	Spirillum Obermeieri Pathogenic.
		Spirillum cholerae Asiaticæ	{ Pathogenic in man (?), possibly only saprophytic. Pathogenic in animals.
In animals	{	Spirillum Finkleri { Saprophytic in man. Pathogenic in animals.
		Spirillum sputigenum { Saprophytic. Pathogenic in animals.
		Spirillum tyrogenum }

UNASSOCIATED WITH DISEASE :—

Spirillum rubrum	Chromogenic saprophyte.
Spirillum plicatile	} Simple saprophytes.
Spirillum serpens	
Spirillum tenue	
Spirillum undula	
Spirillum volutans	
Spirillum Rosenbergi	
Spirillum attenuatum	
Spirillum leucomelaneum	

Spirillum Obermeieri (*Spirochaete Obermeieri*, Cohn. *Spirillum of Relapsing Fever*).—Threads similar to the *Spirillum plicatile*. In length mostly 16—40 μ , with screw-curves regular (Plate I., Fig. 19). They move very rapidly, and exhibit peculiar wave-like undulations. They have been observed in the blood of patients suffering from relapsing fever,* but never in the secretions. They only occur during the relapses, and are absent during the non-febrile intervals. Their number is variable, but usually is strikingly great. Outside the body, in blood serum and 50 per cent. salt solution, the threads preserve their movements. From analogy to the *Spirillum plicatile* it is presumed that these threads are composed of articulated rods and cocci. Monkeys have been inoculated with success from man,† but inoculations of mice, rabbits, sheep, and pigs give negative results.

The spirilla were found in the blood of the inoculated monkeys in great numbers, and also in the brain, lung, liver, kidney, spleen, and skin; and are believed to be the cause of the disease.

METHODS OF STAINING THE SPIRILLUM OBERMEIERI.

In cover-glass preparations of blood the spirilla stain strongly with fuchsine, methyl-violet, gentian-violet, or bismarck-brown.

In sections, brown aniline stains have been recommended.

* Obermeier, *Med. Centralb.* 1873.

† Carter, *Lancet.* 1879 and 1880. Koch, *Cohn's Beiträge.*

Spirillum cholerae Asiaticae (*Comma-bacillus*, Koch).—Curved rods, spirilla, and threads (Plate I., Fig. 18). The curved rods or commas are about half the length of a tubercle-bacillus. They occur isolated, or attached to each other, forming S-shaped organisms or longer screw-forms; the latter resembling the spirilla of relapsing fever. Finally they may develop into spirilliform threads. In old cultivations threads are found with bulgings or irregularities, which are called involution-forms (Plate I.,




FIG. 85.—COVER-GLASS PREPARATION OF THE EDGE OF A DROP OF MEAT INFUSION, containing a pure cultivation of comma-bacilli, with (a) spirilliform threads, $\times 600$ [after Koch].

Fig. 35).* The commas are actively motile; their movements and development into spirilla may be studied in drop-cultivations (Fig. 85). In plate-cultivations, at a temperature of from 16° — 20° C., the colonies develop as little specks (Fig. 86), which begin to be visible after about twenty-four hours. Examined with a low-power, and a small diaphragm, these colonies have the following characteristics. They appear as little masses, granular, and of a very faintly yellowish-red tinge, which have liquefied

* Compare also Van Ermengem, *Recherches sur le Microbe du Chol. Asiat.* 1885.

the gelatine, and sunk down to the bottom of the resulting excavations (Fig. 87).

In test-tubes of slightly alkaline nutrient gelatine (10 per cent.), the appearance of the growth




FIG. 86.—COLONIES OF COMMA-BACILLI ON NUTRIENT GELATINE, NATURAL SIZE [after Koch].

is very striking. It commences to be visible in about twenty-four hours. Liquefaction sets in very slowly, commencing at the top of the needle track around an enclosed bubble of air, and forming a funnel continuous with the lower part of the



FIG. 87.—COLONIES OF KOCH'S COMMA-BACILLI, $\times 60$; from a nutrient-gelatine plate-cultivation.

growth (Plate. IV., Fig. 1); the latter preserves for several days its resemblance to a white thread (Figs. 93 and 94).^{*} In about eight days, however, liquefaction takes place along the whole of the needle track.

On a sloping surface of agar-agar the cultivation

^{*} From *Remarks on Comma-bacillus of Koch*. *Lancet*. 1885.

develops as a white, semi-transparent layer, with well-defined margin. In potato-cultivations the microbe will only grow at the temperature of the blood (37° C.), forming a slightly brown, transparent layer. Inoculation of a cultivation of the bacillus in the duodenum of guinea-pigs, with* and without †




Fig. 88.



Fig. 89.

FIG. 88.—COVER-GLASS PREPARATION FROM THE CONTENTS OF A CHOLERA INTESTINE, $\times 600$. (a) Remains of the epithelial cells; (b) Comma-bacillus; (c) Group of comma-bacilli [after Koch].

FIG. 89.—COVER-GLASS PREPARATION OF CHOLERA DEJECTA IN DAMP LINEN (two days old), $\times 600$. Great proliferation of the bacilli with spirilla (a) [after Koch].

ligation of the bile-duct, has given positive results. More recently these results have been confirmed by the following method:—Five ccm. of a 5 per

* Nicati et Rietsch, *Com. à l'Académie de Médecine*. 1884.

† Van Ermengem, *Le Microbe du Choléra Asiatique*. 1885.

cent. solution of potash were injected into the stomach of a guinea-pig, and twenty minutes after 10 ccm. of a cultivation of comma-bacilli diffused in broth were similarly introduced. Simultaneously with the latter, an injection of tincture of opium was made into the abdominal cavity, in the propor-




FIG. 90.—SECTION OF THE MUCOUS MEMBRANE OF A CHOLERA INTESTINE, $\times 600$. A tubular gland (*a*) is divided transversely; in its interior (*b*) and between the epithelium and the basement membrane (*c*) are numerous comma-bacilli [after Koch].

tion of 1 ccm. for every 200 grammes' weight of the animal. Those who have had success with inoculation experiments maintain that choleraic symptoms were produced without any trace of peritonitis or putrid infection, and that the comma-bacilli of Koch were again found in the intestinal contents, and fresh cultivations established.

On the other hand, these results have been

disputed, the fatal effects of the inoculation attributed to septicæmic poisoning, and the proliferation of the bacilli considered to be dependent upon an abnormal condition of the intestines induced by the injection of tincture of opium.* It is, however, very




Fig. 91.

Fig. 92.

Fig. 93.

Fig. 94.

PURE CULTIVATIONS IN NUTRIENT GELATINE.

Fig. 91. Finkler's bacillus, twenty-four hours old.

Fig. 92. " " two days old.

Fig. 93. Koch's cholera-bacillus, twenty-four hours old.

Fig. 94. " " " two days old.

probable that these organisms, like several others which have been isolated from intestinal discharges, are truly pathogenic in the lower animals. The comma-bacilli were found in the superficial necrosed

* Klein, *Brit. Med. Journal, and Micro-organisms and Disease*. 1885. Lankester, *Nature*, xxxi.; *Nineteenth Century*. July, 1885. Klein and Gibbes, "An Inquiry into the Etiology of Asiatic Cholera." *Bluebook*, 1885.

layer of the intestine, in the mucous flakes and liquid contents of the intestinal canal of cases of Asiatic cholera* (Figs. 88, 90). It is stated that they were also detected in a tank which contained the water supply to a neighbourhood where cholera cases occurred; but comma-shaped organisms are commonly present in sewage-contaminated water (Fig. 95). The comma-bacilli are ærobie, and their development is arrested by deprivation of oxygen. They are destroyed by drying and the presence of various antiseptic substances. They are distinguished from all other comma-shaped organisms by the test of cultivation. The entirely different results obtained



FIG. 95.—COMMA-SHAPED ORGANISMS WITH OTHER BACTERIA IN SEWAGE-CONTAMINATED WATER, X 1200.

* At a meeting of the Physiological Society, May 15th, 1886, at Cambridge, a preliminary communication was made upon the investigations in Spain referred to in the first edition of this work. The observations made by Roy, Brown, and Sherrington rather tend, in the opinion of the author, to confirm Koch's views. Comma-bacilli were found to be present, in some cases, in enormous numbers, and the frequency of their occurrence led these observers to believe that they must bear some relation to the disease. At the same time, as they failed to find them in all cases, they regarded the existence of a causal relation as not proven. They failed to find the Naples bacterium or the small straight bacillus noted by Klein; but they drew attention to certain peculiar mycelium-like threads in the mucous membrane of the intestines. These organisms, however, judging from a preparation stained with methylene-blue which was exhibited at the meeting, appeared to the author to much more closely resemble some of the involution-forms of the comma-bacillus, *filaments à masses globuleuses*, figured by Van Ermengen, than anything else he had seen. Yet assuming these peculiar structures to belong as described to some species of Chytridiaceæ, it is very

in the case of the comma-bacilli of cholera nostras (Figs. 91 to 94), renders a thorough acquaintance with these bacilli of the greatest importance as an aid in diagnosis.

METHODS OF STAINING THE COMMA-BACILLI OF KOCH.

In cover-glass preparations they may be well stained in the ordinary way with an aqueous solution of methyl-violet or fuchsine, or by the rapid method, without passing through the flame (p. 67, Babes' method).

*Nicati and Rietsch's method.**

A small quantity of the stools or of the scraping of the intestinal mucous membrane is spread out on a glass slide and dried, then steeped during some seconds in sublimate solution or in osmic acid (1—100). It is then stained by immersion in fuchsine-aniline solution (1 or 2 grammes of Bâle fuchsine dissolved in a saturated aqueous solution of aniline), washed, dried, and mounted in Canada balsam.

In sections of the intestine their presence may be demonstrated by

(a) *Koch's method.†*

Sections of the intestine, which must be well hardened doubtful whether they can be considered to have any significance. Methylene-blue has been employed by Koch and others, including the author, for staining sections of the intestine from cholera cases, and had they been constantly present it is hardly possible that such striking objects could have been overlooked. Again, we must bear in mind that hyphomycetous fungi occasionally have been found to occur saprophytically in the intestinal canal as well as in the lungs, external auditory meatus, and elsewhere. We must wait, before expressing a more definite opinion, until the report of these observers is published in full.

* *Brit. Med. Journal*, Sept., 1885.

† *Berliner Klinische Woch.*, No. 31.

in absolute alcohol, are left for twenty-four hours in a strong watery solution of methylene-blue, or for a shorter time if the colour solution is warmed. Then treated in the usual way.

(b) *Babes' method*.*

Sections, preferably from a recent case of cholera, and made as soon as possible after death, are left for twenty-four hours in a watery solution of fuchsine (fabrique de Bâle), then washed in distilled water faintly acidulated with acetic acid, or in sublimate solution (1—1000), passed rapidly through alcohol and oil of cloves, dried with filter paper, and preserved in Canada balsam.

Spirillum Finkler-Prior (*Comma-bacillus in Cholera nostras*).—Curved rods thicker than the comma-bacillus of Koch, and spirilla. The colonies on plate-cultivations (Plates XII and XIII.) are very much larger than those of the comma-bacillus of Koch of the same age. They have the faintest yellowish-brown tinge, a well-defined border, and a distinctly granular appearance. They liquefy nutrient gelatine very rapidly, so that the first plate of a series is, as a rule, completely liquefied on the day following inoculation, and the second plate in two or three days more. In a test-tube cultivation in nutrient gelatine the appearances are especially characteristic; the gelatine is very rapidly liquefied along the whole track of the needle, so that the cultivation resembles a conical sack, or the finger of a glove turned inside out

* Cornil and Babes, *Les Bactéries*, p. 458. 1885.

(Figs. 96 and 97). On a sloping surface of nutrient agar-agar a white moist layer forms very quickly.




Fig. 96.

Fig. 97.

PURE CULTIVATIONS OF THE SPIRILLUM
FINKLER-PRIOR IN NUTRIENT GELATINE.

Fig. 96. In twenty-four hours.

Fig. 97. In thirty-six hours.

On potatoes they grow at the ordinary temperature of the air, producing a brownish layer and corrosion of the surface of the potato. They were discovered in the evacuations of cases of cholera nostras, and were claimed at first to be identical with the comma-bacillus of Koch. By the test of cultivation they are now ascertained to be distinct. They also have been shown to be pathogenic.*

Spirillum sputigenum, Lewis.—

Curved rods very similar to the comma-bacilli of Koch. Many have failed with repeated attempts to cultivate these bacilli, and, therefore, maintain that they are quite distinct biologically from the spirilla

* Finkler and Prior, *Ergänzungshefte zum Centralblatt für Allgemeine Gesundheitspflege*, Erster Band. 1885.

associated with Asiatic cholera. Klein asserts that they can be cultivated in an acid nutrient gelatine, and that they are identical with Koch's comma-bacilli in their mode of growth. This has not



FIG. 98.—*SPIRILLUM SPUTIGENUM*. Occurring with *spirochæta denticola*, leptothrix-filaments, micrococci, and bacteria in a scraping from a carious tooth, $\times 1200$.

been confirmed. They occur with other bacteria in saliva, and in scrapings from carious teeth (Fig. 98).

Spirillum tyrogenum, Deneke.—Curved rods, slightly smaller than Koch's comma-bacilli, with a great tendency to form long spirillar threads



FIG. 99.—*SPIRILLUM TYROGENUM*. From a cultivation in nutrient gelatine, $\times 1200$.

(Fig. 99). The colonies on plate-cultivations are sharply defined, and of a greenish-brown colour. After a time they liquefy the gelatine, but the liquefaction is much more marked than in colonies of Koch's commas of the same age, though not so rapid as in the case of the commas of *cholera nostras*. In test-tubes of nutrient gelatine a turbid

liquefaction occurs along the needle track, and on the surface of nutrient agar-agar a yellowish-white layer develops. Inoculation of potatoes gives no result. Administration of the bacilli by the mouth, in the manner employed for testing the pathogenic effect of Koch's bacilli, produced a fatal result in a few cases; on the other hand, injection into the duodenum failed entirely. The pathogenic properties may be, therefore, considered as not yet established. They were isolated from old cheese.

Spirillum rubrum, Esmarck.—Curved rods, spirilla, and spirochetæ. They are actively motile.




FIG. 100.—**SPIRILLUM PLICATILE** (Marsh-spirochæte). From sewage-contaminated water, $\times 1200$.

The growth on artificial nutrient media is extremely slow. In test-tube cultivations they grow along the track of the needle, forming a filament of a wine-red colour, and without causing liquefaction. The growth on the surface of the gelatine is colourless. In broth-cultivations long spirillar threads are formed. They were isolated from the putrid tissues of a mouse.

Spirillum plicatile, Ehrenberg (*Marsh-spirochæte*).—Thin threads, 2.25μ in breadth, with numerous narrow windings, $110-125 \mu$ long, occurring also in spirular forms. The threads have

primary and secondary windings; the former are in each example of equal size, but the latter are often irregular; their ends are cut off bluntly, and they exhibit rapid movement. They occur abundantly in marsh-water in summer, and can be obtained by allowing algæ to decompose in water (Fig. 100). On cultivation the threads break up into long rods, short rods, and finally cocci. This change is rendered visible by making cover-glass preparations, and staining with aniline dyes.



FIG. 101.—SPIRILLUM UNDULA, $\times 1500$.

The following may be provisionally described as distinct species, though they are probably the spiral phase-forms of protean species.

Spirillum serpens, Müller (*Vibrio serpens*).—Threads $11-28 \mu$ long, $.8-1.1 \mu$ thick, with three or four windings. They are actively motile, often united into chains, or forming swarms, and are abundant in stagnant liquids.

Spirillum tenue.—Very thin threads, with at least $1\frac{1}{2}$, usually 2—5 spirals. Height of a single screw is 2—3 μ , and the length of spiral, therefore, 4—15 μ . They are very swiftly motile, and often occur in felted dense swarms in vegetable infusions.

Spirillum undula.—Threads $1.1-1.4 \mu$ thick, 9—12 μ long; spirals 4.5 μ high; each thread has $1\frac{1}{2}-3$ spirals. They are actively motile, and possess at each end a flagellum. They occur in various infusions (Fig. 101).

Spirillum volutans, Ehrenberg. — Threads $1.5-2 \mu$ thick, 25—30 μ long; tapering towards their extremities, which are rounded off. They possess dark granular contents. Each thread has $2\frac{1}{2}-3\frac{1}{2}$ windings or spirals, whose height is 9—13 μ . They have a flagellum at each end, and are sometimes motile, sometimes not. They are found in various infusions and water of marshes.

Spirillum Rosenbergii.—Threads with $1-1\frac{1}{2}$ windings; $4-12$ μ long; $1.5-2.6$ μ thick. They are colourless, but the contents include strongly refractive sulphur granules. Also spirals $6-7.5$ μ in height, which are actively motile, are found in brackish water.

Spirillum attenuatum, Warming.—Threads much attenuated at the ends, which consist usually of three spirals. The middle spiral is about 11 μ high, and 6 μ in diameter; and the end ones 10 μ high, and 2 μ in diameter. They are found in brackish water.

Spirillum leucomelaneum, Koch.—A rare form observed in water covering rotting algæ. Dark and glass-like spaces alternate in the spirillum, resulting from a regular arrangement of the dark granular contents.

Genus III.—*Leuconostoc*.

SPECIES.

UNASSOCIATED WITH DISEASE:—

Leuconostoc mesenteroides . . . Zymogenic saprophyte.

Leuconostoc mesenteroides, Cienkowski (*Gomme de sucrerie*, *Froschlaichpilz*, *Frogspawn fungus*).—Cells singly, in chains, and in zooglœa, surrounded by a thick gelatinous envelope (Fig. 102). The life-history has been very thoroughly investigated.* The spores, $1.8-2$ μ in diameter, are of a round or ellipsoidal form, with thick membrane and shining contents. The outer membrane-layer bursts, and a middle lamella oozes out, and forms a thick gelatinous envelope, while the inner layer remains adherent to the plasma. Thus the spore-germination leads to the formation of a

* Cienkowski, *Die Gallertbildungen d. Zuckerrübensaftes*, 1878; and Van Tieghem, "Sur la Gomme de sucrerie," *Ann. Sc. Nat.* 1879.

coccus with a gelatinous envelope. The coccus then elongates into a short rod-form, and the gelatinous envelope becomes ellipsoidal. The rod divides into two cocci, and each of these lengthens into a rod and divides. By repetition of this process




FIG. 102.—*LEUCONOSTOC MESENEROIDES*.

1. Spores. 2. Spores after germination, showing gelatinous envelope.
- 3, 4, 5, 6. Increase by division. 7. Glomerular form of zooglœa.
8. Section of an old mass of zooglœa. 9. Cocci chains with arthrospores [after Tieghem and Cienkowski].

a chain of cocci results, encased in a cylindrical or ellipsoidal envelope. The chains increase in length, become twisted up, and eventually fall apart into pieces of various lengths. In nourishing liquids a great number of little masses are formed, which adhere together, and produce

pseudo-parenchymatous structures. These latter may join together, forming still larger agglomerations. The masses of zooglœa are of almost a cartilaginous consistency, and admit of sections being made with a razor. After a long time the envelope liquefies, and the cocci are set free; the latter introduced into fresh nourishing media develop new colonies. In the chains some of the cocci become enlarged without changing their form. These acquire the properties of spores, and are called *arthrospores* (p. 161).

This micro-organism occurs occasionally in beet-root juice and the molasses of sugar-makers, forming large gelatinous masses resembling frog-spawn. The vegetation is so rapid, that forty-nine hectolitres of molasses, containing 10 per cent. of sugar, were converted within twelve hours into a gelatinous mass; consequently, it is a formidable enemy of the sugar manufacturers.

Genus IV.—Bacillus.

SPECIES.

ASSOCIATED WITH DISEASE :—

	Bacillus lepræ	Pathogenic.
	Bacillus in syphilis	Pathogenic (?) ; possibly only saprophytic.
	Bacillus typhosus	"
	Bacillus malarie	Pathogenic in animals.
	Bacillus of choleraic diarrhoea from meat-poisoning	Pathogenic.
In man	Bacillus pyrogenes foetidus	Saprophytic in man, pathogenic in animals.
	Bacillus in septicæmia in man	Saprophytic.
	Bacillus in gangrenous septicæmia	"
	Bacillus tuberculosis	Pathogenic.
	Bacillus anthracis	"
	Bacillus mallei	"
	Bacillus malignant oedema	"
In animals	Bacillus of septicæmia of mice	"
	Bacillus of ulcerative stomatitis in the calf	"
	Bacillus in swine-typhoid	"
	Bacillus of swine-erysipelas	"
	Bacillus in tetanus	"
	Bacillus alvei	"

UNASSOCIATED WITH DISEASE :—

	Bacillus pyocyaneus	} Chromogenic saprophytes.
	Bacillus ianthinus	
	Bacillus cyanogenus	} Zymogenic saprophytes.
	Bacillus acidi lactici	
	Bacillus Fitzianus	} Simple saprophytes.
	Bacillus subtilis	
	Bacillus ulna	
	Bacillus tumescens	
	Bacillus megaterium	
	Bacillus figurans	
	Bacillus mycoides	
	Bacillus tremulus	
	Bacillus of jequirity	
	Bacillus caucasicus	
	Bacillus dysodes	
	Bacillus Hansenii	
	Bacillus erythrosporus	
	Bacillus septicus	
	Bacillus saprogenes	
	Bacillus foetidus	
	Bacillus putrificus coli	
	Bacillus coprogenes foetidus	
	Bacillus aerophilus	
	Bacillus mesentericus fuscus	
	Bacillus mesentericus vulgatus	
	Bacillus phosphorescens	

Bacillus lepræ, Hansen—Fine slender rods, 4—6 μ long, and less than 1 μ wide, occasionally

pointed at both ends, some clearly motile, and others not. In tissue sections they have a beaded appearance (Fig. 103). Spore-formation has been described. They have been cultivated artificially on blood serum and glycerine-agar-agar. Inoculation experiments on monkeys and other animals have failed to produce the disease; though in cats and rabbits there have been indications of success.* The bacilli occur in enormous numbers in tubercular leprosy in the nodules of the skin (Plate XX.,



FIG. 130.—LEPROSY BACILLI FROM A SECTION OF SKIN, $\times 1200$.

Figs. 1 and 2), and of the mucous membrane of the mouth, palate, larynx, etc.† They occur also in the liver, spleen, testicles, lymphatic glands, and kidneys (Plate III., Fig. 2); and in the interstitial tissue of the nerves in anæsthetic leprosy. They probably spread by the lymphatics, and are not found in the blood. In their behaviour to staining reactions they are similar to the bacillus of tubercle, except that they stain much more readily.

METHODS OF STAINING THE BACILLUS OF LEPROSY.

Cover-glass preparations may be made in the ordinary way, or by a special method, which consists in clamping a

* Damsch, *Virchow's Archiv*, Bd. 92, Heft 1.

† Thin, *Med. Chir. Trans. Lond.*, 1883; *Brit. Med. Journal*, No. 129, 1884, and Steven, *Brit. Med. Journal*, No. 1281, 1885.



Fig 1.




Fig 2.

nodule with a pile-clamp, until a state of anæmia of the tissue is produced. On pricking with a needle or sharp knife a drop of clear fluid exudes, from which cover-glass preparations may be made.* Cover-glass preparations and sections may be stained by Ehrlich's method (p. 78), or with Neelsen's solution and decolorising with dilute sulphuric acid, or sections by the following process:—

Method of Babes.†—Preparations are stained in a solution of rosaniline hydrochlorate in aniline-water. Decolorise in 33 per cent. hydrochloric acid, and after-stain with methylene-blue.

Bacillus in syphilis, Lustgarten.‡—Rods resembling the bacilli of leprosy and tuberculosis, 3—4 μ long, $\cdot 8 \mu$ thick. Two or more colourless, ovoid points in the course of the rod are visible with a high power; it is thought that they are possibly spores. The bacilli are always found in the interior of nucleated cells which are more than double the size of leucocytes. They have been observed in the discharge of the primary lesion, and in hereditary affections of tertiary gummata. Some observers state that an identical bacillus is found in normal secretions,§ and others || have described a bacillus associated with specific lesions, which is stated to differ from the above in its behaviour towards staining reagents.

METHOD OF STAINING THE BACILLUS OF SYPHILIS.

Method of Lustgarten:—

Sections are placed for from twelve to twenty-four hours in the following solution, at the ordinary temperature of

* Manson, *Lancet*. 1884.

† Babes, *Compt. Rend. de l'Acad. d. Sc.* 1883.

‡ Lustgarten, *Die Syphilisbacillen*. Mit 4 Tafeln. 1885.

§ Alvarez et Tavel, "Recherches sur le Bacille de Lustgarten," *Archiv de Phys. Norm. et Path.*, 17; Klemperer, "Ueber Syphilis und Smegma Bacillen," *Deutsche Med. Woch.* 1885.

|| Eves and Lingard, *Lancet*, April 10th, 1886.

the room, and finally the solution is warmed for two hours at 60° C. :—

Concentrated alcoholic solution of gentian-violet	11
Aniline water	100

The sections are then placed for a few minutes in absolute alcohol, and from this transferred to 1·5 per cent. solution of permanganate of potash. After ten minutes they are immersed for a moment in a pure concentrated solution of sulphurous acid. If the section is not completely decolorised, immersion in the alcohol and in the acid bath must be repeated three or four times. The sections are finally dehydrated with absolute alcohol, cleared with clove-oil, and mounted in Canada balsam.

Method of De Giacomi :—

Cover-glass preparations are stained with hot solution of fuchsine containing a few drops of perchloride of iron. They are then decolorised in strong perchloride of iron and after-stained with vesuvin or bismarck-brown.

Method of Doutrelepont and Schütz :—

Sections are stained in a weak aqueous solution of gentian-violet and after-stained with safranin.

Bacillus typhosus, Eberth (*Bacillus in typhoid fever*).—Rods, 2 μ broad, and forming filaments up to 50 μ long; * or † rods, short, rounded at their ends, and occasionally constricted in the middle; some exhibiting spore-formation. These bacilli have been observed in inflamed Peyer's glands, in the spleen, mesenteric glands, and the lungs in fatal cases of typhoid fever. More recently ‡ a bacillus has been cultivated on several plates of

* Kleb's *Arch. f. Experimental Pathol.* 1880.

† Eberth, *Virchow's Archiv*, Bd. 83.

‡ Gaffky, *Mittheil. a. d. K. Gesundheitsamte.* 1884.

gelatine which were inoculated from different spleens. After twenty-four hours the course of the inoculation streak became visible, and in forty-eight hours a distinct whitish growth had developed. With a low power this was found to consist of numerous colonies of a yellow-brownish colour. The gelatine was not liquefied. The rods varied in length (Fig. 104), were capable of development into threads,



FIG. 104.—BACILLUS TYPHOSUS FROM A POTATO-CULTIVATION, $\times 1500$.

and were motile. They can be cultivated on potatoes at 37° C. They grow well also on blood serum, forming a whitish-grey, somewhat transparent layer. Spore-formation occurs at the ends of the rods. It is stated that inoculation experiments have been made in some cases with success.*

METHODS OF STAINING THE BACILLUS OF TYPHOID FEVER.

To stain the bacilli in the tissues the method of Gram can be employed, or the sections may be left for twenty-four hours in methylene-blue. Koch recommends bis-marck-brown. To colour the spores cover-glass preparations and sections must be left for several days in the fuchsine solution employed in the method of Ehrlich (p. 78); or the solution may be warmed, and in the case of cover-glasses, even raised to boiling-point. They

* Fränkel and Simmonds, *Die Ätiolog. bedeutung des Typhus-bacillus*. 1886.

are then decolorised with nitric acid, and after-stained with methylene-blue.

Bacillus malarix, Klebs (*Bacillus in intermittent fever*).—Rods, 2—7 μ long, which grow into twisted threads. Spore-formation takes place in the centre, or at either end (Plate I., Fig. 14).

Inoculated in rabbits they were stated to produce a febrile disorder analogous to malarial fever,* and in the spleen and marrow the threads and spores of the bacilli were found in abundance. Bacilli with end-spores have been discovered also in the blood of patients suffering from malaria.†

The bacilli were first described as present in the soil of the Roman Campagna.

According to more recent observations, more importance is to be attached to peculiar amœboid bodies and motile filaments present in the blood of cases of malaria (see Appendix D).

Bacillus of choleraic diarrhœa from meat-poisoning, Klein.‡—Rods from 3—9 μ in length, 1.3 μ wide, rounded at their extremities, singly or in chains of two. Spore-formation occurs, the spores being 1 μ thick, oval, and situated in the centre or at the end of the rod.

Feeding with the bacilli and inoculation produced positive results. At the autopsy, pneumonia, peritonitis, pleuritis, enlargement of the liver and spleen, and hæmorrhage were observed, and bacilli were present in the blood and exudations of the animal. They occurred in the blood and juices, and especially in the glomeruli of the kidneys, of several fatal cases of choleraic diarrhœa.

* Klebs and Tommasi Crudeli, *Archiv f. Experimental Pathol.* 1879.

† Marchiafava, *ibid.*

‡ Klein, p. 87.

Bacillus pyogenes fœtidus, Passet.—Small rods, with rounded ends of about 1.45μ in length, and $.58 \mu$ in width; often in twos, or linked together in chains. They are motile, and spore-formation occurs. When cultivated in nutrient gelatine, a greyish, veil-like growth forms on the surface. In plate-cultivations white points appear after twenty-four hours, which develop into greyish spots, and these enlarging coalesce into a layer. In nutrient agar-agar the cultivation resembles the growth on gelatine. On blood serum a moderately thick greyish-white streak develops, and on sterilised potato an abundant, shining, brownish culture. From all these media a putrid odour emanates, but no smell is detected from a cultivation in milk. Inoculated into mice and guinea-pigs abscesses are produced or death from septicæmia results. They were isolated from putrid pus.

Bacillus in septicæmia of man, Klein.*—Rods singly or in chains, $1-2.5 \mu$ long, $.3-.5 \mu$ wide, which were observed in the blood-vessels of the swollen lymphatic glands. They are possibly identical with the following:—

Bacillus in gangrenous septicæmia, Arloing and Chauveau.—Short rods, possessing spores, were observed around wounds in gangrenous septicæmia, and considered to be the cause of the gangrene.

Bacillus tuberculosis, Koch.†—Rods, $2-4 \mu$ and occasionally 8μ long, very thin, and rounded at the ends. They are straight or curved, and frequently beaded (Fig. 105), and occur singly in pairs, or in bundles. They are found in the cells of tubercles, especially in the interior of giant cells. In the latter they are often accompanied

* Klein, *Micro-organisms and Disease*. 1885.

† Koch, *Berl. Klin. Woch.*, No. 15, 1882; and *Mittheil. aus dem Kaiserlich. Gesundheitsamte*, "Ätiologie der Tuberkulose."

with grains which exhibit the same colour reaction (Plate XXI., Fig. 1). They are non-motile. Spore-formation has been described (see p. 164). A good medium for cultivation is solid blood serum of cow or sheep, with or without the addition of gelatine; and the most favourable temperature for their development is 37° — 38° C. The growth takes place very slowly, and only between the temperatures of 30° and 41° C. In about eight




FIG. 105.—*BACILLUS TUBERCULOSIS*, FROM TUBERCULAR SPUTUM, STAINED BY EHRLICH'S METHOD, $\times 2500$. From Photographs.

or ten days the growth appears as little whitish or yellowish scales and grains (Plate XVIII., Fig. 1).

The bacillus can also be cultivated in a glass capsule on blood serum, and the appearances of the growth studied under the microscope. The scales or pellicles are then seen to be made up of colonies of a perfectly characteristic appearance, which may be still further studied by making a cover-glass impression (p. 69, and Plate XVIII., Fig. 4). They are then seen to be composed of bacilli, arranged more or less with their long axis

corresponding with that of the colony itself, and with an appreciable interval between the individual bacilli. The colonies themselves appear as fine curved lines, the smallest being mostly S-shaped. Longer colonies have serpentine twistings and bendings, which often recall the curves of fancy lettering. The ends of the lines run to sharp points, but the middle of the growth is spindle-formed. The youngest colonies are extremely delicate and narrow, but the older colonies increase in size, are thicker across, and, blending with each other, gradually obliterate the characteristic appearances; a lamellated growth results, which increases, and gives the appearance to the naked eye of the scale or pellicle already described. The blood serum is not liquefied unless putrefactive bacteria contaminate the culture. A fresh tube can be inoculated with one of these little scales, and a new generation started. The scales gradually increase in size, and consist entirely of bacilli. In about three to four weeks the cultivation ceases to increase, and it is then necessary to inoculate a fresh tube. Glycerine-agar-agar is even a better medium than blood serum (Fig. 106). A relatively small portion of a cultivation inoculated into the subcutaneous tissue, into the peritoneal or pleural cavity, into the anterior chamber of the eye, or directly into the blood stream, produces after three or more weeks artificial tuberculosis in guinea-pigs and rabbits. Dogs and cats can also be infected by experimental inoculation.

The appearances observed at the autopsy are, swollen lymphatic glands in the neighbourhood of the inoculation, followed by softening and abscess; enlargement of the spleen and liver, with formation of caseous tubercles; tuberculosis of the lungs, bronchial glands, and peritoneum. After inoculation of the eye, grey tubercles appear on the iris, and undergo enlargement and caseation, followed by tuberculosis of the eyeball and organs generally. The bacilli appear to be the direct cause of tuberculosis, and the presence of the bacillus in the sputum of patients is regarded as a distinctive sign of the existence of this disease. The detection of the bacillus has, consequently, become a test which is daily applied by physicians in forming clinical diagnoses.



FIG. 106.

From a photograph of a cultivation of the tubercle-bacillus on glycerine-agar; six months old.

The bacilli are found in all tubercular growths of man, monkeys, cattle (*Perlsucht*), birds, and many other animals, and in cases of artificial tuberculosis, in rabbits, guinea-pigs, cats, etc. (Plate XXI., Fig. 2). In man the bacillus can be detected in the tissues, in the sputum, in the blood, and in the urine.*

* Babes, *Centralbl. f. d. Med. Wissensch.*, 1883, p. 145.




Fig 1.



Fig 2.




Fig 1.



Fig 2.

Tuberculosis may also be produced by inhalation and feeding experiments (p. 137). The channels of infection in man are also most probably the pulmonary or intestinal mucous membranes. The possibility of inoculation of skin wounds is open to doubt. The bacilli or their spores are inhaled from the air, or taken in with food. As a relatively high temperature is required for their growth, they cannot thrive outside the animal body in cold climates. Morphologically identical bacilli have also been observed, but very sparsely, in sections of lupus.

METHODS OF STAINING THE TUBERCLE BACILLUS.

Numerous methods have been recommended for staining the *bacillus tuberculosis*, each of which will be given in detail.

Neelsen's method is preferred by the author for both sections and cover-glass preparations.

Koch's original method.—Cover-glass preparations or sections are laid in Koch's solution (No. 23, c.) for twenty-four hours, or for one hour if the solution is warmed to 40° C. Rinse in water; immerse in a watery solution of vesuvin for two minutes; rinse again in water, and examine; or, after rinsing in water, treat with alcohol, clove-oil, and Canada balsam.

Ehrlich's method.—Cover-glass preparations are allowed to float in a watch-glass, containing a solution of gentian-violet or fuchsine, added to aniline water. A saturated alcoholic solution of the dye is added till precipitation commences (10 ccm. aniline water, and 10—20 drops of the colour solution). The cover-glasses are left in the solution for about half an hour; then washed for a few seconds in strong nitric acid (one part commercial nitric

acid to two of distilled water), and rinsed in distilled water. *After-stain* with vesuvin or methylene-blue, rinse in water, dry and preserve in Canada balsam (Plate III., Fig. 1).

Sections and cover-glass preparations may be stained by this method, as described by Koch.*

Saturated alcoholic solution of methyl-violet or fuchsine	11
Aniline water	100
Absolute alcohol	10

Preparations are left for twelve hours in this solution (colouring of the cover-glass preparations can be expedited by warming the solution).

Treat the preparations with (1—3) solution of nitric acid a few seconds.

Wash in alcohol (60 per cent.) for a few minutes (cover-glass preparations need only be rinsed a few times). After-stain with diluted solution of vesuvin or methylene-blue for a few minutes.

Wash again in 60 per cent. alcohol, dehydrate in absolute alcohol. Clear with cedar-oil, mount in Canada balsam.

Rindfleisch's method.—Prepare a solution composed of

Saturated alcoholic solution of fuchsine	10 drops
Aniline water	2 drams.

Pour it into a watch-glass, and float the cover-glass; warm the watch-glass over a spirit-lamp until steam rises. Remove it from the flame, and set it aside for five minutes. Take out the cover-glass, and transfer it for a few seconds to acidulated alcohol (two drops of nitric acid in a watch-glass full of alcohol). Wash in distilled water, dry, and preserve in balsam.

* *Mittheil. aus dem Gesundheitsamte, Zweiter Band, 1884, p. 10.*

After-stain, if necessary, with bismarck-brown, or methylene-blue.

Weigert-Ehrlich method (*vide* p. 78).

Orth's modification of Ehrlich's method.—Stain by the method of Ehrlich, but decolorise with acidulated alcohol (one of hydrochloric to one hundred parts of 70 per cent. alcohol).

*Gibbes' method.**—Stain cover-glass preparations in magenta solution (No. 22) for 15—20 minutes. Wash in (1—3) solution of nitric acid, until the colour is removed. Rinse in distilled water. After-stain with methylene-blue, methyl-green, iodine-green, or watery solution of chrysoidin, five minutes. Wash in distilled water till no more colour comes away. Transfer to absolute alcohol for five minutes; dry, and preserve in Canada balsam. Leave sections in the stain for half an hour, then treat with nitric acid, and wash with distilled water. Transfer to methylene-blue till deeply stained, wash again in distilled water, and then in spirit. Pass through absolute alcohol and clove-oil, and preserve in Canada balsam.

Gibbes' new method.—Cover-glass preparations are placed in the double staining solution (No. 16), which has been warmed in a test-tube, and, as soon as steam rises, poured into a watch-glass. They are allowed to remain for five minutes, and then are washed in methylated spirit till no more colour comes away, dried in the air or over a spirit-lamp, and mounted in Canada balsam. If the solution is used without warming, the cover-glasses must be left in it for an hour. Sections are treated on the same principles, but must be left in the solution for several hours. The crumpling of the sections by the action of nitric acid is avoided.

Baumgarten's method.—Cover-glass preparations of sputum are made as already described (p. 65), and immersed in a very dilute solution of potash (1—2 drops

* Gibbes, *Practical Pathology*. 1883.

of a 33 per cent. solution of potash in a watch-glass of distilled water). The cover-glass is pressed down on a slide, and examined with a high power. The bacilli can be thus examined in the unstained condition, and to avoid any mistake from confusion with other species, the cover-glass can be removed, dried, passed through the flame, and stained with a drop of an aqueous solution of fuchsine, or gentian-violet. The putrefactive bacteria are stained, but the tubercle bacilli remain absolutely colourless.

Baumgarten's new method.—A solution is prepared as follows:—Drop 4—5 drops of concentrated alcoholic methyl-violet solution into a small watch-glass full of water. (a) Stain the sections in this solution, wash them in water, and decolorise in absolute alcohol (five to ten minutes), or, before treating with alcohol, immerse the sections for five minutes in a half-saturated solution of carbonate of potash. Pass through clove-oil, and mount in a mixture of Canada balsam, free from chloroform, and clove-oil (equal parts). The object of this process is to differentiate the tubercle bacilli from chance bacteria, inasmuch as the tubercle bacilli gradually are decolorised by the clove-oil. (b) Sections stained in the above solution are placed for five minutes in alcohol, and then in a concentrated solution of bismarck-brown in 1 per cent. solution of acetic acid. The after-treatment may be conducted as already described.

Neelsen's method.—Cover-glass preparations may be quickly stained in Neelsen's solution (No. 25) warmed in a watch-glass till steam rises. Sections are left for from five to ten minutes in the solution, and then washed in a watery solution of sulphuric acid (25 per cent.); rinsed in distilled water, and immersed in methylene-blue solution. After two or three minutes they are passed through alcohol and oil of cloves, and mounted in Canada balsam.

Balmer-Früntzel method.—Dissolve two grammes of freshly powdered gentian-violet in 100 grammes of

aniline-water. Immerse the sections for twenty-four hours, and treat as in Ehrlich's method.

Ziehl's method.—Stain with Ehrlich's method, but omit the nitric acid; after-stain with methylene-blue. The latter replaces the stain of all bacteria except the tubercle bacillus.

Lichtheim's method.—Concentrated solution of fuchsine or gentian-violet is diluted with distilled water, and the sections stained for thirty-six hours.

Peters' method.—Sections are stained for half an hour in fresh aniline-gentian-violet solution. Transfer to 20 ccm. of absolute alcohol for eighteen hours, the alcohol being renewed two or three times. Rinse in distilled water for one minute, and immerse for three minutes in a watery solution of aniline-yellow (aniline-yellow .2 dissolved in distilled water 10, filter). Wash in absolute alcohol, clarify with clove-oil, and preserve in Canada balsam.

Fränkel's method.—Sputum preparations are rapidly double-stained by the following method:—Prepare a solution by adding concentrated alcoholic methyl-violet or fuchsine solution, drop by drop till opalescence arises, to 5 ccm. of aniline-water heated to 100° C. Float the prepared cover glasses two minutes in the warmed solution. The process of after-staining and decolorisation is effected by placing the preparation for one to two minutes in one of the following solutions: for fuchsine-stained preparations, a saturated solution of methylene-blue in a mixture of

Alcohol	50
Distilled water	30
Nitric acid	20

which is filtered before use; for preparations stained in methyl-violet, a saturated solution of vesuvin may be used in

Alcohol	70
Nitric acid	30

which must be filtered before use. The sections are washed in water (or weakly acidified 50 per cent. alcohol), dried and mounted in the usual way.

Pfuhl-Petri's method.—The colouring solution consists of 10 ccm. of a saturated alcoholic solution of fuchsine added to 100 ccm. of water. Float the cover-glasses for two minutes in the solution, heated till steam rises. Wash for one minute in glacial acetic acid, rinse in water, and after-stain in an alcoholic or watery solution of malachite-green for a half or one minute. Rinse again in water. Dry, and examine in glycerine, or preserve in Canada balsam.

Senkewitsch's method.—Stain cover-glass preparations in concentrated fuchsine solution. When strongly coloured, wash out the stain for one to two minutes in alcohol, to which one drop of nitric acid has been added for every 10 ccm. Rinse in water, dry, and mount in Canada balsam.

Kaatzer's method.—Place the cover-glass preparations for twenty-four hours in a solution of over-saturated alcoholic gentian-violet, or, if warmed to 80° C., for three minutes. Decolorise in a solution consisting of

Alcohol 90 per cent.	100 ccm.
Water	20 ccm.
Strong hydrochloric acid	20 drops.

Rinse in 90 per cent. alcohol, and after-stain with concentrated watery solution of vesuvin for two minutes; wash again in distilled water, dry, and mount in Canada balsam.

Ehrlich's method and eosin.—The author has found that after sections have been stained with methyl-violet and bismarck-brown by Ehrlich's method, as described by Koch (p. 310), they may with advantage be immersed in a weak alcoholic solution of eosin, then rinsed in clean absolute alcohol, clarified with clove-oil, and mounted in Canada balsam. The giant cells are then stained pink, while

their nuclei are brown and the bacilli blue (Plate XXI., Fig. 1).

Bacillus anthracis (*Bactéridie du charbon*, *Bacillus of splenic fever*, *wool-sorter's disease*, or *malignant pustule*).—Rods, 5—20 μ long and 1—1.25 μ broad, and threads; spore-formation present. As a thorough knowledge of the life-history of this bacillus is of the greatest importance, inasmuch as it is without any doubt the actual cause of widespread disease, the various steps to be followed in a practical study of it will be successively treated in detail. Its morphological and biological charac-



FIG. 107.—BACILLUS ANTHRACIS, $\times 1200$. From a preparation of blood from the spleen of an inoculated mouse.

teristics have been very completely worked out, and it serves as an excellent subject for gaining an acquaintance with the various methods that should be employed in studying micro-organisms. It is found that a mouse inoculated with the bacillus or its spores will die in from twenty-four to forty-eight hours, or more rarely in from forty-eight to about sixty hours.

Examination after death.—The details to be observed in the autopsy have already been described (p. 141). The spleen is found to be considerably enlarged, and may be removed (p. 142), and

examined by making cover-glass preparations, inoculations, and subsequently sections.

Cover-glass preparations.—In cover-glass preparations of the blood of the spleen the bacilli are found in enormous numbers. Preparations should be made similarly with blood from the heart and exudations from the lungs, etc. In the last-mentioned the bacilli are present in very small numbers, or altogether absent. They should be examined both unstained and stained (p. 65). The rods are straight, or sometimes curved, rigid, and motionless, and vary in size in different animals. They stain intensely with aniline dyes, and are then seen to be composed of segments with their extremities truncated at right angles; between the segments a clear linear space exists, which gives them a characteristic appearance (Fig. 107). By double staining (p. 67), the rods are seen to consist of a membrane or hyaline sheath with protoplasmic contents (Fig. 46).

Drop-cultures—A little of the blood from the spleen or heart is employed to inoculate the liquid medium, bouillon or blood serum. Several of these cultures should be prepared, and some of them placed in the incubator. Examined from time to time it will then be observed that the rods grow into long homogeneous filaments, which are twisted up in strands, and then untwisted in long and graceful curves. In a few hours they begin to swell, become faintly granular, and finally, bright, oval spores develop (Plate I., Fig. 28). The



Fig 1.



Fig 2.




Fig 1.




Fig 2.

cultures in the incubator develop rapidly, a temperature of 25°—40° C. being most favourable for the growth of the bacillus. The spores are eventually set free, and by making a fresh cultivation, or by injecting them into a mouse or guinea-pig, they germinate again into the characteristic bacilli, which in their turn grow into filaments and spores. When the spore germinates it swells, the outer layer becomes jelly-like, and giving way at one or other pole, the contents escape and grow into a rod. With the precautions previously described (p. 141) cultivations should be established in nutrient gelatine, nutrient agar-agar, and on sterilised potatoes.

Test-tube cultivations in nutrient gelatine.—Typically characteristic appearances are obtained by inoculating a 5 to 8 per cent. nutrient gelatine. A whitish line develops in the track of the inoculating needle, and from it fine filaments spread out in the gelatine* (Fig. 108). Occasionally a little isolated spot develops, from which rays extend in all directions, like the silky filaments of thistle-down. The filaments are more easily observed with a magnifying glass. In a more solid nutrient gelatine the growth appears only as a thick white thread. As liquefaction of the gelatine progresses, these appearances rapidly disappear, and the growth subsides as a white flocculent mass (Plate VI., Fig. 3). In exhausted culture-media,

* The Author, *Lancet*. 1885.

and sometimes in the blood, filaments are seen in a state of degeneration. This has also been observed in sections of the kidney, etc., of a rabbit inoculated with the anthrax bacillus, which had died of septicæmia the following morning.




FIG. 108.

PURE CULTIVATION OF THE BACILLUS ANTHRACIS IN NUTRIENT GELATINE.

Test-tube cultivations in nutrient agar-agar.—Cultivated upon a sloping surface of nutrient agar-agar a viscous snow-white layer is developed (Plate X., Fig. 1). Without access of air no cultivation can be obtained, the bacilli being ærobic. This can be demonstrated by embedding a piece of lung or spleen pulp containing bacilli in nutrient agar-agar (p. 167). No growth of the bacilli takes place.

Potato-cultivations.—A very characteristic growth results from the inoculation of sterilised potatoes. The damp chamber containing the potatoes is placed in the incubator, and in about thirty-six to forty-eight hours a creamy, very faintly yellowish layer forms over the inoculated surface, with usually a peculiar translucent edge (Plate XI., Fig. 1). On removing the cover of the damp chamber a strong, penetrating odour of sour milk is encountered.

Plate-cultivations.—From the spleen or blood of the heart, cultivations must be established in

nutrient gelatine on plates. The colonies develop in about two days, according to the temperature of the room. They appear to the naked eye as little white spots or specks, which, on examination with a low power of the microscope and small diaphragm, exhibit two distinct forms. One form, on careful focussing, has the appearance of a little compact ball of twisted thread; in the other, liquefaction of the gelatine has commenced, and the thread



FIG. 109.—COLONIES IN A PLATE-CULTIVATION, $\times 70$.

bundles are spreading out like locks or plaits of hair in the neighbouring gelatine. These appearances are perfectly characteristic (Fig. 109).

Cover-glass impressions.—The plate-cultivations should be also examined as soon as the colonies appear, by making cover-glass impressions (p. 69), and staining them with aniline dyes. The filaments, examined with a high power, will then be seen to consist of a number of rods or segments (Plate I., Fig. 30). On the other hand, filaments from a tube-cultivation in a solid medium will be found

to be composed, not only of rods, but here and there of torula-like involution-forms (Plate I., Fig. 30). In a cover-glass impression from a potato-culture (Plate I., Fig. 29) the individual segments have a great tendency to be isolated one from the other, and there is copious spore-formation.

Preservation of spores.—Spores may be preserved simply by allowing anthracic blood to dry and sealing it in a tube. The spores from a potato-




FIG. 110.—COVER-GLASS IMPRESSION-PREPARATION, $\times 70$.

cultivation are treated as follows:—The inoculated surface containing the creamy cultivation is sliced off in a thin layer, and is mashed up with distilled water in a glass capsule. Sterilised silk-thread is cut up into lengths of about a quarter of an inch, and allowed to soak in the paste for some hours, under a bell-glass. The threads are then picked out with a pair of forceps, and laid upon a sterilised glass plate, covered with a bell-glass, and allowed to dry. From the plate, when perfectly dry, they

are transferred to a small test-tube, which can be plugged with cotton-wool, or sealed in the Bunsen burner.

Examination in the tissues.—The organs must be hardened in absolute alcohol, cut, and stained (pp. 75-78). The method of Gram is the most instructive, and eosin a very satisfactory contrast stain. The capillaries all over the body, lungs, liver, kidney, spleen, skin, mucous membrane, etc., will be found to contain bacilli. In some cases the bacilli are so numerous (*e.g.*, in the capillaries of the kidney, Plate XXIII., Fig. 2), that examination with a low power gives the appearance of an injected specimen.

Inoculation of animals.—A thread containing spores, a drop of blood from an infected animal, or a minute portion of a cultivation, introduced under the skin of a mouse or guinea-pig, causes its death, as a rule, in from twenty-four to forty-eight hours. Sheep fed upon potatoes which have been the medium for cultivating the bacillus, die in a few days. Goats, hedgehogs, sparrows, cows, horses, and swine are all susceptible. Rats are infected with difficulty. Dogs, cats, white rats, and Algerian sheep have an immunity from the disease. Frogs and fish have been rendered susceptible by raising the temperature of the water in which they lived.

Dissemination of the disease and mode of infection.—It has been stated that when carcasses of animals which have died of anthrax are buried under the soil, the

development of the bacilli into spores can take place. The spores were supposed to be taken up by earth worms, carried to the surface, and deposited in their castings; animals then grazing or sojourning on the soil are thus liable to be infected.* This has not been borne out by experiment.† Bacilli, however, occur in large numbers in the blood and discharges from the nose and mouth of the moribund animals, and in the urine and fæces. They find a nourishing soil in decaying vegetable and animal matter, and having free access of oxygen form copious spores, so that the grass is extensively contaminated.

In warm and marshy districts the spore-formation is still more active, and the spores may be carried by floods over adjacent meadows. As to the mode of infection, the animals may be directly infected through buccal wounds caused by siliceous grasses, or by wounds of insects; the intestinal and pulmonary mucous membranes are also regarded as pathways of infection. In animals the disease is known as "splenic fever."

In man the mode of infection is by inhalation of spores, and ingress by the pulmonary or intestinal mucous membrane, or by direct inoculation of a wound or abrasion. The spores are derived from the wool or hides of animals which have died of anthrax, and the resulting disease is known as "wool-sorter's disease," and "pustula maligna."

* Pasteur, *Bulletin de l'Académie de Médecine*. 1880.

† Koch, *Mittheil. a. d. Gesundheitsamte*. 1881.

Bacilli are found in the serum of the pustule, and in sputum, urine, fæces, and sweat; and if the disease prove fatal, in the capillaries throughout the body.

Attenuation of the virus.—By cultivating the bacillus in neutralised bouillon at 42°—43° C. for about twenty days, the infecting power is weakened, and animals inoculated with it (*premier vaccin*) are protected against the disease.* To obtain a still more perfect immunity, they are inoculated a second time with material (*deuxième vaccin*) which has been less weakened. The animals are then protected against the most virulent anthrax, but only for a time. From such a culture, however, new cultures of virulent bacilli can be started, and a culture that is “vaccin” for sheep kills a guinea-pig, and then yields bacilli that are fatal to sheep.† Exposure to a temperature of 55° C., or treatment with .5 to 1 per cent. carbolic acid, deprives the bacilli of their virulence. The virulence of the bacillus is also altered by passing the bacillus through different species of animals. The bacillus of sheep or cattle is fatal when re-inoculated into sheep or cattle; but if inoculated in mice, the bacilli then obtained lose their virulence for sheep or cattle; only a transitory illness results, and the animals are protected for a time against virulent anthrax.‡ The possibility of mitigating the virus

* Pasteur, *Compt. Rend.*, 1861, and *Revue Scientifique*, 1883.

† Klein, *Micro-organisms and Disease*. 1885.

‡ Klein, *Reports of the Medical Officer of the Local Government Board*. 1882.

depends upon the species of animals ; rodents cannot be rendered immune by any known "vaccin."

METHODS OF STAINING THE BACILLUS ANTHRACIS.

Cover-glass preparations of blood, etc., can be stained with a watery solution of any of the aniline dyes. They may be rapidly stained with a drop of fuchsine or gentian-violet (p. 65), but more satisfactorily by floating the cover-glasses for twenty-four hours. The preparations may be dried and mounted in Canada balsam, but the typical appearances are best observed in freshly stained specimens examined in water.




FIG. 111.—SPORES OF BACILLUS ANTHRACIS UNSTAINED, $\times 1500$.

The sheath and protoplasmic contents can be differentiated by staining with eosin after the method of Gram.

The spores (Fig. 111) are not stained by the ordinary methods. The cover-glass preparations must be raised to a high temperature in the incubator, or treated with sulphuric acid (p. 68), or passed about twelve times through the flame of the Bunsen burner, or floated on hot solution of the dye.

To double-stain spore-bearing bacilli.—Float the cover-glasses for about twenty minutes on hot alcoholic solution of fuchsine. Decolorise in weak hydrochloric acid, and after-stain with methylene-blue (Fig. 113).

Tissue sections are best stained by the method of Gram, and *after-stained* with eosin, picrocarminate of ammonia, or picro-lithium-carmin.

A more rapid double stain is obtained by immersing the sections in a watery solution of gentian-violet, rinsing in alcohol, and then staining by the method of Orth (p. 78).




FIG. 112.—SPORES OF *BACILLUS ANTHRACIS*, $\times 1200$; stained with gentian-violet, after passing the cover-glass twelve times through the flame.

Weigert's method.—Place the sections from two to five minutes in a 1 per cent. watery solution of gentian-violet. Wash in alcohol, rinse in water, and transfer to picrocarmine solution (Weigert) for from half an hour to an



FIG. 113.—FROM A DOUBLE-STAINED PREPARATION OF *BACILLUS ANTHRACIS*, $\times 1200$.

hour. Treat with alcohol till the colour is almost washed out, and finally clear in oil of cloves and mount in Canada balsam.

Bacillus mallei (*Bacillus of glanders*).—Rods about the size of tubercle bacilli (Fig. 114). When cultivated on solid sterile blood serum at 38° C., the growth appears in the form of minute, transparent drops consisting entirely of the characteristic bacilli.

On sterilised potatoes they form, in a week to ten days at 37° C., a brown gelatinous layer. Pure cultivations after several generations produce the following results when inoculated into horses, rabbits, guinea-pigs, and field-mice. A spreading ulcer with indurated base appears at the site of inoculation while smaller ulcers break out in its vicinity. The lymphatics become swollen, and general infection follows in the form of nodules in the internal organs,



FIG. 114.—*BACILLUS MALLEI*, $\times 1200$; from a section of a glanders' nodule.

and nodules and ulcers on the nasal septum. In guinea-pigs a characteristic tumour of the testis, or ovary and vulva, frequently results, and should be prepared for microscopical sections. The bacilli are found in the nodules of the nasal mucous membrane, the lung, spleen, liver, and other organs in horses and sheep affected with glanders.

METHODS OF STAINING THE BACILLUS OF GLANDERS.

The bacilli of glanders are extremely difficult to demonstrate. The most satisfactory results are obtained as follows:—

Method of Schütz.—The sections are placed for twenty-four hours in a mixture of

Potash solution (1 in 10,000);	} Equal parts.
Concentrated alcoholic methylene-blue solution;	

Wash the sections in a watch-glass with water acidulated with four drops of acetic acid. Transfer for five minutes to 50 per cent. alcohol, fifteen minutes to absolute alcohol, clarify in clove oil, and mount in Canada balsam.

Bacillus œdematis maligni, Koch (*Pasteur's Septicæmia*).—Rods from 3—3.5 μ long and 1—1.1 μ wide; they mostly lie in pairs, and then appear to be double this length. The rods are rounded at their ends, and form threads which are sometimes straight, but more commonly curved. In stained preparations they have a somewhat granular appearance. The bacilli are distinguished from anthrax bacilli by their being somewhat thinner, by their rounded ends, and by their being motile. Anthrax bacilli also never appear as threads in fresh blood, and are differently distributed throughout the body. They are anærobic, and can be cultivated on blood serum and on neutral solution of Liebig's meat extract in an atmosphere of carbonic acid. By embedding material containing bacilli in nutrient agar-agar and nutrient gelatine, characteristic cultivations are obtained. The following process may be adopted to obtain a pure cultivation.* A mouse inoculated subcutaneously with dust, as a rule, dies in one to two days. It is then pinned out, back uppermost, on a slab of wood (p. 141), and the hair singed with a Paquelin's cautery from one hind leg up to the neck, across

* Hesse, *Deutsch. Med. Woch.*, No. 14. 1885.

the latter, and down again to the opposite hind leg. Following the cauterised line, the skin is cut through with sterilised scissors, and the flap turned back and pinned out of the way. With curved scissors little pieces of the subcutaneous œdematous tissue, in the neighbourhood of the inoculated spot, are cut out, and sunk with a platinum needle in a 1 per cent. nutrient agar-agar, or 5 per cent. nutrient gelatine. Fragments of tissue may also be embedded by the method already described (p. 167).

The inoculated tubes are placed in the incubator. In a few hours a whitish turbidity spreads out from the piece of tissue, and upwards in the needle track. Examined microscopically, the turbidity is found to be due solely to the development of bacilli of œdema. The surface exposed to the air exhibits no trace of the bacilli.

To investigate the tubes microscopically, a sterilised glass tube with a capillary end may be used, with its neck plugged with sterilised cotton-wool, and provided at the mouth with a suction ball. The capillary end is thrust into the cultivation, and a small fragment removed by aspiration. In the course of the first day the bacilli spread throughout a great part of the agar-agar in such a way that a more or less equally diffused cloudiness of the medium ensues, with subsequent appearance of strongly marked clouds or lines of turbidity. At the same time gas-bubbles develop along the needle

track, and a collection of liquid takes place, while spore-formation also commences. The following day these appearances are more marked, the opacity is more pronounced, the development of gas increases, and the liquid contains more spore-forming bacilli and numerous free spores.

The nutrient-gelatine cultures during the first day show no macroscopic change, but after a few days the piece of tissue is surrounded with a white halo. This gradually spreads in all directions, and is apparently beset with hairs. The gelatine liquefies, and the fragment of tissue, degenerated bacilli, and spores, sink to the bottom. The cultivation is also very characteristic in $\frac{1}{2}$ per cent. nutrient agar-agar. If placed in the incubator, in a few hours a cloudiness forms around the piece of embedded tissue, which is caused by bacilli gradually spreading in all directions in the nutrient medium. Mice inoculated from these cultivations die more quickly than from the original infection from dust. On potatoes they are cultivated by introducing a piece of liver or other tissue containing the bacilli, into the interior of a sterilised potato (p. 143), incubated at 38° C. The bacillus is not deprived of its virulence by cultivation. The spores of the œdemabacilli appear to be very widely distributed. They are found in the upper cultivated layer of the soil, in hay dust, in decomposing liquids, and especially in the bodies of suffocated animals, which are left to decompose at a high temperature. From any of

these sources animals can be successfully inoculated. If a guinea-pig, for example, be subcutaneously inoculated with earth, putrid fluid, or hay dust, death frequently occurs in from twenty-four to forty-eight hours. At the autopsy the most characteristic symptom is a widespread subcutaneous œdema, which originates from the point of inoculation, accompanied with air-bubbles, and contains a clear reddish liquid full of motile and non-motile bacilli. The internal organs are little changed, the spleen is enlarged and of a dark colour, and the lungs are hyperæmic, and have hæmorrhagic spots. Examined immediately after death, few or no bacilli are detected in the blood of the heart, but in that of the spleen, liver, lungs, and other organs, in the peritoneal exudation, and in and upon the serous coating of abdominal organs they are present in large numbers. If, on the other hand, the animal is not examined until some time after death, then the bacilli are found in the blood of the heart, and distributed all over the body.

Bacillus of septicæmia of mice, Koch.—Extremely minute bacilli, $\cdot 8$ — 1μ long, and $\cdot 1$ — $\cdot 2 \mu$ broad, often in pairs, seldom in chains of four. On cultivation they do not appear to make threads, but the bacilli lie together in masses. Spores have been observed. The bacilli are probably non-motile. They are most commonly in the interior of white blood corpuscles. In these they increase, and in many cases a white cell is only represented

by a mass of bacilli. The bacilli, or rather their spores, occur in putrid liquids. If a number of mice be inoculated with a minimum quantity of putrid fluid, about a third of them die of septicæmia. They rapidly sicken, their eyes inflame, their eyelids stick together, they become soporific, and die in from about forty to sixty hours. At the autopsy one finds slight œdema at the seat of inoculation, and enlargement of the spleen; the bacilli are found both free and lodged in the white corpuscles, in the œdematous tissue, and in the blood capillaries. A minimal quantity of this blood produces the disease if inoculated in house-mice or sparrows. Field-mice have an immunity. Rabbits




Fig. 115.

Fig. 116.

PURE CULTIVATIONS OF THE BACILLUS OF SEPTICÆMIA OF MICE IN NUTRIENT GELATINE.

Fig. 115. In two days.

Fig. 116. In five days.

and guinea-pigs inoculated in the ear suffer from only a local erythema, which disappears after five or six days, and renders them for a time immune. Rabbits inoculated in the cornea suffer from an intense

inflammation of the eyes. The bacilli are easily cultivated outside the body on a mixture of aqueous humour and gelatine, and especially on nutrient gelatine rendered slightly alkaline with sodium phosphate. They grow also very well on the ordinary nutrient gelatine, forming in plate-cultivations scarcely perceptible cloud-like specks, and in a test-tube of nutrient gelatine they form a delicately clouded cultivation along the needle track (Figs. 115 and 116). A small quantity of pure cultivation carried through many generations reproduces the disease when inoculated into mice. The organs should be hardened in absolute alcohol, and sections stained preferably by the method of Gram (Plate XXV., Figs. 1 and 2).

Bacillus of ulcerative stomatitis in the calf, Lingard and Batt.—Rods $4\ \mu$ — $8\ \mu$, or more in length; $1\ \mu$ in width. Spores are frequently present. Injected into the rabbit or mouse they produce a fatal result. They were observed in ulcerations on the tongue and mucous membrane of the mouth of calves.

METHOD OF STAINING THE BACILLI OF ULCERATIVE STOMATITIS.

Sections through the ulcerations of the calf's tongue, or of the inoculated tissue of the rabbit, were stained by immersion in a mixture of magenta and methylene-blue. They were then washed in spirit, cleared in olive-oil, and mounted in Canada balsam.




Fig 1.



Fig 2.

Bacillus swine-typhoid (*Bacillus in swine-plague or swine-fever. Bacillus in pneumo-enteritis of the pig*, Klein*).—Rods 2—3 μ long, actively motile (Fig. 117); spore-formation described (Fig. 119). They can be cultivated in broth and hydrocele-fluid, and carried on through successive generations. A drop of any of these cultivations produces the disease in pigs, mice, and rabbits (Fig. 118); the animals die with a characteristic




FIG. 117.—FROM A PREPARATION OF BRONCHIAL MUCUS OF A PIG
[after Klein].

swelling of the spleen, coagulative necrosis of tracts of the liver tissue, and inflammation of the lungs; it is said that pigs inoculated with artificial cultures are protected against a fatal attack.

The bacillus was observed in the diseased organs of pigs that had died of swine-fever, and of animals that had died from the inoculated disease.

Bacillus of swine-erysipelas (*Bacillus des erysipelas malignum, of Rothlauf, or Rouget du porc*).—Extremely minute bacilli, bearing a close

* Klein, *Report to Med. Offic. Loc. Govt. Board.* 1877—1878.

resemblance to the bacilli of septicæmia of mice.* In test-tubes of nutrient gelatine they develop a cloudy growth in the needle track (Plate XXVI., Fig. 1),



FIG. 118.—BLOOD OF FRESH SPLEEN OF A MOUSE, AFTER INOCULATION WITH SWINE-FEVER [after Klein].

and in plate-cultivations characteristic, thread-like, branching, or star-like colonies are formed (Plate XXVI., Fig. 2). Inoculated into mice and rabbits, a fatal result is produced; but experiments with



FIG. 119.—BACILLI FROM AN ARTIFICIAL CULTURE, WITH SPORES [after Klein].

pigs were unsuccessful. Pigeons were also susceptible, and the bacilli were detected in their blood (Plate XXVI., Fig. 3).

Bacillus in tetanus, Nicolaier and Kitasato.†

* Löffler und Schütz, *Arbeiten aus dem Kaiserlichen Gesundheitsamte*, vol. i. 1885.

† Kitasato. *Zeitschrift für Hygiene*, 1889.




Fig 1.



Fig 2.




Fig 3.

Long slender motile rods. They exhibit terminal (*drumstick*) spore-formation. They were found associated with other bacteria in abscesses resulting from the inoculation of mice and rabbits with garden earth. Inoculation of earth subcutaneously in these animals induces fatal tetanus. They have been isolated from tetanus in man, and pure-cultivations obtained in different media. In grape-sugar-gelatine they produce a characteristic growth composed of excessively fine filaments. The gelatine is slowly liquefied. Mice, guinea pigs, and rabbits, inoculated with a pure-cultivation die of tetanus.

Bacillus alvei, Cheshire and Cheyne.* — Rods varying in size, and forming large oval spores. When cultivated in nutrient gelatine in test-tubes a delicate, ramifying growth appears on the surface, and irregular whitish masses arise along the needle track. Processes shoot out from these masses, and extend through the gelatine for long distances. They are thickened at points in their course, and clubbed at the ends. The gelatine is gradually liquefied, and the bacilli form a loose, white, flocculent deposit at the bottom of the tube. The liquid in the tube becomes yellowish in colour after a time, and gives off an odour of stale but not ammoniacal urine. The colour and odour are distinctive also of the disease attributed to the bacilli. In plate-cultivations the bacilli grow out in series of rods in single file, or in rows of several side by side. The processes which are formed, tend to curve, and at a short distance from the track of

* Cheshire and Cheyne, *Journ. Royal Microscopical Society*, 1885, pp. 582-601.

the needle-streak form a distinct circle, from which another process grows out, and a fresh circle is developed. The gelatine in the vicinity of the bacilli gradually liquefies, and channels are formed in the gelatine in which the bacilli move backwards and forwards. On nutrient agar-agar a whitish layer develops, consisting of bacilli arranged side by side, which in a few days are replaced by rows of spores similarly arranged. On potatoes they form a dryish yellow layer, and in milk a tremulous jelly. A cultivation of the bacillus in milk, sprayed over a honey-comb containing a healthy brood of bee larvæ, produced the disease known as "foul-brood." Adult bees fed on material containing bacilli became affected; inoculation of mice and rabbits with the bacillus gave doubtful results. The bacilli were isolated from the diseased larvæ of bees.

Bacillus pyocyaneus (*Micrococcus pyocyaneus*, Gessard. *Bacterium æruginosum*, Schröter. *Bacillus fluorescens*).—Slender rods sometimes linked in twos or threes, or collected in irregular masses. Spore-formation present. On plate-cultivations white colonies with indistinct contour appear in twenty-four hours, and the whole of the gelatine has a greenish shimmer. In test-tube cultivations the gelatine is liquefied, and coloured green by reflected light, and a deep orange by transmitted light (Plate VI., Fig. 1). On nutrient agar-agar they form a white layer, and colour the medium a pea-green. The pigment formed by the rods is a definite principle—pyocyanin.* It can be extracted with chloroform from pus, and from washings

* Gessard, *De la Pyocyanine et de Son Microbe*. Paris thèse. 1882.

of bandages ; it is soluble in acidulated water, which it colours red. In neutral solution it becomes blue. It crystallises in chloroform in long needles ; and forms sometimes lamellæ and prisms. The rods occur in the pus of those cases in which the wounds and pus-stained bandages exhibit a greenish-blue colour.

Bacillus ianthinus (*Bacterium ianthinum*, Zopf. *Bacillus violaceus*).—Slender rods, about four times their width in length, with rounded ends. They also form threads, and are actively motile. Spore-formation present in the rods. On plate cultivations the colonies occur as circumscribed liquefied areas, in the centre of which is a collection of the coloured growth. In test-tubes a funnel-shaped liquefaction takes place, while a granular-looking violet mass subsides to the bottom. On agar-agar and potatoes a beautiful violet growth rapidly develops. They were observed on pieces of pigs' bladder floating on the surface of water rich in bacteria. They occurred only on the surface of the bladder exposed to the air, and never on the part under water. They occasionally occur in common tap water. The colouring matter is soluble in alcohol.

Bacillus yanogenus, Fuchs. (*Bacterium syncyanum*. *Bacillus of Blue Milk*).—Motile rods, 2·5—3·5 μ in length, and ·4 μ wide (Fig. 120). The rods after division may remain linked together, and form chains. Non-motile rods

enveloped in a gelatinous capsule, and involution-forms, have also been described. Spore-formation present.

Cultivated in a test-tube of nutrient gelatine, the bacilli grow principally upon the free surface, in the form of a white layer. The surface of the




FIG. 120. *BACILLUS CYANOGENUS*, $\times 650$.

A. Active rods. B. Rods in zooglœa. C. Chain of short rods. D. Chain of cocci. E. Cocci stage. F, G. Spore-forming rods. H. Involution-forms [after Neelsen].

gelatine becomes cupped, and a peculiar greenish-brown colour develops in the medium, especially in proximity to the growth.

On a sloping surface of nutrient agar-agar, they grow as a white layer, and colour the upper part of the medium a smoky brown (Plate VIII., Fig. 2).

The bacilli can also be cultivated in milk and on

various other substrata, as potatoes (Plate XVII.), boiled rice, and starch. A pure cultivation in neutralised sterile milk develops a weak alkaline reaction. The colouring matter which is formed, varies with the nourishing medium; for example, in milk a slate-blue coloration is produced, but if the milk has become acid by the growth of the *Bacillus acidi lactici*, then the colour is an intense blue.

The micro-organism occurs occasionally in cow's milk, producing a blue colour. It has been observed, especially in the north of Germany, during the warm months; and where milk is kept in hot rooms, in the winter also. The blueing was originally attributed to a diseased condition of the cows, or to their eating certain meadow plants.

Bacillus acidi lactici.—Long and short rods, 1—2·8 μ long, .3—·4 μ thick, and thread-forms; no cocci; spore-formation.* Cultivated on nutrient gelatine the breadth of the rods is lessened. They grow best between 35 and 42° C., and cease under 10° C. Cultivated at a temperature over 45·5°, they are no longer able to produce acidity. Probably several micro-organisms are able to produce an acid reaction in milk.

They occur with various other bacteria in sour milk, and a pure cultivation, isolated by plate-cultivations, turns sterilised milk sour.

Bacillus Fitzianus, Zopf.—Cocci, short rods, long rods, and threads. This bacillus, cultivated in meat extract and glycerine at 36° C., causes an active fermentation with the production of ethyl alcohol. Spore-formation occurs in the rods. Observed in unboiled hay infusion, accompanying the hay bacillus.

* Hueppe, *Mittheil a.d. Gesundheitsamt*, N. Band.

Bacillus subtilis (*Hay bacillus*).—Cylindrical rods as much as 6 μ in length, and about three times as long as broad. Single forms grow to double their length, and then undergo division. They also form threads which may be composed of long rods, short rods, and cocci. They are motile, and provided with a flagellum at each end. If the nourishing medium is impoverished, the multiplication of the rods by division gradually ceases, and spore-formation commences. The rods become motionless, and a dark spot is visible, either in the middle or towards one end. This gradually develops into a shining spore with a dark contour. The rods swell slightly during this process, their contour becomes undefined, and soon disappears entirely, so that the spores are set free in about twenty-four hours. The spores are 1.2 μ long, and .6 μ broad. They develop into rods in the following way. On one side of the spore a swelling appears, at the summit of which an opening in the spore-membrane results, and the germ escapes. This lengthens into a rod, and remains for a time attached to the empty spore-membrane. These spores are widely distributed, and occur in the air, soil, dust, etc. On the excrement of herbivorous animals the bacilli form a white efflorescence, and on infusion of horse-dung a thick crumpled skin. They flourish equally in liquids and upon damp, solid, nourishing media. On potatoes they grow as a yellowish-white skin; on ordinary nutrient liquids

they develop a thin, and subsequently a thick, dense, crumpled pellicle, with copious spore-formation. They are ærobic; deprivation of oxygen causes the growth of the bacilli to cease, and the rods degenerate. They may be cultivated in various other nourishing media, such as blood-serum, nutrient gelatine, and nutrient agar-agar (Plate XI., Fig. 2).

The simplest way to obtain a culture of the bacillus is to make a decoction of hay. The hay is chopped into small pieces, and boiled with distilled water in a flask for a quarter of an hour; it is then filtered into a beaker, which must be covered with a glass plate, and set aside in a warm place. In two or three days the liquid swarms with the bacilli, the spores of which exist in great numbers in ordinary hay. A more sure method for obtaining a pure cultivation is as follows:—

(a) Add only a small quantity of water to some finely chopped hay, and set aside for four hours at 36° C.

(b) Pour off the extract, and dilute it to the Sp. Gr. 1.004.

(c) Boil gently for one hour in a bulb plugged with cotton wool.

(d) Set aside 500 ccm. of the extract at 36° C.

In about twenty-four hours, as a rule, a pellicle has commenced to develop upon the surface of the liquid. If the reaction is definitely acid, carbonate of soda solution must be added to the decoction.

METHODS OF STAINING HAY BACILLUS.

To demonstrate the flagella of the bacilli, they may be stained with hæmatoxylin solution (Koch).

The linking together of cocci, long rods, and short rods in the threads, is shown by treating with alcoholic solution of fuchsine, or with iodine solution (Zopf).

To stain the spores the cover-glass preparations must be heated to a very high temperature (210° C.), in the hot-air steriliser for half an hour, or they may be exposed for a few seconds to the action of concentrated sulphuric acid (Büchner), or floated for twenty minutes on hot solution of the dye.

Bacillus ulna, Cohn.—Cocci, short rods, long rods, and threads. Diam. of the cocci 1.5 — 2.2 μ . Spore-formation in both short and long rods. No septic odour is produced by this bacillus in a nourishing liquid. Cloudy masses are found on the surface of the liquid, which later form a thick dry pellicle. The latter consists of bundles of threads matted together. The formation of ellipsoidal spores occurs in the usual way; they measure 2.5 — 2.8 μ long, and more than 1 μ wide. The bacillus is found in rotting eggs, and can be cultivated on boiled white of egg. It is closely allied to *Bacillus subtilis*.

Bacillus tumescens, Zopf.*—Cocci, long and short rods. They form a jelly-like disc, $.5$ — 1 cm. in diam., on slices of boiled carrot, with the appearance of a rather tough crumpled skin of a whitish colour. Examination of this pellicle shows that it

* Zopf, *Die Spaltpilze*. 1885.

is formed of rows of rods lying closely together. These rods can be observed to divide into short rods and cocci. Spore-formation occurs in two stages of development, viz., in the cocci and in the short rods. A cultivation is obtained by exposing slices of boiled carrot, slightly moistened, to the air at the temperature of the room.

Bacillus megaterium, De Bary.—Large rods 2.5μ wide, and four to six times as long. They




FIG. 121.—BACILLUS MEGATERIUM.

a. A chain of rods, $\times 250$. The rest $\times 600$.

b. Two active rods.

d to *f.* Successive stages of germination.

h and *l.* Successive stages of germination.

[After De Bary.]

are usually somewhat curved. Transverse division occurs, each segment attaining the length of the original rod. In the fresh state they appear non-articulated, but when treated with a dehydrating agent (tincture of iodine, alcohol), they are seen to be composed of short segments. The rods are

motile, and form irregular chains, of a disjointed appearance. They can be cultivated on nutrient agar-agar and nutrient gelatine. The latter is slowly liquefied, but the appearances are not characteristic.



FIG. 122.
PURE CULTIVATION OF *BACILLUS FIGURANS* ON THE SURFACE OF NUTRIENT AGAR-AGAR.

Spore-formation occurs in the usual way (Fig. 121). It was first observed on boiled cabbage.

***Bacillus figurans* (Crookshank).—**

Rods, with rounded ends, varying in length. Spore-formation present. In plate-cultivations they cause a cloudy growth, spreading from various points; if a cover-glass impression be made, this is found to consist of the regularly-arranged parallel rods. The chains of rods become twisted at intervals into curious convolutions, from which offshoots are continued in various directions. These long shoots or processes are again twisted at intervals into varying shapes and patterns (Plate XXVII., Figs. 1 and 2). Cultivated in nutrient gelatine, the bacilli form on the surface visible windings, from which fine filaments grow down into the gelatine.

They spread out also in almost parallel lines transversely from the needle track. The gelatine is not liquefied. On an oblique surface of nutrient agar-agar the filaments spread downwards into the substance of the jelly, and outwards from the central



Fig 1.



Fig 2.

streak on the surface, forming a feather-like cultivation* (Fig. 122). Cultivated from the air. Identical with *Bacterium Zopfii*, (p. 278).

Bacillus mycoides.—Rods with rounded ends, and varying in length. Spore-formation present. In plate-cultivations the colonies have a characteristic appearance like tufts of wool or masses of mycelium. In test-tube cultivations there is the same mycelial appearance. Nutrient gelatine is liquefied. The bacillus is found in garden earth.

Bacillus tremulus.—Rods shorter and thinner than those of *Bacillus subtilis*. They are provided with a flagellum at both ends, and exhibit characteristic trembling and rotatory movements. Spores thicker than the bacillus, and often placed laterally. They were observed on rotting plant infusions, forming a thick slimy skin.

Bacillus of jequirity, Sattler.—Rods 2—4.5 μ long and .58 μ thick. They can be cultivated on nutrient gelatine and blood serum. Infusion of jequirity containing the bacilli, or an artificial cultivation of the bacilli, inoculated into the conjunctiva of healthy rabbits produces severe ophthalmia. The poisonous principle is, however, believed to be a chemical ferment, *abrin*, and not the bacillus. Boiling, which does not destroy the spores of the bacillus, destroys the ferment, and cultivations started with these spores, though teeming with jequirity bacilli, are quite harmless.† The bacilli occur in infusions of the beans of *Abrus precatorius*, or jequirity.

Bacillus caucasicus, Kern.—Rods forming two spores, one at each end, otherwise similar to *Bacillus subtilis*. They occur in the form of whitish lumps in company

* Described by the author, "Notes from a Bacteriol. Laboratory," *Lancet*. 1885.

† Klein, *Micro-organisms and Disease*. 1885.

with *Saccharomyces mycoderma* in the production of a drink "kephir" from cow's milk. The fermentation is not due to the bacillus.

Bacillus dysodes, Zopf.—Cocci, long and short rods, and spores. They were observed in bread, making it greasy and unfit for food, and generating a penetrating odour resembling a mixture of peppermint and turpentine. A great loss may result to bakers if the fungus is introduced with the yeast.

Bacillus Hansenii, Rasmussen.—Rods 2·8—6 μ long, ·6—·8 μ wide. Cultivated on sterilised potato in four days they form a chrome-yellow layer with an agreeable fruitlike smell. Two or three days later the growth dries, and changes to orange-yellow in colour; later it passes to yellowish or brown, and forms at the same time spores 1·7 μ long, 1·1 μ wide. The colouring matter is insoluble in most reagents.

The bacilli occur on nourishing solutions, malt infusion, broth, wine, which have been kept at 31 to 33° C., as a yellow or whitish skin.

Bacillus erythrosporus, Cohn.—Motile rods and threads; rods exhibiting spore-formation. They grow well in nutrient gelatine, colouring the medium green by transmitted light. They were found to form a pellicle on meat-extract-solutions and on rotting albuminous liquids.

Bacillus septicus, Klein.*—Rods varying in size, non-motile. They form threads or leptothrix filaments, and are rounded at the ends. They are anærobic, and form spores independently of access of air. In a nourishing fluid they are overcome by the presence of *micrococci*, *Bacterium termo* or *Bacillus subtilis*. They occur in the soil, in putrid blood, and many putrid albuminous fluids, and occasionally in the blood-vessels of man and animals after death.

Bacillus saprogenes, Rosenbach. — Three rod-

* Klein, *Micro-organisms and Disease*. 1885.

formed organisms have been described by Rosenbach as intimately associated with putrefactive processes.

No. 1.—Large rods (Fig. 123), which form an irregular sinuous streak with a mucilaginous appearance, when cultivated on nutrient agar-agar. Spore-formation present. They grow also very readily on blood serum, and all cultivations yield the odour of rotting kitchen refuse. They are not pathogenic.

No. 2.—Rods shorter and thinner than No. 1. They develop very rapidly on agar-agar, forming transparent drops, which become grey. They were isolated from a patient suffering from profusely-sweating feet. The cultivations yield a characteristic odour similar to the



FIG. 123.—BACILLUS SAPROGENES, NO. 1. [After Rosenbach.]

last. They are pathogenic in rabbits. They appear to be identical with *Bacillus fœtidus* (*Bacterium fœtidum*, Thin).

No. 3.—See *Bacterium saprogenes*.

Bacillus fœtidus (*Bacterium fœtidum*, Thin).—Cocci, short rods, long rods, and leptothrix. The cocci, 1.25—1.4 in diam., occur singly or in pairs. Spore-formation present in the rods. They were isolated from the exudation in a case of profuse sweating of the feet, and the odour was noticeable in the cultivation (*vide Bacillus saprogenes*).

Bacillus putrificus coli, Bienstock.—Slender, motile rods, 3 μ in length, often less, sometimes forming long threads. Spore-formation present. Cultivations in gelatine are iridescent. They are constantly present in fæces.

Bacillus coprogenes fœtidus, Schottelius.—

Rods with rounded ends, shorter, but about same width as the hay-bacillus. Immotile; spore-formation present. On nutrient gelatine the colonies are yellowish, and do not liquefy the medium. On potatoes they form a pale grey layer. They develop a strong rotting odour. They were isolated from the organs and intestinal contents of pigs reputed to be ill with swine-erysipelas.

Bacillus aerophilus, Liborius.—Slender rods, two-thirds the width of the hay-bacillus, and thread-forms. Spore-formation present. In nutrient gelatine they form dot-like colonies of greenish-yellow colour, which liquefy the gelatine. In test-tubes a somewhat funnel-shaped liquefaction results. On potatoes they develop a yellowish layer. Powerfully ærobic. Found as a contamination.

Bacillus mesentericus fuscus, Flügge.—Small, short, actively-motile bacilli, often linked in twos and fours. Spore-formation present. They form white colonies on plate-cultivation, which later stream out in rays at the periphery, and liquefy the gelatine. In test-tube cultivations a funnel-shaped turbidity is produced, and then a stratum of liquefied gelatine with subsiding flocculi. On potatoes they develop a smooth yellowish layer, which soon becomes folded and wrinkled, forming a delicate veil over the nutrient surface. They occur on unsterilised potatoes.

Bacillus mesentericus vulgatus, Flügge (*Potato bacillus*).—Rods, longer and thicker than the above, and sometimes thread-forms; spore-formation present. The colonies, at first somewhat transparent, have later an opaque centre, and liquefy the gelatine. In test-tubes of nutrient gelatine a funnel-shaped turbidity results, and then an upper-stratum is completely liquefied, while a skin floats on the surface, and flocculent masses subside to the bottom of the liquid layer. They occur on potatoes.

Bacillus phosphorescens, Fischer.—Motile rods. Nutrient gelatine quickly liquefied. Cultivated from seawater (West Indies).

Genus V.—Vibrio.

SPECIES.

UNASSOCIATED WITH DISEASE:—

Vibrio rugula Zymogenic saprophyte.

***Vibrio rugula*, Müller.**—Rods and threads, 6—16 μ long, about $\cdot 5$ — $2\cdot 5$ μ thick. The rods are either simply bowed, or possessed of one shallow




FIG. 124.—*VIBRIO RUGULA*, $\times 1020$. A. Bowed threads. B. Slightly-curved rods. C. Rods swollen preparatory to spore-formation. D. Rods swollen at the spore-forming end. E. Various stages of the developing spores. [After Prazmowski.]

spiral (Fig.). They bear a flagellum at each end. The rods form swarms when causing decomposition, and then, or after, grow out into threads, curved in a screw-like manner. In the next stage of development the rods cease to move, and become swollen with granular contents. One

extremity develops an enlargement, giving the rod the appearance of a pin. The spore formed by the contraction of the plasma in the swollen end finally becomes globular. The vibrios appear in vegetable infusions, causing fermentation of cellulose.

Genus VI.—Clostridium.

ASSOCIATED WITH DISEASE IN ANIMALS :—

Clostridium of symptomatic anthrax. Pathogenic.

UNASSOCIATED WITH DISEASE :—

Clostridium butyricum . . . Zymogenic saprophytes.

Clostridium polymyxa . . . " "

Clostridium of symptomatic anthrax (*Rauschbrand, Charbon symptomatique*.*)—Rods rounded at the ends, mostly with a shining spore at one end. They are especially distinguished from the bacilli of anthrax by being motile. Cultivated on blood-serum, threads develop, consisting of both rods and cocci. From blood-serum they can be cultivated on nutrient gelatine, and vegetable albumen.

Cultivation does not deprive the micro-organism of its virulence, but heating the spores to 85° C. renders them harmless.

Inoculation in the subcutaneous tissue of guinea-pigs, rabbits, calves, and sheep proves fatal. White rats, dogs, and fowls have an immunity. Injection into the veins in small quantity produces a febrile

* Arloing, Cornevin et Thomas, *Bull. de l'Acad. de Méd.* 1881.

disorder, in larger quantities death. Animals in the former case suffer an abortive illness, which protects them against further inoculation. The micro-organism is the cause of a disease in cattle, "*black-leg*," "*quarter-evil*," or "*Rauschbrand*." At the autopsy the micro-organisms are found in the subcutaneous connective tissue, in the lymph glands, kidneys, spleen, and lungs. An irregular tumour is formed in the skin, which develops rapidly, and gives crepitus on palpation. The tumour, which is hæmorrhagic effusion, occurring in the extremities, impedes the animal's movements. The cattle infected die in thirty-six to forty-eight hours.

Clostridium butyricum, Prazmowski (*Bacillus amylobacter*, Van Tieghem; *Bacillus butyricus*. *Bacillus of butyric acid fermentation*).—Rods 3—10 μ long, and under 1 μ wide, often indistinguishable from *Bacillus subtilis*. They grow out into long, apparently unjointed threads. They are mostly actively motile, but also occur in zooglœa. The rods and threads are sometimes slightly bent, like vibrios. They are anærobic. The shorter rods as a rule swell in the middle, becoming ellipsoidal, lemon, or spindle-shaped; the long rods, and sometimes the short ones, swell at one end; in either case ellipsoidal spores are developed (Fig. 125).

If they be cultivated in nutrient gelatine, the medium is liquefied, and a scum formed on the surface. They grow best between 35° and 40° C. The spores are widely distributed in nature, and

grow readily on fleshy roots, old cheese, etc. They convert the lactic acid in milk into butyric acid, and produce the ripening of cheese. They occur




FIG. 125.—*CLOSTRIDIUM BUTYRICUM*.

A. Active stage. (a, b) Bent rods (vibrio-form) and threads. (c) Short rods. (d) Long rods.

B. Spore-formation. C. Spore-germination. [After Prazmowski.]

also in solutions of starch, dextrine, and sugar, and are the active agents in the fermentation of sauerkraut and sour gherkins.

METHOD OF STAINING THE BACILLUS OF BUTYRIC
ACID FERMENTATION.

Treat the bacilli with iodine-solution. At certain stages of the fermentation process the plasma takes a blue or violet-black coloration. The young rods give the former appearance, and the older ones the latter. It is most easily observed when the bacillus is cultivated in a substance containing starch, or, if starch is wanting, in the presence of cellulose, calcium-lactate, or glycerine; in bacilli cultivated in sugar solutions the reaction seldom appears.

Clostridium polymyxa, Prazmowski.—Threads consisting of rods which vary in length; cocci, involution-forms, and spores are also present; cultivated on nourishing solutions they develop a thick skin on the surface. On boiled beet and other roots they form a gelatinous scum, which often consists of crinkled, tough masses, several cm. in diam., somewhat similar to the *Ascococcus Billrothii*. They cause fermentation in solutions of dextrine, and more actively in potato or bean paste. Some cells give the iodine reaction weakly, as in *Clostridium butyricum*.

GROUP III.—LEPTOTRICHEÆ.

Genus I. *Crenothrix*.—Threads articulated; cells sulphurless; habitat water.

Genus II. *Beggiatoa*.—Threads unarticulated; cells with sulphur granules; habitat water.

Genus III. *Phragmidiothrix*.—Threads jointless; successive subdivision of cells is continuous; cells sulphurless; habitat, water.

Genus IV. *Leptothrix*.—Threads articulated or unarticulated; successive subdivisions of cells not continuous; cells sulphurless.

Genus I.—*Crenothrix*.

SPECIES.

UNASSOCIATED WITH DISEASE:—

Crenothrix Kühniana Simple saprophyte.

Crenothrix Kühniana, Rabenhorst.—Cocci, rods, and thread-forms. The cocci are globular, 1—6 μ in diam. The threads are colourless, 1.5—5 μ thick, and club-shaped at the extremity, reaching a diam of 6—9 μ . The threads form colonies with a brick-red, olive-green, or dark-brown to brown-black coloration caused by impregnation with oxide of iron. The threads are distinctly articulated, and ensheathed. The segments are set free when the sheath bursts, and develop into new threads. In other cases the segments remain enclosed, and subdivide into discs, which, by vertical fission, break up into globular forms (cocci). These again develop into new threads, either within the sheath eventually penetrating it, or after they are set free (Fig. 126).

The micro-organism appears in little whitish or brownish tufts in wells and drain-pipes, and it not only renders drinking-water foul, but may stop up the narrower pipes.




FIG. 126.—CRENOTHRIX KÜHNIANA.

a, b, c, d, e. Cocci in various stages of fission, $\times 600$.

f. Zooglæa of cocci, $\times 600$.

g. Various forms of zooglæa, natural size.

h. Colony of threads composed of rods grown out of a zooglæa of cocci.

i-r. Thread-forms; some straight, others spiral, with more or less differentiation between base and apex. (*r*) is composed of short rods at the base and above these of cylindrical segments, and at the apex these segments have divided into cocci, $\times 600$. [After Zopf.]

Genus II.—Beggiatoa.

SPECIES.

UNASSOCIATED WITH DISEASE:—

<i>Beggiatoa alba</i>	Simple saprophyte.
<i>Beggiatoa mirabilis</i>	" "
<i>Beggiatoa roseo-persicina</i>	Chromogenic saprophyte.

***Beggiatoa alba*, Vauch.**—Cocci, rods, spirals, and threads (Fig. 127). The threads are longer and thicker than leptothrix, indistinctly articulated, actively oscillating, and colourless; their protoplasm contains numerous strongly refractive granules consisting of sulphur. They occur as greyish or chalk-white gelatinous threads, 3—3.5 μ thick, in sulphur springs and marshes.

***Beggiatoa mirabilis*, Cohn.**—Threads distinguished from others of this genus by their breadth, which may reach 30 μ . They are motile, bent and curled in various ways, and rounded at the ends. Around the threads isolated cells have been observed, "macrococci," but spiral forms are as yet unknown. The threads are filled with sulphur granules. They occur on sea water, forming a white gelatinous scum on decomposing algæ, etc.

METHODS OF EXAMINING SPECIES OF BEGGIATOA.

The articulation of the threads is best demonstrated by staining with an alcoholic solution of methyl-violet, fuchsine, or vesuvin; or by treating with sodic sulphate, or warm glycerine.

Beggiatoa roseo-persicina (*Cohnia roseo-persicina*. *Bacterium rubescens*, or *Peach-coloured bacterium*, Lankester).—Cocci, rods, spirals, and threads (Fig.




FIG. 127.—*BEGGIATOA ALBA*.

A. Threads at base distinctly linked, partly spiral. B. A thread, spiral in its whole length. C, D. Fragments detached from threads; immotile. E. Active spirillum-forms, with a flagellum at either end. F, G. Thin and short spiral forms. H. A spiral showing the individual links. $\times 540$. [After Zopf.]

128). The cocci, globular or oval, reach 2.5μ in diam. They form at first solid families, bound

together by gelatinous substance. Later they become larger, globular or ovoid in shape, and hollow, containing watery fluid in their interior. The families reach a diameter of $660\ \mu$, in which the cocci form simply a peripheral layer. The hollow families or vesicles are often perforated, presenting a delicate




FIG. 128.—SEVERAL PHASE-FORMS OF BEGGIATOA ROSEO-PERSICINA.
[After Warming.]

reticulated appearance, which finally may become broken up into irregular structures. The red colouring matter can be distinguished from other red pigments, and it is designated by the name bacteriopurpurin. It is quite distinct from the pigment produced by *Bacterium prodigiosum*, being peach-blossom red, and insoluble in water, alcohol, etc. Examined spectroscopically it shows a strong

absorption in the yellow, and a weaker band in the green and blue, as well as a darkening in the more refrangible half of the spectrum. In the cocci, especially of the older vesicles, dark granules are to be seen, which consist of sulphur. The micro-organisms occur on the surface of marshes, or on water in which algæ are rotting. They form a rose-red, blood-red, violet-red, or violet-brown scum; and sometimes in such quantity that whole marshes and ponds may be coloured blood-red by them.

All the following, hitherto described as distinct species, are probably only phase-forms of Beggiatoa roseo-persicina.

Spirillum sanguineum, Cohn (*Ophidomonas sanguinea*).—Threads $3\ \mu$ and more in thickness with $2-2\frac{1}{2}$ spirals, each $9-12\ \mu$ high, with their ends provided with flagella. Their colour is due to the presence of reddish granules contained in the cells. They were observed in brackish water with putrefying substances.

Spirillum rosaceum, Klein.—Resembles *Spirillum undula*, but is reddish in colour; the colouring matter is insoluble in water, alcohol, or chloroform.

Spirillum violaceum, Warming.—Threads, crescent-shaped, or possessing $1\frac{1}{4}$ or 1 spiral, with their ends broad, rounded, and provided with flagella. The colour is due to the contents, which are violet.

Monas vinosa.—Round or oval cells of about $2.5\ \mu$ in diameter, often united in pairs. Their motion is slow and tremulous, and the cell substance pale-red, with dark grains interspersed. Flagella have not been observed. They were observed in water with decaying vegetable matter.

Monas Okenii.—Short cylindrical cells, $5\ \mu$ wide, $8-15\ \mu$ long, with rounded ends. They undergo lively movements, each end being provided with a flagellum twice as long as the cell itself. They have pale-red cell-substance with dark grains. They occur in stagnant water.

Rhabdomonas rosea.—Spindle-form cells, $3.8-5.0\ \mu$ broad, $20-30\ \mu$ long. They exhibit slow, trembling movements, having at

each end of the cell a flagellum. The cell substance is very pale, with dark grains interspersed. In brackish water.

Monas Warmingii.—Cylindrical cells, rounded at the ends; 15 μ long, 5–8 μ broad. They are possessed of a flagellum at each end of the cell, and exhibit rapid, irregular movements. The cell-substance is pale-red, and studded at the rounded ends with dark-red grains.

Genus III.—Phragmidiothrix.

SPECIES.

UNASSOCIATED WITH DISEASE :—

Phragmidiothrix multiseptata . . . Simple saprophyte.

Phragmidiothrix multiseptata.—Cocci and threads. The latter, 3–6 μ in breadth, are separated by transverse partitions into short cylindrical discs, whose height is a fourth or sixth of their breadth. Repeated transverse and longitudinal division takes place in the discs, resulting in the formation of cocci. The cocci have not been observed isolated from the threads in a free state, but they develop *in situ* into slender threads. In addition to this continuous subdivision, *Phragmidiothrix* differs from *Beggiatoa* in the absence of sulphur, and from *Crenothrix* by its wanting a sheath. They occur attached to crabs (*Gammarus locusta*) in sea water.

Genus IV.—Leptothrix.

SPECIES.

ASSOCIATED WITH DISEASE :—

Leptothrix buccalis	. . .	Saprophytic.
Leptothrix gigantea	. . .	"

Leptothrix buccalis, Robin. — Long, thin threads, $\cdot 7$ — 1μ broad, colourless, often united in thick bundles or felted together. Masses of cocci occur with the threads, and the threads themselves are composed of long rods, short rods, and cocci. The threads may break up into spiral-, vibrio-, and spirochæta-forms. The last-named occur in large numbers in the mouth, and have been named *Spirochæte buccalis*. The *Leptothrix buccalis* is found in teeth slime, and is believed to be intimately connected with dental caries. The threads penetrate the tissue of the teeth, after the enamel has been acted upon by acids generated by the fermentation of food. The short rods, long rods, cocci, leptothrix-forms, and screw-forms are found in the dental canals.

METHODS OF STAINING THE LEPTOTHRIX BUCCALIS.

The threads of *Leptothrix buccalis* have a special staining reaction (Leber). They become coloured if in an acid medium with iodine ; if the medium be alkaline, it must be acidified with very dilute hydrochloric acid or acetic acid, and the filaments then stained with iodine. The contents are stained violet, and contrast with the sheath and septa, which remain uncoloured.

Leptothrix gigantea, Miller. — Long rods, short rods, and cocci, can be observed in the same thread. There are also screw-threads, which may take the form of spirals, vibrios, or spirochætæ. The threads increase in diameter from base to apex, and corresponding with the thickness of the threads, the rods and cocci show different dimensions. They have been observed in the diseased teeth of dogs, sheep, cats, and other animals.

GROUP IV.—CLADOTRICHEÆ.

Genus I.—Cladotrix.

SPECIES.

UNASSOCIATED WITH DISEASE:—

Cladotrix dichotoma . . . Saprophytic.

Cladotrix Fœrsteri . . . "

ASSOCIATED WITH DISEASE:—

Actinomyces . . . Pathogenic.

Cladotrix dichotoma, Cohn. — Threads resembling those of *leptothrix*; slender, colourless, not articulated, straight or slightly undulated, and in places twisted in irregular spirals with pseudo-branchings. The development can be traced from the cocci to rods and threads. The latter are at the beginning simple threads, which were formerly described as *Leptothrix parasitica*, or if coloured by impregnation with iron, as *Leptothrix ochracea*. Later they form false branches by single rods turning aside, which by repeated division

lengthen into threads. A thread appears to be first composed of long rods, then of short rods,




FIG. 129.—CLADOTHRIX DICHOTOMA.

- A. Branching schizomycete:—(a) Vibrio-form; (b) Spirillum-form [slightly magnified].
 B. A screw-form with (a) Spirillum-form; (b) Vibrio-form.
 C. Long spirochæta-form.
 D. Fragment with spirillum-form at one end, vibrio-form at the other.
 E. Screw-forms: (a) continuous; (b) composed of rods; (c) composed of cocci.
 F. Spirochæta-form: (a) continuous; (b) composed of long rods; (c) short rods; (d) cocci. [After Zopf.]

and lastly of cocci. The iodine reaction must be applied to distinguish these forms, especially when

the sheath of the threads has a yellow, rust-red, olive-green, or dark-brown coloration. The cocci may grow into rods while still in the sheath, and finally become leptothrix threads, surrounded by a delicate gelatinous sheath, from which the false branching proceeds. Fragments may break off, which are actively motile, and appear as vibrios, spirilla, and spirochæta-forms (Fig. 129). They may also occur in zooglœa.

They are the commonest of all bacteria in both still and running water, in which organic substances are present. They are observed also in the waste water of certain manufactures, such as sugar. Artificially they can be cultivated on infusions of rotting algæ and animal substances, forming on these media small tufts, about 1—3 μ and floating masses.

Cladothrix Fœrsteri (*Streptothrix Fœrsteri*, Cohn).—Cocci, rod-forms, and leptothrix-threads. The threads are twisted in irregular spirals, and branch sparingly and irregularly. Screw-forms are produced by the threads breaking up into small pieces. They occur in the lachrymal canals of the human eye, in the form of closely felted masses.

There are some little-known species, which possibly belong to this group.

Sphærotilus natans.—Cells 4—9 μ long, 3 μ thick, united in a gelatinous sheath to form threads. The cells comprise rods and cocci-forms ; the cocci are set free, and

develop into rods, which again form threads. In the last a false branching has been observed. The plasma of the cells breaks up into minute, strongly refractive portions, which develop into round spores, at first of a red, and afterwards a brown colour. They occur in stagnant and flowing water contaminated with organic matter, and form floating flakes of a white, yellow, rust-red, or a yellow-brown colour.

Myconostoc gregarium, Cohn.—The threads are very thin, colourless, unarticulated, but fall apart into short cylindrical links when dried. They form gelatinous masses, 10—17 μ in diameter, singly or heaped into slimy drops on water in which algæ are decomposing.

Spiromonas volubilis, Perty.—Colourless, transparent cells, 15—18 μ long. Rapidly motile, and revolving round a longitudinal axis. They occur in marsh water and putrefying infusions.

Spiromonas Cohnii.—Colourless cells, consisting of $1\frac{1}{4}$ spirals, with both ends acutely pointed and provided with a flagellum. Breadth of the cells, 1.2—4 μ . They occur in water containing decomposing matter.

APPENDIX.




Fig. 1.



Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.




Fig. 7.



Fig. 8.



Fig. 9.




Fig. 10.



Fig. 11.

YEAST-FUNGI OR SACCHAROMYCETES AND MOULD-FUNGI OR HYPHOMYCETES.



APPENDIX A

YEASTS AND MOULDS.

Yeast-fungi and *mould-fungi*, like *bacteria* or *fission-fungi*, are *achlorophyllous Thallophytes*. They belong to two separate orders, the *Saccharomycetes* and *Hyphomycetes*, which are intimately related to each other, but quite distinct from *bacteria*. Their germs occur widely distributed in air, soil, and water, and are constantly encountered in bacteriological investigations. In addition many species are of hygienic and pathological interest or importance in being either accidentally associated with, or actually the cause of various morbid processes. For a complete account of all the described species and full details of the various forms of development,* reference must be made to botanical treatises. A description of certain species is appended here, and may afford some useful information to the worker in a bacteriological laboratory.

YEAST-FUNGI OR SACCHAROMYCETES.

***Saccharomyces cerevisiæ* (*Torula cerevisiæ*).**—

Cells round or oval, 8—9 μ long, singly or united in small chains. Spores occur three or four together in a mother-cell, 4—5 μ in diameter (Plate XXVIII., Fig. 2).

***Sacch. ellipsoideus*.**—Elliptical cells, mostly 6 μ long, singly or united in little branching chains. Two to

* Sachs, *Text-book of Botany*. 1882.

four spores found in a mother-cell, 3—3.5 μ in diam. It is widely distributed, and is the principal agent in accidental fermentation.

Sacch. conglomeratus.—Cells round, united in clusters, consisting of numerous cells produced by budding from one or a few mother-cells. There are 2 to 4 spores in each mother-cell. They occur on rotting grapes and in wine at the commencement of fermentation.

Sacch. exiguus.—Conical or top-shaped cells, 5 μ long, and reaching 2.5 μ in thickness, in slightly branching colonies. Spore-forming cells are isolated, each containing 2 or 3 spores in a row. Present in the after-fermentation of beer.

Sacch. pastorianus.—Cells oval or club-shaped. Colonies consist of primary club-shaped links, 18—22 μ long, which build lateral, secondary round or oval daughter-cells, 5—6 μ long. Spores 2 to 4. In the after-fermentation of wine, fruit-wines, or fermenting beer.

Sacch. apiculatus.—Cells lemon-shaped, both ends bluntly pointed, 6—8 μ long, 2—3 μ wide. Budding occurs only at the pointed ends. Rarely united in colonies. Spores unknown. They occur with other yeasts in various accidental fermentations.

Sacch. sphæricus.—Cells varying in form; the basal ones of a colony oblong or cylindrical, 10—15 μ long, 5 μ thick; the others round, 5—6 μ in diam. United in ramified families. Spores unknown.

Sacch. mycoderma (*Mycoderma cerevisiæ et vini*).—Cells oval, elliptical, or cylindrical, 6—7 μ long, 2—3 μ thick, united in richly branching chains. Spore-forming cells reaching 20 μ long. Spores 1 to 4 in each mother-cell. Forms the so-called "mould" on fermented liquids, and develops on the surface without exciting fermentation. When forced to grow submerged, a little alcohol is produced, but the fungus soon dies. They occur on wine, beer, fruit-juices, and sauerkraut.

Sacch. albicans (*Oidium albicans*).—Cells partly

round, partly oval or cylindrical, $3\cdot5$ — $5\ \mu$ thick, the cylindrical cells 10—20 times as long as they are thick. The bud-colonies mostly consist of rows of cylindrical cells, from the ends of which oval or round cells shoot out. Spores form singly in roundish cells. They occur on the mucous membrane of the mouth, especially of infants, in greyish-white patches which consist of epithelium, bacteria, yeasts, and the mycelia of various moulds. They can be easily cultivated in a nutrient solution containing sugar and ammoniac tartrate.* The cells germinate according to the richness of the fluid in sugar; they either grow into long threads, or, in a very strongly saccharine solution, many daughter-cells are formed, budding out in various directions (Plate XXVIII., Fig. 3).

Sacch. glutinis.—Cells round, oval, or short cylinders, 5 — $11\ \mu$ long, $4\ \mu$ wide, isolated, or united in twos. Cell-membrane and contents are colourless in the fresh state, but when dried and remoistened possess a pale-reddish nucleus in the middle. Spore-formation unknown. Forms rose-coloured, slimy spots on starch, paste, or on sterilised potatoes. The colouring matter is not changed by acids or alkalies.

Sacch. rosaceus (*Pink Torula*).—Cells 9 — $10\ \mu$ in diam. Forms a coral-pink growth in nutrient gelatine, nutrient agar-agar (Plate VII., Fig. 3) or on sterilised potatoes (Plate XV., Fig. 2). They are present in the air.

Sacch. niger (*Black Torula*).—Cells also present in the air. Cultivated in nutrient gelatine they form a black crust (Plate V., Fig. 3).

MOULD-FUNGI OR HYPHOMYCETES.

The mould-fungi have been divided into five orders: † *Hypodermii*, *Phycomycetes*, *Ascomycetes*, *Basidiomycetes*, and

* Grawitz, *Virch. Archiv*, vol. 70.

† Flügge, *Fermente u. Mikroparasiten*. 1883.

Myxomycetes. The following species, with the orders to which they belong, are of especial interest :—

HYPODERMII.

Ustilago carbo (mildew, smut).—Spores, brown, circular; episporium smooth; sporidia, ovoid cells. The spores or conidia occur as a black powder in the ears and panicles of wheat, barley, and oats.

Tilletia caries.—Spores round, pale, brown; episporium with reticulated thickenings. In germinating sporidia grow out radially from the end of the promycelium; these, at their lower part, conjugate by a cross branch, and separate from the promycelium, and at some point of the pair a hypha grows out, on which abundant secondary sporidia develop. The latter are long, oval cells, which can in turn germinate. The fungus occurs in the form of a stinking powder in grains of wheat, which renders the meal impure, and gives it a disagreeable smell.

Urocystis occulta.—The spores consist of several cells united together; partly large, dark-brown cells in the interior, and outside, several flat, semicircular, colourless cells. Spores, $\cdot 024$ mm. Promycelium germinates as in *Tilletia*, but the cylindrical cells produce a hypha, without, as a rule, previous conjugation. Occurs as a black powder in rye straw, in long disintegrated stripes, which are at first greyish. The affected plant produces abortive ears.

Empusa muscæ—Spores, $\cdot 011$ mm. in diam. A spore or conidium alighting upon the white area of the under surface of the body of the house-fly, germinates into a hypha. The latter, penetrating the skin, forms toruloid cells, which multiply by germination, and are disseminated in the blood throughout the body of the fly. These cells again grow into hyphæ, which penetrate the skin, each forming a conidium, which is cast off with considerable force. The parasite is fatal to flies, especially

in the autumn. They are often observed attached to the walls or window-panes, surrounded by a powdery substance, consisting of the extruded conidia.

Empusa radicans.—The spores form long hyphæ, which pierce the transparent skin of the caterpillar of the cabbage-white butterfly. The terminal cells ramify, and fill the body of the caterpillar with a network of mycelial filaments. The caterpillars attacked become restless, then motionless, and death ensues.

Tarichium megaspermum.—The spores are $\cdot 05$ mm. in diam., black in colour, and provided with a thickened episporium. They occur at the sides and ends of mycelial threads, attacking caterpillars (*Agrotis segetum*).

PHYCOMYCETES.

Saprolegnia.—Colourless threads, forming dense radiating tufts, occur on living and dead animal and vegetable matter in fresh water. The filaments penetrate into the substratum, and branch more or less in the surrounding water. The cylindrical ends of threads are shut off by a septum-forming zoosporangia, or mother-cells, in the interior of which a number of spherical spores, zoospores, develop. These are set free through an apical opening in the thread, and, after a time coming to rest, give rise to new plants (Plate XXVIII., Fig. 4). In the sexual mode of reproduction, a spherical bud, the *oogonium*, develops at the end of a mycelial thread; from the thread, small processes or *antheridia* sprout out laterally towards the oogonium, and blend with its protoplasm (Plate XXVIII., Fig. 4). The latter breaks up into a number of *oospores*, which clothe themselves with a membrane, while still within the mother-cell, and, eventually being set free, grow into fresh mycelial filaments. The parasite attacks fish and tritons, and produces a diseased condition of the skin, which may be ultimately fatal. In salmon it produces the common disease of salmon.

Peronospora infestans.—Mycelium, $\cdot 005$ mm. in

thickness. Twigs with as many as five branches, each bearing an egg-shaped conidium. The contents of the conidia falling off and reaching a drop of moisture, break up into a number of swarming zoogonidia, which in turn develop upon plants. Fixing themselves to the cuticle of the host, they throw a germinating filament into an epidermal cell; piercing first its outer wall, and then its inner wall, the filament reaches an intercellular space, where the mycelium develops. This continues to grow and spread throughout the plant. In tubers it can hibernate and develop in the young shoots in the following spring. The parasites appear in the form of brown patches on the green parts of the plants, especially the leaves. The attacked parts wither and turn yellow or brown in colour. If the under surface of a diseased leaf be examined, a corresponding dark spot may be observed, accompanied with a faint greyish-white bloom which covers it. The latter consists of the conidia-bearing branches of the fungus.

Pilobolus.—Hyphæ, 1—2 mm. high. Fruit-hyphæ, possessing spherical receptacles containing conidia. When ripe the receptacles with their conidia are detached at their bases, and spring by their elasticity to some distance. The mould occurs as glassy tufts on the excrement of cows, horses, etc. A cultivation can generally be obtained by keeping fresh horse-dung under a bell-glass.

Mucor mucedo.—Hyphæ colourless, simple or branched, 1—15 cm. long; sporangia are yellowish-brown or black. Spores ovoid, .008 mm. long and .0037 wide. Occurring as the familiar white mould on fruits, bread, potatoes, excreta, and penetrating into the interior of nuts and apples. A network of fibrils develops in the substance of nutrient gelatine, with formation of sporangia on the free surface. The germination of the spores and development into hyphæ can be observed in a few hours, if the fungus be cultivated in a decoction of horse-dung.

Mucor racemosus.—Hyphæ, at most 1.5 cm. long;

sporangia, yellowish to pale-brown; spores round. By continued cultivation in liquids saturated with carbonic acid, the hypha becomes shorter, and exhibits a yeast-like sprouting. These yeast-like or toruloid cells can, when the carbonic acid is withdrawn, germinate into normal mycelium.

Mucor stolonifer, Lichtheim.—Mycelium grows in the air, and then bends down and re-enters the nutrient substratum; sporangia black and spores globular. The mycelium can penetrate through the shell of eggs, and form conidiophores within them.

Mucor aspergillus, Lichtheim.—Fruit-hyphæ, thinned at the base, and with many fork-like divisions; dark-brown spores.

Mucor phycomyces, Lichtheim.—Mycelium thick-walled; olive-green fruit-hyphæ, black sporangia, and oblong spores.

Mucor macrocarpus, Lichtheim.—Spindle-formed, pointed spores.

Mucor fusiger, Lichtheim.—Ovoid spores.

Mucor mellittophorus, Lichtheim.—Spores elliptical. Found in the stomach of bees.

Mucor corymbifer, Lichtheim.—Forms branched fruit-hyphæ; sporangium has a smooth membrane. Found in the external auditory meatus; occurring also upon bread. Pathogenic in rabbits.

Mucor rhizopodiformis, Lichtheim.—Occurs on bread. The spores of *Mucor rhizopodiformis* and *Mucor corymbifer*, when introduced into the vascular system of rabbits, can germinate in the tissues, especially in the kidneys, where they set up hæmorrhagic inflammation. Dogs are immune, and only artificial mycosis is known.*

* Lichtheim, *Zeitschr. f. Klin. Med.*, vii.; Hückel, *Beitr. z. Anat. u. Phys.*, herausgeg. v. Ziegler u. Nauwerck. 1885.

ASCOMYCETES.

Oidium Tuckeri.—Fruit-hyphæ, bearing single ovoid conidia. Observed in the form of brown patches, covered with a white mildew-like layer on the leaves, branches, and young fruit of the vine, producing a "grape-disease."

Oidium lactis.—Fruit-hyphæ, simple, erect, and colourless, bearing at their ends a series or chain of conidia (Plate XXVIII., Fig. 5). In some cases the fruit-hypha branches beneath the chain of spores. Spores are short cylinders, $\cdot 0077$ — $\cdot 0108$ mm. long. The fungus is sometimes found as a whitish mould on milk, bread, paste, potato, and excrement, and is believed to be identical* with the fungus of certain human skin diseases, *Favus* (*Achorion Schoenleinii*), *Herpes tonsurans* (*Tricophyton tonsurans*), and *Pityriasis versicolor* (*Microsporon furfur*) (Plate XXVIII., Fig. 6). Cultivated artificially in nutrient gelatine, the conidia germinate into filaments of varying length, which by subdivision form septate mycelial hyphæ; these and their branches give rise in turn to spores or conidia. The differences observed in various diseases are attributed to differences in the nutrient substratum. Others† maintain that, in artificial cultivations of the spores of *Tricophyton tonsurans*, the fructification is identical with *Penicillium*.

Oidium albicans.—Vide *Saccharomyces albicans*.

Aspergillus glaucus (*Eurotium aspergillus glaucus*).—Mycelium, at first whitish, becoming grey-green or yellow-green. Spores grey-green, thick-walled, $\cdot 009$ — $\cdot 015$ mm. in diam. Sometimes found on various substances, chiefly cooked fruit (Plate XXVIII., Fig. 10).

Aspergillus repens (*Eurotium repens*, De Bary).—Fruit-heads fewer than in the above, which are at first pale and then blue-green to dark-green in colour; conidia mostly oval, smooth, $\cdot 005$ — $\cdot 008$ mm. long, colourless or pale to grey-green.

* Grawitz, *Virchow's Archiv*, vol. 70.

† Morris and Henderson, *Journ. Royal Microsc. Society*. 1883.

Aspergillus flavus.—Gold-yellow, greenish, and brown tufts; fruit-heads round; yellow, olive-green, or brown. Conidia round, seldom oval; sulphur-yellow to brown in colour, .005—007 mm. in diam. Saprophytic in man, pathogenic in rabbits.

Aspergillus fumigatus.—Greenish, bluish, or grey tufts. Fruit-heads long and conical. Conidia round, seldom oval, smooth, mostly pale and colourless. Diam. .0025 to .003 mm. Observed saprophytically in human lungs, external auditory meatus, and middle ear. The spores introduced into the vascular system of rabbits, or into the peritoneal cavity, establish metastatic foci in the kidneys, liver, intestines, lungs, muscles, and sometimes in the spleen, bones, lymphatic glands, nervous system, and skin.

Aspergillus niger (*Eurotium aspergillus niger*, De Bary).—Dark chocolate-brown tufts. Conidia round, black-brown, or grey-brown, when ripe; .0035 to .005 mm. This mould can be cultivated readily on bread moistened with vinegar, on slices of lemon, and on acid fruits and liquids. It flourishes best of all, according to Raulin,* in a liquid of the following composition:—

	Grammes.
Water	1500'
Sugar-candy	70'
Tartaric acid	4'
Nitrate of ammonia	4'
Phosphate6
Carbonate of potassium6
" " magnesium4
Sulphate of ammonia25
" " zinc07
" " iron07
Silicate of potassium07

It was also found that the fungus grew best when the quid was spread out in a layer 2 or 3 cm. in depth in a

* Duclaux, *Health Exhibition Handbook*. London: 1884.

shallow dish, and a temperature of 35° C. proved to be the most favourable. The abstraction of zinc from the nutritive liquid reduced the weight of a crop from 25 (the average) to 2 grammes, and the presence of $\frac{1}{1000000}$ part of nitrate of silver, or $\frac{1}{50000}$ part of corrosive sublimate, stopped the growth altogether. It is saprophytic in the living body.

METHOD OF EXAMINING ASPERGILLUS NIGER.

Species of aspergillus stain intensely with carmine, fuchsine, or methyl-violet ; but to examine *Aspergillus niger* with a high-power, a little special technique is employed, as follows:—A drop of glycerine is placed on a clean slide, and a drop of alcohol on a cover-glass. With a fine pair of forceps a few of the fruit-hyphæ with their black heads are immersed in the alcohol. The cover-glass is then turned over on to the drop of glycerine, and the slide held in the flame of a Bunsen burner till the spores or conidia are dispersed. To make a permanent preparation, remove the cover-glass, and transfer the fruit-hyphæ so treated to a mixture of glycerine and water (1 to 5) ; a drop may be conveniently placed ready on a slide provided with a ring of Canada balsam. The specimen is then permanently mounted by employing a circular cover-glass, and surrounding it with a ring of cement in the usual way (Plate XXVIII., Figs. 8 and 9).

Aspergillus ochraceus.—At first flesh-coloured, and then ochre-yellow heads.

Aspergillus albus.—Pure white fruit-heads.

Aspergillus clavatus.—Club-shaped fruit-heads on long stems.

Penicillium glaucum.—Occurs as a white, and later a blue-green mould, on which dew-like drops of liquid may appear (Plate XIV., Fig. 2). Its spores are present in large numbers in the air, and are liable to contaminate cultivations. Diam. of the spores '0035 mm. ; threads vary in diameter between '004 and '00071 mm.,

according to the nourishing material; the fruit-hypha bears terminally a number of branched cylindrical cells, from which chains of greenish conidia are developed (Plate XXVII., Fig. 7). It is the commonest of all moulds.

Botrytis Bassiana.—Hyphæ and spores colourless. Hyphæ usually simple, but sometimes united in arborescent stems (Plate XXVIII., Fig. 11). It is the cause of muscardine, a fatal disease of silkworms, and occurs also in various other caterpillars and insects.

UNCLASSIFIED.

Chionyphe Carteri.—Mycelium, penetrating the skin and subcutaneous tissue, sets up suppuration and ulceration. Described as the cause of a disease known in India as "madura-foot."

APPENDIX B.

ACTINOMYCES.

Actinomyces is a fungus associated with a disease occurring occasionally in man and very commonly in cattle. The micro-organism appears in the form of a rosette of pyriform or club-shaped elements, and hence the name "ray-fungus."

The fungus is believed to commonly effect an entrance by the mouth, being taken in with the food, possibly through the medium of a wound of the gum or a carious tooth. In whatever manner it has gained access to the living organism, it sets up inflammation in its neighbourhood, resulting in the formation of a neoplasm, composed chiefly of round cells. The nodules may break down and suppurate, or go on increasing in size. Fibrous tissue develops between the nodules, and large tumours often result, containing purulent cavities and excavations. In

the slimy detritus the little pale-yellow grains of fungus can be detected. In cattle the lower jaw is usually affected, and then the upper jaw and neighbouring parts. The organism may also occur in nodular tumours in the lung, subcutaneous and intermuscular tissues. It is the cause of "wens" and "wooden tongue," and also of other diseases of cattle which have been variously described before their true nature was understood as bone-canker, bone-tubercle, osteo-sarcoma, etc.

In man the pulmonary formations tend to break down early, forming fistulæ and sinuses, with the clinical character of empyema. In some cases there are symptoms of chronic bronchitis with fœtid expectoration. In other cases the disease, originating in the lung, spreads to the prævertebral tissues. If the actinomyces invade bones, as has been especially observed in the bodies of the vertebræ, caries results. The organism has been known to produce disease of the intestinal canal. The fungus has also been detected in the crypts of the tonsils of healthy pigs, and a similar, if not identical, one in the spermatic duct of the horse.*

The disease has been transmitted from cattle to cattle by inoculation,† and rabbits have been infected by means of pieces of actinomycotic tumours from human subjects, introduced into the peritoneal cavity.

THE FUNGUS IN CATTLE.

Examination in the Fresh State.

The fungus may be detected with the naked eye in the muco-purulent discharge, or in a scraping from the cut surface of a growth. The tufts of the fungus vary in size under different circumstances, from that of a grain of fine sand, to that of a pin's head. If the pus or scraping

* Johne, *Bericht über das Veterinärwesen im Königreich Sachsen für das Jahr*. 1884.

† Johne, *Deutsche Zeitschr. f. Thiermedizin*. 1881.

be spread out on a slide and examined against a dark background, the grains appear to be white or yellowish-white in colour; but if examined by transmitted light, they appear distinctly brownish. On pressing the cover-glass on the slide the grains readily flatten out, being of a soft tallowy consistency; or in the process of gently pressing the cover-glass on the slide with slight lateral movement, a distinct gritty sensation is transmitted to the finger, owing to the presence of calcareous matter. On examination with a low power the fungus will be recognised in the form of irregular patches scattered over the field, which might readily be regarded as collections of granular *débris* of a brownish or yellowish-brown colour, but on careful examination this apparent granular *débris* is observed to have a more or less characteristic appearance. On examining with a higher power, spherical, ovoid, or reniform bodies are to be seen which are either typical rosettes of clubs or granular masses, with here and there a club-shaped body at the periphery. Pus cells, round cells, fat granules, and minute spherical bodies may also be distinguished. If the grains consist of typical rosettes, and be merely covered with the cover-glass, and examined without being flattened out between the cover-glass and the slide, they will recall to mind, on focussing in turn the centre and the periphery, the appearance of the capitulum of a composite flower. The central portion appears to consist of spherical forms; these are the extremities of the component elements, and as we focus the edge of the rosette these elements are seen laterally, and their characteristic club-form is readily distinguished. The central portion may be flattened against the cover-glass, and as the individual clubs vary considerably in size, the appearance of an irregular mosaic is produced (Plate XXVIII., Fig. 1).

By pressing upon the cover we break up the rosette, and then the clubs are recognised either singly or in twos, or attached together in the form of wedge or fan-shaped

segments. Calcareous material, if present, may readily be demonstrated by the action of acids. It will be found that on the addition of hydrochloric acid, nitric acid, or acetic acid, the calcareous deposit is dissolved while the clubs are not affected, and even with the addition of the strongest acids the only result will be to dissolve out the calcareous matter and clarify the tufts of the fungus, the form of the clubs being still recognisable. They are not affected by ether or potash, and thus the effects of chemical reagents clearly distinguish them from fat crystals or calcareous particles. By breaking up the growth into small fragments, we may readily study the shape of the individual club-like elements. By using high power objectives, and properly arranging the illumination, various forms will be clearly delineated. In some cases the club will be found to be bifid at the extremity; in other cases there are lateral offshoots or daughter-clubs. Here and there will be found clubs closely pressed together like a bunch of bananas, and in other cases the broken-off pieces have a palmate form. By teasing out the grains in water, and pressing them apart between the slide and cover-glass, we find that the central portion is composed, as a rule, of a structureless core. More rarely there are the delicate filaments which are found in the disease in man.

If the grains be mounted in glycerine, the appearance of the organism in the fresh state may be preserved.

The granules may be stained by picking them out with needles and transferring to a watch-glass containing alcohol, to which a few drops of concentrated alcoholic solution of eosin have been added. They remain in the solution until distinctly stained, and they are then placed on a glass slide in a drop of glycerine.

The muco-pus may be spread out into as thin a film as possible on a cover-glass, allowed to dry, fixed by warming slightly over the flame, and stained by the methods of Plaut or Gram. The characters of the fungus can be so readily recognised in the perfectly fresh state

that methods of staining are in diagnosis of secondary importance, though there are certain minute points which can only be satisfactorily determined by means of suitable dyes.

Cultivation Experiments.

Repeated attempts by the author to cultivate the fungus from bovine specimens have failed. A section of a tongue, for example, may be made with a sterilised knife, a small portion of the growth transplanted on such media as blood serum, nutrient gelatine, and nutrient agar-agar. The little pieces of tissue remain without showing any visible growth; and though on microscopical examination the presence of the clubs can still be demonstrated, there is no distinct evidence of any increase of the fungus.

PREPARATION AND EXAMINATION OF TISSUES.

In order to examine the microscopical appearances, the tissues should be hardened in absolute alcohol and embedded in celloidin. The sections, when stained, are to be dehydrated as a rule in strong spirit instead of absolute alcohol, as the latter dissolves the celloidin. If the sections are very friable, they can be cleared with clove-oil on the slide. By these means the little fungus tufts, which have a great tendency to fall out of the sections, may be preserved *in situ* after passing through the various staining processes. To cut the sections, we can use either Jung's microtome cutting in alcohol, or the freezing microtome. In the latter case, after the celloidin has hardened it is necessary to shave off all that surrounds the piece of tissue. It is placed in water until it sinks, and then transferred to gum and frozen and cut in the ordinary way.

Staining Methods.

There are several methods by which the organism can be stained in the tissues, but it is best to employ for

this purpose Gram's method and modifications of Plaut's method.

Gram's Method.—By Gram's method the clubs in the bovine disease are distinctly stained, especially if the sections contain the fungus at a suitable stage. Use freshly prepared staining solution. A few drops of aniline-oil are placed in a test-tube, which is filled up with distilled water, the mouth of the tube closed with the thumb, and the mixture shaken up thoroughly. An emulsion forms, which is then filtered until a perfectly clear solution of aniline-water is obtained. To this is added, drop by drop, an alcoholic solution of gentian-violet until precipitation commences. About fifteen to twenty drops in a small capsule of aniline-water will be sufficient. Sections are floated in this dye for about ten minutes, then transferred to iodine-potassic-iodide solution until they turn brown like a tea-leaf. They are then decolorised in alcohol; then stained in a weak alcoholic solution of eosin, dehydrated in strong commercial alcohol, cleared in clove-oil, and mounted in balsam. It will be found that the clubs are stained blue, and that there is a central area, which is, as a rule, tinged by the eosin. There are various modifications of the method, and some of them are extremely successful in affording not only a picture of the fungus, but also the structure of the surrounding tissue. Very instructive results may be obtained by combining the method of Gram with Ehrlich's histological stain. In this case, after the section has been decolorised in alcohol, it is ready to be transferred to logwood and treated as described below (p. 385).

Weigert's Method.—This also gives very beautiful results. The sections are placed for an hour in Wedl's solution of orseille, which is prepared as follows:—Add liquid extract of orseille to a mixture of absolute alcohol 20 parts, strong acetic acid 5 parts, distilled water 40 parts, until a dark-red liquid results. This must be filtered before use. The sections are left in this solution




Fig 1.




Fig 2.



Fig 1.




Fig 2.

for an hour, then just rinsed in alcohol, and transferred to a solution of gentian-violet. Such sections show the nuclei of a violet-blue colour, and the peripheral part of the central core in the larger masses of the fungus also takes a blue colour, while the club-shaped structures are stained a striking wine-red colour (Plate XXIX.).

Plaut's Method and Modifications.—This is one of the most valuable methods for staining the clubs. The original method was to float sections for ten minutes in *magenta solution* warmed to 45° C. This solution consisted of magenta 2 parts, aniline-oil 3 parts, alcohol of specific gravity 0.830, 20 parts, distilled water 20 parts (Gibbes). The sections were then rinsed in water, stained in concentrated alcoholic solution of picric acid for from five to ten minutes, immersed in water five minutes, 50 per cent. alcohol fifteen minutes, passed through absolute alcohol and clove-oil, and preserved in Canada balsam. The clubs are stained a brilliant red and the tissue yellow. Instead of employing the magenta solution, we now use Ziehl-Neelsen solution (Plate XXX.).

By removing the picric acid in Plaut's method by prolonged immersion in alcohol, and then staining with gentian-violet or methylene-blue, a very successful contrast can be obtained. The most instructive histological picture can be obtained by first staining with Neelsen's solution, removing the stain from the tissue in the way that has been already described, and then transferring the sections to distilled water, and subsequently staining with Ehrlich's histological stain.

Ehrlich's New Histological Stain.—This is a combination of Ehrlich's logwood with orange-rubin. It is of especial value for sections of actinomycosis, and particularly in combination with carbolised fuchsine. It is employed in the following way:—The sections must be placed in alcohol or distilled water, and then in Ehrlich's logwood for about half a minute. From this solution they are transferred to distilled water, washed to remove

the excess of stain, and then placed in a large dish of tap-water, where they are left for half an hour or more, until the sections turn blue; if preferred, they may be left overnight. They are then stained for one or two minutes in a solution of rubin. S, and orange, and washed again in distilled water to remove the excess. They must then be dehydrated in alcohol, cleared in clove-oil, and mounted in balsam.

Preparation of Large Sections.

The method of cutting large sections of organs, which has somewhat recently been introduced, is one which may be very advantageously employed in studying actinomycosis. The value of these sections depends, not only upon their affording an instructive picture of the naked-eye appearances, but they can also be studied with a pocket lens, or under the microscope with a $\frac{1}{4}$ or $\frac{1}{8}$ -in. objective. By staining the sections the relation of the morbid to the healthy structures is brought out in greater contrast, and thus the topography of the disease can be studied more minutely than by simply observing the cut surfaces of organs or growths. And, further, it affords a means of rendering permanent many of the instructive appearances observed at the autopsy, without preserving the whole structure in the form of a museum specimen. Very satisfactory results can be obtained with material hardened either in spirit or Müller's fluid. The fresh material is cut with a large keen-edged knife into slices about a quarter of an inch, or less, in thickness. These slices are placed between filter-paper in large porcelain dishes, such as are employed for photographic purposes, and well covered with the hardening solution, which should be frequently changed. By covering the slice with a small pane of glass, which is lightly weighted, any curling or turning up of the edges is prevented, and the slice not only kept flat, but hardened with smooth surfaces. Several weeks are required for hardening in

Müller's fluid. The slices, after a short time in water, are placed in gum, and then frozen and cut; the slices which are hardened in alcohol are soaked in water until all trace of the spirit has been removed. A large microtome on the Bruce model is used to freeze and cut the sections. But in some cases it will be found better to embed the slices in celloidin, and cut under alcohol with a large microtome of Jung's pattern. The sections are carefully removed from the blade of the knife with a large camel's-hair brush, and in the case of frozen sections floated in water.

The next process is to float a section out in spirit, and with the camel's-hair brush to unfold it and spread it out on a sheet of glass. The glass with the section is lifted out and examined, and if the section is sufficiently thin, transferred to the staining solution. In the same way the section is passed through the various stains, as it should be prevented from rolling up or folding in the dye, or it may not be evenly stained throughout. Modifications of this process will suggest themselves, such as pouring off the dye and leaving the section spread out at the bottom of the dish, and then using the same dish for the next process. The sections are so easily injured, that it is better, as much as possible, to avoid handling them. If the sections are only a few inches in diameter, such as transverse sections of the anterior portion of the tongue of an ox, they can readily be transferred from dish to dish by means of a large spatula made by soldering a piece of sheet German silver to thick copper wire.

To stain them employ carbolised fuchsine and picric acid, or alum cochineal, or logwood and orange-rubin. The processes of staining are precisely the same as with ordinary sections; but, from their unusual size, experience and practice are required in their manipulation.

When the section is dehydrated, it is ready to be cleared in clove-oil. The glass on which it is to be permanently mounted should be selected without scratches or flaws,

and thoroughly cleaned and polished. It is slipped under the section, which is evenly spread out upon it, and then lifted out of the dish. The excess of spirit is drained off, the glass placed on a level surface, and clove-oil poured on the section. It is left until completely clarified; the clove-oil, as much as possible, drained off, and the rest entirely removed by gentle pressure with several thicknesses of best filter-paper. In this way several large sections can be cleared at the same time. But when only one or two sections are dealt with, they are cleared in clove-oil in a dish, and the mounting glass at this stage passed underneath them as already described. Another plan which will be found of advantage is as follows:—A piece of clean thick filter-paper rather larger than the section is slipped underneath it, and then raised with the section upon it. After allowing the excess of clove-oil to drain back into the dish, it is carefully laid on the glass with the section downwards, and gently pressed down. By taking up a corner, the filter-paper is peeled off, and the section left behind on the glass. Any creases or folds are adjusted with needles. After removal of the clove-oil, balsam is run over the section, and a cover-glass gently and dexterously lowered, so as to avoid the presence of air-bubbles. The preparations are set aside to harden in a warm place and on a level surface, and are then ready for fixing in suitable frames.

Naked-eye Appearances of Large Sections.

In the sections of an actinomycotic tongue it is at once apparent that the new growth is more or less limited to the periphery of the section. In parts there are dense clusters of little nodular neoplasms, *the fungus systems*, each having a rounded form, and averaging in size that of a small pea. In other parts small nodules, varying in size from a millet-seed to a hemp-seed, have a linear arrangement between bundles of muscular fibres. The

appearance is suggestive of an invasion of the tongue along the lymphatics.

In many of the nodules the largest tufts of the fungus can be seen, with the naked eye, to occupy a more or less central position. In parts the muscular fibres are replaced by fibrous tissue.

If now these sections be placed under the microscope, the minute structure may be examined; but as it is obvious that still better results may be obtained by small sections, any part which it is necessary to examine with high powers can be selected from a corresponding part of the growth and prepared in the ordinary way.

In the case of a "wen" the whole growth can be excised with the surrounding tissues, sliced and treated in the way already described, and sections stained by different methods.

The nature of the growth is at once recognisable as actinomycosis from the characteristic honeycombed appearance produced by the trabeculæ of fibrous tissue which form a spongy structure, from the loculi of which the fungus tufts and caseous matter have for the most part dropped out. In other parts this structure is intact, and the tufts of the fungus can be detected with the naked eye, and readily recognised with a pocket lens.

A fungus system may be studied more minutely in ordinary sections of the tongue. Each nodule is composed of the actinomyces surrounded by round cells and epitheloid cells, and beyond this fibrous tissue often forming a distinct capsule. In some specimens the fungus is surrounded by a single row of large multinucleated cells, and in other specimens the fungus is found in the interior of large oval giant cells.

THE FUNGUS IN MAN.

Examination in the Fresh State.

A close examination of pus from a case of actinomycosis reveals the yellowish grains which a casual observer

might mistake for grains of iodoform. On collecting some of the discharge in a test-tube, and holding it between the light and the eye, the tufts of fungi appear as brownish or greenish-brown grains, embedded in a mucopurulent matrix.

On spreading some of the discharge on a glass slip, and examining in the same way described for the bovine disease, the largest tufts of the fungus are found to be about the size of the head of a small pin. They have a distinctly sulphur-yellow colour by reflected light, but appear of a yellowish or greenish-brown tint by transmitted light. With a needle or a platinum wire flattened at the end into a miniature spatula, the grains can be readily picked out of the discharge, or taken off the dressing, transferred to a clean slide, and gently covered with a cover-glass. Examined with an inch objective, they have the appearance of more or less spheroidal masses of a pale greenish-yellow colour. On removing the preparation from the microscope, and gently pressing down the cover-glass with the finger, the grains flatten out like specks of tallow, and on again examining with the same power they are found to have fallen apart into a number of irregular and sometimes wedge-shaped fragments of a faintly brown colour, affording a characteristic appearance. By preparing another specimen, and covering it with a cover-glass without completely flattening out the grains, the spherical, oblong, and reniform masses of which the tufts are composed can be recognised with a $\frac{1}{8}$ th in. objective, like the rosettes of clubs which are observed in cattle. By examining the peripheral part of a rosette with a $\frac{1}{12}$ th in., and especially after pressing the grains into a thin layer, with or without the addition of a drop of glycerine, the characteristic clubs are most readily demonstrated, and the most varied shapes observed by carefully examining the form of the individual elements. As in the bovine fungus, every variation in form is found from single clubs to clubs with lateral offshoots, clubs




Fig. 1.




Fig. 2.

bifid at the extremity, palmate or fan-shaped groups, and banana-like bunches. In many cases the clubs are divided by transverse fission into two, three, or more segments. As a rule, the clubs are irregular in shape, and of about equal size, while a few are conspicuous by their length. In other parts of a preparation the clubs are replaced by long slender forms, which are sometimes transversely divided into a number of short links. With suitable illumination many clubs are seen to taper off into slender filaments. In addition there are free filaments, which are twisted, branched, and sometimes distinctly spirilliform. Many of the clubs are composed of layers differing in their refractive power, and many have the appearance of a central channel. There are also in the preparation small highly refractive bodies, fat granules, granular detritus, round cells, pus cells, and sometimes blood corpuscles (Plate XXXI.).

The grains differ, as a rule, from those from a bovine source, in the absence of that sensation of grittiness so often transmitted to the finger when pressing the cover-glass upon them, and in the slightly greater tendency of the tufts to retain their compact form. By teasing the grains in a drop of water on a slide, and examining the preparation with a sixth or a twelfth objective, the explanation of the latter is forthcoming, for by this process the clubs are gradually washed away, and a central core remains which is composed entirely of a dense network of filaments. This can readily be observed by using a small diaphragm, and it will be found that the rosettes of clubs are now replaced by tangled masses having some resemblance to miniature tufts of cotton-wool. These filaments constitute the mycelial network which is seen in sections stained by the method of Gram; this can be readily verified by making a cover-glass preparation of the grains and staining by that method. The characters of the fungus can readily be studied by proper illumination without staining. The clubs have a faintly greenish tint, and

in form and arrangement are quite characteristic and easily recognisable. Permanent preparations may be made by mounting the fungus in glycerine.

THE FUNGUS IN STAINED SPECIMENS.

The fungus may be stained in alcoholic solution of eosin, in the manner already described for the bovine organism, or in orange-rubin, and in either case mounted in glycerine. But although the fungus can be detected without any staining process, there may sometimes be doubtful appearances, and then cover-glass preparations should be made and stained by the method of Gram and eosin. The filaments can be readily recognised, and this is of great value, as it forms an additional means for the diagnosis of the disease. In combination with orange-rubin we have a test that is as characteristic and useful as staining for tubercle bacilli. The discharge, scraping from a growth, sputum, or the isolated fungus is squeezed between two cover-glasses, which are then slid apart; they are allowed to dry, passed through the flame in the ordinary manner, and then stained. The cover-glasses can be cleared in clove-oil, the excess of clove-oil being removed by gentle pressure between pieces of blotting-paper, and then the preparation can be mounted in balsam and rendered permanent. On examination of these specimens the masses of filaments will be found to be stained blue, and the tissue elements pink. These filaments vary very much in extent and character in different preparations. In some cases there are masses of short threads which are straight, or sinuous, or twisted, and sometimes branched. In other parts the field is occupied by very short, straight, or curved and sometimes spiral fragments; in others, again, there are comparatively long strands. On examination with a high power, and with careful illumination, some filaments will be observed to be moniliform, while others are provided with a terminal oval body. There are also free spherical




Fig. 1.




Fig. 2.

round bodies stained blue, which possibly represent the spores of the organism. If orange-rubin be used instead of eosin, the clubs will be stained and easily recognised. And this method enables one to determine the exact relation of the threads to the club-shaped bodies. This is an interesting point, as it has been suggested that the threads are not connected with the clubs, but are merely an adventitious micro-organism growing in the track of the ray-fungus. The threads are stained blue and the clubs crimson (Plate XXXII.). In the younger clubs the protoplasm of the thread can be traced into the interior of the club. In some of the older clubs the central portion takes a yellowish stain, and in others the protoplasm is not continued as a thread, but is collected into a spherical, ovoid, or pear-shaped mass. In others, again, irregular grains, stained blue, are scattered throughout the central portion. The sheath of the thread is stained pink; and the protoplasm, stained blue, fills the sheath, or consists of small spherical and irregular grains, giving a distinctly beaded appearance.

The effect of various reagents should be tried upon the isolated grains. The grains are picked out of the pus and transferred to watch-glasses containing strong potash, xylol, and benzole. If returned to a slide and covered with a cover-glass, the clubs are found unaltered. Water or weak potash washes away the clubs, and the filaments become easily distinguished; ether and strong acids have no effect upon them. Corallin soda, Hanstein's violet, and iodine zinc-chloride fail in giving any particular reaction. Hoffman's blue stains the clubs, but without bringing out any structural details which could not be observed in the unstained specimens.

CULTIVATION EXPERIMENTS.

Boström succeeded in cultivating the fungus outside the animal body; and finding that his cultures consisted of what appeared to be cocci forms, short rods, and threads,

but no clubs, he advanced the theory that the fungus belonged to the bacteria, forming one of the *Cladothrix* group, and possibly closely allied to the *Streptothrix Færsteri* of Cohn. In accordance with this view, Boström regarded the clubs as comparable to the degenerate or involution forms, which are often found in old cultivations of bacteria. They constituted, in his opinion, the lifeless organism.

According to the old theory the actinomyces was a hyphomycetous fungus, and the clubs were the flask-shaped structures or gonidia. In the hope of throwing some light on this point, the author has made attempts at cultivation upon a number of different media. Glycerine agar-agar, a favourable medium for the growth of the tubercle bacillus, suggested itself as a suitable soil for actinomyces. An actinomycotic abscess was opened, the discharge collected in sterilised tubes, and cultivations prepared with as little delay as possible. Some of the discharge was spread out on a sterilised glass slide, and the grains isolated with sterilised needles and quickly transplanted on the surface of the nutrient medium. The tubes were placed in the incubator at 37° C., and the result watched from day to day. For several days there was no promise of success to the naked eye, but gradually the grains began to change, and by the end of a fortnight there was an appreciable increase in size. Numerous cover-glass preparations were made from what was originally a single grain, and on examination by the method of Gram the appearance was very striking. There could be no doubt as to the increase of the mycelial structure. The dense masses of filaments covered almost the whole area of the preparation. In parts less thickly covered there were a vast number of oval bodies and rod-like segments with terminal enlargements. These "crocus" forms corresponded with the appearances previously described as met with in the interior of certain clubs. From this it would appear that

some other condition is necessary for the development of the fully formed club, this being the result of the sheath undergoing some change, possibly mucilaginous, resulting in the formation of a thick investment of the clubbed mass of protoplasm at the end of the thread.

These club-shaped bodies represent organs of fructification, rather than the results of degeneration or death. The difficulty in accepting the view of their being entirely lifeless forms lies in the fact that one can observe daughter clubs growing from the mature clubs; and, further, in the bovine fungus the author has been able to trace by this process the stages in the development of a single club to a completely formed rosette.

In the unstained condition, the clubs are found, on the whole, to be very regular in their form and arrangement, while by certain staining methods they can be shown to have a somewhat complex structure. If we take all the characters into account, and particularly the minute structure and the relation to each other of the threads and clubs, we are justified in the opinion that the club in the early stages is an integral part of the living fungus, and that these characters bring the fungus into relation with a higher group of micro-fungi, the Basidiomycetes, although the filaments, regarded by themselves, correspond with the characters of streptothrix.

LIFE-HISTORY.

It has not been possible to trace every step in the life-history of many of the Basidiomycetes; but if we regard the ray-fungus in the light of what is known to occur in many species, we may explain its life-history as follows:—The spores sprout into hyphæ, which form excessively fine, straight, or sinuous, and sometimes distinctly spirilliform threads, which branch irregularly and sometimes dichotomously. The extremities of the branches develop the club-shaped bodies. The clubs are closely packed together, so that a more or less globular body is formed,

with a central core composed of a dense mycelium. The threads can be differentiated by the method of Gram into an external sheath, and protoplasmic contents. The club-shaped body externally appears to be mucilaginous, while internally it is continuous with the protoplasm of the thread. It is difficult to say what further changes occur in the club-shaped bodies; in all probability they represent the organs of fructification. If so, the protoplasm in the interior of the club may possibly undergo changes leading to the development of spores, which are ultimately set free; in some cases the terminal segment of a club is separated by transverse fission in the form of a globular body, a process resembling the formation of spores by abjunction. In others, the forms sprouting from the club are suggestive of teleutospores. In whatever way they may be formed, there can be little doubt that spores are set free in the vicinity of a rosette, and give rise to fresh individuals; the ultimate result recalling, as has previously been suggested, the appearance of "fairy rings." There can be little doubt that spores and young fungi are taken up by wandering cells, and conveyed to a distance from the parent fungus, and thus fresh centres of growth are established.

According to the view here propounded, the filaments or threads may be regarded as extremely delicate hyphæ forming a dense mycelium, and the more familiar clubs as the basidia of the hymenial layer. There are occasionally long slender forms, very different from the ordinary clubs; they possibly represent *paraphyses* or abortive elements.

APPENDIX C.

FLAGELLATED PROTOZOA IN THE BLOOD.*

WHEN examining blood the bacteriologist must be prepared to meet with minute organisms which at the first glance under moderate amplification may be mistaken for vibronic or spiral forms of bacteria. The organisms referred to belong not to the vegetable but to the animal kingdom. They may occur associated with disease, but




FIG. 130.—PARASITES IN THE BLOOD OF RATS [after Lewis].

they appear to be more commonly found in the blood of apparently perfectly healthy animals.

Flagellated organisms in the blood of rats and hamsters.—Lewis† described peculiar organisms in the blood of healthy rats in India. When first noticed they were thought to be vibrios or spirilla. A drop of blood under examination appeared to quiver with life, and on diluting the blood motile filaments could be seen

* Abstract of paper by the Author, *Journ. Roy. Micros. Soc.*, read November 10th, 1886.

† Lewis, *Microscopic Organisms in the Blood of Man and Animals*. Calcutta, 1879 (with photographs); and *Quart. Journ. Micr. Sci.*, lxxiii. (1879), pp. 109-14, and xxiv. (1884), pp. 357-69.

rushing through the serum, and tossing the blood-corpuscles about in all directions. Under careful examination the filaments were found to consist of a thicker portion or body, with at one end a flagellum (Fig. 130).

After fixing with osmic acid, they measured $0\cdot8-1\ \mu$ in width, and $20-30\ \mu$ in length; the flagellum was about as long as the body, so that the total length of the organism was about $50\ \mu$. Lewis detected these parasites in 29 per cent. of the species *Mus decumanus* and *Mus rufescens*. Though they had many features in common with motile organisms of vegetable origin, they appeared to approach much more closely to the Protozoa, more particularly several of the species of Dujardin's *Cercomonas*.

Wittich* discovered, in the blood of hamsters, whip-like bodies with lively movements. They resembled frog's spermatozoa, possessing a thick portion continued into a long lash-like thread. Wittich considered them identical with the organisms described by Lewis, and they also were observed in apparently healthy animals. Koch† later met with the same organisms.

Flagellated organisms in the blood of horses, mules, and camels.—In India a fatal disease, known by the natives as *Surra*, occurs in horses, mules, and camels. The malady is described as a blood disease, characterised by fever, accompanied by jaundice, petechiæ of mucous membranes, great prostration, and rapid wasting, terminating in death. Evans‡ observed the presence of a parasite in the blood, and by means of subcutaneous inoculation, and by the introduction into the stomach of blood containing the parasites, the disease was transmitted to healthy animals.

* "Spirillen im Blut von Hamstern," *Centralbl..f. Med. Wiss.* 1881, No. 4.

† *Mittheilungen aus dem Kaiserlich. Gesundheitsamt.* 1881.

‡ Evans, *Report published by the Punjab Government Military Department*, No. 439. 1880.

Steel,* who was deputed to investigate this disease in British Burma, also found the parasite in all cases, and further observed that it appeared as the temperature rose and disappeared during the apyrexial periods. This observer concluded that the organism was a spiral bacterium, and named it, after its discoverer, *Spirochæta Evansi* (see p. 402).




FIG. 131.—HEMATOMONAS COBITIS.

- a*, First variety; *b*, second variety; *c*, third variety.
d, First variety in a state of diminished activity.
e, The same after treatment with osmic acid.

[After Mitrophanow.]

Flagellated organisms in the blood of fish.

—In the blood of mud-fish (*Cobitis fossilis*) Mitrophanow† observed the presence of peculiar micro-parasites (Fig. 131). As a 1 per cent. salt solution had been added to the blood under examination, it occurred to Mitrophanow

* Steel, A.V.D., *An Investigation into an Obscure and Fatal Disease among Transport Mules in British Burma*. 1885.

† "Beiträge zur Kenntniss der Hämatozoen," *Biol. Centralbl.*, iii., 1883, pp. 35-44.

that they were possibly the cytozoa described by Gaule ; but this idea was dismissed by the fact that they were found in blood to which no salt solution was added. Their size varied from 30 to 40 μ in length, and 1 to 1.5 μ in width. At first their rapid movements baffled examination, but as the rapidity lessened there was the appearance of a curling movement in the body portion, and a swinging movement of the lash. The organism moved in the




FIG. 132.—ORGANISMS IN THE BLOOD OF THE CARP.

a, b, c, Hæmatomonas Carassii. *d, e, f, g, h,* Other organisms in the same blood [after Mitrophanow].

direction of the lash, the anterior end of the body being more pointed than the posterior, and gradually fining off into the lash. When the body seemed to rest, the lash might be seen to whip out in all directions. As the movement of the body gradually diminished, it appeared to have a complicated screw-form, the axis of the screw corresponding to the body, to which an undulating membrane is fastened spirally. This could be distinguished when the organism was dying, because the body in death

contracted, and the membrane then looked like a spiral addition. Thus the organism consisted of a body, a spiral membrane, and a flagellum.

With higher magnification the organism appeared to consist of a refractive, strongly contractile, protoplasmic substance, which, when death occurred, formed a shapeless mass. In the same blood two other forms were observed, one without a membrane, but having two highly refractive spherules in the protoplasm, and another with neither membrane nor flagellum, consisting of very granular protoplasm with several refractive spherules, and capable of protruding processes like pseudopodia.

In the blood of the German carp (*Cyprinus Carassius*) Mitrophanow describes a parasite which is perceptibly larger, and possesses an undulating membrane fastened along the edge of the long body (Fig. 132). When the body bent first towards one side and then to the other, a wave-like movement was observable at the free edge of this membrane.

In *Cobitis fossilis* these parasites were found in all the fish examined except one, and in great numbers in the hot months. In *Cyprinus Carassius* they were only found occasionally. Mitrophanow described other varieties, which he considered were possibly not complete organisms, but developmental forms. He considered that these organisms were infusoria between the genera *Cercomonas* and *Trichomonas*, with great similarity to the *Trichomonas* described in the Lieberkühn's glands of fowls and ducks (Eberth*).

On account of their special habitat Mitrophanow suggested a new genus (*Hæmatomonas*), defining this genus as follows:—Parasites of normal fish-blood, worm-like, actively moving organisms, with indistinct differentiation of body parenchyma. Bodies pointed at both ends, 30 to 40 μ long and 1 to 1.5 μ wide. May possess in front a flagellum, and on one side an undulating membrane.

* *Vide* Leuckart, *The Parasites of Man*, translated by Hoyle p. 248.

Species :—

Hæmatomonas cobitis.—Body provided with a spiral membrane and a flagellum at the fore-end. Parenchyma of body homogeneous. Second variety, body and flagellum only. Movement undulatory, body containing highly refractive spherules. Third variety, plasma-like body, without membrane or flagellum; quickly changes form by sending out processes laterally, and contains two to four refractive spherules. Blood of *Cobitis fossilis*.

Hæmatomonas Carassii.—Long bodies, with narrow membrane attached along the whole length; less actively




FIG. 133.—“SURRA” PARASITES OCCURRING SINGLY AND FUSED.
From preparations stained with magenta, $\times 1200$.

motile. Several forms also observed strikingly smaller than the above; many disc-shaped. Often seen attached to a red corpuscle, setting them in motion by their movements. Blood of *Cyprinus Carassius*.

Quite recently the author has investigated the parasites found in the disease known as Surra, and came to the following conclusions :—

In stained preparations the somewhat tapering central portion, or body, of the parasite is found to be continuous at one end with a whip-like lash, and at the other end to terminate in an acutely pointed stiff filament, or spine-like process. Here and there, possibly from injury or want of development, the spine-like process appears to be blunted

or absent. By very careful focussing on the upper edge of the central portion, the author discovered the existence, much more markedly in some of the parasites than in others, of a *longitudinal membrane* with either a straight or undulating margin (Fig. 133). The membrane is attached along the body, arising from the base of the rigid filament, and becomes directly continuous at the opposite end with the flagellum. In some cases the edge only is deeply stained, giving the appearance of a *thread continuous with the flagellum*, so that one might be easily led to overlook the membrane, and imagine that the flagellum arose from the opposite end of the body, at the base of the spine-like process.

Close to the base of the spine-like process a clear unstained spot is, in many parasites, easily distinguished, and at the opposite end there is, in some, the appearance of the deeply stained protoplasmic contents having contracted within the faintly stained cell-wall. Where the longitudinal membrane has a wavy outline, the undulations are much more marked in some cases than in others. Here and there the wavy outline appears first on one side of the central portion, and then on the other, but there never is any waving outline on both sides of the same part of the body, and this is explained by careful examination, which shows that in dying the somewhat ribbon-like parasite has become doubled on itself.

Owing to the somewhat curved and twisted shape of the parasite and the curling of the flagellum, in the stained preparations, it was difficult to make exact measurements; but the average width, according to whether the membrane was visible or not, varied from 1 to 2 μ , and the length of the body from 20 to 30 μ . The flagellum was about the same length as the body.

From these observations (especially the discovery of the undulating longitudinal membrane) the author recognised a very close resemblance to Mitrophanow's descriptions, and concluded that, if we followed the classification

adopted by Mitrophanow, the genus *Hæmatomonas* must not be restricted to organisms in fish-blood. It must be expanded to include this parasite of mammalian blood, which should in that case be named *Hæmatomonas Evansi*, rather than *Spirochæta Evansi*, as proposed by Steel.

In the course of this investigation the author was led to examine the blood of rats obtainable in this country. Organisms were discovered in the blood of about 25 per cent. of common brown rats; and after examining them with various objectives, from a $\frac{1}{8}$ dry to a $\frac{1}{25}$ oil-immersion of Powell and Lealand, the following conclusions were arrived at:—That they are polymorphic,



FIG. 134.—A MONAD IN RAT'S BLOOD, $\times 3000$.

The organism is represented at partial rest, with its posterior filament impinging on a corpuscle, and showing the undulating longitudinal membrane, the long flagellum, and the refractive spherules in the granular protoplasm.

presenting for the most part slightly tapering bodies, which terminate at one end in a stiff, immotile, acutely pointed, flexible filament or spine-like process, and at the opposite end are provided with a long flagellum, while longitudinally attached a delicate undulating, fin-like membrane can be traced, which starts from the base of the posterior filament, and becomes directly continuous with the flagellum (Fig. 134).

With careful illumination the body is found to be distinctly granular, with one or more highly refractive spherules. When the rapid movement is arrested, the undulating membrane is distinctly visible. The best opportunity occurs for seeing this when the organism comes to partial rest with its stiff filament against a

corpuscle, as if to obtain a *point d'appui*, while lashing its flagellum in all directions (Fig. 135, *b*). At other times, when the parasite has impinged with its posterior extremity against a corpuscle, or the stiff filament is apparently entangled in *débris*, the movements of the organism give one the idea of its endeavouring to set itself free.

In the active state the thicker portion, or body, appears to twist and bend from side to side with great activity. The organism can turn completely round with lightning rapidity, so that the flagellum, at one moment lashing




FIG. 135.—MONADS IN RAT'S BLOOD, $\times 1200$.

a, A monad threading its way among the blood-corpuscles; *b*, another with pendulum movement attached to a corpuscle; *c*, angular forms; *d*, encysted forms; *e* and *f*, the same seen edgewise.

in one direction, is suddenly observed working in the opposite direction. Then suddenly the organism makes progression, and it can be distinctly seen to move in the direction of the flagellum, the flagellum threading its way between the corpuscles and drawing the rest of the organism after it. By treating cover-glass preparations with osmic acid, the appearances corresponded exactly with photographs of the organisms observed by Lewis in India, so that the author has no doubt of their identity, in spite of the descriptions not completely according. A great likeness to the organisms described by Mitrophanow, and to the Surra parasite, as just described, was obvious; and

after staining the rat parasites, the closest examination confirmed the belief that they were morphologically identical with the stained parasites of Surra.

The cover-glasses with a thin layer of blood should be passed through the flame of a Bunsen burner in the way commonly employed for examining micro-organisms, and stained with an aqueous solution of fuchsine, methyl-violet, or bismarck-brown. The following method will, however, be found most instructive:—Use freshly prepared saturated solution of fuchsine or methyl-violet in absolute alcohol, and put a drop with a pipette on the centre of




FIG. 136.—MONADS IN RAT'S BLOOD, showing membrane under different aspects, blood-corpuscles some crenated, and stained discs, $\times 1200$.

the preparation; do not disturb the drop-form for a few moments; then, before the alcohol has evaporated, wash off the excess of stain. It will be found that where the drop rested the organisms will be very deeply stained, while in the surrounding area the colour will vary in intensity.

By the effect of the different degrees of staining much may be learnt (Fig. 136). In one organism the body and entire membrane will be equally stained; in another the margin of the membrane only. In some the posterior stiff filament is stained, and at its base a darkly stained speck is very striking; and in other cases, again, the posterior filament is only faintly tinged, or an unstained spot occurs near its base.

The morphological identity of the rat and Surra parasites is thus established, and both seem morphologically identical with the organism of Mitrophanow. If we follow Mitrophanow, we must, therefore, enlarge his genus of *Hæmatomonas*. The author ventures, however, to disagree with Mitrophanow in the advisability of adopting this entirely new generic name. Mitrophanow suggested this new term because of the special habitat—normal fish-blood—of the species he discovered. But the characteristic features of these organisms are the characteristic marks of the genus *Trichomonas*.* They are, therefore, embraced by the genus *Trichomonas*, and there is no need to create a new one. If it were not for the different description given by Mitrophanow of the organism in the blood of *Cobitis fossilis*, the author would be inclined to say that all these organisms belonged to one and the same species, which might well be named *Trichomonas sanguinis*. The monads in the rat and the Surra parasite have been shown to be morphologically identical with each other, and both, as far as one can judge from the description, are morphologically identical with the monad in the blood of the carp. We have, however, seen that the organism in Surra is believed to be pathogenic, and too much stress must not be laid on morphological identity. There is strong evidence in favour of believing in its pathogenic properties; but, at the same time, it must be borne in mind that the organism has never been *isolated apart from the blood*, and the disease then produced by its introduction into healthy animals. It is quite possible that the parasites in Surra are only associated with the disease, the impoverished blood affording a suitable nidus for their development, while the contaminated water may be the common source of the organism and of the disease. On the other hand, the organism in the rat is found in apparently perfectly healthy, well-nourished animals.

* *Vide* Leuckart, *The Parasites of Man*, translated by Hoyle, 1886.

APPENDIX D.

HÆMATOZOA OF MALARIA (Laveran).

IN 1880, Laveran in Algiers noticed the existence of peculiar structures in the blood of a patient suffering from malaria. It occurred to him that he had discovered the "*Microbe du paludism.*" Laveran followed up his observation by researches which were communicated to the Academy of Medicine in Paris, in 1881 and 1882, and they were subsequently published *in extenso* in a treatise on the subject in 1884.*

Laveran described various bodies which he was led to regard as different stages in the life-history of the same micro-parasite. The most striking forms were cylindrical elements with pointed extremities. They were crescent-shaped and pigmented in the middle. There were other forms, more frequently found, which were either free in the serum or in contact with the red blood-corpuscles. They were more or less spherical, pigmented, and endowed with amœboid movement. Other forms, again, were provided with motile filaments three or four times as long as the diameter of a red blood-corpuscle. And, lastly, there were little masses of hyaline material, which Laveran regarded as dead forms.

These observations at first attracted little attention; but they have been confirmed and extended by Richard,† Councilman and Abbot,‡ Marchiafava and Celli,§ Golgi,||

* Laveran, *Traité des Fièvres Palustres.* 1884.

† Richard, *Comptes Rendus.* 1882.

‡ Councilman and Abbot, *American Journal of Medical Sciences.* 1885.

§ Marchiafava and Celli, *Archivio per le Scienze Mediche.* 1885, 1886. *Atti della R. Accademia Medica.* 1886—1887. *Archiv. Ital. de Biolog.* 1887—1888.

|| Golgi, *La Riforma Medica.* 1888. *Archivio per le Scienze Mediche.* 1886. *Fortschritte der Medicin.* 1889.

Sternberg,* Osler,† Crookshank, and Vandyke Carter,‡ and their importance fully recognised.

The different forms assumed by the hæmatozoon of malaria may be described in two groups: those within the red blood-corpuscles and those free in the serum.

Intra-corpuscular bodies.—These are of three kinds:—*First*, structureless protoplasmic bodies, much smaller than, and within or attached to, the red blood-corpuscles (Fig. 137). These rapidly change their shape,



FIG. 137.—NON-PIGMENTED AMŒBOID FORMS [after Marchiafava and Celli].

exhibiting amoeboid movement. They were first described by Marchiafava and Celli, and possibly represent the first stage in the life-history of the hæmatozoon. Marchiafava and Celli suggested the name *Plasmodium malarie*. *Second*, minute masses of finely granular or of hyaline



FIG. 138.—PIGMENTED AMŒBOID FORMS [after Golgi].

protoplasm enclosing granules of pigment (Fig. 138). These forms are sometimes present in large numbers, and at other times can be found only with difficulty. They are more or less spherical, but exhibit amoeboid movement, and rapidly change their form. The pigment

* Sternberg, *Medical Record*. 1886.

† Osler, *British Medical Journal*. 1887.

‡ Carter, *Scientific Memoirs Med. Officers Army of India*. 1888.

granules are also in active movement. There may be one or more of these amœboid bodies to a blood-corpuscle, and they vary in size; one may occupy the whole of the corpuscle. In cases of pernicious malaria similar bodies may be seen in tissue sections, in the corpuscles filling the capillaries. *Third*, forms which appear like isolated grains, and larger homogeneous bodies surrounded by clear spaces which change in outline.

Extra-corpuseular bodies.—These are the most striking and perhaps the most interesting forms. *First*, the semi lunar bodies of Laveran. These are crescent-shaped bodies, sometimes pointed at the extremities, but more usually rounded off (Fig. 139). They are not



FIG. 139.—SEMI LUNAR BODIES OF LAVERAN [after Golgi]

always curved; some, indeed, are almost spherical, and others sausage-shaped. They are motionless. In many specimens a delicate line is visible on the concave side of the crescent, connecting the extremities. On careful examination this is found to be the edge of a very delicate membrane. The body is composed of homogeneous protoplasm. Centrally placed is a collection of pigment granules, which on careful examination can be distinctly seen to be in movement. The semi lunar bodies vary in number in different cases. Sometimes several can be seen in the field at the same time, and in other cases they are only observed after a long and patient search. They are, as a rule, free in the serum; but they have also been seen within the red blood-cells. *Second*, finely granular masses of protoplasm,

which arise, according to Golgi, from the intra-corpuscular pigmented bodies. The pigment is collected in a rosette, and the protoplasm by segmentation gives rise to a number of small spherical forms, which are




FIG. 140.—ROSETTE FORMS WITH SEGMENTATION [after Golgi].

ultimately set free (Fig. 140). Golgi believes that these changes occur in definite relation to the development of the paroxysm. *Third*, spherical, pear-shaped, or ovoid bodies, rather smaller than the red blood-corpuscles, and provided with one or more actively motile flagella




FIG. 141.—FLAGELLATED FORMS [after Vandyke Carter].

1. A flagellated spherule; (a) the same in the interior of a phagocyte; (b) free motile filaments.

(Fig. 141). These flagella are long lash-like filaments, which by their activity set the neighbouring blood-corpuscles in motion. Free filaments in active movement have also been observed. *Fourth*, small spherical pigmented bodies about one quarter the size of a red blood-corpuscle, which exhibit amœboid movement.

Inoculation experiments. — Marchiafava and Celli assert that inoculation of a healthy subject with blood containing the parasites will produce a paroxysm of ague with development of the hæmatozoa. The pathogenic power of these parasites, however, has not been established. As in *Surra* disease, there has been no cultivation of the parasite outside the animal body, and reproduction of the disease with a pure cultivation. In favour of its being a pathogenic organism, Laveran points out its invariable presence in some form or other in cases of malaria; the marked changes it effects in the red blood-cells; the increase in the number of the parasites in proportion to the severity of the attack; and, lastly, their disappearance after the administration of quinine. Others, again, have doubted the parasitic nature of these bodies, and have looked upon them as representing pathological changes in the blood-cells.*

Laveran first of all suggested the name *Oscillaria malariae*, but subsequently he recognised that these bodies belonged to the animal, not to the vegetable kingdom. Osler has suggested that, temporarily at any rate, the organism should be placed in the genus *Hæmatomonas* of Mitrophanow, thus: "Genus, *Hæmatomonas*; species, *Hæmatomonas malariae*. Definition—Body plastic; ovoid or globose; no differentiation of protoplasm, which contains pigment grains; flagella variable from one to four, highly polymorphic, occurring in (1) amœboid form, (2) crescents, encysted form, (3) sporocysts, (4) cellular free pigmented bodies."

EXAMINATION OF THE HÆMATOZOA OF LAVERAN.

In the living condition.—Select a patient by preference who has had several attacks of malaria, and is markedly anæmic. Examine before the invasion of the febrile paroxysm. Take two perfectly clean cover-glasses

* Cattaneo and Monti, *Archivio per le Scienze Mediche*. 1888.

and two clean slides. Wash one of the fingers of the patient with soap and water, and then cleanse with alcohol. Apply a ligature, and with a clean needle puncture the thin skin near the root of the nail. Touch the drop of blood which collects with a clean slide. Cover quickly with a cover-glass, and gently press it if the layer of blood be too thick. Examine with a $\frac{1}{12}$ o.i.

In stained preparations.—Puncture the finger again if necessary, touch the droplet of blood with a clean cover-glass, apply another cover-glass, press them gently together, and then slide them apart. Stain with two or three drops of alcoholic solution of methylene-blue, wash off excess and examine in water, or allow the preparation to dry, and mount in balsam.

APPENDIX E.

CHRONOLOGICAL BIBLIOGRAPHY.

- A. METHODS.
- B. MORPHOLOGY AND CLASSIFICATION.
- C. GENERAL BIOLOGY.
- D. ZYMOGENIC SAPROPHYTES AND FERMENTATION.
- E. CHROMOGENIC SAPROPHYTES.
- F. SIMPLE SAPROPHYTES.
- G. PTOMAINES AND PUTREFACTION.
- H. ANTISEPTICS AND DISINFECTANTS.
- I. IMMUNITY.
- J. BACTERIA ASSOCIATED WITH DISEASES IN MAN AND ANIMALS :—
- | | |
|-------------------------------|------------------------------|
| I. ACTINOMYCOSIS. | XIX. OPHTHALMIC DISEASES. |
| II. ACUTE YELLOW ATROPHY. | XX. OSTEO-MYELITIS. |
| III. ANTHRAX. | XXI. PLEURO-PNEUMONIA. |
| IV. CATTLE PLAGUE. | XXII. PNEUMONIA. |
| V. CEREBRO-SPINAL MENINGITIS. | XXIII. PUERPERAL FEVER. |
| VI. CHICKEN-CHOLERA. | XXIV. PYÆMIA AND SEPTICÆMIA. |
| VII. CHOLERA. | XXV. RELAPSING FEVER. |
| VIII. DENTAL CARIES. | XXVI. RHINOSCLEROMA. |
| IX. DIPHTHERIA. | XXVII. SCARLATINA. |
| X. ERYSIPELAS. | XXVIII. SWINE-ERYSIPELAS. |
| XI. ENDOCARDITIS. | XXIX. SWINE-TYPHOID. |
| XII. GLANDERS. | XXX. SYMPTOMATIC ANTHRAX. |
| XIII. GONORRHOEA. | XXXI. SYPHILIS. |
| XIV. HYDROPHOBIA. | XXXII. TETANUS. |
| XV. LEPROSY. | XXXIII. TUBERCULOSIS. |
| XVI. MALARIA. | XXXIV. TYPHOID FEVER. |
| XVII. MALIGNANT ŒDEMA. | XXXV. VARIOLA AND VACCINIA. |
| XVIII. MEASLES. | XXXVI. YELLOW FEVER. |
- K. BACTERIA IN THE AIR, IN SOIL, AND IN WATER.

(A) *METHODS.*

- 1852 Perty.—Zur Kenntniss Kleinster Lebensform.
- 1867 Pasteur.—Etudes sur la Bière.
- 1873 Klebs.—Ueber Fractionirte Cultur. Archiv f. Exp. Pathol., Bd. I.
- 1875 Brefeld.—Verhandl. d. Physik. Med. Ges. in Würzburg.
Cohn.—Beiträge zur Biologie der Pflanzen, Bd. I., Heft 3.
- 1876 Cohn.—Beiträge zur Biologie der Pflanzen, Bd. II., Heft 2.
- 1877 Schäfer.—Course of Practical Histology.
Koch.—Verfahren z. Untersuch., z. Conservir. und Photographir. der Bakterien. Beiträge z. Biol. d. Pflanzen, Bd. II., Heft 3.
- 1878 Koch.—Untersuchungen über Wundinfections Krankheiten.
- 1879 Salomonsen.—Zur Isolation Differenter Bakterien. Bot. Zeitung, No. 39.
Blanchard.—Rev. Inter. Sci., III.
- 1880 Salomonsen.—Eine Einfache Meth. z. Reincultur Versch. Fäulniss Bakterien. Bot. Zeit., No. 28.
Ehrlich.—Zeitschr. f. Klin. Med., Bd. I.
- 1881 Ehrlich.—Zeitschr. f. Klin. Med., Bd. II., Heft 3.
Carpenter.—The Microscope and its Revelations. 6th Edition.
Koch.—Zur Untersuchung von Pathogenen Organismen. Mitth. aus d. K. Gesundheits Amt, Bd. I.
Brefeld.—Bot. Untersuch. über Schimmelpilze, Bd. IV.
Duclaux.—Ferments et Maladies.
- 1882 Koch.—Biol. Klin. Woch., No. 15.
Ehrlich.—Deut. Med. Woch., No. 19.
Fehleisen.—Ueber Neue Method. der Untersuch. u. Cultur Pathogen. Bakterien. Physik. Med. Ges. zu Würzburg.
Büchner.—In Nägeli's Untersuch. über Niedere Pilze. Munich.
- 1883 Almquist.—Hygeia, XLV. Stockholm.
Behrens.—Hilfsbuch zur Ausführung Mikroskop. Untersuch.
Hazlewood.—American Monthly Microscop. Journ.
Brefeld.—Die Künstl. Kultur Parasitischer Pilze. Botan. Unters. über Hefenpilze, Bd. V.
Flügge.—Handbuch der Hygiene und der Gewerbe Krankheiten.
- 1884 Magnin and Sternberg.—Bacteria.
Friedländer.—Microscopische Technik. 1st Edition.
Büchner.—Ueber das Verhalten der Spaltpilzsporen zu den Anilinfarben. Aerztl. Intelligenzbl., No. 33.
Orth.—Patholog. Anatom. Diagnostik.
Plaut.—Färbungs Methode z. Nachweis. der Micro-organismen. 1st Ed.
Baumgarten.—Beitr. zur Darstellungsmethode d. Tuberkel Bacillen. Zeitschr. f. Wissensch. Mikr.
Koch.—Mitth. a. d. Kais. Ges. Amt, Bd. II.
Gram.—Ueber die Isolirte Färbung der Schizomyceten. Fortsch. d. Med., II., No. 6.

- 1885 **Hueppe**.—Die Methoden der Bakterien Forschung.
Cornil and Babès.—Les Bactéries.
Gibbes.—Practical Histology and Pathology.
Hauser.—Ueber Fäulnisbakterien. Mit 15 Tafeln in Lichtdruck.
Klein.—Micro-organisms and Disease.
Lee.—The Microtomist's Vade Mecum.
Malley.—Photomicrography.
Plaut.—Färbungs Methoden. 2nd Edition.
Woodhead and Hare.—Pathological Mycology.
Bizzozzo and Firket.—Manuel de Microscopie Clinique.
Johne.—Ueber die Kochschen Reinculturen.
Bordoni Uffreduzzi.—Microparasitici.
Banti.—Manuale di Tecnica Batteriologica.
Dolley.—Technology of Bacteria Investigation.
- 1886 **Hueppe**.—Die Methoden der Bakterien Forschung. 3rd Edition.
Hueppe.—The Methods of Bacteriological Investigation. Translated by Biggs.
Hüber and Breker.—Die Path. Histolog. und Bacteriologischen Untersuch. Methoden.
Klein.—Micro-organisms and Disease. 3rd Edition.
Cornil and Babès.—Les Bactéries. 2nd Edition.
Crookshank.—Manuel Pratique de Bactériologie, traduit par Bergeaud. Avec 4 Photomicrographies.
Plaut.—Fortschr. d. Med.
Eisenberg.—Bakteriologische Diagnostik.
Esmarch.—Zeitschrift f. Hygiene.
Hueppe.—Bakteriologische Apparate. Deut. Med. Woch.

(B) *MORPHOLOGY AND CLASSIFICATION.*

- 1838 **Ehrenberg**.—Die Infusionsthierchen als Volkom. Organism.
1841 **Dujardin**.—Histoire Naturelle des Zoophytes.
1866 **De Bary**.—Morph. und Phys. der Pilze.
Hallier.—Die Pflanzlichen Parasiten.
1872 **Cohn**.—Beiträge zur Biologie der Pflanzen, Bd. I.
1873 **Lister**.—Quart. Journ. Microscop. Science.
Lankester.—Quart. Journ. Microscop. Science.
1874 **Billroth**.—Unters. über d. Veg. Form der Coccobacteria septica.
1875 **Cohn**.—Beiträge z. Biol. d. Pflanzen.
Dallinger and Drysdale.—Monthly Microscop. Journ.
1877 **Leunis**.—Synopsis d. Pflanzenkunde. Hanover.
Nägeli.—Die Niederen Pilze.
1879 **van Tieghem**.—Compt. Rend.
1881 **Brefeld**.—Botanische Untersuch. über Schimmelpilze, Heft 1.
Koch.—Mitheil aus d. K. Ges. Amt, Bd. I.
Rabenhorst.—Kryptogamenflora. Bd. I., Pilze, von Winter.

- 1882 **Sachs**.—Text-Book of Botany.
- 1883 **Flügge**.—Handbuch der Hygiene.
Neelsen.—Neuere Ansichten über d. Systematik der Spaltpilze. Biol. Centralbl., III., No. 18.
van Tieghem.—Traité de Botanique.
- 1884 **De Bary**.—Verg. Morph. und Biolog. der Pilze, Mycetozen, und Bacterien.
Grove.—Synopsis of the Bacteria and Yeast Fungi.
Marpmann.—Die Spaltpilze.
Miller.—Deut. Med. Woch.
- 1885 **Cornil and Babès**.—Les Bactéries.
Fisch.—Die Systemat. Stellung der Bacterien. Biolog. Centralbl., V.
Zopf.—Die Spaltpilze.
A. de Bary.—Vorlesungen über Bacterien.
Biedert.—Virch. Archiv, Bd. 100.
Hauser.—Ueber Fäulniss Bacterien.
- 1886 **Hueppe**.—Die Formen der Bakterien.
Baumgarten.—Lehrbuch der Pathologischen Mykologie.
Flügge.—Die Micro-organismen. 2nd Edition.
Lutz.—Fortschr. d. Med.

(C) *GENERAL BIOLOGY.*

- 1877 **Nägeli**.—Die Niederen Pilze. München.
- 1879 **Cohn and Mendelsohn**.—Ueber Einwirkung des Elektrischen Stromes auf die Vermehrung d. Bakterien. Beitr. z. Biol. d. Pflanzen, Bd. III., Heft 1.
Nencki.—Virchow's Archiv.
- 1880 **Nencki**.—Beiträge zur Biol. der Spaltpilze. Leipzig.
- 1881 **Tumas**.—St. Petersb. Med. Wochenschr.
Engelmann.—Arch. f. d. Ges. Physiologie, Bd. 26.
- 1882 **Engelmann**.—Botan. Zeitg.
Nägeli.—Unters. über Niedere Pilze. München.
- 1884 **Regnard**.—Rech. sur l'Influence de très-hautes Pressions sur les Org. Vivants. Compt. Rend., T. 98, p. 744.
Cortes.—De l'Action des Hautes Pressions sur les Phénomènes de la Putréfaction, etc. Compt. Rend., T. 99, p. 385.
Nencki.—Ueber das Eiweiss der Milzbrandbacillen. Ber. d. Deutschen Chem. Gesellsch. S. 2605.
- 1885 **Duclaux**.—Influence de la Lumière sur la Vitalité des Germes de Microbes. Compt. Rend.
Mittenzweig.—Die Bakt. Aetiolog. der Infect. K.
- 1886 **v. Fodor**.—Bakterien im Blute lebender Thiere. Archiv f. Hygiene.
Hauser.—Archiv f. Exper. Patholog. u. Pharmacologie.
Wysokowitsch.—Zeitschrift. f. Hygiene.
Pavy.—Relation of Bacteria to Disease. Harveian Oration. Brit. Med. Journ.

- 1886 **Bordoni-Uffreduzzi.**—Fortschr. d. Med.
Arloing.—Archiv de Physiol.
Watson-Cheyne.—Brit. Med. Journ.
Liborius.—Zeitschrift. f. Hygiene.
Downes.—Action of Sunlight on Micro-org. Proc. Roy. Soc.
- (D) *ZYMOGENIC SAPROPHYTES AND FERMENTATION.*
- 1860 **Müller.**—Journ. f. Prakt. Chem.
Pasteur.—Annal. de Chim. et de Phys., III. Sér., T. 58.
Pasteur.—Compt. Rend., T. 50.
- 1861 **Liebig.**—Verhandl. der Münchener Akad. d. Wiss.
Pasteur.—Bull. de la Soc. Chim.
Pasteur.—Compt. Rend., T. 52.
- 1862 **Pasteur.**—Ann. de Chim. et de Phys., T. 64.
Monoyer.—Thèse de Strassburg.
- 1863 **Pasteur.**—Compt. Rend., T. 56.
- 1864 **Pasteur.**—Compt. Rend.
van Tieghem.—Compt. Rend., T. 58, p. 210.
Tyndall.—Compt. Rend., T. 58.
- 1865 **Duclaux.**—Thèses présentées à la Faculté de Paris.
Béchamp.—Compt. Rend., T. 60, p. 445.
Trécul.—Ueber Bacillus Amylobacter. Compt. Rend., T. 61.
- 1866 **Pasteur.**—Etudes sur le Vin.
- 1867 **Hallier.**—Gährungserscheinungen.
Trécul.—Compt. Rend., T. 65.
Trécul.—Ann. des Sc., Sér. 7, T. 7.
- 1869 **Karsten.**—Chemismus der Pflanzenzelle.
Liebig.—Verhandl. der Münchener Akad. d. Wiss., 5 Nov.
- 1870 **Liebig.**—Ueber Gährung, Quelle der Muskelkraft und Ernährung.
 Leipzig u. Heidelberg.
- 1871 **Pasteur.**—Compt. Rend.
Dubrunfaut.—Compt. Rend., T. 73.
Hoppe-Seyler.—Medic.-Chem. Untersuchungen, Heft 4.
- 1872 **Pasteur.**—Compt. Rend.
Dumas.—Compt. Rend., T. 75, Nr. 6.
Lex.—Centralbl. f. d. Med. Wiss., S. 291.
- 1873 **Fitz.**—Berichte d. Chem. Ges., Bd. 6, S. 48.
Hofmann.—Aerztl. Verein zu Wien, Mai.
Hofmann.—Allgem. Med. Centralbl., S. 605.
- 1874 **Schützenberger.**—Die Gährungserscheinungen.
Dumas.—Ann. de Chim. et de Phys.
Brefeld.—Landwirthsch. Jahresber., Bd. 3.
Scheibler.—Ueber die Natur des Froschlaich. Zeitschr. f. Rübenzuckerindustrie.

- 1874 Feltz and Bitter.—Journ. de l'Anat. et Phys.
 Hiller.—Centralbl. f. d. Med. Wiss., S. 53.
 van Tieghem.—Compt. Rend., T. 58.
 Musculus.—Ber. d. Chem. Ges., S. 124.
 Musculus.—Compt. Rend., T. 78.
- 1875 Popoff.—Botan. Jahresber.
 Brefeld.—Landwirthsch. Jahresber., Bd. 4.
 Colin.—Bull. de l'Acad. de Méd.
- 1876 Pasteur.—Bull. de l'Acad. de Méd., No. 27.
 Pasteur and Joubert.—Compt. Rend., T. 83.
 Pasteur.—Etudes sur la Bière.
 A. Mayer.—Lehrbuch der Gährungschemie. 2 Aufl.
 Fleck.—Ber. d. Chem. Centralst. Dresden.
 Brefeld.—Landwirthsch. Jahresber., Bd. 5.
- 1877 Lister.—The Cause of Putrefaction and Lactic Fermentation. The
 Pharmac. Journ. and Transact.
 Harz.—Grundzüge der alkoholischen Gährungslehre.
- 1878 Cienkowski.—Die Gallertbildungen d. Zuckerrübensaftes.
 Boutroux.—Sur la Fermentation Lactique. Compt. Rend., T. 86.
 Fitz.—Berichte d. Chem. Ges., Bd. 10, p. 216.
- 1879 Fitz.—Berichte d. Chem. Ges., Bd. 11, pp. 42 and 498; and Bd. 12,
 p. 474.
 Richet.—Compt. Rend., T. 88.
 Nägeli.—Theorie der Gährung. München.
 van Tieghem.—Développement du Spirillum Amyliférum. Bull. de la
 Soc. Bot. de France, and Sur la Fermentation de la Cellulose,
 pp. 25—30.
 van Tieghem.—Compt. Rend., T. 88.
 van Tieghem.—Identité du Bacillus Amylobacter et du Vibrion
 Butyrique de Pasteur. Compt. Rend., T. 89.
- 1880 Fitz.—Berichte d. Chem. Ges., Vol. 13, p. 1309.
 Prazmowski.—Untersuchungen über die Entwicklungsgeschichte und
 Fermentwirkung einiger Bakterien.
 van Tieghem.—Compt. Rend.
- 1881 Béchamp.—Compt. Rend., T. 93.
 Tyndall.—Essays on the Floating Matter of the Air.
 Eriksson.—Unters. aus d. Botan. Institut in Tübingen, Heft 1.
 Kern.—Ueber ein neues Milchferment aus dem Kaukasus. Bull. de la
 Soc. Imp. des Naturalistes de Moskau, No. 3.
 v. Jacksch.—Studien über den Harnstoffpilz. Zeitschr. f. Physiol.
 Chemie, Bd. 5.
- 1882 Fitz.—Berichte d. Chem. Ges., Bd. 15, p. 857.
- 1883 Guiard.—Thèse de Paris.
 Tappeiner.—Celluloseverdauung. Fortschr. d. Med., I., 151; II., 377,
 416.
 Fitz.—Berichte d. Chem. Ges., Bd. 16, p. 844.

- 1884 **Hueppe**.—Unters. üb. die Zersetzungen der Milch durch Mikroorganismen. Mitth. a. d. Ges. Amt, Bd. II.
Hueppe.—Deut. Med. Woch.
Krannhals.—Ueber das Kumysähnliche Getränk Kephir. Deut. Arch. f. Klin. Med., Bd. 35.
Fitz.—Berichte d. Chem. Ges., Bd. 17, p. 1188.
Ladureau.—Sur le Ferment Ammoniacal. Compt. Rend., T. 99, p. 877.
van Tieghem.—Développement de l'Amylobacter dans les Plantes à l'Etat de Vie Normale. Compt. Rend.
1885 **Lépine and Roux**.—Compt. Rend., T. 101.
Leube.—Ueber die Ammoniakalische Harnsäure. Virch. Arch., Bd. 100, S. 540.
Billet.—Sur le Bacterium Ureae. Ib., Bd. 100, p. 1252.
Sheridan Lea.—Journ. of Physiology.

(E) *CHROMOGENIC SAPROPHYTES.*

- 1833 **Hermstädt**.—Ueber die Blaue und Rothe Milch.
1838 **Steinhoff**.—Ueber das Blauwerden der Milch. Neue Ann. d. Mecklenb. Landw. Ges.
1839 **Ehrenberg**.—Micr. Prodigiosus. Verhandl. d. Berl. Acad.
1850 **Chabert and Fromage**.—D'une Altération du Lait de Vache, désignée sous le Nom du Lait Bleu.
1852 **Gielen**.—Mag. f. Ges. d. Thierheilkunde.
1860 **Fordos**.—Compt. Rend. de l'Acad. de Sc.
1862 **Lücke**.—Arch. f. Klin. Chir.
1863 **Eberth**.—Centralbl. f. d. Med. Wissensch.
1868 **Mosler**.—Ueber Blaue Milch und durch deren Genuss herbeigeführte Krankheiten. Virchow's Archiv, Bd. 43.
1872 **Schröter**.—Ueber einige von Bakterien gebildete Pigmente. Cohn, Beitr. z. Biol. der Pflanzen, Bd. I., Heft 2.
1873 **Lankester**.—On a Peach-coloured Bacterium. Quart. Journ. of Micr. Sc., Vol. 13.
1875 **Frank**.—Cohn's Beitr. z. Biol. d. Pflanzen, Bd. I., Heft 3.
Girard.—Unters. über Blauen Eiter. Chirurg. Centralbl., II.
Klein.—Quart. Journ. of Micr. Sc., Vol. 15.
1876 **Lankester**.—Further Observations on a Peach- or Red-coloured Bacterium. Quart. Journ. of Micr. Sc.
1877 **Giard**.—Revue des Sc., T. 5.
1879 **Wernich**.—Cohn's Beiträge zur Biol. d. Pflanzen, Bd. III., Heft 1.
Cohn and Miflet.—Cohn's Beiträge zur Biol. d. Pflanzen, Bd. III., Heft 1.
1880 **van Tieghem**.—Bull. de la Soc. Bot. de France.
Neelsen.—Cohn's Beitr. z. Biologie d. Pflanzen, Bd. III., Heft 2.
1882 **Babès**.—Vom Rothen Schweiss. Biolog. Centralbl., Bd. 2.
Gessard.—De la Pyocyanine et de son Microbe.
1884 **Charrin**.—Communication faite à la Société Anatomique.
Hugues.—Le Lait Bleu. Echo Vétérinaire.

(F) *SIMPLE SAPROPHYTES.*

- 1833 **Kützing**.—*Sphaerotilus Natans*. Linnæa, 8.
 1840 **Oersted**.—*Naturhist. Tidsskrift*, Bd. III.
 1842 **Goodsir**.—*Edinb. Med. and Surg. Journ.*
 1845 **Weisse**.—*Monas Okenii*. Bull. Phys.-Mathemat. de St. Pétersbourg III.
 1847 **Heller**.—*Heller's Arch. f. Chemie.*
 1858 **Eberth**.—*Virchow's Archiv*, Bd. 13.
 Itzigsohn.—*Virchow's Archiv*, Bd. 13.
 1859 **Welcker**.—*Zeitschr. f. Ration. Med.*
 1861 **Munk**.—*Virch. Arch.*, Bd. 22.
 1862 **Pasteur**.—*Ann. de Chim. et de Phys.*, T. 64.
 1864 **Friedreich**.—*Beiträge zur Kenntniss der Sputa*. Virch. Arch., Bd. 30.
 Munk.—*Med. Centralbl.*
 1865 **Cohn**.—*Ueber zwei Neue Beggiatoen*. Hedwigia.
 1866 **Suringar**.—*Arch. Neerland.*
 Suringar.—*Ueber den Zellenbau der Sarcina*. Bot. Zeit.
 1867 **Lüders**.—*Ueber Abstammung u. Entwicklung des Bacterium Termo.*
 1871 **Losdorfer**.—*Med. Jahrb.*, Heft 3.
 1872 **Cohn**.—*Beitr. z. Biol. d. Pflanzen*, Bd. I.
 1873 **Lankester**.—*Quart. Journ. of Microscop. Sc.*
 1874 **Caspary**.—*Schriften der Physik. Oekon. Ges. zu Königsberg*, Bd. 15.
 Ewart.—*Proceedings of the Roy. Soc.*
 1875 **Eidam**.—*Cohn's Beitr. zur. Biol. d. Pflanzen.*
 Dallinger and Drysdale.—*Monthly Microsc. Journ.*
 1876 **Lankester**.—*Quarterly Journ. of Microscop. Sc.*
 1877 **Heimer**.—*Ueber Pneumonomycosis Sarcinica*. Deut. Arch. f. Klin. Med.
 1878 **Dallinger**.—*Journ. of the Roy. Micro. Soc. London.*
 Geddes and Ewart.—*On the Life-History of Spirillum*. Proceed. of
 the Roy. Soc. of London.
 Brefeld.—*Bacillus Subtilis*. Ges. Nat. Freunde.
 1879 **Zopf**.—*Ueber Crenothrix Polyspora*. Berlin.
 1882 **Zopf**.—*Bacterium Merismopedioïdes*. Sitz.-Ber. d. Botan. Ver. d.,
 Prov. Brandenb. Juni.
 Engelmann.—*Bacterium Photometricum*. Unters. aus. d. Phys.
 Laborat. zu Utrecht. Pflüger's Arch. f. d. Ges. Physiol., Bd. 30
 S. 95.
 Giard.—*Sur le Crenothrix Kühniana*. Compt. Rend.
 1883 **Kurth**.—*Bacterium Zopfii*. Ber. d. Deutsch. Bot. Ges. Botanische
 Zeitg.
 1884 **Vandevelde**.—*Studien zur Chemie des Bacillus Subtilis*. Zeitschr. f.
 Phys. Chemie, Bd. 8.
 Mülhåuser.—*Ueber Spirillen*. Virch. Arch., Bd. 97.
 1885 **Falkenheim**.—*Arch. f. Exp. Patholog. u. Pharmakol.*, Bd. 19.
 Billet.—*Sur la Formation des Spores chez le Cladotrix Dichotoma*.
 Compt. Rend., T. 100.

- 1886 **Bordoni-Uffreduzzi**.—Ueber d. biol. Eigensch. d. norm. Hautmikrophyt. Fortsch. d. Med.

(G) *PTOMAINES AND PUTREFACTION.*

- 1866 **Hemmer**.—Exper. Studien über d. Wirkung Faulender Stoffe.
Schweninger.—Ueber d. Wirkung Faulender Org. Substanzen.
v. Raison.—Zur Kenntniss der Putriden Intoxication.
Bergmann.—Das Putride Gift.
- 1868 **Bergmann and Schmiedeberg**.—Med. Centralbl.
- 1869 **Zülzer and Sonnenschein**.—Berl. Klin. Woch.
- 1872 **Bergmann**.—Deut. Zeitschr. f. Chirurgie, Bd. I.
Ravitsch.—Zur Lehre von der Putriden Infection.
- 1873 **Clementi and Thin**.—Unters. üb. d. Putride Infection. Wien. Med. Jahrb.
- 1874 **Panum**.—Virchow's Arch., Bd. 60.
Frisch.—Exper. Studien üb. d. Verbreitung d. Fäulnisorganismen.
Kehrer.—Archiv f. Exper. Pathol., Bd. I.
- 1876 **Hiller**.—Centralbl. f. Chirurgie.
Nencki.—Ueber die Zersetzung der Gelatine und des Eiweisses bei der Fäulnis mitt Pancreas.
- 1877 **Salomonsen**.—Die Fäulnis des Blutes.
- 1878 **Selmi**.—Chemische Ber., Bd. 6, 7, 12, Sulle Ptomaïne ad Alcaloide Cadaverici.
Kaufmann.—Zersetzung des Blutes durch Bacillus Subtilis. Journ. f. Prakt. Chemie, Bd. 17.
- 1879 **Hiller**.—Die Lehre von der Fäulnis.
- 1880 **Husemann**.—Arch. d. Pharmac.
- 1881 **König**.—Massenerkrankung von Menschen nach dem Genuss von Fleisch einer an Putrider Metritis Verendeten Kuh. Ber. üb d. Veterinärwesen im Königreich Sachsen.
Tanret.—Compt. Rend., T. 92.
Brouardel and Boutmy.—Compt. Rend., T. 92, p. 1056.
- 1882 **Husemann**.—Arch. d. Pharmac.
Schiffer.—Arch. f. Anat. u. Physiol., Physiol. Abtheil.
Bocci.—Centralbl. f. d. Med. Wiss.
Groebner.—Beiträge z. Kenntniss der Ptomaïne.
Bergmann and Angerer.—Das Verhältniss der Fermentintoxication zur Septicämie. Würzburger Jubil. Festschr.
Bouchard.—Compt. Rend. de Biol.
Nencki.—Journ. f. Pract. Chem., Bd. 26.
Gautier.—Compt. Rend., T. 94.
Gautier and Etard.—Compt. Rend., T. 94.
Rosenbach.—Giebt es Verschiedene Arten von Fäulnis? Deut. Zeitschrift für Chir., XVI., S. 342.
Etard and Olivier.—De la Réduction des Sulfates par les Etres Vivants. Compt. Rend.

- 1883 **Husemann.**—Arch. d. Pharmac.
Guareschi and Mosso.—Arch. Ital. de Biolog.
Brieger.—Zur Kenntniss der Fäulnissalkaloide. Zeitschr. f. physiol. Chemie, Bd. 7.
- 1884 **Maas.**—Ueber Fäulnissalkaloide des gekochten Fleisches und des Fischfleisches. Fortschr. d. Med., 11., 729.
Brieger.—Ber. d. Deutsch. Chem. Ges., Bd. 17.
Vandevelde.—Les Ptomaïnes. Arch. de Biol. par van Beneden.
Willgerodt.—Ueber Ptomaïne.
Brieger.—Ueber Giftige Producte der Fäulnissbakterien. Berl. Klin. Woch., Nr. 14.
Otto.—Anleitung zur Ausmittlung der Gifte.
- 1885 **Tappeiner.**—Med. Centralbl.
Blumberg.—Experimenteller Beitrag zur Kenntniss der Putriden Intoxication. Virch. Arch., Bd. 100, S. 377.
Offinger.—Die Ptomaïne.
Brieger.—Ueber Ptomaïne.
Brieger.—Weitere Untersuchungen über Ptomaïne.
Backlisch.—Ber. d. Deutsch. Chem. Gesellsch., Bd. 18.
Hauser.—Ueber Fäulnissbakterien.
- 1886 **Brieger.**—Ueber Ptomaïne. Berl. Klin. Woch.

(H) *ANTISEPTICS AND DISINFECTANTS.*

- 1875 **Eidam.**—Einwirkung Versch. Temperaturen a. d. Entwicklung von Bacterium Termo. Cohn's Beitr. zur. Biol., Bd. I., Heft 3.
Schröter.—Cohn's Beitr. zur. Biol. der Pflanzen, Bd. I., Heft 3.
- 1876 **Koch.**—Cohn's Beitr. zur. Biol. der Pflanzen, Bd. 11., Heft 2.
Buchholz.—Ueber das Verhalten von Bakterien zu einigen Antiseptics.
- 1877 **Nägeli.**—Die Niederen Pilze.
Buchholz.—Arch. f. Exp. Pathol., Bd. 7.
Frisch.—Ueber den Einfluss nied. Temp. auf. d. Lebensfähigkeit der Bakterien. Sitzungsber. d. Wiener Akad., Bd. 75 u. 80.
Tyndall.—Philos. Transact. of the Roy. Soc.
Vallin.—Ann. d'Hyg.
- 1879 **Soyka.**—Ber. d. Bayr. Akad. d. Wissensch.
Schwartz.—Sitzungsber. d. Dorpater Naturf. Ges.
Wernicke.—Diss. Dorpat.
Haberkorn.—Das Verhalten von Harnbakterien gegen einige Antiseptica. Dissert. Dorpat.
Wernich.—Die Aromatischen Fäulnissproducte in ihrer Einwirkung auf Spalt und Sprosspilze. Virchow's Archiv, Bd. 78
- 1880 **Merke.**—Virchow's Archiv, Bd. 81.
Mörschell.—Deut. Med. Woch.
Lassar.—Deut. Med. Woch.
Pasteur.—Ann. d'Hyg.

- 1880 **Wernich.**—Grundriss der Desinfectionslehre.
Meyer.—Ueb. d. Milchsäureferment u. sein Verhalten gegen Antiseptica.
Toussaint.—Bull. de l'Acad.
- 1881 **Wolffhügel.**—Ueber den Werth der schwefligen Säure als Desinfectionsmittel. Mittheilg. a. d. Kais. Ges. Amt, Bd. I., S. 188.
Wolffhügel and **Knorre.**—Zu der verschiedenen Wirksamkeit von Carbolöl u. Carbolwasser. Mitth. a. d. Kaiserl. Ges. Amt, Bd. I., S. 352.
Koch.—Ueber Desinfection. Mitth. a. d. Ges. Amt, Bd. I., S. 234.
De la Croix.—Das Verhalten der Bakterien d. Fleischwassers gegen einige Antiseptica. Arch. f. Exp. Pathol., Bd. 13.
Koch, Gaffky, and **Löffler.**—Versuche über die Verwerthbarkeit heisser Wasserdämpfe zu Desinfectionszwecken. Mitth. a. d. Kaiserl. Ges. Amt, Bd. I., S. 322.
Koch and **Wolffhügel.**—Unters. üb. die Desinfection mit heisser Luft. Mitth. a. d. Kaiserl. Ges. Amt, Bd. I., S. 301.
Hueppe.—Ueber das Verhalten ungeformter Fermente gegen hohe Temperatur. Mittheilg. a. d. Kaiserl. Ges. Amt, Bd. I., S. 341.
- 1882 **Lebedeff.**—Desinfectionsversuche am Malignen Oedem. Arch. de Physiol. Norm. et Pathol.
W. Cheyne.—Antiseptic Surgery.
Lebedeff.—Arch. de Phys. Norm. et Pathol.
Hueppe.—Ueber die Hitze als Desinfectionsmittel. Deut. Militärärztl. Zeitschr.
- 1883 **Fischer.**—Unters. üb. d. Wirkung des Naphtalins. Berl. Klin. Woch.
Perroncito.—Sur la Tenacité de Vie du Virus Charbonneux. Arch. Ital. de Biol.
Vallin.—Traité des Désinfectants et de la Désinfection.
Miquel.—Antiseptiques et Bactéries. Semaine Médicale.
Chauveau.—Compt. Rend.
Chamberland and **Roux.**—Compt. Rend.
Laillier.—Du Gaz Acide Sulfureux. Ann. d'Hygiène.
Arloing, Cornevin, and **Thomas.**—Lyon Méd.
Vallin.—Les Nouvelles Etuves à Désinfection. Revue d'Hygiène.
Larrivé.—L'Eau Oxygénée. Thèse de Paris.
Maly and **Emich.**—Antisept. Wirkung der Gallensäuren. Sitzungsber. d. Kais. Akad. d. Wiss. zu Wien. Jan.
Schultz.—Die Antiseptischen Eigenschaften der Citronensäure. Deut. Med. Woch., Nr. 17.
- 1884 **Schill** and **Fischer.**—Ueber die Desinfection des Auswurfs der Phthisiker. Mitth. a. d. Kaiserl. Ges. Amt, Bd. II.
Plaut.—Desinfection der Viehställe.
Steinmeyer.—Ueber Desinfectionslehre.
Colin.—Compt. Rend., T. 99.
Chairy.—Action des Agents Chimiques sur les Bactéries du Genre Tyrothrix. Compt. Rend.

- 1884 **Chauveau**.—Compt. Rend.
Mignet.—Annuaire de l'Observatoire de Montsouris.
Schnetzler.—Les Propriétés Antiseptiques de l'Acide Formique. Archiv de Genève.
Rosbach.—Einfluss des innerl. Naphthalingebruchs auf die Harnfäulniss. Berl. Klin. Woch.
Hoffmann.—Experimentelle Untersuchungen über die Wirkung der Ameisensäure Diss. Greifswald.
Fischer and Proskauer.—Mitth. a. d. Kaiserl. Ges. Amt, Bd. II.
Heydenreich.—Sur la Stérilisation des Liquides au moyen de la Marmite de Papin. Compt. Rend., T. 98.
Pictet and Young.—De l'Action du Froid sur les Microbes. Compt. Rend., T. 98.
Vallin.—Ann. d'Hygiène.
Rocheftort, Herscher.—Revue d'Hygiène.
Dujardin-Beaumetz.—Bull. de l'Acad. de Méd. de Paris.
Marié-Davy.—Revue d'Hygiène.
- 1885 **M. Wolff**.—Zur Desinfectionsfrage. Centralbl. f. d. Med. Wiss., Nr. 11.
Reinl.—Zur Theorie der Heilwirkung des Franzensbader Moores. Prager Med. Woch., Nr. 10 u. 11.
Thol.—Ueber d. Einfluss nicht aromat. organ. Säuren. auf Fäulniss u. Gährung. Diss. Greifswald.
Schulz.—Die Ameisensäure als Antisepticum. Deut. Med. Woch., Nr. 24.
Schede.—Die Antiseptische Wundbehandlung mit Sublimat. Sammlung Klin. Vorträge, Nr. 25.
König.—Sublimatdämpfe. Chirurg. Centralbl.
Frank.—Ueber Desinfection von Abtrittsgruben.
Gärtner and Plagge.—Deut. Med. Woch.
- 1886 **Handford**.—Brit. Med. Journ.
Koch and Gaffky.—Arbeit. a. d. K. Gesundh. Amt.
Winter Blyth.—Studies of Disinfectants by New Methods. Proc. Roy. Soc.

(1) *IMMUNITY.*

- 1876 **Oemler**.—Arch. f. Wiss. u. Pract. Thierheilk.
1879 **Ollive**.—Sur la Résistance des Moutons de la Race Berberine à l'Inoculation du Charbon. Compt. Rend., T. 89.
Chauveau.—De la Prédisposition et de l'Immunité Pathologique. Compt. Rend., T. 89.
1880 **Pasteur**.—Bull. de l'Acad. de Méd. and Gaz. Méd. de Paris, Nr. 18.
Semmer and Krajewski.—Centralbl. f. d. Med. Wiss.
Toussaint.—Bull. de l'Acad. de Méd.; and Compt. Rend.
1881 **Toussaint**.—Compt. Rend.
Loeffler.—Zur Immunitätsfrage. Mitth. a. d. Ges. Amt, Bd. I.
Semmer.—Virchow's Arch., Bd. 83.
Grawitz.—Die Theorie der Schutzimpfung. Virchow's Arch., Bd. 48.

- 1881 **Toussaint**.—Gazette Médicale de Paris, Nr. 32.
Oemler.—Arch. f. Wiss. u. Pract. Thierheilk.
- 1882 **Rószahegyí**.—Pester Med.-Chir. Presse.
Koch.—Ueber die Milzbrandimpfung.
- 1883 **Frank**.—Jahresber. d. K. Thierarzneischule in München.
Arloing, Cornevin, and Thomas.—Du Charbon Bactérien; Pathogénie et Inoculations Préventives.
Chauveau.—Compt. Rend., T. 96, Nr. 9.
Chauveau.—Compt. Rend., T. 96, Nr. 10.
Chauveau.—Du Rôle de l'Oxygène de l'Air dans l'Atténuation quasi-instantanée des Cultures Virulentes par l'Action de la Chaleur. Compt. Rend., T. 96, Nr. 11.
Pasteur.—Sur la Vaccination Charbonneuse. Compt. Rend.
Pasteur.—La Vaccination Charbonneuse.
Pasteur.—Réponse au Doct. Koch. Revue Scientifique.
Pasteur.—Bull. de l'Acad. de Méd.
Büchner.—Eine neue Theorie über Erzielung v. Immunität gegen Infektionskrankheiten.
Perroncito.—Sull' Attenuazione del Virus Carbonchioso. Atti R. Acc. d. Lincei.
Chamberland and Roux.—Compt. Rend., T. 96, Nr. 15.
Chamberland.—Le Charbon et la Vaccination Charbonneuse d'après les Travaux Récents de M. Pasteur.
Massé.—Des Inoculations Préventives dans les Maladies Virulentes.
- 1884 **Feltz**.—De la Durée de l'Immunité Vaccinale Anticharbonneuse chez le Lapin. Compt. Rend., T. 99, p. 246.
Chauveau.—De l'Atténuation des Cultures Virulentes par l'Oxigène Comprimé. Gaz. Hebdom. de Méd. et de Chir., 22.
Koch, Gaffky, and Loeffler.—Exper. Studien über d. Künstl. Abschwächung der Milzbrandbacillen. Mitth. a. d. Ges. Amt, Bd. II.
Blazekovic.—Zur Präventiv-Inoculation Pasteur's. Oesterr Monatschr. f. Thierbeilk.
Pütz.—Vorträge f. Thierärzte, Ser. 7, Heft 1.
- 1885 **Hess**.—Vorl. Mitth. ü. die Schutzimpfung gegen Milzbrand im Kanton Bern nach der Methode Chauveau. Schweiz. Arch. f. Thierheilk., Bd. 27.
Strebel.—Zur Rauschbrandimpfung. Schweiz. Arch. f. Thierheilk.
Bouley.—L'Inoculation Préventive de la Fièvre Jaune. Compt. Rend., T. 100.

(J) *BACTERIA ASSOCIATED WITH DISEASES
 IN MAN AND ANIMALS.*

(I.) Actinomycosis.

- 1877 **Bollinger**.—Centralbl. f. d. Med. Wiss.
 1878 **Israël**.—Virchow's Arch., Bd. 74.

- 1879 **Israël.**—Virchow's Arch., Bd. 78.
- 1881 **Ponfick.**—Die Actinomykose. Berlin.
Johne.—Deutsche Zeitschr. f. Thiermed.
- 1882 **Hink.**—Centralbl. f. d. Med. Wiss.
Pflug.—Centralbl. f. d. Med. Wiss.
Gannet.—Boston Med. and Surg. Journ.
- 1883 **Pusch.**—Arch. f. Wiss. u. Pr. Thierheilk.
Bang.—Tidskrift for Veterinaerer.
Zemann.—Wien. Med. Jahrb., S. 477.
Fleming.—Actinomycosis.
- 1884 **Israël.**—Virchow's Arch., Bd. 96.
Mitteldorpf.—Deut. Med. Woch.
Karsten.—Deut. Med. Woch.
Chiari.—Prager Med. Woch., Nr. 10.
Treves.—Lancet.
Firket.—Rev. de Méd.
- 1885 **Israël.**—Kenntniss der Actinomykose des Menschen. Klinische Beiträge.
Boström.—Verh. d. Congr. f. Inn. Med. Wiesbaden.
Baumgarten.—Berl. Klin. Woch.
Johne.—Bericht ü. d. Veter.-Wesen i. K. Sachsen.
Murphy.—New York Med. Journ.
Ponfick.—Breslauer Aerztl. Zeitschr.
Soltmann.—Breslauer Aerztl. Zeitschr.
Magnussen.—Beiträge zur Diagnostik u. Casuistik der Actinomykose.
Diss. Kiel.
- 1886 **Hertwig.**—Archiv f. Wiss. u. Prakt. Thierheilk.
O'Neill.—Lancet.
Acland.—Brit. Med. Journ. and Trans. Path. Soc.
Roser.—Deut. Med. Woch.
Israël.—Archiv f. Klin. Chir.

(II.) Acute Yellow Atrophy.

- 1875 **Eppinger.**—Prag. Viertelj.
1882 **Hlava.**—Prag. Med. Wochenschr.
Balzer.—Archiv de Phys. Norm. et Path.

(III.) Anthrax.

- 1855 **Pollender.**—Vierteljahrschr. f. Ger. Med., Bd. 8.
1857 **Brauell.**—Virchow's Arch., Bd. 11.
1858 **Brauell.**—Virchow's Arch., Bd. 14.
1863 **Davaine.**—Compt. Rend. Paris. T. 57. Ibid., T. 59.
1865 **Davaine.**—Compt. Rend. Paris. T. 60.
1866 **Davaine.**—Compt. Rend. Paris. T. 61.
1872 **Bollinger.**—Centralbl. f. d. Med. Wiss., Bd. 10.

- 1873 Colin.—Bull. Acad. de Méd. Paris. T. 2.
 Davaine.—Compt. Rend. Paris. T. 77.
- 1876 Koch.—Beiträge zur. Biologie der Pflanzen, Bd. II., Heft 2.
- 1877 Pasteur.—Compt. Rend. Paris. T. 84.
 Toussaint.—Compt. Rend., T. 85.
 Davaine.—Rec. de Méd. Vét., T. 4.
 Davaine.—Compt. Rend.
 Bert.—Compt. Rend. Soc. de Biol., T. 4.
 Pasteur.—Bull. Acad. de Méd.
- 1878 Colin.—Bull. Acad. de Méd. Paris. T. 7.
 Toussaint.—Compt. Rend. Paris.
 Oemler.—Archiv f. Wiss. u. Pract. Thierheilk., Bd. 4.
 Ewart.—Quart. Journ. of Microsc. Sc.
 Bert.—Compt. Rend. Soc. de Biol., T. 5.
 Koch.—Wundinfectionskrankheiten.
- 1879 Schmidt.—Milzbrand bei Wildschweinen. Deut. Zeitschr. f. Thiermed.
 u. Vergl. Pathol.
 Toussaint.—Recherches Expérimentales sur la Maladie Charbonneuse.
 Bert.—Compt. Rend. Soc. de Biol., T. 6.
 Greenfield.—Quart. Journ. Micr. Sc. London.
 Oemler.—Archiv f. Wiss. u. Pract. Thierheilk.
 Colin.—Bull. Acad. de Méd. Paris. T. 8.
 Pasteur.—Bull. Acad. de Méd. Paris.
- 1880 Pasteur.—Compt. Rend. Paris. T. 90 and 91.
 Chauveau.—Compt. Rend., T. 90 and 91.
 Toussaint.—Compt. Rend.
 Colin.—Bull. Acad. de Méd. Paris. T. 9.
 Pasteur.—Bull. Acad. de Méd. Paris.
 Bouley.—Bull. Acad. de Méd. Paris. T. 9.
 Wachenheim.—Etude Expérimentelle sur la Septicité et la Virulence
 du Sang Charbonneux.
 Büchner.—Ueber die Exper. Erzeugung des Milzbrandcontagiums
 aus den Heupilzen.
 Büchner.—Versuche über die Entstehung des Milzbrands durch
 Einathmung. Sitzungsber. d. K. Bayer. Akad. d. Wissensch.
 Fokker.—Centralbl. f. d. Med. Wissensch., Bd. 18.
 Semmer.—Centralbl. f. d. Med. Wissensch., Bd. 18.
 Oemler.—Archiv f. Wissensch. u. Pract. Thierheilk.
 Szpilman.—Zeitschr. f. Physiol. Chemie. Strassburg. Bd. 4.
 Greenfield.—Proc. Roy. Soc. London.
- 1881 Pasteur.—Compt. Rend. Paris. T. 92.
 Chauveau.—Compt. Rend., T. 92.
 Bouley.—Compt. Rend., T. 92. Ibid., T. 93.
 Colin.—Bull. Acad. de Méd., T. 10.
 Bouley.—Bull. Acad. de Méd., T. 10.
 Rodet.—Contribution a l'Etude Expérimentelle du Carbon Bactérien.

- 1881 Koch.—Mitth. aus. d. Ges. Amt, Bd. I.
 Fokker.—Centralbl. f. d. Med. Wiss.
 Büchner.—Vorträge im Aerztl. Verein zu München.
 Huber.—Deut. Med. Woch., Bd. 7.
 Klein.—Rep. of the Medical Officer of the Local Govt. Board.
 Sternberg.—Am. Monthly Micr. Journ.
- 1882 Chauveau.—Compt. Rend. Paris. T. 94.
 Rodet.—Compt. Rend., T. 94.
 Pasteur.—Compt. Rend. Paris. T. 95.
 Feltz.—Compt. Rend., T. 95.
 Fokker.—Virchow's Archiv, Bd. 88.
 Archangelski.—Centralbl. f. d. Med. Wiss.
 Semmer.—Der Milzbrand und das Milzbrandcontagium.
 Esser and Schütz.—Zur Casuistik des Milzbrands. Mitth. a K. Preuss
 Amtl. Vet. Sanitätsbericht.
- 1883 Chauveau.—Compt. Rend. Paris. T. 96.
 Archangelski.—Centralbl. f. d. Med. Wiss.
 Roloff.—Ueber die Milzbrandimpfung und d. Entwicklung d. Milz-
 brandbakterien. Archiv f. Wissensch. u. Pract. Thierheilk., Bd. 9.
 Roloff.—Der Milzbrand. Berlin.
 Büchner.—Die Umwandlung der Milzbrandbakterien in unschädliche
 Bakterien. Virchow's Arch., Bd. 91.
 Toepper.—Die Neueren Erfahrungen über d. Aetiologie d. Milzbrands.
 Klein.—Quart. Journ. Micr. Sc.
 Dowdeswell.—Rep. Med. Off. Local Gov. Board.
- 1884 Bleuler.—Milzbrand beim Menschen. Correspondenzbl. d. Schweiz
 Aerzte.
 Schrakamp.—Zur Aetiologie des Milzbrandes. Archiv f. Hygiene,
 Bd. 2.
 v. Chelchowsky.—Zur Charakteristik des Milzbrandvirus. Der
 Thierarzt.
 Semmer.—Centralbl. f. d. Med. Wiss., Bd. 22.
 Prazmowski.—Acad. d. Wissensch. in Krakau.
 Prazmowski.—Biol. Centralbl., Bd. 4.
- 1885 Bollinger.—Zur Aetiologie des Milzbrands. Sitzungsber. d. Ges. f.
 Morphol. Physiol. zu München.
 Kitt.—Sitzungsber. d. Ges. f. Morphol. u. Physiol. zu München.
 Friedrich.—Zur Aetiologie des Milzbrands.
 Osol.—Experiment. Untersuch. ü. das Anthraxgift. Inaug. Diss.
 Dorpat.
- 1886 W. Koch.—Milzbrand und Rauschbrand. Stuttgart.
 Hoffa.—Die Natur des Milzbrandgiftes. Wiesbaden.
 v. Fodor.—Deut. Med. Woch.
 Johne.—Ber. ü. d. Veter. Wesen. i. K. Sachsen.
 Bollinger.—Arbeit. a. d. Patholog. Inst. zu München.
 Frank.—Zeitschr. f. Hygiene.



(IV.) Cattle Plague.

- 1883 **Semmer and Archangelski.**—Ueber das Rinderpestcontagium und dessen Mitigation. Centralbl. f. d. Med. Wiss.

(V.) Cerebro-Spinal Meningitis.

- 1833 **Leyden.**—Die Mikrokokken der Cerebrospinal-Meningitis. Centralbl. f. Klin. Med., Nr. 10.
1885 **Leichtenstern.**—Deut. Med. Woch., Nr. 23 u. 31.

(VI.) Chicken-Cholera.

- 1877 **Ioannès and Mégnin.**—Journ. d'Acclimatation.
1878 **Semmer.**—Hühnerpest. Deut. Zeitschr. f. Theirmed. u. Vergl. Path.
1879 **Perroncito.**—Ueber das Epizootische Typhöid der Hühner. Arch. f. Wiss. u. Prakt. Thierheilk.
1880 **Pasteur.**—Sur le Choléra des Poules. Compt. Rend., T. 90.
1882 **Zürn.**—Die Krankheiten des Hausgeflügels. Cornil.—Arch. de Physiol., Bd. 10.
1883 **Barthélémy.**—De l'Incubation des Œufs d'une Poule atteinte du Choléra des Poules. Compt. Rend., T. 96, No. 18.
Babès.—Arch. de Physiol.
Babès.—Compt. Rend. de l'Acad. d. Sc.
1884 **Petri.**—Centralbl. f. d. Med. Wiss.
1885 **Kitt.**—Mitth. über die Typhöidseuche des Geflügels. Allg. Deut. Geflügelzeitung.

(VII.) Cholera.

- 1883 **Strauss, Roux, Thuillier, and Nocard.**—Compt. Rend. Soc. de Biol., T. 4.
1884 **Finkler and Prior.**—Ueber den Bacillus der Cholera Nostras und seine Cultur. Naturforscherversammlung Magdeburg.
Pfeiffer.—Der bisherige Verlauf der Cholera in Thüringen, etc. Corresp.-Bl. des Allgem. Aerztl. Ver. in Thüringen, Nr. 9.
Koch.—Ueber die Cholera-bakterien. Deut. Med. Woch.
Emmerich.—Die Cholera in Neapel. Deut. Med. Woch., Nr. 50.
Finkler and Prior.—Unters. über Cholera Nostras. Deut. Med. Woch., Nr. 36.
Doyen.—Mikro-organismen in Leber und Niere von Choleraleichen. Soc. de Biol. de Paris. December 13th.
Büchner.—Ueber Cholera-bacillen. Münchn. Aerztl. Intelligenzbl., S. 549.
Bianchi.—Lancet.
Macnamara.—Brit. Med. Journ.
Hunter.—Brit. Med. Journ.
Carter.—Lancet.
Cameron.—Brit. Med. Journ.

- 1884 Bochefontaine.**—Expér. pour servir à l'Etude des Phénomènes déterminés chez l'Homme par l'Ingestion Stomacale du Liquide Diarrhéique du Choléra. *Compt. Rend.*
- Lewis.**—*Med. Times and Gazette.* September 20th.
- 1885 John.**—Einiges über die sogen. Choleraeure im. K. Ges. Amt. *Deutsche Zeitschr. f. Thiermed.,* Bd. XI.
- Finkler and Prior.**—Forschungen über Cholera-bakterien. *Ergänzungshefte zum Centralbl. f. Allg. Gesundheitspflege,* Bd. I., Heft 5 u. 6.
- Watson Cheyne.**—*Brit. Med. Journ.*
- Babès.**—*Virchow's Archiv.*
- Ferran.**—Sur l'Action Pathogène et Prophylactique du Bacillus-virgule. *Compt. Rend.,* T. 100, p. 959.
- Héricourt.**—Sur la Nature Indifférente des Bacilles-virgules. *Compt. Rend.*
- Villers.**—Sur la Formation des Ptomaines dans le Choléra. *Ibid.,* p. 91.
- Flügge.**—Kritik der Emmerich'schen Untersuchungen über Cholera. *Deut. Med. Woch.,* Nr. 2.
- Pfeiffer.**—Ueber die Cholera in Paris. *Deut. Med. Woch.,* Nr. 2.
- Deneke.**—Ueber eine Neue den Choleraspirlen ähnliche Spaltpilzart. *Deut. Med. Woch.,* Nr. 3.
- Miller.** Kommaförmiger Bacillus aus der Mundhöhle. *Deut. Med. Woch.,* Nr. 9.
- Klebs.**—Ueber Cholera Asiatica.
- Schottelius.**—Zum Mikrosk. Nachw. v. Cholera-bacillen in Dejectionen. *Deut. Med. Woch.,* Nr. 14.
- Drasche.**—*Allg. Wien. Med. Zeit.*
- van Ermengem.**—Die Ferran'schen Impfungen. *Deut. Med. Woch.,* Nr. 29.
- Gibier and van Ermengem.**—Rech. Exper. sur le Choléra. *Compt. Rend.,* T. 101.
- van Ermengem.**—Recherches sur le Microbe du Choléra Asiatique.
- Nicati and Rietsch.**—*Arch. de Physiol.*
- Nicati and Rietsch.**—*Revue d'Hygiène.*
- Nicati and Rietsch.**—*Revue de Médecin,* T. 5.
- Brunetti.**—Fatti Considerazioni Conclusioni sul Coléra.
- Klein.**—*Brit. Med. Journ. and Proc. Roy. Soc. London,* No. 38.
- Klein and Gibbes.**—An Inquiry into the Etiology of Asiatic Cholera. *Bluebook.*
- Büchner.**—*Archiv f. Hygiene.*
- Büchner and Emmerich.**—*Münch. Med. Woch.*
- Crookshank.**—Remarks on the Comma Bacillus of Koch. *Lancet.*
- 1886 Koch.**—Etiology of Cholera. Berlin Cholera Conference. Translated by Laycock, in *Microparasites and Disease (New Syd. Soc.).*
- Pettenkofer** (abstracted by Koplik).—*Lancet.*
- Cantani.**—*Deut. Med. Woch.*

- 1886 **Weisser**.—Zeitschrift f. Hygiene.
Bitter.—Archiv f. Hygiene.
Pfeiffer.—Deut. Med. Woch.
Klein.—Bacteria of Asiatic Cholera. Practitioner.
Weisser and Frank.—Zeitschr. f. Hygiene.

(VIII.) Dental Caries.

- 1867 **Leber and Rottenstein**.—Unters. über Caries der Zähne. Berlin.
1880 **Arndt**.—Beob. an Spirochæta Denticola. Arch. f. Pathol. Anat.,
Physiol. u. Klin. Med., Bd. 79.
1882 **Miller**.—Der Einfluss der Mikroorganismen auf die Caries der Zähne.
Arch. f. Exp. Path., XVI.
1883 **Miller**.—Ueber einen Zahnpilz *Leptothrix Gigantea*. Ber. d.
Deutsch. Bot. Ges., Heft 5.
1884 **Miller**.—Deutsche Med. Woch., Nr. 25 u. 36.
Miller.—Correspondenzbl. f. Zahnärzte, Bd. 13.
Rosenbach.—Mikroorganismen bei den Wundinfektionskrankheiten,
S. 77.

(IX.) Diphtheria.

- 1867 **Buhl**.—Micrococci of Diphtheria. Zeitschr. für Biol.
1868 **Hueter and Tommasi**.—Centralbl. f. d. Med. Wissensch.
1871 **Oertel**.—Deut. Arch. f. Klin. Med., Bd. 8.
1872 **Letzerich**.—Virch. Arch., Bd. 55.
Birch-Hirschfeld.—Archiv für Heilkunde.
1873 **Eberth**.—Der Diphtheritische Process. Med. Centralbl., XI, Nr. 8.
1874 **Letzerich**.—Mikrochemische Erscheinungen des Diphtheritispilzes.
Berl. Klin. Woch., XI.
1875 **Klebs**.—Arch. f. Exp. Pathol., Bd. 4.
1876 **Letzerich**.—Virch. Arch., Bd. 68.
1878 **Zahn**.—Beiträge zur Pathol. u. Histol. der Diphtherie.
1879 **Friedberger**.—Ueber Croup u. Diphtheritis beim Hausgeflügel. Deut.
Zeitschr. für Thiermed. u. Vergl. Pathol.
Nicati.—Compt. Rend., T. 88.
1880 **Neumayer**.—Neue Thesen zur Diphtheritisfrage.
1881 **Everett**.—Med. and Surg. Reporter.
Talamon.—Bull. de la Soc. Anat. de Paris, T. 56.
Cornil.—Arch. de Physiol.
Oertel.—Zur Aetiologie der Infectionskrankheiten.
Forster.—Wien. Med. Woch.
Lumner.—Aerztl. Int. Bl., No. 31.
1882 **Wood and Formad**.—Bull. Nat. Board of Health, Wash. and Med.
Times and Gazette.
Salisbury.—Gaillard's Med. Journ. New York.
Gerhardt and Klebs.—Verhandlung des Congresses für Innere Med.

- 1883 Fürbringer.—Virch. Arch., Bd. 91.
 Heubner.—Die Experimentelle Diphtherie.
 Gerhardt and Klebs.—Verhandl. d. Congresses f. Inn. Med.
 Francotte.—La Diphthérie.
 1884 Emmerich.—Deut. Med. Woch.
 Emmerich.—Compt. Rendus et Mémoires du V. Congrès Internat.
 d'Hygiène.
 Rivolta.—Die Diphtherie der Hühner im Vergleich zu der des
 Menschen. Giornale de Anat. Fisiol. e Patol. delli Anim.
 Loeffler.—Mittheil. a. d. Kais. Ges. Amt, Bd. II.
 1886 Loeffler.—Microparasites and Disease (New Syd. Soc.).

(X.) Erysipelas.

- 1868 Hüter.—Med. Centralbl., Nr. 34.
 1873 Raynaud.—Union Méd.
 Orth.—Archiv f. Exper. Pathol. u. Pharmacol., Bd. I.
 1874 Recklinghausen and Lankowski.—Ueber Erysipelas. Virchow's Arch.,
 Bd. 60.
 Lukomsky.—Virchow's Archiv, Bd. 60.
 1875 Trosier.—Bull. Soc. Anat. de Paris.
 Baader.—Zur Aetiologie des Erysipels. Schweiz. Naturf. Gesellsch.
 Klebs.—Archiv f. Exper. Pathol. u. Pharmacol., Bd. 4.
 1878 Tillmanns.—Verhandl. d. Deutsch. Ges. f. Chirurgie.
 1879 Tillmanns.—Experimentelle u. Anatomische Unters., etc. Archiv f. Klin.
 Chirurgie, Bd. 23.
 1880 Wolff.—Virchow's Arch., Bd. 81.
 1881 Dupeyrat.—Recherches Cliniques et Expérimentales sur la Pathogénie
 de l'Erysipèle.
 Fehleisen.—Ueber den Erysipelaspilz. Würzburger Phys. Med. Ges.
 1882 Fehleisen.—Deut. Zeitschr. f. Chir., Bd. 16.
 1883 Fehleisen.—Die Aetiologie des Erysipels. Berlin.
 1884 Janicke and Neisser.—Exitus Lethalis nach Erysipelas. Centralbl. f.
 Chir., Nr. 25.
 Rheiner.—Beitr. z. Path. Anat. des Erysipels bei Gelegenheit der
 Typhusepid. in Zürich. Virchow's Arch., Bd. 100, Heft 2.
 1885 Nepven.—Des Bactéries dans l'Erysipèle.
 Denuce.—Etude sur la Pathogénie et l'Anatomie pathologique de
 l'Erysipèle.

(XI.) Endocarditis.

- 1875 Koester.—Virchow's Arch., Bd. 72.
 1876 Birch-Hirschfeld and Gerber.—Archiv d. Heilkunde.
 1877 Wedel.—Berl. Klin. Wochenschr.
 1878 Klebs.—Archiv f. Exper. Pathol., Bd. 9.
 O. Rosenbach.—Ueber Artificielle Herzklappenfehler. Archiv fur Exper.
 Pathol., Bd. 9

- 1880 **Hamburg**.—Ueber Acute Endocarditis. Berlin. Inaug.-Diss.
Goodhart.—Trans. Path. Soc., Vol. XXXI.
- 1881 **Oberbeck**.—Casuistische Beiträge zur Lehre von der Endocarditis
 Ulcerosa. Inaug.-Diss.
Leyden.—Zeitschr. f. Klin. Med.
Weigert.—Virchow's Arch., Bd. 84.
Koch.—Mittheil. a. d. Kais. Ges. Amt, Bd. I.
Osler.—Trans. Int. Med. Congress.
- 1882 **Wilks**.—Brit. Med. Journ.
- 1883 **Kundrat**.—Sitz.-Ber. d. Kais. Acad. d. Wissensch. zu Wien.
- 1884 **Bramwell**.—Diseases of the Heart.
Bristowe.—Brit. Med. Journ.
Gibbes.—Brit. Med. Journ.
- 1885 **Wyssokowitsch**.—Centralbl. f. d. Med. Wissensch., Nr. 33.
Coupland.—Brit. Med. Journ.
Orth, J.—Versammlung Deutscher Naturf. zu Strasburg
Nocard.—Recueil de Méd. Vét.
Osler.—Brit. Med. Journ.
Weichselbaum.—Wien. Med. Woch.
- 1886 **Ribbert**.—Fortsch. d. Med.

(XII.) Glanders.

- 1882 **Löffler and Schütz**.—Ueber den Rotzpilz. Deut. Med. Woch., Nr. 52.
Boucharde, Capitan, and Charrin.—Bull. de l'Acad. d. Sc., Nr. 51.
- 1883 **Israël**.—Berl. Klin. Woch., Nr. 11.
Wassilieff.—Deut. Med. Woch., Nr. 11.
Molkentin.—Zur Sicherstellung der Diagnose von Rotz. Inaug.-Diss.
Struck.—Deut. Med. Woch., Nos. 51 u. 52.
Vulpian and Bouley.—Bull. de l'Acad. de Méd.
Fröhner.—Rotzige Elephantiasis des Kopfes beim Pferde. Rep. d.
 Thierheilk.
- 1884 **Kitt**.—Versuche über d. Züchtung des Rotzpilzes. Jahresber. d.
 München. Thierarzneisch.
Grünwald.—Zur Differentialdiagnose des Rotzes. Oesterr. Monatschr.
 für Thierheilk., Nr. 4.
Weichselbaum.—Zur Aetiologie der Rotzkrankheit des Menschen.
 Wiener Med. Woch., 21—24.
- 1886 **Löffler**.—Arbeit. a. d. K. Gesundh. Amt.

(XIII.) Gonorrhœa.

- 1879 **Neisser**.—Centralbl. f. d. Med. Wiss., Nr. 28.
Beter.—Centralbl. f. d. Med. Wiss.
- 1880 **Bücker**.—Ueber Polyarthrits Gonorrhœica. Diss.
Bokai.—Ueber das Contagium der Acuten Blennorrhœ. Allgem. Med.
 Centralzeitung, Nr. 74.
Bokai and Finkelstein.—Prager Med. Chir. Presse.

- 1880 Weiss.—Le Microbe du Pus Blennorrhagique.
- 1881 Aufrecht.—Pathologische Mittheilungen.
 Haab.—Der Mikrokokkus der Blennorrhœa Neonator.
 Hirschberg and Krause.—Zur Pathologie der Ansteckenden Augenkrankheiten. Centralbl. f. Pract. Augenheilk.
- 1882 Martin.—Rech. sur les Inflamm. Métast. à la suite de la Gonorrhée.
 Krause.—Die Mikrokokken der Blenorrhœa Neonator.
 Leistikow.—Ueber Bakterien bei den Venerischen Krankheiten. Charité-Annalen, 7 Jahrg., S. 750.
 Bockhart.—Sitzungsbericht d. Phys. Med. Ges. zu Würzburg.
- 1883 Bockhart.—Beitrag z. Aetiologie und Pathol. des Harnröhrentrippers. Viertelj. f. Dermatol. und Syph.
 Arning.—Gonokokken bei Bartolinitis. Viertelj. f. Dermatol. u. Syph., S. 371.
 Eschbaum.—Beitr. zur Aetiologie der Gonorrh. Secrete. Deut. Med. Woch., S. 187.
 Newberry.—Maryland Med. Journ.
 Campona.—Italia Medica.
 Aufrecht.—Mikrokokken i. d. Inneren Organen bei Nabelvenenentzündung Neugeborener. Centralbl. f. d. Med. Wiss., Nr. 16.
 Petrone.—Sulla Natura dell' Artrite Blenorragica. Rivista Clin., No. 2.
- 1884 Chameron.—Progrès Medical, 43.
 Sternberg.—Med. News, Vol. 45, Nr. 16.
 Kammerer.—Ueber Gonorrh. Gelenkentzündung. Centralbl. f. Chirurgie, Nr. 4.
 Welander.—Gaz. Med. de Paris.
 Kroner.—Zur Aetiologie der Ophthalmoblennorrhœa Neonator. Naturforschervers. in Magdeburg, Arch. f. Gyn., XXV., S. 109.
 Sanger.—Ueber Gonorrh. Erkrankung der Uterusadnexe. Naturforschervers. in Magdeburg. Ibid., S. 126.
 Oppenheimer.—Naturforschervers. in Magdeburg. Ibid., S. 51.
- 1885 Fränkel.—Deut. Med. Woch., S. 22.
 Bumm.—Der Mikroorganismus der Gonorrhöischen Schleimhauterkrankungen.
 Lundström.—Studier öfver Gonokokkus. Diss. Helsingfors.
 E. Fränkel.—Mikrokokken bei Colpitis. Deut. Med. Woch., Nr. 2.
- 1886 Smirnoff.—Vrach.

(XIV.) Hydrophobia.

- 1881 Colin.—Bull. Acad. de Méd. Paris, T. 10.
 Doléris.—Gaz. Méd. de Paris, T. 3. Tribune Méd. Paris, T. 14.
 Pasteur.—Compt. Rend.
- 1882 Pasteur, Chamberland, Roux, and Thuiller.—Compt. Rend.
 Bert.—Compt. Rend.
- 1883 Gibier.—Compt. Rend.

- 1884 **Pasteur**.—Nouvelle Communication sur la Rage. Ann. de Méd. Vétérin.
Percheron.—La Rage et les Expériences de M. Pasteur.
Pasteur.—Compt. Rend.
 Fol.—Acad. des Sciences.
- 1886 **Babès**.—In Les Bactéries.
Dowdeswell.—Journ. Roy. Micro. Soc. and Lancet
Dolan.—Hydrophobia. M. Pasteur and his Methods.
Pasteur.—Compt. Rend.
Vignal.—Brit. Med. Journ.
Lankester.—Nineteenth Century, No. 114.
Kerr.—Brit. Med. Journ.
Bauer.—Münch. Med. Woch.
Frisch.—Wien. Med. Woch.
Hime.—Experimental Researches concerning Pasteur's Prophylactic. Lancet.

(XV.) Leprosy.

- 1879 **Neisser**.—Breslauer Aerztl. Zeitschr.
Neisser.—Jahresber. d. Schles. Ges. für Vaterl. Cultur.
- 1880 **Hansen**.—Virchow's Archiv.
Gaucher and Hillairet.—Progrès Méd.
- 1881 **Neisser**.—Virchow's Archiv, Bd. 84.
Cornil and Suchard.—Ann. de Dermat. et Syph.
Vossius.—Uebertragungsversuche von Lepra auf Kaninchen. Ber. über d. Ophthalmologengcongress in Heidelberg.
John Hills.—On Leprosy in British Guiana.
- 1882 **Köbner**.—Uebertragungsversuche von Lepra auf Thiere. Virchow's Arch.
Köbner.—Virchow's Arch.
Hausen, Armauer.—Virchow's Arch., Bd. 90.
- 1883 **Damsch**.—Uebertragungsversuche von Lepra auf Thiere. Virch. Arch., Bd. 92. Centralbl. f. d. Med. Wissensch., Bd. 21.
Kaposi.—Wiener Med. Woch.
Hillis.—Trans. Path. Soc.
Babès.—Etude Comparative des Bactéries de la Lèpre et de la Tuberculose. Compt. Rend.
Moretti.—Il Primo Caso di Lebbra nelle Marcho Confermato dalla Presenza del Bacillus Lepræ.
Müller.—Deut. Archiv f. Klin. Med., Bd. 34.
- 1884 **Vidal**.—La Lèpre et son Traitement.
Arning.—Ueber das Vorkommen des Bacillus Lepræ bei Lepra Anæsthetica s. Nervorum. Virchow's Archiv, Bd. 97.
Thin.—Brit. Med. Journal.
- 1885 **Virchow**.—Berl. Klin. Woch., N. 12.
Unna.—Ueber Leprabacillen. Deut. Med. Woch., Nr. 32.

- 1885 Steven.—Brit. Med. Journ.
 1886 Touton.—Fortsch. d. Med.
 Unna.—Deut. Med. Woch.
 Neisser.—Virchow's Archiv.
 Unna and Lutz.—Dermatologische Studien, Heft I.
 Melcher and Ortman.—Berl. Klin. Woch.

(XVI.) Malaria.

- 1879 Klebs and Tommasi-Crudeli.—Arch. f. Exp. Pathol., Bd. 2.
 1880 Tommasi-Crudeli.—Der Bacillus Malariae im Erdboden von Selinunte u. Campobello. Arch. f. Exp. Pathol., Bd. 12.
 Cuboni and Marchiafava.—Neue Studien üb. d. Natur der Malaria. Arch. f. Exp. Pathol., Bd. 13.
 1881 Sternberg.—Bull. Nat. Board of Health, Washington.
 Cuboni and Marchiafava.—Atti della R. Acad. dei Lincei.
 Laveran.—Compt. Rend.
 1882 Marchand.—Zur Aetiologie der Malaria. Virch. Archiv, Bd. 88.
 Ziehl.—Deutsch. Med. Woch., Nr. 48.
 Roszahegyi.—Von der Ursache des Wechselfiebers. Biol. Centralbl., Bd. 2.
 Tommasi-Crudeli.—Die Malaria von Rom.
 Richard.—Bestätigung der Laveranschen Beob. Compt. Rend., No. 8.
 Laveran.—Les Parasites du Sang dans l'Impaludisme. Compt. Rend., No. 17.
 Bardels.—Gaz. des Hôpit.
 1883 Torelli.—La Malaria in Italia.
 Maurel.—L'Étiologie et la Nature du Paludisme. Ann. d'Hygiène.
 1884 Kotelmann.—Der Bacillus Malariae im Alterthum. Virchow's Arch., Bd. 97.
 Gerhardt.—Zeitschr. f. Klin. Med., Bd. 7.
 Sehlen.—Fortschr. d. Med., Bd. II.
 Leoni.—Gazetta Medica di Roma.
 Marchiafava and Celli.—Atti della R. Accademia dei Lincei.
 Mariotti and Ciarrocchi.—Lo Sperimentale, T. 54.
 Tommasi-Crudeli.—La Production Naturelle de la Malaria. Conférence faite à la 8, Sess. du Congrès Intern. Méd. à Copenhague.
 Laveran.—Traité des Fièvres Palustres.
 1885 Marchiafava and Celli.—Neue Untersuchungen über die Malariainfection. Fortschr. d. Med., Bd. 3.
 1886 Camillo Golgi.—Archivio per le Scienze Mediche.

(XVII.) Malignant Œdema.

- 1862 Davaine.—Bull. de l'Acad. de Méd.
 1877 Pasteur.—Vibron. Septique. Bull. de l'Acad. de Méd.
 1881 Koch.—Mitth. aus dem Ges. Amt, I., S. 54.
 Gaffky.—Mitth. as. d. K. Ges. Amt.
 Pasteur.—Vibron. Septique. Bull. de l'Acad. de Méd.

- 1882 Brieger and Ehrlich.—Berl. Klin. Wochenschr., N. 44.
 Lebedeff.—Versuche über Desinfection bei Malignen Oedembacillen.
 Arch. de Phys. Norm. et Path.
- 1884 Lustig.—Zur Kenntniss Bakteriämischer Erkrankungen bei Pferden
 (Malignes Oedem). Jahresber. d. K. Thierarzneischule zu Hannover.
 Kitt.—Unters. über Malignes Oedem und Rauschbrand bei Hausthieren.
 Jahresber. der K. Thierarzneischule in München.
- 1885 Hesse, W. and R.—Ueber Züchtung der Bacillen des Malignen Oedems.
 Deut. Med. Woch.

(XVIII.) Measles.

- 1882 Keating.—Phil. Med. Times.
 1883 Cornil and Babès.—Archiv de Phys.

(XIX.) Ophthalmic Diseases.

- 1872 Röth.—Retinitis Septica. Virchow's Archiv, Bd. 55.
- 1879 Balogh.—Sphærobakterien in der entzündeten Hornhaut. Med. Centralbl., XIV.
- 1882 Kahler.—Ueber Septische Retinitis. Prager Zeitschr. f. Prakt. Heilk.,
 Bd. I.
 Sattler.—Unters. über das Trachom. Ber. üb. d. Ophthalmologen
 Congress zu Heidelberg.
 Michel.—Graefe's Archiv f. Augenheilkunde.
- 1883 Schleich.—Verh. des Ophthalmologen Congr. zu Heidelberg.
 Bock.—Ueber die Miliare Tuberkulose der Uvea. Virchow's Arch.,
 Bd. 91.
 Kuschbert and Neisser.—Zur Pathologie und Aetiologie der Xerosis
 Conjunctivæ. Bresl. Aerztl. Zeitschr., Nr. 4.
 Sattler.—Ueber d. Natur der Jéquirity-Ophthalmie. Zehender's Klin.
 Monatsblatt; and Wien. Med. Woch., Nr. 17.
 Sattler and de Wecker.—L'Ophthalmie Jéquiritique.
 Cornil and Berlioz.—Sur l'Empoisonnement par le Jéquirity. Compt.
 Rend. de l'Acad. d. Sc.
- 1884 v. Reuss.—Pilzconcretionen i. d. Thränenröhrchen. Wien. Med.
 Presse.
 Goldzieher.—Streptothrix Færsteri im unteren Thränenröhrchen. Cen-
 tralblatt f. Prakt. Augenheilk.
 Kroner.—Zur Aetiologie der Ophthalmoblennorrhœ. Verh. d. Natur-
 forschers-Vers. Magdeburg.
 Kuschbert.—Deutsch. Med. Woch., Nr. 21.
 Vossius.—Berl. Klin. Wochenschr., Nr. 17.
 Klein.—Centralbl. f. d. Med. Wiss., Nr. 8.
 Vennemann and Braylants.—Le Jéquirity et son Principe Pathogène.
 Neisser.—Fortschr. d. Med., Bd. 2, S. 73.
 Salomonsen.—Fortschr. d. Med., Bd. 2, S. 78.

- 1885 **Widmark**.—Hygeia. Stockholm.
Deutschmann.—v. Graefe's Archiv, Bd. XXXI.
 1886 **Gifford**.—Archiv f. Augenheilkunde.
v. Zehender.—Bowman Lecture. Lancet.

(XX.) Osteomyelitis.

- 1873 **Collmann**.—Bakterien im Organismus eines an einer Verletzung am Oberschenkel verstorbenen Mädchens. Göttingen.
 1875 **Eberth**.—Virchow's Arch., Bd. 65.
 1876 **Friedmann**.—Fall von Primärer Infectiöser Osteomyelitis. Berl. Klin. Woch.
Schüller.—Zur Kenntniss der Mikrokokken bei Acuter Infectiöser Osteomyelitis. Mikrokokkenherde im Gelenkknorpel. Centralbl. f. Chirurgie, Nr. 12.
 1882 **Fehleisen**.—Phys. Med. Ges. Würzburg.
 1883 **Becker**.—Vorl. Mitth. über den die Acute Infectiöse Osteomyelitis erzeugenden Mikroorganismus. Deut. Med. Woch., and Berl. Klin. Woch.
 1884 **Peyroud**.—Compt. Rend.
Krause.—Ueber einen bei der Acuten Infectiösen Osteomyelitis vorkommenden Mikrokokkus. Fortschr. d. Med., Bd. 2.
Rosenbach.—Vorl. Mitth. über die Acute Osteomyelitis beim Menschen erzeugenden Mikroorganismen. Centralbl. f. Chirurgie.
Rodet.—Compt. Rendus, T. 99.
 1885 **Garré**.—Fortschr. d. Med.

(XXI.) Pleuropneumonia.

- 1879 **Sussdorff**.—Ueber d. Lungenseuche des Rindes. Deutsche Zeitschr. f. Thiermed. u. Vergl. Pathol.
 1880 **Bruylants and Verriers**.—Bull. de l'Acad. Belg.
 1882 **Pasteur**.—Note sur la Péripleurite Contagieuse des Bêtes à Cornes. Recueil de Méd Vét.
 1883 **Cornil and Babès**.—Arch. de Physiol. Norm. et Path., T. 2.
 1884 **Poels and Nolen**.—Centralbl. f. d. Med. Wiss., Nr. 9.
Mayrwieser.—Ueber Infectiösen Bronchialcroup bei Rindern. Woch. f. Thierheilk. u. Viehzucht, 19.
 1886 **Poels and Nolen**.—Das Contagium der Lungenseuche. Fortsch. d. Med.
Poels.—Septische Pleuro-pneumonia der Kälber. Fortsch. d. Med.

(XXII.) Pneumonia.

- 1878 **Kühn**.—Die Contagiöse Pneumonic. Deutsch. Arch. f. Klin. Med.
 1880 **Bruylant and Verriers**.—Bull. de l'Acad. Belge.
 1881 **Kühn**.—Die Uebertragbarkeit Endemischer Pneumonieformen auf Kaninchen. Berl. Klin. Wochenschr., Nr. 38.
 1882 **Friedländer**.—Virchow's Arch., Bd. 87.

- 1883 Salvioli and Zäslein.**—Ueber den Micrococcus und die Pathogenese der Croupösen Pneumonie. Centralbl. f. d. Med. Wissensch.
- Matray.**—Wien. Med. Presse, Nr. 23.
- Ziehl.**—Ueber das Vorkommen der Pneumoniekokken im Pneumonischen Sputum. Centralbl. f. d. Med. Wissensch.
- Friedländer.**—Fortschr. d. Med., Bd. I.
- Talamon.**—Progr. Médic.
- Giles.**—Brit. Med. J., Vol. II.
- Friedländer and Frobenius.**—Berl. Klin. Woch.
- Griffini and Cambria.**—Centralbl. f. d. Med. Wiss.
- 1884 Friedländer.**—Fortschr. d. Med., Bd. II.
- Ziehl.**—Ueber das Vorkommen der Pneumoniekokken im Pneumonischen Sputum. Centralbl. f. d. Med. Wissensch.
- Klein.**—Centralbl. f. d. Med. Wissensch.
- Nauwerck.**—Ueber Morbus Brightii bei Croupöser Pneumonie. Beitr. zur Pathol. Anat. von Ziegler.
- Jürgensen.**—Berl. Klin. Wochenschr., Bd. 22.
- Mendelsohn.**—Zietschr. f. Klin. Med., Bd. 7.
- Platonow.**—Ueber die Diagnostische Bedeutung d. Pneumoniekokken. Inaug.-Diss. Würzburg.
- Korányi and Babès.**—Pester. Med. Chirurg. Presse.
- A. Fränkel.**—Verhandl. d. Congr. f. Innere. Med., Fortschr. d. Med. Nov.
- Germain-Sée.**—Compt. Rend. Acad. de Sc. Paris.
- Afanassiew.**—Compt. Rend. Soc. de Biol. Paris, T. 5.
- Salvioli and Zäslein.**—Arch. pour les Sc. Méd., T. 8.
- Maguire.**—Brit. Med. Journ., Vol. II.
- 1885 Dreschfeld.**—Ueber Wanderpneumonie u. ihre Beziehung zur Epidemischen Pneumonie. Fortschr. d. Med., Bd. 3, 389.
- Schou.**—Untersuchungen über Vaguspneumonie. Fortschr. der Medicin, Bd. 3, Nr. 15.
- Ribbert.**—Zur Färbung d. Pneumoniekokken. Deut. Med. Wochenschr., Nr. 9.
- Germain-Sée.**—Des Maladies Spécifiques du Poupon.
- de Blasi.**—Rivista Internaz. di. Med. e. Chir.
- Pawlowsky.**—Berl. Klin. Woch.
- Platanow.**—Mitth. a. d. Würzburg. Med. Klinik.
- Sternberg.**—Amer. Journ. Med. Sciences.
- Peterlein.**—Bericht ü. d. Vet.-Wesen. i. K. Sachsen.
- Friedländer.**—Beste Färbung zur Darstellung der Kapseln der Pneumoniekokken. Fortschritt, Bd. 3, 92.
- 1886 Sternberg.**—Micrococcus Pasteuri. Journ. Royal Microscop. Soc.
- Neumann.**—Berl. Klin. Woch.
- Pipping.**—Fortschr. d. Med., Nos. 10 and 14.
- Fränkel.**—Zeitschr. f. Klin. Med., Bd. X. and XI.
- Fränkel.**—Deut. Med. Woch.

- 1886 **Foa and Bordoni-Uffreduzzi.**—Deut. Med. Woch.
Wechselbaum.—Wien. Med. Woch.
Manfredi.—Fortsch. d. Med.
Jaccoud.—La France Médicale.
Lancereaux and Besançon.—Archiv. Gén. de Méd.
Thost.—Deut. Med. Woch.

(XXIII.) Puerperal Fever.

- 1865 **Mayrhofer.**—Vibrionen als Krankheitsursache des Puerperalfiebers.
 Monatsschr. f. Geburtsk. u. Frauenkrankheiten, Bd. 25.
 1872 **Waldeyer.**—Ueber Vorkommen von Bakterien bei der Diphtheritischen
 Form des Puerperalfiebers. Arch. f. Gynäkologie, III.
 1873 **Heiberg.**—Die Puerperalen und Pyämischen Prozesse.
 Orth.—Virchow's Arch., Bd. 58.
Recklinghausen and Lankowski.—Ueber Erysipelas. Virchow's Arch.,
 Bd. 60.
 1879 **Laffter.**—Gehirnerweichungsherde durch Mikrokokkenembolien bei
 Puerperaler Pyämie. Bresl. Aerztl. Ztg.
 1880 **Dolérís.**—La Fièvre Puerpérale et les Organismes Infect.
Pasteur.—Bull. de l'Acad. de Méd., T. 9.
 1881 **Karewski.**—Exper. Unters. über die Einwirkung Puerperaler Secrete
 auf den Thierischen Organismus. Zeitschr. f. Geburtsh. u. Gynä-
 kologie, Bd. 7.
 1884 **Aufrecht.**—Die Exper. Erzeugung der Endometritis Diphtherica Puer-
 peralis. Naturforsch. Versamml.

(XXIV.) Pyæmia, Septicæmia, and Suppuration.

- 1866 **Rindfleisch.**—Lehrb. der Pathol. Gewebelehre. 1. Aufl., S. 204.
 1867 **Waldeyer.**—Zur Pathol. Anatomie der Wundinfectionskrankheiten.
 Virchow's Arch., Bd. 40.
Lister.—Lancet.
 1870 **Ainstie.**—Lancet.
 1871 **Tiegel.**—Ueber d. Fiebererregenden Eigenschaften des Microsporon
 Septicum. Bern. Diss.
Waldeyer.—Mikrokokkencolonien in den Parenchymatischen Organen.
 Vortrag i. d. Med. Ges. zu Breslau. 4 Aug.
 1872 **Burdon-Sanderson.**—Trans. Path. Soc.
 1873 **Birch-Hirschfeld.**—Untersuchungen über Pyämie.
Heiberg.—Die Puerperalen u. Pyämischen Prozesse.
 1874 **Tiegel.**—Virchow's Archiv, Bd. 60.
 1875 **Barthold.**—Pyämisch-Metast. Dissert. Berlin.
Burdon-Sanderson.—Brit. Med. Journ.
 1877 **Balfour.**—Edinb. Med. Journ.
Lister.—Med. Times and Gazette.
 1878 **Dowdeswell.**—Quart. Journ. Micr. Sc. London, Vol. 18.

- 1878 Koch.—Wundinfektionskrankheiten. Leipzig.
 Bastian.—Brit. Med. Journ.
- 1880 Perret.—De la Septicémie. Paris.
 Beck.—Rep. Loc. Govt. Board.
 Drysdale.—Pyrexia.
- 1881 Koch.—Mittheil. d. Kais. Ges. Amts, Bd. I.
 Oertel.—Zur Aetiologie der Infektionskrankheiten.
 Ogston.—Brit. Med. Journ., Vol. I.
 Béchamp.—Compt. Rend.
 Béchamp.—Trans. Internat. Med. Cong. London.
 Horsley.—Rep. Med. Officer Loc. Govt. Board.
 Lister.—Quart. Journ. Microscop. Science.
- 1882 Ogston.—Journ. of Anat. and Phys., Vol. 17.
 Braidwood and Vacher.—Brit. Med. Journ.
 Sternberg.—Amer. Journ. Med. Sc.,—John Hopkins' Univ. Stud.
 Biol. Lab.
- 1883 Dowdeswell.—Proc. Roy. Soc. London, Vol. 34.
 Babès.—Compt. Rend.
 Dreschfeld.—Brit. Med. Journ.
 Sutton.—Trans. Path. Soc.
- 1884 Rosenbach.—Mikroorganismen bei den Wundinfektionskrankheiten des
 Menschen. Wiesbaden.
 Arloing.—Recherches sur les Septicémies.
 Watson-Cheyne.—Trans. Path. Soc., XXXV.
 Steven.—Glasgow Med. Journal.
- 1885 Passet.—Ueber Mikroorganismen der Eitrigen Zellgewebsentzündung
 des Menschen. Fortschritte d. Med., Nr. 2.
 Garré.—Zur Aetiologie acut Eitriger Entzündungen. Fortschritte d.
 Med., 165.
 Passet.—Fortschritte d. Medicin, Bd. 3.
 Klemperer.—Zeitschr. f. Klin. Med.
 Hoffa.—Fortsch. d. Med.
- (XXV.) Relapsing Fever.
- 1873 Obermeier.—Vorkommen Feinster, Eigene Bewegung zeigender Fäden
 im Blute von Recurrenkranken. Med. Centralbl., 11. Berl. Med.
 Ges. Berl. Klin. Wochenschr.
- Engel.—Ueber die Obermeier'schen Recurrensspirillen. Berl. Klin.
 Woch.
- 1875 Laptshinsky.—Centralbl. f. d. Med. Wissensch., Bd. 13.
- 1876 Weigert.—Deut. Med. Woch.
 Heydenreich.—St. Petersb. Med. Woch.
 Manassem.—St. Petersb. Med. Woch., No. 18.
- 1877 Heydenreich.—Der Parasit des Rückfalltyphus.
- 1879 Carter.—Lancet.
 Albrecht.—St. Petersb. Med. Woch.

- 1879 **Cohn**.—Deut. Med. Woch.
Koch.—Deut. Med. Woch.
 1880 **Guttman**.—Virchow's Arch.
Mühlhäuser.—Virchow's Arch., Bd. 97.
Jaksch.—Wien. Med. Wochenschr. Juli.
 1885 **Günther**.—Fortschr. d. Med.

(XXVI.) Rhinoscleroma.

- 1885 **Cornil**.—B. de la Soc. Anatom. 15 Fev.
Cornil and Alvarez.—Sur les Micro-organismes du Rhinosclérome.
 Acad. de Méd. et Archiv. de Phys. Norm. et Path.
 1886 **Paltauf and Eiselsberg**.—Fortsch. d. Med.
Davies.—Brit. Med. Journ.

(XXVII.) Scarlatina.

- 1872 **Coze and Feltz**.—Les Maladies Infectieuses.
McKendrick.—Brit. Med. Journal.
 1876 **Klein**.—Report of the Medical Officer of the Privy Council.
 1882 **Hahn**.—Berl. Klin. Woch., No. 38.
 1883 **Roth**.—Münchener Aerztl. Intelligenzbl.
Pohl-Pincus.—Centralbl. f. d. Med. Wiss., No. 36.
Crooke.—Lancet.
 1884 **Heubner and Bahrdt**.—Zur Kenntniss der Gelenkeiterungen bei
 Scharlach. Berl. Klin. Woch., Nr. 44.
 1885 **Crooke**.—Fortsch. d. Med.
 1886 **Laure**.—Lyon Médical.
Klein.—Nature, XXXIV.

(XXVIII.) Swine-Erysipelas.

- 1881 **Salmon**.—Report Depart. Agricul. Washington.
Detmers.—Science.
 1882 **Pasteur**.—Sur le Rouget ou Mal Rouge des Porcs. Compt. Rend., T. 95.
 1883 **Cornil and Babès**.—Arch. de Physiol.
Pasteur and Thuillier.—Bull. de l'Acad. de Méd. de Paris. Compt.
 Rend., T. 97.
Eggeling.—Ueber den Rothlauf der Schweine. Fortschr. d. Med.
 1884 **Salmon**.—Rep. Dept. Agricult. Washington.
 1885 **Löffler**.—Experimentelle Untersuchungen über Schweinerothlauf
 Arbeiten aus dem Kaiserl. Gesundheits Amt, Bd. I.
Schütz.—Ueber den Rothlauf der Schweine und die Impfung desselben.
 Ibid.
Lydtin and Schottelius.—Der Rothlauf der Schweine. Wiesbaden.

(XXIX.) Swine-Typhoid.

- 1879 **Klein**.—Rep. of the Med. Offic. of the Privy Council [1877-1878].
 1884 **Klein**.—Virchow's Archiv.

(XXX.) Symptomatic Anthrax.

- 1878 **Bollinger and Feser.**—Wochenschr. f. Thierheilkunde.
 1880 **Arloing, Corneving, and Thomas.**—Compt. Rend.
 1881 **Arloing, Corneving, and Thomas.**—Bull. de l'Acad. de Méd. and Revue de Méd.
 1883 **Arloing, Corneving, and Thomas.**—Du Charbon Bactérien, Charbon Symptomatique, etc.
 1884 **Kitt.**—Unters. über Malignes Oedem und Rauschbrand bei Hausthieren. Jahresber. der K. Thierarzneisch. in München.
Babès.—Journ. de l'Anatomie.
Neelsen and Ehlers.—Ueber den Rauschbrand. Ber. d. Naturforsch. Ges. zu Rostock.
Ehlers.—Unters. üb. d. Rauschbrandpilz. Inaug. Diss. Rostock.
Arloing, Corneving, and Thomas.—Revue de Méd.
 1886 **Arloing, Corneving, and Thomas.**—Chabert's Disease. Transl. by Dawson Williams in Microparasites and Disease. (New Syd. Soc.)
Hess.—Bericht über die entschädigten Rauschbrand u. Milzbrandfälle im Canton Bern.
Strebel.—Schweiz. Archiv f. Thierheilk.

(XXXI.) Syphilis.

- 1872 **Lostorfer.**—Arch. f. Dermat. u. Syph.
 1879 **Klebs.**—Arch. f. Exp. Pathol., Bd. 10.
 1881 **Aufrecht.**—Centralbl. f. d. Med. Wissensch., Bd. 19.
 1882 **Birch-Hirschfeld.**—Bakterien in Syphilitischen Neubildungen. Centralbl. f. d. Med. Wissensch., Nrs. 33, 34.
Peschel.—Centralbl. f. Augenheilk.
Martineau and Hamonic.—De la Bactéridie Syphilitique. Compt. Rend., p. 443.
Morison.—Maryland Med. Journ.
 1883 **Letnik.**—Wien. Med. Wochenschr.
Morison.—Maryland Med. Journ. Baltimore.
Morison.—Wiener Med. Wochenschr.
Heyden.—Préservation de la Syphilis, etc. Traduit par Roberts.
 1884 **Petrone.**—Gaz. Medica Ital.
Torney and Marcus.—Compt. Rend., p. 472.
Lustgarten.—Wien. Med. Woch., Nr. 47.
Königer.—Deut. Med. Woch., S. 816.
 1885 **Lustgarten.**—Die Syphilisbacillen.
Doutrelepont and Schütz.—Deut. Med. Woch., Nr. 19.
Alvarez and Tavel.—Bull. de l'Acad. de Méd. et Archiv. de Phys. Norm. et Path.
de Giacomi.—Neue Färbungsmethode der Syphilisbacillen. Correspondenzbl. f. Schweizer Aerzte, Bd. 15.
Gottstein.—Fortschr. d. Med., Bd. 3, S. 543.

- 1885 Klemperer.—Deut. Med. Woch.
 1886 Eve and Lingard.—Brit. Med. Journ.
 Bienstock.—Fortsch. d. Med.
 Kassowitz and Hochsinger.—Wien. Med. Blätter.
 Disse and Taguchi.—Deut. Med. Woch.

(XXXII.) Tetanus.

- 1884 Carle and Rattone.—Studio Sperimentale sull' Etiologia del Tetano.
 Giorn. della R. Acad. di Medicina di Torino.
 Vogel.—Drei Fälle von Infectiösem Tetanus. Deut. Med. Woch.,
 Nr. 31.
 Nicolaier.—Deut. Med. Woch., Nr. 52.
 1886 Rosenbach.—Archiv f. Klin. Chir.

(XXXIII.) Tuberculosis.

- 1868 Villemin.—Etude sur la Tuberculose.
 Klebs.—Virchow's Arch., Bd. 44.
 1873 Klebs.—Arch. f. Exp. Pathol. u. Pharmakol., Bd. I.
 1877 Cohnheim.—Uebertragbarkeit der Tuberculose. Berlin.
 1879 Baumgarten.—Berl. Klin. Woch., Bd. 17.
 1881 Toussaint.—Compt. Rend., T. 93.
 Baumgarten.—Centralbl. f. d. Med. Wiss. Berlin. Bd. 19.
 1882 Baumgarten.—Centralbl. f. d. Med. Wiss. Berlin. Bd. 20.
 Baumgarten.—Deut. Med. Woch., Bd. 8.
 Wahl.—Zur Tuberculosefrage. Deut. Med. Woch., Nr. 46.
 van Ermengem.—Le Microbe de la Tuberculose. Ann. de la Soc.
 Belge de Microscopie.
 Hiller.—Deut. Med. Woch., Bd. 8.
 Balogh.—Wien. Med. Woch.
 Koch.—Die Aetiologie der Tuberculose. Berl. Klin. Woch.
 Formad.—The Bacillus Tuberculosis. The Philad. Medical Times.
 Green.—Lancet.
 Creighton.—Trans. Path. Soc.
 Rindfleisch.—Phys. Med. Ges. zu Würzburg, Nr. 8.
 Ewart.—Lancet.
 Ehrlich.—Deut. Med. Woch.
 Fränzel and Palmers.—Berl. Klin. Woch.
 Aufrecht.—Centralbl. für d. Med. Wiss.
 1883 Schuchardt and Krause.—Fortschr. d. Med.
 Klebs.—Arch. f. Exp. Pathol. u. Pharmakol., Bd. 17.
 Ehrlich.—Deut. Med. Woch., Bd. 9.
 Fränzel and Palmers.—Berl. Klin. Woch.
 de Giacomi.—Fortschr. d. Med., Bd. I., S. 145.
 Meisels.—Wien. Med. Woch.
 Raymond.—Arch. Gén. de Méd., T. 11.

- 1883 Demme.**—Jahresber. d. Jennerschen Kinderspitals. Bern.
Lydtin.—Badische Thierärztl. Mittheil.
Ribbert.—Ueber d. Verbreitungsweise der Tuberkelbacillen bei Hühnern. Deut. Med. Woch., Nr. 28.
Wechselbaum.—Wien. Med. Jahrb.
Johne.—Ber. üb. d. Veterinärwesen im Königr. Sachsen.
Bollinger.—Münch. Aerztl. Intelligenzbl., Nr. 16.
Johne.—Die Geschichte der Tuberculose.
Pütz.—Ueber die Beziehungen der Tuberculose des Menschen zu der Thiere.
Celli and Guarnieri.—Arch. pour les Sciences Médic.
Weigert.—Deut. Med. Woch., Nr. 24ff.
Nauwerck.—Deut. Med. Woch.
Smith.—Bristol Med. Chir. Journ.
Williams.—Lancet.
Veraguth.—Arch. f. Exp. Path. u. Pharmakol., Bd. 16.
Schottelius.—Virchow's Archiv, Bd. 91.
Brouilly.—Note sur la Présence des Bacilles dans les Lésions Chirurgicales Tuberculeuses. Rev. de Chir., T. 3.
Schlegtendal.—Fortschr. d. Med., Bd. I.
Küssner.—Beitr. z. Impftuberculose. Deut. Med. Woch., Nr. 36.
Aufrecht.—Centralbl. f. d. Med. Wiss., Bd. 21.
Bollinger.—Centralbl. f. d. Med. Wiss., Bd. 21, S. 600.
Cheyne.—Brit. Med. Journ., Vol. I.
Cheyne.—Practitioner. London. Vol. XXX.
Bock.—Virchow's Archiv, Bd. 91.
Dettweiler.—Berl. Klin. Woch., Bd. 21.
Damsch.—Deut. Med. Woch., Nr. 17.
Babès.—Der erste Nachweis des Tuberkelbacillus im Harn. Centralbl. f. d. Med. Wissensch.
Babès.—Compt. Rendus.
Rosenstein.—Centralbl. f. d. Med. Wiss.
Levinsky.—Deut. Med. Woch., Bd. 9.
Lichtheim.—Fortschr. d. Med., Bd. I.
West.—Lancet, Vol. I.
West.—Trans. Path. Soc.
Ziehl.—Bedeutung der Tuberkelbacillen für Diagnose und Prognose. Deut. Med. Woch., Nr. 5.
Dieulafoy and Krishaber.—Arch. de Physiol. Norm. et Path., T. I.
Malassez and Vignal.—Ibid., T. II. Compt. Rend., T. 97.
Koch.—Kritische Besprechung der gegen die Bedeutung der Tuberkelbacillen gerichteten Publicationen. Deut. Med. Woch., Nr. 10.
Vignal.—Compt. Rend. Soc. de Biol., T. 5.
Gibbes.—Lancet.
Kundrat.—Wien. Med. Presse.
Landouzy and Martin.—Rev. de Méd., T. 3.

- 1883 **Chiari**.—Wien. Med. Presse.
Cochet.—Compt. Rend. Soc. de Biol. Paris. T. 5.
Pfeiffer.—Berlin. Klin. Woch., Bd. 21.
Leube.—Sitzungsber. der Phys.-Med. Soc. zu Erlangen.
Spina.—Studien über Tuberculose. Wien.
Crämer.—Sitzungsber. d. Phys. Med. Soc. zu Erlangen.
Marchand.—Deut. Med. Woch., Nr. 15.
Heron.—Lancet.
Green.—Brit. Med. Journ.
Baumgarten.—Centralbl. f. d. Med. Wissensch. Berlin. Bd. 21.
Baumgarten.—Zeitschr. f. Klin. Med., Bd. 6.
Somari and Brugnatelli.—Redii R. Instit. Lombardo.
- 1884 **Malassez and Vignal**.—Sur le Microorg. de la Tuberculose Zoogloëique.
 Compt. Rend., T. 99, p. 200.
Doutrelepont.—Die Aetiologie des Lupus Vulgaris. Vierteljahrschr. f. Dermatologie u. Syphilis.
Cornil and Leloir.—Recherches, etc., sur la Nature du Lupus. Arch. de Physiol. Norm. et Pathol.
Babès and Cornil.—Note sur les Bacilles de la Tuberculose. Journ. de l'Anat. et de la Physiol. Norm. et Pathol.
Bouley.—La Nature Vivante de la Contagion, Contagiosité de la Tuberculose. Paris.
Déjérine.—Rev. de Méd. Paris. T. 4.
Arloing.—Compt. Rend.
Biedert.—Virchow's Archiv, Bd. 98.
Leyden.—Klinisches über Tuberkelbacillen. Zeitschr. f. Klin. Med., VIII.
Müller.—Ueber den Befund von Tuberkelbacillen bei Fungösen Knochen u. Gelenkaffectionen. Centralbl. f. Chir., 3.
Albrecht.—Arch. f. Kinderheilk., Bd. 5.
Schill and Fischer.—Mitth. a. d. Kaiserl. Ges. Amt, Bd. II.
Lustig.—Ueber Tuberkelbacillen im Blut bei an Allg. Acuter Miliartub. Erkrankten. Wien. Med. Woch., Nr. 48.
Voltolini.—Ueber Tuberkelbacillen im Ohr. Deut. Med. Woch., Nr. 31.
Strassmann.—Virchow's Archiv, Bd. 96.
Koch.—Mittheilungen aus dem Kais. Ges. Amt, Bd. II.
Baumgarten.—Centralbl. f. d. Med. Wissensch. Berlin. Bd. 22.
Gaffky.—Verhalten der Tuberkelbacillen im Sputum. Mitth. a. d. Kaiserl. Gesundheitsamt, Bd. II.
Negri.—Colorations des Spores dans les Bacilles de la Tuberculose. Journ. de Microsc., T. 8.
Sutton.—Trans. Path. Soc. London, Vol. XXXV.
Andrew.—Lancet.
Williams.—Journ. Royal Microscop. Soc.
- 1885 **Doutrelepont**.—Fall von Meningitis Tuberculosa nach Lupus, Tuberkelbacillen im Blut. Deut. Med. Woch., Nr. 7.
Karg.—Centralbl. f. Chir.

- 1885 **Spina.**—Casopis Lekaru Ceskych, Nr. 4.
Obrzut.—Prof. Spina's Neue Färbungsmethode der Fäulnismikroorganismen und ihre Beziehung zu den Tuberkelbacillen. Deut. Med. Woch., Nr. 12.
Creighton.—Brit. Med. Journ.
Creighton.—Lancet.
Fütterer.—Ueber das Vorkommen u. die Vertheilung der Tuberkelbacillen in den Organen. Virch. Arch., Bd. 100, Heft 2.
Johne.—Ein Zweifelh. Fall von Congenitaler Tuberculose. Fortschr. d. Med., Bd. 3, 198.
Biedert.—Berl. Klin. Woch.
Ribbert.—Deut. Med. Woch.
Harris.—St. Barthol. Hosp. Reports.
Sticker.—Centralbl. f. Klin. Med.
Wesener.—Fütterungstuberculose.
Tscherning.—Inoculationstuberculose beim Menschen. Fortschr. d. Med., Bd. 3, 65.
- 1886 **Middeldorpf.**—Fortsch. d. Med.
Müller.—Centralbl. für Chir.
Goldenblum.—Vrach., Nos. I. and XI.
Black.—Lancet.
Baumgarten.—Zeitschrift. f. Klin. Med.
Kirstein.—Deut. Med. Woch.
Max Bender.—Ueber die Beziehungen des Lupus Vulg. zu Tuberc. Deut. Med. Woch.
Harries and Campbell.—Lupus. London.

(XXXIV.) Typhoid Fever.

- 1874 **Birch-Hirschfeld.**—Unters. zur Pathologie des Typhus Abdominalis. Zeitschr. f. Epidemiologie, I.
Klein.—Med. Centralbl., XII.
- 1876 **Letzerich.**—Virchow's Archiv, Bd. 68.
- 1877 **Feltz.**—Compt. Rend., T. 85.
- 1878 **Letzerich.**—Archiv f. Exper. Pathol., Bd. 9.
- 1880 **Eppinger.**—Beitr. zur Pathol. Anatomie aus d. Patholog. Institut Prag.
Fischel.—Beitr. zur Pathol. Anat. aus d. Pathol.-Anat. Inst. Prag.
- 1880 **Tizzoni.**—Studi. di Pat. Sperim. sulla Gen. d. Tifo. Milano.
Klebs.—Arch. f. Exper. Pathol. u. Pharmakol.
Eberth.—Arch. f. Pathol. Anat., Bd. 81.
- 1881 **Letzerich.**—Archiv f. Exper. Pathol., Bd. 10.
Brautlecht.—Pathogene Bakterien im Trinkwasser bei Epidemien. Virchow's Arch., Bd. 84.
Meyer.—Unters. über den Bacillus des Abdominaltyphus. Inaug. Diss.
Rappin.—Contrib. à l'Etude des Bact. de la Bouche, à l'Etat Normal et dans la Fièvre Typhoïde.
- 1882 **Coats.**—Eberth's Typhoid Bacillus. Brit. Med. Journ.

- 1882 **Wernich.**—Zeitschr. f. Klin. Med., Bd. 6.
Maraghano.—Centralbl. f. d. Med. Wiss., Bd. 15.
Almqvist.—Typhöidfeberus-Bakterie.
Crooke.—Brit. Med. Journ.
- 1883 **Boens.**—Acad. Roy. de Méd. de Belgique. Bull., 3 Sér., T. 17.
Eberth.—Der Typhus-Bacillus und die Intestinelle Infection. Volk-
mann, Klin. Vorträge.
Letzerich.—Experimentelle Untersuchungen über die Aetiologie des
Typhus Abdominalis. Leipzig.
- 1884 **Gaffky.**—Zur Aetiologie des Addominaltyphus. Mitth. a. d. Ges. Amt,
Bd. II.
Tayon.—Le Microbe de la Fièvre Typhoïde de l'Homme. Compt. Rend.,
T. 99, p. 331.
- 1885 **Pfeiffer.**—Ueber den Nachweis der Typhusbacillen im Darminhalt u.
Stuhlgang. Deut. Med. Woch., Nr. 29.
- 1886 **Michael.**—Typhusbacillen im Trinkwasser. Fortsch. d. Med.
Meisels.—Wien. Med. Woch.
Seitz.—Bakteriolog. Studien z. Typhusätiologie.
Lucatello.—Bollet. d. R. Accad. Med. di Genova.
Fraenkel and Simmonds.—Die Aetiologische Bedeutung des Typhus-
Bacillus.
Neuhauss.—Berl. Klin. Woch.
Sirotinin.—Zeitschr. f. Hygiene.
Beumer and Peiper.—Zeitschr. f. Hygiene.

(XXXV.) Variola and Vaccinia.

- 1868 **Chauveau.**—Nature du Virus-Vaccin. Compt. Rend.
- 1871 **Weigert.**—Ueber Bakterien in der Pockenheit.
- 1872 **Cohn.**—Virchow's Archiv, Bd. 55.
Zülzer.—Berl. Klin. Wochenschr.
- 1873 **Luginbuhl.**—Der Micrococcus der Variola. Verhdl. d. Phys. Med. Ges.
in Würzburg.
- 1874 **Weigert.**—Anat. Beitr. z. Lehre v. d. Pocken.
Pissin.—Berl. Klin. Wochenschr.
Stropp.—Vaccination u. Mikrokokken.
Klebs.—Arch. f. Exp. Path. u. Pharmakol., Bd. 10.
- 1880 **Ischamer.**—Ueber das Wesen des Contag. der Variola Vaccine u.
Varicella. Aerztl. Verein. Steiermark.
- 1881 **Tappe.**—Aetiologie u. Histologie der Schafpocken.
- 1882 **Pohl-Pincus.**—Vaccination.
Pfeiffer.—Ueber die Rückimpfung auf Kühe und Kälber. Jahrb. f.
Kinderheilk.
Marchand.—Les Virus-Vaccins. Revue Mycologique.
Strauss.—Vaccinal Micrococci. Soc. de Biol. de Paris.
- 1883 **Wolf.**—Zur Impffrage. Berl. Klin. Wochenschr., Nr. 4.
Plaut.—Das Organism. Contagium der Schafpocken.

- 1883 **Cornil and Babès.**—Soc. Médicale des Hôpitaux.
Quist.—St. Petersburg Medic. Woch., Nr. 46.
- 1884 **Hamerink.**—Ueber die sog. Vaccination u. Variola.
 (XXXVI.) Yellow Fever.
- 1883 **Babès.**—Sur les Microbes trouvés dans le Foie et dans le Rein d'Individus morts de la Fièvre Jaune. Compt. Rend., 17 Sept.
- 1884 **Domingos Freire and Rebougeon.**—Compt. Rend., T. 99, p. 804.
Domingos Freire.—Recherches sur la Cause de la Fièvre Jaune.
- 1885 **Bouley.**—L'Inoculation Préventive de la Fièvre Jaune. Compt. Rend., T. 100, p. 1276.
Carmona y Valle.—Leçons sur l'Étiol. et la Prophylax. de la Fièvre Jaune.
- 1886 **Cerecedo.**—El Siglo Medico.

(K.) *BACTERIA IN THE AIR, IN SOIL, AND IN WATER.*

- 1858 **Pouchet.**—Compt. Rend., T. 47.
- 1860 **Pasteur.**—Compt. Rend., T. 50.
- 1861 **Pasteur.**—Compt. Rend., T. 52.
- 1862 **Pasteur.**—Ann. de Chim. et de Phys., T. 64.
- 1863 **Pasteur.**—Compt. Rend., T. 56.
Lemaire.—Compt. Rend., T. 57.
- 1870 **Tyndall.**—Med. Tim. and Gaz.
Maddox.—Month. Microscop. Journal.
- 1874 **Cunningham.**—Micr. Exam. of the Air. Calcutta.
Tissandier.—Compt. Rend., T. 78.
- 1877 **Pasteur.**—Compt. Rend., T. 85.
Miquel.—Annuaire de l'Observat. de Montsouris.
Tyndall.—Brit. Med. Journ.
- 1878 **Miquel.**—Compt. Rend., T. 86.
Miquel.—De la Présence dans l'Air du Ferment de l'Urée. Bull. de la Soc. Chim.
- 1879 **Miflet.**—Unters. über die in der Luft Suspendirten Bakterien. Cohn's Beitr. z. Biol. d. Pflanzen, Bd. III.
Wernich.—Cohn's Beiträge z. Biol. d. Pflanzen, Bd. III., S. 105.
Klebs and Tommasi-Crudeli.—Unters. der Luft auf die Mikroorganismen der Malaria. Archiv f. Exper. Path., Bd. II.
Miquel.—Ann. d'Hygiène.
- 1881 **Koch.**—Milzbrandbacillen im Boden. Mitth. a. d. Ges. Amt, Bd. I.
Soyka.—Sitz.-Ber. der K. Bayr. Akad. d. Wiss. Math. Physik. Classe.
Soyka.—Vorträge im Aerztl. Verein in München.
Büchner.—Die Bedingungen des Uebergangs von Pilzen in die Luft. Vorträge im Aerztl. Verein zu München.
Tyndall.—Essays on the Floating Matter of the Air.
Winnacker.—Ueber die in Rinnsteinen beobacht. Nied. Organismen. Göttinger Inaug.-Diss. Frankfurt a. M.

- 1882 v. Fodor.—Hygienische Unters. über Luft, Boden u. Wasser.
 Nägeli.—Unters. über Niedere Pilze.
 Soyka.—D. Vierteljsch. f. Oeff. Ges., Bd. 14.
 Miquel.—Annuaire de l'Observat. de Montsouris.
- 1883 Wolny.—Ueber die Thätigkeit Niederer Organismen im Boden. Viert.
 f. Oeff. Ges., S. 705.
 Letzerich.—Exp. Unters. üb. die Aetiologie des Typhus mit bes.
 Berücksichtigung der Trink- u. Gebrauchswässer.
 Zander.—Centrabl. f. Allg. Ges.
 Gunning.—Beitr. z. Hygienischen Untersuchung des Wassers, Arch. f.
 Hyg., 3.
 Miquel.—Les Organismes Vivants de l'Atmosphère.
 Olivier.—Les Germes de l'Air. Thèse. Rev. Scientif.
 Angus Smith.—Sanitary Record.
 Smart.—Germs, Dust, and Disease.
 Torelli.—La Malaria in Italia.
- 1884 Hesse.—Ueber Quantitative Best. der in der Luft enthaltenen Mikro-
 organismen. Mitth. a. d. Ges. Amt, Bd. II.
 Hesse.—Ueber Abscheidung d. Mikroorganismen aus d. Luft. Deut-
 sche Med. Wochenschr., 2.
 Hesse.—Weitere Mitth. über Luftfiltration. Deut. Med. Woch., Nr. 51.
 Schrakamp.—Archiv f. Hygiene, Bd. II.
 Miquel and Freudenreich.—La Semaine Médicale.
 Sehlen.—Fortschr. d. Med., Bd. II., S. 585.
 Chamberland.—Sur un Filtre Donnant de l'Eau Physiologiquement
 Pure. Compt. Rend., T. 99, p. 247.
 Moreau and Plantymausion.—La Semaine Médicale.
 Angus Smith.—On the Examination of Waters. Rep. to the Loc. Gov.
 Board.
- 1885 Cramer.—Die Wasserversorgung von Zürich.
 Crookshank.—Notes from a Bact. Labor., Lancet.
 Becker.—Reichsmedicinalkalender.
 Hesse.—Ueber Wasserfiltration. Deut. Med. Woch.
 Soyka.—Prager Med. Woch.
 Frankland.—Removal of Micro-organism from Water. Proc. Roy. Soc.
- 1886 Soyka.—Fortschr. d. Med.
 Laurent.—Journal de Pharmacie et de Chimie.
 Wolffhügel and Riedel.—Arbeit. a. d. K. Ges. Amt.
 Heræus.—Zeitschr. f. Hygiene.
 C. Frankel.—Zeitschr. f. Hygiene.
 Meade Bolton.—Zeitschr. f. Hygiene.
 Hesse.—Zeitschr. f. Hygiene.
 Beumer.—Deut. Med. Woch.
 Frankland.—Proc. Roy. Soc.
 Bischof.—Journ. Soc. Chem. Industry.
 Pfeiffer.—Zeitschr. f. Hygiene.

APPENDIX F.

TABLE SHOWING THE MAGNIFYING POWER OF
ZEISS' OBJECTIVES.

Ocular:	1	2	3	4	5	
a_1	7	11	15	22		a_1
a_2	12	17	24	34		a_2
a_3	20	27	38	52		a_3
a^*		4—12	7—17	10—24		a^*
aa	22	30	41	56	75	aa
A, AA	38	52	71	97	130	A, AA
B, BB	70	95	130	175	235	B, BB
C, CC	120	145	195	270	360	C, CC
D, DD	175	230	320	435	580	D, DD
E	270	355	490	670	890	E
F	405	540	745	1010	1350	F
G	260	340	470	640	855	G
H	320	430	590	805	1075	H
J	430	570	785	1070	1430	J
K	570	760	1045	1425	1900	K
L	770	1030	1415	1930	2570	L
$\frac{1}{8}$	260	340	470	640	855	$\frac{1}{8}$
$\frac{1}{12}$	380	505	695	950	1265	$\frac{1}{12}$
$\frac{1}{18}$	605	810	1110	1515	2020	$\frac{1}{18}$
	1	2	3	4	5	

INDEX.

- A
- Abbé condenser, 23
 Abrin, 345
 Abrus precatorius, 345
 Abscesses, 229
 Acetate of potash, 34
 Acetic acid, 27
 Achorion Schoenleinii, 376
 Actinomyces, 377.
 — methods of staining, 383
 Actinomycosis, 379
 Acute infectious osteomyelitis, coccus
 of, 246
 Acute yellow atrophy, 250
 Æroniscopes, 129
 Ærosopes, 129
 Agar-agar, 41
 — glycerine, 87
 — nutrient, preparation of, 86
 Air, examination of, 126
 Aitken's test-tube, 48, 110
 Alcohol, 25
 — acidulated, 27
 Alopecia areata, 266
 Alum carmine, 27
 Ammonia, 28
 Aniline, 28
 — water, 28
 Animals, examination of, 141
 Anthrax, 315
 Antiseptics, 180
 Ascococcus, 257
 — Billrothii, 257
 Ascomycetes, 376
 Asiatic cholera, 284
 Aspergillus albus, 378
 — clavatus, 378
 — flavus, 377
 — fumigatus, 377
 — glaucus, 376
 — niger, 377
 — ochraceus, 378
 — repens, 376
 Asphalte lac, 34
 Attenuation of virus, 193
- B
- Babes' incubator, 53
 — method, 67
 Babes' of staining comma-bacilli, 291
 — of staining bacillus of leprosy,
 301
 Bacillus, 299
 — acidi lactici, 339
 — aërophilus, 348
 — alvei, 335
 — amylobacter, 351.
 — anthracis, 315
 — butyricus, 351
 — caucasicus, 345
 — cavicola, 273
 — columbarum, 267
 — cuniculicida, 269
 — cyanogenus, 337
 — dysodes, 346
 — erythrosporus, 346
 — figurans, 344
 — Fitzianus, 339
 — fluorescens, 336
 — foetidus, 347
 — Hansenii, 346
 — ianthinus, 337
 — indicus, 274
 — in gangrenous septicæmia, 305
 — in septicæmia of man, 305
 — in swine-erysipelas, 333
 — in swine-typhoid, 333
 — in syphilis, 301
 — in tetanus, 334
 — lepræ, 299
 — liordermos, 278
 — luteus, 276
 — malarie, 304
 — mallei, 325
 — megaterium, 343
 — mesentericus fuscus, 348
 — — vulgatus, 348
 — multipediculus, 278
 — mycoides, 345
 — Neapolitanus, 263
 — cedematis maligni, 327
 — of blue milk, 337
 — of choleraic diarrhoea from meat-
 poisoning, 304
 — of diphtheria, 265
 — of glanders, 325
 — of jequirity, 345
 — of pneumo-enteritis of the pig, 333

- Bacillus of rhinoscleroma, 264
 — of septicæmia of mice, 330
 — of splenic fever, 315
 — of ulcerative stomatitis in the calf, 332
 — oxytocus perniciosus, 273
 — parvus ovatus, 252
 — phosphorescens, 348
 — pneumonicus agilis, 272
 — prodigiosus, 275
 — pseudo-pneumonicus, 263
 — putrificus coli, 347
 — pyocyaneus, 336
 — pyogenes fœtidus, 305
 — ramosus liquefaciens, 278
 — saprogenes, No. 1, 347
 — — No. 2, 347
 — — No. 3, 266
 — — fœtidus, 347
 — septicus, 346
 — — agrigenus, 269
 — — sputigenus, 271
 — subtilis, 340
 — tremulus, 345
 — tuberculosis, 305
 — tumescens, 342
 — typhosus, 302
 — ulna, 342
 — ureæ, 276
 — violaceus, 337
 — virens, 147
 — vitulorum, 266
 Bacteria, chromogenic, 172
 — classification of, 205
 — distribution of, 178
 — general morphology of, 152
 — pathogenic, 174
 — saprogenic, 173
 — zymogenic, 172
 Bacteriaceæ, 215
 Bactériedie du charbon, 315
 Bacteridium cyaneum, 253
 Bacterium, 260
 — aceti, 277
 — æruginosum, 336
 — brunneum, 276
 — cavicida, 273
 — chlorinum, 147
 — cholerae gallinarum, 267
 — coli commune, 273
 — crassum sputigenum, 272
 — decalvans, 266
 — fluorescens liquefaciens, 276
 — — putidum, 276
 — fœtidum, 347
 — fusiforme, 285
 — hyacinthi, 274
 — ianthinum, 337
 Bacterium indicum, 274
 — in diphtheria of calves, 266
 — — of man, 265
 — lactis aerogenes, 273
 — lineola, 282
 — liodermos, 278
 — litoreum, 280
 — luteum, 276
 — merismopedioides, 279
 — multipediculum, 278
 — navicula, 280
 — Neapolitanum, 263
 — of Davaine's septicæmia, 270
 — of diphtheria of pigeons, 267
 — of fowl-cholera, 267
 — of rhinoscleroma, 264
 — of septicæmia in rabbits, 269
 — of yellow milk, 274
 — oxytocum perniciosum, 273
 — Pasteuri, 271
 — Pasteurianum, 277
 — Pflügeri, 280
 — photometricum, 280
 — pneumoniæ cruposæ, 261
 — pneumonicum agile, 272
 — prodigiosum, 275
 — pseudo-pneumonicum, 263
 — ramosum liquefaciens, 278
 — rubescens, 357
 — rubrum, 275
 — saprogenes, 266
 — septicum agrigenum, 269
 — — sputigenum, 271
 — syncyanum, 337
 — synxanthum, 274
 — termo, 281
 — ureæ, 276
 — violaceum, 276
 — viride, 147
 — xanthinum, 274
 — Zoppii, 278
 Balance and weights, 40
 Balmer-Frëntzel method, 312
 Baumgarten's method, 311
 — new method, 312
 Bees, bacillus in disease of, 335
 Beggiatoa, 356
 — methods of examining species of, 356
 — alba, 356
 — mirabilis, 356
 — roseo-persicana, 357
 Bergamot oil, 25
 Bière malade, 256
 Bismarck-brown, 28
 Blackleg, 350
 Black torula, 371
 Bleeding host, 275

- Blood rain, 275
 — serum, liquid, 107
 — — sterile, 104
 Blue milk, bacillus of, 337
 Borax carmine, 28
 Botrytis Bassiana, 379
 Bouillon, 106
 Bread-paste, 103
 Brush, 46
 Bulbed tubes, 49
 Butyric acid fermentation, bacillus of, 351
- C
- Camera-lucida, 34
 Canada-balsam, 34
 Caoutchouc caps, 42
 Carbonate of soda, 41
 Carious teeth, 361
 Cattle plague, 236
 Cedar oil, 28
 Cell-contents, 149
 Cell-wall, 148
 Celloidin, 25, 74
 Cerebro-spinal meningitis, 238
 Charbon symptomatique, 350
 Chemical composition, 147
 — disinfectants, 183
 Cheshire's trough, 89
 Chionyphe Carteri, 379
 Cholera, comma-bacillus of, 284
 — fowl, 267
 — nostras, comma-bacillus in, 291
 Choleraic diarrhoea from meat-poisoning, bacillus of, 304
 Cladothrix, 362
 — dichotoma, 362
 — Fœrsteri, 364
 Cladotricheæ, 216
 Classification, 205
 — Flügge's, 210
 — Hueppe's, 217
 — Zopf's, 215
 Clostridium, 350
 — butyricum, 351
 — of symptomatic anthrax, 350
 — polymyxa, 353
 Coccaceæ, 215
 Cocci, methods of staining, 258
 Cohn-Mayer fluid, 108
 Cohnia roseo-persicina, 357
 Collection of water samples, 133
 Comma-bacillus of Finkler, 291
 — of Koch, 284
 — in cholera nostras, 291
 Cork, 25
 Cornil and Alvarez, method of, 265
 Cotton-wool, 40
- Cover-glass impressions, 69
 — preparations, 65
 — — double coloration, 67
 Crenothrix, 354
 — Kühniana, 354
 Cutaneous inoculation, 138
 Cutting tissues, 72
- D
- Damp chambers, 26, 45, 92
 D'Arsonval's incubator, 49
 Decalcifying preparations, 71
 Dental caries, 361
 Desiccator, 60
 Diphtheria, 265, 266
 Diplococcus albicans tardissimus, 242
 Disinfectants, 180
 Dissecting boards, 58
 — case, 58
 Dissection, 141
 Distribution of bacteria, 178
 Double coloration, 67
 Double-stain spore-bearing bacilli, 68
 Double-staining, 67
 Dressing-case, 58
 Drinking water, 131
 Drop-cultures, 112
 Drop-culture slides, 49
- E
- Ebner's solution, 25
 Ehrlich's method, 67
 — and eosin, 314
 Electricity, application of, 121
 Embedding tissues, 72
 Empusa muscæ, 372
 — radicans, 373
 Endocarditis ulcerosa, 236
 Eosin, 28
 Erysipelas, 230
 — malignum, bacillus des, 333
 Ether, 29
 Eurotium aspergillus glaucus, 376
 — aspergillus niger, 377
 — repens, 376
 Examination of plate-cultivations, 96
 — of test-tube cultivations, 89
 Experiments on living animals, 137
- F
- Farrant's solution, 34
 Favus, 376
 Filter, making, 85
 — paper, 41
 Fire blight, 253

Flagella, to stain, 69
 Flagellated protozoa in blood, 397
 Flagellum, 157
 Flannel, 41
 Folded filter, 85
 Foot-and-mouth disease, 235
 Form, 152
 Foulbrood, 335
 Fowl-cholera, bacterium of, 267
 Fränkel's method, 313
 Friedländer, method of, 262
 Frogspawn fungus, 296
 Fuchsine, 29

G

Gangrene, 250
 Gas burners, 52
 — chambers, 120
 — pressure regulator, Moitessier's, 54
 Gases, effect of, 170
 Gelatine, 26, 40
 — nutrient, preparation of, 82
 — plates, 94
 Gelatinous envelope, 150
 Gentian-violet, 29
 Gibbes' first method, 311
 — magenta solution, 29
 — new method, 311
 — solution for double-staining, 29
 Glanders, bacillus of, 325
 Glass bells, 43
 — benches, 44
 — capsules, 48
 — dishes, 43, 45
 — jar, 46
 — plates, 44
 — rods, 44
 — vessels, 39
 Glycerine agar-agar, 87
 — gelatine, 26
 — gum, 34
 — pure, 29
 Gomme de sucrerie, 296
 Gonorrhœa, 241
 Gram's method, 76
 — solution, 30
 Growth, circumstances affecting, 169
 — products of, 171

H

Hæmatomonas carassii, 400
 — cobitis, 399
 — Evansi, 404
 Hæmatozoa of malaria, 408
 Hæmatoxylin solution, 30
 Hæmophilia neonatorum, 249
 Hardening preparations, 71

Hay-bacillus, 340
 — methods of staining, 342
 Heat regulator (Meyer's), 57
 — (Reichert's), 55
 — (Schlösing's), 51
 Herpes tonsurans, 376
 Hesse's apparatus, 127
 His' method, 68
 Hollis' glue, 34
 Hot air and steam, 189
 Hot-air steriliser, 37
 Hot-water filter, 39
 Hydrophobia, 250
 Hyphomycetes, 371
 Hypodermii, 372

I

Immunity, 192
 Impression-preparations, 69
 Incubators, 49
 — Babes', 53
 — D'Arsonval's, 49
 Inoculation of test-tubes, 88
 — cutaneous and subcutaneous, 138
 — of potatoes, 101
 — protective, 193
 Intermittent fever, bacillus of, 304
 Iodine solution, 30
 — (Gram), 30
 Iron box, 44
 Isolation of micro-organisms, 90
 Israel's case, 44
 — warming apparatus, 118

J

Japanese isinglass, 41
 Jequirity, bacillus of, 345

K

Kaatzner's method, 314
 Kephir, 345
 Kleinenberg's solution, 26
 Koch's apparatus for examination of
 air, 127
 — method for staining comma-
 bacilli, 290
 — original method for staining the
 tubercle-bacillus, 309
 — postulates, 17
 — solution (methylene-blue), 31
 — solution (methyl-violet), 31

I

Laboratory requisites, 23
 Lactic acid, 41

Leprosy, 299
 Leptothrix, 361
 — buccalis, 361
 — gigantea, 362
 — ochracea, 362
 — parasitica, 362
 Leptotricheæ, 215
 Leuconostoc, 296
 — mesenteroides, 296
 Leukæmia, 239
 Levelling apparatus, 91
 Lichtheim's method, 313
 Light, 171
 Liquid media, 106
 Lister's flasks, 48, 109
 Lithium-carminic solution, 30
 Litmus papers, 41
 Lochial discharges, 242
 Löffler's solution, 31
 Lustgarten, method of, 301

M

Madura-foot, 379
 Magenta solution (Gibbes'), 30
 Malaria, 304
 Malignant œdema, bacillus of, 327
 — pustule, 315
 Marsh-spirochæte, 294
 Measles, 249
 Merismopedia, 241
 — gonorrhœa, 241
 Methylene-blue, 30
 Methyl-violet, 31
 Metschnikoff's theory, 201
 Meyer's thermo-regulator, 57
 Micrococcus, 245
 — albicans amplius, 242
 — amyliworus, 253
 — aurantiacus, 253
 — bombycis, 253
 — candicans, 254
 — candidus, 254
 — cereus albus, 248
 — — flavus, 248
 — chlorinus, 254
 — cholerae gallinarum, 267
 — cinnabareus, 255
 — citreus conglomeratus, 242
 — coronatus, 256
 — crepusculum, 255
 — cyaneus, 253
 — flavus desidens, 256
 — — liquefaciens, 255
 — — tardigradus, 255
 — foetidus, 255
 — fulvus, 256

Micrococcus hæmatodes, 254
 — indicus, 274
 — in acute yellow atrophy, 249
 — in gangrene, 249
 — in hæmophilia neonatorum, 249
 — in measles, 249
 — in rabies, 249
 — in scarlatina, 249
 — insectorum, 253
 — in typhus, 249
 — in whooping-cough, 249
 — lacteus faviformis, 256
 — luteus, 254
 — of progressive suppuration in rabbits, 251
 — of pyæmia in mice, 252
 — of pyæmia in rabbits, 251
 — of septicæmia in rabbits, 250
 — parvus ovatus, 252
 — perniciosus, 252
 — prodigiosus, 275
 — pyocyaneus, 336
 — pyogenes albus, 247
 — — aureus, 246
 — — citreus, 247
 — — tenuis, 249
 — radiatus, 256
 — rosaceus, 254
 — subflavus, 242
 — tetragonus, 241
 — ureæ, 276
 — versicolor, 255
 — violaceus, 254
 — viscosus, 256
 — viticulosus, 255
 Micro-organisms in the blood, 397
 — inhalation of, 137
 — isolation of, 90
 Micro-photographic apparatus, 35
 Microscope, 23
 Microsporon furfur, 376
 Microtome, 24
 — freezing, 25
 Microzyma bombycis, 253
 Microzyme test, 131
 Milk, 107
 Miquel's bulbs, 111
 Moist chambers, 113
 Moitessier's gas-pressure regulator, 54
 Monas crepusculum, 255
 — hæmorrhagicum, 249
 — Okenii, 359
 — vinosa, 359
 — Warmingii, 360
 Morphology, 147
 Moulds, 369
 Mouse-cages, 58
 Movement, 154

Mucor aspergillus, 375
 — *corymbifer*, 375
 — *fusiger*, 375
 — *macrocarpus*, 375
 — *melittophorus*, 375
 — *mucedo*, 374
 — *phycomyces*, 375
 — *racemosus*, 374
 — *rhizopodiformis*, 375
 — *stolonifer*, 375
 Müller's fluid, 26
Mycoderma cerevisiæ et vini, 370
Myconostoc gregarium, 365

N

Neelsen's method, 312
 — solution, 31
 Nicati and Rietsch's method, 290
 Nitric acid, 31
Nosema bombycis, 273
 Nutrient agar-agar, 86
 — gelatine, 82
 — media, 80
 Nutrition, 167

O

Oedema, 327
Oidium albicans, 376
 — *lactis*, 376
 — Tuckeri, 376
Ophidomonas sanguinea, 359
 Orseille (Wedl), 31
 Orth's modification of Ehrlich's
 method, 311
 — solution, 32
 Osmic acid, 26
 Osteomyelitis, 246

P

Pagè's thermo-regulator, 56
 Panhistophyton ovatum, 273
 Paper trays, 26
 Paraffine, 26
 Paste-cultivations, 102
 Pasteur's apparatus, 111
 — fluid, 108
 — septicæmia, 327
 Peach-coloured bacterium, 357
 Pébrine, 273
Penicillium glaucum, 378
Peptonum siccum, 41
 Perlsucht, 308
Peronospora infestans, 373
 Peters' method, 313
 Pfuhl-Petri's method, 314

Phagocytes, 175, 201
 Phosphorescence, 280
 Photo-micrographic apparatus, 35
Phragmidiothrix, 360
 — *multiseptata*, 360
 Phycomycetes, 373
 Picric acid, 31
 Picro-carmin (Ranvier), 31
 — lithium-carmin (Orth), 31
Pilobolus, 374
 Pink torula, 371
 Pityriasis versicolor, 376
 Plate-cultivations, 90
 Platinum needles, 42
 Plaut's method, 385
 Pleomorphism, 212
Pneumococcus, 261
 Pneumo-enteritis of the pig, 333
 Pneumonia, 261
 — method of staining bacteria of,
 262
 Potash solution, 31
Potato-bacillus, 348
 — bacterium, 278
 — cultivations, 99
 Potatoes, disease of, 374
 Potato-knives, 45
 — paste, 102
 Pouchet's æroscope, 129
 Pravaz' syringe (Koch's modification,
 58
 Products of growth, 171
 Protected burners, 52
 Protective inoculation, 193
Proteus mirabilis, 281
 — *vulgaris*, 280
 — Zenkeri, 281
 Ptomaines, 174
 Puerperal fever, 231
 Pus, 229, 246, 247
 Putrefaction, 280
 Pyæmia, 229
 — in mice, 252
 — in rabbits, 251
 Pyocyanin, 336

Q

Quarter-evil, 350

R

Rabies, 249
 Ranvier's picro-carmin, 32
 Rauschbrand, 350
 Ray-fungus, 380

Reichert's thermo-regulator, 55
 Relapsing fever, spirillum of, 283
 Reproduction, 158
 Respiration, 166
Rhabdomonas rosea, 359
 Rhinoscleroma, 264
 Rindfleisch's method, 310
 Rothlauf, 333
 Rouget du porc, 333

S

Saccharomyces albicans, 370
 — *apiculatus*, 370
 — *cerevisiæ*, 370
 — *conglomeratus*, 370
 — *ellipsoideus*, 370
 — *exiguus*, 370
 — *glutinis*, 371
 — *mycoderma*, 370
 — *niger*, 371
 — *pastorianus*, 370
 — *rosaceus*, 371
 — *sphaericus*, 370
Saccharomyces, 369
 Safety-burner, Koch's, 53
 Safranine, 33
 Salmon, disease of, 373
 Salt solution, 33
Saprolegnia, 373
Sarcina, 243
 — *alba*, 245
 — *aurantiaca*, 244
 — *hyalina*, 244
 — *intestinalis*, 244
 — *litoralis*, 244
 — *lutea*, 243
 — *Reitenbachii*, 244
 — *urinæ*, 244
 — *ventriculi*, 244
 Scalpels, 46
 Scarletina, 232
 Schäfer's warm stage, 117
Schizomyces, 224
 Schlösing's membrane regulator, 51
 Schütz, method of, 325
 Senkewitsch's method, 314
 Septicæmia, bacillus of gangrenous,
 305
 — in rabbits, 269
 — of man, 305
 — of mice, 330
 Serum inspissator, 47
 — steriliser, 46
 Silkworms, 273
 Siphon apparatus, 59
 Soil, examination of, 130

Spermacti, 27
Sphærotilus natans, 364
Spirillum, 283
 — *attenuatum*, 296
 — *cholerae Asiaticæ*, 284
 — *Finklerii*, 291
 — *leucomelaneum*, 296
 — *Obermeieri*, 283
 — of relapsing fever, 283
 — *plicatile*, 294
 — *rosaceum*, 359
 — *Rosenbergii*, 296
 — *rubrum*, 294
 — *sanguineum*, 359
 — *serpens*, 295
 — *sputigenum*, 292
 — *tenue*, 295
 — *tyrogenum*, 293
 — *undula*, 295
 — *violaceum*, 359
 — *volutans*, 295
 Spirit-level, 43
Spirochæte buccalis, 361
 — *Obermeieri*, 283
Spiromonas Cohnii, 365
 — *volubilis*, 365
 Splenic fever, 315
 Spores, method of staining, 68
 — preservation of, 320
 Sputum, 65
 — *septicæmia*, 271
 Staining bacteria, methods of, 75
 — *actinomyces*, 384
 — — *bacilli of ulcerative stomati-*
tis, 332
 — — *bacillus anthracis*, 315
 — — — of butyric acid fer-
mentation, 351
 — — — of glanders, 325
 — — — of leprosy, 399
 — — — of rhinoscleroma, 264
 — — — of syphilis, 301
 — — — of typhoid fever, 302
 — — *bacteria in diphtheria*, 265
 — — — of pneumonia, 261
 — — — *comma-bacilli*, 290
 — — — *hay-bacillus*, 340
 — — — *leptothrix buccalis*, 361
 — — — *spirillum Obermeieri*, 283
 — — — *tubercle-bacilli*, 309
Staphylococcus cereus albus, 248
 — *cereus flavus*, 248
 — *pyogenes albus*, 247
 — — *aureus*, 246
 — — *citreus*, 247
 Steam, steriliser, 36
 Sterilisation apparatus, 36
 Sternberg's bulbs, 48, 109

Streptococcus, 224
 — articularum, 231
 — bombycis, 253
 — coronatus, 256
 — erysipelatosus, 230
 — flavus desidens, 256
 — in abscesses, 229
 — in bilious fever, 236
 — in cerebro-spinal meningitis, 238
 — in contagious mammitis, 236
 — in diphtheria, 231
 — in endocarditis, 236
 — in foot-and-mouth disease, 235
 — in measles, 236
 — in pneumonia, 237
 — in puerperal fever, 231
 — in scarlet fever, 232
 — insectorum, 253
 — in small-pox, 235
 — in spreading gangrene, 231
 — in strangles, 236
 — in surgical fever, 231
 — in typhoid fever, 237
 — in yellow fever, 236
 — of cattle plague, 236
 — of progressive tissue necrosis in mice, 238
 — of septicæmia consecutive to anthrax, 238
 — perniciosus, 252
 — pyogenes, 225
 — viscosus, 256
 Streptothrix Fœrsteri, 364
 Stricker's warm stage, 118
 Sulphuric acid, 33
 Surra, 402
 Swine-erysipelas, 333
 — typhoid, 333
 Symptomatic anthrax, 350
 Syphilis, 301

T

Table salt, 41
 Tarichium megaspermum, 373
 Teeth, carious, 361
 Test-tube cultivations, 87
 — stands, 42
 — water-bath, 39
 Tetanus, 334
 Thermo-regulator, 57
 Tilletia caries, 372
 Torula cerevisiæ, 369
 Trichomonas sanguinis, 407
 Tricophyton tonsurans, 376
 Tripod levelling-stand, 43

Tuberculosis, bacillus of, 305
 Turpentine, 35
 Typhoid fever, 302
 Typhus, 249

U

Ulcerative stomatitis in the calf, 332
 Urine, 107
 Urocystis occulta, 372
 Ustilago carbo, 372

V

Vaccinia, 235
 Variola, 235
 Vaseline, 49
 Vegetables, 103
 Vegetable infusions, 108
 Vesuvin, 33
 Vibrio, 349
 — regula, 349
 — serpens, 295

W

Warm stages, 115
 Warming apparatus, 115
 Water, 33
 — bath, 38
 — examination of, 131
 Wedl's orseille, 32
 Weigert-Ehrlich method, 78
 Weigert's method, 76
 Wens, 380
 Whooping-cough, 249
 Wire cages, 41
 Wooden-tongue, 380
 Wool-sorter's disease, 31
 Wurzel-bacillus, 345

X

Xylol, 27

Y

Yeast fungi, 369
 Yeasts, 369
 Yellow fever, 236

Z

Ziehl's method, 313
 Zinc-white, 34

February, 1892.

CATALOGUE OF WORKS

PUBLISHED BY

H. K. LEWIS

136 GOWER STREET, LONDON, W.C.

Established 1844.

SIR WILLIAM AITKEN, KNT., M.D., F.R.S.

Professor of Pathology in the Army Medical School.

ON THE ANIMAL ALKALOIDS, THE PTOMAINES, LEUCOMAINES, AND EXTRACTIVES IN THEIR PATHOLOGICAL RELATIONS. Second edition, Crown 8vo, 3s. 6d.

H. ALDER-SMITH, M.B. LOND., F.R.C.S.

Resident Medical Officer, Christ's Hospital, London.

RINGWORM: Its Diagnosis and Treatment.

Third Edition, enlarged, with Illustrations, fcap. 8vo, 5s. 6d.

E. CRESSWELL BABER, M.B. LOND.

Surgeon to the Brighton and Sussex Throat and Ear Dispensary.

A GUIDE TO THE EXAMINATION OF THE NOSE, WITH REMARKS ON THE DIAGNOSIS OF DISEASES OF THE NASAL CAVITIES. With Illustrations, small 8vo, 5s. 6d.

JAMES B. BALL, M.D. LOND., M.R.C.P.

Physician to the Department for Diseases of the Throat and Nose, and Senior Assistant Physician, West London Hospital.

I.
A HANDBOOK OF DISEASES OF THE NOSE AND NASO-PHARYNX. Large post 8vo, with Illustrations, 6s.

II.
INTUBATION OF THE LARYNX. With Illustrations, demy 8vo, 2s. 6d.

G. GRANVILLE BANTOCK, M.D., F.R.C.S. EDIN.

Surgeon to the Samaritan Free Hospital for Women and Children.

I.
ON THE USE AND ABUSE OF PESSARIES. Second Edition, with Illustrations, 8vo, 5s.

II.
ON THE TREATMENT OF RUPTURE OF THE FEMALE PERINEUM IMMEDIATE AND REMOTE. Second Edition, with Illustrations, 8vo, 3s. 6d.

III.
A PLEA FOR EARLY OVARIOTOMY. Demy 8vo, 2s.

ARTHUR E. J. BARKER, F.R.C.S.

Hunterian Professor of Surgery and Pathology; Surgeon to University College Hospital.

HUNTERIAN LECTURES ON INTRA-CRANIAL INFLAMMATIONS STARTING IN THE TEMPORAL BONE, THEIR COMPLICATIONS AND TREATMENT. 8vo, 3s. nett.

FANCOURT BARNES, M.D., M.R.C.P.

Physician to the Chelsea Hospital for Women; Obstetric Physician to the Great Northern Hospital, &c.

A GERMAN-ENGLISH DICTIONARY OF WORDS AND TERMS USED IN MEDICINE AND ITS COGNATE SCIENCES.
Square 12mo, Roxburgh binding, 9s.

JAMES BARR, M.D.

Physician to the Northern Hospital, Liverpool; Medical Officer of Her Majesty's Prison, Kirkdale, &c.

THE TREATMENT OF TYPHOID FEVER, and reports of fifty-five consecutive cases with only one death. With Introduction by W. T. GAIRDNER, M.D., LL.D., Professor of Medicine in the University of Glasgow. With Illustrations, demy 8vo, 6s. [Now ready.]

ASHLEY W. BARRETT, M.B. LOND., M.R.C.S., L.D.S.E.

Dental Surgeon to, and Lecturer on Dental Surgery in the Medical School of, the London Hospital.

DENTAL SURGERY FOR MEDICAL PRACTITIONERS AND STUDENTS OF MEDICINE. Second edition, With Illustrations, cr. 8vo, 3s. 6d. [Now ready.]

[LEWIS'S PRACTICAL SERIES.]

ROBERTS BARTHOLOW, M.A., M.D., LL.D.

Professor of Materia Medica and Therapeutics in the Jefferson Medical College of Philadelphia, &c., &c.

I.
A PRACTICAL TREATISE ON MATERIA MEDICA AND THERAPEUTICS. Seventh Edition, Revised and Enlarged, 8vo, 18s. [Just published.]

II.
A TREATISE ON THE PRACTICE OF MEDICINE, FOR THE USE OF STUDENTS AND PRACTITIONERS. Fifth Edition, with Illustrations, large 8vo, 21s.

H. CHARLTON BASTIAN, M.A., M.D., F.R.S.

Examiner in Medicine at the Royal College of Physicians; Professor of the Principles and Practice of Medicine in University College, London; Physician to University College Hospital, &c.

PARALYSES: CEREBRAL, BULBAR, AND SPINAL.
A MANUAL OF DIAGNOSIS FOR STUDENTS AND PRACTITIONERS. With numerous Illustrations, 8vo, 12s. 6d.

GEO. M. BEARD, A.M., M.D.

AND

A. D. ROCKWELL, A.M., M.D.

Formerly Professor of Electro-Therapeutics in the New York Post Graduate Medical School; Fellow of the New York Academy of Medicine, &c.

I.
ON THE MEDICAL AND SURGICAL USES OF ELECTRICITY. Eighth Edition. With over 200 Illustrations, roy. 8vo, 28s. [Just published.]

II.
NERVOUS EXHAUSTION (NEURASTHENIA) ITS HYGIENE, CAUSES, SYMPTOMS AND TREATMENT. Second edition, 8vo, 7s. 6d.

W. M. BEAUMONT.

Surgeon to the Bath Eye Infirmary.

THE SHADOW-TEST IN THE DIAGNOSIS AND ESTIMATION OF AMETROPIA. Post 8vo, 2s. 6d.

E. H. BENNETT, M.D., F.R.C.S.I.

Professor of Surgery, University of Dublin,

AND

D. J. CUNNINGHAM, M.D., F.R.C.S.I.

Professor of Anatomy and Chirurgery, University of Dublin.

THE SECTIONAL ANATOMY OF CONGENITAL CÆCAL HERNIA. With coloured plates, sm. folio, 5s. 6d.

A. HUGHES BENNETT, M.D., M.R.C.P.

Physician to the Hospital for Epilepsy and Paralysis, Regent's Park, and Assistant Physician to the Westminster Hospital.

I.

A PRACTICAL TREATISE ON ELECTRO-DIAGNOSIS IN DISEASES OF THE NERVOUS-SYSTEM. With Illustrations, 8vo, 8s. 6d.

II.

ILLUSTRATIONS OF THE SUPERFICIAL NERVES AND MUSCLES, WITH THEIR MOTOR POINTS; a knowledge of which is essential in the Art of Electro-Diagnosis. 8vo, cloth, 2s.

HORATIO R. BIGELOW, M.D.

Permanent Member of the American Medical Association; Fellow of the British Gynaecological Society, &c.

I.

GYNÆCOLOGICAL ELECTRO-THERAPEUTICS. With an Introduction by Dr. GEORGES APOSTOLI. With Illustrations, demy 8vo, 8s. 6d.

II.

PLAIN TALKS ON ELECTRICITY AND BATTERIES WITH THERAPEUTIC INDEX, FOR GENERAL PRACTITIONERS AND STUDENTS OF MEDICINE. Crown 8vo, with Illustrations, 4s. 6d.

DR. THEODOR BILLROTH.

Professor of Surgery in Vienna.

GENERAL SURGICAL PATHOLOGY AND THERAPEUTICS. With additions by Dr. ALEXANDER VON WINIWARTER, Professor of Surgery in Luttich. Translated from the Fourth German edition, and revised from the Tenth edition, by C. E. HACKLEY, A.M., M.D. 8vo, 18s.

DRS. BOURNEVILLE AND BRICON.

MANUAL OF HYPODERMIC MEDICATION.

Translated from the Second Edition, and Edited, with Therapeutic Index of Diseases, by ANDREW S. CURRIE, M.D. Edin., &c. With Illustrations, crown 8vo, 6s.

RUBERT BOYCE, M.B., M.R.C.S.

Assistant in the Pathological Laboratory of University College, London.

A TEXT-BOOK OF MORBID HISTOLOGY. With coloured Plates, royal 8vo. [Nearly ready.]

GURDON BUCK, M.D.

CONTRIBUTIONS TO REPARATIVE SURGERY:
Showing its Application to the Treatment of Deformities, produced by Destructive Disease or Injury; Congenital Defects from Arrest or Excess of Development; and Cicatricial Contractions from Burns. Large 8vo, 9s.

MARY BULLAR & J. F. BULLAR, M.B. CANTAB., F.R.C.S.
RECEIPTS FOR FLUID FOODS. 16mo, 1s.

STEPHEN SMITH BURT, M.D.

Professor of Clinical Medicine and Physical Diagnosis in the New York Post-graduate School and Hospital.

EXPLORATION OF THE CHEST IN HEALTH AND DISEASE. With Illustrations, crown 8vo, 6s.

DUDLEY W. BUXTON, M.D., B.S., M.R.C.P.

Administrator of Anæsthetics at University College Hospital and the Hospital for Women, Soho Square.

ANÆSTHETICS THEIR USES AND ADMINISTRATION. Second edition, with illustrations, crown 8vo. [*In the press.*]
[LEWIS'S PRACTICAL SERIES.]

HARRY CAMPBELL, M.D., B.S. LOND., M.R.C.P.

Assistant Physician and Pathologist to the North-West London Hospital.

I.
THE CAUSATION OF DISEASE: An exposition of the ultimate factors which induce it. Demy 8vo, 12s. 6d.

II.
FLUSHING AND MORBID BLUSHING: THEIR PATHOLOGY AND TREATMENT. With plates and wood engravings, royal 8vo, 10s. 6d. [*Now ready.*]

III.
DIFFERENCES IN THE NERVOUS ORGANISATION OF MAN AND WOMAN, PHYSIOLOGICAL AND PATHOLOGICAL. Royal 8vo, 15s. [*Now ready.*]

R. E. CARRINGTON, M.D., F.R.C.P.

Late Assistant Physician and Senior Pathologist to Guy's Hospital.

NOTES ON PATHOLOGY. With an Introductory Chapter by JAMES FREDERICK GOODHART, M.D. (ABERD.), F.R.C.P., Physician to Guy's Hospital, and Lecturer on Pathology in its Medical School. Edited, revised and amplified by H. EVELYN CROOK, M.D. (LOND.), F.R.C.S. (ENG.), and GUY MACKESON. Crown 8vo. [*Just ready.*]

ALFRED H. CARTER, M.D. LOND.

Fellow of the Royal College of Physicians; Physician to the Queen's Hospital, Birmingham; late Examiner in Medicine for the University of Aberdeen, &c.

ELEMENTS OF PRACTICAL MEDICINE. Sixth Edition, crown 8vo, 9s. [*Just published.*]

P. CAZEAUX.

Adjunct Professor in the Faculty of Medicine of Paris, &c.

AND

S. TARNIER.

Professor of Obstetrics in the Faculty of Medicine of Paris.

OBSTETRICS: THE THEORY AND PRACTICE; including the Diseases of Pregnancy and Parturition, Obstetrical Operations, &c. Seventh Edition, edited and revised by ROBERT J. HESS, M.D., with twelve full-page plates, five being coloured, and 165 wood-engravings, 1081 pages, roy. 8vo, 35s.

WAYLAND C. CHAFFEY, M.D. LOND.

Physician to the Royal Alexandra Hospital for Sick Children, Brighton.

LYMPH-STASIS, OR RETARDATION OF LYMPH, AS AN ELEMENT IN THE CAUSATION OF DISEASE; Especially in regard to Scrofula and Tuberculosis. 8vo, 3s.

F. H. CHAMPNEYS, M.A., M.D. OXON., F.R.C.P.

Physician-Accoucheur and Lecturer on Obstetric Medicine at St. Bartholomew's Hospital; Examiner in Obstetric Medicine in the University of Oxford, and in the Royal College of Physicians, London, &c.

I.

LECTURES ON PAINFUL MENSTRUATION. THE HARVEIAN LECTURES, 1890. Roy. 8vo, 7s. 6d. [*Just published.*]

II.

EXPERIMENTAL RESEARCHES IN ARTIFICIAL RESPIRATION IN STILLBORN CHILDREN, AND ALLIED SUBJECTS. Crown 8vo, 3s. 6d.

W. BRUCE CLARKE, M.A., M.B. OXON., F.R.C.S.

Assistant Surgeon to, and Senior Demonstrator of Anatomy and Operative Surgery at, St. Bartholomew's Hospital; Surgeon to the West London Hospital; Examiner in Surgery to the University of Oxford.

THE DIAGNOSIS AND TREATMENT OF DISEASES OF THE KIDNEY AMENABLE TO DIRECT SURGICAL INTERFERENCE. Demy 8vo, with Illustrations, 7s. 6d.

JOHN COCKLE, M.A., M.D.

Physician to the Royal Free Hospital.

ON INTRA-THORACIC CANCER. 8vo, 4s. 6d.

ALEXANDER COLLIE, M.D. ABERD., M.R.C.P. LOND.

Secretary of the Epidemiological Society for Germany and Russia, &c.

ON FEVERS: THEIR HISTORY, ETIOLOGY, DIAGNOSIS, PROGNOSIS, AND TREATMENT. Illustrated with Coloured Plates, crown 8vo, 8s. 6d. [LEWIS'S PRACTICAL SERIES.]

M. P. MAYO COLLIER, M.B., M.S. LOND., F.R.C.S. ENG.

Professor of Comparative Anatomy and Physiology at the Royal College of Surgeons, England, &c.

THE PHYSIOLOGY OF THE VASCULAR SYSTEM. Illustrations, 8vo, 3s. 6d.

WALTER S. COLMAN, M.B., M.R.C.P. LOND.

*Pathologist and Registrar to the National Hospital for the Paralysed and Epileptic;
Formerly Assistant to the Professor of Pathology in the University of Edinburgh.*

SECTION CUTTING AND STAINING: A Practical
Guide to the Preparation of Normal and Morbid Histological Specimens.
Crown 8vo, 3s. [Now ready.]

W. H. CORFIELD, M.A., M.D. OXON.

Professor of Hygiene and Public Health in University College, London.

DWELLING HOUSES: their Sanitary Construction and
Arrangements. Third Edition, with Illustrations. Crown 8vo.
[In preparation.]

J. LEONARD CORNING, M.A., M.D.

Consultant in Nervous Diseases to St. Francis Hospital.

A PRACTICAL TREATISE ON HEADACHE, NEU-
RALGIA, SLEEP AND ITS DERANGEMENTS, AND SPINAL
IRRITATION. With an Appendix—Eye Strain, a Cause of Headache.
By DAVID WEBSTER, M.D. Second edition, Demy 8vo, 7s. 6d.

EDWARD COTTERELL, F.R.C.S. ENG., L.R.C.P. LOND.

Late House Surgeon, University College Hospital.

ON SOME COMMON INJURIES TO LIMBS; their
Treatment and After-treatment, including Bone-setting (so-called).
With Illustrations, small 8vo, 3s. 6d.

SIDNEY COUPLAND, M.D., F.R.C.P.

*Physician to the Middlesex Hospital, and Lecturer on Practical Medicine in the Medical
School; Examiner in Medicine at the Examining Board for England.*

NOTES ON THE EXAMINATION OF THE SPUTUM,
VOMIT, FÆCES, URINE, AND BLOOD. Second edition, 12mo,
1s. nett,

CHARLES CREIGHTON, M.D.

I.

ILLUSTRATIONS OF UNCONSCIOUS MEMORY IN
DISEASE, including a Theory of Alteratives. Post 8vo, 6s.

II.

CONTRIBUTIONS TO THE PHYSIOLOGY AND
PATHOLOGY OF THE BREAST AND LYMPHATIC GLANDS.
New Edition with additional chapter, with wood-cuts and plate, 8vo, 9s.

III.

BOVINE TUBERCULOSIS IN MAN: An Account of the
Pathology of Suspected Cases. With Chromo-lithographs and other
Illustrations, 8vo, 8s. 6d.

H. RADCLIFFE CROCKER, M.D. LOND., B.S., F.R.C.P.

Physician, Skin Department, University College Hospital.

DISEASES OF THE SKIN; THEIR DESCRIPTION,
PATHOLOGY, DIAGNOSIS, AND TREATMENT. With 76 Illus-
trations, 8vo, 21s.

EDGAR M. CROOKSHANK, M.B. LOND., F.R.M.S.

Professor of Comparative Pathology and Bacteriology in, and Fellow of King's College London.

I.

HISTORY AND PATHOLOGY OF VACCINATION.

Vol. I., A Critical Inquiry. Vol. II., Selected Essays, (Edited) including works by Jenner, Pearson, Woodville, Henry Jenner, Loy, Rogers, Birch, Bousquet, Estlin, Ceely, Badcock, Auzias-Turenne, Dubreuilh and Layet. Two volumes, illustrated with 22 coloured plates, including reproductions of the plates illustrating Jenner's Inquiry, of selected plates from the work of Ceely and others, and with a reduced facsimile of an engraving of Mr. Jesty, a facsimile of the first folio of the manuscript of Jenner's original paper, a facsimile of an unpublished letter from Jenner to Mr. Head, Royal 8vo, 36s.

II.

MANUAL OF BACTERIOLOGY: Illustrated with Coloured Plates from original drawings, and with other Illustrations in the text. Third Edition, 8vo, 21s. [Now ready.]

RIDLEY DALE, M.D., L.R.C.P. EDIN., M.R.C.S. ENG.

EPITOME OF SURGERY, being a complete compendium of the Science and Art of Surgery. Large 8vo, 10s. 6d.

HERBERT DAVIES, M.D., F.R.C.P.

Late Consulting Physician to the London Hospital.

THE MECHANISM OF THE CIRCULATION OF THE BLOOD THROUGH ORGANICALLY DISEASED HEARTS. Edited by ARTHUR TEMPLER DAVIES, B.A. (Nat. Science Honours), M.D. Cantab., M.R.C.P.; Physician to the Royal Hospital for Diseases of the Chest. Crown 8vo, 3s. 6d.

HENRY DAVIS, M.R.C.S. ENG.

Teacher and Administrator of Anæsthetics to St. Mary's and the National Dental Hospitals

GUIDE TO THE ADMINISTRATION OF ANÆSTHETICS. Second edition, fcap. 8vo. [In the press.]

J. THOMPSON DICKSON, M.A., M.B. CANTAB.

Late Lecturer on Mental Diseases at Guy's Hospital.

THE SCIENCE AND PRACTICE OF MEDICINE IN RELATION TO MIND, the Pathology of the Nerve Centres, and the Jurisprudence of Insanity being a course of Lectures delivered at Guy's Hospital. Illustrated by Chromo-lithographic Drawings and Physiological Portraits. 8vo, 14s.

HORACE DOBELL, M.D.

Consulting Physician to the Royal Hospital for Diseases of the Chest, &c.

I.

ON DIET AND REGIMEN IN SICKNESS AND Health and on the Interdependence and Prevention of Diseases and the Diminution of their Fatality. Seventh Edition, 8vo, 5s. nett.

II.

AFFECTIONS OF THE HEART AND IN ITS NEIGHBOURHOOD. Cases, Aphorisms, and Commentaries. Illustrated by the heliotype process. 8vo, 6s 6d.

JOHN EAGLE.

Member of the Pharmaceutical Society.

A NOTE-BOOK OF SOLUBILITIES. Arranged chiefly for the use of Prescribers and Dispensers. 12mo, 2s. 6d.

ARTHUR W. EDIS, M.D. LOND., F.R.C.P.

Senior Physician to the Chelsea Hospital for Women; Late Obstetric Physician to the Middlesex Hospital.

STERILITY IN WOMEN: including its Caustion and Treatment. With 33 Illustrations, demy 8vo, 6s. [Just published.]

DR. FERBER.

MODEL DIAGRAM OF THE ORGANS IN THE THORAX AND UPPER PART OF THE ABDOMEN. With Letter-press Description. In 4to, coloured, 5s.

J. MAGEE FINNY, M.D. DUBL.

King's Professor of Practice of Medicine in School of Physic, Ireland, &c.

NOTES ON THE PHYSICAL DIAGNOSIS OF LUNG DISEASES. 32mo, 1s. 6d. [Now ready.]

AUSTIN FLINT, M.D., LL.D.

Professor of Physiology and Physiological Anatomy in the Bellevue Hospital Medical College, New York; visiting Physician to the Bellevue Hospital, &c.

A TEXT-BOOK OF HUMAN PHYSIOLOGY. Fourth edition, Illustrated by plates, and 316 wood engravings, large 8vo, 25s.

J. MILNER FOTHERGILL, M.D., M.R.C.P.

Late Physician to the City of London Hospital for Diseases of the Chest, Victoria Park, &c.

I.
A MANUAL OF DIETETICS. Large 8vo, 10s. 6d.

II.
THE HEART AND ITS DISEASES, WITH THEIR TREATMENT; INCLUDING THE GOUTY HEART. Second Edition, entirely re-written, copiously illustrated with woodcuts and lithographic plates. 8vo, 16s.

III.
INDIGESTION AND BILIOUSNESS. Second Edition, post 8vo, 7s. 6d.

IV.
GOUT IN ITS PROTEAN ASPECTS. Post 8vo, 7s. 6d.

V.
THE TOWN DWELLER: His Needs and His Wants. With an Introduction by B. W. RICHARDSON, M.D., LL.D., F.R.S. Post 8vo, 3s. 6d.

FORTESCUE FOX, M.D. LOND.

Fellow of the Medical Society of London.

STRATHPEFFER SPA: Its Climate and Waters. With OBSERVATIONS HISTORICAL, MEDICAL, AND GENERAL DESCRIPTIVE OF THE VICINITY. Crown 8vo, with Map and Illustrations, 2s. 6d., *nett*.

JOHN HENRY GARRETT, M.D.

Licentiate in Sanitary Science and Diplomate in Public Health, Universities of Durham and Cambridge, &c.

THE ACTION OF WATER ON LEAD; being an inquiry into the Cause and Mode of the Action and its Prevention. Crown 8vo, 4s. 6d.

ALFRED W. GERRARD, F.C.S.

Examiner to the Pharmaceutical Society; Teacher of Materia Medica and Pharmacy at University College Hospital.

ELEMENTS OF MATERIA MEDICA AND PHARMACY. Crown 8vo, 8s. 6d.

NEW OFFICIAL REMEDIES, B.P., 1890. Supplement to the above. Crown 8vo, 1s.

HENEAGE GIBBES, M.D.

Lecturer on Physiology and on Normal and Morbid Histology in the Medical School of Westminster Hospital; &c.

PRACTICAL HISTOLOGY AND PATHOLOGY. Third Edition, revised and enlarged, crown 8vo, 6s.

C. A. GORDON, M.D., C.B.

Deputy Inspector General of Hospitals, Army Medical Department.

REMARKS ON ARMY SURGEONS AND THEIR WORKS. Demy 8vo, 5s.

JOHN GORHAM, M.R.C.S.

TOOTH EXTRACTION: a Manual on the proper mode of extracting Teeth. Third Edition, fcap. 8vo, 1s. 6d. [Now ready.]

GEORGE M. GOULD, B.A., M.D.

Ophthalmic Surgeon to the Philadelphia Hospital, &c.

A NEW MEDICAL DICTIONARY: including all the words and phrases used in Medicine, with their proper pronunciation and definitions. 8vo, 12s. 6d.

W. R. GOWERS, M.D., F.R.C.P., M.R.C.S.

Physician to University College Hospital, &c.

DIAGRAMS FOR THE RECORD OF PHYSICAL SIGNS. In books of 12 sets of figures, 1s. Ditto, unbound, 1s.

J. B. GRESSWELL, M.R.C.V.S.

Provincial Veterinary Surgeon to the Royal Agricultural Society.

VETERINARY PHARMACOLOGY AND THERAPEUTICS. With an Index of Diseases and Remedies. Fcap. 8vo, 5s.

SAMUEL D. GROSS, M.D., LL.D., D.C.L. OXON.

Professor of Surgery in the Jefferson Medical College of Philadelphia.

A PRACTICAL TREATISE ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE URINARY BLADDER, THE PROSTATE GLAND, AND THE URETHRA. Third Edition, revised and edited by S. W. GROSS, A.M., M.D., Surgeon to the Philadelphia Hospital. Illustrated by 170 engravings, 8vo, 18s.

SAMUEL W. GROSS, A.M., M.D.

Surgeon to, and Lecturer on Clinical Surgery in, the Jefferson Medical College Hospital and the Philadelphia Hospital, &c.

A PRACTICAL TREATISE ON TUMOURS OF THE MAMMARY GLAND: embracing their Histology, Pathology, Diagnosis, and Treatment. With Illustrations, 8vo, 10s. 6d.

PROF. JOSEF GRUBER.

Professor of Otology in the Imperial Royal University of Vienna, etc.

A TEXT-BOOK OF THE DISEASES OF THE EAR. Translated from the second German edition by special permission of the Author, and Edited by EDWARD LAW, M.D., C.M. EDIN., M.R.C.S. ENG., Surgeon to the London Throat Hospital for Diseases of the Throat, Nose and Ear; and by COLEMAN JEWELL, M.B. LOND., M.R.C.S. ENG., late Physician and Pathologist to the London Throat Hospital. With 150 Illustrations, and 70 coloured figures on 2 lithographic plates, Royal 8vo, 24s. [Just Published.]

ALLAN McLANE HAMILTON, M.D.

THE MODERN TREATMENT OF HEADACHES.
Square 16mo, 2s. 6d.

WILLIAM A. HAMMOND, M.D.

Professor of Mental and Nervous Diseases in the Medical Department of the University of the City of New York, &c.

SPIRITUALISM AND ALLIED CAUSES AND CONDITIONS OF NERVOUS DERANGEMENT. With Illustrations, post 8vo, 8s. 6d.

ALEXANDER HARVEY, M.D.

Late Emeritus Professor of Materia Medica in the University of Aberdeen, &c.,

AND

ALEXANDER DYCE DAVIDSON, M.D., F.R.S. EDIN.

Late Regius Professor of Materia Medica in the University of Aberdeen.

SYLLABUS OF MATERIA MEDICA FOR THE USE OF STUDENTS, TEACHERS AND PRACTITIONERS. Based on the relative values of articles and preparations in the British Pharmacopœia. Ninth edition, 32mo, 1s. 6d.

K. M. HEANLEY.

Matron of Boston Cottage Hospital.

A MANUAL OF URINE TESTING. Compiled for the use of Matrons, Nurses, and Probationers. Post 8vo, 1s. 6d.

C. HIGGENS, F.R.C.S.

Ophthalmic Surgeon to Guy's Hospital; Lecturer on Ophthalmology at Guy's Hospital Medical School

MANUAL OF OPHTHALMIC PRACTICE.

Crown 8vo, illustrations, 6s.

[LEWIS'S PRACTICAL SERIES.]

BERKELEY HILL, M.B. LOND., F.R.C.S.

Professor of Clinical Surgery in University College; Surgeon to University College Hospital and to the Lock Hospital.

THE ESSENTIALS OF BANDAGING. With directions for Managing Fractures and Dislocations; for administering Ether and Chloroform; and for using other Surgical Apparatus; with a Chapter on Surgical Landmarks. Sixth Edition, revised and enlarged, Illustrated by 144 Wood Engravings, crown 8vo, 5s.

BERKELEY HILL, M.B. LOND., F.R.C.S.

Professor of Clinical Surgery in University College; Surgeon to University College Hospital and to the Lock Hospital.

AND

ARTHUR COOPER, L.R.C.P., M.R.C.S.

Surgeon to the Westminster General Dispensary.

I

SYPHILIS AND LOCAL CONTAGIOUS DISORDERS.
Second edition, entirely re-written, royal 8vo, 18s.

II.

THE STUDENT'S MANUAL OF VENEREAL DISEASES. Being a Concise Description of those Affections and of their Treatment. Fourth edition, post 8vo, 2s. 6d.

PROCTER S. HUTCHINSON, M.R.C.S.

Assistant Surgeon to the Hospital for Diseases of the Throat.

A MANUAL OF DISEASES OF THE NOSE AND THROAT; including the Nose, Naso-pharynx, Pharynx, and Larynx. With Illustrations, crown 8vo, 3s. 6d. [Now ready.]

C. R. ILLINGWORTH, M.D. ED., M.R.C.S.

THE ABORTIVE TREATMENT OF SPECIFIC FEBRILE DISORDERS BY THE BINIODIDE OF MERCURY.
Crown 8vo, 3s. 6d.

SIR W. JENNER, Bart., M.D.

Physician in Ordinary to H.M. the Queen, and to H.R.H. the Prince of Wales.

THE PRACTICAL MEDICINE OF TO-DAY: Two Addresses delivered before the British Medical Association, and the Epidemiological Society, (1869). Small 8vo, 1s. 6d.

GEORGE LINDSAY JOHNSON, M.A., M.B., B.C. CANTAB.
Clinical Assistant, late House Surgeon and Chloroformist, Royal Westminster Ophthalmic Hospital, &c.

A NEW METHOD OF TREATING CHRONIC GLAUCOMA, based on Recent Researches into its Pathology. With Illustrations and coloured frontispiece, demy 8vo, 3s. 6d.

JOHN M KEATING,
Fellow of the College of Physicians, Philadelphia, &c.

AND

HENRY HAMILTON.

POCKET MEDICAL LEXICON.

32mo, 3s. nett.

NORMAN KERR, M.D., F.L.S.

President of the Society for the Study of Inebriety; Consulting Physician, Dalrymple Home for Inebriates, &c.

INEBRIETY: its Etiology, Pathology, Treatment, and Jurisprudence. Second edition, Crown 8vo, 12s. 6d.

NORMAN W. KINGSLEY, M.D.S., D.D.S.

President of the Board of Censors of the State of New York; Member of the American Academy of Dental Science, &c.

A TREATISE ON ORAL DEFORMITIES AS A BRANCH OF MECHANICAL SURGERY. With over 350 Illustrations, 8vo, 16s.

F. CHARLES LARKIN, F.R.C.S. ENG.

Surgeon to the Stanley Hospital; late Assistant Lecturer in Physiology in University College, Liverpool,

AND

RANDLE LEIGH, M.B., B.SC. LOND.

Senior Demonstrator of Physiology in University College, Liverpool.

OUTLINES OF PRACTICAL PHYSIOLOGICAL CHEMISTRY. Second edition, with Illustrations, crown 8vo, paper 2s. 6d. nett, or cloth 3s. nett. [Now ready.]

J. WICKHAM LEGG, F.R.C.P.

Late Assistant Physician to Saint Bartholomew's Hospital, and Lecturer on Pathological Anatomy in the Medical School.

I.

ON THE BILE, JAUNDICE, AND BILIOUS DISEASES.

With Illustrations in chromo-lithography, 719 pages, roy. 8vo, 25s.

II.

A GUIDE TO THE EXAMINATION OF THE URINE; intended chiefly for Clinical Clerks and Students. Sixth Edition, revised and enlarged, with Illustrations, fcap. 8vo, 2s. 6d.

ARTHUR H. N. LEWERS, M.D. LOND., M.R.C.P. LOND.

Assistant Obstetric Physician to the London Hospital; Examiner in Midwifery and Diseases of Women to the Society of Apothecaries of London, &c.

A PRACTICAL TEXTBOOK OF THE DISEASES OF WOMEN. Third edition, Illustrations, crown 8vo, 10s. 6d. [Now ready.]

[LEWIS'S PRACTICAL SERIES.]

LEWIS'S POCKET CASE BOOK FOR PRACTITIONERS AND STUDENTS. Designed by A. T. BRAND, M.D. Roan, with pencil, 3s. 6d. *nett*.

LEWIS'S POCKET MEDICAL VOCABULARY.

Second Edition, thoroughly revised, 32mo, roan, 3s. 6d. [*Just ready.*]

T. R. LEWIS, M.B., F.R.S. ELECT, ETC.

Late Fellow of the Calcutta University, Surgeon-Major Army Medical Staff, &c.

PHYSIOLOGICAL AND PATHOLOGICAL RESEARCHES. Arranged and edited by SIR WM. AITKEN, M.D., F.R.S., G. E. DOBSON, M.B., F.R.S., and A. E. BROWN, B.Sc. Crown 4to, portrait, 5 maps, 43 plates including 15 chromo-lithographs, and 67 wood engravings, 30s. *nett*.

* * A few copies only of this work remain for sale.

C. B. LOCKWOOD, F.R.C.S.

Hunterian Professor, Royal College of Surgeons of England; Surgeon to the Great Northern Hospital; Senior Demonstrator of Anatomy and Operative Surgery in St. Bartholomew's Hospital.

HUNTERIAN LECTURES ON THE MORBID ANATOMY, PATHOLOGY AND TREATMENT OF HERNIA. Demy 8vo, 36 illustrations, 5s.

J. S. LOMBARD, M.D.

Formerly Assistant Professor of Physiology in Harvard College.

I.
EXPERIMENTAL RESEARCHES ON THE REGIONAL TEMPERATURE OF THE HEAD, under Conditions of Rest, Intellectual Activity, and Emotion. With Illustrations, 8vo, 8s.

II.
ON THE NORMAL TEMPERATURE OF THE HEAD. 8vo, 5s.

WILLIAM THOMPSON LUSK, M.A., M.D.

Professor of Obstetrics and Diseases of Women in the Bellevue Hospital Medical College, &c.

THE SCIENCE AND ART OF MIDWIFERY.

Third Edition, with numerous Illustrations, 8vo, 18s.

A. W. MACFARLANE, M.D., F.R.C.P. EDIN.

Examiner in Medical Jurisprudence in the University of Glasgow; Honorary Consulting Physician (late Physician) Kilmarnock Infirmary.

INSOMNIA AND ITS THERAPEUTICS.

Medium 8vo, 12s. 6d.

SURGEON-MAJOR C. J. McNALLY, M.D., D.P.H. CAMB.
Fellow of the Madras University; Professor of Chemistry, Madras Medical College.

THE ELEMENTS OF SANITARY SCIENCE.

Plates, Demy 8vo, 8s. 6d.

RAWDON MACNAMARA.

Professor of Materia Medica, Royal College of Surgeons, Ireland; Senior Surgeon to the Westmoreland (Lock) Government Hospital; Surgeon to the Meath Hospital, &c.

AN INTRODUCTION TO THE STUDY OF THE BRITISH PHARMACOPŒIA. Demy 32mo, 1s. 6d. [*Just published.*]

JOHN MACPHERSON, M.D.

Inspector-General of Hospitals H.M. Bengal Army (Retired).

Author of "Cholera in its Home," &c.

I.

ANNALS OF CHOLERA FROM THE EARLIEST PERIODS TO THE YEAR 1817. With a map. Demy 8vo, 7s. 6d.

II.

BATH, CONTREXEVILLE, AND THE LIME SULPHATED WATERS. Crown 8vo, 2s. 6d.

A. COWLEY MALLEY, B.A., M.B., B.CH. T.C.D.

PHOTO-MICROGRAPHY; including a description of the Wet Collodion and Gelatino-Bromide Processes, together with the best methods of Mounting and Preparing Microscopic Objects for Photo-Micrography. Second Edition, with Photographs and Illustrations, crown 8vo, 7s. 6d.

PATRICK MANSON, M.D., C.M.

THE FILARIA SANGUINIS HOMINIS; AND CERTAIN NEW FORMS OF PARASITIC DISEASE IN INDIA, CHINA, AND WARM COUNTRIES. Illustrated with Plates and Charts. 8vo, 10s. 6d.

JEFFERY A. MARSTON, M.D., C.B., F.R.C.S., M.R.C.P. LOND.

Surgeon General Medical Staff (Retired).

NOTES ON TYPHOID FEVER: Tropical Life and its Sequelæ. Crown 8vo, 3s. 6d. [*Now ready.*]

PROFESSOR MARTIN.

MARTIN'S ATLAS OF OBSTETRICS AND GYNÆCOLOGY. Edited by A. MARTIN, Docent in the University of Berlin. Translated and edited with additions by FANCOURT BARNES, M.D., M.R.C.P., Physician to the Chelsea Hospital for Women; Obstetric Physician to the Great Northern Hospital; and to the Royal Maternity Charity of London, &c. Medium 4to, Morocco half bound, 31s. 6d. *nett.*

EDWARD MARTIN, A.M., M.D.

MINOR SURGERY AND BANDAGING WITH AN APPENDIX ON VENEREAL DISEASES. Crown 8vo, 82 Illustrations, 4s.

WILLIAM MARTINDALE, F.C.S.

Late Examiner of the Pharmaceutical Society, and late Teacher of Pharmacy and Demonstrator of Materia Medica at University College,

AND

W. WYNN WESTCOTT, M.B. LOND.

Deputy Coroner for Central Middlesex.

THE EXTRA PHARMACOPŒIA with the additions introduced into the British Pharmacopœia, 1885 and 1890, with Medical References, and a Therapeutic Index of Diseases and Symptoms. Sixth Edition, limp roan, med. 24mo, 7s. 6d. [Now ready

WILLIAM MARTINDALE, F.C.S.

Late Examiner of the Pharmaceutical Society, &c.

COCA, AND COCAINE. Their History, Medical and Economic Uses, and Medicinal Preparations. Second edition, coloured plate, fcap. 8vo, 2s.

MATERIA MEDICA LABELS.

Adapted for Public and Private Collections. Compiled from the British Pharmacopœia of 1885, with the additions of 1890. The Labels are arranged in Two Divisions:—

Division I.—Comprises, with few exceptions, Substances of Organized Structure, obtained from the Vegetable and Animal Kingdoms.

Division II.—Comprises Chemical Materia Medica, including Alcohols, Alkaloids, Sugars, and Neutral Bodies.

On plain paper, 10s. 6d. *nett.* On gummed paper, 12s. 6d. *nett.*

The 24 additional Labels of 1890 only, 1s. *nett.*

* * * Specimens of the Labels, of which there are over 470, will be sent on application.

S. E. MAUNSELL, L.R.C.S.I.

Surgeon-Major, Medical Staff.

NOTES OF MEDICAL EXPERIENCES IN INDIA PRINCIPALLY WITH REFERENCE TO DISEASES OF THE EYE. With Map, post 8vo, 3s. 6d.

J. F. MEIGS, M.D.

Consulting Physician to the Children's Hospital, Philadelphia,

AND

W. PEPPER, M.D.

Lecturer on Clinical Medicine in the University of Pennsylvania.

A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. Seventh Edition, revised and enlarged, roy. 8vo, 28s.

Wm. JULIUS MICKLE, M.D., F.R.C.P. LOND.
Medical Superintendent, Grove Hall Asylum, London, &c.

I.
GENERAL PARALYSIS OF THE INSANE.

Second Edition, enlarged and rewritten, 8vo, 14s.

II.
**ON INSANITY IN RELATION TO CARDIAC AND
 AORTIC DISEASE AND PHTHISIS.** Crown 8vo, 3s. 6d.

ANGEL MONEY, M.D. LOND., F.R.C.P.
*Assistant Physician to University College Hospital, and to the Hospital for Sick Children
 Great Ormond Street; Assistant Professor of Clinical Medicine in University
 College, London, &c.*

I.)
**TREATMENT OF DISEASE IN CHILDREN: EM-
 BODYING THE OUTLINES OF DIAGNOSIS AND THE
 CHIEF PATHOLOGICAL DIFFERENCES BETWEEN CHILD-
 REN AND ADULTS.** Second edition, crown 8vo, 10s. 6d.

[LEWIS'S PRACTICAL SERIES.]

II.
**THE STUDENT'S TEXTBOOK OF THE PRACTICE
 OF MEDICINE.** Fcap. 8vo, 6s. 6d.

A. STANFORD MORTON, M.B., F.R.C.S. ENG.
Surgeon to the Royal South London Ophthalmic Hospital.

**REFRACTION OF THE EYE: Its Diagnosis, and the
 Correction of its Errors.** Fourth Edition, with Illustrations, small 8vo,
 3s. 6d. [Now ready.]

C. W. MANSELL MOULLIN, M.A., M.D. OXON., F.R.C.S. ENG.
*Assistant Surgeon and Senior Demonstrator of Anatomy at the London Hospital; formerly
 Radcliffe Travelling Fellow and Fellow of Pembroke College, Oxford.*

**SPRAINS; THEIR CONSEQUENCES AND TREAT-
 MENT.** Crown 8vo, 5s.

PAUL F. MUNDE, M.D.
*Professor of Gynecology at the New York Polyclinic; President of the New York Obstetrical
 Society and Vice-President of the British Gynecological Society, &c.*

**THE MANAGEMENT OF PREGNANCY, PARTURI-
 TION, AND THE PUERPERAL STATE.** Second edition, square
 8vo, 3s. 6d.

WILLIAM MURRAY, M.D., F.R.C.P. LOND.
Consulting Physician to the Children's Hospital, Newcastle-on-Tyne, &c.

**ILLUSTRATIONS OF THE INDUCTIVE METHOD IN
 MEDICINE.** Crown 8vo, 3s. 6d. [Just ready.]

WILLIAM MURRELL, M.D., F.R.C.P.

Lecturer on Pharmacology and Therapeutics at Westminster Hospital; late Examiner in Materia Medica to the Royal College of Physicians of London, etc.

I.

MASSOTHERAPEUTICS, OR MASSAGE AS A MODE OF TREATMENT. Fifth edition, with Illustrations, crown 8vo, 4s. 6d.

II.

WHAT TO DO IN CASES OF POISONING. Sixth edition, royal 32mo, 3s. 6d.

III.

NITRO-GLYCERINE AS A REMEDY FOR ANGINA PECTORIS. Crown 8vo, 3s. 6d.

IV.

CHRONIC BRONCHITIS AND ITS TREATMENT. Crown 8vo, 3s. 6d.

DR. FELIX von NIEMEYER.

Late Professor of Pathology and Therapeutics; Director of the Medical Clinic of the University of Tübingen.

A TEXT-BOOK OF PRACTICAL MEDICINE, WITH PARTICULAR REFERENCE TO PHYSIOLOGY AND PATHOLOGICAL ANATOMY. Translated from the Eighth German Edition by special permission of the Author, by GEORGE H. HUMPHERY, M.D., and CHARLES E. HACKLEY, M.D. Revised edition, 2 vols. large 8vo, 36s.

GEORGE OLIVER, M.D., F.R.C.P.

I.

THE HARROGATE WATERS: Data Chemical and Therapeutical, with notes on the Climate of Harrogate. Addressed to the Medical Profession. Crown 8vo, with Map of the Wells, 3s. 6d.

II.

ON BEDSIDE URINE TESTING: a Clinical Guide to the Observation of Urine in the course of Work. Fourth Edition, fcap. 8vo, 3s. 6d.

SAMUEL OSBORN, F.R.C.S.

Surgeon to the Hospital for Women, Soho Square; Surgeon to the Royal Naval Artillery Volunteers.

I.

AMBULANCE LECTURES: FIRST AID. Second edition, with Illustrations, fcap. 8vo, 1s. 6d.

II.

AMBULANCE LECTURES: HOME NURSING AND HYGIENE. Second edition, with Illustrations, fcap. 8vo, 2s.

[Just published.]

WILLIAM OSLER, M.D., F.R.C.P. LOND.

Professor of Clinical Medicine in the University of Pennsylvania, &c.

THE CEREBRAL PALSIES OF CHILDREN. A Clinical Study from the Infirmary for Nervous Diseases, Philadelphia. Demy 8vo, 5s. [Just Published.]

KURRE W. OSTROM.

Instructor in Massage and Swedish Movements in the Philadelphia Polyclinic and College for Graduates in Medicine.

MESSAGE AND THE ORIGINAL SWEDISH MOVEMENTS; their application to various diseases of the body. Second edition, with Illustrations, 12mo, 3s. 6d. nett. [Now ready.]

ROBERT W. PARKER.

Senior Surgeon to the East London Hospital for Children; Surgeon to the German Hospital.

I.

DIPHTHERIA: ITS NATURE AND TREATMENT, WITH SPECIAL REFERENCE TO THE OPERATION, AFTER-TREATMENT AND COMPLICATIONS OF TRACHEOTOMY. Third Edition, with Illustrations, 8vo, 6s. [Now ready.]

II.

CONGENITAL CLUB-FOOT; ITS NATURE AND TREATMENT. With special reference to the subcutaneous division of Tarsal Ligaments. 8vo, 7s. 6d.

LOUIS C. PARKES, M.D., D.P.H. LOND. UNIV.

Fellow of the Sanitary Institute, and Member of the Board of Examiners; Assistant Professor of Hygiene and Public Health at University College, London, &c.

HYGIENE AND PUBLIC HEALTH. Second edition, with numerous Illustrations, crown 8vo, 9s. [Just Published.] [LEWIS'S PRACTICAL SERIES.]

JOHN S. PARRY, M.D.

Obstetrician to the Philadelphia Hospital, Vice-President of the Obstetrical and Pathological Societies of Philadelphia, &c.

EXTRA-UTERINE PREGNANCY; Its Causes, Species, Pathological Anatomy, Clinical History, Diagnosis, Prognosis and Treatment. 8vo, 8s.

THEOPHILUS PARVIN, M.D.

Professor of Obstetrics and Diseases of Women and Children at the Jefferson Medical School.

LECTURES ON OBSTETRIC NURSING, Delivered at the Training School for Nurses of the Philadelphia Hospital. Post 8vo, 2s. 6d.

E. RANDOLPH PEASLEE, M.D., LL.D.

Late Professor of Gynæcology in the Medical Department of Dartmouth College; President of New York Academy of Medicine, &c., &c.

OVARIAN TUMOURS: Their Pathology, Diagnosis, and Treatment, especially by Ovariectomy. Illustrations, roy. 8vo, 16s.

HENRY G. PIFFARD, A.M., M.D.

Clinical Professor of Dermatology, University of the City of New York; Surgeon in Charge of the New York Dispensary for Diseases of the Skin, &c.

A PRACTICAL TREATISE ON DISEASES OF THE SKIN. With 50 full page Original Plates and 33 Illustrations in the Text, 4to, £2 12s. 6d. *nett.* [Just published.]

G. V. POORE, M.D., F.R.C.P.

Professor of Medical Jurisprudence, University College; Assistant Physician to, and Physician in charge of the Throat Department of, University College Hospital.

LECTURES ON THE PHYSICAL EXAMINATION OF THE MOUTH AND THROAT. With an Appendix of Cases. 8vo, 3s. 6d.

R. DOUGLAS POWELL, M.D., F.R.C.P., M.R.C.S.

Physician Extra-ordinary to H.M. the Queen; Physician to the Middlesex Hospital and Physician to the Hospital for Consumption and Diseases of the Chest at Brompton.

I.
DISEASES OF THE LUNGS AND PLEURÆ, INCLUDING CONSUMPTION. Third edition, entirely rewritten and enlarged. With coloured plates and wood engravings, 8vo, 16s.

II.
TABLE OF PHYSICAL EXAMINATION OF THE LUNGS—with Note on International Nomenclature of Physical Signs (reprinted from above). On one sheet, 6d.

URBAN PRITCHARD, M.D. EDIN., F.R.C.S. ENG.

Professor of Aural Surgery at King's College, London; Aural Surgeon to King's College Hospital; Senior Surgeon to the Royal Ear Hospital.

HANDBOOK OF DISEASES OF THE EAR FOR THE USE OF STUDENTS AND PRACTITIONERS. Second edition, With Illustrations, crown 8vo, 5s. [LEWIS'S PRACTICAL SERIES.]

CHARLES W. PURDY, M.D. (QUEEN'S UNIV.)

Professor of Genito-Urinary and Renal Diseases in the Chicago Polyclinic, &c., &c.

BRIGHT'S DISEASE AND THE ALLIED AFFECTIONS OF THE KIDNEYS. With Illustrations, large 8vo, 8s. 6d.

DR. THEODOR PUSCHMANN.

Public Professor in Ordinary at the University of Vienna.

A HISTORY OF MEDICAL EDUCATION FROM THE MOST REMOTE TO THE MOST RECENT TIMES. Translated and edited by EVAN H. HARE, M.A. OXON., F.R.C.S. ENG., L.S.A. Demy 8vo, 21s. [Now ready.]

CHARLES HENRY RALFE, M.A., M.D. CANTAB., F.R.C.P. LOND.
Assistant Physician to the London Hospital; Examiner in Medicine to the University of Durham, &c., &c

A PRACTICAL TREATISE ON DISEASES OF THE KIDNEYS AND URINARY DERANGEMENTS. With Illustrations, crown 8vo, 10s. 6d. [LEWIS'S PRACTICAL SERIES.]

FRANCIS H. RANKIN, M.D.
President of the New York Medical Society.

HYGIENE OF CHILDHOOD. Suggestions for the care of Children after the Period of Infancy to the completion of Puberty. Crown 8vo, 3s.

AMBROSE L. RANNEY, A.M., M.D.
Professor of the Anatomy and Physiology of the Nervous System in the New York Post-Graduate Medical School and Hospital, &c.

THE APPLIED ANATOMY OF THE NERVOUS SYSTEM. Second edition, 238 Illustrations, large 8vo, 21s.

H. A. REEVES, F.R.C.S. EDIN.
Senior Assistant Surgeon and Teacher of Practical Surgery at the London Hospital; Surgeon to the Royal Orthopædic Hospital.

BODILY DEFORMITIES AND THEIR TREATMENT: A HANDBOOK OF PRACTICAL ORTHOPÆDICS. Illustrations, crown 8vo, 8s. 6d. [LEWIS'S PRACTICAL SERIES.]

RALPH RICHARDSON, M.A., M.D.
Fellow of the College of Physicians, Edinburgh.

ON THE NATURE OF LIFE: An Introductory Chapter to Pathology. Second edition, revised and enlarged. Fcap. 4to, 10s. 6d.

W. RICHARDSON, M.A., M.D., M.R.C.P.

REMARKS ON DIABETES, ESPECIALLY IN REFERENCE TO TREATMENT. Demy 8vo, 4s. 6d.

SAMUEL RIDEAL, D.SC. (LOND.), F.I.C., F.C.S., F.G.S.
Fellow of University College, London.

I.
PRACTICAL ORGANIC CHEMISTRY; The Detection and Properties of some of the more important Organic Compounds. 12mo, 2s. 6d.

II.
PRACTICAL CHEMISTRY FOR MEDICAL STUDENTS, required at the First Examination of the Conjoint Examining Board in England. Foolscap 8vo, 2s. [Just published.]

E. A. RIDSDALE.

Associate of the Royal School of Mines.

COSMIC EVOLUTION; being Speculations on the Origin
of our Environment. Fcap. 8vo, 3s.

SYDNEY RINGER, M.D., F.R.S.

*Professor of the Principles and Practice of Medicine in University College; Physician to,
and Professor of Clinical Medicine in, University College Hospital.*

I.

A HANDBOOK OF THERAPEUTICS. Twelfth Edition
thoroughly revised, 8vo, 15s.

II.

**ON THE TEMPERATURE OF THE BODY AS
A MEANS OF DIAGNOSIS AND PROGNOSIS IN PHTHISIS.**
Second edition, small 8vo, 2s. 6d.

FREDERICK T. ROBERTS, M.D., B.SC., F.R.C.P.

*Examiner in Medicine at the University of London and for the Conjoint Board; Professor
of Materia Medica and Therapeutics and of Clinical Medicine in University
College; Physician to University College Hospital; Physician to
Brompton Consumption Hospital, &c.*

I.

**A HANDBOOK OF THE THEORY AND PRACTICE
OF MEDICINE.** Eighth edition, with Illustrations, in one volume,
large 8vo, 21s. [Just published.]

II.

THE OFFICINAL MATERIA MEDICA.
Second edition, entirely rewritten in accordance with the latest British
Pharmacopœia, fcap. 8vo, 7s. 6d.

III.

**NOTES ON THE ADDITIONS MADE TO THE BRITISH
PHARMACOPŒIA, 1890.** Fcap. 8vo, 1s. [Now ready.]

R. LAWTON ROBERTS, M.D. LOND., D.P.H. CAMB., M.R.C.S. ENG.
*Honorary Life Member of, and Lecturer and Examiner to, the St. John Ambulance
Association.*

I.

ILLUSTRATED LECTURES ON AMBULANCE WORK.
Fourth edition, copiously Illustrated, crown 8vo, 2s. 6d. [Now ready.]

II.

**ILLUSTRATED LECTURES ON NURSING AND HY-
GIENE.** Second edition, with Illustrations, crown 8vo, 2s. 6d.
[Just ready.]

ROBSON ROOSE, M.D., LL.D., F.C.S.
Fellow of the Royal College of Physicians in Edinburgh.

I.
**GOUT, AND ITS RELATIONS TO DISEASES OF
 THE LIVER AND KIDNEYS.** Sixth Edition, crown 8vo, 3s. 6d.

II.
**NERVE PROSTRATION AND OTHER FUNCTIONAL
 DISORDERS OF DAILY LIFE.** Second edition, demy 8vo, 18s.
 [Now ready.]

III.
**LEPROSY AND ITS PREVENTION: as Illustrated by
 Norwegian Experience.** Crown 8vo, 3s. 6d.

WILLIAM ROSE, M.B., B.S. LOND.), F.R.C.S.
Professor of Surgery in King's College, London, and Surgeon to King's College Hospital.
HARELIP AND CLEFT PALATE. With Illustrations, demy
 8vo, 6s. [Just published.]

BERNARD ROTH, F.R.C.S.
*Fellow of the Medical Society of London; Member of the Clinical and Pathological Societies
 and of the Medical Officers of Schools' Association.*

**THE TREATMENT OF LATERAL CURVATURE OF
 THE SPINE.** With Photographic and other Illustrations, demy 8vo,
 5s.

J. BURDON SANDERSON, M.D., LL.D., F.R.S.
Jodrell Professor of Physiology in University College, London.

**UNIVERSITY COLLEGE COURSE OF PRACTICAL
 EXERCISES IN PHYSIOLOGY.** With the co-operation of F. J. M.
 PAGE, B.Sc., F.C.S.; W. NORTH, B.A., F.C.S., and AUG. WALLER, M.D.
 Demy 8vo, 3s. 6d.

W. H. O. SANKEY, M.D. LOND., F.R.C.P.
Late Lecturer on Mental Diseases, University College, London, etc.

LECTURES ON MENTAL DISEASE. Second Edition, with
 coloured Plates, 8vo, 12s. 6d.

JOHN SAVORY.
Member of the Society of Apothecaries, London.

**A COMPENDIUM OF DOMESTIC MEDICINE AND
 COMPANION TO THE MEDICINE CHEST:** Intended as a
 source of easy reference for Clergymen, Master Mariners, and Tra-
 vellers; and for Families resident at a distance from professional assist-
 ance. Tenth Edition, sm. 8vo, 5s.

E. SCHMIEGELOW, M.D.

Consulting Physician in Laryngology to the Municipal Hospital and Director of the Oto-Laryngological Department in the Polyclinic at Copenhagen.

ASTHMA: Especially in its Relation to Nasal Disease.
Demy 8vo, 4s. 6d.

DR. B. S. SCHULTZE.

Professor of Gynecology; Director of the Lying-in Hospital, and of the Gynecological Clinic at Jena.

THE PATHOLOGY AND TREATMENT OF DISPLACEMENTS OF THE UTERUS. Translated by J. J. MACAN, M.A., M.R.C.S. and edited by A. V. MACAN, M.B., M.Ch., Master of the Rotunda Lying-in Hospital, Dublin. With 120 Illustrations, medium 8vo, 12s. 6d.

JOHN SHAW, M.D. LOND., M.R.C.P.

Obstetric Physician to the North-West London Hospital.

ANTISEPTICS IN OBSTETRIC NURSING. A Text-book for Nurses on the Application of Antiseptics to Gynæcology and Midwifery. Coloured plate and woodcuts, 8vo, 3s. 6d.

WM. JAPP SINCLAIR, M.A., M.D.

Honorary Physician to the Manchester Southern Hospital for Women and Children, and Manchester Maternity Hospital.

ON GONORRHOËAL INFECTION IN WOMEN.

Post 8vo, 4s.

A. J. C. SKENE, M.D.

Professor of Gynecology in the Long Island College Hospital, Brooklyn, New York.

TREATISE ON THE DISEASES OF WOMEN, FOR THE USE OF STUDENTS AND PRACTITIONERS. Nine coloured plates and 251 engravings, large 8vo, 28s.

J. LEWIS SMITH, M.D.

Physician to the New York Foundling Asylum; Clinical Professor of Diseases of Children in Bellevue Hospital Medical College.

A TREATISE ON THE DISEASES OF INFANCY AND CHILDHOOD. Seventh Edition, with Illustrations, large 8vo, 21s. [Just published.]

FRANCIS W. SMITH, M.B., B.S.

THE SALINE WATERS OF LEAMINGTON. Second Edit., with Illustrations, crown 8vo, 1s. nett.

JOHN KENT SPENDER, M.D. LOND.

Physician to the Royal Mineral Water Hospital, Bath.

THE EARLY SYMPTOMS AND THE EARLY TREATMENT OF OSTEO-ARTHRITIS, commonly called Rheumatoid Arthritis, with special reference to the Bath Thermal Waters. Sm. 8vo, 2s. 6d.

LOUIS STARR, M.D.

Clinical Professor of Diseases of Children in the Hospital of the University of Pennsylvania; Physician to the Children's Hospital, Philadelphia, &c.

HYGIENE OF THE NURSERY. Including the General Regimen and Feeding of Infants and Children, and the Domestic Management of the Ordinary Emergencies of Early Life. Second edition, with Illustrations, crown 8vo, 3s. 6d. [Just published.]

JAMES STARTIN, M.B., M.R.C.S.

Surgeon and Joint Lecturer to St. John's Hospital for Diseases of the Skin.

LECTURES ON THE PARASITIC DISEASES OF THE SKIN. VEGETOID AND ANIMAL. With Illustrations, crown 8vo, 2s. 6d.

JOHN LINDSAY STEVEN, M.D.

Assistant Physician and Pathologist, Glasgow Royal Infirmary; Physician for Out-patients, Royal Hospital for Sick Children, Glasgow; Lecturer on Pathology, St. Mungo's and Queen Margaret Colleges, Glasgow, &c.

THE PATHOLOGY OF MEDIASTINAL TUMOURS. With special reference to Diagnosis. With Plates, 8vo, 4s. 6d. [Just published.]

W. R. H. STEWART, F.R.C.S., L.R.C.P. EDIN.

Aural Surgeon to the Great Northern Central Hospital; Surgeon to the London Throat Hospital, &c.

EPITOME OF DISEASES AND INJURIES OF THE EAR, with a Chapter on Naso-Pharyngeal Diseases causing Deafness. Demy 32mo, 2s. 6d.

LEWIS A. STIMSON, B.A., M.D.

Surgeon to the Presbyterian and Bellevue Hospitals; Professor of Clinical Surgery in the Medical Faculty of the University of the City of New York, &c.

A MANUAL OF OPERATIVE SURGERY.

Second Edition, with three hundred and forty-two Illustrations, post 8vo, 10s. 6d.

ADOLF STRÜMPELL.

Director of the Medical Clinic in the University of Erlangen.

A TEXT-BOOK OF MEDICINE FOR STUDENTS AND PRACTITIONERS. Translated from the latest German edition by Dr. H. F. VICKERY and Dr. P. C. KNAPP, with Editorial Notes by Dr. F. C. SHATTUCK, Visiting Physician to the Massachusetts General Hospital, etc. Complete in one large vol., imp. 8vo, with 111 illustrations, 28s.

JUKES DE STYRAP, M.K.Q.C.P., ETC.

Physician-Extraordinary, late Physician in Ordinary, to the Salop Infirmary; Consulting Physician to the South Salop and Montgomeryshire Infirmaries, etc.

I.

THE YOUNG PRACTITIONER: WITH PRACTICAL HINTS AND INSTRUCTIVE SUGGESTIONS, AS SUBSIDIARY AIDS, FOR HIS GUIDANCE ON ENTERING INTO PRIVATE PRACTICE. Demy 8vo, 7s. 6d. *nett.*

II.

A CODE OF MEDICAL ETHICS: WITH GENERAL AND SPECIAL RULES FOR THE GUIDANCE OF THE FACULTY AND THE PUBLIC IN THE COMPLEX RELATIONS OF PROFESSIONAL LIFE. Third edition, demy 8vo, 3s. *nett.*

III.

MEDICO-CHIRURGICAL TARIFFS.

Fourth Edition, fcap. 4to, revised and enlarged, 2s. *nett.*

IV.

THE YOUNG PRACTITIONER: HIS CODE AND TARIFF. Being the above three works in one volume. Demy 8vo, 10s. 6d. *nett.*

C. W. SUCKLING, M.D. LOND., M.R.C.P.

Professor of Materia Medica and Therapeutics at the Queen's College, Physician to the Queen's Hospital, Birmingham, etc.

I.

ON THE DIAGNOSIS OF DISEASES OF THE BRAIN, SPINAL CORD, AND NERVES. With Illustrations, crown 8vo, 8s. 6d.

II.

ON THE TREATMENT OF DISEASES OF THE NERVOUS SYSTEM. Crown 8vo, 7s. 6d.

JOHN BLAND SUTTON, F.R.C.S.

Lecturer on Comparative Anatomy, Senior Demonstrator of Anatomy, and Assistant Surgeon to the Middlesex Hospital; Erasmus Wilson Lecturer, Royal College of Surgeons, England.

LIGAMENTS: THEIR NATURE AND MORPHOLOGY. With numerous Illustrations, post 8vo, 4s. 6d.

HENRY R. SWANZY, A.M., M.B., F.R.C.S.I.

Examiner in Ophthalmic Surgery in the Royal University of Ireland; Surgeon to the National Eye and Ear Infirmary, Dublin; Ophthalmic Surgeon to the Adelaide Hospital, Dublin, etc.

A HANDBOOK OF THE DISEASES OF THE EYE AND THEIR TREATMENT. Third Edition, Illustrated with wood-engravings, colour tests, etc., small 8vo, 10s. 6d. [Just Published.]

EUGENE S. TALBOT, M.D., D.D.S.

Professor of Dental Surgery in the Woman's Medical College; Lecturer on Dental Pathology and Surgery in Rush Medical College, Chicago.

IRREGULARITIES OF THE TEETH AND THEIR TREATMENT. With 152 Illustrations, royal 8vo, 10s. 6d.

H. COUPLAND TAYLOR, M.D.

Fellow of the Royal Meteorological Society.

WANDERINGS IN SEARCH OF HEALTH, OR MEDICAL AND METEOROLOGICAL NOTES ON VARIOUS FOREIGN HEALTH RESORTS. Crown 8vo, with Illustrations, 6s. [Now ready]

JOHN DAVIES THOMAS, M.D. LOND., F.R.C.S. ENG.

Physician to the Adelaide Hospital, S. Australia.

I.

HYDATID DISEASE, WITH SPECIAL REFERENCE TO ITS PREVALENCE IN AUSTRALIA. Demy 8vo, 10s. 6d.

II.

HYDATID DISEASE OF THE LUNGS. Demy 8vo, 2s.

HUGH OWEN THOMAS, M.R.C.S.

CONTRIBUTIONS TO SURGERY AND MEDICINE:—

PART I.—Intestinal Obstruction; with an Appendix on the Action of Remedies. 10s.

„ 2.—The Principles of the Treatment of Joint Disease, Inflammation, Ankylosis, Reduction of Joint Deformity, Bone Setting. 5s.

„ 3.—Fractures, Dislocations, Diseases and Deformities of the Bones of the Trunk and Upper Extremities. 10s.

„ 4.—The Collegian of 1666 and the Collegians of 1885; or what is recognised treatment? Second Edition, 1s.

„ 5.—On Fractures of the Lower Jaw. 1s.

„ 6.—The Principles of the Treatment of Fractures and Dislocations. 10s.

„ 7.—Fractures, Dislocations, Deformities, and Diseases of the Lower Extremities, 10s.

„ 8.—The Inhibition of Nerves by Drugs. Proof that Inhibitory Nerve-Fibres do not exist. 1s.

J. ASHBURTON THOMPSON, M.R.C.S.

Late Surgeon at King's Cross to the Great Northern Railway Company.

FREE PHOSPHORUS IN MEDICINE WITH SPECIAL REFERENCE TO ITS USE IN NEURALGIA. A contribution to *Materia Medica* and Therapeutics. An account of the History, Pharmaceutical Preparations, Dose, Internal Administration, and Therapeutic uses of Phosphorus; with a Complete Bibliography of this subject, referring to nearly 200 works upon it. Demy 8vo, 7s. 6d.

J. C. THOROWGOOD, M.D.

Assistant Physician to the City of London Hospital for Diseases of the Chest

THE CLIMATIC TREATMENT OF CONSUMPTION AND CHRONIC LUNG DISEASES. Third Edition, post 8vo, 3s. 6d.

D. HACK TUKE, M.D., LL.D.

Fellow of the Royal College of Physicians, London.

THE INSANE IN THE UNITED STATES AND CANADA. Demy 8vo, 7s. 6d.

DR. R. ULTZMANN.

ON STERILITY AND IMPOTENCE IN MAN. Translated from the German with notes and additions by ARTHUR COOPER, L.R.C.P., M.R.C.S., Surgeon to the Westminster General Dispensary. With Illustrations, fcap. 8vo, 2s. 6d.

W. H. VAN BUREN, M.D., LL.D.

Professor of Surgery in the Bellevue Hospital Medical College.

DISEASES OF THE RECTUM: And the Surgery of the Lower Bowel. Second Edition, with Illustrations, 8vo, 14s.

RUDOLPH VIRCHOW, M.D.

Professor in the University, and Member of the Academy of Sciences of Berlin, &c., &c.

INFECTION-DISEASES IN THE ARMY. Chiefly Wound Fever, Typhoid, Dysentery, and Diphtheria. Translated from the German by JOHN JAMES, M.B., F.R.C.S. Fcap. 8vo, 1s. 6d.

ALFRED VOGEL, M.D.

Professor of Clinical Medicine in the University of Dorpat, Russia.

A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. Third Edition, translated and edited by H. RAPHAEL, M.D., from the Eighth German Edition, illustrated by six lithographic plates, part coloured, royal 8vo, 18s.

A. DUNBAR WALKER, M.D., C.M.

THE PARENT'S MEDICAL NOTE BOOK.

Oblong post 8vo, cloth, 1s. 6d.

JOHN RICHARD WARDELL, M.D. EDIN., F.R.C.P. LOND.

Late Consulting Physician to the General Hospital Tunbridge Wells.

CONTRIBUTIONS TO PATHOLOGY AND THE PRACTICE OF MEDICINE. Medium 8vo, 21s.

W. SPENCER WATSON, F.R.C.S. ENG., B.M. LOND.

Surgeon to the Throat Department of the Great Northern Hospital; Senior Surgeon to the Royal South London Ophthalmic Hospital.

I.

DISEASES OF THE NOSE AND ITS ACCESSORY CAVITIES. Second edition, with Illustrations, demy 8vo, 12s. 6d.

[*Now ready.*]

II.

DISEASES OF THE LACHRYMAL PASSAGES.

[*In preparation.*]

III.

EYEBALL-TENSION: Its Effects on the Sight and its Treatment. With woodcuts, p. 8vo, 2s. 6d.

IV.

ON ABSCESS AND TUMOURS OF THE ORBIT.

Post 8vo, 2s. 6d.

FRANCIS H. WELCH, F.R.C.S.

Surgeon Major, A.M.D.

ENTERIC FEVER: as Illustrated by Army Data at Home and Abroad, its Prevalence and Modifications, Ætiology, Pathology and Treatment. 8vo, 5s. 6d.

W. WYNN WESTCOTT, M.B.
Deputy Coroner for Central Middlesex.

SUICIDE; its History, Literature, Jurisprudence, and Prevention. Crown 8vo, 6s.

E. G. WHITTLE, M.D. LOND., F.R.C.S. ENG.
Senior Surgeon to the Royal Alexandra Hospital for Sick Children, Brighton.

CONGESTIVE NEURASTHENIA, OR INSOMNIA AND NERVE DEPRESSION. Crown 8vo, 3s. 6d.

JOHN WILLIAMS, M.D., F.R.C.P.
Professor of Midwifery in University College, London; Obstetric Physician to University College Hospital; Physician Accoucheur to H.R.H. Princess Beatrice, &c.

CANCER OF THE UTERUS: Being the Harveian Lectures for 1886. Illustrated with Lithographic Plates, royal 8vo, 1os. 6d.

E. F. WILLOUGHBY, M.D. LOND.

THE NATURAL HISTORY OF SPECIFIC DISEASES OR STUDIES IN ETIOLOGY, IMMUNITY, AND PROPYLAXIS. 8vo, 2s. 6d.

E. T. WILSON, B.M. OXON., F.R.C.P. LOND.
Physician to the Cheltenham General Hospital and Dispensary.

DISINFECTANTS AND HOW TO USE THEM. In Packets of one doz. price 1s.

DR. F. WINCKEL.

Formerly Professor and Director of the Gynæcological Clinic at the University of Rostock.

THE PATHOLOGY AND TREATMENT OF CHILD-BED: A Treatise for Physicians and Students. Translated from the Second German edition, with many additional notes by the Author, by J. R. CHADWICK, M.D. 8vo, 14s.

BERTRAM C. A. WINDLE, M.A., M.D. DUBL.
Professor of Anatomy in the Queen's College, Birmingham; Examiner in Anatomy in the Universities of Cambridge and Durham.

A HANDBOOK OF SURFACE ANATOMY AND LAND-MARKS. Illustrated, post 8vo, 3s. 6d.

EDWARD WOAKES, M.D. LOND.

Senior Aural Surgeon and Lecturer on Aural Surgery at the London Hospital; Surgeon to the London Throat Hospital.

I.

ON DEAFNESS, GIDDINESS AND NOISES IN THE HEAD.

VOL. I.—POST-NASAL CATARRH, AND DISEASES OF THE NOSE CAUSING DEAFNESS. With Illustrations, cr. 8vo, 6s. 6d.

VOL. II.—ON DEAFNESS, GIDDINESS AND NOISES IN THE HEAD. Third Edition, with Illustrations, cr. 8vo. [*In preparation.*]

II.

NASAL POLYPUS: WITH NEURALGIA, HAY-FEVER, AND ASTHMA, IN RELATION TO ETHMOIDITIS. With Illustrations, cr. 8vo, 4s. 6d.

DAVID YOUNG, M.C., M.B., M.D.

Licentiate of the Royal College of Physicians, Edinburgh; Licentiate of the Royal College of Surgeons, Edinburgh, etc.

ROME IN WINTER AND THE TUSCAN HILLS IN SUMMER. A CONTRIBUTION TO THE CLIMATE OF ITALY. Small 8vo, 6s.

HERMANN VON ZEISSL, M.D.

Late Professor at the Imperial Royal University of Vienna

OUTLINES OF THE PATHOLOGY AND TREATMENT OF SYPHILIS AND ALLIED VENEREAL DISEASES. Second Edition, revised by M. VON ZEISSL, M.D., Privat-Dozent for Diseases of the Skin and Syphilis at the Imperial Royal University of Vienna. Translated, with Notes, by H. RAPHAEL, M.D., Attending Physician for Diseases of Genito-Urinary Organs and Syphilis, Bellevue Hospital, Out-Patient Department. Large 8vo, 18s.

Clinical Charts for Temperature Observations, etc.

Arranged by W. RIGDEN, M.R.C.S. 50s. per 1000, 28s. per 500, 15s. per 250, 7s. per 100, or 1s. per dozen.

Each Chart is arranged for four weeks, and is ruled at the back for making notes of Cases; they are convenient in size, and are suitable both for hospital and private practice.

Lewis's Clinical Chart, specially designed for use with the Visiting List. This Temperature Chart is arranged for four weeks and measures 6 X 3 inches. 30s. per 1000, 16s. 6d. per 500, 3s. 6d. per 100, 1s. per 25, 6d. per 12.

Lewis's Nursing Chart.

25s. per 1000, 14s. per 500, 3s. 6d. per 100, 2s. per 50, or 1s. per 20.

These Charts afford a ready method of recording the progress of the case from day to day.

Boards to hold the Charts, price 1s.

LEWIS'S PRACTICAL SERIES.

Under this title Mr. LEWIS is publishing a Series of Monographs, embracing the various branches of Medicine and Surgery.

The volumes are written by well-known Hospital Physicians and Surgeons, recognized as authorities in the subjects of which they treat. The works are intended to be of a THOROUGHLY PRACTICAL nature, calculated to meet the requirements of the practitioner and student, and to present the most recent information in a compact and readable form.

HYGIENE AND PUBLIC HEALTH.

By LOUIS C. PARKES, M.D., D.P.H. LOND. UNIV., Fellow of the Sanitary Institute, and Member of the Board of Examiners; Assistant Professor of Hygiene and Public Health at University College, etc. Second edition, with numerous Illustrations, cr. 8vo, 9s. *[Just published.]*

MANUAL OF OPHTHALMIC PRACTICE.

By C. HIGGENS, F.R.C.S., Ophthalmic Surgeon to Guy's Hospital; Lecturer on Ophthalmology at Guy's Hospital Medical School. With Illustrations, crown 8vo, 6s.

A PRACTICAL TEXTBOOK OF THE DISEASES OF WOMEN.

By ARTHUR H. N. LEWERS, M.D. Lond., M.R.C.P. Lond., Assistant Obstetric Physician to the London Hospital; Examiner in Midwifery and Diseases of Women to the Society of Apothecaries of London, etc. Third Edition, with Illustrations, crown 8vo, 10s. 6d. *[Now ready.]*

ANÆSTHETICS THEIR USES AND ADMINISTRATION.

By DUDLEY W. BUXTON, M.D., B.S., M.R.C.P., Administrator of Anæsthetics in University College Hospital and the Hospital for Women, Soho Square. Second Edition, crown 8vo. *[In the press.]*

TREATMENT OF DISEASE IN CHILDREN: EMBODYING THE OUTLINES OF DIAGNOSIS AND THE CHIEF PATHOLOGICAL DIFFERENCES BETWEEN CHILDREN AND ADULTS. By ANGEL MONEY, M.D., F.R.C.P., Assistant Physician to the Hospital for Sick Children, Great Ormond Street, and to University College Hospital. Second edition, cr. 8vo, 10s. 6d.

ON FEVERS: THEIR HISTORY, ETIOLOGY, DIAGNOSIS, PROGNOSIS, AND TREATMENT. By ALEXANDER COLLIE, M.D. Aberd., Member of the Royal College of Physicians of London; Secretary of the Epidemiological Society for Germany and Russia. Illustrated with Coloured Plates, crown 8vo, 8s. 6d.

HANDBOOK OF DISEASES OF THE EAR FOR THE USE OF STUDENTS AND PRACTITIONERS. By URBAN PRITCHARD, M.D. Edin., F.R.C.S. Eng., Professor of Aural Surgery at King's College, London; Aural Surgeon to King's College Hospital; Senior Surgeon to the Royal Ear Hospital. Second Edition, with Illustrations, crown 8vo, 5s. *[Now ready.]*

A PRACTICAL TREATISE ON DISEASES OF THE KIDNEYS AND URINARY DERANGEMENTS. By CHARLES HENRY RALFE, M.A., M.D. Cantab., Fellow of the Royal College of Physicians, London; Assistant Physician to the London Hospital; Examiner in Medicine to the University of Durham, etc., etc. With Illustrations, crown 8vo, 10s. 6d.

DENTAL SURGERY FOR MEDICAL PRACTITIONERS AND STUDENTS OF MEDICINE. By ASHLEY W. BARRETT, M.B. Lond., M.R.C.S., L.S.D., Dental Surgeon to, and Lecturer on Dental Surgery in the Medical School of, the London Hospital. Second edition, with Illustrations, cr. 8vo, 3s. 6d.

BODILY DEFORMITIES AND THEIR TREATMENT: A HANDBOOK OF PRACTICAL ORTHOPÆDICS. By H. A. REEVES, F.R.C.S. Edin., Senior Assistant Surgeon and Teacher of Practical Surgery at the London Hospital; Surgeon to the Royal Orthopædic Hospital, &c. With numerous Illustrations, cr. 8vo, 8s. 6d.

Further volumes will be announced in due course.

THE NEW SYDENHAM SOCIETY'S PUBLICATIONS.

President :—SIR JAMES PAGET, BART., F.R.S.
Honorary Secretary :—JONATHAN HUTCHINSON, ESQ., F.R.S.
Treasurer :—W. SEDGWICK SAUNDERS, M.D., F.S.A.

Annual Subscription, One Guinea.

The Society issues translations of recent standard works by continental authors on subjects of general interest to the profession.

Amongst works recently issued are "Flügge's Micro-Organisms," "Cohnheim's Pathology," "Henoch's Children," "Spiegelberg's Midwifery," "Hirsch's Historical and Geographical Pathology," "Ewald's Disorders of Digestion," works by Charcot, Duchenne, Begbie, Billroth, Graves, Koch, Hebra, Guttman, etc.

The Society also has in hand an Atlas of Pathology with Coloured Plates, and a valuable and exhaustive "Lexicon of Medicine and the Allied Sciences."

The Annual Report, with full list of works published, and all further information will be sent on application.

PERIODICAL WORKS PUBLISHED BY H. K. LEWIS.

THE BRITISH JOURNAL OF DERMATOLOGY. Edited by H. G. Brooke, H. Radcliffe Crocker, T. Colcott Fox, Malcolm Morris, J. F. Payne and J. J. Pringle. Published monthly, 1s. Annual Subscription 12s. post free.

THE NEW YORK MEDICAL JOURNAL. A Weekly Review of Medicine. Annual Subscription, Thirty Shillings, post free.

THE THERAPEUTIC GAZETTE. A Monthly Journal, devoted to the Science of Pharmacology, and to the introduction of New Therapeutic Agents. Edited by Dr. R. M. Smith. Annual Subscription, 10s., post free.

THE GLASGOW MEDICAL JOURNAL. Published Monthly. Annual Subscription 20s., post free. Single numbers, 2s. each.

LIVERPOOL MEDICO-CHIRURGICAL JOURNAL, including the Proceedings of the Liverpool Medical Institution. Published twice yearly, 3s. 6d. each number.

TRANSACTIONS OF THE COLLEGE OF PHYSICIANS OF PHILADELPHIA. Volumes I. to VI., 8vo, 10s. 6d. each.

MIDDLESEX HOSPITAL, REPORTS OF THE MEDICAL, SURGICAL, AND Pathological Registrars for 1883 to 1888. Demy 8vo, 2s. 6d. *nett* each volume.

* * MR. LEWIS is in constant communication with the leading publishing firms in America, and has transactions with them for the sale of his publications in that country. Advantageous arrangements are made in the interests of Authors for the publishing of their works in the United States.

Mr. Lewis's publications can be procured of all Booksellers in any part of the world.

QR

41

C8

1890

BIOL. LIB.

