




AVERAGES (%) OF FOODS CONTAINING LOW LEVELS OF NEGATIVE FOOD COMPONENTS (TO LIMIT INTAKE OF ANY NEGATIVE FOOD COMPONENT) BASED ON THE PROPOSED METHOD IN FOOD GROUPS



References:

- Forouzes A, Forouzes F, Samadi Foroushani S, Forouzes A. A new method for calculating cholesterol and saturated fat contents and determining appropriate cholesterol levels in foods. SSRN 2022. DOI: 10.2139/ssrn.4131337
- Forouzes A, Forouzes F, Samadi Foroushani S, Forouzes A. A new method for calculating energy content and determining appropriate energy levels in foods. SSRN 2022. DOI: 10.2139/ssrn.4132581
- Forouzes A, Forouzes F, Samadi Foroushani S, Forouzes A. A new method for calculating fat content and determining appropriate fat levels in foods. Iran J Public Health 2023;52:1038–1047. DOI: 10.18502/ijph.v52i5.12722
- Forouzes A, Forouzes F, Samadi Foroushani S, Forouzes A. A new method for calculating saturated fat content and determining appropriate saturated fat levels in foods. SSRN 2022. DOI: 10.2139/ssrn.4133571
- Forouzes A, Forouzes F, Samadi Foroushani S, Forouzes A. A new method for calculating sodium content and determining appropriate sodium levels in foods. SSRN 2022. DOI: 10.2139/ssrn.4133574
- Forouzes A, Forouzes F, Samadi Foroushani S, Forouzes A. A new method for calculating sugars content and determining appropriate sugars levels in foods. SSRN 2022. DOI: 10.2139/ssrn.4133577