



Digitized by the Internet Archive
in 2018 with funding from
Wellcome Library

https://archive.org/details/b30456290_0001

23,769/D

M. Ant. Radneri

A
S Y S T E M
OF THE
A N A T O M Y
OF THE
H U M A N B O D Y.

W. L. G. & C.

W. L. G. & C.

W. L. G. & C.

W. L. G. & C.

W. L. G. & C.

A
S Y S T E M
OF THE
ANATOMY
OF THE
H U M A N B O D Y;

ILLUSTRATED BY

UPWARDS OF TWO HUNDRED TABLES, CONTAINING NEAR A THOUSAND
FIGURES,

COPIED FROM

THE MOST CELEBRATED AUTHORS,

AND

FROM NATURE.

By ANDREW FYFE.

IN THREE VOLUMES.

SECOND EDITION, WITH ALTERATIONS AND IMPROVEMENTS.

VOL. I.

EDINBURGH:

PRINTED BY J. PILLANS & SONS.

SOLD BY THE AUTHOR, EDINBURGH;
BY LONGMAN, HURST, REES, & ORME; AND BY T. KAY, T. COX,
AND J. MURRAY, LONDON.

1805.

МЕТЗУЗ

УМОТАМА

УДОДИАМУН



СЛАВЯНСКАЯ

ИСТОРИЯ МЕДИЦИНЫ

1884

№ 1

ТОМ I

ИЗДАНИЕ

САНКТ-ПЕТЕРБУРГ

1884

ИЗДАТЕЛЬСТВО

С. ПЕТЕРБУРГ

TO THE
GENTLEMEN

attending the MEDICAL CLASSES of the UNIVERSITY of
EDINBURGH,

THE FOLLOWING WORK

IS DEDICATED,

With much respect,

And with best wishes,

By their most obedient,

And very humble Servant,

UNIVERSITY OF EDINBURGH,
NOVEMBER 1. 1805.

ANDREW FYFE.

CONTENTS

OF THE

FIRST VOLUME.

PART I.

OF THE BONES.

	<i>Pag.</i>		<i>Pag.</i>
OF THE BONES IN GENERAL,	3	<i>The FALSE VERTEBRÆ,</i>	51
OF THE SKELETON,	8	BONES of the PELVIS,	52
OF THE HEAD OR SKULL IN GENERAL,	ib.	————— THORAX,	58
OF THE SEPARATE BONES OF THE HEAD,	18	BONES OF THE SUPERIOR EXTREMITY,	62
OF THE BONES OF THE FACE,	33	————— INFERIOR EXTREMITY,	74
OS HYOIDES,	45	<i>Ossa Sesamoidea,</i>	79
BONES OF THE TRUNK,	ib.	<i>Principal Differences between the MALE and FEMALE</i>	
<i>The TRUE VERTEBRÆ,</i>	ib.	SKELETON,	86

PART II.

OF THE MUSCLES.

OF THE MUSCLES IN GENERAL,	101	MUSCLES of the FEMALE PARTS of GENERATION	
MUSCLES of the INTEGUMENTS of the CRANIUM, and		<i>and of the ANUS,</i>	112
<i>of the EYE-LIDS,</i>	ib.	MUSCLE of the Os COCCYGIS,	ib.
————— <i>common to the HEAD and EXTERNAL EAR,</i>	102	MUSCLES situated within the CAVITY of the ABDOMEN,	ib.
————— <i>of the NOSE and MOUTH,</i>	ib.	————— <i>situated upon the Anterior Part of the THORAX,</i>	113
————— <i>of the LOWER JAW,</i>	104	————— <i>situated between the RIBS, and within the THORAX,</i>	114
————— <i>of the Fore and Lateral Part of the NECK,</i>	ib.	————— <i>situated on the Anterior Part of the VERTEBRÆ of the NECK,</i>	115
————— <i>situated between the Os HYOIDES and TRUNK,</i>	105	————— <i>situated on the Posterior Part of the TRUNK,</i>	ib.
————— <i>situated between the LOWER JAW and Os HYOIDES,</i>	ib.	————— of the SUPERIOR EXTREMITY,	119
————— <i>situated upon the Back Part of the PHARYNX,</i>	107	MUSCLES arising from the SCAPULA,	ib.
————— <i>of the GLOTTIS,</i>	ib.	————— <i>chiefly situated on the ARM, serving for the Motion of the FORE-ARM,</i>	120
————— <i>situated on the Anterior and Lateral Parts of the ABDOMEN,</i>	109	MUSCLES	
————— of the MALE PARTS of GENERATION and of the ANUS,	111		

	<i>Pag.</i>		<i>Pag.</i>
MUSCLES on the FORE-ARM and HAND, serving for the Motion of the HAND and FINGERS,	121	ving for the Motion of the THIGH and LEG,	125
MUSCLES of the INFERIOR EXTREMITY,	125	MUSCLES situated on the LEG and FOOT, serving for the Motion of the FOOT and TOES,	128
MUSCLES on the PELVIS and THIGH, ser-			

PART III.

OF THE BURSÆ MUCOSÆ.—OF THE LIGAMENTS AND OTHER PARTS OF THE JOINTS:


OF THE BURSÆ MUCOSÆ,	163	CARPAL BONE of the THUMB, and at the FIRST JOINT of the FINGERS,	174
OF THE LIGAMENTS, &c. OF THE JOINTS,	168	LIGAMENTS at the FIRST and SECOND JOINTS of the THUMB, and SECOND and THIRD JOINTS of the FINGERS,	ib.
SYNOVIAL ORGANS, commonly called GLANDS of the JOINTS,	ib.	retaining the TENDONS of the MUSCLES of the HAND and FINGERS in situ,	ib.
LIGAMENTS of the HEAD and TRUNK,	169	LIGAMENTS of the INFERIOR EXTREMITY,	175
LIGAMENTS of the LOWER JAW,	ib.	LIGAMENTS connecting the Os FEMORIS with the Os INNOMINATUM, ib.	
connecting the HEAD with the FIRST and SECOND VERTEBRÆ of the NECK, and these two VERTEBRÆ with each other,	ib.	of the JOINT of the KNEE,	ib.
of the other VERTEBRÆ,	ib.	connecting the FIBULA to the TIBIA,	176
of the RIBS,	170	connecting the BONES of the TARSUS with those of the LEG,	ib.
of the BONES of the PELVIS,	171	of the TARSUS,	177
of the SUPERIOR EXTREMITY,	172	between the TARSUS and METATARSUS,	ib.
LIGAMENTS of the CLAVICLE,	ib.	connecting the METATAR- SAL BONES to each other,	ib.
proper to the SCAPULA,	ib.	of the PHALANGES of the TOES,	ib.
of the JOINT of the SHOUL- DER,	ib.	and SHIATHS retaining the TENDONS of the MUSCLES of the FOOT and TOES in situ,	ib.
of the JOINT of the ELBOW,	173		
between the BODIES, and be- tween the Under Ends of the RADIUS and ULNA,	ib.		
between the FORE-ARM and WRIST,	ib.		
of the CARPUS,	174		
between the CARPAL and METACARPAL BONES,	ib.		
between the EXTREMITIES of the METACARPAL BONES,	174		
LIGAMENTS at the Base of the META-			

P A R T I.

OF

T H E B O N E S.

OF THE BONES IN GENERAL.



THE Bones are the most hard, compact, and inflexible parts of the body.

They are more or less of a *white* or *red* colour, according to the proportions of *Earth* or *Blood* entering their composition; and are therefore whitest in the Adult, and reddest in the Child, more Earth being found in the former, and more Blood in the latter.

In living Animals they are of a bluish colour, in consequence of the Blood contained in their small vessels appearing through their surface.

Bones are composed of *Lamellæ*, or *Plates*, which are formed of Fibres running longitudinally, or in a radiated manner, according to the natural figure of the Bone.

The lamellated structure may be seen, by exposing them to the heat of a strong fire; or to the weather; or by boiling them under an increased pressure; or by observing their exfoliations when in the diseased state.

A late Author, Scarpa, denies the lamellated structure of the Bones, and endeavours to prove, that they have every where a cellular texture.

The *Plates* of Bones are originally formed by the Vessels of the Periosteum Externum and Membrana Medullaris, and not, as has been supposed by some Authors, from Layers detached from the external Periosteum.

The Plates are connected by *Fibres*, which some have considered as *Claviculi* or *Nails*, and called *Perpendicular*, *Oblique*, &c. according to their different directions.

The *outer Plates* of Bones are firmly compacted, so as to appear like one solid substance.

The *inner Parts* of Bones in general, whether long, round, or flat, have their Plates and Threads running in various directions, intersecting each other, and forming the *Cancelli*, or *Spongy Substance* of the Bones; the *Cancelli* every where communicating with each other.

The *Cancelli*, in the middle of long Bones, are Fibrous, and form the *Reticular Substance* which divides the Bone into large cells.

Towards the extremities of long Bones, the *Cancelli* are lamellated, and much more numerous than in their middle.

Cancelli, of a similar nature to those of the long Bones, are also placed between the Tables of flat, and inner parts of round Bones.

In some of the broad Bones, however, as the Scapula, the solid parts are so much compressed, as to leave little or no room for *Cancelli*.

On the contrary, in the middle of the long Bones, as the Os Humeri, the *Cavities* are so large as to give to the Bone the appearance of a *hollow Cylinder*.

In some of the largest of the long Bones, as the Os Femoris, their solid sides, near their middle, are remarkably thick, and there the *Cancelli* are scarcely perceptible; while at their extremities, their sides are scarcely thicker than writing paper, and their *Cancelli* are so numerous as to occupy the whole space between their sides.

The *Cancelli* of Bones are formed by the internal Plates passing inwards, and decussating each other; and in the long Bones, the sides of the Bone, in consequence of sending off the *Cancelli*, become gradually thinner towards its extremities, while the *Cancelli* in proportion become more numerous.

The *Cancelli*, though extremely minute, exist even in the most solid parts of Bones, as can be seen by exposure to heat, or in Bones enlarged by disease. In either of these cases, small cells may be observed, and are distinguishable from the Canals for containing the Vessels, the former being irregular, and the latter cylindrical.

The *Cancelli* support the Membranes containing the Marrow, as the Cellular Substance does the Fat, and prevent one part of the column of Marrow from gravitating upon another in the various positions of the body. They also furnish a wider surface for the dispersion of the Arteries which secrete the Marrow.

Upon the *surface* of Bones there are numerous *Fissures*, for the more intimate connection of the Periosteum with the Bone, and for lodgement to Blood-vessels, which pass into its Substance.

Many minute *Orifices* are observed upon the surface, and particularly in the Furrows of Bones, for the transmission of Blood-vessels into their substance.

Near the middle of most of the Bones, especially the long ones, there is a slanting *Canal* for the passage of the principal Medullary Vessels.

Numerous *Orifices*, some of them very considerable in size, are observed at the *extremities* of long Bones.

Some serve for the transmission of Blood-vessels, and others for giving attachment to the Fibres of the Ligaments of the Joints.

The principal Vessels pass into the Cancelli, internal Membranes and Marrow, and return to the solid Substance of the Bone, where they meet those sent inwards from the Periosteum.

In some *flat Bones*, as those of the *Cranium*, the Bones are entirely supplied from the Vessels of the surrounding Membranes, and the Vascularity there is uniform.

In the Subject, the Blood-vessels of the Bones can be shewn by a successful injection thrown into them; and in living Animals, when the Bones are cut across, their vascularity appears by the blood which oozes from their divided extremities.

The Vascularity of Bone is also shewn, by feeding an Animal, for some time, on the Rubia Tinctorium, or Madder-root, after which the Bones are found to be completely tinged with the colouring matter of the Madder.

As a person advances in life, the Blood-vessels of the Bones contract in their Diameters, as appears from the Bones of old people having less Blood in them, than those of a person at an early period of life; from Injections being thrown into the Vessels of the Bones of old persons with more difficulty than in youth; from less of the injected matter being received in the former; and from the Bones of old animals receiving less of the tinging matter of Madder than those of young ones.

From comparing the Bones of people of different ages, it is found, that there is a constant waste and renewal of their Substance; that the Bones increase in weight as a person advances to maturity; that they continue nearly of the same weight till old age begins, and then become lighter; that the specific gravity of their solid sides, on the contrary, increases by age; for then they become harder and more compact, but thinner, and have larger cavities than the Bones of young persons.

Bones, like other parts, have their *Lymphatics*, as appears by the absorption of Madder found deposited in the Substance of the Bones of Animals receiving it with their food; by the absorption of part of the Bone itself, when in the diseased state; by the absorption of Bone as a person advances in life; and even by Injection.

The *Nerves* of the Bones are small, but may be observed in certain parts of the Bones, and, it is presumed, they exist in all.

From the minuteness of the Nerves, and rigidity of the parts on which they are dispersed, Bones are not *sensible* in the sound state; and even in the diseased, the pain felt may be owing to the Membranes within them.

The component parts of Bones are, an Earthy Matter, Cartilage, Gelatin, and Marrow, and these varying in proportion in different persons, in different Bones of the same person, and in the same Bone at different ages.

The earthy matter is obtained by Calcination, or by Maceration in a diluted acid, and afterwards precipitating

it, when it is found to consist chiefly of Phosphat of Lime. In either of these Processes, the Bone retains its shape; but in the former it is brittle, and of a pure white Colour, while in the latter it is flexible, consisting principally of Cartilaginous Matter.

By boiling in water for a sufficient length of time, and especially if under an increased pressure, as Papin's Digester, the Fat and Gelatin of Bones are dissolved and separated, and the Bone retaining its earthy matter, preserves also its white Colour.

The general *Use* of Bones is,—to furnish attachment to the Muscles, and to lodge, protect, and support the Bowels.

PERIOSTEUM.

The *Periosteum* derives its name from its furnishing a general Covering to the Bones.

In certain parts, however, it is *perforated* by Muscles, Ligaments, or Cartilages, which are fixed immediately to the surface of the Bones; and at the Joints it separates from the Bones, to give a Covering to the Capsular Ligaments.

It is *formed* of many Fibres, which, in certain parts, can be divided into Layers.

The *outer Surface* of this Membrane is connected to the surrounding parts by cellular Substance.

The *inner Surface* is more uniform than the outer, and its Fibres run, most frequently, in the same direction with those of the subjacent Bone.

The *inner part* of the Periosteum is intimately connected to the surface of the Bones by short Fibres; and this Connection is much stronger in the Child than in the Adult. Some of these Fibres may be considered as ligamentous, but most of them are found by Injection to consist of Blood-vessels.

The Periosteum, as well as other Membranes, must be supplied with *Nerves*; but these are too minute to be readily traced.

The *Sensibility* of the Periosteum, like that of other Membranes, is by no means acute. In the inflamed state, its sensibility is very considerable.

The *principal Uses* of this Membrane are,—to transmit the Vessels which are spread out upon its surface into the Substance of the Bones;—to give attachment to Muscles;—to prevent the effects of Friction between them and the Bones;—to assist in binding the latter together;—to assist in setting limits to the increase, and to check the overgrowth of Bones;—and in young persons, to strengthen the junction of the Bones with their Epiphyses, Cartilages, and Ligaments.

MEMBRANA MEDULLARIS.

This, improperly called Periosteum Internum, is an extremely fine Membrane, which lines the inside of the Bones in general, and is divided into numberless small parts,

parts, which also line the different Cancelli. It forms so many irregular Bags, communicating with each other, and affording a large surface for the dispersion of the secretory Vessels of the Marrow.

MARROW.

The *Marrow* may be considered as an *Appendage* to the general *Corpus Adiposum*. It is deposited by the Arteries in the Cavities of the Bones, at the same time that the rest of the Body is supplied with Fat.

The Blood-vessels of the Marrow, surrounded by the Periosteum, enter the Bones by oblique Canals, which have already been taken notice of in the description of the Bones in general.

When the Arteries have entered the Cavities of the Bones, they divide into Branches, which are spread out upon the Cancelli, *Membrana Medullaris*, and Marrow; from these many minute Branches are reflected outwards, to the Tables of the Bones which communicate with those sent from the inner surface of the Periosteum.

The Veins which return the Blood from the Marrow and Substance of the Bones, are collected into small Trunks, which pass out where the Arteries penetrated the Bones, and discharge their contents into the neighbouring Veins.

The greater degree of Vascularity of the Solids in Children than in Adults, is no where more conspicuous than here; for Injections which pass readily in these Vessels in Children, cannot be made to penetrate so far in those of persons of an advanced time of life. In consequence of which the Marrow is found to be thin and bloody in Children, oily and thick in Adults, and watery in old people.

The Marrow, like the Fat, when viewed through a Microscope, resembles a cluster of Pearls;—or it is contained in spherical Sacs, upon which Vessels are minutely dispersed, but from which no Excretory Ducts have been discovered.

It possesses little *Sensibility* in the sound state; and what it does possess, is considered by the latest Authors as belonging rather to its Membranes than to the Marrow itself.

But that this part of the Body is not without Nerves, seems to be proved by the Experiments made on the Marrow, when the Bones of living Animals are cut, and by the pain a person frequently suffers from Diseases within the Bones.

CARTILAGES.

Cartilages are of a *white Colour*, of an *elastic Substance*, and much softer than Bones, in consequence of the smaller quantity of *Earth* entering their composition.

Their *Structure* is not so evidently Fibrous as that of Bones, yet, by long Maceration, or by tearing them asunder, a Fibrous disposition is perceptible.

Their *Vessels* are extremely small, though they can be readily injected in *Cartilages* where Bone is beginning

to form. The Vessels of the *Cartilages* of the Joints, seem entirely to exclude the red blood. No Anatomist has been able to inject them, and Matter, mixed with the Food of Animals, does not change their colour as it does that of Bones. They have no Cancelli, nor internal Membranes, for lodging Marrow; no Nerves can be traced to them; nor do they possess any Sensibility in the sound state. Yet the Granulations which rise on the surface of *Cartilages* after Amputation at the Joints is very sensible.

Upon their Surface, there is a thin Membrane termed *Perichondrium*, which, in *Cartilages* supplying the place of Bone, as in those of the Ribs, or at the ends of the long Bones in Children, is a continuation of the *Periosteum*, and serves the same general purposes to *Cartilage* as the *Periosteum* does to Bone.

The *Perichondrium* of *Cartilages*, which supply the place of Bone, or by their flexibility possess a degree of motion, has Blood-Vessels, which, like those of the *Periosteum*, can be injected. But the Vessels of this Membrane belonging to other *Cartilages*, particularly those covering the *Articular Cartilages*, cannot be injected.

Upon the surface of *Articular Cartilages*, the *Perichondrium* is a *Reflection* of the inner Surface of the *Capular Ligament*, and is so very thin, and adheres so closely, as to appear like part of the *Cartilage* itself.

They have no internal Cavity, nor Cancelli, nor internal Membrane; their weight is nearly a third less than that of Bone. Their texture is less changed by acids; but a much greater proportion of them than of Bones, is destroyed by the action of a strong fire.

One set of *Cartilages* supply the place of Bone;—or, by their flexibility, admit of a certain degree of motion, while their elasticity recovers their natural position;—as in the *Nose*, *Larynx*, *Cartilages of the Ribs*, *Cartilages supplying Brims to Cavities*, &c.

Another set, in Children, supply the place of Bone, until Bone is formed, and afford a *Nidus* for the *Oseous Fibres* to shoot in;—as in the *long Bones of Children*.

A third set, the most extensive, by the smoothness and slipperiness of their surface, allow the Bones to move readily, without any abrasion;—as, in the *Cartilages of the Joints*. By their elastic nature, they render the motions easier, and lessen the concussion in the more violent motions of the Body, as running, jumping, &c. They also prevent the inordinate growth of Bones at their articulating surfaces, and the coalescence of the *Fibres* of the adjoining Bones.

A fourth set supply the office both of *Cartilages* and *Ligament*, giving the elasticity of the former, and flexibility of the latter; uniting some immoveably together, and allowing to others a small degree of motion;—as in the *Cartilages of the Bones of the Pelvis and Spine*.

OF THE FORMATION OF BONE.

The generality of Bones are originally formed, either
between

between Membranes, or in the substance of Cartilages; the *Teeth* are formed in distinct Bags.

The Ossification of *broad Bones* begins in some, as in those of the Cranium, between Membranes only, and in others, as in the *Offa Ilii*, in Cartilage, and it appears in each Bone in one or more places: There the Osseous Particles are so joined together, as to have a fibrous appearance.

The Fibrous Structure is most distinctly seen in the Cranium of a Fœtus about three months after conception, where the beginning of the Ossification is like a fine irregular net-work, in the middle of which the Fibres are more closely connected than in the circumference.

In viewing the flat Bones of a Fœtus a little more advanced, the Bony particles are observed to be so disposed, as to have a distinct radiated appearance. Tab. XI.

The vacancies between the Fibres, which occasion the radiated appearance, are found by injection to be chiefly passages for Blood-vessels.

As the Fœtus increases in Size, the Osseous Fibres increase in number, till a Lamina is produced; and as the Bone continues to grow, more Laminæ are added, till the more solid part of a Bone is formed.

The Inner Layers of the Bones are observed to be more porous than the Outer, and none of them are found to have the solidity they acquire in the Adult state, till they have arrived at their full growth.

The Ossification of *long Bones* begins between the Periosteum and Membrana Medullaris, by a *central Ring*, from which the Fibres extend towards the ends of the Bones.

The Interior Lamellæ, forming the solid sides of the long Bones, are considerably shorter than the Exterior, because they pass gradually inwards to form the Cancelli, while the exterior parts are continued to the extremities of the Bones.

The Ossification of spherical-shaped Bones, as in the Wrist, begins by one Nucleus, and that of irregular-formed Bones, as in the *Vertebræ*, by different Nu-

clei, and both of these sets of Bones have their origin in Cartilage.

Some of the flat Bones also, as the *Offa Ilii*, and all the Epiphyses, have their original formation in Cartilage.

The Ossification which begins in Cartilage, is considerably later than that which has its origin between Membranes, and this is at very different times in different parts of the Body.

When Ossification is about to begin in a particular Cartilage, and which is most frequently in the centre, the Arteries, which were formerly transparent, become dilated, and receive the red blood from which the Osseous matter is secreted. This matter retains, for some time, the form of the Vessels which give it origin, till, more Arteries being by degrees dilated, and more Osseous matter deposited, the Bone at length attains its complete form.

Some Bones are completely formed at the time of Birth, as the *small Bones of the Ear*.

The generality of Bones are *incomplete* until the age of puberty, or between the fifteenth and twentieth year, and in some few instances not until a later period.

In Children, the greater number of parts in Bones are *Epiphyses* or *Appendices*, which, in Adults, become *Processes*.

The *Epiphyses* begin to appear after the Body of the Bone is ossified, and are themselves ossified at seven or eight years of age, though their external surface is still somewhat Cartilaginous.

In the early period of life, the Body and Ends of long Bones make *three distinct parts*, which can readily be separated by boiling, or by maceration in water.

The Epiphyses are joined to the Body of the Bone by Cartilages, which are thick in Children, but gradually become thinner as Ossification advances, till at last, in the Adult, the external marks of division are not to be seen; though frequently some mark of distinction may be observed in the Cancelli.

DIFFERENT KINDS

OF

CONNECTION OF BONES.

<p><i>SYNARTHROSIS</i>, Or Connection without inter- mediate Substance.</p>	<p><i>Suture</i>, Like a Seam.</p>	<p>{ The Bones of the Cranium, and greater part of those of the Upper Jaw with each other.</p>		
			<p><i>Gomphosis</i>, Like a Nail in a Board.</p>	<p>{ The Teeth in the Alveoli.</p>
<p><i>SYMPHYSIS</i>, Or Connection by interme- diate Substance.</p>	<p><i>Synchondrosis</i>, Or Connection by Cartilage.</p>	<p>{ The Bodies of the Vertebrae to each other: The Ribs to the Sternum: The Offa Innominata to the Os Sacrum, or to each other.</p>		
		<p><i>Syndesmosis</i>, Or Connection by Ligament.</p>	<p>{ The Lower Jaw and Os Hyoides to the Head: The Ribs to the Spine: The Proccesles of the Vertebrae, and also the Bones of the Extremities to each other.</p>	

DIFFERENT KINDS OF MOTION.

<p><i>ARTHRODIA</i>; Where the flat ends of Bones are opposed to each other with little Motion.</p>	<p>{ Between the Clavicle and Scapula. The Bones in the second row of the Car- pus. The Carpus and Metacarpus. The Tibia and Fibula. The greater number of Bones in the Tarsus. The Tarsus and Metatarsus.</p>	
<p><i>GINGLIMUS</i>; The Bones mutually receiving each other; and the Liga- ments admitting of a hing- like Motion.</p>		
	<p><i>Angular</i>. One Bone, in moving, forming an angle with another.</p>	<p>{ The Lower Jaw and Head. The Joint of the Elbow. The First and Second Joints of the Thumb, and Second and Third of the Fingers. The Joint of the Knee. The Ankle. The two last Joints of the Toes.</p>
	<p><i>Lateral or Circular</i>.</p>	<p>{ Between the First Vertebra and Proccesus Dentatus of the Second. Between the Radius and Ulna.</p>
	<p><i>Compound</i>.</p>	<p>{ Between the Occipital Bone and Atlas. Between the different Vertebrae. And between the Ribs and Ver- tebrae.</p>
<p><i>ENARTHROSIS</i>; Or Ball and Socket; the Li- gaments allowing Motion in all directions.</p>	<p>{ Inner end of the Clavicle. Head of the Os Humeri. Between the Fore-arm and Wrist, and between the two Rows of the Carpal-Bones. At the root of the Metacarpal Bone of the Thumb, and root of the first Phalanx of the Fingers. At the Head of the Thigh-Bone. Between the Astragalus and Os Naviculare, and at the root of the first Phalanx of the Toes.</p>	

OF THE SKELETON.

THOUGH the term *Skeleton* is applied to a variety of substances, yet, in Anatomy, it is always understood to signify the Bones of Animals, connected together in their natural situation, after the soft parts of the body in general are removed.

It is termed a *Natural Skeleton*, when the Bones are joined by their own Ligaments;

And an *Artificial Skeleton*, when joined by wire, &c.

Small subjects, and the Bones of those which are not fully ossified, are most conveniently prepared the first way; while the Bones of large Adult Animals, are more readily cleaned when single, and are easily restored to their natural situation.

In viewing the Bones in their natural situation in the Skeleton, scarcely any of them is observed to be placed in a perpendicular direction to another; yet in an erect posture, a perpendicular line from their common centre of gravity, falls in the middle of their common base. On this account, the Body is found to be as firmly supported, as if the axis of all the Bones had been a straight line perpendicular to the horizon, and much greater quickness, ease, and strength is given to the Body, in several of its most necessary motions.

The Human Skeleton is generally divided into *Head, Trunk, Superior and Inferior Extremities*.

OF THE HEAD OR SKULL IN GENERAL.

By the Head is meant all that part of the Skeleton which is placed above the first Bone of the Neck. It therefore comprehends the Cranium, and Bones of the Face.

The Cranium varies in shape in different Persons, according to the original form of the Brain upon which it is moulded, but the variety of shape has been supposed by some Authors to be increased by the different management of the heads of Children at an early period of life. From this the different shapes of the Skulls of People of different nations have been accounted for. The form, however, does not appear to be much affected by the management of the Head at an early period of infancy, since its characteristic marks are found to remain nearly the same, however much the customs in dress and general management may vary.

The Cranium forms a vaulted cavity for lodging and defending the Brain, with its Membranes, Vessels, and Nerves.

The *General Figure* of the upper part of the Cranium, compared to that of an Egg. Tab. VIII.

The Cranium is of a flat form at its sides; partly by the action of the Temporal Muscles. Tab. IV.

The flatness of this part of the Head is found to increase the sphere of vision, and to give a more advantageous situation for the Ears, that they may receive a greater quantity of sound, while they are less exposed to danger.

The *Surface* of the upper and outer part of the Cranium is smooth, where it is little affected by Muscles, and is covered with the Periosteum common to all the Bones, but in the Skull termed Pericranium.

The *under and outer Surface* of the Cranium, is *ir-*

regular, where it gives attachment to Muscles, &c. and passage to Vessels and Nerves. Tab. VI.

The *anterior and under* part of the Cranium *hollow*, to make part of the Orbits. Tab. III.

The *posterior* Surface of the Cranium *marked* by the insertion of Muscles arising from the back-part of the Trunk. Tab. VI.

The *upper and inner* Surface of the Cranium *hollow*, for lodging the Brain. Tab. VIII. Fig. 2.

The *under and inner* Surface of the Cranium, has many *unequal Cavities*, for lodging the Lobes and Appendages of the Brain and Cerebellum, and for allowing passage to the Vessels and Nerves of the Encephalon in general. Tab. VIII.

Along the inner side of the Cranium, are many Furrows, for the reception of the Blood-vessels of the Dura Mater. Tab. VIII. Fig. 2.

Upon the inner Surface of certain Crania, Sinuosities are observed, for lodging Luxuriations of the Brain; and here the Cranium is sometimes so thin, as to be rendered transparent; the two Tables being then closely compacted without any Cancelli.

In some Crania, Pits are seen of different figures and sizes, for lodging *Granulous Bodies* on the Dura Mater, termed *Glands of PACCHIONI*; or sometimes they are occupied by the meeting of large Veins of the Dura Mater. Here also there is often a want of Cancelli. Tab. VIII. Fig. 2.

The Bones of the Cranium are composed of two tables, with intermediate Cancelli, termed here *Diploe*, though nearly of the same nature with the Cancelli in other flat Bones. Tab. VIII.

The *External Table* of the Cranium is somewhat thicker

thicker than the *Internal*, which, from its thinness and consequent brittleness, is called *Vitreæ*. Tab. V. s, s.

The *Diploe*, or *Cancelli*, between the Tables, are more regular between the Bones of the upper than of the under part of the Cranium, where, in several of the hard Bones, they are not observable.

In the skulls of old subjects, the *Diploe* is often so obliterated, that scarcely any vestige of it can be seen.

In certain diseased Bones, the *Diploe* is of great thickness, while the tables of the Skull are thin like paper.

The Cranium is generally composed of *eight Bones*; *six* of which are said to be proper to the Cranium, and *two* common to it and the Face.

The *six* proper to the Cranium are,

The *Os Frontis*, placed in the fore-part of the Cranium. Tab. III. A.

The *two Ossa Parietalia*, placed in the upper and lateral parts of the Cranium. Tab. IV. B.

The *two Ossa Temporum*, placed in the under and lateral parts. Tab. IV. D.

The *Os Occipitis*, which forms the back, and some of the lower part of the Cranium. Tab. VII.

The *two Bones* common to the Cranium and Face are,

The *Os Ethmoides*, placed in the fore-part of the Base of the Cranium. Tab. V. C, C.

The *Os Sphenoides*, situated in the middle of the Base. Tab. V.

SUTURES.

The Bones of the Cranium have Seams or Sutures between them, which are five in number. Of these *three* are termed *True*, from having serrated appearances; and *two* are called *False* or *Squamous* Sutures, from the Bones which form them overlapping each other, as the scales of Fishes do.

The *three True Sutures* are,

The *Coronal Suture*, placed between the Frontal and Parietal Bones, and getting its name from this being the part where the Ancients wore their *Coronæ* or Garlands. It loses its serrated appearance near its terminations. Tab. IV. m.

The *Lambdoid Suture*, situated where the Occipital joins the Parietal and Temporal Bones. It begins some way below the Vertex, or Crown of the Head, from which its two legs extend obliquely downwards and to each side, in form of the Greek Λ . Tab. II. b. Tab. VII.

The parts of the Lambdoid Suture, placed between the Occipital and Temporal Bones, have little of the serrated appearance, and are called *Addimenta Suturae Lambdoidalis*.

The *Sagittal Suture*, situated between the Parietal Bones, and named from being extended between the middle of the Coronal and Lambdoid Sutures, as an Arrow is between the String and Bow. Tab. VIII. Fig. 1. d, d.

The Sagittal Suture is sometimes continued through the middle of the Frontal Bone to the Nose. This is said, by some Authors, to be more frequently in the Female than in the Male.

The *serrated appearance* of the true Sutures is seen distinctly on the outside of the Cranium only: on the

inside the Bones appear almost to be joined as straight lines. Tab. VIII. Fig. 1. 2.

In some skulls, the internal surface is found entire, while the Sutures are manifest without; the inner Plates meeting and coalescing sooner than the external.

As a person advances in life, the True Sutures begin to be obliterated, first on the outer, then on the inner side, till in very old age not a vestige of one of them is to be seen.

The *two False*, called also *Temporal Sutures*, placed between the upper Edge of the Temporal, and under Edge of the Parietal bones. Tab. IV. o.

Each of the Portions of the False Sutures, situated between the under and back part of the Parietal, and the upper and back part of the Temporal Bones, called by some *Additamentum Suturae Squamosæ*, and having in that part the true serrated appearance. Tab. IV.

Besides the Squamous Sutures here taken notice of, it is to be observed, that the term *Squamous* is also applied to all the Sutures on which the Temporal Muscle is placed; it therefore includes part of the Coronal and Sphenoid Sutures.

In the Sutures of the Cranium there are often Additional Bones, called *Ossa Triquetra*, from their being of a triangular form, and *Ossa Wormiana*, from Wormius, who, though not the Discoverer, gave a description of them.

The *Ossa Wormiana* vary much in figure, size, and number, and are occasionally found in the different Sutures, though most frequently in the middle of the Lambdoid.

Wherever they occur, the Sutures surrounding them are observed to be similar to the neighbouring Sutures; of course they are equally with them distinguished from Fractures of the Skull.

Between the Bones of the Cranium, and those of the Face, five Sutures are also found, and they are said to be common to these two sets of Bones. Parts, however, of these Sutures are only between the Bones of the Cranium. The Sutures here are,

The *Ethmoid Suture*, which surrounds the Ethmoid Bone. Tab. V. C, C.

The *Sphenoid Suture*, which surrounds the Sphenoid Bone. Tab. V. f, o.

The Ethmoid and Sphenoid Sutures in some parts assist in forming other Sutures, especially the Squamous and Transverse; and in other parts there is but one Suture common to these two Bones.

The *Transverse Suture*, which runs across the Orbits and Root of the Nose, between the Frontal, Malar, Sphenoid, Ethmoid, superior Maxillary, and Nasal Bones. Tab. III. q, e, f.

The *Zygomatic Sutures*, placed between the Temporal and Cheek Bones, and slanting obliquely downwards and backwards. Tab. IV. t.

The advantages derived from the Cranium being formed of different Bones and Sutures are, that the Sphenoidal figure is sooner completed;—that the Bones, which are at some distance from each other at birth, yield, and conduce to an easier Delivery;—that the Dura Mater, by the Sutures, has a firmer adhesion;—and that Fractures are frequently prevented from extending so far as they would do in one continued bony substance.



T A B L E I.

REPRESENTS a Front View of the MALE SKELETON, with some of the CARTILAGES and LIGAMENTS which connect the BONES to each other.

HEAD and NECK.

- A, The frontal bone.
- B, The parietal bone.
- C, The temporal process of the sphenoid bone.
- D, The squamous part of the temporal bone.
- E, The mastoid process of that bone.
- F, The malar, or cheek-bone.
- G, The nasal bone; behind which is the nasal process of,
- H, The superior maxillary bone.
- I, The lower jaw.
- K, The cervical vertebræ, with their intermediate cartilages and transverse processes.

TRUNK.

- A, The sternum.
- B, The seventh, or last true rib.
- C, The cartilages of the ribs.
- D, The twelfth, or last false rib.
- E, The lumbar vertebræ, with their intervertebral cartilages and transverse processes.
- F, The os sacrum.
- G, The os innominatum, composed of,
 - a, The os ilium,
 - b, The os pubis,
 - c, The os ischium.

UPPER EXTREMITY.

- A, The clavicle.
- B, The inner surface of the scapula.
 - a, The acromion of the scapula.
 - b, The coracoid process of that bone.
- C, The os humeri.
 - c, The head or ball of the os humeri, articulated with the glenoid cavity of the scapula.
 - d, The internal tubercle of the os humeri, and farther out, the groove for lodging the tendon of the long head of the biceps muscle.

- e, The inner, and,
- f, The outer condyle of the os humeri. Between e and f, The hollow for lodging the coronoid process of the ulna in the flexion of the fore-arm.
- D, The radius.
 - g, The head of the radius.
- E, The ulna.
 - b, The coronoid process of the ulna.
- F, The bones of the carpus.
- G, The metacarpal bone of the thumb.
- H, The metacarpal bones of the fingers.
- I, The two bones of the thumb.
- K, The three phalanges of the fingers.

UNDER EXTREMITY.

- A, The os femoris.
 - d, The ball or head of this bone, lodged in the acetabulum.
 - e, The cervix of the bone.
 - f, The large trochanter.
 - g, The small trochanter.
 - b, The inner condyle.
 - i, The outer condyle.
- B, The patella, placed upon the trochlea of the os femoris.
- C, The tibia.
 - k, The head of the tibia. Between the head of the tibia, and condyles of the os femoris, the femilunar cartilages appear.
 - l, The tubercle of the tibia.
- m, The malleolus internus.
- D, The fibula; the upper end of which is connected with the tibia.
 - n, The malleolus externus.
- E, The bones of the tarsus.
 - o, The projection of the os calcis.
- F, The metatarsal bones.
- G, The phalanges of the toes.

T A B L E II.

REPRESENTS a Back-View of the MALE SKELETON, with some of the CARTILAGES and LIGAMENTS which connect the BONES to each other,

HEAD and TRUNK.

- A, THE parietal bone.
a, The sagittal future, and parietal hole.
 B, The occipital bone.
b, b, The lambdoid future.
 C, The joining of the temporal and parietal bones.
 D, The cheek-bone.
 E, F, The inner or back part of the jaws, with the teeth.
 G, The first cervical vertebra.
 H, The second cervical vertebra.
 I, The seventh cervical vertebra.
c, The spinous processes of the cervical vertebræ.
 K, The first dorsal vertebra.
 L, The twelfth dorsal vertebra.
d, The spinous processes of the dorsal vertebræ.
e, Their transverse processes.
 M, The first lumbar vertebra:
 N, The fifth lumbar vertebra.
f, Their spinous, and,
g, Their transverse processes.
 O, The os sacrum.
b, The uppermost spinous process. Farther out are seen the superior oblique processes of this bone, joined to the inferior oblique of the last lumbar vertebra.
i, i, The lateral parts of the os sacrum, joined to the ossa innominata. Between *i* and O, the posterior foramina of the os sacrum.
k, An opening in the under and back part of this bone, covered in the subject by a ligamentous membrane.
 P, The os coccygis, joined by its shoulders to the os sacrum, at the lower part of the opening *k*.
 Q, The os ilium.
 R, The os pubis.
 S, The os ischium.
 T, U, The seven true ribs.
 V, V, The five false ribs.

SUPERIOR EXTREMITY.

- A, The clavicle.
 B, The dorsum scapulæ.
a, The spine of the scapula.
b, The acromion of the scapula.
c, A fossa for lodging the supra-spinatus muscle.


- d*, An irregular surface, occupied by the infra-spinatus muscle.
 C, The os humeri.
e, The ball of the os humeri.
f, The external tubercle of the bone.
g, The external condyle.
h, The internal condyle.
i, The cavity for lodging the olecranon of the ulna.
 D, The radius.
k, The head of the radius articulated with the trochlea of the os humeri.
l, The under end of the radius, grooved by the tendons of muscles.
 E, The ulna.
m, The olecranon of the ulna.
n, The under end of the ulna, with the styloid process.
 F, The bones of the carpus.
 G, The metacarpal bone of the thumb.
 H, The metacarpal bones of the fingers.
 I, The two bones of the thumb.
 K, The three phalanges of the fingers.

INFERIOR EXTREMITY.

- A, The os femoris.
a, Part of the ball of the os femoris.
b, The cervix of the bone.
c, The trochanter major.
d, The trochanter minor.
e, The cavity for lodging the popliteal vessels and nerves.
f, The external condyle.
g, The internal condyle.
h, The femilunar cartilages.
 B, The tibia.
i, The head of the tibia.
k, The malleolus internus.
 C, The fibula.
l, The head of the fibula.
m, The malleolus externus.
 D, The bones of the tarsus.
n, The astragalus.
o, The os calcis.
p, The fore-part of the tarsus.
 E, The bones of the metatarsus.
 F, The phalanges of the toes.



TAB. 3.



T A B L E III.


A FRONT-VIEW of the SKULL.

-
- A, THE frontal bone.
a, The temporal process, or ridge of the frontal bone.
b, The temporal fossa of that bone.
c, The superciliary ridge.
d, The foramen superciliare.
e, e, The external and internal orbital processes.
f, e, e, The transverse suture.
g, The orbital plate.
h, The lacrymal fossa.
 B, The under and fore part of the parietal bone.
 C, The squamous part of the temporal bone.
i, The zygomatic process of that bone.
 D, The pars plana of the ethmoid bone.
k, k, The ossa spongiosa superiora of the ethmoid bone.
 E, The temporal plate of the sphenoid bone.
l, Part of the squamous suture.
m, The orbital plate of the sphenoid bone.
n, The foramen opticum.
o, The foramen lacerum of the sphenoid bone, and the foramen lacerum inferius of the orbit, the former above, and the latter below.
 F, The os nasi; in the middle of which is a hole proper to this bone.
 G, The os unguis.
p, The lacrymal groove of the os unguis.
- H, The os malæ.
q, q, q, q, The four angles, or processes of this bone.
r, The internal orbital process.
s, The external orbital hole.
 I, The fossa of the os maxillare superius.
t, The base of the nasal process of the superior maxillary bone, where there is frequently such a hole as is marked in this figure.
u, The tuberosity at the back-part of the bone.
v, The connection between the os malæ and os maxillare superius.
w, The connection between the superior maxillary bones.
x, The alveolar processes, with the teeth.
y, The fossa nasalis.
z, The foramen infra-orbitarium.
 K, The os spongiosum inferius.
 L, The vomer.
 M, The symphysis or middle of the lower jaw.
 N, The base of the lower jaw;
 1. Its angle.
 2. The ascending plate which sends off the coronoid and condyloid processes.
 3. The alveolar processes and teeth.
 4. The mental hole.

T A B L E IV.

A PROFILE of the SKULL.

-
- | | |
|--|--|
| <p>A, THE os frontis.
 <i>a</i>, The temporal process of this bone.
 <i>b</i>, The temporal fossa of the bone.
 <i>c</i>, The superciliary ridge.
 <i>d</i>, An elevation formed by the frontal sinus.
 <i>e</i>, The external and internal orbital processes.
 <i>f</i>, The foramen superciliare.
 <i>g</i>, <i>b</i>, The orbital plate.
 <i>i</i>, Part of the lacrymal fossa.
 <i>k</i>, <i>l</i>, That part of the transverse suture which unites the os frontis to the os planum and os unguis.
 <i>l</i>, <i>l</i>, The foramina orbitaria interna, anterius et posterius.
 <i>m</i>, Part of the coronal suture.
 B, The os parietale.
 <i>n</i>, The arched impression upon the surface of that bone.
 <i>o</i>, The squamous suture.
 <i>p</i>, Part of the lambdoid suture.
 C, A small portion of the os occipitis.
 D, The pars squamosa of the temporal bone.
 <i>q</i>, Part of the squamous suture between the temporal and sphenoid bones.
 <i>r</i>, The middle of the temporal fossa.
 <i>s</i>, The zygomatic process of the temporal bone.
 <i>t</i>, The zygomatic suture.
 <i>u</i>, The mastoid process of the temporal bone.
 <i>v</i>, The meatus auditorius externus.
 <i>w</i>, Part of the base of the pars petrosa.
 E, The pars plana of the ethmoid bone.
 F, The temporal plate of the sphenoid bone.
 <i>x</i>, The foramen nasale.
 H, The os unguis.
 <i>y</i>, The lacrymal fossa of the os unguis.
 I, The os malæ.
 <i>x</i>, The superior orbital process of that bone.</p> | <p>1. The zygomatic, and,
 2. 2. Maxillary processes of the os malæ.
 3. The orbital plate of that bone.
 4. The cavity for lodging part of the temporal muscle.
 5. The orbital passage of the os malæ.
 K, The maxillary fossa.
 6. 6. The ragged edges of the os nasi and os maxillare superius, to which the cartilages of the nose are fixed.
 7. The angular process of the os maxillare superius.
 8. The lacrymal fossa of that bone.
 9. A depression of the os maxillare at the entrance of the orbit.
 10. A portion of the orbit which belongs to the os maxillare.
 11. The notch at the opening of the nose.
 12. The foramen infra-orbitarium.
 13. 13. Several small holes in the os maxillare, for the passage of blood-vessels and nerves.
 14. The spine or ridge, formed by the union of the ossa maxillaria.
 15. 15. The malar processes to which the cheek-bone is fixed.
 16. The large tuber, or bulge of the os maxillare.
 17. 17. Elevations of the alveoli over the roots of the teeth.
 18. 18. The edge of the alveoli.
 L, The lower jaw.
 19. 19. Its base ;
 20. Its angles.
 21. 21. Muscular prints.
 22. The condyloid process.
 23. The coronoid process.
 24. The notch between these processes.
 25. 25. The sharp edge of the coronoid process.
 26. The mental hole.
 27. 27. The edge of the alveoli.</p> |
|--|--|




Altera fac.

TAB. 4.





TAB. 5.



T A B L E V.

A VIEW of the INNER SURFACE of the BASE of the CRANIUM.

-
- a, a*, THE upper edge of the zygoma. Between the anterior *a*, and the cut edge of the skull, a portion of the external temporal fossa.
b, The mastoid process of the temporal bone.
c, c, The external surface of the occipital bone.
A, The left frontal fossa, marked with ridges and depressions.
d, Part of the frontal spine.
B, The foramen placed at the bottom of the frontal spine.
C, The cribriform plate of the ethmoid bone.
e, The crista galli.
D, The fella turcica.
E, The left anterior clinoid process.
F, The posterior clinoid process.
f, A small process of the sphenoid bone, projecting into the back-part of the ethmoid bone.
g, Part of the sphenoid suture.
h, The processus semi-olivaris.
G, The left temporal fossa of the sphenoid bone.
H, The left transverse spinous process.
I, The foramen opticum.
K, i, A portion of the foramen lacerum.
L, The foramen rotundum.
M, The foramen ovale.
N, The foramen spinale.
k, An impression made by the internal carotid artery.
l, The point of the pars petrosa of the temporal bone; under which is the passage of the internal carotid artery. At the fore-part of the process, in the shaded place, there is an irregular opening, which, in the subject, is filled partly with bone, and partly with a cartilaginous ligament.
O, The future common to the sphenoid and temporal bones.
P, The squamous part of the temporal bone, which completes the temporal fossa for the lateral lobe of the brain.
Q, The ridge which divides the pars petrosa of the temporal bone into anterior and posterior surfaces.
m, The posterior surface of the pars petrosa.
n, The foramen innominatum.
o, A groove which lodges the superior petrosal sinus.
R, The meatus auditorius internus.
S, The anterior part of the foramen lacerum common to the temporal and occipital bones.
T, The posterior part of the same foramen.
U, The fossa for lodging a portion of the lateral sinus.
V, The cuneiform process of the occipital bone.
W, The anterior condyloid foramen of that bone.
X, The foramen magnum.
Y, The inferior occipital fossa, which lodges the corresponding lobe of the cerebellum.
Z, Z, p, A fossa of the occipital bone for the left lateral sinus.
z, z, The inferior limb of the cruciform spine, running backwards from the foramen magnum.
p, Part of the lambdoid suture.
q, A continuation of the fossa for the lateral sinus.
r, A fossa for the inferior petrosal sinus.
s, s, The cut edge of the skull.

T A B L E VI.

THE Outer and Under Surface of the SKULL, turned a little to the Left Side.

-
- A, THE inferior posterior angle of the parietal bone.
 B, B, The lambdoid future.
 C, C, The large transverse arch, ridge, or spine of the occipital bone; the upper and outer part of which gives rise to the occipito-frontalis, and the middle to the trapezii muscles.
 C, *m*, C, *m*, Depressions made by the insertions of the complexi muscles on that bone.
n, The spinous tuberosity, observed only in some skulls.
i, *i*, The perpendicular spine. Between *m*, *m*, and the back-parts of the temporal bones, are impressions made by the splenii muscles.
l, *l*, The smaller transverse ridge or spine.
k, *k*, The cavities where the recti minores muscles are inserted. On the outer side of these cavities, the obliqui superiores and recti majores make impressions.
 D, The foramen magnum.
 E, The cuneiform process.
 F, F, The condyloid processes.
b, *b*, The tuberosities at the roots of the condyles, which give attachment to the capsular ligament of the first vertebra.—The *b* placed at the root of the left condyle, points out the superior condyloid hole.
 G, G, The posterior condyloid holes.
 H, The squamous portion of the temporal bone.
 I, I, The squamous future.
 K, K, The mastoid processes.
t, *t*, The mastoid fissures.
m, The foramen mastoideum.
 L, The root of the zygoma;
p, Its articular process.
 M, The styloid process; behind the root of which the foramen stylo-mastoideum is concealed.
 N, The meatus auditorius externus.
 O, The glenoid cavity, for the articulation of the lower jaw.
q, The glenoid fissure.
 P, The foramen caroticum.
 Q, The thimble-like cavity, or jugular fossa.
 R, R, The pterygoid fossæ, at the outer sides of which are the external pterygoid plates.
 V, The internal pterygoid plate.
 W, The hook-like process, round which the circumflex muscle of the palate moves.
 S, The temporal process of the sphenoid bone.
 T, The spinous process, and spinous hole of that bone.
 7. The osseous mouth of the Eustachian tube.
 Y, The foramen ovale.
s, *s*, Passages common to the temporal and sphenoid bones.
 X, The foramen pterygoideum.
 Z, The inferior orbital fissure.
a, The under part of the tuberosity of the superior maxillary bone.
b, The palate process of that bone.
c, The foramen incisivum.
d, The internal surface of the os malæ, which contains a portion of the temporal muscle.
e, The under edge of the zygomatic process.
o, The zygomatic future.
f, The palate process of the palate bone.
 5. 6. The superior and inferior spongy bones.
g, The posterior edge of the vomer.
 4 The foramen gustativum.
 1. 1. 1. 1. The dentes incisores.
 2. 2. The dentes canini.
 3. &c. The dentes molares.

TAB. 6.



Fig. 1.




Fig. 2.



Fig. 3.



T A B L E VII.

A Back-View of the SKULL, with the Additional Bones called OSSA TRIQUETRA.

FIG. 1.

The SKULL seen from its Posterior and Left Side.

- A, The frontal bone.
 B, Part of the temporal fossa of that bone.
 C, C, The parietal bones.
 D, The coronal future.
 E, The sagittal future.
 F, F, The lambdoid future.
 G, G, Ossa triquetra, between the lambdoid and sagittal futures.
 H, H, The foramina parietalia.
 I, I, The arched impression of the left parietal bone.
 K, The occipital bone.
 L, Part of the large transverse arched ridge of that bone.
 M, The squamous future.
 N, The squamous part of the temporal bone.
 O, The mastoid process.
 P, The zygoma.
 Q, Part of the meatus auditorius externus.
 R, Part of the temporal fossa of the sphenoid bone.
 S, The temporal fossa of the temporal bone.
 T, The outer surface of the orbital process of the cheek-bone.

U, The zygomatic future.

V, The superior orbital process of the cheek-bone.

W, Part of the superior maxillary bone.

X, Part of the outer plate of the pterygoid process.

Y, Y, Some of the teeth.

FIG. 2.

A Portion of the Upper and Back Part of the CRANIUM, with Ossa Triquetra.

- A, A, Part of the parietal bones.
 B, A portion of the sagittal future.
 C, C, The parietal foramina, uncommonly large.
 D, The upper part of the occipital bone.
 E, Part of the lambdoid future.
 The ossa triquetra are seen between the occipital and parietal bones, varying considerably in figure and size.

FIG. 3.

An additional piece of Bone in the Side of the Cranium, inclosed by a double Squamous Suture, and forming a kind of Os Triquetrum.

T A B L E VIII.

VIEWS of the Upper Part of the CRANIUM and of the FRONTAL BONE.

FIG. 1.

Represents the Upper and Outer Surface of the CRANIUM.

- a*, The frontal bone.
- b, b*, The coronal future.
- c, c*, The parietal bone.
- d, d*, The sagittal future.
- e, e*, The parietal holes.
- f*, Part of the occipital bone.
- g, g*, The middle of the lambdoid future.

FIG. 2.

A View of the Upper and Inner Surface of the CRANIUM.

- b*, The outer and fore part of the frontal bone.
- i, i*, Prints made by the blood-vessels of the dura mater.
- k, k*, The finosity where the upper part of the falx is fixed, and the superior longitudinal finus is lodged.
- l, l*, Pits frequently found; the larger occasioned by luxuriances of the brain, and the smaller by the glands of Pacchioni, or by the meeting of blood-vessels of the dura mater.

FIG. 3.

The Outer Surface of the FRONTAL BONE.

- a*, The middle and convex part of the frontal bone.
- b, b*, The elevations of this bone.
- c*, The muscular print of the left side.
- d*, Part of the temporal fossa.
- e, e*, The external and internal angular processes.
- f*, The nasal process.

- g, g*, Eminences and cavities, to which the nasal and maxillary bones are fixed.
- b, b*, The superciliary arches.
- i, i*, The superciliary holes.
- k, k*, The orbital plates.
- l, l*, The lacrymal fossæ.
- m, m*, The foramina orbitaria interna.
- n, n*, The inequalities which unite the frontal to the sphenoid bone.

FIG. 4.

The Inner Surface of the FRONTAL BONE.

- a*, The internal concave part of the frontal bone.
- b*, The cavity which lodges the anterior lobes of the brain.
- c*, The frontal spine.
- d*, A furrow where the falx is fixed, and the superior longitudinal finus is lodged.
- e*, The ragged edge of the bone, which assists in forming the coronal future.
- f, f*, Other inequalities, which join the frontal to the sphenoid bone.
- g, g, g, g*, The inner surface of the four angular processes.
- h*, The posterior surface of the nasal process.
- i, i*, Other inequalities near the nasal process.
- k, k*, The orbital plates.
- l, l*, The lacrymal fossæ.
- m, m*, Cells which correspond with those of the ethmoid bone.
- n, n*, Passages from the frontal sinuses.
- o*, The foramen cæcum.
- p*, The opening which receives the cribriform plate of the ethmoid bone.
- q, q*, The furrows which lodge the blood-vessels of the dura mater.

Fig. 1.



TAB. 8.

Fig. 2.





Fig. 3.



Fig. 4.



OSSA PARIETALIA.

The *situation* of the Ossa Parietalia, in the upper and lateral parts of the Cranium. Tab. IV. B.

The *figure* of each Parietal Bone a Trapezium, or approaching that of a Square. Tab. IX. Fig. 1. 2.

The *upper* Edge, longest. Tab. IX.

The *anterior* Edge, next in length. Tab. IX.

The *posterior* Edge, shorter. Tab. IX.

The *inferior*, shortest, and in form of a ragged arch, to be connected to the upper rounded edge of the Squamous part of the Temporal Bone. Tab. IX.

The *three first Edges* of the Bone *ragged*, where they assist in forming the True Sutures. Tab. IX.

The *corners* of the Bone *obtuse*, excepting the under anterior, which forms a kind of process. Tab. IX.

The external surface of the Bone, *smooth* and *convex*. Tab. IX. Fig. 1.

The Transverse arched *Ridge*, or *Line*, frequently of a whiter colour than the rest of the Bone, placed externally, a little below its middle height, for the origin of the Temporal Muscle. Tab. IV. n.

The *radiated Furrows* at the under part of the Bone, formed by the Fibres of the Temporal Muscle. Tab. IV. o.

Near the semicircular edge, many inequalities, which join similar inequalities on the inside of the Temporal Bone, to form the Squamous Suture.

The *Foramen Parietale*, near the upper and back part of the Bone, for the transmission of a Vein from the Integuments of the Head to the superior longitudinal Sinus; and sometimes of a small Branch from the Temporal or the Occipital Artery, to the Falx of the Dura Mater. Tab. VIII. Fig. 1. 2.

In several skulls, one of the Parietal Holes is wanting; in others, two are found in one side; in others, none in either.

The internal *concave* Surface of the Bone.

The *Furrows* made by the Blood-Vessels of the Dura Mater; the principal of which begin by a Trunk at the under and fore part of the Bone, where frequently a full canal is formed. Tab. IX. Fig. 2.

In their progress upwards, the Furrows divide into many branches, and frequently small passages are seen running from these into the diploe.

The *depression* at the upper Edge of the Bone, for the attachment of the upper part of the Falx, and lodgement of the superior longitudinal Sinus. Tab. VIII. Fig. 2. This is most distinctly seen when the Bones are conjoined.

The Depression for the longitudinal Sinus, like the Sinus itself, becomes large in its course backwards; and frequently it is larger in one Bone than the other.

The *Fossa* at the under and back part of the Bone, for lodging a small part of the lateral Sinus. Tab. IX. Fig. 2.

Numerous *depressions* found on the inside of the Bone, occasioned by the prominences of the Brain.

The *connection* of the Parietal Bones to the Os Frontis by the Coronal, and to each other by the Sagittal Suture. Tab. VIII. Fig. 1. b, b, d, d.

The Parietal Bones have the two Tables and Diploe the completest, and are the most equal and smooth of any of the Cranium.

In the Fœtus, the sides of the Parietal Bones are incomplete, and there is no Parietal Hole. Between the Parietal Bones, and the middle of the divided Os Frontis, there is a large interstice, termed in common language, *opening of the Head*, and by Anatomists, *Bregma*, *Fons*, or *Fontanella*, from its having been supposed by the Ancients that through it the superfluous humours of the Brain are evacuated. Tab. XI. B.

The Bregma is occupied by a strong ligamentous Membrane, which adheres firmly to the ragged Edges of the Bones, and is lined within by the Dura Mater, and covered externally by the Pericranium.

The whole of the Bregma is generally ossified by two, though sometimes not till near seven years of age, and it has sometimes, though very rarely, been found open in the Adult.

OS OCCIPITIS.

THE *situation* of the Occipital Bone in the back and under part of the Cranium. Tab. VI.

Its *rhomboid* figure, with the angle above, generally a little rounded. Tab. IX. Fig. 4.

The two lateral *Angles* obtuse. Tab. IX. Fig. 4.

The external Surface *convex*, and *smooth* at the upper part. Tab. IX. Fig. 3.

The *large Arched Ridge*, running across the Bone, near the middle of the convex Surface, to the centre of which the Trapezii Muscles are fixed, the outer parts giving origin to the Occipito-frontalis Muscle. Tab. IX. Fig. 3. *d, e, e*.

The *smaller Arch*, half-way between the former, and the passage termed *Foramen Magnum*. Tab. IX. Fig. 3. *g, h, h*.

The *depressions* between the middle of the large and small Arches, for the connection of the Complexi Muscles.

The *impressions* between the Arches and the Temporal Bones, for the attachment of the Splenii. Tab. IX. Fig. 3.

Cavities between the smaller Arch and the Foramen Magnum, for the reception of the Recti Minores. Tab. IX. Fig. 3.

The *perpendicular Spine*, running through the middle of the two Arches, and separating the Muscles of the opposite Sides. Tab. IX. Fig. 3.

The *unequal Edges* of the Foramen Magnum, for the insertion of Ligaments, by which the Head is fixed to the Vertebrae of the Neck.

The *inferior Angle*, contrary to the rest of the Bone, flattened and stretched forwards in form of a wedge, hence called *Cuneiform Process*, or, from its situation, *Basilar Process*. Tab. IX. Fig. 3. *p*.

The *unequal Surface* of the Cuneiform Process, for the attachment of the Recti Anteriores Muscles.

The *Condyles* placed at the Base of the Cuneiform Process, and at the fore and lateral parts of the Foramen Magnum, for the Articulation with the Atlas, or first Vertebra of the Neck. Tab. IX. Fig. 3. *l, l*.

The *oval form* and smooth Cartilaginous Surface of the Condyles, corresponding with the superior articulating Processes of the Atlas.

The Condyles run obliquely forwards and inwards, and are deepest at their inner parts; in consequence of which they are prevented from sliding to either side out of the cavities of the Atlas.

In some Subjects, each of the Condyles is more or less divided, giving the appearance of two Prominences.

Round their roots, the surface is unequal, for the attachment of their Capsular Ligaments.

The *rough Prominences* between the Condyles and Mastoid Processes of the Temporal Bones, for the insertion of the Recti Capitis Laterales Muscles; and anterior to these, the semilunar Notches which form part of the Holes common to the Temporal and Occipital Bones. Tab. VI. between F and K.

The Flexion and Extension of the Head are performed at the Condyles, but they are found to be placed behind its centre of gravity, which affords space for the Mouth, Throat, &c. and the Head is prevented from falling forwards by the constant action of the strong Extensor Muscles, placed on the back-part of the Neck.

The *internal Surface* of the Bone, *hollow*, for containing the back-part of the Brain. Tab. IX. Fig. 4.

The *Cruciform Spine* of the inner side, formed by two Ridges, the one placed perpendicularly in the middle of the Bone, the other crossing the first in a horizontal direction. Tab. IX. Fig. 4. *a, f, b, b*.

The *upper Limb* of the perpendicular Spine, *hollow* in the middle, or frequently at one side, for the reception of the superior longitudinal Sinus, and the attachment of the Falx. Tab. IX. Fig. 4. *a—b*.

The *lateral Limbs*, placed opposite to the great external arched Spine, and *hollow* in the middle, for containing the lateral Sinuses, and giving attachment to the Tentorium Duræ Matris. Tab. IX. Fig. 4. *b, b*.

The hollow in one of the lateral Limbs is frequently the continuation of that made in the perpendicular Spine by the longitudinal Sinus, and therefore is often larger than the other.

The *lower Limb*, short, for the attachment of the Falx Minor, and sometimes hollow for the reception of an Occipital Sinus.

The *Fossæ* at the sides of the upper Limb, for containing the posterior Lobes of the Cerebrum. Tab. IX. Fig. 4. *e, e*.

The *Fossæ* at the sides of the lower Limb, for containing the Cerebellum. Tab. IX. Fig. 4. *e, e*.

Anterior to the Fossæ for lodging the Cerebellum, are two Cavities for receiving the lateral Sinuses, previous to their leaving the Cavity of the Cranium.

The *concave Surface* of the Cuneiform Process, for receiving the Medulla Oblongata, and Basilar Artery. Tab. IX. Fig. 4. *n*.

The *depressions* at each side of the Cuneiform Process, where the inferior Petrosal Sinuses are placed. Tab. V. *r, r*.

The *Foramen Magnum*, behind the Basilar Process, and between the Condyles, for the passage of the Medulla Oblongata, Vertebral Blood-Vessels and Accessory Nerves. Tab. V. X.

The

The *superior* or *anterior Condylloid Foramina*, at the sides of the Foramen Magnum, and immediately over the Condyles, for the passage of the Ninth Pair of Nerves. Tab. IX. Fig. 3. *n, n*.

The *posterior Condylloid Foramina*, at the back-part of the root of the Condyles, for the passage of Veins from the Occiput, or from the Vertebral Veins, into the lateral Sinuses, near their terminations. Tab. IX. *m, m*.

Frequently one of the posterior Condylloid Foramina is wanting; sometimes both, when the Veins pass through the Foramen Magnum.

Besides the holes above taken notice of, others are often found, near the edges of the Bone, for the transmission of Veins, the number and size of which are uncertain.

The *connection* of the Bone to the *Ossa Parietalia*, by the Lambdoid Suture. Tab. VII.

This Bone is among the thickest of the Cranium, though very unequal; being thick and strong above, where it is little affected by Muscles, and so thin below, where it is pressed by the weight of the Cerebellum internally, and by the action of the Muscles externally, as to be in many skulls rendered transparent. But the thick Muscles and strong Spine of the Bone assist greatly in preventing injuries from happening here.

In the Fœtus, the Occipital Bone is divided into *four pieces*; the first, which is larger than the other three, forms all the part of the Bone above the Foramen Magnum; the second and third are placed at the sides of that Foramen, and constitute almost the whole of the Condyles; and the fourth forms the Cuneiform Process.

T A B L E IX.

VIEWS of the PARIETAL and OCCIPITAL BONES.

FIG. 1.

The External Surface of the Right PARIETAL BONE.

- a*, The middle convex part of the bone.
b, b, The upper ragged edge of the bone, which, when joined to its fellow, forms the sagittal future.
c, c, The anterior edge, which assists in forming the coronal future.
d, The posterior edge, which joins the occipital bone, and assists in forming the lambdoid future.
e, e, The inferior semilunar edge, which is joined to the pars squamosa of the temporal bone.
f, The parietal hole.
g, g, The arched ridge of the parietal bone, which gives origin to a large share of the temporal muscle.
h, The anterior and superior angle of the bone.
i, The anterior-inferior angle.
k, k, The posterior angles.

FIG. 2.

The Internal Surface of the same PARIETAL BONE.

- a*, The middle internal concave part.
b, b, The inner surface of the upper edge of the bone, where the indentations are more apparent than those of the outer side.
c, The inner orifice of the parietal hole.
d, d, The anterior serrated edge of the bone.
e, e, The posterior edge, more strongly marked with indentations than its anterior edge.
f, f, The superior angles.
g, The inferior anterior angle, where the beginning of the furrow is seen, which lodges the trunk of the principal artery of the dura mater.
h, h, h, h, The ramifications of that furrow.
i, i, Small furrows of the bone, which correspond to its posterior inferior angle.
k, A depression which lodges part of the lateral sinus.
l, The inferior edge of the bone, considerably thinner than the rest.

FIG. 3.

A View of the External Surface of the OCCIPITAL BONE.

- a*, The superior angle of the bone.
b, b, The ragged edge of the bone which assists in forming the lambdoid future.
c, c, The irregularities at the lateral and inferior parts of the bone, where it is joined to the ossa temporum.
d, e, e, The large transverse arched ridge, or spine.
e, e, Muscular prints upon the transverse ridge.

- f*, The perpendicular spine.
g, The smaller arched ridge, running across the spine of the bone.
h, h, Muscular prints, above the foramen magnum.
i, i, The edge of,
k, The foramen magnum.
l, l, The occipital condyles.
m, m, The posterior condyloid foramina.
n, n, The inner side of the left, and outer side of the right anterior condyloid foramina.
o, o, Nitches which assist in forming the holes common to the occipital and temporal bones.
p, The extremity of the cuneiform process, upon which are seen,
q, q, Prints which give origin to some of the flexor muscles of the head.

FIG. 4.

The Internal Surface of the OCCIPITAL BONE.

- a*, The superior angle of the bone.
b, b, The middle, or lateral angles.
c, c, &c. The eminences and cavities which assist in forming the lambdoid future.
d, d, The superior occipital fossæ, which lodge a share of the posterior lobes of the cerebrum.
e, e, The inferior occipital fossæ, which lodge a part of the cerebellum.
f, f, The extremities of the crucial ridge or spine. The undermost, *f*, points out the attachment of the falx minor.
g, The fossa which lodges the superior longitudinal sinus, and has the falx fixed to its edges.
h, The middle of the crucial ridge.
The fossæ which lodge the lateral sinuses, and have the tentorium fixed to their edges, are included between the letters *b, b, b*.
i, i, The openings of the os occipitis, which form part of the foramina lacera, common to the occipital and temporal bones.
k, k, The small processes of the os occipitis, which assist in forming part of the foramina lacera.
l, l, The internal orifice of the posterior condyloid holes.
m, The anterior condyloid hole of the os occipitis of the right side.
n, The concave surface of the cuneiform process.
o, The inequalities of the cuneiform process of the os occipitis, by which it is united with the sphenoid bone.
p, The foramen magnum.

TAB. 9.

Fig. 1.




Fig. 2.



Fig. 3.



Fig. 4.



OSSA TEMPORUM.

THE *situation* of each Temporal Bone in the under part of the Cranium. Tab. IV. D.

The *Squamous Plate*, which forms a part of the Temple, and gives origin to a portion of the Temporal Muscle. Tab. IV. D.

The Squamous Plate appearing equal and smooth externally, with a thin semicircular edge above, which, by overlapping the under edge of the Parietal Bone, gives name to this Process. Tab. VII. Fig. 1. N, N.

The *Mastoid* or *Mammillary Process*, giving insertion to strong Muscles, particularly the Sterno-mastoid; and containing cells which communicate with each other, and with the Cavity of the Ear, called Tympanum. Tab. IV. u.

The *Petrous Process*, remarkably hard, very unequal, and of an oblong form, but becoming smaller in its progress; placed at the base of the Bone, from which it runs obliquely forwards and inwards, and contains the internal Organ of hearing; to be afterwards described. Tab. X. Fig. 6. f, q.

The *Zygomatic Process*, running from the under and fore part of the Squamous Plate, to join the Os Malæ; forming an Arch, on the inner side of which the Temporal Muscle passes to the Lower Jaw. Tab. X. Fig. 5. e. Tab. 4.

A *Tubercle* of an oblong form at the root of this Process, covered with a smooth Cartilage, making part of the Articulation of the Lower Jaw. Tab. IV. behind s.

The *Styloid Process*, placed at the root of the Pars Petrosa, and going obliquely downwards and forwards, to give origin to Muscles which borrow part of their name from it, and belong to the Tongue and Throat. Tab. X. Fig. 5. r. Tab. VI. M.

It is generally about an inch in length, though sometimes a great deal more, and is remarkably slender. It is frequently, even in Adults, not entirely ossified, and is therefore apt to drop off in macerating the Bones.

The *Vaginal Process*, of an inconsiderable size, surrounding the root of the Styloid Process, and deepest at its fore-part. Tab. X. Fig. 5. n.

The *Rough Semicircular Ridge*, at the under part of the external Meatus; sometimes also considered as a Process, and called *Auditory*, for the connection of the Cartilage of the Ear. Tab. X. Fig. 5. n.

A *Groove*, at the inner part of the root of the Mastoid Process, giving origin to the Digastric Muscle; and a little anterior to this, another groove, in which the Occipital Artery runs. Tab. VI. t.

The *Glenoid Cavity*, of great size, surrounded by the roots of the Zygoma, the Auditory and Vaginal Processes, and lined with Cartilage, for the articulation of the Lower Jaw. Tab. IV. O.

The *Glenoid Fissure*, near the back-part of this Cavity, running from the upper to the under side of it, for the attachment of part of the Capsular Ligament, of the articulation of the Lower Jaw. Tab. VI. q. Tab. X. Fig. 5. o.

A *Depression* between the Glenoid Fissure and Styloid Process, for lodging a portion of the Parotid Gland. Tab. VI. between G and the Styloid Process M.

The *Thimble-like Cavity*, or the *Jugular Fossa*, at the inner side of the root of the Styloid Process, for lodging the top of the internal Jugular Vein. Tab. VI. Q.

The *Meatus Auditorius Externus*,—a large Canal, between the Mastoid and Zygomatic Processes, leading inwards and forwards to the Organ of Hearing. Tab. IV. v.

Around the External Meatus, a rough Surface, for the Connection of the Cartilages and Ligaments of the Ear.

The *Foramen Stylo-mastoideum*, or *Aqueduct of Fallopius*, between the Styloid and Mastoid Processes, for the transmission of the Portio Dura of the Seventh Pair of Nerves.

The *Foramen Caroticum*, or *Canalis Caroticus*, at the inner and fore-part of the Jugular Fossa, and also before and at the inside of the Styloid Process, leading upwards, then forwards, through the point of the Pars Petrosa, for the transmission of the Internal Carotid Artery to, and of the great Sympathetic Nerve, from the Brain. Tab. VI. P.

In the upper and back part of the Canal, one, sometimes two, minute Holes are observed, through which the internal Carotid Artery sends one or two Twigs to the Tympanum.

The *Iter a Palato ad Aures*, or *Eustachian Tube*, between the Fissure for the Capsular Ligament of the Lower Jaw, and the Passage of the internal Carotid Artery, extending outwards and backwards in an horizontal direction, till it terminates in the Tympanum.

In the Subject, it is formed, by the addition of Cartilage and Ligament, into a trumpet-like Tube, which is continued forwards and inwards to the back-part of the Nostril, and conveys air from the Nose to the Tympanum of the Ear. Tab. with Views of Ear, Vol. II.

The *Foramen Mastoideum*, occasionally found at the back-part of the Mastoid Process, or in the Lambdoid Suture. When present, it sometimes transmits an Artery to the Dura Mater, but more commonly a Vein from the Integuments of the Head to the lateral Sinus. Tab. VI. u.

Sometimes two or three Foramina Mastoidea are observed, serving the same purpose with that already noticed; but these, like all the other Passages for Veins leading into the Sinuses, are very uncertain.

The *upper* and *inner Edge* of the Squamous Plate formed into *ridges* and *furrows*, where it is connected with the Parietal Bone. Tab. X. 6. *a, a*.

The *inner Surface* of the Squamous Plate, *unequal* where it is marked by the *Convolutions* of the middle part of the Brain, and by the *Arteries* of the Dura Mater. Tab. X. Fig. 6. *b, b*.

The *Pars Petrosa*, of great size, running forwards and inwards, with a sharp angle above, and two flat sides; one facing obliquely forwards and outwards, and the other as much backwards and inwards.

The *anterior* and *outer Surface* of the Pars Petrosa opposed to the lateral Lobes of the Brain. Tab. V. *n, o*.

The *posterior* and *inner Surface* of the Pars Petrosa, opposed to the Cerebellum. Tab. V. *o, m*.

The *Ridge* between the two surfaces of the Pars Petrosa, for the attachment of the Tentorium Duræ Matris. Tab. V. *Q*.

A *Groove* frequently found upon the Ridge of the Pars Petrosa, for lodging the superior Petrosal Sinus. Tab. V. *o*.

A *Fossa*, at the root of the posterior Surface of the Pars Petrosa, and opposite to the Mastoid Process, for lodging the lateral Sinus, where it turns downwards to go out of the Cranium. In this Fossa the passage is observed corresponding with the Foramen Mastoideum. Tab. V. *m*.

The Lateral Sinuses are frequently of unequal size; of course this cavity is then larger in one Temporal Bone than in the other.

The *Meatus Auditorius Internus*, or *Foramen Auditivum*, passing outwards and backwards, in the posterior Surface of the Pars Petrosa, for the passage of the Seventh Pair of Nerves, and the principal Artery belonging to the Inner Ear. Tab. V. *R*.

In the bottom of this passage, there are many Foramina; one above, more conspicuous than the rest, is the beginning of the passage for the Portio Dura of the Seventh Pair of Nerves. The others are the passages of the Branches of the Portio Mollis of that Nerve.

Some way below the Meatus Internus, is the opening of the passage termed, by Cotunnus, *Aquæductus Cochleæ*, and near the same distance behind the Meatus; and on the same side of the Bone, is the mouth of the *Aquæductus Vestibuli*.

The *Foramen Innominatum*, near the middle of the Anterior Surface of the Pars Petrosa, and leading backwards for the passage of the Vidian Nerve, which is reflected from the Second Branch of the Fifth, to the Portio Dura of the Seventh Pair. Tab. V. *n*.

The Orifice of the Canalis Caroticus appearing at the under part of the point of the Pars Petrosa. Tab. V.

The *Foramen Lacerum Posterius*, or *Hole* common to the Pars Petrosa and Cuneiform Process of the Occipital Bone, for the passage of the Lateral Sinus, the Eighth Pair, and Accessory Nerves. Tab. V. *S, T*.—The Nerves pass through the fore-part of the Hole, and are separated from the Sinus by a Process of the Dura Mater, stretched between two small Processes of these Bones. In some Skulls, an Osseous Partition separates the Nerves from the Sinus.

The *Connection* of the Bone, at its upper curved Edge, to the Parietal Bone by the Squamous Suture. Tab. IV. *o*.

To the under and back part of the Parietal Bone, by the Additamentum Suture Squamosæ. Tab. IV.

To the Occipital Bone, by the Additamentum Suture Lambdoidalis. Tab. IV.

The Squamous part of the Temporal Bone is thin, but equal, while the Pars Petrosa is thick and strong, but irregular, having within it several Cavities, Processes, and Bones, which belong to the Organ of Hearing.

In a Fœtus, the Squamous is separated from the Petrous part by a Fissure; there is no appearance of Mastoid or Styloid Process, and instead of an Osseous Meatus Externus, there is only a Ring of Bone, in which the Membrana Tympani is fixed. Tab. XI.

OS ETHMOIDES.

THE *Situation* of the Ethmoid, or Cribriform Bone, in the fore-part of the Base of the Cranium.

Its *Cuboid Figure*.

The *Cribriform Plate*, from which the Bone has its name, placed horizontally, and perforated, excepting at its back-part, with many holes, for the transmission of the Branches of the First or Olfactory Pair of Nerves. Tab. X. Fig. 4. *c*.

In the recent Subject, these Holes are so much filled up by the Processes of the Dura Mater which inclose the Nerves, that they are much less evident than in Bones where the Membranes have been removed.

The *Crista Calli*, arising perpendicularly from the middle of the Cribriform Plate, and highest at the upper and fore part. Tab. X. Fig. 4. *b*.

To

To the Edge of this Process, and to the unimperforated part of the Cribriform Plate, the Falx of the Dura Mater is fixed.

A *Notch* at the fore-part of the root of the Crista Galli, contributing, in a small degree, to the formation of the Foramen Cæcum of the Frontal Bone.

The *Nasal Plate*, extending downwards and forwards from the Base of the Crista Galli, to form the upper and back-part of the Septum Narium, or Partition of the Nostrils. Tab. XIII. Fig. 2. F.

The greater Part of this Process is very thin, but towards its anterior and under edge it becomes thicker, for its firmer junction with the Bones and middle Cartilage of the Nose.

It is frequently bent a little to one side; in such cases, the two Nostrils become unequal in size.

The *Ethmoid Cells*, of an indeterminate number and form, placed under the Cribriform Plate, a little to the outside of the Nasal Lamella, separated from each other by thin Laminae, and serving the same purposes as the Frontal Sinuses. Tab. X. Fig. 4. d. Tab. XIII. Fig. 1. G, H.

Their *Communications* with each other, with the Frontal Sinuses, and also with the Cavity of the Nose. Tab. XIII. Fig. 3. B.

The *Os Spongiosum Superius*, on each side, projecting inwards and downwards from the Ethmoid Cells at the side of the Nasal Lamella, for enlarging the Organ of Smell. Tab. X. Fig. 3. e.

The *Triangular Form*, and *Spongy Texture* of each. Tab. XIII. Fig. 1. k, l, m, n.

In the Quadruped, this Bone is convoluted like a *Tur-*

ban. Hence, in the Human species, it is frequently called *Turbinatum*.

Its *Convexity* towards the Septum, and *Concavity* outwards. Tab. X. Fig. 3. e.

The *Ossa Plana*, or *Orbital Plates*, for covering a large share of the Ethmoid Cells, and forming the greater part of the inner sides of the Orbits. Tab. X. Fig. 4. g. Tab. IV. E.

On the upper Edge of each Os Planum, two small Notches appear, which, with similar Notches in the Frontal Bone, form the internal Orbital Holes. Tab. IV. l, l.

The *Connection* of the Cribriform Plates to the Orbital Plates of the Frontal Bone by the Ethmoid Suture; and to the Sphenoid Bone, by a Suture common to the two Bones, but generally considered as belonging to the latter. Tab. V. C, f.

The *Connection* of the Ossa Plana to the Orbital Plates of the Frontal Bone, by part of the Transverse Suture. Tab. IV. k, l.

The *posterior Edge* of the Nasal Plate, joined to the Processus Azygos of the Sphenoid Bone. Tab. XIII. Fig. 2.

Its *upper Edge*, joined to the Nasal Processes of the Frontal and Nasal Bones. Tab. XIII. Fig. 2.

Its *anterior Edge*, joined to the middle Cartilage of the Nose. Tab. XIII. Fig. 2. I.

In the Fœtus, the Ethmoid Bone is divided into two by a Cartilaginous Partition, which afterwards forms the Nasal Plate and Crista Galli. The other parts of the Bone are completely ossified. Tab. XI.

OS SPHENOIDES.

THE *Situation* of the Sphenoid or Cuneiform Bone, in the middle of the Cranium. Tab. V. g, f, O, F.

Its *Irregular Figure*, compared to that of a Bat with extended wings. Tab. X. Fig. 2.

The *Temporal Plates*, or *Wings*, placed at the sides of the Bone, and each hollow at the upper and outer part, for lodging a share of the Temporal Muscles. Tab. IV. F.

The *Orbital Plates*, at the fore-part of the Temporal Wings, forming a portion of the outside of the Orbits. Tab. III. m.

The *Spinous Process*, at the under and back part of each Temporal Plate. Tab. X. Fig. 2. s.

A *Styloid Process*, frequently observed at the point of the Spinous, from both of which the Circumflexus Palati arises. Tab. X. Fig. 2.

Between the Temporal and Spinous Processes, an *Arch* for receiving the fore-part of the Temporal Bone. Tab. IV. q. Tab. X. Fig. 2. q.

The two *Pterygoid*, or *Aliform Processes*, placed almost perpendicular to the base of the Cranium. The Pterygoid Processes are compared to the wings, though more properly resembling the feet of a Bat. Each is composed of two Plates.

The *External Plate*, broad and hollow without, where the external Pterygoid Muscle has its origin. Between the root of this Plate and that of the temporal one, a large Depression, where the principal part of the external Pterygoid Muscle has its origin. At the fore-part of this is another depression, forming part of the opening common to the Sphenoid, and to the Malar and superior Maxillary Bones.

The *internal Plate*, narrower and longer than the external, and, with its fellow, forming the back-part of the Nose.

A *Hook-like Process* upon the internal Plate, over which the *Circumflexus Palati* moves.

The *Fossa Pterygoidea*, facing backwards between the Pterygoid Plates, giving rise to the internal Pterygoid Muscle. Tab. VI. R.

A *small Depression* at the back-part of the root of the internal Pterygoid Plate, which gives origin to part of the *Circumflex Muscle* of the Palate.

A *Groove* on each side, which extends at the inner part of the Bone, between the root of the Styloid Process, and that of the internal Pterygoid Plate, assisting in the formation of the Eustachian Tube.

The two *Triangular Processes*, which adhere to the under part of the Sphenoid, and to the Ethmoid Bone, and which are considered as two of the Bones of the Face. Tab. X. Fig. 1. b. Fig. 3. k.

The *Processus Azygos*, standing single, and forming a sharp ridge which projects from under the middle and fore part of the Bone. Tab. X. Fig. 1. a.

The *Clinoid Processes*, seen on the inside of the Bone, compared to the supporters of a Bed, of which there are,

Two *Anterior*, projecting from the fore-part of the Body of the Bone, and extending horizontally outwards; each terminating in a point which obtains the name of *Transverse Spinous Process*. Tab. V. E. Tab. X. Fig. 2. c. and,

One *Posterior*, situated transversely, some way behind the anterior Processes, and frequently ending in two knobs, which incline obliquely forwards. Tab. X. Fig. 2. b.

Sometimes one or both of the anterior Clinoid Processes are united with the posterior, forming an Arch over the internal Carotid Artery.

The *Processus Olivaris*, considered by some as a fourth Clinoid Process, lying between, and a little behind, the roots of the anterior Clinoid Processes. Tab. V. b.

Between the anterior Clinoid Processes, a *small-pointed Process* frequently juts forwards, to join the Cribriform Plate of the Ethmoid Bone, from which it is sometimes called Ethmoid Process. Tab. V. f.

The *Temporal Fossæ* of this Bone, which lodge a share of the Lateral Lobes of the Brain. Tab. V. P.

A *Fossa* between the anterior Clinoid Processes, where part of the anterior Lobes of the Brain rests.

A *Depression* before the *Processus Olivaris*, where the conjoined Optic Nerves lie; and a *Fossa* on each side of it, where these Nerves are situated, previous to their entering the Orbits. Tab. X. D.

The *Fossa Pituitaria*, *Sella Turcica*, *Ephippium*, or *Turkish Saddle*, between the *Processus Olivaris* and posterior Clinoid Process, for lodging the *Glandula Pituitaria*.

A *Depression*, running first upwards then forwards, upon

each side of the posterior Clinoid Process and *Sella Turcica*, and terminating in a Pit at the root of the anterior Clinoid Process. These Depressions point out the course of the internal Carotid Arteries, when they have entered the Cranium, and previous to their perforating the *Dura Mater*, to be dispersed upon the Brain. Tab. V. k.

Besides these Impressions, several others may be observed made by Nerves and Vessels leading to or from their respective Holes in the Base of the Cranium.

The *Foramina Optica* at the roots of the anterior Clinoid Processes, for the transmission of the Optic Nerves and Ocular Arteries. Tab. X. Fig. 2. f.

The *Foramina Lacera Superiora*, or *Superior Orbital Fissures*, under the anterior Clinoid Processes and their transverse Spinous parts, for the passage of the Third, Fourth, first part of the Fifth, and the Sixth Pair of Nerves, with the Ocular Veins. Tab. X. Fig. 2. i.

The *Foramina Lacera* are largest at their inner ends. At their outer extremities they are considerably smaller, and are formed there by the *Os Frontis*; hence they may be ranked among the common Holes of the Cranium.

The *Foramina Rotunda*, a little behind the *Foramina Lacera*, for the passage of the second part of the Fifth Pair of Nerves, which are termed also *Superior Maxillary*. Tab. X. Fig. 2. n.

The *Foramina Ovalia*, considerably larger than the *Foramina Rotunda*, and placed farther back, and more externally, for the passage of the third part of the Fifth Pair of Nerves, and, commonly, for the passage of the Veins which accompany the principal Arteries of the *Dura Mater*. Tab. X. Fig. 2. o.

The *Foramina Spinalia*, a little to the outer and back part of the *Foramina Ovalia*, and in the Points of the Spinous Processes, for the transmission of the principal Arteries of the *Dura Mater*. Tab. X. Fig. 2. p.

The *Foramina Pterygoidea*, termed also, after the discoverer, *Foramina Vidiana*, at the roots of the inner Plates of the Pterygoid Processes, for the passage of two reflected branches of the second part of the Fifth Pair of Nerves. Tab. X. Fig. 1. n. Tab. VI. X.

The *Foramina Pterygoidea* are the smallest of the Sphenoid Holes, and cannot be distinctly seen in the entire Skull.

Sometimes one or more small passages are observed in or near the *Sella Turcica*, for the transmission of Blood-vessels into the Sphenoid Sinus, or to the substance of the Bone.

The *Foramina Lacera Anteriora*, common to the points of the *Partes Petrosæ* and the Sphenoid Bone. Tab. V. before l.

In a recent Skull, each of these Holes is filled behind with a thin Plate of Bone, which covers the internal Carotid Artery, and farther forwards, with a Cartilaginous Ligament, which lies over the Eustachian Tube, both of which drop out by Maceration.

The *Sphenoid Sinuses*, in the body of the Bone, at the under

under and fore part of the Sella Turcica, answering the same purpose with the Ethmoid and Frontal Cells. Tab. X. Fig. 1. c.

A *complete Partition* between the right and left Sphenoid Sinuses. Tab. X. between two c's.

A *Passage* from the upper and fore part of each of the Sphenoid Sinuses, descending, in a slanting direction, into the superior-posterior parts of the Nose. See Tab. with the Nerves of the Nose.

The two Sinuses are frequently of unequal size, and sometimes there is but one large Cavity, with an opening into one of the Nostrils. In some subjects, instead

of Sinuses, the body of the Bone is composed of large Cells.

The *Substance* of this Bone is the most unequal of any in the Body, some parts being extremely thin, while others are thicker than most parts of the Cranium.

The *Connection* of the Bone to all the other Bones of the Cranium, by the Sphenoid Suture, though others, as the Transverse and Ethmoid Sutures, are confounded with it.

In the Fœtus, the Temporal Wings are separated from the Body of the Bone by Maceration, and there are no Sphenoid Sinuses. Tab. XI.

T A B L E X.

Represents the SPHENOID, ETHMOID, and TEMPORAL BONES.

FIG. 1.

A View of the Outer and Under Surface of the SPHENOID BONE.

- a*, THE processus azygos of the sphenoid bone.
b, b, The small triangular bones, which, in old people, grow to the body of this bone.
c, c, The orifices of the sphenoid sinuses.
d, d, The foramina lacera.
e, The anterior and superior part of the body of the bone.
f, f, The external surface of the transverse processes.
g, g, The orbital plates.
h, h, The superior extremities of the temporal processes.
i, i, The middle of the temporal processes, which form part of the temporal fossæ.
k, k, The asperities by which this bone is joined to the ossa malarum.
l, l, Gutters in the os sphenoides, which lodge branches of the fifth pair of nerves.
m, m, The foramina rotunda.
n, n, The foramina pterygoidea.
o, o, The anterior openings, which assist in forming the sphenomaxillary fissures.
p, p, The foramina ovalia.
q, q, The spinous processes.
r, r, The roots of the pterygoid processes.
s, s, The internal plates of the pterygoid processes.
t, t, The hook-like processes, at the extremities of the internal plates.
v, v, Small sinuosities in the hook-like processes, over which the tendons of the circumflex muscles of the palate play.
x, x, The external plates of the pterygoid processes.
y, y, The parts of the bone adapted to the ossa palati.
z, z, The posterior openings, common to the sphenoid and temporal bones, over which the internal carotid arteries pass.

FIG. 2.

A View of the Inner and Upper Surface of the SPHENOID BONE.

- a, a*, The superior and anterior part of the sphenoid bone, which is joined to the under and back part of the frontal bone.

- b, b*, The temporal plates, or processes of the bone.
c, c, The transverse processes.
d, The small anterior process, which unites with the ethmoid bone.
e, A protuberance situated before the union of the optic nerves.
f, f, The foramina optica.
g, g, The anterior clinoid processes.
h, h, The posterior clinoid processes.
i, i, Part of the foramina lacera.
k, k, Impressions made by the internal carotid arteries.
l, The Sella Turcica.
m, m, The cavities of the temporal processes, which receive the lateral lobes of the brain.
n, n, The foramina rotunda.
o, o, The foramina ovalia.
p, p, The foramina spinalia.
q, q, The ragged edge of the bone, which assists in forming the sphenoid suture.
r, Part of the sphenoid bone, which joins the cuneiform process of the occipital bone.
s, s, Part of the spinous processes.
t, t, Part of the pterygoid processes.

FIG. 3.

The Outer and Under Surface of the ETHMOID BONE.

- a*, The anterior extremity of the nasal plate, which forms the upper part of the septum narium.
b, The posterior extremity of the nasal plate, which is very thin.
c, c, The ethmoid grooves, or chinks which separate the nasal plate from the ossa spongiosa superiora.
d, d, The passages for the branches of the olfactory nerves.
e, e, The ossa spongiosa superiora.
f, f, The cavities of the ossa spongiosa superiora.
g, g, Part of the ethmoid cells.
h, h, Inequalities of the ethmoid bone, by which it is joined to the frontal bone.
i, i, The posterior extremity, which is joined to the sphenoid bone.
k, k, The small cornua, or triangular bones, which, in adults, are joined to the body of the sphenoid bone.

FIG.

FIG. 4.

The Inner and Upper Surface of the ETHMOID BONE.

- a*, The anterior extremity of the bone, terminating in a small flat process.
b, The upper part of the Crista Galli.
c, c, The cribriform plate, with the different passages of the olfactory nerves.
d, d, Some of the cells of the ethmoid bone.
e, The posterior extremity of the nasal plate, which forms part of the septum narium.
f, f, The posterior margins of the ethmoid bone.
g, A great part of the os planum of the left side.
h, h, The upper part of the triangular bones which are joined to the body of the sphenoid bone.
i, The joining of the triangular bones with the ethmoid bone.
k, Holes formed by the union of the triangular bones with the ethmoid bone.

FIG. 5.

A View of the Outer Surface of the TEMPORAL BONE.

- a*, The upper and squamous part of the temporal bone.
b, The middle of the squamous part.
c, The under part, which lodges a portion of the temporal muscle.
d, That part of the temporal bone, which, when joined to the under and back part of the os parietale, forms the additamentum futuræ squamosæ.
e, The zygomatic process.
f, The base of the zygomatic process.
g, The transverse, or articular process.
h, h, The mastoid process.
i, i, Several small holes which transmit vessels to the substance of the bone, or to the dura mater.
k, k, Two holes at the root of the zygomatic process, for the transmission of vessels to the substance of the bone, or to the dura mater.
l, The meatus auditorius externus.
m, Inequalities at the beginning of the meatus.

- n*, The glenoid cavity for the articulation of the lower jaw.
o, The glenoid fissure, to which part of the articular ligament is fixed.
p, The vaginal process.
q, Part of the mastoid groove.
r, The styloid process.
s, The foramen mastoideum.
t, The base, or upper part of the mastoid process.
v, The inferior and anterior part of the temporal bone, which is joined to the os sphenoides.
x, A portion of the Eustachian tube.
y, A portion of the pars petrosa.

FIG. 6.

A View of the Inner Surface of the TEMPORAL BONE.

- a, a*, The upper edge of the squamous process.
b, b, Depressions which correspond with the circumvolutions of the brain.
c, Part of the temporal bone, which is joined to the os sphenoides.
d, The notch which receives the posterior and inferior angle of the parietal bone.
e, The upper part of the pars petrosa.
f, The groove which lodges the superior petrosal sinus.
g, The fossa which lodges a part of the lateral sinus.
h, The meatus auditorius internus.
i, The notch which assists in forming the foramen lacerum.
k, Part of the fossa of the temporal bone, which lodges the beginning of the internal jugular vein.
l, The posterior part of the bone, which is joined to the occipital one.
m, The inner surface of the foramen mastoideum.
n, A portion of the mastoid process.
o, A considerable part of the mastoid groove.
p, The styloid process.
q, The inner extremity of the pars petrosa, divided into two portions.

TABLE

T A B L E X I.

VIEWS OF THE FOETAL BONES OF THE HEAD.

FIG. 1.

A Lateral View of the SKULL.

- A, THE frontal bone, not yet complete in its middle and upper part.
 B, Part of the fontanella.
 C, The coronal future, which is here in part membranous.
 D, The parietal bone, formed of radiated fibres.
 E, A portion of the left parietal bone.
 F, The sagittal future.
 G, The occipital bone.
 H, The lambdoid future.
 I, The fontanella posterior, seen only in some bones.
 K, The squamous part of the temporal bone.
 L, The zygomatic process of that bone.
 M, The mastoid process.
 N, The squamous future, partly membranous.
 O, The membrana tympani.
 P, The temporal plate of the sphenoid bone.
 Q, The nasal process of the superior maxillary bone.
 R, The body of that bone.
 S, The orbit.
 T, The os malæ.
 U, The lower jaw.

FIG. 2.

The Outer Surface of the Two Pieces which form the FRONTAL BONE.

- A, The right, and,
 B, The left portion of the frontal bone.
 C, C, The orbital plates.

FIG. 3.

The Outer Surface of the Left PARIETAL BONE, in which the radiated Appearance of the Osseous Fibres is seen; the under and middle part appears prominent and more compact than the rest of the Bone.

FIG. 4.

The Outer Surface of the same Bone.

FIG. 5.

The Outer Surface of the four Pieces which form the OCCIPITAL BONE.

- A, The upper and largest portion.
 B, B, The two lateral portions, with the condyles and condyloid foramina.
 C, The portion which forms the cuneiform process.

FIG. 6.

The Outer Surface of the TEMPORAL BONE of the Right Side.

- A, The squamous plate.
 B, The zygomatic process.
 C, The articular cavity.
 D, The osseous circle, in which the membrana tympani is incased.

FIG. 7.

The Inner Surface of the same Bone.

- A, Inequalities which facilitate the union between the squamous and petrous portions.
 B, The fissure of the osseous circle, in which the membrana tympani is incased.

FIG. 8.

The Outer Surface of the Petrous Part of the TEMPORAL BONE.

- A, The cellular texture of the bone.
 B, The part which forms the future mastoid process.
 C, The bottom of the tympanum, with passages belonging to the internal organ of hearing.

FIG. 9.

The Lateral Portions of the ETHMOID BONE.

FIG. 10.

The three Pieces which form the SPHENOID BONE.

- A, A, The temporal wings, &c.
 B, The middle portion, forming the body of the bone, with the clinoid processes and sella Turcica.

Fig. 3.





Fig. 4.



TAB. II.



Fig. 5.




Fig. 8.



Fig. 10.



Fig. 9.




Fig. 2.




Fig. 6.





Fig. 7.





OF THE BONES OF THE FACE.

THEIR *Division* into *Upper and Under Jaws*.

The Upper Jaw, or Maxilla Superior, besides the Teeth, composed of seven Pairs of Bones, and one without a fellow, viz.

Two Ossa Nasi; Two Ossa Unguis; Two Ossa Malorum; Two Ossa Maxillaria Superiora; Two Ossa Palati; Two Ossa Spongiosa Inferiora; Two Ossa Triangularia; and the Vomer.

The Lower Jaw, or Maxilla Inferior, consists of a single Bone, with the Teeth.

The Bones of the Upper Jaw are joined together by Sutures which have no distinct Indentations, like those of the Cranium; but, like them, they are frequently found obliterated in the Skulls of old people. The Bones here, in consequence of the nature of the Sutures, have no motion but what they possess in common with the Cranium. The Sutures shall be taken notice of in the description of the Bones between which they are placed.

OSSA NASI.

Their *Situation* in the upper and fore part of the Nose. Tab. III. F.

The *Oblong Form* of each, though irregularly so. Tab. XII. Fig. 3.

The *thick, ragged*, upper end, where it forms a strong connection with the Frontal Bone. Tab. XII. Fig. 3.

Each narrowest a little below the upper end, and bent backwards.

The inferior extremity, thinner and broader than the rest of the Bone, and unequal where it gives attachment to the Cartilaginous part of the Nose. Tab. XII. Fig. 3.

The under half convex externally, by which, when the Bone is joined to its fellow, a strong Arch is formed, that is fitted for resisting injury. Tab. III. F.

Its internal *Concavity*, where it forms part of the Cavity of the Nose. Tab. XII. Fig. 4.

The *Spinous Process*, which joins the Nasal Lamella of the Ethmoid Bone, and thereby forms part of the partition of the Nose. Tab. XII. Fig. 4. C.

One or more *Holes* externally, for transmitting Vessels into the Substance of the Bone, or to the Membrane of the Nose. Tab. IV. x.

Its *Connection* to the Frontal Bone by the Transverse Suture. Tab. III. f, and,

To its fellow by the anterior Nasal Suture. Tab. III. before F.

In the Fœtus, the Ossa Nasi are proportionally short, but are otherwise complete.

Vol. I.

OSSA UNGUIS, or LACRYMALIA.

Their *Situation* at the inner and fore part of the Orbit. Tab. IV. H.

The *Division* of each externally, into two depressed Surfaces, and a middle Ridge. Tab. XII. Fig. 5. C.

The *posterior Depression*, the larger of the two, forming part of the Orbit. Tab. XII. Fig. 5. B.

The *anterior Depression*, lodging part of the Lacrymal Sac and Duct, and perforated by small Holes, through which Fibres pass, to make a firm connection between the Bone and its investing Membrane. Tab. XII. Fig. 5. A.

The *inner Surface*, composed of a Furrow and two irregular convex Surfaces, corresponding with the anterior Ethmoid Cells. Tab. XII. Fig. 6.

The *Substance* of the Bone, is the *thinnest* and *most brittle* of any in the Body.

Its *connection* to the Frontal Bone, by the Transverse Suture, and to the Os Planum by the Ethmoid Suture.

Internally, it is connected with the Ethmoid Cells.

In the Fœtus, it is fully formed.

OSSA MALARUM.

The *Situation* of each in the outer part of the Cheek. Tab. IV. I.

The *external convex, smooth Surface*. Tab. III. H.

The *posterior hollow Surface*, for lodging part of the Temporal Muscle. Tab. XII. Fig. 8. F.

The *superior Orbital Process*, forming part of the outside of the Orbit. Tab. IV. x.

The *inferior Orbital Process*, forming part of the lower Edge of the Orbit. Tab. IV. uppermost 2.

The *Maxillary Process*, forming the under part of the Prominence of the Cheek. Tab. III. undermost q.

The *Arch* between the Orbital Processes, which forms near a third part of the anterior circumference of the Orbit. Tab. IV. 3.

The *Zygomatic Process*, the most conspicuous, with that of the Temporal Bone, forming an Arch over the Temporal Muscle. Tab. IV. 1.

The *Internal Orbital Plate*, forming the outer and forepart of the Orbit. Tab. III. r.

A *Passage* through the Bone, for the transmission of small Vessels or Nerves from the Orbit to the Face. Tab. III. s.

The *Connection* of the superior Orbital Process and internal Orbital Plate to the Frontal and Sphenoid Bones, by the Transverse Suture. Tab. III. q, r.

E

The

The *Connection* of the Zygomatic Process to the Temporal Bone, by the Zygomatic Suture. Tab. IV. 1.

The substance of the Bone is thick and hard, with some Cancelli.

In the Fœtus, the Bone is fully ossified.

OSSA MAXILLARIA SUPERIORA.

The *Situation* in the fore-part of the Upper Jaw, and sides of the Nose. Tab. III. I.

Their *Size*, the largest of the Bones of the Upper Jaw, on which account the Bones have got their name. Tab. XII. Fig. 1.

The *Nasal*, or *Angular Process* of each, forming part of the side of the Nose, and of the inner part of the Orbit, and overlapping the outer edge of the Os Nasi above, while that Bone covers the Edge of the Nasal Process below. Tab. III. t.

A *Ridge* at the under and inner side of the Nasal Process, for supporting part of the Os Spongiosum Inferius. Tab. XII. Fig. 2. b.

The *Orbital Plate*, forming a large share of the under side of the Orbit. Tab. IV. 10.

The *Malar Process*, *unequal* and *ragged*, where it contributes, with the Os Malæ, to form the Prominence of the Cheek. Tab. XII. Fig. 1. e, e, f, f.

The *Tuberosity*, or *Bulge* at the back-part of the Bone, which forms the posterior boundary of the Cavity called *Antrum Maxillare*, and gives origin to a portion of the External Pterygoid Muscle. Tab. IV. 16.

The *Alveolar Arch*, of a spongy nature, where the Alveoli or Sockets of the Teeth are placed. Tab. III. x.

The *Palate Plate* or *Process*, placed horizontally, forming part of the Roof of the Mouth, and of the Bottom of the Nose. Tab. VI. b.

The *Palate Plate*, *thin* in its middle, and *thick* at its edges; *smooth* towards the Nose, but *rough* and *unequal* below, for the firm connection of the Membrane of the Palate. Tab. XII. Fig. 2.

The *Nasal Spine*, contributing, in a small degree, to the formation of the Septum of the Nose. Tab. XII. Fig. 2. b.

A *Depression* behind the Malar Process, where the under end of the Temporal Muscle plays. Tab. III. a.

A *Depression* at the under and fore part of the Malar Process, where the Muscles which raise the Upper Lip and Corner of the Mouth originate, and where a Branch of the Fifth Pair of Nerves is lodged, and commonly, a large Portion of Fat. Tab. III. I.

An *Arch* formed by the Palate Plate, both above and below, for enlarging the Cavities of the Nose and Mouth. Tab. VI. b.

A *Notch* forming the under and fore part of the Nostril, to the edge of which, and to the corresponding one of the Nasal Process, the Cartilages of the side of the Nose are connected. Tab. III. y.

The *Alveoli*, or *Sockets* for the Teeth, *porous* for the firmer adhesion of the reflected Membrane of the Gums,

and for the transmission of Blood-vessels into the substance of the Bone; the number of Sockets corresponding to the Fangs of the Teeth. Tab. XIV. Fig. 1. A, B, C, D.

The *Lacrymal Fossa*, which, with that of the Os Unguis, forms a passage for the Lacrymal Duct into the Nose. Tab. IV. s.

A *Canal* in the Orbital Plate, terminated anteriorly by the *Foramen Infra-orbitarium*, through which the Infra-orbital branch of the second part of the Fifth Pair of Nerves, with a branch of the internal Maxillary Artery, pass to the Face. Tab. XII. Fig. 1. d. Tab. 3. z.

The *Foramen Incisivum*, or *Palatinum Anterius*, behind the Fore-Teeth, common to both Bones below, but proper to each above, and filled with a Process of the Soft Palate, and with small Vessels and Nerves, which run between the Membranes of the Mouth and Nose. Tab. VI. C.

In some Subjects, there is a distinct *Ductus Incisivus*, leading from one or from each Nostril into the Cavity of the Mouth, similar to that which is always found in the large Quadrupeds.

A small *Hole* commonly found in the Nasal Process, and some *minute Passages* at the back-part of the Tuberosity, for the transmission of Blood-vessels and Nerves into the Substance of the Bone and Teeth, and into the Antrum Maxillare. Tab. III. t.

The *Sinus Maxillaris*, *Antrum Maxillare*, or, from its describer, called *Highborianum*, of great size, occupying the whole inner part of the Bone, situated under the Orbital Plate, and above the large Dentcs Molares, destined for the same purposes as the other Sinuses of the Bones of the Head. Tab. XII. Fig. 2. c, c. Tab. XIII. Fig. 3. L, M, N, O.

The Partition between the Sinus and Sockets of the Teeth is commonly of considerable thickness; but not unfrequently there is only a thin plate interposed, and small prominences, containing the points of the roots of the Teeth, may often be observed in the bottom of this Cavity.

The *Opening* of the Sinus, large in the separated Maxillary Bone, but, in the connected state, so covered by the inferior spongy Bone, and by the Palate-bone and Membranes, as to leave only one, or sometimes two Apertures, little larger than to admit the point of a Surgeon's Probe. The Aperture is situated at the upper part of the Sinus, and descends obliquely backwards, to terminate between the Offa Spongiosa, superius et inferius, in the Cavity of the Nose. Tab. XIII. Fig. 1. q.

The *Connection* of the Os Maxillare Superius, to the Frontal Bone, by the Transverse Suture, Tab. III. between e and p;—to the Os Unguis, by the Lacrymal Suture, Tab. III. between p and G;—to the Os Nasi, by the lateral Nasal Suture, Tab. IV. behind x;—to the Cheek-bone, by the internal and external Orbital Sutures, Tab. IV. 2.;—to the Os Planum, by the Ethmoid Suture, Tab. IV. between E and 10.;—to its fellow, by the Longitudinal Palate Suture, Tab. VI. 4.;—to its fellow

fellow also, between the fore-part of the Nose and Mouth, by the Mafachial Suture. Tab. III. *w.*

The *Substance* of this Bone is hard and denfe, except at the Alveoli, which are remarkably fpongy.

The Ossa Maxillaria form the greater part of the Nose, and Roof of the Mouth, a confiderable part of the Orbits, and contain all the Teeth which belong to the Upper Jaw.

In the Fœtus, there are Six Sockets for the Teeth,—no Tuberoſity, and the Maxillary Sinus is only beginning to form.

OSSA PALATI.

The *Situation* in the back-part of the Palate. Tab. VI. *f.*

The *Oblong Form* of the Palate-plate of each, which forms the back-part of the Offeous Palate. Tab. VI. *f.*

Its *poſterior curved Edge*, where it is connected with the Velum Palati; alfo the *Point* at the inner extremity of the Curve, for the Origin of the Muſcle of the Uvula. Tab. VI. *f.*

Its *thick, ſtrong Substance*, where it joins its fellow. Tab. XII. Fig. 2. *n.*

Its *Spinous Proceſs* at the inner edge of the Palate-plate, joining the under Edge of the Vomer, and contributing to the formation of the Septum Narium. Tab. XII. Fig. 2. between *l* and *n.*

The *Pterygoid Proceſs*, of a *Triangular form*, with *Foſſæ* correſponding to the Pterygoid Plates of the Sphenoid Bone. Tab. XII. Fig. 11. *B.*

The *Nasal Plate*, forming a portion of the ſide of the Nose, and Antrum Maxillare. Tab. XII. Fig. 2. *l.*

A *Ridge* on the inſide of this Plate, upon which the back-part of the inferior Spongy Bone reſts. Tab. XII. Fig. 2. *m.*

The *Two Orbital Proceſſes*, at the upper and back-part of the Naſal Plate, contributing a little in the formation of the Orbit, and of the Ethmoid and Sphenoid Sinuſes; being hollow within. Tab. XII. Fig. 2. *k.*

The *Anterior Orbital Proceſs*, the larger of the two, with its upper Surface appearing in the bottom of the Orbit, behind the back-part of the Os Planum and Os Maxillare.

A *Notch* between the Orbital Proceſſes, forming part of the *Foramen Spheno-palatium*, for the paſſage of the lateral Naſal Veſſels and Nerves. Tab. XII. Fig. 11. *F.* Fig. 2. under *k.*

The *Foramen Palatinum Poſterius*, or *Palato-Maxillare*, at the outer end of the Palate-plate of this Bone, but common to it and the Maxillary Bone, for the tranſmiſſion of the Palatine Veſſels and Nerve. Tab. VI. 4.

A *ſmall Hole* frequently obſerved behind the former, and communicating with it, for the paſſage of a Branch of the Palatine Nerve. Tab. XII. Fig. 2. *o.*

The *Foramen Spheno-Maxillare, Lacerum Inferius*, or

Inferior Orbital Fiſſure, at the under and outer part of the Orbit, and common to the Cuneiform, Maxillary, Malar, and Palatine Bones, for lodging Fat belonging to the Eye, and tranſmitting ſmall twigs of Veſſels and Nerves into the Orbit.

The Palate-plate of this Bone and its Pterygoid Proceſs are firm and ſtrong; but the Naſal Plate and Orbital Proceſs are thin and brittle.

The *Connection* of the Os Palati to the Palate Plate of the Maxillary Bone, by the Tranſverſe Palate Suture, Tab. VI. before *f*;—to the Maxillary Bone, at the ſide of the Nose and bottom of the Orbit, by the Palato-maxillary Suture, Tab. XII. Fig. 2. *f*;—to the Pterygoid Proceſs of the Sphenoid Bone, by the Sphenoid Suture;—to the Os Planum and Ethmoid Cells, by the Ethmoid Suture;—and to its fellow, by the longitudinal Palate Suture. Tab. VI. at the inſide of *f*.

In the Fœtus, the Palate-bone is complete, but there are no Cells in the Orbital Proceſſes.

OSSA SPONGIOSA, or TURBINATA, INFERIORA.

The *Situation* of each in the under part of the ſide of the Nose. Tab. XIII. Fig. 1. between *q* and *r.*

Its *Triangular form* and Spongy appearance, reſembling the Os Spongioſum Superius. Tab. XII. Fig. 9.

Its *Convexity* towards the Septum Naſi, and *Concavity* outwards. Tab. XIII. Fig. 8. *Q.*

The under edge placed horizontally near the under part of the Nose, and ending in a ſharp point behind.

The *two Proceſſes* at the upper part of the Bone, the anterior aſcending and forming part of the Lacrymal Groove, and the poſterior deſcending in form of a Hook, to make part of the Side of the Maxillary Sinus. Tab. XII. Fig. 9. *B.*

The *connection* of this Bone to the Os Maxillare, Os Palati, and Os Unguis, by a diſtinct Suture in the young ſubject; but in the Adult, by a concretion of ſubſtance.

The Oſſa Spongioſa afford a large ſurface for extending the Organ of Smell; by allowing the Membrane of the Nose to be expanded, upon which the Olfactory Nerves are diſperſed.

In the Fœtus, theſe Bones are almoſt complete.

OSSA TRIANGULARIA, or CORNUA SPHENOIDALES.

The *Situation* of each Triangular Bone between the body of the Sphenoid Bone and root of its internal Pterygoid Proceſs, covering the under part of the Sphenoid Sinus. Tab. X. Fig. 1. *b.*

Its *Connection* to the back-part of the Ethmoid Bone. Tab. X. Fig. 3. *k.*—In an old Perſon, it grows ſo firmly to the Sphenoid Bone, as to be conſidered by ſome Authors as one of its Proceſſes.

VOMER.

Its *Situation* in the under part of the Septum Nafi, where it separates the Nostrils from each other. Tab. III. L.

It is frequently bent to one side, in which case the one Nostril is rendered larger than the other.

Its *Form* compared to that of the Plough-share, from which it has its name. Tab. XIII. Fig. 2. H.

The *Superior* and *Posterior* Part, *thick* and *strong*, with a *Furrow* above to receive the Processus Azygos of the Sphenoid Bone. Tab. XII. Fig. 13. A. Tab. XIII. Fig. 2. behind G.

The *Superior Part*, with a *Groove* to receive the Na-

sal Plate of the Ethmoid Bone, and Cartilage of the Nose. Tab. XIII. Fig. 2. before G.

The *Inferior Edge* connected with the Spinous Processes of the Palate and Maxillary Bones, by a small Ridge corresponding with a Groove of these Bones. Tab. XIII. Fig. 2. under H.

The *Posterior Edge*, unconnected with any other Bone, and turned to the Cavity of the Fauces. Tab. XIII. Fig. 2. M. Tab. VI. g.

The Vomer has a smooth Surface, and a dense substance, and consists of two Plates in a young person, but in an old Subject the Plates are compressed together so as to render the Bone transparent.

TABLE

TAB. 12.

Fig. 1.



Fig. 2.



Fig. 3.



Fig. 4.



Fig. 7.




Fig. 8.



Fig. 5.




Fig. 9.



Fig. 11.




Fig. 10.



Fig. 12.



Fig. 15.



T A B L E XII.

Views of the different BONES of the FACE.

FIG. 1.

The Outer Side of the Right Os MAXILLARE SUPERIUS, with a small Portion of the Os PALATI.

- A, The maxillary fossa.
 B, The nasal process of the maxillary bone.
a, Inequalities by which it is joined to the os frontis.
b, The angle which is joined to the under end of the os nasi, and to the cartilage of the nose.
 C, The orbital plate.
c, The edge of the orbit.
d, A groove which belongs to the infra-orbital canal.
e, e, f, f, The malar process.
 D, D, The alveolar process.
 E, The maxillary tuberosity of the bone.
 F, A small portion of the os palati.
g, g, Two of many small holes which penetrate into the substance of the bone.
 G, The fore-part of the nostril.
h, The nasal spine, forming part of the partition of the nose.
i, The letter is placed upon the palate-plate, and points to the upper end of the palate-fissure.
 H, The fore-part of the malar process.
 I, The foramen infra-orbitarium.
 1. 1. The dentes incisores.
 2. The dens caninus.
 3. &c. The dentes molares.

FIG. 2.

Represents the Inner Surface of the same Os MAXILLARE SUPERIUS, with the Os PALATI.

- A, The nasal process, or upper angle.
a, The middle angle, at the base of the nasal process.
b, Inequalities, where the fore-part of the os spongiosum inferius is fixed.
 B, B, The palate-process.
 C, C, The alveolar process.
 D, D, The irregular surface of the palate-process.
c, c, c, c, The edges of the maxillary sinus.
 E, The cavity of the maxillary sinus.
d, d, Small cells in the upper part of this bone.
 F, The lacrymal fossa.
e, The palate-fissure which assists in forming the foramen incisivum.

- f*, The future which unites this bone to the os palati.
g, The part of this bone which forms the largest share of the nasal fossa.
h, The spine, which, by the union of its fellow, forms a small portion of the partition of the nose.
 G, *e*, The irregular surface, by which the fore-part of this bone joins its fellow on the opposite side.
 H, H, H, The os palati.
k, The small sinus, commonly found in this bone.
l, A portion of the palate-bone forming part of the fossa nasalis, and partition of the maxillary sinus.
m, An eminence by which this bone is connected to the os spongiosum inferius.
n, The rough surface where the two palate-bones unite.
o, The hole proper to this bone.
p, The foramen gustativum.
q, The pterygoid process.
 1. 1. The dentes incisores.
 2. The dens caninus.
 3. 3. The dentes molares.

FIG. 3.

The Outer Surface of the Ossa NASI.

- A, A, The upper part, which is joined to the frontal bone.
 B, B, The lower ragged end to which the cartilage of the nose is fixed.
 A, B, A, B, Holes penetrating the bone.

FIG. 4.

The Inner Surface of the Ossa NASI.

- A, A, The upper ragged end.
 B, B, The lower end, broader and thinner than the rest.
 C, C, The inner edge, thick and strong, where it joins its fellow, and sends a spine backwards, to be fixed to the partition of the nose.
 D, D, The cavity which forms part of the arch of the nose.

FIG. 5.

The Outer Surface of the Left Os UNGUIS.

- A, The lacrymal process, perforated by numerous holes.
 B, The orbital process.
 C, The ridge which separates the processes.

FIG.

FIG. 6.

The Inner Surface of the Os Unguis, with Eminences and Cavities which belong to the Ethmoid Cells.

FIG. 7.

The Outer Surface of the Right Os MALÆ.

- A, The superior orbital process.
- B, The inferior orbital process.
- C, The internal orbital plate.
- D, The maxillary process.
- E, The zygomatic process.
- F, The external orbital hole.
- G, G, The under and outer edge of the orbit.
- H, Part of the inner rough surface of the maxillary process.
- I, The zygomatic notch of the os malæ.

FIG. 8.

The Inner Surface of the same Os MALÆ.

- A, B, C, D, E, As in Fig. 7.
- F, The internal fossa, and situation of the external orbital hole.
- G, G, The rough edge which joins the os malæ to the superior maxillary bone at the external orbital foramen.

FIG. 9.

The External Surface of the Os SPONGIOSUM INFERIUS of the Right Side.

- A, The under edge of the bone turning outwards.
- B, The upper edge, sending down a hook-like plate to cover a portion of the maxillary sinus.
- C, The broad anterior extremity, where the connection is chiefly made with the superior maxillary bone.
- D, The posterior extremity, narrow and irregular in its surface.
- E, The external surface, with numerous small holes which mark its porosity.
- F, G, The superior edge, which joins the os unguis to form a share of the lacrymal groove.

FIG. 10.

The Internal Convex Surface of the same Os SPONGIOSUM INFERIUS, which, like the External Surface, is also of a spongy texture.

FIG. 11.

The Posterior, and almost the whole of the External Surface of the Left Os PALATI.

- A, The palate-plate.
- B, The pterygoid process.
- C, The nasal plate.
- D, The orbital process.
- E, A small sinus, corresponding with those of the ethmoid bone.
- F, The notch, which, with the body of the sphenoid bone, forms the foramen sphenopalatinum.
- G, A small hole which penetrates the thickness of the bone.
- H, Part of the groove which helps to form the foramen gustativum.

FIG. 12.

The Anterior, and almost all the External Surface of the same Os PALATI.

- A, The notch which assists in forming the foramen gustativum.
- B, The orbital process.
- C, The palate-plate.
- D, The nasal plate.
- E, The groove which helps to form the foramen gustativum.
- F, The pterygoid process.

FIG. 13.

The Left Side of the VOMER.

- A, The hollow surface which receives the processus azygos of the sphenoid bone.
- B, The anterior and upper edge, which is connected to the nasal plate of the ethmoid bone, and middle cartilage of the nose.
- C, The inferior edge, which is connected to the palate-plates of the superior maxillary and palate-bones.
- D, A ridge on the side of the vomer.

MAXILLA INFERIOR.

THE *Figure* of the Maxilla Inferior, or Lower Jaw, compared to that of the letter U, or it forms half of a long oval, with the convex middle part forwards.

The *Division*, into *Chin*, *Sides*, and *Processes*.

The *Chin*, extending between the holes termed *Mental Foramina*, at the fore-part of the Jaw. Tab. IV.

The *Sides*, reaching from the Mental Foramina to the back-part of the Bone.

A *transverse Ridge* on the fore-part of the Chin, with depressions on each side, for the origin of Muscles of the Under Lip. Tab. III. M.

Small Prominences and *Depressions* on the under and back part of the Chin, for the attachment of the Frænum Linguae, and of several Muscles which belong to the Throat. Tab. XXX. Fig. 13.

The *Base*, or *lowest part*, forming the under boundary of the Face. Tab. IV. XIX.

The *Angle* of the Jaw at the back-part of the Base. Tab. IV. XX.

Impressions made by the Masseter Muscle, upon the outside of the Angle, and also on the Plate, which arises from the back-part of the Bone. Tab. IV. XXI.

The *Plate*, which rises from the Angle of the Jaw, on each side, running upwards and a little backwards, and terminating in two Processes, termed *Condylod* and *Coronoid*. Tab. XIV. Fig. 2. F.

The *Condylod* or *Articular Process*, with an oblong rounded head, covered with Cartilage, and placed almost transversely, though, with respect to each other, the Condyles a little oblique, upon a Cervix at the upper and back part of the Bone.

The *Coronoid Process*, into which the Temporal Muscle is inserted, *situated* a little before the Condylod Process, and in the natural situation of the Jaw, placed on the Inner Side of the Zygoma. Tab. IV. XXIII.

The *Anterior Edge* of this Process, forming a Ridge which goes downwards and forwards, terminating at the outside of the Posterior Alveoli.

From the inner side of the Coronoid Process, another Ridge seen terminating nearly opposite to the former. To these Ridges, the Membranes of the Gums and Muscles belonging to the Mouth are fixed.

The *Alveolar Process*, and *Alveoli*, nearly similar to those of the Upper Jaw. Tab. XIV. II.

The *Alveolar Process*, extending along the Upper Edge of the Bone, from the Coronoid Process of one side to that of the other; and thickest behind, corresponding there with the increased thickness of the Teeth.

The Alveolar Process, composed of two Plates, and divided by cross Partitions, which, as in the Upper Jaw, mark the different Alveoli for the Fangs of the Teeth.

The Posterior Part of the Internal Plate, slanting inwards, and thinner than the External, giving the Jaw a twisted appearance.

Opposite the Alveoli, the External Plate swelling, and

giving a fluted form, which is observed in the whole extent of the Alveolar Process of the Upper Jaw, and in the fore-part of the Lower Jaw.

At the fore-part of the Jaw, the Alveoli are perpendicular, but turn inwards behind, where they are placed nearer the inner than the outer part of the Jaw.

The *Sockets* worn down by old age, in consequence of which the Teeth drop out, the Jaw becomes narrower, and, when the Mouth is shut, appears more prominent. Tab. 14. Fig. 7. A.

The *posterior Maxillary Foramen* at the root of the Condylod and Coronoid Processes, upon the inner side of the Jaw, for the passage of the Third, or inferior Maxillary Branch of the Fifth Pair of Nerves, with corresponding Blood-vessels. Tab. XXX. Fig. 13. 7.

A *small-pointed Process* at the inner edge of this Hole, where a Ligament goes off to be fixed to the Temporal Bone. Tab. XIV. Fig. 2. at the inner side of the Bone, opposite E.

Above the Hole, the Bone *marked* by the passage of the Nerve and Vessels, and below it, commonly, a *small Furrow* pointing out the course of a Nerve which goes to a Muscle and Gland under the Tongue.

Between the posterior Maxillary Foramen and the angle, the Bone *marked* by the insertion of the Internal Pterygoid Muscle. Tab. XXX. Fig. 13. 6.

The *Anterior Maxillary Foramen*, or *Mental Hole*, at the side of the Chin, where the remains of the Inferior Maxillary Nerve and Vessels come out. Tab. IV. XXVI. C.

The *Inferior Maxillary Canal*, running in the substance of the Bone, between the Posterior and Anterior Foramina, a little below the roots of the Teeth, and having many perforations, for the passage of small branches of Vessels and Nerves which supply the Jaw and Teeth.

The *Surface* of the Jaw, remarkably *hard*, and within, having *numerous Cells*, which surround the Maxillary Canals.

The *Articulation* of the Jaw by its Condylod Processes, with the Glenoid Cavity of the Temporal Bone, and also with the Tubercle at the root of its Zygomatic Process.

An *Intermediate moveable Cartilage*, placed in the Articulation of the Lower Jaw, in its gentler motion allowing the Condyle to remain in the Glenoid Cavity, but admitting it to advance upon the Tubercle or root of the Zygoma, when the Mouth is widely opened. Tab. XXX. Fig. 17. r.

In a Fœtus, the Lower Jaw is somewhat of a semicircular figure, and is composed of two Pieces, joined together in the middle of the Chin by the intervention of a Cartilage. This Union, termed *Symphysis*, gradually ossifies, and leaves no mark of any former division. Tab. XXVII.

TEETH.

T E E T H.

THE *Situation* of the Teeth in the Alveoli of the Jaws. Tab. III. and IV.

Their *Number*, Sixteen in each Jaw. Tab. III.

The *Base*, or *Body* of each Tooth, appearing without the Sockets. Tab. XIV.

The *Roots*, or *Fangs*, placed in the Sockets, and of a *Conical* form. Tab. XIV.

The *Cervix*, or *Collar*, between the Base and Roots of the Teeth. Tab. XIV.

The *Roots* of the Teeth covered by a *Vascular Membrane*, reflected from the Gums, and serving as a *Perio-steum* to the Teeth, and a lining to the Alveoli.

The *Cortex*, or *Enamel*, covering the base of each Tooth, and becoming gradually thinner towards the *Cervix*.

The *Fibres* of the Enamel placed perpendicular to the Osseous Substance, to diminish the effects of Friction.

The *Fibres* of the Osseous Part of the Teeth forming *Lamellæ*, which run according to the length of the Teeth.

A *Foramen* at the point of the root of each Tooth, and a Passage leading from it into a common Cavity in the Base of the Tooth, for lodging the Vascular and Nervous Substance called Pulp of the Teeth.

Division of the Teeth into *Three Classes*, viz.

—On each side of each Jaw,—

Two *Incisores*:—One *Cuspidatus*:—Two *Bicuspides*:—and *Three Molares*. Tab. XIV.

The *Incisores*, having their *Bases* formed into *Wedges*, *shaped out behind*.

The *Cuspidus*, having its *Base* in form of a *Wedge*, like the *Incisores*, but pointed in the middle.

The *Bicuspides*, each with *double points*, one external, the other internal, which, in the Upper Jaw, are nearly upon a level, but, in the Under Jaw, highest on the outside of the Teeth.

The *Incisores*, *Caninus*, and small *Molares*, with *single Roots*, excepting the small *Molares* of the *Upper Jaw*, which have frequently *two Roots*.

The first of the three posterior, or large *Molares* of the *Under Jaw*, with *five*, and each of the other two with *four points*.

Each of these three Teeth having *two, three*, or sometimes *four roots*.

In the *Upper Jaw*, the first large *Molaris* having *four*, and each of the other two only *three points*.

In each of these three Teeth, generally *one root more* in those of the *Upper*, than in the corresponding Teeth of the *Under Jaw*.

The last, or backmost *Molaris*, called *Sapiens*, *smaller*, and having generally *fewer roots*.

The Teeth *connected* to the Sockets by *Gomphosis*, and by a firm adhesion to the Gums.

At Birth, the outer Shell only of the five deciduous Teeth, and of the anterior permanent *Molaris*, in each side of each Jaw is found.

These Teeth are situated in Capsules, within the Jaw, and under its surface. At this period there are no roots formed.

For a fuller description of the Teeth, see VOL. II.

TABLE

TAB. 13.

Fig. 1.





Fig. 2.



Fig. 3.



T A B L E XIII.

Gives different VIEWS of the CAVITY of the NOSE.

FIG. 1.

The Left Portion of the BASE of the HEAD divided from the Septum Narium by a Perpendicular Section, proceeding in a straight line from before, backwards.

- A, Part of the os frontis.
 B, The posterior lamina, called Vitrea.
 C, The frontal sinus.
 D, Part of the transverse suture, dividing the frontal from the superior maxillary bone.
 E, Part of the frontal bone, contiguous to the os ethmoides.
 F, The upper part of the ethmoid bone.
 G, The fore-part of the ethmoid cells entire. From a little behind G, to H, the anterior, middle, and posterior ethmoid cells laid open.
 I, Openings of the ethmoid cells into the nose.
 K, The last and uppermost passage of the nostrils.
 L, The left anterior clinoid process of the sphenoid bone.
 M, The posterior clinoid process.
 N, The Sella Turcica.
 O, The left sphenoid sinus.
 P, The part where the sinus opens into the posterior and upper passage of the nostril.
 Q, R, A section of the back-part of the sphenoid, and cuneiform process of the occipital bone.
 S, The spinous process of the occipital bone.
 T, The internal pterygoid plate.
 U, The uncus of the sphenoid bone.
 V, The fore-part of the meatus auditorius.
 W, The superior condyloid foramen, for the passage of the ninth pair of nerves.
 X, The mastoid process of the temporal bone.
 Y, Part of the os occipitis.
 Z, The inner side of the occipital bone.
 a, The cut edge of the occipital bone.
 b, The under and outer part of this bone.
 c, The nasal process of the superior maxillary bone.
 k, The inner side of this bone, which forms the middle passage of the nostril.
 e, Part of the same bone, which forms the beginning of the lower passage of the nostril.
 f, A section of the alveolar process.
 g, A section of the osseous palate.
 h, The upper part of the osseous palate.

- i, The descending, or alveolar part of the palate.
 k, l, m, n, The os spongiosum superius. Between l and n, the part resembling a concha.
 o, p, The middle passage of the nostril.
 q, The opening of the antrum maxillare. Between q and r, the os spongiosum inferius.
 r, A part of the inferior spongy bone, opposite the opening of the lacrymal duct.
 s, t, The lowest passage of the nostril.

FIG. 2.

The Left Surface of the SEPTUM NARIUM.

- A, The os frontis, with its plates and diploe.
 B, The frontal sinus.
 C, The crista Galli.
 D, Part of the os planum of the left side, having no ethmoid cells.
 E, The foramina cribrosa of the ethmoid bone.
 F, The nasal plate of the ethmoid bone, which forms part of the septum narium.
 G, That part of the nasal plate of the ethmoid bone where it is joined to the vomer.
 H, The vomer.
 I, The cartilaginous part of the septum narium.
 K, Part of the upper jaw.
 L, L, The dentes incisivi of the upper jaw.
 M, The posterior edge of the vomer, covered with a membrane.
 N, Part of the cuneiform process of the occipital bone.
 O, The right sphenoid sinus.
 P, The posterior clinoid process of the sphenoid bone.
 Q, The anterior clinoid process of the sphenoid bone.
 R, The Sella Turcica.
 S, A portion of the septum, between the two sphenoid sinuses.
 T, A partition between the sphenoid sinus and the nostrils.


FIG. 3.

The Anterior and Right Portion of the BASE of the HEAD, taken off by a Longitudinal Section separating it from the Septum, and by a Transverse Section separating it from the Posterior Part. The two cut Surfaces of this Preparation are represented.

F

A, The

- A, The frontal sinus.
B, The opening of the frontal sinus into the first ethmoid cell, which is seen on each side of C.
D, The nasal process of the superior maxillary bone.
E, A section of the os spongiosum superius.
F, The wall, or boundary of the middle passage of the nose, by which it is separated from the superior appendix of the maxillary process.
G, Cells in the superior maxillary bone and its orbital part, first described by *Haller*. The uppermost of these opens into the anterior ethmoid cell.
H, The orbital plate of the superior maxillary bone, forming part of the bottom of the orbit, and upper part of the antrum maxillare.
I, The orbital plate of the frontal bone.
K, The os malæ.
L, M, N, O, P; The sinus, or antrum maxillare.—O, The partition by which the antrum maxillare is separated from the cavity of the nostril.—P, The foramen by which the antrum communicates with its appendix.
Q, A section of the os spongiosum inferius.
R, The orifice of the lacrymal groove.
S, Part of the palate.
T, The lateral dens incisivus.
U, The dens caninus.
V, W, The two small molares.



TAB. 14.



T A B L E XIV.

Gives different VIEWS of the BONES of the EAR ;—of the JAWS and TEETH ;—and likewise a VIEW of the Os HYOIDES.

FIG. 1.

Represents the SOCKETS of the Upper Jaw, and OSSEOUS PART of the Palate.

- A, B, C, The sockets of the dentes incisores, dens caninus, and two small molares of the right side, single.
 D, The sockets of the three large molares, each with three cavities.
 E, The palate-plate of the superior maxillary bone.
 F, The palate-plate of the os palati.
 G, The pterygoid plates of the sphenoid bone.

FIG. 2.

A View of the SOCKETS of the Lower Jaw.

- A, B, C, The sockets of the incisores, canini, and small molares, single.
 D, The sockets of the large molares, each with two cavities.
 E, E, The coronoid processes.
 F, F, The condyloid processes.

FIG. 3.

A View of the Fangs of the Teeth in both Jaws, the number of Fangs corresponding to that of the Sockets seen in Fig. 1. and 2.—The External Plate of the Alveolar Process is removed.

FIG. 4.

Views of the different Classes of TEETH of both Jaws.

- A, B, C, The posterior surface of an incisor of the upper jaw ;—A, Its body ;—B, Its cervix, where the enamel ends ;—C, Its straight fang.
 D, An incisor of the lower jaw, smaller than that of the upper one, A, B, C.
 E, The posterior surface of a canine tooth, with its pointed body and large fang.
 F, A small molaris, with its straight single fang.
 G, A large molaris of the under jaw, with its fangs crooked at the points.
 H, A large molaris of the upper jaw, with three diverging fangs.
 I, A molaris of uncommon size, with an appearance of four fangs.

FIG. 5.

A View of the DECIDUOUS, or MILK TEETH, in various degrees of Growth.

- A, The body of an incisor, the fang not yet evolved.
 B, The body of a small molaris.
 C, The body of a canine tooth.
 D, The body of a large molaris.
 E, The body and part of the fangs of a large molaris.
 F, The body and part of the fangs of a large molaris, more advanced than E.
 G, A canine tooth nearly complete.
 H, I, Two incisores nearly complete.

FIG. 6.

A View of the Disposition of the two Sets of TEETH, in both Jaws, at the time of shedding the Milk-Teeth.

1. 1. &c. The deciduous, first set, or milk-teeth.
2. 2. &c. The deciduous, adult, or second set of teeth, proceeding to the edge of the alveoli.

FIG. 7.

A View of the Upper and Under Jaws of an Old Person, without the Teeth, and the Alveoli obliterated, whence the Jaws are narrower, the Chin much more prominent, and the Cavity of the Mouth diminished.

FIG. 8.

A magnified View of the SMALL BONES of the EAR, articulated with each other, and covered with their Periosteum, in which the Blood-vessels appear.

- A, The incus.
 B, The malleus.
 C, The stapes.
 D, The os orbiculare *in situ*.

FIG. 9.

Views of the SMALL BONES of the EAR.

The upper set gives a view of the small bones of the ear, of their natural size, and as they are connected with each other.

The under set shews these bones separated from each other, and somewhat magnified.

- A, The incus, with its body, articular cavity, short posterior, and long inferior branch.
- B, The malleus, with its head, neck, cavity of articulation with the incus, great process or handle, middle process, and long slender one.
- C, The stapes, with its head, crura, and base.
- D, The os orbiculare.

FIG. 10.

The Temporal Bone of a Young Subject, with the Small Bones of the Ear in situ,—the Membrana Tympani being removed.

FIG. 11.

The Temporal Bone, with the Membrana Tympani in situ, the Small Bones shining through it.

FIG. 12.

An External View of the RIGHT LABYRINTH, and Outlines of the PARS PETROSA of a Young Subject.

In the labyrinth are seen,
Anteriorly,—the cochlea.
Posteriorly,—the semicircular canals.
Superiorly,—the foramen ovale.
Inferiorly,—the foramen rotundum.

FIG. 13.

A View of the Upper and Fore Part of the OS HYOIDES.

- A, The body of the os hyoides.
- B, Its cornua.
- C, Its appendices.

OS HYOIDES.

THE *Situation* of the Os Hyoides, at the root of the Tongue and top of the Larynx.

The *Shape*, compared to that of the Greek letter ν . Tab. XIV. Fig. 13.

The *Body* of the Bone, the middle broad part, *convex* before, and *concave* behind.

The *concavity* behind, oblique, to receive the Thyroid Cartilage, when the Os Hyoides and Larynx are pulled towards each other.

Several *Impressions* seen on its Body, occasioned by the numerous muscles fixed to it.

The *Cornua*, extending backwards and upwards from each side of the Body, with their two plain Surfaces slanting from above downwards and outwards, and giving attachment to Muscles and Ligaments of the Tongue and Larynx.

Each of the Cornua becoming gradually smaller in its course backwards, and ending in a round Tubercle, which is connected to the upper Cornu of the Thyroid Cartilage.

Between the Body and Cornua frequently a Furrow, pointing out the former separation in young subjects.

The *Appendices*, placed at the upper part of the Articulation between the Body and Cornua, for the attachment of Muscles.

From each Appendix, a *Ligament* sent up to the Styloid Process of the Temporal Bone. Tab. XXXI. Fig. 1. 2.

The Os Hyoides is not immediately connected to any other Bone, but is kept in its place by numerous Muscles and Ligaments, to be afterwards mentioned.

The Substance of this Bone is cellular, but covered with a firm external Plate, which adds considerably to its strength.

At birth, the greater part of the Bone is in a Cartilaginous state, and the Appendices continue so for many years after the other parts are completely ossified.

The Os Hyoides serves as a Lever for numerous Muscles acting upon the Tongue, Larynx, and Fauces.

T R U N K.

THE Trunk, composed of the Spine, Pelvis, and Thorax.

The *Spine*, reaching from the Condyles of the Occipital Bone, to the lower end of the Os Coccygis. Tab. II. G, P.

The Spine appearing *straight*, when viewed anteriorly or posteriorly. Tab. I. II.

The several *Curvatures* of the Spine, when viewed in a lateral direction; the Curvatures accommodating themselves to the soft parts of the Neck, Thorax, Abdomen, and Pelvis. Tab. XV. XVI.

The Spine, composed of a long upper, and a short under Pyramid, joined together by their Bases. Tab. II. b.

The upper Pyramid, composed of *true Vertebrae*, or Bones which turn upon each other. Tab. G.—N.

The under Pyramid, formed of *false Vertebrae*, or Bones which, at an early period of life, resemble the true Vertebrae, but afterwards grow together, so as not to contribute to the motions of the Trunk of the Body. Tab. II. b.—P.

THE TRUE VERTEBRÆ,

Twenty-four in Number.

EACH of the true Vertebrae composed of a Body and Processes.

The *Body* of a spongy nature, with *upper* and *under Surfaces* placed horizontally. Tab. XXXI. Fig. 11. a.

The anterior *convexity* of the Body, and posterior *concavity*. Tab. XVIII. Fig. 9. 10.

Numerous *small Holes* on the anterior and lateral parts of the Body, for the passage of Blood-vessels into the Substance of the Bone, or for the attachment of Ligamentous Fibres. Tab. XVI. Fig. 8. a.

A *Ring of Bone*, at the upper and under edges of the Body, of a firmer texture than the rest of its Substance, and thereby adding to the general strength of the Bone. Tab. XVIII. Fig. 9. b, b. Tab. XXXI. Fig. 12. r.

The Ring of Bone forming a *superficial Cavity*, which receives the Intervertebral Substance. Tab. XVIII. Fig. 9. a.

The Bodies of the Vertebrae in general smaller and more solid above; as they descend, they become larger and more spongy.

The

The *Intervertebral Substances*, of a *Cartilago-ligamentous nature*, placed between the Bodies of the *Vertebræ*, for fixing them together, and allowing the Spine to be moved in all directions. Tab. I. E. Tab. XXXI. Fig. 15. u.

The *Intervertebral Substances*, composed of *Concentric Lamellæ*, with their edges firmly fixed to the Bodies of the *Vertebræ*.

The *Lamellæ* of these *Substances*, formed of *Oblique Fibres*, which decussate each other, and are very compressible.

The *Centre* of these *Substances* changes from *Lamellæ*, and puts on the appearance of a *Mucus* or *Pulp*, which has little compressibility, and serves as a *pivot* upon which the other parts of the *Ligament* can move with such gradual yielding as to lessen shocks in the Spine in violent motions of the *Body*.

The *Intervertebral Substances*, like the *Vertebræ* themselves, *larger* and *thicker* as they descend, to give greater security to the parts they support. Tab. I.

An *Arch* sent out from the back-part of the *Body* of each *Vertebra*, which, together with the *Body*, forms a *large Hole* for the passage of the *Spinal Marrow*. Tab. XVIII. Fig. 3. b.

A *Notch* at the upper and under edge of each side of the *Arch*, for the passage of the *Spinal Nerves*. Tab. XVIII. Fig. 10. d, d. Fig. 8. c, c.

The *two Superior Oblique*, or *Articulating Processes*, covered with *Cartilage*, placed upon the upper part of the sides of the *Arch*. Tab. XVIII. Fig. 3. b, b.

The *two Inferior Oblique*, or *Articulating Processes*, also covered with *Cartilage*, and placed upon the under part of the sides of the *Arch*. Tab. XVIII. Fig. 4. b, b.

The *two Superior Oblique Processes* of one *Vertebra*, articulated with the *two Inferior Oblique* of the *Vertebra* immediately above it. Tab. XVIII. Fig. 5. e, m.

Round the *Edges* of the *Oblique Processes*, *rough Lines* for the attachment of their *Articulating Ligaments*.

The *two Transverse Processes* projecting from the sides of the *Arch*, and between the *Oblique Processes*. Tab. XVIII. Fig. 10. e, e.

The *Spinous Process*, sent out from the back-part of the *Arch*, which, being sharp and pointed, gives name to the whole chain of *Bones*. Tab. XVIII. Fig. 7. d, i.

The *Edges* of the *Processes*, as well as of the *Body*, *rough*, where *Ligaments* come off which fix them to each other.

The *Substance* of the *Processes* stronger, with a thicker external *Plate* than in the true *Vertebræ*.

The *Vertebræ* are joined to each other by a double *Articulation*; their *Bodies* being connected by the *Intervertebral Substances* already described, and their *Oblique Processes* are so connected by their *Ligaments* as to allow a small degree of motion to all sides.

The uses of the true *Vertebræ* are, to give an erect posture to the *Trunk* of the *Body*; to allow a sufficient

and secure motion to the *Head*, *Neck*, and *Trunk*; and to support and protect the *Bowels* and other soft parts.

In the *Fœtus*, each *Vertebra* consists of three pieces connected by *Cartilages*, viz. the *Body* not fully ossified; a curved *Bone* on each side, on which are a small share of the bony *Arch*, the *Oblique Processes* complete, the beginning *Transverse Processes*, but no *Spinal Processes*.

The *Vertebræ* divided into seven *Cervical*, twelve *Dorsal*, and five *Lumbar*.

The *Cervical Vertebrae*, or *Vertebrae of the Neck*, having their *Bodies smaller*, more flattened before and behind, and more hollowed above and below, than those of the other *Vertebrae*. Tab. XV. A, A, &c.

The *Articulating Processes*, more *Oblique* than the rest. Tab. XVIII. Fig. 5. e, m.

The *Transverse Processes*, perforated for the passage of the *Vertebral Blood-vessels*, and hollowed above for the transmission of the *Spinal Nerves*. Tab. XVIII. Fig. 2. e, e.

The *Spinal Processes*, placed horizontally, shorter than the rest, and forked for the attachment of *Muscles*. Tab. XVIII. Fig. 6. r, r.

The *Cervical Vertebrae* admit of *free motion*, in consequence of the thickness of their *Cartilages*, and the nature of their *Processes*.

The first *Vertebra*, called *Atlas*, from its supporting the *Globe* of the *Head*, having only a small *Arch* instead of a *Body*. Tab. XVIII. Fig. 1. a.

The *Upper* and *Under Surfaces* of the *Arch*, marked by the *Ligaments* which fix it to the *Head* and second *Vertebra*. Tab. XVIII. Fig. 1. 2.

The back-part of the *Arch*, hollow, and covered by a smooth *Cartilage*, where it turns upon the *Processus Dentatus* of the second *Vertebra*. Tab. XVIII. Fig. 2. a.

The *Inner Parts* of the sides of the *Vertebra*, between the *Superior* and *Inferior Oblique Processes*, marked by the *Lateral Ligaments* which go to the *Processus Dentatus*, and by the *Transverse Ligament* which passes behind that *Process*. Tab. XVIII. Fig. 2. c, c. Tab. XXI. Fig. d, d.

An *Arch* upon the back-part of the *Atlas*, instead of a *Spinous Process*, marked by *Muscles* and *Ligaments*. Tab. XVIII. Fig. 2. g.

The *Superior Oblique Processes*, oval and hollow, for receiving the *Condyles* of the *Occipital Bone*. Tab. XXI. Fig. 1. D, D.

A curved *Fossa* under the outer and back part of each *Oblique Process*, for the passage of the *Vertebral Arteries* into the *Head*, and *Tenth Pair* of *Nerves* out of it. Tab. XXI. Fig. 1. f.

The *Transverse Processes*, longer than in any other *Cervical Vertebra*, for the origin of several *Muscles*. Tab. XVIII. Fig. 5. d, d.

The *Connection* of the *Atlas* to the *Occipital Bone*, where the *Head* has its *flexion* and *extension*, but little other motion.

The second Vertebra, called *Dentata*, from the Tooth-like Process on the upper part of its Body.

The *Body* of this Vertebra larger than the rest, and of a *Conical* figure. Tab. XVIII. Fig. 5. *b—l*.

The fore-part of the *Processus Dentatus*, *convex* and covered with *Cartilage* where it turns upon the Atlas. It has the same appearance behind, where it moves upon the Transverse Ligament. Tab. XVIII. Fig. 5. 6.

The *Sides* of this Process, marked by the insertion of the lateral Ligaments, and its *Point* by the insertion of the perpendicular Ligament which is fixed to the Edge of the Foramen Magnum of the Occipital Bone. Tab. XVIII. Fig. 4. *a*.

The *Superior Oblique Processes* placed horizontally, and elevated in the middle, to be received into the hollow Inferior Oblique Processes of the Atlas, where the Head has its principal rotatory motion. Tab. XVIII. Fig. 3. *b; b*. Fig. 5. *m, m*.

The *Spinous Process*, thick and strong, to give origin to the Muscles which assist in the extension and rotation of the Head, and turned down to allow these motions to be readily performed. Tab. XVIII. Fig. 3. *g, g*. Tab. XXXI. Fig. 5. *d, e*.

In the Fœtus, the Vertebra *Dentata* consists of four pieces, three of which are common to all the Vertebrae, the fourth is the *Processus Dentatus*, which is joined by Cartilage to the Body of the Bone.

The *seventh Cervical Vertebra*, approaching to the form of the Dorsal Vertebrae.—The Spinal and Transverse Processes have no bifurcation. Tab. XXXI. Fig. 17.

The *Dorsal Vertebrae*, or *Vertebrae of the Back*, horizontal above and below, having their bodies larger, sharper before, flatter at the sides, and more hollow behind, than those of the Cervical Vertebrae. Tab. XVIII. Fig. 7. *F*. Tab. XXXI. Fig. 10.

A *Pit*, lined with Cartilage at each side of their upper and under Edges, near the Transverse Processes, for the articulation of the Heads of the Ribs. Tab. XXXI. Fig. 15. *t, t, W*.

The *Intervertebral Substances*, *thin*, to admit only of little motion, and *thinnest* anteriorly, to enlarge the Curvature of the Spine, and increase the Cavity of the Thorax. Tab. XV.

The Spinal Canal is here more circular, but corresponding with the size of the Spinal Marrow,—is smaller than in any of the other Vertebrae.

The *Oblique Processes*, having nearly a perpendicular direction, the upper ones slanting forwards, and the under ones backwards. Tab. XXXI. Fig. 9. *b*. Fig. 10. *k*.

The *Transverse Processes*, long, turned obliquely backwards, and enlarged at their outer extremity, where they are faced with Cartilage, to be articulated with the Tubercles of the Ribs. Tab. XVIII. Fig. 7. Fig. 8. *f, f*.

The *Spinous Processes*, long, thick at the roots, but slender near the extremities, and pointing obliquely downwards over each other, by which the Spinal Marrow in this part is well protected. Tab. XVIII. Fig. 7. Tab. XXXI. Fig. 17.

The upper Edge of each of the Spinous Processes of these Vertebrae, formed into a *Ridge*, which, in certain motions of the Spine, is received by a Groove in the under part of the Spinous Process of the Vertebra immediately above it. Tab. XVIII. Fig. 7. 8.

The last peculiarity of Structure, with the others already mentioned, prevent the Dorsal Vertebrae from having much motion.

The *first Dorsal Vertebra* having the whole of the Pit for the Head of the first Rib formed in it.

The *twelfth Dorsal Vertebra* receiving the whole Head of the last Rib, and having no Cartilaginous Surface on its Transverse Process.

The *Lumbar Vertebrae*, or *those of the Loins*, having their Bodies larger and broader than those of the other two classes. Tab. XVIII. Fig. 9. 10.

The *Intervertebral Substances*, the thickest of any, and most so at their fore-part, by which the Spine is rendered convex there, for the support of the Abdominal Bowels. Tab. XVI.

The *Oblique Processes*, remarkably deep, and placed upright, the Superior Oblique Process of one Vertebra facing inwards, and receiving the Inferior Oblique Process of the Vertebra below it, which is turned in the opposite direction. Tab. XXXI. Fig. 12. *b*. Fig. 13. *k*.

The *Transverse Processes*, long, slender, and almost erect, to give origin to large Muscles, and to admit of free motion. Tab. XVIII. Fig. 9. 10.

The *Spinous Processes*, short, large, and strong, and placed horizontally, with narrow edges above and below, and broad flat Sides, giving origin to Muscles of great strength. Tab. XVIII. Fig. 9. 10. Tab. XXXI. Fig. 12. 13.

The *Spinal Canal*, larger than in the back, for the passage of the Cords of the Spinal Marrow which form the Cauda Equina. Tab. XVIII. Fig. 9.

In consequence of the thickness of the Intervertebral Substances, and the situation of the Processes of the Lumbar Vertebrae, the motion of this part of the Spine is extensive, though not so much so as in the Neck.

TABLE

T A B L E XV.


An Anterior and Lateral View of the Upper Part of the TRUNK of the SKELETON.

-
- A, &c. THE bodies of the cervical vertebræ, with their intermediate cartilages.
 B, C, The transverse processes of the cervical vertebræ, with a hole in each of the processes, forming a canal for the vertebral blood-vessels.
 D, The processus dentatus of the second vertebra of the neck.
 E, &c. The bodies of the dorsal vertebræ, with their intermediate cartilages, forming a curve backwards.
 F, &c. The transverse processes of the dorsal vertebræ.
 G, &c. The outer convex surface of the ribs.
 H, &c. The cartilages of the seven upper or true ribs, by which they are joined to the sternum.
 I, &c. The cartilages of the false ribs, the three uppermost of which are joined together.
 K, Head of the first rib, joined to the first dorsal vertebra.
 L, The tubercle of the first rib, joined to the transverse process of the first dorsal vertebra.
 M, Head of the second rib, joined to the first and second dorsal vertebræ, and its tubercle fixed to the transverse process of the latter of these two bones.
 O, The upper triangular piece of the sternum.—P, The middle or long piece.—Q, The lower piece, or ensiform cartilage.
 R, S, T, U, V, The scapula.—R, Glenoid cavity of that bone.—S, Acromion of the scapula.—T, Coracoid process of the scapula.—U, Inferior costa of the scapula.
 W, The clavicle joined to the sternum, and at X, to the scapula.
 Y, The anterior convexity of the clavicle.

TABLE

TAB. 15.

D



A. F. sc.

T A B L E XVI.

AN Anterior and Lateral VIEW of the Under Part of the TRUNK of the SKELETON.


- A, THE body of the last dorsal vertebra.
- B, &c. The anterior extremities of the four lowest ribs of the left, and two lowest of the right side.
- C, &c. The cartilages of the four lowest ribs of the left, and two lowest of the right side.
- D, &c. The bodies of the lumbar vertebræ, forming an arch forward.
- d, &c. The intervertebral cartilages.
- E, &c. F, &c. Four lowest transverse processes on the left, and points of the three lowest on the right side.
- G, &c. The points of the three spinous processes.
- H, The upper piece of the os sacrum, joined to the last lumbar vertebra.
- I, &c. The five original pieces, which compose the os sacrum, grown together, but leaving traces of their former divisions, near the parts where the letters are placed, and forming an arch backwards, whereby the cavity of the pelvis is considerably enlarged.
- K, &c. Slanting holes opposite the original interstices of the pieces of which the os sacrum is composed.
- a, The brim of the pelvis.
- L, The inner hollow side, or venter of the os ilium.
- b, The connection between the os sacrum and os ilium. —A little below *b*, on the right side, the passage for the principal blood-vessels of the bone.
- M, N, The spine of the os ilium.
- O, The anterior-inferior spinous process of the os ilium.
- P, The point of union between the os ilium and os pubis.
- Q, The ischiatic notch.
- R, Part of the outer surface, or dorsum of the os ischium.
- T, The crus of the os ischium, joining the crus of the os pubis.
- U, The tuberosity, forming the lowest part of the trunk of the skeleton.

T A B L E XVII.

A View of the Posterior TRUNK of the SKELETON.

- A**, THE upper part of the first cervical vertebra.
B, One of the oblique, and,
C, One of the transverse processes of this bone.
D, Muscular prints on the back-part of this bone.
E, E, The spinous processes of the six other cervical vertebrae, of which the four first are forked.
F, F, The oblique processes of these vertebrae.
G, G, The transverse processes.
H, H, The spinous processes of the three first vertebrae of the back.
I, I, The spinous processes of the six middle vertebrae, which are long, and sloping downwards over each other.
K, K, The spinous processes of the three last dorsal vertebrae, which are short and straight.
L, L, The transverse processes of all the dorsal vertebrae.
M, M, The oblique processes of all these vertebrae.
N, N, The spinous processes of the lumbar vertebrae.
O, O, The transverse processes of these vertebrae.
P, P, The oblique processes of the same vertebrae.
Q, Q, Part of the bodies of the same bones.
R, R, The arches of these bones which form the back-part of the spinal canal.
S, S, The spinous processes of the os sacrum.
T, One of the lateral and superior tuberosities of this bone.
U, The superior orifice of that part of the spinal canal which belongs to this bone.
V, One of the superior oblique processes of the os sacrum.
W, W, The holes in the back-part of the os sacrum, which transmit small vessels and nerves to the parts adjacent.
X, X, The eminences and cavities at the lateral parts of this bone.
Y, One of the cornua of the os sacrum.
Z, The inferior orifice of the spinal canal.
a, The first or uppermost piece of the os coccygis.
b, b, The posterior extremities of the ribs.
c, c, The necks of the ribs.
d, d, The angles of the same bones.
e, e, The cartilages of the false ribs.
f, The outer surface of the os ilium.
g, g, The posterior spinous processes of this bone.
h, h, The great posterior tuberosity of this same bone.
i, The spine of this bone.
k, A portion of the anterior tuberosity of this bone.
l, The posterior edge of the acetabulum.
m, The ischiatic notch.
n, The spinous processes of the os ischium.
o, A portion of the internal surface of the superior branch of the os pubis.
p, The tuberosity of the os ischium.
q, The internal surface of the branch of this bone.
r, The foramen ovale.

FALSE



FALSE VERTEBRÆ.

THE FALSE VERTEBRÆ, composed of the Os Sacrum and Os Coccygis.

OS SACRUM.

The *triangular form* of the Bone, with its pointed under extremity. Tab. XIX.

The *flat concave anterior Surface*, for enlarging the Cavity of the Pelvis. Tab. XIX.

The *under and fore part*, forming a *turn*, called by some the *Lesser Angle* of this Bone. Tab. with Nerves of Pelvis.

The *convex irregular Surface behind*, where strong Muscles arise, which assist in extending the Spine and Thigh. Tab. XXXI.

Four transverse prominent Lines seen anteriorly, indicating the situation of the Cartilages which originally divided the Bone into five pieces. Tab. XIX.

The *Spinal Canal*, of a triangular form, of great size above, but becoming gradually smaller in its descent; corresponding to the size of the under end of the Spinal Marrow, termed *Cauda Equina*, which goes through it. Tab. XXXI. Fig. 18.

The *under part* of the Spinal passage, commonly *open behind*; the Canal being completed, in the Subject, by the addition of a strong Ligamentous Membrane. Tab. XVII.

The *Arch* at the sides and back-part of the Spinal Canal, much thicker and stronger than in the True Vertebræ. Tab. XXXI. Fig. 18.

The *two Oblique Processes* belonging to this Bone, facing backwards, to correspond with the two inferior Processes of the last Lumbar Vertebra. Tab. XIX.

A *large Oblong Process* on each side of the Bone, formed by the concretion of all the original Transverse Processes. Tab. XIX.

The upper lateral parts of the Bone, which correspond with the three superior Transverse Processes, divided into *two irregular Cavities* on each side, by a *perpendicular Ridge*. Tab. XIX.

The *anterior* of the two cavities lined with *Cartilage*, which glues this Bone to the Os Ilium, and in such a manner as not to allow any motion.

The *posterior Cavity*, *rough and irregular*; and in the recent Subject, full of *Ligamentous Fibres* and *Cellular Substance*, which are included in the general Capsular Ligament, and also assist in fixing the two Bones to each other.

The *Spinous Processes*: The three uppermost commonly *distinct*, but remarkably *short*: There is a great variety, however, in the number and appearance of the Spinous Processes in different Bones, and consequently of

the length of the complete part of the Spinous Canal. Tab. XIX.

Four Pair of large Holes on the *anterior Surface* of the Bone, at the end of the Lines already described, and *Grooves* running out from the Holes, for the passage of the Sacral Nerves. Tab. XIX.

Four Pair of Holes on the *posterior Surface*, not much smaller than those seen anteriorly; but so filled with Cellular Substance, and covered with Membranes in the recent Body, as only to admit small Nerves to pass out to the Muscles on the back-part of the Pelvis, and minute Arteries to enter to the Cauda Equina. Tab. XIX.

A *Notch* at the under end of each side of the Bone, or a *Hole* common to it and the Os Coccygis, for the passage of the last Spinal Nerve. Tab. XIX.

The *Substance* of the Os Sacrum, like that of the other Vertebræ, is very *spongy*, and covered only by a thin external Plate; this, however, is rendered considerably stronger by a Ligamentous Membrane which adheres to it. Tab. with Nerves of Pelvis.

The *Connection* of this Bone above to the last Lumbar Vertebra, just as the other Vertebræ are connected to each other, and the same motions allowed as to these Vertebræ.—The projection formed between these two Bones anteriorly, obtains the name of *Promontory* or *Greater Angle* of the Os Sacrum. Tab. XVII. d.

The Os Sacrum serves as the common base and support of the Trunk of the Body, guards the Nerves issuing from the under end of the Spinal Marrow, defends the back-part of the Pelvis, and gives origin to Muscles moving the Trunk and Thigh.

In the Fœtus, the Os Sacrum is composed of five distinct Vertebræ, which have Intervertebral Substances similar to those of the True Vertebræ. Tab. XXVII.

At this time, each of the Vertebræ of the Os Sacrum, as well as of the True Vertebræ, consists of a Body and two lateral parts, which are joined together by Cartilages.

OS COCCYGIS.

The Os Coccygis, or *Rump-bone*, forming an Appendage to the under end of the Os Sacrum.

The *Situation* of this Bone at the under end of the Os Sacrum. Tab. XX.

Its *Figure*, *broad and flat above*, and *tapering below*, *convex behind*, and forming a *curve forwards*, to defend it from injury when a person is in a sitting posture. Tab. XIX. Tab. with Viscera of Pelvis in Vol. II. and III.

The *four pieces* of which it is composed in young Subjects. Tab. XXVII.

The Bone is considered by some Authors as being formed

formed of *three* pieces; and then the Os Sacrum is said to have *six* pieces.

The *first* or *uppermost* piece the *largest*, with Shoulders reaching farther than the end of the Os Sacrum. This is regarded by some as a proper distinction between the Os Coccygis and Os Sacrum. Tab. XIX.

From the back-part of the Shoulders, *two Cornua* frequently ascend to join the forked Spinous Processes at the end of the Os Sacrum, and form a passage for the transmission of the last Pair of Spinal Nerves. Tab. XIX.

The *three lower Bones* of the Os Coccygis becoming gradually smaller, the *fourth* terminating in a *rough point*. Tab. XX.

Cartilage is interposed between the different pieces of this Bone in Young Subjects, Tab. XXXII. Fig. 15. joining them together, after the manner of the Vertebrae; allowing motion upon each other forwards and backwards, but chiefly between the first and second pieces; and a greater degree of motion there in the Female than in the Male.

In advanced life, but earlier in Men than in Women, the pieces grow together so as to admit of no motion; but this takes place much later between the first and second, than between the other pieces.

The *Substance*, like that of the Os Sacrum, is *spongy*, but differs from it in having no passage for the Spinal Marrow, nor Holes for Spinal Nerves.

The *Connection* of this Bone, in Young Subjects, to the Os Sacrum, by *Cartilage*.—In Old People, by an *union* of Substance.

The Surface of the Bone is covered by a strong Ligament, which adds to its strength; and its sides give rise to numerous Muscular Fibres, which, while they derive their origin from it, serve at the same time to protect it.

The Os Coccygis sustains the Intestinum Rectum, contracts the Inferior Opening of the Pelvis, assists in supporting the Rectum, Bladder, and Uterus.

In the Fœtus, the Os Coccygis is almost entirely composed of Cartilage.

P E L V I S.

THE PELVIS, situated at the lower part of the Trunk, and formed by the Os Sacrum, Os Coccygis, and two Offa Innominata.

OS INNOMINATUM.

The *Situation* of the Os INNOMINATUM, in the fore-part and side of the Pelvis, and in the under and lateral part of the Abdomen. Tab. XVI. c, U.

The *Division* of the Bone, in Children, into *Os Ilium*, *Os Ischium*, and *Os Pubis*. Tab. XXXII. Fig. 15. f, g. Tab. XXVII.

In the Adult, the three Bones are *ossified together*, but retain their original names.

OS ILIUM.

The *Os Ilium*, forming the upper part of the Os Innominatum, and spreading out, to assist in supporting the contents of the Abdomen. Tab. XVI. L.

The *Dorsum*, or *outer convex Surface* of the Bone, raised in some parts and depressed in others, where the Glutei Muscles, or Extensors of the Thigh, have their origin. Tab. XVII.

The *Spine*, or upper semicircular edge of the Bone, for the attachment of the Oblique and Transverse Abdominal Muscles. Tab. XVI. M, N.

The *anterior-superior Spinous Process*, or anterior ex-

trinity of the Spine, for the attachment of the Sartorius and POUPART'S Ligament. Tab. VI. N.

The *anterior-inferior Spinous Process*, a little below the former, for the attachment of the Rectus Femoris. Tab. XVI. O.

The *two posterior Spinous Processes*, at the back-part of the Spine, less considerable than the two anterior; partly for the Origin of Muscles, but chiefly for the attachment of Ligaments which belong to the Joint between this Bone and the Os Sacrum. Tab. XXXI. Fig. 17. v, d.

The *Notch* of the Os Ilium under the posterior-inferior Spinous Process, for the passage of the Pyriform Muscle, the Sciatic Nerve, and Blood-vessels. Tab. XIX.

The *Venter*, or *inner concave Surface* of the Bone, for the attachment of one of the Flexors of the Thigh, termed *Iliacus Internus*, and the support of a portion of the Intestinum Ilium and Colon. Tab. XX. F.

A *Passage* in the Venter for the principal Medullary Vessels of the Bone. Tab. XVI. under b, right side.

A *Depression* at the inside of the anterior-inferior Spinous Process, where the Flexor Muscles of the Thigh, and the anterior Crural Vessels and Nerves pass. Tab. XVI. O, P.

The *Linea Innominata* at the under part of the Venter of the Bone, forming the lateral portion of the Brim of the Pelvis, and the line of division between the Pelvis and Abdomen. Tab. XVI. a.

The

The *inner and back part* of the Bone, very *irregular*, for the origin of some of the large Muscles of the Back, for the attachment of Ligaments which go to the Os Sacrum, and for the firm connection which subsists between this Bone and the Os Sacrum.

The *under, fore, and outer part* of the Bone, forming the upper and back part of the Acetabulum, or Cavity for the articulation of the Thigh-bone. Tab XVI. under R.

OS ISCHIUM.

The *Situation* of the Os Ischium in the lowest part of the Pelvis. Tab. XVI. g, U, T.

Its *Figure, irregular*; its *Size*, next to that of the Os Ilium.

The *upper thick Part* of the Bone, forming the under part of the Acetabulum. Tab. XVI. g.

The *Spinous Process* sent back from the upper part of the Bone, for the attachment of Muscles,—and of the Superior Sacro-sciatic Ligament, which completes the Notch of the Os Ilium into an Iliac Foramen. Tab. XVI. U, right side.

The *Cervix* placed under the Spinous Process, and covered with Cartilage where the Tendon of the Obturator Internus plays. Tab. XIX. Fig. 1. i.

The *Tuberosity, or Tuber Ischii*, which is covered with Cartilage that is separated by macerating the Bone. Tab. XVI. U, left side. Tab. XVII.

The *Upper Part* of the Tuber placed obliquely, and giving attachment to the Geminus Inferior, to the under Sacro-sciatic Ligament, and to the great Flexor Muscles of the Thigh. The thinner and more scabrous part of the Tuber, which has a curved direction, gives attachment to the Crus Penis in the Male, to the Crus Clitoridis in the Female, and to part of the Adductor Muscles of the Thigh. Tab. XVI. I.

OS PUBIS.

The *Situation* of this Bone at the upper and fore part of the Pelvis. Tab. XVI. V, W, X.

Its *Size*, the least of the three portions of the Os Innominatum.

The *thick and strongest part* of the Bone, forming the upper and fore side of the Acetabulum. Tab. XVI.

The *Upper part* of the Bone becoming smaller where it is flattened above, and rendered smooth by the passage of the Flexor Muscles of the Thigh, and of the anterior Crural Vessels and Nerves. Tab. XX. N.

The upper and inner part of the Bone increasing in size, and forming the rough *Crest, or Angle*, where the Rectus and Pyramidalis, and the inner end of POU-PART'S Ligament, are attached. Tab. XVI. W.

A *Ridge, or Spine*, extended from the outer and fore part of the Crest, along the upper and inner edge of the Bone, to form, with a similar Ridge of the Os Ilium, the Brim or upper opening of the Pelvis. Tab. XVI. a.

That part of the Brim of the Pelvis which is formed by this Bone is sometimes so sharp, as to injure the parts which lie immediately contiguous to it.

Another *Ridge*, from the former, extending downwards and backwards towards the breach in the fore-part of the Acetabulum. Tab. XVI. under V.

A *Cavity* below these Ridges, for the origin of the Pectineus. Tab. XVI. outside of c.

At the upper and outer part of the Foramen Thyroideum, the Bone having a *twisted appearance*, and a *Notch* which is formed into a Hole in the Subject, by the addition of the Obturator Ligament, for the passage of the Obturator Vessels and Nerves. Tab. XVI. e.

The *inner end* of the Bone, *rough and unequal*, but covered with a Ligamentous Cartilage, which, in fresh Bones, joins the two Offa Pubis so firmly together, as to prevent them from moving upon each other. Tab. XVI. c.

The inner part of the Bone is broad, and depressed before, where it gives origin to part of the Adductor Muscles of the Thigh. Tab. XVI. between W and. X.

The inner part of the Bone becoming narrower, and ending in the *Crus*, which goes downwards to join the Crus of the Os Ischium, and form, along with that Crus, one side of the Arch of the Pubis. Tab. XVI. X, d.

The *Foramen Thyroideum*, formed by the Os Pubis and Os Ischium, and in the Subject, filled by a Membranous Ligament, excepting at the Notch above mentioned, which gives origin to a large share of the Obturator Muscles. Tab. XIX. O.

The *Acetabulum, or Cavity*, (compared to a Vinegar-measure used by the Ancients), placed farther out than the Foramen Thyroideum, and formed by the three pieces which compose the Os Innominatum, in such a manner, that the Os Ilium constitutes near two-fifths, the Os Ischium more than two-fifths, and the Os Pubis one fifth of that Cavity. Tab. XVI. g.

The *Brim* of the Acetabulum very deep, especially behind, and made still deeper in the Subject, by being tipped with a Cartilaginous Ligament.

Round the outer Edge of the Brim, the Bone *rough*, where the Capsular Ligament of the Joint is fixed. Tab. XVI.

A *Breach* in the inner and fore part of the Acetabulum, which, in the Subject, has a strong Ligament stretched from one end of that Notch to the other, but leaving a Hole behind for containing part of the Substance called *Gland of the Joint*. Tab. XVI.

The *Cavity* of the Acetabulum lined with Cartilage, excepting at its under, inner, and fore part, where there is a rough Surface for containing the greater part of the Substance mentioned above. Tab. LIV. Fig. 3. 6.

The *Brim or upper opening* of the Pelvis, approaching in the Male to a Circular, and in the Female to an oval form. Tab. XX. C, C.

The *Inferior Opening* is large in the Skeleton, and in the

the Subject in a great measure is filled up by Ligaments and Muscles, which support and protect the contained parts, and leave only the passages from the Bladder of Urine and Rectum in the Male, and, together with these, the passage from the Uterus, in the Female.

The *Ossa Innominata*, joined behind to the *Os Sacrum* by a *thin Cartilage* and by *strong Ligaments*, so as to have no motion; the Joint obtaining the name of *Posterior Symphysis*. Tab. XX. A.

The *Connection* of these Bones to each other anteriorly, by a *Ligamentous Cartilage* and *Ligaments*, which also prevent motion here. This *Connection* is termed *Symphysis*, or *Anterior Symphysis Pubis*. Tab. XX. B.

The Substance of the *Iliac* part of the *Os Innominatum*, is cellular, with a thin external *Table*, which, in some Old People, is so much affected by muscular action about its middle, as to become transparent. The other two Portions of the *Os Innominatum* are cellular, as in other

flat Bones, but some parts of the *External Table* are of considerable thickness.

USE OF THE PELVIS.—It constitutes the Basis of the Trunk, and forms Sockets for the Thigh-bones to move in. It contains the Bladder of Urine and the Rectum in the Male, and, together with these, the Uterus in the Female. It gives origin to the Muscles which extend the Trunk, and insertion to those which bend the body. It sends off the principal part of the Muscles which move the Thigh, and gives passage and protection to Blood-vessels, and to some of the largest Nerves of the Body.

In the Fœtus, the Spine of the *Os Ilium*, and that part of the Bone which belongs to the *Acetabulum*, are Cartilaginous. The Spinous Process, the Tuberosity, and Crus of the *Os Ischium*, the Crus of the *Os Pubis*, and that portion of it which forms the *Acetabulum*, are also, at this period, in a Cartilaginous state.

TABLE

T A B L E XVIII.

Gives different VIEWS of the TRUE VERTEBRÆ.

FIG. 1.

A View of the Inferior Surface of the ATLAS, or FIRST VERTEBRA of the NECK.

- a*, The anterior part of the atlas.
- b, b*, The inferior oblique processes.
- c*, A muscular impression on the posterior part of the bone.
- d, d*, Transverse processes, which terminate in tuberosities.
- e, e*, The inferior orifices of the oblique holes.
- f, f*, Inferior notches for the passage of the spinal nerves.
- g*, The large vertebral hole, which forms part of the spinal canal.

FIG. 2.

Represents the ATLAS, seen from its Upper and Back Part.

- a*, The small articular cavity, which receives the odontoid process.
- b, b*, The superior oblique processes, which receive the occipital condyles.
- c, c*, Protuberances below the superior oblique processes, to which the transverse ligaments are fixed.
- d, d*, The posterior fossæ, where the vertebral arteries are reflected in their ascent to the cranium.
- e, e*, The oblique holes at the roots of the transverse processes, for the passage of the vertebral arteries.
- f, f*, Extremities of the transverse processes, each in form of a tuberosity.
- g*, A muscular print on the posterior part of the bone.
- h*, The large vertebral hole.

FIG. 3.

The VERTEBRA DENTATA, viewed superiorly, and a little posteriorly.

- a*, The odontoid process of the second vertebra.
- b, b*, The superior oblique processes.
- c, c*, The transverse processes.
- d, d*, The superior notches of this vertebra for the passage of the spinal nerves.
- e, e*, A portion of the inferior oblique processes.

- f, f*, The extremities of the spinous processes, of a forked shape.
- g, g*, Muscular prints on the two sides of the spinous process.
- h*, The large vertebral hole.

FIG. 4.

The Anterior and Inferior Part of the VERTEBRA DENTATA.

- a*, The upper part of the processus dentatus.
- b*, The anterior and middle surface of the second vertebra.
- c*, The inferior surface, somewhat convex.
- d, d*, The anterior margin of the superior oblique processes.
- e, e*, The extremities of the transverse processes.
- f, f*, The oblique holes at the roots of the transverse processes, for the passage of the vertebral arteries.
- g, g*, The inferior notches of the vertebra for the passage of the spinal nerves.
- h, h*, The inferior oblique processes.
- i, i*, The bifurcation of the spinous process.
- k*, A furrow on the inner surface of the spinous process.
- l*, The large vertebral hole.

FIG. 5.

The Connection of the two First VERTEBRÆ of the NECK with each other, seen anteriorly.

- a*, The eminence, or anterior print of the atlas.
- b, b*, The two superior fossæ of the atlas.
- c, c*, The anterior edge of the superior oblique processes.
- d, d*, The extremities of the transverse processes.
- e, e*, The anterior edge of the inferior oblique processes.
- f, f*, The inferior fossæ, or hollows of the atlas.
- g*, The extremity of the tooth-like process of the second vertebra.
- h*, The root of the tooth-like process.
- i*, A small eminence on the middle of the body of the second vertebra.
- k, k*, Prints upon the lateral parts of the body of the bone.
- l*, The convexity of the inferior part of the body.
- m, m*, The anterior margin of the superior oblique processes.

n, n, The

- n, n*, The inferior oblique processes.
o, o, The transverse processes.
p, p, The inferior hollows of the vertebra.

FIG. 6.

The Connection of the two First VERTEBRÆ of the NECK, seen posteriorly and superiorly.

- a, a*, Cavities of the superior oblique processes of the atlas.
b, b, The ligamentous protuberances at the under and inner part of the superior oblique processes.
c, c, The posterior edge of the inferior oblique processes of the atlas.
d, d, The posterior fossæ of the atlas, through which the vertebral arteries and tenth pair of nerves pass.
e, e, The holes of the transverse processes of the atlas.
f, f, The extremities of the transverse processes of the atlas.
g, The eminence of the atlas, in form of a spinous process.
h, The superior extremity of the tooth-like process of the second vertebra.
i, i, Ligamentous impressions upon the superior extremity of that process.
k, The neck of the tooth-like process.
l, l, The posterior edge of the superior oblique processes of the second vertebra.
m, The middle of the large vertebral hole.
n, n, The posterior orifices of the passages at the roots of the transverse processes of the second vertebra.
o, o, The extremities of the transverse processes of that vertebra.
p, p, The posterior edge of the inferior oblique processes of the second vertebra.
q, A crest projecting from the upper part of the spinous process of the second vertebra.
r, r, The extremities of the spinous process of the second vertebra, on which are muscular prints.

FIG. 7.

A View of the Upper and Back Part of one of the First DORSAL VERTEBRÆ.

- a*, The superior surface of the body of the first dorsal vertebra, which is somewhat triangular.
b, b, The superior oblique processes.
c, Part of the body of the bone, which assists in forming the vertebral hole.
d, The thin sharp edge of the vertebral hole.
e, e, The posterior part of the transverse processes.
f, f, The under edge of the inferior oblique processes.
g, g, The posterior fossæ of this bone.
h, The ridge of the spinous process.
i, The small extremity of the spinous process.

FIG. 8.

A View of the Under and Fore Part of the VERTEBRA represented in the preceding Figure.

- a*, The anterior part of the body of the bone.
b, The inferior surface of the body.
c, c, The superior notches for the passage of the spinal nerves.
d, d, The inferior notches for the passage of the spinal nerves.
e, e, The transverse processes.
f, f, The small articular cavities, which receive the tubercles of the ribs.
g, g, The inferior oblique processes.
h, The large vertebral hole.
i, The interior fossa, or groove of the spinous process of the first dorsal vertebra.
k, The inferior extremity of the spinous process.

FIG. 9.

The Inferior Surface of the Third VERTEBRA of the LOINS.

- a*, The middle of the inferior surface of the third lumbar vertebra.
b, b, The ossicous lamina which borders the whole circumference of the inferior surface.
c, c, The inferior notches of this vertebra.
d, d, The transverse processes.
e, e, The inferior oblique processes.
f, f, The superior notches.
g, The large vertebral hole.
h, A small groove on the inner side of the spinous process.
i, The rounded extremity of the spinous process.

FIG. 10.

The VERTEBRA represented in the preceding Figure, seen from its Upper and Back Part.

- a*, The centre of the body of the vertebra, which is very spongy.
b, b, The small ossicous lamina, which surrounds the spongy surface.
c, A portion of the body of this vertebra, which forms part of the vertebral hole.
d, d, The superior fossæ of this vertebra.
e, e, The extremities of the transverse processes.
f, f, The superior oblique processes.
g, g, The inferior fossæ of this vertebra.
h, h, The posterior fossæ.
i, i, The inferior oblique processes.
k, The large vertebral hole.
l, The exterior ridge of the spinous process.
m, The rounded extremity of the spinous process.

TAB. 19.

Fig. 1.




Fig. 2.




Fig. 6.



Fig. 5.



A. Fyfe fecit.

T A B L E XIX.

VIEWS of the SEPARATE BONES of the PELVIS.

FIG. 1.

The Internal or Anterior Surface of the Os SACRUM, turned a little towards the Left Side.

- A, THE upper part of the os sacrum, which receives the body of the last lumbar vertebra.
- B, The osseous lamina which surrounds this surface.
- C, C, A portion of the oblique processes.
- D, D, The superior notches, for the passage of the twenty-fourth pair of spinal nerves.
- E, E, The large lateral eminences.
- F, F, The pieces of which the bone is originally composed.
- G, G, The transverse lines which indicate the union of these pieces.
- H, H, The oblique passages on each side, for the transmission of the sacral nerves.
- I, A portion of the surface of the os sacrum, by which it is articulated with the os ilium.
- K, K, The inferior notches, where the last pair of spinal nerves pass out.
- L, The point of the bone which is joined to the Os Coccygis.

FIG. 2.

The Posterior Surface of the Os SACRUM, turned a little towards the Left Side.

- A, The upper surface of the os sacrum.
- B, B, Its superior oblique processes.
- C, C, Its superior notches.
- D, The beginning of the spinal canal of this bone.
- E, E, The spinous processes.
- F, F, The appendices, or cornua.
- G, G, The posterior foramina.
- H, The termination of the spinal canal.
- I, I, The inferior notches.
- K, The point of the bone which is united with the os coccygis.
- L, L, Ligamentous and muscular impressions, rendering the back-part of the bone very unequal.

M, A portion of the articular surface, by which this bone is united with the os ilium.

FIG. 3. *The Internal Surface ; and, FIG. 4. The External Surface, of the Os COCCYGIS.*

- A, A, A, The three pieces of which the bone shewn in this figure is composed.
- B, The great notch of this bone.
- C, The cornua.
- D, D, The lateral notches.
- E, E, The lateral processes.
- F, The point, or inferior extremity.

FIG. 5.

The Internal Surface of the LEFT Os INNOMINATUM.

- A, The cavity or venter of the os ilium.
- B, The orifice of the internal iliac canal.
- C, C, The spine or crest of this bone.
- D, The superior-anterior spinous process.
- E, The inferior-anterior spinous process.
- F, F, The posterior spinous processes.
- G, The surface by which the os ilium is articulated with the os sacrum.
- H, H, An irregular surface which also belongs to this joint.
- I, The anterior iliac notch.
- H, The posterior iliac notch.
- L, The ischiatic, or, more properly, the great iliac notch.
- M, The great sinuosity, where the internal iliac muscle passes out of the abdomen.
- N, The ridge of the os ilium, which forms a share of the brim of the pelvis.
- O, The body of the os ischium.
- P, The spinous process of this bone.
- Q, Part of the tuberosity of this bone.
- R, The inner part of the sinuosity which is between the spinous process and tuberosity.

S, The

- S, The notch of the os ischium, which assists in forming the foramen ovale.
 T, The crus of the os ischium.
 U, The eminence which marks the union of the superior branch of the os pubis with the os ilium.
 V, The spine of the os pubis.
 W, The inner surface of the crest of the os pubis.
 X, The inner surface of the superior branch of the os pubis.
 Y, The inferior notch of this branch.
 Z, The inner surface of the body of the os pubis.
a, The crus of this bone.
b, The cartilaginous surface which unites itself with the os pubis of the opposite side.
c, The foramen ovale.
d, The inferior notch of the os pubis, which assists in forming the foramen ovale.


FIG. 6.

The External Surface of the Left Os INNOMINATUM.

- A, The dorsum of the os ilium, raised in some parts, and depressed in others.
 B, B, The crest of this bone.
 C, The superior-anterior, and,
 D, The inferior anterior spinous process.
 E, E, The posterior spinous process.
 F, The anterior notch.
- G, The posterior notch, and,
 H, The great notch of this bone.
 I, Eminences and cavities at the upper part of the acetabulum.
 K, The brim of the acetabulum, tipped with cartilage.
 L, The bottom of this cavity encrusted with cartilage.
 M, A rough surface in the acetabulum, where the substance termed *gland of the joint* is lodged.
 Surrounding the acetabulum, and upon the dorsum of the os ilium, foramina appear, which are the passages of blood-vessels.
 N, The spinous process of the os ischium.
 O, The tuberosity of this bone.
 P, The notch between the spinous process and tuberosity.
 Q, The cervix of the bone.
 R, The breach in the acetabulum, which in the subject has a strong ligament connected to it.
 S, The crus of the os ischium.
 T, The notch which assists in forming the foramen ovale.
 U, The outer end of the superior branch of the os pubis.
 V, The middle of this bone.
 W, The crest of this bone.
 X, The notch at the under part of the branch of this bone.
 Y, The body of the bone.
 Z, The crus of this bone.
a, The notch which assists in forming the foramen ovale.
b, The part where the one os pubis joins the other.
c, The foramen ovale.

TAB. 20.

Fig 4.



T A B L E XX.

A VIEW of the FEMALE PELVIS, from the Upper and Fore Part.

-
- | | |
|--|---|
| <p>A, The connection of the os ilium with the os sacrum.
 B, The symphysis pubis.
 C, &c. The brim of the pelvis.
 D, Articulation of the head of the os femoris with the acetabulum of the os innominatum.
 E, The arch formed by the crura of the ossa pubis.
 F, The cavity of the os ilium.
 G, The spine or arch of the os ilium.
 H, The superior-anterior spinous process.
 I, Ligaments passing between the spine of the ilium, to the transverse process of the last lumbar vertebra.
 K, The tuberosity of the os ischium.
 L, The crus of that bone.
 M, The posterior part, forming a share of the acetabulum.</p> | <p>N, The back part of the os pubis, forming a portion of the acetabulum.
 O, The angle, or crest.
 P, The crus of the os pubis.
 Q, The last lumbar vertebra.
 R, R, The os sacrum; the transverse lines marking its original pieces, with the four pairs of holes for the transmission of nerves.
 S, S, The four pieces composing the os coccygis, with a pair of holes between it and the os sacrum.
 T, T, The cavity of the pelvis.
 U, The foramen thyroideum.
 V, The cervix of the os femoris.
 W, The trochanter major.</p> |
|--|---|

THORAX.

THE *Thorax*, formed of the Sternum before, of the Ribs on each side, and of the Dorsal Vertebrae behind. Tab. XV.

The general *Figure* of the Thorax approaching that of a *Cone*, but left open above for the passages to the Lungs and Stomach, and for the great Blood-vessels.

The *Lower Part* of the Thorax *slanting*; the *fore-part* being considerably shorter than it is behind.

The *Under Margin* on each side, forming a curved Line; the convex side of which is turned downwards.

The under end of the Thorax, occupied, in the Subject, by the *Diaphragm*, which forms a Partition between it and the Abdomen. Tab. XLVIII. Fig. 2.

COSTÆ.

The *Situation* of the Costæ, or Ribs, *slanting* downwards with respect to the Spine. Tab. XV.

Their *Number*, commonly *twelve* on each side, though sometimes thirteen, and at other times only eleven; the number of the Dorsal Vertebrae always corresponding with that of the Ribs.

Their *Figure*, *convex* externally, by which their strength is increased; and *concave* and *smooth* internally, with their flat sides turned towards the lungs which they protect.

The *Head* of each Rib formed into a *Ridge* and *two hollow Surfaces* covered with Cartilage, to be articulated with the bodies of two Vertebrae and their intermediate Cartilage. Tab. XXXI. Fig. 14. a.

Round the Head, the Bone *spongy*, for the attachment of the Capsular Ligament of the Joint.

The *Tubercle* of the Rib, at a little distance from its Head, with a flat Cartilaginous Surface and irregular Edge, to be articulated to the Transverse Process of the undermost of the two Vertebrae, to which the head of the Rib is joined. Tab. XXXI. Fig. 14. b.

The *Cervix* of the Rib, between its Head and Tubercle, of a roundish form. Tab. XXI. Fig. 3. a.

Another *small Tubercle* in most of the Ribs, at the outer side of the former, for the attachment of Ligaments which fix the Ribs to each other and to the Transverse Processes, Tab. XXXI. Fig. 14. d.; and also for the insertion of the outer Slips of the Longissimus Dorsi.

Beyond the Tubercle, the Rib rendered *flat* by the Sacro-lumbalis.

The *Angle* of the Ribs to which the Sacro-lumbalis is fixed, where the Bones are about to bend to form the lateral part of the Thorax. Tab. XXXI. Fig. 14. e.

The Rib becoming *broader* and *flatter* at the lateral part of the Thorax, and the flat Surface opposed to the Lungs.

The *upper Edge* of the Rib, *round* where the Intercostales are fixed.

The *under Edge*, *sharp* where the Intercostalis Externus is fixed.

A *Fossa* at the inside of the under Edge, for lodging the Intercostal Vessels and Nerves.

The *Fossa* *awanting* towards the extremities of the Ribs; the Vessels not being in contact with them behind, and too small to impress them anteriorly.

An *Oval Pit* in the anterior extremity of each Rib, for receiving the Cartilage which runs from it to the Sternum. Tab. XXXI. Fig. 14. b.

The *Cartilages* of the Ribs, placed between them and the Sternum. Tab. XV. H.

The Cartilages, like the Ribs, *flat* on their outer and inner Surfaces, and *smooth* where they are opposed to the Lungs.

The Cartilage of each Rib, forming, with the Rib itself, a *Curve*, the concave part upwards.

And with the Sternum, an *obtuse Angle* above, and an *acute one* below.

The Cartilages yield to the motions of the Ribs, and enable them to return to their former position, when the Muscles cease to act. The Cartilages of the Ribs, in Old People, are frequently ossified.

The Ribs are *connected behind* to the Vertebrae, by a double articulation, and *before* to the Sternum by the Cartilages, or by the Cartilages to each other, in such a manner as to allow motion upwards and downwards, though only a small degree in any single Rib, and that towards its middle; but no motion in any other direction. Tab. XXXI. Fig. 17.

The *first Rib* the *most crooked*; from this downwards the Ribs becoming gradually straighter. Tab. XV. G, G, &c.

The *uppermost Ribs* approaching nearer to the horizontal situation; their obliquity, with respect to the Spine, increasing as they descend, and their anterior extremities becoming more distant from each other. Tab. XV.

The *Cartilages* of the Ribs, like the Ribs themselves, becoming gradually longer, but, contrary to what happens in the Ribs, approaching nearer to each other in their descent. Tab. XV. H, H, &c.

The *length* of the Rib, *increasing* from the first to the seventh, and then *decreasing* to the twelfth Rib. Tab. XV.

The *distance* between the heads of the Ribs and their Angles, increasing to the ninth Rib, corresponding with the

the breadth of the Sacro-lumbalis which covers them. Tab. XLII.

The Ribs divided into *True* and *False*.

The *True Ribs*,—the seven uppermost, having their Cartilages joined to the Sternum, and opposed to the Heart and Lungs, from which they are termed the *True Custodes*, or *Guards of Life*. Tab. XV.

The *False Ribs*,—the five inferior, not reaching the Sternum. Tab. XV.

The *Cartilages* of the False Ribs, *shorter* as they descend, and more flexible than those of the True Ribs. Tab. XV. I, I, &c.

The *posterior Extremity* of the first Rib, *articulated only with the first Vertebra*. Tab. XV. K.

A *flat Surface* upon the upper part of the first Rib, where the *Subclavian Vessels* pass over it to the arm. Tab. XV. N.

The *Fossa* for the Intercostal Vessels wanting at the edge of this Rib.

The *Cartilages* of the two under True Ribs, and three upper False Ribs, *joined* to each other by an union of Substance. Tab. XV.

The *Head* of the eleventh Rib, *having no Tubercle* for articulation; being only loosely joined to the Transverse Process.

The twelfth Rib *much shorter* than the rest. Its Head is only joined to the twelfth Vertebra of the Back. It has no Tubercle, nor articulation with the Transverse Process; neither has it any Fossa at its under edge, the Vessels running some way below it. Tab. XV.

The *Anterior Extremities* of the Eleventh and twelfth Ribs, not joined to each other, nor to any other Rib, but lying loose among the Muscles;—hence sometimes named *Floating Ribs*. Tab. XV.

The Substance of the Ribs, like that of the Vertebrae, is cellular, and only covered with a thin external Plate, which becomes somewhat thicker towards the Vertebrae.

In the Fœtus, the Heads and Tubercles of the Ribs have Cartilages, part of which become thin Epiphyses. After Birth, the Bodies of the Ribs encroach gradually on the Cartilages; hence the Cartilages of the Ribs are proportionally shorter in Adults than Children.

The Ribs give form to the Thorax, cover and defend the Heart and Lungs, and assist the latter in performing respiration.

STERNUM.

The *Situation* of the Sternum in the fore-part of the Thorax. Tab. XXIX. Fig. 10. V.

Three Pieces composing the Sternum, in a person of middle age, and these joined together by Cartilage. Tab. XV. O, P, Q.

The different pieces of this Bone are frequently found *ossified together* in Old People.

The Sternum, *thick and broad above*, and *thin and narrow below*. Tab. XV.

The *outer Surface flat*. Tab. XV.

The *inner Surface slightly hollowed*, to enlarge the Cavity of the Thorax.

Pits upon each edge of the Sternum, to receive the Cartilaginous ends of the seven True Ribs. Tab. XXI. Fig. 2. D—L. Tab. XV.

The *Pits* at a considerable distance from each other above, but becoming gradually nearer as they descend. Tab. XXI.

The *Cancelli* of the Sternum, covered only by a thin external plate; but this rendered stronger by a Tendinous Membrane investing it in the recent state. Tab. LII. Fig. 10.

The *upper piece* of the Sternum, of a somewhat *triangular* figure, compared to that of a heart as painted on playing cards, but cut across below. Tab. XXI. Fig. 2.

The *upper and back part hollowed*, to make way for the Trachea. Tab. XXI. Fig. 2. C.

The *upper Corners*, *thicker and stronger* than the rest of the Bone, with a *Cavity* in each, lined with Cartilage for receiving the ends of the Collar Bones. Tab. XXI. Fig. 2. B, B.

Under these Cavities, the Bone becoming *thinner*, and having a *Pit* upon each side, for receiving the Cartilage of the first Rib. Tab. XXI. Fig. 2. D, D.

Part of the *Pit* in the *under Corner* of the first piece, for the Cartilage of the second Rib. Tab. XXI. Fig. 2. F, F.

The *second piece* of the Sternum, of an *oblong* form, but a little broader below than above, and considerably longer than the former. Tab. XXI. Fig. 2. E, E.

Complete Pits upon the edge of this piece, for the Cartilages of the third, fourth, fifth, and sixth Ribs, and part of the *Pits* for those of the second and seventh. Tab. XXI. Fig. 2. F—L. Tab. XV.

Lines extending between the Pits, denoting the original marks of division of this piece. Tab. XXI. Fig. 2.

The *Connection* of the second piece of the Sternum to the first by Cartilage, which, in the earlier period of life, allows some yielding, but this becoming gradually less as the person advances in life. Tab. XXI. Fig. 2. F, F.

The *third piece* of the Sternum, *cartilaginous* in a young Subject, and pointed like a broad-sword, hence termed *Cartilago Ensiformis*. Tab. XV. Tab. XXX.

The Adult has this piece commonly *ossified* in the middle, and *cartilaginous* at the edges. Tab. XXIX. Fig. 10. NO. 3. 4.

The *Size* of this piece *much less* than that of the other two. Tab. XV.

Only *one half* of the Pit, for the Cartilage of the seventh Rib, formed in the side of this piece. Tab. I.


The *Variations* of the Cartilago *Ensiformis* are considerable in different Subjects;—for, instead of the common form, it is sometimes narrow like the point of a small sword, or turned obliquely to one side, or forwards, or backwards, or forked at the point, or perforated in the middle.

The Sternum is *joined* by Cartilage to the seven upper or True Ribs, and by an inter-articular Cartilage to the anterior ends of the Clavicles. Tab. XV.

In the Fœtus, this Bone is composed of seven or eight pieces, but the number of these varies in different Subjects. By degrees the pieces unite, till at length they form the three Bones already described.

The Sternum gives origin to several Muscles, defends the Heart and Lungs, assists in the formation of the Thorax, sustains the Mediastinum, is a medium of attachment to the Ribs, and serves as a Fulcrum or Point on which the Clavicles roll.

TABLE



T A B L E XXI.

Represents the **ATLAS, STERNUM, FIRST RIB, CLAVICLE, and SCAPULA.**

FIG. 1.

The Superior Surface of the ATLAS, or FIRST VERTEBRA.

- A, THE spinal hole.
- B, The articular notch which receives the processus dentatus of the second vertebra of the neck.
- b, The direction of the ligament which confines this process in its place.
- C, The anterior part of the atlas.
- D, D, The superior oblique processes.
- d, d, Placed behind prominences, to which the lateral ligaments of the head are fixed.
- E, E, The transverse processes.
- e, e, The holes in the transverse processes for the vertebral blood-vessels.
- F, The spinous process.
- f, f, The posterior depressions, where the vertebral arteries are reflected in their way to the cranium.

FIG. 2.

The External Surface of the Sternum.

- A, The upper triangular piece of the sternum.
- a, a, Impressions made by the pectoralis major.
- B, B, Notches which receive the inner ends of the clavicles.
- C, An excavation where the trachea passes into the thorax.
- D, D, The lateral parts which receive the cartilages of the first pair of ribs.
- E, E, The middle and longest part of the sternum.
- e, e, Transverse lines pointing out the union of the different pieces of which this bone is originally composed.
- F, F, The impressions of the cartilages of the second pair of ribs, and the connection between the first and second pieces of the sternum.
- G, H, I, K, The lateral cavities, or impressions of the cartilages of the third, fourth, fifth, and sixth ribs.
- L, L, Impressions of the cartilages of the seventh pair of ribs.
- M, The cartilago ensiformis.

FIG. 3.

The Outer Surface of the Second TRUE RIB of the RIGHT SIDE.

- A, The head.
- a, The cervix.
- B, The tubercle.
- C, The angle.
- D, The upper and inner edge.

FIG. 4.

A View of the Back-Part, or DORSUM of the SCAPULA of the RIGHT SIDE.

- A, The spine.
- B, The acromion.
- C, C, The superior spinous fossa.
- c, The superior angle.
- D, D, The inferior spinous fossa.
- E, E, The base or posterior costa.
- F, The back-part of the posterior costa.
- G, G, The inferior costa.
- H, The glenoid cavity.
- b, b, b, The brim of the glenoid cavity.
- I, I, The cervix of the scapula.
- K, The coracoid process.
- L, The inferior angle.

FIG. 5.

Under Side of the CLAVICLE.

- A, The middle or body of the clavicle, with a depression running along it, for lodging part of the subclavian muscle.
- B, The inner or sternal extremity.
- C, The inferior angle of this extremity.
- D, The superior angle.
- E, The part where the upper edge of the pectoralis major rises.
- F, The spine of the clavicle.
- G, The ligamentous and muscular impressions of the external extremity.
- H, The orifices of several passages for vessels.
- I, The humeral extremity.

SUPERIOR

SUPERIOR EXTREMITY.

THE *Superior Extremity*, composed of the Bones of the *Shoulder, Arm, and Hand*.

The *Shoulder*, consisting of the *Clavicle* and *Scapula*.

CLAVICLE.

The *Situation* of the *Clavicle*, between the upper part of the *Sternum* and top of the *Scapula*. Tab. XV. Y.

The *Sternal*, or *internal Extremity*, *triangular*, and *larger* than the *Body*, with one of the angles elongated backwards, where it gives origin to a *Ligament* extended between the two *Clavicles*. Tab. XXI. B, C, D. Tab. XV. W.

The *Surface* next the *Sternum*, covered with *Cartilage*, and *irregularly hollowed*, to correspond with the *inter-articular Cartilage*, which, with the *Capfular Ligament* of this *Joint*, allows a small degree of motion in all directions. Tab. II. Fig. 5. B.

The *Body* of the *Bone* next the *Sternum* *bent forwards*, and that next the *Shoulder* *turned back*, the whole resembling an *Italic s*, or a *key* used by the *Ancients*; from which, or from the support it gives the *Shoulder*, its name is derived. Tab. XV.

The upper part of the *Clavicle* next the *Sternum*, *rounded*, and that next the *Scapula*, *flat* where it lies over the *Joint* of the *Humerus*. Tab. XV.

Over the *Bone* in general, rough marks are observed, for the attachment of *Muscles* and *Ligaments*.

The under *Surface* *hollow*, for lodging a portion of the *Subclavius*. Tab. XXI. Fig. 5. A.

In the under *Surface*, one or more small *Canals* leading obliquely outwards, for the passage of the *Medullary Vessels*.

The *External* or *Scapulary Extremity*, tipped with *Cartilage*, to be articulated with the *Acromion* of the *Scapula*. Tab. XV. 10.

Near the back-part of the *Scapulary Extremity*, a *Tubercle*, for the attachment of a strong *Ligament*, which connects this *Bone* to the *Coracoid Process* of the *Scapula*.

The *Substance* of this *Bone* is like that of other long round *Bones*, but the external *Table* is of considerable thickness.

The *Clavicle* supports the *Shoulder* at a proper distance from the *Thorax*, and thereby renders the motions of the *Arm* more extensive. It gives origin to several *Muscles*, and defence to large *Vessels* and *Nerves*.

In a *Fœtus*, the *Clavicle* is completely formed.

SCAPULA.

The *Situation* of the *Scapula*, upon the upper and back part of the *Thorax*, at some distance from the *Ribs*, the interval being filled up by a cushion of *Flesh*. Tab. XVI.

The *shape* of the *Scapula* *triangular*, with one of the angles placed downwards. Tab. XVI.

The *Venter*, or inner *Surface*, *concave*, corresponding with the convexity of the *Ribs*, and marked with *Ridges* and *Depressions* by the *Subscapularis*. Tab. XXIII Fig. 1. A.

The *Dorsum*, or outer *Surface* of the *Scapula*, rendered *convex* in some parts, and *concave* in others, by the action of the *Muscles* which cover it. Tab. XVI. Tab. XIX.

The *body* of the *Scapula* is *remarkably thin*, and, in an old person, *transparent*.

The *Edges* of the *Bone* *thick* and *strong*, and termed *Costæ*.

The superior or *Cervical Costa* the *shortest* of the three, and placed nearly opposite to the second *Rib*. Tab. XXI. Fig. 4. F. Tab. XVI.

A *Semilunar Notch* near the fore-part of the superior *Costa*, for the passage of the superior *Scapulary Vessels* and *Nerves*.

The *inferior* or *anterior Costa*, extending obliquely downwards and backwards, between the third and eighth *Ribs*. Tab. II.

The *posterior Costa*, or *Base* of the *Bone*, placed obliquely with respect to the *Spine*, the upper end being considerably nearer to it than the under. Tab. II.

The *upper part* of the *Base*, above the *Spine*, running obliquely forwards to the upper angle, and giving attachment to the *Levator Scapulæ*. Tab. XXI. Fig. 4.

The *inferior Angle* very acute, and marked by the passage of the *Latissimus Dorsi*, and the origin of the *Teres Major*. Tab. XXI. Fig. 4.

The *superior Angle* approaching a right Angle. Tab. XXI. Fig. 4. c.

The *anterior Angle*, forming the *Cervix* which supports the *head* of the *Bone*. Tab. XXI. Fig. 4. I, I.

The *Glenoid Cavity*, placed on the fore-part of the *Head* of the *Bone*, and lined with *Cartilage* for the articulation of the *Os Humeri*. Tab. XXI. Fig. 4. H.

The *Cartilage* lining this *Cavity* thickest at the edges, by which it is rendered deeper for receiving the *Ball* of the *Os Humeri*.

The *shape* of that *Cavity*, resembling an *Egg* cut longitudinally, with the large end undermost, but so shallow as to receive only a small portion of the *Ball* of the *Os Humeri*, the rest of the *Ball* being contained in the *Capfular Ligament*. Tab. XV. R.

The *Spine* running across the *Bone*, dividing it into a small upper, and large under *Surface*, and giving origin to part of the *Spinati*. Tab. XXI. Fig. 4. A, A.

The *Spine*, *small* at its beginning, and becoming *higher* and *broader* in its course forwards. Tab. XXI. Fig. 4.

A *triangular Space*, between the root of the *Spine* and *Base* of the *Bone*, where part of the *Trapezius* is fixed. Tab. XXXI. Fig. 17. between W and the *Base* of the *Scapula*.

Near

Near the Base of the Spine, a passage for the principal Vessels which supply the Substance of the Bone.

The *Fossa Supra-spinata*, or space above the Spine, for the origin of the *Supra-spinatus*. Tab. XXI. Fig. 4. C, C.

The *Fossa Infra-spinata*, for the origin of the *Infra-spinatus*. Tab. XXI. Fig. 4. D.

The Spine becoming broad and flat, and terminating in a point at its anterior extremity, where it is termed *Acromion*, or top of the Shoulder. Tab. XXI. Fig. 4. B.

The under Surface of the *Acromion*, hollow for the passage of the *Spinati*. Tab. XV. S.

The *Situation* of the *Acromion* over the Joint of the *Humerus*, which, together with the Ligaments, contributes to its protection. Tab. XXXI. Fig. 7. v.

The anterior Edge of the *Acromion*, tipped with *Cartilage* for its articulation with the outer end of the *Clavicle*. Tab. XXIV. A. Fig. 10.

The *Coracoid Process*, arising from the neck of the Bone, and making a curvature forwards, so as to leave a hollow at its root for the passage of the *Subscapularis*. Tab. XXI. Fig. 4. b. Tab. XV. T. V.

The *Point of this Process*, giving origin to Muscles, and to a strong Ligament which passes transversely from its side, to be fixed to the *Acromion*, for the protection of the Joint. Tab. XV.

The inferior *Costa* and *Processes*, being thick and strong, while the Body is so pressed by its own Muscles, especially in Old People, as to become in many parts transparent.

The *Scapula* is joined to the *Clavicle* by Ligaments of such strength, as only to allow a small degree of motion, and that chiefly of a twisting nature; but the *Scapula* is so connected by Muscles to the Head, *Os Hyoides*, Trunk, and Arm, as to have motion upwards, downwards, and to either side, and through the Medium of the *Clavicle*, to be rolled upon the top of the *Sternum*.

In the *Fœtus*, the Base, *Acromion*, *Coracoid Process*, and Head of the *Scapula*, are *Cartilaginous*. The three first are afterwards joined as *Epiphyses*; while the Head, with the *Glenoid Cavity*, is gradually produced from the Body of the Bone.

OS HUMERI.

The *Situation* of the *Os Humeri* at the side of the *Thorax*, and under the *Scapula*. Tab. I.

The *Ball*, or *Head* of the *Os Humeri*, forming a small Segment of a large Sphere, and this covered with *Cartilage*, and placed at the upper, inner, and back-part of the body of the Bone, to correspond with the *Glenoid Cavity* of the *Scapula*. Tab. XXII. Fig. 1. a.

The *Cervix*, or *Neck*, surrounding the edge of the *Ball*, and forming a superficial *Fossa*, where the *Capfular Ligament* is fixed. Tab. XXII. Fig. 2. b, b. Tab. I. c.

Numerous Holes round the upper end of the Bone, for the insertion of the Fibres of the *Capfular Ligament*, and for the passage of *Blood-vessels* into the Bone. Tab. XXII. Fig. 1. f.

A *Fossa*, or long *Groove*, lined with a *Cartilaginous* and *tendinous Crust*, in the upper and fore part of the Bone, for lodging the *Tendon* of the long head of the *Biceps*. Tab. XXII. Fig. 1. e.

The *smaller Tubercle*, placed at the upper and inner side of the above-mentioned *Groove*, for the attachment of the *Subscapularis*. Tab. XXII. Fig. 1. d.

The *larger Tubercle*, opposite to the former, and on the outer side of the *Groove*, for the attachment of the Muscles which cover the *Dorsum* of the *Scapula*. Tab. XXII. Fig. 1. c.

A *Ridge* continued down from each *Tubercle* along the sides of the long *Fossa*, for the insertion of Muscles coming from the *Trunk* of the *Body*, or from the *Scapula*. Tab. XXII. Fig. 1. g, b.

A *Passage* slanting downwards in the fore and inner part of the Bone, near its middle, for the *Medullary Vessels*. Tab. XXII. Fig. 2. f.

The *Bone*, marked at the under end of the *Groove* for lodging the long head of the *Biceps*, by the attachment of the *Deltoides* and other Muscles. Tab. XXII. Fig. 1. under i. Fig. 2. at the outer side of g.

The *Body* of the Bone, round near its upper end; but, as it descends, appearing *twisted*, then *flat*, and increasing in breadth at the lower extremity. Tab. XXII. Fig. 1. 2.

From the *Muscular Prints* on the fore-part of the body of the Bone, a *blunt Ridge* continued to the upper part of the *Trochlea*. Tab. XXII. Fig. 1. k, k, k.

The under and back part of the Bone, rendered *flat* and *smooth*, by the motion of the *Triceps Extensor Cubiti*. Tab. XXII. Fig. 2.

A *large Ridge* at the under and outer, and a *small Ridge* at the under and inner edge of the Bone, for the attachment of strong *Tendinous Fasciæ*, which give origin to part of the Muscles of the *Fore-arm*. Tab. XXII. Fig. 1.

The *Ridges* ending below in two *Condyles*.

The *external Condyle*, placed at the under and outer part of the Bone, for the origin of the *Extensor Muscles* of the *Hand* and *Fingers*. Tab. XXII. Fig. 1. n.

The *internal Condyle*, at the under and inner part of the Bone, more prominent than the former, for the origin of the strong *Flexor Muscles* of the *Hand* and *Fingers*. Tab. XXII. Fig. 1. m.

The *articulating Surface* at the under end of the Bone, between the *Condyles*, covered with *Cartilage* for the articulation with the *Bones* of the *Fore-arm*. Tab. XXII. Fig. 2. o, p, q.

The *oblique Situation* of the *articulating Surface*, the inner end being lower than the outer, by which the *Hand* turns more readily to the upper parts of the *Body*. Tab. XXII. Fig. 1. Tab. I.

The

The *inner Part* of the articulating Surface, consisting of a large internal, and small external eminence, with a middle Cavity, or a Trochlea, upon which the Ulna moves. Tab. XXII. Fig. 1. *p, p, q.* Fig. 2. *o, o, p.*

The *outer Part* of the Articular Surface, upon which the head of the Radius moves, of a round form, and considered by some authors as the smooth part of the upper Condyle. Tab. XXII. Fig. 1. *o.*

Round the Edge of the Articular Cavity, the Bone marked by the insertion of the Capsular Ligament of the Joint. Tab. XXII. Fig. 1. *2.*

A *small Cavity* at the under and fore part of the Bone, above the Trochlea, for receiving the Coronoid Process of the Ulna in the Flexion of the Fore-arm. Tab. XXII. Fig. 1. *3.*

A *large Cavity* at the under and back part of the Bone, also above the Trochlea, the under part of it for receiving the Olecranon of the Ulna in the extension of the Fore-arm, and the upper part for containing the Fat of the Joint. Tab. XXII. Fig. 2. *r.*

Between these Cavities, the Bone is pressed so thin as to become transparent, especially in an Old Person.

The Substance and inner Structure of the Os Humeri is the same as in other long round Bones.

The Ball of the Os Humeri is articulated with the Glenoid Cavity of the Scapula, which, from its superficial nature, and the long Ligaments inclosing the Joint, allows the Arm to move in all directions; the Bone even performing a small degree of motion round its own axis. The extent of motion of the Arm, however, is considerably increased by the rolling of the Scapula.

In the Fœtus, the Extremities of the Bone are Cartilaginous; and the Ball with the Tubercles, and the Trochlea with the Condyles, form afterwards Epiphyses, previous to their union with the Body of the Bone.

FORE-ARM,

Consisting of two Bones, the *Ulna* and *Radius*.

ULNA.

The *Situation* of the Ulna at the inner part of the Fore-arm; the Arm being supposed to hang by the side of the Body, with the Palm of the Hand turned forwards. Tab. 1. *E.*

The *Olecranon, Processus Anconæus, or Top of the Cubit*, placed at the upper end of the Bone, and forming the posterior prominent part of the Elbow. Tab. XXII. Fig. 4. *a, e, e, d.*

The upper end of this Process, *rough* where the Triiceps Extensor Cubiti is fixed.

The *Coronoid, or sharp Process*, at the upper and fore part of the Bone, but considerably lower than the Olecranon, for forming a part of the Hinge of the Joint of the Elbow. Tab. XXII. Fig. 4. *b, c, c.*

The *Great Sigmoid, or Semilunar Cavity*, between the

Olecranon and Coronoid Process, lined with Cartilage, and divided into two slanting Surfaces by a middle Ridge; the Cavity being adapted to the Trochlea of the Os Humeri. Tab. XXII. Fig. 4. *d, e, e.*

Across the middle of the great Sigmoid-Cavity, a *little Pit* for lodging part of the Fat of the Joint.

The *Small Sigmoid, or Semilunar Cavity*, lined with Cartilage, at the outer side of the Coronoid Process, where the round head of the Radius plays. Tab. XXII. Fig. 4. *f.*

The *Tubercle* of the Ulna, or small rough spot under the root of the Coronoid Process, for the insertion of the Brachialis Internus. Tab. XXII. Fig. 4. *b.*

At the upper and outer part of the Bone, a *triangular Surface*, where the Anconeus is lodged.

The *Body* of the Ulna, of a *triangular form*, becoming gradually smaller in its descent. Tab. XXII. Fig. 4.

The *sharpest Angle* opposed to the Radius, for the attachment of the Interosseous Ligament. Tab. XXII. Fig. 4. *g, g.*

The sides forming this Angle, *flat*, and *marked* by the Muscles which arise from them. Tab. XXII. Fig. 4.

A *Passage* slanting upwards, about a hand-breadth below the upper end, for the Medullary Vessels. Tab. XXII. Fig. 4. *l.*

The under end of the Bone, forming a *small round Head*, which is covered with Cartilage on that side where the Radius moves upon it, and also on its extremity, where it is opposed to a moveable Cartilage placed between it and the Carpus. Tab. XXII. Fig. 4. *n, p.*

The *Styloid Process*, at the inner side of the small round head, from which a strong Ligament goes off to be fixed to the Bones of the Wrist. Tab. XXII. Fig. 4. *o.*

The Ulna is articulated at its superior extremity with the lower end of the Os Humeri, the Joint at this part forming a complete Hinge, which allows an extensive degree of flexion, and as much extension as to approach a straight line with the Upper Arm; but little or no rotation.

RADIUS.

The *Situation* of the Radius at the outer part of the Fore-arm. Tab. I. *P.*

The *upper end of the Radius*, covered with Cartilage, and formed into a circular head, which is hollowed above, for receiving the outer part of the Articular Surface of the Os Humeri. Tab. I.

The *inner Side of the Head* smooth, and also covered with Cartilage, where it plays in the small Semilunar Cavity, at the outer side of the Ulna. Tab. XXII. Fig. 3.

The *Cervix* of the Radius smaller than the head; in the Subject, surrounded by a circular Ligament, which keeps the Bone in its place, and allows it to roll upon the Ulna. Tab. XXII. Fig. 3.

The

The *Tubercle* of the Radius, at the under and inner part of the Cervix, for the insertion of the Biceps Flexor Cubiti. Tab. XXII. Fig. 3. c.

The *Body* of the Bone larger than that of the Ulna, convex on its outer and back part, and rounded by the Muscles which cover it.

The *Surfaces* next the Ulna *flat*, where Muscles of the Hand take their origin. Tab. XXII. Fig. 3. e, e.

The *anterior* and *posterior Surfaces*, terminating next the Ulna, in a *sharp Ridge*, to which the Interosseous Ligament of the Fore-arm is fixed. Tab. XXII. Fig. 3. d, d.

A *Passage* slanting upwards, on the fore-part of the Bone, and about a hand-breadth below its upper end, for the Medullary Vessels. Tab. XXII. Fig. 3. g.

A *rough Surface* at the outer and middle part of the Bone, for the insertion of the Pronator Radii Teres.

The *lower End of the Radius*, becoming gradually larger, and flat on its fore-part, where it is covered by the Pronator Radii Quadratus. Tab. XXII. Fig. 3. i.

A *Ridge* upon the under and back part of the Radius, with a *Fossa* upon each side of it, where the Tendons of the Extensor Muscles of the Fingers pass. Tab. II.

The outer side of this extremity of the Bone, *hollowed* by the Extensors of the Thumb. Tab. II.

A *semilunar Cavity* at the inner side of the under end of the Radius, lined with Cartilage for receiving the corresponding extremity of the Ulna. Tab. XXII. Fig. 3. m.

The *lower End* of the Bone, formed into a *Cavity* of an *oval* or *navicular form*, and lined with Cartilage for receiving the two first Bones of the Carpus. Tab. XXII. Fig. 3. l.

A small *Transverse Ridge*, frequently found in the middle of this Cavity, which insinuates itself between the two first Bones of the Carpus.

The under and outer part of the Radius, forming a *Process* somewhat similar to the Styloid Process of the Ulna. Tab. XXII. Fig. 3. k. From this Process a Ligament is sent to the Wrist.

The *Head* of the Radius is articulated with the outer part of the articular Surface of the Os Humeri; the Radius is besides joined by a double Articulation to the Ulna, for above, the head of the Radius is received into the small Semilunar Cavity of the Ulna, while the under end of the Ulna is received into the small Semilunar Cavity of the Radius; in consequence of which connection, the Radius accompanies the Ulna in the flexion and extension of the Fore-arm, while the Radius moves round its own axis above, but at the lower end, it turns upon the head of the Ulna, carrying the Hand with it.

The Turning of the Radius with the Hand is termed *Supination*, and *Pronation*; when the Palm is turned upwards it is in a state of supination, and in pronation, when in a contrary direction.

VOL. I.

The Structure of the Radius and Ulna is the same as that of other Long Bones.

In the Fœtus, the Extremities of the Bones of the Fore-arm are Cartilaginous; they afterwards become Epiphyses before they are united to the bodies of the Bones.

HAND,

Composed of the Bones of the *Carpus*, *Metacarpus*, and *Fingers*.

The *posterior Surface* of the Hand, *convex*, which gives it a greater degree of strength.

The *anterior Surface* of the Hand, *concave*, for grasping and holding Substances.

CARPUS.

THE *Carpus* is composed of *eight Bones*, disposed in two Rows, and each Bone being broader on its posterior than anterior Surface; they form an Arch, convex behind, by which it gives security and strength; and concave before, for containing the Muscles, Vessels, and Nerves, which run to the Fingers. The ends of the Arch on the Palm side of the Wrist, form projecting Points, between which the Ligamentum Carpi Annulare is stretched, which confines the Muscles in their places. Tab. XXIV. Fig. 2. Tab. XXXI. Fig. 4.

The posterior or convex Surface of the Carpus, *marked* by the numerous Ligaments attached to it.

The anterior or hollow Surface, also marked by Ligaments.

The Surfaces of the Bones of the Carpus are articulated with each other, or with the neighbouring Bones, and all their articular Surfaces are *covered with Cartilage*, to facilitate the motion of the Joints.

In the First Row of Carpal Bones are,
The *Scaphoides*, *Lunare*, *Cuneiforme*, *Pisiforme*.

In the Second Row,

The *Trapezium*, *Trapezoides*, *Magnum*, *Unciforme*.

The Os SCAPHOIDES, placed at the outer and upper part of the Carpus. Tab. XXIV. Fig. 1. B.

The *upper Surface*, *convex*, and articulated with the Radius. Tab. XXII. Fig. 5. a.

The *under* and *outer Surface*, also *convex*, to be articulated with the Trapezium, and Trapezoides. Tab. XXII. Fig. 5. d.

Between the upper and under Cartilaginous Surfaces, a *rough Fossa* for the insertion of the Capsular Ligament. Tab. XXIV. Fig. 5. c.

The anterior and inner Surface, having an *oval Cavity* which gives name to the Bone, where it is articulated with the Os Magnum. Tab. XXII. Fig. 5. b.

A *Process* upon the outer end of the Bone, for the attachment of part of the anterior Transverse Ligament of the Wrist. Tab. XXII. Fig. 1. under B.

I

The

The Os LUNARE, situated upon the inner side of the former Bone. Tab. XXII. Fig. 1. C.

The *upper Surface, convex*, for its articulation with the Radius. Tab. XXII. Fig. 1. C.

The *outer Edge*, in form of a *Crescent*, from which the Bone is named; articulated with the Os Scaphoides. Tab. XXII. Fig. 6. a.

The *under Surface, hollow*, for its articulation with the Os Magnum. Tab. XXII. Fig. 6. b.

The *inner Surface* of the Bone, articulated with the Os Cuneiforme. Tab. XXIV. Fig. 1.

The Os CUNEIFORME, situated on the inner side of the former Bone. Tab. XXIV. Fig. 1.

The *anterior Edge, thin*, in form of a *wedge*.

The *upper and outer Surface*, articulated with the Os Lunare. Tab. XXIV. Fig. 1.

The upper part forms a slight Convexity, which is included in the Joint of the Wrist. Here the moveable Cartilage already taken notice of is interposed between this Bone and the Ulna.

The *under and outer Surface*, articulated with the Os Unciforme. Tab. XXIV. Fig. 1.

The *anterior and inner Surface*, forming a slight *convexity* for its articulation with the Os Pisiforme. Tab. XXII. Fig. 7. a.

The three first Bones of the Carpus form an oval convexity, by which they are articulated with the lower end of the Bones of the Fore-arm; the Offa Scaphoides and Lunare being received in the Socket formed by the Radius, while the Os Cuneiforme is opposed to the Cartilaginous end of the Ulna. Tab. XXIV. Fig. 1. By this kind of articulation, extensive motion is allowed forwards and backwards, and to each side; and by a succession of these motions the hand is made to move in a circle, but no motion is performed by the Carpus round its own axis, except what it has along with the Radius in the Supination and Pronation of the Hand.

The Os PISIFORME, placed upon the anterior and inner Surface of the Os Cuneiforme, forming a Prominence which is readily felt in the Wrist, and which gives attachment to strong Tendons and Ligaments, particularly to part of the Ligamentum Carpi Annulare. Tab. XXII. Fig. 8. Tab. XXIV. Fig. 1. D.

The Os TRAPEZIUM, named from the four unequal Edges of its posterior Surface.

The *Situation* of this Bone, at the root of the Metacarpal Bone of the Thumb. Tab. XXIV. Fig. 1. A.

The upper part of the Bone, forming a *smooth Pit*, to be articulated with the Os Scaphoides. Tab. XXIV.

The inner side, *hollow*, and articulated with the Os Trapezoides. Tab. XXIV.

The under Surface, forming a *Pulley* on which the Metacarpal Bone of the Thumb moves. Tab. XXXI. Fig. 4.

The anterior Surface, sending out a Process, which is prominent in the Palm, and marked by the Transverse Ligament of the Wrist, by the Flexor Carpi Radialis, and Flexors of the Thumb. Tab. XXII. Fig. 9. b.

The Os TRAPEZOIDES, so named from its being somewhat like the former Bone, though considerably smaller. Tab. XXII. Fig. 10.

The *Situation* of the Os Trapezoides, at the inner side of the Os Trapezium. Tab. XXIV. Fig. 1. E.

The upper Surface, *hollow* where it joins the Os Scaphoides. Tab. XXXI. Fig. 5. m.

The outer Surface, *convex*, and articulated with the Trapezium. Tab. XXXI. Fig. 5. m.

The *inner Surface*, articulated with the Os Magnum. Tab. XXIV. Fig. 1. E.

The *under Surface*, formed into a sort of *Pulley*, to be articulated with the Metacarpal Bone of the Fore-Finger. Tab. XXIV. Fig. 1. E.

The Os MAGNUM, or CAPITATUM, or *largest Bone* of the Carpus, placed at the inner side of the former Bone, and consisting of four oblong sides, with a round head, and triangular under end. Tab. XXIV. Fig. 1. F.

The *head or ball* of the Bone, received into the hollow Surfaces of the Scaphoides and Lunare, like Ball and Socket. Tab. XXIV. Fig. 1.

The *under part* of the outer side, joined to the Os Trapezoides. Tab. XXIV.

The *inner side*, to the Os Unciforme. Tab. XXIV.

The *under end*, opposed to the Metacarpal Bone of the Middle Finger. Tab. XXIV.

The Os UNCIFORME, placed in the under and inner part of the Wrist. Tab. XXIV. Fig. 1. G.

The *upper and inner Surface*, articulated with the Os Cuneiforme. Tab. XXIV.

The *outer Surface*, articulated with the Os Magnum. Tab. XXIV.

The *inferior Surface*, opposed to the Metacarpal Bones of the Ring and Little Fingers. Tab. XXIV.

The anterior Surface, sending out the *Unciform Process*, which gives name to the Bone. Tab. XXIV. Fig. 1. H.

The Unciform Process, *curved* for the passage of the Flexor Muscles of the Fingers. Tab. XXIV.

The Articulation between the first and second Row of Carpal Bones, allows motion to each side, but chiefly forwards and backwards; the motion, however, is less extensive than between the Fore-arm and Wrist.

The

The Connection between the different Bones in each Row, is of such a nature as not to admit of any sensible motion.

The Substance of the Carpal Bones is spongy, but strong in proportion to their size.

The Carpus serves as a Base to the Hand, protects its Tendons, &c. and affords free and extensive motion.

In the Fœtus, the Bones of the Carpus are in a Cartilaginous state.

METACARPUS,

Consisting of *four Bones* for supporting the Fingers, and one for the Thumb. Tab. XXIV. Fig. 1. K, &c. I.

Metacarpal Bones of the Fingers.

Their *bodies* long and round behind, forming part of the convexity of the Hand, before, giving hollowness to the Palm. Tab. XXXII. Fig. 5. 4.

The *extremities* of these Bones, considerably larger than their bodies, in consequence of which they leave spaces for the Interossei. Tab. XXXII.

The *upper ends* or *bases*, *flat*, where they are articulated with the Bones of the Carpus. Tab. XXIV.

Round the edges of the Cartilaginous Surfaces, at the upper ends, the *depressions* where the Capsular Ligaments are fixed. Tab. XXIV. Fig. 1.

The *sides* of the upper ends *flat*, and drawn close together, where they are articulated with each other. Tab. XXIV. Fig. 1.

Their *Bodies* diverging towards their under extremities, by which they regulate the motions of the Fingers. Tab. XXXII.

A *Ridge* at the upper and back part of their Bodies, with a depression on each side of it, formed by the Interossei. Tab. XXXII. Fig. 5.

The under and back part of their Bodies, made *flat* by the motion of the Tendons of the Extensors of the Fingers. Tab. XXXII. Fig. 5.

The anterior Surface of their Bodies *concave*, and rendered *flat at the sides* by the Interosseous Muscles. Tab. XXIV. Fig. 1.

The *lower ends*, or *heads*, formed into *Balls*, which are flattened upon their sides by their motions upon each other. Tab. XXIV. Fig. 1.

At the fore-part of each side of the heads, a little *prominence*, for the attachment of the Ligaments which fix these Bones to each other. Tab. XXIV. Fig. 1.

Round the heads, a *depression* for the insertion of the Capsular Ligaments. Tab. XXIV. Fig. 1.

The Metacarpal Bones are joined by their Base to the Carpus, and to each other by nearly plain Surfaces; in consequence of which, and the strength of their connecting Ligaments, their motions here are inconsiderable.

The *Base* of the *Metacarpal Bones of the Fore-finger*, opposed to, and corresponding with, the Os Trapezoides, and partly with the Os Trapezium. Tab. XXXII. Fig. 5.

The inner part of the Base, forming a *Ridge*, which is articulated with the Os Magnum, and with the next Metacarpal Bone. Tab. XXIV. Fig. 1.

The connection of the Base so firm, that it has little or no motion.

The *Metacarpal Bone of the Mid-finger*, commonly the *second in length*. Tab. XXIV. Fig. 1.

The Base of the Bone generally slanting inwards and downwards, opposed to the Os Magnum. Tab. XXXII. Fig. 5.

The outer and back part of the Base *projecting*, and forming a sort of *Process*, the external Surface of which is connected with the Ridge of the former Bone. Tab. XXXII. Fig. 5.

The motion of this Bone is little more than that of the former one.

The *Metacarpal Bone of the Ring-finger*, shorter than the former Bone. Tab. XXXII. Fig. 5.

Its Base, *semicircular* where it is opposed to the Os Unciforme. Tab. XXXII. Fig. 5.

The motion is something greater than that of the former Bone.

The *Metacarpal Bone of the Little Finger*, the *smallest* of the four. Tab. XXIV. Fig. 1.

The Base, which slants downwards and outwards, opposed to the under and inner part of the Os Unciforme. Tab. XXXII. Fig. 4.

The inner part of the Base having no smooth Surface, not being contiguous to any other Bone.

From the nature of the Joint, the looseness of the Ligaments, and from the existence of a proper Muscle here, this Bone possesses a larger share of motion than any of the rest.

Metacarpal Bone of the Thumb.

Having the general resemblance of those of the Fingers, but differing from them in being placed oblique with respect to the Metacarpal Bones of the Fingers, and in some measure opposing them. Tab. XXIV. Fig. 1. I.

This Bone thicker and stronger, but shorter than those of the Fingers. Tab. XXIV. Fig. 1.

The *Base* of this Bone articulated with the Pulley formed by the Trapezium, the Bone appearing to admit of flexion and extension only; but, from the looseness of the Ligaments, enjoying the same kind of motion with Joints formed after the manner of Ball and Socket. Tab. XXIV. Fig. 1.

The *inferior extremity* of the Bone, considerably *flatter* than those of the other Metacarpal Bones. Tab. XXIV.

The

The *Fingers*, composed each of three Bones; the three Rows of Bones taken transversely, termed *Phalanges*. Tab. XXIV. N, &c. O, &c. P. &c.

The different *Phalanges*, tapering a little as they descend, and their Bases larger than their inferior extremities. Tab. XXIV.

The posterior Surfaces *convex*, and covered chiefly by the tendinous expansions of the Extensors of the Fingers. Tab. XXXII. Fig. 5.

Their anterior Surfaces *flat*, and in some parts *concave*, for lodging the Tendons of the Flexor Muscles. Tab. XXXII. Fig. 4.

Ridges at the sides of their anterior Surfaces, for the attachment of the retaining Ligaments of the Tendons of the Flexor Muscles. Tab. XXIV. Fig. 1.

The first Phalanx *longer* than the second, and the second than the third. Tab. XXIV. Fig. 1.

The *Bases* of the first Phalanx formed into *Sockets*, to receive the Balls of the Metacarpal Bones, and to allow motion to all sides. Tab. XXIV. Fig. 1. *i, i*.

The lower ends of this Phalanx, consisting of *lateral Prominences*, and *middle Cavities* or Pulleys, the Cartilaginous Surfaces of which reach considerably farther up in the fore than in the back part. Tab. XXXII. Fig. 4. 5.

The *Bases* of the second Phalanx, with *lateral Cavities*, and *middle Ridges*, corresponding with the Pulleys of the first Phalanx, and admitting of flexion and extension only. Tab. XXIV. Fig. 1. *m, m*.

The *lower ends* of this Phalanx, similar to those of the first. Tab. XXIV. Fig. 1. *n, n*.

The *Bases* of the *third Phalanx* like those of the second, and the motions also similar. Tab. XXIV. Fig. 1. *p, p*.

The under ends of the third Phalanx, *rough* where the

Pulpy, Vascular, and Nervous Substance of the points of the Fingers is situated. Tab. XXIV. Fig. 1.

The Peculiarities of the Bones of the Fingers consist only in their size.

The Bones of the Mid-finger the largest and longest. Tab. XXIV. Fig. 1.

Those of the Ring-finger next in length. Tab. XXIV.

The Bones of the Fore-finger next to those of the Ring-finger in length, and of the Mid-finger in thickness. Tab. XXIV.

Those of the Fourth Finger the smallest. Tab. XXIV.

The *Thumb*, consisting of only *two* Bones. Tab. XXIV. Fig. 1. L, M.

The *first* Bone like the Bones of the first Phalanx of the Fingers, but *thicker* and *stronger*. Tab. XXIV.

The *Cavity* at the Base of the Bone, longer from one side to the other, and shallower than the Cavities of the corresponding Bones of the Fingers, but, like them, forming a Socket for the Metacarpal Bone. From the flatness of the Joint, however, and strength of the lateral Ligaments, the motions here are confined to flexion and extension only. Tab. XXIV.


The *lower end* of the first Bone of the Thumb like that of the first of the Fingers. Tab. XXIV.

The *second* Bone of the Thumb like the *third* of the Fingers, but broader. Tab. XXIV.

The *Base* of this Bone, like that of the second and third Bones of the Fingers, and like their Joints also, admitting of flexion and extension only.

The Substance of the Metacarpus, and of the Bones of the Fingers, is the same with that of the Long Bones.

In the Fœtus, both extremities of the Metacarpal Bones of the first and second, and upper ends of the third Phalanx, are in a state of Cartilage.



TAB. 22.

Fig. 2.



Fig. 4.



Fig. 3.




Fig. 1.



Fig. 8.



Fig. 7.



Fig. 6.



Fig. 5.



Fig. 12.



Fig. 11.



Fig. 10.



Fig. 9.



T A B L E XXII.

Represents the BONES of the UPPER ARM, FORE-ARM, and CARPUS of the LEFT SIDE.

FIG. 1.

The Fore-Part of the Os HUMERI.

- a*, The middle of the ball of the os humeri.
b, b, The cervix of the os humeri.
c, Part of the large tuberosity.
d, The small tuberosity.
e, e, The groove, or sinuosity which receives the long head of the biceps.
f, f, The orifices of several conduits, for the insertion of the fibres of the capsular ligament, and for the passage of vessels into the bone.
g, The projecting line which answers to the large tuberosity.
h, Another ridge which answers to the small tuberosity.
i, Muscular prints which give insertion to the deltoides.
k, k, The long line which occupies the whole length of the os humeri.
l, The internal condyle.
m, m, m, The three facets, or muscular prints of this condyle.
n, Part of the external condyle.
o, The head, or rounded eminence, which is articulated with the radius.
p, p, Two other articular eminences which correspond to the ulna.
q, The articular cavity, which receives the middle process of the ulna.
r, Another articular cavity, which facilitates the motion of the radius.
s, The cavity which receives the coronoid process of the ulna, upon bending the fore-arm.

FIG. 2.

The Posterior Part of the same BONE.

- a*, The posterior part of the head of the os humeri.
b, b, The cervix of this bone.
c, c, c, The muscular prints of the large tuberosity.
d, d, The orifices of different conduits which open into the substance of the bone.

- v, e*, The superficial triangular cavity, which gives attachment to muscles.
f, The orifice of the passage which communicates with the inner cavity of the os humeri.
g, g, The projecting line which answers to the external condyle.
h, A part of the posterior surface, which answers to the external surface.
i, A second posterior surface, which also answers to the external surface.
k, k, A third surface, which is blended superiorly with the external surface.
l, The external condyle.
m, Several muscular prints on the posterior part of this condyle.
n, A portion of the eminence of the os humeri, which is articulated with the radius.
o, o, The two eminences of the os humeri, which are articulated with the ulna.
p, The internal condyle of the os humeri.
q, The articular cavity of the os humeri, which facilitates the motion of the ulna.
r, The posterior cavity, which receives the olecranon upon the extension of the fore-arm.

FIG. 3.

The Inner, and Part of the Outer Surface of the RADIUS.

- a*, The semicircular eminence of the radius, which is lodged in the sigmoid cavity of the ulna.
b, The cervix of the radius.
c, The tuberosity of the radius.
d, d, The crest, or osseous line.
e, e, The inner surface of the radius.
f, f, A portion of the anterior surface.
g, The small conduit which opens into the inner cavity of the bone.
h, An osseous line, or muscular print.
i, A small fossa at the inferior extremity of the radius.
k, The styloid process.
l, The oval cavity which receives the first of the carpal bones.

m, The

m, The femilunar cavity, which receives the articular process of the ulna.

FIG. 4.

The Outer and somewhat Posterior Surface of the ULNA.

- a*, The olecranon, a little hollow in the middle.
- b*, The coronoid process.
- c, c, c*, Several muscular and ligamentous impressions.
- d*, The middle, or articular process of the ulna.
- e, e*, The articular, or great femilunar cavity.
- f*, The small sigmoid, or femilunar cavity.
- g, g, g*, An osseous line, extending the whole length of the ulna.
- h*, A muscular print below the sigmoid cavity.
- i, i, i*, The inner surface of the ulna.
- k, k*, The outer surface.
- l*, A small conduit which communicates with the cavity of the bone.
- m*, A small fossa, or muscular print of the inferior extremity of the ulna.
- n*, A semicircular eminence articulated with the under end of the radius.
- o*, The styloid process.
- p*, A portion of the cavity which answers to the carpus.

FIG. 5.

The Convex or External Surface of the Os SCAPHOIDES, viewed in a Position most favourable for shewing its NAVICULAR CAVITY.

- a*, The upper part of the os scaphoides, by which it is articulated with the os lunare.
- b*, The navicular cavity, which receives the head of the large carpal bone.
- c*, The middle rough part, to which several ligaments are attached.
- d*, The under end, by which it is articulated with the trapezium and trapezoides.

FIG. 6.

The Outer Surface of the Os LUNARE, the under Part of which is chiefly shewn.

- a*, The articular facet of the os lunare, excavated in form of a crescent, by which it is joined to the os scaphoides.
- b*, The large lunar cavity, by which this bone is articulated with a considerable portion of the large carpal bone.
- c*, Part of the inferior surface, by which it is joined to the os cuneiforme.

FIG. 7.

The Inner Surface of the Os CUNEIFORME.

- a*, The small surface, which is articulated with the os pisiforme.
- b, b*, Different ligamentous impressions.

FIG. 8.

Os PISIFORME.

- a*, The articular surface by which it is connected with the os cuneiforme.

FIG. 9.

The Inferior Part of the TRAPEZIUM, or First of the Second Range of the CARPAL BONES.

- a, a*, The two small surfaces on the under part of the trapezium.
- b*, Part of the oblique process.
- c*, Part of the sinuosity.
- d*, The small surface by which it is joined with the first metacarpal bone.

FIG. 10.

The different Surfaces of the Os TRAPEZOIDES.

- a*, Part of the external surface.
- b*, The inferior surface.
- c*, The anterior surface, by which it is joined with the trapezium.
- d*, Part of the superior surface, by which it is joined with the scaphoides.
- e*, Part of the print of this bone, which is turned towards the palm.

FIG. 11.

The External and somewhat Anterior Surface of the Os MAGNUM.

- a*, The head of this bone incrustated with a smooth cartilage, for facilitating the motions of its articulation with the scaphoides and lunare.
- b*, The cervix of this bone.
- c*, The outer surface.
- d*, Part of its anterior surface.
- e*, Its under edge, by which it is joined with the second metacarpal bone.

FIG. 12.

The Posterior Part of the Os UNCIFORME.

- a*, The upper part, which answers to the os lunare.
- b*, The inner surface, where there is a sinuosity for the passage of several tendons.
- c*, The unciform process, on which there is a sort of groove, which also facilitates the passage of tendons.
- d*, The posterior and superior surface, by which it is joined to the os cuneiforme.
- e, e*, The inferior surface, divided by a small superficial line, the larger part of which corresponds to the third, and the other to the last metacarpal bone.

TABLE

Fig. 1



Fig. 2 A



Fig. 4




Fig. 3



T A B L E XXIII.

Represents the different BONES of the EXTREMITIES, excepting those of the THUMB and FINGERS.

FIG. 1.

An Inside View of the BONES of the SHOULDER and ARM.

- A, The venter, or anterior cavity of the scapula.
- B, The acromion.
- C, The notch of the scapula, which transmits vessels and nerves.
- D, D, The inferior costa.
- E, E, The base.
- F, The anterior angle, which contains the glenoid cavity.
- G, The coracoid process.
- H, The posterior angle.
- I, The inferior angle.
- K, The posterior end of the clavicle, fixed to the acromion scapulæ.
- L, The sternal extremity.
- M, The ball of the os humeri, articulated with the glenoid cavity of the scapula.
- N, The cervix.
- O, The inner, or small tuberosity.
- P, The round body.
- Q, The internal condyle.
- R, The trochlea.

FIG. 2.

A View of the Anterior and Inner SURFACE of the BONES of the FORE-ARM, to shew their Articulation.

- A, The great sigmoid cavity of the ulna, which receives the trochlea of the os humeri.
- B, The olecranon of the ulna.
- C, The coronoid process.
- D, Its triangular body.
- E, The inferior extremity, articulated with the carpus by the intervention of a cartilage.
- F, The styloid process.
- G, The head of the radius, occupying the semilunar cavity of the ulna.

H, The cervix of the radius.

I, The tuberosity for the insertion of the biceps.

K, The triangular body of the radius.

L, The interstice between the radius and ulna, filled chiefly with the interosseous ligament.

M, The inferior broad extremity of the radius, articulated with the carpus and ulna.

N, N, The anterior concave surface of the carpus, composed of

The os scaphoides,

The os lunare,

The os cuneiforme,

The os pisiforme,

The os trapezium.

The os trapezoides,

The os magnum, and

The os unciforme.

O, &c. The metacarpal bones of the thumb and fingers.

FIG. 3.

The Outer and Fore Part of the OS FEMORIS and PATELLA.

- A, The ball of the thigh-bone.
- B, The cervix of this bone.
- C, The trochanter major.
- D, The trochanter minor, which ought not to be seen in this view of the bone.
- E, The curved body of the bone.
- F, The external condyle.
- G, The cartilaginous surface, on which the tibia moves.
- H, The trochlea, which receives the patella.
- I, The patella.

FIG. 4.

A View of the Outer and Fore Part of the BONES of the LEG and FOOT, in their connected State.

- A, The head of the tibia.
- B, The tubercle of this bone.

C, The

TABLE XXIII. CONTINUED.

- C, The body of the tibia.
 D, The lower extremity, at the inner side of which is the malleolus internus.
 E, The head of the fibula, joined to the outer part of the tibia.
 F, The irregular surface of the body of the fibula.
 G, The lower end, or malleolus externus, joined to the outer side of the tibia and tarsus.
 H, The space between the tibia and fibula, filled with the interosseous ligament.
 I, to P, The tarsus, composed of,
 I, The astragalus.
 K, The os calcis.
 L, The projection of this bone forming the heel.
 M, The os naviculare.
 N, The os cuneiforme medium, on the inner side of which is seen a small part of the os cuneiforme internum.
 O, The os cuneiforme internum, and,
 P, The os cuboides.
 Q, &c. The metacarpal bones of the toes.
 R, &c. The first phalanx of the bones of the small toes.
 S, &c. The second phalanx.
 T, &c. The third phalanx.
 U, The first bone of the great toe.
 W, The second bone of the great toe.

TABLE





Fig. 2.

Fig. 1.



T A B L E XXIV.

Represents the BONES of the HAND and FOOT.

FIG. 1.

View of the Inner, or Palm-Side of the BONES of the LEFT HAND.

- B, The os scaphoides.
 C, The os lunare.
 C, G, The os cuneiforme.
 D, The os pisiforme.
 A, The os trapezium.
 E, The os trapezoides, or pyramidale.
 F, The os magnum.
 * H, The os unciforme.
 H, The unciform process.
 I, The metacarpal bone of the thumb.
 a, The base of this bone, sending inwards a coronoid process.
 b, The inferior extremity.
 K, &c. The metacarpal bones of the fingers.
 k, k, &c. The interstices occupied by the interosseous muscles.
 c, c, &c. The upper ends irregular, where they are joined to the carpus, and to each other.
 d, d, The under ends, in form of balls, covered with cartilage.
 L, The first bone of the thumb, concave, for lodging the flexor longus.
 e, The base, forming a glenoid cavity for the articulation with the metacarpal bone.
 f, The lower end, with two lateral protuberances, and a middle cavity.
 M, The second bone of the thumb.
 g, The base, with two lateral cavities, and a middle protuberance corresponding to the end of the former bone.
 h, The under and inner side, rough and irregular, where the soft substance at the end of the thumb is placed.

- N, N, &c. The first phalanx of the bones of the fingers, flat like the surface marked L.
 i, i, The base with a cavity similar to e, but rounder.
 l, l, The inferior extremity, similar to f.
 O, &c. The second phalanx; hollow like the first.
 m, m, The base, similar to g.
 n, n, The under and inner surface, like f.
 P, q, &c. like g, b.

FIG. 2.

A View of the Under Side of the BONES of the LEFT FOOT.

- A, A, The astragalus.
 a, The upper and inner surface of the astragalus.
 B, The body of the os calcis.
 C, That portion of the os calcis which forms the lower part of the heel.
 b, That part to which the tendo Achillis is fixed.
 c, The large sinuosity of this bone, which lodges the principal muscles, tendons, vessels, and nerves of the sole.
 D, The os naviculare.
 E, The os cuneiforme internum.
 F, The os cuneiforme medium.
 G, The os cuneiforme externum.
 H, The os cuboides.
 d, The fossa of the os cuboides, for lodging the tendon of the peroneus longus.
 I, The metatarsal bone of the great toe.
 K, &c. The metatarsal bones of the small toes.
 b, &c. Interstices occupied by the interosseous muscles.
 L, The first bone of the great toe.
 M, The second bone of the great toe.
 N, &c. The bones of the first phalanx of the toes.
 O, &c. The bones of the second phalanx of the toes.
 P, &c. The bones of the third phalanx of the toes.

INFERIOR EXTREMITY.

THE *Inferior Extremity* composed of the Thigh, Leg, and Foot.

The *Thigh* consisting of a single Bone, viz.

OS FEMORIS.

THE Os Femoris the *longest* of the Body, and *thickest* and *strongest* of the Cylindrical Bones.

The *Situation* of the Bone, at the under and outer part of the Pelvis. Tab. XXIX.

The *oblique* Situation of the *body* of the Bone; the under end being considerably nearer its fellow on the other side than the upper one is, which is favourable for the passages at the bottom of the Pelvis, for the origin of Muscles, and for walking. Tab. I.

The *Ball* or *Head* of the Thigh-bone, smooth, covered with Cartilage, and forming almost two-thirds of a Sphere, which is received into the deep Socket formed by the Acetabulum of the Os Innominatum. Tab. XXV. Fig. 1. a. Tab. XX.

A *rough Pit* at the inner part of the Ball, for the attachment of the Ligamentum Rotundum, which is fixed by its other end to the bottom of the Acetabulum. Tab. XXV. Fig. 1. b.

The *Cervix* or *Neck*, much longer than that of any other Bone, passing obliquely downwards and outwards from the Ball, to allow the free motion of the body of the Bone in different directions. Tab. XXV.

Numerous Holes in the Cervix, for the insertion of the Fibres of the Ligament reflected from the Capsular one. Tab. XXV. XXIX.

The *Trochanter Major*, placed at the outer part of the Neck, and upper end of the Body of the Bone, for the insertion of the Extensor, Abductor, and Rotator Muscles of the Thigh. Tab. XXV. Fig. 1. e.

Two Rough Surfaces upon the upper and fore part of the large Trochanter, for the insertion of the Glutei Medius and Minimus. Tab. XXV. Fig. 1. f.

A *Cavity*, placed at the inner side of the root of the large Trochanter, for the insertion of the Rotator Muscles of the Thigh.

The *Trochanter Minor*, at the under and inner part of the Cervix, for the insertion of the Flexor Muscles of the Thigh. Tab. XXV. Fig. 1. b.

The Trochanter Minor is small and pointed, and in the Subject is so much covered by Muscles, as to be out of the reach of the Finger.

A *rough Line* on the fore-part of the Bone, extending

obliquely between the two Trochanters, for the insertion of the Capsular Ligament. Tab. XXV. Fig. 1. g, g.

A *rough Line* between the Trochanters, on the *back-part* of the Bone, for the insertion of the Capsular Ligament, and of the Quadratus Femoris. Tab. XXV. Fig. 2. k, k.

The *Body* of the Thigh-bone, *bent forwards*, and a *little outwards*, of a *roundish form* above, but somewhat *triangular* about its middle. Tab. XXIII. Tab. XXV.

The *fore-part* of the Bone, *flat* where it is covered by the Crureus. Tab. I. A.

The *Sides* of the Bone, *flattened* at its middle and lower part by the two Vasti. Tab. XXV. Fig. 2.

The *Linea Aspera* or *ragged Ridge*, on the back-part of the Bone, extending from the Trochanters, but chiefly from the large one, to the lower part of the Bone, and giving attachment to numerous Muscles which pass from the Pelvis to the Thigh, or from the Thigh to the Leg. Tab. XXV.

The *Linea Aspera* is forked at both its extremities; extending above to the Trochanters, while below, the two lines into which it divides, terminate in the Condyles. Tab. XXV.

The *Canal* for the Medullary Vessels, slanting upwards, a little below the middle of the posterior part of the Bone. Tab. XXV. Fig. 2. g.

The *under and back part* of the Bone, *flat* where the Popliteal Vessels and Nerves are placed.

The *lower End* of the Bone becoming gradually enlarged, and perforated by many Holes, for the insertion of the Capsular Ligament of the Knee, and for the passage of the Nutritious Vessels of the Bone. Tab. XXV.

The lower End, also marked by the insertion of several Muscles. Tab. XXV.

The *Cartilaginous Trochlea* at the under and fore part of the Bone, placed obliquely, with its outer Surface larger and higher than its inner one, to be adapted to the Patella, which moves upon it. Tab. XXV.

The *external and internal Condyles*, continued back from the Trochlea, and also covered with Cartilage, for the motion of the Tibia. Tab. XXV.

The *internal Condyle*, *larger and deeper* than the external, to compensate for the obliquity of the Thigh, and give less obliquity to the Leg. Tab. XXV. Fig. 2. v, v.

A *Notch* between the back-part of the Condyles, for lodging the Popliteal Vessels and Nerves. Tab. XXV. Fig. 2. z, z.

A *semilunar rough Notch*, deeper and lower than the former

former one, for the attachment of the Crucial or internal Ligaments of the Knee. Tab. XXV. Fig. 2. *y*.

The Thigh-bone is articulated above with the Os Innominatum, which allows the free motion of the Body of the Bone in all directions. It is restrained, however, in its motions outwards by the Ligamentum Rotundum, and by the high Brim of the Acetabulum.

The Head and Neck of the Bone can move round their own axis, though its Body possesses little rotatory motion. In consequence of the oblique situation of the Head and Neck when the Ball rolls, the Body of the Bone is only brought forwards or backwards.

In the Fœtus, the different Processes of the Bone are Cartilaginous, and afterwards form large Epiphyses. Tab. XXVII. Tab. XXXII. Fig. 15.

LEG,

Composed of two Bones, the Tibia and Fibula,—to which may be added the Patella.

TIBIA,

Situated at the inner part of the Leg. Tab. I. C.

The upper End of the Tibia, forming a large Head, and that divided on its upper Surface into two superficial Cavities, for receiving the Cartilaginous part of the Condyles of the Thigh-bone. Tab. XXV. Fig. 3. Tab. XXXII. Fig. 7.

A rough Protuberance projecting between the articulating Cavities, and received in the space between the Condyles. It is pitted on its fore and back parts, for the insertion of the anterior and posterior Crucial Ligaments. Tab. XXXI. Fig. 17. No. 4.

The articulating Surfaces at the upper end of the Tibia, rendered deeper in the Subject by the addition of two femilunar Cartilages placed upon their Edges. Tab. XXXII. Fig. 7. *c, c*.

The circumference of the Head of the Bone, rough and porous, for the insertion of the Capsular Ligament. Tab. XXV. Fig. 3. *b, b*.

A Tubercle at the upper and fore part of the Bone, for the insertion of the lower Tendon or Ligament of the Patella. Tab. XXV. Fig. 3. uppermost *f*.

A Cartilaginous Surface under the outer Edge of the Head of the Bone, for the articulation with the upper end of the Fibula. Tab. XXV. Tab. XXIII. Fig. 4. E.

The Body of the Bone, of a triangular form, with the sharpest Angle placed anteriorly. Tab. XXV. Fig. 3. *a*.

The anterior Angle, called Spine or Shin, a little wavy, and extending from the Tubercle to the inner Ankle. Tab. XXV. Fig. 3. *b, a, b*.

The anterior and inner Surface of the Bone, smooth, being covered with skin only. Tab. XXV. Fig. 3. *c, c*.

The anterior and outer Surface, hollowed by one of the Flexor Muscles of the Foot, and by the long Extensors of the Toes. Tab. XXV. Fig. 3. *d, d*.

The Angle at the outer and back-part of the Bone, giving attachment to the Interosseous Ligament. Tab. LI. Fig. 1. 2.

The middle of the posterior Surface, also hollowed by Muscles which assist in extending the Foot, and in bending the Toes. Tab. XXXI. Fig. 17. No. 5.

A Ridge extending obliquely downwards from the upper and outer part of the Bone, posteriorly, to its inner Angle, and giving origin to part of the Muscles which extend the Foot and bend the Toes. Tab. XLIII. Fig. 2.

A flat Surface above the Ridge, indicating the situation of the Popliteus. Tab. XLIII. Fig. 2. under the Head of the Tibia.

The Canal for the Medullary Vessels, slanting downwards at the inner and back part of the Bone, a little above its middle. Tab. II. above B.

The under End of the Tibia, smaller than the upper one, and its inferior Surface hollow, and covered with Cartilage, for the Articulation with the Astragalus. Tab. XXV. Fig. 3.

The Malleolus Internus, or inner Ankle, produced from the inner and fore part of the under end, and covered also with Cartilage where the Astragalus plays. Tab. XXV. Fig. 3. *r, m*.

A Pit in the point of the Malleolus Internus, for the attachment of the internal lateral Ligament, and a Groove behind, where the Tendon of the Tibialis Posticus is placed. Tab. XXV. Fig. 3. *m*.

The semicircular Cavity, at the under and outer side of the Tibia, for receiving the under end of the Fibula. Tab. XXV. Fig. 3. *q*. Tab. XXIII. Fig. 4. G.

Round the edge of the articulating Cavity, the Bone marked by the insertion of the Capsular Ligament. Tab. XXIII. Fig. 4. P.

The Tibia has a strong external Table, with a considerable quantity of spongy substance.

The Articulation of the upper end of the Tibia with the Os Femoris, is of such a nature as to allow flexion to a great degree, but the numerous Ligaments fixed here prevent it from being extended beyond a straight line with the Thigh; and then there is no rotation nor lateral motion, though, when the Joint is bent, the Ligaments are so much relaxed, that the Leg may be made in a small degree to roll or to turn a little to either side.

The Extremities of the Tibia are Cartilaginous in the Fœtus, and become afterwards Epiphyses. Tab. XXVII.

FIBULA,

Placed at the outer side of the Tibia, and by much the smaller of the two Bones, being the most slender Bone, in proportion to its length, of any in the Body. Tab. XXIX. Fig. 10. *i*.

The upper end of the Fibula, formed into a *large Head*, with a *superficial smooth Cavity* towards its inner side, to be articulated with the Tibia, where it is tied by Ligaments of such strength as to allow very little motion. Tab. XXV. Fig. 4. Tab. XXIII. Fig. 4.

The *Head* of the Fibula, *irregular* and *rough* externally, for the insertion of the Biceps Flexor Cruris, and of the external lateral Ligament of the Knee. Tab. XXIII. Fig. 4.

The *Body* of the Bone bent a little inwards and backwards, and unequally triangular, with the Surfaces between the Angles marked by the Muscles which arise from it, or are placed upon it. Tab. XXV. Fig. 4. Tab. XXIII. Fig. 4.

A *Ridge* at the inner side of the Fibula, opposed to one at the outer part of the Tibia, for the insertion of the Interosseous Ligament. Tab. XXIII. Fig. 4.

A *Canal* on the back-part of the Bone, slanting obliquely downwards, a little above its middle, for the passage of the Medullary Vessels. Tab. II. above C.

The *under End* of the Fibula, *broad* and *flat*, to be received by the femoral Cavity of the Tibia. Tab. XXIII. Fig. 4. above G.

The under end of the Bone forming the *Malleolus Externus*, or outer Ankle, which is lower and farther back than the inner Ankle, the obliquity of the two Malleoli in some measure corresponding with the obliquity of the Foot. Tab. II. *m*.

A *convex smooth Surface* on the inner side of the Malleolus Externus, opposed to the outer side of the Astragalus, which moves upon it.

The *Coronoid Process*, sent down from the Malleolus Externus, from which Ligaments go to the Bones at the outer side of the Foot. Tab. XXV. Fig. 4. *i*.

A *Furrow* upon the back-part of the Malleolus Externus, for lodging the Tendons of the Peronei.

The Fibula being articulated with the Tibia at its superior extremity by almost plain surfaces, and tied to it by strong and short Ligaments, only a very little motion is allowed.

At the under end it is joined so firmly by strong Ligaments, that no sensible motion appears in the Subject; though in this Joint, as in several others, where the Bones are firmly fixed by short Ligaments, there may be an elastic yielding in the living body.

In Old People, these two Bones are not unfrequently joined by an union of Substance.

The Fibula affords attachment to Muscles, assists in securing the Articulation of the Foot; adds to the form and strength of the Leg; and, by the head of the Bone being fixed to that of the Tibia, it widens the space for the Interosseous Ligament.

In the Fœtus, the extremities of the Fibula are Cartilaginous, and afterwards become Epiphyses, previous to being united to the Body of the Bone. Tab. XXVII.

PATELLA, vel ROTULA.

The Patella, placed at the fore-part of the Joint of the Knee, and in some respects bearing the same relation to the Tibia as the Olecranon does to the Ulna. Tab. I. B.

The *shape* of the Patella, *triangular* and *flat*, or of the figure of a Heart as painted upon playing-cards. Tab. XXVI. Fig. 1.

The *anterior Surface* of the Bone, *convex*, and perforated by numerous Holes, for the insertion of Tendons and Ligaments which cover it. Tab. XXVI. Fig. 1.

The *posterior Surface*, which corresponds with the Trochlea of the Os Femoris, smooth, covered with Cartilage, and divided by a longitudinal prominent Ridge into two unequal Cavities. Tab. XXVI. Fig. 2.

The circumference of the articular Surface, marked by a *rough Line*, into which the Capsular Ligament of the Joint is fixed. Tab. XXVI. Fig. 2. *a, a*.

The *Base*, or upper part of the Bone, *horizontal*, and marked by the insertion of the Tendons of the Extensors of the Leg. Tab. XXVI. Fig. 1. *a*.

The back-part of the Apex, *rough* and *depressed*, for the attachment of the Ligament which passes from the Patella to the Tubercle of the Tibia. Tab. XXVI. Fig. 2. *d*.

The Ligaments of the Patella allow it to be moved upwards and downwards; and when the Leg is extended, they admit of its motion to either side, or to be rolled.

When the Leg is extended, the Patella is lodged in the Trochlea of the Os Femoris; when the Limb is bent, the Patella is pulled down by the Tibia, and lodged in a hollow at the fore-part of the Knee.

The Patella has a thin, though firm external Table. Its internal Substance is cellular, but the Cells are small, and have so much Osseous Matter employed in their formation, as to give the Bone a considerable degree of strength.

The structure of this Bone, the toughness of the Ligaments which cover it, and the free motion it is allowed, are found to enable it better to resist any common force applied to it, than if it had been a process continued from the Tibia, as the Olecranon is from the Ulna.

The Patella defends the fore-part of the Knee, and serves as a pulley or lever, by enabling the Muscles fixed to it to act with greater advantage in extending the Leg.

It is entirely Cartilaginous at birth, and is later in ossifying than most of the Epiphyses.

FOOT,

Composed of *Tarsus*, *Metatarsus*, and *Toes*.

TARSUS,

TARSUS,

Composed of seven Bones, viz. The *Astragalus*, *Os Calcis*, *Naviculare*, *Cuboides*, *Cuneiforme Externum*, *Cuneiforme Medium*, and *Cuneiforme Internum*.

The upper part of the Tarsus *convex*, the under part *concave*.

In the *Concavity*, numerous *Muscles*, *Vessels*, and *Nerves* are lodged, belonging to the Sole.

The different Bones of the Tarsus have their *rough Surfaces joined together by strong Ligaments*, and their parts of Articulation *covered with Cartilage*, in such a manner as to form part of a strong and elastic Arch, for supporting the weight of the Body, and lessening the shock it would otherwise undergo in the different motions it has to sustain.

The ASTRAGALUS, placed under the Bones of the Leg. Tab. XXIII. Fig. 4. I.

The upper part of the Astragalus, formed into a *large Head*, in form of a Pulley, which is smooth on its upper part and sides, to be articulated with the under end of the Leg-bones. Tab. XXVI. Fig. 5. *a, a, b*.

Each of the Cartilaginous Surfaces of the Head of this Bone *depressed* in its middle, to correspond with the parts of the Leg-bones with which they are articulated.

Round the inferior edge of the articulating Surfaces, a *rough Fossa* for the insertion of the Capsular Ligament; and at the sides of this Surface, the Bone marked by the lateral Ligaments. Tab. XXVI. Fig. 5. *c, d, f*.

The under part of the Bone, consisting of a *deep Fossa*, or sinuous Cavity, which divides it into an anterior and posterior articulating Surface. Tab. XXVI. Fig. 7. *d, c*.

The Fossa in the under Surface, narrower at the inner part of the Bone, *d*, and becoming gradually wider as it goes outwards and forwards, *c*.

The *posterior articulating Surface*, large and *concave*, for its Articulation with the upper and middle part of the Os Calcis. Tab. XXVI. Fig. 7. *e, f*.

The *anterior articulating Surface*, irregular and *convex*, where it plays upon two smooth Cavities at the inner and fore part of the Os Calcis, and upon a Cartilaginous Ligament extended between the Os Calcis and Os Naviculare. Tab. XXIV. Fig. 7. *e, f*.

A *large oblong smooth Head*, at the fore-part of the Bone, for its Articulation with the Os Naviculare. Tab. XXVI. Fig. 7. *g*. Fig. 5. *e*.

The Joint between the Astragalus and Leg-bones forms a *complete Hinge*, which, together with the above-mentioned Ligaments, allows the Foot to bend and extend upon the Leg, but admits of no lateral nor rotatory motion, except in the extended state, when there is a little of each.

In a Fœtus, a considerable portion of this bone is ossified.

The OS CALCIS, the largest of the Tarsal Bones, situated under the Astragalus, and in the back-part of the Foot. Tab. XXVI. Fig. 5. B.

A *large rough Tuberosity* or Knob, projecting behind, to form the Heel, and to make one end of the Arch of the Foot. Tab. XXIII. Fig. 4. L.

A *superficial Cavity* in the upper and back part of this Knob, for the insertion of the *Tendo Achillis*. Tab. XXVI. Fig. 5. above *i*.

A *smooth Convexity* on the upper part of the Bone, for its Articulation with the under and back part of the Astragalus. Tab. XXVI. Fig. 8. *a*.

A *Fossa* or Sinuous Cavity at the fore-part of this articulating Surface, running forwards and outwards, and giving origin to strong Ligaments which are inserted into the corresponding Fossa of the Astragalus. Tab. XXVI. Fig. 8. *c, c*.

Two Prominences at the inner and fore part of the Bone, concave, and smooth above, with a pit between them, for the Articulation with the under and fore part of the Astragalus. Tab. XXVI. Fig. 8. *b, b*.

From the posterior Prominence the Cartilaginous Ligament arises, which is fixed to the Os Naviculare.

A *large Cavity* or Arch at the inner side of the Bone, between the posterior of the two last-mentioned Processes and Projection of the Heel, for lodging the Tendons of the long Flexors of the Toes, together with the Vessels and Nerves of the Sole. Tab. XXVI. Fig. 5. B.

A *Depression* in the external Surface of the Bone, near its fore-part, where the Tendon of the Peroneus Longus runs in its way to the Sole. Tab. XXVI. Fig. 8. *f*.

The under and back part of the Bone, forming *two Prominences*, where it gives origin to the Aponeurosis and to several Muscles of the Sole; and before the Prominences, the Bone *concave*, where it lodges part of these Muscles. Tab. XXXII. Fig. 10. *a, b*.

The anterior Surface *concave*, and somewhat in form of a Pulley placed obliquely, for its articulation with the Os Cuboides. Tab. XXIII. Fig. 4. before K.

The *Os Calcis* is articulated with the Astragalus by Ligaments of such strength, that this part of the Foot, upon which the Body rests, is rendered firm and secure, but enjoys very little motion.

In a Fœtus, a large proportion of this Bone is ossified, and the Projection forming the Heel is afterwards an Epiphysis.

The OS NAVICULARE, situated at the fore-part of the Astragalus, and inner part of the Foot. Tab. XXVI. Fig. 5. C.

The *posterior Surface*, forming a Cavity somewhat like that of a Boat, for receiving the head of the Astragalus in the manner of Ball and Socket. Tab. XXVI. Fig. 6. *c*.

A *Prominence* at the inner side of the Bone, for the insertion

fertion of Tendons, Muscles, and strong Ligaments, particularly for the Ligament stretched between this Bone and the Os Calcis, for the support of the Astragalus. Tab. XXVI. Fig. 5. o.

The *fore-part* of the Bone *convex*, and divided into *three articular Surfaces*, for the articulation with the Ossa Cuneiformia. Tab. XXVI. Fig. 5. p, p, p.

Between the Os Naviculare and Astragalus the foot has its principal lateral and rotatory motions, though each of the other Joints of the Tarsus contributes a little.

The Os CUBOIDES, placed at the fore and outer part of the Tarsus. Tab. XXXII. Fig. 12. i.

The *posterior Surface* of this Bone *smooth, convex* at its *inner*, and *concave* at its *outer* part, corresponding with the anterior extremity of the Os Calcis. Tab. XXVI. Fig. 12. b.

The *inner Side*, articulated with the Os Naviculare and external Os Cuneiforme. Tab. XXIII. Fig. 4. P.

Its *under Surface*, *irregular* where it gives attachment to strong Ligaments, and to the Adductor Pollicis. Tab. XXXII. Fig. 10. k.

A *deep Fossa* in the outer and under part of the Bone, for lodging the Tendon of the Peroneus Longus, where it crosses the Sole. Tab. XXXII. Fig. 10. l.

The *anterior* extremity, divided into a small inner, and large outer plain Surface, to be articulated with the fourth and fifth Metatarsal Bones. Tab. XXXII. Fig. 10.

The Three OSSA CUNEIFORMIA, situated at the fore-part of the Tarsus, and inner side of the Os Cuboides, and applied to each other like the stones of an Arch. Tab. XXXII. Fig. 12. n, o, p.

The *upper part* of these Bones, *flat* where they are covered with Ligaments. Tab. XXXII. Fig. 12.

The *under part irregular*, for the attachment of Muscles and strong Ligaments lying in the Sole. Tab. XXXII. Fig. 10. n, o, p.

The *posterior Surface, flat*, and covered with Cartilage, to be articulated with the Os Naviculare. Tab. XXXII. Fig. 12.

The *anterior Surface* also flat, for the articulation with the Metatarsal Bones. Tab. XXXIV. Fig. 12.

The Os Cuneiforme Externum, or Medium, of a middle size between the next two Bones, and opposed to the Metatarsal Bone of the Third Toe.—The outer side of this Bone articulated with the Os Cuboides. Tab. XXVI. Fig. 9. Tab. XXXII. Fig. 12. p.

Os Cuneiforme Medium, or Minimum, the least of the three, and articulated at its outside with the former Bone, and anteriorly with the second Metatarsal Bone. Tab. XXVI. Fig. 10. Tab. XXXII. Fig. 12. o.

The Os Cuneiforme Internum, or Maximum, the largest of the Cuneiform Bones, and placed obliquely, with its anterior Surface opposed to the Metatarsal Bone of the

great Toe. Tab. XXVI. Fig. 11. Tab. XXXII. Fig. 12. n.

The sharp Edge of this Bone turned upwards, while that of the other two is in the opposite direction. Tab. XXXII. Fig. 12. n.

The Os Naviculare, Os Cuboides, and Ossa Cuneiformia, are almost Cartilaginous at birth.

METATARSUS,

Composed of *five Bones*, which answer to the general characters given to the Metacarpal Bones. Tab. XXIII. Fig. 4. Q; Q, &c.

Their *bodies long, arched upwards*, and *tapering* towards their anterior extremities. Tab. XXXII. Fig. 12.

The *extremities, large* in proportion to their Bodies, and the posterior much larger than the anterior. Tab. XXXII. Fig. 10.

The *Bases, flat*, or a very little *hollowed*, to be articulated with the fore-part of the Tarsal Bones. Tab. XXXII. Fig. 12.

From the flatness of their Bases, and the strength of the Ligaments which fix these Bones to those of the Tarsus, very little motion is allowed to this part of the Foot.

Round the Bases, *rough Surfaces* for the attachment of Ligaments. Tab. XXIII. Fig. 4.

The *Sides of the Bases flat* where they are articulated with each other. Tab. XXXII. Fig. 12.

A *Ridge* above, and a *flat Surface* at each side of their bodies, for the origin of the Interosseous Muscles. Tab. XXXII. Fig. 12.

The *flat Surfaces* turned obliquely outwards, and the obliquity increasing the more externally the Bones are placed.

The *anterior Extremities* forming *Balls*, to be articulated with the Toes;—the Balls much longer from above downwards, than from one side to the other. Tab. XXXII. Fig. 12.

Round the Heads *distinct impressions*, where the Capsular Ligaments are fixed. Tab. XXXII. Fig. 12.

The *Metatarsal Bone of the Great Toe*, by much the thickest and strongest, but shortest of the Metatarsus. Tab. XXXII. Fig. 12.

The *articulating Cavity* of its Base, deeper than the rest. Tab. XXXII. Fig. 12.

The *anterior Extremity*, bearing a greater proportion to the Base than the rest, having a much larger share of the weight of the Body to sustain here, and formed into a *middle Prominence*, with *two lateral Depressions*, where the Bones termed Ossa Sesamoidea move. Tab. XXIV. Fig. 2.

The *Metatarsal Bone of the second Toe*, the longest of the five. Tab. XXXII. Fig. 12.

The *Metatarsal Bone of the Middle Toe*, the second in length, with a Base like that of the former Bone, triangular, but a little larger, to be articulated with the Os Cuneiforme Externum. Tab. XXXII. Fig. 12.

The

The *Metatarsal Bone of the Fourth Toe*, nearly of the same length as the former, but distinguished from it by its Base being thicker below, and its Cartilaginous Surface being more of a square form, corresponding with the anterior and inner part of the Os Cuboides, with which it is articulated. Tab. XXXII. Fig. 10.

The *Metatarsal Bone of the Little Toe* the shortest, with flat Surfaces facing upwards and downwards. Tab. XXXII. Fig. 10. 12.

The *Base* which rests on the Os Cuboides, projecting outwardly into a large Tuberosity, which gives origin to Muscles, and forms one of the points on which the Body rests in standing. Tab. XXXII. Fig. 10.

The Bones of the Metatarsus, with those of the Tarsus, form an irregular Arch for supporting the Body, one end of the Arch being formed by the projection of the Heel, the other by the anterior extremity of the Metatarsal Bones. The different pieces composing this Arch are bound by Ligaments of such strength, as to give security to the whole.

TOES.

The *Bones of the Toes*, the same in number with those of the Fingers, viz. two to the Great Toe, and three to each of the smaller Toes, and the different Bones here, as in the Fingers, disposed in Ranks or Phalanges. Tab. XXXII. Fig. 10. 12.

The *two Bones of the Great Toe* like those of the Thumb, but stronger, and placed in the same row with the Bones of the Toes, for the purpose of walking, and assisting in supporting the Body. Tab. XXXII. Fig. 12.

The *Bones of the smaller Toes* every way less than those of the Fingers. Tab. XXIV.

Their *under Surface depressed*, where the Tendons of their Flexor Muscles are lodged. Tab. XXXII. Fig. 10.

The *Bases* of the first Phalanx, as in the Fingers, forming Sockets to receive the Balls, or Heads of the Metatarsal Bones. Tab. XXXII. Fig. 12.

The *Joints* between the first and second Phalanx, and also between the second and third, as in the Fingers,

forming *Hinges*, and the motions similar, but more confined. Tab. XXXII. Fig. 10. 12.

Of the small Toes, the *first*, or that next the Great Toe, the *largest*, the rest becoming *smaller* the more externally they are placed. Tab. XXXII. Fig. 10. 12.

The second and third Bones, especially of the little Toe, frequently joined by an union of Substance.

The Bones of the Toes allow a free and easy motion in Children, and a considerable degree of it also in People whose feet have not been confined in shoes. In others, especially in advanced life, the Toes are frequently found squeezed together, and some of the smallest Bones of the Toes, as the two last of the little one, having the pieces which originally composed them joined together by an union of Substance.

The structure of the Bones of the Foot is nearly similar to that of the Bones of the Hand.

In the Fœtus, the Bones of the Metatarsus and Toes are in the same condition as those of the Metacarpus and Fingers.

OSSA SESAMOIDEA.

THEIR size, situation, and number, vary in different persons.

They are sometimes found at the roots of the Fingers and small Toes; at the second Joint of the Thumb, and at that of the Great Toe; between the Condyles of the Os Femoris and Gastrocnemius Muscle; between the Tendons of the Peroneus Longus and Os Cuboides, &c.

Those commonly observed are placed in pairs at the roots of the Thumb and great Toe, between the Tendons of their Flexor Muscles and Joints.

They are *convex* on their outer Surface, where they are inclosed by the Tendons and Muscles fixed to them. Tab. XXXII. Fig. 12. a.

And *concave*, and lined with Cartilage next the Joints, where they play upon the Bones with which they are articulated. Tab. XXXII. Fig. 14. a. Tab. XXVI. Fig. 3. 4.

They are considered by Anatomists as serving the same general purpose with the Patella.

TABLE

T A B L E XXV.

Different VIEWS of the OS FEMORIS, TIBIA, and FIBULA.

FIG. 1.

Anterior Surface of the OS FEMORIS of the LEFT SIDE.

- a*, THE head of the os femoris, covered with a smooth and polished cartilage.
b, A portion of the pit, or ligamentous impresson of this head.
c, The upper part of the neck of the os femoris.
d, d, Various openings or fissures, which give passage to vessels.
e, The trochanter major.
f, The blunt point of this process.
g, g, The ridge, or projecting line which extends from the greater to the smaller trochanter.
h, The trochanter minor.
i, The upper and middle part of the os femoris, somewhat flattened.
k, The middle part, which is convex and rounded.
l, The inferior and middle part, more of a triangular form.
m, m, Muscular and ligamentous impressions upon the lateral and under parts of the bone.
n, A triangular cavity, with fissures for the insertion of the capsular ligament, and for the passage of vessels.
o, A cavity, or pulley, at the bottom of the os femoris, covered with cartilage, to facilitate the motion of the patella.
p, p, The eminences which form the sides of the pulley.

- b*, The base of this process.
i, i, Openings in this process for the transmission of vessels.
k, k, The ragged eminence, or crest, which extends from one trochanter to the other.
l, The trochanter minor, on which, as well as on the trochanter major, are a number of muscular prints.
m, The middle of the inner surface of the os femoris.
n, The middle of the outer surface.
o, The middle of the linea aspera, which, through its whole length, is only a continuation of muscular impressions.
p, p, The division of this line into two branches, of which one goes to the larger, and the other to the smaller trochanter.
q, Orifice of the canal for the medullary vessels, in the middle and inner part of the os femoris.
r, r, Division of the linea aspera into two small ridges, which extend from the middle and under part of the bone as far as its condyles.
s, The triangular cavity between these two branches and the condyles, for facilitating the passage of blood-vessels.
t, t, The fissures at the under part of this cavity, through which the vessels of the cancelli pass.
v, v, The condyles, incrusted with a smooth cartilage.
x, x, The tuberosities of the condyles, into which ligaments and muscles are inserted.
y, A cavity between the condyles.
z, z, Ligamentous impressions upon the upper edge of this cavity.

FIG. 2.

The Posterior Surface of the same BONE.

- a*, The posterior part of the head of the os femoris.
b, The cavity, or ligamentous print of the head of the os femoris.
c, c, The unequal edge of the cartilaginous substance of the head.
d, The upper and posterior part of the cervix femoris.
e, The under part of the cervix.
f, The openings in the cervix, for the transmission of vessels, which penetrate the substance of the bone.
g, The blunt point of the trochanter major, on which are muscular prints.

FIG. 3.

The Anterior Surface of the TIBIA of the LEFT SIDE.

- a*, The middle of the crest of the tibia.
b, b, The upper and under parts of the crest or ridge.
c, c, The inner smooth surface.
d, d, The outer surface, generally hollow.
e, e, The edge of the superior cavities of the tibia.
f, f, The tuberosity of this bone divided into two parts, of which the upper gives attachment to the ligament of the patella, and the other insertion to the tendons of muscles.

g, The

TAB. 25.

Fig. 2.




Fig. 3.




Fig. 4.



Fig. 1.



- g*, The small articular process, which answers to the fibula.
b, b, Prints made by the capsular ligament on the upper part of the tibia.
i, i, i, Openings of the spongy substance, which are orifices for the transmission of vessels spread over this part of the bone.
k, k, A porous surface, where the tendons of muscles, with their aponeurosis, are fixed.
l, The under and middle round part of the tibia.
m, The malleolus internus.
n, n, Vestiges of the union of the inferior process to the body of the bone.
o, The eminence which answers to the fibula.
p, p, The articular cavities which correspond with the astragalus.
q, Part of the articular cavity which receives the fibula.
r, r, Ligamentous impressions.

FIG. 4.

The External Surface of the FIBULA of the LEFT SIDE.

- a, a, a*, The length of its external surface.
b, Part of the posterior surface.
c, c, The ridge, or osseous line which separates the posterior from the outer surface.
d, Part of the ridge which separates the outer from the inner surface.
e, The superior process of the fibula.
f, The articular cavity which receives the eminence of the tibia.
g, The articular process of the fibula, which corresponds to the tibia.
h, h, Asperities, or ligamentous and muscular prints.
i, Coronoid process of the malleolus externus.
k, k, The asperities, or ligamentous prints of this process.
l, l, The orifices of several conduits, for the transmission of vessels.

T A B L E XXVI.

VIEWS of the PATELLA, OSSA SESAMOIDEA, and BONES of the FOOT.

FIG. 1.

The External Surface of the PATELLA of the LEFT SIDE.

- a*, A hollow in the upper part of the patella, into which the tendon of the extensor muscles of the leg is fixed.
b, The middle of the bone, somewhat convex.
c, c, The lateral parts, which are so many muscular prints.
d, Fissures on the surface of the patella, with the orifices of the conduits by which the vessels penetrate into the interior of the bone.
e, The inferior extremity of the patella, into which a very strong ligament is fixed.

FIG. 2.

The Inner Surface of the same PATELLA.

- a, a, a*, The circumference of the patella, or margin to which the capsular ligament is fixed.
b, b, The articular cavities of the inner surface, covered with smooth cartilage.
c, The small prominent ridge, or superficial process, which separates the two cavities.
d, An irregular hollow, to which the ligament is fixed which goes to the tibia.

FIG. 3.

One of the Sesamoid Bones of the Thumb.

FIG. 4.

A Sesamoid Bone of the Great Toe.

FIG. 5.

The Inner Surface of the LEFT FOOT, in a Position nearly horizontal.

- A*, The middle and upper part of the astragalus, where it is somewhat concave, and covered with cartilage.
a, a, The semicircular eminences which border the upper part of this bone.

- b*, The oblong cavity of the inner surface, by which it is joined with the malleolus internus.
c, The cervix of the astragalus.
d, Inequalities, or ligamentous prints on the upper and inner part.
e, The head of the astragalus, which is received into the cavity of the os scaphoides.
f, Inequalities on the inner surface, also marked with ligamentous prints.
g, A small eminence on the posterior part of the astragalus, which is articulated with the os calcis.
B, The middle of the inner surface of the os calcis, excavated, in form of an oblique gutter, for the passage of tendons, vessels, and nerves.
b, The upper and posterior eminence of the os calcis, by which it is articulated with the astragalus.
i, The tuberosity of this bone, the upper and back part of which is impressed by the insertion of the tendo Achillis.
k, The inner and upper surface of the os calcis, articulated with a small surface of the astragalus.
l, l, The sinuosity below this eminence, through which the tendon of the flexor longus pollicis passes.
m, The anterior eminence of the os calcis, by which it is joined with the os cuboides.
C, The upper and middle part of the os naviculare, on which several ligamentous prints are seen.
n, n, The navicular cavity, which receives the head of the astragalus.
o, The tuberosity of the os naviculare, to which the tendon of the tibialis posticus is fixed.
p, p, p, The small surfaces by which this bone is joined to each of the cuneiform bones.
D, The middle of the large cuneiform bone.
q, The under end of this bone, where there is a small surface anteriorly, to which the tendon of the tibialis anticus is fixed,—and posteriorly, a tuberosity to which the abductor pollicis is fixed.
r, The upper part of the same bone, by which it is joined with the second os cuneiforme.
E, The upper part of the second cuneiform bone.

F, A

TAB. 26.

Fig. 1.



Fig. 2.



Fig. 4.



Fig. 3.



Fig. 5.




Fig. 7.



Fig. 8.



Fig. 6.



Fig. 9.



Fig. 10.





Fig. 11.



Fig. 12.



- F, A small part of the third cuneiform bone.
 G, The middle and upper part of the large metatarsal bone.
 s, The posterior part of this bone, where it is joined with the first os cuneiforme.
 t, The head of the same bone, which forms an articular articulation with the first phalanx of the great toe.
 u, u, Eminences, or prints, on the lateral parts of this bone.
 H, The middle and upper part of the second metatarsal bone.
 v, The posterior and upper part of this bone.
 w, Its anterior extremity, which terminates in a roundish head.
 I, The upper and middle part of the first bone of the great toe.
 x, The edge of the glenoid cavity of this bone.
 y, The anterior extremity, the articulation of which with the second bone is a complete ginglymus.
 K, Part of the first bone of the second toe.
 L, A portion of the first bone of the third toe.
 M, The middle part of the second bone of the great toe, which is convex.
 z, The upper and posterior part of this bone, by which it is joined to the first.
 Q, The anterior extremity of the second bone, on which are asperities to support the nails, and furnish attachment to tendinous fibres.
 N, N, Portions of the second bones of the second and third toes.
 O, A very small portion of the third metatarsal bone.

FIG. 6.

The Posterior Surface of the Os SCAPHOIDES, seen a little obliquely.

- a, a, a, A large part of the upper surface of the os scaphoides, which is very irregular, and gives attachment to several ligaments.
 b, A portion of the tuberosity of this bone, on which are muscular and ligamentous prints.
 c, The navicular cavity which receives the head of the astragalus.

FIG. 7.

The Under Part of the ASTRAGALUS.

- a, The outer surface of the astragalus, seen a little fore-shortened.
 b, Its great cavity, which corresponds to the large upper eminence of the os calcis.
 c, Edge of the large fossa of the astragalus.

- d, d, The bottom of this fossa, on which are several ligamentous prints.
 e, A small oval surface, which is joined with a part of the oblong cavity on the eminence of the inner surface of the os calcis.
 f, Another small eminence, which is joined with another part of the same cavity of the os calcis.
 g, The under part of the head of the astragalus.

FIG. 8.

Shews the Upper Part, and External Surface of the Os CALCIS.

- a, The upper eminence of the middle part of the os calcis, which is articulated with the great cavity of the astragalus.
 b, b, Another eminence of the inner surface of this bone, in which there is an oblong cavity, which is articulated with another part of the astragalus.
 c, c, Irregular hollows, into which the principal ligaments which unite the astragalus with the os calcis are inserted.
 d, d, d, The posterior part, or edge of the large tuberosity of the os calcis.
 e, The edge of the large hollow of this bone.
 f, The external sinuosity, through which the tendon of the peroneus longus passes.
 g, The small tuberosity to which part of the ligaments of the peronei are fixed.
 h, Anterior process of the os calcis, by which it is articulated with the os cuboides.
 i, The middle external part, which is slightly convex.

FIG. 9.

The Upper and Outer Surface of the External Os CUNEIFORME.

- a, The upper surface, to which several small ligaments are fixed.
 b, A portion of the anterior surface.
 c, That part of its outer surface which is joined to the cuboides.
 d, The depressions of this surface, to which strong ligaments are fixed.
 e, The point of the bone which answers to the sole.

FIG. 10.

The Upper, Posterior, and Outer Surfaces of the Middle Os CUNEIFORME.

- a, The upper surface, on which are asperities, or ligamentous prints.

b, Par.

- b*, Part of the posterior surface.
- c*, The outer surface, on which is an eminence joined to the third cuneiform bone.
- d*, The under part, which answers to the sole.

FIG. 11.

Part of the Inferior, and Part of the Anterior Surface of the Internal Os CUNEIFORME.


- a, a*, The small surface, by which it is articulated with the second bone of the same name.
- b*, A portion of the under part of the large cuneiform bone, which terminates in a thick tuberosity, to which the tendon of the tibialis anticus is fixed.
- c*, The anterior surface, a little hollow, for the adaptation of this with the large metatarsal bone.

FIG. 12.

The Superior, and somewhat External Surface of the Os CUBOIDES.

- a*, A part of the inner edge of this bone, which is joined to the third cuneiform bone.
- b*, The posterior surface, on which there is an oblique process, adapted to the os calcis, with which it is articulated.
- c*, The anterior surface, which receives the two last metatarsal bones.
- d*, The upper surface, the numerous asperities of which are so many ligamentous prints.
- e*, Part of the inferior oblique process.
- f*, A portion of the sinuosity, through which the tendon of the peroneus longus passes.

TABLE



T A B L E XXVII.

Represents the SKELETON of a FOETUS at the Full Time.

IN this figure are seen the following, among other peculiarities of structure ; viz.

A portion of the fontanella.

A membranous substance, in form of a future, uniting the two pieces which form the frontal bone.

A ring of bone surrounding the outer edge of the tympanum.

The symphysis of the lower jaw, formed of cartilage.

The os sacrum, composed of distinct vertebræ, with intervertebral substances.

The different points of ossification upon the sternum.

The cartilaginous border of the os ilium.

The cartilaginous union of the three pieces which compose the os innominatum.

The ends of the long bones in general of the superior extremities, in the state of epiphyses.

The bones of the carpus cartilaginous.

The ends of the long bones of the inferior extremities, in the form of epiphyses.

The patellæ in a state of cartilage.

The bones of the tarsus partly cartilaginous, and partly osseous.

TABLE

T A B L E X X V I I I .

Represents a well-formed YOUNG ADULT FEMALE SKELETON, the different Parts of which may be understood, by comparing it with the SKELETONS already described.

PRINCIPAL DIFFERENCES BETWEEN THE MALE AND FEMALE SKELETON.

The greater part of which may be observed in Tab. I. II. XXIX. XXXI. representing the Male Skeleton, and Tab. XXVIII. XXX. the Female Skeleton. See also Tab. XV. XVI. XVII. XX.

THE Female Skeleton is observed, in general, to be smaller and more slender throughout than that of the Male.

A ripe Female Bone, of the same size with a Male Bone, is usually distinguished by the Ridges, Depressions, rough Surfaces, and other Inequalities, being less conspicuous in the former.

The Circumference of the Female Skull is said by Soemmering to be larger.

The Os Frontis is found to be more frequently divided by a continuation of the Sagittal Suture.

The Frontal Sinuses are observed to be narrower ;

All the Bones of the Face more delicate ;

The Bodies of the Vertebrae longer, and the Vertebral Canal, according to the Author quoted above, larger ;

The Intervertebral Substances deeper or thicker ;

The upper part of the Thorax in proportion wider ;

The under part narrower, or the whole Thorax less conical ;

The Cartilages of the True Ribs longer in proportion to the Ossous part, and broader and flatter to support the Breasts ;

The Sternum more raised, and the whole Thorax more distant from the Pelvis ;

The length of the Sternum less, and terminating below on a line nearly opposite to the plane of the fourth pair

of Ribs, but in the Male Skeleton terminating opposite to that of the fifth pair ;

The Cartilago Eniformis oftener perforated in the middle, or Bifurcated ;

The length of the Loins greater ;

All the Diameters of the Pelvis larger ;

The Spines and Processes of the Ossa Innominata farther distant from each other ;

The Os Sacrum, broader, and turned more backwards, to enlarge the Cavity of the Pelvis ;

The Os Coccygis, more slender, turned more backwards, and having a greater degree of motion ;

The Ossa Iliæ, flatter, and more reflected outwards, by which the under part of the Abdomen is rendered more capacious ;

The Notches of the Ossa Iliæ wider, and the conjoined Surfaces of the Ossa Innominata and Os Sacrum less ;

The space between the Ossa Pubis larger ; of course the Ligamentous Cartilage of the Symphysis broader, though shorter ;

The Angle formed by the Crura of the Ossa Pubis with the Symphysis Pubis larger ; that of the Male being acute, while in the Female the Angle extends to 80 or 90 degrees ;

The Tuberosities of the Ossa Ischia, flatter, and at a greater distance from each other ;

The



The Brim of the Pelvis, wider, and of an oval form, corresponding with the Head of a Child, and the longest Diameter extending between the Ossa Iliæ ;

In the Male, the Brim of the Pelvis observed to have more of a circular appearance, and to have the greatest extent between the Ossa Pubis and Os Sacrum ;

The opening at the under part of the Pelvis, in the Female, much wider, and of an oval form ; but the oval the reverse of that at the Brim ;

The Foramina Ovalia wider ;

All the Openings at the under part of the Pelvis, being wider, leave a large passage for the Birth of the Child ;

The Acetabula farther distant from each other, in

consequence of which, Women who are very broad at this part of the Body waggle when they walk ;

The Ossa Femorum more curved : The Neck of the Thigh-bone forming a greater Angle with its Body : The Body of the Thigh-bone placed more obliquely : The internal Condyle larger ;

The Feet smaller ;

The Clavicles less crooked ;

The Scapulæ smaller, and their Angles more acute ;

The Superior Extremities shorter ;

The Ossa Carpi narrower ; and,

The Fingers more tapering towards their Extremities.

TABLE

T A B L E XXIX.

Represents the General Structure of the BONES, and a Front View of the MALE SKELETON.

FIG. 1.

Part of the Os FRONTIS, covered with its PERIOSTEUM, the Arteries of which are Injected.

A, The branches which come from the orbit;—the other small trunks of the arteries observed in several places are sent from the common integuments to the periosteum.

FIG. 2.

The Parietal Bone of a Fetus, to shew the Radiated Fibres of a Flat Bone, proceeding from the first ossified point A.

FIG. 3.

The THIGH-BONE of a Fetus, to shew the Longitudinal Parallel Fibres of a CYLINDRICAL BONE.

A, The part which first ossifies.

B, B, The two extremities in a cartilaginous state.

FIG. 4.

Section of Part of the Os FEMORIS, to shew the PLATES and CANCELLI of Long BONES in general.

A, A, A, A, The plates of the thigh-bone separated.

B, B, The cancelli.

FIG. 5.

Transverse Section of a Bone burnt, to shew the Cavities for containing the Marrow and Vessels.

FIG. 6.

The Appearance of the Marrow, viewed with a Microscope.

FIG. 7.

The THIGH-BONE, sawed Longitudinally through the middle.

A, A, A, A, The cancelli.

B, B, The union of the bone with its extremities, which are here in a state of epiphyfis.

C, C, The reticular substance.

D, D, The sides, or tables, which are thick and strong in the middle of the bone, and thinner towards the extremities.

FIG. 8.

The Os Ilium, sawed through the middle to shew the Cancelli.

FIG. 9.

The HIP-JOINT of a CHILD, opened to shew,

A, The head of the thigh-bone.

B, The round ligament connecting it to the acetabulum.

C, The capsular ligament of the joint, with its arteries injected, and,

D, The numerous vessels of the fatty glandular-like substance of the joint also injected.

FIG. 10.

A Front View of the MALE SKELETON,

A, The frontal bone.

δ, Its superciliary hole.

β, The external orbital process.

γ, The internal orbital process.

B, The parietal bone.

Between A and B, the coronal future.

C, The temporal bone.

D, ε, The occipital bone.

E, The bones of the nose.

F, The os malæ.

G, The superior maxillary bone.

H, The lower jaw.

I, The teeth.

K, The seven cervical vertebræ, with their intermediate cartilages.

ζ, Their transverse processes.

L, L, &c. The twelve dorsal vertebræ.

M, The five lumbar vertebræ.

N, Their intermediate cartilages.

κ, Their transverse processes.

O, The

FIG. I.




FIG. IO.




FIG. 2.





FIG. 6.



FIG. 5.





FIG. 7.




FIG. 4.



FIG. 3.




FIG. 8.





FIG. 9.



A. Cooper fecit

- O, The os sacrum.
 P, The os coccygis.
 Q, The os ilium.
 λ, Its spine.
 μ, Its anterior-superior spinous process.
 ν, Its inferior-anterior spinous process.
 ξ, The venter of the ilium.
 ο, The brim of the pelvis.
 R, The os pubis.
 S, The os ischium.
 π, The crura of the ossa ischia.
 ς, The foramen thyroideum.
 ϕ, The acetabulum.
 T, The seven true ribs.
 U, The five false ribs.
 η, η, The joining of the ribs with their cartilages.
 θ, The cartilages of the 6th, 7th, and 8th ribs, united with each other.
 V, The sternum.
 1. Its upper piece.
 2. Its middle piece.
 3. 4. Its cartilago ensiformis.
 X, The clavicle.
 Y, The concave surface of the scapula of the right side.
 φ, The superior costa of the scapula, with its femilunar notch.
 τ, The coracoid process of the left side.
 υ, The acromion of the left side.
 ϑ, The anterior inferior costa of the scapula of the right side.
- ζ, The head of the os humeri under the acromion.
 ψ, A groove for the tendon of the biceps.
 Z, The body of the os humeri.
 α, The trochlea.
 α, The external,
 β, The internal condyle of the os humeri.
 δ, The head of the radius of the left side.
 ε, The olecranon of the ulna of the same side.
 α, The ulna of the right side.
 β, The radius.
 c, The carpus.
 d, The metacarpus.
 e, The phalanges of the fingers.
 f, The right os femoris.
 λ, The internal,
 l, The external condyle,
 g, The patella.
 h, The tibia.
 i, The fibula.
 k, The tarsus.
 l, The metatarsus.
 m, The phalanges of the toes.
 r, The malleolus externus.
 t, The malleolus internus.
 f, The ball of the left thigh-bone.
 g, The great trochanter.
 b, The cervix.
 i, The small trochanter.
 p, The tubercle of the tibia.
 q, The os calcis.

T A B L E XXX.

Represents a Front-View of the FEMALE SKELETON, with the BONES of the HEAD.

FIG. 1.

By comparing this Figure with Fig. 10. of the former Table, the different Proportions of the Bones of the two Sexes are seen, and the Letters to the several Bones of the MALE SKELETON, explained in Fig. 10. may guide the Eye to the like Bones of the FEMALE SKELETON represented here.

THE Letters added here to the BONES of the HEAD, are,

- a, The coronal future.
- b, The squamous future.
- o, The lambdoid future.
- c, c, The transverse future.
- d, The zygomatic future.
- e, The external orbitar future.
- f, The lateral nasal future.
- δ, The superciliary hole of the frontal bone.
- ξ, The orbitar process of the frontal bone.
- H, The os planum of the ethmoid bone.
- r, The lacrymal groove of the os unguis.
- π, The external orbitar hole of the maxillary bone.
- ϕ, The tuber of the maxillary bone.
- 1. The chin.
- 2. The base of the lower jaw.
- 3. Its angle.
- 4. Its coronoid process.
- 5. Its condyle.
- 6. The mental hole.

SUPERIOR EXTREMITIES.

- 5. The right os humeri.
- 6. The head of the radius.
- 7. The olecranon.
- 8. The ulna.
- 9. The under end of the radius, marked by muscles.
- τ, The coracoid process of the scapula of the left side.
- υ, The acromion of the scapula.
- φ, The femilunar notch on the upper edge of the scapula.
- 10. The coronoid process of the ulna.
- 11. The tubercle of the radius.
- 12. The under end of the radius.
- 13. Its styloid process.
- 14. The styloid process of the ulna.

PELVIS, AND INFERIOR EXTREMITY.

- λ, The spine of the os ilium.
- μ, Its anterior-superior spinous process.
- ν, Its anterior-inferior spinous process.
- R, The joining of the os ilium and os pubis.

- S, The os ischium.
- 1. The spinous process of the os ischium.
- 2. The joining of the os sacrum with the os ilium.
- 3. The symphysis pubis.
- 4. The pelvis.
- f, The ball of the thigh-bone.
- g, The trochanter major.
- b, The cervix of the thigh-bone.
- 15. The head of the fibula.
- 16. The spine, and,
- 17. The inner edge of the tibia.
- 18. The under end of that bone.
- r, The malleolus externus.
- t, The malleolus internus.

FIG. 2.

Back-View of the FRONTAL BONE.

- η, The frontal sinus.
 - ι, The sagittal future; continued in this figure to the root of the nose.
- The other parts seen in this View have been described in Tab. VIII. Fig. 4.

FIG. 3.

The Inner Side of the Left Parietal Bone.—See Tab. IX. Fig. 2.

FIG. 4.

A Fore-View of the OCCIPITAL BONE.—See Tab. IX. Fig. 4.
To which add here,

- π, An os triquetrum.
- ξ, The extremity of the cuneiform process, where it joins the sphenoid bone.
- σ, The exterior surface of the cuneiform process.
- τ, τ, The condyles.
- α, Part of the hole common to the occipital bone, and right temporal bone.
- β, The hole for one of the nerves of the ninth pair.

FIG. 5.

The Inner Side of the Right Temporal Bone.—See Tab. X. Fig. 6.

FIG. 6.

Internal View of the SPHENOID BONE.—See Tab. X. Fig. 2.
To which add here,

- 3. A connection which is sometimes observed between the anterior and posterior clinoid processes.

FIG.

FIG. 15.



FIG. 1.





FIG. 16.



TAB. 30.

FIG. 2.



FIG. 3.



FIG. 13.




FIG. 4.




FIG. II.






FIG. 5.

FIG. 6.



FIG. 7.



FIG. 8.




FIG. 9.




FIG. 10.



FIG. 11.



FIG. 7.

Exterior View of the Ethmoid Bone.—See Tab. X. Fig. 3.

FIG. 8.

Posterior View of the two Nasal Bones.—See Tab. XII. Fig. 3. 4.

FIG. 9.

The Side of the Os Unguis next the Nose.—See Tab. XII. Fig. 6.

FIG. 10.

Posterior View of the Os Mala.—See Tab. XII. Fig. 8.

FIG. 11.

A View of the Lower Part, and Side next the Nose, of the Left Os Maxillare, with the Palate Bone, and α , the Os Turbinatum Inferius.—See Tab. XII. Fig. 2.

FIG. 12.

The Left PALATE BONE inverted.

A, B, C, D, The palate plate.

The other letters refer to parts belonging to the nose and orbit.—See Tab. XII. Fig. 11. 12.

FIG. 13.

A View of the Inner SURFACE of the Right Side of the LOWER JAW.

1. A section of the chin.
2. The base of the jaw.
3. The angle.
4. The coronoid process.
5. The condyle.
6. The rough print of the internal pterygoid muscle.
7. The orifice of the passage for the nerve and blood-vessels.
8. The five molares.

FIG. 14.

A TOOTH cut perpendicularly, magnified.

- A, The fibres of the cortical part.
 B, The bony part.
 C, The entry for the vessels and nerve.
 D, The cavity of the tooth.

FIG. 15.

A View of the Interior Surface of the Base of the Skull.—See Tab. V.

FIG. 16.

Interior View of the Right Side of the CRANIUM, and BONES of the UPPER JAW.

- D, D, The two tables and diploe of the frontal and occipital bones.
 a, The coronal future.
 α , The serrated edges of the parietal bone, for forming the sagittal future.
 o, The lambdoid future.
 b, The squamous future.
 ξ , The furrows made by the vessels of the dura mater.
 η , The frontal sinus.
 E, The crista Galli.
 F, The nasal lamella of the ethmoid bone.
 1. The hollow wing of the sphenoid bone.
 6. The sella Turcica.
 19. The sphenoid sinus.
 18. The nasal plate of the sphenoid bone.
 20. The spongy substance of the sphenoid and occipital bones.
 β , The hole for the passage of the ninth pair of nerves.
 c, The squamous part of the temporal bone.
 e, The ridge of the os petrosum, with the print of a small sinus.
 f, The internal meatus auditorius.
 λ , The dentes incisores.
 μ , The dens caninus.
 ν , The dentes molares.
 π , The foramen incisivum of the maxillary bone.
 ξ , The rough spine of the superior maxillary bone.
 ϕ , The joining of it to the vomer.
 ϵ , The broad hollow base of the vomer.
 H, The posterior edge of the vomer.
 G, The body of the vomer.
 τ , The conjunction of it with the thin plate of the sphenoid and ethmoid bones.
 α , Its hollow anterior part, which receives the middle cartilage of the nose.
 P, The anterior edge of the nasal bone.

FIG. 17.

The External SURFACE of the Base of the CRANIUM and UPPER JAW.—See Tab. VI.

- q, The tubercle of the root of the zygoma.
 r, The concave moveable cartilage placed on that tubercle.
 t, t, Its ligaments.
 2. The hole for the portio dura of the 7th pair of nerves.
 5. The bony part of the Eustachian tube.

T A B L E XXXI.

Represents the LARYNX, the BONES of the TRUNK of the BODY, and a Posterior View of the MALE SKELETON.

FIG. 1.

Anterior View of the CARTILAGES of the LARYNX, with the Os HYOIDES.

- a*, THE anterior surface of the base of the os hyoides.
b, Its superior surface.
c, A ligament connecting the os hyoides, thyroid cartilage, and epiglottis.
d, d, The two appendices of the os hyoides.
d, The ligament sent out from the appendix of the left side, to the styloid process of the left temporal bone.
e, The union of the base with the cornu.
f, f, The two cornua.
g, g, Tubercles at their extremities.
h, h, Ligaments going from the tubercles to the superior cornua of the thyroid cartilage.
i, k, &c. The thyroid cartilage; the greater part of which is ossified in the preparation from which this Figure is taken.
i, The anterior middle part of the thyroid cartilage.
k, Its right side.
l, l, Two unossified cartilaginous pieces on that side.
m, The right superior cornu.
n, The right inferior cornu, connected to the cricoid cartilage.
o, A strong ligament.
p, The narrow anterior part of the cricoid cartilage.
q, Its right side.
r, The first cartilage of the trachea arteria, divided into two at the sides.
s, The second, third, and fourth cartilages of the trachea.

FIG. 2.

Back-View of the PARTS represented in FIG. 1.

- d, d, f, g, l, m, n*, The same parts pointed out by these letters in Fig. 1.
t, The epiglottis.
G, G, The two arytenoid cartilages.
v, The middle unossified part of the cricoid cartilage.

W, W, The bony sides of that broad posterior part.

X, The membranous back-part of the trachea.

N. B. These two Figures are as large as life; whereas the Figures of particular bones in this and the preceding Table, are represented only one half as large as Nature.

FIG. 3.

A View of the Upper Part of the First VERTEBRA of the NECK.

a, The body of the bone.

The other parts are described in Tab. XVIII. Fig. 2.

FIG. 4.

The Under and Back Part of the same VERTEBRA.

i, The smooth depression for the anterior part of the tooth-like process of the second vertebra.

For the other parts consult Tab. XVIII. Fig. 1.

FIG. 5.

A Side-View of the Second VERTEBRA of the NECK.

a, The body.

b, The superior oblique process.

c, The transverse process, and farther back, the inferior oblique process.

d, The plate extended to,

e, The spinous process.

g, The spinous hole.

h, The passage in the transverse process.

l, The point of the tooth-like process.

m, Its anterior smooth surface.

n, Its posterior smooth surface.

FIG. 6.

The First and Second VERTEBRÆ of the NECK, with part of the OCCIPITAL BONE, and the LIGAMENTS of the TOOTH-LIKE PROCESS..

a, Part of the occipital bone.

b, b, Its








FIG. I6.




FIG. I5.




FIG. I3.

FIG. I2.



FIG. I0.





FIG. I4





- b, b*, Its broken extremities.
1. c, b, Part of the first vertebra.
2. e, Part of the second vertebra.
n, The point of the tooth-like process.
o, Its transverse ligament.
p, p, Its two oblique, or moderator ligaments.
q, Its perpendicular ligament.
 See Tab. XVIII. Fig. 6.

FIG. 7.

Upper Part of the Fourth Vertebra of the Neck.

FIG. 8.

Upper Part of the Seventh Vertebra of the Neck.

FIG. 9.

Upper Part of the Seventh Vertebra of the Back.—See Tab. XVIII. Fig. 7.

FIG. 10.

Under Part of the Sixth Vertebra of the Back.—See Tab. XVIII. Fig. 8.

FIG. 11.

Side-View of the Twelfth VERTEBRA of the BACK.

- a*, The body.
c, The transverse process.
e, The spinous process.
k, The inferior oblique process.
t, The depression for the head of the rib.

FIG. 12.

View of the Upper Part of the Fourth Vertebra of the Loins.—See Tab. XVIII. Fig. 10.

FIG. 13.

Under and Lateral Part of the Third Vertebra of the Loins.

- a*, The under part of the body.
b, The superior oblique process.
c, The transverse process.
e, The spinous process.
g, The spinal hole.
r, The process round the body of the Bone.

FIG. 14.

The Seventh TRUE RIB of the LEFT SIDE.

- a*, Its head.
b, Its smooth surface, which was joined to the transverse process of the vertebra.
c, The depression.
d, The tubercle.

- e*, The angle.
f, The furrow at the inferior edge.
g, The smooth internal side.
h, The anterior extremity.

FIG. 15.

The Sixth and Seventh Vertebra of the Back, with Part of the Seventh Rib of the Left Side.

- a, c, e, k, t*, As in Fig. 11.
u, The cartilage between the vertebrae.
v, The depression made by the tubercle of the sixth rib.
W, The seventh rib articulated with the vertebra.
x, The beginning of the furrow on the under edge.

FIG. 16.

The STERNUM, with the CARTILAGES of the RIBS, and the Internal MAMMARY ARTERIES.

- A*, The first or upper bone of the sternum.
B, The second.
C, The third, or cartilago ensiformis.
D, D, D, D, The cartilages of the four superior ribs.
E, The conjoined cartilages of the fifth, sixth, and seventh ribs.
F, F, F, The radiated ligaments connecting the cartilages to the sternum.
G, The internal mammary artery.

FIG. 17.

Posterior View of the MALE SKELETON. HEAD.

- a*, The coronal suture.
b, The squamous.
o, The lambdoid.
A, The frontal bone.
B, The right parietal bone.
C, The right temporal bone.
D, The occipital bone.
E, The nasal bone.
F, The os malæ.
3, The angle of the lower jaw.
4, Its right coronoid process.
5, Its right condyle.

TRUNK.

- K*, The seven vertebrae of the neck.
L, L, The twelve vertebrae of the back.
M, The five vertebrae of the loins.
b, The transverse processes of the os sacrum.
c, The posterior holes of that bone.
e, Its spinous processes.
g, The open part of the canal for the cauda Equina.
P, The os coccygis.
α, The dorsum of the os ilium.

β, Its

- β , Its spine.
 γ , The superior-posterior spinous process.
 δ , The inferior-posterior spinous process.
 ϵ , The great notch.
 μ , The superior-anterior spinous process.
 ν , The inferior-anterior process.
 σ , The brim of the acetabulum.
 ζ , The spinous process of the os ischium.
 η , The tuberosity of that bone.
 S , Its branch.
 R , The os pubis.
 π , Its crus.
 ς , The great thyroid hole.

RIGHT SUPERIOR EXTREMITY.

- d , The dorsum of the scapula.
 f , Its posterior costa.
 g , Its superior angle.
 p , The anterior or inferior costa.
 q , The inferior angle.
 t , The cervix of the bone.
 w , The acromion, to which the outer end of the clavicle is joined.
 K , The spine.
 W , The fossa above the spine.
 ϕ , The superior costa, with the femilunar notch.
 r , The inner end of the clavicle joined to the sternum.
 X , The body of the clavicle.
 a , The ulna.
 b , The radius.
 6. The head and neck of the radius.
 7. The olecranon of the ulna.
 9. The under end of the radius.

RIGHT INFERIOR EXTREMITY.

- f , The ball of the os femoris.
 g , The trochanter major.
 b , The cervix of the bone.
 i , The trochanter minor.
 k , The upper part of the body of the bone.
 l , Its outer condyle.
 t , The malleolus externus of the fibula.

LEFT SUPERIOR EXTREMITY.

- ω , The inner condyle of the os humeri.
 7. The olecranon of the ulna.


LEFT INFERIOR EXTREMITY.

1. The linea aspera.
 2. k , The inner condyle, and,
 3. The outer condyle of the os femoris.
 4. The head of the tibia.
 5. The body of the bone.
 t , The malleolus internus of the tibia.
 15. The head of the fibula.
 r , The malleolus externus of that bone.
 10. The os calcis.

FIG. 18.

Posterior View of the BONES and LIGAMENTS of the PELVIS.

- a , The fifth lumbar vertebra.
 b , Its superior oblique process.
 d , The bony plate extended to its spinal process e .
 $c, c, \&c.$ The posterior holes of the os sacrum.
 g , The channel for the cauda Equina.
 $\alpha, \beta, \gamma, \delta, \epsilon, \zeta, \eta$, As in Fig. 17.
 A , The posterior sacro-ischiatic ligament, extended from the tuber of the os ischium to the os ilium, os sacrum, and os coccygis.
 B , The anterior sacro-ischiatic ligament, proceeding from the spinous process of the os ischium, to the os sacrum and os coccygis.
 ς , The notch of the os ilium, for the passage of the posterior crural vessels and nerves, and the pyriform muscle.
 C , Passage of the obturator internus.
 D , The fibrous ligamentous substance, connecting the os innominatum and sacrum.
 E , The capsular ligament of the joint of the thigh.
 g , The greater,
 i , The lesser trochanter of the thigh-bone.






FIG. 1.




FIG. 2.




FIG. 3.




FIG. 4.



FIG. 5.




FIG. 6.



FIG. 8.




FIG. 9.

FIG.15.



FIG.16.





FIG.14.



FIG.13.




FIG.12.




FIG.11.






FIG.10.





T A B L E XXXII.

Represents the BONES and some of the Principal LIGAMENTS of the EXTREMITIES ; including the SKELETON of two YOUNG SUBJECTS.

FIG. 1.

A Posterior View of Part of the STERNUM and CLAVICLES, with the LIGAMENT connecting the CLAVICLES to each other.

- a*, THE posterior surface of the sternum.
- b, b*, The broken ends of the two clavicles.
- c, c, c, c*, The two tubercles near the extremity of each clavicle.
- d*, The ligament connecting the clavicles.

FIG. 2.

Outer and Fore View of the LEFT SCAPULA, and part of the CLAVICLE, with their LIGAMENTS.

- a*, The spine of the scapula.
- b*, The acromion.
- c*, The inferior angle.
- d*, The inferior costa.
- e*, The cervix.
- f*, The glenoid cavity, covered with cartilage for the articulation with the os humeri.
- g, g*, The cut edge of the capsular ligament of the joint of the arm.
- h*, The coracoid process.
- i*, The point of that process.
- k*, The broken end of the clavicle.
- l*, Its extremity joined to the acromion.
- m*, A ligament stretched obliquely from the clavicle to the coracoid process.
- n*, A ligament coming out single from the acromion, and dividing into two, which are fixed to the coracoid process.

FIG. 3.

The JOINT of the ELBOW of the LEFT ARM, with the LIGAMENTS.

- a*, The os humeri.
- b*, Its internal condyle.
- c, c*, The two prominent parts of its trochlea, appearing through the capsular ligament of the joint.
- d*, The ulna.
- e*, The radius.
- f*, That part of the ligament including the head of the radius.
- g*, The end of that ligament, like a ring, surrounding the neck of the radius, but connected very loosely to it.

FIG. 4.

Anterior View of the BONES of the RIGHT HAND.

- a*, The radius.
- e*, Its flat anterior part.
- c*, Its styloid process.
- b*, The ulna.
- f*, Its flattened extremity.
- d*, Its styloid process.
- g*, The os scaphoides of the carpus.
- h*, The os lunare.
- i*, The os pisiforme.
- k*, The cuneiforme.
- l*, The trapezium.
- m*, The trapezoides.
- n*, The capitatum.
- o*, The unciforme.
- p*, Its unciform process.
- q, r, s, t*, The metacarpal bones of the fingers.
- u*, Their bases.
- v*, Their heads.
- α*, The metacarpal bone of the thumb.
- β*, The first bone of the thumb.
- γ*, The second bone.
- δ*, The first phalanx of the fingers.
- ζ*, Their second phalanx.
- η*, Their third phalanx.

FIG. 5.

Posterior View of the BONES of the LEFT HAND.

The explanation of Fig. 4. will serve for this Figure, the same letters pointing to the same bones, though in a different view.

- α*, The ridge of the radius, between the grooves made by the tendons of the extensor muscles.

FIG. 6.

View of the Anterior, or Palm-Side of the RIGHT HAND, with its LIGAMENTS.

- i*, The pisiform bone.
- p*, The hook-like process of the unciform bone.
- θ*, The annular ligament, under which the tendons of the flexor muscles pass in the cavity *α*.
- q—t*, The metacarpal bones of the fingers.
- γ*, Their bases, with the ligaments connecting them to the bones, pointed out by *m, n, o*, in Fig. 4.

α, β, γ,

- α, β, γ , The metacarpal bone, and the two bones of the thumb, with the ligaments of their articulations.
 χ , The fore-fingers, with the sheath for the tendons of the flexor muscles entire.
 λ , The ligament connecting the head of its metacarpal bone to that of the middle finger.
 μ , The middle finger, with the sheath of the tendons cut open.
 ν , The ligaments on the back-part of the second joint of the ring and little fingers.

FIG. 7.

The Upper Extremity of the TIBIA, with the semilunar CARTILAGES of the KNEE-JOINT and some LIGAMENTS.

- a , The strong ligament which connects the patella to the tubercle of the tibia.
 b, b , The parts of the extremity of the tibia, covered with cartilage, which appear within the semilunar cartilages.
 c, c , The semilunar cartilages.
 d , Part of the crucial ligaments.

FIG. 8.

Posterior View of the JOINT of the RIGHT KNEE.

- a , A section of the os femoris.
 b , Its internal condyle.
 c , Its external condyle, both covered with cartilage.
 d , The cavity between the condyles.
 e, e , The back part of the tibia.
 f , The superior extremity of the fibula.
 g , The edge of the internal semilunar cartilage.
 h , An oblique ligament.
 i , A small perpendicular ligament.
 k , A larger perpendicular ligament.
 l , The external lateral ligament, connecting the femur and fibula.
 m , A ligament between the tibia and fibula.

FIG. 9.

Anterior View of the JOINT of the RIGHT KNEE.

- b , The internal condyle.
 c , The external.
 d , Part of the os femoris, on which the patella moves.
 e , A perpendicular ligament.
 f, f , The crucial ligaments.
 g, g , The edges of the two semilunar cartilages.
 h , The tibia.
 i , The strong ligament of the patella.
 k , The back-part of it, where some of the fat of the joint has been dissected away.
 l , The external depression.
 m , The internal one, on the posterior surface of the patella.
 n , A section of the tibia.

FIG. 10.

A View of the Inferior Part of the BONES of the RIGHT FOOT.

- a , The great knob of the os calcis.
 b , A prominence on its outside.
 c , The interior thin process, bearing the print of the tendon of the flexor pollicis longus.
 d , The hollow, for the tendons, nerves, and blood-vessels.
 e , The anterior extremity of the os calcis.
 f , Part of the astragalus.
 g , Its head, covered with cartilage.
 h , The internal prominence of the os naviculare.
 i , Its hollow in the sole of the foot.
 k , The os cuboides.
 l , Its hollow, for the tendon of the peroneus longus.
 m , Its anterior extremity.
 n , The os cuneiforme internum.
 o , The medium.
 p , The externum.
 q, r, s, t , The metatarsal bones of the four lesser toes.
 u , Their bases.
 v , Their heads.
 α , The metatarsal bone of the great toe.
 β , Its first,
 γ , Its second bone.
 2 , The depressions on the head of the metatarsal bone, for the two sesamoid bones.
 δ , The first,
 ζ , Second, and,
 η , Third phalanx of the four lesser toes.

FIG. 11.

The Inferior Surface of the Two Large Sesamoid Bones, at the First Joint of the Great Toe.

FIG. 12.

Upper View of the BONES of the RIGHT FOOT.

- a , The posterior knob of the os calcis.
 b , Its exterior process.
 k , Its anterior extremity.
 c , The superior head of the astragalus.
 d , A depression made by the tendon of the flexor pollicis longus.
 f , The rough hollow part.
 g , The anterior head.
 h , The os naviculare.
 i , The os cuboides.
 l , The hollow for the peroneus longus.
 n , The internal,
 o , The middle,
 p , The external cuneiform bones.
 $u, v, \alpha, \beta, \gamma, \delta, \zeta, \eta$, The same as in Fig. 10.

FIG.

FIG. 13.

View of the SOLE of the FOOT, with its LIGAMENTS.

- a, d,* As in Fig. 10.
e, The sheaths of the flexores longi pollicis et digitorum opened.
f, The strong cartilaginous ligament supporting the head of the astragalus.
g, h, Two ligaments which join into one, to be fixed to the metatarsal bone of the great toe.
i, j, k, l, m, Other ligaments.
n, o, The ligaments of the joints of the five metatarsal bones.

FIG. 14.

The Superior Concave SURFACE of the SESAMOID BONES at the First JOINT of the GREAT TOE, with their LIGAMENTS.

- a,* Three sesamoid bones.
b, The ligamentous substance in which they are formed.

FIG. 15.

Front View of the SKELETON of a BOY of nine years of age.

- ** &c.* The most remarkable epiphyses.
f, The joining of the os ilium and pubis, and,
g, Of the os ischium and pubis.

This Figure is executed upon a scale only half as large as that of the Skeleton in Tab. XXIV.

FIG. 16.

The SKELETON of a NEW-BORN CHILD, where the shades and shrivellings in the Figure represent the parts which are Cartilaginous at Birth, and which are contracted in the Skeleton.

- a,* The fontanelle.

This Figure is too small, even in proportion to Fig. 15.

P A R T II.

OF

T H E M U S C L E S.

И Т Р А Ч

03

Т Н Р М И С Г Е К С

OF THE MUSCLES IN GENERAL.

THE MUSCLES serve for the motion of the different parts of the Body, and derive their general name from their power of contracting.

Of Muscles in general, the following things are to be attended to.

The *Cellular Substance*, which surrounds the Muscles, and allows them to move upon each other, and upon the adjacent parts.

The *Cellular Substance*, condensed in certain parts of the Body, and giving an appearance of *Membrane*, formerly called *Tunica Propria Muscularum*.

The *Division* of a Muscle into

The *Origin*, or *Head*;—or that Extremity of the Muscle which arises from the most fixed part, and towards which the contraction is made;

The *Belly*, or *thickest part*, which swells when the Muscle is in action;

The *Insertion*, or *Termination*, or that Extremity which is implanted into the part to be moved, and which is commonly smaller than the Origin.

The Division of a Muscle into Fleshy and Tendinous Parts.

The *Fleshy part* distinguished by being *soft, sensible*, generally of a *red colour*,—from the great quantity of Blood in it—and possessing *contractility*.

The *Fleshy part*, having numerous *Blood-vessels, Lymphatics, and Nerves*.

The *Division* of Muscles into *Rectilineal*, as in the *Sarto-*

rius;—*Simple Penniform*, as in the *Peroneus Longus*;—*Complete Penniform*, as in the *Rectus Femoris*;—*Compound Penniform*, as in the *Fore-part of the Soleus*;—*Radiated*, as in the *Pectoralis Major*;—*Hollow*, as in the *Heart, Intestines, Bladder of Urine, &c.*

The *particular Names* of Muscles are taken from their *shape, size, situation, direction, composition, use, and attachment*.

The *Tendon*, is distinguished from the Fleshy part, by being generally *smaller, firmer, stronger*;—of a *white glistening colour*, having *no contractility*, and little or no *sensibility* in the sound state.

Tendons have very few *Blood-vessels*, and no evident *Nerves*.

Tendons connect Muscles to Bones, and by the smallness of their size compared to the Belly of the Muscle, preserve the elegance and symmetry of the Joints, &c.

The *Appendages* of Muscles, viz.

Aponeuroses, or Fasciæ, which are the Tendons expanded upon a wide Surface, serving to give insertion to Muscular Fibres, to keep them in their proper situation, and to brace them in their action.

Annular Ligaments, to keep Tendons from starting.

Trochleæ, or Pulleys, to alter the direction of Tendons.

Bursæ Mucosæ, placed where Tendons play over hard Substances; serving to contain Synovia, and prevent Abrasion.

MUSCLES OF THE INTEGUMENTS OF THE CRANIUM, AND OF THE EYE-LIDS.

OCCIPITO-FRONTALIS,

Vel Occipitalis and Frontalis, vel Epicranius, &c.

Origin: Fleshy from near the middle of the upper arched Ridge of the Occipital Bone, Tab. XL. Fig. 1. a; and Tendinous from the extremity of that Ridge, where it joins the Temporal Bone.—It arises after the same manner on the other side. From the Fleshy Origins, and also from between them, a Tendinous expansion is extended along the upper part of the Cranium, adhering firmly to the skin, and but loosely to the Pericranium, Tab. XL. Fig. 1. b, c. Tab. XXXIV. Fig. 1. a.—At

the upper part of the Forehead it becomes Fleshy, and, descends with straight Fibres.

Insertion: Into the Skin and parts under it belonging to the Eye-brows, and to the Frontal Bone at the inner part of the Orbit. Tab. XXXIV. Fig. 1. A.

Action: To move all that part of the Skin which covers it, and particularly the Skin of the Brow and Eye-brows.

From the under and middle part of the Muscle, a *Slip* is continued down upon the Root of the Nose, to be connected with the Compressor Naris, and Levator Labii Superioris Alæque Nasi. Tab. XXXIV. Fig. 1. b.

This

This Slip may either assist the Nasal Muscles connected with it, or antagonize the Occipito-frontalis.

CORRUGATOR SUPERCILII.

Origin: From the internal angular Process of the Os Frontis, above the joining of that Bone with the Os Nasi. From thence it runs upwards and outwards, in the direction of the Superciliary Ridge, and behind the inferior part of the Occipito Frontalis.

Insertion: Into the inner part of the Occipito-frontalis and Orbicularis Palpebrarum, where these two Muscles join each other, as far out as the middle of the Superciliary Ridge. Tab. XXXV. Fig. 1. A.

Action: To assist its fellow in drawing the Eye-brows downwards and inwards, and corrugating or wrinkling the Skin between them into longitudinal folds.

ORBICULARIS OCULI, vel Palpebrarum.

Origin: From the Orbital Process of the superior Maxillary Bone; from the internal Angular Process of the Frontal Bone; and, by a small round Tendon, from the Nasal Process of the superior Maxillary Bone.

From these Origins the Muscle passes outwards, under the Skin of the Eye-lids, surrounding the Orbit in a circular manner; extending somewhat beyond it, and covering the upper part of the Cheek. Tab. XXXIV. Fig. 1. D.

The outer Surface of the Muscle adheres to the Skin of the Eye-lids; its upper and inner Edge is intimately connected with the Frontal and Corrugator Muscles.

Action: To close the Eye by bringing the Eye-lids together, to press the Ball of the Eye inwards, and act upon the Lacrymal Organs, so as to assist them in the production and direction of the Tears.

That part of the Orbicularis-Oculi which covers the Cartilages of the Eye-lids, and which is remarkably thin, is the *Musculus Ciliaris* of some Authors. Tab. XXXIV. Fig. 1. c.

A *Fleshy Slip* frequently passes down from the under and outer part of the Orbicularis, to join the Levator Labii Superioris Alæque Nasi. Tab. XXXIV. Fig. 1. between F and G. When present, it may draw a little towards each other those parts to which it is attached.

LEVATOR PALPEBRÆ SUPERIORIS.

Origin: From the upper margin of the Foramen Opticum of the Sphenoid Bone. It runs forwards within the Orbit over the Levator Oculi, where it becomes gradually broader, its anterior extremity passing under the Orbicularis Oculi.

Insertion: By a broad thin Tendon, into nearly the whole length of the Cartilage of the upper Eye-lid. Tab. XXXV. Fig. 1. a.

Action: To open the Eye by raising the upper Eye-lid.

MUSCLES COMMON TO THE HEAD AND EXTERNAL EAR.

ATTOLLENS AUREM, vel Superior Auris.

Origin: By a broad Tendinous expansion, from the Tendon of the Occipito-frontalis. It goes down over the Aponeurosis of the Temporalis. In its passage, it forms a thin Fleshy Slip, which becomes gradually narrower.

Insertion: Into the upper part of the root of the Cartilage of the Ear. Tab. XXXIV. Fig. 1. B.

Action: To give tension to the part into which it is inserted, and, in some Persons, to raise the Ear.

Insertion: By a narrow Tendon into the back-part of the beginning of the Helix. Tab. XXXIV. Fig. 1. C.

Action: To stretch that part of the Ear to which it is fixed.

RETRAHENTES AUREM, vel Posteriores Auris.

Origin: By two, and sometimes by three distinct Muscles, from the upper and outer part of the Mastoid Process.

Insertion: By small Tendons into the back-part of the Concha. Vol. II. First Table of the Ear, Fig. 2.

Action: To stretch the Concha, and, in some persons, to draw back the Ear.

ANTERIOR AURIS.

Origin: Thin and Membranous, near the posterior part of the Zygoma; the middle part being mixed with Fleshy Fibres.

MUSCLES OF THE NOSE AND MOUTH.

COMPRESSOR NARIS.

Origin: By a narrow beginning from the root of the Ala Nasi, where it is connected with the Levator Labii Superioris Alæque Nasi. It spreads into a number of thin scattered Fibres, which cross the Ala Nasi, and run towards the Dorsum Nasi, where it joins its fellow.

Insertion: Into the anterior extremity of the Nasal

Bones, and to the Slip which descends from the Frontal Muscle. Tab. XXXIV. Fig. 1. d.

Action: To press the Ala towards the Septum, as in smelling; or if the Fibres of the Frontal Muscle, which are connected to it, act, they pull the Ala outwards. It also corrugates the Skin of the Nose, and assists in expressing certain Passions.

LEVATOR

LEVATOR LABII SUPERIORIS ALÆQUE NASI.

Origin: By two thin Flethy Slips; the first from the external part of the Orbital Process,—the second from the upper part of the Nasal Process, of the Superior Maxillary Bone.

Insertion of the first part of the Muscle into the Upper Lip, and of the second into the Upper Lip and outer part of the Wing of the Nose. Tab. XXXIV. Fig. 1. F, E.

Action: To raise the Upper Lip in opening the Mouth, and to dilate the Nostril.

DEPRESSOR LABII SUPERIORIS ALÆQUE NASI.

Origin: Thin and Flethy, from the Alveoli of the Dentes Incisivi and Caninus of the Upper Jaw; running upwards, at the side of the Furrow of the Lip.

Insertion: Into the Upper Lip, and root of the Ala Nasi.

Action: To draw the Upper Lip and Ala Nasi downwards. Tab. XXXVI. Fig. 1. E.

LEVATOR ANGULI ORIS,

Vel *Levator Labiorum Communis*, vel *Caninus*.

Origin: Thin and Flethy, from the superior Maxillary Bone, immediately under the Foramen Infra-orbitarium;—running deeper down and farther out than the Levator Labii Superioris.

Insertion: Into the Angle of the Mouth, and the Cheek, where it joins its Antagonist. Tab. XXXV. Fig. 1. D.

Action: To raise the corner of the Mouth;—as in expressing Joy.

DEPRESSOR LABII INFERIORIS, vel *Quadratus Genæ*.

Origin: Broad and Flethy, from the under part of the Lower Jaw, at the side of the Chin;—from thence it runs obliquely upwards and inwards, till it becomes contiguous to its fellow in the middle of the Lip.

Insertion: Into one half of the Edge of the Under Lip. Tab. XXXV. Fig. 1. G.

Action: To assist in opening the Mouth, by depressing the Under Lip, and pulling it a little outwards.

LEVATOR LABII INFERIORIS, vel *Levator Mentis*.

Origin: From the roots of the Alveoli of the Dentes Incisores and Dens Caninus of the Lower Jaw.

Insertion: Into the Under Lip, and Skin of the Chin. Tab. XXXVI. Fig. 1. H.

Action: To raise the parts into which it is inserted.

DEPRESSOR ANGULI ORIS, vel *Triangularis*.

Origin: Broad and Flethy, from the under edge of the Lower Jaw, at the side of the Chin.—It runs over the

Origin of the Depressor Labii Inferioris, becoming gradually narrower.

Insertion: Into the angle of the Mouth, where it intermixes with the Levator Anguli Oris. Tab. XXXIV. Fig. 1. K.

Action: To depress the corner of the Mouth;—as in expressing Anger.

ZYGOMATICUS MAJOR.

Origin: Flethy from the Os Malæ, near the Zygomatic Suture.—Descending obliquely forwards.

Insertion: Into the angle of the Mouth; its Fibres intermixing with those of the Depressor Anguli Oris and Orbicularis Oris. Tab. XXXIV. Fig. 1. H.

Action: To raise the angle of the Mouth, in the direction of its Fibres, and to make the Cheek prominent;—as in Laughing.

ZYGOMATICUS MINOR.

Origin: Higher on the Os Malæ than the former Muscle. It is situated before it, and takes the same course, but is much more slender.

Insertion: Into the Upper Lip, along with the Levator Anguli Oris. Tab. XXXIV. Fig. 1. G.

Action: To raise the corner of the Mouth, and draw it obliquely outwards.

This Muscle is often wanting.

By the frequent action of the Muscles which raise the corners of the Mouth and Upper Lip, that Furrow is formed which extends between the outer corner of the Nose and Mouth, and which is so conspicuous in the Face of a person advanced in life.

BUCCINATOR, vel *Retractor Anguli Oris*.

Origin: From a Ridge, extending between the last Dens Molaris and Coronoid Process of the Lower Jaw;—from the Upper Jaw, between the last Dens Molaris and Pterygoid Process of the Sphenoid Bone; from the extremity of which it has also part of its origin. Thence going forwards with straight Fibres, it adheres closely to the Membrane which lines the Mouth.

Insertion: Into the corner of the Mouth, along with the Orbicularis Oris. Tab. XXXVI. Fig. 1. G.

Action: To draw the angle of the Mouth backwards and outwards, and to contract its Cavity by pressing the Cheek inwards, by which the Food is thrust between the Teeth in Manducation.—It is likewise active in blowing Wind-instruments,—as a Trumpet;—hence its name.

ORBICULARIS ORIS, vel *Sphincter Labiorum*.

This is a complete Sphincter surrounding the Mouth; great

and composing the principal part of the Lips, and is in a great measure formed by the Muscles which terminate in it.—At the corners of the Mouth, the Fibres decussate each other, so as to make it resemble two semicircular Muscles, from which it has been named by some Authors, *Semi-Orbicularis Superior*, and *Semi-Orbicularis Inferior*. Tab. XXXVI. Fig. 1. F.

Action: To shut the Mouth, and to counteract the different Muscles inserted into it.

Nasalis Labii Superioris of Albinus,—part of the former Muscle, running up to be connected to the Septum Nasi, and serving as a Levator of the Upper Lip, or a Depressor of the under part of the Nose. Tab. XXXIV. Fig. 1. above L.

MUSCLES OF THE LOWER JAW.

APONEUROSIS TEMPORALIS.

This is a strong Tendinous Membrane, proper to be taken notice of before describing the Temporalis. It arises from the Bones which give origin to the upper semicircular part of the Temporal Muscle, and, descending over it, is fixed to the Zygoma.

Use: To brace the whole, and to give origin to part of the Temporal Muscle. Tab. XLIV.

TEMPORALIS.

Origin: Semicircular and Fleishy, from the lower half of the Parietal Bone, and Temporal Fossa of the Frontal Bone; and from the Squamous part of the Temporal, and Temporal Plate of the Sphenoid Bones.—It arises likewise from the Aponeurosis covering it;—from these Origins the Fibres descend like Radii, and the Muscle sends off a strong Tendon, which passes under the Jugum.

Insertion: Into the whole of the Coronoid Process of the Lower Jaw, which it incloses as in a sheath, and is continued to near the last Dens Molaris. Tab. XXXV. Fig. 1. B, b.

Action: To pull the Lower Jaw upwards and a little backwards against the Upper Jaw.

MASSETER.

Origin: By strong Tendinous and Fleishy Fibres from the Superior Maxillary Bone, where it joins the Os Malaæ, and from the whole length of the under and inner edge of the Zygoma;—the outer part of the Muscle slanting backwards, the inner part forwards, and in some measure decussating the other. In its descent, it covers the Coronoid Process and under end of the Temporal Muscle.

Insertion: Into the angle of the Lower Jaw, and from that upwards to the outside of the Coronoid Process. Tab. XXXV. Fig. 1. C.

Action: To raise the Lower Jaw, and to pull it a little forwards or backwards, according to the direction of the Fibres of the Muscle.

PTERYGOIDEUS INTERNUS, vel *Major*.

Origin: From the Fossa Pterygoidea of the Sphenoid and Palate Bones; passing downwards and outwards.

Insertion: Into the inner side of the Angle of the Lower Jaw, and continued as far as the Groove for the inferior Maxillary Nerve. Tab. XXXVII. Fig. 1. inner side of G.

Action: To raise the Jaw, and draw it obliquely towards the opposite side.

PTERYGOIDEUS EXTERNUS, vel *Minor*.

Origin: From the outer side of the Pterygoid Process of the Sphenoid Bone; from the Tuberosity of the Superior Maxillary Bone; and from the root of the Temporal Process of the Sphenoid Bone. From these origins it passes, almost horizontally, outwards and a little backwards.

Insertion: Into the Cervix and Capsular Ligament of the Lower Jaw. Tab. XXXIX. Fig. 1. k.

Action: To pull the Lower Jaw to the opposite side, and, if both Muscles act, to bring it forwards, so as to make the Fore-Teeth project beyond those of the Upper Jaw. The Muscle, in its different motions, acts also upon the Inter-articular Cartilage.

MUSCLES ON THE FORE AND LATERAL PART OF THE NECK.

PLATYSMA MYOIDES, vel *Cutaneus*.

Origin: By a number of separate Fleishy Slips, from the Cellular Substance, which covers the upper parts of the Pectoral and Deltoid Muscles.—In their ascent, they unite to form a thin Muscular Expansion, which runs obliquely upwards along the side of the Neck, adhering to the skin, and is similar to the Cutaneous Muscle of Quadrupeds. Tab. XXXIV. Fig. 1. l, M.

Insertion: Into the side of the Lower Jaw and the Depressor Anguli Oris, and into the skin which covers the under part of the Masseter and Parotid Gland. Tab. XXXIV. Fig. 1. i.

Action: To assist in depressing the Lower Jaw, the corner of the Mouth, and the Skin of the Cheek;—when the Jaws are shut, to raise all that part of the Skin connected with it under the Lower Jaw.

STERNOC.

STERNO-CLEIDO-MASTOIDEUS, vel *Sterno-Mastoideus*.

Origin: From the top of the Sternum, and the anterior end of the Clavicle, by two distinct Heads; the first of which is round, tendinous, and a little fleshy; the other is broad and fleshy. A little above the Clavicle, the two Heads unite to form a strong Muscle, which runs obliquely upwards and outwards; the greater part

of it being covered by the *Platysma Myoides*. Tab. XXXV. Fig. 1.

Insertion: By a thick strong Tendon, into the Mastoid Process, which it surrounds; and becoming thinner, the Insertion extends as far back as the Lambdoid Suture.

Action: To turn the Head to one side, and assist in rolling it. When both Muscles act, they bow the Head.

MUSCLES SITUATED BETWEEN THE OS HYOIDES AND TRUNK.

STERNO-HYOIDEUS.

Origin: From the edge of the upper Bone of the Sternum internally, and from the adjacent parts of the Clavicle and Cartilage of the first Rib;—ascending upon the fore-part of the Trachea and following Muscle.

Insertion: Into the Base of the Os Hyoides. Tab. XXXV. Fig. 1. I.

Action: To depress the Os Hyoides.

STERNO-THYROIDEUS.

Origin: From the upper and inner part of the Sternum, and partly from the Cartilage of the first Rib; running along the fore-part and side of the Trachea and Thyroid Gland.

Insertion: Into the under and lateral part of the Thyroid Cartilage. Tab. XXXVI. Fig. 1. K.

Action: To depress the Larynx.

THYRO-HYOIDEUS, vel *Hyo-Thyroideus*.

Origin: Where the former Muscle terminates, having the appearance of being continued from it.

Insertion: Into part of the Base, and almost all the Cornu of the Os Hyoides. Tab. XLVII. Fig. 1. k.

Action: To depress the Os Hyoides, or to raise the Thyroid Cartilage.

CRICO-THYROIDEUS.

Origin: From the side and fore-part of the Cricoid Cartilage; running obliquely upwards.

Insertion: By two portions; the one into the under part of the Thyroid Cartilage, the other into its inferior Cornu. Tab. XLVII. Fig. 1. m.

Action: To depress and pull forwards the Thyroid Cartilage, or to raise and draw backwards the Cricoid Cartilage.

OMO-HYOIDEUS.

Origin: From the superior Costa of the Scapula, near the Semilunar Notch. It goes obliquely upwards and forwards, and is of a very slender form. Under the Sterno-mastoideus, it becomes Tendinous, and again grows Fleshy.

Insertion: Into the Base of the Os Hyoides, at the side of the Sterno-Hyoideus. Tab. XXXV. Fig. 11. k.

Action: To depress the Os Hyoides, and pull it to one side; or, when both act, to draw it directly down.

MUSCLES SITUATED BETWEEN THE LOWER JAW AND OS HYOIDES.

DIGASTRICUS, vel *Biventer Maxillæ Inferioris*.

Origin: By a Fleshy Belly, from the Groove at the root of the Mastoid Process of the Temporal Bone. It runs downwards and forwards, and forms a strong round Tendon, which passes through the Stylo-Hyoideus;—it is then fixed by a Ligament to the Os Hyoides, and, having received an addition of Tendinous and Muscular Fibres, it runs obliquely upwards and forwards, forming another Fleshy Belly.

Insertion: Into a rough Sinuosity at the under part of the Symphysis of the Lower Jaw. Tab. XLIX. Fig. 1. b, c.

Action: To open the Mouth by pulling the Lower Jaw downwards and backwards; and when the Jaws are

shut, to raise the Os Hyoides, and, of course, the Throat,—as in Swallowing.

MYLO-HYOIDEUS.

Origin: Fleshy, broad, and thin, from the inside of the Lower Jaw, between the last Dens Molaris and the middle of the Chin, where it joins its fellow; running down behind the Digastricus.

Insertion: Into the lower edge of the Body of the Os Hyoides, and joined to its fellow by the intervention of a white Tendinous Line. Tab. XLIX. Fig. 2. a.

Action: To pull the Os Hyoides forwards, upwards, and to a side.

GENIO-

GENIO-HYOIDEUS.

Origin: From a Tubercle on the under and inner part of the Symphysis of the Lower Jaw, by a slender beginning, which descends and gradually becomes broader.

Insertion: Into the Body of the Os Hyoides, under the former Muscle. Tab. XLIX. Fig. 3. *a*.

Action: To draw the Os Hyoides towards the Chin, when the Jaws are shut; or the Chin towards the Os Hyoides, when the latter is fixed by the Muscles which come from the Sternum.

GENIO-HYO-GLOSSUS.

Origin: From the same Tubercle with the former Muscle; its Fibres spreading out like a Fan.

Insertion: Into the whole length of the Tongue, and Base of the Os Hyoides. Tab. LXVIII. No. 57.

Action: According to the direction of its Fibres,—to draw the Tongue forwards, or backwards,—to pull it downwards, and render its Dorsum concave,—and to make the Os Hyoides advance towards the Chin.

HYO-GLOSSUS.

Origin: From the whole length of one half of the Os Hyoides; running upwards, and a little outwards.

Insertion: Into the side of the Tongue, near the Stylo-Glossus. Tab. XLVII. Fig. 3. *e*.

Action: To depress the edge of the Tongue, and thereby render its upper Surface convex.

LINGUALIS.

Origin: From the root of the Tongue, laterally. It advances between the Genio-Hyo-Glossus and Hyo-Glossus.

Insertion: Into the tip of the Tongue. Tab. XLVII. Fig. 4. *k*.

Action: To raise the point of the Tongue; to contract its substance, and bring it backwards.

STYLO-GLOSSUS.

Origin: From the Styloid Process of the Temporal Bone, and from a Ligament which connects that Process to the angle of the Lower Jaw. It goes downwards and forwards, and is of a slender form.

Insertion: Into the root of the Tongue, near the Hyo-Glossus; and, running along its side, is insensibly lost near the Apex. Tab. XLVII. Fig. 2. *d*.

Action: To draw the Tongue backwards, and to one side.

STYLO-HYOIDEUS.

Origin: From the under half of the Styloid Process. It goes downwards and forwards, splitting for the passage of the Digastricus.

Insertion: Into the Os Hyoides, at the junction of the Base and Cornu. Tab. XLIX. Fig. 1. *d*.

Action: To pull the Os Hyoides to one side, and a little upwards.

STYLO-HYOIDEUS ALTER.

When present, it is a more slender Muscle than the former; but, like it, has nearly the same Origin, Insertion, and Action.

STYLO-PHARYNGEUS.

Origin: From the root of the Styloid Process;—it goes downwards and forwards.

Insertion: Into the side of the Pharynx, along which it expands.—It is also fixed to the back part of the Thyroid Cartilage. Tab. XLVII. Fig. 8. *g, b, i, k, l*.

Action: To dilate and raise the Pharynx, and thereby prepare it to receive the Morfel from the Mouth. It at the same time elevates the Thyroid Cartilage.

CIRCUMFLEXUS PALATI, vel *Tensor Palati*.

Origin: From the Spinous Process of the Sphenoid Bone, from the Osseous and Cartilaginous parts of the Eustachian Tube, and from the root of the Internal Pterygoid Process. It runs along the Pterygoideus Internus, passes over the Hook of the Internal Plate of the Pterygoid Process, and playing on it by a round Tendon, as on a Pulley, it spreads out into a broad Membrane.

Insertion: Into the Velum Palati, and femilunar edge of the Os Palati, extending as far as the Suture which joins the two Bones. Generally some of its posterior Fibres join the Constrictor Pharyngis Superior and Palato-Pharyngeus. Tab. XLVII. Fig. 14. *b, c*.

Action: To depress and stretch the Velum.

LEVATOR PALATI, vel *Levator Palati Mollis*.

Origin: From the point of the Pars Petrofa of the Temporal Bone, and also from the Membranous Portion of the Eustachian Tube. From these parts it descends.

Insertion: By a broad Expansion, into the Velum Palati, extending as far as the root of the Uvula, and uniting with its fellow. Tab. XXVII. Fig. 14. *a*.

Action: To raise the Velum in the time of Swallowing, and prevent the food or drink from passing into the Nose,

Nose, by pressing the Velum against the back part of the Nostrils.

CONSTRUCTOR ISTHMI FAUCIUM.

Origin : From the side of the root of the Tongue. It runs in the doubling of the Skin, which forms the anterior Arch of the Palate.

Insertion : Into the middle of the Velum Palati, at the root of the Uvula, where it is connected with its fellow.

Action : To draw the Palate and root of the Tongue towards each other, and thereby to shut the Opening into the Fauces.

PALATO-PHARYNGEUS.

Origin : From the middle of the Velum Palati, at the root of the Uvula; and from the insertion of the Constrictor Isthmi Faucium and Circumflexus Palati. The Fibres proceed within the posterior Arch of the Palate, and run to the upper and lateral part of the Pharynx, where they spread, and mix with those of the Stylo-pharyngeus.

Insertion : Into the Edge of the upper and back part of the Thyroid Cartilage; some of its Fibres being lost between the Membrane and inferior Constrictors of the Pharynx. Tab. XLVII. Fig. 11. c, c.

Action : To draw the Velum and Uvula downwards; the Larynx and Pharynx being at the same time raised; along with the Constrictor Superior and Tongue, to assist in shutting the passage into the Nostrils, and, in Swallowing, to convey the food from the Fauces into the Pharynx.

The SALPINGO-PHARYNGEUS of ALBINUS is composed of a small portion of the former Muscle, which arises from the Eustachian Tube, and which, when acting, may affect it. Tab. XLVII. Fig. 11. e, e.

AZYGOS UVULÆ.

Origin : From the posterior extremity of the longitudinal Palate Suture. It runs in the middle of the Velum Palati, and goes through the whole length of the Uvula, adhering in its passage to the Circumflexi.

Insertion : Into the point of the Uvula. Tab. XLVII. Fig. 10. b.

Action : To shorten the Uvula.

MUSCLES SITUATED UPON THE BACK PART OF THE PHARYNX.

CONSTRUCTOR PHARYNGIS INFERIOR.

Origin : From the sides of the Thyroid and Cricoid Cartilages. The superior Fibres running obliquely upwards, cover the under part of the following Muscle, and terminate in a point; the inferior Fibres run more transversely, and cover the beginning of the Œsophagus.

Insertion : Into its fellow, by the medium of a longitudinal Tendinous line in the middle of the back part of the Pharynx. Tab. XLVII. Fig. 6. a, b.

Action : To compress the lower part of the Pharynx, and to draw it and the Larynx a little upwards.

CONSTRUCTOR PHARYNGIS MEDIUS.

Origin : From the Appendix and Cornu of the Os Hyoides, and also from the Ligament which connects the Cornu to the Thyroid Cartilage. In its passage, it spreads out, and terminates in a point above and below; the upper part covering the following Muscle.

Insertion : Into the Cuneiform Process of the Occipi-

tal Bone, before the Foramen Magnum, and to its fellow on the opposite side by a Tendinous Line, in a similar manner to the former Muscle. Tab. XLVII. Fig. 7. a, b.

Action : To compress the middle and upper part of the Pharynx.

CONSTRUCTOR PHARYNGIS SUPERIOR.

Origin : From the Cuneiform Process of the Occipital Bone, before the Foramen Magnum; from the Pterygoid Process of the Sphenoid Bone, and from both Jaws, near the last Dentures Molares: It is likewise connected with the Buccinator, and with the root of the Tongue and Palate. From these Origins, it runs almost horizontally.

Insertion : Into its fellow, by the intervention of a Tendinous line, as in the former Muscle. Tab. XLVII. Fig. 8. a, d.

Action : To compress the upper part of the Pharynx, and, with the assistance of the other Constrictors, to thrust the Food down into the Œsophagus.

MUSCLES OF THE GLOTTIS.

CRICO-ARYTENOIDEUS POSTICUS.

Origin : Broad and Fleishy, from the back part of the Cricoid Cartilage.

Insertion : By a narrow extremity into the back part of the Base of the Arytenoid Cartilage. Tab. XLVII. Fig. 11. p.

Action : To pull back the Arytenoid Cartilage, by which

which the Ligament of the Glottis is made tense, and the Glottis itself longer.

CRICO-ARYTENOIDEUS LATERALIS.

Origin : From the side of the Cricoid Cartilage, where it is covered by the Thyroid.

Insertion : Into the side of the Base of the Arytenoid Cartilage. Tab. XLVII. Fig. 16. *b*.

Action : To open the Glottis, by separating the Arytenoid Cartilages, and, with them, the Ligaments of the Glottis.

THYRO-ARYTENOIDEUS.

Origin : From the under and back part of the middle of the Thyroid Cartilage, from which it runs backwards and a little upwards, in a double order of Fibres, upon the side of the Glottis and Ventricle of the Larynx.

Insertion : Into the fore-part of the Arytenoid Cartilage. Tab. XLIX. Fig. 7. *k*.

Action : To pull the Arytenoid Cartilage outwards and forwards, and so to widen the Glottis, and relax its Ligaments. It may also affect the Ventricle of the Larynx.

A small slip, termed by ALBINUS *Thyro-Arytenoides Alter Minor*, arises from the upper and back part of the middle of the Thyroid Cartilage, and is inserted into the Arytenoid Cartilage, above the insertion of the Crico-Arytenoideus Lateralis. *Use* : To assist the former in shortening and relaxing the Ligaments of the Glottis.

ARYTENOIDEUS OBLIQUUS, vel *Minor*.

Origin : From the root of one of the Arytenoid Cartilages; crossing its fellow obliquely.

Insertion : Near the point of the other Arytenoid Cartilage. Tab. XLVII. Fig. 11. *n*.

Action : To draw the Arytenoid Cartilages towards each other, and assist in closing the Aperture of the Glottis.

Frequently one of the oblique Arytenoid Muscles is a-wanting.

ARYTENOIDEUS TRANSVERSUS, vel *Major*.

Origin : From almost the whole length of the back part of one of the Arytenoid Cartilages, and runs transversely.

Insertion : In a similar manner, into the other Arytenoid Cartilage. Tab. XLVII. Fig. 11. *o*.

Action : To close the Glottis, by drawing together the two Arytenoid Cartilages and the Ligaments of the Glottis.

THYRO-EPIGLOTTIDEUS.

Origin : By a few scattered Fibres, from the Thyroid Cartilage.

Insertion : Into the side of the Epiglottis. Tab. XLVII. Fig. 16. *c, e, f*.

Action : To assist its fellow in drawing the Epiglottis towards the Glottis.

ARYTENO-EPIGLOTTIDEUS.

Origin : By a number of small Fibres from the Arytenoid Cartilage. It runs along the outer side of the external Opening of the Glottis.

Insertion : Into the Epiglottis, along with the former Muscle.

Action : To assist its fellow, in drawing the Epiglottis immediately down upon the Glottis.

It is counteracted by the elasticity of the Epiglottis.

The two last-mentioned Muscles are obscurely seen, excepting in robust Bodies.

MUSCLES SITUATED ON THE ANTERIOR AND LATERAL PARTS OF THE ABDOMEN.

OBLIQUUS DESCENDENS EXTERNUS,

Vel *Obliquus Externus Abdominis*.

Origin : In a serrated manner, from the lower edge of the eight inferior Ribs, near their Cartilages. The Serræ intermix with the Indentations of the Serratus Major Anticus, and it is commonly connected with the Pectoralis Major, Intercostales, and Latissimus Dorsi; the last of which covers the edge of a portion of it, extending from the twelfth Rib to the Spine of the Os Ilium, from the anterior half of which it has also part of its origin.— From these attachments, the Fibres of the Muscle run

obliquely downwards and forwards, and terminate (sometimes by distinct Indentations) in an Aponeurosis, which, near its margin, is firmly connected with the Aponeurosis of the following Muscle, Tab. XXXIV. Fig. 1. G, G, where it forms a curved line, called *Linea Semilunaris*. From this the Tendinous Fibres are continued in the same direction with the Fleishy Fibres, to the middle of the Abdomen.

Insertion : Into its fellow of the opposite side, by the medium of the *Linea Alba*, Tab. XXXIV. Fig. 1. E, E, which extends from the Cartilago Eniformis to the Pubes. This white Line is formed by the meeting of the Tendons

Tendons of the Oblique and Transverse Muscles of the Abdomen, and is perforated in the middle by the Umbilicus—originally a passage for the Umbilical Cord, and now formed into a Cicatrix. Tab. XXXIV. Fig. 1. F.

The Tendon of this Muscle is strengthened by other Tendons of a more delicate nature, which decussate it in a curved direction, upwards and inwards.

The under part of the Tendon, thicker and stronger than the rest of it, extends from the superior-anterior Spinous Process of the Os Ilium, over the Flexor Muscles and great Vessels and Nerves of the Thigh, to the upper part of the Os Pubis, to which it is fixed. Tab. XXXIV. Fig. 1. g.

This part of the Tendon, which was formerly known by the name of POUPART'S, or FALLOPIUS'S, or Inguinal Ligament, forms a curve behind, over the Blood-vessels, and therefore is now known by the name of *Crural Arch*.

Somewhat higher, and farther out, than the Symphysis Pubis, or about an inch and a half in a full-sized Adult, the Crural Arch divides into an upper and under column.

The upper is fixed to the Symphysis Pubis, and to the Os Pubis of the opposite side. The under is twisted or doubled in, and inserted into the upper part of the Os Pubis, from the Femoral Vessels, as far as the Crest of the Bone, forming the Crural Ring, of late so frequently mentioned by Surgeons.

The under column is loose, and more slender, in the Female than in the Male; and the space between the Femoral Vessels and the insertion of this part of the Ligament is larger; in consequence of which, Protrusions of the Bowels happen here more frequently in Women.

Where the columns separate, a space of an oval form is left, about an inch in length in the Male, but less in the Female; the direction running upwards and outwards, or somewhat in a line between the Pubis and Spine of the Ilium. This is the Ring of the External Oblique Muscle, Tab. XXXIV. Fig. 1. I, for the transmission of the Spermatic Cord in the Male, and the round Ligament of the Uterus in the Female.

Surrounding the exit of the Cord, or the round Ligament, from the Ring, there is a quantity of Cellular Substance, and some tendinous Fibres, which assist in filling that passage, and in preventing any communication between the outer and inner parts.

The place where the columns separate to form the Ring varies in different Subjects. In some, the separation is considerably farther out than the part already described, though more generally the division is at the outer end of the Ring. At this end of the Ring, the Columns are joined by Tendinous Fibres, which arise from the Os Ilium, and from the Inguinal Ligament; and are part of the Fibres mentioned above, as decussating the Tendon of the External Oblique Muscle.

From the under end of the Inguinal Ligament, and from the Offa Pubis, an Aponeurosis descends and forms part of the general Fasciæ of the Thigh. This Expansion stretches the Ligament in the extended state of the Thigh, but allows it to be relaxed when the Limb is bent, or when the Aponeurosis, by being cut across, loses hold of the Thigh.

Besides the Aponeurosis, there is a thin expansion which descends upon the Abdominal Muscles, and vanishes in the Fat, and over the Inguinal Glands. Part of it also descends over the Cord, as far as the Scrotum.

At the upper and fore-part of the Thigh, the Aponeurosis sends off a process, which passes over the Femoral Vessels, and is fixed to the Os Femoris, at their inner side.

Through the Abdominal Ring, there is no direct opening into the Cavity of the Abdomen; the passage being shut by the Obliquus Internus, and Transversalis Abdominis.

OBLIQUUS ASCENDENS INTERNUS,

Vel *Obliquus Internus Abdominis.*

Origin: From the back part of the Os Sacrum;—from the Spinous Processes of the three lowest Lumbar Vertebrae, by a Tendon common to it and the Serratus Posticus Inferior;—from the whole length of the Spine of the Os Ilium;—and from the inside of POUPART'S Ligament, at the middle of which it sends off the Cremaster. From these Origins the Fibres are disposed in a radiated manner; but the greater part of them run in a slanting direction upwards. At the Linea Semilunaris, the Muscle becomes Tendinous, and adheres firmly to the Tendon of the Obliquus Externus. Here its Tendon divides into two Layers: The anterior Layer, with the greater part of the inferior portion of the posterior Layer, joins the Tendon of the External Oblique, and goes over the Rectus, to be inserted into the whole length of the Linea Alba: The posterior Layer joins the Tendon of the Transversalis, and goes behind the Rectus; and this union is continued down, till it reaches about half-way between the Umbilicus and Os Pubis. Lower than this, only a few scattered Fibres of the posterior Layer are to be found behind the Rectus; the principal part of it passing before that Muscle, to be inserted into the Linea Alba.

Insertion: Into the Cartilages of all the False Ribs, into the Cartilago Ensisiformis, and whole length of the Linea Alba. Tab. XXXV. Fig. 1. Trunk, *f, i, k, l.*

Action: To assist the former Muscle. It bends the Body, however, in the same direction with the Obliquus Externus of the opposite side.

TRANSVERSALIS,

TRANSVERSALIS, vel *Transversus Abdominis*.

Origin: Flethy from the inner Surface of the Cartilages of the six or seven Lower Ribs, where it intermixes with the Digitations of the Diaphragm, and with the Intercostal Muscles; from the Transverse Processes of the twelfth Dorsal and four superior Lumbar Vertebrae; from the whole inner edge of the Spine of the Os Ilium; and anterior to this, it is connected to the under edge of the Obliquus Externus. At the Linea Semilunaris, the Muscle becomes Tendinous, and the Tendon is continued across, adhering to the Obliquus Internus in the manner already mentioned.—In the whole of its course, it is closely connected to the Surface of the Peritoneum.

Insertion: Into the Cartilago Eniformis, and Linea Alba. Tab. XXXVI. Fig. 1. Trunk, C, D, E.

Action: To support, and immediately to compress the Abdominal Bowels.

From the inside of the Crural Arch, and from the Spine of the Ilium, a Tendinous Aponeurosis is sent off, which is reflected over the Iliacus Internus, and part of the Psoas Magnus, which it braces and protects.

From the Crural Arch, and reflected also from the under part of the Expansion covering the Iliacus Internus, another Aponeurotic Expansion is sent upwards, which lines the inner side of the Transversalis, lies between it and the Peritoneum, and becomes gradually thinner in its ascent.

The Angle of reflection between these two Expansions being formed of strong Tendinous Fibres, the Abdomen at this place is fortified, and the Bowels are prevented from protruding between the Spine of the Ilium and the Iliac Blood-vessels.

From the Aponeurosis of the Internal Oblique Muscle, a Process descends behind the External Iliac Vessels, to be inserted at the outer edge of the Pectineus. This, with the Process sent off from Poupart's Ligament already taken notice of, forms a Sheath which incloses the upper part of the Femoral Vessels.

Half-way between the Spine of the Ilium and Symphyfis Pubis, the Expansion lining the Transversalis leaves an opening for the passage of the Spermatic Cord, or for the round Ligament of the Uterus; the beginning of which passage may be considered as the Internal Abdominal Ring.

The under part of this opening is formed of Poupart's Ligament, strictly so called; the upper by the Transversalis and Obliquus Internus.

From this opening there is no direct passage outwards, the part being shut by the Tendon of the Obliquus Externus.

The inner Ring is of the same form and size with the outer Ring, and is directed in the same manner with it.

Between the Internal and External Abdominal Rings, the passage is oblique, like the Rings themselves, and has also a quantity of Cellular Substance, which is considered by some Authors as forming a distinct Canal. This surrounds the Cord, or the round Ligament, and assists these in completely filling the whole of this passage.

Between the inner part of the External Iliac Vein and the intortion of the under column of Poupart's Ligament into the Os Pubis, a triangular space is left, at the outer part of which there is a small aperture occupied by Cellular Substance. The triangular Cavity and Aperture are more considerable in the Female than in the Male. Through these the Bowels protrude in Femoral Hernia.

RECTUS.

Origin: Tendinous from the fore and upper part of the Symphyfis Pubis. It soon becomes Flethy, and runs upwards in form of a flat Band, the whole length of, and parallel to, the Linea Alba. In its course, it is divided by three Tendinous Intersections, at and above the Umbilicus; and there is generally a half intersection below it. These seldom penetrate through the whole thickness of its Substance. They adhere firmly to the anterior part of the Sheath which incloses the Muscle, but slightly to the posterior Layer.

Insertion: Into the Cartilages of the three inferior True Ribs and extremity of the Sternum. It frequently intermixes with the under edge of the Pectoralis Major. Tab. XXXV. Fig. 1. D, D.

Action: To compress the fore-part of the Abdomen; to draw down the Ribs in Expiration; and to bend the Body forwards, or to raise the Pelvis. By means of its Sheath and Tendinous Intersections, it is kept in its place, and allowed to act more equally.

PYRAMIDALIS.

Origin: By a broad Base, from the upper part of the Symphyfis Pubis. It runs upwards within the same Sheath with the Rectus, tapering to a point in its ascent.

Insertion: Near half-way between the Pubis and Umbilicus, in the Linea Alba and inner edge of the Rectus. Tab. XXXIV. Fig. 1. p.

Action: To assist the under part of the Rectus in drawing down the Ribs, or to compress the under part of the Abdomen.

It is frequently a-wanting in both sides, and then the under end of the Rectus is larger, thus in some measure supplying its place.

MUSCLES OF THE MALE PARTS OF GENERATION, AND OF THE ANUS.

CREMASTER, vel *Musculus Testis*.

Origin: From the under edge of the Obliquus Internus Abdominis. Passing through the Ring of the Obliquus Externus, it surrounds the Spermatic Cord as far as the Testicle, where the Fibres separate and expand.

Insertion: Into the Tunica Vaginalis Testis, and Cellular Substance of the Scrotum. Tab. XXXIV. Fig. 1. K.

Action: To suspend and elevate, and to compress and evacuate the Testicle.

ERECTOR PENIS, vel *Ischio-Cavernosus*.

Origin: Tendinous from the inner side of the Tuberosity of the Os Ischium.—It runs upwards, fleshy, increasing in breadth, and embracing the whole Crus Penis.

Insertion: By a thin Tendon into the Elastic Membrane which covers the Corpora Cavernosa Penis, as far as the union of the Crura. Tab. XLVIII. Fig. 6. d.

Action: To compress the Crus Penis, by which means the Blood is pushed from it into the fore-part of the Corpora Cavernosa, and the Penis is thereby more completely distended.

ACCELERATOR URINÆ, vel *Ejaculator Seminis*.

Origin: Fleshy from the Sphincter Ani, and Membranous part of the Urethra; and Tendinous from the Crus and beginning of the Corpus Cavernosum Penis.—In its course it forms a thin fleshy Layer, the inferior Fibres of which run more transversely than the superior, which descend in an oblique direction; the Muscles on the opposite sides completely inclosing the Bulb of the Urethra.

Insertion: Into its fellow, by a Tendinous line running longitudinally on the middle of the Bulb. Tab. XLVIII. Fig. 6. a.

Action: To propel the Urine or Semen forwards, and, by compressing the Bulb, to push the Blood into, and thereby distend the Corpus Cavernosum Urethrae and Glans Penis.

TRANSVERSUS PERINEI, vel *Transversalis Urethrae*.

Origin: From the inside of the Tuberosity of the Os Ischium, close to the Erector Penis; running transversely.

Insertion: Into the back part of the Accelerator Urinae, and adjoining part of the Sphincter Ani. Tab. XLVIII. Fig. 6. b.

Action: To dilate the Bulb for the reception of the Semen or Urine; and to assist the Levator Ani in retracting the Anus, after the discharge of the Faeces.

There is frequently another Muscle, termed *Transversalis Perinei Alter*, running along with the former, and having nearly the same Origin, Insertion, and Action, but going more obliquely upwards.

SPHINCTER ANI.

Origin: From the extremity of the Os Coccygis, running forwards within the Skin and Fat which cover the verge of the Anus, and in its passage forming a broad, flat, oval Muscle, which surrounds the extremity of the Intestinum Rectum.

Insertion: By a narrow point, into the Acceleratores Urinae and Transversi Perinei. Tab. XLVIII. Fig. 9. a.

Action: To shut the Anus, and also to pull down the Bulb of the Urethra, by which it assists in ejecting the Urine and Semen.

The *Sphincter Internus* of some Authors is merely the Circular Muscular Coat of the end of the Rectum.

LEVATOR ANI.

Origin: By a semicircular edge, from the Os Pubis, within the Pelvis, at the upper edge of the Foramen Thyroideum; from the Aponeurosis which covers the Obturator Internus and Coccygeus; and from the Spinous Process of the Os Ischium.—Its Fibres descend like rays from a circumference, to meet those of its fellow, and with it to form a kind of inverted Funnel.

Insertion: Into the Sphincter Ani, Accelerator Urinae, and under and fore part of the Os Coccygis.—It surrounds the extremity of the Rectum, neck of the Bladder, Prostate Gland, and part of the Vesiculæ Seminales. Tab. XLVIII. Fig. 9. b.

Action: To support the contents of the Pelvis; to retract the end of the Rectum, after the evacuation of the Faeces; and to assist in the evacuation of the Rectum, Bladder, Vesiculæ Seminales, and Prostate Gland.—It is likewise considered by some as a principal agent in the distension of the Penis, by pressing upon its Veins.

Part of the Levator Ani, which arises from the Os Pubis, between the lower part of the Symphysis and the upper part of the Foramen Ovale, and assists in inclosing the Prostate Gland, is called by SOEMMERING *Compressor Prostatæ*.

MUSCLES OF THE FEMALE PARTS OF GENERATION, AND OF THE ANUS.

ERECTOR CLITORIDIS.

Origin: As in the Erector Penis in the Male, but the Muscle smaller.

Insertion: Into the Crus and Body of the Clitoris.

Action: To draw the Clitoris downwards and backwards; and by pushing the Blood into it from its Crus, it may render the Body of the Clitoris more tense.

SPHINCTER VAGINÆ.

Origin: From the Sphincter Ani, and, near the Perineum, from the posterior side of the Vagina. It passes along the outer end of the Vagina, covers the Corpus Cavernosum Vaginæ; going behind the Nymphæ.

Insertion: Into the union of the Crura Clitoridis.—See Lateral View of Female Parts of Generation.

Action: To contract the external Orifice of the Vagina, by compressing its Corpus Cavernosum, from which it likewise pushes the blood into the Nymphæ and Clitoris.

TRANSVERSUS PERINEI.

Origin: As in the Male.

Insertion: Into the upper part of the Sphincter Ani, and into a tough white Substance in the Perineum.

Action: Upon the Perineum and Anus, as in the Male.

SPHINCTER ANI.

Origin and course, as in the Male.

Insertion: Into the tough white Substance in the Perineum.

Action: To shut the Anus, and, by pulling down the Perineum, to assist in contracting the external Orifice of the Vagina.

LEVATOR ANI.

Origin: As in the Male. In its descent, it embraces the inferior part of the Vagina and Rectum.

Insertion: Into the Perineum, Sphincter Ani, extremity of the Vagina, and Rectum.

Action: Upon the Bladder and Rectum, as in the Male.—It also assists in supporting and contracting the Vagina, and may, by pressing upon the Veins, contribute to the distension of the Cells of the Clitoris and Corpus Cavernosum Vaginæ.

MUSCLE OF THE OS COCCYGIS.

COCCYGEUS.

Origin: By a narrow point, from the Spinous Process of the Os Ischium.—In its passage, it gradually expands, and covers the inside of the posterior Sacro-Ischiatic Ligament.

Insertion: Into the whole length of the side of the Os Coccygis, and into the under end of the Os Sacrum. Tab. XLVIII. Fig. 3.

Action: To move the Os Coccygis forwards, by which it assists the Levator Ani in supporting or raising the end of the Rectum.

MUSCLES SITUATED WITHIN THE CAVITY OF THE ABDOMEN.

DIAPHRAGMA.

The Diaphragm forms a Flethy and Tendinous Partition, which separates the Cavity of the Abdomen from that of the Thorax, and is perforated by several Holes, for the passage of Vessels and Nerves which go into, or come out from the Abdomen. It is concave below, and convex above; the middle of it reaching as high within the Thorax as the fourth pair of Ribs. Above, it is covered by the Pleura, and below, by the Peritoneum; and is commonly divided into two portions, called Superior and Inferior Muscles of the Diaphragm.

SUPERIOR, or Greater Muscle of the Diaphragm.

Origin: By Flethy Indentations, from the Cartilago Eniformis, and from the Cartilages of the seventh, and of all the inferior Ribs on both sides. From these different Origins, the Fibres run in a radiated manner.

Insertion: Into a Cordiform Tendon, placed in the middle of the Diaphragm, and in which the Fibres of the opposite sides are interlaced.—Towards the right side, the Tendon is perforated by a triangular Hole for the passage of the Vena Cava Inferior; and to the upper convex

convex part of it, the Pericardium and Mediastinum are connected.

INFERIOR, or LESSER MUSCLE, or *Appendix of the Diaphragm.*

Origin: By four Pair of Heads, of which one Pair in the middle, commonly called its Long, or Tendinous *Crura*, is the longest. The long *Crura* arise from the fore-part of the fourth Lumbar Vertebra, and adhere to the Bodies of all the Vertebrae of the Loins above this, by the intervention of the Ligamentum Commune Anteriorius covering these Bones. In their ascent, they leave an oval opening for the passage of the Aorta and Thoracic Duct. The other Heads arise from the third, and also from the second Lumbar Vertebra, and are placed farther out. From the different Heads the Muscular Fibres run upwards, and form, in the middle, two Fleishy Columns, or *Crura*, which decussate, and leave an opening for the passage of the Œsophagus.

Insertion: By strong Fleishy Fibres, into the posterior edge of the Cordiform, or middle Tendon. Tab. XLVIII. Fig. 2.

Action: To enlarge the Cavity of the Thorax in Inspiration, by its Fleishy part contracting, and bringing its two sides down from a convex to a plane Surface; the Abdominal Muscles at the same time yielding, but the Tendinous part of the Diaphragm remaining nearly in the same situation. In Expiration, the Diaphragm is replaced, chiefly by the action of the Abdominal Muscles. It is the Antagonist of the Abdominal Muscles in Inspiration, but acts in concert with them in Dejection and Vomiting.

QUADRATUS LUMBORUM.

Origin: Broad, Tendinous, and Fleishy, from the posterior half of the Spine of the Os Ilium, and from a Ligament extended between it and the transverse Process of the last Lumbar Vertebra.

Insertion: Into the transverse Processes of all the Lumbar Vertebrae; into the last Rib, near the Spine; and, by a small Tendon, into the side of the last Dorsal Vertebra. Tab. XXXVII. Trunk, D.

Action: To move the Loins to one side, to pull down the last Rib in laborious expiration, and, when both act, to bend the Loins forwards.

MUSCLES SITUATED UPON THE ANTERIOR PART OF THE THORAX.

PECTORALIS MAJOR, vel *Pectoralis.*

Origin: From the Sternal half of the Clavicle; from the fore-part of the edge of almost the whole length of the upper and middle bone of the Sternum, where it is connected with its fellow; and from the Cartilages of

PSOAS PARVUS.

Origin: Fleishy, from the side of the last Vertebra of the Back, and from that of one or two of the upper Vertebrae of the Loins. It sends off a slender Tendon which runs down by the inner side of the Psoas Magnus, and an Aponeurosis which expands upon the neighbouring Muscles.

Insertion: Into the Brim of the Pelvis, at the joining of the Ilium and Pubis. Tab. XXXVII. Fig. 1. C, S.

Action: To assist in bending the Spine upon the Pelvis, and in particular positions, in raising the Pelvis.

This Muscle is frequently wanting.

PSOAS MAGNUS.

Origin: From the side of the Bodies, and from the transverse Processes of the last Dorsal, and of all the Lumbar Vertebrae, by an equal number of Fleishy Slips, which, uniting, form a thick strong Muscle, which bounds the upper part of the side of the Pelvis; passing down over the Os Pubis, behind *POUPART'S Ligament.*

Insertion: Tendinous and Fleishy, into the Trochanter Minor, and part of the Body of the Os Femoris. Tab. XXXVII. Fig. 1. A.

Action: To bend the Thigh, and turn it a little outwards, or, when the Inferior Extremity is fixed, to assist in bending the Body.

ILIACUS INTERNUS.

Origin: Fleishy, from the Transverse Process of the last Lumbar Vertebra; from all the inner edge of the Spine of the Os Ilium; from the edge of that Bone, between its anterior-superior Spinous Process and the Acetabulum, from most of the hollow part of the Os Ilium, and from an Aponeurosis covering it, which is sent up from the Spine of the Bone and inner side of *POUPART'S Ligament.* It joins the Psoas Magnus, where it begins to become Tendinous on the Os Pubis.

Insertion: Along with the Psoas Magnus. Tab. XXXVII. Fig. 1. B.

Action: To assist the Psoas in bending the Thigh.

the fifth and sixth Ribs, where it mixes with the Obliquus Externus. The Fibres from thence converge towards the Axilla, where they decussate, and send off a flat twisted Tendon.

Insertion: Into the Ridge at the outer edge of the Groove

Groove for lodging the Tendon of the long Head of the Biceps. Tab. XXXIV. Fig. 1. A.

Action : To draw the Arm towards the Sternum.

Between the Portions of the Muscle arising from the Clavicle and Sternum, there is a slight Separation, in consequence of which these Portions have been considered by some Authors as two distinct Muscles.

PECTORALIS MINOR,

Vel Serratus Minor Anticus.

Origin : Tendinous and Flethy, in a serrated manner, from the third, fourth, and fifth Ribs, near their Cartilages. Passing obliquely outwards, it becomes gradually narrower.

Insertion : Tendinous, into the point of the Coracoid Process of the Scapula. Tab. XXXV. Fig. 1. B.

Action : To bring the Scapula downwards and forwards, or in laborious respiration to raise the Ribs.

SUBCLAVIUS.

Origin : Tendinous, from the Cartilage of the first Rib. It soon becomes Flethy, and runs outwards, under the Clavicle, increasing in breadth.

Insertion : Into the under Surface of the Clavicle, from near its Head, as far outwards as the Coracoid Process of the Scapula. Tab. XXXV. Fig. 1. A.

Action : To pull the Clavicle, and with it the Scapula, downwards and forwards.

SERRATUS MAGNUS,

Vel Serratus Major Anticus.

Origin : From the nine superior Ribs, by an equal number of Flethy Digitations. It runs obliquely upwards and backwards upon the side of the Thorax, and between it and the Subscapularis.

Insertion : Flethy, into the whole length of the Base of the Scapula, and in a manner folded round it, between the Insertion of the Rhomboides and the Origin of the Subscapularis. Tab. XXXV. Fig. 1. C, C. Tab. XLI. Fig. 1. E.

Action : To move the Scapula forwards or downwards, according to the direction of its different Digitations, and when the Scapula is forcibly raised, as in violent inspiration, to assist in dilating the Thorax, by elevating the Ribs.

MUSCLES SITUATED BETWEEN THE RIBS, AND WITHIN THE THORAX.

INTERCOSTALES EXTERNI.

Origin : From the under edge of each Rib, excepting the twelfth. They run obliquely downwards and forwards, from the Spine to the joining of the Ribs with their Cartilages, from which, to the Sternum, they are discontinued; that place being occupied by an Aponeurosis.

Insertion : Into the upper edge of each Rib, immediately below that from which they take their respective Origins. Tab. XXXVI. Fig. 13. B, &c.

Portions of the Intercostales Externi, which arise from the Transverse Processes of the Vertebrae, and terminate in the Ribs immediately below, are termed by ALBINUS, *Levatores Costarum Breviares*, Tab. XLIII. Fig. 1. C, C. —Other Portions, which arise in the same manner, but pass over one Rib, and terminate in the next below it, are named by the same Author, *Levatores Costarum Longiores*. Tab. XLIII. Fig. 1. D, D.

INTERCOSTALES INTERNI.

Origin : The same with that of the Externi; but they begin at the Sternum, and run downwards and backwards, decussating the former Muscles like the strokes of the letter X, and continuing as far as the Angle of the Ribs, from which to the Spine they are wanting.

Insertion : In the same manner as the Externi. Tab. XLIII. Fig. 1. a, &c.

Portions of the Intercostales Interni, near the under part of the Thorax, which pass over one Rib, and terminate in the next below it, are called, by DOUGLAS, *Costarum Depressores Proprii*. Tab. XLIX. Fig. 8. L, L.

Action of the Intercostales Interni, as well as of the Externi: To enlarge the Cavity of the Thorax, by elevating the Ribs in the time of Inspiration; and the obliquity of the one set balancing that of the other, allows them to be raised more immediately upwards.

From the obliquity of their Fibres, they are found to possess a greater power in raising the Ribs, than Fibres going in a perpendicular direction.

The Intercostales Externi cease near the Sternum, and the Interni near the Spine, to admit the ready motion of the Ribs; for, had the former been continued to the Sternum, and the latter to the Spine, the parts of these Muscles supposed to be thus fixed, would of course have become Antagonists to the rest.

The Portions called *Levatores* and *Depressores Costarum* assist in raising the Ribs, in the same manner as the rest of the Intercostales.

STERNO-

STERNO-COSTALIS, vel *Triangularis Sterni*.

Origin: From the edges of the Cartilago Eniformis, and lower half of the middle Bone of the Sternum, within the Thorax. It runs upwards and outwards, behind the Cartilages of the Ribs.

Insertion: Generally by three Angular Terminations into the Cartilages of the third, fourth, and fifth Ribs,

and sometimes, also, by a fourth Termination into the corresponding part of the Cartilage of the second or sixth Rib, near the union of the Cartilaginous with the Osseous part of the Ribs. Tab. XXXIX. Fig. 1. *i, i, &c.*

Action: To depress the Ribs into which they are fixed, and, of course, to assist in contracting the Cavity of the Thorax during Expiration.

MUSCLES SITUATED ON THE ANTERIOR PART OF THE VERTEBRÆ OF THE NECK.

LONGUS COLLI.

Origin: Tendinous and Flethy, from the side of the Bodies of the three superior Vertebrae of the Back, and from the Transverse Processes of the four inferior Vertebrae of the Neck.

Insertion: Into the fore-part of the Bodies of all the Vertebrae of the Neck, by as many small Tendons, which are covered with Flesh. Tab. XLIX. Fig. 8. D, E.

Action: It bends the Neck forwards and to one side, or, when both Muscles act, they immediately bend the Neck.

RECTUS CAPITIS ANTERIOR MAJOR,

Vel *Rectus Anterior Longus*.

Origin: From the fore-part of the Transverse Processes of the third, fourth, fifth, and sixth Vertebrae of the Neck. It runs upwards, and a little inwards, covering the outer edge of the Longus Colli.

Insertion: Into the Cuneiform Process of the Occipital

Bone, near its joining with the Os Sphenoides. Tab. XLIX. Fig. 8 C.

Action: To bend the Head forwards.

RECTUS CAPITIS ANTERIOR MINOR,

Vel *Rectus Anterior Minor*.

Origin: From the fore-part of the Atlas, opposite to its Superior Oblique Process. It runs obliquely inwards behind, and a little to the outside of the former Muscle.

Insertion: Into the Cuneiform Process of the Occipital Bone, immediately before the Condyles. Tab. XLIX. Fig. 8. B.

Action: To assist the Rectus Major.

RECTUS CAPITIS LATERALIS.

Origin: From the anterior part of the Transverse Process of the Atlas.—It goes obliquely outwards.

Insertion: Into the Occipital Bone, directly behind the Jugular Fossa. Tab. XLV. Fig. 14. C.

Action: To incline the Head a little to one side.

MUSCLES SITUATED UPON THE POSTERIOR PART OF THE TRUNK.

TRAPEZIUS, vel *Cucullaris*.

Origin: From the middle of the great arched Ridge of the Occipital Bone; from its fellow over the Spinous Processes of the Cervical Vertebrae, by the intervention of a strong Tendon, called *Ligamentum Nuchæ*, vel *Colli*; from the Spinous Processes of the two inferior Vertebrae of the Neck, and from all those of the Back, adhering Tendinous to its fellow the whole length of its Origin.

Insertion: Flethy, into the Scapular half of the Clavicle; Tendinous and Flethy, into the Acromion, and into the Spine of the Scapula. Tab. XL. Fig. 1. I, I.

Action: To move the Clavicle and Scapula, according to the directions of its different Fibres. The superior Fibres, descending, raise the Shoulder; the middle, running transversely, pull it backwards; and the inferior Fibres, ascending, depress it. The whole acting together bring it immediately back.—When the Scapula is fixed, the Muscle assists in moving the Head backwards.

LATISSIMUS DORSI.

Origin: By a broad Tendinous Expansion, from the posterior part of the Spine of the Os Ilium; from all the Spinous Processes of the Vertebrae extending between the under end of the Os Sacrum and sixth Dorsal Vertebra, and, by three or four Tendinous or Flethy Slips, from an equal number of inferior Ribs. The Tendon by degrees changes into a Muscle of great breadth, the inferior Fibres of which run upwards and outwards, and the superior transversely over the inferior Angle of the Scapula, receiving a small Slip from it in their way to the Axilla, where the Fibres of the Muscle in general are

are collected, twisted, and folded, like those of the Pectoralis Major.

Insertion: By a strong thin Tendon, into the inner edge of the Groove for lodging the Tendon of the long Head of the Biceps. Tab. XL. Fig. 1. K, K.

Action: To pull the Arm downwards and backwards, and to roll the Os Humeri inwards, by which the Palm of the Hand is made to face backwards. When the Pectoralis Major acts at the same time with this one, the Arm is brought immediately down towards the Trunk.

The Latissimus Dorsi and Pectoralis Major form the Axilla, in which the great Vessels and Nerves, and likewise the Glands, lie which belong to the Arm.

SERRATUS POSTICUS INFERIOR.

Origin: By the same common Tendon with the Latissimus Dorsi, from the two inferior Dorsal, and from the three Lumbar Vertebrae.

Insertion: By four Flethy Slips, into the same number of Inferior Ribs, near their Cartilages. Tab. XLI. Fig. 1. Trunk, D.

Action: To depress the Ribs into which it is inserted, and thereby, during Expiration, to assist in contracting the Cavity of the Thorax.

RHOMBOIDEUS.

Origin: Tendinous, from the Spinous Processes of the four or five Superior Dorsal, and the three Inferior Cervical Vertebrae, and from the Ligamentum Nuchae; descending obliquely.

Insertion: Into the whole length of the Base of the Scapula. Tab. XLI. Fig. 1. Trunk, AB.

Action: To draw the Scapula upwards and backwards.

This Muscle is frequently divided by an indistinct Line into two unequal Portions: The part, A, arising from the Dorsal Vertebrae, and fixed to the Base of the Scapula under the Spine, is commonly called *Rhomboides Major*, and the other part of the Muscle, B, *Rhomboides Minor*.

SPLenius.

Origin: Tendinous, from the four Superior Spinous Processes of the Dorsal, and Tendinous and Flethy, from the five Inferior of the Cervical Vertebrae. It adheres firmly to the Ligamentum Nuchae, and at the third Cervical Vertebra, it recedes from its fellow, so that part of the Complexus is seen.

Insertion: By as many Tendons, into the five Superior Transverse Processes of the Cervical Vertebrae, and by a Tendinous and Flethy Portion, into the posterior part of the Mastoid Process, and into the Os Occipitis, where it joins with that Process. Tab. XLI. Fig. 1. Neck, C.

Action: To antagonize the Sterno-mastoideus, by bringing the Head, and upper Cervical Vertebrae, obliquely backwards and to one side. When the Splenii act together, they draw the Head directly backwards.

This Muscle is divided by ALBINUS into *Splenius Capitis*, or that which arises from the Neck, and goes to the Head; and *Splenius Colli*, or that which arises from the Back, and is fixed to the Neck.

SERRATUS POSTICUS SUPERIOR.

Origin: By a broad thin Tendon, from the Ligamentum Nuchae, over the Spinous Processes of the three last Cervical, and two uppermost Dorsal Vertebrae; going obliquely downwards.

Insertion: By four Flethy Slips into the second, third, fourth, and fifth Ribs, under the upper and back part of the Scapula. Tab. XLI. Fig. 1. Trunk, c.

Action: To elevate the Ribs, and thus to dilate the Thorax in violent Inspiration.

SACRO-LUMBALIS.

Origin: In common with the Longissimus Dorsi, Tendinous without, and Flethy within, from the side, and all the Spinous Processes of the Os Sacrum; from the posterior part of the Spine of the Os Ilium; and from all the Spinous and Transverse Processes of the Lumbar Vertebrae. The common Head fills up the space between the Os Ilium and Os Sacrum, and also the hollow of the Loins. At the under part of the Thorax, the Muscle begins to send off Tendons, which lie flat upon the Ribs, and become gradually longer the nearer they are to the Spine.

Insertion: Into the Angles of all the Ribs, by an equal number of Tendons. Tab. XLII. Fig. 1. Trunk, C.

From six or eight of the lower Ribs arise an equal number of Flethy Portions, which terminate in the inner side of this Muscle, and get the name of *Musculi Accessorii*, vel *Additamentum ad Sacro-lumbalem*.

Action: To assist in raising and keeping the Trunk of the Body erect. It also assists the Serratus Inferior, and Quadratus Lumborum, in depressing the Ribs during laborious Expiration.

From the upper part of this Muscle, a Flethy Slip, called *Cervicalis Descendens*, runs up to be fixed to the Transverse Processes of the fourth, fifth, and sixth Cervical Vertebrae, by three distinct Tendons. It turns the Neck obliquely backwards and to one side. Tab. XLII. Fig. 1. Neck, L.

LONGISSIMUS DORSI.

Origin: In common with the Sacro-lumbalis. It forms a large, thick, and strong Muscle, which fills the Hollow

Hollow between the Spine and Angles of the Ribs; becoming gradually smaller in its ascent.

Insertion: Into the Transverse Processes of all the Dorsal Vertebrae, chiefly by small double Tendons; and, by a Tendinous and Fleishy Slip, into the lower edge of each of the Ribs, excepting the two inferior, near their Tubercles. Tab. XLII. Fig. 1. Trunk, B.

Action: To extend the Trunk, and keep it erect; the outer part may assist in depressing the Ribs during laborious Expiration.

From the upper part of this Muscle, a round Fleishy Slip runs up to join the Cervicalis Descendens.

SPINALIS DORSI.

Origin: By five Tendinous Slips, from the Spinous Processes of the two upper Lumbar, and the three lower Dorsal Vertebrae. In its ascent, it is incorporated with the Longissimus Dorsi.

Insertion: Into the Spinous Processes of the eight or nine uppermost Dorsal Vertebrae, excepting the first, by as many Tendons. Tab. XLII. Fig. 1. Trunk, A.

Action: To fix the Vertebrae, and to assist in extending the Trunk, and keeping it erect.

COMPLEXUS.

Origin: By distinct Tendons, from the Transverse Processes of the seven Superior Dorsal, and four Inferior Cervical Vertebrae, and by a Fleishy Slip, from the Spinous Process of the first Dorsal Vertebra. In its passage upwards, it is intermixed with Tendinous and Fleishy parts.

Insertion: Into a Depression, under the large arched Ridge of the Occipital Bone. Tab. XLII. Fig. 1. Neck, c, d.

Action: To draw the Head backwards, and to one side; and when both act, to draw the Head directly backwards.

The long Portion of this Muscle, which lies next the Spinous Processes, is more loose than the rest, and has a roundish Tendon in the middle of it, with a Fleishy Belly at each end, on which account it is called, by ALBINUS, *Biventer Cervicis*.

TRACHELO-MASTOIDEUS,

Vel Complexus Minor, vel Mastoideus Lateralis.

Origin: From the Transverse Processes of the three uppermost Dorsal, and five lowest Cervical Vertebrae, where it is connected to the *Transversalis Cervicis*, by as many thin Tendons, which unite into a slender Belly, and run up under the Splenius.

Insertion: Into the posterior Margin of the Mastoid Process, by a thin Tendon. Tab. XLII. Fig. 1. Neck, C.

Action: To assist the Complexus; but pulling the Head more laterally.

LEVATOR SCAPULÆ,

Vel Levator Proprius, vel Musculus Patientiæ.

Origin: From the Transverse Processes of the five Superior Cervical Vertebrae, by the same number of distinct Heads, which soon unite to form a flat Muscle, running downwards and outwards.

Insertion: Into the Superior Angle of the Scapula. Tab. XLI. Fig. 1. A, d.

Action: To pull the Scapula upwards, and a little forwards, as in shrugging the Shoulder; and, when the Scapula is fixed, to pull the Neck a little to one side.

SEMI-SPINALIS DORSI,

Vel Transverso-spinalis Dorsi.

Origin: From the Transverse Processes of the seventh, eighth, ninth, and tenth Dorsal Vertebrae, by as many distinct Tendons, which soon grow Fleishy, and then again become Tendinous.

Insertion: Into the Spinous Processes of the six or seven uppermost Dorsal, and two lowest Cervical Vertebrae, by as many Tendons. Tab. XLIII. Fig. 1. A, A.

Action: To extend the Spine obliquely backwards.

MULTIFIDUS SPINÆ,

Formerly described as three distinct Muscles, viz. *Transverso-Spinalis Lumborum, Transverso-Spinalis Dorsi,* and *Transverso-Spinalis Colli.*

Origin: From the side and Spinous Processes of the Os Sacrum, and from that part of the Os Ilium which joins with the Sacrum; from all the Oblique and Transverse Processes of the Lumbar Vertebrae; from all the Transverse Processes of the Dorsal, and of the four Inferior Cervical Vertebrae, by as many distinct Tendons, which soon become Fleishy, and run obliquely upwards and inwards.

Insertion: By distinct Tendons, into all the Spinous Processes of the Lumbar, Dorsal, and Cervical Vertebrae, excepting the Atlas. Tab. XLIII. Fig. 1. Trunk, B, B.

Action: To extend the Spine obliquely, and pull it to a side. When both Muscles act, they draw the Spine directly backwards.

SEMI-SPINALIS COLLI,

Vel Transverso-Spinalis Colli.

Origin: From the Transverse Processes of the six uppermost Dorsal Vertebrae, by an equal number of distinct

unct Tendons, which run obliquely under the Complexus.

Insertion: Into the Spinous Processes of all the Cervical Vertebrae, except the first and last. Tab. XLIII. right side of Neck, I, I.

Action: To extend the Neck obliquely backwards, and to a side.

TRANSVERSALIS COLLI.

Origin: From the Transverse Processes of the five uppermost Dorsal Vertebrae, by the same number of Tendinous and Fleishy Slips. It runs between the Trachelo-Mastoideus, Splenius Colli, and Cervicalis Descendens.

Insertion: Into the Transverse Processes of all the Cervical Vertebrae, except the first and last. Tab. XLII. Fig. 1, Neck, I, I.

Action: To turn the Neck obliquely backwards, and a little to one side.

RECTUS CAPITIS POSTICUS MINOR,

Vel Rectus Minor.

Origin: Tendinous, close to its fellow, from a small Protuberance which is instead of the Spinous Process of the first Cervical Vertebra; spreading out in its ascent.

Insertion: Fleishy, into a Depression between the smaller Arch and Foramen Magnum of the Occipital Bone. Tab. XLIII. Fig. 1. A.

Action: To assist the following Muscle in drawing the Head backwards.

RECTUS CAPITIS POSTICUS MAJOR,

Vel Rectus Major.

Origin: Fleishy, from the external part of the Spinous Process of the second Cervical Vertebra. It becomes gradually broader, and goes obliquely upwards and outwards.

Insertion: Tendinous and Fleishy, into the Os Occipitis, at the outside of the Insertion of the Rectus Minor, part of which it covers. Tab. XLIII. Fig. 1. B.

Action: To pull the Head backwards, and to assist a little in its rotation.

OBLIQUUS CAPITIS INFERIOR.

Origin: Fleishy, from the Spinous Process of the second Cervical Vertebra, at the outside of the Rectus Major. It forms a thick Belly, which runs upwards and outwards.

Insertion: Into the Transverse Process of the first Cervical Vertebra. Tab. XLIII. Fig. 1. D.

Action: To roll the Head.

OBLIQUUS CAPITIS SUPERIOR.

Origin: From the Transverse Process of the first Cervical Vertebra, passing upwards and a little inwards.

Insertion: Into the Occipital Bone, at the outer part of the Insertion of the Rectus Major. Tab. XLIII. Fig. 1. C.

Action: To assist in drawing the Head backwards, and a little to one side.

SCALENUS ANTIQUS.

Origin: Tendinous and Fleishy, from the upper part of the first Rib, near its Cartilage.

Insertion: Into the Transverse Processes of the fourth, fifth, and sixth Cervical Vertebrae, by as many Tendons. Tab. XXXVI. Fig. 1. L.

SCALENUS MEDIUS.

Origin: From the upper and outer part of the first Rib, from its Root to near its Cartilage.

Insertion: Into the Transverse Processes of all the Cervical Vertebrae, by as many strong Tendons. Tab. XXXVII. Neck, M, N.

The Subclavian Artery, and the Nerves which form the Brachial Plexus, pass between this and the former Muscle.

SCALENUS POSTICUS.

Origin: From the upper edge of the second Rib, near the Spine.

Insertion: Into the Transverse Processes of the fifth and sixth Cervical Vertebrae. Tab. XLII. Neck, E.

Action of the three Scaleri: To bend the Neck to one side; or, when the Neck is fixed, to raise the Ribs, and dilate the Thorax, as in violent Inspiration.

INTERSPINALES COLLI.

The spaces between the Spinous Processes of the Cervical Vertebrae, most of which are forked, are occupied by double Fleishy Portions.

Origin: From each Inferior Spinous Process.

Insertion: Into each Superior. Tab. XLIII. Neck, G, G.

Action: To draw these Processes nearer to each other, and of course the Neck a little backwards.

INTERTRANSVERSALES COLLI.

The spaces between all the Transverse Processes of the Cervical Vertebrae, which are also forked, are filled up in like manner with double Fleishy Portions.

Action: To draw these Processes towards each other, and

and turn the Neck a little to one side. Tab. XXXVII.
Neck, *f, f.*

INTERSPINALES LUMBORUM.

They are of the same nature with the Interspinales and Intertransversales Dorfi.

INTERSPINALES ET INTERTRANSVERSALES DORSI.

These are rather small Tendons than Muscles, serving to connect the Spinal and Transverse Processes. Tab. XLIII. Trunk, *c, c, d, d.*

INTERTRANSVERSALES LUMBORUM.

These are five distinct Muscles which occupy the Spaces between the Transverse Processes of the last Dorsal and of all the Lumbar Vertebrae, and serve to draw them a little towards each other. Tab. XLIII. Loins, H, H.

MUSCLES OF THE SUPERIOR EXTREMITY.

MUSCLES ARISING FROM THE SCAPULA.

SUPRA-SPINATUS.

Origin: Flethy, from the whole Fossa Supra-Spinata, and from the Spine and Superior Costa of the Scapula; passing under the Acromion, and adhering to the Capsular Ligament of the Joint.

Insertion: Tendinous, into the large Tubercle on the Head of the Os Humeri. Tab. XLI. Shoulder, B.

Action: To raise the Arm, and at the same time to pull the Capsular Ligament from between the Bones, so as to prevent it from being pinched.

INFRA-SPINATUS.

Origin: Flethy, from all that part of the Dorsum Scapulæ which is below its Spine; and from the Spine itself, as far as the Cervix Scapulæ. The Fibres run obliquely towards a Tendon in the middle of the Muscle, which runs forwards, and adheres to the Capsular Ligament.

Insertion: By a flat thick Tendon, into the upper and outer part of the large Protuberance on the Head of the Os Humeri. Tab. XLI. Shoulder, C.

Action: To roll the Os Humeri outwards; to assist in raising, and in supporting it when raised; and to pull the Ligament from between the Bones.

These two Muscles are covered by an Aponeurosis, which extends between the Costæ and edges of the Spine of the Scapula, and gives rise to many of the Muscular Fibres.

TERES MINOR.

Origin: Flethy, from the inferior Costa of the Scapula.

It ascends along the under edge of the Infra-Spinatus, and adheres to the Capsular Ligament.

Insertion: Tendinous, into the back part of the large Protuberance on the Head of the Os Humeri, a little below the Infra-Spinatus. Tab. XLI. Shoulder, D.

Action: To roll the Os Humeri outwards, to draw it backwards, and to prevent the Ligament from being pinched between the Bones.

TERES MAJOR.

Origin: Flethy, from the Dorsal side of the inferior Angle of the Scapula, and from a small part of its inferior Costa. It is situated at the under part of the Teres Minor, and sends off a broad flat Tendon, which accompanies that of the Latissimus Dorfi.

Insertion: Along with the Latissimus Dorfi, into the Ridge at the inner side of the Groove for lodging the Tendon of the long Head of the Biceps. Tab. XLI. Shoulder, E.

Action: To roll the Humerus inwards, and to draw it backwards and downwards.

DELTOIDES.

Origin: Flethy, from all the outer part of the Clavicle unoccupied by the Pectoralis Major, from which it is separated by a small Fissure; Tendinous and Flethy, from the Acromion, and lower Margin of almost the whole Spine of the Scapula, opposite to the insertion of the Trapezius.

From these Origins it runs, under the appearance of three Muscles going in different directions, and separated from each other by slight Fissures; viz. from the Clavicle outwards,

outwards, from the Acromion downwards, and from the Spine of the Scapula forwards; and is composed of a number of Fasciculi, forming a strong Flešhy Muscle, which covers the Joint of the Os Humeri.

Insertion: Below that of the Pectoralis Major, by a short and strong Tendon, into a rough Surface, on the outer side of the Os Humeri, near its middle, where the Fibres of this Muscle intermix with part of the Brachialis Externus. Tab. XL. Shoulder, A.

Action: To pull the Arm directly outwards and upwards, and a little forwards or backwards, according to the different directions of its Fibres.

CORACO-BRACHIALIS.

Origin: Tendinous and Flešhy, from the fore-part of the Coracoid Process of the Scapula, in common with the short Head of the Biceps, to which it adheres through the greater part of its length.

Insertion: Tendinous and Flešhy, into the internal part

of the Os Humeri, near its middle, where it sends down an Aponeurosis to the internal Condyle of that Bone. Tab. XXXVI. Arm, C.

Action: To bring the Arm obliquely upwards and forwards.

SUBSCAPULARIS.

Origin: Flešhy, from the three Costæ, and whole inner Surface of the Scapula. It is composed of a number of Tendinous and Flešhy portions, which run in a radiated manner, and make prints on the Bone; in its passage outwards, adhering to the Capsular Ligament. Tab. XXXVII. Shoulder, A.

Insertion: Tendinous, into the upper part of the Internal Protuberance at the Head of the Os Humeri.

Action: To roll the Arm inwards, draw it to the side of the Body, and to prevent the Capsular Ligament from being pinched.

MUSCLES CHIEFLY SITUATED ON THE ARM, SERVING FOR THE MOTION OF THE FORE-ARM.

APONEUROSIS,

Of the Superior Extremity. Tab. XXXIII.

Fig. 1. 2.

The greater part of the Superior Extremity is covered by a Tendinous Membrane or Aponeurosis, which arises from the Bones of, and muscles on, the Shoulder.

On the Humerus, it incloses the Flexor and Extensor Muscles of the Fore-Arm, and is connected to the Ridges and Condyles at the under end of the Os Humeri.

At the bending of the Elbow, it receives considerable additions from the Tendons of the Biceps and Triceps of the Fore-Arm, where the Fibres from the opposite sides decussate each other.

It becomes thicker and stronger on the Fore-Arm, and forms a firm covering to the Muscles there.

In its descent, it gives off Partitions among the Muscles, and these are fixed to the Radius and Ulna; the Membrane itself being lost insensibly upon the Hand.

It is thicker and stronger on the outer than upon the inner side of the Extremity, particularly on the Fore-Arm, at the under and back part of which it forms a thick and strong Band, which, running transversely, gets the name of *Ligamentum Carpi Annulare Posterius*.

The Use of this Aponeurosis, like that in other parts of the Body, is to brace the Muscles, by keeping them in their proper place while in action, and to give Origin to many of the Muscular Fibres which lie immediately under it.

BICEPS FLEXOR CUBITI, vel *Biceps*.

Origin: By two Heads; the outer one, called its

Long Head, begins by a slender Tendon from the upper edge of the Glenoid Cavity of the Scapula, passes over the Ball of the Os Humeri within the Joint, and, in its descent without the Joint, is inclosed in a Groove upon the upper and fore part of the Bone, by a Ligament which proceeds from the Capsular Ligament and adjacent Tendons: The inner one, called its *Short Head*, arises, Tendinous and Flešhy, from the Coracoid Process of the Scapula, in common with the Coraco-Brachialis. A little below the middle of the fore-part of the Os Humeri, the two Heads unite, and form a thick Flešhy Belly.

Insertion: By a strong roundish Tendon, into the Tubercle at the upper and inner part of the Radius, and by a Tendinous Expansion into the Aponeurosis of the Fore-Arm, which it likewise assists in forming. Tab. XXXV. Arm, A.

Action: To bend the Fore-Arm, and to assist the Supinator Muscles in rolling the Radius outwards, and, of course, to turn the Palm of the Hand upwards.

BRACHIALIS INTERNUS.

Origin: Flešhy, from the middle of the Os Humeri, at each side of the Deltoides, covering all, and attached to most, of the under and fore part of the Bone. It runs over the Joint, adhering firmly to the Capsular Ligament.

Insertion: By a strong short Tendon, into the Coronoid Process of the Ulna. Tab. XXXVI. Arm, D.

Action: To bend the Fore-Arm, and to prevent the Ligament of the Joint from being pinched.

TRICEPS

TRICEPS EXTENSOR CUBITI.

Origin: By three Heads: The first, or *long* Head, broad and Tendinous, from the inferior Costa of the Scapula, near its Cervix: The second, or *short* Head, by an acute, Tendinous, and Fleshy beginning, from the outer and back part of the Os Humeri, a little below its upper extremity: The third, called *Brachialis Externus*, arises, by an acute beginning, from the back part of the Os Humeri, near the Insertion of the Teres Major. The three Heads unite about the middle of the Humerus, and cover the whole posterior part of that Bone, adhering to it in their descent.

Insertion: Into the upper and outer part of the Olecranon of the Ulna, and partly into the Condyles of the

Os Humeri, adhering closely to the Ligament. Tab. XLI. Arm, F, g, b, i.

Action: To extend the Fore-Arm.

ANCONEUS.

Origin: Tendinous, from the posterior part of the external Condyle of the Os Humeri. It descends under a triangular form, soon becomes Fleshy, and part of its Flesh is likewise continued from the third Head of the Triceps.

Insertion: Fleshy and thin, into a Ridge on the outer and back edge of the Ulna, a little below the Olecranon. Tab. XLI. Fore-Arm, I.

Action: To assist the Triceps in extending the Fore-Arm.

MUSCLES ON THE FORE-ARM AND HAND, SERVING FOR THE MOTION OF THE HAND AND FINGERS.

To prevent confusion in the application of the terms *Outer* and *Inner*, when the Muscles are described in the prone state of the Hand,—the Arm is here supposed to hang by the side of the Body, with the Palm turned forwards, so that the Radius and Thumb are upon the outer, and the Ulna and Little Finger upon the inner side.

PALMARIS LONGUS.

Origin: Tendinous from the internal Condyle of the Os Humeri; soon becoming Fleshy, and sending off a long slender Tendon.

Insertion: Into the Ligamentum Carpi Annulare Anterior, and into the Aponeurosis Palmaris. Tab. XXXVIII. Fore-Arm, B. γ.

Action: To bend the Hand, and stretch the Aponeurosis Palmaris.

This Muscle is frequently a-wanting; but the Aponeurosis is always to be found.

APONEUROSIS PALMARIS.

This Tendinous Expansion begins at the Ligamentum Carpi Annulare Anterior; and, after spreading out and covering the greater part of the Palm of the Hand, is fixed to the Roots of all the Fingers by an equal number of double Slips. Tab. L. Fig. 1.

PALMARIS BREVIS.

Origin: By small bundles of Fleshy Fibres, from the Ligamentum Carpi Annulare, and Aponeurosis Palmaris.

Insertion: Into the Skin and Fat which cover the Abductor Minimi Digiti, and into the Os Pisiforme. Tab. L. Fig. 1. o.

VOL. I.

Action: To assist in contracting the Palm of the Hand.

FLEXOR CARPI RADIALIS, vel *Radialis Internus*.

Origin: Tendinous and Fleshy, from the inner Condyle of the Os Humeri, and from the fore and upper part of the Ulna, between the Pronator Radii Teres and Flexor Sublimis, to which it firmly adheres. It forms a long Tendon, which passes down near the Radius, goes through a Fossa in the Os Trapezium, and becomes flat at its inferior extremity.

Insertion: Into the fore and upper part of the Metacarpal Bone which sustains the Fore-Finger. Tab. XXXVIII. Arm, α, α.

Action: To bend the Wrist, and to assist in the pronation of the Hand.

FLEXOR CARPI ULNARIS, vel *Ulnaris Internus*.

Origin: Tendinous, from the internal Condyle of the Os Humeri; and, by a small Fleshy beginning, from the corresponding side of the Olecranon. It passes along the inner side of the Ulna, from which also it derives part of its origin for a considerable way down. A number of its Fleshy Fibres likewise arise from the Aponeurosis of the Fore-Arm.

Insertion: By a strong Tendon, into the Os Pisiforme. Tab. XXXVIII. Right Arm, ε.

Action: To assist the former Muscle in bending the Wrist.

EXTENSOR CARPI RADIALIS LONGIOR, vel *Radialis Externus Longior*.

Origin: Broad, thin, and Fleshy, directly below the Supinator Longus, from the lower part of the Ridge of the Os Humeri, above its external Condyle. It sends off

a long flat Tendon, which passes down, first upon the outer, and then upon the back part of the Radius, descending in a Groove there, and going under the Ligamentum Carpi Annulare.

Insertion: Into the upper, back, and outer part of the Metacarpal Bone of the Fore-Finger. Tab. XLV. Right Arm, F.

Action: To extend the Wrist, and bring the Hand backwards.

EXTENSOR CARPI RADIALIS BREVIOR, vel Radialis Externus Brevior.

Origin: Tendinous, in common with the Extensor Longior, but farther down, from the external Condyle of the Os Humeri, and from the Ligament which connects the Radius to it.—Passing down upon the back part of the Radius, its Tendon goes under the Ligamentum Annulare, in the same Channel with the Tendon of the Extensor Longior.

Insertion: Into the upper and back part of the Metacarpal Bone of the Middle Finger. Tab. XLV. Right Arm, G.

Action: To assist the former Muscle in extending the Wrist; or, with it and the Flexor Carpi Radialis, to draw the Hand to the side next the Thumb.

EXTENSOR CARPI ULNARIS, vel Ulnaris Externus.

Origin: Tendinous, from the external Condyle of the Os Humeri; and, in its progress, Fleishy, from the middle of the Ulna where it passes over that bone. Its round Tendon is inclosed by a Membranous Sheath, in a Groove at the back part of the extremity of the Ulna.

Insertion: Into the posterior and upper part of the Metacarpal Bone of the Little Finger. Tab. XLIV. Right Arm, Q.

Action: To assist the two former Muscles in extending the Wrist; or, with the assistance of the Flexor Ulnaris, to draw the Hand towards the side next the Little Finger.

FLEXOR DIGITORUM SUBLIMIS, vel PERFORATUS.

Origin: Tendinous and Fleishy, from the internal Condyle of the Os Humeri; Tendinous, from the root of the Coronoid Process of the Ulna; and Membranous and Fleishy, from the middle of the fore-part of the Radius. Its Fleishy Belly sends off four round Tendons before it passes under the Ligamentum Carpi Annulare. In their course, they are connected to those of the following Muscle by fine Membranous Webs, and upon the Fingers are inclosed in strong Tendinous Sheaths.

Insertion: Into the anterior and upper part of the second Phalanx of the Fingers, being near the under part of the first Phalanx, split and twisted to form a Passage,

and at the same time a kind of Sheath for the Tendons of the Flexor Profundus. Tab. XXXVIII. Left Arm, z, A.

Action: To bend the second, and then the first Phalanx of the Fingers.

FLEXOR DIGITORUM PROFUNDUS, vel PERFORANS.

Origin: Fleishy, from the external side and upper part of the Ulna, for some way down; and from a large share of the Interosseous Ligament. It descends behind the Flexor Sublimis, and, like it, splits into four Tendons, a little before it passes under the Ligamentum Annulare, and these pass through the Slits in the Tendons of the Flexor Sublimis. Tab. XXXIX. Left Arm, V, W, X.

Insertion: Into the anterior and upper part of the third Phalanx of the Fingers.

Action: To bend the last Joint of the Fingers.

LUMBRICALES.

Origin: Thin and Fleishy, from the outside of the Tendons of the Flexor Profundus, a little above the lower edge of the Ligamentum Carpi Annulare. At the under ends of the Metacarpal Bones, they send off long slender Tendons.

Insertion: Into the outer sides of the broad Tendons of the Interossei Muscles, about the middle of the first Phalanx. Tab. L. Fig. 3. m, &c.

Action: To bend the first Phalanx, and increase the Flexion of the Fingers, while the long Flexors are in full action.

EXTENSOR DIGITORUM COMMUNIS.

Origin: Tendinous and Fleishy, from the external Condyle of the Os Humeri, where it adheres to the Supinator Radii Brevis. It passes down upon the back part of the Fore-Arm, and before it goes under the Ligamentum Carpi Annulare Posterius, it splits into three or four Tendons, some of which may be divided into smaller ones. Upon the back of the Metacarpal Bones, the Tendons become broad and flat, and near the Heads of these Bones send Aponeurotic Expansions to each other.

Insertion: Into the posterior part of all the Bones of the four Fingers, by a Tendinous Expansion. Tab. XLV. Right Arm, S, T.

Action: To extend all the Joints of the Fingers.

SUPINATOR RADII LONGUS.

Origin: By an acute Fleishy beginning, from the Ridge of the Os Humeri, above the external Condyle, nearly as high as the middle of the Bone. It forms a thick Fleishy

Fleshy Belly, which covers the upper part of the Extensor Carpi Radialis Longior; and about the middle of the Fore-Arm sends a tapering Tendon along the edge of the Radius.

Insertion: Into the outer side of the under end of the Radius. Tab. XXXVIII. Fore-Arm, W.

Action: To roll the Radius outwards, and, of course, to turn the Hand into a supine situation, or with the Palm forwards.

SUPINATOR RADII BREVIS.

Origin: Tendinous from the external Condyle of the Os Humeri, and Tendinous and Fleshy from the outer and upper part of the Ulna, and from the Interosseous Ligament. It passes over the external edge of the Radius.

Insertion: Into the upper and fore part of the Radius. Tab. XLIV. Fore-Arm, O.

Action: To assist the Supinator Longus.

PRONATOR RADII TERES.

Origin: Fleshy from the internal Condyle of the Os Humeri, and Tendinous from the Coronoid Process of the Ulna. It goes obliquely across the upper end of the Flexor Muscles of the Wrist, and is of a tapering form.

Insertion: Thin, Tendinous, and Fleshy, into the middle of the posterior part of the Radius. Tab. XXXIV. Fore-Arm, G.

Action: To roll the Radius inwards, by which it brings the Palm of the Hand backwards, or into a state of Pronation.

PRONATOR RADII QUADRATUS.

Origin: Broad, Tendinous, and Fleshy, from the under and inner part of the Ulna. The Fibres run transversely.

Insertion: Into the under and fore part of the Radius. Tab. L. Fig. 4. a.

Action: To assist the Pronator Teres.

FLEXOR LONGUS POLLICIS MANUS, Vel Flexor Tertii Internodii.

Origin: By an acute Fleshy beginning, from the fore-part of the Radius and Interosseous Ligament, the Origin extending from the Tubercle of the Bone, as far as the Pronator Quadratus. It has frequently another Origin, by a distinct Fleshy Slip, from the internal Condyle of the Os Humeri.

Insertion: Into the last Joint of the Thumb, after its Tendon has passed under the Ligamentum Carpi Annulare Anterius. Tab. XXXIX. Left Fore-Arm, α , β .

Action: To bend the last Joint of the Thumb.

FLEXOR BREVIS POLLICIS, Vel Flexor Secundi Internodii.

Origin: From the Offa Trapezoides, Magnum, et Unciforme. It is divided into two Portions, which form a Groove for the Tendon of the Flexor Longus Pollicis.

Insertion: Into the Offa Sefamoidea, and Base of the first Bone of the Thumb. Tab. L. Fig. 2. m , n .

Action: To bend the first Joint of the Thumb.

OPPONENS POLLICIS,

Vel Flexor Ossis Metacarpi Pollicis, vel Flexor Primi Internodii.

Origin: Fleshy, from the Os Trapezium and Ligamentum Carpi Annulare Anterius. It lies immediately under the Abductor Pollicis.

Insertion: Tendinous and Fleshy, into the under and fore part of the Metacarpal Bone of the Thumb. Tab. L. Fig. 2. l .

Action: To bring the Thumb inwards, so as to make it oppose the Fingers; from which circumstance it has derived its name.

EXTENSOR OSSIS METACARPI POLLICIS.

Origin: Fleshy, from the middle of the posterior parts of the Ulna, Radius, and Interosseous Ligament. It runs obliquely over the Radius, sending one, or more frequently two Tendons, through an Annular Sheath.

Insertion: Into the Os Trapezium, and upper and back part of the Metacarpal Bone of the Thumb. Tab. XLIV. Right Fore-Arm, i .

Action: To extend the Metacarpal Bone of the Thumb, and draw it from the Fingers.

EXTENSOR PRIMI INTERNODII POLLICIS, Vel Extensor Minor.

Origin: Fleshy, from the back part of the Ulna, and from the Interosseous Ligament, near the former Muscle, by the side of which it runs.

Insertion: Tendinous, into the posterior part of the first Bone of the Thumb: A portion of it may be traced as far as the second Bone. Tab. XLIV. Right Fore-Arm, b .

Action: To extend the first Joint of the Thumb.

EXTENSOR SECUNDI INTERNODII, Vel Extensor Major.

Origin: By an acute, Tendinous, and Fleshy beginning, from the middle of the back part of the Ulna, and from the Interosseous Ligament. Its Tendon runs through a small Groove at the under, inner, and back part of the Radius.

Insertion:

Insertion : Into the last Bone of the Thumb. Tab. XLIV. Left Fore-Arm, R.

Action : To extend the last Joint of the Thumb.

ABDUCTOR POLLICIS.

Origin : Broad, Tendinous, and Fleshy, from the Ligamentum Carpi Annulare, and from the Os Trapezium. It lies immediately under the Skin, and over the Opponens Pollicis.

Insertion : Tendinous, into the outer side of the root of the first Bone of the Thumb. Tab. L. Fig. 1. i.

Action : To draw the Thumb from the Fingers.

A particular portion on the inner side of this Muscle is called, by ALBINUS, *Abductor Brevis Alter*.

ADDUCTOR POLLICIS.

Origin : Fleshy, from almost the whole length of the Metacarpal Bone of the Middle Finger; going across the Metacarpal Bone of the Fore-Finger, its Fibres converge, and send off a short Tendon.

Insertion : Into the inner part of the root of the first Bone of the Thumb. Tab. XXXIX. Fig. 1. Right Arm, g.

Action : To pull the Thumb towards the Fingers.

INDICATOR,

Vel Extensor Indicis Proprius.

Origin : By an acute Fleshy beginning, from the middle of the posterior part of the Ulna, at the inner side of the Extensor Secundi Internodii Pollicis. Its Tendon passes under the same Ligament with the Extensor Digitorum Communis.

Insertion : Along with part of the Extensor Digitorum Communis, into the posterior part of the Fore-Finger. Tab. XLIV. Left Fore-Arm, F.

Action : To assist the Extensor Communis in extending all the Joints of this Finger, as in pointing at any thing; hence called *Indicator*.

ABDUCTOR INDICIS.

Origin : From the Os Trapezium, and from the upper part and inner side of the Metacarpal Bone of the Thumb.

Insertion : By a short Tendon, into the outer and back part of the first Bone of the Fore-Finger. Tab. L. Fig. 6. i.

Action : To bring the Fore-Finger towards the Thumb.

ABDUCTOR MINIMI DIGITI.

Origin : Fleshy, from the Os Pisiforme, and from

that part of the Ligamentum Carpi Annulare Anterior next it.

Insertion : Tendinous, into the inner side of the Base of the first Bone of the Little Finger. Tab. L. Fig. 2. v.

Action : To draw the Little Finger from the rest.

FLEXOR PARVUS MINIMI DIGITI.

Origin : From the Ulnus of the Os Unciforme, and adjacent part of the Annular Ligament. It passes obliquely over the under end of the former Muscle.

Insertion : By a roundish Tendon, into the inner part of the Base of the first Bone of this Finger. Tab. L. Fig. 3. b.

Action : To bend the Little Finger, and assist the Adductor.

ADDUCTOR MINIMI DIGITI, vel Metacarpeus.

Origin : Fleshy, from the edge of the hook-like Process of the Os Unciforme, and from that part of the Ligamentum Carpi Annulare next it.

Insertion : Tendinous, into the inner side, and anterior or under extremity of the Metacarpal Bone of the Little Finger. Tab. L. Fig. 2. u.

Action : To bend the Metacarpal Bone, and bring this Finger towards the rest.

INTEROSSEI.

Origin : From the sides of the Metacarpal Bones. They fill up the spaces between these, and are something similar to the Lumbricales, but larger.

Insertion : By slender Tendons, along with those of the Lumbricales, into the sides of the Tendinous Expansions of the Extensor Digitorum Communis. Tab. L. Fig. 4. e—l.

Action : To give the Fingers their lateral motions, and to assist a little, according to their situations, in bending or extending the first Phalanx of the Fingers.

Of the Interossei, three, seen in the Palm of the Hand, arise with single Heads, and are called *Interni*; and four on the back of the Hand, with double Heads, termed *Externi*, or *Bicipites*. Part of the Externi, however, are also seen in the Palm of the Hand.

INTEROSSEI INTERNI.

PRIOR INDICIS.

Origin : From the outer or Radial side of the Metacarpal Bone of the Fore-Finger.

Insertion : Into the outside of the Tendon on the back of the Fore-Finger.

Action : To draw that Finger outwards, towards the Thumb.

POSTERIOR

POSTERIOR INDICIS.

Origin : From the inner or Ulnar side of the Metacarpal Bone of the Fore-Finger.

Insertion : Into the inside of the Tendon on the back of the Fore-Finger.

Action : To draw the Fore-Finger inwards.

PRIOR ANNULARIS.

Origin : From the outside of the Metacarpal Bone of the Ring-Finger.

Insertion : Into the outside of the Tendon, on the back of the Ring-Finger.

Action : To draw the Ring-Finger outwards.

INTEROSSEUS AURICULARIS.

Origin : From the outside of the Metacarpal Bone of the Little Finger.

Insertion : Into the outside of the Tendon on the back of the Little Finger.

Action : To draw the Little Finger outwards.

INTEROSSEI EXTERNI.

PRIOR MEDII DIGITI.

Origin : From the corresponding sides of the Metacarpal Bones of the Fore and Middle Fingers.

Insertion : Into the outside of the Tendon on the back of the Middle Finger.

Action : To draw the Middle Finger outwards.

POSTERIOR MEDII DIGITI.

Origin : From the corresponding sides of the Metacarpal Bones of the Middle and Ring Fingers.

Insertion : Into the inside of the Tendon on the back of the Ring Finger.

Action : To draw the Middle Finger inwards.

POSTERIOR ANNULARIS.

Origin : From the corresponding sides of the Metacarpal Bones of the Ring and Little Fingers.

Insertion : Into the inside of the Tendon on the back of the Ring Finger.

Action : To draw the Ring-Finger inwards.

MUSCLES OF THE INFERIOR EXTREMITY.

MUSCLES ON THE PELVIS AND THIGH, SERVING FOR THE MOTION OF THE THIGH AND LEG.

APONEUROSIS OF THE INFERIOR EXTREMITY.

Previous to the description of the Muscles of the Inferior Extremity, it is proper to take notice of a Tendinous Expansion, which, as in the Superior Extremity, forms a general Covering to the Muscles, and sends off Partitions between them, to be connected to the Ridges and Processes of the Bones.

It is thick and strong on the outside of the Thigh and Leg, but towards the inner side of both, particularly on the former, it gradually turns thinner, and has rather the appearance of Cellular Membrane.

It descends from the Processes and other Projections on the outside of the Bones of the Pelvis, especially from the Tendons of the external Layers of Muscles of the Loins and Abdomen.

A little below the Trochanter Major, it is intimately connected to the Linea Aspera. At the Joint of the Knee it receives additions from the Tendons of the Extensors of the Leg, and is there connected with the outer and inner sides of the Head of the Tibia and Fibula. In the Leg, it is firmly fixed to the Spine of the Tibia; and at the under end, to the Bones of the Ankle, where part of it, thicker and stronger than the rest, forms the Ligamentum Annulare. It is lost at last upon the Foot.

It serves the same general purposes with the Aponeurosis of the Superior Extremity.

PSOAS MAGNUS. } See p. 113.
ILIACUS INTERNUS. }

PECTINALIS, vel *Pectineus*.

Origin : Broad and Flethy, from the upper and fore part of the Os Pectinis vel Pubis, immediately above the Foramen Thyroideum. It runs downwards and outwards at the inner side of the Psoas Magnus.

Insertion : By a flat and short Tendon, into the Linea Aspera of the Os Femoris, a little below the Trochanter Minor. Tab. XXXIV. Thigh, E.

Action : To pull the Thigh upwards and inwards, and to give it, and, of course, the Foot, a degree of rotation outwards.

TRICEPS ADDUCTOR FEMORIS.

Under this appellation are comprehended three distinct Muscles, viz. *Adductor Longus*, *Adductor Brevis*, and *Adductor Magnus*.

ADDUCTOR

ADDUCTOR LONGUS.

Origin: By a strong roundish Tendon, from the upper and fore part of the Os Pubis, and Ligament of the Synchondrosis, at the inner side of the Pectinalis: It runs downwards and outwards.

Insertion: By a broad flat Tendon, into the middle of the Linea Aspera. Tab. XXXVIII. Thigh, c.

ADDUCTOR BREVIS.

Origin: Tendinous, from the Os Pubis, at the side of its Symphysis, below and behind the former Muscle. It runs obliquely outwards.

Insertion: By a short flat Tendon, into the inner and upper part of the Linea Aspera, from a little below the Trochanter Minor, to the beginning of the Insertion of the Adductor Longus. Tab. XXXIX. Left Thigh, e.

ADDUCTOR MAGNUS.

Origin: From the side of the Symphysis Pubis, a little lower than the former. The Origin is continued downwards from the Crus and Tuberosity of the Os Ischium. The Fibres run outwards and downwards, spreading out wide, and forming a very large Muscle.

Insertion: Into the whole length of the Linea Aspera; the under part of the Muscle extending along the Ridge which leads to the inner Condyle of the Os Femoris. It is also fixed by a roundish Tendon, into the upper part of that Condyle, a little above which the Femoral Artery, taking a Spiral turn towards the Ham, passes between the Tendon of this Muscle and the Bone. Tab. XXXVII. Fig. 1. Thigh, D. Fig. 2. A, B, C.

Action of the three Adductors: To bring the Thigh inwards and upwards, according to the different directions of their Fibres, and to assist a little in rolling it outwards.

OBTURATOR EXTERNUS.

Origin: By a semicircular Margin, from the parts of the Ossa Pubis and Ischium, which form the anterior half of the Foramen Thyroideum, and from the Membrane which fills up that Foramen. The Fibres are collected like rays towards a centre, and pass outwards over the back part of the Cervix of the Os Femoris.

Insertion: By a strong round Tendon, into the Cavity at the inner and back part of the root of the Trochanter Major, adhering in its course to the Capsular Ligament of the Thigh-Bone. Tab. XXXVII. Fig. 1. Thigh, C.

Action: To roll the Thigh-Bone obliquely outwards, and to prevent the Capsular Ligament from being pinched.

GLUTEUS MAXIMUS.

Origin: Fleishy, from the back part of the Spine of

the Os Ilium; from the under and outer part of the Os Sacrum; from the Os Coccygis; and from the posterior Sacro-Sciatic Ligaments over which part of the inferior edge hangs in a Flap. The Fibres run obliquely forwards, and a little downwards, to form a thick broad Muscle, which is composed of distinct coarse Fasciculi. The upper part of it covers almost the whole of the Trochanter Major, and it is intimately connected with the broad Tendon of the *Tensor Vaginae Femoris*.

Insertion: By a strong, thick, and broad Tendon, into the upper and outer part of the Linea Aspera, along which it is continued for some way down. Tab. XL. Pelvis and Thigh, A, A.

Action: To extend the Thigh, and pull it backwards and a little outwards.

GLUTEUS MEDIUS.

Origin: Fleishy, from all that part of the Spine of the Os Ilium which is unoccupied by the Gluteus Maximus; from the upper part of the Dorsum of that Bone; and from an Aponeurosis which covers the Muscle, and joins the Fascia of the Thigh. It sends off a broad Tendon.

Insertion: Into the outer and back part of the Trochanter Major. Tab. XLI. Pelvis, A.

Action: To pull the Thigh outwards, and a little backwards. The fore-part of the Muscle assists in rolling it inwards.

GLUTEUS MINIMUS.

Origin: Fleishy, from the lower half of the Dorsum of the Os Ilium. Its Origin is continued from the superior-anterior Spinous Process, along a rising of the Bone, as far as the great Sciatic Notch; and the Muscle runs in a radiated manner to a strong flat Tendon.

Insertion: Into the fore and upper part of the Trochanter Major. Tab. XLII. Pelvis, A.

Action: To assist the former in pulling the Thigh outwards, and a little backwards; and, along with other Muscles, in rolling it inwards.

PYRIFORMIS.

Origin: Within the Pelvis, by three Tendinous and Fleishy Heads, from the second, third, and fourth pieces of the Os Sacrum; and, becoming round and tapering, it passes out of the Pelvis, along with the Sciatic Nerve, through the great Notch of the Ilium, from which it receives the addition of a few Fleishy Fibres.

Insertion: By a roundish Tendon, into the upper part of the Cavity, at the inner side of the root of the Trochanter Major. Tab. XLI. Pelvis, B.

Action: To assist in the Abduction of the Thigh, and in its rotation outwards,

GEMINI,

GEMINI, vel *Gemelli*.

Origin: By two distinct Heads; the superior from the Spinous Process, and the inferior from the Tuberosity of the Os Ischium, and from the Sacro-Sciatic Ligament. The two Heads are united by a Tendinous and Flethy Membrane, and form a Sheath for the reception of the Tendon of the Obturator Internus.

Insertion: Tendinous and Flethy, into the Cavity at the inner side of the root of the Trochanter Major, on each side of the Tendon of the Obturator Internus, to which they firmly adhere. Tab. XLI. Pelvis, C.

Action: To roll the Thigh outwards, and to prevent the Tendon of the Obturator Internus from starting out of its place while the Muscle is in action.

OBTURATOR INTERNUS, olim *Marsupialis*.

Origin: Within the Pelvis, by a semicircular Flethy margin, from the anterior half of the Foramen Thyroideum, and, in part, from the Obturator Ligament. Its Fibres converge, and send off a round Tendon which passes over the Os Ischium, between the Spine and Tubercle of that Bone, as a rope passes over a pulley.—Where it goes over the Capsular Ligament of the Thigh-Bone, it is inclosed in the Sheath of the Gemini.

Insertion: By a round Tendon, along with the Gemini, into the large Pit at the root of the Trochanter Major. Tab. XLII. Fig. 1. Pelvis and Inferior Extremity, B.

Action: To roll the Thigh obliquely outwards.

QUADRATUS FEMORIS.

Origin: Tendinous and Flethy, from the outer side of the Tuberosity of the Os Ischium; running transversely outwards.

Insertion: Flethy, into a rough Ridge continued from the root of the great, to that of the small Trochanter. Tab. XLI. Pelvis, E.

Action: To roll the Thigh outwards.

The Piriformis, Gemini, Quadratus, and Obturatores, which are the Rotators of the Thigh, when it is in a line with the Body, become its Abductors when it is in the bended state.

TENSOR VAGINÆ FEMORIS.

Origin: By a narrow, Tendinous, and Flethy beginning, from the external part of the anterior-superior Spinous Process of the Os Ilium. It goes downwards, and a little backwards, forming a thick Flethy Belly, which is inclosed in a doubling of the Aponeurosis or Vagina of the Thigh.

Insertion: A little below the Trochanter Major, into the inner Surface of the Aponeurosis which covers the outside of the Thigh. Tab. XXXIV. Thigh, A.

Action: To stretch the Aponeurosis, and to assist in the Abduction of the Thigh, and in its rotation inwards.

SARTORIUS.

Origin: Tendinous, from the superior-anterior Spinous Process of the Os Ilium. It soon becomes Flethy, runs obliquely downwards over the Muscles situated upon the fore and inner side of the Thigh, and is the longest Muscle of the Body.

Insertion: By a broad and thin Tendon, into the inner side of the Tibia, near the inferior part of its Tubercle. Tab. XXXVIII. Right Inferior Extremity, g, q.

Action: To bend the Knee, and bring one Leg obliquely inwards across the other.

GRACILIS, vel *Rectus Internus*.

Origin: By a thin Tendon, from the Os Pubis, near the Symphysis; soon becoming Flethy, and descending in a direct course by the inside of the Thigh.

Insertion: Tendinous, into the Tibia, under the Sartorius. Tab. XXXVI. Fig. 1. Thigh, G. Fig. 2. D.

Action: To assist the Sartorius in making the full Flexion of the Knee, after it has been bent to a certain degree by the Flexors on the back part of the Thigh.

RECTUS, vel *Gracilis Anterior*.

Origin: Flethy from the inferior-anterior Spinous Process of the Os Ilium; and Tendinous, from the Dorsum of that Bone, a little above the Acetabulum. It runs down over the anterior part of the Cervix of the Os Femoris, and, in its passage along the fore-part of the Thigh, becomes gradually larger as far down as its middle, after which it decreases towards its lower extremity. In the middle of the Muscle, there is a longitudinal Tendinous Line, from which the Muscular Fibres run off like the Plumage of a Feather; the Tendon itself being most conspicuous behind.

Insertion: Tendinous, into the upper part of the Patella. Tab. XXXIV. Fig. 2. A.

Action: To extend the Leg.

CRURALIS, vel *Crureus*.

Origin: Flethy, from between the two Trochanters of the Os Femoris, but near the Minor; and from the fore-part of the Thigh-Bone to near its under extremity. Its sides are connected to both Vasti Muscles, and, below, it sends off a Tendon which joins that of the former Muscle.

Insertion: Into the upper and back part of the Patella, behind the Rectus. Tab. XXXV. Fig. 2. A.

Action: To assist in the extension of the Leg.

VASTUS EXTERNUS.

Origin: Broad, Tendinous, and Flethy, from the outer part of the Root of the Trochanter Major. Its Origin is continued from the Trochanter, along the whole

whole outer side of the *Linea Aspera*, to near the external Condyle of the *Os Femoris*, by *Fleshy Fibres*, which run obliquely forwards to a middle Tendon, where they terminate.

Insertion: Into the upper and outer part of the *Patella*, at the edge of the Tendon of the *Rectus*, with which it is connected. Part of it ends in an *Aponeurosis*, which is fixed to the Head of the *Tibia*, and afterwards is continued to the Leg. Tab. XXXV. Fig. 2. C.

Action: To extend the Leg.

VASTUS INTERNUS.

Origin: Tendinous and *Fleshy*, from between the fore-part of the *Os Femoris*, and root of the *Trochanter Minor*. The Origin is also continued along the whole inside of the *Linea Aspera*, by *Fibres* running obliquely forwards and downwards.

Insertion: Tendinous, at the side of the *Crureus* with which it is connected, into the upper and inner edge of the *Patella*, continuing *Fleshy* lower than the *Vastus Externus*. Part of it likewise ends in an *Aponeurosis*, which is fixed to the upper part of the *Tibia*, and afterwards is continued to the Leg. Tab. XXXV. Fig. 2. B.

Action: To assist the three former Muscles in extending the Leg; then the *Patella*, fixed to the Tubercle of the *Tibia* by a strong Ligament, supplies the office of a Pulley.

SEMITENDINOSUS.

Origin: Tendinous and *Fleshy*, in common with the long Head of the *Biceps*, from the posterior part of the Tuberosity of the *Os Ischium*. Its *Fleshy Belly* runs down the back part of the Thigh, and sends off a long roundish Tendon, which passes by the inner side of the Knee, and ends flat.

Insertion: Into the inside of the Ridge of the *Tibia*, a little below its Tubercle, and connected to the under edge of the *Gracilis*. Tab. XLI. Thigh, E.

Action: To bend the Leg, and, when bended, to roll it inwards.

SEMIMEMBRANOSUS.

Origin: By a broad flat Tendon, from the upper and back part of the Tuberosity of the *Os Ischium*. The *Fibres* composing the *Fleshy Belly*, run in an oblique direction towards a Tendon at the inner and under part of the Muscle, which is situated behind the *Semitendinosus*.

Insertion: Into the inner and back part of the Head of the *Tibia*. Tab. XLII. Thigh, C.

Action: To bend the Leg, and bring it directly backwards.

BICEPS FLEXOR CRURIS.

Origin: By two distinct Heads. The first, or *Long Head*, arises in common with the *Semitendinosus*, from the upper and back part of the Tuberosity of the *Os Ischium*. The second, or *Short Head*, arises from the *Linea Aspera*, a little below the termination of the *Gluteus Maximus*, by a *Fleshy* acute beginning, which soon grows broader, as it descends to join the first Head a little above the external Condyle of the *Os Femoris*.

Insertion: By a strong Tendon, into the upper part of the Head of the *Fibula*. Tab. XLI. Thigh, b, c.

Action: To bend the Leg.

The *Semitendinosus* and *Semimembranosus* form the *inner Ham-string*, and the *Biceps* the *outer Ham-string*; between the *Ham-strings* the great Vessels and Nerves are situated, which run to the Leg.

POPLITEUS.

Origin: By a small round Tendon, from the outer and under part of the external Condyle of the *Os Femoris*, and from the back part of the Capsular Ligament of the Joint. In passing the Joint, it becomes *Fleshy*, and spreads out, the *Fibres* running obliquely inwards and downwards, covered with a *Tendinous Membrane*.

Insertion: Thin and *Fleshy*, into a Ridge at the upper and inner edge of the *Tibia*, a little below its Head. Tab. XLII. Leg, G.

Action: To assist in bending the Leg, and, when bent, to roll it inwards. The Muscle also prevents the Capsular Ligament from being pinched.

MUSCLES SITUATED ON THE LEG AND FOOT, SERVING FOR THE MOTION OF THE FOOT AND TOES.

GASTROCNEMIUS EXTERNUS, vel *Gemellus*.

Origin: By two distinct Heads; one from the upper and back part of the internal Condyle of the *Os Femoris*, and from a little above the Condyle, by two separate beginnings: The other, *Tendinous* from the upper and back part of the external Condyle. A little below the Joint, their *Fleshy Bellies* meet in a middle Tendon, the union giving the appearance of a longitudinal *Raphè*. Below the middle of the *Tibia*, the Muscle sends off a

broad thin Tendon, which, becoming gradually narrower, joins that of the *Gastrocnemius Internus*, a little above the Ankle. Tab. XL. Leg, K, K, M.

GASTROCNEMIUS INTERNUS, vel *Soleus*.

Origin: By two Heads. The first from the back part of the Head, and upper and back part of the Body of the *Fibula*: The other from the back part of the *Tibia*, running inwards along the under edge of the *Popliteus*, towards

wards the inner part of the Bone, from which it receives Flethy Fibres for some way down. The Fleth of this Muscle, covered by the Tendon of the Gastrocnemius Externus, descends nearly as far as the extremity of the Tibia, a little above which the Tendons of both Gastrocnemii unite, and form a strong round Cord, called *Tendo Achillis*.

Insertion: Into the upper and back part of the Os Calcis, by the projection of which the Tendo Achillis is at a considerable distance from the Tibia. Tab. XLI. Leg, H, L.

Action: To extend the Foot, by raising the Heel.

PLANTARIS.

Origin: Thin and Flethy, from the upper and back part of the external Condyle of the Os Femoris, and from the Capsular Ligament of the Joint. A little below the Head of the Fibula, it sends off a long slender Tendon, which descends obliquely inwards, between the inner Heads of the Gastrocnemii, and afterwards runs along the inner edge of the Tendo Achillis.

Insertion: Into the inside of the posterior part of the Os Calcis, below the Tendo Achillis. Tab. XLI. Leg, I, f.

Action: To assist the Gastrocnemii, and to pull the Capsular Ligament of the Knee from between the Bones.

This Muscle is sometimes, though very seldom, wanting.

TIBIALIS ANTICUS.

Origin: Tendinous, from the upper part of the Tibia, between its Tubercle and Articulation with the Fibula. It then runs down, Flethy, on the outside of the Tibia, adhering to it and to the upper part of the Interosseous Ligament. Towards the under part of the Leg, it sends off a strong round Tendon, which passes under the Ligamentum Tarsi Annulare, near the inner Ankle.

Insertion: Tendinous, into the middle of the Os Cuneiforme Internum and Base of the Metatarsal Bone of the Great Toe. Tab. XXXIV. Leg, G, e.

Action: To bend the Foot.

TIBIALIS POSTICUS.

Origin: Flethy, from the upper and fore part of the Tibia, under the Process which joins it to the Fibula: Then, passing through a Fissure in the upper part of the Interosseous Ligament, it continues its Origin from the back part of the Fibula, next the Tibia, and from near one half of the upper part of the last-named Bone, as also from the Interosseous Ligament; the Fibres running towards a middle Tendon, which, in its descent, becomes round, and passes in a Groove behind the Malleolus Internus.

Insertion: Tendinous, chiefly into the upper and inner

part of the Os Naviculare, and partly into the under Surface of the Tarsal Bones by separate Slips, the last of which goes to the root of the Metatarsal Bone of the Middle Toe. Tab. XLIII. Leg, B.

Action: To extend the Foot, and, with the assistance of the Tibialis Anticus, to turn the Toes inwards, and the outer edge of the Foot downwards.

PERONEUS LONGUS, vel *Primus*.

Origin: Tendinous and Flethy, from the fore-part of the Head of the Fibula; and Flethy, from the outer part of that Bone, down to within a hand-breadth of the Ankle. The Fibres run in a Penniform manner towards a long Tendon, which becomes round, and, inclosed in a Sheath, passes through a Channel behind the Malleolus Externus. It is then reflected to the Sinuosity of the Os Calcis, runs along a Groove in the Os Cuboides, and goes obliquely across the Bones in the middle of the Sole.

Insertion: Tendinous, into the outside of the root of the Metatarsal Bone of the Great Toe, and partly into the Os Cuneiforme Internum. Tab. XLII. Leg, L, b.

Action: To extend the Foot a little, to draw it outwards, and to turn the inner edge of it downwards.

PERONEUS BREVIS, vel *Secundus*.

Origin: Flethy, from the outer part of the Fibula, beginning some way above the middle height of the Bone, and continuing its adhesion as far as the Malleolus Externus. The Fibres run, like those of the former Muscle, to an External Tendon, which becomes round, passes behind the outer ankle, where it is included in the same Sheath with the Tendon of the preceding Muscle, and there crossing behind that Tendon, runs forwards in a Sheath proper to itself.

Insertion: Tendinous, into the root and external part of the Metatarsal Bone of the Little Toe. Tab. XLII. Leg, M, C. Tab. XLIII. Leg, C.

Action: To assist the former Muscle in pulling the Foot outwards, its outer edge outwards, and to extend it in a small degree.

EXTENSOR LONGUS DIGITORUM.

Origin: Tendinous and Flethy, from the upper and outer part of the Head of the Tibia, and from the Head and almost the whole length of the anterior Spine of the Fibula. It arises, also, Flethy, from the Aponeurosis which covers the upper and outer part of the Leg, and from the Interosseous Ligament. Under the Ligamentum Tarsi Annulare, it splits into four round Tendons, which pass along the upper part of the Foot.

Insertion: Into the Base of the first Phalanx of the four small Toes, by flat Tendons which are expanded over the

the upper side of the Toes as far as the root of the last Phalanx. Tab. XXXV. Leg, *k, b*.

Action: To extend all the Joints of the four small Toes.

A portion of this Muscle is called by ALBINUS,

PERONEUS TERTIUS.

Origin: From the middle of the Fibula, in common with the Extensor Longus Digitorum. It continues down to near the Malleolus Externus, and sends its Fleishy Fibres forwards to a Tendon which passes under the Ligamentum Annulare.

Insertion: Into the root of the Metatarsal Bone of the Little Toe. Tab. XLVI. Left Leg, W, V.

Action: To assist in bending the Foot.

EXTENSOR BREVIS DIGITORUM.

Origin: Fleishy and Tendinous, from the outer and fore part of the Os Calcis; soon forming a Fleishy Belly, which is divided into four Portions. These send off an equal number of Tendons, which pass over the upper part of the Foot, crossing under those of the former Muscle.

Insertion: By four slender Tendons, into the Tendinous Expansion continued from the Long Extensors of all the Toes, excepting the little one. Tab. XXXVI. Foot, O.

Action: To assist in the extension of the Toes.

APONEUROSIS PLANTARIS.

This, like the Aponeurosis Palmaris, is a strong Tendinous Expansion, which covers the Muscles, Vessels, and Nerves of the Sole.

It arises from the Tuberosity at the under and back part of the Os Calcis, and is divided into three Portions, which run forwards to be connected to the Heads of the Metatarsal Bones of all the Toes. The middle Portion is subdivided into five Slips, which split at the roots of the Toes, and embrace the Tendons of the Flexor Muscles. Tab. L. Fig. 8. *a, b, c*.

Besides serving the general purpose of Aponeuroses, it performs the office of a Ligament, by binding the two ends of the Arch of the Foot together.

FLEXOR BREVIS DIGITORUM,

Vel Flexor Sublimis, vel Perforatus.

Origin: Narrow and Fleishy, from the inferior-anterior part of the Tuberosity of the Os Calcis, and from the Aponeurosis Plantaris. It forms a thick Fleishy Belly, which sends off four small Tendons, and these split for the passage of the Tendons of the Flexor Longus.

Insertion: Into the second Phalanx of the four small Toes. Tab. L. Fig. 9. *a*.

Action: To bend the first and second Joints of the Toes, but particularly the second.

The Tendon of the Little Toe is frequently a-wanting.

FLEXOR LONGUS DIGITORUM,

Vel Flexor Profundus, vel Perforans.

Origin: By an acute Tendon, which soon becomes Fleishy, from the back part of the Tibia, at the under edge of the Popliteus; and this beginning is continued down the inner edge of the Bone, by short Fleishy Fibres ending in its Tendon. It arises also by Tendinous and Fleishy Fibres, from the outer edge of the Tibia; and between this double order of Fibres the Tibialis Posticus lies inclosed. Having gone under two Annular Ligaments, it passes through a Sinuosity at the inside of the Os Calcis; and about the middle of the Sole, receives a Tendon from the Flexor Longus Pollicis. It then divides into four Tendons which run through the Slits of the Perforatus.

Insertion: Into the Base of the third Phalanx of the four smaller Toes; the Tendons of this, as well as of the Flexor Brevis, being inclosed upon the Toes by Annular Ligaments. Tab. XLII. Leg, I. Tab. L. Fig. 10. *a, f, f*.

Action: To bend the different Joints of the Toes, particularly the last one.

FLEXOR DIGITORUM ACCESSORIUS,

Vel Massa Carnea Jacobi Sylvii.

Origin: By two Portions; the inner Fleishy, from the Sinuosity of the Os Calcis; the outer Tendinous, but soon becoming Fleishy, from the fore and outer part of that Bone.

Insertion: Into the Tendon of the Flexor Longus Digitorum, before it divides into smaller Tendons. Tab. L. Fig. 10. *b, c, d*.

Action: To assist the Flexor Longus.

LUMBRICALES.

Origin: By four Tendinous and Fleishy beginnings, from the Tendon of the Flexor Profundus, just before its division. They run forwards, under the same general appearance with those in the Hand, but are somewhat smaller.

Insertion: By four slender Tendons, at the inside of the first Joint of the four small Toes, into the Tendinous Expansion sent from the Extensors to cover the upper part of the Toes. Tab. L. Fig. 10. *g, h, i, k*.

Action: To increase the Flexion of the Toes, and to draw them inwards.

EXTENSOR

EXTENSOR PROPRIUS POLLICIS,

Vel *Extensor Longus*.

Origin: By an acute, Tendinous, and Flešhy beginning, from the fore-part of the Fibula, some way below its Head. It continues its Origin from the same Bone to near the outer Ankle, by Flešhy Fibres, which descend obliquely towards a Tendon.

Insertion: Tendinous, into the posterior part of both the Bones of the Great Toe. Tab. XXXV. Leg, *d, e, f*.

Action: To extend the Great Toe.

FLEXOR LONGUS POLLICIS.

Origin: Tendinous and Flešhy, from the back part of the Fibula, some way below its Head; being continued down the same Bone, almost to its under end, by a double order of oblique Flešhy Fibres. Its Tendon passes under an Annular Ligament at the inner Ankle.

Insertion: Into the last Joint of the Great Toe. Tab. XLII. Leg, K.

Action: To bend the Great Toe; particularly the last Joint.

FLEXOR BREVIS POLLICIS.

Origin: Tendinous, from the under and fore part of the Os Calcis, and from the Os Cuneiforme Externum. It is inseparably united with the Abductor and Adductor Pollicis.

Insertion: Into the external Os Sefamoideum, and root of the first Bone of the Great Toe. Tab. L. Fig. 11. *i, k*.

Action: To bend the first Joint of the Great Toe.

ABDUCTOR POLLICIS.

Origin: Flešhy, from the anterior and inner part of the Protuberance of the Os Calcis, and Tendinous from the same Bone, where it joins with the Os Naviculare.

Insertion: Tendinous, into the internal Os Sefamoideum, and root of the first Bone of the Great Toe. Tab. L. Fig. 9. *c*.

Action: To pull the Great Toe from the rest.

ADDUCTOR POLLICIS.

Origin: By a long thin Tendon, from the under part of the Os Calcis, from the Os Cuboides, from the Os Cuneiforme Externum, and from the root of the Metatarsal Bone of the Second Toe. The Muscle is divided into two Flešhy Portions.

Insertion: Into the external Os Sefamoideum, and root of the Metatarsal Bone of the Great Toe. Tab. L. Fig. 11. *g, h*.

Action: To pull the Great Toe towards the rest.

ABDUCTOR MINIMI DIGITI.

Origin: Tendinous and Flešhy, from the edge of a Cavity on the under part of the Protuberance of the Os Calcis, and from the root of the Metatarsal Bone of the Little Toe.

Insertion: Into the outer part of the root of the first Bone of the Little Toe. Tab. L. Fig. 9. *d, e*.

Action: To draw the Little Toe outwards.

FLEXOR BREVIS MINIMI DIGITI.

Origin: Tendinous, from the Os Cuboides, near the Groove for lodging the Tendon of the Peroneus Longus, and Flešhy, from the outer and back part of the Metatarsal Bone of this Toe.

Insertion: Into the anterior extremity of the Metatarsal Bone, and root of the first Bone of the Little Toe. Tab. XXXV. Fig. 11. *f*.

Action: To bend this Toe.

TRANSVERSALIS.

Origin: Tendinous, from the under and fore part of the Metatarsal Bone of the Great Toe, and from the internal Os Sefamoideum of the first Joint. It forms a Flešhy Belly, which runs transversely between the Metatarsal Bones and Flexor Muscles of the Toes.

Insertion: Tendinous, into the under and outer part of the anterior extremity of the Metatarsal Bone of the Little Toe, and Ligament of the next Toe. Tab. L. Fig. 11. *l*.

Action: To contract the Foot, by bringing the roots of the outer and inner Toes towards each other.

INTEROSSEI.

The Interossei arise Tendinous and Flešhy from, and fill the spaces between, the Metatarsal Bones. Three, called *Interni*, arise with single Heads, and are placed in the Sole; and four, termed *Externi* or *Bicipites*, arise with double Heads, and appear on both sides of the Foot.

The Insertion of all the Interossei is by slender Tendons, into the Expansion sent off from the Tendons of the Lumbricales and of the Extensor Muscles of the Toes.

INTEROSSEI INTERNI.

Tab. L. Fig. 12.

PRIOR, vel *Abductor Medii Digiti*.

Origin: From the inside of the Metatarsal Bone of the Middle Toe.

Insertion: Into the inside of the root of the first bone of the Middle Toe.

Action: To pull the Middle Toe inwards.

PRIOR, vel *Abductor Tertii Digiti*.

Origin: From the inner and under part of the Metatarsal Bone of the Third of the small Toes.

Insertion:

Insertion : Into the inside of the root of the first Bone of the Third Toe.

Action : To pull the Third Toe inwards.

PRIOR, vel *Adductor Minimi Digiti*.

Origin : From the inside of the Metatarsal Bone of the Little Toe.

Insertion : Into the inside of the root of the first Bone of the Little Toe.

Action : To pull the Little Toe inwards.

INTEROSSEI EXTERNI, vel *Bicipites*.

Tab. L. Fig. 12. Tab. XXXVII. Foot, F.

PRIOR, vel *Abductor Indicis*.

Origin : From the contiguous sides of the Metatarsal Bones of the Great and Fore Toes.

Insertion : Into the inside of the root of the first Bone of the Fore-Toe.

Action : To pull the Fore-Toe inwards.

POSTERIOR, vel *Adductor Indicis*.

Origin : From the contiguous sides of the Fore-Toe and Second of the small Toes.

Insertion : Into the outside of the root of the first Bone of the Fore-Toe.

Action : To pull the Fore-Toe outwards.

POSTERIOR, vel *Adductor Medii Digiti*.

Origin : From the contiguous sides of the Metatarsal Bones of the Second and Third of the small Toes.

Insertion : Into the outside of the root of the first Bone of the Second of the small Toes.

Action : To pull this Toe outwards.

POSTERIOR, vel *Adductor Tertii Digiti*.

Origin : From the contiguous sides of the Metatarsal Bones of the Third and Fourth of the small Toes.

Insertion : Into the outside of the root of the first Bone of the Third of the small Toes.

Action : To pull this Toe outwards.

TABLE

TAB. 33.

Fig. 3.

Fig. 4.

Fig. 1.

Fig. 2.



T A B L E XXXIII.

Represents the APONEUROSES OR TENDINOUS MEMBRANES, which appear upon removing the Common INTEGUMENTS, and which cover the MUSCLES of the EXTREMITIES.

FIG. 1.

Gives a View of the APONEUROSIS which covers the MUSCLES on the Fore-part of the SUPERIOR EXTREMITY.

- A, The deltoid muscle.
- B, The aponeurosis sent off from the tendons of the muscles on the shoulder to cover the flexor muscles of the fore-arm.
- C, The aponeurosis continued from the fore-side of the arm, joined to a thicker and stronger one sent off from the tendon of the biceps flexor cubiti, to cover the muscles on the anterior part of the fore-arm.
- D, The continuation of this aponeurosis, covering the tendons of the flexor muscles of the hand and fingers, which, on account of its thinness, are seen shining through it.
- E, The aponeurosis palmaris, which is connected above, chiefly to the tendon of the palmaris longus, and to the anterior transverse ligament of the wrist, and below, to the roots of the four fingers by an equal number of double slips.
- F, The musculus abductor pollicis.
- G, The palmaris brevis.

FIG. 2.

Shews the APONEUROSIS upon the Back Part of the SUPERIOR EXTREMITY.

- A, The deltoid muscle.
- B, An aponeurosis covering the infra-spinatus, similar to one which covers the supra-spinatus.
- C, The aponeurosis which covers the back part of the arm, sent down from the tendons of the muscles on the shoulder, and with the aponeurosis on the fore-side of the arm, forming a sheath, which incloses the muscles on this part.
- D, The aponeurosis of the back part of the fore-arm, continued from that which covers the back part of the upper arm, and likewise from the tendon of the triceps extensor cubiti; many of the fibres intermixing with, and decussating each other, at the opposite sides of the fore-arm.
- E, A thick and strong portion of the aponeurosis on the

back part of the fore-arm, forming the ligamentum carpi annulare posterius.

The tendinous sheaths which cover the Fingers are here but faintly represented.

FIG. 3.

Gives a View of the APONEUROSIS upon the Fore-part of the INFERIOR EXTREMITY.

- A, The thick and strong aponeurosis at the fore and outer part of the thigh, sent down from the muscles about the pelvis, and from the under end of the external oblique muscles of the abdomen.
- B, An aponeurosis covering the muscles upon the fore and inner part of the thigh, much thinner and weaker than that farther out.
- C, An attachment of the aponeurosis of the thigh to the head of the tibia.
- D, The aponeurosis of the inside of the thigh, fixed to the corresponding side of the tibia.
- E, F, The aponeurosis sent from the fascia of the thigh, and from the extensor muscles of the leg, to cover the muscles on the fore-part of the leg. This aponeurosis, like that on the thigh, is thick and strong at the outer, and becomes gradually thinner towards the inner part of the leg.
- G, H, Parts of this aponeurosis thicker and stronger than the rest, forming the superior and inferior ligaments of the tarsus.

FIG. 4.

APONEUROSIS on the Back Part of the INFERIOR EXTREMITY.

- A, The gluteus magnus.
- B, The aponeurosis of the back part of the thigh, arising from the tendons of the glutei, and from those of the loins, fixed to the linea aspera of the os femoris.
- C, Continuation of this aponeurosis covering the muscles, nerves, &c. of the ham.
- D, That part of the aponeurosis which covers the gemelli. From this part, the aponeurosis is continued down, and lost upon the Foot.

TABLE

T A B L E XXXIV.

A VIEW of the First LAYER of MUSCLES on the Anterior Part of the BODY, after the INTEGUMENTS and APONEUROSES have been removed.

FIG. 1.

HEAD and NECK.

- A, The frontal part of the occipito-frontalis.
a, The tendon of the occipito-frontalis.
b, A fleshy slip sent from the occipito-frontalis over the root of the nose.
 B, The attollens aurem.
 C, The anterior auris.
 D, The orbicularis palpebrarum.
c, The ciliary part of the orbicularis.
d, The compressor naris.
 E, F, The levator labii superioris alæque nasi.
 G, The zygomaticus minor.
 H, The zygomaticus major.
e, The levator anguli oris.
 I, The cartilage of the nose.
 K, The depressor anguli oris.
f, The depressor labii inferioris.
g, The buccinator.
 L, The orbicularis oris.
h, The masseter.
 M, The platysma myoides.
i, The upper end of that muscle.
k, The under end.
l, The sterno-cleido-mastoideus.

TRUNK.

- A, The pectoralis major.
 B, B, Part of the serratus magnus.
 C, The edge of the latissimus dorsi.
 D, D, The obliquus descendens externus.
 E, E, The linea alba.
 F, The umbilicus.
 G, G, The linea femilunaris.
m, The tendon at the outer side of the linea femilunaris.
n, The obliquus internus, shining through the tendon of the obliquus externus.
o, o, The rectus abdominis, also shining through the tendon.
 H, H, The tendinous interfections of the rectus.

- p*, The pyramidalis.
 I, The ring of the external oblique muscle, transmitting the spermatic cord.
 K, The cremaster, covering the spermatic cord.
 L, The under end of that muscle, covering the tunica vaginalis testis.
q, The lower edge of the external oblique muscle, called Poupart's Ligament.

SUPERIOR EXTREMITY.

- A, Part of the trapezius.
 B, The deltoides.
 C, C, The biceps flexor cubiti; at the lower C, the aponeurosis cut off.
s, The round tendon of the biceps.
 D, The long head of the triceps extensor cubiti.
 E, The edge of the brachialis internus.
t, The third head of the triceps extensor cubiti, called brachialis externus.
 F, The supinator longus.
 G, The pronator radii teres.
 H, The flexor carpi radialis.
 I, The flexor sublimis.
 K, The extensor ossis metacarpi pollicis.
 L, The extensor primi internodii.
 M, The extensor secundi internodii.
 N, The ligamentum carpi annulare posterius crossing the extensors of the hand and fingers.—See the muscles on the hand in Tab. XXXV. Fig. 1.

INFERIOR EXTREMITY.

- A, The tensor vaginæ femoris.
 B, The anterior edge of the gluteus maximus.
 C, The lower end of the iliacus internus.
 D, The lower end of the psoas magnus.
 E, The pectinalis.
 F, The adductor longus femoris.
 G, The gracilis.
 H, The sartorius.
 I, The rectus.

FIG.

Fig 1

Fig 2





FIG. 2.

- A, Continuation of the rectus femoris,
 B, ————— of the sartorius,
 C, ————— of the adductor longus femoris,
 D, ————— of the gracilis,
 E, ————— of the vastus externus, and,
 F, ————— of the vastus internus.
a, Part of the patella.
b, The ligament fixing the patella to the tibia.
c, The tendons of the sartorius, gracilis, and femimem-
 branosus.
d, The under end of the biceps flexor cruris.
 G, The tibialis anticus ;
- e*, Its tendon.
 H, The peroneus longus.
 I, The extensor longus digitorum pedis.
f, The tendons of the extensor longus digitorum.
g, The extensor proprius pollicis.
 K, The gastrocnemius externus.
 L, L, The gastrocnemius internus.
 M, The flexor longus digitorum pedis.
h, The tibialis posticus.
i, The tendo Achillis and tendon of the plantaris.
 N, N, The upper and under portions of the ligamentum
 tarfi annulare.
k, The ligaments retaining the tendons at the inner ankle.
 O, O, The abductor pollicis.

TABLE

T A B L E XXXV.

A VIEW of the Second LAYER of MUSCLES on the Anterior Part of the Body.

FIG. 1.

HEAD and NECK.

- A, The corrugator supercilii.
a, The levator palpebræ superioris.
 B, The temporalis.
b, Its tendon passing under the zygoma.
 C, The masseter.
 D, The levator anguli oris.
 E, The buccinator.
 F, The orbicularis oris.
f, The nasalis labii superioris, at the upper side of which is a portion of the depressor labii superioris aëque nasi.
 G, The depressor labii inferioris.
 H, The sterno-cleido-mastoideus.
 I, The sterno-hyoideus.
c, The trachea seen obscurely.
 K, The omo-hyoideus.
 L, The hyo-thyroideus.
d, The os hyoides.
 M, The levator scapulæ.
e, The scalenus medius.

TRUNK.

- A, The subclavius.
 B, The pectoralis minor.
 C, C, The serratus magnus.
g, &c. Intercostrales interni, the tendinous fascia being removed.
 D, D, The rectus abdominis, with the tendon of the obliquus internus covering its outer edge;—on the right side it is entirely exposed.
b, b, Tendinous interfections of the rectus abdominis.
 E, The pyramidalis.
 F, The obliquus internus.
i, k, The tendon of the obliquus internus. Between *i* and *k*, the tendon splits into two layers, which inclose the rectus; from *k* to the pubis, the whole tendon goes before the rectus.
l, l, A portion of the tendon of the obliquus internus remaining upon the outer edge of the rectus.
 G, The cremaster testis.

SUPERIOR EXTREMITY.

- A, The biceps flexor cubiti;
a, Its short head;

- b*, Its long head.
c, A section of the aponeurotic tendon of the biceps;
d, Its round tendon.
 B, The coraco-brachialis.
e, The subscapularis of the right side.
f, The teres major of the right side.
 C, The under end of the brachialis internus.
 D, The long head of the triceps extensor cubiti.
g, That part of the triceps called brachialis externus.
 E, Extensor carpi radialis, longior et brevior.
 Upper F, Extensor ossis metacarpi pollicis.—Lower F, Extensor primi internodii.
b, Extensor secundi internodii pollicis.
 G, Flexor sublimis perforatus.

See also Tab. XXXV. Fig. 2.

PELVIS, and INFERIOR EXTREMITY.

- A, The under end of the iliacus internus.
 B, The under end of the psoas magnus.
 C, The pectinalis.
 D, The cut end of the rectus femoris.
 E, The anterior edge of the gluteus medius.

FIG. 2.

- A, The cruralis, with its tendinous fascia.
 B, The vastus internus.
 C, ——— externus.
 D, The cut tendon of the rectus fixed to the patella.
 E, The adductor longus femoris.
 F, The gracilis.
 G, The tendons of the gracilis and semitendinosus.
 H, The tendon of the biceps flexor cruris.
 I, The peroneus longus.
a, The peroneus brevis.
 K, The extensor longus digitorum pedis.
b, The tendons of the extensor longus digitorum.
c, The peroneus tertius.
d, The extensor proprius pollicis.
e, The tendon of the extensor proprius pollicis.
f, A branch of the tendon of the extensor proprius pollicis, not always found.
 L, The edge of the gastrocnemius internus.
 M, The edge of the flexor longus digitorum pedis.
 N, The tendons of the tibialis posticus, and flexor longus digitorum pedis.
 O, The flexor brevis digitorum pedis.

TABLE

Fig 1



Fig 2



Fig. 1

TAB. 36.

Fig. 2



T A B L E XXXVI.

A VIEW of the Third LAYER of MUSCLES on the Anterior Part of the BODY.

FIG. 1.

HEAD and NECK.

- A, The infertion of the abductor oculi.
a, The adductor oculi of the right side.
 B, The infertion of the levator oculi.
 C, The trochlea, and part of the tendon of the obliquus superior.
 D, The obliquus inferior oculi, immediately above which is the infertion of the depressor oculi.
 E, The depressor labii superioris alæque nasi.
 F, The orbicularis oris.
 G, The buccinator.
 H, The levator labii inferioris.
b, Part of the pterygoideus externus.
c, Part of the pterygoideus internus.
 I, The sterno-thyroideus.
 K, The thyro-hyoideus.
d, The os hyoides.
e, The thyroid cartilage.
f, The cricoid cartilage, with the two crico-thyroid muscles arising from it.
g, The trachea.
h, Part of the pleura.
 L, The scalenus anticus.
 M, N, The scalenus medius.
i, A portion of the trachelo-mastoideus.
 O, The rectus capitis anterior major.
k, The longus colli.
l, The constrictor pharyngis inferior.

TRUNK.

- A, A, &c. Anterior portions of the intercostales externi.
 B, B, &c. Anterior part of the intercostales interni.
 C, The fleshy part of the transversalis abdominis;
 F, Its tendon.
 D, D, The cut edge of that part of the tendon of the transversalis muscle which joins the obliqui, and passes before the rectus and pyramidalis.
 E, E, That part of the tendon which passes behind the rectus, and is covered by,
 G, G, The posterior layer of the tendon of the obliquus internus.
m, m, The remains of the tendons of the oblique muscles, forming the linea alba.
 H, The umbilicus.
 I, The spermatic vessels passing under the edge of the transverse muscle.
n, The peritoneum.

- o, o, p*, A print of the two umbilical arteries *o, o*, and of the urachus *p*, upon the peritoneum.
y, The penis cut across, in which are seen the corpora cavernosa penis et urethræ.

SUPERIOR EXTREMITY.

- A, The subscapularis;
a, Its tendon.
 B, The teres major of the right side;
b, Its tendon.
 C, The coraco-brachialis.
 D, H, The brachialis internus.
 E, The brachialis externus.
 F, The extensor carpi radialis brevior.
 G, The extensor carpi radialis longior.
 I, The flexor digitorum profundus.
 K, The flexor longus pollicis.
 L, The flexor brevis pollicis.

See also Tab. L. Fig. 3.

PELVIS, and INFERIOR EXTREMITY.

- A, The gluteus minimus.
 B, The iliacus internus.
 C, The psoas magnus.
 D, The obturator externus.
 E, The adductor brevis.
 F, The adductor magnus.
 G, The gracilis.

FIG. 2.

- A, The adductor brevis.
 B, C, The adductor magnus.
 D, The gracilis.
a, The femimembranosus;
b, Its infertion into the tibia.
 E, The under end of the biceps flexor cruris.
 F, F, The os femoris.
 G, The patella.
 H, The tubercle of the tibia.
 I, I, The edges of the femilunar cartilages.
 K, The peroneus longus.
c, The peroneus brevis.
 L, The tibialis posticus.
 M, The flexor longus digitorum pedis.
 N, The tendon of the tibialis posticus.
d, The tendon of the flexor longus digitorum pedis.
e, The flexor digitorum accessorius.
 O, The extensor brevis digitorum pedis.

S

TABLE

T A B L E XXXVII.

A VIEW of the FOURTH LAYER of MUSCLES situated on the Anterior Part of the Body.

FIG. 1.

HEAD and NECK.

- A, The levator palpebræ superioris.
a, The levator oculi.
 B, The adductor oculi.
 C, The abductor oculi.
 D, The depressor oculi.
 E, The obliquus superior.
 F, The obliquus inferior.
 G, The pterygoideus internus.
 H, The obliquus superior capitis.
 I, The scalenus medius.
 K, *b, c, d, e*, The longus colli.
f, f, f, Intertransversales colli.

TRUNK.

- A, A, A, Intercostales interni.
a, a, Intercostales externi.
 B, B, *b*, The convex part, or thoracic side of the diaphragm;
b, The anterior point of its middle tendon.
c, d, e, f, g, h, The fleshy origins of the diaphragm, from the cartilago ensiformis, peritoneum, seventh, eighth, ninth, tenth, and eleventh ribs.
i, i, i, i, The first heads, or tendinous crura of the inferior muscle of the diaphragm;
k, The passage for the aorta, between these heads;
 The second and third heads of the inferior muscle of the diaphragm are situated between the upper ends of the first heads and the psoæ muscles, but are not represented here.
l, The fourth head;
m, Another head, sometimes found connected with the quadratus lumborum;
n, n, The fleshy crura from the joining of these heads.
o, Fibres crossing each other, under,
p, The passage for the œsophagus.

- q*, The middle tendon on the left side, with its fibres de-
 cuffating.
r, Origin of the diaphragm, from the twelfth rib.
 C, The psoas parvus on the right side, that on the left
 being removed.
s, The tendon of the psoas parvus passing down to be
 fixed to the brim of the pelvis.
 D, The quadratus lumborum.
 E, A section of the penis.
t, The corpus cavernosum penis of the left side.
u, The corpus cavernosum urethræ.
v, The erector penis.
w, The accelerator urinæ.
x, The sphincter ani.
y, The transversalis perinei.

SUPERIOR EXTREMITY.

- A, The subscapularis.
 B, The supinator brevis.
 C, The flexor brevis pollicis.
 D, The adductor pollicis.

See also Tab. L. Fig. 4.

PELVIS, and INFERIOR EXTREMITY.

- A, The psoas magnus.
z, Its origin, chiefly from the lumbar vertebræ.
 C, Its passage out of the abdomen, along with,
 B, The iliacus internus.
 C, The obturator externus.
 D, The upper part of the adductor magnus.



FIG. 2.

- A, B, C, The continuation of the adductor magnus.
a, The insertion of the psoas magnus and iliacus internus.
 D, The tibialis posticus, the interosseous ligament being
 removed;
b, Its tendon.
 E, The peroneus brevis.
 F, The interossei externi.

TABLE

Fig. 1

Fig. 2






FIG. 8.




FIG. 9.





FIG. I.



FIG. 10.





T A B L E XXXVIII.

Represents the Common INTEGUMENTS, some of the MUSCLES and GLANDS of the HEAD and NECK, with the First LAYER of MUSCLES on the Right, and Second Layer on the Left Side of the Anterior Part of the BODY.

FIG. 1.

A HAIR, viewed with a Microscope.

- A, The root.
B, The bearded body.
C, The small extremity.

FIG. 2.

The Cuticle of the Hand, with the Nails adhering to it.

FIG. 3.

A Toe, with the Cuticle taken off, to shew the villous appearance of the Exterior Surface of the Skin.

FIG. 4.

A Piece of Skin, according to RUYSCH, with the Papillæ Pyramidales, as they appear to the naked Eye.

FIG. 5.

The piece of Skin, Fig. 4. seen with a Microscope.

FIG. 6.

The Corpus Reticulare of the Skin, seen with the naked Eye.

FIG. 7.

The Corpus Reticulare, Fig. 6. viewed with a Microscope.

FIG. 8.

The MUSCLES, GLANDS, &c. of the Left Side of the FACE and NECK, after the Common INTEGUMENTS and PLATYSMA MYOIDES have been removed.

- a*, The frontal muscle.
b, The temporal muscle, on which the larger branches of the temporal artery are seen.
c, c, The orbicularis palpebrarum.
d, The orbicularis oris.
e, The levator labii superioris.

- f*, The levator anguli oris.
g, The zygomaticus major.
h, The depressor anguli oris.
i, The depressor labii inferioris.
k, The buccinator.
l, The masseter.
m, The parotid gland;
n, Its duct.
o, o, The facial artery.
p, p, The facial vein.
q, The anterior heads of the digastrici.
r, The inferior maxillary gland.
s, The sterno-hyoidei.
t, t, The omo-hyoideus.
u, Continuation of the facial vein.
v, The sterno-mastoideus.
w, The trapezius.
x, The levator scapulæ.
y, The scalenus medius.
z, The scalenus anticus.
α, One of the nerves of the superior extremity.

FIG. 9.

On the RIGHT SIDE, the MUSCLES immediately under the Common Integuments, on the Anterior Part of the BODY, are represented; on the LEFT SIDE, the MUSCLES are seen, which come in view when the Exterior Set have been removed.

For the explanation of these Muscles, see Tab. XXXIV. and XXXV. to which add here the First Layer of Muscles on the

RIGHT FORE-ARM AND HAND.

- W*, The supinator radii longus;
X, Its tendon.
Y, The pronator radii teres.
Z, The flexor carpi radialis;
α, Its tendon.
β, The palmaris longus;

- γ, Its tendon.
 δ, δ, Parts of the flexor digitorum sublimis.
 ε, The tendon of the flexor carpi ulnaris.
 ζ, The flexor longus pollicis;
 η, Its tendon, inserted into the last joint of the thumb.
 θ, Part of the pronator radii quadratus.
 ι, Part of the extensor primi et secundi internodii pollicis;
 κ, Their tendons;
 λ, Their annular ligament.
 μ, The tendinous aponeurosis of the palm.
 ν, The transverse ligament of the wrist.
 ξ, The palmaris brevis.
 ο, Part of the flexor primi internodii pollicis.
 π, The abductor pollicis;
 ς, Its tendon, forming an aponeurosis with the extensors.
 ς, Part of the flexor secundi internodii.
 τ, The annular sheath of the tendon of the flexor longus pollicis.
 υ, Part of the adductor pollicis.
 φ, The tendon of the adductor indicis, and first lumbricalis.
 χ, The abductor minimi digiti.
 ψ, The flexor brevis minimi digiti.
 1. 2. 3. The annular sheaths of the tendons of the flexors of the fore-finger. These ligaments are also represented in the other fingers.
 4. The tendons of the lumbricales and interossei, which may also be seen on the sides of the other fingers.

FIG. 10.

The Second LAYER of MUSCLES of the FACE and NECK, after the First has been removed.

See them described Tab. XXXV.

LEFT FORE-ARM AND HAND of Fig. 9.

- t, The extensor carpi radialis longior.
 u, Part of the extensor carpi radialis brevior.
 v, The supinator radii brevis.
 w, The cut extremity of the pronator radii teres.
 x, Part of the flexor carpi ulnaris;
 y, Its tendon.
 z, The flexor digitorum sublimis;
 A, Its tendons.
 B, Part of the pronator radii quadratus.
 C, The extensors of the thumb.
 D, The flexor pollicis longus;
 E, Its tendon, near its insertion.
 F, The flexor ossis metacarpi pollicis.
 G, The flexor brevis pollicis.
 H, The flexor parvus minimi digiti.
 I, The abductor minimi digiti.
 K, K, The first and second lumbricales; the third and fourth are also in view, but unlettered.
 L, The tendons of the lumbricales and interossei, which may also be seen on the sides of the other fingers.
 M, The tendons of the flexor digitorum sublimis, divided near their insertion, for the passage of the tendons of the flexor profundus, marked N.

TABLE

FIG. 1.



FIG. 2.




FIG. 3.




FIG. 4.




FIG. 8.






FIG. 7.




FIG. 6.




FIG. 9.



FIG. 5.



R. Cooper, Sculp.

T A B L E XXXIX.

Represents the Parts situated under those shewn in Fig. 9. Tab. XXXVIII. together with the EYE-LIDS, LACRYMAL GLAND and DUCTS, and MUSCLES of the EYE.

FIG. 1.

The Muscles represented here are explained Tab. XXXVI. and XXXVII. excepting those on the right side of the Trunk, and on the under part of the Fore-Arms and Hands.

LEFT FORE-ARM AND HAND.

- S, The extensor carpi radialis longior.
- T, Part of the extensor carpi radialis brevior.
- U, The supinator radii brevis.
- V, The flexor digitorum profundus;
- W, Its tendons;
- X, X, Their insertions into the last joint of each of the fingers.
- Z, The transverse ligament of the wrist.
- y, y, y, y, The four lumbricales.
- α, The flexor longus pollicis;
- β, Its tendon, inserted into the last joint of the thumb.
- δ, The flexor brevis pollicis.
- ε, The adductor ossis metacarpi minimi digiti.
- ζ, The os pisiforme.

RIGHT SIDE OF THE TRUNK.

- A, A, The intercostales externi.
- B, B, The internal intercostals.
- θ, The mammary artery and vein.
- i, i, i, i, The triangularis, or sterno-costalis.
- κ, κ, The surface of the lungs appearing through the pleura.
- λ, λ, λ, The peritoneum, through which the bowels appear obscurely.—Between this and the linea alba, the vestige of the epigastric artery is seen.
- μ, The spermatic cord, coming out behind the peritoneum.

RIGHT FORE-ARM AND HAND.

- c, The supinator radii brevis.
- d, The pronator radii quadratus.
- e, The flexor brevis pollicis.
- f, The sesamoid bones into which it is inserted.

- g, The adductor pollicis.
- b, i, k, l, The seven interossei.

FIG. 2.

Described Tab. LXXII. Fig. 9.

Shews the Lacrymal Canals, the Teguments and Bones being cut away.

FIG. 3.

Described Tab. LXXII. Fig. 8.

Represents the Palpebræ inverted, to obtain a view of the Lacrymal Canals.

FIG. 4.

The two EYE-LIDS cut from each other, at the exterior CANTHUS.

- A, A, The interior membrane of the eye-lids.
- B, The caruncula lacrymalis.
- C, D, The edges of the eye-lids, with the small orifices of the excretory ducts of the sebaceous glands.
- E, The puncta lacrymalia.
- F, F, The eye-lashes.

FIG. 5. 6. 7. 8. and 9.

Shew the MUSCLES of the EYE.

- A, A, The two optic nerves before they meet;
- B, These nerves conjoined.
- C, The nerve of the right eye.
- D, The attollens palpebram;
- d, Its tendon.
- E, The attollens oculum.
- F, The abductor.
- G, The obliquus superior, or trochlearis;
- H, I, Its tendon.
- K, The adductor.
- L, The deprimens.
- M, The obliquus minor.
- Z, The ball of the eye.
- X, Part of the frontal bone.
- y, Part of the maxillary bone.

TABLE

T A B L E XL.

A VIEW of the First LAYER of MUSCLES on the Posterior Part of the BODY, after the INTEGUMENTS and APONEUROSES have been removed.

FIG. 1.

HEAD, NECK, and TRUNK.

- A, The occipital part of the occipito-frontalis;
a, The fleshy,
b, The tendinous part of this muscle;
c, A tendinous membrane joining its two sides;
d, Part of the tendinous membrane, covering the upper end of the temporal muscle.
 B, The attollens aurem.
 C, The anterior auris.
 D, A small part of the retrahentes aurem.
 E, The back part of the orbicularis palpebrarum.
 F, The zygomaticus major.
 G, The masseter.
e, The pterygoideus internus.
f, The platysma myoides.
 H, The sterno-cleido-mastoideus.
 I, I, I, The trapezius.
g, *h*, Its insertion into the spine of the scapula and outer end of the clavicle.
i, *i*, Its tendinous portion, in the nape of the neck, called *Ligamentum Nuchæ*.
 K, K, K, The latissimus dorsi;
k, *k*, Its tendon.
 L, Part of the obliquus externus abdominis.
m, Part of the insertion of the rhomboides.
n, Part of the sacro-lumbalis.

SUPERIOR EXTREMITY.

- A, The deltoides;
l, Its insertion into the os humeri.
 B, The infra-spinatus.
 C, The teres minor.
 D, The teres major.
 E, The triceps extensor cubiti.
o, The long,
p, The short head of the triceps.
q, The third head, called *Brachialis Externus*.
r, The common tendon of these three heads.
s, The brachialis internus.
 F, The anconens.
 G, The supinator radii longus.
 H, The extensor carpi radialis longior.
 I, The extensor carpi radialis brevior.
 K, The extensor digitorum communis.

- L, The extensor ossis metacarpi pollicis.
 M, The extensor primi internodii pollicis.
 N, The extensor proper to the little finger.
 O, The extensor carpi ulnaris.
 P, The palmaris longus.
t, The flexor sublimis perforatus.
 Q, The flexor carpi ulnaris.
 R, *u*, Part of the flexor profundus perforans.
v, The ligamentum carpi annulare posterius.
 See the Muscles on the Hand, Tab. L. Fig. 5.

PELVIS, and INFERIOR EXTREMITY.

- A, The gluteus maximus.
 B, Part of the gluteus medius.
 C, The edge of the tensor vaginæ femoris.

FIG. 2.

- A, The under part of the gluteus maximus.
 B, The vastus externus.
a, The adductor magnus femoris.
 C, C, The gracilis.
b, Part of the sartorius.
 D, The long head of the biceps flexor cruris;
 E, Its short head.
c, The insertion of the biceps into the fibula.
 F, The femitendinosus.
 G, H, The femimembranosus.
 I, The edge of the vastus internus.
d, Part of the plantaris.
 K, K, The gastrocnemius externus.
 L, L, L, The edge of the gastrocnemius internus.
 M, The tendo Achillis.
e, The tendon of the plantaris.
 N, The peroneus longus.
 O, The peroneus brevis.
 P, The flexor longus pollicis pedis.
 Q, The tendon of the peroneus brevis.
f, The tendon of the peroneus longus passing into the sole.
 R, The tendon of the extensor longus digitorum.
g, The tendon of the peroneus tertius.
 S, The abductor minimi digiti.
h, The ligament common to the long and short peronei muscles.
i, The ligament proper to each of these two muscles.
k, The ligamentum tarfi annulare.

TABLE

Fig 1

TAB. 40.

Fig 2





Fig. 1

TAB. 41.

Fig. 2



T A B L E X L I.

A VIEW of the SECOND LAYER of MUSCLES on the Posterior Part of the BODY.

FIG. 1.

HEAD.

- A, The temporalis exposed, by removing its tendinous aponeurosis.
a, The tendon of the temporalis, passing under the zygoma.
b, The pterygoideus internus.
 B, The masseter.
c, The mylo-hyoideus.

NECK.

- A, *d*, The levator scapulæ.
 C, The splenius capitis et colli.
 D, The upper end of the complexus.

TRUNK.

- A, *e*, The rhomboides major.
 B, The rhomboides minor.
 C, The serratus posticus superior of the left side.
 D, *f*, The serratus posticus inferior.—*f*, The part from which the latissimus dorsi was cut.
 E, The under part of the serratus magnus.
 F, Part of the sacro-lumbalis.
 G, Part of the longissimus dorsi.
 H, Part of the spinalis dorsi.
 I, I, The broad tendon common to the latissimus dorsi and serratus posticus inferior.
 K, The back part of the obliquus internus abdominis.
 L, L, The intercostales externi.
 M, The coccygeus.
 N, The levator ani.
 O, The sphincter ani.

SUPERIOR EXTREMITY.

- B, The supra-spinatus.
 C, The infra-spinatus.
 D, The teres minor.
 E, The teres major.
 F, The triceps extensor cubiti;
g, Its long head;
h, Its short head.
 G, G, Part of the third head, named *Brachialis Externus*.
i, The common tendon of the triceps, inserted into the olecranon.
 H, Part of the brachialis internus.
 I, The anconeus.
 K, The extensor carpi radialis longior.

- L, The extensor carpi radialis brevior.
 M, The supinator radii brevis.
 N, The extensor ossis metacarpi pollicis.
 O, The extensor primi internodii pollicis.
 P, The extensor secundi internodii pollicis.
 Q, The indicator.
 R, The flexor profundus.
 S, The flexor carpi ulnaris.
 T, A small share of the flexor sublimis.
 See also Tab. L. Fig. 6.

PELVIS, and INFERIOR EXTREMITY.

- A, The gluteus medius.
 B, The pyriformis.
 C, The gemini.
 D, The tendon of the obturator internus, passing between the gemini.
 E, The quadratus femoris.
 F, The vastus externus.
 G, G, The adductor magnus femoris.
 H, The semitendinosus.
 I, The gracilis.

FIG. 2.

- a, a*, The continuation of the adductor magnus femoris, and of,
 A, A, The vastus externus.
 B, The biceps flexor cruris;
b, Its long head;
c, c, Its short head.
d, The common tendon of the two heads.
 C, A small portion of the vastus internus.
 D, Continuation of the gracilis, and of,
 E, The semitendinosus.
 F, F, The semimembranosus.
e, e, The cut heads of the gastrocnemius externus.
 G, The popliteus.
 H, H, The soleus.
 I, *f*, The plantaris;—*f*, Its tendon.
 K, The cut tendon of the gastrocnemius externus.
 L, The tendo Achillis, with that of the plantaris adhering to it, fixed to the os calcis.
 M, The peroneus longus passing to the sole.
 N, The peroneus brevis.
 O, The tendons of the extensor digitorum longus.
 P, The tendon of the peroneus brevis.
g, The tendon of the peroneus tertius.
h, The extensor brevis digitorum.
 Q, The flexor brevis digitorum.

TABLE.

T A B L E XLII.

A VIEW of the Third LAYER of MUSCLES on the Posterior Part of the BODY.

FIG. 1.

HEAD and NECK.

- A, The back part of the buccinator.
a, The pterygoideus internus.
b, The mylo-hyoideus.
 B, B, *c*, *d*, The complexus.—*d*, A fleshy slip from the spinous process of the first dorsal vertebra.
 C, The trachelo-mastoideus.
 D, The scalenus medius.
 E, The scalenus posticus.
 F, The semi-spinalis colli.
 G, G, The interspinales colli.
 H, The obliquus capitis superior.
 I, I, The transversalis colli.
 K, The upper end of the longissimus dorsi, joining the trachelo-mastoideus and cervicalis descendens.
 L, The fleshy slip from the sacro-lumbalis, called *Cervicalis Descendens*.

TRUNK.

- A, E, E, The spinalis dorsi.—Between the spinous processes of the dorsal and lumbar vertebræ, the interspinales muscles appear.
a, *b*, Part of the semi-spinalis dorsi.
 B, The longissimus dorsi.
 C, The tendons of the sacro-lumbalis.
c, A tendon covering, and partly giving origin to the common head of the longissimus dorsi and sacro-lumbalis.
 D, Part of this tendon running across the longissimus dorsi.
 F, The transversalis abdominis.
 G, The inner layer of the aponeurosis common to the ferratus posticus inferior, and obliquus internus abdominis.
 H, H, H, The intercostales externi.
 I, I, Portions of the intercostales externi, called by ALBINUS *Levatores Costarum*.

SUPERIOR EXTREMITY.

- A, The teres major.
a, Part of the subscapularis.

- B, Part of the coraco-brachialis.
 C, Part of the brachialis internus.
 D, The brachialis externus, or third head of the triceps extensor cubiti.
 E, The extensor carpi radialis longior.
 F, The extensor carpi radialis brevior.
 G, The flexor profundus.
 H, The supinator radii brevis.
 I, Part of the flexor longus pollicis.
 K, The pronator radii quadratus.

See also Tab. L. Fig. 7.

PELVIS, and INFERIOR EXTREMITY.

- A, The gluteus minimus.
 B, The obturator internus.
 C, The tendon of the obturator externus.
a, The insertion of the iliacus internus and psoas magnus.
 D, The upper end of the gracilis.
 E, The femimembranosus.
 F, F, The adductor magnus femoris.


FIG. 2.

- A, A, A continuation of the gracilis,
 B, B, ————— of the adductor magnus femoris,
 and,
 C, ————— of the femimembranosus.
 D, The short head of the biceps flexor cruris.—The latter is placed over the part from which the long head was cut,
 E, E, The cut heads of the gastrocnemius externus.
 F, The origin of the plantaris.
 G, The popliteus.
 H, The tibialis posticus.
 I, The flexor longus digitorum pedis.
 K, The flexor longus pollicis pedis.
 L, The peroneus longus.
a, The tendon of the tibialis posticus.
b, The tendon of the peroneus longus passing to the sole.
 M, The peroneus brevis.
c, The tendon of the peroneus brevis.
 N, The extensor brevis digitorum pedis.
 O, Part of the flexor longus digitorum pedis.

TABLE

Fig. 1

TAB. 42. Fig. 2






Fig. 1

TAB. 43 Fig. 2



T A B L E XLIII.

A VIEW of the Fourth LAYER of MUSCLES on the Posterior Part of the BODY.

FIG. 1.

HEAD and NECK.

- A, The rectus capitis posterior minor.
 B, _____ major.
 C, The obliquus capitis superior.
 D, _____ inferior.
 E, The scalenus medius.
 F, The upper end of the multifidus spinæ.
 G, G, The inter-spinales colli.
 H, H, The inter-transversales colli posteriores.
 I, I, I, The semi-spinalis colli.

TRUNK.

- A, A, The semi-spinalis dorfi.
 B, B, The multifidus spinæ.—On the left side of the neck and trunk, the semi-spinalis colli and semi-spinalis dorfi are raised, by which a full view of this muscle is obtained.
 C, C, &c. The levatores costarum breviores.
 D, D, The levatores costarum longiores.
 E, E, &c. The intercostales externi.
 a, a, a, The intercostales interni of the left side.
 b, b, The pleura.

- c, c, c, The inter-transversales dorfi.
 d, d, The inter-spinales dorfi.
 F, The quadratus lumborum.
 G, G, The inter-transversales lumborum.
 H, H, The inter-spinales lumborum.

SUPERIOR EXTREMITY.

- A, The subscapularis.
 B, The supinator radii brevis.
 C, The pronator radii quadratus.

INFERIOR EXTREMITY.

- A, The iliacus internus.
 a, The psoas magnus.
 B, The obturator externus.
 C, The tendon of the iliacus internus and psoas magnus.
 D, D, The adductor magnus femoris.

FIG. 2.

- A, A, A, The continuation of the adductor magnus femoris.—The shaded part in the middle of the muscle represents the impression made by the femi-membranosus.
 B, The tibialis posticus.
 C, The peroneus brevis.

T A B L E XLIV.

A VIEW of the First LAYER of MUSCLES on the Right, and Second LAYER of MUSCLES on the Left Side of the Posterior Part of the BODY, and of the MUSCLES of the External Parts of GENERATION.

FIG. 1.

On the Left Side of the HEAD, and Right Side of the Posterior Part of the TRUNK and EXTREMITIES, the MUSCLES immediately under the Common INTEGUMENTS are shewn; on the Left Side of the Posterior Part of the TRUNK and EXTREMITIES, are seen the MUSCLES which come in view when the Exterior SET has been removed.

All the Muscles represented in this Figure are explained in Tab. XL. and XLI. excepting those on the under parts of the Fore-Arm, and on the Hand.

RIGHT FORE-ARM and HAND.

- H, The supinator radii longus.
 I, The extensor carpi radialis longior.
 K, ————— brevior.
 L, Part of the anconeus.
 M, Part of the flexor profundus which comes from the ulna.
 N, Part of the palmaris longus.
 O, Part of the flexor sublimis.
 P, The flexor carpi ulnaris.
 Q, The extensor carpi ulnaris.
 R, The tendons of the extensor digitorum communis, belonging to the little finger.
 S, The extensor digitorum communis;
 T, Its tendons, going to the other fingers;
 U, U, Their joinings by cross tendons.
 V, The tendon of the indicator, going to join with the tendon of the extensor.
 W, W, The tendons of the extensor communis, joined with those of the lumbricales and interossei;
 X, X, The extremities of these tendons, joined to the bones of the second phalanx.
 Y, The abductor minimi digiti.
 Z, Z, Z, The interossei externi.

- a, a, The tendons of the anterior interossei, joining with the lumbricales.
 b, b, b, The tendons of the posterior interossei.
 c, The abductor indicis.
 d, The tendon of the extensor secundi internodii pollicis.
 e, The annular ligament of the wrist.
 f, The ligament of the extensor carpi ulnaris.
 g, A ligament for the extensores offis metacarpi, et primi internodii pollicis.
 h, The extensor primi internodii pollicis.
 i, The extensor offis metacarpi pollicis.
 k, The tendons of the three extensors of the thumb.
 l, The adductor pollicis.
 m, m, The tendons of the interossei and lumbricales, after joining with the tendons of the extensor digitorum communis, and receiving additions from it, fixed to the third phalanx.

LEFT FORE-ARM and HAND.

- L, L, The extensor carpi radialis longior.
 M, M, ————— brevior.
 N, The anconeus.
 O, The supinator radii brevis.
 P, The extensor offis metacarpi pollicis.
 Q, ————— primi internodii pollicis.
 R, ————— secundi internodii pollicis.
 S, The conjoined tendons of the extensors, fixed to the last bone of the thumb.
 T, The indicator.
 U, The flexor profundus perforans.
 V, The flexor carpi ulnaris.
 W, W, The tendons of the extensor communis, cut off where they are about to join with those of the lumbricales and interossei.
 X, X, The tendons of the extensor communis, fixed to the second phalanx.
 Y, The abductor minimi digiti.
 Z, Z, Z, The tendons of the anterior interossei, joining with those of the lumbricales.

a, The




FIG. 1.



TAB. 44.

FIG. 2.






FIG. 3.



- a*, The tendon of the first lumbricalis.
- b, b, b*, The tendons of the posterior interossei.
- c, c, c*, The interossei.
- d*, The prior indicis.
- e*, The abductor indicis.
- f*, The adductor pollicis.
- g, g*, The tendons of the interossei and lumbricales, after joining with the tendons of the extensor communis, fixed to the third phalanx.

FIG. 2.

MUSCLES about the Root of the PENIS and Under End of the INTESTINUM RECTUM of a CHILD.

These muscles are described in Tab. XLVIII. Fig. 9.

FIG. 3.

MUSCLES of the External Parts of GENERATION, &c. in the FEMALE.

- a, a*, A section of the thighs.
- b*, The clitoris.
- c, c*, The crura clitoridis.
- d, d*, The erectores clitoridis.
- e, e*, The sphincter vaginae.
- f*, The sphincter ani, connected with the sphincter vaginae.

See also the Muscles in a Lateral View of the Female: Parts of Generation, Vol. II.

FIG. 1. TAB. 45.




FIG. 2.



FIG. 3.






FIG. 6.




FIG. 5.




FIG. 8.




FIG. 9.




FIG. 7.





FIG. 4.






FIG. 1. TAB. 46:



FIG. 2.



FIG. 18.



FIG. 3.




FIG. 16.



FIG. 17.




FIG. 4.




FIG. 5.




FIG. 6.




FIG. 7.



FIG. 15.




FIG. 8.




FIG. 9.



FIG. 10.



FIG. 11.




FIG. 12.





FIG. 13.



T A B L E XLVI.

The MUSCLES seated about the THROAT ; with a VIEW of the First LAYER of MUSCLES upon the Lateral Parts of the Body.

FIG. 1.—17.

Explained in Tab. XLVII.

FIG. 18.

A View of the First LAYER of MUSCLES on the Lateral Parts of the BODY.

HEAD and TRUNK.

- A, The occipito-frontalis ;
 B, The aponeurosis joining the two sides of this muscle.
 C, The attollens aurem.
 D, The anterior auris.
 E, The retrahentes aurem.
 a, The helix major.
 b, ——— minor.
 c, The tragicus.
 d, The anti-tragicus.
 F, The orbicularis palpebrarum.
 G, The zygomaticus major.
 H, The buccinator.
 I, The masseter.
 K, The depressor anguli oris.
 L, The pterygoideus internus.
 M, The platysma myoides.
 N, The sterno-cleido-mastoideus.
 O, The complexus.
 P, The splenius.
 Q, The scalenus medius.
 R, The levator scapulæ.
 S, S, T, The trapezius.
 U, The teres minor.
 W, ——— major.
 X, X, Y, The latissimus dorsi.
 Z, The pectoralis minor.
 a, a, b, The pectoralis major.
 c, c, c, The serratus magnus.
 d, d, e, e, The obliquus externus abdominis ;—d, d, The fleshy ; e, e, The tendinous parts.
 f, The cremaster testis.

LEFT SUPERIOR EXTREMITY.

- A, The deltoïdes.
 B, The biceps flexor cubiti.
 C, The brachialis internus.
 D, The triceps extensor cubiti.
 E, The flexor carpi ulnaris.
 F, The supinator radii longus.
 G, The flexor carpi radialis.
 H, H, The extensor carpi radialis longior.
 I, ——— brevior.
 K, The extensor carpi ulnaris.
 L, The extensor digitorum communis ;
 M, Its tendon.
 N, The extensor ossis metacarpi pollicis ;
 O, Its tendon.
 P, The extensor primi internodii pollicis ;
 Q, Its tendon.
 R, The tendo secundi internodii.
 S, The ligamentum carpi annulare posterius.
 T, The ligament confining the tendons of the extensor ossis metacarpi, and extensor primi internodii pollicis.
 W, The adductor pollicis.
 X, The opponens pollicis.

RIGHT SUPERIOR EXTREMITY.

- A, B, The triceps extensor cubiti ;—A, The part called *Extensor Longus* ;—B, The part called *Extensor Brevis*.
 C, The brachialis internus.
 D, The biceps flexor cubiti.
 E, The supinator longus.
 F, The pronator teres.
 G, The flexor carpi radialis.
 H, The palmaris longus.
 I, The flexor sublimis perforatus.
 K, The flexor carpi ulnaris.
 L, The extensor carpi ulnaris.
 M, The flexor brevis pollicis.
 N, The tendon of the flexor longus, with its retaining ligaments.

O, The

- O, The palmaris brevis, and, on the outside of it, the abductor minimi digiti.
 P, The tendons of the extensor digitorum communis;
 Q, The aponeuroses of these tendons, stretched over the back of the four fingers.


LEFT INFERIOR EXTREMITY.

- A, The adductor longus femoris.
 B, The pectinalis and psoas magnus.
 C, The sartorius.
 D, The tensor vaginæ femoris.
 E, The gluteus medius.
 F, ————— maximus.
 G, The semi-tendinosus.
 H, The biceps flexor cruris.
 I, The vastus externus.
 K, The rectus.
 L, The vastus internus.
 M, The ligament connecting the patella to the tibia.
 N, The outer head of the gastrocnemius externus.
 O, O, The gastrocnemius internus.
 P, The tendo Achillis.
 Q, The peroneus longus.
 R, The peroneus brevis.
 S, Ligaments binding the tendons of the peronei.
 T, The extensor longus digitorum, inseparably connected with,
 U, The peroneus tertius.
 V, The tendon of the peroneus tertius, inserted into the metatarsal bone of the little toe.
 W, The tendon of the extensor longus, splitting into four smaller tendons;
 X, X, Their insertions into the toes.
 Y, The extensor proprius pollicis;
 Z, Its tendon.
 a, a, The tibialis anticus.
 b, The upper and under portions of the ligamentum tarfi annulare.

- c, The extensor brevis digitorum pedis; its tendons are inserted into all the toes, excepting the smallest.
 d, d, Part of the interossei pedis externi.
 e, The abductor minimi digiti pedis;
 f, Its tendon.
 g, g, The flexor brevis minimi digiti pedis.

RIGHT INFERIOR EXTREMITY.

- A, The rectus;
 B, Its insertion into the patella.
 C, The ligament which fixes the patella to the tibia.
 D, The vastus internus.
 E, The sartorius;
 F, Its tendon, fixed to the tibia.
 G, The gracilis.
 H, The semi-membranosus.
 I, I, The semi-tendinosus.
 K, The gastrocnemius externus;
 L, Its tendon.
 M, The gastrocnemius internus.
 N, The tendo Achillis.
 O, The tendon of the plantaris.
 P, The flexor proprius pollicis pedis.
 Q, The ligament binding the tendon of the flexor longus.
 R, The flexor longus digitorum pedis.
 S, The tendon of the tibialis posticus.
 T, The ligament covering the tendon of the flexor longus digitorum pedis, and tibialis posticus.
 U, The ligament which retains the tibialis posticus.
 V, The tibialis anticus;
 W, Its tendon.
 X, X, The upper and under portions of the ligamentum tarfi annulare.
 Y, The tendon of the extensor proprius pollicis pedis;
 Z, An aponeurosis joining this tendon.
 a, The abductor pollicis pedis.
 b, The flexor digitorum accessorius.



TAB. 47.

Fig. 2.

Fig. 1.

Fig. 3.






Fig. 6.

Fig. 5.

Fig. 4.






Fig. 7.

Fig. 8.

Fig. 9.

Fig. 10.

Fig. 11.







Fig. 12.


Fig. 13.

Fig. 14.

Fig. 15.

Fig. 16.

Fig. 17.



T A B L E XLVII.

MUSCLES seated about the THROAT.

FIG. 1.

Presents a Lateral View of the MUSCLES seated under the HEAD, and before the VERTEBRÆ of the NECK.

- a*, The pterygoideus externus.
- b*, ————— internus.
- c*, The mylo-hyoideus.
- d*, The stylo-hyoideus.
- e, f*, The digastricus.
- g, h*, The hyo-glossus.
- i*, The os hyoides.
- k*, The thyro-hyoideus.
- l*, The thyroid cartilage.
- m*, The crico-thyroideus.
- n*, The cricoid cartilage.
- o*, A section of the œsophagus.
- p, p*, The constrictor pharyngis inferior.
- q*, ————— medius.
- r*, ————— superior.

FIG. 2.

Represents the MUSCLES under those shewn in the preceding Figure, which, together with the Right Side of the LOWER JAW, are here removed.

- a*, The upper jaw.
- b*, A section of the lower jaw.
- c*, The tongue.
- d*, The stylo-glossus.
- e*, The hyo-glossus.
- f*, The genio-glossus.
- g*, The stylo-pharyngeus.
- h, h*, The constrictor pharyngis superior.
- i*, ————— medius.
- k, k*, ————— inferior.
- l*, The thyroid cartilage.
- m*, The cricoid cartilage.
- n*, A section of the œsophagus.

FIG. 3.

Represents the next Order of MUSCLES, the Outermost of those in the preceding Figure being removed.

- a*, The under half of the stylo-glossus, the upper half being removed.
- b, c*, The genio-glossus.
- d*, The constrictor pharyngis superior.
- e*, ————— medius.

- f*, The constrictor pharyngis inferior.
- g*, The os hyoides.
- h*, The thyroid cartilage.
- i*, The cricoid cartilage.
- k*, A section of the œsophagus.

FIG. 4.

Represents the next Order of MUSCLES, after the Outermost of Fig. 3. are removed.

- a*, The circumflexus palati, immediately behind which is the levator palati.
- b*, The stylo-pharyngeus.
- c, c*, The palato-pharyngeus, covering a part of the membrane of the pharynx.
- d*, The constrictor isthmi faucium.
- e*, The tonsil.
- f*, The stylo-glossus, where it joins the tongue.
- g*, A section of the hyo-glossus.
- h*, The lingualis.
- i, k*, The genio-hyo-glossus;—*i*, Its origin from the lower jaw.
- l*, The os hyoides.
- m*, The ligament which joins the cornu of the os hyoides and thyroid cartilage.
- n*, The body of the thyroid cartilage.
- o*, The cricoid cartilage.
- p*, The ligament by which the thyroid and cricoid cartilages are joined together.
- q*, A section of the œsophagus.

FIG. 5.

In this Figure, some of the Outer MUSCLES shewn in the Fourth are removed, the PHARYNX is laid open longitudinally, and the Right Part of it cut off, to shew its Cavity, with the Root of the TONGUE and EPIGLOTTIS.

- a*, The circumflexus palati.
- b*, The levator palati.
- c*, The tonsil.
- d*, The constrictor isthmi faucium.
- e*, The tongue.
- f*, The under part of the lingualis.
- g*, The genio-hyo-glossus.
- h*, The epiglottis.
- i*, The os hyoides.
- k*, The thyroid cartilage.
- l*, The cricoid cartilage.
- m*, The pharynx laid open.
- n*, The upper part of the œsophagus.

FIG.

FIG. 6.

Presents a Posterior View of the PHARYNX, and the Under Part of the BONES of the HEAD, to which the PHARYNX is connected.

- a*, The upper point of the constrictor pharyngis inferior on each side.
- b*, The under end of the pharynx;—the letter points also at the inner transverse fibres of the œsophagus, which are laid bare.
- c, c*, The outer fibres of the œsophagus descending obliquely backwards on each side.
- d*, A section of the œsophagus.
- e, e*, A section of the trachea.
- f, f*, The ends of the cornua of the os hyoides.
- g, g*, The ligaments which join the upper processes of the thyroid cartilage to the ends of the cornua of the os hyoides.
- h, h*, The constrictor medius pharyngis, on each side.
- i, i*, The constrictor superior pharyngis, on each side.
- k, k*, The naked membrane of the pharynx.
- l, l*, The stylo-pharyngeus, on each side.
- m, m*, The styloid processes of the temporal bones.
- n, n*, The pterygoid processes of the sphenoid bone.
- o, o*, The backmost tooth of the upper and under jaws, on each side.

FIG. 7.

Presents the next View, after the removal of the Lower CON- STRICTORS of the PHARYNX. The BONES of the HEAD are not added; but the STYLOID PROCESSES are left, to shew the ORIGIN of the STYLO-PHARNGEI.

- a, b, k*, The constrictor pharyngis medius.
- c*, The upper constrictor of the pharynx, cut off from the buccinator.
- d*, The naked membrane of the pharynx.
- e*, The styloid process of the temporal bone, cut off at its root.
- f*, The stylo-pharyngeus, arising, tendinous, from the styloid process.
- g*, The common end of the stylo-pharyngeus and palato-pharyngeus.
- h*, Part of the stylo-pharyngeus and palato-pharyngeus, fixed to the edge of the thyroid cartilage.
- i*, The naked membrane of the lower part of the pharynx, continued to the œsophagus.
- k*, The cornu of the os hyoides.
- l*, The superior cornu of the thyroid cartilage.
- m*, The posterior edge of the thyroid cartilage;
- n*, Its inferior cornu.
- o*, The tubercle on the outer side, at the root of the superior cornu.
- p*, The cricoid cartilage.
- q*, A section of the trachea.

FIG. 8.

The next View of the MUSCLES, after the Middle CON- STRICTORS of the PHARYNX are removed.

- a, c, b, d*, The constrictores pharyngis superiores.
- e*, The levator palati.
- f*, The circumflexus palati.
- g*, The tendinous origin of the stylo-pharyngeus, where it is cut off from the styloid process.
- h*, That part of the stylo-pharyngeus which forms two fasciculi, passing separately under the fibres of the upper constrictor.
- i*, The under and larger part of the stylo-pharyngeus.
- k*, Part of the common end of the stylo-pharyngeus and palato-pharyngeus, fixed to the thyroid cartilage.
- l*, Part of the common end of the stylo-pharyngeus, joined to its fellow on the back of the pharynx.

FIG. 9.

Represents the next Order of MUSCLES, after the Upper CONSTRUCTORS of the PHARYNX are removed.

- a*, The naked membrane of the pharynx.
- b*, The small hook of the pterygoid process.
- c*, The palato-pharyngeus.
- d, e*, Part of the common end of the stylo-pharyngeus and palato-pharyngeus.

FIG. 10.

Represents the Inner and Fore Part of the PHARYNX, the whole Posterior Part being removed.

- a*, The Eustachian tube;—its orifice opening laterally into the posterior foramen of the nostril.
- b*, The septum narium.
- c*, The cavity of the nostril, with the lower os spongiosum, covered with the mucous membrane.
- d, d*, The palatum molle.
- e*, The uvula.
- f*, The posterior arch, which descends laterally from the soft palate through the side of the pharynx.
- g*, The tonsil.
- h*, The tongue.
- i*, The epiglottis.
- k*, The membranous side of the glottis.
- l*, The rima, or slit of the glottis.
- m*, The back part of the tube of the larynx, projecting within the pharynx.

FIG. 11.

Represents the MUSCLES lying immediately under the MEM- BRANE of the PHARYNX, which, with the ŒSOPHAGUS and TRACHEA, are removed.

- a*, The levator palati.
- b*, The azygos uvulae.

c, The

- c*, The palato-pharyngeus ;
- d*, The part which afterwards passes under the levator palati.
- e*, Part of the palato-pharyngeus, called by ALBINUS, *Salpingo Pharyngeus*.
- f*, Part of the common end of the palato-pharyngeus and stylo-pharyngeus.
- g*, The posterior edge of the velum palati.
- h*, The uvula.
- i*, The tonsil, projecting before the palato-pharyngeus.
- k*, The tongue.
- l*, The epiglottis.
- m*, The point of the arytenoid cartilage.
- n*, The arytenoideus obliquus.
- o, o*, ————— transversus.
- p*, The crico-arytenoideus posticus.
- q*, The cricoid cartilage.

FIG. 12.

Represents the MUSCLES deeper seated than those shewn in the former Figure.

- a*, The Eustachian tube opening laterally into the posterior foramen of the nostril.
- b*, The os spongiosum inferius, covered with the mucous membrane.
- c*, The levator palati.
- d*, The circumflexus.
- e*, The small hook of the pterygoid process.
- f*, Part of the palato-pharyngeus, which passes through the soft palate, under the end of the levator.
- g*, Part of the common end of the stylo-pharyngeus and palato-pharyngeus, produced more particularly from the stylo-pharyngeus.
- b, b*, The arytenoid cartilage.

FIG. 13.

Represents the MUSCLES which appear upon the Removal of the LEVATORES PALATI, the ANNULAR and ARYTE-NOID CARTILAGES, and their Appendages.

- a*, The circumflexus palati.
- b*, The aponeurosis of the circumflexi.
- c*, The hook-like process of the pterygoid plate.
- d*, The palato-pharyngeus.
- e*, Part of the stylo-pharyngeus inserted into the thyroid cartilage.
- f*, The thyroid cartilage.
- g*, A prominence upon the inner side of the thyroid cartilage.
- h*, The under end of the epiglottis, fixed to the thyroid cartilage.

FIG. 14.

MUSCLES of the PALATE, viewed on the Under Side.

- a*, The levator palati.
- b, c*, The circumflexus palati;—*c*, Its tendon, passing over the hook-like process of the pterygoid plate.
- d*, The membrane of the palate.
- e*, The Eustachian tube.
- f, f, f*, The circumference, from which the membrane of the palate is cut off.

FIG. 15.

The MOUTH and FAUCES open, to shew the MUSCLES of the PALATUM MOLLE, on the Under and Fore Side; the investing MEMBRANE being removed.

- a*, The posterior arch, and,
- b*, The anterior arch of the palate.—Between this and the posterior arch *a*, is the seat of the amygdala.
- c*, The edge of the soft palate.
- d*, The uvula.
- e*, The tongue.
- f, f*, The fauces.
- g*, The constrictor isthmi faucium.
- h*, The palato-pharyngeus.

FIG. 16.

Shews the LARYNX, with its Posterior MUSCLES, and those at the Side of the THYROID CARTILAGE, the Right Part of which is removed.

- a*, The crico-arytenoideus posticus.
- b*, ————— lateralis.
- c, e, f*, The thyro-epiglottideus.
- d, g*, The thyro-arytenoideus.
- h*, The arytenoideus transversus.
- i*, ————— obliquus, with its continuation to the epiglottis.

FIG. 17.

Represents the same VIEW of the CARTILAGES of the LARYNX with the preceding Figure, but wholly freed from the MUSCLES and MEMBRANES.

- a, b, c*, The inside of the left half of the thyroid cartilage.
- d*, The superior cornu of the thyroid cartilage.
- f*, The cricoid cartilage.
- g*, The right arytenoid cartilage.
- h*, The left arytenoid cartilage.
- i, k*, The epiglottis;—*k*, Its concave part.

T A B L E XLVIII.

VIEWS of various MUSCLES, some of which are not sufficiently shewn in former FIGURES.

FIG. 1.

The Sterno-Costalis.

See Tab. XXXIX. Right-side of the Trunk.

FIG. 2.

A View of the CAVITY of the DIAPHRAGM, or of the Side next the ABDOMEN.

- A, The cartilago enfiformis.
- B, The cartilage of the seventh rib.
- C, The point of the twelfth rib.
- D, E, F, The first, second, and third lumbar vertebræ.
- G, A section of the aorta.
- H, The mouth of the cœliac artery.
- I, The superior mesenteric artery, and, on each side of it, the renal arteries.
- K, A section of the vena cava.
- L, A section of the œsophagus.
- M, The psoas magnus.
- N, The quadratus lumborum.
- O, The great intercostal nerve.
- P, The last of the dorsal nerves.
- Q, Q, The interior arch, or fleshy boundary of the diaphragm, to which the peritoneum adheres.
- R, S, S, The cordiform tendon of the diaphragm.
- T, T, The fleshy parts of the diaphragm which come from the ribs.
- U, U, The fleshy pillars of the diaphragm.
- V, V, The tendinous crura, or long heads of the diaphragm.
- W, W, Fleshy columns on each side, where the œsophagus passes.

FIG. 3.

The Fore-Part of the COCCYGEUS.

- a, The tendinous origin of the coccygeus, from the spinous process of the os ischium.
- b, Insertion of part of the coccygeus into the os sacrum.
- c, d, Insertion of the greater part of the coccygeus into the os coccygis.

FIG. 4.

A Posterior View of the LEVATORES, and SPHINCTER INTERNUS ANI.

- a, The anterior portion of the levator ani, viewed on its inner side within the pelvis;
- b, Its tendinous origin, from the inner side of the os pubis;
- c, ————— from the spinous process of the os ischium;
- d, The posterior and outer part of the muscle;
- e, Its insertion into the under part of the os coccygis;
- f, The tendinous end meeting its fellow below the coccyx.
- g, The muscular coat of the rectum, called by ALBINUS, *Sphincter Internus Ani.*
- h, The anus.

FIG. 5.

Anterior View of the LEVATORES, with the SPHINCTER INTERNUS ANI.

- c, The bulb of the urethra.
- d, A section of the urethra and corpus spongiosum.
- e, The sphincter internus ani.
- f, f, f, f, The place from whence a portion of the os pubis is cut out, to obtain a view of the levatores which lie behind.
- g, The levator ani, arising from the inner side of the os pubis.
- h, The thin portion which comes out from the angle where the crus penis joins the corpus spongiosum urethræ.

FIG. 6.

The ACCELERATORES URINÆ, TRANSVERSI PERINEI, and ERECTORES PENIS.

- a, The accelerator urinæ, investing the bulb of the urethra.
- b, The transversus perinei.
- c, ————— alter.
- d, e, The erector penis.
- f, The corpus cavernosum penis.
- g, ————— urethræ.

FIG. 7.

The Back Part of the SPHINCTER EXTERNUS ANI.

- a, The point by which it adheres to the extremity of the os coccygis.
- b, The anus, from which to a the fibres meet in angles which point upwards, and become more acute as they ascend.

FIG. 8.

The Fore-Part of the SPHINCTER EXTERNUS ANI.

- a, The fibres which meet from each side, forming angles pointing upwards, and becoming more acute as they ascend.
- b, The termination of the sphincter ani in the perineum.
- c, The corpus spongiosum urethræ.
- d, The bulb of the urethra.

FIG. 9.

Muscles about the Root of the Penis, and Under End of the Intestinum Rectum in a Child.

- a, a, The sphincter ani.
- b, The levator ani.
- c, The transversalis perinei.
- d, The erector penis.
- e, The accelerator urinæ.
- f, The corpus cavernosum penis.
- g, The corpus spongiosum urethræ.
- h, The scrotum turned up.
- i, Part of the thigh.
- k, The cut edge of the integuments.

FIG. 1.



TAB. 49.

FIG. 2.



FIG. 3.




FIG. 6.



FIG. 5.



FIG. 4.



FIG. 7.



FIG. 8.

FIG. 9.







FIG. 10.



T A B L E X L I X .

Represents the SALIVARY GLANDS, Parts about the THROAT, and certain deep-seated MUSCLES, in the Interior Part of the BODY, not sufficiently shewn in former FIGURES.

FIG. 1.

Part of the MUSCLES of the OS HYOIDES, together with the SUBMAXILLARY GLAND.

- a*, Part of the masseter.
- b*, The posterior head of the digastricus ;
- c*, Its anterior head.
- d*, The stylo-hyoideus, through which the tendon of the digastricus passes.
- e, e*, The sterno-hyoidei.
- f*, The omohyoideus.
- g*, The pharynx.
- h*, The submaxillary gland.

FIG. 2.

MUSCLES deeper seated than the former, and the SUBMAXILLARY GLAND raised.

- a, a*, The mylo-hyoidei.
- b*, The hyo-glossus.
- c*, The sterno-thyroideus.
- d*, The thyro-hyoideus.
- e*, The submaxillary gland raised from its place, behind the angle of the lower jaw.
- f*, The stylo-glossus.
- g*, The stylo-pharyngeus.
- h*, The pharynx.

FIG. 3.

MUSCLES deeper seated than the former.

- a*, The genio-hyoideus.
- b*, The genio-hyo-glossus.
- c*, The stylo-glossus.
- d*, The stylo-pharyngeus.
- e*, The submaxillary gland raised, by which its duct is seen in its passage under the tongue, to its termination at the side of the frænum linguæ.
- f*, The sublingual gland.
- g*, The os hyoides.
- h*, The thyroid cartilage.
- i*, The cricoid cartilage, with the crico-thyroidei.
- k*, The thyroid gland.
- l*, The trachea.
- m*, The pharynx.

FIG. 4.

The Inferior Surface of the TONGUE, with its MUSCLES dissected.

- a, a*, The genio-hyo-glossus ;
- b*, Its origin, cut from the inner part of the lower jaw.
- c, c, c*, The hyo-glossus.
- d*, The stylo-glossus.
- e*, The tip of the tongue pinned out, at each side of which the papillæ appear.
- f, f*, The basis, or root of the tongue.
- g*, The membrane, with its mucous glands, continued from the tongue to the epiglottis.

FIG. 5.

Shews the TONGUE, OS HYOIDES, and LARYNX, separated from the Left Side of the HEAD, and turned over upon the Right,—the HEAD being inverted.

- a*, The inner side of the lower jaw.
- b*, Part of the glandulæ palatinæ.
- c*, The uvula, with its muscle hanging over the openings into the back part of the nose.
- d*, The right side of the pharynx remaining entire.
- e*, The tongue, at the anterior edges of which the papillæ are seen.
- f*, The salivary glands of the tongue.
- g*, One of the amygdalæ.
- h*, The os hyoides, with its left cornu joined to the left superior cornu of the thyroid cartilage.
- i*, The thyroid cartilage.
- k*, The back part of the cricoid cartilage.
- l, l*, The arytenoid cartilages.
- m*, The epiglottis.
- n*, The cartilages of the trachea.
- o*, The membranous part of the trachea.

FIG. 6.

The Back Part of the PHARYNX, and its Connections with the LARYNX.

- A*, The cartilages of the trachea.
- B*, The membranous back part of the trachea.
- C, C*, That part of the pharynx which arises from the pterygoid processes, levatores palati, and os occipitis.

- D, D, Parts of the pharynx which arise from the lower jaw.
 E, E, Fibres of the pharynx, from the root of the tongue;
 F, ————— from the os hyoides;
 G, ————— from the thyroid cartilage;
 and,
 H, ————— from the cricoid cartilage.
 I, The os hyoides.
 K, The thyroid cartilage.
 L, L, The styloid processes.
 M, M, The ligaments from the styloid processes, fixed to the appendices of the os hyoides.
 N, N, The stylo-hyoidei.
 O, O, The stylo-pharyngei.
 P, The back part of the œsophagus;
 Q, Q, Its external surface.
 R, The sterno-thyroideus.
 S, The thyro-hyoideus.
- F, The scalenus anticus.
 G, ————— medius.
 H, The trachelo-mastoideus.
 I, I, &c. The intercostales externi.
 K, K, &c. ————— interni.
 L, L, &c. Portions of the internal intercostals, called *Depressores Proprii Cowperii*.
 M, The transversus abdominis.
 N, The quadratus lumborum.
 O, The psoas parvus.
 P, ————— magnus.
 Q, A portion of the psoas magnus; the upper part which lay over the quadratus lumborum is cut off.
 R, The iliacus internus.
 S, The pyriformis.
 T, The obturator externus.
 U, The adductor brevis femoris.

FIG. 7.

View of the Right and Back Part of the LARYNX.

- a, The cricoid cartilage.
 b, The epiglottis;
 c, Its root cut from the base of the tongue, where many small glands appear.
 d, The tips of the arytenoid cartilage freed from their membrane.
 e, The concave surface of the thyroid cartilage, and its superior cornua: The right half of the cartilage is turned back.
 f, The inferior cornu of the thyroid, cut from
 g, Its connection to the cricoid cartilage.
 h, The crico-arytenoideus posticus.
 i, ————— lateralis.
 k, The thyro-arytenoideus.
 l, The arytenoideus transversus.
 m, The trachea;
 n, Its membranous part.

FIG. 8.

A View principally of the MUSCLES in the Interior Part of the BODY, next the SPINE.

- A, The rectus capitis lateralis.
 B, ————— anterior minor.
 C, ————— major.
 D, E, The longus colli.

FIG. 9.

Explained in Tab. XXXVII.

FIG. 10.

The ABDOMEN opened, and its Contents removed, to shew the DIAPHRAGM and MUSCLES of the LOINS.

- A, A, The containing parts of the abdomen cut and turned back.
 B, B, The cut ends of the ribs.
 C, The origin of the superior, or greater muscle of the diaphragm, from the cartilago ensiformis.
 D, D, Origins from the ribs.
 E, E, E, The cordiform tendon of the diaphragm.
 F, The perforation in that tendon, for the passage of the vena cava inferior.
 G, G, The long crura of the inferior, or lesser muscle of the diaphragm.
 H, The passage of the aorta between the long crura.
 I, I, Shorter heads.
 K, K, The fleshy columns from the joining of these heads.
 L, Fibres crossing under
 M, The passage of the œsophagus:
 N, The quadratus lumborum.
 O, The psoas parvus.
 P, P, The large psoæ, the right of which is turned outwards at its lower end.

TAB. 50

Fig. 1.




Fig. 2.



Fig. 3.



Fig. 4.



Fig. 5.




Fig. 6.



Fig. 7.



Fig. 8.



Fig. 9.



Fig. 10.



Fig. 11.



Fig. 12.



T A B L E L.

The different Orders of MUSCLES on the Under Part of the FORE-ARM and HAND, and on the SOLE of the FOOT.

FIG. 1.

The First Order of MUSCLES on the Under and Anterior Part of the FORE-ARM and PALM.

- a*, The tendon of the supinator longus,
b, ————— flexor carpi radialis,
c, ————— palmaris longus,
d, ————— flexor sublimis, and,
e, ————— flexor carpi ulnaris.
f, The flexor longus pollicis.
g, The ligament under which the extensores ossis metacarpi et primi internodii pollicis pass.
h, The flexor ossis metacarpi pollicis.
i, The abductor pollicis.
k, The tendon of the extensor primi internodii pollicis.
l, Part of the flexor brevis pollicis.
m, The tendon of the flexor longus pollicis bound by ligaments.
n, The aponeurosis palmaris, slightly distinguished into four portions, the extremities of which afterwards become more distinct, and are strengthened by transverse tendinous fibres.
o, The palmaris brevis.
p, The abductor minimi digiti.
q, The adductor metacarpi minimi digiti.
r, The flexor parvus minimi digiti.
s, t, u, Three small annular ligaments, which retain the tendons of the sublimis and profundus in their places on the fore-finger.
v, w, x, The tendon of the flexor sublimis, with the tendon of the flexor profundus passing through it.
w, The insertion of the tendon of the flexor profundus into the third bone of the mid-finger.—The same parts with those marked *s, t, u, v, w, x*, are seen on the other fingers.

FIG. 2.

The Second Order of MUSCLES on the Under and Fore Part of the FORE-ARM and PALM.

- a*, The tendon of the flexor carpi ulnaris.
b, b, A portion of the flexor digitorum sublimis;
c, c, d, The tendons of that muscle passing behind the anterior annular ligament of the wrist.

- e*, The flexor pollicis longus;
f, Its tendon.
g, h, The extensor ossis metacarpi pollicis.
i, ————— primi internodii pollicis.
k, The ligamentum carpi annulare anterius.
l, The flexor ossis metacarpi pollicis.
m, n, The anterior and posterior portions of the flexor brevis pollicis, with the tendon of the flexor longus pollicis between them.
o, The adductor pollicis.
p, The abductor indicis.
q, q, &c. The lumbricales.
r, The tendon of the flexor sublimis, perforated by that of the flexor profundus, and inserted into the second bone of the fore-finger.
s, The tendon of the flexor profundus.
t, The insertion of the flexor profundus into the third bone of the fore-finger. The tendons of the flexor sublimis and profundus are seen on the other fingers also.
u, The flexor parvus minimi digiti.
v, The abductor minimi digiti.

FIG. 3.

The Third Order of MUSCLES on the Under and Fore Part of the FORE-ARM and PALM:

- a*, The flexor longus pollicis;
b, Its tendon.
c, d, e, The flexor profundus.
f, The tendon of the flexor profundus to the fore-finger. A tendon of that muscle is also sent to each of the other fingers.
g, The ligamentum carpi annulare.
h, The adductor metacarpi minimi digiti.
i, k, The flexor brevis pollicis.
l, The adductor pollicis.
m, m, &c. The lumbricales.
n, The prior indicis.

FIG. 4.

The Fourth Order of MUSCLES on the Under and Fore Part of the FORE-ARM and PALM.

- a, a*, The pronator radii quadratus.

b, c, The

b, c, The flexor brevis pollicis, with the ossa sesamoidea into which it is inserted.
d, The adductor pollicis.

Interossei interni et externi.	{	<i>e</i> , The prior indicis.
		<i>f</i> , The posterior indicis.
		<i>g</i> , The prior medii digiti.
		<i>h</i> , The posterior medii digiti.
		<i>i</i> , The prior annularis.
		<i>l</i> , The interosseus auricularis.

FIG. 5.

The First Order of MUSCLES of the Under and Back Part of the FORE-ARM and BACK of the HAND.

a, The extensor ossis metacarpi pollicis.
b, ———— primi internodii pollicis.
c, The tendon of the extensor primi internodii pollicis.
d, ———— secundi internodii pollicis.
e, f, The extensor digitorum communis.
g, ———— carpi ulnaris.
h, The flexor carpi ulnaris.
i, The ligament which confines the two first extensors of the thumb.
k, The ligamentum carpi annulare posterius.
l, ———— extensoris carpi ulnaris.
m, The tendons of the extensor digitorum communis, divided by longitudinal fissures upon the back of the hand.
n, The tendon of the indicator.
o, Aponeurotic slips joining the tendons of the extensor digitorum to each other.
p, Part of the abductor minimi digiti.
q, The adductor pollicis.
r, Tendinous expansions continued from the tendons of the extensor digitorum communis, and of the interossei and lumbricales, adhering closely to the bones.
s, The division of these expansions continued from the tendons of the extensor digitorum communis, and of the interossei and lumbricales,—for the readier motion of the joints;
t, Their termination at the last joint of the fingers.

FIG. 6.

The Second Order of MUSCLES on the Under and Back Part of the FORE-ARM and BACK of the HAND.

a, The extensor ossis metacarpi pollicis.
b, ———— primi internodii pollicis.
c, The tendon common to the two muscles.
d, The extensor secundi internodii pollicis;
e, Its tendon.
f, The tendon of the indicator.
g, The flexor carpi ulnaris.
h, The cut tendon of the extensor digitorum communis. Sections of the tendons of this muscle are also seen upon the ring and little finger.
i, The abductor indicis.

k, The prior medii digiti.
l, The posterior medii digiti.
m, ———— annularis.

FIG. 7.

The Third Order of MUSCLES on the Under and Back Part of the FORE ARM and BACK of the HAND.—One of the Heads of each External INTEROSSEOUS MUSCLE is removed, to obtain a View of the Internal.

a, The tendon of the extensor carpi radialis longior.
b, ———— brevior.
c, The cut tendon of the extensor digitorum communis.
d, The adductor pollicis.
D, The flexor brevis pollicis.
e, The prior indicis.
f, The posterior indicis.
g, Part of the prior medii digiti.
h, Part of the posterior medii digiti.
i, The prior annularis.
k, Part of the posterior annularis.
l, The interosseus auricularis.

FIG. 8.

Shews the APONEUROSIS, and Part of the First Order of MUSCLES and LIGAMENTS of the SOLE, after removing the Common INTEGUMENTS.

a, b, c, The aponeurosis plantaris, connected behind to the os calcis, and before to the first joint of all the toes.—
a, The middle part divided into five slips, which split at the roots of the toes, and embrace the tendons of the flexor muscles.—
b, The portion which covers the abductor minimi digiti.—
c, The portion which covers a part of the abductor pollicis.
d, The abductor pollicis.
e, Part of the flexor brevis pollicis.
f, The common tendinous ends of the abductor, and short flexor of the great toe.
g, Part of the transversalis pedis.
h, The abductor minimi digiti pedis.—A small part of the lumbricales, the abductor minimi digiti, the short flexors, the tendons of the long and short flexors, with the ligaments which confine them, are seen upon the other toes,—nearly as in the hand.

FIG. 9.

The First Order of MUSCLES on the SOLE, after the APONEUROSIS, and most of the LIGAMENTS shewn in the former Figure, are removed.

a, The flexor brevis digitorum, sending tendons to the second phalanx of the four small toes.
b, The tendon of the flexor longus pollicis.
c, The abductor pollicis.

- d, e*, The abductor minimi digiti, composed of two parts, and fixed by a common tendon to the first bone of the little toe.
- f*, The flexor brevis minimi digiti.

FIG. 10.

The Second Order of MUSCLES in the SOLE.

- a*, The tendon of the flexor longus digitorum.
- b, c, d*, The flexor digitorum pedis accessorius;—*b, c*, Its two heads arising from the os calcis;—*d*, Its insertion into the tendon of the flexor longus digitorum pedis.
- e*, The connection between the tendons of the flexor longus digitorum and flexor longus pollicis.
- f, f*, The insertion of the tendons of the flexor longus digitorum into the last bone of the small toes.
- g, h, i, k*, The lumbricales.
- l*, The tendon of the flexor longus pollicis.
- m*, The insertion of the flexor longus pollicis into the last joint of the great toe.
- n*, The insertion of the tibialis posticus.
- o*, ————— anticus.
- p, q*, The two parts of the flexor brevis pollicis.
- r*, The insertion of the peroneus brevis.
- s*, The tendon of the peroneus longus passing to the sole.
- t, u*, The ligaments connecting the bones at this part of the sole, and giving origin to muscles.

FIG. 11.

The Third Order of MUSCLES in the SOLE.

- a*, The insertion of the peroneus brevis.

- b*, The tendon of the peroneus longus.
- c*, The insertion of the tibialis posticus.
- d, e*, A ligament binding the os cuboides to the os calcis, and giving origin to muscles.
- f*, The flexor brevis minimi digiti.
- g, h*, The adductor pollicis.
- i, k, m*, The flexor brevis pollicis pedis;—*k*, Its tendinous origin from the os calcis and os cuneiforme externum;—*i*, Its connection with the adductor pollicis;—*m*, Its connection with that part from which the abductor pollicis was cut.
- l*, The transversalis pedis.

FIG. 12.

The Fourth Order of MUSCLES in the SOLE.

- a*, The insertion of the peroneus brevis.
- b*, The tendon of the peroneus longus joined to the metatarsal bone of the great toe, and sending tendons to the os cuneiforme internum.
- c*, The insertion of the tibialis anticus.
- d—i*, The interossei.
- d*, The abductor, and,
- e*, The adductor indicis pedis.
- f*, The abductor and adductor medii digiti.
- g*, The abductor, and,
- h*, The adductor tertii digiti.
- i*, ————— minimi digiti.

END OF PART SECOND.



P A R T III.
OF THE
BURSÆ MUCOSÆ.
OF THE
LIGAMENTS
AND
OTHER PARTS OF THE JOINTS.

OF THE BURSAE MUCOSAE IN GENERAL.

THE BURSAE MUCOSAE belong chiefly to the Extremities, and are found between Tendons and Bones, where these play upon each other, as at the Insertion of the Biceps Flexor Cubiti :

Or, completely surrounding the Tendons, as around the Tendons of the Flexores Digitorum :

Or, where Tendons rub on each other, as between those of the Extensores Carpi Radiales and Extensores Pollicis :

Or, between Tendons and External Parts, as over the Tendons of the Flexores Digitorum, in the Palm of the Hand :

Or, between Tendons and Ligaments of the Joints, as between the Tendons of the Flexores Digitorum and Ligamentum Carpi Capfulare.

They are found in a few places where Processes play upon Ligaments, as between the Acromion and Capfular Ligament of the Humerus :

Or, where the Bones play on each other, as between the Clavicle and Coracoid Process of the Scapula.

Some of the Bursae of contiguous Tendons communicate with each other ; as between the Extensor Carpi Radialis and Extensor Secundi Internodii Pollicis.

Others communicate, not only in Adults, but often also in Children, with the Cavity of the Joints ; as behind the Tendon of the Extensors of the Leg ; though this is more frequently the case in advanced age.

Their Structure is the same with the inner Layer of the Capfular Ligament of the Joints.

Like that, they are formed of a thin Pellucid Membrane, possessing little Sensibility, and joined to the surrounding parts by Cellular Substance, which, in many places, is intermixed with Fat.

Like the Capfular Ligament, they have commonly a thin Layer of Cartilage, or of tough Membrane, between them and the Bone.

Like it, they have reddish-coloured Masses of Fat projecting into their Cavities, from the edges of which Fringes are sent off ; as behind the Ligament of the Patella, or at the Insertion of the Tendo Achillis.

Like it also, the inside of the Bursae is remarkably smooth, being lubricated with the same kind of Gelatinous Mucus which is found in the Cavities of the Joints ; —the Mucus serving the same general purpose with that of the Joints, viz. to lessen Friction, and prevent the consequences which would otherwise result from it.

T A B L E L I.

Represents the BURSÆ MUCOSÆ of the SUPERIOR EXTREMITY.

FIG. 1.

The BURSÆ MUCOSÆ on the Fore-Part of the SUPERIOR EXTREMITY.

- A, The clavicle.
 B, The acromion scapulæ.
 C, The fore-part of the scapula.
 D, The ligament of the semilunar notch of the scapula.
 E, The coracoid process of the scapula.
 F, F, Two ligaments which tie the clavicle to the coracoid process.
 G, The head of the os humeri.
 H, The body of that bone.
 I, A bursa under the clavicle, where it plays upon the coracoid process of the scapula.
 J, A strong ligament which joins the acromion to the coracoid process.
 K, A large bursa between the acromion and ligament J, and the capsular ligament of the humerus.
 L, A small bursa, sometimes absent, between the point of the coracoid process, and capsular ligament of the humerus.
 M, The tendon of the subscapularis.
 N, A bursa between it and the capsular ligament of the humerus, which frequently communicates with the cavity of that joint.
 O, A bursa, not constant, between the origin of the coraco-brachialis and short head of the biceps, and capsular ligament of the humerus.
 P, The tendon of the teres major turned outwards.
 Q, A bursa between the tendon of the teres major and the os humeri, and upper part of the tendon of the latissimus dorsi.
 R, The tendon of the latissimus dorsi turned outwards.
 S, A small bursa between the tendon of the latissimus dorsi and os humeri.
 T, A bursa between the tendon of the long head of the biceps flexor cubiti and the body of the humerus.
 U, The radius.
 V, The ulna.
 W, The tendon of the biceps flexor cubiti turned inwards.
 X, A bursa, with a *peloton* of fat, between the tendon of the biceps and tubercle of the radius.
 Y, The origin of the extensor carpi radialis longior.
 Z, A small bursa, between the tendon common to the extensor carpi radialis brevis, extensor digitorum communis, and the round head of the radius.
a, The sheath of the flexor longus pollicis, slit open.
b, ————— flexors of the fore-finger, entire.
c, The thick part of the sheath of the tendons of the flexors of the middle finger.
d, The sheath of the tendons of the flexors of the ring-finger, slit open.
e, The sheath of the tendons of the flexors of the little finger slit open, and the tendons drawn forwards to shew them fully. Each of the sheaths of the flexors of the fingers is lined with a bursa.
f, A very large bursa surrounding the tendon of the flexor pollicis longus.
g, b, i, k, Four short burse on the fore-part of the tendons of the flexor sublimis digitorum in the palm.
l, A probe introduced into a large bursa between the tendon of the flexor pollicis longus, and the fore-part of the radius, and between the capsular ligament of the wrist and the os trapezium.
m, A probe put into a large bursa behind the tendons of the flexor digitorum profundus, and on the fore-parts of the end of the radius, and capsular ligament of the wrist. These two last-mentioned burse are sometimes found to communicate with each other.
n, A bursa between the tendon of the flexor carpi radialis and os trapezium.
o, A bursa between the tendon of the flexor carpi ulnaris and os pisiforme.

FIG. 2.

The BURSÆ MUCOSÆ on the Back of the SUPERIOR EXTREMITY.

A, The dorsum scapulæ.

B, The

FIG. 1.





FIG. 2.





- B, The cervix scapulæ.
 C, The acromion.
 D, The capsular ligament of the humerus.
 E, The tendon of the infra-spinatus turned outwards.
 F, The tendon of the teres minor turned outwards.
 I, The os humeri.
 J, The external, and,
 K, The internal condyle of the humerus.
 L, The radius.
 M, The ulna.
 N, The olecranon.
 O, The tendon of the triceps extensor cubiti turned down.
 P, A small bursa between it and the olecranon.
 Q, A bursa between the tendon of the extensor ossis metacarpi pollicis and the radius.
 R, R, A large bursa common to the extensores carpi radiales, where they cross behind the extensor ossis metacarpi pollicis.
- S, S, Another bursa common to the extensores carpi radiales, where they cross behind the extensor secundi internodii pollicis.
 T, A third bursa at the insertion of the tendon of the extensor carpi radialis brevior.
 U, A bursa for the tendon of the extensor secundi internodii pollicis, which communicates with the bursa S, S.
 V, Another bursa between the tendon of the extensor secundi internodii pollicis and metacarpal bone of the thumb.
 W, The tendons of the extensor of the fore, mid, and ring fingers.
 X, A bursa between these tendons and the ligament of the wrist.
 Y, Another bursa for the extensor of the little finger.
 Z, A bursa between the tendon of the extensor carpi ulnaris and ligament of the wrist.

TABLE

T A B L E LII.

Represents the BURSÆ MUCOSÆ of the INFERIOR EXTREMITY.

FIG. 1.

The BURSÆ MUCOSÆ on the Fore-Part of the INFERIOR EXTREMITY.

- A, The spine of the os ilium.
 B, The inner side of the os ilium.
 C, The os pubis.
 D, The neck of the thigh-bone.
 E, The root of the great trochanter.
 F, The thigh-bone.
 G, The iliacus internus.
 H, The psoas magnus.
 I, The insertion of the iliacus internus and psoas magnus into the trochanter minor.
 K, A very large bursa mucosa, between these two muscles and the capsular ligament of the thigh-bone.
 L, The pectineus.
 M, A bursa between the tendon of the pectineus and thigh-bone.
 N, The adductor brevis femoris.
 O, The gluteus minimus.
 P, The tendinous part of the gluteus medius.
 Q, A small bursa between the gluteus medius and trochanter major. Behind it the tendon of the pyriformis is inserted.
 R, A bursa between the tendon of the gluteus minimus and trochanter major.
 S, Part of the gluteus maximus, which is joined to the gluteus medius.
 T, Part of the vastus externus.
 U, A bursa between the gluteus maximus and vastus externus.
 V, The patella.
 W, The capsular ligament of the knee.
 X, The tendon of the extensors of the leg, cut and turned up.
 Y, A large bursa behind the tendon of the extensors of the leg.
 Z, A communication frequently found between this bursa and the cavity of the knee-joint.
a, a, The ligament which joins the patella to the tibia, cut and turned outwards;
b, A bursa behind it.

- c*, A fatty *peloton* hanging down into the cavity of the bursa.
d, The tibia.
e, The fibula.
f, The back part of the insertion of the sartorius, turned forwards.
g, The tendon of the gracilis.
h, ————— femi-tendinosus.
i, A large bursa between the tendons of the sartorius, gracilis, femi-tendinosus, and tibia.
k, The internal lateral ligament of the knee.
l, A bursa between the tendon of the tibialis anticus, and under part of the tibia and ligament of the ankle.
m, A bursa between the tendon of the extensor proprius pollicis pedis, and the tibia and capsular ligament of the ankle.
n, A bursa between the tendons of the extensor digitorum longus, and ligament of the ankle.

FIG. 2.

The BURSÆ MUCOSÆ on the Back Part of the INFERIOR EXTREMITY.

- A, The dorsum of the os ilium.
 B, The os sacrum.
 C, The os coccygis.
 D, The tuber of the os ischium.
 E, The large trochanter.
 F, The middle of the thigh-bone.
 G, The gluteus minimus.
 H, The pyriformis.
 I, A bursa mucosa between the gluteus medius and pyriformis.
 K, K, The obturator internus cut across.
 L, L, The gemini.
 M, A bursa between the obturator internus and os ischium.
 N, A probe put into a bursa, which is continued to a dotted line between the obturator internus, gemini, and capsular ligament of the thigh-bone.
 O, The quadratus femoris.
 P, The origin of the femi-membranosus, and long head of the biceps flexor cruris.
 Q, A small bursa mucosa.
 R, The origin of the femi-tendinosus turned back.

S, A


FIG. 1.





FIG. 2.



FIG. 3.







- S, A small bursa mucosa.
 T, The tendon of the gluteus maximus.
 U, A large bursa between the tendon of the gluteus maximus and root of the trochanter major.
 V, V, Two small bursæ between the tendon of the gluteus maximus and os femoris.
 W, The back part of a large bursa, between the tendinous part of the gluteus maximus and vastus externus.
 X, The internal condyle, and,
 Y, The external condyle of the thigh-bone.
 Z, The tibia.
 a, The fibula.
 b, The biceps flexor cruris turned downwards.
 d, The inner head of the gastrocnemius externus turned up.
 e, The semi-membranosus turned down.
 f, A bursa between the tendons of the semi-membranosus and gastrocnemius, and the ligament of the knee.
 g, A probe passed into a small bursa within the bursa f, from which there is a passage into the cavity of the joint of the knee.
 h, A probe passed into a bursa between the tendon of the semi-membranosus and the internal lateral ligament of the knee, from which there is a passage communicating with the joint of the knee.
 i, A small portion of the popliteus.
 l, A bursa under it, communicating with the cavity of the joint of the knee.
 n, The tendon of the peroneus longus.
 o, ————— brevis.
 p, A large bursa common to the tendons of the peronei.
 q, A bursa proper to the tendon of the peroneus brevis.
 r, A ligament which ties the fibula to the os calcis.
 s, The tendo Achillis turned down.
 t, A bursa between the tendo Achillis and os calcis.
 u, A *peloton*, or mass of fat, which projects into the cavity of that bursa.
 v, The tendon of the plantaris.
 w, A bursa between the os calcis and flexor pollicis longus.
 x, The flexor digitorum longus.
 y, A bursa between that flexor and the tibia and os calcis.
 z, The tendon of the tibialis posticus.
 &, A bursa between this tendon and the tibia and astragalus.

FIG. 3.

A View of the BURSÆ MUCOSÆ in the SOLE.

- A, The os calcis.
 B, The tendo Achillis.
 C, C, The abductor minimi digiti cut across.
 D, The tendon of the peroneus longus.
 E, A second bursa for that tendon.
 F, An oblong *peloton* of fat within this bursa.
 G, The fleshy,
 H, The tendinous part of the flexor longus pollicis pedis.
 I, A bursa common to this tendon, and the tendon of the flexor digitorum profundus.
 K, A fatty *peloton* at the upper end of this bursa.
 L, An imperfect septum between the two last-named tendons, containing some fat.
 M, The tendon of the tibialis posticus in its bursa.
 N, The place at which the flexor digitorum sublimis is cut off.
 O, The massa carnea Jacobi Sylvii, or flexor tertius.
 P, The abductor pollicis pedis cut off from the os calcis.
 Q, R, S, T, U, The bursæ mucosæ of the flexor tendons slit open nearly their whole length.

The BURSÆ MUCOSÆ of this and the preceding TABLE are all represented as slit open, and several of them inflated.

OF THE LIGAMENTS, &c. OF THE JOINTS.

LIGAMENTS are *white, strong, flexible* bodies, of an intermediate firmness between Cartilage and common Membrane.

They are *composed* of Fibres variously disposed; the greater part of them, however, running in a longitudinal direction.

The Ligaments of moveable Joints arise, for the most part, at the junction of the Bodies of the Bones with their Epiphyses, from the *Cervix*, and beyond the edges of the articulating Cartilage of one Bone, and are fixed, in a similar manner, into the corresponding parts of the Bone adjoining.

The Ligaments thus fixed are called *Capsular*, from their forming a *Purse* or *Bag*, which includes the Joint.

Where *variety of Motion* is allowed, the Capsular Ligament is nearly of equal strength round the whole circumference of the Joint; but, where the Joint is of the nature of a *Hinge*, the Ligament is strongest at the sides of that Hinge.

The *outer* part of the Capsular Ligament is formed of a continuation of the *Periosteum*, which is connected to the surrounding parts by Cellular Substance; while the *inner* Layer,—remarkably thin and dense,—is reflected over the Bones and Cartilages which the Ligament includes; one part of it thus forming *Periosteum*, and the other *Perichondrium*.

In certain parts of the Body, there are, besides the Capsular Ligaments, others for the firmer connection of the Bones, or for confining the motion to one particular side; as the *Round Ligament* of the Thigh, or *Crucial* or *Lateral* Ligaments of the Knee.

Wherever the Ligaments are few, long, and weak, the Motions are more extensive; and, on the contrary, where the Ligaments are numerous, short, and strong, the Motions are more limited.

In some parts of the Body, Ligaments supply the place of Bones, as in the *Pelvis*: In others, they give origin to Muscles, as between the *Radius* and *Ulna*: In some parts, they assist in connecting immoveable Bones, as at the *Os Sacrum* and *Os Innominatum*: In others, they form a Socket in which moveable Bones play, as where part of the *Astragalus* moves on the Ligament stretched between the *Os Calcis* and *Os Scaphoides*.

Ligaments have numerous *Blood-vessels*, which can be readily injected.

Upon the inner Surface of the Capsular Ligaments, their *Arteries* secrete a Liquor which assists in the lubrication of the Joints.

The *Nerves* of Ligaments are very minute, but in some parts can be easily traced upon their Surface.

The *Sensibility* of Ligaments, in the sound state, is in-

considerable; when in a state of Inflammation, however, they are found to occasion extreme pain.

Use: The Capsular Ligaments connect Bones together, assist in the Secretion of the Synovia, which they contain, and prevent the other parts from being pinched in the Joint. The other Ligaments join Bones together, and preserve them in their proper situation. In many parts, they give attachment to Tendons, and, in some, to the Fleishy parts of Muscles.

SYNOVIAL ORGANS,

Commonly called GLANDS of the JOINTS.

These are *Masses of Fat* found in the greater number of the Joints, covered with a continuation of the inner Layer of the Capsular Ligament, and projecting in such a manner as to be gently pressed, but not bruised, by the motions of the Joint; and, in proportion as these motions are more or less frequent, the Liquor which they secrete is discharged in a greater or smaller quantity.

In some Joints, they have the same appearance with the common Fat of the Body; in others, they are of a redder colour, from the numerous Blood-vessels dispersed upon them.

They have been generally considered as *Glands* lodged within Masses of Fat; but, upon a minute inspection, no knotty or glandular bodies are to be found in them; nor have they the appearance of Glands, farther than in being Secreting Substances; which circumstance alone assimilates them to the nature of Glands.

From the edges of these Fatty Bodies, *Fimbriae* hang loose, and convey a lubricating Liquor, called *Synovia*, into the Cavity of the Joints.

From the extremities of these *Fringes*, the Liquor can be readily squeezed out by pressure; but their Cavities and Orifices are so minute, or are otherwise of such a nature, as to have hitherto eluded discovery.

The *Fimbriae* have been generally considered as *Excretory Ducts of Glands* within the Joints. Dr MONRO, however, in his Work upon the *Bursæ Mucosæ*, supposes them to be of the nature of the *Follicles of the Urethra*, which prepare a Mucilaginous Liquor without the assistance of any knotty or glandular Organ.

The *Arteries* which supply these Bodies with Blood for their Secretions, and the *Veins* which return the Blood after the Secretion has been performed, can be readily seen; but no *Nerves* can be traced into them; nor does it appear that they possess a higher degree of sensibility than the other parts of the Joints already described; although, when they inflame and suppurate, they have in some

some instances been observed to occasion the most excruciating pain.

The *Synovia*, which is a thin Mucilaginous Liquor, resembling the glair of an egg, appears to be furnished, not only by the Substances already mentioned, but also by the inner Surface of the Capsular Ligaments in general, and serves for the lubrication of the Joints.

LIGAMENTS OF THE HEAD AND TRUNK.

LIGAMENTS of the LOWER JAW.

The *Capsular Ligament*, on each side, which arises from the whole margin of the Articular Cavity of the Temporal Bone, and is inserted, first into the edge of the Inter-articular Cartilage, formerly taken notice of, and afterwards round the Cervix of the Lower Jaw. This Ligament, like others which belong to Joints of the hinge-kind, is thickest and strongest at the sides of the Joint, to confine the lateral motion of the Jaw. Tab. LIII. Fig. 1. *f*. Fig. 2. *d, d*.—Tab. XXX. Fig. 17. *r, t, t*.

By it the Jaw is allowed to move upwards, downwards, or a little forwards or backwards, or to a side; and the motions are rendered easier by the intervention of the Inter-articular Cartilage, which follows the Condyle in its different motions.

The *Suspensory Ligament* of the Stylo-Glossus, which is attached by one end to the Styloid Process, and to a Ligament running from that Process to the *Os Hyoides*, and, by the other end, to the Angle of the Lower Jaw;—serving to support the Stylo-Glossus, and to give origin to part of it. Tab. LIII. Fig. 1. *k*.

The *Lateral Ligament*, which arises from the margin of the Articular Cavity of the Temporal Bone, and is inserted into the inner Surface of the Angle of the Lower Jaw, near its posterior Foramen;—assisting to keep the Jaw *in situ*, and to prevent the inferior Maxillary Vessels and Nerve from being injured by the action of the Pterygoid Muscle. Tab. LIII. Fig. 1. *g*.

LIGAMENTS connecting the HEAD with the First and Second VERTEBRÆ of the NECK, and these two VERTEBRÆ with each other.

The two *Capsular Ligaments*, which arise from the margin of the superior Articulating Processes of the Atlas, and are inserted into the Base of the Condyles of the Occipital Bone, where the Head has its flexion and extension without rotation. Tab. LIII. Fig. 3. *e, f*.

The *Circular Ligament*, which arises from the edge of the Spinal Hole of the first Vertebra, is connected with the Capsular Ligaments of the superior Articulating Processes of the Atlas, and is inserted into the edge of the Foramen Magnum of the Occipital Bone. Tab. LIII. Fig. 3. *d*.

The two *Capsular Ligaments*, which fix the inferior Oblique Processes of the Atlas to the superior Oblique of the Vertebra Dentata, and admit of the rotation of the Head, with a small degree of flexion to either side. Tab. LIII. Fig. 3. *n, n*.

The *Perpendicular Ligament*, which fixes the point of the Processus Dentatus of the second Vertebra to the edge of the anterior part of the Foramen Magnum, and is twisted in the rotation of the Head. Tab. XXXI. Fig. 6. *q*.

The two *Lateral, or Moderator Ligaments*, which arise each from the side of the Processus Dentatus, and run outwards and upwards, to be fixed to the inner part of the side of the Atlas, and to the inner edge of the Foramen Magnum. Tab. XXXI. Fig. 6. *p, p*. They are short, but of great strength, and prevent the Head from turning too far round.

The *Transverse Ligament*, which arises from the inner side of the Atlas, and, going across behind the Processus Dentatus, is fixed to the opposite side of the Atlas. Tab. XXXI. Fig. 6. *o*.

The edges of this Ligament extend upwards and downwards, and form two Processes, called its *Appendices*, which are fixed to the Foramen Magnum and Processus Dentatus. The middle of the Ligament is remarkably firm where that Process plays upon it. It keeps the Processus Dentatus in its place, and prevents it from injuring the Spinal Marrow in the different motions of the Head.

LIGAMENTS of the Other VERTEBRÆ.

The *Anterior Common Ligament* of the Vertebrae, which is a strong Tendinous Band, extending along the convex or outer part of the Vertebrae, from the upper to the under region of the Spine. Tab. LIII. Fig. 4. *a, b*. It begins at the second Cervical Vertebra, and descends as far as the Os Sacrum, where it spreads out, becomes thinner, and vanishes about the under part of this Bone. It is much thicker upon the fore-part than on the sides of the Vertebrae, by which the Bones are more firmly united anteriorly, and is thinnest in the Neck and Loins, where the motions of the Spine are greatest. Internally, it is blended with the Periosteum, and through its whole course, it sends off small Processes to be fixed to the Bodies of the Vertebrae, by which their connection is made more secure. While it assists in binding the Vertebrae together, it prevents the Spine from being stretched too much backwards.

The *Crucial Inter-vertebral Ligaments*, which are numerous and short, but strong, situated behind the Ligamentum Commune Anterior; crossing each other obliquely, they join the Bodies of the Vertebrae together, upon the outer edges of the Inter-vertebral Substances, to which also they firmly adhere. Tab. LIII. Fig. 5. *c, d*.

The *Inter-vertebral Substances*, already described in

p. 46. which join the Bodies of the Vertebrae together, and allow an yielding motion in all directions.

These Substances are so compressible as to yield to the weight of the upper part of the Body; so that, after having been in the erect posture through the course of the day, the height of a person in the prime of life, and of middle stature, is diminished from half an inch to an inch in the evening, but, after a night's rest in the usual attitude, it is found to be restored.

The *Ligaments* which run from the edge of the Bony Arch and Spinous Process of one Vertebra to that of the next, so as to assist in filling up the Interstices, and in fixing the Vertebrae together.

A *Ligamentous Cord* which fixes the points of the Spinous Processes together. Tab. LIII. Fig. 9. f, f.

The *Cervical Ligament*, termed *Ligamentum Nuchae*, vel *Colli*, which arises from the perpendicular Spine of the Occipital Bone, and descends on the back part of the Neck, adhering to the Spinous Processes of the Cervical Vertebrae, and giving origin to part of the Trapezius

Ligaments between the Transverse Processes of the Vertebrae of the Back, fixing these Processes to each other. Tab. XLIII. c, c.

The *Capsular Ligaments*, which join the Articulating Processes to each other.

The *Posterior* or *Internal Common Ligament* of the Vertebrae, somewhat similar to the Anterior one. Tab. LIII. Fig. 6. 7.

It begins at the anterior edge of the Foramen Magnum, and passes along the inner or concave part of the Bodies of the Vertebrae, becoming broader over each of the Inter-vertebral Substances. It adheres firmly to their upper and under edges, and terminates at the lower part of the Os Sacrum.—It prevents the Spine from being too much bent forwards.

LIGAMENTS of the RIBS.

The *Capsular Ligaments of the Heads of the Ribs*, which arise from these Heads, and are fixed to the Circumference of the Pits in the sides of the Bodies of the Vertebrae and Inter-vertebral Cartilages. The outer part of each Ligament sends off, or is connected with, radiated Fibres which are spread out upon the sides of the Vertebrae. Tab. LIII. Fig. 8. c, c.

The *Capsular Ligaments of the Tubercles of the Ribs*, which arise round the Articular Pits on the points of the Transverse Processes of the Vertebrae of the Back, and are fixed round the Tubercles of the Ribs.

The *Internal Ligaments of the Back of the Ribs*, called *Ligamenta Transversaria Interna*, which arise from the

inferior Surfaces of the Transverse Processes, and are fixed to the Superior Margins of the Necks of the nearest Ribs. Tab. LIII. Fig. 8. d, d.

The *External Ligaments of the Necks of the Ribs*, called *Ligamenta Transversaria Externa*. They arise from the points of the Transverse Processes externally, and are fixed to the back part of the Necks of the Ribs. Tab. LIII. Fig. 9. c, c.

Ligamenta Cervicis Costarum Externa, or *External Ligaments of the Necks of the Ribs*, which arise from the external Margins of the inferior Oblique Processes, and descend obliquely outwards, to be fixed to the upper and outer part of the Necks of all the Ribs. Tab. LIII. Fig. 9. e, e.

The Ligaments at this end of the Ribs, together with the situation of the Transverse Processes, admit of their motion upwards and downwards, but prevent them from moving in any other direction.

Short Ligamentous Fibres, which run from the Margins of the anterior extremities of the Ribs to the Margins of their corresponding Cartilages; the Cartilages and Ribs being joined by an union of Substance. Tab. LIII. Fig. 13. d.

Radiated Ligaments, which go from the anterior Surfaces of the Capsular Ligaments upon the external Surface of the Sternum. Tab. LIII. Fig. 12. d. 13. e.

Many of the Fibres of these Ligaments intermix with their fellows on the opposite side.

The *Capsular Ligaments of the Cartilages of the Ribs*, which arise from the Margins of the Articular Cavities of the Sternum, and are fixed round the extremities of the seven true Ribs. Tab. LIII. Fig. 13.

Membrane proper to the Sternum, which is a firm Expansion composed of Tendinous Fibres running in different directions, but chiefly in a longitudinal one, and covering the anterior and posterior Surfaces of the Bone, the Membrane itself being confounded with the Periosteum. Tab. LIII. Fig. 12. a, b.

Ligaments of the Cartilago Ensiformis, Tab. LIII. Fig. 12. g, g, which are part of the proper Membrane of the Sternum, divided into strong Bands running obliquely from the under and fore part of the second Bone of the Sternum, and from the Cartilages of the seventh pair of Ribs, to be fixed to the Cartilago Ensiformis.—The Ligaments covering the Sternum serve considerably to strengthen it.

Thin Tendinous Expansions, which run over the Intercostales at the fore-part of the Thorax, and connect the Cartilages of the Ribs to each other. They are chiefly seated in the spaces unoccupied by the intercostales Externi. Tab. LIII. Fig. 12. f, f.

LIGAMENTS OF THE BONES OF THE PELVIS.

THE two *Transverse Ligaments of the Pelvis*, which arise from the posterior part of the Spine of the Os Ilium, and run transversely. The one is *superior*, and is fixed to the Transverse Process of the last Vertebra of the Loins; the other *inferior*, and is connected to the first Transverse Process of the Os Sacrum. Tab. LIII. Fig. 4. *p, q*.

The *Ileo-Sacral Ligaments*, which arise from the posterior Spinous Process of the Os Ilium, descend obliquely, and are fixed to the first, third, and fourth spurious Transverse Processes of the Os Sacrum. Tab. LIII. Fig. 10. *f, g, b*.

These, with the two Transverse Ligaments, assist in binding the Bones together to which they are connected.

The *Capsular Ligament of the Symphysis of the Os Ilium and Os Sacrum*, which surrounds the Joint, and assists in connecting the two Bones to each other.

A *very thin Cartilage* within this Joint, which cements the two Bones strongly together, and which constantly adheres to the Os Sacrum, when the Joint is opened. Tab. LIII. Fig. 11. *f*.

A *Ligamentous and Cellular Substance* containing *Mucus*, which forms the back part of this Joint, also assisting to fix the two Bones to each other, in such a manner as to allow no motion; the Joint, however, along with its fellow, and that between the Os Pubis, being useful in diminishing the effects which might result from concussion. Tab. LIII. Fig. 11. *g*.

The two *Sacro-Ischiatic Ligaments*, situated in the under and back part of the Pelvis. They arise in common from the Transverse Processes of the Os Sacrum, from the under and lateral part of that Bone, and from the upper part of the Os Coccygis. The first, called the *Large, External, or Posterior Sacro-Ischiatic Ligament*, descends obliquely, to be fixed to the Tuberosity of the Os Ischium. Tab. LIII. Fig. 10. *k*. The other, called the *Small, Internal, or Anterior Sacro-Ischiatic Ligament*, runs transversely to be fixed to the Spinous Process of the Os Ischium. Tab. LIII. Fig. 10. *m*. These two Ligaments assist in binding the Bones of the Pelvis, in supporting its Contents, and in giving Origin to part of its Muscles. By the External, the Notch of the Ilium is formed into a Hole for the passage of the Pyriform Muscle, the Sciatic Nerve, and the Blood-vessels which belong to the outside of the Pelvis. Between the two Sacro-Sciatic Ligaments, an opening is left for the passage of the Obturator Internus.

The two *Membranous Productions* which are connected with the large Sacro-Ischiatic Ligament, termed by WEITBRECHT the *Superior and Inferior Appendices* of the large Sacro-Ischiatic Ligament.

The *Superior Appendix*, which is Tendinous, arises from the back part of the Spine of the Os Ilium, and is

fixed along the outer edge of the Ligament, which it increases in breadth. Tab. LIII. Fig. 10. *l*.

The *Inferior or Falciform Appendix* is situated within the Cavity of the Pelvis; the back part of it is connected with the middle of the large External Ligament, and the remainder is extended round the Curvature of the Os Ischium. Tab. LIII. Fig. 11. *n, z*.

These two Productions assist the large Sacro-Ischiatic Ligament in furnishing a more commodious situation for, and attachment of, part of the Gluteus Maximus and Obturator Internus.

Besides the Ileo-Sacral and Sacro-Ischiatic Ligaments, several other *Slips* are observed upon the back of the Os Sacrum, which descend in an irregular manner, and strengthen the connection between that Bone and the Os Illia. Tab. LIII. Fig. 10. *i, i*.

The large Holes upon the back part of the Os Sacrum are also surrounded with various *Ligamentous Expansions* projecting from one Tubercle to another, and giving origin to Muscular Fibres, and protection to small Vessels and Nerves which creep under them. Tab. LIII. Fig. 10.

A *General Covering* sent down from the Ligaments of the Os Sacrum, which spreads over and connects the different pieces of the Os Coccygis together, allowing considerable motion, as already mentioned in the description of this Bone. Tab. LIII. Fig. 10.

Longitudinal Ligaments of the Os Coccygis, which descend from those upon the Dorsum of the Os Sacrum, to be fixed to the back part of the Os Coccygis. Tab. LIII. Fig. 10. *n*. The Ligaments of this Bone prevent it from being pulled too much forward by the action of the Coccygeus, and they restore the Bone to its natural situation, after that Muscle has ceased to act.

The *Inguinal, or POUPART'S, or FALLOPIUS'S Ligament, or Cruval Arch*, which runs transversely from the anterior-superior Spinous Process of the Os Ilium to the Crest or Angle of the Os Pubis. It has been already described as the inferior margin of the Tendon of the External Oblique Muscle of the Abdomen. Tab. XXXIV. Trunk, *q*.

The *Capsular Ligament of the Symphysis of the Os Pubis*, which joins the two Bones to each other externally.

The *Ligamentous Cartilage*, Tab. LIII. Fig. 11. *u*, which unites the two Os Pubis so firmly together as to admit of no motion, excepting in the state of pregnancy, when it is frequently found to be so much thickened as to be capable of yielding a little in the time of Delivery.

The *Obturator Membrane, or Ligament of the Foramen Thyroideum*, Tab. LIII. Fig. 11. *r*. Tab. LIII. Fig. 1. *q, r, s, u, z*, which adheres to the margin of the Foramen

men Thyroideum, and fills the whole of that opening, excepting the Oblique Notch at its upper part for the passage of the Obturator Vessels and Nerve. It assists in supporting the Contents of the Pelvis, and in giving

origin to the Obturators. By yielding a little in the time of Labour, it contributes in a small degree to an easier Delivery.

LIGAMENTS OF THE SUPERIOR EXTREMITY.

LIGAMENTS of the CLAVICLE.

The *Radiated Ligaments*, which arise from the outer Surface of the inner end of the Clavicle, and are fixed round the edge of the corresponding Articular Cavity of the Sternum. Tab. LIV. Fig. 1. *f*.

The *Capsular Ligament* which lies within the former. Tab. LIV. Fig. 2. *i*.

The *Inter-articular Cartilage*, which divides the Joint into two distinct Cavities, and accommodates the Articulating Surfaces of the Clavicle and Sternum. Tab. LIV. Fig. 1. *f*.

The *Inter-clavicular Ligament*, joining the Clavicles together behind the top of the Sternum, and partly formed by a continuation of the radiated Ligaments. Tab. LIV. Fig. 1. *b*.—Tab. XXXII. Fig. 1. *d*.

The *Ligamentum Rhomboideum*, which arises from the inferior rough Surface at the anterior extremity of the Clavicle, and is fixed to the Cartilage of the first Rib. Tab. LIV. Fig. 1. *g*.

By the Ligaments of this Joint, with the assistance of the intervening Cartilage, the Shoulder is allowed to move in different directions, as upon a centre.

The *Ligaments* which join the posterior extremity of the Clavicle to the Acromion, having a Capsular Ligament within, and sometimes an Inter-articular Cartilage. Tab. LIV. Fig. 5. *b*.

The *Ligamentum Conoideum*, which arises from the root of the Coracoid Process, and is fixed to the Tubercle at the outer end of the Clavicle.

The *Ligamentum Trapezoideum*, which arises from the point of the Coracoid Process, and is fixed to the under edge of the Clavicle. Tab. LIV. Fig. 5. *g*.

A *thin Ligamentous Slip*, which comes from the Tendon of the Subclavius, or from the Clavicle, and joins the Trapezoid Ligament. Tab. LIV. Fig. 3. *o*.

The Ligaments fixing the Clavicle to the Scapula are of such strength, as to allow only a small degree of motion, and that chiefly of a rolling or twisting nature.

LIGAMENTS proper to the SCAPULA.

The *Proper Anterior Triangular Ligament of the Scapula*, which arises broad from the external Surface of the Coracoid Process, and becomes narrower where it is fixed

to the posterior Margin of the Acromion. Tab. LIV. Fig. 5. *f*.

This Ligament forms one continued Surface. It is thickest, however, on each side; and these thicker parts are united by a thin intermediate Ligamentous Membrane, which, when removed, gives to the Ligament the appearance of being double, Tab. XXXII. Fig. 2. *n*. It confines the Tendon of the Supra-spinatus, and assists in protecting the upper and inner part of the Joint of the Humerus.

The *Proper Posterior Ligament of the Scapula*, which is sometimes double, and is stretched across the Semilunar Notch of the Scapula, forming that Notch into one or two Holes for the passage of the superior-posterior Scapular Vessels and Nerve. Tab. LIV. Fig. 3. *g*.

LIGAMENTS, &c. of the JOINT of the SHOULDER.

The *Capsular Ligament*, which arises from the Cervix of the Scapula, behind the Margin of the Glenoid Cavity, and is fixed round the Neck of the Os Humeri, loosely inclosing the ball of that Bone. Tab. LIV. Fig. 3. *i*. Fig. 6. *d*.

A *small Fimbriated Organ* within the Capsular Ligament, for the Secretion of Synovia. Tab. LIV. Fig. 7. *e*.

A *Sheath* sent down from the fore-part of the Capsular Ligament, between the Tuberosities of the Os Humeri, which incloses the Tendon of the long Head of the Biceps Flexor Cubiti, and prevents it from starting out of its place. Tab. LIV. Fig. 5. *r*.

Additional Ligamentous Bands of the Capsular Ligament, which adhere to its anterior Surface. Tab. LIV. Fig. 5. *m*.

What gives most strength to this Joint, as well as to several other Joints of the Body, is the covering from the surrounding Muscles.

From the shallowness of the Glenoid Cavity, from the extent and looseness of the Capsular Ligament, and from the structure of the other parts of the Joint, more extensive motion is allowed to the Os Humeri than to any other Bone of the Body; as it can not only move freely to every side, but also possesses a considerable degree of motion upon its own axis.

LIGAMENTS,

LIGAMENTS, &c. of the JOINT of the ELBOW.

The *Capsular Ligament*, which arises round the Margin of the Articular Surface, at the lower end of the Os Humeri, and is fixed about the edge of the Articular Surface of the Ulna, and also to the Coronary Ligament of the Radius. Tab. LIV. Fig. 8. *l, l.*—Tab. XXXII. Fig. 3.

The sides of the Elbow-joint are strengthened by two *Ligamentous Bands*, which adhere so firmly to the outer Surface of the Capsular Ligament, that they appear to be part of its Substance, viz.

The *Brachio-Cubital*, or *Internal Lateral Ligament*, which arises from the fore-part of the inner Condyle of the Os Humeri, and spreads out, in a radiated manner, to be fixed to the inside of the Coronoid Process of the Ulna. Tab. LIV. Fig. 8. *m.*—And

The *Brachio-Radial*, or *External Lateral Ligament*, which is like the former, but larger, Tab. LIV. Fig. 9. *b.* It arises from the external Condyle of the Os Humeri, and is expanded upon the Coronary Ligament into which it is inserted.

The *Coronary, Annular, or Orbicular Ligament* of the Radius, which approaches to the firmness of a Cartilage. It arises from one side of the small Semilunar Cavity of the Ulna, and, after surrounding the Neck of the Radius, is fixed to the other side of that Cavity. The upper edge of it is incorporated with, and may be considered as a part of the Capsular Ligament, while its under edge is fixed round the Neck of the Radius, allowing that Bone to move freely round its own axis, upon the Articular Surface of the Os Humeri, and in the small Semilunar Cavity of the Ulna. Tab. LIV. Fig. 8. *m.*

Besides the Ligaments already described, there are others which run in various directions upon the fore and back parts of the Joint, contributing to its strength, and having the names of *Anterior* and *Posterior Accessory Ligaments*. Tab. LIV. Fig. 8. *o, p.* Fig. 9. *l.*

There are also two Tendinous Substances, termed *Inter-muscular Ligaments of the Os Humeri*, which extend along the under and lateral parts of this Bone, giving origin to part of the Muscles situated at this part of the Arm.

The Ligaments and Bones of the Joint of the Elbow form a complete Hinge, which allows the Fore-arm to have free flexion and extension upon the Os Humeri, but no rotation when the Arm is in the extended state, though a small degree of it is perceptible when the Joint is moderately bent, and the Ligaments thereby relaxed.

Within the Capsular Ligament, and chiefly in the upper part of the Pit of the Os Humeri in which the Olecranon plays, the *Fatty Substance* is lodged for the lubrication of the Joint. Tab. LI.

A *similar Substance*, but much smaller in quantity, is also found in the Depression in which the Coronoid Process of the Ulna moves.

LIGAMENTS between the Bodies, and between the Under Ends of the RADIUS and ULNA.

The *Interosseous Ligament*, which extends between the sharp Ridges of the Radius and Ulna, filling up the greater part of the space between these two Bones. It is broadest in the middle, in consequence of the bones here being largest at their extremities, and is composed of small *Fasciculi*, which run obliquely downwards and inwards. Two or three of these Slips, however, go in the opposite direction; and one of them, termed *Oblique Ligament*, and *Chorda Transversalis Cubiti*, is stretched between the Tubercle of the Ulna and under part of the Tubercle of the Radius.

In different parts of the Interosseous Ligament there are Perforations for the passage of Blood-vessels from the fore to the back part of the Fore-arm, and a large Opening is found at the upper edge of it, which is occupied by Muscles. Tab. LIV. Fig. 8. *r.*

This Ligament assists in binding the Ulna and Radius together, prevents the Radius from rolling too much outwards, and furnishes a commodious attachment for Muscles.

The *Capsular, or Sacciform Ligament*, which arises from the edges of the Semilunar Cavity at the under end of the Radius, and surrounds the Head of the Ulna, allowing the Radius to turn upon the Ulna, in performing the different motions of Pronation and Supination of the Hand. Tab. LIII. Fig. 10. *e.*

LIGAMENTS, &c. between the FORE-ARM and WRIST.

The *Capsular Ligament*, which arises from the Margin of the Glenoid or Navicular Cavity of the Radius, and from the edge of the moveable Cartilage at the Head of the Ulna, and is fixed to the Cartilaginous edges of the three first Bones of the Carpus. Tab. LIV. Fig. 11. *i.*

The *Inter-articular Cartilage*, placed between the Head of the Ulna and Os Cuneiforme, and which is a continuation of the Cartilage covering the end of the Radius. It is concave above and below, and is connected loosely to the end of the Styloid Process. Tab. LIV. Fig. 10. *c.*

The *Two Lateral Ligaments*, one of which arises from the Styloid Process at the under end of the Radius, and is fixed to the Os Naviculare, and the other from the Styloid Process of the Ulna, and is fixed to the Cuneiform and Pisiform Bones.

The Ligaments of this Joint allow extensive motion forwards and backwards; and a considerable degree of it to either side.

The *Mucous Ligament*, which lies within the Joint, Tab. LIV. Fig. 11. *o.* and extends from the Groove between the two first Bones of the Carpus, to the corresponding part of the Radius. It is supposed to regulate the Mucous Organ connected with it.

LIGAMENTS of the CARPUS.

The *Anterior, Annular, or Transverse Ligament*, which is stretched across from the projecting Points of the *Ossa Pisiforme and Unciforme*, to the *Scaphoides* and *Trapezium*, and forms an Arch which covers and preserves in their places the Tendons of the Flexor Muscles of the Fingers. Tab. XXXII. Fig. 6. *θ*.

The *Capsular Ligament*, which arises from the Cartilaginous edge of the upper Row of the Carpus, and is fixed in a similar manner to that of the under Row, chiefly admitting of flexion and extension, and that in a smaller degree than in the former Joint. Tab. LIII. Fig. 15. *k*.

The *Short Ligaments* of the Bones of the Carpus, which are *small Ligamentous Slips* running in various directions, joining the different Bones of the Carpus,—first of the same Row, then of the two Rows together, Tab. LIV. Fig. 12. 13. 14. They are termed *Oblique, Transverse, Capsular, and Proper* Ligaments of the Bones of the Wrist, and admit only of a small degree of yielding between the different Bones in the same Row.

LIGAMENTS between the CARPAL and METACARPAL BONES.

The *Articular Ligaments*, which arise from the Margins of the second Row of the Carpal Bones, and are fixed to the Margins of the adjoining Bones of the Metacarpus, Tab. LIV. Fig. 12. 16. *Other Ligaments* run in a radiated manner from the Carpal to the Metacarpal Bones; the whole getting the names of *Articular, Lateral, Straight, Perpendicular, &c.* according to their different directions.

From the flatness of the Articular Surfaces, and strength of the Connecting Ligaments, very little motion is allowed between the Carpus and Metacarpus.

LIGAMENTS between the Extremities of the METACARPAL BONES.

The *Interosseous Ligaments* at the Bases of the Metacarpal Bones, Tab. LIV. Fig. 12. 14. They are short Slips, which run transversely, and join these Bones to each other, obtaining the names of *Dorsal, Lateral, or Palmar*, according to their different situations.

The *Interosseous Ligaments* at the Heads of the Metacarpal Bones, which run transversely in the Palm, and connect the Heads of these Bones to each other. Tab. LIV. Fig. 12. *t*.—Tab. XXXII. Fig. 6. *λ*.

LIGAMENTS at the Base of the METACARPAL BONE of the THUMB, and at the First JOINT of the FINGERS.

These consist of the *Capsular Ligaments* which inclose the Joints, and the *Lateral Ligaments* which are situated at the sides of the Joints, adhering to and strengthening them; the whole admitting of flexion, extension, and lateral motion. Tab. XXXII. Fig. 6.

LIGAMENTS of the First and Second JOINTS of the THUMB, and Second and Third JOINTS of the FINGERS.

The *Capsular Ligaments* inclosing the Joints. Tab. LIV. Fig. 17.—Tab. XXXII. Fig. 6.

The *Lateral Ligaments* placed at the sides of the Joints, and adhering to the Capsular Ligaments, confining the motion to flexion and extension. Tab. LIV. Fig. 17. *e*.

LIGAMENTS retaining the TENDONS of the MUSCLES of the HAND and FINGERS, in situ.

The *Anterior, Transverse, or Annular Ligament* of the Wrist,—already described.

The *Vaginal Ligaments of the Flexor Tendons*, which are fine Membranous Webs connecting the Tendons of the *Sublimis*, first to each other, then to those of the *Profundus*, and forming, at the same time, *Bursæ Mucosæ*, which surround the Tendons. Tab. LI. Fig. 1. *g—k*.

The *Vaginal or Crucial Ligaments of the Phalanges*, which arise from the Ridges on the concave side of the Phalanges, and run over the Tendons of the Flexor Muscles of the Fingers. Tab. L. Fig. 1. *s, t, u*.—Tab. LI. Fig. 1. *b, c, d*. Upon the Body of the Phalanges, they are thick and strong, to bind down the Tendons, while their Muscles are in action; but over the Joints they are thin, and have, in some parts, a Crucial appearance, to allow the ready motion of the Joints.

The *Accessory Ligaments of the Flexor Tendons of the Fingers*, which are *small Tendinous Fræna*, arising from the first and second Phalanges of the Fingers. They run obliquely forwards within the Vaginal Ligaments, terminate in the Tendons of the two Flexor Muscles of the Fingers, and assist in keeping them in their places. Tab. LIV. Fig. 18. *f, g, h*.

The *External Transverse or Posterior Annular Ligament of the Wrist*, which is part of the Aponeurosis of the Fore-Arm, extending across the back of the Wrist, from the inner side of the extremity of the Ulna and *Ossa Pisiforme* to the outer side of the extremity of the Radius. Tab. XXXIII. Fig. 2. *E*. It is connected with the small Annular Ligaments which tie down the Tendons of the *Extensores Ossis Metacarpi et Primi Internodii Pollicis*, and the *Extensor Carpi Ulnaris*. Tab. L. Fig. 5. *i, l*.

The *Vaginal Ligaments* which adhere to the former Ligaments, and serve as Sheaths and *Bursæ Mucosæ* to the Extensor Tendons of the Hand and Fingers. Tab. LI. Fig. 2.

The *Transverse Ligaments* of the Extensor Tendons, which are Aponeurotic Slips running between the Tendons of the *Extensor Digitorum Communis*, near the Heads of the Metacarpal Bones, and retaining the Tendons in their places. Tab. L. Fig. 5. *o*.

LIGAMENTS OF THE INFERIOR EXTREMITY.

LIGAMENTS, &c. connecting the Os FEMORIS with the Os INNOMINATUM.

The *Capsular Ligament*, the largest and strongest of the Articular Ligaments. It arises round the outside of the Brim of the Acetabulum, embraces the Head of the Thigh-Bone, and incloses the whole of its Cervix as far as the root or outer extremity, round which it is firmly connected. Tab. XXXI. Fig. 18. E.

The *outer part* of the Capsular Ligament is extended farther down than the *inner*, which is reflected back upon the Neck of the Bone, and in certain parts forms *Retinacula*. Tab. LV. Fig. 1. 2.

It is not every where of the same strength. It is thickest at its anterior and outer part; thinner where it is covered by the Iliacus Internus; and thinnest posteriorly, where the adjacent Quadratus is opposed to it.

It is strengthened on its outer Surface by various *Accessory* or *Additional Slips*, which run down from the Fascia Lata and surrounding Muscles; but the strongest of these Slips arises with diverging Fibres from the inferior anterior Spinous Process of the Os Ilium. Tab. LV. Fig. 1. m, n, o.

The Capsular Ligament allows the Thigh-Bone to be moved to every side, and when its Body is moved forwards or backwards, a small degree of rotation is performed round the Cervix of the Bone.

The *Internal*, commonly called the *Round Ligament*, which arises by a broad flat beginning from the under and inner part of the Cavity of the Acetabulum, and is connected with the Substance termed *Gland of the Joint*. From this it runs backwards and a little upwards, becoming gradually narrower and rounder, to be fixed to the Pit upon the inner Surface of the Ball of the Os Femoris. Tab. LV. Fig. 2. g—k.

The Round Ligament prevents the Bone from being dislocated upwards or inwards, and assists also in agitating the Mucous Substance within the Joint.

A *Cartilaginous Ligament* surrounding the Brim of the Acetabulum, and thereby increasing the depth of that Cavity for the reception of the Head of the Thigh-Bone. Tab. LV. Fig. 2. c.

A *Double Cartilaginous Ligament*, Tab. LV. Fig. 3. d, stretched from one end of the Breach, in the under and fore part of the Acetabulum, to the other, but leaving a Hole behind it for containing part of the Substance called *Gland of the Joint*, and for the passage of the Vessels of that Substance.

This Ligament allows the Thigh-Bone to be moved inwards, and the Glandular-looking Substance to be agitated with safety.

The Substance called *Gland of the Joint*, covered with a Vascular Membrane, and lodged in a Depression in

the under and inner part of the Acetabulum. Tab. LV. Fig. 2. m.

At the edges of this Substance, *Fringes* are sent out, which furnish part of the Synovia for the lubrication of the Joint.

The edges of this Substance are fixed to those of the Pit in the Acetabulum, by small Ligamentous Bridles, termed *Ligamenta Mucosa*, or *Ligamentula Massæ Adiposo-Glandulosæ*.

LIGAMENTS, &c. of the JOINT of the KNEE.

The *Lateral Ligaments* which lie at the sides of the Joint, and adhere to the outer Surface of the Capsular Ligament.

The *Internal Lateral Ligament*, which is of considerable breadth, arising from the upper part and Tubercle of the internal Condyle of the Os Femoris, and inserted into the upper and inner part of the Tibia; the Fibres passing obliquely forwards, till they have reached a little below the Head of the Bone. Tab. LV. Fig. 4. k.

The *Long External Lateral Ligament*, which is narrower, but thicker and stronger than the former, arising from the Tubercle above the external Condyle of the Os Femoris, and fixed to the Fibula, a little below its Head. Tab. LV. Fig. 5. f.—Tab. XXXII. Fig. 8. l.

Behind the long external Lateral Ligament, there is an *Expansion* attached nearly in the same manner as this Ligament, which has been termed the *Short External Lateral Ligament*. Tab. LV. Fig. 5. g.

These Ligaments are placed a little behind the middle of the Joint, in consequence of which, when the Leg is extended, they are stretched, and thereby prevent lateral and rotatory motion; but when the Knee is bent, they are relaxed, and then admit of a small degree of rotation and motion to either side.

The *Posterior Ligament* of WINSLOW, Tab. LV. Fig. 5. b, formed of irregular Bands which arise from the upper and back part of the external Condyle of the Os Femoris, and descend obliquely over the Capsular Ligament, to be fixed to the Tibia under the inner and back part of its Head. It prevents the Leg from being pulled farther forwards than to a straight line with the Thigh, and also furnishes a convenient situation to the beginnings of the Gastrocnemius and Plantaris.

When this Ligament is wanting, which is sometimes the case, its place is supplied by a *Membranous Expansion*.

The *Ligament of the Patella*, which arises from a Depression behind the Apex of that Bone, and is fixed to the Tuberosity of the upper and fore part of the Tibia. Tab. XXXIV. Leg, b.—Tab. XXXII. Fig. 9. i. By the

the intervention of this Ligament, the Muscles inserted into the Patella are enabled to extend the Leg.

The *Capsular Ligament*, which arises from the whole Circumference of the under end of the Thigh-Bone, some way above the Margin of the Articulating Cartilage, and above the posterior part of the Great Notch between the Condyles. From this it descends to be fixed round the Head of the Tibia, and into the whole Margin of the Articulating Surface of the Patella, in such a manner that the Patella forms part of the Capsule of the Joint. Tab. LV. Fig. 6. 7.

The Capsular Ligament is of itself remarkably thin, but so covered by the Ligaments already mentioned, by the general Aponeurosis of the Limb, and by the Tendons of Muscles which surround the Joint, as to acquire a considerable degree of strength.

The Capsular Ligament, along with the other Ligaments of this Joint, admits of the flexion and extension of the Leg, but of no lateral nor rotatory motion in the extended state, though of a small degree of each when the Limb is fully bent.

The *Ligamentum Alare, Majus et Minus*, which are Folds of the Capsular Ligament, running like *Wings* at the sides of the Patella, to which, and to the Fatty Substance of the Joint, they are attached. Tab. LV. Fig. 6. e, d.

The *Ligamentum Mucosum*, continued from the joining of the Ligamenta Alaria to the Os Femoris, immediately above the Anterior Crucial Ligament. It preserves the Synovial Substance in its proper place, during the various motions of the Joint. Tab. LV. Fig. 6. c.

The *two Crucial or Internal Ligaments*, which arise from the Semilunar Notch between the Condyles of the Os Femoris, and decussate each other within the Cavity of the Joint. Tab. XXXII. Fig. 9. f, f.

The *Anterior Crucial Ligament* runs downwards and forwards, to be fixed to a Pit before the rough Protuberance in the middle of the Articulating Surface of the Head of the Tibia. Tab. LV. Fig. 8. e.

The *Posterior Crucial Ligament* descends to be fixed to a Pit behind the above-mentioned rough Protuberance. Tab. LV. Fig. 8. f.—Tab. XXXII. Fig. 8. b.

These Ligaments, in the extended state of the Leg, prevent it from going forwards beyond a straight Line with the Thigh. When the Knee is bent, they allow the Foot to be turned outwards, but not in the contrary direction.

The *two Inter-articular Cartilages*, called *Semilunar* from their shape like a Crescent, placed upon the top of the Tibia. Tab. LV. Fig. 9. c, d.—Tab. XXXII. Fig. 7. c, c.

The outer convex edge of each of these Cartilages is thick, while the inner concave edge becomes thin and sharp like a knife or sickle, and being concave above, the Sockets for the Condyles of the Os Femoris are rendered deeper, and this Bone and the Tibia are more accurately adapted to each other.

Each of these Cartilages is broad in the middle, their extremities becoming narrower and thinner as they approach each other. Each covers about two thirds of the Superficial Cavity of the top of the Tibia, leaving one third bare in the middle. The extremities are termed *Cornua*, and are fixed by Ligaments to the Protuberance of the Tibia. The anterior Cornua are joined to each other by a *Transverse Ligament*. Tab. LV. Fig. 9. b.

The convex edge of these Cartilages is fixed to the Capsular and other Ligaments, in such a manner as to allow them to play a little upon the Cartilaginous Surface of the Tibia, by which the motions of that Bone upon the Condyles of the Os Femoris are facilitated.

The *Mucous or Fatty Substances* of this Joint, which are the largest of any in the Body, are situated in the different interstices of the Joint, but chiefly round the edges of the Patella. Tab. LV. Fig. 13. b, b, b. They are covered by a fine Membrane reflected from the inner Surface of the Capsular Ligament.

Fimbriae project from the edges of these Fatty Substances, which discharge Synovia for the lubrication of the Joint. Tab. LV. Fig. 13. c, c.

LIGAMENTS connecting the FIBULA to the TIBIA.

The *Capsular Ligament* of the superior extremity of the Fibula, which ties it to the outer part of the Head of the Tibia, and which is strengthened by the external Lateral Ligament of the Knee, and by the Tendon of the Biceps which is fixed to the Fibula. Tab. LV. Fig. 7. i. Fig. 8. o.

The *Interosseous Ligament*, one edge of which is fixed to the Ridge or Angle at the outer and back part of the Tibia, the other to the corresponding Ridge at the inner side of the Fibula. It fills the space between the Tibia and Fibula, like the Interosseous Ligament of the Fore-Arm, and is of a similar structure, being formed of oblique Fibres, and perforated in various places for the passage of Vessels and Nerves.

At its upper part there is a large Opening, where the Muscles of the opposite sides are in contact, and where Blood-vessels pass to the fore-part of the Leg.

It serves chiefly for the Origin of part of the Muscles which belong to the Foot, and thereby supplies the place of Bone.

The *Ligaments* of the inferior extremity of the Fibula, which are called *Anterior-superior*, and *Posterior-superior*, according to their situations, arising from the edges of the Semilunar Cavity of the Tibia, and fixed to the Malleolus Externus of the Fibula. Tab. LV. Fig. 10. b. Fig. 11. e.

The Ligaments between the ends of the Tibia and Fibula join the two Bones so firmly together, as to admit of no sensible motion.

LIGAMENTS connecting the BONES of the TARSUS with those of the LEG.

The *Anterior Ligament of the Fibula*, which arises from

from the anterior part of the Malleolus Externus, and passes obliquely forwards, to be fixed to the upper and outer part of the Astragalus. Tab. LV. Fig. 10. *k*.

The *Middle*, or *Perpendicular Ligament* of the Fibula, which arises from the point of the Malleolus Externus, and descends almost perpendicularly, to be inserted into the outside of the Os Calcis. Tab. LV. Fig. 10. *i*.

The *Posterior Ligament of the Fibula*, which arises from the under and back part of the Malleolus Externus, and runs backwards, to be joined to the outer and posterior part of the Astragalus. Tab. LV. Fig. 11. *b*.

The *Ligamentum Deltoides* of the Tibia, which arises from the Malleolus Internus, and descends in a radiated form, to be attached to the Astragalus, Os Calcis, and Os Naviculare. Tab. LV. Fig. 11. *f*.

The *Capsular Ligament*, which lies within the former Ligaments, and is remarkably thin, especially before and behind, for the readier motion of the Joint. It arises from the Margin of the Articular Cavity of the Tibia and Fibula, and is fixed round the edge of the Articular Surface of the Astragalus.

The Ligaments and the other constituent parts of the Ankle-joint form it into a complete Hinge, which allows flexion and extension, but no rotation or lateral motion, in the bended state of the Foot, though a small degree of each when it is fully extended.

LIGAMENTS of the TARSUS.

The *Capsular Ligament*, which joins the Articular Surface of the Os Calcis to that of the Astragalus.

A number of *Short Ligaments*, lying in the Fossa of the Astragalus and of the Os Calcis, and forming the *Ligamentous Apparatus* of the Sinuous Cavity, which assists in fixing the two Bones strongly together. Tab. LV. Fig. 10. *l, m*.

The *Capsular*, the *Broad Superior*, and the *Internal Lateral Ligaments*, connecting the Astragalus to the Os Naviculare, and admitting of the lateral and rotatory motions of the Foot. Tab. LV. Fig. 10.

The *Superior*, the *Lateral*, and the *Inferior Ligaments*, fixing the Os Calcis to the Os Cuboides, where a small degree of motion is allowed to every side. The inferior Ligaments consist of a *Long*, an *Oblique*, and a *Rhomboid Ligament*, which are the longest and strongest of the Sole. Tab. LV. Fig. 12.

The *Superior Superficial*, the *Interosseous*, and the *Inferior Transverse Ligaments*, which fix the Os Naviculare and Os Cuboides to each other. Tab. LV. Fig. 10. Fig. 12.

The *Superior Lateral*, and *Plantar Ligaments*, which fix the Os Naviculare to the Os Cuneiforme. Tab. LV. Fig. 10. Fig. 12.

The *Superior Superficial*, and the *Plantar Ligaments*, which connect the Os Cuboides to the Os Cuneiforme Externum. Tab. LV. Fig. 10. Fig. 12.

VOL. I.

The *Dorsal* and *Plantar Ligaments*, which unite the *Ossa Cuneiformia* to each other. Tab. LV. Fig. 10. Fig. 12.

Besides the Capsular Ligaments of the Tarsus already mentioned, each of the other Joints of these Bones is furnished with its *proper Capsular Ligament*.

From the strength of the Ligaments which unite these Bones to each other, and from the plainness of their articulating Surfaces, no more motion is allowed than to prevent the effects of concussion in walking, leaping, &c.

LIGAMENTS between the TARSUS and METATARSUS.

The Bones of the Metatarsus are fixed to those of the Tarsus by *Capsular*, and numerous other *Ligaments*, which are called *Dorsal*, *Plantar*, *Lateral*, according to their situations;—and *Straight*, *Oblique*, or *Transverse*, according to their directions. The nature of this Joint is the same with that between the Carpus and Metacarpus. Tab. LV. Fig. 10. Fig. 12.

LIGAMENTS connecting the METATARSAL BONES to each other.

The *Dorsal*, *Plantar*, and *Lateral Ligaments*, which connect the Bases of the Metatarsal Bones with each other. Tab. LV. Fig. 10. Fig. 12.

The *Transverse Ligaments*, which join the Heads of these Bones together.

LIGAMENTS of the PHALANGES of the TOES.

The *Capsular* and *Lateral Ligaments*;—resembling those of the Fingers.

LIGAMENTS and SHEATHS retaining the TENDONS of the MUSCLES of the FOOT and TOES, in situ.

The *Annular Ligament of the Tarsus*, which is a thickened part of the Aponeurosis of the Leg, splitting into superior and inferior portions, which bind down the Tendons of the Extensors of the Toes, upon the fore-part of the Ankle. Tab. XXXIV. Fig. 2. N, N.

The *Vaginal Ligament of the Tendons of the Peronei*, which, behind the Ankle, is common to both, but at the outer part of the Foot becomes proper to each, Tab. LII. Fig. 2 *p, q*; preserving, each, its respective Tendon in its proper place, and forming the *Bursa* of that Tendon.

The *Lacinated Ligament*, which arises from the inner Ankle, and spreads in a radiated manner, to be fixed partly in the Cellular Substance and Fat, and partly to the Os Calcis, at the inner side of the Heel. Tab. XXXIV. Fig. 2. *k*. It incloses the Tibialis Posticus and Flexor Digitorum Longus.

The *Vaginal Ligament of the Tendon of the Extensor Proprius*

Z

Proprius Pollicis, which runs in a Crucial Direction. Tab. XXXVIII. Right Foot, C.

The *Vaginal Ligament of the Tendon of the Flexor Longus Pollicis*, which furrounds this Tendon in the Hollow of the Os Calcis. Tab. LIV. Fig. 3. H.

The *Vaginal and Crucial Ligaments of the Tendons of the Flexors of the Toes*, which inclose these Tendons on the Surfaces of the Phalanges, and form their Bursæ

Mucosæ. Tab. LIV. Fig. 3. Q, R, S, T, U.—Tab. L. Fig. 8.

The *Accessory Ligaments of the Flexor Tendons of the Toes*, which, as in the Fingers, arise from the Phalanges, and are included in the Sheaths of the Tendons in which they terminate.

The *Transverse Ligaments of the Extensor Tendons*, which run between these Tendons, and preserve them in their places behind the roots of the Toes.

TABLE




Fig. 3.




Fig. 2.



Fig. 4.




Fig. 7.



Fig. 6.




Fig. 1.




Fig. 5.




Fig. 8.




Fig. 9.



Fig. 10.




Fig. 12.




Fig. 11.




Fig. 13.




Fig. 14.



T A B L E LIII.

LIGAMENTS of the BONES of the HEAD and TRUNK.

FIG. 1.

LIGAMENTS of the LOWER JAW, and adjacent Parts,
viewed on the Right Side.

- a*, The angle of the lower jaw ;
- b*, Its condyloid process.
- c*, The edge of the zygomatic process of the temporal bone.
- d*, A section of the ear.
- e*, The styloid process.
- f*, The capsular ligament of the lower jaw.
- g*, The lateral ligament.
- h*, The connection of the suspensory ligaments of the os hyoides, and of the stylo-glossus, to the styloid process.
- i*, The suspensory ligament of the os hyoides.
- k*, The suspensory ligament of the stylo-glossus.
- l*, The masseter.
- m*, The stylo-glossus.
- n*, The stylo-hyoides.
- o*, The stylo-pharyngeus, upon which
- p*, A small nerve rests in its way to the pharynx.
- q*, A section of the digastricus.
- r*, The sterno-mastoideus.
- s*, The lingual nerve.

FIG. 2.

The CAPSULAR LIGAMENT of the LOWER JAW, and Part
of the TEMPORAL BONE.

- a*, The zygomatic process of the temporal bone.
- b*, The tubercle of that process.
- c*, The glenoid or articular cavity.
- d, d*, The capsular ligament surrounding the whole margin of the glenoid cavity.

FIG. 3.

An Anterior View of the CONNECTION of the Os OCCIPITIS
with the First and Second VERTEBRÆ.

- a*, A portion of the os occipitis.
- b, b*, The transverse processes of the first vertebra ;

- c, c*, Those of the second.
- d*, The capsular membrane of the last joint of the occipital bone.
- e*, The membrane of the right joint cut open, to obtain a view of the joint, and,
- f*, The distance of the origin of the membrane.
- g*, The membrane which fills up the anterior opening between the occiput and first vertebra.
- h*, A slip inserted into the middle of this membrane.
- i*, The origin of this slip in the occipital bone.
- k*, Its termination in the tubercle of the first vertebra.
- l*, The slip which unites the first and second vertebra.
- m*, The ligament proper to the first vertebra.
- n*, The ligament of the articulation of the first vertebra with the second.

FIG. 4.

LIGAMENTS on the Fore-Part of the SPINE, and Upper
and Fore Part of the Inner Side of the Bones of the
PELVIS.

- a, b*, The bodies of the dorsal vertebræ.
- c, d*, ————— lumbar vertebræ.
- e, f*, A section of the ribs.
- g, h*, The anterior common ligament of the bodies of the vertebræ.
- i, i*, The crura of the diaphragm.
- k, k*, Part of the longi colli.
- l, l*, Two of the transverse processes of the lumbar vertebræ.
- m*, The psoas magnus.
- n*, The quadratus lumborum.
- o*, The os ilium.
- p*, The superior transverse ligament of the pelvis, of a triangular form.
- q*, The inferior transverse ligament, of a round form.
- r*, The longitudinal ligamentous fibres, belonging to,
- s*, The symphysis of the ilium with the os sacrum.
- t*, The intestinum rectum.
- u*, The vesica urinaria.
- v*, A ligament which forms a sort of sac between the sides of the bladder and rectum.

FIG. 5.

The CRUCIAL INTERVERTEBRAL LIGAMENTS of two of the LUMBAR VERTEBRÆ, seen Anteriorly.

- a, The first,
- b, The second lumbar vertebra.
- c, d, The crucial intervertebral ligament, formed of different strata.

FIG. 6.

The Posterior, or Internal Common LIGAMENT of the VERTEBRÆ, seen in the NECK and beginning of the BACK. The CRURA of the SPINOUS PROCESSES are removed.

- a, a, &c. The transverse processes of the vertebræ.
- b, b, The vertebral arteries.
- c, c, &c. Sections of the crura of the spinous processes.
- d, d, The whole breadth of the posterior or internal common ligament of the vertebræ.

FIG. 7.

The CONTINUATION of the Posterior Common LIGAMENT in the LOINS.

- a, The first, and,
- b, The second lumbar vertebra.
- c, c, Vestiges of the crura of the spinous processes.
- d, d, The posterior, or internal common ligament of the vertebræ.
- e, e, Its expansion over the cartilaginous interstices.

FIG. 8.

Three of the DORSAL VERTEBRÆ, with Portions of the three corresponding RIBS and their LIGAMENTS, seen within the THORAX.

- a, a, a, The bodies of the three vertebræ.
- b, b, b, Portions of the three ribs.
- c, c, c, The ligaments which fix the heads of the ribs to the bodies of the vertebræ.
- d, d, d, The internal ligaments of the cervix of the ribs, by which they are fixed to the tuberosities of the next superior transverse processes.
- e, e, Part of the intercostal muscles.
- f, The spinous process.

FIG. 9.

The same Parts shewn in the preceding Figure, but viewed Posteriorly.

- a, a, a, The crura of the spinous processes.
- b, b, b, Portions of the three ribs.

- c, c, c, The external transverse ligaments.
- d, d, The internal ligaments of the cervix of the ribs.
- e, e, The external ligaments of the cervix of the ribs.
- f, f, A ligamentous cord between the apices of the spinous processes.

FIG. 10.

LIGAMENTS on the Outer and Right Side of the PELVIS.

- a, The os sacrum.
- b, ——— coccygis.
- c, ——— ilium.
- d, The tuber of the os ischium.
- e, The notch of the os ilium, which, together with the ligaments, &c. under it, form the foramen magnum.
- f, The posterior long ileo-sacral ligament.
- g, The posterior short ileo-sacral ligament.
- h, The posterior lateral ileo-sacral ligament.
- i, i, The ligamenta accessoria vasa, on the back of the os sacrum.
- k, The large sacro-ischiatic ligament.
- l, The appendix, or superior membranous production.
- m, The small sacro-ischiatic ligament.
- n, The longitudinal ligaments of the os coccygis.
- o, p, The origin of some of the muscles of the thigh.

FIG. 11.

LIGAMENTS seen in the LEFT CAVITY of the PELVIS; the RIGHT OS INNOMINATUM being removed.

- a, The os sacrum.
- b, ——— coccygis.
- c, ——— ilium.
- d, ——— pubis.
- e, ——— ischium.
- f, The cartilaginous surface of the os sacrum, for the articulation with the os ischium.
- g, A protuberance covered with ligamentous villi for the articulation with the os ischium.
- h, The linea alba, which marks the articulation of the os ilium and sacrum.
- i, The spinous process of the os ischium.
- k, The small internal sacro-ischiatic ligament.
- l, The remains of the coccygeus.
- m, A portion of the large external sacro-ischiatic ligament.
- n, n, The inferior falciform production of WINSLOW.
- o, The foramen magnum, for the transmission of the pyriformis, &c.
- p, The foramen minus for the transmission of the obturator internus.
- q, The superior oblique sinus of the foramen thyroideum.
- r, The membrana obturans.

s, The

- s*, The transverse ligament of the membrana obturans.
t, The tendon of the psoas magnus.
u, The cartilago-ligamentous substance of the os pubis.

FIG. 12.

LIGAMENTS connecting the CARTILAGES of the RIBS, and
 CARTILAGO ENSIFORMIS, to the Body of the STER-
 NUM.

- a*, The upper end of the sternum.
b, The cartilago ensiformis.
c, The cartilage of the eighth rib.
d, d, The ligaments of the cartilages of the ribs, distributed in the form of radii.
e, e, The intercostal muscles.
f, f, The tendinous ligaments of the cartilages of the ribs.
g, g, The ligaments of the cartilago ensiformis.

FIG. 13.

Represents the Connection of the Osseous Part of a RIB
 with its CARTILAGE, and of the CARTILAGE with the

STERNUM; the INVESTING MEMBRANES being removed
 Anteriorly.

- a*, The osseous,
b, The cartilaginous part of the rib.
c, A part of the sternum.
d, Short ligamentous fibres connecting the margin of the osseous and cartilaginous parts of the rib.
e, The capsular ligament which joins the cartilage of the rib to the sternum; the external radiated fibres of this joint being removed.

FIG. 14.

A SECTION of the Parts represented above, to shew their
 Internal Structure.

- a*, The cancelli of the rib.
b, ————— sternum.
c, The connection of the inner part of the rib with its cartilage by a firm union of substance.
d, The cavity of the joint formed between the cartilage and sternum.

TABLE

T A B L E L I V.

LIGAMENTS of the LEFT SUPERIOR EXTREMITY.

FIG. 1.

LIGAMENTS of the Upper Part of the STERNUM, viewed Anteriorly.

- a, a*, Part of the clavicles.
- b*, The inter-clavicular ligament.
- c, c*, The insertions of the sterno-mastoidei into the clavicles.
- d, d*, The cartilages of the first ribs.
- e*, A section of the sternum.
- f, f*, Ligaments connecting the clavicles to the inter-clavicular ligaments, to the sternum, and to the cartilages of the first ribs.
- g, g*, The ligamentum rhomboides, on each side, connecting the clavicle to the cartilage of the first rib.

FIG. 2.

A View of the ARTICULATION between the STERNUM and CLAVICLES.

- a*, A section of the sternum.
- b*, The cartilage of the first rib.
- c*, The head of the right clavicle turned back.
- d*, The inter-clavicular ligament.
- e*, The articular sinus of the sternum.
- f, f*, The inter-articular cartilage covering the articular sinus and head of the clavicle.
- g*, The part to which the head of the clavicle is fixed.
- h*, The prolongation of the inter-articular cartilage.
- i, i*, The capsular ligament.

FIG. 3.

An Anterior View of the LIGAMENTS between the CLAVICLE, SCAPULA, and OS HUMERI.

- a*, The upper part of the scapula.
- b*, The point of the acromion.
- c*, ————— coracoid process.
- d*, A portion of the clavicle.
- e*, ————— os humeri.
- f*, The edge of the anterior proper ligament of the scapula.
- g, g*, The posterior proper ligament of the scapula.
- h*, A portion of the common conoid ligament.
- i, i*, The capsular ligament of the head of the os humeri.
- k*, The accessory membrane of the capsular ligament.
- l, l*, The edge of the oval hole in the capsular membrane, for the passage of,
- m*, The tendon of the sub-scapularis, which fills the hole, and makes part of the capsule.

- n*, The outer edge of the fleshy portion of this muscle.
- o*, A ligament which arises from the sheath of the sub-clavius.
- p*, The remains of the tendon of the pectoralis.
- q*, The tendon of the biceps.

FIG. 4.

A Posterior View of the LIGAMENTS between the CLAVICLE, SCAPULA, and OS HUMERI.

- a*, The dorsum ;
- b*, The spine ;
- c*, The upper edge ;
- d*, The cervix, and,
- e*, The acromion of the scapula.
- f*, A section of the clavicle.
- g*, ————— os humeri.
- h*, Ligaments connecting the clavicle with the acromion.
- i, i*, The posterior proper ligament of the scapula.
- k*, The common conoid ligament of the scapula.
- l, l*, The capsular ligament inclosing the head of the os humeri.

FIG. 5.

LIGAMENTS between the SCAPULA, CLAVICLE, and OS HUMERI, viewed Anteriorly.—Some of the Parts are twisted to one side.

- a*, The upper edge of the scapula, twisted to one side.
- b*, The point of the acromion.
- c*, ————— coracoid process.
- d*, A section of the clavicle.
- e*, ————— os humeri.
- f*, The anterior proper triangular ligament of the scapula.
- g*, The common trapezoid ligament of the scapula.
- h*, Ligaments connecting the clavicle with the acromion.
- l*, The capsular ligament of the head of the os humeri ;
- m*, Its appendix.
- n*, The edge of the oval hole, for the transmission of,
- o*, The tendon of the sub-scapularis.
- p*, The remains of the tendon of the supra-spinatus.
- q*, The tendon of the biceps.
- r*, Part of the sheath of the tendon.







FIG. 6.

Gives a View of the GLENOID CAVITY of the SCAPULA, with the ORIGIN of the CAPSULAR LIGAMENT of this JOINT.

- a*, The glenoid cavity.
- b, b*, The cartilaginous brim of the cavity.

c, The

TAB. 54.



- c*, The origin of the long head of the biceps.
d, d, The inner surface of the capsular ligament.
e, The edge of the oval hole.

FIG. 7.

The HEAD of the Os HUMERI, with a Portion of the CAPSULAR LIGAMENT which incloses it.

- a*, The head of the os humeri.
b, The tendon of the long head of the biceps continued through its sheath.
c, The retinaculum in the beginning of the sheath of the tendon of the biceps.
d, d, The inner surface of the capsular ligament.
e, A fimbriated organ for the secretion of synovia, situated immediately under the ball of the os humeri, within the capsular ligament.

FIG. 8.

The CONNECTION of the Os HUMERI and BONES of the FORE-ARM, and of the latter with each other; viewed Anteriorly.

- a*, A portion of the os humeri.
b, The inner,
c, The outer round process of articulation.
d, The inner acute prominent process of articulation between the os humeri and bones of the fore-arm.
e, The ulna.
f, The olecranon of the ulna.
g, The coronoid process.
h, The inferior head of the ulna.
i, The radius.
k, The inferior extremity of the radius.
l, l, The capsular ligament.
m, The internal lateral ligament.
n, The coronary, annular, or orbicular ligament of the radius.
o, The accessory ring.
p, The posterior accessory ligament.
q, The chorda transversalis cubiti.
r, r, The interosseous membrane or ligament of the fore-arm.
s, An opening left by this membrane, and filled with muscles.

FIG. 9.

The CONNECTION of the Os HUMERI with the BONES of the FORE-ARM, viewed Posteriorly.

- a*, A portion of the os humeri;
b, Its external condyle.
c, The outer round process of articulation.
d, A portion of the ulna.
e, The olecranon.
f, The lateral surface of the olecranon, covered by the anconeus.
g, A portion of the radius.
h, The external lateral ligament.

- i*, The orbicular ligament of the radius.
k, The accessory ring.
l, The posterior accessory ligament.

FIG. 10.

The CONNECTION of the Under End of the RADIUS with that of the ULNA.

- a*, The extremity of the radius.
b, b, The double glenoid cavity of the radius.
c, The intermediate triangular cartilage between the ulna and os cuneiforme.
d, The extremity of the ulna.
e, The ligament called Sacciform capsular membrane.
f, f, The cut edge of the capsular ligament between the fore-arm and wrist.
g, The mucous ligament within the joint.

FIG. 11.

A View of the CAVITY of the JOINT between the FORE-ARM and WRIST; the CAPSULAR LIGAMENT being cut Posteriorly, and the BONES of the CARPUS turned back.

- a*, The extremity of the radius.
b, ————— ulna.
c, Part of the glenoid cavity of the radius, receiving,
d, The first bone of the carpus.
e, The other portion of the glenoid cavity of the radius, and,
f, The intermediate triangular cartilage, receiving,
g, The second, and,
h, The third bone of the carpus.
i, The capsular ligament between the bones of the fore-arm and wrist.
k, The palmar slip intermixed with this membrane.
l, The palmar accessory ligament from the radius;
m, That from the fourth, and,
n, That from the third bone of the carpus.
o, The mucous ligament.
p, The ligament by which the first and second bone of the carpus are attached.

FIG. 12.

Shows certain LIGAMENTS on the PALM-SIDE of the HAND, after the INTEGUMENTS and TENDONS of the EXTENSORS have been removed.

- a*, The extremity of the radius.
b, ————— ulna.
c, The os pisiforme, with the tendon of the flexor carpi ulnaris fixed to it.
d, The os trapezium, placed at the root of the thumb.
e, The sharp process of the unciform bone.
f, The metacarpal bone of the thumb.
g, g, The metacarpal bones of the fingers.
h, h, The first phalanx of the fingers.
i, The tendon of the flexor carpi radialis.
k, The tendon of one of the extensors of the thumb.

l, l, l, The

- l, l, l,* The internal interosseous muscles.
m, The ligaments between the ulna and os pisiforme.
n, o, p, q, The capsular ligament investing the extremities of the bones of the fore-arm and bones of the wrist, intermixed with numerous accessory slips.
r, s, Ligaments between the carpal and metacarpal bones.
t, Ligaments joining the bases of the metacarpal bones to each other.
u, u, u, Ligaments connecting the heads of the metacarpal bones to each other.

FIG. 13.

A View of the LIGAMENOUS BANDS which assist in forming the CAPSULAR LIGAMENT on the Back Part of the WRIST.

- a,* The extremity of the radius.
b, ————— ulna.
c, The first,
d, The third, and,
e, The fifth bone of the carpus.
f, The tuber of the seventh carpal bone.
g, The two tendons of the extensores carpi radiales.
h, The tendon of the extensor carpi ulnaris.
i, The ligamentum rhomboides.
k, A ligamentous cord, which extends from the styloid process of the ulna to the third bone of the carpus.
l, The common oblique slip.
m, The ligament between the third and eighth carpal bone.
n, A section of the metacarpal bones.

FIG. 14.

The Anterior LIGAMENTS which bind the BONES of the FORE-ARM to the BONES of the CARPUS, these to each other, and also to the BONES of the METACARPUS.

- a,* The extremity of the radius.
b, ————— ulna.
c, c, The bones of the carpus.
d, d, d, d, The bones of the metacarpus.
e, The capsular ligament of the wrist, with its accessory slips.
f, The anterior annular ligament of the carpus. Besides the annular ligament, numerous ligamentous slips are seen running in various directions, which unite the bones of the carpus to each other, and likewise to those of the metacarpus.
g, h, Two of the interosseous ligaments, which connect the bases of the last-mentioned bones.

FIG. 15.

The CAVITY of the JOINT in the middle of the CARPUS; the CAPSULAR LIGAMENT being divided in the Back of the HAND, and the two Rows of CARPAL BONES separated from each other.

- a, b, c, d,* The first row, and,
e, f, g, h, The second row of the carpal bones.
i, The inside of the palmar part of the capsular ligament.
k, The inside of the dorsal part of that ligament.
l, A ligament joining the first and second carpal bones together, where they are articulated with the end of the radius.
m, A ligament connecting the third and seventh carpal bones to each other.
n, The ligamentum mucosum connected with the second carpal bone.
o, A small frænum of the capsular ligament.

FIG. 16.

LIGAMENTS between the Inferior Row of the CARPAL BONES and those of the METACARPUS, on the Back of the HAND.

- a, b, c, d,* The four bones of the second row of the carpus.
e, The metacarpal bone of the thumb.
f, f, The metacarpal bones of the fingers.

The ligaments which unite these bones are seen running in various directions.

FIG. 17.

The LIGAMENTS of the JOINTS of one of the FINGERS.


- a,* The metacarpal bone.
b, c, d, The first, second, and third bones of the fingers.
e, e, e, The lateral ligaments of the joints of the fingers.
f, Part of the capsular ligament.

FIG. 18.

The MIDDLE FINGER, with the ACCESSORY LIGAMENTS which bind the TENDONS of the PERFORANS and PERFORATUS.

- a,* A section of the vaginal ligaments.
b, The tendon of the perforatus.
c, The end of this tendon inserted into the second bone of the finger.
d, Decussating branches of the tendon of the perforatus.
e, The tendon of the perforans.
f, The short accessory ligament of the perforans.
g, h, Long accessory ligaments of the tendon of the perforans.

The tendon of the perforatus has accessory ligaments similar to those of the perforans, but they could not be represented in this view.



TAB. 55.





Fig. 6.




Fig. 7.



Fig. 8.




Fig. 5.



Fig. 9.




Fig. 11.



Fig. 12.



Fig. 10.



T A B L E LV.

LIGAMENTS of the LEFT INFERIOR EXTREMITY.

FIG. 1.

A View of the CAPSULAR LIGAMENT of the HEAD of the Os FEMORIS, and of the LIGAMENTS filling the FORAMEN THYROIDEUM.

- a*, The spine of the right os ilium ;
 - b*, Its inferior spinous process.
 - c*, The os pubis.
 - d*, ——— femoris.
 - e*, The trochanter major.
 - f*, ——— minor.
 - g*, The place of the insertion of the pectineus.
 - h, i*, The seat of the under end of the iliacus internus.
 - k*, The seat of the gluteus minimus.
 - l*, The upper end of the rectus femoris cut and turned back.
- The space between *m* and *k* is occupied by the capsular ligament, which arises from the acetabulum.
- m, n, o*, The accessory slips of the capsular ligament, placed between the surrounding muscles, and strengthening the general capsule.
 - p*, The oblique termination of the capsular ligament between the two trochanters.
 - q, r, s, u, x*, The membrane which shuts up the foramen thyroideum.
 - r, s*, The external and internal ligamentous slips of this membrane, which form fulci.
 - t*, The place where the obturator externus adheres to the capsular ligament.
 - u*, An opening in the upper part of the obturator ligament, for the passage of the obturator vessels and nerve.

FIG. 2.

The HEAD of the Os FEMORIS taken out of the ACETABULUM, and still adhering by means of the LIGAMENTUM ROTUNDUM.

- a*, The ball of the os femoris.
- b*, The cavity of the acetabulum.
- c, c*, The cartilaginous brim of the acetabulum.
- d, d, d, d*, The capsular ligament cut and turned back, to shew,
- e, e*, Its density and thickness.
- f*, Retinacula reflected about the cervix of the os femoris.
- g, h, i, k*, The ligamentum rotundum.
- l, l*, Small ligaments which bind the fatty glandular mass *m* to the bottom of the acetabulum.

VOL. I.

FIG. 3.

Gives a View of the ACETABULUM, with its LIGAMENTS.

- a*, The bottom of the acetabulum, with its cartilaginous surface.
- b*, The sinus for the synovial fat.
- c, c, c*, The cartilaginous brim of the acetabulum.
- d, d*, The external transverse ligament.

FIG. 4.

A Side-View of the KNEE, shewing the Internal Lateral LIGAMENT.

- a*, The os femoris.
- b*, The patella.
- c*, The inner side of the head of the tibia.
- d*, A portion of the vastus internus.
- e*, Aponeurosis of the vastus internus.
- f*, The remains of the inner tendon of the gastrocnemius.
- g*, The tendon of the semi-membranosus.
- h*, The tendons of the gracilis and semi-tendinosus, turned back.
- i*, The popliteus.
- k, k*, The broad internal lateral ligament, enlarged by,
- l*, An accessory branch.
- m*, A circular ligamentous margin adhering to the internal femilunar cartilage.

FIG. 5.

A Posterior View of the LIGAMENTS of the KNEE.

- a*, The os femoris.
- b*, The external condyle.
- c*, The internal condyle.
- d*, The tibia.
- e*, The fibula.
- f*, The long external lateral ligament.
- g*, The short external lateral ligament.
- h*, The ligamentum posticum Winsloii.
- i*, The tendon of the semi-membranosus.
- k*, Irregular membranous expansions.

FIG. 6.

An Anterior View of the JOINT of the KNEE, [the CAPSULAR LIGAMENT being opened, and turned down along with the PATELLA.

- a, a*, The condyles of the os femoris covered with cartilage.
- b*, The

A a

- b*, The patella, with the capsular ligament at the sides of it, cut from the os femoris, and turned down.
c, The ligamentum mucosum, supporting the fat at the under edge of the patella.
d, e, Folds of the inner side of the capsular ligament, called by WEITBRECHT, *Ligamentum Alare externum minus*, and *Ligamentum Alare internum majus*.
f, g, The anterior edges of the femilunar cartilages.
b, Part of the posterior crucial ligament.

FIG. 7.

A View of the CRUCIAL LIGAMENTS, as seen in the back Part of the JOINT of the KNEE; the CAPSULAR LIGAMENT being laid open.

- a*, The os femoris;
b, c, Its condyles: above which is seen the cut edge of the capsular ligament.
d, The tibia.
e, The fibula.
f, The posterior crucial ligament.
g, The insertion of the anterior crucial ligament into the os femoris.
h, The edge of the external femilunar cartilage.
i, Ligamentous fibres strengthening the joint at the head of the fibula.

FIG. 8.

Anterior View of the CRUCIAL LIGAMENTS.

- a, b*, The condyles of the os femoris.
c, The tibia.
d, The fibula.
e, The anterior crucial ligament.
f, f, The insertion of the posterior crucial ligament into the os femoris.
g, The ligament of the posterior cornu of the external femilunar cartilage, connected with the posterior crucial ligament, and with it fixed to the os femoris.
h, The ligament of the anterior cornu of the external femilunar cartilage.
i, The ligament of the anterior cornu of the internal femilunar cartilage.
k, The transverse ligament connecting the anterior cornua of the femilunar cartilages to each other.
l, A slip fixed to the transverse ligament, and connected with the mucous ligament.
m, The external lateral ligament of the knee.
n, The insertion of the biceps muscle of the thigh.
o, The anterior ligament of the fibula.

FIG. 9.

A View of the Upper SURFACE of the TIBIA and SEMILUNAR CARTILAGES.

- a, b*, The glenoid cavities for lodging the condyles of the os femoris.

- c*, The external, and,
d, The internal femilunar cartilages.
e, The adhesion of the anterior cornu of the external femilunar cartilage to the fore-part of the tuberosity on the top of the tibia.
f, The superior ligament of the posterior cornu of this cartilage, connected with the posterior crucial ligament.
g, The inferior adhesion of the posterior cornu of the external femilunar cartilage.
h, The adhesion of the anterior cornu of the internal femilunar cartilage to the fore-part of the margin of the head of the tibia.
i, The adhesion of the posterior cornu to the back part of the tuberosity of the head of the tibia.
k, k, The common transverse ligament of the femilunar cartilages.
l, A slip fixing the transverse ligament, and intermixing with the mucous ligament.
m, The posterior crucial ligament.

FIG. 10.

An Outer and Fore View of the LIGAMENTS connecting the BONES of the TARSUS to those of the METATARSUS and LEG.

- a*, The extremity of the tibia.
b, The malleolus externus of the fibula.
c, The astragalus.
d, The os naviculare.
e, ——— cuboides.
f, ——— cuneiforme internum.
g, The metatarsal bone of the great toe, and, farther out, those of the other toes.
h, The anterior superior ligaments of the malleolus externus.
i, The middle perpendicular ligament of the malleolus externus.
k, The anterior ligament between the fibula and astragalus.
l, m, Irregular ligaments forming the ligamentous apparatus of the sinuous cavity of the astragalus and os calcis.
n, The superior ligament connecting the astragalus and os naviculare.
o, p, q, Numerous ligaments joining the bones of the tarsus to each other, and likewise to those of the metatarsus, and obtaining the names of Perpendicular, Oblique, Lateral, Dorsal, &c. according to their different directions and situations.
r, s, t, u, Ligaments which connect the bases of the metatarsal bones to each other.

FIG. 11.

LIGAMENTS between the Under and Back Part of the BONES of the LEG, and those of the FOOT.

- a*, A portion of the tibia.
b, ————— fibula.
c, The os calcis.

d, The

- d*, The upper part of the astragalus, upon which the tibia moves.
- e*, The posterior superior ligament of the malleolus externus.
- f*, A portion of the deltoid ligament.
- g*, The inferior posterior ligament of the malleolus externus.
- h*, A fibrous ligament between the fibula and astragalus.
- i*, A ligamentous slip proper to the astragalus.
- k*, A fibrous slip connected with the capsular ligament.
- l*, The frænulum of the capsular membrane, between the astragalus and os calcis.
- m*, The middle perpendicular ligament of the malleolus externus.
- n*, The insertion of the tendo Achillis, with its frænulum.

FIG. 12.

LIGAMENTS *in the SOLE of the FOOT.*

- a*, The os calcis.
- b*, ——— naviculare.
- c*, ——— cuneiforme magnum.

- d, d*, A section of the metatarsal bones.
- e, f*, The ligaments between the os calcis and naviculare.
- g*, The oblique, and,
- h*, The long ligament of great strength, which connect the os calcis to the os cuboides.
- i*, The transverse ligament between the cuboid and external cuneiform bones.
- k*, The ligaments between the navicular and internal cuneiform bones.
- l—q*, The ligaments connecting the bones of the tarsus to each other, and those again to the metatarsus.
- r, s, t*, The ligaments connecting the bases of the metatarsal bones.

FIG. 13.

The MASSES of FAT, with their FIMBRIÆ, which are situated round the Edges of the PATELLA.

- a*, The inner side of the patella.
- b, b, b*, The masses of fat.
- c, c, c*, The fimbriæ which project from the edges of these fatty substances.

END OF PART THIRD,

AND OF

VOLUME FIRST.



EDINBURGH:

PRINTED BY J. PILLANS & SONS,
LAWNMARKET.

