


This is a reproduction of a library book that was digitized by Google as part of an ongoing effort to preserve the information in books and make it universally accessible.

Google™ books

<https://books.google.com>





1

2

*Bulletin - United
States National Museum*

United States National Museum, Smithsonian Institution,
United States. Dept. of the Interior

for Doc




SI 3.3!

No. 58

Dark mean

SI 3.31
V SR



SKETCH MAP OF JAPAN AND ADJACENT TERRITORY.

SMITHSONIAN INSTITUTION
UNITED STATES NATIONAL MUSEUM
Bulletin 58

HERPETOLOGY OF JAPAN AND ADJACENT TERRITORY

BY
LEONHARD STEJNEGER
CURATOR, DIVISION OF REPTILES AND BATRACHIANS

With 35 Plates and 409 Figures
in the text



WASHINGTON
GOVERNMENT PRINTING OFFICE

1907

Published July 22, 1907.

11

ADVERTISEMENT.

The scientific publications of the National Museum consist of two series—the *Bulletin* and the *Proceedings*.

The *Bulletin*, publication of which was begun in 1875, is a series of more or less extensive works intended to illustrate the collections of the United States National Museum, and, with the exception noted below, is issued separately. These bulletins are monographic in scope and are devoted principally to the discussion of large zoological and botanical groups, faunas and floras, contributions to anthropology, reports of expeditions, etc. They are usually of octavo size, although a quarto form, known as the Special Bulletin, has been adopted in a few instances in which a larger page was deemed indispensable.

This work forms No. 58 of the Bulletin series.

Since 1902 the volumes of the series known as "*Contributions from the National Herbarium*," and containing papers relating to the botanical collections of the Museum, have been published as bulletins.

The *Proceedings*, the first volume of which was issued in 1878, are intended as a medium of publication of brief original papers based on the collections of the National Museum, and setting forth newly acquired facts in biology, anthropology, and geology derived therefrom, or containing descriptions of new forms and revisions of limited groups. A volume is issued annually, or oftener, for distribution to libraries and scientific establishments, and in view of the importance of the more prompt dissemination of new facts a limited edition of each paper is printed in pamphlet form in advance.

CHARLES D. WALCOTT,

Secretary of the Smithsonian Institution.

WASHINGTON, U. S. A., June 15, 1907.

TABLE OF CONTENTS.

	Page.
Table of contents	v
List of illustrations.....	xi
Preface.....	xvii
Herpetology of Japan and adjacent territory.....	1
Class Amphibia	1
Order Caudata.....	2
Suborder Mutabilia	3
Family Cryptobranchidae.....	3
Genus 1. MEGALOBATRACHUS.....	4
1. japonicus.....	6
Family Salamandridæ.....	11
Genus 2. TYLOTOTRITON.....	12
2. andersoni.....	12
Genus 3. DIEMICTYLUS.....	15
3. pyrhogaster.....	16
4. ensicauda.....	22
Family Ambystomide.....	24
Genus 4. HYNوبيUS.....	25
5. naevius.....	26
6. leechii.....	29
7. nebulosus.....	30
8. peropus.....	32
9. nigrescens.....	34
10. lichenatus.....	36
Genus 5. SALAMANDRELLA.....	37
11. keyserlingii.....	37
Genus 6. Onychodactylus.....	42
12. japonicus.....	42
Genus 7. GEOMOLGE.....	47
13. fischeri.....	47
Order Salientia.....	47
Suborder Costata.....	50
Family Discoglosside.....	50
Genus 8. BOMBINA.....	50
14. orientalis.....	51
Suborder Linguata.....	54
Family Bufonidæ.....	55
Genus 9. BUFO.....	55
15. formosus.....	60
16. smithi.....	64
17. sachalinensis.....	65
18. bufo asiaticus.....	66
19. bufo japonicus.....	69
20. raddei.....	70
21. melanostictus.....	72

Herpetology of Japan and adjacent territory—Continued.	Page.
Class Amphibia—Continued.	
Order Salientia—Continued.	
Suborder Linguata—Continued.	
Family Hylidae.....	75
Genus 10. <i>HYLA</i>	75
22. <i>arborea japonica</i>	76
[<i>arborea immaculata</i>].....	82
23. <i>stepheni</i>	84
24. <i>chinensis</i>	86
Family Engystomidae.....	87
Genus 11. <i>MICROHYLA</i>	87
25. <i>fissipes</i>	88
26. <i>okinavensis</i>	89
Family Ranidae.....	92
Genus 12. <i>RANA</i>	93
27. <i>nigromaculata</i>	94
28. <i>plancyi</i>	101
29. <i>okinavana</i>	102
30. <i>longicrus</i>	104
31. <i>holsti</i>	105
32. <i>japonica</i>	107
33. <i>temporaria</i>	113
34. <i>tsushimensis</i>	116
35. <i>amurensis</i>	119
36. <i>ijimæ</i>	121
37. <i>rugosa</i>	123
38. <i>limnocharis</i>	127
39. <i>swinhoana</i>	132
40. <i>ishikawæ</i>	132
41. <i>narina</i>	134
42. <i>namiyæi</i>	136
43. <i>tigerina</i>	139
Genus 13. <i>POLYPEDATES</i>	143
44. <i>schlegelii</i>	145
45. <i>viridis</i>	147
46. <i>owstoni</i>	149
47. <i>buengeri</i>	150
48. <i>eiffingeri</i>	153
49. <i>japonicus</i>	155
50. <i>leucomystax</i>	157
Tadpoles.....	159
Introductory.....	159
Key.....	161
Class Reptilia.....	161
Order Squamata.....	162
Suborder Sauria.....	162
Family Gekkonidae.....	164
Genus 1. <i>GEKKO</i>	165
1. <i>japonicus</i>	165
Genus 2. <i>PTYCHOZOON</i>	170
2. <i>horsfieldii</i>	170

Herpetology of Japan and adjacent territory—Continued.

Page.

Class Reptilia—Continued.

Order Squamata—Continued.

Suborder Sauria—Continued.

Family Gekkonidae—Continued.

Genus 3. HEMIDACTYLUS	172
3. <i>frenatus</i>	172
<i>marmoratus</i>	175
4. <i>bowringii</i>	176
Genus 4. COSYMBOTUS	178
5. <i>platyurus</i>	178
Genus 5. PEROPUS	180
6. <i>mutilatus</i>	180

Family Agamidae..... 182

Genus 6. JAPALURA	182
7. <i>swinhonis</i>	184
8. <i>polygonata</i>	188
9. <i>mitsukurii</i>	190
Genus ?.....	191
<i>species ?</i>	191

Family Scincidae..... 193

Genus 7. EUMECEUS	193
10. <i>laticutatus</i>	195
11. <i>laticutatus okadae</i>	200
12. <i>elegans</i>	202
13. <i>marginatus</i>	205
14. <i>chinensis</i>	208
15. <i>kishinouyei</i>	210

Genus 8. MABUYA	213
16. <i>longicaudata</i>	214

Genus 9. SPHENOMORPHUS	216
17. <i>igdicus</i>	216

Genus 10. LEIOLOPISMA	218
18. <i>laterale</i>	218

Genus 11. LYGOSAUROS	221
19. <i>pellopleurus</i>	222

Genus 12. CRYPTOBLEPHARUS	225
20. <i>boutonii nigropunctatus</i>	225

Family Lacertidae..... 228

Genus 13. TAKYDROMUS	229
21. <i>dorsalis</i>	229
22. <i>septentrionalis</i>	232
23. <i>formosanus</i>	235
24. <i>smaragdinus</i>	236
25. <i>tachydromoides</i>	238
26. <i>amurensis</i>	245
27. ? <i>wolteri</i>	247

Genus 14. EREMIAS	247
28. <i>argus</i>	248

Genus 15. LACERTA	251
29. <i>vivipara</i>	251

Herpetology of Japan and adjacent territory—Continued.	Page.
Class Reptilia—Continued.	
Order Squamata—Continued.	
Suborder Serpentes.....	254
Family Typhlopidae.....	259
Genus 16. TYPHLOPS.....	260
30. braminus.....	260
Family Natricidae.....	262
Subfamily Natricinae.....	263
Genus 17. NATRIX.....	263
31. vibakari.....	266
32. tigrina.....	272
33. tigrina lateralis.....	278
34. stolata.....	280
35. pryeri.....	284
36. piscator.....	288
37. annularis.....	291
38. swinhonis.....	293
Genus 18. ACHALINUS.....	294
39. spinalis.....	297
Subfamily Homalopsinae.....	299
Genus 19. ENHYDRIS.....	300
40. plumbea.....	300
41. bennettii.....	302
Genus 20. HURRIA.....	304
42. rynchops.....	304
Subfamily Coronellinae.....	306
Genus 21. ELAPHE.....	307
43. carinata.....	308
44. rufodorsata.....	310
45. schrenckii.....	313
46. dione.....	315
47. tenuirus.....	319
48. schmackeri.....	322
49. climacophora.....	324
50. quadrivirgata.....	327
51. conspicillata.....	334
Genus 22. LIOPELTIS.....	337
52. major.....	338
53. semicarinata.....	340
54. herminae.....	343
Genus 23. PTYAS.....	345
55. mucosus.....	345
56. korros.....	348
Genus 24. ZAMENIS.....	349
57. spinalis.....	349
Genus 25. ZAOCYS.....	352
58. dhumnades.....	352
Genus 26. HOLARCHUS.....	353
59. formosanus.....	354
Genus 27. DINODON.....	356
60. rufozonatum.....	358
61. rufozonatum walli.....	364

Herpetology of Japan and adjacent territory—Continued.	Page.
Class Reptilia—Continued.	
Order Squamata—Continued.	
Suborder Serpentes—Continued.	
Family Natricidæ—Continued.	
Subfamily Coronellinæ—Continued.	
Genus 27. <i>DINODON</i> —Continued.	
62. <i>semicarinatum</i>	366
63. <i>septentrionale rubistrati</i>	370
64. <i>orientale</i>	372
Genus 28. <i>CALAMARIA</i>	375
65. <i>berezowskii</i>	376
66. <i>pfefferi</i>	378
Subfamily Boigine.....	381
Genus 29. <i>BOIGA</i>	381
67. <i>kraepelini</i>	381
Genus 30. <i>PSAMMODYNASTES</i>	383
68. <i>pulverulentus</i>	383
Family Elapidae.....	386
Subfamily Elapine.....	386
Genus 31. <i>HEMIBUNGARUS</i>	387
69. <i>japonicus</i>	387
70. <i>boettgeri</i>	389
Genus 32. <i>CALLOPHIS</i>	391
71. <i>maclellandii</i>	391
Genus 33. <i>NAJA</i>	394
72. <i>naja atra</i>	394
Genus 34. <i>BUNGARUS</i>	397
73. <i>multicinctus</i>	397
Subfamily Hydrine.....	400
Genus 35. <i>LATICAUDA</i>	402
74. <i>laticaudata</i>	402
[<i>colubrina</i>].....	406
75. <i>semifasciata</i>	409
Genus 36. <i>EMYDOCEPHALUS</i>	413
76. <i>ijime</i>	413
Genus 37. <i>DISTEIRA</i>	418
77. <i>melanocephala</i>	421
[<i>fasciata</i>].....	427
[<i>gracilis</i>].....	427
78. <i>cyanocincta</i>	428
79. <i>godeffroyi</i>	430
[<i>viperina</i>].....	434
[Genus <i>LAPEMIS</i>].....	435
[<i>hardwickii</i>].....	435
[Genus <i>ESHYDRINA</i>].....	437
[<i>valakadyi</i>].....	437
Genus 38. <i>HYDRUS</i>	438
80. <i>platurus</i>	439
Family Cobridæ.....	442
Genus 39. <i>COLUBER</i>	443
81. <i>berus</i>	445

Herpetology of Japan and adjacent territory—Continued.		Page.
Class Reptilia—Continued.		
Order Squamata—Continued.		
Suborder Serpentes—Continued.		
Family Crotalidæ.....		448
Genus 40. AGKISTRODON.....		449
82. blomhoffii.....		457
83. blomhoffii ? affinis.....		461
84. blomhoffii brevicaudus.....		463
85. blomhoffii intermedius.....		464
Genus 41. TRIMERESURUS.....		465
86. mucrosquamatus.....		467
87. elegans.....		470
88. flavoviridis.....		475
89. okinavensis.....		479
90. gramineus.....		480
Order Testudinata.....		483
Suborder Athecæ.....		485
Family Dermochelidæ.....		485
Genus 42. DERMOCHELYS.....		485
91. schlegelii.....		485
Suborder Laminiæ.....		488
Family Testudinidæ.....		488
Subfamily Emydinæ.....		488
Genus 43. OCADIA.....		489
92. sinensis.....		489
Genus 44. CLEMMYS.....		492
93. japonica.....		492
Genus 45. GEOCLEMYS.....		496
94. reevesii.....		497
Genus 46. GEOEMYDA.....		500
95. spengleri.....		501
Genus 47. CYCLEMYS.....		503
96. flavomarginata.....		503
Family Cheloniidæ.....		506
Genus 48. CARETTA.....		507
97. olivacea.....		507
Genus 49. CHELONIA.....		509
98. japonica.....		509
Genus 50. ERETMOCHELYS.....		511
99. squamosa.....		511
Suborder Chilotæ.....		513
Family Trionychidæ.....		514
Genus 51. AMYDA.....		514
100. japonica.....		515
101. sinensis.....		524
102. schlegelii.....		526
103. ? maackii.....		529
[Genus DOGANIA].....		530
[subplana].....		531
Bibliography.....		533
List of Japanese and Formosan localities.....		545
Explanation of plates.....		553
Index.....		557

LIST OF ILLUSTRATIONS.

TEXT FIGURES.	Page.
Figs. 1-6.— <i>Tylotriton andersoni</i> . 1, top of head; 2, side of head; 3, open mouth; 4, section of tail at end of basal third; 5, hand; 6, foot	13
Figs. 7-8.— <i>Diemictylus pyrrhogaster</i> . 7, underside of head; 8, section of tail at end of basal third	17
Figs. 9-15.— <i>Diemictylus ensicauda</i> . 9, top of head; 10, side of head; 11, underside of head; 12, open mouth; 13, hand; 14, foot; 15, section of tail at end of basal third	22
Fig. 16.—Diagram of arrangement of vomerine teeth in <i>Onychodactylus</i> , <i>Salamandrella</i> , and <i>Geomolge</i>	24
Figs. 17-22.— <i>Hynobius navius</i> . 17, top of head; 18, side of head; 19, open mouth; 20, section of tail at end of basal third; 21, hand; 22, foot	26
Figs. 23-26.— <i>Hynobius nebulosus</i> . 23, top of head; 24, side of head; 25, open mouth; 26, section of tail at end of basal third	30
Figs. 27-32.— <i>Hynobius nigrescens</i> . 27, top of head; 28, side of head; 29, open mouth; 30, section of tail at end of basal third; 31, hand; 32, foot	34
Figs. 33-38.— <i>Salamandrella keyserlingii</i> . 33, top of head; 34, side of head; 35, open mouth; 36, section of tail at end of basal third; 37, hand; 38, foot	40
Figs. 39-43.— <i>Bombina orientalis</i> . 39, top of head; 40, side of head; 41, open mouth; 42, underside of right hand showing nup.ial excrescences; 43, foot	52
Figs. 44-48.— <i>Bufo formosus</i> . 44, top of head; 45, side of head; 46, underside of hand; 47, underside of foot; 48, upper side of foot to show extent of web	61
Figs. 49-52.— <i>Bufo bufo asiaticus</i> . 49, top of head; 50, side of head; 51, underside of hand; 52, underside of foot	67
Figs. 53-57.— <i>Bufo raddei</i> . 53, top of head; 54, side of head; 55, open mouth; 56, underside of hand; 57, underside of foot	71
Figs. 58-61.— <i>Bufo melanostictus</i> . 58, top of head; 59, side of head; 60, underside of hand; 61, underside of foot	73
Figs. 62-67.— <i>Hyla arborea japonica</i> . 62, top of head; 63, side of head; 64, open mouth; 65, underside of hand; 66, top of finger, side view; 67, underside of foot	78
Figs. 68-70.— <i>Hyla arborea japonica</i> . 68, top of head; 69, side of head; 70, underside of foot	80
Figs. 71-75.— <i>Microhyla okinawensis</i> . 71, top of head; 72, side of head; 73, open mouth; 74, underside of hand; 75, underside of foot	90
Figs. 76-80.— <i>Rana nigromaculata</i> . 76, head from above; 77, head from side; 78, open mouth; 79, underside of hand; 80, underside of foot	96
Figs. 81-83.— <i>Rana longicrus</i> . 81, top of head; 82, side of head; 83, underside of foot	104

	Page.
Fig. 84.— <i>Rana holsti</i> . Underside of left hand.....	106
Figs. 85-89.— <i>Rana japonica</i> . 85, top of head; 86, side of head; 87, open mouth; 88, underside of hand; 89, underside of foot.....	109
Figs. 90-93.— <i>Rana temporaria</i> . 90, top of head; 91, side of head; 92, open mouth; 93, underside of foot.....	114
Figs. 94-98.— <i>Rana tsushimensis</i> . 94, top of head; 95, side of head; 96, open mouth; 97, underside of hand; 98, underside of foot.....	117
Figs. 99-103.— <i>Rana ijima</i> . 99, top of head; 100, side of head; 101, open mouth; 102, underside of hand; 103, underside of foot.....	122
Figs. 104-106.— <i>Rana rugosa</i> . 104, top of head; 105, side of head; 106, open mouth.....	124
Figs. 107-110.— <i>Rana limnocharis</i> . 107, top of head; 108, side of head; 109, underside of hand; 110, underside of foot.....	128
Figs. 111-116.— <i>Rana ishikawa</i> . 111, top of head; 112, side of head; 113, open mouth; 114, underside of hand; 115, underside of foot; 116, tip of digit.....	133
Figs. 117-121.— <i>Rana varina</i> . 117, top of head; 118, side of head; 119, open mouth; 120, underside of hand; 121, underside of foot.....	135
Figs. 122-125.— <i>Rana namiyai</i> . 122, top of head; 123, side of head; 124, underside of hand; 125, underside of foot.....	137
Fig. 126.— <i>Rana namiyai</i> . Open mouth.....	137
Figs. 127-131.— <i>Rana tigerina</i> . 127, top of head; 128, side of head; 129, open mouth; 130, underside of hand; 131, underside of foot.....	140
Figs. 132-136.— <i>Polypedates schlegelii</i> . 132, top of head; 133, side of head; 134, open mouth; 135, underside of hand; 136, underside of foot.....	145
Figs. 137-141.— <i>Polypedates viridis</i> . 137, top of head; 138, side of head; 139, open mouth; 140, underside of hand; 141, underside of foot.....	148
Figs. 142-146.— <i>Polypedates buergeri</i> . 142, top of head; 143, side of head; 144, open mouth; 145, underside of hand; 146, underside of foot.....	151
Fig. 147.— <i>Polypedates eifingeri</i> . Vomerine teeth.....	154
Figs. 148-153.— <i>Polypedates japonicus</i> . 148, top of head; 149, side of head; 150, open mouth; 151, underside of hand; 152, underside of foot; 153, tip of digit.....	155
Fig. 154.—Open mouth of tadpole of <i>Rana temporaria</i> . <i>bo</i> , buccal orifice; <i>lp</i> , labial papillae; <i>m'</i> , upper, <i>m''</i> , lower mandible; <i>l'</i> , series of teeth on upper lip; <i>l''</i> , series of teeth on lower lip.....	159
Figs. 155-156.—Lower view of tadpoles, showing location of spiraculum and anal opening. 155, Discoglossoid (costate) toad; 156, <i>Hyla sp.</i> , spiraculum; <i>an.</i> , anus.....	160
Figs. 157-159.—Mouth of tadpoles. 157, <i>Bombina salsu</i> ; 158, <i>Hyla arborea</i> ; 159, <i>Bufo bufo</i>	160
Figs. 160-163.— <i>Gekko japonicus</i> . 160, top of head; 161, side of head; 162, chin; 163, underside of foot.....	167
Figs. 164-166.— <i>Pychozoon horsfieldii</i> . 164, side of head; 165, underside of right hind leg and foot; 166, underside of tail, middle and terminal portions.....	171
Figs. 167-168.— <i>Hemidactylus frenatus</i> (type of <i>H. inornatus</i>). 167, side of head; 168, chin.....	173
Figs. 169-170.— <i>Hemidactylus bowringii</i> . 169, chin; 170, underside of foot..	176
Figs. 171-173.— <i>Peropus mutilatus</i> . 171, side of head; 172, chin; 173, underside of foot.....	180

	Page.
Figs. 174-175.— <i>Japalura polygonata</i> . 174, top of head; 175, side of head....	189
Figs. 176-178.—Head shields of Scincid lizards. 176, side view of head of <i>Leiolopisma</i> ; 177, upper view of head of <i>Leiolopisma</i> ; 178, upper view of head of <i>Eumia</i> . <i>cs</i> , chin-shields; <i>d</i> , disk on lower eyelid; <i>e</i> , ear-opening; <i>f</i> , frontal; <i>fn</i> , fronto-nasal; <i>fp</i> , fronto-parietal; <i>i</i> , interparietal; <i>il</i> , lower labials; <i>l</i> , loreals; <i>lb</i> , upper labials; <i>m</i> , mental; <i>n</i> , nasal; <i>nc</i> , nuchals; <i>p</i> , parietals; <i>pf</i> , prefrontal; <i>pn</i> , postnasal; <i>r</i> , rostral; <i>sc</i> , supraciliaries; <i>sn</i> , supranasal; <i>so</i> , supraocular; <i>t</i> , temporal.....	194
Figs. 179-180.— <i>Eumeces latiscutatus</i> . 179, side of head, normal postnasal; 180, underside of hind foot.....	197
Fig. 181.— <i>Eumeces latiscutatus okadae</i> . Side of face.....	201
Figs. 182-183.— <i>Eumeces elegans</i> . 182, side of head; 183, chin.....	203
Fig. 184.— <i>Eumeces marginatus</i> . Side of head.....	206
Fig. 185.— <i>Eumeces chinensis</i> . Side of head.....	208
Figs. 186-189.— <i>Eumeces kishinouyei</i> . 186, top of head; 187, side of head; 188, underside of head; 189, underside of foot.....	211
Figs. 190-192.— <i>Leiolopisma laterale</i> . 190, top of head; 191, side of head; 192, underside of head.....	219
Figs. 193-195.— <i>Lygosaurus pellopleurus</i> . 193, top of head; 194, side of head; 195, underside of head.....	223
Figs. 196-198.— <i>Cryptoblepharus boutonii nigropunctatus</i> . 196, top of head; 197, side of head; 198, underside of head.....	226
Figs. 199-202.— <i>Takydromus dorsalis</i> . 199, top of head; 200, underside of head; 201, middle of back; 202, inguinal region.....	230
Figs. 203-204.— <i>Takydromus smaragdinus</i> . 203, underside of head; 204, inguinal region.....	236
Figs. 205-208.— <i>Takydromus tachydromoides</i> . 205, top of head; 206, side of head; 207, underside of head; 208, inguinal region.....	240
Figs. 209-212.— <i>Takydromus tachydromoides</i> . 209, top of head; 210, middle of back; 211, top of head; 212, chin-shields.....	242
Figs. 213-216.— <i>Takydromus amurensis</i> . 213, top of head; 214, side of head; 215, middle of back; 216, inguinal region.....	246
Figs. 217-219.— <i>Eremias argus</i> . 217, top of head; 218, side of head; 219, femoral pores and anal region.....	249
Figs. 220-222.— <i>Lacerta vivipara</i> . 220, top of head; 221, side of head; 222, femoral pores and anal region.....	252
Figs. 223-228.—Posterior dorsal vertebrae of snakes. 223, back view; 224, lower view; 225, side view of vertebra with hypapophysis; 226, back view; 227, lower view; 228, side view of vertebra without hypapophysis.....	256
Figs. 229-231.—Head shields of <i>Typhlops</i> . <i>f</i> , frontal; <i>ip</i> , interparietal; <i>l</i> , supralabials; <i>n</i> , nasal; <i>o</i> , ocular; <i>p</i> , parietals; <i>pf</i> , prefrontal; <i>pro</i> , preocular; <i>pto</i> , postocular; <i>r</i> , rostral; <i>so</i> , supraocular.....	259
Figs. 232-235.— <i>Typhlops braminus</i> . 232, top of head; 233, side of head; 234, underside of head; 235, tail.....	261
Figs. 236-238.—Head shields of a natricid snake. <i>cs'</i> , anterior chin-shields; <i>cs''</i> , posterior chin-shields; <i>f</i> , frontal; <i>il</i> , lower labials; <i>in</i> , internasals; <i>l</i> , loreal; <i>lb</i> , supralabials; <i>m</i> , mental; <i>n'</i> , anterior nasal; <i>n''</i> , posterior nasal; <i>p</i> , parietal; <i>pf</i> , prefrontals; <i>pro</i> , preocular; <i>pto</i> , postocular; <i>r</i> , rostral; <i>so</i> , supraocular; <i>t</i> , temporals; <i>v</i> , ventrals.....	262

	Page.
Figs. 239-241.— <i>Natrix vibakari</i> . 239, top of head; 240, side of head; 241, rostral from front	267
Figs. 242-243.— <i>Natrix tigrina</i> . 242, top of head; 243, side of head.....	273
Figs. 244-246.— <i>Natrix stolata</i> . 244, top of head; 245, side of head; 246, underside of head	281
Figs. 247-249.— <i>Natrix pryeri</i> . 247, top of head; 248, side of head; 249, underside of head	284
Figs. 250-252.— <i>Natrix piscator</i> . 250, top of head; 251, side of head; 252, underside of head	288
Figs. 253-255.— <i>Natrix amularis</i> . 253, top of head; 254, side of head; 255, rostral from front	292
Fig. 256.— <i>Natrix swinhonis</i> . Side of head.....	294
Figs. 257-259.— <i>Achalina spinalis</i> . 257, top of head; 258, side of head; 259, underside of head	297
Figs. 260-262.— <i>Euhydria plumbea</i> . 260, top of head; 261, side of head; 262, underside of head	301
Figs. 263-265.— <i>Euhydria bennettii</i> . 263, top of head; 264, side of head; 265, underside of head	303
Figs. 266-268.— <i>Elaphe carinata</i> . 266, top of head; 267, side of head; 268, underside of head	309
Figs. 269-271.— <i>Elaphe rufolorsata</i> . 269, top of head; 270, side of head; 271, underside of head	311
Fig. 272.— <i>Elaphe diome</i> . Top of head	317
Figs. 273-274.— <i>Elaphe tenuis</i> . 273, top of head; 274, side of head.....	320
Figs. 275-276.— <i>Elaphe schmackeri</i> . 275, top of head; 276, side of head.....	323
Figs. 277-278.— <i>Elaphe climacophora</i> . 277, top of head; 278, side of head....	325
Figs. 279-281.— <i>Elaphe quadrivirgata</i> , adult. 279, top of head; 280, side of head; 281, color pattern around the middle of the body.....	329
Figs. 282-283.— <i>Elaphe quadrivirgata</i> , young. 282, color pattern of top of head and neck; 283, color pattern around the middle of the body	330
Figs. 284-286.— <i>Elaphe conspicillata</i> . 284, top of head, with color pattern; 285, side of head; 286, underside of head	335
Figs. 287-289.— <i>Liopeltis major</i> . 287, top of head; 288, side of head; 289, underside of head.....	339
Figs. 290-292.— <i>Liopeltis semicarinata</i> . 290, top of head; 291, side of head; 292, underside of head.....	341
Figs. 293-295.— <i>Liopeltis semicarinata</i> . 293, top of head; 294, side of head; 295, underside of head.....	342
Figs. 296-297.— <i>Liopeltis herminia</i> . 296, top of head; 297, side of head	344
Figs. 298-300.— <i>Ptyas mucosus</i> (type of <i>trifrenatus</i>). 298, top of head; 299, side of head; 300, underside of head.....	346
Figs. 301-303.— <i>Zamenis spinalis</i> . 301, top of head; 302, side of head; 303, underside of head	350
Figs. 304-306.— <i>Holarchus formosanus</i> . 304, top of head; 305, side of head; 306, underside of head.....	355
Figs. 307-309.— <i>Dinodon rufozonatum walli</i> . 307, top of head; 308, side of head; 309, underside of head.....	364
Figs. 310-311.— <i>Dinodon semicarinatum</i> . 310, top of head; 311, side of head..	368
Fig. 312.— <i>Calamaria berezowskii</i> . Top of head	377
Figs. 313-316.— <i>Calamaria psiffieri</i> . 313, top of head; 314, side of head; 315, underside of head; 316, color pattern around middle of body.	379
Figs. 317-319.— <i>Psammodonastes pulverulentus</i> . 317, top of head; 318, side of head; 319, underside of head	384

	Page.
Figs. 320-322.— <i>Hemibungarus boettgeri</i> . 320, top of head; 321, side of head; 322, color pattern around middle of body	390
Figs. 323-324.— <i>Calliophis macclellandii</i> . 323, top of head; 324, side of head..	392
Figs. 325-327.— <i>Bungarus multicinctus</i> . 325, underside of head; 326, side of head; 327, top of head.....	398
Figs. 328-330.— <i>Laticauda laticaudata</i> . 328, top of head; 329, side of head; 330, underside of head.....	405
Figs. 331-333.— <i>Laticauda semifasciata</i> . 331, top of head; 332, side of head; 333, underside of head.....	410
Figs. 334-337.— <i>Emydocephalus ijima</i> . 334, top of head; 335, side of head; 336, underside of head; 337, scutellation at middle of body..	415
Figs. 338-340.— <i>Disteira melanocephala</i> . 338, top of head; 339, side of head; 340, rostral	422
Figs. 341-344.— <i>Disteira melanocephala</i> . 341, top of head; 342, side of head; 343, underside of head; 344, rostral.....	422
Figs. 345-347.— <i>Disteira melanocephala</i> . 345, top of head; 346, side of head; 347, underside of head.....	423
Figs. 348-350.— <i>Disteira melanocephala</i> . 348, side of head; 349, scutellation at middle of body; 350, side of tail.....	424
Fig. 351.— <i>Disteira cyanocincta</i> . Side view of head.....	429
Figs. 352-354.— <i>Disteira godeffroyi</i> . 352, top of head; 353, side of head; 354, underside of head	432
Figs. 355-357.— <i>Hydruis platurus</i> . 355, top of head; 356, side of head; 357, underside of head	440
Figs. 358-360.— <i>Coluber berus</i> . 358, top of head; 359, side of head; 360, underside of head.....	446
Fig. 361.—Poison apparatus of a Crotalid snake. Right side. <i>a</i> , External pterygoid bone (ectopterygoid or transpalatine); <i>b</i> , internal pterygoid bone (entopterygoid); <i>c</i> , palatal bone; <i>d</i> , maxillary bone; <i>e</i> , lachrymal bone (prefrontal).....	449
Figs. 362-364.— <i>Agkistrodon blomhoffii</i> . 362, top of head; 363, side of head; 364, underside of head.....	458
Fig. 365.— <i>Agkistrodon blomhoffii?</i> <i>affinis</i> . Side of head.....	462
Figs. 366-368.— <i>Trimeresurus elegans</i> . 366, top of head; 367, side of head; 368, underside of head	472
Fig. 369.— <i>Trimeresurus flavoviridis</i> . Side of head.....	476
Figs. 370-372.— <i>Trimeresurus gramineus</i> . 370, top of head; 371, side of head; 372, underside of head.....	476
Fig. 373.— <i>Dermochelys coriacea</i> , young. Entire animal from above	486
Figs. 374-376.— <i>Dermochelys coriacea</i> , young. 374, underside of shell; 375, side of head; 376, underside of head.....	487
Figs. 377-378.— <i>Ocacia sinensis</i> . 377, carapace; 378, plastron	490
Figs. 379-381.— <i>Ocacia sinensis</i> . 379, top of head; 380, side of head; 381, head from front	491
Figs. 382-383.— <i>Clemmys japonica</i> . 382, carapace; 383, plastron.....	493
Fig. 384.— <i>Clemmys japonica</i> , young.....	495
Figs. 385-386.— <i>Geoclemmys reevesii</i> . 385, carapace; 386, plastron.....	498
Figs. 387-388.— <i>Geoclemmys reevesii</i> . 387, top of head; 388, side of head.....	498
Fig. 389.— <i>Caretta caretta</i> , young. Entire animal from above	508
Figs. 390-392.— <i>Caretta caretta</i> , young. 390, shell from below; 391, head from side; 392, underside of head.....	508
Figs. 393-395.— <i>Chelonia japonica</i> , young. 393, entire animal from above; 394, side of head; 395, plastron.....	510

	Page.
Figs. 396-399.— <i>Eretmochelys imbricata</i> . 396, shell from below; 397, top of head; 398, side of head; 399, right fore flipper, dorsal view . . .	512
Fig. 400.— <i>Eretmochelys imbricata</i> . Shell from above	513
Fig. 401.—Elements of the trionychid plastron, showing plastral callosities. <i>c</i> , commissure; <i>ep</i> , entoplastron; <i>epi</i> , epiplastron; <i>hp</i> , hypoplastron; <i>hyp</i> , hypoplastron; <i>pa</i> , anterior inner process; <i>pm</i> , median process; <i>pp</i> , posterior inner process; <i>xp</i> , xiphiplastron	518
Figs. 402-403.— <i>Amyda japonica</i> . 402, side view; 403, view from above	520
Figs. 404-405.— <i>Amyda japonica</i> . 404, underside; 405, side of head	521
Fig. 406.— <i>Amyda schlegelii</i> . Side view	522
Fig. 407.— <i>Amyda schlegelii</i> . View from above	527
Fig. 408.— <i>Amyda schlegelii</i> . Underside	528
Fig. 409.— <i>Amyda schlegelii</i> . Side view	528

PLATES.

	Facing page.
I. Sketch map of Japan and adjacent territory	title page
II. <i>Diemictylus purrhogaster</i> . (From Schlegel)	16
III. <i>Hymobius navius</i> ; <i>H. nebulosus</i> . (From Schlegel)	26
IV. <i>Hymobius lichenatus</i> ; <i>H. peropus</i> ; <i>H. lechii</i> . (From Boulenger)	32
V. <i>Ouzchodactylus japonicus</i> ; <i>Geomolge fischeri</i> ; <i>Salamandrella keyserlingii</i> . (From Schlegel, Boulenger, and Strauch)	42
VI. <i>Ouzchodactylus japonicus</i> . (From Schlegel)	42
VII. <i>Bombina orientalis</i> . (From Boulenger)	50
VIII. <i>Bufo formosus</i> . (From Boulenger)	60
IX. <i>Hyla arborea japonica</i> ; <i>H. chinensis</i> . (From Boulenger, Schlegel, and Guenther)	76
X. <i>Rana nigromaculata</i> ; <i>R. rugosa</i> . (From Schlegel)	96
XI. <i>Rana japonica</i> ; <i>R. amurensis</i> . (From Schlegel and Boulenger)	108
XII. <i>Polypedates buergeri</i> . (From Schlegel)	152
XIII. <i>Gekko japonicus</i> . (From Schlegel)	166
XIV. <i>Japalura polygonata</i> . (From Boulenger)	188
XV. <i>Eumeces latiscutatus</i> . (From Schlegel)	194
XVI. <i>Mabuya longicaudata</i> . (From Fischer)	214
XVII. <i>Sphenomorphus indicus</i> ; <i>Lygosaurus pellopleurus</i> . (From Boulenger)	216
XVIII. <i>Takydromus tachydromoides</i> . (From Schlegel)	240
XIX. <i>Natrix tigrina</i> . (From Temminck and Schlegel)	272
XX. <i>Natrix pryeri</i> . (From Boulenger)	284
XXI. <i>Elaphis quadrivirgata</i> . (From Temminck and Schlegel)	328
XXII. <i>Laticauda semifasciata</i> . (From Temminck and Schlegel)	408
XXIII. <i>Distenia melanocephala</i> . (From Boulenger)	420
XXIV. <i>Lopemis hardwickii</i> . (From Jan)	434
XXV. <i>Euhydriina calakadjan</i> . (From Jan)	436
XXVI. <i>Agkistrodon blomhoffii</i> . (From Temminck and Schlegel)	456
XXVII. <i>T. inermis</i> ; <i>T. flavoviridis</i> . (From Hilgendorf)	476
XXVIII. <i>Ovalia sinensis</i> . (From Gray)	490
XXIX. <i>Clemmys japonica</i> . (From Schlegel)	492
XXX. <i>Geoclemys reversii</i> . (From Gray)	496
XXXI. <i>Geoclemys spengleri</i>	500
XXXII. <i>Geoclemys spengleri</i>	500
XXXIII. <i>Cyclemys flavomarginata</i> . (From Guenther)	504
XXXIV. <i>Caretta olivacea</i> . (From Eschscholtz)	506
XXXV. <i>Amyda japonica</i> . Young	522

PREFACE.

The present work is the result of a critical study of more than 1,500 specimens from Japan and its dependencies, as well as from portions of the adjacent territory on the mainland of eastern Asia. By far the larger number are contained in the herpetological collection of the United States National Museum, and in addition to these I have had here in Washington for direct comparison the extensive collections of the Science College Museum of the Imperial University in Tokyo. For this privilege I wish to express my sincere gratitude to the authorities of the university, and most especially to Prof. Isao Ijima, in whose immediate charge these collections are. Thanks to the kindness of Prof. C. Ishikawa, I have also had some specimens belonging to the Imperial Museum in Tokyo. Moreover, it was my good fortune during two brief visits in Tokyo in 1896 and 1897 to be able to examine there a number of specimens in the two institutions mentioned.

The American museums are not rich in material from the countries covered by this work, but several important specimens are preserved in the museum of the Philadelphia Academy of Sciences, and the collections made by Doctors Furness and Heller in the Riu Kiu Islands are deposited in the Wistar Institute, Philadelphia. A few specimens collected by Mr. Olinger in Korea are in the museum of the State University of Michigan. For permission to study this material, I am greatly indebted to the authorities of these institutions.

It would have been impossible to prepare the present volume without an examination of the important material of types and other specimens contained in the various museums of Europe. I need only mention Leiden, where the material brought home by Buerger and Von Siebold, and upon which Schlegel and Temminck based their account of the reptiles and batrachians in the *Fauna Japonica*, is preserved, and the British Museum with its vast material and numerous types. It has been my privilege during three visits to Europe to study these and numerous other collections in various cities, and it gives me particular pleasure to acknowledge with sincere thanks the liberality of the authorities of the Smithsonian Institution and the National Museum in enabling me to undertake these visits for that specific purpose. It was thus possible for me to study the collections in the Hamburg

Museum of Natural History brought home by Lenz, Warburg, and Dieckmann from the Riu Kiu Islands and Amurland; in the Museum Senckenbergianum, Frankfort on the Main, the depository of Bernhard Schmacker's collection from China and the Riu Kiu Islands; in Stockholm's Riksmuseum where are the Japanese collections made by Dr. Osc. Nordquist during the *Vega* expedition; and in the museums of Bergen and Christiania, some highly important collections from Formosa. To the directors and curators of all these institutions I wish to extend my thanks for facilities and assistance received.

The material in the United States National Museum consists of some very important collections. Only a few specimens from the Perry expedition which opened Japan to foreign intercourse are still in existence; there were probably not very many to start with. The next expedition, the Rodgers North Pacific Exploring Expedition, of which Dr. W. Stimpson was the naturalist, secured rather large collections, and these were reported upon by Doctor Hallowell. Unfortunately, he died before the printing of the report, the editing of which was intrusted to Dr. E. D. Cope, then quite young and inexperienced. Many new species were described, but the descriptions were very deficient and the whole publication was marred by numerous serious errors and lapses. A large number of the specimens are lost, many were apparently never returned to the museum, while the few that are left have suffered from neglect and most of them are in poor condition. With their aid, however, and that of Doctor Stimpson's manuscript field catalogue, which is still in existence, I have been enabled to straighten out many dubious questions. Of later noteworthy collections made for the museum, or acquired by it, I may mention the excellent material gathered by my late friend Pierre Louis Jouy in Japan and Korea. Doctor Ferrebee also sent in some specimens from the latter country. During my visit to Japan in 1896, as one of the Fur Seal Commissioners, I was enabled to secure numerous specimens of the commoner species from the surroundings of Yokohama. One of the most extensive and valuable collections of reptiles and batrachians was made by Dr. Hugh M. Smith, United States Deputy Fish Commissioner, in 1901, in southern Japan, especially in the island of Shikoku, hitherto almost unknown herpetologically. Dr. S. Nozawa, of the Agricultural College, Sapporo, has presented to the museum an excellent collection of Yezo reptiles. In 1904 the National Museum acquired from Mr. Alan Owston his splendid collection of Japanese reptiles embracing large series of species from Hondo, the central and southern groups of the Riu Kiu Islands and from Formosa. Finally, the Science College of the Imperial University in Tokyo has presented to the U. S. National Museum a valuable set of duplicates of Japanese and Formosan reptiles.

The territory embraced in the present work (see Plate I) consists of what is known as Japan proper—i. e., the four main islands and their immediately adjacent islets, the Riu Kiu Archipelago, Formosa, the Bonin Islands, Korea, Sakhalin, Amurland, and the Russian Coast Province, as well as the Chinese provinces of Manchuria and Pechili. A few species, mostly marine snakes, not hitherto recorded from within this territory have been included because of the probability of their occurrence; but these have been inclosed in brackets.

Analytical keys for the easy and sure identification of the groups and species are provided. It must be understood, however, that reptiles and batrachians are often subject to individual variation far in excess of the true specific (or subspecific) and even generic limits. In all cases, therefore, the specimens should be compared with the detailed descriptions. These are not generalized diagnoses, however, but minute descriptions of individuals, the deviations of other specimens under observation being noted under the heading "Variation." In selecting specimens for description or for illustration I have chosen individuals collected inside the territorial limits of this work whenever possible. Preferably the type-specimen has been described, or, if not available, a specimen from the type-locality or from the nearest locality to that of the type. If more than one specimen of this character were available, the one showing the species in its typical and normal development has been selected.

Full synonymy of each genus and species has been given, with references to nearly all the published records of the species within the territorial limits. At the end a bibliography is provided, giving full titles of publications specifically referring to the herpetological fauna of the territory.

Special attention has been given to the geographical distribution of the various forms. Unfortunately, a large number of specimens in the old collections are credited to "Japan," or "Loochoo," or "Formosa," without further particulars, and a good many others have wrong localities attached to them. Much critical work has been done to clear the records in the latter cases. Nevertheless, the range of very many species is as yet only very imperfectly known. Large portions of the territory included are still unexplored, and many islands are as yet unvisited by a scientific collector. With regard to the vertical distribution, it is known only imperfectly in the great majority of cases. To help resident and visiting naturalists remedy these defects in our knowledge this work has been largely undertaken. It was at first intended to accompany this account by a general tabulation of the geographical distribution of the various forms and a full discussion of their origin and dispersal, but this will be treated of by the author in a separate paper.

With regard to the nomenclature of families, genera, and species, the author adheres strictly to the "International Rules of Zoological Nomenclature" adopted by the International Congresses of Zoology.^a Changes in nomenclature necessitated by these rules, therefore, must not be laid to any desire of the author to alter names, but to the necessity of conforming strictly to the laws now generally accepted by the working zoologists of the world. As for my guiding principle in selecting group names higher than families, see pages 48 to 49.

It has not been found expedient to accompany this work by a map of the region giving the various localities mentioned, since good maps of Japan and adjacent territory are nowadays easily available. The different names employed in the literature and on the maps for the same localities and the often radically different spelling of the same names have made necessary a synonymical and briefly descriptive list of Japanese localities which will be found at the end of the book.

In the preparation of the present work the author has received kind assistance from many friends and colleagues which he wishes to acknowledge with gratitude. First of all, he must mention his Japanese friends connected with the Imperial University in Tokyo, Professor Mitsukuri, Professor Ijima, Professor Ishikawa, and Mr. Namiye. To Dr. G. A. Boulenger, British Museum; Dr. O. Boettger, Senckenberg Museum; Doctors Kraepelin and Pfeffer, Hamburg Museum; Doctor Jentink, Leiden Museum; Dr. E. Loennberg, Stockholm Museum; Dr. R. Collett, Kristiania Museum; and Messrs. Witmer Stone and Rehn, Philadelphia Academy of Sciences, he is indebted for loan of specimens, much information, and numerous courtesies.

Most of the outline drawings illustrating the text have been prepared by Mr. R. G. Paine.

The plates are mostly reproductions of important illustrations, more or less inaccessible to those for whom this work is chiefly intended. Very often these illustrations represent type-specimens, and in nearly every instance are based on specimens collected in the region covered by this work. The expensive *Fauna Japonica* is long since out of print, and the reproduction of the best figures from this classic will be welcome to the majority of the students of Japanese herpetology.

For the original drawings of two of the plates (Plates XXXI-XXXII) I am under great obligations to Prof. Isao Ijima.

^a Règles Internationales de la Nomenclature Zoologique adoptées par les Congrès Internationaux de Zoologie. (Paris, F. R. de Rudeval, Éditeur, 1905. 8^{vo}. 64 pp.)

HERPETOLOGY OF JAPAN AND ADJACENT TERRITORY.

By LEONHARD STEJNEGER,

Curator, Division of Reptiles and Batrachians, U. S. National Museum.

INTRODUCTION.

Until a comparatively recent date reptiles and amphibians were generally considered members of the same class, intermediate between the old class Pisces and the birds. True, the Amphibia were elevated to class rank early in the last century, but the proposition was not generally accepted until the subdivision of the vertebrates into Anamnia and Amniota had been gaining ground. It is now generally conceded that the Amphibia, or Batrachia, as they are often called, are more nearly allied to the fishes, and the reptiles more so to the birds than the two classes are among themselves, some authors even going so far, and not without good reasons, as to include the reptiles with the birds in the same class, Monocotylia. Yet, chiefly for the same reasons which cause the Lancelets and the Lampreys to be embraced with the true fishes in the term "ichthyology," the old name for the study of the compound class Amphibia or Reptilia, "herpetology," is still in common use, and in faunistic works, like the present, the two classes are usually treated of together.

Class AMPHIBIA.

1758. *Amphibia* LINNÆUS, Syst. Nat., 10 ed., I, p. 194.
1802. *Batrachia* MACARTNEY, in Ross's Translat. Cuvier's Lect. Comp. Anat., I, tab. III.
1803. *Batrachii* DAUDIN, Hist. Nat. Rept., V, tab. p. 8.
1806. *Batrachii* DUMÉRIL, Zool. Anal., p. 90.
1807. *Calamita* LINK, Besch. Nat. Samml. Rostock, II, p. 53.
1813. *Achelata* FISCHER, Zoognosia, 3 ed., I, p. 57.
1814. *Ranacea* WILBRAND, Classif. Thiere, p. 117.
1820. *Nuda* HEMPRICH, Grundr. Naturg., p. 111.
1821. *Dipnoa* LEUCKART, Isis, 1821, Litt. Anz., p. 259.
1847. *Malacopoda* MAYER, Rheinl. und Westphal. Verhandl., VI (p. 177).
1855. *Psiloderma* VAN DER HOEVEN, Handb. Dierk., 2 ed., II, p.—: Handb. Zool., II, p. 251.

The amphibians, or batrachians, as they are also called, are anamniote, archæcraniate, and stomatophysous vertebrates possessing a well-developed skull provided with a lower jaw and articulating with the vertebral column by means of two occipital condyles; limbs, when not atrophied, consisting of humerus or femur followed by two propodials (radius and ulna, tibia and fibula), metapodials (carpals and metacarpals, tarsals and metatarsals), and digits (phalanges); heart with three chambers; internal nares; respiration, at least during part of life, by means of gills; skin naked. Young, usually after leaving egg, undergoes a metamorphosis.

The term "Batrachia," usually applied to this class, has to be rejected and "Amphibia" accepted, since it has been shown that the former name originally applied only to the order of tailless frogs, and that consequently it is a synonym pure and simple of the much older term Salientia.^a

The class includes three recent orders, namely, the Cæcilians (Apoda), the Salamanders (Caudata), and the Frogs (Salienta). Of these only the two latter orders are represented within our present limits.

Order CAUDATA.

1806. *Caudati* DUMÉRIL, Zool. Anal., p. 94.
 1811. *Caudata* OPPEL, Ordn. Rept., p. 72.
 1813. *Urodéli* FISCHER, Zoognosia, 3 ed., I, p. 58.
 1820. *Gradientia* MERREM, Syst. Amph., p. 166.
 1825. *Urodela* LATREILLE, Fam. Nat. Règne Anim., p. 105.
 1828. *Cercopi* WAGLER, Isis, 1828, p. 859.
 1833. *Sozura* VAN DER HOEVEN, Handb. Dierk., II, Pt. 2, p. 304.
 1835. *Homomorpha* FITZINGER, Ann. Wien. Mus., I, p. 107.
 1838. *Urophora* HOGG, Mag. Nat. Hist. (n. s.), III, p. 270.
 1855. *Sauvobatrachi* VAN DER HOEVEN, Handb. Dierk., 2 ed., II, p. 461.
 1857. *Delesura* JAN, Cenni Mus. Milan., p. 54.

This order is often called Urodeles or Urodela, and Duméril (Zoologie Analytique, 1806) is quoted as authority. This is a mistake, however, for Duméril only uses the French term "Urodèles," adding in parentheses the Latin word by which he designates the group, namely, *Caudati*. From an inspection of page 94 this is plain enough, but to clinch the matter one needs only examine the two indexes at the end of the book, the "Table Française" containing the word "Urodèles" (p. 330), and the "Table Latine" the *Caudati* (p. 333). The Latin forms "Urodela" and "Urodeles" have been introduced much later by other authors.

This order may be divided into three suborders, viz, the Mudpuppies (Proteida), the Sirens (Meantes), and the true Salamanders (Mutabilia), only the latter being found in the territory here included.

^a See Stejneger, Science (n. s.), XX, Dec. 30, 1904, pp. 924-925.

Suborder MUTABILIA.

1820. *Mutabilia* MERREM, Syst. Amph., p. 166.
 1820. *Salamandra* GOLDFUSZ, Handb. Zool., II, p. 129.
 1828. *Morphuromolgei* RITGEN, Nova Acta Acad. Leop. Carol., XIV, p. 277.
 1828. *Molge* RITGEN, Nova Acta Acad. Leop. Carol., XIV, p. 277.
 1832. *Derotremata*+*Salamandrina* MUELLER, Isis, 1832 (p. 504).
 1831. *Urodela* BONAPARTE, Saggio Distrib. Met., p. 77.
 1840. *Deiretremata*+*Aphanobranchiata* LEUCKART, Froriep's Neue Notizen, XIII, p. 20.
 1866. *Sozura* HÆCKEL, Gen. Morphol., II, p. CXXXI.
 1866. *Caducibranchiata* COPE, Journ. Phila. Acad. Sci., VI, Pt. 1, p. 102.
 1889. *Pseudosauria* COPE, Amer. Natural., XXIII, 1889, p. 861.

In the true salamanders we recognize two superfamilies, viz, the Amphiumoideæ and the Salamandroideæ. For our present purpose it is enough to separate them by the presence of well-developed eyelids in the latter and the absence of eyelids in the former. It is stated that the absence of eyelids is concurrent with the absence of a first epibranchial and with the connection of stapes with the quadrate arch in the Amphiumoideæ, while in the Salamandroideæ a first epibranchial is present and stapes not connected with the quadrate arch in the adult. The Amphiumoideæ correspond to Cope's Trematodera and Amphiumoidea, and to Boulenger's Amphiumidæ, while the Salamandroideæ are the latter's Salamandridæ and Cope's Pseudosauria (1889, N. Am. Batr., p. 33; not of 1898, Syllabus, p. 48, where the term corresponds strictly to our Mutabilia).

Both superfamilies occur within our limits. The genera which belong to them may be referred to their respective groups as follows:

ANALYTICAL KEY TO FAMILIES.

- α^1 (AMPHIUMOIDEÆ.) No eyelids.....CRYPTOBRANCHIDÆ, p. 3.
 α^2 (SALAMANDROIDEÆ.) Eyelids developed.
 β^1 Vomero-palatine teeth in two longitudinal series diverging backward; vertebræ opisthocæalous.....SALAMANDRIDÆ, p. 11.
 β^2 Vomero-palatine teeth in transverse series, or converging backward; vertebræ amphiocæalous.....AMBYSTOMIDÆ, p. 24.

Superfamily AMPHIUMOIDEÆ.

Two families compose the superfamily Amphiumoideæ, the Amphiumidæ and the Cryptobranchidæ, differing in many anatomical characters, the former represented by the eel-like *Amphiuma* with at most three digits on the rudimentary legs, the latter by the hellbender, with its more salamander-like body, 4-5 digits and well-developed limbs. The former occurs only in North America.

Family CRYPTOBRANCHIDÆ.

This family embraces three genera, *Proteocordylus* (*Andrias*), known only from the miocene of western Europe, *Cryptobranchus*, which only

occurs in eastern North America, and *Megalobatrachus*, known from Japan and China. They are so closely allied that osteologists and paleontologists prefer to regard the two latter at least as congeneric, but the closing of the branchial fissure in the adult *Megalobatrachus* seems to be sufficient reason for adopting the latter genus. In the American genus the fissure remains open throughout life.

Genus MEGALOBATRACHUS^a Tschudi.

1837. *Megalobatrachus* TSCHUDI. Neues Jahrb. Mineral. Geol. Paläont., (Stuttgart) 1837, Pt. 5, September, p. 547 (type, *M. sieboldi*).
1838. *Sieboldia* BONAPARTE in Gray, Ann. Mag. Nat. Hist., May, 1838, p. 413 (same type).
1838. *Sieboldia* AGASSIZ, in Anhang to Tschudi's Classif. Batr. (err. typogr.).
1840. *Hydrosalamandra* LEUCKART. Foriep's Neue Notizen, XIII, No. 2, Jan. 1840, p. 20 (same type).
1854. *Tritomegas* DUMÉRIE and BIBRON, Erpét. Gén., IX, p. 163 (same type).
1904. *Sieboldiana* ISHIKAWA, Proc. Nat. Hist. Tokyo Imp. Mus., I, No. 2, p. 21 (emendation).

Unless it can be shown that Bonaparte published his *Sieboldia* before September, 1837, it will have to give way to Tschudi's *Megalobatrachus*. Agassiz (l. c.), on September 29, 1838, speaks of "the established priority" of the former,^b but, like many later authors, he evidently regarded *Megalobatrachus* as first proposed by Tschudi in his "Classification der Batrachier," which Agassiz himself published in October, 1838. As will be seen from the above synonymy Tschudi had already published it in September, 1837, a fact of which Agassiz was apparently unaware.

Few animals can boast such an extensive literature as the Japanese giant salamander. The bibliography attached to this work and the synonymic list of quotations under the head of the species, greatly defective as they are, give a fair indication of the interest this animal and especially its anatomy has aroused among naturalists. It is only recently, however, that accurate and detailed observations on its habits and propagation have been made. A good account based upon ample experience in the field has been published by Prof. C. Sasaki,^c and recently Prof. C. Ishikawa has supplemented his notes by very important observations on the eggs and newly hatched young.^d Finally, Dr. C. Kerbert, in Amsterdam, has succeeded in making the giant salamander breed in captivity, thus being able to follow the process from the laying of the eggs to the hatching of the larvæ and their subsequent growth and change. A brief summary of their observations may be included here. According to them the giant

^a From μέγανος = μέγας, great; βάτραχος, frog, or, in this case, batrachian.

^b "Die constatirte Priorität seines Namens Sieboldia."

^c Journ. Coll. Sci. Tokyo, I, Pt. 3, 1887, pp. 269-274.

^d Proc. Nat. Hist. Tokyo Imp. Mus., I, No. 2, 1904, pp. 19-37.

salamander inhabits high mountain valleys, where it frequents the swift-running brooks of clear cold water from the mountain springs. These streams are quite shallow, have stony beds, seldom attain a width of more than a few meters and are for the most part shaded with shrubs and trees. The temperature of the water in the middle of August was found to be between 17° and 23° C. Here they keep themselves habitually in dark holes under rocks, along the banks or the middle of the stream, and cavities inhabited by one of these animals may be recognized by the bottom at the entrance being kept clean. They feed on fishes, frogs, crustaceans, earthworms, etc, and are easily captured by a baited fishhook thrust into the retreat of the sluggish animal. The flesh, which is eaten by the Japanese, is said to be delicious, and is also used for medicinal purposes. Another mode of capturing the salamander during the breeding season is to throw into the stream a strong-smelling mixture of various animal ingredients made into small balls, thereby enticing the animals out of their holes. As a result, the "hanzaki" is constantly becoming scarcer. Only the smaller and medium-sized individuals stay in the small, shallow streams, the big old monsters of 3 feet and more seek deeper water lower down. The eggs are deposited in August and September in the deep holes. They vary as to size and number according to the size of the female, and consist of an outer gelatinous envelope or capsule 16 to 20 mm. long, in which the spheroid egg floats in a clear fluid. Each capsule is connected with the next by means of a comparatively short string about equaling the length of the larger axis of the embryo, the whole deposit resembling a rosary in form. Kerbert estimates the number of eggs deposited by the female in the Amsterdam Aquarium at about 500. Ishikawa found 60 to 80 in the various holes examined by him. The entire development from the deposition of the eggs to the escape of the larva from the capsule lasted in captivity from fifty-two to sixty-eight days, the escaping larvæ measuring about 30 mm. in length. At this time they have external gills; the anterior extremities show indications of two fingers and the posterior limbs are indicated; the mouth which is still plainly ventral becomes gradually terminal. The gills seem to disappear when the young are between 200 and 250 mm. in length. Both Ishikawa and Kerbert maintain that the adult animal takes care of the eggs, wrapping the egg string around itself and by its movements keeping the mass in motion so as to facilitate the respiration process of the eggs and embryos; but they differ in opinion as to whether it is the male or the female who thus undertakes the care of the offspring, Ishikawa from observations in nature maintaining the latter, Kerbert from the behavior of the animals in the aquarium the former.

MEGALOBATRACHUS JAPONICUS^a (Temminck).

HANZAKI (Ishikawa).

1837. *Triton japonicus* TEMMINCK, Fauna Japon., Pt. 3, Coup d'œil, p. xxvi (Japan).—WAGNER, Muenchen Gelehrte Anzeiger, V, July 8, 1837, p. 55.—VAN DER HOEVEN, Tijdschr. Natuurl. Geschied., IV, Feb. 1838, p. 375.—*Cryptobranchus japonicus* VAN DER HOEVEN, Tijdschr. Natuurl. Geschied., IV, Feb. 1838, p. 384, pl. va, figs. 5-6 (Japan): Bull. Sci. Phys. Natur. Néerlande (Leiden), 1838, p. 91; Proc. Zool. Soc. London, 1838, p. 25; Ann. Mag. Nat. Hist., II, Nov. 1838, p. 230; Ann. Sci. Nat., (2) Zool., XI, 1839 (p. 63); Mém. Soc. Hist. Nat. Strasbourg, III, 1840, author's separate, p. 11, pls. —; Ann. Sci. Nat., (2) Zool., XV, 1841, p. 251 (151 by typogr. err.); Tijdschr. Natuurl. Geschied., VIII, 1841, p. 270 (blood corpuscles).—SCHMIDT, GODDARD, and VAN DER HOEVEN, Nat. Verh. Maatsch. Wet. Haarlem, XIX, 1862, p. 3 (anatomy).—HYRTL, De Cryptobr. Jap., 1865, p. 9 (descr.; anat.).—REIN and von RORETZ, Zool. Garten, XVII, 1876, p. 33, pl. — (distrib.; habits, etc.).—BOETTGER, Zool. Garten, XVII, 1876, p. 432 (young).—DAMBECK, Natur (Halle) (n. s.), III, 1877, (p. 685).—WIEDERSHEIM, Kopfskel. Urodel., 1877, p. 56, pl. II, figs. 21-22 (skull).—CORNALIA, Atti. Soc. Ital. Sci., XXI, 1878, p. 207 (dimensions).—HILGENDORF, Sitz. Ber. Naturf. Freunde, Berlin, 1880, p. 121.—BIELETSKI, [Rem. Physiol. Sal. Gig., 1882] p. — pl.—GEERTS, Nouv. Arch. Mus. Paris, (2) V, 1883, p. 274, pl. xvii.—REIN, Japan, Engl. ed., 1884, p. 188 (Hida to Iwami).—SASAKI, Journ. Coll. Sci. Tokyo, I, Pt. 3, 1887, p. 269 (Iga; Ise; Yamato; habits, eggs, etc.).—MOESCH, Neujahrsblatt Naturf. Ges. Zuerich, 1887, p. 1, figs. 1-4 (full fig. and skel.).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, 1891, p. 239 (absent in Yezo).—MEURON, Bull. Soc. Sci. Nat. Neuchâtel, XXI, 1893, p. 186.—GADOW, Camb. Nat. Hist., VIII, 1901, p. 98 (fig.).—OSAWA, Mitth. Med. Fac. Univers. Tokio, V, no. 4, 1902 (pp. 221-427+44 pls.) (anatomy).—*Andrias japonicus* LAPARENT, Traité Geol., 4 ed., 1900, p. 1532.—*Megalobatrachus japonicus* BEDDARD, Proc. Zool. Soc. London, 1903, II, Pt. 2 (publ. Apr. 1904) p. 298 (anatomy).
1837. *Megalobatrachus sieboldi* TSCHUDI, Neues Jahrb. Mineral. Geol. Palaeont., 1837, Pt. 5, Sept., p. 547 (type-locality, Japan; type in Leiden Mus.; Siebold collector); Classif. Batr., Oct. 1838, p. 96, pl. vi (Japan).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1876, p. 384 (southern Japan).—*Tritomegas sieboldii* DUMÉRIL and BIBRON, Érpét. Gén., IX, 1854, p. 164.—BLEEKER, Tijdschr. Nederland. Indië, XVI, 1858, p. 205.
1837. *Salamandra maxima* SCHLEGEL in Tschudi, Neues Jahrb. Mineral. Geol. Palaeont., 1837, Pt. 5, Sept., p. 546 (not of Barton, 1808); Fauna Japon. Rept., pp. 127, 139. Saur. et Batr., pls. vi, vii, viii (type-locality, "Suzuga yama near Sakanosta",^b Okude mountains).—SIEBOLD, Fauna Japon., Rept., 1838, p. xv.—POMPE VAN MEERDERVOORT, Natuurk. Tijdschr. Nederland. Indië, XX, Pts. 4-6, 1860, p. 386.—*Sieboldia maxima* GRAY, Cat. Batr. Grad. Brit. Mus., 1850, p. 52 (Japan).—PHISALIX, Compt. Rend. Acad. Sci. Paris, CXXV, 1897 (p. 121) (poisonous secretions); Bull. Mus. Hist. Nat. Paris, 1897 (p. 242).—*Megalobatrachus maximus* BOULENGER, Cat. Batr. Grad. Brit. Mus., 1882, p. 80 (Japan).—OKADA, Cat.

^a Signifying Japanese.^b Evidently Sudsaka Yama near Sakanoshita, of Hassenstein's Atlas of Japan, the mountain where the boundaries between the provinces of Iga, Ise, and Ōmi meet.

- Vert. Japan, 1891, p. 65 (Mimasaku; Ise; Iga; Hida; Mino; Ivami; Tamba; Tajima).—BOETTGER, Kat. Batr. Mus. Senckenberg., 1892, p. 60 (Iga).—ANDRES, Atti Soc. Ital. Sci. Nat., XXXV, fasc. 3-4, 1896, p. 201, pl. 1 (descr.; general).—SCHNEE, Natur und Haus, VIII, 1900 (p. 246) (notes on living spec.).—ISHIKAWA, Mitth. Deutsch. Ges. Ost-Asiens, IX, Pt. 1, 1902, p. 81 (Hondo from Mino to Suwo and Nagato; habits, propagation, etc.); Proc. Nat. Hist. Tokyo Imp. Mus., I, No. 2, 1904, p. 19, pls. VIII-XI (general; distrib.; propag.; develop.).—KERBERT, Tijdschr. Nederland. Dierk. Vereen. (2), VIII, 1903 (pp. XXVIII-XXIX) (eggs); Zool. Anz., XXVII, 1904, Feb. 23, p. 305 (propagation, etc.); Compt. Rend. Sixth Int. Zool. Congr. Berne, 1904. (1905) p. 463 (propagation).—*Cryptobranchus maximus* CHAPMAN, Proc. Phila. Acad., 1893, p. 227 (anat.).
1840. *Hydrosalamandra siboldi* LEUCKART, Foriep's Neue Notizen, XIII, p. 20 (err. typ.).
1854. *Salamandra gigas* "Schlegel" DUMÉRIL and BIBRON, Erpét. Gén., IX, p. 164 (*lapsus*).
1854. *Tritomegas sieboldii* DUMÉRIL and BIBRON, Erpét. Gén., IX, p. 426 (err. typ.).
1907. *Cryptobranchus sieboldia* CALMETTE, Les Venins, p. 330, fig. 123.

The above quotations refer to Japanese specimens. A *Sieboldia davidiana* has been described by Blanchard^a from China, which Boulenger, however, regards as identical with the Japanese species. I have no means of verifying this identification. P. Krefft (in Verh. Ges. Deutsch. Naturf. Aerzte, 69 Vers. Braunsch., 1897, II, Pt. 1, 1898, p. 187) treats of specimens believed to be from Amoy and Canton, and refers them "auf Grund allerdings oberflächlicher Untersuchung" to *Cryptobranchus japonicus*. Gray, in 1873 (Ann. Mag. Nat. Hist. (4), XII, 1873, p. 188) mentions a skin including the bones of head and feet sent by Swinhoe from Shanghai, which he could not distinguish from a Japanese specimen.

There has been almost as much confusion about the specific name of the giant salamander as there has been with the generic term.

The oldest name is Temminck's *Triton japonicus*, which was published in his "Coup d'oeil," the introductory chapter to the Fauna Japonica, probably as early as March or April, 1837, and certainly not later than June of that year (see Bibliography, p. 542). In that paper which was penned as early as November, 1835, Temminck described the giant salamander sufficiently to give the name *Triton japonicus* a status in zoological nomenclature.^b

Description.—*Half grown*: U.S.N.M. No. 11349; Japan; Prof. E. S. Morse, collector. Vomerine teeth in an arched series between the

^a Compt. Rend. Acad. Paris, LXXIII, 1871, p. 79.—GRAY, Ann. Mag. Nat. Hist. (4), XII, 1873, p. 188.—*Sieboldia davidi* DAVID, Journ. Trois. Voy. Emp. Chinois, II, 1875, p. 20 (emendation) (S. W. Shensi).

^b La Salamandre gigantesque, *Triton japonicus* porte des formes bizarres: une tête extrêmement large et déprimée, un corps aplati muni d'une queue en forme de large aviron, le tout porté par des pieds très-courts; elle vit dans les eaux limpides des torrens. Coup d'oeil, p. xxvi.

choanæ, parallel with the maxillary and premaxillary series; whole body strongly depressed except terminal two-thirds of tail which is strongly compressed; head broad, flat; nostrils anterior, near the tip of the snout and the lips, their distance from each other less than one-half the distance between the eyes, which are without eyelids and very small, their diameter being less than one-tenth of the distance between them; legs short, depressed; fore legs with four fingers slightly webbed at base, the outer one with an outer dermal fold which continues along the entire length of the arm; hind feet with five toes webbed at base, the two outer ones with a distinct dermal flap on the outer side; a broad cutaneous expansion on the posterior aspect of the leg; tail short, slightly more than one-third the length of head and body, compressed, with a high dorsal fin beginning at the insertion of the hind legs, and a much lower ventral fin; a strong lateral fold on the neck from behind the angle of mouth to above the shoulder; another prominent undulating lateral dermal fold from the axilla to above the hind legs; skin rough, with numerous grooves, transverse on underside of body and on the sides, longitudinal on the throat; head above covered with closely set round tubercles which are somewhat smaller than the eyes, but not arranged with any degree of regularity or symmetry; lips, tips of snout, and a limited area along the median line of the top of head comparatively smooth; similar tubercles, though more scattered on sides of throat and on upper side of neck, a few occurring irregularly along the entire side of the body above the lateral fold. Color (in alcohol) burnt umber in various shades, paler below, irregularly blotched and marbled with dusky spots, limbs similarly spotted, tips of digits light orange brown.

Dimensions.

	<i>mm.</i>
Total length.....	490
Tip of snout to vent.....	322
Anterior border of vent to tip of tail.....	168
Width of head.....	69
Distance between eyes.....	40*
Fore legs.....	52
Hind legs.....	59
Greatest height of tail.....	48

Few salamanders have been oftener described in detail, or oftener figured than the present species as will be seen from the synonymy given above, where ample references are cited. A more detailed description therefore seems superfluous. It may be added, however, that there is no apparent external difference between the sexes, except that during the breeding season the borders of the vent are swollen in the male, but flat in the female. The young ones differ chiefly in being less rough and lighter in color.

Habitat.—The giant salamander of Japan seems to be restricted to the higher altitudes of the island of Hondo west of longitude $137^{\circ} 30'$ east of Greenwich. Rein states that it lives in clear running mountain streams in the granite and schist ranges, at a height of from 400 to 1,000 meters above the sea, while Geerts mentions 200 to 800 meters as its true habitat. According to these two authorities the principal localities where it is found are as follows:

1. The mountain streams and sources of the Yodogawa and Lake Biwa drainage, especially the tributaries of the Kidzugawa in the provinces of Iga, Yamato, and western Ise. The type-specimen—which von Siebold brought home alive to the Leiden Museum—was taken in this district, namely, in one of the streams of Sudzaka Yama, the mountain where the boundaries of the three provinces of Ise, Iga, and Omi meet, and Mr. Sasaki, who, in 1880 and 1881, visited this part of Japan in search of this species, obtained all his specimens (71) in Iga, Ise, and Yamato.

2. The streams of the border mountains between Hida and Mino provinces, especially the sources and tributaries of Rokagawa. Mr. Geerts quotes Keisuke Ito (*Nihon Sanbutu Si*, II, p. 39) as authority for the statement that it occurs in the province of Owari, though judging from the maps accessible to me it seems doubtful whether there are any localities high enough in that province.

3. The streams on the watershed between the districts of Sanindo and Sanyodo. Geerts mentions several streams by name, such as Miyadugawa and Itigawa, in the province of Tamba, Toyokagawa, in Tazima, and Osakigawa in Mimasaku and Harima, while Rein also mentions the province of Iwami. Okada likewise names the province of Iwami, in addition to those of Mimasaku, Tamba, Tajima, Ise, Iga, Mino, and Hida, but upon what authority I do not know.

Professor Ishikawa, in a lecture given before the German East Asiatic Society in Tokyo, 1900,^a gives the distribution of the giant salamander as follows: Mountain streams in the lower half of Hondo, i. e., from Mino to Iwami, Nagato, and Suwo. The species is consequently to be found in the mountain chain southwest from Mino to Suwo and Nagato, and also in the mountains of Iga and Ise, a branch of the main mountain chain. With regard to the provinces in which it occurs, the following are to be mentioned: Mino, Omi, Iga, Ise, Tamba, Tango, Tajima, Inaba, Mimasaku, Harima, Hoki, Bizen, Bitchu, Bingo, Idzumo, Iwami, Aki, Suwo, and Nagato. It has not been found as yet in Kii nor in Shikoku or Kiusiu. It occurs most frequently, so far as now known, in the mountain streams which come from the famous volcano Daisen, and in the streams on the south side of the Hiruzen Mountains and their neighborhood, espe-

^aMitth. Deutsch. Ges. Ost-Asiens, IX, p. 79.

cially in the sources of the rapid Asahigawa or river of the rising sun. For many years he collected the "hanzaki" in this part of the island, and he mentions the localities of Tsuyama, Onaru Valley, Kuginuki Valley, Mikamo, Maniwagori, all apparently in western Mimasaku. To his very interesting article, Contributions to the Knowledge of the Giant Salamander, in the Proceedings of the Department of Natural History, Tokyo Imperial Museum, I, 1904, he has added a colored map (Plate VIII) showing the distribution of the species. In this map the extension westward to the provinces of Nagato and Suwo is shown, and it also includes a large portion of the province of Idzumo. The text affords, however, no opportunity of judging upon what material this is based. This is particularly unfortunate as the testimony as to the occurrence of this species so far west is quite conflicting. According to a memorandum kindly furnished me by Dr. Hugh M. Smith, he was informed by Dr. T. Kitahara, zoologist of the Imperial Japanese Fisheries Bureau in Tokyo, that the western limit is the province of Bingo. The main drainage of the provinces of Tamba, Tazima, and Iwami is toward the Sea of Japan, the others drain to the inland sea, or to the Pacific, but the maps accessible to me are not accurate and detailed enough to decide whether some of the upper valleys of Iwami and Tazima do not in reality drain southward to the Inland Sea. At any rate it can not be taken for granted that the giant salamander occurs north of the watershed until its occurrence in streams of the Sea of Japan drainage is proven beyond a shadow of a doubt by actual specimens.

In this connection it may be added that, according to Doctor Smith's memorandum alluded to above, "Mr. I. Shishido, teacher of zoology in the Third High School, Kyoto, states that the species is reported from near Funatsa, province of Hida, in a river flowing into the Japan Sea (Toyama Bay), but no specimens from that locality are known to be in collections. The people in the vicinity of Funatsu, especially the women, eat this animal as a medicine." It is of the greatest importance that this question, whether the giant salamander occurs originally (not introduced) and regularly in the Japan Sea drainage, should be definitely solved. The qualification "regularly" is added because an isolated occurrence would not be convincing, since it might be explained either as an accidental introduction, or as the result of a Japan Sea stream having captured the sources of a Pacific Coast stream.

The capture of specimens in the Takadagawa, province of Musashi, and at Kyoto, in 1832, in the canal of the castle Nizyono siro, as related by Handangensei (Keisuke Ito.^a) probably refers to animals escaped from captivity or deliberately transplanted.

^aNihon Sanbutu, Sci. II, p. 39.

List of specimens of *Megalobatrachus japonicus*.

U. S. N. M. No.	Age.	Locality.	When collected.	By whom collected or from whom received.
6163.....	Young.....	Japan.....		Colonel Taylor.
6543.....	do.....	do.....		C. E. Smidt.
11349.....	Half grown ^a	do.....		E. S. Morse.
34214.....	Adult ^b	Province Mino.....		A. Owston.
34215.....	do.....	do.....		Do.
34216.....	do.....	do.....		Do.
34217.....	do.....	do.....		Do.

^a Description, p. 7.

^b 618 mm. long.

Superfamily SALAMANDROIDEÆ.

The Salamandroids fall into three families, namely, the Ambystomidæ, Plethodontidæ, and Salamandridæ. The Plethodontidæ are characterized by the presence of parasphenoid teeth. They do not occur within our area, being confined, with one exception, to the New World.

The two families found in Japanese territory may be distinguished as follows:

KEY TO FAMILIES.

- a*¹ Vomero-palatines prolonged posteriorly over the parasphenoid as two long processes, the teeth placed on the inner side of these forming two parallel or backward diverging series SALAMANDRIDÆ, p. 11.
- a*² Vomero-palatines not prolonged backward over the parasphenoid, the teeth placed on the posterior margin in transverse or posteriorly converging series. AMBYSTOMIDÆ, p. 24.

Family SALAMANDRIDÆ.

The salamanders, with the vomero-palatine teeth extending far backward in two parallel or posteriorly diverging series, are characteristic of the palearctic region, especially the Mediterranean subdivision. Three genera only occur in eastern Asia, two of which have been found within our territory. Of these, one, *Diemictylus*, extends into North America.

The Japanese genera may be distinguished as follows:

KEY TO GENERA.

- a*¹ Vomero-palatine teeth commencing in front of the line of the choanæ; series of isolated, knob-like glands on sides of back, some of these warts pierced by the extremity of the rib; maxillary bone reaching the quadrate; tip of pterygoid in contact with maxillary *Tylototriton*, p. 12.
- a*² Vomero-palatine teeth commencing on a line with the choanæ; no series of isolated knob-like glands on the sides of the back; maxillary bone separated from quadrate and from pterygoid by large intervals *Diemictylus*, p. 15.

Genus TYLOTOTRITON ^a Anderson.

1871. *Tylototriton* ANDERSON, Proc. Zool. Soc. London, 1871, p. 423 (type, *T. verrucosus*).
1885. *Tylotriton* BOETTGER, Offenbach. Ver. Naturk. 24-25 Ber., p. 165 (emendation).
1889. *Glossolega* COPE, Bull. U. S. Nat. Mus., No. 34, p. 201 (in part).

Thus far only two species of this genus are known, namely, *T. verrucosus*, which is known from the eastern Himalayas to the mountains of Yunnan, and the present species, which has been found only in Okinawa Shima. Doctor Anderson has given a detailed description, with illustrations, of the anatomy^b of the type species.

The nearest related genus appears to be *Pleurodeles*, the only species of which, *P. waltl*, is confined to the southwestern portion of the Pyrenean peninsula and the northern portion of Morocco. The two genera agree not only in the singularly pointed ribs protruding through the sides of the body, sometimes even perforating the skin, but also in the forward extension of the vomero-palatine teeth.

The present distribution of these forms—one in Spain, one in the Himalayan region, and one in the Riu Kiu Archipelago—is suggestive of the antiquity of this type of urodeles.

TYLOTOTRITON ANDERSONI ^c Boulenger.

1892. *Tylototriton andersoni* BOULENGER, Ann. Mag. Nat. Hist. (6), X, Oct. 1892, p. 304 (type-locality, Okinawa, Riu Kiu; type, Brit. Mus. No. 92. 9. 3. 30; Holst, collector).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 865; author's separate p. 16.—BOETTGER, Offenbach. Ver. Naturk., 33-36 Ber., 1895, p. 107 (Okinawa).


Description.—*Adult*; Sci. Coll. Tokyo, No. 67; Okinawa (figs. 1-6). Vomero-palatine teeth in two longitudinal series, meeting in front and commencing some distance anterior to a line through the anterior border of the choanae, then proceeding backward nearly parallel, but considerable distance apart, then at the beginning of the posterior third sharply diverging; tongue nearly circular, its diameter more than half the width of the mouth, extensively free on the sides, less behind and least in front; nostrils near the tip of the snout, and half-way between the top and the edge of the lip, their distance from each

^a From "τυλωτός, knobbed." "Along the body a lateral line of equidistant, large, rounded, knob-like, porous, glandular tubercles, terminating at the root of the tail. The second to fifth epipleural processes and the extremities of the remaining ribs terminate in the knob-like lateral glands." Anderson, Proc. Zool. Soc. London, 1871, p. 423.

^bZool. Res. Yunnan, p. 848, pl. LXXVII.

^c"Named after Dr. J. Anderson, to whom science is indebted for the discovery of the remarkable newt on which he established the genus *Tylototriton* in 1871." John Anderson was born in Edinburgh, October 4, 1833; superintendent of Indian Museum, Calcutta, from 1865 to 1886; then occupied with the study of the fauna of Egypt till his death on August 15, 1900. He visited Japan in 1884.

other greater than their distance from the eye, but smaller than the interorbital space; upper eyelid small, its width less than one-third the interorbital space; top of head surrounded on the sides and anteriorly by a rather rough, bony crest which covers the tip of the snout, canthus rostralis, supraorbital edge, and fronto-squamosal arch; parotoid gland scarcely as long as the width of the interorbital space, narrow, rather prominent; distance from tip of snout to gular fold contained less than three times in distance from gular fold to vent; fingers and toes interdigitating when legs are pressed against the sides of the body; a rough, bony ridge, similar to the one on the head, along the middle of the back, from occiput to the tail and continuous with the upper ridge of the latter: from the shoulder to the side of



FIGS. 1-6.—*TYLOSTOTRITON ANDERSONI*. $1\frac{1}{2} \times$ NAT. SIZE. 1, TOP OF HEAD; 2, SIDE OF HEAD; 3, OPEN MOUTH; 4, SECTION OF TAIL AT END OF BASAL THIRD; 5, HAND; 6, FOOT. No. 67, SCI. COLL., TOKYO.

the tail a dorso-lateral series of knob-like glands on the body terminating the pointed end of the ribs, which protrude as hard spines, though not perforating the skin; this series continues some distance on the side of the tail, on which it gradually disappears; between this series and the median dorsal ridge another series of prominent, conical, bony knobs, about eight on each side, each one located over a rib and connected along the latter with the terminal-rib knob by a series of smaller, less conspicuous, tubercles; fingers and toes rather short, depressed, with indication of a slight web at the base; fifth toe very short, rudimentary on the right foot, entirely missing on the left; no metacarpal or metatarsal tubercles; tail compressed, tapering from

the base, ending in a blunt point, upper and lower edge compressed, ridge-like; vent a short longitudinal slit with transverse folds or papillæ along the edges which are not raised or swollen; gular fold a slight depression; the whole body, including throat and legs above and below roughly granulated or tuberculated, the only smooth places being the lips, underside of hands and feet, and the lower edge of the tail. Color (in alcohol) above and below dark "mummy-brown," the bony ridges on head and back, as well as the two series of knobs on each side of the back, pale, like old ivory; underside of hands and feet, as well as lower edge of tail, yellowish.

Dimensions.

	<i>mm.</i>
Total length.....	159
Tip of snout to gular fold.....	21
Gular fold to anterior angle of vent.....	61
Anterior angle of vent to tip of tail.....	77
Width of head.....	21.5
Fore leg.....	25
Hind leg.....	27
Height of tail in the middle.....	6

Variation.—The specimen described above being only the second one known, it is obviously impossible to indicate the extent of individual variation in this species or even of sexual differences, as both specimens appear to be females. There are several points in which the two specimens differ, however. Thus, in the type the vomero-palatine series of teeth converge somewhat about the middle before the final flaring divergence. In the proportions there is, moreover, this difference, that in the type the tip of the toes reaches the elbow of the fore leg when the legs are pressed against the side; the length of the tail of the type is also proportionally greater. That the type is said to have an oval tongue and to be of a black color are differences of but slight consequence, but it is rather remarkable that the type does not appear to have the inner row of dorsal knobs differentiated to such a degree as to call for special mention in the original description.

Habitat.—The type of this species was discovered by Mr. Holst in Okinawa Shima, in 1892, and is in the British Museum. A second specimen, the one described above, belongs to the Science College Museum in Tokyo, the only ones thus far on record. This latter specimen is from the same locality as the type.

List of specimens of Tylototriton andersoni.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.
Sci. Coll. Tokyo.....	67	Adult fem. ^a	Okinawa shima.....		

^a Description, p. 12; figs. 1-6.

Genus DIEMICTYLUS^a Rafinesque.

1820. *Diemictylus* RAFINESQUE, Annals of Nature, 1820, No. 1, p. 5 (type, *T. viridescens*).
1820. *Notophthalmus* RAFINESQUE, Annals of Nature, 1820, No. 1, p. 5 (type, *T. miniatus*).
1838. *Cynops* TSCHUDI, Classif. Batr., p. 94 (type, *C. subcristatus*).
1850. *Taricha* GRAY, Cat. Batr. Grad. Brit. Mus., p. 25 (type, *T. torosa*).
1858. *Diemictylus* HALLOWELL, Journ. Phila. Acad. (2), III, p. 362 (emendation).
1878. *Triturus* BOULENGER, Bull. Soc. Zool. France, III, p. 307 (type, *T. viridescens*) (not of Rafinesque, 1815).
1879. *Pelonictes* LATASTE, Rev. Internat. Sci., III (p. 275), (type, *P. boscai*) (not of Fitzinger, 1843).

Professor Cope (Bull. U. S. Nat. Mus., No. 34, p. 203) has raised the question whether Rafinesque's *Diemictylus* has the priority over Merrem's *Molge* or not, both having been proposed in 1820. The question is immaterial, however, since *Molge* is a strict synonym of *Triturus*, Rafinesque (1815), both the latter names being assuredly only substitute names for *Triton*, preoccupied; hence they are in every respect identical and can not be used independently for different sections of the original genus *Triton*. *Molge* can never be used unless *Triturus* should become unavailable for some reason not now apparent. To the latter belong the species without a bony fronto-squamosal arch.

Two species of *Diemictylus* occur in Japan, one peculiar to Kiusiu, Hondo, and Shikoku, the other confined to the northern and central Riu Kius. The latter is a broader-headed form with shorter digits and longer tail, and Boulenger also indicates certain differences in the coloration of the live animals. The structural differences are very difficult to express in definite terms on account of the sexual and individual variability of these forms. It appears that the males have relatively longer digits than the females; consequently, in the female *D. pyrrogaster* the digits are of about the same relative length as in the male *D. ensicauda*. On the other hand, the length of the tail—though undoubtedly averaging considerably longer in *D. ensicauda*—varies to such an extent individually as to baffle all my attempts at using it diagnostically. The head of the latter is also decidedly larger, but the variability of the other proportions renders the use of it illusory as a unit for comparative measurements. Nevertheless, the two forms are quite distinct, and it may be that a larger series of *D. ensicauda* in good condition (many of my specimens being rather hard and dry) will yield better results in the future.

^a Derivation and meaning obscure. Two derivations suggest themselves, namely, *διαμυκτος*, from *διαμυγμα*, or *δι-ἡμικοτυλος*, but the application of neither is obvious.

In the meantime the following key must suffice:

KEY TO JAPANESE SPECIES OF DIEMICTYLUS.

- a*¹ Head smaller; digits longer; tail shorter; color in life, underneath red, underside of digits blackish..... *D. pyrrohogaster*, p. 16.
*a*² Head larger; digits shorter; tail longer; color in life, underneath, including underside of digits, yellowish..... *D. ensicauda*, p. 21.

DIEMICTYLUS PYRRHOGASTER ^a (Boie).



IMORI.

Plate II.

1826. *Molge pyrrohogaster* BOIE, Isis, 1826, p. 215 (type-locality, Japan; Blomhoff, collector); Bijdr. Natuurk. Wetensch., II, Pt. 1, 1827, p. 27.—*Molge pyrrohogastra* BOULENGER, Cat. Batr. Grad. Brit. Mus., 1882, p. 19 (Nagasaki; Kiukiang Mts. China).—OKADA, Cat. Vert. Jap., 1891, p. 65 (Tokyo; Kyoto; Kaga; Suwo).—BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 101; Kat. Batr. Mus. Senckenberg., 1892, p. 56 (Yokohama; Nagasaki).—SCLATER, List Batr. Indian Mus., 1892, p. 35 (Hakone; Lake Biwa).—*Cynops pyrrohogaster* GRAY, Cat. Batr. Grad. Brit. Mus., 1850, p. 25 (Japan).—BOULENGER, Bull. Soc. Zool. France, 1880, p. 39.—HILGENDORF, Sitz. Ber. Naturf. Fr. Berlin, 1880, p. 120 (Tokyo; prov. Kazusa; Hakone Mts.).—*Triton pyrrohogaster* STRAUCH, Mém. Acad. Sci. St. Pétersb. (7), XVI, No. 4, 1870, p. 51 (Japan; China?).—IWAKAWA, Quart. Journ. Microsc. Sci., XXX, 1882, p. —; Zool. Anz., V, 1882, p. 10.—*Salamandra pyrrohogastra* GEERTS, Nouv. Arch. Mus. Paris, (2) V, 1881, p. 275 (Japan).—*Diemyctylus pyrrohogaster* COPE, Bull. U. S. Nat. Mus., No. 34, 1889, p. 201.
1838. *Salamandra suberistata* SCHLEGEL, Fauna Jap. Rept., pp. 125-139; Saur. et Batr. pl. iv, figs. 1-3; *b* pl. v, figs. 7-8 (type-locality, Japan; types in Leiden Mus.; Siebold and Buerger, collectors); Abbild. Amph., 1844, p. 122, pl. XL, figs. 1-3 (color from life).—*Cynops suberistatus* TSCHUDI, Classif. Batr., 1838, p. 94, pl. II, figs. *a-c*.—*Triton suberistatus* DUMÉRIL and BIBRON, Erpét. Gén., IX, 1854, p. 140 (Japan).—HALLOWELL, Proc. Phila. Acad., 1860, p. 494 (in part: "Japan").—MARTENS, Preuss. Exped. Ost-Asien. Zool., I, 1866, pp. 109, 116; 1876, p. 384 (Yokohama).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, 1891, p. 239 (Yezo?).
1898. *Triton suberistatus* var. *typica* KREFFT, Verh. Ges. Deutsch. Naturf. Aerzte, 69 Vers. Braunsch., 1897, II, Pt. 1, p. 187 (Tokyo; Kyoto).
1898. *Triton suberistatus* var. *innaculiventris* KREFFT, Verh. Ges. Deutsch. Naturf. Aerzte, 69 Vers. Braunsch., 1897, II, Pt. 1, p. 187 (Kiusiu).
1906. *Triton pyrrohogaster* subsp. *typica* WOLTERSTORFF, Abh. Mus. Magdeburg, 1906, p. 162.

Description.—*Adult male*; U.S.N.M. No. 31796; Kyoto, Hondo, Japan; April 20, 1903; Dr. Hugh M. Smith, collector. Vomeropalatine teeth in two closely approximated longitudinal series beginning between the choanae and continued posteriorly to beyond a line between the corners of the mouth, in the anterior half nearly parallel, then strongly diverging; tongue small, its width about one-half that of the mouth; nostrils anterior, barely visible from above, their distance from each other considerably less than their distance from the

^aFrom *πυρρός*, fiery red; *γαστήρ*, belly. *b* Reproduced in this work, Plate II.



DIEMICTYLUS PYRRHOGASTER. (FROM SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 553.

eyes and only about two-thirds the width of the interorbital space, which is nearly twice that of the upper eyelid; body moderate, the distance from tip of snout to gular fold (fig. 7) being contained three times in the distance from gular fold to anterior border of vent; legs rather long, tip of longest finger reaching knee and tip of longest toe reaching elbow when the limbs are adpressed to the sides of the body; digits long, depressed, without webs, third finger longest, third and fourth toes nearly equal and longest, first finger and first toe shortest, shorter than fourth finger and fifth toe, respectively; tail strongly compressed (fig. 8), with a fin above and below, the outline of the fins being straight and nearly parallel, toward the tip tapering to a fine point, which is drawn out as a soft filament nearly 4 mm. long; skin strongly and mostly transverse rugose, except on anterior portion of head, on limbs, and tail; a low, smooth, vertebral ridge from nape to tail, being contiguous with the upper fin of the latter and bifurcating on the occiput; parotoid gland large, prominent, compressed, flap-like, almost continuous with a dorso-lateral glandular ridge cut by transverse grooves into a series of bead-like glands; a similar ventro-lateral ridge from axilla to groin; a large, smooth, globular gland on each side of the neck at each end of the gular fold; a somewhat smaller and flatter but well-defined circular gland above the insertion of the fore leg between the latter and the dorso-lateral ridge; vent a longitudinal slit surrounded by a large, globular swelling, the surface of which is coarsely pustular, the pustules near the posterior end of the vent being drawn out into long, pointed, conical papillæ. Color (in alcohol), above, uniform blackish brown; below, including the edge of upper lip, ventro-lateral ridge, and the anterior portion of the lower edge of the tail, yellowish white, with a series of irregular blackish blotches on each side of the belly and a few paler ones unsymmetrically on throat and fore neck; underside of hands and feet, dusky.



FIGS. 7-8.—DIEMICTYLUS. PYRRHOGASTER. 2X NAT. SIZE; 7, UNDERSIDE OF HEAD; 8, SECTION OF TAIL AT END OF BASAL THIRD. No. 6272, U.S.N.M.

Dimensions.

	<i>mm.</i>
Total length.....	93
Tip of snout to gular fold.....	12
Gular fold to anterior angle of vent.....	32
Anterior angle of vent to tip of tail exclusive of filament.....	47
Length of caudal filament.....	4
Width of head.....	10
Fore leg.....	18
Hind leg.....	20
Longest toe.....	7
Height of tail in the middle.....	7

The *adult female* (U.S.N.M. No. 31894; Kochi, Shikoku; May 11, 1903, Dr. Hugh M. Smith, collector) is larger, total length being 123 mm., and differs from the breeding male in the lesser development of the parotoid glands as well as the lateral glandular ridges, absence of the scapular gland, and the caudal filament. The body is also longer, the distance from tip of snout to gular fold being contained more than three times in the distance from gular fold to vent; so that the fingers and toes barely touch when the limbs are pressed against the side, the toes themselves also being shorter than in the male; finally, the vent is quite different, being a nearly circular opening with long radiating papillæ forming an oval swelling on a compressed elevated base.

The *nonbreeding male* and *female* (U.S.N.M. Nos. 30735, 30736; Miyazaki, Kiusiu) differ from the breeding ones chiefly in the lesser development of the glands and of the papillæ surrounding the vent, the male also by the absence of the caudal filamentous appendix.

Color of living specimens.—Dr. Hugh M. Smith has kindly furnished the following note on the color of the living specimens collected by him in Kiusiu and Shikoku during the breeding season. The back is black or blackish green; the underside of head, body, legs, and tail is blood red, with dark green mottling or streaks inclined to be symmetrical. In some specimens the back is lighter green and the sides of the tail may be steel blue.

Variation.—The chief variation is in the amount of dusky or blackish color on the underside. The normal markings seem to consist of a series of irregular spots on each side of the belly sometimes confluent to longitudinal streaks. These markings are often so heavy as to make the sides nearly uniform black, but the center of the belly is usually unspotted. On the other hand, some specimens are nearly devoid of spots. These form Kreff's variety *immaculiventris*, but they are not confined to any particular locality and represent only the extreme individual variation.

A male specimen taken in May, 1898, near Mount Fuji (U.S.N.M., No. 34308), is of a light grayish brown, almost unspotted underneath and seems to be a sort of albino.

Boulenger and Wolterstorff (Zool. Anz., XXX, 1906, p. 562) express the opinion that a more detailed study of *D. pyrrhogaster* might show the desirability of breaking it up into several subspecies, and the former mentions several breeding males and females from the island of Iki as being very small, as well as some Kobe specimens as having very weak parotoid glands. Whether the former island contains a particular dwarfed race it is impossible to say, as no measurements are given, but I do not believe it will be possible to split the specimens inhabiting the various large islands into geographical forms. The size and prominence of the parotoid glands seem to be more or less dependent on the season.

Habitat.—Widely distributed and common, inhabiting streams, ponds, reservoirs, and roadside drains, but probably confined to the lower altitudes.

Siebold^a reported it from Hondo and Shikoku as well as from Kiusiu. Doctor Fritze has stated that he found no specimens in Yezo, but that he saw some in the Sapporo Museum. Doctor Nozawa, however, has emphatically asserted (in letter dated February 27, 1901) that this is a mistake, that "there is no record of its presence [in Yezo] and, moreover, no specimen in the Sapporo Museum."

There are numerous specimens from Nagasaki in the various museums—for instance, in British Museum, the Hamburg Museum, and Museum Senckenbergianum. The United States National Museum has it from Miyazaki, province of Hiuga, Kiusiu, and Dr. Hugh M. Smith reports having observed it in the province of Satsuma.

The latter gentleman collected a fine series at Kochi, province of Tosa, Shikoku.

From Hondo we have numerous records. Thus, from Tokyo, Kyoto, Kaga, and Suwa (Okada); from Yokohama (von Martens; Mus. Senckenberg), Hakone (Indian Mus.; Hilgendorf), Tokyo, and province of Kazusa (Hilgendorf), Lake Biwa (Indian Museum). Jouy collected it on the way between Yokohama and Kobe, and the *Challenger* expedition brought home specimens from the latter place. In the Hamburg Museum there are specimens collected by T. Lenz in Kawachi Province, and Prof. R. Hitchcock sent us a specimen from Osaka. Dr. Hugh M. Smith collected three fine specimens in spring ponds near Kyoto and observed the species in the provinces of Ise and Shima, and from Mr. A. Owston we have a large series collected at Mount Fuji. Finally, British Museum has two specimens from "northern parts of Nippon."

There is no record of this species from Tsu-shima, but specimens from Iki-shima, between the latter and Kiusiu are in British Museum.^b

It would be highly interesting to know how far north this species goes in Hondo and also what its distribution is on the west coast. It is also of the greatest importance that its vertical range in the various provinces should be ascertained and recorded.

Diemictylus pyrrhogaster is said to occur in China, but the alleged Chinese specimens of this species are either of very dubious origin^c

^a Introd. Rept. Fauna Japon., p. xvi.

^b Boulenger, Zool. Anz., XXX, 1906., p. 562.

^c Thus the specimen in the museum in St. Petersburg referred to by Strauch (Mém. Acad. Sci. St. Pétersbourg, (7) XVI, No. 4, 1870, p. 51) and obtained from Doctor Schaufuss, the Dresden dealer, who was unable, however, to guarantee the correctness of the locality. The habitat "China" is the more suspicious in this case, as the museum in question also has a specimen of *Hynobius navius* from the same source and same alleged locality. (See Strauch, same reference as above, p. 55.)

or else belong to the Chinese representative, *Diemictylus orientalis* David^a recently redescribed by Woltersdorff.

Another nearly related species, *Diemictylus sinensis* (Gray) is also a native of China, the type having been collected "inland of Ningpo."

Boettger mentions four specimens from Hongkong in Museum Senckenbergianum. For difference from *D. pyrrhogaster* consult Boettger, Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 100.

Lately a third species has been discovered at Yunnan^{fu}, province of Yunnan, at an elevation of about 6,000 feet, namely, *Molge wolterstorffi* Boulenger.^b

List of specimens of Diemictylus pyrrhogaster.

U.S.N.M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Total length.
					<i>mm.</i>
6272a	Male ^a	Japan			80
6272b	do.	do.			88
6272c	do.	do.			87
6272d	do.	do.			83.5
6272e	Female	do.			98
14865	do.	Osaka		R. Hitchcock	114
17811	Male	Kobe to Yokohama	Aug. --, 1886	P. L. Jouy	78
17812	Female	do.	do.	do.	111
30735	Male ^b	Miyazaki, Kiusiu		C. A. Clark	77.5
30736	Female ^b	do.		do.	92.5
30737	do.	do.		do.	88.5
30738	do.	do.		do.	81
31794	do.	Kyoto, Hondo	Apr. 20, 1903	H. M. Smith	98
31795	Male	do.	do.	do.	93
31796	do.	do.	do.	do.	93
31894	Female ^b	Kochi, Shikoku	May 11, 1903	do.	123
31897	do.	do.	do.	do.	113
31898	do.	do.	do.	do.	104
31899	do.	do.	do.	do.	109
31900	Male	do.	do.	do.	83.5
31901	do.	do.	do.	do.	80
34288	Female	Mount Fuji, Hondo	May, 1898	A. Owston	119.5
34289	do.	do.	do.	do.	126
34290	do.	do.	do.	do.	116
34291	Male	do.	do.	do.	104
34292	do.	do.	do.	do.	100
34293	Female	do.	do.	do.	118.5
34294	do.	do.	do.	do.	122
34295	Male	do.	do.	do.	100
34296	Female	do.	do.	do.	116
34297	Male	do.	do.	do.	101

^a Figs. 7-8.

^b Description, p. 18.

^c Description, p. 16

^a *Triton orientalis* DAVID. Jour. Trois. Voy. Emp. Chinois. I. 1875, p. 32 (type-locality, Kioutchéou, Prov. Che-kiang, China); II, pp. 215, 238 (Tsitou, Prov. Kiangsi) (types in Paris Mus.).—*Triton pyrrhogaster* subsp. *orientalis* WOLTERSTORFF, Zool. Anz., XXX, Aug. 28, 1906, p. 558; Abh. Mus. Magdenburg, 1906, p. 153, pl. 1, figs. 3-6 (Chee-chou, Prov. Hupeh).

^b Proc. Zool. Soc. London, 1905. I. p. 177, pl. xvii.

List of specimens of *Diemictylus pyrrhogaster*—Continued.

U.S.N.M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Total length.
34298	Female	Mount Fuji, Hondo	May, 1898	A. Owston	103
34299	do	do	do	do	100
34300	Male	do	do	do	95.5
34301	do	do	do	do	93.5
34302	Male	Mount Fuji, Hondo	May, 1898	A. Owston	93
34303	do	do	do	do	95.5
34304	do	do	do	do	97
34305	Female	do	do	do	118
34306	Male	do	do	do	94
34307	do	do	do	do	92
34308	do ^a	do	do	do	110

^a Albino?

DIEMICTYLUS ENSICAUDA^a (Hallowell).

1860. *Triton subcristatus* HALLOWELL, Proc. Phila. Acad., 1860, p. 494 (in part; not of Schlegel).
1860. *Triton ensicauda* HALLOWELL, Proc. Phila. Acad., 1860, p. 494 (type-localities: Amakarima I. and "Ralousima . . . the northern half of Ousima proper;" types nos. 7410 and 7456, U.S.N.M.; Stimpson, collector).
1887. *Molge pyrrhogaster* var. *ensicauda* BOULENGER, Proc. Zool. Soc. London, 1887, p. 150 (Riu Kiu).—BOETTGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 106 (Okinawa).
1891. *Molge pyrrhogastra* var. *ensicaudatus* OKADA, Cat. Vert. Japan, p. 65 (Okinawa shima).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 865; author's separate, p. 16 (Okinawa).
1902. *Molge pyrrhogastra ensicauda* BROWN, Proc. Phila. Acad., 1902, June 11, p. 186 ("Iax) Choo Islands," probably Okinawa shima; Furniss and Hiller coll.).


Description.—*Adult male*: U.S.N.M. No. 7410a; cotype; O-shima, Riu Kiu Archipelago; William Stimpson, collector (figs. 9-15). Very similar to the male of *D. pyrrhogaster*, but proportions different, the head being larger, toes shorter, and tail longer; distance from tip of snout to gular fold contained less than three times in distance from gular fold to vent; fingers and toes interdigitate when limbs are pressed against the side; glands and glandular ridges much less developed. Color (in alcohol) underneath uniform yellowish; under-side of digits also yellowish.

Dimensions.

	mm.
Total length	111
Tip of snout to gular fold	14
Gular fold to anterior angle of vent	38
Anterior angle of vent to tip of tail	59
Width of head	12
Fore leg	20
Hind leg	20
Longest toe	4
Height of tail in the middle	6

^a From the Latin words, *ensis*, sword; *cauda*, tail.

This specimen is apparently not taken in the breeding season. Another *adult male* (U.S.N.M. No. 36554) appears to be in the breeding stage, however, as there is a distinct indication of a terminal caudal appendix and the globular swelling around the vent is very great with highly developed papillæ. Notwithstanding these signs the glands and glandular ridges are very poorly developed; the parotoids are



FIGS. 9-15.—*DIEMICTYLUS ENSICAUDA*. $2 \times$ NAT. SIZE. 9, TOP OF HEAD; 10, SIDE OF HEAD; 11, UNDER-SIDE OF HEAD; 12, OPEN MOUTH; 13, HAND; 14, FOOT; 15, SECTION OF TAIL AT END OF BASAL THIRD. NO. 7410a, U.S.N.M.

small, the gular spherical gland, the scapular circular gland, and the ventro-lateral ridge are not indicated.

The *females* appear to differ from the males much as do the *D. pyrrhogaster* females.

Color of living specimens.—It is stated that this species is yellow or

orange underneath and not blood-red, as *D. pyrrhogaster*. (See Stimpson's notes as given in the footnote.⁴)

Variation.—Our O-shima specimens seem to be browner, though this may be due to fading, and are unspotted underneath. On the other hand, the Okinawa specimens are blotched with blackish after the fashion of *D. pyrrhogaster*. Whether there is any constant difference in the coloration of the specimens from the two islands can not be decided from the specimens at hand, nor whether any structural difference exists between them. Large series of fresh material representing both species in the corresponding stages would be required for the purpose.

Habitat.—This salamander was first collected during Commodore Perry's expedition in May, 1853, by Mr. W. Heine, in the Loo Choo Islands, but it was first recorded from specimens brought home by the Rodgers North Pacific Surveying and Exploring Expedition. According to Hallowell's account of the herpetology of this expedition, specimens were collected by Messrs. Squires and Macomb in April, 1855, on Amakarima Isle (Kerama shima), while others were from the northern half of Oushima—i. e., Amami-o-shima. The latter were collected by W. Stimpson, but, as shown below, they probably came from Kakeroma shima. Since then various collectors have obtained it in the Riu Kius. Thus the U. S. National Museum has received two specimens (48a) from the Science College Museum, and there are three in the British Museum—one from one of Pryer's collectors, the two others from Holst. It has not been recorded

^a Hallowell's statements and Stimpson's manuscript notes are at variance, and it is impossible now to straighten out the tangle. Only one set of specimens brought home by the expedition has been preserved, viz, three specimens numbered 7410 U.S.N.M., which are uniform underneath without black blotches.

These, according to Hallowell (p. 494), should be the No. 53 of the (original) catalogue and come from "Ralousima" "the northern half of Oushima proper." Besides these he mentions "specimens marked No. 5, found in the paddy fields at the Amakarima Isle Loo Choo, April, 1855, by Mr. Squires and Mr. Macomb."

Now, turning to Doctor Stimpson's manuscript catalogue we find the following entries:

"Ou. 53. Salamander? found in muddy pond holes among the mountains, and in damp places in thickets [Oushima scratched out by W. S.] April [and substituted] Katonasima I. (the S. half of Oushima) (Wright) [here follows a sketch of the two islands—the northern inscribed Oushima, the southern Katonasima]. Above black, below deep vermilion with black spots and patches."

"Am.-5. Salamander? found in the paddy fields at the Amakirrima Isles (Loo Choo) April, 1855 (Mr. Squires and Mr. Macomb)."

In the first place, I can find on no map and in no gazetteer such a name as "Ralousima," and one wonders if Hallowell's manuscript can have had "real Oushima." In the second place, Stimpson's "Katonasima" is plainly Kakeroma shima. It then seems probable that Hallowell got mixed up and wrote northern where he should have written southern.

from the Yaeyama group, and consequently appears to be confined to the central group of the Riu Kiu.

List of specimens of Diemictylus ensicauda.

U. S. N. M. No.	Sex.	Locality.	When collected.	By whom collected	Total length.
					<i>mm.</i>
7410a	Male ^a	O-shima group Riu Kiu		W. Stimpson	111
7410b	do ^a	do		do	106
7410c	Female ^a	do		do	
7451a	Male	[Okinawa] Riu Kiu		W. Heine	105
7451b	Female	do		do	112
36554	Male ^b	Okinawa shima		Sci. Coll. Tokyo	104
36555	do	do		do	110

^a Cotypes; description, p. 21, and figures, p. 22.

^b Description, p. 21.

Family AMBYSTOMIDÆ.

Two subfamilies based upon cranial and hyoid characters may be recognized, the Hynobiinæ, which are peculiar to the eastern part of the Old World, and the Ambystominæ, which are almost entirely confined to the New World.

A word may here be said as to the name Ambystomidæ for this family. It is derived from the generic name *Amblystoma*, given by




FIG. 16.—DIAGRAM OF ARRANGEMENT OF VOMERINE TEETH IN ONYCHODACTYLUS, SALAMANDRELLA, AND GEOMOLGE. (FROM NIKOLSKI, ANN. MUS. ZOOL. ACAD. SCI. ST. PÉTERSBOURG, I, 1896, P. 78.)

Tschudi in 1838, generally emended to *Amblystoma* under the erroneous impression that the original spelling was due to a misprint, and that Tschudi intended to derive it from *αμβλυς*, blunt. This emendation is utterly unwarranted for not only does Tschudi use the word *Amblystoma* four times, but it is plainly a legitimate contraction for *Anamblystoma*, with allusion to the phrase *ανα στόμα βνειν*, to cram into the mouth. The shorter form of Ambystomidæ for Ambystomatidæ is justified by the perfectly correct practice in Greek to form composite words with nouns ending in *-μα* (genitive, *-ματος*) by leaving out the *τ* of the root (Kretschmer, *Sprachregeln*, 1899, p. 19, § 15), for instance, *στόμαργος*.

KEY TO THE GENERA INHABITING JAPAN AND EASTERN SIBERIA (fig. 16).

- a¹. Median angle of vomero-palatine series of teeth extending farther back than the lateral ends.
 - b¹. Toes five..... *Hynobius*, p. 25.
 - b². Toes four..... *Salamandrella*, p. 37.
- a². Vomero-palatine series not extending farther back in the middle than on the sides.

- b*¹. Vomero-palatine series continuous; digits (except of nonbreeding females) with black claws.....*Onychodactylus*, p. 42.
*b*². Vomero-palatine series interrupted in the middle; no claws.....*Geomolge*, p. 47.

Genus HYNوبيUS^a Tschudi.

1838. *Pseudosalamandra* TSCHUDI, Classif. Batr., pp. 56, 91 (type, *H. navius*).
 1838. *Hynوبيus* TSCHUDI, Classif. Batr., pp. 60, 94 (type, *H. nebulosus*).
 1839. *Molge* BONAPARTE, Icon. Fauna Ital., II, fasc. xxvi, fol. 131** (type, *M. navius*) (not of Merrem).
 1848. *Hydroscoptes* GISTEL, Naturg. Thierr., p. xi (substitute for *Pseudosalamandra*).
 1854. *Ellipsoglossa* DUMÉRIE and BIBRON, Erpét. Gén., IX, p. 97 (same type).

As will be seen from the above synonymy, *Pseudosalamandra* of Tschudi has a few pages precedence of his *Hynوبيus*. As E. D. Cope, the first author to formally combine the two genera under one of Tschudi's names, selected *Hynوبيus* for the greater group, the name so selected by him must stand, according to International Code of Zoological Nomenclature, art. 28.

The genus *Hynوبيus*, although far from peculiar to Japan, is nevertheless quite characteristic. Of the seven known species, four have not been found outside of the islands of Hondo, Shikoku, Kiusiu, and Tsu-shima; one has been found in Korea; and one is of uncertain locality, it being doubtful whether the only specimen known came from Japan or China. Thus far only one species is known with certainty from the latter country, namely, *H. chinensis* Guenther.^b

KEY TO THE JAPANESE AND KOREAN SPECIES OF HYNوبيUS.

- a*¹ Tail subcylindrical for more than the basal half, more or less compressed toward the tip, not keeled.
*b*¹ Fifth toe well developed.
*c*¹ "Body more than four times the length of the head; length of the [vomero] palatine series [of teeth] from anterior border to posterior angle, equals two-thirds or three-fourths the width of the tongue".....*H. navius*, p. 26.
*c*² "Body thrice and two-thirds the length of the head;" length of vomero-palatine series "from anterior to posterior angle equals one-half of the width of the tongue.".....*H. leechii*, p. 29.
*b*² Fifth toe rudimentary.....*H. peropus*, p. 32.
*a*² Tail compressed behind the vent, keeled.
*b*¹ Thirteen costal grooves.....*H. nebulosus*, p. 30.
*b*² Eleven costal grooves.
*c*¹ Fifth toe well developed; vomero-palatines with the median angle extending backward a considerable distance beyond the lateral arms.
H. nigrescens, p. 34.
*c*² Fifth toe rudimentary; vomero-palatines with the median angle not extending backward beyond the lateral arms.....*H. lichenatus*, p. 36.

^a From *ὄπις*, a ploughshare; *βίος*, life.


^b Ann. Mag. Nat. Hist. (6), IV, 1889, p. 222, from Ichang.

HYNOBIUS NÆVIUS^a (Schlegel).

Plate III, figs. 1-3.

- 1838.—*Salamandra nævia* SCHLEGEL. Fauna Jap., Rept., pp. 122, 129, Saur. et Batr., pl. iv, figs. 4-6;^b pl. v, figs. 9-10 (type-locality, Hondo and Shikoku; types in Rijksmuseum, Leiden; Siebold, collector); Abbild. Amphib., 1844, p. 122, pl. xxxix, fig. 4 (color from life).—GEERTS. Nouv. Arch. Mus. Paris, (2)V, 1883, p. 275.—*Ellipsoglossa nævia* DUMÉRI. and BIBRON, Erpét. Gén., IX, 1854, p. 99, pl. ci, fig. 5 (Japan, Siebold coll.).—STRAUCH, Mém. Acad. Sci. St. Pétersb. (7), XVI, 1870, p. 55 (Japan; China?).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, 1891, p. 239 (absent in Yezo).—*Hynobius nævius* COPE. Proc. Phila. Acad., 1859, p. 125.—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1876, p. 384 (Nagasaki).—OKADA, Cat. Vert. Jap., 1891, p. 65 (Kyoto).
- 1850.—*Molge striata* GRAY, Cat. Batr. Grad. Brit. Mus., p. 31 (not of Merrem 1820).

Description (figs. 17-22).—*Adult female*; U.S.N.M. No. 23901; Kumamoto, province Higo, Kiusiu; Nakagawa, collector. Vomero-







FIGS. 17-22.—HYNOBIUS NÆVIUS. 2 X NAT. SIZE. 17, TOP OF HEAD; 18, SIDE OF HEAD; 19, OPEN MOUTH; 20, SECTION OF TAIL AT END OF BASAL THIRD ($\frac{1}{4}$ NAT. SIZE); 21, HAND; 22, FOOT. SCI. COLL., TOKYO.

palatine series long, forming a very acute angle posteriorly, length of inner principal branch about equaling three-fourths the width of the tongue, the outer, short, recurring branch forming an acute angle; head rounded anteriorly, flat, somewhat concave behind the eyes; nostrils situated halfway between eye and tip of snout, their distance from each other equaling the interorbital space; eyes large, prominent, their longest diameter nearly equaling the interorbital space, which is

^a Latin, signifying spotted.

^b Reproduced in this work on Plate III, figs. 1-3.



Figs. 1-3, *HYNOBIUS NÆVIUS*; Figs. 4-6, *H. NEBULOSUS*. (FROM SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 553.

less than twice the width of the upper eyelid; distance from tip of snout to gular fold contained three times in distance from gular fold to anterior border of vent; limbs rather short, fingers and toes when laid against the body not meeting by about two costal folds; fingers and toes moderate, somewhat depressed, without webs; fifth toe well developed, about size of third finger, but slightly shorter than first toe; tail very short, its length from anterior end of vent only one-half the distance from the same point to tip of snout, subcylindrical, becoming compressed only a short distance from the tip, no keel above or below; skin everywhere smooth and glossy with numerous minute pits all over; fourteen distinct costal grooves including the axillar and inguinal ones; a median dorsal groove; parotoid gland flat, large, distinct, with a sharp, rounded upper edge; a distinct groove from eye to parotoid; gular fold strong; vent (female) a single longitudinal slit, borders not swollen. Color (in alcohol) pale cinnamon-rufous above, on sides and underneath gradually fading to pale pinkish buff, everywhere densely marbled and clouded with dusky, the markings being larger and coarse on the upper surface.

Dimensions.

	<i>mm.</i>
Total length.....	86
Tip of snout to anterior border of vent.....	57
Anterior border of vent to tip of tail.....	29
Tip of snout to gular fold.....	14
Width of head.....	10
Long diameter of eye.....	3
Fore leg.....	12
Hind leg.....	15
Width of tail at middle.....	5
Height of tail at middle.....	6

The *male* is said to have the vent consisting of a longitudinal slit "with a small papilla and a short transverse slit anteriorly."

Variation.—The length of the tail in the specimens measured by me varies to such an extent that I am not quite sure we here have to deal with only one species. The coloration also is very different in different specimens, and gives rise to similar doubts. But, unfortunately, the variation in the ratio between the length of the tail and that of the head and body is so gradual, as shown in the table below, that I do not see my way clear to draw a line, while on the other hand the two specimens with the greatest difference in tail proportion agree in color and differ from the specimens with intermediate proportions. The relative width and height of the tail is similarly varied, and is equally unsuited to furnish a demarcation line. One of the specimens, the one in the Science College, Tokyo, collection, has thirteen costal grooves, the others have fourteen.

The difference in the coloration is considerable. The Science College specimen is paler (more faded?) with less coarse marblings than the one described above, and the underside nearly uniform pale brownish. The three specimens in the Copenhagen Museum, however, are uniformly brown above, with large pale spots and coarse marblings on sides only. These specimens were probably received from the Leiden Museum, and thus verify Schlegel's remarks in the original description as follows: The system of coloration in this species seems to us to be rather constant in the large number of specimens which we have examined. This salamander is of a slaty or plumbeous color, uniform above, but varied with large whitish spots confluent like marblings on the flanks and on the lower parts, which often are paler or tending to yellowish. Finally, the colored figure reproduced in Schlegel's "Abbildungen," pl. xxxix, fig. 4, from a sketch from a living specimen, is black above with very distinct irregular whitish blotches on sides of head, body, and tail, and upper side of limbs. Boulenger's description of specimens in British Museum having the same origin as the above is similar, "blackish gray, lighter beneath; sides variegated with light gray."

In view of this uniformity it is difficult to understand the variation in the two specimens before me, and, I may add, in a third one in Science College from the same locality as ours described above.

Habitat.—Judging from Siebold's preface to the reptile volume of *Fauna Japonica* (p. xvi) he obtained the present species only in Hondo and Shikoku, but not in Kiusiu. It should be noted, however, that von Martens (*l. c.*) records it as obtained at Nagasaki by Doctor Pompe, and that the specimen in our museum (No. 23901) is from Kumamoto, province Higo, Kiusiu. These are the only definite records of locality which I have been able to find except Kyoto, as given by Okada. It seems certain, however, that it is absent in Yezo.

List of specimens of Hynobius navius.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Snout to vent.	Vent to tip of tail.	Snout to gular fold.	Width of head.	Width of tail.	Height of tail.	Ratio of head and body to tail.
						mm	mm	mm	mm	mm	mm	
U.S.N.M.	23901	Fem. ad. ^a	Kumamoto, Kiusiu	1884	Nakagawa.	57	29 14	10	5 6	1.96:1		
Copenh.	34	do.	"Japan"			71	45 16.5	12	5 7.5	1.58:1		
Do.	35	do.	do.			80	48 19.5	13	6 8	1.67:1		
Do.	36	do.	do.			82	61 18.5	14	7 11.5	1.34:1		
Sci. Coll.		do. ^b				58	49 14	11	5 7	1.18:1		

^a Description, p. 26.

^b Figs. 17-22.

HYNOBIUS LEECHII^a Boulenger.

Plate IV, fig. 8.

1887. *Hynobius leechii* BOULENGER, Ann. Mag. Nat. Hist. (5). XIX, p. 67 (type-locality, Gensan, Korea; type in Brit. Mus.; Leech, collector); Proc. Zool. Soc. London, 1890, p. 326.

Hynobius leechii is undoubtedly closely allied to *H. navius*. Boulenger hints at a difference in the proportion between head and body which I have been unable to appreciate while examining the type with the kind permission of the authorities of the British Museum.

The chief difference is perhaps in the shorter vomero-palatine series. It should be noted, however, that one of the British Museum specimens of *H. navius* approaches the *H. leechii* very closely in this respect. Boulenger's original description is herewith appended.

Original description of type specimen.—British Museum No. 86. 12. S. 14; Gensan, Korea; J. H. Leech, collector. Palatine teeth forming a V-shaped figure, which is broader than long; the length of one of the series, from anterior to posterior angle, equals one-half of the width of the tongue; head depressed, a little longer than broad; snout short, rounded; body thrice and two-thirds the length of the head; the distance from snout to gular fold contained nearly thrice in the distance from latter to cloaca; limbs not meeting when adpressed; fifth toe well developed; tail nearly as long as the distance between gular fold and vent, thick, compressed, not heeled, with vertical grooves, obtusely pointed posteriorly; skin smooth; thirteen costal grooves; a vertebral groove. Blackish brown, above speckled with pale brownish; upper surface of tail pale brownish, with a few black dots.

Dimensions.

	<i>mm.</i>
Total length.....	83
From snout to cloaca.....	47
Head.....	10
Width of head.....	8.5
Fore limb.....	12
Hind limb.....	14
Tail.....	36 ^b

Habitat.—A single specimen only has thus far been obtained. It formed part of a collection made by Mr. J. H. Leech at Gensan, Korea, and by him presented to British Museum in 1886.^c

^a For J. H. Leech, esq., who collected the type.

^b Boulenger, Ann. Mag. N. H. (5), XIX, 1887, p. 67.

^c While there is nothing intrinsically improbable in this species being Korean, it may be well (until the find is corroborated) to refer to what is said about a specimen of *Elaphe madrivirgata* alleged to have been collected by Mr. Leech at Gensan (see p. 331).

List of specimens of *Hynobius leechii*.

Brit. Mus. No.	Sex and age.	Locality.	When collected.	By whom collected.
86.12.8.140		Gensan, Korea.		J. H. Leech.


^a Type.

HYNOBIUS NEBULOSUS^a (Schlegel).

Plate III, figs. 4-6.

1838. *Salamandra nebulosa* SCHLEGEL, Fauna Jap., Rept., pp. 127, 139, Saur. et Batr., pl. iv, figs. 7-9^b (type-locality, "Mitsujama" near Nagasaki, Kiusiu; types in Rijksmuseum, Leiden; Siebold, collector); Abbild. Amphib., 1844, p. 126, pl. xl, figs. 7-10 (color from life).—GEERTS Nouv. Arch. Mus. Paris (2), V, 1883, p. 275.—*Hynobius nebulosus* TSCHUDI, Classif. Batr., 1838, pp. 60, 94.—GRAY, Cat. Batr. Grad. Brit. Mus., 1850, p. 30 (Japan).—BOULENGER, Cat. Batr. Grad. Brit. Mus., 1882, p. 32 (Nikko).—OKADA, Cat. Vert. Japan, 1891, p. 65 (Nikko).—SCLATER, Batr. Ind. Mus., 1892, p. 36 (Yezo(?)).—BOETTGER, Kat. Batr. Mus. Senckenberg., 1892, p. 52 (Nagasaki).—*Ellipsoglossa nana* DUMÉRIER and BIBRON, Expét. Gén., IX, 1854, p. 100 (Japan).—STRACH, Mém. Acad. Sci. St. Pétersbourg (7), XVI, 1870, p. 55 (Japan).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, 1891, p. 239 (absent in Yezo).

Description.—*Adult male*; Science College Museum, Tokyo, No. 51; Nikko (figs. 23-26). Vomero-palatine series of teeth long, forming



FIGS. 23-26.—*HYNOBIUS NEBULOSUS*. 2 × NAT. SIZE. 23, TOP OF HEAD; 24, SIDE OF HEAD; 25, OPEN MOUTH; 26, SECTION OF TAIL AT END OF BASAL THIRD. No. 51, SCI. COLL. TOKYO.

an acute angle posteriorly, length of inner principal branch equaling one-half the width of the tongue, the outer, short, recurving branch forming an arch with the inner one; head flat, broad; nostrils situated much nearer the tip of snout than the eyes, their distance from each other equaling the interorbital space; eyes moderate, their long diameter about equaling two-thirds the width of the interorbital space, which is at least twice

the width of the upper eyelid; distance from tip of snout to gular fold contained three and one-fourth times in distance from gular fold to anterior border of vent;

^a Latin, signifying clouded.

^b Reproduced in this work on Plate III, figs. 4-6.

limbs moderate, fingers and toes when laid against the body overlapping; fingers and toes rather long, depressed, without webs; fifth toe well developed, nearly equaling first toe, though not as long as third finger; tail long, equaling the length of head and body, compressed from the base, tapering, pointed at end, with a dorsal fin beginning at the basal first third and a lower ventral fin in the terminal fourth; 13 distinct costal grooves including the axillar and inguinal ones; a median dorsal groove; parotoid gland indistinct; postocular groove indistinct; gular fold well marked; vent (male) a longitudinal slit bifurcating anteriorly, with a papilla in the fork, borders swollen. Color (in alcohol) above dull isabella-color, the upper surface with indistinct paler marblings; underneath pale drab-gray; the entire surface above and below dusted over with minute dusky specks visible under the magnifying glass.

Dimensions.

	<i>mm.</i>
Total length.....	135
Tip of snout to anterior border of vent.....	68
Anterior border of vent to tip of tail.....	67
Tip of snout to gular fold.....	16
Width of head.....	12
Long diameter of eye.....	3
Fore leg.....	22
Hind leg.....	20
Width of tail one head-length from vent.....	4
Height of tail one head-length from vent.....	6

Variation.—An adult specimen in British Museum (No. 92. 1. 11. 18) collected by Holst in Tsu-shima, October 6, 1891, has a somewhat longer series of vomero-palatines, inasmuch as the length from angle to angle is about two-thirds the width of the tongue. The proportions of head and body are nearly the same—viz, snout to gular fold 16 mm. and gular fold to vent 52 mm., or the latter distance three and one-fourth times the former. The tail, however, is much shorter, being less than the body without head. The color (in alcohol) is also different, being above yellowish brown with numerous obscure dusky spots; underneath uniform pale brownish, this pale effect being produced by a uniform dusting of minute, round, yellowish specks on a dark brownish gray ground. The coloration of this species is usually described as more strongly marbled and variegated than the above individuals, in fact much more like the two specimens described above as *H. navius*. It should be noted, however, that the colored reproduction of the drawing from a living specimen in Schlegel's "Abbildungen," pl. XI, figs. 7 and 8, shows an animal uniformly dark brown above, more grayish and paler underneath.

Remarks.—This species differs chiefly from *H. navius* in the shorter body with longer limbs, so that the fingers and toes overlap on the side, and in the more pointed tail, which is compressed all the way from the base and keeled and finned above and below toward the tip.

Aberration?.—A single specimen (No. 33) in the Zoological Museum in Copenhagen, which I had the opportunity to examine in 1905 owing to the kindness of Inspector Jensen, was received in 1847, probably from Schlegel in Leiden, and is labeled "Japan." It is in every respect a typical *H. nebulosus*, except that it has only 4 well-developed toes, the rudiment of the fifth toe being scarcely noticeable. It consequently holds the same relation to *H. nebulosus*, as *H. peropus* to *H. nævius*; in fact, the rudiment of the fifth toe is even smaller than in Boulenger's figure of *H. peropus*.^a It has the vomeropalatine series very long and 13 costal grooves.

The lack of the toe being the only character, I do not regard this specimen as representing a distinct form or species.

Habitat.—This species was originally described from specimens collected by von Siebold at Mitsuyama, near Nagasaki. A specimen from the latter place is also in the Senckenberg Museum. Siebold also indicates its occurrence in Hondo and Shikoku, though somewhat indirectly.^b A young specimen collected by C. Maries at Nikko, Hondo, is in British Museum, and Nikko specimens are also in the Science College Museum in Tokyo (No. 51).

Selater records three specimens in the Indian Museum as collected by Dr. J. Anderson in Yezo, but this is probably an error for Yedo, as there is no corroboration of its occurrence in the northern island.

British Museum has specimens (No. 92. 1. 11. 18; female and larvæ) collected by Holst in Tsu-shima, the female on October 6, 1891, the larvæ on August 7.

List of specimens of Hynobius nebulosus.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Total length.	Snout to vent.	Vent to tip of tail.	Snout to gular fold.	Width of head.
						mm	mm	mm	mm	mm
Sci. Coll. Tokyo.	51	Male ^a	Nikko.			135	68	67	16	12
Copenhagen	33	do. b.	"Japan"			95	54	41	12	9
Brit. Mus.	92.1.11.18	Female c.	Tsu-shima.	Oct. 6, 1891	P. Holst.	117	68	49	16

^a Description, p. 30; figs. 23-26.

^b See p. 32.

^c See p. 31.

HYNوبيUS PEROPUS c Boulenger.

Plate IV, figs. 4-7.


1882. *Hynobius peropus* BOULENGER, Cat. Batr. Grad. Brit. Mus., p. 33, pl. II, figs. 1-1a^d (type-locality uncertain, China or Japan; type in Brit. Mus.; A. Adams, collector).

^a Reproduced on Plate IV (figs. 4-5) of this work.

^b Introd. Rept. Fauna Japon., p. xvi.

^c From *πηρος*, mutilated; *πούς*, foot.

^d Reproduced in this work on Plate IV, figs. 4-7.



FIGS. 1-3, *HYNOBIUS LICHENATUS*; FIGS. 4-7, *H. PEROPUS*; FIG. 8, *H. LEECHII*.
(FROM BOULENGER.)

FOR EXPLANATION OF PLATE SEE PAGE 553.

The exact locality of the type-specimen is unknown, but a specimen in the Hamburg Museum (No. 1020), collected by T. Lenz, in 1896 in Yamato, shows that the species is Japanese. Thanks to the kindness of the authorities of the British Museum, I have had the opportunity to examine the type. As stated by Boulenger, the fifth toe is quite rudimentary, even more so than in *H. lichenatus*. It has nothing to do with the latter species, however, as it has thirteen costal grooves, longer series of vomerine teeth, and a cylindrical tail. This latter character places it near to *H. nævius*. The chief difference from the latter in the shape of the tail consists in its being slightly longer, and the terminal portion more compressed and pointed. The vomerine dental series is also shorter, and in *H. nævius* the fifth toe is well developed.

Having nothing further to add to Boulenger's original description, it is here reproduced.

Original description of type specimen.—*Adult male*; British Museum; China or Japan; A. Adams, collector. The length of the palatine series, from anterior border to posterior angle, equals half the width of tongue; head depressed, longer than broad; snout short, rounded; eyes rather large, prominent; no labial lobes; body cylindrical, slightly depressed, nearly four times the length of the head; the distance from snout to gular fold three times in the distance from latter to cloaca; limbs moderate, not meeting when adpressed; fingers and toes more elongate than in the preceding species; fifth toe rudimentary, tubercle-like; carpal and tarsal tubercles indistinct; tail shorter than head and body, strongly compressed and keeled in its posterior half, ending in a point; vent (of male) three slits meeting in front, the median longitudinal and longest, the two others obliquely directed forward, forming an angle; the borders not swollen; skin smooth; thirteen costal grooves; a median dorsal groove; parotoids indistinct; gular fold distinct; no fold from eye to gular fold. Reddish brown above, speckled and minutely marbled with dark brown; tail dark brown, with lichen-like gray variegation; yellowish brown beneath; belly speckled with lighter.

Dimensions.

	<i>mm.</i>
Total length.....	109
From snout to cloaca.....	59
Head.....	12
Width of head.....	9
Fore limb.....	15
Hind limb.....	17
Tail.....	50 ^a

Habitat.—As already stated, the exact locality where Mr. Adams collected the unique specimen which Boulenger described and figured

^a Boulenger, Cat. Batr. Grad. Brit. Mus., 1882, p. 33.

is not known, and as he collected both in China and Japan it may have come from either of these countries. The probability is, however, that it really came from Japan, since a specimen in the Hamburg Museum was collected in the province of Yamato, Hondo.


List of specimens of Hynobius peropus.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.
Brit. Mus.	(a)	Male ^a	"China or Japan"		A. Adams.

^a Type; description, p. 33; plate IV, figs. 4-7.

HYNOBIUS NIGRESCENS, ^a new species.

Diagnosis.—Series of palatine teeth typical, the length equaling four-sevenths the width of the tongue; fifth toe well developed; tail



FIGS. 27-32. *HYNOBIUS NIGRESCENS*. 2 × NAT. SIZE. 27, TOP OF HEAD; 28, SIDE OF HEAD; 29, OPEN MOUTH; 30, SECTION OF TAIL AT END OF BASAL THIRD; 31, HAND; 32, FOOT. NO. 57A, SCI. COLL. TOKYO.

long, strongly compressed, keeled above and below; eleven costal grooves.

Habitat.—Sendai, Hondo, Japan.

Type.—Science College Museum, Tokyo, No. 57A.

Description of type-specimen.—*Adult male* (figs. 27-32). Shape of palatine series of teeth typical, though more U-shaped than V-shaped,

^a Latin, signifying blackish.

the posterior branches but slightly converging, the posterior end of each making a sudden bend to meet its fellow on the other side; the extreme length of the series equals four-sevenths the width of the tongue; head large, depressed, as broad as long; eyes rather large, prominent; body medium, distance from snout to gular fold contained less than three times in distance from gular fold to anterior end of vent; limbs well developed, fingers and toes interlocking when laid against the body; fingers and toes moderate, depressed, fifth toe well developed; tail long, the distance from tip to anterior end of vent exceeding that from the latter point to end of snout, strongly compressed, pointed, keeled above and below, the upper keel originating almost opposite the posterior end of the vent; skin smooth, except on the tail, where it is rather strongly rugose; sides with eleven very distinct costal grooves; a medium dorsal groove; parotoids indistinct; gular fold strongly marked underneath, but not on the sides; no groove from eye to gular fold; no groove behind angle of the jaws; vent (of male) a longitudinal slit meeting two short, lateral, converging slits anteriorly; a large papilla at the anterior end; a distinct transverse groove at the posterior end; borders greatly swollen. Color uniform blackish above, the whole surface finely dusted over with minute pale specks, below considerably paler, the specks predominating over the black ground; tail uniformly dark, including the keels.

Dimensions.

	<i>mm.</i>
Total length.....	150
From snout to gular fold.....	20
From gular fold to vent.....	54
Vent to tip of tail.....	76
Width of head.....	14
Fore limb.....	25
Hind limb.....	25

An adult *female* in the Science College Museum (No. 52A) agrees with this species in many points especially in the number of costal folds, which are only eleven. It differs, however, in lacking the fin on the tail. This may be seasonal, and in the absence of precise information as to the seasonal changes of the various species of *Hynobius*, I am inclined to regard it as belonging to the present species. It measures, total length, 108 mm.; snout to vent, 59; vent to tip of tail, 49; tip of tail to gular fold, 15.5; axilla to groin, 28; width of head, 11.

Habitat.—The only locality where this species has been found thus far is Sendai, Hondo, whence came the type and four other specimens in the Science College Museum. A specimen doubtfully referred to this species is from Nikko.

Remarks.—In its compressed tail and only eleven costal grooves this new species approaches *H. lichenatus*. The latter has a very short vomerine series and the fifth toe rudimentary.

List of specimens of Hynobius nigrescens.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Total length.	Snout to vent.	Vent to tip of tail.	Snout to gular fold.	Width of head.
						mm.	mm.	mm.	mm.	mm.
Sci. Coll. Tokyo	57A	Male ^a	Sendai, Hondo.			150	54	76	20	14
Do.	52A	Female ^b	Nikko, Hondo.			108	59	49	15.5	11

^a Type; description, p. 34; figs. 27-32.^b P. 35.**HYNOBIUS LICHENATUS^a (Boulenger).**

Plate IV, figs. 1-3.

1883. *Hynobius lichenatus* BOULENGER, Ann. Mag. Nat. Hist. (5), XII, p. 165, pl. v, fig. 1^b (type-locality, Aomori, Hondo; type in Brit. Mus.; G. Lewis, collector); Proc. Zool. Soc. London, 1886, p. 413.

With the permission of the authorities of the British Museum I examined the hitherto unique type of this species in 1898. It is chiefly characterized by possessing only eleven costal grooves and only a rudiment of the fifth toe, which is but slightly larger than the one in *H. peropus*. Moreover, the vomerine dentition is quite peculiar in forming only a slight median posterior prolongation. If this character is normal, *H. lichenatus* is the most distinct species in the genus. Its eleven grooves and compressed tail otherwise bring it near to *H. nigrescens*, which, however, has a well-developed fifth toe.

Boulenger's original description is herewith appended.

Description of type-specimen.—British Museum; Aomori, Hondo; George Lewis, collector.—In the shape of the series of palatine teeth intermediate between *Hynobius* and *Onychodactylus*; these series form a zigzag row, the central or posterior angle not extending posteriorly beyond the extremity of the outer branches; head large, depressed, as broad as long; snout short, rounded; eyes rather large, prominent; no labial lobes; body short, a little more than thrice the length of head; the distance from snout to gular fold contained a little more than twice and a half in the distance from latter to cloaca; limbs moderate; when laid against the body the fingers cross the toes; fingers and toes moderate, depressed; fifth toe rudimentary, as in *Hynobius peropus*; tail about as long as head and body, strongly compressed and keeled superiorly and inferiorly, ending in a blunt point; skin smooth, shining; eleven costal grooves; a median dorsal groove; parotoids rather indistinct, porous; gular fold strongly marked; a distinct groove behind the angle of the jaws. Brown

^a Covered with lichens, referring to the "lichen-like grayish variegation" of color.^b Reproduced in this work on Plate IV, fig. 1-3.

above, lighter beneath, with lichen-like grayish variegation; whitish dots on the sides of the body.

Dimensions.

	<i>mm.</i>
Total length.....	74
From snout to cloaca.....	38
Head.....	9
Width of head.....	8.5
Fore limb.....	13
Hind limb.....	14
Tail.....	36 ^a

Habitat.—Thus far only a single specimen has been collected at Aomori, near the northern extremity of Hondo.

List of specimens of Hynobius lichenatus.

Brit. Mus. No.	Sex and age.	Locality.	When collected.	By whom collected.
(a)		Aomori, Hondo.....		George Lewis.

^aType; description, p. 36; plate VI, figs. 1-3.

Genus SALAMANDRELLA ^b Dybowski.

1870. *Salamandrella* DYBOWSKI, Verh. Zool.-Bot. Ges. Wien, XX, 1870, p. 237 (type, *S. keyserlingii*).

1870. *Isodactylum* STRAUCH, Mém. Acad. Sci. St. Pétersbourg, (7) XVI, No. 4, p. 55 (type, *I. schreckii*=*S. keyserlingii*).

This genus is not represented in Japanese territory, unless it should be found on some of the northern Kuril islands, such as Paramushir or in south Sakhalin. So far as known it occurs only in south-eastern Siberia from Lake Baikal to Ussuri and also in southern Kamchatka.

Salamandrella differs from *Hynobius* chiefly in the complete absence of a fifth toe. In view of the rudimentary state of this member in several species of the latter genus, and its occasional total suppression in individuals, it is a great question whether the genus can be maintained in the future.

SALAMANDRELLA KEYSERLINGII ^c Dybowski.

Plate V, figs. 7-8.

1859. *Triton* nov. sp. MAACK, Putesh. Amur (p. 154).

1870. *Salamandrella keyserlingii* DYBOWSKI, Verh. Zool.-Bot. Ges. Wien, XX, 1870, p. 237, pl. VII (type-locality, southwest corner of Lake Baikal).—
STRAUCH, Mém. Acad. Sci. St. Pétersbourg, (7) XVI, No. 4, 1870, p. 110.—

^a Boulenger, Ann. Mag. Nat. Hist. (5), XII, 1883, p. 165.

^b Diminutive form of *Salamandra*.

^c For Count Alexander Keyserling, the Russian explorer and zoologist. Born August 15, 1815, at Kabillen, Kurland; died May 25, 1891, in Dorpat. With Blasius joint author of "Die Wirbelthiere Europas" (1840).

- BOULENGER, Cat. Batr. Grad. Brit. Mus., 1882, p. 34 (Shilka; Baikal); Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 144 (Khabarovka, Doerries collector).—MUELLER, Verh. Naturf. Ges. Basel, VIII, Pt. 2, p. 250 (Khabarovka).—BEDRIAGA, Wiss. Result. Przewalski Zentralasiat. Reise, Zool., III, Pt. 1, lief. 1, Amph., 1898 (p. 8) (Ksernzo, Prov. Sze-Chuan, China).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, (8) XVII, No. 1, 1905, p. 436 (Yekaterinburg to Kamchatka).—*Salamandrella keyserlingi* WOLTERSTORFF, Verh. V Internat. Zool. Congress, Berlin, 1901 (1902), p. 585 (Verkhoyansk).
1870. *Isodactylum schrenkii* STRAUCH, Mém. Acad. Sci. St. Pétersbourg (7), XVI, No. 4, 1870, pp. 56, 110, pl. II, fig. 1 (type-locality, east Siberia: Ussuri, Shilka, and Lake Baikal; types in St. Petersburg Mus.; Schrenck, collector).—SABANEFF, Bull. Soc. Nat. Moscou, XLIV, Pt. 2, 1872, p. 275 (Yekaterinburg, Ural).—*I. schrenki* SHITKOV, Zool. Anz., XVIII, May 6, 1895, p. 165 (Yekaterinburg, Ural Mts.; reproduction and development).
1870. *Isodactylum wosnessenskyi* STRAUCH, Mém. Acad. Sci. St. Pétersbourg (7), XVI, No. 4, 1870, p. 58, pl. II, figs. 2a-da (type-locality, Javina, southern Kamchatka; types in St. Petersburg Mus.; I. G. Wosnessensky, collector).—*Salamandrella wosnessenskyi* BOULENGER, Cat. Batr. Grad. Brit. Mus., 1882, p. 34 (Kamchatka).
1896. *Salamandrella uralensis* NIKOLSKI, Annuaire Mus. Zool. St. Pétersbourg, I, p. 13 (*nomen nudum*).
1905. *Salamandrella keyserlingii* var. *tridactyla* NIKOLSKI, Zap. Imp. Acad. Nauk, S. Peterburg (8), XVII, No. 1, p. 491 (type-locality, Vladivostok; type Mus. Acad. St. Petersburg, No. 2279; Palchevski, collector).

Until recently *S. keyserlingii* and *S. wosnessenskyi* have been considered distinct species. According to Strauch the main differences between them as recognized by him are as follows:

Tail at most seven times as long as its height at the middle; the outer branch of each vomerine row of teeth bent straight backward.....	<i>S. keyserlingii</i>
Tail at least nine times, or even ten times, as long as its height at the middle; the outer branch of each vomerine row of teeth not bent backward but directed outward.....	<i>S. wosnessenskyi</i>

Neither of these characters holds in a large series of specimens.

In seven, mostly well preserved, Kamchatkan specimens which consequently represent *S. wosnessenskyi* I find both extremes of palatine dentition as described above and figured by Strauch. It may even be stated that the extreme obtuseness of the lateral angles as shown in his figure of the skull^b of the Kamchatkan species is not equaled by any of the specimens at hand, while one, at least (U.S.N.M. No. 31716), has an open median angle and acute anterior angles in excess even of Strauch's figure of *S. keyserlingii*.^c

The distinction derived from the alleged lesser elevation of the tail in *S. wosnessenskyi* also falls to the ground, as an inspection of the subjoined table of dimensions (p. 41) shows. None of my seven specimens has the tail more than nine times as long as high, and only two

^a Reproduced in this work on Plate V, figs. 7-8.

^b Mém. Acad. Sci. St. Pétersbourg (7), XVI, No. 4, pl. II, fig. 2d.

^c Idem., fig. 1.

reach this proportion; in two it is between seven and eight times, while in three it is only seven, the proportion given by Strauch for *S. keyserlingii*.


Strauch, in the description of *S. wosnessenskyi*, mentions several other slight characters such as can only be appreciated with specimens of both species for actual comparison. Thus the snout is said to be shorter and much more bluntly rounded; the tongue somewhat shorter in proportion to its width; the lateral and ventral folds to be better developed. The alleged difference in their number, viz, 12 in *S. keyserlingii* and 13 in *S. wosnessenskyi* he admits is inconstant, and as a matter of fact I find the latter number to be the exception in my series. Strauch also intimates that the Kamchatkan form has longer extremities inasmuch as the fore limbs when bent forward and pressed to the side of the body reach beyond the anterior border of the eye, but this proportion is reached by only one of my specimens (No. 31716); in the others it falls considerably short of this point, and in one (No. 31713) it reaches barely beyond the posterior border of the eye.

Finally, the coloration seems to be identical in both forms, as I have found nothing in the description of true *S. keyserlingii* which would separate it from the Kamchatkan form as far as color is concerned.

While thus unable to distinguish the two species from the descriptions, I was unwilling to unite them, as I have only specimens of one of them. I now learn, however, from Professor Nikolski that he has convinced himself not only that *S. uralensis*, which was published by him as a *nomen nudum* but also *S. wosnessenskyi* are identical with *S. keyserlingii*, and that he has so considered them in his new work "Herpetologia Rossica" (1905). Under these circumstances I can see no reason for maintaining two distinct headings for this species. Prof. B. Shitkov has also informed me (letter of May 18, 1906) that he regards the *S. wosnessenskyi* at most as a variety.

Description.—*Adult*; U.S.N.M. No. 22594; Rakovaya Bukhta, Avatcha Bay, Kamchatka; L. Stejneger, collector (figs. 33–38). Vomeropalatine teeth in an angular series, the median angle being directed backward about 45°, and its arms about as long as half the width of the tongue, the lateral retroverted angles much more obtuse, or about 60°; head flat; nostrils about halfway between eye and tip of snout, their distance from each other more than the interorbital width; eye large, upper eyelid longer than interorbital width and slightly wider than one-half the same; body elongate, the head from tip of snout to gular fold being contained three and one-half times in the distance from gular fold to anterior border of vent; fingers and toes of adpressed limbs do not meet by a distance of four costal folds; limbs rather slender, digits short, four on each foot, scarcely webbed at base; tail less than distance from tip of snout to vent, compressed from behind the vent, not keeled, the upper outline straight to the

tip, the lower one parallel with the upper until near the tip, toward which it ascends rapidly; thirteen very distinct costal grooves which are continuous across the belly; median dorsal groove not distinct; parotoid gland large, flat, distinct; a distinct groove from posterior angle of eye to the groove in front of the parotoid, continuing over the gland and behind the latter, joining the gular fold; vent a longitudinal slit with several lateral folds posteriorly, borders swollen. Color (in alcohol), top of head and a broad dorsal band extending the whole length of the tail, as well as the legs above, tawny-olive; sides and entire under surface dirty pale olive-buff; an irregular blackish streak bordering the dorsal band from behind the eye to the tip of the tail; sides of body and tail below this streak blotched and variegated



FIGS. 33-38. SALAMANDRELLA KEYSERLINGII. 2 X NAT. SIZE. 33, TOP OF HEAD; 34, SIDE OF HEAD; 35, OPEN MOUTH; 36, SECTION OF TAIL AT END OF BASAL THIRD; 37, HAND; 38, FOOT. No. 22504 U.S.N.M.

with blackish; on the middle line of the back a more or less continuous series of blackish dots bifurcating on the nape; on the head numerous similar dots; underside irregularly mottled with dusky.

Dimensions.

	<i>mm.</i>
Total length.....	124
Tip of snout to gular fold.....	15
Gular fold to anterior angle of vent.....	54
Anterior angle of vent to tip of tail.....	55
Width of head.....	10
Fore leg.....	16
Hind leg.....	18
Height of tail a head-length from vent.....	6
Width of tail a head-length from vent.....	4

Variation.—The differences in dentition and proportions of the tail have been alluded to above (pp. 38–39). The coloration is fairly uniform throughout the series, except that in the lot received from Dr. Vladimir Tushoff the median dorsal area is of a frosted silvery gray rather than tawny.

Habitat.—Extends from the Ural Mountains in the west to the Ussuri country and Kamchatka in the east, north to Verkhoyansk, Siberia, and south into the Chinese province of Sze-chuan.

Dybowski collected the types at the southwest corner of Lake Baikal, while von Schrenck brought home specimens from the Shilka, from Ussuri, and also from Lake Baikal. Boulenger has recorded specimens collected by Mr. Doerries at Khabarovka. Dieckmann, in 1894, also obtained it at the latter place and near Vladivostok, the specimens being in the Hamburg Museum (Nos. 865–866). Sabaneef, in 1872, recorded it from Yekaterinburg, on the east slope of the Ural Mountains, where he found it to be common, and Shitkov, in 1895, studied its reproduction and development there.

The Kamchatkan specimens were at one time considered a distinct species. The types were collected by Wossnessensky at the Javina River, on the west coast, not far from Cape Lopatka, the southern end of the peninsula. I myself obtained specimens from the neighborhood of Petropaulski, on the eastern shore, while von Ditmar records the species from the interior, from the valley of Kamchatka, near Tolbatcha, and also from the Uson volcano, near Lake Kronotskoi.^a

According to Nikolski (1905) this salamander has been collected in Sakhalin by Doctor Saprunenko in 1890, the specimen being in the St. Petersburg Museum (No. 1941).

List of specimens of Salamandrella keyserlingii.

Museum.	No.	Age.	Locality.	When collected.	By whom collected.	Total length.		Gular fold to vent.	Vent to tip of tail.	Width of head.		Height of tail at base.	Height of tail in the middle.
						mm.	mm.			mm.	mm.		
St. Petersburg		Adult ^a	Kamchatka			108	13	41	54	7.5	5.5	5	5
U.S.N.M.	22594	do. b.	Avaicha Bay, Kamchatka.	1895	L. Stejneger	124	15	54	54	10	6	6	6
Do.	23531	do.	Paratunka, Kamchatka.	1896	do.	110	13.6	47	55	9.5	6	7	7
Do.	31712	do.	Kamchatka		Doctor Tushoff.	112	14	50	48	8.5	7	7	7
Do.	31713	do.	do.		do.	117	14.5	49	53.5	9	6	6	6
Do.	31714	do.	do.		do.	120	15	53	52	10	7.5	7.5	7.5
Do.	31715	do.	do.		do.	110	15	45	50	9	6	6.5	6.5
Do.	31716	do.	do.		do.	99	13	41.5	44.5	8	5	6	6

^a Typical of *S. wossnessenskyi*.

^b Description, p. 39, figs. 33–38.

^a Beitr. Kenntn. Russ. Reich., VII, 1890, pp. 400, 751.

List of specimens of *Salamandrella keyserlingii* Continued.

Museum.	No.	Age.	Locality.	When collected.	By whom collected.	Total length.	Snout to gular fold.	Gular fold to vent.	Vent to tip of tail.	Width of head.	Height of tail at base.	Height of tail in the middle.
						mm.	mm.	mm.	mm.	mm.	mm.	mm.
U.S.N.M.	36414	Adult	Paratunka, Kamchatka.	1887	N. A. Grobniński.
Do.	36415	do.	do.	1887	do.
Do.	36416	do.	do.	1887	do.
Do.	36417	do.	do.	1887	do.
Do.	36418	do.	do.	1887	do.
Do.	36419	do.	do.	1887	do.
St. Petersb.	do. ^a	Lake Baikal.	L. v. Schrenck.	107	14	44	19	9	6.5	7

^a Typical of *S. schrenckii*.Genus *ONYCHODACTYLUS*^a Tschudi.1838. *Onychodactylus* TSCHUDI, *Classif. Batr.*; pp. 57, 92 (type, *O. schlegeli*).1839. *Dactylonyx* BIBRON, in Bonaparte, *Icon. Fauna Ital.*, II, fasc. xxvi, fol. 131**2 (variant).1854. *Onychopus* DUMÉRIL and BIBRON, *Erpét. Gén.*, IX, p. 113 (variant).

The most remarkable character of this genus is the presence of distinct, compressed, blackish claws, which are even better developed, hooked, and pointed in the larvæ than in the adult. Troschel, in a very elaborate paper on this animal,^b made the statement that the claws of the adult animal are confined to the males and that the females do not possess them. This is a mistake, as clearly proven by over forty specimens, sixteen of which are females, all possessing as well-developed claws as the males. Apparently Troschel's specimens were not taken during the breeding season, as shown by his description of the vent being a mere slit with swollen lips. It appears then that the males are provided with claws at all seasons of the year, the females only during the breeding time. Only to this extent then is the possession of the claws a sexual character.


ONYCHODACTYLUS JAPONICUS (Houttuyn).

HAKONE SANSHOUWO.

Plate V, figs. 1-4; plate VI.


1782. *Salamandra japonica* HOUTTUYN, *Verh. Genootsch. Wetensch. Vlissingen*, IX, p. 329, pl. facing p. 336, fig. 3 (type-locality, Japan; Thunberg, collector).—*Salamandra japonicus* SCHNEIDER, *Hist. Amph.*, I, 1799, p. 73.—*Lacerta japonica* THUNBERG, *Svensk. Vet. Akad. Nya Handl.*, VIII, 1787, p. 124, pl. iv, fig. 1 (type-locality, Hakone Mts., "Faconiebergerget," Hondo; type in Upsala Acad. Mus.; Thunberg, collector).—

^a From *ὄνυξ*, *ονυχος*, nail and *δάκτυλος*, digit.^b *Arch. Naturg.*, 1877, I.



FIGS. 1-4, ONYCHODACTYLUS JAPONICUS; FIGS. 5, 6, GEOMOLGE FISCHERI; FIGS. 7, 8, SALAMANDRELLA KEYSERLINGII. (FROM SCHLEGEL, BOULENGER, AND STRAUCH.)

FOR EXPLANATION OF PLATE SEE PAGE 553.



ONYCHODACTYLUS JAPONICUS. (FROM SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 553.

Onychodactylus japonicus BONAPARTE, Icon. Fauna Ital., II, fasc. XXVI, 1839, fol. 131**, 2.—GRAY, Cat. Batr. Grad. Brit. Mus., 1850, p. 33, pl. III, fig. 15 (Japan).—STRAUCH, Mém. Acad. Sci. St. Pétersbourg (7), XVI, No. 4, 1870, p. 60 (Japan).—TROSCHEL, Arch. Naturg., 1877, I, p. 199, pl. xv (elaborate descriptions of adult and larva).—HILGENDORF, Sitz. Ber. Naturf. Freunde Berlin, 1880, p. 121 (Hatta village, Hakone Mts.; Oyama, SW. from Yokohama).—BOULENGER, Cat. Batr. Grad. Brit. Mus., 1882, p. 35 (Oyama, near Tokyo; Hakone).—REIN, Japan, Engl. ed., 1884, p. 188 (Hakone Mts.).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V (1891), p. 239.—OKADA, Cat. Vert. Japan, 1891, p. 65 (Iwaki; vicinity of Oyama; Hakone).—SLATER, List. Batr. Indian Mus. (1892), p. 37 (Hakone).—BOETTGER, Kat. Batr. Mus. Senckenberg. (1892), p. 58 (Yokohama).

1798. *Lacerta thunbergii* DONNDORF, Zool. Beytr., III, p. 132 (based on Thunberg, 1787).
1820. *Molge striata* MERREM, Tent. Syst. Amph., p. 185 (Japan; based on Houttuyn and Thunberg).
1838. *Salamandra unguiculata* SCHLEGEL, Fauna Japon., Rept., pp. 123, 139, Saur. et Batr., pl. v, figs. 1-6^a (Hondo and Shikoku).—GEERTS, Nouv. Arch. Mus. Paris, (2) V, 1883, p. 275 (Prov. Iga; Mountains of Hakone and Hata).
1838. *Onychodactylus schlegeli* TSCHUDI, Class. Batr., p. 92 (based on Fauna Japonica).—DUMÉRIL and BIBRON, Erpét. Gén., IX (1854), p. 114; Atlas, pl. XCIII, fig. 1.
1841. *Onycopus sieboldii* DUMÉRIL and BIBRON, Erpét. Gén., VIII, p. 4 (*lapsus*).

Duméril and Bibron^b assert that Schlegel's identification of Houttuyn's *Salamandra japonica* as the present species is erroneous, but to anyone who examines the original description and figures it must be perfectly plain that Schlegel was correct. There can be absolutely no doubt as to the identity of Houttuyn's species, and the mistake of the authors of the Erpétologie Générale can only be explained by their having confounded figures 2 and 3 on Houttuyn's plate. The latter states that his specimen was collected by Thunberg in Japan. Thunberg himself described and figured the same species under the same name, though apparently without knowing Houttuyn's description, and he expressly gives the Hakone Mountains (Fakoniebergerget) as the type-locality, consequently it is in all probability also the locality where he obtained the specimen Houttuyn described.

Description.—*Adult male*; U.S.N.M. No. 34240; Hakone Lake, Sagami, Hondo; July, 1898; A. Owston, collector. Vomero-palatine teeth in a doubly arched series behind the choanae, the two arches extending forward to a line through the anterior edge of the choanae; head flat, rounded anteriorly; nostrils nearer the eye than the tip of the snout, much wider apart than the width of the interorbital space, which is greater than the the upper eyelids; eyes greatly protruding; body elongate, the head from tip of snout to gular fold being con-

^a Reproduced in this work on Plates V-VI.

^b Erpét. Gén., IX, p. 116.

tained three and one-third times in the distance from gular fold to anterior angle of vent; adpressed limbs do not meet by a distance of two costal folds; fore legs slender, the four fingers short, but well developed, each with a raised, compressed, brownish black claw; fourth finger slightly longer than first, third longest; hind legs strong and stout; toes depressed, widened at base, each with a black claw like the fingers; first finger very short, fourth longest, fifth equaling third; a thick dermal expansion from the outer side of fifth finger along the outer side of the foot and tibia, giving the foot a very wide under surface; no tubercles; tail long, longer than head and body, nearly cylindrical in the basal half, then gradually more and more compressed to the end, which is rounded; 13 costal grooves which are rather indistinct; a flat parotoid gland on the side of the neck, bordered above and behind by a groove which begins behind the eye and joins the strong gular fold behind the parotoid; a shallow groove along the median line of upper neck and anterior half of back; vent arrow-shaped, the lips of the oblique anterior slits forming the arrows' head distinctly swollen. Color (in alcohol): top of head, the whole median portion of upper neck and back, as well as the upper edge of the tail, "wood brown," sprinkled more or less with blackish brown, most densely on the head; sides of neck, body, and tail blackish brown, the upper outline against the pale band of the back irregular, but very distinct, the lower outline gradually merging into the pale color of the under surface; legs above dark brown, like the lower portion of the flanks, and, like these, with irregular roundish spots of pale wood-brown.

Dimensions.

	<i>mm.</i>
Total length.....	161
Tip of snout to gular fold.....	16
Gular fold to anterior angle of vent.....	53
Anterior angle of vent to tip of tail.....	92
Width of head.....	10
Fore leg.....	18
Hind leg.....	21
Height of tail a head-length from vent.....	8
Width of tail a head-length from vent.....	7

The *adult breeding female* (U.S.N.M. No. 34245; Hakone Lake; July, 1898; Owston, collector) differs from the male chiefly in the shorter tail and the slenderer hind legs, which, moreover, are devoid of the lateral tarso-tibial enlargement. The vent is also of a somewhat different aspect, inasmuch as the lateral oblique folds, which in the male converge at the anterior end, in the female meet at the middle of the median cleft; the lips of the vent anterior to the lateral folds are much swollen; all the toes provided with black nails.

Dimensions.

	<i>mm.</i>
Total length.....	141
Tip of snout to gular fold.....	15
Gular fold to anterior border of vent.....	54
Anterior border of vent to tip of tail.....	72
Width of head.....	11
Fore leg.....	18
Hind leg.....	23
Height of tail one head-length from vent.....	7.5

The *nonbreeding female* is apparently distinguished by the absence of nails, and the vent is a simple longitudinal slit.

The *larvæ* are characterized by the presence of strong, hooked, black claws.

Variation.—The variation in structural characters is unimportant and presents no difficulties with regard to the identification of this species, being chiefly confined to slight deviations in the position of the vomero-palatine teeth relative to the choanæ, and to the number of costal folds, which varies between 12 and 14. The coloration is also rather uniform, the chief difference consisting in the varying degree of fineness of the dark sprinkling in the dorsal pale area, where these minute dots often congregate near the median line so as to form an irregular dusky stripe.

Habitat.—In the *Fauna Japonica* (p. 125) it is stated that this salamander is found in abundance in the mountainous parts of Hondo and Shikoku, particularly in the provinces of Sagami, Shinano, Tamba, Tazima, and Tosa, and Siebold himself (p. xvi) clearly intimates that he did not obtain it in Kiusiu. Nor am I aware of any record of its having been taken in the latter island. Doctor Fritze states that it is absent in Yezo. Doctor Nozawa, however, in a letter to me enumerates it among the species found there, though as yet he has been unable to secure specimens for me.

To the provinces specifically enumerated above Geerts adds Iga, and Okada adds Iwaki. Most of the specimens in the museums are from Hakone, the type-locality of the species and the locality from which it also takes its Japanese name. Another specific locality from which there are specimens in the British and Berlin museums is Oyama in Sagami province.

The species does not occur outside of Japan proper. It is unknown in the Riu Kiu islands.

List of specimens of Onychodactylus japonicus.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Total length. <i>m m.</i>
34218	Male	Hakone Lake, Hondo	July, 1898	A. Owston	162
34219	do	do	do	do	155
34220	Female	do	do	do	159
34221	Male	do	do	do	168
34222	do	do	do	do	167
34223	Female	do	do	do	144
34224	Male	do	do	do	149
34225	Female	do	do	do	141
34226	do	do	do	do	146
34227	do	do	do	do	128
34228	do	do	do	do	144
34229	Male	do	do	do	144
34230	do	do	do	do	158
34231	do	do	do	do	151
34232	Female	do	do	do	157
34233	do	do	do	do	
34234	do	do	do	do	130
34235	Male	do	do	do	151
34236	do	do	do	do	151
34237	do	do	do	do	155
34238	do	do	do	do	149
34239	Female	do	do	do	140
34240	Male ^a	do	do	do	161
34241	do	do	do	do	148
34242	do	do	do	do	141
34243	do	do	do	do	131
34244	Female	do	do	do	147
34245	do ^b	do	do ^b	do	141
34246	do	do	do	do	150
34247	do	do	do	do	141
34248	Male	do	do	do	153
34249	do	do	do	do	144
34250	Female	do	do	do	145
34251	Male	do	do	do	154
34252	Female	do	do	do	141
34253	Male	do	do	do	155
34254	Female	do	do	do	138
34255	Male	do	do	do	152
34256	do	do	do	do	137
34257	do	do	do	do	155
34258	} Larvæ	do	do	do	
to					
34287					

^a Description p. 43.^b Description p. 44.

Genus *GEOMOLGE*^a Boulenger.

1886. *Geomolge* BOULENGER, Proc. Zool. Soc. London, p. 416 (type, *G. fischeri*).

Geomolge appears to be very closely allied to *Onychodactylus*, the chief differences being that in the former the vomero-palatine series of teeth is interrupted in the middle and that it lacks the black, horny claws.

GEOMOLGE FISCHERI^b Boulenger.

Plate V. fig. 5 6.

1886. *Geomolge fischeri* BOULENGER, Proc. Zool. Soc. London, 1886, p. 416, pl. xxxix, fig. 2 c (type-locality, Khabarovka, Ussuri; types in Brit. Mus. no. 86. 5. 15. 11-12; Doerries, collector); Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 144.—NIKOLSKI, Annuaire Mus. Zool. St. Pétersbourg, I, 1896, p. 77 (Valley of Sutchan River, Ussuri district; St. Petersb. Mus., No. 1901; Busse, collector); Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 440 (Ussuri).

There being no specimen of this species in the U. S. National Museum, an abstract of Boulenger's original description is given:

Physiognomy that of *Onychodactylus japonicus*: limbs in every respect similar to those of *Onychodactylus japonicus* save the absence of claws; the male likewise with tibio-tarsal dilatation; anal opening subcruciform, as in *Onychodactylus*; skin smooth; 14 or 15 costal grooves; parotoids and gular fold as in *Onychodactylus*. Brown above, with blackish variegations, most crowded on the sides, which also bear some whitish spots; lower surface brownish white.

	Male.	Female.
	mm.	mm.
From snout to vent.....	70	80
Head.....	12	14
Tail.....	93d

Habitat.—This species was collected by Mr. Doerries near Khabarovka, in the Ussuri. Nikolski has since recorded a specimen from the valley of Sutchan River, also in the Ussuri district. Beyond this nothing is known of the distribution of this species.

Order SALIENTIA.

- 1768. *Salientia* LAURENTI, Syn. Rept., p. 24.
- 1777. *Ecaudata* SCOPOLI, Introd. Hist. Nat., p. 464.
- 1788. *Batrachi* BATSCH, Anleit. Kenntn. Thiere Mineral., I (p. 437).
- 1802. *Rana* SHAW, Gen. Zool., III, Pt. 1, p. 5.
- 1806. *Ecaudati* DUMÉNIL, Zool. Anal., p. 92.

^a From γῆ, the earth, land, and the generic term *Molge* Merrem, a synonym of *Triturus* Rafinesque, derived from the German word *Molch*, a salamander.

^b Named in honor of Dr. Johann Gustav Fischer, the well-known Hamburg herpetologist. Born on March 1, 1819; died on January 27, 1889.

^c Reproduced in this work on Plate V.

^d Boulenger, Proc. Zool. Soc. London, 1886, p. 416.

List of specimens of Onychodactylus japonicus.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Total length. <i>mm.</i>
34218	Male	Hakone Lake, Hondo	July, 1888	A. Owston	162
34219	do	do	do	do	155
34220	Female	do	do	do	159
34221	Male	do	do	do	168
34222	do	do	do	do	167
34223	Female	do	do	do	144
34224	Male	do	do	do	149
34225	Female	do	do	do	141
34226	do	do	do	do	146
34227	do	do	do	do	128
34228	do	do	do	do	144
34229	Male	do	do	do	144
34230	do	do	do	do	158
34231	do	do	do	do	151
34232	Female	do	do	do	137
34233	do	do	do	do	
34234	do	do	do	do	130
34235	Male	do	do	do	151
34236	do	do	do	do	151
34237	do	do	do	do	155
34238	do	do	do	do	149
34239	Female	do	do	do	140
34240	Male ^a	do	do	do	161
34241	do	do	do	do	148
34242	do	do	do	do	141
34243	do	do	do	do	131
34244	Female	do	do	do	147
34245	do ^b	do	do ^b	do	141
34246	do	do	do	do	150
34247	do	do	do	do	141
34248	Male	do	do	do	153
34249	do	do	do	do	144
34250	Female	do	do	do	145
34251	Male	do	do	do	154
34252	Female	do	do	do	141
34253	Male	do	do	do	155
34254	Female	do	do	do	138
34255	Male	do	do	do	152
34256	do	do	do	do	137
34257	do	do	do	do	155
34258	} Larvæ	do	do	do	
to					
34287					

^a Description p. 43.^b Description p. 44.

Genus *GEOMOLGE*^a Boulenger.

1886. *Geomolge* BOULENGER, Proc. Zool. Soc. London, p. 416 (type, *G. fischeri*).

Geomolge appears to be very closely allied to *Onychodactylus*, the chief differences being that in the former the vomero-palatine series of teeth is interrupted in the middle and that it lacks the black, horny claws.

GEOMOLGE FISCHERI^b Boulenger.

Plate V, fig. 5-6.

1886. *Geomolge fischeri* BOULENGER, Proc. Zool. Soc. London, 1886, p. 416, pl. xxxix, fig. 2 c (type-locality, Khabarovka, Ussuri; types in Brit. Mus. no. 86. 5. 15. 11-12; Doerries, collector); Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 144.—NIKOLSKI, Annuaire Mus. Zool. St. Pétersbourg, I, 1896, p. 77 (Valley of Sutchan River, Ussuri district; St. Petersb. Mus., No. 1904; Busse, collector); Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 440 (Ussuri).

There being no specimen of this species in the U. S. National Museum, an abstract of Boulenger's original description is given:

Physiognomy that of *Onychodactylus japonicus*: limbs in every respect similar to those of *Onychodactylus japonicus* save the absence of claws; the male likewise with tibio-tarsal dilatation; anal opening subcruciform, as in *Onychodactylus*; skin smooth; 14 or 15 costal grooves; parotoids and gular fold as in *Onychodactylus*. Brown above, with blackish variegations, most crowded on the sides, which also bear some whitish spots; lower surface brownish white.

	Male.	Female.
	mm.	mm.
From snout to vent.....	70	80
Head.....	12	14
Tail.....	93	d

Habitat.—This species was collected by Mr. Doerries near Khabarovka, in the Ussuri. Nikolski has since recorded a specimen from the valley of Sutchan River, also in the Ussuri district. Beyond this nothing is known of the distribution of this species.

Order SALIENTIA.

- 1768. *Salientia* LAURENTI, Syn. Rept., p. 24.
- 1777. *Ecaudata* SCOPOLI, Introd. Hist. Nat., p. 464.
- 1788. *Batrachi* BATSCH, Anleit. Kenntn. Thiere Mineral., I (p. 437).
- 1802. *Rana* SHAW, Gen. Zool., III, Pt. 1, p. 5.
- 1806. *Ecaudati* DUMÉRIL, Zool. Anal., p. 92.

^a From γῆ, the earth, land, and the generic term *Molge* Merrem, a synonym of *Triturus* Rafinesque, derived from the German word Molch, a salamander.

^b Named in honor of Dr. Johann Gustav Fischer, the well-known Hamburg herpetologist. Born on March 1, 1819; died on January 27, 1889.

^c Reproduced in this work on Plate V.

^d Boulenger, Proc. Zool. Soc. London, 1886, p. 416.

1813. *Anuri* FISCHER, *Zoognosia*, 3 ed., I, p. 58.
 1825. *Anoura* LATREILLE, *Fam. Nat. Règne Anim.*, p. 104.
 1828. *Pygomolgan* RITGEN, *Nova Acta Acad. Leop. Carol.*, XIV, p. 278.
 1828. *Acerei* WAGLER, *Isis*, 1828, p. 859.
 1832. *Batrachia* MUELLER, *Isis*, 1832 (p. 504).
 1833. *Miura* VAN DER HOEVEN, *Handb. Dierk.*, II, Pt. 2, p. 307
 1835. *Heteromorpha* FITZINGER, *Ann. Wien Mus.*, I, p. 107.
 1839. *Raniformia* HOGG, *Mag. Nat. Hist. (n. s.)*, III, p. 271.
 1855. *Batrachii* VAN DER HOEVEN, *Handb. Dierk.*, 2 ed., II, p. 468.
 1866. *Anura* HAECKEL, *Gen. Morphol.*, II, p. CXXXII
 1878. *Acaudata* KNAUER, *Naturg. Lurche*, p. 100.

The name adopted for this order varies greatly with the various authors. Thus Salientia, Anura, and Ecaudata are used almost indiscriminately. Of these Salientia is the oldest.

It is perfectly true that the various rules for zoological nomenclature do not compel a strict application of the law of priority to groups higher than genera, and that the A. O. U. code specifically proclaims this, but the principle underlying the modern way of fastening a name to a genus is very different from that involved in naming an order or a class. The ends in view are different, and the *modus operandi* must also be different. Nobody nowadays in applying a generic term cares the least whether the name, as originally proposed, covers the same aggregate of species, even approximately, as the genus to which he attaches it, or not. It is enough that a single species, the type species, is included in the original group and in the one adopted. This, however, is a comparatively modern view. In the early days, when generic terms were few, and long before the term "type-species" or "genotype"(!) was invented, it happened quite frequently that authors selected a new generic name for a group which contained the essential elements of one previously named, simply because his grouping was not coextensive with that of his predecessor's. But as new discoveries and closer studies multiplied the genera, and the views as to their compass became more and more divergent, it was found impossible to adhere to such a principle as far as generic terms were concerned.

Not so with the terms for orders and classes. It has always been the recognized custom to employ a name embracing, at least approximately, an assembly of genera, or orders, coextensive with it when first proposed. The inconvenience caused by the coining of new names of groups higher than genera has been small compared to the convenience of knowing approximately the extent of a group by the name applied to it. A good reason why it is inconvenient and inexpedient to apply strictly the law of priority to such higher groups is the fact that we know no such thing as a "type genus" or a "type order" by the aid of which the name of the order or the class can be arbitrarily fixed, even if it were desirable to do so.

As a general proposition it may be stated that there is but little divergence among authors with regard to the names of classes and orders. Common usage has in most cases been too strong for the extremists, who have either tried to apply names according to priority or who have inflicted new ones every time the "rank" of the group was changed or a trifling component part shifted from one group to another, or whose fine linguistic sensitiveness compelled them to improve the name and render its form "classic." There are names which it has been impossible to dislodge, which are employed almost unanimously. These it would be folly to attempt to disturb.

The principle which has guided me in selecting the names of orders and classes may be stated briefly as follows: In cases where a term has not received a practically universal acceptance the oldest known group name (above a generic name and later than 1758) is applied, which covers approximately the same aggregate of genera, irrespective of the comparative rank assigned to the group by the original namer, provided its form is not the plural of a current genus in the same class, nor that now universally adopted for superfamilies, families, and subfamilies.

The application of this principle may be shown by a few examples. Thus Salientia is adopted as being the oldest known group name for the present order, because when originally proposed it covered the same aggregate of genera as now, and none of the later names of the same compass have received universal acceptance. On the other hand, Gradientia has been rejected, although it is older than Caudata, and the latter adopted, because the term Gradientia, as originally proposed, embraced the saurian reptiles as well. Oppel's Squamata has been accepted for the reptilian order containing the lizards and snakes, although it included the crocodiles, a discrepancy too slight to affect the general applicability of the term. Again, Testudinata, although proposed four years later than Link's Cataphractæ, has been accepted in preference to the latter (after the elimination of the older terms Testudines and Chelonia as conflicting with the earlier or synchronous generic names *Testudo* and *Chelonia*) because of the comparative currency of that term, as against the almost total obscurity of the one coined by Link. Finally, the suborder containing the soft-shelled turtles is called Chilotæ in spite of the fact that the term Trionychoidea is both earlier and commonly adopted, the reason for the rejection of the latter name being that its form is identical with that of the superfamily.

The Salientia may be divided into three suborders, viz, Aglossa, Linguata, and Costata, of which the first one is distinguished by having the eustachian tubes united into a single pharyngeal opening and no tongue, while the last one is amply characterized by the presence

of ribs. The Aglossa are only found in Africa and South America; the other two suborders are represented within our area.

Suborder COSTATA.

1879. *Costati* LATASTE, Act. Soc. Linn. Bordeaux, XXXIII, 1879, p. 339.

Salientia having a tongue; eustachian tubes, when present, with two pharyngeal openings; ribs present and a pair of transverse processes near the proximal end of the coccyx. Larvæ with one spiraculum in the middle of the thoracic region.

The Costata, which embrace only one family, the Discoglossidæ, have a very wide and disconnected distribution showing their great antiquity and fully bearing out the verdict of Professor Howes, based upon anatomical investigations, that they are "by far the least modified of living anura."^a

The principal center of distribution at present is central and southern Europe, where three genera together with five species are at home, one of these species extending its range into northwestern Africa and another into central Russia. At the other extreme of the Palearctic region another species of one of the European genera maintains its isolated existence, namely, the species to be treated of in this work, while in the Yunnan region another one has recently been discovered. Two more genera belong to this family, both widely removed from the above, namely, *Liopelma*, the single species of which is the only batrachian occurring in New Zealand to which antipodal island group it is peculiar and where it seems to be very rare, and *Ascaphus*, represented thus far by two specimens only, of a single species, *A. truei*, which has recently been discovered in the State of Washington, not far from the Pacific coast. The extreme rarity of the last two species would seem to indicate that they are nearing extinction.^b

Family DISCOGLOSSIDÆ.

Only this family is included in the suborder.

Genus BOMBINA^c Oken.

1816. *Bombina* OKEN, Lehrb. Zool., II, p. 207 (type, *Rana bombina*).



1820. *Bombinator* MERREM, Tent. Syst. Amph., p. 178 (same type).

1830. *Bombitator* WAGLER, Nat. Syst. Amph., p. 206 (emendation).

^a Proc. Zool. Soc. London, 1888, pp. 178, 506.

^b See Stejneger, A Résumé of the Geographical Distribution of the Discoglossoid Toads in the Light of Ancient Land Connections, in Bull. Amer. Geogr. Soc., XXXVII, No. 2, Feb. 1905, pp. 91-93; and The Geographical Distribution of the Bell-Toads, in Science (n. s.), XXII, Oct. 20, 1905, p. 502.

^c From the medieval Latin verb *bombinare*, to utter a *bombus*, a loud hollow sound, such as the stroke of a bell.



BOMBINA ORIENTALIS. (FROM BOULENGER.)

FOR EXPLANATION OF PLATE SEE PAGE 554.

The genus *Bombina* has a very curious distribution. Two species, *Bombina bombina* (Linnaeus) and *Bombina salsa*^a (Schrank), occur in Europe, the former in northern and eastern central Europe. The latter, which seems to embrace several local forms, such as *B. salsa pachypus* (= *Bombinator apenninicus* Gistel and Bronne, 1850), in Italy, *B. salsa nigriventris* (Dueringen) in Bosnia and Montenegro, *B. salsa kolombatovici* (Bedriaga) in Dalmatia, etc., occupies France, parts of southern central Europe, Italy, and the Balkan peninsula. None of these species extends eastward beyond the Volga. A third species of the genus occurs in eastern Asia, though probably not to the west of a line through Lake Baikal. A fourth species, *Bombina maxima*, has recently been described from Yunnan at an altitude of 6,000 feet.

BOMBINA ORIENTALIS^b (Boulenger).

Plate VII.

1878. *Bombinator igneus* MARTENS, Preuss. Exped. Ost-Asien, Zool., I, p. 384 (not of Laurenti) (Chefu, N. China).—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 447 (part: Chefu, China).—BOETTGER, Offenbach. Ver. Naturk. 24-25 Ber., 1885, p. 164 (Chefu).
1890. *Bombinator orientalis* BOULENGER, Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 143, pl. IX, fig. 2^c (type-localities, N. China, Korea, Khabarovka; types in Brit. Mus.); Proc. Zool. Soc. London, 1890, p. 326 (Chefu; Korea; Ussuri).—WERNER, Abh. Bayer. Akad. Wiss., (Muenchen) II Klasse, XXII, Pt. 2, 1904, p. 359 (Tsingtau, China).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, (8) XVII, No. 1 (1905) p. 415 (Vladivostok; Korea, etc.).


Description (figs. 39-43).—*Adult male*; U.S.N.M. No. 17522; Fusan, Korea; 1885; P. L. Jouy, collector. Vomerine teeth in two

^aIn his Tailless Batrachians of Europe, p. 322, footnote, Doctor Boulenger, in adopting the sixteen years younger name *Rana agilis* in preference to the older *R. dalmatina*, as required by "the strict application of the law of priority," pleads that "this is one of those cases in which, it appears to me, conservatism is desirable," and he continues: "Similar considerations have guided me in the naming of the two species of the genus *Bombinator*, and I hope, in the interest of stability of nomenclature they will commend themselves to future workers." It is indeed not uncommon to hear the breaking of this law termed "conservatism," and certainly the conservatives ought to know if that is a correct definition, but we radicals prefer to submit to the law as the only means of obtaining the desired "stability of nomenclature." If Boulenger, in the two cases cited, had followed the law we would by this time have reached stability in their cases, while now it has been postponed probably for a generation. Who is to decide in which cases it is allowable to break the law and substitute "conservatism?" A simple "it appears to me" applied here and there by a dozen herpetologists would cause a fine "stability!"

^bEastern.

^cReproduced in this work on Plate VII.

small transverse oval groups behind the level of the choanæ; tongue oval, truncate behind, large, adherent all around; snout flat, rounded, without canthus rostralis; nostrils on top of snout about halfway between eye and tip of snout, distance from each other equaling the interorbital space, which is narrower than the upper eyelids; no tympanum; parotoid gland long, narrow, indistinct; no gular sac; a very faintly indicated gular fold; fore leg well developed, longer than half the length of head and body; fingers short, depressed, first shortest, third longest, second and fourth of about equal length; no subarticular tubercles, but a large and distinct inner palmar pad and a smaller indistinct outer one; a large patch of horny nuptial excrescences on the inner side of the fore arm, and smaller patches on the inner side of the large palmar pad, and of first, second, and third toes; when hind leg is carried forward along the side, the heel reaches



FIGS. 39-43. --*BOMBINA ORIENTALIS*. $1\frac{1}{2} \times$ NAT. SIZE. 39, TOP OF HEAD; 40, SIDE OF HEAD; 41, OPEN MOUTH; 42, UNDERSIDE OF RIGHT HAND SHOWING NUPTIAL EXCRESCENCES; 43, FOOT. No. 17522. U.S.N.M.

the corner of the mouth and tarso-metatarsal joint slightly beyond the tip of the snout; tibia equals distance from base of inner metatarsal tubercle to tip of longest toe; heels just meet when the femurs are placed at right angles to the axis of the body; toes short, webbed nearly to their tips, webs deeply excised, third toe longer than fifth; no subarticular tubercles; inner metatarsal tubercle small and weakly developed, no outer tubercle; no nuptial excrescences on toes; upper surface covered with closely set and irregularly arranged prominent tubercles of varying size, all sprinkled over with minute, dark brownish points and each surmounted by one or more horny spines, the base of which is pale and the tip blackish brown; underside smooth, with the posterior part of the belly and proximal parts of femurs somewhat rugose; outer side of sole of feet and underside of base of fifth toe with spines like the upper side. Color (in alcohol) above dark drab-

gray, with irregular obscure dusky spots and dusky cross-markings on the limbs; a pale sage-green spot of irregular outlines across the back between the fore legs and two smaller ones, still more indefinite on each side of the back about the middle; a pale longitudinal stripe on the tip of the snout from the level of the nostrils to the lip; lips dusky; underside pale buff (red in life), with numerous blackish, sharply defined, irregular spots; palms buff (red), separated from the similarly colored under surface of the fore arm by a black wristband; a large buff (red) spot on the inner side of the plantar surface, involving the entire first toe, below and above, and continuous with the similarly colored patch on the inner side of the tarsus; tips of all the digits abruptly and bright buff (red) above and below.

Dimensions.

	<i>mm.</i>
Tip of snout to vent.....	44
Tip of snout to gular fold.....	16
Width of head.....	16
Width of interorbital space.....	3
Fore leg.....	25
Hind leg, from vent.....	62
Foot from base of inner metatarsal tubercle to tip of longest toe.....	18

Sexual variation.—The males of this species during the breeding season possess groups of blackish horny excrescences on the inner side of fore arm, palmar tubercle and first, second, and third fingers exactly as in *Bombina salsa*; but, unlike the latter, there are none on the posterior extremities. In the female the fingers are longer and slenderer, especially the outer one. The dorsal spines and asperities are smaller.

Color variation.—In the series before me there is very little deviation from the coloration of the underside as described above. In two specimens, Nos. 17527 and 17530, the blackish markings are larger, especially in the latter, in which the underside may be described as black with buff (red) spots; in both, the plantar light patch is separated from the tarsal patch by a narrow black bridge, in No. 17530 only on one foot, however. The upper surface is considerably more olive in some specimens and the dusky spots and markings above much better defined and darker.

Habitat.—The eastern bell-toad is confined to northeastern China, Korea, and the Ussuri country. Specimens from all of these localities are in the British Museum. The U. S. National Museum possesses nine specimens collected in 1885 by P. L. Jouy near Fusan, Korea. Pleske's specimens from Vladivostok are in the Academy Museum in St. Petersburg, and the Hamburg Museum also has it from the same locality (No. 862), collected by Dieckmann in 1894. In the St. Petersburg Museum there are also specimens collected by Poljakow at the Novgorodski Port, Possiet Bay.

List of specimens of Bombina orientalis.

U. S. N. M. No.	Sex.	Locality.	When col- lected.	By whom col- lected.
17522	Male a	Fusan, Korea	1885	P. L. Jouy.
17523	do.	do.	1885	Do.
17524	do.	do.	1885	Do.
17525	do.	do.	1885	Do.
17526	do.	do.	1885	Do.
17527	do b	do.	1885	Do.
17528	do.	do.	1885	Do.
17529	Female	do.	1885	Do.
17530	Male b	do.	1885	Do.

a Description, p. 51; figs. 39-43.

b P. 53.

Suborder LINGUATA.

1845. *Linguata* GRAVENHORST, Thierreich, p. 43.1847. *Phaneroglossa* GERVAIS, Dict. d'Hist. Nat., IX, p. 721.

Salientia having a tongue; eustachian tubes with two pharyngeal openings; lacking ribs and transverse process to urostyle. Larvæ with one spiraculum on left side only.

The term *Ranæ phaneroglossæ* used in 1830 by Wagler (and usually quoted simply as "*Phaneroglossa*") included the majority of both Salientia and Caudata. Duméril and Bibron first restricted the "*Phanéroglosses*" (first latinized by Gervais in 1847, so far as I have been able to ascertain) to the Salientia with a tongue, therein including also the Discoglossidæ. Those who still regard these as belonging to the same suborder as the other tongued Salientia will probably retain the name *Phaneroglossa*. Gravenhorst's *Linguata* may then be restricted to the latter without the genera having ribs. Originally, it is true, it was meant to be equivalent to the *Phanéroglosses*, but it has never been accepted for them, and the Discoglossidæ form so small a portion that it is better to retain it as here defined than to create a new name.

The families to which the toads and frogs inhabiting our region belong may be distinguished as follows:

KEY TO FAMILIES.

- a¹ Clavicle and coracoid of each side connected by a longitudinal arched cartilage, which overlaps that of the opposite side.
- b¹ Both jaws toothless.....BUFONIDÆ
- b² Upper jaw toothed.....HYLIDÆ
- a² Clavicles and coracoids of both sides firmly united by a single median cartilage.
- b¹ Both jaws toothless; sacral diapophyses dilated.....ENGYSTOMIDÆ
- b² Upper jaw toothed; sacral diapophyses not or but slightly dilated.....RANIDÆ

The characters of the two groups indicated by a¹ and a² correspond to those of Cope's *Arcifera* and *Firmisternia*, respectively, divisions which Boulenger calls "series."

As the employment of the systematic arrangement given above for purposes of identification would necessitate dissection of the specimens, I add the following artificial Key, by which the Salientia occurring within our limits may be referred to their respective families by external characters alone:

KEY TO THE JAPANESE FAMILIES OF TAILLESS BATRACHIANS.

- a¹ Digits without subarticular tubercles; pupil triangular (Suborder COSTATA).....
DISCOGLOSSIDÆ, p. 50
- a² Digits with subarticular tubercles; pupil round or horizontal (Suborder LINGUATA).
b¹ A parotoid gland.....BUFONIDÆ, p. 55
- b² No parotoid gland.
c¹ Tongue without posterior prolongations or "horns."
d¹ Upper jaw toothed.....HYLIDÆ, p. 75
- d² Upper jaw toothless.....ENGYSTOMIDÆ, p. 87
- c² Tongue with two posterior horn-shaped prolongations.....RANIDÆ, p. 92

Family BUFONIDÆ.

A large family having no maxillary teeth, a large parotoid gland behind the head, shoulder-girdle arciferous, and lateral sacral processes dilated. Distributed over the entire globe, except the polar regions, New Zealand, and Polynesia. Some forms are aquatic, some arboreal, some burrowing in their habits, but most are terrestrial. They feed largely on insects, and are among the best friends of the farmer and gardener, by whom they ought to be protected instead of persecuted and exterminated, as they usually are, on account of their often unattractive appearance.

Only one genus is represented within the territory covered by this work.

Genus BUFO^a Laurenti.

1768. *Bufo* LAURENTI, Syn. Rept., p. 25 (type, *B. vulgaris*).
1788. *Bufo* LACÉPÈDE, Hist. Nat. Quadr. Ovip., I, Syn. Meth. (emendation).
1814. *Batrachus* RAFINESQUE, Specchio Sci., (Palermo), II, fasc. 7 (substitute for *Bufo*).
1815. *Bufotes* RAFINESQUE, Analyse Nat., p. 78 (substitute for *Bufo*).
1816. *Calamita* OKEN, Lehrb. Zool., II, p. 209 (type, *B. calamita*; not of Schneider 1799).
1828. *Chusca* RITGEN, Nova Acta Acad. Carol. Leop., XIV, p. 278 (type, *Bombinator strumosus*).
1843. *Phryne* FITZINGER, Syst. Rept., p. 32 (type, *B. vulgaris*).
1843. *Phrynoïdis* FITZINGER, Syst. Rept., p. 32 (type, *B. asper*).
1864. *Epidalea* COPE, Proc. Phila. Acad., 1864, p. 181 (type, *B. calamita*).

The above synonyms are based upon species occurring within our limits or are additional to those found in previously published synonymies. The other synonyms of *Bufo* are too numerous to be reproduced here.

The status of the Japanese toads is still very unsatisfactory.

^aThe Latin word for toad.

List of specimens of *Salamandrella keyserlingii*—Continued.

Museum.	No.	Age.	Locality.	When collected.	By whom collected.	Total length.	Snout to gular fold.	Gular fold to vent.	Vent to tip of tail.	Width of head.	Height of tail at base.	Height of tail in the middle.
						mm.	mm.	mm.	mm.	mm.	mm.	mm.
U.S.N.M.	36414	Adult	Paratunka, Kamchatka.	1887	N. A. Grebnitski.							
Do.	36415	do.	do.	1887	do.							
Do.	36416	do.	do.	1887	do.							
Do.	36417	do.	do.	1887	do.							
Do.	36418	do.	do.	1887	do.							
Do.	36419	do.	do.	1887	do.							
St. Petersb.		do. ^a	Lake Baikal.		L. v. Schrenck.	107	14	44	49	9	6.5	7

^a Typical of *S. schrenckii*.Genus *ONYCHODACTYLUS*^a Tschudi.1838. *Onychodactylus* TSCHUDI, *Classif. Batr.*, pp. 57, 92 (type, *O. schlegelii*).1839. *Dactylonyx* BIBRON, in Bonaparte, *Icon. Fauna Ital.*, II, fasc. xxvi, fol. 131**2 (variant).1854. *Onychopus* DUMÉRIL and BIBRON, *Erpét. Gén.*, IX, p. 113 (variant).

The most remarkable character of this genus is the presence of distinct, compressed, blackish claws, which are even better developed, hooked, and pointed in the larvæ than in the adult. Troschel, in a very elaborate paper on this animal,^b made the statement that the claws of the adult animal are confined to the males and that the females do not possess them. This is a mistake, as clearly proven by over forty specimens, sixteen of which are females, all possessing as well-developed claws as the males. Apparently Troschel's specimens were not taken during the breeding season, as shown by his description of the vent being a mere slit with swollen lips. It appears then that the males are provided with claws at all seasons of the year, the females only during the breeding time. Only to this extent then is the possession of the claws a sexual character.


ONYCHODACTYLUS JAPONICUS (Houttuyn).

HAKONE SANSHOUWO.

Plate V, figs. 1-4; plate VI.

1782. *Salamandra japonica* HOUTTUYN, *Verh. Genootsch. Wetensch. Vlissingen*, IX, p. 329, pl. facing p. 336, fig. 3 (type-locality, Japan; Thunberg, collector).—*Salamandra japonicus* SCHNEIDER, *Hist. Amph.*, I, 1799, p. 73.—*Lacerta japonica* THUNBERG, *Svensk. Vet. Akad. Nya Handl.*, VIII, 1787, p. 124, pl. iv, fig. 1 (type-locality, Hakone Mts., "Faconiebergct," Hondo; type in Upsala Acad. Mus.; Thunberg, collector).—

^a From *Ορυζ*, *ορυχος*, nail and *δάκτυλος*, digit.^b *Arch. Naturg.*, 1877, I.




FIGS. 1-4, ONYCHODACTYLUS JAPONICUS; FIGS. 5, 6, GEOMOLGE FISCHERI; FIGS. 7, 8, SALAMANDRELLA KEYSERLINGII. (FROM SCHLEGEL, BOULENGER, AND STRAUCH.)
FOR EXPLANATION OF PLATE SEE PAGE 553.



1



2



3

ONYCHODACTYLUS JAPONICUS. (FROM SCHLEGEL.)
FOR EXPLANATION OF PLATE SEE PAGE 553.

- Onychodactylus japonicus* BONAPARTE, Icon. Fauna Ital., II, fasc. XXVI, 1839, fol. 131** 2.—GRAY, Cat. Batr. Grad. Brit. Mus., 1850, p. 33, pl. III, fig. 15 (Japan).—STRAUCH, Mém. Acad. Sci. St. Pétersbourg (7), XVI, No. 4, 1870, p. 60 (Japan).—TROSCHEL, Arch. Naturg., 1877, I, p. 199, pl. xv (elaborate descriptions of adult and larva).—HILGENDORF, Sitz. Ber. Naturf. Freunde Berlin, 1880, p. 121 (Hatta village, Hakone Mts.; Oyama, SW. from Yokohama).—BOULENGER, Cat. Batr. Grad. Brit. Mus., 1882, p. 35 (Oyama, near Tokyo; Hakone).—REIN, Japan, Engl. ed., 1884, p. 188 (Hakone Mts.).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V (1891), p. 239.—OKADA, Cat. Vert. Japan, 1891, p. 65 (Iwaki; vicinity of Oyama; Hakone).—SLATER, List. Batr. Indian Mus. (1892), p. 37 (Hakone).—BOETTGER, Kat. Batr. Mus. Senckenberg. (1892), p. 58 (Yokohama).
1798. *Lacerta thunbergii* DONNDORF, Zool. Beytr., III, p. 132 (based on Thunberg, 1787).
1820. *Molge striata* MERREM, Tent. Syst. Amph., p. 185 (Japan; based on Houttuyn and Thunberg).
1838. *Salamandra unguiculata* SCHLEGEL, Fauna Japon., Rept., pp. 123, 139, Saur. et Batr., pl. v, figs. 1-6^a (Hondo and Shikoku).—GEERTS, Nouv. Arch. Mus. Paris, (2) V, 1883, p. 275 (Prov. Iga; Mountains of Hakone and Hata).
1838. *Onychodactylus schlegelii* TSCHUDI, Class. Batr., p. 92 (based on Fauna Japonica).—DUMÉRIL and BIBRON, Erpét. Gén., IX (1834), p. 114; Atlas, pl. XCIII, fig. 1.
1841. *Onycopus sieboldii* DUMÉRIL and BIBRON, Erpét. Gén., VIII, p. 4 (*lapsus*).

Duméril and Bibron^b assert that Schlegel's identification of Houttuyn's *Salamandra japonica* as the present species is erroneous, but to anyone who examines the original description and figures it must be perfectly plain that Schlegel was correct. There can be absolutely no doubt as to the identity of Houttuyn's species, and the mistake of the authors of the Erpétologie Générale can only be explained by their having confounded figures 2 and 3 on Houttuyn's plate. The latter states that his specimen was collected by Thunberg in Japan. Thunberg himself described and figured the same species under the same name, though apparently without knowing Houttuyn's description, and he expressly gives the Hakone Mountains (Fakonieberget) as the type-locality, consequently it is in all probability also the locality where he obtained the specimen Houttuyn described.

Description.—*Adult male*; U.S.N.M. No. 34240; Hakone Lake, Sagami, Hondo; July, 1898; A. Owston, collector. Vomeropalatine teeth in a doubly arched series behind the choanae, the two arches extending forward to a line through the anterior edge of the choanae; head flat, rounded anteriorly; nostrils nearer the eye than the tip of the snout, much wider apart than the width of the interorbital space, which is greater than the the upper eyelids; eyes greatly protruding; body elongate, the head from tip of snout to gular fold being con-

^a Reproduced in this work on Plates V-VI.

^b Erpét. Gén., IX, p. 116.

tained three and one-third times in the distance from gular fold to anterior angle of vent; adpressed limbs do not meet by a distance of two costal folds; fore legs slender, the four fingers short, but well developed, each with a raised, compressed, brownish black claw; fourth finger slightly longer than first, third longest; hind legs strong and stout; toes depressed, widened at base, each with a black claw like the fingers; first finger very short, fourth longest, fifth equaling third; a thick dermal expansion from the outer side of fifth finger along the outer side of the foot and tibia, giving the foot a very wide under surface; no tubercles; tail long, longer than head and body, nearly cylindrical in the basal half, then gradually more and more compressed to the end, which is rounded; 13 costal grooves which are rather indistinct; a flat parotoid gland on the side of the neck, bordered above and behind by a groove which begins behind the eye and joins the strong gular fold behind the parotoid; a shallow groove along the median line of upper neck and anterior half of back; vent arrow-shaped, the lips of the oblique anterior slits forming the arrows' head distinctly swollen. Color (in alcohol): top of head, the whole median portion of upper neck and back, as well as the upper edge of the tail, "wood brown," sprinkled more or less with blackish brown, most densely on the head; sides of neck, body, and tail blackish brown, the upper outline against the pale band of the back irregular, but very distinct, the lower outline gradually merging into the pale color of the under surface; legs above dark brown, like the lower portion of the flanks, and, like these, with irregular roundish spots of pale wood-brown.

Dimensions.

	<i>mm.</i>
Total length.....	161
Tip of snout to gular fold.....	16
Gular fold to anterior angle of vent.....	53
Anterior angle of vent to tip of tail.....	92
Width of head.....	10
Fore leg.....	18
Hind leg.....	21
Height of tail a head-length from vent.....	8
Width of tail a head-length from vent.....	7

The *adult breeding female* (U.S.N.M. No. 34245; Hakone Lake; July, 1898; Owston, collector) differs from the male chiefly in the shorter tail and the slenderer hind legs, which, moreover, are devoid of the lateral tarso-tibial enlargement. The vent is also of a somewhat different aspect, inasmuch as the lateral oblique folds, which in the male converge at the anterior end, in the female meet at the middle of the median cleft; the lips of the vent anterior to the lateral folds are much swollen; all the toes provided with black nails.

Dimensions.

	<i>mm.</i>
Total length.....	141
Tip of snout to gular fold.....	15
Gular fold to anterior border of vent.....	54
Anterior border of vent to tip of tail.....	72
Width of head.....	11
Fore leg.....	18
Hind leg.....	23
Height of tail one head-length from vent.....	7.5

The *nonbreeding female* is apparently distinguished by the absence of nails, and the vent is a simple longitudinal slit.

The *larvæ* are characterized by the presence of strong, hooked, black claws.

Variation.—The variation in structural characters is unimportant and presents no difficulties with regard to the identification of this species, being chiefly confined to slight deviations in the position of the vomero-palatine teeth relative to the choanæ, and to the number of costal folds, which varies between 12 and 14. The coloration is also rather uniform, the chief difference consisting in the varying degree of fineness of the dark sprinkling in the dorsal pale area, where these minute dots often congregate near the median line so as to form an irregular dusky stripe.

Habitat.—In the Fauna Japonica (p. 125) it is stated that this salamander is found in abundance in the mountainous parts of Hondo and Shikoku, particularly in the provinces of Sagami, Shinano, Tamba, Tazima, and Tosa, and Siebold himself (p. XVI) clearly intimates that he did not obtain it in Kiusiu. Nor am I aware of any record of its having been taken in the latter island. Doctor Fritze states that it is absent in Yezo. Doctor Nozawa, however, in a letter to me enumerates it among the species found there, though as yet he has been unable to secure specimens for me.

To the provinces specifically enumerated above Geerts adds Iga, and Okada adds Iwaki. Most of the specimens in the museums are from Hakone, the type-locality of the species and the locality from which it also takes its Japanese name. Another specific locality from which there are specimens in the British and Berlin museums is Oyama in Sagami province.

The species does not occur outside of Japan proper. It is unknown in the Riu Kiu islands.

List of specimens of Onychodactylus japonicus.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Total length.
					<i>mm.</i>
34218	Male	Hakone Lake, Hondo	July, 1888	A. Owston	162
34219	do	do	do	do	155
34220	Female	do	do	do	159
34221	Male	do	do	do	168
34222	do	do	do	do	167
34223	Female	do	do	do	144
34224	Male	do	do	do	149
34225	Female	do	do	do	141
34226	do	do	do	do	146
34227	do	do	do	do	128
34228	do	do	do	do	144
34229	Male	do	do	do	144
34230	do	do	do	do	158
34231	do	do	do	do	151
34232	Female	do	do	do	137
34233	do	do	do	do	
34234	do	do	do	do	130
34235	Male	do	do	do	151
34236	do	do	do	do	151
34237	do	do	do	do	155
34238	do	do	do	do	149
34239	Female	do	do	do	140
34240	Male ^a	do	do	do	161
34241	do	do	do	do	148
34242	do	do	do	do	141
34243	do	do	do	do	131
34244	Female	do	do	do	147
34245	do ^b	do	do ^b	do	141
34246	do	do	do	do	150
34247	do	do	do	do	141
34248	Male	do	do	do	153
34249	do	do	do	do	144
34250	Female	do	do	do	145
34251	Male	do	do	do	154
34252	Female	do	do	do	141
34253	Male	do	do	do	155
34254	Female	do	do	do	138
34255	Male	do	do	do	152
34256	do	do	do	do	137
34257	do	do	do	do	155
34258	} Larvæ	do	do	do	
to					
34287					

^a Description p. 43.^b Description p. 44.

Genus *GEOMOLGE*^a Boulenger.

1886. *Geomolge* BOULENGER, Proc. Zool. Soc. London, p. 416 (type, *G. fischeri*).

Geomolge appears to be very closely allied to *Onychodactylus*, the chief differences being that in the former the vomero-palatine series of teeth is interrupted in the middle and that it lacks the black, horny claws.

GEOMOLGE FISCHERI^b Boulenger.

Plate V, fig. 5-6.

1886. *Geomolge fischeri* BOULENGER, Proc. Zool. Soc. London, 1886, p. 416, pl. XXXIX, fig. 2^c (type-locality, Khabarovka, Ussuri; types in Brit. Mus. no. 86. 5. 15. 11-12; Doerries, collector); Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 144.—NIKOLSKI, Annuaire Mus. Zool. St. Pétersbourg, I, 1896, p. 77 (Valley of Sutchan River, Ussuri district; St. Petersb. Mus., No. 1904; Busse, collector); Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 440 (Ussuri).

There being no specimen of this species in the U. S. National Museum, an abstract of Boulenger's original description is given:

Physiognomy that of *Onychodactylus japonicus*: limbs in every respect similar to those of *Onychodactylus japonicus* save the absence of claws; the male likewise with tibio-tarsal dilatation; anal opening suberuciform, as in *Onychodactylus*; skin smooth; 14 or 15 costal grooves; parotoids and gular fold as in *Onychodactylus*. Brown above, with blackish variegations, most crowded on the sides, which also bear some whitish spots; lower surface brownish white.

	Male.	Female.
	mm.	mm.
From snout to vent.....	70	80
Head.....	12	14
Tail.....	93 ^d

Habitat.—This species was collected by Mr. Doerries near Khabarovka, in the Ussuri. Nikolski has since recorded a specimen from the valley of Sutchan River, also in the Ussuri district. Beyond this nothing is known of the distribution of this species.

Order SALIENTIA.

- 1768. *Salientia* LAURENTI, Syn. Rept., p. 24.
- 1777. *Ecaudata* SCOPOLI, Introd. Hist. Nat., p. 464.
- 1788. *Batrachi* BATSCHE, Anleit. Kenntn. Thiere Mineral., I (p. 437).
- 1802. *Rana* SHAW, Gen. Zool., III, Pt. 1, p. 5.
- 1806. *Ecaudati* DUMÉRIL, Zool. Anal., p. 92.

^a From γῆ, the earth, land, and the generic term *Molge* Merrem, a synonym of *Triturus* Rafinesque, derived from the German word Molch, a salamander.

^b Named in honor of Dr. Johann Gustav Fischer, the well-known Hamburg herpetologist. Born on March 1, 1819; died on January 27, 1889.

^c Reproduced in this work on Plate V.

^d Boulenger, Proc. Zool. Soc. London, 1886, p. 416.

1813. *Anuri* FISCHER, Zoognosia, 3 ed., I, p. 58.
 1825. *Anoura* LATREILLE, Fam. Nat. Règne Anim., p. 104.
 1828. *Pygomolgar* RITGEN, Nova Acta Acad. Leop. Carol., XIV, p. 278.
 1828. *Accrei* WAGLER, Isis, 1828, p. 859.
 1832. *Batrachia* MUELLER, Isis, 1832 (p. 504).
 1833. *Miura* VAN DER HOEVEN, Handb. Dierk., II, Pt. 2, p. 307.
 1835. *Heteromorpha* FITZINGER, Ann. Wien Mus., I, p. 107.
 1839. *Raniformia* HOGG, Mag. Nat. Hist. (n. s.), III, p. 271.
 1855. *Batrachii* VAN DER HOEVEN, Handb. Dierk., 2 ed., II, p. 468.
 1866. *Anura* ILAECKBL, Gen. Morphol., II, p. CXXXII
 1878. *Acaudata* KNAUER, Naturg. Lurche, p. 100.

The name adopted for this order varies greatly with the various authors. Thus Salientia, Anura, and Ecaudata are used almost indiscriminately. Of these Salientia is the oldest.

It is perfectly true that the various rules for zoological nomenclature do not compel a strict application of the law of priority to groups higher than genera, and that the A. O. U. code specifically proclaims this, but the principle underlying the modern way of fastening a name to a genus is very different from that involved in naming an order or a class. The ends in view are different, and the modus operandi must also be different. Nobody nowadays in applying a generic term cares the least whether the name, as originally proposed, covers the same aggregate of species, even approximately, as the genus to which he attaches it, or not. It is enough that a single species, the type species, is included in the original group and in the one adopted. This, however, is a comparatively modern view. In the early days, when generic terms were few, and long before the term "type-species" or "genotype"(!) was invented, it happened quite frequently that authors selected a new generic name for a group which contained the essential elements of one previously named, simply because his grouping was not coextensive with that of his predecessor's. But as new discoveries and closer studies multiplied the genera, and the views as to their compass became more and more divergent, it was found impossible to adhere to such a principle as far as generic terms were concerned.

Not so with the terms for orders and classes. It has always been the recognized custom to employ a name embracing, at least approximately, an assembly of genera, or orders, coextensive with it when first proposed. The inconvenience caused by the coining of new names of groups higher than genera has been small compared to the convenience of knowing approximately the extent of a group by the name applied to it. A good reason why it is inconvenient and inexpedient to apply strictly the law of priority to such higher groups is the fact that we know no such thing as a "type genus" or a "type order" by the aid of which the name of the order or the class can be arbitrarily fixed, even if it were desirable to do so.

As a general proposition it may be stated that there is but little divergence among authors with regard to the names of classes and orders. Common usage has in most cases been too strong for the extremists, who have either tried to apply names according to priority or who have inflicted new ones every time the "rank" of the group was changed or a trifling component part shifted from one group to another, or whose fine linguistic sensitiveness compelled them to improve the name and render its form "classic." There are names which it has been impossible to dislodge, which are employed almost unanimously. These it would be folly to attempt to disturb.

The principle which has guided me in selecting the names of orders and classes may be stated briefly as follows: In cases where a term has not received a practically universal acceptance the oldest known group name (above a generic name and later than 1758) is applied, which covers approximately the same aggregate of genera, irrespective of the comparative rank assigned to the group by the original namer, provided its form is not the plural of a current genus in the same class, nor that now universally adopted for superfamilies, families, and subfamilies.

The application of this principle may be shown by a few examples. Thus *Salientia* is adopted as being the oldest known group name for the present order, because when originally proposed it covered the same aggregate of genera as now, and none of the later names of the same compass have received universal acceptance. On the other hand, *Gradientia* has been rejected, although it is older than *Caudata*, and the latter adopted, because the term *Gradientia*, as originally proposed, embraced the saurian reptiles as well. Oppel's *Squamata* has been accepted for the reptilian order containing the lizards and snakes, although it included the crocodiles, a discrepancy too slight to affect the general applicability of the term. Again, *Testudinata*, although proposed four years later than Link's *Cataphractæ*, has been accepted in preference to the latter (after the elimination of the older terms *Testudines* and *Chelonia* as conflicting with the earlier or synchronous generic names *Testudo* and *Chelonia*) because of the comparative currency of that term, as against the almost total obscurity of the one coined by Link. Finally, the suborder containing the soft-shelled turtles is called *Chilotæ* in spite of the fact that the term *Trionychoidea* is both earlier and commonly adopted, the reason for the rejection of the latter name being that its form is identical with that of the superfamily.

The *Salientia* may be divided into three suborders, viz, *Aglossa*, *Linguata*, and *Costata*, of which the first one is distinguished by having the eustachian tubes united into a single pharyngeal opening and no tongue, while the last one is amply characterized by the presence

of ribs. The Aglossa are only found in Africa and South America; the other two suborders are represented within our area.

Suborder COSTATA.

1879. *Costati* LATASTE, Act. Soc. Linn. Bordeaux, XXXIII, 1879, p. 339.

Salientia having a tongue; eustachian tubes, when present, with two pharyngeal openings; ribs present and a pair of transverse processes near the proximal end of the coccyx. Larvæ with one spiraculum in the middle of the thoracic region.

The Costata, which embrace only one family, the Discoglossidæ, have a very wide and disconnected distribution showing their great antiquity and fully bearing out the verdict of Professor Howes, based upon anatomical investigations, that they are "by far the least modified of living anura."^a

The principal center of distribution at present is central and southern Europe, where three genera together with five species are at home, one of these species extending its range into northwestern Africa and another into central Russia. At the other extreme of the Palearctic region another species of one of the European genera maintains its isolated existence, namely, the species to be treated of in this work, while in the Yunnan region another one has recently been discovered. Two more genera belong to this family, both widely removed from the above, namely, *Liopelma*, the single species of which is the only batrachian occurring in New Zealand to which antipodal island group it is peculiar and where it seems to be very rare, and *Ascaphus*, represented thus far by two specimens only, of a single species, *A. truei*, which has recently been discovered in the State of Washington, not far from the Pacific coast. The extreme rarity of the last two species would seem to indicate that they are nearing extinction.^b

Family DISCOGLOSSIDÆ.

Only this family is included in the suborder.

Genus BOMBINA^c Oken.

1816. *Bombina* OKEN, Lehrb. Zool., II, p. 207 (type, *Rana bombina*).



1820. *Bombinator* MERREM, Tent. Syst. Amph., p. 178 (same type).

1830. *Bombitator* WAGLER, Nat. Syst. Amph., p. 206 (emendation).

^a Proc. Zool. Soc. London, 1888, pp. 178, 506.

^b See Stejneger, A Résumé of the Geographical Distribution of the Discoglossoid Toads in the Light of Ancient Land Connections, in Bull. Amer. Geogr. Soc., XXXVII, No. 2, Feb. 1905, pp. 91-93; and The Geographical Distribution of the Bell-Toads, in Science (n. s.), XXII, Oct. 20, 1905, p. 502.

^c From the medieval Latin verb *bombinare*, to utter a *bombus*, a loud hollow sound, such as the stroke of a bell.



BOMBINA ORIENTALIS. (FROM BOULENGER.)
FOR EXPLANATION OF PLATE SEE PAGE 554.

The genus *Bombina* has a very curious distribution. Two species, *Bombina bombina* (Linnæus) and *Bombina salsaa* (Schrank), occur in Europe, the former in northern and eastern central Europe. The latter, which seems to embrace several local forms, such as *B. salsaa pachypus* (= *Bombinator apenninicus* Gistel and Bronne, 1850), in Italy, *B. salsaa nigriventris* (Ducringen) in Bosnia and Montenegro, *B. salsaa kolombatovici* (Bedriaga) in Dalmatia, etc., occupies France, parts of southern central Europe, Italy, and the Balkan peninsula. None of these species extends eastward beyond the Volga. A third species of the genus occurs in eastern Asia, though probably not to the west of a line through Lake Baikal. A fourth species, *Bombina maxima*, has recently been described from Yunnan at an altitude of 6,000 feet.

BOMBINA ORIENTALIS *b* (Boulenger).

Plate VII.

1878. *Bombinator igneus* MARTENS, Preuss. Exped. Ost-Asien, Zool., I, p. 384 (not of Laurenti) (Chefu, N. China).—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 447 (part: Chefu, China).—BOETTGER, Offenbach. Ver. Naturk. 24-25 Ber., 1885, p. 164 (Chefu).
1890. *Bombinator orientalis* BOULENGER, Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 143, pl. ix, fig. 2^c (type-localities, N. China, Korea, Khabarovka; types in Brit. Mus.); Proc. Zool. Soc. London, 1890, p. 326 (Chefu; Korea; Ussuri).—WERNER, Abh. Bayer. Akad. Wiss., (Muenchen) II Klasse, XXII, Pt. 2, 1904, p. 359 (Tsingtau, China).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, (8) XVII, No. 1 (1905) p. 415 (Vladivostok; Korea, etc.).


Description (figs. 39-43).—*Adult male*; U.S.N.M. No. 17522; Fusan, Korea; 1885; P. L. Jouy, collector. Vomerine teeth in two

^a In his Tailless Batrachians of Europe, p. 322, footnote, Doctor Boulenger, in adopting the sixteen years younger name *Rana agilis* in preference to the older *R. dalmatina*, as required by "the strict application of the law of priority," pleads that "this is one of those cases in which, it appears to me, conservatism is desirable," and he continues: "Similar considerations have guided me in the naming of the two species of the genus *Bombinator*, and I hope, in the interest of stability of nomenclature they will commend themselves to future workers." It is indeed not uncommon to hear the breaking of this law termed "conservatism," and certainly the conservatives ought to know if that is a correct definition, but we radicals prefer to submit to the law as the only means of obtaining the desired "stability of nomenclature." If Boulenger, in the two cases cited, had followed the law we would by this time have reached stability in their cases, while now it has been postponed probably for a generation. Who is to decide in which cases it is allowable to break the law and substitute "conservatism?" A simple "it appears to me" applied here and there by a dozen herpetologists would cause a fine "stability!"

^b Eastern.

^c Reproduced in this work on Plate VII.

small transverse oval groups behind the level of the choanæ; tongue oval, truncate behind, large, adherent all around; snout flat, rounded, without canthus rostralis; nostrils on top of snout about halfway between eye and tip of snout, distance from each other equaling the interorbital space, which is narrower than the upper eyelids; no tympanum; parotoid gland long, narrow, indistinct; no gular sac; a very faintly indicated gular fold; fore leg well developed, longer than half the length of head and body; fingers short, depressed, first shortest, third longest, second and fourth of about equal length; no subarticular tubercles, but a large and distinct inner palmar pad and a smaller indistinct outer one; a large patch of horny nuptial excrescences on the inner side of the fore arm, and smaller patches on the inner side of the large palmar pad, and of first, second, and third toes; when hind leg is carried forward along the side, the heel reaches



FIGS. 39-43. —*BOMBINA ORIENTALIS*. $1\frac{1}{2} \times$ NAT. SIZE. 39. TOP OF HEAD; 40. SIDE OF HEAD; 41. OPEN MOUTH; 42. UNDERSIDE OF RIGHT HAND SHOWING NUPTIAL EXCRESCENCES; 43. FOOT. No. 17522. U.S.N.M.

the corner of the mouth and tarso-metatarsal joint slightly beyond the tip of the snout; tibia equals distance from base of inner metatarsal tubercle to tip of longest toe; heels just meet when the femurs are placed at right angles to the axis of the body; toes short, webbed nearly to their tips, webs deeply excised, third toe longer than fifth; no subarticular tubercles; inner metatarsal tubercle small and weakly developed, no outer tubercle; no nuptial excrescences on toes; upper surface covered with closely set and irregularly arranged prominent tubercles of varying size, all sprinkled over with minute, dark brownish points and each surmounted by one or more horny spines, the base of which is pale and the tip blackish brown; underside smooth, with the posterior part of the belly and proximal parts of femurs somewhat rugose; outer side of sole of feet and underside of base of fifth toe with spines like the upper side. Color (in alcohol) above dark drab-

gray, with irregular obscure dusky spots and dusky cross-markings on the limbs; a pale sage-green spot of irregular outlines across the back between the fore legs and two smaller ones, still more indefinite on each side of the back about the middle; a pale longitudinal stripe on the tip of the snout from the level of the nostrils to the lip; lips dusky; underside pale buff (red in life), with numerous blackish, sharply defined, irregular spots; palms buff (red), separated from the similarly colored under surface of the fore arm by a black wristband; a large buff (red) spot on the inner side of the plantar surface, involving the entire first toe, below and above, and continuous with the similarly colored patch on the inner side of the tarsus; tips of all the digits abruptly and bright buff (red) above and below.

Dimensions.

	<i>mm.</i>
Tip of snout to vent.....	44
Tip of snout to gular fold.....	16
Width of head.....	16
Width of interorbital space.....	3
Fore leg.....	25
Hind leg, from vent.....	62
Foot from base of inner metatarsal tubercle to tip of longest toe.....	18

Sexual variation.—The males of this species during the breeding season possess groups of blackish horny excrescences on the inner side of fore arm, palmar tubercle and first, second, and third fingers exactly as in *Bombina salsa*; but, unlike the latter, there are none on the posterior extremities. In the female the fingers are longer and slenderer, especially the outer one. The dorsal spines and asperities are smaller.

Color variation.—In the series before me there is very little deviation from the coloration of the underside as described above. In two specimens, Nos. 17527 and 17530, the blackish markings are larger, especially in the latter, in which the underside may be described as black with buff (red) spots; in both, the plantar light patch is separated from the tarsal patch by a narrow black bridge, in No. 17530 only on one foot, however. The upper surface is considerably more olive in some specimens and the dusky spots and markings above much better defined and darker.

Habitat.—The eastern bell-toad is confined to northeastern China, Korea, and the Ussuri country. Specimens from all of these localities are in the British Museum. The U. S. National Museum possesses nine specimens collected in 1885 by P. L. Jouy near Fusan, Korea. Pleske's specimens from Vladivostok are in the Academy Museum in St. Petersburg, and the Hamburg Museum also has it from the same locality (No. 862), collected by Dieckmann in 1894. In the St. Petersburg Museum there are also specimens collected by Poljakow at the Novgorodski Port, Possiet Bay.

List of specimens of Bombina orientalis.

U. S. N. M. No.	Sex.	Locality.	When collected.	By whom collected.
17522	Male ^a	Fusan, Korea	1885	P. L. Jouy.
17523	do	do	1885	Do.
17524	do	do	1885	Do.
17525	do	do	1885	Do.
17526	do	do	1885	Do.
17527	do ^b	do	1885	Do.
17528	do	do	1885	Do.
17529	Female	do	1885	Do.
17530	Male ^b	do	1885	Do.

^a Description, p. 51; figs. 39-43.

^b P. 53.

Suborder LINGUATA.

1845. *Linguata* GRAVENHORST, Thierreich, p. 43.

1847. *Phaneroglossa* GERVAIS, Dict. d'Hist. Nat., IX, p. 721.

Salientia having a tongue; eustachian tubes with two pharyngeal openings; lacking ribs and transverse process to urostyle. Larvæ with one spiraculum on left side only.

The term *Ranæ phaneroglossæ* used in 1830 by Wagler (and usually quoted simply as "Phaneroglossa") included the majority of both Salientia and Caudata. Duméril and Bibron first restricted the "Phanéroglosses" (first latinized by Gervais in 1847, so far as I have been able to ascertain) to the Salientia with a tongue, therein including also the Discoglossidæ. Those who still regard these as belonging to the same suborder as the other tongued Salientia will probably retain the name Phaneroglossa. Gravenhorst's *Linguata* may then be restricted to the latter without the genera having ribs. Originally, it is true, it was meant to be equivalent to the Phanéroglosses, but it has never been accepted for them, and the Discoglossidæ form so small a portion that it is better to retain it as here defined than to create a new name.

The families to which the toads and frogs inhabiting our region belong may be distinguished as follows:

KEY TO FAMILIES.

- a¹ Clavicle and coracoid of each side connected by a longitudinal arched cartilage, which overlaps that of the opposite side.
 - b¹ Both jaws toothless..... BUFONIDÆ
 - b² Upper jaw toothed..... HYLIDÆ
- a² Clavicles and coracoids of both sides firmly united by a single median cartilage.
 - b¹ Both jaws toothless; sacral diapophyses dilated..... ENGYSTOMIDÆ
 - b² Upper jaw toothed; sacral diapophyses not or but slightly dilated..... RANIDÆ

The characters of the two groups indicated by a¹ and a² correspond to those of Cope's Arcifera and Firmisternia, respectively, divisions which Boulenger calls "series."

As the employment of the systematic arrangement given above for purposes of identification would necessitate dissection of the specimens, I add the following artificial Key, by which the Salientia occurring within our limits may be referred to their respective families by external characters alone:

KEY TO THE JAPANESE FAMILIES OF TAILLESS BATRACHIANS.

- a¹ Digits without subarticular tubercles; pupil triangular (Suborder COSTATA).....
DISCOGLOSSID.E. p. 50
- a² Digits with subarticular tubercles; pupil round or horizontal (Suborder LINGUATA).
- b¹ A parotoid gland.....BUFONID.E. p. 55
- b² No parotoid gland.
- c¹ Tongue without posterior prolongations or "horns."
- d¹ Upper jaw toothed.....HYLID.E. p. 75
- d² Upper jaw toothless.....ENGYSTOMID.E. p. 87
- c² Tongue with two posterior horn-shaped prolongations.....RANID.E. p. 92

Family BUFONID.E.

A large family having no maxillary teeth, a large parotoid gland behind the head, shoulder-girdle arciferous, and lateral sacral processes dilated. Distributed over the entire globe, except the polar regions, New Zealand, and Polynesia. Some forms are aquatic, some arboreal, some burrowing in their habits, but most are terrestrial. They feed largely on insects, and are among the best friends of the farmer and gardener, by whom they ought to be protected instead of persecuted and exterminated, as they usually are, on account of their often unattractive appearance.

Only one genus is represented within the territory covered by this work.

Genus BUFO^a Laurenti.

1768. *Bufo* LAURENTI, Syn. Rept., p. 25 (type, *B. vulgaris*).
1788. *Bufo* LACÉPÈDE, Hist. Nat. Quadr. Ovip., I, Syn. Meth. (emendation).
1814. *Batrachus* RAFINESQUE, Specchio Sci., (Palermo), II, fasc. 7 (substitute for *Bufo*).
1815. *Bufotes* RAFINESQUE, Analyse Nat., p. 78 (substitute for *Bufo*).
1816. *Calamita* OKEN, Lehrb. Zool., II, p. 209 (type, *B. calamita*; not of Schneider 1799).
1828. *Chascax* RITGEN, Nova Acta Acad. Carol. Leop., XIV, p. 278 (type, *Bombinator strumosus*).
1843. *Phryne* FITZINGER, Syst. Rept., p. 32 (type, *B. vulgaris*).
1843. *Phrynoïdis* FITZINGER, Syst. Rept., p. 32 (type, *B. asper*).
1864. *Epidalea* COPE, Proc. Phila. Acad., 1864, p. 181 (type, *B. calamita*).

The above synonyms are based upon species occurring within our limits or are additional to those found in previously published synonymies. The other synonyms of *Bufo* are too numerous to be reproduced here.

The status of the Japanese toads is still very unsatisfactory.

^a The Latin word for toad.

It will be remembered that Schlegel, in *Fauna Japonica*, described the toad of Japan as a variety of the European toad, to which he attached the name *Bufo vulgaris japonicus*, suppressing Boie's name *B. prætextatus*, as he says, "because of the great affinity which this toad has to that of Europe, of which it is the representative in Japan." The differences indicated by him are chiefly those of coloration.

Camerano, in 1879, found the differences to be of such a character as to induce him to regard the Japanese toad as a species, *Bufo japonicus*, his principle being that any well-defined local form should be so considered. His material consisted of 18 specimens in the Zoological Museum at Turin, nearly all being from Tokyo and Yokohama.

In the following year, 1880, we find both Lataste and Boulenger treating of the subject. The former, who examined fourteen specimens from Peking and two specimens from Japan (lent him by Camerano), expresses himself in favor of recognizing a *Bufo vulgaris japonicus*, while the latter, with sixteen specimens from various Chinese localities and five Japanese, pronounced in favor of an undivided *Bufo vulgaris*. The chief characters, he stated, are the more prominent and spinous warts and the black horny layer on various parts of the body, the rather larger head, and the blackish stripe on the flanks. None of these characters appears to him to be constant, and he adds that if the Japanese form should be separated from the European it should certainly also be separated from the Chinese: "But I do not think that distinction necessary." This conclusion he adhered to in his Catalogue of *Batrachia Salientia*, issued two years later. It should be noted, however, that in the description of the species he was obliged repeatedly to make exceptions for Chinese or Japanese specimens.

Additional material did not alter his view, for, in 1890, in a paper on the reptiles and batrachians of Amurland, he treats of *Bufo vulgaris* in the following language:

Its eastern range extends over nearly the whole of China and Amoorland and Japan. The differences between European and Japanese specimens, which consist chiefly in the greater size and perfect distinctness of the tympanum, the black lateral stripe, and the deep black spots or marblings of the lower parts in the latter, are completely bridged over by the Chinese and Manchurian specimens. Specimens from Ichang, on the Yangtse Kiang, and Ningpo come nearest the Japanese, from which they do not differ in coloration; but the tympanum, although as distinct, is not so large. Specimens from Shanghai, Chefoo, Peking, and Korea are intermediate between the latter and the European; the tympanum is always very distinct, but varies considerably in size; the dark lateral stripe is usually ill-defined or absent, and the belly may be either largely spotted with black or almost immaculate. Judging from the two specimens collected at Chabarowka by Hr. Dörries, the northernmost form is still nearer the European; the tympanum is rather small, but perfectly distinct, the belly is immaculate, and the coloration might be said to be identical with that of European specimens, but for the presence of traces of a light vertebral line, as is often found in specimens from Japan, Korea, and Northern China.

This statement is almost literally repeated in his latest work on the Tailless Batrachians of Europe (1898).

In the meantime (1883) he had described and figured a *Bufo formosus* from two specimens collected by the *Challenger* Expedition at Yokohama, which "in its general characters agrees with *Bufo vulgaris*, especially the Japanese form." The main distinctions relied on are the large size of the tympanum, the greater length of the hind leg, the more elongate digits, more excised webs of the hind foot, and the shortness of the first finger.

Nothing more is heard of this species. In the subsequent discussions of the "Japanese form" of *Bufo vulgaris* there is not the slightest reference to it. The fact, however, that the name is not included in Boulenger's synonymy of *Bufo vulgaris* in his Tailless Batrachians of Europe, nor the types in the list of specimens appended, seems to indicate that he still maintains its distinctness, although some doubt arose in my mind upon seeing that in the appendix to the latter work he enumerates a specimen from Dagoshima,^a collected by Doctor Anderson in May, 1884, under *Bufo vulgaris*. To me this specimen, which the authorities of the British Museum kindly allowed me to examine in 1898, is a strongly marked *Bufo formosus* with the types of which I compared it.

Much of the uncertainty is probably due to the confusion of these two toads in Japan. Thus, on examining the large series in the Leiden Museum upon which Schlegel founded his *Bufo vulgaris japonicus*, I discovered that he had both forms before him. I think it is also certain that Camerano based his conclusion of the specific distinctness of *Bufo japonicus*, or *B. pratertatus*, as he called it in his elaborate paper of 1901, upon true *Bufo formosus*.

The status of the Japanese toad resolves itself into three queries:

- (1) Is there more than one species of *Bufo* in Japan?
- (2) Is one of these identical with the species on the mainland?
- (3) In case of different forms being recognized, what names are to be employed?

First, then, are there more than one species of *Bufo* in Japan proper? In the material which I have had an opportunity to examine, viz, 13 specimens in the Leiden Museum, 9 specimens in the British Museum, and 36 in our own National Museum, there appear to be three forms. One has larger tympanum, longer limbs, slenderer digits, much more deeply excised webs, longer and narrower parotoids, and, as a rule, the first finger shorter than the second. This is the so-called *B. formosus*. That we have not to do with sexual or seasonal differences seems plain. Thus the males with the nuptial asperities on their fingers, which under this assumption would have the greatest extent

^a Probably Dogoshima, near Miyanoshta.

of web, on the contrary show a minimum size of the interdigital membranes.

A very similar form, possibly confined to Shikoku, having the deeply excised webs, long parotoids, long limbs, and slender digits, is very near to *B. formosus*, but has a much smaller tympanum and first and second fingers subequal.

The third form is characterized by small tympanum, stouter and shorter limbs, much larger webs, the true *B. japonicus*.

In spite of the general similarity in coloration and otherwise it is difficult to escape the conclusion that the Japanese toads represent three more or less distinct forms, and as it is at present impossible to establish their interrelationship clearly it is thought best to designate them by binominal appellations. The great difficulty arises in ascribing definite geographical limits to them. Unfortunately, a large number of the specimens examined are without indication of their exact origin. Thus all the specimens in the Leiden Museum collected by Buerger and von Siebold are only labeled "Japon." Several in the British Museum are similarly deficient. The only true *B. japonicus* with a definite locality attached seems to be one from Kobe in the British Museum and one from Miyakoshima in Science College Museum. On the other hand, it would almost appear as if all the *B. formosus* of which the exact locality are known came from near Tokyo and Yokohama (Camerano's specimens; types in British Museum; specimens in United States National Museum) and Dogoshima, wherever that locality may be.^a We have thus an indication of the *B. formosus* being eastern and northern, while the one with the larger webs, etc., may be more western.

Finally, the only authentic locality from which I have seen specimens of the new form, further on described as *B. smithi*, is the island of Shikoku. There is a specimen in the Imperial Museum in Ueno Park, Tokyo, apparently belonging to this form, which was in a bottle with the type specimen of *Polypedates ishikawæ* and said to have come from Okinawa shima, Riu Kiu. As no toad has been recorded by any collector from that archipelago, the locality given becomes very doubtful.

The next question then is whether any one of the forms which we thus recognize as occurring in Japan is identical with the toads occurring on the mainland. I have already quoted Boulenger to the effect that there are several forms, though he does not recognize them by name because they intergrade. Let it be understood at once that the form in Japan most remote from the mainland forms is *B. formosus*. It is consequently the so-called *B. vulgaris japonicus*, i. e., the one with the greater webs, shorter parotoids, shorter legs, etc., which

^a If Dogashima is meant, as surmised by me, the locality agrees well with the others.

comes nearest to the mainland forms. Boulenger then maintains that this form approaches the Chinese specimens from Ningpo and Yangtse-Kiang, while the Shanghai, Chefu, Manchurian, Korean, and Ussuri specimens are intermediate between the Japanese form and the true European *Bufo vulgaris*. The differences are chiefly those of color, and the relative size and distinctness of the tympanum. Thus Boulenger, in discussing the characters of *Bufo formosus* asserts that a black line bordering the lower lip "is constant in the Japanese specimens of *B. vulgaris*, absent in the continental specimens." On the whole, my own observations corroborate those of Boulenger; yet while it is admitted that the separating characters are somewhat vague and uncertain, it is considered best to recognize the various local forms for the present at least until more abundant and more reliably labeled material shall have been accumulated. It may then be expedient to recognize three oriental forms of the common European toad, one from eastern South China, one from northeastern China, Korea and Amurland, and one from Japan. The status of a fourth one, *B. sachalinensis*, is more uncertain yet.

The third and final question now arises as to the names to be applied to these forms. As for the Japanese form, the question is somewhat intricate because of the occurrence of several forms or species in the islands. Unfortunately there is nothing in Boie's description of *Bufo praterstatus* by which it can be decided which of these he had before him, and as the type is not in existence, his name may be left out as unidentifiable. Schlegel's material upon which he based his *Bufo vulgaris japonicus*, as has been shown, was composed of specimens of both forms. The illustrations in *Fauna Japonica*, although very good, fail to throw much light on the question, for while the full figure apparently represents the short-webbed form (*B. formosus*), the profile of the head seems to be taken from a specimen with a smaller tympanum, possibly from a *B. smithi*. The illustrations as well as the description and the type-material consequently are composite. It would then seem that Boulenger's action in describing the short-webbed form as *Bufo formosus* and relegating *B. vulgaris japonicus* to *B. vulgaris* should not now be undone, and that the names must stand as he restricted them.

The South China form must stand as *Bufo bufo gargarizans* (Cantor)^a while the North China form probably is properly named *Bufo bufo asiaticus* (Steindachner), as Boulenger refers the Shanghai specimens to the same category as the Chefu and Manchurian ones.

^a1842. *Bufo gargarizans* CANTOR, Ann. Nat. Hist., IX, Aug. 1842, p. 483 (type-locality, Chusan, China; types in Brit. Mus.).

1860. ? *Bufo griseus* HALLOWELL, Proc. Phila. Acad., 1860, p. 506 (type-locality, marshes of Whampoa, Hongkong; Stimpson, collector).

The Japanese and North China forms and species may be distinguished as follows:^a

KEY TO THE SPECIES OF BUFO IN JAPAN, FORMOSA, KOREA, AND SIBERIAN COAST PROVINCE.

- a*¹ Top of head without bony ridges.
- b*¹ Inner edge of tarsus rounded, tubercular, with no sharp cutaneous fold.
- c*¹ Web between toes extending only as far as first basal phalanx of third toe and one-half the first basal phalanx of fourth (fig. 48).
- d*¹ Diameter of tympanum about twice as large as its distance from the eye.
Bufo formosus, p. 60
- d*² Diameter of tympanum about equals the distance from the eye.
Bufo smithi, p. 64
- c*² Web between toes extends as far as end of the two basal phalanges of third toe and first basal phalanx of fourth.
- d*¹ Underside uniform pale, or with small black spots.
Bufo bufo asiaticus, p. 66
- d*² Underside marked with large black blotches.
Bufo bufo japonicus, p. 69
- b*² Inner edge of tarsus with a sharp cutaneous fold.
Bufo raddi, p. 70
- a*² Top of head with bony ridges.
Bufo melanostictus, p. 72

BUFO FORMOSUS^b Boulenger.

Plate VIII.


1838. *Bufo vulgaris japonicus* SCHLEGEL, Fauna Jap., Rept., pp. 106, 139 (Japan; specimens in Leiden Mus.; part only); Sauri et Batr., pl. II, fig. 6.
1876. *Bufo vulgaris* var. *pratextatus* MARTENS, Preuss. Exped. Ost-Asien, I, p. 384 (Yokohama) (not of Boie?).—HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 120.
1879. *Bufo japonicus* CAMERANO, Atti Accad. Torino, XIV, Pt. 5, Apr., 1879, p. 884 (Yokohama; Tokyo; Mus. Zool. Turin; Aimonin and Fontanesi, collectors).
1883. *Bufo formosus* BOULENGER, Proc. Zool. Soc. London, 1883, p. 140, pl. xxiii. c (type-locality, Yokohama; types in British Museum; *Challenger* Exped. collect.)—OKADA, Cat. Vert. Jap., 1891, p. 66.—BOETTGER, Kat. Batr. Mus. Senckenberg., 1892, p. 37 (Nikko; Nippon; Japan).
1900. *Bufo pratextatus* CAMERANO, Mem. Accad. Torino, (2) L, p. 113; author's separate, p. 33, pls. I-II (not of Boie?).

Bufo formosus is most easily distinguished from *B. japonicus* by its longer fore limbs, longer and slenderer digits, and much shorter webs between the toes. The alleged difference in the relative size of first and second finger, does not hold in every instance, for I have two specimens of undoubtedly the same species, from Gensan, Korea, in which the relation is reversed, but as a rule it may be said that in *B. formosus* the first finger is shorter than the second, while in *B. japonicus* it is equal to or longer than the second. In *B. formosus*, moreover, the interorbital space appears to be proportionally narrower.

^a*Bufo sachalinensis* is not given a place in this key, as the characters utilized in the latter are not indicated in the only description of this form thus far published.


^bSignifying Handsome.

^cReproduced in this work on Plate VIII.



BUFO FORMOSUS. (FROM BOULENGER.)
FOR EXPLANATION OF PLATE SEE PAGE 554.

Description (figs. 44–48).—*Adult male*; U.S.N.M. No. 23540; Yokohama; September, 1896; L. Stejneger, collector. Snout somewhat pointed, rising steeply from the labial edge, the canthus rostralis fairly well marked and the loreal region slightly concave; nostrils about halfway between eyes and tip of snout, their distance from each



FIGS. 44–48. —BUFO FORMOSUS. NAT. SIZE. 44, TOP OF HEAD; 45, SIDE OF HEAD; 46, UNDERSIDE OF HAND; 47, UNDERSIDE OF FOOT; 48, UPPER SIDE OF FOOT, TO SHOW EXTENT OF WEB. No. 23540, U.S.N.M.

other greater than their distance from the eyes, but less than their distance from the labial edge or than the upper eyelids, and only two-thirds the width of the interorbital space; interorbital space very slightly concave, considerably wider than the upper eyelid; tym-

panum very distinct, circular, nearly as large as the diameter of the eye, its distance from the latter less than half its own diameter, its distance from the labial edge slightly less than the diameter; parotoid glands rather long and narrow, more than two and a half times as long as wide; fore legs long and stout; fingers with slight webs at the base, tips rounded, third longest, second shortest, and considerably shorter than first; two large palmar tubercles, the inner one particularly prominent with a cutting edge, resembling the inner metatarsal tubercle; nuptial asperities on the inner side of this tubercle, on the inner and upper sides of first and second fingers and on the inner side of the third finger; heels not meeting when folded hind legs are placed perpendicular to the axis of the body; toes one-third webbed, the web extending only as far as the first basal joint of the third toe and one-half the first basal joint of the fourth toe; inner metatarsal tubercle large, with prominent cutting edge, longer than the first toe; outer metatarsal tubercle smaller, rounded, prominent, but less so than the inner one; no tarsal fold; some of the subarticular tubercles double both under fingers and toes; back covered by numerous round tubercles of various sizes, none as large as the tympanum, all rather smooth and not arranged symmetrically, though the larger ones are more numerous toward the sides of the back; under side divided up into small areas by innumerable transverse furrows, each area having one or more small granules or tubercles with a small brown horny tip, these tips becoming larger and almost spiny on the hind legs. Color (in alcohol) on upper side of body nearly uniform sepia; from the posterior corner of the eye backward, margining the lower edge of the parotoid and along the sides of the body to the groin a wide, sharply contrasted, black band; above the latter on the side a smoke-gray border fading into the sepia of the back; a narrow black streak on the lower jaw along its whole length; underside of body and limbs dirty whitish with large strongly defined vermiculations and anastomizing black spots, contiguous with similar though less sharply defined spots on the upper surface of hands and feet; palmar and metatarsal tubercles as well as tips of fingers and toes raw umber.

Dimensions.

	<i>mm.</i>
Tip of snout to vent.....	110
Tip of snout to anterior border of tympanum.....	24
Nostril to tympanum.....	18
Internarial width.....	6
Interorbital width.....	10
Diameter of tympanum.....	8
Distance of tympanum from eye.....	3
Width of head.....	46
Fore leg.....	66
Hind leg from vent to tip of longest toe.....	147

	<i>mm.</i>
Tibia	42
Hind foot from base of inner metatarsal tubercle to tip of longest toe.	52
Length of parotoid	21
Width of parotoid.....	8

Variation.—The variation of this and allied species has been mentioned above under the general discussion of their status. It may be repeated here that the relative length of the first and second fingers is not constant. Thus, in the type, “the first finger being laid against the second does not extend quite so far as the latter.” The same relation holds in over No. 11348, but in my two topotypes these proportions are reversed. The second of these has the whole underside almost uniform whitish with a few scattered dusky spots.

The females differ from the males in this and allied forms in the less-developed fore legs and somewhat larger webs between the toes. They also lack the nuptial pads of asperities on the fingers described above which characterize the males not only during the breeding season but for a long time after, as shown by my specimens which were taken in September.

Habitat.—As indicated above, the only positive locality whence we have undoubted specimens of this species is the neighborhood of Tokyo and Yokohama, to which must be reckoned “Mount Fuji.” The locality “Dagoshima” attached to a specimen in British Museum which was collected by Doctor Anderson on May 24, 1884, I have failed to find on any map.^a A great amount of careful collecting and critical identification will be necessary before the range of this and allied forms in Japan can be determined.

List of specimens of Bufo formosus.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Di-	Dis-
						am-	tan-
						<i>mm.</i>	<i>mm.</i>
U. S. N. M.	11348	Adult.	Japan.....		E. S. Morse.....	5	2.5
Do.	23540	Male ^a	Yokohama.....	Sept. 1896	L. Stejneger.....	8	3
Do.	23541do.....do.....do.....do.....	8	3.5
Do.	34309do.....	Mount Fuji.....	Aug. 1898	A. Owston.....	9	4
Do.	34310	Female.....do.....do.....do.....	8.5	3
Do.	34311do.....do.....do.....do.....	9	3.5
Do.	34312do.....do.....do.....do.....	9	4
Do.	34313	Male.....do.....do.....do.....	8.5	3
Do.	34314	Female.....do.....do.....do.....	9.5	3.5
Do.	34315	Male.....do.....do.....do.....	8.5	3
Do.	34316	Female.....do.....do.....do.....	8.5	4
Do.	34317do.....do.....do.....do.....	9	3.5

^a Topotype; description, p. 61, figs. 44-48.

^a Unless it be meant for Dogashima, near Miyanoshta, in Hakone, about 1,300 feet above the sea, consequently not far from the other localities from which we have specimens.

List of specimens of *Bufo formosus*—Continued.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Diameter of tympanum.	
						mm.	mm.
U.S.N.M.	34318	Male	Mount Fuji	Aug., 1898	A. Owston	8.5	4
Do.	34319	do.	do.	do.	do.	8.5	3
Do.	34320	do.	do.	do.	do.	9.5	3.5
Do.	34321	Female	do.	do.	do.	8	4
Do.	34322	Male	do.	do.	do.	8	3
Do.	34323	do.	do.	do.	do.	10	3
Do.	34324	Female	do.	do.	do.	7.5	2.5
Do.	34325	do.	do.	do.	do.	9	3.5
Do.	34326	Male	do.	do.	do.	9	3
Do.	34327	Female	do.	do.	do.	8	3.5
Do.	34328	do.	do.	do.	do.	6	3
Brit. Mus.	82.7.14.27	Male ^a	Yokohama		Challenger Exped.		
Do.	82.7.14.28	Female ^a	do.		do.		
Do.	86.6.30.31	Male	Dagoshima (?)	May 24, 1884	J. Anderson		

^a Types.**BUFO SMITHI**,^a new species.

Diagnosis.—Like *Bufo formosus*, but with much smaller tympanum, its diameter about equaling its distance from the eye.

Habitat.—Shikoku Island, Japan.

Type.—U.S.N.M. No. 31951; Kochi, Province of Toza, Shikoku Island; May 11, 1903; Dr. Hugh M. Smith, collector.

Remarks.—This form is so closely allied to *Bufo formosus* that no further description is necessary, the only difference which I have been able to detect being the smaller tympanum which is situated farther away from the eye than in that species. For the sake of completeness a set of measurements of the type is added.

Dimensions.

Male adult; No. 31951 U.S.N.M.; Kochi, Toza, Shikoku; May 11, 1903; H. M. Smith, collector.

	mm.
Tip of snout to vent	110
Tip of snout to anterior border of tympanum	28
Nostril to tympanum	21
Internarial width	7
Interorbital width	9.5
Diameter of tympanum	6
Distance of tympanum from eye	6
Width of head	43
Fore leg	74

^a Named after Dr. Hugh M. Smith, United States Deputy Fish Commissioner, who collected the type.

	<i>mm.</i>
Hind leg from vent to tip of longest toe	154
Tibia	44
Hind foot from base of inner metatarsal tubercle to tip of longest toe	54
Length of parotoid.....	25
Width of parotoid.....	8

Habitat.—The only definite locality whence we have specimens of this form is Kochi, in the island of Shikoku, where Dr. Hugh M. Smith collected a fine series of specimens in the spring of 1903.

List of specimens of Bufo smithi.

U. S. N. M. Nos.	Sex.	Locality.	When collected.	By whom collected.	Diameter of tympanum.	Distance of tympanum from eye.
					<i>mm.</i>	<i>mm.</i>
31851	Male.....	Suzaki, Toza, Shikoku.....	May 8, 1903	Dr. H. M. Smith....	3.5	3.5
31929	Young.....	Kochi, Toza, Shikoku.....	May 11, 1903do.....	2	2
31942	Female.....do.....	May 7, 1903do.....	5.5	5.5
31943do.....do.....	May 11, 1903do.....	5	4
31944	Male.....do.....do.....do.....	5.5	5.5
31945	Young.....do.....do.....do.....	3.5	3.5
31946do.....do.....do.....do.....	2.5	2.5
31947do.....do.....do.....do.....	2.5	2.5
31948do.....do.....do.....do.....	2	2
31949do.....do.....do.....do.....	2	2
5260	Female.....do.....do.....do.....	6	6
31951	Male ^ado.....do.....do.....	6	6

^a Type.

BUFO SACHALINENSIS ^a (Nikolski).

1905. *Bufo vulg[ar]is* var. *sachalinensis* NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, No. 1, p. 389 (type-locality, Sakhalin; types, St. Petersburg Mus., Nos. 1934-1936; Doctor Suprunenko, collector).

Not having seen any specimens referable to this form, I append a translation of Nikolski's original diagnosis:

Longitudinal diameter of eye contained one and one-half times in length of snout; width of upper eyelid one and three-fourths to two times in interorbital space; forehead smooth (not warty); tympanum very distinct, its diameter equaling two-thirds to three-fourths the longitudinal diameter of the eye; internal metatarsal tubercle very large, its length equaling or exceeding the longitudinal diameter of the eye; a black line always marking the outer margin of the parotoids, as in *B. japonica* Schlegel.

This diagnosis, based upon six specimens, is too brief and does not allude to the crucial characters of the webbing of the toes and to the distance of the tympanum from the eye. Comparing it, however, with specimens of the next species from Korea and with *B. formosus* it almost looks as if the Sakhalin toad may belong to the latter rather

^a From the island of Sakhalin, or Krafto, in the Okhotsk Sea.

than to the former. This, from a zoogeographical standpoint, is rather unexpected, and a comparison of specimens from central Hondo, Yezo, and Sakhalin should be highly instructive.

Habitat.—Described from, and thus far recorded only from, the island of Sakhalin, where it was collected by Doctor Suprunenko in 1890.

BUFO BUFO^a ASIATICUS^b (Steindachner).


1869. *Bufo vulgaris* var. *asiatica* STEINDACHNER, Novara Exped., Zool., I, Amph., p. 39 (Shanghai; Vienna Mus.; Zelebor and Frauenfeld, collectors).
1880. *Bufo vulgaris japonicus* LATASTE, Bull. Soc. Zool. France, V, 1880, p. 506 (Peking, China; de Plancy, collector) (not of Schlegel).
1880. *Bufo vulgaris* BOULENGER, Proc. Zool. Soc. London, 1880, p. 569 (part, Peking); Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 141 (Khabarovka, Ussuri; Korea); Taill. Batr. Europe, Pt. 2, 1898, pp. 213, 347 (part: Khabarovka; Seoul, Chemulpo, Gensan, Korea; Chefoo, Shanghai, China).—BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 145 (Shanghai).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 358 (Tsingtau, Shanghai).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, No. 1, 1905, p. 387 (part: Amur, Ussuri, etc.).
1892. *Bufo vulgaris* var. *japonica* BOETTGER, Kat. Batr. Mus. Senckenberg., p. 37 (part: Shanghai).
1899. *Bufo japonicus* STONE, Proc. Phila. Acad., 1899, p. 183 (Tore River, Eastern Mongolia).

Description (figs. 49–52).—*Adult male*; U.S.N.M. No. 17516; Gensan, Korea; July, 1886; P. L. Jouy, collector. Head wide, snout rather rounded with well-marked canthal angle, loreal region scarcely concave; nostrils slightly nearer the eye than the tip of the snout, their distance from each other greater than their distance from the eye, equaling that from the labial edge and the upper eyelid, and somewhat greater than one-half the width of the interorbital space, which is rather concave; tympanum very distinct, nearly circular, about one-half the diameter of the eye, its distance from the latter nearly equaling its own diameter, and its distance from the labial border twice its diameter; parotoid glands large, their width more than one-half the length; fore legs long; fingers free at base, third longest, second shortest, slightly shorter than first and fourth, which are equal; two very distinct palmar tubercles, the outer more than twice as large as the inner, the latter with nuptial asperities; similar but larger pads on upper and inner side of first and second fingers, and a narrow band on inner side of third; heels not meeting when folded hind legs are placed perpendicular to the axis of the body; toes nearly fully webbed, the web extending as far as the end of the two basal phalanges of the third toe and the first basal phalanx of the fourth, but slightly excised; inner metatarsal tubercle moderate, with prominent cutting edge, considerably shorter than first toe; outer metatarsal tubercle

^a *Rana bufo* Linnæus, Syst. Nat., 10 ed., 1758, I, p. 210.

^b Signifying Asiatic.

slightly smaller, less prominent; no tarsal fold; double subarticular tubercles under all the fingers and under third and fourth toes; upper surface covered with numerous rounded tubercles of various sizes, which show a tendency to a symmetrical arrangement, one longitudinal series extending obliquely backward from the parotoid, while on the back between the parotoids there is a chevron-shaped figure pointing forward and a similar one on the lower back; the tubercles, many of which are larger than the tympanum, are surmounted by a conical brown point, becoming spiny on lower back and upper surface of legs; underside strongly granular or tubercular, with similar,



FIGS. 49-52.—*BUFO BUFO ASIATICUS*. NAT. SIZE. 49, TOP OF HEAD; 50, SIDE OF HEAD; 51, UNDERSIDE OF HAND; 52, UNDERSIDE OF FOOT. No. 17517, U.S.N.M.

though less developed, pointed asperities. Color (in alcohol) above pale grayish olive, with some ill-defined longitudinal vermiculating markings of an olive-buff color on both sides of back; a light band similarly colored from parotoids obliquely backward to groin and upper surface of limbs, with cross bands of the same tint; a dark slate-colored band from posterior angle of eye curving down over anterior lower portion of parotoid to insertion of arm, and a similarly colored spot on the posterior outer aspect of the gland, continuing backward below the lateral pale band and fading gradually into the

general grayish olive; tubercles on upper surface topped with umber brown, and the upper surface of the parotoids vermiculated with the same color; underside uniform, unspotted pale ochre yellow, palmar and metatarsal tubercles, as well as nuptial pads, brown; edge of upper lip and lower edge of mandible slightly tinged with the same brown color, but no dusky line on the lower jaw.

Dimensions.

	<i>mm.</i>
Tip of snout to vent.....	84
Tip of snout to anterior border of tympanum.....	19
Nostril to tympanum.....	14
Internarial width.....	4
Interorbital width.....	7
Diameter of tympanum.....	3
Distance of tympanum from eye.....	2.5
Width of head.....	30
Fore leg.....	52
Hind leg from vent to tip of longest toe.....	110
Tibia.....	30
Hind foot from base of inner metatarsal tubercle to tip of longest toe.....	39
Length of parotoid.....	16
Width of parotoid.....	9

The *adult female* (U.S.N.M. No. 17517; same locality, date, and collector) (figs. 49-52) is larger; total length from snout to vent, 96 mm.; differs otherwise from the male chiefly in the somewhat shorter legs and in the tubercles of the skin being more pointed and more extensively pigmented, even on the underside all the tubercles being tipped with brown; that she lacks the nuptial excrescences on the fingers is a matter of course.

Another *adult female* (U.S.N.M. No. 21214; Seoul, Korea; P. L. Jouy, collector) is darker, the ground color above being brownish, and the pale markings clay color; the lower edge of the parotoids involved in a narrow black band, which continues for some distance backwards on the side, and a crescent-shaped branch descending from the middle of the parotoid to in front of the insertion of the arm; on the sides of the belly there are a number of small, black, isolated spots, none as large as the tympanum.

A *young* specimen, 41 mm. long, also from Korea (Chemulpo; Jouy, collector; U.S.N.M. No. 21215) is uniformly pinkish brown above and unspotted below, except a few blackish specks on throat and chest; there is a dusky bar descending from the eye to the edge of the lip and some indication of dusky markings on the upper lip and below the nostrils.

Habitat.—The northeastern continental representative of the European common toad (*Bufo bufo*) is found in eastern Mongolia, Manchuria, Amurland at least as far north as Khabarovka, Korea, north-eastern China from Peking and Chefu to Shanghai, beyond which is found the southeastern representative, *B. bufo gargarizans*.

In Korea it has been found at Gensan, as well as at Chemulpo and Seoul. From Fusan we have thus far no records.

List of specimens of Bufo asiaticus.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected.
17516	Male ^a	Gensan, Korea	July, 1886	P. L. Jouy.
17517	Female ^b	do	do	Do.
21214	do. ^c	Seoul, Korea		Do.
21215	Young ^c	Chemulpo, Korea	Aug., 1883	Do.

^a Description, p. 66.

^b P. 68; figs. 49-52.

^c P. 68.

BUFO BUFO JAPONICUS ^a (Schlegel).

- 1826. ?*Bufo praetertatus* BOIE, Isis, 1826, p. 215; Bijdr. Natuurk. Wetensch., II, Pt. 1, 1827, p. 271 ("Nangasacki,"² Blomhoff, collector).
- 1838. *Bufo vulgaris japonicus* SCHLEGEL, Fauna Japon., Rept., pp. 106, 139 (type-locality, Japan; types in Mus. Leiden; Buerger and von Siebold, collectors; part).—GUENTHER, Cat. Batr. Sal. Brit. Mus., 1858, p. 56 (Japan).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1866, p. 109.
- 1858. *Bufo vulgaris* BLEEKER, Natuurk. Tijdschr. Nederland Indië, XVI, p. 205 (Japan) (not of Laurenti).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1876, p. 384 (Nagasaki).—BOULENGER, Proc. Zool. Soc. London, 1880, p. 569 (part: Japan); Cat. Batr. Sal. Brit. Mus., 1882, p. 303 (part: Japan); Taill. Batr. Europe, Pt. 2, 1898, pp. 213, 347 (Kobe).—OKADA, Cat. Vert. Jap., 1891, p. 66 (Japan).—?FRITZE, Mitth. Deutsch. Gesell. Ost-Asiens, V, 1891, p. 239 (Yezo).

In structural characters and general proportions this island form is apparently identical with the foregoing and needs no special detailed description. Its status has been discussed above, and it was there pointed out that the only appreciable difference from the continental form is one of coloration, inasmuch as the Japanese form has the underside more or less strongly blotched and coarsely mottled with blackish, with a blackish line along the whole lower jaw, as in *Bufo formosus*.

Habitat.—The exact distribution of this form in Japan is highly problematical at the present time. Specimens from Kobe and Nagasaki seem to belong to it, but beyond this statement all is conjecture.

The occurrence of a toad in Yezo is reported by Doctor Fritze, and Doctor Nozawa writes me that he believes the toad is found in the southern part of this island, but it is impossible to say to what form it may belong. A comparison of Yezo and Sakhalin specimens would be highly interesting and instructive.

List of specimens of Bufo japonicus.

Museum.	No.	Age.	Locality.	When collected.	By whom collected.
Sci. Coll. Tokyo....	74	Adult.....	Miyakoshima (?).....		

^a Signifying Japanese.


BUFO RADDEI^a Strauch.

1859. ? *Bufo variabilis* var. *amurensis* MAACK, Putesh. na Amur (p. 153) (Amur-land).
1859. *Bufo calamita* MAACK, Putesh. na Amur, (p. 153) (Amur Valley), (not of Laurenti).
1876. *Bufo raddei* STRAUCH in Przewalski's Mongoliya i Strana Tungutov, II, Pt. 3, p. 53 (Chinese province of Ordos, and Alashan Desert).—LATASIE, Rev. Intern. Sci. (Paris), I, 1878 (p. 437); Bull. Soc. Zool. France, V, 1880, p. 69 (Peking; Chefu).—BOULENGER, Proc. Zool. Soc. London, 1880, p. 551 (Chefu; Amur).—PRZEWALSKI, Reisen in Tibet, 1884 (pp. 254, 257) (Gobi Desert, eastern part).—BOETTGER, Offenbach. Ver. Naturk. 24-25 Ber., 1885, p. 163.—MUELLER, Verh. Naturf. Ges. Basel, VIII, Pt. 2, 1887, p. 258 (Khabarovka).—STONE, Proc. Phila. Acad., 1899, p. 183 (east. Mongolia).—BEDRIAGA, Przewalski Reise, Zool., III, Pt. 1, 1899 (p. 42, pl. I, figs. 1, 1a, 6).—MÉHELY, Zichy's Dritte Asiat. Forschungsr., II, 1901, p. 67 (North China between Daba and Khalgan; Peking).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, (8) XVII, No. 1, 1905, p. 372 (Eastern Siberia, Tibet, Ordos, Amur province, etc.).
1882. *Bufo raddei* BOULENGER, Cat. Batr. Sal. Brit. Mus., p. 294 (Sungatchi; Chefu); Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 141 (Valley of the Amur; Dauria; Peking; Chefu).—WERNER, Abh. Bayer. Akad. Wiss., (Muenchen) II Kl., XXII, Pt. 2, 1904, p. 359 (Tsingtau, China).

Description.—*Adult male*; U.S.N.M. No. 21394; Zagan Olui, Eastern Siberia (figs. 53-57).—Snout rounded, with indistinct canthus rostralis; loreal region but very slightly concave; nostrils nearly equidistant between eyes and tip of snout, their distance from each other slightly less than the width of the interorbital space; upper eyelid slightly wider than interorbital space; tympanum distinct, circular, its diameter about one-half that of the eye, distant from the eye about two-thirds its diameter; parotoid glands large, flat, of somewhat irregular outline, about twice as long as broad; fingers rather short, first and second subequal, fourth very short, not extending beyond the distal end of the basal phalanx of the third; two large palmar tubercles, the outer one especially very large and with a conspicuous central prominence; inner palmar tubercle as well as the first, second, and third fingers with pads of dark brown nuptial asperities; subarticular tubercles single, except one under first finger, which is semidivided; hind legs rather short; heels scarcely meeting when folded legs are placed at right angles to axis of body; toes about half webbed; subarticular tubercles single; inner metatarsal tubercle very prominent, much shorter than its distance from tip of first toe; outer metatarsal tubercle small but very distinct, prominent; a sharp cutaneous fold along the inner side of the tarsus from the inner metatarsal tubercle to the heel; a long prominent oval gland covering nearly the whole upper

^a For Dr. Gustav Ferdinand Richard Radde, late director of the museum in Tiflis, who in 1855-60 explored southeastern Siberia. Born in Dantzig, November 27, 1831; died in Tiflis, March 15, 1903.

aspect of the tibia; skin coarsely granular above and below with numerous irregular glands of varying size on the back, those on either side of the median line and on the sides being rather elongate; an interior gular sac with one large opening on each side of the rather narrow tongue. Color (in alcohol) above brown with a paler median stripe; upper surface of legs with large dusky spots having narrow whitish margins; upper tip with a whitish line continued backwards to the glands back of the corner of the mouth; a conspicuous, sharply defined, longitudinal, oval blackish spot on the underside of the forearm: underside uniform pale.



FIGS. 53-57.—*BUFO RADDEI*. $1\frac{1}{2} \times$ NAT. SIZE. 53, TOP OF HEAD; 54, SIDE OF HEAD; 55, OPEN MOUTH; 56, UNDERSIDE OF HAND; 57, UNDERSIDE OF FOOT. No. 21394, U.S.N.M.

Dimensions.

	<i>mm.</i>
Tip of snout to vent.....	55
Tip of snout to anterior border of tympanum.....	14
Nostril to tympanum	11
Internarial width.....	3.5
Interorbital width.....	4
Diameter of tympanum.....	3
Distance of tympanum from eye.....	2
Width of head.....	20
Fore leg.....	32
Hind leg from vent to tip of longest toe.....	66
Tibia.....	19
Hind foot from base of inner metatarsal tubercle to tip of longest toe	22
Length of parotoid.....	12
Width of parotoid.....	6

A *young* specimen, 27 mm. long, from Lake Khanka, Ussuri (U.S.N.M. No. 14343) also received from the St. Petersburg Academy, has the interorbital space much narrower, scarcely more than two-thirds the width of the upper eyelid, and the tympanum is much smaller, its diameter scarcely more than one-third the diameter of the eye. The color is very bright and the pattern very distinct; a brown line runs from nostril over canthus rostralis and the extreme outer edge of the upper eyelid, bordered above by a pale olive line which on the tip of the snout joins a similar one occupying the entire median line of head and body above; a similarly pale line crosses the middle of the eyelids, and another one the occiput from tympanum to tympanum; parotoids pale olive with an irregular brownish figure in the outer posterior portion; all the larger glands on the back pale olive with a brownish outer edge forming ocelli; underside uniform pale with a few dusky spots on the belly.

The *females* lack the subgular sac and the nuptial asperities on the fingers. These excrescences persist in the male some time after the end of the breeding season.

Habitat.—This species which seems to occur from the eastern part of the desert Gobi, Alashan, and Ordos in Mongolia to Peking and Chefu in the south and north to Dauria, the valley of the Amur and the Ussuri country, is regarded by Boulenger as representing the European *Bufo calamita* in eastern Asia, and as a form intermediate between the latter and *Bufo viridis*. In the Hamburg Museum there is a specimen collected by Dieckmann in 1894 at Khabarovka, Amurland, and another from the same locality is in the Basel Museum. Stone has recorded it from a lake near Lama-mio and various other localities in eastern Mongolia.

List of specimens of Bufo raddi.

U.S. N.M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
14343	Young ^a	Lake Khanka, Ussuri.....		Acad. Sci. St. Petersburg.
21394	Male ^b	Zagan, East Siberia.....		Do.

^a P. 72.

^b Description, p. 70; figs. 53-57.


BUFO MELANOSTICTUS^a Schneider.

1799. *Bufo melanostictus* SCHNEIDER, Hist. Amph., I, p. 216 (India orientalis).—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 306 (South China; East Indies); Fauna Brit. India, Rept., 1890, p. 505, fig. p. 506.—BOETTGER, Offenbach, Ver. Naturk. 24-25 Ber., 1885, p. 131 (Canton; Prov. Quangtung, China).—FLOWER, Proc. Zool. Soc. London, 1896, p. 911, pl. XLIV, fig. 3 (tadpole).—STEFNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 216 (*meranostictus* err. typ.) (Taipa and Gillan, Formosa.)

^a From μέλας, black; στικτός, spotted.

The numerous synonyms based upon Indian and Malayan specimens are omitted here.

Description (figs. 58-61).—*Adult female*; U.S.N.M. No. 36498; Sic. Coll. Mus., Tokyo, No. 26a; Taipa, Formosa; September, 1896; Tsunasuke Tada, collector. Head with strongly developed bony crests, involving the upper lip, a rostral ridge on end of snout from lip to between nostrils, there bifurcating and continuing on canthus rostralis, along supraorbital border to above the tympanum, forming there a knob-like prominence and sending a branch downward along the anterior border of the tympanum; a slight parietal spur indicated, also a short anteorbital spur or crest; nostrils nearer the tip of the



FIGS. 58-61.—*BUFO MELANOSTICTUS*. NAT. SIZE. 58, TOP OF HEAD; 59, SIDE OF HEAD; 60, UNDERSIDE OF HAND; 61, UNDERSIDE OF FOOT. No. 36498, U.S.N.M.

snout than the eye; interorbital space deeply concave, much wider than upper eyelid, which is strongly tubercular and with thickened glandular edges; tympanum vertical, oval or almost pear-shaped, separated from the orbit by a very narrow space, its longest diameter about three-fourths the diameter of the eye; parotoids large, swollen, kidney-shaped; first finger reaches considerably beyond second which does not quite reach as far as fourth; an elongate inner and a much larger outer palmar tubercle; subarticular tubercles single, prominent; palm, underside, and edges of fingers with numerous pointed tubercles, which, like all the other digital tubercles and the

horny tip of the fingers, are blackish brown; heels just meeting when hind legs are placed at right angles to axis of body; metatarsal tubercles reaching posterior corner of eye when hind legs are placed forward along the sides of the body; toes less than one-half webbed, the webs being greatly excised and their edges sharply denticulate; inner metatarsal tubercle prominent, oval, shorter than first toe, outer one large, rounded, less prominent; the single subarticular tubercles, plantar tubercular asperities, and tip of toes like those on the fore foot; skin strongly tubercular, the tubercles on the back larger, more distant, glandular, rounded, those on sides and limbs more prominent, almost conical, all with a horny dark-brown tip, which on sides and limbs becomes spinous. Color (in alcohol), clay-colored above, more pure pale ocher-yellow underneath, all the crests and nearly all the tips of the tubercles dark brown, as are also the elevated fine vermiculations on the surface of the parotoids; indistinct dusky spots and coarse vermiculations on limbs, sides, throat, and breast.

Dimensions.

	<i>mm.</i>
Tip of snout to vent.....	84
Tip of snout to anterior border of tympanum.....	20
Nostril to tympanum.....	16
Internarial width.....	5
Interorbital width.....	8.5
Longest diameter of tympanum.....	5.5
Width of head.....	31
Fore leg.....	50
Hind leg from vent to tip of longest toe.....	99
Tibia.....	26
Hind foot from base of inner metatarsal tubercle to tip of longest toe.....	30
Length of parotoid.....	20
Width of parotoid.....	8

The *adult male* appears to be smaller, U.S.N.M. No. 36499, measuring only 61 mm. from snout to vent, but in proportions and otherwise similar to the female, except that the webs between the toes are considerably smaller, and the outer metatarsal tubercle is smaller than the inner. It has also distinct traces of nuptial asperities on top of the first finger, probably only remnants from the breeding season, during which such excrescences are present on first and second fingers. A small internal subgular sac, entrance slits to which are seen on each side of the tongue. The dark color markings are somewhat more distinct.

A *halfgrown* specimen (Sci. Coll. No. 26*e*) measuring 46 mm. is very similar, but the skin is much less rough except on hands and feet; the interorbital space is flat, but all the cranial crests are plainly developed except the preauricular; the tips of the fingers are pale, and the dusky brown spots and vermiculations on throat, chest, and anterior portion of abdomen very strong and well defined; lips similarly blotched with dark brown.

Habitat.—Widely distributed through India, Ceylon, Burma, the Malayan peninsula and archipelago to southern China and Formosa. Boulenger states that in the Sikhim Himalayas it ascends to about 10,000 feet and in the Nilgiris to 7,000 feet.

In Formosa it seems to be common, as Mr. Tada collected many specimens at Taipa and Giilan during September and October, 1896. There are two specimens in the British Museum from Taiwan fu, collected by Holst.

List of specimens of Bufo melanostictus.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
Sci. Coll. Tokyo.....	2ic	Male.....	Taipa, Formosa.....	Sept.—, 1896	T. Tada.
Do.....	2id	Female.....	do.....	do.....	Do.
Do.....	2ie	Halfgr. a	do.....	do.....	Do.
Do.....	25a	Giilan, Formosa.....	Oct., 1896	Do.
Do.....	25b	do.....	do.....	Do.
Do.....	25c	do.....	do.....	Do.
Do.....	25d	do.....	do.....	Do.
U. S. N. M.	34329	Taipa, Formosa.....	Mar., 1903	A. Owston.
Do.....	36498	Female b	do.....	Sept., 1896	T. Tada.
Do.....	36499	Male a	do.....	do.....	Do.
Do.....	36500	Giilan, Formosa.....	Oct., 1896	Do.

aDescription, p. 74.

bDescription, p. 73, figs. 58-61.

Family HYLIDÆ.

The tree-toads are characterized by the presence of teeth in the upper jaw, or in both upper and lower jaws, combined with an arciferous shoulder-girdle and dilated sacral lateral processes; bone of last toe-joint claw-shaped.

A large family of wide distribution, though not occurring in the Ethiopian region. The greatest number of species is found in tropical America and in Australia, but only a few in Asia and Europe. Most of the forms are arboreal with the tip of the digits enlarged into a disk, which is more or less sticky. Some are modified to such an extent for life among the trees that they lay their eggs in water collected in holes in trunks and branches of trees.

Only one genus is represented in eastern Asia.

Genus HYLÆ^a Laurenti.

1768. *Hyla* LAURENTI, Synops. Rept., p. 32 (type, *H. viridis*).

1799. *Calamita* SCHNEIDER, Hist. Amph., I, p. 151, comp. p. 174 (type, *C. arboreus*).

^a Not derived from the Greek word ὕλη, wood-land, copse, as commonly stated, but from the vocative of Hylas, in Greek mythology the favorite of Hercules, who lost him in Bithynia, the crying of *hyla, hyla* being part of the religious ceremonies instituted in his honor. The croaking of the tree-toad suggested to Laurenti the fanciful idea of its being Hyla's priest, and thus meriting his name, "hæc quasi Hylæ sacerdos nomen ejusdem merita est."

1814. *Hylaria* RAFINESQUE, Specchio Sci. (Palermo), II, fasc. 7 (substitute for *Hyla*).
1825. *Boana* GRAY, Ann. Philos. (n. s.), X, p. 214 (type, *Rana boans*).
1826. *Calamita* FITZINGER, Neue Classif. Rept., p. 38 (type, *Rana caerulea* White).
1827. *Rhacophorus* "Kuhl" SCHLEGEL, Isis, 1827, p. 294 (type, *H. palmata*; not of Tschudi, 1838).
1830. *Calamites* WAGLER, Syst. Amph., p. 200 (emendation of Fitzinger's *Calamita*).
1830. *Auletris* WAGLER, Syst. Amph., p. 201 (type, *H. boans*).
1830. *Hyas* WAGLER, Syst. Amph., p. 201 (type, *H. arborea*; not of Leach, 1815).
1830. *Scinar* WAGLER, Syst. Amph., p. 201 (type, *H. aurata*).
1830. *Dendrohyas* WAGLER, Syst. Amph., p. 342 (substitute name for *Hyas*, preoccupied).
1843. *Lobipes* FITZINGER, Syst. Rept., p. 80 (type, *H. palmata*).
1843. *Dryophytes* FITZINGER, Syst. Rept., p. 31 (type, *H. versicolor*).
1885. *Epedaphus* COPE, Proc. Amer. Philos. Soc., 1885, p. 383 (type, *H. gratiosa*).
1899. *Hyliola* MOCQUARD, Nouv. Arch. Mus. Paris (4), I, p. 337 (type, *H. regilla*).

The list of synonyms is considerably larger than the above, which includes mostly such names only as are based upon species occurring within the faunal area here treated of, or constitute corrections or additions to the synonymy as commonly given.

The tropical genus *Hyla*, as now understood, is American and Australian in its distribution, and embraces considerably over 200 species, less than 3 per cent of which occur in the Old World. The species, or forms, which are found in the mainland of Eurasia, viz, *H. annectens* and *H. simplex*, with the ones treated of in this work, constitute a closely related group ^a extending from Great Britain to Japan, south to northeastern India and Annam.

The forms which are to occupy us in this connection may be distinguished as follows:

KEY TO THE JAPANESE, KOREAN, AND FORMOSAN SPECIES OF HYLA.

a¹ No deep black spots on sides or legs.

b¹ Digital disks equaling inner metatarsal tubercle.

c¹ A distinct blackish stripe between nostril and eye.

H. arborea japonica, p. 76

c² No blackish stripe between nostril and eye. *H. arborea immaculata*, p. 82

b² Digital disks much smaller than inner metatarsal tubercle. . . . *H. stephensi*, p. 84

a² Sides of body and posterior aspect of hind limbs with deep black spots.

H. chinensis, p. 86

HYLA ARBOREA ^b **JAPONICA** ^c **Guenther.**

Plate IX, figs. 1-3.

1838. *Hyla arborea* SCHLEGEL, Fauna Jap. Rept., pp. 112, 136, 139; Saur. et Batr., pl. III, figs. 5^d, 6 (not of Linnaeus) (Japan).—HILGENDORF, Sitz. Ber. Berlin Ges. Naturf. Fr., 1880, no. 8, p. 120 (Hondo; Yezo).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, 1891, p. 239 (Yezo).

^a In turn related to *Hyla crimia* of Mexico.

^b Signifying arboreal.—*Rana arborea* LINNÆUS, Syst. Nat., 10 ed., I, 1758, p. 213.

^c Signifying Japanese.

^d Reproduced in this work on Plate IX, fig. 3.



1



2



3



4

FIGS. 1-3, *HYLA ARBOREA JAPONICA*; FIG. 4, *H. CHINENSIS*. (FROM BOULENGER, SCHLEGEL, AND GUENTHER.)

FOR EXPLANATION OF PLATE SEE PAGE 554.

1858. *Hyla viridis* var. *japonica* GUENTHER, Cat. Batr. Sal. Brit. Mus., p. 81.
1858. *Hyla arborea* var. *japonica* GUENTHER, Cat. Batr. Sal. Brit. Mus., p. 109 (type-locality, Japan; types in Brit. Mus.).—BOETTGER, Offenbach. Ver. Naturk., 17-18 Ber., 1878, p. 8 (Japan); Kat. Batr. Mus. Senckenberg., 1892, p. 43 (Hakone; Goto Isl.); Ber. Senckenberg. Naturf. Ges., 1894, p. 141.—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 381 (Japan); Proc. Zool. Soc. London, 1887, p. 578, pl. LI, fig. 2 (Port Hamilton, Korea).—BEDRIAGA, Bull. Soc. Natural. Moscou (n. s.), III, no. 3, 1889, pp. 475, 487 (Tokyo; Yezo).—OKADA, Cat. Vert. Jap., 1891, p. 66 (Tokyo).—SCLATER, Batr. Ind. Mus., 1892, p. 32.
1860. *Hyla viridis*? HALLOWELL, Proc. Phila. Acad., 1860, p. 500 (Simoda).
1878. *Hyla arborea* var. *meridionalis* BOETTGER, Offenbach. Ver. Naturk. 17-18 Ber., p. 8 (Japan) (not of 1874).
1879. *Hyla japonica* CAMERANO, Atti Accad. Torino, XIV, Pt. 5, Apr. 1879, p. 895 (Tokyo).
1887. *Hyla arborea* var. *savignyi* BOULENGER, Ann. Mag. Nat. Hist. (5), XIX, 1887, p. 67 (Gensan, Korea) (not of Audouin); Taill. Batr. Europe, Pt. 2, 1898, p. 250 (part), pl. xv, fig. 5 (Tsu-shima), fig. 6 (Daibutsu).^a
1905. *Hyla stepheni* NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, p. 401 (part: Yezo) (not of Boulenger?).

Schlegel's fig. 5 on Plate III^b of the Saurii and Batrachii of Fauna Japonica is usually referred to *Polypedates schlegelii*, but I think erroneously. The webbing between the digits and the shape of the snout, as well as the coloration, indicate *Hyla arborea*. Moreover, an examination of the specimen in Leiden Museum shows that this view is correct.

In his catalogue of the tailless batrachians in the British Museum (1882) Boulenger recognized four varieties of *Hyla arborea*, of which *H. arborea savignyi* was credited to various islands in the Mediterranean, parts of western Asia, and to the island of Hainan, South China (a single half-grown specimen), while *H. arborea japonica* was represented as being confined to Japan. In 1887 (July) he recorded^c a specimen of the latter from Port Hamilton, a small island at the south end of Korea, opposite Kiusiu, found together with another specimen which he described as *Hyla stepheni*. Shortly before (January, 1887) he had referred a specimen from Gensan, Korea, to *H. arborea* var. *savignyi*,^d a reference, by the way, which appears to have escaped Doctor von Bedriaga.^e While the latter author, as well as Boettger^f still recognize *H. arborea japonica*, Boulenger in his latest important work^g after the study of a large series of specimens (34 from the Mediterranean and western Asia as far as the Persian Gulf and 18 from Japan, Korea, and China)

^a Reproduced in this work on Plate IX, figs. 2, 1.

^b Reproduced in this work on Plate IX, fig. 3.

^c Proc. Zool. Soc. London, 1887, p. 578.


^d Ann. Mag. Nat. Hist. (5), XIX, Jan. 1887, p. 67.

^e Bull. Soc. Nat. Moscou, (n. s.) III, 1889.

^f Kat. Batr. Mus. Senckenb., 1892, and Ber. Senckenb. Naturf. Ges., 1894.

^g Taill. Batr. Europe, II, 1897.

pronounce them all *H. arborea savignyi*. From the text (pp. 250-251) it is evident that his reason for this is that there are spotted specimens found in the western series as well as unspotted ones among those from eastern Asia. It is quite possible that the unspotted specimens may be somewhat difficult to distinguish, but in the spotted ones the angular shoulder spots of the Japanese and Korean specimens seem to be almost diagnostic. In our plain-colored specimens, moreover, I can always detect dusky bars across the tarsus, and I would also call attention to the dark spot on the upper lip under the anterior portion of the eye which is found in many of the eastern specimens. Under these circumstances I think it preferable to follow Boettger in maintaining the old status of *H. arborea japonica* as a distinguishable form.^a



FIGS. 62-67.—*Hyla arborea japonica*. $1\frac{1}{2} \times$ NAT. SIZE. 62, TOP OF HEAD; 63, SIDE OF HEAD; 64, OPEN MOUTH; 65, UNDERSIDE OF HAND; 66, TOP OF FINGER, SIDE VIEW; 67, UNDERSIDE OF FOOT. No. 23542, U.S.N.M.

Description (figs. 62-67).—*Adult female*; U.S.N.M., No. 31926; Kochi, Province of Tosa, Shikoku; May 11, 1903; Dr. Hugh M. Smith, collector. Vomerine teeth in two nearly straight, short series between the choanæ; tongue large, broadly oval, somewhat truncate behind, with a slight median incision; nostrils somewhat nearer the tip of the snout than the eye, their distance from each other shorter than their distance from the labial edge, and still shorter than the inter-

^a What is *Hyla cyanea* Hallowell. Proc. Phil. Acad., 1860, p. 494, from "Loo Choo"? All he says is: "*Hyla cyanea* Daudin (young). One specimen." There is great confusion in this paper of Hallowell's (for instance, *Polypedates viridis* and *burgerii*, p. 501, under Japan instead of Loo-Choo; *Ophthalmidium tenue*, p. 497, from Hongkong, China, under Japan), and it is quite possible that this reference to a young *Hyla cyanea* is only a duplication of the Australian record, p. 490: "*Hyla cyanea* Daud. One specimen. Sidney (young). Taken Dec., 1853," which got misplaced. I may add that Boulenger refers it with a query to *Rhacophorus viridis* (Proc. Zool. Soc. London, 1887, p. 149).

orbital width, which is considerably greater than the upper eyelid; tympanum nearly circular, its diameter about one-half that of the eye, its distance from the latter greater than from the corner of mouth; fingers distinctly webbed at base, first shortest, fourth longer than second, third longest; disks of fingers slightly larger than those of toes, nearly as large as tympanum; hind feet rather long, tarso-metatarsal joint reaching considerably beyond tip of snout and heel joint reaching to middle of eye when legs are carried forward along the side of the body; heels just touch when folded legs are placed vertical to the axis of the body; toes webbed slightly more than one-half; third toe as long as fifth; inner metatarsal tubercle oval, prominent, about the size of the disk of the longest toe; no outer metatarsal tubercle; sub-articular tubercles under fingers and toes prominent; skin above smooth; below granular, except on inner side of tibia; a strong glandular fold from eye, curving downward behind the tympanum to the insertion of the fore leg; a strong dermal fold across the chest from axilla to axilla. Color (in alcohol) about smoke gray (in life bright green), with dusky markings as follows: A Y-shaped rostral stripe on the end of the snout, connected below with an undulating blackish line, forming the edge of the upper jaw, the diverging upper branches reaching the nostrils and continuing behind these through the eye and over the tympanum as a dark stripe, edged above with whitish; on the tympanum it widens so as to embrace the whole of this organ, and continues backward over the shoulder along the flanks as an irregular broad band, which in the groin is replaced by coarse vermiculation of blackish and white; a dusky spot between lower anterior portion of eye to edge of lip; a pair of dusky spots, one on each upper eyelid, converging, though not meeting, backward; a large angular, L-shaped mark on each shoulder, and a smaller, more irregular spot on each side of the lower back, descending toward the groin, but not connected with the dusky band of the flanks; on the coccyx several smaller irregular blotches; a spot in front of the upper end of humerus, a narrow dusky line along the posterior outline of the arm; posterior aspect of femur brownish gray with distinct rounded white spots; upper aspect of fore leg and hind legs, including tarsus, with distinct dusky crossbars; lower lip whitish, with a series of dark spots; underside whitish, with numerous small dusky spots on throat.


Dimensions.

	<i>mm.</i>
Tip of snout to vent.....	37
Tip of snout to anterior border of tympanum.....	9.5
Nostril to anterior border of tympanum.....	8
Internarial width.....	3
Interorbital width.....	4
Diameter of tympanum.....	2

	mm.
Width of head.....	12
Fore leg.....	23
Hind leg from vent to tip of longest toe.....	57
Tibia.....	16.5
Foot from base of inner metatarsal tubercle to tip of longest toe....	18
Inner metatarsal tubercle.....	2

The *adult male* has a large external, dark-pigmented, vocal sac under the throat, folding up with longitudinal folds when not inflated. The males of U.S.N.M. No. 7428, which were collected by Dr. Wm. Stimpson, at Simoda, in May, 1855, consequently during the breeding season, have a large and distinct pad of nuptial asperities on the upper side of the base of the first finger, but they are not distinguished from the adjacent skin by any different coloration. The color of these specimens, which have the back unspotted, is described from life by Stimpson as being "bright light green above; a golden band along each side of the head, including the eye; sides minutely sprinkled with copper color; belly white."

Variation.—In the series before me the structural and proportional



FIGS. 68-70.—*Hyla arborea japonica*. $1\frac{1}{2} \times$ NAT. SIZE. 68, TOP OF HEAD; 69, SIDE OF HEAD; 70, UNDERSIDE OF FOOT. No. 21211, U.S.N.M.

characters, such as the size and shape of the vomerine series of teeth, the size of tympanum, digital disks, and metatarsal tubercle, vary within certain limits. The length of the foot is found to vary considerably in a few specimens, apparently without being correlated with any other difference. It is in the coloration, however, that we meet with the greatest amount of variation. Not only are many of the specimens entirely uniform above, but the lateral stripe is scarcely indicated in a large number, and the mark on the tip of the snout is absent in most.

The dusky spot beneath the eye is present in a majority, and the crossbars on the tarsus are visible in all the specimens which are not so faded from age as to have lost nearly all trace of markings. In the specimens with dorsal spots the angular shoulder mark is usually present, but in No. 23542, at least, there is a longitudinal stripe on each side of the back, caused by a fusion of the two dorsal spots and the obliteration of the transverse branch of the anterior mark.

Habitat.—The tree-toad, a name probably preferable to that of tree-frog, which may properly be restricted to the genus *Polypedates* (or *Rhacophorus* of many authors), occurs all over Japan proper in suitable stations. It is hardly necessary to enumerate specific locali-

ties, though the following are represented by specimens in various museums: In Kiusiu: Yamagawa and Miyazaki (U.S.N.M.), Mount Onsen in Shimabara (Brit. Mus.); in Shikoku: Kochi (U.S.N.M.); in Hondo: Nara, Koriyama, Simoda, Mount Fuji, and Yokohama (U.S.N.M.), Kobe, Kiga, Daibutsu (Brit. Mus.), Setsu (Hamburg Mus.), Hakone (Senckenberg Mus.); in Yezo: Hakodate and Sapporo (U.S.N.M.). There are also in British Museum specimens from Tsu-shima collected by Holst, and in Museum Senckenbergianum one specimen from Goto Island, near Nagasaki.

There is no record of the species in any of the Riu Kiu Islands or Formosa.

Outside of Japan it occurs in Korea, whence specimens are in the United States National Museum (Fusan) and in the British Museum (Gensan and Seoul); the Hamburg Museum also has specimens from southern Korea, collected by Doctor Warburg. One specimen was brought from Port Hamilton by Mr. G. Stephen, together with the type of *Hyla stephensi*.

The status of the Chinese specimens which have been referred to this form is too uncertain to detain us here. It may be sufficient to refer to what is said under the heading of *Hyla arborea immaculata*, with the additional remark that the Hainan specimen belongs to the recently described *Hyla simplex*^a as suggested by Doctor Boettger.

List of specimens of Hyla arborea japonica.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
7428a	Male a.	Simoda, Izu, Hondo	May, 1855	Wm. Stimpson.
7428b	do.	do.	do.	Do.
7428c	do.	do.	do.	Do.
7428d	do.	do.	do.	Do.
7428e	Female	do.	do.	Do.
7428f	Young.	Hakodate, Yezo.		Do.
7428g	do.	do.		Do.
11345	do.	Japan	1878	E. S. Morse.
11346	do.	do.	1878	Do.
17511	Female	Fusan, Korea	1885	P. L. Jouy.
21211	do. b	(Seoul?), Korea	1883	Do.
21212	Young.	do.	1883	Do.
23441	do.	Sapporo, Yezo.	1896	Nozawa.
23442	do.	do.	1896	Do.
23443	do.	do.	1896	Do.
23444	do.	do.	1896	Do.
23445	do.	do.	1896	Do.
23542	Female c	Yokohama, Hondo	Sept., 1896	L. Stejneger.
23588	do.	do.	do.	Do.
23607	do.	Onogami, Gunba	do.	I. Ijima.
23608	Halfgrown	do.	do.	Do.

^a Description, p. 80.

^b Figs. 68-70; see also p. 85.

^c Figs. 62-67; see also p. 85.

^a *Hyla chinensis* var. *simplex* Boettger, Ber. Senckenberg. Naturf. Ges., 1901, Abh., p. 53.

List of specimens of *Hyla arborea japonica*—Continued.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
30742	Female.....	Miyazaki, Hiuga, Kiusiu.....		C. A. Clark.
30743	Male.....	do.....		Do.
31803	Halfgrown ..	Nara, Yamato, Hondo	Apr., 1903	H. M. Smith.
31805do.....	Koriyama, Yamato, Hondo	Apr. 26, 1903	Do.
31806do.....	do.....	do.....	Do.
31807do.....	do.....	do.....	Do.
31825	Male.....	Yamagawa, Satsuma, Kiusiu.....	June 14, 1903	Do.
31826	Young.....	do.....	do.....	Do.
31827do.....	do.....	do.....	Do.
31913	Male.....	Kochi, Toza, Shikoku.....	May 11, 1903	Do.
31914	Female.....	do.....	do.....	Do.
31915do.....	do.....	do.....	Do.
31916	Male.....	do.....	do.....	Do.
31917	Female.....	do.....	do.....	Do.
31918	Male.....	do.....	do.....	Do.
31919do.....	do.....	do.....	Do.
31920do.....	do.....	do.....	Do.
31921do.....	do.....	do.....	Do.
31926	Female ^a	do.....	do.....	Do.
34374	Male.....	Mount Fuji, Hondo	Summer, 1898	A. Owston.
34375do.....	do.....	do.....	Do.
34376	Female.....	do.....	do.....	Do.
34377do.....	do.....	do.....	Do.
34378	Male.....	do.....	do.....	Do.
34379	Female.....	do.....	do.....	Do.
34380do.....	do.....	do.....	Do.
34381	Male.....	do.....	do.....	Do.
34382	Female.....	do.....	do.....	Do.
34383do.....	do.....	do.....	Do.
34384do.....	do.....	do.....	Do.
34385do.....	do.....	do.....	Do.
34386do.....	do.....	do.....	Do.
34387do.....	do.....	do.....	Do.
34388do.....	do.....	do.....	Do.
34389do.....	do.....	do.....	Do.

^a Description, p. 78.[HYLA ARBOREA IMMACULATA^a Boettger.]

1888. *Hyla chinensis* var. *immaculata* BOETTGER, Ber. Senckenberg. Naturf. Ges., 1888, Abh., p. 189 (type-locality, Shanghai; type in Mus. Senckenberg.; Schmacker, collector).—STONE, Proc. Phila. Acad., 1899, p. 184 (Tore River, eastern Mongolia).

1892. *Hyla arborea* var. *immaculata* BOETTGER, Kat. Batr. Mus. Senckenberg., p. 43 (Shanghai; Lueshan Mts., near Kiukiang); Ber. Senckenberg. Naturf. Ges., 1894, p. 140 (Lueshan).

This form which Boettger originally referred to *Hyla chinensis* as a variety he has since associated with *H. arborea*. According to his revised view of 1894^b it is most nearly related to *H. arborea japonica*,

^a Signifying Unspotted.^b Ber. Senckenberg. Naturf. Ges., 1894, p. 141.

from which "it may be easily separated by the following characters: No web between the outer fingers; toes one-third webbed; no dark loreal stripe," while *H. arborea japonica* has "one-fourth web between third and fourth finger; toes one-half webbed, and an always distinct dark loreal stripe. In the proportion between the length of the tibia to that of the femur there may also be a slight difference. In var. *immaculata* the tibia appears to be somewhat longer than the femur; in var. *japonica*, on the contrary, the femur is longer than the tibia."

I have not seen any specimens of this form from the type locality and therefore can not express any opinion as to its validity, but if the three specimens from Mongolia, which Mr. Witmer Stone somewhat doubtfully referred to this form, really belong to it, it would appear that the characters assigned to it are somewhat unstable and unreliable.

Thanks to the kindness of the authorities of the Philadelphia Academy, I was recently permitted to examine the three specimens alluded to above, namely, Nos. 4943-4945, collected by Doctors Smith and Furness at Tore River, Mongolia.

No. 4945, which is adult, differs from a German specimen, with which I could compare it, by being entirely immaculate, with no trace of canthal or postocular streaks or dusky crossbars on the legs. The foot is somewhat shorter and the webs slightly larger; the heels do not meet by a space of 2 mm. when the folded hind legs are placed vertical to the axis of the body, and they do not reach the axilla when the leg is extended along the side of the body.

No. 4944, also adult, has relatively longer hind legs, as the heel reaches the posterior border of the eye. A canthal and postocular line can be distinctly traced, also some spots on the upper lip, and apparently faint cross markings on the legs.

No. 4943, which is quite young, is nearly unspotted like No. 4945, but with trace of a subocular spot on one side and of cross markings on legs.

The following *measurements* were taken of the two adults to show the difference in the relative length of the hind legs:

	No. 4945.	No. 4944.
	mm.	mm.
Snout to vent.....	40	39
Vent to tip of longest toe.....	49	59
Heel to tip of longest toe.....	25	27

The status of the tree-toad of northern China is consequently still doubtful. It seems to be fairly distinct from the Japanese form, however, in the *relatively* immaculate coloration.

Habitat.—Boettger originally recorded this form from Shanghai and later from the Lueshan Mountains near Kiukiang on the Yangtse-kiang, where it occurs together with *Hyla chinensis*. Three Mongolian specimens, from the Tore River, have been referred to this form doubtfully, as detailed above.

List of specimens of Hyla immaculata.

Phila. Acad. No.	Age.	Locality.	When col- lected.	By whom collected.
4943	Young ♂	Tore River, Mongolia		Dr. Smith and Furness.
4944	Adult ♂	do		Do.
4945	do, ♀	do		Do.

* P. 83.

HYLA STEPHENI^a Boulenger.

1887. *Hyla stephensi* BOULENGER. Proc. Zool. Soc. London, 1887, p. 579, pl. LI, fig. 1 (type-locality, Port Hamilton, Korea); Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 142 (Ussuri River); Proc. Zool. Soc. London, 1890, p. 326.—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, No. 1, 1905 (part: Ussuri, etc.).

This species was described by Boulenger from a specimen collected at Port Hamilton, together with a specimen of true *H. arborea japonica*. It is professedly nearly allied to *H. arborea*, and no comparison with the latter accompanies the description.

Later on (1890) he records two specimens from the Ussuri River of which he gives the measurements and says that the species "is easily distinguished from *H. arborea* by the much larger and more prominent metatarsal tubercle." The length of the inner toe is given as 4 mm. and that of the metatarsal tubercle as 2.5 mm.

Through the kindness of the authorities of the British Museum, I was able to examine these specimens during my visit in 1898.

The type (Brit. Mus. 89.11.8.5) is in very poor condition (dried up) and nothing can be made of it. The two other specimens, however, referred by Boulenger to this species (Brit. Mus. 89.12.16.197.7), collected by Doerries at the Middle Ussuri in 1886 (Fischer collection Nos. 754-5), are in very good condition and, compared with the Hondo specimens of *H. japonica* then in the British Museum, show a difference in the relative size of disks and metatarsal tubercle as follows:

Digital disks much smaller than inner metatarsal tubercle *H. stephensi*
 Digital disks equaling metatarsal tubercle *H. japonica*

In the latter, moreover, the ventral granulation is considerably finer.

Whether the Ussuri specimens are identical with the Port Hamilton type is now hard to determine, but we will have to accept Boulenger's identification as correct.

^a For Mr. George Stephen, R. N., who collected the type.

A single specimen in the United States National Museum (No. 21211, Figs. 68-70), collected by Jouy in Korea in 1883, probably in Seoul, has a somewhat larger inner metatarsal tubercle than the other specimens from Korea and Japan, and while the snout is possibly a trifle shorter, the hind legs are not short, the tibio-tarsal articulation reaching the front of the eye; finally, the ventral granulation is exactly as in the typical *Hyla japonica*. At first I thought that the hind foot of this individual might afford a good character, it being considerably larger than in most of my Japanese specimens, but one collected by myself at Yokohama, in 1896, is so nearly like it that I do not venture to separate them.

Altogether I am not assured of the distinctness of *H. stephensi*, my faith in it having been considerably shaken by the last-mentioned specimen (U.S.N.M. No. 23542, figs. 62-67). However, having been favorably impressed by the examination of the Ussuri specimens, and not now having an opportunity to reexamine them in connection with my own material, I do not feel warranted in disturbing the status given them by Boulenger.

Nikolski, in 1905, has referred the tree-toads in the St. Petersburg Museum from Yezo, collected by Albrecht in 1865 and by Maximovitch, to *H. stephensi*, apparently as distinct from the tree-toads of southern Japan. I have not been able to accept this view, as I have been unable to discover any tangible difference between the *Hylas* on both sides of Tsugaru Strait, but Nikolski's action goes to show how unsatisfactory the whole question is as yet. Much more material will be necessary before it can be settled.

There being no undoubted specimen of this form in our museum, I reproduce herewith Boulenger's original description and his measurements of the two Ussuri specimens.

Original description of type-specimen.—Brit. Mus. No. 89.11.8.5; Port Hamilton, Korea; G. Stephen, collector.—Near *H. arborea*. Tongue subcircular, slightly nicked, and free behind; vomerine teeth in two slightly oblique series between the choanæ; head a little larger than that of *H. arborea*, broader than long; snout rounded, shorter than the diameter of the orbit; canthus rostralis distinct; loreal region slightly concave; interorbital space as broad as the upper eyelid; tympanum distinct, half the diameter of the eye; fingers with a short web at the base; no projecting rudiment of pollex; toes hardly two-thirds webbed; disks a little smaller than the tympanum; subarticular tubercles moderate; inner metatarsal tubercle large and very prominent, oval, about two-thirds the length of the inner toe; a fold along the inner edge of the tarsus; the tibio-tarsal articulation reaches the tympanum; tibia not half the length of the head and body; skin smooth above; lower surfaces, throat of male included, granular. Color varying from grayish or brownish to green, with

darker, blackish-edged permanent symmetrical insuliform spots on the body and cross bands on the limbs; a dark brown streak from nostril to eye; lower parts whitish; lower belly and limbs carneous; sides of thighs carneous; iris dark bronzy brown. Male with an external subgular vocal sac.

From snout to vent 35 mm. ^a

Dimensions.—British Museum, No. 89.12.16.197.7; middle Ussuri; Doerries, collector.

	Male.	Female.
	<i>mm.</i>	<i>mm.</i>
From snout to vent.....	35	40
Length of head.....	12	13
Width of head.....	13	15
Diameter of the tympanum.....	2	3
Fore limb.....	20	23
Hind limb.....	51	57
Tibia.....	15	17
Inner toe.....	4	4
Inner metatarsal tubercle.....	2.5	2.5

^a Boulenger, Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 142.

Habitat.—The only specimens until recently recorded as belonging to this form are the ones mentioned above, one from Port Hamilton, off the south end of Korea, and two from the middle Ussuri. In the Hamburg Museum there are specimens from Khabarovka (No. 863) as well as from Vladivostok (No. 864), collected by Dieckmann, which are there referred to this species. Nikolski, in 1905, refers tree-toads from as far west as Nertshinsk and the river Shilka, as well as the Yezo specimens in the St. Petersburg Museum to this species.

List of specimens of Hyla stepheni.

Brit. Mus. No.	Sex.	Locality.	When collected.	By whom collected.
89. 11. 8. 5.....	(a)	Port Hamilton, Korea.....		G. Stephen.
89. 12. 16. 197. 7	Male ^b	Middle Ussuri.....		Doerries.
.....	Female ^b	do.....		Do.

^a Type, description, p. 85.

^b P. 86.

HYLA CHINENSIS ^b Guenther.

Plate IX, fig. 4.

1858. *Hyla arborea* var. *chinensis* GUENTHER, Cat. Batr. Sal. Brit. Mus., p. 108, pl. IX, fig. C^c (type-locality, China; types in Brit. Mus.).

^a Boulenger, Proc. Zool. Soc., 1887, p. 579.

^b Signifying Chinese.

^c Reproduced in this work on Plate IX.

1864. *Hyla chinensis* GUENTHER, Rept. Brit. India, p. 436 (southern China and Formosa).—STEINDACHNER, Novara Exped., Zool., I, Amph., 1867, p. 59 (Amoy).—CAMERANO, Atti Accad. Sci., Torino, XIV, Pt. 5, Apr. 1879, p. 894 (China).—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 381 (China; Formosa).—BOETTGER, Kat. Batr. Mus. Senckenberg., 1892, p. 43 (Dalanshan and Lueshan Mts., China); Ber. Senckenberg. Naturf. Ges., 1894, p. 141 (Lueshan Mts. near Kiukiang); p. 149 (Dalanshan and Chinhai, near Ningpo).

This species is closely allied to *Hyla arborea*, but the vomerine teeth are more posterior. The coloration is also peculiar and easily recognizable, the color above being uniform and immaculate bright green, the sides beautifully sulphur yellow (according to Boettger), white in alcohol, with round spots of a deep black, and extensive markings of the same color on the posterior aspect of the thighs.

Habitat.—This species occurs in southern and eastern China at least as far north as Shanghai, and as far west as the Tsin Ling Mountains, Shensi, whence the U. S. National Museum has specimens (Cat. nos. 19230 and 35529).

Swinhoe obtained it in Formosa and sent specimens to the British Museum, which also has a specimen from Taiwan fu, received from the Formosa Museum, and several other specimens (males, females, and young) collected by Mr. Holst in West Taiwan fu.

List of specimens of Hyla chinensis.

Museum.	No.	Age.	Locality.	When collected.	By whom collected or from whom obtained.
U.S.N.M.....	19230	Adult.....	Shanghai, China.....	Brit. Mus.
Do.....	35529do.....	Shensi, China.....	1904	E. Blackwelder.
Brit. Mus.....do.....	Taiwan fu, Formosa.....	P. Holst.

Family ENGYSTOMIDÆ.

Jaws without teeth; shoulder-girdle, firmisternian and sacral lateral processes dilated. Widely distributed over the tropical portions of the globe. Most of the forms are terrestrial, some burrow in the ground, some are aquatic, but none are arboreal. Many of the species are ant-eaters.

Only one genus is represented within the area covered by this work.

Genus MICROHYLA^a Tschudi.

- 1838. *Microhyla* TSCHUDI, Classif. Batr., pp. 28, 71 (type, *Hylaplesia achatina*).
- 1841. *Microhyla* DUMÉRIL and BIBRON, Erpét. Gén., VIII, p. 613 (emendation).
- 1843. *Hylaplesia* FITZINGER, Syst. Rept., p. 31 (type, *H. achatina*) (not of Boie, 1827).
- 1843. *Siphneus* FITZINGER, Syst. Rept., p. 33 (type, *Engystoma ornatum*) (not of Brants, 1827).

^a From μικρος, small; and *Hyla*, the generic name of the common tree-toad.

1848. *Dendromanes* GISTEL, Naturg. Thierr., p. xi (substitute for *Microhyla*).
 1858. *Diplopelma* GUENTHER, Cat. Batr. Sal. Brit. Mus., p. 50 (type, *Engystoma ornatum*).
 1861. *Scaptophryne* FITZINGER, Sitz. Ber. Akad. Wiss. Wien, XLII (p. 146) (type, *S. labyrinthica*=*Engystoma pulchrum*).
 1872. *Ranina* DAVID, Nouv. Arch. Mus. Hist. Nat. Paris, VII, fasc. 4, Bull., p. 76 (type, *R. symmetrica*=*E. pulchrum*) (not of Lamarck, 1801).

The genus *Microhyla*, though belonging to an entirely different family, agrees with *Bufo* in having no teeth, but is easily distinguishable by the smooth skin, roundish pupil, and absence of parotoid glands.

This genus embraces at least nine species peculiar to China and various parts of the East Indies, including Ceylon and some of the islands in the Malay Archipelago, such as Sumatra, Java, and Borneo. Two forms allied to the widely distributed *Microhyla ornata*, which also occurs in China, are found within our territory, namely, one in Formosa and the other in the Riukiu Archipelago.

They may be distinguished as follows:

- a¹ Toes entirely free; metatarsal tubercles rather small. *M. fissipes*, p. 88.
 a² Toes with a distinct rudiment of web at base; metatarsal tubercles rather large. *M. okinavensis*, p. 89.

MICROHYLA FISSIPES^a Boulenger.

1884. *Microhyla fissipes* BOULENGER, Ann. Mag. Nat. Hist. (5), XIII, p. 397 (type-locality, Taiwan fu, southern Formosa; type in British Museum).

The original description by Boulenger is reproduced herewith. For further notes see the remarks under *Microhyla okinavensis* further on (pp. 89-90).

Original description of type-specimen.—British Museum; Taiwan fu, south Formosa; R. Swinhoe, collector. Habit slender; snout truncate, slightly longer than the orbital diameter; interorbital space broader than the upper eyelid; fingers slender, first much shorter than second; toes long and slender, free, with a slight lateral fringe; tips of fingers and toes not swollen; subarticular tubercles distinct; two rather small, obtuse, metatarsal tubercles; the hind limb being carried forward along the body, the tibio-tarsal articulation reaches the eye; skin nearly smooth above, with small warts on the sides. Olive-brown above, the small warts tinged with red; a darker lateral band from the tip of the snout, passing through the eye down to the middle of the side; an elongate X-shaped darker marking commencing between the eyes, and another, V-shaped, on sacral region; limbs with dark crossbars. From snout to vent 26 mm. (Boulenger, Ann. Mag. Nat. Hist. (5), XIII, 1884, pp. 397-398.)

Habitat.—This species seems to be restricted to Formosa. There are three specimens in British Museum, viz, the type collected by

^a From *Fissu*, split; *pes*, foot.

Swinhoe at Taiwan fu and two specimens secured by Holst in central Formosa. The United States National Museum has recently obtained a small specimen from "Daïpe or Daihoku," north Formosa, through Mr. A. Owston.

List of specimens of Microhylla fissipes.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	34469		Daïpe, north Formosa	Mar. —, 1903	A. Owston.
Brit. Mus.	(a)		Taiwan fu		R. Swinhoe.
Do.	94. 11. 14. 9		Central Formosa		P. Holst.
Do.	94. 11. 14. 10		do.		Do.

^a Type; description p. 88.

MICROHYLLA OKINAVENSIS ^a Stejneger.

1887. *Microhylla fissipes* BOULENGER, Proc. Zool. Soc. London, 1887, p. 150 (not of 1884) (Riu Kiu Islands).—OKADA, Cat. Vert. Jap., 1891, p. 66 (Okinawa).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 865, author's separate, p. 16.—BOETTGER, Offenbach. Ver. Naturk. 33–36 Ber., 1895, p. 106 ("Oshshima or Okinawa").
1901. *Microhylla okinavensis* STEJNEGER, Proc. Biol. Soc. Washington, XIV, 1901, Dec. 12, p. 189 (type-locality, Okinawa shima; type, Sci. Coll. Mus. No. 25a = U.S.N.M. No. 36553).
1902. *Microhylla undulata* BROWN, Proc. Phila. Acad., 1902, June 11, p. 186 (type-locality, "Loo Choo Islands;" types, Wistar Instit. Phila., Nos. 5726–8; Furness and Hiller, collectors).

I was so fortunate as to be able to compare five specimens of the present species with the type and three other Formosan specimens of *M. fissipes* in British Museum. The former consisted of two specimens brought by Pryer's collector from the Riu Kius (Brit. Mus. 87.1. 31. 43–44), and three collected by Holst in the mountain regions of Motobu, about 30 miles north of Nafa, Okinawa shima (Brit. Mus. 92. 9. 3. 24–26). The chief difference between the Formosan specimens and those from Riu Kiu consists in the presence of a distinct, though small, basal web between the toes and in the appreciably greater development of the metatarsal tubercles in the latter. Other differences which I thought that I had detected between my type-specimen and the original description of *M. fissipes* did not hold. Thus, the smoothness of the sides or presence of slight warts does not seem to be of any moment, and the coloration is essentially alike. The X and V shaped marks of the type, as described by Boulenger, are only due to the somewhat darker outlining anteriorly and posteriorly of the median dorsal patch; the median white dorsal stripe of the type of *M. okinavensis* is only shown by one of Pryer's specimens.


^a From Okinawa shima.

How unreliable a character is furnished by the comparative length of the hind legs is shown by Holst's Riukiu specimens. In the smallest of these the tibio-tarsal joint just reaches the snout; in the medium-sized specimens it goes considerably beyond the snout, and in the largest one only to the front of the eye.

The United States National Museum has now a large series from Ishigakishima, the first record from this group. They agree distinctly with the Okinawa type and not with the Formosa specimen, which we also now possess.

Brown's *Microhyla undulata* is the same as the present species and is published six months later. It is stated in the original description that the toes are "half-webbed," but that is a mistake, as an examination of the types in the Wistar Institute in Philadelphia showed me. The web in these specimens, as in the others, is between the metatarsals, while only a small portion of it extends beyond the base of the toes.

Description of type.—*Adult female*; U.S.N.M. No. 36553. Okinawa shima; Sci. Coll. Mus. Tokyo No. 25a (figs. 71-75).—Snout



FIGS. 71-75.—*MICROHYLA OKINAVENSIS*. $1\frac{1}{2} \times$ NAT. SIZE. 71, TOP OF HEAD; 72, SIDE OF HEAD; 73, OPEN MOUTH; 74, UNDERSIDE OF HAND; 75, UNDERSIDE OF FOOT. No. 36553, U.S.N.M.

rather pointed, slightly longer than diameter of orbit; inter-orbital space broader than upper eyelid; first finger much shorter than second; toes with a distinct though slight web at base and a mere indication of a fringe along the toes; tips of fingers and toes not widened; subarticular tubercles very distinct; metatarsal tubercles rather large, the inner measuring more than one-half the orbital diameter; the hind

limb being carried forward along the body, the tibio-tarsal joint barely reaches the eye; skin smooth above and on sides. Color (in alcohol) of a purplish gray above, with a light, very narrow median line from tip of snout to above anus; a spear-shaped dark brownish mark on occiput, from between the eyes, with the tip pointing backward and widening into two wavy dark bands which diverge toward the groins, contracting again on the sacrum, ending on the upper side of the thighs and embracing a space on the back somewhat darker than the ground color; outside of these bands and parallel with them are several faint, wavy lines of a dusky color; a dark brownish band of considerable width starts at the tip of the snout, covering the sides of face and neck, runs backward above the arm, is lost on the groin but reappears on the anterior aspect of the thigh;

the opposite edges of the dorsal and lateral dark bands are very sharply defined and emphasized by a very narrow light line; a dark band on the posterior aspect of the thighs, joining in an arch over the arms; a pale band from eye to the insertion of the fore leg; limbs with dark crossbars; underside whitish, heavily marbled with dark brownish gray on chin and throat, less so on anterior and lateral parts of belly; only a small portion of the latter posteriorly is immaculate.

Dimensions.

	<i>mm.</i>
Total length from snout to vent.....	28
Greatest width of body.....	20
Width of head at posterior angle of eye.....	8
Fore limb.....	13
Hind limb.....	43
Tibia.....	14

The *adult male* is somewhat smaller than the female, with slightly longer hind legs; throat distinctly blackish.

Variation.—The pale line down the middle of the back is often lacking and the faint wavy lines between the broad, heavy dorsal and lateral bands are often very faint or absent.

Habitat.—This species inhabits the islands between Japan and Formosa, from Okinawa shima southward, as shown by numerous specimens collected in the latter island and Ishigaki shima, of the Yaeyama subgroup.

Whether the species also occur on Miyako shima is not known.

List of specimens of Microhyla okinavensis.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.....	23906		Kunshiau, Okinawa shima.		I. Ijima.
Do.....	34472	Female	Ishigaki shima.....	April - June, 1899.	A. Owston.
Do.....	34473	do.	do.....	do.....	Do.
Do.....	34474	do.	do.....	do.....	Do.
Do.....	34475	do.	do.....	do.....	Do.
Do.....	34476	do.	do.....	do.....	Do.
Do.....	34477	do.	do.....	do.....	Do.
Do.....	34478	do.	do.....	do.....	Do.
Do.....	34479		do.....	do.....	Do.
Do.....	34480		do.....	do.....	Do.
Do.....	34481	Male	do.....	do.....	Do.
Do.....	34482		do.....	do.....	Do.
Do.....	34483		do.....	do.....	Do.
Do.....	34484	Male	do.....	do.....	Do.
Do.....	34485		do.....	do.....	Do.
Do.....	34486	Male	do.....	do.....	Do.
Do.....	34487	Female	do.....	do.....	Do.
Do.....	34488		do.....	do.....	Do.

List of specimens of *Microhyla okinavensis*—Continued.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	34489		Ishigaki shima	April - June, 1899.	A. Owston.
Do.	34490		do.	do.	Do.
Do.	34491	Male	do.	do.	Do.
Do.	36553	Female a.	Okinawa shima.		Sci. Coll.
Sci. Coll. Tokyo.	(1) P. 3		Shuri, Okinawa shima		Shiraishi.
Brit. Mus.	87. 1. 31. 43		Riu Kiu		J. S. Pryer.
Do.	87. 1. 31. 44		do.		Do.
Do.	92. 9. 3. 24		Motobu, Okinawa shima		P. Holst.
Do.	92. 9. 3. 25		do.		Do.
Do.	92. 9. 3. 26		do.		Do.
Wistar Inst.	5726	(b)	Riu Kiu		Furness and Hiller.
	5727	(b)	do.		Do.
	5728	(b)	do.		Do.

a Type; description, p. 90; figs. 71-75.

b Types of *M. undulata*.

Family RANIDÆ.

The frogs proper are characterized by teeth in the upper jaw combined with a firmisternian shoulder-girdle.

They constitute the largest family of the tailless batrachians, embracing considerably over one-fourth of all the species, and are distributed over nearly all the temperate and tropical regions of the globe, except Australia, New Zealand, eastern Polynesia, and Hawaii. Most of the species are more or less aquatic in their habits, but many are arboreal; some are terrestrial, and a few burrow in the ground. Several of the larger species are used for food in various countries and are considered great delicacies. Like all the other batrachians they are highly useful in destroying injurious insects.

Many of the frogs belonging to this family, especially those with large digital disks, resemble the tree-toads to a high degree; so much so in fact that until a comparative recent date they were referred to the latter, from which they differ radically, however, in their osteological characters. Nevertheless, there is an "external" character—i. e., one that can be ascertained without dissection—which will enable us to properly place them, viz, the lateral or median prolongation of the posterior free edge of the tongue so as to form two lateral "horns" with a deep median incision, or a single median "horn," as in the genus *Oxyglossus*. All the species known to occur within our territory are provided with two lateral posterior horns on the tongue.

Two genera only are represented in Japan and adjacent territory.

KEY TO THE GENERA OF JAPANESE FROGS.

- a¹ Distal and penultimate phalanges of digits continuous, without any kink separating them, there being no bone intercalated between them:^a (species occurring in Japan proper have tips of digits simple, without regular disks).....*Rana*, p. 95
- a² A well-defined kink, or transverse notch, on upper surface of digits between the distal and penultimate phalanges, due to an intercalated small bone:^a (tips of digits with large disks).....*Polypedates*, p. 143

Genus *RANA* ^b Linnæus.

1758. *Rana* LINNÆUS, Syst. Nat., 10 ed., I, p. 210 (type, *R. temporaria*); 12 ed., 1766, I, p. 354.
1814. *Ranaria* RAFINESQUE, Specchio Sci. (Palermo), II, fasc. 7 (substitute name for *Rana*).
1843. *Pelophylax* FITZINGER, Syst. Rept., p. 31 (type, *Rana esculenta*).
1891. *Crotaphitis* SCHULZE, Jahresber. Naturw. Ver. Magdeburg, 1890, p. 176 (type, *Rana temporaria*).
1891. *Baliopygus* SCHULZE, Jahresber. Naturw. Ver. Magdeburg, 1890, p. 177 (type, *Rana esculenta*).

Only such synonyms are here enumerated as refer directly to Japanese species or are additional to those quoted in recent synonymies.

The numerous species may be identified by the following:

SYNOPSIS OF THE SPECIES OF *RANA* RECORDED FROM JAPAN AND ADJACENT TERRITORY.

- a¹ A distinct dorso-lateral fold.
- b¹ Heels not overlapping.
- c¹ Vomerine teeth between choanæ.
- d¹ Dorso-lateral fold narrower than interorbital space; numerous distinct longitudinal ridges between folds.....*R. nigromaculata*, p. 94
- d² Dorso-lateral fold as broad as interorbital space; no longitudinal glandular ridges between dorso-lateral folds.....*R. planeyi*, p. 101
- c² Vomerine teeth behind choanæ.....*R. okinavana*, p. 102
- b² Heels overlapping.
- c¹ Tips of digits without regular disks.
- d¹ Hind legs twice as long as head and body.....*R. longicrus*, p. 104
- d² Hind legs much less than twice as long as head and body.
- e¹ A very prominent knob, rudiment of pollex, on inner side of first finger.
R. holsti, p. 105
- e² No prominent knob (not to be confounded with copulatory excrescences) on inner side of first finger.
- f¹ Snout long, pointed, distance from orbit to tip of snout longer than width between black stripes at anterior border of orbit.....*R. japonica*, p. 107
- f² Snout short, blunt; distance from orbit to tip of snout not longer than width between black stripes at anterior border of orbit.
- g¹ Vomerine teeth between choanæ, only their posterior end projecting backwards beyond them; webs large.....*R. temporaria*, p. 113
- g² Vomerine teeth behind level of choanæ, or nearly so; webs between toes small.
- h¹ Inner metatarsal tubercle rather large.....*R. tsushimensis*, p. 116
- h² Inner metatarsal tubercle small.....*R. amurensis*, p. 119
- c² Tips of digits dilated into distinct disks.....*R. ijima*, p. 121

^a It is well not to rely upon the "kink," but to ascertain the presence or absence of the intercalated phalanx by dissection.

^b Latin name for frog.

- a*² No dorso-lateral fold.
- b*¹ No tooth-like protuberance in lower jaw.
- c*¹ Distance of nostril from eye much less than twice that from edge of lip.
- d*² Tips of digits not dilated into disks.
- e*¹ Width of head two and a half times in length of head and body; heels scarcely overlapping; forehead concave; two terminal joints of fourth toe free *R. rugosa*, p. 123
- e*² Width of head about three times in length of head and body; heels overlapping considerably; forehead not concave; three terminal joints of fourth toe free *R. limnocharis*, p. 127
- d*² Tips of fingers dilated into distinct disks.
- e*¹ Skin above smooth *R. swinhoana*, p. 132
- e*² Skin above very rough, with warts in circular clusters of smaller surrounding larger ones *R. ishikawae*, p. 132
- c*² Distance of nostril from eye twice that from edge of lip *R. narina*, p. 134
- b*² Two large tooth-like protuberances in lower jaw.
- c*¹ Interorbital width much greater than width of upper eyelid. . . . *R. namiyei*, p. 136
- c*² Interorbital width less than upper eyelid *R. tigerina*, p. 139

RANA NIGROMACULATA^a Hallowell.

Plate X, fig. 1.

1765. ?*Rana chinensis* OSBECK, Reise Ostind. China, I (p. 244); Voy. China (Engl. ed.), I, 1771, p. 299 (Canton, China).
1838. *Rana esculenta* SCHLEGEL, Fauna Japon., Rept., pp. 109, 139, Saur. et Batr., pl. III, fig. 1^b (Japan) (not of Linnæus).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1866, p. 111 (Yokohama); 1876, p. 384 (Yokohama and Nagasaki).—HILGENDORF, Sitz. Ber. Berlin. Naturf. Fr., 1880, p. 119 (Tokyo).—FRITZE, Mitt. Deutsch. Ges. Ost-Asiens, V, 1891, p. 239 (Yezo).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, (8), XVII, No. 1, 1905, p. 337 (part: Tokyo, Amurland, etc.).
1858. *Rana viridis* BLEEKER, Natuurk. Tijdschr. Nederland. Indië, XVI, p. 204 (Japan) (not of Linnæus).
1859. *Rana esculenta* var. *japonica* MAACK, Putesh. na Amur (p. 153) (not *R. temporaria* var. *japonica* GUENTHER, 1858).—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 40 (Japan; Chifu; Chusan; Shanghai; Ningpo).—GIGLIOLI and SALVADORI, Proc. Zool. Soc. London, 1887, p. 595 (Fusan, Korea).—BOETTGER, Offenbach. Ver. Naturk. 26–28 Ber., 1888, p. 93 (Peking; Shanghai).—OKADA, Cat. Vert. Japan, 1891, p. 67 (Suwo; Yamaguchi; Tokyo; Mino; Gifu; Awaji).
1860. *Rana marmorata* HALLOWELL, Proc. Phila. Acad., 1860, p. 500 (type-locality, Simoda, Japan; type, U.S.N.M. No. 12159; Dr. Morrow, collector) (not of Massalongo, 1854).—CAMERANO, Atti Accad. Sci. Torino, XIV, Pt. 5, Apr., 1879, p. 871 (Tokyo; Yokohama).—Fox, Science (n. s.), XII, Nov. 9, 1900, p. 717.
1860. *Rana nigromaculata* HALLOWELL, Proc. Phila. Acad., 1860, p. 500 (type-locality, Japan; Dr. Morrow, collector).
1860. *Rana rugosa* HALLOWELL, Proc. Phila. Acad., 1860, p. 499 (part only: specimens from Simoda) (not of Schlegel).
1867. *Hoplobatrachus reinhardtii* PETERS, Mon. Ber. Berlin Akad. Wiss., 1867, p. 711 (type-locality, Malacca or China).—*Rana reinhardtii* MOELLEN-DORFF, Journ. N. China, Br. R. Asiat. Soc. (n. s.), XI, 1877, p. 105 (Peking).

^a From the Latin, signifying black-spotted.

^b Reproduced in this work on Plate X, fig. 1.

1868. *Tomopterna porosa* COPE, Proc. Phila. Acad., 1868, p. 139 (type-locality, Kanagawa; type in Mus. Comp. Zool. Camb. No. 305; Dr. Gulick, collector).—*Rana porosa* BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 40.
1880. *Rana esculenta marmorata* LATASTE, Bull. Soc. Zool. France, V, 1880, p. 61 (Peking); Le Natural., 1880 (p. 210) (not of Massalongo, 1854).
1891. *Rana esculenta* var. *nigromaculata* BOULENGER, Proc. Zool. Soc. London, 1891, pp. 376, 377, 383. (Yokohama; Hakone Lake; Okinawa shima, Riu Kiu).—BOETTGER, Kat. Batr. Mus. Senckenberg, 1892, p. 6 (China and Japan).—BEDRIAGA, Przewalski Reise, Zool., III, Pt. 1, 1899 (p. 11).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II, Klasse, Pt. 2, 1904, p. 358 (Shanghai and Hankow, China).
1897. *Rana esculenta* var. *chinensis* BOULENGER, Tailless Batr. Eur., II, p. 272; fig. D, p. 273 (Korea and Japan to southern China and Siam).—MÉHELY, Zichy's Dritte Asiat. Forschungsr., II, 1901, p. 62 (Peking).
1899. *Rana chinensis* STONE, Proc. Phila. Acad., 1899, p. 183 (Khingian mountains and Lama-mio, eastern Mongolia).

In an article on "The names of animals, published by Osbeck in 1765,"^a Mr. W. J. Fox states that Osbeck's *Rana chinensis* is "without doubt a *Bufo*," and that "Boulenger has erred in referring it to *Rana*, as Osbeck says in his description, 'the body above warty,' which sufficiently indicates that the species is not a *Rana*." Now, if Mr. Fox had known *R. tigerina* and *R. rugosa*, he would not have said this, since they are a great deal more warty than many a *Bufo*. There are points in Osbeck's description which show that the animal was "without doubt" a *Rana* and that Boulenger did not err in so referring it.

Whether Boulenger was right, however, in referring *Rana chinensis* Osbeck to the eastern representative of *R. esculenta* is another question, and as I have grave doubts as to the correctness of this view I prefer to leave the name with a ? as unidentifiable. The expression "the body is warty at the top" (to use the English translation of 1771) might be applied to the frog in question, but it applies much better to *Rana tigerina*, which is also sold in the markets of Hongkong and Canton. With this agrees also the character: "The throat is white, speckled with black." On the other hand, Osbeck's allusion to the "sixth or innermost" finger seems to fit better the larger metatarsal tubercle of *R. nigromaculata*. As the name can not therefore be applied with absolute certainty to either of these species I think it is better to dispense with it altogether.


Mr. Fox's plea for *Rana marmorata* Hallowell, 1860, as the proper name for the eastern frog can not be entertained either, as that name was disposed of by Massalongo six years earlier^b for one of the European forms of *Rana esculenta*.

^a Science (n. s.), XII, Nov. 9, 1900, pp. 716-718.

^b *Rana esculenta* var. *marmorata* Massalongo, Saggio di un' erpetol. popol. Verones., 1854 (p. 47).

There seems to be no good reason for applying a trinomial to this eastern representative of *Rana esculenta*. The character of the numerous short dermal ridges on the back between the dorso-lateral folds seems absolutely diagnostic. Moreover, in proportions it differs more from its nearest neighbor, *Rana ridibunda*, than from the more remote forms of *R. esculenta* inasmuch as the heels do not overlap. Finally, it has not been shown as yet that the range of our species meets that of any of the western forms.

The types of *R. nigromaculata* of Hallowell (collected by Doctor Morrow) are not in the National Museum, but the description fits our species fairly well. The types of his *R. marmorata*, however, are still



FIGS. 76-80.—*RANA NIGROMACULATA*. NAT. SIZE. 76, HEAD FROM ABOVE; 77, HEAD FROM SIDE; 78, OPEN MOUTH; 79, UNDERSIDE OF HAND; 80, UNDERSIDE OF FOOT. No. 7415, U.S.N.M.

in good state of preservation (U.S.N.M. No. 12159), being apparently two females very much like the specimen figured by Schlegel (*l. c.*) as *R. esculenta*.^a Hallowell's measurements were apparently taken (in a rather careless manner) from the smaller specimen. It is worthy of note that Hallowell referred several other specimens, males with large vocal vesicles (U.S.N.M. No. 7415, figs. 76-80), to *Rana rugosa*, thus enumerating the specimens of the same species, and practically from the same locality, under three different specific names.

A young specimen in the collection of the Philadelphia Academy of Sciences (No. 15256) collected by Dr. W. H. Jones, at Chemulpo,

^a Plate IX, fig. 1 of this work.




FIG. 1, *RANA NIGROMACULATA*; FIGS. 2-4, *R. RUGOSA*. (FROM SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 554.

Korea, August, 1890, is almost without spots on the back and has an oblique black line behind the tympanum suggesting the ear spot of *R. japonica*. Such a specimen is superficially much like the latter species, but the much shorter tarsus and femur, the much larger inner metatarsal tubercle, and the longitudinal dermal ridges on the back between the dorso-lateral ridges at once indicate *R. nigromaculata*.

Description.—*Adult male*; U.S.N.M. No. 23579; Yokohama; September, 1896; L. Stejneger, collector. Vomerine teeth in two slightly oblique groups between the choanæ, their posterior end extending somewhat behind the latter; nostrils somewhat nearer the eye than the tip of the snout; distance between nostrils equals their distance from orbits; interorbital space narrow, scarcely more than half the width of upper eyelid; tympanum slightly longer than high, its longest diameter four-fifths that of the eye, and more than twice its distance from the latter; first finger considerably longer than second, its basal phalanx and the metacarpal on the inner side provided with a large swollen pad very minutely granulated; tips of digits tapering, obtusely pointed, the terminal phalanx being set off from the rest of the digit by a fold across the joint which is markedly more distinct than on the other joints; toes about three-fourths webbed, the deepest incision of the webs on both sides of fourth toe scarcely reaching the end of its basal phalanx; subarticular tubercles small but distinct; inner metatarsal tubercle large, projecting, sharp, its length nearly four-fifths the length of the first toe from the anterior end of the tubercle a distinct outer metatarsal tubercle; no tarsal fold; hind limb being carried forward along the side of the body the tibio-tarsal joint reaches the anterior border of the eye; heels do not meet when hind limbs are bent at right angles to axis of body; a strongly marked dorso-lateral fold from posterior border of eye to insertion of hind limb; between these folds about four series of somewhat irregular but sharply marked longitudinal dorsal folds, the longest about as long as upper eyelid; skin above indistinctly shagreened, more coarsely granular on the sides and on the posterior aspect of the femur, transversely rugose on the belly, nearly smooth on chest and throat; a large external vocal sac on each side below the angle of the mouth. Color (in alcohol) above brownish olive with longitudinal blackish spots, one surrounding each dorsal glandular fold; dorso-lateral folds paler brownish; a narrow pale bluish line from the snout down the median line of the back; sides with irregular blackish spots which encroach upon the lower edge of the dorso-lateral fold; upper surface of limbs with dusky spots which form crossbars on femur; upper lip with a few small dusky spots; underside uniform whitish.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent.....	60
Width of head.....	22
Length of fore leg.....	34
Length of hind leg, tip of longest toe to vent.....	92
Length of tibia.....	25
Diameter of tympanum.....	5
Length of inner metatarsal tubercle.....	5

The *adult female* (U.S.N.M. No. 23576) lacks the vocal sacs and the swollen pad at the base of the first finger, but agrees in other respects closely with the description of the male given above.

Variation.—My September specimens from Yokohama are all essentially like the ones described above. The specimens collected by Doctor Smith in Shikoku on May 11, as well as Owston's from Mount Fuji, also collected in May, differ chiefly in coloration, the black pigment being much more concentrated, the pale areas consequently lighter, and the whole appearance brighter and with greater contrasts. Thus the pale median dorsal line is greatly widened and the ground color between it and the dorso-lateral folds so darkened as to entirely obscure the dorsal spots; a black line runs from tip of snout through nostril and eye bending down sharply behind the tympanum; the dark blotches on the hind legs form distinct and strongly contrasted cross-bars. With the elongation of the blackish coloration on the back there is associated a distinct lengthening of the dorsal glandular folds so that they run together into long irregular ridges. Jouy's specimens from near Seoul, which were taken in June, are very much like the ones from Shikoku, and so are Hallowell's types of *R. marmorata*. The difference is evidently seasonal.

Habitat.—Widely distributed in eastern Asia from Vladivostok (Hamburg Mus., No. 871, collected by Dieckmann) in the North to Bangkok, Siam (Brit. Mus.), in the South. Maack collected it on the Amur and Ussuri rivers and Radde in the Khingan Mountains. It is recorded from numerous localities in China, such as Peking (Moellendorff and Lataste); Shanghai, Kiukiang, Chapu, Hankow (Boettger); Chefu, Chusan, Ningpo (Boulenger); Ordos (Przewalski). There are specimens in British Museum from Broughton Bay and Gensan, Korea; in the Philadelphia Academy from Chemulpo (No. 15256), and others in the U. S. National Museum from Seoul and Fusan. Stone records it from the Khingan Mountains and Lama-mio in eastern Mongolia.

On the map in Boulenger's *Tailless Batrachia of Europe* (at p. 263), showing the distribution of the *Rana esculenta* group, the present species is also indicated as inhabiting Formosa, though I am unable to state upon what evidence this is done.

In Japan proper *R. nigromaculata* is recorded from the provinces of Suwo and Mino and on the island of Awaji (Okada); from Naga-

saki and Yokohama (von Martens); Tokyo (Okada and Hilgendorf); Miyazaki on Kiusiu, Sumiyoshi, near Osaka, Koriyama Province, Yamato, Yokohama and Simoda in Hondo (U.S.N.M.); Hakone Lake (Brit. Mus.). The United States National Museum also has specimens collected in Shikoku by Dr. H. M. Smith. Buerger's and von Siebold's numerous specimens in the Leiden Museum (Nos. 1946-1947) are only labeled "Japon." The museum in Stockholm has specimens brought home by the Vega Expedition from Biwa, Yokohama, and Hirosami.

The only record of its occurrence in Yezo is by Doctor Fritze, who states that it is "common in every pond." I am not aware that authentic specimens were identified, and the statement needs corroboration.

There is a specimen (male) in British Museum, received from G. E. Mason, said to be from "Great Loo Choo Island," but this locality is also open to question. Whether the specimens from Oo-shima (Amami-o-shima) recorded by Hallowell as *R. rugosa* really belong to this species must remain unsettled until that island can be thoroughly explored.

List of specimens of Rana nigromaculata.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	a 7415	Male a	Simoda, Hondo.	May, 1855	W. Stimpson.
Do.	b 7416	Male.	do	do	Do.
Do.	a 8852	Female	Japan.		?
Do.	b 8832	Halfgrown	do		?
Do.	c 8852	do	do		?
Do.	a 12159	Female b	Simoda, Hondo.		Doctor Morrow.
Do.	b 12159	do	do		Do.
Do.	22188		Japan.		?
Do.	22189		do		?
Do.	23576	Female c	Yokohama, Hondo.	Sept., 1896	L. Stejneger.
Do.	23577	do	do	do	Do.
Do.	23578	do	do	do	Do.
Do.	23579	Male d	do	do	Do.
Do.	23580	Female	do	do	Do.
Do.	23581	do	do	do	Do.
Do.	23582	do	do	do	Do.
Do.	23583	do	do	do	Do.
Do.	23584	do	do	do	Do.
Do.	23585	do	do	do	Do.
Do.	23586	do	do	do	Do.
Do.	26045	Halfgrown	do	do	Do.
Do.	30739	Male	Miyazaki, Kiusiu.		C. A. Clark.
Do.	31799	do	Sumiyoshi, near Osaka.	Apr. 11, 1903	H. M. Smith.
Do.	31800	do	do	do	Do.
Do.	31808	Young	Koriyama, Yamato.		Do.
Do.	31815	Halfgrown	do	Apr. 26, 1903	Do.
Do.	31816	do	do	do	Do.

^a Figs. 76-80; recorded by Hallowell as *R. rugosa*.

^b Types of *R. marmorata*.

^c Description, p. 98.

^d Specimen described, p. 97.

List of specimens of *Rana nigromaculata*—Continued.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	31902	Female...	Kochi, Shikoku.....	May 11, 1903	H. M. Smith.
Do.	31903	Male.....	do.....	do.....	Do.
Do.	31927	Halfgrown.....	do.....	do.....	Do.
Do.	31928	do.....	do.....	do.....	Do.
Do.	31933	Female.....	do.....	do.....	Do.
Do.	31934	do.....	do.....	do.....	Do.
Do.	31935	Male.....	do.....	do.....	Do.
Do.	31936	Female.....	do.....	do.....	Do.
Do.	31937	do.....	do.....	do.....	Do.
Do.	31938	do.....	do.....	do.....	Do.
Do.	31939	Halfgrown.....	do.....	do.....	Do.
Do.	31940	do.....	do.....	do.....	Do.
Do.	31941	Male.....	do.....	do.....	Do.
Do.	34413	Halfgrown.....	Mount Fuji, Hondo.....	May, 1898	A. Owston.
Do.	34414	do.....	do.....	do.....	Do.
Do.	34415	do.....	do.....	do.....	Do.
Do.	34416	do.....	do.....	do.....	Do.
Do.	34417	do.....	do.....	do.....	Do.
Do.	34418	do.....	do.....	do.....	Do.
Do.	34419	do.....	do.....	do.....	Do.
Do.	34420	do.....	do.....	do.....	Do.
Do.	34421	Young.....	do.....	do.....	Do.
Do.	34422	Halfgrown.....	do.....	do.....	Do.
Do.	34423	do.....	do.....	do.....	Do.
Do.	34424	Young.....	do.....	do.....	Do.
Do.	34425	Halfgrown.....	do.....	do.....	Do.
Do.	34426	do.....	do.....	do.....	Do.
Do.	34427	do.....	do.....	do.....	Do.
Do.	34428	Young.....	do.....	do.....	Do.
Do.	34429	do.....	do.....	do.....	Do.
Do.	34430	do.....	do.....	do.....	Do.
Do.	34431	do.....	do.....	do.....	Do.
Do.	34432	do.....	do.....	do.....	Do.
Do.	34468	Halfgrown.....	Yokohama, Hondo.....	Apr. 2, 1901	Do.
Phila. Acad.	14421	(a)	Simoda, Hondo.....	1855	Dr. Morrow.
U.S.N.M.	17512	Young.....	Fusan, Korea.....	1885	P. L. Jouy.
Do.	21195	Female.....	Seoul, Korea.....	May, 1883	Do.
Do.	21196	do.....	do.....	do.....	Do.
Do.	21197	do.....	do.....	do.....	Do.
Do.	21198	do.....	do.....	do.....	Do.
Do.	21199	do.....	do.....	1883	Do.
Do.	21200	do.....	do.....	do.....	Do.
Do.	21201	do.....	do.....	do.....	Do.
Do.	21202	Halfgrown.....	do.....	do.....	Do.
Do.	21203	do.....	do.....	do.....	Do.
Do.	21204	do.....	do.....	do.....	Do.
Do.	21205	do.....	do.....	do.....	Do.
Do.	21206-10	do.....	do.....	do.....	Do.
Phila. Acad.	15256		Chemulpo, Korea.....	Aug., 1890	W. H. Jones.
Hamburg.	871		Vladivostok.....	1894	Dieckmann.

^a Cotype of *R. marmorata*.

RANA PLANCYI ^a Lataste.

1880. *Rana plancyi* LATASTE. Bull. Soc. Zool. France, V, 1880, p. 64 (type-locality, Peking, China); Le Naturaliste, 1880 (p. 210).—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 37 (Shanghai; Chusan; Ningpo; Formosa).

Description.—British Museum (one of Nos. 84. 3. 11. 1–5); Taiwan fu, Formosa; presented by Formosa Museum. Vomerine teeth in two slightly oblique groups, their posterior edge barely extending beyond the hinder edge of the choanæ; snout rather long, the distance from the tip to eye longer than that from the latter point to tympanum; lores concave; tympanum nearly as large as the eye; interorbital space narrower than upper eyelid; first finger extending slightly beyond second; toes fully webbed; inner metatarsal tubercle rather large and strong; hind leg being carried forward the tibio-tarsal joint reaches the anterior corner of the eye; heels not overlapping; dorso-lateral glandular fold prominent, scarcely as wide as the interorbital space; no longitudinal glandular ridges between the dorso-lateral folds. Color (in alcohol): Upper parts with indistinct blackish spots; underside distinctly mottled with dusky, especially on throat, chest, and thighs; marblings on posterior aspect of thigh confluent into an ill-defined band; a fairly distinct dusky band from axilla to groin.

<i>Dimensions.</i>	<i>mm.</i>
Snout to vent.....	44
Snout to posterior edge of tympanum	18
Width of head.....	16
Diameter of eye.....	5.5
Width of upper eyelid	4
Interorbital width.....	2.5
Snout to eye.....	8
Diameter of tympanum.....	5
Fore leg.....	25
Tibia.....	20.5
Width of dorso-lateral fold.....	1.75

Variation.—Formosa specimens in British Museum do not have the dorso-lateral fold as broad as specimens from Shanghai (Brit. Mus.), in which it is fully as broad as the eyelid, while in the Formosan ones it barely reaches the width of the interorbital space; yet it is considerably wider than in *R. nigromaculata*. There are no distinct glandular ridges between the folds. The vomerine teeth are between the choanæ, fully as far forward as in *R. nigromaculata*, while in the Shanghai specimens their position is somewhat more posterior. Boulenger ^b

^a For Mr. V. Collin de Plancy, attaché at the French legation in Peking, who collected the types. He is author of a Catalogue des Reptiles et Batraciens du Département de l'Aube et Étude sur la Distribution Géographique des Reptiles et Batraciens de l'est de la France, 1877, and other herpetological papers.

^b Cat. Batr. Sal. Brit. Mus., p. 37.

says, "Seldom a light vertebral line," but of the 5 specimens from Taiwan fu, since received, 4 have a very distinct broad whitish line down the middle of the back. Inner metatarsal tubercle variable; in most specimens it is strong and large, while in a female from Formosa (by Dickinson) it is comparatively small.

Habitat.—Plancy's frog was originally described from Peking, and specimens from Shanghai, Chusan, and Ningpo are in British Museum.

The specimens recorded from Formosa are all in British Museum, the only ones with a definite locality being from Taiwan fu.

List of specimens of Rana plancyi.

Museum.	Number.	Sex.	Locality.	When collected.	By whom collected.
Brit. Mus.	65. 10. 29. 37		Formosa		R. Swinhoe.
Do.	65. 10. 29. 38		do.		Do.
Do.	79. 6. 4. 25	Female	do.		Dickinson.
Do.	84. 3. 11. 1		Taiwan fu, Formosa.		Formosa Mus.
Do.	84. 3. 11. 2		do.		Do.
Do.	84. 3. 11. 3		do.		Do.
Do.	84. 3. 11. 4		do.		Do.
Do.	84. 3. 11. 5		do.		Do.

RANA OKINAVANA *a* Boettger.

1895. *Rana okinavana* BOETTGER, Zool. Anz., XVIII, July 8, 1895, p. 266 (type-locality, Okinawa shima, Riu Kiu; types, Mus. Senckenberg. No. 1072, 1a); Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 103.

This species is not represented in any of the collections which have reached the U. S. National Museum or the Science College Museum in Tokyo. I owe the inspection of the two type specimens in the Senckenberg Museum in Frankfort on the Main to the kindness of Doctor Boettger, whose original description I give below in translation. I have only to add that the heels do not overlap. A third specimen also from the late Mr. B. Schmacker, whose Japanese collector obtained them in the Riu Kius, is in the Bremen Museum according to Doctor Boettger.

Original description by Doctor Boettger.—*Adult female*; Senckenberg Museum, No. 1072; Okinawa shima. In general similar to *R. lateralis* Boulenger, but with shorter snout and more robust; vomerine teeth in two rounded bundles beginning on a line with the posterior border of the choanæ and separated from them as well as from each other by equal intervals; head moderately large; snout short, anteriorly slightly pointed, and somewhat produced, as long as the diameter of the eye; nostrils equidistant from tip of snout and eye; canthus rostralis angular, loreal region slightly depressed longitudinally; interorbital space scarcely broader than the upper eyelid; tympanum very distinct, three-fourths the size of the eye; fingers moderately

a From Okinawa shima.

long, first longer than second; toes three-fifths webbed and in addition with narrow dermal margins to the last phalanx; tips of digits truncate but without distinct disks; subarticular tubercles well developed; inner metatarsal tubercle oval, prominent, but less than half the length of the inner toe; no outer metatarsal tubercle; the adpressed hind leg reaches between anterior border of eye and nostril; skin on back and underside rather smooth, on the sides with large flat warts; a narrow, high, strongly swollen dorso-lateral glandular fold; a second fold beginning beneath the eye and ending above the shoulder in two round or oval glandular aggregations. Upper side gray, frequently with a fine, somewhat lighter median dorsal line; a black wedge-shaped spot from nostril through eye extending over the temporal region; a blackish line bordering the upper lip and separated from the freno-temporal spot by a pure white band which ends on the glandular protuberances above the insertion of the arm; tympanum brown; dorso-lateral fold externally edged with black, the sides being often entirely blackish; posterior extremities with dark cross bands; posterior aspect of femur whitish yellow with black spots and marblings; underside whitish yellow, spotted and dotted all over very densely with blackish brown, the underside of the head, however, being mostly uniform blackish gray.

Dimensions.

	Female.	Female.	Female.
	<i>mm.</i>	<i>mm.</i>	<i>mm.</i>
Length of head and body.....	40	42	46
Length of head.....	16	16.5	17
Width of head.....	15	15.5	16
Diameter of tympanum.....	3.5	3.5	4
Fore legs.....	22.5	24.5	25
Hind legs.....	67	73	76
Tibia.....	20.5	22.5	a 23.5

a Boettger, Zool. Anz., XVIII, 1895, pp. 266-267.

Remarks.—The *Rana lateralis* Boulenger with which Boettger compares this species is described from Tenasserim and is related to *R. malabarica* from Malabar and the Bombay hills. *R. okinavana* appears, then, to be of decided Indian affinities.

Habitat.—There is no record of this species having been observed beyond the type locality, Okinawa shima in the Riu Kiu Archipelago.

List of specimens of Rana okinavana.

Museum.	Number.	Sex.	Locality.	When collected.	By whom collected or from whom obtained.
Senckenberg.....	1072, 1	Female a.	Okinawa shima.....		B. Schmacker.
Do.....	1072, 1	do.....	do.....		Do.


a Type; description, p. 102.

RANA LONGICRUS^a Stejneger.

1898. *Rana longicrus* STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 216 (type-locality, Taipa, Formosa; type, Sci. Coll. Mus. Tokyo, No. 26; T. Tada, collector).

The head and body of this species resembles greatly *R. japonica*, the snout being fully as pointed and elongated, but the most remarkable character is the excessively lengthened hind legs which are actually more than twice as long as the head and body together. Even the fore legs partake of this elongation, though less markedly. The coloration is that characteristic of the group.

Description of type-specimen.—*Adult*; Science College Museum, Tokyo, No. 26; Taipa, Formosa; September, 1896; T. Tada, collector (figs. 81–83). Vomerine teeth in two oblique groups



FIGS. 81–83.—*RANA LONGICRUS*. NAT. SIZE: 81, TOP OF HEAD; 82, SIDE OF HEAD; 83, UNDER SIDE OF FOOT. NO. 26, SCI. COLL. TOKIO.

behind a line through the center of the choanæ, and projecting backward beyond the latter; snout long, pointed, distance from tip to eye much greater than horizontal diameter of orbit; nostril nearer the tip of the snout than the eye; interorbital space much wider than upper eyelid; tympanum two-thirds the diameter of the eye; first finger not extending beyond second; no knob-like protuberance on inner side of thumb; tip of digits not tapering, obtusely rounded; webs of feet deeply excised, the incision of the web on both sides of fourth toe reaching the middle of the basal

phalanx; subarticular tubercles well developed; inner metatarsal tubercle strongly developed, outer one barely indicated; hind legs extremely elongate, the distance from anus to tip of longest toe twice as great as length of head and body; tibio-tarsal joint of extended hind leg reaching beyond the tip of the snout by the width of the head at the center of the eye; kneejoint of adpressed limb reaches the axilla; heels overlap by the length of the forearm; skin smooth, back and sides indistinctly shagreened with a few irregular tubercles on the sacrum; posterior aspect of femurs coarsely granulated; a narrow, but very distinct dorso-lateral fold on each side from posterior border of eye to insertion of hind leg, the nearest approximation of the two folds being equal to the distance between

^a Signifying long-leg.

the anterior corners of the eyes. Color (in alcohol) above pale drab,^a with a few distinct and irregular small dusky spots between eyes, on the middle of the back and along the outer edge of the dorso-lateral fold; a distinct dark brown line from tip of snout through nostril to eye, the upper edge being sharply defined, the lower fading gradually into the loreal region; from eye to shoulder a large, sharply defined oblong spot of dark brown covering the tympanum; upper and lower lips marbled brown and whitish; pale indications of the usual mark on the upper arm as well as cross-bars on fore and hind limbs; an ill-defined brown line along outer aspect of femur and tibia, underside of tarsus, and fifth toe.

Dimensions.

	<i>mm.</i>
From snout to vent.....	52
From snout to posterior edge of tympanum.....	17
From snout to corner of mouth.....	16.5
Width of head.....	15
Diameter of eye.....	4.5
Width of upper eyelid.....	3
Interorbital width.....	4
From eye to nostril.....	4.7
From eye to end of snout.....	8
Diameter of tympanum.....	3
From eye to tympanum.....	2
Fore limb.....	34
Elbow to tip of longest finger.....	24
Hind limb.....	105
Thigh.....	29
Tibia.....	36
Metatarsal tubercle.....	1.5

Habitat.—This species is only known from Formosa, Mr. T. Tada having collected a single specimen at Taipa, September, 1896.

List of specimens of Rana longicrus.

Museum.	No.	Age.	Locality.	When collected.	By whom collected.
Sci. Coll., Tokyo.....	26	Adult ^a	Taipa, Formosa.....	Sept., 1896	T. Tada.

^a Type; description, p. 104; figs. 81-83.

RANA HOLSTI^b Boulenger.

1892. *Rana holsti* BOULENGER, Ann. Mag. Nat. Hist. (6), X, October, 1892, p. 302 (type-locality, Okinawa shima, Riu Kiu; type, Brit. Mus. No. 92.9.3.19; Holst, collector).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 865; author's separate, p. 16.

^a Ridgway, Nomencl. Col., Plate III, fig. 18.

^b For the late Mr. P. A. Holst, who collected the type.

As stated by Boulenger, this species, which is one of the largest of the group, is particularly well characterized by the knob on the inner side of the first finger, and by the elongate inner metatarsal tubercle. The nostrils are considerably nearer the end of the snout than the eye, and there is no outer metatarsal tubercle. The interorbital width is also rather unusual. I have had the privilege of examining the unique type specimen, but I have nothing to add to Boulenger's original description, which is therefore reproduced herewith.

Original description of type specimen.—*Adult female*; British Museum No. 92.9.3.19; Okinawa shima, Riukiu; Holst, collector (fig. 84). Near *R. temporaria*; vomerine teeth in two well-developed oblique groups behind the level of the choanæ; head broader than long; snout rounded, slightly prominent, as long as the diameter of the




FIG. 84.—*RANA HOLSTI*. NAT. SIZE. UNDERSIDE OF LEFT HAND. No. 92.9.3.19, BRIT. MUS.

orbit; loreal region nearly vertical, slightly concave; nostrils a little nearer the end of the snout than to the eye, the distance between them equal to the interorbital width, which equals the width of the upper eyelid; tympanum very distinct, circular, measuring two-thirds diameter of eye and about once and a half its distance from orbit; fore limb longer than tibia; first finger extending considerably beyond second; tips of fingers blunt, subarticular tubercles strong; a very prominent knob (rudiment of pollex) on inner side of first finger; tibio-tarsal articulation reaching the eye; tibia slightly longer than foot, half length of head and body; toes three-fourths webbed, the two distal phalanges of fourth toe free, but with the membrane prolonged as a narrow fringe on each side; subarticular tubercles strong; inner metatarsal tubercle blunt, elliptical, not very prominent, three-fifths length of inner toe; no outer metatarsal tubercle; no tarsal fold; back with a few scattered small warts, sides and hind limbs with numerous warts; body and limbs with whitish pearl-like excrescences; glandular lateral folds prominent, broken up into warts behind, nearly parallel, the distance between them on the scapular region two-ninths length of head and body. Olive-brown above, sides with blackish spots; a blackish temporal spot; tympanum reddish brown; a light streak from below the eye to the angle of the mouth; limbs with dark crossbars; hinder side of thighs marbled with black; throat spotted with brown; belly with a few brown dots.

Dimensions.

	mm.
From snout to vent.....	120
Length of head.....	37
Width of head.....	43
Diameter of eye.....	12
Interorbital width.....	11
From eye to nostril.....	9
From eye to end of snout.....	17
Tympanum.....	8
From eye to tympanum.....	5
Fore limb.....	70
Hind limb.....	170
Tibia.....	58
Foot.....	55
Inner toe.....	14
Inner metatarsal tubercle.....	8 ^a

Habitat.— Only the type-specimen having been obtained thus far, we know nothing of the distribution of this species beyond the fact that it was collected by Holst in Okinawa shima.

List of specimens of Rana holsti.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.
Brit. Mus.	92.9.3.19	Female ^a ...	Okinawa shima, Riu Kiu....	P. A. Holst.

^a Type; description, p. 106; fig. 84.

RANA JAPONICA ^b (Guenther).

Plate XI, fig. 1.

1838. *Rana temporaria* SCHLEGEL, Fauna Japon., Rept., pp. 109, 139; Saur. et Batr. pl. III, fig. 2^c (Japan) (not of Linnaeus).—BLEEKER, Natuurk. Tijdschr. Nederland. Indië, XVI, 1858, p. 205 (Japan).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1866, p. 111 (Yokohama); 1876, p. 380 (Tokyo and Yokohama).—HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 119 (Tokyo).
1858. *Rana temporaria* var. *japonica* GUENTHER, Cat. Batr. Sal. Brit. Mus., p. 17 (Ningpo and Chusan, China; Japan.)
1860. *Rana sylvatica* HALLOWELL, Proc. Phila. Acad., 1860, p. 500 (Simoda, Hondo) (not of Le Conte).—CAMERANO, Atti Accad. Torino, XIV, Pt. 5, Apr., 1879, p. 876 (Tokyo; Yokohama).
1879. *Rana japonica* BOULENGER, Bull. Soc. Zool. France, 1879, p. 190 (Japan); 1886, p. 599 (Japan; China); Cat. Batr. Sal. Brit. Mus., 1882, p. 47 (Japan; Szechuen and Ningpo, China).—BOETTGER, Offenbach. Ver. Naturk. 26–28 Ber., 1888, p. 96 (Peking); Kat. Batr. Mus. Senckenberg., 1892, p. 9 (Nikko and Yokohama, Japan; Dalanshan, near Ningpo, Da-dschiao-sy, near Peking, Lueshan, near Kiukiang, Lake Tahoo, near Shanghai, China); Offenbach. Ver. Naturk. 33–36 Ber., 1895, p. 102 (Oho

^a Boulenger, Ann. Mag. Nat. Hist. (6), X, 1892, pp. 302–303.

^b Signifying Japanese.

^c Reproduced in this work on Plate XI, fig. 1.

- shima or Okinawa).—OKADA, Cat. Vert. Japan, 1891, p. 67 (Tokyo; Awaji; Suwo).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 358 (Shanghai).
1886. *Rana martensi* BOULENGER, Proc. Zool. Soc. London, 1886, p. 414 (type-locality, Tokyo; types, Berlin Mus. Nos. 4410-4411; Martens, collector).—BOETTGER, Kat. Batr. Mus. Senckenberg., 1892, p. 10 (Ekingami, near Yokohama).
1886. *Rana martensii* BOULENGER, Bull. Soc. Zool. France, 1886, p. 599; author's separate (p. 5).
1904. *Rana japonica* var. *ornativentris* WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, p. 383 (type-locality, Nikko, Hondo, Japan; type in Zool. Mus. Akad. Wiss. Munich; Doctor Haberer, collector).

R. japonica seems chiefly characterized by the pointed snout. This character is seconded by rather long legs, the tibio-tarsal joint as a rule reaching beyond the snout, though in several specimens falling short of it. The inner metatarsal tubercle is rather strong.

From this I can not separate *R. martensi*, based by Boulenger upon Yedo (now Tokyo) specimens. I have examined the cotype in the British Museum, a rather short-legged female, but I can find no character, neither in the position of nostrils, metatarsal tubercle, fingers, nor coloration, which will separate this specimen from a large series of true *R. japonica*; it has also the pointed snout of the latter.^a

Description.—*Adult male*; U.S.N.M. No. 23549; Yokohama, Hondo; Sep. 1896; L. Stejneger, collector (figs. 85-89). Vomerine teeth in two oblique patches on a line with the center of the choanæ and extending slightly behind the latter; snout long, pointed, the distance from orbit to tip of snout longer than width between black stripes at the anterior border of orbit; nostrils nearer the tip of snout than the eyes; interorbital space equals the width of upper eyelid; tympanum about two-thirds the diameter of eye, and slightly more

^a The frogs of the *Rana temporaria* group present some of the most difficult problems to the student, even with authentic specimens of all the forms at hand, and nowhere do the perplexities seem greater than with the species inhabiting the area of which we treat. The principal difficulty consists in the fact that several of the described species are said to occur in the same localities, and are based upon characters which seem vanishing in a large series. Thus *R. japonica* and *R. martensi* are presumably both from central Japan, yet in no publication have I been able to find their characters contrasted. In a vague way it is intimated that the former is more nearly related to *R. dalmatina* (= *agilis* Thomas), the latter to *R. temporaria*, but that is in itself but scant consolation inasmuch as both *R. japonica* and *martensi* agree with *R. temporaria* in the character in which it is stated that the latter differs from *R. dalmatina*. That I am not the only herpetologist disconcerted by this state of affairs is plain from Dr. O. Boettger's remarks upon a number of specimens from Chinhai, northeast of Ningpo, China (Ber. Senckenberg. Naturf. Ges., 1894, pp. 146-147), which he refers to the three species *R. amurensis*, *martensi*, and *japonica* in spite of a number of discrepancies between the specimens and the descriptions, which, it seems to me, break down the barriers between them.





FIG. 1, RANA JAPONICA; FIGS. 2, 3, R. AMURENSIS. (FROM SCHLEGEL AND BOULENGER.)

FOR EXPLANATION OF PLATE SEE PAGE 554.

than twice its distance from the latter; fingers obtuse, first extending beyond second, and provided with a large double swelling on the inner side, the surface of which is minutely granulated; tip of toes tapering; webs considerably excised, the incision on both sides of fourth toe extending to the distal third of the basal phalanx; subarticular tubercles rather weak; inner metatarsal tubercle rounded, projecting; no outer metatarsal tubercle; tibio-tarsal joint of extended hind leg reaches the snout; total length more than one and three-fourths times the length of tibia; heels overlapping; skin above obscurely shagreened with numerous small rounded tubercles on the back mostly irregularly placed, except those on the shoulder, which form a V-shaped figure with the apex pointing forward; a narrow



FIGS. 85-89.—*RANA JAPONICA*. $1\frac{1}{2} \times$ NAT. SIZE. 85, TOP OF HEAD; 86, SIDE OF HEAD; 87, OPEN MOUTH; 88, UNDERSIDE OF HAND; 89, UNDERSIDE OF FOOT. No. 23549, U.S.N.M.

dorso-lateral glandular ridge on each side from eye to insertion of hind leg, the shortest distance between the two being considerably greater than the distance between the anterior angles of the eyes; underside smooth, except posterior aspect of femurs, which is coarsely granular; a small vocal sac below each corner of the mouth. Color (in alcohol) pale clay-color (Ridgway, Nomencl. Col., pl. v, fig. 8) the dorso-lateral ridge but slightly paler; the latter is marked on the outside by a series of black spots almost forming a continuous line; all the tubercles on the back and sides including the interscapular V marked with a black spot; a blackish brown line from tip of snout through nostril to eye, and a large patch of the same color behind the eye descending obliquely toward the insertion of the fore limb and covering the tympanum; a

blackish line on the anterior face of the upper arm and dusky cross-bars on the hind legs with round blackish spots on the light interspaces; underside uniform whitish.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent	60
Width of head	19
Fore leg	32
Hind leg, tip of longest toe to vent	107
Tibia	34

The *adult female* (U.S.N.M. No. 23548, same locality, date, and collector) is essentially like the male described above, except that the body is broader and the dorso-lateral ridges consequently farther apart, the fore legs less robust, and the swelling on the first finger, which is characteristic of the male sex, absent; the web between the toes is also somewhat more excised. The color is more reddish brown, with scarcely any black spots on the back except those marking the inverted V on the shoulder. Both upper and lower lips are blotched with dark brown.

The *young* specimens are mostly perfectly smooth above, without any tubercles between the dorso-lateral folds.

Variation.—This species offers a great amount of individual variation. The most variable character is the length of the hind legs, for, while in most specimens the tibio-tarsal articulation reaches just beyond the snout, in a few it falls short, while in others it extends considerably farther. The longest hind legs in our collection are possessed by an adult female (No. 31828) collected by Dr. Hugh M. Smith at Yamagawa, Satsuma, Kiusiu, but other specimens in the same lot (for instance, No. 31829) can be exactly matched in every respect by the general run of specimens of corresponding size from central Japan (for instance, No. 34465). The situation of the nostrils relative to the tip of the snout and the correlated length and pointedness of the snout also vary considerably, but in no case have I been able to trace any correlation with length of hind legs or geographic distribution. The color varies greatly; the ground color with regard to the amount and tinge of the brownish or yellowish admixture, the dark spots and bands with regard to intensity and number. Spots are frequently absent, and the male described is rather unusually dotted with black above. The underside is not always uniform white; in many specimens there are brown spots and mottlings on throat or even on chest, but no connection with distribution or structural characters can be discovered.

There is a young specimen in the collection of the Philadelphia Academy of Sciences (No. 15257), collected by Dr. W. H. Jones at Chemulpo, Korea, August, 1890, which appears even more slender and long-snouted than the Japanese specimens, but this may be due

to its somewhat dried-up condition. The web between the toes is very deeply excised, however, more so than in any Japanese specimen examined by me.

Habitat.—*Rana japonica*, besides inhabiting Japan, is said to occur also in eastern China. Specimens are on record from Szechuen, Ningpo, the neighborhood of Peking, Kiukiang, and Shanghai, and I have already referred to the specimen from Chemulpo, Korea, in the Philadelphia Academy.

The specimens in the Leiden Museum collected by von Siebold and Buerger (No. 1930) are only labeled as from "Japan." In Museum Senckenbergianum there are specimens from Nikko and Yokohama, in Berlin from Tokyo, in Hamburg from Yamato and Kawachi. In the United States National Museum there are numerous specimens from Yamagawa, Province of Satsuma, Kiusiu; from Shikoku, Yokohama, and from Mount Fuji. Okada has recorded it from the island of Awaji and from the Province of Suwo, Hondo.

The Senckenberg Museum, in Frankfurt on the Main, in 1889 received from the late Mr. B. Schmacker, of Shanghai, several specimens said to have come from Yezo. One of these is now in British Museum (No. 91. 8. 26. 16), where I have had the opportunity of examining it. It is identical with Yokohama specimens and, if the locality is correct, then both *R. japonica* and *R. temporaria* occur in Yezo. It would be very important to have this question thoroughly investigated. The species has not been found there by others and, as it is not claimed that Mr. Schmacker collected the specimens in Yezo personally, confirmation is necessary.^a

Doctor Boettger has referred two half-grown specimens which were collected by Mr. Schmacker's Japanese collector, either in "Ohoshima or Okinawa," to this species, the only record for the Riu Kius. One might be tempted to regard these as possibly young *Rana holsti* were it not for certain discrepancies in the description, Boettger's specimens having a distinct outer metatarsal tubercle and the tibio-tarsal articulation extending far beyond the tip of the snout. Under these circumstances they are not likely to be from Okinawa or Ohoshima

List of specimens of Rana japonica.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U. S. N. M.	11336	Halfgr.	Japan	1878	E. S. Morse.
Do.	11341do.....do.....	1878	Do.
Do.	11342	Young.do.....	1878	Do.
Do.	11343do.....do.....	1878	Do.
Do.	11344do.....do.....	1878	Do.

^a It is to be noted that several other species from the same source are under similar suspicion, as, for instance, *Eumeces laticutatus* (p. 199) and *Natrix tigrina* (p. 139).

List of specimens of Rana japonica—Continued.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	23548	Female	Yokohama, Hondo	Sept., 1896	L. Steinger.
Do.	23549	Male ^b	do	do	Do.
Do.	23550	Male	do	do	Do.
Do.	23551	do	do	do	Do.
Do.	23552	Female	do	do	Do.
Do.	23553	do	do	do	Do.
Do.	23554	do	do	do	Do.
Do.	23555	do	do	do	Do.
Do.	23556	do	do	do	Do.
Do.	23557	Male	do	do	Do.
Do.	23558	Female	do	do	Do.
Do.	23559	do	do	do	Do.
Do.	23560	do	do	do	Do.
Do.	23561	Male	do	do	Do.
Do.	23562	do	do	do	Do.
Do.	23563	do	do	do	Do.
Do.	23564	do	do	do	Do.
Do.	23565	do	do	do	Do.
Do.	23566	do	do	do	Do.
Do.	23567	Female	do	do	Do.
Do.	23568	do	do	do	Do.
Do.	23569	Male	do	do	Do.
Do.	23570	do	do	do	Do.
Do.	23571	do	do	do	Do.
Do.	23572	Female	do	do	Do.
Do.	23573	do	do	do	Do.
Do.	23574	Male	do	do	Do.
Do.	23575	do	do	do	Do.
Do.	31828	Female	Yamagawa, Satsuma, Kiusiu.	June 14, 1903	H. M. Smith.
Do.	31829	Adult ^c	do	do	Do.
Do.	31830	Young	do	do	Do.
Do.	31831	do	do	do	Do.
Do.	31907	Male	Kochi, Tosa, Shikoku.	May 11, 1903	Do.
Do.	31908	Female	do	do	Do.
Do.	31909	do	do	do	Do.
Do.	31910	do	do	do	Do.
Do.	31911	Halfgr.	do	do	Do.
Do.	31912	do	do	do	Do.
Do.	31922	Young	do	do	Do.
Do.	31923	do	do	do	Do.
Do.	31924	do	do	do	Do.
Do.	31925	do	do	do	Do.
Do.	31931	do	do	do	Do.
Do.	34448	Female	Mount Fuji, Hondo	May, 1908	A. Owston.
Do.	34449	do	do	do	Do.
Do.	34450	do	do	do	Do.
Do.	34451	Male	do	do	Do.
Do.	34452	do	do	do	Do.
Do.	34453	do	do	do	Do.
Do.	34454	Halfgr.	do	do	Do.
Do.	34455	do	do	do	Do.
Do.	34456	do	do	do	Do.

^a Description, p. 110.

^b Description, p. 108, figs. 85-89.

^c Referred to, p. 110.

List of specimens of *Rana japonica*—Continued.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	34457	Halfgr.	Mount Fuji, Hondo	May, 1898	A. Owston.
Do.	34458	do	do	do	Do.
Do.	34459	do	do	do	Do.
Do.	34460	do	do	do	Do.
Do.	34461	do	do	do	Do.
Do.	34462	do	do	do	Do.
Do.	34463	do	do	do	Do.
Do.	34464	do	do	do	Do.
Do.	34465	do	do	do	Do.
Do.	34466	do	do	do	Do.
Do.	34467	do	do	do	Do.
Phila. Acad.	15275	Young ^a	Chemulpo, Korea		W. H. Jones.
Leiden	1930		Japan		Siebold and Buerger.
Brit. Mus.	44. 2. 22. 10		do		Do.
Do.		(b)	Tokyo		von Martens.
Do.	91. 8. 26. 16	(c)	Yezo[?]		B. Schmacker.
Hamburg.	1005		Yamato	1896	T. Lenz.
Do.	1006		Kawachi	1896	Do.

^a P. 110.^b Cotype of *R. martensi*.^c P. 111.RANA TEMPORARIA^a Linnæus.


1758. *Rana temporaria* LINNÆUS, Syst. Nat., 10 ed., I, p. 212 (type-locality, Europe); 12 ed., 1766, I, p. 357.—MIDDENDORFF, Sibir. Reise, II, Pt. 2, 1853, p. 247, pl. xxvi, figs. 2-4 (River Aldan to Udskoi Ostrog).—DOBROROVSKI, Izvest. Sibirsk. Otd. Geograf. Obstch., I, 1870 (p. 23) (southern Sakhalin).—BOULENGER, Cat. Batr. Brit. Mus., 1882, p. 44 (Europe; Abrek Bay, Amur Prov.); Bull. Soc. Zool. France, 1886, p. 594 (Yezo; Kasakevitch and Abrek Bay, Amur Prov.; Wiloni river; Mongolia); Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 140 (Amurland; Yezo); Taill. Batr. Europe, II, 1898, p. 301, pls. xx-xxi.—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, LX, Prilozh., no. 5, 1889, p. 292 (Sakhalin); Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 348 (Russia to Amurland, Sakhalin, etc.).
1768. *Rana muta* LAURENTI, Synops. Rept., p. 30.
1788. *Rana rufa* LACÉPÈDE, Hist. Nat. Quadr. Ovip., I, Syn. meth.
1853. *Rana cruenta* MIDDENDORFF, Sibir. Reise, II, Pt. 2, p. 249, pl. xxvi, figs. 5-7 (southern coast of Okhotsk Sea) (not of Pallas?).—МАЛАК, Putesh. na Amur, 1859 (p. 153) (Amurland).
1855. *Rana fusca* THOMAS, Ann. Sci. Nat. (4), IV (p. 365).
1876. *Rana dybowskii* GUENTHER, Ann. Mag. Nat. Hist. (4), XVII, May, 1876, p. 387 (type-locality, Abrek Bay, near Vladivostok; type in Brit. Mus.; B. Dybowski, collector).—STONE, Proc. Phila. Acad., 1899, p. 183 (Khin-gan Mountains, eastern Mongolia).
1902. *Rana muta johansenii* KASTCHENKO, Izvest. Tomsk Univers., 1902 (p. 24).

^a Signifying temporary, in the present case, perhaps, in the sense of changeable. Application somewhat uncertain, but possibly meant for *temporalis*, with reference to the blackish temporal spot.

Various synonyms referable only to some European form or phase are here omitted.

Rana temporaria, as compared with *R. japonica*, is a much more short-legged and short-snouted species with the dorso-lateral folds apparently wider. Altogether it is larger and stouter, the body especially between head and pelvis being relatively longer. Thus in my specimens of *R. temporaria* from Yezo, the length of the foot, measured from inner metatarsal tubercle to tip of longest toes, exactly equals one-half the distance from anterior border of eye to vent, while in *R. japonica* the length of the foot, similarly measured, is about two-thirds the same distance.

I have compared our three Japanese specimens from Yezo with specimens of corresponding sex and size from Norway (for instance, No. 25339, Bergen) and find no tangible differences. The web



FIGS. 90-93. — *RANA TEMPORARIA*. $1\frac{1}{2} \times$ NAT. SIZE. 90, TOP OF HEAD; 91, SIDE OF HEAD; 92, OPEN MOUTH; 93, UNDERSIDE OF FOOT. No. 23440, U.S.N.M.

between the toes is perhaps a trifle more excised, and the inner metatarsal tubercle a trifle stronger, but these differences are clearly within the range of individual variation.

Description.—*Adult female*: U.S.N.M. No. 23440; Sapporo, Yezo; S. Nozawa, collector (figs. 90-93). Vomerine teeth in two slightly oblique groups, the anterior border of which extends forward slightly beyond a line through the posterior border of the choanæ; snout short, rounded, the distance from orbit to tip of snout equaling the distance between black stripes at anterior border of orbit; nostrils nearer the tip of snout than the eye; interorbital space as wide as the upper eyelid and a little narrower than distance between the nostrils; tympanum vertically oval, its greatest diameter about three-fourths the diameter of the eye; distance of tympanum from eye less than

one-half the former's greatest diameter; fingers blunt, first finger extending slightly beyond second; incision of web on both sides of fourth toe a trifle deeper than the distal end of the basal phalanx; inner metatarsal tubercle slightly compressed, its length equaling its distance from subarticular tubercle of first toe; no outer metatarsal tubercle; tibio-tarsal joint of extended hind limb reaches center of eye; tibia less than half the length of head and body; heels overlapping by less than interorbital width; skin above and below smooth, except posterior aspect of femur, which is coarsely granulated; the dorso-lateral glandular folds but slightly raised, their shortest distance contained five times in total length. Color (in alcohol) above brownish gray, slightly grayer on snout, with obscure and ill-defined dark-brown mottlings, which on the back and sides occasionally surround a light spot so as to form ocelli; dorso-lateral folds lighter, externally margined by an ill-defined brown line; indications of a dusky band across the top of the head through the center of the upper eyelids; upper lip mottled with brown; a brown line from tip of snout through nostril to eye expanding behind the latter to form the usual dark-brown ear-patch; brown stripe on anterior face of over arm long and parallel with the axis; limbs cross-banded; underside whitish, the whole surface coarsely marbled with brown.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent.....	56
Width of head.....	17
Fore limb.....	31
Hind limb, vent to tip of longest toe.....	87
Tibia.....	27

Males differ from *females* in the stronger, more muscular fore limbs, which acquire an extraordinary development during the pairing season; in the pad-like swelling of the inner side of the first finger, which becomes covered with black horny spinules during the same season; in the greater development of the web between the toes; and in the presence of a pair of internal vocal sacs, situated at the sides of the throat. ^a

Variation.—Apparently this species is much less variable than *R. japonica*, especially if we consider its great geographical distribution. The presence of more or less numerous glandular warts on the upper surface and sides, so conspicuously absent in the specimen described above, depends on the season, they being best developed during the breeding time, in museum specimens also to some extent on the state of preservation and the preserving fluid employed. The color, however, varies greatly both as to the tint of the ground color and as to the amount and distinctness of the dark markings, some

^a Boulenger, Tailless Batrach. Europe, p. 305.

specimens being nearly uniform above, others densely blotched with blackish.

Habitat.—The typical *R. temporaria* occurs in the more northern portions of the Palearctic region, from the British Islands and Scandinavia in the west, to Japan in the east.

In the St. Petersburg Museum there are numerous specimens from the Amur and Ussuri valleys. The Berlin Museum has a specimen from Kasakevitcha near the junction of the Ussuri and Amur rivers, and British Museum has another from Abrek Bay not far from Vladivostok, the type of *R. dybowskii*. In the Hamburg Museum there are specimens from Khabarovka, Poprovka and Nikolayesk, Amurland, collected in 1894 by Dieckmann. Nikolski reports that this frog is not common in Sakhalin, where he found it in the river Duiki and Dobrotvorski in the southern part of the island.

The occurrence of this species in Yezo, therefore, is only what might be expected. British Museum has five specimens from this island, collected by Dr. John Anderson, and our museum has also three Hokkaido specimens collected by Dr. S. Nozawa near Sapporo. In the St. Petersburg Museum there is a specimen from Nemuro, collected by Grigoriew.

For the possible occurrence of *R. japonica* in Yezo see under the latter species. Here I may emphasize only the fact that the specimens there alluded to certainly belong to this species and not to *R. temporaria*, and the only question is whether the locality is correctly given or not.

List of specimens of Rana temporaria.

Museum,	No.	Sex.	Locality.	When collected.	By whom collected.
U.S.N.M.....	23438	Female ...	Sapporo, Yezo		S. Nozawa.
Do.....	23439	do.....	do.....		Do.
Do.....	23440	do. a.....	do.....		Do.
Brit. Mus.....	86. 6. 30. 18		Yezo.....		J. Anderson.
Do.....	19		do.....		Do.
Do.....	20		do.....		Do.
Do.....	21		do.....		Do.
Do.....	22		do.....		Do.

a Description and figs., p. 114.

RANA TSUSHIMENSIS, a new species.

Diagnosis.—*Rana temporaria* group; vomerine teeth behind level of choanæ; snout short, the distance from orbit to tip of snout not greater than width between black stripes at anterior border of orbit; tibio-tarsal articulation reaching beyond eye or tip of snout; distance between dorso-lateral glandular folds contained five or six times in


a Signifying from Tsushima, in the Straits of Korea, between the south end of Korea and Kiusiu, Japan.

length from snout to vent; inner metatarsal tubercle oval, less than one-half its distance from tip of inner toe; first finger extending beyond second; tibia shorter than fore limb; tympanum about one-half the diameter of eye, from which it is separated by about one-half its own diameter; distance between nostrils considerably greater than interorbital width; webs between toes deeply excised, three terminal phalanges of fourth toe being free.

Type.—U.S.N.M. No. 17519; Tsushima; May, 1885; P. L. Jouy, collector.

Habitat.—Tsushima, Japan.

This new species is probably nearest related to *Rana amurensis*. It shares with it the short, rounded snout, and the deeply excised webs, but the inner metatarsal tubercle is considerably stronger and larger.



FIGS. 94-98.—*RANA TSUSHIMENSIS*. $1\frac{1}{2} \times$ NAT. SIZE. 94, TOP OF HEAD; 95, SIDE OF HEAD; 96, OPEN MOUTH; 97, UNDERSIDE OF HAND; 98, UNDERSIDE OF FOOT. No. 17519, U.S.N.M.

In this respect it agrees with *R. temporaria*, but has more deeply excised webs, narrower interorbital space, narrower dorso-lateral folds, and slightly longer legs, being on the whole of slenderer build. From *R. japonica* it differs by its short, rounded snout and its much shorter hind foot. The vomerine teeth are decidedly further back than in *R. temporaria* and *japonica* though not quite so far as in *R. amurensis*.

Description of type-specimen (figs. 94-98).—Vomerine teeth in two oblique groups their anterior border on a line with the posterior border of the choanæ; snout short, the distance from orbit to tip of snout equals the width between the black stripes at anterior border of orbit; nostrils half-way between eye and tip of snout, their mutual

distance being much greater than their distance from the eye or than the interorbital space, which is narrower than upper eyelid; vertical diameter of tympanum greater than horizontal, a trifle more than one-half the diameter of eye, and more than twice the distance of tympanum from eye; first finger slightly extending beyond second; webs of toes deeply excised, barely reaching the middle of basal phalanx of fourth toe; inner metatarsal tubercle oval, projecting, slightly shorter than its distance from subarticular tubercle of first toe; a strong conical outer metatarsal tubercle; subarticular tubercles strongly developed; tibio-tarsal articulation of extended hind leg reaches almost to the nostril; tibia one-half the total length; heels overlapping by the width of the interorbital space; skin above with numerous small scattered granules, those on the sides larger; dorso-lateral glandular folds very narrow and sharply defined, their shortest distance being contained about five times in the total length; lower surface smooth except the proximal half of the posterior aspect of the femur which is granular. Color (in alcohol) above uniform pale drab, with a few minute dusky spots on the back; the brown stripe from tip of snout to eye, very narrow, and the tympanic dark patch rather pale and ill-defined between eye and tympanum; humeral stripe and crossbars on limbs obscure; underside uniform whitish.

Dimensions.

	<i>mm.</i>
Total length, snout to vent.....	52
Width of head.....	16.5
Fore limb.....	30
Hind limb, tip of longest toe to vent.....	83
Tibia.....	26

The *male* (U.S.N.M. No. 17520; Tsushima; May, 1885; P. L. Jouy, collector) is smaller (total length 37 mm.), with relatively longer tibia (20 mm.). The snout is somewhat shorter, with the nostrils nearer to the tip than to the eye; tympanum slightly larger and nearer the eye; first finger has the swollen pad set with asperities, characteristic of the breeding male; webs less excised, reaching to the anterior third of the basal phalanx of fourth toe; outer metatarsal tubercle barely indicated; tibio-tarsal articulation reaching tip of snout, overlapping by more than interorbital space; shortest distance between dorso-lateral folds one-sixth the total length; lower back and sides with numerous small, white-tipped, pointed tubercles, and in addition on sides large glandular warts. Color like the female, but canthal stripe and ear-patch darker and better defined; underside densely sprinkled with minute brown dots.

Remarks.—Most of the differences between the two specimens as pointed out here are sexual, the relative shorter body, the pad on the first finger, the greater extent of the webs indicating the male. The

most notable difference, however, is the presence of the strongly developed outer metatarsal tubercle in the female specimen.

There are a number of specimens from Tsushima, in British Museum, collected by Holst, and in 1898 I was given the privilege of examining them, but I find no reference to the presence of an outer metatarsal tubercle, which may therefore be regarded as problematical. These specimens were referred partly to *R. japonica* and partly to *R. martensi*. My own examination led me to the conclusion that they belong to one species only, as I could not correlate any two of the various differences. They were all short-snouted, with deeply incised webs and narrow dorso-lateral folds.

Habitat.—Only known from Tsushima, in the channel between Japan and Corea, where specimens were collected by P. L. Jouy, in May, 1885, and later by P. A. Holst.

List of specimens of Rana tsushimensis.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.
U.S.N.M.	17519	Female ^a .	Tsushima, Japan.....	May, 1885	P. L. Jouy.
Do.	17520	Male ^bdo.....do.....	Do.
Brit. Mus.		(c)do.....do.....	P. A. Holst.

^a Type; description, p. 117; figs. 94-98.

^b Description, p. 118.

^c P. 119

RANA AMURENSIS ^a Boulenger.

Plate XI, figs. 2-3.

1853. ? *Rana temporaria* MIDDENDORFF, Sibir. Reise, II, ii, Pt. 1, p. 247, pl. xxvi, fig. 3 *a-c* (Eastern Siberia).

1886. *Rana amurensis* BOULENGER, Bull. Soc. Zool. France, 1886, p. 598; author's separate (p. 4) (type-locality, Kazakevitch, Amur Province; type, Berlin Mus. No. 9864); Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 140, pl. ix, fig. 1 *b* (Lake Khanka, Doerries, collector; Chemulpo, Korea); Proc. Zool. Soc. London, 1890, p. 324.— ? BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 146 (Chinghai, near Ningpo, China).—BEDRIAGA, Przewalski Reise, Zool., III, Pt. 1, 1899, (p. 27).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, No. 1, 1905, p. 369 (Kuku-Nor, Tibet).

Rana amurensis seems to be closely related to true *R. temporaria*, being round-snouted and short-legged. It differs, however, in the more posterior position of the vomerine teeth which are almost entirely behind the level of the choanae and, apparently, in having the inter-orbital space narrower. The inner metatarsal tubercle is also unusually weak. As the two forms occur in the same locality, the Berlin Museum having also true *R. temporaria* (No. 9865) from Kazakevitch, whence came the type of *R. amurensis*, there seems to be no doubt as to the specific validity of the latter.

Having no specimen at command, though it was my good fortune to be allowed to examine the Ussuri and Korean specimens in the Brit-

^a Signifying from Amurland.

^b Reproduced in this work on Plate XI, figs. 2-3.

ish Museum, I print herewith a translation of Boulenger's original description.

Original description of type specimen.—*Adult male*; Berlin Museum, No. 9864; Kazakevitch, Amur Province. Vomerine teeth in two oval groups forming a chevron behind the opening of the inner nares; head rather depressed, as long as broad, or a little longer than broad; snout long and rounded, projecting but little beyond the lip; loreal region rather abruptly bent; nostril halfway between the end of the snout and the anterior corner of the eye; width of interorbital space equaling that of the upper eyelid; tympanum measuring two-thirds the diameter of the eye, separated from the latter by a space equal to one-half of its diameter; first finger not extending beyond the second when placed alongside of it; tibio-tarsal articulation reaching eye when the hind limb is carried forward; tibia much shorter than the fore limb; inner metatarsal tubercle small, oval, blunt; no outer tubercle; subarticular digital tubercles small; toes scarcely two-thirds webbed; back smooth; flanks, lower abdomen, and posterior aspect of the femurs with very large granulations; dorso-lateral folds narrow and but little prominent. Grayish brown above with the ordinary black spots; a blackish band bordered below by whitish along the lip, as in *R. arvalis*; a pale vertebral band rather indistinct; humeral spot very elongated; lower parts spotted with grayish. Male without vocal sacs.

Dimensions.

	<i>mm.</i>
From snout to vent.....	55
Length of head.....	18
Width of head.....	17
Diameter of eye.....	5.5
Interorbital space.....	3
Anterior corner of eye to nostril.....	3.5
Anterior corner of eye to tip of snout.....	7.5
Diameter of tympanum.....	3.5
Posterior border of eye to anterior border of tympanum ..	1.5
Fore limb.....	34
Hind limb.....	90
Tibia.....	28
Inner toe.....	7
Inner metatarsal tubercle.....	^a 1.5

The changes in the description made by him upon the examination of nine specimens from Lake Khanka, are chiefly verbal and qualifying, the most important being the "interorbital space a little narrower than the upper eyelid," an observation which I can corroborate from my own inspection of the same material in 1898. He also adds that the male is provided with black nuptial excrescences on the thumb.

^a Boulenger, Bull. Soc. Zool. France, 1886, p. 598.

Habitat.—Originally described from two specimens collected at Kazakevitch, not far from Khabarovka, a little above the junction of the Ussuri River with the Amur, this species has since been recorded from Lake Khanka and from Chemulpo, Korea. In the Hamburg Museum there are two specimens from Nikolayevsk, Amurland, collected by Dieckmann in 1894 (Nos. 873, 875).

Whether the grown specimens which Boettger (Ber. Senckenberg. Naturf. Ges., 1894, pp. 146–147) records from Chinhai, near Ningpo, China, under the name *Rana amurensis* really belong here is rather doubtful, as he himself does not seem to be quite satisfied with their identity.

Nikolski has recently (1905) identified seven specimens in the St. Petersburg Museum collected at Lake Kuku-Nor, northeastern Tibet, by Przevalski and Grum-Grzymailo as belonging to this species.

List of specimens of Rana amurensis.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.
Brit. Mus.	(α)	Lake Khanka, Amurland	Doerries.

α Nine specimens.

RANA IJIMÆ^a Stejneger.

1901. *Buergeria ijimæ* STEJNEGER, Proc. Biol. Soc. Washington, XIV, Dec. 12, 1901, p. 190 (type-locality, Okinawa shima; type, Sci. Coll. Mus., Tokyo, No. 19 (914)).


The outward conformation of the terminal digital joint of this species led me to refer it to *Polypedates* (or *Buergeria*). I have since examined more closely into the matter and can find no interpolated bone, so that its proper place seems to be in the genus *Rana*.

The exact nature of the dorso-lateral gland in the unique type specimen is doubtful. The specimen is somewhat soft and the light-colored dorso-lateral lines appear to be glandular and somewhat raised, but whether we have to do with a continuous fold or only a series of longitudinal glands is not certain, although the former condition is more probable.

Description.—*Adult*.—Science College Museum, Tokyo, No. 19 (914); Tanebimura, Okinawa shima, Riu Kiu (figs. 99–103). Vomerine teeth in two slightly oblique series on a line through the posterior border of the choanæ and about equidistant from the latter and from each other; tongue without free, conical papilla; snout somewhat projecting, the nostrils much nearer the tip of the snout than the eyes and nearly vertical over the tip of the mandible; interorbital space

^a Named in honor of Dr. Isao Ijima, professor of zoology in the Imperial University, Tokyo.

slightly narrower than the upper eyelids; canthus rostralis rather abruptly bent; lores concave; tympanum one-half the diameter of the eye; fingers free, first extending slightly beyond second; disks distinct, but small, on third and fourth finger largest, less than half the diameter of tympanum; toes moderately webbed, the two terminal phalanges of fourth toe being free and the excision reaching to the terminal third of the basal phalanx of that toe; disks well developed but small, slightly smaller than those of the fingers; sub-articular tubercles very prominent; inner metatarsal tubercle oval, not prominent, small, contained about two and a half times in length



FIGS. 99-103.—*RANA IJIMÆ*. NAT. SIZE. 99, TOP OF HEAD; 100, SIDE OF HEAD; 101, OPEN MOUTH; 102, UNDERSIDE OF HAND; 103, UNDERSIDE OF FOOT. No. 19 (914) SCI. COLL. TOKYO.

of first toe, measured from the anterior border of the tubercle; a distinct outer metatarsal tubercle at the base of fourth toe; no outer dermal fringe on fifth toe, nor a tarsal fold; tibio-tarsal articulations reach considerably beyond tip of snout when hind legs are stretched forward, and overlap by as much as the length of the snout when folded legs are placed at right angles to axis of body; tibia more than one-half the total length of head and body; skin above obscurely shagreened, with a few scattered pustules on the back and numerous ones on the flanks; sides of face and neck and upper side of tibia with numerous white-tipped, pointed asperities, which also stud the

tympanum like a string of pearls, as well as three large glandular warts behind the corner of the mouth; from the posterior angle of the eye a somewhat indistinct dorso-lateral fold, which on the shoulder seems broken up into a series of shorter longitudinal glands; under-side smooth, except posterior aspect of femur, which is granular. Color (in alcohol) dark chocolate brown, apparently nearly uniform, the dorso-lateral folds and upper side of limbs paler, more cinnamon; a whitish line under eye and on posterior part of upper lip, which is mottled with brown along the edge; flanks pale brownish, with irregular blackish spots; legs cross-banded with dark brown and upper posterior aspect of femur mottled with the same color; under-side whitish, with very faint brown mottlings on throat, chest, and hind legs.

Dimensions.

	mm.
Total length, tip of snout to vent.....	68
Width of head.....	23
Distance between nostrils.....	7
Distance between nostril and eye.....	5
Diameter of eye.....	9
Diameter of tympanum.....	4.5
Interorbital space.....	5
Fore leg.....	44
Largest finger disk.....	2
Hind leg, vent to tip of longest toe.....	118
Tibia.....	38
Inner metatarsal tubercle.....	2.25

Habitat.—Thus far this species is only known from Okinawa shima, where the type specimen was collected.

List of specimens of Rana ijima.

Museum.	No.	Age.	Locality	When collected.	By whom collected.
Sci. Coll. Tokyo.	19:914	Adult σ .	Tanabimura, Okinawa shima.		

σ Type; description, p. 121; figs. 99-103.

RANA RUGOSA σ Schlegel.


Plate X, figs. 2-4.

1838. *Rana rugosa* SCHLEGEL, Fauna Japon., Rept., p. 110, Sauri et Batr., pl. III, figs. 3, 4^b (type-locality, Japan, probably Nagasaki; types, Leiden Museum No. 2064; von Siebold, collector).—DUMÉRIL and BIBRON, *Erpét. Gén.*, VIII, 1841, p. 368 (Japan).—GUENTHER, *Cat. Batr. Sal. Brit. Mus.*, 1858, p. 11 (Japan).—BLEEKER, *Natuurk. Tijdschr. Nederland Indië*, XVI, 1858, p. 204 (Japan).—HALLOWELL, *Proc. Phila. Acad.*, 1860, p. 499 (part: Simoda).—MARTENS, *Preuss. Exped. Ost-Asien, Zool.*, I, 1866, p. 111 (Yokohama); 1876, p. 384 (Nagasaki).—CAMERANO, *Atti. Accad. Torino*, XIV,

σ Latin, signifying wrinkled. b Reproduced in this work on Plate X, figs. 2-4.

Pt. 5, Apr. 1879, p. 868 (Tokyo, Yokohama).—HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 119 (Yokohama; Hakone Mts.).—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 35 (Japan).—OKADA, Cat. Vert. Jap., 1891, p. 67 (Tokyo; Suwo).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, 1891, p. 239 (Yezo).—BOETTGER, Kat. Batr. Mus. Senckenberg., 1892, p. 4 (Hakone Lake; Nikko; Yokohama).—IKADA, Annot. Zool. Japon., I, Pt. 3, Aug. 1897, p. 114 (breeding season).

Hallowell's *R. rugosa*^a belongs only partly to this species, being evidently composed of two species. The small ones "marked 13 in the catalogue . . . found along the mountain streams at Simoda, Japan, May, 1855," which he describes as "presenting the conical pustulations upon the longitudinal elevations upon the back," are in all probability true *R. rugosa*,^b but the others from "Ousima," and the two larger ones from Simoda, of which the last-mentioned ones are in our collection (U.S.N.M. No. 7415) belong to *R. nigromaculata*. This blunder of Hallowell's is the more remarkable, as *R. rugosa* is one of the most easily recognized species, because of its



FIGS. 104-106.—*RANA RUGOSA*. $1\frac{1}{2} \times$ NAT. SIZE. 104, TOP OF HEAD; 105, SIDE OF HEAD; 106, OPEN MOUTH. No. 23543, U.S.N.M.

excessive roughness, which makes it appear more like a toad than a frog. The hollow snout is also quite characteristic.

Description.—*Adult female*; U.S.N.M. No. 23543; Yokohama, Hondo; September, 1896; L. Stejneger, collector (figs. 104-106). Vomerine teeth in two small groups, their anterior border on a line through the center of the choanæ and more distant from the latter than from each other; nostrils much nearer the tip of snout than the eye; snout and lores concave, the canthus rostralis forming a ridge separating the two concavities; interorbital space less than width of upper eyelid; tympanum nearly circular, fully three-fourths the diameter of the eye distant from the latter less than one-half its own diameter; fingers not tapering, rounded at end, first somewhat opposable and longer than second, with a small swollen pad at the base; webs of toes full, the excision opposite the distal end of the first phalanx of fourth toe; inner metatarsal tubercle small and rather

^a Proc. Phila. Acad., 1860, p. 499.

^b They are not in the U. S. National Museum.

weak; a weak outer metatarsal tubercle; subarticular tubercles rather weak, the distal one triangular, with the base forward; a decided tarsal fold; tibio-tarsal articulations reach center of tympanum when hind legs are carried forward, and do not overlap when legs are bent at right angles to axis of body; tibia less than one-half the total length; skin above and on sides densely granular, with prominent tubercles in addition to a number of irregular series of short longitudinal ridges, which are largest on the middle of the back, smallest on the head, all these ridges tubercular like the rest of the skin; upper eyelids coarsely tubercular, as also the loreal region; even the tympanum is studded with smaller tubercles; no dorso-lateral fold; a number of prominent tubercular glands behind tympanum; underside smooth, but transversely creased, throat and posterior aspect of femurs very coarsely granular. Color (in alcohol) dark brownish olive, with obscure dusky spots and dark cross bands on legs; underneath pale, uniformly and densely mottled with dusky.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent.....	56
Width of head.....	22.5
Fore leg.....	30
Hind leg, vent to tip of longest toe.....	83
Tibia.....	25

The adult male in the breeding season (U.S.N.M. No. 34436; Mount Fuji, Hondo; May, 1898) is smaller, with relatively shorter body, somewhat fuller webs, and a large swelling at the base of the first finger covered with asperities, though not so large as in the species previously described.

There are no vocal sacs, internal or external. Boulenger^a says: "Male with external vocal vesicles," but as there were then no male specimens in the British Museum he probably had this information from Hallowell, who attributes large vocal vesicles to this species. As I have already shown, the specimens with such, which Hallowell took for *R. rugosa*, were in reality *R. nigromaculata*.

Variation.—On the whole, this species is very uniform, the greater or lesser distinctness of the blackish markings mostly due to a varying shade of the ground color being the chief source of variation. Hilgendorf mentions that one of his specimens (Berlin Mus. No. 4312) lacks the vomerine teeth.

Habitat.—This very distinct species appears to be confined to Japan proper,^b i. e., to the islands of Hondo, Kiusiu, and Shikoku. Doctor

^a Cat. Batr. Sal. Brit. Mus., p. 35.

^b Von Martens (Preuss. Exped. Ost-Asien, Zool., I, 1876, p. 384) has recorded it from Hongkong, China. The identification, which seems to be by Peters, is probably correct, but the locality is most likely erroneous. Hallowell records two specimens from Ooshima, but they were probably *Rana nigromaculata*, like the male specimens from Simoda, which he also identified as *R. rugosa*.

Fritze is responsible for the statement that it is common in Yezo, but as he is the only writer who records it from that island it may be well to await the arrival of undoubted Yezo specimens. I am not aware that it has been found on Tsushima nor in Korea or any of the Riukiu islands.

The following definite localities from where we have reliable records or have seen specimens may be mentioned: Mountain streams at Simoda (Hallowell); mountains of Hakone (Hilgendorf); Hakone Lake and Nikko (Boettger); Yokohama (von Martens; Hilgendorf; Camerano; Boettger; Stejneger); Tokyo (Camerano); Mount Fuji (Owston); Kobe and Enoshima (Brit. Mus.); Seta, near south end of Lake Biwa, Province of Omi, collected by Dr. T. Lenz (Hamburg Mus. No. 1052); Province of Suwo, Hondo (Okada); Nagasaki (Dr. E. A. Mearns; von Martens). Dr. Hugh M. Smith obtained specimens at Koriyama, Kiusiu, and Kochi, Shikoku; also at Nara, Province of Yamato, Hondo.

The types in the Leiden Museum, which were collected by von Siebold, are only inscribed "Japon."

List of specimens of Rana rugosa.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected.
11335		Japan.....	1878	E. S. Morse.
11337		do.....	1878	Do.
11338		do.....	1878	Do.
11339		do.....	1878	Do.
11340		do.....	1878	Do.
22190		do.....		(?)
22191		do.....		(?)
23543	Female ^a	Yokohama, Hondo.....	Sept. 1896	L. Stejneger.
23544	Adult	do.....	do.....	Do.
23545	do	do.....	do.....	Do.
23546	do	do.....	do.....	Do.
23547	do	do.....	do.....	Do.
23587	do	do.....	do.....	Do.
26044	do	do.....	do.....	Do.
31801	Female	Nara, Yamato, Hondo.....	1903	H. M. Smith.
31802	do	do.....	1903	Do.
31813	do	Koriyama, Yamato, Hondo.....	Apr. 26, 1903	Do.
31814	do	do.....	do.....	Do.
31930	do	Kochi, Toza, Shikoku.....	May 11, 1903	Do.
31932	do	do.....	do.....	Do.
34433	do	Mount Fuji, Hondo.....	May 1898	A. Owston.
34434	do	do.....	do.....	Do.
34435	do	do.....	do.....	Do.
34436	Male ^b	do.....	do.....	Do.
34437	Female	do.....	do.....	Do.
34438	Male	do.....	do.....	Do.
34439	do	do.....	do.....	Do.
34440	do	do.....	do.....	Do.
34441	Female	do.....	do.....	Do.

^a Description, p. 124, figs. 104-106.

^b Description, p. 125.

List of specimens of *Rana rugosa*—Continued.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected.
3442	Female	Mount Fuji, Iiondo	May, 1898	A. Owston.
3443	do	do	do	Do.
3444	do	do	do	Do.
3445	Male	do	do	Do.
3446	Female	do	do	Do.
3447	Halfgrown	do	do	Do.
35275	Adult	Nagasaki, Kiusiu		E. A. Mearns.
35276	do	do		Do.
35277	do	do		Do.
35278	do	do		Do.
35279	do	do		Do.
35280	do	do		Do.
35281	do	do		Do.
35282	do	do		Do.
35283	do	do		Do.
35284	Halfgrown	do		Do.
35285	Young	do		Do.
35286	do	do		Do.

RANA LIMNOCHARIS^a Wiegmann.


1835. *Rana limnocharis* WEIGMANN, Nova Acta Acad. Leop. Carol., XVII, Pt. 1, p. 255 (no type-locality; apparently based on specimens in the Leiden Museum from Java, collected and named by H. Boie).—BOULENGER, Fauna Brit. India, Rept., 1890, p. 450 (China and southern Japan to India and Malay Archipelago).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 865; author's separate, p. 16 (Okinawa shima).—BOETTGER, Offenbach. Ver. Naturk. 33–36 Ber., 1895, p. 103 ("Ohoshima or Okinawa").—STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 217 (Taipa and Giilan, Formosa; Botel Tobago Il.; Pescadores IIs).
1835. *Rana gracilis* WIEGMANN, Nova Acta Acad. Leop. Carol., XVII, Pt. 1, p. 257 (type-locality, Cape Syngmore, China; Meyen, collector) (not of Gravenhorst, 1829).—HALLOWELL, Proc. Phila. Acad., 1860, p. 505 (Whampoa, China).—GUENTHER, Rept. Brit. India, 1864, p. 409 (Ningpo; Hongkong; Riu Kiu; Himalayas; Siam; Madras).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1876, p. 383 (Tamsui, Formosa).—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 28 (East Indies; Chusan, Ningpo, Shanghai, Szechuen, Hongkong, Hainan, China; Formosa; Riu Kiu); Proc. Zool. Soc. London, 1887, p. 149 (Riu Kiu, Pryer, coll.).—BOETTGER, Offenbach. Ver. Naturk. 24–25 Ber., 1885, p. 159 (Japan).—OKADA, Cat. Vert. Japan, 1891, p. 67 (Okinawa shima).—BROWN, Proc. Phila. Acad., 1902, June 11, p. 185 (Riu Kiu, probably Okinawa shima, Furness and Hiller, coll., 1896).
1858. *Rana vittigera* GUENTHER, Cat. Batr. Sal. Brit. Mus., p. 9 (Ningpo; Hongkong; Riu Kiu; Ceylon; India; Java) (not of Wiegmann, 1835).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1866, p. 163 (Tamsui, Formosa).

Several names based on Indian specimens and usually referred to this species are omitted from this synonymy.

^a From *λίμνη*, swamp, pond; *χάρις*, grace, one of the Graces.

Boulenger^a regards this species as closely allied to *R. tigerina*, differing in smaller size, half-webbed toes, slight development of the fringe on the fifth toe and usually in the presence of a small outer metatarsal tubercle. One of the chief differences I find, however, is in the situation of the vomerine teeth. In *R. tigerina* they are anterior, approximated very closely to the choanæ, while in *R. limnocharis* they are posterior and quite distant from the choanæ, their distance from the latter being twice as great as their distance from each other.

Description (figs. 107-110).—*Adult female*; U.S.N.M. No. 31798; Onomichi, Prov. Aki, Hondo; June 4, 1903; Dr. H. M. Smith, collector. Vomerine teeth in two oblique groups, the anterior border of which is on a line with the posterior border of the choanæ, their distance from the latter twice as great as their distance from each other posteriorly; nostrils a little nearer the tip of the snout than the eyes; interorbital space about two-thirds the width of upper eyelid; snout convex in cross-section, canthus rostralis rounded; tympanum



FIGS. 107-110.—*RANA LIMNOCHARIS*. NAT. SIZE. 107, TOP OF HEAD; 108, SIDE OF HEAD; 10., UNDERSIDE OF HAND; 110, UNDERSIDE OF FOOT. No. 36501, U.S.N.M.

circular about one-half the diameter of the eye, its distance from the latter more than half of its own diameter; fingers slightly tapering, rounded, first extending much beyond second, and even beyond fourth; a small smooth swelling at base of first finger; web between toes deeply excised to the middle of the basal phalanx of the fourth toe; inner metatarsal tubercle strong, oval, longer than one-half the first toe; a strong conical outer metatarsal tubercle; a narrow dermal fold along the outer side of fifth toe; a fold from inner metatarsal tubercle halfway up the tarsus; tibio-tarsal articulations reach the tympanum when hind legs are stretched forward, and only touch, without overlapping when legs are bent at right angles to axis of body; skin above finely shagreened, with numerous narrow, elongated glandular ridges which become shorter on the sides; no dorso-lateral fold; a deep crease on the chest in front of each arm joining another crease which runs from axilla to axilla across the chest; skin under-

^a Cat. Batr. Sal. Brit. Mus., p. 28, and Fauna Brit. India, Rept., p. 450.

neath nearly smooth becoming coarsely granular posteriorly on the sides, lower belly and posterior half of femurs. Color (in alcohol) above dark reddish gray, with numerous irregular, light-edged dusky blotches; a dark chevron across the head at the center of the upper eyelids, the angle pointing backwards, and a straight dark line across the head at the anterior corner of the eyes; a narrow dark line over canthus rostralis and upper part of tympanum; several broad dark vertical marks on the upper lip and crossing to the lower, one near the tip of the snout, one below the nostrils, two below the eye; some irregular black marks on white ground in the groin; upper posterior aspect of femur marbled black and white in strong contrast; legs cross-banded; underside whitish, throat and chest densely sprinkled with dark gray.

Dimensions.

	<i>mm.</i>
Total length of snout to vent	51
Width of head	18
Fore leg	22
Hind leg, vent to tip of longest toe	68
Tibia	22

The *adult male* (U.S.N.M. No. 23902; Kumamoto, province of Higo, Kiusiu) is smaller (38 mm.), but does not seem to differ materially in proportions; the first finger is provided with copulatory asperities which cover two separate, slightly swollen pads, one basally on the side and below, the other distally and on the upper side of the metacarpal, no asperities on the phalanges; the excision of the web is not quite so deep as in the female; a pair of external vocal sacs on the side of the throat, the openings being behind the tongue near the corner of the mouth. A broad gray band across the throat involving the vocal sacs. A very narrow white line on the upper side along the whole median line is also shown in this specimen, but this character is not sexual.

Variation.—No remarkable variation is found among the Japanese specimens as regards structural characters or proportions. There is some difference in the amount of roughness of the skin on the upper surface, but this is probably due to some extent, at least, to the different state of preservation; one young specimen (No. 30744) from Miyazaki, Kiusiu, is so rough, however, even to the inclusion of the upper eyelids, that it might easily be mistaken for *R. rugosa* were it not for the convex snout, the narrow interorbital space, deeply excised webs, etc. In color there is greater divergence, though in the specimens from Japan proper this is chiefly confined to the presence of a narrow median white line on the back in a few specimens; none of them have the wide pale median line found in some Formosan specimens. This narrow white line is present in all the Ishigaki

Island specimens before me, and in two it is almost as broad as in those from Formosa. In the latter all three styles of coloration are present, irrespective of locality or sex.

Habitat.—This species is distributed over a wide area in southeastern Asia from Japan and China, and according to Boulenger extending throughout India, Ceylon, and Burma to the Malay peninsula and archipelago, ascending to an altitude of 7,000 feet in the Himalayas. Over this range it is probably represented by several subspecies.

It is the commonest species of frog in Formosa and adjacent islands. Von Martens reported it from Tamsui, long ago, and I have recently recorded it from Taipa and Giilan, where this species was collected by Mr. Tada in September and October, 1896. Numerous species collected by him in the Pescadores and in Botel Tobago, April and May, 1897, were reported upon at the same time.

Its occurrence in the Riu Kius was first known through a specimen in British Museum presented by Mr. A. Collie. It was obtained later by Pryer's collectors, and Doctor Fritze found it very common on Okinawa shima. From this island our museum also possesses a specimen (No. 36552) obtained from the Science College Museum in Tokyo. In the Hamburg Museum there are specimens (Nos. 530-531) collected by Doctor Warburg in Myako shima and Iriomote shima, while the U. S. National Museum has a large series from Ishigaki shima, all of the Saki shima group.

It was unknown to the authors of *Fauna Japonica*, but has since been attributed to Japan in a general way, though with no authentic specimens or definite locality to prove it. We have now in the U. S. National Museum numerous specimens from Kiusiu, two collected at Kumamoto, Province of Higo, on the west side, and one from Miyazaki, Province of Iiuga, on the east side. Doctor Smith brought us two specimens from Yamagawa, Satsuma, and five from Hondo, viz, one from Onomichi, in Aki, and four from Koriyama, in Yamato. Dr. E. A. Mearns obtained several young specimens above Nagasaki. Dr. O. Nordquist, of the Vega Expedition, also obtained it in Nagasaki during October, 1879, and at Hirosami on the 19th of the same month (Mus. Stockholm, Nos. 1034-1035 and 158-165). It almost looks as if this species is extending its range in Japan rapidly, and the inquiry seems legitimate whether it may not owe its presence in Japan proper to a comparatively recent introduction. It seems almost incredible that von Siebold should have missed this species at Nagasaki, if it had really existed there during the time of his residence near that city.

List of specimens of *Rana limnocharis*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	22192	Adult	Japan		(?)
Do.	22193	do.	do.		(?)
Do.	23302	Male ^a	Kumamoto, Higo, Kiusiu	1884	I. Ijima.
Do.	23303	Female	do.	1884	Do.
Do.	30744	Young	Miyazaki, Kiusiu		C. A. Clark.
Do.	31823	Female	Yamagawa, Satsuma, Kiusiu	June 14, 1903	H. M. Smith.
Do.	31824	do.	do.	do.	Do.
Do.	31798	Female ^b	Onomichi, Aki, Hondo	June 4, 1903	Do.
Do.	31809	Young	Koriyama, Yamato, Hondo	1903	Do.
Do.	31810	do.	do.	1903	Do.
Do.	31811	do.	do.	1903	Do.
Do.	31812	do.	do.	1903	Do.
Do.	34393	Male	Ishigakishima, Riu Kiu	Apr., 1899	A. Owston.
Do.	34394	do.	do.	do.	Do.
Do.	34395	do.	do.	do.	Do.
Do.	34396	Female	do.	do.	Do.
Do.	34397	do.	do.	do.	Do.
Do.	34398	Male	do.	do.	Do.
Do.	34399	do.	do.	do.	Do.
Do.	34400	do.	do.	do.	Do.
Do.	34401	Female	do.	do.	Do.
Do.	34402	Male	do.	do.	Do.
Do.	34403	do.	do.	do.	Do.
Do.	34404	do.	do.	do.	Do.
Do.	34405	Female	do.	do.	Do.
Do.	34406	do.	do.	do.	Do.
Do.	34407	do.	do.	do.	Do.
Do.	34408	Male	do.	do.	Do.
Do.	34409	do.	do.	do.	Do.
Do.	34410	do.	do.	do.	Do.
Do.	34411	do.	do.	do.	Do.
Do.	34412	do.	do.	do.	Do.
Do.	34470	do.	Northern Formosa	Mar. -, 1903	Do.
Do.	34471	do.	do.	do.	Do.
Do.	35273	Young	Nagasaki, Kiusiu		E. A. Mearns.
Do.	35274	do.	do.		Do.
Do.	36501	Female ^c	Taipa, Formosa	Sept. -, 1896	T. Tada.
Do.	36502	(^d)	Gilan, Formosa	Oct. -, 1896	Do.
Do.	36503	(^e)	Botel Tobago II	May -, 1897	Do.
Do.	36504	(^f)	Pescadores Is	Apr. -, 1897	Do.
Do.	36552	Male ^g	Okinawa shima		Sci. Coll., Tokyo.
Sci. Coll., Tokyo.	26a-c		Taipa, Formosa	Sept. -, 1896	T. Tada.
Do.	25a-g		Gilan, Formosa	Oct. -, 1896	Do.
Do.	27a-k		Botel Tobago II	May -, 1897	Do.
Do.	28a-d		Pescadores Is	Apr. -, 1897	Do.

^a Description, p. 129.^d Sci. Coll. No. 25.^f Sci. Coll. No. 28.^b Description, p. 128.^e Sci. Coll. No. 27.^g Sci. Coll. No. A. 1.^c Sci. Coll. No. 26, figs. 107-110.

RANA SWINHOANA ^a Boulenger.

1903. *Rana swinhoana* BOULENGER. Ann. Mag. Nat. Hist. (7), XII, Nov., 1903, p. 556 (type-locality, Bangkimtsing, Formosa; types, Brit. Mus. Nos. 99. 4. 24. 104 105; J. D. La Touche, collector).

Description of types.—*Female.* Vomerine teeth in two oblique series between the choanæ; head much depressed, slightly broader than long; snout as long as the diameter of the orbit, rounded; canthus rostralis obtuse; loreal region not very oblique, concave; nostril equally distant from the end of the snout and from the eye; interorbital space as broad as the upper eyelid; tympanum distinct, one-half to three-fifths the diameter of the eye; fingers moderate, first not extending beyond second; toes entirely webbed; tips of fingers and toes dilated into well-developed disks, which are much smaller than the tympanum; subarticular tubercles rather small; a small, oval inner metatarsal tubercle; the tibio-tarsal articulation reaches the tip of the snout; skin smooth, granular on the side; no dorso-lateral fold. Brown above, with small darker spots; a dark canthal streak and a dark temporal blotch; a whitish streak on the upper lip; limbs with rather indistinct dark crossbars; lower parts white.

Dimensions.—From snout to vent 80 mm. (Boulenger's original description).

Remarks.—Doctor Boulenger states that the nearest ally to this species is *Rana livida* Blyth (= *R. chloronota* Guenther), the range of which extends from Darjeeling, in the Himalayas, through the mountains of Assam and Tenasserim to Hongkong.

Habitat.—Thus far only recorded from Bangkimtsing, Formosa, where the types were collected by J. D. La Touche.

RANA ISHIKAWÆ ^b Stejneger.

1901. *Buergeria ishikawa* STEJNEGER. Proc. Biol. Soc. Washington, XIV, Dec. 12, 1901, p. 190 (type-locality, Okinawa shima; type, Tokyo Imp. Mus. No. 30).


The outward appearance of the last joint of fingers and toes together with the large disks misled me into believing that this species had an interpolated phalanx between the regular ultimate and the penultimate ones, but dissection has failed to discover such a bone, so that it appears safest to refer it to the genus *Rana*.

Description.—*Adult*; Tokyo Imperial Museum, No. 30; Okinawa shima. (Figs. 111–116.) Vomerine teeth in two short series behind the level of the choanæ; flat above, blunt, descending vertically at the tip; nostril much nearer the tip of snout than eye; canthus rostralis

^a "Named in memory of the late Robert Swinhoe, the first explorer of the reptile fauna of Formosa." See p. 184.

^b Named in honor of Prof. C. Ishikawa, Imperial University, Tokyo, director of the Department of Natural History of the Tokyo Imperial Museum.

well marked; lores concave; interorbital space slightly wider than upper eyelid; tympanum one-half the diameter of eye and scarcely more than its distance from the latter; fingers free, first extending beyond second almost the length of the disk, and with smooth swelling at base; disks large, about two-thirds the diameter of tympanum, rounded anteriorly; toes moderately webbed, two distal phalanges of fourth toe being free and the excision of the web reaching to the distal third of the basal phalanx of the same toe; subarticular tubercles prominent; a flat, weak, inner metatarsal tubercle, less than one-half the length of the first toe measured from the tubercle; no outer tubercle; a tarsal fold; specimen too hardened to be unbent or legs stretched out, but tibio-tarsal articulation would probably reach



FIGS. 111-116.—*RANA ISHIKAWÆ*. ♂ \times NAT. SIZE. 111, TOP OF HEAD; 112, SIDE OF HEAD; 113, OPEN MOUTH; 114, UNDERSIDE OF HAND; 115, UNDERSIDE OF FOOT ($\frac{1}{4} \times$ NAT. SIZE), 1; 116, TIP OF DIGIT. No. 30, IMP. MUS. TOKYO.

between eye and tip of snout; tibio-tarsal articulations of bent hind legs overlapping slightly; skin of upper surface exceedingly rough, even on upper eyelids and sides of face, the tubercles assuming a very extraordinary pattern caused by the smaller tubercles surrounding concentrically a number of large round pustules which are wrinkled radially from the central prominence; sides very roughly studded with closely set, large, wrinkled tubercles; underside finely granular, except posterior aspect of femur, which is coarsely so. Color (in alcohol) above brownish, a network of chocolate brown surrounding the large insular tubercles, which are ochraceous-buff, with the central prominence darker brown; lips pale, blotched with

dark brown; limbs crossbarred with dark brown and drab; upper posterior aspect of femur mottled with the same colors; underside pale russet, with a network of obscure whitish mottlings.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent (approximately <i>a</i>).....	115
Width of head.....	40
Distance between nostrils.....	10
Distance between nostril and eye.....	8.5
Diameter of eye.....	12
Diameter of tympanum.....	6
Interorbital space.....	61
Width of upper eyelid.....	9
Fore leg <i>a</i>	61
Largest finger disk.....	4
Hind leg, vent to tip of longest toe <i>a</i>	178
Tibia.....	53
Inner metatarsal tubercle.....	5.5

Remarks.—This large species is easily recognized by its very peculiar pattern of glandular pustules on the back, which is still more enhanced by the coloration, emphasizing the odd arrangement which suggests a map of the moon, only that there are no craters visible on the top of the volcanoes.

Habitat.—The only specimen thus far known is the type in the Imperial Museum, Uyeno Park, Tokyo, which is said to have come from Okinawa shima.

List of specimens of Rana ishikawæ.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.
Imp. Mus. Tokyo.....	30	(<i>a</i>)	Okinawa shima.....		

a Type; description, p. 132; figs. 111-116.

RANA NARINA *b* Stejneger.


1901. *Rana narina* STEJNEGER, Proc. Biol. Soc. Washington, XIV, Dec. 12, 1901, p. 189 (type-locality, Okinawa shima. Riu Kiu; type, Sci. Coll. Mus. Tokyo, No. 19*a*).

Description of type specimen.—Adult; Science College Museum, Tokyo, No. 19*a*; Okinawa shima (figs. 117-121). Vomerine teeth in two distinct, nearly transverse series, which do not extend beyond a line between the posterior margin of the choanæ; the distance between the series a little less than between the latter and the choanæ, which are very large; head longer than wide; snout long, nearly squarish

a The specimen has the back very much bent and is so hardened that it can not be stretched out to its normal length. For the same reason the legs can not be unbent and their measurements are, therefore, also approximate only.

b Having reference to the nostrils, *nares*, on account of their extreme forward position in this species.

truncate, vertical, the nostrils being situated near the outer extremity, their distance from edge of lip being about one-half their distance from the eyes; lores subvertical, very concave; eyes large; inter-orbital space narrower than upper eyelids; tympanum very distinct, its diameter about one-half the diameter of the orbit, its distance from the orbit being more than one-half its diameter; several large glandular warts behind the angle of the mouth; fingers rather long, first extending beyond second; terminal disks distinct, but small, much smaller than tympanum; subarticular tubercles large; toes broadly webbed, reaching to the disks of the third and fifth toes; disks about the size of those of the fingers; subarticular tubercles strong; inner



FIGS. 117-121.—*RANA NARINA*. NAT. SIZE. 117, TOP OF HEAD; 118, SIDE OF HEAD; 119, OPEN MOUTH; 120, UNDERSIDE OF HAND; 121, UNDERSIDE OF FOOT. No. 199, SCI. COLL. TOKYO.

metatarsal tubercle but slightly prominent, narrow, less than one-half the length of the inner toe; no outer metatarsal tubercle; fore limbs longer than tibia; tibio-tarsal joint extends considerably beyond the snout; heels overlap considerably when thighs are bent at right angles to the axis of the body; no dorsal or dorso-lateral folds; no tarsal fold; both surfaces smooth. Color (in alcohol) above brownish, below whitish; a narrow whitish line from under the eye to and including the postoral glands; sides and posterior surface of thighs coarsely marbled with dark brown; chin and throat clouded with dusky. The specimen appears to have faded considerably, so that the description of the coloration is probably very defective.

Dimensions.

	<i>mm.</i>
From snout to vent.....	66
From snout to corner of mouth.....	22
Length of head.....	22
Width of head.....	21
Diameter of eye.....	7
Width of upper eyelid.....	6
Interorbital width.....	5
From eye to nostril.....	7
From nostril to edge of lip.....	3.5
From eye to end of snout.....	8
Diameter of tympanum.....	4
From eye to tympanum.....	2.7
From snout to fore limb.....	23
Fore limb.....	43
Hind limb.....	120
Tibia.....	38
Foot.....	51

Remarks.—I know of no species to which the present one is particularly closely allied. In some respects it appears to resemble *R. everetti* Boulenger, from the Philippines, particularly in the position of the nostrils, but in detail there are more differences than similarities. The forward position of the nostrils in *R. everetti* is even exceeded in our species, and the vomerine teeth, digital disks, length of hind limb, etc., are totally different.

Habitat.—Only known from the unique type specimen which hails from Okinawa shima, Riu Kiu.

List of specimens of Rana narina.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.
Sci. Coll., Tokyo....	19 ^a	Adult ^a	Okinawa shima, Riu Kiu.....		

^a Type; description p. 134; figs. 117-121.

RANA NAMIYEI^a Stejneger.


1901. *Rana namiyei* STEJNEGER, Proc. Biol. Soc. Washington, XIV, Dec. 12, 1901, p. 190 (type-locality, Okinawa shima, Riu Kiu; type, Sci. Coll. Mus. Tokyo, No. 31A).

Description.—*Adult:* Science College Museum, No. 31A; Okinawa shima, Riu Kiu (Figs. 122-126).—Vomerine teeth in two^b rather large, very distinct, and very oblique series, the anterior end on a level with the posterior border of the choanæ, the distance from the latter being about one-half that between the posterior ends of the vomerines; lower jaw with two greatly developed tooth-like prominences in front, fitting into two deep pits in the upper jaw; head very large,

^a Named for Mr. M. Namiye, of the Science College, Imperial University, Tokyo.

^b The right one lost in this specimen.

its width at the angle of mouth equals the distance from snout to insertion of fore limb; snout short, truncate, without canthus ros-



FIGS. 122-125.—*RANA NAMIYEI*. $\frac{1}{4}$ NAT. SIZE. 122, TOP OF HEAD; 123, SIDE OF HEAD; 124, UNDERSIDE OF HAND; 125, UNDERSIDE OF FOOT. No. 31A, SCI. COLL. TOKYO.

tralis; nostril situated half way between eye and median point of upper lip; eyes directed upward; interorbital space one and one-half times as wide as upper eyelid; tympanum hidden; fingers short, slightly dilated at tip, the second one more so than the others, subarticular tubercles well developed; second finger extending considerably beyond first, and falling a trifle short of fourth, which reaches to the penultimate phalanx of third; toes broadly webbed to the tip of the toes, which are somewhat dilated; fourth toe about one-third longer than fifth, which extends only to the middle of third phalanx from the tip of the former, third reaching to the base of the penultimate phalanx; subarticular tubercles well developed, single; inner metatarsal tubercle prominent, with a free outer edge, the width




FIG. 126.—*RANA NAMIYEI*. NAT. SIZE. OPEN MOUTH. No. 31A, SCI. COLL. TOKYO.

contained two and one-half times in the length, which about equals the diameter of the eye; no outer tubercle; a well-developed membranaceous fold along the outer edge of the fifth toe; a tarsal fold; the tibio-tarsal joint does not reach the eye; the heels do not meet by a distance greater than the diameter of the eye when the hind limbs are bent at right angles to the axis of the body; skin above nearly smooth, very loose, and apparently transversely wrinkled; a well-marked fold from posterior corner of eye to above fore limb and another less distinct from under the eye backward parallel with the former. Color above brown, with a broad dark band between and including the posterior half of the upper eyelids, with the faintest trace of a lighter band in front of it; lower surface whitish, clouded with dusky, especially on the throat and chin.

Dimensions.

	<i>mm.</i>
From snout to vent.....	116
From snout to corner of mouth.....	40
Width of head.....	53
Diameter of eye.....	9
Width of upper eyelid.....	8
Interorbital width.....	12
From eye to nostril.....	9
From eye to end of snout.....	18
From snout to fore limb.....	48
Fore limb.....	62
Hind limb.....	138
Tibia.....	45
Foot.....	60
Inner toe.....	12
Inner metatarsal tubercle.....	9

Variation.—Another large specimen is in the Science College Museum, collected by Mr. S. Ungawa in Okinawa shima. It has numerous longitudinal tubercular ridges, and the skin is transversely wrinkled on the posterior part of the back only. The interocular light band is very pronounced.

A much smaller specimen in the same museum (No. 18), also from Okinawa shima, and collected by Mr. Nakagawa, differs chiefly in having the head proportionally less broad, the bony tubercles on the mandible less developed, vomerines placed a little more forward, tympanum pretty well defined, and first finger slightly extending beyond second. The skin of the back, eyelids, and thighs is strongly rugose with warts more or less connected by longitudinal ridges; no transverse plaiting. Upper lips with large blackish spots; transverse dark band between eyes very distinct, the light one less so.

At this age it resembles distantly some specimens of *R. tigerina*, from which it differs, however, in most essential points. Thus the nostrils are placed more forward; the interorbital space is much broader;

the vomerine series much shorter and more anterior; shorter tibia; larger inner metatarsal tubercle, etc., not to mention the different coloration.

Remarks.—This species is evidently most nearly related to *R. corrugata*, from Ceylon, and *R. kuhlii*, from southern China and the Malay archipelago, but it shows marked differences from both. From *R. kuhlii* it is easily distinguished by the much longer fifth toe, while it differs from *R. corrugata* in the much larger metatarsal tubercle and the large and prominent groups of vomerine teeth.

Habitat.—Thus far only known from Okinawa shima, Riu Kiu, from which island I have examined three specimens.

List of specimens of Rana namiyai.

Museum.	No.	Age.	Locality.	When collected.	By whom collected.
Sci. Coll. Tokyo.....	31A	Adult a....	Okinawa shima, Riu Kiu.....		
Do.....		do. b.....	do.....		S. Ungawa.
Do.....	18	Halfgr b.....	do.....		Nakagawa.

a Type: description, p. 136; figs. 122-126.

b Description, p. 138.

RANA TIGERINA *a* Daudin.


- 1803. *Rana tigerina* DAUDIN, Hist. Nat. Rain., fol. ed., p. 42; quarto ed. (p. 64), pl. xx (type-locality, Bengal; type in Mus. Paris; Massé, collector); Hist. Nat. Rept., VIII, 1803, p. 125.
- 1820. *Rana tigrina* MERREM, Tent. Syst. Amphib. (p. 174) (emendation).—HALLOWELL. Proc. Phila. Acad., 1860, p. 504 (Hongkong).—BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 26 (East Indies; Shanghai, Ningpo, China; Formosa).—PARENTI and PICAGLIA, Atti Soc. Natural. Modena, Mem. (3) V, 1886, p. 90 (Hongkong market).
- 1835. *Rana vittigera* WIEGMANN, Nova Acta Acad. Leop. Carol., XVII, Pt. 1, p. 255, pl. XXI, fig. 1 (type-localities, Laguna del Bay, Luzon, Philippine Islands, and Macao, China; Meyen, collector).
- 1835. *Rana rugulosa* WIEGMANN, Nova Acta Acad. Leop. Carol., XVII, Pt. 1, p. 258, pl. XXI, fig. 2 (type-locality, Cape Syng-more, China; Meyen, collector).

Several other names, in the synonymies credited to this species, are omitted here as they may be applicable to possible subspecies.

Description (figs. 127-131).—*Adult female*; British Museum, No. 68. 1. 27. 24; Formosa; Doctor Collingwood, collector. Vomerine teeth in two very large oblique series, their anterior border on a level with the anterior border of the choana, contiguous with the latter and nearly meeting on the median line; lower jaw with two fairly well-developed tooth-like prominences in front, fitting into pits in the upper jaw; width of head at angle of mouth equaling the distance between tip of snout and insertion of fore leg; snout rather short and rounded without canthus rostralis; nostrils situated about

a Careless form for *tigrinus*, spotted or barred like a tiger, *tigris*.

halfway between eye and tip of snout; interorbital space much narrower than upper eyelid; tympanum distinct, about three-fourths the diameter of the eye; fingers tapering, not dilated at tips; second finger shorter than first, about equaling fourth; toes nearly entirely webbed, not dilated at tips; a rather wide flap of skin bordering the outer side of the fifth toe, the flap being particularly wide along the metatarsal bone; subarticular tubercles moderate; inner metatarsal tubercle small, its length scarcely one-half the diameter of the eye; no outer tubercle; a tarsal fold; tibio-tarsal joints reach eyes when



FIGS. 127-131.—*RANA TIGERINA*. NAT. SIZE. 127, TOP OF HEAD; 128, SIDE OF HEAD; 129, OPEN MOUTH; 130, UNDERSIDE OF HAND; 131, UNDERSIDE OF FOOT. No. 7435, U.S.N.M.

hind legs are stretched forward along the side and meet when bent at right angles to the axis of the body; skin above with numerous, very distinct, longitudinal, short, glandular ridges, but no transverse fold, and no dorso-lateral fold; a well-marked fold from posterior corner of eye around upper edge of tympanum to above fore leg. Color brown above with numerous roundish dark spots; no vertebral light line; lower surface whitish with brownish marblings on throat and chest; thighs strongly marbled with dark brownish.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent	88
Width of head.....	32
Snout to corner of mouth.....	30
Diameter of eye.....	8
Diameter of tympanum.....	6
Width of upper eyelid	8
Width of interorbital space.....	5
Eye to nostril.....	8
Eye to tip of snout.....	14.5
Snout to fore leg.....	34
Fore leg.....	46
Tibia.....	39
Inner metatarsal tubercle.....	4

The *adult male* (U.S.N.M. No. 34392; northern Formosa) differs chiefly in the presence of copulatory excrescences on the side of the first finger, which is somewhat swollen at base, and by the two large external vocal sacs, located one on each side of the throat.

The *tadpoles* of this species have been described and figured by Mr. Stanley S. Flower from Siamese specimens.^a His description, drawn up from specimens obtained in June, 1897, at Ayuthia, Siam, is herewith given in full.

Length of body once and a half its width, rather more than half length of tail; nostrils a little nearer to the eyes than to the end of the snout; eyes on the upper surface of the body, nearer the end of the snout than the spiraculum, the distance between the eyes twice the distance between the nostrils, and about equal to the width of the mouth; spiraculum on the left side, directed backward and upward, a little nearer the anus than the end of the snout, visible from above and from below; anus opening on the right side. Tail from three and one-half to four times as long as deep; acutely pointed; upper crest convex, a little deeper than the lower, not extending on to the back; depth of the muscular portion at its base about half the greatest total depth. Mouth: the large powerful beak is entirely black; the upper mandible terminates in front in a long sharp tooth-like prominence; the lower mandible is bicuspid, each "tooth" being long and sharp; the lips are bordered with very short fleshy papillæ; inside the upper lip are five series of fine, black teeth; the first series is uninterrupted by the individual teeth being "grouped with intervals" about the center of the line; the remaining series are broadly interrupted, the fifth being very short and difficult to distinguish; the lower lip has also five series of teeth; the first is short and uninterrupted, the second long and uninterrupted, the remainder broadly interrupted and very short. Color (in life), above yellowish brown, mottled with darker brown, a very distinct dark brown crescent-

^a Proc. Zool. Soc. London, 1899, pp. 892-893, pl. LIX, figs. 2-2a.

shaped mark above each nostril; below white, purplish gray about the chin and throat; tail yellow, mottled with brown, a horizontal dark line along the median line of the basal third of the muscular portion; iris golden. Total length 52 mm.; length of body 18.5; width of body 12.5; length of tail 33.5; depth of tail 9.

Variation.—This is the largest species of frog in that part of the world which is included in the present work. The female described above is small. Another Formosan specimen in our collection (U.S.N.M. No. 34390), measures 131 mm. from snout to vent, and specimens from other localities as large as 165 mm. are on record. The males are smaller, our largest from Formosa (U.S.N.M. No. 34392) measuring only 94 mm. in total length. The hind legs appear also to be shorter, inasmuch as the tibio-tarsal articulation does not reach even the tympanum. The ground color in our specimens is not brownish but olive, and the color in life is probably some shade of the latter tint.

Remarks.—Specimens from Borneo, which I had the privilege of examining in British Museum, have the vomerine groups of teeth more oblique and separated from the choanæ by a space fully equal to that between the two patches. I can not, therefore, agree in regarding *Rana schlueteri* Werner as a synonym. Chinese specimens in our museum (U.S.N.M. No. 7435; Hongkong; W. Stimpson, collector) agree with the Formosan specimens as described.

Habitat.—Widely distributed from India and Ceylon through Burma and Pegu to the Malay peninsula and archipelago,^a Celebes, and the Philippine Islands; eastward it extends into southern China, where it occurs as far as Ningpo and Shanghai.

Its occurrence in Formosa is attested by four female specimens in British Museum, collected at various times by Swinhoe, Dickson, and Collingwood, and by three specimens, male and female, in the United States National Museum from northern Formosa, obtained through Mr. A. Owston.

List of specimens of Rana tigerina.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	7435	(^a)	Hongkong, China.		W. Stimpson.
Do.	34390	Female	Northern Formosa	May —, 1903	A. Owston.
Do.	34391	Male	do	do	Do.
Do.	34392	do. ^b	do	do	Do.
Brit. Mus.	68. 1. 27. 24	Female ^c	Formosa		Doctor Collingwood.
Do.	68. 1. 27. 25	do.	do		Do.

^a Figs. 127-131.

^b P. 141.

^c Description, p. 139.

^a Note, however, the observations made above under the heading of "Remarks."

Genus POLYPEDATES^a Tschudi.

1838. *Rhacophorus* TSCHUDI, *Classif. Batr.*, pp. 32, 73 (type, *R. reinwardtii*; not of Schlegel 1827).
1838. *Theloderma* TSCHUDI, *Classif. Batr.*, pp. 32, 73 (type, *T. leprosa*).
1838. *Buergeria* TSCHUDI, *Classif. Batr.*, pp. 34, 75 (type, *Hyla buergeri* Schlegel).
1838. *Polypedotes* TSCHUDI, *Classif. Batr.*, p. 34 (type, *P. leucomystax*).
1838. *Polypedates* TSCHUDI, *Classif. Batr.*, p. 75 (correction).
1843. *Trachyhyas* FITZINGER, *Syst. Rept.*, p. 31 (type, *Polypedates rugosus*=*P. leucomystax*).
1848. *Dendricus* GISTEL, *Naturg. Thierr.*, p. viii (substitute for *Buergeria*).
1890. *Polypeletes* COUES, *Century Dictionary*, IV, p. 4606 (emendation).

Authors who reject Fitzinger's generic names of 1843 as *nomina nuda*, in spite of the fact that they are accompanied by an explicit statement of type, are not always consistent, as for instance in the present case.

In Oken's *Isis* for 1827 Schlegel made a number of perfunctory remarks about various herpetological matters, in which he mentioned several manuscript names of genera instituted by Kuhl or H. Boie. Among these is also *Rhacophorus*, which is introduced (p. 294) in the following words:

“Nach *Hyla* reiht Kuhl sein neues Geschlecht: *Racophorus* ein. Sp.: *Rac. reinwardtii* Kuhl n. sp., *palmatus* Kuhl (*Hyla* Daud.), *moschatus* Kuhl n. sp., und noch zwey unbenannte Species vom Herrn Professor Reinwardt mitgebracht.”

That is all. A generic name without the slightest trace of a definition or description accompanied by three specific names, two of which are *nomina nuda*, as one (*R. reinwardtii*) was only described many years after, the other (*R. moschatus*) apparently never. Only one of the species mentioned could be recognized when the generic name was published, viz, Daudin's *Hyla palmata*, which of necessity must be the type. And as this species is a typical *Hyla*, *Racophorus* of 1827 becomes a synonym of this genus. Any other result is preposterous, for certainly a generic name without diagnosis to be tenable can not well have for type an undescribed species.^b And yet *Rhacophorus* is accepted by those who even refuse to quote Fitzinger's genera in the synonymy. Even the rehabilitated *Rhacophorus* of Tschudi (1838) is not in a much better position, since the species assigned to it were not described even then, as the part of Schlegel's "Abbildungen" containing the first description of *Rhacophorus reinwardtii* was published after Tschudi's work^c appeared, but it is plainly untenable in view of the different application of the name in 1827.

^a From *πολύ*, much; *πηδάω*, I jump.

^b That the genus *Rhacophorus* was understood in this sense by herpetologists before Tschudi's time is shown by van der Hoeven's use of *Racophorus* in 1833 (*Handb. Dierk.*, II, Pt. 2, p. 311) for *Hyla palmata*, *faber* and *crepitans*.

^c Tschudi, *Classif. Batr.*, p. 32.

I have not alluded to Kuhl's own application of *Rhacophorus* in 1822,^a because it was certainly unidentifiable at the time. He refers, it is true, to the dermal flap at the heel,^b but in so uncertain a way that the genus without reference to a described or figured species (and no species was described until twenty-five years afterwards) must have been completely unintelligible. The fact that we know now what he meant is not sufficient.

Polypedates, or *Polypedotes*, *Buergeria*, and *Theloderma* were proposed by Tschudi simultaneously for two different species of the same genus, *Theloderma* being mentioned first. Duméril and Bibron, however, in 1841, in combining the first two genera, selected *Polypedates* for the greater group, and Guenther, in 1887, merged *Theloderma* in *Polypedates*, hence this name must be adopted according to article 28 of the International Zoological Code of Nomenclature.

The genus *Polypedates*, as here understood, differs from the genus *Rana* chiefly in the presence of a small bone intercalated between the last phalanx of the digit and the normally penultimate one. Externally this arrangement is marked on the upper surface by a kink or notch just before the widening of the disk, thus rendering dissection in most cases unnecessary. However, it is much safer to dissect, as shown by the erroneous reference of *Rana ijimæ* and *Rana ishikawæ* to the present genus.

KEY TO THE JAPANESE AND FORMOSAN SPECIES OF POLYPEDATES.

- a**¹ Fingers considerably webbed; color above green (blue or purplish in alcohol), immaculate.
- b**¹ A distinct outer metatarsal tubercle; dermal margin of fingers much narrower than the bone.....*P. schlegelii*, p. 145.
- b**² No outer metatarsal tubercle; dermal margin of fingers equaling the bone in width.
- c**¹ Distance from tip of coccyx to end of sacral diapophysis equals the width of the head.....*P. viridis*, p. 147.
- c**² Distance from tip of coccyx to end of sacral diapophysis much less than width of head.....*P. owstoni*, p. 149.
- a**² Fingers free, or with only a rudiment of web at base; color above brownish, more or less marked with darker.
- b**¹ Tibia about one-half the length of head and body.
- c**¹ Vomerine teeth in two long straight series between the choanæ, each series considerably longer than interval between them.....*P. buergeri*, p. 150.
- c**² Vomerine teeth in two small oval series, each series considerably shorter than the interval between them.....*P. cijfingeri*, p. 153.
- b**² Tibia more than one-half the length of head and body.
- c**¹ Interorbital space narrower than upper eyelid; tympanum less than one-half the diameter of the eye.....*P. japonicus*, p. 155.
- c**² Interorbital space wider than upper eyelid; tympanum more than one-half the diameter of the eye.....*P. leucomystax*, p. 157.


^a Isis, 1822, p. 476.

^b This dermal heel flap is also found in *Hyla palmata*, and consequently is not diagnostic. As a matter of fact, the latter species being the only one known at the time, the genus was at first restricted to it, as shown above.

POLYPEDATES SCHLEGELII^a Guenther.

1858. *Polypedates schlegelii* GUENTHER, Cat. Batr. Sal. Brit. Mus., p. 81, pl. vi, fig. c (type-locality, Japan; types in Brit. Mus.).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1866, p. 111; 1876, p. 383 (Yokohama).—HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 120 (Tokyo; Tsukuba Mt., Hitatchi Prov.).—*Rhacophorus schlegelii* BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 86, pl. ix, fig. 1 (Japan).—OKADA, Cat. Vert. Japan, 1891, p. 67 (Aizu, prov. Iwashiro; Tadami; Tokyo).—*Racophorus schlegelii* IKEDA, Annot. Zool. Jap., I, Pt. 3, Aug. 1897, p. 113 (Aizu, prov. Iwashiro; Tokyo; breeding habits).
1892. *Rhacophorus schlegelii* BOETTGER, Kat. Batr. Mus. Senckenberg., p. 16 (Hakone Mts.; Yokohama).

Description.—*Adult female*; U.S.N.M. No. 23589; Yokohama, Hondo; September, 1896; L. Stejneger, collector (figs. 132–136).



FIGS. 132–136.—POLYPEDATES SCHLEGELII. $1\frac{1}{4} \times$ NAT. SIZE. 132, TOP OF HEAD; 133, SIDE OF HEAD; 134, OPEN MOUTH; 135, UNDERSIDE OF HAND; 136, UNDERSIDE OF FOOT. No. 23589, U.S.N.M.

Vomerine teeth in two nearly straight series between and close to the choanæ, each series shorter than the distance between them; snout declivous; nostrils slightly nearer the eye than the tip of snout; interorbital space more than one and a half times the width of upper eyelid; tympanum circular, about two-thirds the diameter of the eye; fingers webbed, between third and fourth to distal end of basal phalanx, the others at base only; first finger much shorter than second, not even reaching the disk of the latter; disks of second, third, and fourth fingers large, about three-fourth the diameter of the tympanum, that of

^a Named in honor of Dr. Hermann Schlegel, director of the Rijksmuseum at Leiden, and the chief author of the herpetological portion of "Fauna Japonica." Born, January 19, 1804, in Altenburg, Germany; died in Leiden, January 17, 1884.

the first small, all slightly rounded anteriorly; toes nearly entirely webbed; disks large, but smaller than those of fingers, gradually decreasing in size from fourth toe; a pear-shaped inner metatarsal tubercle, as large as one-half the first toe, rounded; a large, but not prominent outer one; tibio-tarsal articulations reach posterior border of eye, and do not meet when hind legs are placed vertical to the axis of the body; skin above very finely shagreened; no dorso-lateral folds; a strong fold from posterior border of eye over tympanum to shoulder; skin on underside granulated, the granulation increasing in coarseness from the throat backward. Color above in life green, in alcohol dull marine blue; beneath yellowish, as are also the upper surface of the two inner fingers, the three inner toes, and disks of all the digits; a few dusky spots on posterior aspect of femur; no cross-bars on legs.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent.....	49
Width of head.....	18
Interorbital space.....	6
Upper eyelid.....	3.5
Diameter of eye.....	5
Diameter of tympanum.....	3.3
Diameter of largest finger disk.....	3
Fore leg.....	29
Hind leg, vent to tip of largest toe.....	68
Tibia.....	20

Variation.—There is apparently very little external difference between the sexes. The *male* is said to have an internal vocal sac, and No. 34373, a May specimen, has the throat suffused with dark gray, which the others collected in the same locality and at the same time do not.

There is some variation in the length of the vomerine series, so that the internal is not often so great as in the specimen described. The tympanum is often smaller and the declivity of the snout is not always equally great.

Habitat.—This species is apparently confined to Japan. Thus far it is only recorded from Kiusiu (Miyazaki, U. S. Nat. Mus.) and Hondo as far north as 37° N. L. (Aizu, prov. Iwashiro, according to Okada). It is not uncommon in the woods near Yokohama (von Martens, Boettger, Stejneger) and Tokyo (Hilgendorf, Okada). Hilgendorf also records it from the Tsukuba Mountain, northeast of Tokyo in the province of Hitatchi, while Boettger has it from the mountains near Hakone. We also have four specimens from Mount Fuji, and in the Stockholm Museum there is one specimen collected by Dr. O. Nordquist at Enoshima. It is not found in Yezo, and is replaced in the Riu Kius by the next form.

List of specimens of Polypedates schlegelii.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	23389	Female ^a	Yokohama, Hondo.	Sept., 1896	L. Stejneger.
Do.	23390do.....do.....do.....	Do.
Do.	23591do.....do.....do.....	Do.
Do.	30741	Halfgrown	Miyazaki, Kiusiu.		C. A. Clark.
Do.	34370	Male	Mount Fuji, Hondo.	May, 1898	A. Owston.
Do.	34371do.....do.....do.....	Do.
Do.	34372do ^bdo.....do.....	Do.
Do.	34373do.....do.....do.....	Do.
Leiden.	c 1604		Japan.		v. Siebold.

^a Description, p. 145; figs. 132-136.^b P. 146.^c About 10 specimens.**POLYPEDATES VIRIDIS** ^a Hallowell.


1860. *Polypedates viridis* HALLOWELL, Proc. Phila. Acad., 1860, p. 500 (type-locality, Okinawa shima, Riu Kiu; type, U.S.N.M. No. 25397; W. Stimpson, collector). *Rhacophorus viridis* BOULENGER, Proc. Zool. Soc. London, 1887, p. 149 ("Loo Choo Islands," Pryer collection).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 865; author's separate, p. 16 (Okinawa shima).—BOETGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 106 (Okinawa shima).—BROWN, Proc. Phila. Acad., 1902, June 11, p. 185 ("Loo Choo Islands," probably Okinawa shima; Furness and Hiller, collectors).

The relationship of this form to *P. schlegelii* is very close indeed, so close that all my attempts to draw a hard and fast line based upon differences in proportion between hind legs and total length, distance of nostrils from eye, relative size of tympanum and large finger disks, etc., have failed in a large series. It seems, however, that the narrowness of the dermal margin of the digits in the northern form is a fairly good characteristic, inasmuch as the fingers appear slenderer and the disks consequently more differentiated, though in reality they are relatively smaller than in the southern forms. It is also possible that a larger series of the Okinawa form would show it to have slightly longer hind limbs, although in the only specimen before me the tibia is less than half the total length, notwithstanding the fact that the specimen is very much hardened, with the back bent at nearly 60 degrees. This specimen has also a very weak inner metatarsal tubercle, and the outer one is scarcely noticeable, and as Boulenger has also commented upon the smallness and flatness of the former in the specimen examined by him, it may be that we here have an additional character.

Under these circumstances it is unnecessary to submit a detailed description of this form, but a set of measurements of Science College

^a Latin word, signifying green.

Museum, No. 27 (figs. 137-141) from Okinawa shima is here submitted for comparison with those of a *P. schlegelii* given on page 146.



FIGS. 137-141.—POLYPEDATES VIRIDIS. NAT. SIZE. 137, TOP OF HEAD; 138, SIDE OF HEAD; 139, OPEN MOUTH; 140, UNDERSIDE OF HAND; 141, UNDERSIDE OF FOOT. No. 27, SCI. COLL. TOKYO.

Dimensions.

	<i>mm.</i>
Total length (back strongly bent).....	67
Width of head.....	25
Interorbital space.....	8.5
Upper eyelid.....	5.5
Diameter of eye.....	8
Diameter of tympanum.....	5
Diameter of largest finger disk.....	5
Fore leg.....	44
Hind leg (approximate).....	110
Tibia.....	31

According to the manuscript notes of Doctor Stimpson the color during life (of specimens taken in December) is "grass green^a above, below pale red; no dark line of separation between these colors."

Habitat.—Originally described from Okinawa shima, this species has been collected there by nearly all the travelers who have visited that island, where, according to Doctor Fritze, it is not rare.

^aNot "pale green" as rendered by Hallowell.

List of specimens of Polypedates viridis.

Museum.	No.	Age.	Locality.	When collected.	By whom collected.
U.S.N.M.	25397	(a)	Okinawa shima	Dec., 1854	W. Stimpson.
Sci. Coll. Tokyo.	27	Adult ^b	do.		

^a Type; poor condition.^b Dimensions, p. 148; figs. 137-141.**POLYPEDATES OWSTONI, a new species.**

Diagnosis.—Similar to *Polypedates viridis* and like it having the dermal margin of fingers equaling the bone in width; nostrils at least twice as far from the eye as the latter is from the edge of the lip; legs shorter, the tibio-tarsal joint not reaching beyond the center of the eye; distance from end of coccyx to end of sacral diapophysis much less than width of head and not more than distance from tip of snout to posterior rim of tympanum; no outer metatarsal tubercle.

Type.—U.S.N.M. No. 34333; Ishigaki shima, Yaeyama group, Riu Kiu Archipelago.

Remarks.—This new form is closely related both to *P. viridis*, from Okinawa shima, and to *P. schlegelii*, from Japan proper. With the former it shares the greater width of the fingers relative to the disks, due to the greater width of the lateral dermal margins, and also the greater distance of the nostrils from the eye, if this indeed is a character to be relied upon, while with the latter it has in common the comparatively shorter hind legs, inasmuch as the tibio-tarsal joint does not reach beyond the center of the eye when the hind leg is stretched forward along the side of the body. Correlated with this character is the relative shortness of the sacrum. In *P. owstoni* the distance from the end of sacral diapophysis to the end of the coccyx, or so-called urostyle, equals the distance from tip of snout to center of tympanum, being much less than the width of the head, while in the specimen of *P. viridis* examined by me that distance is much greater than from tip of snout to posterior rim of tympanum and at least as great as the width of the head. The inner metatarsal tubercle is larger than in *P. viridis*, but an outer tubercle is entirely absent. Most of the specimens have the sides, the anterior and posterior aspects of femur as well as the underside of the tibia, dotted with small roundish spots of a dark purplish brown, but this character, unfortunately, is not without exceptions.

As with *P. viridis* a detailed description is superfluous, but a list of measurements of the type specimen is appended for comparison with the others.

^a Named for Alan Owston, esq., of Yokohama, to whom science is indebted for numerous interesting additions to the Japanese fauna.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent.....	65
Width of head.....	23
Interorbital space.....	7.5
Upper eyelid.....	5
Diameter of eye.....	6.5
Diameter of tympanum.....	4
Diameter of largest finger disk.....	5
Fore leg.....	38
Hind leg, vent to tip of longest toe.....	90
Tibia.....	27

Habitat.—Thus far found only on the southern group of the Riu Kius. The United States National Museum has a fine series from Ishigaki shima. The specimen collected by Doctor Warburg, in 1891, in Miyako shima, and now in the Hamburg Museum (No. 534) belongs probably to the present form.

List of specimens of Polypedates owstoni.

U. S. N. M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.
34330	Adult	Ishigaki shima, Riu Kiu.		A. Owston.
34331	do.	do.		Do.
34332	do.	do.		Do.
34333	Female ^a	do.		Do.
34334	Adult	do.		Do.
34335	do.	do.		Do.
34336	Male	do.		Do.
34337	Adult	do.		Do.
34338	do.	do.		Do.
34339	do.	do.		Do.
34340	do.	do.		Do.
34341	do.	do.		Do.
34342	do.	do.		Do.
34343	do.	do.		Do.
34344	do.	do.		Do.
34345	do.	do.		Do.
34346	do.	do.		Do.
34347	do.	do.		Do.
34348	do.	do.		Do.
34349	do.	do.		Do.

^a Type.

POLYPEDATES BUERGERI^a (Schlegel).

Plate XII.

1838. *Hyla buergeri* SCHLEGEL, Fauna Japon. Rept., p. 113, Sauri. et Batr., pl. III, figs. 7-8^b (type-locality, Japan; types in Leiden Mus.; Buerger, collector).—SCHLEGEL, Abbild. Amph., 1844, p. 140, pl. I, fig. 5 (Japan; colored


^a To Doctor Buerger, the companion and successor of von Siebold in the exploration of the fauna of Japan.

^b Reproduced in this work on Plate XII.

original sent by Buerger).—BOETTGER, Offenbach. Ver. Naturk. 17–18 Ber., 1878, p. 8 (Japan).—*Polypedates buergeri* DUMÉRIL and BIBRON, Erpét. Gén., VIII, 1841, p. 521 (Japan).—GUENTHER, Cat. Batr. Sal. Brit. Mus., 1858, p. 80 (Japan).—*Rana buergeri* BOULENGER, Cat. Batr. Sal. Brit. Mus., 1882, p. 73 (Japan); Proc. Zool. Soc. London, 1886, p. 412 (Japan).—OKADA, Cat. Vert. Japan, 1891, p. 67 (Hakone, Musashi, Nikko, etc.).—BOETTGER, Kat. Batr. Mus. Senckenberg., 1892, p. 13 (Japan).—KREFFT, Verh. Ges. Deutsch. Naturf. Aerzte, 69 Vers., Braunschw., 1897, II, Pt. 1, 1898, p. 187 (habits).—*Rhacophorus buergeri* BOULENGER, Proc. Zool. Soc. London, 1888, p. 205.

1838. *Buergeria subversicolor* TSCHUDI, Classif. Batr., pp. 34, 75 (Japan; substitute name for *buergeri*).

Description (figs. 142–146).—*Adult female*; U.S.N.M. No. 31904; Kochi, Tosa, Shikoku: May 11, 1903; Dr. H. M. Smith, collector.



FIGS. 142–146.—POLYPEDATES BUERGERI. 14 X NAT. SIZE; 142, TOP OF HEAD; 143, SIDE OF HEAD; 144, OPEN MOUTH; 145, UNDERSIDE OF HAND; 146, UNDERSIDE OF FOOT. No. 23904, U.S.N.M.

Vomerine teeth in two long, nearly straight series between the choanæ, starting at the inner front corner of the latter, and separated from each other by an interspace less than the length of a series; nostril nearer the tip of snout than the eye; interorbital space about equaling the width of the eye; tympanum about one-half the diameter of the eye; fingers perfectly free; first finger much shorter than second, which falls considerably short of fourth; disks of third and fourth finger large, as large as tympanum, of second and first successively smaller; toes fully webbed; largest disks of toes equaling in size disk of second finger; subarticular tubercles strong; inner metatarsal tubercle small, weak, slightly projecting; no outer tubercle; tibio-tarsal articulations reach center of eye and overlap slightly when hind legs are placed at right angles to axis of body; skin above, including eyelids, snout, sides of face, and upper surface of limbs warty; underside nearly

smooth except lower belly and basal portion of posterior aspect of femur granular. Color (in alcohol) above dull brownish gray with obscure dark spots which become sharply defined and blackish on the sides and hind legs; a chevron mark of dusky on the upper eyelids and interorbital space, the convexity pointing backward, and two longitudinal curved lines between the shoulders with their convexity toward the median line; legs crossbarred and marbled with blackish; upper posterior aspect of femur marbled with black; underside whitish, lower lips and adjacent portions of throat, as well as lower portions of the thigh and the tibia, with numerous small black spots.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent.....	70
Width of head.....	24
Fore leg.....	42
Hind leg, vent to tip of longest toe.....	115
Tibia.....	33

The *adult male* is apparently much smaller, U.S.N.M. No. 31906, same locality, date, and collector, measuring only 45 mm. in total length. The hind legs are much longer, as the tibio-tarsal joint reaches the nostril, and the tibia is slightly longer than half the total length, viz. 23.5 mm. There is a large swelling at the base of the first finger, with a pad of copulatory asperities, as in the *Ranas*, covering the side of the metacarpal and the basal phalanx. There is also an internal vocal sac with an opening in the mouth on either side of the tongue. The coloration is essentially as in the female, only the dusky spots on the throat are much smaller and much more numerous.

Variation.—Our series is rather uniform, showing but little variation, except that there is a great deal of difference in the amount of dark markings on the underside, the above descriptions representing the maximum.

Habitat.—None of the records thus far published, except Okada's Catalogue, gives any definite localities for this strictly Japanese species. The original types collected by Buerger have only "Japan" for habitat, presumably southern Japan, or more strictly Kiusiu. However, in the United States National Museum we have specimens from Miyazaki, in Kiusiu, collected by Rev. C. A. Clark, and from the province of Iga, in Iondo, about 30 miles east of Osaka, through Professor Ijima. Dr. T. Lenz, in 1896, also collected it in the same general region, specimens from the provinces of Yamashiro, Setsu, and Yamato, surrounding Kyoto and Osaka, being in the Hamburg Museum (Nos. 1000, 1009, 1011). Osaka, besides recording it from the provinces of Mino, Hida, and Yamashiro, mentions Hayakawa, near Hakone, Chichibu in Musashi, and Yumoto and Chuzenji Lake, at Nikko, as places where Buerger's tree-frog occurs. United States



POLYPEDATES BUERGERI. (FROM SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 554.

National Museum also has a large number of specimens from Mount Fuji. Whether this species goes farther north than Nikko I do not know. In Shikoku it has been collected by Dr. H. M. Smith. In the Riu Kius it is replaced by the next species.

List of specimens of Polypedates buergeri.

U. S. N. M. No.	Sex.	Locality.	When col- lected.	By whom col- lected.
23004	Male ^a	Province of Iga, Hondo		I. Ijima.
23005	do.	do.		Do.
30745	Female	Miyazaki, Kiusiu		C. A. Clark.
31904	do. ^b	Kochi, Tosa, Shikoku	May 11, 1903	H. M. Smith.
31905	Male	do.	do.	Do.
31906	do. ^c	do.	do.	Do.
34330	do.	Mount Fuji, Hondo	May, 1898	A. Owston.
34351	do.	do.	do.	Do.
34352	do.	do.	do.	Do.
34353	do.	do.	do.	Do.
34354	do.	do.	do.	Do.
34355	do.	do.	do.	Do.
34356	do.	do.	do.	Do.
34357	do.	do.	do.	Do.
34358	do.	do.	do.	Do.
34359	do.	do.	do.	Do.
34360	do.	do.	do.	Do.
34361	do.	do.	do.	Do.
34362	do.	do.	do.	Do.
34363	do.	do.	do.	Do.
34364	do.	do.	do.	Do.
34365	do.	do.	do.	Do.
34366	do.	do.	do.	Do.
34367	do.	do.	do.	Do.
34368	do.	do.	do.	Do.
34369	do.	do.	do.	Do.

^a Figs. 142-146 p. 151.

^b Description, p. 151.

^c Description, p. 152.

POLYPEDATES EIFFINGERI ^a (Boettger).

1860. *Polypedates burgeri* HALLOWELL, Proc. Phila. Acad., 1860, p. 501 ("Loo Choo Islands") (not of Schlegel).

1895. *Rana eiffingeri* BOETTGER, Zool. Anz., XVIII, July 8, 1895, p. 267 (type-locality, "either, and probably, from Okinawa, of the middle group, or from Ohoshima, of the northern group;" type, Mus. Senckenberg., No. 1074a); Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 104.

As I have no specimen of *P. eiffingeri*, I give below a translation of Doctor Boettger's original description of the type (fig. 147) which, thanks to his courtesy, I had the pleasure of examining in 1898. Like *P. buergeri*, the specimen in question has the first finger much shorter than second and the tibia about one-half the total length of head and body.

^a Named for Mr. Georg Eiffinger, of Frankfort on the Main.

Original description of type specimen.—*Adult female*: Museum Senckenbergianum, No. 1074a: Riu Kiu Islands.—Vomerine series between the choana, feebly developed, small, oval, separated by a



FIG. 147.—POLYPEDATES EIFFINGERI. ABOUT 1½ X NAT. SIZE. VOMERINE TEETH. SKETCH BY AUTHOR FROM TYPE IN SENCKENBERG MUSEUM.

very wide interval; head broader than body, short, very broad; snout short triangular, anteriorly truncated; eye larger and more protruding than in *P. buergeri* (Schlegel); all other characters as in the latter, but the back less warty, a small white tubercle on the tibio-tarsal joint and a series of smaller, isolated white tubercles along the posterior edge of the lower arm and the tarsus, and the posterior legs shorter; the adpressed hind leg reaches with the tibio-tarsal joint to the anterior border of the eye (in *P. buergeri* to the nostril or to the tip of the snout); the digital disks somewhat smaller than the tympanum. Color quite similar, but the posterior aspect of the femur lighter, brownish yellow, with large blackish spots and marblings.

Dimensions.

	<i>mm.</i>
Length of head and body.....	35
Length of head	12.5
Width of head	14
Tympanum.....	2.25
Fore leg	22
Hind leg	57
Tibia.....	18
Largest digital disk	2 ^a

Habitat.—Thus far only known as occurring in the Riu Kiu Archipelago at large. Hallowell reports *Polypedates burgerii* from “Ousima” and the “Loo Choo Islands,” a record probably referable to the present species. The origin of the type is not known exactly, as Mr. Schmacker’s Japanese collector visited all three groups in the Riu Kius, but Doctor Boettger regards it as certain that the specimen in question came either from the northern or the middle group, with the probability in favor of the latter.

List of specimens of Polypedates eiffingeri.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected or from whom received.
Senckenberg ...	1074a	Female a....	Riu Kiu Islands.....	B. Schmacker.


^a Type: description, p. 154; fig. 147.

^a Boettger, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 105.

POLYPEDATES JAPONICUS (Hallowell).

1860. *Ixalus japonicus* HALLOWELL, Proc. Phila. Acad., 1860, p. 501 (type-locality, Oshima, northern Riu Kiu; type, U.S.N.M. No. 7313; Stimpson, collector).—OKADA, Cat. Vert. Japan, 1891, p. 66 (Okinawa shima).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 865; author's separate, p. 16.
1886. *Rana macropus* BOULENGER, Proc. Zool. Soc., London, 1886, p. 414 (substitute name; "Oho shima"); Ann. Mag. Nat. Hist. (6), X, Oct., 1892, p. 302 (Okinawa; Holst, collector).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 865; author's separate, p. 16.—BOETTGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 103 ("Ohoshima or Okinawa.")

I have convinced myself by a careful dissection of the fingers of one of Hallowell's cotypes of *Ixalus japonicus* that there is an intercalated bone between the penultimate and the ultimate phalanges, and that this species, therefore, belongs to the genus *Polypedates* and not to *Rana*. This arrangement of the tip of the digits is also quite easily



FIGS. 148-153.—POLYPEDATES JAPONICUS. $\frac{1}{4}$ X NAT. SIZE. 148, TOP OF HEAD; 149, SIDE OF HEAD; 150, OPEN MOUTH; 151, UNDERSIDE OF HAND; 152, UNDERSIDE OF FOOT; 153, TIP OF DIGIT. No. 7313, U.S.N.M.

discernible from the outside, the peculiar notch between the disk and the penultimate phalanx being very conspicuous.

The vomerine teeth are so insignificant that at a first inspection they seem to be absent. Under a very powerful lens, however, a transverse series of very minute teeth is discovered close to the anterior border of the choanæ. This justifies the reference of this species to *Polypedates* rather than to *Ixalus*, or *Philautus*^a Gistel, as the name should stand, *Ixalus* Duméril and Bibron, 1841, being preoccupied by *Ixalus* Ogilby, Proc. Zool. Soc. London, 1836, p. 119.

Description.—*Adult female*: U.S.N.M. No. 7313 a; Amami-o-shima, Riu Kiu; May, 1855; W. Stimpson, collector (figs. 148-153). Vomerine teeth in two oblique, very indistinct series between and very

^a*Philautus* GISTEL, Naturg. Thierr., 1848, p. x (substitute name for *Orchestes* Tschudi, 1838, which is also preoccupied by Illiger, 1807). See Stejneger, Proc. U. S. Nat. Mus., XXVIII, 1905, p. 346.

close to the inner border of the choanæ; nostril nearer the tip of snout than the eye; interorbital space narrower than upper eyelid; horizontal diameter of tympanum less than one-half the diameter of the eye; fingers free, first much shorter than second; disks of third fingers largest, less than half the diameter of tympanum; webs of toes excised to distal end of first phalanx of fourth toe; subarticular tubercles not very prominent; inner metatarsal tubercle small, oval; no outer tubercle; tibio-tarsal articulations reach beyond the tip of the snout more than half the width of the head when hind legs are extended forward along the body, and when bent at right angles to the axis of the body the same joints overlap more than the width of the interorbital space; skin above smooth with scattered tubercles, which are more numerous on the sides, and various very narrow and not conspicuous longitudinal folds which anastomose more or less and form two shallow crescents on the shoulders, with the convexity toward the median line; throat and chest smooth; rest of underside of body, as well as posterior aspect of the femur near the base, coarsely granular; a fold from posterior border of eye over tympanum to above insertion of fore leg. Color (in alcohol), above pale drab with brown markings; a chevron mark on top of head across the middle of upper eyelids with the angle pointing backward; on the shoulders two angular marks corresponding to the glandular folds, a dark vertical band from front of eye to edge of lip, and a longitudinal one from eye over upper part of tympanum to shoulder along the lower edge of the glandular fold; legs and feet cross-barred; lower lip with some dark brown spots; a dark brown longitudinal line in front of upper part of overarm; underside otherwise white.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent.....	33
Width of head.....	11
Fore leg.....	19
Hind leg, vent to tip of longest toe.....	62
Tibia.....	21

The *adult male* is smaller; U.S.N.M. No. 7313 *d*, same locality, date, and collector, measures only 26 mm.; a copulatory pad of minute asperities forms a swelling on top and side of the metacarpal of first finger; there is apparently an internal vocal sac with openings on both sides of the mouth back of the tongue.

Variation.—In structure and proportions there is very little variation observable. The coloration is also on the whole very uniform, though the extent and intensity of the dark markings vary somewhat; thus there is often a dark line connecting the angles of the scapular marks, and in some there is an elongated lozenge-shaped mark on the lower back.

Habitat.—Although originally recorded by Hallowell as from "Japan," this species is confined to the Riu Kiu, for the original label, with the locality "Ousima, Japan," is still in the bottle with Hallowell's types of *Iralus japonicus*. Boulenger, in referring the species to *Rana*, redescribed and renamed it (as there was already another *Rana japonica*) from a specimen in the British Museum, obtained by Lieut. Alfred Carpenter, R. N., at "Oho Shima," and he has since recorded its occurrence in Okinawa shima.^a Two specimens in the Science College Museum (Nos. 24 A and (1) P 2) from the latter locality have been examined by me and compared with Hallowell's types.

List of specimens of Polypedates japonicus.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.
U.S.N.M.	7313 a	Female a.	Amami-o-shima, Riu Kiu.	May, 1855	W. Stimpson.
Do.	7313 b	do	do	do	Do.
Do.	7313 c	do	do	do	Do.
Do.	7313 d	Male b.	do	do	Do.
Do.	7313 e	do	do	do	Do.
Do.	7313 f	do	do	do	Do.
Sci. Coll., Tokyo.	24 A	Female	Okinawa shima, Riu Kiu.		
Do.	(1) P 2	do	Naha, Okinawa		Shiraiishi.

^a Type; description, p. 155; figs. 148-153.

^b Description, p. 156.

POLYPEDATES LEUCOMYSTAX *b* (Gravenhorst).

1829. *Hyla leucomystax* GRAVENHORST, Delic. Mus. Vratislav., p. 26 (type-locality, Java; type in Breslau Mus.; Kuhl, collector).—*Rhacophorus leucomystax* BOULENGER, Proc. Zool. Soc. London, 1889, p. 29 (southern China and eastern Himalayas to Malay peninsula and archipelago).
1829. *Hyla servigata* GRAVENHORST, Delic. Mus. Vratislav., p. 28 (type-locality, Java; type in Breslau Mus.; Reinwardt, collector).
1835. *Hyla quadrilineata* WIEGMANN, Nova Acta Acad. Leop. Carol., XVII, Pt. 1, p. 260, pl. xxii, fig. 1 (type-locality, Manila, P. I.; Meyen, collector).
1841. *Polypedates rugosus* DUMÉRIE and BIBRON, Erpét. Gén., VIII, p. 520 (type-localities, Java and Manila; types in Paris Mus.).
1858. *Polypedates maculatus* GUENTHER, Cat. Batr. Sal. Brit. Mus., p. 78 (part: China; Philippines).
1860. *Polypedates megacephalus* HALLOWELL, Proc. Phila. Acad., 1860, p. 507 (type-locality, Hongkong).
1882. *Rhacophorus maculatus* var. *quadrilineata* BOULENGER, Cat. Batr. Sal. Brit. Mus., p. 84 (Singapore, Java, Philippines, Formosa).
1885. *Hylorana longipes* FISCHER, Archiv Naturg., LI, Pt. 1, p. 47 (type-locality, Pagat, southeast Borneo; type in Brit. Mus.; Grabowsky, collector).
1889. *Rhacophorus leucomystax* var. *servigata* BOULENGER, Proc. Zool. Soc. London, 1889, p. 30.

Description.—*Adult female*; Brit. Mus. No. 79.6.20.27; Formosa; Matthew Dickson, collector. Vomerine teeth in two nearly straight

^a Proc. Zool. Soc. London, 1892, p. 302.

^b From λευκός, bright, white; μύστραξ, upper lip.

rows between the choanae, contiguous with the latter and separated from each other by a space as wide as the choanae; nostrils nearer tip of snout than eyes; upper eyelids only two-thirds the interorbital space; diameter of tympanum about three-fourths that of the eye; fingers with barely a rudiment of web; first finger shorter than second by half the diameter of the disk, which is more than half that of the tympanum; toes about half webbed, inner metatarsal tubercle much smaller than digital disks; no outer tubercle; tibio-tarsal joint extends beyond the tip of the snout; skin nearly smooth above and on throat and chest, strongly granular on belly and underside of thighs; a sharp, narrow fold from posterior corner of eye in a straight line over tympanum, ending rather abruptly some distance past the insertion of the fore limb. Color (in alcohol), pale brownish gray above, with four obscure longitudinal brownish bands, the lateral ones broader and originating on the anterior portion of the upper eyelids, the median ones narrower, originating on the snout and apparently confluent on the sacrum; a narrow blackish line runs from the tip of snout through the nostrils, middle of eyes, and over the tympanum just underneath the postocular fold; on the flanks a few distinct blackish reticulations; edge of jaws whitish, accentuated by a very narrow line of dark brownish.

Dimensions.

	<i>mm.</i>
Total length, tip of snout to vent.....	58
Snout to eye.....	11
Snout to posterior border of tympanum.....	22
Nostril to eye.....	8
Diameter of eye.....	7
Diameter of tympanum.....	5
Interorbital width.....	7
Fore leg.....	36
Tibia.....	32
Digital disks.....	2.75

Habitat.—*Polypedates leucomystax* is said to inhabit "Southern China and Eastern Himalayas to the Malay peninsula and archipelago." The striped form, supposed to be only a color variety,^a has been found in Java, Sumatra, Singapore, the Philippines, and in Formosa, a specimen from the latter island, collected by Mr. Matthew Dickson, being in the British Museum.

^aMr. S. S. Flower, who studied *Polypedates leucomystax* alive in the Malay peninsula, says that the striped specimens do not represent "even a true variety, as the dark lines appear conspicuously and disappear entirely in the same individual. If killed with or without the lines visible they remain so in spirit. In Singapore at different times I noticed many young frogs which had just left the water, all of which had the dark lines visible; these disappear as the animal grows, only to reappear temporarily in the adult." (Proc. Zool. Soc. London, 1896, p. 906.)

List of specimens of *Polypedates leucomystax*.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.
Brit. Mus.....	79.6.20.27	Female ^a	Formosa.....		M. Dickson.

^a Description, p. 157.

TADPOLES.

The characters of the larvæ of the tailless batrachians of Japan and adjoining territory have received no attention thus far. Boulenger has given the study of the European forms special attention, and Flower has described several eastern Asiatic species. From Boulenger's key to the European tadpoles^a and Flower's descriptions it is possible to draw up a general scheme of the Japanese forms, by which at least the genera may be recognized. Upon this basis it will then be comparatively easy for the resident naturalists to gradually extend the key to include all the species.

It will be necessary for this purpose to give a brief account of the morphological characters which are utilized for the distinction of these larvæ. The following is therefore condensed from Boulenger's excellent chapter on the tadpoles in the work alluded to above.

Tadpoles for identification are preferably selected after the budding of the hind legs and before the appearance of the front legs.

By *body* is meant head and body together, and its longitudinal measurement is taken to the origin of the hind limbs. Depth of tail includes crests.

By *mouth* (fig. 154) is understood the opening with its horny beak, not unlike that of a cuttlefish, as well as the surrounding, funnel-shaped lip. This may be entirely bordered by fleshy papillæ, or these may be restricted to the sides, or to the sides and the lower border. The inner surface of the lip is furnished with ridges armed with series of minute, bristle-like, erect, horny teeth. By drawing an imaginary line across the mandibles, the lip may be divided into an upper and a lower portion, the series of teeth above the upper mandible being termed *upper labial*, those below the lower mandible





FIG. 154.—OPEN MOUTH OF TADPOLE OF *RANA TEMPORARIA*. 15 X NAT. SIZE. *bo*, BUCCAL ORIFICE; *lp*, LABIAL PAPILLÆ; *m*, UPPER, *m'* LOWER MANDIBLE; *l*, SERIES OF TEETH ON UPPER LIP; *l'*, SERIES OF TEETH ON LOWER LIP. (ADAPTED FROM PROC. ZOOLOG. SOC. LONDON, 1891, PL. XLV, FIG. 3A.)

^a *Taill. Batr. Europe*, pp. 105-109.

being *lower labial*. These are described as first, second, third, etc., proceeding from the outer border toward the beak in both the upper and lower sections of the lip. The first series, either in the upper or




FIGS. 155-156.—LOWER VIEW OF TADPOLES, SHOWING LOCATION OF SPIRACULUM AND ANAL OPENING. 155, DISCOGLOSSOID (COSTATE) TOAD; 156, HYLA. *sp.*, SPIRACULUM; *an.*, ANUS. (FROM BOULENGER, PROC. ZOOL. SOC. LONDON, 1891, P. 506.)

lower section, may be marginal or may be within the border, which is then occupied by fleshy papillæ. The arrangement of the series is expressed by a formula: $\frac{3}{2}$, for instance, indicating the number in the upper and lower sections, the line corresponding to the beak.

By *spiraculum* (figs. 155-156) is understood the opening or tube by which the water is expelled from the bronchial chambers. In our species it is located either on the left side or in the median line on the breast.

The *anal* opening (figs. 155-156) is also a tube situated at the beginning of the tail either in the middle line or laterally beside the origin of the lower caudal crest.

The openings of numerous *sensory ducts* form more or less complicated systems of lines on head and body, and are of some systematic value. A glandular streak extending from the nostril toward the eye is the *lachrymal canal*.



FIGS. 157-159.—MOUTH OF TADPOLES. 10 X NAT. SIZE. 157, BOMBINA SALSA; 158, HYLA ARBOREA. 15 X NAT. SIZE. 159, BUFO BUFO. (FROM BOULENGER, PROC. ZOOL. SOC. LONDON, 1891, PL. XLVII, FIG. 5A; PL. XLVI, FIGS. 1A AND 4A.)

PRELIMINARY KEY TO THE TADPOLES.

- a¹ Mouth with horny beak; series of horny teeth, more or less surrounded by papillæ.
- b¹ Spiraculum in the mid-ventral line (fig. 155); labial teeth in two or more rows in nearly every series *Bombina (bombina, salsa*, fig. 157)
- b² Spiraculum on the left side of the body (fig. 156); labial teeth in one row in each series.
- c¹ Anus median (fig. 155); spiraculum directed backward; lower lip only papillose at the corners.
 Bufo (bufo, fig. 159; *viridis; calamita; melanostictus* [Flower, Proc. Zool. Soc., 1896, pl. XLIV, fig. 3].)
- c² Anus dextral, fig. 156; spiraculum directed backward and upward; papilla continuous along lower lip or only narrowly interrupted in the middle.
- d¹ Anus opening above the lower edge of the tail; upper caudal crest extending far forward on the back almost to between the eyes. *Hyla (arborea*, fig. 158)
- d² Anus opening close to the lower edge of the tail; upper caudal crest not extending forward beyond the vertical of the spiraculum.
- e¹ Papillose series bordering lower lip uninterrupted in the middle. *Rana*.
- f¹ Upper lip not bordered by papillæ; beak white, more or less broadly bordered with black.
- g¹ Series of labial teeth $\frac{2}{3}$ or 3
 R. macrodon [Flower P. Z. S., 1899, pl. LIX, fig. 1]; *esculenta; arvalis*.
- g² Series of labial teeth $\frac{3 \text{ to } 5}{4}$ *R. temporaria*, fig. 154; *agilis*.
- f² Upper lip bordered by papillæ like lower one; beak entirely black.
 R. tigerina [Flower, P. Z. S., 1899, pl. LIX, fig. 2]
- e² Papillose border of lower lip narrowly interrupted in the middle.
 Polydectes leucomystax [Flower, P. Z. S., 1896, pl. XLIV, fig. 2; 1899, pl. LIX, fig. 3]
- a² Mouth without hard beak, labial teeth, or papillæ, consisting of a simple upper lip and a contractile lower one. *Microhyla ornata* [Flower P. Z. S., 1899, pl. LX, fig. 1]

Class REPTILIA.

- 1768. *Reptilia* LAURENTI, Synops. Rept., p. 19 (compare p. 6).
- 1820. *Pholidota* MERREM, Tent. Syst. Amph., p. 5.
- 1826. *Monopnoa* FITZINGER, Neue Classif. Rept., p. 3.
- 1841. *Haplopnœa* LEUCKART, Zool. Bruchst., II, p. 31.

The existing reptiles are divided by Dr. H. F. Osborn into two sub-classes, *Diapsida* and *Synapsida*. To the latter belongs only one existing order, viz, the turtles (*Testudinata*), which are characterized by having the scapular arch internal to the ribs, while in the three existing orders of the *Diapsida* it is external. These orders are: *Crocodylini* (also called *Loricata* or *Emydosauria*), characterized by two-headed ribs; *Rhynchocephalia*, represented by a single surviving genus in New Zealand (*Sphenodon*), superficially resembling a lizard, but distinguished from the next order by having, among other characters, the quadrate bone immovably fixed to the adjacent cranial elements

by suture, while in the *Squamata*, embracing snakes and lizards, it is loosely articulated with the cranium at the proximal end; the last two orders have one-headed ribs.

As stated, recent *Rhynchocephalia* occur only in New Zealand, and the crocodiles are not known with certainty to enter our territory, in which consequently only the *Squamata* and the *Testudinata* are found. These two orders are always easily distinguished by their external appearance alone, the solid encasement of the turtles in their bony shell, covered by horny plates or a soft skin, being a character quite sufficient. Moreover, the presence of teeth in the squamate order contrasts easily with the toothless horn-sheathed jaws of the turtles.

Subclass DIAPSIDA.

1903. *Diapsida* H. F. OSBORN, Science (n. s.), XVII, Feb. 10, 1903, p. 276.

Order SQUAMATA.

1811. *Squamata* OPPEL, Ordn. Rept., p. 14.

1825. *Squamosa* LATREILLE, Fam. Nat. Règne Anim., p. 93.

1831. *Ophidosaurii* EICHWALD, Zool. Specialis, III, p. 168.

1854. *Streptostylica* STANNIUS, Handb. Zootom., II, Wirbelth., Pt. 2, Amph., p. 5.

1866. *Lepidosauria* HAECKEL, Gen. Morphol., II, p. cxxxvi.

1898. *Sauria* GADOW, Classif. Vertebr., p. 24.

Besides the chameleons, which form a suborder by themselves (*Pren dentia*, or *Rhìptoglossi*), but do not occur east of India, the order *Squamata* consists of two suborders—the lizards (*Sauria*) and the snakes (*Serpentes*). The anatomical characters which distinguish them consist chiefly in the separate condition of the rami of the lower jaw in the snakes, while they are solidly united in the lizards; in the total absence even of vestiges of a pectoral arch in the snakes; and in the closing of the brain case anteriorly in the latter. There are no external characters which will in all cases separate a snake from a limbless lizard, except that in the latter the tongue is not retractile into a basal sheath, while in most cases they possess distinct eyelids and ear-openings, both wanting in the snakes. All the lizards within the region here treated of have four limbs and are thus easily distinguished from the snakes.

Suborder SAURIA.

1788. *Lacertæ* BATSCH, Anleit. Kenntn. Thiere Mineral., I (p. 437).

1802. *Sauria* MACARTNEY, in Ross' Transl. Cuvier's Lect. Comp. Anat., I, tab. iii.

1803. *Sauri* DAUDIN, Hist. Nat. Rept., V, tabl., p. 8.

1804. *Sauru* LATREILLE, Nouv. Dict. d'Hist. Nat., XXIV, tabl. méth., p. 61.

1828. *Pneumaturomolgaci* RITGEN, Nova Acta Acad. Leop. Carol., XIV, p. 274.

1828. *Sauræ* WAGLER, Isis, 1828, p. 860.

1831. *Lacertina* MUELLER, Tiedem. et Trevir. Zeitschr. Phys., IV, p. —.

1839. *Saures* SWAINSON, Nat. Hist. Class. Fish. Amph. Rept., II (Lardner's Cab. Encycl.), p. 148.
 1842. *Lacertilia* OWEN, Rep. Brit. Ass. Adv. Sci. Plymouth Meet., 1841 (p. 144).
 1845. *Saura* GRAY, Cat. Liz. Brit. Mus., p. 3.
 1849. *Porpopoda* MAYER, Rheinlaend. und Westphal. Verhandl., VI (p. 177).
 1857. *Lepidota* JAN, Cenni Mus. Milano, p. 37.
 1886. *Eriglossa* GILL, Rep. Smithson. Inst., 1885, p. 801.
 1898. *Autosauri* HAECKEL, in Gadow, Classif. Vertebr., 1898, p. 24.

The saurian fauna of Japan is very fragmentary, containing at most 30 species belonging to only 4 families. It is very suggestive that the bulk of this number belongs to the two families *Gekkonidæ* and *Scincidæ*, which are nearly cosmopolitan in their distribution. Members of these families, moreover, seem to be more easily transported to distant islands than almost any other lizards. Thus, among the species occurring in Japanese territory, *Hemidactylus frenatus* occurs throughout the islands and along the shores of the western Pacific, from Korea to Australia, the entire Indian Ocean, and even in St. Helena, in the Atlantic Ocean, while the *Cryptoblepharus pacilopleurus* has a distribution even more extended. The most remarkable range, however, is that of *Leiolopisma laterale*, which furnishes one of the most extraordinary instances of disconnected distribution, as the specimens of the East Asiatic colony seem to be practically indistinguishable from those occurring in North America east of the Rocky Mountains.

The herpetological fauna of Formosa is as yet too little known to allow us to make any generalization as to the route by which the Indo-Malayan species have reached the Riu Kiu Islands.

Of the rather numerous superfamilies into which the lizards may be divided, only three are represented in Japan and neighboring countries, viz, the *Gekkonoidæ*, the *Agamoidæ*, and the *Lacertoidæ*, the first two with one family each, the latter with two, the *Lacertidæ* and the *Scincidæ*. Some of their main characters are included in the following:

TABLE OF SAURIAN FAMILIES IN JAPAN, ETC.

- α^1 Tongue smooth, or with villose papillæ.
 b^1 (GEKKONOIDEÆ). No postorbital or postfronto-squamosal arches; clavicle dilated proximally GEKKONIDÆ, p. 164
 b^2 (AGAMOIDEÆ). Postorbital and postfronto-squamosal arches present; clavicle not dilated proximally AGAMIDÆ, p. 182
 α^2 (LACERTOIDEÆ). Tongue covered with imbricate scale-like papillæ or with oblique papillose folds.
 b^1 Premaxillary double; body with osteodermal plates SCINCIDÆ, p. 193
 b^2 Premaxillary single; no osteodermal plates LACERTIDÆ, p. 228

Taking cognizance of external, easily verified characters only, the species occurring within our area may be referred to their respective families by the following:

ARTIFICIAL KEY TO THE FAMILIES OF LIZARDS OCCURRING IN JAPAN AND ADJACENT TERRITORY.^a

a¹ No large symmetrical shields on top of head.

b¹ Digits dilated; no movable eyelids; tympanum exposedGEKKONIDÆ, p. 164

b² Digits compressed; movable eyelids; tympanum concealedAGAMIDÆ, p. 182

a² Top of head with large symmetrical shields.

b¹ Ventral scales cycloid, arranged quincunciallySCINCIDÆ, p. 193

b² Ventral scales squarish, in longitudinal and transverse seriesLACERTIDÆ, p. 228

Family GEKKONIDÆ.

The Geckos are lizards with smooth or papillose tongue, dilated clavicles and biconcave vertebræ; mostly without movable eyelids and with vertical pupil; toes variously modified or dilated into more or less adhesive organs.

A large family they are represented in all the hotter portions of the globe, but are most numerous in the Indian and Australian regions. In the area covered by this work only four or five genera have been found. Owing to the ease with which the geckos may be accidentally transported from place to place, they are often introduced unintentionally into new localities by human agency.

Probably most of the species are nocturnal, and owing to the construction of the toes are enabled to run with ease on smooth perpendicular surfaces. Some species emit a chuckling sound resembling the word "gecko," hence their name. They lay round eggs with a hard, white shell.

Swinhoe has given a spirited account of the habits of the common Japanese house gecko as observed by him in Formosa, which deserves to be reproduced here.^b

In running over the perpendicular face of the wall they keep so close and their movements are made so quickly with one leg in advance of the other that they have the appearance at a distance of gliding rather than running. The tail is somewhat writhed as the body is jerked along, and much so when the animal is alarmed and doing its utmost to escape; but its progress even then is in short runs, stopping at intervals and raising the head to look about it. If a fly perch on the wall it cautiously approaches to within a short distance, then suddenly darts forward, and with its quickly protruded glutinous tongue fixes it. Apart from watching its curious maneuvers after its insect food, the attention of the most listless would be attracted by the singular series of loud notes these creatures utter at all hours of the day and night, more especially during cloudy and rainy weather. These notes resemble the syllables "chuck-chuck" several times repeated, and, from their more frequent occurrence during July and August, are, I think, the call notes of the male to the female. During the greater part of the day the little creature lies quiescent in some cranny among the beams of the roof or in the wall of the house where, however, it is ever watchful for the incautious fly that approaches its den, upon whom it darts forth

^a For nomenclature of head-shields of lizards, see p. 194, figs. 176-178.

^b Swinhoe, Ann. Mag. Nat. Hist. (3), XI, 1863, pp. 222-223.

with but little notice. But it is by no means confined to the habitations of men. Every old wall and almost every tree possesses a tenant or two of this species. It is excessively lively, and even when found quietly ensconced in a hole, generally manages to escape—its glittering little eyes (black, with yellow ochre iris) appearing to know no sleep; and an attempt to capture the runaway seldom results in more than the seizure of an animated tail, wrenched off with a jerk by the little fellow as it slips away, without loss of blood. The younger individuals are much darker than the larger and older animals, which are sometimes almost albinos. * * *

I have found the eggs of this gecko in holes in walls or among mortar rubbish. They usually lie several together, are round, and did not seem to me to offer any appearance other than those of ordinary lizards. The young, when first hatched, keep much to themselves under stones in dark cellars, where they live until they attain two-thirds the size of the adults. At this stage they begin to show out in conspicuous places, but always evince alarm at the approach of their older brethren; for what reason, I could not make out.

KEY TO THE GENERA OF GEKKONIDÆ INHABITING JAPAN AND ADJACENT TERRITORY.

- a¹ Infradigital plates in a single series; the free distal joint at the extremity of the digital expansion.
- b¹ Digits with but a slight rudiment of web at base; no lateral parachute-like expansion *Gekko*, p. 165
- b² Digits entirely webbed; wide lateral parachute-like dermal expansion to body, limbs, and tail *Ptychozoon*, p. 170
- a² Infradigital plates in a double series, the distal joint rising from within the extremity of the digital expansion.
- b¹ Inner digit with a compressed clawed phalanx.
- c¹ Digits free; body without lateral dermal expansion *Hemidactylus*, p. 172
- c² Digits half webbed; body with a much developed lateral dermal expansion. *Cosymbotus*, p. 178
- b² Inner digit clawless *Peropus*, p. 180

Genus GEKKO^a Laurenti.

1768. *Gekko* LAURENTI, Synops. Rept., p. 43 (type, *G. verticillatus*).
1800. *Gekko* CUVIER, Leçons d'Anat. Comp., I, tabl. III (emendation).
1810. *Gecus* RAFINESQUE, Caratteri, p. 9 (emendation).
1826. *Platydictylus* FITZINGER, Neue Classif. Rept., p. 13 (same type) (not of Oken 1817).
1833. *Lomatodactylus* VAN DER HOEVEN, Handb. Dierk., II, Pt. 2, p. 342 (type, *L. vittatus*).
1843. *Scelotretus* FITZINGER, Syst. Rept., p. 101 (type, *G. vittatus*).

GEKKO JAPONICUS (Duméril and Bibron).

YAMORI.

Plate XIII.

1836. *Platydictylus japonicus* DUMÉRIL and BIBRON, Erpét. Gén., III, p. 337 (type-locality, Japan; types in Paris Mus.).—DUMÉRIL, Cat. Méth. Rept. Mus. Paris, I, 1851, p. 37.—*Gekko japonicus* GUENTHER, Rept. Brit. India, 1864, p. 103 (southern Japan; Chusan, China; Formosa).—HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 113.—BOULENGER, Cat.

^a A name derived from the sound made by several species.

- Liz. Brit. Mus., I, 1885, p. 188 (Japan; Kiukiang Mts., Chefu, Shanghai, Szechuen, Ningpo, Hongkong, China; Formosa).—OKADA, Cat. Vert. Jap., 1891, p. 71 (Tokyo; Chukuzen, etc., Okinawa).—BOERTGER, Kat. Rept. Mus. Senckenberg., I, 1893, p. 34 (Japan; Hongkong, Canton, Shanghai, Prov. Guan-dung, China).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 860; author's separate, p. 11 (Tokyo; Okinawa).—*Gekko japonicus* STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 218 (Taipa, Formosa).
1838. *Platydaetylus jamori* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., p. 103 (type-locality, southern Japan; types in Leiden Mus.; Siebold, collector).—BLEEKER, Natuurk. Tijdschr. Nederland. Indië, XVI, 1858, p. 204 (name only).—MARTENS, Preuss. Exped. Ost-Asien. Zool., I, 1866, pp. 109, 112 (Nagasaki).—*Gekko (Platydaetylus) jamori* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., 1838, pp. 136, 139, Saur. Batr., pl. II, figs. 1-4.^a
1842. *Gekko chinensis* GRAY, Zool. Misc., p. 57 (type-locality, China; type in Brit. Mus.; J. Reeves, collector).—*Platydaetylus chinensis* MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1876, p. 374 (Nagasaki).
1842. *Hemidaetylus nanus* CANTOR, Ann. Mag. Nat. Hist., IX, Aug. 1842, p. 482 (type-locality, Chusan, China; types in Brit. Mus.; Cantor, collector).
1863. *Gekko swinhonis* SWINHOE, Ann. Mag. Nat. Hist. (3), XII, p. 222 (Taiwan fu, Formosa; habits) (*nomen nudum*, not of Guenther 1864).
1891. *Platydaetylus jamori* FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, p. 239 (Tokyo; not in Yezo).
1902. *Gehyra intermedia* BROWN, Proc. Phila. Acad., 1902, June 11, p. 183 (type-locality, probably Okinawa shima; type in Wistar Inst. Phila. No. 572, orig. No. 46; Furness and Hiller, collectors).

The specimens from Tsushima and Fusan, Korea, are identical with the Japanese specimens, showing no tendency toward *Gekko swinhonis*^b from Northern China. The latter differs in having very few dorsal tubercles, median pair of chin-shields smaller, and scarcely any trace of web between the toes.


Description (figs. 160-163).—*Adult male*; U.S.N.M. No. 31821; Yamagawa, Province of Satsuma, Kiusiu; June 14, 1903; Hugh M. Smith, collector.—Rostral squarish, about twice as wide as high, bounded above by two large internasals and two small scales between the latter, the superior angles entering the nostrils; nostril between first supralabial, rostral, large internasal, and three smaller shields above and behind; distance between nostril and eye equals distance between eye and ear-opening; diameter of eye a little more

^a Reproduced in this work on Plate XIII.

^b 1864. *Gekko swinhonis* GUENTHER, Rept. Brit. India, p. 104, pl. XII, fig. A (type-locality, Peking; type in Brit. Mus.; Swinhoe, collector).—BOULENGER, Cat. Liz. Brit. Mus., I, 1885, p. 189 (Peking).—MUELLER, Verh. Naturf. Ges. Basel, VII, Pt. 3, 1885 (Tien-tsin).

1871. *Gekko japonicus* MOELLENDORFF, Journ. N. China, Br. R. Asiat. Soc. (n. s.), XI, p. 104 (Prov. Chili, China) (not of Duméril and Bibron).—MÉHELY, Zichy's Dritte Asiat. Forschungsr., II, 1901, p. 44, pl. vi, figs. 1-2 (Peking).


1885. *Gekko swinhoci* BOERTGER, Offenbach. Ver. Naturk. 24-25 Ber., p. 139; 26-28 Ber. 1888, p. 61 (Miau-feng-shan, near Peking); Kat. Rept. Mus. Senckenberg., I, 1893, p. 35.



GEKKO JAPONICUS. (FROM SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 554.

than half the distance between eye and tip of snout; ear-opening small, oval, less than one-half the diameter of the eye; eleven upper labials; mental pentagonal, scarcely larger than the adjacent lower labials; nine lower labials; behind mental a pair of median, somewhat elongated chin-shields, on either side of which another similar shield of about the same size; behind these several polygonal smaller shields; whole upper surface covered with granules, those on the snout considerably larger; among the granules, from the eyes backward to the root of the tail, numerous small, rounded tubercles, their mutual distance averaging about one-half the diameter of the eye, not arranged in regular longitudinal series, though about twelve tubercles can be counted in a line across the back; upper surface of



FIGS. 160-163.—*GEKKO JAPONICUS*. 2 × NAT. SIZE. 160, TOP OF HEAD; 161, SIDE OF HEAD; 162, CHIN; 163, UNDERSIDE OF FOOT. No. 17521, U.S.N.M.

limbs covered with granules like the back; lower surface of body and limbs covered with imbricate scales except on the throat and anterior portion of neck, which are covered with granules of the size of those on the back; first toe with 9 lamellæ, fourth with 15 underneath; second, third, and fourth toes connected by a basal web; an angular series of six preanal pores (by an anomaly two inner pores on left side not developed); base of tail swollen, with a single large tubercular scale on each side; tail suddenly tapering from the basal swelling, slightly depressed, oval in section, covered above with small scales, the annular arrangement of which is chiefly marked by the straight posterior outline of every ninth row, and a pair of median tubercles

on the latter; underneath scales larger with a median series of wide plates, every third of which is a little wider, its posterior edge corresponding to the posterior outline of the annulus. Color (in alcohol) drab gray above with a double series of obscure dusky spots on back and crossbars on limbs and tail; an obscure dusky band from nostril through eye to occiput meeting its fellow from the other side on the latter; a similar band from eye to ear-opening, and ill-defined dusky spots on both upper and lower labials.

Dimensions.

	<i>mm.</i>
Total length.....	106
Snout to vent.....	60
Vent to tip of tail.....	46
Snout to ear-opening.....	16
Greatest width of head.....	13
Fore leg, from axilla.....	18
Hind leg, from groin.....	26

The *adult female* (U.S.N.M. No. 31822; same locality and collector) differs chiefly in the absence of the preanal pores and of the basal swelling of the tail; the spur-like tubercle on each side of the base of the latter is present, but is smaller than in the male; the tail itself is slightly broader and more depressed.

Variation.—Besides slight differences in the number of labials, in the relative size and shape of the chin-shields, and in the number of tubercles on the occiput, it may be mentioned that some specimens, for instance, U.S.N.M. No. 13563, from Nagasaki, have plainly developed tubercles among the granules on the upper side of forearm and tibia, especially on the latter. The number of preanal pores in the males varies somewhat. In some specimens the series is narrowly interrupted on the middle line by a single poreless scale, in others the series is continuous. Thus, No. 13563, from Nagasaki, and No. 13751, from Yokohama, have three pores on each side of the median line; Sci. Coll. No. 30, from Formosa, has 9 pores in a continuous line; No. 14866, from Osaka, has 7 altogether, like a specimen from Shikoku, No. 31895.^a The spur-like tubercle on each side of the base is very variable. Often there is a group of three large tubercles, and in No. 13563 these are supplemented by two additional large scales at their base.

When the tail is broken off, the new tail grown, as in No. 13563, approximates the old one in size and shape, but the scales are larger somewhat imbricate, and not arranged in transverse series; neither is there any trace of annuli nor of tubercles; a few of the scales on the under side are widened, but there is no median series of regular shields.

^a Boulenger's statement (Cat. Liz. Brit. Mus., I, 1885, p. 188), that there are "six to sixteen on each side" is due to a lapse.

Habitat.—The center of its distribution seems to be south-central and eastern China. On the coast it occurs from Hongkong to Chefu. To the north it is replaced by *G. swinhonis*. Outside of China it is found in Formosa, the Riu Kius, Korea, Tsushima, and southern Japan.

In Formosa it is said to be "common everywhere," though the only localities from which specimens have been recorded are Taipa and Taiwan fu. Doctor Fritze reports it equally common in Okinawa. I have before me specimens from Miyako shima, Tsushima, and from Fusan, Korea, collected by Jouy. They are probably a recent introduction in the latter place.

In Japan proper it occurs at least as far north as Tokyo, and specimens from Yokohama, Nagasaki, and Osaka are in the various museums, while Okada enumerates it from the province of Kawachi, Suwo, and Nagato, in Hondo, and Chikuzen, in Kiusiu. Dr. Hugh M. Smith collected it in Satsuma and Shikoku. It does not seem to be found in the mountain districts, and Fritze specifically denies its occurrence in Yezo. Doctor Nozawa writes me that he has "heard of its presence in south Yezo," but if specimens have been taken there they have probably only been introduced in cargoes from farther south. In this connection I may cite Mr. E. Klocke's experience, who once observed a *Gekko* on the walls of a hut in the Chitose forests near Lake Shikotko, which he suggests might have been unintentionally introduced from Hondo.^a

List of specimens of Gekko japonicus.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	13563	Male	Nagasaki	May —, 1881	Jouy and Dale.
Do.	13751	do. ^a	Yokohama (?)		P. L. Jouy
Do.	14866	(^a)	Osaka		R. Hitchcock.
Do.	17513		Fusan, Korea	1885	P. L. Jouy.
Do.	17514		do.	1885	Do.
Do.	17515		do.	1885	Do.
Do.	17521	Female ^b	Tsushima	May —, 1885	Do.
Do.	21362	Young	Fusan, Korea	May 28, 1886	Do.
Do.	31821	Male ^c	Yamagawa, Kiusiu	June 14, 1903	H. M. Smith.
Do.	31822	Female ^a	do.	do.	Do.
Do.	31895	Male ^a	Kochi, Shikoku	May 11, 1903	Do.
Do.	31896	Young	do.	do.	Do.
Do.	34186		Kanagawa, Yokohama	Dec. 14, 1898	A. Owston.
Do.	34187	do.	do.	do.	Do.
Do.	34189	Female	do.	May 14, 1902	Do.
Do.	34190	do.	do.	do.	Do.
Do.	36528	Male	Miyakoshima, Riu Kiu		
Sci. Coll. Tokyo	30	do.	Taipa, Formosa	Sept. —, 1896	T. Tada.
Wistar Inst.	572	(^d)	Riu Kiu		Furness and Hiller

^a P. 168.

^b Figs. 160-163.

^c Description, p. 166.

^d Type of *Gehyra intermedia*.

^a Annot. Zool. Japon., V., Pt. 2, 1904, p. 85.

Genus *PTYCHOZOON*^a Kuhl.

1822. *Ptychozoon* KÜHL, Isis, 1822, p. 475 (*nomen nudum*).

1826. *Ptychozoon* FITZINGER, Neue Classif. Rept., p. 13 (type, *P. homalocephalum*).

1827. *Pteropleura* GRAY, Philos. Mag. (2), II, p. 56 (type, *P. horsfieldii*).

1893. *Ptychozoon* BOETTGER, Kat. Rept. Mus. Senckenberg., I, p. 35 (emendation).

There are two species of *Ptychozoon*, namely, *P. kuhli*^b and *P. horsfieldii*, which differ considerably. The more essential characters consist in the absence of enlarged tubercles among the dorsal granules in *P. horsfieldii*, the shorter series of preanal pores and the presence of a distinct series of femoral pores in the males of this species. Also in the much feebler segmentation of the tail, as well as the much smaller development of the caudal lobes, which do not join to form a terminal large flap as in *P. kuhli*.

PTYCHOZOON HORSFIELDII^c (Gray).

1827. *Pteropleura horsfieldii* GRAY, Philos. Mag. (2), II, July, 1827, p. 56 (type-locality, India: Singapore; type in Brit. Mus.; Hardwicke, collector).—*Platydaetylus horsfieldii* FITZINGER, Syst. Rept., 1843, p. 101.—*Ptychozoon horsfieldii* MÜLLER, Verh. Naturf. Ges. Basel, X, 1892, p. 209, pl. iv (Penang).

1845. *Ptychozoon homalocephala* GRAY, Cat. Liz. Brit. Mus., p. 164 (part: Singapore) (not of Creveld).—*P. homalocephalum* BOULENGER, Cat. Liz. Brit. Mus., I, 1885, p. 190 (part: Singapore); Proc. Zool. Soc. London, 1887, p. 146 (Riu Kiu Islands).—OKADA, Cat. Vert. Japan, 1891, p. 71 (Okinawa).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 860; author's separate, p. 11.

Thanks to the courtesy of the authorities of British Museum, I have been able to compare Pryer's specimen, said to have come from "Loo-Choo," with the type of Gray's *P. horsfieldii* (which is a female and consequently lacks the pores) from Singapore, and found them to agree completely. A male from Mount Dulit, Borneo, in the same museum, differs only in having three more preanal pores.




Description (figs. 164-166).—*Adult male*; British Museum, No. S7. 1. 31. 5; "Loo-Choo Islands;" presented by Mr. Pryer. Rostral almost quadrangular, large, at upper corners entering nostrils; the latter surrounded by rostral, first labial, one postnasal, and two supranasals; between the anterior supranasals a large nearly quadrangular shield separating them entirely and joining the rostral; twelve supralabials; mental small triangular, followed by a pair of

^a From πτυξ, a fold; ζωον, an animal.

^b *Ptychozoon kuhli* is Creveld's *Lacerta homalocephala*, the name of which is preoccupied. See Stejneger, Proc. Biol. Soc. Washington, XV, March 5, 1902, p. 37.

^c Named for Dr. Thomas Horsfield. Born May 12, 1773, at Bethlehem, Pennsylvania; died in London, July 14, 1859. Lived a long time in Java. Author of Zoological Researches in Java, etc. (1821-1824), and Plantae Javanicae, 3 vols. (1838-1852).

elongated median postmentals; top of head, back, and throat covered with uniform granules, without tubercles, belly with scales several times larger than the dorsal granules; parachute covered above with larger quadrangular scales in horizontal rows alternating like bricks in a wall, below apparently entirely without scales; an angular series of eight preanal pores, and a series of about twelve smaller pores near the distal end of the femur; the whole animal is strongly depressed and laterally surrounded by a series of dermal expansions or flaps, the one on the body between axilla and groin assuming the character of a parachute, being more than one-half the width of the whole body; a small flap behind angle of mouth underneath the ear-opening, one in the angle between upper and lower arms, and a small one on the posterior side of each of these; similar flap in angle between femur and tibia behind, and a small one on the outer side of the tibia, but none on the femur; tail with a lateral series of small rounded lobes, gradually decreasing in size toward the tip which is not expanded; the segments thus indicated by each pair of lobes are further marked by a couple of slightly enlarged flat scales on the upper surface of the tail near the median line;



FIGS. 164-166.—PTYCHOZOON HORSFIELDII. $\frac{1}{4}$ X NAT. SIZE.
164, SIDE OF HEAD; 165, UNDERSIDE OF RIGHT HIND LEG AND FOOT; 166, UNDERSIDE OF TAIL, MIDDLE AND TERMINAL PORTIONS. No. 24630, U.S.N.M.

hands and feet webbed to the tips of the digits. Color (in alcohol) light drab gray above with four narrow, wavy, dusky cross bands on the back and six wider bars on the tail; a line of ill-defined circular spots of same color from eye to shoulder; underneath whitish.

Dimensions.

	<i>mm.</i>
Total length.....	130
Snout to vent.....	68
Vent to tip of tail.....	62
Snout to eye.....	8
Snout to ear-opening.....	18
Width of head.....	15
Fore limb.....	32
Hind limb.....	62

The *female* lacks preanal and femoral pores.

Habitat.—This remarkable species is an inhabitant of the Malayan peninsula, the Natuna Islands, and Borneo.

A single specimen presented by Mr. Pryer to the British Museum as having been obtained by his Japanese collector in the Riu Kiu Islands, is the only one thus far recorded east and north of the region indicated above. As no other collectors have found it in the Riu Kius or the intervening regions, I may perhaps be justified in expressing a doubt as to the correctness of the locality. It may be remembered that Pryer himself did some collecting in Borneo in 1880, and it is possible that the specimen in question may have become mixed up with the Riu Kiu collection.

List of specimens of Ptychozoon horsfieldii.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.
Brit. Mus.	87. 1. 31. 5	Male ^a	Loo Choo Islands [?]		H. J. S. Pryer.
U. S. N. M.	24030	Female ^b	Trong, Lower Siam	1897	Dr. W. L. Abbott.

^a Description, p. 170.

^b Figs. 164-166.

Genus HEMIDACTYLUS^a Oken.

1817. *Hemidactylus* OKEN, Isis, 1817, p. 1183 (based on Cuvier's "Hemidactyles," type "Gecko tuberculeux Daudin").

1843. *Tachybates* FITZINGER, Syst. Rept., p. 105 (type, *H. mabouia*).

1843. *Pnoëpus* FITZINGER, Syst. Rept., p. 106 (type, *H. javanicus*=*H. frenatus*).

The above synonymy refers chiefly to the species occurring within our limits. For others see my "Land Reptiles of the Hawaiian Islands."^b

Only one species of true *Hemidactylus* has been reported from our territory thus far. Another is here recorded for the first time. They may be distinguished as follows:

^{a1} Inner digit very small, with a short sessile terminal joint, the claw of which is usually very minute; males with the femoral pores continued across the belly.

H. frenatus, p. 172.

^{a2} Inner digit not disproportionately small, well developed, with a free erect terminal joint, which is provided with a well-developed claw; males with femoral series of pores interrupted on the middle of the belly *H. bowringii*, p. 176.

HEMIDACTYLUS FRENATUS^c Duméril and Bibron.

1836. *Hemidactylus frenatus* DUMÉRIL and BIBRON, Erpét. Gén., III, p. 366 (type-localities, Java and Timor).—BOULENGER, Cat. Liz. Brit. Mus., I, 1885, p. 120 (Taiwan fu, South Formosa; Korea).—BROWN, Proc. Phila. Acad., 1902, p. 183 (Loo Choo Islands, probably Okinawa shima).

1860. *Hemidactylus inornatus* HALLOWELL, Proc. Phila. Acad., 1860, p. 492 (type-locality, Okinawa shima; type, U.S.N.M. No. 7429; Stimpson, collector).

1868. *Hemidactylus longiceps* COPE, Proc. Phila. Acad., 1868, p. 320 (type-locality, Manila, Philippine Islands.).

^a From ἡμι, half; δακτυλος, digit.

^b Proc. U. S. Nat. Mus., XXI, 1899, p. 791.

^c Signifying bridled; alluding to the dusky streak on the lores (*frenum*).

Description (figs. 167-168).—*Adult male*; U.S.N.M. No. 34202; Ishigaki shima, Riu Kiu Archipelago; June, 1899; A. Owston, collector. Rostral squarish, broader than high, upper outer corner entering nostril, a median cleft behind; nostril between first supralabial, rostral, and three nasals, of which the upper and the lower are larger than the one between; distance from nostril to eye scarcely longer than from eye to ear-opening; diameter of eye a little more than half its distance from tip of snout; ear-opening small, oval; 11 supralabials; mental large, triangular followed by a median pair of large oblique chin-shields broadly in contact; an outer pair of equally large shields extends a short distance back of the median pair, in contact with first and second lower labials; granules of throat extending close up to the chin-shields; upper surface covered with small granules, those on occiput, supraocular and interorbital region smallest, those on lores, rostral canthus, and anterior portion of snout largest; among the granules a few isolated rounded tubercles on lower back; first digit very small, the terminal compound joint scarcely free and with a very minute claw; six pairs of lamellæ, one terminal and two basal single lamellæ under fourth toe; under surface covered with small, flat, slightly imbricate



FIGS. 167-168.—HEMIDACTYLUS FRENATUS (TYPE OF *H. inornatus*).
2 X NAT. SIZE. 167, SIDE OF HEAD; 168, CHIN. No. 7419, U.S.N.M.

scales except on throat, where the scales are reduced to small granules; a continuous series of 31 femoral pores meeting at an angle in front of anus; tail gradually tapering, cylindric toward the tip, more flattened toward the base, covered above with small scales with indication of annulation, the posterior outline of the annuli being marked by three large pointed tubercles on each side and forming longitudinal series; underside of tail with a median series of wide plates. Color (in alcohol) above pale drab gray, with faint indications of dusky mottlings and cross marks; a pale broad band from nostril through eye and over temporal region, bordered above and below by an obscure narrow dusky line; labials mottled with dusky brown; underside whitish.

<i>Dimensions.</i>	<i>mm.</i>
Total length.....	99
Snout to vent.....	45
Vent to tip of tail.....	54
Snout to ear-opening.....	13
Greatest width of head.....	9.5
Fore limb.....	15
Hind limb.....	18

The adult *female* (No. 34196, same locality and collector) differs chiefly in the absence of femoral pores.

Variation.—The greatest amount of individual variation is seen in the number of tubercles on the back. In most specimens the tubercles are confined to the posterior half of the back, but in some examples scattered tubercles are found as far forward as the shoulder, while in others, as in the specimens described above, they are nearly absent. Sex does not seem to have anything to do with it, for we have males and females with numerous tubercles, as, for instance, Nos. 34206 and 34200, and likewise males and females almost devoid of tubercles, as Nos. 34199 and 34196.

The number of femoral pores in the males varies in the series before me between 28 and 35.

The size and mutual arrangement of the postnasals, and also of the chin-shields, are subject to some variation from what is described above; thus the outer pair of chin-shields are often excluded from contact with the first lower labial.

The reproduced tail is somewhat shorter and stouter at the base than the original one, the scales are somewhat larger and less regularly arranged, and there are no tubercles. The transverse plates on the underside are well developed, however.

Habitat.—One of the most widely distributed species of geckos, which has been carried in cargoes to many distant localities. It has recently been found in western Mexico, and it occurs also in St. Helena, in the South Atlantic. From South Africa and Madagascar it extends through the various groups of islands, India, Malayan Peninsula and Archipelago, northern Australia, Philippine Islands, Hainan, southern and eastern coast of China, to Formosa, Riu Kiu Islands, and Korea.

From the latter country there is a specimen in the British Museum presented by Sir E. Belcher, and in the same museum several from Taiwan fu, southern Formosa. Another specimen from Formosa is in the museum at Bergen, collected by Capt. von der Ohe. It has been collected in the "Loochoo Islands," as recorded by Dr. A. E. Brown, and specimens similarly labeled are in the Hamburg Museum from Doctor Warburg (No. 1152). Doctor Lenz, March 13, 1897, collected four specimens on Iriomote Island, of the Yaeyama group (Hamb. Mus., No. 1899), and the United States National Museum has 22 specimens from Ishigaki Island, of the same group. Finally, the Science College Museum, in Tokyo, has a number of specimens (No. 40) from Miyako shima, in the same group, and a single one (No. 41) from Naha, Okinawa shima, all collected by Tashiro.

List of specimens of Hemidactylus frenatus.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Femoral pores.
U.S.N.M.	7429a	Female a.	Okinawa shima.....	Nov. —, 1854	W. Stimpson.....	
Do.	7429b	Young.....	Okinawa shima (?)	do	W. Stimpson (?)	
Do.	7429c	do.	do.	do	do	
Do.	34191	Male.....	Ishigaki shima.....	June —, 1899	A. Owston.....	28
Do.	34192	do.	do.	do	do	28
Do.	34193	do.	do.	do	do	31
Do.	34194	do.	do.	do	do	29
Do.	34195	do.	do.	do	do	28
Do.	34196	Female.....	do.	do	do	
Do.	34197	Male.....	do.	do	do	30
Do.	34198	do.	do.	do	do	31
Do.	34199	do.	do.	do	do	33
Do.	34200	Female.....	do.	do	do	
Do.	34201	Male.....	do.	do	do	31
Do.	34202	do. b.	do.	do	do	31
Do.	34203	do.	do.	do	do	29
Do.	34204	do.	do.	do	do	30
Do.	34205	do.	do.	do	do	26
Do.	34206	do.	do.	do	do	35
Do.	34207	Female.....	do.	do	do	
Do.	34208	Male.....	do.	do	do	33
Do.	34209	do.	do.	do	do	19
Do.	34210	Female.....	do.	do	do	
Do.	34212	do.	do.	do	do	
Do.	34213	Male.....	do.	do	do	31
Do.	36529	do. c.	Miyakoshima.....	do	do	33
Do.	36530	do. c	do.	do	do	29
Imp. Mus. Tokyo	2	do.	Okinawa shima.....			31
Do.		Female.....	do.			
Do.		Male.....	do.			31

^a Type of *H. inornatus* (figs. 167-168).

^b Description, p. 173.

^c Sci. Coll. No. 40

HEMIDACTYLUS MARMORATUS^a Hallowell.

1860. *Hemidactylus marmoratus* HALLOWELL, Proc. Phila. Acad., 1860, p. 491 (type-locality, Loo Choo).—BOULENGER, Cat. Liz. Brit. Mus., I, 1885, p. 113.

The type has apparently been lost, and no species of *Hemidactylus* answering to Hallowell's description, which is very defective, to say the least, has been brought from the Riu Kiu Archipelago. Under these circumstances it may be useful to reproduce here Hallowell's original description in full, the obvious errors being corrected, as explained in the footnotes.

Description.—Mental plate pentangular, small; a transverse row of plates immediately behind the mental and first infra-labials; the two middle much larger than the lateral ones; behind these several

^a Signifying marbled; with dark markings like the veins in marble.

series of unequal scales, larger than those upon the throat;^a eleven supra-labials on each side; rostral^b plate pentangular, much broader than high, presenting an angle posteriorly; scales upon the muzzle and in front of the orbits, nearly equal, small; head covered with small granulations; five rows of very small circular tubercles on each side of the back above; a row of seven pores in front of the anus. Color grayish, marked all over with black; abdomen white; chin and throat white, marked with brown.

Dimensions.—Length of head 7 lines; greatest breadth $4\frac{1}{2}$ lines; length of head and body 1 inch 8 lines.

Habitat.—Loo Choo. One specimen.

It is just possible that Hallowell had before him a *Gekko japonicus* so shriveled or mutilated that the subdigital lamellæ appeared to be divided.

HEMIDACTYLUS BOWRINGII^c (Gray).


1845. *Doryura bowringii* GRAY, Cat. Liz. Brit. Mus., p. 156 (type-locality unknown; types in Brit. Mus.).—*Hemidactylus bowringii* BOULENGER, Cat. Liz. Brit. Mus., I, 1885, p. 139, pl. xii, fig. 2 (Burma; Northern and Central India); Fauna Brit. India, Rept., 1890, p. 93 (Eastern India, Burma, and Tenasserim).

1853. *Leiurus bermorei* BLYTH, Journ. Asiat. Soc. Bengal, XXII (p. 646) (type-locality. —).

1860. *Hemidactylus pumilus* HALLOWELL, Proc. Phila. Acad., 1860, p. 502 (type-locality. Hongkong).

1872. *Hemidactylus coctai* GUENTHER, Ann. Mag. Nat. Hist. (4), IX, Jan. 1872, p. 86 (part; not of Duméril and Bibron).—MUELLER, Kat. Amph. Rept. Mus. Basel, 1878 (p. 637) (Lilong and other localities in province of Canton).—BOETTGER, Offenbach. Ver. Naturk. 24-25 Ber., 1885, p. 138.

Description.—*Adult male*; U.S.N.M. No. 34188; northern Formosa; June 23, 1903; A. Owston, collector (figs. 169-170). Rostral squarish,



FIGS. 169-170.—HEMIDACTYLUS BOWRINGII. 3 X NAT. SIZE. 169, CHIN; 170, UNDERSIDE OF FOOT. No. 34188, U.S.N.M.

broader than high, with a median cleft behind, the upper corners entering the nostrils; a pair of internasals separated by a small scale; nostril surrounded by first supralabial, rostral, internasal and two small scales; eye equidistant from nostril and ear-opening; supra-

labials 12 on left side, 10 on right; mental large, triangular, or rather subpentagonal, followed by a median pair of large chin-shields, on the side of which a lateral pair of much smaller shields which

^aThe last paragraph in the original text follows immediately after the first sentence describing the mental plate; it is evidently misplaced and must be inserted in the position here given to it.

^bThe original text has "ventral," evidently a slip for rostral.

^cNamed after Charles Bowring, esq., who presented the types to British Museum.

do not extend backward beyond the median pair; granules on throat immediately join the chin-shields; upper surface covered with uniform granules (without trace of interspersed tubercles) those on vertex and on supraocular region smaller, those on snout and lores larger; first digit well-developed with a well-developed, free, and raised terminal joint provided with a full-sized claw; six lamellæ, of which three in pairs, under first toe; ten lamellæ, of which six in pairs, under fourth toe; throat covered with uniform granules, rest of under surface with small, rounded, imbricate scales; a well-developed series of 13 pores on each femur, not meeting on the middle of the abdomen, nor reaching the knee joint; tail subcylindric, more depressed near base, tapering, covered with uniform small scales of about the same size as the large granules on the snout; underside of tail covered with a median series of wide transversal plates. Color (in alcohol) above dark drab gray with obscure brown marblings; scarcely discernible dark band from lores through eyes, above ear-opening to side of neck, and a similar one on parietal region; underside whitish.

Dimensions.

	<i>mm.</i>
Total length.....	88
Snout to vent.....	50
Vent to tip of tail (reproduced).....	38
Snout to ear-opening.....	13
Greatest width of head.....	10
Fore limb.....	15
Hind limb.....	20

The adult *female* (U.S.N.M. No. 34211; Ishigaki Island, Riu Kiu; Owston collection) differs in the absence of femoral pores.

The most marked deviation, due to individual variation, from the specimen described above is the presence of a second lateral pair of chin-shields, smaller than the others, but well differentiated.

Remarks.—The specimens here recorded from Formosa and Riu Kiu have not been compared with Indian specimens, but they agree so completely with the published descriptions that I have no doubt that they are correctly identified. The well-developed inner toe with long terminal free phalanx and large claw, the series of 13 to 14 femoral pores on each side interrupted on the middle of the abdomen, and the tail devoid of tubercles indicate *H. bowringii* beyond a doubt.

The fact that this species has hitherto been recorded only from India, Burma, and Tenasserim is of no weight now, since Dr. Jean Roux, curator of the herpetological collection in the Basel Museum, informs me that the *H. coctæi* recorded by Mueller in the catalogue of that museum as from "Liolong and other localities in the province of Canton," China, is in reality *H. bowringii*, a fact of which he has assured himself by repeated and careful examinations and comparisons with allied species.

Habitat.—This species is known to inhabit the mainland of India, Burma, and Tenasserim, and as shown above, it is now recorded from the province of Canton in China. A single specimen collected in June, 1899, in "northern Formosa" and obtained from Mr. Alan Owston, is now in the United States National Museum. It is the first record of this species in the island.

Another specimen, a female, from Ishigaki shima, of the Yaeyama group, Riu Kiu Archipelago, was found among a lot of 22 specimens of *H. frenatus*. This species is probably much rarer than *H. frenatus*, or has different habits which result in its being more rarely caught.

List of specimens of Hemidactylus bouringii.

U. S. N. M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Femoral pores.
34188	Male ^a	North Formosa.....	June 23, 1903	A. Owston.....	13
34211	Female.....	Ishigaki shima, Riu Kiu.....	June —, 1899	do.....	

^a Description, p. 176; figs. 100-170.

Genus COSYMBOTUS^a Fitzinger.

1836. *Platyurus* OKEN, Allgem. Naturgesch., VI, p. 641 (not *Platurus* Latreille, 1802).
 1843. *Cosymbotus* FITZINGER, Syst. Rept., p. 104 (type, *H. platyurus*).
 1858. *Crossurus* GIRARD, Herpet. U. S. Expl. Exped., p. 281 (same type; not of Wagler 1830).
 1864. *Nycteridium* GUENTHER, Rept. Brit. India, p. 111 (same type).

COSYMBOTUS PLATYURUS^b (Schneider).

1792. *Stellio platyurus* SCHNEIDER, Amph. Physiol., II, p. 30 (no type-locality given; type in coll. Rebeft).—*Hemidactylus platyurus* BOULENGER, Cat. Liz. Brit. Mus., I, 1885, p. 143 (India, South China, Malay Peninsula and Archipelago; Philippines).
 1798. *Lacerta platyuros* DONNDORF, Zool. Beytr., III, p. 133 (emendation).
 1802. *Lacerta schneideriana* SHAW, Gen. Zool., III, Pt. 1, p. 1278 (substitute name).
 1829. *Gecko marginatus* CUVIER, Règne Anim., 2 ed., II, p. 54.
 1864. *Nycteridium schneideri* GUENTHER, Rept. Brit. India, p. 111 (emendation).

Description.—*Adult male*; Bergen Museum; Formosa; von der Ohe, collector. Rostral broader than high, in contact behind with two supranasals and a small scale between the latter; first supralabial in contact with a postnasal and reaching nostril; nostril between rostral, first supralabial and three smaller scales behind and above; 10 supralabials bordered above by a series of enlarged granules; scales on top of head granular, somewhat larger on snout, still larger on the loreal region; eye nearer ear-opening than nostril; large, about two-thirds the distance from ear-opening, which is small, about the size of one of the larger supralabials; mental shield large, penta-

^a From κοσμβωτός, fringed, tasseled.

^b From πλατύς, flat; ὀυρά, tail.

gonal, in contact behind with first lower labials and one pair of large postmentals, which are followed by another pair of equal size not in contact with each other; 7 larger lower labials which are bordered by a series of enlarged granules; body and legs above covered with small uniform granules; body and extremities below, as well as upper surface of feet, covered with larger imbricate scales, those on throat and underside of lateral fold, however, smaller, scarcely larger than the dorsal granules; from axilla to groin a thin skin flap about 2 mm. wide, and a similar fold along the posterior edge of thigh and tibia; five pairs of plates under shortest toe and eight pairs under longest; four toes nearly half webbed, hallux free; tail flat, with sharp, almost fin-like, and serrated edges, and a strong basal constriction; it is covered with somewhat pointed imbricate scales and underneath with a median series of about 40 wide subcaudal plates; at the basal origin of the flap-like lateral expansion of the tail a number of well-differentiated elongated spines; femoral pores continuous from knee across the belly, 20 on each side. Color (in alcohol) above drab with indistinct marblings of darker and lighter color; a distinct dark band from loreal region through lower part of eye and above ear to shoulder; below whitish.

Dimensions.

	<i>mm.</i>
Total length.....	88
Snout to vent.....	46
Vent to tip of tail.....	42
Snout to ear-opening.....	13
Greatest width of head.....	9.5
Fore leg, from axilla.....	16
Hind leg, from groin.....	23

Remarks.—From the above description it will be seen that the Formosan specimen agrees in all essential points with those from other localities. The second pair of postmentals is unusually large, but I have before me (in Bergen Museum) a specimen from Luzon, in the Philippines, which is halfway in this respect between normal specimens and the Formosan specimen.

Habitat.—This gecko is widely distributed from India and the East Indian Archipelago to southern China and the Philippines. So far as I know it is here recorded from Formosa for the first time. It was collected in the early sixties by Captain von der Ohe, from Bergen, who during his eastern cruises collected extensively for the museum in his native city.

List of specimens of Cosymbotus platyurus.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.
Bergen.....	Male ♂	Formosa.....	Capt. von der Ohe.

♂ Description, p. 178.

Genus *PEROPUS*^a Wiegmann.

1835. *Peropus* WIEGMANN. Nova Acta Acad. Leop. Carol., XVII, Pt. 1, p. 238 (type, *H. mutilatus*).
 1843. *Dactyloperus* FITZINGER. Syst. Rept., p. 103 (type, *H. variegatus*).
 1845. *Peripia* GRAY, Cat. Liz. Brit. Mus., p. 158 (type, *H. peronii*=*mutilatus*).
 1883. *Chalinocnemis* DUGÈS, La Natureza, VI, p. 312 (type, *H. navarri*=*mutilatus*).
 1883. *Spasmocnemis* DUGÈS, La Natureza, VI, p. 312 (same type).


Wiegmann's genus *Peropus* of 1835 antedates that of Lay and Bennett for a genus of fishes, as the Zoology of the *Blossom*, in which the latter occurs for the first time, was not published until 1839.^b

PEROPUS MUTIATUS^c (Wiegmann).

1834. *Hemidactylus mutilatus* WIEGMANN. Herpet. Mex., I, p. 54 (type-locality, Manila) (corrected for *H. pristiurus*, op. cit., p. 20).—*Hemidactylus* (*Peropus*) *mutilatus* WIEGMANN. Nova Acta Acad. Leop. Carol., XVII, Pt. 1, 1835, p. 238.—*Gehyra mutilata* BOULENGER. Cat. Liz. Brit. Mus., I, 1885, p. 148 (Mascarene Islands to Mexico).—*Peropus mutilatus* GIRARD, Herp. U. S. Expl. Exped., 1858, p. 277 (Philippines).—STEJNEGER, Proc. U. S. Nat. Mus., XXI, no. 1174, 1899, p. 796 (Hawaiian Islands).
 1857. *Dactyloperus insulensis* GIRARD. Proc. Phila. Acad., 1857, p. 195; author's separate, p. 5 (type-locality, Sandwich Islands).
 1883. *Hemidactylus navarri* DUGÈS, La Natureza, VI, p. 311, pl. VIIA (type-locality, Tangancicuara, Michoacan, Mexico).

For complete synonymy see Stejneger, Land Reptiles of the Hawaiian Islands.^d

Description (figs. 171–173).—*Adult female*; Bergen Museum; Formosa; von der Ohe, collector. Rostral broader than high, with a



FIGS. 171–173.—*PEROPUS MUTIATUS*, 2 × NAT. SIZE. 171, SIDE OF HEAD; 172, CHIN; 173, UNDERSIDE OF FOOT. No. 26395, U.S.N.M.

median cleft above, in contact with two supranasals and a minute granule between the latter; first supralabial in contact with the lower postnasal and reaching nostril; nostril between rostral, first supralabial and three scales behind and above, the upper one of which is large and in contact behind the rostral with the corresponding scale of the other side; 9 supralabials; scales on top of head finely granular, slightly larger on snout; eye equidistant from nostril

and ear-opening, large, more than two-thirds its distance from ear-opening which is moderate, somewhat larger than one of the largest

^a From *πρῶτος*, mutilated; and *ποῦς*, foot.

^b Liter. Gaz. and Journ. Belle Lett., London, no. 1179, p. 542, Aug. 24, 1839, where, according to T. S. Palmer, Index Gen. Mamm., p. 16, it is first mentioned in the list of new books. A review printed in Mag. Nat. Hist., London, March, 1840, begins thus: "The long-looked for and long-despaired of zoology of Beechey's Voyage is at last before us."

^c Signifying mutilated.

^d Proc. U. S. Nat. Mus., XXI., 1899, p. 796.

supralabials; mental shield rather small, triangular, followed by six chin-shields in one transverse series, their posterior outline forming together a straight line, the median pair being large, elongate pentagonal, the others gradually smaller toward the sides, the largest pair longer than diameter of eye; 9 large lower labials; body and legs above covered with small uniform granules like those on head; throat with granular scales of same size as those on back, those on rest of under surfaces, including those of the legs, larger, distinctly imbricate, and rounded behind; on the flanks a barely indicated longitudinal fold, and a somewhat better developed fold along the posterior edge of thigh and tibia; toes slightly webbed at base; about 8 pairs of lamellæ under longest toe and 6 under the shortest; tail (reproduced) rather flat, with sharp, finely serrated lateral edge, covered, with small, somewhat pointed, imbricate scales above and with a median series of wide plates bordered on each side by a somewhat irregular series of enlarged scales; about 5 mm. from the base the tail has a distinct lateral indentation; no femoral pores. Color (in alcohol) above brownish drab with faint indications of darker marblings, below whitish.

Dimensions.

	<i>mm.</i>
Total length.....	60
Snout to vent.....	33
Vent to tip of tail.....	27
Snout to ear-opening.....	8.5
Greatest width of head.....	7
Hind leg, from groin.....	12

Remarks.—The specimen agrees in most particulars with other specimens of this species.

The *males* have femoral pores in a continuous angular series across the abdomen, 14 to 19 on each side.

For further remarks as to variation, etc., see Proceedings United States National Museum, XXI, 1899, p. 798.

Habitat.—Irregularly distributed over various islands and coasts in the Indian Ocean, from the Mascarenes and Seychelles to the Philippines. This species seems very easily transported in ships, as evidenced by an apparently recent colony in western Mexico, and another in Hawaii. Its occurrence in Formosa and the Riu Kius is therefore not surprising. It was collected in Formosa during the early sixties by Capt. von der Ohe, who brought a single specimen to the museum at Bergen, Norway. This is the only record of its having been taken in Formosa.

Another specimen is in the Hamburg Museum (Saur., No. 1151), collected by Doctor Warburg in the Riu Kius, without further data.

List of specimens of Peropus mutilatus.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.
U.S.N.M.	26395	(a)	Manila, P. I.		G. C. Lewis.
Bergen		Female ^b	Formosa		Capt. von der Ohe.
Hamburg	1151		Riu Kiu Islands		Dr. O. Warburg.

^a Figs. 171-173.^b Description, p. 180.

Family AGAMIDÆ.

Lizards with a thick villose tongue and acrodont teeth; no large symmetrical plates on head or abdomen.

Distributed over the tropical and subtropical regions of the Old World, principally the Indian region; they are absent in Madagascar. The family contains both terrestrial and arboreal species, the former usually with a depressed body, the latter of a compressed shape. Some are insectivorous, others again herbivorous.

Only one genus occurs in Japanese territory.

Genus JAPALURA^a Gray.

1853. *Japalura* GRAY, Ann. Mag. Nat. Hist. (2), XII, p. 387 (type, *J. variegata*).

1853. *Biancia* GRAY, Ann. Mag. Nat. Hist. (2), XII, p. 387 (type, *B. niger*=*J. variegata*).

1860. *Diploderma* HALLOWELL, Proc. Phila. Acad., 1860, p. 490 (type, *D. polygonatum*).

1863. *Japalura* SWINHOE, Ann. Mag. Nat. Hist. (3), XII, p. 225 (emendation?).

1890. *Pelturagonia* MOQUARD, Le Naturaliste, XII (p. 144) (type, *P. cephalum*).

Species of this genus occur from the Himalayas and eastern Bengal through Assam to western China. One species each is known from Borneo, Formosa, Botel Tobago, and the Riu Kiu Archipelago.

KEY TO THE JAPANESE SPECIES OF JAPALURA.

^a Width between outer superciliary edges at center of eye exceeding length of third toe, without claw.

^b Fourth toe, without claw and measured from base of third, longer than distance from center of eye to tip of snout; throat dark with numerous roundish white (in alcohol) spots..... *J. swinhonis*, p. 184.

^b Fourth toe, without claw and measured from base of third, usually not longer than distance from center of eye to tip of snout; throat light colored, uniform or with converging dark lines..... *J. polygonata*, p. 188.

^a Width between outer superciliary edges at center of eye not exceeding length of third toe, without claw..... *J. mitsukurii*, p. 190.

As already indicated by Boulenger, *Japalura swinhonis*, from Formosa, and *J. polygonata*, from the Riu Kius, are very closely related, but the recent acquisition of a large series of specimens from Ishigaki shima shows a relationship even more close than was supposed

^a Derivation obscure; possibly the J is a misprint (?) for H, from ἀπάλος, fine, and οὐρα, tail.

by him. Not that the two forms are not distinct. On the contrary, there are quite a number of distinctive characters, but it is exceedingly difficult to express them in words and it is doubtful whether there is any *single* character which will separate the two forms in all cases.

Boulenger found the difference to be chiefly one of proportions and coloration. He stated^a that *J. polygonata* "differs in having the tibia constantly shorter than the skull, fewer labials as a rule, and in coloration," the inference being that in *J. swinhonis* the tibia is constantly at least as long as the skull. Unfortunately, however, the inference is not borne out by our material, since in several of the Formosa specimens of *J. swinhonis*, thus in U.S.N.M. Nos. 34084 and 34089, both old males, the tibia is decidedly shorter than the skull, as much so as in several Ishigaki shima specimens. It is true, nevertheless, that the Formosan form has the hind limbs longer as a rule, and this is more especially marked when we examine specimens of the male sex, since the females as a rule are somewhat more short-legged. *J. swinhonis* has a relatively longer fourth toe, the latter extending farther beyond the third than in *J. polygonata*, but the difference is so small that it defies a definite expression which would make it possible to use it as a criterion in a "key" for identification. A better result is obtained if we compare the length of the fourth toe (without claw) from the base of the third with the distance from the center of the eye to the tip of the snout. In most cases *J. swinhonis* has the fourth toe decidedly longer than this distance, while in *J. polygonata* it is usually equal or smaller.

The number of supralabials offers no tangible diagnosis. There are 7 or 8 labials in about equal proportion among over 20 Ishigaki shima specimens, not one with six.

The coloration furnishes possibly the best distinction, at least among the adults. While there are several of the adult males of the Ishigaki specimens which show a whitish lateral band quite distinctly and thus differ from the uniform green color of some of the Okinawa adult males, there is one color character in which all the old Formosan specimens agree, viz, in having the throat dark with large distinct white spots. This pattern is not found in any of our Riu Kiu specimens in which the throat is whitish more or less suffused with rust-red, uniform in the males but in the females marked with irregular longitudinal dusky lines converging posteriorly. In the latter there is also a more or less conspicuous longitudinal pale line under the eyes parallel with the upper lip, while in the Formosan specimens there are distinct blackish lines running from the eye to the lip.

There are several other minor characters of more or less constancy which are real enough though hard to express. Thus, the outline of the head is somewhat different especially when viewed from below,

^a Proc. Zool. Soc. London, 1887, p. 146.

inasmuch as in *J. swinhonis* the rami of the jaw are more convex than in *J. polygonata* in which they are more straight. In the latter the ventral scales are possibly a trifle larger, but I have been unable to establish a ratio. It also seems as if the spines of the nuchal crest were more conical and pointed, while in *J. swinhonis* they are wider and more rounded toward the apex.

When establishing *J. mitsukurii*, I had only one Formosan specimen for comparison. The latter was then unhesitatingly taken for a typical *J. swinhonis*, but the 17 specimens from Formosa since received put a somewhat different aspect on the whole question. The single specimen referred to shows such aberrant characters that I have become dubious as to its Formosan origin, or if it really comes from that island whether it may not represent a different species, questions which will be discussed farther on under the head of *J. swinhonis*. An examination of the large series of typical specimens of the latter now shows that the principal character upon which I relied for the diagnosing of the Botel Tobago form, viz, the absence of a specially differentiated series of larger scales between the eye and the upper labials, does not hold. But there are other differences separating it from both the other forms. In coloration *J. mitsukurii* agrees most with *J. swinhonis*, but the throat is much paler and the white markings on it, instead of being isolated roundish spots, form more or less distinct bands across the throat. In some of its proportions it also comes close to *J. swinhonis*. Thus, for instance, the distance from center of eye to tip of snout is very much less than the length of the fourth toe from the base of the third. Both feet and tail are longer. On the other hand, the head is much narrower and its lateral outline much straighter, in this respect resembling *J. polygonata*, only exceeding it. This narrowness of the head compared with the elongation of the toes furnishes a good character for distinguishing *J. mitsukurii*.

JAPALURA SWINHONIS^a Guenther.

1864. *Japalura swinhonis* GUENTHER. Rept. Brit. India, p. 133, pl. xiv, fig. B (type-locality, Tamsui, Formosa; type in Brit. Mus.; Swinhoe collector; see Ann. Mag. Nat. Hist. (3), XII, 1863, p. 225).—BOULENGER, Cat. Liz. Brit. Mus., I, 1885, p. 309 (Formosa); Proc. Zool. Soc. London, 1887, p. 146.—MUELLER, Verh. Naturf. Ges. Basel, VIII, Pt. 2, 1887, p. 294 (southern Formosa).—? STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 218 (Taipa, Formosa).
1885. *Japalura swinhori* BOETTGER, Offenbach. Ver. Naturk. 24–25 Ber., 1885, p. 140 (emendation).

^a Named for Robert Swinhoe, then British consul in Formosa, known from his investigation of the Chinese and Formosan faunas. He was born in Calcutta, September 1, 1836, and died in London, October 28, 1877. He was in the British consular service in China from 1854 to 1873.

A single adult female specimen (Sci. Coll. Mus. No. 22) said to have been collected by Mr. Tada at Taipa, Formosa, during July, 1896, has been recorded by me as *Japalura swinhonis*^a with the statement that "it agrees well with the published description." At that time I had no other Formosan specimens for comparison, and only three from Okinawa. The fine series of typical *J. swinhonis* from Formosa, and of *J. polygonata* from Ishigaki shima, now at hand, make the specimen in question appear in a somewhat different light. In fact, it presents so many peculiarities differing from all the three forms known from our territory that I do not know whether to regard it as an individual freak or a new species, in which case one might be tempted to question the correctness of the alleged habitat.

It has the dark dorsal cross-bars of the Formosan *J. swinhonis* as well as the white lateral longitudinal band, only the latter is unusually wide, but the throat is entirely uniform white, showing none of the characteristic pattern found in all our adult specimens of the latter. In addition it has the uninterrupted white longitudinal streak found in the *J. polygonata* of the Yaeyama group.

The proportions are very different from those of the normal Formosan specimens, and would even be considered extreme among the *J. polygonata*. The hind legs are exceedingly short, particularly the tibia, which is not longer than the distance from tip of snout to the posterior edge of the orbit, consequently very much shorter than the skull. The third toe, without claw, equals only three-fourths of the width between the outer superciliary edges. The distance from center of eye to tip of snout is much greater than the fourth toe measured from the base of the third. If the hind leg be carried forward along the side, the tip of the longest toe does not reach farther than the posterior edge of the (concealed) tympanum. It has seven supralabials. In addition to the above characters it has a row of enlarged scales from under the eye to the nasal, separated from the labials by a row of very narrow ones. Some of these enlarged scales are as large as some of the supralabials and three times as large as the other scales below and in front of the eye. Finally there is a very distinct row of enlarged submalar scales along the entire length of the lower edge of the mandible. In the other specimens, especially those from Formosa, only the two or three scales on each side of the mental are thus enlarged. In some of the Ishigaki shima specimens, however, a tendency to such a submalar series may be traced, but the full development as shown in the Taipa specimen seems to be unique.

It does not seem probable that the above deviations from the typical Formosan form can be due to individual variation of the

^a Journ. Sci. Coll. Tokyo, XII, Pt. 2, 1898, p. 218.

ordinary kind. It may represent a freak or a sport, but in view of the many points in which it differs, this explanation does not appear plausible. That two separate species so closely related should occur in the same locality in Formosa would scarcely be accepted by herpetologists without incontrovertible proof. The question then naturally suggests itself whether by some accident a wrong locality may not have been assigned to it. To the neck of the specimen a large paper tag is firmly tied bearing the following inscription in Latin letters: "No. 22, Taipa, Formosa. 7-96," but there are good reasons for believing that this is not the original collector's field label. However, if not from Formosa proper, where could it have come from? The specimen, as we have seen, is not identical with any from the middle or southern groups of the *Riukius*. On the other hand, that it was part of the collection made by Mr. Tsunasuke Tada seems unquestionable. He collected in Formosa, Botel Tobago, and the Pescadores Islands. Our specimen agrees even less with *J. mitsukurii*, from Botel Tobago, than with the other two forms. Could it, then, possibly have come from the Pescadores or any other outlying island?

The question can not be decided now, and is thus left for future explorers in that region.

Description.—*Adult male*; U.S.N.M. No. 34082; northern Formosa; June 23, 1903; Owston collection. Rostral low, separated from nasal by one scale; nostril round, in a single, somewhat swollen nasal, which is in contact with first supralabial; canthus rostralis very sharp, covered with about five scales; superciliary ridge interrupted by a deep notch at the posterior end of the orbit, followed by a single low spine; various isolated spines on occiput and temporals; eight supralabials, eighth very long and narrow; all head scales wrinkled and keeled; back covered by imbricated keeled scales of at least three different sizes; a low nuchal and dorsal crest of pointed scales, those of the former slightly higher, seven in number, flattened, conical, the latter triangular, 33 in number; a similar crest on the base of the tail, the three crests being continuous, the spines over the shoulder and between the hind legs, however, being lower than the others; scales on flanks not much different from dorsals, except those back of the axilla and near the groin which are much smaller; scales on underside strongly imbricate and keeled, pointed but not mucronate; digits underneath with a double series of spiny keels; third and fourth finger nearly equal; fourth toe, without claw, from base of third considerably longer than distance from center of eye to tip of snout; third toe, without claw, from base of fourth, shorter than distance between superciliary edges at center of eye; distance from upper side of knee of hind leg, bent at right angles, to tip of

longest toe without claw, longer than distance from tip of snout to inner angle of elbow; tail slightly compressed, swollen at base, covered above and below with imbricate, keeled and pointed scales. Color (in alcohol) above brownish, with a broad irregular, black-edged, pale gray band on each side of back, connected across the back by several indistinct pale chevron bands pointing backward, these chevrons occupying corresponding series of enlarged scales; head with numerous black spots, which on temples and under eye form interrupted lines radiating from the latter; throat and chest dark slate gray, the former with large, rounded, white spots; rest of underside pale gray; tail with alternating broad bands of pale gray and dark brown.

Dimensions.

	<i>mm.</i>
Total length.....	236
Snout to vent.....	70
Vent to tip of tail.....	166
Snout to center of eye.....	12
Greatest width of head.....	15.5
Fore leg.....	35
Hind leg.....	62

The *females* differ chiefly in lacking the basal swelling of the tail underneath, in the less development of the nuchal and dorsal crests, and in having the throat brown with white spots instead of dark slate gray.

The *young* have the throat uniform whitish or with faint gray marblings.

Variation.—The variation in the size and relative position of the head scales is considerable but has no immediate significance in so far as the identification of the species is concerned. The extent of individual variation of proportions and coloration has been alluded to above under the general discussion of the related forms (p. 183).

Habitat.—This species, which is restricted to the main island of Formosa, was first discovered at Tamsui by Robert Swinhoe, whose name it bears. It has since been obtained by most collectors in various parts of the island, and judging from the large number received by Mr. Owston it does not seem to be rare even in the northern part as formerly supposed. It is replaced by closely allied forms in the adjacent islands, thus by *J. polygonata* in the Riu Kius and *J. mitsukurii* in Botel Tobago.

On the Chinese mainland there is a nearly related species, *Japalura yunnanensis*, which occurs from western Yunnan to Ichang, on the Yangtse Kiang, and north into the province of Shen-si, whence we have a specimen from the Chin-ling Mountains collected by Dr. E. Blackwelder (U.S.N.M. No. 35524).

List of specimens of Japalura swinhonis.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
U.S.N.M.	34082	Male ^a	Northern Formosa	June 23, 1903	A. Owston.
Do.	34083	do.	do.	do.	Do.
Do.	34084	do. ^b	do.	do.	Do.
Do.	34085	do.	do.	do.	Do.
Do.	34086	do.	do.	do.	Do.
Do.	34087	do.	do.	do.	Do.
Do.	34088	do.	do.	do.	Do.
Do.	34089	do. ^b	do.	do.	Do.
Do.	34090	Female	do.	do.	Do.
Do.	34091	Male	do.	do.	Do.
Do.	34092	Female	do.	do.	Do.
Do.	34093	Male	do.	do.	Do.
Do.	34094	do.	do.	do.	Do.
Do.	34095	Female	do.	do.	Do.
Do.	34096	Young	do.	do.	Do.
Do.	34097	do.	do.	do.	Do.
Do.	34098	do.	do.	do.	Do.
Sci. Coll., Tokyo	22	Female ^c	Taipa, Formosa	Sept., 1896	T. Tada.

^a Description, p. 186.^b P. 183.^c See pp. 184, 185.**JAPALURA POLYGONATA** ^a (Hallowell).

Native name "ATAKAB."

Plate XIV.

1860. *Diploderma polygonatum* HALLOWELL, Proc. Phila. Acad., 1860, p. 490 (type-locality, Kerama shima, "Amakarima," Riu Kiu Archipelago; type U.S. N.M. No. 12203).—*Japalura polygonata* GUENTHER, Rept. Brit. India, 1864, p. 134.—BOULENGER, Cat. Liz. Brit. Mus., I, 1885, p. 310; III, 1887, p. 493 ("Loo Choo"); Proc. Zool. Soc. London, 1887, p. 146, pl. xvii, fig. 1^b ("Loo Choo Islands").—OKADA, Cat. Vert. Japan, 1891, p. 71 (Okinawa shima).—BOETTGER, Ber. Senckenberg. Naturf. Ges., 1888, Abhandl., p. 187 (O-shima, Riu Kiu); Kat. Rept. Mus. Senckenberg., I, 1893, p. 46 (O-shima); Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 107 (Okinawa shima).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 860; author's separate, p. 11 (Okinawa).—BROWN, Proc. Phila. Acad., 1902, June 11, p. 184 ("Loo Choo Islands").

This species agrees so closely with the preceding one that a detailed description is unnecessary. The differences, which consist chiefly in the shorter legs and longer and narrower head of *J. polygonata*, have been sufficiently indicated above under the general discussion of the genus (pp. 182-184).

The colors of the Okinawa shima specimens are said to be green in life. Thus Doctor Stimpson in his manuscript catalogue describes the type as being "dark green, back shaded, and tail annulated and still darker." In the alcoholics there are distinct traces of the dorso-lateral pale band as well as the pale line under the eye.

^a From *πολύγωνος*, with many angles, referring to the polygonal scutes of the head.^b Reproduced in this work on Plate XIV.




JAPALURA POLYGONATA. (FROM BOULENGER.)

FOR EXPLANATION OF PLATE SEE PAGE 554.

In the specimens from Ishigaki shima there are indistinct dark cross bands on the back. The males have the throat more or less uniform pale brownish gray, but in the females it is more whitish with irregular though distinct reddish brown lines running obliquely from the sides backward toward the median line. The coloring of the throat is therefore very different from what it is in *J. swinhonis*.

Dimensions.—*Adult male*; U.S.N.M. No. 12203; Kerama shima, Riu Kiu; type (figs. 174-175).



FIGS. 174-175.—*JAPALURA POLYGONATA*. 2 × NAT. SIZE. 174, TOP OF HEAD; 175, SIDE OF HEAD. No. 12203, U.S.N.M.

	<i>mm.</i>
Total length (tail defective).....	189
Snout to vent.....	66
Vent to tip of tail (defective).....	123
Snout to center of eye.....	11
Greatest width of head.....	15
Fore leg.....	32
Hind leg.....	55

A full-grown male, with perfect tail, U.S.N.M. No. 34102, from Ishigaki shima measures 61 mm. from snout to vent, and 152 mm. from vent to tip of tail; total length 213 mm.

Habitat.—Originally described from a single specimen collected by the North Pacific Exploring Expedition on "Amakarima" Island, *i. e.* Kerama shima, opposite Okinawa shima, this species has since been found common on the latter island by Doctor Fritze and by Mr. Tashiro. The Science College in Tokyo has several specimens (No. 26) collected there by the latter, two of which have been presented to the United States National Museum. Mr. Pryer's specimens, now in the British Museum, reported upon by Boulenger probably also came from Okinawa.

Doctor Boettger records a specimen presented to the Senckenberg Museum by Maj. H. von Schoenfeldt as from O-shima.

From the Yaeyama group the United States National Museum has a fine series collected on Ishigaki shima by one of Mr. Owston's men, and the Science College Museum in Tokyo has three specimens from Miyakoshima (No. 27) and one from Iriomote shima (No. 32), all collected by Tashiro in 1887.

List of specimens of Japalura polygonata.

U. S. N. M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.
12203	Male ^a	Kerama shima	Apr. —, 1855	North Pac. Expl. Exp.
34089	do.	Ishigaki shima		A. Owston.
34100	do.	do.		Do.
34101	do.	do.		Do.
34102	do. ^b	do.		Do.
34103	do.	do.		Do.
34104	do.	do.		Do.
34105	do.	do.		Do.
34106	do.	do.		Do.
34107	Female	do.		Do.
34108	do.	do.		Do.
34109	Male	do.		Do.
34110	do.	do.		Do.
34111	do.	do.		Do.
34112	do.	do.		Do.
34113	Female	do.		Do.
34114	do.	do.		Do.
34115	do.	do.		Do.
34116	do.	do.		Do.
34117	Male	do.		Do.
34118	Female	do.	Do.	
36519	(c)	Okinawa shima	Tashiro.	
36520	(c)	do.		Do.

^a Type, p. 189, figs. 174-175.

^b P. 189.

^c Sci. Coll. No. 26.

JAPALURA MITSUKURII^a Stejneger.

1898. *Japalura mitsukurii* STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 218 (type-locality, Botel Tobago Island; type, U.S.N.M. No. 36505; T. Tada, collector).

Japalura mitsukurii differs from *J. swinhonis* chiefly in proportions and coloration. Thus, the head, and especially the snout, is longer and narrower; the hind legs are somewhat longer, and the tail very much so. As a rule, the scales are somewhat smaller,^b with less contrast between the extremes, but the nuchal and dorsal crests are better developed, the latter continuous with the caudal, the keels on underside sharper and the scales more pointed. The coloration

^a Dedicated to Prof. Kakichi Mitsukuri, of the Imperial University of Tokyo.

^b Thus, as a rule, the nasal is smaller and separated from the supralabial by a narrow scale, while in *J. swinhonis* they are as often in contact. The percentage is not great enough to make this character available for diagnostic purposes.

is marked by about seven dark bars across the back separated by narrow pale cross stripes, which combine with the strongly defined light dorso-lateral band into a ladder pattern; the throat in the males is pale bluish (in alcohol) with large whitish spots, which form about three broad bars across the throat; the sides below the dorso-lateral band are dark with numerous whitish spots; the white longitudinal stripe under the eye is conspicuous, but mostly crossed by narrow black lines radiating from the eye.

For further discussion of this form see previous remarks under the genus (p. 184). I may also refer to the remarks accompanying my original diagnosis of the species, but it should be noted that the comparison there made was chiefly with the abnormal female specimen alleged to have come from Taipa and described above, p. 185.

Dimensions.—*Adult male*: type, U.S.N.M., No. 36505; Botel Tobago Island, May, 1899; T. Tada, collector.

	<i>mm.</i>
Total length.....	289
Snout to vent.....	76
Vent to tip of tail.....	213
Snout to center of eye.....	14
Greatest width of head.....	15
Fore leg.....	39
Hind leg.....	70

Habitat.—This form is only known from the island of Botel Tobago, where, according to Mr. Tada, it is abundant.

List of specimens of Japalura mitsukurii.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.
U.S.N.M.....	36505	Male ^a	Botel Tobago Island.....	May —, 1899	T. Tada.
Do.....	36506	Female.....do.....do.....	Do.
Sci. Coll. Tokyo.....	24 a	Male.....do.....do.....	Do.
Do.....	24 b	Female.....do.....do.....	Do.
Do.....	24 c	Male.....do.....do.....	Do.
Do.....	24 ddo.....do.....do.....	Do.
Do.....	24 edo.....do.....do.....	Do.

^a Type, p. 191.

Genus — ?

Species — ?

The young Agamoid lizard designated by Hallowell (Proc. Phila. Acad., 1860, p. 491) simply as "*Otocryptis*," though practically admitting that it does not belong to Wiegmann's genus, is still in the museum (No. 12252). When Hallowell wrote his notes on this specimen, it was already "much mutilated and otherwise injured," and the intervening 47 years have not improved its condition. In fact, several

of the characters noted by him are now obliterated. Yet, what is left seems to show that the specimen does not belong to any now recognized genus and species, and also that Hallowell's description, which I reproduce verbatim below, is fairly good.

The dentition is now the most prominent feature, the comparatively enormous tricuspid molars standing out in bold relief. They are eight in number on each side in both jaws, gradually increasing in size backward; in the upper jaw there are, besides, three "incisors," the median largest, and a small hardly differentiated "canine" on each side of these; in the lower jaw there are three conical "incisors" on each side in front of the molars, the middle tooth on each side being largest. The dental formula consequently looks like this:

$$\frac{8 + 1 + 3 + 1 + 8}{8 + 3 + 3 + 8}$$

For comparison I append that of the type of *Japalura polygonata*:

$$\frac{14 + 2 + 5 + 2 + 14}{14 + 3 + 3 + 14}$$

The small number of molars in both jaws is probably due to the youth of the specimen, as Siebenrock has shown that in some *Agamidae* the number varies between 8 and 21, according to age,^a but the difference in the number of incisors is not so easily explained, for it seems, according to the same authority,^b as if the young animals have even more incisors than the adult, a condition reversed in the *Japalura* and the uncertain young Agamoid before us.

A young specimen of *Japalura swinhonis* (U.S.N.M. No. 34097), from Formosa, though more than once and a half as large as Hallowell's specimen, corroborates the above. In the upper jaw it has nine "molars," two "canines," and four "incisors," consequently essentially like the adult, only fewer molars.

Hallowell's description is as follows:

OTOCRYPTIS Weig.

Among the saurians in the Loo-Choo collection is a very small Acrodont Iguanian, no doubt a young animal, which appears to belong to a genus very near if not identical with the *Otocryptus* of Weigmann. The form of the head, however, is different from that of *Otocryptus*, as described by Dumeril & Bibron, and the number of molar teeth is less, viz.:—eight instead of twelve. The specimen, small as it is, is much mutilated, and otherwise injured, but the following characters can be distinctly made out.

Gen. char. Head rounded, and high posteriorly, the anterior slope more gradual than the posterior, the front and vertex being nearly on a line; snout covered with polygonal scales; supraorbital ridges distinct; eyelids covered with carinated scales of nearly equal size; occiput covered with polygonal scales, partly smooth, partly carinate; temples covered with carinated scales; eyelids covered with small scales.

^a Sitz. Ber. Akad. Wiss. Wien, Math.-Naturw. Cl., CIV, Pt. 1, p. 1116 (Nov., 1895.)

^b Idem, p. 1121.

The incisor teeth on each side separate, a conical canine and eight tricuspid, the first three the smallest, the last two the largest; the same number in the lower jaw; ears concealed by the integument; body covered with scales arranged in transverse rows; five fingers without serratures, the fourth finger a little the longest, all armed with recurved nails; tail slender.

Dimensions. Length of head 4 lines; of neck and body $6\frac{1}{2}$ lines; of tail (mutilated) 10 lines; length of anterior extremities 6 lines; of posterior

Habitat. Loo-Choo. One specimen. Caught Nov. 1854, by McKnight.

In addition to Hallowell's description, I note that the scales on the sides of the neck and body appear to be very small, almost granular.

No lizard corresponding to the above description has been brought from the Riu Kius by later travelers, and the possibility is, therefore, that a most interesting novelty may yet be found in those islands.

Family SCINCIDÆ.

Lizards with scaly tongue and lower surface covered with rounded, imbricate scales.

Distributed over all the warmer and temperate regions of the globe. Many species have an enormous geographic range owing to the ease with which they may be accidentally transported.

Throughout the family there is a strong tendency toward the suppression of the limbs, and although in the higher forms they are well developed, there are numerous genera in which they become rudimentary or even disappear entirely. None of these occur in Japanese or neighboring territory, all the species here represented belonging to six genera, having the normal number of digits.

KEY TO THE GENERA OF SKINKS OCCURRING IN JAPAN AND ADJACENT TERRITORY.

*a*¹ Movable lower eyelid; fronto-parietals and interparietals distinct.

*b*¹ Supranasals present.

*c*¹ Palatine bones separated on the median line..... *Eumeces*, p. 193.

*c*² Palatine bones in contact on the median line..... *Mabuza*, p. 213.

*b*² Supranasals absent.

*c*¹ Fronto-parietals in contact; dorsals smooth or weakly bicarinate; claws not retractile.

*d*¹ Lower eyelid scaly *Sphenomorphus*, p. 216.

*d*² Lower eyelid with an undivided transparent disk.... *Laciotropis*, p. 218.

*c*² Fronto-parietals not in contact; dorsals tricarinate; claws retractile.

Lygosaurus, p. 221.

*a*² No movable lower eyelid; fronto-parietals and interparietal united into a single shield *Cryptoblepharus*, p. 225.

Genus EUMECES^a Wiegmann.

1834. *Eumeces* WIEGMANN, Herpet. Mex., p. 36 (type, *Scincus pavimentatus* = *E. schneiderii*).


1839. *Plestiodon* DUMÉRIE and BIBRON, Erpét. Gén., V. p. 697 (same type).

^a From *εὐμήκης*, elongated.

1852. *Lamprosaurus* HALLOWELL, Proc. Phila. Acad., 1852 (p. 206) (type, *L. guttulatus*).


1871. *Plistodon* COPE, Sec. and Third Ann. Rep. Peabody Acad., 1871 (p. 82) (emendation).

The characters relied upon in the genus *Eumeces* for the proper separation and identification of the species are unfortunately subject to a wide range of variation, at least in some of the species. Such criterions as the absence or presence of a postnasal, of one or two postmentals, the number of scales around the body, etc., are apt to fail when most needed. Yet they are the only ones which the systematist has to rely on, and though he may be convinced that he has to deal with separate forms, nevertheless he may occasionally have to fall back on locality. Take, for instance, *Eumeces elegans* of Boulenger, from China. The only character by which I can distinguish it from *E. latiscutatus*, from Japan, is by the absence of a postnasal. Yet, among



FIGS. 176-178.—HEAD SHIELDS OF SCINCID LIZARDS. 176, SIDE VIEW OF HEAD OF LEILOLOIPISMA; 177, UPPER VIEW OF HEAD OF LEILOLOIPISMA; 178, UPPER VIEW OF HEAD OF EMOIA. *cs*, CHIN-SHIELDS; *d*, DISK ON LOWER EYELID; *e*, EAR-OPENING; *f*, FRONTAL; *fn*, FRONTO-NASAL; *fp*, FRONTO-PARIETAL; *i*, INTERPARIETAL; *il*, LOWER LABIALS; *l*, LOREALS; *lb*, UPPER LABIALS; *m*, MENTAL; *n*, NASAL; *nc*, NUCHALS; *p*, PARIETALS; *pf*, PREFRONTAL; *pn*, POSTNASAL; *r*, ROSTRAL; *sc*, SUPRACILIARIES; *sn*, SUPRANASAL; *so*, SUPRAOCULAR; *t*, TEMPORAL. (ALL FIGURES ENLARGED.)

sixty-two Japanese specimens examined by me, five lack the postnasal on both sides and two on one side. It does not seem to be to the best interest of science to make *E. elegans* a synonym of *E. latiscutatus*. Much less would it do to call the five Japanese specimens *E. elegans* and enumerate both species as occurring together in the island of Hondo. Very likely some of these specimens without postnasals are brothers or sisters of the others which have this shield. Moreover, what are we to do with those which have it on one side but not on the other? It would probably be a fit case for trinominals were we to find a similar proportion of postnasals present in Chinese *E. elegans*, but unfortunately we have not sufficient material of the latter to determine the point. The most conservative proceeding therefore is to leave the two forms their binominal nomenclature at least until we shall know the Chinese form better.



EUMECES LATISCUTATUS. (FROM SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 554.

The use of the following key in determining the skinks within the territory covered by this work must then be subject to the above remarks. The stability of the various characters employed may best be understood by a perusal of the remarks on *variation* following the descriptions of each species.

KEY TO THE JAPANESE SPECIES OF EUMECES.

- a*¹ One unpaired postmental only (normally); upper temporal of the second row largest, wedge-shaped; soles of hind feet granular with only a few large tubercles on heel.
- b*¹ Posterior loreal nearly as high as long, normally in contact with two supralabials.
- c*¹ Postnasal present (normally).
- d*¹ 24 to 26 (exceptionally 28) scales round the middle of the body.
- E. latiscutatus*, p. 195.
- d*² 28 to 32 scales round the middle of the body . . . *E. latiscutatus okadae*, p. 200.
- e*¹ No postnasal. *E. elegans*, p. 202.
- b*² Posterior loreal much longer than high, in contact with three supralabials.
- E. marginatus*, p. 205.
- a*² Two unpaired postmentals; lower temporal of the second row largest, wedge-shaped; soles of hind feet with two series of enlarged tubercles between heel and base of third and fourth toes.
- b*¹ No postnasal; normally two pairs of nuchals. *E. chinensis*, p. 208.
- b*² Normally a postnasal and three pairs of nuchals. *E. kishinouyei*, p. 210.

EUMECES LATISCUTATUS *a* (Hallowell).

TOKAGE.

Plate XV.

1838. *Scincus quinquelineatus* SCHLEGEL, Fauna Japon., Rept., pp. 99, 139; Sauri et Batr., pl. 1, figs. 1-4^b (Leiden Mus.; Siebold and Buerger, collectors) (not of Linnæus).—*Plestiodon quinquelineatum* BLEEKER, Natuurk. Tijdschr. Nederland. Indië, XVI, 1858, p. 204 (Japan).—*Eumeces quinquelineatus* HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 113.—FRITZE, Mitth. Deutsch. Ges. Ost.-Asiens, V, 1891, p. 239 (Yezo).
1860. *Plestiodon latiscutatus* HALLOWELL, Proc. Phila. Acad., 1860, p. 496 (type-locality, Simoda, Japan; collected by the Rodger Exped.).
1864. *Eumeces (Plestiodon) quinquelineatus* var. *japonicus* PETERS, Mon. Ber. Berlin Akad. Wiss., 1864, p. 57 (type-locality, Nagasaki; type in Berlin Mus.; von Martens, collector).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1876, p. 376 (Nagasaki).
1878. *Eumeces (Plestiodon) japonicus* BOETTGER, Offenbach. Ver. Naturk. 17 18 Ber., Mitth., p. 4 (Japan).
1879. *Eumeces japonicus* BOCOURT, Miss. Sci. Mexique, Rept., livr. 6, p. 423.
1887. *Eumeces marginatus* BOULENGER, Cat. Liz. Brit. Mus., III, p. 371 (part: Miyanoshta; Nikko) (not of Hallowell).—OKADA, Cat. Vert. Japan, 1891, p. 70 (part: Tokyo; Hakone; Nikko; Awaji; Suwa).—BOETTGER, Kat. Rept. Mus. Senckenberg., I, 1893, p. 111 (part: Nikko; Yezo).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 860; author's separate, p. 11 (part: Hondo; Yezo).

a Signifying with broad shields.

b Figs. 1, 3, 4, reproduced in this work on Plate XV.


As the name *E. marginatus* belongs to the Riu Kiu species, *Eumeces latiscutatus* of Hallowell must be taken for that of Japan proper. The type appears to be lost, but there can be no doubt as to the pertinency of the name in spite of the fact that Hallowell apparently overlooked the presence of a postnasal. He describes his specimen as having "a plate above the anterior frenal," which, however, he considers an individual abnormality. The fact is that the plate alluded to is the real anterior frenal, or first loreal, while the scale he so designates is only the abnormally enlarged postnasal crowding the loreal out of contact with the supralabial. In specimens from the main islands of Japan this state of affairs is not uncommon (14 out of 47 specimens), and in extreme cases it looks at a first glance as if the postnasal were missing.

Doctor Hallowell does not state the locality in Japan where the type specimen of *E. latiscutatus* was collected, but the only entry in Dr. W. Stimpson's manuscript catalogue which can have reference to this specimen is his "Sm.—15. Lizard found on hillsides at Simoda, Japan, May, 1855. Black above, lineate with light blue. Below pale blue." Simoda may therefore be regarded as the type locality.

One may still occasionally see this species quoted as *E. quinquelineatus*, it being asserted upon Schlegel's authority that there is no difference between the Japanese and the North American species. A careful comparison will show that there exist material differences in the number of scale rows, size, and shape of rostral and nasals, size of median dorsal scales, shape of claws, etc., as already pointed out by Hallowell, Peters, and Boulenger. To their statements I may add that the two species can be very readily distinguished by the differences shown in the relative size and shape of the large temporal shields between the parietals and the seventh labial, the North American species having an arrangement very much like that of the Yaeyama species *E. kishinouyei* as distinguished from *E. marginatus*, *E. latiscutatus*, and their immediate allies and illustrated on figs. 179, 182, 184.

Description (figs. 179–180).—*Adult male*; U.S.N.M. No. 12715; Yokohama; April, 1881; P. L. Jouy, collector. Rostral high, bent over on the snout, the portion visible from above being nearly as large as the fronto-nasal; supranasals in contact behind rostral; nostril in the anterior portion of a single nasal, which is higher anteriorly than posteriorly; a small postnasal in contact with supranasal, nasal, first and second labials (on left side with first labial only) and anterior loreal; fronto-nasal broader than long, not in contact with rostral nor with frontal; prefrontals nearly as large as fronto-nasal in contact behind the fronto-nasal, also in contact with both loreals and with upper preocular and anterior supraocular; anterior loreal high and narrow, in contact with second supralabial (on left side also with first); posterior loreal pentagonal, nearly as high as wide, in contact with

second and third supralabials; frontal as long as parietals, longer than its distance from tip of snout, anteriorly a little wider than behind, in contact with three supraoculars; four supraoculars, second largest; seven superciliaries; fronto-parietals somewhat smaller than interparietal; parietals large, in contact behind interparietal; one pair of nuchals; seven supralabials, seventh largest, fifth under the eye; two temporals in the first row, lower much the larger; upper temporal in the second row very large, much larger than lower one, wedge-shaped with the apex pointing forward; lower temporal of the second row elongate squarish, with upper and lower edges nearly parallel; a single large scale between seventh supralabial and ear-opening which is bordered anteriorly by a few projecting granules; mental small, followed by a single pentagonal, unpaired postmental; seven lower labials, the sixth being band-like, long and narrow; body covered above and below with cycloid scales in 26 rows around the middle, those on the



FIGS. 179-180.—*EUMECES LATISCUTATUS*. 2 X NAT. SIZE. 179, SIDE OF HEAD, NORMAL POSTNASAL; (23 X NAT. SIZE) 180, UNDERSIDE OF HIND FOOT. No. 6166, U.S.N.M.

middle of the back slightly wider than the others, laterals not essentially smaller than ventrals; a pair of large preanal plates; length of hind leg contained twice and a half times in the distance from snout to vent; the adpressed limbs overlap by the length of the fingers; sole of hind foot granular with a single larger tubercle and a few others at the heel; a specialized scale with a claw-like pointed keel behind the vent at each corner; tail cylindric, tapering, slightly swollen underneath at base, with a series of wide transverse plates on the underside. Color (in alcohol), above pale olive gray with a dorso-lateral whitish band about one scale wide, bordered below by a broad dark-brown band from the eye over the ear-opening, above the legs and continued on the tail nearly to the tip; head nearly uniform olive clay color; whole underside, including lower half of flanks, whitish, chin and throat suffused with clay color, the rest with bluish gray.

Dimensions.

	<i>mm.</i>
Total length.....	186
Snout to vent.....	64
Vent to tip of tail.....	122
Snout to ear-opening.....	15
Greatest width of head.....	12
Axilla to groin.....	34
Fore leg.....	18
Hind leg.....	25

The *female* differs in lacking the swelling at the base of the tail and apparently also the differentiated keeled scale behind each corner of the vent.

The *younger* individuals differ in not having the temporal region swollen and also differ greatly in color. As exemplified by a large series from Mount Fuji, they are (in alcohol) nearly blackish brown above, becoming dark brownish slate on the terminal half of the tail, with five pale longitudinal stripes, viz, one median; one on each side proceeding from the outer edge of the supraoculars down the sides of the back, two scale widths from the median stripes; finally, a lateral stripe on each side originating on the supralabials, through the ear and above both fore and hind legs; all continued on the tail where they gradually disappear; the median stripe bifurcates on the interparietal, each branch following the outer edge of the frontal to the snout, which is pale brownish; underside bluish gray, except chin, throat, limbs underneath, anal plate, and the subcaudal transverse plates, which are pale buff. According to Stimpson they are black in life with light blue stripes and pale blue underside.

Variation.—There is considerable variability in the head scales and their various relations, but as it would be of little use to note all the deviations from the specimen described above, it will be sufficient to call attention to those which are of systematic value.

Out of 62 specimens examined by me only one (No. 34123) has the azygous postmental divided, and the abnormality of the specimen is further shown by a small portion of the side of the posterior postmental being also separated by a suture.

In 3 specimens (Nos. 34120, 34130, 34138) the posterior loreal is in contact with three supralabials on both sides; in one (No. 30734) on one side only, and in one (No. 34126) it barely touches the third labial on one side.

Five specimens (Nos. 34119, 34121, 34126, 34127, 34133) have no postnasal at all; two lack it on one side only (Nos. 34128, 34139), while in 37 specimens it is present on both sides.

Curiously enough the variation in the postnasal is most pronounced in a single series of specimens collected near Mount Fuji in September, 1898. Out of the 21 specimens five have no postnasals at all, two on

one side only, and of the fourteen others which have it on both sides, no less than nine have the postnasal so large as to crowd the anterior loreal away from the supralabials, so that only in five does it come in contact with the supralabials behind the postnasal and that not even always on both sides.

Of 39 specimens, the only ones examined for this character, the scale rows around the middle of the body are 28 in two only, 26 in twenty-four, and 24 in thirteen.

Habitat.—Apparently confined to the large islands of Japan proper.

The types came from the hills back of Simoda, and the species is apparently quite common in that part of Hondo. We have a large number from around Mount Fuji and from Yokohama; British Museum has it from Miyanoshta and Nikko, whence specimens are also in the Senckenberg Museum, and Okada records it from Tokyo and Hakone. From farther south the latter also indicates the provinces of Awaji and Suwa; the Hamburg Museum has it from Kiyotaki toge in the province of Kawachi (No. 1833), and the United States National Museum from Koriyama, province of Yamato. From northern Hondo there are no records. From Yezo, however, there is said to be a specimen in the Senckenberg Museum received from Mr. B. Schmacker,^a and Doctor Fritze also indicates it from that island, without giving his authority, however. If occurring there, it must be rare or local. Dr. Hugh M. Smith brought us a fine series from Kochi, on Shikoku, the only record from that island. He also collected it in the extreme southern end of Kiusiu at Yamagawa, province of Satsuma. Our museum also has it from Miyazaki, and von Martens obtained it at Nagasaki, from which locality the Riksmuseum in Stockholm also has a specimen (No. 2319), collected by Dr. Oscar Nordquist in October, 1879, during the *Vega* expedition.

List of specimens of Euneces latiscutatus.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Postmentals.	Postnasal.
6166a	(a)	Japan.....				1	1
6166b	do.....				1	1
6166c	do.....				1	1
6166d	do.....				1	1
6166e	do.....				1	1
12715	(b)	Yokohama.....	Apr., 1881	Dale and Jouy.....	26	1	1
13750	do.....		P. L. Jouy.....		1	1
30734		Miyazaki, Kiusiu.....		C. A. Clark.....	24	1	1
31804		Koriyama, Yamato.....	Apr. 26, 1903	H. M. Smith.....	26	1	1
31832		Yamagawa, Kiusiu.....	June 14, 1903do.....	24	1	1
31833	do.....do.....do.....	26	1	1

^a Figs. 179-180.

^b Description, p. 196.

^a See, however, remarks on alleged Yezo specimens by the same collector, pp. 111, and 275.

List of specimens of Eumeces latiscutatus—Continued.

U.S. N.M. No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Post-mentals.	Post-nasal.
31834	Yamagawa, Kiusiu.....	June 14, 1903	H. M. Smith.....	26	1	1
31835	do.	do.	do.	26	1	1
31836	do.	do.	do.	26	1	1
31837	do.	do.	do.	26	1	1
31879	Fochi, Shikoku.....	May 11, 1903	do.	26	1	1
31880	do.	do.	do.	26	1	1
31881	do.	do.	do.	26	1	1
31882	do.	do.	do.	26	1	1
31883	do.	do.	do.	26	1	1
31890	do.	do.	do.	26	1	1
31891	do.	do.	do.	24	1	1
31892	do.	do.	do.	28	1	1
31893	do.	do.	do.	28	1	1
34119	Mount Fuji.....	Sept., 1898	A. Owston.....	24	1	0
34120	do.	do.	do.	26	1	1
34121	do.	do.	do.	24	1	0
34122	do.	do.	do.	24	1	1
34123	do.	do.	do.	26	2	1
34124	do.	do.	do.	24	1	1
34125	do.	do.	do.	24	1	1
34126	do.	do.	do.	26	1	0
34127	do.	do.	do.	24	1	0
34128	do.	do.	do.	26	1	1-0
34129	do.	do.	do.	26	1	1
34130	do.	do.	do.	26	1	1
34131	do.	do.	do.	24	1	1
34132	do.	do.	do.	26	1	1
34133	do.	do.	do.	26	1	0
34134	do.	do.	do.	26	1	1
34135	do.	do.	do.	24	1	1
34136	do.	do.	do.	26	1	1
34137	do.	do.	do.	24	1	1
34138	do.	do.	do.	24	1	1
34139	do.	do.	do.	26	1	1-0

EUMECES LATISCUTATUS OKADÆ,^a new subspecies.

Diagnosis.—Scales round the body 28–30; first upper labial forming a suture with the nasal and the postnasal; a single postmental.

Habitat.—Miyakeshima, Idzu Seven Islands, Japan (not to be confounded with Miyakoshima, Yaeyama Islands).

Type.—U.S.N.M. No. 23891; Miyakeshima; May 3; N. Okada, collector; topotypes in Science College Museum, Tokyo.

Description of type-specimen.—Rostral heptagonal, rather high, and broad at top, though not bent much over on the upper surface; nasal small with the nostril pierced in front of the center; a small postnasal in contact with first and second supralabial (fig. 181);

^a Named for Mr. S. Okada, the author of the Catalogue of Vertebrated Animals of Japan (Tokyo, 1891).

supranasals broadly in contact, larger than prefrontals; frontonasal longer than wide, behind broadly in contact with frontal; anterior loreal forming a suture with frontonasal; four supraoculars, the three anterior in contact with frontal; six superciliaries; one pair of nuchals in contact with posterior border of parietals and interparietal; seven supralabials, fifth bordering the orbit, seventh largest; one unpaired postmental; ear-opening slightly larger than largest dorsal scales and bordered anteriorly by about 5 distinct obtuse tubercles; 30 scale rows round the middle of the body, the laterals very much smaller than either ventrals or dorsals, the two median rows of the latter slightly larger than the others; two very large preanal scales; length of hind limb (from groin to tip of fourth toe) contained twice in the distance from snout to vent; limbs overlap considerably when pressed against the body; no continuous row of tubercles on soles from heel to base of fourth toe; median series of subcaudals transversely enlarged anteriorly. Color (in alcohol) above uniform dark olive; an obscure brownish band proceeds from eye backward above the limbs disappearing at the base of tail; this band is bordered above by an equally obscure pale band, these bands so obscure as to be imperceptible in the specimen when taken out of the alcohol; under-side bluish, chin and throat whitish.




FIG. 181.—*EUMECES LATISCUTATUS* OKADA. 2 X
NAT. SIZE. SIDE OF FACE.
No. 23895, U.S.N.M.

<i>Dimensions.</i>	<i>mm.</i>
Total length.....	130
Snout to vent.....	56
Snout to ear-opening.....	13
Greatest width of head.....	9
Axilla to groin.....	29
Fore limb.....	20
Hind limb.....	28

Remarks.—The present species is most nearly related to *E. latiscutatus* from the main island opposite, and is probably a direct offspring of the latter. It differs chiefly in the greater number of scale rows round the middle of the body and apparently also in a somewhat different coloration, inasmuch as even young specimens of the island form are of a more or less uniform olive color without the strongly marked pale lines so characteristic of the main island specimens of corresponding size. They are only plainly marked in the smallest specimen from Niishima, measuring 40 mm. from snout to vent, and even in this specimen the median line does not reach the head.

The additional number of scale rows makes the diagnosis of the new species read even more like that of *E. fasciatus* of the United

States of North America than *E. latiscutatus*, but aside from the single unpaired postmental common to both the Japanese species, the shape of the rostral as well as the position of the nostril in the nasal in the latter are characters in themselves sufficient to distinguish them from their North American relative, not to speak of the difference in the temporal scales as mentioned above.

The segregation of the present form from that of the main island is no more remarkable than that of the very distinct thrush, *Turdus celanops*, of the same island, from *T. chrysolaus* of the mainland, as well as *Acanthopneuste ijimæ* from *A. coronata*, and *Parus owstoni* from *P. varius*.

Doctor Ijima informs me that he has five more specimens of the same form "all of the same or similar coloration."

Habitat.—This form probably occurs on all the islands of the Seven Islands Archipelago of Idzu, but I have only seen specimens from Miyakeshima and Niishima.

List of specimens of Eumeces okada.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scales.		
						Snout-vent.	Postmentals.	Postnasal.
U.S.N.M.	23891	(a)	Miyakeshima, Idzu.	May 3, 1887	Okada	30	1	1
Do.	23892		do.	do.	do.	30	1	1
Do.	23895	(b)	Niishima, Idzu.		do.	28	1	1
Do.	23896		do.		do.	28	1	1
Do.	36533		do.		do.	30	1	1
Do.	36531		do.		do.	28	1	1
Do.	36532		do.		do.	28	1	1
Do.	36534	(c)	Miyakeshima, Idzu.	May 3, 1887	do.	30	1	1
Sci. Coll. Tokyo.			do.		Hirota	28	1	1

^a Type; description, p. 200.

^b Sci. Coll. No. 46, fig. 181.

^c Sci. Coll. No. 47.



EUMECES ELEGANS^a Boulenger.

1863. *Mabouia chinensis* SWINHOE, Ann. Mag. Nat. Hist. (3), XII, p. 225 (Tamsui, Formosa) (not of Gray 1838).—GUENTHER, Rept., Brit. India, 1864, p. 83 (part), pl. x, fig. *F* (Ningpo).—BOETTGER, Offenbach, Ver. Naturk. 24-25 Ber., 1885, p. 144.
1879. *Eumeces pulchra* BOCOURT, Miss. Sci. Mexique, Zool., Rept., livr. 6, p. 423 (not of Duméril and Bibron).
1887. *Eumeces elegans* BOULENGER, Cat. Liz. Brit. Mus., III, p. 371 (type-locality, Ningpo, China; type in Brit. Mus.; Formosa, Shanghai, Kiukiang, Pescadores).—BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 146, Chapu, near Ningpo).—STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 220 (Taipa, Formosa; Pescadores).—*Lygosoma elegans* GUENTHER, Ann. Mag. Nat. Hist. (6), I, 1888, p. 169 (Mts. north of Kiukiang, China).

^a Signifying elegant.

A direct comparison of typical *Eumeces elegans* from China (which our museum owes to the kindness of Dr. O. Boettger) with *E. latiscutatus*, from Japan, and *E. marginatus*, from the Riukius, shows that it is more nearly related to the former than to the latter, the second loreal being as high and short as in the most typical *E. latiscutatus*. In fact, the only difference I can see consists in the presence of a postnasal in the latter, in this respect agreeing with *E. marginatus*. The latter I regard as less closely related to the other two than these are among themselves. For further remarks on this subject see above (p. 194).

Description (figs. 182-183).—*Male*; U.S.N.M. No. 36507 (Science College Museum, Tokyo, No. 20a); Taipa, northern Formosa; October, 1896; T. Tada, collector. Rostral high, rather narrow, the portion visible from above slightly smaller than fronto-nasal; supranasals broadly in contact behind rostral; nostril in the anterior portion of a single nasal which is much higher in front than behind; no postnasal; fronto-nasal much broader than long, not in contact with rostral, but with frontal; prefrontals smaller than fronto-nasal, not in contact with each other, but with both loreals, upper preocular and anterior supraocular; anterior loreal higher than the posterior, in contact with supranasal, nasal, first and second supralabials; posterior loreal pentagonal, as high as wide, in contact with second and third supralabials (on left side also barely touching fourth); frontal longer than parietals, longer than its distance from tip of snout, anteriorly a little wider than behind, in contact with three supraoculars; four supraoculars, second largest; six or seven superciliaries; fronto-parietals smaller than interparietal, which is as long as the parietals and broadly in contact behind with the nuchals; parietals not in contact behind the interparietal; one pair of nuchals; seven supralabials, seventh largest, fifth under the eye; two temporals in first row, lower many times larger than the upper; upper temporal in the second row very large, much larger than the lower one, wedge-shaped, or conical, with the apex forward; lower temporal of the second row long and narrow with nearly parallel upper and lower edges; a single scale row between seventh supralabial and ear-opening, which is bordered anteriorly by several projecting lobules; mental followed by a single pentagonal, unpaired postmental; seven lower labials, sixth longest; 28 scale rows around the middle



FIGS. 182-183.—EUMECES ELEGANS. 2X NAT. SIZE. 182, SIDE OF HEAD; 183, CHIN. No. 20B, SCI. COL., TOKYO.

of the body, those on the sides apparently smaller than those on back and abdomen; a pair of large preanal plates; length of hind leg contained nearly two and a half times in the distance from snout to vent; the adpressed limbs overlap by the length of the fingers; soles of hind foot granular with a single larger tubercle and several others at the heel; a small keeled scale behind the vent at each corner; tail cylindrical, tapering, with a series of wide transverse plates underneath. Color (in alcohol) above dark olive brown on the middle of the back, dark brown on the sides, becoming slaty toward the terminal half of the tail, with five pale greenish, longitudinal stripes, viz., one median, somewhat broader than the others; one on each side proceeding from the outer edge of the supraoculars down the sides of the back two scale widths from the median stripe; finally, a lateral stripe on each side originating on the upper lip proceeding backward through the ear-opening and above both fore and hind legs, all continued on the tail, where they gradually disappear; the median stripe bifurcates on the interparietal, each branch following the outer edge of the frontal to the snout which is pale buff; underside pale, suffused with buff on throat, lower neck, limbs, and tail, but with pale bluish on abdomen and flanks.

Dimensions.

	<i>mm.</i>
Total length.....	179
Snout to vent.....	68
Vent to tip of tail.....	111
Snout to ear-opening.....	13.5
Greatest width of head.....	10
Axilla to groin.....	35
Fore leg.....	19
Hind leg.....	28

Variation.—In younger specimens the ground color is darker, brownish black, and the median dorsal stripe narrower; otherwise as the specimen described above. In older specimens the back becomes more olive and the median stripe disappears altogether, the head swells greatly in the temporal region and becomes red. A male from the Pescadores Islands, collected by Mr. Tada in January, 1897, has this coloration of the fully adult specimens, the brown longitudinal band on the flanks even being quite pale, notwithstanding the fact that it is of the same size (snout to vent, 68 mm.) as the adolescent specimen described above.

The scutellation of the head in these specimens is fairly constant. Thus all four have the parietals widely separated by the interparietal. In three the frontal is in contact with the fronto-nasal, but in the fourth, the largest specimen, the prefrontals are in contact behind the fronto-nasal, thus separating it from the frontal. The number of scale rows seems to be very variable, as two have 26, one 24, and one 28.

Habitat.—Swinhoe collected two half-grown specimens in Formosa, which are in the British Museum. The Science College, Tokyo, has several striped specimens from Taipa, Formosa (No. 20), collected in October, 1896, and one nearly uniform specimen (No. 19) from the Pescadores Islands, January, 1897.

The species is known from the mainland of China between Shanghai and the province of Fokien as far inland as Kiukiang, on the Yangtse River.

List of specimens of Eumeces elegans.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Postmentals.	Postnasal.
U.S.N.M.	36507	(a)	Taipa, Formosa	Oct., 1896	T. Tada	28	1	0
Sci. Coll. Tokyo	20	(b)	do	do	do	24	1	0
Do.	20		do	do	do	26	1	0
Do.	19	(c)	Pescadores Islands	Jan., 1897	do	25	1	0

^a Description, p. 204.

^b Figs. 182-183.

^c P. 205.

EUMECES MARGINATUS ^a (Hallowell).

1860. *Plestiodon marginatus* HALLOWELL, Proc. Phila. Acad., 1860, p. 492 (type-locality, "Loo Choo Island;" type, U.S.N.M. No. 11713; W. Stimpson, collector).—*Eumeces marginatus* BOULENGER, Cat. Liz. Brit. Mus., III, 1887, p. 371 (part: Nara); Ann. Mag. Nat. Hist. (6), X, Oct. 1892, p. 302 (Okinawa; Holst, collector).—BOERTGER, Ber. Senckenberg. Naturf. Ges., 1888, Abh. p. 188 (O-shima, Riu Kiu).—OKADA, Cat. Vert. Japan, 1891, p. 70 (part: Okinawa).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 860; author's separate, p. 11 (part: Okinawa).—BROWN, Proc. Phila. Acad., 1902, June 11, p. 185 ("Loo Choo Island;" probably Okinawa shima).

The larger of Hallowell's two types, said to have been collected on "Ousima" (i. e., Amami-o-shima), appears to have been lost, and as it would be utterly impossible from his description now to decide one way or the other, all we can do is to accept the other specimen, viz, the one which Dr. W. Stimpson collected in "Loo Choo Island" April, 1855, and which is now before me, United States National Museum No. 11713, as *the* type upon which to base the name.

This specimen agrees with numerous specimens from Okinawa shima in every respect and a comparison with over sixty specimens from Japan proper proves them to be unquestionably distinct. Not only do the Okinawa specimens lack the postnasal, a character which is usually conceded to be of sufficient importance, but there are material differences in other points, notably in the size and shape of the large second loreal. This shield in the Riu Kiu specimens is much lower and longer with less difference in the length of the ante-

^a Having a margin or edge; application obscure, but probably the word is here used loosely as meaning striped.

rior and the posterior borders, and is in contact with three supralabials, viz, the second, third, and fourth. In all specimens from Japan proper I have found the second loreal much shorter and higher, with the anterior border much higher than the posterior one, and except in a few cases mentioned above, only in contact with two supralabials, viz, the second and third (if there are seven supralabials in all, but with the third and fourth in exceptional cases of eight supralabials).

Description.—*Adult male*; U. S. N. M. No. 11713; "Loo Choo Island" = Okinawa shima; April, 1855; W. Stimpson, collector; type




FIG. 184.—*EUMECES MARGINATUS*.
2 X NAT. SIZE. SIDE OF HEAD.
No. 11713, U.S.N.M.

(fig. 184).—Rostral pentagonal, the portion visible from above much smaller than fronto-nasal; supranasals in contact behind rostral; nostril in the anterior portion of a single nasal which is somewhat higher in front than behind; no postnasal; fronto-nasal somewhat broader than long, not in contact with rostral, but broadly so with frontal; prefrontals much smaller than fronto-nasal, in contact with both loreals, upper preocular and anterior supraocular, but not with each other; anterior loreal high and narrow, in contact with nasal, first and second supralabials; posterior loreal pentagonal, much wider than high, in contact with second, third, and fourth supralabials; frontal longer than parietals and much longer than its distance from tip of snout, anteriorly wider than behind, in contact with three supraoculars; four supraoculars, none unusually developed; fronto-parietals scarcely more than half as large as interparietal; interparietal as long as parietals, broadly in contact behind with nuchals; parietals rather small, not in contact; one pair of nuchals; seven supralabials, seventh largest, fifth under the eye; two temporals in first row, lower one several times larger than upper one; two temporals in second row, the upper one largest, wedge-shaped with the apex forward, the lower narrower with nearly parallel upper and lower edges; a single large scale between seventh supralabial and ear-opening which is bordered anteriorly by several projecting lobules; mental followed by a single narrow, pentangular, unpaired post-mental; seven lower labials, sixth very long and narrow; 26 scales around the middle of the body, the lateral but very slightly smaller than the others; a pair of large preanal plates; length of hind leg contained two and a half times in the distance from snout to vent; the adpressed limbs overlap by the length of the longest toe; sole of hind foot with small scales or granules, but no enlarged tubercles except a few at heel; a strongly keeled scale behind the vent at each corner; tail cylindric, swollen at base underneath, with a series of wide

transverse plates on the underside. Color (in alcohol) above pale olive gray with faint indications of a median and two dorso-lateral pale stripes as well as of a darker brownish band on upper half of flanks; head yellowish buff (probably red in life); underside pale buffy white, abdomen and flanks grayish.

Dimensions.

	<i>mm.</i>
Total length.....	138
Snout to vent.....	65
Vent to tip of tail.....	74
Snout to ear-opening.....	14
Greatest width of head.....	11
Axilla to groin.....	35
Fore leg.....	19
Hind leg.....	26

The *female* and *younger* stages are exactly like the corresponding ones in *Eumeces latiscutatus* and *E. elegans*.

Variation.—In the series of nine specimens before me the variation is very slight. All lack the postnasal; all have one postmental only, supranasals in contact behind rostral; fronto-nasal in contact with rostral, parietals separated by interparietal, and in all the second loreal is in contact with three supralabials. All have 26 scale rows, except No. 34185 which has 28.

No. 23894 presents a very unusual abnormality in the scutellation of the upper parts of the head, in as much as the fronto-parietals have been fused in such a way with the surrounding scutes that the frontal has come into contact with the fourth supraocular, parietals, and interparietal, the latter shield separating the parietals in their entire length.

Habitat.—This skink as far as known is confined to the Riu Kiu islands, and if our surmise as to Hallowell's second specimen is correct, it occurs in all three groups of the archipelago.^a In Okinawa it appears from all accounts to be common, and in the Yaeyama group it has been collected both by Doctor Warburg and by Doctor Lenz on Iriomote Island (Hamburg Mus. Nos. 1182 and 1900). Moreover, the United States National Museum has a specimen from Ishigaki shima (No. 34185).

^a Boettger also refers to *Eumeces marginatus* from O-shima, but as he did not distinguish between *E. marginatus* and *E. latiscutatus* no conclusion can be drawn from that record without a reexamination of the specimen.

List of specimens of *Eumeces marginatus*.

Museum.	No.	Age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Postnasal.	Postnasal.
						mm	mm	mm
U.S.N.M.	11713	Adult a ..	Okinawa shima	Apr. —, 1855	W. Stimpson ..	26	1	0
Do.	23893	do	do	26	1	0
Do.	23894	Adollesc. b	do	1887	Tashiro	26	1	0
Do.	34185	Adult	Ishigaki shima	May —, 1899	A. Owston	28	1	0
Do.	36522	Adollesc. ..	Okinawa shima	1887	Tashiro	26	1	0
Do.	36523	do	do	do	do	26	1	0
Do.	36524	Young	do	do	do	1	0
Do.	36525	do	do	do	do	1	0
Sci. Coll. Tokyo.	3	Adollesc. ..	Shuri, Okinawa	July —, 1895	Shiraishi	26	1	0

a Type; description, p. 206, fig. 184.

b P. 207.

EUMECES CHINENSIS^a (Gray).

1838. *Tiliqua chinensis* GRAY, Ann. Nat. Hist., II, Dec., 1838, p. 289 (type-locality, China).—*Mabouia chinensis* SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 225 (Tamsui, Formosa).—*Eumeces chinensis* BOULENGER, Cat. Liz. Brit. Mus., III, 1887, p. 375 (Ningpo; Chusan; Si Kiang; Canton; Hongkong).—BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, pp. 132 (Hainan), 143 (Shanghai), 146 (Dalanshau and Chinhai, near Ningpo).—STREJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 220 (Taipa, Formosa).

1839. *Plestiodon sinense* DUMÉRIL and BIBRON, Erpét. Gén., V, p. 704 (Canton; Paris Mus.; Gernaert, collector).

1839. *Plestiodon pulchrum* DUMÉRIL and BIBRON, Erpét. Gén., V, p. 710 (China).

1842. *Tiliqua rufo-guttata* CANTOR, Ann. Mag. Nat. Hist., IX, 1842, p. 482 (type-locality, Chusan; type in Brit. Mus.; Cantor, collector).

Description.—*Halfgrown*; Science College Museum, Tokyo, No. 21; Taipa, Formosa; October, 1896; T. Tada, collector (fig. 185). Por-




FIG. 185.—EUMECES CHINENSIS. 2
× NAT. SIZE; SIDE OF HEAD. NO.
21, SCI. COLL. TOKYO.

tion of nostril visible from above considerably smaller than fronto-nasal; supranasals broadly in contact behind rostral; nostril in the center of a single nasal which is nearly triangular; no postnasal; first supralabial in contact with supranasal behind nasal; fronto-nasal somewhat broader than long, not in contact with rostral but with frontal; prefrontals smaller than fronto-nasal, in contact with anterior supraocular, upper preocular, and anterior loreal, but not with each other; anterior loreal nearly twice as high as wide, squarish, in contact with first and second supralabials (on left side also with third), supranasal, fronto-nasal, prefrontal, and upper preocular, but not with nasal; posterior loreal low, small, not larger than

^a Signifying Chinese.

anterior, nearly twice as wide as high, in contact with second, third, and fourth supralabials (on left side only with second and third), anterior loreal, and upper preocular, but not with prefrontal; upper preocular greatly elongated, extending along the entire upper edge of the posterior loreal so as to enter into broad contact with the anterior loreal; frontal much longer than parietals, longer than its distance from tip of snout, anteriorly somewhat wider than behind, in contact with three supraoculars, the sutures with first and third being very small; four supraoculars, first smallest, third very narrow, second unusually large; fronto-parietals larger than interparietal, slightly larger than prefrontals; interparietal pointed, behind separating the parietals which are comparatively short, much shorter than frontal; 2 pairs of nuchals; seven supralabials, fifth under the eye, seventh longest; but very low behind; two temporals in first row, lower much the larger; two temporals in second row, the upper rather narrow and elongated with parallel edges, the lower much larger, wedge-shaped with the apex forward; two scales between seventh supralabial and ear-opening, which is bordered anteriorly by three small lobules; mental small, followed by two unpaired postmentals, the anterior narrow, with parallel edges, the posterior larger, pentagonal; 7 lower labials; 24 smooth scales around the middle of the body, laterals not appreciably smaller; a pair of large preanal plates; length of hind leg contained two and two-thirds times in distance from snout to vent; the adpressed limbs overlap by about the length of the fingers; sole of hind foot granular, with two series of enlarged tubercles from heel to base of third and fourth toes; tail with a median series of transverse plates underneath. Color (in alcohol) above pale clay-color, with a network of brownish black color, each scale being marked with the dark and the light color in such a way as to bring out a median dorsal and on each side a dorso-lateral pale black-edged stripe; flanks and sides of neck similarly colored, but with about three irregular series of whitish spots instead of stripes; head above pale clay-color, with traces of the pale dorso-lateral stripe from supraoculars backwards, each shield outlined posteriorly or superiorly with blackish brown; underside uniform whitish.

Dimensions.^a

	<i>mm.</i>
Snout to vent	72
Snout to ear-opening	14.5
Greatest width of head	10.5
Axilla to groin	38
Fore leg	18.5
Hind leg	27

^a Tail missing.

Variation.—The specimen described above is probably abnormal in the exclusion of the anterior loreal from the nasal. In the descriptions of Chinese specimens 26 scale rows are sometimes indicated. In adult specimens the adpressed limbs are described as not overlapping.

The *young* are described as "dark brown above, with three longitudinal light bands on the back, the median broadest and not bifurcating anteriorly."

Habitat.—Eastern China from the island of Hainan northwards to Shanghai. It has also been recorded from Formosa. Swinhoe found it at Tamsui, and an apparently somewhat abnormal specimen collected by Mr. Tada at Taipa has been described and commented upon above.

List of specimens of Eumeces chinensis.

Museum.	No.	Age.	Locality.	When collected.	By whom collected.	Scale rows.	Postnasal.	Postnasal.
Sci. Coll. Tokyo.	21	Adolesc. ♀.	Taipa Formosa.	Oct. —, 1896	T. Tada.	24	2	0

^a Description, p. 208; fig. 185.


EUMECES KISHINOUEI^a Stejneger.

1901. *Eumeces kishinouyei* STEJNEGER, Proc. Biol. Soc. Washington, XIV, Dec. 12, 1901, p. 190 (type-locality, Miyakoshima, Sakishima group, Riu Kiu; type Sci. Coll. Mus. Tokyo, no. 22).

Description.—*Adult*: Science College Museum, Tokyo, No. 22; Miyakoshima, Riu Kiu Archipelago; Tashiro, collector, type (figs. 186–189).—Rostral small, nearly triangular, the portion visible from above much smaller than fronto-nasal; supranasals in contact behind rostral; nostril in the center of a single nasal; a narrow, pentangular postnasal, in contact with supranasal, first and second supralabials; fronto-nasal broader than long, not in contact with rostral, but with frontal; prefrontals somewhat smaller than fronto-nasal, in contact with both loreals, upper preocular and anterior supraocular, but not with each other; anterior loreal about half the size of the posterior, in contact with second supralabial, supranasal, fronto-nasal, and prefrontal; posterior loreal nearly as high as anterior, and twice as large, in contact with second and third supralabials, anterior loreal, prefrontal, and upper preocular; upper preocular smaller than anterior supraocular, widely separated from anterior loreal; frontal much longer than parietals and its distance from tip of snout, with parallel lateral edges being of equal width in front and behind, in contact

^a Named in honor of Dr. K. Kishinouye, chief of the Fisheries Bureau, Imperial Department of Agriculture and Commerce, Tokyo.

with anterior three supraoculars, the suture with third supraocular very small; four supraoculars, second largest; fronto-parietals somewhat smaller than interparietal, about the same size as prefrontals; interparietal obtusely pointed behind, separating the parietals, which are comparatively small, much shorter than frontal; three pairs of nuchals; seven supralabials, fifth under the eye, seventh very large, but low behind; two temporals in first row, lower many times larger than upper; two temporals in second row, the upper rather narrow,



FIGS. 186-189.—*EUMECES KISHINOUEI*. $1\frac{1}{2} \times$ NAT. SIZE. 186, TOP OF HEAD; 187, SIDE OF HEAD; 188, UNDERSIDE OF HEAD ($2 \times$ NAT. SIZE); 189, UNDERSIDE OF FOOT. No. 22, SCI. COLL. TOKYO.

with slightly diverging upper and lower edges, the lower much larger, wedge-shaped, with the apex forward; a single, rather long scale between seventh supralabial and ear-opening, which is bordered anteriorly by three small lobules; mental small, followed by two unpaired postmentals, the anterior very narrow, band-like, the posterior pentagonal; 24 scales round the middle of the body, nearly subequal, each dorsal scale with about four distinct striæ; a pair of large preanal plates; length of hind leg contained about two and two-

thirds times in distance from snout to vent; the adpressed limbs overlap by about half the length of the fingers; sole of hind foot with two somewhat irregular series of enlarged tubercles from heel to base of third and fourth toes; tail with a series of transversely enlarged plates underneath. Color (in alcohol) above nearly uniform olive, head more clay color (in life probably red), underneath whitish.

Dimensions.

	<i>mm.</i>
Total length (tip of tail reproduced).....	259
Snout to vent.....	148
Snout to tip of tail (tip reproduced).....	111
Vent to ear-opening	31.5
Greatest width of head.....	24
Axilla to groin.....	71
Fore leg.....	38
Hind leg.....	55

The type described above is in very poor state of preservation and is probably a female. A fully adult *male* in excellent condition (U.S.N.M. No. 34080) from Ishigaki shima agrees in all essential details with the type, except that the head is considerably more swollen at the temples. This specimen measures: Total length, 357 mm.; snout to vent, 159 mm.; vent to tip of tail, 198 mm.; snout to ear, 39 mm.; greatest width of head, 34 mm.

I find in none of these specimens any specially differentiated, keeled scale behind the vent near each corner as in *Eumeces latiscutatus* and its allies.

The half-grown specimens have a median dorsal pale band bordered with dark-brown lines, a dorso-lateral whitish stripe on each side similarly bordered above, and with a broad brown lateral band, below which again a whitish line.

A *young* specimen (Science College Museum, Tokyo, No. 35), from Iriomote shima, measuring 60 mm. from tip of snout to vent, differs in having the frontal shield somewhat broader in front than behind. The striae on the dorsals show almost as plainly as in the adult. It has three pairs of nuchals, but is abnormal in lacking a postnasal. The coloration is indicated below.

Variation.—I have six very large adult and adolescent specimens before me, two from Ishigaki shima, two from Iriomote shima, and two from Miyakoshima. In scutellation they are practically identical, and the postnasal is present and well developed in all. Its presence may therefore be regarded as normal in this species. Five of them in addition have three nuchals, three specimens have them on both sides, two on one side, while the sixth has two only on both sides (No. 34081); there is consequently reason to believe that three pairs is the normal number. In addition to these full-grown and nearly full-grown specimens, there is a seventh, quite young, black with yellowish stripes. It

agrees in every structural character with them, except in possessing no postnasal, thus closely approaching *E. chinensis*. As it comes from Iriomote shima I have but little doubt that it is an abnormal specimen of *E. kishinouyei*, especially as it has the dorsal scales striated and three pairs of nuchals, while *E. chinensis* normally has but two. Moreover, it seems to differ somewhat in coloration from true *E. chinensis*, at least as it is described by Boulenger.^a He states that the young is "dark brown above, with three longitudinal light bands on the back, the median broadest and not bifurcating anteriorly; sides with light spots," while in the Yaeyama specimen the spots on the sides have united to form two additional longitudinal bands on each side, thus making seven in all, the median not broader than the others and distinctly bifurcating.

Remarks.—This form, which assumes gigantic proportions, our specimen (No. 34080) measuring 159 mm. from snout to vent, is very closely allied to *Eumeces chinensis*, from which it differs chiefly in the presence of a postnasal, three pairs of nuchals, and possibly in the coloration of the young.

Habitat.—Apparently confined to the southern group of the Riu Kiu chain of islands. I have examined specimens from all the principal islands, namely Miyakoshima, Ishigaki shima, and Iriomote or Nishiomote shima.

List of specimens of Eumeces kishinouyei.

Museum.	No.	Age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Postnasal.	Nuchals.
U. S. N. M.	34080	Adult ^a	Ishigaki shima, Riu Kiu	June —, 1899	A. Owston	24	2	1 3-3
Do.	34081	do.	do.	do.	do.	26	2	1 2-2
Sci. Coll. Tokyo.	22	do. ^b	Miyakoshima.		Tashiro	24	2	1 3-3
Do.	22	Halfgr.	do.		do.	26	2	1 3-2
Do.	23	do.	Iriomote shima.				2	1 3-3
Do.	24	do.	do.			24	2	1 3-2
Do.	35	Young ^a	do.		Tashiro	26	2	0 3-3
Hamburg.	1601		Miyakoshima.		Dr. Warburg.	24	2	1

^a P. 212.

^b Type: description, p. 210 figs. 186-189.

Genus MABUYA ^b Fitzinger.

1826. *Spondylurus* FITZINGER, Neue Class. Rept., p. 23 (type, *Scincus sloanci* Daudin).

1826. *Mabuya* FITZINGER, Neue Class. Rept., p. 23 (type, *M. carinata* Fitzinger).

1830. *Euprepis* WAGLER, Nat. Syst. Amph., p. 161 (type, *Scincus multifasciatus* Kuhl).

^a Cat. Liz. Brit. Mus., III, p. 375.

^b A Caribbean name attributed to several lizards.

1834. *Euprepes* WIEGMANN, Herpet. Mex., p. 36 (emendation).
 1845. *Mabouya* GRAY, Cat. Liz. Brit. Mus., p. 93 (emendation).
 1862. *Mabuia* COPE, Proc. Phila. Acad., 1862, p. 185 (emendation).

With reference to the names cited in the above synonymy it is true that *Spondylurus* occurs before *Mabuya*, but of two names having the same date that one selected by the reviser is to be retained. *Mabuya* has been so selected, not only by the first reviser, but by all subsequent authors. This rule is a direct application of article 28 of the International Rules of Zoological Nomenclature.

MABUYA LONGICAUDATA ^a (Hallowell).

Plate XVI.

1856. *Euprepis longicaudata* HALLOWELL, Trans. Amer. Philos. Soc. (2), XI, p. 77, pl. IV, fig. 1 [*Euprepes* on plate] (type-locality, Siam; type in Mus. Phila. Acad.; Doctor Ruschenberger, collector).—*Mabuia longicaudata* BOULENGER, Cat. Liz. Brit. Mus., III, 1887, p. 189.
 1864. *Eumeces siamensis* GUENTHER, Rept. Brit. India, p. 91 (type-locality, Siam; type in Brit. Mus.).—*Mabuia siamensis* BOULENGER, Cat. Liz. Brit. Mus., III, 1887, p. 188 (Siam; Hainan).
 1867. *Euprepes (Tiliqua) bicarinatus* PETERS, Mon. Ber. Berlin Akad. Wiss., 1867, p. 22 (type-locality, Hongkong; type in Berlin Mus.; Jagor, collector).—*Tiliqua bicarinata* MUELLER, Verh. Naturf. Ges. Basel, VIII, Pt. 2, 1887, p. 287 (South Formosa).
 1886. *Euprepes (Tiliqua) ruhstrati* FISCHER, Abh. Naturw. Ver. Hamburg, IX, no. 6, p. 7, pl. I, figs. 2a-d^b (type-locality, South Formosa; type in Oldenburg Mus.; Ruhstrat, collector).


It turns out, as has long been suspected, that Guenther's *Eumeces siamensis* is the same as Hallowell's *Euprepis longicaudata*, described eight years earlier, both from Siam. The separation of the two was undoubtedly due to Hallowell's description, which states that the lower eyelid is "scaly with a transparent disc." Mr. James A. G. Rehn, of the Philadelphia Academy Museum, whom I requested to examine the type for me, writes to the contrary: "The lower eyelid of the type of *Euprepis longicaudata* appears to have a series of four subquadrate scales of rather large size, surrounded by much smaller scales." This removes any doubt concerning the identity of the alleged two species.

There being no specimens of this species accessible to me, I have translated and condensed Doctor Fischer's original description of the type of *Tiliqua ruhstrati* from Formosa. The identity of the latter with the Siamese species is on the authority of Doctor Boulenger.

Description of type specimen.—*Adult*; Oldenburg Museum; South Formosa; Ruhstrat, collector.—Lower eyelid with a row of four to five large quadrangular scales in the middle; the azygos internasal somewhat broader than long, barely touching the rostral and widely sepa-

^a Signifying Long-tailed.

^b Reproduced in this work as Plate XVI.



MABUYA LONGICAUDATA. (FROM FISCHER.)

FOR EXPLANATION OF PLATE SEE PAGE 554.

rated from the frontal by the prefrontals, which are broadly in contact; frontal shield half as wide as long; two fronto-parietals broadly in contact medially; interparietal of same length as fronto-parietals, separating the parietals completely; one pair of nuchals; supraorbitals four, the second in contact with the entire outer edge of the frontal, as large as third and fourth together; nasal elongate, with a semicircular groove behind the nostril; supranasals narrow, barely separated by the internasal; anterior loreal quadrangular, slightly longer than high, in contact with second and third supralabial; second loreal in contact with third supralabial only; seven supralabials, third twice as long as second, fourth largest with parallel vertical edges, situated under the eye; seventh small, scale-like; a single unpaired postmental; ear-opening moderate, with projecting scales overhanging the anterior border; scales in 28 rows around the body, and 26 in a line between axilla and groin, dorsals with two faint keels; a series of preanal scales about twice as large as the preceding series; adpressed limbs overlap with their palms and soles, tip of fourth toe falling considerably short of the axilla. Color above chestnut brown with seven black^a longitudinal lines extending down the back along the edges of the scales; a broad black lateral band, sprinkled with a few white spots, from the loreal region, through eye and over ear and shoulder to the beginning of the tail, above narrowly edged with white; below the lateral band scattered black spots; underside greenish yellow.

Dimensions.

	<i>mm.</i>
Tip of snout to vent	100
Tip of snout to posterior edge of ear-opening	23
Fore leg	31
Hind leg	43 ^b

Remarks.—Boulenger gives the following measurements of one of the specimens in British Museum: Total length, 340 mm.; tail, 225 mm. His description, apparently based on Siamese and Hainan specimens, agrees in most essentials with Fischer's as given above, the main discrepancy being that the ear-opening is "without projecting lobules," while Fischer repeatedly states the opposite ("Ohroeffnung maessig, ihr Vorderrand von vorragenden Schuppen ueberragt"). Mr. Rehn kindly informs me that in the type of Hallowell's *E. longicaudata* he found "that no projecting scales are present on the anterior border of the ear-opening." The question then arises whether the Formosan specimens may not represent a distinct species after all. The next question, then, is to which Peters' *E. bicarinatus* from Hongkong belongs. The original description of Peters contains no reference to the scales surrounding the ear-opening.

^a In the diagnosis "brown."

^b Fischer, Abh. Naturw. Ver. Hamburg, IX, no. 6, pp. 7-8.

Habitat.—Southeastern Asia, from Siam to southern China, at least as far as Hongkong, including the islands of Hainan and Formosa.

From the latter it is known through the type specimen of *Euprepes ruhstrati*, which was collected by Herr Ruhstrat in the southern part of the island, a specimen in the Basel Museum recorded by F. Mueller, and a specimen in the British Museum, collected by Mr. La Touche at Bangkimtsing.

Genus SPHENOMORPHUS^a Fitzinger.

1843. *Sphenomorphus* FITZINGER, Syst. Rept., p. 23 (type, *Lygosoma melanopogon* Duméril and Bibron.)
 1845. *Hinulia* GRAY, Cat. Liz. Brit. Mus., p. 74 (type, *H. navia*=*L. melanopogon*).
 1845. *Otosaurus* GRAY, Cat. Liz. Brit. Mus., 93 (type, *O. cunningii*).

Only one species of this genus has hitherto been found in the territory included in this work.

SPHENOMORPHUS INDICUS^b (Gray).

Plate XVII, figs. 1-2.


1853. *Hinulia indica* GRAY, Ann. Mag. Nat. Hist. (2), XII, Dec., 1853, p. 388 (type-locality, Sikkim, Himalayas; types in Brit. Mus.; Sir J. Hooker, collector).—*Lygosoma indicum* BOULENGER, Cat. Liz. Brit. Mus., III, 1887, p. 241, pl. XVI, figs. 1-1^c (Eastern Himalayas, Assam, Burma); Fauna Brit. India, Rept., 1890, p. 195; Proc. Zool. Soc. London, 1899, p. 162 (Kuatun, prov. of Fokien).—GUENTHER, Ann. Mus. Zool. St. Pétersbourg, I, 1896, p. 203 (Sungpan, Lun-ngan-fu, prov. of Sze-Chuen, China).
 1893. *Lygosoma (Liolepisma) laterale* var. *reversi* BOETTGER, Kat. Rept. Mus. Senckenberg., I, p. 104 (part: Dalanshan, near Ningpo) (not of Gray).
 1894. *Lygosoma (Hinulia) indicum* BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 145 (Dalanshan).

Description.—*Young*: British Museum No. 99. 4. 24. 101; Bangkimtsing, Formosa; J. D. La Touche, collector. Rostral large, broadly in contact with fronto-nasal, which is in contact with anterior loreal, prefrontals, and broadly with the frontal; no supranasals; frontal long, tapering backward almost to a point, in contact with the anterior two large supraoculars; four supraoculars; fronto-parietals and interparietal distinct, the former in contact with second, third, and fourth supraoculars; parietals short, with a short suture behind interparietal; no nuchals; nostril in a single nasal, no postnasal; anterior loreal high, in contact with internasals; 7 supralabials, fifth and sixth largest;

^a From σφήν, a wedge; μορφή, shape.

^b Signifying from India.

^c Reproduced in this work on plate XVII, figs. 1-2.



FIGS. 1-2, SPHENOMORPHUS INDICUS; FIG. 3, LYGOSAURUS PELLOPLEURUS. (FROM BOULENGER.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

upper large temporal of second row larger than lower; lower eyelid scaly; ear-opening large, though smaller than eye, without lobules; one azygous postmental, or chin-shield; 38 scale rows round the body; scales smooth; 4 preanal scales, median pair largest; longest toe of hind limb reaches elbow of arm stretched backward; soles nearly uniformly granular. Color (in alcohol) above pale olive brown, with numerous small blackish spots arranged in a median series and a lateral row on each side, the spots of the latter being wider and bordering the upper dorso-lateral white band; from eye to above hind legs a broad blackish band with zigzag edges, and a median series of small white dots; above this black band a narrower whitish band and below it a broader white band from snout through ear-opening to groin; below this again a dark-brown band with median larger white spots; underside whitish; tail buff, darker above and with pale brownish spots in continuation of the dorsal and lateral bands; legs above brown with rounded pale spots; sutures of labials marked with dark brown spots.

Dimensions.^a

	<i>mm.</i>
Tip of snout to vent.....	37
Snout to ear.....	9
Fore leg.....	12
Hind leg.....	19

In an *adult* Chinese specimen (U.S.N.M. No. 35527) before me, with perfect tail, the latter measures 97 mm. from vent to tip, while the distance from tip of snout to vent is 75 mm. The tail of the latter is covered underneath with cycloid scales, the median series of which is irregularly widened transversely some distance behind the vent.

Variation.—Judging from Boulenger's description this species has sometimes a postnasal, while the number of scale rows round the middle of the body varies between 38 and 34, the latter number occurring in our Chinese specimen. The specimen described may be abnormal in having the frontal in contact only with first and second supraocular and consequently the fronto-parietal in contact with second supraocular, since Boulenger describes the species as having the frontal in contact with three supraoculars, a condition also found in our Chinese specimen (No. 35527).

Habitat.—This is a characteristic Himalayo-Chinese species which extends from the eastern Himalayas, where according to Boulenger it is common in Darjeeling, through the hills of Assam and Burma to China and Tonkin. United States National Museum has it from the province of Shensi, collected by Dr. Eliot Blackwelder, in 1904. It reaches its extreme eastern limit in Formosa where a specimen, now in the British Museum, was collected at Bangkimtsing by Mr. J. D. La

^a Tail broken.

Touche. I owe the opportunity to examine and describe this addition to the Formosan fauna to the kindness of Dr. G. A. Boulenger.

List of specimens of Sphenomorphus indicus.

Museum.	No.	Age.	Locality.	When collected.	By whom collected.
U. S. N. M.	35527		Shih Chuen Hsien, Shensi, China	1904	E. Blackwelder.
Brit. Mus.	99.4.24.101	Young	Bangkimsing, Formosa		J. D. La Touche.

^a Description, p. 216.

Genus **LEIOLOPISMA** ^a Duméril and Bibron.

1839. *Leiolopisma* DUMÉRIL and BIBRON, *Erpét. Gén.*, V, p. 742 (type, *L. telfairii*).
 1843. *Leiolepisma* FITZINGER, *Syst. Rept.*, p. 22 (emendation).
 1843. *Lampropholis* FITZINGER, *Syst. Rept.*, p. 22 (type, *M. guichenoti*).
 1845. *Mocoo* GRAY, *Cat. Liz. Brit. Mus.*, p. 80 (type, *M. guichenoti*).
 1845. *Lipinia* GRAY, *Cat. Liz. Brit. Mus.*, p. 84 (type, *L. pulchella*).
 1857. *Oligosoma* GIRARD, *Proc. Phila. Acad.*, 1857, p. 196; author's separate, p. 4 (type, *O. mocoo*).
 1864. *Liiolepisma* PETERS, *Mon. Ber. Berlin Akad. Wiss.*, 1864, p. 387 (emendation).

The genus, as here understood, is somewhat more restricted than Boulenger's section of the comprehensive genus *Lygosoma* having the same name. The species belong to the Old World, except one which occurs also in North America. This is the only species which occurs in the territory embraced in this work.

LEIOLOPISMA LATERALE ^b (Say).

1823. *Scincus lateralis* SAY, in Long's Exped. Rocky Mts., II, p. 324 (type-locality, bank of the Mississippi River between Cape Girardeau, Missouri, and New Orleans, Louisiana).—*Lygosoma lateralis* DUMÉRIL and BIBRON, *Erpét. Gén.*, V, 1839, p. 719.—*L. laterale* BOULENGER, *Cat. Liz. Brit. Mus.*, III, 1885, p. 263 (southern North America; Ningpo, and Szechuen, China); *Proc. Zool. Soc. London*, 1899, p. 162 (Kuatun, Prov. Fokien).—*Leiolopisma laterale* JORDAN, *Man. Vertebr. Anim. North. U. S.*, 8th ed., 1899, p. 201.
 1825. *Scincus unicolor* HARLAN, *Journ. Phila. Acad. Sci.*, V, Pt. 1, p. 156 (type-locality, "Southern States"); Pt. 2, 1827, p. 221.
 1838. *Tiliqua reevesi* GRAY, *Ann. Nat. Hist.*, II, Dec. 1838, p. 292 (type-locality, China; types in Brit. Mus.; Reeves, collector).
 1864. *Eumeces modestus* GUENTHER, *Rept. Brit. India*, p. 87, pl. x, fig. g (type-locality, near Ningpo, China; types in Brit. Mus.).
 1878. *Lygosoma nigropunctatum* BOCOVERT, *Ann. Sci. Nat., Zool.* (6), VII, art. 16, p. 2 (type-locality, Whampoa, China; types in Paris Mus.; Doctor Vesco, collector).
 1888. *Lygosoma (Leiolopisma) laterale* var. *modesta* BOETTGER, *Ber. Senckenberg. Naturf. Ges.*, 1888, *Abhandl.*, p. 187 (Napier Isl.); 1894, p. 145 (Chapu, near Ningpo).

^a From *λεῖος*, smooth; *λόπισμα*, covering envelope.


^b Signifying lateral; referring to the dark lateral band.

1893. *Lygosoma (Liolepisma) laterale* var. *reevesi* BOETTGER, Kat. Rept. Mus. Senckenberg., I, p. 104 (Peking; Chapu; Napier Island; Hongkong; Canton); Ber. Senckenberg. Naturf. Ges., 1901, p. 41 (Tsushima).

Boulenger^a failed to find any distinctive characters by which to separate Chinese specimens (the so-called *L. reevesi*) from North American ones. In the same manner I have been unable, upon the most searching comparison, to discover any difference between the Riu Kiu specimens and typical *L. laterale*.

Description.—*Adult:* Science College Museum, Tokyo, No. 37; Miyakoshima, Riu Kiu Archipelago (figs. 190–192.) Rostral broadly in contact with fronto-nasal, the portion visible from above much smaller than the latter shield; no supranasals; nostril in the center of a single shield; no postnasal; fronto-nasal, in contact with frontal, broader than long; prefrontals not in contact; anterior loreal in touch with first supralabial, in contact with second supralabial, fronto-parietal and prefrontals; frontal much longer than its distance from tip of snout, shorter than length of fronto-parietals and interparietal together, very wide in front, tapering nearly to a point behind, in contact with first and second supraoculars; four supraoculars, second largest; fronto-parietals long and narrow, longer than interparietal,

which is shaped like the frontal but not so long; parietals long and narrow, as long as frontal, in contact behind interparietal; four pairs of nuchals, first not in contact; lower eyelid granular with a small, transparent disk; seven supralabials, fifth under the eye, sixth largest; a very large upper temporal, fan-shaped with the apex forward, in contact with the entire outer edge of the parietal; ear-opening large, as large as eye, without lobules along the anterior border; a single unpaired pentagonal shield behind the mental; 28 smooth scales around the middle of the body, those on the sides but slightly smaller; two large preanal plates with a small one on each side; hind leg contained about three times in distance from snout to vent; fore and hind legs fail to meet when pressed along the side; 16 lamellæ under longest toe; tail cylindrical, pointed, with transversely widened plates underneath only towards the tip (in reproduced part). Color (in alcohol) above uniform olive brown; a blackish-brown line from nostril through eye, widening on tympanic region and extending above the ear-opening



FIGS. 190–192. —*LEIOLOPISMA LATERALE*. 2X NAT. SIZE. 190, TOP OF HEAD; 191, SIDE OF HEAD; 192, UNDERSIDE OF HEAD. No. 37, SCI. COLL. TOKYO.

^a Cat. Liz. Brit. Mus., III, 1887, p. 264.

backward along sides above fore and hind legs to side of tail as a broad dark-brown band above and below narrowly and indistinctly edged with whitish; on the flanks below this edge a paler brown, more indistinct band; underside pale.

Dimensions.

	<i>mm.</i>
Total length (tail reproduced).....	74
Snout to vent.....	44
Vent to tip of tail (reproduced).....	30
Snout to ear-opening.....	8.5
Greatest width of head.....	5
Axilla to groin.....	25
Fore leg.....	10
Hind leg.....	15

In specimens with unbroken tail, the length of the latter greatly exceeds the distance from snout to vent; thus in a North American specimen (U.S.N.M. No. 18012; St. Louis, Mo.) measuring 43 mm. from snout to vent, the distance from vent to tip of tail is 61 mm.; hind leg 14.5 mm.

Variation.—Considering the extent of the range of this species the amount of variation is small. The number of scales around the body is said to vary between 26 and 32. In the Asiatic specimens 28 seems to be normal, two specimens recorded by Boettger and two examined by myself having this number. Boulenger records one from Fokien having 26. The most usual color variation is the presence of minute dark-brown spots on the back in more or less regular longitudinal rows.

Habitat.—On the supposition that the American and Asiatic specimens really are identical, the present species has a very unique distribution. In North America it is known to inhabit the lower Austral life zone east of the Rocky Mountains, and is not found west of the latter at all. In Asia it occurs over a large area in China along the coast from near Ningpo to Canton, in the interior to the province of Szechuen, or (if Anderson's *Mocoa erigua*,^a from Momien really is the same species, which seems quite likely) to the extreme west end of the province of Yunnan, while northward it extends its range to the neighborhood of Peking, if the locality which attaches to a specimen in the Museum Senckenbergianum, purchased from the late Dr. Otto Herz in St. Petersburg, really be correct, which I doubt.

From the territory covered by the present work it has been recorded by Doctor Boettger as occurring in Tsushima. A single specimen (No. 37) in the Science College Museum, in Tokyo, is from Miyakoshima, southern group of the Riu Kius, and one in the Ham-

^aZool. West. Yunnan, I, 1879, p. 797.

burg Museum (No. 1195) was collected by Doctor Warburg in Okinawa shima.

List of specimens of Leiolopisma laterale.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.
Sci. Coll. Tokyo.	37	Adult a.	Miyakoshima		Tashiro (?).....	28
Hamburg.....	1195do.	Okinawa shima.....		Dr. Warburg.....	28

^a Description, p. 219, figs. 190-192.

Genus **LYGOSAURUS**^a Hallowell.

1860. *Lygosaurus* HALLOWELL. Proc. Phila. Acad., 1860, p. 496, (type, *L. pellopleurus*).

Only one species known, which is confined to the Riu Kiu Archipelago.

Hallowell's *Lygosaurus pellopleurus* has been relegated to that vast mass of species (more than 150) which some authors call *Lygosoma*, and more particularly to the section or subgenus *Homolepida*. The peculiar scutellation on top of the head, as well as the retractility of the claws, make the retention of the genus *Lygosaurus* imperative.

In the first place, the parietals have been reduced so in size as to differ but very little in appearance from ordinary nuchals; furthermore, the interparietal, with the parietal foramen, is pushed so far forward that the latter is on a line with or anterior to the posterior edge of the orbit; the frontal is also pushed ahead of its normal position so as to separate the prefrontals widely and reduce them to small scutes on the canthus rostralis. Its greater portion overlies, in fact, the nasal bones. Moreover, instead of a single frontal there are two, the posterior frontal being apparently composed of the posterior portion of the original frontal and the anterior portion of the fronto-parietals. It thus comes to pass that the posterior frontal and the interparietal are widely in contact, while the fronto-parietals are as widely separated. My preference to regard this unpaired shield which lies between the supraoculars as a second frontal rather than a third and unpaired fronto-parietal is due to the fact that the two frontals together are in contact with only the two anterior supraoculars, which is the normal arrangement in allied genera with only one frontal. Another reason is the fact that in two specimens examined by me the two frontals are actually fused. This extraordinary arrangement of the cephalic plates will be best understood by a glance at the appended figure (fig. 193). Hallowell's original description should be read in the light of the above interpretation of the scutes, so that his internasal becomes the fronto-nasal; his fronto-nasal is the anterior frontal; his frontal becomes the posterior frontal; his "inter-

^a From λύγος, flexible twig; σαῦρος, lizard.

parieto-fronto-parietal" is the interparietal; and his fronto-parietals are the true parietals.^a

The digits also show peculiarities which make it inadmissible to incorporate this genus with the typical *Lygosoma*. The claws appear to be retractile, as in the genus *Ristella*, though not within a single shield as in the latter, but between a large upper and a lower shield.^b The digits, moreover, are not covered above with plates, but with imbricate alternating scales on each side of the middle line.

Lygosaurus, so far as recorded, is a genus peculiar to the Riukiu Archipelago, and does not seem to possess any known near relative. Among the skinks reported from the adjacent countries it stands quite isolated. At present one can only suggest that possibly in the future Formosa may prove to possess some related form. It should be noted, however, that *Lygosaurus* does not seem to belong to the Himalayo-Chinese fauna. Various points in its structure recall south Indian forms rather than any genus or species peculiar to the northern mountains.

LYGOSAURUS PELLOPLEURUS c Hallowell.

Plate XVII, fig. 3.

1860. *Lygosaurus pellopleurus* HALLOWELL, Proc. Phila. Acad., 1860, p. 496 (type-localities, Amami-o-shima and Okinawa, Riu Kiu Archipelago).—*Lygosoma pellopleurum* BOULENGER, Cat. Liz. Brit. Mus., III, 1887, pp. 319, 512 ("Loo Choo Islands"); Proc. Zool. Soc. London, 1887, p. 147, pl. xviii, fig. 2d ("Loo Choo Islands," Pryer collection).—OKADA, Cat. Vert. Japan, 1891, p. 70 (Amami-o-shima; Okinawa shima).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 860; authors' separate, p. 11 (Okinawa).—BROWN, Proc. Phila. Acad., 1902, June 11, p. 185 ("Loo Choo Islands").—*Lygosoma (Homolepida) pellopleurum* BOETTGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 107 (Okinawa shima).


^a Boulenger offers a homology of the cephalic shields somewhat different from the above (Proc. Zool. Soc. London, 1887, p. 148, footnote), inasmuch as he regards the posterior frontal as an azygous fronto-parietal, and the parietals as split up into an anterior (my fronto-parietals) and a posterior pair. I can not follow him in this. In Science College Museum No. 4 (the one here figured, fig. 193) the suture between the anterior and posterior frontals is obliterated, and so it is in our No. 36527. A single shield formed by a combination of these two would have a striking analogy in the frontal of *Chalcidoeops*, from Ceylon (Boulenger, Cat. Liz. Brit. Mus., III, 1887, p. 423, pl. xxxvii, fig. 1a) in which it is "angularly emarginate on each side by the first supraocular." In the nearly related *Scopphis*, from southern India, (Boulenger, Cat. Liz. Brit. Mus., XII, p. 423, pl. xxxvii, fig. 2a), the frontal is actually split much in the same way as in *Lygosaurus*, and Boulenger himself in the diagnosis of the genus speaks of "two frontals." In this genus "fronto-parietals," much in the same relative position as in *Lygosaurus*, are "present," while the interparietal is nearly identical in shape and relation.

^b I have no specimen of *Ristella*. The arrangement of the claws is described by Boulenger (Cat. Liz. Brit. Mus., III, p. 357) as follows: "Claws completely retractile in a large compressed sheath formed of one large scale cleft inferiorly." Stoliczka first called attention to this character (Journ. Asiat. Soc. Bengal, XLI, Pt. 2, 1872, p. 129), but indicates "retractile claws lying between two terminal enlarged shields."

^c From *πρᾶλλός*, blackish; *πλευρά*, side.

^d Reproduced in this work on Plate XVII.

Description.—*Adult:* Science College Museum, Tokyo, No. 4; Shuri, Okinawa shima; July, 1895, Shiraiishi, collector (figs. 193–195). Rostral very broadly in contact with fronto-nasal; no supranasals; nostril in a single nasal; no postnasal; fronto-nasal broader than long, broadly in contact with frontal; prefrontals very small, smaller than posterior supraocular and interparietal, widely separated; frontal undivided, very long, twice as long as its distance from tip of snout, angularly emarginate laterally by the anterior supraocular; the anterior portion (corresponding to the anterior frontal in such specimens in which there is a transverse suture between the lateral emarginations) with the longer sides converging posteriorly and in contact with first supraocular, the posterior portion (corresponding to the posterior frontal) with the longer sides diverging posteriorly and in contact with first and second supraoculars, also in contact behind with interparietal; four supraoculars, anterior largest, first and second in contact with frontal (anterior and posterior frontals); fronto-parietals not in contact with each other, about as large as third supraocular; interparietal about as wide as long, slightly larger than fronto-parietals, in contact with (posterior) frontal; parietals very short, not larger than fronto-nasal, in contact behind interparietal; no nuchals; loreals small, narrow, their lower edge coextensive with the upper edge of second supralabial; lower eyelid scaly; six supralabials, fifth longest; between fourth and fifth a large subocular, the lower angle of which nearly reaches the edge of the lip; temporals small, scale-like; ear-opening rather large, as large as both loreals together, horizontally oval, without projecting lobules; a single narrow, quadrangular shield, with nearly parallel anterior and posterior edges, behind the mental; submandibulars small, scarcely differentiated; 26 scale rows around the middle of the body, the scales of nearly equal size, those on back strongly tricarinate, the median ones even with two more though fainter keels; preanal scales not enlarged; legs short, hind leg being contained about three and three-fourths times in distance from snout to vent, while fore and hind legs fail to meet by the length of the fore leg; digits very short, first especially so, covered above with imbricate, alternating scales, one on each side of the median line and terminating above in a large nail-shaped scale, under which the claw can be retracted (in the present specimen nearly



FIGS. 193–195.—LYGOSAURUS PELLOPLEURUS. 2 X NAT. SIZE. 193, TOP OF HEAD; 194, SIDE OF HEAD; 195, UNDERSIDE OF HEAD. NO. 4, SCI. COLL. TOKYO.

temporals small, scale-like; ear-opening rather large, as large as both loreals together, horizontally oval, without projecting lobules; a single narrow, quadrangular shield, with nearly parallel anterior and posterior edges, behind the mental; submandibulars small, scarcely differentiated; 26 scale rows around the middle of the body, the scales of nearly equal size, those on back strongly tricarinate, the median ones even with two more though fainter keels; preanal scales not enlarged; legs short, hind leg being contained about three and three-fourths times in distance from snout to vent, while fore and hind legs fail to meet by the length of the fore leg; digits very short, first especially so, covered above with imbricate, alternating scales, one on each side of the median line and terminating above in a large nail-shaped scale, under which the claw can be retracted (in the present specimen nearly

concealed, only the extreme point being visible); tail cylindric, tapering to a point; the caudal scales above with four decreasing posteriorly to two keels, no transversely enlarged plates underneath. Color (in alcohol) above pale brown, with a series of narrow elongate dark spots on each of the two median scale rows; a dark brown band from nostril through eye, high above the ear-opening and along the upper part of the sides, disappearing on the sides of the tail; top of head irregularly spotted with dark brown, with larger spots of similar color on upper and lower labials; whole under surface whitish.

Dimensions.

	<i>mm.</i>
Total length.....	98
Snout to vent.....	52
Vent to tip of tail.....	46
Snout to ear-opening.....	7.5
Greatest width of head.....	6
Axilla to groin.....	30
Fore leg.....	9
Hind leg.....	14

Variation.—The most noteworthy variation seems to be the presence or absence of a suture dividing the frontal into two distinct shields. In Hallowell's specimen as well as in that described by Boulenger and one in our museum (U.S.N.M. No. 36526) the suture is present, while in the other (No. 36527), as well as the one in Science College Museum, Tokyo (No. 4), the suture is absent. It is consequently not possible at the present to say which condition is the normal one. Otherwise there is not much difference in the headshields of the specimens before me, except that in No. 36526 the parietals fail to meet behind the interparietal. The size of the ear-opening is variable, probably due to the state of preservation of the specimen. Two of the specimens before me have 26 scale rows, and one 28 around the body, while Boulenger gives 24 for his specimen, the same number as recorded by Hallowell.

Habitat.—Apparently confined to the middle and northern groups of the Riu Kiu Archipelago, from both of which the Rodgers expedition brought home specimens which now are lost. Later collectors have obtained it in Okinawa shima. Curiously enough it has not been recorded from the southern group.

*List of specimens of *Lygosaurus pellopleurus*.*

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.
U. S. N. M.	36526	Adult <i>a</i>	Okinawa shima		Tashiro	26
Do	36527do. <i>a</i>do	do	28
Sci. Coll., Tokyo	4	Adult <i>b</i>	Okinawa	July, 1895	Shiraishi	26

a Sci. Coll. No. 36.

b Description, p. 223, figs. 193-195.

Genus **CRYPTOBLEPHARUS**^a Wiegmann.

1834. *Cryptoblepharus* WIEGMANN, *Herpet. Mex.*, p. 12 (type, *Ablepharus pocillopleurus*).

Having recently obtained specimens of the type species of the genus *Ablepharus*, I have convinced myself that it is not congeneric with the type of *Cryptoblepharus*.^b The arrangement of the supraoculars in the former is so radically different from that of the latter that even if there were no other distinction I should hesitate to place them in the same genus.

In the key to the genera (p. 193) this genus is said to have "no movable lower eyelid." This expression does not mean that the lower eyelid is absent. On the contrary, it is there, but not movable, having coalesced with the rudimentary upper lid. What appears to be the exposed eye itself is only the very large transparent disk of the lower eyelid.

CRYPTOBLEPHARUS BOUTONII^c **NIGROPUNCTATUS**^d (Hallowell).

1860. *Ablepharus nigropunctatus* HALLOWELL, *Proc. Phila. Acad.*, 1860, p. 489 (type-locality, Port Lloyd, Bonin Islands; Rodgers, collector).

With the number of scale rows of a large series of Hawaiian "snake-eyed skinks" varying between 28 and 30, I can not well refer the ten Bonin Island specimens with 24 and 26 scale rows to the same subspecies, notwithstanding the fact that I find them to agree substantially as to coloration. With the present material insufficient to work out an intelligent and comprehensive review of the various forms clustering around *C. boutonii*, it is safer to keep the groups separate nomenclaturally. How essential it is to follow this rule is shown by Boulenger's recent notes under *Ablepharus boutonii*.^e It is even a question with me whether the Bonin Island form should not be designated binominally on account of the apparently constant presence of a distinct postnasal. The occasional occurrence of such a shield in Hawaiian specimens,^f however, makes it safer to adhere to the provisionally accepted nomenclature of trinomials for this group of forms.

Hallowell's type of *Ablepharus nigropunctatus* appears to be lost. It was obtained at Port Lloyd, on Peel Island (Chichishima), of the middle or Beechey Group, Bonin Archipelago, consequently from the same locality from which came U.S.N.M. Nos. 23897-8 and 23900,

^a From *κρυπτός*, hidden; *βλέφαρον*, eyelid.

^b See Stejneger, *Proc. U. S. Nat. Mus.*, XXI, no. 1174, 1899, p. 810.

^c 1831. *Scincus boutonii* DESJARDIN, *Ann. Sci. Nat.* (1), XXII (p. 298) (type-locality, Island of Mauritius; types in Paris Mus.). Named for Louis Bouton, a French botanist.


^d Spotted with black.

^e *Ann. Mus. Civ. Genova* (2), XVIII, 1898, pp. 702 and 719.

^f See Stejneger, *Proc. U. S. Nat. Mus.*, XXI, No. 1174, 1899, p. 811.

and the three specimens in the Science College Museum, in Tokyo. I have also before me three specimens from Haha shima, in the southern, or Coffin Group, U.S.N.M. Nos. 23899, 33857, and Sci. Coll. Mus. Tokyo No. 48, essentially like the ones from the middle group. These agree pretty well with Hallowell's description,^a except that he does not mention a pale dorso-lateral line which is more or less apparent in all the specimens mentioned above. A single large specimen (Sci. Coll. Mus. No. 50), probably from Chichi shima, is considerably darker, with scarcely a trace of the lateral line, but the duskiess of this specimen may be due to defective preservation. Of the ten specimens mentioned, six have 24 scale rows around the middle of the body, while four have 26.

Description (figs. 196-198).—*Adult*; U.S.N.M. No. 33857; Haha shima, Bonin Archipelago; March, 1904; Owston collection. Rostral in contact with fronto-nasal; nostril round, in the lower portion of a rather large, pentagonal nasal; a small triangular postnasal,



FIGS. 196-198.—CRYPTOBLEPHARUS BOUTONII NIGROPUNCTATUS. 2 X NAT. SIZE. 196, TOP OF HEAD; 197, SIDE OF HEAD; 198, UNDERSIDE OF HEAD. No. 50, SCI. COLL. TOKYO.

in contact with nasal, anterior loreal and first supralabial; no supranasal; fronto-nasal slightly broader than long, not in contact with frontal; prefrontals rather large, in contact with each other, with both loreals, and with anterior supraocular; frontal small, considerably smaller than fronto-parietal, in contact with first and second supraoculars; four supraoculars, second largest; four superciliaries; fronto-parietal (consisting of the fused fronto-parietals and interparietal) quadrangular, in contact with frontal and three supraoculars, very large; parietals long and narrow, broadly in contact behind fronto-parietal; one pair of large nuchals, in contact behind parietals, and followed by a double series of short, but very wide plates, or scales, which gradually decrease in width backward, merging into the two median dorsal scale rows; first loreal very high and narrow, in touch with first supralabial behind postnasal; second loreal much lower, pentagonal, in contact with two preoculars; the upper suture of the

^a In Hallowell's description "fronto-parietals" is evidently a misprint for fronto-parietal.

supralabials from the rostral to the anterior border of the subocular supralabial deeply depressed, forming a slightly arched groove; eight supralabials, seventh very high, sixth longest, forming a long and low subocular; eye surrounded by granules and above by three elongate narrow scales; a large upper temporal adjoining parietal; ear-opening round, smaller than eye, without projecting lobules; mental small; a large, pentagonal postmental followed by three pairs of large submandibulars, the last of which is very long; 24 rows of scales around the middle of the back, laterals smallest, and the two median dorsals largest, the dorsals faintly multicarinate; preanal scales somewhat enlarged; hind leg about two and a half times in distance from snout to vent; fore and hind legs when pressed against the sides overlap by more than length of longest toe; tail cylindrical, with a median series of transversely enlarged scales underneath. Color (in alcohol) above pale olive brown, with numerous irregular blackish brown elongated spots, many of which are edged with paler spots; flanks dark brown, with similar pale spots which at the upper edge collect so as to form a somewhat irregular pale dorso-lateral line; legs above like the flanks; lower surface bluish white with scattered dark spots on legs and tail.

Dimensions.

	<i>mm.</i>
Total length.....	143
Snout to vent.....	54
Vent to tips of tail.....	89
Snout to ear-opening.....	10.5
Greatest width of head.....	8
Axilla to groin.....	28
Fore leg.....	17
Hind leg.....	23

Variation.- There is a great deal of variation in the size and relation of the head shields. Thus in about one-half of the specimens before me the fronto-nasal and the frontal are in touch; on the other hand, the suture between the prefrontals in No. 23897 is so wide as to nearly equal the length of the fronto-nasal, while the frontal is so reduced in size as to equal the anterior supraocular and to lose contact with the frontal parietal, the second supraoculars being in contact behind it. All have a distinct postnasal, but none show signs of a supranasal. The number of supralabials is also slightly variable. In most specimens there are four supralabials in front of the subocular, but in the specimen described above as well as in one of the Chichi shima specimens in the Science College Museum, Tokyo, there are five on both sides. In our No. 23897, from the latter island, there are four on one side and five on the other. The coloration is very variable, not only the ground color, which varies from pale clay color to dark brown, but also in the amount of spotting, some specimens being nearly unspotted,

while in others the numerous dots are reddish brown instead of blackish. No. 23897 is densely spotted with blackish underneath, especially under the tail and throat.

Habitat.—The species, of which I consider the Bonin Island specimens to represent a separable form, is now distributed in several slightly deviating varieties or subspecies over nearly the entire tropical island world in the Indian and Pacific oceans, including Australia. It is also found locally on both the eastern and western coasts of Africa.

Whether the present form is found outside the Bonin Archipelago it is impossible to say at present. The one found in the Hawaiian Islands is characterized by a greater number of scale rows, normally 28 to 30.

In the Bonins it has been collected in the largest island of each of the two southern groups.

List of specimens of Cryptoblepharus nigropunctatus.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.
U.S.N.M.....	23897	(a)	Chichi shima, Bonin....	Mar., 1884	Hirota and Sekiguchi.	24
Do.....	23898		do.....	do.....	do.....	24
Do.....	23900		do.....	do.....	do.....	24
Do.....	23899		Haha shima, Bonin....	Nov., 1887	26
Do.....	33857	(b)	do.....	Mar., 1904	A. Owston.....	24
Do.....	36535	(c)	do.....	Nov., 1887	24
Sci. Coll. Tokyo..	49		Chichi shima, Bonin....	do.....	26
Do.....	50	(d)	Bonin.....	26
Do.....	?		Chichi shima, Bonin....	Hirota and Sekiguchi.	24
Do.....	?		do.....	do.....	26

a P. 227.

b Description, p. 226.

c Sci. Coll. No. 48.

d Figs. 196-198.

Family LACERTIDÆ.

Pleurodont lizards with scaly or papillose plicate tongue; lower surface with squarish scales in longitudinal and transverse rows.

An Old World family, mostly found in Africa and the western Palearctic region. It is absent in Madagascar and Australia, and in southeastern Asia it is only represented by the genus *Takydromus*, which is peculiar to that part of the world. Peripheral representatives of two other genera occur in the continental area covered by this work, one of these even reaching the island of Sakhalin.

The species of this family are chiefly terrestrial in their habits.

KEY TO THE GENERA OF LACERTIDÆ OCCURRING IN JAPAN AND ADJACENT TERRITORY.

a¹ Inguinal pores only *Takydromus*, p. 229.

a² Femoral pores only.

b¹ Lamellæ on underside of digits keeled..... *Eremias*, p. 247.

b² Lamellæ on underside of digits smooth..... *Lacerta*, p. 251.

Genus TAKYDROMUS^a Daudin.

1803. *Takydromus* DAUDIN, Hist. Nat. Rept., III, p. 251 (type, *T. quadrilineatus* Daudin).
 1804. *Tachydromus* LATREILLE, Nouv. Dict. d'Hist. Nat., XXIV, tabl. méth., p. 63 (emendation).
 1806. *Tachydromus* FROEIER, Dumeril's Anal. Zool. (German transl.), p. 83 (emendation).
 1845. *Tachysaurus* GRAY, Cat. Liz. Brit. Mus., p. 52 (type, *T. japonicus*).

KEY TO JAPANESE, KOREAN, AND FORMOSAN SPECIES OF TAKYDROMUS.

- a*¹ Dorsal scales small, not arranged in longitudinal series..... *T. dorsalis*, p. 229.
*a*² Dorsal scales large, arranged in longitudinal series.
*b*¹ Three pairs of submental shields.
*c*¹ Width of head more than half the length of shielded part of head.
*d*¹ One inguinal pore on each side..... *T. septentrionalis*, p. 232.
*d*² Two inguinal pores on each side..... *T. formosanus*, p. 235.
*c*² Width of head one-half the length of shielded part of head.
T. smaragdinus, p. 236.
*b*² Four pairs of submental shields.
*c*¹ More than one inguinal pore on each side.
*d*¹ Two (very rarely three) inguinal pores on each side; not more than six rows of enlarged dorsals; posterior enlarged throat scales pointed.
T. tachydromoides, p. 238.
*d*² Three inguinal pores on each side; eight rows of enlarged dorsals; posterior enlarged throat scales scarcely pointed..... *T. amurensis*, p. 245.
*c*² One inguinal pore on each side..... *T. wolteri*, p. 247.

TAKYDROMUS DORSALIS^b Stejneger.


1904. *Takydromus dorsalis* STEJNEGER, Smithsonian Quarterly (Misc. Coll., XLVII), Nov. 9, 1904, p. 294 (type-locality, Ishigaki shima, Yaeyama Archipelago, Riu Kiu Islands, Japan; type U.S.N.M. No. 34162).

Description.—*Adult male*: U.S.N.M. No. 34162; Ishigaki shima, Riu Kiu; April–June, 1899; A. Owston collection (figs. 199–202). Rostral not in contact with internasal, its upper lateral corner, formed by the suture with the first supralabial, barely entering the nostril; nostril rounded, between two nasals, the anterior being in contact with that of the other side behind the rostral, the posterior in contact with only one supralabial, the first; two loreals, of which the posterior very large; internasal much longer than broad, about three-fourths the length of the prefrontals, which are nearly as long as the frontal; two large supraoculars in contact with frontal; a very small third one behind, which is barely touching the outer corner of the fronto-parietals; supraoculars separated from the superciliaries by a single row of granules, the anterior of which is somewhat enlarged and separates the first supraocular from the posterior loreal; fronto-parietals slightly longer than the internasal; parietals longer than frontal; interparietal small, narrow; occipital slightly smaller, triangular; six supralabials, the fifth


^a Careless transliteration for *Tachydromus*, from τὰχύς, swift, and δρόμος, running.

^b Signifying dorsal; referring to the unique arrangement of the scales on the back.


very long and under the eye; temporals small, strongly keeled; a long, narrow shield along the anterior half of the outer edge of the parietal; lower labials very long and narrow; four pairs of submandibulars, or chin-shields, increasing in size posteriorly, anterior two pairs entirely in contact with each other, the third pair posteriorly separated by a small, elongated scale; upper surface of body covered with strongly carinated scales, 28-30 across the middle of the body, those on the back larger, about five corresponding to three ventral plates, gradually merging into




201



200



202



FIGS. 199-202. TAKYDROMUS DORSALIS. 2 X NAT. SIZE.
199, TOP OF HEAD; 200, UNDERSIDE OF HEAD; 201, MIDDLE OF BACK; 202, INGUINAL REGION. No. 34162. U.S.N.M.

the laterals, of which seven to eight correspond to three ventrals; scales on upper neck about the size of smallest laterals; scales on upper surface of limbs large, keeled, larger than largest dorsals, those on forearm being smooth and forming broad transverse plates; gular scales large granules posteriorly, gradually becoming larger and keeled toward the collar, 26 in a series on the median line between third pair of chin-shields and collar; scales on collar large, pointed, keeled, the median pair largest; six rows of ventral plates, median faintly keeled, outer row strongly so, 24 on the median line from collar to preanal plate; preanal plate large, with two minute plates on each side; two inguinal pores on each side; limbs very long and slender, with very long digits, the longest toe of hind leg stretched forward, reaching the shoulder joint; tail more than three and

a third times as long as head and body together, swollen at base, covered above and below with strongly keeled scales, which are twice as long as the largest dorsals. Color (in alcohol) bluish slate (probably greenish in life), throat paler; a black line from nostril through middle of eye to center of ear-opening; a pale (probably yellowish) line, below this from nostril through lower eyelid to lower edge of ear-opening, and a dusky line, less distinct, bordering the pale line below from middle of subocular labial to under ear.

Dimensions.

	<i>mm.</i>
Total length	284
Snout to vent.....	64
Vent to tip of tail	220
Snout to edge of collar	22.5
Snout to posterior edge of occipital.....	16
Width of head.....	9
Fore limb.....	25
Hind limb.....	35

The adult *female* (U.S.N.M. No. 34166; same locality and collector) differs chiefly in having the tail tapering from the beginning without the basal swelling and in somewhat shorter legs, the adpressed hind leg not reaching beyond the axilla.

The *young* (No. 34169, measuring 38 mm. from snout to vent) is like the female; occipital is not proportionally larger.

Variation.—In the ten specimens which are in the National Museum very little individual variation is observed. All have four pairs of chin-shields; all have two inguinal pores on each side, except No. 34163, which has three. All have the nasals in contact behind the rostral. There is some difference in the distinctness of the keels on the ventral plates, due apparently to age, since the larger specimens are more distinctly keeled, while in the smaller ones the four median plates are smooth or nearly so.

Remarks.—This exceedingly distinct species does not show any near affinity to the other known species of the genus. It is absolutely unique in the arrangement and size of the dorsal scales.

Another character unexpected in so southern a species is the number of submandibular shields. Four chin-shields are found normally only in the northern forms, namely, *T. tachydromoides* and its allies.

The discovery of this novelty in the southern group of the Riu Kiu Archipelago is the more startling, since we have *T. smaragdinus* from Miyakoshima in the same group, a species which conforms in all respects to the general type of the genus. On the other hand, the fact that we have ten specimens of the present species from Ishigaki shima and none of the regular type seems to indicate that the latter may not occur in that island at all.

Habitat.—Apparently confined to the island of Ishigaki, in the southern group of the Riu Kiu Archipelago, where Mr. A. Owston's collector obtained it during the summer of 1899.

List of specimens of Takydromus dorsalis.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Inguinal pores.
34160	Male.....	Ishigaki shima.....	Apr. June, 1899.	A. Owston.....	2
34161	do.....	do.....	do.....	do.....	2
34162	do. ^a	do.....	do.....	do.....	2
34163	do.....	do.....	do.....	do.....	3
34164	Female.....	do.....	do.....	do.....	2
34165	do.....	do.....	do.....	do.....	2
34166	do. ^b	do.....	do.....	do.....	2
34167	do.....	do.....	do.....	do.....	2
34168	do.....	do.....	do.....	do.....	2
34169	Young ^b	do.....	do.....	do.....	2

^a Type; description, p. 229; figs. 199-202.^b P. 231.**TAKYDROMUS SEPTENTRIONALIS^a Guenther.**

1864. *Tachydromus septentrionalis* GUENTHER, Rept. Brit. India, p. 69, pl. vii, fig. E (type-locality, Ningpo, China; types in Brit. Mus.); Ann. Mag. Nat. Hist. (6), I, 1888, pp. 166, 167, 168 (Kiukiang; Nankin; Ningpo); Ann. Mus. Zool. St. Pétersbourg, I, 1896, p. 203 (Huihsien, Prov. Kansu).—BOERRGER, Kat. Rept. Mus. Senckenberg., I, 1893, p. 79 (Lueshan near Kiukiang; Dalanshan, near Ningpo); Ber. Senckenberg. Naturf. Ges., 1894, p. 139 (Lueshan Mts.), p. 145 (Dalanshan Mts. and Tchinhai, near Ningpo).—BOULENGER, Proc. Zool. Soc. London, 1899, p. 161 (Kuaton, 3,000-4,000 alt, N. W. Fokien; J. D. La Touche, collector).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 354 (Ningpo).

Nine specimens (U.S.N.M. Nos. 34170-78) from Taipe, Formosa, appear to be typical *T. septentrionalis* with one inguinal pore on each side and three submandibular shields. They have a moderately broad head and moderately large temporals, about seven in a row between orbit and ear-opening, and in these points as well as in coloration differ from *T. smaragdinus* which normally also has two more rows of large dorsals. I have compared them with two Chinese specimens collected by Dr. Eliot Blackwelder, in the Chin Ling Mountains in the Province of Shansi (U.S.N.M. Nos. 35525-26).

The relations to *T. formosanus* will be discussed under the latter.

Description.—*Adult male*; U.S.N.M. No. 34172; Taipe, Formosa; March, 1903; A. Owston collection. Rostral not in contact with internasal; nostril rounded, between two nasals, the anterior being in contact with that of the other side behind the rostral, the posterior in contact with only one supralabial, the first; two loreals, posterior very large; internasal about as long as broad, a trifle shorter than the prefrontals which are less than three-fifths the length of the frontal; two large supraoculars in contact with frontal; a small posterior supraocular in contact with fronto-parietals; supraoculars separated from the posterior superciliaries by a single row of granules,

^a Signifying northern.

the anterior, very long, superciliary being in contact with the anterior supraocular throughout its whole length; a small shield in front of first supraocular separating it from the posterior loreal; frontoparietals longer than internasal; parietals slightly longer than frontal; interparietal moderate, about as large as anterior supraocular; occipital much smaller, about the size of third supraocular; seven supralabials, fifth very large, under the eye; temporals moderate, keeled, about 7 in a row between orbit and ear-opening; an elongated shield along the anterior half of the outer edge of the parietal; lower labials long and narrow; three pairs of chin-shields, increasing in size posteriorly, only first pair in contact throughout, second pair posteriorly separated by a wedge of elongated granules; back covered by six series of large keeled scales, three on each side, and two median ones slightly smaller, the keels forming continuous ridges, six of the large scales corresponding in length to five ventral plates; laterals granular, forming a broad longitudinal band on the sides; scales on upper surface of limbs large, keeled, not larger than large dorsals, a series on the forearm being rather widened transversely and smooth; gulars large, granules gradually merging into large, pointed, keeled scales on neck and collar, about 23 granules and scales on the median line between second pair of chin shields and collar; scales on collar long, pointed, and keeled like those on neck and chest; eight rows of ventral plates, which on the posterior part of the abdomen are flanked by two series of larger keeled scales having a row of similar but smaller scales between them; ventral plates all pointed behind and strongly keeled, 29 on the median line from collar to preanal plate; preanal plate large, smooth, with two smaller plates on each side; a single inguinal pore on each side; limbs moderate, the outstretched hind leg reaching the axilla; tail three and one-third times as long as head and body together, swollen at base, covered above and below with strongly keeled scales, which are as large as the largest dorsals. Color (in alcohol) above olive, becoming light tawny-olive on the tail; two narrow blackish brown lines on back along the keels of the first row of larger dorsals on each side of the median line, these lines continuing some distance on the tail; granular area of the sides of the body dark brownish gray, continued on the sides of the tail as a narrow dusky line, which almost disappears halfway from the tip; a black line from nostril through center of eye, behind the latter two blackish lines, one above along the parietals, the other below through the ear-opening, inclosing between them a lighter brown area posteriorly connected with the lateral dark band; an elongated whitish spot on upper eyelid; a narrow whitish line from below nostril through the lower eyelid, over the lower temporal region, through the lower edge of ear-opening, to the shoulder, where it disappears; a black line on the posterior aspect of the femur and indicated on the tibia; lower surface whitish, suffused with tawny-olive on tail.

Dimensions.

	<i>mm.</i>
Total length.....	211
Snout to vent.....	49
Vent to tip of tail.....	162
Snout to collar.....	18
Snout to posterior edge of occipital.....	12
Greatest width of head.....	7
Fore leg.....	17
Hind leg.....	26

The *female* (U.S.N.M. No. 34174, same locality and collector) differs in the tail gradually tapering from the base, and in the shorter hind leg, the tips of the fourth toe scarcely reaching beyond the elbow.

The *young* (U.S.N.M. No. 34178; snout to vent 31 mm.) is essentially like the female. In this specimen the interparietal is comparatively larger, but the occipital is nearly obsolete. In a Chinese specimen of same size (U.S.N.M. No. 35526) both of these shields are uncommonly large, with a small square shield between them.

Variation.—In the Formosan series of nine specimens there is but little individual variation. All have one inguinal pore; three pairs of chin-shields; six rows of enlarged dorsals; and the nasals broadly in contact or at least touching behind the rostral. None of the specimens, including the Chinese, has any outer accessory dorsal scale row. The color is also essentially alike, except that in several specimens a distinct whitish line on the inner half of the outer dorsal row of scales forms the upper edge of the lateral dark band, while the dorsal median black lines are often absent or obscure.

Habitat.—Widely distributed over eastern China in the mountains, on the coast from Fokien to Ningpo, west to the provinces of Kansu, where it was collected by Berezowski in 1892, and Shensi, where Dr. E. Blackwelder obtained it in 1904.

In Formosa it was taken during March, 1903, by Mr. Owston's collector at Taipe, the specimens being in the United States National Museum.

List of specimens of Takydromus septentrionalis.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Inguinal pores.
34170	Female.....	Taipe, Formosa.....	Mar., 1903	A. Owston.....	1
34171do.....do.....do.....do.....	1
34172	Male <i>a</i>do.....do.....do.....	1
34173do.....do.....do.....do.....	1
34174	Female <i>b</i>do.....do.....do.....	1
34175	Male.....do.....do.....do.....	1
34176	Female.....do.....do.....do.....	1
34177	Male.....do.....do.....do.....	1
34178	Young <i>b</i>do.....do.....do.....	1
35525	Female.....	Liang Ho, Shensi, China.....	1904	E. Blackwelder.....	1
35526	Young <i>b</i>	Hsiao Wang, Shensi, China.....	1904do.....	1

^a Description, p. 232.^b P. 234.

TAKYDROMUS FORMOSANUS ^a Boulenger.

1894. *Tachydromus formosanus* BOULENGER, Ann. Mag. Nat. Hist. (6), XIV, Dec. 1894, p. 462 (type-locality, Taiwan fu, Formosa; types in British Museum; Holst, collector).

There being no specimens of this species in the collections now accessible to me, I reproduce the original description.

Original description.—Head moderately elongate; nasals in contact behind the nostral; a series of granules between the supraoculars and the superciliaries; a small shield separates the large anterior supraocular from the loreal; temporal scales very small and keeled; three pairs of chin-shields; six longitudinal series of large strongly keeled shields, separated on the vertebral line by one or two series of smaller shields; eight or ten series of strongly keeled ventral shields, with two or three series of smaller shields on each side; two inguinal pores on each side. Olive brown above; a dark brown or blackish lateral band from the end of the snout to the base of the tail, passing through the eye, edged below, and usually also above by a whitish streak; the upper light streak, if present, originates above the tympanum and runs along the outer series of dorsal shields; the lower extends from the end of the snout to the thigh, passing through the tympanum and following the upper series of ventro-lateral shields; a whitish, black-edged streak along the hinder side of the hind limb; lower parts whitish.

Dimensions.

	mm.
Total length.....	186
Head.....	13
Width of head.....	6
Body.....	44
Fore limb.....	15
Hind limb.....	21
Tail.....	122 ^b

Remarks.—The Formosan grass lizard seems closely related to *Tachydromus septentrionalis* Guenther, if indeed distinct, the chief difference being in the latter having only one inguinal pore. This character may or may not be constant, but as Boulenger had “several” specimens with two pores, while, on the other hand, I have nine with one pore only, it may be better to retain *T. formosanus* separate for the present until examination of a large number of specimens from the type-locality can be made.

Habitat.—Recorded only from Formosa, specimens now in the British Museum having been collected by Mr. Holst at Taiwan fu and in the central portion of the island.


^a Signifying belonging to Formosa.

^b Boulenger, Ann. Mag. Nat. Hist. (6), XIV, 1894, pp. 462–463.

TAKYDROMUS SMARAGDINUS^a Boulenger.

1848. *Tachysaurus japonicus* ADAMS, Narrat. Voy. *Samarang*, II, p. 305 ("Miaco-shimas") (not of Duméril and Bibron, 1839).
1887. *Tachydromus smaragdinus* BOULENGER, Cat. Liz. Brit. Mus., III, p. 509 (type-locality, "Loochoo Islands;" types in Brit. Mus.; Fryer collection); Proc. Zool. Soc. London, 1887, p. 147, pl. xvii, fig. 2; pl. xviii, fig. 1 (Loochoo).—GUENTHER, Ann. Mag. Nat. Hist. (6), I, Mar. 1888, p. 168 (Loochoo Is.).—OKADA, Cat. Vert. Japan, 1891, p. 70 (Okinawa).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 859; author's separate, p. 10 (Okinawa).—BOETTGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 107 (Okinawa shima).—BROWN, Proc. Phila. Acad., 1902, June 11, p. 184 (O-shima or Okinawa shima).—SCHENKEL, Verh. Naturf. Ges. Basel, XIII, Pt. 1, p. 186 (Okinawa).

Description (figs. 203-204).—*Adult male*; U.S.N.M. No. 16361; Great Loo Choo Island; May, 1853; W. Heine, collector. Rostral not in



FIGS. 203-204.—TAKYDROMUS SMARAGDINUS. 2X NAT. SIZE. 203, UNDERSIDE OF HEAD; 204, INGUINAL REGION. No. 3524, U.S.N.M.

contact with internasal, the anterior nasals being in contact behind the rostral; posterior loreal twice as large as anterior; internasal longer than broad, shorter than prefrontals which are more than three-fifths and less than three-fourths the length of the frontal; two large supraoculars in contact with frontal; a small posterior supraocular in contact with fronto-parietals; supraoculars separated from the posterior superciliaries by a single row of granules, the anterior superciliary, which is very long, being in contact with the anterior supraocular throughout its whole length; a small shield in front of first supraocular, separating it from the posterior loreal; fronto-parietals longer than internasal, as long as prefrontals; parietals longer than frontal; interparietal small, very much smaller than anterior supraocular; occipital small, scarcely larger than posterior supraocular; nine supralabials, of which sixth (on right side, fifth on left side) is under the eye, and much longer and higher than the others, its anterior portion overlying the labial anterior to it; temporals small, slightly keeled, about 12 in a row between orbit and ear-opening; an elongated narrow shield along the anterior portion of the outer edge of parietals; three pairs of chin-shields, increasing in size posteriorly, only first pair in contact throughout, second pair posteriorly separated by a single elongated scale; dorsal scales in eight longitudinal series of larger scales, four on each side, and two median rows of smaller scales, all keeled, the keels of the larger scales forming continuous, nearly parallel ridges, five of the large dorsals corresponding in length to four ventral plates; laterals granu-

^a From *σμάργαδιος*, emerald green.

lar, forming a broad band; gulars large granules posteriorly, merging into large, pointed, keeled scales, covering the lower neck and collar, about 26 granules and scales on the median line between second pair of chin-shields and collar; six rows of ventral plates, each strongly keeled and somewhat pointed, especially those in front and behind, 26 on the median line between collar and preanal plate; a series of somewhat enlarged, keeled scales on each side of the abdomen, separated from the outer row of ventral plates by two or three rows of smaller keeled scales or granules; preanal plate large, with two keels, with a small keeled scale on each side; a single inguinal pore on each side; legs long, extended foreleg reaching the shoulder; tail three and a third times as long as head and body together, swollen at base, tapering nearly to a point, and covered above and below with strongly keeled scales, larger than the largest dorsals, about four of the former corresponding to six of the latter. Color (in alcohol) above greenish slate (in life said to be "emerald-green"); a narrow black line from middle of lores through eye and over lower temporal region, through ear to shoulder, bordering above a bright yellow stripe on the upper labials, through lower part of ear over shoulder and on the sides to the anterior side of the femur, following the enlarged series of scales on the flanks; yellow stripe continued behind femur on sides of basal portion of tail; under side greenish yellow, the green occupying lower half of subocular and labial behind it.

Dimensions.

	<i>mm.</i>
Total length.....	150
Snout to vent.....	45
Vent to tip of tail.....	105
Snout to collar.....	16
Snout to posterior edge of occipital.....	12
Greatest width of head.....	5.5
Fore leg.....	17
Hind leg.....	23

The *female* (U.S.N.M. No. 16360; same locality and collector) differs in having the preanal shield divided so that each half forms a large, strongly keeled scale, which is pointed behind; the base of the tail tapering, not swollen.

Variation.—The characters of this species are fairly constant. Thus far three pairs of chin-shields and one inguinal pore have been found in all the specimens examined. The small shield between the first supraocular and the loreal is not always present, and judging from Boulenger's original description the series of granules separating the supraoculars from the superciliaries is not always complete. The number of longitudinal dorsal scale rows is rather variable. In two of the specimens examined by me (male from Okinawa and female from Miyakoshima) there were four large rows on each side with two small ones between, while one (female from Okinawa) has three large

ones and the intervening two rows rather larger than in the other two specimens. Boulenger also describes the species as having only three enlarged dorsal rows on each side. He furthermore describes it as having eight rows of large ventrals, while the three specimens examined by me only have six. From his description it also appears that some of the females have the preanal plate undivided.

In the two females examined by me there are three rows of small, subequal, keeled scales on each side of the outer row of large ventrals instead of the one enlarged row and intervening small scales or granules as described in the male. Whether this difference is sexual or individual I am unable to say.

Remarks.—The present species, by having only three pairs of submental shields or chin-shields, clearly belongs to the southern group of species constituting this genus, such as *T. formosanus*, *T. septentrionalis*, etc., as distinguished from the forms with four chin-shields inhabiting more northern localities, such as *T. tachydromoides* in Japan and *T. amurensis* in Korea. It is also probably most nearly related to the former, its chief distinction from *T. septentrionalis* being its much more slender and elongated build.

Habitat.—Although first collected by W. Heine in May, 1853, during Perry's visit to Okinawa shima, this species was not made known until 1887, when Boulenger described and figured specimens obtained by Pryer in the Riu Kius. In point of fact it was observed even earlier, for Doctor Adams mentions having seen it on the Sakishima group when surveying there in 1845. From the latter islands we have a specimen collected in Miyakoshima obtained from the Science College Museum in Tokyo.

List of specimens of Tachydromus smaragdinus.

C. S. N. M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Inguinal pores.
16360	Female ^a	Great Loo Choo	May, 1853	W. Heine	1
16361	Male ^b	do	do	do	1
36521	Female ^c	Miyakoshima		Sci. Coll., Tokyo	1

^a P. 237.

^b Description, p. 236.

^c Sci. Coll. No. 29, figs. 203-204

TAKYDROMUS TACHYDROMOIDES^a (Schlegel)

KANAHEBI.

Plate XVIII.

1838. *Lacerta tachydromoides* SCHLEGEL, Fauna Jap. Rept., pp. 101, 139; Saur. & Batr., pl. 1, figs. 5-7 ^b (type-locality, Nagasaki; types in Leiden Mus.; Siebold, collector).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, 1891, p. 239 (Yezo).—*Tachydromus tachydromoides* BOULENGER, Cat. Liz. Brit. Mus., III, 1887, pp. 5, 509 (part: Japan; Yokohama; Hakone Lake;

^a From *Tachydromus* and εἶδος, similar, like.

^b Reproduced in this work on Plate XVIII.

Miyanoshta).—GUENTHER, Ann. Mag. Nat. Hist. (6), I, Mar., 1888, p. 169 (Japan).—OKADA, Cat. Vert. Jap., 1891, p. 70 (Tokyo; Oshima; Hakone Mts.).—BOETTGER, Kat. Rept. Mus. Senckenberg., I, 1893, p. 79 (Nagasaki; Yokohama; Yezo).


1839. *Tachydromus japonicus* DUMÉRIL and BIBRON, Erpét. Gén., V, p. 161 (type-locality, Japan; types in Mus. Paris; Siebold, collector).—DUMÉRIL, Cat. Mét. Rept. Mus. Paris, I, 1851, p. 118 (Japan).—HALLOWELL, Proc. Phila. Acad., 1860, p. 495 (Oshima, Idzu).—HILGENDORF, Sitz. Ber. Berlin. Naturf. Fr., 1880, p. 112 (Tokyo; Hakodate).—*Tachysaurus japonicus* GRAY, Cat. Liz. Brit. Mus., 1845, p. 52 (Japan).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1866, p. 111 (Yokohama).
1860. *Tachydromus sexlineatus* HALLOWELL, Proc. Phila. Acad., 1860, p. 495 (not of Daudin) (Simoda).
1895. *Tachydromus holsti* BOULENGER, Proc. Zool. Soc. London, 1894, p. 733, pl. XLIX, figs. 1-1b (type-locality, Mt. Onsen, near Shimabara, Prov. Hizen, Kiusiu; types in Brit. Mus. No. 94. 1. 25. 8. 32; Holst, collector).

Tachydromus sexlineatus Daudin is sometimes doubtfully attributed to Japan, apparently on Hallowell's authority^a who among his Japanese specimens distinguished some as *T. sexlineatus* and others as *T. japonicus*. Only the former are now in the National Museum (No. 7310, five specimens from Simoda), and I can affirm that they are nothing but the regular *T. tachydromoides*. They have the median dorsal scales rather small and the ventral scales are well carinated; hence probably Hallowell's identification.

Description (figs. 205-208).—*Adult male*: U.S.N.M. No. 31844; Yamagawa, Prov. of Satsuma, Kiusiu; June 14, 1904; Dr. Hugh. M. Smith, collector. Rostral scarcely in touch with internasal, not entering nostril; first supralabial enters nostril; anterior nasals barely in touch behind rostral; posterior loreal not much larger than anterior; internasal shorter than prefrontals which are nearly three-fifths the length of the frontal; two large supraoculars, the anterior separated by a small scute from the posterior loreal, the second followed by a small third supraocular; anterior superciliary in contact with first large supraocular, the others separated from the supraoculars by a single series of granules; frontoparietals in contact with posterior supraoculars, slightly longer than the prefrontals; parietals shorter than frontal; interparietal about one-fourth the size of anterior supraocular; occipital small, about the size of third supraocular; eight supralabials, fifth (on right side, sixth on left side) under the eye, its upper, or orbital edge nearly three times as long as its lower, or labial, edge; temporals rather large, keeled, about six in a row between orbit and ear-opening; an elongated shield, followed by two smaller ones along the outer half of the parietal; four pairs of chin-shields, increasing in size posteriorly, first two pairs in contact throughout, the third only in contact anteriorly, separated posteriorly by a wedge of granules; six series of large keeled dorsal scales, three on each side of median line, with a small interrupted median series consisting of keeled scales

^aProc. Phila. Acad., 1860, p. 495.


less than one-sixth the size of the others; keels of large scales forming continuous ridges, four of the large dorsals corresponding in length to about four and a half ventral plates; sides covered with minute keeled scales or large granules; scales on upper surface of limbs like dorsals, those on fore legs smaller, on hind legs of the same size; gulars granular, merging gradually into the pointed and keeled scales which cover lower neck and collar, about 23 granules and scales on the median line between third pair of chin-shields and collar; eight rows of ventral plates, the outer rows on each side smaller, the anterior ones somewhat keeled and pointed posteriorly, 27 on the median line from collar to preanal plate; adjoining the ventral plates on each side of the middle portion of the body three rows of small keeled scales which anteriorly and posteriorly grade into the lateral granules; preanal plate large, smooth, with two smaller plates on each side; two inguinal pores on each side; limbs moderate, the outstretched hind leg reaching the axilla; tail less than three times (about two and



FIGS. 205-208.—TAKYDROMUS TACHYDROMOIDES. 2 X NAT. SIZE. 205, TOP OF HEAD; 206, SIDE OF HEAD; 207, UNDERSIDE OF HEAD; 208, INGUINAL REGION. No. 7310, U.S.N.M.

two-thirds times) as long as head and body together, swollen at base, covered with strongly keeled scales which are about as long as the largest dorsals. Color (in alcohol)* on back olive brown with black tips to the scales forming two interrupted dorsal bands continued on the tail, a median series of ill-defined blotches and a dorso-lateral stripe on each side on the outer half of the outer row of large dorsals; in the inner half of this row a poorly defined pale stripe; a narrow black stripe from nostril through eye, narrowly rimming both eyelids, bifurcating behind the eye, the upper branch continuing backward and joining the dorso-lateral stripe, the lower proceeding obliquely through the ear-opening, over the shoulder and along the sides to the groin; the continuation of this stripe visible behind the femur on the side of the base of the tail as a series of black spots; below this black stripe a somewhat wider white stripe which becomes irregular on the

* In Stimpson's MS. catalogue the *Sinoda* specimens (May, 1855) are described in life as "above dark coppery-brown, below white (like white lead)."



TAKYDROMUS TACHYDROMOIDES. (FROM SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

flanks and stops at the groin, and below this another black stripe from the upper edge of the subocular; upper side of limbs with blackish reticulations; supralabials obscurely marked with dusky; lower side whitish.

Dimensions.

	<i>mm.</i>
Total length.....	207
Snout to vent.....	57
Vent to tip of tail.....	150
Snout to collar.....	20
Snout to posterior edge of occipital.....	13.5
Snout to ear-opening.....	13.5
Greatest width of head.....	8.5
Fore leg.....	21
Hind leg.....	31

The *female* (U.S.N.M. No. 31841, same locality and collector) differs in the tail tapering from the base which is not swollen; the hind legs are also shorter, the tip of the longest toe of extended hind leg barely reaches beyond the elbow.

In the *young* (U.S.N.M. Nos. 13748, 13749, from snout to vent, respectively, 20 and 24 mm.) the interparietal is relatively very large, being as large as or larger than the internasal.


Variation.—The number of chin-shields, or submandibulars, seems to be subject to less variation than almost any other character, there being four pairs in all the 66 specimens examined by me, except No. 31842, which has five irregular pairs, and No. 31889, which has only three, but showing indication of fusion between the two anterior pairs.^a

The number of inguinal pores is also very constant, all the 66 specimens having two on each side. Only two specimens (Nos. 12714 and 31886) show indication of a third pore on one side, the scale next to the outer pore on one side having a minute dusky spot. Doctor Guenther states, however, that he has seen a specimen with three pores, and if the figure of this species in *Fauna Japonica*^b is correct we have

^aBoulenger (Proc. Zool. Soc. London, 1899, pp. 161-162) lays stress upon an alleged exception to the rule that *T. septentrionalis* has three chin-shields only, as "not due to fusion or accidental division, as shown by the figure (p. 162)." The figure quoted shows a specimen with three chin-shields on one side and four on the other. The reason why the abnormality is considered not due to fusion or division is evidently the asymmetric arrangement, inasmuch as the suture between the second shield from behind and the shield anterior to it on each side are not on a line with each other, the assumption being that these sutures always correspond in normal specimens having three or four pairs, respectively. Such is not always the case, however, as will be seen from the accompanying figure from a specimen (No. 23539) brought home by me from Yokohama (fig. 212). If in this specimen the first two shields on the left side were fused the result would be exactly like the one figured by Boulenger (only reversed). The abnormality in his specimen, therefore, may easily be due to division.

^bReproduced on Plate XVIII (fig. 2) of this work.

another to record. The carination of the ventrals, gulars, and temporals is quite variable, but seems to be strongest in the younger specimens; in the one from Sapporo, Yezo (No. 23437), a very large male, even the gulars are smooth. This specimen also has one or two of the central temporals greatly enlarged and smooth. The number of longitudinal rows of dorsals is also subject to considerable variation, the common arrangement being three enlarged series on each side with a single or sometimes double series of very small keeled scales between the two median rows; these scales in case of a single series alternating on the right and left side of the middle line; in a number of specimens, however, there are only two series of very large scales on each side with a double series of smaller ones between them, but in this case the median scales are considerably larger than where there are three large series on each side. There is also considerable variation in the way the rostral, fronto-nasal, and nasals meet (though in most cases the



FIGS. 209-212.—*TAKYDROMUS TACHYDROMOIDES*. 2 × NAT. SIZE. 209, TOP OF HEAD; 210, MIDDLE OF BACK; (No. 23437, U.S.N.M.) 211, TOP OF HEAD; (No. 13753, U.S.N.M.) 212, CHIN-SHIELDS. No. 23539, U.S.N.M.

rostral and fronto-nasal are broadly in contact), as well as in the relation between the fronto-nasal and the frontal, there being sometimes a small unpaired squarish shield interpolated between these two shields and the prefrontals. Finally we may note that the granules between the supraoculars and the superciliaries are subject to considerable variation. While in most specimens the anterior elongate superciliary is followed by a continuous series of granules which separate the second supraocular from the superciliaries, this series is often broken. In one specimen (No. 13753, fig. 211) it is reduced to such an extent that the second supraocular is entirely in contact with the superciliaries, the two granules left only separating the posterior end of the first supraocular from the latter, and in another (No. 34143, one out of a lot of 20 from Mount Fuji) both supraoculars are in contact throughout with the superciliaries, being a perfect *T. holsti* in this respect. On the other hand, in one specimen (No. 23437) the one

from Yezo, and in many of the specimens from Kiusiu and Shikoku, the granules are so much developed as to entirely separate both the first and the second supraoculars from the outer series of elongated superciliaries (fig. 209). In the majority of the specimens from the southern lowland localities I also find several granules bordering the anterior end of the first large supraocular. In the Hondo specimens there is a greater uniformity inasmuch as in nearly every one of these the first large supraocular is broadly in contact with the long supra-ciliary. There is not enough constancy in these characters to base a separation upon them, and it should be noted that the relation of these shields in the southern specimens tends in a direction away from *T. holsti* rather than toward it.^a

The coloration varies considerably; the male described above shows almost a maximum of dark and light markings, the female has the markings nearly obsolete except on the sides of the head; between these extremes there are all transitional stages, though the females are less heavily marked than the males; in many specimens the pale dorso-lateral line is more definitely marked than in the one described; in others it is more or less broken up by dark spots from the adjoining dorso-lateral black line; the sides are often solidly dark brown, in other specimens again more or less marked with pale or whitish spots.

Habitat.—The present form of grass lizard is apparently restricted to Japan proper. The range was formerly thought to include the adjacent portions of the mainland, but these are inhabited by other species, such as *T. septentrionalis* and *T. wolteri*, with which it was at one time confounded.

The types came from the neighborhood of Nagasaki, and it is a curious fact that von Siebold and Buerger failed to get specimens outside of Kiusiu. It is, nevertheless, rather common in Hondo; thus specimens are recorded from Yokohama, Tokyo, Simoda, Oshima, and the region around Hakone Lake. Yezo specimens are also in various museums, and our National Museum has a fine one from Sapporo, Yezo, thanks to the kindness of Doctor Nozawa.

Dr. Hugh M. Smith collected a fine series of specimens both at Yamagawa, at the entrance of the Bay of Kagoshima, Kiusiu, as well as near Kochi, on the southern coast of Shikoku, the first record from this island.

Four specimens (U.S.N.M. Nos. 34179 to 34183), absolutely indistinguishable from those collected near Yokohama, are labeled by Mr. Owston, from whom the National Museum recently acquired

^a Since writing the above I have had an opportunity to examine the type specimens of *T. holsti*, thanks to the kindness of Doctor Boulenger. This examination substantiates what is said above, and Doctor Boulenger agrees now with me that the specimens do not represent a separable form.

them, as coming from "Northern Formosa, June 23, 1903." Needless to say some mistake has been made in labeling and that they never were collected in Formosa. In all probability they became confused with a specimen of *Natrix piscator* credited to "Mount Fiji," and the lizards were really collected near or on this mountain during August, 1898. (See under *Natrix piscator*, p. 290.)

List of specimens of *Takydromus tachydromoides*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Ingenial pores.
U.S.N.M.	7310a	Adult a	Simoda, Hondo	May, 1855	W. Stimpson	2
Do	7310b	do	do	do	do	2
Do	7310c	do	do	do	do	2
Do	7310d	do	do	do	do	2
Do	7310e	do	do	do	do	2
Do	11332	Young	Japan	1878	E. S. Morse	2
Do	11333	do	do	1878	do	2
Do	11334	Male	do	1878	do	2
Do	12714	do ^b	[Yokohama]	Apr., 1881	P. L. Jouy	2-3
Do	13748	Young	do	do	do	2
Do	13749	do	do	do	do	2
Do	13752	Female	do	do	do	2
Do	13753	do ^c	do	do	do	2
Do	23335	do	Yokohama	Sept., 1896	L. Stejneger	2
Do	23437	Male d	Sapporo, Yezo	do	S. Nozawa	2
Do	23539	Young e	Yokohama	Oct., 1896	L. Stejneger	2
Do	30732	Female	Miyazaki, Kiusiu	do	C. A. Clark	2
Do	30733	Young	do	do	do	2
Do	30740	do	do	do	do	2
Do	31797	Female	Onomachi, Aki, Hondo	June 4, 1904	H. M. Smith	2
Do	31820	do	Yamagawa, Kiusiu	June 14, 1904	do	2
Do	31838	Male	do	do	do	2
Do	31839	do	do	do	do	2
Do	31840	do	do	do	do	2
Do	31841	Female ^b	do	do	do	2
Do	31842	Male f	do	do	do	2
Do	31843	Female	do	do	do	2
Do	31844	Male g	do	do	do	2
Do	31845	do	do	do	do	2
Do	31884	do	Kochi, Shikoku	May 11, 1904	do	2
Do	31885	do	do	do	do	2
Do	31886	do ^b	do	do	do	2-3
Do	31887	do	do	do	do	2
Do	31888	Female	do	do	do	2
Do	31889	do ^h	do	do	do	2
Do	34140	do	Mount Fuji	Aug., 1898	A. Owston	2
Do	34141	do	do	do	do	2
Do	34142	Male	do	do	do	2
Do	34143	Female i	do	do	do	2
Do	34144	do	do	do	do	2
Do	34145	do	do	do	do	2

^a P. 240; figs. 205-208.

^b P. 241.

^c P. 242; fig. 211.

^d P. 242; figs. 209, 210.

^e P. 242; fig. 212.

^f 5 pairs of chin-shields.

^g Description, p. 239.

^h 3 pairs of chin-shields.

ⁱ P. 242.

List of specimens of *Takydromus tachydromoides*—Continued.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Inguinal pores.
U. S. N. M.	34146	Female	Mount Fuji	Aug. 1898	A. Owston	2
Do	34147	do	do	do	do	2
Do	34148	do	do	do	do	2
Do	34149	Male	do	do	do	2
Do	34150	do	do	do	do	2
Do	34151	Female	do	do	do	2
Do	34152	Male	do	do	do	2
Do	34153	Female	do	do	do	2
Do	34154	do	do	do	do	2
Do	34155	Male	do	do	do	2
Do	34156	Female	do	do	do	2
Do	34157	Male	do	do	do	2
Do	34158	Female	do	do	do	2
Do	34159	Male	do	do	do	2
Do	34179	Female	do ²	[Aug., 1898 ²]	do	2
Do	34180	do	do	do	do	2
Do	34181	do	do	do	do	2
Do	34182	Male	do	do	do	2
Do	34183	do	do	do	do	2
Do	34184	Female	Yokohama	Aug., 1884	do	2
Do	36536	Male ^a	Hondo		Sci. Coll. Tokyo	2
Do	36537	Young ^a	do		do	2
Do	36538	do. ^a	do		do	2
Do	36539	do. ^a	do		do	2
Brit. Mus.	94. 1. 25. 8. 32	Female ^b	Mount Onsen, Kiusiu		P. Holst	2
Do	94. 1. 25. 8. 32	Young ^b	do		do	2

^a Sci. Coll. No. 53.^b Types of *T. holsti*.TAKYDROMUS AMURENSIS ^a Peters.

1881. *Tachydromus amurensis* PETERS, Sitz. Ber. Berlin Naturf. Fr., 1881, p. 71 (type-locality, Kossakewitcha, Amurland; type, Berlin Mus. No. 9869).—FISCHER, Jahrb. Hamburg. Wiss. Anst., II, 1885, p. 84.—BOULENGER, Cat. Liz. Brit. Mus., III, 1887, p. 6 (Kossakewitcha, Amurland); Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 137 (Khabarovka; Duerries, collector).—MUELLER, Verh. Naturf. Ges. Basel, VIII, Pt. 2, 1887, p. 285 (Khabarovka).—GUENTHER, Ann. Mag. Nat. Hist. (6), I, March 1888, pp. 168, 169 (Amurland).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, No. 1, 1905, p. 92 (Ussuri; Vladivostok).


The introduction of *T. amurensis* into the Korean fauna is based upon a specimen (No. 21185, U.S.N.M., figs. 213–216) collected by the late P. L. Jouy in 1883 (no definite locality). It agrees in every particular, except one, with the various descriptions of *T. amurensis*, widely differing from *T. tachydromoides*. It has 3 inguinal pores on each side; dorsal scales in 8 rows, the median ones but slightly smaller than the others; the toes are shorter, the fourth toe from base of

^a From the Amur country.

third, for instance, is not longer than the width of the head; the larger scales on fore neck, as well as those forming the collar, are wider, rounded at the end and smooth; no row of granules between supraoculars and superciliaries, only one or two isolated granules on the right side; on one side 1 scale between first supraocular and loreal, on the other 2; 4 pairs of submentals; 4 keels on upper surface of tail; color above brownish olive with black spots, below whitish, but with no trace of any white band between eye and ear and beyond.

The only difference between Boulenger's descriptions of the type from Kossakewitcha, on the Amur, and four specimens from Khabarovka, which expressly state that the temporals are "perfectly smooth," consists in these scales being obtusely keeled in our specimen.

On the other hand, were it not for the 3 inguinal pores, our specimens might with equal propriety be referred to *T. wolteri*, originally described from Korea (Chemulpo). The latter has only one inguinal pore on each side. Fischer, in the original description, does not



FIGS. 213-216.—TAKYDROMUS AMURENSIS. 2 × NAT. SIZE. 213, TOP OF HEAD; 214, SIDE OF HEAD; 215, MIDDLE OF BACK; 216, INGUINAL REGION. No. 21185, U.S.N.M.

mention whether the temporals are keeled or smooth, but from the fact that Boulenger, with the type before him, refers it to *T. tachydromoides*, I feel justified in inferring that it has keeled temporals. However, as I find a great difference in the amount of carination of the temporals in typical Japanese *T. tachydromoides*, I am inclined to place but little faith in this character.

The question then arises whether in reality *T. wolteri* is distinct from *T. amurensis*. The material is as yet too scant (3 specimens in British Museum, viz, the type from Chemulpo, one specimen from Kiukiang, China, and one from an unknown locality; Guenther, Ann. Mag. Nat. Hist. (6), I, 1888, p. 168) to permit any authoritative decision, and I therefore abide provisionally by that of Doctor Guenther and recognize it as distinct, although with considerable doubt.

A detailed description seems superfluous after the above remarks.

Habitat.—*T. amurensis* has been recorded so far only from the Amur district. In addition to the specimens from Kossakewitcha

and Khabarovka, it has been collected by Maximowitch in 1864 at the rivers Olakha (Ussuri) and Dadsoshu, and by Pleske at Vladivostok in 1883.

Its occurrence in Korea is mentioned here for the first time.

List of specimens of Takydromus amurensis.

U.S. N.M. No.	Sex.	Locality.	When collected.	By whom collected.	Inguinal pores.
21185	Female ♀	Korea	1883	P. L. Jouy	3

^a Figs. 213-216.

TAKYDROMUS WOLTERI ^a Fischer.

1885. *Tachydromus wolteri* FISCHER, Jahrb. Wiss. Anst. Hamburg., II, p. 82 (type-locality, Chemulpo, Korea; type in Brit. Mus.).—MUELLER, Verh. Naturf. Ges. Basel, VIII, Pt. 2, 1887, p. 285 (Seoul, Korea).—GUENTHER, Ann. Mag. Nat. Hist. (6), I, March 1888, pp. 168, 169 (Kiukiang, China).

For the status of this species see the remarks under *T. amurensis*, from which it is said to differ by having only one inguinal pore on each side and a very distinct white lateral band from eye backward.

T. wolteri was reduced by Boulenger to a synonym of *T. tachydromoides* at a time when in the latter he also included *T. septentrionalis*. He has since admitted the distinctness of *T. septentrionalis*, but there is apparently no record of where he would place *T. wolteri*. It can not well go with *T. septentrionalis* because of its four pairs of chin-shields. On the other hand, it seems impossible to join it to *T. tachydromoides*, as it has eight subequal dorsal rows. This latter character links it distinctly with *T. amurensis*. One of the characters by which the original describer, Doctor Fischer, attempted to separate it from the latter, viz, the comparatively large size of the median dorsal series, is not a valid character, as our Korean specimen of *T. amurensis* has all the dorsal rows subequal concomitant with three inguinal pores.

Habitat.—*T. wolteri* was originally described from a specimen collected at Chemulpo, Korea, and Mueller states that one specimen from Seoul is in the Basel Museum. Doctor Guenther has since recorded another from Kiukiang, east central China, but it is probably permissible to suggest that the latter may be a *T. septentrionalis* with an abnormal number of chin-shields.

Genus EREMIAS ^b Wiegmann.

1834. *Eremias* WIEGMANN, Herpet. Mex., p. 9 (type, *Lacerta velox*).

1841. *Aspidorhinus* EICHWALD, Fauna Casp.-Cauc. (p. 74) (same type).

^a For Mr. C. Wolter, who collected at Chemulpo. ^b From ἔρημος, alone, lonely.

EREMIAS ARGUS^a Peters.

1859. ? *Eremias velox* MAACK, Putesch. na Amur (p. 152) (Amurland) (not of Pallas).
1869. *Eremias argus* PETERS, Mon. Ber. Berlin Akad. Wiss., 1869, p. 61, pl. —, fig. 3 (type-locality, Chefu, China; types, Berlin Mus. No. 4532; Schottmueller, collector).—STEINDACHNER, Sitz. Ber. Wien Akad. Wiss., Math.-Naturw. Cl., LXII, Pt. 1, 1870, p. 336; author's separate, p. 11, pl. II, figs. 1, 2 (Peking).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1876, p. 375 (Chefu).—MOELLENDORFF, Journ. N. China, Br. R. Asiat. Soc. (n. s.), 1877, p. 103 (Peking).—MUELLER, Verh. Naturf. Ges. Basel, VIII, Pt. 2, 1887, p. 286 (Chemulpo, Korea).—BOULENGER, Cat. Liz. Brit. Mus., III, 1887, p. 102 (part: Chefu; Peking; Manchuria?); Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 138 (Manchuria).—BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, pp. 63, 122 (Miau-feng-shan, Peking; Herz, collector); Kat. Rept. Mus. Senckenberg., I, 1893, p. 94.—FISCHER, Jahrb. Hamburg Wiss. Anst., V, 1888, p. 46 (eastern Mongolia).—STONE, Proc. Phila. Acad., 1899, p. 184 (Klingan Mts., eastern Mongolia).—WERNER, Abh. Bayer. Akad. Wiss. Muenchen, II Klasse, XXII, Pt. 2, 1904, p. 354 (Tsingtau and Kiautschou, China).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, (8) XVII, No. 1, 1905, p. 167 (Mongolia, Manchuria, Korea).—*Podarces (Eremias) argus* STRAUCH, in Przevalski's Mongoliya i Strana Tangutov, III, 1876, p. 30 (Ordos).


Whatever opinion one may hold about the status of *Eremias brechleyi*,^b there can be but little doubt that our specimens from Korea are typical *E. argus*, as the subocular rests on several supra-labials without reaching the lip; fronto-parietals are larger than second supraocular; the lower nasal does not reach the rostral, except in No. 21184; the number of scales or granules in a row between parietals and lip is about 10; the number of granules and ventrals in a row around the middle of the body between 65 and 71. Moreover, the coloration is typically that of *E. argus*, with two well developed median dorsal rows of ocelli and no lateral dark band. The scutellation on top of the head is exactly as figured by Peters, except that the smallest specimen has a second scale between the prefrontals, but it is so small as to scarcely be more than a granule. Our specimens also agree closely with Steindachner's figures of a specimen from Peking.

^a From *ἄργος*, the many-eyed Argus, slain by Hermes; referring to the many ocellated spots of this species.

- ^b 1872. *Eremias brechleyi* GUENTHER, Ann. Mag. Nat. Hist. (4), X, 1872, p. 419 (type-locality, Land of Grass, Mongolia; type in Brit. Mus.; Brechley, collector); Brechley's Cruise 'Curaçao,' 1873, p. 396, pl. xxii, fig. A.—FISCHER, Jahrb. Hamburg Wiss. Anst., V, 1888, p. 46, pl. iv, fig. 9 (Tehikiang; Oldenburg Mus.).—MÉHELY, Zichy's Dritte Asiat. Forschungsr., II, 1901, p. 56 (Khalgan, Chinese Wall, 760 m. alt.; Nankou Pass).—*Podarces (Eremias) brechleyi* STRAUCH, in Przevalski's Mongoliya i Strana Tangutov, III, 1876, p. 32 (Ordos; eastern Mongolia).
1887. *Eremias argus* BOULENGER, Cat. Liz. Brit. Mus., III, p. 102 (part: Mongolia).
1888. *Eremias argus* var. *brechleyi* BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., pp. 63, 122 (Miau-feng-shan, Peking; Herz, collector); Kat. Rept. Mus. Senckenberg., I, 1893, p. 94.

Thanks to the kindness of Mr. Witmer Stone, curator of the Philadelphia Academy Museum, I have been enabled to examine two specimens recorded by him from eastern Mongolia. One of them is a typical *E. argus* in every respect. The other has only a very small, single median fronto-nasal and a pair of very long prefrontals. It consequently approaches *E. arguta*, though a close inspection seems to show that the prefrontals have become so long by fusion with a pair of fronto-nasals, the shield which now appears in that rôle being probably only the ordinary azygous interprefrontal pushed forward so as to touch the supranasals. In these specimens the light centers of the outer rows of ocelli are nearly confluent, so as to form longitudinal light lines.

Description.—*Adult female*; U.S.N.M. No. 21182; Seoul, Korea; June, 1883; P. L. Jouy, collector (figs. 217–219). Rostral pentagonal, in contact with first supralabials and supranasals; nostril a round hole between three bulging nasals, of which the supranasals are more than twice as large as the other two together and broadly in



FIGS. 217–219.—*EREMIAS ARGUS*. 2X NAT. SIZE. 217, TOP OF HEAD; 218, SIDE OF HEAD; 219, FEMORAL PORES AND ANAL REGION. No. 21182, U.S.N.M.

contact behind the rostral; postnasal small, in contact with interparietal and first loreal; subnasal long and narrow, in contact with first and second labials, first loreal, and the two other nasals; a pair of internasals behind the supranasals and smaller than the latter; a pair of pentagonal prefrontals separated by a small median azygous spear-shaped prefrontal, the former in contact externally by second loreal and first superciliary; frontal long, slightly longer than its distance from rostral, twice as wide anteriorly as posteriorly, in contact posteriorly with the first supraocular; three supraoculars, the first two large, the third very small, separated from the superciliaries, the first by a single row and the second by a double row of granules; the space in front of first supraocular, between that, frontal, prefrontal, and first superciliary filled with granules, the row nearest frontal and prefrontals being somewhat larger than the others; five superciliaries, the first longest and in contact with prefrontal; frontoparietals larger than second supraoculars, almost as long as parietals,

in contact with third supraocular; parietals shorter than frontal, in contact behind interparietal, which is small, only, little larger than third supraocular; a long and narrow shield along the outer edge of each parietal; two loreals, the anterior rather small, triangular, and in contact with internasal; a long subocular in contact with four supralabials, not reaching the edge of the lip; nine supralabials, fifth longest, sixth directly under the subocular, smaller than seventh; over the seventh labial a large lower postocular; lower temporals large, smooth, upper ones granular, about ten granules and scales in a row between parietals and supralabials; six pairs of enlarged submandibulars, or chin-shields, third pair largest, sixth smallest, first three pairs in contact; upper surface covered with small, uniform granules, about 58 in a row across the middle of the back between the ventrals; gulars large flat granules followed by a few rows of smaller ones gradually merging into the flat scales of the lower neck; collar consisting of about nine large, smooth scales, of which the median one is the largest; fold behind collar covered with small granules; ventral plates subequal, twelve in a row across the abdomen, and not forming longitudinal series, smooth, about 30 shields in a row on the median line between collar fold and groin; anterior aspect of arm with large smooth scales, which on the forearm assume the shape of transverse plates; anterior aspect of femur and lower side of tibia with a series of broad transverse plates; scales on underside of digits sharply keeled; a large median preanal plate with a deep cleft behind; a series of 11 femoral pores on each side; tail covered with elongate, squarish, keeled scales arranged in verticels, about 24 in a verticel, a head length from the vent. Color (in alcohol) dull clay-colored above with about eight longitudinal rows of yellowish spots surrounded by a black ring on the back, these ocelli being continued on the base of the tail; underside whitish.

Dimensions.

	<i>mm.</i>
Total length.....	110
Snout to vent.....	51
Vent to tip of tail.....	59
Snout to collar.....	17
Snout to ear-opening.....	12.5
Snout to posterior edge of parietals.....	11.5
Greatest width of head.....	8
Fore leg.....	18
Hind leg.....	26

Variation.—The two other specimens (same locality and collector) before me are rather young. Both differ in having the paired prefrontals touch behind the azygous shield and in having a longer suture between the parietals behind the interparietal. No. 21183 lacks the posterior pair of submentals and in No. 21184 the lower nasal distinctly reaches the rostral; the former has about 65 granules and

plates around the middle of the body and 10 femoral pores, the latter 70 granules and plates around the body and 11 femoral pores on each leg.

Habitat.—Owing to the doubt whether certain references to *E. argus* may not in reality belong to *E. breuchleyi*, it is at present impossible to define the habitats of these two forms exactly. The difficulty is increased by the fact that both seem to occur together, in some localities, at least.

As far as can be made out at present *E. argus* occurs in northeastern China; thus at Chefu, whence came the types; at Kiautshou and Tsingtau, recorded by Werner; and at Peking, from the neighborhood of which city Moellendorff, Steindachner, and Boettger, have recorded several specimens. Fischer, who also distinguishes between the two forms, mentions specimens of true *E. argus* from "eastern Mongolia." During Count Zichy's third Asiatic voyage it was collected in the same general region, viz, at Shara-murun, in eastern Mongolia, and farther east on the road to Peking between Daba and Khalgan.

It was first recorded from Korea by F. Mueller, who presented a specimen from Chemulpo to the Basel Museum, and four other specimens are in the St. Petersburg Museum from the same locality, collected by Bunge in 1889 and 1891. The late P. L. Jouy, during June and August, 1883, obtained three specimens at Seoul, which are now in the United States National Museum.

List of specimens of Eremias argus.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected.	Femoral pores.
21182	Female <i>a</i>	Seoul, Korea.....	June, 1883	P. L. Jouy.....	11
21183	Young <i>b</i>	do.....	Aug., 1883	do.....	10
21184	do <i>b</i>	do.....	do	do.....	11

a Description p. 249, figs. 217-219.

b P. 250.

Genus LACERTA *a* Linnæus.

1758. *Lacerta* LINNÆUS, Syst. Nat., 10 ed., I, p. 200 (type, *L. agilis*); 12 ed., I, 1766, p. 359.

1830. *Zootoca* WAGLER, Nat. Syst. Amph., p. 155 (type, *L. vivipara*).

1851. *Atropis* GLUECKSELG, Lotos, 1851 (p. 138) (same type).

LACERTA VIVIPARA *b* Jacquin.

1787. *Lacerta vivipara* JACQUIN, Nova Acta Helvet., I (p. 33, pl. 1) (type-locality, Schneeberg, near Vienna, Austria).—DOBROTVOVSKI, Izvest. Sibirsk. Otd. Geogr. Obstch., I, 1870 (p. 23) (southern Sakhalin).—BEDRIAGA, Abh. Senckenberg. Naturf. Ges., XIV, Pt. 2, 1886, p. 338 (Padun, Baikal Lake; Sakhalin).—BOULENGER, Cat. Liz. Brit. Mus., III, 1887, p. 23


a Signifying lizard.

b Signifying bearing living young.

(Stanovoi Mts.; Niklayevsk, Amur; Sakhalin); Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 138 (Amurland).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg. LX, Prilozh., No. 5, 1889, p. 291 (Sakhalin); Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 115 (Europ. and Asiat. Russia; Sakhalin).—*Lacerta (Zootoca) vivipara* MIDDENDORFF, Sibir. Reise, II, II. Pt. 1, 1853, p. 247 (Stanovoi Mts.; Aldan Riv.).—*Zootoca vivipara* МААСК, Путеш. на Amur, 1859 (p. 152) (Amurland); Путеш. на Ussuri, 1861, (p. 189) (Ussuri Valley).

The numerous synonyms based upon European specimens are here omitted.

Description.—*Adult female*; U.S.N.M. No. 14385; Sakhalin Island; St. Petersburg Academy (figs. 220–222). Rostral pentagonal, not entering nostril, not in contact with internasal; nostril round, between two nasals; supranasals in contact behind rostral, also in contact with first supralabial, and anterior loreal; postnasal not in contact with second supralabial; anterior loreal high and narrow, in contact with supranasal, internasal, and prefrontal; posterior loreal twice as large as anterior; internasal much wider than long, not in contact with poste-



FIGS. 220–222.—*LACERTA VIVIPARA*. 2×NAT. SIZE. 220, TOP OF HEAD; 221, SIDE OF HEAD; 222, FEMORAL PORES AND ANAL REGION. No. 14385, U.S.N.M.

rior loreal, prefrontals smaller than fronto-parietals, in contact with first and second supraoculars; frontal hexagonal, with parallel sides, slightly longer than its distance from rostral, and equaling the length of the parietals; four supraoculars, anterior and posterior very small, all in contact throughout with superciliaries; first supraocular in contact with posterior loreal, second with prefrontal, third with frontal, and fourth with fronto-parietal; four superciliaries on one side, six on the other; fronto-parietals larger than prefrontals; interparietal pentagonal, twice as large as posterior supraocular, broadly in contact behind with occipital, which is slightly larger than posterior supraocular, hexagonal, and slightly protruding behind parietals; parietals not in contact, their external edge bordered by three small scutes not specially differentiated from the temporals; seven supralabials on left side, six on right, fifth or fourth, respectively, longer and higher than the others directly under the eye; temporals large, flat irregularly polygonal scutes, about four in a row between orbit and ear-opening.

the one at the upper anterior edge of the latter (the so-called tympanic) larger than the others; six lower labials on left side, five on right; six pairs of submandibulars, or chin-shields, the last pair very small; first two pairs of chin-shields in contact throughout, third pair separated posteriorly by a wedge of gular scales; back covered with bluntly keeled, slightly imbricated scales, the length of six from the middle of the back corresponding to the length of four ventral plates, these dorsals becoming gradually smaller and nearly smooth on upper neck and sides, about 30 in a row across the middle of the back between the ventrals; gular scales smooth, irregularly polygonal with rounded corners, increasing in size backward toward the collar, which consists of 10 large, smooth shields; fold behind collar covered with granules and small scales; ventral plates in six longitudinal rows, the two median rows narrower and the next row on each wider than the outer series, which is bordered externally by a series of large flat scales only slightly smaller than the median series of plates; plates between the arms not in regular longitudinal rows, the series next to the collar fold being oblique, lengthened, more or less pointed behind; 29 transverse rows of plates between collar and preanal plate; preanal plate very large, smooth; arm with a series of large smooth plates above the elbow and two below; anterior and lower surface of femur and tibia also covered with large smooth plates; eight pores on right thigh, ten on left; scales covering underside of digits smooth; tip of longest toe reaching wrist of fore leg; tail covered with squarish, elongated scales, as long as the ventral plates, those on the upper surface keeled, those on the basal half of the lower surface smooth. Color (in alcohol) olive clay-color above with two dorsal series of small whitish dots, each about covering one scale, edged on each side by a dark brown dot of similar size; a lateral series is also indicated as well as a few brown dots on the middle portion of the back; these marks, rather better defined, continued on the tail; underside paler olive gray, posterior ventrals obscurely marked with blackish, hind legs and tail underneath more heavily marked with dusky.

Dimensions.

	<i>mm.</i>
Total length (tail regenerated)	106
Snout to vent	57
Snout to collar	17
Snout to ear-opening	10
Snout to posterior edge of occipital	10
Greatest width of head	8
Fore leg	17
Hind leg	24

The *adult male* has relatively longer legs, especially hind legs, and tail, the latter being nearly twice as long as the head and body in the male, and only about one and a half times as long in the female.

Variation.—The present species is not very variable as far as scutellation is concerned. Usually the fifth supralabial is the subocular; the number of scales on the collar varies between 6 and 10; there are sometimes 8 regular longitudinal series of ventrals; femoral pores vary between 7 and 13.

The coloration is much more variable, at least in European specimens. The underside of the male is in Europe usually more or less orange colored, and in moist localities and higher altitudes melanistic specimens are not uncommon.

The young, as a rule, are much darker than the adults, nearly black with yellowish dots.

Habitat.—This is one of the most widely distributed species of lizards, inasmuch as it occurs in Europe from the Pyrenees and the Alps to the Arctic Ocean in East Finmark, Norway, and from the Atlantic in the west to the Okhotsk Sea and the Sea of Japan in the east. It is recorded from Nikolayevsk on the Amur River, from the valley of the Ussuri, as well as from the island of Sakhalin. The U. S. National Museum possesses a specimen from the latter locality, thanks to the liberality of the St. Petersburg Academy of Sciences. Nikolski reports that he met it not infrequently in the middle part of Sakhalin, that Poljakof collected two specimens near the mouth of the river Tim, and that Dobrotvorski observed it in the southern part of the island. He also enumerates numerous specimens collected by Maaek, von Schrenck, and others, among them a specimen from Decastries Bay collected by P. Semenow.

List of specimens of Lacerta vivipara.

U. S. N. M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Femoral pores.
1485	Female ^a	Sakhalin Island.....		Acad. St. Petersburg.....	8-10

^a Description, p. 252; figs. 220-222.

Suborder SERPENTES.

1758. *Serpentes* LINNÆUS, Syst. Nat., 10 ed., I, p. 214.

1759. *Serpentia* LINNÆUS, Syst. Nat., 11 ed., p. 76.

1802. *Ophidia* MACARTNEY, in Ross' Transl. Cuvier's Lect. Comp. Anat., I, tab. III.

1803. *Ophidii* DAUDIN, Hist. Nat. Rept., V, tab. p. 8.

1826. *Idiophides* FICINUS and CARUS, Uebers. Ges. Thierr., tab.

1828. *Strepsichrotos* RITGEN, Nova Acta Acad. Leop. Carol., XIV, p. 257.

1828. *Ophes* WAGLER, Isis, 1828, p. 860.

1839. *Ophides* SWAINSON, Nat. Hist. Class. Fish. Amph. Rept., II (Lardner's Cab. Encycl.), p. 134.

1849. *Pleuropoda* MAYER, Rheinlaend. und Westphal. Verhandl., VI (p. 177).

The characters separating the snakes from the lizards have been pointed out under the head of the latter, and it is only necessary here to recall the fact that all the reptiles occurring within the boundaries of this work, which have no external limbs are snakes. In southern China, ranging west to the Himalayas, there occurs a genus of legless lizards, *Ophisaurus*, of the family *Anguidæ*, but the species have an external ear-opening which at once distinguishes them from the snakes.

Several families of snakes are not represented in the fauna here treated of. The most important of these is that of the *Boidæ*, which contains the largest snakes, the boas and the pythons,^a the most remarkable external character of which is the rudiment of posterior limbs in the shape of a small claw-like spur on each side near the vent. They are found chiefly in the tropical regions of both the Old and the New World. The *Aniliidæ*, a small family with one representative in America and less than half a dozen species in south-eastern Asia, have also rudiments of hind legs. The *Leptotyphlopidae*, which have a very great external resemblance to the *Typhlopidae*, but with teeth only in the lower jaw, are confined to America and Africa, a few species only entering Asia in the southwestern part. The *Uropeltidæ* and *Xenopeltidæ* are two small families, the former occupying Ceylon and the mountains of southern India, the latter consisting of a single species inhabiting the Malayan Archipelago and peninsula as well as the adjacent portions of the mainland. Finally, the family *Amblycephalidæ* is a very interesting one, because of certain well-marked characters in the anatomy of the skull, together with the absence of the groove under the chin, so characteristic of the bulk of the snakes which the *Amblycephalidæ* resemble in most other external characters. The members of this family belong mostly to Central and South America, but two genera inhabit southeastern Asia, from the Himalayas to Hongkong south to the Malayan Archipelago and the Philippines.


The following table of the superfamilies, families, and subfamilies of snakes ascertained to occur within our limits gives the essential characters by which these divisions have been separated. There is still considerable divergence in the opinions of authors concerning the relative value of some of these divisions, but in breaking up of the aglyph and opisthaglyph "series" I believe that a better recognition of the true relationships of their component parts can be obtained.

^a *Python molurus* is sometimes included in the fauna of Formosa, on the strength of a specimen (skin) sent by Swinhoe. Doctor Guenther (Rept. Brit. India, p. 331), however, states that according to a communication from Swinhoe himself, the specimen had been imported into Formosa from China.

SYNOPSIS OF SUPERFAMILIES, FAMILIES, AND SUBFAMILIES.

- a*¹ (TYPHLOPOIDEÆ). No ectopterygoid (transpalatine); teeth in upper jaw only.
 TYPHLOPIDÆ, p. 259.
- a*² An ectopterygoid (transpalatine); teeth in both jaws.
- b*¹ (NATRICOIDÆ). Maxillary bone horizontal.
- c*¹ None of the anterior maxillary teeth grooved or perforated. NATRICIDÆ, p. 262.
- d*¹ Hypapophyses present throughout the vertebral column (figs. 223-225).
- e*¹ All maxillary teeth solid NATRICINÆ, p. 263.
- e*² Posterior maxillary teeth grooved HOMALOPSINÆ, p. 299.
- d*² Hypapophyses absent in posterior dorsal vertebrae (figs. 226-228).
- e*¹ All maxillary teeth solid CORONELLINÆ, p. 306.
- e*² Posterior maxillary teeth grooved BOIGINÆ, p. 381.
- e*² Anterior maxillary teeth grooved or perforated ELAPIDÆ, p. 386.
- d*¹ Tail cylindrical ELAPINÆ, p. 386.
- d*² Tail compressed HYDRINÆ, p. 400.
- b*¹ (CROTALOIDEÆ). Maxillary bone vertical.
- c*¹ No loreal pit COBRIDÆ, p. 442.
- c*² A deep loreal pit CROTALIDÆ, p. 448.

The above synopsis is chiefly based upon characters which either require more or less dissection or examination of the posterior maxil-



FIGS. 223-228.—POSTERIOR DORSAL VERTEBRÆ OF SNAKES. 223, BACK VIEW; 224, LOWER VIEW; 225, SIDE VIEW OF VERTEBRA WITH HYPAPOPHYSIS; 226, BACK VIEW; 227, LOWER VIEW; 228, SIDE VIEW OF VERTEBRA WITHOUT HYPAPOPHYSIS (COPIED FROM BOULENGER, CAT. SNAKES BRIT. MUS., I 1893, p. 171.)

lary teeth. The latter often presents difficulties, and while in some cases it must be resorted to in order to obtain absolutely reliable identification, a handy artificial key to the genera of snakes occurring within our territory may be useful in determining the greater majority of specimens.

This key is *not* satisfactory for the determination of the last four genera, and great care should be exercised in their identification. Specimens referred to any of these should be tested by the other keys and by comparison with the descriptions. There is no reliable external character by which to distinguish our species belonging to *Natrix* from those of *Elaphe*. However, a specimen with smooth or nearly smooth scales (with pores) is usually an *Elaphe*, and when more strongly keeled, if it has a small scale below the preocular (a sub-preocular), it also belongs to the latter. *Natrix* never has this sub-preocular. *Zamenis* has it, but can be recognized by the very long and narrow shields on top of the head.

The land snakes occurring in the four large islands of Japan can be easily determined by the following artificial key:

ARTIFICIAL KEY FOR THE IDENTIFICATION OF THE SNAKES OF JAPAN PROPER.


- a*¹ A deep hole or pit between nostril and eye.....*Agkistrodon blomhoffii*, p. 457.
*a*² No hole between nostril and eye.
*b*¹ Subcaudals entire; no postoculars, temporals joining eye.
Achalinus spinalis, p. 297.
*b*² Subcaudals in pairs; postoculars, separating temporals from eye.
*c*¹ Pupil vertically elliptic.
*d*¹ Anal entire; a preocular.....*Dinodon semicarinatum*,^a p. 366.
*d*² Anal divided; no preocular.....*Dinodon orientale*, p. 372.
*c*² Pupil round.
*d*¹ Scale rows 19.
*e*¹ Ventrals less than 170.
*f*¹ Parietals equaling the distance from eye to tip of snout.
Natrix tigrina, p. 272.
*f*² Parietals longer than distance from eye to tip of snout.
Natrix vibakari, p. 266.
*e*² Ventrals more than 190.....*Elaphe quadrivirgata*, p. 327.
*d*² Scale rows more than 19.
*e*¹ Scale rows 21.....*Elaphe conspicillata*, p. 334.
*e*² Scale rows 23-25.....*Elaphe climacophora*, p. 324.

The examination of the dentition must be made very carefully in order to avoid mistakes. The safest way is probably to dissect out one of the maxillary bones. This can be done very easily by running the point of a sharp knife between the supralabials and the underlying bone, cutting the tissue along the whole length of the latter. By forcing the point of the knife over the upper edge of the bone in the region of the eye the bone can be easily lifted up and the connecting ligaments severed. The adherent tissue may be carefully removed, though in most cases it is sufficient to let it dry. The teeth can now be examined conveniently. Care must be had not to mistake the space left by a lost tooth for a natural interval; if a tooth has fallen out, a distinct pit or depression is left on the alveolar edge of the max-

^a Occurrence in Japan very doubtful.

illa. In counting the teeth the second inner row of loose teeth which are only the reserve teeth must not be taken into consideration. If the specimen is so hardened that it is difficult to open the mouth it should not be forced open by prying, a procedure apt to ruin the teeth and break the lower jaw, but the thick muscle at the corner of the mouth closing the jaws should be cut through on both sides. If properly done the specimen need show no outward sign of mutilation. The maxilla after being dissected out and cleaned should be placed in a small glass tube or vial and, provided with the same number as the snake, kept in the same bottle, or separately together with other preparations of the same kind.

The apical scale pits are usually distinctly visible under a fairly good magnifying glass. In some cases, however, they are rather difficult to discern, thus, for instance, in *Elaphe rufodorsata*. When the pits are not discovered at once the skin should be allowed to dry and then viewed at different angles to the light. The epidermis of



FIGS. 229-231.—HEAD-SHIELDS OF TYPHLOPS. *f*, FRONTAL; *ip*, INTERPARIETAL; *l*, SUPRALABIALS; *n*, NASAL; *o*, OCULAR; *p*, PARIETALS; *pf*, PREFRONTAL; *pro*, PREOCULAR; *pto*, POSTOCULAR; *r*, ROSTRAL; *so*, SUPRAOCULAR.

scales of different parts of the body should be examined. In very doubtful cases it may even become necessary to remove some of the epidermis and examine it under a more powerful lens.^a

Family TYPHLOPIDÆ.

The blind-snakes are small cylindric snakes of even thickness, head and tail not distinct, the latter extremely short and terminating suddenly. The body is covered with smooth scales somewhat resembling fish scales, above and below, and the eye is only dimly visible under the semitransparent head shields. They are very worm-like in appearance and burrow in the ground.

The family is distributed over the warmer portions of both hemispheres. Only one genus is represented in our territory.

^a For the nomenclature of head shields of Typhlopidae snakes, see figs. 229-231 on the present page, and for that of the Natricid snakes, figs. 236-238, on page 262.

Genus *TYPHLOPS*^a Oppel.

1811. *Typhlops* OPPEL, Ord. Rept., p. 54 (type, *T. lumbricalis*).
 1843. *Aspidorhynchus* FITZINGER, Syst. Rept., p. 24 (type, *Typhlops eschrichtii* Schlegel).
 1843. *Pseudotyphlops* FITZINGER, Syst. Rept., p. 24 (type, *Typhlops polygrammicus* Schlegel).
 1843. *Ramphotyphlops* FITZINGER, Syst. Rept., p. 24 (type, *Typhlops multilineatus* Schlegel).
 1844. *Ophthalmidium* DUMÉRIEUX and BIBRON, Erpét. Gén., VI, p. 262 (type, *O. longissimum*).
 1844. *Onychocephalus* DUMÉRIEUX and BIBRON, Erpét. Gén., VI, p. 272 (type, *O. delalandii*).
 1845. *Argyrophis* GRAY, Cat. Liz. Brit. Mus., p. 136 (type, *A. bicolor*).
 1845. *Anilius* GRAY, Cat. Liz. Brit. Mus., p. 135 (type, *A. australis*) (not *Anilius* Oken, 1816).
 1845. *Meditoria* GRAY, Cat. Liz. Brit. Mus., p. 139 (type, *Typhlops nasutus*).
 1845. *Onychophis* GRAY, Cat. Liz. Brit. Mus., p. 132 (type, *O. franklinii*).

The genus *Typhlops*, of which only one species of wide distribution enters our territorial limits in the extreme south, is a large one widely distributed over the tropical and subtropical countries of both the eastern and the western hemisphere. Only a few species, however, occur in America.

TYPHLOPS BRAMINUS^b (Daudin).

AMAGAKU OR MEKU HABU.

1803. *Eryx braminus* DAUDIN, Hist. Nat. Rept., VII, p. 279 (based on Russell's Ind. Serp., I, pl. XLIII; type-locality, Vizagapatam, India).—*Typhlops braminus* CUVIER, Règne Anim., 2 ed., II, 1829 (p. 73).—JAN, Icon. Ophid. (p. 12), livr. 3, 1864, pls. iv and v, fig. 16 c (East Indies).—BOULENGER, Cat. Sn. Brit. Mus., I, 1893, p. 16 (South Asia; Indian Ocean islands; Africa S. of equator).—WALL, Proc. Zool. Soc. London, 1903, p. 85 (Formosa; Hongkong).
 1820. *Tortrix russellii* MERREM, Tent. Syst. Amph., p. 84 (based on Russell).
 1845. *Argyrophis truncatus* GRAY, Cat. Liz. Brit. Mus., p. 138 (type-locality, Philippines; types in Brit. Mus.; Cuming, collector).
 1845. *Argyrophis braminus* GRAY, Cat. Liz. Brit. Mus., p. 138 (*lapsus* for *braminus*).
 1845. *Onychocephalus capensis* SMITH, Illus., Zool. S. Afr., Rep. (pl. LI, fig. 3; pl. LIV, figs. 9-16) (type-locality, Cape of Good Hope; types in Brit. Mus.).
 1860. ? *Ophthalmidium tenue* HALLOWELL, Proc. Phila. Acad., 1860, p. 497 (type-locality, Hongkong, China).
 1863. *Typhlops inconspicuus* JAN. Elenco Sist. Ofid., p. 11 (*nomen nudum*; Madagascar).
 1882. *Typhlops* (*Typhlops*) *euproctus* BOETTGER, Zool. Anz., 1882, p. 479 (type-locality, Nossi-Bé; type in Mus. Senckenberg.; A. Stumpff, collector).
 1891. ? *Typhlops* sp. OKADA, Cat. Vert. Japan. p. 68 (Miyakoshima).


Description (figs. 232-235).—Science College Museum, Tokyo; Okinawa shima, Riukiu; Nishi, collector. Snout rounded, moder-

^a From τῦφλός, blind; ὤψ, eye.

^b Signifying Brahmin, a Hindu priest.

^c See this work, figs. 232-235.

ately projecting; nostrils lateral, the nasal cleft proceeding from the preocular; preocular as large as ocular, in contact with second and third supralabials and with anterior (lower nasal); rostral very narrow, scarcely one-third the width of the head, in contact with prefrontal, not extending as far back as the eyes; anterior nasal extends to the upper surface of the head; posterior (upper) nasals separated by the prefrontal; prefrontal about same size as supraoculars, slightly larger than frontal, all considerably larger than the body scales; 4 supralabials; eyes distinct; 20 scales round the body; tail ending in a spine and shorter than greatest diameter of body which is contained 38 times in total length. Color (in alcohol) dark brown



FIGS. 232-235.—*TYPHLOPS BRAMINUS*. ENLARGED. 232, TOP OF HEAD; 233, SIDE OF HEAD; 234, UNDERSIDE OF HEAD; 235, TAIL. FROM JAN, *ICON. OPHID.*, LIV. 3, PL. V, FIG. 16.

above, each scale being brown with a pale base; snout and tail whitish; lower surface paler, the base of the scales being dark, the terminal two-thirds pale brown.

Dimensions.

	mm.
Total length.....	152
Snout to vent.....	149
Vent to tip of tail.....	3
Greatest diameter of body.....	4

Habitat.—A species of very wide distribution in the tropics of the Old World, rivaling some of the geckos. It is found from South Africa and Madagascar through India and the Malayan archipelago to South China, the Philippines, and Guam^a, extending as far north as Okinawa.

There are specimens from Formosa in the British Museum collected by Swinhoe and Dickson, and in 1894 the museum in Christiania received a number of specimens collected by Mr. Novara in the same island, one of which is now in the United States National Museum.

^a Originally found there by Quoy and Guaynard (Duméril and Bibron, *Erpét. Gén.*, VI, p. 312). Dr. J. C. Thompson, U. S. N., recently captured a specimen on that island according to letter of September 17, 1906.

I have examined three specimens from the Riu Kius in the museums in Tokyo, viz, two in the Science College Museum, from Okinawa shima, and one in the Imperial Museum, Ueno Park, from Yaeyama Island, southern group. There is also a specimen in the Hamburg Museum (No. 1479) collected by Doctor Warburg in the "Liukius."


List of specimens of Typhlops braminus.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Total length.	Length of tail.	Diameter.
							mm	mm	mm
U.S.N.M.	32970	Adult	Formosa		Mr. Novara	20	147	3	3
Sci. Coll., Tokyo		do.	Okinawa shima		Mr. Nishi	20	152	3	4
Do.		Young	do.		Dobutsu Hyohonsha		76	2	2
Imp. Nat., Tokyo		do.	Miyakoshima		Mr. Sagara	81		2	2.5

^a Description, p. 260.

Family NATRICIDÆ.

This family embraces the great bulk of snakes, in fact only somewhat less than two-thirds of all the species of snakes known.



FIGS. 236-238. HEAD-SHIELDS OF A NATRICID SNAKE. *cs*¹, ANTERIOR CHIN-SHIELDS; *cs*², POSTERIOR CHIN-SHIELDS; *f*, FRONTAL; *il*, LOWER LABIALS; *im*, INTERNASALS; *l*, LOREAL; *lb*, SUPRALABIALS; *m*, MENTAL; *n*¹, ANTERIOR NASAL; *n*², POSTERIOR NASAL; *p*, PARIETAL; *pf*, PREFRONTALS; *pro*, PREOCULAR; *pro*, POSTOCULAR; *r*, ROSTRAL; *so*, SUPRAOCULAR; *t*, TEMPORALS; *v*, VENTRALS.

In this vast horde of species, considerably over one thousand, there is a great uniformity of structure, making it difficult to formulate easily recognizable groups. Our knowledge of the phylogenetic relationships of the various forms has been established only recently, thanks

to the studies of Cope and Boulenger, though much is to be learned yet before a natural classification can be expected.

This family is usually known as the *Colubridæ*, but, as will be shown further on, the generic name *Coluber* belongs to the typical genus of the so-called *Viperidæ*.

The distinctive characters of the four subfamilies recognized among the natricid snakes occurring in Japan and adjacent territory have been tabulated as follows:

- α^1 Hypapophyses present throughout the vertebral column (figs. 223-225).
 b^1 All maxillary teeth solid.....NATRICINÆ, p. 263.
 b^2 Posterior maxillary teeth grooved.....HOMALOPSINÆ, p. 299.
 α^2 Hypapophyses absent in posterior dorsal vertebræ (figs. 226-228).
 b^1 All maxillary teeth solid.....CORONELLINÆ, p. 306.
 b^2 Posterior maxillary teeth grooved.....BOIGINÆ, p. 381.

Boulenger has shown that the question whether the hæmal processes are present on the posterior vertebræ or not can be easily ascertained by making an incision along the belly in the posterior fifth of the body, pushing aside the viscera, and disarticulating the backbone by bending the body dorsally. The two styles of vertebræ, with or without the lower median process, are shown in the accompanying figures. It should be remarked, however, that recently considerable doubt has been thrown on this character as to whether it is of the fundamental importance usually attributed to it, or not.

Subfamily NATRICINÆ.

The Natricinæ are without grooved teeth in the posterior part of the upper jaw.

Our species, although sometimes with the nostrils placed rather high, never have them on top of the snout, and the internasal shields, though often small and narrowed anteriorly, are in pairs and join the rostral. The scales are strongly keeled.

Only two genera are known from our territory, which may be distinguished as follows:

- α^1 Subcaudals divided; postoculars well developed, separating temporals from orbit.
Natrix, p. 263.
 α^2 Subcaudals single; postoculars rudimentary, the temporals apparently entering the orbit.....*Achalinus*, p. 294.

Genus NATRIX ^a Laurenti.

1768. *Natrix* LAURENTI, Syn. Rept., p. 73 (type, *N. vulgaris* = *Coluber natrix*).
 1822. *Tropinotus* KÜHL, Isis, 1822, p. 473 (*nomen nudum*).
 1826. *Tropidonotus* H. BOIE, Isis, 1826, p. 205 (type, *T. natrix*).
 1843. *Rhabdophis* FITZINGER, Syst. Rept., p. 27 (type, *T. subminiatus*, Schlegel).
 1843. *Steiropphis* FITZINGER, Syst. Rept., p. 27 (type, *T. chrysargus* Kuhl).
 1852. *Hydrophilophis* SCHMIDT, Abh. Naturw. Ver. Hamburg, II, Pt. 2 (substitute for *Tropidonotus*).

^a Signifying a water snake; literally, a swimmer.

1853. *Nerodia* BAIRD and GIRARD, Serp. N. Am., p. 38 (type, *N. sipedon*).
 1853. *Regina* BAIRD and GIRARD, Serp. N. Am., p. 45 (type, *R. leberis*).
 1854. *Amphisma* DUMÉRIL and BIBRON, Erpét. Gén., VII, p. 724 (type, *A. stolatum*).
 1886. *Bothrodytes* COPE, Proc. Amer. Philos. Soc., XXIII, 1886 (p. 495) (type, *Amphisma subminiatum*).
 1893. *Ceratophallus* COPE, Amer. Natural., XXVII, May, 1893, p. 483 (type, *Natrix vittata*).
 1893. *Diplophallus* COPE, Amer. Natural., XXVII, May, 1893, p. 483 (type, *N. piscator*).

This genus is quite commonly known as *Tropidonotus* Kuhl, but the above synonymy shows how poor the status of that name is. *Natrix* was established by Laurenti as early as 1768, and no difficulty is encountered as to the type, since Fleming, in 1822, before the name *Tropidonotus* was invented, designated *N. torquatus* (*Coluber natrix* Linnæus, or *Natrix vulgaris* Laurenti) specifically as the type.

The genus *Natrix*, as here understood, has been variously subdivided into genera based upon the syncranterian or diacranterian character of the dentition, the presence, or absence of scale pores, and, lately, upon the structure of the intromittent organ, whether simple or bifurcate, and whether provided with a papilla at the apex or not. These structural characters are only known in a few of the numerous species contained in the present genus, and any rational attempt at subdivision must be postponed until they are investigated in all the species.

That the dentition in itself alone is not a satisfactory guide to the proper segregation of the species into natural groups is shown by *Natrix vibakari*. By Boulenger this species is retained in the first section of the genus, which he restricts as *Tropidonotus* proper, because having the hinder maxillary teeth "gradually enlarged," against the section *Amphiesma* having the "last two or three abruptly enlarged." I have dissected out the maxilla of three specimens of this species and would designate the last two teeth as "abruptly enlarged" in every case. It is therefore impracticable to base a primary subdivision of the group upon this character.

On the other hand, the scale pores have only been recorded in the case of very few species. Thus I can find no record as to whether *Natrix swinhonis* possesses them or not, and inclined as I am to regard this character as indicating natural genera among these snakes, I am unable to employ it at present.

This leads to an inquiry into the closer relationship of the Japanese species of this complex genus. As will be shown further on, *Natrix tigrinus* is only subspecifically distinct from the mainland form *N. lateralis*, which extends westward to the confines of Tibet. *Natrix vibakari* belongs to another group of species which also extend over China to the eastern Himalayas. It is not difficult to point out *Natrix khasiensis*, from Assam, and *Natrix craspedogaster*, from

the province of Fokien, opposite Formosa, as the nearest relatives known. Bearing in mind that the relative size of the posterior maxillary teeth is of little practical importance, I think we may also safely add to this group *Natrix chrysarga*, a species which extends from the eastern Himalayas to southern China, the Indo-China peninsula, and the Malay Archipelago. As will be shown further on, *Natrix pryeri* is a specialized member of this group. From what we now know of the distribution and relationship of the other reptiles in these regions, it is pretty safe to predict that a *Natrix* of the same group, either *N. craspedogaster*, or *N. chrysarga*, or a form intermediate between one of these and *N. pryeri*, will be found to occur in Formosa when the herpetological fauna of this island shall have become thoroughly known.^a

The other species enter our territory only in Formosa and as such their Himalayo-Chinese relationships are easily traced. Both *N. stolata* and *N. piscator* occur from the Himalayas and India to southern China, Indo-China, and the Malayan Islands. *N. annularis* also inhabits the Yangtse drainage and *N. swinhonis* has its nearest relative, *N. nuchalis*, located on the upper Yangtse Kiang.

KEY TO THE JAPANESE, KOREAN, AND FORMOSAN SPECIES OF NATRIX.

*a*¹ Scale rows, 19; supralabials, 7, 8, or 9.

*b*¹ Subcaudals less than 105 pairs.

*c*¹ Suture between rostral and internasal not shorter than between rostral and first supralabial (fig. 241).

*d*¹ Parietals longer than distance from eye to tip of snout . . . *N. vibakari*, p. 266.

*d*² Parietals equaling the distance from eye to tip of snout.

*e*¹ Ventrals plus subcaudals more than 226; subcaudals more than 65.

N. tigrina, p. 272.

*e*² Ventrals plus subcaudals less than 226; subcaudals less than 65.

N. tigrina lateralis, p. 278.

a NATRIX CHRYSARGA (Schlegel).

1837. *Tropidonotus chrysargos* SCHLEGEL, Phys. Serp., II, p. 312 (type-locality, Java; types in Leiden Mus.; Kuhl and Reinwardt, collectors).

1847. *Tropidonotus junceus* CANTOR, Cat. Mal. Rept. (p. 93) (type-locality, Pinang; type in Brit. Mus.; Cantor, collector).

1868. *Amphiesma platyceps* JAN, Icon. Ophid., livr. 29, pl. II, fig. 1 (type-locality, Sikkim; type in Milano Mus.) (not *Tropidonotus platyceps* Blyth, 1854).

Scale rows, 19, all strongly keeled; 9 or 8 supralabials; 60-93 pairs of subcaudals; suture between rostral and internasal not shorter than between rostral and first supralabial; parietal longer than distance from eye to tip of snout; 2 anterior temporals.

NATRIX CRASPEDOGASTER (Boulenger).

1899. *Tropidonotus craspedogaster* BOULENGER, Proc. Zool. Soc. London, 1899, p. 163, pl. XVII, fig. 1 (type-locality, Kuantun, prov. Fokien, China; types in Brit. Mus.; La Touche, collector).

Scale rows, 19, outer faintly keeled; 8 supralabials; 87-97 pairs of subcaudals; anterior temporals 1 or 2 (from original description).

- c*² Suture between rostral and internasal considerably shorter than between rostral and first supralabial (fig. 255).
- d*¹ One outer scale row smooth, or none.
- e*¹ Second row of temporals consisting of 1 or 2 scales; 3 supralabials entering eye..... *N. stolata*, p. 280.
- e*² Second row of temporals consisting of 3 scales; normally only 1 supralabial (rarely 2) entering eye..... *N. annularis*, p. 291.
- d*² Two or three outer scale rows smooth..... *N. piscator*, p. 288.
- b*² Subcaudals more than 105 pairs..... *N. pryleri*, p. 284.
- a*² Scale rows, 15; supralabials, 6..... *N. swinhonis*, p. 293.

NATRIX VIBAKARI^a (Boie).

HIBAKARI.


1826. *Tropidorotus vibakari* BOIE, Isis, 1826, p. 207 (type-locality, Japan; Blomhoff, collector); Bijdr. Natuurk. Wetensch., II, Pt. 1, 1827, p. 254.—TEMMINCK and SCHLEGEL, Fauna Jap., Rept., 1837, p. 87, Ophid., pl. v (Japan, Decima).—SCHLEGEL, Phys. Serp., II, 1837, p. 316.—DUMÉRIL and BIBRON, Erpét. Gén., VII, Pt. 1, 1854, p. 595 (Japan).—BLEEKER, Natuurk. Tijdschr. Nederland. Indië, XVI, 1858, p. 204 (Japan).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1866, p. 111 (Yokohama), 1876, p. 379.—STRAUCH, Mém. Acad. Sci. St. Pétersb. (7), XXI, No. 4, 1873, pp. 174, 277 (Possiet Bay; Baranovski Post, Suifun River, Amurland).—BOULENGER, Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 139 (Khabarovka, Ussuri); Cat. Snakes Brit. Mus., I, 1893, p. 221 (Ussuri; Yokohama; Kobe; Formosa).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 21 (Hondo).—WALL, Proc. Zool. Soc. London, 1903, p. 99 (Nagasaki; Mt. Fuji); 1905, II, p. 513 (Hondo).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, No. 1, 1905, p. 214 (Possiet Bay).
1827. *Tropidonotus vibikari* F. BOIE, Isis, 1827, p. 535 (err. typ).
1876. *Tropidonotus martensii* HILGENDORF, Mitth. Deutsch. Ges. Ost-Asiens, I, heft 10, pp. 30, 31 (substitute name) (Yokohama); Sitz Ber. Ges. Naturf. Fr. Berlin, 1880, p. 115.—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, 1891, p. 239 (not in Yezo).

Doctor Hilgendorf changed the well-known specific name of this species into *T. martensii* because the name Hibakari by its significance, as indicating a deadly snake, is undesirable. This change, of course, is wholly inadmissible under all recognized nomenclatural codes. The principle involved would cause endless confusion and set the law of priority at naught.

Description.—*Adult male*; U.S.N.M. No. 23335; Yokohama; September, 1896; L. Stejneger, collector (figs. 239–241). Rostral much broader than high, just visible from above; internasals rather large, truncate anteriorly, their suture with the rostral longer than half their width and longer than suture between rostral and first supralabial; prefrontals considerably larger than internasals, broadly in contact with supraocular; frontal slightly longer than its distance

^a *Vibakari*, or more correctly *hibakari*, is the name given by the Japanese to this snake, under the impression that it is so poisonous that its victim will live only through the day. Needless to say, it is perfectly harmless.

from tip of snout, nearly equaling the interparietal suture; parietals as long as frontal and prefrontals together, much longer than distance from eye to tip of snout; nostril lateral, between two subequal nasals; loreal somewhat longer than high, its upper edge much shorter than the lower; one preocular, widely separated from frontal; three postoculars; temporals 1 + 2 (normal) on right side, the two of the second row being fused on the left side; 7 supralabials, third and fourth entering eye, sixth largest (normal), on right side, 8 on left side, an additional labial being interpolated in front of the eye; 4 lower labials in contact with anterior chin-shields, which are shorter than the posterior; 19 rows of scales, outer row smooth, the others distinctly keeled, with obscure apical pits best observable on the lateral scales; 146 ventrals; anal divided; 73 pairs of subcaudals. Color (in alcohol) uniform smoke-gray above, top of head darker, also darker along the middle of the back; upper lip whitish, the labials posteriorly edged with blackish; a black postocular band descend-



FIGS. 239-241.—*NATRIX VIBAKARI*. 2 × NAT. SIZE. 239, TOP OF HEAD; 240, SIDE OF HEAD; (3] × NAT. SIZE). 241, ROSTRAL FROM FRONT. NO. 2335, U.S.N.M.

ing on the upper half of the penultimate supralabial and crossing obliquely the suture between it and the last supralabial to the angle of the mouth, cutting off the latter labial from the others and associating it with a whitish postoral band which curves slightly upward from it toward the nape without crossing it; a few blackish spots along the edges of the postoral band; on the middle line of the nape, from the parietal suture a short distance backward a narrow whitish dark-edged line; two pale dots on the parietals near the middle of the interparietal suture; underside uniform whitish, with a series of small dusky spots—or rather, dashes—one near the outer end of each ventral and subcaudal.

Dimensions.

	<i>mm.</i>
Total length.....	426
Snout to vent.....	308
Vent to tip of tail.....	118

This is a small snake, which probably seldom exceeds 600 mm. in total length.

Life colors, according to Dr. H. M. Smith, "light brown above; belly cream color."

The *young*, as shown by a specimen in our collection from Shikoku (U.S.N.M. No. 31850) only 155 mm. in total length, is essentially as the adult; the ground color, however, is darker and there are two distinct series of numerous roundish, pale, dark-edged dots, located on the fifth and sixth scale-rows on each side.

Variation.—A noteworthy variation in this species is the occasional joining of the parietal to one of the supralabials, noteworthy particularly because it was the condition found in Boie's type. In two of our specimens (No. 30730, from Kiusiu, and No. 34028, from Mount Fuji) the contact is between the parietal and the sixth supralabial behind the first temporal, and in another specimen (No. 34029) there is such a contact on one side. The number of preoculars varies to some extent, but one is the normal number, not a single specimen of the 22 before me having more. Strauch, on the other hand, found two out of four specimens. Altogether, his series was extraordinarily variable, as their postoculars varied between two, three, and two and three, while the supralabials showed the unusual variation of coalescing to the extent of being reduced to six, or even five. We have 12 with three postoculars, 4 with two and 3, and 3 with two, while all have 7 supralabials except three, which have 8 on one side, and one (No. 34034a) which has 8 on both sides. This additional labial is interpolated in front of the eye, and in one of the specimens the interpolation occurs on both sides, but the normal number of 7 is restored on one side by the coalescing of two labials behind the eye.^a

Remarks.—The relationship of this species to *Natrix khasiensis*, from Assam, and *N. craspedogaster*, from Fokien, has been alluded to above (p. 264). It is not only shown in the unmistakable pattern of the upper lip and its continuation in the nuchal crescent, but also in the postparietal median light streak and in the series of white spots on the body of *N. craspedogaster* and the young *N. vibakari* described above (p. 268). About the relationships with *N. pryeri*, see under the latter species (p. 286).

Habitat.—Common throughout Japan from Kiusiu to Yezo. Doctor Nozawa has informed me that it occurs in the latter island, but he has sent no specimen as yet. Numerous specimens from the neighborhood of Tokyo, Yokohama, and Nagasaki are in the various museums, as well as some from Kobe, Miyazaki, and other places.

Outside of Japan it has been found in the Russian Coast Province from Possiet Bay, on the Korean frontier (Mus. St. Petersburg), to Vladivostok (Hamburg Mus. No. 2170) and Khabarovka, at the junction of the Ussuri with the Amur (Brit. Mus.). In 1885 Jouy

^a For further variations see Wall, Proc. Zool. Soc. London, 1905, II, p. 513.

obtained a specimen at Fusan, southern extremity of Korea (U. S. Nat. Mus. No. 17510), the first record from that country.

A specimen in British Museum is said to have been collected by Swinhoe in Formosa. It has the extraordinary low number of 127 ventrals, and altogether either its identity or its locality seems open to doubt.^a

It has not been recorded from the Riu Kius.

^aSee also Wall, Proc. Zool. Soc. London, 1905, II, p. 513.

List of specimens of *Natrix vibakari*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Occipitals.	Temporals.	Supra-labials.
U.S.N.M.	11327	Female	Japan	1878	E. S. Morse	19	148	2	71	1,3	1+1	7
Do.	11328	Male	do.	1878	do.	19	151	2	73	1,2-3	1+2	7
Do.	11329	do.	do.	1878	do.	19	146	2	70	1,3	1+1-2	7
Do.	11330	Young	do.	1878	do.	19	147	2	73	1,3	1+2	7
Do.	11331	Female	do.	1878	do.	19	146	2	63	1,3	1+1	7
Do.	17510	do.	Fusan, Korea	1885	P. L. Jouy	19	148	2	58	1,2	1+1	7
Do.	23335	Male a	Yokohama	Sept. —, 1886	L. Stejneger	19	146	2	73	1,3	1+1-2	7-8
Do.	30730	Female	Miyazaki, Kinsiu		C. A. Clark	19	146	2	72	1,3	1+2	7
Do.	31850	Young	Suzaki, Toss, Shikoku	May 8, 1903	Dr. H. M. Smith	19	148	2	79	1,3	2+2	8
Do.	31858	Male	Kochi, Shikoku	May 11, 1903	do.	19	147	2	83	1,3-4	1+2	7-8
Do.	31859	Young	do.	do.	do.	19	141	2	74	1,3	1+2	7
Do.	34028	Female	Mount Fuji	Aug. —, 1888	A. Owston	19	147	2	74	1,2-3	1+2	7
Do.	34029	do.	do.	do.	do.	19	149	2	74	1,3	1+1	7-8
Do.	34030	do.	do.	do.	do.	19	150	2	75	1,3	1+2	7
Do.	34031	do.	do.	do.	do.	19	145	2		1,2-3	1+2	7
Do.	34032	Male	do.	do.	do.	19	148	2	79	1,2	1+1	7
Do.	34033	do.	do.	do.	do.	19	147	2	80	1,2-3	1+2	7
Do.	34034	Young	do.	do.	do.	19	145	2	78	1,3	1+1	7
Do.	34034a	do.	do.	do.	do.	19	145	2	78	1,3	1-2+1-2	8
Do.	36543	Female b	Tokyo	do.	Sci. Coll. Tokyo	19	147	2		1,3	1+2	7
Do.	36544	Male b	do.	do.	do.	19	151	2	70	1,2	1+2	7-8
Do.	36545	do. b	do.	do.	do.	19	150	2	74	1,3	1+1	7
Do.	36546	Young b	do.	do.	do.	19	150	2	74	1,3	1+1	7
Do.	c	do. c	Japan	do.	A. Adams	19	137	2	62			
Do.	d	do. d	do.	do.	do.	19	137	2	65			
Do.	e	Male e	do.	do.	Museum, Leiden	19	148	2	74			
Do.	f	Female e	do.	do.	do.	19	144	2	79			
Do.	h	Male e	do.	do.	Doctor Bleeker	19	148	2				
Do.	i	Female e	Yokohama	do.	Challenger Exped	19	144	2	68			

Brit. Mus.....	k	Young c.....	Kola.....	Challenger Exped.....	19	145	2	73
Do.....	a	Male c.....	Khabarovka, Ussuri.....	Mr. Doerries.....	19	150	2	59
Do.....	b	do. s.....	do.....	do.....	19	151	2	60
St. Petersb.....	1835	(d)	Japan.....	B. Dybowski.....	19	147	2	76
Do.....	2984	(d)	Possiet Bay, Coast Province.....	do.....	19	146	2	64
Do.....	2985	(d)	do.....	do.....	19	148	2	62

d Strauch, p. 277.

c Boulenger, Cat. I, p. 222.

b Sci. Coll. No. 67.

a Description p. 286; figs. 230-241.

NATRIX TIGRINA^a (Boie).

YAMA GAMISHI.

Plate XIX.


1826. *Tropidonotus tigrinus* BOIE, Isis, 1826, p. 205 (type - locality, Japan; Blomhoff, collector); Bijdr. Natuurk. Wetensch., II, Pt. 1, 1827, p. 250.—TEM-MINCK and SCHLEGEL, Fauna Japon., Rept., 1837, pp. 85, 139; Ophid., pl. iv^b (Japan).—SCHLEGEL, Phys. Serp., II, 1837, p. 315 (Japan).—GUENTHER, Cat. Colubr. Sn. Brit. Mus., 1858, p. 71 (part: Japan); Rept. Brit. India, 1864, p. 271 (part: Japan).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1866, p. 110 (Yokohama); 1876, p. 379 (Tokyo, Nagasaki).—STRAUCH, Mém. Acad. Sci. St. Pétersb. (7), XXI, no. 4, 1873, pp. 176, 277 (part: Japan).—HILGENDORF, Mitth. Deutsch. Ges. Ost-Asiens, I, heft 10, 1876, p. 30 (Tokyo); Sitz. Ber. Naturf. Fr. Berlin, 1880, p. 115 (Tokyo).—BOULENGER, Cat. Snakes Brit. Mus., I, 1893, p. 249 (part: Kobe), III, 1896, p. 607 (Shimabara).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V., 1891, p. 239 (not in Yezo).—OKADA, Cat. Vert. Jap., 1891, p. 68 (Tokyo; Nikko; Osaka; Suwo; Awachi; etc.).—BOETTGER, Kat. Schl. Mus. Senckenberg, 1898, p. 25 (part: Nikko; Yokohama; Yezo).—WALL, Proc. Zool. Soc. London, 1903, p. 99 (Japan); 1905, II, p. 513 (Yaku I., Tanega I.).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 222 (Nagasaki; Unzen).—*Amphipsma tigrinum* DUMÉRIL and BIBRON, Erpét. Gén., VII, Pt. 1, 1854, p. 732 (Japan).—BLEEKER, Natuurk. Tijdschr. Nederland. Indië, XVI, 1858, p. 204 (Japan).—HALLOWELL, Proc. Phila. Acad., 1860, p. 499 ([Simoda] Hondo).

Description (figs. 242-243).—*Adult male*; U.S.N.M. No. 31846; Sendai, province of Satsuma, Kiusiu; June 4, 1903; Dr. H. M. Smith, collector. Rostral much broader than high, well visible from above; internasals somewhat smaller than prefrontals, truncate in front, the suture with rostral slightly longer than the suture of rostral with first supralabial; prefrontals in contact with supraoculars; frontal much broader than supraoculars, about as long as its distance from tip of snout and as interparietal suture; parietals about as long as distance of eye from tip of snout; nostril lateral, large, between two nasals, of which the posterior is higher than the anterior; loreal pentagonal, as high as long; two preoculars, the upper not in contact with frontal; three postoculars; temporals 1 + 2, those of the second row being abnormally fused on the left side; 7 supralabials, fifth and sixth largest, fourth and fifth entering eye; 5 lower labials in contact with anterior chin-shields, which are considerably shorter than posterior; 19 rows of long, narrow, strongly keeled scales, each with two large oval apical pits same distance from the tip; 161 ventrals; anal divided; 83 pairs of subcaudals. Color (in alcohol)^c above tawny-

^a Signifying tiger-like, probably with reference to the red and black spots on the sides. According to von Siebold this species is in Japan also called *Torano Kuchinaha*, and according to von Martens, *Torasuhebi*, both meaning tiger snake.

^b Reproduced in this work on Plate XIX.


^c In life, according to Dr. H. M. Smith, these snakes are "greenish above, with black and orange blotches intermingled; head greenish above; belly slate color."



NATRIX TIGRINA. (FROM TEMMINCK AND SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

olivè (light bluish gray where the epidermis has come off), with broad black crossbars on the anterior part of the back, soon breaking up into four series of large squarish and alternating black spots; the outlines of the dark and light colors is very ragged and the keels light colored, even in the black patches as if the light color pattern had been produced by scraping off the black color with a knife from an uneven surface; temporal region dark, with a black line crossing obliquely from upper postocular across anterior temporal, upper portion of sixth supralabial, and middle of seventh to angle of mouth; supralabials yellowish, narrowly edged with black along the posterior suture, the one below the eye on the fourth being heaviest; throat yellowish, a broad band of similar color curving up behind the temporal region so as to form a collar; lower neck yellowish, the under-



FIGS. 242-243.—*NATRIX TIGRINA*. $1\frac{1}{2} \times$ NAT. SIZE. 242, TOP OF HEAD; 243, SIDE OF HEAD. No. 7385 U.S.N.M.

side gradually becoming darker posteriorly, each ventral being marked basally with an increasing amount of dark slate color until the under-side posteriorly is nearly uniform slaty black.

Dimensions.

	mm.
Total length.....	682
Snout to vent.....	507
Vent to tip of tail.....	175

The *young* do not differ materially from the adults.

Our largest specimen (No. 12711) measures about 935 mm. in total length, and probably few grow to be more than 1 meter in length.

Variation.—The scale formula of this species is fairly constant. All the specimens examined and on record have 19 scale rows, a divided anal (except No. 33988), and seven supralabials (two of our specimens have only six on one side). Hilgendorf in a series of 40 found only one in which the two preoculars were fused into one, and all

the 53 specimens examined by me for this character have two, except one, in which they are fused on one side, and two which have three preoculars on one side.^a The temporals are almost always 1 + 2, though one of our specimens (No. 7386a) has 2 + 2 on one side.^b On the other hand, the postoculars vary between 3 and 4 and sometimes only 2; in our large series there are only three specimens which have 4 postoculars on both sides, and eight which have 4 on one side, but Hilgendorf^c found 3 and 4 in about even proportion, viz, 23 with 3 and 28 with 4 on both sides, and 7 with 3 and 4 unsymmetrically (unless his statement involves some misprint, for instance, 8 instead of 28).

The coloration and markings, however, are very variable, especially the size and arrangement of the black spots and the amount of black on the underside. In the latter respect specimens vary without regard to locality between almost solid black and uniformly greenish white. The upper surface is often so dark that the dorsal blotches can be made out only with difficulty, but in most cases the orange-colored spots occupying the second, third, and fourth scale rows on each side are very conspicuous. The orange color is confined to the base and lateral margins of the scales, the median portion and tip being blackish, forming linear spots within the bright area.

The few Korean specimens examined by me have the dorsal blotches less differentiated and scarcely alternating with the lateral ones, thus forming more or less continuous crossbars and approaching what appears to be the common coloration of northern specimens of *Natrix lateralis*.

Habitat.—This species, as here restricted, is confined to Japan and southern Korea, being represented in the coast regions of the mainland by a closely allied form described hereafter.

The only definite locality of this species in Korea is Fusan, whence we have a specimen collected by Jouy. Two other specimens from Korea have also been examined by me, but their exact locality is uncertain, though probably either Fusan or Seoul.

The tiger snake seems to be common in Kiusiu and Hondo. Blomhoff, Siebold, and Buerger collected numerous specimens in the neighborhood of Nagasaki; von Martens obtained it for the Berlin Museum in Nagasaki, Yokohama, and Tokyo; British Museum has specimens from Kobe collected by the *Challenger* expedition and from

^a One specimen (No. 33994) has the lower preocular on both sides fused with the loreal which thus enters the eye. Dr. Wall has examined three specimens with the loreal confluent with the postnasal on both sides, and one similarly abnormal on one side.

^b No. 33999 has a small abnormal scale-like shield separated off from the upper anterior edge of the first temporal and not reaching the second.

^c Sitz. Ber. Naturf. Fr. Berlin, 1880, p. 115.

Shimabara by Holst. Museum Senckenbergianum also has a specimen from Nikko, and finally the U. S. National Museum possesses specimens from Satsuma and Miyazaki, Kiusiu, as well as from Osaka, Oide, Simoda, Suruga, Mori, Yokohama, and Tokyo. Dr. H. M. Smith collected numerous specimens in Shikoku. Dr. Wall has recently recorded it from the islands of Yaku and Tanega, south of Kiusiu.

Its occurrence in Yezo is doubted by Fritze whose collectors failed to obtain it in that island, and Nozawa does not include it in his list of Yezo reptiles communicated to me, so there is apparently no specimen in the Sapporo Museum. Boettger, however, records two adult specimens which the Senckenberg Museum obtained from Schmacker in 1889 as having been collected in Yezo. I must here recall the fact, however, that specimens of *Rana japonica* and *Eumeces latiscutatus* (see pp. 111 and 199) from the same locality and source are not above suspicion. Altogether Schmacker's Yezo records require confirmation.

It is not recorded from the Riu Kiu Islands. There is in the Imperial Museum at Ueno Park, Tokyo, a specimen (No. 20) which is said to have been collected at Naha, Okinawa shima. Beyond a somewhat unusual low number of ventrals plus subcaudals this specimen is exactly like Tokyo specimens, and most likely some error has occurred in labeling.

List of specimens of *Natrix tigrina*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Subcaudals and subcaudals.	Ventrals and subcaudals.	Oculars.	Temporals.	Supralabials.
U.S.N.M.	7385	Female ^a .	Simoda, Japan.	Doctor Morrow.	19	162	73	235	2,3	1+2	7
Do.	7386 ^a	do.	do.	May —, 1885	W. Stimpson.	19	166	74	240	2,3	1-2+2	7
Do.	7386 ^b	do.	do.	do.	do.	19	163	72	235	2,3	1+2	7
Do.	12711	do ^b .	Yokohama, Japan.	Apr. —, 1881	P. L. Jouy.	19	161	69	230	2,3	1+2	7
Do.	12712	Halfgr.	do.	do.	do.	19	160	2,3	1+2	7
Do.	12713	Young.	do.	do.	do.	19	160	73	233	2,3	1+2	7
Do.	14147	Female.	Japan.	19	165	67	232	2,4	1+2	7
Do.	15420	Male.	Olde, Shinshu, Japan.	Oct. —, 1882	19	156	72	228	2,3	1+2	7
Do.	31846	do ^c .	Sendai, Satsuma, Kiusiu.	June 4, 1903	H. M. Smith.	19	161	83	244	2,3	1+1-2	7
Do.	31847	Halfgr.	do.	June —, 1903	do.	19	169	75	244	2,3	1+2	7
Do.	31848	Female.	Suzaki, Tosa, Shikoku.	May 8, 1903	do.	19	159	73	232	2,3	1+2	7
Do.	31849	Halfgr.	do.	do.	do.	19	157	79	236	1-2,3	1+2	7
Do.	31853	do.	Kochi, Tosa, Shikoku.	May 11, 1903	do.	19	159	78	237	2,3	1+2	7
Do.	31854	do.	do.	do.	do.	19	158	85	243	2,3	1+2	7
Do.	31855	do.	do.	do.	do.	19	157	70	227	2,3	1+2	7
Do.	31856	Male.	do.	do.	do.	19	153	2,3	1+2	7
Do.	31857	do.	do.	do.	do.	19	157	72	229	2,3	1+2	7
Do.	31860	Female.	do.	May —, 1903	do.	19	159	76	235	2,3-4	1+2	7
Do.	31861	do.	do.	do.	do.	19	168	90	248	2,2-3	1+2	7
Do.	31862	do.	do.	do.	do.	19	160	72	232	2,3	1+2	7
Do.	31863	do.	do.	do.	do.	19	160	70	230	2,3	1+2	7
Do.	31864	do.	do.	do.	do.	19	162	73	235	2,3	1+2	7
Do.	31865	Halfgr.	do.	do.	do.	19	158	69	227	2,3	1+2	7
Do.	31868	Male.	do.	do.	do.	19	162	75	237	2,3	1+2	7
Do.	31869	Female.	do.	do.	do.	19	166	74	240	2,3	1+2	7
Do.	31870	Young.	do.	do.	do.	19	157	82	239	2,3	1+2	7
Do.	31871	do.	do.	do.	do.	19	157	80	237	2,3	1+2	7
Do.	33682	Female.	Mount Fuji, Hondo.	July —, 1868	A. Owston.	19	164	71	235	2,3	2+2	7
Do.	33683	Male.	do.	do.	do.	19	160	74	234	2,2-3	1+2	7
Do.	33684	Female.	do.	do.	do.	19	165	73	238	2,3	1+2	7

Do.....	do.....	do.....	do.....	do.....	19	162	71	233	2,3-4	1+2	7
Do.....	Male.....	do.....	do.....	do.....	19	162	80	242	2,3-4	1+1-2	7
Do.....	Female.....	do.....	do.....	do.....	19	163	70	233	2,3-4	1+2	7
Do.....	Male <i>d</i>	do.....	do.....	do.....	19	162	76	238	2,3	1+2	6-7
Do.....	Young.....	do.....	do.....	do.....	19	157	76	233	2,4	1+2	7
Do.....	Female.....	do.....	do.....	do.....	19	156	72	228	2,3	1+2	7
Do.....	do.....	do.....	do.....	do.....	19	164	2,3	1+2	7
Do.....	do.....	do.....	do.....	do.....	19	161	2,3	1+2	7
Do.....	Male.....	do.....	do.....	do.....	19	155	74	229	2,3-4	1+2-3	6-7
Do.....	Female <i>e</i>	do.....	do.....	do.....	19	168	71	239	2,3-4	1+2	7
Do.....	Male.....	do.....	do.....	do.....	19	159	76	235	2-3,3	1+2	7
Do.....	do.....	do.....	do.....	do.....	19	162	2,3-4	1+2	7
Do.....	do.....	do.....	do.....	do.....	19	159	82	241	2,3	1+2	7
Do.....	Female.....	do.....	do.....	do.....	19	168	69	237	2,3-4	1+2	7
Do.....	do. <i>f</i>	Mori, near Yokohama.....	do.....	do.....	19	161	66	227	2,3	1+2	7
Do.....	do.....	do.....	do.....	do.....	19	164	69	233	2-3,3	1+2	7
Do.....	Male.....	Suruga, Hondo.....	do.....	do.....	19	160	68	228	2,3	1+2	7
Do.....	Female.....	Tokyo, Hondo.....	do.....	do.....	19	162	70	232	2,4	1+2	7
Brit. Mus.....	<i>a</i> Male <i>g</i>	Japan.....	Sci. Coll. Tokyo.....	do.....	19	162	78	240
Do.....	<i>b</i>	do.....	do.....	do.....	19	159
Do.....	<i>c</i>	do.....	do.....	do.....	19	158	84	242
Do.....	Female <i>g</i>	do.....	do.....	do.....	19	163	74	237
Do.....	<i>e</i>	do.....	do.....	do.....	19	164	71	235
Do.....	Male <i>g</i>	Kobe.....	Challenger Exped.....	do.....	19	160	76	236
Do.....	Female <i>g</i>	do.....	do.....	do.....	19	163	80	243
Do.....	<i>h</i>	do.....	do.....	do.....	19	166	72	238
Do.....	<i>v</i>	Shimabara, Hizen, Kinsiu.....	do.....	do.....	19	166	72	238
St. Petersb.....	(<i>t</i>).....	Japan.....	P. A. Holst.....	do.....	19	161	71	232
Do.....	(<i>t</i>).....	do.....	Leiden Mus.....	do.....	19	164	83	247	2,3
Do.....	(<i>t</i>).....	do.....	do.....	do.....	19	164	71	235	2,3
Do.....	Male.....	Korea.....	Dr. N. M. Ferebu.....	do.....	19	161	73	234	2,3	1+2	7
Do.....	do.....	Fusan, Korea.....	P. L. Jony.....	1885	19	163	67	230	2,3-4	1+2	7
Mich. Univ.....	do.....	Korea.....	G. A. Ollinger.....	19	165	2,2-3	1+2	7

a Figs. 242-243. *b* P. 273. *c* Description p. 272. *d* Anal undivided. *e* Loreal fused with preocular. *f* P. 274. *g* Boulenger Cat. I, p. 250. *h* Boulenger, Cat. III, p. 607. *i* Strauch, p. 277.

NATRIX TIGRINA LATERALIS ^a (Berthold).

1858. *Tropidonotus tigrinus* GUENTHER, Cat. Colubr. Sn. Brit. Mus., p. 71 (part: Ningpo, China); Rept. Brit. India, 1864, p. 271 (part: northern China); Ann. Mus. Zool. St. Pétersbourg, I, 1896, p. 205 (Huihsien, prov. Kansu, China).—STRAUCH, Mém. Acad. St. Pétersbourg (7), XXI, no. 4, 1873, p. 176 (part: Strelok, Bay of Peter the Great, Amurland).—GIGLIOLI and SALVADORI, Proc. Zool. Soc. London, 1887, p. 594 (Olga Bay; Gensan).—BOULENGER, Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 139 (Gensan, Korea); Cat. Snakes Brit. Mus., I, 1893, p. 249 (part: Gensan; Chefu; Peking; Ningpo; Kiukiang; Hainan); III, 1896, p. 607 (part: Chefu); Proc. Zool. Soc. London, 1899, p. 164 (Fokien).—BOERTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 139 (Lueshan Mts., Kiukiang, China); p. 146 (Delanshan, near Ningpo); p. 149 (Weihaiwei and Chefu); Kat. Schl. Mus. Senckenberg., 1898, p. 25 (part: northern China).—WALL, Proc. Zool. Soc. London, 1903, p. 86 (Yangtse Valley).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 354 (Hankow).—NIKOLSKI, Zap. Imp. Akad. Nauk. S. Peterburg (8), XVII, No. 1, 1905, p. 222 (part: Hongkong; Possiet Bay; etc.).—*Amphisma tigrinum* STONE, Proc. Phila. Acad., 1899, p. 184 (Peking; northern Pechili).
1859. *Tropidonotus lateralis* BERTHOLD, Nachrichten Ges. Wiss. Goettingen, 1859, p. 180 (type-locality, China; type in Goettingen Mus.).—JAN, Icon. Ophid., livr. 25, 1867, pl. v, fig. 1 (Bangkok, Siam).
1862. *Tropidonotus orientalis* GUENTHER, Proc. Zool. Soc. London, 1861, p. 391 (type-locality, Peking; types in Brit. Mus.; Swinhoe, collector); Ann. Mag. Nat. Hist. (3), IX, Jan., 1862, p. 59, pl. IX, fig. 3.—STRAUCH, in Przewalski's Mongoliya i Strana Tangutov, III, 1876, p. 48 (Ordos; Peking).

A glance at the list of specimens and their scale formulas under *N. tigrina* (p. 276) and the present form (p. 280) is sufficient to show that the latter is separable from the former by the lower number of scutes covering the underside of the body. There is not a very great difference in the number of ventrals (average 160.5 ventrals in 62 *N. tigrina*, and 155.6 in 20 *N. lateralis*) but the subcaudals are much more numerous in the Japanese form than in the Chinese. Thus in 54 specimens of the former they range between 66 and 85, while in 18 specimens of the latter between 53 and 64, thus leaving a perceptible gap between these two series. The sum of ventrals and subcaudals in doubtful cases will assist in identifying these specimens, for I find that in the above series the total number of scutes in *N. tigrina* varies between 227 and 248, and in *N. lateralis* between 206 and 224, the difference between the extremes in the former being 21 and in the latter 18. The average between these two series, 236.1 for *N. tigrina* and 215.2 for *N. lateralis*, therefore, is probably very nearly exact.

While there is thus a small gap in the formulas available, it can hardly be doubted that larger series would bridge it, and in view of the exceedingly close relationship of the two forms and their ranges unquestionably meeting in Korea, it is thought best to use a trinomial appellation for one of them. According to present usage and

^a Signifying Lateral, referring to the spots on the sides.

the various codes of nomenclature, the last-named form receives the trinomial; thus in this case the Chinese form becomes *Natrix tigrina lateralis*.

In the above discussion I have made no reference to a specimen in the Imperial Museum, Ueno Park, Tokyo (No. 20), alleged to have been collected at Naha, Okinawa Shima. As already mentioned under *N. tigrina*, this locality is so unsupported by other evidence that one is justified in suspecting a mistake in labeling. This specimen, according to the count which I made while in Tokyo in 1895, has 163 ventrals and 64 subcaudals—showing the minimum in the above-mentioned series of Japanese specimens. It is otherwise identical with specimens of the tiger snake collected around Tokyo, and I have but little doubt that it is a Japanese specimen of *N. tigrina*, with a minimum number of scutes, if my count was really correct.

There is no specimen of this form in the United States National Museum, but as the chief difference from *N. tigrina* consists in the lower number of ventrals and subcaudals, as indicated above, a detailed description is not necessary. In all other respects the scale formula agrees with that of the Japanese form. The color appears to differ in the absence of alternation of the dorsal and lateral spots. Doctor Guenther describes the color of the types of his *Tropidonotus orientalis* as follows:

“Greenish olive, with three series of black spots anteriorly, becoming very indistinct on the middle of the trunk; a black suberescient spot on each side of the neck, without yellow; posterior margins of the upper labials and a spot on the temple black. Belly more or less blackish.”

Habitat.—The mainland form of the tiger snake is distributed along the eastern coast of Asia from Olga Bay in the Russian Coast Province to Hainan and, if Jan's localities are correct, to Cochin China and Siam.^a Boettger even seems to think that the specimen in British Museum credited to Hainan may have come from some other locality, or at least from the mountainous interior of that island. Possibly the Siamese origin also refers to specimens obtained in the mountains of the interior. It is not recorded from Formosa. Westward it has been found by Berezowski as far as southern Kansu, the province next to Thibet.

Both Mr. Leech and Doctor Giglioli obtained specimens in Gensan, northern Korea, and the Warshaw Museum has it from Strelok, a military post on the Bay of Peter the Great. Giglioli even obtained

^a It may not be out of place to remark here that in Proc. Zool. Soc. London, 1899, p. 661, Mr. Stanley S. Flower gives a table of 21 specimens from Penang, Kedah, and Bangkok, which has the appearance of belonging to *Tropidonotus tigrinus*. This is due to a mistake in the typography, as the table, which is not self-explanatory, was apparently meant to follow the account of *Tropidonotus piscator*.

it as far north as Olga Bay, also the west side of the Sea of Japan. The types of Guenther's *T. orientalis* were collected by Swinhoe at Peking and the Philadelphia Academy has it from the same locality.

List of specimens of Natrix lateralis.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Subcaudals.	Ventrals + Subcaudals.
Goettingen		(a)	China			19	157	56	213
Brit. Mus.	k	Male ^b	Peking, China		R. Swinhoe	19	153	63	216
Do.	l	Young ^b	do.		do.	19	164	60	224
Do.	m	Male	Chefu, China		do.	19	150	64	214
Do.	n	do.	do.		do.	19	161	57	218
Do.	o	do.	do.		do.	19	154	65	219
Do.	p	Female	do.		do.	19	157		
Do.	t	Young	do.			19	151	64	215
Do.	u	Young ^c	do.			19	153		
Do.	q	Female ^c	Ningpo, China			19	160	55	215
Do.	r	do.	Kiukiang, China		A. E. Pratt	19	15		
Do.	s	do.	Hainan Island, China		J. Neumann	19	160	56	216
Senckenberg	7249 c	Halfg. ^e	Chefu, China		B. Schmacker	19	154	63	217
Do.	7249 e	do.	do.		do.	19	159	53	212
Do.	7249 d	Adult ^f	Dulanshan, China		do.	19	159	58	217
Do.	7249 i	do.	Weihaiwei, China		do.	19	159	54	213
Do.	7249 k	do.	Lueshan Mts., China		do.	19	151	55	206
Munich		Male ^h	Hankow, China		Doctor Haberer	19	150	63	213
St. Petersb.		(i)	Ordos, China		Przewalski	19	147	63	210
Do.		(i)	Peking, China		do.	19	153	59	212
Brit. Mus.	i	Male ^d	Gensan, Korea		J. H. Leech	19	159	63	222

a Type of *lateralis*.

b Type of *orientalis*; Boulenger, Cat. I, p. 250.

c Boulenger, Cat. III, p. 607.

d Boulenger, Cat. I, p. 250.

e Boettger, 1894, p. 149.

f Boettger, 1894, p. 146.

g Boettger, 1894, p. 139.

h Werner, p. 354.


i Strauch, p. 48.

NATRIX STOLATA a (Linnæus).

1758. *Coluber stolatus* LINNÆUS, Syst. Nat., 10 ed., I, p. 219 (type-locality "America; type in Roy. Mus. Stockholm); 12 ed., I, 1766, p. 379 (Asia).—*Natrix stolatus* MERREM, Syst. Amph., 1820, p. 123.—STREJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 221 (Taipa, Formosa).—*Tropidonotus stolatus* BOIE, Isis, 1827, p. 535.—SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 225 (Tamsui, Formosa).—GUENTHER, Rept. Brit. India, 1864, p. 266 (India to Formosa).—BOULENGER, Fauna Ind., Rept., 1890, p. 348, fig. (India to southern China); Cat. Snakes Brit Mus., I, 1893, p. 253 (India to China, Hainan, Formosa).—BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 132 (Hainan).—WALL, Proc. Zool. Soc. London, 1903, p. 86 (Hongkong).—*Amphisma stolatum* DUMÉRIl and BIBRON, Erpét. Gén., VII, Pt. 1, 1854, p. 727.
1801. *Elaps bilineatus* SCHNEIDER, Hist. Amph., II, p. 299 (type-locality, East Indies; type in Mus. Bloch).

a Signifying wearing a stole, with reference to the two light-colored longitudinal bands

Description (figs. 244-246).—*Adult male*: U.S.N.M. No. 34036; Taipa, Formosa; March, 1903; A. Owston collection. Rostral much broader than high, well visible from above; internasals slightly shorter than the prefrontals, very narrow in front, their suture with rostral considerably shorter than suture between rostral and first labial; prefrontals in contact with supraoculars; frontal longer than its distance from tip of snout and interparietal suture; parietals equaling the distance of eye from tip of snout; nostril large, between two subequal nasals; loreal somewhat longer than high, upper edge shorter than and parallel with lower; one preocular, not in contact with frontal; 3 postoculars; temporals 1+2; supralabials 8, third, fourth, and fifth entering eye, sixth and seventh largest; 5 lower labials in contact with anterior chin-shields which are shorter than the posterior; 19 rows of scales, strongly keeled except outer row, and without apical pores; 149 ventrals; anal divided; 81 pairs of sub-caudals. Color (in alcohol) above brownish gray with numerous



FIGS. 244-246.—*NATRIX STOLATA*. 2 × NAT. SIZE. 244, TOP OF HEAD; 245, SIDE OF HEAD; 246, UNDER-SIDE OF HEAD. No. 36510, U.S.N.M.

narrow black crossbars alternating on each side of the median line anteriorly but continuous farther back; across this pattern two longitudinal, dorso-lateral pale bands occupying the whole of sixth scale row and the adjacent halves of fifth and seventh rows; posteriorly these bands are nearly uniformly pale, but anteriorly they exhibit a kind of chain pattern, inasmuch as the outer edges of the middle scale row are black except where the band intersects as black crossbar; below the lateral band many small irregularly alternating black spots; top of head with obscure dusky edges to the shields; no nuchal collar; supralabials whitish, the light color extending upward on the preocular and the lower postoculars, the vertical edges of the labials heavily margined with black as are also the preocular in front and the postoculars behind, the vertical, black edged, white bar in front of the eye being very characteristic; underside uniform whitish, each ventral with a black mark near the outer edge.

Dimensions.

	<i>mm.</i>
Total length.....	522
Snout to vent.....	382
Vent to tip of tail.....	140

This snake is not a large one, and a length of 700 mm. is probably seldom exceeded.

Variation.—Comparatively little variation is noted in the head shields of this species. The ninth supralabial noted in one of the Taipa specimens is merely a narrow wedge interpolated between the second and third supralabials, as shown in fig. 245. The range of ventrals and subcaudals is also restricted in the Formosan series, as shown in the table given below, but in specimens from all over the wide area inhabited by this species they vary between 120 and 161 and between 50 and 89, respectively. It seems, however, as if this wide range is more or less correlated with various geographical subdivisions.

Habitat.—From India to Philippine Islands and China. One of the most widely distributed species in southeastern Asia.

There are five species in the British Museum, collected by R. Swinhoe in Formosa, and I have recorded four specimens from Taipa. Formosa, collected by Mr. Tada. Our museum has since received three more from the same locality through Mr. Owston. In the Hamburg Museum there is also a specimen (No. 1563) collected by Doctor Warburg at Kelung, northern Formosa.

List of specimens of *Natrix stolata*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ven. trais.	Anal.	Subcaudals.	Oculars.	Temporals.	Supra-labials.
U.S.N.M.	34035	Male	Taiipa, Formosa	Mar., 1903	A. Owsfou.	19	148	2	1, 3	1+2	8
Do.	34036	do a	do	do	do	19	149	2	81	1, 3	1+2	8
Do.	34037	Female	do	do	do	19	149	2	1, 3	1+2	8
Do.	36510	do b	do	Sept., 1897	T. Tada.	19	146	2	74	1, 3	1+2	8-9
Sci. Coll. Tokyo	b 14	do	do	do	do	19	150	2	74	1, 3	1+2	8
Do.	c 14	Male	do	do	do	19	146	2	81	1, 3	1+2	8
Do.	d 14	Female	do	do	do	19	147	2	74	1, 3-4	1+2	8
Hamburg	1563	Male	Kelung, Formosa	Doctor Warburg	19	142	2	77	1, 3	1+2	8
Brit. Mus	p	do c	Formosa	R. Swinhoe	19	147	2	65
Do.	r	do e	do	do	do	19	147	2	75
Do.	σ	Female e	do	do	do	19	148	2	75
Do.	τ	do e	do	do	do	19	149	2	72
Do.	v	do e	do	do	do	19	149	2	73

a Description, p. 281.

b Figs. 244-246.

c Boulenger Cat. I, p. 254.


NATRIX PRYERI^a (Boulenger).

GARASU HEBU (Okada).

Plate XX.

1887. *Tropidomotus pryeri* BOULENGER, Proc. Zool. Soc. London, 1887, p. 149, pl. xviii, fig. 3^b (type-locality "Loo Choo Islands;" types in Brit. Mus.).—BOULENGER, Cat. Snakes Brit. Mus, I, (1893), p. 250 (Great Loo Choo I.).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 862; author's separate p. 13 (Nafa, Okinawa shima).—BOETTGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 107 (Okinawa shima: Miyakoshima group); Kat. Schlangen Mus. Senckenberg., (1898), p. 26 (Okinawa shima and Miyakoshima); WALL, Proc. Zool. Soc. London, 1903, p. 100 ("Nawa, Loo Choos"); 1905, II, p. 515 (Okinawa; Amami; Irimote).

Description.—*Adult female*; Science College Museum, Tokyo, No. 6; Irimote shima, southern group, Riu Kiu Archipelago; Tashiro, col-




FIGS. 247-249.—NATRIX PRYERI. $\frac{1}{4}$ X NAT. SIZE. 247, TOP OF HEAD; 248, SIDE OF HEAD; 249, UNDERSIDE OF HEAD. NO. 6, SCI. COLL. TOKYO.

lector (figs. 247-249). Rostral nearly twice as broad as high, just visible from above; internasals fully as long as prefrontals, rather truncate anteriorly, their suture with rostral equaling that of the latter with first supralabial; prefrontals in contact with supraoculars; frontal longer than its distance from tip of snout, wider in front than behind, wider than supraoculars; parietals longer than distance of eye from tip of snout; nostril between two subequal nasals; loreal longer than high; one preocular, not in contact with frontal; three postoculars (normal), four on right side; 2+1 temporals, the lower one of the second row on both sides abnormally

^a For Henry James Stovin Pryer, a well-known naturalist and collector, who lived, as a merchant in Yokohama for many years. He was at one time connected with the Museum of the Educational Department Museum in Tokyo. Born in London, June 10, 1850; died in Yokohama, February 17, 1888.

^b Reproduced in this work on Plate XX.



MATRIX PRYER. (FROM BOULENGER.)
FOR EXPLANATION OF PLATE SEE PAGE 555.

fused with seventh supralabial; eight supralabials, fourth and fifth entering eye, sixth and seventh largest; five lower labials in contact with anterior chin-shields which are shorter than the posterior; 19 rows of scales, with two apical pits, all keeled; 170 ventrals; anal divided; (tail broken). Color (in alcohol) pale clay-color, with long, dark, dorso-lateral, slightly alternating blotches anteriorly separated by whitish, black-edged, vertical bars on the sides, this pattern shortening up posteriorly and resolving itself into smaller dusky spots on the back and a series of whitish spots on the sides; supralabials whitish, the anterior five with black posterior margin; an oblique black line from the middle postocular, across lower portion of lower anterior temporal, sending a short branch downward along upper third of posterior edge of sixth supralabial, crossing the seventh labial nearly diagonally and the lower anterior corner of the eighth labial to the angle of the mouth; the lower, anterior and upper margins of the latter black, isolating a sharply-defined whitish spot; behind the angle of the mouth a round whitish black-edged spot continued backward and upward to the middle of the neck as a similarly edged but slightly narrower whitish band forming a crescent, or rather a chevron, on the nape; underside uniform pale, with a series of small dusky spots on each side near the outer angle of the ventrals and subcaudals. .

Dimensions.

	mm.
Snout to vent.....	675
Tail defective.....	

The tail in this species is nearly one-third the total length. A specimen in British Museum measuring 930 mm. has a tail 300 mm. long. A similar proportion would give a total length for the specimen described above of about 1 meter, probably near the maximum length of this species.

Variation.—Scale formula fairly constant. Temporals are said rarely to be 1+2; the specimen described above has 2+1 instead of 2+2; it has also four postoculars on one side instead of three, the normal number. Dr. Wall has recorded a specimen with two postoculars on one side. Ventrals range between 170 and 181 in thirteen specimens, and subcaudals between 114 and 128 in seven specimens.

Remarks.—The color pattern of head and nape of this species is an enlarged copy of that of *Natrix vabakari*, *craspedogaster*, and *khasiensis*: the black margin to the anterior supralabials; the oblique, black postocular band crossing the upper half of the penultimate labial and the lower margin of the last one, thereby separating it from the others; the dark upper edge to the last thus isolating a well circumscribed whitish spot on the latter; the subsequent postoral light spot and temporo-nuchal crescent edged with dusky; each one

of these features is more or less plainly indicated in all these species. The median nuchal light streak described in *N. vibakari* is not visible in the specimen of *N. pryeri* before me, but it is very distinct in the figure accompanying the original description of the species^a herewith reproduced (Plate XX, fig. 2); the lateral spots on the ventrals are common to all these species, though not by any means confined to them. The white dorso-lateral spots of the young of *N. vibakari* are very marked in *N. craspedogaster* and also on the posterior part of *N. pryeri*. The sudden increase in the size of the posterior maxillary teeth of the latter is no real obstacle, for as I have shown above (p. 264) *N. vibakari* may with propriety be said also to have suddenly enlarged posterior maxillaries. Moreover, *Natrix chrysarga*, which all authors agree in referring to the so-called "*Amphiesma*" group, appears also to belong to the assembly of species here discussed, in spite of the fact that the posterior labials are not quite typically marked, though judging from Jan's figure of his *Amphiesma platyceps*, the Himalayan specimens at least may come pretty close to the characteristic pattern.

Habitat.—This species has now been recorded from all the groups of the Riu Kiu chain. The types were from the "Loo Choos," without further locality, but both Holst and Fritze obtained it in Okinawa shima, from which island specimens also are in the National Museum in Tokyo (no. 8) and in Museum Senckenbergianum. The latter also has a specimen from Miyakoshima in the Saki shima group, both obtained through Mr. Schmacker. Finally, a specimen from Nishiomote shima, also called Iriomote shima, of the same group, is in the Science College Museum, Tokyo. Dr. Wall has recently recorded a specimen from the latter island and three from Amami-o-shima, of the northern group.

^a Proc. Zool. Soc. London, 1887, pl. xviii, fig. 3.

List of specimens of *Natrix pryferi*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected, or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supra-labials.
Sci. Coll. Tokyo.....	6	Female ^a	Iriomote shima, Riu Kiu.	Tashiro.....	19	170	2	1,3-4	2+1	8
Nat. Mus. Tokyo.....	8	Male.....	Okinawa shima, Riu Kiu.	19	178	2	1,3	2+2	8
Do.....		Female.....	do.....	19	178	2	1,3	2+2	8
Do.....		do.....	do.....	19	174	2	121	1,3	2+2	8
Brit. Mus.....	a	Male ^b	"Loo Choo".....	H. J. S. Pryer.....	19	180	2	121	1,3	8
Do.....	b	do.....	do.....	do.....	19	181	2	1,3	8
Do.....	c	do.....	do.....	do.....	19	181	2	128	1,3	8
Do.....	d	Female ^b	do.....	do.....	19	176	2	1,3	8
Do.....	e	do.....	do.....	P. A. Holst.....	19	177	2	114	1,3	8
Do.....	f	Young ^c	Great Loo Choo.	do.....	19	171	2	118	1,3	8
Senckenb.....	7250a	do.....	Okinawa.....	B. Schmaecker.....	19	175	2
Do.....	7250b	Adult ^d	Miyako shima.	do.....	19	176	2	118
Do.....		do.....	do.....	do.....	19	179	2	125

^a Description, p. 284, figs. 247-249.

^b Boulenger, Cat. I, p. 251; types.

^c Boulenger, Cat. I, p. 251.


^d Boettger, 1895, p. 107.

NATRIX PISCATOR^a (Schneider).

1799. *Hydrus piscator* SCHNEIDER, Hist. Amphib., I, p. 247 (type-locality, East Indies).—*Natrix piscator* MERREM, Syst. Amphib., 1820, p. 122.—STENNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 221 (Taipa, Formosa).—*Tropidonotus piscator* BOIE, Isis, 1827, p. 535.—BOULENGER, Cat. Sn. Brit. Mus., I, 1893, p. 230 (India to southern China and Malay Archipelago); Proc. Zool. Soc. London, 1899, p. 163 (Kuatun, Prov. Fokien, China).—BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 132 (Hainan); Kat. Schl. Mus. Senckenberg., 1898, p. 22 (India, China, and Japan).—WALL, Proc. Zool. Soc. London, 1903, p. 86 (Hongkong).
1837. *Tropidonotus quincunciatus* SCHLEGEL, Phys. Serp., II, p. 307, pl. XII, figs. 4-5 (part; type-locality, Java).—JAN, Icon. Ophid., livr. 27, 1868, pl. I, figs. 1-2 (Bengal).—BOETTGER, Offenbach. Ver. Naturk. 24-25 Ber., 1885, p. 150. (China); 26-28 Ber., 1888 (p. 76).
1860. *Amphisma flavipunctatum* HALLOWELL, Proc. Phila. Acad., 1860, p. 503 (type-locality, Canton River, China; type in Phila. Acad.; cotype U. S. Nat. Mus. No. 7387; Hongkong; Brooke, collector).

A large number of synonyms, many based on the various plates in Russell's "Indian Serpents," are here purposely omitted.

Description (figs. 250-252).—*Adult female*; U.S.N.M. No. 36511; Sci. Coll. Tokyo, No. 13; Taipa, Formosa; September, 1897; T.



FIGS. 250-252.—*NATRIX PISCATOR*. $1\frac{1}{2} \times$ NAT. SIZE. 250, TOP OF HEAD; 251, SIDE OF HEAD; 252, UNDER-SIDE OF HEAD. NO. 96, SCI. COLL. TOKYO.

Tada, collector. Rostral much broader than high, visible from above; internasals much narrowed anteriorly, their sutures with the rostral scarcely more than one-half the suture of the latter with the first supralabial; prefrontals broadly in contact with supraoculars; frontal, with almost parallel sides, longer than its distance from tip of snout, and slightly longer than the interparietal suture; parietals longer than the distance of eye from tip of snout; nostril small, semi-lunar, in the upper posterior corner of the anterior nasal; loreal higher

^a Signifying a fisherman. The masculine form is retained notwithstanding the fact that *Natrix* is of the feminine gender, in analogy with the usage of Linnæus himself in writing *Coluber natrix*, a usage followed by all subsequent writers (*Tropidonotus natrix*). A similar case is *Agkistrodon contortrix*.

than long, upper edge considerably shorter than lower; one preocular widely separated from frontal; eye rather small; three postoculars; 2 + 2 temporals, the upper one of the second row fusing with the one behind it so as to form a large irregular shield bordering the parietal laterally; 9 supralabials, fourth and fifth entering eye, seventh and eighth largest; five lower labials in contact with anterior pair of chin-shields, which are much shorter than posterior; 19 rows of scales, without pores, rather feebly keeled, the two outer rows smooth; 140 ventrals; anal divided; 74 pairs of subcaudals. Color (in alcohol) pale tawny-olive, with numerous small black spots in four, alternating, longitudinal rows, the lower ones on the third and fourth scale rows largest and most distinct anteriorly; head nearly uniform above, with a pale black-edged oval spot on the posterior end of the interparietal suture and the scale immediately behind; a similar but smaller dot on each side of the interparietal suture near the frontal; sides of head pale, the anterior supralabials without black margins; a sharply defined narrow black line from the lower edge of the eye obliquely backward following the suture between sixth and seventh supralabials to the commissure; a similar line from the upper postocular parallel with the other across temporals and eight supralabials to the lip, which it crosses where the suture between eighth and ninth supralabials join the commissure; the broad band between these black lines lighter and somewhat yellower than the rest of the head; back of the angle of mouth a similarly colored crescentic nuchal band interruptedly outlined by black; lower side pale, ventrals and subcaudals narrowly margined with blackish.

Dimensions.

	<i>mm.</i>
Total length.....	673
Snout to vent.....	497
Vent to tip of tail.....	176

Specimens of this species measuring 1,200 mm. are on record.

The *young* (U.S.N.M. No. 34051) slightly more than 150 mm. long, is colored essentially as the adult described above, only the pattern is more definite, the spots larger, especially those on the sides which descend to the ventrals and join their blackish margins; there is also a median dorsal row of spots which has become obsolete in the adult, so that the back is regularly checkered; the contrast between the light color of the postocular band and the rest of the sides of the head is very marked.

Variation.—The six Formosan specimens examined by me show great constancy in the scale formula, as all have 19 scale rows, 9 supralabials, 2 + 2 temporals (one has 3 + 2 on one side), and one preocular. Four have 3 postoculars on both sides, one has 3 on one

side and 4 on the other, and one has 4 on both sides; the ventrals vary between 139 and 144, the subcaudals between 72 and 77 pairs. Boulenger notes the occurrence of 2+3 temporals, rarely 1+2; ventrals, 125-150; and subcaudals, 70-90. I may add here that Hallowell's cotype of *Amphiesma flavipunctatum*, from Hongkong (U.S. N.M. No. 7387) has the following scale formula:

sc. 19; v. 126; a 2; c. 85; l. 9; t. 2+2; oc. 1-3.

Doctor Wall, also in Hongkong specimens, found 5 postoculars on one side in one, and four on both sides in another; one had 8 supralabials on both sides, fourth only touching the eye.

The range of variation in coloration is very considerable, viz, from nearly uniform dark with small white spots, due to the great size of the blackish spots, to nearly uniform pale drab without any black markings whatsoever. Two specimens from Formosa (No. 16 a, Sci. Coll. and No. 36512 U.S.N.M.) show the latter peculiarity and in every respect agree with the Himalayan specimens, which have been called *Tropidonotus sanctijohannis*.^a

Habitat.—The known range of this species embraces the whole of southeastern Asia from India to the Malay peninsula and archipelago, as well as Hainan and southern China as far north as Fokien.

Its occurrence in Formosa was first discovered by Mr. Tada, who collected five specimens at Taipa in November, 1897, as previously recorded by me.

In the collection recently acquired by the U. S. National Museum from Mr. A. Owston, there is a very young *N. piscator* (U.S.N.M. No. 34051) the label of which indicated that it has been collected at Mount Fuji. Here is undoubtedly a mistake, and it appears quite certain that the specimen in question has become confused with the four specimens of *Takydromus tachydromoides* (U.S.N.M. Nos. 34179-34183) said to have come from northern Formosa. This suggestion is corroborated by the fact that the latter, although said to have been collected on June 23, 1903, look as if they had been preserved for a much longer period, while the *Natrix*, which is alleged to date from August, 1898, is quite fresh. I therefore conclude that the latter was collected in northern Formosa June 23, 1903. (See p. 243.)

^aBoulenger, Cat. Snakes Brit. Mus., I, p. 230, pl. xv, fig. 1.

List of specimens of *Natrix piscator*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supralabials.
U. S. N. M.	7387	Male ^a . .	Hongkong, China.	1854. May --	Mr. Brooke.	19	126	2	85	1, 2	2+2	9
Do. . . .	34051	Young . .	Northern Formosa.	1903. June 23	A. Owston . .	19	142	2	130	1, 3-4	2-3+2	9
Do. . . .	36511	Female ^b	Taipa, Formosa	1897. Sept. --	T. Tada	19	140	2	74	1, 3	2+2	9
Do. . . .	36512	. . do. ^c do. do. do.	19	139	2	76	1, 3	2+2	9
Sci. Coll. Tokyo.	9a	Female	. . do. do. do.	19	139	2	76	1, 3	2+2	9
Do. . . .	9b	. . do. ^d do. do. do.	19	144	2	77	1, 3	2+2	9
Do. . . .	16a	. . do. do. do. do.	19	142	2	77	1, 4	2+2	9

^a Type of *A. flavipunctatum*. ^b Description p. 288. ^c Sci. Coll. Tokyo No. 16 b. ^d Figs. 250-252.

NATRIX ANNULARIS ^a (Hallowell).

1856. *Tropidonotus annularis* HALLOWELL, Proc. Phila. Acad., 1856, p. 151 (type-locality, Ningpo, China; type in Mus. Phila. Acad.; Doctor McCartee, coll.).—SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 225 (Tamsui, Formosa).—GUENTHER, Rept. Brit. India, 1864, p. 261 (Ningpo; Chikiang; Formosa); Ann. Mag. Nat. Hist. (6), I, 1888 (p. 171) (Mts. N. of Kiukiang).—BOULENGER, Cat. Snakes Brit. Mus., I, 1893, p. 233 (China, Formosa).—BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 146 (Chinhai, near Ningpo).—WALL, Proc. Zool. Soc. London, 1903, p. 86 (Yangtse Valley).
1859. *Tropidonotus chinensis* BERTHOLD, Nachrichten Ges. Wiss. Goettingen, 1859, p. 180 (type-locality, China; type in Goettingen Mus.).

A nearly related form has recently been described by Doctor Werner as *Natrix habereri* ^b from the Ningpo Mountains, near Shanghai. It has practically the same scale formula as *N. annularis* (viz, 163-164 ventrals and 53-65 pairs of subcaudals), but the three outer scale rows are said to be smooth and the eye larger; the keels on the scales are not so strong. Above plumbeous, below red; markings as in *N. annularis*. Boulenger regards them as identical. ^c


Description.—*Adult male*; U.S.N.M. No. 34044; northern Formosa, May 25, 1903; A. Owston, collector (figs. 253-255). Rostral much wider than high, just visible from above; internasals narrow, espe-

^a Signifying with small rings, referring to ventral black half-rings.

^b *Tropidonotus habereri* WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 354, pl. I, figs. 1-2 (type-locality Ningpo Mts., near Shanghai; types in Mus. Munich; Doctor Haberer, collector).

^c Zool. Rec., 1903, Rept. p. 24.

cially in front, their suture with the rostral being shorter than that of the latter with first supralabial; prefrontals broadly in contact with supraoculars; frontal shorter than its distance from tip of snout, scarcely equaling the interprefrontal suture, with nearly parallel sides; supraoculars narrow, scarcely more than half as wide as frontal; parietals much longer than distance of eye from tip of snout, as long as frontal and prefrontals together; nostril small, round behind the middle and near the upper edge of the nasal which is only divided into two unequal parts below the nostril; loreal nearly rectangular, slightly longer than high; one preocular, widely separated from frontal; eye small; three postoculars, the lower one almost horizontal; temporal 2 + 3, those of the first row very long and the upper especially very narrow; eight supralabials, fourth entering eye; five lower labials in contact with anterior pair of chin-shields, which are considerably



FIGS. 253-255.—*NATRIX ANNULARIS*. 2 × NAT. SIZE. 253, TOP OF HEAD; 254, SIDE OF HEAD ($\frac{3}{4}$ × NAT. SIZE); 255, ROSTRAL FROM FRONT. No. 34044, U.S.N.M.

shorter than posterior, chin-shields and labials studded with numerous small, rounded tubercles; 19 rows of scales, with two somewhat obscure apical pits, striated and strongly keeled, except outer row, which is feebly keeled; 163 ventrals; anal divided; 61 pairs of subcaudals. Color (in alcohol) on back dark grayish drab with ill-defined black reticulations or series of small ill-defined, alternating black spots; sides gradually much paler toward the ventrals, with numerous vertical black bars about two scales wide and two scales apart, the bars continued on the ventrals joining the corresponding ones of the other side as half-rings or alternating with them; scales included in black lateral bars with pale centers; shields on head obscurely outlined with dusky; labials whitish, with blackish sutures both on upper and lower lip, except on suture between fifth and sixth supralabials; an obscure, oblique blackish postocular streak; lower parts pale vinaceous-cinnamon, with black half-rings.

Dimensions.

	<i>mm.</i>
Total length.....	517
Snout to vent.....	415
Vent to tip of tail.....	102

Boulenger records a specimen of this species measuring 820 mm. in total length.

The *young* (U.S.N.M. No. 34049; Formosa), total length 212 mm., tail 42 mm., is quite like the adult, only darker with more sharply defined pattern on sides and abdomen; four outer scale rows on each side as light as underside.

Variation.—The number of supralabials varies between 8 and 9, and the one entering the eye is the fourth or the fifth accordingly. Boulenger states that two supralabials rarely enter the eye. There is occasionally only a single anterior temporal. Ventrals vary between 145 and 163 (in twelve specimens), subcaudals between 53 and 69 pairs (in 8 specimens).

Habitat.—Restricted apparently to Formosa and southeastern China, where it has been found as far north as Ningpo and in the Yangtse Valley as far west as Kiukiang.

In Formosa it has been collected by Swinhoe, whose specimens are in British Museum. The United States National Museum also has two specimens from northern Formosa obtained recently from Mr. A. Owston.

List of specimens of Natrix annularis.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supralabials.
						1903.						
U. S. N. M.	34044	Male ^a	Northern Formosa.	May 25	A. Owston.	19	163	2	61	1,3	2 + 3	8
Do.	34049	Young	do.	July 10	do.	19	149	2	53	1,3	7 + 3	9
Brit. Mus.	<i>g</i>	Female ^a	Formosa		R. Swinhoe.	19	147	2				
Do.	<i>h</i>	do. ^b	do.		do.	19	153	2	54			

^aDescription, p. 291, figs. 253-255.

^bBoulenger, Cat. 1, p. 233.

NATRIX SWINHONIS ^a (Guenther).

1868. *Tropiaonotus swinhonis* GUENTHER, Ann. Mag. Nat. Hist. (4), I, 1868, p. 420, pl. XIX, fig. *F*^b (type-locality, Formosa; type in Brit. Mus.: R. Swinhoe, collector).—BOULENGER, Cat. Snakes Brit. Mus., I, (1893), p. 218 (Formosa).

1888. *Tropidonotus swinhoei* BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., p. 137 (emendation).

^a For Robert Swinhoe, who collected the type; see p. 184.

^b See this work, fig. 256.

Only the type specimen of this species being known, the description by Boulenger is here reproduced.

Description of type specimen.—*Adult female*; British Museum; Formosa; R. Swinhoe, collector (fig. 256). Head narrow and elongate; eye moderate; rostral broader than deep, just visible from above; internasals a little broader than long, shorter than the prefrontals; frontal once and a half as long as broad, as long as its distance from the end



FIG. 256.—*NATRIX SWINHONIS*. ENLARGED. SIDE OF HEAD. COPY FROM GUENTHER, ANN. MAG. NAT. HIST., (4), I, 1868, PL. XIX, FIG. F.

of the snout, shorter than the parietals; loreal as long as deep; one preocular and three postoculars; temporals 1+2; 6 supralabials, third and fourth entering eye, fifth very long; 4 lower labials in contact with the anterior chin-shields which are shorter than the posterior; scales strongly keeled, of outer row faintly keeled, in 15 rows; ventrals 150; anal divided; subcaudals *circa* 50. Brown above, anteriorly with irregular blackish spots; occiput and nape reddish, with a broad black collar; a black spot below the eye, and an oblique black band on the temporal region; lower parts yellowish, powdered with brown on the sides.

Dimensions.

	<i>mm.</i>
Total length.....	580
Tail [defective] about.....	100

Habitat.—This species is described from Formosa. Only a single specimen is known, the type, collected by R. Swinhoe, being in the British Museum.

Remarks.—The nearest relative of this species appears to be *Natrix nuchalis* described in 1891^a by Boulenger from specimens in the British Museum, collected by A. E. Pratt at Ichang, on the upper Yangtse. The scale formula is the same, but *N. nuchalis* has the scales very feebly keeled, a shorter head, and a very distinct groove along the middle of the back.

Genus *ACHALINUS*^b Peters.

1869. *Achalinus* PETERS, Mon. Ber. Berlin Akad. Wiss., 1869, p. 436 (type, *A. spinalis*).

1877. *Ophidaps* SAUVAGE, Bull. Soc. Philom. Paris (7), I, p. 108 (type, *O. braconnieri*).

The statement that this genus has no postoculars is scarcely quite correct. It is true that in some of the young specimens no such shield can be detected, but in our large specimen, No. 34046, there is

^a Ann. Mag. Nat. Hist. (6), VII, p. 281.

^b From α , without; $\chi\alpha\lambda\iota\nu\acute{o}\varsigma$, poison fang.

plainly a small triangular, or rather semilunar, scute behind the eye fitting into the blunt angle between the anterior temporals. A similar scute is also discernible in the young specimen No. 34047.

Thus far the genus has only been recorded from southern China and Japan. Its nearest relatives seem to be found in Ceylon and the Malay islands. However, specimens are as yet rare and it has been but recently discovered, so that it is not safe to draw any conclusions from its distribution and relationship as known at present.

Boulenger recognizes three species of this curious genus, viz, *A. rufescens*^a and *A. braconneri*^b, from southern China, and *A. spinalis*, from "Japan (?)." Of these, the first and the last were only known from one specimen each at the time the first volume of his catalogue was published (1893), while of *A. braconneri* he had four specimens. *A. rufescens*, with its very long internasal suture and partly tricarinate scales^c may be regarded as unquestionably distinct. Additional specimens from Japan, however, show so much deviation from the type of *A. spinalis* and so much variability in the direction of *A. braconneri* that it may become difficult to maintain the latter as a distinct species. The characters which are supposed to distinguish the two species break down one by one. *A. braconneri* was characterized as having 23 to 25 scale rows as against 21 in the type of *A. spinalis*, but five additional specimens from Japan seem to indicate that 23 is the normal number in the latter. Furthermore, in *A. braconneri* the suture between the internasals is said to be shorter than that between the prefrontals, while in the type of *A. spinalis* these sutures are described as of equal length.^d This is also approximately the case in our specimen from Kiusiu (U.S.N.M. No. 30731), but in the specimen in the Science College Museum, in Tokyo, also from Kiusiu, the interprefrontal suture is much longer, the one between the internasals being only 0.3 the length of the one between the prefrontals. In the two specimens from Mount Fuji (U.S.N.M. Nos. 34046-7) the relations are 0.4 and 0.6. Finally, *A. braconneri* has

^a BOULENGER, Ann. Mag. Nat. Hist. (6), II, 1888, p. 43, (type-locality, Hongkong); Cat. Sn. Brit. Mus., I, 1893, p. 308, pl. xx, fig. 2; III, 1896, p. 612.

^b *Ophiclaps braconneri* SAUVAGE, Bull. Soc. Philom. Paris, (7), I, 1877, p. 109 (type-locality, eastern Kiangsi, China).—*Achalinus braconneri* BOULENGER, Cat. Sn. Brit. Mus., I, 1893, p. 309; III, 1896, p. 613.

^c The ventrals of the type are given as 136, subcaudals 82, and Boulenger (Cat. Snakes Brit. Mus., III, 1896, p. 612) in another male specimen from Hongkong counts 137 ventrals and 76 caudals. This is greatly at variance with the number of counts given by Doctor Wall for four Hongkong specimens, namely, 150+56, 158+58, 158+61, and 154+58, respectively. In view of the apparent narrow limits of variation in this series as well as in *A. spinalis*, I am at a loss how to explain the above discrepancy.

^d Peters says: "Die Praefrontalia sind nicht länger als die Internasalia." In the figure (fig. 1a), however, the prefrontal suture is considerably longer than the internasal suture. Which is correct?

two pairs of chin-shields, while the type of *A. spinalis* has three. But here, again, our specimen No. 30731 from Kiusiu steps in between as having only two, while the specimen in Tokyo and our two specimens from Fuji have three like the type.^a The slight difference in the number of ventrals and subcaudals is completely bridged over by the additional specimens. The coloration, on the other hand, is the only feature which *thus far* offers a character by which to distinguish the two forms. *A. braconnieri* is described by Boulenger as being uniform blackish, the ventrals having a narrow whitish edge. The Japanese specimens all seem to have the median black dorsal line and often a blackish longitudinal median stripe on the underside of the tail, the latter being absent in No. 34047 and very obscure in No. 30731, which also shows trace of a dorso-lateral dusky line.

It may be that an actual comparison of specimens from Japan and China may demonstrate further differences,^b but it is quite clear that the Japanese species is very closely allied to the Ichang specimens collected by Mr. Pratt and described by Boulenger.

Recently Doctor Wall^c has come to a similar conclusion as to the probable identity of the two species. He examined three Japanese specimens, which all had 23 scale rows, viz, the two specimens now in the U. S. National Museum alluded to above (Nos. 34046, 34047), and a third one from the same locality. Unfortunately he does not mention the coloration nor the number of chin-shields in the third specimen. It is also impossible to make out which of the scale formulas given belong to the the third Japanese specimen.^d

Whether, on the other hand, the Ichang specimens are identical with Sauvage's originals of *A. braconnieri* from eastern Kiangsi is another question. His description is certainly very different from Boulenger's. Thus he speaks of the temporals as being numerous; of the prefrontals as small; his specimens have a small lower postocular, and the number of labials he states to be eight; the color, moreover, is a "brun ardoisé uniforme."

^a I may here call attention to the fact that in Boulenger's figure of *A. rufescens* (Cat. Sn. Brit. Mus., 1. 1893, p. 308, pl. xx, fig. 2) there is only indicated two pairs of chin-shields, while the description speaks of three. This figure also shows only five supralabials as against six in the description.

^b It may be well to point out a difference in the dental formula of the genus and the dentition as I find it in the specimen in the Science College specimen, Tokyo, and in our No. 34046. In these I count 14-16 maxillary teeth, which are subequal, the first three and the last three slightly smaller, while Boulenger for the genus says: "Maxillary teeth 22 to 25, small, equal." Is this character derived from Ichang specimens, and have we here a possible difference in the two forms?


^c Proc. Zool. Soc. London, 1903, p. 88.

^d I presume that his scale formulas 165+44 and 165+48 are meant for our specimens No. 34047 and 34046, respectively. Repeated counts of these specimens give 163+45 and 165+49.

ACHALINUS SPINALIS^a Peters.

1869. *Achalinus spinalis* PETERS, Mon. Ber. Berlin Akad. Wiss., 1869, p. 436, pl. —, fig. 1 (type-locality unknown).—BOULENGER, Cat. Sn. Brit. Mus., I, p. 309 (Japan?).—NAMIYE, Annot. Zool. Japon., II, Pt. 1, 1898, p. 29, fig. (Kiusiu).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 33 (Hakone Mts.).—WALL, Proc. Zool. Soc. London, 1903, pp. 88, 100 (Mt. Fuji).

Description (figs. 257–259).—*Adult female*; U.S.N.M. No. 34046; Mount Fuji, Hondo; August, 1898; A. Owston, collector. Rostral scarcely visible from above, wider than high, and so much hollowed out below that the surface laterally is represented as a ridge only; internasals very small, narrow anteriorly, the suture with rostral less than half that of the latter with anterior nasal; prefrontals very large, the suture between them twice as long as the one between the internasals, broadly in contact with supraoculars, and with second supra-labial; frontal pentagonal, straight anteriorly, posterior angle acute,



FIGS. 257–259.—ACHALINUS SPINALIS. $2\frac{1}{2} \times$ NAT. SIZE. 257, TOP OF HEAD; 258, SIDE OF HEAD; 259, UNDERSIDE OF HEAD. NO. 30731, U.S.N.M.

as wide as long, shorter than distance from tip of snout, scarcely one-half as long as parietals; supraoculars very small, their width less than one-fourth that of the frontal; parietals exceedingly large, twice as long as distance of eye from tip of snout, pointed posteriorly; nostril large, horizontally oval in the anterior of two nostrils which are deeply hollowed out; loreal longer than high, entering eye and in contact with prefrontal and supraocular but not with posterior nasal; no preocular; eye very small, smaller than loreal; a crescentic trace of a postocular in angle between two anterior temporals; 2 + 2 temporals, upper anterior smaller than the others, in contact with supraocular; a large, elongate shield bordering the posterior half of the parietals; six supralabials, first exceedingly small, sixth exceedingly large, as large as the other five together, fourth and fifth entering eye; mental deeply hollowed out, the surface forming a V-shaped ridge; 3 lower labials in contact with anterior chin-shields, which together

^a Pertaining to the spine or backbone, referring to the black line along the middle of the back.

form a heart-shaped figure; 3 pairs of chin-shields, posterior shortest; 23 rows of small, nonimbricate, convex and keeled scales without apical pits, the soft skin mostly visible between the scales; 165 ventrals; anal entire; 49 undivided subcaudals. Color (in alcohol) above uniform "walnut-brown," more rufous toward the middle of back, with a very distinct black longitudinal stripe along the median line involving the median scale row and the adjacent half of the scale on either side; underside paler, more cinnamon colored, each scute with a narrow light edge and a darker submarginal line; a median dusky stripe on the underside of the tail.

Dimensions.

	<i>mm.</i>
Total length.....	412
Snout to vent.....	347
Vent to tip of tail.....	65

The *male* seems to have a relatively longer tail with more subcaudals. Thus the specimen in the Science College Museum, Tokyo, measures 410 mm. in total length, of which the tail is 90 mm. and contains 61 subcaudals.

The *young* is darker above in stronger contrast to the underside and seems to lack the dark median stripe on the underside of the tail.

Variation.—Only a few specimens are in the museums, but these show considerable variability. The six of which the individual scale formulas are recorded, agree in having 6 supralabials; no preoculars and no postoculars (except occasionally a trace); 2 + 2 elongated temporals; the scale rows vary between 21 and 23, six being known with 23 and only one, the type, with 21; ventrals between 146 and 166; subcaudals between 45 and 62; all have an undivided anal and undivided subcaudals. The specimen described above seems to be unusual in having the loreal separated from the posterior nasal by the prefrontal touching the second supralabial. The specimen figured (fig. 259) is also abnormal in possessing only two pairs of chin-shields. The Tokyo museum specimen shows a curious and rare anomaly in having the frontal divided longitudinally by a suture, and it is interesting to note that one of Sauvage's types of his *A. braconnieri*, from China, also has the frontal divided (though he does not say whether longitudinally or transversely); this specimen also has one of the supraoculars divided.

Habitat.—Under the supposition that *A. braconnieri*, which is recorded from eastern Kiangsi and from Ichang, in southern China, represents a distinct form, the present species is restricted to Japan.

Thus far only ten specimens are known, seven of which are from definite localities, which throw some light on the distribution of the species. Of these three are from Kiusiu, viz, the two which Mr. N. Takachiho collected at Hikosan, province of Buzen, near the northern

extremity of the island, and a fine specimen sent our museum by Rev. Cyrus A. Clark from Miyazaki, toward the southern end of the same island. The one in Museum Senckenbergianum in Frankfort on the Main, which was obtained from Mr. B. Schmacker in 1889 came from the Hakone Mountains in Hondo, and Doctor Wall mentions three specimens in Owston's collection from "Mount Fuji." Two of the latter are now in the United States National Museum (Nos. 34046, 34047).

List of specimens of Acalinus spinulalis.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supralabials.	Chin-shields.
U.S.N.M.	30731	Young ^a .	Miyazaki, Hiuga, Kiusiu.	C. A. Clark.	23	146	1	57	0, 0	2+2	6	2
				1808.									
Do.	34046	Female ^b	Mount Fuji, Hondo.	Aug. —	A. Owston	23	165	1	49	0, 0	2+2	6	3
Do.	34047	Young	do.	do.	do.	23	163	1	45	0, 0	2+2	6	3
Berlin		(c)	Japan?	21	149	1	62	0, 0	2+2	6	3
				1895.									
Sci. Coll. Tokyo.		Male ^d .	Hikosan, Buzen, Kiusiu.	June —	N. Takachiho.	23	163	1	61	0, 0	2+2	6	3
Takachiho.		(e)	do.	do.	23	166		51				
Owston.		(f)	Mount Fuji, Hondo.	23							

^a Figs. 257-259.
^b Description, p. 297.

^c Type; Peters, p. 436.
^d Frontal divided.

^e Namiye, p. 29.
^f Wall, p. 88.

Subfamily HOMALOPSINÆ.

The snakes of this group are apparently closely allied to the Natriicine snakes, differing chiefly in the presence of grooved fangs at the posterior end of the maxillary. They are even more thoroughly aquatic, having the nostrils on the upper surface of the snout and provided with valves by means of which they can be closed. The head is usually small and thick, scarcely distinct from neck, and the eyes are small mostly with a vertically elliptic pupil; very often there is only a single internasal shield.

The Homalopsine snakes are confined to southeastern Asia, including India, the Malay Archipelago, Philippines and southern China, New Guinea, and northern Australia. Two genera occur within our limits, inasmuch as two species of the genus *Enhydris* and one of *Hurria* have been recorded from Formosa.

The two genera may be distinguished as follows:

- a¹ scales smooth..... 1. *Enhydris*, p. 300.
- a² scales keeled..... 2. *Hurria*, p. 304.

Genus ENHYDRIS^a Latreille.

1802. *Enhydris* LATREILLE, Hist. Nat. Rept., IV, p. 200 (type, *E. carulea*=*Hydrus enhydris* Schneider) (not of Merrem 1820).
 1830. *Hypsirhina* WAGLER, Syst. Amphib., p. 169 (type, *H. aer*).
 1842. *Ferania* GRAY, Zool. Miscell., p. 67 (type, *H. sieboldii*).
 1843. *Hypiscopus* FITZINGER, Syst. Rept., p. 25 (type, *H. plumbea*).
 1843. *Pythomorphus* FITZINGER, Syst. Rept., p. 25 (type, *H. sieboldii*).
 1853. *Eurostus* DUMÉRIL, Prodröm. Class. Ophid., p. 102 (type, *E. dugsumieri*).
 1853. *Trigonurus* DUMÉRIL, Prodröm. Class. Ophid., p. 103 (type, *H. sieboldii*).

The authors who employ the generic name, "*Enhydris*, Merrem, 1820," for two of the Hydrid snakes which should correctly stand as *Lapemis*, seem to have overlooked entirely the fact that Latreille, as early as 1802, established the genus *Enhydris* for an entirely different set of snakes, embracing all Schneider's species of *Hydrus*, from *H. caspius* on. This name, which can not be discarded, as it is not a synonym of any name previously given, must have for type one of the six species enumerated by Latreille, and as neither he nor any of his successors have designated a type for the name, we are obliged to resort to the method of elimination. The last species mentioned by Latreille, *Enhydris dorsalis*, is of dubious application. The third, *Enhydris rynchops*, is the type of Daudin's *Hurria* (1803), Goldfusz' *Strephon* (1820), and Cuvier's *Cerberus* (1829), while three other species, viz, *E. caspius*, *piscator*, and *palustris* belong to the genus *Natrix* (*Tropidonotus*). A single species is thus left as type, viz, *Enhydris carulea*, which is Schneider's *Hydrus enhydris*. This species, or rather a synonym of this species, was afterwards (1830) made the type of *Hypsirhina*, and there can be no doubt but that the latter name will have to give way to *Enhydris* Latreille. The same result is obtained by those who would fix the type by employing the "principle of tautonymy."

Two species have been recorded as occurring in Formosa and for that reason are here included. They may be distinguished as follows:

- a*¹ Scales in 19 rows; supralabials 8; ventrals less than 150. *E. plumbea*, p. 300.
*a*² Scales in 21 rows; supralabials 7; ventrals more than 150. *E. bennettii*, p. 302.

ENHYDRIS PLUMBEA^b (Boie).

1827. *Homalopsis plumbea* BOIE, Isis, 1827, p. 550 (type-locality, Java).—*Hypsirhina plumbea* GRAY, Zool. Misc., 1842, p. 66.—GUENTHER, Rept. Brit. India, 1864, p. 280 (Java to Formosa).—JAN, Icon. Ophid., livr. 30, 1868, pl. v, figs. 2-3 (Java).—BOETTGER, Offenbach. Ver. Naturk. 24-25 Ber., 1885, pp. 123, 151 (China); Ber. Senckenberg. Naturf. Ges., 1894, p. 134 (Hainan).—BOULENGER, Cat. Snakes Brit. Mus., III, 1896, p. 5 (Formosa; China; Siam; Malay Peninsula; Java).—WALL, Proc. Zool.


^a From ἔνυδρις, a water snake.

^b Signifying lead-colored.

Soc. London, 1903, p. 94 (Hongkong).—*Enhydris plumbea* STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 222 (Taipa, Formosa).

1834. *Hypsirohina hardwickii* GRAY, Ill. Ind. Zool., II (pl. LXXXVII, fig. 1) (type-locality, Pinang; type in Brit. Mus.).

Description.—*Adult female*; Science College Museum, Tokyo, No. 11a; Taipa, Formosa; November, 1897; T. Tada, collector (figs. 260–262). Rostral much broader than high, just visible from above; internasal single, twice as wide as long, separated widely from rostral by nasals, separated from loreal; prefrontals broader than long, broadly in contact with supraocular; frontal longer than its distance from tip of snout, equaling interparietal suture, nearly straight anteriorly, sides parallel, twice as wide as supraocular at center of eye; parietals as long as their distance from rostral, rather narrow posteriorly; nostrils small, round, on upper surface of snout, in the posterior margin of the anterior nostrils which are broadly in contact behind the rostral; posterior nasal less than half the size of the anterior, smaller than



FIGS. 260–262.—*ENHYDRIS PLUMBEA*. 2 X NAT. SIZE. 260, TOP OF HEAD; 261, SIDE OF HEAD; 262, UNDER-SIDE OF HEAD. No. 11a, SCI. COLL. TOKYO.

loreal, which is slightly longer than high; eye small, its diameter less than its distance from edge of lip; one preocular, not in contact with frontal; two postoculars; temporals 1 + 2; 8 supralabials, increasing in height backward to the seventh, which is largest, and much higher than eight, first in contact with loreal, separating posterior nasal from second supralabial, fourth and fifth entering eye; five lower labials in contact with anterior chin-shields; posterior chin-shields slightly shorter than anterior, separated from each other by two scales; 19 rows of smooth scales without apical pits; 130 ventrals; anal divided; 31 pairs of subcaudals. Color (in alcohol) above uniform slate gray; labials and two outer scale rows whitish with gray edges; underside whitish, each scute with lateral edge and base narrowly margined with dark gray and with a median basal spot of the same color; on anal shield and the underside of the tail a sharply defined median line of dark gray.

Dimensions.

Total length.....	mm. 430
Snout to vent.....	382
Vent to tip of tail.....	48

Variation.—Few deviations from the specimens described above are recorded. In the other specimen from the same locality the posterior nasal is in contact with second supralabial. Sometimes only the fourth supralabial enters the eye. The number of ventrals varies between 120 and 134, subcaudals between 29 and 46. Coloration is also rather constant; a median dorsal series of small black spots may be present and the whole underside may be unspotted.

Habitat.—Originally described from the Malay Archipelago this species has since been found on the opposite mainland from Burma to southern China. It occurs in Formosa, there being two specimens in British Museum collected by R. Swinhoe, and two specimens were collected by Tsunasuke Tada in 1897 at Taipa.

List of specimens of *Enhydris plumbea*.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals	Supradentals.
U.S.N.M.	36513	Male ^a	Taipa, Formosa..	1897. Nov. —,	T. Tada ...	19	126	2	...	1, 2	1+2	8
Sci. Coll. Tokyo.	11a	Female ^b	do.....	do.....	do.....	19	130	2	31	1, 2	1+2	8
Brit. Mus.	(a)	Male ^c	Formosa.....	R. Swinhoe...	19	128	2	40
Do...	(b)	Female ^c	do.....	do.....	19	129	2	33

^a Sci. Coll. Tokyo, no. 11b.

^b Description, p. 301; figs. 260-262.

^c Boulenger, Cat., III, p. 5.

ENHYDRIS BENNETTII^a (Gray).


1842. *Hypsirhina bennettii* GRAY, Zool. Miscell., p. 67 (type-locality, China: type in Brit. Mus.; Bennett, collector).—GUENTHER, Rept. Brit. India, 1864, p. 283 (China).—W. L. SCLATER, List Sn. Ind. Mus., 1891, p. 55 (Formosa).
1885. *Hypsirhina bennetti* BOETTGER, Offenbach. Ver. Naturk. 24 and 25 Ber., p. 151 (China); Ber. Senckenberg. Naturf. Ges., 1894, p. 134 (Hainan); Kat. Schl. Mus. Senckenberg., 1898, p. 88 (Hainan; Herz, collector).—BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 8 (China).
1854. *Hypsirhina maculata* DUMÉNIL and BIBRON, Érpét. Gén., VII, Pt. 2, p. 950 (type-locality, China; type in Paris Mus.; Eydoux, collector).
1868. *Hypsirhina enhydris* var. *maculata* JAN, Icon. Ophid., livr. 30, pl. iv, fig. 1.

No specimen from Formosa being accessible to me, I describe one which probably hails from the mainland opposite.

Description.—*Male*; U.S.N.M. No. 22128; Hongkong, China?; Dale and Jouy, collectors (figs. 263-265). Rostral somewhat broader

^a Named after G. Bennett, esq., who collected the type.

than high, visible from above; internasal single, widely separated from rostral by nasals, small, not much broader than long, widely separated from loreal; prefrontals smaller than nasals, in contact with them and with supraoculars; frontal somewhat shorter than its distance from tip of snout, nearly twice as wide as supraoculars at center of eye; parietals rather broad, the lateral outline irregular and showing evidence of fusion with adjacent scales; nostrils semilunar, on top of snout, in a large nasal which on the left side is semidivided by a suture from nostril to first labial; nasals broadly in contact with each other behind rostral with prefrontals and first supralabials; loreal trapezoid, longer than high, in contact with first three supralabials; one preocular, not in contact with frontal; eye small, its diameter less than distance from edge of lip; two postoculars; 1+2 temporals; 8 supralabials, increasing in height backward to the sixth, which is highest, first in contact with loreal, thus exclud-



FIGS. 263-265.—ENHYDRIS BENNETTU. 2 × NAT. SIZE. 263, TOP OF HEAD; 264, SIDE OF HEAD; 265, UNDERSIDE OF HEAD. No. 22128, U.S.N.M.

ing nasal from second, fourth entering eye; three lower labials in contact with anterior chin-shields, which are followed by two pairs of smaller chinshields, both separated by scales; 21 rows of smooth scales without pits; 164 ventrals; anal divided; 64 subcaudals. Color (in alcohol) above drab gray, with two series of irregular dark brownish gray dorsal spots, which anteriorly unite into short crossbars and on the neck to a median longitudinal dark band; outer four scale rows white, the first and fourth having their lower and upper edge respectively margined with dark gray, forming an upper and lower dark indented edge to the lateral white band; labials and whole underside whitish, each scale and scute edged with dark gray, the ventrals with a median series of dark spots formed by a projection of the dark basal edge, and a similar arrangement on the subcaudals, causing a zigzag longitudinal dark band on the median line of the underside of the tail.

Dimensions.

	mm.
Total length.....	377
Snout to vent.....	314
Vent to tip of tail.....	63

Variation.—In the few specimens of this species recorded the number of ventrals varies between 158 and 164, subcaudals between 47 and 54.

Habitat.—Apparently restricted to China. The authority for the occurrence of this species in Formosa is a specimen in the Indian Museum (No. 12693) recorded by Mr. W. L. Sclater (*l. c.*). It was received from the Hongkong Museum in exchange.

List of specimens of Enhydris bennettii.

U. S. N.M. No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supraorbital.
22128	Male ^a ...	Hongkong ?.....	Dale and Jouy	21	164	2	54	1, 2	1+2	8

^a Description, p. 302; figs. 263-265.

Genus HURRIA ^a Daudin.

1803. *Hurria* DAUDIN, Bull. Soc. Philom. Paris, III, no. 72, March, 1803, p. 187 (type, *Hydrus rynchops*).
1813. *Huria* FISCHER, Zoognosia, 3 ed., I, p. 65 (emendation).
1815. *Hurrianus* RAFINESQUE, Anal. Nat., p. 77 (emendation).
1820. *Strephon* GOLDFUSZ, Handb. Zool., II, p. 151 (substitute for *Hurria*).
1829. *Cerberus* CUVIER, Règne Anim., 2 ed., II, p. 81 (type, *Coluber cerberus*).
1839. *Cerchus* SWAINSON, Classif. Fish. Amph. Rept., II (Lardner's Cab. Encycl.), p. 366 (*lapsus*).

The name *Cerberus*, by which this genus is generally known, must give way to *Hurria* of Daudin, 1803, the exclusive type of which is Russell's, Indian Serpents, pl. XL, which Doctor Boulenger identifies as *H. rynchops*.

HURRIA RYNCHOPS ^b (Schneider).

1799. *Hydrus rynchops* SCHNEIDER, Hist. Amphib., I, p. 246 (based on Russell, Ind. Serp., I, pl. XVII; type-locality, Ganjam).
1801. *Elaps boaciformis* SCHNEIDER, Hist. Amphib., II, p. 301 (type-locality not given; type in Mus. Univers. Halle).
1802. *Hydrus cinereus* SHAW, Gen. Zool., III, Pt. 2, p. 567 (based on Russell, pl. XVII).
1803. *Hurria schneideriana* DAUDIN, Hist. Nat. Rept., V, p. 281 (substitute name for *Elaps boaciformis* Schneider).
1803. *Hurria bilineata* DAUDIN, Hist. Nat. Rept., V, p. 284 (based on Russell, Ind. Serp., II, pl. XL; type-locality, Hyderabad).

^a From the Indian name Hurriah, by which Russell designated the snake which Daudin named *Hurria bilineata*.

^b From *Rynchops*, better *rhyrachops*, from $\rho\upsilon\chi\alpha\sigma$, snout, and $\omega\psi$, eye, or face.

1803. *Coluber cerberus* DAUDIN, Hist. Nat. Rept., VII, p. 167 (based on Russell, pl. XVII).
1820. *Python clapiformis* MERREM, Tent. Syst. Amph., p. 89 (substitute for *Elaps boacformis* Schneider).
1820. *Python rhynchops* MERREM, Tent. Syst. Amph., p. 90 (emendation).—*Cerberus rhynchops* GUENTHER, Rept. Brit. India, 1864, p. 279 (East Ind. archipelago, Ceylon, India, Malay peninsula, Siam).—BOULENGER, Cat. Snakes Brit. Mus., III, 1896, p. 16 (India, Malay peninsula and archipelago, Philippines, Pelew Islands).
1826. *Homalopsis molurus* BOIE, Isis, 1826, p. 213 (type-locality, Moluccas).
1837. *Homalopsis schneideri* SCHLEGEL, Phys. Serp., I., p. 171 (Pondichery, Bengal, Java, Timor, Amboina, New Guinea); II, p. 341 (*schneiderii*).
1837. *Coluber obtusatus* REINWARDT, in Schlegel, Phys. Serp., II, p. 341.
1837. *Cerberus russlii* CUVIER, in Schlegel, Phys. Serp., II, p. 342, footnote (Pondichery).
1849. *Cerberus acutus* GRAY, Cat. Snakes Brit. Mus., p. 65 (type-locality, Borneo; type in Brit. Mus.).
1849. *Cerberus unicolor* GRAY, Cat. Snakes Brit. Mus., p. 65 (type-locality, Philippines; type in Brit. Mus.; Cuming, collector).

Description.—*Adult male*; Christiania University Zoological Museum, No. 830; Formosa; Navara collector, 1894. Rostral pentagonal; the large semidivided nasals broadly in contact with each other behind the rostral; nostrils on upper side of snout; internasals small, triangular, separated widely from rostral, in contact with loreal; prefrontals smaller than nasals, in contact with internasals, loreal, prefrontal, supraocular, and frontal; frontal irregularly divided into four smaller scales; parietals broken up into small scales; supraoculars narrow; loreal lozenge-shaped; one high preocular; two suboculars separating eye from labials; one postocular; temporals 2 + 3, second row keeled; supralabials 10 on one side, 9 on the other, very high and narrow, the last three divided horizontally; 11 inferior labials, 4 in contact with anterior chin-shields; two pairs of chin-shields, the posterior nearly entirely lateral to the anterior; 25 scale rows, all keeled, except outer row; 145 ventrals; anal double; caudals, 63 pairs. Color (in alcohol) above ash gray with narrow blackish crossbands which do not reach the ventrals, about 25 between head and anus; a narrow blackish band from nostril, through eye, obliquely backward on the side of the neck; underside, including posterior supralabials and the three or four lower rows of scales whitish with numerous large blackish spots on the sides connected with the corresponding spot by irregular dark gray blotches across the ventrals; lower labials with a few small dusky spots.

Dimensions.

	mm.
Total length.....	570
Snout to vent.....	447
Vent to tip of tail.....	123

Variation.—The head scales of this species are subject to great variation. The frontal shield may be intact or broken up; eye may be surrounded by from four to six shields; internasals may be united, etc. The scale rows vary between 23 and 27, ventrals between 132 and 160, and subcaudals between 49–72.

Habitat.—This species lives in rivers and along the seacoasts of India east to Indo-China and the Philippines, and south to Ceylon and throughout the Malay countries and archipelago to Borneo, Celebes, Ceram, etc.

Its right to a place in this work is based upon a specimen collected in Formosa by Mr. Navara, and now in the zoological museum of the University of Christiania.

List of specimens of Hurria rynchops.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supralabials.
Christiania.....		Male ^a ...	Formosa.....	1894	Mr. Navara.	25	145	2	63	1,2,1	2+3	9-10

^a Description, p. 305.

Subfamily CORONELLINÆ.

The coronelline snakes embrace a great variety of forms of arbo-real or terrestrial habits, and consequently of the most different physiognomy. Having no grooved fangs of any description, they are of course nonpoisonous, although a few, like the *Dinodon*, with their vertical pupil, to some extent resemble various poisonous forms. But apart from the absence of grooves on the posterior maxillary teeth there seems to be no character by which they can be distinguished from the next subfamily.

Most of our snakes belong to this group, the genera of which may be identified by the following:

KEY TO THE GENERA OF CORONELLINE SNAKES KNOWN FROM JAPAN AND ADJACENT TERRITORY.

- a*¹ Scale rows round body in even numbers.....*Zaocys*, p. 352.
*a*² Scale rows round body in uneven numbers.
*b*¹ Anterior temporals present, separating parietals from supralabials.
*c*¹ Median maxillary teeth differentiated from the others by intervals; pupil vertically elliptic.....*Dinodon*, p. 356.
*c*² Median maxillary teeth not differentiated from the others; pupil round.
*d*¹ Scales with apical pits; head distinct from neck.
*e*¹ More than one loreal.....*Ptyas*, p. 345.
*e*² One loreal only.
*f*¹ Maxillary teeth increasing in size posteriorly; tail more than one-fourth of total length.....*Zamenis*, p. 349.
*f*² Maxillary teeth subequal; tail less than one-fourth of total length.
Elaphe, p. 307.

- d*² Scales without pits; head not, or scarcely, distinct from neck.
*e*¹ Maxillary teeth posteriorly strongly enlarged; portion of rostral visible from above as great as its distance from frontal *Holarchus*, p. 353.
*e*² Maxillary teeth equal; portion of rostral visible from above less than its distance from frontal *Liopeltis*, p. 337.
*f*² No anterior temporal, parietal being in contact with supralabials.....
Calamaria, p. 375.

Genus ELAPHE ^a Fitzinger.

1826. *Coluber* BOIE, Isis, 1826, p. 209 (type, *C. flavescens*; not of Linnæus).
 1833. *Elaphe* FITZINGER, in Wagler's Descr. et Icon. Amphib., III, text to pl. xxvii (type, *E. parreysii*).
 1834. *Calopeltis* FITZINGER, in Bonaparte's Iconogr. Fauna Ital., fasc. VII (type, *C. leopardina*).
 1840. *Elaphe* BONAPARTE, Mem. Accad. Sci. Torino (2), II, Sci. Fis. Mat., p. 402 (emendation).
 1843. *Pantherophis* FITZINGER, Syst. Rept., p. 25 (type, *Coluber guttatus*).
 1843. *Calopeltis* FITZINGER, Syst. Rept., p. 26 (emendation).
 1853. *Scotophis* BAIRD and GIRARD, Cat. N. Am. Rept., I, Serp., p. 73 (type, *Coluber alleghaniensis*).
 1860. *Leptophidium* HALLOWELL, Proc. Phila. Acad., 1860, p. 497 (type, *L. dor sale*).
 1860. *Proterodon* HALLOWELL, Proc. Phila. Acad., 1860, p. 498 (type, *P. tessellatus*).
 1862. *Natrix* COPE, Proc. Phila. Acad., 1862, p. 338 (type, *Coluber flavescens*; not of Laurenti).
 1864. *Phyllophis* GUENTHER, Rept. Brit. India, p. 295 (type, *P. carinata*).

The above synonymy refers chiefly to such names as have been employed in connection with species treated of in this work.

The generic name *Coluber* is commonly applied to this genus, but as will be shown under that heading further on, without strict observance of the rules of zoological nomenclature, inasmuch as the authors who have specifically selected types for a restricted genus *Coluber* have invariably selected species unknown to Linnæus at the time he instituted the genus. The first one to mention a specific type was Fleming, in 1822, but he selected a *Python* unknown to Linnæus. The next one was Boie,^b who again selected a post-Linnæan species, namely, Scopoli's *C. flavescens*.

^a This name is apparently a blunder for *Elaphis*. The latter is generally considered as derived from *ἐλάφος*, a deer, referring either to the coloring of the snake, or to its swiftness. Aldrovandi, who seems to be the source whence the modern authors received the name, apparently considered it as of the same origin as *Elaps* and *Elops*. See Aldrovandi Serpentum et Draconum Historia, Chapter XV, "De Elaphe, sive Elape aut Elope."

^b Isis, 1826, p. 209, and 1827, p. 518.

KEY TO JAPANESE, KOREAN, AND FORMOSAN SPECIES OF ELAPHE.

- a*¹ Scale rows 19..... *E. quadrivirgata*. p. 327.
*a*² More than 19 scale rows.
*b*¹ Scale rows 21.
*c*¹ Less than 195 ventrals..... *E. rufolorsata*. p. 310.
*c*² More than 195 ventrals..... *E. conspiciillata*. p. 334.
*b*² More than 21 scale rows.
*c*¹ One or two outer scale rows smooth, the others strongly keeled.
E. carinata. p. 308.
*c*² More than two rows of scales smooth.
*d*¹ More than 85 pairs of subcaudals.
*e*¹ Underside of tail black with a very distinct whitish band along the median suture between the subcaudals.
*f*¹ Scale rows, 23-25; ventrals, 229-258; subcaudals, 90-111 pairs.
E. taniurus. p. 319.
*f*² Scale rows, 25-27; ventrals, 251-260; subcaudals, 104-124 pairs.
E. schmackeri. p. 322.
*e*² Underside of tail not longitudinally striped..... *E. climacophora*. p. 324.
*d*² Less than 85 pairs of subcaudals.
*e*¹ Eleven to thirteen median scale rows distinctly keeled; parietal suture not longer than distance of frontal from rostral; postocular dark stripe crosses commissure in front of angle of jaw; color uniform or with wide dark spots and light cross bands..... *E. schrenckii*. p. 313.
*e*² Scales smooth with only faint traces of keels on the median ones; parietal suture longer than distance of frontal from rostral; postocular dark band reaches commissure at angle of mouth; color light with numerous narrow blackish cross bands..... *E. dione*. p. 315.


ELAPHE CARINATA^a (Guenther).

1858. *Elaphis sauromates* GUENTHER, Cat. Colubr. Sn. Brit. Mus., p. 93 (Ningpo, China) (not of Pallas); Rept. Brit. India, 1864, p. 241 (part: Ningpo), pl. XXI, fig. E.
 1864. *Phyllophis carinata* GUENTHER, Rept. Brit. India, p. 295, pl. XXI, fig. B (type-locality, China; type in Brit. Mus.); Ann. Mag. Nat. Hist. (6), I, 1888, p. 170 (Kiukiang).—*Elaphe carinata* STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 221 (Formosa).
 1891. *Coluber phyllophis* BOULENGER, Ann. Mag. Nat. Hist. (6), VII, May, 1891, p. 280 (type-locality, Kiukiang, China; types in Brit. Mus.; Pratt, collector); Cat. Sn. Brit. Mus., II, 1894, p. 55 (Kiukiang; Ningpo); Proc. in Zool. Soc. London, 1899, p. 165 (Kuatun, Prov. Fokien).—GUENTHER, Ann. Mus. Zool. St. Pétersbourg, I, 1896, p. 205 (Prov. Szechuen).—WALL, Proc. Zool. Soc. London, 1903, p. 92 (China).—WERNER, Abh. Bayer. Akad. Wiss. (Muenich), II Klasse, XII, Pt. 2, 1904, p. 356 (Ningpo Mts. near Shanghai).

Description.—*Adult male*; Science College Museum, Tokyo, No. 5; Taipa, Formosa; December, 1897; T. Tada, collector (figs. 266-268). Rostral about one-third wider than high, largely visible from above; internasals longer than broad, nearly as long as prefrontals, the latter broadly in contact with supraocular; frontal considerably longer than wide, longer than its distance from rostral, but

^a Signifying keeled; with reference to the strong development of the keels of the scales.

shorter than parietals, the sides strongly convergent backward, the width at the posterior lateral angle being less than width of supraocular; nostril large, between two nasals, the posterior of which is less than half as wide as the anterior; loreal much wider than high; one preocular, not in contact with frontal; a subpreocular; eye large, its diameter equaling half its distance from tip of snout; two postoculars; temporals 2 + 3; eight supralabials, seventh largest, fourth and fifth entering eye; five lower labials in contact with anterior chinshields, which are slightly longer than posterior; 23 rows of scales all strongly keeled except outer two rows; 214 ventrals, angularly bent laterally; anal divided; 97 pairs of subcaudals. Color (in alcohol) olive clay color, pale underneath; back with blackish ill-defined cross markings the black color only involving the edge of the scales



FIGS. 266-268.—*ELAPHE CARINATA*. $\frac{1}{3}$ \times NAT. SIZE. 266, TOP OF HEAD; 267, SIDE OF HEAD; 268, UNDERSIDE OF HEAD. No. 5, SCI. COLL. TOKYO.

and the skin between; a longitudinal blackish band on ventrals near the lateral angle and frequently interrupted; each ventral with the posterior edge irregularly marked with black; subcaudals narrowly edged with black so as to form a black median line; all the upper and lateral cephalic sutures marked with black.

Dimensions.

	<i>mm.</i>
Total length.....	1,455
Snout to vent.....	1,155
Vent to tip of tail.....	300
Tip of snout to posterior end of parietals.....	34

The *young* (U.S.N.M. No. 36508) are much paler and underneath uniform white; back with a few scattered black dots which, on the anterior half of the body, form short and narrow cross lines; on posterior half an obscure, pale brown longitudinal line on each side; upturned ends of ventrals with small black dots, forming an irregular longitudinal series; upper and lower labials with dark brown sutures.

Variation.—The commonest variation is apparently in the number of temporals, the extremes recorded (for Chinese specimens) being 1 + 2 and 3 + 3. The scale rows are almost always 23, but Wall mentions

a Chinese specimen with 21. Ventrals vary from 208 to 221 and caudals from 80 to 97. According to Boulenger an undivided anal and nine supralabials are rare features. Four lower labials in contact with anterior chin-shields as often as five. The above figures are derived from 17 specimens, of which only three are from Formosa.

This snake evidently grows to a great size, as Doctor Wall has recorded a specimen measuring 8 feet 1 inch (2.44 meters.).

Habitat.—Peculiar to China, where it has been obtained from near Shanghai, in the north, to Kuatun in Fokien in the south, and as far west as Lung-ngan-fu in the western part of the province of Sze-chuen.

In Formosa three specimens were collected by Mr. Tada in 1897, and a fourth specimen is in the zoological museum in Christiania, collected by Mr. Navara in 1894.

List of specimens of *Elaphe carinata*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Supralabials.	Temporals.
				1897.							
U.S.N.M.	3508	Young a	Formosa	Sept. 29	T. Tada	23	217	2	93	8	2-3
Sci. Coll.	10	do.	do.	Sept. —	do.	23	212	2	90	8	2-3
Tokyo.											
Do.	5	Male b.	Taipa, Formosa	Dec. —	do.	23	214	2	97	8	2-3

a P. 309.

b Description, p. 308; figs. 266-268.


ELAPHE RUFODORSATA ^a (Cantor).

1842. *Tropidonotus rufodorsatus* CANTOR, Zool. Chusan (pl. XIII) (type-locality, Chusan, China; types in Brit. Mus.; Cantor, collector).—*Coluber rufodorsatus* SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 225 (Tamsui, Formosa).—GUENTHER, Rept. Brit. India, 1864, p. 238, pl. xx, fig. G.—STRAUCH, Mém. Acad. Sci. St. Pétersb. (7), XXI, no. 4, 1873, pp. 79, 270 (Khinggan; High Gobi; Dauria; Possiet Bay).—MUELLER, Verh. Naturf. Ges. Basel, VII, Pt. 3, p. 681 (Tientsin).—BOERTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, pp. 70, 130 (Shanghai); Ber. Senckenberg. Naturf. Ges., 1894, p. 140 (Luechan Mts., Hankow); p. 144 (Lake Sitai, near Shanghai); Kat. Schl. Mus. Senckenberg., 1898, p. 50 (Chinhai, near Ningpo).—BOULENGER, Cat. Sn. Brit. Mus., II, 1894, p. 43 (eastern China, Formosa, Hainan).—WALL, Proc. Zool. Soc. London, 1903, p. 91 (China).—WERNER, Abh. Bayer. Akad. Wiss. (Muenich), II Klasse, XXII, Pt. 2, 1904, p. 356 (Shanghai; Hankow).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1 (1905), p. 248 (Possiet Bay; Korea).—*Ablabes rufodorsatus* BOULENGER, Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 138 (eastern Siberia; Peking).

^a Signifying red-backed.

1854. *Ablabes sex-lineatus* DUMÉRIL and BIBRON, *Erpét. Gén.*, VII. Pt. 1, p. 324 (type-locality, China; types in Paris Mus.).—*Coronella sexlineata* JAN, *Icon. Ophid.*, livr. 14, 1865, pl. vi, fig. 2 (China; type).
1886. *Simotes herzi* BOETTGER, *Zool. Anzeig.*, IX, 1886 p. 519, (type-locality, hills near Shanghai; type in Mus. Senckenberg.; Herz, collector).

Description.—*Adult female*; U.S.N.M. No. 21192; Seoul, Korea; October 8, 1883; P. L. Jouy, collector (figs. 269–271). Rostral very much broader than high, just visible from above; internasals small, very narrow in front, suture between them considerably shorter than between prefrontals, which are large and in contact with supraocular; frontal slightly longer than its distance from tip of snout, as long as interparietal suture, slightly broader in front than behind, not in contact with preocular; parietals long; nostril small, semicircular, situated a little above the center of the larger semidivided nasal; loreal large, somewhat longer than high; one preocular, not in contact with frontal; no subpreocular; two postoculars, upper larger



FIGS. 269–271.—*ELAPHE RUFODORSATA*. 2 X NAT. SIZE. 269, TOP OF HEAD; 270, SIDE OF HEAD; 271, UNDERSIDE OF HEAD. No. 21192, U.S.N.M.

than lower; temporals 2 + 3; seven supralabials, third and fourth in contact with eye (on left side eight, fourth and fifth in contact); five lower labials, in contact with anterior chin-shields; posterior chin-shields shorter than anterior; 21 rows of smooth scales; ventrals 177; anal divided; 51 pairs of subcaudals; no ventral lateral keel. Color (in alcohol), above olive gray, with four series of small dark brown spots, the inner two ocellated, the outer solid, which on the posterior half of the body join to form four continuous dark bands, extending to the tip of the tail; top of the head with a very distinct V-shaped mark, angle forward on the frontal and limbs extending far back on the neck, the edges of the mark strongly outlined by blackish brown; a similarly colored fleur-de-lis-shaped figure on the fronto-parietal suture; a blackish brown angular line on the prefrontals, continuing backward obliquely through the eye to angle of mouth and sides of

neck; underside whitish, ventrals outlined with black, and checkered with large square blotches; a black median line on the suture between the subcaudals.

Dimensions.

Total length.....	<i>mm.</i>
Snout to vent.....	561
Vent to tip of tail.....	470
	91

Variation.—The normal number of supralabials is seven, but occasionally eight are found; thus in both of our Korean specimens there are eight on one side, and Doctor Werner found a similar number among five Chinese examples. Twenty-one scale rows seem to be the constant number. Boulenger mentions that two preoculars occur rarely. In the Korean specimen in the museum of the Michigan University the temporals show the unusual number 1 + 2. Doctor Wall mentions that in 15 Chinese specimens examined by him only one had four lower labials in contact with anterior chin-shields. Ventrals range between 162 and 190, subcaudals between 47 and 68.

Habitat.—China, from Peking to Hainan, according to specimens in British Museum.^a It occurs also in eastern Siberia from Lake Baikal to Amurland and Ussuri, where it has been taken at Vladivostok and Possiet Bay, near the Korean boundary, while the United States National Museum has two specimens collected by the late Mr. P. L. Jouy at Seoul, Korea, October 8, 1883. Another specimen from Korea, collected by Mr. G. A. Olinger, is in the museum of the Michigan University, and a third in the St. Petersburg Museum.

Swinhoe obtained it at Tamsui, in Formosa, and sent the specimens to British Museum.

List of specimens of Elaphe rufodorsata.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Supralabials.	Temporals.
				1883.							
U. S. N. M.	21191	Male.....	Seoul, Korea.	Oct. 8	P. L. Jouy	21	169	2	56	7-8	2+2-3
Do.	21192	Female ^a	do.	do.	do.	21	177	2	51	7-8	2+3
Mich. Univ.	30526	Adult.....	Korea		G. A. Olinger.	21	167	2	60	7	1+2
Brit. Mus.	<i>m</i>	Female ^b	Formosa		R. Swinhoe.	21	183	2	54		
Do.	<i>n</i>	do ^a	do.		do.	21	178	2	49		

^a Description, p. 311; figs. 269-271.

^b Boulenger, Cat. 11, p. 44.

^a As for the locality Hainan, see under *Elaphe dione*, p. 318, footnote.

ELAPHE SCHRENCKII^a Strauch.

1873. *Elaphis schrenckii* STRAUCH, Mém. Acad. Sci. St. Pétersbourg (7), XXI, no. 4, pp. 100, 272 (type-locality, Khinggan Mil. Post, eastern Siberia; type, No. 3706 St. Petersburg, Acad. Mus.; Schrenck, collector).—*Coluber schrenckii* BOULENGER, Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 139 (Ussuri River; Brit. Mus.); Cat. Snakes Brit. Mus., II, 1894, p. 48 (Ussuri R.; Seoul, Korea).—WALL, Proc. Zool. Soc. London, 1903, pp. 92, 100 (Amurland; Korea; N. Japan).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 260 (Amurland; Korea, etc.).
1888. *Elaphis virgatus* BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., p. 72 (Korea; Mus. Senckenb.; Moellendorff, collector) (not of Schlegel).
1898. *Coluber schrencki* BOETTGER, Kat. Schl. Mus. Senckenberg., p. 51 (Korea).

Description.—*Male adult*; U.S.N.M. No. 21187; Seoul, Korea; August, 1887; P. L. Jouy, collector. Rostral much broader than high, visible from above; internasals longer than broad, much smaller than prefrontals, which are in contact with supraocular; frontal shorter than its distance from tip of snout, equaling interparietal suture, much wider anteriorly than behind, the sides strongly convergent backward; supraoculars narrower than frontal; parietals rather short, equaling frontal and half of interprefrontal suture; nostril large, round, between two subequal nasals; loreal longer than high; one preocular, not in contact with frontal; one small subpreocular (absent on left side); two postoculars (lower fused with fifth supralabial on right side); temporals 2 + 3; eight supralabials, seventh largest, fourth and fifth entering eye; four lower labials in contact with anterior chin-shields; posterior chin-shields shorter than anterior; 23 rows of scales, the median ones distinctly keeled, the keels becoming more and more indistinct on the sides, disappearing entirely on the outer four or five rows; 214 ventrals, with indistinct lateral angle; anal divided; subcaudals, 70 pairs. Color (in alcohol) above dark isabella-color, with scattered black specks occupying part of a scale; on the neck these spots become more numerous, and toward the tail the edges of the scales become dusky, with indications of longitudinal lines; head above like the back, with irregular dusky spots; supralabials whitish, except the last, all with black posterior edge; a black line obliquely from eye to edge of lip anterior to corner of mouth; indications of a pale black-edged line between nostril and eye; lower labials with dusky posterior edges; underside whitish, the anterior ventrals dark-edged.

^a For Dr. Leopold von Schrenck, the celebrated Russian naturalist and explorer of Amurland. He was born in the Kharkov government on April 24, 1826, and died in St. Petersburg on January 20, 1894, when director of the Anthropological-Ethnographical Museum of the Academy of Sciences. He traveled in eastern Siberia and in Sakhalin during 1854-1856.

Dimensions.

	<i>mm.</i>
Total length.....	1,580
Snout to vent.....	1,330
Vent to tip of tail.....	250

The *young* have a very characteristic head pattern. Thus, in the specimen in the Philadelphia Academy of Sciences (No. 15255), from Seoul, Korea, the ground color is brownish with white black-edged bands and spots, as follows: A broad band from upper postocular over temporals past the last supralabial, confluent with the white underside; a black line from lower postocular over suture between seventh and eighth supralabials to edge of lip; a white spot on frontal anteriorly, and a small one on each of the parietals near the frontal; an interrupted white cross band on supraoculars; a festooned white band across anterior half of prefrontals; a white spot on rostral; a white horizontal band from nostril to preocular; semicircular black markings on anterior supralabials; an inverted Y-shaped white mark on upper neck; back with rhomboidal brown black-edged spots separated by whitish cross bands.

Halfgrown specimens, such as U.S.N.M. No. 21189, show traces of this pattern, especially on the posterior part of the body and on sides of head; this specimen also has indication of a checker pattern on the underside.

Very old specimens are often nearly uniform pale isabella-color.

Variation.—Apparently the most variable feature of the scutellation of this species is the frequent absence of the subpreocular. Thus, out of 11 specimens Strauch found it lacking on both sides in five, while of my five Korean examples it is absent in three on one side. Only in one of our specimens is there a ninth supralabial on one side, while in the Michigan University specimen from Korea the number of supralabials is reduced to six on one side and seven on the other. The number of scale rows appears to be very constant, all the 19 specimens on record or examined by me having 23. The number of ventrals is also confined within narrow limits, namely, 208 to 226. The subcaudals do not exceed 76 and rarely go as low as 61. One specimen (U.S.N.M. No. 21186) has an undivided anal, and I also believe that it is undivided in the young specimen in the Philadelphia Academy of Sciences.

Habitat.—Eastern Asia from the River Argun, the tributary of the Amur, ranging east to the sea. The cotypes were collected in the Khinggan Mountains by Doctor Radde, according to Schrenck, but as I can not find that Radde ever visited the main body of this chain, which is situated in Chinese Manchuria, it is most likely that he obtained his specimens along the borders of the rivers Argun and Amur, which skirt the mountains mentioned. Dybowski collected it at Vladivostok and Posiet Bay, near the Korean frontier, and in Korea

itself it has been collected more recently by Moellendorff, Campbell, Ferebee and Jouy. It seems to be rather common in Korea, especially around Seoul whence Jouy brought us four specimens.

Strauch enumerates two specimens in the St. Petersburg Museum as having been collected by Mr. Goschkewitsch in Japan, and as this gentleman was for some little time Russian consul in Hakodate, Strauch assumes that they were collected in Yezo. The occurrence of this species in Japan is entirely unconfirmed and the probability is that the specimens really came from the mainland opposite, though possibly shipped from Japan. In this connection it is well to remember that *Elaphe dione* is also credited to Yezo by the same authority on material from the same source. Neither the one nor the other species has been found on Japanese territory by other collectors.^a

List of specimens of Elaphe schrenckii.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supralabials.
				1883.							
U. S. N. M.	14609	Male ^a	Korea		Dr. N. M. Ferebee.	23	212	2	62	2+3	8
Do.	21186	Female	Seoul, Korea.	Oct 16	P. L. Jouy	23	226	1	2+3	8-9	
Do.	21187	Male ^b	do	Aug. —	do	23	214	2	70	2+3	8
Do.	21188	Female	do	do	do	23	218	2	72	2+3	8
Do.	21189	Male	do	do	do	23	225	2	70	2+3	8
				1890.							
Phila. Acad.	15255	Young	do	Sept. —	Dr. W. H. Jones	23			70	2+3	7-8
Mich. Univ.	30529	Adult	Korea		G. A. Olinger	23	219	2	72	2+2	6-7
Senckenberg		(c)	do		Moellendorff	23	226	2	70		
Brit. Mus.	b	Young ^d	Seoul, Korea.		C. W. Campbell	23	210	2	68		
Do.	c	Male ^e	do		do	23	216	2	76		
Do.	a	Female ^d	Ussuri River.			23	220	2	68		
St. Petersburg	2907	(f)	Possiet Bay.			23	209	2	70		
Do.	2955	(f)	[Japan?]		Goschkewitsch	23	219	2			
Do.	2956	(f)	do		do	23	211	2	71		

^a Total length, 1,635 mm.
^b Description, p. 313.

^c Boettger.
^d Boulenger, Cat. II, p. 48.

^e Boulenger, p. 49.
^f Strauch, p. 272.

ELAPHE DIONE ^b (Pallas).

1773. *Coluber dione* PALLAS, Reise Russ. Reichs, II, p. 717 (type-locality, "Salt steppes toward the Caspian Sea").—BOULENGER, Ann. Mag. Nat. Hist. (6), V, Feb., 1890, p. 138 (Ussuri River; Peking); Cat. Sn. Brit. Mus., II, 1894, p. 44 (southern Russia to Japan).—BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 149 (Chefu, China); Kat. Schl. Mus. Senckenberg., 1898, p. 51 (Chefu; Turkestan).—STONE, Proc. Phila. Acad., 1899, p. 184

^a See also under *Agkistrodon blomhoffii*, p. 452.

^b From *Διώνη*, a name in Greek mythology, the mother of Aphrodite.

- (northern Pechili, China).—MÉHELY, Zichy's Dritte Asiat. Forschungsr., II, 1901, p. 58 (Peking).—WALL, Proc. Zool. Soc. London, 1903, p. 91 (China).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, No. 1, 1905, p. 249 (Vladivostok, Korea, etc.).—*Elaphis dione* DUMÉRIL and BIBRON, Erpét. Gén., VII, Pt. 1, p. 248.—GUENTHER, Rept. Brit. India, 1864, p. 240 (Kaukasus; Peking).—JAN, Icon. Ophid., livr. 21, 1867, pl. m, fig. A (Sarepta, Russia).—STRAUCH, Mém. Acad. Sci. St. Pétersb. (7), XXI, No. 4, 1873, pp. 82, 270 (eastern Russia, Siberia, Amur, Ussuri, Korea, Japan); in Przevalski's Mongoliya i Strana Tangutov, III, 1876, p. 46 (Ordos).—MOELLENDORFF, Journ. N. China Br. R. Asiat. Soc. (n. s.), XI, 1877, p. 104 (Peking; Manchuria).—BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 131 (East Europe to North China and Korea).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 356 (Tsingtau, China; Peking).
1802. *Coluber diana* LATREILLE, Hist. Nat. Rept., IV, p. 159 (emendation).
 1831. *Coluber eremita* EICHWALD, Zool. Special., III (p. 174).
 1837. *Coluber macoticus* RATHKE, Mém. Sav. Étrang. Acad. St. Pétersb., III, p. 433, pl. 1, figs. 9-11 (type-locality, Caspian Sea at the mouth of Riv. Ural; Goebel, collector).

Description.—*Adult male*; U.S.N.M. No. 14610; Korea; Dr. N. M. Ferebee, collector. Rostral much broader than high, visible from above; internasals much broader than long, only half as long as prefrontals which are broadly in contact with supraocular; frontal nearly as long as its distance from tip of snout, somewhat wider in front than behind; parietals longer than frontal, interparietal suture longer than distance of frontal from rostral; nostril large, between two nasals, the anterior slightly larger than the posterior; loreal longer than high; one preocular, not in contact with frontal; one subpreocular; two postoculars; temporals 2 + 3; eight supralabials, seventh largest, fourth and fifth entering eye; five lower labials in contact with anterior chin-shields, which are as long as the posterior; 25 rows of nearly smooth scales; 204 ventrals; anal divided; 65 pairs of subcaudals. Color (in alcohol) light tawny-olive above, with numerous irregular narrow dusky crossbars, about one to one and a half scale wide; head with slightly darker, black-edged markings consisting of an irregular chevron mark on parietals and upper neck, a cross mark on prefrontals and frontal extending backward obliquely through eye and reaching commissure at the angle of mouth; supralabials, including lower edge of seventh and eighth, pale yellow; underside pale yellowish with irregular dusky spots which near the obtuse lateral angle of the ventrals assume the character of an interrupted longitudinal line.

Dimensions.

	<i>mm.</i>
Total length.....	825
Snout to vent.....	677
Vent to tip of tail.....	148

In the *young* (No. 21193) (fig. 272) the markings are darker and more distinct; the pattern on the head shows a fleur-de-lis figure on the interparietal suture; the lateral dark spots on the underside are blacker, broader, with a convex anterior outline; there are three pale dorsal lines, the median one not well defined and interrupted by the dark crossbars, the lateral ones on the fifth, sixth, and seventh scale rows continuous and cutting off a series of blackish lateral spots from the dorsal dark crossbars.

This pattern is often retained in tolerably large specimens.

Variation.—The variation in our Korean material is quite insignificant; thus all possess a subpreocular, but, as might be expected in a species of so wide distribution, many anomalies are found when a large series is studied. Strauch examined 84 specimens from nearly the entire range, and has enumerated the deviations he found. Thus five specimens lacked the subpreocular on both sides and five on one side; thirteen had three postoculars on both sides and six had three on one side; four specimens had nine supralabials on both sides and two specimens only seven on both sides, while eleven had nine or seven on one side only. The number of scale rows varies in about even proportions between 23 and 25 in 43 specimens.^a As to the subcaudals, Strauch mentions as a curious anomaly only found in some east Siberian and west Chinese specimens that some of the subcaudals are entire.

The number of ventrals varies between 172 and 214. In this a slight tendency to geographical variation can be traced. Thus in 42 specimens from Siberia the ventrals vary between 180 and 214 in those west of the Yenisei River, but between 172 and 195 in those from east of that river. In 17 Chinese specimens they vary again between 180 and 213 ventrals, while in five Korean specimens the number is 203–207.

One of our specimens (No. 21190) shows the anomaly of having on both sides a small shield cut off from the third supralabial between it and the loreal. Another (No. 14331) has two presuboculars on each side.

Habitat.—A species of extraordinarily wide distribution, ranging, as it does, from the valley of the Volga in southeastern Russia through




FIG. 272.—ELAPHE
DIONE. 2 X NAT.
SIZE. TOP OF
HEAD. No. 21193,
U.S.N.M.

^a Curiously enough, in 21 specimens in British Museum enumerated by Boulenger, of which 12 were from China, from which country Strauch had no specimens, 13 have 25 scale rows, 6 have 27, and only 2 have 23. All our Korean specimens have 25, and I am inclined to regard this as the normal number. This agrees with Doctor Wall, who found 23 in 3, 25 in 10, and 27 in 1 Chinese specimens, and with Doctor Werner, who in 4 specimens from China found 3 with 25 and 1 with 27 scale rows, the latter in a specimen from Tsingtau.

temperate Asia to the Amur country in the north and south through eastern China to Kiukiang, and possibly even to the island of Hainan.^a

Dybowski, according to Strauch, collected it on the banks of the Amur between Seljonoï-Less and Khabarovka, and also at Posiet Bay. It is also recorded from the Ussuri, the Sungari, and upper Amur, and the Hamburg Museum has a specimen each from Vladivostok and Pokrovka, collected by Dieckmann (Nos. 2168, 2169). The St. Petersburg Museum also has a specimen from Vladivostok, collected by Doctor Zander.

Schrenck and Doctor Ferebee obtained specimens in Korea and Jouy brought home three specimens from Seoul.

Two specimens obtained in 1861 by the St. Petersburg Academy from the Russian consul in Hakodate, Mr. Goschkewitsch, are credited by Strauch to Japan, but, as explained under the head of *Elaphe schrenckii*, I do not place any faith in their having been collected in any of the Japanese islands.

List of specimens of Elaphe diene.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supralabials.
U.S.N.M.	14321	Young.	Ala Tan, Siberia.		Mus.St. Petersburg.	25	207	2	61	3+4	8
Do.	14331	Female ^a	Amur land		do.	25	192	2	50	2+4	8
Do.	14610	Male ^b	Korea.		Dr. N. M. Ferebee.	25	204	2	65.	2+3	8
				1883.							
Do.	21190	Female ^a	Seoul, Korea.	Oct. 8	P. L. Jouy	25	205	2		2+3	8
Do.	21193	Young ^c	do.	June, —	do.	25	207	2	71	2+3	8
Do.	21194	Female	do.	Aug. —	do.	25	204	2	68	2+2-3	8-9
Brit. Mus.	h.	Male ^d	Ussuri River		Warsaw Mus.	23	179	2	60		
St. Petersburg.	2910	(c)	Bikinda, Ussuri.			25	183	2			
Do.	2911	(c)	Dshare, Amur			25	195	2	61		
Do.	2953	(c)	[Japan?]		Goschkewitsch	23	194	2	58		
Do.	2954	(c)	do.		do.	23	180	2	62		
Do.	3704	(c)	Korea.			23	203	2	59		

^a P. 317. ^b Description, p. 316. ^c Fig. 272. ^d Boulenger, Cat. II, p. 45. ^e Strauch, p. 271.

^a This locality is given on the authority of a specimen in British Museum said to have been collected at Hojhow by J. Neumann. It should be noted in this connection, however, that several rather more northern forms have been attributed to the same locality upon material received from the same source, such as *Natrix tigrina*, *Dinodon rufosonatus*, *Zamenis spinalis*, *Elaphe rufodorsata*, *Agkistrodon blomhoffii*. None of these species have been found in that island by other collectors (see BOETTGER, *Liste der von der Insel Hainan bekannten Kriechthiere*, in Ber. Senckenberg. Naturf. Ges., 1894, pp. 129-136.—COPE, On a Collection of Batrachia and Reptilia from the Island of Hainan, in Proc. Phila. Acad., 1894, pp. 423-428.—BOULENGER, On the Reptiles, Batrachians, and Fishes collected by the late Mr. John Whitehead in the Interior of Hainan, in Proc. Zool. Soc. London, 1899, pp. 956-962, pls. LXVI-LXIX), and the suggestion that these northern species might have come from the mountainous interior is not supported by the result of Whitehead's expedition. It seems most conservative, therefore, to wait for corroboration before the island of Hainan be given as an undoubted locality for those species.

ELAPHE TÆNIURUS^a Cope.

1858. *Elaphis virgatus* GUENTHER, Cat. Colubr. Sn. Brit. Mus., p. 95 (part: specimen c, Chikiang, China) (not of Schlegel).
1860. *Elaphe tæniurus* COPE, Proc. Phila. Acad., 1860, p. 565 (type-locality, Ningpo, China, type in Phila. Acad. Mus.; McCartee, collector).—*Elaphis tæniurus* GUENTHER, Rept. Brit. India, 1864, p. 242 (Chikiang; Ningpo).—STRAUCH, Mém. Acad. Sci. St. Pétersb. (7), XXI, no. 4, 1873, pp. 103, 272 (Possiet Bay).—MOELLENDORFF, Journ. N. China Br. R. Asiat. Soc. (n. s.), XI, 1877, p. 104 (Tachiosse, NW. of Peking).—BOULENGER, Ann. Mag. Nat. Hist. (5), XIX, 1887, p. 170 (part: Chikiang; Shanghai; Peking).—BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, pp. 71, 131 (Shanghai; Formosa).—*Coluber tæniurus* BOULENGER, Fauna Brit. India. Rept., 1890, p. 333 (part: Manchuria; China; Darjeeling); Ann. Mag. Nat. Hist. (6), V, Feb. 1900, p. 139 (hills northwest of Peking); Cat. Snakes Brit. Mus., II, 1894, p. 47 (part: western hills of Peking; Shanghai; Chikiang; mountains of Kiukiang; Darjeeling).—BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 138 (South Cape, Formosa); p. 144 (Wusung near Shanghai); Kat. Schl. Mus. Senckenberg., 1898, p. 51 (Shanghai; Formosa).—WALL, Proc. Zool. Soc. London, 1903, p. 92 (Yangtze Valley; Shanghai).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 259 (Possiet Bay; Novgorodsk).
1879. *Elaphis yunnanensis* ANDERSON, Zool. Res. Yunnan Exped., p. 813 (type-locality, Momien, Yunnan).

Several forms of varying degree of distinctness seem to cluster around *E. tæniurus*^b. The typical form which extends from the Himalayas over China and Korea to the Amur Province is characterized by having 23-25 scale rows, 226-258 ventrals, and 90-111 subcaudals, a formula derived from about 20 specimens. A slightly differentiated form with 25-27 scale rows, 251-260 ventrals, and 104-124 subcaudals inhabits the southern group of the Riu Kiu Islands, as demonstrated by 5 specimens examined by me. This is Boettger's *Coluber schmackeri*. Finally, the form which Fischer called *E. grabowskyi*^c seems clearly entitled to recognition, there being a considerable gap between typical *E. tæniurus* and the specimens

^a From *ραϊνία*, band; *ὄψα*, tail.

^b *Coluber tæniurus ridleyi* BUTLER, Journ. Bombay Nat. Hist. Soc., XII, 1899 (p. 426), is unknown to me, as that publication is not now received in Washington.


^c 1885. *Elaphis grabowskyi* FISCHER, Archiv Naturg., LI, Pt. 1, 1885, p. 59 (author's separate p. 19), pl. IV, figs. 3a-e (type-locality, Batu Hapu, Pengaron, southeast Borneo; type in British Museum; Grabowski, collector).

1887. *Elaphis tæniurus* BOULENGER, Ann. Mag. Nat. Hist. (5), XIX, p. 170 (part: Pajo, Sumatra).—*Coluber tæniurus* BOULENGER, Cat. Sn. Brit. Mus., II, 1894, p. 47 (part: SE. Borneo; Sumatra).—FLOWER, Proc. Zool. Soc. London, 1899, p. 668 (Kuala Lumpur, Malay peninsula).

I have added the last reference on the strength of a specimen in the United States National Museum (No. 26234) collected by Dr. W. L. Abbott in Trong, Lower Siam. It is undoubtedly an *E. grabowskyi* with 25 scale rows, 295 ventrals, and 111 subcaudals; 9 supralabials, and preoculars *broadly* in contact with frontal. Mr. Flower gives no scale formulas of his specimens, but it seems highly probable that they are the same as Abbott's.

from Borneo, Sumatra, and the Malay peninsula so far as the scale formula is concerned, which in four specimens is as follows: 25 scale rows, 278-295 ventrals, 94-111 subcaudals.^a

Description.—*Adult male*; U.S.N.M. No. 14608; Korea; Dr. N. M. Ferebee, collector (figs. 273, 274). Rostral slightly broader than high, largely visible from above; internasals longer than broad, half as large as prefrontals, which are narrowly in contact with supraocular; frontal equaling its distance from rostral, a little shorter than interparietal suture, somewhat broader in front than behind; parietals longer than the distance of frontal from tip of snout, strongly convergent behind; nostril rounded, large, between two nasals, the posterior slightly larger than the anterior; loreal lozenge-shaped, longer than high; one preocular, not in contact with frontal; one small subpreocular; two



FIGS. 273-274.—*ELAPHE TENUIRUS*. $1\frac{1}{2} \times$ NAT. SIZE. 273, TOP OF HEAD; 274, SIDE OF HEAD. No. 14608, U.S.N.M.

postoculars; temporals 2 + 3 (on left side 2 + 4); eight supralabials, posterior ones not specially enlarged, fourth and fifth entering eye; six lower labials in contact with anterior chin-shields (on left side seven), first pair rather elongate; chin-shields subequal; 25 rows of scales, of which the outer seven on each side are smooth or nearly so, the others feebly but distinctly keeled; 229 ventrals, with a distinct lateral angle; anal divided; subcaudals, 111 pairs. Color (in alcohol) pale olive clay-color above; on the middle of the back a black rope-ladder pattern, consisting of two narrow lateral black lines at regular intervals connected by a black cross line of approximately the same width, this pattern gradually disappearing on the posterior third of the body; on the sides two or three ill-defined series of irregular black spots which on the posterior third of the body gradually consolidate

^a Two of the specimens, the scale formulas of which are here included, viz, Doctor Fischer's two original specimens, now in British Museum, have respectively 278 and 282 ventrals and 98 and 94 subcaudals according to Boulenger's count. Fischer himself, in the original description, counted 281-284 ventrals and 101-104 subcaudals.

into a broad (about 6 scale rows) longitudinal blackish band with vertical series of white spots about four scales apart; underside pale yellowish with a dusky spot on each ventral at the lateral angle, these spots posteriorly consolidating into a broad black longitudinal band separated from the lateral broad band by a white one which occupies outer scale-row and the adjacent portion of the upturned ventral; these bands are continued on the tail to the tip, so that the tail is black with four whitish or yellowish bands, one on the median line underneath, one on the median line above, and one on each side; head above uniform clay-color with a broad, horizontal postocular black band which ends above the angle of the mouth but does not reach it; a small dusky spot on preocular; labials and throat uniform unspotted pale yellowish.

Dimensions.

	<i>mm.</i>
Total length.....	1.345
Snout to vent.....	1.030
Vent to tip of tail.....	315

The *young* are essentially colored like the above, except that the markings are more sharply defined; the vertical light marks on the lateral black band are suffused with brownish; top of head without any marks whatever.

Variation.—This species, which appears to be so constant in coloration, on the other hand is rather variable in its scutellation. Thus of about 22 specimens three are recorded as having an undivided anal; the temporals are rather variable, from 1+2 to 2+4; nine supralabials are sometimes noted; Doctor Wall records a specimen in which the loreals were confluent with prefrontals on both sides. As stated before, the number of ventrals ranges between 226 and 258, and the subcaudals between 90 and 111. The majority of specimens recorded seems to have 25 scale rows, but in about 40 per cent the number is 23.

The relation of the preocular to frontal and consequently the prefrontal to supraocular is also variable. As a rule, I believe the latter are in contact, but in one of our specimens (No. 31718) they are separated by the preocular being broadly in contact with the frontal.

Habitat.—Ranging from Darjeeling in the west to Formosa and the Amur Province in the east, this species has the regular Himalayo-Chinese distribution. The extreme northeastern point from which it has been recorded is Possiet Bay, in the Amur Province near the Korean boundary, but it seems to be common in the mountains west and northwest of Peking. A fine large specimen is in the U. S. National Museum (No. 14608) collected in Korea by Doctor Ferebee.

Doctor Boettger reports two specimens in Museum Senckenbergianum from Formosa, though without recording their scale formula. However, as he is the author of *E. schmackeri* and also had specimens from Shanghai for comparison the determination can not well be doubted.

List of specimens of Elaphe tæniurus.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supracaudals.
U.S.N.M.	7499	Young.	China.....		W. Stirpison	25	230	2	95	1+2	8-9
Do.	14608	Male ^a	Korea.....		Dr. N.M. Ferebee	25	227	2	111	2+3-4	8
Do.	31718	Young.	Shanghai, China.....		E. Deschamps	25	233	2	103	2+3	8-9
St. Petersburg.	3192	(b)	Possiet Bay.....		Butakow	23	231	2	98		

^a Description, p. 320; figs. 273-274.

^b Strauch, p. 272.

ELAPHE SCHMACKERI^a (Boettger).

1895. *Coluber schmackeri* BOETTGER, Zool. Anz., XVIII, 8 July, 1895, p. 268 (type-locality, Mijakoshima, Sake shima group, Riu Kiu.; type in Mus. Senckenberg.; Schmacker, collector); Kat. Schl. Mus. Senckenberg., 1898, p. 53.—BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 627.—WALL, Proc. Zool. Soc. London, 1903, p. 101 (Loo Choos); 1905, II, p. 515 (Ishigaki).

Notwithstanding the fact that its original describer, Doctor Boettger, compares this form with *E. moellendorffi*, I can only regard it as an offshoot of *E. tæniurus*. In fact I have hesitated long whether it would not be most consistent to call it *Elaphe tæniurus schmackeri*, as the scale formulas plainly intergrade. However, as long as my colleagues are satisfied with the binominal appellation I can have no objection.

This species, which in Iriomote shima is known as "Ki-tokara" according to Doctor Lenz, is so closely related to *Elaphe tæniurus* that a separate description seems unnecessary. The coloration is identical. The main differences in scutellation are the greater number of scale rows, ventrals, and subcaudals, and the somewhat feebler carination of the dorsal scales. Thanks to the kindness of Doctor Boettger I have been able to examine his type specimen in Museum Senckenbergianum and satisfied myself that it has keels, though very faint ones, on the median scales. In the Hamburg Museum specimen I found very weak keels on the median scales while about eight lateral rows on each side are practically smooth, and in the specimen before

^a For the late Mr. B. Schmacker, of Shanghai.


me now (No. 5, Sci. Coll. Tokyo, figs. 275-276) the outer nine scale rows are smooth, but the seven middle ones have distinct though feeble keels. This specimen measures as follows:

<i>Dimensions.</i>		<i>mm.</i>
Total length.....		1,870
Snout to vent.....		1,445
Vent to tip of tail.....		425

The type as recorded by Doctor Boettger measures 2,080 mm. in total length, of which 445 belong to the tail. This species consequently grows to a considerable size.

Variation.—Very much as in *Elaphe tæniurus*, though the range between the extremes of the number of ventrals and subcaudals is much less, as might be expected from the much more restricted habitat.

In most specimens the preocular is not in contact with the frontal, but in the Hamburg Museum specimen these two shields touch on



FIGS. 275-276.—ELAPHE SCHMACKERI. NAT. SIZE. 275, TOP OF HEAD; 276, SIDE OF HEAD. No. 5, Sci. Coll. Tokyo.

the right side, and in the specimen before me (No. 5) they practically touch on both sides. In the type, on the contrary, they are widely separated.

A single anal is found in No. 89, Science College Museum, and in this the lower temporals of both rows are consolidated so as to make only 2 + 2 temporals, but normally they are 2 + 3.

In the Hamburg Museum specimen there are 10 supralabials and the Ishigaki shima specimen recorded by Dr. Wall has ten on one side.

Habitat.—So far as known, this form of *E. tæniurus* is confined to the southern group of the Riu Kius. The type in Museum Senckenbergianum is supposed to be from Miyakoshima, from which island

there is also a large specimen with mutilated tail in the Imperial Museum, Ueno Park, Tokyo (No. 6), and another in Science College Museum, Tokyo (No. 5). Another large specimen (No. 89) is in the same museum, collected by Mr. Tashiro in Iriomote shima, and a specimen from the same island in the Hamburg Museum (No. 2590) collected by Doctor Lenz on March 13, 1897, was obtained in the mountains of the interior, according to a note attached to it. Dr. Wall has recently recorded a specimen from Ishigaki shima.

List of specimens of Elaphe schmackeri.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supracaudals.
Sci. Coll., Tokyo.	5	Male ^a .	Miyako shima			25	251	2	117		9
Imp. Nat. Tokyo.	6	Adult	do.			27	254	2		2+2-3	9
Sci. Coll., Tokyo.	89	do	Iriomote shima.	1888. June 14	Tashiro	25	259	1	118	2+3	9
Hamburg	2590		do.	1897. Mar. 13	Doctor Lenz	27	254	2	124	2+3	10
Senckenberg	8056.2 ^a	Male ^b	Miyakoshima		B. Schmacker.	27	260	2	104		

^a P. 323, figs. 275-276.

^b Type.


ELAPHE CLIMACOPHORA ^a (Boie).

AODAISHŌ.

1826. *Coluber climacophorus* BOIE, Isis, 1826, p. 210, (type-locality, Japan; Blomhoff, collector).—BOULENGER, Cat. Sn. Brit. Mus., II, 1894, p. 54 (Nagasaki; Nikko; Miyanoshta; Mt. Onsen, Shimabara).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 52 (Hakone Mts.; Miyanoshta; Nikko).—WALL, Proc. Zool. Soc. London, 1903, p. 100 (Nikko); 1905, II, p. 514 (Hondo).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 356 (Tsingtau and Hankow, China).
1837. *Coluber virgatus* SCHLEGEL, Phys. Serp., I, p. 146; II, p. 145 (type-locality, Japan; types in Leiden Mus.; Buerger and Siebold, collectors) (not of Bonnaterre, 1790).—SCHLEGEL and TEMMINCK, Fauna Japon., Rept., 1837, pp. 83, 139 (Japan).—*Elaphis virgatus* DUMÉRILO and BIBRON, Exp. Gén., VII, Pt. 1, 1854, p. 261 (Japan).—BLEEKER, Natuurk. Tijdschr. Nederland. Indië, XVI, 1858, p. 204 (Japan).—JAN, Icon. Ophid., livr. 21, 1867, pl. 1 (Japan).—HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 113 (Tokyo).—OKADA, Cat. Vert. Japan, 1891, p. 68 (Tokyo; Osaka; Awaji; Nagato; Yezo).—WALL, Journ. Bombay Soc. Nat. Hist., XIV, 1902, p. 375 (Arima).—*Elaphis virgata* MARTENS, Preuss. Exped. Ost-Asiens, Zool., I, 1866, p. 111; 1876, p. 379 (Yokohama).—HILGENDORF, Mitth. Deutsch. Ges. Ost-Asiens, I, heft 10, 1876, p. 29 (Tokyo).—FRITZE, Mitth. Deutsch. Ges. Ost-Asiens, V, heft 46, 1891, p. 238 (Yezo).
1837. *Coluber quadrivirgatus* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., Ophid. pl. II (error; not of the text).

^a From κλίμαξ, ladder, φορος, carrying; referring to the pattern on the back.

Description (figs. 277-278).—*Adult male*; U.S.N.M. No. 31874; Kochi, Shikoku; May, 1902; Dr. H. M. Smith, collector. Rostral visible from above, much broader than high; internasals about as broad as long, smaller than prefrontals which are in contact with supraocular; frontal as long as its distance from rostral and as the interparietal suture, much broader in front than behind; parietals longer than distance of rostral from tip of snout, much narrowed and pointed behind; nostril large, rounded, between two subequal nasals; loreal longer than high; one preocular, not in contact with frontal; one subpreocular; two postoculars; temporals 2 + 3; supralabials eight, seventh largest, fourth and fifth entering eye; four lower labials in contact with anterior chin-shields, which are of about the same length as the posterior; 23 rows of scales, most of which are distinctly



FIGS. 277-278.—*ELAPHE CLIMACOPHORA*. 2 × NAT. SIZE. 277, TOP OF HEAD; 278, SIDE OF HEAD. NO. 36551, U.S.N.M.

keeled, but outer two rows, at least, perfectly smooth; 228 ventrals; anal divided; 111 pairs of subcaudals; the ventrals angularly bent laterally. Color (in alcohol) dark umber brown above with four somewhat obscure darker longitudinal bands, the lower ones occupying four scale rows, viz, second to fifth, the upper ones three rows, viz, eighth, ninth, and tenth; head uniform, with a blackish oblique band from eye backward to the edge of the lip crossing and occupying the whole of the last supralabial; other supralabials pale, lower labials and throat whitish; rest of underside bluish slate gray, the ventrals pale-edged behind and with the lateral angle marked by a narrow longitudinal line.

Dimensions.

	<i>mm.</i>
Total length.....	1,365
Snout to vent	1,040
Vent to tip of tail.....	325

The *young*, for instance No. 34050, have no longitudinal dark bands; on the back there are numerous brown transverse, black-edged spots, separated by whitish cross bands and alternating on the sides with a series of brown spots; whole underside whitish, posteriorly with dark edges to the ventrals; an oblique blackish band from eye backward crossing the last supralabial, but otherwise no pattern on head; supralabials narrowly edged with dusky.

Variation.—Only one specimen with an undivided anal is recorded. The subpreocular is more variable; it is absent on one side in our No. 34003; in the Hamburg Museum No. 2427 it is lacking on both sides, though indicated on the right side by a minute granule; Doctor Wall has found it absent in one specimen in a lot of eleven; Hilgendorf in 16 specimens found it absent in one on both sides and on one side in one. He also found three postoculars on both sides in two specimens and on one side in one specimen, while one postocular only was present once on both sides and twice on one side. Doctor Wall records two specimen, with three anterior temporals on both sides. In all specimens examined by me the prefrontals are in contact with supraocular. The two specimens from Shikoku have the parietals unusually pointed behind. In the young the frontal is longer relative to its distance from tip of snout.

The normal number of scale rows appears to be 23, but 25 per cent of the specimens recorded below have 25. Ventrals range between 224 and 244, subcaudals between 97 and 122.

Habitat.—Formerly believed to be confined to Japan proper where it appears to be common from Kiusiu to Yezo, there being numerous specimens from various localities in the museums.^a Doctor Werner has recently recorded two specimens in the Munich Zoological Museum said to have been collected by Doctor Haberer in Tsingtau and Hankow, China. It should be remembered, however, that he also collected in Japan and that he has credited two more Japanese species to China, which have not been obtained there by any other collector (see pp. 331, 336).

There are two specimens (No. 7) in the Imperial Museum, Ueno Park, Tokyo, which are said to be from Okinawa shima, but the locality given is almost certainly erroneous.

^a Tokyo; Mount Fuji; Yokohama; Kochi, Shikoku (U. S. Nat. Mus.). Mount Onsen, Shimabara, Kiusiu; Nagasaki; Nikko; Miyanoshta (Brit. Mus.). Tokyo; Nikko (Sci. Coll. Tokyo). Hakone Mts; Miyanoshta.; Nikko (Mus. Senckenberg.). Nagasaki; Yamato; Kawachi (Hamburg Mus.). Nagasaki (Stockholm Mus.).

List of specimens of Elaphe climacophora.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supraabials.
U.S.N.M.	4678	Halfgr.	Japan		Doctor Morrow.	25	228	2	102	2+3	8
				1903.							
Do.	31872	Male	Kochi, Shikoku.	May	Dr. H. M. Smith.	25	234	2	107	2+3	8
Do.	31874	do. ^a	do.	do.	do.	23	228	2	111	2+3	8
				1902.							
Do.	34002	Female	Yokohama	May 2	A. Owston	23	236	2	103	2+2	8-9
Do.	34003	Male	do.	do.	do.	23	226	2	99	2+3	8
				1898.							
Do.	34027	Young	Mount Fuji	Aug.	do.	23	224	2	101	2+3	8
Do.	34050	do.	do.	do.	do.	23	226	2	110	3-2+3	8
Do.	36551	do. ^b	Tokyo?		Sci. Coll. Tokyo.	25	231	2	102	2+2	8
				1896.							
Hamburg	2427	Male	Kawachi	Jan.	T. Lenz	23	236	2	99		8
Leiden	386	(c)	Japan		von Siebold.		230		114		
Do.	390		do.	1861.	do.		233		98		
Imp. Mus. Tokyo.	7		[Okinawa ? ?]			23	231	2		2+3	8
Brit. Mus.	a	Female ^d	Japan		Leiden Mus.	23	231	2	100		
Do.	b	Halfgr ^d	do.		do.	23	234	2	112		
Do.	c	Young ^d	do.		Sir R. Owen	23	222	2	108		
Do.	d	do. ^d	do.		Mr. Adams	23	232	1			
Do.	e	Female ^d	Nagasaki		Mr. Whitely	23	234	2	103		
Do.	f	Young ^d	do.		do.	23	235	2	116		
Do.	g	do. ^e	Nikko		do.	23	229	2	103		
Do.	h	do. ^e	Miyanoshta		J. H. Leech	25	232	2	100		
Do.	i	do. ^e	do.		do.	23	226	2	97		
Do.	k	Male	Mount Onsen, Shimabara.		P. Holst	25	234	2	99		

^a Description, p. 325.

^b Figs. 277-278.

^c Type of *C. virgatus*.

^d Boulenger, Cat. II, p. 54.

^e Boulenger, Cat. II, p. 55.

ELAPHE QUADRIVIRGATA ^a (Boie).

SHIMAHEBI.

Plate XXI.

1826. *Coluber quadrivirgatus* BOIE, Isis, 1826, p. 209, (type-locality, Japan; Blomhoff, collector); Bijdr. Natuurk. Wetensch., II, Pt. 1, 1827, p. 256.—SCHLEGEL, Phys. Serp., 1837, I, p. 146; II, p. 147, pl. v, figs. 15, 26 (Japan).—TEMMINCK and SCHLEGEL, Fauna Jap., Rept., 1837, pp. 84, 139 (Japan).—BOULENGER, Cat. Sn. Brit. Mus., II, 1894, p. 59 (Yokohama; Nikko; Miyanoshta; Kumamoto; Satsuma; Gensan, Korea).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 53 (Nikko; Yezo).—WALL, Proc. Zool. Soc. London, 1903, p. 100 (Japan); 1905, II, p. 514 (Hondo; Yaku Is.).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 357 (Hankow, China).—NIKOLSKI, Zap. Imp. Akad. Nauk,


^a Signifying four-striped.

- S. Peterburg (8), XVII, No. 1, 1905, p. 263 (Japan: Unzen; Hongkong).—*Compsosoma quadrivirgatum* DUMÉRIE and BIBRON, *Erpét. Gén.*, VII, Pt. 1, 1854, p. 301 (Japan).—*Elaphis quadrivirgatus* GUENTHER, *Cat. Colubr. Sn. Brit. Mus.*, 1858, p. 94 (Japan).—JAN, *Icon. Ophid.*, livr. 24, 1867, pl. I, fig. 1 (Japan).—HILGENDORF, *Sitz. Ber. Ges. Naturf. Fr. Berlin*, 1880, p. 114 (Hakodadi; Tokyo; Nambu, NE. Hondo).—OKADA, *Cat. Vert. Japan*, 1891, p. 68 (Tokyo; Hakone; Nii shima; Koze shima; Awaji; Suwo; Yezo).—*Elaphis quadrivirgata* MARTENS, *Preuss. Exped. Ost-Asien, Zool.*, I, 1866, p. 112 (Nagasaki); 1876, p. 379 (Yokohama).—HILGENDORF, *Mitth. Deutsch. Ges. Ost-Asiens*, I, heft 10, 1876, p. 30 (Yezo; Tokyo).—FRITZE, *Mitth. Deutsch. Ges. Ost-Asiens*, V, heft 46, 1891, p. 238 (Yezo).
1826. *Coluber vulneratus* BOIE, *Isis*, 1826, p. 212 (type-locality, Japan: Blomhoff, collector).
1837. *Coluber virgatus* TEMMINCK and SCHLEGEL, *Fauna Japon.*, Rept., Ophid., pl. 1^a (error; not of text).
1860. *Elaphis bilineatus* HALLOWELL, *Proc. Phila. Acad.*, 1860, p. 497 (type-locality, Volcano Bay, Yezo; Doctor Morrow, collector).
1860. *Leptophidium dorsale* HALLOWELL, *Proc. Phila. Acad.*, 1860, p. 498 (type-locality, Hakodate, Yezo; Stimpson, collector).
1867. *Elaphis quadrivirgatus* var. *interrupta* JAN, *Icon. Ophid.*, livr. 24, pl. I, fig. 2, cover p. 2 (type-locality, Japan; type in Mus. Milano).
1867. *Elaphis quadrivirgatus* var. *atra* JAN, *Icon. Ophid.*, livr. 24, pl. I, fig. 3, cover p. 2 (type-locality, Japan; type in Mus. Milano).—*Coluber quadrivirgatus* var. *atra* BOETTGER, *Kat. Schl. Mus. Senckenberg.*, 1898, p. 53 (Japan).

I have no doubt that *Leptophidium dorsale*, HALLOWELL is correctly referred here as a synonym. The type seems to be lost, but I have before me a young *E. quadrivirgata* from Yezo (U.S.N.M. No. 23435), which agrees with Hallowell's description in every particular, except in the number of ventrals (100, which of course is erroneous and probably a misprint for 200) and in the shape of the pupil which Hallowell says is "ovoid," but the latter statement is easily accounted for.

Description (figs. 279–281).—*Adult female*: U.S.N.M. No. 31819; Yamagawa, province of Satsuma, Kiusiu; June 14, 1903; Dr. Hugh M. Smith, collector. Rostral much broader than high, visible from above; internasals about as long as broad, smaller than prefrontals which are broadly in contact with supraocular; frontal as long as its distance from tip of snout and as the interparietal suture, broader in front than behind; parietals one-fourth longer than distance of frontal from tip of snout; nostril large, roundish between two subequal nasals; loreal squarish, about as high as long; one preocular, not in contact with frontal; one subpreocular; two postoculars; temporals 2+3; supralabials eight, fourth and fifth entering eye; five lower labials in contact with anterior chin-shields which are of the same length as the posterior; 19 rows of scales, the 11 median faintly, but distinctly keeled, the outer four on each side smooth; ventrals 203, obtusely angulate laterally; anal divided; 87

^a Reproduced in this work on Plate XXI.




ELAPHE QUADRIVIRGATA. (FROM TEMMINCK AND SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

pairs of subcaudals. Color (in alcohol) above tawny-olive, with four longitudinal, dark brown stripes, the outer ones involving the third and fourth scale rows, the middle ones the sixth and seventh scale rows; the lower stripes end at the vent, the upper ones continue on the tail to the tip; dark brown stripe from eye obliquely backward


279



280



281



FIGS. 279-281.—*ELAPHE QUADRIVIRGATA*, ADULT. $1\frac{1}{3} \times$ NAT. SIZE. 279, TOP OF HEAD; 280, SIDE OF HEAD; 281, COLOR PATTERN AROUND THE MIDDLE OF THE BODY. No. 36548, U.S.N.M.

across the eighth supralabial and ending at the angle of the mouth; underside and lower half of supralabials light yellowish.


Dimensions.

	<i>mm.</i>
Total length.....	890
Snout to vent.....	695
Vent to tip of tail.....	195

The *young* may, or may not, have a coloration like that of the adult, a feature to be mentioned more in detail under the head of variation. When unlike the adult, as in Nos. 23435, from Yezo (figs. 282-283), and 34026, from Yokohama, the longitudinal stripes are absent, and

instead there are numerous more or less complete narrow, dusky, transverse zigzag lines often with pale edges, or these cross lines are interrupted so as to form a lateral series of dots and a dorsal cross line, often quite oblique; on the posterior half of the upper surface of the head there is a brown, black-edged pattern somewhat resembling an ornate M, and anterior to this a semilunar band across prefrontals and supraoculars, as well as various marks of similar color on top and sides of the snout and on the supralabials; a brown black-edge postocular band crossing the last supralabial; ventrals faintly marked with brownish except on lateral angle which is visible as a whitish line.

Variation.—The number of scale rows and of supralabials seem to be absolutely constant in this species, and only two aberrations with regard to the latter are on record, namely, two specimens examined by Doctor Wall, who in one found on one side three supralabials



FIGS. 282-283.—*ELAPHE QUADRIVIRGATA*. YOUNG. ENLARGED. 282, COLOR PATTERN OF TOP OF HEAD AND NECK; 283, COLOR PATTERN AROUND THE MIDDLE OF THE BODY. NO. 23435, U.S.N.M.

entering the eye instead of two and in another 9 supralabials on one side, fifth and sixth touching the eye. An undivided anal occurs sometimes, and curiously enough it is found in no less than three specimens out of the five on record from Shikoku (Nos. 31876-31878). The subpreocular is also occasionally wanting, sometimes on both sides. Doctor Hilgendorf found three postoculars only once on one side, while several in our series show this anomaly, thus Nos. 23434 and 23435, while No. 4677 has them on both sides. The normal number of temporals is 2+2, but about 35 per cent of the specimens examined by me have 2+3, while I have only found 1+2 in three specimens, namely, Nos. 33978, 33979, and 34016. The number of ventrals varies between 193 and 215, subcaudals between 70 and 99.

The coloration is very variable. In the first place, there are two distinct color phases, the normal pale one with the distinct four stripes, the other a melanistic phase, more or less uniform blackish brown with the stripes more or less obscured according to the darkness of the

ground color. Of 45 specimens examined by me only six are melanistic, while in British Museum there is the same number in a total of 14 specimens. The black specimens are not confined to any one locality, but are found in all the islands, Yezo, Hondo, Shikoku, and Kiusiu, nor is this phase peculiar to the adults, as shown by No. 13754, which is quite young. In most of these black specimens, the labials, throat, anterior ventrals, and several longitudinal series of spots on the sides of the neck are more or less whitish.

In the pale, striped specimens there is also considerable variation. Thus the width of the stripes is not always the same, they being sometimes of equal width, sometimes the outer ones wider than the median ones, sometimes vice versa; sometimes the median stripes involve one row of scales only, the seventh, sometimes two, the sixth and seventh, or even three, viz. the eighth. The underside is also subject to great variation from nearly uniform whitish, or yellowish, to uniform slate-blue, with a distinct whitish line marking the lateral angle.

The most remarkable variation in coloration is probably that of the young, for while the majority appear to have a very distinct pattern of dusky dorsal cross lines and an ornate pattern on the head, as described above, others of exactly the same age are practically colored like the adults. Thus No. 34018, which shows as yet no longitudinal stripes, measures 413 mm. in total length, while the two striped young, Nos. 13754 and 34544, are respectively 375 and 378 mm. long. Of these the latter is pale and shows faint traces of a pattern on top of the head, sometimes also noticeable in larger specimens; the former is melanistic with a pale-brown underside.

Habitat.—This species seems to be confined to Japan proper. Numerous specimens are in the various museums from Kiusiu, Shikoku, Hondo, and Yezo. That it also occurs in the seven islands of Idzu is shown by a specimen from Nii shima in the Science College Museum, Tokyo (No. 92). Dr. Wall has recently recorded two specimens from the island of Yaku, south of Kiusiu.

A specimen in British Museum is said to have been collected by Mr. J. H. Leech at Gensan, Korea, but the locality is uncorroborated and highly suspicious. Mr. Leech also collected in Japan, and there is probably some error involved.^a

Such is probably also the case with regard to a specimen in the Science College Museum, Tokyo (No. 8), said to have been collected by Mr. Tashiro in Okinawa shima. The occurrence of this snake in the Riu Kius is highly improbable.

Similar remarks apply to Doctor Slunin's specimen from Hongkong in the St. Petersburg Museum.

^a See p. 29, under *Hynobius leechii*. It should be noted, however, that Werner has recently referred an aberrantly colored specimen to this species, which Doctor Haberer is said to have collected at Hankow, China, and which is now in the collection of the Academy of Sciences in Munich. The unfortunate part is that Doctor Haberer also collected in Japan (Werner, Bayer, Akal., p. 383), and added two other Japanese snakes to the Chinese fauna (see *E. climacophora*, p. 326. and *conspicillata*, p. 336).

List of specimens of *Elaphe quadrivirgata*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ven. trals.	Annal.	Subcaudals.	Temporals.	Supra-labials.
U. S. N. M.	4677	Male	Japan		Doctor Morrow	19	196	2	74	2+3	8
Do.	13754	Young ^a	do		P. L. Jouy	19	207	2	85	2+2	8
Do.	15418	Male	Otake, Shinshuu	Oct. —, 1882	do.	19	205	2	80	2+3	8
Do.	15419	do	Near Tokyo	Apr. 12, 1883	do.	19	205	2	80	2+2	8
Do.	23433	do	Yezo		S. Nozawa	19	197	2	83	2+3	8
Do.	23434	do. ^b	do		do.	19	202	2	84	2+2-3	8
Do.	23435	Young ^c	do		do.	19	198	2	78	2+3	8
Do.	31819	Female ^d	Yamagawa, Kiinsu	June 14, 1903	Dr. H. M. Smith	19	203	2	87	2+3	8
Do.	31873	Male ^e	Kochi, Shikoku	May —, 1903	do.	19	209	2	80	2+2	8
Do.	31875	do. ^b	do	do.	do.	19	209	2	91	2+2	8
Do.	31876	do	do	do.	do.	19	205	1	87	2+2	8
Do.	31877	do	do	do.	do.	19	207	1	87	2+2	8
Do.	31878	do	do	do.	do.	19	214	1	95	2+2	8
Do.	33867	Female	Mount Fuji	Aug. —, 1898	A. Owston	19	200	2	78	2+3	8
Do.	33908	do	do	do.	do.	19	207	2	87	2+2-3	8
Do.	33909	Male	do	do.	do.	19	206	2	90	2+2	8
Do.	33970	do	do	do.	do.	19	208	2	86	2+2-3	8
Do.	33971	do	do	do.	do.	19	209	2	86	2+2	8
Do.	33972	Female	do	do.	do.	19	209	2	85	2+2	8
Do.	33973	Halfgr.	do	do.	do.	19	201	2	87	2+2-3	8
Do.	33974	do	do	do.	do.	19	202	2	87	2+2	8
Do.	33975	Female	do	do.	do.	19	204	2	81	2+3	8
Do.	33976	do	do	do.	do.	19	203	2	82	2+2	8
Do.	33977	do	do	do.	do.	19	205	2	86	1+2	8
Do.	33978	do	do	do.	do.	19	209	2	81	1+2	8
Do.	33979	do	do	do.	do.	19	206	2	90	1+2	8
Do.	33980	do	do	do.	do.	19	202	2	82	2+2	8
Do.	33981	Halfgr.	do	do.	do.	19	213	1	88	2+2	8
Do.	34004	Female	Yokohama	Apr. —, 1903	do.	19	203	2	84	2+2	8
Do.	34005	Male	do	do.	do.	19	201	2	87	2+2	8

Do.....	34015	do, b	Suruga.....	do.....	19	208	2	2+2-3	8
Do.....	34016	Haligr.	Yokohama.....	Aug. —, 1898	19	204	2	86	1+2	8
Do.....	34017	Female.	do.....	Nov. 2, 1902	19	203	2	83	2+2	8
Do.....	34018	Young.	Mount Fuji.....	Aug. —, 1898	19	202	2	85	2+2	8
Do.....	34019	do.	do.....	do.....	19	205	2	88	2+2-3	8
Do.....	34020	do.	do.....	do.....	19	205	2	88	2+2-3	8
Do.....	34021	do.	do.....	do.....	19	201	2	82	2+2	8
Do.....	34022	do.	do.....	do.....	19	207	2	80	2+3	2
Do.....	34026	do.	Suruga.....	Nov. —, 1903	19	205	2	81	2+2	8
Do.....	34038	Female b.	Nagasaki.....	Aug. —, 1901	19	199	2	87	2+3	8
Do.....	34544	Young e.	Isobe, near Yokohama.....	May 10, 1904	19	207	2	86	2+3	8
Do.....	36548	Male f.	Tokyo.....	Sci Coll. Tokyo.	19	209	2	85	2+2	8
Do.....	36549	do, f	do.....	do.....	19	208	2	2+2	8
Do.....	36550	do, g.	Nii shima, Idzu Islands.....	do.....	19	211	2	87	2+2	8
Sci. Coll. Tokyo.....	8	Male.	[Okina wa shima?].	Tashiro.....	19	215	2	99	2+3	8
Mus. Leiden.....	396	(b)	Japan.....	do.....	19	208	2	90
Brit. Mus.....	a	Male f.	do.....	Leiden Mus.....	19	208	2	88
Do.....	b	do, f	Yokohama.....	Challenger Exped.....	19	204	2	81
Do.....	c	do, f	do.....	do.....	19	204	2	84
Do.....	d	Female f.	do.....	do.....	19	200	2	70
Do.....	e	Young f.	Nikko.....	C. Maries.....	19	209	2	80
Do.....	f	do, f	do.....	do.....	19	200	1	87
Do.....	g	do, f	Miyanoshta.....	J. H. Leech.....	19	195	2	80
Do.....	h	Haligr f.	Satsuma.....	do.....	19	202	2	80
Do.....	k	Female J.	Japan.....	Leiden Mus.....	19	200	1	84
Do.....	l	do, f	do.....	do.....	19	202	2	83
Do.....	m	Male J.	do.....	Zool. Soc.....	19	203	2	85
Do.....	n	Young J.	do.....	Sir R. Owen.....	19	203	2	83
Do.....	o	do, J.	Kumamoto.....	J. H. Leach.....	19	203	2	83
Do.....	p	do, f	do.....	do.....	19	205	2	89

♂ Melanistic, p. 331.
 ♀ Melanistic.

c Fig. 282-283.
 d Description, p. 328.

e P. 231.
 f Sci. Coll. No. 72. Figs. 279-281.

g No subpreocular.
 h Fauna Jap. Ombid., pl. 1.

i Boulenger, Cat. II, p. 60.
 j Boulenger; melanistic.

ELAPHE CONSPICILLATA ^a (Boie).

DJIMEGURI.

1826. *Coluber conspicillatus* BOIE. Isis. 1826, p. 211 (type-locality, Japan; Blomhoff, collector); Bijdr. Natuurk. Wetensch., II, Pt. 1, 1827, p. 262.—SCHLEGEL. Phys. Serp., 1837, II, p. 171 (Japan).—TEMMINCK and SCHLEGEL. Fauna Japon. Rept., 1837, p. 85; Ophid., pl. III (Japan).—HILGENDORF. Mitth. Deutsch. Ges. Ost-Asiens. I, heft 10, 1876, p. 30 (Tokyo).—FRITZE. Mitth. Deutsch. Ges. Ost-Asiens. V, heft 46, 1891, p. 239 (not seen in Yezo).—BOULENGER. Cat. Sn. Brit. Mus., II, 1894, p. 51 (Yokohama; Nikko; Haruna Hills).—BOETTGER. Kat. Schl. Mus. Senckenberg., 1898, p. 52 (Hondo).—WALL. Proc. Zool. Soc. London, 1903, p. 100 (Nagasaki); 1905, II, p. 513 (Hondo, Tanaga II.).—*Elaphis conspicillatus* DUMÉRIL and BIBRON, Erpét. Gén., VII, Pt. 1, 1854, p. 285 (southern islands of Japan).—HILGENDORF. Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 114 (Tokyo).—OKADA. Cat. Vert. Japan, 1891, p. 68 (Tokyo; Osaka; Nikko).—*Elaphis conspicillata* MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1866, p. 112; 1876, p. 379 (Nagasaki).
1860. *Proterodon tessellatus* HALLOWELL, Proc. Phila. Acad., 1860, p. 499 (type-locality, Japan).—*Dinodon* ? *tessellatus* BOULENGER, Cat. Sn. Brit. Mus., I, 1893, p. 364.
1878. *Coronella perspicillata* MUELLER, Verhandl. Naturf. Ges. Basel, VI, Pt. 4, p. 595 (*lapsus*).


The specimen which served Hallowell for the type of the new genus and species *Proterodon tessellatus* was probably never entered in the catalogue of the National Museum, and seems to have been lost. Boulenger has referred it, with a query, to the genus *Dinodon*,^b but I have no doubt that Hallowell described a fully adult *Elaphe conspicillata*, which he failed to recognize as such, because of the disappearance of the characteristic markings on head and back. Scale formula, coloration, and all, agree with *E. conspicillata*. Even the dental formula, as given by him, does not deviate so very far from what I find in my specimens, in which the posterior teeth are decidedly smaller than those preceding them.

Description (figs. 284–286).—*Adult male*; U.S.N.M. No. 34025; Mori, near Yokohama; October, 1902; Owston, collector. Rostral very much broader than high, visible from above; internasals squarish, broader than long, much shorter than prefrontals which are in contact with supraoculars; rostral as long as its distance from tip of snout, and longer than interparietal suture, twice as large as the rather narrow supraocular; parietals much longer than distance of frontal from tip of snout; nostril large, rounded, between two nasals, of which the posterior is considerably larger than anterior; loreal rather small, wider than high; one preocular not in contact with prefrontal; no subpreocular or subocular; two postoculars; temporals large,

^a Signifying spectacled, from *conspicilla*, medieval Latin for a pair of spectacles; referring to the black markings on the head of the young.

^b Cat. Snakes Brit. Mus., I, 1893, p. 364.

1+2; supralabials 7, sixth largest, third and fourth entering orbit; four lower labials in contact with anterior chin-shields which are longer than posterior; 21 rows of scales, with feeble keels visible on the middle nine rows; 207 ventrals, scarcely angulate laterally; anal divided; 74 pairs of subcaudals. Color (in alcohol) above uniform drab-gray, the outer seven or eight scales with black bases and lateral margins, and a black spot on the upturned outer portion of every other or third ventral; head above with a few black marks, viz, an inverted V on the frontal and parietal region with a black line on the posterior third of the interparietal suture; three black dots on prefrontals; a black line on the internasals along the suture with the rostral; on the sides of head a black mark on suture between first and second supralabials and another under eye on suture between third and fourth supralabials, this mark coming to the lower lip on the suture between fourth and fifth lower labials; an oblique black



FIGS. 284-286.—*ELAPHE CONSPICILLATA*. 2 X NAT. SIZE. 284, TOP OF HEAD, WITH COLOR PATTERN; 285 SIDE OF HEAD; 286, UNDERSIDE OF HEAD. No. 36547, U.S.N.M.

line from postoculars to lip across the middle of the sixth (penultimate) supralabial; underside whitish checkered with bold black spots, underside of tail with a narrow, often interrupted, black zigzag line on the median suture.

Dimensions.

	<i>m.m.</i>
Total length.....	771
Snout to vent.....	623
Vent to tip of tail.....	148

The *young*, as a rule, have a much more distinct color pattern, such as No. 6164, consisting of narrow, black, white-edged dorsal cross bands alternating with vertical bars of the same width and color, sometimes divided into two rows of spots; the pattern on the head is much more distinct, and the black lines continuous, the most marked being the inverted V, a curved line on the prefrontals between the eyes, and the black band back of the upper outline of the rostral.

Variation.—There is not any great variation in the scutellation. The number of scale rows seems to be constant, 21^a, and so is the absence of a true subpreocular (the anomalous scute figured in fig. 285 is really only an accidental subocular separated off on one side from the upper posterior corner of the third supralabial). Temporals are also nearly always 1 + 2, though Boulenger mentions the occurrence of 2 + 2. In one of our specimens the anal is undivided; ventrals vary between 200 and 227, subcaudals between 60 and 76. In two specimens there are eight supralabials on one side, in one six. One specimen, No. 34011, is abnormal in lacking the loreal on one side.

In the young there is some apparent variation in the color due to the fact that the strongly marked first color pattern gradually gives way to the uniform tint of the adults. Apparently some specimens retain the early pattern later than others; thus No. 31852, although 18 mm. longer than No. 34013, which has nothing left of the original pattern on the back but numerous irregular and scattered black dots, shows the young pattern in its least modified form. These remarks refer chiefly to the pattern on the back, because that of the head remains strong and plain long after the back has become uniform. Some of the head markings remain longer than others, especially the most characteristic ones, viz, the line crossing the penultimate supralabial, the two lines crossing the lips below the eye and between first and second supralabials, and the black postrostral line.

Habitat.—Apparently confined to Japan, but not so common as *E. quadrivirgata* or *E. climacophora*, and comparatively few specimens with definite localities attached are in the museums. The Leiden Museum specimens collected by von Siebold and Buerger probably came from Kiusiu. The Stockholm Museum has a specimen (No. 536) from Nagasaki, collected by Dr. O. Nordquist. The Berlin Museum has it from Nagasaki and Tokyo, and British Museum from Yokohama, Nikko, and the Haruna Hills. Dr. Wall has recently received a specimen from Tanega shima, south of Kiusiu. Our museum, finally, has it from Tokyo, from Oide, Shinshiu, where Jouy collected it in 1882, and from Mount Fuji. Dr. Hugh M. Smith collected it in Shikoku. The only observation as to the altitude attained by this species in the mountains is furnished with a specimen in British Museum, which Lord Dormer collected in the Haruna Hills, 2,500 feet above the sea.

Giglioli and Salvadori record^b a young specimen as *Callopeltis conspicillatus* taken by the "Vettor Pisani" expedition at the Avahuna River, Olga Bay, Russian Coast Province, during September, 1879.

^a A specimen in the Munich Museum said to have been collected at Hankow, China, by Doctor Haberer, is recently reported by Doctor Werner (Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 357) as having 23 scale rows. Altogether it is a very dubious record (see pp. 326, 331).

^b Proc. Zool. Soc. London, 1887, p. 595.

Until further details relative to this specimen be obtained it is permissible to suspect that a mistake in identification has been made, and that in reality it is an *E. schrenkii*. I must add, however, that there are several specimens in the Hamburg Museum (Nos. 849-852) collected by Doerries in the "Amurgebiet," which are identified there as *E. conspiciata*.

List of specimens of Elaphe conspiciata.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scute rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supralabials.
U.S.N.M.	6164	Young.	Japan.....			21	214	2	72	1+2	7-8
Do.	15417	Male	Oide, Shinshiu....	1882. Oct. —	P. L. Jouy.....	21	212	2	1+2		7
Do.	31852	Half gr. ^a	Kochi, Toza, Shikoku.	1903. May —	Dr. H. M. Smith.	21	219	2	73	1+2	7
Do.	34009	Female	Mount Fuji.....	1898. Aug. —	A. Owston.....	21	227	2	67	1+2	7
Do.	34010	do	do.....	do.....	do.....	21	217	1	68	1+2	7
Do.	34011	Male	do.....	do.....	do.....	21	211	2	67	1+2	6-7
Do.	34012	Female	do.....	do.....	do.....	21	214	2	69	1+2	7-8
Do.	34013	Half gr. ^b	do.....	do.....	do.....	21	206	2	63	1+2	7
Do.	34025	Male	Mori, near Yokohama.	1902. Oct. —	do.....	21	207	2	74	1+2	7
Do.	36547	Young ^d	Tokyo.....		Sci. Coll., Tokyo	21	207	2	68	1+2	7
Do.		Adult ^e	Japan.....		do.....	21	211	2	73	1+2	8?
Brit. Mus.	a	Male f	do.....		Leiden Mus....	21	219	2	67		
Do.	b	Young f	do.....		do.....	21	217	2	66		
Do.	c	Male f	Yokohama.....		Challenger Exped.	21	215	2	67		
Do.	d	Young f	Nikko.....		do.....	21	224	2	63		
Do.	e	do. f	Haruna Hills, 2,500 feet.		Lord Dormer...	21	206	2	68		

^a Length 488 mm.

^b Length 470 mm.

^c Description, p. 334.

^d Figs. 284 2-6; Sci. Coll. No. 71.

^e Hallowe's type of *Proterodon tessellatus*.

^f Boulenger, Cat. II, p. 52.

Genus LIOPELTIS^a Fitzinger.

1843. *Liopeltis* FITZINGER, Syst. Rept., p. 26 (type, *Herpetodryas tricolor* Schlegel).

1860. *Eurypholis* HALLOWELL, Proc. Phila. Acad., 1860, p. 493 (type, *E. semicarinatus*).

1862. *Phragmitophis* GUENTHER, Ann. Mag. Nat. Hist. (3), IX, p. 126 (type, *Cyclophis tricolor*).

1895. *Entechinus* COPE, Proc. Phila. Acad., 1894, p. 427 (type, *Cyclophis major*).

1900. *Entacanthus* COPE, Rep. U. S. Nat. Mus., 1898, p. 780 (type, *Cyclophis major*).

^a From λειος, smooth; πέλτη, a little shield; scale.

Liopeltis is almost coextensive with Boulenger's *Ablabes* (part of *Ablabes* Duméril and Bibron, 1854) for which he selected *A. baliodeirus* as the type (= *Gongylosoma* Fitzinger, 1843), with the exception of the latter species, which Cope says has hypapophyses present throughout the vertebral column. The species are very difficult to distinguish generically from *Elaphe*, the chief character being the absence of apical pits to the dorsal scales. These are very obscure in some of the *Elaphe*, for instance *E. rufodorsata*, and can only be discovered by a very careful examination of the dry scale. This latter species, therefore, has been referred repeatedly to the genus here treated of.

This genus is mostly oriental in its distribution, extending in the west to the Himalayas and in the south to the Malayan Islands. In China it does not reach farther north than Shanghai and is also unknown in Japan proper. The Chinese species extends, however, to Formosa, and there are two species apparently peculiar to the Riukius.

KEY TO THE RIUKIU AND FORMOSA SPECIES OF LIOPELTIS.

*a*¹. Scale rows 15.

*b*¹. Posterior chin-shields much shorter than the anterior.....*L. major*, p. 338.

*b*². Posterior and anterior chin-shields subequal.....*L. semicarinata*, p. 340.

*a*². Scale rows 17*L. herminæ* p. 343.

LIOPELTIS MAJOR ^a (Guenther).

1858. *Cyclophis major* GUENTHER, Cat. Colubr. Snakes Brit. Mus., p. 120 (type-locality, near Ningpo, China; type in Brit. Mus.); Rept. Brit. India, 1864, p. 230 (Ningpo; Hongkong) pl. xvii, fig. *L*; Ann. Mag. Nat. Hist. (6), I, 1888, p. 170 (near Kiukiang, China).—*Ablabes major* BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 140 (Lueshan Mts., Kiukiang).—BOULENGER, Cat. Snakes Brit. Mus., II, 1894, p. 279 (eastern China; Formosa); Proc. Zool. Soc. London., 1899, p. 165 (Prov. of Fokien).—WALL, Proc. Zool. Soc. London, 1903, p. 93 (Hongkong; Formosa).

1860. *Herpetodryas chloris* HALLOWELL, Proc. Phila. Acad., 1860, p. 503 (type-locality, Hongkong; type, U.S.N.M. No. 7335; Brooke, collector).

Description.—*Half grown male*; U.S.N.M. No. 7335; type of *Herpetodryas chloris* Hallowell; Hongkong Island, China; May, 1854; Mr. Brooke, collector (figs. 287–289). Rostral somewhat broader than high, visible from above; internasals very small, much smaller than prefrontals, which are broadly in contact with supraocular; frontal much longer than its distance from tip of snout, equalling the interparietal suture, slightly wider in front than behind, wider than supraoculars; parietals long, twice as long as the frontal's distance from rostral, followed by two enlarged scales or scutes, which are larger than internasals; nostril large, oval, in the anterior portion of an elongated nasal; loreal much longer than high, about the same


^a Signifying larger; rather large.

length as nasal but lower; one preocular widely separated from frontal; two postoculars, upper largest; temporals 1 + 2, large, with straight edges; 8 supralabials, fourth and fifth entering eye, seventh largest, sixth nearly triangular; four lower labials in contact with anterior chin-shields, which are longer than posterior; 15 rows of smooth scales; 163 ventrals, not angulate laterally; anal divided; 86 pairs of subcaudals. Color (in alcohol) above blue, in life bright green; under side white or, in life, yellow.

Dimensions.

	<i>mm.</i>
Total length.....	684
Snout to vent.....	506
Vent to tip of tail.....	178

A *young* specimen from the province of Fokien, China, is described by Boulenger^a as having irregular black transverse spots on the nape and anterior part of the back, and traces of an interrupted black lateral streak.



FIGS. 287-289.—*LIOPELTIS MAJOR*. 2 × NAT. SIZE. 287, TOP OF HEAD; 288, SIDE OF HEAD; 289 UNDEP-
SIDE OF HEAD. No. 7335, U.S.N.M.

Variation.—Boulenger describes this species as having a divided nasal, but I can not make out any such division in our specimen. Doctor Wall mentions a specimen having seven labials, with the third and fourth touching the eye on one side. Ventrals range between 163 and 177; subcaudals between 70 and 90.

Habitat.—Apparently restricted to the lower Yangtse Valley and coasts of eastern China, between Hongkong and Shanghai, as well as to Formosa, from which island Swinhoe brought four specimens to British Museum. Doctor Wall saw two additional specimens from Formosa in the Hongkong City Hall Museum. In the Chinese province of Fokien it has been found as high as 3,000 to 4,000 feet altitude.

^a Proc. Zool. Soc. London, 1899, p. 165.

List of specimens of Liopeltis major.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supradubials.
				1854.							
U.S.N.M.	7335	Male ^a	Hongkong Island	May	Brooke	15	163	2	86	1+2	8
Brit. Mus.	<i>f</i>	Female ^b	Formosa		R. Swinhoe	15	170	2	86		
Do.	<i>g</i>	do. ^b	do.		do.	15	168	2	86		
Do.	<i>h</i>	do. ^b	do.		do.	15	176	2	85		
Do.	<i>i</i>	do. ^b	do.		do.	15	171	2	83		

^a Description, p. 338; figs. 287-289. Type of *H. chloris*.^b Boulenger, Cat. II, p. 279.**LIOPELTIS SEMICARINATA** ^a (Hallowell).

1860. *Eurypholis semicarinatus* HALLOWELL, Proc. Phila. Acad., 1860, p. 493 (type-locality, Nafa, Okinawa shima, Riu Kiu; types, U. S. Nat. Mus., No. 7316; Wright, collector).—*Ablabes semicarinatus* BOULENGER, Proc. Zool. Soc. London, 1887, p. 148 ("Loochoo Islands," Pryer, collector); Cat. Sn. Brit. Mus., II, 1894, p. 278 (Great Loochoo Island; Holst, collector; Nagasaki!).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 862; author's separate, p. 13 (Okinawa shima), p. 864; author's separate, p. 15 (Amami-o-shima).—BOETTGER, Offenbach. Ver. Naturk., 33-36 Ber., 1895, p. 115 (Miyakoshima, Yaeyama group); Kat. Schl. Mus. Senckenberg., 1898, p. 78.—WALL, Proc. Zool. Soc., London, 1903, p. 101 (Japan; Loo Choos); 1905, II, p. 515 (Okinawa; Amami).
1868. *Cyclophis nebulosus* GUENTHER, Ann. Mag. Nat. Hist. (4), I, p. 418, pl. xix, fig. c (type-locality, Nagasaki!; type in Brit. Mus.; Whitely, collector).


The two specimens which served Hallowell as types for this species are still in the United States National Museum (No. 7316) in a fair state of preservation. The scale formulas, as given by him and as counted by me, agree pretty well, but his dimensions do not. He says: Total length, 1 foot 7 $\frac{3}{4}$ inches; of tail, 4 inches, five lines [Pied du Roi], the equivalents in mm being about 536 and 118. The two specimens, as measured by Mr. Paine, my assistant, are: Total length, 492 and 564 mm.; tail, 107 and 122 mm., a considerable discrepancy. The means of the two specimens, however, viz, 528 and 115, are so near Hallowell's figures that it may be regarded as certain that the latter are not intended to represent the dimensions of the individual snake.

An examination of the type of Guenther's *Cyclophis nebulosus*, which I was permitted to make in 1898, fully confirms Boulenger's view that it is only the young of the present species. The young specimen collected by Holst in Okinawa shima has the frontal shield considerably larger than Guenther's type, but is otherwise indistinguishable, and a young specimen from Miyakoshima of the Saki shima group now before me and described below is also similar. The alleged

^a Signifying half-keeled, the scales of the body being only keeled in their anterior half.

habitat of the type of *C. nebulosus*, which is said to have been "obtained by Mr. Whitely at Nagasaki" is undoubtedly erroneous, and it is almost certain that all the birds and reptiles recorded as obtained by him at that place were not actually collected there. They are of much more southern origin, most probably from some group of the Riu Kius.

Description.—*Young*—Science College Museum, Tokyo, Japan, No. 15; Miyakoshima, Saki shima group; figs. 290–292. Rostral about as broad as high, visible from above; internasals very small, the suture between them less than one-half the suture between the prefrontals; frontal much longer than its distance from the tip of the snout and equally suture between parietals; nostril large, extending nearly the entire height of the nasal; loreal trapezoidal higher behind, slightly longer than high; one preocular not in contact with frontal; two post-



FIGS. 290–292.—*LIOPELTIS SEMICARINATA*. 2 X NAT. SIZE. 290, TOP OF HEAD; 291, SIDE OF HEAD; 292, UNDERSIDE OF HEAD. No. 15, SCI. COLL. TOKYO.

oculars; temporals 1 + 2, large; eight supralabials, seventh and eighth largest, sixth rather smaller than fifth, fourth and fifth entering eye; four lower labials in contact with anterior chin-shields, which are as long as the posterior; 15 scale rows, median nine rows with a slight indication of keel at the base, outer three on each side smooth, all without apical pits; ventrals, 189; anal divided; subcaudals, 80 pairs. Color (in alcohol) above olive gray, with numerous obscure "nebulous" spots of dusky on the anterior third of body posteriorly grading into two indistinct dark longitudinal lines on each side, including a paler band about two scales wide, a dusky narrow line frequently interrupted marking the boundary between scales and ventrals; top and sides of head, except labials, uniform olive gray with a minute black spot on each parietal near the middle of the suture between them; entire lower surface and supralabials uniform whitish, the last two with posterior margins blackish.

Dimensions.

	<i>mm.</i>
Total length.....	285
Snout to vent.....	225
Vent to tip of tail.....	60


The *adult* male (U.S.N.M., No. 7316a; type: Naha, Okinawa shima; November, 1854; Wright, collector, figs. 293-295) is in all essential features of scutellation like the one described above, except that the snout is proportionally more elongated, with resulting longer internasals and longer loreal. The coloration is more uniform above, with faint indications of longitudinal dusky streaking, which is very marked on the side of the neck as interrupted longitudinal black streaks; last two supralabials marked as in the young.

Dimensions.

	<i>mm.</i>
Total length.....	492
Snout to vent.....	385
Vent to tip of tail.....	107

This snake grows to be more than 800 mm. long.

Variation.—Variation both as to scutellation and coloration is rather slight in this species. The most noteworthy abnormality is presented by U.S.N.M., No. 11127, which has the fifth supralabial on



FIGS. 293-295.—*LIOPELTIS SEMICARINATA*. 2 × NAT. SIZE. 293, TOP OF HEAD; 294, SIDE OF HEAD; 295, UNDERSIDE OF HEAD. No. 7316a, U.S.N.M.

both sides divided by a horizontal suture so as to cut off a large posterior subocular. The number of scale rows, oculars, temporals, and supralabials seems to be absolutely constant, the ventrals and subcaudals vary between narrow limits. The greatest deviation from the normal is shown by the type of *C. nebulosus*, which is said to have only 174 ventrals, while in the other recorded specimens this number oscillates between 185 and 192; the subcaudals vary from 70 to 82.

The most aberrant coloration is shown by one of five large individuals in the Science College Museum, Tokyo. It has four very distinct brown longitudinal stripes on the whole length of the body from nape to vent, while the upper two stripes continue to the end of the tail. These specimens were collected by Mr. Tashiro in Okinawa shima.

Habitat.—Restricted to the Riu Kiu Archipelago. The types were collected during the Rodgers North Pacific Exploring Expedition at

Naha, Okinawa shima, from which island there are now specimens in various museums, thus in Science College Museum, Tokyo, and in the Imperial Museum (Ueno Park), Tokyo, British Museum, and Hamburg Museum (No. 1491), and Doctor Fritze states that it is not rare there in summer. That it also occurs in the southern group is shown by a Science College specimen (No. 15) and two in Museum Senckenbergianum, all from Miyakoshima. Finally, Doctor Fritze enumerates it as one of the species occurring in Amami-ō-shima of the northern group, a statement possibly based upon specimens collected by Doctor Doederlein. Doctor Wall also records three specimens from this island as having been secured by Owston's collectors.

Its alleged occurrence in Japan proper is entirely unsupported. Whitely's specimen, upon which Guenther founded his *Cyclopis nebulosus* is said to have been obtained in Nagasaki, but all his birds and reptiles credited to that locality most certainly came from some of the Riu Kiu Islands.^a

List of specimens of *Liopeltis semicarinata*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Labials.
1854.											
U.S.N.M.	7316a	Male a	Near Naha, Okinawa shima.	Nov. —	Wright	15	186	2	77	1+2	8
Do.	7316b	Female b	do.	do.	do.	15	190	2	78	1+2	8
Do.	11127	Male c	"Japan"		Perry exped.	15	191	2	77	1+2	8
Do.	26518	do. d	Okinawa shima.		Tashiro.	15	187	2	77	1+2	8
Sci. Coll. Tokyo.	15	Young e	Miyakoshima			15	189	2	80	1+2	8
Brit. Mus.	a	Male f	"Loo Choo Islds"		Pryer collection	15	190	2	82	1+2	8
Do.	b	do. f	do.		do.	15	190	2	76	1+2	8
Do.	c	do. f	do.	Nov. —	do.	15	190	2	77	1+2	8
Do.	d	Female f	do.		do.	15	192	2	70	1+2	8
Do.	e	do. f	do.		do.	15	187	2	72	1+2	8
Do.	f	Young f	Great Loo Choo I		Holst	15	185	2	76	1+2	8
Do.	g	do. g	Nagasaki (?)		Whitely	15	174	2	74	1+2	8

a Type; description p. 342, figs. 293-295.

b Type.

c P. 342.

d Sci. Coll. No. 7.

e Description, p. 341, figs. 290-292.

f Boulenger, Cat. II, p. 279.

g Boulenger; type of *C. nebulosus*.


LIOPELTIS HERMINÆ b (Boettger).

1895. *Ablabes herminae* BOETTGER, Zool. Anz., XVIII, July 8, 1895, p. 269 (type-locality, Miyakoshima, Sake shima group. Riu Kius; type in Mus. Senckenberg.); Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 110 (Miyakoshima); Kat. Schl. Mus. Senckenberg., 1898, p. 78 (Miyakoshima).—BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 643.—WALL, Proc. Zool. Soc. London, 1903, p. 101 (Loo Choos); 1905, II, p. 515 (Iriomote).

a See also Wall, Proc. Zool. Soc. London, 1905, II, p. 515, footnote.

b Named for —?

Description.—*Adult male*; U.S.N.M.No. 34014; Ishigaki shima, Riu Kiu Archipelago; June, 1899; Owston collection (figs. 296–297). Rostral as high as broad, just visible from above; internasals very small, much less than one-half the size of the prefrontals, which are broadly in contact with supraoculars; frontal as long as its distance from tip of snout and as interparietal suture, broader than supraoculars, considerably broader in front than behind; parietals broad, much longer than frontal; nostril large between two nasals, of which the posterior is much the larger; loreal, long and low, twice as long as high; one preocular widely separated from frontal; two postoculars; 1 + 2 large temporals; 8 supralabials, fourth and fifth entering eye, seventh largest, twice as high as eighth, which is very small, smaller



FIGS. 296–297.—*LIOPELTIS HERMIN.E.* $1\frac{1}{2} \times$ NAT. SIZE. 296, TOP OF HEAD; 297, SIDE OF HEAD. No. 34014, U.S.N.M.

than the temporal immediately above it; four lower labials in contact with anterior chin-shields which are longer than the posterior (on one side, shorter on the other); 17 rows of scales, which are smooth except on the posterior fourth of the body, where they show a faint keel at the base; 162 ventrals; anal divided; 52 pairs of subcaudals. Color (in alcohol) above, olive gray, each scale with an ill-defined median dusky spot near the tip, thus forming faint dusky longitudinal lines; underside whitish, some ventrals with a gray line along the base, and the subcaudals more or less mottled with gray; four median supralabials white, with dark gray sutures, the others olive-gray, like the top and sides of the head.

Dimensions.

	mm.
Total length.....	745
Snout to vent.....	610
Vent to tip of tail.....	135

Variation.—The only noteworthy deviation from the above description shown by the type specimen in the Senckenberg Museum is the two undivided subcaudals following the first two pairs; the ventrals are practically identical in number, while the subcaudals are slightly more

numerous in the type; in the latter the subcaudals have a dark gray longitudinal stripe on each side. Doctor Wall in one specimen counts 47 subcaudals only, and in one he records 19 scale rows "in mid-body; anteriorly and posteriorly the normal 17 scales were present." This specimen, from Iriomote shima, has 159 ventrals and 60 subcaudals.

Habitat.—Thus far only known from the Saki shima group, Riu Kiu Archipelago. Mr. B. Schmacker's Japanese collector obtained a single specimen in Miyakoshima. Two more specimens in Mr. Owston's collection were recorded by Doctor Wall, one of which is now in the United States National Museum. It was collected in Ishigaki shima, June, 1899. Another is now recorded by him from Iriomote shima.

List of specimens of *Liopeltis herminx*.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supralabials.
U.S.N.M.	34014	Male ^a . . .	Ishigaki shima . . .	1899. June —	A. Owston	17	162	2	52	1+2	8
Senckenberg .	8277a	...do. ^b . . .	Miyakoshima		B. Schmacker . . .	17	163	2	60	1+2	8

^a Description, p. 344; figs. 296-297.

^b Type.

Genus **PTYAS**^a Fitzinger.

1843. *Ptyas* FITZINGER, Syst. Rept., p. 26 (type, *Coluber blumenbachii*=*C. mucosus*).

1854. *Coryphodon* DUMÉRIL and BIBRON, Erpét. Gén., VII, Pt. 1, p. 180 (part).

The two species, in the East known as rat-snakes, may be easily distinguished thus:

*a*¹ Scale rows 17; scale rows a head length anterior to vent 14; ventrals more than 185..... *P. mucosus*, p. 345.

*a*² Scale rows 15; scale rows a head length anterior to vent 11; ventrals less than 185..... *P. korros*, p. 348.

PTYAS MUCOSUS ^b (Linnæus).


1758. *Coluber mucosus* LINNÆUS, Syst. Nat., 10 ed., I, p. 226 (type-locality, "in Indiis;" type in Roy. Mus. Stockholm); 12 ed., I, 1766, p. 388.—*Ptyas mucosus* COPE, Proc. Phila. Acad., 1860, p. 563.—GUENTHER, Rept. Brit. India, 1864, p. 249 (South-continental Asia; Formosa).—*Zamenis mucosus* BOULENGER, Cat. Sn. Brit. Mus., I, 1893, p. 385 (south. Asia; Formosa).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 41 (Taiwan fu, Formosa).—ANDERSON, Bih. Svensk. Vet. Akad. Handl., XXIV, Pt. 4, no. 6, 1899, p. 25 (type).—WALL, Proc. Zool. Soc. London, 1903, p. 90 (Hongkong).

^a From *πρῶς*, a kind of snake, so named by the ancient Greeks from its hissing or spitting (*πρῶω*).

^b Signifying slimy.—The gender of *Ptyas* is feminine, but as Cope, and all the authors after him, have treated it as masculine, it is here so considered.

1820. *Coluber blumenbachii* MERREM, Syst. Amph., p. 119 (type-locality, Bengal).—*Coryphodon blumenbachi* JAN, Elenc. Ofid., 1863, p. 64 (Bengal; Ceylon; Japan); Icon. Ophid., livr. 24, 1867, pl. III, figs. 2-3 (Ceylon; Bengal).
1839. *Coluber dhumna* CANTOR, Proc. Zool. Soc. London, 1839, p. 51 (type-localities, Carnatic, Orissa, Bengal, Nepal, Assam, Arracan, Tenasserim).
1860. *Leptophis trifrenatus* HALLOWELL, Proc. Phila. Acad., 1860, p. 503 (type-locality, Hongkong; type, U.S.N.M., No. 7510).
1867. *Coryphodon blumenbachi* var. *carinata* JAN, Icon. Ophid., livr. 24, pl. III, fig. 4 (type-locality, "Japan;" type in Mus. Genova) (not *C. carinatus* GUENTHER, 1858).

Description.—Adult male; U.S.N.M. No. 7510, type of *Leptophis trifrenatus*; island of Hongkong; April, 1854; Lieutenant Brooke,



FIGS. 298-300.—*PTYAS MUCOSUS* (TYPE OF *TRIFRENATUS*). NAT. SIZE. 298, TOP OF HEAD; 299, SIDE OF HEAD; 300, UNDERSIDE OF HEAD. NO. 7510, U.S.N.M.

collector (figs. 298-300). Rostral broader than high, somewhat projecting; internasals as long as broad, much smaller than prefrontals, which are in contact with supraocular; frontal as long as its distance from rostral and as the interparietal suture, much broader in front than behind, the lateral edges distinctly concave; supraocular large, as wide as the frontal at its lateral convexity; parietals broad, as long as the distance of frontal from tip of snout; nostril large, vertically elliptic between two subequal nasals; three loreals, one larger anterior followed by two smaller, one above the other; one preocular not in contact with frontal; a subpreocular; eye rather large, its diameter

equaling its distance from nostril; two postoculars; 2+2 temporals; eight supralabials, seventh and eighth largest, fourth and fifth entering eye, fifth narrow and high, in contact with lower postocular and lower anterior temporal; five lower labials in contact with anterior chin-shields which are shorter than the posterior; one pair of postgenials; 17 rows of scales, the middle three of which are feebly keeled, the others quite smooth; 190 ventrals; anal divided; 111 pairs of subcaudals. Color (in alcohol) tawny with irregular narrow cross lines of black which is confined to the edges and tips of the scales; head above uniform, sides paler, labials and temporals edged behind with brownish black; underside whitish, the ventrals marked with a narrow brownish line at the posterior margin, becoming darker and broader behind, and in addition every fifth ventral has on each side a sharply defined, narrow dark brown bar along the base or anterior margin.

Dimensions.

	<i>mm.</i>
Total length.....	1.665
Snout to vent.....	1.250
Vent to tip of tail.....	415

The *young* are said to have usually light cross bands on the front half of the body.

Variation.—Ventrals are recorded as varying between 187 and 208, subcaudals between 95 and 140. The number of loreals is somewhat variable, and Doctor Wall records one specimen with nine supralabials on one side, fifth and sixth touching the eye.

This species grows to be more than 2 meters long.

Habitat.—The rat-snake has a wide distribution over the entire southern continental Asia. In the Himalayas it is found at an altitude of over 5,000 feet. In southeastern China it occurs from Hainan to Chusan. Swinhoe collected two specimens in Formosa, where, according to Guenther, the species is not rare. These specimens are in British Museum. A third specimen, from Taiwan fu, South Formosa, was sent by Mr. Schmacker to the Senckenberg Museum.

The locality of the specimen in the Genova Museum, upon which Jan based his variety *carinata* and which is said to have come from Japan, is undoubtedly erroneous.

List of specimens of Ptyas mucosus.

U. S. N. M. No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supralabials.
75:0	Male ^a	Hongkong, China.....	Apr. 1854.	Lieutenant Brooke..	17	190	2	111	2+2	8

^a Type of Hallowell's *Leptophis trifrenatus*. Description, p. 346, figs. 298-300.

PTYAS KORROS^a (Schlegel).

1837. *Coluber korros* SCHLEGEL, Phys. Serp., II, p. 139 (type-locality, Java; types in Leiden Mus.; Reinwardt, collector); Abbild. Amph., 1839, p. 99, pl. XXVII; pl. XXVIII, figs. 1-6.—*Coryphodon korros* JAN, Icon. Ophid., livr. 24, 1867, pl. IV, fig. 2 (Oude, India).—*Ptyas korros* GOPE, Proc. Phila. Acad., 1860, p. 563 (Siam).—STEINDACHNER, Novara Exped., Zool., I, Rept., 1867, p. 65 (Amoy).—*Zamenis korros* BOULENGER, Fauna Brit. India, Rept., 1890, p. 324; Cat. Sn. Brit. Mus., I, 1893, p. 384 (Himalayas to Southern China, Malay Peninsula, Sumatra, and Java).—WALL, Proc. Zool. Soc. London, 1903, p. 90 (Hongkong; Formosa).
1837. *Coluber cancellatus* OPPEL in Schlegel's Phys. Serp., II, p. 139 (specimen label in Paris Museum).

Description.—*Adult*; U.S.N.M. No. 26233; Trong, Lower Siam, Malay Peninsula; Dr. W. L. Abbott, collector. Rostral slightly broader than high, visible to a considerable extent from above; internasals slightly broader than long, smaller than prefrontals, which do not quite touch the supraoculars; frontal longer than its distance from rostral, and longer than the interparietal suture, much broader in front than behind, the lateral outline quite concave; supraoculars narrower than frontal at its narrowest point; parietals broad and truncate behind, their length equaling the distance of the frontal from tip of snout; nostril large, vertically elliptic, between two nasals, the anterior of which is slightly larger; two loreals, the anterior largest, as high as long, in contact with preocular above the posterior loreal, which is rather small; one preocular, barely in touch with frontal; one subpreocular; eye large, its diameter equaling its distance from center of nostril; two postoculars; 2+2 temporals; eight supralabials, seventh largest, fourth and fifth entering eye, fifth not in contact with temporal; five lower labials in touch with anterior chinshields, which are shorter than the posterior; 15 rows of smooth scales; 161 ventrals; anal divided; 140 pairs of subcaudals. Color (in alcohol) above anteriorly olive, each scale with a faint submarginal dusky line, on the posterior half of the body becoming gradually tawny olive, the dark lines becoming marginal and black; no distinct marks on head; sides of snout and six anterior supralabials pinkish buff; whole underside uniformly whitish suffused with pinkish buff.

Dimensions.—The specimen described is a skin in alcohol, the total length measuring about 1,210 mm.

Boulenger measures one specimen in British Museum as follows: Total length, 1,500 mm.; tail, 600 mm.

The *young* are said to have transverse series of pearl-colored spots forming cross bands.

Variation.—The number of loreals is commonly two or three, but Doctor Wall reports one Chinese specimen having only one loreal on

^a*Korros*, according to Schlegel himself, is the Javanese name for another snake (*Elaphe melanurus*), and was wrongly applied by him to the present species.

both sides. The number of ventrals varies between 161 and 177, the subcaudals between 122 and 145. Doctor Wall gives the latter as 110-116, but some error seems to be involved in this statement, as our five specimens from the Malay Peninsula corroborate Boulenger's figures, there being from 133 to 140 subcaudals. In our six specimens the preocular touches the frontal in only one specimen, namely, the one described above, in all the others the prefrontals and supraoculars are in decided contact.

Habitat.—This snake belongs to the group of species which extend from the eastern Himalayas eastward through southern China, in this case at least as far north as Amoy, and southward through the Malay Peninsula to Sumatra and Java.

It is admitted to the fauna of Formosa on the strength of a specimen stated by Doctor Wall to be in the Hongkong City Hall Museum. The records of this collection, as explained by him, can not now be trusted, and while it is true that no other collector has obtained it there, the general distribution of the species is such as to make its occurrence in Formosa probable.

List of specimens of *Ptyas korros*.

U. S. N. M. No.	Age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supralabials.
26233	Adult ^a	Trong, Lower Siam		Dr. W. L. Abbott	15	161	2	140	2+2	8
22877	do	do		do	15	174	2	133	2+2	8
22878	do	do		do	15	175	2	133	2+2	8
22879	do	do		do	15	174	2	138	2+2	8
22880	do	do		do	15	172	2	138	2+2	8
22853	do	do		do	15	168	2	137	2+2	8

^a Description, p. 348.

Genus ZAMENIS^a Wagler.

- 1830. *Zamenis* WAGLER, Nat. Syst. Amph., p. 188 (type, *Coluber viridiflavus*).
- 1843. *Eremiophis* FITZINGER, Syst. Rept., p. 25 (type, *C. trabalis*).
- 1843. *Hierophis* FITZINGER, Syst. Rept., p. 26 (type, *C. viridiflavus*).
- 1862. *Tyria* COPE, Proc. Phila. Acad., 1862, p. 338 (substitute for *Zamenis*) (not of FITZINGER, 1826).

ZAMENIS SPINALIS^b (Peters).

- 1866. *Masticophis spinalis* PETERS, Mon. Ber. Berlin Akad. Wiss., 1866, p. 91 (type-locality, "Mexico;" type in Berlin Mus.).—MOELLENORFF, Journ. N. China Br. R. Asiat. Soc. (n. s.), XI, 1877, p. 104 (Peking).—*Zamenis spinalis* GUENTHER, Ann. Mag. Nat. Hist. (4), IX, 1872, p. 22 (N. China).—


^a From ζαμενής, irascible.

^b Signifying pertaining to the spine, or backbone, with reference to the yellow, black-edged vertebral line.

- STRAUCH, in Przewalski's Mongoliya i Strana Tangutov, III, 1876, p. 46, pl. VIII (Ala-shan Desert).—MUELLER, Verh. Naturf. Ges. Basel, VII, Pt. 3, 1885, p. 681 (Mongolia).—BOULENGER, Cat. Snakes Brit. Mus., I, 1893, p. 394 (Korea; north China; Hainan).—WALL, Proc. Zool. Soc. London, 1903, p. 91 (Yangtse Valley; Huangtsun, N. China).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 355 (Tsingtau, China).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (S.), XVII, No. 1, 1905, p. 226 (Ala-shan; Ordos).
1866. *Zamenis bocourti* JAN, Nouv. Arch. Mus., Paris, II, 1866, Bull., p. 6 (type-locality, Siam; type in Paris Mus.; Bocourt, collector).
1867. *Zamenis cataphoranotus* JAN, Icon. Ophid., livr. 23, pl. II, fig. 2^a (type-locality, Bangkok, Siam; type in Paris Mus.).

There being no specimen of this species in our collection, I give the following somewhat condensed account of the type specimen derived from Doctor Peters's original description.

Description (figs. 301-303).—*Adult*; Berlin Museum; type; "Mexico" (erroneously). Snout projecting, rounded; prefrontals broader than long, not longer than internasals; frontal very long, anteriorly broad and very obtusely angulate, the lateral outline forming a shallow concavity, posteriorly forming an acute angle with rounded apex; supra-



FIGS. 301-303.—*ZAMENIS SPINALIS*. ENLARGED. 301, TOP OF HEAD; 302, SIDE OF HEAD; 303, UNDERSIDE OF HEAD. FROM JAN, ICON. OPHID., LIV. 23, PL. II, FIG. 2.

oculars anteriorly narrower than behind, where they are broader than the frontal at the same point; parietals scarcely longer than frontal, rounded behind; nostril between the upper part of the two nasals; loreal trapezoid, longer than high; preocular not in contact with frontal; one subpreocular; two postorbitals; temporals 2+2 on left side, 1+2 on right, the upper anterior temporal on this side being coalesced with the parietal; eight supralabials, fourth and fifth entering eye; five lower labials in contact with chin-shields, of which the posterior are slightly longer than the anterior; 17 rows of smooth scales; 203 ventrals; anal divided; 96 pairs of subcaudals. Color, above olive greenish brown; a yellow black-bordered longitudinal line beginning near the anterior end of the frontal along the middle of

^a See this work, figs. 301-303.

the back nearly to the end of the tail, occupying the median dorsal scale row and the adjacent halves of the rows on either side; an indistinct cross band at the posterior edge of the prefrontals, preoculars and postoculars, an indistinct narrow longitudinal band on the temporals, the greater part of the supralabials and the whole underside yellow; on the upper part of the supralabials a black irregularly serrated band which is interrupted on the fifth supralabial and again appears, though narrower, on the posterior portion of the neck; a black spot anteriorly on the lateral angle of each ventral; an indistinct blackish line on the end of the ventrals and that part of the first scale row lying between them, the scales themselves white in the middle so as to form a pale longitudinal line above the black one.

Dimensions.

	<i>mm.</i>
Total length.....	835
Snout to vent.....	615
Vent to tip of tail.....	220

Variation.—This species seems to be subject to the usual variations, such as “preocular separated from or just touching the frontal,” “temporals 1 + 2, 2 + 2, or 2 + 3.” Boulenger says: “Eight (exceptionally nine) upper labials, fourth and fifth (or fifth and sixth) entering the eye,” basing his diagnosis on four specimens, while Doctor Wall, who apparently examined 11 specimens, probably mostly from the Yangtse Valley, says that the normal arrangement is nine, with the fifth and sixth touching the eye, and that only two specimens had eight supralabials. He also mentions one specimen with two loreals, one behind the other, and one with the subpreocular absent on one side; two specimens had four lower labials in contact with anterior chin-shields on both sides. All his specimens had 2 + 2 temporals. With regard to the number of supralabials, it may be added that Doctor Werner in the two females, from Tsingtau, examined by him found eight. It is greatly to be regretted that the exact localities of the other specimens recorded have not been given with each individual scale formula. Minimum of ventrals recorded by Doctor Wall is 179; maximum by Doctor Werner is 211; subcaudals both minimum and maximum by Doctor Wall is 84 and 100.

Habitat.—This species was originally described as from Mexico, but it has since been discovered that its real habitat is eastern Asia, where it is recorded from Siam, the island of Hainan, the Yangtse Valley, and from northern China, Mongolia, and Korea. A specimen from the latter country is in British Museum.

List of specimens of *Zamenis spinalis*.

Museum.	No.	Sex.	Locality.	When collected.	By whom recorded.	Scale rows.	Ventrals.	Anal.	Caudals.
St. Petersburg.			Ali-shan Desert.		Strauch, l. c. p. 46	17	180	2	98
Berlin.		(a)	"Mexico"		Peters.	17	203	2	96
British.	a	Female	Korea.		Boulenger, Cat. I, p. 394.	17	198	2	85
Do.	b	Male	Near Peking.		do.	17	189	2	99
Do.	c	do	North China.		do.	17	192	2	90
Do.	d	do	Hainan.		do.	17	198	2	90
Munich.		Female	Tsingtau.		Werner, l. c.	17	210	2	90
Do.		do	do.		do.	17	211	2	91

^a Type; description, p. 350.

Genus ZAOCYS^a Cope.

1860. *Zaocys* COPE, Proc. Phila. Acad., 1860, p. 563 (type, *Z. dhumnades*).

1864. *Zapyrus* GUENTHER, Rept. Brit. India, p. 256 (type, *Z. fuscus*).

1891. *Zaocys* W. L. SCLATER, Journ. Asiatic Soc. Bengal, Nat. Hist., LX, p. 238 (error typogr.).

ZAOCYS DHUMNADES^b (Cantor).

1842. *Coluber dhumnades* CANTOR, Ann. Mag. Nat. Hist., IX, p. 483 (type-locality, Chusan, China; type in Brit. Mus.; Cantor, collector).—*Zaocys dhumnades* COPE, Proc. Phila. Acad., 1860, p. 563 (Ningpo).—GUENTHER, Rept. Brit. India, 1864, p. 256, pl. xxii, fig. A (Chusan; Ningpo).—BOULENGER, Cat. Snakes Brit. Mus., I, 1893, p. 375, pl. xxvi, fig. 1 (Southern China).—WALL, Proc. Zool. Soc. London, 1903, p. 90 (Shanghai; Yangtse Valley).—WERNER, Abh. Bayer. Akad. Wiss. (Murnchen), Klasse II, XXII, Pt. 2, 1904, pp. 357–364 (China), 372, footnote (Tamsui, Formosa).

1854. *Ablabes vittatus* DUMÉRIE and BIBRON, Erpét. Gén., VII, Pt. 1, 1854, (p. 326) (type-locality, China; type in Paris Mus.; Montigny, collector).

1858. *Coryphodon carinatus* GUENTHER, Cat. Colubr. Snakes Brit. Mus., p. 112 (part: Chusan).

Having no specimen of this species for description, I submit the following diagnosis taken from Boulenger:^c

Description.—Rostral broader than deep, visible from above; internasals shorter than the prefrontals; frontal once and one-fourth to once and a half as long as broad, as long as or a little shorter than its distance from the end of the snout, a little shorter than the parietals; loreal longer than deep; one preocular, with a subocular below it; two postoculars; temporals 2+2 or 2+1; eight upper labials,

^a From ζα, a prefix meaning very; and ὠκύς, swift.

^b Cantor gives no explanation of the name. Perhaps he meant to indicate a snake being like his *Coluber dhumna* (= *Ptyas mucosus*), the name of which in Hindu is said to be *dhamin*.

^c Cat. Snakes Brit. Mus., III, p. 375.

fourth and fifth entering the eye; five lower labials in contact with the anterior chin-shields, which are shorter than the posterior; scales in 16 rows, the two median strongly keeled; a second dorsal pair may be feebly keeled; ventrals 187-194; anal divided; subcaudals 108-116. Color, anterior half of body olive above, with black edges to the scales and a yellow or orange, black-edged vertebral stripe; upper lip and belly yellow, the olive of the upper parts extending down to the ends of the ventrals; on the posterior part of the body the upper surfaces turn to black and the lower to dark olive or gray.

	<i>Dimensions.</i>	<i>mm.</i>
Total length.....		1,950
Snout to vent.....		1,420
Vent to tip of tail.....		530

Variation.—Doctor Werner has counted as many as 120 subcaudals in a Ningpo specimen and Doctor Wall as few as 96; the latter also gives 199 as maximum of ventrals observed by him. He also notes the following variations: Loreal, in one specimen two superposed shields on one side; temporals, a single anterior in one specimen on one side; anterior chin-shields in contact with four lower labials on both sides in one specimen.

Habitat.—Southern China, from Shanghai to Hongkong on the coast west to Lun-ngan-fu, province of Sze-chuen, in the interior.

Doctor Werner has recorded its occurrence in Formosa on the strength of a specimen in the Museum of the Munich Academy of Sciences, said to be from Tamsui.

Genus *HOLARCHUS*^a Cope.

1854. *Simotes* DUMÉRIl and BIBRON, *Erpét. Gén.*, VII, Pt. 1, p. 624 (type, *S. russelii* = *arnensis*) (not of Fischer, 1817).

1887. *Holarchus* COPE, *Bull. U. S. Nat. Mus.*, No. 32, p. 54 (no species mentioned).

1893. *Dicraular* COPE, *Amer. Natural.*, 1893, p. 480 (type, *Holarchus trinotatus* = *Simotes purpurascens*.)

The generic name *Simotes*, by which the snakes of this genus have long been designated is preoccupied by *Simotes* of Fischer for a group of mammals as early as 1817. It has consequently to be replaced. Cope proposed *Holarchus*, in 1887, as a term for those species of the genus which have an undivided anal. It is not believed that this character alone, which moreover is not always constant, is sufficient ground for a division of the genus, and as *Holarchus* is the name next in date after *Simotes* it must stand for the combined genus.

^a From ὅλος, whole, undivided; ἀρχός, anus.

HOLARCHUS FORMOSANUS^a (Guenther).

1863. *Simotes swinhonis* SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 225 (nomen nudum; not of Guenther, 1864) (Tamsui, Formosa).
1872. *Simotes formosanus* GUENTHER, Ann. Mag. Nat. Hist. (4), IX, Jan. 1872, p. 20 (type-locality, Takao, Formosa; type in Brit. Mus.; Swinhoe, collector).—FISCHER, Abh. Naturw. Ver. Hamburg, IX, 1886, p. 12 (south Formosa; Oldenburg Mus.; Ruhstrat, collector).—MUELLER, Verh. Naturf. Ges. Basel, VIII, Pt. 2, 1887, p. 263 (south Formosa).—BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 129; Kat. Schl. Mus. Senckenberg., 1898, p. 73 (Hainan I., China).—BOULENGER, Cat. Sn. Brit. Mus., II, 1894, pp. 222, 359, pl. VIII, fig. 2 (Takao, Formosa; Swatow, China); III, 1896, p. 640 (central Formosa).—STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 221 (Taipa, Formosa).—SCHENKEL, Verh. Naturf. Ges. Basel, XIII, Pt. 1, 19—, p. 161.
1878. *Simotes bicatenatus* MUELLER, Verh. Naturf. Ges. Basel, VI, Pt. 4, 1878, p. 595 (Fumun, Prov. Kanton, not of Guenther).
1894. *Simotes hainanensis* BOETTGER, Ber. Senckenberg. Naturf. Ges., 1894, p. 133, pl. III, figs. 2a-c (type-locality, Hainan; type in Mus. Senckenberg.; Schmacker, collector).

Sauvage in 1877 has described a *Simotes* from China as *S. vaillanti*.^b The chief points of difference seem to be the double anal and single preocular, but as these characters are not absolutely constant in *S. formosanus* it may possibly turn out to be the same species.

This species seems also closely allied to *H. violaceus* which occurs on the continent from Bengal to southern China, and which differs chiefly in having only 17 scale rows and fewer subcaudals. The two species appear to occur together in some localities, thus, for instance, in Hainan.^c


Description.—*Adult male*: Science College Museum, Tokyo, No. 17A.; Taipa, Formosa; September, 1897; T. Tada, collector (figs. 304-306). Rostral higher than broad, projecting, turned over on top of the snout, the portion visible from above nearly as long as the internasal and prefrontal sutures; internasals oblique, much broader than long, smaller than prefrontals, which are in contact with supraocular; frontal very large, twice as broad as supraoculars, nearly as broad as long, longer than distance from tip of snout, and nearly as long as parietals, which are broad and truncate behind; nostril round, between two nostrils of which the anterior is much the larger; loreal higher than long; one preocular, not in contact with frontal; one subpreocular; two postoculars, the lower ones on both sides in this specimen abnormally coalesced with fifth supralabial; temporals 1 + 2, on right side an additional small anterior temporal; eight supralabials, all higher than long, the first three lower, the fourth suddenly higher, sixth,

^a From Formosa.

^b Bull. Soc. Philom. Paris (7), I, p. 107.

^c Cope's *Holarchus dolleyanus* (Proc. Phila. Acad., 1894 (publ. 1895), p. 423, pl. x, fig. 1), with 17 scale rows and 36 subcaudals, seems to belong to *Holarchus violaceus*.

seventh, and eighth largest, fourth and fifth entering eye; three lower labials in contact with anterior chin-shields, which are much longer than posterior; 19 rows of perfectly smooth scales; 167 ventrals, strongly angulate laterally; anal undivided; 46 pairs of subcaudals. Color (in alcohol) above pale drab with numerous narrow and serrate chocolate-brown crossbars dissolved laterally into small spots; a somewhat indistinct longitudinal, median whitish stripe more or less interrupted by the dorsal crossbars; head above more clay-colored with a somewhat intricate symmetrical pattern of brown dark-edged figures which are again outlined by a light edge, viz, an inverted V on the nape with the apex on the interparietal suture; a median figure on frontal like an exclamation mark; an oblique band on each side across parietal, posterior temporals, and hind edge of last supralabial; a pale dark-edged line across anterior part of frontal and supraoculars through the eye to the upper lip, crossing the suture between fifth and sixth supralabials; an angular mark with apex toward the nostral on internasals



FIGS. 304-306.—*HOLARCHUS FORMOSANUS*. 2 X NAT. SIZE. 304, TOP OF HEAD; 305, SIDE OF HEAD; 306, UNDERSIDE OF HEAD. No. 17A. SCI. COLL. TOKYO.

and prefrontals; and a few dark marks on some of the supralabials; underside whitish shaded with pale drab toward the lateral ventral angle, which is thus marked off as a fairly distinct white line.

Dimensions.

	<i>mm.</i>
Total length.....	557
Snout to vent.....	449
Vent to tip of tail.....	108

The *younger* specimens, for instance No. 34045, are much darker in coloration, the color above being dark drab with a rufous median dorsal line interrupted by very short, blackish crossbars; sides posteriorly nearly uniform, with a narrow dusky longitudinal line on the fourth scale row; markings on head essentially as in adults.

Variation.—Occasionally specimens are found with a divided anal.

The parietals in the three specimens examined by me show the curious characteristic of being in contact with the lower postocular

behind the upper one, except on one side in two examples in which the outer part of the parietal is separated off so as to form a small abnormal upper temporal of the first row.

The ventrals, in twelve recorded specimens, vary between 162 and 173, the subcaudals between 46 and 55.

Habitat.—Originally described from the island of Formosa, where it has been recorded from Tamsui, Takao, Taipa, and now from Byoritsu, on the strength of a specimen in the U. S. National Museum recently acquired from Mr. A. Owston. It has since been found in the province of Kwangtung, southern China, and in the adjacent island of Hainan.

List of specimens of Holarchus formosanus.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supralabials.
U. S. N. M. . . .	34045	Halfgr. . .	Byoritzu, Formosa. . .	1903. May 25	A. Owston	19	167	1	46	1-2+2	7-8
Do.	36509	Female.	Taipa, Formosa	1897. Sept. —	T. Tada . .	19	165	2	47	1+2	8
Sci. Coll., Tokyo.	17A	Male σ . .	do.	do.	do.	19	163	1	53	1-2+2	8

^a Description, p. 354; figs. 304-306.

Genus *DINODON* ^a Duméril.

1853. *Dinodon* DUMÉRIL, Mém. Acad. Sci., Paris, XXIII (p. 463), author's separate, p. 67.—DUMÉRIL and BIBRON, Erpét. Gén., VII, Pt. 1, p. 447 (type, *D. cancellatum*).

1860. *Eumesodon* COPE, Proc. Phila. Acad., 1860, p. 262 (type, *E. semicarinatus*).

1860. *Lepidocephalus* HALLOWELL, Proc. Phila. Acad., 1860, p. 498 (same type).

1893. *Dianodon* COPE, Amer. Natural., XXVII, May, 1893, p. 481 (emendation).

The genus *Dinodon* differs chiefly from *Ophites* Wagler ^b in the dentition, inasmuch as the former has a double interruption of the maxillary series, while in the latter only the anterior maxillaries are separated from the posterior series by a single interval.

^a From $\delta\iota\varsigma$, double; $\nu\omega\delta\acute{o}\varsigma$, toothless; referring to the two toothless spaces in each upper jaw.

^b This is the correct name for the genus which many authors, including Boulenger, call *Lycodon* Boie. The latter name appears first in Ferrussac's Bulletin des Sciences Naturelles, 1826, p. 238, unaccompanied by any diagnosis, but embracing with certainty only three species then known, namely: "*Col. audax* Daud.—*C. Hebe* Daud.—*C. aulicus* Linn." One of these is the type. In the following year Boie's diagnosis of the genus was published in the Isis, 1827, p. 521, and the type is expressly given as *Col. audax*. This certainly fixes the name *Lycodon* beyond a doubt, and it will consequently take precedence over *Lycognathus* for the South American snakes of the latter genus. The name next in time for the Asiatic genus is *Ophites* Wagler (Syst. Amph., 1830, p. 186), which was based on the single species *L. subcinctus* Boie.

Ophites, which, according to what has been shown, is equivalent to *Lycodon*, as hitherto limited by Boulenger and others, is not represented with certainty within the territory covered by this work. However, reference must be made to two species which have been credited to it, namely, *Ophites albofuscus*, of which Guenther says^a that it "occurs also in Formosa," the other being Mueller's "*Eumesodor. carinatus* Cope (?)" from Japan.^b

As to *Ophites albofuscus*, I have found no further corroborative evidence, and as this species seems to be confined to Sumatra and Borneo, Guenther's specimens probably belonged to some other species. It is then permissible to guess that he referred to the two specimens of *Dinodon septentrionale ruhstrati*, which Swinhoe sent from Formosa.

Mueller's "*Eumesodon carinatus* Cope (?),^c Japan, juv." in the Basel Museum,^d judging from his description of the dentition ("the second tooth and the one before the last are somewhat stouter and longer than the others") is not a "*Eumesodon*" (*Dinodon*) at all, but in all probability an *Ophites*. The description does not exactly tally with any of the known species, but comes probably nearest to *O. subcinctus*, from which it differs, however, in having smooth scales "except in the sacral region [Lendentheil], where there are three rows of tectiform scales" and two anterior temporals. The most noteworthy similarity is probably that the "upper preocular is fused with prefrontal," a feature quite characteristic of *O. subcinctus*, with which the rest of the scale formula also agrees, viz, Sc. 17; v. 211; a. 1 (exceptional); c. 80. As for the locality, "Japan," it should be noted that it is simply a surmise of Mueller's, based upon the fact that the snake was found in a jar which also contained a Japanese mole (*Talpa vogura*). The jar, however, was labeled "Java," and Java is just the type-locality of *Ophites subcinctus*. It should be noted, however, that Mueller later^e referred the specimen to *Dinodon orientale* (*Ophites japonicus*).

Dinodon is a very compact little genus of four species. Its center of distribution seems to be China. One species may extend as far west as the Himalayas, one is thus far found only in Hondo, Japan, another in Kiusiu and the Riu Kius, while a third extends from Hainan in the south to Vladivostok in the north.

^a Ann. Mag. Nat. Hist. (4), I, 1868, p. 426.

^b As to *Ophites* [*Lycodon*] *aulicus*, see footnote on p. 358.

^c Cope has given no such name, so far as I know; probably Mueller meant *E. semicarinatus*.

^d Verh. Naturf. Ges. Basel, VII, 1885, p. 685.

^e Idem, VIII, Pt. 2, 1887, p. 270.

The genus is not found in any portion of the Malayan region. In the latter it is represented by the genus usually known as *Lycodon*.^a

KEY TO THE SPECIES OF DINODON.

a¹ Anal entire; a preocular.

b¹ Scales smooth; less than 210 ventrals.

c¹ Ventrals 189 (185)–209 (average 198); cross-bars on body and tail more than 60 *D. rufozonatum*, p. 358

c² Ventrals 178–195 (average 189); cross-bars on body and tail less than 60.

D. rufozonatum walli, p. 364

b² Scales feebly keeled; more than 210 ventrals.

c¹ Portion of rostral visible from above half as long as the interprefrontal suture.

D. semicarinatum, p. 366

c² Rostral just visible from above *D. septentrionale*, p. 370

a² Anal divided; no preocular *D. orientale*, p. 372

DINODON RUFOZONATUM b (Cantor).

1840. *Lycodon rufozonatus* CANTOR, Zool. Chusan (pl. xi) (type-locality, Chusan, China; type in Brit. Mus.; Cantor, collector).—GUENTHER, Rept. Brit. India, 1864, p. 319 (Chusan); Ann. Mag. Nat. Hist. (4), I, 1868, p. 426 (Formosa).—JAN, Icon. Ophid., livr. 36, 1870, pl. iv, fig. 3 (Chusan).—STEINDACHNER, Sitz. Ber. Wien Akad. Wiss. Math. Naturw. Cl., LXII, Pt. 1, 1870, p. 338; author's separate, p. 13 (Peking).—MUELLER, Verh. Naturf. Ges. Basel, X, Pt. 1, 1892, p. 205 (Vladivostok).—*Dinodon rufozonatus* PETERS, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1881, p. 89.—BUTLENGER, Cat. Snakes Brit. Mus., I, 1893, p. 361 (Hainan; Formosa; Tsushima; northern China).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 38 (part: Formosa).—WALL, Proc. Zool. Soc. London, 1903, pp. 89, 100 (part: Formosa; Korea; Tsushima).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 354 (Ningpo Mts. near Shanghai).

1854. *Dinodon cancellatum* DUMÉRIl and BIBRON, Erpét. Gén., VII, Pt. 1, p. 477 (type-locality unknown; type in Paris Mus.).

1856. *Coronella striata* HALLOWELL, Proc. Phila. Acad., 1856, p. 152 (type-locality, Ningpo, China; types in Phila. Acad. Mus.; McCarter, collector).—*Eumecesodon striatus* COPE, Proc. Phila. Acad., 1860, p. 263.

1885. *Dinodon rufozonatus* var. *formosana* BOETTGER, Offenbach. Ver. Naturk., 24–25 Ber., p. 125 (type-locality, Formosa; type in Mus. Senckenb.); 26–28 Ber., 1888, p. 144 (Formosa).

^a *Lycodon aulicus* (*Coluber aulicus* LINNÆUS, Syst. Nat., 10 ed., I, 1758, p. 220) is credited to Formosa by Doctor Wall (Proc. Zool. Soc. London, 1903, p. 88) on the strength of a specimen in the Hongkong City Hall Museum now so labeled. He states that the specimen from Hongkong, which Boettger, upon the authority of von Moellendorff, recorded as being in that museum (Offenbach. Ver. Naturk. 26–28 Ber., 1888, p. 84) is found there no longer, but, on the contrary, that the only specimen of *L. aulicus* in that institution is from Formosa. It is greatly to be apprehended that the specimen in question, or rather its label, has been a victim of the typhoon mentioned by Doctor Wall in the introduction to his paper (p. 84). He there expressly states that the circumstances render the accuracy of some of the records open to question. In view of the probability that the original Hongkong specimen and the one now labeled as from Formosa are one and the same, I find it inadvisable to admit *Lycodon aulicus* to the fauna of Formosa on the strength of this record.

^b Signifying with red belts, or bands.

Description.—*Adult male*; U.S.N.M. No. 14614; Korea?; Dr. N. M. Ferebee, collector. Rostral about one and a half times as high, the portion visible from above equaling about one-third its distance from frontal; internasals very much smaller than prefrontals, which are broadly in contact with supraocular; frontal short and broad, its length equaling its distance from rostral and shorter than interparietal suture, its width anteriorly nearly equaling its length, and its width at a line through the center of the eyes twice as great as that of the supraoculars at the same line; parietals large, twice as long as prefrontals; nostril large, between two nasals, of which the posterior is the larger; loreal twice as long as high, entering eye below the small preocular, which is widely separated from frontal; two post-oculars; temporals 2 + 3, the upper one of the second row bordering the parietal very large; 8 supralabials, third, fourth, and fifth entering eye, sixth and seventh subequal, largest; five lower labials in contact with anterior chin-shields, which are considerably longer than the posterior; 17 rows of scales, which are smooth except a few median dorsal rows on the posterior fourth of the body, which show faint keels at the base of the scales; 197 ventrals; anal entire; 71 pairs of sub-caudals. Color (in alcohol) pinkish buff (in life said to be coral red) with about 58 broad, dark brown cross-bars on upper side of body and 23 on tail, the bars being separated by narrow bands, about the width of a scale, of the ground color; a series of alternating dark blotches on the sides involving the four outer scale rows and the upturned end of the ventrals; top of head isabella-color with ill-defined pale borders to many of the shields; labials and underside uniform pale, that of the tail with irregular dusky blotches.

Dimensions.

	<i>mm.</i>
Total length.....	1,063
Snout to vent.....	863
Vent to tip of tail.....	200

In a *younger* specimen, from the country between Tientsin and Peking (No. 29701), all the dark markings are solid black, even the top of the head and the light edges to the head shields are better contrasted. There is, moreover, a somewhat ill-defined black transocular spot from nostril to the temporal region with radiating black edges to the supralabials.

Variation.—With the exception of the scutellation of the sides of the head this snake is not subject to great variation. Thus of 29 recorded specimens only one has 2 scale rows over the normal number, 17.^a The number of supralabials 8 is fairly constant, 7 occurring only

^a Dr. J. S. Thompson, U. S. N., in a letter dated February 9, 1907, informs me that he has a specimen from Formosa with 19 scale rows.

rarely. Temporals also are nearly always 2 + 3, but Boettger records a Formosa specimen with 2 + 2 and Doctor Wall one specimen with a single temporal in the first row. He also mentions a specimen with three postoculars on both sides and one with three on one side. The greatest variation is seen in the relation of the loreal and the consequent relation of the supralabials to the eye. In our specimens it reaches the eye in one, but not in the other. In 16 Chinese specimens examined by Doctor Wall it touched the eye in 11 and did not reach it in 5, while in the large series of Chinese specimens recorded by Boulenger, viz, 23, the proportion must have been quite different, as in the description of the species he says "loreal elongate, sometimes entering the eye." The resultant variations in the relation of the supralabials to the eye are given by Boulenger as rare, viz, second, third, and fourth, or only fourth and fifth entering eye. The number of lower labials in touch with anterior chin-shield is also subject to some variation, Doctor Wall having found four on both sides in one, and six on one side in two specimens. The normal range of variation in the number of ventrals seems to be between 189 and 209, 185 as recorded by Boulenger in a young Shanghai specimen being quite abnormal; 57 subcaudals in another specimen of the same origin is also considerably below the normal range, which appears to be 62 to 83. It will be noted that the Formosa specimens average a greater number of subcaudals than the specimens from the mainland of China, but the overlapping is not only very great but appears to be uncorrelated with other characters.

The number of dark cross-bars on body and tail varies considerably within the limits of 68 and 99 in the specimens in which they have been recorded, the greatest range being on the body, viz, 50 to 75, as against 18 to 24 on the tail.

Habitat.—The present species extends over a large portion of eastern China, including the islands of Hainan,^a Formosa, and Chusan, ranging northward at least as far as Vladivostok, from which locality a specimen has been recorded by Mueller as being in the Basel Museum. It also occurs in Korea, but no definite locality has been recorded, the only specimen of certain Korean origin being in the Michigan University museum.

Holst sent two specimens from Tsushima to British Museum.

It has not been recorded from Japan proper. A fine specimen of the continental type, in the United States National Museum (No. 14614), collected by Dr. N. M. Ferebee, U. S. Navy, is credited on the record book to Nagasaki, but there is no original label or any other original document to corroborate this locality of this specimen which came with a collection all the other specimens of which are from Korea. As

^a Regarding this locality, see under *Elaphe dione*, p. 318, footnote.

the entries of that particular time when this specimen was received are not to be entirely depended on, it seems most probable that this specimen was really collected in Korea.

In Formosa the species seems to be common and to agree with the continental form in the number of dorsal spots. British Museum has numerous specimens from the island, the Senckenberg Museum has one, and the Hamburg Museum several collected by Doctor Warburg both in the southern and the northern part of Formosa (Nos. 1526, 1527). Similarly, the museum in Christiania has two specimens collected by Mr. Navara.

List of specimens of *Dinodon rufosomatum*.

Museum.	No.	Sex and age.	Locality.	When collected or received.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supratrabials.	Dark cross-bars.
U. S. N. M.	14614	(c)	Korea.		Dr. N. M. Ferebee.	17	197	1	71	2+3	8	81
Do.	29701	Male	Peking, China.		M. L. Robb.	17	200	1	68	2+3	8	68
Mich. Univ.	30527	do.	Korea.		G. A. Olinger.	17	208	1	75	2+3	7-8	99
Brit. Mus.	w	do. b.	Tsu shima		P. A. Holst.	17	204	1	68			
Do.	x	do. b.	do.		do.	17	202	1	68			
Do.	y	do. b.	Formosa.		R. Swinhoe.	17	189	1	76			
Do.	b	do. b.	do.		do.	17	190	1	83			
Do.	c	Young b.	do.		do.	17	199	1	71			
Do.	d	Female b.	do.		Taiwanfu Mus.	17	199	1	71			
Do.	e	do. b.	do.		do.	17	198	1	73			
Do.	f	Halfgr. b.	do.		do.	17	204	1	75			
Do.	g	Young b.	do.		do.	17	193	1	73			
Do.	h	do. b.	do.		do.	17	199	1	81			
Do.	i	do. b.	do.		do.	17	192	1	76			
Do.	k	do. b.	do.		do.	17	190	1	79			
Senckenberg.	7361 b	Adult c.	do.		F. Knoblauch.	17	205	1	76	2+2	7-8	79
Paris.		do. d.	?		Doctor Smith.	17	194	1	68		8	70
Brit. Mus.	l	Halfgr. e.	Chusan, China.		Doctor Cantor.	17	193	1	72			80
Do.	m	Male f.	Shanghai, China.		R. Swinhoe.	17	190	1	65			
Do.	n	Young f.	do.		do.	17	198	1	57			
Do.	o	do. f.	do.		do.	17	201	1	68			
Do.	p	do. f.	do.		do.	17	185	1	69			
Do.	q	Halfgr. f.	Kiukiang Mountains, China.		C. Maries.	17	203	1	67			
Do.	r	Female f.	Mountains north of Kiukiang, China.		A. F. Pratt.	17	203	1	72			
Do.	s	Male f.	Provinces of Szechuen, China.		do.	17	204	1	72			
Do.	t	Female f.	West Hills, Peking, China.		S. W. Bushell.	17	194	1	65			

Brit. Mus.	u	Female /	West Hills, Peking, China.	S. W. Bushell	17	163	1	65
Do.	r	do. /	do.	do.	17	204	1	62
Do.	a	Male ♀	Hainan Island, China	J. Neumann	17	202	1	73
Munich		Female ♀	Ningpo Mountains, China.	Doctor Haberer	19	198	1	65

♀ Boulenger, Cat. II. p. 361.
 ♂ Werner, p. 354.

♂ Boulenger, Cat. II. p. 362; type.
 ♀ Boulenger, Cat. II. p. 362.

a Description, p. 339.
 b Boulenger, Cat. II, p. 362.
 c Boettger, p. 125; type of *D. var. formosana*.
 d Duméril and Bibron, p. 449; type of *D. cancellatum*.

DINODON RUFOZONATUM WALLI,^a new subspecies.


1895. *Dinodon rufozonatus* BOETTGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 108 (Miyakoshima group, Riu Kius) (not of Cantor); Kat. Schl. Mus. Senckenberg., 1898, p. 38 (part: Miyakoshima).—WALL, Proc. Zool. Soc. London, 1903, pp. 89, 100 (part: Japan, Loo Choos; 1905, II, p. 515 (Miyako; Ishigaki; Iriomote).

Diagnosis.—Similar to typical *Dinodon rufozonatum*, but with fewer ventrals, viz, 178-195 (average 189), and fewer dark cross-bars on body and tail, viz, 22 to 32 on body and 15 to 20 on tail, or together on both, 41 to 52 (average 45); a pale inverted V on the nape; underside more or less mottled with brown (figs. 307-309).

Type.—U.S.N.M. No. 34007; Ishigaki shima, Yaeyama group; June, 1899; Owston collection.

Habitat.—Riu Kiu Archipelago.

Remarks.—While in Frankfort on the Main in 1898 Doctor Boettger had the great kindness to show me the specimens of this species in the



FIGS. 307-309.—*DINODON RUFOZONATUM WALLI*. 2 × NAT. SIZE. 307, TOP OF HEAD; 308, SIDE OF HEAD; 309, UNDERSIDE OF HEAD. No. 11, SCI. COLL. TOKYO.

Senckenberg Museum. Unfortunately, I did not have time to examine them in detail, but I made the note that the Formosa specimen appeared to have about twice as many dorsal spots (56 according to Boettger) as those from the Riu Kius. This observation I have had an opportunity to test on only eight additional specimens, but in these I find a similar proportion inasmuch as in five specimens from the southern Riu Kius,^b the dorsal spots are only 32, 29, 29, 25, and 26, respectively, while in two specimens from Korea and one from northern China, Tientsin, or Peking, the numbers are 75, 58, and 50.^c

Doctor Wall, who examined a large number of Chinese specimens and four from Ishigaki shima (three of which are the ones now in the

^a Named for Capt. F. Wall, of the Indian Medical Service, author of a *Prodromus* of the snakes hitherto recorded from China, Japan, and the Loo Choo Islands, as well as several papers on Indian snakes.

^b Science College Museum, Nos. 11, 12, and U.S.N.M. Nos. 34006-8.

^c For further color distinctions, see remarks under *Variation*, p. 365.

U.S.N.M. mentioned above), as well as one alleged to be from "Japan," made similar observations, and also called attention to the lower number of ventrals and higher number of subcaudals in the latter. The numbers of these plates as given by him do not overlap, but from the tables presented by me (pp. 362 and 366) it will be seen that there is considerable overlapping. While it may thus be inexpedient to give the Riu Kiu form a binominal appellation, there can be no doubt that it should be recognized nomenclatorially.

It appears that all the names in the synonymy of *D. rufozonatum* are based upon Chinese or Formosan specimens with more than 192 ventrals and less than 73 subcaudals. The type localities of *D. rufozonatum*, *D. r.* var. *formosana*, and *C. striata* are known and agree in the scale formula. The type of *D. cancellatum* is from an unknown locality, but the scale formula as well as the number of black bars on back and tail are conclusive. Hence the necessity of giving a new name.

The number of ventrals and dorsal dark cross-bars being the chief distinctive characters of this subspecies, as indicated above, it appears superfluous to present a detailed description of a Riu Kiu specimen, but a few remarks on the individual variation within the form peculiar to the archipelago may not be out of place.

Variation.—In the five specimens before me the loreal is in every case excluded from the eye by the preocular. That this relation is not constant, however, is shown by Boettger's statement that in some of the four Miyakoshima specimens examined by him there is found two preoculars on one side, or on both sides in some, while mostly there is only one. Other variations noticed by him are three postoculars on one side in one (also found in one of the specimens before me), and only two supralabials entering the eye in another. Ventrals vary between 178 and 195 in nine recorded specimens; subcaudals between 73 and 87. The largest number of dark cross-bars in five specimens is 52, the smallest 41.

In the youngest specimen before me (Sci. Coll. No. 11) the dark markings on the upper side are nearly solid, though even in this many of the scales in the interior of the blotches have narrow pale edges. In all the other specimens, however, which are much larger, these pale edges are much broader, so that the cross-bars appear lighter brown with dark margins, the paler interior having mostly a streaky appearance.

The typical *D. rufozonatum* from the mainland seems to have the cross-bars solid-colored.

Habitat.—This race seems to be confined to the southern and middle groups of the Riu Kiu Archipelago.

Mr. Tashiro has collected specimens in the Yaeyama group, both on Iriomote shima and on Miyakoshima. The Senckenberg Museum

also has specimens from the latter island and the Hamburg Museum from the former (No. 2579). United States National Museum has three specimens from Ishigaki shima, and Doctor Wall examined three more from the same island. Finally, Doctor Warburg also collected it in Okinawa shima. (Hamburg Mus. No. 1545.)

Doctor Wall also mentions a specimen in Owston's collection as from "Japan." No other locality is given, and until further evidence is produced it is permissible to question the occurrence of this species in Japan proper.

List of specimens of Dinodon walli.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supraorbitals.	Dark cross-barr.
				1899.								
U.S.N.M.	34006	Male	Ishigaki shima	June	A. Owston	17	195	1	82	2+3	8	45
Do.	34007	Male ^a	Riukiu	do.	do.	17	190	1	87	2+3	8	43
Do.	34008	Female	do.	do.	do.	17	186	1	81	2+3	8	41
Sci. Coll., Tokyo.	11	Young ^b	Iriomote shima, Riukiu.	June 14	Tashiro.	17	192	1	85	2+3	8	52
Do.	12	Male	Miya koshima, Riukiu.	do.	do.	17	189	1	73	2+3	8	44
Senckenb.	7361	^a (c)	do.	do.	B. Schmacker	17	178	1				
Do.		(c)	do.	do.	do.	17	181	1				
Do.		(c)	do.	do.	do.	17	192	1	83			
Do.		(c)	do.	do.	do.	17	195	1	81			

^a Type, p. 364.

^b Figs. 307-309.

^c Boettger, p. 108.

DINODON SEMICARINATUM ^a (Cope).

AKAMATAH (on OKINAWA SHIMA, According to Mr. Tashiro).

MATSUTABU (on AMAMI-O-SHIMA, According to Mr. Nakagawa).

1860. *Eumesodon semicarinatus* COPE, Proc. Phila. Acad., 1860, p. 263 (type-locality, "Loo Choo;" type in Phila. Acad. Mus.; Heine, collector); 1861, p. 75 (identity with *L. fasciatus*).—*Dinodon semicarinatus* BOULENGER, Ann. Mag. Nat. Hist. (6), X, Oct. 1892, p. 302 (Okinawa shima); Cat. Snakes Brit. Mus., I; 1893, p. 362 (Oho shima; Great Loo Choo Island).—BROWN, Proc. Phila. Acad., June 11, 1902, p. 185 ("Loo Choo Islands").—WALL, Proc. Zool. Soc. London, 1903, p. 100 (Loo Choos); 1905, II, p. 515 (Okinawa; Amami).

1860. *Lepidocephalus fasciatus* HALLOWELL, Proc. Phila. Acad., 1860, p. 498 (type locality, "Japan and Loo Choo;" types, U. S. Nat. Mus. No. 7354).

The Perry expedition and the Rodgers expedition, according to Hallowell, brought home four specimens of this species, namely, two from "Loo Choo" by the former, two from "Japan" by the latter. One

^a Signifying half-carinated, with reference to the scales being keeled only in their basal half.

of the former was apparently donated to the Philadelphia Academy and served as type of Cope's *E. semicarinatus*; the other three are still in the U. S. National Museum as No. 7354, but the identity of each specimen was not maintained, and it is now impossible to point out with certainty their individual origin. So much is certain, however, that the specimen now designated as No. 7354c is the type of Hallowell's *L. fasciatus*, his description being based on that one alone. I venture the guess that this is the second of the Perry expedition specimens and that the other two are Stimpson's specimens from "Japan."

There is indirect evidence that Hallowell was mistaken about these two specimens of *Lepidocephalus fasciatus* being from "Japan" as distinct from "Loo Choo." Altogether he recorded only the following land snakes as being collected by the Rodgers expedition (W. Stimpson):

From "Loo Choo:"

P. 492. *Bothrops flavoviridis*, 1 specimen from Amakarima Island, and

P. 493. *Eurypholis semicarinatus*, 2 specimens from near Napa, Loo Choo, caught by Mr. Wright, Nov., 1855.

From "Japan:"

P. 498. *Leptophidium dorsale*, 1 specimen, "taken on a hillside near Hakodadi, Island of Jesso, June, 1855, by W. Stimpson."

P. 498. *Lepidocephalus fasciatus*, 2 specimens.

P. 499. *Amphiesma tigrinum*, 2 specimens "caught at Nippon, Japan, May, 1855, by Mr. Stimpson."^a

Dr. W. Stimpson, in his manuscript catalogue of specimens brought home by the Rodgers Expedition, enumerates the following specimens:

From "Loo Choo":

No. Am. 2. Amakirrima, April, 1855.

γ -200. Snakes taken near Napa, Loo Choo, Nov., 1854 (Wright).

γ -199. Snake shot in a paddyfield, Loo Choo, Nov., 1854 (W. S.).

γ -187. Snake caught at Loo Choo, under a stone in low land, Nov., 1854 (W. S.).

From Japan:

No. γ -249. Snake taken on a hillside near Hakodadi, I. of Jesso, Japan, June 1855 (W. S.).

γ -275. Simoda, Island of Nippon, Japan, May, 1855 (Mr. Brooke).


γ -277. Simoda, Island of Nippon, Japan, May, 1855 (W. S.).

It is plain that No. Am. 2 is *Bothrops flavoviridis*; No. γ -200 is *Eurypholis semicarinatus*; No. γ -249 is *Leptophidium dorsale*; and Nos. γ -275 and γ -277 are the two specimens of *Amphiesma tigrinum*. Thus all the specimens enumerated by Hallowell and Stimpson are accounted for except the latter's Nos. γ -199 and γ -187 from "Loo Choo" or Okinawa shima, and the former's two specimens of

^a The other snakes recorded are *Elaphis bilineatus*, collected by Doctor Morrow, and *Proterodon tessellatus*, without any record of locality or collector.

Lepidocephalus fasciatus. There can scarcely be the slightest doubt, therefore, that they are identical, and that Hallowell's locality, "Japan," for this species, is erroneous.

Description (figs. 310-311).—*Adult male*; U.S.N.M., No. 7354a: paratype of *Lepidocephalus fasciatus*; Okinawa shima; November, 1854?; W. Stimpson, collector? Rostral a little less than twice as broad as high, the portion visible from above equaling about two and a half times its distance from frontal and about one-half the interprefrontal suture; internasals small, less than half as large as the prefrontals which are broadly in contact with supraoculars; frontal broad and short, slightly longer than broad, its length less than interparietal suture and equaling its distance from rostral, the anterior outline nearly straight, the lateral outline curved to the posterior apex; supraoculars narrow, much narrower in front than behind; parietals long, nearly as long as frontal and prefrontal together;



FIGS. 310-311.—*DINODON SEMICARINATUM*. $\frac{1}{4}$ X NAT. SIZE. 310, TOP OF HEAD; 311, SIDE OF HEAD. No. 7354c, U.S.N.M.

nostril large, between two nasals of which the posterior is much the larger; loreal long and narrow, twice as long as high, upper and lower edges parallel; one small preocular widely separated from frontal; two postoculars; temporals 2 + 3; the upper temporal of the third row very large, bordering the posterior half of the parietal externally; 8 supralabials, third, fourth, and fifth entering eye, the upper portion of the third being greatly extended posteriorly for the purpose, sixth and seventh largest; five lower labials in contact with anterior chin-shields which are somewhat longer than the posterior; 17 rows of scales, the four outer rows on each side smooth, the other with a feeble though distinct keel on the basal half of each scale; 229 ventrals with a strongly marked lateral angle; anal entire; 100 pairs of subcaudals. Color (in alcohol) pale yellowish with 40 dark cross-bars on the body and about twenty on the tail, these bars having solid blackish margins while the scales inside the bars have yellow

centers; alternating with these cross-bars there is a series of lateral blackish spots mostly on the four outer scale rows and more or less connected with an irregular series of single or double spots on the upturned ends of the ventrals; nape black with two yellowish spots on the outer posterior angle of the parietals and the adjacent scales; top of head blackish with ill-defined yellowish markings; sides of head yellowish, most of the shields, including the supralabials, narrowly edged with blackish; underside uniform pale yellowish, that of the tail with irregular blackish brown spots medially.

Dimensions.

	mm.
Total length.....	880
Snout to vent.....	690
Vent to tip of tail.....	190

The *younger* specimen, Sci. Coll. Mus., Tokyo, No. 10, is essentially like the adult. It grows to a considerable size, as a specimen from O-shima with defective tail measures 1,510 mm. in length.

Variation.—This snake does not seem to be subject to great variation, as the only abnormalities in the specimens recorded are two posterior temporals on both sides in one, nine supralabials on one side in another specimen and only two supralabials, namely, fourth and fifth, touching the eye in a third specimen. The ventrals vary between 211 and 234, and the subcaudals between 92 and 105 in the perfect specimens on record. The lowest extreme of the latter is given by Boulenger as 65, but this figure is derived from Hallowell's account of the type of *Lepidocephalus fasciatus* in which, however, the tail is mutilated.

Habitat.—Both the Perry Expedition and the Rodgers Expedition brought home specimens of this species, those from the latter being said to hail from "Japan" in contradistinction to the former as being from the Riu Kius, but as shown above this statement is undoubtedly erroneous. Later collectors, including Mr. Tashiro, have also obtained it in Okinawa shima, and in British Museum there is a specimen from "Oho shima" collected by Lieut. A. Carpenter, R. N., from which island Mr. Nakagawa also brought a specimen in 1891 to the Science College Museum. Recently other specimens from Okinawa shima and Amami-o-shima have been recorded by Dr. Wall.

In the National Museum, Ueno Park, Tokyo, there is a specimen (No. 23) of this species said to be from the province of Hiuga, southern Kiusiu. If the locality given is correct, it is the only record of the species in Japan proper, but the occurrence there needs confirmation.

List of specimens of *Dinodon semicarinatum*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supratuberc.
U.S.N.M.	7354a	Male a	Okinawa shima.	1854? Nov. —	Rodgers Exped?	17	229	1	100	2+3	8
Do.	7354b	Female	do.	do.	do.	17	215	1	100	2+3	8
Do.	7354c	do. b	do.	1853?	Perry Exped?	17	218	1	100	2+3	8
Sci. Coll. Tokyo.	4	Male c	O-shima.	1891	Nakagawa.	17	233	1	100	2+3	8-9
Do.	10	Young.	Okinawa shima.	1887	Tashiro.	17	229	1	105	2+2	8
Do.	54	Adult.	do.	Feb. —	do.						
Do.	54	do.	do.	Apr. —	do.			1			
Nat. Mus. Tokyo.	23		Hiuga, Kiusiu(?)			17		1			
Brit. Mus.	a	Male	Amami-o-shima		Lieut. A. Carpenter.	17	234	1	97		
Do.	b	do.	Okinawa shima.		Holst.	17	228	1	101		
Do.	c	Young.	do.		do.	17	211	1	93		
Phila. Acad.		(d)	"Loo Choo".		Perry Exped. (W. Heine).	17	221	1	92	2+	8

a Description p. 368.

b Type of *L. fasciatus*; figs. 310-311.

c Total length 1,510 mm.; tail defective.

d Cope, 1860, p. 263; type.

DINODON SEPTENTRIONALE^a RUHSTRATI^b (Fischer).

1868. *?Ophites albofuscus* GUENTHER, Ann. Mag. Nat. Hist. (4), I, 1868, p. 426 (Formosa; Brit. Mus.) (not of Duméril and Bibron).
1886. *Ophites ruhstrati* FISCHER, Abh. Naturw. Ver. Hamburg, IX, Pt. 1, no. 6, p. 16, pl. II, fig. 6 (type-locality, south Formosa; types in Mus. Oldenburg; Ruhstrat, collector).
1893. *Dinodon septentrionalis* BOULENGER, Cat. Sn. Brit. Mus., I, p. 363 (part: Formosa).
1899. *Dinodon septentrionalis* var. *ruhstrati* BOULENGER, Proc. Zool. Soc. London, 1899, p. 165 (Formosa).

Doctor Guenther, as quoted above, in a paper dealing with new additions to the British Museum snake collection, mentions *Ophites albofuscus* with the statement: "Occurs also in Formosa; feeds on lizards." In Boulenger's catalogue there is no indication of any specimens of this species from Formosa being in British Museum. I take it, therefore, that Guenther mistook Swinhoe's specimens from Formosa for this species, a supposition the more probable as he afterwards in describing *Ophites septentrionalis* referred it to the same genus. *O. albofuscus* is restricted to Sumatra and Borneo (see p. 357).

^a *Ophites septentrionalis* GUENTHER, Proc. Zool. Soc. London, 1875, p. 233 (type-locality, unknown, but supposed to be "northern India," Boulenger giving "Himalayas or Khasi Hills (?);" Jerdon, collector).

From *Septentrionalis*, signifying northern.

^b For Mr. Ruhstrat, who collected the types in southern Formosa.

The collective species *Dinodon septentrionale* is recorded from the eastern Himalayas or Khasi Hills, in Assam, and the Karen Hills, in Burma, to eastern China and Formosa.

The exact relation of the specimens from the various localities is not very well known, but there seem to be several color forms, of which the easternmost and the westernmost are the extremes, viz, the typical subspecies *D. septentrionale* from Assam or Himalaya, and Burma, on the one side, and *D. septentrionale ruhstrati*, from Formosa, on the other. Specimens from the Chinese coast province of Fokien, according to Boulenger,^a are "intermediate in the pattern of coloration" between these two forms. The status of the form which occurs in the mountains of Kiukiang is still uncertain on account of scantiness of the material.

It is unfortunate that the full scale formulas of the three Fokien specimens has not been published, as altogether the formulas of only seven specimens have been recorded, and the more so since the latter seem to indicate a possible difference in the number of subcaudals in one of the races. I find, namely, that in the three Formosan specimens in which the tail was complete the subcaudals number, respectively, 103, 104, and 97, while in the three continental specimens on record they are stated to be 83, 87, and 88.

Under these circumstances it seems best to maintain, for the present at least, the subspecific distinctness of the Formosan specimens.

A partial translation of Doctor Fischer's original description of the scutellation and coloration of the Formosan specimens is subjoined, as I have not seen any specimens of this form.

Description.—South Formosa; Mus. Oldenburg; Ruhstrat, collector. Rostral just reaching the upper side of the snout; internasals slightly broader than long, longer than the prefrontals; frontal pentagonal, as broad as long, shorter than the interparietal suture; nostril between two nasals, of which the posterior is much higher than the anterior; loreal pentagonal, twice as long as high, its posterior angle reaching under the preocular; latter well developed, quadrangular, twice as high as long, resting on the third supralabial and part of the loreal, extending to the upper surface of the head, though not reaching the frontal; two postoculars, the lower resting on fifth and sixth supralabials; temporals 2 + 3, the two anterior in contact with both postoculars; eight supralabials, of which third, fourth, and fifth join the eye; ten infralabials, the first six in contact with the chin-shields; the posterior chin-shields somewhat narrower and but slightly shorter than the anterior; between these and the ventrals two or three rows of elongated throat scales; scales in 17 rows, only the six or eight median rows^b weakly keeled; ventrals

^a Proc. Zool. Soc. London, 1899, p. 165.

^b Must be "seven or nine," as the number of scale rows is uneven.—L. S.

221 to 223, laterally with a distinct but not sharp angle; anal single; 103 pairs of subcaudals, followed by a horny tip. Head to the middle of the frontal deep black, thence to the third scale row on the nape gray in the old specimen, pure white in the quite young one; ground color above reddish gray; numerous (47 to the anus in the old, 36 in the young) deep brown, light-edged crossbands descending on the sides almost to the ventral surface, the spots as in *O. orientalis*, cut on the posterior two-thirds of the body by a light longitudinal line on the fourth scale row, so that the portions thus cut off form a separate row of spots; alternating with these the upturned portion of many ventrals have a small brown spot, thus forming another series of spots on each side; the tail above also has a series of dark brown transverse spots, which in the young specimen extend to the ventrals, forming (18) complete rings; the brown dorsal spots gradually diminish in width posteriorly, the first one on the nape extending over 17 scales, while those following become gradually shorter, until on the second third of the body the light interspaces exceed them in width (especially in the young specimen); underside of head and body yellowish white, the tail of the old specimen grayish brown, of the young divided by complete dark rings.

Dimensions.—Young specimen: Total length, 250 mm.; tail, 60 mm.

Habitat.—The distribution of the species has been given above under the preliminary remarks.

The subspecies, *D. ruhstrati*, as here understood, is confined to the island of Formosa, where it has been collected by Mr. R. Swinhoe, and later by Mr. Ruhstrat, the specimens brought home by the former being in British Museum, while those by the latter are in the natural history museum in Oldenburg.

List of specimens of Dinodon ruhstrati.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.
Oldenburg.....	a	Adult ^a	South Formosa.....		Ruhstrat.....	17	221	1
Do.....	b	Young ^a	do.....		do.....	17	223	1	103
Brit. Mus.....	b	Female ^b	Formosa.....		R. Swinhoe.....	17	216	1	104
Do.....	c	do. ^b	do.....		do.....	17	224	1	97

^a Cotypes; Fischer, p. 16.

^b Boulenger, Cat. 1, p. 363.

DINODON ORIENTALE ^a (Hilgendorf).

SHIROMADARA (Doctor IJIMA, In Letter).

1880. *Ophites orientalis* HILGENDORF, Sitz. Ber. Ges. Naturf. Fr., Berlin, 1880, p. 115, pl. —, figs. 1-5 (type-locality, Tokyo; type, Berlin Mus. No. 9419; Hilgendorf, collector).

^a Signifying eastern.

1880. *Ophites japonicus* GUENTHER, Ann. Mag. Nat. Hist. (5), VI, Dec., 1880, p. 462 (type-locality, Nikko, Hondo; types, in Brit. Mus.; Maries, collector).—MUELLER, Verh. Naturf. Ges. Basel, VIII, Pt. 2, 1887, p. 270 (Japan).—*Dinodon japonicus* BOULENGER, Cat. Snakes Brit. Mus., I., 1893, p. 363 (Nikko, Hondo).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 38 (Hondo).—WALL, Proc. Zool. Soc. London, 1903, p. 100 (Kiusiu); 1905, II, p. 513 (Yamanashi, Hondo).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, No. 1 (1905), p. 224 (Nagasaki).

Hilgendorf's *Ophites orientalis* was described in the Sitzungs-Bericht of the meeting held October 19, 1880, which was probably published very shortly after that date. Guenther's *O. japonicus*, on the other hand, appeared in the December number of the Annals and Magazine of the same year. There is, therefore, apparently a very brief priority in favor of Hilgendorf's description. It may be interesting to note that Hilgendorf as early as 1876 indicated this species, but refrained from naming it.^a

Description.—*Young*; U.S.N.M. No. 34052; Mount Fuji; August, 1898; Owston collection. Rostral broader than high, the portion visible from above scarcely as long as suture between internasals; the latter small, less than half as long as prefrontals, the suture between the former only half as long as that between the latter; prefrontals broadly in contact with supraoculars and with eye; frontal as broad as long, and as long as its distance from rostral, shorter than interparietal suture; supraoculars small, less than half the size of frontal; parietals large, nearly twice as long as frontal; nostril large, round, between two subequal nasals; loreal narrow, twice as long as high, entering eye below prefrontal; no preocular; two postoculars; temporals, 2+3; a very large upper temporal of the third row bordering the posterior half of parietal externally; 8 supralabials, sixth largest, fourth and fifth entering eye; 5 lower labials in contact with anterior pair of chin-shields, which are somewhat larger than posterior; 17 rows of scales, all of which are smooth on the anterior half of the body, but on the posterior half the median 5 to 7 rows have a faint keel at the base of each scale; 207 ventrals, distinctly angulate laterally; anal divided; 74 pairs of subcaudals. Color (in alcohol) above dull ecru-drab^b with dark brown cross-bars, 38 on the body, 15 on the tail, these bars being broader than the intervening light ground color on the anterior part of the body, but becoming gradually narrower posteriorly so as to be narrower than the light interspaces; the latter are paler colored along the edges of the dark cross-bars; on the posterior half of the body the lower part of the bars are separated off by a narrow light line between fourth and fifth scale row; on the sides of the middle third of the body there is an

^a Mitth. Deutsch. Ges. Ost-Asiens, I, heft 10, July, 1876, p. 30.

^b Hilgendorf says: "Oberseite in frischem Zustande, schmutzig braunroth, ziemlich hell."

alternating series of vertical brown spots between the lower end of the cross-bars; top of head uniform dark brown; lower part of supralabials and entire temporal region, including the posterior lateral portion of the parietals, whitish; underside whitish with a few dusky marblings on the middle and occasional dark spots on the sides of the ventrals of the posterior half; dorsal cross-bars nearly meeting on underside of tail.

Dimensions.

Total length.....	<i>mm.</i> 290
Snout to vent.....	235
Vent to tip of tail.....	55

The *adult* specimens are essentially like the one described above, but this snake does not seem to reach any considerable size. Boulenger measures a total length of 660 mm. and the species probably does not grow much larger.

Variation.—Comparatively little variation is shown in the scutellation of this species, and then mostly in the temporals, the second row occasionally having only two scales. There seems also to be some variability in the number of supralabials in touch with the eye, inasmuch as both Hilgendorf and Boulenger state that in their specimens the third also enters into relation with it. In all those specimens in our museum, however, the third supralabial is excluded from the eye, the loreal being in broad contact with the fourth supralabial. Ventrals vary between 199 and 208; subcaudals between 68 and 75 pairs.

Habitat.—Apparently restricted to Japan proper. Hilgendorf in describing the species mentioned seven specimens from the neighborhood of Tokyo, mostly from the Botanical Garden, and for this reason he suspected that the species had been accidentally introduced. This suspicion seems to be unfounded in view of the many specimens which have been found in various localities both in Hondo and Kiusiu. The two specimens in the British Museum were collected by Mr. Maries at Nikko; Doctor Nordquist found a specimen at Murayama, and United States National Museum has a specimen each from Isobe near Yokohama and from "Mount Fuji;" a specimen (No. 5) from the province of Iga is in the Imperial Museum, Ueno Park, Tokyo. Finally Doctor Wall mentions one specimen from Yamanashi, Hondo, and one from Kiusiu. It was collected there as early as 1879 by Dr. O. Nordquist near Nagasaki, where another specimen, now in the St. Petersburg Museum, was collected by Doctor Slunin in 1888. Both the Nordquist specimens are in the Riksmuseum at Stockholm, where I was permitted to examine them in 1905, thanks to the kindness of the curator, Dr. Einar Loennberg.

List of specimens of *Dinodon orientale*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supralabials.
U. S. N. M.	34052	Young a	Mount Fuji	1898. Aug. —	A. Owston	17	207	2	74	2+3	8
Do	34543	Halfgr.	Isobe, near Yokohama.	1904. May 10	do	17	201	2	—	2+2	8
Do	36542	do. b.	Tokyo	—	Sci. Coll., Tokyo	17	203	2	73	2+2	8
Sci. Coll., Tokyo.	64	Female	do	—	—	17	199	2	75	2+3	8
Stockholm	418	do	Murayama, Hondo	1879. Nov. 27	Dr. O. Nordquist	17	201	2	72	2+3	8
Do	419	Male	Nagasaki, Kiusiu	—	do	17	203	2	74	2+2	8
Berlin	9419	Young c.	Tokyo	1874. May —	Doctor Hilgendorf.	17	200 or 208	2	74	2+2	8
Brit. Mus.	a	Female d	Nikko	—	C. Maries	17	202	2	68	2+3	8
Do	b	do	do	—	do	17	205	2	70	2+3	8

a Description, p. 373.

b Sci. Coll., No. 64a.

c Type, Hilgendorf, p. 116.

d Cotypes of *O. japonicus*, Boulenger, Cat. I, p. 364.Genus CALAMARIA^a Boie.1827. *Calamaria* BOIE, Ferrusac, Bull. Sci. Nat., IX, p. 236 (type, *Coluber calamarius*).1834. *Changulia* GRAY, Ill. Ind. Zool., II, (pl. LXXXVI), (type, *C. albiventer*).

The range of the genus embraces the Malayan archipelago, extending into Burma, southern China, and the Philippine archipelago. One species is from the southern and middle group of the Riu Kiu archipelago, and one is here recorded from Formosa for the first time.

Of the Calamarias with four supralabials there is a small compact group containing *C. brevis*, from an unknown locality; *C. pavimentata*, from Java; *C. siamensis* from the Indo-China peninsula and southern China; *C. berezowskii* from the upper Yangtse Valley and Formosa; *C. septentrionalis* from the lower Yangtse and the coast of China; and *C. pfefferi* from the Riu Kiu Archipelago. Boulenger has united *C. pavimentata* and *C. siamensis* under the former name, but while I have no specimens for comparison, the figures and descriptions show such contradictions that I can not consider the question finally settled. A long narrow head is characteristic of *C. brevis* and *C. pfefferi*, which also agree in the lack of a yellow collar and light spots. They are probably nearly related. On the other hand *C. siamensis*, *berezowskii*, and *septentrionalis* are very closely allied and I have some doubts whether they are properly separated and whether the specimens from

^a Signifying like a reed, from *καλαμος*, a tube, a straw.

China and Indo-China may not be grouped to better advantage and more consonant with their geographical distribution, questions I am unable to solve at present on account of lack of material.

The two species which are found within our territory may be distinguished as follows:

- a*¹ Head wide; the width of parietals together more than half the distance from tip of snout to their posterior end; suture of frontal with supraocular shorter than with prefrontal; a yellowish, more or less interrupted nuchal collar; ventrals uniform pale *C. berezowskii*, p. 376.
- a*² Head narrow; the width of the parietals together equals one-half their distance from tip of snout to their posterior end; suture of frontal with supraocular longer than with prefrontal; no collar; ventrals strongly spotted with black.

C. pfefferi, p. 378.

CALAMARIA BEREZOWSKII^a Guenther.

1896. *Calamaria berezowskii* GUENTHER, Annuaire Mus. Zool. St. Pétersbourg, I, p. 205, pl. I, fig. A (type-locality, Luu-ngan-fu, Prov. Sze-chuen, China; types in Mus. St. Petersb.; Berezowski, collector).

A specimen of a *Calamaria* which seems to belong to this species, collected by Mr. Tsunasuke Tada at Taipa, Formosa, was accidentally omitted from my account of the collection made by him,^b and is here introduced into the Formosan fauna for the first time.

There is nothing in Doctor Guenther's description or figures of *C. berezowskii* by which I can separate this specimen from it. One would naturally expect the Formosa specimen to agree with *C. septentrionalis* which occurs from Hongkong, on the mainland opposite, to Chusan in the north and to the mountains north of Kiukiang on the Yangtse River, in the interior, but this form is described as having the rostral scarcely visible from above and the tip of tail rounded. Our specimen has the same wide frontal as *C. berezowskii*, which according to Guenther is the character distinguishing it from *C. siamensis*. The latter has been taken as far north as Canton.

Curiously enough, the Formosa specimen shows no special inclination towards *C. pfefferi* from the Riu Kius, and it is quite likely that a form of the latter type may eventually turn up in Formosa also.

Description.—*Adult male*; Science College Museum, Tokyo, No. 12; Taipa, Formosa; September, 1897; T. Tada, collector (fig. 312). Rostral high as broad, the portion visible from above more than half as long as suture between prefrontals; no internasals; prefrontals slightly smaller than frontal, in contact with first and second supra-labials; frontal slightly longer than broad, the width equaling its distance from tip of snout and the interparietal suture, about three times

^a Named after Mr. M. Berezowski, the Russian explorer, who in 1892-1894 traveled in the Chinese provinces of Kansu and Sze-chuen. During this journey he collected the types of this species.

^b Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, pp. 215-225.

as wide as supraoculars, six-sided forming an obtuse angle in front and a nearly right angle behind, sutures with supraoculars shorter than those with prefrontals; parietals in contact with fourth supralabial, shorter than their distance from tip of snout, rather broad; nostril small, in a very small and narrow triangular nasal between rostral, first supralabial and prefrontal; one preocular; one postocular; no temporals; a large shield bordering posterior half of parietal behind fourth supralabial; four supralabials, first and third subequal, small, second twice as large as third, fourth as long as second and third together, second and third entering eye; first lower labials in contact with each other behind mental; three lower labials in contact with anterior chin-shields, which are in contact with each other and as long as the posterior; 13 rows of smooth scales, without pits; 160 ventrals; anal undivided; 23 pairs of subcaudals; tail rather tapering, pointed at tip. Color (in alcohol) above cinnamon-brown, each scale darker edged so that the whole upper surface appears finely reticulated, the margins on the sides more emphasized on the lateral scales so as to almost form narrow longitudinal lines; on upper neck two large yellowish spots, or an interrupted yellowish collar, four scales from the head, and on the posterior angle of each parietal a similar but smaller round spot; top of head and occiput between parietals and collar much darker brown; upper and lower labials irregularly marked with dark brown; underside uniform whitish, ventrals with the lateral edge brownish like the scales; subcaudals marked with dark brown so as to form an ill-defined median band; no pale spots on tail.




FIG. 312.—CALAMARIA BEREZOWSKII. 4 × NAT. SIZE. TOP OF HEAD. No. 12, SCI. COLL. TOKYO.

Dimensions.

	<i>mm.</i>
Total length.....	195
Snout to vent.....	177
Vent to tip of tail.....	18

The *female* has shorter tail with fewer subcaudals.

Habitat.—This species is only known from the two types collected by Mr. Berezowski in the Chinese province of Sze-chuen, and the specimen brought home by Mr. T. Tada from Taipa, Formosa, where he obtained it in September, 1897. It is probably also to be found in the intervening territory.

List of specimens of Calamaria berezowskii.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Supralabials.
Sci. Coll., Tokyo.	12	Male ^a	Taipa, Formosa	1897. Sept. —	T. Tada	3	160	1	23	1, 1	4
St. Petersb.		(b)	Lun-ngan-fu, Szechuen.	1893.	M. Berezowski		155		22	1, 1	4
Do.		(b)	do.	1893.	do.		167		14	1, 1	4

^a Description p. 376.^b Type. Guenther, p. 205.**CALAMARIA PFEFFERI^a Stejneger.**

1901. *Calamaria pfefferi* STEJNEGER, Proc. Biol. Soc. Washington, XIV, Dec. 12, 1901, p. 191 (type-locality, Miyakoshima, Sake shima group, Riu Kiu; type, Science Coll. Mus. Tokyo, No. 14).

1903. *Calamaria parimentata* WALL, Proc. Zool. Soc. London, 1903, p. 101 (not of Duméril and Bibron).


Doctor Wall^b expresses the opinion that my *C. pfefferi* is nothing but *C. parimentata* [Boulenger, not Duméril and Bibron, = *siamensis* Guenther], but nothing shows better than this case how dangerous it is to "lump" species without having seen the material upon which they are based or without having at least material from the type locality. As a matter of fact, it is not to the form mentioned by Doctor Wall, but to Boulenger's *C. brevis* that our Riu Kiu species is most nearly related. With the latter it shares the long and narrow head as well as the plain upper surface and spotted underside, and the chief differences appear to be the shorter parietals and the low number of ventrals in the latter. Unfortunately, the habitat of the unique type of *C. brevis* is not known, and the exact relationship of these two forms must consequently remain unsettled until more material shall have been accumulated and direct comparisons between typical specimens instituted.

Description.—*Halfgrown female*; Science College Museum, Tokyo, No. 14. Miyakoshima, Riu Kiu archipelago; type (figs. 313–316). Rostral about as high as broad, the portion visible from above nearly as long as suture between prefrontals; no internasals; prefrontals smaller than frontal, in contact with first and second supralabials; frontal considerably longer than broad, the width equaling the distance from tip of snout but shorter than interparietal suture, about three and a half times as wide as supraoculars, six-sided, forming an

^a Named in honor of Dr. George Pfeffer, curator in the Natural History Museum, Hamburg.

^b Proc. Zool. Soc. London, 1903, p. 93, footnote.

obtuse angle in front and an acute one behind, sutures with supra-oculars longer than those with prefrontals; parietals in contact with fourth supralabial, longer than their distance from tip of snout; the suture between them twice as long as that between prefrontals, rather narrow, especially posteriorly; nostril small, in a small nasal between rostral, first supralabial and prefrontal; one preocular; one postocular; no temporals; a large single shield bordering posterior half of parietal behind fourth supralabial; four supralabials, fourth largest, three times as large as second which is somewhat larger than first and third, second and third entering eye; first pair of lower labials forming a suture behind mental: two pairs of subequal chin-shields in con-



FIGS. 313-316.—*CALAMARIA PFEFFERI*. 4 × NAT. SIZE. 313, TOP OF HEAD, WITH COLOR PATTERN; 314, SIDE OF HEAD; 315, UNDERSIDE OF HEAD, ENLARGED; 316, COLOR PATTERN AROUND MIDDLE OF BODY. No. 14, SCI. COLL. TOKYO.

tact with each other; scales smooth, in 13 rows without apical pits; 160 ventrals; anal undivided; 15 pairs of subcaudals; tail ending in a very short spine. Color (in alcohol) above brownish gray with 9 parallel darker stripes commencing directly behind the parietals; of these stripes the median one covers the entire width of the vertebral scale row, while the others only occupy the lines between the scale rows, viz, on each side, one between the ventrals and first scale row, and one between first and second and one between second and third rows, and finally one between fourth and fifth rows; head slightly paler than the body with a few rather irregular and ill-defined darker markings; an ill-defined dark band from nostril through eye and over upper portion of fourth supralabial; labials below this band and entire

underside pale yellowish, each ventral usually marked with two well-defined, but irregularly placed, dark brown spots which, however, show a tendency to an arrangement in two longitudinal rows; under the tail a broad well-defined median band of similar color occupying the suture between the subcaudals and also extending down the middle of the anal; no nuchal collar, nor any yellowish spots at base of tail.

Dimensions.

	<i>mm.</i>
Total length.....	162
Snout to vent.....	152
Vent to tip of tail.....	10

The *male* is distinguished by a longer tail with a larger number of subcaudals.

Variation.—A specimen from Okinawa shima in the Hamburg Museum (No. 1567) collected by Doctor Warburg is very similar in scale formula, viz, sc. 13; v. 158; a. 1; c. 15; oc. 1-1; l. 4; 1st pair of lower labials in contact behind mental; second pair of chin-shields in contact; frontal four times as broad as supraocular; parietals longer than frontal. Coloration is also essentially like the one described above, there being no collar, and no yellow spots at base of or at end of tail, but there are two dorsal stripes more; the median dark stripe is two scales wide and separated by an interspace of equal width on both sides from the other stripes, of which there are five on each side; belly irregularly spotted with dark brown; a broad dark stripe along middle of underside of tail.

Science College Museum No. 13, from Okinawa shima is also similar in coloration, lacking collar and spots on tail, but has 26 pairs of subcaudals, being a male.

Habitat.—Thus far this *Calamaria* is only known from the Riu Kiu Islands, where it has been collected both in the southern group, viz, on Miyakoshima, and in the middle group, on Okinawa shima, a specimen from each of these localities being in the Tokyo Museum. A third specimen, from the latter island where Doctor Warburg collected it, is in the Hamburg Museum.

List of specimens of Calamaria pfefferi.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Supralabials.
Sci. Coll. Tokyo.	14	Female	Miyakoshima, Riu Kiu.			13	160	1	15	1-1	4
Do.....	13	Male	Okinawa shima, Riu Kiu.						26		
Hamburg..	1567	(b)	do.....		Doctor Warburg	13	158	1	15	1-1	4

a Type; description p. 378; figs. 313-316.

b P. 380.

Subfamily BOIGINÆ.

There does not seem to be any convincing proof that these snakes, although provided with grooved fangs at the posterior end of the maxilla, are particularly close allies of the other opisthoglyph snakes, such as the *Homalopsinæ*. Nor do the *Langahinæ* which, like the latter and the Natricine snakes, have the hypopophyses present throughout the vertebral column, and which form a group chiefly confined to Madagascar, appear to belong here.

Like the coronelline subfamily the present one contains forms of very varied habits, arboreal, terrestrial, and subterranean, diurnal and nocturnal, etc.

Their distribution is chiefly tropical and southern both in the Old and the New World. The subfamily is therefore represented within our limits by only two genera, each with a single species, both recorded from Formosa.

They may be distinguished as follows:

- a¹ Maxillary teeth subequal in front of the enlarged posterior grooved fangs; nostril between two nasals; scales with apical pits; scale rows 21; more than 200 ventrals; anal double *Boiga kræpelini*, p. 381.
- a² Third or third and fourth maxillary teeth much enlarged; nostril in an undivided nasal; scales without pits; scales in 17, or rarely 19, rows; less than 180 ventrals; anal entire *Psammodynastes pulverulentus*, p. 383.

Genus BOIGA^a Fitzinger.

1826. *Boiga* FITZINGER, Neue Classif. Rept., pp. 29, 31 (type, *Coluber irregularis*).
1843. *Dipsadomorphus* FITZINGER, Syst. Rept., p. 27 (type, *Dipsas trigonata*).
1843. *Macrocephalus* FITZINGER, Syst. Rept., p. 27 (type, *Dipsas drapiezii*).
1843. *Gonyodipsas* FITZINGER, Syst. Rept., p. 27 (type, *Dipsas irregularis*).
1843. *Eudipsas* FITZINGER, Syst. Rept., p. 27 (type, *Dipsas cynodon*).
1843. *Cephalophis* FITZINGER, Syst. Rept., p. 27 (type, *Dipsas dendrophila*).
1853. *Opetiodon* DUMÉRIL, Prodr. Class. Ophid., p. 98 (type, *O. cynodon*).
1853. *Triglyphodon* DUMÉRIL, Prodr. Class. Ophid., p. 111 (type, *T. irregularis*).
1857. *Toxicodryas* HALLOWELL, Proc. Phila. Acad., 1857, p. 60 (type, *T. blandingii*).
1877. *Pappophis* MACCAY, Proc. Linn. Soc. New S. Wales, II, p. 39 (type, *P. laticeps*).
1895. *Liohallus* COPE, Proc. Phila. Acad., 1894, p. 427 (type, *Dipsas fusca*).

Fitzinger's *Boiga*, of 1826, clearly takes precedence over his *Dipsadomorphus* of 1843, as shown by me.^b

BOIGA KRÆPELINI^c Stejneger

1902. *Boiga kræpelini* STEJNEGER, Proc. Biol. Soc. Washington, XV, p. 16 (type-locality, Kelung, Formosa: type in Naturh. Hist. Mus., Hamburg, No. 1565; Doctor Warburg, collector).—*Dipsadomorphus kræpelini* WALL, Proc. Zool. Soc. London, 1903, p. 94 (Formosa).

^a "La Boiga" is a name given by the early French ophiologists to some brilliantly colored tropical snake. The name is probably of barbaric origin. Agassiz suggests its connection with *Boa*, but apparently with no good reason.

^b Proc. Biol. Soc., Washington, XV, 1902, p. 16.

^c Dedicated to Prof. K. Kræpelin, director of the Natural History Museum in Hamburg.

Description of type specimen.—Naturh. Mus. Hamburg, No. 1565; Kelung, Formosa; Doctor Warburg, collector. About 11 solid maxillary teeth, subequal, slightly separated from grooved fangs, which are not much enlarged; head very distinct from neck; eye large, equaling its distance from anterior border of nostril, with vertically elliptical pupil; rostral wider than high, barely visible from above; internasals much broader than long, much shorter than prefrontals; frontal as long as broad, as long as its distance from rostral and as the interparietal suture; nasals large, posterior concave; loreal higher than long; 2 preoculars, the upper one reaching the upper surface of the head, but separated widely from frontal; 2 postoculars; temporals 4+5, scale-like, irregular; 9 supralabials, of which the third, fourth, and fifth enter the eye; 5 (on one side 6) lower labials in contact with anterior chin-shields, which are much larger than the posterior pair; body compressed, about twice as high as wide; scales in 21 oblique rows, smooth, with apical pits, the vertebral row scarcely enlarged, the scales pointed behind; ventrals, 245, flat underneath, obtusely angulate laterally; anal double; subcaudals, 142 pairs. Color brownish gray, with about 57 darker cross-bars composed of black-edged scales from neck to anus, these cross-bars extending on the sides to about 4 scale rows from the ventrals; only faint indications of alternating lateral spots; top of head uniform brown; underside pale with a median area more grayish and laterally bordered by an irregular dusky line following the ventral angle.

Variation.—The color description of specimen No. 1569, Hamburg Mus., which was also collected by Doctor Warburg near South Cape, Formosa, is as follows: Ground color paler and more grayish than the type (No. 1565), with better defined cross bands, which alternate with a row of lateral spots approximately covering the third, fourth, and fifth rows from the ventrals; the ventral median area darker and better defined; head with a median dark line on internasal and prefrontal sutures and middle of frontal, reappearing on the anterior part of upper neck as a median elliptical, brown spot; a similar brownish band from posterior half of supraoculars posteriorly to side of neck, where it joins another originating on the upper part of the rostral and running obliquely through nostril and eye over posterior supralabials to side of neck; between these lines a pale gray band with whitish edges; supralabials also pale, more or less marked with dusky and with a dusky spot on the suture below the center of the eye.

The specimen in the Hongkong City Hall Museum examined by Doctor Wall had six temporals on one side and ten labials on one side; five lower labials in contact with anterior chin-shields on both sides.

Remarks.—Kræpelin's *Boiga* differs from most of the other species of the genus in having a double anal and a scarcely enlarged vertebral scale series; also in the very short posterior chin-shields, and especially in the numerous small temporals.

Habitat.—This species is only known from Formosa, where two specimens were collected by Doctor Warburg, one at Kelung, in the northern part, the other near South Cape. A third specimen from Formosa is in the Hongkong Museum, as reported by Doctor Wall.

List of specimens of Boiga kræpelini.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Temporals.	Supracaudals.
Hamburg ^a .	1565	Kelung, Formosa.....	Dr. O. Warburg.	21	245	2	142	4+5	9
Do. ^b	1569	Near South Cape, Formosa.....	do.....	21	232	2	143	5+4	9-10
Hongkong ^c	Formosa.....	21	244	2	140	9-10

^a Type: description, p. 382.

^b P. 382.

^c Wall, p. 94.

Genus PSAMMODYNASTES^a Guenther.

- 1858. *Psammodynastes* GUENTHER, Cat. Colubr. Sn. Brit. Mus., p. 140 (type, *P. pulverulentus*).
- 1904. *Thamnodynastes* WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, p. 372 (*lapsus*; not of Wagler, 1830).
- 1905. *Anisodon* ROSÉN, Ann. Mag. Nat. Hist. (7), XV, Feb., 1905, p. 176 (type, *A. lilljeborgi*) (not of Lartet, 1849).
- 1905. *Anisodontes* ROSÉN, Ann. Mag. Nat. Hist. (7), XVI, Aug., 1905, p. 128 (substitute).

PSAMMODYNASTES PULVERULENTUS^b (Boie).


- 1827. *Psammophis pulverulenta* BOIE, Isis, 1827, p. 547 (type-locality, Java).—SCHLEGEL, Phys. Serp. II, (1837), p. 211, pl. VIII, figs. 10-11 (Java).—*Psammodynastes pulverulentus* GUENTHER, Cat. Colubr. Sn. Brit. Mus., 1858, pp. 140, 251 (Philippines; Sumatra; India).—BOULENGER, Cat. Sn. Brit. Mus., III (1896), p. 172 (Eastern Himalayas, etc.; Formosa).
- 1839. *Dipsas ferruginea* CANTOR, Proc. Zool. Soc. London, 1839, p. 53 (type-locality, Assam; type? in Brit. Mus.; Cantor, collector).
- 1867. *Lycodon bairdii* STEINDACHNER, Novara Exped., Zool., I, Rept., p. 90 (type-locality, Philippines; types in Vienna Mus.).
- 1905. *Anisodon lilljeborgi* ROSÉN, Ann. Mag. Nat. Hist. (7), XV, p. 176, pl. XI, fig. 3 (type-locality, Tjibodas, Java; types in Lund Mus.; Hj. Moeller, collector).

Description (figs. 317-319).—*Adult female*; Brit. Mus. No. 94. 1. 25. 9.; near Taiwan fu, Formosa; Holst, collector. Rostral higher than wide, visible from above; internasals small, only one-third the size of the prefrontals, nearly triangular; prefrontals very large, broadly in contact with preocular and loreal; frontal

^a From ψάμμος, sand; δυνάστης, lord, ruler.

^b Signifying dusty.

small, narrow, urceolate, the posterior outline rounded, the lateral sutures greatly concave; supraoculars excessively large, much larger than frontal, very broadly in contact with prefrontals; parietals large, the suture between them longer than the frontal; nostril a round hole in the middle of a single nasal; canthus rostralis sharp, overhanging the very concave loreal region; a small lozenge-shaped loreal; a single, large, concave preocular; three postoculars; temporals 2+(1 small)+3, the lower anterior on left side coalesced with supralabial; mouth very deeply cleft, with 8 supralabials, first long, in contact with loreal, 4, 5, and 6 in contact with eye; 8 lower labials, 3 in contact with first pair of chin-shields, of which there are three pairs; scales smooth, without pits, in 17 rows; -ventrals 175;^a anal single; subcaudals 60.^a Color (in alcohol) brownish gray densely



FIGS. 317-319.—*PSAMMODYNASTES PULVERULENTUS*. 2× NAT. SIZE. 317, TOP OF HEAD; 318, SIDE OF HEAD; 319, UNDERSIDE OF HEAD. No. 34707, U.S.N.M.

spotted with dusky and pale ochraceous, forming a very obscure pattern; a dark narrow band from tip of snout to posterior end of frontal, followed by a V-shaped mark, the opening resting on the upper edge of the preoculars, the apex on the posterior end of the interparietal suture; a similar band, from upper postocular along outer edge of parietal; all the bands on the head composed of narrower dusky lines with a pale line between them; from apex of parietal V a solid dark line along the median line of the upper neck; an obscure dusky transocular band; underside brownish gray with two narrow dusky lines, one on each side of the median line; the grayish ground color is produced by innumerable dusky specks powdered all over the surface.

Dimensions.

	mm.
Total length.....	382
Snout to vent.....	313
Vent to tip of tail.....	69

^a According to Boulenger (MS.).

Variation.—The lateral head-shields are subject to considerable variation. Thus there is often a second loreal underneath the ordinary one; the temporals are more or less irregular in many instances; the number of preoculars is sometimes two, while the postoculars vary between two and four. In a large series of specimens from the whole range of the species the minima and maxima of the ventrals have been found to be 146 and 175, the latter in the Formosa specimen described above, and of the subcaudals respectively 44 and 66.

The coloration is very variable, especially the ground color, which varies from nearly uniform raw umber to nearly blackish, with intermediate shades spotted with darker and paler in great profusion.

Habitat.—Widely distributed from the eastern Himalayas to the Malay peninsula and archipelago, Indo-China, Philippine Islands, and Formosa.

The only record of its occurrence in the latter island is the female specimen in the British Museum, collected by Mr. Holst at Taiwan fu, and described above. I owe the opportunity to make this description to the kindness of Doctor Boulenger.

As yet *Psammodynastes pulverulentus* has not been recorded from the mainland of China and would thus appear to be an exception to the rule that the Formosan snakes have reached that island from China. On the other hand, its wide distribution in the Philippine Archipelago might tempt one to conclude that this species had arrived in Formosa from the south. With our present defective knowledge of the herpetology of China, such a conclusion would be too hasty, however. It is moreover a significant fact that this snake is at home in the eastern Himalayas, and in the mountains of Assam and Burma, where so many other Formosan species, also occurring in China, have their nearest relatives and probable origin. There is every reason to believe that *P. pulverulentus* may yet be discovered in the mountains of southern and western China.

List of specimens of Psammodynastes pulverulentus.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal	Subcaudals.	Temporals.	Supralabials.	Loreal.
Brit. Mus.	94.1.25.9	Female ^a	Near Taiwan fu, Formosa.	1896.	Mr. Holst.	17	175	1	60	2+3	8	1
U. S. N. M.	23753	do	Trong, Lower Siam.	Sept. —	Dr. W. L. Abbott.	17	161	(?)	57	2+2	8	1
Do.	26220	do	do	do	do	17	162	1	59	2+3	8	1
Do.	34707	Male ^b	Mount Apo, Mindanao, P. I.	1904. July —	Dr. E. R. Mearns.	17	156	1	60	2+3	8	2
Do.	34708	do	do	do	do	17	160	1	61	2+3	8	2
Do.	34709	do	do	July 9	do	17	163	1	55	2+3	8	2

^a Description, p. 383.

^b Figs. 317-319.

Family ELAPIDÆ.

This family contains all the so-called "proteroglyph" snakes, or snakes with a permanently erect grooved poison fang in the anterior portion of the horizontal maxillary bone.

The members of the family are divided into two subfamilies, one containing the terrestrial species which have retained the general physiognomy of the majority of snakes, the other embracing forms which have become strongly modified for marine life, as follows:

- a*¹ Tail cylindrical ELAPINÆ. p. 386.
*a*² Tail strongly compressed, oar-shaped HYDRINÆ p. 400.

Subfamily ELAPINÆ.

As stated above, the general physiognomy of these snakes is that of the great majority of snakes, the *Natricidæ*, and the resemblance of some of the species in each family is so complete as to have deceived experts on a superficial examination. The generally small eye with vertical pupil, frequent absence of a loreal, and even width of head with body will serve to characterize many of the Elapines, but the presence of the poison fang is the only reliable criterion.

It is the more to be regretted that these snakes can not be readily and at a glance distinguished from the numerous harmless species, as they are the deadliest and often the most dangerous of all snakes. Thus the Indian cobra, found in Formosa, and the so-called hamadryad belong to this subfamily. The "krait," which grows to be over 4 feet, and which is represented by a separate form in Formosa (*Bungarus multicinctus*) is also very destructive of life where it is common. On the other hand, the smaller species, although their poison in proportion is equally deadly, are not very dangerous to man, partly because of the smallness of their mouths, partly, it seems, because of the supposed gentleness of some of them.

The Elapine snakes are truly tropical in their distribution. Only one genus, *Elaps*, occurs in America, while on the other hand the majority of the Australian snakes belong to this family.

Only four species, belonging to four genera, come within our limits, viz, three in Formosa and one in the Riu Kius, although it would not be surprising if additional species were discovered in the former island.

They may be distinguished as follows:

KEY TO THE FORMOSAN AND RIU KIUAN GENERA.

- a*¹ Median dorsal scale row not enlarged; subcaudals all, or mostly, in pairs.
*b*¹ Neck not dilatible, not surrounded by more scale rows than middle of body; internasal not bordering the nostril.
*c*¹ One to three small teeth on maxillary behind fangs *Hemibungarus*, p. 387.
*c*² No teeth on maxillary behind fangs *Calliophis*, p. 391.
*b*² Neck dilatible; more scale rows around neck than around middle of body; internasal bordering the nostril *Naja*, p. 394.
*a*² Median dorsal scale row enlarged; subcaudals single *Bungarus*, p. 397.

Genus HEMIBUNGARUS^a Peters.

1843. *Brachyrhynchus* FITZINGER, Syst. Rept., p. 28 (type, *Elaps calligaster*) (not of LAPORTE, 1833).
 1862. *Hemibungarus* PETERS, Mon. Ber. Berlin Akad. Wiss., 1862, p. 637 (type, *Elaps calligaster*).

The only difference between the genera *Hemibungarus* and *Calliophis* appears to be the presence of several (1-3) small solid teeth on the maxillary bone behind the grooved fang in the former, and it is questionable whether these so-called genera really represent genetic groups. The two species occurring within our limits, namely, *Calliophis macclellandii* in Formosa and *Hemibungarus japonicus* in the Riu Kiu Islands seem to be so closely related to each other that the latter appears less nearly allied to any of the other species of *Hemibungarus*. The only difference, besides that of dentition, consists in the greater length of the parietals in *H. japonicus* and in a slightly different pattern of coloration. Thus far no *Hemibungarus* has been recorded from Formosa, but this island is as yet so imperfectly known that no conclusion can be based on this fact.

The relationship of these snakes (through *Calliophis*) is undoubtedly Himalayo-Chinese, but beyond this statement it would be hazardous to generalize for the present.

The two forms described in the following pages under the names of *H. japonicus* and *H. boettgeri* are evidently very closely allied in spite of their very different coloration. The scutellation is almost identical, though it must be pointed out that in the former (9 specimens) the number of ventrals averages higher, namely, 208.5 (maximum 216, minimum 196), the corresponding figures for the latter (11 specimens) being 193.5 (204 and 183). While the scale formulas thus show some overlapping, the difference in coloration is so striking and apparently so constant, without any intermediary specimens, that I do not feel compelled to use trinominals for these forms, especially in view of their geographical isolation, the former being confined to the Oshima subgroup, the latter to the Okinawa subgroup of the Riu Kiu Archipelago.

They may be distinguished as follows:

- a¹ Red (white in alcohol) with one, or three, narrow dark-brown longitudinal stripes, and distant rings of the same color; ventrals 196 to 216. *H. japonicus*, p. 387.
 a² Black, with four narrow longitudinal red stripes (whitish in alcohol) on back, and distant white half-rings; ventrals 184 to 200 *H. boettgeri*, p. 389.

HEMIBUNGARUS JAPONICUS^b (Guenther).

HAI (according to Dr. Ijima).

1868. *Calliophis japonicus* GUENTHER, Ann. Mag. Nat. Hist. (4), I, 1868, p. 428, pl. xvii, fig. c (type-locality, Nagasaki [?]; type in Brit. Mus.; Whitely, collector).— *Hemibungarus japonicus* BOULENGER, Cat. Snakes Brit. Mus., III, 1896, p. 395 (part: Oo-shima).— WALL, Proc. Zool. Soc. London, 1905, II, p. 515 (part: Anami-o-shima).

^a From *hū-*, half-; and *Bungarus*, a nearly related genus of snakes.

^b Signifying Japanese.

Description.—*Young:* U.S.N.M. No. 31817; Amami-o-shima, Riu Kiu Archipelago; Dr. H. M. Smith collection. Rostral broader than high, easily visible from above; internasals broader than long, the suture between them scarcely more than one-half that between the prefrontals which are broadly in contact with supraoculars; frontal longer than its distance from tip of snout and than interparietal suture, much broader than supraoculars; parietals very long, longer than frontal; nostril large, round, at the posterior margin of the anterior nostril, which is larger than the posterior and in contact with second supralabial; no loreal; a single preocular situated above the center of the eye in contact with posterior nasal; eye rather large, the diameter greater than its distance from edge of lip; two postoculars; temporals 1 + 1, second one very large; a large shield bordering the outer posterior half of parietal; 7 supralabials, sixth and seventh largest, third rather high, third and fourth entering eye; four lower labials in contact with anterior chin-shields which are as long as the posterior, fourth lower labial very large; 13 rows of smooth scales without apical pits; 196 ventrals; anal divided; 31 pairs of subcaudals. Color (in alcohol) pale cream-color (in life said to be red) with a narrow dark-brown median line, one scale wide, from frontal to tip of tail; 16 dark-brown rings around the body and tail, about 3 scales broad on back and two ventrals broad below, the intervals between them about three to four times as wide; top of head dark brown except the median portion of each parietal, the brown color on the sides of the head extending some distance past the last supralabial; on the underside a large, roundish blotch of dark brown extending over four ventrals between the rings.

Dimensions.

	<i>mm.</i>
Total length.....	221
Snout to vent.....	200
Vent to tip of tail.....	21

Judging from the *adult female* in British Museum, the full-grown specimens do not differ essentially from the young described above.

Variation.—The scutellation seems to be very constant, though the ventrals range between 196 and 216 in the nine specimens recorded. The subcaudals on the other hand only vary between 28 and 31 pairs.

The coloration is also rather constant, except that some specimens, for instance the type and the young specimen in British Museum, collected by M. Ferrié, have an additional dark-brown stripe along each side of the body. Its presence is not a sign of youth as shown by the specimen described above. The number of rings around the body varies slightly; thus in the specimen in the Imperial Museum in Ueno Park, Tokyo, there are 12 rings on the body, one at the vent and one on tail. In this specimen the dark bands on the back are plainly edged with whitish.

Since the above was written Doctor Wall has reported upon four specimens from Amami-o-shima, obtained by Mr. Owston's collectors. Of these two agreed in having three longitudinal dorsal black lines narrower than the white intervals. One of these had indications of another line in the flanks on each side. One had 12 and the other 14 black annuli. The ventrals and subcaudals were 208 + 29 and 198 + 28. Two others were pale pink with a single, narrow, vertebral black line. One of these had 15 and the other 13 black annuli, and the ventrals and subcaudals were 215 + 28 and 214 + 30.

Habitat.—This form appears to be restricted to the O-shima subgroup of the Riu Kiu Archipelago, inasmuch as all the specimens supplied with exact data have been taken in "Oo-shima," or as we prefer to call it, Amami-o-shima, in order to avoid confusion with the many other O-shimas.

Whitely's specimen, upon which Guenther founded the species, was said to have come from Nagasaki. We know now that all his specimens credited to that locality came from other places, the present one almost certainly from Amami-o-shima.^a

List of specimens of Hemibungarus japonicus.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supratemporals.
Brit. Mus.	a	Young. ^a	"Nagasaki" [error].	Whitely.....	13	205	2	29	7
Do.....	d	do. ^a	Amami-o-shima.	Ferrié.....	13	211	2	28
Do.....	e	Male ^a	do.....	do.....	13	216	2	28
Imper. To-kyo.	24	Young.	do.....	13	214	2	27	1, 2	1+1	7
U.S.N.M.	31817	do. ^b	do.....	H. M. Smith.....	13	196	2	31	1, 2	1+1	7

^a Type.

^b Description, p. 388.

HEMIBUNGARUS BOETTGERI ^b (Fritze).

HAI (according to Dr. Ijima).

1892. *Callophis japonicus* BOULENGER, Ann. Mag. Nat. Hist. (6), X, Oct. 1892, p. 302 (Okinawa) (not of Guenther).—BOETTGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 115 (Okinawa).—*Hemibungarus japonicus* BOULENGER, Cat. Snakes Brit. Mus., III, 1896, p. 395 (part: Okinawa).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 123 (Okinawa).—WALL, Proc. Zool. Soc. London, 1905, II, p. 515 (part: Okinawa).
1894. *Callophis boettgeri* FRITZE, Zool. Jahrb. Syst., VII, p. 861; author's separate p. 12 (type-locality, Tokuchimura, Okinawashima; type in Mus. Senckenberg.; Fritze, collector).—BOETTGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 115 (Okinawa).

^a Boulenger, Cat. Sn. Brit. Mus., III, p. 395, foot note, has already intimated that this specimen may have come from Riu Kiu.

^b For Dr. Oscar Boettger, the distinguished herpetologist and malacologist of Museum Senckenbergianum, in Frankfurt am Main.


Description.—*Adult female*: Fifth Higher Middle School collection, Kumamoto, No. B 25; Okinawa shima, Riu Kiu archipelago; 1891 (figs. 320-322). Rostral broader than high; frontal longer than its



320



322



FIGS. 320-322.—HEMIBUNGARUS BOETTGERI. $\frac{2}{3}$ X NAT. SIZE. 320, TOP OF HEAD; 321, SIDE OF HEAD. ENLARGED. 322, COLOR PATTERN AROUND MIDDLE OF BODY. NO. B25, HIGH SCHOOL, KUMAMOTO.

distance from end of snout; parietals very long, as long as frontal and prefrontals together, and more than twice the distance from frontal to rostral; one preocular; two postoculars; temporals 1 + 1; 7 supralabials, third and fourth entering eye; 4 lower labials in contact with anterior chin-shields; 13 scale rows; ventrals 193; anal divided; subcaudals 29 pairs. Color (in alcohol) above iridescent blackish blue with four longitudinal light bands, the two median bands being apparently reddish, the two outer ones white, the former commencing on inner anterior corner of parietals and continuing to end of tail, the latter beginning on sides of neck and ending on sides in front of anus;

across this longitudinal pattern there is a transverse one consisting of about fourteen (of which one on the tail) irregular cross bands of bluish black, edged with white, more or less interrupted by the longitudinal stripes so as to form two superimposed ocelli; these black cross bands are carried across the belly on a single ventral; underside whitish with numerous large and irregular blackish blue blotches; labials of both jaws mottled light and blackish.

Dimensions.

	<i>mm.</i>
Total length	518
Snout to vent	474
Vent to tip of tail	44

Variation.—With the exception of one specimen having 8 supralabials on one side, the head-shields show no noteworthy variation. In the specimens recorded the ventrals range between 183 and 204, the subcaudals between 27 and 30 only.

The coloration is also very constant. The only variation is a slight one in the number of the cross bands. Thus Science College Museum No. 3 has only 8 complete bands on body, while No. 25 in the Imperial Museum, Tokyo, has 10, both having one at the vent and one on tail.

Doctor Wall, since the above was written, has reported upon six specimens obtained in Okinawa by Owston's collectors, and states

that they were all alike in having five longitudinal black dorsal lines broader than the pink intervals. There were 9-12 narrow black annuli around the body and 1-2 (2 usually) on the tail. The ventrals and subcaudals are 204+28, 183+28, 194+30, 202+?, 197+27, 129 [192?] +29.

Habitat.—This richly and uniquely colored form seems to be confined to the middle group of the Riu Kiu archipelago, inasmuch as all the specimens of which we have detailed record are from Okinawa shima or "Great Loo Choo." Doctor Fritze collected the type on the west side of that island.

List of specimens of *Hemibungarus boettgeri*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supralabials.
				1891.								
Senckenberg Imper. Tokyo Sci. Coll. Tokyo.	9395a 25 3	Adult a.	Okinawa shima 'Riu Kiu'	Aug. 5	Doctor Fritze	13	184	2	30	1,2	1+2	7
Kumamoto School.	B 25.	Female ^b	do.	1891.		13	193	2	29	1,2	1+1	7
Brit. Mus.	9.9.3-14	Male	do.		P. Holst.	13	190	2	28			

^a Type.

^b Description p. 300; figs. 320-322.

Genus *CALLIOPHIS* ^a Gray.

1834. *Calliophis* GRAY, Indian Zool., II, (C. fig.) (type, *C. gracilis*).

1859. *Callphis* GUENTHER, Proc. Zool. Soc. London, 1859, p. 81 (emendation).

With regard to the genetic relationship of the snakes at present embraced in this genus it is sufficient to refer to the remarks under *Hemibungarus* (p. 387).

Calliophis univirgatus occupies the eastern Himalayas, being represented farther east, in Burma and South China, by *C. macclellandii*. Two species inhabit the mountains in southern India, one the eastern Indian peninsula, from Burma to Cochin China, while one occupies the Malay peninsula and Sumatra.

CALLIOPHIS MACCLELLANDII ^b (Reinhardt).

1844. *Elaps macclellandii* REINHARDT, Calcutta Journ. Nat. Hist., IV (p. 532) (type-locality, Assam; type in Mus. Copenhagen).—*Calliophis macclellandii* GUENTHER, Rept. Brit. India, p. 349 (part; var. γ , Assam).—BOULENGER, Fauna Brit. India, Rept., 1890, p. 385 (part; Assam; Burma;

^a From κάλλος, beauty; ὄφης, snake.

^b Named in honor of Dr. John McClelland, then editor of the Calcutta Journal of Natural History, who presented the type to the Royal Museum in Copenhagen. McClelland is the author of a large number of papers on Indian palaeontology and ichthyology.

Southern China); Cat. Sn. Brit. Mus., III, 1896, p. 399 (part: forma *typica*, Assam; Pegu; Formosa; South China); Proc. Zool. Soc. London, 1899, p. 166 (Kuatun, Fokien, China).


1854. *Elaps personatus* BLYTH, Journ. Assiat. Soc. Bengal, XXIII, no. 3, p. 298 (type-locality, Assam).

1861. *Elaps maclellandi* REINHARDT, Vid. Meddel. Naturh., For. Kjoebenhavn, 1860, p. 247 (part: Assam).

1864. *Callophis annularis* GUENTHER, Rept. Brit. India, p. 350, pl. xxiv, fig. 1 (type-locality, "India").^a

Calliophis univirgatus,^b which is said to differ only in coloration, having a narrow black line down the middle of the back and no regular dorsal crossbars, is the western representative of the present species. It inhabits Sikkim and Nepal, apparently to the exclusion of typical *C. maclellandii*.

Description (figs. 323-324).—*Adult male*; British Museum; Formosa; R. Swinhoe, collector. Rostral barely visible from above;



FIGS. 323-324.—*CALLIOPHIS MACLELLANDII*. (FROM BOULENGER, FAUNA BRIT. INDIA, REPT., p. 383.)
323, TOP OF HEAD; 324, SIDE OF HEAD.

internasals nearly entering nostrils; nostril in posterior edge of a large nasal, followed by a somewhat smaller postnasal; prefrontals large, broadly in contact with supraoculars; frontal about as long as its distance from tip of snout, slightly shorter than parietals; one large preocular in contact with postnasal, prefrontal, supraocular, and third supralabial; two postoculars; temporals 1 + 1; 7 supra-labials, third and fourth entering eye, first small, third large, as large

^a In Boulenger's catalogue (III, p. 399) it is stated that the type in British Museum is from South China and presented by Mr. J. C. Bowring.

^b 1858. *Elaps univirgatus* GUENTHER, Cat. Col. Sn. Brit. Mus., p. 231 (type-locality, Nepal; types in British Museum; Hodgson, collector).—*Calliophis univirgata* GUENTHER, Proc. Zool. Soc. London, 1859 (p. 83, pl. xvii Nepal).

1861. *Elaps maclellandi* REINHARDT, Vid. Meddel. Naturh., For. Kjoebenhavn, 1860, p. 247 (part: Darjeeling).

1861. *Callophis maclellandii* GUENTHER, Proc. Zool. Soc. London, 1861, p. 219 (Nepal).

1896. *Callophis maclellandii* var. *univirgatus* BOULENGER, Cat. Sn. Brit. Mus., III, p. 399 (Nepal; Darjeeling).

as sixth; 6 lower labials, four in contact with anterior chin-shields, fourth very large; posterior chin-shields about as large as anterior; 13 rows of smooth scales; 240 ventrals^a; anal divided; 34 pairs of subcaudals.^a Color (in alcohol) above dull reddish gray, with 31 yellow-edged black rings on the body and 4 on the tail, the bands being one scale wide on the back and two ventrals wide on the under-side; halfway between the rings on the back two black spots about 5 scales apart to within 4 rings from the vent, and on the under-side a single rounded large black blotch on the median line; a black collar about three scales wide just back of temporals and including the extreme posterior angle of the parietals; another black band from edge of lip through eye and involving anterior third of frontal and the prefrontals; snout in front of this band, as well as the rest of the head between the two black bands yellow.

Dimensions.

	mm.
Total length.....	400
Snout to vent.....	363
Vent to tip of tail.....	37

Variation.—The only feature worthy of notice is the great range in the number of ventrals, as shown by the Indian and Chinese specimens in British Museum, the extremes being 182 and 240. It should be noted, however, that the latter figure is that of the only specimen thus far recorded from Formosa, and that there is a wide gap between it and the other specimens in this respect. It is not beyond the bounds of probability that the Formosan specimen may represent a separate form characterized by a greater number of ventrals.

Habitat.—Extending from Assam and Burma in the west to southern China and Formosa in the east. It is recorded from the mountains north of Kiukiang, and also from high altitudes (3,000 to 4,000 feet) in the province of Fokien.

A single specimen collected by Swinhoe and now in the British Museum is known from Formosa.

List of specimens of Calliophis macclendani.

Brit. Mus. No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows,		Anal.	Subcaudals.
					Ventrals.			
d	Male ^a	Formosa.....	R. Swinhoe.....	13	240	2	34
c	Half grown ^b	Kiukiang Mountains, China.....	A. E. Pratt.....	13	215	2	26
e	Male ^b	Southern China.....	J. C. Bowring.....	13	212	2	32
.....	do. c.....	Kuatun, Fokien.....	J. D. La Touche.....	13	193	2	36

^a Description, p. 392. ^b Boulenger, Cat. 111, p. 399. ^c Boulenger, Proc. Zool. Soc., 1899, p. 166.

^a According to Boulenger's count.

Genus NAJA^a Laurenti.

1768. *Naja* LAURENTI, Synops. Rept., p. 90 (type, *Naja naja*).
 1830. *Uraus* WAGLER, Nat. Syst. Amph., p. 173 (type, *Naja haji*).
 1830. *Aspis* WAGLER, Nat. Syst. Amph., p. 173 (type, *Naja naja*) (not of Laurenti, 1768).
 1831. *Tomurus* EICHWALD, Zool. Spec., III (p. 171) (type, *T. oriana*).
 1836. *Hamadryas* CANTOR, Asiat. Research., XIX, p. 187 (type, *H. hannah*) (not of Huobner, 1806).
 1843. *Dendraspis* FITZINGER, Syst. Rept., p. 28 (type, *N. bungarus*).
 1858. *Pseudohaje* GÜENTHER, Cat. Colubr. Snakes Brit. Mus., p. 222 (type, *P. nigra*).
 1864. *Ophiophagus* GÜENTHER, Rept. Brit. India, p. 340 (type, *O. elaps*).

One species of these deadly snakes occurs within our territory, inasmuch as the eastern representative of the terrible Indian cobra has been found in the extreme south of Formosa.

NAJA NAJA^b ATRA^c (Cantor).

1842. *Naja atra* CANTOR, Ann. Nat. Hist., IX, 1842, p. 482 (type-locality, Chusan, China).
 1859. ?*Naja tripudians* var. *scopinucha* COPE, Proc. Phila. Acad., 1859, p. 343 (type-locality, Canton River; type in Mus. Acad. Phila.; Doctor Ruschenberger, collector).
 1860. *Naja haji* HALLOWELL, Proc. Phila. Acad., 1860, p. 504 (Hongkong) (not *N. haji* (Linnaeus)).
 1866. *Naja tripudians* var. *atra* MARTENS, Preuss. Exped. Ost-Asien, Zool., I, p. 177 (Chusan Archipel).
 1867. *Naja lutescens* var. *larcata* STEINDACHNER, Novara Exped., Zool., I, Rept. p. 84 (Hongkong) (not of Cantor).
 1876. *Naja tripudians* var. *unicolor* MARTENS, Preuss. Exped. Ost-Asien, Zool., I, p. 382 (substitute name).
 1885. *Naja tripudians* BOETTGER, Offenbach, Ver. Naturk., 24-25 Ber., pp. 127, 154 (Canton) (not of Merrem); Ber. Senckenberg. Naturf. Ges., 1894, p. 134 (Hainan); p. 137 (Hongkong).—WALL, Proc. Zool. Soc. London, 1903, p. 97 (Hongkong; Hainan).
 1896. *Naja tripudians* var. *fasciata* (part) BOULENGER, Cat. Snakes Brit. Mus., III, p. 383 (Kiukiang; Canton; Hainan; Siam).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 121 (Canton; Hainan) (not of Gray?).
 1896. *Naja tripudians* var. *sputatrix* (part) BOULENGER, Cat. Snakes Brit. Mus., III, p. 384 (Chusan Ids.) (not of Boie).

In treating of the association of subspecies constituting the species *Naja tripudians* (= *Naja naja*) Boulenger^d has attempted a subdivision based chiefly upon color differences. A glance at the list of 71 specimens given by him, shows that the "subspecies," or "varieties," or "forms" thus distinguished are not correlated with geographical distribution. In view of Doctor Boulenger's statement that "the

^a Said to be derived from *Nag*, the Indian word for snake.

^b 1758. *Coluber naja* LINNÆUS, Syst. Nat., 10 ed., I, p. 221 (type-locality, India orientalis); 12 ed., I, 1766, p. 382.

^c Signifying black.

^d Cat. Snakes Brit. Mus., III, 1896, p. 381.

forms enumerated [by him] might be regarded as distinct species but for the absence of any sharp demarcation-lines between them," this lack of correlation might be explained. Nevertheless, even if we regard them as species, their geographical distribution becomes disconnected, confused, and unintelligible.

It seems that a better subdivision might have been obtained by taking the scale formula as a primary basis and color only as a secondary character.

It might then be possible to separate out a form *Naja naja oxiana* (Eichwald), from Transcaspia, Afghanistan, and Gilgit, characterized by a large number (averaging 268) of ventrals and caudals together, the sum of these giving more definite results than each separately, owing to the variable location of the vent.

The specimens from India proper, Ceylon, and possibly Burma and the Malay peninsula seem to be fairly well characterized by the large number of scales around the neck, 29 to 35, rarely 27, and should probably stand as *Naja naja*.

The cobras from the Malay Archipelago have about the same average number of ventrals plus caudals as the Indian cobras (235 in the former as against 238 in the latter), but the number of scales around the neck is smaller, viz, only 21 to 25. These may be Boie's *Naja naja sputatrix*, or they may fall into several more subspecies.

Finally, the Chinese and Siamese specimens seem to hold an intermediary position so far as neck scales are concerned, viz, 25 to 27, but with a minimum of ventrals plus caudals (average of 9 specimens 216.) These are also fairly uniform in color and appear entitled to the subspecific appellation *Naja naja atra* (Cantor).

The demarcation-lines here drawn up are not very sharp, but they seem better than those based primarily on color and have the advantage of a more rational geographic distribution than the others. Moreover, like Boulenger's scheme, this is only a provisional arrangement which needs to be tested with a very large material. In this connection I would earnestly warn against using specimens, to the origin of which there attaches the least suspicion. Only by utilizing material which in every detail is authentic and beyond dispute will it be possible to reach a satisfactory solution.

Description.—*Young*: British Museum, No. 99.4.24.102; South Cape, Formosa; J. D. La Touche, collector. Rostral very broad, the height only two-thirds of the width; internasals somewhat longer than prefrontals; frontal pentagonal, broader than supraoculars; parietals as long as frontal and prefrontals together; nostril separating the two large nasals; no loreal; one prefrontal in contact with supraocular, prefrontal, internasal, posterior nasal, and third supralabial; diameter of eye as large as its distance from anterior rim of

nostril; two postoculars, upper much the larger; large temporals. $2 + 1$; a large scale in the posterior angle between the parietals, and a larger one on each side of the parietals in continuation of the larger upper temporal; seven supralabials, third and fourth entering eye, third very high, broadly in contact with posterior nasal, fifth high, touching upper postocular; four lower labials in contact with anterior chin-shields which are larger than posterior pair; 25 scales round the neck, and 21 round the body;^a 178 ventrals;^b anal divided; 42 pairs of subcaudals.^a Color (in alcohol) uniform blackish brown, except sides and lower part of head, as well as 16 anterior ventrals, which are whitish; on the upper neck a pair of white-rimmed "spectacles" and 13 very narrow yellowish cross lines, the anterior of which is faintly indicated on the ventrals also.

Dimensions.

	<i>mm.</i>
Total length.....	352
Snout to vent.....	304
Vent to tip of tail.....	48

As to how large the cobra grows to be in Formosa there is no information, but specimens from other localities are known to exceed 1½ meters in length.

Variation.—The species is subject to a considerable amount of variation as might be expected from its wide distribution. The more essential features of this have been touched upon in the preliminary remarks following the synonymy (see p. 394).

Distribution.—The species *Naja naja* has a wide distribution in southeastern Asia, from Transcaaspa to the Philippines, but the present subspecies, as defined above, seems to be confined to China south of Shanghai, including Hainan, French Indo-China, continental Siam, and Formosa.

Its occurrence in Formosa is now demonstrated by a young specimen in British Museum collected near South Cape by Mr. J. D. La Touche. I owe the opportunity to examine and describe this specimen to the courtesy of Dr. G. A. Boulenger.

^a According to Boulenger's count.

Last of specimens of Naja naja atra.

Museum.	Number.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows, neck.	Scale rows, body.	Ventrals.	Subcaudals.	Ventrals plus subcaudals.
1854.										
U.S.N.M.	7517	Female	Hongkong, China.	May	W. Stimpson	27	21	170	45	215
Brit. Mus.	99.4.24.102	Young	South Cape, Formosa.		J. D. La Touche	25	21	178	42	220
Do.	Da	Male <i>b</i>	Chusan Islands.		J. J. Walker	25	21	167		
Do.	Db	Young <i>t</i>	China.		R. Lindsay	25	21	177	47	224
Do.	Cba	Female <i>c</i>	Kiukiang, China.		J. Walley	25	21	166	48	214
Do.	Cbb	do. <i>c</i>	Canton.			25	21	170		
Do.	Cbc	do. <i>c</i>	Haiman Island.		J. Neumann	25	21	168	43	211
Senckenberg.	9383a	Adult <i>d</i>	Canton.		O. Fr. v. Moellendorff.	(e)	21	162	47	209
Do.	9383b	do. <i>d</i>	do.			(e)	21	164	50	214
(?)		(<i>d</i>)	do.		do.	(e)	21	165		
(?)		(<i>d</i>)	do.		do.	(e)	21	169		
Brit. Mus.	Cbd	Male <i>c</i>	Siam.			27	21	170	48	218
Do.	Cbc	Female <i>c</i>	do.			27	21	174	47	221

a Description, p. 395.

d Boettger, 1885, p. 127.

b Boulenger, Cat. III, p. 384.

e 25 to 27 scale rows round the neck, Boettger, *l. c.*

c Boulenger, Cat. III, p. 383.

Genus BUNGARUS *a* Daudin.

- 1803. *Bungarus* DAUDIN, Bull. Soc. Philom. Paris, III, No. 72, March, 1803, p. 187 (type, *B. fasciatus*).
- 1811. *Pseudo-boa* OPPEL, Ordn. Rept., p. 68 (type, *B. fasciatus*) (not of Schneider, 1801).
- 1828. *Aspidoclonion* WAGLER, Icon. Amph., I, fol. 2 (type, *A. semifasciatum*).

BUNGARUS MULTICINCTUS *b* Blyth.

- 1858. *Bungarus semifasciatus* GUENTHER, Cat. Col. Snakes Brit. Mus., p. 221 (China) (not of Boie, 1827); Rept. Brit. India, 1864, p. 344 (China; Formosa).—SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 225 (Formosa; Amoy, China).—BOETTGER, Offenbach. Ver. Naturk. 24–25 Ber., 1885, p. 128 (Canton).
- 1861. *Bungarus multicinctus* BLYTH, Journ. Asiat. Soc. Bengal, XXIX, p. 98 (type-locality, Formosa; type in Calcutta Mus.; Swinhoe, collector).—BOETTGER, Offenbach. Ver. Naturk. 26–28 Ber., 1888, pp. 86, 148 (South China; Formosa).—STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 222 (Taipa, Formosa).
- 1896. *Bungarus candidus* var. *multicinctus* BOULENGER, Cat. Sn. Brit. Mus., III, p. 369 (China; Formosa; Burma); Proc. Zool. Soc. London, 1899, p. 164 (Kuatun, Fokien, China).—*Bungarus candidus* var. *multicincta* BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 119 (Canton).

Three closely-related forms of the “Krait” differing in color but apparently occupying distinct geographic areas are united by Bou-


a Derived from “Bungarum-pamah,” which according to Russell’s “Indian Serpents” is the vernacular name for *B. fasciatus* on the coast of Coromandel, India.

b Signifying many-belted, i. e., with many crossbars.

lenger under the oldest name, *B. candidus*. The typical form seems to be confined to islands in the Malayan Archipelago; another, var. *cæruleus*, extends over entire India (exclusive of the Himalayas and Ceylon) and the Malay Peninsula; the third, the one here treated of, is recorded from Toungoo, in lower Burma, not far from the Karennee Hills, and from various localities in southern China. The first is considered most distinct, and there seems to be no good reason for connecting it nomenclaturally with our Chinese form. Of the Indian tropical form Boulenger says that it is "almost completely connected" with *B. multicinctus*. This "almost" only prevents me at present from speaking of the latter as *Bungarus cæruleus multicinctus*.

Another very closely allied form, differing chiefly in the less-pronounced enlargement of the median dorsal scale row, *B. lividus*, inhabits Assam. The Ceylon krait *B. ceylonicus*, is another nearly related form.

Description.—*Adult male*: Science College Museum, Tokyo, No. 15; Taipa, Formosa; November, 1897; T. Tada, collector (figs. 325-327). Rostral broader than high; visible from above; internasals,



FIGS. 325-327.—*BUNGARUS MULTICINCTUS*. NAT. SIZE. 325, UNDERSIDE OF HEAD; 326, SIDE OF HEAD; 327, TOP OF HEAD. No. 15, SCI. COLL. TOKYO.

broader than long, half as large as prefrontals which are broadly in contact with supraoculars; frontal as long as its distance from tip of snout and interparietal suture twice as wide as supraoculars at center of eye; parietals much longer than frontal, nearly as long as their distance from tip of snout; nostril large, vertically elliptic, between two nasals, the posterior narrowed behind so as to meet the preocular in a very short suture; no loreal; one long preocular, anteriorly in narrow contact with posterior nasal; eye rather small, its vertical diameter shorter than its distance from edge of lip; two postoculars; temporals 1 + 2; seven supralabials, last three largest, second in contact with preocular, third and fourth entering eye; four lower labials in contact with anterior chin-shields which are longer than the posterior pair; 15 rows of smooth scales, without apical pits, the median row greatly enlarged, more than twice as large as the other dorsals; 210 ventrals;

anal entire; 46 undivided subcaudals. Color (in alcohol) above dark seal brown with 45 narrow whitish cross lines or bars, 31 on the body and 14 on the tail, about one scale wide on the back, widening near the belly so as to equal two of the outer scale rows, the light-colored lateral scales often with a median dark line; head uniform dark, somewhat paler on labials; underside whitish more or less clouded with brown, and the dark brown of the sides descending on the body over the ends of the corresponding ventrals, while forming continuous, though somewhat irregular, rings on the tail.

Dimensions.

	<i>mm.</i>
Total length.....	860
Snout to vent.....	743
Vent to tip of tail.....	117

The *young* (U.S.N.M. No. 36514) is essentially like the adult described above, but has a whitish crescentic mark on each side of the neck behind the head.

This species reaches a length of more than a meter.

Variation.—The scale formula varies but little though 1+1 temporals occur besides 1+2. The ventrals vary in this form between 198 and 221, the subcaudals between 44 and 52.

The variation in coloration has reference chiefly to the number of white cross lines. Thus, the adult described above has 45, and the young 51 (40 on body 11 on tail), while as many as 60 have been recorded.

Habitat.—Southern China, from the provinces of Kiangsi and Fokien to Kwangtung and Kwangsi. Farther west it has been collected at Toungoo in Lower Burma, by Mr. E. W. Oates.

In Formosa it was first found by Swinhoe, and Mr. Tashiro has since informed us that it is found both in the northern and southern part of the island. He states that it was common around Taipa, whence he brought two specimens.

List of specimens of Bungarus multicinctus.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows		Anal.	Subcaudals.	Oculars.	Temporals.	Supralabials.
						Ventrals.						
U.S.N.M...	36514	Young ^a	Taihoku (Taipa), Formosa.	1897, Sept. —	T. Tada...	15	210	1	48	1, 2, 1+2		7
Sci. Coll., Tokyo.	15	Male ^b	Taipa, Formosa.	do	do	15	210	1	46	1, 2, 1+2		7
Brit. Mus..	Bc..	Female ^c	Formosa.....		R. Swinhoe	15	207	1	46			

^a Sci. Coll. No. 3.

^b Figs. 325-327; description, p. 398.

^c Boulenger, Cat. III, p. 369.

Subfamily HYDRINÆ.

The high, compressed tail, shaped like the blade of an oar, distinguishes these snakes at once from all others and stamps them as inhabitants of the water. As a matter of fact these snakes are not only aquatic, but truly marine, some of them to an extent that renders it correct to designate them as pelagic.

Not only has the shape of the tail become specialized in these snakes, but the whole body is strongly compressed in some; in others the general shape has been greatly modified, so that the whole posterior portion of the body is abnormally enlarged while the front part is drawn out into an excessively elongated and narrow neck finished off with a small head not wider than the neck. Owing to the compressed form of the body the ventrals have been reduced in most of the forms to an extent which makes it difficult to recognize them as such. Even the scales of the body have become highly altered in many forms, being polygonal and juxtaposed with short keels or knobs or spines, sometimes in pairs, and totally unlike the scales of other snakes. That the nostrils are placed on top of the snout and provided with valves it is scarcely necessary to mention.

Not all the species are as greatly differentiated as above mentioned. A small group, the habits of which are still partly terrestrial, live near the shores and occasionally and temporarily climb among the rocks or even sometimes venture farther inland. These have still fairly well-developed ventrals to some extent fit for terrestrial locomotion, and their scales are imbricate. To this group belong the genera *Laticauda* and *Emydocephalus*.

Being "proteroglyph," all these marine snakes are poisonous. Nevertheless, several forms are perfectly harmless to man, such as the *Laticauda* occurring so frequently around the Riu Kiu Islands which the fishermen handle without fear and without accident.

The sea snakes, broadly speaking, inhabit all the tropical and subtropical seas outside of the Atlantic Ocean. In spite of much good work of late in this group, the species and their distribution are as yet but poorly understood. The lack of definite boundaries to their habitat and their liability to be carried beyond their natural limits by currents make it difficult to decide how many species to include in the present work. It has therefore been thought best to take cognizance of a few more species than have actually been recorded from the coasts of the territory covered by this work, when they are known to occur in the adjacent seas. Such species, however, have been included in brackets.

Doctor Boettger, in 1888 (*Zool. Anz.*, pp. 395 seq.), published a paper on the external sexual characters of the sea snakes based upon an examination of 46 specimens belonging to two genera (*Disteira* and *Lapemis*), all from the Philippine Islands. He came to the con-

clusion that the males differ in having fewer ventrals, fewer scale rows around the neck, more anals, longer tail and smaller circumference of body relative to total length, and much more developed spines on the ventrals. In all three species examined by him he found a difference of about 20 ventrals between the maximum of the males and the minimum of the females. Thus in *Lapemis hardwickii* (*Hydrophis loreatus*) he found in the males 135-168 ventrals (average 153) and in the females 186-237 (average 202), the difference between the averages being 49 ventrals. His result was calculated to carry conviction as it was based upon a series of no less than 31 specimens. Nevertheless, if we inquire into the data furnished by Boulenger regarding the specimens of the same species in British Museum ^a we find in a much smaller series (14 specimens, 5 males and 9 females) proportions essentially different. In his list the ventrals of the males range between 130 and 155 (average 140), those of the females 136-200 (average 156), consequently an overlap of 19 ventrals instead of a gap of 18. The British Museum specimens are not from one locality, however, but cover the whole area from India to the Philippines, and it might be supposed that this discrepancy had a geographical significance. Such is not the case, however, for three females from Negros, Philippine Archipelago, consequently from practically the same locality as Boettger's specimens, have 140, 156, and 180 ventrals, respectively, thus nearly covering the whole range of Boettger's 18 males (135-168) without reaching the minimum of his 13 females (186).

These apparent discrepancies seem at present inexplicable, and indicate one of the many difficulties with which the student of these snakes has to deal.

My own observations bear out Boettger's conclusions only partly. In our four specimens of *Disteira godeffroyi* the female has more scale rows, more ventrals, and fewer subcaudals. On the other hand, in 20 specimens of *Disteira melanocephala* there is apparently no difference among the sexes except that the males average a few more subcaudals than the females.

KEY TO THE GENERA OF HYDRINE IN JAPANESE AND CHINESE WATERS.

- a*¹ Ventrals wide, at least four times as wide as long.
*b*¹ Nasals widely separated by internasals; supralabials normal. *Laticauda*, p. 402.
*b*² Nasals broadly in contact on median line of snout; middle supralabials fused into a large plate on each side *Emydophalus*, p. 413.
*a*² Ventrals rudimentary, less than twice as wide as long.
*b*¹ Two pairs of chin-shields, anterior always, posterior nearly always in contact. *Disteira*, p. 418.
*b*² Chin-shields scarcely differentiated, not in contact.

^a Cat. Sn. Brit. Mus., III, p. 301.

- c*¹ Snout, from eye to tip shorter than width of head at posterior corner of supraoculars.
*d*¹ Mental not concealed in a deep groove [*Lapemis*], p. 435.
*d*² Mental concealed in a deep groove of the chin [*Enhydrina*], p. 437.
*c*² Snout, from eye to tip equaling or longer than width of head at posterior corner of supraoculars *Hydrus*, p. 438.

Genus *LATICAUDA*^a Laurenti.

1768. *Laticauda* LAURENTI, Syn. Rept., p. 109 (type, *L. scutata*).
 1802. *Platurus* LATREILLE, Hist. Nat. Rept. IV, p. 183 (type, *P. fasciatus*).

Five species of this genus are provisionally recognized, of which *Laticauda muelleri*^b seems to be a *L. laticaudata* with a ventral keel. A *Laticauda frontalis*^c has recently been described as differing from *L. semifasciata* in having an undivided rostral and shield-like temporals.

Only two species are known with certainty to have been taken on the coasts of the territory covered by this work, but a third may be reasonably expected to occur and its characters are therefore given in conjunction with the others. They may be distinguished by the following:

KEY TO THE SPECIES OF *LATICAUDA* IN JAPANESE AND CHINESE SEAS.

- a*¹ Rostral undivided horizontally; belly without a median keel; two pairs of contiguous chin-shields; temporals 1+2; top of snout yellow.
*b*¹ Scale rows 19; no unpaired shield between the prefrontals; upper lip dark brown. *L. laticaudata*, p. 402.
*b*² Scale rows 21 to 25; normally an unpaired shield between the prefrontals; upper lip yellow [*L. colubrina*, p. 406.]
*a*² Rostral divided horizontally; belly with a median keel on posterior half; one pair of contiguous chin-shields; temporals 2+3; top of snout dark brown. *L. semifasciata*, p. 409.

LATICAUDA LATICAUDATA^d (Linnaeus).

1758. *Coluber laticaudatus* LINNAEUS, Syst. Nat., 10 ed., I, p. 222 (part: type-locality, the Indies; type in Mus. Stockholm); 12 ed., I, 1766, p. 383.—ANDERSSON, Bihang Svensk. Vet. Akad. Handl., XXIV, 1899, Pt. 4, no. 6, p. 18 (type).—*Platurus laticaudatus* GIRARD, Herpet. U. S. Expl. Exped., 1858, p. 180 (Fiji Is.).—PETERS, Mon. Ber. Berlin Akad. Wiss., 1877, p. 417 (part: Fiji, Tonga, Philippine Is.).—BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 307 (Bay of Bengal to western Pacific; Loo Choo Islands).—WALL, Proc. Zool. Soc. London, 1903, pp. 96, 101 (Formosa; Loo Choo Islands); 1905, II, p. 517 (Okinawa).
 1768. *Laticauda scutata* LAURENTI, Syn. Rept., p. 109 (substitute name for *C. laticaudatus* LINNAEUS).

^a From *latus*, broad; *cauda*, tail.

^b *Platurus muelleri* BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 309 (South Pacific).

^c *Platurus frontalis* DE VIS, Ann. Queensland Mus., No. 6, 1905, p. 48 (New Guinea).

^d From *latus*, broad; *caudatus*, tailed.

1802. *Platurus fasciatus* LATREILLE, Hist. Nat. Rept., IV, p. 185 (Indies).—FISCHER, Abh. Naturw. Ver. Hamburg, III, 1856, pp. 28, 70 (part: specim. Hamburg Mus., Indian Ocean).—HALLOWELL, Proc. Phila. Acad., 1860, p. 493 (Naha, Okinawa shima).—BOULENGER, Proc. Zool. Soc. London, 1887, p. 149 (Lao Choo Islands).
1817. *Platurus laurenti* RAFINESQUE, Amer. Month. Mag., I, p. 432 (substitute name; not of Daudin, 1803).
1836. *Coluber platycaudatus* OKEN, Allgem. Naturg., VIII, p. 566 (*lapsus*).
1858. *Platurus laticaudatus* var. GUENTHER, Cat. Colubr. Sn. Brit. Mus., p. 272 (New Guinea, Bengal, Siam, Van Diemen's Land).
1859. *Platurus fischeri* JAN, Rev. Mag. Zool., 1859 (p. 149); author's separate, p. 25, pl. D (type-locality, Indian Ocean; type in Mus. Milano); Icon. Ophid., livr. 40, 1872, pl. 1, fig. 2 (type).—GUENTHER, Rept. Brit. India, 1864, p. 356, pl. xxv, fig. A (Bengal; Siam; New Guinea; New Hebrides).—ANDERSON, Proc. Zool. Soc. London, 1871, p. 189 (Calcutta).
1871. *Platurus affinis* ANDERSON, Proc. Zool. Soc. London, 1871, p. 190 (type-locality, Fallahs Mullah, Calcutta; type in Ind. Mus., Calcutta).

The *Coluber laticaudatus* of Linnaeus (Syst. Nat., 10th and 12th editions) was based upon his account of the same species in Museum Adolphi Frederici, which was published in 1754 and not binominal. Two specimens were apparently described in that work. It now turns out from Andersson's examination of the types^a that Linnaeus had before him a specimen of each of the two species lately known as *Platurus laticaudatus* and *P. colubrinus*, the former with 19 scale rows, the latter with 23. His *C. laticaudatus* is consequently a composite species, and the application of the name by the first reviser who separated the component forms must stand.

The two species were steadily confounded by all authors, just as Linnaeus had done, though under various names, such as *Platurus fasciatus* and *Hydrophis colubrinus*, until 1858, when Girard, for the first time (and not Peters in 1877, as occasionally stated), clearly separated, described, and named them. He tied the name *laticaudatus*, which had scarcely been used at all in the herpetological literature since the time of Linnaeus, to the species with 19 scale rows, and applied *Hydrus colubrinus* of Schneider correctly to the other, this being the oldest name used for a specimen of the latter, though Schneider himself did not realize its distinctness from the Linnaean species, of which, as we have seen, in reality it only formed a part. Girard's nomenclature has been followed by nearly all subsequent writers.

The fact that Linnaeus in the Systema Naturæ only gives the scale formula (i. e., number of ventrals and subcaudals) of the specimen with 23 scale rows (220-42) is not sufficient reason for fastening the name *C. laticaudatus* to the species represented by this specimen, in view of the above argument, and is, moreover, sufficiently set off by

^a Bihang Svensk. Vet. Akad. Handl., XXIV, 1899, Pt. 4, No. 6, p. 18.

his reference to the plate in Museum Adolphi Frederici, which is said to represent the other specimen.

Whether the above synonymy includes one or more distinguishable forms may perhaps yet be open to question; at least I have not been able to fully make up my mind from our material which is entirely too insufficient, however. As far as it goes it seems to indicate two color forms. At least our specimens can be divided according to the following concurrent color characters, thus:


- a*¹ Light horseshoe-shaped mark on top of head bent down behind the eye, reaching the lip; none, or one, light ring on neck incomplete below; light rings on belly 4-5 ventrals wide *L. laticaudata* (Nos. 8994, Java; 29448, Nicobars).
- a*² Light horseshoe-shaped mark not bent down behind the eye; usually two light rings on neck incomplete below; light rings on belly 1-3 ventrals wide.
L. laticaudata affinis (Nos. 30761, Pulo Mansular, W. Sumatra; 7381, Riu Kius.; 21399, China Seas; 32669, Formosa; 33937-41, Riu Kius).

It should be noted that our eastern specimens belong to *L. affinis*, and it would be interesting to know to what extent the characters mentioned above hold good in a large series. In this connection it may be stated that the character of the incomplete light rings on the neck below is the least constant, as might be expected. It holds in seven of our nine specimens of *L. affinis*, but two of the Riu Kiu specimens (Nos. 33937 and 33939) have only one incomplete ring.

Since the above was written I have examined two specimens from Formosa in the Christiania Museum (No. 832) which in the main bear out the above.

Description (figs. 328-330).—*Adult male*; U.S.N.M. No. 32669; Formosa; 1894; Mr. Navara, collector. Rostral undivided, higher than broad, just visible from above; internasals broader than long, nearly triangular, their suture with rostral very short; prefrontals in contact throughout, anteriorly very broad with an acute outer angle descending on side of face between nasal and preocular without quite reaching the second supralabial, broadly in contact behind with supraocular; frontal long, pointed behind, much longer than distance from tip of snout; as long as parietals, broader than supraoculars; parietals short, not longer than frontal, the suture between them shorter than distance between frontal and rostral; nostril in the posterior half of an undivided nasal which is three times as long as high, slightly lower in front and narrowly in contact with preocular beneath the prefrontal; one preocular; eye small, its diameter less than its distance from edge of lip; 2 postoculars; temporals 1+2, slightly larger than the adjoining labials; 7 supralabials, first much wider above than below, third and fourth entering eye; mental very small, smaller than first lower labials; 7 lower labials, the last four exceedingly narrow, three anterior only in contact with anterior chin-shields which are somewhat smaller than the posterior, the latter broadly in contact; 19 rows of smooth scales without apical pits; 243 ventrals; anal

divided; 47 pairs of subcaudals. Color (in alcohol) bluish gray above, yellowish underneath, with broad blackish brown rings around the body, there being 47 on neck and body and 5 on tail, the first on neck incomplete below; width of black rings above about four scale rows and about three ventrals underneath, the light interspaces



FIGS. 328-330.—*LATICAUDA [LATICAUDATA]*. $2 \times$ NAT. SIZE. 328, TOP OF HEAD; 329, SIDE OF HEAD; 330, UNDERSIDE OF HEAD. NO. 7381, U.S.N.M.

respectively $2\frac{1}{2}$ above and 2 below; head blackish brown, rostral, snout, and eyebrow region as far as anterior angle of parietal and first temporal, yellow; labials brownish black as well as throat and lower neck, the latter with a median line of yellow.

Dimensions.

	mm.
Total length.....	769
Snout to vent.....	659
Vent to tip of tail.....	110

The *female* and *young* are essentially like the above, the former somewhat larger with a shorter tail, No. 33937, for instance, our largest specimen, measuring 890 mm. with a tail 95 mm. long.

Variation.—The scale formula is fairly constant in this species, oculars, temporals, supralabials, and anal showing no deviation from the normal in a large series. In the specimens recorded from the waters adjacent to our territory the ventrals vary between 230 and 243, the subcaudals between 30 and 47.

As for the color variation of this species it is sufficient to refer to the preliminary remarks (on p. 404). The number of black rings on the body in ten specimens from Japanese waters ranges between 39 and 55.

Habitat.—Widely distributed along the coasts from India to the South Pacific islands and Tasmania. The species seems common about the Fijis and the Friendly Islands, and Fischer mentions a specimen from the Society Islands.

On the coasts of southeastern Asia it is not rare—for instance, in the Philippines—and to the north it has been found as far as Okinawa shima. A specimen from Naha, collected by Heine, of the Perry

Expedition, was recorded by Hallowell (U.S.N.M. No. 7381), and two specimens obtained by Pryer's collector, probably in the same island, are in British Museum. Two other specimens, also from Okinawa shima, are in the Imperial Museum, Ueno Park, Tokyo (Nos. 11, 12), as well as one from the Yaeyama group (No. 13). In the latter it must be common, for we have five specimens from Ishigaki shima through Mr. Owston, and British Museum has one from the same locality and collector through Doctor Wall. Similarly on the coasts of Formosa, whence I have examined two in the University Museum of Christiania, collected by Mr. Navara, and one from the same source in our own museum, obtained through the kindness of Prof. Robert Collett.

List of specimens of Laticauda laticaudata.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom obtained.	Scale rows.	Ventrals.	Subcaudals.	Black rings on body.	Oculars.	Temporals.	Supratibials.
U.S.N.M.	7381	Young ^a	Naha, Okinawa shima.	1853. June	W. Heine	19	232	45	55	1, 2, 1+2		7
Do.	21399	do.	China Sea (?)			19	242	43	48	1, 2, 1+2		7
Do.	32699	Male ^b	Formosa	1894	Navara	19	243	47	46	1, 2, 1+2		7
Do.	33937	Female	Ishigaki shima	1899. May-June	A. Owston	19	237	34	48	1, 2, 1+2		7
Do.	33938	do.	do.	do.	do.	19	237	33	51	1, 2, 1+2		7
Do.	33939	do.	do.	do.	do.	19	234	33	46	1, 2, 1+2		7
Do.	33940	do.	do.	do.	do.	19	238	32	50	1, 2, 1+2		7
Do.	33941	Male	do.	do.	do.	19	238	38	45	1, 2, 1+2		7
Christiania.	832 ^a	Young	Formosa	1894	Navara	19	235	44	39			
Do.	832 ^b	Male	do.	do.	do.	19	237		43			
Brit. Mus.	n		Ishigaki shima.		Doctor Wall	19	230	41				
Do.	c	Female ^c	Loo Choo Islands.		H.J.S. Pryer	19	233	30				
Do.	d	Young ^c	do.		do.	19	232	31				

^a Figs. 328-330.

^b Description, p. 404.

^c Boulenger, Cat. III, p. 308.

[*LATICAUDA COLUBRINA* ^a (Linnæus).]

1758. *Coluber laticaudatus* LINNÆUS, Syst. Nat., 10 ed., I, p. 222 (part); 12 ed., I, 1766, p. 383.—*Platurus laticaudatus* PETERS, Mon. Ber. Berlin Akad. Wiss., 1877, p. 417 (part: Philippines; Amboina; Fiji Is.; New Hanover).
1799. *Hydrus colubrinus* SCHNEIDER, Hist. Amph., I, p. 238 (type-locality not stated; type, no. 9078, Berlin Mus.).—*Hydrophis colubrinus* SCHLEGEL, Phys. Serp., II, 1837, p. 514, pl. xviii, figs. 21-22 (Moluccas, Timor, Celebes; China Sea).—*Platurus colubrinus* GIRARD, Herpet. U. S. Expl. Exped., 1858, p. 183 (Fiji and Tonga Is.).—BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 308.—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 114 (Philippines: Luzon, Cebu; Ternate; Tonga Tabu; Australia).—WALL, Proc. Zool. Soc. London, 1903, p. 101.

^a Signifying colubrine, like a *Coluber*; so named by Schneider because Linnæus had referred the species to that genus.

1803. *Platurus fasciatus* DAUDIN, Hist. Nat. Rept., VII, p. 226 (part), pl. LXXXV, fig. 1.—JAN, Icon. Ophid., livr. 40, 1872, pl. I, fig. 1 (New South Wales).
1847. *Laticauda scutata* CANTOR, Journ. Asiat. Soc. Bengal, XVI (p. 1046) (not of LAURENTI).—*Platurus scutatus* GUENTHER, Rept. Brit. India, 1864, p. 356 (Bay of Bengal to Chinese seas and New Zealand).
1856. *Platurus fasciatus* var. *colubrina* FISCHER, Abh. Naturw. Ver. Hamburg, III, p. 30.
1858. *Platurus laticaudatus* var. B GUENTHER, Cat. Col. Sn. Brit. Mus., p. 272 (Fiji Is.; Amboina; Sulu I.; Isle of Pines; India).
1885. *Platurus laticaudatus* var. *colubrina* BOETTGER, Offenbach. Ver. Naturk. 24-25 Ber., p. 155 (China Sea).

Description.—*Halfgrown*; U.S.N.M. No. 15236; China Seas; J. G. Swan, collector.—Rostral one-third higher than broad, scarcely visible from above; a pair of internasals narrowly in contact with rostral; three prefrontals, the median unpaired shield pentagonal, entirely separating the lateral ones, broadly in contact with the internasals in front and the frontal behind; frontal long, as long as parietals, much longer than its distance from tip of snout, wider than supraoculars; parietals short and broad, the two dimensions about equal; nostril large, behind the middle of a long and narrow undivided nasal which is broadly in contact behind with preocular and third supralabial; one high preocular; diameter of eye equaling its distance from edge of lip; two postoculars; 1 + 2 temporals, a small supernumerary shield being interpolated on the left side between anterior temporal and fifth and sixth supralabials; 7 supralabials, first somewhat broader above than below, sixth largest, third and fourth entering eye; mental very small and narrow; only anterior two lower labials reaching chin-shields, the posterior ones being very low, scarcely more than covering the labial edge; two pairs of chin-shields in contact throughout, posterior pair larger than anterior; 25 rows of smooth scales without apical pits; 230 ventrals, without median keel; anal divided; 32 pairs of subcaudals. Color (in alcohol) gray above, each scale pale at tip, yellow on sides and below, with 28 solid blackish brown rings around the body and 3 on tail, the rings being widest (about 5 scale rows) on middle of back, narrowest (about 1½ ventral) on belly, the pale interspaces above averaging about 4 scale rows in width, and 6 to 7 ventrals on underside; a broad yellow band from sides of neck across occiput, two scale rows behind parietals; head in front of this nearly to anterior edge of frontal blackish brown; snout, a superciliary line, entire upper lip and tip of lower lip yellow; a black band narrowly surrounding eye and extending over lower postocular and temporals; lower lip also blackish brown, the black of top of head, postocular band, and lower labial band united by a vertical black band behind angle of mouth.

Dimensions.

	mm.
Total length.....	692
Snout to vent.....	626
Vent to tip of tail.....	66

Adult specimens of this species are known having reached a length of nearly 1,300 mm.

Variation.—As in the other species of this genus the scale formula is very constant. A specimen (No. 149) from Queensland, in the Zoological Museum in Christiania, which I was allowed to examine through the kindness of Prof. Robert Collett, is remarkable for lacking the azygous shield between the prefrontals, thus resembling *L. laticaudata* in this respect, though otherwise a normal *L. colubrina*, as shown by its 23 scale rows and yellow supralabials.^a

Habitat.—No specimen has been recorded from any definite locality within the limits of this work.^b There are in collections, however, several specimens said to have been taken in the "China Seas," or "China Sea." As the latter is bounded by Formosa in the north, while on the other hand the range of this species mostly coincides with that of *L. laticaudata*, it seems quite probable that *L. colubrina* may be found in the seas about Formosa and the Riu Kius now that attention has been called to it.

As stated above, its general distribution is almost identical with that of *L. laticaudata* inasmuch as it is found from the Bay of Bengal to the western part of the South Pacific at least as far east as the Friendly Islands, extending southward to Australia and New Zealand.


List of specimens of *Laticauda colubrina*.

U.S. N. M. No.	Sex and age.	Locality.	When col- lected.	By whom collected or from whom re- ceived.	Scale	Ventrals.	Subcaudals.	Dark rings on body.	Oculars.	Temporals.	Supralabials.
					rows.						
15236	Halfgr. ^a	China Seas.....		J. G. Swan.....	25	230	32	28	1.2	1-2+2	7

^a Description, p. 407.

^a Dr. Franz Werner (Mitt. Zool. Samml. Mus. Naturk. Berlin, I, Pt. 4, 1900, p. 104) mentions a similar specimen in the collection of the Vienna university, but he regards it as a hybrid between *L. laticaudata* and *L. colubrina*, a view I can not accept, at least for the specimen in the Christiania Museum, which is in every respect a typical *L. colubrina*, but lacking the unpaired prefrontal.

^b Doctor Wall enumerates the species as No. 25 of his "List of Japanese and Loo Choo Islands Ophidia," but gives no reference to any record in substantiation.



LATICAUDA SEMIFASCIATA. (FROM TEMMINCK AND SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

LATICAUDA SEMIFASCIATA^a (Reinwardt).

ERABU-UNAGI.

Plate XXII.

1837. *Platurus semifasciatus* REINWARDT, in Schlegel, Phys. Serp., II, p. 516 (type-locality, Moluccas; type, Leiden Mus. No. 1468; Reinwardt, collector).
1837. *Hydrophis colubrina* SCHLEGEL, Phys. Serp., Atlas, pl. XVIII, figs. 18-20; Fauna Jap., Rept., 1837, p. 92 (Liukius); Ophid., pl. x^b (probably Reinwardt's type).—*Platurus colubrinus* FISCHER, Jahrb. Wiss. Anst. Hamburg, V, 1888, p. 18 (Okinawa shima; Tonga; Society Is.).
1856. *Platurus fasciatus* var. *semifasciata* FISCHER, Abh. Naturw. Ver. Hamburg, III, p. 30, p. 70 (Reinwardt's type).
1860. *Platurus fasciatus* var. HALLOWELL, Proc. Phila. Acad., 1860, p. 489 (Loo Choo; Cleopatra I.).
1874. *Platurus schistorhynchus* GUENTHER, Proc. Zool. Soc. London, 1874, p. 297, pl. XLV, fig. B (type-locality, Savage Island; types in Brit. Mus.).—BOULENGER, Fauna Brit. India, Rept., 1890, p. 395 (China Seas); Cat. Sn. Brit. Mus., III, 1896, p. 309 (Savage I.; Loo Choo Is.).—WALL, Proc. Zool. Soc. London, 1903, p. 101 (Loo Choos); 1905, II, p. 517 (Miyako; Okinawa; Amami).
1876. *Platurus fasciatus* HILGENDORF, Mitth. Deutsch. Ges. Ost-Asiens, I, heft 10, pp. 30, 31 (Prov. Satsuma) (not of Latreille).—DOEDERLEIN, Mitth. Deutsch. Ges. Ost-Asiens, III, heft 24, July 1881, p. 149 (Amami-o-shima); heft 25, Dec. 1881, p. 210 (Liukiu Is.; Erabu shima).—OKADA, Cat. Vert. Japan, 1890, p. 69. (Okinawa shima).
1876. *Platurus laticaudatus* PETERS, Mon. Ber. Berlin Akad. Wiss., 1876, p. 534 (Matuka, Fiji Is.) (not of Linnaeus); 1877, p. 417 (part: Berlin Mus. No. 9663, Tonga; 9071, Samoa).

Reinwardt's *P. semifasciatus* must take precedence over Guenther's *P. schistorhynchus*. I have examined Reinwardt's type in the Leiden Rijksmuseum (No. 1468; probably the same individual which Schlegel has figured, Fauna Japonica, pl. x, especially fig. 2) from the "Moluques," and find it an unquestionable specimen of this species, with divided rostral, one pair of adjacent chin-shields, and keeled posterior ventrals.

Concurrent with the two apparently disconnected habitats of this species there seems to be an average difference in the number of ventrals. Thus in the eastern group (Tonga, Savage I.), in six recorded specimens the ventrals vary between 177 and 202 (average 187), while in eight western specimens (Riu Kius, Moluccas), they vary between 197 and 212 (average 205). Unfortunately there are two other specimens recorded which to some extent spoil this harmony, viz, one from the Riu Kius recorded by Boulenger^c with only 188 ventrals, the other, from the Society Islands, recorded by Fischer,^d with no less than 229 ventrals. With regard to the latter, I am


^a From *semi*-, half; *fasciatus*, banded.

^b Reproduced in this work on Plate XXII.

^c Cat. Sn. Brit. Mus., III, p. 309: specimen *c*.

^d Jahrb. Wiss. Anst. Hamburg, V, p. 19: Hamburg Mus. No. 384.

strongly inclined to doubt either the accuracy of the count, the locality, or the identification, the number being so out of all proportion. I may finally remark that in the only Savage Island specimen examined by me (U.S.N.M. No. 28194) the number of light body rings is only 23. Guenther, in the original description of his Savage Island specimens, also gives 22 as the number of the body rings. In our Riu Kiu specimens the number of body rings varies between 35 and 43, and in the type of *L. semifasciata* the number is apparently about 38. No detailed record of the other specimens is accessible. It is consequently at the present moment impossible to decide whether we really have to do with two separable races, one with an average number of 205 ventrals and 38 body rings, typical *L. semifasciata*, from the Riu Kius and the Moluccas, and another form averaging 187 ventrals and 23 body rings, *L. semifasciata schistorhynchus*, from the Friendly Islands.



FIGS. 331-333. *LATICAUDA SEMIFASCIATA*. $1\frac{1}{2} \times$ NAT. SIZE. 331, TOP OF HEAD; 332, SIDE OF HEAD; 333, UNDERSIDE OF HEAD. No. 7515, U.S.N.M.

Description (figs. 331-333).—*Halfgrown*; U.S.N.M. No. 7336; Yokoshima (Cleopatra Island), Tokara group, Riu Kiu Archipelago; May, 1855; Captain Stevens, collector. Rostral broader than high, upper edge broad and truncate, scarcely visible from above; three internasals, one unpaired anterior adjoining the rostral, of which it is in reality only a detached portion, and two posterior normal ones broadly in contact; three prefrontals, a median pentagonal one, posteriorly broadly in contact with frontal, and two lateral ones, broadly in contact with frontal and with supraocular; frontal large, much longer than its distance from tip of snout and than the parietals, supraoculars as broad as frontal at the middle; parietals very short, not longer than broad, much shorter than frontal; nostril large, semilunar, near the middle of the long and narrow undivided nasal; no loreal; one preocular, broadly in contact with nasal; eye rather small, its vertical diameter less than its distance from edge of lip; two postoculars;

temporals 2 + 3, only slightly differentiated from the adjacent scales; seven supralabials, third and fourth largest and entering eye, first as wide above as below; seven lower labials, of which the first pair behind the small mental does not reach the edge of the lip, the labials from the third backward very low, only the first three in contact with chin-shields, of which only the anterior pair is clearly differentiated, the posterior being represented by two scales separated by one of nearly the same size; 23 rows of smooth scales without apical pits; 205 ventrals, on the posterior half of the body by a median blunt keel and a corresponding notch in the posterior edge of each scute; anal divided; 40 pairs of subcaudals. Color (in alcohol) bluish gray, darker above, paler underneath, with 43 dark brown rings around the body and seven on the tail, the bands being widest on the median line of the back, viz, about $3\frac{1}{2}$ scales wide, and there separated by a pale interval only two scales wide; the rings are about $2\frac{1}{2}$ ventrals wide on the underside and the light intervals about the same width; head uniform dark brown, with a yellowish horseshoe-shaped mark, the convexity of which rests on the prefrontals extending backward on the outer edge of supraoculars, upper postocular and upper temporals to and joining the first pale cross line on occiput a scale row behind the parietals; snout and labials dark brown like the rest of the head.

Dimensions.

	<i>mm.</i>
Total length.....	582
Snout to vent.....	507
Vent to tip of tail.....	75

The *young* (in alcohol) are of a light bluish gray with blackish brown rings and markings. The latter as the snake grows larger become lighter and the former darker and browner, while the demarcation between them becomes more obscure until in very large specimens the markings become almost obliterated. In the larger specimens therefore the dark gray cross markings correspond to the whitish cross markings in the young.

This species grows to a considerable size. The largest specimen in our collection (No. 5546) measures 1,097 mm. in total length, with a tail 136 mm. long, while the type measures, respectively, 1,118 mm. and 140 mm.

Variation.—There is very little variation in the scale formula proper, for only in one specimen (No. 5546) have I seen 4 temporals on one side, the normal number of 3 occurring on the other. In eastern specimens the number of ventrals is rarely as low as 188, but ranges usually between 197 and 212, while the subcaudals vary between 32 and 43 pairs. Sometimes anomalies are found in the internasals; thus in No. 106 of the Imperial Museum, Tokyo, there is a small unpaired shield behind the detached part of the rostral, broadly in contact with it and with the unpaired median prefrontal, and in our No. 7515

there are two unsymmetrical shields detached from the left internasal, as shown in fig. 331.

The variation in color and pattern has been referred to above (pp. 410 and 411).

Habitat.—Thus far this species has been recorded from only two disconnected areas very far apart, namely, from the Malay Archipelago (Amboina, according to Peters, and the type from the "Moluccas") and the Riu Kius in the west and from the Fiji, Samoa, and Tonga (Friendly) islands, the neighboring isolated Savage Island, and possibly the Society Islands in the east. No specimen from the intervening seas have been recorded, so far as I know.

In the Riu Kius this species appears to be common, as it can scarcely be doubted that this is the species of which Doederlein speaks as *Platurus fasciatus*, and which according to him is caught in considerable numbers at Erabu shima, in the northern group, so numerous in fact that the snake takes its name, Erabu-unagi, from that island, whence it is shipped to Japan to be used as a medicine. The Rogers Expedition brought home three specimens from Yoko shima (Cleopatra Island), which lies between the Tokara (Linschoten) group and Amami-o-shima, where Doederlein collected it. The Imperial Museum in Tokyo has specimens from Okinawa shima, where Doctor Warburg and Pryer's collector also obtained it. Doctor Wall mentions 29 specimens collected at Miyako shima. Finally, I have examined a specimen in the Hamburg Museum (No. 2633), collected by Doctor Lenz in Iriomote shima of the Yaeyama group.

List of specimens of Laticauda semifasciata.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received	Scale rows.	Ventrals.	Subcaudals.	Dark rings on body.	Ocellars.	Temporals.	Supraorbials.
U.S.N.M.	5546	Female.	Cleopatra Isl.	U. S. Expl. Exped.(?)	23	197	32	1, 2+3	7	7
Do	7336	Halfgr. ^ado.....	1855.	Capt. Stevens	23	205	40	43	1, 2	2+3	7
Do	7380	Young.	Loo Choo	W. Stimpson ^b	23	204	40	37	1, 2	2+3	7
Do	7515	Female ^b	Cleopatra Isl.	May	Capt. Stevens	23	201	37	35	1, 2	2+3	7
Imp. Mus. Tokyo.	10 ^c	Halfgr.	Okinawa shima	23	212	43	37	1, 2	2+3	7
Do	10 ^c	do	do	23	208	41	35	1, 2	2+3	7
Hamburg.	(c)	(c)	Okinawa shima	Dr. O. Warburg.	23	210	41
Do	2633	Adult	Iriomote shima	1897.	Doctor Lenz.	23
Brit. Mus.	(d)	Female	Loo Choo Is.	H. Pryer	188	36

^a Description p. 410.

^b Probably "No. 7-198. *Hydrophis?* Taken off the south end of Loo Choo, December, 1854, by Captain Rodgers" of Stimpson's MS. Catalogue. In the Museum register the entry is only as above.

^c Figs. 331-333.

^d Fischer, 1888, p. 18.

Genus *EMYDOCEPHALUS*^a Kreffft.

1869. *Emydocephalus* KREFFFT, Proc. Zool. Soc. London, 1869, p. 321 (type, *E. annulatus*).

Boulenger, with specimens of another species of this genus before him, still persists (Willey, Zool. Results, 1898, p. 57) in placing these snakes in the genus *Aipysurus* which he has diagnosed as having the "maxillary a little longer than the ectopterygoid [transpalatine];" "poison-fangs . . . followed, after a short interspace, by 6 to 10 grooved teeth."

In the genus *Emydocephalus*, on the contrary, the maxillary bone is shorter than the ectopterygoid, very much as in *Laticauda* (see Boulenger, Fauna Brit. India, Rept., p. 394, fig. 116A) *even slightly more reduced*. Behind the fang there is no trace of another tooth, such as is found in *Laticauda*, not even an impression on the surface of the bone. Add to this the remarkable and apparently constant fusion of the median labials, which some authors might regard as a good generic character in itself, and the status of *Emydocephalus* seems assured. So far as I can see, it is more closely allied to *Laticauda* than to *Aipysurus*, the chief difference being the presence of internasals and the consequent more lateral position of the nostrils.

The genus, which is as yet but very poorly known, seems to contain three species, the one here described, from the Riu Kiu and Formosan waters; *E. chelonicephalus*, from the Loyalty Islands; and *E. annulatus*, with single anal, and strongly tuberculated scales, from a locality, probably somewhere in Australian waters.

In discussing the specific characters of the species constituting the genus *Aipysurus*, Doctor Guenther^b speaks of the head shields being much subdivided in old examples, implying that they are not so much subdivided in the young. While our present species is not strictly congeneric with those treated of by Doctor Guenther, it seems nearly enough related to make it proper to question the above inference. I have before me an adult female with two embryos ready to be expelled, and in all three the scutellation of the head is so identical that the conclusion seems justified, that no change takes place as the snake grows older, and that where a difference occurs it is due either to individual or to specific variation.

EMYDOCEPHALUS IJIMÆ^c Stejneger.

1898. *Emydocephalus ijima* STEJNEGER, Journ. Sci. Coll. Tokyo, XIII, Pt. 3, p. 223 (type-locality, Riu Kiu Sea; type, U.S.N.M. No. 36517; Tashiro, collector).

1899. *Aipysurus annulatus* BOULENGER, Zool. Record for 1898, Rept. Brit., p. 21 (not of Kreffft).—WALL, Proc. Zool. Soc. London, 1903, pp. 95, 101; 1905, II, p. 517 (Okinawa).

^a From *ἐμύς*, turtle; *κεφαλή*, head.

^b Rept. Brit. India, p. 357.


^c Named in honor of Prof. Isao Ijima, Science College, Imperial University, Tokyo.

It has been intimated by Boulenger (Zool. Rec. for 1898, Rept. Batr., p. 21) that this species is identical with Krefft's *E. annulatus* (of unknown habitat) and Bavay's *E. chelonicephalus*, from the Loyalty Islands. I shall express no opinion here as to whether he is right in thinking it "advisable to consider, provisionally, at least," the latter which has a rostral bearing a conical, spine-like tubercle to be the same species as Krefft's snake,^a but I must dissent most emphatically from having *E. ijimæ* included with any of them. Its four prefrontals, divided anal, enlarged posterior vertebrals and smooth scales seem at once to forbid such a procedure, notwithstanding the fact that all the specimens of this genus show a remarkably small range in the number of their ventrals. That we have not to do with individual or sexual variations is proven by the fact that I have seven specimens, mature ones of both sexes as well as young ready to be born, which agree in all essential points. They all possess the enlarged vertebrals and four prefrontals, thus making out a pretty conclusive case for the distinctness of this species. Doctor Wall, who examined three of the specimens now in the National Museum, remarks that "an examination of the specimens [of *E. chelonicephalus*?] in British Museum shows that a certain slight enlargement of the vertebrals is present at some spots, but no specimen has four prefrontals." It should be mentioned, however, that in a specimen recently received by British Museum from Mr. Owston (No. 1903.6.29.23) as collected in the Yaeyama Islands, the two outer prefrontals are fused with the inner ones, and that Doctor Wall has since recorded another with similarly fused prefrontals.

Description.—*Adult female*; Science College Museum, Tokyo, No. 1; Riu Kiu Seas; 1888; Tashiro, collector (figs. 334-337). Head short; body rather short and stout; rostral broader than high, pentagonal, the labial border with a posterior projection which fits into the mental groove; nasals as long as frontal, twice as long as suture between prefrontal, and wider than long, in contact on median line and with very large semilunar nostrils situated on top near posterior border, the concavity directed forward; two pairs of prefrontals, the median pair large, pentagonal, in contact with nasals, frontals, supraocular and outer prefrontals which are small and nearly square and in contact with median prefrontals, nasals, preocular, and supraocular; frontal rather small, hexagonal, about as broad as long; supraoculars similar, but slightly longer; parietals rather large, about as

^aOne of our specimens of *E. ijimæ*, U.S.N.M. No. 33942, has also a very pointed conical tubercle on the rostral as well as numerous other pointed asperities or tubercles on the anterior facial shields. Another (No. 33943) has tubercles on other head shields, though the one on the rostral is not so prominent and pointed as in the other. Both of these specimens are males. The third is a female with entirely smooth shields. This character then seems to be sexual and not specific.

broad as long; one preocular, pentagonal; two postoculars, the upper largest, followed by two similar, but slightly larger, temporals; three supralabials, only the first and last being of normal size, the second forming a large plate corresponding to at least four ordinary supralabials, and bordering the eye below; mental deeply notched in front and divided by a deep median groove, followed behind by a pair of narrow shields which do not reach the commissure; behind these a pair of large chin-shields in contact, followed by another pair separated by an unpaired shield slightly smaller; three lower labials, on right side, four on left, first and third (and fourth) quite small, second forming a large plate corresponding to the second supralabial; dorsal



FIGS. 334-337.—*EMYDOCEPHALUS IJIMEI*. NAT. SIZE. 334, TOP OF HEAD; 335, SIDE OF HEAD; 336, UNDERSIDE OF HEAD; 337, SCUTELLATION AT MIDDLE OF BODY. NO. 1, SCI. COLL. TOKYO.

scales large, strongly imbricate, 15 rows around neck, 17 around middle of body, subequal, except median row which on the posterior four-fifths of the body and tail is greatly enlarged, twice as wide as the others, hexagonal and gastrostege-like; the scales are smooth, but under the magnifying glass show a few slight tubercle-like elevations; ventrals 140, wide with a rounded median keel in the posterior half; anal large, regular, divided; subcaudals, 23, undivided; basal half of tail with four scale rows on each side between the marginals above and below, the terminal half with three such rows; a large terminal rounded shield, irregularly frayed along the edges. Color (in alcohol)

of a rich dark chestnut with narrow, bright ochraceous yellow cross-bands round the entire body, one to two scales wide on the back and two to three scales on the sides, and about four scales distant; a similar yellow band from angle of mouth through anterior temporals, upper postocular, outer half of supraocular to median prefrontals; a few irregular yellowish marks on second supralabial and on throat.

Dimensions.

	<i>mm.</i>
Total length.....	890
Snout to vent.....	778
Vent to tip of tail.....	112

Variation.—A *young* specimen (U.S.N.M. No. 36517), ready to be born and taken from the abdomen of the above described specimen, is a perfect counterfeit of the old snake differing only in the following particulars: (1) It has 17 scale rows around the neck as well as around the middle of the body; (2) the mental is large, nearly heart-shaped, with a deep median groove, and the pair of narrow shields between the mental and the first pair of chin-shields is wanting. In other respects, notably the size and shape of the head-shields, the remarkable consolidation of the labials in both jaws, in the great width of the median dorsal scale row as well as in color, the two specimens agree completely. Enough of another young, still attached to the mother was freed from the egg membranes to ascertain that these points are equally developed in this specimen.

A fourth specimen is the *male* from Botel Tobago, mentioned by me when first describing the species. Although about a third smaller than the mature female described above, it appears to be adult, and agrees in almost every particular with that specimen. The only point of difference is that the rostral has a slightly elevated point, somewhat indicative of the spine in *E. chelonicephalus*. This character seems to be confined to the male sex. The specimen in question has 17 scale rows; 142 ventrals; a double anal; and 28 undivided subcaudals. Total length 613 mm.; tail, 93 mm. The fact that this specimen, as well as the three others in the National Museum, agrees with the adult female in having a pair of narrow shields immediately behind the small grooved mental and before the first pair of regular chin-shields, proves that the condition in the young specimens in which these pseudo-chin-shields are fused with the mental is an individual aberration only.

These "postmentals" or "pseudo-chin-shields," however, seem to be the first pair of sublabials which have been crowded away from the lip. The posterior lower projection of the rostral fits snugly into the anterior notch and groove of the mental and closes the mouth effectively.

The number of scale rows varies between 17 and 19, though the former seems to be the normal number; ventrals between 134 and 143, subcaudals between 23 and 28.

A specimen in British Museum, as previously mentioned (p. 414), has the outer prefrontals fused with the inner ones.

Since the above was written Doctor Wall has reported upon six examples obtained by Mr. Owston's collectors in Okinawa. Four had a sharp spine on the rostral, which was absent on the other two. The prefrontals were very irregular. There were four in three specimens, two in one specimen, and three (2 right, 1 left) in two specimens. In all one or both parietals were split by a suture behind. The scales were 17 in mid-body when the vertebral row was enlarged, which was usually the case, though often to a variable degree. In two specimens the scales were 19, the vertebrae being divided into three rows subequal to the other dorsals. Ventrals numbered 138 to 143.

Habitat.—This species is only known from the "Riu Kiu Sea," where Tashiro collected the gravid female in 1888, and from Botel Tobago Island, east of the south end of Formosa. A specimen was caught there on the rocks at ebb tide by Mr. Tsunasuke Tada, May 1897. Three specimens obtained by Mr. A. Owston's collector, on Ishigaki Island of the Yaeyama group, during the month of May, 1899, are now in the United States National Museum, and two are in British Museum, one from Yaeyama Island and one from Iriomote. Doctor Wall now records six specimens from Okinawa.

List of specimens of Emydocephalus ijima.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.
				1899.					
U. S. N. M.	33042	Male....	Yaeyama Islands...	Apr.-May	A. Owston...	19	139	2	28
Do.....	33943	do.....	do.....	do.....	do.....	19	138	2	28
Do.....	33944	Female	do.....	do.....	do.....	17	140	2	24
Do.....	36517	Young ^a	Riu Kiu Seas.....	1888.	Tashiro.....	17	138	2	23
Sci. Coll. Tokyo.	1	Female ^b	do.....	1888.	1888.	17	139	2	23
				1897.					
Do.....	1a	Male....	Botel Tobago Island.	May	T. Tada.....	17	142	2	28
Brit. Mus.	1901. 10. 23. 8	do.....	Iriomote shima.....		Doctor Wall.	19	136	2	27
Do.....	1903. 6. 29. 23	Female.	Yaeyama Island.....		A. Owston...	19	134	2

^a Type, p. 416.

^b Description, p. 414; figs. 334-337.

Genus *DISTEIRA*^a Lacépède.

1803. *Hydrophis* DAUDIN, Hist. Nat. Rept., VII, p. 372 (part: not of Latreille, 1802).
1804. *Disteira* LACÉPÈDE, Ann. Mus. d'Hist. Nat., Paris, IV, p. 210 (type, *D. dohata*).
1804. *Leioselasma* LACÉPÈDE, Ann. Mus. d'Hist. Nat., Paris, IV, p. 210 (type, *L. striata*).
1830. *Enhydris* WAGLER, Nat. Syst. Amph., p. 166 (type, *Hydrophis cyanocincta*) (not of Latreille, 1802).
1834. *Microcephalophis* LESSON, in Bélanger's Voy. Indes Orient., Rept. (p. 320): Atlas, Rept., pl. III (type, *M. gracilis*).
1842. *Liopala* GRAY, Zool. Misc., p. 60 (type, *Hydrus gracilis* Shaw).
1842. *Liopala* GRAY, Zool. Misc., p. 60 (err. typogr.).
1842. *Aturia* GRAY, Zool. Misc., p. 61 (type, *A. ornata*).
1848. *Noterophis* GISTEL, Naturg. Thierr., p. IX (substitute for *Enhydris* Wagler).
1849. *Chitalia* GRAY, Cat. Sn. Brit. Mus., p. 56 (type, *Ch. inornata*).
1849. *Kerilia* GRAY, Cat. Sn. Brit. Mus., p. 57 (type, *K. jerdonii*).
1852. *Thalussophis* SCHMIDT, Abh. Naturw. Ver. Hamburg, II, Pt. 2, p. 75 (substitute for *Hydrophis* Daudin).
1887. *Distira* COPE, Bull. U. S. Nat. Mus. No. 32, p. 61 (emendation).

The large central group of Hydrid snakes, about 40 species, has been gathered into two genera by Boulenger according to whether the maxillary teeth following the first pair of poison fangs are grooved (*Disteira*) or "solid" (*Microcephalophis* = *Hydrophis* Daudin, 1803, not Latreille, 1802). Recently some doubt has been thrown on the validity of this separation, inasmuch as Mr. West, in a typical *Disteira*, viz, *D. cyanocincta*, found that "the first two or three only of these teeth showed any indications of a groove."^b

Relying, however, on the division established by Boulenger, I identified some of my specimens in which no grooved teeth were observable under a magnification of 10 diameters, as *Microcephalophis melanocephalus*.^c Three years later some similar snakes came under my notice, which, investigated by the same means, showed distinct grooves in the maxillary teeth, and as their other characters did not correspond to those of any other *Disteira*, I described them as a new species, *D. orientalis*.^d While preparing detailed descriptions for the present work, however, I was forcibly struck by the fact that in all external characters the two species were absolutely identical. Mr. West's experience with *D. cyanocincta* and other parallel cases with various opisthoglyph snakes then occurred to me. The specimens at my command were consequently subjected to much greater magnifications, with the result that I find all the teeth grooved in all the specimens examined by me. To see the groove plainly it was often

^a From $\delta\iota\varsigma$, double: $\sigma\tau\epsilon\iota\rho\alpha$, keel.

^b Proc. Zool. Soc. London, 1895, p. 823, pl. XLVI, fig. 8.

^c Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 224.

^d Proc. Biol. Soc. Washington, XIV, 1901, p. 191.

necessary to use a binocular microscope magnifying 45 diameters, and to make sure of the presence of grooves on the anterior face of the maxillary teeth of *Disteira gracilis* it was even necessary to raise the power to 65 diameters. From my experience it is then plain that there is no real difference between the "grooved" and the "solid" maxillary teeth in these snakes, that it is merely a question of degree.

Under these circumstances there seems to be no other course than to unite the two supposed genera (and species!), the name *Disteira* being available for the thus consolidated genus. The term *Hydrophis* must disappear from our nomenclature, as Latreille, in 1802, framed it as a substitute pure and simple for *Hydrus*, while Daudin, in 1803, substituting *Pelamis* for the latter reemployed the former for the bulk of the Hydrid snakes.

Within the enlarged genus the limits and characters of the species offer some of the most difficult problems to the ophiologist. These snakes, in many respects so excessively specialized, often show a correspondingly excessive amount of individual variation, and as the series available for study is insignificant except in the case of a few species the result must of course be in great doubt.

If, for instance, we take the latest monographic account of these snakes, namely, Boulenger's treatment of them in the Catalogue of the Snakes in British Museum, by far the ablest treatise on the genus, based upon the largest material in any collection, we find that of 22 species recognized by him in the genus *Hydrophis* he had 77 specimens, or an average of 3.5 specimens to each species.^a If we analyze this material for the variability of the number of ventrals we come to the result that the range of the ventrals, that is, the difference between the greatest and smallest number of ventrals—of the specimens of each species in British Museum is greater in proportion as the series of specimens examined increases. This is startlingly shown in the following table:

Number of specimens in British Museum.	Range of ventrals.	Name of species.	Number of specimens in British Museum.	Range of ventrals.	Name of species.
13	147	<i>fasciatus.</i>	4	40	{ <i>nigrocinctus.</i> <i>coronatus.</i>
12	128	<i>obscurus.</i>			
10	74	<i>gracilis.</i>	3	35	<i>elegans.</i>
8	24	<i>carulescens.</i>	2	36.5	{ <i>mamillaris.</i> <i>cantoris.</i>
5	42.5	<i>spiralis.</i>			
		<i>torquatus.</i>			

The only exception is the series of eight *H. carulescens* in which the range of variability of the ventrals is unusually low. These specimens however, cover a relatively small geographic range.

^a In no instance did he have more than five specimens from one locality.

The 27 specimens of *D. melanocephala* examined by me are practically from the same locality, and so is the Miyakoshima specimen recorded by Boettger. Even the type of the species collected by Sir. E. Belcher may well have been picked up at the same island. In these 29 specimens the range of the ventrals only amounts to 56, while in the 13 specimens enumerated by Boulenger as *H. fasciatus* it amounts to 147. His 13 specimens, however, cover a range from Malabar to China. If we eliminate the two types of Guenther's *Hydrophis atriceps*, with 364 and 353 ventrals, respectively, which I am inclined to refer to *D. melanocephala*, the variation in the ventrals is reduced to only 98.

The rational explanation seems to be that the individual variation in a fairly large series from the same locality is within reasonable bounds, and that the so-called "species" must be broken up on different lines from those which have been established by Boulenger.

This question is particularly important to us in the present connection, as some of the species which are reported from the waters adjacent to the region here treated of are very closely related, their differences consisting mostly in slight gaps between their scale formulas. The three species alluded to agree in having a very small head and a long body very slender anteriorly. As a rule they have a single large anterior temporal, and two pairs of small chin-shields in contact.

Carefully noting what is said in the following under the discussion of each species the various forms treated of may be distinguished by the following:


KEY TO THE SPECIES OF *DISTEIRA* IN CHINESE ^a AND JAPANESE WATERS.

- a*¹ Frontal shield longer than broad.
*b*¹ Ventrals more than 295.
*c*¹ A single anterior temporal (normally).
*d*¹ Scale rows around the neck more than 23 (25-33).
*e*¹ Scale rows around body 38, or less (38-32); ventrals less than 371 (301-370).
D. melanocephala, p. 421.
*e*² Scale rows around body 40, or more (40-56); ventrals more than 399 (400-500).....[*D. fasciata*] p. 427.
*d*² Scale rows around neck less than 23 (19-21)[*D. gracilis*] p. 427.
*e*² Two anterior temporals.....*D. cyanocincta*, p. 428.
*b*² Ventrals less than 295.....*D. godeffroyi*, p. 430.
*a*² Frontal shield as broad as long.....[*D. viperina*] p. 434.

^a The six snakes mentioned by Peters (Mon. Ber. Berlin Akad. Wiss., 1872, p. 859, under the name of *Hydrophis diadema* Guenther, from the coast of China and Siam (Berlin Mus. Nos. 4428, 7428, and 4747) can scarcely belong to that species, if it is identical with *D. obscura*, for which Boulenger gives the following scale formula: Scales around neck, 33-40; around body, 40-50; ventrals, 310-438. Peters's specimens have 31-33 neck scales, 37-40 body scales, and the one specimen of which the ventrals were given has 296. This scale formula does not agree with any of the species as diagnosed by Boulenger.



1



2



4



3



5

DISTEIRA MELANOCEPHALA. (FROM BOULENGER.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

DISTEIRA MELANOCEPHALA^a (Gray).

Plate XXIII.

1849. *Hydrophis sublaevis*, var. *melanocephala* GRAY, Cat. Sn. Brit. Mus., p. 53 (part: type-locality, "Indian Ocean;" type in Brit. Mus.; Sir E. Belcher, collector).
- 1864 ?*Hydrophis atriceps* GUENTHER, Rept. Brit. India, p. 371, pl. xxv, figs. I-I' (type-locality, Siam; types in Brit. Mus.).
1895. *Hydrophis fasciatus* BOETTGER, Offenbach. Ver. Naturk. 33-36 Ber., 1895, p. 111 (Miyakoshima, Riu Kiu); Kat. Schl. Mus. Senckenberg., 1898, p. 113 (Miyakoshima) (not of Schneider).
1896. *Hydrophis melanocephalus* BOULENGER, Cat. Sn. Brit. Mus., III, p. 283, pl. xv^b (Indian Ocean; type).—WALL, Proc. Zool. Soc. London, 1903, pp. 95, 101 ("Loo Choos"); 1905, II, p. 516 (Ishigaki; Iriomote).—*Microcephalophis melanocephalus* STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 224 (Pescadores Is.).
1901. *Disteira orientalis* STEJNEGER, Proc. Biol. Soc. Washington, XIV, Dec. 12, 1901, p. 191 (type-locality, Okinawa shima; type No. 29, Imper. Mus., Tokyo).
1903. *Disteira subcineta* WALL, Proc. Zool. Soc. London, 1903, p. 95 (part: "Japan, Loo Choos;" not of Gray).


The line drawn between this and the following species, *D. fasciata*, viz, at the 370-399th scale rows, is possibly an arbitrary one, but I am inclined to believe that in this way we obtain better results and come nearer the truth. It is not certain, however, that other characters coincide with such a division, and possibly it might be better to separate the two forms according to the scale rows around the body, but the records are very defective on this point. On the other hand, there may even be three forms, one more or less confined to the coasts of India and the Bay of Bengal, with the maximum average of ventrals and body scales; another intermediate form from the Malay island world and adjacent coasts of the mainland, and a third, with a more eastern habitat, including the China Seas, having the minimum number of ventrals and scales.^c

^a From μέλας, black; κεφαλή, head.

^b Reproduced in this work on Plate XXIII.

^c There is a very serious obstacle to the defining of these forms from the records, viz, the difficulty in counting the scales and scutes of these snakes exactly. In the records we find the most divergent figures for the same specimens. Thus among the snakes included by Boulenger in this species, British Museum Nos. *a*, *d*, and *e* seem to be three of the four specimens mentioned by Guenther (Rept. Brit. India, p. 371) as *H. lindsayi*. These four he gives as having, respectively, 345, 390, 424, and 449 ventrals, while Boulenger's count is 402, 412, and 452. Guenther states that the number of ventrals of the type of his *H. atriceps* is 376; Boulenger counts 364. Guenther describes Gray's *H. ocellata* as having 296-334 ventrals, mentioning three specimens, one of these the type; according to Boulenger (p. 291), these three specimens (including the type) have 274, 275, and 290 ventrals; Guenther examined four *H. ornata*, among which the type and that of *H. inornata* and gives the ventrals of the species as 252-260; according to Boulenger the ventrals of the two types mentioned are 232 and 240.

The case would perhaps be less complicated if we knew all the data relating to the types of Guenther's *H. atriceps*. They seem to have a sufficiently low number of ventrals to be included within *D. melanocephala*, but I can find no details relating to the number of scales round the body. Guenther gives the number of scales around the neck as 26 and 28, respectively, consequently well within the limits of *D. melanocephala*, and I have consequently referred them, provisionally at least, to the latter rather than to *D. fasciata*.




FIGS. 338-340.—*DISTEIRA MELANOCEPHALA*. 2 × NAT. SIZE. 338, TOP OF HEAD; 339, SIDE OF HEAD; 340, ROSTRAL. NO. 4, SCI. COLL. TOKYO.

Boettger's *H. fasciatus*, from "Miyakoshima," with 27 scales round the neck, 38 round the body, and 355 ventrals, is clearly identical with the twenty-one specimens from the same locality now before me.

With regard to my *Disteira orientalis*, which was based upon specimens with posterior maxillary teeth plainly showing grooves

under a low magnification, and consequently believed then to belong to a different genus, see remarks above under the genus (p. 418).

Description (figs. 338-347).—*Adult female*; U.S.N.M. No. 33945; Ishigaki shima, Yaeyama group, Riu Kiu archipelago; April-May,




FIGS. 341-344.—*DISTEIRA MELANOCEPHALA*. 2 × NAT. SIZE. 341, TOP OF HEAD; 342, SIDE OF HEAD; 343, UNDERSIDE OF HEAD. 2½ × NAT. SIZE. 344, ROSTRAL. NO. A40, HIGH SCHOOL, KUMAMOTO

1899; A. Owston collection. Rostral much wider than high, well visible from above; nasals very long, the suture between them twice as long as that between the prefrontals; the semicircular valvular nostril situated near the outer posterior corner of the nasal; frontal hexag-

onal, slightly longer than its distance from rostral, shorter than parietals, half as wide as long, wider than supraoculars; parietals as long as frontal and prefrontals together; no loreal; one preocular; one postocular; one large anterior temporal; 7 supralabials, second in contact with prefrontal, third and fourth entering eye, sixth very small; 3 lower labials in contact with anterior chin-shields; posterior chin-shields nearly as large as anterior, separated by a small scale; 25 rows of smooth, imbricate scales

round the neck, 35 rows round the middle of the body, the scales on the body being imbricate and provided with a short keel which reaches neither base nor tip of scale; about 340

ventrals which often are so reduced as to be distinguished with difficulty from the scales, each ventral with two incomplete keels; anal divided, with an elongated smaller shield on each side; 40 subcaudals; terminal scale of tail small, only differentiated on one side. Color (in alcohol) above cinereous, white below, with 50 blackish rings round the body and 5 on the tail; head black with superciliary region and snout irregularly pale, the posterior supralabials being likewise light colored with a brownish tinge.



FIGS. 345-347.—*DISTEIRA MELANOCEPHALA*. NAT. SIZE. 345, TOP OF HEAD; 346, SIDE OF HEAD; 347, UNDERSIDE OF HEAD. No. 14, IMPER. MUS. TOKYO.

Dimensions.


Head very small; neck and anterior part of body very slender, increasing in thickness backward, the greatest height of the body being nearly 4 times the diameter of the neck.

	<i>mm.</i>
Total length.....	1, 125
Snout to vent.....	1, 027
Vent to tip of tail.....	98
Width of head.....	15
Diameter of neck.....	11
Greatest height of body.....	42
Greatest height of tail.....	26

The *male* is relatively much slenderer posteriorly.

Adult male; Imperial Museum, Ueno Park, Tokyo, No. 29; Okinawa shima, Riu Kiu Archipelago (figs. 348-350). Rostral nearly as high as broad, somewhat projecting, the portion visible from above longer than interprefrontal suture; nasals long, the suture between them twice as long as the one between the prefrontals, nostrils semilunar close to the outer posterior angle; frontal long and narrow, the length equaling distance from tip of snout and length of parietals, the width at the middle equaling that of the supraoculars; parietals long,

the suture between them slightly longer than internasal suture; no loreal; one preocular; one postocular; one anterior temporal; 7 supralabials, sixth smallest, second largest, broadly in contact with prefrontal third and fourth entering eye; 3 lower labials in contact with anterior chin-shields which are somewhat shorter than the posterior, the latter broadly in contact anteriorly; 25 rows of smooth scales on the neck, 33 shortly keeled or tuberculated ones on the middle of body; 323 bicarinate ventrals, often scarcely differentiated; 4 subequal, elongate anals in a line. 43 subcaudals; terminal scales of tail



FIGS. 348-350.—*DISTEIRA MELANOCEPHALA*. NAT. SIZE. 348, SIDE OF HEAD; 349, SCUTELLATION AT MIDDLE OF BODY ($\frac{1}{4}$ \times NAT. SIZE); 350, SIDE OF TAIL. No. 23, IMPER. MUS. TOKYO.

but slightly differentiated. Color (in alcohol) very pale yellow, more ochraceous on back, with 50 blackish rings on back, six on tail, wider on the back and belly and confluent on the anterior third of the latter into a black ventral band, the dark rings narrower than the pale interspaces; head blackish with an irregular yellow spot on most of the shields, except posterior supralabials; lower labials, chin, and throat blackish; tip of tail black.

Dimensions.

Head small; neck and body slender, especially anteriorly.	mm.
Total length.....	1,080
Snout to vent.....	985
Vent to tip of tail.....	95
Width of head.....	8
Diameter of neck.....	7
Greatest height of body.....	27
Greatest height of tail.....	21

The *young* resemble the adults except that the black color is more extended, so that the coloration of the snake may be described as

black with narrow whitish rings which are mostly interrupted on the underside, the whole lower neck and a more or less complete longitudinal band on the ventrals being black; head entirely black with a whitish postocular band and a pale spot on each of the prefrontals; tail black with a few lateral, vertical, white bars or spots.

Variation.—The extreme specialization of these snakes is accompanied by a correspondingly great variability. Thus the reduced ventrals are so irregular and so often subdivided to the size of the adjacent scales that their number in a large series shows an excessive range, a feature alluded to above under the preliminary discussion of the species (p. 420). The head shields also show considerable variation, though to a less degree. The supralabials are 7, as a rule, but 8 are occasionally found, and 6 very seldom; three of them may enter the eye; two postoculars occur, but one is the normal number; posterior chin-shields in contact as a rule, but occasionally separated by a scale. The most notable variation, however, is the occasional, though rare, presence of two anterior temporals, because the scale formula of such specimens reads like that of *D. cyanocincta*, in which two anterior temporals is the rule. The arrangement, however, is different, for while in the former the two anterior temporals are produced by the horizontal division of the larger temporal located above the very small sixth supralabial, in the latter it is caused by the horizontal division of the fifth supralabial. In the former case we have the very small sixth supralabial between the much larger fifth and seventh, while in the latter the three are subequal or the sixth intermediate in size between the other two.

In the Pescadores Island specimens the small sixth supralabial on the left side is abnormally fused with the fifth, as shown in fig. 339, but on the other side it is normal.

Habitat.—The locality whence came the type of this species is given as "Indian Ocean," but as it is one of specimens brought home by Sir E. Belcher, this probably means only that it was obtained during his cruises in the "Samarang" in which case it is just as likely to have been collected in the China Sea or even in the waters about the Riu Kius, where it is evidently a common snake, as I have examined 24 specimens from the southern group (Saki shima), one from Okinawa, and one from the Pescadores in the Formosa Channel. Here also belongs the *H. fasciatus* recorded by Boettger from "Miyakoshima." Finally there is the specimen in the Rijksmuseum in Leiden (No. 1483). It is labeled "*Hydrophis striatus*, von Siebold, Japan." However, it is not the specimen figured under that name in Fauna Japonica (Ophid., pl. VII), but most probably the specimen referred to in the text (p. 89) as "harponné près des îles Lioukiou au 27^m degré de lat. bor." apparently the only specimen collected by

von Siebold.^a It would therefore almost appear as if this species were confined to the neighborhood of the Riu Kiu Islands.

List of specimens of Distira melanocphala.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows on neck.	Scale rows on body.	Ventrals.	Subcaudals.	Supranals.	Anterior temporals.
1899.											
U.S.N.M.	33945	Female ^a	Ishigaki shima, Riu Kiu.	Apr.-May.	A. Owston.	25	35	340	40	7	1
Do.	33946	Male	do.	do.	do.	27	35	323	35	7	1
Do.	33947	do.	do.	do.	do.	25	35	318	48	7	1
Do.	33948	Female	do.	do.	do.	27	37	328	42	7	1
Do.	33949	Male	do.	do.	do.	27	33	314	42	7	1
Do.	33950	do.	do.	do.	do.	25	35	315	44	7-8	1
Do.	33951	Female	do.	do.	do.	31	35	322	43	7-8	1
Do.	33952	do.	do.	do.	do.	29	35	345	40	7	1
Do.	33953	Male	do.	do.	do.	27	35	313	40	7	1
Do.	33954	Female	do.	do.	do.	29	35	345	39	7	1
Do.	33955	Young.	do.	do.	do.	25	33	336	43	7-8	1
Do.	33956	do.	do.	do.	do.	27	33	331	44	7-8	2
Do.	33957	do.	do.	do.	do.	25	35	327	42	7	1
Do.	33958	do.	do.	do.	do.	25	35	339	44	7-8	1
Do.	33959	Male	do.	do.	do.	25	35	357	45	8	1
Do.	33960	Female	do.	do.	do.	27	35	311	35	7	1
Do.	33961	Male	do.	do.	do.	25	35	317	47	7	1
Do.	33962	Female	do.	do.	do.	27	33	356	37	6-7	1
Do.	33963	Male	do.	do.	do.	27	35	353	49	8	1
Kumamoto	A 40	Young ^b	do.	1891		27	33	324	42	7	1
Senckenberg.		(c)	Miyakoshima			27	38	355	53	7	1
Brit. Mus.	a	Female ^d	"Indian Ocean"	1897		25	35	329			1
Do.	b	do.	Iriomote shima			25	32	321			
Sci. Coll., Tokyo.	4	Male ^f	Pescadores Islands.	Feb.	T. Tada.	27	37	301	49	8	1
Imp. Mus., Tokyo.	14	Female ^g	Miyakoshima			27	35	330	39	7	1 2
Do.	29	Male ^h	Okinawa shima, Riu Kiu.			25	33	323	43	7	1
1897.											
Hamburg	2574a		Iriomote shima, Riu Kiu.	Mar. 13	Lenz.	23	34				1
Do.	2574b		do.	do.	do.	24	32				1
Leiden	1483		"Japan"			25	35	341	51		1

^a Description, p. 427.

^b Figs. 341-344.

^c Boettger, p. 111.

^d Type.

^e Boulenger, MS. cat.

^f Figs. 338-340.

^g Figs. 345-347.

^h Type of *D. orientalis*. Description, p. 423; figs. 348-349.

^a The specimen in question (No. 1483) was examined by me in 1898; for the permission to do so thanks are due to the director, Doctor Jentink. It was labeled as stated above, and the count of ventrals and subcaudals gave respectively 341 and 51 scutes. These figures, however, agree only with the third specimen mentioned by Schlegel (Phys. Serp., II, p. 504), who states that this is the one collected by Buerger. Whatever error may have crept in, there can be but little doubt that the specimen was collected practically in the same waters as the others.

[**DISTEIRA FASCIATA** ^a (Schneider).]

1799. *Hydrus fasciatus* SCHNEIDER, Hist. Amph. I, p. 240 (type-locality, unknown; types in Berlin Mus., Nos. 2836-2837).—*Disteira fasciata* FITZINGER, Neue Class. Rept., 1826, p. 55 (India).—*Hydrophis fasciatus* PETERS, Mon. Ber. Berlin Akad. Wiss., 1872, p. 849, pl. I, fig. 1 (types, description).—BOULENGER, Cat. Snakes Brit. Mus., III, 1896, p. 281 (coasts of India to China and New Guinea).
1837. *Hydrophis gracilis* SCHLEGEL, Phys. Serp., II, p. 507 (part: Padang, Sumatra) (not *Hydrus gracilis* SHAW, 1802).—JAN, Icon. Ophid., livr. 41, 1872, pl. IV, fig. 2 (Java).
1842. *Aturia lindsayi* GRAY, Zool. Misc. (p. 61) (type-locality, China; type in Brit. Mus.; W. Lindsay, coll.).
1864. *Hydrophis chloris* GUENTHER, Rept. Brit. India, p. 370 (not *H. cloris* DAUDIN, 1803) (Madras; Pinang).

For the characters supposed to separate this species see the "key" (p. 420) and the discussion under the foregoing species (p. 421). In view of the statements there made a detailed description is deemed superfluous.

Habitat.—Boettger has recorded *Hydrophis fasciatus* from Miyakoshima, but I have given my reasons elsewhere (p. 422) for referring this specimen to *D. melanocephala*. Under these circumstances this species has no standing in the Japanese fauna, but I have included its synonymy here because of the uncertainty of the status of this and allied forms.

As generally understood, *D. fasciata* has a wide distribution from the coasts of India to the Philippines.

[**DISTEIRA GRACILIS** ^b (Shaw).]

1802. *Hydrus gracilis* SHAW, Gen. Zool., III, Pt. 2, p. 560 (type-locality not given).—*Disteira gracilis* FITZINGER, Neue Classif. Rept., 1826, p. 55 (India).—*Microcephalophis gracilis* LESSON, in Bélanger's Voy. Ind. Orient., Rept. (p. 321), Atlas, Rept., 1834, pl. III (Bay of Bengal).—*Hydrophis gracilis* GUENTHER, Rept. Brit. India, 1864, p. 373 (Madras).—BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 280 (Persia to Malay Archipelago).—*Hydrophis (Microcephalophis) gracilis* BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 90 (Hainan).
1827. *Hydrophis kadell-nagam* BOIE, Isis, 1827, p. 554 (based on Russell's Ind. Serp., II, pl. XIII).
1852. *Thalassophis microcephala* SCHMIDT, Abh. Naturw. Ver. Hamburg, II, Pt. 2, 1852, p. 78, pl. II (type-locality, Java; types in Hamburg Mus.).—*Hydrophis microcephalus* DUMÉRII and BIBRON, Expét. Gén., VII, Pt. 2, 1854, p. 1356 (Pondichéry).—*Hydrophis microcephala* FISCHER, Abh. Naturw. Ver. Hamburg, III, 1856, p. 52 (Java).
1856. *Hydrophis leprogaster* DUMÉRII and BIBRON, in Fischer, Abh. Naturw. Ver. Hamburg, III, 1856, p. 53 (substitute for *Th. microcephala*; type-locality, Pondichéry; type in Paris Mus.).

The differentiating characters of this species being chiefly those of proportion and scale formula, a detailed description is not necessary

^a Signifying banded.

^b Signifying slender.

in the present connection. The essential points may be gathered from the "key" (p. 420).

Habitat.—The slender sea snake has a wide distribution from the Gulf of Omar, between Arabia and Persia, to the Malay Archipelago and the South China Sea.

It has been recorded as far north as the island of Hainan and Macao, but not yet from Japanese waters. It is included here, however, to help identification in case its range extends that far.

DISTEIRA CYANOCINCTA ^a (Daudin).

1803. *Hydrophis cyanocinctus* DAUDIN, Hist. Nat. Rept., VII, p. 383 (type-locality, Bengal; based on Russell's Ind. Serp., II, pl. ix).—BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 88 (Hainan).—*Hydrophis cyanocincta* HILGENDORFF, Mitth. Deutsch. Ges. Ost-Asiens, I, heft 10, July, 1876, p. 31.—*Dustira cyanocincta* BOULENGER, Fauna Brit. India, Rept., 1890, p. 410; Cat. Sn. Brit. Mus., III, 1896, p. 294 (Persian Gulf to Japan; Formosa).—WEST, Proc. Zool. Soc. Lond., 1895, p. 823, pl. LXVI, figs. 1, 8, 17 (poison gland; maxilla).—WALL, Proc. Zool. Soc. London, 1903, pp. 96, 101 (Hongkong).
1804. *Leoselasma striata* LACÉPÈDE, Ann. Mus. d'Hist. Nat., Paris, IV, p. 210, pl. LVII, fig. 1 (type-locality, "Nouvelle Hollande").—*Hydrophis striata* SCHLEGEL, Phys. Serp., II, 1837, p. 502, pl. XVIII, figs. 4-5 (Japan Seas); Fauna Japon., Rept., 1837, p. 89 (part), Oph. et Saur., pl. VII (Japan Seas).—OKADA, Cat. Vert. Japan, 1891, p. 69 (Okinawa Sea).
1817. *Hydrophis chittal* RAFINESQUE, Amer. Month. Mag., I, p. 432 (India; based on Russell).
1849. *Hydrophis subannulata* GRAY, Cat. Sn. Brit. Mus., p. 54 (type-locality, India).
1849. *Hydrophis aspera* GRAY, Cat. Sn. Brit. Mus., p. 55 (type-locality, Singapore; type in Brit. Mus.).
1859. *Hydrophis westermanni* JAN, Rev. Mag. Zool., 1859, p. —; author's separate, p. 26, pl. E (type-locality, Indian Ocean; type in Mus. Milano).
1863. *Hydrophis westermanni* JAN, Elenco Sist. Ofid., p. 111 (emendation); Icon. Ophid., livr. 39, 1872, pl. v, fig. 1 (type).
1870. *Hydrophis trachiceps* THEOBALD, Cat. Rept. (p. 70).
1871. *Hydrophis crassicollis* ANDERSON, Journ. Asiat. Soc. Bengal, XI, 1871, (p. 19).
1887. *Hydrophis taprobanica* ITALY, Taprobanian, II, (p. 107).
1887. *Hydrophis phipsoni* MURRAY, Journ. Bombay Nat. Hist. Soc., II, (p. 32, pl. —).

Description.—*Adult female*; U.S.N.M. No. 33931; Inatori, Idzu, Hondo, Japan; December 22, 1903; A. Owston collection (fig. 351). Rostral nearly as high as broad, the portion visible from above about as long as interprefrontal suture; nasals large, the suture between them twice as long as that between prefrontals; nostril in outer posterior corner of nasal; prefrontals not larger than supraoculars with which they are broadly in contact; frontal long, longer than its

^a Signifying blue-belted.

distance from rostral, but shorter than that from tip of snout, slightly broader than supraoculars; parietals longer than frontal, the interparietal suture equaling the distance of frontal from rostral; no loreal; one preocular; eye small, not larger than preocular; two postoculars; two anterior oblique temporals, the lower one being the upper cut-off portion of fifth (sixth on right side) supralabial; 7 supralabials on left side, 8 on right, second largest, fifth, sixth, and seventh (sixth, seventh, and eighth on right side), very low, subequal, third and fourth (third and fifth on right side, the interpolated fourth not reaching the orbit) entering eye; 3 lower labials in contact with anterior chin-shields which equal the posterior pair, the latter in contact throughout; 33 rows of scales on neck, 41 on middle of body, the latter with a bituberculate keel; 329 ventrals which are nearly smooth; four elongate, subequal anals; 38 subcaudals; tip of tail with a slightly enlarged scale. Color (in alcohol) pale yellowish, back suffused with gray, the skin between scales black; 53 dark gray crossbands, broadest and darkest, nearly black, on back, 5 on tail; top of head olive gray fading into a paler tint on labials and throat.




FIG. 351.—*DISTEIRA CYANOCINCTA*.
14 × TIMES NAT. SIZE. SIDE VIEW
OF HEAD. No. 33931, U.S.N.M.

Dimensions.

Head small, neck fully as wide, body very elongate, its greatest height not far from the vent, posteriorly considerably compressed.

	<i>mm.</i>
Total length.....	1, 290
Snout to vent.....	1, 180
Vent to tip of tail.....	110
Width of head.....	14
Diameter of neck.....	15
Greatest height of body.....	46
Greatest height of tail.....	24

The *male* is less elevated behind and the scales and ventrals strongly tuberculate.

Variation.—Of the head shields probably the supralabials are subject to the greatest variation, 7 being the normal, while the eighth may be interpolated at various points and thus make the formula appear more unstable than it really is. The arrangement of the anterior temporals is characteristic; the difference from that of *D. melanocephala* was pointed out under the latter species. The coloration is rather variable inasmuch as occasionally specimens have been found having the upper surface uniformly dark without cross-bands or rings, while on the other hand individuals having a black longitudinal band along the underside uniting the dark rings are common.

Habitat.—This is one of the most widely distributed and commonest of the sea snakes, extending as it does from the mouth of the Euphrates

in the Persian Gulf and Maskat in Arabia to Japan and, according to Peters, even to the Kingsmill Islands (Gilbert Archipelago) in the east.

Whether the specimen figured in Fauna Japonica (Plate VII) be the one collected by von Siebold or by Buerger (see p. 426 under *D. melanocepala*) there can be but little doubt that it was taken in Japanese or rather in Riu Kiuian waters. That it is rather common around Formosa seems proven by five specimens in British Museum collected by Swinhoe.

Stray specimens may occasionally drift far enough north to be taken in Japan proper as is shown by the specimen in the U. S. National Museum which was taken off Inatori, Idzu, on December 22, 1903.

List of specimens of *Disteira cyanocincta*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows on neck.	Teale rows on body.	Ventrals.	Subventrals.	Suprahyals.	Anterior temporals.
U.S.N.M.	33031	Female ^a	Inatori, Idzu, Hondo.	1903. Dec. 22	A. Owston	33	41	329	38	7-8	2
Brit. Mus.	Ag	Female ^b	Formosa		R. Swinhoe			310			
Do.	Ah	Young ^b	do.		do.			308			
Do.	Ai	do. ^b	do.		do.			305			
Do.	Ak	do. ^b	do.		do.			303			
Do.	Ch	Female ^c	do.		do.			330			

^a Description, p. 428.

^b Boulenger, Cat. III, p. 295.

^c Boulenger, Cat. III, p. 296.

DISTEIRA GODEFFROYI ^a (Peters).

1872. *Hydrophis godeffroyi* PETERS, Mon. Ber. Berlin Akad. Wiss., 1872, p. 856, pl. I, fig. 3 (type-locality, Kingsmill Islands; types in Berlin Mus.).—*Distira godeffroyi* BOULENGER, Cat. Snakes Brit. Mus., III, 1896, p. 291 (2 specimens Brit. Mus., loc. unknown).

1903. *Distira ornata* WALL, Proc. Zool. Soc. London, 1903, pp. 95, 101 ("Loo Choos") (not of Gray, 1842); 1905, II, p. 517 (Okinawa).

Four specimens from Ishigaki shima I refer to *D. godeffroyi* with considerable hesitation.^b They have the maxillary teeth following the poison fang distinctly grooved, and no doubt they are properly located among the species included by Boulenger in the genus *Disteira*. Doctor Wall, who examined them while yet in Mr. Owston's possession, identified them as *D. ornata* and recorded them under that name in

^a In honor of the German merchant, Johann Cesar Godeffroy (born in Kiel, July 1, 1813; died in Hamburg, Feb. 9, 1885) who in 1861 founded the Museum Godeffroy. The museum which was finally dispersed after his death was devoted mostly to Polynesian zoology and anthropology.

^b This hesitation does not refer to the specimens in British Museum from an unknown locality and named *D. godeffroyi* by Boulenger, but to the species as originally described by Peters from the Kingsmill Islands.

the Proceedings of the Zoological Society of London for 1903. They can not well be referred to that species because of the small number of scale rows, especially around the neck. Their scale formula agrees well with that of *D. major*, which seems to have the same distribution as *D. ornata*, but their rostral is much broader than high, as is the case in the latter and not as broad as high. They also agree in many respects with *D. cyanocincta*, which occurs as far north as Formosa at least, but the maximum of their ventrals is 246, while the minimum of ventrals in the latter species is said to be 281, and, save in exceptional cases, above 300. In Boulenger's synopsis of the species,^a *D. cyanocincta* on the one hand and *D. ornata*, *major*, and *godeffroyi* on the other are distinguished by the latter having the "second pair of chin-shields, if distinct, separated by several scales," while in the former they are said to be "in contact on the middle line" "or separated by a single scale," but evidently this distinction does not hold, at least not for the present species. Peters, in the original description of *D. godeffroyi*, expressly states that in the larger specimen of the types both pairs of chin-shields are in contact, while in the smaller one the posterior pair is separated by two scales which are pointed anteriorly. In our four specimens scarcely two are exactly alike, though there can not be the faintest doubt that they all belong to the same species. One (No. 33934) has the two posterior chin-shields in contact; in two (Nos. 33933, 33935) they are separated by one, and to some extent also by a second scale, while in the fourth (No. 33936) they are separated by two scales exactly shaped as described by Peters.

The distance from the type locality, the Kingsmill, or Gilbert Islands, is certainly considerable, but so little has been collected in the intermediate region that no adverse argument can be derived from the remoteness of the Riu Kiu Islands.

In one sense our specimens are intermediate between all the species mentioned. This has already been demonstrated with regard to the relations of the posterior chin-shields. In the number of scales around the neck (viz, 30-34) they assume a position between *D. cyanocincta*, with 27-33, and *D. ornata*, with 35-42. In scales around the body (33-43) they agree better with *D. cyanocincta* (39-43) than with *D. ornata* (40-50), while in the number of ventrals (227-246) they come within the limits of the latter (210-300) and not within the former (281-426).^b It is just possible that the case is similar to the one pointed out under *D. fasciata* and *melanocephala* (p. 421) and that the limits of the three species are not properly defined.


It may be that the distinction is not so much one of scale formula as of other characters. Thus, according to Boulenger, *D. major*,

^a Cat. Sn. Brit. Mus., III, 1896, pp. 286-287.

^b The type specimens of *D. godeffroyi* are said to have 294 and 284 ventrals, respectively, consequently more than the exceptional minimum of a few *D. cyanocincta*.

with which our species agrees best in scale formula, has the rostral "as deep as broad," and only one pair of chin-shields, while ours has the rostral much broader than high and the second pair of chin-shields well developed. In *D. ornata* this feature is apparently not so marked, for Boulenger speaks of the "posterior chin-shields, if distinct, separated by two or three scales." In *D. cyanocincta* the rostral is but "slightly broader than deep," and the scales "subimbricate,"^a while in our specimens the difference between breadth and depth of the rostral is considerable and the scales can not be termed subimbricate on the posterior part of the body at least. I do not know whether the size of the terminal scale of the tail is of any consequence. Boulenger does not mention it in his descriptions, but in Jan's figure^b which Boulenger cites under *D. ornata* it is quite large, while in our specimens no such enlarged terminal scale is differentiated. Peters expressly mentions that in *D. godeffroyi* the terminal tail scales are small.

I may add that our specimens agree on the whole very well with Peters's figures, except that the first supralabial is considerably higher



FIGS. 352-354.—*DISTIEIRA GODEFFROYI*. $1\frac{1}{4} \times$ NAT. SIZE. 352, TOP OF HEAD; 353, SIDE OF HEAD; 354, UNDERSIDE OF HEAD. NO. 33933, U.S.N.M.

than shown in his fig. 3. The anal shield, or scales (fig. 3d), agree exactly with the arrangement in our female (No. 33933), while in the three males there are four rows of small scales between the lateral large scutes instead of two. This arrangement of the preanals, together with the greater relative shortness of the body, seems to be among the best characters of this species.

Description.—*Adult female*; U.S.N.M. No. 33933; Ishigaki shima, Yaeyama group, Riu Kiu archipelago; April to May, 1899; A. Owston collection (figs. 352-354). Rostral much broader than high, well visible from above; nasals long, with the crescentic valvular nostril close to the posterior edge; internasal suture as long as interparietal suture and three times as long as the one between the prefrontals:

^a "Scales all imbricate" in key, Cat. Sn. Brit. Mus., III, 1896, p. 287.

^b Icon. Ophid., livr. 40, pl. vi, fig. 1.

frontal as long as its distance from rostral, once and a half as long as wide, shorter than parietals; parietals as long as frontal and interprefrontal suture together; no loreal; one preocular; two postoculars; 8 supraoculars, seventh and eighth very small and low, fifth a low triangular piece separated off from sixth, second in contact with prefrontal, very high and large; 3 lower labials in contact with anterior pair of chin-shields which are larger than posterior, the latter separated by two scales; 34 rows of slightly tuberculated scales on neck, 43 on middle of body, the tubercles on the latter more strongly developed; 246 bituberculated ventrals; preanal region covered with scales, four of which form the anterior edge of the vent, the outer ones on each side slightly longer than the others; 39 subcaudals; terminal scale of tail but slightly enlarged. Color (in alcohol) buffy white with a tinge of gray on the back and throat; 16 slaty-gray crossbars on neck and body and 8 on tail, broadest on middle of back, where broader than the light interspaces, gradually narrowing to the middle of the sides whence they extend as narrow, paler-colored bands to the abdomen, which many of them cross; head nearly uniformly isabella-colored; the last three bands on the tail blackish.

Dimensions.

Head rather large; neck thick; body short, high, and very compressed.

	<i>mm.</i>
Total length.....	790
Snout to vent.....	690
Vent to tip of tail.....	100
Width of head.....	19
Diameter of neck.....	19
Greatest height of body.....	51
Greatest height of tail.....	27

The *males* are much slenderer posteriorly, the greatest height of body being less than twice as high as the neck.

Variation.—Under the preliminary discussion of this species (p. 431) reference is made to the variation of chin-shields and number of ventrals in this species. The supralabials are also subject to considerable variation, chiefly caused by the irregular breaking up of the posterior shields and the consequent uneven arrangement of the fragments edging the lip. This irregularity in the posterior supralabials also affects the adjacent temporals inasmuch as a shield becomes a labial or a temporal according to whether the lower portion of it is cut off so entirely as to exclude the upper portion from the edge of the lip, or not.

Habitat.—The distribution of *D. godeffroyi*, as here understood, is a curiously disconnected one, inasmuch as the only specimens referred to it are known from the Kingsmill Islands in Polynesia, and from

Ishigaki shima in the Riu Kiu Archipelago, while the other two specimens in British Museum have no locality attached to them. Doctor Wall now records three from Okinawa.

List of specimens of Distira godffroyi.

U.S.N.M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Scale rows on neck.	Scale rows on body.	Ventrals.	Subcaudals.	Supraabials.	Anterior temporals.
			1889.							
33933	Female a.	Ishigaki I., Riu Kiu....	April-May.	A. Owston....	34	43	246	39	7-8	1
33934	Male.....	do.....	do.....	do.....	30	35	228	45	7-8	2
33935	do.....	do.....	do.....	do.....	32	33	235	44	7	2
33936	do.....	do.....	do.....	do.....	33	33	227	42	8	2-1

a Description, p. 432; figs. 352-354.


[**DISTEIRA VIPERINA** a (Schmidt).]

1852. *Thalassophis viperina* SCHMIDT, Abh. Naturw. Ver. Hamburg, II, Pt. 2, p. 79, pl. III (type-locality, Java; type in Hamburg Mus.).—*Hydrophis* (*Thalassophis*) *viperinus* BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 91 (Hainan).—*Distira viperina* BOULENGER, Fauna Brit. India. Rept., 1890, p. 413; Cat. Sn. Brit. Mus., III, 1896, p. 298 (Maskat to Hongkong).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 114 (Hainan).
1854. *Distira præscutata* DUMÉRIL and BIBRON, Érpét. Gén., VII, Pt. 2, p. 1331 (type-locality unknown; type in Paris Mus.).
1856. *Hydrophis doliata* FISCHER, Abh. Naturw. Ver. Hamburg, III, p. 56 (Schmidt's type of *T. viperina*; not of Lacépède, 1804).
1887. *Hydrophis jayakari* BOULENGER, Ann. Mag. Nat. Hist. (5), XX, p. 408 (type-locality, Maskat, Arabia; type in Brit. Mus.; Doctor Jayakar, collector).
1887. *Hydrophis plumbea* MURRAY, Journ. Bombay Nat. Hist. Soc., II (p. 34).

No specimen of this species being at hand, I submit the diagnosis given by Boulenger in the Catalogue of Snakes.

Diagnosis.—Head rather small; body moderately elongate, rather slender anteriorly; rostral as deep as broad; nasals as long as or a little shorter than the frontal; suture between the prefrontals very short; frontal as broad as long, as long as its distance from the rostral or end of the snout, shorter than the parietals; one or two pre- and one or two postoculars; 7 or 8 upper labials, second largest, fourth or third and fourth entering the eye; one or two anterior temporals; two pairs of chin-shields, in contact with each other, posterior longest; 27 to 29 scales round the neck, 37 to 43 round the body; scales juxtaposed, obtusely keeled; ventrals relatively large anteriorly, bituberculate in the male, 235-267. Slaty gray above, with rhomboidal transverse black spots, white or pink on the sides and inferiorly; some specimens uniform dark gray above; in the young, the black spots are continued down the sides of the body; end of tail black.

a Signifying viper-like.



LAPEMIS HARDWICKII. (FROM JAN.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

Dimensions.

	mm.
Total length.....	920
Snout to vent.....	820
Vent to tip of tail.....	100

Habitat.—Distributed from Arabia in the west to China in the east, though apparently rather rare.

British Museum has one specimen from Hongkong, and Herz collected one at Hainan, which is now in the Senckenberg Museum.

It has not yet been recorded from any locality within our territory, but it seems highly probable that it occurs around Formosa at least occasionally.

[Genus **LAPEMIS**^a Gray.]

1820. *Enhydris* MERREM, Syst. Amph., p. 140 (part; not of Latreille, 1802).
 1834. *Lapemis* GRAY, Ill. Ind. Zool., II (pl. LXXXVII, fig. 2) (type, *L. hardwickii*).
 1861. *Pelamidoïdis* FITZINGER, Sitz. Ber. Akad. Wiss. Wien, Math.-Nat. Cl., XLII, p. 409 (type, *Hydrophis pelamidoïdis* Schlegel).

Apart from the fact that Merrem's *Enhydris* of 1820 is preoccupied by Latreille for an entirely different kind of snake, that name can not be used for a genus with *Hydrus curtus* for type, as restricted by Boulenger in 1890, because Wagler, as early as 1830, restricted it to *H. cyanocinctus*, one of the species included by Merrem in his *Enhydris*.

[**LAPEMIS HARDWICKII**^b Gray.]

Plate XXIV.

1834. *Lapemis hardwickii* GRAY, Ill. Ind. Zool., II (pl. LXXXVII, fig. 2) (type-locality, India; type in Brit. Mus.; Hardwicke, collector).—*Hydrophis hardwickii* GUENTHER, Rept. Brit. India, 1864, p. 380, pl. xxv, fig. w (type-locality, Pinang?).—MUELLER, Verh. Naturf. Ges. Basel, VII, Pt. 1, 1882, p. 170; author's separate (p. 7) (China Sea).—*Enhydris hardwickii* BOULENGER, Fauna Brit. India, Rept., 1890, p. 397; Cat. Sn. Brit. Mus., III, 1896, p. 301 (Bay of Bengal to Chinese Sea and New Guinea).—WALL, Proc. Zool. Soc. London, 1903, p. 96 (Manila; Bangkok).
 1837. *Hydrophis pelamidoïdes* SCHLEGEL, Phys. Serp., I, p. 187; II, p. 512 (Bay of Bengal, and seas of China and of the Moluccas); Atlas, pl. xviii, figs. 16–17; Fauna Jap., Rept., 1837, p. 91; Ophid., pl. ix (seas south of Japan).—JAN, Icon. Ophid., livr. 41, 1872, pl. iii, fig. 1c (Indian Ocean).—BOETTGER, Offenbach. Ver. Naturk. 17–18 Ber., 1878, p. 8 (Japan).—OKADA, Cat. Vert. Japan, 1891, p. 69 (Okinawa Sea).
 1843. *Lapemis loreatus* GRAY, Ann. Mag. Nat. Hist., XI (p. 46) (type-locality, Borneo; type in Brit. Mus.; Belcher, collector).
 1856. *Hydrophis (Pelamis) pelamidoïdes* var. *annulata* FISCHER, Abh. Naturw. Ver. Hamburg, III, p. 67, pl. iii (Java; Hamburg Mus.).
 1863. *Hydrophis abbreviatus* JAN, Elenco Sist. Ofid., p. 109 (type-locality, Manila; type in Essex Inst., Salem, Mass.); Icon. Ophid., livr. 40, 1872, pl. iv, fig. 2.
 1863. *Hydrophis brevis* JAN, Elenco Sist. Ofid., p. 109 (type-locality, Manila; type in Mus. Milano); Icon. Ophid., livr. 40, 1872, pl. v, fig. 2.

^a Anagram of *Pelamis*.

^b For Gen. Thomas Hardwicke, who collected the type.

^c Reproduced in this work on Plate XXIV.

1871. *Hydrophis fayerriana* ANDERSON, Journ. Asiat. Soc. Bengal, XI, Pt. 2, 1871 (p. 19).
 1876. *Hydrophis pelamoides* HUGENDORF, Mitth. Deutsch. Ges. Ost-Asiens, I, heft 10, p. 31 (*lapsus*).
 1885. *Hydrophis hardwicki* BOETTGER, Offenbach. Ver. Naturk. 24-25 Ber., p. 155 (emendation).
 1888. *Hydrophis hardwicki* BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., p. 150 (emendation).

Jan's *Hydrophis problematicus* (Rev. Mag. Zool., 1859, author's separate, p. 25) with 27 scale rows on the highest part of the body and 230 ventrals, can not well be this species. It is said to be from Manila.

There being no specimen from the waters adjacent to our territory available for description a modified copy of Boulenger's diagnosis in his Catalogue of Snakes is deemed sufficient for identification.


Diagnosis.—Rostral as deep as broad or slightly broader than deep; parietals large; one pre- and one to three postoculars; a loreal sometimes present; two or three superposed anterior temporals; 7 supralabials, fourth or third and fourth entering eye; chin-shields very small and separated by scales, or quite indistinct; 25 to 33 scales round the anterior part of the body, 34 to 37 round the middle, with tubercles [very feebly developed or a short keel]; ventrals usually very indistinct, 130-200. Coloration [above with dark transverse bands, broadest in the middle; these bands] often forming complete rings round the body.

Dimensions.

	<i>mm.</i>
Total length.....	750
Snout to vent.....	670
Vent to tip of tail.....	80

Habitat.—The claim of *Lapemis hardwickii* to a place in the Japanese fauna is doubtful. Schlegel enumerates and figures the species in Fauna Japonica, as *Hydrophis pelamidoïdes*, but the only reason given in the text is as follows: "A specimen of this curious reptile, collected in the seas south of Japan, has recently been brought to Holland. We hesitate the less in admitting this *Hydrophis* among the number of animals belonging to the fauna of Japan, inasmuch as it furnishes us an opportunity for publishing an exact figure of this species." The implication is that the specimen was not actually taken in Japanese waters, and so far as I know no later capture is on record. However, Mueller reports recently a specimen in the Basel Museum from the China Sea, and as the species appears to be common on the west side of Luzon, there is every chance that it may eventually be taken within our limits.

To the west the species is known in the Bay of Bengal and in the south to the coasts of New Guinea and Java.



1

ENHYDRINA VALAKADYN. (FROM JAN.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

[Genus ENHYDRINA^a Gray.]

1834. *Polyodontes* LESSON, in Bélanger's Voy. Indes Orient., Rept. (p. 325); Atlas, Rept., pl. iv (type, *P. annulatus*) (not *Polyodontes* Montfort, 1810, nor *Polyodonta* Megerle, 1812).
1849. *Enhydrina* GRAY, Cat. Sn. Brit. Mus., p. 47 (type, *Hydrus valakadyen*).

[ENHYDRINA VALAKADYN^b (Boie).]

Plate XXV.

1827. *Hydrus valakadyen* BOIE, Isis, 1827, p. 554 (type-locality, Tranquebar, India).
1827. *Distera russelii* FITZINGER, Isis, 1827, p. 733 (based on Russell's Ind. Serp., II, pl. xi).
1834. *Polyodontes annulatus* LESSON, in Bélanger's Voy. Indes Orient., Rept. (p. 325); Atlas, Rept. pl. iv (type-locality, ———).
1837. *Hydrophis schistosa* SCHLEGEL, Phys. Serp., II, p. 500; Atlas, pl. xviii, figs. 1-3 (not of Daudin, 1803).—*Hydrophis schistosus* JAN, Icon. Ophid., livr. 41, 1872, pl. ii, fig. 1 c ("Japan;" Mus. Milano).
1842. *Hydrophis bengalensis* GRAY, Zool. Miscell., p. 62 (type-locality, Bengal; type in Brit. Mus.; Hardwicke, collector).
1842. *Hydrophis subfasciata* GRAY, Zool. Miscell., p. 62 (type-locality, Bengal; type in Brit. Mus.).
1849. *Enhydrina valakadyen* GRAY, Cat. Sn. Brit. Mus., p. 48 (emendation).
1872. *Hydrophis fasciatus* JAN, Icon. Ophid., livr. 41, pl. iii, fig. 2 (Bay of Bengal) (not of Schneider).
1890. *Enhydrina valakadien* BOULENGER, Fauna Brit. India, Rept., p. 406 (emendation); Cat. Sn. Brit. Mus., III, 1896, p. 302 (Persian Gulf to Papuasias).
1892. *Enhydrina vikadien* BOETTGER, Offenbach. Ver. Naturk. 29-32 Ber., p. 89 (*lapsus*) (Madras).
1899. *Enhydrina velakadien* FLOWER, Proc. Zool. Soc. London, 1899, p. 688 (*lapsus*) (Siam).

This is possibly the snake which von Siebold meant when mentioning "*Hydrophis flaviventris* mihi," referring as he does to Bosc's "*Hydrophis ardoisé*" (Nouv. Diet. d'Hist., Nat., XV, p. 491). The reference is found in a scarce pamphlet published by von Siebold in Batavia in 1824 under the title "De historiae naturalis in Japonia statu," etc. A reprint in pamphlet form was issued in Wuerzburg by some of his friends in 1826, and reference is also made to it in Oken's Isis for 1827. From these latter two issues has been taken the above reference, which is found on page 18 and page 142, respectively, the heading under which it occurs being *Spicilegia Faunae Japonicae*.

There is no specimen in our museum from Chinese or Japanese waters, and a detailed description of a specimen from a distant locality is not deemed satisfactory. For the purposes of identification a transcript of Boulenger's diagnosis in his Catalogue of Snakes is given.

^a Signifying like *Enhydris*, the generic name given by Latreille to another water snake.

^b *Valakadyen* is the native name according to Russell, its first describer.

^c Reproduced in this work on Plate XXV.

Diagnosis.—Rostral deeper than broad; frontal small, longer than broad, shorter than its distance from the end of the snout or than the parietals; one pre- and one or two postoculars; 7 or 8 supralabials, fourth or third and fourth entering the eye; usually a single anterior temporal; [symphysial shield narrow, partly concealed in a deep groove in the chin]; chin-shields small or indistinct; 40 to 60 scales round the neck, 50 to 70 round the body; scales feebly imbricate, with a small tubercle or short keel, which is stronger in the males; ventrals very slightly enlarged, 230–314. Young olive or gray above, with black transverse bands, broadest in the middle, and tapering to a point on the sides; in the adult these bands are usually less distinct, some specimens being uniform dark gray above; sides and lower parts whitish.

Dimensions.

	<i>mm.</i>
Total length.....	1.300
Snout to vent.....	1.110
Vent to tip of tail.....	190

Habitat.—The Valakadyen is introduced here mainly on the strength of a specimen recorded by Jan as being in the Milano Museum with the locality "Japan." This is ordinarily a very doubtful authority, and the chances are that the specimen was not taken in Japanese waters. Yet, with the possibility that it may be the species which von Siebold referred to in 1824 (see p. 437), I have thought it best to include a brief characteristic, as it may occasionally, at least, drift as far as Formosa, or farther, notwithstanding the fact that it has not been reported by others north of Cochin China.

This species is known from Arabia in the west to the Malay Archipelago, Papuasia, and Siam in the east. It is said to be "incredibly numerous" in the Bay of Bengal, at Pinang and Singapore.

Genus HYDRUS^a Schneider.

- 1799. *Hydrus* SCHNEIDER. Hist. Amph., I, p. 233 (type, *H. bicolor*).
- 1802. *Hydrophis* LATREILLE, Hist. Nat. Rept., IV, p. 193 (substitute for *Hydrus* Schneider).
- 1803. *Pelamis* DAUDIN, Hist. Nat. Rept., VII, p. 357 (substitute for *Hydrophis* Latreille).
- 1830. *Pelamys* WAGLER, Syst. Amph., p. 166 (emendation).
- 1848. *Elaphrodites* GISTEL, Naturg. Thierr., p. ix (substitute for *Hydrus*).

The type of *Hydrus*, as ascertained by the process of elimination, is *H. platurus*. Latreille changed the name *Hydrus* into *Hydrophis* simply because the French name "hydre" was preoccupied for another animal, the *Hydra*. It is, therefore, a synonym pure and simple and must have the same type as Schneider's *Hydrus*. Daudin's *Pelamis* of the following year has precisely the same status,

^a From ὕδρος, a water snake.

as he invented another name for no other reason than "elles habitent dans la mer." *Hydrophis* and *Pelamis* are consequently unavailable as names for other genera.


HYDRUS PLATURUS ^a (Linnæus).

1766. *Anguis platura* LINNÆUS, Syst. Nat., 12 ed., I, p. 391 (type-locality unknown; type originally in Fr. Ziervogel's pharmacy, Stockholm).—*Hydrophis platura* LATREILLE, Hist. Nat. Rept., IV, 1802, p. 197 (islands of the Pacific Ocean).—*Hydrus platurus* BOULENGER, Fauna Brit. India, Rept., 1890, p. 397; Cat. Sn. Brit. Mus., III, 1896, p. 267 (Indian and Pacific oceans: Ooshima, Loo Choo Is.; Formosa).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 112.—STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 223 (Kilung, Formosa; Pescadores Is.).—WALL, Proc. Zool. Soc. London, 1903, pp. 95, 101 (Hongkong; Formosa; Loo Choos); 1905, II, p. 516 (Okinose, Sagami).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 291 (Possiet Bay).
1788. *Anguis platurus* GMELIN, Syst. Nat., I, Pt. 3, p. 1122 (Pine Isle, Pacific Ocean; Forster, collector) (emendation).
1799. *Hydrus bicolor* SCHNEIDER, Hist. Amph., I, p. 242 (type-locality not specified; specimen in Mus. Blochianum).—*Pelamis bicolor* DAUDIN, Hist. Nat. Rept., VII, 1803, p. 366 (Bengal).—SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 225 (Kelung, Tamsui River, Formosa).—STRAUCH, Mém. Acad. Sci. Pétersb. (7), XXI, no. 4, 1873, p. 199 (Possiet Bay, Siberian Coast Province).—HILGENDORF, Mitth. Deutsch. Ges. Ost-Asiens, I, heft 10, July, 1876, p. 30 (Prov. Kaga, Hondo; Setanai, Prov. Kunashiri, Yezo).—MARTENS, Preuss. Exped. Ost-Asien, Zool., I, 1876, p. 382 (Strait of Formosa).—DOEDERLEIN, Mitth. Deutsch. Ges. Ost-Asiens, III, heft 25, Dec. 1881, p. 210 (west coast of Hondo and Yezo).—BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 91 (Hongkong).—*Hydrophis (Pelamis) bicolor* FISCHER, Abh. Naturw. Ver. Hamburg, III, 1856, p. 61 (Indian Ocean; west coast Mexico).—*Hydrophis bicolor* JAN, Icon. Ophid., livr. 40, 1872, pl. II, fig. 4; pl. III, fig. 2 (China Sea).—OKADA, Cat. Vert. Japan, 1891, p. 69 (Noto; Hoki; Totomi; Idzumo).
1817. *Pelamis schneideri* RAFINESQUE, Amer. Month. Mag., I, p. 432 (substitute name for *Hydrus bicolor* Schneider).
1837. *Hydrophis pelamis* SCHLEGEL, Phys. Serp., I, p. 187; II, p. 508; Atlas, pl. XVIII, figs. 13-15 (substitute name); Fauna Jap., Rept., 1837, p. 90 (voyage between Java and Japan, north to 27° N. lat.).—BOETTGER, Offenbach. Ver. Naturk. 17-18 Ber., 1878, p. 8 (Japan).
1837. *Hydrophis pelamis* var. SCHLEGEL, Fauna Jap., Rept., Ophid., pl. VIII (Borneo).
1842. *Pelamis ornata* GRAY, Zool. Miscell., p. 60 (type-locality, India; type in Brit. Mus.).
1854. *Pelamis bicolor* var. *variegata* DUMÉRIL and BIBRON, Erpét. Gén., VII, Pt. 2, p. 1337 (Macassar, Celebes).—JAN, Icon. Ophid., livr. 40, 1872, pl. III, fig. 3.
1854. *Pelamis bicolor* var. *sinuata* DUMÉRIL and BIBRON, Erpét. Gén., VII, Pt. 2, p. 1338 (locality unknown).
1856. *Hydrophis (Pelamis) bicolor* var. *alternans* FISCHER, Abh. Naturw. Ver. Hamburg, III, 1856, p. 63 (substitute for var. *variegata*).
1863. *Hydrophis bicolor* var. *maculata* JAN, Eleuco Sist. Ofid., p. 109 (Indian Ocean); Icon. Ophid., livr. 40, 1872, pl. III, fig. 1.

^a From πλάτυς, flat; ὄψα, tail.

There are several well-marked color varieties which have received varietal names. Their exact status has not been made clear, but while it is true that they do not seem to represent local races, there is nevertheless a certain regularity in their distribution. Thus the ordinary black-backed form seems to be almost exclusive in the Pacific Ocean; spotted individuals, such as the one figured in *Fauna Japonica*, plate VIII (from the west coast of Borneo) appear to occur mostly in the Indian Ocean. All the specimens from eastern China, Formosa, Riu Kiu, and Japan, so far as the records show, are of the black-backed form.

Description (figs. 355-357).—*Adult female*; U.S.N.M. No. 17518; harbor of Nagasaki, Kiusiu; December 15, 1883; P. L. Jouy, collector. Head very depressed; snout elongate, the distance from eye to tip of snout equaling the width of head at posterior end of supra-



FIGS. 355-357.—HYDRUS PLATURUS. $1\frac{1}{2}$ \times NAT. SIZE. 355, TOP OF HEAD; 356, SIDE OF HEAD; 357, UNDERSIDE OF HEAD. NO. 8, SCI. COLL. TOKYO.

oculars; rostral nearly as high as broad, well visible from above; nasals not much longer than prefrontals, the suture between the former about twice as long as between the latter; frontal much longer than its distance from tip of snout and nearly as long as parietals, much wider than supraoculars; parietals long, with irregular outlines, a small intercalated shield in the suture between them; no loreal; a long preocular, pointed anteriorly; two suboculars; two postoculars; three anterior temporals; ten supralabials, second largest, in contact with prefrontals, fourth to tenth small and low, none entering eye; no differentiated chin-shields; 61 scale rows around middle of body, the scales hexagonal, juxtaposed, smooth; no differentiated ventrals or subcaudals; two slightly enlarged scales, one at each side in front of vent; terminal scale on tail somewhat enlarged. Color (in alcohol) on back dark seal-brown, sides and underparts abruptly clay-colored, the

boundary between the two colors being a straight line until a short distance in front of the vent where it becomes wavy; top of head like back, supralabials like underside, lower labials washed with brown; tail straw-yellow, with a series of about six dark brown roundish blotches along the upper and the lower edge and a median series of smaller ones on the sides.

Dimensions.

	<i>mm.</i>
Total length.....	609
Snout to vent.....	528
Vent to tip of tail	81
Width of head across center of eyes.....	14
Length of shielded portion of head.....	31
Greatest height of body.....	38
Greatest height of tail.....	20

The *males* have a somewhat slenderer body, fewer scale rows, and the scales have a central tubercle.

Variation.—This species is exceedingly variable; especially the shields of the sides of the head are subject to so numerous modifications that it is hardly profitable to enumerate them in view of the striking appearance of this snake, which makes the identification of the species one of the easiest. It may be mentioned, however, that the supralabials may vary between 7 and 11; one or more loreals may be present; suboculars may be present or not; two anterior chin-shields may be sometimes recognized in a pair of enlarged scales never in contact, however. In our specimens the number of scale rows varies between 49 and 61, the higher figures belonging to the females, while Boulenger gives 45 to 47 as the normal number.

The coloration is also very variable, and mention has already been made of the spotted form from the Indian Ocean (p. 440). In the black-backed form which is the only one found in the eastern waters, the tail is most subject to variation, inasmuch as the spots on the upper and lower edges may join so as to form blackish cross-bands, or they may disappear on the lower edge entirely. In many specimens there may be seen on the sides between the blackish color of the back and the more or less dirty clay-color of the belly a broad yellow band.

Habitat.—Probably the most widely distributed species of snake, occurring as it does on all the coasts and islands of the Indian and Pacific oceans within the Tropics and even extending its range in many places considerably beyond the tropical waters, thus in the southern hemisphere to the Cape of Good Hope and to New Zealand. In the northern hemisphere it has been found on the eastern coasts of Asia as far north as Yezo and the Russian Coast Province

Beyond the tropics, however, these snakes are only carried occasionally by the warm currents. Thus while they are yet of regular occurrence around Formosa, they become gradually scarcer in the

Riu Kius, although stray individuals are picked up rather frequently on the coasts of Japan proper. A Formosan specimen is in British Museum collected by Swinhoe, and Mr. Tada also obtained it in that island and recorded it from the Pescadores Islands; British Museum furthermore has a specimen collected by Mr. Ferrié in Amami-o-shima, Riu Kiu, and Doctor Wall saw two or three more specimens in Mr. Owston's collection from the shores of the Riu Kius. Authentic records for Japan proper show specimens obtained on the eastern and the western shores. Thus Jouy collected one in the harbor of Nagasaki on December 15, 1883; Okada mentions it as having been found in the provinces of Idzumo, Hoki, and Noto, and Hilgendorf records a specimen from the province of Kaga, all in Hondo, in the Sea of Japan; the latter author also mentions having seen a specimen from "Setanai (Prov. Kunasiri)"^a in the exhibition of the government of Hokkaido. Doederlein mentions another specimen from Yezo, probably also from the west coast.

From the east coast we have in the U. S. National Museum two specimens from the Bay of Sagami, one from the Bay of Suruga, and one from Inatori, province of Idzu, opposite O-shima; Doctor Wall mentions another specimen from Sagami Bay, and Okada records the species from Omayá Saki, province of Totomi.

On the mainland side of the sea of Japan, Dr. B. Dybowski collected a specimen in Possiet Bay, near the Korean boundary, the specimen, according to Strauch, being in the museum in Warsaw.

List of specimens of Hydrus platurus.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.
U.S.N.M.	17518	Female a.	Nagasaki, Kiusiu	Dec. 15, 1883	P. L. Jouy	61
Do.	33932	Male	Inatori, Idzu, Hondo	Dec. 22, 1903	A. Owston	49
Do.	33964	do	Sagami Bay, Hondo	Apr. --, 1902	do	52
Do.	33965	Female	do	do	do	57
Do.	33966	Male	Suruga Bay, Hondo	Jan. 12, 1904	do	53
Sci. Coll. Tokyo.	8	do. b.	Kilung, Formosa	Nov. --, 1897	T. Tada	50

^a Description p. 440.

^b Figs. 355-357.

Family COBRIDÆ.

As will be seen from the explanation under the generic name of the only species of this family treated of in this work, the term *Vipera*, from which the family name *Viperidæ* is derived, must be superseded by *Coluber*. The confusion necessarily resulting from this transfer of

^a Kunashiri is an island off the northeast extremity of Yezo, but I have been unable to find on my maps any locality of that name on Yezo itself. There is, however, a Setanai in the province of Shiribeshi, on the west coast of Yezo (about 42° 30' N. lat.) which is possibly the place meant.

the latter name would be still more far-reaching were we compelled to derive the new family name from it.^a Under these circumstances it seems preferable to give a new family name derived from the generic term of the African vipers nearest related to the genus *Coluber*. This genus is Laurenti's *Cobra*, which is equivalent to *Bitis* given much later. To have adopted the family name *Causidæ* and giving it a much wider significance than intended by its originator would also lead to confusion.

The chief characteristic of this group is the shortness of the maxillaries on which the large poison fangs are placed immovably. The maxillary bone is so connected with the other bones of the skull as to be readily movable in a vertical direction so that the fangs can be raised or folded back at will. This apparatus is essentially the same as that of the Crotalids and will be described more in detail under the latter, the chief difference being that in the vipers the maxillary is not scooped out, while in the Crotalids it has a deep hollow for the loreal pit.

The vipers are confined to the Old World and are found mostly in Africa. A number of forms also inhabit the palearctic region, one species thus coming within our province, while only a single viper inhabits the Indian region proper, and none the Malay archipelago or Australia.

Genus COLUBER ^b Linnæus.

1758. *Coluber* LINNÆUS, Syst. Nat., 10 ed., I, p. 216 (type, *C. berus*).
 1768. *Vipera* LAURENTI, Syn. Rept., p. 99 (type, *V. illyrica*).
 1820. *Pelias* MERREM, Syst. Amph., p. 148 (type, *Coluber berus*).
 1820. *Berus* GOLDFUSZ, Handb. Zool., II, p. 144 (type, *Vipera berus*; not of Oken, 1816).
 1822. *Chersea* FLEMING, Philos. Zool., II, p. 295 (type, *C. vulgaris*=*Coluber berus*).

The name *Coluber* was instituted by Linnæus in 1758 for the great majority of snakes and the "genus" so designated embraced poisonous as well as nonpoisonous snakes. The term since then has been applied to widely different groups of serpents, and there is hardly a name in ophiology which has been more in dispute, or attributed to more kinds of snakes than that of *Coluber*. It has been restricted so as to fall entirely within the *Pythonidæ*, the *Viperidæ*, or the aglyphodont Natricoids. All sorts of schemes for ascertaining the type to which the name might be fastened have been tried,^c but without satisfactory results, and even the same author has at various times applied it to widely different genera.

^aAccording to the International Code, art. 5, "the name of a family or subfamily is to be changed when the name of its type genus is changed."

^bGeneral Latin term for snake.

^cSee, for instance, Cope, Proc. U. S. Nat. Mus., XI, p. 389.

In order to settle the question according to the International Code of Zoological Nomenclature we must turn to its article 30, which reads as follows:

If the original type of a genus was not indicated, the author who first subdivides the genus may apply the name of the original genus to such restricted genus or subgenus as may be judged advisable, and such assignment is not subject to subsequent change. In no case, however, can the name of the original genus be transferred to a group containing none of the species originally included in the genus; nor can a species be selected as type which was not originally included in the genus, or which the author of the generic name doubtfully referred to it.

Linnaeus, of course, did not indicate a type, and "the author who first subdivides the genus may apply the name of the original genus to such restricted genus or subgenus as may be judged advisable, and such assignment is not subject to subsequent change." The first one to do this is unquestionably Laurenti, in 1768. He dismembered the enormous genus *Coluber* of Linnaeus and left comparatively few species in it. Five of these are additional to those included by Linnaeus in his original *Coluber* of 1758, and are consequently ineligible as types, because a species can not "be selected as type which was not originally included in the genus." The type must of necessity, therefore, be selected from among the remaining Linnaean species, viz, *Coluber stolatus*, *buccatus*, *sibon*, *chersea*, and *berus*, the two last-mentioned being in reality only one species. It now so happens that only one of these has been designated by a subsequent writer as type of the genus *Coluber*, viz, *C. berus*, which was so designated by Prof. Robert Collett, in 1878,^a and in the words of article 30 "such assignment is not subject to subsequent change."^b The result is fully consonant with the "recommendation" accompanying article 30, more particularly that part of paragraph C, which provides that "if the genus contains both exotic and nonexotic species, from the standpoint of the original author, the type is to be selected from the nonexotic species," inasmuch as *C. berus* is the only nonexotic species among the eligible species. That this result also agrees with Laurenti's "intention" has been well shown by Cope,^c who expressly remarks that "the poisonous species are, then, the types of the *Coluber* of Laurenti."

In a large collection of Russian and Siberian reptiles received by the United States National Museum in 1885 from Dr. Alexander Strauch there are three specimens named *Vipera berus* which in the invoice are marked thus: No. 32, Dui auf Sachalin; No. 84, Smei-nogorsk (Altai); No. 88?, Dui auf Sachalin.

^a Christiania Vidensk. Selsk. Forhandl., 1878, no. 3, p. 6.

^b It might even be maintained that the same selection was made as early as 1820 by Merrem (Syst. Amphib., p. 148). He subdivided also the genus *Coluber*, but unfortunately he did not retain the name for any of the subdivisions. But in instituting the genus *Pelias* for *Coluber berus* he expressly mentions Laurenti's *Coluber* as the equivalent.

^c Proc. U. S. Nat. Mus., XI, p. 389.

The original parchment tags with the numbers are still attached to the specimens which represent two species as follows: No. 32 (U.S.N.M. No. 14325) *Coluber berus*; No. 84 (14324), *Coluber berus*; and No. 88 (14318), *Coluber renardi*. It should be noted that the question mark following 88 in the invoice is not found on the parchment tag; also that there is another No. 88, viz, a *Natrix hydrus*, from Mangyshlak, Transcaspia (U.S.N.M., No. 14327). This question mark in the invoice throws some doubt upon the accuracy of the whole entry rendering the locality Dui on the island of Sakhalin for *Coluber renardi* very dubious in the face of the fact that this species hitherto has not been recorded from any locality east of the Altai Mountains. On the other hand, it must not be forgotten that *Coluber berus* and *C. renardi* occur in the same localities in western Siberia, and that Strauch himself did not distinguish between the two species. Nevertheless, *C. renardi* can not be admitted to the fauna of Sakhalin upon the strength of our specimen alone, the locality of which must be regarded as erroneous until corroborated by additional material.

Coluber renardi^a is quite distinct from *C. berus*. Our specimen (female) has 21 scale rows, 145 ventrals, and 25 subcaudals and shows all the characteristic features of the species, such as: Snout pointed, hollow with swollen canthal edge; one apical scale only in contact with the rostral; upper preocular long, anteriorly touching nasal; nostril small, located in the lower half of the nasal. The scutellation of the top of the head is almost identical with Boulenger's figure c on Plate LXIV.^b

COLUBER BERUS^c Linnæus.

1758. *Coluber berus* LINNÆUS, Syst. Nat., 10 ed., I, p. 217 (type-locality, Europe); 12 ed., I, 1766, p. 377.—LAURENTI, Syn. Rept., 1768, p. 97.—*Vipera berus* DAUDIN, Hist. Nat. Rept., VI, 1803, p. 89.—MIDDENDORFF, Sibir. Reise, II, II, Pt. 1, 1853, p. 247 (Udskoi Ostrog; sources of the Riv. Tugur).—STRAUCH, Mém. Acad. Sci. St. Pétersb. (7), XXI, no. 4, 1873, p. 206, 279 (Ussuri; Hadshi Bay; Udskoi Ostrog; Uisut Island; Sakhalin; Nikolayevsk; Possiet Bay).—JAN, Icon. Ophid., livr. 45, 1874, pl. II.—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, LX, Prilozhen. no. 5, 1889, p. 290 (Sakhalin); Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, 1905, p. 303 (Russia to Sakhalin).—BOULENGER, Ann. Mag. Nat. Hist. (6),

^a 1861. *Pelias renardi* CHRISTOPH, Bull. Soc. Nat. Moscow, XXXIV, Pt. 2, p. 599 (type-locality, Sarepta, SE. Russia).—*Vipera renardi* BOULENGER, Proc. Zool. Soc. London, 1893, pp. 598, 757, pl. LXIV (Southeastern Russia to eastern Turkestan); Cat. Sn. Brit. Mus., III, 1896, p. 475 (southern Russia and central Asia).

Named for councilor of State, Doctor Renard, then editor of the Bulletin of the Imperial Society of Naturalists of Moscow.

^b Proc. Zool. Soc. London, 1893.


^c *Berus* as the name of a water snake, probably *Natrix natrix*, is said to be used first by such medieval writers as Albertus Magnus, Vincent de Beauvais, etc.

V, Feb. 1890, p. 139 (Possiet Bay to Sakhalin); Cat. Sn. Brit. Mus., III, 1896, p. 476 (Sakhalin).—*Pelias berus* DOBROTVORSKI, Izvest. Sibir. Otd. Geogr. Obstchest., I, 1870 (p. 23) (Sakhalin).—DOEDERLEIN, Mith. Deutsch. Ges. Ost-Asiens, III, heft 22, Dec. 1880, p. 89 (Sakhalin).

1822. *Chersa vulgaris* FLEMING, Philos. Zool., II, p. 295.—*Berus vulgaris* SWAINSON, Classif. Fish. Amph. Rept., II, (Lardner's Cab. Encycl.), p. 362.

The numerous synonyms based upon European color varieties are here omitted.

No distinctions have been pointed out between east Asiatic and European specimens, and no tangible ones seem to exist. It is perhaps worth noticing that the number of ventrals of all the eastern specimens recorded is rather high, viz, between 143 and 158, averaging about 150, while in a large number of western specimens (167) the average is about 145. This tendency toward a lower number



FIGS. 358-360.—*COLUBER BERUS*. $14 \times$ NAT. SIZE. 358, TOP OF HEAD; 359, SIDE OF HEAD; 360, UNDER-SIDE OF HEAD. No. 14325, U.S.N.M.

of ventrals in the west is shown by a still lower average of 143 ventrals in 54 British specimens or even 137.5 in 11 specimens from the Pyrenean peninsula.

Description.—*Adult female*: U.S.N.M. No. 14325; Dui, island of Sakhalin (figs. 358-360). Rostral slightly higher than wide, well visible from above; snout above covered by about 16 scales, the fragments of internasals and prefrontals, of which two join the rostral and two larger ones cover the canthus rostralis between naso-rostral and supraocular; frontal as long as its distance from tip of snout, wider than supraoculars, from which it is separated by a series of scales; parietals irregularly broken up, the two large shields adjoining the frontal somewhat smaller than the latter; nostril large, in the middle of the posterior nasal, the anterior nasal, or prenasal, not entering nostril and therefore also known as naso-rostral; three loreals behind posterior nasal, one above, two below; eye on sides and below surrounded by nine scales, ten on right side; temporals numerous scales, more or less irregular, those of the anterior row not larger than the postoculars; nine supralabials, fourth and fifth largest,

situated under the eye and separated by a single row of scales; four lower labials in contact with anterior chin-shields which are much larger than the posterior; 21 rows of scales, all except outer row keeled and with two apical pits; 146 ventrals; anal entire; subcaudals in pairs, 22+ (tail defective). Color (in alcohol) above tawny-olive with a brownish black, very distinct zigzag band along the middle of back and tail, and a series of obscure, alternating, lateral spots; top of head brownish black continuous with a similarly colored chevron mark opening backward on occiput; a brownish black band from nostril through eye over lower temporals to the side of neck passing over the last supralabial at angle of mouth; canthus rostralis and supralabials yellowish, the latter mostly with a black line along the posterior suture; chin and throat yellowish, mottled with dark slate color; whole underside uniform slate black, each ventral narrowly edged with pale olive.

Dimensions.

	<i>mm.</i>
Total length (tail defective)	568
Snout to vent.....	520

The proportion of tail to total length in the female averages as 1 to 9.33.

In the *male* the tail is considerably longer, with more numerous subcaudals, the proportion to total length averaging as 1 to 7.33.

The viper very rarely exceeds 700 mm. in total length.

Variation.—The specimen described above represents the average of this species, but both scutellation and coloration are very variable. The degree of breaking up of the normal shields of the head is thus very different in different individuals, so that two specimens exactly alike can rarely be found; thus the frontal may be in contact with the supraoculars, or on the other hand it may be separated from the parietals by a series of scales; number of scales around the eye varies between 6 and 13, of supralabials between 6 and 10. The variation in number of ventrals and its correlation with the geographical distribution has been alluded to above (p. 446).

The coloration is also exceedingly variable, at least in European specimens, the ground color varying through all the shades of brown and olive from silvery gray to black, and the pattern showing endless modifications of the one described above which may be said to be typical.

Habitat.—The common viper extends from western Europe to the Japanese and Okhotsk seas in the East.

Numerous specimens were collected in the island of Sakhalin by Prof. F. Schmidt, in 1863. Nikolski says that this snake is very common on the island, that Dobrotvorski noted it in the southern part and that Poljakof collected a specimen in the neighborhood of Korsakovski.

Maack obtained it in Ussuri, and von Schrenck in various parts of Amurland. Dybowski sent to the museum in Warsaw, Poland, a specimen taken at Possiet Bay near the boundary between Korea and the Russian Coast Province, according to Strauch, who also mentions its occurrence at Udskoi Ostrog.

The United States National Museum has one specimen from Sakhalin obtained from the St. Petersburg Academy. The museum in Hamburg has three specimens collected by Dieckmann at Nikolayevsk (No. 2162).

List of specimens of Coluber berus.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.		Subcaudals.	Ocellars.	Supracaudals.
						Ventrals.	Anal.			
U.S.N.M.	14325	Female ^a	Dui, Sakhalin Island		St. Petersburg Academy.	21	146	1	9-10	9
Brit. Mus.	<i>a</i> ⁴	do. ^b	Sakhalin Island		do.	21	158	1	33	
Do.	<i>b</i> ⁴	do. ^b	do.		do.	21	143	1	32	
St. Petersburg.	1031	(c)	Dui, Sakhalin		F. Schmidt.	21	152		32	
Do.	1032		do.		do.	21	149		39	
Do.	1028		Ussuri.		Maack.	21	153		33	
Do.	1029		Bay Hadshi, Amurl.		L. v. Schrenck.	21	158		30	
Do.	1030		Uisut Island, Liman.		do.	21	151		32	
Do.	1035		Nikolayevsk.		do.	21	146		38	

^a Description p. 446; figs. 358-360.

^b Boulenger, Cat., III, p. 481.

^c Strauch, p. 279.

Family CROTALIDÆ.

The "pit vipers" receive their name from the deep pit or hole on the side of the face between the nostril and the eye, as shown in fig. 363. This cavity sinks deep into the maxillary bone and represents a "blind" sac lined with epidermis and not connected with any of the other cavities or organs in the head by any inside opening or canal.

The maxillary bone, into the lower end of which the large hollow fang is immovably fastened like the knife in a handle, is extremely shortened and higher than long so as to appear to be in a vertical position (fig. 361). On the outer face of this bone there is the deep cavity forming the bony walls of the pit already referred to, which separates two articular surfaces. The upper one at the top of the maxillary forms with the corresponding concave face of the lachrymal (prefrontal) bone, which projects from and articulates with the frontal bone, a hinge-like joint allowing considerable freedom of motion. The lower articular surface receives the flattened anterior end of the external pterygoid bone (ectopterygoid or transpalatine). It will be

seen from the accompanying cut (fig. 361) that if the latter bone (*a*) be moved forward or backward, the maxillary (*d*) hinges on the lachrymal (prefrontal) (*e*) and that if the ectopterygoid be pushed forward the fang is erected.

For a more detailed description of the poison apparatus, the poison, etc., see my Poisonous Snakes of North America.^a

The Crotalid snakes are found in both hemispheres. They are entirely absent, however, in Africa and the southwestern corner of Asia, as well as in Australia. They are practically unknown in Europe also, except that one species enters the extreme south-eastern corner.

The general impression that the *Crotalidæ* are overwhelmingly American in their distribution is entirely erroneous, as nearly 40 per cent of the known species occur in Asia, and I am strongly of the opinion that the family originated in the Old World and spread to America afterward. The rattlesnakes which are peculiar to the latter continent are the latest specialization of the type.




FIG. 361.—POISON APPARATUS OF A CROTALID SNAKE. RIGHT SIDE. *a*, EXTERNAL PTERYGOID BONE (ECTOPTYERYGOID OR TRANSPALATINE); *b*, INTERNAL PTERYGOID BONE (ENTOPTERYGOID); *c*, PALATAL BONE; *d*, MAXILLARY BONE; *e*, LACHRYMAL BONE (PREFRONTAL). (AFTER MITCHELL.)

KEY TO JAPANESE GENERA OF CROTALIDÆ.

- a*¹ Top of head with large symmetrical shields *Agkistrodon*, p. 449.
*a*² Top of head covered with scales *Trimercsurus*, p. 465.

Genus *AGKISTRODON* ^b Beauvois.

1799. *Agkistrodon* BEAUVOIS, Trans. Amer. Philos. Soc., IV, p. 381 (type, *A. mokasen*).
 1802. *Scytale* LATREILLE, Hist. Nat. Rept., III, p. 158 (same type).
 1803. *Cenchris* DAUDIN, Bull. Soc. Philom. (Paris), III, No. 72, March 1803, p. 188 (same type).
 1819. *Scytalus* RAFINESQUE, Am. Journ. Sci., I, p. 84 (emendation).
 1826. *Tisiphone* FITZINGER, Neue Class. Rept., pp. 34, 63 (type, *T. cupreæ*).
 1826. *Trigonocephalus* BOIE, Isis, 1826, p. 214 (not of Oppel, 1811).
 1836. *Acontias* TROOST, Ann. Lyc. Nat. Hist. N. Y., III, p. 190 (type, *A. leucostoma*; not of Cuvier, 1829).
 1836. *Toxicophis* TROOST, Ann. Lyc. Nat. Hist. N. Y., III, p. 190 (same type).
 1849. *Halys* GRAY, Cat. Snakes Brit. Mus., p. 14 (type, *Trigonocephalus halys* Boie; not of Fabricius, 1803).
 1854. *Ancistrodon* BAIRD, Serp. N. Y., p. 13 (emendation) (not of Roemer, 1849).

Among the species of the genus *Agkistrodon* which extends from the northern borders of the Caspian Sea throughout most of the Asiatic

^a Rept. U. S. Nat. Mus. for 1893, pp. 365-380 and 457-480.

^b From ἄγκιστρον, hook; δδών, tooth. The correct transliteration would be *Ancistrodon*.

mainland and in North and Central America there is a small compact group consisting of the species *A. halys*, *blomhoffii* and *himalayanus* occupying the vast territory from the Caspian Sea in the west to the Pacific Ocean in the east, and from Lake Baikal in the north to the Himalayas in the south. They are closely interrelated, in fact so nearly allied that their descent from a common ancestor can not have taken place at a very distant period. Of these, *A. blomhoffii*, which is the only species occurring within our limits, shows certain differentiations according to locality, making it advisable to discriminate between several forms.

The *Aglistrodon intermedius* which Strauch described in 1868 and characterized as an *A. blomhoffii* with 23 scale rows, Boulenger, on the other hand, diagnoses as an *A. halys* without the snout turned up. Practically both seem to be right, for as far as I can see *A. intermedius*, as accepted by them, is not a well-defined form (much less a species as this term is used conventionally), but an aggregation of more or less intermediate specimens. In the sense in which they have been adopted hitherto neither have a definable range. According to Strauch not only does *A. blomhoffii* occur in the same localities on the mainland as *A. intermedius*, but the latter occurs also in Japan promiscuously with the former, as individuals with 23 scale rows occur both in southern and northern Japan. The trouble is that the number of scale rows is not constant, the inconstancy apparently varying to some extent with the locality. On the other hand, to what extent the turning up of the end of the snout may serve in all instances as a character to separate *A. halys* I can not say for lack of material, and for that reason I shall at present treat the latter as a good species.

For the study of *A. intermedius* and *A. blomhoffii* I have collected about 100 more or less complete scale formulas of individual specimens, all of which will be found in the appended tables.

Table Ia is a record of 27 specimens of undoubted Japanese origin.^a They show *A. blomhoffii* in its purity with a normal scale formula of 21 scale rows; 132-146 ventrals (average 141.3); 44-56 pairs of subcaudals (average 49.2); and 7 supralabials. Two of the specimens in the United States National Museum, viz, Nos. 31866 and 34040, have 23 scale rows, and Strauch mentions that in the Leiden Museum he found some specimens from Japan also with 23 scale rows, while Hilgendorf records one of his Tokyo specimens as having 23 rows. On the other hand, the number of supralabials (7) seems to be absolutely constant in Japan.

Table II contains formulas of 20 specimens from Korea, southeastern China, and Formosa, viz, 21-23 scale rows; 138-151 ventrals

^a For reasons given farther on (p. 452), I have not included in this table certain specimens in the St. Petersburg Academy Museum, recorded by Strauch as being from Japan.

(average 143.8); 29–46 subcaudals (average 35). The supralabials are not given in detail, but judging from Boulenger's account all the specimens examined by him had 7. One of my Korean specimens has 8 supralabials on one side. It will be seen that these formulas agree tolerably with those from Japan *except* that the subcaudals are much fewer.

Tables III and IV embrace the specimens recorded from the region lying to the north and northwest of those contained in Table II—i. e., from the Russian Coast Province and the River Amur to eastern Turkestan. They represent mostly the specimens attributed to *A. intermedius* by Strauch and Boulenger. They agree in having an intermediate number of subcaudals, viz, 36–49 (average 41) and [32] 37–53 (average 43.7), respectively. They also agree fairly well in the number of ventrals, viz, 151–163 (average 155.8) and [147] 154–179 (average 163.5), respectively, but it will be seen that these figures greatly exceed the corresponding ones of Tables I and II. There is, however, one marked difference between Tables III and IV, since it will be found that among the specimens collected east of the Khingan Mountains in Manchuria (Table III) the great majority (70.6 per cent in a total of 17 specimens) have 21 scale rows, while of those from the Khingan Mountains and the country to the west, viz, Mongolia, Transbaikalia, and eastern Turkestan, the greater number (76 per cent, in a total of 29 specimens) have 23 rows or more.

Before proceeding farther a few words may be said regarding Table 1*b*. The two specimens from Yaeyama in the United States National Museum agree structurally and in their scale formulas exactly with the Japanese specimens, but they differ exceedingly from the latter in color. Their ground color (in alcohol) is a "tawny olive,"^a uniform on top of head, but indistinctly varied and marbled on back and sides with blackish and with indications of pale cross bands; a band slightly darker than the ground color through eye and temples; lips whitish, as is also the whole underside; each ventral sprinkled with irregular blackish brown dots and spots on the basal half. It seems probable that they represent a permanent local color race, and are so recognized further on.

The differences between the specimens listed in Tables I–IV may be synoptically expressed as follows:

- a*¹ Ventrals 151 or less.
*b*¹ Subcaudals 43 or more.....Tables I and 1*a* (Japan and Yaeyama Islands).
*b*² Subcaudals 46 or less.....Table II (SE. China; Korea; Formosa).
*a*² Ventrals 147 or more.
*b*¹ Scale rows (in 70.6 per cent) 21..Table III (Coast Province; Amurland, west to Khingan Mountains).
*b*² Scale rows (in 79 per cent) 23....Table IV (Khingian Mountains to East Turkestan).

^a Ridgway, Nomencl. Colors, 1886, pl. III, No. 17.

It will be seen that a considerable amount of intergradation occurs. Thus it would be impossible to say to which of the three forms (Tables II, III, or IV) a specimen with 151 ventrals and 45 subcaudals were to be referred, unless it had 8 supralabials, in which case it would probably be correct to refer it to either Tables III or IV, with the chances in favor of Table III, if it had 21 scale rows, and in favor of Table IV if it had 23 rows. Unfortunately, the number of supralabials is not given in detail in the recorded scale formulas, or we might perhaps have been able to make a more definite statement. We only know in a general way that quite a large number of the specimens included in Tables III and IV have 8 supralabials, and I hold it to be probable that there are more having 8 supralabials in Table IV than in Table III.

I now call attention to Table V, containing the records of five specimens in the Zoological Museum of the St. Petersburg Academy of Sciences, which are reported to be from Japan. These, with a number of other snakes, were sent to the academy in 1861 by Mr. Goschkewitsch, who was Russian consul at Hakodate, Japan; hence Strauch concluded that they were from Yezo. Of the five *Agkistrodon*s, he referred No. 2225, with 23 scale rows, to *A. intermedius*; the other four, with only 21 scale rows, to *A. blomhoffii*, the typical Japanese species. Upon this evidence rests the admittance of *A. intermedius* into the fauna of Japan proper. A glance at Table V shows at once, however, that the five specimens did not come from the island empire at all. They belong most assuredly to Table III, within the limits of which not only the extremes are easily accommodated but with the averages of which their own averages nearly coincide (ventrals, respectively, 156.4 and 155.8; subcaudals 42 and 41). If we consider, furthermore, that Mr. Goschkewitsch sent two other species, which nobody else has found in Japan (namely, *Elaphe dione*, see p. 318, and *E. schrenckii*, see p. 315), but which occur in the Amur Province, I think there can not be a shadow of a doubt that all the specimens sent by him were only shipped from Japan but collected somewhere on the mainland, either in the Coast Province or in Amurland.

TABLE Ia.--Japan.

Museum.	No.	Sex and age.	Locality.	Authority.	Scale rows.	Ventrals.	Subcaudals.	Supracaudals.
U. S. N. M.	15421	Female ..	Omachi, Hondo	Stejneger	21	139	53	7
Do.	15422	Young ..	do	do	21	142	48	7
Do.	15423	do	do	do	21	132	47	7
Do.	15424	do	do	do	21	144	49	7
Do.	15425	do	do	do	21	139	53	7
Do.	15426	do	do	do	21	141	54	7
Do.	17847	do	Japan	do	21			7
Do.	23436	Halfgr.	Yezo	do	21	143	52	7
Do.	31866	Female ..	Shikoku	do	23	140	47	7
Do.	31867	Male	do	do	21	142	45	7
Do.	34023	Female ..	Yokohama	do	21	140	49	7
Do.	34024	Male	do	do	21	144	56	7
Do.	34040	Female ..	do	do	23	137	45	7
Do.	34041	Male	do	do	21	144	55	7
Do.	34042	do	do	do	21	146	49	7
Do.	34043	Female ..	do	do	21	143	47	7
		(a)	Japan	Boie	21	142	45
Sci. Coll. Tokyo	59	Female ..	Hachijo shima	Stejneger	21	139	46	7
St. Petersb.	2231	do	Japan	Strauch	21	140	46
Leiden		do	do	Schlegel		136	46
Do.		do	do	do		142	56
Brit. Mus.	a	Male	do	Boulenger	21	140	48	7
Do.	b	Female ..	Yokohama	do	21	145	47	7
Do.	c	Male	do	do	21	142	55	7
Do.	e	do	Tsu-shima	do	21	146	46	7
Do.	f	Halfgr.	do	do	21	138	44	7
Do.	g	Young ..	do	do	21	145	45	7
			Average			141.3	49.2

^a Type of *blomhoffii*.

TABLE Ib.--Yae-yama Islands.

U. S. N. M.	34038	do	Yae-yama	Stejneger	21	139	49	7
Do.	34039	do	do	do	21	138	53	7
Brit. Mus.	r	Male ^a ..	(?)	Boulenger	21	137	46	7
			Average			138.0	49.3

^a Type of *afinis*.

TABLE II.—(*East China; Korea; Formosa.*)

Museum.	No.	Sex and age.	Locality.	Authority.	Scale rows.	Ventrals.	Subcaudals.	Supracaudals.
U. S. N. M.	14612		Korea	Stejneger	21	142	34	7.8
Do.	14613		do.	do.	21	146	37	7
Do.	17507	Male ^a	Fusan, Korea	do.	21	148	31	7
Do.	17508	do.	Seoul, Korea	do.	21	146	34	7
Mich. Univ.	1	Young	Korea	do.	21	150	42	7
Brit. Mus.	o	Male	Hang-Chau, China	Boulenger	21	144	41	7
Do.	p	do.	Ichang, China	do.	21	144	35	7
Do.	q	Female	do.	do.	21	140	30	7
Do.	r	Male	do.	do.	21	143	38	7
Do.	s	Female	Mountains north of Kiu-kiang, China.	do.	23	145	31	7
Do.	t	do.	do.	do.	21	142	32	7
Do.	u	do.	do.	do.	21	138	29	7
Do.	v	Young	do.	do.	21	138	32	7
Do.	w	Female	do.	do.	21	143	31	7
Do.	h	Male	Formosa	do.	23	151	40	7
Do.	i	do.	do.	do.	21	146	46	7
Do.	k	Female	Hainan	do.	21	142		7
Do.	l	do.	do.	do.	21	144	32	7
Do.	m	Young	do.	do.	21	144	39	7
Do.	n	do.	do.	do.	21	141	32	7
			Average			143.8	35.0	

^a Type of *brevicaudus*.TABLE III.—(*Coast Province; Amurland, west to Khingan Mountains.*)

St. Petersb.	2219		Ussuri	Strauch	23	163	36	
Do.	2221		Cape Tyr.	do.	23	161		
Do.	2232		Possiet Bay	do.	21	156	40	
Do.	2233		Upper Amur	do.	21	153	41	
Do.	2234		Komar	do.	21	154	37	
Do.	2236		Amur	do.	21	151	47	
Do.	2237		Ana, Ussuri	do.	21	157	44	
Do.	2238		Amur	do.	21	151	49	
Do.	2240		Bay Guerin	do.	21	153	41	
Do.	2241		Adi, Amur	do.	21	154	41	
Do.	2242		Noor, Ussuri	do.	21	158	41	
Do.	3723		Tschianka, Amur	do.	21	155	36	
Do.	3723		Ussuri	do.	21	158	42	
Do.	3724		do.	do.	21	155	43	
Brit. Mus.	c	Male	Khabarovka	Boulenger	23?	155	40	
Do.	d	do.	do.	do.	23?	156	40	
Do.	e	Female	Ussuri River	do.	23?	158	39	
			Average			155.8	41.0	

TABLE IV. - *Khingan Mountains to East Turkestan.*

Museum.	No.	Sex and age.	Locality.	Authority.	Scale rows.	Ventrals.	Subcaudals.	Supralabials.
St. Petersb.	2216		Padun.....	Strauch.....	23	162	45
Do.	2217		do.....	do.....	23	164	39
Do.	2218		Nikolski Zavod.....	do.....	23	161	43
Do.	2222		Khingan.....	do.....	23	157	42
Do.	2223		Daurian Steppes.....	do.....	23	163	32
Do.	2224		Tarei Noor.....	do.....	23	167	41
Do.	2226		Argun River.....	do.....	25	160	43
Do.	3719		Bukukun.....	do.....	23	163	40
Do.	3720		Khingan.....	do.....	23	161	37
Do.	2235		Argun.....	do.....	21	160	42
Do.	2239		Ust Strielka.....	do.....	21	154	46
Do.	2243		Gobi.....	do.....	21	147	51
Do.	3721		Khingan.....	do.....	21	155	49
Do.			Ordos.....	do.....	23	179	44	7-8
Do.			do.....	do.....	23	169	42	8
Do.			Ala-shan Governm.....	do.....	21	163	48	8
Do.			do.....	do.....	21	168	37	7
Stockholm.	1806	Female	Altai.....	Stejneger.....	25	172	43	8
Do.	1807	Male	do.....	do.....	21	175	51	7-8
Do.	1808	Female	do.....	do.....	23	163	47	8
Do.	1809	do	do.....	do.....	23	171	46	7-8
Do.	1810	Male	do.....	do.....	23	161	53	8
Do.	1811	do	do.....	do.....	23	166	47	7
Do.			Southern Mongolia.....	Méheli.....	23	163	37
Do.			do.....	do.....	23	170	43
Brit. Mus.	a	Halfgr	East Turkestan.....	Boulenger.....	23?	161	49
Do.	b	Female	Smeinogorsk.....	do.....	23?	166	45
Do.	f	Halfgr	Lob Nor.....	do.....	23?	157	42
Do.	g	do	do.....	do.....	23?	164	44
			Average.....			163.5	43.7

TABLE V. - "*Japan*" *Goschkewitsch, 1861.*

St. Petersb.	2225		"Japan" (?).....	Strauch.....	23	162	38
Do.	2227		do.....	do.....	21	153	42
Do.	2228		do.....	do.....	21	157	42
Do.	2229		do.....	do.....	21	157	41
Do.	2230		do.....	do.....	21	153	47
			Average.....			156.4	42.0

It may now be regarded as demonstrated that the only form found in the islands of Japan proper is *Agkistrodon blomhoffii* which, so far as the accessible material and records go, is the best differentiated form.

The Korean and Chinese forms differ only in the fewer subcaudals. There is no absolutely sharp line between the two forms inasmuch as 44

is the minimum of the former and 46^a the maximum of the latter. Yet, of the 36 specimens enumerated, only one (Brit. Mus. No. *h*), or less than 3 per cent would be unidentifiable by that test. The averages 49 and 35.3 are much more significant, and it is scarcely to be doubted that we have to do with a form which it will be advantageous to recognize by name.


Contiguous to this South Chinese and Korean form we have another occupying the mainland to the north and northwest. Its chief characteristic consists in the greater number of ventrals, which in 36 specimens vary between [147] 151-170 (average 158.4). In addition numerous specimens have eight supralabials and about one-half 23 scale rows instead of 21. With regard to the latter character it is possible to divide the localities of these specimens into two classes, the great majority of them occupying the territory east of the Khingan Mountains having 21 scales, those farther west 23. The percentage of exceptions in each group, namely, respectively 29 and 24 per cent—is too great to make it advantageous to recognize two forms. Unfortunately, the published records do not indicate whether the number of supralabials is more constant. For the present at least they must be united under the name of *A. blomhoffii intermedius*, a term consequently somewhat more comprehensive than Strauch's original *A. intermedius*.

We have thus three, or possibly four, local forms which may be recognized by the following key:

- a*¹. Ventrals 151, or less.
*b*¹. Subcaudals 44, or more.
*c*¹. Underside black, more or less blotched with whitish. *A. blomhoffii*, p. 457.
*c*². Underside whitish, sprinkled with blackish. *A. blomhoffii*? *affinis*, p. 461.
*b*². Subcaudals 46, or less. *A. blomhoffii brevicaudus*, p. 463.
*a*². Ventrals (147) 151, or more. *A. blomhoffii intermedius*, p. 464.

The most interesting conclusion to be drawn from the above is that the typical Japanese form is least closely related to the northern *A. intermedius*, which occurs on the opposite side of the Sea of Japan, and that their connection is only effected through the shorttailed southern Chinese form, a conception entirely different from the one formerly held, according to which there were two distinct species, both occurring on either side of the Sea of Japan. The latter would involve a previous northern line of intercommunication between the two territories, while now it is seen that the connection between these forms lies toward the south.

^a It would appear that the number may exceptionally reach 50, which is the maximum noted by Doctor Werner for some specimens from Hankow. I will here again call attention to the fact that possibly some of Doctor Haberer's specimens alleged to have been collected at Hankow in reality may be from Japan, where he also collected. (See under *Elaphe climacophora*, p. 326; *quadrivirgata*, p. 331; and *conspicillata*, p. 336.)



AGKISTRODON BLOMHOFFII. (FROM TEMMINCK AND SCHLEGEL.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

AGKISTRODON BLOMHOFFII^a (Boie).

MAMUSHI.

Plate XXVI.


1826. *Trigonocephalus blomhoffii* BOIE. Isis, 1826, p. 214 (type-locality, Japan; Blomhoff, collector); Bijdr. Natuurk. Wetensch., II, Pt. 1, 1827, p. 268.— TEMMINCK and SCHLEGEL. Fauna Jap., Rept., 1837, pp. 88, 96, 139; Ophid., pl. vi b (Kiusiu; Hondo).—DUMÉRIL and BIBRON. Erpét. Gén., VII, Pt. 2, 1854, p. 1496 (Japan).—BLEEKER. Natuurk. Tijdschr. Nederland. Indië, XVI, 1858, p. 204 (Japan).—STRAUCH. Mém. Acad. Sci. St. Pétersb. (7), XXI, No. 4, 1873, pp. 251, 282 (part: spm. No. 2231; Japan, Maximowitch, collector).—HILGENDORF, Mitth. Deutsch. Ges. Ost-Asiens, I, heft 10, July 1876, p. 30 (Tokyo).—EASTLAKE. Trans. Asiatic Soc. Japan, XIII, 1885, p. 72 (Japan); Nature, XXXI, April, 1885, p. 587.—OKADA, Cat. Vert. Japan, 1891, p. 69 (Kaga; Kyoto; Tokyo; Awa; Hondo; Hachijo shima; Echigo; Awaji; Yezo; Toza).—*Halys blomhoffii* GUENTHER. Rept. Brit. India, 1864, p. 393 (part: Japan).—HILGENDORF. Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 117 (Tokyo).—*Ancistrodon blomhoffii* BOULENGER. Cat. Snakes Brit. Mus., III, 1896, p. 525 (part: Yokohama; Tsushima).—WALL Proc. Zool. Soc. London, 1903, p. 102 (part: Japan); 1905, II, p. 514 (Hondo; Yezo; Tanega I.; Yaku I.).
1837. *Trigonocephalus blomhoffi* SCHLEGEL, Phys. Serp., II, p. 552 (Japan).—FRITZE. Mitth. Deutsch. Ges. Ost-Asiens, V, heft 46, 1891, p. 238 (Yezo).—*Ancistrodon blomhoffi* BOETTGER. Kat. Schl. Mus. Senckenberg., 1898, p. 137 (Hondo; Yezo).—SCHNEE. Zool. Garten, XLI, 1900, p. 395 (color description from life).
1859. *Trigonocephalus blomhoffii* var. *megaspilus* COPE. Proc. Phila. Acad., 1859, p. 336 (type-locality, unknown; type in Phila. Acad. Mus.; collected by the Perry Expedition).
1862. *Halys blomhoffii* PETERS. Mon. Ber. Berlin Akad. Wiss., 1862, p. 671 (*lapsus*).

Description.—*Adult male*; U.S.N.M. No. 31867; Kochi, province of Toza, Shikoku, May, 1903; Dr. Hugh M. Smith, collector (figs. 362–364). Rostral as high as broad scarcely visible from above; internasals small, subtriangular, their suture half that between prefrontals which are broadly in contact with supraoculars; frontal slightly longer than broad, as long as its distance from rostral and as interparietal suture, nearly straight in front; supraoculars longer than frontal and nearly as broad at the center of eye; parietals as long as supraoculars, longer than distance of frontal from tip of snout; nostril roundish in the posterior edge of the anterior nasal which is twice as large as the posterior; two loreals one above the other, the lower one bordering the pit anteriorly; a narrow subfoveal bordering the pit below posteriorly, not reaching the eye; two preoculars, of which the lower borders the pits above posteriorly; pit nearer eye than nostril; two postoculars, of which the lower is long, narrow, and

^a Blomhoff was director of the Dutch factory on Decima for more than ten years during the early part of the nineteenth century. His collections were brought to Batavia, where H. Boie described the reptiles and batrachians.

^b Reproduced in this work on Plate XXVI.

crescentic, nearly reaching the lower preocular below the eye and separating the latter from fourth supralabial; 2+4 temporals, of which the lower ones in each row are large hexagonal flat shields, those above being small scales, keeled on the row nearest the parietals: the lower temporal of the third row large and shaped like those in front of it; these three lower temporals form a series of large shields larger than the adjoining labials; 7 supralabials, second smallest, third and fourth largest, and those behind gradually decreasing in size. third just entering eye; five lower labials in contact with anterior chin-shields, which are much larger than the second pair, the latter being scarcely differentiated from the scales behind and between them; 21 rows of strongly keeled scales with a pair of apical pits; 142 ventrals; anal undivided; 45 subcaudals, all divided. Color (in alcohol) above brownish gray with a series of large, rhomboid, darker



FIGS. 362-364.—*AGKISTRODON BLOMHOFFII*. $1\frac{1}{2} \times$ NAT. SIZE. 362, TOP OF HEAD; 363, SIDE OF HEAD; 364, UNDERSIDE OF HEAD. No. 31867, U.S.N.M.

brown, black-edged blotches on each side close to the median dorsal line, alternating, often confluent, descending on the sides to the outer scale row, separated by a narrow paler grayish band the edges of which nearest the black margin of the dark blotches incline to whitish; top of head of lighter brown, with a dark irregular marking on each shield; a sharply marked blackish brown stripe on lower preocular through eye where it widens to a very broad band on the temples, passing across the last supralabial at the corner of the mouth, the sharp edges of this band emphasized by a narrow white line above and below, the one below having a small sharply defined white spot on the anterior angle of the long crescentic lower postocular; underside irregularly mottled with whitish and blackish in about equal proportions; tip of tail brownish white.

Dimensions.

	<i>mm.</i>
Total length.....	513
Snout to vent.....	438
Vent to tip of tail.....	75

The *young* have clearer, brighter colors, more contrasting pattern, and the tail always light colored toward the tip both above and below.

Variation.—The most obvious instability in the scale formula is the varying number of temporals, but this variability is only confined to the upper scales near the parietals. As a matter of fact the outer edges of the later shields seem to be on the verge of breaking up into small scales as evidenced by the many nicks and half-finished sutures on their outer margins. The prefrontals also show sign of breaking up, and there is a distinct regularity in this process. In nearly all the specimens the inner anterior corner of these shields is more or less convex, and in a number of cases this slightly elevated region is cut off by regular sutures so as to form an additional anterior pair of prefrontals adjoining the internasals. Such is the adult female, No. 34043, from Yokohama. Quite similar in this respect is the adult female No. 15421, from Omachi, province of Shinshiu, and what is even more interesting, no less than three of her five young just born have these accessory prefrontals exactly like the mother, while only two of them are normal.

The number of supralabials, 7, is remarkably constant. Less so is the number of scale-rows, namely, 21, but 23 is rather of rare occurrence. The ventrals range between 132 and 146; subcaudals between 44 and 56 pairs.

The coloration is very variable. While specimens lighter than the one described above do occur, individuals much darker seem to be more frequent, and it is not uncommon to find specimens with the whole underside uniformly black or nearly so. While the tip of the tail seems to be always light yellowish in the young I find among the adults before me an even proportion of specimens with dark and light tail ends, there being apparently no distinction as to sex or locality.

The specimen from Hachijo shima (U.S.N.M. No. 36540), the southernmost island of the Idzu group, kindly submitted to me by Doctor Ijima, differs greatly in coloration from the normal specimens. The ground color is of a rich reddish brown with nearly all traces of blackish gone. In all other respects the specimen seems to be normal. Doctor Ijima writes me that he has six more specimens from the same locality of a similar coloration, but adds that the Science College Museum also has a specimen (No. 58) from the same island of a dark color similar to those from Hondo. In view of this fact I refrain from imposing any systematic name on this color variety.

Habitat.—As here understood *A. blomhoffii* is restricted to the islands of Japan, being represented on the mainland opposite by closely allied forms, in Korea by *A. blomhoffii brevicaudus*, in the Russian Coast Province by *A. blomhoffii intermedius*.

The mamushi occurs more or less frequently on the four principal islands. Eastlake says that it "is most frequent in the neighborhood of Kyoto, also the environs of Yokohama and the Hakone Mountains; tolerably frequent in the north." In Yezo it was observed by Doctor Fritze on the road to Kamigawa in the neighborhood of Sorachi-buto, and the United States National Museum, thanks to the kindness of Doctor Nozawa, has a specimen, which probably came from near Sapporo.

It seems to be common in the seven islands of Idzu, at least on Hachijo shima, from which island the Science College has six specimens, the majority of an unusual bright reddish color.

Doctor Wall records two specimens from the islands south of Kiusiu, namely, one from Yaku shima and one from Tanega shima, the former with 143, the latter with 142 ventrals, and both with 23 scale rows.

Holst collected specimens on Tsu-shima, which are now in British Museum.

List of specimens of Agkistrodon blomhoffii.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supraorbitals.
			1882.								
15421	Female ^a .	Omachi, Shinshiu	Oct. —	P. L. Jouy	21	143	1	48	2, 2	2+4	7
15422	Young ^b	do.	do.	do.	21	142	1	48	2, 2	2+3	7
15423	do. ^b	do.	do.	do.	21	132	1	47	2, 2	2+4	7
15424	do. ^b	do.	do.	do.	21	144	1	49	2, 2	3+4	7
15425	do. ^c	do.	do.	do.	21	139	1	53	2, 2	2+4	7
15426	do. ^c	do.	do.	do.	21	141	1	54	2, 2	3+4	7
17847		Japan		H. Loomis	21						7
23436	Half gr.	Yezo		S. Nozawa	21	143	1	52	2, 2	2+3-4	7
			1903.								
31866	Female.	Kochi, Shikoku	May —	Dr. H. M. Smith	21	140	1	47	2, 2	2+4	7
31867	Male. ^d	do.	do.	do.	21	142	1	45	2, 2	2+4	7
			1902.								
34023	Female.	Yokohama	June 20	A. Owston	21	140	1	49	2, 2	2+3	7
34024	Male.	do.	do.	do.	21	144	1	56	2, 2	2+4	7
			1894.								
34040	Female.	do.	Aug. —	do.	23	137	1	45	2, 2	2+3-4	7
			1901.								
34041	Male.	do.	July 24	do.	21	144	1	55	2, 2	2+4	7
			1902.								
34042	do.	do.	June 6	do.	21	146	1	49	2, 2	2+3	7
			1901.								
34043	Female. ^e	do.	Aug. —	do.	21	143	1	47	2, 2	2-3+4	7
			1887.								
36540	do.	Hachijo shima	May —	N. Okada	21	139	1	46	2, 2	2+3	7

^a 2 pairs of prefrontals.

^b Young of 15421; 2 pairs of prefrontals.

^c Young of 15421.

^d Description, p. 457; figs. 362-364.

^e 2 pairs of prefrontals.

AGKISTRODON BLOMHOFFII ?AFFINIS^a (Gray).

1849. ? *Trigonocephalus affinis* GRAY, Cat. Sn. Brit. Mus., p. 14 (type-locality unknown; type in Brit. Mus.; Capt. Sir Edward Belcher, collector).

Allusion has already been made (p. 451) to two specimens from Yaeyama, in the United States National Museum, which in every respect except color agree with the specimens from Japan proper. As there stated their ground color (in alcohol) is a "tawny-olive," uniform on top of head, but indistinctly varied and marbled on back and sides with blackish, and with indications of pale cross bands; a band slightly darker than the ground color through eye and temples; lips whitish, as is also the whole underside, each ventral sprinkled with irregular blackish brown dots and spots on the basal half.

I venture to suggest that the type specimen of Gray's *Trigonocephalus affinis* in British Museum may belong to this form. At least his brief description seems to justify such a suggestion. The specimen has no locality attached to it, but it was presented to the museum by Capt. Sir Edward Belcher. As noted later under *Trimeresurus elegans* (p. 471.) Belcher surveyed and collected on Ishigaki shima, in the Yaeyama group for three weeks, and there is consequently a possibility that the type of *Trigonocephalus affinis* came from that island. Under these circumstances I adopt Gray's name provisionally and with the necessary reservation.

Finally I wish to call attention to specimen *d* in Boulenger's enumeration of the specimens of *A. blomhoffii* in the British Museum.^b It is a female with 142 ventrals and 43 subcaudals, presented by Mr. M. K. Rokugo, and said to have been collected in "Okinawa, Loo Choo Islands." Nobody else has recorded the mamushi from the middle group or Okinawa shima proper, and it is therefore rather probable that the specimen in question came from some other locality, possibly from the southern group.

It is not in the least probable that the Yaeyama *Agkistrodon* is directly connected, in a genetic sense, with the typical *A. blomhoffii* in Japan. It seems much more likely that the former is a slight modification of the Formosan stock which has developed in the same direction as the Japanese form, viz, toward an increased number of subcaudals, assuming at the same time a peculiar pale coloration. As a slight indication of their relationship with the mainland form rather than with the one from Japan, I may mention the absence of the white spot on the anterior corner of the long lower postocular.

As previously intimated, the Yaeyama specimens are mainly distinguished from the typical *A. blomhoffii* by the coloration and a detailed description of the lepidosis is therefore unnecessary. I wish, however, to call attention to a peculiarity in their scutellation (fig. 365),

^a Signifying related.

^b Cat. Sn. Brit. Mus., III, p. 526.

which may or may not indicate a real difference. This feature consists in the consolidation of the two scales above the large lower temporals of the second and third rows into a single elongate shield which is in contact anteriorly with the large lower shield of the first row without extending above it; at the same time the two lower shields of the second and third rows are reduced in size and their upper edges form a straight line parallel with the edge of the lip. The arrangement does not appear abnormal and is absolutely identical in both specimens, though in No. 34039 it is only found on the right side, while on the left it is as in typical *A. blomhoffii*.

Color description.—*Adult male*; U.S.N.M., No. 34038; Yaeyama, Riu Kiu Archipelago; October 12, 1900; A. Owston collection.




FIG. 365.—*AGKISTRODON BLOMHOFFII*? AFFINIS. $\frac{1}{4}$ X NAT. SIZE. SIDE OF HEAD. NO. 34038. U.S.N.M.

The predominating color (in alcohol) above is a pale drab, with very faint indications of slightly browner cross-bars with traces of darker gray edges, the cross-bars separated by narrow slightly paler interspaces; on the outer row of scales and the adjoining corner of the ventrals a series of irregular cinnamon-colored spots about 2–3 ventrals apart: head uniform, the temporal region darker, especially on the upper border of the adjoining supralabials on which the contrast with the pale color of the rest of these shields is fairly well contrasted; only faint traces of a whitish line above and below the dusky area on the temporal region; lower postocular uniform drab without anterior white spot; underside whitish with slight and irregular dark gray markings at the base and outer corner of the ventrals.

Dimensions.

	<i>mm.</i>
Total length.....	404
Snout to vent.....	340
Vent to tip of tail.....	64

Variation.—No. 34039 differs chiefly in being still paler and more cinnamon-colored with even less traces of a pattern on the upper side and on sides of head; on the other hand, the dark gray markings on the underside are distinctly darker but not larger.

The difference in the scutellation of the temporal region on the two sides of this specimen has been alluded to above.

Habitat.—Uncertain as the whole status of this form is, the only statement which can be made with certainty refers to the two specimens in the U. S. National Museum, which were collected October 12, 1900, on Yaeyama, according to Mr. A. Owston.

Whether it occurs on any of the other islands it is impossible to say now, but it appears probable that it is confined to the southern group of the Riu Kiu Archipelago.

List of specimens of *Agkistrodon blomhoffii*? *affinis*.

U.S. N.M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supralabials.
34038	Male ^a ...	Yaeyama, Riu Kiu.....	1900. Oct. 12	A. Owston.....	21	139	1	49	2, 2	2, 4	7
34039	do.....	do.....	do.....	do.....	21	138	1	53	2, 2, 2+3	4	7

^a Description, p. 462; fig. 365.

AGKISTRODON BLOMHOFFII BREVICAUDUS^a, new subspecies.

1856. *Trigonocephalus blomhoffii* HALLOWELL, Proc. Phila. Acad., 1856, p. 153 (Ningpo, China) (not of Boie, 1826).—*Halys blomhoffii* GUENTHER, Rept. Brit. India, 1864, p. 393 (part: Formosa).—SWINHOE, Proc. Zool. Soc. London, 1870, p. 412 (Ichang, China).—GIGLIOLI and SALVADORI, Proc. Zool. Soc. London, 1887, p. 594 (part: Fusan, Korea).—*Ancistrodon blomhoffii* BOULENGER, Cat. Snakes Brit. Mus., III, 1896, p. 525 (part: Formosa: Hainan, Chekiang, Ichang, Mts. north of Kiukiang, China).—WALL, Proc. Zool. Soc. London, 1903, p. 98 (part: Shanghai: Yangtse Valley).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 357 (Ningpo Mts. and Hankow, China).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, (8) XVII, no. 1, 1905, p. 329 (part: Korea).

1863. *Halys blomhoffii* SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 225 (Tamsui, Formosa).—BOETTGER, Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 153 (part: China).

Diagnosis.—Similar to *Agkistrodon blomhoffii*, but number of subcaudals averages only 35 (known minimum 29, known maximum 46); scale rows 21, rarely (about 10 per cent) 23; supralabials 7, very rarely (about 2½ per cent) 8; ventrals 138-151.

Dimensions of type specimen (adult male).

Total length.....	mm. 595
Snout to vent.....	532
Vent to tip of tail.....	63

Type.—U.S.N.M. No. 17507; Fusan, Korea; 1885; P. L. Jouy, collector.

Remarks.—This form is so much like typical *Agkistrodon blomhoffii*, from Japan, that no detailed description is deemed necessary. Of the four adult Korean specimens before me, three have uniform black belly. If this series can be trusted the dark postorbital band descends slightly farther down on the supralabials than in Japanese specimens. All lack the white spot at the anterior angle of the crescentic lower postocular.

Habitat.—Eastern China, Korea, and Formosa.

Our knowledge of the distribution of this form is very defective. In China it has thus far been found in the eastern part, although

^a Signifying short-tailed.

apparently always some distance back from the coast; for instance, at Ningpo, Ichang, Chekiang, Kiukiang, etc. It seems probable that it is confined to the more mountainous districts without coming down on the alluvial plains. Four specimens said to be from Hoi How, island of Hainan, were presented by J. Neumann to the British Museum. This locality needs corroboration ^a (see pp. 326, 331 and 336). In Korea it has been collected both at Seoul and Fusan by the late P. L. Jouy.

Two male specimens from Formosa collected by Swinhoe are in the British Museum.

List of specimens of *Agkistrodon blomhoffii brevicaudus*.

Museum.	No.	Sex and age.	Locality.	When collected	By whom collected.	Snout-vent.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supraductals.
U.S.N.M.	14612	Male	Korea		N. M. Fensholt.	21	142	1	34	2-2-3	2+4	7-8
Do	14613	Female	do		do	21	146	1	37	2-2-3-2-4-4	7	
Do	17307	Male ^a	Fusan, Korea	1885	P. L. Jouy.	21	148	1	31	2-2-3	2+4	7
Do	17508	do	Seoul, Korea	1883	do	21	146	1	34	2-3	2+4	7
Mich. Univ.	30525	Young ^b	Korea		G. A. Olinger.	21	150	1	42	2-3	2+4	7

^a Type; p. 463.

^b Total length, 335 mm.

AGKISTRODON BLOMHOFFII INTERMEDIUS ^b (Strauch).

1859. *Trigonocephalus blomhoffii* MAACK, Putesh. na Amur (p. 152) (Amurland (not of Boie); Putesh. na Ussuri, 1861 (p. 190) (valley of the Ussuri.—STRAUCH, Trudi Perv. Siezda Russkikh Yestestv., Zool., 1868 (p. 296).—Mém. Acad. Sci. St. Pétersb. (7), XXI, no. 4, 1873, pp. 251, 282 (Ussuri and Amurland; Possiet Bay and Vladivostok; De Castries Bay); in Przewalski's Mongoliya i Strana Tangutov, III, 1876, p. 52 (Ala-shan Government).—GIUGLIOLI and SALVADORI, Proc. Zool. Soc. London, 1887, p. 594 (part: Olga Bay, Russ. Coast Prov.).—*Ancistrodon blomhoffii* BOULENGER, Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 140 (Amurland; Ussuri).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, (8) XVII, no. 1, 1905, p. 329 (Amurl.; Ussuri, etc.)

1868. *Trigonocephalus intermedius* STRAUCH, Trudi Perv. Siezda Russkikh Yestestv., Zool., (p. 294); Mém. Acad. Sci. St. Pétersb. (7), XXI, no. 4, 1873, pp. 245, 282 (type-locality, Governm. Irkutsk, East Siberia; types 1, St. Petersb. Acad.); in Przewalski's Mongoliya i Strana Tangutov, III, 1876, p. 52 (Ordos).—*Halys intermedia* PETERS, Mon. Ber. Berlin Akad. Wiss., 1877, p. 735 (Smeinogorsk, Tomsk Governm.).—*Halys intermedius* BOETTGER, Offenbach. Ver. Naturk., 26-28 Ber., 1888, p. 153 (Nan-shan, near Kuku-nor, south. Gobi; Ala-shan Desert, Prov. Ordos; Russ. Turkestan; east. Siberia).—*Ancistrodon intermedius* BOULENGER, Ann. Mag. Nat. Hist. (6), V, Feb. 1890, p. 140 (Kunges, east. Turkestan;

^a Bocourt (Nouv. Arch. Mus. Paris, II, 1866, Bull., p. 9), in a list of the reptiles brought home by him from the southern provinces of Siam, mentions "*Trigonocephalus blomhoffii*." Locality and identification need corroboration.

^b Signifying intermediate, i. e., in characters between *A. blomhoffii* and *A. halys*.

Smeinogorsk; Khabarovka, Ussuri); Cat. Sn. Brit. Mus., III, 1896, p. 525 (Tarim Riv, Lob Nor).—MÉHELY, Zichy's Dritte Asiat. Forschungsr., II, 1902, p. 58 (Southern Mongolia: Chalute; Chere-muchor).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg, (8) XVII, no. 1, 1905, p. 326 (Amurland, Ussuri, etc.).

No specimen of this form is available for detailed description, which, moreover, seems to be unnecessary, as the chief difference is believed to consist in the greater number of ventrals and greater average of scale rows and supralabials, features previously discussed (pp. 450-456).

Habitat.—This form has a wide distribution from the Gulf of Tartary and the Sea of Japan, in the East, through southern Siberia and Mongolia to the Government of Tomsk and eastern Turkestan, in the West consequently extending over more than 50 degrees of longitude. At its eastern boundary it has been found as far north as Cape Tyr (above Nikolayevsk), on the Amur, and Decasties Bay, on the Gulf of Tartary. On the west shore of the Sea of Japan it goes southward at least to Possiet Bay. It seems to be common all over the Ussuri and Amur country, as well as along the coast, where, in addition to the localities already mentioned, it has been collected at Vladivostok and Olga Bay.

List of specimens of Agkistrodon blomhoffii intermedius.

Stockholm Mus. No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Oculars.	Temporals.	Supralabials.
1806	Female....	Altai, Siberia.....	1875	Slowzoff.....	25	172	1	43	2,2	2+4	8
1807	Male.....	do.....	1875	do.....	21	175	1	51	2,2	2+3	7-8
1808	Female....	do.....	1875	do.....	23	163	1	47	2,2	2+3	8
1809	do.....	do.....	1875	do.....	23	171	1	46	2,2	2+3	7-8
1810	Male.....	do.....	1875	do.....	23	161	1	53	2,2	2+3	8
1811	do.....	do.....	1875	do.....	23	166	1	47	2,2	2+3	7

Genus TRIMERESURUS,^a Lacépède.

1804. *Trimeresurus* LACÉPÈDE, Ann. Mus. d'Hist. Nat. Paris, IV, p. 209 (type, *T. viridis*).
1822. *Trimeresura* FLEMING, Philos. Zool., II, p. 291 (emendation).
1830. *Megara* WAGLER, Syst. Amph., p. 174 (type, *Vipera trigonocephala*).
1830. *Atropos* WAGLER, Syst. Amph., p. 175 (type, *Trigonocephalus puniceus*) (not of Oken, 1815).
1830. *Tropidolæmus* WAGLER, Syst. Amph., p. 175 (type, *Cophias wagleri*).
1839. *Trimeresurus* SWAINSON, Classif. Fish. Amph. Rept., II (Lardner's Cab. Encycl.), p. 363 (emendation).
1843. *Botrophis* FITZINGER, Syst. Rept., p. 28 (type, *T. viridis*).

^a From *τριμερής*, divided into three parts; *ὀψρά*, tail.

The snakes of this genus are distinguished from *Agkistrodon* at the first glance by the absence of the regular shields on top of the head, which are here replaced by scales more or less similar to those of the body.

It has not been demonstrated conclusively that they are generically identical with the numerous American pit-vipers of a similar head scutellation, and which are usually known as *Trigonocephalus* or *Bothrops*. The South American genus *Lachesis* is sufficiently characterized by the peculiar scutellation of the tail.

It is not even certain that all the snakes here enumerated as *Trimeresurus* really belong to the same genus. There are various indications that *T. gramineus* represents a group of species which might profitably be separated from the others. As a matter of fact, until the cranial structure of all the various forms which make up the bulk of the *Crotalidæ* is known, generic combinations must be very uncertain in this family.

KEY TO THE SPECIES OF TRIMERESURUS OCCURRING IN FORMOSA AND THE RIU KIU.

- α^1 A squarish loreal shield between nasal and upper preocular; nasal divided or semi-divided; tail normal; color brownish.
- b^1 A single elongate subocular; more than 175 ventrals; more than 65 subcaudals.
- c^1 Less than 30 scale rows.
- d^1 More than 195 ventrals.....*T. mucrosquamatus*, p. 467.
- d^2 Less than 195 ventrals.....*T. elegans*, p. 470.
- c^2 More than 30 scale rows.....*T. flavoviridis*, p. 475.
- b^2 A series of several small suboculars; less than 155 ventrals; less than 60 subcaudals.....*T. okinavensis*, p. 479.
- α^2 An elongate canthal shield between nasal and upper preocular; nasal flat, undivided; tail prehensile; color green.....*T. gramineus*, p. 480.

The five forms here recognized occur within our territory only in Formosa and the Riu Kiu Archipelago. *T. gramineus* extends from the eastern Himalayas to Formosa, but has not as yet been found in the islands to the northeast of the latter. It represents a group of species not very closely allied to the other four. These fall again into two well-defined groups, viz. *T. okinavensis* on the one hand, and *T. mucrosquamatus*, *elegans*, and *flavoviridis* on the other. The last three are plainly of common origin, and *T. mucrosquamatus*, which is supposed to occur in Assam as well as in Formosa, may consequently be regarded as the mother species,^a *T. elegans* having developed from it on the southern group of the Riu Kiu by a decrease in the number of scales and scutes, while *T. flavoviridis* which became isolated in the middle and northern groups developed in the other direction, viz, by an increase in scale rows. *T. okinavensis*, finally, has thus far only been found in the central and northern groups, and no near relative is known either from the southern Riu Kiu or from

^a See pp. 467-468 for a discussion of this species and its relation to *T. jerdonii*.

Formosa. Nevertheless, it is possible to trace the origin of this form also to the eastern Himalayan region, as its nearest and apparently quite close relative is *T. monticola* which ascends to 8,000 feet in the Himalayas and extends east into China at least as far as the province of Szechuen. That this species, or closely allied forms, have not as yet been found in Formosa and the intervening territory is not so surprising when we consider how little we know of the fauna of this region, and I have but little doubt that such a link connecting the habitat of *T. monticola* and *T. okinavensis* will be discovered some day.

Reverting to the *T. mucrosquamatus-elegans-flavoviridis* group, it is evident that while *T. elegans* is geographically intermediate it is not so located systematically and phylogenetically, but that this form and *T. flavoviridis* have developed independently and in several respects in opposite directions—from *T. mucrosquamatus*—in other words, that in extending eastward and northward after having reached Formosa, the ancestor of *T. flavoviridis* did not invade the central Riu Kiu group (Okinawa-Oshima) by way of the southern (Yaeyama) group. The difference in the geological structure of these two groups, as demonstrated by Doctor Yoshiwara recently, probably accounts for this apparent incongruity, as their connection with Formosa may have occurred at different geological periods.

TRIMERESURUS MUCROSQUAMATUS^a (Cantor).

1839. ?*Trigonocephalus mucrosquamatus* CANTOR, Proc. Zool. Soc. London, 1839, p. 32 (type-locality, Naga Hills, Assam).—*Trimeresurus mucrosquamatus* GUENTHER, Rept. Brit. India, 1864, p. 390 (type lost).
1870. *Trimeresurus mucrosquamatus* SWINHOE, Proc. Zool. Soc. London, 1870, p. 411, pl. XXXI (Formosa).—FISCHER, Abh. Naturw. Ver. Hamburg, IX, 1886, No. 6, p. 18 (South Formosa); Jahrb. Wiss. Anst. Hamburg, V, 1888, p. 21 (South Formosa).—BOULENGER, Fauna Brit. India, Rept., 1890, p. 428 (Assam; Formosa).—STEJNEGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 225 (Taipa, Formosa).—*Lachesis mucrosquamatus* BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 552 (Formosa).—WALL, Proc. Zool. Soc. London, 1903, p. 99 (Formosa).

Cantor's original specimen which came from Naga Hills, Assam,^b has been lost, and no specimens from that locality have been recorded since. As the description is exceedingly brief it is consequently not absolutely certain that the Formosan specimens have been correctly identified as *T. mucrosquamatus*. However, the number of ventrals and subcaudals, respectively 218 and 91, agrees very well, and in view of the close relationship of many Formosan snakes with those

^aSignifying *mucro*, a sharp point, *squamatus*, scaly; hence, with pointed scales.

^bBoettger (Offenbach. Ver. Naturk. 26-28 Ber., 1888, p. 153) suggests a confusion with "Nafa Hills," Okinawa shima, in which case Cantor's name would antedate Hallowell's *T. flavoviridis*. There is not the slightest probability, however, that Cantor had any Riu Kiu specimen before him.

of the Himalayan region it would be too rash to deny the possibility of their identity.

Nevertheless, I wish to put on record my suspicion that the Naga Hills and the Formosa specimens represent separate local developments of *Trimeresurus jerdonii* of Guenther,^a also from Assam, but occurring all the way from Tibet to Ichang on the upper Yangtse-kiang. The scale formula of Cantor's specimen, the type of *T. mucrosquamatus*, precludes its being identical with *T. jerdonii*, but the latter comes pretty close to *T. elegans*, from the southern Riu Kius, with which it also agrees very well in coloration.

Description.—*Adult male*; Science College Museum, Tokyo, No. 6; Taipa, Formosa; October, 1897; T. Tada, collector. Rostral somewhat broader than high, barely visible from above; three small scales behind the truncate upper edge of the rostral between the anterior nasals; six scales, including the turned-over upper anterior corner of anterior nasal, covering the canthus rostralis, the one adjoining the latter being elongate and largest; supraoculars rather narrow, about one-third as wide as interocular space; 15 scales in a line between supraoculars; nostril in the posterior margin of the anterior nasal which is much larger than the posterior, its upper anterior corner being turned over on the canthus rostralis; two small loreals, one behind the other, on a line with and of the same width as the upper preocular; two long preoculars, of which the lower is much narrower than the upper, but as long as the latter and posterior loreal together, and forming the upper edge of the pit; three postoculars, including a long, narrow, crescentic subocular which anteriorly is separated by one scale from the subfoveal shield; subocular separated from third supralabial by one scale and from fourth by two series of scales; temporals numerous, the upper smaller, scale-like, convex, the lower somewhat larger, flat; 9 supralabials (on left side only 8 by fusion of fifth and sixth), second forming the anterior border of pit, and with third, fourth, and fifth the largest supralabials, sixth to tenth suddenly much smaller, not larger than the adjacent temporals; first supralabial which is separated from anterior loreal by two scales is of the same size as the last ones; two lower labials only in contact with anterior chin-shields, third and following ones being separated from them by scales; posterior chin-shields scarcely differentiated and separated by a pair of scales; 27 rows of pointed scales, all strongly keeled and with two faint apical pores, except the outer one which is only feebly keeled at base; 206 ventrals; anal entire; 95 pairs of subcaudals. Color (in alcohol) above drab with a median series of irregular transverse, dark brown, black-rimmed and very narrowly

^a Proc. Zool. Soc. London, 1875, p. 233, pl. xxxiv (type-locality, Khasi Hills, Assam; types in Brit. Mus.; Dr. T. C. Jerdon, collector).

white-edged spots, consisting of two halves, one on each side of median line and not always evenly joined, in which case sometimes a zigzag band is forming by connection with succeeding spots; directly under each dorsal spot a smaller lateral spot of the same color, these lateral spots disappearing on sides of neck and tail; top of head with a distinct pattern of dark brown lines diverging from snout backward and forming a lozenge-shaped figure on occiput in the center of which an elongate dark line; a narrow dark line from upper corner of the crescentic subocular over the temporal region to beyond the angle of mouth passing above it on the scale above the last supralabial; lips uniform pale; underside pale drab with several irregular series of large roundish, dark-edged whitish spots, often transversely oblong, and not extending over more than one shield.

Dimensions.

	<i>mm.</i>
Total length.....	780
Snout to vent.....	588
Vent to tip of tail.....	192

This snake undoubtedly grows to be more than 1 meter long.

The *young* (Sci. Coll. Mus. Tokyo, No. 18) is colored essentially as the adult, but the contrast between ground-color and the dark spots is greater, and on the former, between the latter, there is on the sides a shadowy pale brownish gray vertical line; the sides of head, including labials, are not pale, but dark brown, especially a broad vertical band on second supralabial, though not as dark as the postocular stripe which is bounded below by a whitish line and several of the labials and temporals are bordered behind by a narrow whitish mark.

Structurally the only difference from the adults consists in the upturned outer edge of the supraocular.

Variation.—The number of scale rows in this form is fairly constant, nine out of eleven specimens having 27 and only two 25 rows; in the same number of specimens the ventrals vary only between 200 and 216, and the subcaudals between 76 and 95. The supralabials are more variable, namely, 8 to 11. U.S.N.M. No. 36515 has 10 on each side, a small labial having been interpolated between third and fourth.

Habitat.—*T. mucrosquamatus*, as here understood, is only known with absolute certainty from Formosa, where it seems to be fairly common in the lowlands, since nearly all who have collected there obtained specimens. It is assumed that it is the same form which was originally described by Cantor, in 1839, from the Naga Hills in Assam, but it has not been collected there since that time, nor has it been found in the intermediate territory of the Chinese mainland.

It is unquestionably related, however, to a species which occurs from Assam and Tibet as far east as Ichang on the Yangtse, viz,

T. jerdonii. It is at least equally closely allied to *T. elegans* on the islands of the southern Riu Kiu group, while *T. flavoviridis* from the Okinawa-Oshima group is slightly more differentiated.

List of specimens of *Trimeresurus mucrosquamatus*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scafe rows.	Ventrals.	Anal.	Subcaudals.	Interorbital.	Supraorbital.
U. S. N. M.	36515	Male a.	Taipei, Formosa	1897, Nov.—	T. Tada	27	205	1	91	15	10
Sci. Coll. Tokyo.	6	do. b	do.	Oct. —	do.	27	206	1	95	15	8-9
Do.	18	Young	do.	Oct. —	do.	27	202	1	78	14	8
Oldenburg.	a	Adult c.	South Formosa.	1888.	Ruhstrat.	27	216		90		
Do.	b	do. c.	do.	1888.	do.	27	216		90		
Do.	c	Young c	do.	1886.	do.	27	214		81		
Brit. Mus.	a	Male d.	Takow, Formosa		R. Swinhoe.	25	200	1	92		
Do.	b	Female d	Formosa		do.	27	214	1	76		
Do.	c	Young d	Central Formosa.		Holst.	27	211	1	89		
Do.	d	do. d.	South Formosa		J. G. Fischer.	27	210	1	90		
Do.	g	Male c.	South Cape, Formosa.		J. D. La Touche.	25	211	1	91		

a Total length, 932 mm.

b Description, p. 468.

c Fischer, p. 18.

d Boulenger, Cat. III, p. 552.

e Boulenger, MS.

TRIMERESURUS ELEGANS^a (Gray).

1849. *Craspedocephalus elegans* GRAY, Cat. Sn. Brit. Mus., p. 7 (type-locality, "west coast of (North?) America?"; types, Brit. Mus., Nos. 47, 3, 4, 62 and 64; Sir E. Belcher, collector) (not *Trimeresurus elegans* Gray, 1853).
1895. *Trimeresurus luteus* BOETTGER, Zool. Anz., XVIII, July 8, 1895, p. 269 (type-locality, "Miyakoshima," Riu Kius; type in Mus. Senckenberg.; Schmaeker, collector); Offenbach, Ver. Naturk., 33-36 Ber., 1895, p. 111 ("Miyakoshima.")—*Lachesis luteus* BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 553.—*Lachesis lutea* BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 139 ("Miyakoshima.")
1906. *Lachesis mucrosquamatus* WALL, Proc. Zool. Soc. London, 1905, II, p. 516 (Miyako; Iriomote) (not of Cantor).

The types of *Craspedocephalus elegans* in British Museum are enumerated under *Lachesis mucrosquamatus* by Boulenger in his catalogue, but their lower number of ventrals precludes their annexation to that species. Boulenger does not seem to have known *T. luteus* from autopsy, and in his key to the species on page 532 he fails to point out any difference between the two species. The chief difference shown in his descriptions (pp. 552 and 553) consists in the "temporal scales smooth" in *T. mucrosquamatus* and "temporal scales keeled" in *T. luteus*, but this character is not reliable.

^a Signifying elegant.

The locality whence Captain Belcher obtained the specimens upon which Gray established the new species is not known. Gray suggested "west coast of (North?) America?" By a study of Belcher's itineraries both in the *Sulphur* and in the *Samarang* I have come to the conclusion that the only likely place where he can have obtained the specimens in question is Ishigaki shima, an island in the southern group of the Riu Kiu, where he stayed some time and undertook considerable surveying on shore^a, the very same island whence come two of the specimens of this species examined by me. In fact I have but little doubt that Ishigaki shima is the true type-locality.

With regard to the probability of this island also being the type-locality of *T. luteus*, see remarks farther on under *habitat* of this present species (p. 474).

Thanks to the courtesy of Doctor Boulenger, I have been able to examine Gray's types of *C. elegans*. They consist of two young specimens, No. 47.3.4.62, with the upper edge of snout and supraoculars turned up, and with the temporals slightly but distinctly keeled, and No. 47.3.4.64, an adult male, with temporals scarcely keeled. There is, therefore, no doubt in my mind that Boettger's *T. luteus* is the same.

T. elegans is very closely allied to the Formosan *T. mucrosquamatus*, and while there are various average characters which separate the two, the number of ventrals is the only one which seems to offer an absolutely reliable diagnosis. In all the specimens of *T. elegans* examined by me the supraoculars have the outer edge sharply turned up, but in the only small Formosan specimen before me (Sci. Coll. No. 18) this edge is similarly turned up. Neither the keeling of the temporals, nor the size of the canthal scales offer any stable characteristics.

In the three specimens of each species now before me—the only ones I can now examine with regard to this character—I find another difference, viz, that in the three Formosan *T. mucrosquamatus* there are two loreals, one behind the other, followed by a long horizontal upper preocular, while in the three Riu Kiu *T. elegans* there is only a single loreal between the nasal and the elongate preocular.


Additional characters may be found in the lower number of supralabials, the somewhat wider supraoculars, and resultant narrower interorbital space in *T. elegans* as well as in the lower number of scale rows, characters, however, which intergrade and offer no hard and fast line of separation. The coloration is essentially the same, with a

^aThe "examination of Pa-tchung-san" [Ishigaki shima] "occupied us twenty-one days," Belcher, Narrative Voy. Samarang, I, 1848, p. 75.

See also Doctor Adams's reference to a species of "*Trigonocephalus*" in the Meiacosima group in his "Notes from a Journal of Research into the Natural History of the Countries visited during the Voyage of H. M. S. *Samarang* under the command of Capt. Sir E. Belcher, C. B.," in Belcher's Narrative Voy. Samarang, II, 1848, pp. 305 and 306. This may be an allusion, however, to *Agkistrodon affinis*.

tendency, however, to a less definite arrangement of light and dark spots on the underside and to a more or less complete obliteration of the lateral spots.

Description (figs. 366-368).—*Young*; Science College Museum, Tokyo, No. 19; Iriomote shima; 1887; Tashiro, collector. Rostral broader than high, barely visible from above, bordered behind by two scales which separate the anterior nasals, the latter not turned over the canthus rostralis to the upper surface of the snout; two larger scales on canthus rostralis, the anterior separated from the rostral and from the posterior one respectively by a small scale and the latter by a similar scale from the supraocular, the outer edges of these scales being raised like that of the supraoculars; upper head scales small, smooth, 13 in a row between supraoculars, which are somewhat more than one-third as wide as interorbital space; posterior nasal smaller than anterior, followed by a single squarish loreal, behind which an equally broad, elongate upper preocular; a second, much narrower and somewhat longer preocular forms the upper edge of the pit; three postoculars,



FIGS. 366-368.—*TRIMERESURUS ELEGANS*. NAT. SIZE. 366, TOP OF HEAD; 367, SIDE OF HEAD; 368, UNDERSIDE OF HEAD. NO. C 40, HIGH SCHOOL, KUMAMOTO.

including the long, narrow crescentic subocular which reaches the subfoveal, and is separated from fourth supralabial by two rows of scales; upper temporals weakly keeled, lower ones smooth, those bordering the supralabials larger than the latter; 7 supralabials, first small; second forming the anterior border of the pit, third largest, fourth long, but perceptibly lower than third, and not much higher than those following; three lower labials in contact with anterior chin-shields, which are larger than the posterior ones; scales pointed, in 23 rows, all strongly keeled, except outer row which is smooth; 188 ventrals; anal entire; 72 pairs of subcaudals. Color (in alcohol) pale brownish gray above; a series of darker brownish blotches, with paler centers, along the middle of the back, the respective halves of each blotch on each side of the median line not always fitting together, but sometimes even completely alternating and thus forming in places an interrupted zigzag band; tip of tail uniform light-colored (yellow?); on the sides obscure traces of a dark spot below each dorsal blotch; a somewhat ill-defined

brownish band on top of head covering the two scale rows nearest to the supraoculars and extending backward to the neck; a well-defined dark-brown band or line, one scale wide, starting from the posterior margin of the orbit below the center of the eye and extending backward beyond the angle of the mouth passing the latter on the scale row above the last supralabial; a dark brown vertical line on the posterior half of the second supralabial from the pit to the edge of the lip, lower surface pale buff with ill-defined markings of pale brownish gray.

Dimensions.

	<i>mm.</i>
Total length.....	370
Snout to vent.....	305
Vent to tip of tail.....	65

In the *adult male* from Ishigaki shima the coloration is essentially as described above, but the lateral series of spots, one below each dorsal blotch, is as well marked and dark colored as the latter; tip of tail is without dark cross-bars but is not paler than the basal portion. This specimen (Sci. Coll. Mus. Tokyo, No. 18) measures 655 mm. in total length, of which the tail is 123 mm. This is considerable less than the length reached by the species, as one of the types, according to Doctor Boettger, measures 945 mm.

The edge of the supraocular is turned up in all the specimens, but according to my observations on the specimens in the Hamburg Museum less so in the old specimens than in the young ones.

Variation.—The present species is rather variable so far as the scutellation of the head is concerned. In Boettger's types of *T. luteus*, which he kindly allowed me to examine in 1898, the scales covering the canthus rostralis are small and irregular. Of four specimens in the Hamburg Museum two have one large canthal scale followed by three small irregular ones, while two have two large scales followed by a small one, and of the three specimens before me two have one and one two enlarged canthal scales. In his types, the temporal scales are rather strongly keeled; in the Hamburg specimens three of the specimens have the upper temporals more or less strongly keeled, in the fourth one the keels are only faintly indicated, and so they are in my specimens. In one of the types of *T. elegans* the temporals are slightly, but distinctly keeled, as already stated. Usually there are two rows of scales between the subocular and the labials, in one Hamburg specimen (No. 2582) only one such row. The number of scales between the supraoculars varies between 10 and 13. Supralabials probably most commonly 7, but one of my specimens has 8 on both sides, and two have 7 on one side, 8 on the other. In the specimen in the Kumamoto Fifth Higher Middle School the long upper preocular is divided vertically near the eye on both sides.

Of 13 specimens eight have 25 scale rows and five have 23. Doctor Wall records one with 24 scale rows. Ventrals vary between 183 and 191, subcaudals between 66 and 77.

The variation in color seems to be less great and consists in the more or less perfect obliteration of the lateral series of spots, and in the greater or less definite outline of the light and dark spots on the underside.

Remarks by collector.—Mr. Tashiro has furnished the following note with the larger specimen collected by him on Ishigaki shima:

"This species is not found on Okinawa shima, but is confined to the Yaeyama Islands. It is known by the name of 'Habu,' the general name applied to the snakes of this genus, of which there are about five varieties, each with a special name. It lives in crevices in stone-walls, hollow trees, etc. Its habits are more sluggish than the real 'habu' (*Trimeresurus flavoviridis*). The natives say that it is less poisonous than the latter, yet I think it is by no means inferior in this respect."

Habitat.—This species is confined to the southern group of the Riu Kius, and seems to be fairly common both on Iriomote shima and on Ishigaki shima. Boettger's types of *T. luteus* are said to have come from "Yaeyama auf Mijako shima," a somewhat confusing statement which might have been suspected of meaning "Miyako shima of the Yaeyama group," as formerly at least that island was considered part of the group, though now excluded as a special subgroup, but in the introduction to his paper^a Doctor Boettger says:

"Aber die Thiere scheinen leider nicht genau nach den Fundorten getrennt worden zu sein * * * Die Schlangen sollen zum grössten Theil von Yaeyama oder Patchungsan, einer der Miyako shima Inseln, herstammen."

But Patchungsan is the Chinese name for Ishigaki shima, and Doctor Boettger evidently uses Miyako shima as the collective name of the two southern subgroups. It is therefore exceedingly dubious if the types came from Miyako shima proper, and the probability is that they really originate from Ishigaki shima. That the same species occurs on Miyako shima has now been shown by Doctor Wall who records three specimens from that island under the name of *Lachesis mucrosquamatus*.

^a Offenbach. Ver. Naturk. 33-36 Ber., 1895. p. 101.

List of specimens of *Trimeresurus elegans*.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Interorbitals.	Supraorbitals.
Sci. Coll. Tokyo.	19	Young ^a	Iriomote shima.	1887.	Tashiro.	23	188	1	72	13	7
Do.	18	Male	Ishigaki shima.	1887.	do.	25	183	1	72	12	8
Kumamoto	C(40)	Young	do.	1891	do.	25	185	1	76	10	7
Hamburg	2578a	do	Iriomote shima	1897	Doctor Lenz.	23	187	1	72	11	7-8
Do.	2578b	do	do	1897	do.	25					7-8
Do.	2582	Female	do	1897.	do.	23	191	1		11	7
Brit. Mus.	47.3.4.64	Male ^b	Ishigaki shima")	Mar.13	Capt. E. Becher.	25	187	1	76		
Do.	47.3.4.62	Young ^b	do		do.	25	183	1	72		
Senckenberg.	9553a	Adult ^c	"Miyako shima."		B. Schmaeker.	23	182	1	72		
Do.		do. ^c	do		do.	25	186	1	74	12-13	7-8

^a Description, p. 472.

^b Types of *T. elegans*.

^c Types of *T. luteus*.

TRIMERESURUS FLAVOVIDRIS ^a (Hallowell).

HABU.

Plate XXVII.

1860. *Bothrops flavoviridis* HALLOWELL, Proc. Phila. Acad., 1860, p. 492 (type-locality, "Amakarima" Island = Kerama shima, middle group, Riu Kiu).—BOULENGER, Proc. Zool. Soc. London, 1887, p. 149 ("Loo Choo" Islands).—? OKADA, Cat. Vert. Japan, 1891, p. 70 (Okinawa shima).—? FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 860; author's separate, p. 11 (Okinawa).—*Trimeresurus flavoviridis* BOULENGER, Fauna Brit. India, Rept., 1890, p. 425 (Loo Choo).—*Lachesis flavoviridis* BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 550 (Okinawa).—BOETTGER, Kat. Schl. Mus. Senckenberg., 1898, p. 139 (O-shima, "Liu Kiu gruppe").—BROWN, Proc. Phila. Acad., 1902, June 11, p. 185 ("Loo Choo Islands").—SCHENKEL, Verh. Naturf. Ges. Basel, XIII, Pt. 1, —, p. 179 (Okinawa).—WALL, Proc. Zool. Soc. London, 1903, p. 102 (Loo Choo); 1905, II, p. 516 (Amami; Okinawa).

1880. *Trimeresurus riukiuanus* HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 118, pl. —, figs. 6 10 ^b (type-locality, District Nase, O-shima; types, Nos. 9767-9768, Berlin Mus.).—DOEDERLEIN, Mith. Deutsch. Ges. Ost-Asiens, III, 1881, p. 149 (Amami-o-shima).—LANDOIS, Westfal. Prov. Ver. 16 Ber., 1887 (p. 45).—FISCHER, Jahrb. Wiss. Anst. Hamburg, V, 1888, p. 20 (Okinawa shima).—OKADA, Cat. Vert. Japan, 1891, p. 70 (Okinawa; O-shima).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 860; author's separate p. 11 (Amami-o-shima; Okinawa shima).

The type of Hallowell's *Bothrops flavoviridis* which, according to his statement, was much mutilated (only 61 ventrals could be counted)

^a Signifying yellowish green.

^b Reproduced in this work on Plate XXVII.

has apparently been lost. From Doctor Stimpson's MS. catalogue it is learned that the "viper," as he calls it, was "killed on one of the Amakirrima Isles (Loo Choo) by Mr. Macomb of the *Hancock*, April 1855," consequently on one of the small islands composing the Kerama shima group just west of the south end of Okinawa shima.

Hilgendorf's *Trimeresurus riukiuanus* on the other hand was described from specimens obtained in the Nase district on Amami-o-shima, the principal island of the northern group. The scale formulas of the specimens from both groups agree very well, and a direct comparison of a few specimens does not reveal any essential difference. The material at hand, viz, one from the northern group and two from the middle group, is not conclusive, however. The former seems to have a somewhat longer snout with more, or smaller, scales between the supraoculars, but until verified in large series from both groups the two names must be regarded as synonymous. The low number of scale-rows assigned to his specimen by Hallowell, viz, 31, is evidently due to the fact that the specimen was mutilated.

Okada in his List of the Vertebrates of Japan enumerated both *T. flavoviridis* and *T. riukiuanus* as separate species, though possibly he included the former with *T. okinawensis* which was not then described. Doctor Fritze, however, enumerates all three from Okinawa. Inasmuch as he seems actually to have examined specimens of two species it is probable that the one from Tokushimura was really a *T. okinawensis* although enumerated as a *T. flavoviridis*.

Description (fig. 369).—*Half-grown female*; U.S.N.M. No. 31818; Amami-o-shima, Riu Kiu; Dr. H. M. Smith collection. Rostral






FIG. 369. — TRIMERESURUS FLAVOVIDIDIS. NAT. SIZE. SIDE OF HEAD. No. 16. SCI. COLL. TOKYO.


almost as high as broad, nearly triangular, the sutures with first supralabials and internasals being very short, just visible from above; a single small scale behind the rostral separating the anterior nostrils; four or five shields, including the anterior nostrils, forming the raised canthus rostralis between rostral and supraocular, the shield second from the supraocular being the largest; supraocular long, pointed anteriorly, half as wide as interocular space; upper head scales smooth, small, about 13 in a line between the supraoculars; nostril round, in the posterior margin of the anterior nasal which is turned over the canthus rostralis to the upper side of the head; loreal nearly rectangular, slightly longer than high; two preoculars, very elongated, both in contact with loreal, the upper nearly twice as long as the latter, the lower, which forms the upper posterior border of the pit, much narrower than the upper; a long narrow shield bordering the pit below, separated from long subocular by a couple of small scales; eye rather large, vertical diameter equaling its distance from edge of lip; a very long narrow crescentic subocular




1



2



3



4

TRIMERESURUS FLAVOVIRIDIS. (FROM HILGENDORF.)

FOR EXPLANATION OF PLATE SEE PAGE 555.

separated from the lower preocular and from the subfoveal by several small scales; from the fourth supralabial by two rows of scales, and from the temporals by a single row of smaller scales; one postocular on left side, three on right, in addition to suboculars; lower temporals large, upper ones small, smooth scales; 8 supralabials, first subtriangular, second forming the anterior border of the pit, third largest, fifth to eighth small, low; three lower labials in contact with anterior chin-shields, fourth and fifth being separated from them by scales; anterior pair of chin-shields much larger than posterior, which are scarcely differentiated from the other postgenials and separated by a pair of smaller scales; 35 rows of narrow pointed scales, of which all except the outer are keeled and provided with two apical pits; 226 ventrals; anal undivided; tail defective, subcaudals 73+, very few missing. Color (in alcohol) above very pale isabella-color, on each side of the median line with a series of elongated darker brownish spots, which are nearly in pairs anteriorly but alternating on the posterior two-thirds, mostly anastomosing across the median line; on the sides a similar but very obscure series; on the upper side of the tail the spots become united to regular cross-bars; occiput with a very obscure pattern of several elongate divergent spots; sides of head whitish with a narrow brown line from upper corner of subocular across the temples to the sides of neck, one scale above last supralabial; all the brown markings are dark-edged with paler centers; underside uniform whitish.

Dimensions.

	<i>mm.</i>
Total length.....	612
Snout to vent.....	517
Vent to tip of tail.....	95

This species grows to a considerable size. Hilgendorf's types of *T. riukiuanus* are 1.380 and 1.555 m., and Doctor Fischer records two specimens measuring 1.433 and 1.285 m., respectively. A snake of this kind over 5 feet long must be a pretty dangerous reptile.

Variation.—The number of scale rows shows considerable variation, viz, from 33 to 37, though 35 seems to be the normal number, inasmuch as in 13 specimens more than two-thirds have this number. Doctor Wall records a specimen with 40 scale rows. Ventrals vary between 222 and 234, subcaudals between 75 and 90. Supralabials usually 8, but a specimen in the Hamburg Museum has 7 on one side, and other specimens are on record as having 9.

The size of the small scales on top of the head is not so variable as the figures given for the number of scales on a line between the supraoculars, namely, from 7 to 13 would indicate, since the discrepancy depends chiefly upon the irregular size of these scales on the interocular region.

The two specimens from Okinawa before me are smaller than the one described above and show a more definite color pattern. The lateral markings are so light as to be almost lost while the dorsal blotches are shorter, with a greater tendency to combine into median transverse marks across the middle line, thus forming angular cross-bars deeply notched in front and behind and with a large pale spot on each side of the median line. The pattern on the top of the head is also very sharply defined and formed as shown in Hilgendorf's figure of his type of *T. riukiuanus* (Plate XXVII, fig. 1 of this work).

Habitat.—This dangerous snake is confined to the islands of the Okinawa and Ōshima subgroups of the Riu Kiu Archipelago, in some of which it appears to be rather common. Amami-ōshima, and—according to Doederlein, also Kome shima, are particularly notorious for the occurrence of the "habu," while both he and Fritze assert that it is much less common in Okinawa shima.

It is represented by closely allied species in the southern group and in Formosa.

List of specimens of Trimeresurus flavoviridis.

Museum.	No.	Sex and age.	Locality.	When collected.	By whom collected or from received.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Interorbitals.	Supraorbitals.
		(a)	"Amakarima Is-land."	1855. April	Mr. Maccomb.	31			81		8
U.S.N.M.	31818	Female ^b	Amami-ōshima.		Dr. H. M. Smith.	35	226	1		13	8
Sci. Coll. Tokyo.	16	Young ^c	Oshiro mura, Okinawa shima.	1887. May 10	Tashiro.	35	222	1	87	9	9
Imp. Mus. Tokyo.	9	do	Okinawa shima			35	223	1	86	11	8
Hamburg.	1540	do	do		Doctor Warburg.	35	227	1	80	9	7-8
Brit. Mus.	a	Female ^d	do		Dr. J. G. Fischer.	35	225	1	80		
Do.	b	do, d	do			37	225	1	76		
Do.	c	Male ^e	do		Mr. Holst.	35	231	1	90		
Do.	d	Young ^f	do		do	33	229	1	79		
Do.	e	Male ^g	"Loo Choo Is."		H. Pryer.	33	223	1	86		
Do.	f	Young ^g	do		do	35	222	1	83		
Do.	g	do, c	do		do	35	230	1	75	7-8	8
Berlin.	9767	Adult ^h	Nase, Amami-ōshima.		Doctor Hilgendorf.	35	226	1	84		8
Do.	9768	do, t	do		do	37	228	1	86		8

^a Type, specimen mutilated, the number of scale rows given is therefore probably not correct.

^b Description, p. 476.

^c Fig. 3-9.

^d Boulenger, Cat. III, p. 570.

^e Boulenger, Proc. Zool. Soc., 1887, p. 149.

^f Types of *T. riukiuanus*, Hilgendorf, p. 118. The scale formulas as here given for each individual may not be exactly correct, as Hilgendorf's statement is not quite explicit, however, they are correct for the two specimens together.

TRIMERESURUS OKINAVENSIS^a Boulenger.

KUFUHA (native name in OKINAWA SHIMA according to TASHIRO).

1892. *Trimeresurus okinavensis* BOULENGER. Ann. Mag. Nat. Hist. (6), X, Oct. 1892, p. 302 (type-locality, Okinawa shima; types in Brit. Mus.; Holst, collector).—FRITZE. Zool. Jahrb. Syst., VII, 1894, p. 861; author's separate, p. 12 (Okinawa).—*Lachesis okinavensis* BOULENGER, Cat. Sn. Brit. Mus., III, 1896, p. 549, pl. xxv, fig. 2 (Okinawa).—WALL, Proc. Zool. Soc. London, 1905, II, p. 516 (Okinawa; Amami; Yaku).
1894. ? *Bothrops flavoviridis* FRITZE. Zool. Jahrb. Syst., VII, p. 860; author's separate, p. 11 (Tokuchimura, Okinawa) (not of Hallowell?).

Description.—*Adult female*: Science College Museum, Tokyo, No. 17; Okinawa shima; March, 1887; Tashiro, collector. Rostral nearly as high as broad, scarcely visible from above; behind the truncated upper edge three scales of the same size as the other upper head scales in a line between the anterior nasals and the first canthal scale; four canthal scales between anterior nasal and supraocular decreasing in size posteriorly; supraoculars wider than one-third the interocular space; 8 scales on a line between the supraoculars; head scales flat, smooth anteriorly, faintly tubercular or keeled on parietal and occipital regions; nostril large, rounded, in posterior margin of anterior nasal which is much larger than posterior, its upper anterior edge just visible from above, but not turned over the canthal edge; a small loreal, nearly twice as long as high below second and third canthal scales, in contact behind with the two preoculars, of which the upper is twice as wide as the lower; pit very large; subfoveal shield entering eye below preoculars; eye rather small; three small suboculars separated from supralabials by two scale rows; two postoculars; temporal scales numerous, the upper ones small, keeled, the lower series next to the supralabials larger, smooth; 8 supralabials, first small, triangular, separated from loreal by one scale, second broad, forming anterior border of pit, third largest; two lower labials in contact with anterior chin-shield; posterior chin-shield scarcely differentiated; 23 rows of rather obtuse scales with obscure apical pits, all keeled, except outer row, but keels not reaching the tip of the scales; 128 ventrals; anal entire; 51 pairs of subcaudals. Color (in alcohol) above brownish gray, with large transverse blotches of brown anteriorly and posteriorly narrowly edged with dusky; a series of lateral spots on the middle of the sides, each one below the corresponding dorsal blotch of which it is a detached portion and with which it is not alternating; below and alternating with these another row of dark brown spots situated on the two outer scale rows and the adjacent part of the ventrals; a broad dark brown, black-edged band three scales wide from eye over temporal region to posterior end of lower jaw, its anterior border crossing the last supralabial at the corner of the mouth; above this a narrower whitish band, and a similar one below it on the

^a From Okinawa.

last four supralabials, crossing to the lower jaw; sides of face anterior to this brown with two darker blotches across both lips at the level of second and fourth supralabial; on the lower lip a third similar dark brown blotch near the corner of the mouth; underside pale obscurely clouded with brownish.

Dimensions.

	mm.
Total length.....	445
Snout to vent.....	371
vent to tip of tail.....	74

Variation.—Supralabials vary between 8 and 7, ventrals between 127 and 134, subcaudals between 42 and 51 pairs in the three specimens which have thus far been recorded. Doctor Wall reports one specimen with 24 scale rows.

Habitat.—The few specimens known of this species have nearly all come from Okinawa shima. According to a recent paper by Doctor Wall the collector of Mr. Owston obtained not only four specimens in Okinawa shima, but also three in Amami-o-shima, and one in Yaku shima. The latter occurrence is so extraordinary, however, that it would be well to await corroborative evidence, as some mistake in the labeling may be possible.

As I have stated elsewhere, there is no near relative of this species known from Formosa, but as it is allied to the Himalayo-Chinese *Trimeresurus monticola*, the latter or a related form may be expected to occur in that island.

List of specimens of Trimeresurus okinawensis.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Interorbital.	Supranal.
				1887.							
Sci. Coll. Tokyo...	17	Female ^a	Okinawa shima.	Mar.	Tashiro.....	23	128	1	51	8	8
Brit. Mus.....	a	do. ^b	do.....		P. V. Holst....	23	130	1	43	9	8
Do.....	b	do. ^b	do.....		do.....	21	129	1	47	6	7

^a Description, p. 479.

^b Types; Boulenger, Cat. III, p. 549


TRIMERESURUS GRAMINEUS^a (Shaw).

1802. *Coluber gramineus* SHAW, Gen. Zool., III, Pt. 2, p. 420 (type-locality, Vizagapatam, India; based on Russell's Ind. Serp., I, pl. IX).—*Trimeresurus gramineus* GUENTHER, Rept. Brit. India, 1864, p. 385. BOETTGER, Ber. Senckenberg. Naturf. Ges., 1888, Abh. p. 188 (South Formosa); 1894, p. 135 (Hainan).—BOULENGER, Fauna Brit. India, Rept., 1890, p. 429 (southern China, India, Malay Peninsula and Archipelago, etc.).—STRENGER, Journ. Sci. Coll. Tokyo, XII, Pt. 3, 1898, p. 225 (Taipa, Formosa).—*Lachesis gramineus* BOULENGER, Cat. Sn. Brit. Mus., III, 1896.

^a Signifying of grass, grassy; referring to its green color.

- p. 554 (southeastern Asia; Formosa).—WALL, Proc. Zool. Soc. London, 1903, p. 99 (Hongkong; Formosa).—*Lachesis graminea* BOETGER, Kat. Schl. Mus. Senckenberg., 1898, p. 139 (south Formosa, etc.)
1802. *Coluber viridis* BECHSTEIN, Lacépède's Naturg. Amph., IV, p. 252, pl. xxxix, fig. 1 (type-locality, Vizagapatam, India; based on Russell's Ind. Serp., I, pl. ix).—*Trimeresurus viridis* LACÉPÈDE, Ann. Mus. Paris IV, 1804, p. 209.
1839. *Trigonocephalus erythrurus* CANTOR, Proc. Zool. Soc. London, 1839, p. 31 (type-locality, Ganges Delta, India; type in Brit. Mus.; Cantor, collector).—*Trimeresurus erythrurus* GUENTHER, Rept. Brit. India, 1864, p. 386.—SWINHOE, Proc. Zool. Soc. London, 1870, p. 412 (Takow, Formosa).
1842. *Trimeresurus albolabris* GRAY, Zool. Miscell., (p. 48) (type-locality, China; types in Brit. Mus.; Reeves, collector).
1853. *Trimeresurus elegans* GRAY, Ann. Mag. Nat. Hist. (2), XII, p. 391 (type-locality, Sikkim; type in Brit. Mus.; Hooker, collector) (not *Craspedocephalus elegans* Gray, 1849).

Description (figs. 370–372).—*Adult male*; U.S.N.M. No. 36516; Taihoku (Taipa), Formosa; September, 1897; T. Tada, collector. Rostral as high as broad, very narrow above, nearly triangular



FIGS. 370-372.—TRIMERESURUS GRAMINEUS. $\frac{1}{3}$ \times NAT. SIZE. 370, TOP OF HEAD; 371, SIDE OF HEAD; 372, UNDERSIDE OF HEAD. No. 29 SCI. COLL. TOKYO.

bordered behind by a single scale between the upturned anterior corners of the nasal, just visible from above; canthus rostralis sharp, formed anteriorly by the upturned edge of the nasal, the upturned edge of an elongated shield corresponding to the loreal in the other species here described and the upturned portion of the upper preocular; head shields small, smooth anteriorly, keeled on parietal and occipital regions; supraoculars very narrow, occupying only the outer edge of the supraocular region, their width being scarcely more than one-fifth the distance between them; about 12 scales on a line between the supraoculars; nasal large, smooth, undivided, with a round nostril pierced near the lower edge; behind it above, on the canthal ridge an elongated shield, being the loreal of the other species; below it, separating it from the first supralabial, and between nasal and the upper portion of second supralabial which

enters the pit, two small scales, one above the other, two elongate anterior preoculars, the upper, which is somewhat wider anteriorly turned up over the canthal ridge, the lower forming the upper border of the large pit; the subfoveal as large as the lower preocular, posteriorly entering eye between the latter and the subocular; a long, narrow, crescentic subocular anteriorly in contact with subfoveal, separated from fourth and subsequent supralabials by one and posteriorly two rows of scales; three small postoculars on left side, two on right; temporals numerous, lower ones largest, upper ones smaller, keeled; 10 supralabials, first small, triangular, second very high, forming anterior border of pit, third largest, fourth slightly larger than fifth to tenth, which are subequal; 3 lower labials in contact with anterior chin-shields, posterior chin-shields scarcely differentiated; 21 rows of narrow, pointed, keeled scales without apical pits; 161 ventrals; anal entire; 69 pairs of subcaudals; tip of tail rather blunt. Color (in alcohol) above saturated uniform "parrot-green;" from the subocular, under the center of the eye, a narrow, strongly defined, pale-yellowish line, the lower row of temporals and across the last supralabial to the side of neck and from there to near the tip of the tail on the middle line of the outer scale row, the lower edge of which is somewhat darker than the rest of the body; underside paler green, washed with blue so as to be almost "beryl-green" toward the sides; tips of tail colored like the rest of the body.

Dimensions.

	<i>mm.</i>
Total length.....	618
Snout to vent.....	488
Vent to tip of tail.....	130

The *female* appears to have a relatively much shorter tail. No. 2a, Science College Museum, measuring 525 mm. in total length, has a tail only 92 mm. long.

Variation.—The Formosan specimens examined by me are very uniform, both in scutellation and coloration, as will be seen from the table at the end, but in individuals from all over the wide range of the species there is displayed a corresponding range of variation. Thus, 19 and 23 scale rows are occasionally, though rarely, recorded; supralabials vary between 8 and 12, ventrals between 145 and 175, subcaudals between 53 and 75 pairs. The color variations consist in more or less distinct blackish dorsal cross bands and the absence of the lateral yellow line. The terminal portion of the tail is frequently yellow or reddish, a color phase which until recently was regarded as a distinct species under the name of *Trimeresurus erythrus*.

Habitat.—This species is widely distributed from the Himalayas in the north and west, through India, Indo-China, southern China, to the Malayan peninsula, Sumatra, Java, and Timor in the south, and Formosa in the east.

In Formosa it was first collected by Swinhoe, who found it at Takow, no less than seven specimens from him being in British Museum. Both the Hamburg and the Senckenbergian Museum, in Frankfort on the Main, have specimens from south Formosa, the former (No. 1541) through Doctor Warburg, the latter through Schmacker. Mr. Tada, who collected in Formosa for the Japanese Government in 1896 and 1897, brought four specimens from Taihoku (Taipa), and stated that it is common, especially in northern Formosa.

List of specimens of *Trimeresurus gramineus*.

Museum.	No.	Sex.	Locality.	When collected.	By whom collected.	Scale rows.	Ventrals.	Anal.	Subcaudals.	Interorbitals.	Supraorbitals.
1897.											
U.S.N.M.	36516	Male a	Taipa, Formosa	Sept.	T. Tada	21	161	1	69	12	10
Sci. Coll.	2a	Female b	do	do	do	21	158	1	68	12	9-10
Tokyo.											
Do.	2b	Male	do	do	do	21	158	1	69	11	10
Do.	2c	do	do	do	do	21	159	1	73	11	9-10
Brit. Mus.	z	Male, c	Formosa		R. Swinhoe	21	163	1	67		
Do.	a	do, c	do		do	21	159	1	71		
Do.	β	do, c	do		do	21	156	1	68		
Do.	γ	do, c	do		do	21	162	1	68		
Do.	δ	do, c	do		do	21	155	1	65		
Do.	ε	do, c	do		do	21	152	1	66		
Do.	ζ	do, c	do		do	21	151	1	60		

a Description, p. 481.

b Figs. 370-372.

c Boulenger, Cat. 111, p. 558.

Subclass SYNAPSIDA.

1903. *Synapsida* H. F. OSBORN. Science (n. s.) XVII, Feb. 13, 1903, p. 276.

Order TESTUDINATA.

- 1788. *Testudines* BATSCH. Anleit. Kennzn. Thiere, Mineral., I (p. 437).
- 1802. *Chelonia* MACARTNEY in Ross' Transl. Cuvier's Lect. Comp. Anat., I, tab. III.
- 1804. *Chelonii* LATREILLE. Nouv. Dict. d' Hist. Nat., XXIV, tabl. méth. p. 61.
- 1807. *Cutaphractæ* LINK, Besch. Nat. Samml. Rostock, II, p. 51.
- 1811. *Testudinata* OPPEL. Ordn. Rept., p. 3.
- 1814. *Perostia* RAFINESQUE. Specchio Sci. (Palermo) II, no. 9, 1 Sett., 1814, p. 66.
- 1822. *Chelonca* FLEMING, Philos. Zool., II, p. 268.
- 1825. *Fornicata* HA WORTH. Philos. Mag., 1825, (p. 372).
- 1828. *Sterrichroles* RITGEN, Nova Acta Acad. Leop. Carol., XIV, p. 269.
- 1828. *Chelyna* WAGLER, Isis, 1828, p. 861.
- 1837. *Chelonites* BURMEISTER, Handb. Naturg., p. 730.
- 1839. *Chelonides* SWAINSON, Nat. Hist. Classif. Fish. Amph. Rept., II (Lardner's Cab. Encycl.), p. 112.
- 1845. *Testudinea* GRAVENHORST, Thierwelt, p. 45.
- 1849. *Tylopada* MAYER, Rheinland und Westphal. Verhandl., VI (p. 177).

The proper interrelationship between the various types of living Chelonians has not been finally determined as yet. The greatest stumbling block has been the *Dermochelys*, which some authors, such as Baur and Fuerbringer, regard as a highly specialized form more or less closely allied to the other marine turtles, while others have taken the opposite view, regarding it as one of the most primitive types and so radically different from all the other chelonians as to make it expedient to collect the latter in one great suborder and to keep *Dermochelys* by itself in another. The former view is scarcely tenable, if Goette^a is correct in regarding the bony carapace of the majority of turtles as an integral part of the skeleton as opposed to the dermal origin of the dorsal mosaic of *Dermochelys*. On the other hand, the soft-shelled turtles exhibit so many differences from those with a horny covering that it seems inexpedient to include them in the same category. For these reasons it has been thought best to adhere to the division first proposed by Seeley, in 1882, but to substitute subordinal names for those chosen then by him, these having the form of family names, as understood by most zoologists.

The characters which define the three suborders may be tabulated as follows:

- a*¹ No solid carapace, the vertebrae and ribs being separated from a shell consisting of a mosaic of numerous small polygonal bony plates imbedded in a leathery skin; no descending process of the parietal bone; limbs without claws. . . 1. *ATHECÆ*, p. 485.
- a*² A solid carapace of a few large symmetrical bony plates not separated from the underlying vertebrae and ribs; parietals with descending processes; limbs with at least one claw each.
- b*¹ Body covered with horny scutes arranged differently from the bony plates beneath; epiplastra and hyoplastra in contact, not separated by entoplastron; center of last cervical and first dorsal vertebrae articulating with each other; fourth digit never with more than three phalanges; jaws covered by horny sheaths not concealed under fleshy lips. 2. *LAMINIFERA*, p. 488.
- b*² Body covered by an undivided leathery skin without scutes; epiplastra separated by entoplastron from hyoplastra; last cervical vertebra articulating with first dorsal by zygapophyses only; fourth digit with more than three phalanges; jaws concealed under fleshy lips. 3. *CHILOTEÆ*, p. 513.

The turtles occurring within our limits may be easily referred to their respective families by the following:

ARTIFICIAL KEY TO FAMILIES OF TURTLES.

- a*¹ Limbs clawless; back with five longitudinal keels or ridges (*Athecæ*).
DERMOCHELIDÆ, p. 485
- a*² Limbs with at least one claw each; back, if keeled, with at most three longitudinal ridges.
- b*¹ Outer body covering a soft skin without horny plates (*Chilota*).
TRIONYCHIDÆ, p. 513.
- b*² Outer body covering consisting of symmetrical horny plates (*Laminifera*).
- c*¹ Limbs not paddle-shaped; claws four or five on each limb. TESTUDINIDÆ, p. 488
- c*² Limbs paddle-shaped; claws two or one on each limb. CHELONIDÆ, p. 500

^aZeitschr. Wissensch. Zool., LXVI, 1899, pp. 407-434.

Suborder ATHECEÆ^a

1871. *Atheca* COPE, Proc. Am. Assoc. Adv. Sci., XIX, p. 235.
 1880. *Dermatochelyida* SEELEY, Quart. Journ. Geol. Soc., XXXVI, p. 412.
 1889. *Atheca* LYDEKKER, Cat. Foss. Rept. Brit. Mus., III (p. 223).
 1890. *Atheca* STRAUCH, Mém. Acad. St. Pétersbourg, (7) XXXVIII, No. 2, p. 38.
 1891. *Antheca* OKADA, Cat. Vert. Japan, p. 71 (misprint).

Family DERMOCHELIDÆ.

Limbs paddle-shaped, clawless; shell with five longitudinal dorsal keels and covered with a leathery skin.

The leather-back turtles inhabit the oceans between the Tropics, occasionally straying to temperate coasts. When full grown they reach an enormous size. Only one genus is known.

Genus DERMOCHELYS^b Blainville.

1816. *Dermochelys* BLAINVILLE, Bull. Soc. Philom. Paris, 1816, p. 119 (type, *T. coriacea*).
 1820. *Sphargis* MERREM, Syst. Amphib., p. 19 (same type).
 1822. *Coriudo* FLEMING, Philos. Zool., II (p. 271) (same type).
 1828. *Scytina* WAGLER, Isis, 1828, p. 861 (substitute for *Sphargis*).
 1830. *Dermatochelys* WAGLER, Nat. Syst. Amphib., p. 133 (emendation).
 1832. *Chelyra* RAFINESQUE, Atlantic Journ., I, No. 2, p. 64 (type, *T. coriacea*).

DERMOCHELYS SCHLEGELII^c (Garman).

1835. *Sphargis mercurialis* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., pp. 10 (part), 76, 139, pl. I: pl. II, figs. 3-5; pl. III (Japan) (not of Merrem).—OKADA, Cat. Vert. Japan, 1891, p. 71 (Tango).
 1850. *Sphargis coriacea* BLEEKER Natuurk. Tijds. Nederl. Indië, XV, (p. 260), Padang, Sumatra (not of Linnæus).—TICKEL, Journ. Asiat. Soc. Bengal, Nat. Hist., 1862, No. IV, p. 367, pl. — (Tenasserim).—McCoy, Nat. Hist. Victoria, II, Dec. XI, 1885, p. 1, pl. CI (Portland, Australia).—*Dermatochelys c.* GUENTHER, Rept. Brit. India, 1864, p. 55. *Dermochelys c.* BOULENGER, Cat. Chel. Brit. Mus., 1889, p. 10 (part); Fauna Brit. India, Rept. 1890, p. 50.—BURNE, Proc. Zool. Soc. London, 1905, I, pp. 291 seqv. (Japan; anatomy).
 1884. *Sphargis coriacea* var. *schlegelii* GARMAN, Bull. U. S. Nat. Mus., No. 25, p. 303 (Tropical Pacific and Indian Oceans).
 1884. *Sphargis schlegelii* GARMAN, Bull. U. S. Nat. Mus., No. 25, p. 295.

Dr. R. A. Philippi has recently described^d and roughly figured a new species *Sphargis angustata* from the coast of Chile. It is not possible at present to say whether it is identical with *D. schlegelii* or not.

The exact status of this form is not known and no specimen from the Pacific Ocean has come under my observation. Under these cir-

^a From the Greek *ἀ*, without, and *θήκη*, a closed box.

^b From *δέρμα*, skin; *χέλυς*, turtle.

^c Named after Dr. Hermann Schlegel. See p. 145.

^d Anales de Universidad, Mem. Cient. Liter., CII CIV, Oct. 1899, p. 730.

cumstances it does not seem necessary to submit any general description of this easily identified, gigantic animal. The young may be found some day in the southern waters adjacent to our territory and may then be referred to their proper place in the system by a comparison with the appended text figures, which are taken from an Atlantic specimen (figs. 373-376).

A specimen was recently captured in Japan and acquired for the Royal College of Surgeons in London, where the skeleton is now




FIG. 373.—DERMOCHELYS CORIACEA, YOUNG. NAT. SIZE. ENTIRE ANIMAL FROM ABOVE. No. 19796, U.S.N.M.


preserved. A detailed account of the muscular and visceral anatomy of this specimen has been published recently by Mr. R. H. Burne.

Mr. Burne gives the following description of some of the external parts of this animal.

Description. —*Young female:* Museum Roy. College Surgeons, London; Japan.

In color the animal was black above, blotched with irregular white spots, each of which measured on an average 1-2 cm. in diameter. The ventral surface of the body, limbs, and tail was dirty white, marked with irregular longitudinal bands and blotches of black.

The six longitudinal areas into which the carapace is divided by seven bony-ridges are apparently of equal breadth—11 cm. in the middle of the trunk—gradually narrowing toward the tail.



FIGS. 374-376.—DERMOCHELYS CORIACEA, YOUNG. NAT. SIZE. 374, UNDERSIDE OF SHELL; 375, SIDE OF HEAD; 376, UNDERSIDE OF HEAD. No. 19796, U.S.N.M.

There are six rows of scutes half embedded in the thick plastral integument, a double row along the mid-line, with two single rows about 11 cm. apart on either side.

Dimensions.

	<i>mm.</i>
Total length (following the curve of the carapace).....	1,350
Length between the bases of the flippers.....	680
Girth (under fore limb).....	1,350
Girth (midway between the limbs).....	1,400
Girth (at base of hind limb).....	910
Length of fore limb from its point of emergence from the body (following the outer curve).....	820
Greatest breadth of hand.....	200
Length of hind limb (tibial border).....	330
Girth of head at hinder extremity of the gape.....	530
From point of snout to inner canthus of eye.....	85
From point of snout to nostril.....	25

Habitat.—Tropical seas between the east coast of Africa and the west coast of South America, occasionally straggling farther north. Thus von Siebold obtained the specimen, from which the figure in *Fauna Japonica* is taken, in Japan. In the Imperial Museum, Ueno Park, Tokyo, there is one from the coast of the province of Tango, on the Sea of Japan. A young female was recently acquired by the Royal College of Surgeons, in London, from A. Owston.

Suborder LAMINIFERA. ^a1820. *Laminifera* HEMPRICH, Grundr. Naturg., p. 102.1880. *Aspidochelyida* SEELEY, Quart. Journ. Geol. Soc., XXXVI, p. 412.1895. *Cerachelya* HECKEL, Syst. Phylog. Wirbelth., p. —.

This suborder, which comprises the *Cryptodira* and *Placodira* of Cope and many later authors and which equals Dollo's *Thecophora* minus the soft-shelled turtles, as well as Baur's *Paradiacostoidea* minus the leather-back turtle, is trenchantly characterized by the horny plates which externally cover the shell.

Hemprich, as early as 1820, established the present division as opposed to that of the *Athecæ* plus *Chilotæ*, which he designated collectively as *Coriacea*.^b

The horny-shelled turtles belong to two different superfamilies, the *Testudinoidea*, corresponding to the group *Cryptodira*, and the *Chelydoidea*, equaling the *Placodira*, but only members of the former enter our limits, as at the present day the latter are confined to the southern hemisphere. Of the numerous families composing the *Testudinoidea* only two are intralimital, viz, the marine turtles, with paddle-shaped limbs having less than three claws, and the subfamily *Emydina* of the family *Testudinidæ*, having normal limbs with more than two claws.

Family TESTUDINIDÆ.

Subfamily EMYDINÆ.

Web-footed turtles having the nuchal plate without costiform lateral processes.

The terrapins constitute the bulk of the species and genera of turtles, widely distributed in the temperate and tropical countries. They live in streams, lagoons, or on land, and are both vegetable and animal feeders. Some species are highly esteemed as delicacies.

SYNOPSIS OF THE EMYDINE GENERA OCCURRING IN JAPAN, FORMOSA, AND KOREA.

*a*¹ Plastron not hinged; plastron and carapace joined by suture.

*b*¹ Triturating surface of upper jaw with a longitudinal median ridge. *Ocadia*, p. 489.

*b*² Triturating surface of upper jaw without a median ridge.

*c*¹ Choanae behind level of eyes; skin of hinder part of head divided into small shields *Grooclemys*, p. 496.

*c*² Choanae between the eyes; upper surface of head with undivided skin.

*d*¹ Hexagonal neural plates with the anterior lateral edge shorter than the posterior lateral edge *Clemmys*, p. 492.



*d*² Some of the hexagonal neural plates with the anterior lateral edge longer than the posterior lateral edge *Geocmyda*, p. 500.

*a*² Plastron hinged; plastron and carapace joined directly by ligament. *Cyclemys*, p. 503.

^a From *lamina*, a thin plate [of horn, in this case], and *fero*, I bear.

^b Grundr. Naturg., 1820, p. 101.


entire top of head smooth; body rather depressed, the depth being but slightly more than half the width of the carapace; carapace with



FIGS. 377-378. - *OCADIA SINENSIS*. NAT. SIZE. 377, CARAPACE; 378, PLAGSTRON. No. 29, SCI. COLL. TOKYO.

three keels, a median, continuous, broad, prominent, and one on each side consisting of four narrow, low keels, one on each costal, in a discontinuous series inasmuch as being placed obliquely across the costal shields the posterior end of each keel extends externally beyond the anterior end of the one following; nuchal broader than long; first vertebral squarish, somewhat broader in front than behind; all vertebrals broader than long, about as broad as the adjacent costals; margin slightly turned up, not serrated behind, though the posterior corner of each marginal extends slightly beyond the one following; eighth and ninth marginals broadest; plastron flat, emarginate behind sharply bent at the bridge, truncate anteriorly; posterior lobe as long as the width of the bridge, narrower than the shell; abdominal seam longest, equaling gular and humeral seams together, the latter being the shortest; pectoral seam as much longer than the femoral seam as the latter is longer than the anal seam; inguinal shield twice as large as axillary; toes webbed to the tips; tail long, a little more than one-half the length of the carapace, tapering to a point.


Color (in alcohol): Carapace raw umber, each shield with a large, polygonal, excentric, well-defined reddish brown spot; each marginal



OCADIA SINENSIS. (FROM GRAY.)

FOR EXPLANATION OF PLATE SEE PAGE 556.

anteriorly narrowly edged with bright yellow; underside of shell ochre yellow, each shield with a large reddish brown blotch at the outer posterior corner, each marginal and each shield on bridge with a narrow circle of dark brown, the reddish brown blotches being placed



FIGS. 379-381.—*OCADIA SINENSIS*. $1\frac{1}{2} \times$ NAT. SIZE. 379, TOP OF HEAD; 380, SIDE OF HEAD; 381, HEAD FROM FRONT. NO. 29, SCI. COLL. TOKYO.

excentrically to this ring; crown uniform olive-brown, the rest of head and neck olive, with narrow longitudinal black-edged, pale yellow stripes between pale olive stripes of nearly the same width; legs, feet, and tail similarly striped.

Dimensions.

	<i>mm.</i>
Greatest length of carapace.....	73
Greatest width of carapace.....	58
Greatest length of plastron.....	66
Length of hind lobe of plastron.....	24
Width of hind lobe of plastron.....	32
Width of opening of shell posteriorly.....	38
Width of bridge.....	24
Depth of shell.....	31
Width of head.....	12
Length of tail from vent.....	38

The *adults* are said to have shorter tails, the females shorter than the males. In very young specimens the tail is said to be about two-thirds the length of the shell. The lateral keels usually disappear in the adults, according to Boulenger, who records an adult specimen with a length of shell of 230 mm.

Variation.—The edges of both jaws are described by Boulenger as finely denticulated, and the pectoral seam as equaling or exceeding the gular and humeral seams together; nuchal may be longer than broad, and second and third vertebrae may be as long as broad.

Habitat.—The present species seems to be restricted to southern China and the island of Formosa. On the latter it has been recorded both from the northern and from the southern part. Swinhoe had it from Taiwan fu and the Senckenberg Museum has it from Takao, while I have examined a specimen from Taipa, and Mr. Tada notes it as common in the Tamsui River, though he may possibly have confounded it with *Cyclemys flavomarginata*.

List of specimens of Ocadia sinensis.

Museum.	No.	Age.	Locality	When col- lected.	By whom col- lected.
Sci. Coll. Tokyo.....	29	Young ^a	Taipei, Formosa.....	Sept. 1896	T. Tada.

^a Description, p. 489; figs. 377-381.

Genus CLEMMYS^a Ritgen.

1828. *Clemmys* RITGEN, Nova Acta Acad. Leop. Carol., XIV, Pt. 1, p. 272 (type, *Emys punctata*).
1832. *Chelopis* RAFINESQUE, Atlantic Journ., I, no. 2, p. 64 (same type).
1857. *Nanemys* AGASSIZ, Contr. Nat. Hist. U. S., I, p. 442 (same type).
1857. *Calemys* AGASSIZ, Contr. Nat. Hist. U. S., I, p. 443 (type, *C. mohlbergii*).
1857. *Glyptomys* AGASSIZ, Contr. Nat. Hist. U. S., I, p. 443 (type, *G. insculpta*).
1857. *Actinemys* AGASSIZ, Contr. Nat. Hist. U. S., I, p. 444 (type, *A. marmorata*).
1869. *Mauremys* GRAY, Proc. Zool. Soc. London, 1869 (p. 499) (type, *M. lanaria*).
1870. *Geoclemmys* GRAY, Suppl. Cat. Shield Rept. Brit. Mus., p. 26 (type, *G. guttata*; not of 1855).
1870. *Soculia* GRAY, Suppl. Cat. Shield Rept. Brit. Mus., p. 35 (type, *S. bealii*).
1870. *Emmenia* GRAY, Suppl. Cat. Shield Rept. Brit. Mus., p. 38 (type, *E. grayi*).
1870. *Eryma* GRAY, Suppl. Cat. Shield Rept. Brit. Mus., p. 44 (type, *E. laticeps*).

CLEMMYS JAPONICA (Temminck and Schlegel).



ISHIGAME.

Plate XXIX.

1835. *Emys vulgaris* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., p. 53 (Japan; not of Gray 1831).
1835. *Emys vulgaris japonica* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., p. 139 (Japan). SCHLEGEL, Abbild. Amph., 1840, p. 126, pl. XII.
1835. *Emys palustris* var. *japonica* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., pl. VIII, figs. 1-4; b pl. IX.
1844. *Emys japonica* GRAY, Cat. Tortois. Brit. Mus., p. 19 (Japan); Proc. Zool. Soc. London, 1869, p. 190 (Japan).—OKADA, Cat. Vert. Japan, 1891, p. 71 (Awaji; Bingo; Suwa; Chikuzen).—*Clemmys japonica* STRAUCH, Chelon. Studien, 1862 (p. 32); Mém. Acad. Sci. St. Pétersb. (7), XXXVIII, no. 2, 1890, p. 70 (Tsu-shima; Hondo).—HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 111 (Tokyo).—BOULENGER, Cat. Chel. Brit. Mus., 1889, p. 106 (Kobe; Yokohama).—BOETTGER, Kat. Rept. Mus. Senckenberg., I, 1893, p. 5 (Tokyo).
1851. *Emys caspica* var. DUMÉRIL, Cat. Méth. Rept. Mus. Paris, p. 8 (Japan).
1852. *Emys caspica* var. *japonica* DUMÉRIL, Arch. Mus. Paris, VI, p. 219.
1878. *Cistudo lutaria* BOETTGER, Offenbach. Ver. Naturk. 17 and 18 Ber., p. 8 (Japan) (not of Linnaeus, Bell, nor Schneider).

Judging from the above synonymy one might suppose that the Japanese *Clemmys* is a particularly close ally to one or both of the west-palaearctic species, since it has been described as a variety of *Emys vulgaris* by Temminck and Schlegel, and as a variety of *E. caspica* by Duméril. This is not so, however, and the only character which points in that direction is the short anal seam, a character


^a From *κλέμυς*, turtle. ^b Figs. 2-3 reproduced in this work on Plate XXIX.



CLEMMYS JAPONICA. (FROM SCHLEGEL.)
FOR EXPLANATION OF PLATE SEE PAGE 556.

which is not constant, as shown later on. Nor does it appear to be very closely allied to any of the other species of *Clemmys* known with certainty to occur on the Chinese mainland, viz, *C. bealii* and *C. nigricans*. Its nearest relative seems to be *C. schmackeri*, described not long ago by Doctor Boettger from a young specimen, the locality of which is given as "China, probably Hainan." It is permissible to suggest that this species may occur in various parts of southern China, and that the unconnected position of *C. japonica*, so curious in view of the close relationship of the other two Japanese turtles, *Amyda japonica* and *Geoclemys reevesii*, may be more apparent than real.

Clemmys japonica and *Ocadia sinensis* are very easily told apart by color alone as indicated in the "key" above (p. 489), but in general proportions, relations of shields, etc., they show many similarities,



FIGS. 382-383.—CLEMMYS JAPONICA. } NAT. SIZE. 382, CARAPACE; 383, PASTRON. No. 23522, U.S.N.M.

although a more close examination of details will reveal a great number of important differences. Apart from the difference in the triturating surface of the upper jaw, with a longitudinal ridge in the latter and without one in the former, *C. japonica* has no median notch to the edge of the upper jaw; its shell is more depressed and the hind lobe of the plastron relatively broader; the posterior outline of the carapace is also strongly serrated.

Description (figs. 382-383).—*Female*, third year; U.S.N.M., No. 34066; Tokyo; February 23, 1904; A. Owston collection. Snout but slightly projecting, vertical profile slightly oblique, straight; edges of jaws not denticulated, upper jaw without hooks or median notch; triturating surface of upper jaw narrow without any longitudinal ridge; mandibular symphysis nearly equaling length of eye slit; skin on entire top of head smooth; body rather depressed, the

depth being considerably less than half the length of carapace and but slightly more than half its width; carapace with a single, median, undulating keel; shields with deeply cut concentric lines toward the border, the vertebrals and costals with well-marked bosses from which issue radiating lines; nuchal rather large, almost as broad as long; first vertebral slightly wider than the others, which are as broad as or broader than the adjacent costals, and all much broader than long; edge of marginals from third to eighth turned up, behind the latter each marginal extending considerably beyond the next with its posterior corner, so that the outline is strongly serrate; eighth and ninth marginals broadest; plastron flat, angularly notched behind, obtusely emarginate in front, bridge angle gently rounded, each shield with deeply cut parallel lines mostly along the anterior and interior margins; posterior lobe fully as long as the bridge, as wide as the opening of the shell; abdominal seam much the longest, longer than that of humerals and gulars together, femoral seam longer than anal; inguinal shield somewhat larger than axillary; toes webbed to the tips; tail rather long, more than one-third the length of the carapace, tip rather blunt, base studded with numerous high conical tubercles or blunt spines. Color (in alcohol): Carapace tawny-olive, obscurely spotted with dusky; dorsal keel burnt umber, this color spreading sidewise on the transverse seams of the vertebrals; costals paler buff toward the outer margin, the dusky or blackish maculations being more distinct on this pale belt; underside uniformly black; head above uniform olive; a narrow black line from nostril to eye, and a narrow, pale, black-edged supratemporal band from the posterior corner of eye; neck olive gray with numerous raised, slightly tuberculated longitudinal ridges which appear to be paler; legs and feet blackish, with a pale band along the inner edge of forearm and tibia; tail with obscure, pale, longitudinal bands, one on each side of the median line.

Dimensions.

	<i>mm.</i>
Greatest length of carapace	124
Greatest width of carapace	88
Greatest length of plastron.....	117
Length of hind lobe of plastron.....	45
Width of hind lobe of plastron.....	60
Width of bridge.....	42
Depth of shell.....	47
Width of head.....	19
Length of tail from vert.....	45

The *adult* is more elongate in proportion to width and depth than the one described above. The *male* differs but slightly from the female, the plastron being scarcely concave, but the hind lobe of the plastron is slightly narrower and the depth of the shell somewhat



GEOEMYDA SPENGLERI.

FOR EXPLANATION OF PLATE SEE PAGE 556.

provinces of Awaji, Bingo, Suwa, and Chikuzen. The Academy of Sciences in St. Petersburg has it from Tsushima.

List of specimens of Clemmys japonica.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.	Length of carapace.	Width of carapace.	Length of plastron.	Width of hind lobe.	Width of bridge.	Depth of shell.	Tail, from vent.
					mm	mm	mm	mm	mm	mm	mm
9551	Female ^a	Japan.....	1896.	E. S. Morse.....	182	128	175	103	68	73	69
23522	Young ^b	Yokohama.....	Oct.	L. Stejneger.....							
23523do. ^cdo.do.do.							
23524do. ^cdo.do.do.							
23525do. ^cdo.do.do.							
23526do. ^ddo.do.do.	35	31	29	16	9	14	32
23527do. ^cdo.do.do.							
			1904.								
34066	Female ^e	Tokyo.....	Feb. 23	A. Owston.....	124	88	117	60	42	47	45
34067	Male ^fdo.do.do.							
34068do. ^fdo.do.do.	112	81	100	53	35	40	31
34069	Young ^gdo.do.do.							
34070do. ^gdo.do.do.							
34071do. ^gdo.do.do.	87	66	80	41	29	33	34
34072do. ^gdo.do.do.							
34073do. ^gdo.do.do.							
34074do. ^gdo.do.do.							
34075do. ^gdo.do.do.							

^a P. 495.

^d Recently hatched; fig. 384.

^f Third year.

^b Second year; figs. 382-383.

^c Third year; description, p. 493.

^g Second year.

^e Recently hatched.

Genus *GEOCLEMYS*^a Gray.

1855. *Geoclemys* GRAY, Cat. Shield Rept. Brit. Mus., 1855, p. 17 (type, *G. hamiltonii*).

1869. *Damonia* GRAY, Proc. Zool. Soc. London, 1869, p. 193 (type, *D. macrocephala*=*G. subtrijuga*).

In his Catalogue of Shielded Reptiles (1855) Gray first established the genus *Geoclemys* for the following species, without indicating a type:

- G. hamiltonii.* *G. seba.* *G. muhlenbergii.*
- G. reevesii.* *G. pulchella.* *G. guttata.*


Two years later Agassiz^b made each of the last three species a type of separate genera, respectively *Glyptemys*, *Clemmys*, and *Nanemys*.

G. seba is a synonym of *Emys trijuga* Gray (1855) which he himself in 1869^c made the type of his new genus *Melanochelys*.


^a From *γῆ*, earth, land; and *clemys*, erroneous for *κλεμμυς*, a turtle.

^b Contr. Nat. Hist. U. S., I. 1857.


^c Proc. Zool. Soc. London, 1869, p. 187.



1



2



3

GEOCLEMYS REEVESII. (FROM GRAY.)

FOR EXPLANATION OF PLATE SEE PAGE 556.

This leaves only the species *G. hamiltonii* and *G. reevesii* in the genus *Geoclemmys*, and both being strictly congeneric it matters not which one we select as the type. It is, consequently, quite irrelevant that Gray himself in 1869^a included these in his new genus *Damonia*. As the latter only includes species congeneric with *G. hamiltonii* and *reevesii*, it becomes a synonym of Gray's *Geoclemmys* of 1855. His *Geoclemmys* of 1869, as he then spelt it, on the other hand becomes a synonym of *Clemmys*.

GEOCLEMYS REEVESII *b* (Gray).

Plate XXX.

1831. *Emys reevesii* GRAY, Synops. Rept., p. 73 (type-locality China; types in Brit. Mus.; J. Reeves, jr., collector).—RÜETIMYER, Verh. Naturf. Ges. Basel, VI, Pt. 1, 1873, p. 48 (Japan).—*Geoclemmys reevesii* GRAY, Cat. Shield Rept. Brit. Mus., I, 1855, p. 18, pl. v (China).—*Damonia reevesii* GRAY, Proc. Zool. Soc. London, 1869, p. 194.—BOULENGER, Cat. Chel. Brit. Mus., 1889, p. 95 (China; Japan).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, 1904, p. 353 (Shanghai and Hankow, China).—*Clemmys* *v.* STRAUCH, Chenol. Stud., 1862 (p. 104); Mém. Acad. Sci. St. Pétersb. (7), XXXVIII, No. 2; 1890, p. 74 (China; Chemulpo, Korea).—SCLATER, Proc. Zool. Soc. London, 1873, p. 517 (Ningpo).
1840. *Emys vulgaris picta* SCHLEGEL, Abbild. Amph., p. 127, pl. XLII (type-locality, Japan; types in Leiden Mus.; Buerger, collector) (not *E. picta* of Schweigger).
1851. *Emys japonica* DUMÉRIL, Cat. Méth. Rept. Mus. Paris, I, p. 8 (Japan); Arch. Mus. Paris, VI, 1852, p. 220 (not *E. vulgaris japonica* of Schlegel).
1873. *Damonia unicolor* GRAY, Ann. Mag. Nat. Hist. (4), XII (p. 78) (type-locality, Shanghai, China; types in Brit. Mus.; R. Swinhoe, collector).—*Clemmys unicolor* SCLATER, Proc. Zool. Soc. London, 1873, p. 517, pl. XLIV (correct type-locality, Ningpo).—HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 111 (Japan).
1889. *Damonia reevesii* var. *unicolor* BOULENGER, Cat. Chel. Brit. Mus., p. 96 (Shanghai: [probably Ningpo, Sclater, Proc. Zool. Soc. London, 1873, p. 577]).—WERNER, Abh. Bayer. Akad. Wiss. (Muenchen), II Klasse, XXII, Pt. 2, p. 353 (Hankow, China).
1891. *Emys* ? *sincensis* OKADA, Cat. Vert. Japan, p. 72 (Osaka) (not of Gray).


Description.—*Male*; U.S.N.M. No. 21181; Seoul, Korea; August, 1883; P. L. Jouy, collector (figs. 385–388). Snout rather pointed, strongly projecting, vertical profile oblique; edges of jaws not denticulated, upper jaw without hooks or median notch; triturating surface of upper jaw broad, without any longitudinal ridge; mandibular symphysis slightly longer than eye slit; anterior part of head covered by a large smooth shield on crown and snout, another on each side covering the whole temporal region, and the usual rostro-labial shield; parietal and auricular regions covered by small polygonal

^a Proc. Zool. Soc. London, 1869, p. 193.

^b For Mr. John Russell Reeves, jr., who collected the type.


^c Reproduced in this work on Plate XXX.

shields, most of those on the former with a central rounded tubercle; body moderately depressed, its depth being less than half the length, but considerably more than half the width of the carapace; carapace with nearly parallel sides, and three strong keels; shields with deeply



FIGS. 385-386.—*Geoclemys reevesii*. $\frac{1}{2}$ \times NAT. SIZE. 385, CARAPACE; 386, PASTRON. No. 21181, U.S.N.M.

cut concentric lines, radiating lines also fairly strong; nuchal small, narrower in front, emarginate behind, as long as broad; first vertebral pentagonal, broader in front than behind, not broader than the others



FIGS. 387-388.—*Geoclemys reevesii*. $\frac{1}{4}$ \times NAT. SIZE. 387, TOP OF HEAD; 388, SIDE OF HEAD. No. 21181, U.S.N.M.

which are of subequal width and of the same width as the anterior three costals; edge of marginals from third to eighth turned up; posterior marginals not projecting their corners so that the posterior outline of the shell is nearly even, not serrated; ninth marginal broad-

est; plastron very slightly concave, angularly notched behind, truncate in front, bridge angle rounded, shields nearly smooth; posterior lobe slightly longer than bridge, nearly as wide as opening of the shell; abdominal seam longest, but only slightly longer than femoral, less than humeral and gular together; femoral longer than pectoral; gular seam longer than anal, humeral shortest; inguinal shield larger than axillary; toes webbed to the tips; tail tapering, somewhat compressed at tip. Color (in alcohol): Carapace chestnut brown, the median keel blackish brown; a small darker brown pentagonal spot on the areola of each costal above the lateral keel; vertebrals and transverse edges of costals narrowly outlined with yellow; shields of underside blackish brown with all seams and outer edge of hind lobe of plastron edged with buff, the median seams broadly so; head, neck, and limbs tawny olive; a number of yellow, black-edged stripes and vermiculations on side of head, the principal being supratemporal line; a medio-temporal line; a preauricular line; a line posteriorly on both lips and surrounding the angle of mouth; a small spot in front of eye; several irregular lines and spots on lower jaw; neck with numerous longitudinal lines of ocellated, dull buff spots; limbs and tail without distinct markings, though on underside of thigh there are indications of lines of spots similar to those on neck.

Dimensions.

	<i>mm.</i>
Greatest length of carapace.....	122
Greatest width of carapace.....	87
Greatest length of plastron.....	119
Length of hind lobe of plastron.....	44
Width of hind lobe of plastron.....	53
Width of bridge.....	42
Depth of shell.....	53
Width of head.....	21
Length of tail from vent.....	37

Variation.—The specimens examined by me offer but very insignificant variations in structure as well as in coloration. The latter, however, is not constant, and specimens, more or less uniform black, occur both in China, where they have given rise to "*Damonia unicolor*," and in Japan. The specimens from the latter country, which Schlegel described as *Emys vulgaris picta* seem to belong to this color variety, but the yellowish marks on side of head and neck were very conspicuous. That perfectly uniform black ones occur in Japan also seems certain from Hilgendorf's notes on a specimen of "*Chelmys unicolor*," which he purchased alive in Tokyo, and which had "neck and head . . . plain dusky without any kind of markings."

Habitat.—Eastern and southern China, Korea, and southern Japan.

In China it is known from Tientsin to Canton and in the interior at least as far as Hankow. It is also reported from Cochin China.

Its occurrence in Korea was first recorded by Strauch from a specimen collected by Doctor Bunge at Chemulpo in 1890 and now in the museum of the St. Petersburg Academy (No. 7907). Two other specimens from Korea, only one of which has a definite locality, are in our museum collected by Jouy and Bernadou, respectively.

In Japan it seems to be rare and is probably restricted to the southern part. It was the first addition to the herpetological fauna of the country after the publication of the Fauna Japonica, and was made public by Schlegel himself (as *Emys vulgaris picta*) from three specimens collected by Mr. Buerger, probably in Kiusiu. Since then but few Japanese specimens have been recorded, mostly without definite localities. I myself have only examined one from Kagoshima, Satsuma, in Kiusiu. In the Hamburg Museum there are several specimens collected by Doctor Lenz, in 1896, in the province of Setsu, Hondo (No. 188). Okada records a turtle from Osaka as *Emys sinensis* which probably is the present species. British Museum has it from Tsushima collected by Holst.

List of specimens of Geoclemys reevesii.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected.	Length of carapace.	Width of carapace.	Length of plastron.	Width of hind lobe.	Width of bridge.	Depth of shell.	Tooth from vent.
					mm	mm	mm	mm	mm	mm	mm
14512	Male	Korea	1883.	J. B. Bernadou	101	73	93	41	34	43	30
21181	do. ^a	Seoul, Korea	Aug. --	P. L. Jouy	122	87	119	53	42	53	37
21249	Young ^b	(Bering Island)	Sept. *	N. A. Grebnitski							

^a Description, p. 497; figs. 385-388.

^b Thrown out by the sea.

Genus GEOEMYDA ^a Gray.

1834. *Geomyda* GRAY, Proc. Zool. Soc. London, 1834, p. 100 (type, *Testudo spengleri*).

1836. *Geomyds* BONAPARTE, Chelon. Tab. Anal., p. 6 (emendation; same type).

1855. *Nicoria* GRAY, Cat. Shield Rept. Brit. Mus., I, p. 17 (same type).

1869. *Melanochelys* GRAY, Proc. Zool. Soc. London, 1869, p. 187 (type, *M. trijuga*).

1876. *Chaibassia* THEOBALD, Cat. Rept. Brit. India (p. 6) (type, *Ch. tricarinata*).

The reason for adopting the name *Geomyda* instead of *Nicoria* for the present genus is the fact that *G. spengleri* was specifically designated as the type when the genus was first established as demonstrated by me in detail on a former occasion.^b


^a From $\gamma\eta$, earth, land; and *myda*, for *emys*, from $\epsilon\mu\nu\varsigma$, turtle.

^b Proc. Biol. Soc. Washington, XV, Dec. 16, 1902, pp. 237-238.



GEOEMYDA SPENGLERI.

FOR EXPLANATION OF PLATE SEE PAGE 556.



2

1

GEOMYDA SPENGLERI.

FOR EXPLANATION OF PLATE SEE PAGE 556.

GEOEMYDA SPENGLERI^a (Gmelin).

YAMA GAME (in Okinawa dialect, according to Okada).

Plates XXXI and XXXII.

1789. *Testudo spengleri* GMELIN, Syst. Nat., I, Pt. 3, p. 1043 (no locality given).—*Geomyda spengleri* GRAY, Proc. Zool. Soc. London, 1834, p. 100 (China).—*Nicoria spengleri* GRAY, Cat. Shield Rept. Brit. Mus., I, 1855, p. 17 (China).—BOULENGER, Cat. Chel. Brit. Mus., 1889, p. 120 (Sumatra; Borneo; southern China?); Ann. Mag. Nat. Hist. (6), X, Oct. 1892, p. 302 (Okinawa); Proc. Zool. Soc. London 1893, p. 237 (Okinawa).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 859; author's separate, p. 10 (Tokuchimura, Okinawa).—*Emys? spengleri* OKADA, Cat. Vert. Japan, 1891, p. 72 (Okinawa shima).
1802. *Testudo serrata* SHAW, Gen. Zool., III, Pt. 1, p. 51, pl. ix, fig. 2 (no locality given).
1804. *Testudo tricarinata* BORY DE ST. VINCENT, Voy. Iles d'Afrique, II (p. 308, pl. xxxvii, fig. 1).

Description.—*Female*; U.S.N.M. No. 34053; Naha, Okinawa shima; A. Owston collection. Snout short, not projecting, lateral profile vertical, straight; edges of jaws not denticulated, upper jaw hooked medially, without lateral notches; triturating surface of upper jaw narrow without any longitudinal ridge; mandibular symphysis shorter than length of eye slit; head above covered with smooth skin; body rather depressed, its depth being somewhat more than half the width; carapace slightly wider behind, considerably emarginate in front; three well-developed keels; shields (very badly eroded) with distinct concentric lines; nuchal large, broader behind than in front, broader than long; first vertebral pentagonal, slightly broader in front than behind, as long as second and slightly longer than third and fourth, all somewhat broader than long, and as broad as the adjacent costals; edge of marginals from third to eighth slightly turned up; anterior and posterior marginals with their posterior corners greatly projecting, so as to make the anterior and posterior outlines of the shell strongly serrate; first and second marginals broadest; plastron flat, deeply emarginate behind, very obtusely angulate in front with outer anterior corners of gulars projecting; bridge angle gently rounded, shields smooth; posterior lobe slightly longer than bridge, as wide as opening of shell; abdominal and pectoral seams subequal, longest, as long as humeral and gular together, the latter shortest; femoral seam longer than anal, which in turn is longer than humeral; axillary shield distinct, moderate; inguinal shield wanting; forearm anteriorly with large scales the tips of which are pointed and projecting; web between toes emarginate; tail short, rather broad, depressed, with seven pairs of flat square shields on upper surface toward the tip; base of tail and posterior aspect of

^a For Lorentz Spengler, intendant of the Royal "Kunstskammer" in Copenhagen, from whom Wahlbaum received a shell of this species. Spengler was a conchologist of note. He was born in 1720 and died in 1807.

femur with numerous long, blunt spines. Color (in alcohol): Carapace tawny ochraceous more or less clouded with darker brown; underside black, margins of plastral lobes, bridge angle as well as larger or smaller areas on underside of marginals wax-yellow; head on top dark tawny olive; an indistinct band on canthus rostralis, lower eyelid and on supratemporal region as well as entire neck above pale isabella-color; sides and underside of head and neck blackish with several more or less well-defined, pale isabella-colored markings, viz, a narrow postocular line over tympanic region to sides of neck, a line on posterior half of upper lip crossing the lower jaw at angle of mouth and an irregular broad mark on throat with branches on lower jaw and anterior neck; horny jaws pale brownish; limbs with a couple of ill-defined stripes of same color, and tail above similarly pale.

Dimensions.

	<i>mm.</i>
Greatest length of carapace.....	128
Greatest width of carapace.....	84
Greatest length of plastron.....	113
Length of hind lobe of plastron.....	41
Width of hind lobe of plastron.....	59
Width of bridge.....	38
Depth of shell.....	49
Width of head.....	21
Length of tail from vent.....	24

Variation.—Boulenger, in Chinese specimens, notes the following sequence in the length of the plastral shields, beginning with the largest: Abdominals, femorals, pectorals, humerals, anals, gulars, while our specimen from Okinawa shima and the specimen from Ishigaki shima in the Kumamoto school show the following sequence with regard to their plastral seams, beginning with the longest: Abdominal = pectoral, femoral, anal, humeral, gular. Boulenger also notes the axillary shield as absent, while in ours it is very distinct and well-developed.

Habitat.—The distribution of this species is still of considerable uncertainty, as specimens with undoubted and explicit localities are rare in collections. It is said to occur in Borneo and Sumatra, as well as in southern China. It is not recorded from Formosa, but on the other hand has been received from the Riu Kius.

Holst sent two specimens from Okinawa to the Zoological Society in London, and Doctor Fritze obtained three specimens at Tokuchimura on the west side of Okinawa shima. The Imperial Museum, Ueno Park, Tokyo, also has a specimen from Okinawa, and recently the United States National Museum has obtained, through Mr. Ows-ton, a specimen from Naha (No. 34053).

Professor Ijima has sent me excellent drawings of a specimen belonging to the Kumamoto Higher Middle School. It was collected in Ishigaki shima of the Yaeyama subgroup in February, 1893.

List of specimens of Geoemyda spengleri.

U. S. N. M. No.	Sex.	Locality.	When col- lected.	By whom col- lected or from whom received.	Length of carapace.	Width of carapace.	Length of plastron.	Width of hind lobe.	Width of bridge.	Depth of shell.	Tail, from vent.
34053	Female a.	Naha, Okinawa.....		A. Owston.....	mm. 128	mm. 84	mm. 113	mm. 59	mm. 38	mm. 49	mm. 24

a Description, p. 501.

Genus CYCLEMYS^a Bell.

1830. *Sternothaerus* WAGLER, Nat. Syst. Amph., p. 137 (type, *S. trifasciatus*) (not of Bell, 1825).
1834. *Cyclemys* BELL, Proc. Zool. Soc. London, June, 1834, p. 17 (type, *C. orbiculata*).
1855. *Cuora* GRAY, Cat. Shield Rept. Brit. Mus., I, p. 41 (type, *C. amboinensis*).
1863. *Cistoclemmys* GRAY, Proc. Zool. Soc. London, 1863, p. 175 (type, *C. flavomarginata*).
1863. *Pyciclemmys* GRAY, Proc. Zool. Soc. London, 1863, p. 176 (type, *Cuora trifasciata*).
1870. *Cystoclemmys* GRAY, Suppl. Cat. Shield Rept. Brit. Mus., p. 20 (emendation).

CYCLEMYS FLAVOMARGINATA^b Gray.

Plate XXXIII.

1863. *Cistoclemmys flavomarginata* GRAY, Proc. Zool. Soc. London, 1863, p. 175 (type-locality, Tamsui, Formosa; type in Brit. Mus.; Swinhoe, collector).—SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 220 (Tamsui).—*Cuora flavomarginata* GUENTHER, Rept. Brit. India, 1864, p. 13, pl. v, fig. A^c (Tamsui).—*Cystoclemmys flavomarginata* GRAY, Suppl. Cat. Shield Rept. Brit. Mus., 1870, p. 20 (China; Formosa).—*Cyclemys flavomarginata* BOULENGER, Cat. Chel. Brit. Mus., 1889, p. 135 (Tamsui, Formosa; China?).
1891. *Cuora amboinensis* OKADA, Cat. Vert. Japan, p. 72 (Yaeyama) (not of Daudin).—*Emys amboinensis* FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 864; author's separate, p. 15 (Yaeyama).

Description.—*Adult female*: U.S.N.M. No. 34077; Ishigaki shima, Riu Kiu Archipelago; 1899; A. Owston collection. Snout short, not projecting, lateral profile vertical, straight; edges of jaws not denticulated, upper jaw feebly hooked mesially, without notches; triturating surface of upper jaw moderately wide, without any longitudinal ridge; mandibular symphysis shorter than length of eye slit; head above covered with smooth skin; body not at all depressed, its depth being considerably more than one-half its width; carapace slightly

a From κύκλος, circle; ἔμυς, river turtle.



b Signifying margined with yellow.

c Reproduced in this work on Plate XXXIII.

wider behind the middle, not emarginate in front; a well-developed vertebral keel and an interrupted keel on each side dividing the costal areoles; shields with small areoles and wide margins with numerous deeply cut concentric lines; nuchal rather large, broader behind than in front, longer than broad; first vertebral somewhat narrower than the others, and like these, with the exception of fifth, angularly emarginate behind; second vertebral slightly longer than broad, longest, the others broader than long, all except fifth narrower than the adjacent costals; sixth to tenth marginals broadest, first slightly narrower; lateral edge slightly turned up; none of the marginals extending beyond the others, the whole outline being entire, not serrate; plastron flat, each shield with deep cut lines, rounded behind and in front; a ligamentous hinge between pectoral and abdominal shields; plastron fastened to carapace with ligament and as wide as opening of shell; abdominal seam longest; anal seam slightly longer than pectoral, which equals humeral and gular seams together; humeral and femoral seams very short, the last one shortest and less than one-fourth the anal seam; axillary present, but inguinal shield wanting; forearm anteriorly with a few large scales, with free, not pointed, edges; digits with a slight web at base; base of tail and posterior aspect of femur with numerous blunt tubercles. Color (in alcohol): Carapace blackish brown, each areole bright chestnut, median keel from middle of second vertebral backward bright buff; plastron uniform blackish brown, marginals underneath and a narrow outer margin on abdominal and pectoral shields pale buff; top of head dark drab, parietal region exteriorly and posteriorly narrowly edged with black, a narrow black-edged bridge of drab on the median line posteriorly joining with the isabella-color of the neck; sides of head pale reddish gray; a narrow black line from upper corner of eye widening backward and bordering beneath a broad pale band, which surrounds the sides and posterior outline of the dark parietal region; the pale, dark-edged band is faintly indicated in front of the eye on the canthus rostralis; neck isabella-color, with indications of longitudinal buff, dark-edged stripes; legs dull ochraceous underneath, palms and soles dusky; tail isabella-colored, with a dusky, longitudinal band on each side and a dusky line on the middle line above and below.

Dimensions.

	<i>mm.</i>
Greatest length of carapace	124
Greatest width of carapace	92
Greatest length of plastron	121
Width of hind lobe of plastron	69
Depth of shell	58
Width of head	21
Length of tail from vent	20



CYCLEMYS FLAVOMARGINATA. (FROM GUENTHER.)

FOR EXPLANATION OF PLATE SEE PAGE 556.

Variation.—As in many other turtles with deeply carved shields there is great variation in the intensity of the sculpturing. Thus No. 34076, also a female but a few millimeters longer than the one described above, is entirely smooth underneath and nearly so above. Otherwise there is but little noteworthy variation in the four specimens before me. In the largest one just mentioned, the femoral seam is slightly longer than the humeral, and the posterior half of the anal seam has become obliterated; in the two younger specimens the pectoral seam is relatively somewhat shorter than in the two adults inasmuch as it does not equal the length of the humeral and gular seams together. All four are practically alike in color.

Habitat.—This species has been recorded from southern China as well as from Formosa, though some doubt has been expressed as to the correctness of the Chinese habitat.

Swinhoe found it to be frequent in the Tamsui River, northern Formosa, where according to him it is the prevailing species.^a

Four specimens recently acquired by the U. S. National Museum from Mr. A. Owston show that this species also occurs in Ishigaki shima of the Yaeyama subgroup of the Riu Kiu archipelago.

The *C. amboinensis* of Okada and Doctor Fritze, said to occur in the Yaeyama group, is undoubtedly meant for the present species. Doctor Fritze states that from the latter islands it is occasionally brought to Okinawa shima.

List of specimens of Cyclemys flavomarginata.

U. S. N. M. No.	Sex and age.	Locality.	When col- lected.	By whom col- lected or from whom received.	Length of carapace.		Length of plastron.	Width of hind lobe.		Tail, from vent.
					mm.	mm.		mm.	mm.	
34076	Female	Ishigaki shima, Riu Kiu	1899	A. Owston	131	93	125	73	56	19
34077	do. ^a	do.	1899	do.	124	92	121	69	58	20
34078	Halfgr.	do.	1899	do.	92	78	92	58	47
34079	Young	do.	1899	do.	69	58	65	38	31	13

^a Description, p. 533.

^a "Swinhoe has frequently seen the Tamsui tortoise showing its head and the top of its back on the surface of the water in ponds about the rice-fields, and has watched them basking, several at a time, on the top of large stones in such ponds." (Gray, Suppl. Cat. Shield Rept. Brit. Mus., p. 20).

GEOEMYDA SPENGLERI^a (Gmelin).**YAMA GAME** (in Okinawa dialect, according to Okada).

Plates XXXI and XXXII.

1789. *Testudo spengleri* GMELIN, Syst. Nat., I. Pt. 3, p. 1043 (no locality given).—*Geoemyda spengleri* GRAY, Proc. Zool. Soc. London, 1834, p. 100 (China).—*Nicoria spengleri* GRAY, Cat. Shield Rept. Brit. Mus., I, 1855, p. 17 (China).—BOULENGER, Cat. Chel. Brit. Mus., 1889, p. 120 (Sumatra; Borneo; southern China?); Ann. Mag. Nat. Hist. (6), X, Oct. 1892, p. 302 (Okinawa); Proc. Zool. Soc. London 1893, p. 237 (Okinawa).—FRITZE, Zool. Jahrb. Syst., VII, 1894, p. 859; author's separate, p. 10 (Tokuchimura, Okinawa).—*Emys? spengleri* OKADA, Cat. Vert. Japan, 1891, p. 72 (Okinawa shima).
1802. *Testudo serrata* SHAW, Gen. Zool., III, Pt. 1, p. 51, pl. ix, fig. 2 (no locality given).
1804. *Testudo tricarinata* BORY DE ST. VINCENT, Voy. Iles d'Afrique, II (p. 308, pl. xxxvii, fig. 1).

Description.—*Female*: U.S.N.M. No. 34053; Naha, Okinawa shima; A. Owston collection. Snout short, not projecting, lateral profile vertical, straight; edges of jaws not denticulated, upper jaw hooked medially, without lateral notches; triturating surface of upper jaw narrow without any longitudinal ridge; mandibular symphysis shorter than length of eye slit; head above covered with smooth skin; body rather depressed, its depth being somewhat more than half the width; carapace slightly wider behind, considerably emarginate in front; three well-developed keels; shields (very badly eroded) with distinct concentric lines; nuchal large, broader behind than in front, broader than long; first vertebral pentagonal, slightly broader in front than behind, as long as second and slightly longer than third and fourth, all somewhat broader than long, and as broad as the adjacent costals; edge of marginals from third to eighth slightly turned up; anterior and posterior marginals with their posterior corners greatly projecting, so as to make the anterior and posterior outlines of the shell strongly serrate; first and second marginals broadest; plastron flat, deeply emarginate behind, very obtusely angulate in front with outer anterior corners of gulars projecting; bridge angle gently rounded, shields smooth; posterior lobe slightly longer than bridge, as wide as opening of shell; abdominal and pectoral seams subequal, longest, as long as humeral and gular together, the latter shortest; femoral seam longer than anal, which in turn is longer than humeral; axillary shield distinct, moderate; inguinal shield wanting; forearm anteriorly with large scales the tips of which are pointed and projecting; web between toes emarginate; tail short, rather broad, depressed, with seven pairs of flat square shields on upper surface toward the tip; base of tail and posterior aspect of

^a For Lorentz Spengler, intendant of the Royal "Kunstammer" in Copenhagen, from whom Wahlbaum received a shell of this species. Spengler was a conchologist of note. He was born in 1720 and died in 1807.

Family CHELONIIDÆ.

Hard-shelled turtles with paddle-shaped limbs.

A small group of half a dozen species inhabiting the tropical and subtropical seas, often found hundreds of miles from land, but resorting to sandy beaches in order to deposit their numerous round eggs. Full-grown individuals reach an enormous size, 7 feet in length and weighing 800 to 900 pounds. Some species—for instance, the green turtle—are highly esteemed as food, while another furnishes the true "tortoise shell" of commerce.


Our knowledge of the marine turtles has advanced very little since the time when Doctor Guenther treated of them in his Reptiles of British India (1864). Boulenger, in his Catalogue of the Chelonians in the British Museum (1889), regards the species which are treated of in the present work as identical with the Atlantic forms, though under the loggerhead turtle he makes the admission that the enormous variation in a large series "leaves no alternative but to further multiply the number of species or to admit only one." It is not impossible that he found similar alternatives in the other species. My own material is too limited to decide the status of these forms, and under these circumstances I consider it much more rational to enumerate the West Pacific forms under names corresponding to those employed by Guenther and Garman. Such a treatment is much less calculated to cause confusion than the opposite one of joining together those which are not with certainty known to belong together.

Dr. R. A. Philippi, of Santiago, Chile, has recently described a number of new marine turtles from that country, but the descriptions are not sufficiently explicit to justify any judgment as to their actual status. Under these circumstances no attempt has been made to correlate any of these new names with those of the marine turtles of the western and northern Pacific.

Only one species of each genus being known to occur within our territory, the key to the genera becomes a key to the species at the same time.

KEY TO THE GENERA AND SPECIES OF CHELONIIDÆ OCCURRING IN JAPANESE WATERS.

- a*¹ Costal shields 5 pairs, or more *Caretta olivacea*, p. 507.
*a*² Costal shields 4 pairs.
 *b*¹ One pair of prefrontal shields *Chelonia japonica*, p. 509.
 *b*² Two pairs of prefrontal shields *Eretmochelys squamosa*, p. 511.



CARETTA OLIVACEA. (FROM ESCHSCHOLTZ.)

FOR EXPLANATION OF PLATE SEE PAGE 556.

Genus CARETTA^a Rafinesque.

1814. *Caretta* RAFINESQUE, Specchio Sci. (Palermo), II, no. 9, 1 Sett., 1814, p. 66 (type, *C. nasuta*=*Testudo caretta*).
1835. *Thalassochelys* FITZINGER, Ann. Wien Mus., I, p. 121 (type, *Testudo caouana*=*T. caretta*).
1838. *Caouana* COCTEAU in Sagra's Hist. Fis. Pol. Nat. Cuba, IV, Rept., p. 31 (type, *Chelonia cephalo*=*T. caretta*).
1843. *Lepidochelys* FITZINGER, Syst. Rept., p. 30 (type, *Chelonia olivacea*).

The necessity for adopting Rafinesque's *Caretta* for the present genus has already been pointed out by me in my Herpetology of Porto Rico.^b

CARETTA OLIVACEA^c (Eschscholtz).

AKAUMIGAME.

Plate XXXIV.

1829. *Chelonia olivacea* ESCHSCHOLTZ, Zool. Atlas, Pt. 1, p. 2, pl. iii^d (type-locality, Manila Bay, P. I.)—*Caretta olivacea* RUEPPELL, Neue Wirbelth. Abyssin., 1835 (p. 7, pl. iii).—*Thalassochelys olivacea*, STRAUCH, Chelon. Stud., 1862, (p. 63).—GARMAN, Bull. U. S. Nat. Mus. No. 25, 1884, p. 301 (Tropical Pacific and Indian oceans).—*Caouana olivacea* GRAY, Cat. Tortois. Brit. Mus., 1844, p. 53 (Philippine Is.).—GUENTHER, Rept. Brit. India, 1864, p. 52 (seas of Philippine Is. and of China).—*Lepidochelys olivacea* GIRARD, Herpet. U. S. Expl. Exped., 1858, p. 435.
1831. *Chelonia caretta* β *olivacea* GRAY, Synops. Rept., I, p. 54 (China).
1835. *Chelonia dussumieri* DUMÉRIL and BIBRON, Erpét. Gén., II, p. 557 (coast of Malabar; types in Paris Mus.).—DUMÉRIL, Cat. Méth. Rept. Mus. Paris, I, 1851, p. 25.
1857. *Lepidochelys dussumieri* GIRARD, Herpet. U. S. Expl. Exped., p. 437.
1880. *Thalassochelys corticata* HILGENDORF, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, p. 112 (Japan) (not of Girard, 1858).
1891. *Chelonia caouana* OKADA, Cat. Vert. Japan, p. 72 (not of Schweigger, 1814) (Awa, Hondo; Toza; Bonin Is.).—MITSUKURI, Zool. Mag., Tokyo, III, no. 35, Sept. 1891, p. 364 (breeding habits, etc.); Journ. Sci. Coll. Tokyo, VI, 1893, p. 227 (Sagara, prov. Totomi), X, Pt. 1, 1896, pp. 11, etc. (embryology).

The Asiatic loggerhead turtle is easily distinguished from the other marine turtles of the Far East by having more than four pairs of costal shields. Like the hawksbill it has two pairs of prefrontals, but the shields on the carapace are not imbricate. The figure of the young Atlantic loggerhead will help placing any specimen of the group in the proper genus, but it should be remembered that the vertebral and costal shields in *C. olivacea* are apt to be more numerous. No

^a The name Caret (new Latin *Caretta*) according to Lacépède is the one by which the hawksbill turtle is generally known in the country it inhabits. Carey is Spanish for tortoise shell.

^b Report U. S. Nat. Mus., 1902, p. 714.

^c Signifying olive-colored.

^d Reproduced in this work on Plate XXXIV.

authentic specimen from Asiatic waters is in our collection, hence no detailed description is feasible at present. (Figs. 389-392.)

Habitat.—Notwithstanding the fact that the present species is not included in the list of Japanese reptiles in the *Fauna Japonica*, it is





FIG. 389.—*CARETTA CARETTA*, YOUNG. NAT. SIZE. ENTIRE ANIMAL FROM ABOVE. No. 14823, U.S.N.M.

probably the most common of the marine turtles in that part of the world. Professor Mitsukuri writes that this species “deposits its eggs



FIGS. 390-392.—*CARETTA CARETTA*, YOUNG. NAT. SIZE. 390, SHELL FROM BELOW; 391, HEAD FROM SIDE; 392, UNDERSIDE OF HEAD. No. 14823, U.S.N.M.

on almost every suitable stretch of sandy beach in the southern half of Japan during the summer months of the year.” Special localities are mentioned by Okada as Awa, Hondo, and Toza, also

Bonin Islands, and Professor Mitsukuri collected a large number of eggs at Sagara, province of Totomi, for embryological studies. Dr. S. Nozawa writes me that the Pacific loggerhead occasionally straggles to Yezo.

Genus CHELONIA^a Latreille.

1800. *Chelonia* BRONGNIART, Bull. Soc. Philom. Paris, II, p. 89 (*nomen nudum*).
 1802. *Chelonia* LATREILLE, Hist. Nat. Rept., I, p. 22 (type, *C. mydas*).
 1806. *Chelone* BRONGNIART, Mém. Sav. Étrang., I, p. 610 (emendation).
 1814. *Chelonias* RAFINESQUE, Specchio Sci. (Palermo), II, No. 9, 1 Sett., 1814, p. 66 (emendation).
 1838. *Mydas* COCTEAU in Sagra's Hist. Fis. Pol. Nat. Cuba, IV, Rept., p. 22 (type *C. mydas*) (not of Fabricius, 1799).
 1843. *Mydasca* GERVAIS, Dict. d'Hist. Nat., III, p. 457 (same type).
 1845. *Euchelonia* TSCHUDI, Fauna Peruana (p. 22) (same type).
 1848. *Megomys* GISTEL, Naturg. Thier., p. viii (substitute for *Chelonia*).
 1858. *Euchelys* GIRARD, Herpet. U. S. Expl. Exped., p. 447 (type, *E. macropus*).

CHELONIA JAPONICA (Thunberg).

ARAUMIGAME.

1787. *Testudo japonica* THUNBERG, Svensk. Vetensk. Acad. Nya Handl., VIII, p. 178, pl. VII, fig. 1 (Japan).—*Chelonia japonica* SCHWEIGGER, Prodr. Mon. Chelon., 1814, p. 21.
 1814. *Chelonia virgata* SCHWEIGGER, Prodr. Mon. Chelon., p. 21 (seas of the Torrid Zone).—DUMÉRIL, Cat. Méth. Rept. Mus. Paris, I, 1851, p. 24 (Indian Ocean; New Guinea).—SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 221 (Formosa).—GUENTHER, Rept. Brit. India, 1864, p. 53 (Formosa).—GARMAN, Bull. U. S. Nat. Mus., No. 25, 1884, p. 302 (tropical portions of Western Pacific and Indian oceans).
 1820. *Caretta thunbergii* MERREM, Syst. Amph., p. 19.
 1831. *Chelonia mydas* ♂ *japonica* GRAY, Synops. Rept., I, p. 53 (Japan).
 1835. *Chelonia viridis* TEMMINCK and SCHLEGEL, Fauna Japon., Rept. (part; not of Schneider), p. 18 (part; Indian Archipelago to Japan), p. 139 (Japan); pl. IV, figs. 4, 5, 6 (Indian Ocean), pl. VI, figs. 1, 2 (Moluccas).—NAMIBE, Cat. Spec. Vert. Educ. Mus. Tokyo, 1881, p. 84 (Ogasawara shima).—OKADA, Cat. Vert. Japan, 1891, p. 72 (Bonin Is.: Koge shima).
 1860. *Chelonia mydas* HALLOWELL, Proc. Phil. Acad. 1860, p. 486 (not of Linnaeus) (Bonin Islands).—*Chelone mydas* BOULENGER, Cat. Chel. Brit. Mus., 1889, p. 180 (part; Malay Peninsula; Formosa; Bonin Islands).

An examination of Thunberg's plate^b leaves no doubt as to the identity of the sea-turtle he describes, the four pairs of costal shields and one pair of prefrontals being plainly shown and quite conclusive. The figure is not a bad one for the time and is easily identified. It has nothing to do with *Caretta olivacea*.


The "green" turtle is easily identified by the single pair of long prefrontal shields. The carapace has only four pairs of costal shields, like the hawksbill, but in the latter the dorsal shields are more or less imbricate, so that no confusion between the two species should occur.

^a From *χελώνη*, turtle.

^b Svensk. Vetensk. Acad. Nya Handl., VIII, 1787, pl. VII.

Several views of a young specimen from the Bonin Islands, just hatched (figs. 393-395), will help in the identification of this species.

Habitat.—Widely distributed in the tropical part of the Pacific



FIGS. 393-395. *CHELONIA JAPONICA*, YOUNG. NAT. SIZE. 393, ENTIRE ANIMAL FROM ABOVE; 394, SIDE OF HEAD; 395, PLASTRON. NO 7706, U.S.N.M.

Ocean, but apparently less common on the coasts of Japan than the other species, judging from the scant records available. It breeds on the Bonin Islands, and may be the prevailing species there.

List of specimens of Chelonia japonica.

U. S. N. M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
7705	Young.....	Port Lloyd, Bonin Islands.....	1854	W. Stimpson.
7706do. ^ado.....	1854	Do.
7707do.....do.....	1854	Do.
33858do.....	Baha shima, Bonin Islands.....	Mar., 1904	A. Owston.

^a Figs. 393-395.

Genus ERETMOCHELYS^a Fitzinger.

1828. *Caretta* RITGEN, Nova Acta Acad. Leop. Carol., XIV, p. 270 (type, *Chelonia imbricata*) (not of Rafinesque, 1814).
 1843. *Eretmochelys* FITZINGER, Syst. Rept., p. 30 (same type).
 1873. *Onychochelys* GRAY, Proc. Zool. Soc. London, 1873, p. 397 (type, *O. kraussi*).

ERETMOCHELYS SQUAMOSA^b (Girard).

TAIPEI.


1769. *Testudo imbricata* PENNANT, Indian Zool., (p. 87) (not of Linnaeus 1766).—*Chelonia imbricata* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., 1835, pp. 13, 139, pl. v, figs. 1-2 (Japan); pl. vi, fig. 4 (Moluccas).—OKADA, Cat. Vert. Japan, 1891, p. 72 (Okinawa shima).—*Chelone imbricata* STRAUCH, Chelon. Stud., 1862 (p. 181) (part); Mém. Acad. Sci. St. Petersb. (7), XXXVIII, No. 2, 1890, p. 121 (part: Carmen Isl. etc.).—BOULENGER, Cat. Chel. Brit. Mus., 1889, p. 183 (part: Formosa, etc.).
 1857. *Eretmochelys squamata* AGASSIZ, Contr. Nat. Hist. U. S. Am., I, p. 382 (Indian and Pacific oceans; Japan) (not *Testudo squamata* Gmelin).—GARMAN, Bull. U. S. Nat. Mus. No. 25, 1884, p. 300 (tropical Pacific and Indian oceans).—*Caretta squamata* SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 221 (Tamsui, Formosa).—GUENTHER, Rept. Brit. India, p. 54 (Formosa).
 1858. *Caretta squamata* GIRARD, Herpet. U. S. Expl. Exped., p. 442, pl. xxx, figs. 1-7 (Sulu Seas and Indian Ocean).
 1858. *Caretta rostrata* GIRARD, Herpet. U. S. Expl. Exped., p. 446, pl. xxx, figs. 8-13 (Fiji Is.; type in U. S. Nat. Mus.).

The hawksbill, or tortoise-shell turtle, has two pairs of prefrontals, like *Caretta olivacea*, but only four pairs of costal shields, like *Chelonia japonica*, the vertebral shields being more or less pointed behind and imbricate. The details of structure are shown in the figures of *E. imbricata* (figs. 396-400), and will aid in the identification in default of description and figures of Japanese specimens, of which there are none in our collection.

Habitat.—The capture of this species is an important fishery in the Riu Kius, and the tortoise-shell industry is one of considerable magnitude in Nagasaki. That the hawksbill occasionally straggles farther north is evidenced by its having been taken once off Nemuro, north-east Yezo, according to Doctor Nozawa.

^a From ἐρετυόν, oar; χέλυς, turtle.

^b Signifying scaly, with reference to the imbrication of the dorsal shields.



FIGS. 396-399.—*ERETMOCHELYS IMBRICATA*. $\frac{1}{2} \times$ NAT. SIZE. 396, SHELL FROM BELOW. $\frac{1}{2} \times$ NAT. SIZE. 397, TOP OF HEAD; 398, SIDE OF HEAD; 399, RIGHT FORE FLIPPER, DORSAL VIEW; (No. 25645). U.S.N.M.

Being a tropical species of wide distribution in the eastern Pacific, it is unnecessary to point out that it also occurs in Formosa.




FIG. 400.—ERETMOCHELYS IMBRICATA. $\frac{1}{2}$ \times NAT. SIZE. SHELL FROM ABOVE. No. 28645, U.S.N.M.

Suborder CHILOTÆ.^a

1826. *Trionychoidea* FITZINGER, Neue Class. Rept., p. 7.
 1832. *Chilotæ* WIEGMANN, Handb. Zool., p. 167.
 1835. *Labiata* FITZINGER, Ann. Wien Mus., I, p. 107.
 1837. *Chiloti* BURMEISTER, Handb. Naturg., p. 731.
 1880. *Peltochelyidæ* SEELEY, Quart. Journ. Geol. Soc., XXXVI, p. 412.
 1887. *Diacostoidea* BAUR, Zool. Anz., 1887, p. 99.
 1889. *Trionychia* ZITTEL, Palæozool., III, p. 513.
 1894. *Mecraspedita* VAILLANT, Ann. Sc. Nat. (7), Zool. XVI, Nos. 4-6, p —.

This suborder consists of only one family, the soft-shelled turtles.

^a From the Greek χείλος, lip.

Family TRIONYCHIDÆ.

Turtles covered with a soft, leathery skin; lips fleshy; nostrils at the end of a flexible proboscis; toes webbed, with three claws.

Distributed over the warm and temperate regions of the earth, except Australia and South America, inhabiting lakes, rivers, and swamps. They are exceedingly voracious and fierce, so that they are often known as "snapping turtles." Some species reach a large size and several are highly esteemed as food. There are about 30 species known.

Genus AMYDA ^a Oken.

1816. *Amyda* OKEN, Lehrb. Zool., II, p. 348 (type, *Trionyx euphraticus*).
1830. *Aspidonectes* WAGLER, Nat. Syst. Amphib., p. 134 (type, *T. ægyptiacus* = *T. triumquis*).
1835. *Gymnopus* DUMÉRIEIL and BIBRON, Erpét. Gén., II, p. 472 (substitute for *Aspidonectes*).
1835. *Platypeltis* FITZINGER, Ann. Wien Mus., I, pp. 120, 127 (type, *T. feror*).
1835. *Pelodiscus* FITZINGER, Ann. Wien Mus., I, pp. 120, 127 (type, *T. sinensis*).
1843. *Potamochelys* FITZINGER, Syst. Rept., p. 30 (type, *T. javanicus*).
1844. *Tyroc* GRAY, Cat. Tortois. Brit. Mus., p. 47 (type, *T. nilotica* = *T. triumquis*).
1844. *Trionyx* GRAY, Cat. Tortois. Brit. Mus., p. 49 (type, *T. feror*) (not of Geoffroy-Oken).
1864. *Rafetus* GRAY, Proc. Zool. Soc. London, 1864 (p. 81) (type, *T. euphraticus*).
1864. *Aspilus* GRAY, Proc. Zool. Soc. London, 1864 (p. 83) (type, *T. cariniferus* = *T. cartilagineus*).
1869. *Landemania* GRAY, Proc. Zool. Soc. London, 1869, p. 215 (type, *L. irrorata* = *T. sinensis*).
1869. *Fordia* GRAY, Proc. Zool. Soc. London, 1869, p. 219 (type, *F. africana* = *T. trionyx*).
1869. *Callinia* GRAY, Proc. Zool. Soc. London, 1869, p. 221 (type, *T. spiniferus*).
1873. *Isola* GRAY, Proc. Zool. Soc. London, 1873, p. 51 (type, *I. pequensis* = *T. formosus*).
1873. *Ida* GRAY, Proc. Zool. Soc. London, 1873, p. 55 (type, *I. ornata* = *T. cartilagineus*).
1873. *Oscaria* GRAY, Ann. Mag. Nat. Hist. (4), XII, (p. 157) (type, *O. swinhoci*).
1880. *Yuen* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 18 (type, *Y. leprosus* = *T. swinhoci*).
1880. *Psilognathus* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 24 (type, *P. lavis*).
1880. *Temnognathus* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 25 (type, *T. mordax*).
1880. *Gomphopelta* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 27 (type, *G. officina*).
1880. *Coclognathus* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 29 (type, *C. noremcostatus*) (not of Hessling, 1852).
1880. *Tortisternum* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 31 (type, *T. noremcostatum*).
1880. *Ceramopelta* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 33 (type, *C. latirostris*).
1880. *Coptopelta* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 34 (type, *C. septemcostata*).
1880. *Cinetisternum* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 36 (type, *C. bicinctum*).

^aThis name is of uncertain origin, but apparently a variant of ἔμυς, a river turtle.

The reasons for the adoption of the generic term *Amyda* in preference to *Aspidonectis* or *Trionyx* I have advanced in my article on "Generic names of soft-shelled turtles."^a

To complete the history of the name *Amyda* it may be added that Geoffroy St. Hilaire, in 1809, in the article creating the generic term *Trionyx*, says that Schweigger, in a manuscript communicated to the French "Institut," had given the name *Amyda javanica* to the species called *Trionyx javanicus* by himself. This reference might suggest to some the propriety of crediting *Amyda* to Schweigger, 1809, with *A. javanica* (= *cartilaginea*) for type. The result, however, will be the same as here arrived at, since the latter species is congeneric with the species above adopted as the type for Oken's *Amyda*.

AMYDA JAPONICA (Temminck and Schlegel).

SUPPON.

Plate XXXV.

1835. *Trionyx stellatus* var. *japon[icus]* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., p. 32, pl. v, fig. 7; pl. vii (Japan).
1835. *Trionyx japonicus* TEMMINCK and SCHLEGEL, Fauna Japon., Rept., p. 139.—SCHLEGEL, Abb. Amph., 1840, p. 108, pl. XXXI (rivers of southern Japan).—HILGENDORF, Sitz. Ber. Ges. Natur. Fr. Berlin, 1880 p. 112 (southern Japan).—MITSUKURI and ISHIKAWA, Quart. Jour. Micr. Sci. (n. s.), XXVII, Aug., 1886, p. 17.—OKADA, Cat. Vert. Japan, 1891, p. 72 (Kagoshima; Chikugogawa).—MITSUKURI, Zool. Mag. Tokyo, VII, 1895, p. 143 (oviposition); Bull. Fish. Bur. Washington, XXIV, 1905, p. 260, pls. i-iii (turtle farm).
1866. *Trionyx schlegelii* MARTENS, Preuss. Exped. Ost-Asien, Zool., I, p. 112 (Yokohama) (not of Brandt, 1857).

The status of the soft-shelled turtles inhabiting China and Formosa (*A. sinensis* and *schlegelii*), Japan (*A. japonica*) and Amurland (*A. maackii*) has not been worked out for lack of material. Boulenger who unites all three, had apparently no Japanese specimens at hand when he published his catalogue of the Chelonians. Hilgendorf, however, who had both Chinese and Japanese specimens, declares them to be different. He says that the young of *A. sinensis*, compared with Japanese specimens of the same age, have "a much stronger median dorsal keel and different proportions of length between the broader (basal) part of the ribs and the narrower (apical) end. The basal portion in *sinensis* is shorter than the apical part, but longer in *japonicus*."

My own material is none too conclusive, for while I have quite a number of young Japanese specimens I have only one specimen from the mainland and two from Formosa. The Chinese specimen before me is from the country between Tientsin and Peking, consequently

^a Science (n. s.), XXI, Feb. 10, 1905, pp. 228-229.

practically from the same locality whence came Brandt's type of *Trionyx schlegelii*. It follows that if Brandt's *T. maackii* from the Amur region is a distinct form, it is different from our Chinese specimen also.

The absence of specimens from the drainage of the Yangtse River, which may be supposed to represent the true *Amyda sinensis*, is particularly to be regretted, as it prevents me from arriving at any but the most inconclusive and preliminary results.

That my specimens represent three separable forms, however, I have but little doubt.

In the first place, in the Formosan specimens, which I call *A. sinensis*, the shell is evenly arched, without any depression on either side of the neural plates and without any median raised keel. The north China specimen, as well as all my Japanese specimens, agree, on the other hand, in having a well-marked depression on each side of the neural plates of the upper shell, so that a broad and blunt but very distinct keel is formed along the median line. In addition, the Formosan specimens seem to have the dermal ridges on the back smooth and continuous, while in the north China and Japan specimens the great majority of these ridges are dissolved into lines of tubercles.

To demonstrate the difference between the north China specimen, which I call *A. schlegelii*, and the Japanese specimens (*A. japonica*) the following table is presented:

Comparative measurements of Japanese and Chinese soft-shelled turtles.

U. S. N. M. No.	Sex.	Age.	Locality.	Species.	Direct measurements.				Proportionate measurements.			
					Length of carapace.	Breadth of carapace.	Length of plastron.	Depth of shell.	Length of carapace.	Breadth of carapace.	Length of plastron.	Depth of shell.
21179	Male	4 years	Japan	<i>A. japonica</i>	170	133	124	43				
21178	do	3 years	do	do	112	98	90	28				
34057	do	do	do	do	136	115	102	33				
34058	do	do	do	do	133	111	104	35				
34060	do	do	do	do	128	112	95	33				
Average					136	114	103	34.5	100	84	76	25
34056	Female	3 years	Japan	<i>A. japonica</i>	130	116	100	38				
34059	do	do	do	do	140	116	103	43				
Average					135	116	101.5	40.5	100	86	75	30
34054	Male	4 years	Formosa	<i>A. sinensis</i>	165		136	45	100		82	27
34055	Female	do	do	do	150	120	118	51	100	80	79	34
29700	Male	3 years	N. China	<i>A. schlegelii</i>	119	95	93	41	100	80	78	34

In the first four columns I have given the exact measurements of the membranous shell, and in the last four columns I have reduced the same dimensions to per cent of the length of the membranous carapace. Although the series is small, the figures probably deserve some confidence because of their remarkable uniformity and harmony. If we examine the table, the first observation to impress itself upon us is that the males and females from the same locality differ from each other but very little in their proportions, except in the depth of the shell—i. e., the greatest distance between ventral and dorsal surfaces on the middle line measured with curved calipers, the difference amounting to 5 and 7 per cent, respectively. At the same time it should be noted that the difference between the various individuals of the same sex does not reach $2\frac{1}{2}$ per cent.

If we now compare the north China specimens with the Japanese specimens, we observe that in the males the depth of the shell of the former is 9 per cent greater than in the latter. Unfortunately we have no female specimen from northern China; but the difference between the island specimens and the mainland male is so great that the latter even exceed the females of the former by 4 per cent. The difference may be very strikingly formulated by stating that in the males of *A. japonica* before me the depth of the shell is one-fourth the length of the carapace, while in the male *A. schlegelii* it is one-third the length of the carapace.

The Formosan specimens appear to be slightly narrower and also somewhat deeper than the Japanese without even approaching the north China specimen in this respect, inasmuch as the latter, a male, presents exactly the same proportions as the female of the more southern form.

The Formosan and north China specimens agree in the lateral outline of the shell, for while in the Japanese specimens the greatest width of the body is situated more nearly at the middle, in the former the greatest width is farther back.

The material is so scant that I shall pass over another point wherein the Formosan specimens and the north China one seem to differ from the Japanese, viz, in the somewhat shorter plastron of the latter. Compared with the width of the body, the difference amounts to nearly 10 per cent; but this measurement is less satisfactory than the the others, and, as stated, the series is too small to give positive results.

It would then appear that the Japanese soft-shelled turtles are neither identical with the north Chinese nor with the Formosan forms. The question therefore naturally arises, whether it may not agree with the south Chinese species. As said before, I have no specimens for comparison, and the literature offers very little of a definite nature. Both Gray and Boulenger, examining Formosan as well as Chinese

specimens, declared them to be identical, and in default of material I accept their decision. In that case, the Japanese form must be regarded as distinct. On the other hand, I must call attention to a point which later may have a bearing upon these questions. Doctor Siebenrock, in a very interesting paper on the systematic arrangement of the *Trionychidæ*^a figures the plastral arrangement of a *Trionyx sinensis* (p. 821), which shows a very short plastron, especially the posterior lobe, and he calls particular attention to this feature as characteristic of *T. sinensis* (p. 822). The figure (see fig. 401) and




FIG. 401.—ELEMENTS OF THE TRIONYCHID PLAGSTRON, SHOWING PLAGSTRAL CALLOSITIES. $\frac{1}{4}$ X NAT. SIZE. (FROM SIEBENROCK.) *c.* COMMISSURE; *ep.* ENTOPLAGSTRON; *cyp.* EPIPLAGSTRON; *hp.* HYOPLAGSTRON; *hyp.* HYPOPLAGSTRON; *pa.* ANTERIOR MEDIAN PROCESS; *pm.* MEDIAN PROCESS; *pp.* POSTERIOR MEDIAN PROCESS; *xp.* XIPHIPLAGSTRON.

description fit exactly my Japanese specimens, but unfortunately Doctor Siebenrock does not give the locality of the specimen figured. Now, on the other hand, compare the plastron of the specimens figured by Gray.^b It is an exact counterpart of my Formosan male (No. 34054) in every respect, even to the close approximation of the epiplastra. It is not easy to believe that Gray's and Siebenrock's figures represent the same species.

However, with no certainty as to the exact locality of the specimens figured, it is idle to speculate upon the status of the soft-shelled turtles in China, though it would not be surprising if each of the three main river systems of that empire had its own form. This view is strengthened by some remarks by Doctor Schnee^c to the effect that it is possible to distinguish between a northern and a southern type, the latter being much darker in China and with more strongly developed tubercles than the former, peculiarities which he verified on large series of living specimens in Canton and Shanghai.^d

Doctor Schnee in the same article (p. 207) has a note which will

^a Zur Systematik der Schildkrötenfamilie Trionychidæ Bell. nebst der Beschreibung einer neuen Cyclanorbis-Art, von Friedrich Siebenrock, in Sitz. Ber. Akad. Wiss. Wien, Math.-Naturw. Cl., XCI, Pt. 1, Oct. 1902, pp. 807-846.

^b Cat. Shield. Rept. Brit. Mus., 1855, pl. xxxi.

^c Zeitschr. Naturwiss., LXXII, 1899, pp. 203-204.

^d This difference in color may be due to the character of the rivers and lakes, as already suggested by Heude (Mém. sur les Trionyx, p. 1)

have to be taken into account in all future examinations and comparisons of Japanese specimens. He says that the price of the soft-shelled turtle in Japan is so much higher than in China that a regular importation takes place from China to Japan, and that therefore he is not quite sure that the specimen which was shown him there really came from Japan. It is plain then that in order to settle the question of the status of the Japanese *Amyda* it will be necessary to show that specimens identical with those from China are in reality indigenous to Japan.^a

Before leaving this theme of the characterization of various forms or species confounded under *A. sinensis*, I wish to call the attention of future investigators to the somewhat fugitive pattern of the plastron as a possible means of distinguishing the young 2 years old and under.



Doctor Siebenrock^b has described such a pattern on six specimens of *A. sinensis*, from Annam, scarcely a year old: "Anteriorly in the middle there is an unpaired gular spot, which is the smallest or even may be absent. The gular edge, moreover, has a black margin. Furthermore, on the hyo-hyoplastral interspace there are two spots which usually unite into a short transverse band, and behind this there is again a large unpaired spot between the xiphiplastra. Laterally there is on both sides a spot in the axillary and inguinal regions, one in front of the hyoplastron and another behind the axillary one."

I find this pattern repeated in my Japanese specimens, of which I have a large series, with one very notable exception, viz, there is a single unpaired median mark on the hyo-hyoplastral interspace, and not two lateral marks separated or united. I refer to Plate XXXV illustrating this series, and it will be noted that instead of "two spots, usually uniting into a short transverse band on the hyo-hyoplastral interspace," there is a large, unpaired median, triangular or broadly spear-shaped spot on the same interspace. The essential element of this large spot, the largest by far, is the median portion, which forms the anterior angle of the triangle, since this part persists even in cases where the lateral extensions have not been fully developed. The arrangement, consequently, appears to be quite the reverse of that of the Annamese specimens and is not without considerable significance.

^a In this connection it may be well to refer to an observation by Dr. John Anderson, (Zool. Res. Exped. West. Yunnan, I. 1879, p. 792) to the effect that "there are apparently two species of *Trionyx* found in Japan, one corresponding to the supposed foregoing variety of *T. stellatus*, Geoff., but which is identical with *T. procellatus*, which is the *T. sinensis*, Wieg., and which appears to have been more recently redescribed by Brandt under the name of *T. schlegeli*. . . . The other *Trionyx* found in Japan besides *T. sinensis*, Wieg., is a form allied to *T. javanicus*, Geoffr. and which Gray first referred to *D. subplana*, and which has been figured by Schlegel under the name of *T. japonicus*.

^b Sitz. Ber. Akad. Wiss. Wien, Math. Naturw. Cl., CXII, Pt. 1, May, 1903, p. 349.


Description (figs. 402-405).—*Male*, 4 years old; U.S.N.M. No. 21179; Japan; Imperial Government collection. Head moderate pointed, the proboscis as long as width of interorbital space and one-half the upper eyelid; nostrils separated by a narrow septum, on each



FIGS. 402-403.—*AMYDA JAPONICA*. $\frac{1}{3}$ X NAT. SIZE. 402, SIDE VIEW; 403, VIEW FROM ABOVE. No. 21177, U.S.N.M.

side of which a papilla is visible near the opening; interorbital space equals one-half the horizontal diameter of orbit; carapace oval; the anterior margin of the revolute nuchal border somewhat rugose with two small tubercles on the median line; along the median line of the

shell a raised keel anteriorly as broad as interorbital space and eyelids together, posteriorly not wider than this space plus half an eyelid; on each side of the keel a perceptible depression, the skin of the carapace with numerous more or less disrupted, raised, longitudinal lines studded with tubercles, these lines being more wavy and parallel with the axis of the body near the middle of the back, and straighter, less tubercular, and more parallel with the periphery toward the edge of the carapace; the tubercles gather into a large cluster on the nuchal



FIGS. 404-405.—*AMYDA JAPONICA*. ♀ NAT. SIZE. 404, UNDERSIDE (14 × NAT. SIZE); 405, SIDE OF HEAD. No. 21177, U.S.N.M.

portion of the keel, and a still greater number of larger tubercles congregate near the posterior end of the bony disk and the adjoining posterior portion of the soft flap; plastron flat, smooth, the posterior lobe short, its width at base equaling one-half the length of the plastron; greatest height of body equals one-fourth the length of carapace. Color (in alcohol) above clear greenish olive, with a few indistinct brownish blotches on the carapace, underneath uniform pale buff; head with a number of narrow blackish lines, as follows: A transverse

line across anterior third of upper eyelid and interorbital space to anterior corner of eye and from there obliquely downward across the loreal region to the upper lip at the base of the proboscis; a slightly curved line from below center of eye obliquely backward above the angle of the mouth disappearing some distance behind and below the latter; a slightly curved line from posterior angle of eye above the temporal region; a short blackish streak below the eye between the last two lines; two broken lines across the middle and the posterior




FIG. 406.—*AMYDA SCHLEGELII*. SIDE VIEW. FROM GRAY'S CAT. SHIELD REPT. PL. XXXI.

third of upper eyelid; a few broken longitudinal lines on parietal and occipital regions and a few dusky spots on snout and proboscis; lips, chin, and throat with large round buff spots surrounded by a network of blackish brown lines; sides of neck with a broad longitudinal buff band, more or less interrupted anteriorly and faintly edged with dusky; neck below this band suffused with olive and dotted with indistinct roundish pale buff spots.

Dimensions.

	<i>mm.</i>
Length of carapace.....	170
Greatest width of carapace.....	133
Length of plastron.....	124
Greatest depth of shell.....	43
Greatest width of head.....	27
Distance from plastron to tip of tail.....	73
Free portion of tail from connection with carapace.....	32

The *females* differ chiefly in proportions, being much deeper through the body and having a much shorter tail.

Young recently hatched; U.S.N.M. No. 23531; Yokohama; October, 1896; L. Stejneger, collector (plate XXXV, fig. 3). Essentially as the one described above; the anterior border of the carapace turned over as in the older specimens, but the median portion, instead of being merely rugose, is more or less regularly segmented, and the posterior edge of several of the segments tuberculated, the median segment being pushed a little ahead of the others and the tubercle occupying the upper surface instead of the posterior edge; dorsal keel very prominent. Color essentially as above, but extreme edge of



1



4



8



2



5



9



6



3



7



10

AMYDA JAPONICA. YOUNG

FOR EXPLANATION OF PLATE SEE PAGE 556.

carapace devoid of color, though without a well-defined pale margin; underside with a number of symmetrical blackish spots, with slightly paler centers, as follows: (1) On plastron, a small longitudinal spot anteriorly on the median line between the epiplastra; a large triangular median spot on the hyo-hypoplastral interspace, broader than high, the apex turned forward; a heart-shaped median spot, deeply notched in front, on the anterior processes of the xiphiplastra; a pair of oval spots, one on each side in front of the bend of the hyoplastra; a pair of roundish spots, one on each side in the outer angle between the hyoplastra and the hypoplastra; (2) on the soft skin, a pair of oblique elliptic spots, one on each side behind the bend of the hypoplastra; a pair of circular spots at the base of the tail; (3) on the underside of the flap of the carapace: a longitudinal, ill-defined broad band interrupted on the median line; (4) on the extremities: palms and soles dark gray.

Dimensions.

	<i>mm.</i>
Length of carapace	36
Greatest width of carapace	34
Length of plastron	29
Greatest depth of shell	12

The rate of growth is well illustrated by a table presented by Professor Mitsukuri,^a as follows:

Age.	Length.	Breadth.	Weight.
	<i>Cm.</i>	<i>Cm.</i>	<i>Grams.</i>
Just hatched.....	2.5	2.5	23
First year.....	4.5	4.2	23
Second year.....	10.5	8.8	169
Third year.....	12.5	10.5	300
Fourth year.....	16.0	13.5	563
Fifth year.....	17.5	15.1	750

Habitat.—This species, as here understood, has only been found in Japan proper, where it seems to occur originally in the southern rivers only. It has probably been introduced in many places, and a detailed account of the localities in which it is undoubtedly original would be a very interesting contribution to our knowledge of these animals.

Professor Mitsukuri has given a very interesting and instructive description of Mr. Hattori's turtle farm at Fukagawa, a suburb of Tokyo, which it was my good fortune to inspect in 1896 under his guidance.

^a Bull. Fish. Bur., Washington, XXIV, 1905, p. 265.

List of specimens of *Amyda japonica*.

U.S. N.M. No.	Sex and age.	Locality.	When collected.	By whom collected or from whom received.
21171	Young <i>a</i>	Japan		Imp. Jap. Governm.
21172	do. <i>a</i>	do.		Do.
21173	do. <i>a</i>	do.		Do.
21174	do. <i>a</i>	do.		Do.
21175	do. <i>a</i>	do.		Do.
21176	do. <i>b</i>	do.		Do.
21177	do. <i>c</i>	do.		Do.
21178	Male <i>d</i>	do.		Do.
21179	do. <i>c</i>	do.		Do.
23528	Young <i>a</i>	Yokohama	Oct. —, 1896	L. Stejneger.
23529	do. <i>f</i>	do.	do.	Do.
23530	do. <i>g</i>	do.	do.	Do.
23531	do. <i>h</i>	do.	do.	Do.
23532	do.	do.	do.	Do.
23533	do.	do.	do.	Do.
23534	do.	do.	do.	Do.
23535	do.	do.	do.	Do.
23536	do.	do.	do.	Do.
23537	do.	do.	do.	Do.
23538	do.	do.	do.	Do.
34056	Female <i>d</i>	Tokyo	Mar., 1903	A. Owrston.
34057	Male <i>d</i>	do.	do.	Do.
34058	do. <i>d</i>	do.	do.	Do.
34059	Female <i>d</i>	do.	do.	Do.
34060	Male <i>d</i>	do.	do.	Do.
34061	do. <i>b</i>	do.	Feb. 23, 1904	Do.
34062	Young <i>b</i>	do.	do.	Do.
34063	do. <i>a</i>	do.	do.	Do.
34064	do. <i>a</i>	do.	do.	Do.
34065	do. <i>a</i>	do.	do.	Do.

a First year.*b* Second year.*c* Second year; figs. 402-405.*d* Third year.*e* Probably 4 years old; description, p. 520.*f* First year; pl. xxxv, fig. 9.*g* First year; pl. xxxv, fig. 2.*h* First year; pl. xxxv, fig. 3; description, p. 522.AMYDA SINENSIS *a* (Wiegmann).

1834. *Trionyx (Aspidonectes) sinensis* WIEGMANN, Nova Acta Acad. Leop. Carol. XVII, 1834, p. 189. (type-locality, paddyfield on a small island in the Tiger River, near Macao, China; MEYER, collector).—*Trionyx sinensis* STRAUCH, Chelon. Studien, 1862 (p. 177).—SWINHOE, Ann. Mag. Nat. Hist. (3), XII, 1863, p. 219 (S. W. Formosa).—SCHNEE, Zeitschr. Naturwiss., LXXII, 1899, p. 202 (Shanghai; Canton).—NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, No. 1, 1905, p. 20 (part).
1842. *Trionyx tuberculatus* CANTOR, Ann. Mag. Nat. Hist., IX, p. 482 (Chusan, China).
1844. *Tyrse perocellata* GRAY, Cat. Tortois. Brit. Mus., p. 48 (type-locality Canton, China; types in Brit. Mus.; Cantor, collector).—*Trionyx perocellatus*

a Signifying Chinese, Sina being another form for China.

- GRAY, Cat. Shield Rept. Brit. Mus., 1855, p. 65 (China).—*Landemannia ? perocellata* GRAY, Proc. Zool. Soc. London, 1869, p. 216 (Formosa, Swinhoe, collector); 1873, pp. 53, 54 (fig. of mandibles).
1869. *Landemannia irrorata* GRAY, Proc. Zool. Soc. London, 1869, p. 216 (type-locality, Shanghai, China; type in Brit. Mus.).
1880. *Psilognathus larvis* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 24, pl. II (type-locality, mountain and streams south of the town of Ning-kouo fou.)
1880. *Tennognathus mordax* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 26, pl. III (type-locality, environs of Shanghai).
1880. *Gomphopelta officina* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 27, pl. IV (type-locality, River Houai, where it leaves the province Ho-nan).
1880. *Cælognathus novem-costatus* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 29, pl. V (type-locality, eastern extremity of lake Tch'ao).
1880. *Tortisternum novem-costatum* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 31, pl. VI (type-locality, lake Tch'ao, department of Lu-tcheou).
1880. *Ceramopelta latirostris* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 33, pl. VII (type-locality, environs of Ngan-k'ing fou).
1880. *Coptopelta septem-costata* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 35, pl. VIII (type-locality, lakes of Tong-lieou (Ngan-houé)).
1880. *Cinctisternum bicinctum* HEUDE, Mém. Hist. Nat. Emp. Chinois, I, p. 37, pl. IX (type-locality, irrigation ditches in the marsh of Ngan-k'ing).

The interesting question as to the relationship of the individuals with narrow or wide alveolar surfaces of the jaws, as illustrated by Père Heude's subdivision of the south Chinese species, can not be discussed here. It must be studied on the spot with proper regard to the possibility of complications, due to local differentiations in the various drainage systems as well as to the subsequent mixing of the latter through irrigation canals. Boulenger has suggested "a case of dimorphism caused by a difference of diet," the fish-eating individuals retaining the sharp beak of the young, while those feeding on mussels with hard shells or on other hard substances are supposed to gradually develop a broad crushing mandibular surface. He furthermore supposes such individuals to interbreed, thus preventing the characters from becoming permanent.^a

With regard to the identity of the Formosan and south Chinese specimens and the related question of the distinctness of the latter I refer to the discussion under *A. japonica* (pp. 517-519).

Description.—*Female*, 4 years old; U.S.N.M. No. 34055; Tamsui River, Formosa; March, 1903; A. Owston collection. Carapace short ovate in outline; the anterior margin of the revolute nuchal border segmented toward the median line, a single tubercle on the latter in front of the border; no median keel, nor lateral depression, the carapace descending evenly to both sides from the rounded back; skin of carapace smooth, with about 24 longitudinal raised lines, which are entirely untuberculated, and nearly all straight and uninterrupted, except about three midway on each side, which are wavy and occasionally interrupted; these lines terminate near the

^a Cat. Chel. Brit. Mus., 1889, pp. 242-244.

posterior end of the bony disk in a small tubercle behind which to the end of the carapace there are numerous short raised lines similarly terminated; a couple of rows of blunt tubercles in front of the bony disk on the soft flap behind the overturned nuchal border; plastral bones more or less discernible on underside as well as the callosities, of which there is a pair on the hyo-hyoplastra and one on each of the xiphiplastra. Color (in alcohol) above dark olive slate, underneath drab; head with a few scattered blackish dots and traces of a oculo-supratemporal black line, a subocular and an anterior oculo-labial line; neck below dusky with indistinct pale spots on throat and traces of a lateral pale band; underside with a few dark blotches, the one filling the outer angle between the hyoplastra and the hypoplastra as well as the median blotch on the xiphiplastra being most conspicuous; palms and soles dusky.

Dimensions.

	<i>mm.</i>
Length of carapace.....	150
Greatest width of carapace	120
Length of plastron.....	118
Greatest depth of shell.....	51
Greatest width of head.....	29
Distance from plastron to tip of tail.....	48
Free portion of tail, from connection with carapace.....	18

The *male* of the corresponding age (U.S.N.M. No. 34054) differs mostly in less depth of shell and longer tail. Eight pairs of costal plates can be made out; in addition to the callosities on hyo-hyoplastra and xiphiplastra there are indications of rugosities on each arm of the entoplastron. No trace of keel or depression on carapace.

Habitat.—It is impossible at the present stage of our knowledge to indicate the precise range of this species in China. It seems to occur in all the southern rivers and in Formosa; but how far north it extends or where it meets the next species must be left to the future to clear up.

List of specimens of Amyda sinensis.

U.S. N.M. No.	Sex.	Locality.	When collected.	By whom collected or from whom received.
34054	Male ^a	Formosa.....	May 10, 1896	A. Owston.
34055	Female ^b	Tamsui Riv., Formosa.....	Mar. —, 1903	Do.

^a Probably four years old; p. 526. ^b Probably four years old; description p. 525.

AMYDA SCHLEGELII (Brandt).

1855. *Trionyx perocellatus* GRAY, Cat. Shield Rept. Brit. Mus., pl. xxxi^a (China; (not of 1844).

1857. *Trionyx schlegelii* BRANDT, Mém. Biol. Acad. Sci. St. Pétersbourg. II, p. 610 (type-locality, Peking, China; type Mus. St. Petersb., no. 177; A. Bunge collector).

^a See fig. 406, on p. 522, of this work.

Reference has already been made to the specimen from northern China, to which I apply the above name more because the locality is the same as that of the type than because the description applies better to it than any of the others. It is quite likely that eventually *A. maackii* may turn out to be the same thing as the Peking form, in which case that name will take precedence. At present, however, I deem it the wisest course to enumerate them separately.

If Ford's elegant figures of the half-grown specimen in British Museum, from "China,"^a are as accurate as they are artistic, they




FIG. 407.—*AMYDA SCHLEGELII*. $\frac{1}{2}$ \times NAT. SIZE. VIEW FROM ABOVE. NO. 29700, U.S.N.M.

probably represent an individual of the present species. The dorsal keel, the tubercular ridges, and the coloration agree completely with our north China specimen, and especially the great depth of the body as shown in the upper figure. Expressed in per cent of the total length of the carapace, the depth is nearly 36, or in excess of the corresponding percentage in our male specimen of *A. schlegelii* (see table p. 516).

Description.—*Male, third year*; U.S.N.M. No. 29700; north China, between Tientsin and Peking; 1901; M. L. Robb, collector (figs.

^a Cat. Shield Rept. Brit. Mus., 1855, pl. xxxi. Outline of upper figure reproduced in this work, fig. 406, on p. 522.

407-409). Carapace short ovate, with a broad, raised median keel and a well-marked depression on each side of the latter; anterior border turned over backward, the outer (anterior) edge being smooth,




FIG. 408.—*AMYDA SCHLEGELII*. $\frac{1}{2}$ \times NAT. SIZE. UNDERSIDE. No. 29700, U.S.N.M.

the inner edge near the median line somewhat tuberculate with a couple of rounded tubercles on top of the rim at the middle; carapace with numerous strongly tuberculated longitudinal lines, the outer




FIG. 409.—*AMYDA SCHLEGELII*. $\frac{1}{2}$ \times NAT. SIZE. SIDE VIEW. No. 29700, U.S.N.M.

ones more or less continuous and parallel with the outer edge, the inner ones very irregular and interrupted; posterior end densely tuberculated, as is also the nuchal region; depth of shell about one-

third the length of carapace; plastron smooth. Color (in alcohol) above olive gray with a few small blackish spots, below pale buff, unspotted; on head a narrow black line across the interorbital space anteriorly continued below as a slightly oblique line between anterior angle of eye and edge of upper lip; a similar line behind the eye over the supratemporal region; sides of neck and throat olive with indistinct pale spots and traces of a longitudinal, lateral pale band on neck.

Dimensions.

	<i>mm.</i>
Length of carapace	119
Greatest width of carapace.....	95
Length of plastron	93
Greatest depth of shell	41
Greatest width of head	21
Distance from plastron to tip of tail.....	43
Free portion of tail, from connection with carapace.....	25

Habitat.—Northern China, the specimen described being from the province of Pechili. It is impossible at the present time to determine its range even approximately.

List of specimens of Amyda schlegelii.

U. S. N. M. No.	Sex.	Locality.	When collected.	By whom collected.
29700	Male σ	North China.....	1901	M. L. Robb.

σ Third year; description, p. 527; figs. 407-409.

AMYDA MAACKII σ (Brandt).

1857. *Trionyx maackii* BRANDT, Mém. Biol. Acad. Sci. St. Petersburg, II, p. 609 (type-locality, Southern Amur R. and Ussuri R.; types, Mus. St. Petersburg., Nos. 3690, 4661, 5725, 5726; Maack, collector).

1905. *Trionyx sinensis* NIKOLSKI, Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, p. 20 (part: Ussuri, Amurland).

This form, or species, being known only from the original description of Brandt, I have appended the following translation:

Description.—Upper part of head more convex than in *Trionyx javanicus*, olive colored with very numerous yellow dots which are smallest on the upper surface of the face, a little larger on the sides and much larger on the throat (but with no bands or dusky spots); sides of neck with a few rough places; dorsal shield [carapace] in the middle longitudinally depressed, but subcarinate, convex at the sides, with very smooth skin [“cute glaberrima”], olivaceous with minute yellow or more or less orange dots like the feet; anterior

σ For R. Maack, a Russian, who traveled in the Amur region in the early fifties, and who collected the type.

margin of dorsal shield thickened, not tuberculated; posterior margin of sternal shield [plastron] short, semilunar, rounded.

Dimensions.—Length of carapace, 353 mm; greatest width, 306 mm.

Habitat.—Found in the southern tributaries of the Amur River, especially in the Sungari and Ussuri rivers, as well as in the Amur River itself between the mouths of the tributaries mentioned.

According to Nikolski there are in the St. Petersburg Museum specimens from the above rivers and also from Lake Khanka. Whether the specimen from Korea (No. 9447) belongs to this form it is impossible to say.

[Genus *DOGANIA* ^a (Gray).]

1836. *Amyda* BONAPARTE, Chelon. Tab. Anal., p. 8 (type, *Trionyx subplanus*) (not of Oken, 1816.)

1844. *Dogania* GRAY, Cat. Tortois. Brit. Mus., p. 49 (type, *Trionyx subplanus*).

1869. *Sarberia* GRAY, Proc. Zool. Soc. London, 1869 (p. 200) (type, *Trionyx frenatus*).

The genus *Dogania* differs from *Amyda* in having all the pleuralia separated by the neuralia, while in *Amyda* the last pair of pleuralia are in contact behind the neuralia. Doctor Siebenrock has recently shown^b that *D. subplana* is unique in lacking the median process of the hypoplastron. The head is also comparatively much larger and the body much more depressed, the body being almost concave with no median keel.

The occurrence of *Dogania subplana*—or a related species—in Formosa is very problematical. I can not find that any specimens have been examined and the only foundation for the record seems to be the following statement by Mr. Gray:^c "Mr. Swinhoe informs me that this animal is common in the rivers of China and Formosa." Swinhoe most likely confounded it with *Amyda sinensis*. Before that time Gray had also catalogued a specimen received from the Leiden Museum as from "Japan,"^d but in Boulenger's Catalogue^e this specimen is enumerated as from "Java," which is most likely correct.

A description and references are appended, however, in order to facilitate its identification should it really occur in Formosa.

I may add that there is a specimen in the Hamburg Museum (No. 110), acquired in 1877 and identified as "*Trionyx subplanus*." It is said to be from "Japan," but no reliance can be placed on the record.

^aA coined word without meaning.

^bSitz. Ber. Akad. Wiss. Wien, Math. Naturw. Cl., XCI, 1902, p. 818.

^cProc. Zool. Soc. London, 1862, p. 265; Ann. Mag. Nat. Hist. (3), XII, 1863, p. 158.

^dCat. Shield Rept. Brit. Mus., 1855, p. 69.

^eCat. Chel. Brit. Mus., 1889, p. 247.

[*DOGANIA SUBPLANA* ^a (Geoffroy-Saint-Hilaire).]

1809. *Trionyx subplanus* GEOFFROY-SAINT-HILAIRE, NOUV. Bull. Soc. Philom. Paris, I, No. 22, July 1809, p. 366; Ann. Mus. Hist. Nat. Paris, XIV, 1809, p. 11, pl. v, fig. 2 (type-locality "unknown").—BOETGER, Offenbach. Ver. Naturk. 24-25 Ber., 1885, p. 135.—BOULENGER, Cat. Chel. Brit. Mus., 1889, p. 246 (Malay Peninsula, Sumatra, Borneo, Java).—FLOWER, Proc. Zool. Soc. London, 1899, p. 619, pl. xxxvi (Penang; Singapore).—SIEBENROCK, Sitz. Ber. Akad. Wiss. Wien, Math.-Naturw. Cl., XCI, 1902, p. 817, fig. 2 (plastron).—*Dogania subplana* GRAY, Cat. Tortois. Brit. Mus., 1844, p. 49; Cat. Shield Rept. Mus., 1855, p. 69, pl. xxxiii ("Japan"); Proc. Zool. Soc. London, 1862, p. 265 (China, Formosa); 1873, p. 57, fig. 3 (mandible); Ann. Mag. Nat. Hist. (3), XII, 1863, p. 158.

There being no specimen of this species at hand from the territory covered in this work, I append below the diagnosis given by Boulenger in his Catalogue of Chelonians in British Museum, pp. 246-247.

Diagnosis.—Carapace very flat; costal plates eight pairs, the last well developed and separated by the last neural plate; a single neural between the first pair of costals; these plates very finely granulate and vermiculate; a large fontanelle, till late in life, between the nuchal and dorsal plates; epiplastra narrowly separated from each other in front of the entoplastron, which forms an obtuse or a right angle; plastral rugosities scarcely developed, on the hyo- and hypoplastra and on the xiphiplastra; dorsal skin of young with interrupted longitudinal ridges; head very large; snout (on the skull) about as long as the diameter of the orbit, which is more than double the interorbital width; postorbital arch extremely narrow, forming a keel on its outer surface; mandibular symphysis shorter than the diameter of the orbit, without median ridge. Brown above, yellowish beneath, and on the sides of the neck; head and neck with yellowish dots; some young with an oblique dark streak behind each eye.

Length of dorsal disk 25 centimeters.

Doctor Siebenrock describes the plastral arrangement of this species as follows (op. cit., pp. 817-819): Entoplastron forms an obtuse angle; epiplastra separated, the anterior straight arm as long as the posterior oblique one; not a trace of a median process of the hypoplastron, so that an undivided membranous fontanelle is formed between the entoplastron and the hyo-, hypo- and xiphiplastra. The plastron of this species is distinguished thereby from that of all other species of *Trionyx* [*Amyda*]. The absence of the median process is a very significant character for the determination of this species. Xiphiplastra with short and narrow posterior angles which are separated by a rather broad interspace; xiphiplastral commissure short, on both sides connected by two teeth.

^a Signifying somewhat flat.

The callosities appear to develop rather late and only to a slight degree, as they are not even indicated on the hyo-hyoplastra of the young, while in equally large specimens of other species they have already reached a considerable size.

Habitat.—Known with certainty only from the Malay peninsula and archipelago. As for its alleged occurrence within our territory see remarks on p. 530.

BIBLIOGRAPHY.

General herpetological works published in book form are not included. References in parenthesis have not been verified by me.

1896. ANDRES ANGELO. La Salamandra gigantesca del Giappone. (*Megalobatrachus maximus* Boul.). Cenni descrittivi dell' esemplare esistente vivo nel Civico Museo di Milano. Atti Soc. Ital. Sci. Nat. Mus. Civ. (Milano), XXXV, fasc. 3-4, Jan. 1896, pp. 201-218, pl. 1.
1877. BASTIDE, L. Reptiles du Japon. Mém. Soc. d'Ethnogr., Sess., 1877: Revue Orientale et Américaine, Paris, 1877.
Not seen.
1904. BEDDARD, FRANK E. On some Points in the Anatomy, chiefly of the Heart and Vascular System, of the Japanese Salamander, *Megalobatrachus japonicus*. Proc. Zool. Soc. London, 1903, II, Pt. 2, (Apr., 1904), pp. 298-315.
1882. BIELETSKI, N. T. Remarques Physiologiques sur le Salamandre gigantesque, *Cryptobranchus japonicus* ["En russe; Kharkow, 1882. 8^{vo} avec planche"]
Not seen.
1871. BLANCHARD, ÉMILE. Note sur une nouvelle Salamandre gigantesque (*Sieboldia Davidiana* Blanch.) de la Chine occidentale. Comptes Rendus Acad. Sci. Paris, LXXIII, pp. 79-80.
1858. BLEEKER, P. [Reptiliën van Japan] Natuurk. Tijdschr. Nederland. Indië, XVI, pp. 204-205.
A nominal list of 12 common reptiles and batrachians collected by Mr. P. Knuttel and Dr. J. K. van den Broek in "Japan" (most likely near Nagasaki).
1876. BOETTGER, OSKAR. Ueber die aeusseren Kiemenöffnungen bei jungen Exemplaren des japanischen Riesenmolchs. Zool. Garten (Frankf. a. M.) XVII, No. 12, Dec., 1876, pp. 432-435.
1878. BOETTGER, O. Reptilien und Amphibien Japans. Offenbach. Ver. Naturk. 17-18 Ber., p. 8.
A mere nominal list of 3 marine turtles, 2 fresh water turtles, 10 snakes, 3 lizards, and 12 batrachians.
1888. BOETTGER, OSKAR. Aufzählung einiger neu erworbener Reptilien und Batrachier aus Ost-Asien. Ber. Senckenberg. Naturf. Ges., 1888 Abh., pp. 187-190.
Japalura polygonata and *Eumeces marginatus* from O-shima, Riu Kiu Archipelago; *Trimeresurus gramineus*, South Formosa; *Hyla chinensis* var. nov. *immaculata* from Shanghai, p. 189.
1895. BOETTGER, O. Neue Frösche und Schlangen von den Liukiu-Inseln. Zool. Anz., XVIII, 8 Juli, 1895, pp. 266-270.
Rana okinavana, p. 266; *R. eifingeri*, p. 267; *Coluber schmackeri*, p. 268; *Ablabes herminae*, p. 269; *Trimeresurus luteus*, p. 269.

1895. BOETTGER, O. Neue Frösche und Schlangen von den Liukiu-Inseln. Offenbach. Ver. Naturk. 33 36 Bericht, 1895, pp. 101-117.
The diagnoses of the new species described in Zool. Anz. 1895, repeated. On pp. 113-116, a list of the batrachians and reptiles of the Liukiu Islands, 24 species.
1826. BOIE, H. Merkmale einiger japanischen Lurche. Isis, 1826, pp. 203-216.
Based on the collections made by Blomhoff at Decima. *Tropidonotus tigrinus*, p. 206; *Tropidonotus vibakari*, p. 207; *Coluber quadricinctatus*, p. 209; *Coluber climacophorus*, p. 210; *Coluber conspicillatus*, p. 211; *Coluber vulneratus*, p. 212; *Trigonocephalus blomhoffii*, p. 214; *Bufo praetextatus*, p. 215; *Molge pyrrhogaster*, p. 215.
In addition to these species described as new he enumerates a few additional ones which Blomhoff evidently had mixed in from his Java collection. Boie himself suspected that the locality of these was wrong, but Blomhoff "convinced him by their manner of preparation that such a mistake had not taken place" (p. 204), a sentence which proves that the specimens were not properly labeled.
1827. BOIE, H. Kenteekenen van eenige Japansche Amphibien. Bijdr. Natuurk. Wetensch. (Amsterdam), II, Pt. I, 1827, pp. 243-272.
A translation of the paper in Isis, 1826.
1827. BOIE, H. Caractères de quelques Reptiles du Japon. Féruss. Bull. Sci. Nat., X, 1827 (pp. 160-162).
The same into French.
1883. BOULENGER, G. A. Description of a new species of *Bufo* from Japan. Proc. Zool. Soc. London, 1883, pp. 139-140, pl. xxiii.
Bufo formosus described as new; type-locality, Yokohama.
1886. BOULENGER, G. A. First Report on Additions to the Batrachian Collection in the Natural-History Museum. Proc. Zool. Soc. London, 1886, pp. 411-416, pl. xxxix.
Among others the following species are described as new: *Rana martensi*, p. 414 (Yedo); *Rana macropus*, p. 414 (Oho shima); *Geomolge fischeri*, p. 416, pl. xxxix, figs. 2-26 (Chaborowska).
1887. BOULENGER, G. A. On a Collection of Reptiles and Batrachians made by Mr. H. Pryer in the Loo Choo Islands. Proc. Zool. Soc. London, 1887, pp. 146-150, pls. xvii-xviii.
Tachydromus smaragdinus, new species, p. 147, pl. xvii, fig. 2; pl. xviii, fig. 1.
Tropidonotus pryeri, new species, p. 149, pl. xviii, fig. 3.
1887. BOULENGER, G. A. On a new species of *Hyla* from Port Hamilton, Corea, based on an example living in the Society's Gardens. Proc. Zool. Soc. London, 1887, pp. 578-579, pl. li.
Hyla stepheni, new species, p. 579, pl. li, fig. 1.
1890. BOULENGER, G. A. A List of the Reptiles and Batrachians of Amoorland. Ann. Mag. Nat. Hist. (6), V, Feb. 1890, pp. 137-144, pl. x.
20 species. *Bombinator orientalis*, new species, p. 143.
1892. BOULENGER, G. A. Descriptions of new Reptiles and Batrachians from the Loo Choo Islands. Ann. Mag. Nat. Hist. (6), X, Oct. 1892, pp. 302-304.
New species: *Trimeresurus okinawensis*, p. 302; *Rana holsti*, p. 302; *Tytotriton andersoni*, p. 304.
1894. BOULENGER, G. A. Descriptions of a new Lizard and a new Fish obtained in Formosa by Mr. Holst. Ann. Mag. Nat. Hist. (6), XIV, Dec., 1894, pp. 462-463.
Tachydromus formosanus, new species, p. 462.
1904. BROHL, J. F. Zur Fortpflanzung des japanischen Riesensalamanders (*Megalobatrachus maximus*). Natur und Haus, XII (p. 149).

1902. BROWN, ARTHUR ERWIN. A collection of Reptiles and Batrachians from Borneo and the Loo Choo Islands. *Proc. Phila. Acad.*, 1902 (June 11), pp. 175-186.
Loo-choo Islands Reptilia, pp. 183-185; Batrachia, pp. 185-186. *Gehyra intermedia*, new species, p. 183; *Microhyla undulata*, new species, p. 186.
1905. BURNE, R. H. Notes on the Muscular and Visceral Anatomy of the Leathery Turtle (*Dermochelys coriacea*). *Proc. Zool. Soc. London*, 1905, I, pp. 291-324.
1904. BUSSY, L. P. DE. Eerste Ontwikkelingsstadien van *Megalobatrachus maximus* Schlegel. Amsterdam, 1904 (8°, 112 pp., 10 pls.)
Not seen.
1905. BUSSY, L. P. DE. Die ersten Entwicklungsstadien des *Megalobatrachus maximus*. *Zool. Anz.*, XXVIII, pp. 523-536, 18 figs.
1879. CAMERANO, LORENZO. Di alcune specie di anfibi anuri esistenti nelle collezioni del R. Museo Zoologico di Torino. *Atti Accad. Sci. Torino*, XIV, Pt. 5, April, 1879, pp. 866-898.
To a great extent devoted to Japanese species in the Zoological Museum at Torino. The specimens were mostly collected by Prof. Antonio Fontanesi in 1877 and 1888 at Tokyo, and by Mr. Aimoin and the "Magenta" expedition near Yokohama.
1900. CAMERANO, L. Ricerche intorno alla variazione del 'Bufo vulgaris' Laur. *Mem. Accad. Sci. Torino* (2), L, pp. 81-153, 2 pls.
Considers the Japanese toad a distinct species under the name of *Bufo praterstatus* Bole.
1893. CHAPMAN, HENRY C. Observations on the Japanese Salamander, *Cryptobranchus maximus* (Schlegel). *Proc. Phila. Acad.*, 1893, pp. 227-233, pls. v-vii.
Anatomical.
1870. DOBROTVORSKI. Yuzhnaya tchast Ostrova Sakhalina. *Izvest. Sibirsk. Otd. Imp. Ross. Geograf. Obshchestva*, I, Nos. 2 and 3 (pp. 18-34).
'The Southern Part of the Island of Sakhalin.' *Proc. Siber. Branch Imp. Russ. Geogr. Soc.*, etc.
1880. DOEDERLEIN, L. [Pelias berus auf Saghalien]. *Mitth. Deutsch. Ges. Ost-Asiens*, III, heft 22, December, 1880, p. 89.
1881. DOEDERLEIN, L. Die Liu-Kiu-Insel Amami Oshima. *Mitth. Deutsch. Ges. Ost-Asiens*, III, heft 24, pp. 140-156.
The fauna is considered on pp. 146-150 and the reptiles in particular, pp. 147-149.
This paper was also issued separately as a pamphlet of 31 pp. bearing the following imprint: Yokohama. Buchdruckerei des "Echo du Japon," 1881.
1881. DOEDERLEIN [L.]. Japanische Seeschlangen. *Mitth. Deutsch. Ges. Ost-Asiens*, III, heft 25, pp. 209-210.
1870. DYBOWSKI, BENEDIKT. Beitrag zur Kenntniss der Wassermolche Sibiriens. *Verh. Zool. Bot. Ges. Wien*, XX, 1870, pp. 237-242, pl. vii.
Salamandrella, new genus and *S. keyserlingii*, new species p. 237.
1885. EASTLAKE, W. C. DE LANO. The Mamushi. *Trans. Asiat. Soc. Japan*, XIII, pp. 69-81, 2 pls.
Abstract in "Nature," XXXI, Apr. 23, 1885, p. 587.
1886. FISCHER, J. G. Herpetologische Notizen. *Abh. Naturw. Ver. Hamburg*, IX, 1886, pp. 51-67, pls. I-II.
Describes several species from south Formosa in the Museum in Oldenburg, collected by Ruhstrat.
Author's separates paged pp. 1-19, pls. I-II.
1888. FISCHER, J. G. Ueber zwei von der Liukiu-Insel Okinawa stammende Schlangen. *Ber. Naturh. Mus. Hamburg*, 1887, pp. 18-22.

1894. FRITZE, ADOLF. Die Fauna der Liu-Kiu-Insel Okinawa. Zool. Jahrb. Syst., VII, 1894, pp. 852-926. ~
 Reptilia, pp. 859-865. Amphibia, pp. 865-866.
 Also issued separately with separate pagination (1-77) and title page as follows: Die Fauna der Liu-Kiu-Insel Okinawa. Habilitationsschrift | zur | Erlangung der venia legendi fuer Zoologie | der | Hohen philosophischen Facultaet der Albert-Ludwigs-Universitaet zu Freiburg i. B. | vorgelegt von | Dr. phil. Adolf Fritze. | Jena, | Gustav Fischer. | 1894.
 8vo. 77 pp. Reptilia, pp. 10-16. Amphibia, pp. 16-17.
1883. GEERTS, A. J. C. Notice sur la Grande Salamandre du Japon. Cryptobranchus japonicus v. d. Hoeven. Nouv. Arch. Mus. (Paris), (2) V, pp. 273-290, pl. xvii.
1873. GRAY, J. E. On a Salamander (Sieboldia) from Shanghai. Ann. Mag. Nat. Hist. (4), XII, Aug. 1873, p. 188.
 A skin with bones of head and feet sent by Swinhoe from Shanghai.
1868. GUENTHER, ALBERT. Sixth Account of New Species of Snakes in the Collection of the British Museum. Ann. Mag. Nat. Hist. (4), I, 1868, pp. 413-429, pls. xvii-xix.
 Described as new from Nagasaki [erroneously] *Callophis japonicus*, p. 428, pl. xvii, fig. C.
1876. GUENTHER, ALBERT. Description of a new Frog from North-eastern Asia. Ann. Mag. Nat. Hist. (4), XVII, May, 1876, p. 387.
 Described as new *Rana dybowskii* from Abrek Bay.
1896. GUERNE, JULES DE. Mort de la grande Salamandre du Japon (du Muséum d'Histoire naturelle de Paris). Étangs et Rivières, X, no. 230 (p. 207).
1876. HILGENDORF, F. Die Japanischen Schlangen. Mitth. Deutsch. Ges. Ost-Asiens, [I], heft 10, pp. 29-34.
1880. HILGENDORF [F.]. Bemerkungen ueber die von ihm in Japan gesammelten Amphibien nebst Beschreibung zweier neuer Schlangenarten. Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, no. 8, pp. 111-121, pl.
Ophites orientalis, new species, p. 115.
Trimeresurus riukiuanus, new species, p. 118.
1895. HIROTA, S. Notes on a Skink with an Accessory Tail. Zool. Mag. Tokyo, VII, July, 1895, pp. 107-115, pl. xv, figs. 1-3.
 The species of skink is *Ablepharus boutonii*, from the Bonin Islands.
1838. HOEVEN, J. VAN DER. Iets over den Grooten zoogenoemden Salamander van Japan. Tijdschr. Natuurl. Geschied. Physiol., IV, pp. 375-386, pls. va-vb.
 Also as separate pamphlet of 12 pp. bearing the following imprint: Leiden, S. en J. Luchtmans, 1838. Review in Isis, 1840, pp. 384-385.
1838. HOEVEN, J. VAN DER. Sur une nouvelle espèce de Cryptobranchus du Japon. Bull. Sci. Phys. Natur. Neerlande (Leiden), 1838, pp. 90-91 (Abstract of the above).
1841. HOEVEN, J. VAN DER. Groote Bloedschijfjes bij Cryptobranchus japonicus: eene bijdrage tot de vroegere mededeelingen over dit dier. Tijdschr. Natuurl. Geschied. Physiol., VIII, pp. 270-272.
1782. HOUTTUYN, M[ARTINUS].—Het onderscheid der salamanderen van de haagdissen in 't algemeen, en van de gekkoos in 't byzonder. Verhandel. Genootsch. Wetensch. Vlissingen, IX, pp. 305-336, pl.
1871. HUMPHRY, GEORGE MURRAY. The Muscles and Nerves of the *Cryptobranchus japonicus*. Journ. Anat. Physiol., VI, (pp. 1-61, pls. i-iv).

1865. HYRTL, JOSEPHUS. *Cryptobranchus Japonicus*. |—*Schediasma anatomicum*, | quod | *almae et antiquissimae* | Universitati Vindobonensi, | ad solennia saecularia quinta, pie celebranda, | dicat, dedicat, | Josephus Hyrtl, | Rector. | Accedunt quatuordecim tabulae. |—Vindobonae, | apud Guilielmum Braumüller, | Bibliopolamaulae C.R. et Universitatis. | MDCCCLXV.
Quarto. XII+132 pp.+XIV pls.
1897. IKEDA, S. Notes on the Breeding Habit and Development of *Racophorus Schlegelii*, Günther. *Annot. Zool. Japon.*, I, Pt. 3, Aug. 10, 1897, pp. 113-122.
1902. ISHIKAWA, C.—Ueber den Riesen-Salamander Japan's. *Mitth. Deutsch. Ges. Ost-Asiens*, IX, p. 1, Oct. 1902, pp. 79-94.
1903. ISHIKAWA, C. Distribution, oviposition, habits, &c. of the Japanese giant-salamander. (Japanese) Tōkyō Teishitsu Hakubutsukwan, 1903 (II+27+3 with pl.) 26 cm.
Not seen. Title from *Int. Cat. Sci. Lit., Zool.*, 1903, Pt. 3, pp. 120, 123.
1904. ISHIKAWA, C[HIYOMATSU]. Beiträge zur Kenntniss des Riesen-Salamanders. (*Megalobatrachus maximus Schlegel*). *Proc. Nat. Hist. Tokyo Imp. Mus.*, I, no. 2, pp. 19-37, pls. VIII-XI.
The first part of the paper is a general account of the history and habits of this species essentially as in the same author's paper in the *Mitth. Deutsch. Ges. Ost-Asiens*, 1900, while the second part relates to the egg, egg deposition, and care of eggs.
1882. IWAKAWA, TOMOTARO.—The genesis of the Egg in *Triton*. *Zool. Anz.*, V, Jan. 91, 882, pp. 10-12.
From studies on the egg of *Triton pyrrhogaster* Boie.
1882. IWAKAWA, TOMOTARO. The Genesis of the Egg in *Triton*. *Quart. Journ. Micr. Sci.*, XXX (pp. 260-277, pls. XXII-XXIV).
1900. JACOBI, ARNOLD. Verarbeitung und Herkunft der höhern Thierwelt Japans. *Zool. Jahrb. Syst.*, XIII, pp. 463-478.
On p. 474 is a brief allusion to the reptiles and batrachians.
1903. KERBERT [C.]. Eieren van *Megalobatrachus maximus*. *Tijdschr. Nederland. Dierk. Vereen.* (2), VIII (pp. XXVIII-XXIX).
1904. KERBERT, C. Zur Fortpflanzung von *Megalobatrachus maximus* Schlegel. *Zool. Anz.*, XXVII, Feb. 23, 1904, pp. 305-320.
1905. KERBERT, C. Ueber die Eier und Larven von *Megalobatrachus maximus* Schl. *Compt. Rend. Sixth Congr. Intern. Zool.*, 1904, pp. 289-294.
1898. KREFFT, P. Demonstration lebender süd-und ost-asiatischer Amphibien. *Verh. Ges. Deutsch. Naturf. Aerzte*, 69 Vers. Braunschweig, 1897, II, Pt. 1, pp. 187-188.
Habits of Rana buergeri; Triton subcristatus var. *immaculiventris*, new variety from Klusiu; *Triton subcristatus* var. *typica* (Tokyo; Kyoto).
1902. LAUBER, HANS. Anatomische Untersuchungen des Auges von *Cryptobranchus japonicus*. *Anat. Hefte*, XX, (pp. 1-18+2 pls.).
1859. MAACK, R. Puteshestnie na Amur, sovershennoye po rasporyazheniyu Sibirskago I. Russk. Geograficheskago Obshchestva v 1855 g. S.-Peterburg, 1859.
Not seen, quoted from Nikolski. Mentioned by Troschel, *Arch. Naturg.*, 1863, II, p. 623, as *Reise nach dem Amur* von Richard Maack, and according to him there is a list of 14 species of "amphibians" on p. 152.
1861. MAACK, RICHARD. *Reise im Thal des Flusses Usura*. St. Petersburg, 1861.
Entirely in Russian. Not seen. Quoted from Troschel, *Arch. Naturg.*, 1863, II, p. 623, according to whom there occur 13 species of "reptiles" in the Valley of the Usura (Ussuri), mentioned on pp. 189-193. Two specific names which I have not seen

quoted elsewhere are "*Trionocephalus vittatus* n. sp. Brandt" and "*Coluber amurensis* n. sp. Brandt." They are possibly *Akistrodon intermedius* and *Elaphe schrenckii*, but I presume they are *nomina nuda*. Neither the work nor the names are quoted by Nikolski in his *Herpetologia Rossica* (1905).

1866. MARTENS, EDUARD VON. *Japanische Reptilien*. Preuss. Exped. Ost-Asien, Zool. I, pp. 109-116.

Reptiles and Batrachians: On pp. 152-153 a few "Anmerkungen" (4-6) to the above. The chapter "Die japanische Fauna im Ganzen," pp. 148-151 has a few herpetological remarks on p. 149.

In the "Verzeichniss der gesammelten oder beobachteten Wirbelthiere," pp. 32-40 (not published until 1876) there is reference to the reptiles collected in Japan on pp. 373-383 and to the batrachians on pp. 383-384.

1893. MEURON, PIERRE DE. *Sur la grande Salamandre du Japon*. Bull. Soc. Sci. Nat. Neuchâtel, XXI, pp. 186-187.

Specimen, 0.75 m. long collected by de Meuron, presented to the Natural History Museum in Neuchâtel.

1853. MIDDENDORFF, A. TH. VON. *Amphibien*. *Sibirische Reise*, II, Pt. 2, pp. 247-251, pl. xxvi.

1886. MITSUKURI, K. and C. ISHIKAWA. *On the Formation of the Germinal Layers in Chelonia*. *Quart. Journ. Micr. Sci.* (n. s.), XXVII, Aug. 1886, pp. 17-48, pls. II-V.

1891. MITSUKURI, K. *Zool. Mag.*, Tokyo, III, no. 35, Sept. 1891, pp. 362-368.

1893. MITSUKURI, K. *On the Process of Gastrulation in Chelonia*. (Contribution to the Embryology of Reptilia, IV.) *Journ. Coll. Sci. Tokyo*, VI, Pt. 3, pp. 227-277, pls. VI-VIII.

1895. MITSUKURI, K. *How Many Times does the Snapping Turtle lay Eggs in One Season?* *Zool. Mag.*, Tokyo, VII, no. 85, pp. 143-147.

1896. MITSUKURI, K. *On the Fate of the Blastophore, the Relations of the Primitive Streak, and the Formation of the Posterior End of the Embryo in Chelonia, together with Remarks on the Nature of Mesoblastic Ova in Vertebrates*. (Contributions to the Embryology of Reptilia, V.) *Journ. Coll. Sci. Tokyo*, X, Pt. 1, pp. 1-118+pls. I-XI.

1905. MITSUKURI, K. *The Cultivation of Marine and Fresh-water Animals in Japan*. *Bull. Bur. Fish. Washington*, XXIV, June 17, 1905, pp. 257-289, pls. I-XI.

"The Snapping Turtle, or Soft-shell Tortoise, 'Suppon,'" pp. 260-266, pls. I-III. Account of Hatter's "turtle farm."

1887. [MOESCH, C.] *Der japanische Riesensalamander (Cryptobranchus japonicus) und der fossile Salamander von Oeningen (Andrias scheuchzeri)*. *Neujahrsblatt Naturf. Ges. Zürich*, 1887, 4^o, 12 pp.+pl.

The third chapter, p. 9, is headed: *Der Lebende japanische Riesensalamander in der zoologischen Sammlung im eidgen. Polytechnikum*.

1881. [NAMIYE, MOTOYOSHI].— . . . *Classified Catalogue | of the | Specimens of Vertebrates | in the | collections | of | Kiyoiku Hakubutsukuan | (Educational Museum) | Tokio. . . . | Published by the Educational Museum | . . . | 1881.*

8vo, 4 + 116, pp.

1898. NAMIYE, M. *Ueber eine wenig bekannte einheimische Schlange der Gattung Achalinus (A. spinalis, Peters)*. *Annot. Zool. Japon.*, II, Pt. 1, pp. 29-31.

1903. NAMIYE, MOTOYOSHI. *Shikoku-san Hacky-urui oyobi Ryôsai-rui*. [Batrachia and Reptilia indigenous to Shikoku.] *Dôbutsugaku Zasshi*, Tokyo, XV, 1903 (pp. 284-286).

Not seen. Title from *Int. Cat. Sci. Lit. Zool.*, 1903, Pt. 1, p. 308.

1889. NIKOLSKI, A. M. Ostrof Sakhalin i ego fauna pozvonotchnikh zhivotnikh. =Zap. Imp. Akad. Nauk, S. Peterburg, LX, Prilozhenie. no. 5. xxv+334 pp.
Reptilia, pp. 290-291; Amphibia, p. 292.
1896. NIKOLSKI, A. M. Geomolge fischeri Blgr., nazemnij triton iz Ussuriiskago kraya. Annuaire Mus. Zool. St. Pétersbourg, I, 1896, pp. 77-80.
"Sur le Geomolge fischeri Blgr., triton terrestre de la région du l'Oussouri."
1905. NIKOLSKI, A. M. Presmikayustchiyasya i zemnovodniya Rossiiskoi Imperii. (Herpetologia rossica.) Zap. Imp. Akad. Nauk, S. Peterburg (8), XVII, no. 1, pp. II+518, 2 pls.
Reptilia, pp. 1-331; Amphibia, pp. 332-442; General account of the herpetological fauna and geographical distribution, pp. 442-461; Addenda, pp. 462-495; Bibliography, pp. 496-509.
1881. OKABE RYOYEI [The treatment of snake bite]. Tokei Zusshi Osaka, Sept. 5, 1881. Not seen. Title from Mitchell and Reichert's Bibliography.
1891. OKADA, S. Catalogue | of | Vertebrated Animals | of | Japan | by | S. Okada. | Tôkyô, Japan: | Kinkôdô. | 2551 (1891).
8vo, 22 + 128 pp.
Batrachia, pp. 65-67, —21 species; Reptilia, pp. 68-72, —32 species.
1902. OSAWA, G. Beiträge zur Anatomie des japanischen Riesensalamanders. Mitth. Medic. Facult. Tokyo, V, No. 4 (pp. 221-427, 44 pls.).
Also separate. Tokyo, 1902. 207 pp., quarto + 44 partly colored plates.
1881. PETERS, W. Ueber eine neue Art von Tachydromus aus dem Amurlande. Sitz. Ber. Ges. Naturf. Fr. Berlin, 1881, pp. 71-72.
Tachydromus amurensis, new species, p. 71.
1897. PHISALIX, C. Action physiologique du venin de salamandre du Japon (*Sieboldia maxima*). Atténuation par la chaleur et vaccination de la Grenouille contre ce venin. Compt. Rend. Acad. Sci. Paris, CXXV (pp. 121-123).
Also in Bull. Mus. Hist. Nat. Paris, 1897 (pp. 242-244).
1897. PHISALIX, C. Action physiologique du venin de salamandre du Japon (*Sieboldia maxima*). Atténuation par la chaleur et vaccination de la Grenouille contre ce venin. Compt. Rend. Soc. Biol. Paris (10), IV (pp. 723-725).
1897. PHISALIX, C. Propriétés immunisantes du venin de salamandre du Japon vis-à-vis du venin de vipère. Compt. Rend. Soc. Biol. Paris (10), IV (pp. 822-823).
1860. POMPE VAN MEERDERVOORT, J. L. C. [Over den grooten Japanschen Salamander.] Natuurk. Tijdschr. Nederland. Indië, XX, Pts. 4-6, p. 386.
1884. REIN, J. J. Reptiles and Batrachians [of Japan]. Japan: Travels and Researches. Translated from the German. New York, Armstrong and Son, 1884, pp. 186-189.
1876. REIN, J. J. and RORETZ, A. VON. Beitrag zur Kenntniss des Riesensalamanders (*Cryptobranchus japonicus*). Zool. Garten (Frankf. a. M.), XVII, No. 2, Feb. 1876, pp. 33-37, color. pl.
1897. REJSEK, JOS. L'histologie de l'œil de *Cryptobranchus japonicus*. Bibliogr. Anat. Nancy, V (pp. 139-146, 1 pl).
1887. SASAKI, C. Some notes on the Giant Salamander of Japan (*Cryptobranchus japonicus*, Van der Hoeven). Journ. Coll. Sci. Tokyo, I, Pt. 3, pp. 269-274.

1837-1844. SCHLEGEL, H. Abbildungen | neuer oder unvollständig bekannter | Amphibien, | nach der Natur oder dem Leben entworfen | herausgegeben | und mit erläuternden Texte begleitet | von | Dr. H. Schlegel. | Conservator des Niederländischen Reichs-Museums etc. etc. | Düsseldorf, | Verlag v. Arnz & Comp^{nie} | 1837-1844.

Fol. 50 col. plates. Text 8vo, xiv + 141 pp.

The following Japanese species are included: *Trionyx japonicus*, pl. xxxi; *Salamandra naxia*, pl. xxxix, fig. 4; *S. subcristata*, pl. xl, figs. 1-3; *S. nebulosa*, pl. xl, figs. 7-10; *Emys vulgaris japon.*, pl. xli, figs. 1-3; *Hyla buergeri*, pl. l, fig. 5. All these were probably published in 1844.

The work was issued in parts between the years 1837 and 1844.

First part consisting of 10 plates (pls. 1-X) and two signatures of text (pp. 1-32) appeared early in 1837, and was reviewed by Rudolph Wagner in *Münchener Gelehrte Anzeigen*, IV, No. 127, June 28, 1837, pp. 1043-1044.^a

The second part, containing the second decade of plates (pls. 21-40) and pp. 33-64, was published apparently in 1839. Sundevall (*Årsber. Zool. Arbet.*, 1837-1840, pp. 183-184; published 1841) speaks of the eleventh decade containing a monograph of the genus *Typhlops* having been published in 1839, but he has evidently read Decas II instead of Decas I.

The rest of this publication was probably issued together in 1844, at least I find no contemporaneous reference to matter contained in this part previous to that year, and Wiegmann, in his "Leistungen" for 1840, 1841, and 1842, does not mention it. In the "Leistungen" for 1843, however, the work is referred to as completed, and the years given as "1837-1844," the same as on the title-page. The book, therefore, may have been issued complete shortly after the beginning of 1844, as the preface is dated January, 1844. That it did not appear in 1843 seems plain, moreover, from the fact that there is no reference to Schlegel's monographic synopsis of the genus *Draco* (pp. 81-96) in Fitzinger's *Systema Reptilium* published in that year.

1899. SCHNEE [PAUL K. G.]. Einige Notizen über Weichschildkröten. II. *Trionyx sinensis*, Wiegmann. *Zeitschr. Naturwiss.*, LXXII, pp. 202-208.

Remarks on specimens seen in Canton, Shanghai, and Kobe.

1900. SCHNEE [PAUL K. G.]. *Ancistrodon blomhoffi* Boie. *Zool. Garten (Frankf. a. M.)*, XLI, Dec., 1900, pp. 395-396.

Description of color of living specimens.

1904. SCHOENICHEN, WALTHER. Die Brutpflege bei den Amphibien und besonders bei dem japanischen Riesensalamander (*Megalobatrachus maximus*). *Prometheus*, XVI, 1904 (pp. 37-40; pp. 52-54. 15 figs.).

1895. SHERBORN C. DAVIES, and F. A. JENTINK. On the dates of the parts of Siebold's "Fauna Japonica" and Giebel's "Allgemeine Zoologie" (first edition). *Proc. Zool. Soc. London*, 1895, pp. 149-150.

1895. SHITKOV, B. [M.]. Ueber die Fortpflanzung des *Isodactylum Schrenki* Strauch. *Zool. Anz.*, XVIII, May 6, 1895, pp. 165-168.

1900. SHITKOV, [B. M.]. Sibirskiya salamandri (*Isodactylum Schrenkii* Str.) i ikh zhizn na vodie i v akvarii. *Dnevnik Otdiela Ikhtiologii Imp. R. Obshch. Akklimat. Zhivotn. i Rast.*, II, 1900, pp. 41-42.

Not seen.

1902. SHITKOV, B. M. Kakogo paltsa nedostaet na konetsnostyakh *Isodactylum* i na perednikh konetsnostyakh Urodela. *Izv. M. Obshch. Lyub. Est. Antr. i T. D.*, XCVIII. *Tr. Zool. Otd.*, XIII, *Dnevn. Zool. Otd.*, III, No. 4, p. 45.

Not seen.

1906. SIEBENROCK, F. Zur Kenntnis der Schildkrötenfauna der Insel Hainan. *Zool. Anz.*, XXX, Aug. 28, 1906, pp. 578-586.

On p. 584 there are three lists of the turtles of Japan, of the Riu Kiu Islands, and of Formosa.

^aI am indebted to Dr. Charles W. Richmond for calling my attention to this review.

1824. SIEBOLD, P. F. DE.^a De Historiae naturalis in Japonia statu, nec non de augmento emolumentisque in decursu perscrutationum exspectandis. Dissertatio, cui accedunt spicilegia faunae Japonicae, auctore P. F. de Siebold, Med. Doct. complurium societatum membro. Bataviae, 1824.

Original not seen. Title from the reprint in Oken's *Isis*, 1827, pp. 135-143. It is stated to be in 4° and to contain 16 pp.

1826. SIEBOLD, PHILIPPUS FRANC. DE. De | Historiae Naturalis | in | Japonia | Statu | nec non | de augmento emolumentisque in | decursu perscrutationum ex- | spectandis | Dissertatio, | cui accedunt | Spicilegia Faunae Japonicae | auctore | Philippo Franc. de Siebold, | Medicinae Doctore, | Complurium Societatum Membro. | — Ex Officina Literaria | Car. Phil. Bonitas, Commerc. Assessoris | Wirceburgi MDCCCXXVI.

8vo, 20 pp.

Among the Amphibia, p. 18, reference is only made to *Hydrophis flaviventris* "mblt."

1898. STEJNEGER, LEONHARD. On a collection of batrachians and reptiles from Formosa and adjacent islands. Journ. Sci. Coll. Tokyo, XII, Pt. 3, pp. 215-225.

Rana longicrus, new species, p. 216; *Japalura mitsukurii*, new species, p. 218; *Emydocephalus ijimæ*, new species, p. 223.

1901. STEJNEGER, LEONHARD. Diagnoses of eight new batrachians and reptiles from the Riukiu Archipelago, Japan. Proc. Biol. Soc. Washington, XIV, Dec. 12, 1901, pp. 189-191.

Microhyla okinawensis, new species, p. 189; *Rana narina*, new species, p. 189; *Rana namiyai*, new species, p. 190; *Buergeria ijimæ*, new species, p. 190; *Buergeria ishikawæ*, new species, p. 190; *Eumeces kishinouyei*, new species, p. 190; *Calamaria pfefferi*, new species, p. 191; *Disteria orientalis*, new species, 191.

1902. STEJNEGER, LEONHARD. A new opisthoglyph snake from Formosa. Proc. Biol. Soc. Washington, XV, Feb. 18, 1902, pp. 15-17.

Boiga kræpelini, new species, p. 16.

1904. STEJNEGER, LEONHARD. A new species of lizard from the Riukiu Archipelago, Japan. Smithson. Misc. Coll. (Quart. Iss.) XLVII, Nov. 9, 1904, pp. 294-295.

Takydromus dorsalis, new species, p. 294; type, U.S.N.M. No. 34162.

1899. STONE, WITMER. A small collection of reptiles and batrachians from eastern Mongolia. Proc. Phila. Acad. Sci., 1899, pp. 183-184.

Nine species.

1870. STRAUCH, ALEXANDER. Revision der Salamandriden-Gattungen nebst Beschreibung einiger neuen und weinger bekannten Arten dieser Familie. Mém. Acad. Sci., St. Pétersbourg (7), XVI, No. 4, 110 pp., 2 pls.

1873. STRAUCH, ALEXANDER. Die Schlangen des Russischen Reichs, in systematischer und zoogeographischer Beziehung. Mém. Acad. Sci. St. Pétersbourg (7), XXI, No. 4, 288 pp., 6 pls.

Particularly important in this connection for its exhaustive treatment of the specimens from Amurland and the Coast Province. As for the specimens alleged to have been collected in Japan by Consul Goshkewitsch see pp. 315, 318, and 452 of this work.

^a Dr. C. L. Reuvens, in his doctor dissertation "Die Myoxidae or Schlaefcr" (Leiden, 1890), p. 66, footnote (compare p. 7), makes a curious mistake in ascribing the "De Historiae Naturalis in Japonia Statu" to a G. T. de Siebold as distinct from P. F. de Siebold, the author of *Fauna Japonica* ("G. T. de Siebold ist nicht zu verwechseln mit P. F. de Siebold, dem Autor der *Fauna Japonica*"). In the Würzburg reprint of the pamphlet in question now before me it is plainly attributed to the latter: "Auctore Philippo Franc. de Siebold, Medicinae Doctore."

1876. STRAUCH, ALEXANDER. Opisanie presmikayustchikhsya i zemnovodnikh, sobrannikh ekspeditsiei podpolkovnika Przewalskago. In Przewalski's Mongoliya i Strana Tangutov. S.-Peterburg, 1876.
1890. STRAUCH, ALEXANDER. Bemerkungen über die Schildkröten-sammlung im Zoologischen Museum der kaiserlichen Akademie der Wissenschaften zu St. Petersburg. Mém. Acad. Sci. St. Pétersbourg (7), XXXVIII, No. 2, 127 pp., 4 pls.
1863. SWINHOE, R. A list of the Formosan reptiles; with notes on a few of the species, and some remarks on a fish (*Orthogoriscus*, sp.). Ann. Mag. Nat. Hist. (3), XII, Sept. 1863, pp. 219-226.
- 1835-1838. TEMMINCK, C. J., and SCHLEGEL, H. Fauna Japonica | auctore | Ph. Fr. de Siebold. | Reptilia | elaborantibus | C. J. Temminck et H. Schlegel. Cum mappa geographico-zoologica et tabulis lithogr. XXVIII. | Lugduni Batavorum. | Ex officin. lithogr. auctoris et typis J. G. Lalau. | 1838.

Fol. xxii + 144 pp. + double page map + ix + 10 + viii plates.

The "Reptilia" of the Fauna Japonica was apparently issued in three separate parts, namely, I *Chelonii*, II *Ophidii*, and III *Saurii et Batrachii*, the plates being numbered separately in each part, but the pagination of the text running on continuously. The publication appears to have been as follows:

1835. Fauna Japonica auctore Ph. Fr. de Siebold. Chelonii. Elaborantibus C. J. Temminck et H. Schlegel. Lugd. Batav. (1835). 80 S. mit 9 unkolorirten Steindrucktafeln in Fol.

The above is the title of the first part, as given in München Gelehrte Anzeiger, II, No. 90, 5 May, 1836, p. 743. It is there stated that Fitzinger's paper on the Chelonians (Ann. Wien Mus., I, 1835) "gleichzeitig mit diesem Theil der Fauna Japonica erschien." On p. 751 a list of the plates 1-9 is given, but no reference is made to the map representing the "Distribution géographique des Chéloniens" which accompanies the bound volume of the Reptilia.

Also reviewed in Wiegmann's Archiv., 1836, II, p. 259, in the "Leistungen" for 1835.

1837. Fauna Japonica auctore Ph. Fr. de Siebold. Ophidii. Elaborantibus C. J. Temminck et H. Schlegel. Lugd. Batav. (1837) S. 81-93 und 1-xxx mit 10 unkolorirten Steindrucktafeln in Fol.

The second part of the Reptilia (third of the Fauna Japonica) is thus quoted by A. Wagner, in the München Gelehrte Anzeiger, V, No. 134, July 7, 1837, p. 41. This part, according to him, contained two different memoirs, the first one, by Schlegel, treating of the Japanese snakes (pp. 81-93 + pls. 1-x); the other, by Temminck, giving a review of the faunas of the Sunda Islands and of Japan (pp. 1-xxx).

The latter, which is dated November, 1835, but first distributed through the book trade during the Easter "Messe" at Leipzig in 1837,^a is evidently Temminck's "Coup d'œil."^b

Wagner (on p. 55) calls special attention to the *Triton japonicus* mentioned by Temminck in this work.

This part is also reviewed in Magazine of Zoology and Botany (ed. by Jardine, Selby and John ton), II, 1837, No. IX, p. 296 (published August 1, 1837).

1838. The title page of the entire reptile volume bears this date and may be taken as correct for the publication of the third part, embracing the Saurii and Batrachii (pp. 85-144, pls. 1-viii). On p. 140, the last page of the text (exclusive of index), is printed

^a "Diese Abhandlung ist vom November 1835 datirt; durch den Buchhandel ist sie indes erst seit des letzten Ostermesse angezeigt und verbreitet." Wagner. München Gelehrte Anzeiger, V, p. 43.

^b I have seen only two copies of this memoir, the full title of which is as follows:

Coup d'œil sur la Faune des îles de la Sonde | et de | l'empire du Japon. | Discours préliminaire | destiné à servir d'introduction à la Faune du Japon.

It is a folio of xxx pp., with Temminck's name at the end only. The copy which I examined in the library of the Jardin des Plantes, in Paris, was bound with the mammals of the Fauna Japonica.

the date "Janv. 1838." In this part was also included Siebold's "Préface suivie d'un aperçu historique et physique sur les reptiles du Japon" (pp. 1-xxi), which is signed "Mai, 1838." As Wiegmann, in his "Leistungen" for 1838 (Arch. Naturg., 1839, II, p. 386), complains that the Royal Library in Berlin had not as yet received the part in question, it is plain that he knew of its publication in 1838. It was reviewed in the "Isis" for October, 1838 (pp. 852-854), hence its appearance must have been between May and October; probably at the "Michaelismesse" (September).

It should be added, for the sake of completeness, that Tschudi, in a letter written not later than August, 1837, from Leiden (printed in Leonhard and Bronn's Neues Jahrbuch für Mineralogie, etc., 1837, Pt. 5, Sept., pp. 545-547) says that "Im 7 Hefte der Fauna japonica ist er schon abgebildet," meaning Schlegel's *Salamandra maxima*, which is contained in the third part of the reptile volume. There is no doubt that the plates and possibly also the text of this part were finished in 1837, and we know that Schlegel distributed a few copies of the plate in question during that year, but Tschudi's statement does not invalidate the contention that the part as such was not published until the Michaelismesse of 1838.

It will be seen that the dates of publication of the herpetological parts of Fauna Japonica differ somewhat from those given by Sherborn and Jentink.^a

They may be tabulated as follows:

<i>Chelonii</i>	pp. 1-80, pls. 1-IX.....	1835 (Ostermesse?)
<i>Ophidii</i> (+ Coup d'œil)	pp. 81-93, pls. 1-X..}	1837 (Ostermesse)
	pp. 1-XXX.....}	
<i>Saurii et Batrachii</i>	pp. 85-144, pls. 1-VIII.....	1838 (Michaelismesse?)

1877. TROSCHEL [F. H.]. Ueber *Onychodactylus japonicus* Bonap. Arch. Naturg. XLIII, Pt. 1, pp. 199-215, pl. xv.

Elaborate history, synonymy, and descriptions of male and female adults, male and female larvæ.

1837. TSCHUDI, J. J. Ueberden Homo diluvii testis, Andrias Scheuchzeri. Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefactenkunde (Leonhard and Bronn, editors, Stuttgart), 1837, Pt. 5, Sept., pp. 545-547.

The generic names *Megalobatrachus* and *Andrias* are first proposed in this paper.

1902. WALL, F. Extraordinary magnitude of a snake's meal. Journ. Bombay Nat. Hist. Soc., XIV [?], p. 375, pl.

Elaphis virgatus swallowing a *Lepus brachiuurus* near Arima, Japan.

1903. WALL, F. A Prodrômus of the snakes hitherto recorded from China, Japan, and the Loo Choo Islands; with some notes. Proc. Zool. Soc. London, 1903, pp. 84-102.

Part 1. List of Chinese Ophidia [including Formosan], pp. 85-99.

Part 2. List of Japanese and Loo Choo Islands Ophidia, pp. 99-102.

1906. WALL, F. Notes on a Collection of Snakes from Japan and the Loo Choo Islands. Proc. Zool. Soc. London, 1905, II, pp. 511-517.

1864. WALLACE, ALFRED R. On some Anomalies in Zoological and Botanical Geography. Natural History Review (London and Edinburgh), IV, No. xiii, Jan., 1864, pp. 111-123.

On pp. 114-115 Mr. Wallace endeavors to explain how it is "that the snakes of Japan are Indian and the batrachians Palearctic."

1904. WERNER, FRANZ. Ueber Reptilien und Batrachier aus Guatemala und China in der Zoologischen Staats-Sammlung in München. II. China. Abh. Bayer. Akad. Wiss. (München), II Klasse, XXII, Pt. 2, pp. 353-380.

1903. YAMADA, YOSHISABURŌ. Hatakedojo ni tsukite. [Ethology and description of *Hynobius nebulosus*.] Dōbutsugaku Zasshi, Tokyo, XV, 1903 (pp. 73-75).

Not seen. Title from Int. Cat. Sci. Lit., Zool., 1903, Pt. 1, p. 429, and Pt. 3, p. 119.

^a Proc. Zool. Soc. London, 1895, p. 149.

LIST OF LOCALITIES IN JAPAN PROPER, RIU KIU ARCHIPELAGO, AND FORMOSA, MENTIONED IN THIS WORK, WITH THEIR PRINCIPAL SYNONYMS.

The spelling of many localities is often so different in the various zoological and geographical writings quoted in this work that it seems important to give a list of the various names and their different synonyms, the more so since there is as yet no generally adopted standard spelling, Tokyo and Tokio, Kiusiu and Kyushu, etc., occurring indiscriminately often in the same publication. For this reason no attempt has been made at uniformity in the text of this work.

AGINCOURT; small island of the northern extremity of Formosa.

AIZU; district in the province of Iwashiro, around the town of Wakamatsu.

AKI; province, near western extremity of Hondo, facing the Inland Sea; capital Hiroshima.

AMAKIRIMA, AMAKERRIMA, or AMAKIRIMA ISLAND; same as KERAMASHIMA.

AMAMI-Ō-SHIMA; large northern island of the middle group of the Riu Kiu Archipelago, also known as Ōshima (Satsuma); Oho-shima; Oushima, etc.

AMAMI-Ō-SHIMA; subgroup of the Riu Kiu Archipelago consists of the large Amami-ō-shima, from which Kakeroma shima is separated by a narrow strait, Tokuno shima, Okinoerabu shima, Yoron shima, and a few smaller islets.

AOMORI; town at the head of Aomori Bay, near Tsugaru Strait, the terminus of the Northern Railway.

ASAHIGAWA; river in the San-yōdō.

AWA; there are two provinces of this name, one, also called Boshū, on Hondo, occupying the end of the peninsula east of the Bay of Tokyo, the other on the east side of Shikoku.

AWAJI; province embracing the large island at the eastern entrance to the Inland Sea, between Shikoku and Hondo.

AWOMORI; same as AOMORI.

BANGKIMTSING; locality in southern Formosa.

BINGO; province near the western end of Hondo, facing the Inland Sea.

BITCHŪ; province near the western end of Hondo, east of Bingo, facing the Inland Sea.

BIWA, LAKE; in the province of Omi, Hondo, not far from Kyōto. It is 36 miles long by 12 miles wide. The altitude above the sea is 328 feet.

BIZEN; province near the western end of Hondo, facing the Inland Sea, east of Bitchū.

BONIN ISLANDS; Japanese Ogasawara shima, a small archipelago of volcanic origin, situated on the 142d meridian east of Greenwich and between 26° 30' and 27° 45' north lat. It consists of three groups, Mukoshima, Chichi shima, and Haha shima.

BORODINO ISLANDS; small group in the Northern Pacific, southeast of the Okinawa group, Riu Kiu.

BŌSHŪ; alternate name for Awa, Hondo.

BOTEL TOBAGO; high island situated due east from the southern end of Formosa.

Type-locality of *Japalura mitsukurii* Stejneger.

- BUZEN; province in northeastern part of Kiūsū.
- BYŌRITSU; town in northern Formosa near the west coast, 68 miles from Taihoku.
- CÉCILIA ARCHIPELAGO; same as TOKARA GROUP, Riu Kiu.
- CHICHIBU; district in the western corner of the province of Musashi.
- CHICHI SHIMA; middle group of the Bonin Islands.
- CHIKUGOGAWA; river in northern Kiūsū forming the boundary between the provinces of Chikugo and Hiizen.
- CHIKUZEN; province in northern Kiūsū facing the strait of Tsushima.
- CHISHIMA, the thousand islands; Japanese name for the Kuril Islands.
- CHITOSE FOREST; on the road between Sapporo and Shikotsu Lake, Yezo.
- CHUSENJI, or CHŌZENJI LAKE; near Nikko; it is about 7 miles long by $2\frac{1}{2}$ miles wide, its level being 4,375 feet above the sea.
- CLEOPATRA ISLAND; same as YOKO SHIMA.
- COFFIN ISLANDS; same as HABA SHIMA.
- DAGOSHIMA; probably error for DŌGASHIMA.
- DAIBUTSU; colossal bronze statue of Buddha at Kamakura, province of Sagami.
- DAIHOKU; same as TAIHOKU.
- DAIPE; same as TAIPA.
- DAISEN; volcano in the province of Ioki, 6,050 feet high, the loftiest mountain on the southwest coast.
- DECIMA; same as DESHIMA.
- DESHIMA; small island in the harbor of Nagasaki, the seat of the Dutch factory.
- DEZIMA; same as DESHIMA.
- DŌGASHIMA; village near Miyanoshta, Hakone, about 1,300 feet above the sea.
- ENOSHIMA; small island in Sagami Bay, connected at low tide by a sand spit with the main island, Hondo.
- ERABU SHIMA; one of the islands of the Miyako subgroup, Riu Kiu.
- ETCHŪ; province in the island of Hondo, bordering the sea of Japan just east of the peninsula of Noto.
- FAKONE; same as HAKONE.
- FATSIZIO; same as HACHJŌ.
- FUJI-SAN, also called FUJI-YAMA, or Mount Fuji; in the province of Suruga, 12,390 feet high.
- FUSI-YAMA; same as FUJI.
- GILAN, or GIRAN; town on the east coast of Formosa, near the northern end.
- GOTŌ ISLANDS; a group of islands directly west from Nagasaki, the largest of which are known as Fukue and Nakadori shima.
- GREAT LIUKIU, or LOO CHOO; same as OKINAWA SHIMA.
- HACHJŌ SHIMA; southernmost of the Seven Islands of Idzu. It is located in the Kuroshiwō, is about 12 miles long, and the highest point 2,840 feet above sea level.
- HABA SHIMA; southern group of the Bonin Islands.
- HAKONE; mountainous district at the base of the peninsula of Idzu, southeast of Mount Fuji. Lake Hakone, with the village at its south end, is 2,378 feet above the sea and about $4\frac{1}{2}$ miles long. Type-locality of *Onychodactylus japonicus* (Thunberg).
- HARBOR ISLAND; same as KIKAIKA SHIMA.
- HARIMA; province in the southwestern part of Hondo, at the eastern end of the Inland Sea.
- HATA; village in the Hakone Mountains.
- HATERUMA SHIMA; southernmost island of the Yaeyama subgroup, Riu Kiu.
- HAYAKAWA; river originating in Lake Hakone.
- HIDA; province in central Hondo, south of Etchū.
- HIGO; province in western Kiūsū facing Shimabara Bay, north of Satsuma.

- HIKOSAN**; village and mountain in the province of Buzen, Kiusiu. The altitude of the village above the sea is 1,850 feet, that of the mountain 3,850 feet.
- HIRUZEN**; mountains in the San-yōdō.
- HITACHI**; province on the east coast of Hondo, northeast of Tokyo.
- HIZEN**; province in northwestern Kiūsū, in which Nagasaki is situated.
- HIŪGA**; province on the southeast coast of Kiūsū.
- HOA-PIN-SU**; one of the islets forming the Pinnacle group, northeast of Formosa.
- HOKI**; province in the western end of Hondo, facing the Sea of Japan, opposite the Oki Islands.
- HOKKAIDŌ**; Japanese name for the island of Yezo and its adjoining administrative dependencies, such as the Kuril Islands.
- HŌKO-TŌ**; Japanese name for the Pescadores Archipelago.
- HŌKWATO**; Japanese name of Agincourt Island.
- HONDO**, or **HONSHŪ**; main island of Japan, often called Nippon, or Nippon, by earlier writers.
- IDZU**, or **IZU**; province on the island of Hondo, forming the peninsula east of Tokyo between the bays of Sagami and Suruga.
- IDZUMO**, or **IZUMO**; province in the western extremity of Hondo, facing the Sea of Japan, west of Hoki.
- IGA**; province situated inland in southern Hondo at the base of the peninsula between the bays of Owari and Ōsaka.
- IHEYA ISLANDS**; small cluster of islands northwest of the north end of Okinawa shima, Riu Kiu.
- IKI SHIMA**; large island off the northwest coast of Kiūsū, between the latter and Tsushima.
- INABA**; province in southwestern Hondo, facing the Sea of Japan, east of Hoki.
- INATORI**; town in the province of Idzu, opposite Ōshima, less than 10 miles northeast of Shimoda.
- IRIOMOTE SHIMA**; largest island of the Yaeyama subgroup, Riu Kiu.
- ISE**; province in southern Hondo, on the east side of Owari Bay.
- ISHIGAKI SHIMA**; easternmost island of the Yaeyama subgroup, Riu Kiu.
- ITIGAWA**; river in the province of Tamba.
- IWAMI**; province in the extreme western portion of Hondo, facing the Sea of Japan.
- IWASHIRO**; interior province of northern Hondo, between 37° and 38° north latitude.
- IZU**; same as IDZU.
- IZUMO**; same as IDZUMO.
- JESO**, **JESSO**, or **JEZO**; same as YEZO.
- JOKOHAMA**; same as YOKOHAMA.
- KAGA**; province on the west coast of Hondo, facing the Sea of Japan south of Noto.
- KAKEROMA SHIMA**, or **KAGERUMA SHIMA**; island south of Amami-ōshima, Riu Kiu, separated from it by a narrow strait.
- KARAFUTO**; Japanese name for Sakhalin.
- KATONA SHIMA**; same as KAKEROMA SHIMA.
- KAWACHI**; province of southern Hondo, east of Ōsaka.
- KAZUSA**; province in Hondo, occupying the basal portion of the peninsula east of the Bay of Tokyo.
- KELUNG**; port situated near the northern extremity of Formosa.
- KERAMA SHIMA**; small cluster of islands due west of the south end of Okinawa shima, Riu Kiu, the largest of which is Tokashiki shima.
- KIDZUGAWA**, or **KIZUGAWA**; river in the province of Yamashiro, joining the river Yodogawa a short distance south of Kyōto.
- KII**; province in southern Hondo, occupying the southern extremity of the peninsula between the bays of Owari and Ōsaka.
- KIIRUN**; Japanese name for Kelung.

- KIKAIGA SHIMA; one of the islands of the Amami-ō-shima subgroup, Riu Kiu.
- KILUNG; same as KELUNG.
- KIŌTO; same as KYŌTO.
- KIŪSHŪ, or KYŪSHŪ; extreme southwestern island of the four large ones constituting Japan proper.
- KIYOTAKI TOGE; mountain in the province of Kawachi.
- KŌBE; large city in province of Setsu, on Ōsaka Bay.
- KŌCHI; on the southern coast of Shikoku, the capital of the province of Tosa.
- KŌRIYAMA; town in the province of Yamato, 3 miles from Nara.
- KŌTŌ-SHO; Japanese name for Botel Tobago.
- KRAFŌ; same as KARAFUTO.
- KUCHINOERABU SHIMA; one of the islands of the Ōsumi group, Riu Kiu.
- KUKIEN SAN; Chinese name for Iromote shima.
- KUMAMOTO; capital of the province of Hiigo, Kiūsiū.
- KUNASHIRI; island off the northeast extremity of Yezo.
- KURO SHIMA; (1) one of the islands of the Ōsumi group, Riu Kiu; (2) one of the islands of the Yaeyama subgroup, Riu Kiu.
- KYŌTO; old capital of the Mikado, province of Yamashiro.
- KYŪSHŪ; same as KIŪSHŪ.
- LEW-CHEW ISLANDS; same as RIU KIU.
- LINSCHŌTEN ARCHIPELAGO or GROUP; same as TOKARA GROUP, RIU KIU.
- LIŌUKIŌU; French spelling for Linkiu; same as RIU KIU.
- LIUKIU; Chinese pronunciation of Riu Kiu.
- LOO CHOO; same as RIU KIU.
- LOO CHOO ISLAND, or GREAT LOO CHOO; same as OKINAWA SHIMA.
- LUCHŪ; same as RIU KIU.
- MEIACOSHIMA; same as MIYAKOSHIMA.
- MIACOSHIMA ISLANDS; same as MIYAKOSHIMA ISLANDS as a collective name for the southern Riu Kiu. Same as SAKI SHIMA.
- MIMAZAKA; a province in the central portion of the western extension of Hondo.
- MINO; province in central Hondo between Owari and Echizen.
- MIYADUGAWA; river in province of Tamba.
- MIYAKESHIMA; middle island of the Seven Islands of Idzu. Not to be confounded with Miyakoshima.
- MIYAKO ISLANDS, as a group name often applied to the whole southern Riu Kiu group, or synonymously with Sakishima.
- MIYAKO SHIMA; the eastern large island of the Saki shima group, Riu Kiu, with Erabu shima, Tarama shima, and Mīnua shima, forming the Miyako subgroup.
- MIYAZAKI; town on the east coast of Kiūsiū in the province of Hiūga.
- MUKOSHIMA; northern group of Bonin Islands.
- MURAYAMA; village at the foot of Mount Fuji, on the south side of the volcano.
- MUSASHI; province in Hondo, in which Tōkyō is situated.
- NAFA; same as NAHA.
- NAGASAKI; principal city of Kiūsiū, in the province of Hiizen.
- NAGATO; province in Hondo, forming the western extremity of this island.
- NAHA; capital city of Okinawa shima.
- NANGASACKI, and NANGASAKI; same as NAGASAKI.
- NAPA; same as NAHA.
- NARA; city in the province of Yamato, not far from Kyōto and Ōsaka.
- NASE, or NAZE; a town and district on Amami-ō-shima, the type locality of *Trimerisurus riukianus* Hilgendorf.
- NEMURO; town at the northeastern extremity of Yezo.
- NII SHIMA; one of the northern islands of the Seven Islands of Idzu.

- NIKKO**; mountain district about 100 miles north of Tōkyō, in the province of Shimo-zuke. In a more restricted sense the name applies to the locality of the mausoleums of the Tokugawa shoguns. The altitude of the latter is about 2,000 feet above the sea.
- NIPON**, or **NIPPON**; often used by older writers to designate the main island, now known as Hondo.
- NIHIOMOTE SHIMA**; same as **IRIOMOTE SHIMA**.
- NOTO**; province on the west coast of Hondo, forming a curved peninsula in the Sea of Japan, about the middle of the island.
- OGASAWARA**; Japanese name for the Bonin Archipelago.
- OHOSHIMA**; same as **AMAMI-Ō-SHIMA**.
- OKINAWA SHIMA**; by earlier writers called **GREAT LOOCHOO**, is the largest island of the Riu Kiu Archipelago, situated about halfway in the chain of islands between Kiūsū and Formosa.
- OKINAWA SUBGROUP**, Riu Kiu, comprises the southern islands of the middle group, of which Okinawa shima is the largest. Many other islands belong to this subgroup, some of them forming minor groups such as the Kerama cluster and the Iheya cluster.
- OKINOERABU SHIMA**; one of the islands of the Amami-ō-shima subgroup, Riu Kiu.
- ŌMACHI**; village in the northern part of the province of Shinshū, not far from the boundary of the province of Etchū.
- OMAYA SAKI**, or **OMAEZAKI**; province of Totomi, the promontory marking the western end of Suruga Bay.
- ŌMI**; province in southern Hondo, surrounding Lake Biwa.
- ONOMICHI**; city on the inland sea in the province of Bingo.
- ONSEN**; mountain near Shimabara, province of Hizen, Kiusiu. Type-locality of *Tachydromus holsti*.
- OOSHIMA**; same as **AMAMI-Ō-SHIMA**.
- ŌSAKA**; large city in southern Hondo at the mouth of the Yodogawa.
- ŌSAKIGAWA**; river in southern Hondo, in the provinces of Mimazaka and Harima.
- ŌSEN**, **MOUNT**; erroneous for **MOUNT ONSSEN**.
- Ō-SHIMA**; common name for various islands, the principal ones being Vries Island or Idzu-no-Ōshima; and Amami-ō-shima, also known as Ō-shima (Satsuma) and by older authors as Oushima, Ohoshima, etc.
- ŌSHIMA-OKINAWA GROUP**; large middle group of the Riu Kiu Archipelago, subdivided into the northern Amami-ō-shima subgroup and the southern Okinawa subgroup.
- ŌSUMI**, a province in Kiūsū at the southeastern extremity of the island.
- ŌSUMI GROUP** of the Riu Kiu curve comprises the larger islands south of and nearest to Kiūsū, to which they seem to belong biotically and geologically, the principal ones being Tanega shima, Yaku shima, Kuchinoerabu shima, and Kuro shima.
- OUSIMA**; same as **AMAMI-Ō-SHIMA**.
- OWARI**; province on the south coast of Hondo, at the head of the deep Owari Bay.
- ŌYAMA**; province of Sagami, a celebrated mountain, 4,150 feet high, about 22 miles due west of Yokohama. Also the village at the foot of the mountain.
- PARRY GROUP**; same as **MUKOSHIMA**.
- PA-TCHUNG-SAN**; Chinese name for Ishigaki shima.
- PEEL ISLAND**; same as **CHICHI SHIMA**.
- PESCADORES**; small group of islands on the west coast of Formosa, on the Tropic of Cancer. Japanese name Hōko-tō.
- PINNACLE GROUP**; number of small islands northeast of Formosa and north of the Saki shima group, Riu Kiu, the principal island being Hoa-pin-su.
- PORT LLOYD**; same as **CHICHI SHIMA**.
- RALOUSIMA**; see p. 23 of this work.

RIU KIU; Japanese pronunciation of the name of the chain of islands between Kiūsū and Formosa, known otherwise as Liukiu, Loo Choo, Luchu, Lioukiou, etc. The Riu Kiu curve naturally falls into four main groups from north to south as follows: Ōsumi group, nearest Kiūsū; Tokara group, or Linschoten Archipelago; Ōshima-Okinawa, or middle, group; and Saki shima, or southern group. The Ōshima-Okinawa group falls again into two subgroups, the Amami-ōshima subgroup and the Okinawa subgroup. The Saki shima group also subdivides into the Miyako subgroup and the Yaeyama subgroup. In addition some outlying groups and islets are referred to the Riu Kiu Archipelago, such as the Pinnacle group and the Borodino group.

For the islands composing these groups see under name of each of the latter.

ROKAGAWA; river in western Mino.

SAGALIEN; same as SAKHALIN.

SAGAMI; province in central Hondo, at the western entrance to Tōkyō Bay.

SAGARA; Totomi province.

SAGHALIEN; same as SAKHALIN.

SAKHALIN; Russian name for the large island to the north of **Yezo**, the southern half of which again belongs to Japan since 1905.

SAKI SHIMA GROUP; southern group of islands of the Riu Kiu curve, composed of the Miyako and Yaeyama subgroups.

SAN-INDŌ; name of the district occupying the Sea of Japan slope of the western extension of Hondo, in contradistinction to San-yōdō, embracing the Inland Sea provinces.

SAN-YŌDŌ; district occupying the Inland Sea slope of the western extension of Hondo.

SAPPORO; capital of Hokkaidō, in the interior of **Yezo**.

SASAYAMA; village in the province of Tamba, 60 kilometers west-northwest of **Kyōto**; altitude 250 meters above sea.

SATSUMA; province occupying the southwestern extremity of **Kiūsū**.

SENDAI; large city in the province of Rikuzen, northeastern Hondo.

SENDAI; small town in the province of Satsuma, **Kiūsū**.

SETA; village near south end of Lake Biwa.

SETANAI; village in the province of Shiribeshi, west coast of **Yezo**.

SETSU, or SETTSU; province in Hondo, at the head of Ōsaka Bay. The cities of **Kōbe** and **Ōsaka** are located in this province.

SEVEN ISLANDS OF IDZU; series of volcanic islands stretching north-south from the province of Idzu. The principal islands comprising this archipelago are: Ōshima or Vries Island; Nii shima; Miyakeshima; and Hachijō shima.

SHIKOKU; fourth of the principal islands of Japan, bounding the Inland Sea on the south.

SHIKOTSU LAKE; crater lake about 20 to 30 miles in circumference, about 39 miles from Sapporo, **Yezo**.

SHIMA; province in Hondo at the western entrance to Owari Bay.

SHIMABARA; town in the province of Hizen, **Kiūsū**, at the entrance of Shimabara Bay, not far from **Nagasaki**.

SHINSHŪ, or SHINSHŪ, also called **SHINANO**; inland province in central Hondo, east of Hida and Mino.

SHIRIBESHI; province on the southwest coast of **Yezo**.

SHURI; town on Okinawa shima.

SIMODA, or SHIMODA; town in the province of Idzu.

SUDZAKA YAMA; mountain in southern Hondo, where the boundaries of the three provinces of Ise, Iga, and Omi meet. Type-locality of *Megalobatrachus japonicus* (Temminck).)

SULPHUR ISLAND; same as **TORI SHIMA**.

SUMIYOSHI; famous temple $3\frac{1}{2}$ miles from Ōsaka.

- SURUGA BAY**; deep indentation on the south coast of Hondo, west of the peninsula of Idzu.
- SUWŌ**; province near the extreme western point of Hondo, facing the Inland Sea.
- SUZUGA YAMA**; same as **SUDZAKA YAMA**.
- TAIHOKU**; Japanese name for Taipa or Taibe, especially the inner walled portion of the town.
- TAINAN**; same as **TAIWAN-FU**.
- TAIPA, or TAIBE**; Japanese **TAIHOKU**, the capital of Formosa, near the northern extremity of that island.
- TAIWAN**; Chinese and Japanese name for Formosa.
- TAIWANFU**; former name of **TAINAN**, the capital of southern Formosa, on the west coast of the island under 23° North latitude.
- TAJIMA**; same as **TAZIMA**.
- TAKAO**; port in south Formosa, south of Taiwanfu.
- TAKOW**; same as **TAKAO**, Formosa.
- TAMBA**; province at the base of the western extension of Hondo, not far from Kyōto and Ōsaka.
- TAMSUI**; seaport near the northern extremity of Formosa.
- TANEGA SHIMA**; one of the islands of the Ōsumi group, Riu Kiu.
- TANGO**; province near the base of the western extension of Hondo, facing the Sea of Japan.
- TAZIMA**; province on the western extension of Hondo, west of Tango, and facing the sea of Japan.
- TCHI-TCHI-SHIMA**; same as **CHICHI SHIMA**.
- TOKARA GROUP**; a series of small islands in the Riu Kiu Archipelago between Yaku shima and Amami-ō-shima. The southernmost island of this group is Yoko shima.
- TOKUCHIMURA**; village on the west side of Okinawa shima, the type-locality of *Callophis boettgeri* Fritze.
- TOKUNO SHIMA**; one of the islands of the Amami-ō-shima subgroup, Riu Kiu.
- TŌKYŌ**; capital of Japan.
- TORI SHIMA**; small, volcanic island, lying due west of Tokuno shima, of the Amami-ō-shima subgroup, but geologically reckoned to the Okinawa subgroup, Riu Kiu.
- TOSA, or TOZA**; province in Shikoku, occupying nearly the entire southern coast of the island.
- TOTOMI**; province on the south coast of central Hondo, between Suruga Bay and Owari Bay.
- TOYOKAGAWA**; river in the province of Tazima.
- TOZA**; Same as **TOSA**.
- TSUKUBA MOUNTAIN**; northeast of Tokyo, in the province of Hitachi.
- TSU-SHIMA**; two large islands in the strait separating Japan from Korea.
- UNZEN-DAKE**; same as **MOUNT ONSEN**.
- VRIES ISLAND**; same as **Ō-SHIMA, IDZU**.
- WAHEIZAN**; Japanese name for Hoa-pin-su.
- YAEYAMA ISLAND**; same as **ISHIGAKI SHIMA**.
- YAEYAMA ISLANDS**; subgroup of the Saki shima group, consisting of the Ishigaki shima, Iriomote shima, Yonaguni shima, Hateruma shima, etc. Sometimes used in the same sense as Saki shima, thus including *M'yakoshima*.
- YAKU SHIMA**; one of the islands of the Ōsumi group, Riu Kiu.
- YAMAGAWA**; town in the province of Satsuma, at the western entrance to Kagoshima Bay near the south end of Kiūsū.
- YAMASHIRO**; province in southwestern Hondo, in which Kyōto is situated.
- YAMATO**; province in the center of the large peninsula of southwestern Hondo, between the Bays of Owari and Ōsaka.
- YAYEYAMA**; same as **YAEYAMA**.

- YEDO**, or **YEDDO**; old name for Tokyo. By a curious mistake the locality Yedo on old labels, or in museum catalogues has several times been corrected to Yezo, causing great confusion.
- YEZO**, or **YESSO**; northernmost of the four principal islands of Japan. It is a geographical term applied to the island alone and is consequently not exactly synonymous with Hokkaidō.
- YODOGAWA**; river in southwestern Hondo, draining Lake Biwa, and emptying into the Bay of Ōsaka.
- YOKOHAMA**; large city at the entrance of Tōkyō Bay.
- YOKO SHIMA**, or **YOKOATE SHIMA**; southernmost island of the Tokara group, Riu Kiu.
- YONAGUNI SHIMA**; westernmost island of Yaeyama group, Riu Kiu.
- YORON SHIMA**; southernmost island of the Amami-ō-shima subgroup, Riu Kiu.
- YUMOTO**; watering place, near Nikko, north of Chūzenji Lake, about 5,000 feet above the sea.

EXPLANATION OF PLATES.

PLATE I.

Sketch map of Japan and adjacent territory.

PLATE II.

Figs. 1-3. *Diemictylus pyrrhogaster*, fig. 1, animal from above; fig. 2, from the side; fig. 3, from below. Natural size. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. iv, figs. 1-3.

PLATE III.

Figs. 1-3. *Hynobius nævius*, fig. 1, animal from side; fig. 2, anterior portion of animal from below; fig. 3, from above. Natural size.
4-6. *Hynobius nebulosus*, fig. 4, anterior portion of animal from below; fig. 5, from above; fig. 6, entire animal from side. Natural size. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. iv, figs. 4-9.

PLATE IV.

Figs. 1-3. *Hynobius lichenatus*, fig. 1, animal from above; fig. 2, from below; fig. 3, open mouth. From Boulenger, Ann. Mag. Nat. Hist. (5), XII, 1883, pl. v, figs. 1-1b.
4-7. *Hynobius peropus*, fig. 4, animal from above; fig. 5, from below; fig. 6, head from side; fig. 7, open mouth. From Boulenger, Cat. Batr. Grad. Brit. Mus., 1882, pl. II, figs. 1-1c.
8. *Hynobius leechii*, open mouth. From Boulenger, Ann. Mag. Nat. Hist. (5), XIX, 1887, p. 67.

PLATE V.

Figs. 1-4. *Onychodactylus japonicus*, fig. 1, larva; fig. 2, fore foot; fig. 3, hind foot; fig. 4, toe with nail, from side. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. v, fig. 4.
5-6. *Geomolge fischeri*, fig. 5, adult animal from above; fig. 6, open mouth. From Boulenger, Proc. Zool. Soc. London, 1886, pl. xxxix, figs. 2 and 2b.
7-8. *Salamandrella keyserlingii*, fig. 7, animal from above; fig. 8, from side. From Strauch, Mém. Acad. Sci. St. Pétersbourg (7), XVI, No. 4, pl. II, figs. 2a-b. Natural size.

PLATE VI.

Figs. 1-3. *Onychodactylus japonicus*, adult, fig. 1, animal from side; fig. 2, from above; fig. 3, from below. Natural size. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. v, figs. 1-3.

PLATE VII.

Figs. 1-2. *Bombina orientalis*, fig. 1, animal from above; fig. 2, from below. Natural size. From Boulenger, Ann. Mag. Nat. Hist. (6), V, 1890, pl. ix, fig. 2.

PLATE VIII.

Bufo formosus. Natural size. From Boulenger, Proc. Zool. Soc. London, 1883, pl. xxiii.

PLATE IX.

Figs. 1-3. *Hyla arborea japonica*. Fig. 1, Daibutsu, Hondo; fig. 2, Tsushima. From Boulenger, Taill. Batr. Europe, pl. xv, figs. 6, 5. Fig. 3, Japan. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. iii, fig. 5.
4. *Hyla chinensis*, China. From Guenther, Cat. Batr. Sal. Brit. Mus. 1858, pl. ix, fig. c.

PLATE X.

Fig. 1. *Rana nigromaculata*, animal from above.
2-4. *Rana rugosa*, fig. 2, animal from above; fig. 3, side of head; fig. 4, animal from below. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. iii, figs. 1, 3, 4.

PLATE XI.

Fig. 1. *Rana japonica*, animal from above. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. iii, fig. 2.
2-3. *Rana amurensis*, fig. 2, animal from above; fig. 3, from below. From Boulenger, Ann. Mag. Nat. Hist. (6), V, pl. ix, fig. 1.

PLATE XII.

Figs. 1-3. *Polypedates buergeri*, fig. 1, animal from above; fig. 2, head from side; fig. 3, animal from below. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. iii, figs. 7-8.

PLATE XIII.

Figs. 1-3. *Gekko japonicus*, fig. 1, animal from above; fig. 2, from below; fig. 3, head from side. Natural size. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. ii, figs. 1-4.

PLATE XIV.

Japalura polygonata. Natural size. From Boulenger, Proc. Zool. Soc. London, 1887, pl. xvii, fig. 1.

PLATE XV.

Figs. 1-3. *Eumeces latiscutatus*, fig. 1, adult from above; fig. 2, from below; fig. 3, young from above. Natural size. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. i, figs. 1, 3, 4.

PLATE XVI.

Figs. 1-5. *Mabuya longicaudata*, fig. 1, animal from above natural size; fig. 2, top of head; fig. 3, underside of head; fig. 4, side of head, $1\frac{1}{2}$ ×natural size; fig. 5, dorsal scales, 2×natural size. From Fischer, Abh. Naturw. Ver. Hamburg, IX, pl. i, figs. 2a-e.

PLATE XVII.

- Figs. 1-2. *Sphenomorphus indicus*, fig. 1, animal from side; fig. 2, top of head of same. Natural size. From Boulenger, Cat. Liz. Brit. Mus., III, 1887, pl. xvi, fig. 1-1a.
3. *Lygosaurus pellopleurus*, animal from above. Natural size. From Boulenger, Proc. Zool. Soc. London, 1887, pl. xviii, fig. 2.

PLATE XVIII.

- Figs. 1-3. *Takydromus tachydromoides*, fig. 1, animal from above; fig. 2, from below; fig. 3, from side. Natural size. From Schlegel, Fauna Japonica, Saurii et Batrachii, pl. i, figs. 5-7.

PLATE XIX.

- Figs. 1-2. *Natrix tigrina*, fig. 1, entire animal; fig. 2, top of head. Reduced from Temminck and Schlegel, Fauna Japonica, Ophidii, pl. iv, fig. 1.

PLATE XX.

- Figs. 1-4. *Natrix pryeri*, fig. 1, head and neck from side; fig. 2, from above; fig. 3, lumbar region from above; fig. 4, from below. Natural size. From Boulenger, Proc. Zool. Soc. London, 1887, pl. xviii, fig. 3.

PLATE XXI.

- Figs. 1-2. *Elaphe quadrivirgata*, fig. 1, entire animal; fig. 2, top of head and neck. Reduced from Temminck and Schlegel, Fauna Japonica, Ophidii, pl. i, fig. 1.

PLATE XXII.

- Laticauda semifasciata*. Reduced from Temminck and Schlegel, Fauna Japonica, Ophidii, pl. x, fig. 1.

PLATE XXIII.

- Figs. 1-5. *Disteira melanocephala*. Natural size. Figs. 1-3, side, upper and lower views of head and anterior part of body; fig. 4, side view of middle of body; fig. 5, side view of tail. From Boulenger, Cat. Snakes Brit. Mus., III, 1896, pl. xv.

PLATE XXIV.

- Lapemis hardwickii*. Reduced from Jan, Icon. Ophid., livr. 41, pl. iii, fig. 1.

PLATE XXV.

- Figs. 1-3. *Enhydrina valakadyn*, fig. 1, entire animal; fig. 2, top of head; fig. 3, underside of head. Reduced from Jan, Icon. Ophid., livr. 41, pl. ii, fig. 1.

PLATE XXVI.

- Figs. 1-2. *Agkistrodon blomhoffii*, fig. 1, entire animal; fig. 2, head from side. Reduced from Temminck and Schlegel, Fauna Japonica, Ophidii, pl. vi, fig. 1.

PLATE XXVII.

- Figs. 1-4. *Trimeresurus flavoviridis*, figs. 1-3, side, upper, and lower views of head; fig. 4, side view of middle of body. From Hilgendorf, Sitz. Ber. Ges. Naturf. Fr. Berlin, 1880, pl. —, figs. 6-9.

PLATE XXVIII.

Ocadia sinensis. From Gray, Cat. Shield Rept. Brit. Mus., pl. x.

PLATE XXIX.

Figs. 1-2. *Clemmys japonica*, fig. 1, animal from above; fig. 2, from below. Reduced from Schlegel, Fauna Japonica, Chelonii, pl. VIII, figs. 2-3.

PLATE XXX.

Figs. 1-3. *Geoclemmys reevesii*, fig. 1, animal from side; fig. 2, from above; fig. 3, from below. From Gray, Cat. Shield Rept. Brit. Mus., pl. v.

PLATE XXXI.

Geoemyda spengleri. Natural size. Specimen from Ishigaki shima, in Kumamoto Higher Middle School.

PLATE XXXII.

Figs. 1-2. *Geoemyda spengleri*, fig. 1, carapace; fig. 2, plastron. Natural size. Same specimen as figured on preceding plate.

PLATE XXXIII.

Figs. 1-2. *Cyclemys flavomarginata*, fig. 1, animal from above; fig. 2, shell from below. $\frac{1}{2}$ \times natural size. From Guenther, Rept. Brit. India, pl. v, figs. A-A'.

PLATE XXXIV.

Caretta olivacea. $\frac{1}{2}$ \times natural size. From Eschscholtz, Zool. Atlas, pl. III.

PLATE XXXV.

Figs. 1-10. *Amyda japonica*. Young. Underside, to show variation of black markings on plastron. $\frac{2}{3}$ \times natural size. Nos. 23529-23538, U.S.N.M.

INDEX.

The black-faced numbers indicate generic or specific heading.

A.	Page.		Page.
abbreviatus, Hydrophis	435	albofuscus, Ophites	357, 370
Ablabes	338	albolabris, Trimesurus	481
buliodeirus	338	alleghaniensis, Coluber	307
herminæ	343	alternans, Hydrophis bicolor	439
major	338	Amblycephalidæ	255
rufodorsatus	310	Amblystoma	24
semicarinatus	340	amboinensis, Cuora	503, 505
sexlineatus	311	Emys	503
vittatus	352	Ambystoma	24
Ablepharus boutonii	225	Ambystomidae	3, 11, 24
nigropunctatus	225	Amphibia	1
pocilopleurus	225	Amphiesma	264, 286
Acanthopneuste coronata	202	flavipunctatum	288, 290
lijimæ	202	platyceps	286
Acaudata	48	stolatum	264, 280
Aceri	48	subminiatum	264
Achalinus	257, 263, 294	tigrinum	272, 278, 367
braconnieri	295, 296, 298	Amphiumoidea	3
rufescens	295, 296	anurensis, Bufo variabilis	70
spinalis	258, 294, 295, 296, 297	Rana	93, 117, 119, 121
achatina, Hyalapsia	87	Tachydromus	245
Achelata	1	Takydromus	229, 238, 245, 246, 247
Acontias	419	Amyda	514, 515, 519, 530, 531
leucostoma	449	cartilaginea	515
Actinemys	492	japonica	493, 515, 517, 525
marmorata	492	javanica	515
acutus, Cerberus	305	maackii	515, 527, 529
ægyptiacus, Trionyx	514	schlegelii	515, 516, 517, 526, 527
aer, Hypsirhina	300	sinensis	515, 516, 519, 524, 530
affinis, Agkistrodon blomhoffii	456, 461	Anabystoma	24
Laticauda	404	Aneistrodon	419
laticaudata	404	blomhoffi	457
Platurus	503	intermedius	464
Trigonocephalus	461	andersoni, Tylostrotion	12, 14
africana, Fordia	514	Andrias	3
Agamidae	163, 164, 182, 192	japonicus	6
Agamidae	163	Anguilla	255
agilis, Lacerta	251	Anguis platyrus	439
Rana	161	platurus	439
Agkistrodon	449, 461, 466	angustatus, Sphargis	485
blomhoffii	258, 450, 452, 455, 456, 457, 459, 461, 462, 463, 464	Aniliidae	255
affinis	456, 461	Anilius	260
breviceaudus	456, 459, 463	australis	260
intermedius	456, 459, 464	Anillus	260
halys	450	Anisodon	383
himalayanus	450	lilljeborgi	383
intermedius	450, 451, 452	Anisodontes	383
mokasen	449	annectens, Hyla	76
Aipysurus	413	annularis, Callophis	392
annulatus	413	Natrix	265, 266, 291
albiventer, Changulia	375	Tropidonotus	291
		annulata, Hydrophis pelamidoides	435

	Page.		Page.
annulatus, <i>Alpysurus</i>	413	bealii, <i>Clemmys</i>	493
<i>Emydocephalus</i>	413, 414	<i>Socalia</i>	492
<i>Polyodontes</i>	437	bengalensis, <i>Hydrophis</i>	457
Anoura	48	bennetti, <i>Hypsirhina</i>	392
Anthece	485	bennettii, <i>Emys</i>	489
Anura	48	<i>Enhydria</i>	300, 302
Anuril	48	berdmorei, <i>Leirus</i>	176
Aphanobranchiata	3	berezowskii, <i>Calamaria</i>	375, 376
arborea chinensis, <i>Hyla</i>	86	Berus	443
<i>Hyla</i>	76, 77, 82, 84, 85, 87, 161	<i>vulgaris</i>	146
<i>immaculata</i> , <i>Hyla</i>	76, 81, 82, 83	berus, <i>Coluber</i>	443, 444, 445
<i>japonica</i> , <i>Hyla</i>	76, 77, 78, 82, 83, 84	<i>Pelias</i>	446
<i>meridionalis</i> , <i>Hyla</i>	77	<i>Vipera</i>	443, 444, 445
<i>savignyi</i> , <i>Hyla</i>	77, 78	Biancia	182
arboreus, <i>Calamita</i>	75	<i>niger</i>	182
argus, <i>Eremias</i>	248, 249, 251	bicarinata, <i>Tiliqua</i>	214
<i>Polarces</i>	248	bicarinatus, <i>Euprepes</i>	214, 216
arguta, <i>Eremias</i>	249	<i>Tiliqua</i>	214
Argyrophis	260	bicatenatus, <i>Simotes</i>	354
<i>bicolor</i>	260	bicinctum, <i>Cinetisternum</i>	514, 528
<i>truncatus</i>	260	bicolor alternans, <i>Hydrophis</i>	439
arnensis, <i>Simotes</i>	353	<i>Argyrophis</i>	260
arvalis, <i>Rana</i>	120	<i>Hydrophis</i>	439
Ascephus	50	<i>Hydrus</i>	438, 439
<i>truci</i>	50	<i>maculata</i> , <i>Hydrophis</i>	439
asiatica, <i>Bufo vulgaris</i>	66	<i>Pelamis</i>	439
asiaticus, <i>Bufo bufo</i>	59, 60, 66	<i>sinuata</i> , <i>Pelamis</i>	439
asper, <i>Bufo</i>	55	<i>variegata</i> , <i>Pelamis</i>	439
aspera, <i>Hydrophis</i>	428	<i>bilineata</i> , <i>Hurria</i>	304
Aspidocheilyda	488	bilineatus, <i>Elaphis</i>	325
Aspidoclonion	397	<i>Elaps</i>	280
<i>semifasciatum</i>	397	Bitis	443
Aspidoneetes	514, 515	blandingii, <i>Toxteodryas</i>	381
<i>sincensis</i>	521	blomhoffi, <i>Ancistrodon</i>	457
Aspidorhinus	247	<i>Halys</i>	463
Aspidorhynchus	260	<i>Trigonocephalus</i>	457
Aspilus	514	blomhoffii affinis, <i>Agkistrodon</i>	456, 461
Aspis	391	<i>Agkistrodon</i>	258,
Athece	485	450, 452, 455, 456, 457,	
Athece	484, 485	459, 461, 462, 463, 464	
Athecata	485	<i>brevicaudus</i> , <i>Agkistrodon</i>	456, 459, 463
atra, <i>Coluber quadrivirgatus</i>	328	<i>Halys</i>	457, 463
<i>Elaphis quadrivirgatus</i>	328	<i>intermedius</i> , <i>Agkistrodon</i>	456, 459, 464
<i>Naja</i>	394	<i>megaspilus</i> , <i>Trigonocephalus</i>	457
<i>naja</i>	394, 395	<i>Trigonocephalus</i>	457, 463, 464
<i>tripudians</i>	394	blumenbachi carinata, <i>Coryphodon</i>	346
atriceps, <i>Hydrophis</i>	420, 421, 422	<i>Coryphodon</i>	346
Atropis	251	blumenbachi, <i>Coluber</i>	345, 346
Atropos	465	boeiformis, <i>Elaps</i>	304, 305
Aturia	418	Boana	76
<i>lindsayi</i>	427	boans, <i>Rana</i>	76
<i>ornata</i>	418	bocourti, <i>Zamenis</i>	350
Auletris	76	boettgeri, <i>Callophis</i>	389
aurata, <i>Hyla</i>	76	<i>Hemibungarus</i>	387, 389
australis, <i>Anilius</i>	260	Boide	255
Autosauri	163	Boiga	257, 281
		<i>krapelini</i>	281
		Boiginae	256, 263, 281
		Bombina	50, 51
		<i>maxima</i>	51
		<i>orientalis</i>	51
		<i>salsa nigriventris</i>	51
		<i>pachypus</i>	51
		bombina, <i>Bombina</i>	51
		Bombinator	50
		<i>igneus</i>	51

B.

bairdii, <i>Lycodon</i>	383
baliodeirus, <i>Ablabes</i>	338
Baliopygus	93
Batrachia	1, 48
Batrachi	47
Batrachii	1, 48
Batrachus	55
Batracii	1

	Page.
Bombinator orientalis.....	51
strumosus.....	55
Bombinator.....	50
boscai, Pelonectes.....	15
Bothrodytes.....	261
Bothrops.....	466
flavoviridis.....	367, 475, 479
Botrophis.....	465
boutonii, Ablepharus.....	225
Cryptoblepharus.....	225
nigropunctatus, Cryptoblepharus.....	225
boweringii, Doryura.....	176
Hemidactylus.....	172, 176, 177
Brachyrhynchus.....	387
braconieri, aehalinus.....	295, 296, 298
Ophielaps.....	294, 295, 296, 297, 298
braminus, Eryx.....	260
Typhlops.....	280
brenchleyi, Eremias.....	248, 251
brevicaudus, Agkistrodon blomhoffii.....	456, 459, 463
brevis, Calamaria.....	375, 378
Hydrophis.....	435
buccatus, Coluber.....	444
Buergeria.....	121, 143, 144
ijimae.....	121
ishikawa.....	132
subversicolor.....	151
buergeri, Hyla.....	143, 150
Polypedates.....	144, 150, 151, 153, 154
Rana.....	151
Rhacophorus.....	151
Bufo.....	55
Bufo.....	55, 88, 95
asper.....	55
bufo.....	68, 161
bufo asiaticus.....	59, 66
gargarizans.....	59, 68
japonicus.....	60, 69
calamita.....	55, 70, 72, 161
formosus.....	57, 58, 59, 60, 64, 65, 69
japonica.....	65
japonicus.....	56, 57, 58, 60, 66
melanostictus.....	60, 72, 161
meranostictus.....	72
pratextatus.....	56, 57, 59, 60, 69
raddei.....	60, 70
sachalinensis.....	59, 65
smithi.....	59, 60, 64
variabilis amurensis.....	70
viridis.....	72, 161
vulgaris.....	55, 56, 57, 59, 66, 69
asiatica.....	66
japonica.....	66
japonicus.....	56, 57, 58, 59, 60, 66, 69
sachalinensis.....	65
bufo asiaticus, Bufo.....	59, 60, 66
Bufo.....	68, 161
gargarizans, Bufo.....	59, 68
japonicus, Bufo.....	60, 69
Bufoidea.....	54, 55
Bufoidea.....	55
Bungarus.....	257, 386, 397
caeruleus multineatus.....	398
candidus.....	398
caeruleus.....	398
multineatus.....	397

	Page.
Bungarus candidus multineatus.....	397
Bungarus ceylonicus.....	398
fasciatus.....	397
lividus.....	398
multineatus.....	386, 397, 398
semifasciatus.....	377
bungarus, Naja.....	394
C.	
Caducibranchiata.....	3
caerulea, Enhydryis.....	300
Rana.....	76
caerulescens, Hydrophis.....	419
caeruleus, Bungarus candidus.....	398
multineatus, Bungarus.....	398
Calamaria.....	257, 307, 375, 376
berezowskii.....	375, 376
brevis.....	375, 378
pavimentata.....	375, 378
pfefferi.....	375, 376, 378
septentrionalis.....	375, 376
siamensis.....	375, 376, 378
calamarius, Coluber.....	375
Calamita.....	55, 75, 76
arborescens.....	75
calamita, Bufo.....	55, 70, 72, 161
Calamita.....	1
Calamites.....	76
Calemys.....	492, 496
muhlenbergii.....	492
calligaster, Elaps.....	387
Callinia.....	514
Calliphis.....	257, 386, 387, 391
gracilis.....	391
maclellandii.....	387, 391, 392
univirgatus.....	391, 392
Callopeltis conspiciellatus.....	336
leopardina.....	307
Callophis.....	391
annularis.....	392
boettgeri.....	389
japonicus.....	387, 389
Calopeltis.....	307
cancellatum, Dinodon.....	356, 358, 365
cancellatus, Coluber.....	348
candidus, Bungarus.....	398
caeruleus, Bungarus.....	398
multineatus, Bungarus.....	397
multineatus, Bungarus.....	397
Caouana olivacea.....	507
caouana, Chelonia.....	507
Testudo.....	507
capercies, Onychoccephalus.....	260
Caretta.....	507, 510
nasuta.....	507
olivacea.....	506, 507, 509, 511
rostrata.....	511
squamata.....	511
squamosa.....	511
thunbergii.....	509
caretta, Testudo.....	507
Thalassochelys.....	507
carinata, Coryphodon blumenbachi.....	346
Elaphe.....	306
Mabuya.....	213
Phyllorhis.....	307, 308

	Page.		Page.
carinatus, Coryphodon	346, 352	chersea, Coluber	44
Eumesolon	357	Chilote	484, 488, 513
cariniferus, Trionyx	514	Chiloti	513
cartilaginea, Amyda	515	chinensis, Emys	489
cartilagineus, Trionyx	514	Eumeces	195, 196, 208, 213
caspica, Emys	492	Gecko	166
japonica, Emys	492	Hyla	76, 82, 81, 86, 87
caspius, Henhydrius	300	arborea	86
Hydus	300	Hynobius	25
cataphoranotus, Zamenis	350	immaculata, Hyla	82
Cataphractæ	483	Matouia	202, 308
Caudata	2	Platydaetylus	166
Caudati	2	Rana	94, 95
Causide	443	esculenta	95
celanops, Turdus	202	Tiliqua	306
Cenchrus	449	Tropidonotus	291
cephalo, Chelonia	507	chittal, Hydrophis	428
Cephalophis	381	Chitulia	418
cephalum, Pelturagonia	182	inornata	418
Cerachelya	488	chloris, Herpetodryas	338
Ceramopelta	514	Hydrophis	427
latirostris	514, 525	chloronota, Rana	132
Ceratophallus	264	chrysauga, Natrix	265, 286
Cerberus	300, 304	Tropidonotus	265
acutus	305	chrysolans, Turdus	302
rhynchops	305	Cinetisternum	514
russellii	305	bicinctum	514, 528
unicolor	305	cinereus, Hydus	304
cerberus, Coluber	304, 305	Cistoolemmys	501
Cereopi	2	flavomarginata	505
Cerebus	304	Cistudo lutaria	492
ceylonicus, Bungarus	398	Clemmys	488, 492, 493, 497
Chalibassia	500	bealii	493
tricarinata	500	japonica	489, 492, 493, 495
Chalinocnemis	180	nigricans	493
Changulia	375	schmackeri	495
albiventer	375	unicolor	497, 499
Chaseax	55	clinacophorus, Coluber	324
Chelone	509	Elaphe	258, 308, 324, 326
imbricata	511	cloris, Hydrophis	427
mydas	509	Cobra	443
Chelonea	483	Cobridæ	256, 442
Chelonia	49, 483, 509	cocta-i, Hemidactylus	176, 177
caouana	507	Coelognathus	514
carecta olivacea	507	novemcostatus	514, 525
cephalo	507	Coluber	257, 263, 307, 442, 443, 444
dussumierii	507	alleghaniensis	307
imbricata	511	berus	443, 444, 445
japonica	506, 509, 511	blumenbachii	345, 346
mydas	509	buccatus	444
japonica	509	calamarius	375
olivacea	507	cancellatus	348
virgata	509	cerberus	304, 305
viridis	509	chersea	441
Chelonias	509	clinacophorus	324
cheloniccephalus, Emydocephalus	413, 414, 416	conspicillatus	334
Chelonides	483	dhumna	346
Chelonii	483	dhumnades	352
Cheloniidæ	484, 506	diana	316
Chelonites	483	dione	315
Cheloptus	492	eremita	316
Chelydoideæ	488	flavescens	307
Chelynae	483	gramineus	480
Chelyra	485	guttatus	307
Chersea	443	irregularis	381
vulgaris	443, 446	korros	348

	Page.
<i>Coluber laticaudatus</i>	402, 403, 406
<i>mucotius</i>	316
<i>mucosus</i>	345
<i>natrrix</i>	263, 264
<i>obtusatus</i>	305
<i>phylophis</i>	308
<i>platycaudatus</i>	403
<i>quadrivirgatus</i>	324, 327
<i>atra</i>	328
<i>renardi</i>	445
<i>rufodorsatus</i>	310
<i>schmackeri</i>	319, 322
<i>schrenkii</i>	313
<i>sibon</i>	444
<i>stolatus</i>	280, 444
<i>teniurus</i>	319
<i>trabalis</i>	349
<i>virgatus</i>	324, 328
<i>viridiflavus</i>	349
<i>viridis</i>	481
<i>vulneratus</i>	328
Colubridæ	263
<i>colubrina</i> , <i>Hydrophis</i>	409
<i>Latienda</i>	402, 406, 408
<i>colubrinus</i> , <i>Hydrophis</i>	403, 406
<i>Hydrus</i>	406
<i>Platurus</i>	433, 406, 409
<i>Composoma quadrivirgatum</i>	328
<i>conspicillata</i> , <i>Elaphe</i>	258, 308, 334, 337
<i>Elaphe</i>	334
<i>conspicillatus</i> , <i>Callopeltis</i>	336
<i>Coluber</i>	334
<i>Elaphe</i>	334
<i>Cophias wagleri</i>	465
<i>Coptopelta</i>	514
<i>septemcostata</i>	514, 525
<i>Coriacea</i>	488
<i>coriacea</i> , <i>Dematochelys</i>	485
<i>schlegelii</i> , <i>Sphargis</i>	485
<i>Sphargis</i>	485
<i>Coriudo</i>	485
<i>coronata</i> , <i>Acanthopneuste</i>	202
<i>Coronella perspicillata</i>	334
<i>sexlineata</i>	311
<i>striata</i>	358, 365
<i>Coronelline</i>	256, 263, 306
<i>corrugata</i> , <i>Rana</i>	139
<i>corticata</i> , <i>Thalassochelys</i>	507
<i>Coryphodon</i>	345
<i>blumenbachi</i>	346
<i>carinata</i>	316
<i>carinatus</i>	316, 352
<i>korros</i>	348
<i>Costata</i>	50, 55
<i>Costati</i>	50
<i>Cosymbotus</i>	165, 178
<i>platyrus</i>	178
<i>Craspedocephalus elegans</i>	470, 471, 481
<i>craspedogaster</i> , <i>Natrix</i>	264, 268, 285, 286
<i>crassicoilis</i> , <i>Hydrophis</i>	428
<i>Crocodylini</i>	161
<i>Crossurus</i>	178
<i>Crotalide</i>	256, 257, 418, 449, 466
<i>Crotaloideæ</i>	256
<i>Crotaphitis</i>	93

	Page.
<i>cruenta</i> , <i>Rana</i>	113
<i>Cryptoblepharus</i>	193, 225
<i>boutoni</i>	225
<i>nigropunctatus</i>	225
<i>peccilopleurus</i>	163
<i>Cryptobranchiæ</i>	3
<i>Cryptobranchus</i>	3
<i>japonicus</i>	6, 7
<i>maximus</i>	7
<i>sieboldii</i>	7
<i>Cryptodira</i>	488
<i>cumingii</i> , <i>Otosaurus</i>	216
<i>Cuora</i>	503
<i>amboinensis</i>	503, 505
<i>flavomarginata</i>	503
<i>trifasciata</i>	503
<i>cuprea</i> , <i>Tisiphone</i>	449
<i>curtus</i> , <i>Hydrus</i>	435
<i>cyanocincta</i> , <i>Disteira</i>	418, 420, 425, 428, 431, 432
<i>Hydrophis</i>	428
<i>cyanocinctus</i> , <i>Hydrophis</i>	418, 428, 435
<i>Cyclemys</i>	488, 503
<i>flavomarginata</i>	489, 491, 503
<i>orbiculata</i>	503
<i>Cyclophis major</i>	337, 338
<i>nebulosus</i>	340, 341, 342, 343
<i>tricolor</i>	337
<i>cynodon</i> , <i>Dipsas</i>	381
<i>Opetiodon</i>	381
<i>Cynops</i>	15
<i>pyrrhogaster</i>	16
<i>suberistatus</i>	16
<i>Cystoclemmys</i>	503
<i>flavomarginata</i>	503

D.

<i>Dactylonyx</i>	42
<i>Dactyloperus</i>	180
<i>insulensis</i>	180
<i>Damonia</i>	496, 497
<i>macrocephala</i>	496
<i>reevesii</i>	497
<i>unicolor</i>	497
<i>unicolor</i>	497, 499
<i>delalandii</i> , <i>Onychocephalus</i>	260
<i>Delesura</i>	2
<i>Dendraspis</i>	394
<i>Dendricus</i>	143
<i>Dendrohyas</i>	76
<i>Dendromanes</i>	88
<i>dendrophila</i> , <i>Dipsas</i>	381
<i>Dermatochelyidae</i>	485
<i>Dermatochelys</i>	485
<i>coriacea</i>	485
<i>Dermocheleide</i>	484, 485
<i>Dermochelys</i>	484, 485
<i>schlegelii</i>	485
<i>Derotremata</i>	3
<i>dhumna</i> , <i>Coluber</i>	346
<i>dhumnades</i> , <i>Coluber</i>	352
<i>Zaoceys</i>	352
<i>Dinacostoidæ</i>	513
<i>diana</i> , <i>Coluber</i>	316
<i>Dianodon</i>	356
<i>Diapsida</i>	161, 162

	Page.		Page.
Dieraulax	353	drapiezii, Dipsas	381
Diemictylus	11, 15, 16	Dryophytes	76
ensicauda	16, 21, 22, 24	dussumieri, Chelonia	307
orientalis	20	Eurostus	300
pyrrhogaster	16, 18, 19, 20, 21, 23	Lepidocheilus	307
sinensis	20	dybowskii, Rana	113, 116
Diemictylus	15		
pyrrhogaster	16	E.	
Dinodon	257, 306, 356, 357	Ecaudata	47
cancellatum	356, 358, 365	Ecaudati, Polypedates	144, 153
japonicus	373	Rana	153
orientale	258, 357, 358, 372	Elaphe	257, 258, 306, 307, 388
rufozonatum	358, 364, 365	carinata	308
walli	358, 364	cinacophora	258, 308, 324, 326
rufozonatus	364	conspicillata	258, 308, 334, 337
formosana	358, 365	dione	308, 315, 452
rubstrati	372	moellendorffii	322
semicarinatum	258, 358, 366	parreysii	307
semicarinatus	366	quadrivirgata	258, 308, 327, 328, 336
septentrionale	358, 371	rufodorsata	259, 308, 310, 328
rubstrati	357, 370, 371	schmackeri	308, 322
septentrionalis	370	schrenckii	308, 313, 318, 337, 452
rubstrati	370	teniurus	308, 319, 322, 323
tessellatus	334	schmackeri	322
dione, Coluber	315	Elaphis	307
Elaphe	308, 315, 452	bilineatus	328
Elaphis	316	conspicillata	334
Diploderma	182	conspicillatus	334
polygonatum	182, 188	dione	316
Diploelma	88	quadrivirgata	328
Diplophallus	264	quadrivirgatus	328
Dipnoa	1	atra	328
Dipsadomorphus	381	interrupta	328
kraepelini	381	sauromates	308
Dipsas cynodon	381	schrenckii	313
dendrophila	381	teniurus	319
drapiezii	381	virgata	324
ferruginea	383	virgatus	313, 319, 324
fusa	381	yunnanensis	319
irregularis	381	Elaphrolytes	47
trigonata	381	Elapide	256, 326
Discoglossidae	50, 55	elapiformis, Python	305
Distira	400, 401, 418, 419, 430	Elapine	256, 326
cyanocincta	418, 420, 425, 428, 431, 432	Elaps	26
doliata	418	bilineatus	20
fasciata	420, 421, 427, 431	boeiformis	304, 305
godeffroyi	401, 420, 430, 431, 432, 433	calligaster	387
gracilis	419, 420, 427	maclellandii	392
major	431	maclellandii	391
melanocephala	420, 421, 422, 427, 429, 430, 431	personatus	392
orientalis	418, 421, 422	elaps, Ophiophagus	994
praescutata	434	elegans, Eumeces	194, 195, 202, 203, 207
russellii	437	Craspedocephalus	470, 471, 481
subcincta	421	Lygosoma	202
viperina	420, 434	Trimeresurus	461, 466, 467, 468, 470, 471, 473
Distira	418	Trimeresurus	470, 481
ornata	430, 431, 432	Ellipsoglossa	25
Dogania	530	mevia	26, 30
subplana	530, 531	Emmenia	492
doliata, Distira	418	grayi	492
Hydrophis	434	Emydine	488
dorsale, Leptophidium	307, 328, 367	Emydocephalus	400, 401, 413
dorsalis, Enhydridis	300	annulatus	413, 414
Takydromus	229	cheloniccephalus	413, 414, 416
Doryura bowringii	176		

	Page.		Page.
<i>Emydocephalus ijimæ</i>	413, 414	<i>Euchelonia</i>	509
<i>Emydosauria</i>	161	<i>Euchelys</i>	509
<i>Emys amboinensis</i>	503	<i>Eudipsas</i>	381
<i>bennettii</i>	489	<i>Eumeces</i>	198, 194
<i>caspica</i>	492	<i>chinensis</i>	195, 208, 213
<i>japonica</i>	492	<i>elegans</i>	194, 195, 202, 203, 207
<i>chinensis</i>	489	<i>fasciatus</i>	201
<i>japonica</i>	492, 497	<i>japonicus</i>	195
<i>palustris japonica</i>	492	<i>kishinouyei</i>	195, 196, 210, 213
<i>punctata</i>	492	<i>latiscutatus</i>	194,
<i>reevesii</i>	497	195, 196, 201, 202, 203, 207, 212	
<i>sinensis</i>	489, 497, 500	<i>okadae</i>	195, 200, 275
<i>spengleri</i>	501	<i>marginatus</i>	195, 196, 203, 205
<i>trijuga</i>	496	<i>modestus</i>	218
<i>vulgaris</i>	492	<i>pulehra</i>	202
<i>japonica</i>	492, 497	<i>quinquelineatus</i>	195, 196
<i>picta</i>	497, 499, 500	<i>japonicus</i>	195
<i>Engystoma ornatum</i>	87, 88	<i>schnelderi</i>	193
<i>pulchrum</i>	88	<i>siamensis</i>	214
<i>Engystomidae</i>	54, 55, 87	<i>Eumesodon</i>	356, 357
<i>Enhydryna</i>	402, 437	<i>carinatus</i>	357
<i>valakadien</i>	437	<i>semicarinatus</i>	356, 366, 367
<i>vikadien</i>	437	<i>euphraticus, Trionyx</i>	514
<i>Enhydria</i>	257, 299, 300, 418, 435	<i>Euprepes</i>	214
<i>bennettii</i>	300, 302	<i>bicarinatus</i>	214, 216
<i>cærulea</i>	300	<i>ruhstrati</i>	214
<i>caspicus</i>	300	<i>Euprepis</i>	213
<i>dorsalis</i>	300	<i>longicaudata</i>	214, 215
<i>hardwickii</i>	435	<i>euproctus, Typhlops</i>	260
<i>palustris</i>	300	<i>Eurastus</i>	300
<i>piscator</i>	300	<i>dussumierii</i>	300
<i>plumbea</i>	300, 301	<i>Eurypholis</i>	337
<i>rynchops</i>	300	<i>semicarinatus</i>	337, 340, 367
<i>enhydria, Hydrus</i>	300	<i>everetti, Rana</i>	136
<i>maculata, Hypsishina</i>	302	<i>exigua, Mocoia</i>	219
<i>ensicauda, Diemictylus</i>	15, 16, 21, 24		
<i>Molge pyrrogaster</i>	21	F.	
<i>Triton</i>	21	<i>fasciata, Distera</i>	420, 421, 427, 451
<i>ensicaudatus, Molge pyrrogastra</i>	21	<i>Naja tripudians</i>	394
<i>Entacanthus</i>	337	<i>fasciatus, Bungarus</i>	397
<i>Entechinus</i>	337	<i>Eumeces</i>	201
<i>Epedaphus</i>	76	<i>Hydrophis</i>	420, 421, 422, 425, 427, 437
<i>Epidalea</i>	55	<i>Hydrus</i>	427
<i>Eremias</i>	228, 247	<i>Lepidocephalus</i>	366, 367, 368, 369
<i>argus</i>	248, 249, 251	<i>Platurus</i>	402, 403, 409, 412
<i>arguta</i>	249	<i>semifasciata, Platurus</i>	409
<i>brenchleyi</i>	248, 251	<i>fayreriana, Hydrophis</i>	436
<i>velox</i>	248	<i>Ferania</i>	300
<i>Eremiophis</i>	319	<i>ferox, Trionyx</i>	514
<i>eremita, Coluber</i>	316	<i>ferruginea, Dipsas</i>	383
<i>Eretmochelys</i>	511	<i>fischeri, Geomolge</i>	47
<i>imbricata</i>	511	<i>Platurus</i>	403
<i>squamata</i>	511	<i>fissipes, Microhyla</i>	88, 89
<i>squamosa</i>	506, 511	<i>flavescens, Coluber</i>	307
<i>Eriglossa</i>	163	<i>flavipunctatum, Amphiesma</i>	288, 290
<i>Eryma</i>	492	<i>flaviventris, Hydrophis</i>	437
<i>laticeps</i>	492	<i>flavomarginata, Cystoclemmys</i>	503
<i>erythrurus, Trionocephalus</i>	481	<i>Cuora</i>	503
<i>Trimeresurus</i>	481, 482	<i>Cyclemys</i>	489, 491, 503
<i>Eryx braminus</i>	260	<i>Cystoclemmys</i>	503
<i>eschrichtii, Typhlops</i>	260	<i>flavoviridis, Bothrops</i>	367, 475, 479
<i>esculenta, chinensis, Rana</i>	95	<i>Sachesis</i>	475
<i>japonica, Rana</i>	94	<i>Trimeresurus</i>	466, 467, 470, 474, 475, 476
<i>marmorata, Rana</i>	95	<i>Fordia</i>	514
<i>nigromaculata, Rana</i>	95	<i>africana</i>	514
<i>Rana</i>	93, 94, 95, 96, 98, 161	<i>formosana, Dinodon rufozonatus</i>	358, 365

	Page.		Page.
formosanus, <i>Holarchus</i>	354	<i>guttata</i> , <i>Geoclemmys</i>	492, 496
<i>Simotes</i>	354	<i>guttulatus</i> , <i>Lamprosaurus</i>	194
<i>Takydromus</i>	229, 232, 235 , 238	<i>Gymnopus</i>	314
formosus, <i>Bufo</i>	57, 58, 59, 60 , 64, 65, 69		
<i>Trionyx</i>	514	H.	
<i>Fornicata</i>	483	<i>habereri</i> , <i>Natrix</i>	291
<i>franklinii</i> , <i>Onychophis</i>	260	<i>hainanensis</i> , <i>Simotes</i>	354
<i>frenatus</i> , <i>Hemidactylus</i>	163, 172 , 178	<i>haje</i> , <i>Naja</i>	394
<i>Trionyx</i>	530	<i>haji</i> , <i>Naja</i>	394
<i>fusca</i> , <i>Dipsas</i>	381	<i>Halys</i>	449
<i>Rana</i>	113	<i>blomhoffi</i>	463
<i>fuscus</i> , <i>Zapyrus</i>	352	<i>blomhoffii</i>	457, 463
		<i>intermedia</i>	464
G.		<i>intermedius</i>	464
<i>gargarizans</i> , <i>Bufo bufo</i>	59, 68	<i>halys</i> , <i>Agkistrodon</i>	450
<i>Gecko</i>	165	<i>Trigonocephalus</i>	449
<i>chinensis</i>	166	<i>Hamadryas</i>	394
<i>japonicus</i>	165	<i>hannah</i>	394
<i>marginatus</i>	178	<i>hamiltonii</i> , <i>Geoclemys</i>	496, 497
<i>swinhonis</i>	166	<i>hannah</i> , <i>Hamadryas</i>	394
<i>Gecus</i>	165	<i>Haplophoa</i>	161
<i>Gehyra intermedia</i>	166	<i>hardwickel</i> , <i>Hydrophis</i>	436
<i>mutilata</i>	180	<i>hardwicki</i> , <i>Hydrophis</i>	436
<i>Gekko</i>	165	<i>hardwickii</i> , <i>Enhydris</i>	435
<i>japonicus</i>	165, 166, 176	<i>Hydrophis</i>	435
<i>swinhonis</i>	166, 169	<i>Hypsirhina</i>	301
<i>Gekkonidae</i>	163, 164	<i>Lapemis</i>	401, 435 , 436
<i>Gekkonoidea</i>	163	<i>Hemibungarus</i>	257, 386, 387 , 391
<i>Geoclemmys</i>	492, 497	<i>boettgeri</i>	387, 389
<i>guttata</i>	492, 496	<i>japonicus</i>	387 , 389
<i>Geoclemys</i>	488, 496 , 497	<i>Hemidactylus</i>	165, 172
<i>hamiltonii</i>	496, 497	<i>bowringii</i>	172, 176 , 177
<i>muhlenbergii</i>	496	<i>coctei</i>	176, 177
<i>pulchella</i>	496	<i>frenatus</i>	163, 172 , 178
<i>reevesii</i>	489, 493, 496, 497	<i>inornatus</i>	172
<i>seba</i>	496	<i>longiceps</i>	172
<i>subtrijuga</i>	496	<i>marmoratus</i>	175
<i>Geoemyda</i>	488, 500	<i>mutilatus</i>	180
<i>spengleri</i>	489, 500, 501	<i>nanus</i>	166
<i>Geoemys</i>	500	<i>navarri</i>	180
<i>Geomolge</i>	25, 47	<i>peronii</i>	180
<i>fischeri</i>	47	<i>pristiurus</i>	180
<i>gigas</i> , <i>Salamandra</i>	7	<i>pumilus</i>	176
<i>Glossolega</i>	12	<i>variegatus</i>	180
<i>Glyptemys</i>	492, 496	<i>herminae</i> , <i>Liopeltis</i>	37 , 343
<i>insculpta</i>	492	<i>Herpetodryas chloris</i>	338
<i>godeffroyi</i> , <i>Disteira</i>	401, 420, 430 , 431, 432, 433	<i>tricolor</i>	337
<i>Hydrophis</i>	430	<i>herzi</i> , <i>Simotes</i>	311
<i>Gomphopelta</i>	514	<i>Heteromorpha</i>	48
<i>officina</i>	514, 525	<i>Hierophis</i>	349
<i>Gongylosoma</i>	338	<i>himalayanus</i> , <i>Agkistrodon</i>	450
<i>Gonyodipsas</i>	381	<i>Hinulia</i>	216
<i>gracilis</i> , <i>Calliophis</i>	391	<i>indica</i>	216
<i>Disteira</i>	419, 420, 427	<i>indicum</i>	216
<i>Hydrophis</i>	427	<i>naevia</i>	216
<i>Hydrus</i>	418, 427	<i>Holarchus</i>	257, 307, 353
<i>Microcephalophis</i>	418, 427	<i>formosanus</i>	354
<i>Rana</i>	127	<i>trinotatus</i>	353
<i>Gradientia</i>	2	<i>violaceus</i>	354
<i>graminea</i> , <i>Lachesis</i>	481	<i>holsti</i> , <i>Rana</i>	93, 105 , 111
<i>gramineus</i> , <i>Coluber</i>	480	<i>Tachydromus</i>	239, 242, 243
<i>Lachesis</i>	480	<i>homalocephala</i> , <i>Ptychozoon</i>	170
<i>Trimeresurus</i>	466, 480	<i>homalocephalum</i> , <i>Ptychozoon</i>	170
<i>gratiosa</i> , <i>Hyla</i>	76	<i>Homalopsis</i>	256, 263, 299 , 381
<i>grayi</i> , <i>Emmema</i>	492	<i>molurus</i>	366
<i>guichenoti</i> , <i>Mocoo</i>	218	<i>plumbea</i>	300

	Page.
Homalopsis schneideri	305
schneiderii	305
Homolepida	221
pellopleurum	222
Homomorpha	2
Hoplobatrachus reinhardtii	94
horsfieldii, Platydaetylus	170
Pteropleura	170
Ptychozoon	170
Huria	304
Hurria	257, 299, 300, 304
bilineata	304
rynehops	304
schneideriana	304
Hurrianus	304
Hyas	76
Hydra	438
Hydrinae	256, 257, 386, 400
Hydrophilopsis	263
Hydrophis	418, 419, 436, 438, 439
abbreviatus	435
aspera	428
atriceps	420, 421, 422
bengalensis	437
bicolor	439
alternans	439
maculata	439
brevis	435
caerulescens	419
chittal	428
chloris	427
cloris	427
colubrina	409
colubrinus	403, 406
crassicolis	428
cyanocincta	428
cyanocinctus	418, 428, 435
doliata	434
fasciatus	420, 421, 422, 425, 427, 437
fayreriana	436
flaviventris	437
godeffroyi	430
gracilis	427
hardwickel	436
hardwicki	436
hardwickii	435
jayakari	434
kadell-nagam	427
leprogaster	427
loreatus	401
melanocephalus	421
microcephala	427
microcephalus	427
pelamidoides	435, 436
annulata	435
pelamis	439
pelamoides	436
phipsoni	428
platara	439
plumbea	434
problematicus	436
schistosa	437
schistosus	437
striata	428
striatus	425
subannulata	428

	Page.
Hydrophis subfaciata	437
sublævis melanocephala	421
tabrobanica	428
trachyceps	428
viperinus	434
westernmani	428
westernmanni	428
Hydrosalamandra	4
siboldi	7
Hydroscopes	25
Hydrus	300, 402, 419, 438
bicolor	438, 439
caspius	300
cinereus	304
colubrinus	406
curtus	435
enhydrius	300
fasciatus	427
gracilis	418, 427
piscator	288
platurus	438, 439
rynchops	304
valakadyi	437
hydrus, Natrix	445
Hyla	75, 76, 143
annectens	76
arborea	76, 77, 82, 84, 85, 87, 161
chinensis	86
immaculata	76, 81, 82, 83
japonica	76, 77, 78, 82, 83, 84
meridionalis	77
savignyii	77, 78
aurata	76
buergeri	143, 150
chinensis	76, 82, 84, 86, 87
immaculata	82
gratiosa	76
japonica	77, 84, 85
leucomystax	157
palmata	76, 143
quadrilineata	157
regilla	76
sexvirgata	157
simplex	76, 81
stephani	76, 77, 81, 84, 85
versicolor	76
viridis	75, 77
japonica	77
Hylaplesia	87
achatina	87
Hylaria	76
Hyllidæ	54, 55, 75
Hylliola	76
Hylorana longipes	157
Hynobius	24, 25
leechii	25, 29
lichenatus	25, 33, 35, 36
nævius	25, 26, 29, 31, 32, 33
nebulosus	25, 30, 32
nigrescens	25, 24, 36
peropus	25, 32, 36
Hypsirhina	300
aer	300
bennetti	302
bennettii	302
enhydrius maculata	302

	Page.		Page.
Hypsihrina hardwickii	301	japonica, Bufo	65
maculata	302	vulgaris	66
plumbea	300	Chelonia	506, 509, 511
sieboldii	300	mydas	509
Hypsiscopus	300	Clemmys	489, 492, 493, 495
plumbea	300	Emys	492, 497
I.			
Iapalura	182	caspica	492
Ida	514	palustris	492
ornata	514	vulgaris	492, 497
Idiophides	254	Hyla	77, 81, 85
Igneus, Bombinator	51	arborea	76, 77, 78, 82, 83, 84
Ijima, Acanthopneuste	202	viridis	77
Buergeria	121	Lacerta	42
Emydocephalus	413, 414	ornativentris, Rana	108
Rana	93, 121, 144	Rana ... 93, 97, 104, 107, 108, 111, 114, 115, 116, 117, 119, 157, 275	94
Ilyrica, Vipera	433	esculenta	94
Imbricata, Chelone	511	temporaria	94, 107
Chelonia	511	Salamandra	42, 43
Eretmochelys	511	Testudo	509
Testudo	511	japonicus, Andrias	6
immaculata, Hyla arborea	76, 81, 82, 83	Bufo	56, 57, 58, 60, 66
chinensis	82	bufo	60, 69
immaculiventris, Triton suberistatus	16, 18	vulgaris	56, 57, 58, 59, 60, 66
inconspicuus, Typhlops	260	Callophis	387, 389
indica, Hinulia	216	Cryptobranchus	6
indicum, Lygosoma	216	Dinodon	373
indicus, Sphenomorphus	216	Eumeces	185
inornata, Chitulia	418	quinquelineatus	185
inornatus, Hemidactylus	172	Gecko	163
insepulta, Glyptemys	492	Gekko	165, 166, 176
insulensis, Dactyloperus	180	Hemibungarus	387, 389
intermedia, Gehyra	166	Ixalus	155, 157
Halys	464	Megalobatrachus	6, 11
intermedius, Agkistrodon	450, 451	Onychodactylus	42, 43, 47
blomhoffii	456,	Ophites	357, 373
Ancistrodon	459, 464	Platydaetylus	165
Halys	464	Polypedates	144, 155
Trigonocephalus	464	Salamandra	42
interrupta, Elaphis quadrivirgatus	328	Tachydromus	289
irregulare, Triglyphodon	381	Tachysaurus	236, 239
irregularis, Coluber	381	Trionyx	515
Dipsas	381	stellatus	515
irrorata, Laudemania	514, 525	Triton	6
ishikawa, Buergeria	132	javanica, Amyda	515
Polypedates	58	javanicus, Trionyx	514, 515, 329
Rana	94, 132, 144	jayakari, Hydrophis	434
Isodactylum	37	jerdonii, Kerilla	418
schrenkii	38	Trimeresurus	468, 470
wosnessenskyi	38	johanseni, Rana muta	113
Isola	514	K.	
peguensis	514	kadell-nayam, Hydrophis	427
Ixalus	155	Kerilla	418
japonicus	155, 157	jerdonii	418
J.			
jamori, Platydaetylus	166	keyserlingi, Salamandrella	38
Japalura	182, 192	tridactyla, Salamandrella	38
mitsukurii	182, 184, 187, 190	keyserlingii, Salamandrella	37
polygonata	182, 183, 184, 185, 187, 188, 192	khaslensis, Natrax	264, 268, 285
swinhonis	182, 183, 184, 185, 189, 190, 192	kishinouyei, Eumeces	195, 210, 213
variegata	182	kolombatovici, Bombina salsa	51
yunnanensis	187	korros, Coluber	348
japonica, Amyda	493, 515, 517, 525	Coryphodon	348
		Ptyas	345, 348
		Zamenis	348
		krapelini, Boiga	381

	Page.
krapelini, <i>Dipsadomorphus</i>	381
kraussi, <i>Onychochelys</i>	511
kuhlii, <i>Ptychozoon</i>	170
kuhlii, <i>Rana</i>	139
L.	
Labiata	513
<i>labyrinthica</i> , <i>Scaptophryne</i>	88
Lacerta	228, 251
<i>agilis</i>	251
<i>japonica</i>	42
<i>platyuros</i>	178
<i>schneideriana</i>	178
<i>tachydromoides</i>	238
<i>thunbergii</i>	43
<i>vivipara</i>	251, 252
Lacerte	162
Lacertidae	163, 164, 228
Lacertilia	163
Lacertina	162
Lacertoidea	163
Lachesis	466
<i>flavoviridis</i>	475
<i>graminea</i>	481
<i>gramineus</i>	480
<i>lutea</i>	470
<i>luteus</i>	470
<i>mucrosquamatus</i>	467, 470, 471
<i>okinavensis</i>	479
<i>lævis</i> , <i>Psilognathus</i>	514, 525
Laminifera	484, 488
Lampropholis	218
Lamprosaurus	194
<i>guttulatus</i>	194
lanaria , <i>Mauremys</i>	492
Landemania	514
<i>irrorata</i>	514, 525
Landemania perocellata	525
Languhina	381
Lapemis	300, 400, 402, 435
<i>hardwickii</i>	401, 435, 436
<i>loreatus</i>	435
larvata , <i>Naja lutescens</i>	394
laterale , <i>Leiolopisma</i>	163, 218
<i>modesta</i> , <i>Lygosoma</i>	218
<i>reevesi</i> , <i>Leiolopisma</i>	219
<i>Lygosoma</i>	216, 219
lateralis , <i>Lygosoma</i>	218
<i>Natrix</i>	265, 274
<i>tigrina</i>	265, 278, 279
<i>Rana</i>	102, 103
<i>Scincus</i>	218
<i>Tropidonotus</i>	278
Laticauda	400, 401, 402, 413
<i>affinis</i>	404
<i>colubrina</i>	402, 406, 408
<i>laticaudata</i>	402, 404, 408
<i>affinis</i>	404
<i>muelleri</i>	402
<i>scutata</i>	402
<i>semifasciata</i>	402, 400, 410
<i>schistorhynchus</i>	410
laticaudata affinis , <i>Laticauda</i>	404
<i>Laticauda</i>	402, 401, 408
laticaudatus , <i>Coluber</i>	402, 403, 406
<i>Platurus</i>	402, 403, 406

	Page.
laticeps , <i>Eryma</i>	492
<i>Pappophis</i>	381
latirostris , <i>Ceramopelta</i>	514, 525
laticutatus , <i>Eumeces</i>	194,
195, 196, 201, 202, 203, 207, 212, 275	
<i>okada</i> , <i>Eumeces</i>	195, 200
<i>Plestiodon</i>	195
laurenti , <i>Platurus</i>	403
leberis , <i>Regina</i>	264
leechii , <i>Hynobius</i>	29
Leiolepisma	218
Leiolopisma	193, 218
<i>laterale</i>	163, 218
<i>teffairii</i>	218
Leioselasma	418
<i>striata</i>	418, 428
Leiurus berdmorei	176
leopardina , <i>Callopetalis</i>	307
Lepidocephalus	356
<i>fasciatus</i>	366, 367, 368, 369
Lepidochelys	507
<i>dussumieri</i>	507
<i>olivacea</i>	507
Lepidosauria	162
Lepidota	163
leprogaster , <i>Hydrophis</i>	427
leprosa , <i>Theloderma</i>	143
leprosus , <i>Yuen</i>	514
Leptohidium	307
<i>dorsale</i>	307, 328, 367
Leptophis trifrenatus	346
Leptotyphlopidae	255
leucomystax , <i>Hyla</i>	157
<i>Polypedates</i>	144, 157, 158, 161
<i>Polypedotes</i>	143
<i>Rhacophorus</i>	157
<i>sexvirgata</i> , <i>Rhacophorus</i>	157
leucostoma , <i>Acontias</i>	449
lichenatus , <i>Hynobius</i>	25, 33, 35, 36
lilljeborgii , <i>Anisodon</i>	383
limnocharis , <i>Rana</i>	94, 127, 128
lindsayi , <i>Aturia</i>	427
Linguata	54, 55
Liolepisma	218
<i>laterale reevesi</i>	219
Liopula	418
Liopelma	50
Liopeltis	257, 307, 337, 338
<i>hermine</i>	338, 343
<i>major</i>	338
<i>semicarinata</i>	338, 340
Liophallus	381
Liopola	418
Lipinia	218
<i>pulchella</i>	218
livida <i>Rana</i>	132
lividus , <i>Bungarus</i>	398
Lobipes	76
Lomatodactylus	165
<i>vittatus</i>	165
longicaudata , <i>Euprepis</i>	214, 215
<i>Mabuya</i>	214
longiceps , <i>Homidaetylus</i>	172
longicrus , <i>Rana</i>	93, 104
longipes , <i>Hylorana</i>	157
longissimum , <i>Ophthalmidium</i>	260

	Page.		Page.
loreatus, Hydrophis	401	maxima, Bombina	51
Lapemis	435	Salamandra	6
Loricata	161	Sieboldia	6
lumbricaleis, Typhlops	260	maximus, Cryptobranchus	7
lutaria, Cistudo	492	Megalobatrachus	6
lutea, Lachesis	470	Meeraspedota	513
lutescens larvata, Naja	394	Meditoria	280
Lachesis	470	megacephalus, Polypedates	157
luteus, Trimeresurus	470, 471, 473, 474	Megara	465
Lycodon	357, 358	Megalobatrachus	4
bairdii	383	japonicus	6
rufozonatus	358	maximus	6
Lygosaurus	193, 221, 222	sieboldi	6
pellopleurus	221, 222	megaspilus, Trigocephalus blomhoffii	457
Lygosoma	218, 221, 222	Megemys	309
elegans	202	melanocephala, Disteira	401, 420,
indicum	216	421, 422, 427, 429, 430, 431	
laterale modesta	218	sublaevis	421
lateralis	218	melanocephalus, Hydrophis	421
melanopogon	216	Microcephalophis	418, 421
nigropunctatum	218	Melanochelys	496, 500
pellopleurum	222	trijuga	500
reevesi	216, 219	melanopogon, Lygosoma	216
reevesii	219	melanostictus, Bufo	60, 72, 161
		meranostictus, Bufo	72
		mercurialis, Sphargis	485
		meridionalis, Hyla arborea	77
		microcephala, Hydrophis	427
		Thalassophis	427
		microcephalophis	418
		gracilis	418, 427
		melanocephalus	418, 421
		microcephalus, Hydrophis	427
		Microhyla	87, 88
		fissipes	88, 89
		okinawensis	88, 89
		ornata	88, 161
		undulata	89, 90
		mitsukurii, Japalura	182, 184, 187, 190
		Miura	48
		moco, Oligosoma	218
		Mocoo	218
		exigua	219
		guichenoti	218
		modesta, Lygosoma laterale	218
		modestus, Emmeceps	218
		moellendorffii, Elaphe	322
		mokasen, Agkistrodon	449
		Molgæ	3
		Molge	15, 25
		pyrrhogaster	16
		ensicauda	22
		pyrrhogastra	16
		ensicauda	22
		ensicaudatus	22
		striata	26, 43
		wolterstorffii	20
		moltus, Homalopsis	306
		Monopnoa	161
		monticola, Trimeresurus	467, 480
		mordax, Temnognathus	514, 525
		Morphuromolgaei	3
		moschatus, Rucophorus	143
		mucosus, Coluber	345
		Ptyas	345
		Zamenis	345
		microsquamatus, Lachesis	457, 470, 474
		Trigocephalus	467
M.			
maackii, Amyda	515, 527, 529		
Trionyx	516, 529		
Mabouia chinensis	202, 208		
Mabouya	214		
Mabouia longicaudata	214		
siamensis	214		
Mabuya	193, 213, 214		
carinata	213		
longicaudata	214		
maclellandi, Elaps	392		
maclellandi, Calliophis	387, 391, 392		
Elaps	391		
macrocephala, Dumotia	496		
Macrocephalus	381		
macrodon, Rana	161		
macropus, Rana	155		
maculata, Hydrophis bicolor	439		
Hypsirhina	302		
enhydris	302		
Polypedates	157		
quadrilineata, Rucophorus	157		
major, Ablabes	338		
Cyclophis	337, 338		
Disteira	431		
Liopeltis	338		
malabarica, Rana	103		
Malacopoda	1		
marginatus, Emmeceps	195, 196, 203, 205		
Gecko	178		
Plestiodon	205		
marmorata, Actinemys	492		
Rana	94, 95, 96, 98		
esculenta	95		
marmoratus, Hemidactylus	175		
martensi, Rana	108, 119		
martensii, Rana	108		
Tropidonotus	266		
Masticophis spinalis	349		
Mauremys	492		
lanaria	492		

	Page.
<i>mucrosquamatus Trimeresurus</i>	466
	467, 468, 469, 470, 471
<i>muelleri</i> , Laticauda.....	402
<i>muhlenbergii</i> , <i>Calamys</i>	492
<i>Geoclemys</i>	496
<i>multicincta</i> , <i>Bungarus candidus</i>	397
<i>multicinctus</i> , <i>Bungarus</i>	386, 397 , 398
<i>aeruleus</i>	398
<i>candidus</i>	397
<i>multifasciatus</i> , <i>Scincus</i>	213
<i>multilineatus</i> , <i>Typhlops</i>	260
<i>mutata</i> , <i>johanseni</i> , <i>Rana</i>	113
<i>Rana</i>	113
<i>Mutabilia</i>	3
<i>mutilata</i> , <i>Gehyra</i>	180
<i>mutilatus</i> , <i>Hemidactylus</i>	180
<i>Peropus</i>	180
<i>Mydas</i>	509
<i>mydas</i> , <i>Chelone</i>	509
<i>Chelonia</i>	509
<i>japonica</i> , <i>Chelonia</i>	509
<i>Mydasea</i>	509
N.	
<i>naevia</i> , <i>Ellipsoglossa</i>	26, 30
<i>Hinulia</i>	216
<i>Salamandra</i>	26
<i>navinus</i> , <i>Hynobius</i>	25, 26 , 29, 31, 32, 33
<i>Naja tripudians</i>	394
<i>sputatrix</i>	394
<i>Naja</i>	257, 386, 394
<i>atra</i>	394
<i>naja atra</i> , <i>Naja</i>	394 , 395
<i>naja oxiana</i> , <i>Naja</i>	395
<i>sputatrix</i> , <i>Naja</i>	395
<i>Naja tripudians atra</i>	394
<i>fasciata</i>	394
<i>scopinucha</i>	394
<i>unicolor</i>	394
<i>bungarus</i>	394
<i>haje</i>	394
<i>lutescens larvata</i>	394
<i>naja</i>	391, 395, 396
<i>atra</i>	394 , 395
<i>oxiana</i>	395
<i>sputatrix</i>	395
<i>namiyai</i> , <i>Rana</i>	94, 136
<i>Nannemys</i>	492, 496
<i>nanus</i> , <i>Hemidactylus</i>	166
<i>narina</i> , <i>Rana</i>	94, 134
<i>nasuta</i> , <i>Caretta</i>	507
<i>Typhlops</i>	260
<i>Natricidae</i>	256, 262, 386
<i>Natricinae</i>	256, 263
<i>Natricoidae</i>	256
<i>Natrix</i>	257, 258, 263 , 264, 290, 300, 307
<i>annularis</i>	265, 266, 291
<i>chrysurga</i>	265, 286
<i>craspedogaster</i>	264, 268, 285, 286
<i>habereri</i>	291
<i>hydrus</i>	445
<i>khasiensis</i>	264, 268, 285
<i>lateralis</i>	264, 274
<i>nuchalis</i>	265, 294
<i>pisicator</i>	241, 264, 265, 266, 288 , 290
<i>pryeri</i>	265, 266, 284 , 286
<i>stolata</i>	265, 266, 280

	Page.
<i>Natrix atolatus</i>	280
<i>swinhonis</i>	264, 265, 266, 298
<i>tigrina</i>	258, 265, 272 , 279
<i>lateralis</i>	265, 278 , 279
<i>tigrinus</i>	264
<i>torquatus</i>	264
<i>vibakari</i>	258, 264, 265, 266 , 268, 285, 268
<i>vittata</i>	264
<i>vulgaris</i>	263, 264
<i>natrix</i> , <i>Tropidonotus</i>	263
<i>Coluber</i>	263, 264
<i>navarri</i> , <i>Hemidactylus</i>	180
<i>nebulosa</i> , <i>Salamandra</i>	30
<i>nebulosus</i> , <i>Cyclophis</i>	340, 341, 342, 343
<i>Hynobius</i>	25, 30 , 32
<i>Nerodia</i>	264
<i>sipedon</i>	264
<i>Nicoria</i>	500
<i>spengleri</i>	501
<i>niger</i> , <i>Bianca</i>	182
<i>nigra</i> , <i>Pseudohaje</i>	394
<i>nigrescens</i> , <i>Hynobius</i>	25, 34 , 36
<i>nigricans</i> , <i>Clemmys</i>	493
<i>nigriventris</i> , <i>Bombina salsa</i>	51
<i>nigromaculata</i> , <i>Rana</i>	93,
94, 95, 96, 97, 98, 101, 124, 125	
<i>esculenta</i>	95
<i>nigropunctatum</i> , <i>Lygosoma</i>	218
<i>nigropunctatus</i> , <i>Ablepharus</i>	225
<i>Cryptoblepharus bontoni</i>	225
<i>nilotica</i> , <i>Trionyx</i>	514
<i>Noterophis</i>	418
<i>Notophthalmus</i>	15
<i>novemcostatus</i> , <i>Tortisternum</i>	514, 525
<i>novemcostatus</i> , <i>Coleognathus</i>	514, 525
<i>nuchalis</i> , <i>Natrix</i>	265, 294
<i>Nuda</i>	1
<i>Nycteridium</i>	178
<i>schneideri</i>	178
O.	
<i>obtusatus</i> , <i>Coluber</i>	305
<i>Ocadia</i>	488, 489
<i>sinensis</i>	489 , 493
<i>officinae</i> , <i>Gomphopelta</i>	514, 525
<i>okada</i> , <i>Eumeces latiscutatus</i>	195, 200
<i>okinavana</i> , <i>Rana</i>	93, 102 , 103
<i>okinavensis</i> , <i>Lachesis</i>	479
<i>Microhyla</i>	88, 89
<i>Trimeresurus</i>	466, 467, 476, 479
<i>Oligosoma</i>	218
<i>moco</i>	218
<i>olivacea</i> , <i>Cauana</i>	507
<i>Caretta</i>	506, 507 , 509, 511
<i>Chelonia</i>	507
<i>caretta</i>	507
<i>Lepidochelys</i>	507
<i>Thalassochelys</i>	507
<i>Onchodactylus</i>	36, 47
<i>Onychocephalus</i>	260
<i>capensis</i>	260
<i>delalandii</i>	260
<i>Onychochelys</i>	511
<i>kraussi</i>	511
<i>Onychodactylus</i>	25, 42
<i>japonicus</i>	42 , 43, 47
<i>schlegelii</i>	42, 43

	Page.		Page.
Onychophis	260	pavimentatus, Scincus	193
franklinii	260	peguensis, Isola	514
Onychopus	42	pelamidoidea annulata, Hydrophis	435
Onycopus sieboldii	43	Hydrophis	435, 436
Opetiodon	381	pelamidoidea, Hydrophis	435
cynodon	381	Pelamis	419, 438
Ophes	254	bicolor	439
Ophides	254	sinuata	439
Ophidia	254	variegata	439
Ophidii	254	ornata	439
Ophidosaurii	162	schneideri	439
Ophielaps	294	pelamis, Hydrophis	439
braconneri	294, 295, 296, 297, 298	pelamoides, Hydrophis	436
Ophiophagus	394	Pelamyoidea	435
elaps	394	Pelamyx	438, 439
Ophisaurus	255	Pelias	446
Ophites	356, 357	berus	446
albofuscus	357, 370	pellopleurum, Homolepida	222
japonicus	357, 373	Lygosoma	221
orientalis	372, 373	pellopleurus, Lygosaurus	221, 222
ruhrstrati	370	Pelodiscus	514
septentrionalis	370	Pelonectes	15
subcinctus	357	Pelophylax	93
Ophthalmidium	260	Peltocheilydia	512
longissimum	260	Pelturagonia	182
tenue	260	cephalum	182
orbiculata, Cyclemys	503	Peripia	180
orientale, Dinodon	258, 357, 358, 372	perocellata, Landemania	525
orientalis, Bombina	51	Tyse	521
Bombinator	51	perocellatus Trionyx	521, 526
Diemictylus	20	peronii, Hemidactylus	180
Disteira	418, 421, 422	Peropus	165, 180
Ophites	372, 373	nutilatus	180
Tropidonotus	278, 279, 280	peropus, Hynobius	25, 32, 36
ornata, Aturia	418	Perostia	483
Distira	430, 431, 432	personatus, Elaps	392
Ida	514	pfefferi, Calamaria	375, 376, 378
Microhyla	88, 161	Phaneroglossa	54
Pelamis	439	phaneroglossa, Rana	54
ornativentris, Rana japonica	108	Philautus	136
ornatum, Eugystoma	87, 88	phipsoni, Hydrophis	428
Osearia	514	Pholidota	161
swinhoei	514	Phragmitophis	337
Otoeryptis	191, 192	Phryne	55
Otosaurus	216	Phrynoidis	55
cumingii	216	Phyllophis	307
owstoni, Parus	202	carinata	307, 308
Polypedates	144, 149	phyllophis, Coluber	398
oxiana, Naja naja	395	picta, Emys vulgaris	497, 499, 500
Tommyris	394	piscator, Euhydrix	300
Oxyglossus	92	Hydrus	278
		Natrix	244, 264, 265, 266, 288, 292
		Tropidonotus	278
		plancyi, Rana	93, 101
		platura, Anguis	439
		Hydrophis	439
		platurus, Anguis	439
		Platurus	402
		affinis	403
		colubrinus	403, 406, 409
		fasciatus	402, 403, 409, 412
		semifasciata	409
		fischeri	406
		laticaudatus	402, 403, 403
		laurenti	403
		schistorhynchus	409

P.

pachypus, Bombina salsu	51
palmata, Hyla	76, 143
palmatus, Racomphorus	143
palustris, Euhydrix	300
japonica, Emys	492
Pantherophis	307
Pappophis	381
laticeps	381
Paradiacostioidea	488
parreysii, Elaphe	307
Parus owstoni	202
varius	202
pavimentata, Calamaria	375, 378

	Page.
Platurus, semifaciatus.....	409
platurus, Hydrus.....	438, 439
platycaudatus, Coluber.....	403
platyceps, Amphiesma.....	286
Platydaetylus.....	165
chineseus.....	166
horstfieldii.....	170
jamori.....	166
japonicus.....	165
yamori.....	166
Platypeltis.....	514
platyuros, Lacerta.....	178
Platyurus.....	178
platyurus, Cosymbotus.....	178
Stellio.....	178
Plestiodon.....	193
latiscutatus.....	195
marginatus.....	205
pulchrum.....	208
quinquellineatum.....	195
sinense.....	208
Plethodontidae.....	11
Pleurodeles.....	12
Pleurodira.....	488
Pleuropoda.....	254
Plistodon.....	194
plumbea, Enhydris.....	300 , 301
Homalopsis.....	300
Hydrophis.....	434
Hypsiscopus.....	300
Pneumatromolgaei.....	162
Pnoëpus.....	172
Podarces argus.....	248
Polydotes.....	437
polygonata, Japalura.....	182, 183, 184, 185, 187, 188 , 192
polygomatum, Diploderma.....	182, 188
polygrammicus, Typhlops.....	260
Polyodonta.....	437
Polyodontes.....	437
annulatus.....	437
Polypedates.....	80, 93, 121, 143 , 144, 155
buergeri.....	144, 150 , 151, 153, 154
eiffingeri.....	144, 153
ishikawae.....	58
japonicus.....	144, 155
leucomystax.....	144, 157 , 158, 161
maculatus.....	157
megacephalus.....	157
owstoni.....	144, 149
rugosus.....	143, 157
schlegelii.....	77, 144, 145 , 147, 148, 149
viridis.....	144, 147 , 149
Polypedetes.....	143
Polypedotes.....	143, 144
leucomystax.....	143
porosa Rana.....	95
Tomopterna.....	95
Poropoda.....	163
Potamochelys.....	514
præcutata, Distira.....	434
prætextatus, Bufo.....	56, 57, 59, 60, 69
Prædenia.....	162
pristiurus, Hemidaetylus.....	180
problematicus, Hydrophis.....	136
præcipleurus, Cryptoblepharus.....	163
Proteocordylus.....	3

	Page.
Proterodon.....	307
tessellatus.....	307, 334
pryeri, Natrix.....	265, 266, 284 , 286
Tropidonotus.....	284
Psammodynastes.....	257, 383
pulverulentus.....	381, 383 , 385
Psammophis pulverulenta.....	383
Pseudo-boa.....	397
Pseudohaje.....	394
nigra.....	394
Pseudosalamandra.....	25
Pseudosauria.....	3
Pseudotyphlops.....	260
Psiloderma.....	1
Psilognathus.....	514
lavis.....	514, 525
Pteropleura.....	170
horstfieldii.....	170
Ptyas.....	257, 306, 345
korros.....	345, 348
mucosus.....	345
Ptychozoon.....	165, 170
homalocephala.....	170
homalocephalum.....	170
horstfieldii.....	170
kuhli.....	170
Ptychozoum.....	170
pulehella, Geoclemys.....	496
Lipinia.....	218
pulehra, Eumeces.....	202
pulchrum, Engystoma.....	88
Plestiodon.....	208
pulverulenta, Psammophis.....	383
pulverulentus, Psammodynastes.....	381, 383 , 385
pumilus, Hemidaetylus.....	176
punctata, Emys.....	492
puniceus, Trigonocephalus.....	465
purpurascens, Simotes.....	353
Pygomolgaei.....	48
pyrrhogaster, Cynops.....	61
Diemietylus.....	15,
16 , 18, 19, 20, 21, 23	
ensicauda, Molge.....	21
Molge.....	16
Triton.....	16
typica, Triton.....	16
pyrrhogastra ensicauda, Molge.....	21
ensicaudatus, Molge.....	21
Molge.....	16
Salamandra.....	16
Pythomorpius.....	300
Python.....	307
clapiformis.....	305
rhynchops.....	305
Pythonide.....	443
Pyxielemmys.....	503

Q.

quadrilineata, Hyla.....	157
Rhacophorus maculatus.....	157
quadrivirgata, Eiaphe.....	258, 308, 327 , 328, 336
quadrivirgatum, Comptosoma.....	328
quadrivirgatus atra, Coluber.....	328
Elaphis.....	328
Coluber.....	324, 327
Elaphis.....	328

	Page.		Page.
quadrivirgatus Interrupta, Elaphis	328	Rana rugosa	94, 95, 96, 99, 123, 124, 125, 192
quincunciatus, Tropidonotus	288	rugulosa	139
quinquelineatum, Plestiodon	195	swinhoana	94, 132
quinquelineatus, Eumeces	195, 199	sylvatica	107
japonicus, Eumeces	195	temporaria	93, 106,
Seincus	195	107, 111, 113, 114, 116, 117, 119, 161	
		japonica	95, 107
R.			
Racophorus	143	tigerina	94, 95, 128, 138, 139
maschatus	143	tigrina	139, 161
palmatus	143	tsushimensis	93, 116
reinwardtii	143	viridis	94
schlegelii	145	vittigera	127, 139
raddel, Bufo	60, 70	Ranacea	1
Rafetus	514	Ranae	47
Ramphotyphlops	260	phaneroglossae	54
Rana	93, 95, 121, 132, 144, 155, 157	Ranaria	93
agilis	161	Ranidae	54, 55, 92
amurensis	93, 47, 119, 121	Raniformia	48
arvalis	120	Ranina	88
boans	76	symetrica	88
buergeri	151	reevesi, Liolepisma laterale	219
caerulea	76	Lygosoma laterale	216, 219
chinensis	94, 95	reevesii, Damaonia	497
chloronota	132	Emys	497
corrugata	139	Geoclemys	489, 493, 496, 497
eremita	113	Lygosoma	219
dybowskii	113, 116	Tiliqua	218
effingeri	153	unicolor, Damaonia	497
esculenta	93, 94, 95, 96, 98, 161	regilla, Hyla	76
chinensis	95	Regina	264
japonica	94	leberis	264
marmorata	95	reinhardti, Rana	94
nigromaculata	95	reinhardti, Hoplobatrachus	94
everetti	136	reinwardtii, Racophorus	143
fusca	113	Racophorus	143
gracilis	127	renardi, Coluber	445
holsti	93, 105, 111	Reptilia	161
ijimae	93, 121, 144	Rhabdophis	263
ishikawae	94, 132, 144	Rhacophorus	76, 80, 143, 144
japonica	93, 97, 104, 107, 108,	buergeri	151
111, 114, 115, 116, 117, 119, 157, 275		leucomystax	157
ornativentris	108	sexvirgata	157
kuhli	139	maculatus quadrilineata	157
lateralis	102, 103	reinwardtii	143
limnocharis	94, 127, 128	schlegeli	145
livida	132	schlegelii	145
longicrus	93, 104	Rhoptoglossi	162
macrodon	161	Rhynchocephalia	161, 162
macropus	155	rhynchops, Cerberus	305
malabarica	103	Python	305
marmorata	94, 95, 96, 98	ridibunda, Rana	96
martensi	108, 119	Ristella	222
martensii	108	riukiuanus, Trimeresurus	475, 476, 477, 478
muta	113	rostrata, Caretta	511
johanseni	113	rufa, Rana	113
namiyei	94, 136	rufescens, Achallnus	295, 296
marina	94, 134	rufolorsata, Elaphe	259, 308, 310, 338
nigromaculata	93,	rufodorsatus, Ablabes	310
94, 95, 96, 97, 98, 101, 124, 125		Coluber	310
okinavana	93, 102, 103	Tropidonotus	310
planeyi	93, 101	rufo-guttata, Tiliqua	208
porosa	95	rufozonatum, Dinodon	358, 364, 365
reinhardti	94	walli, Dinodon	358, 364
ridibunda	96	rufozonatus, Dinodon	364
rufa	113	formosana, Dinodon	358, 365
		Lyceon	358

	Page.		Page.
rugosa, Rana	94, 95, 96, 99, 123, 124, 125, 129	schlegelii, Trionyx	515, 516, 526
rugosus, Polypedates	143, 157	schmackeri, Clemmys	493
rugulosa, Rana	139	Coluber	519, 322
ruhstrati, Dinodon	372	Elaphe	308, 322
septentrionale	357, 370, 371	teniurus	322
septentrionalis	370	schneideri, Homalopsis	305
Euprepes	214	Nycteridium	178
Ophites	370	Pelamis	439
Tiliqua	214	schneideriana, Hurria	304
russelli, Cerberus	305	Lacerta	178
Disteira	437	schneiderii, Eumeces	193
Simotes	353	Homalopsis	305
Tortrix	260	schrencki, Coluber	313
rynclops, Enhydria	300	schrenckii, Elaphe	308, 313, 318, 337, 452
Hurria	304	Elaphis	313
		schrenki, Isodaetylium	38
		schrenkii, Coluber	313
		Isodaetylium	38
		Scinax	76
		Scincide	163, 164, 193
		Scincus lateralis	218
		multifasciatus	213
		pavimentatus	193
		quinque-lineatus	195
		sloanei	213
		unicolor	218
		scopinucha, Naja tripudians	394
		Scotophis	307
		scutata, Laticauda	402
		Scytale	449
		Scytalus	449
		Scytina	485
		seba, Geoclemys	496
		semicarinata, Liopeltis	338, 340
		semicarinatum, Dinodon	258, 358, 366
		semicarinatus, Ablabes	340
		Dinodon	366
		Eumesodon	356, 366, 367
		Eurypholis	337, 340, 367
		semifasciata, Laticauda	402, 409, 410
		Platurus fasciatus	409
		schistorhynchus, Laticauda	410
		semifasciatum, Aspidoclonion	397
		semifasciatus, Bungarus	397
		Platurus	409
		septemcostata, Coptopelta	514, 528
		septentrionale, Dinodon	358, 371
		ruhstrati, Dinodon	357, 370, 371
		septentrionalis, Calamaria	375, 376
		Dinodon	370
		Ophites	370
		ruhstrati, Dinodon	370
		Tachydromus	232, 238
		Takydromus	229, 232, 235, 243, 247
		Serpentes	162, 254
		Serpentia	254
		serrata, Testudo	501
		sexlineata, Coronella	311
		sexlineatus, Ablabes	311
		Tachydromus	239
		Takydromus	239
		sexvirgata, Hyla	157
		Rhacophorus pleucomystax	157
		siamensis, Calamaria	375, 376, 378
		Eumeces	214
		Mabuia	214

	Page.		Page.
siboldi, Hydrosalamandra	7	Steiropphis	263
sibon, Coluber	444	stellatus-japonicus, Trionyx	515
siboldii, Megalobatrachus	6	Stello platyrus	178
Sieboldia	4	stepheni, Hyla	76, 77, 81, 84, 85
maxima	6	Sternotherus	503
sieboldia, Cryptobranchus	503	trifasciatus	503
Sieboldiana	4	Sterrichrotes	483
sieboldii, Hypsirhina	300	stolata, Natrix	265, 266, 280
Onycopus	43	stolatum, Amphiesma	284, 280
Tritomegas	6, 7	stolatus, Coluber	280, 444
Sieboldtia	4	Natrix	280
Simotes	353, 354	Tropidonotus	280
arnensis	353	Strephon	300, 304
bifasciatus	354	Strepsichrotes	254
formosanus	354	Streptostylia	162
hainanensis	354	striata, Coronella	358, 365
herzi	311	Hydrophis	428
purpurascens	353	Leioselasma	418, 428
russellii	353	Molge	26, 43
swinhonis	354	striatus, Hydrophis	425
vaillantii	354	strumosus, Bombinator	55
simplex, Hyla	76, 81	subannulata, Hydrophis	428
sinense, Plesiodon	208	subcincta, Disteira	421
sinensis, Amyda	515, 516, 519, 524, 530	subcinctus, Ophites	357
Aspidonectes	524	subcristata, Salamandra	16
Diemictylus	20	subcristatus, Cynops	16
Emys	489, 497, 500	immaculiventris, Triton	16, 18
Oealia	489, 493	Triton	16, 22
Trionyx	514, 518, 524, 529	typica, Triton	16
sinuata, Pelamias bicolor	439	subfasciata, Hydrophis	437
sipedon, Nerodia	264	sublavis melanocephala, Hydrophis	421
Siphneus	87	subminiatus, Amphiesma	264
sloanei, Scincus	213	subminiatus, Tropidonotus	263
smaragdinus, Takydromus	229, 231, 232, 236	subplana, Dogania	530, 531
smithi, Bufo	59, 60, 64	subplanus, Trionyx	530, 531
Socalla	492	subtrijuga, Geoclemys	496
bealii	492	subversicolor, Buergeria	151
Sozura	2, 3	swinhoana, Rana	94, 132
Spasmocnemis	180	swinhoei, Osearia	514
spengleri, Emys	501	Trionyx	514
Geoemyda	489, 500, 501	Tropidonotus	293
Nicroia	501	swinhonis, Gecko	166
Testudo	500, 501	Gekko	166, 169
Sphargis	485	Japalura	182, 183, 184, 185, 189, 190, 192
angustata	485	Natrix	264, 265, 266, 293
coriacea	485	Simotes	354
schlegelii	485	Tropidonotus	293
mercurialis	485	sylvatica, Rana	107
schlegelii	485	symetrica, Ranina	88
Sphenodon	161	Synapsida	161, 483
Sphenomorphus	193, 216		
indicus	216		
spinalis, Achalinus	258, 294, 295, 296, 297		
Masticophis	349		
Zamenis	349		
spiniferus, Trionyx	514		
Spondylurus	213, 214		
sputatrix, Naja naja	395		
tripudians	394		
Squamata	162		
squamata, Caretta	511		
Eretmochelys	511		
Testudo	511		
Squamosa	162		
squamosa, Caretta	511		
Eretmochelys	506, 511		
		T.	
		Tachydromus	229
		Tachybates	172
		tachydromoides, Lacerta	238
		Takydromus	229,
		231, 238, 239, 245, 246, 247, 290	
		Tachydromus	229
		amurensis	245
		formosanus	235
		holsti	239, 242, 243
		japonicus	239
		septentrionalis	232, 238
		sexlineatus	239
		tachydromoides	238

	Page.		Page.
Tachydromus wolteri	247	Tilliqua chinensis	208
Tachysaurus	229	reevesii	218
japonicus	236, 239	rufo-guttata	208
Tadpoles	159	ruhstrati	214
taniurus, Coluber	319	Tisiphone	449
Elaphe	308, 319, 322, 323	cuprea	449
Elaphis	319	Tomopterna porosa	95
schmackerti, Elaphe	322	Tomyris	394
Takydromus	228, 229	oxiana	394
amurensis	229, 238, 245, 246, 247	torquatus, Natrix	264
dorsalis	229	Tortisternum	514
formosanus	229, 232, 235, 238	novemcostatum	114, 525
septentrionalis	229, 232, 235, 243, 247	Tortrix russelii	260
sexlineatus	239	Toxicodryas	381
smaragdinus	229, 231, 232, 236	blandingii	381
tachydromoides	229,	Toxicophis	449
231, 238, 239, 245, 246, 247, 290		trabalis, Coluber	349
wolteri	229, 243, 246, 247	trachyceps, Hydrophis	428
Talpa wogura	357	Trachyhyas	143
taprobatica, Hydrophis	428	tricarinata, Chalbassia	500
Taricha	15	Testudo	501
telfairii, Leiopisma	218	tricolor, Cyclophis	337
Temnognathus	514	Herpetodryas	337
mordax	514, 525	tridactyla, Salamandrella keyserlingii	38
temporaria japonica, Rana	94, 107	trifasciata, Cuora	503
Rana	93,	trifasciatus, Sternotherus	503
106, 107, 111, 113, 114, 116, 117, 119, 161		trifrenatus, Leptophis	346
tenue, Ophthalmidium	260	Triglyphodon	381
tessellatus, Dinodon	331	irregulare	381
Proterodon	307, 334	trigonata, Dipsas	381
Testudinata	161, 162, 483	trigonocephala, Vipera	465
Testudinea	483	Trigonocephalus	449, 466
Testudines	483	affinis	461
Testudinidae	484, 488	blomholii	457
Testudinoidea	488	blomhoffii	457, 463, 464
Testudo	49	megaspilus	457
caouana	507	erythrurus	481
caretta	507	halys	449
imbricata	511	intermedius	464
japonica	509	mucrosquamatus	467
serrata	501	punicus	465
spengleri	500, 501	Trigonurus	300
squamata	511	trijuga, Emys	496
tricarinata	501	Melanocheilus	500
Thalassochelys	507	Trimeresura	465
caretta	507	Trimeresurus	419, 465, 466
corticata	507	elegans	461, 466, 467, 468, 470, 471, 473
olivacea	507	erythrurus	481, 482
Thalassophis	418	flavoviridis	466,
microcephala	427	467, 470, 474, 475, 476	
viperina	434	gramineus	466, 480
Thamnodynastes	383	jerdonii	468, 470
Thecophora	488	luteus	470, 471, 473, 474
Theloderma	143, 144	monticola	467, 480
leprosa	143	mucrosquamatus	466,
thunbergii, Carretta	509	467, 468, 469, 470, 471	
Lacerta	43	okinavensis	466, 467, 476, 479
tigerina, Rana	94, 95, 128, 138, 139	riukiuanus	475, 476, 477, 478
tigrina lateralis, Natrix	265, 278, 279	viridis	465, 481
Natrix	258, 265, 272, 279	Trimesurus	465
Rana	139, 161	albolabris	481
tigrinum, Amphilesma	272, 278, 367	elegans	470, 481
tigrinus, Natrix	264	triotatus, Holarchus	353
Tropidonotus	272, 278	Trionychia	513
Tilliqua bicarinata	214	Trionychidae	484, 514, 518
bicarinatus	214	Trionychoidea	513

	Page.		Page.
Trionyx	514, 515, 531	Tylopoda	483
egyptiacus	514	Tylotriton	11, 12
cariniferus	514	andersoni	12
cartilagineus	514	Tylositron	12
euphraticus	514	Typhlopidae	255, 256, 259
ferox	514	Typhlopoidea	256
formosus	514	Typhlops	257, 260
frenatus	530	braminus	260
japonicus	515	eschrichtii	260
javanicus	514, 515, 529	euproctus	260
maackii	516, 529	inconspicuus	260
niloticus	514	lumbicalis	260
perocellatus	524, 526	multilineatus	260
schlegelii	515, 516, 526	nasutus	260
sinensis	514, 518, 524, 529	polygrammicus	267
spiniferus	514	typica. Triton pyrrogaster	16
stellatus japonicus	515	suberistatus	16
subplanus	530, 531	Tyria	349
swinhoel	514	Tyrse	514
trionyx	514	perocellata	524
triunguis	514		
tuberculatus	524	U.	
tripudians atra. Naja	394	undulata, Microhyla	89, 90
fasciata, Naja	394	unguiculata, Salamandra	43
Naja	394	unicolor, Cerberus	305
scopinucha, Naja	394	Clemmys	497, 499
sputatrix, Naja	394	Damonia	497, 499
unicolor, Naja	394	reevesii	490
Tritomegas	4	Naja tripudians	394
sieboldii	6, 7	Scincus	218
Triton	37	univirgatus, Calliophis	391, 392
ensicauda	21	Uraeus	394
japonicus	6, 7	uralensis, Salamandrella	38, 39
pyrrogaster typica	16	Urodela	2
pyrogaster	16	Urodela	2
suberistatus	16, 21	Uropeltidae	255
immaculiventris	16, 18	Urophora	2
Triturus	16		
triunguis, Trionyx	514	V.	
Tropidolemus	465	vallanti, Simotes	354
Tropidonotus	263, 264, 300	valakadien, Enhydrina	437
annularis	291	valakadien, Enhydrina	437
chinensis	291	valakadien, Hydrus	437
chryseus	263	variabilis amurensis, Bufo	70
lateralis	278	variegata, Japalura	182
martensii	266	Pelamis bicolor	439
matrix	263	variegatus, Hemidaetylus	180
orientalis	278, 279, 280	varius, Parus	202
piscator	288	velakadien, Enhydrina	437
pryeri	284	velox, Eremias	248
quincunciatus	288	verrucosus, Tylotriton	12
rufodorsatus	310	versicolor, Hyla	76
sancti-johannis	290	vibakari, Natrix	258, 264, 265, 266, 268, 285, 286
stolatus	280	vibakari, Tropidonotus	266
subminiatus	263	vikadien, Enhydrina	437
swinhoel	293	violaceus, Holarchus	354
swinhonis	293	Vipera	412
tigrinus	272, 278	berus	443, 444, 445
vibakari	266	illyrica	433
Tropinotus	263	trigonocephala	405
truei, Ascephus	50	Viperidae	263, 442, 443
truncatus, Argyrophus	260	viperina, Dististra	420, 424
tsushimensis, Rana	93, 116	Thalassophis	434
tuberculatus, Trionyx	524	viperinus, Hydrophis	434
Turdus celanops	202	virgata, Chelonia	509
chrysolus	202	Elaphis	324

	Page		Page.
<i>virgatus</i> , Coluber	324, 328	<i>westerni</i> , Hydrophis	428
<i>Elaphis</i>	313, 319, 324	<i>westerni</i> , Hydrophis	428
<i>viridescens</i> , Triturus	15	<i>wogura</i> , Talpa	357
<i>viridiflavus</i> , Coluber	349	<i>wolteri</i> , Takydromus	229, 243, 246, 247
<i>viridis</i> , Bufo	72, 161	<i>wolterstorffi</i> , Molge	20
<i>Chelonia</i>	509	<i>wosnessenskyi</i> , <i>Isodaetyluim</i>	38
<i>Coluber</i>	481	<i>Salamandrella</i>	38
<i>Hyla</i>	75, 77		
<i>japonica</i> , <i>Hyla</i>	77	X.	
<i>Polypedates</i>	144, 147, 149	<i>Xenopeltide</i>	255
<i>Rana</i>	94		
<i>Trimeresurus</i>	465, 481	Y.	
<i>vittata</i> , <i>Natrix</i>	264	<i>yamori</i> , <i>Platydaetylus</i>	166
<i>vittatus</i> , <i>Ablabes</i>	352	<i>Yuen</i>	514
<i>Lomatodaetylus</i>	165	<i>leprosus</i>	514
<i>vittigera</i> , <i>Rana</i>	127, 139	<i>yunnanensis</i> , <i>Elaphis</i>	319
<i>vivipara</i> , <i>Lacerta</i>	251, 252	<i>Japalura</i>	187
<i>Zootoca</i>	252		
<i>vulgaris asiatica</i> , <i>Bufo</i>	66	Z.	
<i>Berus</i>	446	<i>Zamenis</i>	257, 258, 306, 349
<i>Bufo</i>	55, 56, 57, 59, 66, 69	<i>bocourti</i>	350
<i>Chersea</i>	443, 446	<i>cataphoranotus</i>	350
<i>Emys</i>	492	<i>korros</i>	348
<i>japonica</i> , <i>Emys</i>	492, 497	<i>mucosus</i>	345
<i>Bufo</i>	56, 57, 58, 59, 60, 66, 69	<i>spinalis</i>	349
<i>Natrix</i>	263, 264	<i>Zaocys</i>	352
<i>pieta</i> , <i>Emys</i>	497, 499, 500	<i>Zaocys</i>	306, 352
<i>sachalinensis</i> , <i>Bufo</i>	65	<i>dhumnades</i>	352
<i>vulneratus</i> , <i>Coluber</i>	328	<i>Zapyrus</i>	352
		<i>fuscus</i>	352
W.		<i>Zootoca</i>	251
<i>wagleri</i> , <i>Cophias</i>	465	<i>vivipara</i>	252
<i>walli</i> , <i>Dinodon rufozonatum</i>	358		
<i>walti</i> , <i>Pleurodeles</i>	12		

UX 002 021 429



UX 002 021 429



UX 002 021 429

